数学猜想与开放问题

第三卷：多源补充与历史提及记录

20,360

条补充资料，另附 9,342 个参数型开放问题实例

对照前两卷 98,151 条记录筛查 / 原始出处 / 分类导航 / 排除与保留审计

来源：OEIS 官方文本、arXiv 目录、学术论坛、研究者问题集与专业数值表

由中国跨学科学者 沈阳　　124739259@qq.com　　调度 AI 生成

整理日期：2026 年 10 月 25 日（Asia/Shanghai）

收录历史提及，不等同于仍未解决；未匹配前卷，不等同于已证明数学上独立。

01 本卷结论与计数口径

本轮继续扫描前两卷未整体处理的公开来源，并以第一卷 17,614 条、第二卷 80,537 条，共 98,151 条记录为排除基准。最终保留 20,360 条补充资料；另将 9,342 个具体参数的开放取值或存在性问题放入附录。两部分合计 29,702 条来源记录。

章节	条数	实际内容
A 明示猜想与研究问题	326	来源明示的问题块、目录整理表述或有相应复用许可的 arXiv 猜想块。
B OEIS 历史与当前提及	17,579	序列名称、注释或公式中命中猜想等关键词的原段；一条序列可能涉及多个命题。
C 论坛与文献定位	2,455	1,637 个问题帖 + 818 条定位线索；帖文不等同于公认开放问题，定位项不冒充全文。
主目录合计	20,360	用于补充前两卷的资料记录数，不是经数学语义归并后的独立猜想数。
D 参数型问题实例附录	9,342	来自六个问题族的具体参数、界限或存在性实例，不计为 9,342 个命名猜想。

“人类只要提到过”的边界

历史上已证明、已反驳、曾被撤回或仅被讨论的猜想也可能有收录价值。因此，OEIS 提及和论坛问题不因“已解决”就自动删除。反过来，提到一个猜想不等于新提出了一个猜想；也不意味着来源中的推测仍然成立或仍未解决。

无法证明已经穷尽全网、所有语言、所有纸本文献和所有历史版本。本卷的“前卷之外”意为按公开列明的规则未匹配到前卷；不同措辞的等价命题仍可能漏检，保守的题目族匹配也可能排除真正的新变体。

02 本轮实际扫描的范围

来源与取得方式	本次处理范围	需要保留的限定
OEIS 官方 Git 文本导出 [S1]	400,652 个 .seq 序列文件	仅在名称 N、注释 C、公式 F 字段检索 conjectur*、hypothesis/hypotheses、open problem/question；不以参考文献标题单独命中。
OpenConjecture 公开发布集 [S2]	4,415 个候选块	采用来源标为 real_open_conjecture 的 3,531 条进入比对；该标签由模型生成。上游的 26,747 篇论文并非本报告逐篇通读。
ConjectureBench [S3]	14,865 条记录	含 302 条编目、5,221 条扩展目录、9,342 个参数实例；此前已处理的集合镜像先排除。
图论论文/书籍定位集 [S4]	768 条论文记录 + 38 条书籍定位	来自第三方自动抽取或整理。本卷只保留定位资料，不用其自动状态评估认证数学结果。
Epoch / OEIS 元数据 [S5]	492 条历史定位	包含由模型生成的概述；本卷采用序列号、提出者、日期和记录名作为回查线索。
MathOverflow / Math Stack Exchange [S6]	2,269 个不同问题帖	完整翻页取得 MO 的 open-problems、conjectures 两标签及 MSE 的 conjectures 标签；不声称涵盖未打这些标签的讨论。
Randomstrasse 101 [S7]	18 篇博客，38 个明示问题块	按网页标注的 conjecture / openproblem 块提取；完整上下文可回到原文。

输入版本：OEIS 导出时间为 2026-10-24T03:00:17-04:00；OpenConjecture 发布集日期为 2026-07-12；ConjectureBench 参数表观察日期多为 2026-07-23。论坛和博客为本轮获取版本。上述快照时间不同，不能合称为同一时点的“当前全部未解问题”。

03 如何与前两卷排除重合

比对使用前两卷正文记录，以及第一卷中 417 条可映射回主记录的归档别名。先做固定序列号、论文号、来源页面、规范化题名和表述匹配，再做文本相似度筛查，最后合并第三卷内部的相同条目与来源线索。

阶段	数量	处理解释
进入前卷比对的候选	38,527	已去除旧集合镜像、来源标记不适合的候选及部分非数学提问。
匹配前卷或触及已知题目族	7,279	含同一 OEIS 序列、同一 arXiv 论文、相同来源页、题名、文本和已知命名猜想提及；是保守排除，不全是已证明的语义重复。
疑似重合，暂不收入正文	628	标题加权 TF-IDF 词/二元词余弦相似度不低于 0.50；保存最相近前卷编号。
第三卷内部初次去重	748	同一序列、相同文本、题名或相同源文件定位。
来源之间追加归并	54	例如论坛帖子链接到本卷同一 OEIS 项；或历史定位对应已暂缓的题目族。
题名与人工编辑规则复查	116	检查短题名是否包含在前卷较长题名中、同名提出者题目族，以及投稿流程等非数学帖子。
本卷全部保留记录	29,702	其中主目录 20,360；参数实例附录 9,342。

可复算：38,527 - 7,279 - 628 - 748 - 54 - 116 = 29,702。机器相似度不是数学等价判定器；本卷不把“未命中规则”称作已确认的新猜想。按整篇论文、同一序列或命名题目族排除会牺牲一些收录率，目的是降低重印前卷内容的风险。

04 来源贡献与新增分布

来源	保留记录数	收录形式
OEIS 官方序列文本	17,579	参见正文中的类型与完整度标记
ETH Randomstrasse 101	18	参见正文中的类型与完整度标记
ConjectureBench 编目	41	参见正文中的类型与完整度标记
OpenConjecture / arXiv	661	参见正文中的类型与完整度标记
MathOverflow	603	参见正文中的类型与完整度标记
Mathematics Stack Exchange	1,034	参见正文中的类型与完整度标记
TOPP 来源定位	3	参见正文中的类型与完整度标记
参数型问题实例	9,342	参数实例，另列附录
Epoch / OEIS 历史定位	42	参见正文中的类型与完整度标记
图论论文定位	372	参见正文中的类型与完整度标记
Bondy-Murty 附录定位	7	参见正文中的类型与完整度标记

EGRES Open 的 95 条候选也参与了扫描，但经本次比对没有作为新条目保留。发现一个新入口，并不意味着里面的问题是前两卷没有的。

本卷没有给出人类猜想总数的新估计：三个文档都是有边界的记录汇编，计数单位和文本完整度并不统一，不能把三个数字直接当成互不相同的数学猜想总量。

05 如何使用本卷

正文采用 V3-000001 等本卷编号；附录采用 V3P-000001。原来源编号、提出者或帖作者、日期、来源链接同时保留。PDF 目录可点击，阅读器书签可快速跳转；全文搜索支持英文题名、OEIS A 编号和论坛问题号。

A 章：明确表述

保留可复用的原始问题段、LaTeX 猜想块或目录整理文本。它们仍可能依赖文章其他部分的定义。来源若只是自动标注为开放，本报告不会把该标注提升为数学界确认的状态。

B 章：历史与当前提及

同一个序列条目中的所有关键词命中段合并为一条，段落顺序按名称、注释、公式分组。正文不复制整个序列页；完整可用字段保存在 PDF 附件，定义、作者更新、相关序列与外链可据此回查。不能把一条序列记录自动等同于一个独立猜想。

C 章：论坛和来源指针

论坛保留问题帖文字，未复制图像或回答，亦未核验是否已被回答推翻。来源指针则只给出论文、源文件行号或书籍位置；它明确缺少正文，不是“完整收集了该数学猜想”的证明。

D 章：参数问题

代码界、覆盖设计、差集和 Ramsey 数的每一组参数可能是一个有意义的开放任务，却未必对应人类单独命名的猜想。附录单列实例数，避免用大量参数行扩张“猜想总量”。数值界限沿用来源观察日，不代表整理日最新最佳界限。

附件 volume3_records_and_audit.zip 收录全部 29,702 条保留资料、完整排除日志、页码映射、归属说明、源文件哈希与筛查脚本。原始 Unicode 和 LaTeX 保留；极少数字形无法显示时，以 [U+编码] 明示，附件保留原字符。

06 来源、署名与内容许可

S1 OEIS Foundation Inc. 与逐条贡献者；官方 oeisdata

https://github.com/oeis/oeisdata

CC BY-SA 4.0。保留逐条来源和作者；将 N/C/F 字段节选并重新编排，属于本卷所作改动。

S2 Robert Davis / OpenConjecture 项目

https://github.com/davisrbr/conjectures-arxiv

猜想抽取和模型标签由该项目提供。正文只刊本轮选定的 CC BY / CC BY-SA 文本；其他项只保留元数据定位。

S3 Anirudha Ramesh、Shreyas Pimpalgaonkar / Bespoke Labs，ConjectureBench

https://github.com/bespokelabsai/conjecture-bench

其原创编目文字 CC BY 4.0；上游材料并未因此重新授权。参数附录保留数值事实和归属，不复制原表格图像、构造或完整文献。

S4 graph-theory-AI / graph-conjectures

https://github.com/graph-theory-AI/graph-conjectures

仅使用论文、作者、问题标签和书页位置等定位信息；抽取集与原论文均可能需要进一步核验。

S5 Epoch / LeanOpenProblems-results

https://github.com/epoch-research/LeanOpenProblems-results

OEIS 提出者、日期与声明名作为历史线索；其模型生成概述不作为新数学证明。

S6 MathOverflow、Mathematics Stack Exchange 的原帖作者

https://api.stackexchange.com/docs/questions

各帖按返回的 content_license 标注 CC BY-SA 版本；保留作者和页面链接。HTML 转为文字，图片与回答未重印。

S7 Randomstrasse 101 研究博客及逐篇作者

https://randomstrasse101.math.ethz.ch/archives/

网页声明 CC BY 4.0；抽取明示的猜想/开放问题块，保留论文与背景的回查入口。

各来源部分仍受各自许可约束，尤其 OEIS 与论坛的 BY-SA 要求；不对整本文档作统一的宽松再许可。封面的“调度 AI 生成”说明指本报告的检索整理与制版，不替代原始数学贡献的署名。

目录与页码

A 明示猜想与研究问题：326 条 10

A01 ETH Randomstrasse 101 | 18 条 11

A02 ConjectureBench 编目 | 41 条 13

A03 OpenConjecture / arXiv | 267 条 17

B OEIS 历史与当前提及：17,579 条 54

B01 A000000 - A049999 | 1,218 条 55

B02 A050000 - A099999 | 1,917 条 170

B03 A100000 - A149999 | 1,397 条 347

B04 A150000 - A199999 | 1,627 条 472

B05 A200000 - A249999 | 2,749 条 618

B06 A250000 - A299999 | 3,804 条 859

B07 A300000 - A349999 | 2,414 条 1189

B08 A350000 - A401236 | 2,453 条 1410

C 论坛与文献定位：2,455 条 1637

C01 MathOverflow | 603 条 1638

C02 Mathematics Stack Exchange | 1,034 条 1776

C03 OpenConjecture / arXiv | 394 条 2026

目录与页码（续）

C04 图论论文定位 | 372 条 2064
C05 Epoch / OEIS 历史定位 | 42 条 2101
C06 TOPP 来源定位 | 3 条 2106
C07 Bondy-Murty 附录定位 | 7 条 2107
D 参数型开放问题实例：9,342 行 2108
D01 二元常权码 | 250 行 2109
D02 二元线性码 | 965 行 2122
D03 非二元线性码 | 3,535 行 2169
D04 覆盖设计 | 1,480 行 2338
D05 阿贝尔差集 | 3,064 行 2409
D06 二色 Ramsey 数 | 48 行 2557
E 可追溯性、排除日志与后续边界 2560

A 明示猜想与研究问题：326 条

本章刊入来源问题块或有相应许可的提取文本；标签、条件和表达仍需对照原文。未在这里提供证明，也未对源状态逐条审定。

A01 ETH Randomstrasse 101 | 18 条

V3-000001 | R101-DirectionalConvexity-1

Directional Convexity (problems 34 and 35) | Conjecture 34 (Existence of wild D-convex hulls)

The D-convex hull of a finite set of points is not necessarily semialgebraic.

来源明示猜想 | source_label_at_retrieval; not independently reviewed | Chiara Meroni | 2026-02-23

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000002 | R101-DirectionalConvexity-2

Directional Convexity (problems 34 and 35) | Open Problem 35 (Conditions for semialgebraic D-convex hulls)

Determine under which conditions on $D \subset \mathbb{R}^d$ the D-convex hull of finitely many points is a semialgebraic set.

来源明示开放问题 | source_label_at_retrieval; not independently reviewed | Chiara Meroni | 2026-02-23

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000003 | R101-FramesMUBZauner-1

Mutually Unbiased Bases, ETFs, and Zauner’s Conjecture (problems 22-24) | Conjecture 22 ($\mathrm{MUB}(6)<7$)

Let $\mathrm{MUB}(d)$ denote the largest possible number of (simultaneously) mutually unbiased bases (MUBs) in $\mathbb{C}^d$. We have $\mathrm{MUB}(6)<7$.

来源明示猜想 | source_label_at_retrieval; not independently reviewed | Afonso S. Bandeira | 2025-12-14

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000004 | R101-FramesMUBZauner-2

Mutually Unbiased Bases, ETFs, and Zauner’s Conjecture (problems 22-24) | Open Problem 23 (Sum-of-Squares proof of $\mathrm{MUB}(6)<7$?)

Is there a Sum-of-Squares degree 4 proof that there are no 7 Mutually Unbiased Bases in $\mathbb{C}^6$?

来源明示开放问题 | source_label_at_retrieval; not independently reviewed | Afonso S. Bandeira | 2025-12-14

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000005 | R101-GraphBisections-1

Bisections of large random sparse regular graphs (problems 32 and 33) | Conjecture 32 (Min and Max Bisections)

Let G_d be a uniformly random d-regular graph on n nodes (n even). We have, for any $d\geq 3$: \(\frac1n\mathbb{E}\,\mathrm{MinBis}(G_d) + \frac1n\mathbb{E}\,\mathrm{MaxBis}(G_d) = \frac{d}2+o(1).\)

来源明示猜想 | source_label_at_retrieval; not independently reviewed | Afonso S. Bandeira | 2026-02-16

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000006 | R101-GraphBisections-2

Bisections of large random sparse regular graphs (problems 32 and 33) | Conjecture 33 (Max Cut and Max Bisections)

Let G_d be a uniformly random d-regular graph on n nodes (n even). We have, for any $d\geq 3$: \(\frac1n\mathbb{E}\,\mathrm{MaxCut}(G_d) = \frac1n\mathbb{E}\,\mathrm{MaxBis}(G_d) + o(1).\)

来源明示猜想 | source_label_at_retrieval; not independently reviewed | Afonso S. Bandeira | 2026-02-16

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000007 | R101-Graphmatrices-1

Sharp Bounds for Graph Matrices (problem 31) | Open Problem 31 (Sharp bounds for graph matrices)

There are functions f and g depending only on the shape α, such that \begin{equation}\label{eq:sharp_gmtx_bound} \mathbb{E}\|M_\alpha\| = \Theta\left(n^{f(\alpha)}\,(\log n)^{g(\alpha)}\right). \end{equation}

来源明示开放问题 | source_label_at_retrieval; not independently reviewed | Petar Nizic-Nikolac | 2025-12-30

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000008 | R101-HowManyDeviations-2

Did just a couple of deviations suffice all along? (problems 10-14) | Open Problem 11 (How many deviations are needed afterall?)

What is the value of the following quantity? \(\\ \sup_{n\in\mathbb{N}}\sup_{A\in\{\pm1\}^{n\times n}}\frac1{\sqrt{n}}\mathrm{disc}(A) \\\)

来源明示开放问题 | source_label_at_retrieval; not independently reviewed | Afonso S. Bandeira | 2024-12-19

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000009 | R101-HowManyDeviations-3

Did just a couple of deviations suffice all along? (problems 10-14) | Conjecture 12

Given the definitions above, we have \(\\ \limsup_{n\to\infty}\sup_{A\in\{\pm1\}^{n\times n}}\frac1{\sqrt{n}}\mathrm{disc}(A) > 1. \\\)

来源明示猜想 | source_label_at_retrieval; not independently reviewed | Afonso S. Bandeira | 2024-12-19

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000010 | R101-PaleyGraph-2

On the clique number of the Paley Graph (problems 25-29) | Conjecture 26 (1-Localization of the Paley Graph)

We have, for $p \equiv 1 \pmod{4}$ prime, $\vartheta(\overline{G_{p,1}}) \sim \sqrt{p/2}$, where $G_{p,1}$ is the 1-localization of the Paley Graph described above.

来源明示猜想 | source_label_at_retrieval; not independently reviewed | Afonso S. Bandeira | 2025-12-15

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000011 | R101-PaleyGraph-4

On the clique number of the Paley Graph (problems 25-29) | Conjecture 28 (Sum-of-Squares bound for Paley clique number)

There exists $\varepsilon>0$ such that the Sum-of-Squares relaxation of degree 4 for the clique number of the Paley graph gives a bound of $O(p^{\frac12-\varepsilon})$.

来源明示猜想 | source_label_at_retrieval; not independently reviewed | Afonso S. Bandeira | 2025-12-15

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000012 | R101-StablePhaseRetrieval-2

Injectivity and Stability of Phase Retrieval (problems 19-21) | Conjecture 20 (Stability of Phase Retrieval at $2M-1$)

There exist universal constants $C>0$ and $0<\beta<1$ such that, for any $M>1$ and $A\in\mathbb{R}^{2M-1\times M}$ a matrix for which any subset of M rows spans $\mathbb{R}^M$, \(\\ \omega(A) \leq C \max_{k\in[N]}\|A_k\|\beta^M, \\\) where A_k is the k-th row of A.

来源明示猜想 | source_label_at_retrieval; not independently reviewed | Afonso S. Bandeira | 2025-06-04

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000013 | R101-StablePhaseRetrieval-3

Injectivity and Stability of Phase Retrieval (problems 19-21) | Open Problem 21

What is the value of $\omega(A)$ for A with iid standard gaussian entries? Does it satisfy the bound in the conjecture? If so, what is β?

来源明示开放问题 | source_label_at_retrieval; not independently reviewed | Afonso S. Bandeira | 2025-06-04

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000014 | R101-clique-low-degree-1

Low-degree hardness of finding the largest clique in Erdős-Rényi graph (problems 36-38) | Open Problem 35

What are the conditions on the distributions $\mathbb P_n$ and $\mathbb Q_n$ such that if the low-degree advantage remains bounded for $D = (\log n)^{1+\Omega(1)}$ as $n\to\infty$, it rigorously implies that no polynomials of a certain degree can distinguish between $\mathbb P_n$ and $\mathbb Q_n$?”

来源明示开放问题 | source_label_at_retrieval; not independently reviewed | Anastasia Kireeva | 2026-06-05

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000015 | R101-clique-low-degree-2

Low-degree hardness of finding the largest clique in Erdős-Rényi graph (problems 36-38) | Conjecture 35

Let $G \sim G(n, 1/2)$ and $\varepsilon > 0$. For all polynomial functions $f:\{0,1\}^{n \choose 2} \to \mathbb R^n$ of degree $o(\log n)$ \[\mathbb E \max_{S: k\text{-clique in } G} \langle 1_S, f \rangle \le C \cdot \mathbb E \max_{\substack{S: |E(S)| \le (1 - \varepsilon) {k \choose 2} ,\\ |S|=k}}\langle 1_S, f \rangle,\] where $0 < C < 1$ is a constant.

来源明示猜想 | source_label_at_retrieval; not independently reviewed | Anastasia Kireeva | 2026-06-05

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000016 | R101-lovasz-circulant-1

The Lovasz number for random graphs (problems 17 and 18) | Conjecture 17

Let G be distributed as $G(n, 1/2)$. Then, \begin{equation} \mathbb{E} \vartheta(G) = (1 + o(1))\sqrt{n}. \end{equation}

来源明示猜想 | source_label_at_retrieval; not independently reviewed | Daniil Dmitriev | 2025-05-21

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000017 | R101-lovasz-circulant-2

The Lovasz number for random graphs (problems 17 and 18) | Conjecture 18

Let G be distributed as a random dense circulant graph. Then, \begin{equation} \mathbb{E} \vartheta(G) = (1 + o(1))\sqrt{n}. \end{equation}

来源明示猜想 | source_label_at_retrieval; not independently reviewed | Daniil Dmitriev | 2025-05-21

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

V3-000018 | R101-matrix-discrepancy-1

Welcome, and Matrix Discrepancy (problems 1-2) | Conjecture 1 (Matrix Spencer)

There exists a positive universal constant C such that, for all positive integers n, and all choices of n self-adjoint $n\times n$ real matrices $A_1,\dots,A_n$ satisfying, for all $i\in[n]$, $\|A_i\|\leq 1$ (where $\|\cdot\|$ denotes the spectral norm) the following holds \begin{equation}\label{eq:matrixspencerbound} \min_{\varepsilon \in \{\pm1\}}^n \left\| \sum_{i=1}^n\varepsilon_iA_i\right\|\leq C\sqrt{n}. \end{equation}

来源明示猜想 | source_label_at_retrieval; not independently reviewed | Afonso S. Bandeira | 2024-10-15

来源明示问题块；所需定义及上下文见原文 | 原始来源 | CC BY 4.0

A02 ConjectureBench 编目 | 41 条

V3-000019 | CB-cb-0221

OEIS A231201 power-of-two prime-sum conjecture

Every integer n>1 can be written as n=x+y with positive integers x,y such that 2^x+y is prime.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000020 | CB-cb-0222

OEIS A232174 two-prime representation conjecture

Every integer n>1 can be written as n=x+y with positive integers x,y such that x+ny and x^2+ny^2 are both prime.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000021 | CB-cb-0225

OEIS A303656 squares-and-powers conjecture

Every integer n>1 can be written as $n=a^2+b^2+3^c+5^d$ for nonnegative integers a,b,c,d.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000022 | CB-cb-0227

OEIS A308734 restricted power-square conjecture

Every integer n>1 can be written as $n=(2^a3^b)^2+(2^c5^d)^2+x^2+y^2$ for nonnegative integers a,b,c,d,x,y.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000023 | CB-cb-0072

Complete EFX for additive goods

Every instance of four agents with nonnegative additive valuations admits a complete EFX allocation: for every pair of agents i,j and every good g in j's bundle with positive value to i, v_i(A_i)≥v_i(A_j\{g}).

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-23

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000024 | CB-cb-0105

Rule 115 even-period formula

For reversible second-order cellular automaton rule 115, the configuration family 1(01)^k has period P(2i)=(2·4^(2i+2)-7·4^(i+1)+5)/3 for every natural number i.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-23

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000025 | CB-cb-0263

Popovych polynomial-congruence conjecture

Let n>1 and let r be a prime not dividing n. If both $(X-1)^n \equiv X^n-1$ and $(X+2)^n \equiv X^n+2$ in $(\mathbb{Z}/n\mathbb{Z})[X]/(X^r-1)$, then n is prime or $n^2 \equiv 1 \pmod r$.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000026 | CB-cb-0275

Büchi five-squares problem

For all integers x and a, if $(x+n)^2+a$ is a perfect square for n=0,1,2,3,4, then a=0.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000027 | CB-cb-0184

Five-dimensional kissing-number conjecture

The kissing number in Euclidean dimension five is 40.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-23

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000028 | CB-cb-0185

Six-dimensional kissing-number conjecture

The kissing number in Euclidean dimension six is 72.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-23

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000029 | CB-cb-0186

D_4 sphere-packing optimality conjecture

The D_4 checkerboard packing is a densest unit-sphere packing in Euclidean dimension four.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-23

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000030 | CB-cb-0188

E_6 sphere-packing optimality conjecture

The E_6 root-lattice density is optimal among unit-sphere packings in Euclidean dimension six.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-23

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000031 | CB-cb-0108

\(\operatorname{ex}(Q_8,C_4)=680\)

The maximum number of edges in a C_4-free subgraph of the eight-dimensional cube Q_8 is 680.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-22

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000032 | CB-cb-0253

Tight union-closed family cardinality conjecture

If every ground-set element of a nontrivial union-closed family occurs in exactly half its members, then the family has 2^k members for some k.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000033 | CB-cb-0302

Polychromatic plane-graph coloring at face size five

Every plane graph whose shortest facial boundary has length five has a not-necessarily-proper vertex 3-coloring in which all three colors occur on the boundary of every face.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000034 | CB-cb-0159

Three-colouring great-circle arrangements

The arrangement graph of any set of great circles with no triple intersection is vertex three-colourable.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-23

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000035 | CB-cb-0109

Integer complexity of \(2^k3^\ell\)

For all nonnegative k and ℓ, the integer complexity of 2^k3^ℓ equals 2k+3ℓ.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-22

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000036 | CB-cb-0218

OEIS A034693 bounded-multiplier conjecture

For every integer n>1 there is an integer k<n such that nk+1 is prime.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000037 | CB-cb-0228

OEIS A034693 three-quarter-power strengthening

For every positive integer n there is a nonnegative integer k<1+n^(3/4) such that nk+1 is prime.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000038 | CB-cb-0231

Sun's square-plus-one primitive-root conjecture

For every prime p, there is an integer k such that $k^2+1<p$ and k^2+1 is a primitive root modulo p.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000039 | CB-cb-0232

A056777 prime-quadruple characterization conjecture

Every composite n satisfying φ(n+12)=φ(n)+12 and σ(n+12)=σ(n)+12 has the form n=p(p+8), where p,p+2,p+6,p+8 are all prime.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000040 | CB-cb-0233

A063880 residue-108 conjecture

Every positive integer n satisfying σ(n)=2·usigma(n), where usigma is the sum of unitary divisors, is congruent to 108 modulo 216.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000041 | CB-cb-0234

A063880 unique primitive-term conjecture

The only n satisfying σ(n)=2·usigma(n) with no proper divisor satisfying the same identity is n=108.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000042 | CB-cb-0235

A067720 unique composite-successor conjecture

If φ(k²+1)=k·φ(k+1) and k≠8, then k+1 is prime.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000043 | CB-cb-0236

Arithmetic sum S positivity conjecture

For every odd prime k, S(k)>0, where S(k)=∑_{h=1}^{k-1}∑_{j=1}^{k-1}(-1)^(j+1+⌊hj/k⌋).

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000044 | CB-cb-0237

Arithmetic sum S greater-than-k conjecture

For every odd prime k>5, S(k)>k, where S(k)=∑_{h=1}^{k-1}∑_{j=1}^{k-1}(-1)^(j+1+⌊hj/k⌋).

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000045 | CB-cb-0238

Arithmetic sum S greater-than-2k conjecture

For every odd prime k>233, S(k)>2k, where S(k)=∑_{h=1}^{k-1}∑_{j=1}^{k-1}(-1)^(j+1+⌊hj/k⌋).

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000046 | CB-cb-0239

Arithmetic sum S greater-than-3k conjecture

For every odd prime k>3119, S(k)>3k, where S(k)=∑_{h=1}^{k-1}∑_{j=1}^{k-1}(-1)^(j+1+⌊hj/k⌋).

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000047 | CB-cb-0243

Surányi factorial equation conjecture

The only integers n,a,b>1 satisfying n!=a!b!, b≤a, and a+1≠n are (n,a,b)=(10,7,6).

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000048 | CB-cb-0246

Second cuboid polynomial irreducibility conjecture

For all distinct positive coprime integers a,b, the associated monic degree-ten second cuboid polynomial Q_(a,b)(t) is irreducible over the integers.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000049 | CB-cb-0247

Third cuboid polynomial irreducibility conjecture

For positive pairwise-distinct integers a,b,c with gcd(a,b,c)=1, bc≠a², and ac≠b², the associated monic degree-twelve third cuboid polynomial P_(a,b,c)(t) is irreducible over the integers.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000050 | CB-cb-0249

No positive (2,5)-perfect number

There is no positive integer n such that σ(σ(n))=5n, where σ is the sum of positive divisors.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000051 | CB-cb-0036

Collatz, finite-cycle route

The Collatz map T(n)=n/2 for even n and T(n)=3n+1 for odd n has no positive-integer cycle other than 1→4→2→1.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-22

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000052 | CB-cb-0044

Giuga

If an integer n≥2 satisfies sum(a^(n-1), a=1,...,n-1)≡-1 (mod n), then n is prime.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-22

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000053 | CB-cb-0049

Goormaghtigh

For integers x>y>1 and m,n>2, the equation (x^m-1)/(x-1)=(y^n-1)/(y-1) has only the solutions (x,y,m,n)=(5,2,3,5) and (90,2,3,13).

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-22

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000054 | CB-cb-0106

Largest fifth pivot under Gaussian elimination with complete pivoting

The largest fifth pivot growth factor under Gaussian elimination with complete pivoting is the unique real root in [4,5] of the explicit degree-61 polynomial P_5 in Equation (2.15) of the source.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-22

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000055 | CB-cb-0111

\(R(5,5)=43\)

The diagonal Ramsey number R(5,5) equals 43.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-22

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000056 | CB-cb-0101

Li-Guo Laplacian degree-sequence inequality

For every graph G other than a cycle C_(4k+1), the sum over Laplacian eigenvalues $\mu_i \geq 2$ of $(\mu_i-2)^2$ is at most $(1-1/d_1)$ times the sum of $d_i(d_i-1)$.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-23

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000057 | CB-cb-0104

Nordhaus-Gaddum product for the \(k\)-th Laplacian eigenvalue

For every n-vertex graph and $k \leq 3n/4$, the k-th Laplacian eigenvalues of G and its complement satisfy $\mu_k(G)\mu_k(\text{complement}(G)) \leq n(n-k)$.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-23

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000058 | CB-cb-0190

Square peg conjecture

Every Jordan curve in the plane contains four points that are the vertices of a square.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-23

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

V3-000059 | CB-cb-0264

Finite magma equation 677→255 implication

Every finite magma satisfying $x = y \diamond (x \diamond ((y \diamond x) \diamond y))$ for all x,y also satisfies $x = ((x \diamond x) \diamond x) \diamond x$ for all x.

目录整理表述 | open | Anirudha Ramesh; Shreyas Pimpalgaonkar; 原数学作者见来源 | 2026-07-24

摘录，定义可能依赖原文 | 原始来源 | CC BY 4.0 for curated text; upstream rights retained

A03 OpenConjecture / arXiv | 267 条

V3-000060 | OC-1087

Computing the Skyscraper Invariant | main.tex:637-639

\label{obs:factor_dimension} Modules coming from the first persistent homology group of a density-scale bifiltration (\autoref{subsec:bifiltr}) of a generic pointset in $\bbR^2$ do not contain indecomposable uniquely generated submodules of dimension larger than 1. In particular, all semistable factors of the Harder-Narasimhan filtration are interval modules.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Fersztand, Marc; Jendrysiak, Jan | 2026-03-24

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by_sa

V3-000061 | OC-1091

Energy-Morawetz estimates for the wave equation in perturbations of Kerr | scalarKerr-arxiv-v2.tex:588-590

The maximal Cauchy development of any initial data set for EVE, that is sufficiently close to a subextremal Kerr initial data in a suitable sense, has a complete future null infinity and a domain of outer communication\footnote{The domain of outer communication is the complement of the black hole region.} which is asymptotic to a nearby member of the subextremal Kerr family.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Ma, Siyuan; Szeftel, Jérémie | 2024-10-03

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000062 | OC-1098

Cambrian lattices are fractionally Calabi-Yau via 2-cluster combinatorics | main.tex:1710-1713

Let $A=kL$ be the incidence algebra of a lattice L. Then A is combinatorially Serre formal if and only if A is periodic Serre formal.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Kleinau, Markus | 2026-03-24

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000063 | OC-1126

Limit Values of Character Sums in Frobenius Formula of Three Permutations | 1_wholepaper8.tex:1418-1420

If the triple of conjugacy classes \(C_1, C_2, C_3\) of the symmetric group ${\frak S}_n$ are all 3-derangements, then the branching data \(C_1,C_2,C_3\) is realizable.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Liang, Dun; Xu, Bin; Yang, Wenyan | 2026-03-24

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by_sa

V3-000064 | OC-1181

Ties in function field prime race | tie_ff.tex:1579-1585

Let $m \in \mathbb{F}_q[T]$ and let $a, b \in (\mathbb{F}_q[T]/m)^\times$ be two distinct congruence classes modulo m. Then there are finitely many N such that \[\sum_{n=1}^{N} \pi(a, m, n) = \sum_{n=1}^{N} \pi(b, m, n). \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Bates, Graeme; Jesubalan, Ryan; Lee, Seewoo; Lu, Jane; Shim, Hyewon | 2026-03-22

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000065 | OC-1188

Composition Theorems for Multiple Differential Privacy Constraints | main.tex:1132-1137

Let $\boldeps = (\eps_i)_{i=1}^n \in (\R^*_+)^n$ and $\bolddelta = (\delta_i)_{i=1}^n \in [0,1]^n$ such that all $(\eps_i,\delta_i)$-DP constraints are active. Then the privacy region of the composition of k mechanisms which are $(\eps_i, \delta_i)$-DP for $i \in \lintv{1,n}$ is\begin{align*} \regcomposedp{\boldeps}{\bolddelta}{k} = \Tilde{\delta} \regdp{0}{1} + (1-\Tilde{\delta}) \sum_{\substack{\boldsymbol{j} \in \lintv{0,k}^n\\ \sum_{i=1}^n j_i = k}} \omega_{\boldsymbol{j}}\regheterdpsingleparam{\boldeps}{\boldsymbol{j}} \end{align*} for $\omega_{\boldsymbol{j}} = \binom{k}{j_1,\ldots,j_n}\prod_{i=1}^n\alpha_i^{j_i}$ for some functions α_i, $\Tilde{\delta}$ of $\boldeps$, $\bolddelta$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Cadir, Cemre; Najib, Salim; Shkel, Yanina Y. | 2026-03-21

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000066 | OC-1721

Restricted Projections to Hyperplanes in $\mathbb{R}^n$ | Projection-to-hyperplane.tex:223-239

\label{conjfo} Let Z be an analytic set in $\R^3$ and $\gz \in C^2([0,1], S^2)$ be a non-degenerate curve, that is \begin{equation}\label{nondeg0} {\rm span}\{ \gamma(\theta), \dot \gamma(\theta), \ddot \gamma(\theta)\}=\mathbb{R}^3 \text{ for all } \theta \in [0,1]. \end{equation} Then \begin{enumerate} \item[(i)] \begin{equation*} \dim [\pi_{{\rm span}\{\gamma(\theta)\}}(Z)] = \min\{\dim Z, 1 \}, \quad \calH^1\mbox{-\rm a.e.} \ \theta\in [0,1]; \end{equation*} \item[(ii)] \begin{equation*} \dim [\pi_{T_{\gamma(\theta)}S^2}(Z)] = \min\{\dim Z, 2 \}, \quad \calH^1\mbox{-\rm a.e.} \ \theta\in [0,1]. \end{equation*} \end{enumerate}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Liu, Jiayin | 2026-04-16

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000067 | OC-1722

Restricted Projections to Hyperplanes in $\mathbb{R}^n$ | Projection-to-hyperplane.tex:253-259

\label{conjI} For $n \ge 4$, let Z be an analytic set in $\R^n$ and $\Sigma \subset S^{n-1}$ be an $(n-2)$-dimensional C^2 manifold with sectional curvature >1. Then \begin{equation*} \dim [\pi_{T_{x}S^{n-1}}(Z)] = \min\{\dim Z, n-1 \}, \quad \calH^{n-2} \ a.e. \ x\in \Sigma. \end{equation*}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Liu, Jiayin | 2026-04-16

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000068 | OC-1724

A cord algebra for tori in three-space | main.tex:1278-1287

\label{conj_diag} Let $\varepsilon\in(0, \varepsilon_{diag})$. Let $\widehat{M}_{K, \varepsilon}$ denotes the restriction of $M_{K, \varepsilon}$ to the set $\lbrace(s_1, \theta_1, s_2, \theta_2)\in(\R/T\mathbb{Z}\times S^1)^2\,|\,\varepsilon\leq\widetilde{d}(s_1, s_2)\leq\delta_K\rbrace$. Then the following holds \begin{itemize} \item[$(i.)$]$\widehat{M}_{K, \varepsilon}$ is a 4-manifold with corners and the projection $\pi_{s_1}:\widehat{M}_{K, \varepsilon}\rightarrow\R/T\mathbb{Z}$ induce a locally trivial fibration. In more detail, $\widehat{M}_{K, \varepsilon}=(F^{[\varepsilon]})^{-1}(Q)$, where $F^{[\varepsilon]}=(F^{[\varepsilon]}_1, F^{[\varepsilon]}_2, F^{dist})$ and $Q=[0, \infty)\times[0, \infty)\times\lbrace[-\delta_K, -\varepsilon]\cup[\varepsilon, \delta_K]\rbrace$. \item[$(ii.)$]Let $\overline{s}_1, \overline{s}_2\in\R/T\mathbb{Z}$ be such that $\overline{s}_1=\overline{s}_2$, i.e. $(\overline{s}_1, \overline{s}_2)$ is a diagonal pair. Then $\pi_{s_1}^{-1}(\overline{s}_1)$ is a 3-manifold with corners and the projection $\pi_{s_2}:\pi_{s_1}^{-1}(\overline{s}_1)\rightarrow (\overline{s}_2-\delta_K, \overline{s}_2-\varepsilon)\cup(\overline{s}_2+\varepsilon, \overline{s}_2+\delta_K)$ induce a broken fibration. In more detail, $\pi_{s_1}^{-1}(\overline{s}_1)=(\widetilde{F}^{[\varepsilon]})^{-1}(\widetilde{Q})$, where $$\widetilde{F}^{[\varepsilon]}=(F^{[\varepsilon]}_1\vert_{\pi_{s_1}^{-1}(\overline{s}_1)}, F^{[\varepsilon]}_2\vert_{\pi_{s_1}^{-1}(\overline{s}_1)}, Id)$$ and $$\widetilde{Q}=[0, \infty)\times[0, \infty)\times\lbrace[\overline{s}_2-\delta_K, \overline{s}_2-\varepsilon]\cup[\overline{s}_2+\varepsilon, \overline{s}_2+\delta_K]\rbrace.$$ And the only critical points of π_{s_2} are two critical points in $(\widetilde{F}^{[\varepsilon]})^{-1}(\widetilde{\mathcal{Q}}_1)$, each of them has Morse index equal to 1. \end{itemize}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Poppr, Marián | 2026-04-15

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000069 | OC-1725

A cord algebra for tori in three-space | main.tex:2152-2154

\label{conj_grad_diag}Let $x_\varepsilon \in M_{K, \varepsilon}\setminus \Delta_\varepsilon$ such that $\pi_{s_1, s_2}(x_\varepsilon)$ is an almost diagonal pair for $\varepsilon>0$ small (recall Definition \ref{lemma_aux_diag} for almost diagonal pairs). If $$F^{[\varepsilon]}_1(x_\varepsilon)=0\vee F^{[\varepsilon]}_1(x_\varepsilon)=0,$$ then $-\nabla E_\varepsilon(x_\varepsilon)$ is strictly outward-pointing from $M_{K, \varepsilon}$. I.e. $-\nabla E_\varepsilon(x_\varepsilon)\neq 0$ and there is $\delta>0$ such that $x_\varepsilon\cdot_{E_\varepsilon}(0, \delta]\cap M_{K, \varepsilon}=\emptyset$ and $x_\varepsilon\cdot_{E_\varepsilon}[-\delta, 0)\subset Int(M_{K, \varepsilon})$. See also Example \ref{exm_diag_grad}.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Poppr, Marián | 2026-04-15

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000070 | OC-1726

A cord algebra for tori in three-space | main.tex:2234-2260

\label{conj_eating_cusp} For every $\delta>0$ small there is a subset $\Delta_{\varepsilon, \delta}\subset\Delta_{\varepsilon}^{cusp}$ which is $O(\delta)$-close and diffeomorphic to $\Delta_{\varepsilon}^{cusp}$ and the following holds. If $p\in\Delta_{\varepsilon, \delta}$, then the set $$ A _p=\left\{ x_\varepsilon\in\nu_\delta(\Delta_{\varepsilon, \delta})\setminus\Delta_{\varepsilon, \delta} \ \middle\vert \begin{array}{l} x_\varepsilon\cdot_{E_\varepsilon}[0, \infty)\subset M_{K, \varepsilon}, \\ p\hbox{ is the omega limit of }x_\varepsilon \end{array}\right\} $$ is flow invariant and diffeomorphic to a disjoint union of two open intervals. See also Figure \ref{figure_eat_trajectory}. \begin{figure}[!htbp] \labellist \pinlabel p at 80 744 \pinlabel $\pi_{E_\varepsilon}^{-1}(p)$ at 30 828 \pinlabel p at 380 744 \pinlabel $\pi_{E_\varepsilon}^{-1}(p)$ at 320 828 \pinlabel $\textcolor{blue}{M_{K, \varepsilon}}$ at 435 774 \pinlabel $\textcolor{red}{A_p}$ at 422 630 \endlabellist \centering \includegraphics[scale=0.75]{eat_trajectory} \vspace{0.3cm} \caption{\textit{On the left:} the fiber $\pi_{E_\varepsilon}^{-1}(p)$ of the fibration $\widehat{W}^s_{E_\varepsilon}(\Delta_{full})\xrightarrow{\pi_{E_\varepsilon}}\Delta_{full}$ inside $\nu_\delta(\Delta_{full})$. The gradient flow trajectories of $-\nabla E_\varepsilon$ are in grey. \textit{On the right:} the restriction of the gradient flow trajectories from $\pi_{E_\varepsilon}^{-1}(p)$ to $\pi_{E_\varepsilon}^{-1}(p)\cap M_{K, \varepsilon}$. The set A_p is depicted in red.} \label{figure_eat_trajectory} \end{figure}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Poppr, Marián | 2026-04-15

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000071 | OC-1727

A cord algebra for tori in three-space | main.tex:5226-5240

\label{cor_moduli_boundary_phenomena}Let $\bm{a}$ be a Reeb chord on $\mathcal{L}^\ast_+ T_K$ and J be generic. Then the moduli space $\mathcal{M}_{T_K}(\bm{a}; \tfrac{1}{2},\dots, \tfrac{1}{2})$ with 2ℓ switches

admits a natural compactification to a compact oriented $(\vert\bm{a}\vert+\ell)$-dimensional manifold $\overline{\mathcal{M}}_{T_K}(\bm{a}, \tfrac{1}{2},\dots, \tfrac{1}{2})$ with corners. The codimension 1 strata consist of: \begin{itemize} \item[$(i.)$] moduli spaces $\mathcal{M}_{T_K}(\bm{a}, \tfrac{1}{2},\dots, \tfrac{1}{2}, \tfrac{3}{2}, \tfrac{1}{2},\dots \tfrac{1}{2})$ with 2ℓ switches, where one $\tfrac{1}{2}$ switch was replaced by a $\tfrac{3}{2}$ switch. \item[$(ii.)$] moduli spaces $\mathcal{M}_{T_K}(\bm{a}, \tfrac{1}{2},\dots, \tfrac{1}{2}, 2, \tfrac{1}{2},\dots \tfrac{1}{2})$ with $2\ell+1$ switches, where we added one 2 switch from $L^\ast_+ T_K$ back to $L^\ast_+ T_K$ (i.e. not from $\R^3$ to $\R^3$). \item[$(iii.)$] products of moduli spaces $$\mathcal{M}^{sy}_{T_K}(\bm{a}, \bm{b}_1,\dots, \bm{b}_m)/\R\times\prod_{i=1}^m\mathcal{M}_{T_K}(\bm{b}_i; \tfrac{1}{2},\dots, \tfrac{1}{2}),$$ where the Reeb chords $\bm{b}_i$ have total degree $\vert \bm{a}\vert-1$ and the moduli spaces $\mathcal{M}_{T_K}(\bm{b}_i; \tfrac{1}{2},\dots, \tfrac{1}{2})$ have together 2ℓ switches. \item[$(iv.)$] moduli spaces $\mathcal{M}_{T_K}(\bm{a}, \tfrac{1}{2},\dots, \tfrac{1}{2}, 1, \tfrac{1}{2},\dots \tfrac{1}{2})$ with $2\ell-1$ switches, where two consecutive switches $\tfrac{1}{2}$ were replaced by a single 1 switch, provided that the first replaced $\tfrac{1}{2}$ switch was switching from $\R^3$ to $L^\ast_+ T_K$. See also Figure \ref{figure_string_degeneration}. \end{itemize} The codimension 2 strata consist of corners given by the combinations of phenomena $(i.)-(iv.)$ and \begin{itemize} \item[$(v.)$] moduli spaces $\mathcal{M}_{T_K}(\bm{a}, \tfrac{1}{2},\dots, \tfrac{1}{2}, \tfrac{5}{2}, \tfrac{1}{2},\dots \tfrac{1}{2})$ with 2ℓ switches, where one $\tfrac{1}{2}$ switch was replaced by a $\tfrac{5}{2}$ switch. \item[$(vi.)$] moduli spaces $\mathcal{M}_{T_K}(\bm{a}, \tfrac{1}{2},\dots, \tfrac{1}{2}, 2, \tfrac{1}{2},\dots \tfrac{1}{2})$ with $2\ell-1$ switches, where two consecutive switches $\tfrac{1}{2}$ were replaced by a single 2 switch, provided that the first replaced $\tfrac{1}{2}$ switch was switching from $L^\ast_+ T_K$ to $\R^3$. See also Figure \ref{figure_string_degeneration}. \end{itemize}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Poppr, Marián | 2026-04-15

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000072 | OC-1728

A cord algebra for tori in three-space | main.tex:5500-5508

\label{conj_gradient_string}Let K be generic and $\varepsilon>0$ small. Let c be a path from $\partial\mathcal{P}_{K, \varepsilon}$. Then $c\in\partial^-\mathcal{P}_{K, \varepsilon}$ iff c can be reparametrized such that one of the following conditions on normal jets σ of c holds: \begin{itemize} \item[(S)] At the starting point we have $\big(\sigma^{(1)}(0), \sigma^{(2)}(0), \sigma^{(3)}(0)\big)\in0\times0\times(-\infty, 0)$. \item[$(E)$] At the endpoint we have $\big(\sigma^{(1)}(T_0), \sigma^{(2)}(T_0), \sigma^{(3)}(T_0)\big)\in0\times0\times(0, \infty)$. \end{itemize} Note that the curve c satisfying (S) or (E) can be viewed as a Q-string in $\partial^Q\Sigma_1$. Moreover, the analogous holds for paths in $\mathcal{P}_{K, \varepsilon}^{chord}$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Poppr, Marián | 2026-04-15

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000073 | OC-1787

Cohomology of the pure symmetric automorphisms of right-angled Artin groups | main.tex:184-193

\label{ConjA} Let γ_A^i be a one-dimensional class in $H^*(\PAut(A_\G))$, where A is a connected component in $\G-\st(v_i)$. For each pair v_i,v_j of two non-adjacent vertices in $\G$, let D^j, D^i be the respective dominant components and $\mathcal{C}$ a shared component. Then $H^*(\PAut(A_\G))$ is generated by the one-dimensional classes γ_A^i, modulo the following relations for each shared component $\C$ of v_i,v_j. \begin{itemize} \item[(1)] $\gamma_A^i\wedge\gamma_A^i=0$ \item[(2)] $\gamma_A^i\wedge\gamma_B^j=-\gamma_B^j\wedge\gamma_A^i$ \item[(3)] $\gamma_{D^j}^i\wedge \gamma_{D^i}^j=0$ \item[(4)] $\gamma_{\mathcal{C}}^i\wedge \gamma_{\mathcal{C}}^j=\gamma_{\mathcal{C}}^i\wedge \gamma_{D^i}^j+ \gamma_{D^j}^i\wedge\gamma_{\mathcal{C}}^j$ \end{itemize}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Ardaiz-Gale, Peio | 2026-04-15

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000074 | OC-1788

Cohomology of the pure symmetric automorphisms of right-angled Artin groups | main.tex:1373-1375

The map ϕ of Corollary \ref{cor:H2_1} is an isomorphism.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Ardaiz-Gale, Peio | 2026-04-15

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000075 | OC-1813

On a conjecture concerning the property of chromatic polynomials with negative variables | conj3.tex:107-112

\label{con} Let \(G \) be a graph of order \(n \). Then \[\frac{d^k}{dx^k} (\ln[(-1)^n P(G, x)]) < 0 \] holds for all \(k \geq 2 \) and \(x \in (-\infty, 0) \).

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Yang, Yan | 2026-03-08

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000076 | OC-1816

Reconstructing a graph from its Bell colouring graph | main.tex:354-357

\label{conj:mainlower} Let~G_1 and~G_2 be graphs and let~$k_i>\chi(G_i)$ for~$i\in \{1,2\}$. Then $\mathcal{B}_{k_1}(G_1)$ and~$\mathcal{B}_{k_2}(G_2)$ are isomorphic if and only if the graphs obtained from~G_1 and~G_2 by removing all universal vertices are isomorphic, and either~$k_i\geq \ord{V(G_i)}$ for~$i\in \{1,2\}$, or~$\ord{V(G_1)}-k_1=\ord{V(G_2)}-k_2$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Hearn, Brian | 2026-04-14

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000077 | OC-1856

Wheel Classes in Kontsevich Graph Complex and Merkulov's Low-Valence Conjecture | low_valence_wheels_submission.tex:64-71

\label{conj:merkulov} Let $\GC_d^{\mathrm{Me}}$ be the maximal subcomplex of $\GC_d$ spanned by linear combinations of graphs all of whose vertices have valence 3 or 4. Then the inclusion \[i \colon \GC_d^{\mathrm{Me}}\hookrightarrow \GC_d \] is a quasi-isomorphism.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Andersson, Assar | 2026-04-13

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000078 | OC-1857

Coarse Balanced Separators in Fat-Minor-Free Graphs | main.tex:138-140

\label{conj:fatminor-sep} For every graph H and $d\in \N$, there exist constants $c,r\in \N$ such that every n-vertex weighted graph G that excludes H as a~d-fat minor has a~$(c\sqrt{n},r)$-coverable balanced separator.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Bonnet, Édouard; Le, Hung; Pilipczuk, Marcin; Pilipczuk, Michał | 2026-04-13

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000079 | OC-1858

Coarse Balanced Separators in Fat-Minor-Free Graphs | main.tex:661-663

\label{conj:better-clustering} For every graph H there are constants $p,d\in \N$ such that every graph G that excludes H as a 3-fat minor admits a connected partition $\Xx$ of weak diameter at most d such that for every $u\in V(G)$, $B_G(u,2)$ intersects at most p clusters of $\Xx$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Bonnet, Édouard; Le, Hung; Pilipczuk, Marcin; Pilipczuk, Michał | 2026-04-13

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000080 | OC-1859

Coarse Balanced Separators in Fat-Minor-Free Graphs | main.tex:669-671

\label{conj:indminor-sep} For every graph H there exist constants $c,r\in \N$ such that every n-vertex weighted graph~G that excludes H as an induced minor has a balanced separator that is $(c\sqrt{n},r)$-coverable.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Bonnet, Édouard; Le, Hung; Pilipczuk, Marcin; Pilipczuk, Michał | 2026-04-13

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000081 | OC-1860

Coarse Balanced Separators in Fat-Minor-Free Graphs | main.tex:675-677

For every graph H there exist constants $c,q\in \N$ such that every n-vertex graph G that excludes H as an induced minor admits a connected partition $\Xx$ such that every cluster of $\Xx$ is $(q,1)$-coverable and the quotient graph $G/\Xx$ has at most $c\cdot n$ edges.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Bonnet, Édouard; Le, Hung; Pilipczuk, Marcin; Pilipczuk, Michał | 2026-04-13

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000082 | OC-1869

On two conjectures for generalized off-diagonal Schur numbers | main.tex:253-258

{\upshape \cite{AhSc16}}\label{conjecture-2} For $4 \leq s \leq t \leq u$, we have $$ S(3 ; s, t, u)=s t u-t u-u-1. $$

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Song, Yanyan; Mao, Yaping | 2026-04-13

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000083 | OC-1877

Beatty solutions of almost Golomb equations | paper54.tex:1347-1352

\label{conj:defect} The gap sequence of~$\mathcal{D}$ on the alphabet $\{3,4,7\}$ is a coding of the fixed point of a primitive substitution; equivalently,~$\mathcal{D}$ is a morphic set.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Cloitre, Benoit | 2026-04-12

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000084 | OC-1878

Polynomial Time Enumeration of t-Stack-Sortable Permutations Ending in Their Least Entry | stacksortingdiagrams.tex:871-873

\label{conj} For all $n\in\mathbb{N}$, the value $|\mathcal{W}_2'(n)|$ is the nth Motzkin number.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Zhang, Jerry | 2026-04-12

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000085 | OC-1889

An Inductive Strategy Towards a Solution to the Generalized Homotopy Hypothesis | An_Inductive_Guide_to_Homotopy_Hypothesis.tex:2621-2640

\label{the_key_conjecture} Let $n\geq 1$ and assume the canonical model structure exists on $\Mod(IC_{n+1})$. Let $0\leq k<n$, $X:IC_{n+2}\to\Set$ be a model, $f:D^k\to UX$ be a map, and \[\begin{tikzpicture}[node distance=2.5cm] \node(A) {D^k}; \node (B) [right of=A]{UX}; \node(C)[below of=A]{D^{k+1}}; \node (D)[right of=C]{$(UX)_+$}; \draw[->](B) to node[black,right=3]{p}(D); \draw[->](A) to node {}(B); \draw[->] (A) to

node {}(C); \draw[->](C) to node {}(D); \end{tikzpicture} \] be a pushout in $\Mod(IC_{n+1})$. Then \[\pi_{n+1}(\free(p)):\pi_{n+1}(\free UX,x)\to \pi_{n+1}(\free((UX)_+),\free(p)(x)) \] is an isomorphism of groups for all $x\in \free UX_0$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Taylor, Johnathon | 2026-04-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000086 | OC-1891

Gertsch quotient living in the "poor man's adele ring" $\mathcal{A}$: Kurepa-Bell-Wilson congruence | sn-article.tex:333-339

\cite{gertsch1999congruences, kurepa1971left} Let p be an odd prime, then \begin{enumerate} \item $K_p=\sum_{0\leq n\leq (p-1)} n!= 0! + 1! + 2! + \cdots + (p-2)! + (p-1)!$ is not congruent to zero modulo $p.$ \item the $(p-1)^{th}$ Bell number $\mathbf{Bell}_{p-1}$ is not congruent to 1 modulo $p.$ \end{enumerate}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Howard, Francis Atta | 2026-04-08

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000087 | OC-1892

Gertsch quotient living in the "poor man's adele ring" $\mathcal{A}$: Kurepa-Bell-Wilson congruence | sn-article.tex:720-724

\label{conj9}[Agoh \cite{agoh1995giuga}] An integer $p>1$ is an odd prime number if and only if $$p\mathbf{B}_{p-1}\equiv -1 \pmod{p}$$ where $\mathbf{B}_{p-1}$ is the even Bernoulli number.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Howard, Francis Atta | 2026-04-08

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000088 | OC-1894

Graph Energy Maximisation for Integral Circulant Graphs of Order $n = p^2q^3$ | icg_p2q3_arxiv__1_.tex:224-229

\label{conj:universal} For every pair of distinct odd primes (p,q), the set $\Dstar=\{1,p^2,pq,q^2,p^2q^2,pq^3\}$ is the unique maximiser of $E(p^2q^3,\mathcal{D})$ over all nonempty $\mathcal{D}\subseteq \mathrm{div}(p^2q^3)\setminus\{p^2q^3\}$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Roldan, Diego | 2026-04-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000089 | OC-1900

Free Field Realizations of Superelliptic Affine Lie Algebras | SuperellipticFFA_main.tex:1030-1039

\label{conj:critical} For each $1\leq l\leq m-1$, the σ^l-twisted intertwining operator construction realizes the Type~II bracket of $\gtil$ exactly at the critical level $k=k_{\mathrm{crit}}(l,m)=m/(l(m-l)+m)$. In particular, all sectors of a fixed m are realized simultaneously only if $k_{\mathrm{crit}}(l,m)$ is independent of l, which occurs if and only if $l(m-l)$ is constant --- i.e.\ m is arbitrary and $l=1$ (or $l=m-1$), giving $k_{\mathrm{crit}}=m/(m-1+m)=m/(2m-1)$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Santos, Felipe Albino dos | 2026-04-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000090 | OC-2062

Separating Matchings in Cubic Graphs | main.tex:689-692

\label{conj:Funk} Let G be a 2-factor Hamiltonian k-regular bipartite graph. Then, either $k = 2$ and G is a circuit or $k = 3$ and $G \in \mathcal{F}$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Gutiérrez, Juan; Gómez, Renzo | 2026-04-19

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000091 | OC-2085

Moment angle complexes and duality for tight manifolds | ms.tex:305-307

The triangulation of an n-manifold is tight if and only if it's strongly minimal.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Kishimoto, Daisuke; Stanley, Donald; Ruiz, Carlos Gabriel Valenzuela | 2026-04-17

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000092 | OC-2105

Rainbow Separating Path Systems | main.tex:1124-1126

Any class of graphs of bounded treewidth is rainbow separable.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Clifton, Alexander; Kontogeorgiou, George; Taruni, S; Trujillo-Negrete, Ana | 2026-04-17

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000093 | OC-2126

Sequential KV Cache Compression via Probabilistic Language Tries: Beyond the Per-Vector Shannon Limit | KV.tex:1359-1383

\label{conj:cross_session} For sessions s, s' drawn i.i.d.\ from $P_{\mathcal{M}}$, the conditional entropy of session s's KV cache given the cluster centroid s^*'s full cache satisfies: \[H\bigl(\mathrm{KV}_{\leq n}(s) \mid \mathrm{KV}_{\leq n}(s^*)\bigr) \;\leq\; \sum_{i=\bar{d}+1}^{n} H\bigl(t_i^{(s)} \mid t_{<i}^{(s)}\bigr) \;+\; H\bigl(t_{\bar{d}+1}^{(s)} \mid t_{\bar{d}+1}^{(s^*)}, t_{<\bar{d}+1}^{(s)}\bigr), \] where $\bar{d}$ is the divergence position of s and s^*. \emph{Partial proof sketch.} The first term bounds the ``tail'' entropy of session $s$ after divergence: positions $i > \bar{d}$ are bounded by the sequential entropy bound exactly as in Theorem~\ref{thm:seq_bound}. The second term bounds the ``divergence cost'': knowing $t_{\bar{d}+1}^{(s^*)}$ provides partial information about which subtree s is in (via the cluster structure), reducing the entropy of $t_{\bar{d}+1}^{(s)}$. The bound formalizes the

Layer~1 gain: prefix deduplication saves bits on the tail (first term, bounded by per-token surprisal) and reduces divergence cost (second term, bounded by the trie metric $d_{\mathcal{T}}(s, s^*)$). A formal proof requires bounding the conditional mutual information between the centroid's divergence token and the session's divergence token, which depends on the cluster radius δ.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Magarshak, Gregory | 2026-04-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000094 | OC-2127

Sequential KV Cache Compression via Probabilistic Language Tries: Beyond the Per-Vector Shannon Limit | KV.tex:1385-1414

\label{conj:strict_dominance} For any per-vector compression scheme $\mathcal{C}$ (including TurboQuant, KIVI, KVQuant, and all future schemes bounded by the per-vector Shannon limit), there exists a context length $n^*(\mathcal{C})$ such that for all $n > n^*(\mathcal{C})$: \[R_{\mathrm{seq}}(n) \;<\; R_{\mathcal{C}}(n), \] where $R_{\mathrm{seq}}(n)$ is the total bits used by sequential compression and $R_{\mathcal{C}}(n)$ is the total bits used by scheme $\mathcal{C}$, both for a context of length n. \emph{Partial proof.} $R_{\mathcal{C}}(n) \geq B_{\rm pv} \cdot n$ for some $B_{\rm pv} > 0$ (the per-vector Shannon limit is a hard floor for any per-vector scheme). $R_{\mathrm{seq}}(n) = \sum_{i=1}^n H(\mathrm{KV}_i \mid \mathrm{KV}_{<i}) \leq \sum_{i=1}^n H_i$. Under the monotone entropy condition (Corollary~\ref{cor:asymptotic}), $\frac{1}{n}\sum_{i=1}^n H_i = \bar{H}_n \to H_\infty < B_{\rm pv}$ as $n \to \infty$ (since H_∞ is the model's entropy rate, which is strictly less than any fixed per-vector bit budget for a model with perplexity $< 2^{B_{\rm pv}/d_{\rm head}}$). Thus $R_{\mathrm{seq}}(n)/n \to H_\infty < B_{\rm pv} \leq R_{\mathcal{C}}(n)/n$, giving the stated strict dominance for large n. The threshold $n^*(\mathcal{C}) = O(H_\infty / (B_{\rm pv} - H_\infty))$. The main gap in this argument: the monotone entropy condition holds only for coherent text; for mixed-topic sessions with high surprisal variance, $\bar{H}_n$ may not converge below $B_{\rm pv}$. Bounding n^* for realistic session distributions is the key open problem.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Magarshak, Gregory | 2026-04-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000095 | OC-2128

A torsion-intersection proof of perfect-cuboid nonexistence on 1,072 explicit master-tuple fibers | paper3.tex:145-148

\label{conj:B} No perfect cuboid exists in $\N^3$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Peschmann, René | 2026-04-30

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000096 | OC-2152

Hirzebruch χ_{y}-genus of compact almost K\"{a}hler manifold with negative sectional curvature | Hirzebruch_genus_of_AKM-HZ.tex:163-166

Let X be a closed $2n$-dimensional Riemannian manifold with negative sectional curvature $\mathrm{sec}<0$. Then the Euler number of X satisfies \[(-1)^{n}\chi(X)>0. \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Huang, Teng; Zhang, Pan | 2026-04-30

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000097 | OC-2166

Convex Hull Volumes in Hyperbolic 3-Space | main.tex:101-103

Let G be a co-compact Kleinian group. Suppose $K \subset \hat{\comp}$ is a continuum which is disjoint from all of its G-translates. Then K consists of a single point.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | MacMahon, Cameron | 2026-04-30

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000098 | OC-2230

B'-orbits on flag varieties and symmetry breaking | main_SIGMA.tex:989-991

Whenever Ψ is injective, it induces a poset isomorphism onto its image.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Massicot, Valentin | 2026-04-28

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000099 | OC-2231

Balanced and pluriclosed metrics on real semisimple Lie groups | main.tex:171-173

A compact complex manifold admitting both a balanced metric and a (possibly distinct) pluriclosed metric also admits a K\"ahler metric.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Kwong, Joseph | 2026-04-28

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000100 | OC-2237

Families of Eliahou semigroups linked to Farey intervals | paper.tex:1155-1157

\label{conj:gamma2} If~$\gamma_{\ell+1}\ge c_{\max} + m - 2\gamma_2$ and~$E\ge0$, then~$E'\ge0$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Bacher, Axel | 2026-04-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000101 | OC-2292

Pair-Trace Absorption Certificates for Regular Induced Subgraphs | paper.tex:306-312

\label{conj:polynomial-lifting} There are constants \(C,a,C_0>0\) such that: \begin{enumerate}[label=\textup{(\arabic*)}, leftmargin=*] \item every \(2\)-modular witness of size at least \(C_0m\) contains a \(4\)-modular witness of exactly \(m\) vertices;

\item for every power of two \(q\ge 4\), every \(q\)-modular witness of size at least \(Cq^a m\) contains a \(2q\)-modular witness of exactly \(m\) vertices. \end{enumerate}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Ramos, Arthur F.; Hulak, David Barros; de Queiroz, Ruy J. G. B. | 2026-04-26

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000102 | OC-2293

Pair-Trace Absorption Certificates for Regular Induced Subgraphs | paper.tex:318-320

\label{conj:first-bit} There exists a constant \(C_0>0\) such that every \(2\)-modular witness of size at least \(C_0m\) contains a \(4\)-modular witness of exactly \(m\) vertices.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Ramos, Arthur F.; Hulak, David Barros; de Queiroz, Ruy J. G. B. | 2026-04-26

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000103 | OC-2294

Pair-Trace Absorption Certificates for Regular Induced Subgraphs | paper.tex:322-324

\label{conj:higher-bit} There are constants \(C,a>0\) such that for every \(q=2^j\ge 4\), if \(A\) is a \(q\)-modular witness of size at least \(Cq^a m\), then some subset \(W\subseteq A\) with exactly \(m\) vertices satisfies the affine lift criterion of Lemma~\ref{lem:affine-lift}.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Ramos, Arthur F.; Hulak, David Barros; de Queiroz, Ruy J. G. B. | 2026-04-26

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000104 | OC-2312

Asymptotic behaviour of analytic torsion and cohomological torsion for $\mathbb{Q}$-rank 1 arithmetic groups | main.tex:126-132

Let $\Gamma\eqqcolon \Gamma_0 \supset \Gamma_1\supset\dots\supset \Gamma_N\supset\dots$ be a sequence of congruence subgroups satisfying $\bigcap_N \Gamma_N = \lbrace 0\rbrace$. Then \begin{align*} \lim_{N\to\infty}\frac{\log|H^i(\Gamma_N,L)_{\text{tors}}|}{[\Gamma:\Gamma_N]} \end{align*} exists for any i, and is equal to a computable constant $c_{\Gamma,L}$. Furthermore, $c_{\Gamma,L}$ is non-zero exactly if $\delta(G)=1$ and i is the top of the cuspidal range.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Berland, Tim | 2026-04-24

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000105 | OC-2313

Asymptotic behaviour of analytic torsion and cohomological torsion for $\mathbb{Q}$-rank 1 arithmetic groups | main.tex:728-733

\label{globalcoeffconj} There exists constants $b,c>0$ such that for all N, M and $\mathcal{O}_\gamma\in\mathcal{U}_M(\Q)$ we have \begin{align*} \left\vert a^M(S_N,\mathcal{O}_\gamma)\right\vert \leq c(1+\log N)^b. \end{align*}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Berland, Tim | 2026-04-24

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000106 | OC-711

Vigemers: on the number of k-mers sharing the same XOR-based minimizer | main.tex:414-418

Let $g : \Sigma^m\to\Sigma^m$ be such that $g(a_1\cdots a_m) = g_1(a_1)\cdots g_m(a_m)$ where $g_i:\Sigma\to\Sigma$. Then, by defining the minimizer $\min_g(x)$ of a \kmer x as its leftmost \mmer w that minimizes $\texttt{rank}(g(w))$ (for the lexicographical order), we can compute $\pi_k^g(w) = \left\lbrace x\in\Sigma^k : \min_g(x) = w\right\rbrace$ in $O(|\Sigma|\cdot km^2)$ time and $O(km)$ space by adapting the equations presented in this article.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Florian Ingels; Antoine Limasset; Camille Marchet; Mikaël Salson | 2026-02-03

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000107 | OC-2415

Nearly-Tight Bounds for Zonotope Containment and Beyond | submission.tex:443-449

\label{conj:zonoid-sparsification} Given $\varepsilon >0$ and $V ∈ ℝ^{ d × k}$, there exists a matrix $ W \in \R^{d \times n}$ such that the following holds for the zonotopes $Z =\{ Vx : x ∈ [-1,1]^k\}\subseteq \R^d$ and $Z^\prime = \{ Wx : x ∈ [-1,1]^n\}$. \begin{enumerate} \item The number n of generators of Z' is bounded by $n \in O(\nicefrac{d}{\varepsilon^2})$, and \label{item:1-conj} \item $(1-\varepsilon)Z \subseteq Z^\prime \subseteq (1+\varepsilon)Z$. \label{item:2-conj} \end{enumerate}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Eisenbrand, Friedrich; Rothvoss, Thomas; Russo, Matteo; Skorupinski, Ruben | 2026-05-05

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000108 | OC-3884

An Information-Theoretic Analysis of Threshold Group Testing | Paper.tex:794-809

\label{Conj} For any integer $t \geq 3$ and $\alpha \in [0,1]$, let the random variables $\vE^*$, $\vB^*$, $\vX_1$, and $\vX_2$ be independent with distributions \begin{equation*} \vE^* \sim \Po(r^*), \qquad \vB^* \sim \Po(\alpha r^*), \qquad \vX_1, \vX_2 \sim \Po((1-\alpha)r^*). \end{equation*} Define the maximum of $\vX_1$ and $\vX_2$ to be $\vM^* = \max\{\vX_1, \vX_2\}$. Let the probabilities $q(r^*)$ and $p(r^*,\alpha)$ be given by \begin{equation*} q(r^*) = \Pr(\vE^* \leq t-1) \quad \text{and} \quad p(r^*,\alpha) = \Pr(\vB^* + \vM^* \leq t-1). \end{equation*} Then, for all $c>c_{\mathrm{inf}}^{\mathrm{TGT}}$, the function \begin{align} \alpha \mapsto 1 - \alpha + c \bc{q(r^*) \log\bcfr{p(\alpha)}{q(r^*)} + (1-q(r^*)) \log\bcfr{1-2q(^*)+p(r^*,\alpha)}{1-q(r^*)}} \end{align} is maximised in $\alpha=0$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Remco van der Hofstad; Noela Müller; Connor Riddlesden | 2026-06-09

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000109 | OC-800

Functional Properties of the Focal-Entropy | AISTATS_arXiv_v2.tex:2755-2759

\label{conj:S3} Let $|\cS|=3$. Then, for all $\gamma>0$, it holds that $p_{\min} > p_{\gamma,a}$, i.e., $d_3 < 0$ (i.e., the over-suppression regime is absent for $|\cS|=3$).

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Jaimin Shah; Martina Cardone; Alex Dytso | 2026-03-03

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000110 | OC-972

The DNA Coverage Depth Problem: Duality, Weight Distributions, and Applications | journal_paper_arxiv.tex:236-238

Let $\mC \subseteq \mathbb{F}_q^n$ be the q-ary simplex code of dimension k, where $n = (q^k-1)/(q-1)$. Then $\mC$ solves Problem~\ref{second_problem}.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Matteo Bertuzzo; Alberto Ravagnani; Eitan Yaakobi | 2026-03-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000111 | OC-1373

The Geometry of Polynomial Group Convolutional Neural Networks | main.tex:894-899

\label{conj: intersection_of_general_secant_is_trivial} Let $G=\{g_1, \dots, g_n\}$ as a set, and let the activation degree $r\geq 2$. For general filters $\theta=(\theta_1, \dots, \theta_L)$, where $L\geq 1$ we have \[<\sigma_r(\Phi_\theta)(g_1), \dots, \sigma_r(\Phi_\theta)(g_n)> \cap\ VP_{n, r, r^{L-1}} = \{\sigma_r(\Phi_\theta)(g_1), \dots, \sigma_r(\Phi_\theta)(g_n)\}. \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Yacoub Hendi; Daniel Persson; Magdalena Larfors | 2026-03-31

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000112 | OC-3721

Can Neural Networks Achieve Optimal Computational-statistical Tradeoff? An Analysis on Single-Index Model | main.tex:627-630

\label{conj:s^star = 1} For $d^\iota < k < o(d)$ with $s^\star=1$, \Cref{alg:sparse-warm-up} succeeds with sample complexity $n=\tilde O(k)$. Furthermore, the same guarantee applies even without perturbing the weights.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Siyu Chen; Beining Wu; Miao Lu; Zhuoran Yang; Tianhao Wang | 2026-06-13

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000113 | OC-4431

Dynamics of Gradient Descent with Large Step Size Near a Manifold of Flat Minima | neurips_2026.tex:1947-1949

\label{conj:parabolicfoliation} For ρ sufficiently small, the invariant set $V_{\rho}\cap\{y_{2:q}=0\}$ is invariantly Lipschitz-foliated by leaves parametrised by the points $(z,0,u,y_1,0)\in W_{\rho}$, and tangent at any such point to the v-axes.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Lachlan Ewen MacDonald; René Vidal | 2026-07-09

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000114 | OC-4114

Structural properties of the implicit function defined by an integral self-consistency equation | main.tex:623-634

\label{conj:uniqueness} Assume \textup{(D0)--(D3)}, hypotheses \textup{(H1), (H2)} of Theorem~\ref{thm:structure}(e), and that $h(\eta):=\eta\rho(\eta)$ is positive on $(0,M)$ and strictly log-concave: \begin{equation}\label{eq:logconcave} \log h\bigl(\lambda x + (1-\lambda) y\bigr) \;>\; \lambda \log h(x) + (1-\lambda)\log h(y) \qquad (x\ne y\text{ in }(0,M),\ \lambda\in(0,1)). \end{equation} Then the critical point $m^\star$ produced by Theorem~\ref{thm:structure}(e) is the unique critical point of β on Ω.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Ivan Viakhirev | 2026-06-02

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by_sa

V3-000115 | OC-1644

Poisson Vertex Algebra of Seiberg-Witten Theory | pvaofsw-final.tex:679-681

The map $ \varphi: A \to \text{Obs}_{\text{HT}}(\mathfrak{sl}_2)$ is an isomorphism of Poisson vertex algebras.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Ahsan Z. Khan | 2026-04-03

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000116 | OC-2825

A general proof of integer Rényi QNEC | main.tex:1010-1016

\label{conj:RQNEC} Under the hypothesis of Theorem~\ref{thm:modulartranslations}, for $\alpha\in (1,\I)$, with $\omega\in(M_A)_*^+$ the faithful normal vacuum state, and any non-zero $\psi\in (M_A)_*^+$ such that $D_\a(\psi\|\omega;M_A)<\I$, the function \[c\mapsto S_\a(c)= D_\a(\psi_c\|\omega;M_A) \] is convex.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Tanay Kibe; Pratik Roy | 2026-05-14

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000117 | OC-1599

From BV-BFV Quantization to Reshetikhin-Turaev Invariants | RT_paper.tex:1616-1621

\label{conj:E2-equiv} The $\EE_2$-monoidal category $\mathcal{B}_{\CS}^{(k)}$ obtained from BV-BFV quantization of Chern--Simons theory at level k with gauge group G is equivalent, as an $\EE_2$-category, to the ribbon category $\overline{\Rep}_q(G)$ at $q = e^{i\pi/(k+h^\vee)}$: \begin{equation} \mathcal{B}_{\CS}^{(k)} \;\simeq_{\EE_2}\; \overline{\Rep}_q(G). \end{equation}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Nima Moshayedi | 2026-04-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000118 | OC-1600

From BV-BFV Quantization to Reshetikhin-Turaev Invariants | RT_paper.tex:1652-1677

\label{conj:main} There exists a natural equivalence of $(3\text{-}2\text{-}1)$-extended topological quantum field theories \begin{equation} Z_{\mathrm{BV,np}}^{(k)} \;\simeq\; Z_{\mathrm{RT}}^{(k)} \;:\; \Bord_3^{\mathrm{ext}} \longrightarrow 2\text{-}\Vect, \end{equation} where $Z_{\mathrm{BV,np}}^{(k)}$ denotes the non-perturbative completion of the BV-BFV partition function at level k and $2\text{-}\Vect$ is the 2-category of $\CC$-linear categories, functors, and natural transformations. At each categorical level, the equivalence identifies: \begin{enumerate}[label=(\roman*)] \item \emph{Circles:} the $\EE_2$-category $\mathcal{B}_{\CS}^{(k)} \simeq \overline{\Rep}_q(G)$ (Conjecture \ref{conj:E2-equiv}); \item \emph{Closed surfaces:} the factorization homology categories $\iint_{\Sigma_g} \overline{\Rep}_q(G) \simeq (\mathbf{H},\, u_{\Sigma_g})$, with $u_{\Sigma_g} = V_{\mathrm{RT}}(\Sigma_g)$ (Theorem \ref{thm:AKZ}); \item \emph{3-cobordisms:} the linear maps between RT state spaces $\Zrt(M): V_{\mathrm{RT}}(\Sigma_1) \to V_{\mathrm{RT}}(\Sigma_2)$. \end{enumerate} More precisely, for every closed oriented surface Σ_g, the BV-BFV state space is identified with the RT state space via the chain \begin{equation*} \begin{tikzcd}[column sep=2.5em, row sep=2.5em] \mathcal{H}_{\Sigma_g}^{\BV} \arrow[d, dashed, leftrightarrow, "\sim"'] & \Quant\big(\Loc_G(\Sigma_g)\big) \arrow[l, "\textnormal{Step 2}"'] \arrow[r, "\sim", "\textnormal{BBJ}"'] & \displaystyle\iint_{\Sigma_g} \Rep_q(G) \arrow[d, "\textnormal{semisimplify}"] \\ V_{\mathrm{RT}}(\Sigma_g) & (\Vect,\, u_{\Sigma_g}) \arrow[l, "u_{\Sigma_g} \,\mapsto"'] \arrow[r, leftarrow, "\sim"', "\textnormal{AKZ}"] & \displaystyle\iint_{\Sigma_g} \overline{\Rep}_q(G) \end{tikzcd} \end{equation*} where the distinguished object u_{Σ_g} is the RT state space $V_{\mathrm{RT}}(\Sigma_g)$. For every oriented 3-dimensional cobordism $M: \Sigma_1 \to \Sigma_2$, the cobordism induces a functor $\iiint_M$ between the factorization homology categories $\iint_{\Sigma_i} \mathcal{B}_{\CS}^{(k)}$. After the identification with $\overline{\Rep}_q(G)$ via Conjecture \ref{conj:E2-equiv}, and the collapse $\iint_{\Sigma_i} \overline{\Rep}_q(G) \simeq (\Vect,\, u_{\Sigma_i})$ via AKZ, this functor induces a linear map between the distinguished objects. The conjecture asserts that this linear map agrees with the RT partition function: \begin{equation}\label{eq:main-diagram} \begin{tikzcd}[column sep=4em] \iint_{\Sigma_1} \overline{\Rep}_q(G) \arrow[r, "\iiint_M"] \arrow[d, "\textnormal{AKZ}"', "\sim"] & \iint_{\Sigma_2} \overline{\Rep}_q(G) \arrow[d, "\sim"', "\textnormal{AKZ}"] \\ (\Vect,\, u_{\Sigma_1} = V_{\mathrm{RT}}(\Sigma_1)) \arrow[r, "\Zrt(M)"'] & (\Vect,\, u_{\Sigma_2} = V_{\mathrm{RT}}(\Sigma_2)) \end{tikzcd} \end{equation} Here the vertical equivalences are given by the Ai--Kong--Zheng theorem (Theorem \ref{thm:AKZ}), applied after identifying $\mathcal{B}_{\CS}^{(k)} \simeq \overline{\Rep}_q(G)$ (Conjecture \ref{conj:E2-equiv}). The top horizontal arrow is the functor on factorization homology categories induced by the cobordism M; the bottom horizontal arrow is the linear map between distinguished objects, which is the RT partition function $\Zrt(M): V_{\mathrm{RT}}(\Sigma_1) \to V_{\mathrm{RT}}(\Sigma_2)$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Nima Moshayedi | 2026-04-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000119 | OC-1601

From BV-BFV Quantization to Reshetikhin-Turaev Invariants | RT_paper.tex:1713-1715

\label{conj:quant-agree} The BV-BFV quantization of Chern--Simons theory on $D^2 \times [0,1]$ (producing the $\EE_1$-category $\mathcal{B}_{\CS}^{(k)}$ of boundary conditions on the lateral circle $S^1 = \partial D^2$, as in Construction \ref{constr:BCS}) agrees with the shifted deformation quantization of the 1-shifted symplectic stack $\Loc_G(S^1) = [G/G]$, as constructed in the framework of CPTVV \cite{CPTVV2017}, Safronov \cite{Safronov2017b}, and Etingof--Kazhdan \cite{EK1996}.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Nima Moshayedi | 2026-04-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000120 | OC-1602

From BV-BFV Quantization to Reshetikhin-Turaev Invariants | RT_paper.tex:1778-1784

\label{conj:Lagrangian-quant} The Lagrangian morphism \begin{equation} \Loc_G(M) \longrightarrow \Loc_G(\Sigma_1) \times \Loc_G(\Sigma_2)^- \end{equation} of Theorem \ref{thm:AKSZ-PTVV} (where the superscript $(-)$ denotes the opposite symplectic structure) quantizes to a bimodule over the quantized algebras of Σ_1 and Σ_2. This bimodule, viewed as a linear map between the quantized state spaces, is the BV-BFV partition function $\Zbv(M)$. Under the identifications of Step 3, it equals the RT map $\Zrt(M)$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Nima Moshayedi | 2026-04-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000121 | OC-1603

From BV-BFV Quantization to Reshetikhin-Turaev Invariants | RT_paper.tex:1869-1871

\label{conj:cellular} The cellular BV-BFV partition function on a triangulated 3-manifold (M, Δ), computed by iteratively gluing finite-dimensional BV-BFV data along simplices (Construction \ref{constr:cellular}), is equivalent to the value $Z_{\mathcal{B}_{\CS}^{(k)}}(M)$ assigned to M by the $(3\text{-}2\text{-}1)$-extended TQFT determined by the $\EE_2$-category $\mathcal{B}_{\CS}^{(k)}$. More precisely, the iterated BV pushforward along simplices of Δ should reproduce the same linear map between state spaces that the extended TQFT assigns to the cobordism M, with the factorization homology of surfaces providing

the intermediate state-space identifications.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Nima Moshayedi | 2026-04-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000122 | OC-1604

From BV-BFV Quantization to Reshetikhin-Turaev Invariants | RT_paper.tex:1985-1991

\label{conj:koszul} The category of Chern--Simons boundary conditions is equivalent to ind-coherent sheaves on the formal completion of the character stack, i.e., we have \begin{equation} \mathcal{B}_{\CS} \;\simeq\; \IndCoh\big(\Loc_G(M)^{\wedge}\big), \end{equation} where $\IndCoh$ denotes the category of ind-coherent sheaves in the sense of Gaitsgory--Rozenblyum \cite{GR2017} and $\Loc_G(M)^{\wedge}$ is the formal completion of the derived character stack along its classical truncation. Concretely, this means that the BV-BFV partition function around each flat connection determines (and is determined by) the local structure of an ind-coherent sheaf at the corresponding point of the character stack, and the global sheaf encodes all of the perturbative data simultaneously, together with the non-perturbative ``gluing data'' that relates the different perturbative expansions to one another.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Nima Moshayedi | 2026-04-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000123 | OC-1605

From BV-BFV Quantization to Reshetikhin-Turaev Invariants | RT_paper.tex:2102-2114

\label{conj:resurgence-koszul} The Stokes data governing the resurgent structure of the perturbative Chern--Simons partition function are identified with the transition maps in the Koszul duality reconstruction \eqref{eq:koszul-reconstruction}. Specifically, the following hold: \begin{enumerate}[label=(\roman*)] \item The alien derivative $\Delta_{\omega_{AB}}$, which measures the non-perturbative tunneling between flat connections A and B, corresponds to a morphism in $\IndCoh(\Loc_G(M)^\wedge)$ from the skyscraper sheaf $\delta_{[A]}$ to the skyscraper sheaf $\delta_{[B]}$, given by the residue of an ind-coherent sheaf at the connecting singularity of the character stack. \item The Stokes automorphism $\mathfrak{S}_\theta$ associated to a ray θ in the Borel plane corresponds to the wall-crossing automorphism of $\IndCoh(\Loc_G(M)^\wedge)$ associated to the corresponding stability condition on the derived category. \item The full resurgent structure, which is the totality of alien derivatives and their compositions, forming the ``alien algebra'', is equivalent to the A_∞-structure on the Ext-algebra \begin{equation} \bigoplus_{[A],\,[B] \in \Flat_G(M)/G} \mathrm{Ext}^\bullet_{\IndCoh}\!\big(\delta_{[A]},\; \delta_{[B]}\big). \end{equation} \end{enumerate}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Nima Moshayedi | 2026-04-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000124 | OC-1607

From BV-BFV Quantization to Reshetikhin-Turaev Invariants | RT_paper.tex:2321-2323

\label{conj:categorification} Khovanov homology of a link $L \subset S^3$ is the value of a fully extended 4-dimensional TQFT on the link complement $S^3 \setminus L$, viewed as a 3-manifold with boundary. The relevant 4-dimensional theory is associated to an $\EE_4$-algebra obtained from BV-BFV quantization of 4-dimensional Chern--Simons theory (in the sense of Costello) or Kapustin--Witten theory.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Nima Moshayedi | 2026-04-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000125 | OC-1929

Yang-Baxter Integrability and Exceptional-Point Structure in Pseudo-Hermitian Quantum Impurity Systems | main.tex:1784-1789

\label{conj:broken_string} In the PT-broken phase, physical rapidities form strings of the form $k_j^{R,(n)}=k_0+\ii\kappa+\frac{\ii}{2}(n+1-2j)$ with imaginary center shift $\kappa=\sqrt{\beta^2-\gamma^2}>0$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Vinayak M. Kulkarni | 2026-04-23

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000126 | OC-4390

Level lines of the Gaussian free field and $c=1$ degenerate conformal blocks | GFF_level_lines_arXiv.tex:1892-1894

\label{conj:existence_fused_multiple_sle} In each polygon, there exists a unique global $\aSLE_\kappa$ for each $\multii$-valenced link pattern $\alpha \in \LP_\multii$ and for each $\kappa \in (0,8)$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Alex Karrila; Eveliina Peltola; Lukas Schoug | 2026-06-20

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000127 | OC-722

Equilibrium measures for higher dimensional rotationally symmetric Riesz gases | Riesz_equilibrium.tex:1012-1014

\label{Cj1} The equilibrium measure is supported in $\R^d$ if and only if $a \ge a_{ \rm cri }$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Sung-Soo Byun; Peter J. Forrester; Satya N. Majumdar; Gregory Schehr | 2026-02-03

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000128 | OC-4602

Ordered alternating paths and the depth of symbolic powers of cover ideals of graphs | Depth_Powers_Cover_Tree.tex:370-374

Let G be a weakly chordal graph on n vertices. Then $$\operatorname{depth}(S/J(G)^{(t)}) = n - 1 - \alpha_t(G)$$ for all $t \ge 1$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Nguyen Thu Hang; Nguyen Thi Thanh Tam; Thanh Vu | 2026-07-05

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000129 | OC-843

The Generators of a Colon Ideal with an Application to the Weak Lefschetz Property for Monomial Almost Complete Intersections in Three Variables | main.tex:155-161

\label{cook-nagel} Consider the ideal $I=(x^{a_1+t}, y^{a_2+t}, z^{a_3+t}, x^{a_1}y^{a_2}z^{a_3})\subseteq\F[x, y, z]$ where a_1, a_2, a_3, t satisfy the following three conditions: \begin{equation}\label{conditions} 0<a_1\le a_2 \le a_3\le 2(a_1+a_2),\ 3\ \text{divides}\ a_1+a_2+a_3, \ \text{and} \ t\ge \frac{a_1+a_2+a_3}{3}. \end{equation} If (a_1, a_2, a_3, t) is not $(2, 9, 13, 9)$ or $(3, 7, 14, 9)$, then $\F[x, y, z]/I$ fails WLP if and only if t is even, $a_1+a_2+a_3$ is odd, and $a_1=a_2$ or $a_2=a_3$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Matthew Davidson Booth; Adela Vraciu | 2026-03-12

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000130 | OC-109

Moduli of sheaves on hyperkähler manifolds | colorado-notes-2025.tex:2750-2754

\label{cnj:atomicoliscio} Let $\cF$ be an h-slope stable atomic vector bundle on a polarized (projective) HK manifold (X,h). Then the deformation space $\Def(X)$ is smooth.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Kieran G. O'Grady | 2026-02-26

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000131 | OC-127

On a conjecture of Fujino and Sato | Note_FS.tex:362-368

\label{conj.1.4} Let X be a complete $\Q$-factorial toric variety. Then there exists a finite sequence consisting of flips, flops, or anti-flips \begin{equation*} X=:X_0\dashrightarrow X_1\dashrightarrow\cdots\dashrightarrow X_m=:X' \end{equation*} such that X' is a projective $\Q$-factorial toric variety.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Michele Rossi | 2026-02-26

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000132 | OC-1566

The Pandharipande-Thomas rationality conjecture for superpositive curve classes on projective complex 3-manifolds | pt.tex:371-392

\label{pt1conj1} {\bf(a)} $PT_\be\bigl(\ts\prod_{i=1}^m\tau_{k_i}(\eta_i),q\bigr)$ in \eq{pt1eq4} is the Laurent expansion in q of a rational function $F(q)\in\Q(q)$. \smallskip \noindent{\bf(b)} The poles of\/ $F(q)$ occur only at\/ $q=0$ and at roots of unity. \smallskip \noindent{\bf(c)} The rational function $PT_\be\bigl(\ts\prod_{i=1}^m\tau_{k_i}(\eta_i),q\bigr)$ satisfies \begin{equation*} PT_\be\bigl(\ts\prod_{i=1}^m\tau_{k_i}(\eta_i),q^{-1}\bigr)= (-1)^{\sum_{i=1}^mk_i}q^{-c_1(X)\cdot\be}PT_\be\bigl(\ts\prod_{i=1}^m\tau_{k_i}(\eta_i),q\bigr). \end{equation*} \noindent{\bf(d)} More generally, $PT^G_\be\bigl(\ts\prod_{i=1}^m\tau_{k_i}(\eta_i),q\bigr)$ in \eq{pt1eq6} is the Laurent expansion in q of a rational function $F^G(q)\in H^*_G(*,\Q)(q)$. \smallskip \noindent{\bf(e)} The poles of\/ $F^G(q)$ occur only at\/ $q=0$ and at roots of unity. \smallskip \noindent{\bf(f)} The analogue of\/ {\bf(c)} holds for $PT^G_\be\bigl(\ts\prod_{i=1}^m\tau_{k_i}(\eta_i),q\bigr)$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Reginald Anderson; Dominic Joyce | 2026-04-07

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000133 | OC-159

Topology of projective Tate-Shafarevich twists | main.tex:68-70

\label{saccaconj} Total spaces of projective Lagrangian fibrations that are related by a Tate-Shafarevich twist are deformation-equivalent.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | David Zhiyuan Bai | 2026-02-25

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000134 | OC-195

On moduli of Fano varieties: an introduction to K-stability and K-moduli | bootcamp.tex:1239-1241

GIT = K stability for cubic hypersurfaces.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Kristin DeVleming | 2026-02-23

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000135 | OC-2143

Towards Bigness equivalence | towads-bigness-equivalence3.tex:117-118

$Q^a_s $ on $ \mathcal{F}l_s(E)$ is big if only if $\pi_*(Q^a_s)=S_a(E)$ on X is big.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | F. Laytimi; D. S. Nagaraj | 2026-04-30

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000136 | OC-2553

The strong monodromy conjecture for hyperplane arrangements | main.tex:275-281

\label{conj:tmc} If c is a pole of the topological zeta function $Z_f(s)$, then: \begin{itemize} \item $($Weak version$):$ $\exp (2\pi i c)$ is an eigenvalue of the monodromy action on the cohomology of the Milnor fiber. \item $($Strong version$):$ c is a root of the Bernstein--Sato polynomial of f. \end{itemize}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Lei Wu | 2026-05-25

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000137 | OC-2619

Functional codes arising from rank n Hermitian varieties and hypersurfaces in low dimensions | main.tex:531-543

\cite[Conjecture 2 (ii)]{ELX}\label{ELXc} Let $n\geq 3$ and $\U_n$ be a non-degenerate Hermitian variety in $\PP^n(\Fqt)$. If $V(F)$ is a hypersurface of degree $d\leq q$, then $$M_n(d)=\max_{V(F)} |V(F)(\Fqt)\cap \U_n|=\begin{cases} d|\U_{n-1}(\Fqt)|-(d-1)|\U_{n-2}(\Fqt)| \ \text{if} \ n \ \text{is even}\\ (dq^2-d+1)|\U_{n-2}(\Fqt)|+d\ \text{if} \ n \ \text{is odd}, \end{cases}$$ and this maximum is attained if and only if the hypersurface $V(F)$ is union of d distinct hyperplanes $\Sigma_1, \Sigma_2,\dots, \Sigma_d$ such that each of the hyperplanes contains a common $(n-2)$-dimensional linear space Π_{n-2} and $\Pi_{n-2}\cap \U_n$ is a non-degenerate Hermitian variety and: --- If n is even, the d hyperplanes $\Sigma_1,\dots, \Sigma_d$ are non-tangent to $\U_n$. --- If n is odd, the d hyperplanes $\Sigma_1,\dots, \Sigma_d$ are tangent to $\U_n$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Subrata Manna | 2026-05-22

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000138 | OC-2643

On Cox Rings of Calabi-Yau hypersurfaces | final.tex:2276-2284

There exists a finite rational polyhedral cone $\Pi \subseteq \overline{\rm Mov}^{e}(X)$ such that \[\overline{\rm Mov}^{e}(X) = {\rm Bir}(X)\cdot \Pi, \] and such that Π is a fundamental domain for the action of ${\rm Bir}(X)$ on $\overline{\rm Mov}^{e}(X)$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Michela Artebani; Antonio Laface; Luca Ugaglia | 2026-05-21

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000139 | OC-2704

Smooth categories in a 6 functor formalism and compact generation for nuclear categories in analytic geometry | main.tex:741-743

{ramzi2024dualizablepresentableinftycategories}}]\label{conj: congettura Einfty} Let R be an $\mathbb{E}_\infty$-ring specturm and $\mathscr{C}$ be a smooth, stable, presentable R-linear ∞-category. Then $\mathscr{C}$ admits a single compact generator.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Matteo Montagnani | 2026-05-20

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000140 | OC-2718

Infinite transitivity and polynomial vector fields | version-1.tex:535-537

If $\lie(D_1,\ldots D_k)$ has finite codimension in $N(X)$, then the group G acts on $\OOO$ infinitely transitively.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Rafael B. Andrist; Ivan Arzhantsev | 2026-05-19

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000141 | OC-3621

Naive atoms of blowups: examples | ForDummies_arXivVersion.tex:419-421

\label{cIritani} Let $\widetilde{X}$ be the blowup of X in a smooth codimension c subvariety Z. Suppose that the inclusion $Z \hookrightarrow X$ induces an embedding $\mathrm{N}_1 (Z, \mathbb{Z}) \hookrightarrow \mathrm{N}_1 (X, \mathbb{Z})$. Then the naive atomic decomposition of $\widetilde{X}$ is the sequence obtained by concatenating and sorting the naive atomic decompositions of X and $(c-1)$ times the naive atomic decomposition of Z.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Christian Böhning; Hans-Christian Graf von Bothmer; Zac Su'a | 2026-06-16

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000142 | OC-3622

Naive atoms of blowups: examples | ForDummies_arXivVersion.tex:588-590

\label{cIritaniMorePrecise} Let $\widetilde{X}$ be the blowup of a smooth projective variety X in a smooth codimension c subvariety Z. Suppose that the inclusion $Z \hookrightarrow X$ induces an embedding $\mathrm{N}_1 (Z, \mathbb{Z}) \hookrightarrow \mathrm{N}_1 (X, \mathbb{Z})$. Then the naive atomic decomposition of $\widetilde{X}$ is the sequence obtained by concatenating and sorting the naive atomic decompositions of X and $(c-1)$ copies of the naive atomic decomposition of Z.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Christian Böhning; Hans-Christian Graf von Bothmer; Zac Su'a | 2026-06-16

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000143 | OC-381

New irreducible components of $\mathcal{B}(0,c_2)$ and Computation of the Dimension of its tangent space | main.tex:883-890

\label{conjecture} Let \(a\) and \(k\) be positive integers. For each integer \(b\in\{0,1,\dots,k-1\}\) with \(a>b\), there exists a stable rank-2 vector bundle \(\mathcal{E}\) arising as the cohomology of a minimal monad of the form \begin{equation}\label{general-monad} \cM(a, b,k) : 0 \lra

{\mathcal{O}_{\mathbb{P}^{3}}(-a)}^{\oplus k} \stackrel{\alpha}{\lra} \begin{matrix} {\mathcal{O}_{\mathbb{P}^{3}}(a-1)}^{\oplus k} \oplus \mathcal{O}_{\mathbb{P}^{3}}(b) \\ \displaystyle \oplus \\ {\mathcal{O}_{\mathbb{P}^{3}}(1-a)}^{\oplus k}\oplus\mathcal{O}_{\mathbb{P}^{3}}(-b) \end{matrix} \stackrel{\beta}{\lra} {\mathcal{O}_{\mathbb{P}^{3}}(a)}^{\oplus k} \lra 0. \end{equation} Moreover, for all sufficiently large integers \(a \gg 0\), the corresponding family of such bundles belongs to an irreducible component of \(\mathcal{B}(0,k(2a-1)-b^2)\), whose dimension exceeds the expected dimension.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Aislan Fontes; Maxwell Santos | 2026-02-13

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000144 | OC-3832

Functorial stratifications of singularities in characteristic 0 | main.tex:240-242

Let X be a reduced scheme of finite type over a field k of characteristic 0 and $x\in X$ a closed point. Then, we have a canonical inclusion $$\mathrm{rtsp}_x(X)\subset \mathrm{Rid}_x(X).$$

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Vicente Monreal | 2026-06-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by_sa

V3-000145 | OC-3979

Algebraic Hodge generic points are dense | main.tex:525-528

\label{perconj} For all points $s \in S(\Qbar)$ outside the Hodge locus, the $\Qbar$-Zariski closure of $\mathbf{P}(s)$ is $B_{\mathbf{P},s}$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Gregorio Baldi; Gal Binyamini; David Urbanik | 2026-06-07

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000146 | OC-3980

Algebraic Hodge generic points are dense | main.tex:530-532

\label{degdeltaperconj} For all points $s \in S(\Qbar)$ outside the Hodge locus, and for a fixed integer δ, the ideal defining the $\Qbar$-Zariski closure of $\mathbf{P}(s)$ has no elements of degree at most δ defined over $\kappa(s)$ which do not lie in the ideal defining $B_{\mathbf{P},s}$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Gregorio Baldi; Gal Binyamini; David Urbanik | 2026-06-07

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000147 | OC-3982

Algebraic Hodge generic points are dense | main.tex:577-582

(ℓ-Mumford-Tate) \label{MTconj} Let Y be a smooth projective algebraic variety defined over a number field $L \subset \mathbb{C}$, and let ℓ be a prime. Then for each pair of integers (j, k) the Artin comparison map \[H^{2j}_{B}(Y^{k}_{\mathbb{C}}, \mathbb{Q}) \otimes_{\mathbb{Q}} \mathbb{Q}_{\ell}(j) \xrightarrow{\sim} H^{2j}_{\textrm{\'et}}(Y^{k}_{\overline{L}}, \mathbb{Q}_{\ell}(j)) \] identifies the $\mathbb{Q}_{\ell}$-vector spaces spanned by Hodge and Tate classes.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Gregorio Baldi; Gal Binyamini; David Urbanik | 2026-06-07

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000148 | OC-4134

Bondal's conjecture in dimension five | bondal5d.tex:341-343

Let $\X$ be a Fano manifold, and let σ be a Poisson structure on $\X$. Then $\Dgn[2k]{\sigma}$ has an irreducible component of dimension at least $2k+1$ for all integers k such that $0 \leq 2k < \dim \X$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Stéphane Druel; Jorge Vitório Pereira; Brent Pym; Frédéric Touzet | 2026-06-02

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000149 | OC-476

A proof of Dolbeault geometric Langlands for GL_2 with reduced spectral curves | WHcompati.tex:306-313

\emph{(\cite{PTlim})}\label{conj:intro} There is a $\rB$-linear equivalence \begin{align}\label{intro:IndL} \IndCoh_{\mathcal{N}}(\Hig_{{}^{L}G}(w)^{\mathrm{ss}})_{-\chi} \stackrel{\sim}{\to} \IndL_{\mathcal{N}}(\Hig_G(\chi))_w \end{align} which is compatible with Wilson/Hecke operators.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Yukinobu Toda | 2026-02-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000150 | OC-478

A proof of Dolbeault geometric Langlands for GL_2 with reduced spectral curves | WHcompati.tex:1195-1200

\emph{(\cite[Conjecture~1.1]{PTlim})}\label{conj:dl} For $G=\GL_r$ and $(\chi, w)\in \mathbb{Z}^2$, there is a $\rB$-linear equivalence \begin{align*} \IndCoh_{\mathcal{N}}(\Hig_G(w)^{\mathrm{ss}})_{-\chi} \simeq \IndL_{\mathcal{N}}(\Hig_G(\chi))_w. \end{align*}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Yukinobu Toda | 2026-02-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000151 | OC-479

A proof of Dolbeault geometric Langlands for GL_2 with reduced spectral curves | WHcompati.tex:2652-2657

\label{conj:norm} There is an isomorphism \begin{align*} \Phi(\mathcal{O}_{\cH(w)^{\mathrm{ss}}}) \cong s_{!}\mathcal{O}_{\cB} \in \LL_{\mathcal{N}}(\cH(\chi_0))_{w'}. \end{align*}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Yukinobu Toda | 2026-02-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000152 | OC-760

Relative $\mathbb{A}^1$-Contractibility of Smooth Schemes and Exotic Motivic Spheres | Chapters/main.tex:3346-3348

Let S be any Noetherian scheme with characteristic zero residue fields. Let $f: X\to S$ be a smooth scheme of relative dimension 2. Then if X is $\AA^1$-homotopic to $\AA^2_S\bs \{0\}$, then X is isomorphic to $\AA^2_S\bs \{0\}$ as a S-scheme.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Krishna Kumar Madhavan Vijayalakshmi | 2026-03-04

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000153 | OC-797

Cohomological Hall algebras of one-dimensional sheaves on surfaces and Yangians | COHAs_Yangians.tex:3025-3027

The map Φ induces an isomorphism $\Y_\qv^-\otimes \mathsf{Frac}\big(\Hbullet_{\Ttilde}\big) \stackrel{\sim}{\longrightarrow} \cohaqv^{\Ttilde} \otimes \mathsf{Frac}\big(\Hbullet_{\Ttilde}\big)$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Duiliu-Emanuel Diaconescu; Mauro Porta; Francesco Sala; Olivier Schiffmann; Eric Vasserot | 2026-03-03

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000154 | OC-798

Cohomological Hall algebras of one-dimensional sheaves on surfaces and Yangians | COHAs_Yangians.tex:7696-7698

Equality~\eqref{eq:commutations-relations-rank-0-rank-1-i} holds also without the assumption on $\widehat{S}_{C_i}$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Duiliu-Emanuel Diaconescu; Mauro Porta; Francesco Sala; Olivier Schiffmann; Eric Vasserot | 2026-03-03

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000155 | OC-965

On linear $α_p$-quotients | main.tex:782-786

{Tonini_Yasuda_Motivic_McKay_cor_for_alpha_p}}]\label{conj:Tonini_Yasuda} Assume that $\bD_\mathbf{d}=\sum_{\lambda}\frac{d_\lambda(d_\lambda+1)}{2}\geq 2$. Then: $$\mst\big(\bA/(\bZ/p,\mathbf{d})\big) = \mst\big(\bA/(\alpha_p,\mathbf{d})\big).$$

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Quentin Posva; Takehiko Yasuda | 2026-03-07

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000156 | OC-966

On linear $α_p$-quotients | main.tex:2156-2164

\label{conj:equality_of_multisets} Assume that $\bD_\mathbf{d}\geq p$. Define $F=F_{\max\mathbf{d}}$ and N_r as in \autoref{thm:mst_alpha_p_qt}. Then we have the following equality of multi-sets: \begin{align*} &\left \{ \left\{j -\mathrm{sht}(j)\mid 1\le j\le p-1 \right\}\right\} \uplus \left \{ \left\{0,-1,\dots,-\gamma \right\}\right\}\\ &= \left \{ \left\{\theta\left(\frac{s}{r}\right)+i \mid \frac{s}{r} \in F,\, 0\le i \le N_r-1\right\}\right\} \end{align*}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Quentin Posva; Takehiko Yasuda | 2026-03-07

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000157 | OC-1650

Neumann's nodal line may be closed on doubly-connected planar domains | template-article.tex:372-375

Consider the Laplacian on a bounded regular planar domain Ω with Dirichlet or Robin boundary conditions. Then the nodal line of any second eigenfunction cannot be closed if\/ Ω is simply-connected, while there exist doubly-connected domains for which the nodal line is closed and does not touch the boundary.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Pedro Freitas; Roméo Leylekian | 2026-04-03

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000158 | OC-2362

Liouville Theorems for Stationary Navier-Stokes Equations via the Radial Velocity Component | v1.tex:219-225

Show that any solution $\vu$ of (\ref{SNS}) fulfilling the conditions \begin{equation}\label{Conjecture} \vu \in \dot{H}^1(\R^3) \qquad\mbox{and}\qquad \vu(x)\to 0 \mbox{ as } |x|\to +\infty, \end{equation} is identically equal to zero.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Vergara-Hermosilla, Gaston | 2026-05-07

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000159 | OC-2521

Robin nullity and asymptotic geometry of the critical hyperbolic catenoid | RobinNull_CritHyperCatenoid.tex:1345-1348

\label{conj:uniqueness} There exists a unique $a^\sharp\in(1,\infty)$ with $r'(a^\sharp)=0$. Moreover, $r'<0$ on $(1,a^\sharp)$ and $r'>0$ on $(a^\sharp,\infty)$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Pigazzini, Alexander | 2026-05-01

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000160 | OC-3825

Elliptic criticality versus Volterra memory in indirect chemotaxis cascades | arxiv.tex:951-964

\label{conj:sharp-PES-threshold} In the four-dimensional critical regime, the sharp logarithmic interaction constant should yield the threshold \begin{equation} M_*=\frac{64\pi^2\tau}{\chi}. \label{conjectured-critical-mass} \end{equation} Equivalently, one expects \(\mathcal F_{\mathrm{PES}}\) to be bounded below on \(\mathcal A_M(\Omega)\) for \(M<M_*\), to lose compactness at \(M=M_*\), and to be unbounded below for \(M>M_*\). This statement is conjectural unless the corresponding sharp \(K_\tau\)-adapted logarithmic HLS/Adams inequality is proved.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Louis Shuo Wang | 2026-06-11

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000161 | OC-3826

Invisible Defect Cascades for Navier-Stokes Regularity | invisible_defect_cascades_refs_checked.tex:4893-4908

Let \((u,p)\) be a suitable weak solution and let \((0,0)\) have a non-CKN dyadic branch: \[\Psi(r_n)>\eps_{\CKN} \qquad\text{for all sufficiently large } n. \] Then there exist relative filter scales \(\ell_n\in[\ell_-,\ell_+]\subset(0,1)\), a subsequence, and a constant \(c>0\) such that \begin{equation} \Vort_{n,\ell_n} + \Press^{\act}_{n} + \Flux^{\absop}_{n,\ell_n} \ge c. \end{equation}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Runlong Yu | 2026-06-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000162 | OC-4279

A Structural Audit of Navier-Stokes Obstruction Calculus | structural_audit_navier_stokes_final_cleaned.tex:1261-1275

\label{conj:mechanism} If \[\Psi(r_n)\ge\eps_{\CKN} \] for all sufficiently large n, then there exist relative filter scales ℓ_n in a compact subinterval of $(0,1)$ and a subsequence such that \begin{equation}\label{eq:mechanism-conjecture} \limsup_{n\to\infty} \left(V_{n,\ell_n}^+ +P_n^{\rm act} +F_{n,\ell_n}^{\rm abs} \right)>0. \end{equation}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Runlong Yu | 2026-06-24

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000163 | OC-4832

Strichartz Estimates for the Liouville Equation on Euclidean Tori and Applications to Kakeya | v1-main-liouville.tex:344-354

\label{conjecture1} There holds \[\| \rho \|_{L^p_{t,x}(\mathbb{R} \times \mathbb{R}^d)} \lesssim \| f_0 \|_{L^a_v (\mathbb{R}^d, L^b_x(\mathbb{R}^d))} \] if \[a'=dp, \;\;\;\;\;\; b = \frac {dp}{d+1}, \;\;\;\;\;\; p > \frac{d+1}{d}. \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Pierre Germain; Mickaël Latocca | 2026-06-29

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000164 | OC-721

The Small-Dispersion Limit of the Intermediate Long Wave Equation via Semiclassical Soliton Ensembles | main.tex:458-489

\label{conj:WeylLaw} Let u_0 be an admissible initial condition according to Definition \ref{def:AdmisInitCond} and let $R^\wyl: [1/2\delta, \zeta_\maxrm]_{\mathcal{Q}_\delta} \to \R_+$ be defined by \begin{equation} R^\wyl(\zeta) := \frac{1}{4 \delta} \int_{x_-(\zeta)}^{x_+(\zeta)} W_{0}\left(-2 \delta \zeta \ee^{-2\delta(\zeta + u_0(x))} \right) - W_{-1}\left(-2 \delta \zeta \ee^{-2\delta(\zeta + u_0(x))}\right) \, \dd x \, . \label{eq:RWeylLawDef} \end{equation} where $x_\pm : (1/2\delta, \zeta_\maxrm]_{\quadratrix{\delta}} \to \R$ are the turning point functions defined implicitly by \begin{equation} u_0(x_\pm(\zeta)) = E(\zeta) \label{eq:TurningPointFunctDef} \end{equation} with E as in \eqref{eq:TurningPointFromELevel} and such that $x_-(\zeta) \leq x_\maxrm \leq x_+(\zeta)$. Additionally, W_n for $n \in \Z$ are the standard branches of the Lambert-W function \cite[Equation 4.13.1_1]{DLMF}. The asymptotics of the eigenvalues satisfy the ``ILW Weyl law'' \begin{equation} R^\wyl(\zeta_n) = \left(n - \frac{1}{2} \right) \pi \epsilon + \bigO{\epsilon^2} \label{eq:WeylLaw} \end{equation} for $n \in \N$. Defining the tail integrals $\theta_\pm: [1/2\delta, \zeta_{\maxrm}]_{\quadratrix{\delta}} \to \R$ \begin{equation} \theta_\pm(\zeta) := \im{\zeta} x_\pm(\zeta) + \int^{\pm\infty}_{x_\pm(\zeta)} \im{ \zeta + \frac{1}{2 \delta} W_{-1}\left(-2\delta \zeta \ee^{-2\delta(\zeta + u_0(x))}\right)} \, \dd x \, . \label{eq:TailIntegralsDef} \end{equation} Then the norming constants have the asymptotics \begin{align} c_n &= \exp \Bigg(\frac{2}{\epsilon} \theta_+(\zeta_n) \Bigg) \left| \frac{(2 \delta)^2 \zeta_n}{1 -2 \delta \zeta_n} \right| \nonumber \\ &\Quad[2] \times \left(\int_{x_-(\zeta_n)}^{x_+(\zeta_n)} \frac{2\delta W_{-1}(-2\delta \zeta_n \ee^{-2\delta(\zeta_n + u_0(x))})}{1+W_{-1}(-2\delta \zeta_n \ee^{-2\delta(\zeta_n + u_0(x))})} - \frac{2\delta W_0(-2\delta \zeta_n \ee^{-2\delta(\zeta_n + u_0(x))})}{1+W_0(-2\delta \zeta_n \ee^{-2\delta(\zeta_n + u_0(x))})} \, \dd x \right)^{-1} \nonumber \\ &\Quad[2] \times \left(1 + \bigO{\epsilon^{1/6}} \right) \, . \label{eq:NormingConstAsymptoticFormula} \end{align} Finally, the reflection coefficient is small beyond all orders in ϵ.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Matthew Dominique Mitchell | 2026-02-03

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000165 | OC-1374

Poincaré duality spaces related to the Joker | PD-Joker-v1.tex:1111-1114

\label{conj:Manifold} There is a 2-local homotopy equivalence $\mathrm{G}_2/\SO(4)\xrightarrow{\;\simeq\;}J^8$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Andrew Baker | 2026-03-31

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000166 | OC-2533

Unraveling the Bott spiral | BOTT_SPIRAL.tex:1792-1819

\label{direct_SM_conj} With notation as in \cref{interacting_symm_mon}, there is a symmetric monoidal functor $\mathit{TW}\colon\cat{FermGrp}\to\Top_{/B\O/B\Spin}$ of ∞-categories such that \begin{subequations} \begin{enumerate} \item TW of a fermionic group G_f with twisting data (θ,ω) is homotopy equivalent to the map \begin{equation} BG_b\xrightarrow{(\theta,\omega)} B\Z/2\times B^2\Z/2\xrightarrow[\eqref{faux_whitney}]{\simeq} B\O/B\Spin. \end{equation} \item The following diagram of symmetric monoidal functors commutes. \begin{equation} \begin{tikzcd} {\cat{Rep}^{\mathrm{cpt}}} & {\cat{Top}_{/B\O}} & {\cat{Spectra}} \\ {\cat{FermGrp}} & {\cat{Top}_{/B\O/B\Spin}} & {\cat{Mod}_{\MTSpin}.} \arrow["\xi", "{\eqref{xi_defn}}"', from=1-1, to=1-2] \arrow["\zeta"', "{\eqref{zeta_defn}}", from=1-1, to=2-1] \arrow["M", from=1-2, to=1-3] \arrow["q", from=1-2, to=2-2] \arrow["{\wedge\MTSpin}", from=1-3, to=2-3] \arrow["{\mathit{TW}}"', color={BrickRed}, from=2-1, to=2-2] \arrow["T"', from=2-2, to=2-3] \end{tikzcd} \end{equation} Here q is induced from the usual map $B\O\to B\O/B\Spin$ and $T\colon\cat{Top}_{/B\O/B\Spin}\to \cat{Mod}_{\MTSpin}$ is the $\MTSpin$-module Thom spectrum functor: see, e.g., \cite[Theorem 1.13]{DY23}, based on~\cite[Lemma IV.2.6]{MQRT77} and~\cite{ABGHR14a, ABGHR14b}. \end{enumerate} \end{subequations}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Debray, Arun; Krulewski, Cameron; Stehouwer, Luuk | 2026-05-01

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000167 | OC-391

On the equivariant triangulation of some small covers | main.tex:181-183

\cite{Lutz16vertex}\label{RP4conjecture} The minimal triangulation of $\RP^4$ is unique with 16-vertices.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Raju Kumar Gupta; Soumen Sarkar | 2026-02-13

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000168 | OC-2633

A conjecture for arithmetic spherical maximal functions | 20260521-arithmetic_spherical_averages.tex:163-170

\label{conj:2016} Let \(\Lambda \subset \N\). Let \(d \geq 5\). \(M_{\Lambda} \) is bounded on \(\ell^p(\Z^\dimension)\) for all \(p>\eta(\Lambda)\).

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Kevin Hughes | 2026-05-21

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by_sa

V3-000169 | OC-1002

Graph labellings and external difference families | AngHucMcC_arxiv.tex:376-378

\label{nearalphatree} Every tree has a near α-valuation.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Gavin Angus; Sophie Huczynska; Struan McCartney | 2026-03-05

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000170 | OC-1304

Deficit and (q,t)-symmetry in triangular partitions | draft.tex:3497-3504

\label{conj:lattice} If λ is a triangular partition such that its top-down tableau θ is a sim-sym tableau, there exists a Schur-positive symmetric function λ such that \begin{equation} A_\lambda(q,t,1) = \sum_{\tau \preceq \mu} q^{\dist(\tau, \mu)} t^{\ssim_\theta(\tau, \mu)} \end{equation} where the sum runs over intervals in the ν-Tamari lattice of λ.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Viviane Pons; Loïc Le Mogne | 2026-03-31

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000171 | OC-150

On the expansion formulas of cluster varieties from surfaces and their combinatorial properties | DinhIp_Final__edit.tex:5797-5798

Fix a non-$\cT$ diagonal $\gamma = AB$ in a general n-triangulated m-gon with $\cT$-triangulation of type $\mathcal{P}(m_1,n_1,\ldots,m_k,n_k)$. Then for $l = 1,2,...,n$ and $i,j \in \mathbb{N}$ such that $i+j \leq n$, every function $D^{[m_1,n_1,\ldots,m_k,n_k]}_l$ and $I^{[m_1,n_1,\ldots,m_k,n_k]}_{(i,j)}$ is a Laurent polynomial of variables $x^{(r)}_{(a,b,c)}$ where $r = 1,2,...,(m-2)$ and $(a,b,c) \in \Gamma_{n+1}$ with each coefficient being 1.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Vu Tung Lam Dinh; Ivan Chi-Ho Ip | 2026-02-25

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000172 | OC-1546

Helly Theorems for Generalized Turán Problems | arXivHellyTuran.tex:1500-1504

\label{optimal path conjecture} Let k,t be integers with $k\ge 2$ and $t\ge 2k+1$. If $0\le r\le k-2$ is the unique integer such that $t-3\equiv r \mod k-1$, then every family of graphs $\c{F}$ either satisfies \[\ex(n,P_t,\c{F})=O(\ex(n,\c{F})^{r+2}n^{k-2-r}),\] or $\ex(n,P_t,\c{F})=\Omega(n^{k+1})$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Sean English; Sam Spiro | 2026-04-07

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000173 | OC-157

Equivariant log concavity and the $\operatorname{FI^\sharp}$-module structure on $H^i(\operatorname{Conf}(n,\mathbb{R}^d))$ | main.tex:277-282

\label{mainConjecture} Given positive integers with $i<j\leq k <\ell$ and $i+\ell = j+k$, there is an inclusion of $\FIS$-modules: \[A^i\otimes A^\ell \hookrightarrow A^j\otimes A^k,\] and \[C^{i}\otimes C^{\ell}\hookrightarrow C^{j}\otimes C^{k}.\]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Benjamin Homan | 2026-02-25

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000174 | OC-1966

Improving the Even-Size Threshold in Spectral Extrema for $H(4,3)$-Free Graphs | Improving_the_Even-Size_Threshold_in_Spectral_Extrema_for_H4,3-Free_Graphs.tex:607-609

The sharp threshold in \cref{thm:main} is 18.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Shreyhaan Sarkar | 2026-04-21

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000175 | OC-2504

Families without s-matchings: the other end | main__1_.tex:80-85

For all n,s, \[e(n,s)=\max\bigl\{|\mathcal P(m,s,\ell)|,\ |\mathcal P'(m,s,\ell)|,\ |\mathcal Q(m,s,\ell)|,\ |\mathcal W(m,s,\ell)|\bigr\}. \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Kupavskii, Andrey; Sokolov, Georgy | 2026-05-01

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000176 | OC-2612

A Balancing Theorem for Spanning Trees of Rectangular Grid Graphs | main.tex:100-105

\label{conj:induced-grid} Let $n\ge 1$ be an integer. Let $S \subset \mathbb{Z}^2$ be a finite set of vertices such that $|S|=n^2$, and suppose that the induced subgraph $\mathcal{L}[S]$ is connected. Then \[\tau(\mathcal{L}[S]) \le \tau(R_{n,n}). \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Jiechen Zhang | 2026-05-22

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000177 | OC-2824

Boolean--Eulerian numbers | boolean-euler-May14.tex:612-615

\label{conj:vertical} For fixed $k \geq 2$, the sequence $\boel(n,k)$ for $n \geq k$ is log-concave.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Miklós Bóna; Vincent Vatter | 2026-05-14

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000178 | OC-2882

Dyck Symmetric Functions and Applications to \(q,t\)-Catalan Polynomials | arxiv_submission.tex:4849-4857

\label{conj:middle-coefficients} Let \(M=\binom n2\). For any \(n\ge 1\) and any \(0\le d\le \lfloor M/3\rfloor\), the coefficient of \(q^j t^{M-d-j}\) in \(C_n(q,t)\) is the same for all integers \(j\) satisfying \[d\le j\le M-2d. \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Graham Hawkes | 2026-05-13

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000179 | OC-2885

Temperley-Lieb Immanants of Ribbon Decomposition Matrices | main.tex:152-154

\label{conj:KL} Given an infinite ribbon R, a skew shape λ/μ compatible with R. All Kazhdan--Lusztig immannants of the ribbon decomposition matrix $A_{\lambda/\mu,R}$ are Schur positive.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Son Nguyen; Pavlo Pylyavskyy | 2026-05-13

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000180 | OC-2917

Critical Slow Growth in Averaged Meta-Fibonacci Recursions | main.tex:1040-1052

For every fixed \(m\ge2\), there exists a nonempty interval \[1<\alpha<\alpha_m^\ast \] such that the recursion is globally well-defined and satisfies \[Q_{\alpha,m}(n) \sim \left(1-\frac1\alpha\right)n. \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Marco Mantovanelli | 2026-05-11

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000181 | OC-2918

Critical Slow Growth in Averaged Meta-Fibonacci Recursions | main.tex:1432-1447

Let \(p>0\), and replace the arithmetic mean in the recursion by the \(L^p\)-mean \[M_p(x_1,\dots,x_m) = \left(\frac1m\sum_{j=1}^m x_j^p \right)^{1/p}. \] At criticality \(\alpha=1\), the resulting recursion still satisfies \[Q(n)\sim \sqrt{2n}. \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Marco Mantovanelli | 2026-05-11

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000182 | OC-3468

A Degree-Preserving Builder--Chooser Game | DPG_clique_game.tex:901-907

\label{conj:universality} For every $k\ge 3$, \[\rhoK(k)<\infty. \] Equivalently, every clique can be forced from some triangle-free seed.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | András London | 2026-05-26

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000183 | OC-3469

A Degree-Preserving Builder--Chooser Game | DPG_clique_game.tex:955-962

\label{conj:amplifier} For every $r\ge 3$, if there exists a triangle-free graph G such that \[\tau_{K_r}(G)<\infty, \] then there exists a triangle-free graph G' such that Builder can force two compatible supported copies of K_r from G'.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | András London | 2026-05-26

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000184 | OC-3564

Quasi-random graphs, subgraph counts and graph limits, again | sj403.tex:2489-2491

\label{Conj:r>2} If $r\ge3$, then no $k\ge2$ is bad.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Svante Janson | 2026-06-18

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000185 | OC-3565

Quasi-random graphs, subgraph counts and graph limits, again | sj403.tex:2586-2591

\label{CB} Suppose that $\cK=\set1$, and that $e(F)>1$, Then $\tcPm(F;\gaar)$ is a \qrp.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Svante Janson | 2026-06-18

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000186 | OC-3566

Quasi-random graphs, subgraph counts and graph limits, again | sj403.tex:2647-2652

For any $\cK$, if $\tcPm(F;\gaar)$ is {not} \qr, then there is a counterexample with a 2-type graphon.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Svante Janson | 2026-06-18

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000187 | OC-3567

Quasi-random graphs, subgraph counts and graph limits, again | sj403.tex:2656-2660

\label{CC} Suppose that $r>1$ and $e(F)>1$, Then $\tcPm(F;\gaar)$ is a \qrp.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Svante Janson | 2026-06-18

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000188 | OC-3641

Skew column RSK dynamics and the box-ball system | main.tex:3018-3021

\label{conj:general_energy_change_one} The assumption $\ell(\mu) \leq n$ in \cref{th:leading_f0} can be reduced to simply $n > 2$ or $\ell(\mu) \leq 2$ for some other element $\overline{H_{\hw}}$ when $n = \max(H_\hw)$ in the same classical component as $H_{\hw}$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Takashi Imamura; Matteo Mucciconi; Tomohiro Sasamoto; Travis Scrimshaw | 2026-06-16

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000189 | OC-3676

Quasi-kernels in Hereditary Classes and Applications to Break | main-qk.tex:976-979

Every break digraph on \(n\) vertices has a quasi-kernel of size at most \(\frac n2\).

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Jiangdong Ai; Tianyu Huang; Xiangzhou Liu; Chaoliang Tang; Zongqun Xu | 2026-06-15

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000190 | OC-3708

A sharp p-biased product bound for r-cross-intersecting families | cross-int-Jun12.tex:76-83

Let $\max_{i\in[r]}p_i\le \frac{r-1}{r},$ and let \(\F_1,\ldots,\F_r\subseteq2^{[n]}\) be \(r\)-cross-intersecting. Then \[\prod_{i=1}^r \mu_{p_i}(\F_i)\le \prod_{i=1}^r p_i. \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Fan Chang; Hong Liu; Miao Liu | 2026-06-14

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000191 | OC-3814

Critical Sets in Latin Squares and Associated Structures | rb12.tex:492-494

\ \ \lcs{n} $ \leq n^2 - n^{3/2}$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Richard Winston Bean | 2026-06-11

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000192 | OC-3815

Critical Sets in Latin Squares and Associated Structures | rb12.tex:501-503

\ \ \lcs{n} $\leq (1 - (\frac{3}{4})^{log_2{n}}) n^{2}$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Richard Winston Bean | 2026-06-11

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000193 | OC-385

Between proper and square coloring of planar graphs, hardness and extremal graphs | main.tex:64-71

For every planar graph G, \begin{align*} &\bullet~\chi_2(G) \leq 7 & \textrm{if } \Delta(G) = 3, \\ &\bullet~\chi_2(G) \leq \Delta(G) + 5 & \textrm{if } 4 \leq \Delta(G) \leq 7, \\ &\bullet~\chi_2(G) \leq \lfloor 3\Delta(G)/2\rfloor + 1 & \textrm{otherwise.} \end{align*}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Thomas Delépine | 2026-02-13

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000194 | OC-3873

A parity Erdős-Hajnal theorem for t-intersecting curves | mighty_EH.tex:266-269

For every integer \(k\ge 2\), every \(k\)-quasi-planar topological graph on \(n\) vertices has \(O_k(n)\) edges.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Andrew Suk; Su Zhou | 2026-06-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000195 | OC-4055

Equivariant Quantum Cohomology of Grassmannians via the Clifford algebra | arxivsubmission.tex:1557-1559

The coefficients in \eqref{triple} are positive, i.e. $C^{\nu,d}_{\lambda \mu}(y,z) \in \Z_{\geq 0}[y_i - z_j]$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Christian Korff; Mikhail Vasilev | 2026-06-04

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000196 | OC-4069

Frayed Demazure weaves for Poisson-compatible cluster structures on Bott--Samelson charts | weaves.tex:2553-2560

For any word $\underline{w}$, the frayed weave consisting only of fraying vertices \begin{align*} \mathfrak{w}_{\mathrm{rev}}:(i_1,i_2,\dots,i_{\ell-1},i_\ell) &\to (i_1,i_2,\dots,i_{\ell-1},-i_\ell) \to (i_1,i_2,\dots,-i_{\ell-1},-i_\ell),\\ &\to \cdots \to (i_1,-i_2,\dots,-i_{\ell-1},-i_\ell)\to (i_1,-i_2,\dots,-i_{\ell-1},-i_\ell), \end{align*} corresponding to the reversed opening sequence $(\ell, \ell-1,\dots,2,1)$ has its moduli of flags $M(\mathfrak{w}_{\mathrm{rev}})$ equal to the the intersection of all cluster tori in $\cO^{\underline{w}}$ reached along the maximal green sequence $\overrightarrow{\mu_{\underline{w}}}$ applied to $\seed(\underline{w})$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Jon Cheah | 2026-06-04

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000197 | OC-4116

A Structural Approach to Burning Number | A_Structural_Approach_to_Burning_Number.tex:144-145

\label{Conjecture} \emph{(\cite{Bonato1})} If G is a connected graph of order n, then $b(G) \leq \lceil \sqrt{n} \rceil. $

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Jean Guillaume; Tyriana Williams | 2026-06-02

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000198 | OC-4327

Exact Enumeration of Phylogenetic Networks: The Tree-Child, Reticulation-Visible and Orchard Hierarchy | full_paper_arxiv.tex:896-909

\label{conj:pattern} For all $k\geq2$ and $\ell\geq k+1$, \begin{equation}\label{eq:pattern} \Delta_k(\ell) \;=\; A_k(\ell)\,(2\ell-3)!! \;-\; B_k(\ell)\,2^{\ell-k+1}\,\ell!, \end{equation} where $A_k(\ell)$ and $B_k(\ell)$ are polynomials in ℓ with rational coefficients satisfying: \begin{enumerate}[label=(\roman*),leftmargin=*] \item $\deg A_k = 2k-1$ and $\operatorname{lead}(A_k) = 2^k$; \item $\deg B_k = 2k-3$; \item $A_k(\ell)\in\mathbb{Z}$ and $B_k(\ell)\in\mathbb{Z}$ for all $\ell\in\mathbb{Z}_{\geq1}$. \end{enumerate}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Josep Batle | 2026-06-23

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000199 | OC-4328

Exact Enumeration of Phylogenetic Networks: The Tree-Child, Reticulation-Visible and Orchard Hierarchy | full_paper_arxiv.tex:1890-1904

\label{conj:orch_asymp} For each fixed $k\geq2$ there exists a constant $C_k>1$ such that \[\frac{|\mathrm{Orch}_{\ell,k}|}{|RV_{\ell,k}|} \;\longrightarrow\; C_k \qquad (\ell\to\infty). \] From the data: $C_2\approx1.148$ (converged within the available range) and $C_3>1$ (the ratio $1.607,1.572,1.531$ is decreasing toward a limit above~1). If Conjecture~\ref{conj:orch_asymp} holds, then $|\mathrm{Orch}_{\ell,k}|\sim C_k|TC_{\ell,k}|$ by Theorem~\ref{thm:universal}, meaning the orchard and RV generating functions share the dominant singularity at $x=\tfrac{1}{2}$ but with amplitudes differing by~C_k.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Josep Batle | 2026-06-23

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000200 | OC-4346

Strong chromatic index of bipartite graphs | BiStrongColoring2.tex:79-85

\label{conj:BQ} For any bipartite \(G \) with partite sets A and B, \(\chi'_s(G) \le \Delta_A \Delta_B, \) where $\Delta_A=\max\{d_G(a):a\in A\}$ and $\Delta_B=\max\{d_G(b):b\in B\}$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Yanli Hao; Tianchi Yang; Xingxing Yu | 2026-06-22

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000201 | OC-4372

Excluding an induced star in dense random graphs | main.tex:577-581

Let H be a critical graph (as defined in \Cref{subsec:col-num}) of coloring number $\tau(H)\geq3$ with no isolated vertices. If $\gamma\in(0,1)$ is fixed and $m\sim\gamma\binom{n}{2}$ then $$\lim_{n\to\infty}\frac{1}{\binom{n}{2}}\log N^\ast_{n,m}(H)=\max_{s,t}\,\Phi_{s,t}(\gamma)\,,$$ where the maximum is over witnessing pairs (s,t) of H.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Sam van der Poel | 2026-06-21

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000202 | OC-4427

Spectral Radius Conditions for 3-Uniform Intersecting Families | Version00616.tex:136-140

[Erd\H{o}s Matching Conjecture \cite{Erd65}] Let $n\geq (k+1)r$. If H be an M_{k+1}-free r-graph on n vertices, then $$|E(H)|\leq \max\{|\mathcal{A}^r_1|,|\mathcal{A}^r_r|\}=\max\left\{\binom{n}{r}-\binom{n-k}{r}, \binom{rk+r-1}{r}\right\},$$ where $\mathcal{A}^r_i=\left\{e\in \binom{[n]}{r}:|e\cap[ik+i-1]\geq i\right\}$ for $1\leq i\leq r$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Lusheng Fang; Guorong Gao; An Chang | 2026-07-09

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by_sa

V3-000203 | OC-4596

Counting even cycles and even paths with bounded circumference | even_cycles_and_even_paths_long_cycles_and_paths.tex:182-187

\label{conj:ZGHLSX} For all integers n,k,L such that $k\ge4$, $L>k$, and n is sufficiently large as a function of k and L, one has \[\ex(n,C_k,C_{\ge L+1})=C_k(H(n,L)). \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Xiamiao Zhao; Yuanpei Wang | 2026-07-05

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000204 | OC-4608

Bergeron's conjecture & a tale of two binomial coefficients | bergeron.tex:150-154

{\it If $1\leq a<b<c<d$ are integers with $ad=bc$, then $$\gauss{b+c}b \ \geq \ \gauss{a+d}a \, ,$$ where the inequality is understood coefficient-wise.}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Tewodros Amdeberhan; Matthias Beck | 2026-07-04

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000205 | OC-4677

Robustness and hyperstability for the Erdős-Gallai theorem | main.tex:642-644

\label{conj: robus erdos sos alt} For every $c>0$ the following holds for all large enough K. Let G be a graph on n vertices with average degree d and let T be a tree on $(1-c)d$ vertices with maximum degree at most Δ. Let $p\geq K \Delta /d$. Then, with probability tending to 1 as $n\rightarrow\infty$, G_p contains a copy of T.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Micha Christoph; Alp Müyesser; Yuval Wigderson | 2026-07-02

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000206 | OC-4808

A Polynomial Improvement of Naslund--Sawin Bound for Sunflower-Free Families Using Triangular Tensors | main.tex:92-98

For $k\ge 3$, let $\mathcal{F}$ be a k-sunflower free family of m-subsets of $[n]$, i.e., each set is of size m. Then \[|\mathcal{F}|\le \mu^m \] for a constant μ depending only on k.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Omran Ahmadi; Hassan Norouzi | 2026-06-29

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000207 | OC-4867

Improved Domination--Packing Bounds in Claw-Free Cubic Graphs and Unit Disk Graphs | main.tex:242-247

\label{conj:cubic-75} For every bridgeless claw-free cubic graph G, $ \gamma(G) \leq \frac{5}{4}\rho(G). $

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Juan Gutiérrez; Kaustav Paul | 2026-06-28

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000208 | OC-579

Maximally probable tree topologies with r-furcation | appendix_paper.tex:470-477

\label{conj:max_r_sim} Consider the set of rooted r-furcating unlabeled topologies with n leaves, permitting simultaneous r-furcations. \begin{enumerate}[label = (\roman*)] \item The unlabeled topology whose labelings have the largest number of tie-permitting labeled histories takes the form in Theorem~\ref{thm:r_furcating_max_shape}. This topology is unique in having the maximal value. \item Further, this same unlabeled topology has the largest number of tie-permitting labeled histories with exactly z events, $\lceil \log_r n \rceil \leq z \leq (n-1)/(r-1)$. This topology is not necessarily unique in having the maximal value. \end{enumerate}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Emily H. Dickey; Noah A. Rosenberg | 2026-02-07

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000209 | OC-714

Uniformity of extremal graph-codes | Extremal-graph-codes.tex:989-991

For every pair d,k of positive integers with $k\meg 2$ and every\, $0<\delta\mik 1$, there exists a positive integer $\mathrm{DPHJ}(d,k,\delta)$ such that if $n\meg \mathrm{DPHJ}(d,k,\delta)$, then any subset of $W(n,d,k)$ with cardinality at least $\delta\, k^{n^d}$ contains a polynomial combinatorial line of\, $W(n,d,k)$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Noé de Rancourt; Pandelis Dodos; Konstantinos Tyros | 2026-02-03

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000210 | OC-864

Adjoints of Morphisms of Neural Codes | BMFandNeuralRings.tex:1118-1120

Let $\bC$ be a binary matrix with k redundant neurons of which $\ell \leq k$ are not simply redundant. Then $\brank(\bC) = \brank(\rho(\bC)) + \ell$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Juliann Geraci; Alexander B. Kunin; Alexandra Seceleanu | 2026-03-11

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000211 | OC-885

Binomial Random Matroids | BinMat.tex:238-240

\label{conj:paving} As $n\rightarrow \infty$, the proportion of matroids on $[n]$ that are paving tends to 1.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Patrick Bennett; Alan Frieze | 2026-03-11

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000212 | OC-946

Patrolling cop vs omniscient robber | arxiv.tex:830-832

\label{th:chordal} Let G be a connected chordal graph and $\rho\ge 2$. Its range $\range{G} \ge \rho$ if and only if there is a triod T with $\ell_3(T) \ge 2\rho$ or a clique-triod G with $\ell_3(G) \ge 2\rho -1$ that is a weak retract of G.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Nina Chiarelli; Paul Dorbec; Miloš Stojaković; Andrej Taranenko | 2026-03-09

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000213 | OC-952

On an infinite sequence of strongly regular digraphs with parameters $(9(2n+3), 3(2n+3), 2n+4, 2n+1, 2n+4)$ | article.tex:392-394

For the constructed sequence of strongly regular digraphs with parameters $(9(2n+3), 3(2n+3), 2n+4, 2n+1, 2n+4)$, the automorphism group is $C_2 \times (C_2^{2n+2} \rtimes C_{2n+3})$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Viktor A. Byzov; Igor A. Pushkarev | 2026-03-08

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000214 | OC-957

On a conjecture concerning the property of chromatic polynomials with negative variable | conj2.tex:107-112

\label{con} Let \(G \) be a graph of order \(n \). Then \[\frac{d^k}{dx^k} (\ln[(-1)^n P(G, x)]) < 0 \] holds for all \(k \geq 2 \) and \(x \in (-\infty, 0) \).

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Yan Yang | 2026-03-08

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000215 | OC-286

A type theory for invertibility in weak $ω$-categories | main.tex:2311-2317

The functor \(\loc\) admits a right adjoint, given by: \begin{align*} \radj \ty \markedomegacat &\to \Mod(\cattinv) \\ (\radj X)(\Gamma) &= \markedomegacat(\loc\minterp{\Gamma},X) \end{align*}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Thibaut Benjamin; Camil Champin; Ioannis Markakis | 2026-02-18

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000216 | OC-4173

A Cohesive ∞-Topos with a Quantum Modality from Finite-Dimensional C^{*}-Algebras | main.tex:406-408

\label{conj:class} Let \(\mathcal{C}\) be a small \(\infty\)-category and \(\mathbf{H}\) a cohesive \(\infty\)-topos. Suppose \(Q\) is an accessible, idempotent, product-preserving comonad on \(\Fun(\mathcal{C},\mathbf{H})\) that commutes with the pointwise cohesive modalities and restricts to the identity on the full subcategory of constant discrete objects. Then there exists a coreflective full subcategory \(\mathcal{D}\hookrightarrow\mathcal{C}\) such that \(Q \cong r^{*}\) for the coreflector \(r\), and the coalgebra category is \(\Fun(\mathcal{D},\mathbf{H})\).

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Joey Woo | 2026-06-01

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000217 | OC-4174

A Cohesive ∞-Topos with a Quantum Modality from Finite-Dimensional C^{*}-Algebras | main.tex:410-412

\label{conj:channels} The centre comonad is insufficient to faithfully embed CPTP maps. There exists a site \(\mathcal{S}\) (e.g., algebras with a completely positive structure, or Hilbert bimodules) and an idempotent strong monoidal comonad \(K\) on \(\mathcal{S}\) whose Kleisli category contains \(\mathbf{CPTP}\) as a full subcategory, with Day convolution modelling the tensor product of channels, and whose coalgebras are the commutative algebras.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Joey Woo | 2026-06-01

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000218 | OC-4175

A Cohesive ∞-Topos with a Quantum Modality from Finite-Dimensional C^{*}-Algebras | main.tex:414-420

\label{conj:derived} There exists an \(\infty\)-category \(\mathcal{C}\) of ``derived finite-dimensional \(C^{*}\)-algebras'' and a derived centre functor \(\mathbb{R}\mathrm{Center}\) that is an idempotent strong monoidal comonad satisfying the Seely isomorphism \[\mathbb{R}\mathrm{Center}(A\otimes B) \;\cong\; \mathbb{R}\mathrm{Center}(A) \otimes^{\mathbb{L}} \mathbb{R}\mathrm{Center}(B), \] yielding a fully non-degenerate cohesive linear \(\infty\)-topos.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Joey Woo | 2026-06-01

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000219 | OC-4176

A Cohesive ∞-Topos with a Quantum Modality from Finite-Dimensional C^{*}-Algebras | main.tex:581-589

\label{conj:channels-refined} There exists a site \(\mathcal{S}\) built from finite-dimensional \(C^{*}\)-algebras (e.g., the category of algebras and CPTP maps, or a suitable category of Hilbert bimodules) and an idempotent strong monoidal comonad \(K\) on \(\mathcal{S}\) such that: \begin{enumerate} \item[(i)] The Kleisli category of the induced comonad on \(\Fun(\mathcal{S}, \mathbf{H}_{\mathrm{sm}})\) (with Day convolution) contains \(\mathbf{CPTP}\) as a full subcategory. \item[(ii)] The tensor product of quantum channels corresponds to the Day convolution. \item[(iii)] The coalgebras of \(K\) are precisely the commutative algebras, so that the classical core remains the topos of discrete classical field theories. \end{enumerate} The centre comonad does not satisfy (ii) because it preserves finite products. A non-product-preserving modality is required.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Joey Woo | 2026-06-01

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000220 | OC-4529

Polynomials interpolated totally positive sequences | Polynomials.interpolated.TP.04.07.26.tex:749-750

For all $k\in \mathbb{N}$ we have $M_{2k} = [-c_{2k}, 1+ c_{2k}], 0 < c_{2k} <2,$ and $c_2 \leq c_4 \leq c_6 \leq \ldots$

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Anna Vishnyakova | 2026-07-07

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000221 | OC-469

The Carathéodory metric on Teichmüller space of genus two surface | Caratheodory_Feb10.tex:153-156

\label{conj:equl} A Teichm\"uller disk is a holomorphic retract if and only if it is generated by a holomorphic quadratic differential all of whose zeros are of even order.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Kejie Lin; Weixu Su | 2026-02-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000222 | OC-1976

Allard Regularity for Abelian Yang--Mills--Higgs Equation | Allard_Abelian_YMH_Arxiv_version.tex:359-369

\label{Gibbon} An entire critical point $\begin{pmatrix}u, A\end{pmatrix}^T$ on $\mathbb{R}^{n+2}$ satisfying \begin{align*} \lim_{R \rightarrow \infty} \frac{1}{\left|B_R^n\right|} \int_{B_R^{n+2}} e_{\e}\begin{pmatrix}u\\ A\end{pmatrix}=2 \pi \end{align*} and, writing any $x \in \mathbb{R}^{n+2}$ as $x=(y, z) \in \mathbb{R}^2 \times \mathbb{R}^n$, also satisfying \begin{align*} \lim_{|z| \rightarrow \infty}|u(y, z)|=1, \quad \text{ uniformly in } z, \end{align*} is necessarily two-dimensional. More precisely, it is the pullback through the projection $\mathbb{R}^{n+2} \rightarrow \mathbb{R}^2$ of the standard solution in $\mathbb{R}^2$ with degree ± 1, up to translation and change of gauge.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Huy The Nguyen; Shengwen Wang | 2026-04-21

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000223 | OC-464

Geometric eigenvalue estimates of Kuttler-Sigillito type on differential forms | main.tex:303-310

Let (M^n,g) be a compact Riemannian manifold, whose sectional curvature K_g satisfies $\kappa_1\leq K_g \leq \kappa_2$. Assume that M is star-shaped with respect to $x_0\in M$ then, for all $k\in\mathbb{N}$, $p\geq 1$, there exists a constant $C_1= C_1(p,n,\kappa_1,\kappa_2)>0$ such that: \begin{equation}\label{eq:th1.4(i,1)} \qquad \lambda_{k,p}\geq \frac{C_1 q_{{m_k},p}}{r_{\max}+h_{\max}} . \end{equation} Here m_k is the multiplicity of $\lambda_{k,p}$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Rodolphe Abou Assali | 2026-02-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000224 | OC-620

Quantitative partitioned index theorem and noncompact band-width | bandwidth_Hochs_Wang.tex:230-235

\label{con band width cpt} Let N be a compact manifold without boundary, of dimension $n-1 \geq 5$. Suppose that N does not admit Riemannian metrics of positive scalar curvature. Consider a Riemannian metric g on $N \times [-1,1]$. Let ℓ be the distance

between $N \times \{-1\}$ and $N \times \{1\}$ with respect to this metric. Then \[\inf_{x\in N \times [-1,1]}\Sc_g(x)\le\frac{4\pi^2(n-1)}{n\ell^2}. \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Peter Hochs; Jinmin Wang | 2026-02-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000225 | OC-171

Structure and Spectrum of Nonergodic Nilsystems | Structure_and_Spectrum_of_Nonergodic_Nilsystems.tex:286-292

\label{Conj} Let $(X=G/\Gamma,\mu,T_1,\ldots,T_d)$ be an ergodic $\Z^d$-nilsystem. Then the Koopman representation of $(T_1,\ldots,T_d)$ on $L^2(X,\mu)$ decomposes as \[L^2(X,\mu)=E_{\mathrm{ds}}\oplus E_{\mathrm{Leb}}, \] where $E_{\mathrm{ds}}\cong L^2(G/G_2\Gamma)$ has discrete spectrum and E_{Leb}, if nontrivial, splits into a countable orthogonal sum of invariant subspaces, each having maximal spectral type absolutely continuous with respect to the Haar measure of a nontrivial subtorus of $\bS^d$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Felipe Hernández | 2026-02-24

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000226 | OC-4123

A Graph-Based Method for Invariant Densities of Multidimensional Continued Fractions | main.tex:1531-1533

Assume that $\Leb(D_{\text{AL}}) > 0$, where D_{AL} is the invariant domain described in~(\ref{eq:0}). Then there exists a solution $\big(\Omega_{v}\big)_{v\in V}$ to the system of equations~(\ref{eq:4}) such that every matrix Ω_v has full rank.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | David Siukaev | 2026-06-02

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000227 | OC-4124

A Graph-Based Method for Invariant Densities of Multidimensional Continued Fractions | main.tex:1955-1966

The invariant domain for the Triangle algorithm in dimension $d=3$ is nonconvex and fractal. For the associated win-lose induction, the invariant domain is contained in $$\bigsqcup_{\sigma\in S_4,i}\{v_{\sigma,\sigma_i}\}\times\Big(\Omega^1_{\sigma,\sigma_i}\mathbb{R}^4_+\sqcup \Omega^2_{\sigma,\sigma_i}\mathbb{R}^4_+\sqcup \Omega^3_{\sigma,\sigma_i}\mathbb{R}^4_+\Big),$$ where \begin{center} $\displaystyle \Omega^1_{\sigma,\sigma_i} = C_{\sigma'}\cdot\begin{bmatrix} 1 & 0 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 2 & 1 \\ 1 & 1 & 3 & 2 \end{bmatrix} \cdot C^{-1}_{\sigma'}, \quad\Omega^2_{\sigma,\sigma_i} = C_{\sigma'}\cdot\begin{bmatrix} 1 & 0 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 2 & 2 \\ 1 & 1 & 2 & 3 \end{bmatrix} \cdot C^{-1}_{\sigma'},$ \\ $ $ \\ $\displaystyle\Omega^3_{\sigma,\sigma_i} = C_{\sigma'}\cdot\begin{bmatrix} 1 & 0 & 0 & 1 \\ 1 & 1 & 0 & 2 \\ 2 & 1 & 1 & 3 \\ 2 & 1 & 1 & 4 \end{bmatrix} \cdot C^{-1}_{\sigma'},$ \\ $ $ \\ \end{center} $\sigma' = \sigma\cdot(\sigma_{i-d}\ldots\sigma_{i-1}\sigma_{i})$ and $C_{\sigma'}$ is the matrix for the permutation σ'.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | David Siukaev | 2026-06-02

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000228 | OC-536

Dynamical Mordell-Lang conjecture for split self-maps of affine curve times projective curve | A1P1.tex:122-124

Let $f:X\to X$ be an endomorphism of a quasi-projective variety over a field K of characteristic 0, and V be a closed subvariety of X. Then for every $x\in X(K)$, the return set $\{ n\in \mathbb{N}|\ f^n(x) \in V(K)\}$ is a finite union of arithmetic progressions.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Junyi Xie; She Yang; Aoyang Zheng | 2026-02-09

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000229 | OC-1412

Non-existence of abelian maximal subgroups in cyclic division algebras | hvkhanh_AbelianMaximalSubgroups.tex:43-45

If D is a non-commutative division ring, then D^* contains no nilpotent maximal subgroups.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Huynh Viet Khanh | 2026-03-30

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000230 | OC-2656

Aldous-type Spectral Gaps in Generalized Symmetric Groups | main.tex:165-170

\label{conj:Caputo} For all hypergraphs $\Gamma=([n],c)$, \[\ls(\Gamma,\regSn) = \ls(\Gamma,\std) \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Niv Levhari; Doron Puder | 2026-05-21

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000231 | OC-3548

Lacunary hyperbolic groups with fast injectivity radius growth and enough loxodromic elements are selfless | main_arxiv.tex:642-644

There exists a finitely generated group whose reduced $\Cstar$-algebra is $\Cstar$-selfless but not $\Cstar$-exact.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Goulnara Arzhantseva; Martin Finn-Sell | 2026-06-18

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000232 | OC-3599

Surjectivity of Engel Words on $\mathrm{SL}_2(\mathcal{O})$ and $\mathrm{PSL}_2(\mathcal{O}_2)$ | main.tex:145-147

Every element of $\mathrm{SL}_n(\mathbb{Z}_p)$ where p is a prime and $n\geq 2$ ($p>3$ when $n=2$) is a commutator.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Ayon Roy; Anupam Singh | 2026-06-17

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000233 | OC-2657

Closed geodesics in short intervals for random hyperbolic surfaces | geodesics_vs_primes_short_intervals_v6.tex:207-213

\label{GM conjecture} For every fixed $0<\delta<1$, \begin{equation*} \Var(\psi)(X;H):=\frac 1X\int_1^X \left|\psi(x;H)- H\right|^2 dx \sim H \log \frac XH \end{equation*} holds uniformly for $1\leq H<X^{1-\delta}$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Zeev Rudnick | 2026-05-21

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000234 | OC-4395

Isometric free finite group actions on non-positively curved 3-manifolds | main.tex:355-357

\label{conj:npc_dimension_3} Let M be a closed orientable 3-manifold admitting an NPC metric, and let G be a finite group acting on M by orientation-preserving diffeomorphisms. Then M admits a G-invariant NPC metric.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Zhengyu Zou | 2026-06-20

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000235 | OC-4881

A solution to Ditor's problem | Ditor_s_Problem_arXiv_v.1.tex:945-947

CH implies the existence of a 4-ladder of cardinality $\aleph_2$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Lorenzo Notaro | 2026-06-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000236 | OC-742

Decidability of Interpretability | main_LICS_arxiv_version.tex:152-154

\label{conj:bp} Let $\bC$ be a first-order reduct of a finitely bounded homogeneous structure $\bA$. Then $\CSP(\bC)$ is either in P or NP-complete.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Roman Feller; Michael Pinsker | 2026-02-02

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000237 | OC-958

The reals as a subset of an ultraproduct of finite fields | main.tex:691-693

External partition cuts are never \(\delta\).

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Roee Sinai | 2026-03-08

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by_sa

V3-000238 | OC-1539

A characterization of the sphere in terms of the stereographic projection | stereographic_projection.tex:183-191

\label{conjeeli} Let $K\subset \mathbb{E}^3$ be a convex body and let $N \in K$. If, for all plane Γ such that $\Gamma\cap \inte K\not= \emptyset$ and $N\notin \Gamma$, there exists a rotation $R^{\Gamma} $ with property that \begin{eqnarray}\label{camita} R^{\Gamma} (C_N(K_{\Gamma}))=C_N(K_{\Gamma}), \end{eqnarray} where $K_{\Gamma}:=\Gamma\cap K$, then K is an ellipsoid of revolution.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Efrén Morales-Amaya | 2026-04-08

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000239 | OC-1627

A note on the spectral distribution of non-Hermitian block matrices with Toeplitz blocks | Mean_Block_GLT.tex:824-850

\label{conj_final} Let \begin{equation*} \mathcal{A}_n = \begin{bmatrix} T_n(f_{1,1}) & T_{n}(f_{1,2}) & \dots & T_n(f_{1,k}) \\ T_n(f_{2,1}) & T_n(f_{2,2}) & \ddots & \vdots \\ \vdots & \ddots & \ddots & T_n(f_{k-1 , k}) \\ T_n(f_{k,1}) & \dots & T_n(f_{k, k-1}) & T_n(f_{k, k}) \end{bmatrix}, \end{equation*} where, for every $i,j=1,\dots, k,$ $f_{i,j} \in L^{\infty}([-\pi,\pi])$ and real-valued almost everywhere with $T_n(f_{i,j}) $ Hermitian positive definite whenever $i \neq j$, or $f_{i,j}=0$ almost everywhere. Assume that, for every cycle $i_1,\dots,i_k,i_{k+1}$, we have \begin{equation*} \prod_{l=1}^{k} f_{i_l,i_{l+1}} (\theta) = \prod_{l=1}^{k} f_{i_{l+1},i_l} (\theta) \end{equation*} for a.e. $\theta \in [-\pi,\pi]$, where $i_{k+1}=i_1$. Then, \begin{equation*} \{\mathcal{A}_n\}_n \sim_{\lambda} \begin{bmatrix} f_{1,1} & f_{1,2} & \dots & f_{1,k}\\ f_{2,1} & f_{2,2} & \ddots & \vdots \\ \vdots & \ddots & \ddots & f_{k-1 , k} \\ f_{k,1} & \dots & f_{k, k-1} & f_{k, k} \end{bmatrix}. \end{equation*}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Andrea Adriani; Giacomo Tento | 2026-04-04

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000240 | OC-190

The largest 5th pivot may be the root of a 61st degree polynomial | main.tex:655-658

\label{g5conjecture} The largest possible growth factor for a 5×5 real matrix under complete pivoting is exactly the lower bound proven in this paper, i.e., the unique real root g in the interval $[4,5]$ of the 61^{st} degree polynomial $P_5(g)$ in Equation (\ref{eq:61}).

自动抽取的猜想块 | source_machine_label_real_open_conjecture | James Chen; Alan Edelman; John Urschel | 2026-02-23

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000241 | OC-2132

Flexible GMRES converges in two phases | main.tex:742-744

Let $0 < \mu \leq \frac{1}{2}$. For any right-hand side vector $b \in \C^N \setminus \{0\}$ and any choice of matrix preconditioners $(M_j)_{1\leq j < N}$, there exists an invertible matrix $A \in \C^{N \times N}$ such that, for all $1 \leq j < N$, it holds that $\|r_j^{\mathrm{P}}\| \leq \mu$ and the bound in Theorem~\ref{thm:explicit_bound} holds as an equality.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Stefan Güttel; Lauri Nyman | 2026-04-30

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000242 | OC-1266

Towards the p-adic Hodge parameters in semistable representations of $\mathrm{GL}_n(\mathrm{Q}_p)$ | HodgeParameterPrism.tex:2072-2077

\label{descibeimagetDGL2} Let $u\in\sW_n^{S_0}$.\;Let $\iota_{S_0(u)}:\overline{\ext}^{1}_{S_0(u)}(\Dpik,\Dpik)\hookrightarrow \overline{\ext}^{1}(\Dpik,\Dpik)$ be the natural inclusion.\;Recall the surjection $g^+_{\Dpik}:\bigoplus_{u\in\sW_n^{S_0}}\overline{\ext}^{1}_{S_0(u)}(\Dpik,\Dpik)\rightarrow \overline{\ext}^{1}(\Dpik,\Dpik)$ in Corollary \ref{surforgD}.\; \begin{itemize} \item[(1)]Under the splitting in Proposition \ref{splitingforext1gps},\;$t^+_{D}\circ\gamma^+_{\Dpik}|_{\overline{\ext}^{1}_{S_0(u)}(\Dpik,\Dpik)}$ is equal to $f_{\Dpik}\circ\iota_{S_0(u)}$.\;Moreover,\;$g^+_{\Dpik}$ factors through $\gamma_{\Dpik}$,\;and $g^{+}_{\Dpik}=t^+_{D}\circ\gamma^+_{\Dpik}$.\; \item[(2)] $\ker(t^+_{D})$ determines $\fil_{H}^{\bullet}(D)$.\; \end{itemize}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Yiqin He | 2026-04-02

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000243 | OC-144

Goldfeld conjecture for non-hyperelliptic direction | Goldfeld_another.tex:281-283

The average analytic rank of $\lcrc{C_d}$ for positive square-free integers d is $\frac{1}{4}$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Keunyoung Jeong; Junyeong Park | 2026-02-25

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000244 | OC-1456

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:2217-2221

\label{c:gr} The object $$\Tr(\Frob,\IndCoh_\bO(\LS_\cG^{\on{restr}}))\in \Vect_{\ol\BQ_\ell}$$ is concentrated in cohomological degree 0.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000245 | OC-1457

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:2838-2840

\label{c:filtr rat} The filtration \eqref{e:filtr} is defined over $\BQ$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000246 | OC-1458

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:2864-2869

\label{c:temp rat} The subspace $$\sFunct_c(\Bun_G(\BF_q),\ol\BQ_\ell)_{\on{temp}} \overset{\text{\corref{c:temp=nondeg}}}= \sFunct_c(\Bun_G(\BF_q),\ol\BQ_\ell)_{\on{non-degen}}$$ of $\sFunct_c(\Bun_G(\BF_q),\ol\BQ_\ell)$ is defined over $\BQ$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000247 | OC-1459

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:2887-2891

\label{c:exc rat prel} The algebra $\Gamma({}'\!\LS^{\on{arithm}}_\cG,\CO_{'\!\LS^{\on{arithm}}_\cG})$ possesses a rational structure, uniquely characterized by the property that the action of a rational element sends $\poinc^{\on{Vac}}_!$ to a rational element of $\sFunct_c(\Bun_G(\BF_q),\ol\BQ_\ell)$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000248 | OC-1460

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:2903-2905

\label{c:exc rat} The action of $\on{Exc}(X,\cG)$ on $\sFunct_c(\Bun_G(\BF_q),\ol\BQ_\ell)$ is compatible with the rational structures.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000249 | OC-1461

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3023-3032

\label{c:Arthur} There exists an orthogonal decomposition \begin{equation} \label{e:Arthur} L^2(\Bun_G(\BF_q))\simeq \underset{\bO}\oplus\, L^2(\Bun_G(\BF_q))_\bO, \end{equation} uniquely characterized by the property that for every $x\in |X|$, the spectrum of $\Sph(G)_{x,\BC}$ acting on $L^2(\Bun_G(\BF_q))$ is contained in the subset $$\sR_{q_x,\bO,\BC}\subset (\cG/\!/\on{Ad}(\cG))(\BC),$$ where q_x is the cardinality of the residue field at x.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000250 | OC-1462

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3097-3101

\label{c:Ram-Pet} The inclusion $$\underset{\bO}\oplus\, \sFunct_{\on{cusp}}(\Bun_G(\BF_q),\BC)_\bO\to \sFunct_{\on{cusp}}(\Bun_G(\BF_q),\BC)$$ is an isomorphism.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000251 | OC-1463

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3120-3123

\label{c:Arth alg} The closure of $\sFunct_c(\Bun_G(\BF_q),\BC)_Y$ in $L^2(\Bun_G(\BF_q))$ equals $$\underset{\bO\subset Y}\oplus\, L^2(\Bun_G(\BF_q))_\bO.$$

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000252 | OC-1464

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3127-3145

\hfill \label{c:Arth alg prime} \smallskip \noindent{\em(a)} The subspace $\sFunct_c(\Bun_G(\BF_q),\BC)_Y\subset \sFunct_c(\Bun_G(\BF_q),\BC)$ equals the intersection $$ \sFunct_c(\Bun_G(\BF_q),\BC) \cap \left(\underset{\bO\subset Y}\oplus\, L^2(\Bun_G(\BF_q))_\bO\right),$$ viewed as subsets of $L^2(\Bun_G(\BF_q))$. \smallskip \noindent{\em(b)} For every nilpotent orbit $\bO$, the map $$\sFunct_c(\Bun_G(\BF_q),\BC)_\bO=\sFunct_c(\Bun_G(\BF_q),\BC)_{\ol\bO}/\sFunct_c(\Bun_G(\BF_q),\BC)_{\ol\bO-\bO}\to L^2(\Bun_G(\BF_q))_\bO$$ has a dense image.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000253 | OC-1465

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3190-3192

\label{c:alg discr} The map \eqref{e:Arth alg} is an isomorphism.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000254 | OC-1466

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3201-3206

\label{c:filtr and Arthur and cusp C} Let $Y\subset \Nilp$ be a closed Ad-invariant subset. Then the subspace $$\sFunct_{\on{cusp}}(\Bun_G(\BF_q),\BC)\cap \sFunct_c(\Bun_G(\BF_q),\BC)_Y\subset \sFunct_{\on{cusp}}(\Bun_G(\BF_q),\BC)$$ equals $$\underset{\bO\subset Y}\oplus\, \sFunct_{\on{cusp}}(\Bun_G(\BF_q),\BC)_\bO,$$ where $\sFunct_{\on{cusp}}(\Bun_G(\BF_q),\BC)_\bO$ is as in \secref{sss:cusp O}.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000255 | OC-1467

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3325-3328

\label{c:Ramanujan Q} The inclusion $$\underset{\bO}\oplus\, \sFunct_{\on{cusp}}(\Bun_G(\BF_q),\BQ)_\bO\to \sFunct_{\on{cusp}}(\Bun_G(\BF_q),\BQ)$$ is an isomorphism.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000256 | OC-1468

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3334-3338

\label{c:Ramanujan Q ell} The inclusion $$\underset{\bO}\oplus\, \sFunct_{\on{cusp}}(\Bun_G(\BF_q),\ol\BQ_\ell)_\bO\to \sFunct_{\on{cusp}}(\Bun_G(\BF_q),\ol\BQ_\ell)$$ is an isomorphism.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000257 | OC-1469

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3360-3364

\label{c:filtr and Arthur and cusp} Let $Y\subset \Nilp$ a closed Ad-invariant subset. Then the intersection $$\sFunct_{\on{cusp}}(\Bun_G(\BF_q),\BQ)\cap \sFunct_c(\Bun_G(\BF_q),\BQ)_Y\subset \sFunct_{\on{cusp}}(\Bun_G(\BF_q),\BQ)$$ equals $$\underset{\bO\subset Y}\oplus\, \sFunct_{\on{cusp}}(\Bun_G(\BF_q),\BQ)_\bO.$$

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000258 | OC-1470

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3371-3375

\label{c:filtr and Arthur and cusp ell} $$\sFunct_{\on{cusp}}(\Bun_G(\BF_q),\ol\BQ_\ell)\cap \sFunct_c(\Bun_G(\BF_q),\ol\BQ_\ell)_Y\subset \sFunct_{\on{cusp}}(\Bun_G(\BF_q),\ol\BQ_\ell)$$ equals $$\underset{\bO\subset Y}\oplus\, \sFunct_{\on{cusp}}(\Bun_G(\BF_q),\ol\BQ_\ell)_\bO.$$

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000259 | OC-1471

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3381-3385

\label{c:mock Weil} The inclusion $$\sFunct_{\on{cusp}}(\Bun_G(\BF_q),\BQ)^{\on{Weil}}_\bO \subset \sFunct_{\on{cusp}}(\Bun_G(\BF_q),\BQ)_\bO$$ is an isomorphism.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000260 | OC-1472

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3391-3395

\label{c:mock Weil Q-ell} The inclusion $$\sFunct_{\on{cusp}}(\Bun_G(\BF_q),\ol\BQ_\ell)^{\on{Weil}}_\bO \subset \sFunct_{\on{cusp}}(\Bun_G(\BF_q),\ol\BQ_\ell)_\bO$$ is an isomorphism.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000261 | OC-1473

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3420-3423

\label{c:support O discr C} For every $x\in X$, the support of $\sFunct_{c,\on{discr}}(\Bun_G(\BF_q),\BC)_\bO$ in $(\cG/\!/\on{Ad}(\cG))(\BC)$ is contained in $\sR_{q_x,\bO,\BC}$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000262 | OC-1474

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3427-3431

\label{c:support O discr Q} For every $x\in X$, the support of $\sFunct_{c,\on{discr}}(\Bun_G(\BF_q),\BQ)_\bO$ in $$\on{Specm}(\Sph(G)_{x,\BQ})\simeq \on{Specm}(\CO_{\cG/\!/\on{Ad}(\cG),\BQ})$$ is contained in $\sR_{q_x,\bO,\BQ}$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000263 | OC-1475

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3454-3458

\label{c:support O discr Q ell} For every $x\in X$, the support of $\sFunct_{c,\on{discr}}(\Bun_G(\BF_q),\ol\BQ_\ell)_\bO$ in $$\on{Specm}(\Sph(G)_{x,\ol\BQ_\ell})\simeq (\cG/\!/\on{Ad}(\cG))(\ol\BQ_\ell)$$ is contained in $$\sR_{q_x,\bO,\BQ}\subset \on{Specm}(\CO_{\cG/\!/\on{Ad}(\cG),\BQ}) \subset (\cG/\!/\on{Ad}(\cG))(\ol\BQ_\ell).$$

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000264 | OC-1476

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3601-3615

\label{c:locally finite O} \hfill \medskip \noindent{\em(a)} The support of $\sFunct_{c,\on{discr}}(\Bun_G(\BF_q),\ol\BQ_\ell)_\bO$ in $\LS^{\on{arithm,coarse}}_\cG(\ol\BQ_\ell)$ is contained in the (finite) set consisting of Langlands parameters associated to \emph{elliptic} Arthur parameters whose SL_2 component corresponds to $\bO$. \medskip \noindent{\em(b)} For a given \emph{elliptic} Arthur parameters, the direct summand of $\LS^{\on{arithm,coarse}}_\cG(\ol\BQ_\ell)$ set-theoretically supported at it is at most one-dimensional.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000265 | OC-1477

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3633-3645

\label{c:cusp O} Assume that \conjref{c:locally finite O}(a) holds. Then: \hfill \smallskip \noindent{\em(a)} Conjectures \ref{c:Ramanujan Q ell} and \ref{c:filtr and Arthur and cusp ell} hold. \smallskip \noindent{\em(b)} The support of $\sFunct_{\on{cusp}}(\Bun_G(\BF_q),\ol\BQ_\ell)_\bO$ in $\LS^{\on{arithm,coarse}}_\cG(\ol\BQ_\ell)$ is contained in the (finite) set consisting of Langlands parameters associated to \emph{elliptic} Arthur parameters whose SL_2 component corresponds to $\bO$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000266 | OC-1478

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3666-3669

\label{c:motivic O Q-ell} The support of $\sFunct_{c,\on{discr}}(\Bun_G(\BF_q),\ol\BQ_\ell)_\bO$ in $\LS^{\on{arithm,coarse}}_\cG(\ol\BQ_\ell)$ is contained in $\LS^{\on{mot,coarse}}_\cG(\ol\BQ_\ell)$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000267 | OC-1479

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3676-3679

\label{c:cusp in mot} The support of $\sFunct_{\on{cusp}}(\Bun_G(\BF_q),\ol\BQ_\ell)$ in $\LS^{\on{arithm,coarse}}_\cG(\ol\BQ_\ell)$ is contained in $\LS^{\on{mot,coarse}}_\cG(\ol\BQ_\ell)$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000268 | OC-1480

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3695-3699

\label{c:motivic Q} The support of $\sFunct_{\on{cusp}}(\Bun_G(\BF_q),\ol\BQ)$ in $\LS^{\on{arithm,coarse}}_\cG(\ol\BQ)$ is contained in $$\LS^{\on{mot,coarse}}_\cG(\ol\BQ)\subset \LS^{\on{arithm,coarse}}_\cG(\ol\BQ).$$

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000269 | OC-1481

Tempered vs generic automorphic functions and the canonical filtration on automorphic functions | text.tex:3709-3728

\hfill \smallskip \noindent{\em(a)} Assume that \conjref{c:locally finite O}(a) holds. Then the space $L^2_{\on{discr}}(\Bun_G(\BF_q))$ splits as a direct sum $$\underset{\sigma\in \LS^{\on{mot,coarse}}_\cG(\BC)}\oplus\, L^2_{\on{discr}}(\Bun_G(\BF_q))_\sigma,$$ where for every $x\in X$, the support of $L^2_{\on{discr}}(\Bun_G(\BF_q))_\sigma$ in $$\on{Specm}(\Sph(G)_{x,\BC})\simeq (\cG/\!/\on{Ad}(\cG))(\BC)$$ is a single point equal to the image of σ under $$\LS^{\on{mot,coarse}}_\cG(\BC)\to \LS^{\on{arithm,coarse}}_\cG(\BC) \to (\cG/\!/\on{Ad}(\cG))(\BC),$$ where the last arrow corresponds to the restriction along $x\to X$. \smallskip \noindent{\em(b)} Assume additionally that \conjref{c:locally finite O}(b) holds. Then each $L^2_{\on{discr}}(\Bun_G(\BF_q))_\sigma$ is at most one-dimensional.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Dennis Gaitsgory; Vincent Lafforgue; Sam Raskin | 2026-03-27

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000270 | OC-1568

There are infinitely many Hilbert cubes of dimension 3 in the set of squares | hilbert_cubes_BEU_MathComp_submission.tex:1202-1204

For each $n\in\N_{+}$, there is a Hilbert cube $H(a_{0}; a_{1}, a_{2}, a_{3})\subset\cal{S}$ of dimension 3 with $a_{0}=n^2$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Andrew Bremner; Christian Elsholtz; Maciej Ulas | 2026-04-07

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000271 | OC-1638

Series for $1/π$ arising from Cauchy product | main.tex:43-46

Let $p>3$ be a prime. Then, \[\sum_{n=0}^{p-1} \frac{3n-1}{2^n}\sum_{k=0}^n\binom{-1/3}{k}\binom{-2/3}{n-k}\binom{-1/6}{k}\binom{-5/6}{n-k}\equiv-p\legendre{-6}{p}\pmod{p^2}.\]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Roman Le Lan | 2026-04-02

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000272 | OC-2608

On the Harris-Viehmann conjecture for Hodge-Newton reducible local Shimura data of abelian type | arxiv.tex:475-485

(Harris--Viehmann conjecture) \cite[Conjecture 8.5]{RapoportViehmann}\label{cnj: HV OG} Fix a prime $\ell \neq p$. Let $(G, [b], \{\mu\})$ be a non-basic unramified local Shimura datum. Let L be a Levi subgroup of G such that $b \in L(\breve{\Q}_p)$ for some $b \in [b]$, and let P be a standard parabolic containing L. Let J_b be an inner form of a Levi subgroup contained in L. Let $I_{b,\{\mu\}, L}$ be the set of $L(\breve{\Z}_p)$-conjugacy classes of L with cocharacter representative μ' such that: \begin{enumerate} \item $\mu' \in \{\mu\}$. \item $[b] \cap L(\breve{\Z}_p)\mu'(p)L(\breve{\Z}_p)$ is non-empty. \end{enumerate} Then for any admissible $\overline{\Q}_\ell$-representation ρ of $J_b(\Q_p)$, the virtual representation of $G(\Q_p)\x \mathcal{W}_E$ is expressed by the following parabolic inductive formula: \begin{align*} \mathrm{H}^\bullet(\RZ^\infty_{G, b})_\rho = \mathrm{Ind}_{P(\Q_p)}^{G(\Q_p)}\Big(\sum_{\{\mu'\}\in I_{b,\{\mu\}, L}}\mathrm{H}^\bullet(\RZ^\infty_{L, b})_\rho \Big). \end{align*}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Sandra Nair; Xinyu Zhou | 2026-05-22

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000273 | OC-2623

Distributions of Iwasawa $λ$-invariants of $\mathbf{Z}_p$-towers over supersingular isogeny graphs | main.tex:1104-1106

For an arbitrary non-negative integer \(n\), there exists a prime number \(r\) such that there exists a subset \(W\subset \{1,\dotsc,s\}\) that gives an equality \(n = \sum_{k\smallin W}|O(k)|\), where \(O(1),\dotsc,O(s)\) are the newform orbits of \(S_2\big(\Gamma_0(r)\big)\).

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Taiga Adachi; Kosuke Mizuno; Ryosuke Murooka; Sohei Tateno | 2026-05-22

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000274 | OC-2758

On optimal p-adic uniformization of unitary Shimura curves | KRZII-server.tex:1007-1013

The isomorphism in Theorem \ref{unifor-ab-intro} is the base change by $\Spf O_{E(\varphi_0)_\nu}\to \Spf O_{K_{v_{0}}}$ of an isomorphism \begin{equation*} (\CA_{\bK_{\U}})^\wedge \simeq {J}_{\U}(\BQ)\bs\big[\big(\widehat{\Omega}_{F_{v_{0}}}\times_{\Spf O_{F_{v_{0}}}} \Spf O_{{K_{v_0}}}\big) \times \U(\BA_f)/\bK_{\U}\big] . \end{equation*}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Michael Rapoport; Haining Wang | 2026-05-19

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000275 | OC-2765

Fibonacci Numbers and Vieta Jumping for a Rational Diophantine Equation | FibNumbVietaJumpRatDiophEq_MNS_v10.tex:939-955

Let $(a,\,b)$ be a positive integer solution of \begin{equation*} \frac{a+2}{b}\,+\,\frac{b+2}{a}\ = \ k. \end{equation*} Then \begin{equation*} \frac{2(a+b)}{\gcd(a,b)^2} \,\in\,\left\{ 1 , \; 2 , \; 3 , \; 4 , \; 5 , \; 8 \right\}. \end{equation*} Moreover, \begin{itemize}\itemsep0.5em \item if $k \ = \ 3$, the values are 1 and 5; \item if $k \ = \ 4$, the values are 2 and 3; \item if $k \ = \ 6$, the values are 4 and 8. \end{itemize}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Steven J. Miller; Dimitrios Nikolakopoulos; Anitha Srinivasan | 2026-05-18

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000276 | OC-2969

The variation of zeros of the Miller basis | miller-cj.tex:199-203

(Conjecture \ref{conj}) \label{cl:conj} For each $\delta>0$, the zeros of $g_{k,m}$ asymptotically approach the curve $\mathcal{C}_\delta$, defined as the upper hull of the union of $\mathcal{A}$ and $\mathcal{S}_\delta$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Liubomir Chiriac; Andrei Jorza | 2026-05-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000277 | OC-2970

The variation of zeros of the Miller basis | miller-cj.tex:1069-1195

\label{conj} For each $\delta>0$, the zeros of $g_{k,m}$ asymptotically approach the curve $\mathcal{C}_\delta$, defined as the upper hull of the union of $\mathcal{A}$ and $\mathcal{S}_\delta$. \begin{figure}[H] \centering \begin{subfigure}[b]{0.3\textwidth} \centering \includegraphics[width=\textwidth]{vm_12k_60.png} \subcaption{$\delta=0.6, k=12000$} \end{subfigure} \hfill \begin{subfigure}[b]{0.3\textwidth} \centering \includegraphics[width=\textwidth]{vm_12k_75.png} \subcaption{$\delta=0.75, k=12000$} \end{subfigure} \hfill \begin{subfigure}[b]{0.3\textwidth} \centering

\includegraphics[width=\textwidth]{vm_24k_85.png} \subcaption{$\delta=0.85, k=24000$} \end{subfigure} \vspace{0.5cm} \begin{subfigure}[b]{0.3\textwidth} \centering \includegraphics[width=\textwidth]{vm_36k_94.png} \subcaption{$\delta=0.94, k=36000$} \end{subfigure} \hfill \begin{subfigure}[b]{0.3\textwidth} \centering \includegraphics[width=\textwidth]{vm_120k_96.png} \subcaption{$\delta=0.96,k=120000$} \end{subfigure} \hfill \begin{subfigure}[b]{0.3\textwidth} \centering \includegraphics[width=\textwidth]{vm_60k_98.png} \subcaption{$\delta=0.98,k=240000$} \end{subfigure} \caption{Roots and $\mathcal{S}_\delta$ for several values of $\delta\in (0,1)$} \end{figure} \begin{figure}[H] \centering \begin{subfigure}[b]{0.3\textwidth} \centering \includegraphics[width=\textwidth]{vw_12k_1.2.png} \subcaption{$\delta=1.2, k=-12000$} \end{subfigure} \hfill \begin{subfigure}[b]{0.3\textwidth} \centering \includegraphics[width=\textwidth]{vw_24k_1.15.png} \subcaption{$\delta=1.15, k=-24000$} \end{subfigure} \hfill \begin{subfigure}[b]{0.3\textwidth} \centering \includegraphics[width=\textwidth]{vw_24k_1.13.png} \subcaption{$\delta=1.13, k=-24000$} \end{subfigure} \vspace{0.5cm} \begin{subfigure}[b]{0.3\textwidth} \centering \includegraphics[width=\textwidth]{vw_36k_1.1.png} \subcaption{$\delta=1.1, k=-36000$} \end{subfigure} \hfill \begin{subfigure}[b]{0.3\textwidth} \centering \includegraphics[width=\textwidth]{vw_60k_1.06.png} \subcaption{$\delta=1.06, k=-60000$} \end{subfigure} \hfill \begin{subfigure}[b]{0.3\textwidth} \centering \includegraphics[width=\textwidth]{vw_120k_1.03.png} \subcaption{ for $\delta=1.03, k=-120000$} \end{subfigure} \caption{Roots and $\mathcal{S}_\delta$ for several values of $\delta\in (1,\infty)$} \end{figure}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Liubomir Chiriac; Andrei Jorza | 2026-05-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000278 | OC-3869

Multivariate Period Rings | main.tex:1052-1054

\label{conj} $\bD_B(V)$ is finitely presented.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Rohit Pokhrel | 2026-06-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000279 | OC-3990

Uncertainty Principles for the Number Theoretic Transform | main.tex:587-593

\label{conj:descartes} There exist constants $\eps,\delta<1$ such that the following holds. Let p,q be sufficiently large primes with $p\equiv1\pmod q$, let $\gamma\in\F_p^\ast$ have order q, and let $G\le\F_p^\ast$ be the subgroup of order q. If $k\le q^\delta$, if $\alpha_1,\ldots,\alpha_k\in\Z_q$ are distinct, and if $\beta_1,\ldots,\beta_k\in\F_p$, then \[A(z):=\sum_{j=1}^k \beta_j z^{\alpha_j} \] has at most $\eps q$ roots in G, unless all β_j are zero.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Giulio Malavolta; Alon Rosen | 2026-06-07

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000280 | OC-4101

Representations of positive integers by three almost-prime squares | main.tex:190-200

Let n be a positive integer. If $ n\equiv10\pmod{24} $, then n can be expressed as \[n = x^{2} + y^{2} + (2^{a}3^{b})^{2}, \] where x,y,a,b are non-negative integers and $b>0$. If $ n\equiv5\pmod{12} $, then n can be expressed as \[n = x^{2} + y^{2} + (2^{a}5^{b})^{2}, \] where x,y,a,b are non-negative integers and $a>0$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Yue-Feng She; Yu-Chen Sun; Guang-Liang Zhou | 2026-06-03

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000281 | OC-4102

Representations of positive integers by three almost-prime squares | main.tex:203-209

Every integer $n \geq 2$ can be expressed as \[n = x^{2} + y^{2} + (2^{a}3^{b})^{2} + (2^{c}5^{d})^{2}, \] where \(x,y,a,b,c,d\) are non-negative integers.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Yue-Feng She; Yu-Chen Sun; Guang-Liang Zhou | 2026-06-03

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000282 | OC-4147

On the Argument of the Lerch, Chowla--Selberg Formula and CM Values of $η(τ)$ | etacm2.tex:412-433

Let $\tau=(-b+\sqrt{D})/(2a)$ with $D=b^2-4ac<0$, where as usual $a>0$ and $\gcd(a,b,c)=1$, and write $D=D_0f^2$ with D_0 fundamental. As above, set $$\al(\tau)=\dfrac{\eta(\tau)^4}{a\cdot c(D_0,f)\CS(D_0)/(2\pi D)}$$ and let $\be(\tau)$ as in the above definition. \begin{enumerate} \item The numbers $a\cdot\al(\tau)$ and $a/\al(\tau)$ are algebraic \emph{integers}. \item The norm of $\al(\tau)$ as an algebraic number (i.e., the product of its algebraic conjugates) is equal to ±1 (since it is usually not an algebraic integer, this of course does not mean that it is a unit), and in fact perhaps always equal to 1. \item Write $h(D)=2^vm$ with m odd. The minimal polynomial of $a\cdot\be(\tau)$ is of the form $B(x^{jm})$ with $j=1$ if $v=0$, and $j\mid3\cdot2^{v-1}$ if $v\ge1$, and where the degree of B is equal to $h(D)$ if $a=1$ and to $h(D)$ or $2h(D)$ if $a>1$. \item The number $\al(\tau)^{\lcm(6,h(D))}$ belongs to the ring class field of the quadratic order of discriminant D, of degree $2h(D)$, and when $b=0$ or $a=b$ it even belongs to its subfield of degree $h(D)$ fixed by the automorphism $\sqrt{D}\mapsto-\sqrt{D}$. \end{enumerate}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Henri Cohen | 2026-06-01

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000283 | OC-4156

Explicit generators of the space of modular forms | main.tex:1015-1017

{evenperiods}}]\label{conj:even} Let $k_1>k_2>\cdots>k_n>\ell_n>\ell_{n-1}>\cdots>\ell_1\geq4$ be even integers such that $k_i+\ell_i+2=\kappa$ for $1\leq i\leq n$. If $n\leq \dim S_{\kappa}$, then the set of even periods $\{r_{\ell_i}\}_{i=1}^n$ in $S_{\kappa}^\ast$ is linearly independent. Equivalently, if $n\leq \dim S_{\kappa}$, then the set $\{[E_{k_i},E_{\ell_i}]_1\}_{i=1}^n$ is linearly independent.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Tianyu Ni; Ashley Song; Yanhui Su; Hui Xue; Amanda Yin | 2026-06-01

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000284 | OC-4157

Explicit generators of the space of modular forms | main.tex:1018-1020

{oddperiods}}]\label{conj:odd} Let $k_1>k_2>\cdots>k_n\geq\ell_n>\ell_{n-1}>\cdots>\ell_1\geq4$ be even integers such that $k_i+\ell_i=\kappa$ for $1 \leq i \leq n$. If $n\leq \dim S_{\kappa}$, then the set of odd periods $\{r_{\ell_i-1}\}_{i=1}^n$ in $S_{\kappa}^\ast$ is linearly independent. Equivalently, if $n\leq \dim S_{\kappa}$, then the set $\{E_{k_i}E_{\ell_i}-E_{\kappa}\}_{i=1}^n$ is linearly independent.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Tianyu Ni; Ashley Song; Yanhui Su; Hui Xue; Amanda Yin | 2026-06-01

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000285 | OC-4183

On the unit equation $\varepsilon+δ=n$ in cubic fields | Unit.tex:386-391

\label{conj:diostab} Let K be a number field and $\cX/\OO_K$ a model for a hyperbolic curve. Let $\cL$ be an allowable family of extensions L/K. Then $\cX(\OO_L)_{\mathrm{new}}=\emptyset$ for 100\% of $L \in \cL$, when ordered by norms of discriminants.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Maleeha Khawaja; Samir Siksek | 2026-06-01

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000286 | OC-4530

Correlations of zeros of a family of L-functions in function fields with symplectic symmetry | CORRELATIONS_OF_ZEROS.tex:178-206

\label{Andrade&Keating Ratios Conjecture}\label{ratios} Suppose that the real parts of α_k and γ_m are positive and that q odd is the fixed cardinality of the finite field $\mathbb{F}_q$. Then we have \begin{equation*} \begin{split} \sum_{D\in\h}& \frac{\prod_{k=1}^KL\left(\tfrac{1}{2}+\alpha_k,\x_D\right)}{\prod_{m=1}^QL\left(\tfrac{1}{2}+\gamma_m,\x_D\right)}\\ & =\sum_{D\in\h} \sum_{\varepsilon\in\{-1,1\}^K} |D|^{\frac{1}{2} \sum_{k=1}^K(\varepsilon_k-1)\alpha_k} \prod_{k=1}^{K} \xx\left(\tfrac{1}{2} +\tfrac{(1-\varepsilon_k)}{2}\alpha_k\right)\\ &\ \ \ \times Y_{D}(\varepsilon_1\alpha_1,\cdots,\varepsilon_K\alpha_K;\gamma) A_{D}(\varepsilon_1\alpha_1,\cdots,\varepsilon_K\alpha_K;\gamma)+o\left(|D|\right), \end{split} \end{equation*} where $\xx(s)=q^{-\frac{1}{2}+s}$ and $L(s,\chi_D)$ is the quadratic Dirichlet L-function associated to D, with D being a monic and square-free polynomial in $\mathbb{F}_q[T]$, \begin{equation*} Y_D(\alpha;\gamma) = \frac{\underset{j\leqslant k\leqslant K}{\prod} \z(1+\alpha_j+\alpha_k) \underset{m<r \leqslant Q}{\prod} \z(1+\gamma_m+\gamma_r)} {\prod_{k=1}^K\prod_{m=1}^Q \z\left(1+\alpha_k+\gamma_r\right)}, \end{equation*} and \begin{equation*} \begin{split} A_D(\alpha,\gamma) &= Y_D(\alpha;\gamma)^{-1} \prod_{\substack{P \text{ monic}\\\text{irreducible}}}\left(1+\frac{1}{|P|}\right)^{-1} \\ &\ \ \ \times \left(\frac{1}{2} \frac{\prod_{q=1}^Q\left(1-\frac{1}{|P|^{\frac{1}{2}+\gamma_q}}\right)} {\prod_{k=1}^K\left(1-\frac{1}{|P|^{\frac{1}{2}+\alpha_k}}\right)}+ \frac{1}{2} \frac{\prod_{q=1}^Q\left(1+\frac{1}{|P|^{\frac{1}{2}+\gamma_q}}\right)} {\prod_{k=1}^K\left(1+\frac{1}{|P|^{\frac{1}{2}+\alpha_k}}\right)} +\frac{1}{|P|}\right), \end{split} \end{equation*} and $\zeta_{\mathbb{A}}(s)$ is the zeta-function associated to $\mathbb{F}_q[T]$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Julio Andrade; Asmaa Shamesaldeen | 2026-07-07

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000287 | OC-4532

Correlations of zeros of a family of L-functions in function fields with symplectic symmetry | CORRELATIONS_OF_ZEROS.tex:460-472

\label{Our Ratios Conjecture2.} Let $A=\{\alpha_1,\cdots,\alpha_K\}$ and $B=\{\gamma_1\cdots,\gamma_Q\}$ such that α_k is such that, $1\leqslant k \leqslant K$, and γ_m is such that, $1\leqslant m \leqslant Q$ satisfying the restrictions in (\ref{comditions}). Then \begin{equation}\label{df} \begin{split} R_{DL}(A,B)&=\sum_{D\in\h} \frac{\prod_{k=1}^K L\left(\tfrac{1}{2}+\alpha_k,\x_D\right)}{\prod_{m=1}^Q L\left(\tfrac{1}{2}+\gamma_m,\x_D\right)}\\ & = \sum_{D\in\h} \sum_{\D\subseteq A} \left|D\right|^{- \sum_{\delta\in \D}\delta } \prod_{\delta\in \D} \xx\left(\tfrac{1}{2}+\delta\right) Y_{DL}(A,B,\D) A_{DL}(A,B,\D)\\ &\ \ \ + o\left(|D|\right), \end{split} \end{equation} where $Y_{DL}(A,B,\D),A_{DL}(A,B,\D)$ are defined as in equations (\ref{Y_{DL}}) and (\ref{A_{DL}}) respectively.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Julio Andrade; Asmaa Shamesaldeen | 2026-07-07

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000288 | OC-4757

Finiteness for Étale Fundamental Groups of Néron Models | fundgroupsofneron.tex:102-104

\label{conj: abelian variety uniformity} Given a positive integer $g \geq 1$ and a number field $K,$ there exists a constant $C,$ depending only on K and $g,$ such that the following holds. Let A/K an abelian variety over K with dimension $g.$ Then, if

$\mathcal{A}$ is its N\'{e}ron model over $\mathcal{O}_K,$ then $|\pietgeo(\mathcal{A})| \leq C.$

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Frank Lu | 2026-06-30

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000289 | OC-4828

Harder's conjecture and Hermitian automorphic forms | harder_hermitian.tex:809-826

\label{conj:harder} Assume that \(p_\frakp>2k+j-2\) and \(\frakp\mid L^{\alg}(k+j,f)\). Then there exist a Hecke cusp eigenform \[F\in \calS_{{\det}^k\Sym^j}(\Sp_2) \] and a prime ideal \(\frakP\) of \(\bbQ(f)\bbQ(F)\) lying above \(\frakp\) such that, for every prime number \(\ell\), we have \[L_\ell(X,F,\Spin) \equiv L_\ell(X,f) \left(1-\ell^{k-2}X\right) \left(1-\ell^{k+j-1}X\right) \pmod{\frakP}. \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Hidenori Katsurada; Nobuki Takeda | 2026-06-29

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000290 | OC-886

On the discrete convolution of the Liouville and Möbius functions | mobius-liouville7A.tex:510-515

\label{conj:simple zeros}(SZ) We have that \[\sum_{0<\gamma<T}\frac{1}{\left|\zeta^{\prime}\left(\rho\right)\right|}\ll T(\log T)^{1/2}. \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Marco Cantarini; Alessandro Gambini; Alessandro Zaccagnini | 2026-03-10

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000291 | OC-4203

Inexactly Smooth Performance Estimation and New Optimized Gradient Methods | main.tex:1181-1183

For any $\kappa, D > 0$, $q \in [0,1]$, $N, d \in \N$ with $d \geq N+2$, and any starting point $x_0 \in \R^d$, there exists a convex, $L(\delta) = \kappa/\delta^q$-inexactly smooth $f : \R^d \to \R$ with $\|x_0 -x_\star\| \leq D$ and subgradient oracle such that for any subgradient span method, $$f(x_N)-f(x_\star) \geq \left(\frac{(q+1)^\frac{q-1}{q+1}}{q^\frac{q}{q+1}}+o(1)\right)\frac{\kappa^\frac{1}{q+1}D^\frac{2}{q+1}}{(N+1)^\frac{2-q}{q+1}}.$$

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Aaron Zoll; Benjamin Grimmer | 2026-06-01

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000292 | OC-2336

$\mathsf{GL}_N(\mathbb{C})$ Brownian motion and stochastic PDE on entire functions | GLNentireSPDE.tex:417-427

\label{ConjHP} Let $\mathfrak{s} > -\tfrac{1}{2}$ and $\mathfrak{hp}_0\overset{\mathrm{d}}{=}\mathbb{HP}_\mathfrak{s}$. Then, the reciprocals $(\mathsf{y}_{i}(\bullet))_{i\in \mathbb{Z}\backslash\{0\}}$ of zeroes of $\mathfrak{hp}_{\bullet}$ satisfy the equation \begin{equation*} \mathrm{d}\mathsf{y}_i(t) = \sqrt{2}\mathsf{y}_i(t)\mathrm{d}\mathsf{w}_i(t) - 2\mathfrak{s}\mathsf{y}_i(t)\mathrm{d}t + 2 \lim_{\varepsilon\to 0} \sum_{\substack{|\mathsf{y}_j(t)|>\varepsilon\\ j\in \mathbb{Z}\setminus\{0,i\}}} \frac{\mathsf{y}_i(t)\mathsf{y}_j(t)}{\mathsf{y}_i(t)-\mathsf{y}_j(t)}\mathrm{d}t, \ \ \forall {i}\in \mathbb{Z}\backslash \{0\}. \end{equation*}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Assiotis, Theodoros; Mirsajjadi, Zahra Sadat | 2026-05-07

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000293 | OC-2746

Sharp Spectral Thresholds for Multi-View Spiked Wigner Models | draft.tex:3238-3257

We conjecture that this choice of AMP algorithm will converge to equilibrium, as $t\to\infty$. By equilibrium, we mean the following two statements: \begin{enumerate} \item The state evolution parameters converge $q_t \to q_{\ast}\in\RR^L$, where $q_{\ast}$ is the solution to the following fixed-point equation \begin{equation} q_{\ast} = 1 - \diag\sbr{\rbr{\Diag\rbr{\lambda\odot q_{\ast}}+B^{-1}}^{-1}}. \end{equation} In comparison with our self-consistent equations \eqref{eq: self-consistent eq matrix}, it holds that $q_{\ast}^{(l)}=1-M_{ll}(1)$. \item Let $m_t\in\RR^{nL}$ be the concatenation of $m_t^{(1)},\ldots,m_t^{(L)}$. Then it also converges to equilibrium, as $m_t\to m_{\ast}\in\RR^{nL}$, where $m_{\ast}$ satisfies that \begin{equation}\label{eq:AMP variable equilibrium} m_{\ast}^{(l)}=\sqrt{\frac{\lambda_l}{n}} A^{(l)} \cE_\ast^{(l)}(m_{\ast})-\lambda_l\rbr{1-q_{\ast}^{(l)}}\cE_\ast^{(l)}(m_{\ast}), \end{equation} where we use $\cE_{\ast}$ to denote the denoiser with the limiting state evolution parameters $q_{\ast}$. Moreover, we have the following distributional result \begin{equation}\label{eq:AMP distributional} m_{\ast}^{(l)} \overset{d.}{\approx} \lambda_lq_{\ast}^{(l)} X^{(l)}+\sqrt{\lambda_l q_{\ast}^{(l)}}\cN(0,I_n),\quad l\in[L]. \end{equation} Moreover, we also have $X^{(l)\top}\cE^{(l)}_\ast\rbr{m_{\ast}}/n \overset{d.}{\approx} q_{\ast}^{(l)}$. \end{enumerate}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Xiaodong Yang; Subhabrata Sen; Yue M. Lu | 2026-05-19

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000294 | OC-321

On the depth of depth-weighted trees | main.tex:423-428

\label{conj:exp} For every weight function f satisfying assumption~\eqref{eq:ExpG} for some $c > 1$, there is $\nu=\nu(c)\in(0,1)$ such that \be \frac{d(T_n)}{n}\toas \nu. \ee

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Lyuben Lichev; Amitai Linker; Bas Lodewijks; Dieter Mitsche | 2026-02-17

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000295 | OC-417

Half-plane non-coexistence without FKG | main2__6_.tex:615-617

Let $k \in \Z_{\geq 1}$. Let μ be a translation-invariant, ergodic, finite-energy, k-dependent edge percolation measure on $\Z^2$. Then the trichotomy of Theorem~\ref{thm:main} holds in the full plane.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Frederik Ravn Klausen; Noah Kravitz | 2026-02-12

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by_sa

V3-000296 | OC-418

Half-plane non-coexistence without FKG | main2__6_.tex:633-635

Let μ be a translation-invariant, ergodic percolation measure which almost surely has infinitely many infinite clusters. Then almost surely both $\mu_{\hp}$ and $\mu_{\hp}^*$ have infinitely many infinite clusters.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Frederik Ravn Klausen; Noah Kravitz | 2026-02-12

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by_sa

V3-000297 | OC-4414

Uncountably many extremal Series--Sinai states for the Ising model on Lobachevsky lattices | Vereecke_260709.tex:419-429

[Extremality of all Series--Sinai states, \cite{SS90}] \label{conj:ss_conjecture} Let \(p,\,q\ge{3}\) such that \(1/p+1/q<1/2\). For \(\beta \in (0, \infty)\) large enough, \begin{equation} \mathrm{SS}_{\beta}(p,q) \subset \exG_\beta. \end{equation}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Jean Vereecke | 2026-07-09

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000298 | OC-4462

Loop Equations Characterize Random Matrix Statistics | Loop_eq_characterization.tex:1017-1019

For any \(\beta>0\), not necessarily rational, the following statements hold. Under \Cref{a:bulk}, the poles of \(s\) define a point process on \(\bR\), and this point process has the same law as the Sine_{β} point process. Under \Cref{a:edge}, the poles of \(s\) define a point process on \(\bR\), and this point process has the same law as the \(\mathrm{Airy}_{\beta}\) point process.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Paul Bourgade; Jiaoyang Huang | 2026-07-08

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000299 | OC-4463

Loop Equations Characterize Random Matrix Statistics | Loop_eq_characterization.tex:2820-2824

For every fixed integer \(d\geq 3\), the local statistics of the bulk eigenvalues of random \(d\)-regular graphs on \(N\) vertices converge, as \(N\to\infty\), to those of the {\rm GOE}.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Paul Bourgade; Jiaoyang Huang | 2026-07-08

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000300 | OC-4481

Norm Bounds for Sparse Random Tensors and Spectral Gap of Random Hypergraphs | VJ26main.tex:290-297

\label{conj:seginerTensor} Let $T\in \R^{n_1\times \ldots \times n_r}$ be a random tensor with i.i.d.\ mean-zero entries. Then, \[\E \normin{T} \underset{r} \asymp \max_{k\in [r]} \E \max_{\substack{i_1\in [n_1]\\\ldots\\i_{r}\in [n_{r}]}} \left(\sum_{j=1}^{n_k} T^2_{i_1, \ldots, i_{k-1}, j, i_{k+1}, \ldots, i_r}\right)^{1/2}\,. \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Kevin Lucca; Lucas Pesenti | 2026-07-08

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000301 | OC-4681

On the range of competing random walks | main.tex:1158-1166

If $X^1,\dots,X^N$ are in the domain of attraction of a β-stable process U with scale function $b(n)$ and that $d/\beta<1$, then \begin{equation*} \frac{1}{b(n)}\R_n^k\distribution\lambda\left(\left\{x\in\RR, \tau_x^k\le 1\wedge\min_{j\ne k}\tau_x^j\right\}\right), \end{equation*} where $U^1,\dots,U^N$ are $i.i.d.$ copies of U and where $\tau_x^\ell\eqdef\inf\{t>0, U_t^\ell=x\}$. Furthermore, joint convergence in $k\in[\![N]\!]$ also holds.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Maxence Baccara | 2026-07-02

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000302 | OC-626

Excursion decomposition of the XOR-Ising model | main.tex:417-419

\label{conj:AT} The excursion decomposition of the continuum Ashkin-Teller polarisation field with parameter U is given by those in Theorem \ref{thm:cont-general} for the appropriate value $\alpha=\alpha(U)$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Tomás Alcalde López; Avelio Sepúlveda | 2026-02-05

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000303 | OC-857

An asymptotically optimal bound for the concentration function of a sum of independent integer random variables | discrete_inverse.tex:430-434

(Juškevičius~\cite{tjp})\label{conj.main} Let $X_1, \dots, X_n$ and $Y_1, \dots, Y_n$ be as in Theorem~\ref{thm.tse}. Then there are $a_1, \dots, a_n \in \{-1,1\}$ such that $\conc(\sum_{i=1}^n X_i) \le \conc(\sum_{i=1}^n a_i Y_i)$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Valentas Kurauskas | 2026-03-11

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000304 | OC-2806

Shifted quantum toroidal algebra of type $\mathfrak{gl}_{1|1}$ and the Pieri rule of the super Macdonald polynomials | superMac-v1.tex:288-307

\label{Pieri-con} \begin{equation} E_{1,0} = \pi_1, \qquad E_{2,0} = \sum_{k=1}^\infty c_k[p] \frac{\partial}{\partial \pi_k}, \end{equation} \begin{equation} F_{1,+1} = \frac{\partial}{\partial \pi_1}, \qquad F_{2,-1} = -\sum_{k=1}^\infty \pi_k \cdot \widetilde{c}_k \left[\partial/ \partial p \right], \end{equation} where the generating functions for the coefficients are \begin{equation}\label{c-gen} \sum_{k=0}^\infty c_k[p] z^k = \exp \left(\sum_{r=1}^\infty \frac{1-t^r}{r} p_r z^r \right) = \prod_{i=1}^\infty \frac{1-tu x_i}{1-ux_i}, \end{equation} and \begin{equation}\label{tc-gen} \sum_{k=0}^\infty \widetilde{c}_k \left[\partial/ \partial p \right] z^{-k} = \exp \left(\sum_{n=1}^\infty (q^{-n}-1) \frac{\partial}{\partial p_n} z^{-n} \right). \end{equation}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Hiroaki Kanno; Ryo Ohkawa; Jun'ichi Shiraishi | 2026-05-16

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000305 | OC-2807

Shifted quantum toroidal algebra of type $\mathfrak{gl}_{1|1}$ and the Pieri rule of the super Macdonald polynomials | superMac-v1.tex:371-381

\label{Pieri-con2} \begin{align} u^{-1} (q/t)^{\frac{1}{2}} E_{2,1} &= -\sum_{k=1}^\infty \oint \frac{dw}{w} w^{-2k} V_B^{(-)}(w) V_B^{(+)}(w) \langle \varnothing \vert \widetilde{V}_F^{(-)}(w) \widetilde{V}_F^{(+)}(w) \vert \varnothing\rangle_{F} q^{k} \frac{\partial}{\partial \pi_k}, \\ u^{-1} F_{1,2} &= \sum_{k=1}^\infty\oint \frac{dw}{w} w^{-2k+2} V_B^{(-)}(w) V_B^{(+)}(w) \langle \varnothing \vert \widetilde{V}_F^{(-)}(w) \widetilde{V}_F^{(+)}(w) \vert \varnothing\rangle_{F} q^{k-1} \frac{\partial}{\partial \pi_k}. \end{align}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Hiroaki Kanno; Ryo Ohkawa; Jun'ichi Shiraishi | 2026-05-16

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000306 | OC-2808

Shifted quantum toroidal algebra of type $\mathfrak{gl}_{1|1}$ and the Pieri rule of the super Macdonald polynomials | superMac-v1.tex:1677-1688

\label{E21-F12-con} \begin{align} u^{-1} (q/t)^{\frac{1}{2}} E_{2,1} &= -\sum_{k=1}^\infty \oint \frac{dw}{w} w^{-2k} V_B^{(-)}(w) V_B^{(+)}(w) \langle \varnothing \vert \widetilde{V}_F^{(-)}(w) \widetilde{V}_F^{(+)}(w) \vert \varnothing\rangle_{F} q^{k} \frac{\partial}{\partial \pi_k}, \\ u^{-1} F_{1,2} &= \sum_{k=1}^\infty\oint \frac{dw}{w} w^{-2k+2} V_B^{(-)}(w) V_B^{(+)}(w) \langle \varnothing \vert \widetilde{V}_F^{(-)}(w) \widetilde{V}_F^{(+)}(w) \vert \varnothing\rangle_{F} q^{k-1} \frac{\partial}{\partial \pi_k}. \end{align}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Hiroaki Kanno; Ryo Ohkawa; Jun'ichi Shiraishi | 2026-05-16

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000307 | OC-4214

Zonal Spherical Functions of Quantum Symmetric Pairs | main.tex:1753-1758

\label{conj-FII} Lemma~\ref{lem-rk1-nonletzter-macdonald} also holds for FII, for parameters \[(a,b,c,d) = (q_\beta^{11},-q_\beta^7,q_\beta^5,-q_\beta). \]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Philip Schlösser | 2026-06-26

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by_sa

V3-000308 | OC-2783

Weakly Noetherian Lie Algebra and the Sierra-Walton Conjecture | May15.tex:463-465

\label{conjecture:enveloping-Noetherian} \cite{Sierra-Walton} If $\fg$ is infinite dimensional, then $U(\fg)$ is not Noetherian.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Olivier Mathieu | 2026-05-18

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000309 | OC-3725

The graded Grothendieck group ${K}_0^{\mathrm{gr}}$ is full for weighted Leavitt path algebras | RJHazratNamJune2026.tex:253-265

\label{conjalg} Let E and F be finite graphs and $\K$ a field. \begin{enumerate}[\upshape(1)] \item For any order preserving $\Z[x,x^{-1}]$-module homomorphism $\phi: K^{\gr}_0(L_\K(E)) \rightarrow K^{\gr}_0(L_\K(F))$ with $\phi([L_\K(E)])=L_\K(F)$, there exists a unital $\mathbb Z$-graded $\K$-homomorphism $\psi: L_\K(E) \rightarrow L_\K(F)$ such that $K^{\gr}_0(\psi) = \phi$. \medskip \item If in \textnormal{(1)}, the ϕ is an isomorphism, then there exists a unital $\mathbb Z$-graded $\K$-isomorphism $\psi: L_\K(E) \rightarrow L_\K(F)$ such that $K^{\gr}_0(\psi) = \phi$. \end{enumerate}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Remarl Joseph Damalerio; Roozbeh Hazrat; Tran Giang Nam | 2026-06-13

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000310 | OC-2339

A Sugawara-Legendre mechanism for the hyperelliptic Heisenberg algebra | P4_RepMechanism_main.tex:830-838

\label{conj:irreducibility-general} For all $m \geq 3$, the equivalence of Theorem~\textup{\ref{thm:irreducibility}} extends to the superelliptic Heisenberg algebra $\Hm$ with multiple cocycle components $\omega_1, \ldots, \omega_{\lfloor m/2 \rfloor}$. The p-admissibility condition generalizes to requiring that $\varphi(b_0^{(j)})$ avoids a discrete set of singular values determined by $\psi_{mn}^{(ij)}(a)$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Santos, Felipe Albino dos | 2026-05-07

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000311 | OC-2641

Categorical Lusztig cycles and weave schobers | main.tex:2088-2090

\label{conj:filtered_by_weighted_braid_words} Let $\bw$ be a weighted braid word. Then there exists a unique $\bw$-filtered object $\lbw\in\D$ whose geometric realization is rigid.\qed

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Roger Casals; Merlin Christ | 2026-05-21

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000312 | OC-3635

Spectral Intertwining Operators | main.tex:154-165

\label{conjecture-Langlands} For irreducible representations π, the normalized intertwining operators \[J_{Q|P}^\circ(\pi):= \gamma(0, \pi, \ad_{\fn_{Q^\vee}/\fn_{P^\vee}}) J_{Q|P}(\pi) := \epsilon(0, \pi, \ad_{\fn_{Q^\vee}/\fn_{P^\vee}})\frac{L(1, \pi, \ad_{\fn_{Q^\vee}/\fn_{P^\vee}})}{L(0, \pi, \ad_{\fn_{Q^\vee}/\fn_{P^\vee}})}J_{Q|P}(\pi), \] where the ϵ and L's denote the conjectural ϵ-factor and L-functions for automorphic representations, satisfy the transitivity \[J^\circ_{R|Q} \circ J^\circ_{Q|P} = J^\circ_{R|P}.\]

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Qiyuan Chen | 2026-06-16

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by_sa

V3-000313 | OC-3636

Spectral Intertwining Operators | main.tex:3414-3461

\label{categorical} Let $G = (G^*)_b$ be as above. For a fixed Whittaker datum of G^*, there is an equivalence of categories \[\widetilde{\operatorname{LL}}_{G^*} \colon D_{\text{lis}}(\operatorname{Bun}_{G^*})^\omega \simeq D\Coh^{\text{qc}}(\Par_{G^\vee}), \] where ω denotes the full subcategory of compact object and qc denotes the full subcategory of objects with compact supports, satisfying the following properties: \begin{enumerate} \item Under this functor, the parabolic induction operator corresponds to the Eisenstein operator. More precisely, let P be a parabolic subgroup of G with Levi subgroup M, denote by $\bar{P}$ the opposite parabolic subgroup of P. We have $M=(M^*)_b$ for the quasi-split inner form M^* of M and the Whittaker datum for G induces a Whittaker datum for M^*. Then the diagram \[\begin{tikzcd} D(\mathsf{Rep}(M(F))) & D_{\text{lis}}(\operatorname{Bun}_{M^*})^\omega & D\Coh^{\text{qc}}(\Par_{M^\vee}) \\ D(\mathsf{Rep}(G(F))) & D_{\text{lis}}(\operatorname{Bun}_{G^*})^\omega & D\Coh^{\text{qc}}(\Par_{G^\vee}) \arrow["i_{b!}",from=1-1, to=1-2] \arrow["\widetilde{\operatorname{LL}}_{M^*}",from=1-2, to=1-3] \arrow["i_{b!}",from=2-1, to=2-2] \arrow["\widetilde{\operatorname{LL}}_{G^*}",from=2-2, to=2-3] \arrow["{i_{\bar{P}}^G}"', from=1-1, to=2-1] \arrow["{\Eis^a_{\bar{P}!}}", from=1-2, to=2-2] \arrow["{\Eis_{P^\vee}}", from=1-3, to=2-3] \end{tikzcd} \] commutes up to isomorphisms, where $\Eis^a_{\bar{P}!}$ is the Eisenstein operator for Bun's, defined in \cite{hamann}, Definition 2.1.8. In this notation, ``$a$'' stands for ``automorphic side''. \item This functor is compatible with automorphisms: for an automorphism σ of G^* that preserves b and a pinning, it induces an isomorphism $\sigma^\vee$ of $G^\vee$. Then the diagram \[\begin{tikzcd} D(\mathsf{Rep}(G(F))) & D_{\text{lis}}(\operatorname{Bun}_{G^*})^\omega & D\Coh^{\text{qc}}(\Par_{G^\vee}) \\ D(\mathsf{Rep}(G(F))) & D_{\text{lis}}(\operatorname{Bun}_{G^*})^\omega & D\Coh^{\text{qc}}(\Par_{G^\vee}) \arrow["i_{b!}",from=1-1, to=1-2] \arrow["\widetilde{\operatorname{LL}}_{G^*}",from=1-2, to=1-3] \arrow["i_{b!}",from=2-1, to=2-2] \arrow["\widetilde{\operatorname{LL}}_{G^*}",from=2-2, to=2-3] \arrow["\sigma"', from=1-1, to=2-1] \arrow["\sigma", from=1-2, to=2-2] \arrow["\sigma^\vee", from=1-3, to=2-3] \end{tikzcd} \] commutes up to isomorphisms. Such a functor is called categorical local Langlands correspondence. \item The correspondence is compatible with the Kottwitz isomorphism. More precisely, for any $V \in \mathsf{Rep}(G(F))$, The $Z(G^\vee)^{W_F}$-character of the fiber of $\widetilde{\operatorname{LL}}_{G^*} \circ i_{b!}(V)$ at any point in $\Par_{G^\vee}$ is $\kappa(b)$ under the Kottwitz isomorphism $\kappa\colon B(G^*)_{bas} \simeq X^*(Z(G^\vee)^{W_F})$. \end{enumerate}

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Qiyuan Chen | 2026-06-16

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by_sa

V3-000314 | OC-2604

On the fundamental spectral gap of weighted Schrodinger operators | ON_THE_FUNDAMENTAL_SPECTRAL_GAP_OF_WEIGHTED_SCHR__DINGER_OPERATORS.tex:372-387

We consider the weighted Schr\"{o}dinger operators \begin{equation}\label{eq12} \left\{ \begin{array}{lr} H(V,w) u := -u^{\prime \prime} +V(x) u = \lambda w(x) u \\ u(0)=u(\pi)=0 & \end{array} \right. \end{equation} For a given bounded measurable, strictly positive weight function w, any gap-minimizing optimal potential $V_{*}\in \mathcal{C}_M $ is constant.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Mohammed Ahrami | 2026-05-23

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000315 | OC-4827

Inequalities between Dirichlet and Neumann eigenvalues in large dimensions | Larged2.tex:242-248

\label{conj15} There exist two absolute positive constants C_*, $\al_*$ such that $$ \Psi(d,k,\O) \ge C_* e^{\al_* d} \qquad \forall \ k \in \N, \quad \forall \ \O \subset \R^d. $$

自动抽取的猜想块 | source_machine_label_real_open_conjecture | N. Filonov | 2026-06-29

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000316 | OC-4016

Revisiting the Behrens-Fisher Problem: Validity-First Optimality | behrens_fisher_martin_liu_im_rev2.tex:1505-1514

\label{conj:prior-free} The symmetric-interval-projection restriction in the shortest uniformly valid projection theorem and in Corollary~\ref{cor:proj-admiss} is inessential. Among all prior-free procedures that are exactly and uniformly valid for \(\psi\) over the variance ratio, none can have plausibility interval length no greater than that of the IM/Hsu interval at every \(\xi\) and strictly smaller at some \(\xi\). Equivalently, the IM/Hsu interval is admissible in length, uniformly in the variance ratio, within the entire prior-free class.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Xiao Wang; Chuanhai Liu | 2026-06-05

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000317 | OC-1946

Random Access Codes: Explicit Constructions, Optimality, and Classical-Quantum Gaps | main.tex:1171-1174

\label{conjecture:worst} An optimal set S of the problem Eq.~\eqref{eq:worst-prob} can be chosen so that $S\subset\{0,1\}^L$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Ruho Kondo; Yuki Sato; Hiroshi Yano; Yota Maeda; Kosuke Ito; Naoki Yamamoto | 2026-04-23

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000318 | OC-601

Symmetry and localisation in causally constrained quantum operator dynamics | main.tex:1325-1327

An irreducible k-wall on qubits starts and ends in an element of a conditional unitary class with a fixed gauge.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Marcell D. Kovács; Christopher J. Turner; Lluís Masanes | 2026-02-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000319 | OC-603

Symmetry and localisation in causally constrained quantum operator dynamics | main.tex:1337-1339

A qudit wall begins and ends with a generalised conditional unitary.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Marcell D. Kovács; Christopher J. Turner; Lluís Masanes | 2026-02-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000320 | OC-605

Symmetry and localisation in causally constrained quantum operator dynamics | main.tex:1367-1369

For a probability distribution chosen so that $\mathds{P}(\text{k-wall})$ is finite, the $n\rightarrow\infty$ chain has extensively many fragment subspaces and therefore an exponentially large localised operator subspace.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Marcell D. Kovács; Christopher J. Turner; Lluís Masanes | 2026-02-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000321 | OC-606

Symmetry and localisation in causally constrained quantum operator dynamics | main.tex:1384-1386

almost every local floquet circuit of finite depth and finite gate width has super-logarithmic operator spreading

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Marcell D. Kovács; Christopher J. Turner; Lluís Masanes | 2026-02-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000322 | OC-607

Symmetry and localisation in causally constrained quantum operator dynamics | main.tex:1387-1389

almost every local floquet circuit of finite depth and finite gate width does not have a complete generating set of conserved quantities that are exponentially localised

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Marcell D. Kovács; Christopher J. Turner; Lluís Masanes | 2026-02-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000323 | OC-608

Symmetry and localisation in causally constrained quantum operator dynamics | main.tex:1390-1392

local perturbation of bounded light cone almost always produces a linear light cone

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Marcell D. Kovács; Christopher J. Turner; Lluís Masanes | 2026-02-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000324 | OC-609

Symmetry and localisation in causally constrained quantum operator dynamics | main.tex:1393-1395

local perturbation of a circuit with an $O(x')$ light cones where $x = o(x')$ always produces a $\Omega(y)$ light cone where y is the light cone resulting almost always from local perturbation from a x light cone.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Marcell D. Kovács; Christopher J. Turner; Lluís Masanes | 2026-02-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000325 | OC-610

Symmetry and localisation in causally constrained quantum operator dynamics | main.tex:1397-1399

There is no circuit MBL.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Marcell D. Kovács; Christopher J. Turner; Lluís Masanes | 2026-02-06

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

V3-000326 | OC-3575

Establishing an $Ω(\sqrt{d})$ complexity lower bound for PDMP samplers and how to break it: a sub-$\sqrt{d}$ algorithm for Gaussian-tailed targets | main.tex:484-486

For any PDMP sampling algorithm satisfying Assumption 1 (Target-independence of the flow and auxiliary space), Assumption 2 (Locality of the auxiliary distribution), and Assumption 3 (Local jumps), there exists a one-dimensional distribution π_1 (satisfying reasonable assumptions) such that the sampling complexity for the i.i.d. target distribution $\pi(x) = \prod_{i=1}^d \pi_1(x_i)$ on $\mathbb{R}^d$ is bounded below by $\Omega(\sqrt{d})$.

自动抽取的猜想块 | source_machine_label_real_open_conjecture | Augustin Chevallier | 2026-06-18

源 LaTeX 猜想块，外部定义见论文 | 原始来源 | cc_by

B OEIS 历史与当前提及：17,579 条

每条为一个序列页的关键词命中段集合。作者署名与段内更新者署名均保留。历史猜想、被证明的旧猜想、条件性假设或对猜想的提及可能同时出现。

B01 A000000 - A049999 | 1,218 条

V3-000327 | OEIS-A000009

Expansion of Product_{m >= 1} (1 + x^m); number of partitions of n into distinct parts; number of partitions of n into odd parts.

Conjecture: Each positive integer n can be written as a_1 + ... + a_k, where a_1,...,a_k are strict partition numbers (i.e., terms of the current sequence) with no one dividing another. This has been verified for n = 1..1350. - _Zhi-Wei Sun_, Apr 14 2023 Conjecture: For each integer n > 7, a(n) divides none of p(n), p(n) - 1 and p(n) + 1, where p(n) is the number of partitions of n given by A000041. This has been verified for n up to 10^5. - _Zhi-Wei Sun_, May 20 2023 [Verified for n <= 2*10^6. - _Vaclav Kotesovec_, May 23 2023]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000328 | OEIS-A000033

a(n) = A058087(n,n-2).

Conjecture: a(n) + 2*a(n+p) + a(n+2*p) is divisible by p for any prime p. - _Mark van Hoeij_, Jun 10 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000329 | OEIS-A000041

a(n) is the number of partitions of n (the partition numbers).

Conjecture: starting with offset 1 represents the numbers of ordered compositions of n using the signed (++--++...) terms of A001318 starting (1, 2, -5, -7, 12, 15, ...). - _Gary W. Adamson_, Apr 04 2013 (this is true by the pentagonal number theorem, _Joerg Arndt_, Apr 08 2013) Conjecture: No a(n) has the form x^m with m > 1 and x > 1. - _Zhi-Wei Sun_, Dec 02 2013 Conjecture: Each integer n > 2 different from 6 can be written as a sum of finitely many numbers of the form a(k) + 2 (k > 0) with no summand dividing another. This has been verified for n <= 7140. - _Zhi-Wei Sun_, May 16 2023 G.f.: (conjecture) (r(x) * r(x^2) * r(x^4) * r(x^8) * ...) where r(x) is A000009: (1, 1, 1, 2, 2, 3, 4, ...). - _Gary W. Adamson_, Sep 18 2016; _Doron Zeilberger_ observed today that "This follows immediately from Euler's formula 1/(1-z) = (1+z)*(1+z^2)*(1+z^4)*(1+z^8)*..." _Gary W. Adamson_, Sep 20 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000330 | OEIS-A000043

Mersenne exponents: primes p such that 2^p - 1 is prime. Then 2^p - 1 is called a Mersenne prime.

Conjecture: for k > 1, 2^k-1 is (a Mersenne) prime or k = 2^(2^m)+1 (is a Fermat number) if and only if (k-1)^(2^k-2) == 1 (mod (2^k-1)k^2). - _Thomas Ordowski_, Oct 05 2023 Conjecture: for p prime, 2^p-1 is (a Mersenne) prime or p = 2^(2^m)+1 (is a Fermat number) if and only if (p-1)^(2^p-2) == 1 (mod 2^p-1). - _David Barina_, Nov 25 2024 Conjecture: a(1) = 2; and then, for n > 0, a(n+1) is the least k such that lpf(2^k - 1) > lpf(2^a(n) - 1), where lpf(m) is the least prime factor of m. See A049479 (look at the record value positions). - _Thomas Ordowski_, Sep 03 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000331 | OEIS-A000048

Number of n-bead necklaces with beads of 2 colors and primitive period n, when turning over is not allowed but the two colors can be interchanged.

Conjecture: Also the number of caliber-n cycles of Zagier-reduced indefinite binary quadratic forms with sum invariant equal to s, where (s-1)/n is an odd integer. - _Barry R. Smith_, Dec 14 2014 The comment above is true for discriminants of the form a^2+4 (as Smith stated it, Conjecture 7 of his paper). Proved by an explicit bijection that turns kneading into the cyclic rotation of binary words of odd weight; Smith's Conjecture 6 is also proved in full (see link). - _Roberto Blanco Gómez_, Oct 07 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000332 | OEIS-A000071

a(n) = Fibonacci(n) - 1.

We can establish _Gerald McGarvey_'s conjecture mentioned in the Formula section, however we require n > 4. We need the following 4 prerequisites. (1) a(n) = F(n) - 1, with {F(n)}_{n >= 1} the Fibonacci numbers A000045. (2) (Binet form) F(n) = (d^n - e^n)/sqrt(5) with d = phi and e = 1 - phi, de = -1 and d + e = 1. It follows that a(n) = (d(n) - e(n))/sqrt(5) - 1. (3) To prove floor(x) = y is equivalent to proving that x - y lies in the half-open interval [0, 1). (4) The series {s(n) = c1 x^n + c2}_{n >= 1}, with -1 < x < 0, and c1 and c2 positive constants, converges by oscillation with s(1) < s(3) < s(5) < ... < s(6) < s(4) < s(2). If follows that for any odd n, the open interval (s(n), s(n+1)) contains the subsequence {s(t)}_{t >= n + 2}. Using these prerequisites we can analyze the conjecture. Conjecture: for k a positive odd integer, the sequence {a(k^n): n >= 1} is a strong divisibility sequence; that is, for n, m >= 1, gcd(a(k^n), a(k^m)) = a(k^gcd(n,m)). - _Peter Bala_, Dec 05 2022 Conjecture: for all c such that 2*(2 - Phi) <= c < (2 + Phi)*(2 - Phi) we have a(n) = floor(Phi*a(n-1) + c) for n > 4. - _Gerald McGarvey_, Jul 22 2004. This is true provided n > 3 is changed to n > 4, see proof in Comments section. - _Russell Jay Hendel_, Mar 15 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000333 | OEIS-A000089

Number of solutions to x^2 + 1 == 0 (mod n).

The Dirichlet inverse, 1, -1, 0, 1, -2, 0, 0, -1, 0, 2, 0, 0, -2, 0,.. seems to equal A091400, apart from signs. - _R. J. Mathar_, Jul 15 2010 [This conjecture is correct. - _Isaac Saffold_, Feb 02 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000334 | OEIS-A000105

Number of free polyominoes (or square animals) with n cells.

Conjecture: Almost all polyominoes are holey. In other words, A000104(n)/a(n) tends to 0 for increasing n. - _John Mason_, Dec 11 2021 (This is true, a consequence of Madras's 1999 pattern theorem. - _Johann Peters_, Jan 06 2024)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000335 | OEIS-A000110

Bell or exponential numbers: number of ways to partition a set of n labeled elements.

When n is prime, a(n) == 2 (mod n), but the converse is not always true. Define a Bell pseudoprime to be a composite number n such that a(n) == 2 (mod n). W. F. Lunnon recently found the Bell pseudoprimes 21361 = 41*521 and C46 = 3*23*16218646893090134590535390526854205539989357 and conjectured that Bell pseudoprimes are extremely scarce. So the second Bell pseudoprime is unlikely to be known with certainty in the near future. I confirmed that 21361 is the first. - _David W. Wilson_, Aug 04 2007 and Sep 24 2007 Conjecture: No a(n) has the form x^m with m > 1 and x > 1. - _Zhi-Wei Sun_, Dec 02 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000336 | OEIS-A000111

Euler or up/down numbers: e.g.f. sec(x) + tan(x). Also for n >= 2, half the number of alternating permutations on n letters (A001250).

G.f.: conjecture: 1 + T(0)*x/(1-x), where T(k) = 1 - x^2*(k+1)*(k+2)/(x^2*(k+1)*(k+2) - 2*(1-x*(k+1))*(1-x*(k+2))/T(k+1)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000337 | OEIS-A000122

Expansion of Jacobi theta function theta_3(x) = Sum_{m=-oo..oo} x^(m^2) (number of integer solutions to k^2 = n).

Bala's conjecture is true (see Fried link). - _Sela Fried_, Dec 04 2025 Conjecture: Sum_{k>=0} a(k) / exp(k*Pi) = Pi^(1/4)/Gamma(3/4) = A175573. - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000338 | OEIS-A000123

Number of binary partitions: number of partitions of 2n into powers of 2.

Conjecture: Limit_{n ->infinity} (log(n)*a(2n))/(n*a(n)) = c = 1.63... - _Benoit Cloitre_, Jan 26 2003 [The constant c is equal to 2*log(2) = 1.38629436... =A016627. - _Vaclav Kotesovec_, Aug 07 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000339 | OEIS-A000126

A nonlinear binomial sum.

Conjecture. Let S(1)={1} and, for n>1, let S(n) be the smallest set containing x+1 and 2x+1 for each element x in S(n-1). Then a(n) is the number of the elements in S(n). (See A122554 for a sequence defined in this way.) - _John W. Layman_, Nov 21 2007 [edited by _Sela Fried_, Feb 18 2026]. Layman's conjecture is true (see Fried link). - _Sela Fried_, Feb 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000340 | OEIS-A000139

a(n) = 2*(3*n)! / ((2*n+1)!*(n+1)!).

Conjecture: a(n) is odd iff n is a term of A022341. - _Peter Bala_, Jul 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, entry revised Apr 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000341 | OEIS-A000142

Factorial numbers: n! = 1*2*3*4*...*n (order of symmetric group S_n, number of permutations of n letters).

Conjecturally, 1, 6, and 120 are the only numbers which are both triangular and factorial. - Christopher M. Tomaszewski (cmt1288(AT)comcast.net), Mar 30 2005 Conjecture: a(n) = 1/(2*Pi)*Integral_{x=-oo..oo}(n+i*x+1)!/(i*x+1)-(n+i*x-1)!/(i*x-1)dx. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000342 | OEIS-A000155

Nearest integer to modified Bessel function K_n(1).

Conjecture: a(n) = Integral_{x=0..oo} ChebyshevU(n - 1, x)/exp(x) dx, where ChebyshevU(n, x) is the n-th Chebyshev polynomial of the second kind with variable x. - _Velin Yanev_, Jan 14 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000343 | OEIS-A000159

a(n) = A058087(n,n-3).

Conjecture: 2*(-252307*n + 1041077)*a(n) + (504614*n^2 - 3362985*n + 5118150)*a(n-1) + (1280831*n^2 - 7397886*n + 6461565)*a(n-2) + (746598*n^2 - 2913543*n - 1336090)*a(n-3) + (-405481*n^2 + 6175011*n - 15469320)*a(n-4) + (-375862*n^2 + 4098537*n - 8846430)*a(n-5) + 2*(-187931*n + 560630)*a(n-6) = 0. - _R. J. Mathar_, Nov 02 2015 Conjecture: a(n) + 2*a(n+p) + a(n+2*p) is divisible by p for any prime p except 3. - _Mark van Hoeij_, Jun 10 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000344 | OEIS-A000166

Subfactorial or rencontres numbers, or derangements: number of permutations of n elements with no fixed points.

Conjecture: a(n) with n > 2 is a perfect power only for n = 4 with a(4) = 3^2. This has been verified for n <= 1000. - _Zhi-Wei Sun_, Jan 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000345 | OEIS-A000170

Number of ways of placing n nonattacking queens on an n X n board.

Strong conjecture: there is a constant c around 2.54 such that a(n) is asymptotic to n!/c^n; weak conjecture: lim_{n -> infinity} (1/n) * log(n!/a(n)) = constant = 0.90.... - _Benoit Cloitre_, Nov 10 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000346 | OEIS-A000172

The Franel number a(n) = Sum_{k = 0..n} binomial(n,k)^3.

An identity of V. Strehl states that a(n) = Sum_{k = 0..n} C(n,k)^2 * binomial(2*k,n). Zhi-Wei Sun conjectured that for every n = 2,3,... the polynomial f_n(x) = Sum_{k = 0..n} binomial(n,k)^2 * binomial(2*k,n) * x^(n-k) is irreducible over the field of rational numbers. - _Zhi-Wei Sun_, Mar 21 2013 Conjecture: a(n) == 2 (mod n^3) iff n is prime. - _Gary Detlefs_, Mar 22 2013 Bala's conjecture is true - up to a missing factor (-1)^n. See Fried link. - _Sela Fried_, Oct 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000347 | OEIS-A000181

a(n) = A058087(n,n-4).

Conjecture: 3*(2111*n-8303)*(n-4)*(-9+2*n)^2*a(n) - (n-3)*(25332*n^4 - 377236*n^3 + 1898681*n^2 - 3320738*n + 484000)*a(n-1) - 2*(n-4)*(12140*n^4 - 118152*n^3 + 337063*n^2 - 377436*n + 225720)*a(n-2) + (1052*n^5 - 40656*n^4 + 266063*n^3 - 549153*n^2 + 49850*n + 655200)*a(n-3) +(263*n+640)*(n-3)*(-7+2*n)^2*a(n-4) = 0. - _R. J. Mathar_, Nov 02 2015 Conjecture: a(n) + 2*a(n+p) + a(n+2*p) is divisible by p for any prime p > 3. - _Mark van Hoeij_, Jun 10 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000348 | OEIS-A000182

Tangent (or "Zag") numbers: e.g.f. tan(x), also (up to signs) e.g.f. tanh(x).

Conjectures: Conjectured o.g.f.: x*(1 + 2*x/(1 + x - 9*x/(1 - 8*x/(1 + x - 25*x/(1 - 24*x/(1 + x - 49*x/(1 - 48*x/(1 + x - ...)))))))). - _Peter Bala_, Dec 15 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000349 | OEIS-A000185

a(n) = A058087(n,n-5).

Conjecture: +4*(210968408*n^2 -1603518486*n +2343057493) *a(n) +(-843873632*n^3 -4039256254*n^2 +144382575631*n -553812368850) *a(n-1) +(10453330198*n^3 -175111274403*n^2 +798927275864*n -639098546595) *a(n-2) +(10059264970*n^3 -98879552663*n^2 +170576803994*n -134993524720) *a(n-3) +(470894110*n^3 -5178116941*n^2 +108179055193*n -215961878286) *a(n-4) +(1708832970*n^3 -29554327949*n^2 +137453332457*n -152801514054) *a(n-5) +3*(569610990*n^2 -3742686463*n +4740040723) *a(n-6)=0. - _R. J. Mathar_, Nov 02 2015 Conjecture: (241*n-1066) *(2*n-11) *(-5+n)^2 *a(n) +(-482*n^5 +10099*n^4 -79756*n^3 +285961*n^2 -426904*n +149292) *a(n-1) -(2*n-9) *(n-3) *(248*n^3 -2229*n^2 +5065*n -7134) *a(n-2) +(-14*n^5 -49*n^4 -619*n^3 +13174*n^2 -51690*n +61248) *a(n-3) -(n-3) *(n-4) *(7*n-87) *(2*n-7) *a(n-4)= 0. - _R. J. Mathar_, Nov 02 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000350 | OEIS-A000188

(1) Number of solutions to x^2 == 0 (mod n). (2) Also square root of largest square dividing n. (3) Also max_{ d divides n } gcd(d, n/d).

Conjecture: a(n) = Sum_{k=1..n} A010052(n*k). - _Velin Yanev_, Jul 04 2021 Yanev's conjecture is true (see Fried link). - _Sela Fried_, Dec 01 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000351 | OEIS-A000195

a(n) = floor(log(n)).

The conjectured formula is true, as can be seen from a series expansion in x = log(n) which gives [Sum_{k>=0} x^(k+1)/((k+1)!*(3*exp(2*x))^k)] for the R.H.S., with [.] = floor. Considering n = [e^m] where 1-{x} is minimal ({x} = fractional part of x), results on the irrationality measure of powers of e give an O(1/n) decrease for 1-{x}, while the tail of the sum decreases like O((log n)^2/n^2). - _M. F. Hasler_, Aug 31 2026 Conjecture: a(n) = floor(3*n^2*(n^(1/(3*n^2))-1)), checked for n <= 10^6. - _Joseph M. Shunia_, Aug 02 2024 [Proved, see comment above.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000352 | OEIS-A000204

Lucas numbers (beginning with 1): L(n) = L(n-1) + L(n-2) with L(1) = 1, L(2) = 3.

For n>=3, a(n) = 4*A054886(n-1) - A022388(n-3) - 2*A022120(n-2) (conjectured by _Creighton Dement_, Nov 27 2004; edited and proved by _Sela Fried_, Dec 01 2025. See Fried link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000353 | OEIS-A000212

a(n) = floor(n^2/3).

Conjecture: a(n) is the number of intersection points of n cevians that cut a triangle into the maximum number of pieces (see A007980). - _Anton Zakharov_, May 07 2017 Zakharov's conjecture is true (see Fried link). - _Sela Fried_, Sep 25 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000354 | OEIS-A000215

Fermat numbers: a(n) = 2^(2^n) + 1.

It is conjectured that just the first 5 numbers in this sequence are primes. Conjecture: let the smallest prime factor of Fermat number F(n) be P(F(n)). If F(n) is composite, then P(F(n)) < 3*2^(2^n/2 - n - 2). - _Arkadiusz Wesolowski_, Aug 10 2012 Conjecture: the numbers > 5 in this sequence, i.e., 2^2^k + 1 for k>1, are exactly the numbers n such that (n-1)^4-1 divides 2^(n-1)-1. - _M. F. Hasler_, Jul 24 2015 Layman's conjecture is true. Additionally, it is true that if n > 5 is a Fermat number, then (n-1)^4-1 divides 2^(n-1)-1 (confirming one implication of Hasler's conjecture). It is not true that if phi(F-2) = (F-1)/2, then F is a Fermat prime (disproving one implication of Cloitre's conjecture). See Fried link. - _Sela Fried_, Oct 31 2025 Conjecture: F is a Fermat prime if and only if phi(F-2) = (F-1)/2. - _Benoit Cloitre_, Sep 15 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000355 | OEIS-A000217

Triangular numbers: a(n) = binomial(n+1,2) = n*(n+1)/2 = 0 + 1 + 2 + ... + n.

Conjecturally, 1, 6, 120 are the only numbers that are both triangular and factorial. - Christopher M. Tomaszewski (cmt1288(AT)comcast.net), Mar 30 2005 Conjecture: For n > 0, there is always a prime between A000217(n) and A000217(n+1). Sequence A065383 has the first 1000 of these primes. - _Ivan N. Ianakiev_, Mar 11 2013 Conjecture: the sequence is the genus/deficiency of the sinusoidal spirals of index n which are algebraic curves. The value 0 corresponds to the case of the Bernoulli Lemniscate n=2. So the formula conjectured is (n-1)(n-2)/2. - _Wolfgang Tintemann_, Aug 02 2015 Conjecture: Let m be any positive integer. Then, for each n = 1,2,3,... the set {Sum_{k=s..t} 1/k^m: 1 <= s <= t <= n} has cardinality a(n) = n*(n+1)/2; in other words, all the sums Sum_{k=s..t} 1/k^m with 1 <= s <= t are pairwise distinct. (I have checked this conjecture via a computer and found no counterexample.) - _Zhi-Wei Sun_, Sep 09 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000356 | OEIS-A000225

a(n) = 2^n - 1. (Sometimes called Mersenne numbers, although that name is usually reserved for A001348.)

Conjecture: except for a(2)=3, numbers m such that 2^(m+1) - 2^j - 2^k - 1 is composite for all 0 <= j < k <= m. - _Chai Wah Wu_, Sep 08 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000357 | OEIS-A000241

Crossing number of complete graph with n nodes.

It was conjectured by A. Hill in 1958 (see Guy 1960 and Harary-Hill 1963) that a(n) = floor(n/2)*floor((n-1)/2)*floor((n-2)/2)*floor((n-3)/2)/4 (see A028723). This is also sometimes referred to as Guy's conjecture. - _N. J. A. Sloane_, Jan 21 2015 Another version of the conjecture is that a(n)=C(floor(n/2),2)*C(floor((n-1)/2),2). We conjecture that this sequence is also given by one half of the third coefficient of the denominator polynomial of the n-th convergent to the g.f. of n!. From the Pan and Richter reference: 0.8594 Z(n) <= a(n) <= Z(n), where Z(n) is the conjectured formula (Richter and Thomassen 1997, de Klerk et al. 2007). - _Danny Rorabaugh_, Mar 12 2015 Empirical g.f.: -x^5*(1+x+x^2)/(x+1)^3/(x-1)^5, which is the same as the conjectured formula of A. Hill. - _Simon Plouffe_, Feb 06 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000358 | OEIS-A000246

Number of permutations in the symmetric group S_n that have odd order.

This conjecture has been confirmed, see Bernardi, Duplantier, Nadeau link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000359 | OEIS-A000247

a(n) = 2^n - n - 2.

(Conjecture) Let a(2)=0. For n > 2, let N = 2*n + 1. For each n, define the n X n tridiagonal unit-primitive matrix (see [Jeffery]) A_{N,1}=[0,1,0,...,0; 1,0,1,0,...,0; 0,1,0,1,0,...,0; ...; 0,...,0,1,0,1; 0,...,0,1,1] associated with N. Define the n-dimensional column vectors V_N = [v_1,v_2,...,v_n]^T = [A_{N,1}]^n*[1,1,...,1]^T, where [.]^T denotes matrix transpose and [1,...,1] is the n-dimensional unit vector. Let (v_k)_N denote the k-th element of V_N, k in {1,...,n}. Then a(n) = (v_(n-2))_N. - _L. Edson Jeffery_, Jan 20 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000360 | OEIS-A000254

Unsigned Stirling numbers of first kind, s(n+1,2): a(n+1) = (n+1)*a(n) + n!.

Detlefs' conjectures above and below are true (see Fried link). - _Sela Fried_, Oct 26 2025 The following are all conjectural:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000361 | OEIS-A000262

Number of "sets of lists": number of partitions of {1,...,n} into any number of lists, where a list means an ordered subset.

We conjecture that for k = 2,3,4,..., the difference a(n+k) - a(n) is divisible by k: if true, then for each k the sequence a(n) taken modulo k is periodic with period dividing k. - _Peter Bala_, Nov 14 2017 The above conjecture is true - see

the Bala link. - _Peter Bala_, Jan 20 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000362 | OEIS-A000265

Remove all factors of 2 from n; or largest odd divisor of n; or odd part of n.

a(n) = Sum_{k=0..n} A127793(n,k)*floor((k+2)/2) (conjecture). - _Paul Barry_, Jan 29 2007 Barry's conjecture is true (see Fried link in A127793). - _Sela Fried_, Nov 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000363 | OEIS-A000267

Integer part of square root of 4n+1.

a(n) = A070939(A227368(n+1)), conjectured. - _Antti Karttunen_, Dec 28 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000364 | OEIS-A000274

Number of permutations of length n with 2 consecutive ascending pairs.

Mathar's conjecture is true and the conjectured formula a(n) = (1/2)*([n!/e] - [(n-1)!/e]) is true (see Fried link). - _Sela Fried_, Sep 27 2025 a(n) = (1/2)*([n!/e] - [(n-1)!/e]) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000365 | OEIS-A000275

Coefficients of a Bessel function (reciprocal of J_0(z)); also pairs of permutations with rise/rise forbidden.

Conjectural formula: 1 = Sum_{n>=0} a(n)*x^n*Sum_{k>=0} binomial(n+k,k)^2*(-x)^k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000366 | OEIS-A000312

a(n) = n^n; number of labeled mappings from n points to themselves (endofunctions).

Conjecture: a(n) = Sum_{i=0..n} A048994(n, i) * A048993(n+i, n) for n >= 0; proved by Mike Earnest, see link at A354797. - _Werner Schulte_, Jun 19 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000367 | OEIS-A000330

Square pyramidal numbers: a(n) = 0^2 + 1^2 + 2^2 + ... + n^2 = n*(n+1)*(2*n+1)/6.

Conjecture: Every integer can be represented as a sum of three generalized square pyramidal numbers. A related conjecture is given in A336205 corresponding to pentagonal case. A stronger version of these conjectures is that every integer can be expressed as a sum of three generalized r-gonal pyramidal numbers for all r >= 3. In here "generalized" means negative indices are included. - _Altug Alkan_, Jul 30 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000368 | OEIS-A000332

Binomial coefficient binomial(n,4) = n*(n-1)*(n-2)*(n-3)/24.

... (as a conjectured infinite set). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000369 | OEIS-A000350

Numbers m such that Fibonacci(m) ends with m.

Conjecture: Other than 1 and 5, there is no m such that Fibonacci(m) in binary ends with m in binary. The conjecture holds up to m=50000. - _Ralf Stephan_, Aug 21 2006 The conjecture for binary numbers holds for m < 2^25. - _T. D. Noe_, May 14 2007 Conjecture is true. It is easy to prove (by induction on k) that if Fibonacci(m) ends with m in binary, then m == 0, 1, or 5 (mod 3*2^k) for any positive integer k, i.e., m must simply be equal to 0, 1, or 5. - _Max Alekseyev_, Jul 03 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000370 | OEIS-A000364

Euler (or secant or "Zig") numbers: e.g.f. (even powers only) sec(x) = 1/cos(x).

Conjecture: taking the sequence [a(n) : n >= 1] modulo an integer k gives a purely periodic sequence with period dividing phi(k). For example, the sequence taken modulo 21 begins [1, 5, 19, 20, 16, 2, 1, 5, 19, 20, 16, 2, 1, 5, 19, 20, 16, 2, 1, 5, 19, ...] with an apparent period of 6 = phi(21)/2. - _Peter Bala_, May 08 2023 It appears that a(n) = 3*A076552(n -1) + 2*(-1)^n for n >= 1. Conjectural congruences: a(2*n) == 5 (mod 60) for n >= 1 and a(2*n+1) == 1 (mod 60) for n >= 0. - _Peter Bala_, Jul 26 2013 Peter Bala's conjectured congruences, a(2n) == 5 (mod 60) for n >= 1 and a(2n + 1) == 1 (mod 60), hold due to the results of Stern (mod 4) and Knuth & Buckholtz (mod 3 and 5). - _Charles R Greathouse IV_, Mar 23 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000371 | OEIS-A000373

Conjectured dimension of a module associated with the free commutative Moufang loop with n generators.

Conjectured dimension of a module associated with the free commutative Moufang loop with n generators. There is an explicit formula for the n-th term of this sequence (see Eq. (8.4) of Smith (1982)). It is conjectured that this gives the answer to a question of Manin about the dimension of a certain module associated with the free commutative Moufang loop with n generators. - _N. J. A. Sloane_, May 21 2014 The underlying hypothesis has been disproven, see Grishkov & Shestakov (2008-2011). - _Matthew House_, Sep 03 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000372 | OEIS-A000392

Stirling numbers of second kind S(n,3).

Conjecture. Let S(1)={1} and, for n > 1, let S(n) be the smallest set containing x, 2x and 3x for each element x in S(n-1). Then a(n+2) is the sum of the elements in S(n). (It is easy to prove that the number of elements in S(n) is the n-th triangular number given by A001952.) See A122554 for a sequence defined in this way. - _John W. Layman_, Nov 21 2007; corrected (a(n) to a(n+2) due to offset change) by _Fred Daniel Kline_, Oct 02 2014 John W. Layman's conjecture that a(n+2) is the sum of elements in S(n) follows from the identification of S(n) with the first n rows of A036561, whose row sums are A001047. - _Fred Daniel Kline_, Oct 02 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000373 | OEIS-A000425

Coefficients of ménage hit polynomials.

Conjecture: (n-1)*(n-3)*a(n) -n*(n-2)*(n-3)*a(n-1) -n*(n-1)*(n-3)*a(n-2) -n *(n-1)*a(n-3)=0. - _R. J. Mathar_, Nov 02 2015 Conjecture: a(n) = 2*n*A000271(n). - _R. J. Mathar_, Nov 02 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000374 | OEIS-A000459

Number of multiset permutations of {1, 1, 2, 2, ..., n, n} with no fixed points.

Conjecture: a(n+k) - a(n) is divisible by k. - _Mark van Hoeij_, Nov 15 2023 [Proved in Kallat (2006) (Theorem 62). - _Ahaan Kallat_, Oct 14 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000375 | OEIS-A000464

Expansion of e.g.f. sin(x)/cos(2*x).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000376 | OEIS-A000483

Associated Stirling numbers: second-order reciprocal Stirling numbers (Fekete) a(n) = [[n, 3]]. The number of 3-orbit permutations of an n-set with at least 2 elements in each orbit.

Conjecture: (n-2)*(n-4)*a(n) -(n-1)*(3*n^2-21*n+35)*a(n-1) +(n-1)*(n-2)*(3*n^2-24*n+47)*a(n-2) -(n-5)*(n-1)*(n-2)*(n-3)*(n-4)*a(n-3)=0. - _R. J. Mathar_, Jul 18 2015 Conjecture: 3*(-n+4)*a(n) +(9*n^2-59*n+90)*a(n-1) +(-9*n^3+96*n^2-348*n+436)*a(n-2) +(n-3)*(3*n^3-45*n^2+237*n-430)*a(n-3) +5*(n-5)*(n-6)*(n-3)*(n-4)*a(n-4)=0. - _R. J. Mathar_, Jul 18 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000377 | OEIS-A000521

Coefficients of modular function j as power series in q = e^(2 Pi i t). Another name is the elliptic modular invariant J(tau).

It is an open question if there are infinitely many k such that a(k) is prime. The known such indices are listed in A339429. See the paper by Fredrik Johansson. - _Peter Luschny_, May 05 2021 Conjecture: a(n) with n >= 0 can never be a perfect power. Also, the sequence (a(n+1)/a(n))_{n>=0} is strictly decreasing (to the limit 1). - _Zhi-Wei Sun_, Dec 27 2024 Conjecture: Sum_{k>=-1} a(k) / exp(k*Pi) = 66^3 = 287496. - _Simon Plouffe_, Apr 01 2026 The above conjecture is correct. Note that the LHS is the expansion of j(I/2), and the section "Definition" of the Wikipedia page on the j-invariant contains the example j(2*I) = 66^3. Since the j-invariant is an automorphism form for SL(2,Z), we have j(I/2) = j(-1/(2*I)) = j(2*I) = 66^3. - _Jianing Song_, Apr 08 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000378 | OEIS-A000522

Total number of ordered k-tuples (k=0..n) of distinct elements from an n-element set: a(n) = Sum_{k=0..n} n!/k!.

G.f.: conjecture: T(0)/(1-2*x), where T(k) = 1 - x^2*(k+1)^2/(x^2*(k+1)^2 - (1 - 2*x*(k+1))*(1 - 2*x*(k+2))/T(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 18 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000379 | OEIS-A000531

From area of cyclic polygon of 2n + 1 sides.

A conjectured definition: Let 0 < a_1 < a_2 <...<a_{2n} < 1. Then how many ways are there in which one can add or subtract all the a_i to get an odd number. For example, take n = 2. Then the options are a_1+a_2+a_3+a_4 = 1 or 3; one can change the sign of any of the a_i's and get 1; or -a_1-a_2+a_3+a_4 = 1. That's a total of 7, which is the 2nd number of this sequence. One of the definitions of the sequence (which was how I came across it) is the degree of the equation giving the area of a cyclic polygon in terms of the sides. I conjectured that for any set of side lengths there is a unique way of fitting them together for any possible winding number and any possible subset of sides which go round the circle in a retrograde manner. - Simon Norton (simon(AT)dpmms.cam.ac.uk), May 14 2001 a(n) is also the number of walks of length 2n in the quarter plane starting and ending at the origin using steps {(1,1),(1,0),(-1,0),(-1,-1)} (which appear in Gessel's conjecture) in which the steps (1,0) and (-1,0) appear exactly once each. - _Arvind Ayyer_, Mar 02 2009 (Conjecture) a(n)=2^(2*n)*sum_{k=1..n} cos(k*Pi/(2*n+1))^2*n. - _L. Edson Jeffery_, Jan 21 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000380 | OEIS-A000579

Figurate numbers or binomial coefficients C(n,6).

Conjecture: a(n+3) = Sum_{0 <= k, L, m <= n; k + L + m <= n} k*L*m. - _Ralf Stephan_, May 06 2005 The conjecture is true (see Fried link). - _Sela Fried_, Nov 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000381 | OEIS-A000629

Number of necklaces of partitions of n+1 labeled beads.

Row sums of A154921 as conjectured above by Granvik. a(n) gives the number of outcomes of a race between n horses H1,...,Hn, where if a horse falls it is not ranked. For example, when n = 2 the 6 outcomes are a dead heat, H1 wins H2 second, H2 wins H1 second, H1 wins H2 falls, H2 wins H1 falls or both fall. - _Peter Bala_, May 15 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Don Knuth_, Nick Singer (nsinger(AT)eos.hitc.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000382 | OEIS-A000695

Moser-de Bruijn sequence: sums of distinct powers of 4.

Conjecture: The sums of distinct powers of k > 2 can be constructed as the following (k-1)-ary rooted tree. For each n the tree grows and a(n) is then the total number of nodes. For n = 1, the root of the tree is added. For n > 1, if n is odd one leaf of depth n-2 grows one child. If n is even all leaves of depth >= (n - 1 - A000225(A001511(n/2))) grow the maximum number of children. An illustration is provided in the links. - _John Tyler Rascoe_, Oct 09 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000383 | OEIS-A000698

A problem of configurations: a(0) = 1; for n>0, a(n) = (2n-1)!! - Sum_{k=1..n-1} (2k-1)!! a(n-k). Also the number of shellings of an n-cube, divided by 2^n n!.

The o.g.f. B(x) = Sum_{n>=1} a(n)*x^(2*n-1) = x + 2*x^3 + 10*x^5 + 74*x^7 + ... satisfies the Riccati differential equation y'(x) = -1/x^2 + (1/x^3)*y(x) - (1/x^2)*y(x)^2 with initial condition y(0) = 0 (cf. A005412). The solution is B(x) = 1/z(x) + 1/x, where z(x) = -Sum_{n>=0} A001147(n) * x^(2*n+1) = -(x + x^3 + 3*x^5 + 15*x^7 + ...). The function b(x) = -B(1/x) satisfies b'(x) = -1 - (x + b(x))*b(x). Hence the differential operator (D^2 + x*D + 1), where D = d/dx, factorizes as (D - a(x))*(D - b(x)), where a(x) = -(x + b(x)), as conjectured by [Edgar, Problem 4.32]. For a refinement of this sequence see A053979. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Richard Ehrenborg_, _Gabor Hetyei_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000384 | OEIS-A000700

Expansion of Product_{k>=0} (1 + x^(2k+1)); number of partitions of n into distinct odd parts; number of self-conjugate partitions; number of symmetric Ferrers graphs with n nodes.

Conjecture: Sum_{k>=0} a(k) / exp(k*Pi) = 2^(1/4) * exp(-Pi/24) = A292828. - _Simon Plouffe_, Sep 09 2025 See the Bala link for a proof of the above conjecture. - _Peter Bala_, May 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000385 | OEIS-A000704

Number of degree-n even permutations of order dividing 2.

Conjecture: a(n) -3*a(n-1) +3*a(n-2) -a(n-3) -(n-1)*(n-3)*a(n-4) +(n-3)*(n-4)*a(n-5)=0. - _R. J. Mathar_, Jun 03 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _J. H. Conway_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000386 | OEIS-A000727

Expansion of Product_{k >= 1} (1 - x^k)^4.

Ramanujan (see the link, pp. 155 and 157 Nr. 23.) conjectured the expansion coefficients called Psi_4(n) of eta^4(6*z) in powers of q = exp(2*Pi*i*z), Im(z) > 0, where i is the imaginary unit. In the Finch link on p. 5, multiplicity is used and Psi_4(p^r), called f(p^r), is given (see also b(p^e) formula given by Michael Somos, Aug 23 2006). Mordell proved this conjecture on pp. 121-122 based on Klein-Fricke, Theorie der elliptischen Modulfunktionen, 1892, Band II, p. 374. The product formula for the Dirichlet series, Mordell, eq. (7) for a=2,is used to find Psi_4(n), called f_2(n), from f_2(p) for primes p. The primes p = 2 and 3 do not appear in the product. - _Wolfdieter Lang_, May 03 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000387 | OEIS-A000769

No-3-in-line problem: number of inequivalent ways of placing 2n points on an n X n grid so that no 3 are in a line.

It is conjectured that a(n)=0 for all sufficiently large n. In Canad. Math. Bull. 11 (1968) 527-531, MR 39 #129, Guy & Kelly conjecture that, for large n, at most (c + eps)*n points can be selected, where 3*c^3 = 2*Pi^2, i.e., c ~ 1.87.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000388 | OEIS-A000781

a(n) = 3*Catalan(n) - Catalan(n-1) - 1.

Conjecture: 3*(n+1)*a(n) + (-19*n+7)*a(n-1) + (33*n-53)*a(n-2) + (-21*n+61)*a(n-3) + 2*(2*n-9)*a(n-4) = 0. - _R. J. Mathar_, Aug 20 2013 The conjecture of _R. J. Mathar_ is true. Since a(n) = 3*A000108(n) - A000108(n-1) - 1 and (n+1)*A000108(n) = 2*(2*n-1)*A000108(n-1), substituting into the recurrence gives zero for the Catalan part, and the coefficients sum to 0, which disposes of the constant term. - _Roberto Blanco Gómez_, Oct 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000389 | OEIS-A000797

Numbers that are not the sum of 4 tetrahedral numbers.

It is an open problem of long standing ("Pollock's Conjecture") to show that this sequence is finite. More precisely, Salzer and Levine conjecture that every number is the sum of at most 5 tetrahedral numbers and in fact that there are exactly 241 numbers (the terms of this sequence) that require 5 tetrahedral numbers, the largest of which is 343867.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000390 | OEIS-A000807

Quadratic invariants.

Conjecture: a(n) = 2*exp(-2) * Sum_{k >= 1} k^(2*n)*BesselI(k, 2) for n > 0. - _Velin Yanev_, Aug 29 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000391 | OEIS-A000907

Second-order reciprocal Stirling number (Fekete) a(n) = [[2n+2, n]]. The number of n-orbit permutations of a (2n+2)-set with at least 2 elements in each orbit. Also known as associated Stirling numbers of the first kind (e.g., Comtet).

Conjecture: n*(4*n+5)*a(n) -(2*n+3)*(n+2)*(4*n+9)*a(n-1)=0. - _R. J. Mathar_, Apr 30 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000392 | OEIS-A000918

a(n) = 2^n - 2.

For n >= 1, conjectured to be the number of integers from 0 to (10^n)-1 that lack 0, 1, 2, 3, 4, 5, 6 and 7 as a digit. - _Alexandre Wajnberg_, Apr 25 2005 Conjecture: For n>0, a(n) is the least m such that A007814(A000108(m)) = n-1. - _L. Edson Jeffery_, Nov 27 2015 Actually this follows from the procedure for determining the multiplicity of prime p in C(n) given in A000108 by _Franklin T. Adams-Watters_: For p=2, the multiplicity is the number of 1 digits minus 1 in the binary representation of n+1. Obviously, the smallest k achieving "number of 1 digits" = k is 2^k-1. Therefore C(2^k-2) is divisible by 2^(k-1) for k > 0 and there is no smaller m for which 2^(k-1) divides C(m) proving the conjecture. - _Peter Schorn_, Feb 16 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000393 | OEIS-A000926

Euler's "numerus idoneus" (or "numeri idonei", or idoneal, or suitable, or convenient numbers).

It is conjectured that the list given here is complete. Chowla showed that the list is finite and Weinberger showed that there are at most two further terms. See Corollary 23 and Remark 24 of Kani (2011). [Statement corrected by _David Radcliffe_, Aug 22 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000394 | OEIS-A000928

Irregular primes: primes p such that at least one of the numerators of the Bernoulli numbers B_2, B_4, ..., B_{p-3} (A000367) is divisible by p.

"The pioneering mathematician Kummer, over the period 1847-1850, used his profound theory of cyclotomic fields to establish a certain class of primes called 'regular' primes. ... It is known that there exist an infinity of irregular primes; in fact it is a plausible conjecture that only an asymptotic fraction 1/Sqrt(e) ~ 0.6 of all primes are regular." [Ribenboim]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000395 | OEIS-A000946

Euclid-Mullin sequence: a(1) = 2, a(n+1) is the largest prime factor of 1 + Product_{k=1..n} a(k).

Booker's abstract claims: "We consider the second of Mullin's sequences of prime numbers related to Euclid's proof that there are infinitely many primes. We show in particular that it omits infinitely many primes, confirming a conjecture of Cox and van der Poorten."

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000396 | OEIS-A000952

Numbers k == 2 (mod 4) that are the orders of conference matrices.

If k == 2 (mod 4) then a necessary condition is that k-1 is a sum of 2 squares (A286636). It is conjectured that this condition is also sufficient. If k == 2 (mod 4) and k-1 is a prime or prime power the condition is automatically satisfied.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000397 | OEIS-A000966

n! never ends in this many 0's.

The g.f. conjectured by _Simon Plouffe_ in 1992 dissertation is not correct; the first discrepancy is a(31) = 155, his g.f. gives 160. In fact, the g.f. for this sequence is not rational; the first differences are bounded but not periodic. - _Franklin T. Adams-Watters_, Jul 03 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000398 | OEIS-A000974

Conjecturally the number of even integers the sum of two primes in exactly n ways.

Conjecturally the number of even integers the sum of two primes in exactly n ways.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bill Gosper_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000399 | OEIS-A000975

a(2n) = 2*a(2n-1), a(2n+1) = 2*a(2n)+1 (also a(n) is the n-th number without consecutive equal binary digits).

Conjecture: {a(n)} also gives all j for which A048702(j) = A000217(j); i.e., if we take the a(n)-th triangular number (a(n)^2 + a(n))/2 and multiply it by 3, we get a(n)-th even-length binary palindrome A048701(a(n)) concatenated from a(n) and its reverse. E.g., a(4) = 10, which is 1010 in binary; the tenth triangular number is 55, and 55*3 = 165 = 10100101 in binary. - _Antti Karttunen_, circa 1999. (This

has been now proved by Paul K. Stockmeyer in his arXiv:1608.08245 paper.) - _Antti Karttunen_, Aug 31 2016 Conjectured record values of A264784: a(n) = A264784(A155051(n-1)). - _Reinhard Zumkeller_, Dec 04 2015. (This is proved by Paul K. Stockmeyer in his arXiv:1608.08245 paper.) - _Antti Karttunen_, Aug 31 2016 Conjecture: a(n+1) is the number of compositions of n with two kinds of parts, n and n', where the order of the 1 and 1' does not matter. For n=2, a(3) = 5 compositions, enumerated as follows: 2; 2'; 1,1; 1',1 = 1',1; 1',1'. - _Gregory L. Simay_, Sep 02 2017 Conjecture proved by recognizing the appropriate g.f. is x/((1 - x)*(1 - x)*(1 - 2*x^2 - 2*x^3 - ...)) = x/(1 - 2*x - x^2 + 2*x^3). - _Gregory L. Simay_, Sep 10 2017 Conjecture: For n >= 3, a(n) is the second largest natural number whose Collatz orbit has length n+3. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mira Bernstein_, _N. J. A. Sloane_, _Robert G. Wilson v_, Sep 13 1996 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000400 | OEIS-A000978

Wagstaff numbers: numbers k such that (2^k + 1)/3 is prime.

There is a new conjecture stating that a Wagstaff number is prime under the following condition (based on DiGraph cycles under the LLT): Let p be a prime integer > 3, N(p) = 2^p+1 and W(p) = N(p)/3, S(0) = 3/2 (or 1/4) and S(i+1) = S(i)^2 - 2 (mod N(p)). Then W(p) is prime iff S(p-1) == S(0) (mod W(p)). - _Tony Reix_, Sep 03 2007 Conjecture: a(1) = 3; and then, for n > 0, a(n+1) is the least odd k such that lpf((2^k + 1)/3) > lpf((2^a(n) + 1)/3), where lpf(w) is the least prime factor of w. - _Thomas Ordowski_, Sep 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000401 | OEIS-A000984

Central binomial coefficients: binomial(2*n,n) = (2*n)!/(n!)^2.

Erdős, Graham et al. conjectured that a(n) is never squarefree for sufficiently large n (cf. Graham, Knuth, Patashnik, Concrete Math., 2nd ed., Exercise 112). Sárközy showed that if s(n) is the square part of a(n), then s(n) is asymptotically (sqrt(2)-2) * (sqrt(n)) * zeta(1/2). Granville and Ramare proved that the only squarefree values are a(1)=2, a(2)=6 and a(4)=70. - _Jonathan Vos Post_, Dec 04 2004 [For more about this conjecture, see A261009. - _N. J. A. Sloane_, Oct 25 2015] The MathOverflow link contains the following comment (slightly edited): The Erdős squarefree conjecture (that a(n) is never squarefree for n>4) was proved in 1980 by Sárközy, A. (On divisors of binomial coefficients. I. J. Number Theory 20 (1985), no. 1, 70-80.) who showed that the conjecture holds for all sufficiently large values of n, and by A. Granville and O. Ramaré (Explicit bounds on exponential sums and the scarcity of squarefree binomial coefficients. Mathematika 43 (1996), no. 1, 73-107) who showed that it holds for all n>4. - Fedor Petrov, Nov 13 2010. [From _N. J. A. Sloane_, Oct 29 2015] Straub, Amdeberhan and Moll: "... it is conjectured that there are only finitely many indices n such that C_n is not divisible by any of 3, 5, 7 and 11." - _Jonathan Vos Post_, Nov 14 2008 Conjecture: For any positive integer n, the polynomial Sum_{k=0}^n a(k)x^k is irreducible over the field of rational numbers. In general, for any integer m>1 and n>0, the polynomial f_{m,n}(x) = Sum_{k=0..n} (m*k)!/(k!)^m*x^k is irreducible over the field of rational numbers. - _Zhi-Wei Sun_, Mar 23 2013 Number of vertices of the subgraph of the (2n-1)-dimensional hypercube induced by all bitstrings with n-1 or n many 1s. The middle levels conjecture asserts that this graph has a Hamilton cycle. - _Torsten Muetze_, Feb 11 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000402 | OEIS-A000986

Number of n X n symmetric matrices with (0,1) entries and all row sums 2.

Conjecture: 2*a(n) +2*(-2*n+1)*a(n-1) +2*(n^2-2*n-1)*a(n-2) -2*(n-2)*(n-4)*a(n-3) +(n-1)*(n-2)*(n-3)*a(n-4) -(n-2)*(n-3)*(n-4)*a(n-5)=0. - _R. J. Mathar_, Aug 04 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000403 | OEIS-A001006

Motzkin numbers: number of ways of drawing any number of nonintersecting chords joining n (labeled) points on a circle.

Conjecture: (2/n)*Sum_{k=1..n} (2k+1)*a(k)^2 is an integer for each positive integer n. - _Zhi-Wei Sun_, Nov 16 2017 The Rubey and Stump reference proves a refinement of a conjecture of René Marczinzik, which they state as: "The number of 2-Gorenstein algebras which are Nakayama algebras with n simple modules and have an oriented line as associated quiver equals the number of Motzkin paths of length n." - _Eric M. Schmidt_, Dec 16 2017 Conjecture: If p is a prime of the form 6m+1 (A002476), then a(p-2) is divisible by p. Currently, no counterexample exists for p < 10^7. Personal communication from _Robert Gerbicz_: mod such p this is equivalent to A066796 with comment: "Every A066796(n) from A066796((p-1)/2) to A066796(p-1) is divisible by prime p of form 6m+1". - _Serge Batalov_, Feb 08 2022 The conjecture is proved in the 2017 paper by Rob Burns in the Links below. The result is contained in Tables 4 and 5 of the paper, which show that a(p-2) == 0 (mod p) when p == 1 (mod 6) and a(p-2) == -1 (mod p) when p == -1 (mod 6). The 2024 paper by Nadav Kohen also proves the conjecture. Proposition 6 of the paper states that a prime p divides a(p-2) if and only if p = (1 mod 3). (End) Conjectures: (1) For prime p == 1 (mod 6) and n, r >= 1, a(n*p^r - 2) == -A005717(n-1) (mod p), where we take A005717(0) = 0 to match Batalov's conjecture above. Let b(n) = 1 for n <= 1, otherwise b(n) = Sum_{k=2..n} b(k-1) * b(n-k), then a(n) = b(n+1) (conjecture). - _Joerg Arndt_, Jan 16 2023 Arndt's conjecture is true (see Fried link). - _Sela Fried_, Dec 01 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000404 | OEIS-A001013

Jordan-Polya numbers: products of factorial numbers A000142.

Except for the numbers 2, 9 and 10 this sequence is conjectured to be the same as A034878.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000405 | OEIS-A001030

Fixed under 1 -> 21, 2 -> 211.

a(n) = -1 + floor(n*(1+sqrt(2))+1/sqrt(2))-floor((n-1)*(1+sqrt(2))+1/sqrt(2)). - _Benoit Cloitre_, Jun 26 2004. [I don't know if this is a theorem or a conjecture. - _N. J. A. Sloane_, May 14 2008]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000406 | OEIS-A001032

Numbers k such that sum of squares of k consecutive integers >= 1 is a square.

These conditions can be used to establish the conjecture of Ralf Stephan, above, that all the terms are == 0, 1, 2, 9, 11, 16, or 23 (mod 24). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000407 | OEIS-A001039

a(n) = (p^p-1)/(p-1) where p = prime(n).

Let r be a root of the trinomial x^p-x-1 in a fixed algebraic closure F of the finite field F_p. Radoux conjectured in 1975 (see References) that a(n) equals the multiplicative order of r in F. The conjecture seems still open.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000408 | OEIS-A001040

a(n+1) = n*a(n) + a(n-1) with a(0)=0, a(1)=1.

Conjecture: 2*n!*a(n) is the number of open tours by a rook on an (n X 2) chessboard which ends at the opposite line of length n. - _Mikhail Kurkov_, Nov 19 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000409 | OEIS-A001043

Numbers that are the sum of 2 successive primes.

Conjecture: For any positive even number m, each positive rational number can be written as a sum of finitely many distinct unit fractions whose denominators are sums of m consecutive primes. If m is an odd number greater than one, then for any positive integers c and d with d odd, we can write c/d as a sum of finitely many distinct unit fractions whose denominators are sums of m consecutive primes. - _Zhi-Wei Sun_, Nov 03 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000410 | OEIS-A001045

Jacobsthal sequence (or Jacobsthal numbers): a(n) = a(n-1) + 2*a(n-2), with a(0) = 0, a(1) = 1; also a(n) = nearest integer to 2^n/3.

Copeland's conjecture is true (see Fried link). - _Sela Fried_, Dec 03 2025 Fossaceca's conjecture is true (see Fried link). - _Sela Fried_, Nov 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000411 | OEIS-A001053

a(n+1) = n*a(n) + a(n-1) with a(0)=1, a(1)=0.

Conjecture: 2*n!*a(n) is the number of open tours by a rook on an (n X 2) chessboard which starts and ends at the same line of length n. - _Mikhail Kurkov_, Nov 19 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000412 | OEIS-A001067

Numerator of Bernoulli(2*n)/(2*n).

Ramanujan incorrectly conjectured that the sequence contains only primes (and 1). - _Jud McCranie_. See A112548, A119766. Conjecture: If there is a prime p such that 2*n+1 < p and p divides a(n), then p^2 does not divide a(n). This conjecture is true for p < 12 million. - _Seiichi Manyama_, Jan 21 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Richard E. Borcherds (reb(AT)math.berkeley.edu) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000413 | OEIS-A001076

Denominators of continued fraction convergents to sqrt(5).

Conjecture: 2a(n+1) = a(n+2) - A001077(n+1). - _Creighton Dement_, Nov 28 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000414 | OEIS-A001083

Length of one version of Kolakoski sequence {A000002(i)} at n-th growth stage.

Conjecture: a(n) is asymptotic to c*(3/2)^n where c=0.5819.... - _Benoit Cloitre_, Jun 01 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000415 | OEIS-A001089

Number of permutations of [n] containing exactly 2 increasing subsequences of length 3.

Noonan and Zeilberger conjectured that a(n) = ((59*n^2+117*n+100) /(2*n*(2*n-1)*(n+5))) *binomial(2*n,n-4). This was proved by Fulmek.

猜想/问题提及段 | historical_or_current_mention_unreviewed | John Thomas Noonan [noonan(AT)euclid.math.temple.edu] | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000416 | OEIS-A001101

Moran numbers: k such that k/(sum of digits of k) is prime.

Witno conjectures that a(n) ~ c*n log(n)^2 for some c. - _Charles R Greathouse IV_, Jul 26 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | Bill Moran (moran1(AT)llnl.gov) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000417 | OEIS-A001108

a(n)-th triangular number is a square: a(n+1) = 6*a(n) - a(n-1) + 2, with a(0) = 0, a(1) = 1.

Conjecture: For n>1 a(n) is the index of the first occurrence of -n in sequence A123737. - _Vaclav Kotesovec_, Jun 02 2015 The squares of NSW numbers (A008843) interleaved with twice squares from A084703, where A008843(n) = A002315(n)^2 and A084703(n) = A001542(n)^2. Conjecture: Also numbers n such that sigma(n) = A000203(n) and sigma(n-th triangular number) = A074285(n) are both odd numbers. - _Jaroslav Krizek_, Aug 05 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000418 | OEIS-A001110

Square triangular numbers: numbers that are both triangular and square.

Conjecture: No a(2^k), where k is a nonnegative integer, can be expressed as a sum of a positive square number and a positive triangular number. - _Ivan N. Ianakiev_, Sep 19 2012 For n=2k, k > 0, a(n) is divisible by 12 and is therefore abundant. I conjecture that for n=2k+1 a(n) is deficient [true for k up to 43 incl.]. - _Ivan N. Ianakiev_, Sep 30 2014 The conjecture is true for all k > 0 because: For n=2k+1, k > 0, a(n) is odd. If a(n) is a prime number, it is deficient; otherwise a(n) has one or two distinct prime factors and is therefore deficient again. So for n=2k+1, k > 0, a(n) is deficient. - _Muniru A Asiru_, Apr 13 2016 The conjecture of Sep 19 2012 is false, since a(16) = 98286503002057414584576 = 303363259914^2 + (111868096760*111868096761)/2. This term is A218273(11). For k = 0, 1, 2, 3 there is no such representation, so k = 4 gives the smallest counterexample. - _Clément Garrot_, Aug 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000419 | OEIS-A001156

Number of partitions of n into squares.

Conjecture (Jan Bohman, Carl-Erik Fröberg, Hans Riesel, 1979): a(n) ~ c * n^(-alfa) * exp(beta*n^(1/3)), where c = 1/18.79656, beta = 3.30716, alfa = 1.16022. - _Vaclav Kotesovec_, Aug 19 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000420 | OEIS-A001171

From least significant term in expansion of E(tr (X'*X)^n), X rectangular and Gaussian. Also number of types of sequential n-swap moves for traveling salesman problem.

Conjecture (for n>=5): (n+1)*a(n) = -(4*n-1)*a(n-1) + (5*n^3 - 16*n^2 + 13*n - 1)*a(n-2) + (10*n^3 - 68*n^2 + 150*n - 107)*a(n-3) - (n-4)*(n-2)^2*(2*n-7)^2*a(n-4). - _Vaclav Kotesovec_, Aug 07 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Mallows_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000421 | OEIS-A001176

Number of zeros in fundamental period of Fibonacci numbers mod n.

a(n) = 2 for n == 0, 3, 7, 8, 12, 15 (mod 20). a(p^e) = 1 if primes p == 11, 19 (mod 20); 4 if p == 13, 17 (mod 20). Conjecture: 1/6 of the primes congruent to 1 or 9 mod 40 satisfy a(p^e) = 1, 2/3 of them satisfy a(p^e) = 2 and 1/6 of them satisfy a(p^e) = 4; also, 1/2 of the primes congruent to 21 or 29 mod 40 satisfy a(p^e) = 1 and 1/2 of them satisfy a(p^e) = 4. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000422 | OEIS-A001177

Fibonacci entry points: a(n) = least k >= 1 such that n divides Fibonacci number F_k (=A000045(k)).

In the formula, the relation a(p^e) = p^(e-1)*a(p) is called Wall's conjecture, which has been verified for primes up to 10^14. See A060305. Primes for which this relation fails are called Wall-Sun-Sun primes. - _T. D. Noe_, Mar 03 2009 Conjecture: Sum_{d|n} phi(d)*a(d) = A232656(n). - _Logan J. Kleinwaks_, Oct 28 2017 a(1) = 1, a(2) = 3, a(4) = 6 and for e > 2, a(2^e) = 3*2^(e-2); a(5^e) = 5^e; and if p is an odd prime not 5, then a(p^e) = p^max(0, e-s)*a(p) where s = valuation(A000045(a(p)), p) (Wall's conjecture states that s = 1 for all p). If (m, n) = 1 then a(m*n) = lcm(a(m), a(n)). See Posamentier & Lahmann. - _Robert G. Wilson v_, Jul 07 2007; corrected by _Max Alekseyev_, Oct 19 2007, Jun 24 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000423 | OEIS-A001187

Number of connected labeled graphs with n nodes.

"Based on experimental data obtained using the software LattE [14] and the Online Encyclopedia of Integer Sequences [19], we make the following conjecture: Conjecture 11. For j >= 2, Vol(C_j) is equal to the number of labeled connected graphs on j - 1 vertices." [Beck et al., 2011]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000424 | OEIS-A001250

Number of alternating permutations of order n.

G.f.: conjecture: 2*T(0)/(1-x) -1, where T(k) = 1 - x^2*(k+1)*(k+2)/(x^2*(k+1)*(k+2) - 2*(1-x*(k+1))*(1-x*(k+2))/Q(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 19 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000425 | OEIS-A001277

Number of permutations of length n by rises.

Conjecture: a(n) = A177265(n) if n even, = A177265(n)-1 if n odd. - _R. J. Mathar_, Jun 27 2022 Conjecture: D-finite with recurrence a(n) -n*a(n-1) +(n-1)*a(n-3)=0. - _R. J. Mathar_,

Jul 01 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000426 | OEIS-A001292

Concatenations of cyclic permutations of initial positive integers.

In 1996, Kenichiro Kashihara conjectured that there is no prime power of an integer (A093771) belonging to this sequence (disregarding the trivial case 1); a direct search from 12 to a(100128) has confirmed the conjecture up to 10^1035. There are no perfect powers among terms t which are permutations of 123_...(m - 1)_m for m == {2, 3, 5, 6} (mod 9). This is since 10 == 1 (mod 9) and also (1 + 0) == 1 (mod 9), so digit position has no effect. Hence, t == A134804(m) (mod 9). Now, if m is such that A134804(m) = {3, 6}, there is a lone factor of 3, which is not a perfect power (indeed).

猜想/问题提及段 | historical_or_current_mention_unreviewed | R. Muller | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000427 | OEIS-A001317

Sierpiński's triangle (Pascal's triangle mod 2) converted to decimal.

G.f.: Product_{k>=0} (1 + z^(2^k) + (2*z)^(2^k)). - conjectured by _Shamil Shakirov_, proved by _Vladimir Shevelev_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000428 | OEIS-A001338

-1 + Sum (k-1)! C(n,k), k = 1..n for n > 0, a(0) = 1.

Conjecture: a(n) +(-n-1)*a(n-1) +2*(n-1)*a(n-2) +(-n+2)*a(n-3)=0. - _R. J. Mathar_, Feb 16 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000429 | OEIS-A001354

Coordination sequence for hyperbolic tessellation 3^7 (from triangle group (2,3,7)).

Conjecture: For k>1, the interlaced polynomials b(2*k-1) = a(k)/7 and b(2*k) = (a(k+1) - a(k)) / 7 are the Fibonacci numbers (A000045). - _Avi Friedlich_, May 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000430 | OEIS-A001358

Semiprimes (or biprimes): products of two primes.

Conjecture: a(n)/n ~ (log(n)/log(log(n)))*(1-(M/log(log(n)))) as n -> oo, where M is Mertens's constant (A077761). - _Alain Rocchelli_, Feb 02 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000431 | OEIS-A001370

Sum of digits of 2^n.

It is believed that a(n) ~ n*9*log_10(2)/2, but this is an open problem. - _N. J. A. Sloane_, Apr 21 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000432 | OEIS-A001387

Binary "look and say" sequence.

I conjecture that the ratio r(n) of the number of "1"s to the number of "0"s in a(n) converges to 5/3 (or some nearby limit). - _Joseph L. Pe_, Jan 31 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas L. York_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000433 | OEIS-A001405

a(n) = binomial(n, floor(n/2)).

Let C_n(R, H) denote the transition matrix from the ribbon basis to the homogeneous basis of the graded component of the algebra of noncommutative symmetric functions of order n. Letting I(2^(n-1)) denote the identity matrix of order 2^(n-1), it has been conjectured that the dimension of the kernel of C_n(R, H) - I(2^(n-1)) is always equal to a(n-1). - _John M. Campbell_, Mar 30 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000434 | OEIS-A001419

Number of n-celled polyominoes with holes.

Conjecture: Almost all polyominoes are holey. In other words, a(n)/A000105(n) tends to 1 for increasing n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000435 | OEIS-A001462

Golomb's sequence: a(n) is the number of times n occurs in the sequence, starting with a(1) = 1.

We can interpret this sequence as a triangle: start with 1; 2,2; 3,3; and proceed by letting the row sum of row m-1 be the number of elements of row m. The partial sums of the row sums give 1, 5, 11, 38, 272, ... Conjecture: this proceeds as Lionel Levine's sequence A014644. See also A113676. - _Floor van Lamoen_, Nov 06 2005 Conjecture: a(n) >= n^(phi-1) for all n. - _Jianing Song_, Aug 19 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000436 | OEIS-A001498

Triangle a(n,k) (n >= 0, 0 <= k <= n) of coefficients of Bessel polynomials y_n(x) (exponents in increasing order).

G.f.: conjecture: T(0)/(1-x), where T(k) = 1 - x*y*(k+1)/(x*y*(k+1) - (1-x)^2/T(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 13 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000437 | OEIS-A001518

Bessel polynomial y_n(3).

Conjecture: (Sum_{n >= 0} a(n)*x^n/n!)/(1 + Sum_{n >= 0} a(n)*x^(n+1)/(n+1)!) = Sum_{n >= 0} A011781(n)*x^n/n!. - _Peter Bala_, Dec 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000438 | OEIS-A001519

a(n) = 3*a(n-1) - a(n-2) for n >= 2, with a(0) = a(1) = 1.

Proof: (M_2)^2 = 3*M + 1_2 (with the 2 X 2 unit matrix 1_2) from the characteristic polynomial of M_2 (see a comment in A322602) and the Cayley-Hamilton theorem. The recurrence M^n = M*M^(n-1) leads to (M_n)^n = S(n, 3)*1_2 + S(n-a, 3)*(M - 3*1_2), for n >= 0, with S(n, 3) = F(2(n+1)) = A001906(n+1). Hence ((M_2)^n)[1,1] = S(n, 3) - 2*S(n-1, 3) = a(n) = F(2*n-1) = (1/(2*r+1))*r^(2*n-1)*(1 + (1/r^2)^(2*n-1)), with r = rho(5) = A001622 (golden ratio) (see the first Aug 31 2004 formula, using the recurrence of S(n, 3), and the _Michael Somos_ Oct 28 2002 formula). This proves a conjecture of _Gary W. Adamson_ in A322602.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000439 | OEIS-A001522

Number of n-stacks with strictly receding walls, or the number of Type A partitions of n in the sense of Auluck (1951).

Conjecture: Also the number of integer partitions y of n with a fixed point y(i) = i. These partitions are ranked by A352827. The conjecture is stated at A238395, but Resta tells me he may not have had a proof. The a(1) = 1 through a(8) = 10 partitions are: The above conjecture is true. See Section 4 of the Blecher-Knopfmacher paper in the Links section. - _Jeremy Lovejoy_, Sep 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000440 | OEIS-A001542

a(n) = 6*a(n-1) - a(n-2) for n > 1, a(0)=0 and a(1)=2.

Conjecture: Numbers k such that c/m < k for all natural a^2 + b^2 = c^2 (Pythagorean triples), a < b < c and a+b+c = m. Numbers which correspondingly minimize c/m are A002939. - _Lorraine Lee_, Jan 31 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000441 | OEIS-A001551

a(n) = 1^n + 2^n + 3^n + 4^n.

O.g.f.: 2*(2-5*x)*(1-5*x+5*x^2)/(Product_{j=1..4} (1-j*x)) (via Laplace transformation of the o.g.f., and partial fraction decomposition backwards). See the Maple Program for the o.g.f. conjecture by _Simon Plouffe_. This has now been proved. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000442 | OEIS-A001554

a(n) = 1^n + 2^n + ... + 7^n.

Conjectures for o.g.f.s for this type of sequences appear in the PhD thesis by _Simon Plouffe_. See A001552 for the reference. These conjectures are proved in a link given in A196837. - _Wolfdieter Lang_, Oct 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000443 | OEIS-A001555

a(n) = 1^n + 2^n + ... + 8^n.

Conjectures for o.g.f.s for this type of sequence appear in the PhD thesis by _Simon Plouffe_. See A001552 for the reference. These conjectures are proved in a link given in A196837. [_Wolfdieter Lang_, Oct 15 2011]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000444 | OEIS-A001556

a(n) = 1^n + 2^n + ... + 9^n.

Conjectures for o.g.f.s for this type of sequences appear in the PhD thesis by _Simon Plouffe_. See A001552 for the reference. These conjectures are proved in the link given in A196837. - _Wolfdieter Lang_, Oct 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000445 | OEIS-A001566

a(0) = 3; thereafter, a(n) = a(n-1)^2 - 2.

Starting with 7, the terms end with 7,47,07,47,07,..., of the form 8a+7 where a = 0,1,55,121771,... Conjecture: Every a is squarefree, every other a is divisible by 55, the a's are a subset of A046194, the heptagonal triangular numbers (the first, 2nd, 3rd, 6th, 11th, ?, ... terms). - _Gerald McGarvey_, Aug 08 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000446 | OEIS-A001567

Fermat pseudoprimes to base 2, also called Sarrus numbers or Poulet numbers.

The name "Sarrus numbers" is after Frédéric Sarrus, who, in 1819, discovered that 341 is a counterexample to the "Chinese hypothesis" that n is prime if and only if 2^n is congruent to 2 (mod n). - _Alonso del Arte_, Apr 28 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000447 | OEIS-A001581

Winning moves in Fibonacci nim.

In his 1992 dissertation, Simon Plouffe conjectured that the generating function of this sequence is given by 2(1 + z)(3z^5 + 2z^3 + z^2 + z + 2) / ((z^6 + z^5 + z^4 + z^3 + z^2 + z + 1)(z - 1)^2). This agrees with the sequence up to the z^26 term in the expansion (138z^26), but disagrees at z^27 with coefficient 144 instead of 150. - _Boon Suan Ho_, Oct 07 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000448 | OEIS-A001595

a(n) = a(n-1) + a(n-2) + 1, with a(0) = a(1) = 1.

G.f.: (1-x+x^2)/(1-2x+x^3) = 2/(1-x-x^2) - 1/(1-x). [Conjectured by _Simon Plouffe_ in his 1992 dissertation; this is readily verified.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000449 | OEIS-A001605

Indices of prime Fibonacci numbers.

These primes are occurring with about the same normalized frequency as Repunit primes (see Generalized Repunit Conjecture Ref). Assuming a base=1.618 (ratio of sequential terms), then the best fit coefficient is 0.60324 for the first 56 terms, which is already approaching exp(-gamma) = 0.56145948 (A080130). - _Paul Bourdelais_, Aug 23 2024 [Corrected by _Alper Ferudun_, Oct 20 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000450 | OEIS-A001651

Numbers not divisible by 3.

Conjectured (and like even easily proved) to be the graph bandwidth of the complete bipartite graph K_{n,n}. - _Eric W. Weisstein_, Apr 24 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000451 | OEIS-A001652

a(n) = 6*a(n-1) - a(n-2) + 2 with a(0) = 0, a(1) = 3.

Conjecture: For n>1 a(n) is the index of the first occurrence of n in sequence A123737. - _Vaclav Kotesovec_, Jun 02 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000452 | OEIS-A001653

Numbers k such that 2*k^2 - 1 is a square.

(a(n), a(n+1)), with a(0):= 1, give all proper positive solutions m1 = m1(n) and m2 = m2(n), with m1 < m2 and n >= 0, of the Markoff triple (m, m1, m2) (see A002559) for m = 2, i.e., m1^2 - 6*m1*m2 + m2^2 = -4. Hence the unique Markoff triple with largest value m = 2 is (1, 1, 2) (for general m from A002559 this is the famous uniqueness conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000453 | OEIS-A001668

Number of self-avoiding n-step walks on honeycomb lattice.

mu^n <= a(n) <= mu^n alpha^sqrt(n) for mu = A179260 and some alpha. It has been conjectured that a(n) ~ mu^n * n^(11/32). - _Charles R Greathouse IV_, Nov 08 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000454 | OEIS-A001700

a(n) = binomial(2*n+1, n+1): number of ways to put n+1 indistinguishable balls into n+1 distinguishable boxes = number of (n+1)-st degree monomials in n+1 variables = number of monotone maps from 1..n+1 to 1..n+1.

Conjectured: 4^n GaussHypergeometric(1/2,-n; 2; 1) -- Solution for the path which stays in the first and second quadrant. - _Benjamin Phillabaum_, Feb 20 2011 Conjecture: a(n) = Sum_{k=2^n..2^(n+1)-1} A178244(k). - _Mikhail Kurkov_, Feb 20 2021 Kurkov's conjecture is true (see Fried link). - _Sela Fried_, Oct 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000455 | OEIS-A001711

Generalized Stirling numbers, [n+4,4]_3.

Conjectural D-finite with recurrence: a(n) + (-2*n-5)*a(n-1) + (n+2)^2*a(n-2)=0. - _R. J. Mathar_, Feb 16 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000456 | OEIS-A001712

Generalized Stirling numbers, [n+5,5]_3.

Conjecture: a(n) + 3*(-n-3)*a(n-1) + (3*n^2 + 15*n + 19)*a(n-2) - (n+2)^3*a(n-3)=0. - _R. J. Mathar_, Jun 09 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000457 | OEIS-A001722

Generalized Stirling numbers, [n+7,7]_5.

Conjecture D-finite with recurrence a(n) +3*(-n-5)*a(n-1) +(3*n^2+27*n+61)*a(n-2) -(n+4)^3*a(n-3)=0. - _R. J. Mathar_, Jul 20 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000458 | OEIS-A001723

Generalized Stirling numbers, [n+8,8]_5.

Conjecture D-finite with recurrence a(n) +2*(-2*n-11)*a(n-1) +(6*n^2+60*n+151)*a(n-2) -(2*n+9)*(2*n^2+18*n+41)*a(n-3) +(n+4)^4*a(n-4)=0. - _R. J. Mathar_, Jul 20 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000459 | OEIS-A001724

Generalized Stirling numbers, [n+9,9]_5.

Conjecture D-finite with recurrence a(n) +5*(-n-6)*a(n-1) +5*(2*n^2+22*n+61)*a(n-2) -5*(n+5)*(2*n^2+20*n+51)*a(n-3) +(5*n^4+90*n^3+610*n^2+1845*n+2101)*a(n-4) -(n+4)^5*a(n-5)=0. - _R. J. Mathar_, Jul 20 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000460 | OEIS-A001752

Expansion of 1/((1+x)*(1-x)^5).

Conjecture: For n > 2, a(n-3) is the absolute value of the coefficient of the term [x^(n-2)] in the characteristic polynomial of an n X n square matrix M(n) defined as the n-th principal submatrix of the array A010751 whose general element is given by M[i,j] = floor((j - i + 1)/2). - _Stefano Spezia_, Jan 12 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000461 | OEIS-A001764

a(n) = binomial(3*n,n)/(2*n+1) (enumerates ternary trees and also noncrossing trees).

a(n) is (conjecturally) the number of permutations of [n+1] that avoid the patterns 4-2-3-1 and 4-2-5-1-3 and end with an ascent. For example, a(4)=55 counts all 60 permutations of [5] that end with an ascent except 42315, 52314, 52413, 53412, all of which contain a 4-2-3-1 pattern and 42513. - _David Callan_, Jul 22 2008 Number of sum-indecomposable (4231,42513)-avoiding permutations. Conjecturally, number of sum-indecomposable (2431,45231)-avoiding permutations. - _Alexander Burstein_, Oct 19 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000462 | OEIS-A001784

Second-order reciprocal Stirling number (Fekete) a(n) = [[2n+3, n]]. The number of n-orbit permutations of a (2n+3)-set with at least 2 elements in each orbit. Also known as associated Stirling numbers of the first kind (e.g., Comtet).

Conjecture: 480*(n+1)*a(n) +30*(-32*n^2-14821*n+42287)*a(n-1) +(878700*n^2-403433*n+5134227)*a(n-2) +(911423*n-656446)*(2*n-3)*a(n-3)=0. - _R. J. Mathar_, Jul 18 2015 Conjecture: (n-2)*(20*n^2-5*n-3)*a(n) -n*(2*n+1)*(20*n^2+35*n+12)*a(n-1)=0. - _R. J. Mathar_, Jul 18 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000463 | OEIS-A001845

Centered octahedral numbers (crystal ball sequence for cubic lattice).

G.f.: (1+x)^3 /(1-x)^4. [conjectured (correctly) by _Simon Plouffe_ in his 1992 dissertation]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000464 | OEIS-A001900

Successive numerators of Wallis's approximation to Pi/2 (unreduced).

Conjecture: a(n) - a(n-1) - n*(n-1)*a(n-2) = 0. - _R. J. Mathar_, Jun 07 2013 [The proof, for n >= 2, follows from the bisection recurrence given below. - _Wolfdieter Lang_, Dec 07 2017]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000465 | OEIS-A001917

a(n) = (p-1)/x, where p = prime(n) and x = ord(2,p), the smallest positive integer such that 2^x == 1 (mod p).

Conjecture: k*a(n) = A006694(((p_n)^k-1)/2). (End) This conjecture fails for k >= 2 exactly when p_n is a Wieferich prime (A001220): At p_183 = 1093, ord(2,1093^2) = ord(2,1093) = 364 gives A006694((1093^2-1)/2) = 3282, not 2*a(183) = 6; likewise at p_490 = 3511. - _Andrew R. Epstein_, Aug 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000466 | OEIS-A001936

Expansion of q^(-1/4) * (eta(q^4) / eta(q))^2 in powers of q.

In the math overflow link is a conjecture that a(n) == a(9*n + 2) (mod 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000467 | OEIS-A001951

A Beatty sequence: a(n) = floor(n*sqrt(2)).

Stephan's conjecture is true (see Fried link). - _Sela Fried_, Oct 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000468 | OEIS-A001969

Evil numbers: nonnegative integers with an even number of 1's in their binary expansion.

Conjecture, checked up to 10^6: a(n) is also the sequence of numbers k representable as k = ror(x) XOR rol(x) (for some integer x) where ror(x)=A038572(x) is x rotated one binary place to the right, rol(x)=A006257(x) is x rotated one binary place to the left, and XOR is the binary exclusive-or operator. - _Alex Ratushnyak_, May 14 2016 Conjecture is true: ror(x) and rol(x) have an even number of 1 bits in total (= 2 * A000120(x)), and XOR preserves the parity of this total, so the resulting number must have an even number of 1 bits. An x can be constructed corresponding to a(n) like so:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000469 | OEIS-A002000

a(n+1) = a(n)*(a(n)^2 - 3) with a(0) = 7.

Conjecture: a(n+1) == a(n) (mod 3^(n+r+2)) for n >= r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000470 | OEIS-A002003

a(n) = 2 * Sum_{k=0..n-1} binomial(n-1, k)*binomial(n+k, k).

For fixed m = 1,2,3,..., we conjecture that the sequence b(n) := a(m*n) satisfies a recurrence of the form P(2*m,n)*b(n+1) + P(2*m,-n)*b(n-1) = Q(2*m,n)*b(n), where the polynomials P(2*m,n) and Q(2*m,n) have degree 2*m. Conjecturally, the polynomial Q(2*m,n) is an even function of n; its 2*m zeros seem to belong to the interval [-1, 1] and 2*m - 2 of these zeros appear to lie close to the rational numbers of the form +-(2*k + 1)/(2*m), where 0 <= k <= m - 2. Cf. A103885. (End) Conjecture: a(n) = - [x^n] (1 - F(x))^n, where F(x) = 2*x + 6*x^2 + 34*x^3 + 238*x^4 + ... is the o.g.f. of A108424. Equivalently, a(n) = -[x^n](G(x))^(-n), where G(x) = 1 + 2*x + 10*x^2 + 66*x^3 + 498*x^4 + ... is the o.g.f. of A027307.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000471 | OEIS-A002034

Kempner numbers: smallest positive integer m such that n divides m!.

Conjecture: a(n) = 1 + n - Sum_{k=1..n} Sum_{m=1..n} cos(-2*Pi*k/n*m!)/n. Formula verified for the first 500 terms. - _Mats Granvik_, Feb 26 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000472 | OEIS-A002049

Prime numbers of measurement.

Andrews conjectures that a(n) ~ (1/2) n^2 log n / loglog n. - _N. J. A. Sloane_, Dec 01 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000473 | OEIS-A002050

Number of simplices in barycentric subdivision of n-simplex.

Conjecture: Let k be a positive integer. The sequence obtained by reducing a(n) modulo k is eventually periodic with the period dividing phi(k) = A000010(k). For example, modulo 9 we obtain the sequence [0, 1, 5, 7, 5, 1, 5, 4, 5, 7, 5, 1, 5, 4, 5, 7, 5, 1, 5, 4, 5, 7, 5, ...], with an apparent period of 6 = phi(9) beginning at a(5). - _Peter Bala_, Aug 03 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000474 | OEIS-A002062

a(n) = Fibonacci(n) + n.

Let _x indicate the sequence offset. Then a(n+2)_0 = A006355(n+4)_0 - A066982(n+1)_1 (conjecture); (a(n)) = em[K*]seq(.25'i - .25'j - .25'k - .25i' + .25j' - .75k' - .25'ii' - .25'jj' - .25'kk' - .25'ij' - .25'ik' - .75'ji' + .25'jk' - .25'ki' - .75'kj' + .75e), apart from initial term. - _Creighton Dement_, Nov 19 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000475 | OEIS-A002063

a(n) = 9*4^n.

These are squares that can be expressed as sum of exactly two distinct powers of two. For instance, a(4) = 9*4^4 = 2304 = 2^11 + 2^8 . It is conjectured that these are the only squares with this characteristic (tested on squares up to (10^7)^2). - _Andres Cicuttin_, Apr 23 2016 Conjecture is true. It is equivalent to prove that the Diophantine equation m^2 = 2^k*(1+2^h), where h>0, has solutions only when h=3. Dividing by 2^k we must obtain an odd square on the left, since 1+2^h is odd, so we can write (2*r+1)^2 = 1+2^h. Expanding, we have 4*r*(r+1) = 2^h, from which it follows that r must be equal to 1 and thus h=3, since r and r+1 must be powers of 2. - _Giovanni Resta_, Jul 27 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000476 | OEIS-A002067

a(n) = Sum_{k=0..n-1} binomial(2*n,2*k)*a(k)*a(n-k-1).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000477 | OEIS-A002070

Coefficient of x^p (p = n-th prime) in x * Product_{k>=1} (1-x^k)^2*(1-x^11k)^2.

Conjecture: a(n) = Sum_{k=1..prime(n)} Sum_{y=1..prime(n)} Sum_{x=1..prime(n)} (A023900(k)/prime(n))[GCD(f(x,y), prime(n)) = k], where f(x,y) = x^3 - x^2 - y^2 - y. - _Mats Granvik_, Oct 09 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000478 | OEIS-A002093

Highly abundant numbers: numbers k such that sigma(k) > sigma(m) for all m < k.

Conjecture: (a) Every highly abundant number > 10 is practical (A005153). (b) For every integer k there exists A such that k divides a(n) for all n > A. Daniel Fischer proved that every highly abundant number greater than 3, 20, 630 is divisible by 2, 6, 12 respectively. The first conjecture has been verified for the first 10000 terms. - _Jaycob Coleman_, Oct 16 2013 Conjecture: For each term k: (1) Let p be the largest prime less than k (if one exists) and let q be the smallest prime greater than k; then k-p is either 1 or a prime, and q-k is either 1 or a prime. (2) The closest prime number p < k located to a distance d = k-p > 1 is also always at a prime distance. These would mean that the even highly abundant numbers greater than 2 always have at least a Goldbach pair of primes. h=p+d. Both observations verified for the first 10000 terms. - _David Morales Marciel_, Jan 04 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000479 | OEIS-A002104

Logarithmic numbers.

Conjecture: a(n) +(-n-1)*a(n-1) +2*(n-1)*a(n-2) +(-n+2)*a(n-3)=0. - _R. J. Mathar_, Dec 02 2012 From the first one, we have the recurrence reported below by R. R. Forberg. From the second one, we have the recurrence conjectured above. (End) G.f.: conjecture: T(0)*x/(1-2*x)/(1-x), where T(k) = 1 - x^2*(k+1)^2/(x^2*(k+1)^2 - (1 - 2*x*(k+1))*(1 - 2*x*(k+2))/T(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 18 2013 Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000480 | OEIS-A002105

Reduced tangent numbers: 2^n*(2^{2n} - 1)*|B_{2n}|/n, where B_n = Bernoulli numbers.

Conjecture: taking the sequence modulo an integer k gives an eventually purely periodic sequence with period dividing phi(k). For example, the sequence taken modulo 10 begins [1, 1, 4, 4, 6, 6, 4, 4, 6, 6, 4, 4, 6, 6, ...] with an apparent period [4, 4, 6, 6] of length 4 = phi(10) beginning at a(3). - _Peter Bala_, May 08 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000481 | OEIS-A002106

Number of transitive permutation groups of degree n.

It is conjectured that this is the number of Galois groups for irreducible polynomials of order n. (All such Galois groups are transitive.) - _Charles R Greathouse IV_, May 28 2014 Let G be a transitive permutation groups of degree n. Then G occurs as a Galois group for an irreducible polynomial of degree n with coefficients K if and only if K admits a Galois extension with Galois group G. ("=>" is true by definition of the Galois group for an irreducible polynomial; for "<=", see user631's answer in the Math Overflow link). Hence the conjecture above is equivalent to the inverse Galois problem. Every finite group can be realized as a Galois group of some extension L/K, but for a fixed base field K (for example, K = Q is the field of rational numbers) the question is usually open. - _Jianing Song_, May 26 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000482 | OEIS-A002107

Expansion of Product_{k>=1} (1 - x^k)^2.

Conjecture: C = sqrt(8*Pi*log(2 + sqrt(3))/3), D = log(7 + 4*sqrt(3)). - _Vaclav Kotesovec_, Sep 26 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000483 | OEIS-A002111

Glaisher's G numbers.

Conjecturally, a(n) = the unsigned numerator of B(2*n+1,1/3). Cf. A033470.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000484 | OEIS-A002114

Glaisher's H' numbers.

Conjecture: more generally, reducing the sequence modulo a positive integer k produces an eventually periodic sequence with period dividing phi(k). For example, modulo 13 the sequence becomes [1, 11, 2, 5, 7, 0, 1, 11, 2, 5, 7, 0, 1, 11, 2, 5, 7, 0, ...], with an apparent period of 6 = phi(13)/2. Cf. A000364 and A000436. [The conjecture is true when k = p is a prime since the congruence a(n+p-1) == a(n) (mod p) holds for all n >= 1. Cf. A215066. - _Peter Bala_, Oct 13 2026] (End) Conjectural e.g.f.: Sum_{n >= 1} (-1)^n*Product_{k = 1..n} (1 - exp(A007310(k)*z)) = z + 11*z^2/2! + 301*z^3/3! + ... - _Peter Bala_, Dec 09 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000485 | OEIS-A002143

Class numbers h(-p) where p runs through the primes p == 3 (mod 4).

Conjecture: For any prime p > 3 with p == 3 (mod 8), we have 2*h(-p)*sqrt(p) = Sum_{k=1..(p-1)/2} csc(2*Pi*k^2/p). - _Zhi-Wei Sun_, Aug 06 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000486 | OEIS-A002146

Smallest prime == 7 (mod 8) where Q(sqrt(-p)) has class number 2n+1.

Conjecture: a(n) < A002148(n) for all n >= 1. - _Jianing Song_, Jul 20 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Mira Bernstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000487 | OEIS-A002149

Largest prime p==3 (mod 8) such that Q(sqrt(-p)) has class number 2n+1.

Most of these values are only conjectured to be correct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000488 | OEIS-A002162

Decimal expansion of the natural logarithm of 2.

Conjecture: for k >= 0 and r even with r - 1 <= k, the series Sum_{n >= 1} (-1)^n * (d/dn)^r (P(n,k)) / (P(n,k)^2 = A(r,k)*log(2) + B(r,k), where A(r,k) and B(r,k) are both rational numbers. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000489 | OEIS-A002190

Sum_{n>=0} a(n)*x^n/n!^2 = -log(BesselJ(0,2*sqrt(x))).

Conjecture: G.f.: 1 = Sum_{n>=0} a(n+1)*A000108(n)*x^n*Sum_{k>=0} C(2*n+k,k)^2*(-x)^k. Compare with the following g.f of the Catalan numbers (A000108): 1 = Sum_{n>=0} A000108(n)*x^n*Sum_{k>=0} C(2*n+k,k)*(-x)^k. - _Paul D. Hanna_, Oct 10 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000490 | OEIS-A002193

Decimal expansion of square root of 2.

sqrt(2) = 2 + n*A167199(n-1)/A167199(n) as n -> infinity (conjecture). - _Mats Granvik_, Oct 30 2009 Conjecture: Equals Sum_{k>=0} A014969(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000491 | OEIS-A002285

Decimal expansion of common logarithm of e.

Conjecture by Eric Weisstein: Equals lim_{n->oo} b(n)/10^(n-1), for b=A114467 or b=A114468 (i.e., is the limit of the decimal expansion of the number of decimal digits in both the numerator and denominator of the (10^n)th harmonic number). More generally, log_k(e) seems to equal lim_{n->oo} floor(log_k(b(k^n)))/k^(n-1), for b=A001008 or b=A002805 and k >= 2. - _Natalia L. Skirrow_, Feb 12 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000492 | OEIS-A002293

Number of dissections of a polygon: binomial(4*n, n)/(3*n + 1).

Conjecturally, a(n) is the number of 3-uniform words over the alphabet [n] that avoid the patterns 231 and 221 (see the Defant and Kravitz link). - _Colin Defant_, Sep 26 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000493 | OEIS-A002294

a(n) = binomial(5*n, n)/(4*n + 1).

Conjecturally, a(n) is the number of 4-uniform words on the alphabet [n] that avoid the patterns 231 and 221 (see the Defant and Kravitz link). - _Colin Defant_, Sep 26 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000494 | OEIS-A002312

Arc-cotangent reducible numbers or non-Størmer numbers: largest prime factor of k^2 + 1 is less than 2*k.

Also numbers such that k^2 + 1 has no primitive divisor, hence (by Everest & Harman, Theorem 1.4) 2.138n < a(n) < 10.6n for large enough n. They conjecture that a(n) ~ cn where c = 1/(1 - log 2) = 3.258.... - _Charles R Greathouse IV_, Nov 15 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000495 | OEIS-A002376

Least number of positive cubes needed to sum to n.

The g.f. conjectured by Simon Plouffe in his 1992 dissertation,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000496 | OEIS-A002385

Palindromic primes: prime numbers whose decimal expansion is a palindrome.

Conjecture: The only primes with palindromic prime indices that are palindromic primes themselves are 3, 5 and 11. Tested for the primes with the first 8000000 palindromic prime indices. - _Ivan N. Ianakiev_, Oct 10 2014 It follows from the above conjecture that 2 is the only k such that k, prime(k), prime(m) = k + prime(k) and m are all palindromic primes. - _Ivan N. Ianakiev_, Mar 17 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000497 | OEIS-A002387

Least k such that H(k) > n, where H(k) is the harmonic number Sum_{i=1..k} 1/i.

It is conjectured that, for n > 1, a(n) = floor(exp(n-gamma) + 1/2). - _Benoit Cloitre_, Oct 23 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000498 | OEIS-A002391

Decimal expansion of natural logarithm of 3.

Conjecture: log(3) = Sum_{k>=1} A191907(3,k)/k. - _Mats Granvik_, Jun 19 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000499 | OEIS-A002414

Octagonal pyramidal numbers: a(n) = n*(n+1)*(2*n-1)/2.

G.f.: x*(1+5*x)/(1-x)^4. [Conjectured by _Simon Plouffe_ in his 1992 dissertation.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000500 | OEIS-A002439

Glaisher's T numbers.

We make the following conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000501 | OEIS-A002448

Expansion of Jacobi theta function theta_4(x).

Conjecture: Sum_{k>=0} a(k) / exp(k*Pi) = (1/2)*Pi^(1/4)*2^(3/4)/Gamma(3/4) = A248557. - _Simon Plouffe_, Sep 09 2025 The above conjecture is correct. The series equals theta_4(exp(-Pi)) = phi(-exp(-Pi)) = (1/2)^(1/4)*Pi^(1/4)/Gamma(3/4) by Berndt, Entry 1.(ii), p. 325. - _Peter Bala_, Jun 02 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000502 | OEIS-A002457

a(n) = (2n+1)!/n!^2.

Conjecture: a(n) = floor(sqrt((16^n + 1)^(2*n+1))) mod 16^n, for n > 0. - _Joseph M. Shunia_, Sep 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000503 | OEIS-A002464

Hertzsprung's problem: ways to arrange n non-attacking kings on an n X n board, with 1 in each row and column. Also number of permutations of length n without rising or falling successions.

Conjecture: a(n) = Sum_{k=1..n} k!*A080246(n-1, k-1) for n > 0. - _John Keith_, Nov 02 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000504 | OEIS-A002467

The game of Mousetrap with n cards (given n letters and n envelopes, how many ways are there to fill the envelopes so that at least one letter goes into its right envelope?).

For n > 0, a(n) = {(1-1/exp(1))*n!}, where {x} is the nearest integer. - _Simon Plouffe_, conjectured March 1993, added Feb 17 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Jeffrey Shallit_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000505 | OEIS-A002485

Numerators of convergents to Pi.

I conjecture the following identity below, which represents a generalization of Stephen Lucas's experimentally obtained identities:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000506 | OEIS-A002487

Stern's diatomic series (or Stern-Brocot sequence): a(0) = 0, a(1) = 1; for n > 0: a(2*n) = a(n), a(2*n+1) = a(n) + a(n+1).

Piesk's conjecture above is true: the nim-product 2^x*2^y depends only on x XOR y and x AND y, which together form a hyperbinary representation of n = x + y, and different representations give different products; see the Blanco Gómez link. - _Roberto Blanco Gómez_, Oct 05 2026 Let f(m,n) be the frequency of the integer n in the interval [a(2^(m-1)), a(2^m-1)]. Let phi(n) be Euler's totient function (A000010). Conjecture: for all integers m,n n<=m f(m,n) = phi(n). - _Yosu Yurramendi_, Sep 08 2014 The conjecture is true; in fact 1 <= a(n) < A007306(n+1) and A007306(n+1) divides a(n)*A007305(n+1) - 1 for n >= 1 (see Epstein link). - _Andrew R. Epstein_, Sep 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000507 | OEIS-A002491

Smallest number of stones in Tchoukaillon (or Mancala, or Kalahari) solitaire that make use of n-th hole.

Unfortunately, I believe this paper erred when it claimed to have proved the conjecture of Erdős and Jabotinsky about the sharpness of approximation to n^2/pi. When they computed the sum of I_M in the proof of Theorem 9, they expressed it as f(M-1)^2/4M + O(f(M-1)). That's correct; but the error gets quite large, and when summed gives O(n^(3/2)), not O(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000508 | OEIS-A002496

Primes of the form k^2 + 1.

It is conjectured that this sequence is infinite, but this has never been proved. It is a result that goes back to Mirsky that the set of primes p for which p-1 is squarefree has density A, where A = A005596 denotes the Artin constant. More precisely, Sum_{p <= x} mu(p-1)^2 = A*x/log x + o(x/log x) as x tends to infinity. Conjecture: Sum_{p <= x, mu(p-1)=1} 1 = (A/2)*x/log x + o(x/log x) and Sum_{p <= x, mu(p-1)=-1} 1 = (A/2)*x/log x + o(x/log x). - Pieter Moree (moree(AT)mpim-bonn.mpg.de), Nov 03 2003 There are O(sqrt(n)/log(n)) terms of this sequence up to n. But this is just an upper bound. See the Bateman-Horn or Wolf papers, for example, for the conjectured for what is believed to be the correct density.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000509 | OEIS-A002559

Markoff (or Markov) numbers: union of positive integers x, y, z satisfying x^2 + y^2 + z^2 = 3*x*y*z.

Assuming that each solution (x,y,z) is ordered x <= y <= z, the open problem is to prove that each z value occurs only once. There are no counterexamples in the first 1046858 terms, which have z values < Fibonacci(5001) = 6.2763...*10^1044. - _T. D. Noe_, Mar 19 2009 Kaneko extends a parameterization of Markoff numbers, citing Frobenius, and relates it to a conjectured behavior of the elliptic modular j-function at real quadratic numbers. - _Jonathan Vos Post_, May 06 2012 Riedel (2012) claims a proof of the unicity conjecture: "it will be shown that the largest member of [a Markoff] triple determines the other two uniquely." - _Jonathan Sondow_, Aug 21 2012 Bourgain, Gamburd, and Sarnak have announced a proof that almost all Markoff numbers are composite--see A256395. Equivalently, the prime Markoff numbers A178444 have density zero among all Markoff numbers. (It is conjectured that infinitely many Markoff numbers are prime.) - _Jonathan Sondow_, Apr 30 2015 According to Sarnak on Apr 30 2015, all claims to have proved the unicity conjecture have turned out to be false. - _Jonathan Sondow_, May 01 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _J. H. Conway_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000510 | OEIS-A002586

Smallest prime factor of 2^n + 1.

Conjecture: a(8+48*k) = 257 and a(40+48*k) = 257, where k is a nonnegative integer. - _Thomas König_, Feb 15 2017 Conjecture is true: 257 divides 2^(8+48*k)+1 and 2^(40+48*k)+1 but no prime < 257 ever does. Similarly, a(24+48*k) = 97. - _Robert Israel_, Feb 17 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000511 | OEIS-A002596

Numerators in expansion of sqrt(1+x). Absolute values give numerators in expansion of sqrt(1-x).

Observation/conjecture: If a term of this sequence is divisible by a prime p, then that term is in a block of exactly (p^k-3)/2 consecutive terms all of which are divisible by p. Furthermore, if a(n) is the term preceding such a block then a(p*n-(p-1)/2) also precedes a block of (p^(k+1)-3)/2 terms all divisible by p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000512 | OEIS-A002618

a(n) = n*phi(n).

The conjecture by Layman is correct. Obviously any such permutation must have an increment that is prime to n, and almost as obvious that any such increment will work, with any starting value; hence phi(n) * n total. - _Franklin T. Adams-Watters_, Jun 09 2009 Conjecture: All the rational numbers Sum_{i=j..k} 1/a(i) with 0 < min{2,k} <= j <= k have pairwise distinct fractional parts. - _Zhi-Wei Sun_, Sep 24 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000513 | OEIS-A002627

a(n) = n*a(n-1) + 1, a(0) = 0.

Conjecture: a(n) + (-n-1)*a(n-1) + (n-1)*a(n-2) = 0. - _R. J. Mathar_, Feb 16 2014 a(n+1) = a(n) + n! + Sum_{k=1..n} (n!/k!)*a(k). The above conjectured recurrence can be obtained from the original recurrence or from the differential equation satisfied by f(x). - _Emanuele Munarini_, Jun 20 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000514 | OEIS-A002660

a(n) = Sum_{d|n, d <= 3} d^2 + 3*Sum_{d|n, d>3} d.

Conjectured: Inverse Moebius transform of g.f.: (x + 2x^2 + 2x^3 - 2x^4) / (1 - x)^2. - _Sean A. Irvine_, May 16 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000515 | OEIS-A002691

a(n) = (n+2) * (2n+1) * (2n-1)! / (n-1)!.

Conjecture: a(n) + 4*(-n-1)*a(n-1) + 4*(-2*n+1)*a(n-2) = 0. - _R. J. Mathar_, Jun 07 2013 Conjecture is true and can be proved by direct substitution of the definition. - _Chai Wah Wu_, Jul 22 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000516 | OEIS-A002716

An infinite coprime sequence defined by recursion.

Conjecture: a(2n+1)=A001146(n+1)+1. - _R. J. Mathar_, May 15 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000517 | OEIS-A002720

Number of partial permutations of an n-set; number of n X n binary matrices with at most one 1 in each row and column.

Conjecture: a(n)*k is the sum of the largest parts in all integer partitions containing their own first differences with n + 1 parts and least part k. - _John Tyler Rascoe_, Feb 28 2024 The conjecture is true. Let k = x_0 < ... < x_n be such a partition. For each i>0 there exists a unique j in [0..i-1] so that x_i = x_{i-1} + x_j. Conversely, each choice of the j's recursively determines an admissible partition, so there are n! such partitions. Let b(n) be the mean of x_n/k over these partitions. Then b(0) = 1 and b(n) = b(n-1) + (1/n) * Sum_{j=0..n-1} b(j), which is the recurrence given below for b(n) = a(n)/n!. Hence the sum of the largest parts is n!*E(x_n) = n!*k*b(n) = k*a(n). - _David Radcliffe_, Jul 20 2026 Asymptotically, a(n)/n! ~ (1/2)*Pi^(-1/2)*exp(-1/2 + 2*n^(1/2))/n^(1/4) and so a(n) ~ C*Bessell(0, 2*sqrt(n))*n! with C = exp(-1/2) = 0.6065306597126334236... - Alec Mihailovs, Sep 06 2005, establishing a conjecture of _Franklin T. Adams-Watters_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000518 | OEIS-A002742

Logarithmic numbers.

Conjectures: Let k be a positive integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000519 | OEIS-A002791

a(n) = Sum_{d|n, d <= 4} d^2 + 4*Sum_{d|n, d>4} d.

Conjectured: Inverse Moebius transform of g.f.: (x + 2x^2 + 2x^3 + 2x^4 - 3x^4) / (1 - x)^2. - _Sean A. Irvine_, May 16 2014 Conjectured: a(n) = 4 * sigma(n) - f(n mod 6) where f(0) = 10, f(1) = 3, f(2) = 7, f(3) = 6, f(4) = 7, f(5) = 3. - _Sean A. Irvine_, May 17 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000520 | OEIS-A002804

(Presumed) solution to Waring's problem: g(n) = 2^n + floor((3/2)^n) - 2.

It is known (Kubina and Wunderlich, 1990) that g(n) = 2^n + floor((3/2)^n) - 2 for all n <= 471600000. This formula is conjectured to be correct for all n (see A174420). This sequence (which corresponds to Waring's original conjecture) is much easier to compute than A079611, the problem of finding the minimal s = G(n) for almost all (= sufficiently large) integers. See Wikipedia for a one-line proof that this value for g(n), conjectured by J. A. Euler in 1772, is indeed a lower bound; it is known to be tight if 2^n*frac((3/2)^n) + floor((3/2)^n) <= 2^n, and no counterexample to this inequality is known. - _M. F. Hasler_, Jun 29 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000521 | OEIS-A002814

For n > 1: a(n) = a(n-1)^3 + 3*a(n-1)^2 - 3; a(0) = 1, a(1) = 2.

Conjecture: The sequence {a(n) - 2: n >= 1} is a strong divisibility sequence, that is, gcd(a(n) - 2, a(m) - 2) = a(gcd(n, m)) - 2 for n, m >= 1. [added Jul 26 2026: This is correct - see the Bala link. More generally, for k >= 1, the sequence {a(n+k) - a(k) : n >= 1} is a strong divisibility sequence.] - _Peter Bala_, Dec 08 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000522 | OEIS-A002815

a(n) = n + Sum_{k=1..n} pi(k), where pi() = A000720.

Conjectured g.f.: (Sum_{N>=1} x^A008578(N))/(1-x)^2 = (x + x^2 + x^3 + x^5 + x^7 + x^11 + x^13 + ...)/(1-x)^2. - _L. Edson Jeffery_, Nov 25 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Robert G. Wilson v_, _Mira Bernstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000523 | OEIS-A002839

Number of simple perfect squared rectangles of order n up to symmetry.

In "A Census of Planar Maps", p. 267, William Tutte gave a conjectured asymptotic formula for the number of perfect squared rectangles where n is the number of elements in the dissection (the order): Conjecture: a(n) ~ n^(-5/2) * 4^n / (243*sqrt(Pi)). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000524 | OEIS-A002843

Number of partitions of n into parts 1/2, 3/4, 7/8, 15/16, etc.

The g.f. (z^2+z+1)*(z-1)^2/(1-2*z-z^3+3*z^4) conjectured by _Simon Plouffe_ in his 1992 dissertation is wrong. Conjecture: a(n) ~ A102375 * a(n-1). - _Bill McEachen_, Feb 05 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000525 | OEIS-A002857

Number of Post functions of n variables.

Conjecture: a(n) = A055621(n) - A055152(n). - _R. J. Mathar_, Oct 14 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000526 | OEIS-A002865

Number of partitions of n that do not contain 1 as a part.

Conjecture: Also the number of integer partitions of n - 1 that have a consecutive subsequence summing to each positive integer from 1 to n - 1. For example, (32211) is such a partition because we have consecutive subsequences: There is a sufficient and necessary condition to characterize the partitions defined by Gus Wiseman. It is that the largest part must be less than or equal to the number of ones plus one. Hence, the number of partitions of n with no part greater than the number of ones is the same as the number of partitions of n-1 that have a consecutive subsequence summing to each integer from 1 to n-1. Gus Wiseman's conjecture can be proved bijectively. - _Andrew Yezhou Wang_, Dec 14 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000527 | OEIS-A002867

a(n) = binomial(n,floor(n/2))*(n+1)!.

Conjecture: a(n) - 2*a(n-1) - 4*n*(n-1)*a(n-2) = 0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000528 | OEIS-A002878

Bisection of Lucas sequence: a(n) = L(2*n+1).

Conjecture: for n > 0, a(n) = sqrt(Fibonacci(4*n+3) + Sum_{k=2..2*n} Fibonacci(2*k)). - _Alex Ratushnyak_, May 06 2012 The conjecture is true and was proved by Davala (see comments to A000045). - _Sela Fried_, Nov 29 2025 Conjecture: except for the number 3, a(n) are the numbers such that a(n)^2+2 are Lucas numbers. - _Michel Lagneau_, Jul 22 2014 Comment on the preceding conjecture: It is clear that all a(n) satisfy a(n)^2 + 2 = L(2*(2*n+1)) due to the identity (17 c) of Vajda, p. 177: L(2*n) + 2*(-1)^n = L(n)^2 (take n -> 2*n+1). - _Wolfdieter Lang_, Oct 10 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000529 | OEIS-A002893

a(n) = Sum_{k=0..n} binomial(n,k)^2 * binomial(2*k,k).

Conjecture: For each n=1,2,3,... the polynomial g_n(x) = Sum_{k=0..n} binomial(n,k)^2*binomial(2k,k)*x^k is irreducible over the field of rational numbers. - _Zhi-Wei Sun_, Mar 21 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000530 | OEIS-A002895

Domb numbers: number of 2n-step polygons on diamond lattice.

Conjecture: Let D(n) be the (n+1) X (n+1) Hankel-type determinant with (i,j)-entry equal to a(i+j) for all i,j = 0..n. Then the number D(n)/12^n is always a positive odd integer. - _Zhi-Wei Sun_, Aug 14 2013 The conjecture of _Zhi-Wei Sun_ (Aug 14 2013) was proved in Theorem 1.2 of Zhu and Sun (2018). - _Eiver Rodríguez_, Sep 08 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000531 | OEIS-A002966

Egyptian fractions: number of solutions of 1 = 1/x_1 + ... + 1/x_n where 0 < x_1 <= ... <= x_n.

T(.,2) = A018892. T(.,3) = A004194. T(.,4) = A020327, T(.,5) = A020328. T(2,6) is computed by _D. S. McNeil_, who conjectures that the 2nd row is A003167. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000532 | OEIS-A002984

a(0) = 1; for n > 0, a(n) = a(n-1) + floor(sqrt(a(n-1))).

Conjecture: a(n) ~ n^2/4. - _José María Grau Ribas_, Feb 13 2024 Proof of Ribas's conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000533 | OEIS-A003001

Smallest number of multiplicative persistence n.

For any multiplicative persistence, except multiplicative persistence 2, the set of class representative numbers with that multiplicative persistence is conjectured to be finite. It is observed from this that for the reduced numbers with multiplicative persistence 1, the primes 11, 13, 17 and 19, will not occur in any trajectory of another (larger) number; i.e., all numbers represented by the reduced numbers 11, 13, 17 and 19 have a prime factor of at least 11 (conjectured from the observations). In fact all trajectories can be projected to a trajectory in one of the ten trees with reduced numbers with roots 0..9, and the numbers represented by the reduced number of each leaf have a prime factor of at least 11 (as conjectured from the observations).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000534 | OEIS-A003024

Number of acyclic digraphs (or DAGs) with n labeled nodes.

Also the number of n X n real (0,1)-matrices with all eigenvalues positive. - Conjectured by _Eric W. Weisstein_, Jul 10 2003 and proved by McKay et al. 2003, 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000535 | OEIS-A003056

n appears n+1 times. Also the array A(n,k) = n+k (n >= 0, k >= 0) read by antidiagonals. Also inverse of triangular numbers.

Fan, Yang, and Yu (2012) prove a conjecture of Amdeberhan on the generating function of a(n). - _Jonathan Sondow_, Dec 17 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000536 | OEIS-A003067

Problimes (second definition).

The g.f. (z^2+2+z^9+z^5)/(z-1)^2 conjectured by _Simon Plouffe_ in his 1992 dissertation is wrong.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000537 | OEIS-A003075

Minimal number of comparisons needed for n-element sorting network.

It is conjectured that the sequence continues (after 39) 45, 51, 56, 60, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000538 | OEIS-A003108

Number of partitions of n into cubes.

The g.f. 1/(z+1)/(z**2+1)/(z**4+1)/(z-1)**2 conjectured by _Simon Plouffe_ in his 1992 dissertation is wrong.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Herman P. Robinson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000539 | OEIS-A003116

Expansion of the reciprocal of the g.f. defining A039924.

Conjecture: a(n) is the number of compositions p(1) + p(2) + ... + p(m) = n with p(i)-p(i-1) <= 1, see example; cf. A034297. - _Vladeta Jovovic_, Feb 09 2004 Jovovic's conjecture can be proved as follows. There is a sign-changing involution defined on pairs (L1,L2) where L1 is a partition with difference >= 2 between consecutive parts and L2 is the number of compositions described by Jovovic, with the sign (-1)^(Number of parts of L1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Herman P. Robinson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000540 | OEIS-A003120

Number of rooted trees with n nodes and omega-valency 1.

The g.f. -(z-1)*(3*z**2+z-1)/(-1+3*z+z**2-7*z**3+3*z**4) conjectured by _Simon Plouffe_ in his 1992 dissertation is wrong (starting from index 11).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000541 | OEIS-A003157

A self-generating sequence (see Comments in A003156 for the definition).

These are the positions of 1 in A286044; complement of A286045; conjecture: a(n)/n -> 4. - _Clark Kimberling_, May 07 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000542 | OEIS-A003159

Numbers whose binary representation ends in an even number of zeros.

A conjectured identity relating to the partition sequence, A000041 as polcoeff p(x); A003159, and its characteristic function A035263: (1, 0, 1, 1, 1, 0, 1, 0, 1, 0, 1, 1, 1, ...); and A036554 indicating n-th terms with zeros in A035263: (2, 6, 8, 10, 14, 18, 22, ...). The conjecture states that p(x) = A(x) = A(x^2) when A(x) = polcoeff A174065 = the Euler transform of A035263 = 1/((1-x)*(1-x^3)*(1-x^4)*(1-x^5)*...) = 1 + x + x^2 + 2*x^3 + 3*x^4 + 4*x^5 + ... and the aerated variant = the Euler transform of the complement of A035263: 1/((1-x^2)*(1-x^6)*(1-x^8)*...) = 1 + x^2 + x^4 + 2*x^6 + 3*x^8 + 4*x^10 + ... The conjecture above was proved by Jean-Paul Allouche on Dec 21 2013. - _Gary W. Adamson_, Jan 22 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000543 | OEIS-A003171

Negated discriminants of orders of imaginary quadratic fields with 1 class per genus (a finite sequence).

It is conjectured that a(101) = 7392 is the last term. If it would exist, a(102) > 10^6. - _Hugo Pfoertner_, Dec 01 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Mira Bernstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000544 | OEIS-A003185

a(n) = (4*n+1)*(4*n+5).

Conjecture: a(n+1)-a(n) = A063164(n+2). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000545 | OEIS-A003192

Length of uncrossed knight's path on an n X n board.

I used ZDD techniques to show that a(9)=47. (This is the longest uncrossed knight's path on a 9 X 9 board; thus I confirmed the conjecture that the paths of length 47, found by hand long ago, are indeed optimum.) - _Don Knuth_, Jun 24 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000546 | OEIS-A003237

Number of partially achiral planted trees with n nodes.

The g.f. z*(1-z**2-z**3-z**4+z**5)/(1-z-2*z**2+3*z**5) conjectured by _Simon Plouffe_ in his 1992 dissertation is wrong.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000547 | OEIS-A003238

Number of rooted trees with n vertices in which vertices at the same level have the same degree.

According to the MathOverflow link, log(a(n)) ~ log(n)^2/log(4), and a more precise asymptotic expansion is similar to that of A018819 and hence A000123, so the conjecture in the Formula section is partly correct. - _Andrei Zabolotskii_, Jan 22 2017 Conjecture: log(a(n)) is asymptotic to c*log(n)^2 where 0.4 < c < 0.5 - _Benoit Cloitre_, Apr 13 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000548 | OEIS-A003243

Number of partially achiral trees with n nodes.

The g.f. (1-z**2-2*z**3-8*z**4+7*z**5+4*z**6)/(1-z-z**2-2*z**3-6*z**4+14*z**5) was conjectured by _Simon Plouffe_ in his 1992 dissertation, but this is incorrect.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000549 | OEIS-A003258

The number m such that c'(m) = A005206(A003231(n)), where c'(m) = A249115(m) is the m-th positive integer not in A003231.

Conjecture: a(n) = A078489(n) + n - 1. - _Ralf Stephan_, Feb 24 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000550 | OEIS-A003277

Cyclic numbers: k such that k and phi(k) are relatively prime; also k such that there is just one group of order k, i.e., A000001(k) = 1.

Any divisor of a Carmichael number (A002997) must be odd and cyclic. Conversely, G. P. Michon conjectured (c. 1980) that any odd cyclic number has at least one Carmichael multiple (if the conjecture is true, each of them has infinitely many such multiples). In 2007, Michon & Crump produced explicit Carmichael multiples of all odd cyclic numbers below 10000 (see link, cf. A253595). - _Gerard P. Michon_, Jan 08 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Richard Stanley_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000551 | OEIS-A003278

Szekeres's sequence: a(n)-1 in ternary = n-1 in binary; also: a(1) = 1, a(2) = 2, and thereafter a(n) is smallest number k which avoids any 3-term arithmetic progression in a(1), a(2), ..., a(n-1), k.

Conjecture: a(n) = (A191107(n) + 2)/3 = (A055246(n) + 5)/6. - _L. Edson Jeffery_, Nov 26 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Richard Stanley_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000552 | OEIS-A003285

Period of continued fraction for square root of n (or 0 if n is a square).

Apparently the generating function of the sequence for the denominators of continued fraction convergents to sqrt(n) is always rational and of form p(x)/[1 - C*x^m + (-1)^m * x^(2m)], or equivalently, the denominators satisfy the linear recurrence b(n+2m) = C*b(n+m) - (-1)^m * b(n), where a(n) is equal to m for each nonsquare n, or 0. See A006702 for the conjecture regarding C. The same conjectures apply to the sequences of the numerators of continued fraction convergents to sqrt(n). - _Ralf Stephan_, Dec 12 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000553 | OEIS-A003313

Length of shortest addition chain for n.

It is conjectured by D. E. Knuth, K. Stolarsky et al. that for all n: floor(log_2(n)) + ceiling(log_2(v(n))) <= a(n), where v(n) = A000120(n). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000554 | OEIS-A003319

Number of connected permutations of [1..n] (those not fixing [1..j] for 0 < j < n). Also called indecomposable permutations, or irreducible permutations.

Conjecture: a(n) = A370380(n-2, 0) = A370381(n-2, 0) for n > 1 with a(0) = a(1) = 1. - _Mikhail Kurkov_, Apr 26 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000555 | OEIS-A003324

A nonrepetitive sequence.

Conjecture: a(2*n) = (A292077(n)+1)*2. Confirmed for first 1000 terms. - _John Keith_, Apr 18 2021 [This conjecture is correct. Write n = (2*k+1) * 2^e. If k+e is even, then we have A292077(n) = 0 and a(2n) = 2; if k+e is odd, then we have A292077(n) = 1 and a(2n) = 4. - _Jianing Song_, Nov 27 2021]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000556 | OEIS-A003325

Numbers that are the sum of 2 positive cubes.

It is conjectured that this sequence and A052276 have infinitely many numbers in common, although only one example (128) is known. [Any further examples are greater than 5 million. - _Charles R Greathouse IV_, Apr 12 2020] [Any further example is greater than 10^12. - _M. F. Hasler_, Jan 10 2021]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000557 | OEIS-A003327

Numbers that are the sum of 4 positive cubes in 1 or more way.

It is conjectured that every number greater than 7373170279850 is in this sequence. [See the paper of the same name. - _T. D. Noe_, May 25 2017] - _Charles R Greathouse IV_, Jan 14 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000558 | OEIS-A003401

Numbers of edges of regular polygons constructible with ruler (or, more precisely, an unmarked straightedge) and compass.

If the well-known conjecture that there are only five prime Fermat numbers F_k=2^{2^k}+1, k=0,1,2,3,4 is true, then we have exactly: Sum_{n>=1} 1/a(n)= 2*Product_{k=0..4} (1+1/F_k) = 4869735552/1431655765 = 3.40147098978.... - _Vladimir Shevelev_ and _T. D. Noe_, Dec 01 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000559 | OEIS-A003416

Sociable numbers: smallest member of each cycle (conjectured).

Sociable numbers: smallest member of each cycle (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000560 | OEIS-A003435

Number of directed Hamiltonian circuits on n-octahedron with a marked starting node.

Conjecture: a(n) -(4*n^2 - 2*n + 5)*a(n-1) + 2*(n-1)*(4*n-17)*a(n-2) + 12*(n-1)*(n-2)*a(n-3) = 0. - _R. J. Mathar_, Oct 02 2013 The conjecture is true. Zeilberger's method applied to the sum in the entry (certificate R(n,k) = -k*(2*k^2*n^2 - 3*k^2*n - 4*k*n^3 + 2*k*n^2 + 14*k*n - 10*k + 16*n^3 - 54*n^2 + 55*n - 18)/(n*(k-2*n+1)*(k-2*n+2)*(k-2*n+3))) proves the recurrence of V. Kotesovec, M(n) := (2*n-3)*a(n) - 2*n*(4*n^2-8*n+5)*a(n-1) - 4*n*(n-1)*(2*n-1)*a(n-2) = 0 for n >= 4. If L(n) denotes the left side of the conjectured recurrence, (2*n-3)*L(n) = M(n) - 3*M(n-1). - _Roberto Blanco Gómez_, Oct 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000561 | OEIS-A003440

Number of binary vectors with restricted repetitions.

It is important to consider these two hypotheses to interpret the series. I also think that the second constraint is not useful and could usefully be deleted. In this case, the series should be doubled from the second term and would become 1, 2, 6, 14, 34, 84, ..., i.e., A177790. G.f.: {(1-x)^2 * sqrt[(1+x+x^2)/(1-3x+x^2)] + x^2 - 1}/(2x^2) (conjectured). - _Ralf Stephan_, Mar 28 2004 Conjecture: (n+2)*a(n) +3*(-n-1)*a(n-1) +(n-2)*a(n-2) +(-n+1)*a(n-3) +3*(n-4)*a(n-4) +(-n+5)*a(n-5)=0. - _R. J. Mathar_, Jun 07 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000562 | OEIS-A003459

Absolute primes (or permutable primes): every permutation of the digits is a prime.

Conjecture: for n >= 23, a(n) = A004022(n-21). - _Max Alekseyev_, Oct 08 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000563 | OEIS-A003477

Expansion of 1/((1-2x)(1+x^2)(1-x-2x^3)).

Conjecture: a(n) = A077854(n) + 2*(a(n-3) + a(n-4) + ... + a(1)). - _Arie Bos_, Nov 29 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000564 | OEIS-A003500

a(n) = 4*a(n-1) - a(n-2) with a(0) = 2, a(1) = 4.

Tsumura shows that, for prime p, a(p) is composite (contrary to a conjecture of Juricevic). - _Charles R Greathouse IV_, Apr 13 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000565 | OEIS-A003558

Least number m > 0 such that 2^m == +-1 (mod 2n + 1).

Start a sequence with numbers 1 and n. For next numbers, add previous numbers going backwards until the sum is even. Then the new number is sum/2. I conjecture that the sequence returns to 1,n and a(n) is the cycle length. Proof of the above conjecture: Let n = -1/2; thus 2n + 1 = 0, so operations are performed mod (2n + 1). When the member is even, it is divided by 2. When it is odd, multiply by n, so effectively divide by -2. This is all well-defined in the sense that new members m are 1 <= m <= n. Now see what happens starting from an odd member m. The next member is -m/2. As long as there are even members, divide by 2 and end up with an odd -m/(2^k). Now add all the members starting with m. The sum is m/(2^k). It's divided by 2, so the next member is m/(2^(k+1)). That is the same as (-m/(2^k))/(-2), as with the definition. Conjecture of Kappraff and Adamson in "Polygons and Chaos", p. 13 Section 7, "Chaos and Number": Given the cycle length for N = 2n + 1, the same cycle length is present in bases 4, 9, 16, 25, ..., m^2, for the expansion of 1/N.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000566 | OEIS-A003592

Numbers of the form 2^i*5^j with i, j >= 0.

Conjecture: Each positive integer n not among 1, 4 and 12 can be written as a sum of finitely many numbers of the form 2^a*5^b + 1 (a,b >= 0) with no one dividing another. This has been verified for n <= 3700. - _Zhi-Wei Sun_, Apr 18 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000567 | OEIS-A003603

Fractal sequence obtained from Fibonacci numbers (or Wythoff array).

Conjecture: a(n) = abs(floor(n/phi) - floor(n*(1/phi + 1/(-phi)^(A035612(n) + 1)))) where phi = (1+sqrt(5))/2. - _Alan Michael Gómez Calderón_, Oct 27 2023 The conjecture is true (see Epstein link). - _Andrew R. Epstein_, Sep 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Mira Bernstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000568 | OEIS-A003630

Inert rational primes in Q(sqrt(3)).

Conjecture: Primes congruent to {5, 7} mod 12. - _Vincenzo Librandi_, Aug 06 2012 The above conjecture is correct. In fact, this is the sequence of primes p such that Kronecker(12,p) = -1 (12 is the discriminant of Q[sqrt(3)]), that is, odd primes that have 3 as a quadratic nonresidue. - _Jianing Song_, Nov 21 2018 Conjecture: Let r(n) = (a(n) - 1)/(a(n) + 1) if a(n) mod 4 = 1, (a(n) + 1)/(a(n) - 1) otherwise; then Product_{n>=1} r(n) = (2/3) * (4/3) * (8/9) * (10/9) * (14/15) * ... = sqrt(3)/2. (See A010527.) We see that the sum of the numerator and denominator of each fraction equals the corresponding term of the sequence: 2 + 3 = 5, 4 + 3 = 7, 8 + 9 = 17, ... - _Dimitris Valianatos_, Mar 26 2017 Conjecture: for n >= 5, a(n) is the sum of three distinct earlier terms. Verified for n up to corresponding prime 10^11 by _Jeff Sponaugle_, Oct 15 2026. - _Yves Debeuret_, Oct 15 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Mira Bernstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000569 | OEIS-A003679

Numbers that are not the sum of 3 pentagonal numbers.

Guy's paper says that the sequence probably contains exactly 210 terms, six of which require five pentagonal numbers: 9, 21, 31, 43, 55 and 89. The last term is conjectured to be 33066. - _T. D. Noe_, Apr 19 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Mira Bernstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000570 | OEIS-A003722

Expansion of e.g.f. sin(sinh(x)) (odd powers only).

Conjecture: a(n) = -2 * Sum_{k=1..oo} (-1)^k * (2*k - 1)^(2*n + 1) * BesselI(2*k - 1, 1). - _Velin Yanev_, Feb 20 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000571 | OEIS-A003824

Numbers that are the sum of two 4th powers in more than one way (primitive solutions).

Conjecture: a(n) is congruent to 1,2,10 or 17 mod 24. - _Mason Korb_, Oct 07 2018 [Proof: a^4 + b^4 = 1 or 2 (mod 16, 3 or 5) if a, b have no common factor. - _Martin Fuller_, Feb 01 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000572 | OEIS-A003959

If n = Product p(k)^e(k) then a(n) = Product (p(k)+1)^e(k), a(1) = 1.

Sum_{n>0} a(n)/n^s = Product_{p prime} 1/(1-p^(-s)-p^(1-s)) (conjectured). - _Ralf Stephan_, Jul 07 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marc LeBrun_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000573 | OEIS-A003961

Completely multiplicative with a(prime(k)) = prime(k+1).

Meyers (see Guy reference) conjectures that for all r >= 1, the least odd number not in the set {a(i): i < prime(r)} is prime(r+1). - _N. J. A. Sloane_, Jan 08 2021 Meyers' conjecture would be refuted if and only if for some r there were such a large gap between prime(r) and prime(r+1) that there existed a composite c for which prime(r) < c < a(c) < prime(r+1), in which case (by Bertrand's postulate) c would necessarily be a term of A246281. - _Antti Karttunen_, Mar 29 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marc LeBrun_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000574 | OEIS-A003989

Triangle T from the array A(x, y) = gcd(x,y), for x >= 1, y >= 1, read by antidiagonals.

Imagine a torus having regular polygonal cross-section of m sides. Now, break the torus and twist the free ends, preserving rotational symmetry, then reattach the ends. Let n be the number of faces passed in twisting the torus before reattaching it. For example, if n = m, then the torus has exactly one full twist. Do this for arbitrary m and n (m > 1, n > 0). Now, count the independent, closed paths on the surface of the resulting torus, where a path is "closed" if and only if it returns to its starting point after a finite number of times around the surface of the torus. Conjecture: this number is always gcd(m,n). NOTE: This figure constitutes a group with m and n the binary arguments and gcd(m,n) the resulting value. Twisting in the reverse direction is the inverse operation, and breaking & reattaching in place is the identity operation. - _Jason Richardson-White_, May 06 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marc LeBrun_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000575 | OEIS-A003994

Sequence b_4 (n) arising from homology of partitions with even number of blocks.

Colin Barker's conjecture confirmed by Sean A. Irvine's formula. - _Ray Chandler_, Jul 06 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | Sheila Sundaram (sheila(AT)paris-gw.cs.miami.edu) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000576 | OEIS-A004000

RATS: Reverse Add Then Sort the digits applied to previous term, starting with 1.

It is conjectured that no matter what the starting term is, repeatedly applying RATS leads either to this sequence or into a cycle of finite length, such as those in A066710 and A066711.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000577 | OEIS-A004001

Hofstadter-Conway $10000 sequence: a(n) = a(a(n-1)) + a(n-a(n-1)) with a(1) = a(2) = 1.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000578 | OEIS-A004044

The classic football pool problem: size of minimal covering code in {0,1,2}^n with covering radius 1.

Conjecture: a((3^m+1)/2) = 3^((3^m+1)/2 - m) for m > 0; i.e., a((3^m-1)/2 + 1) = 3 * a((3^m-1)/2) for m > 0. - _Thomas Ordowski_, Jul 10 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000579 | OEIS-A004061

Numbers k such that (5^k - 1)/4 is prime.

With the addition of the 19th prime in the sequence, the new best linear fit to the sequence has G=0.4723, which is slightly closer to the conjectured limit of G=0.56145948, A080130 (see link for Generalized Repunit Conjecture). - _Paul Bourdelais_, Apr 30 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000580 | OEIS-A004062

Numbers k such that (6^k - 1)/5 is prime.

With this 16th prime, the base 6 repunits have an average (best linear fit) occurrence rate of G = 0.4948 which seems to be converging to the conjectured rate of 0.56146 (see ref).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000581 | OEIS-A004081

a(n) = n-th positive integer such that only one integer lies between exp(s(m)) and exp(s(m+1)), where s(m) = 1 + 1/2 + 1/3 + . . . + 1/m.

Conjecturally, a(n) = floor(n/(2 - exp(g)) - 1/2 + exp(g)/(24n)), where g is the Euler-Mascheroni constant. - David W. Cantrell (DWCantrell(AT)sigmaxi.net), Aug 11 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000582 | OEIS-A004108

Number of n-node unlabeled connected graphs without endpoints.

Also number of n-node unlabeled connected mating graphs, cf. A006024 and A092430 (conjectured by Vladeta Jovovic, proved by G. Kilibarda). - _Vladeta Jovovic_, Oct 07 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000583 | OEIS-A004109

Number of connected trivalent (or cubic) labeled graphs with 2n nodes.

Conjecture: a(n) ~ 2^(n + 1/2) * 3^n * n^(3*n) / exp(3*n+2). - _Vaclav Kotesovec_, Feb 17 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000584 | OEIS-A004123

Number of generalized weak orders on n points.

Conjecture: Let k be a positive integer. The sequence obtained by reducing a(n) modulo k is eventually periodic with the period dividing phi(k) = A000010(k). For example, modulo 7 we obtain the sequence [1, 2, 3, 4, 2, 0, 0, 2, 3, 4, 2, 0, 0, 2, 3, 4, 2, 0, 0, ...] with an apparent period of 6 = phi(7) starting at a(2). Cf. A000670. More generally, we conjecture that the same property holds for integer sequences having an e.g.f. of the form G(exp(x) - 1), where G(x) is an integral power series. (End) Conjectural o.g.f. as a continued fraction of Stieltjes type: 1/(1 - 2*x/(1 - 3*x/(1 - 4*x/(1 - 6*x/(1 - ... - 2*n*x/(1 - 3*n*x/(1 - ...))))))). - _Peter Bala_, Jul 08 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000585 | OEIS-A004129

Postage stamp problem: largest m such that there exists an n-subset S of nonnegative integers such that 1,...,m can be expressed as a sum of two distinct elements of S.

The g.f. (z^4+z^3+2*z^2+2*z+1)*(z^2+z+1)/((z-1)*(z^5+z^4+z^3-z-1)) conjectured by _Simon Plouffe_ in his 1992 dissertation is wrong. - _Sean A. Irvine_, Nov 21 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000586 | OEIS-A004130

Numerators in expansion of (1-x)^{-1/4}.

a(n) = prod(k=1, n, (4k-3)/k * 2^A007814(k)), proved by _Mitch Harris_, following a conjecture by _Ralf Stephan_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000587 | OEIS-A004134

Denominators in expansion of (1-x)^{-1/4} are 2^a(n).

a(n) = 3*n - A000120(n). Recurrence: a(2n) = a(n) + 3n, a(2n+1) = a(n) + 3n + 2. Proved by _Mitch Harris_, following a conjecture by _Ralf Stephan_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000588 | OEIS-A004137

Maximal number of edges in a graceful graph on n nodes.

If the conjecture is true that an optimal ruler with more than 12 segments is a Wichmann ruler then the sequence continues 232, 251, 270, 289, 308, 327, ... - _Peter Luschny_, Oct 09 2011 [updated to take the verifications of Robison into account, Oct 01 2015]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000589 | OEIS-A004138

From a counter moving problem.

G.f. = -(-1+z^2-4*z^3+2*z^4)/((z-1)*(2*z^4-z^3+z^2+z-1)). [Conjectured (correctly) by _Simon Plouffe_ in his 1992 dissertation.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000590 | OEIS-A004215

Numbers that are the sum of 4 but no fewer nonzero squares.

Conjecture: The number of terms < 2^n is A023105(n) - 2. - _Tilman Neumann_, Sep 20 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _J. H. Conway_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000591 | OEIS-A004277

1 together with positive even numbers.

Conjecturally, the differences of two consecutive primes (without repetition). - _Juri-Stepan Gerasimov_, Nov 09 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000592 | OEIS-A004395

Ratios of successive terms are 1,1,2,3,3,4,5,5,6,7,7,...

Conjecture D-finite with recurrence 243*a(n) -216*a(n-1) +54*(-2*n+3)*a(n-2) +6*(-12*n^3+60*n^2-87*n+28)*a(n-3) +4*(2*n-5)*(2*n-7)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Sep 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000593 | OEIS-A004402

Expansion of 1 / Sum_{n=-oo..oo} x^(n^2).

Conjecture: Sum_{k>=0} a(k) / exp(k*Pi) = (1/Pi^(1/4))*Gamma(3/4) = A327995. - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000594 | OEIS-A004403

Expansion of 1/theta_3(q)^2 in powers of q.

Conjecture: Sum_{k>=0} a(k) / exp(k*Pi) = (1/Pi^(1/2))*Gamma(3/4)^2 = A096427. - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000595 | OEIS-A004405

Expansion of 1 / (Sum_{n=-oo..oo} x^(n^2))^4.

Conjecture: Sum_{k>=0} a(k) / exp(k*Pi) = (1/Pi)*Gamma(3/4)^4 = A293238. - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000596 | OEIS-A004432

Numbers that are the sum of 3 distinct nonzero squares.

Conjecture: a number n is a sum of 3 squares, but not a sum of 3 distinct nonzero squares (i.e., is in A004432 but not A000408), if and only if it is of the form 4^j*s, where j >= 0 and s in {1, 2, 3, 5, 6, 9, 10, 11, 13, 17, 18, 19, 22, 25, 27, 33, 34, 37, 43, 51, 57, 58, 67, 73, 82, 85, 97, 99, 102, 123, 130, 163, 177, 187, 193, 267, 627, 697}. - _Jeffrey Shallit_, Jan 15 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000597 | OEIS-A004484

Sprague-Grundy values for game of Wyt Queens.

Conjectures from _Chai Wah Wu_, Apr 05 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000598 | OEIS-A004607

Infinitary sociable numbers (smallest member of cycle).

It is conjectured that there are no missing terms in the data, but it was not proven. For example, it is not known that the infinitary aliquot sequence that starts at 840 does not reach 840 again (see A361421). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000599 | OEIS-A004642

Powers of 2 written in base 3.

Sloane (1973) conjectured a(n) always has a 0 between the most and least significant digits if n > 15 (see A102483 and A346497). Erdős (1978) conjectured that for n > 8 a(n) has at least one 2 (see link to Terry Tao's blog). - _Dmitry Kamenetsky_, Jan 10 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000600 | OEIS-A004735

Denominator of average distance traveled by n-dimensional fly.

a(2n) = A001803(n) (conjectured). - _Ralf Stephan_, Mar 10 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000601 | OEIS-A004737

Concatenation of sequences (1,2,...,n-1,n,n-1,...,1) for n >= 1.

Let U_1(t)=1, U_2(t)=2*t, and U_r(t)=2*t*U_(r-1)(t)-U(r-2)(t), r>2, be Chebyshev polynomials of the second kind. For q>1 an integer, let N=2*q and x_k=cos((2*k-1)*Pi/N), and define the ordered column vectors V_k=[U_k(x_1), U_k(x_2), ..., U_k(x_q)]^T, k=1,...,q, where A^T denotes the transpose of matrix A. Let E_N=[V_1, V_2, ..., V_q] be the q X q matrix formed from the ordered components of the V_k. E_N contains the joint spectra of the Danzer basis (see [Jeffery]) associated with N. Let M_N=(1/q)*[E_N]^T*E_N. For the trivial case q=1, let M_2=[1]. CONJECTURE: E_N and M_N are always integral and symmetric, with M_N having diagonal entries {1,2,...} beginning at entries 1,j (j odd) in the first row and i,1 (i odd) in the first column and with zeros elsewhere. If N is allowed to increase without bound, and assuming the conjecture is true, then triangle A004737 emerges in its entirety from the successive antidiagonals containing those entries [M_N]_(i,j) such that i+j=2*v, for each v in {1,2,...,floor((q+1)/2)}. For example, for N=18 and q=9 (omitting the zeros for clarity),

猜想/问题提及段 | historical_or_current_mention_unreviewed | R. Muller | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000602 | OEIS-A004793

a(1)=1, a(2)=3; a(n) is least k such that no three terms of a(1), a(2), ..., a(n-1), k form an arithmetic progression.

a(n) = (3-n)/2 + 2*floor(n/2) + Sum_{k=1..n-1} 3^A007814(k)/2 = A003278(n) + [n is even], proved by Lawrence Sze, following a conjecture by _Ralf Stephan_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000603 | OEIS-A004977

Sum of digits of n-th term in Look and Say sequence A005150.

It appears that the ratio of consecutive terms approaches Conway's constant 1.303.. (A014715). The terms divided by the numbers of added digits also would tend to a constant, i.e. A004977(n)/A005341(n)->const. If the digits in A005150

occur with constant probabilities c1, c2, c3 then A004977(n)=A005341(n)*(c1+2*c2+3*c3) and this conjecture entails the convergences noted here. - _Alexandre Losev_, Aug 31 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000604 | OEIS-A004981

a(n) = (2^n/n!) * Product_{k=0..n-1} (4*k + 1).

Conjecture: a(p*n) = a(n) (mod p^2) for prime p = 1 (mod 4) and all positive integers n. Cf. A004982 and A298799. - _Peter Bala_, Dec 22 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | Joe Keane (jgk(AT)jgk.org) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000605 | OEIS-A004982

a(n) = (2^n/n!) * Product_{k=0..n-1} (4*k + 3).

Conjecture: a(p*n) = a(n) (mod p^2) for prime p == 1 (mod 4) and all positive integers n. Cf. A004981. - _Peter Bala_, Dec 22 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | Joe Keane (jgk(AT)jgk.org) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000606 | OEIS-A005007

Number of cubic (i.e., regular of degree 3) generalized Moore graphs with 2n nodes.

A generalized Moore graph is a regular graph of degree r where the counts of vertices at each distance from any vertex are 1, r, r(r-1), r(r-1)^2, r(r-1)^3, ... with the last distance having every other vertex. That is, all the levels are full except possibly the last which must have the rest. Alternatively, the girth is as great as the naive bound allows and the diameter is as little as the naive bound allows. Or, the average distance between pairs of vertices achieves the naive lower bound. As far as I know, it is an open problem if there are infinitely many generalized Moore graphs of each degree. - _Brendan McKay_, Oct 06 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000607 | OEIS-A005043

Riordan numbers: a(n) = (n-1)*(2*a(n-1) + 3*a(n-2))/(n+1).

The Rubey and Stump reference proves a refinement of a conjecture of René Marczinzik, which they state as: "The number of 2-Gorenstein algebras which are Nakayama algebras with n simple modules and have an oriented line as associated quiver equals the number of Motzkin paths of length n. Moreover, the number of such algebras having the double centraliser property with respect to a minimal faithful projective-injective module equals the number of Riordan paths, that is, Motzkin paths without level-steps at height zero, of length n." - _Eric M. Schmidt_, Dec 16 2017 Conjecture: a(n) = A002426(n) - A005717(n), n > 0. - _Mikhail Kurkov_, Feb 24 2019 [The conjecture is true. - _Amiram Eldar_, May 17 2024] Conjectures: for n >= 1, n divides a(2*n+1) and 2*n-1 divides a(2*n). (End) The conjectures of Bala are true (see Fried link). - _Sela Fried_, Nov 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000608 | OEIS-A005045

Number of restricted 3 X 3 matrices with row and column sums n.

G.f.: -x^2*(-x^5+x^6-x^3+x+1)/((x^2+1)*(x^2+x+1)*(x+1)^2*(x-1)^5). This was conjectured by _Simon Plouffe_ in his 1992 dissertation and is now known to be correct, although it may be that all the details of the proof have not been written down. See the Mathar link for details.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000609 | OEIS-A005046

Number of partitions of a 2n-set into even blocks.

Conjecture: Taking the sequence modulo an integer k gives an eventually periodic sequence. For example, the sequence taken modulo 10 is [1, 1, 4, 1, 9, 6, 9, 1, 4, 1, 9, 6, 9, 1, 4, 1, 9, 6, 9, ...], with an apparent period [1, 4, 1, 9, 6, 9] beginning at a(1), of length 6. Cf. A006154. - _Peter Bala_, Apr 12 2023 Conjecture: a(n) = 2 * exp(-1) * Sum_{k=1..oo} k^(2*n) * Bessel(k, 1) for n > 0. - _Velin Yanev_, Feb 09 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000610 | OEIS-A005103

Minimal determinant of n-dimensional norm 3 lattice.

The g.f. (1 + z + 2*z^2 + 2*z^3 + 6*z^4)/(1 - 2*z + 2*z*^3) conjectured by _Simon Plouffe_ in his 1992 dissertation is almost certainly wrong.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000611 | OEIS-A005133

Number of index n subgroups of modular group PSL_2(Z).

Grady conjectures that: I) if n is odd, then a(n) and a(2*n) have the same parity and II) a(4*n) is even (cf. p. 94 from his article).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000612 | OEIS-A005148

Sequence of coefficients arising in connection with a rapidly converging series for Pi.

Conjecture: The following 2 definitions give the same sequence: (1) numbers k such that 8^m is the highest power of 2 dividing a(k), and (2) numbers k such that k has exactly (m+1) 1's in its binary representation. A018900 is the special case m=1. - _Benoit Cloitre_, Jun 22 2002, edited by _Hugo Pfoertner_, Aug 21 2021 Conjecture: There are polynomials P_k(x) such that P_k(m) = the constant term of j_m(tau)^k where j_m is modular for the Hecke group G(lambda_m), j_3 is the Klein invariant j with constant term 744, and P_k(x) = a(k+1) times a product of monic polynomials. - _Barry Brent_, Nov 25 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ and _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000613 | OEIS-A005150

Look and Say sequence: describe the previous term! (method A - initial term is 1).

Conjecture: DC(a(n)) ~ k * (Conway's constant)^n where k is approximately 1.021... and DC denotes the number of digit changes in the decimal representation of n (e.g., DC(13112221)=4 because 1->3, 3-1, 1->2, 2->1). - _Bill McEachen_, May 09 2025 Conjecture: lim_{n->infinity} (c2+c3-c1)/(c1+c2+c3) = 0.01 approximately, where ci is the number of appearances of 'i' in a(n). - _Ya-Ping Lu_, Jun 05 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000614 | OEIS-A005152

Rotation distance between binary trees on n nodes.

Sleator et al. conjecture that a(n) = 2n-6 for all n >= 11.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000615 | OEIS-A005157

Number of totally symmetric plane partitions that fit in an n X n X n box.

Conjectures: if p == 1 (mod 6) is prime then a(p) == 2^((p+5)/6) (mod p^2); if p == 5 (mod 6) is prime then a(p) == 2^((p+1)/6) (mod p^2) (checked up to p = 1009). - _Peter Bala_, Feb 17 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000616 | OEIS-A005158

Number of alternating sign n X n matrices invariant under a half-turn.

Robbins gives simple (conjectured) formulas related to this sequence in Section 3.3. a(n) = a(n-1) * (1 + [n even]/3) * C(n\2*3, n\2) / C(n\2*2, n\2) for all n > 1, where C(.,.) are the binomial coefficients, n\2 := floor(n/2) and [n even] = 1 if n is even, 0 else (Iverson bracket). [From Robbins conjectured(!) formulas.] - _M. F. Hasler_, Jun 15 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000617 | OEIS-A005161

Number of alternating sign 2n+1 X 2n+1 matrices symmetric with respect to both horizontal and vertical axes (VHSASM's).

Robbins gives a simple (conjectured) formula, which was proven by Okada.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000618 | OEIS-A005162

Number of alternating sign n X n matrices symmetric with respect to both diagonals.

Robbins gives a simple (conjectured) formula.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000619 | OEIS-A005164

Number of alternating sign 2n+1 X 2n+1 matrices invariant under all symmetries of the square.

Hagendorf and Liénardy give a (conjectured) formula in terms of multiple contour integrals. - _Jean Liénardy_, Aug 15 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000620 | OEIS-A005165

Alternating factorials: n! - (n-1)! + (n-2)! - ... 1!.

Conjecture: for n > 2, smallest prime divisor of a(n) > n. - _Gerald McGarvey_, Jun 19 2004 Conjecture: For n>1, a(n) is the number of lattice paths from (0,0) to (n+1,0) that do not cross above y=x or below the x-axis using up-steps +(1,a) and down-steps +(1,-b) where a and b are positive integers. For example, a(3) = 5: [(1,1)(1,1)(1,1)(1,-3)], [(1,1)(1,-1)(1,3)(1,-3)], [(1,1)(1,-1)(1,2)(1,-2)], [(1,1)(1,-1)(1,1)(1,-1)] and [(1,1)(1,1)(1,-1)(1,-1)]. - _Nicholas Ham_, Aug 23 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000621 | OEIS-A005206

Hofstadter G-sequence: a(0) = 0; a(n) = n - a(a(n-1)) for n > 0.

a(n) = A293688(n)/(n+1) for n >= 0 (conjectured). - _Enrique Navarrete_, Oct 15 2017 Conjectured by _Chunqing Liu_, Aug 01 2023 and proved by _Roman Le Lan_, Oct 26 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000622 | OEIS-A005234

Primes p such that 1 + product of primes up to p is prime.

Conjecture: if p# + 1 is a prime number, then the next prime is less than p# + exp(1)*p. - _Arkadiusz Wesolowski_, Feb 20 2013 Conjecture: if p# + 1 is a prime, then the next prime is less than p# + p^2. - _Thomas Ordowski_, Apr 07 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000623 | OEIS-A005251

a(0) = 0, a(1) = a(2) = a(3) = 1; thereafter, a(n) = a(n-1) + a(n-2) + a(n-4).

Conjecture: a(n+1) + |A078065(n)| = 2*A005314(n+1). - _Creighton Dement_, Dec 21 2004 The conjecture is true (see Fried link). - _Sela Fried_, Nov 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000624 | OEIS-A005309

Fermionic string states.

The g.f. -(1-2*z+2*z^2)/(-1+2*z) conjectured by _Simon Plouffe_ in his 1992 dissertation is not correct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000625 | OEIS-A005310

Fermionic string states.

The g.f. 2*(1-2*z+2*z^2)/(-1+2*z)/(z-1) conjectured by _Simon Plouffe_ in his 1992 dissertation is unlikely to be correct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000626 | OEIS-A005313

Maximal sum of inverse squares of the singular values of triangular anti-Hadamard matrices of order n.

G.f.: [x(1-x-3x^2+x^3)]/[(1-3x+x^2)(1+x)(1-x)^2]. - Conjectured by _Simon Plouffe_ in his 1992 dissertation. [This is in fact the correct g.f. - _N. J. A. Sloane_, Jan 28 2022]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000627 | OEIS-A005318

Conway-Guy sequence: a(n + 1) = 2a(n) - a(n - floor(1/2 + sqrt(2n))).

Conway and Guy conjecture that the set of k numbers {s_i = a(k) - a(k-i) : 1 <= i <= k} has the property that all its subsets have distinct sums - see Guy's book. These k-sets are the rows of A096858. [This conjecture has apparently now been proved by Bohman. - I. Halupczok (integerSequences(AT)karimmi.de), Feb 20 2006]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000628 | OEIS-A005319

a(n) = 6*a(n-1) - a(n-2).

(-1)^(n+1) = A090390(n+1) + A001542(n+1) + A046729(n) - a(n) (conjectured). - _Creighton Dement_, Nov 17 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000629 | OEIS-A005321

Upper triangular n X n (0,1)-matrices with no zero rows or columns.

Two conjectural continued fractions for the o.g.f.:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000630 | OEIS-A005329

a(n) = Product_{i=1..n} (2^i - 1). Also called 2-factorial numbers.

Conjecture: this sequence is the inverse binomial transform of A075272 or, equivalently, the inverse binomial transform of the BinomialMean transform of A075271. - _John W. Layman_, Sep 12 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000631 | OEIS-A005333

Number of 2-colored connected labeled graphs with n vertices of the first color and n vertices of the second color.

Conjecture: if n > 1, then a(n) is the number of labeled digraphs D (allowing self-loops) with n vertices such that D|D' and D'|D are (strongly) connected (see preliminaries of Broere et al.). - _Lorenzo Sauras Altuzarra_, Sep 17 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000632 | OEIS-A005374

Hofstadter H-sequence: a(n) = n - a(a(a(n-1))).

Conjecture: a(n) = floor(c*n) + 0 or 1, where c is the real root of x^3+x-1 = 0, c=0.682327803828019327369483739... - _Benoit Cloitre_, Nov 05 2002 [Proved by Letouzey, see Letouzey link. - _Pierre Letouzey_, Feb 20 2025], [Also proved in Shallit (2025). - _Jeffrey Shallit_, Mar 09 2025]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000633 | OEIS-A005375

a(0) = 0; a(n) = n - a(a(a(a(n-1)))) for n > 0.

a(n) = floor(c*n) + (-1) or 0 or 1 or 2, where c is the positive real root of x^4+x-1 = 0, c=0.724491959000515611588372282... (Conjectured with just 0 or 1 by _Benoit Cloitre_, Nov 05 2002; fixed and proved by Letouzey, see Letouzey link]. NB: see for instance a(120) = 88 for a difference of 2 and a(243) = 175 for a difference of -1). - _Pierre Letouzey_, Feb 20 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000634 | OEIS-A005376

a(n) = n - a(a(a(a(a(n-1))))).

Conjecture: a(n) is approximately c*n, where c is the real root of x^5+x-1 = 0, c=0.754877666246692760049508896... - _Benoit Cloitre_, Nov 05 2002 a(n) = c*n + O(ln(n)), with c conjectured by Benoit Cloitre above; see Letouzey link and Dilcher 1993. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000635 | OEIS-A005377

Number of low discrepancy sequences in base 4.

G.f.: z^4 * (z^2+1) * (z^4-z^2+1) / (z-1)^2; [Conjectured by _Simon Plouffe_ in his 1992 dissertation, but is incorrect.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000636 | OEIS-A005379

The male of a pair of recurrences.

The g.f. -z^2*(-1-z^3-z^6-z-z^4-z^7+z^8)/(z+1)/(z^2+1)/(z^4+1)/(z-1)^2, conjectured by _Simon Plouffe_ in his 1992 dissertation is incorrect: the coefficient of z^33 in the g.f. is 21, but a(33) = 20. (Discovered by Sahand Saba, Jan 14 2013.) - _Frank Ruskey_, Jan 16 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000637 | OEIS-A005381

Numbers k such that k and k-1 are composite.

Conjecture: pi(n)=Sum_{k=1..n} k mod a(m) mod a(m-1) ... mod a(1) mod 2, for all values 1<n<=a(m), where the mod are evaluated from left to right. Verified for first 10000 a(n). - _Benedict W. J. Irwin_, May 04 2016 For an attempt at a proof for the conjecture above, see the link. If it is true, then for n>2, isprime(n)=(n mod x) mod 2, where x is the largest a(n)<=n. - _Benedict W. J. Irwin_, May 06 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000638 | OEIS-A005385

Safe primes p: (p-1)/2 is also prime.

Conjecture: there is a strengthening of the Bertrand postulate for n >= 24: the interval (n, 2*n) contains a safe prime. It has been tested by _Peter J. C. Moses_ up to n = 10^7. - _Vladimir Shevelev_, Jul 06 2015 From the fourth entry onward, do these correspond to Smarandache's problem 34 (see A007931 link), specifically values which cannot be used (do not meet conditions) to confirm the conjecture? - _Bill McEachen_, Sep 29 2016 It is conjectured that there are infinitely many safe primes, and their estimated asymptotic density ~ 2C/(log n)^2 (where C = 0.66... is the twin prime constant A005597) converges to the actual value as far as we know. - _M. F. Hasler_, Jun 14 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000639 | OEIS-A005408

The odd numbers: a(n) = 2*n + 1.

Conjecture on primes with one coach (A216371) relating to the odd integers: iff an integer is in A216371 (primes with one coach either of the form 4q-1 or 4q+1, (q > 0)); the top row of its coach is composed of a permutation of the first q odd integers. Example: prime 19 (q = 5), has 5 terms in each row of its coach: 19: [1, 9, 5, 7, 3] ... [1, 1, 1, 2, 4]. This is interpreted: (19 - 1) = (2^1 * 9), (19 - 9) = (2^1 * 5), (19 - 5) = (2^1 - 7), (19 - 7) = (2^2 * 3), (19 - 3) = (2^4 * 1). - _Gary W. Adamson_, Sep 09 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000640 | OEIS-A005411

Number of non-vanishing Feynman diagrams of order 2n for the electron or the photon propagators in quantum electrodynamics.

Conjectural o.g.f. A(x) as a continued fraction:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000641 | OEIS-A005412

Number of non-vanishing Feynman diagrams of order 2n for the vacuum polarization (the proper two-point function of the photon) and for the self-energy (the proper two-point function of the electron) in quantum electrodynamics (QED).

Conjectural o.g.f. as a continued fraction:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000642 | OEIS-A005427

Josephus problem: numbers m such that, when m people are arranged on a circle and numbered 1 through m, the final survivor when we remove every 4th person is one of the first three people.

Is this the same as A072493 with its first 8 terms removed? See also the similar conjecture concerning A005428 and A073941. I do not have a general proof of this conjecture though I strongly believe that Schuh's (1968) way of solving the case k = 3 (see pp. 373-375 and 377-379, where he provides two methods of solution) may provide clues for proving the conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000643 | OEIS-A005439

Genocchi medians (or Genocchi numbers of second kind).

For any positive integer n, a(n) is (-1)^n times the permanent of the 2n X 2n matrix M with M(j, k) = floor((2*j-k-1)/(2*n)). This former conjecture of Luschny, inspired by a conjecture of _Zhi-Wei Sun_ in A036968, was proven by Fu, Lin and Sun (see link). - _Peter Luschny_, Sep 07 2021 [updated Sep 24 2021]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000644 | OEIS-A005465

Number of n-dimensional hypotheses allowing for conditional independence.

Number of n-dimensional hypotheses allowing for conditional independence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000645 | OEIS-A005479

Prime Lucas numbers (cf. A000032).

Conjectures: 7, 47 and 2207 are the only a(n) mod 10 = 7. They are also the only a(n) values where the Lucas index is not a prime. See A001606 for the Lucas index values of these primes. See A266587 for the divisibility of Lucas numbers by powers of primes. - _Richard R. Forberg_, Mar 24 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000646 | OEIS-A005517

Smallest label f(T) given to a rooted tree T with n nodes in Matula-Goebel labeling.

G.f.: z*(1+2*z+3*z^2-z^4)/(1-5*z^3) (conjectured by _Simon Plouffe_).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000647 | OEIS-A005520

Smallest number of complexity n: smallest number requiring n 1's to build using + and *.

The prime 353942783 = A189125(1) = 2*3 + (1 + 2^2*3^2)*(2 + 3^4(1 + 2*3^10)) is the smallest counterexample to Guy's first hypothesis on integer complexity. - _Jonathan Vos Post_, Mar 30 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000648 | OEIS-A005522

a(n) = 1 + L(n) + F(2*n-1) with {L(n)}_{n>=0} the Lucas numbers (A000032) and F(2*n-1)_{n>=0} the bisected Fibonacci numbers (A001519).

(Comment VI) We can now comment on the g.f., P(x), conjectured by _Simon Plouffe_ and mentioned in the MAPLE section below. Expanding P(x) as a series, we obtain the following g.f. of the a(n): 3 + 6x + 10x^2 + 21x^3 ... . In fact, it is easily checked that P(x) = 1/x (A(x)-4). This settles the Plouffe conjecture as follows: (i) A(x) is the rational function g.f. of a(n)_{n>=0}, (ii) P(x) is the rational function g.f. of a(n)_{n>=1}, and (iii) C(x)+L(x)+B(x) is the partial fraction g.f. of a(n)_{n>=0}. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000649 | OEIS-A005528

Størmer numbers or arc-cotangent irreducible numbers: numbers k such that the largest prime factor of k^2 + 1 is >= 2*k.

Also numbers k such that k^2 + 1 has a primitive divisor, hence (by Everest & Harman, Theorem 1.4) 1.1n < a(n) < 1.88n for large enough n. They conjecture that a(n) ~ cn where c = 1/log 2 = 1.4426.... - _Charles R Greathouse IV_, Nov 15 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _J. H. Conway_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000650 | OEIS-A005555

A finite sequence associated with the Lie algebra E_8.

The g.f. (5-6*z+2*z**2)/(z-1)**4 conjectured by _Simon Plouffe_ in his 1992 dissertation is wrong.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000651 | OEIS-A005558

a(n) is the number of n-step walks on square lattice such that 0 <= y <= x at each step.

Number of n-step walks that start at the origin, constrained to stay in the first octant (0 <= y <= x). (Conjectured) - _Benjamin Phillabaum_, Mar 11 2011, corrected by _Robert Israel_, Oct 07 2015 E.g.f (conjectured): Bessell(1,2*x)*(Bessell(0,2*x)+Bessell(1,2*x))/x. - _Benjamin Phillabaum_, Feb 25 2011 Conjecture: (2*n+1)*(n+3)*(n+2)*a(n) - 4*(2*n^2+4*n+3)*a(n-1) - 16*n*(2*n+3)*(n-1)*a(n-2) = 0. - _R. J. Mathar_, Apr 02 2017 Conjecture: (n+3)*(n+2)*a(n) - 4*(n^2+3*n+1)*a(n-1) + 16*(-n^2+n+1)*a(n-2) + 64*(n-1)*(n-2)*a(n-3) = 0. - _R. J. Mathar_, Apr 02 2017 a(n) = Sum_{k=0..floor(n/2)} n!/(k!*k!*(floor(n/2)-k)!*(floor((n+1)/2)-k)!*(k+1)) (conjectured). - _Roger Ford_, Aug 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000652 | OEIS-A005559

Number of walks on square lattice. Column y=1 of A052174.

Conjecture: -(n-1)*(2*n+1)*(n+4)*(n+3)*a(n) +4*(n+1)*(2*n^2+4*n+9)*a(n-1) +16*n*(n-1)*(2*n+3)*(n+1)*a(n-2)=0. - _R. J. Mathar_, Apr 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000653 | OEIS-A005560

Number of walks on square lattice. Column y=2 of A052174.

Conjecture: (n-1)*(n-2)*(2*n+1)*(n+5)*(n+4)*a(n) -4*n*(n+1)*(2*n^2+4*n+19)*a(n-1) -16*n^2*(n-1)*(2*n+3)*(n+1)*a(n-2)=0. - _R. J. Mathar_, Apr 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000654 | OEIS-A005561

Number of walks on square lattice. Column y=3 of A052174.

Conjecture: (n-2)*(n-3)*(2*n+1)*(n+6)*(n+5)*a(n) - 4*n*(n+1)*(2*n^2+4*n+33)*a(n-1) - 16*n^2*(n-1)*(2*n+3)*(n+1)*a(n-2) = 0. - _R. J. Mathar_, Apr 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000655 | OEIS-A005562

Number of walks on square lattice. Column y=4 of A052174.

Conjecture: (n-3)*(n-4)*(2*n+1)*(n+7)*(n+6)*a(n) - 4*n*(n+1)*(2*n^2+4*n+51)*a(n-1) - 16*n^2*(n-1)*(2*n+3)*(n+1)*a(n-2) = 0. - _R. J. Mathar_, Apr 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000656 | OEIS-A005563

a(n) = n*(n+2) = (n+1)^2 - 1.

Erdős conjectured that n^2 - 1 = k! has a solution if and only if n is 5, 11 or 71 (when k is 4, 5 or 7). Conjecture: When using the Sieve of Eratosthenes and sieving (n+1..a(n)), with divisors (1..n) and n>0, there will be no more than a(n-1) composite numbers. - _Fred Daniel Kline_, Apr 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000657 | OEIS-A005572

Number of walks on cubic lattice starting and finishing on the xy plane and never going below it.

Conjecture: a(n) is even except for n of the form 2*(2^k - 1). [added Feb 03 2024: the conjecture follows from the formula a(n) = Sum_{k = 0..n} 2^(n-k)*binomial(n, k)*Catalan(k+1) given above.] (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000658 | OEIS-A005573

Number of walks on cubic lattice (starting from origin and not going below xy plane).

Conjecture: a(n)= 6^n * hypergeom([1/2,-n],[2], 2/3). - _Benjamin Phillabaum_, Feb 20 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000659 | OEIS-A005574

Numbers k such that k^2 + 1 is prime.

Hardy and Littlewood conjectured that the asymptotic number of elements in this sequence not exceeding n is approximately c*sqrt(n)/log(n) for some constant c. - _Stefan Steinerberger_, Apr 06 2006 Apparently Goldbach conjectured that any a > 1 from this sequence can be written as a=b+c where b and c are in this sequence (Lemmermeyer link below). - _Jeppe Stig Nielsen_, Oct 14 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000660 | OEIS-A005598

a(n) = 1 + Sum_{i=1..n} (n-i+1)*phi(i).

Conjectured to be the number of regions in a Farey fan of order n. See A360042 for further details. - _Scott R. Shannon_, Jan 24 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000661 | OEIS-A005653

Lexicographically least increasing sequence, starting with 2, such that the sum of two distinct terms of the sequence is never a Fibonacci number.

Jon E. Schoenfield conjectured, and Jeffrey Shallit proved (using the Walnut theorem prover) the characterization in the title. - _Jeffrey Shallit_, Nov 19 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ and _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000662 | OEIS-A005654

Number of bracelets (turn over necklaces) with n red, 1 pink and n-1 blue beads; also reversible strings with n red and n-1 blue beads; also next-to-central column in Losanitsch's triangle A034851.

Conjecture: n*(n-2)*a(n) - (5*n-3)*(n-2)*a(n-1) + 4*(n-2)*a(n-2) + 4*(5*n^2-27*n+37)*a(n-3) - 8*(2*n-7)*(n-4)*a(n-4) = 0. - _R. J. Mathar_, Nov 09 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000663 | OEIS-A005669

Indices of primes where largest gap occurs.

Conjecture: log a(n) ~ n/2. That is, record prime gaps occur about twice as often as records in an i.i.d. random sequence of comparable length (see arXiv:1709.05508 for a heuristic explanation). - _Alexei Kourbatov_, Mar 28 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000664 | OEIS-A005685

Number of Twopins positions.

G.f.: (-x^4*(x^7+x^6+x^5+2*x^4-x^3+x^2-1))/((x^4+x^2-1)*(x^2-x+1)*(x^2+x-1)). - Conjectured by _Simon Plouffe_ in his 1992 dissertation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000665 | OEIS-A005726

Quadrinomial coefficients.

Conjecture: 2*(n-1)*(2*n+1)*(13*n-14)*a(n) +(-143*n^3+297*n^2-148*n+12) *a(n-1) -4*(n-1)*(26*n^2-41*n+9)*a(n-2) -16*(n-1)*(n-2)*(13*n-1) *a(n-3)=0. - _R. J. Mathar_, Nov 13 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000666 | OEIS-A005777

Rook polynomials.

Conjecture: G.f.: x + 4*x^2 - 15*x^3 + x^3*((33 - 69*x + 16*x^2 - 14*x^3 + 82*x^4 - 8*x^5 - 20*x^6 - 38*x^7 + 9*x^8 + 9*x^9 + 8*x^10) / ((1 - x)^2 * (1 - x - x^2 - x^3)^3)). - _Vaclav Kotesovec_, Aug 27 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000667 | OEIS-A005799

Generalized Euler numbers of type 2^n.

Conjectures: Cf. A000657. [added April 18 2024: for a proof of this conjecture see Fu et al., Section 4.3.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000668 | OEIS-A005819

Number of words of length n in a certain language.

a(n) = -1 + Sum_{i=1..n} (n-i+1)*phi(i) for n>=2, where phi is Euler's totient function (this is conjectured in the Duluck & Gouyou-Beauchamps paper, p. 399).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000669 | OEIS-A005820

3-perfect (triply perfect, tri-perfect, triperfect or sous-double) numbers: numbers such that the sum of the divisors of n is 3n.

If there exists an odd perfect number m (a famous open problem) then 2m would be 3-perfect, since sigma(2m) = sigma(2)*sigma(m) = 3*2m. - _Jens Kruse Andersen_, Jul 30 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000670 | OEIS-A005840

Expansion of (1-x)*e^x/(2-e^x).

Conjecture: A285868 (with offset 1) shows the associated connected threshold graphs. - _R. J. Mathar_, Apr 29 2019 Re: above conjecture - the number of connected threshold graphs on n labeled vertices is A317057 (see also A053525). [_David Galvin_, Oct 18 2021]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000671 | OEIS-A005842

a(n) = minimal integer m such that an m X m square contains non-overlapping squares of sides 1, ..., n (some values are only conjectures).

a(n) = minimal integer m such that an m X m square contains non-overlapping squares of sides 1, ..., n (some values are only conjectures).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000672 | OEIS-A005867

a(0) = 1; for n > 0, a(n) = (prime(n)-1)*a(n-1).

Conjecture: The g.f. for the prime(n+1)-rough numbers (A000027, A005408, A007310, A007775, A008364, A008365, A008366, A166061, A166063) is x*P(x)/(1-x-x^a(n)+x^(a(n)+1)), where P(x) is an order a(n) polynomial with symmetric coefficients (i.e., c(0)=c(n), c(1)=c(n-1), ...). - _Benedict W. J. Irwin_, Mar 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000673 | OEIS-A005908

a(n) = floor(phi*a(n-1)) + floor(phi*a(n-2)), where phi is the golden ratio.

Conjectures from _Chai Wah Wu_, Jun 13 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000674 | OEIS-A005943

Factor complexity (number of subwords of length n) of the Golay-Rudin-Shapiro binary word A020987.

Terms a(0)..a(63) were computed using the first 2^36 terms of the GRS sequence, and are consistent with Arndt's conjectured g.f. - _Sean A. Irvine_, Oct 12 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Jeffrey Shallit_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000675 | OEIS-A006012

a(0) = 1, a(1) = 2, a(n) = 4*a(n-1) - 2*a(n-2), n >= 2.

Conjecture: With offset 1, a(n) is the number of permutations on [n] with no subsequence abcd such that (i) bc are adjacent in position and (ii) max(a,c) < min(b,d). For example, the 4 permutations of [4] not counted by a(4) are 1324, 1423, 2314, 2413. - _David Callan_, Aug 27 2014 The conjecture of David Callan above is correct - with offset 1, a(n) is the number of permutations on [n] with no subsequence abcd such that (i) bc are adjacent in position and (ii) max(a,c) < min(b,d). - _Yonah Biers-Ariel_, Jun 27 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000676 | OEIS-A006014

a(n+1) = (n+1)*a(n) + Sum_{k=1..n-1} a(k)*a(n-k).

Conjecture: a(n) = Sum_{k=0..2^(n-1) - 1} A379818(k). - _Mikhail Kurkov_, Nov 19 2021 [Rewritten by _Mikhail Kurkov_, Mar 27 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000677 | OEIS-A006018

List of periods for game of Third One Lucky.

Conjectures from _Aidan Hennessey_, Apr 27 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000678 | OEIS-A006063

A card-arranging problem: values of n such that there exists a permutation p_1, ..., p_n of 1, ..., n such that i + p_i is a cube for every i.

Conjecture: a(n) = n + 124 for n >= 173, i.e. there is such a permutation for every n >= 173. Verified for 173 <= n <= 1000. - _Robert Israel_, Aug 28 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000679 | OEIS-A006064

Smallest junction number with n generators.

a(4) = 10^24 + 102 was conjectured by Narasinga Rao. a(5) = 10^1111111111124 + 102. - Conjectured by Narasinga Rao, confirmed by _Max Alekseyev_ and _N. J. A. Sloane_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000680 | OEIS-A006077

(n+1)^2*a(n+1) = (9n^2+9n+3)*a(n) - 27*n^2*a(n-1), with a(0) = 1 and a(1) = 3.

Conjecture: Let W(n) be the (n+1) X (n+1) Hankel-type determinant with (i,j)-entry equal to a(i+j) for all i,j = 0,...,n. If n == 1 (mod 3) then W(n) = 0. When n == 0 or 2 (mod 3), W(n)*(-1)^(floor((n+1)/3))/6^n is always a positive odd integer. - _Zhi-Wei Sun_, Aug 21 2013 Conjecture: Let p == 1 (mod 3) be a prime, and write 4*p = x^2 + 27*y^2 with x, y integers and x == 1 (mod 3). Then W(p-1) == (-1)^{(p+1)/2}*(x-p/x) (mod p^2), where W(n) is defined as the above. - _Zhi-Wei Sun_, Aug 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000681 | OEIS-A006093

a(n) = prime(n) - 1.

Conjecture: All the sums Sum_{k=s..t} 1/a(k) with 1 <= s <= t are pairwise distinct. In general, for any integers d >= -1 and m > 0, if Sum_{k=i..j} 1/(prime(k)+d)^m = Sum_{k=s..t} 1/(prime(k)+d)^m with 0 < i <= j and 0 < s <= t then we must have (i,j) = (s,t), unless d = m = 1 and {(i,j),(s,t)} = {(4,4),(8,10)} or {(4,7),(5,10)}. (Note that 1/(prime(8)+1)+1/(prime(9)+1)+1/(prime(10)+1) = 1/(prime(4)+1) and Sum_{k=5..10} 1/(prime(k)+1) = 1/(prime(4)+1) + Sum_{k=5..7} 1/(prime(k)+1).) - _Zhi-Wei Sun_, Sep 09 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000682 | OEIS-A006134

a(n) = Sum_{k=0..n} binomial(2*k,k).

a(n) = A013580(2n+1,n+1) (conjectured). a(n) = A231147(2n+1,n+1) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000683 | OEIS-A006141

Number of integer partitions of n whose smallest part is equal to the number of parts.

Plouffe in his 1992 dissertation conjectured that this has g.f. = (1+z+z^4+2*z^5-z^3-z^8+3*z^10-z^7+z^9)/(1+z-z^4-2*z^3-z^8+z^10), but _Michael Somos_ pointed out on Jan 22 2008 that this is false.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000684 | OEIS-A006154

Number of labeled ordered partitions of an n-set into odd parts.

Conjecture: taking the sequence modulo an integer k gives an eventually periodic sequence. For example, the sequence taken modulo 10 is [1, 1, 2, 7, 2, 1, 2, 7, 2, 1, 2, 7, 2, ...], with an apparent period [1, 2, 7, 2] beginning at a(1), of length 4. Cf. A000670. - _Peter Bala_, Apr 12 2023 a(n) ~ (sqrt(2)/2)*n!/log(1+sqrt(2))^(n+1). - Conjectured by _Simon Plouffe_, Feb 17 2007.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000685 | OEIS-A006165

a(1) = a(2) = 1; thereafter a(2n+1) = a(n+1) + a(n), a(2n) = 2a(n).

Conjectures: for n >= 2 there holds (vii) for k >= 1, the sequence {n - a^(k)(n) : n >= 1} has first differences either 0 or 1. We conjecture that the repeated values of the sequence are of the form (2^k - 1)*2^m. The number of repeated values appears to always be 2, 3, 5, 9, 17, 35, ..., independent of k, conjecturally A000051. Two examples are given below.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000686 | OEIS-A006172

a(n) = 1 + F(2*n+1) + (-1)^n*L(n).

The g.f. -(-2 -3*x + 19*x^2 - 17*x^3 + 4*x^4)/((x-1)*(x^2 - x - 1)*(x^2 - 3*x + 1)) conjectured by _Simon Plouffe_ in his 1992 dissertation is probably wrong. [The correct g.f. is (4 - 10*x - x^2 + 9*x^3 - 3*x^4)/((x - 1)*(x^2 - x - 1)*(x^2 - 3*x + 1)). From _Bruno Berselli_, Mar 09 2015] (Comment VII) We can now comment on the g.f., P(x), given by _Simon Plouffe_ and mentioned at the beginning of this Comment section. Expanding P(x) as a series, we obtain 2 + 9x + 10x^2 + 42x^3 + ..., the g.f. of {H_k}_{k>=1}. In fact, it is easily checked that P(x) = 1/x (H(x)-4). This settles the Plouffe conjecture as follows: (i) H(x) is the rational function g.f. of {H_k}_{k>=0}, (ii) P(x) is the rational function g.f. of {H_k}_{k>=1}, and (iii) C(x)+L(-x)+B2(x) is the partial fraction g.f. of {H_k}_{k>=0}. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000687 | OEIS-A006186

Number of pair-coverings with largest block size 4.

The g.f. (1 + 5*z + 6*z^2 + z^3)/((z-1)*(z^2-z-1)) conjectured by _Simon Plouffe_ in his 1992 dissertation is incorrect.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000688 | OEIS-A006218

a(n) = Sum_{k=1..n} floor(n/k); also Sum_{k=1..n} d(k), where d = number of divisors (A000005); also number of solutions to x*y = z with 1 <= x,y,z <= n.

It appears that the partial sums give A078567. - _N. J. A. Sloane_, Nov 24 2008 [This conjecture is true (see Fried link). - _Sela Fried_, Oct 31 2025]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000689 | OEIS-A006231

a(n) = Sum_{k=2..n} n(n-1)...(n-k+1)/k.

Let P_n be the set of probability distributions over orderings of n objects that can be obtained by drawing n real numbers from independent probability distributions and sorting. Then a(n) is conjectured to be the dimension of P_n, as a semi-algebraic subset of R^(n!). - _Jamie Tucker-Foltz_, Jul 29 2024 Conjecture: a(n) + (-n-2)*a(n-1) + (3*n-2)*a(n-2) + 3*(-n+2)*a(n-3) + (n-3)*a(n-4) = 0. - _R. J. Mathar_, Aug 06 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000690 | OEIS-A006234

a(n) = n*3^(n-4).

Number of spanning trees in the book graph of order n-2, i.e., S_{n-2} X P_2 (S_k = the star graph on k nodes) (conjectured). This conjecture is true - see Doslic (2013). - _N. J. A. Sloane_, Dec 28 2013 Conjecture: a(n+2) is the total number of parts used in the compositions of n if the parts can be runs of any length from 1 to n, and contain any integers from 1 to n. (The number of such compositions is given by A000244(n-1).) - _Gregory L. Simay_, May 27 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000691 | OEIS-A006255

R. L. Graham's sequence: a(n) = smallest m for which there is a sequence n = b_1 < b_2 < ... < b_t = m such that b_1*b_2*...*b_t is a perfect square.

Conjecture: Let b(n) = A006530(A007913(n)). If b(n)^2 >= 2 * n then a(n) = n + b(n) except for n = 2, 3, 10, and 171. [Corrected by Jaideep Padhi_, Jun 15 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000692 | OEIS-A006308

Coefficients of period polynomials.

Conjecture: a(n) = A008837(n) = p*(p-1)/2 = Sum_{k=0..p-1} mod(k^3, p) where p = prime(n). - _Michael Somos_, Feb 17 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000693 | OEIS-A006325

4-dimensional analog of centered polygonal numbers: a(n) = n*(n-1)*(n^2-n+1)/6.

The conjecture by David J. Seal is true and easily proved using MacMahon's Omega operators via the "Omega" package for Mathematica authored by Axel Riese (obtaining (up to an offset) the generating function listed in the formula section below). See the second Mathematica program in which the edges of LOOP X C_4 are indexed as in the example below. The Omega package can be downloaded from the link provided in the article by G. E. Andrews et al. - _L. Edson Jeffery_, Oct 15 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | Albert Rich (Albert_Rich(AT)msn.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000694 | OEIS-A006340

An "eta-sequence": [(n+1)*tau + 1/2] - [n*tau + 1/2], tau = (1 + sqrt(5))/2.

Conjecture: A006340 = continued fraction expansion of (2.729967741... = sup{f(n,1)}), where f(1,x) = x + 1 and thereafter f(n,x) = x + 1 if n is in the lower Wythoff sequence (A000201), else f(n,x) = 1/x. The first 12 values of f(n,1) are given in Example at A245216. - _Clark Kimberling_, Jul 14 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | D. R. Hofstadter, Jul 15 1977 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000695 | OEIS-A006343

Arkons: number of elementary maps with n-1 nodes.

Conjecture: D-finite with recurrence: -(n+3)*(n-1)*a(n) +(11*n^2-2*n-45)*a(n-1) -(37*n+29)*(n-3)*a(n-2) +(29*n^2-125*n+78)*a(n-3) +(61*n-106)*(n-3)*a(n-4) -155*(n-3)*(n-4)*a(n-5)=0. - _R. J. Mathar_, Feb 20 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000696 | OEIS-A006351

Number of series-parallel networks with n labeled edges. Also called yoke-chains by Cayley and MacMahon.

Conjecture: a(n) = A379459(n-2,0) = A379460(n-1,0) for n > 1 with a(1) = 1. - _Mikhail Kurkov_, Jan 16 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000697 | OEIS-A006356

a(n) = 2*a(n-1) + a(n-2) - a(n-3) for n >= 3, starting with a(0) = 1, a(1) = 3, and a(2) = 6.

For a k-glass sequence, say a(n,k), a(n,k) is always asymptotic to z(k)*w(k)^n where w(k) = (1/2)/cos(k*Pi/(2*k+1)) and it is conjectured that z(k) is the root 1 < x < 2 of a polynomial of degree Phi(2k+1)/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000698 | OEIS-A006364

Numbers k with an even number of 1's in binary, ignoring last bit.

Conjecture: a(n) = 2*n + (-1)^(n-A000120(n-1)) - (3+(-1)^n)/2. - _Velin Yanev_, Dec 21 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000699 | OEIS-A006368

The "amusical permutation" of the nonnegative numbers: a(2n) = 3n, a(4n+1) = 3n+1, a(4n-1) = 3n-1.

There is a famous open question concerning the closed trajectories under this map - see A217218, A028393, A028394, and Conway (2013).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _J. H. Conway_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000700 | OEIS-A006415

Number of nonseparable toroidal tree-rooted maps with n + 3 edges and n + 1 vertices.

Conjecture: a(n) = 4 * binomial(n + 4, n) + 84 * binomial(n + 4, n - 1) + 456 * binomial(n + 4, n - 2) + 996 * binomial(n + 4, n - 3) + 950 * binomial(n + 4, n - 4) + 330 * binomial(n + 4, n - 5). - _Sean A. Irvine_, Apr 03 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000701 | OEIS-A006428

Number of loopless tree-rooted planar maps with 3 vertices and n faces and no isthmuses.

Conjecture (for n > 1): a(n) = n*(n+1)*(n^3+6*n^2+11*n-42) / 24. - _Sean A. Irvine_, Apr 10 2017 The above conjectures are true. - _Andrew Howroyd_, Apr 03 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000702 | OEIS-A006451

Numbers k such that k*(k+1)/2 + 1 is a square.

G.f.: x*(-2-3*x+2*x^2+x^3)/(x-1)/(x^2+2*x-1)/(x^2-2*x-1). Conjectured (correctly) by _Simon Plouffe_ in his 1992 dissertation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Jeffrey Shallit_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000703 | OEIS-A006468

Number of rooted planar maps with 4 faces and n vertices and no isthmuses.

G.f.: -x*(x^3-4*x^2+2*x+5)/(x-1)^7, equivalent to a(n) = n*(n+1) *(n+2) *(2*n^3 +33*n^2 +142*n +123) /360, conjectured in _Simon Plouffe_'s Master's thesis, 1992. The above conjecture is true. - _Andrew Howroyd_, Apr 02 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000704 | OEIS-A006472

a(n) = n!*(n-1)!/2^(n-1).

Conjecture: For n > 1, n divides 2*a(n-1) + 4 if and only if n is prime. - _Werner Schulte_, Oct 04 2020 For a proof of the above conjecture see Himane. The list of primes p such that p^2 divides (2*a(p-1) + 4) (analog of A007540 - Wilson primes) begins [239, 24049, ...]. - _Peter Bala_, Nov 06 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000705 | OEIS-A006479

From variance of Fibonacci search.

(Conjecture equivalent to Plouffe g.f.): a(n) = -3 - 2*A001629(n+2) - 3*A001629(n+1) + 2*A001628(n-1) + A020701(n+1). - _R. J. Mathar_, Dec 06 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000706 | OEIS-A006501

Expansion of (1+x^2) / ((1-x)^2 * (1-x^3)^2).

Conjecture: a(n) = A144677(n) + A144677(n-2). - _R. J. Mathar_, Mar 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000707 | OEIS-A006502

Related to Fibonacci numbers.

Conjecture: g.f. (1-7*x+18*x^2-10*x^3+5*x^4-15*x^5-39*x^6-20*x^7-3*x^8) / ((x^2+3*x-1) *(x^2+2*x-1)^2 *(x^2+x-1)^3). - _R. J. Mathar_, Jun 18 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000708 | OEIS-A006531

Number of semiorders on n labeled elements.

Conjecture: for fixed k = 1,2,..., the sequence a(n) (mod k) is eventually periodic with the exact period dividing phi(k), where phi(k) is Euler's totient function A000010. For example, modulo 10 the sequence becomes (1, 1, 3, 9, 3, 1, 3, 9, 3, ...) with an apparent period 1, 3, 9, 3 of length 4 = phi(10) beginning at a(1). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000709 | OEIS-A006562

Balanced primes (of order one): primes which are the average of the previous prime and the following prime.

Conjecture: Limit_{n->oo} n*(log(a(n)))^2 / a(n) = 1/2. - _Alain Rocchelli_, Mar 21 2024 Conjecture: The asymptotic limit of the average of a(n+1)-a(n) is equivalent to 2*(log(a(n)))^2. Otherwise formulated: 2 * Sum_{n=1..N} (log(a(n)))^2 ~ a(N). - _Alain Rocchelli_, Mar 23 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000710 | OEIS-A006579

a(n) = Sum_{k=1..n-1} gcd(n,k).

Conjecture: a(n) != -1 (mod n) for a composite n. - _Thomas Ordowski_, Jun 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marc LeBrun_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000711 | OEIS-A006605

Number of modes of connections of 2n points.

Conjecture: 3*n*(3*n+2)*(3*n+1)*a(n) + (-275*n^3 + 475*n^2 - 328*n + 68)*a(n-1) + 2*(2*n-3)*(139*n^2 - 507*n + 398)*a(n-2) + 180*(2*n-5)*(n-2)*(2*n-3)*a(n-3) = 0. - _R. J. Mathar_, May 30 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000712 | OEIS-A006689

Number of deterministic, completely-defined, initially-connected finite automata with 2 inputs and n unlabeled states.

Conjecture: a(n) = n * P(n+1, n+1) where P(n, k) = P(n-1, k) + (n-1) * P(n, k-1) for 1 < k < n with P(n, 1) = 1, P(n, n) = (n-1) * P(n, n-1). - _Mikhail Kurkov_, Jun 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000713 | OEIS-A006694

Number of cyclotomic cosets of 2 mod 2n+1.

Conjecture: a((3^n-1)/2) = n. - _Vladimir Shevelev_, May 26 2008 [This is correct. 2*((3^n-1)/2) + 1 = 3^n and the polynomial (x^(3^n) - 1) / (x - 1) factors over GF(2) into Product_{k=0..n-1} x^(2*3^k) + x^(3^k) + 1. - _Joerg Arndt_, Apr 01 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 25 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000714 | OEIS-A006697

Number of subwords of length n in infinite word generated by a -> aab, b -> b.

G.f.: 1 + 1/(1-x) + 1/(1-x)^2 * [1/(1-x) - sum(k>=1, x^(2^k+k-1))] (conjectured). - _Ralf Stephan_, Mar 05 2004 Conjectures: partial sums of A103354, also equal to A094913(n) + 1. - _Vladeta Jovovic_, Sep 19 2005 a(n) = Sum_{k=0..n} min(2^k,n-k+1) = 2^(m+1)-1 + (n-m)*(n-m+1)/2, with m = [n+1-LambertW(2^(n+1)*log(2))/log(2)] = integer part of the solution to 2^m = n+1-m. (conjectured). - _M. F. Hasler_, Dec 14 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Jeffrey Shallit_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000715 | OEIS-A006702

Solution to a Pellian equation: least x such that x^2 - n*y^2 = +- 1.

Apparently the generating function of the sequence of the denominators of continued fraction convergents to sqrt(n) is always rational and of form p(x)/[1 - C*x^m + (-1)^m * x^(2m)], or equivalently, the denominators satisfy the linear recurrence b(n+2m) = C*b(n+m) - (-1)^m * b(n). If so, then it seems that a(n) is half the value of C for each nonsquare n, or 1. See A003285 for the conjecture regarding m. The same conjectures apply to the sequences of the numerators of continued fraction convergents to sqrt(n). - _Ralf Stephan_, Dec 12 2013 The conjecture is true, cf. link. - _Jan Ritsema van Eck_, Mar 08 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000716 | OEIS-A006716

Squares with digits 1, 4, 9.

This is probably a finite sequence, but that is only a conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, revised Jul 10 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000717 | OEIS-A006794

Primorial -1 primes: primes p such that -1 + product of primes up to p is prime.

Conjecture: if p# - 1 is a prime number, then the previous prime is greater than p# - exp(1)*p. - _Arkadiusz Wesolowski_, Jun 19 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000718 | OEIS-A006848

Number of extreme points of the set of n X n symmetric doubly-substochastic matrices.

The lgdegf is (2+x+x^4+x^5-2*x^3-2*x^2)/(x-1)^2/(x+1)^2, conjectured in Simon Plouffe's Master's Thesis, Uqam 1992. Lgdegf is the logarithmic derivative of f(x), the g.f. is exponential.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000719 | OEIS-A006863

Denominator of B_{2n}/(-4n), where B_m are the Bernoulli numbers.

Michael Lugo (see link) conjectures, and Peter McNamara proves, that a(n) = gcd_{ primes p > 2n+1 } (p^(2n) - 1). - _Tanya Khovanova_, Feb 21 2009 [edited by _Charles R Greathouse IV_, Dec 03 2014]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Jeffrey Shallit_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000720 | OEIS-A006883

Long period primes: the decimal expansion of 1/p has period p-1.

Emil Artin conjectured that the proportion of primes that belong to this sequence can be expressed as:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Munafo_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000721 | OEIS-A006946

Independence number of de Bruijn graph of order n on two symbols.

Proposition 4.3 (b) in Lichiardopol's paper (see links) can be formulated as a(n) <= 2^(n-1) - A000031(n)/2 + 1 for odd n. For even n, Proposition 5.4 says that a(n) <= (a(n+1) + 1)/2 <= 2^(n-1) - A000031(n+1)/4 + 1. For n<=13, equality holds in both cases, and I conjecture that it holds for all n. If this is true, the sequence would continue a(14)=7645, a(15)=15289, a(16)=30841, a(17)=61681, ... - _Pontus von Brömssen_, Feb 29 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Herb Taylor | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000722 | OEIS-A006950

G.f.: Product_{k>=1} (1 + x^(2*k - 1)) / (1 - x^(2*k)).

It appears that this sequence is related to the generalized hexagonal numbers (A000217) in the same way as the partition numbers A000041 are related to the generalized pentagonal numbers A001318. (See the table in comments section of A195825.) Conjecture: this is 1 together with the row sums of triangle A195836, also column 1 of A195836, also column 2 of the square array A195825. - _Omar E. Pol_, Oct 09 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Warren D. Smith_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000723 | OEIS-A006960

Reverse and Add! sequence starting with 196.

196 is conjectured to be the smallest initial term which does not lead to a palindrome. John Walker, Tim Irvin and others have extended the trajectory of 196 to millions of digits without finding a palindrome.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000724 | OEIS-A006972

Lucas-Carmichael numbers: squarefree composite numbers k such that p | k => p+1 | k+1.

Conjecture: if k = p*q*r, p = a*d - 1, q = b*d - 1, r = c*d - 1 are distinct odd primes, with d = gcd(p + 1, q + 1, r + 1) and a*b*c*d divides k + 1, then k is a Lucas-Carmichael number. - _Davide Rotondo_, Dec 23 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Pinch_ and _Jeffrey Shallit_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000725 | OEIS-A007016

Number of permutations of length n with 1 fixed and 1 reflected point.

Conjecture D-finite with recurrence
(365968635435167109808*n^2
-5566069866485493251505*n
+20525522573033552369132)*a(n)
+(-12153690443264305423 11*n^2
+19103429957352794982854*n
-73690801030090785944295)*a(n-1)
+(-365968635435167109808*n^4
+66639757727909945809 29*n^3
-35836353442786038818589*n^2
+34878550744402035813586*n
+12404354247282100776320 4)*a(n-2)
+(48343177345609632269 5*n^4
-10754417727097457203127*n^3
+85154149458907095778621*n^2
-27768396799472258420606 7*n
+28625487034283575775185 2)*a(n-3)
+2*(-39324190911348388473 8*n^4
+91423349518392650433 83*n^3
-78427160779754271402777*n^2
+30928396816086256758 0813*n
-46505742234427714197792 3)*a(n-4)
+2*(-74504454750258020991 9*n^4
+21471238686323774026196*n^3
-22206783254369019378925 5*n^2
+94469895493204983008423 2*n
-13727325318596191197939 78)*a(n-5)
+4*(365968635435167109808*n^4
-52273745047286429166 27*n^3
+19793104565012302929789*n^2
+39183481600793992708 2*n
-57365695502678698166 146)*a(n-6)
+4*(-48343177345609632269 5*n^4
+75922143123143953797 33*n^3
-45284933032689911393913*n^2
+11753588508890910344916 5*n
-84799883220517633629252)*a(n-7)
+8*(n-7)*(39324190911348388473 8*n^3
-47894006779126255363 35*n^2
+17834478528905815208536*n
-23668675533486426523455)*a(n-8)
+8*(n-7)*(n-8)*(74504454750258020991 9*n^2
-60869159628160735051 21*n
+12854159797389104313178)*a(n-9)=0. - _R. J. Mathar_, Feb 27 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000726 | OEIS-A007030

Non-Hamiltonian simplicial polyhedra with n nodes.

a(18) = 1879665 was conjectured by Dillencourt and verified by direct computation by _Sean A. Irvine_, Sep 26 2017.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000727 | OEIS-A007051

a(n) = (3^n + 1)/2.

The (corrected) conjecture regarding the sequence s(n) is true (see Fried link). - _Sela Fried_, Oct 26 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Mallows_, _N. J. A. Sloane_, _Simon Plouffe_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000728 | OEIS-A007059

Number of balanced ordered trees with n nodes.

Conjecture: for n > 0, a(n+1) is the number of "numbral" divisors of (4^n-1)/3 = A002450(n) (see A048888 for the definition of numbral arithmetic). This has been verified computationally up to n=15. - _John W. Layman_, Dec 18 2001 [This conjecture follows immediately from Proposition 2.3 of Frosini and Rinaldi. - _N. J. A. Sloane_, Apr 29 2011]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Mira Bernstein_, R. Kemp | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000729 | OEIS-A007061

The Ehrenfeucht-Mycielski sequence (1,2-version): a maximally unpredictable sequence.

Sutner continues: All words of length k seem to appear in the first 2^{k+2} bits. This is true for the first billion bits of the sequence, but no proof is known. The main open problem is whether the limiting density of 0's is 1/2. It seems to require a large amount of effort just to show that it is bounded away from 0, never mind some of the more exotic properties of the sequence (see the Sutner reference).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Mira Bernstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000730 | OEIS-A007063

Main diagonal of Kimberling's expulsion array (A035486).

O for "outer", i.e., farthest from expelled. For example, the array A035486 (and diagonal A007063) are coded as RILI. For the eight codes see Example and Mathematica. It is conjectured that six of the eight diagonal sequences are permutations of the positive integers. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Mira Bernstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000731 | OEIS-A007086

Next term is uniquely the sum of 3 earlier terms.

G.f.: (22*x^18 -21*x^17 +x^16 -2*x^13 -7*x^12 -15*x^9 +2*x^8 +2*x^7 -2*x^5 -2*x^4 -x^3 -x^2 -x) / (-x^4+x^3+x-1). Conjectured and verified for n<=1100 - _Alois P. Heinz_, Jan 04 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000732 | OEIS-A007091

Numbers in base 5.

Sigrist's conjecture is true. It is essentially Shepherd's statement. See Fried link. - _Sela Fried_, Nov 03 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000733 | OEIS-A007097

Primeth recurrence: a(n+1) = a(n)-th prime.

Conjecture: log(a(1))*log(a(2))*...*log(a(n)) ~ a(n). - _Thomas Ordowski_, Mar 26 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000734 | OEIS-A007126

Number of connected rooted strength 1 Eulerian graphs with n nodes.

It is not difficult to prove that a(n) = A000088(n-1) - Sum_{k=1..n-1} a(k)*A002854(n-k), n>1, with a(1) = 1, which is equivalent to the conjecture that the Euler transform of A158007(n) gives A007126(n+1) (see A158007).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000735 | OEIS-A007223

Number of distinct perforation patterns for deriving (v,b) = (n+2,n) punctured convolutional codes from (2,1).

Conjecture: Expansion of [A(x)^2 + A(x^2)]/2, where A(x) = A000108(x) - 1. - _R. J. Mathar_, Jul 19 2016 It follows from my comments in A275206 that _R. J. Mathar_'s conjecture is correct and that

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000736 | OEIS-A007228

a(n) = 3*binomial(4*n,n)/(n+1).

Conjecture: for n >= 1, a(n) is odd iff n = 4*A263133(k) + 3 for some k. - _Peter Bala_, Mar 13 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000737 | OEIS-A007283

a(n) = 3*2^n.

a(n) is the upper bound for the "X-ray number" of any convex body in E^(n + 2), conjectured by Bezdek and Zamfirescu, and proved in the plane E^2 (see the paper by Bezdek and Zamfirescu). - _L. Edson Jeffery_, Jan 11 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000738 | OEIS-A007293

Dimension of space of weight systems of chord diagrams.

Broadhurst gives a conjectured g.f.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000739 | OEIS-A007300

a(1)=2, a(2)=5; for n >= 3, a(n) is smallest number which is uniquely of the form a(j) + a(k) with 1 <= j < k < n.

Comments from _Joshua Zucker_, May 24 2006: "Concerning the conjecture about periodicity mod 126. Out of the first 300 terms, only the 2 and 12 are even. But if you neglect those first 6 terms, mod 2 they're all odd, mod 9 it goes: 0 4 6 1 7 4 6 8 7 2 4 6 8 5 0 8 1 2 5 7 0 2 4 6 1 5 0 2 8 1 5 7 which appears to repeat indefinitely and mod 7 it goes: 0 2 6 1 3 0 2 6 5 4 6 1 2 6 1 3 5 4 1 2 4 0 5 0 2 4 6 1 5 2 6 1 which also appears to repeat indefinitely. "The Mathworld link cites a proof that sequences of this type (2,n) have only two even terms and another proof that sequences with only finitely many even terms must eventually have periodic first differences. So I think the period 32 difference of 126 conjecture may be proved in those references."

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Mira Bernstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000740 | OEIS-A007306

Denominators of Farey tree fractions (i.e., the Stern-Brocot subtree in the range [0,1]).

Conjecture: a(n) = a(n-1) + b(n-1) - 2*(a(n-1) mod b(n-1)) for n > 1 with a(0) = a(1) = 1 where b(n) = a(n) - b(n-1) for n > 1 with b(1) = 1. - _Mikhail Kurkov_, Mar 13 2022 The conjecture is true (see Fried link). - _Sela Fried_, Nov 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000741 | OEIS-A007310

Numbers congruent to 1 or 5 mod 6.

Numbers n such that (Sum_{k = 1..n} k^14) mod n = 0. (Conjectured) - _Gary Detlefs_, Dec 27 2011 The above conjecture is true. Apply Ireland and Rosen, Proposition 15.2.2. with m = 14 to obtain the congruence 6*(Sum_{k = 1..n} k^14)/n = 7 (mod n), true for all n >= 1. Suppose n is coprime to 6, then 6 is a unit in Z/nZ, and it follows from the congruence that (Sum_{k = 1..n} k^14)/n is an integer. On the other hand, if either 2 divides n or 3 divides n then the congruence shows that (Sum_{k = 1..n} k^14)/n cannot be integral. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | C. Christofferson (Magpie56(AT)aol.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000742 | OEIS-A007323

Number of numerical semigroups of genus n; conjecturally also the number of power sum bases for symmetric functions in n variables.

Number of numerical semigroups of genus n; conjecturally also the number of power sum bases for symmetric functions in n variables. The theorem alluded to in the email by Zagier is due to Kakeya, not Kakutani (see references.) The theorem states that if a sequence of n positive integers k1, k2,..., kn forms the complement of a numerical semigroup, then the power sums p_k1, p_k2,..., p_kn forms a

basis for the rational function field of symmetric functions in n variables. Kakeya conjectures that every power sum basis of the symmetric functions has this property, but this is still an open problem. Thanks to user Gjergji Zaimi on Math Overflow for the references. - _Trevor Hyde_, Oct 18 2018 Conjectures: A) a(n) >= a(n-1)+a(n-2); B) a(n)/(a(n-1)+a(n-2)) approaches 1; C) a(n)/a(n-1) approaches the golden ratio; D) a(n) >= a(n-1). Conjectures A, B, C, D were presented by M. Bras-Amorós in the seminar Algebraic Geometry, Coding and Computing, in Segovia, Spain, in 2007, and at IMNS 2018 in Granada, Spain, in 2008. Conjectures A, B, C were then published in the Semigroup Forum, 76 (2008), 379-384. Conjectures B and C are proved in the paper by Zhai, 2011. - _Maria Bras-Amorós_, Oct 24 2007, corrected Aug 31 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | Don Zagier (don.zagier(AT)mpim-bonn.mpg.de), Apr 11 1994 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000743 | OEIS-A007335

MU-numbers: next term is uniquely the product of 2 earlier terms.

Conjecture: Sum_{n>=1} 1/a(n) = 181/144. - _Amiram Eldar_, Jul 31 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000744 | OEIS-A007356

Apocalyptic powers: 2^a(n) contains 666.

Conjecture: for n > 26069, a(n) = n + 3715. [_Charles R Greathouse IV_, Sep 28 2011]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000745 | OEIS-A007366

Numbers k such that phi(x) = k has exactly 2 solutions.

The smaller of the solutions to phi(x) = a(n) is given by A271983(n). It is conjectured that the larger solution is 2*A271983(n); or equivalently, all terms in A271983 are odd. - _Jianing Song_, Nov 08 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Mira Bernstein_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000746 | OEIS-A007374

Smallest k such that phi(x) = k has exactly n solutions, n>=2.

The Carmichael Totient Conjecture is that there is no k such that phi(x) = k has a unique solution x. So a(1) does not exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Mira Bernstein_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000747 | OEIS-A007377

Numbers k such that the decimal expansion of 2^k contains no 0.

It is an open problem of long standing to show that 86 is the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000748 | OEIS-A007384

Number of strict 3rd-order maximal independent sets in path graph.

Conjecture: a(n)= 3*a(n-2) -3*a(n-4) +a(n-5) +a(n-6) -2*a(n-7) +a(n-9) with g.f. -x^5/((x^5+x^2-1)*(x-1)^2*(1+x)^2). [From _R. J. Mathar_, Oct 30 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Mira Bernstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000749 | OEIS-A007390

Number of strict (-1)st-order maximal independent sets in cycle graph.

Conjectures from _Colin Barker_, Jun 14 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Mira Bernstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000750 | OEIS-A007392

Number of strict 3rd-order maximal independent sets in cycle graph.

Conjecture: a(n) = 3*a(n-2) - 3*a(n-4) + a(n-5) + a(n-6) - 2*a(n-7) + a(n-9) with g.f. x^10*(-5+3*x^2)/((x^5+x^2-1)*(x-1)^2*(1+x)^2). - _R. J. Mathar_, Oct 30 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Mira Bernstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000751 | OEIS-A007405

Dowling numbers: e.g.f.: exp(x + (exp(b*x) - 1)/b) with b=2.

Conjecture: Let M_n be an n X n matrix whose elements are m_ij = 1 for i < j - 1, m_ij = -1 for i = j - 1, and m_ij = binomial(n - j,i - j) otherwise. Then a(n - 1) = Det(M_n). - _Benedict W. J. Irwin_, Apr 19 2017; corrected by _Sela Fried_, Dec 30 2025 Irwin's conjecture is true (see Fried link). - _Sela Fried_, Dec 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000752 | OEIS-A007413

A squarefree (or Thue-Morse) ternary sequence: closed under 1->123, 2->13, 3->2. Start with 1.

Conjecture. The sequence is formed by the numbers of 1's between every pair of consecutive 2's in A076826. - _Vladimir Shevelev_, May 31 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000753 | OEIS-A007427

Moebius transform applied twice to sequence 1,0,0,0,....

Dirichlet g.f. of |a(n)|: zeta(s)^2/zeta(2s)^2 (conjectured). - _Ralf Stephan_, Jul 05 2013. The conjecture is correct because 1+Sum_{e>=1} binomial(2,e)/p^(e*s) = (p^s+1)^2/p^2s, whose product over p is zeta(s)^2/zeta(2s)^2. - _Michael Shamos_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000754 | OEIS-A007434

Jordan function J_2(n) (a generalization of phi(n)).

a(n) = A001065(n) - A134675(n). - Conjectured by _John Mason_ and proved by _Max Alekseyev_, Jan 07 2015 a(n) = lim_{k->oo} (n^(2*(k + 1)))/A001157(n^k), conjectured by _Velin Yanev_, Dec 04 2024 [See my link for proof. - _Roman Le Lan_, Sep 12 2025]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000755 | OEIS-A007473

Dimension of space of Vassiliev knot invariants of order n.

Broadhurst gives a conjectured explicit g.f. (different from A014595).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000756 | OEIS-A007478

Dimension of primitive Vassiliev knot invariants of order n.

Broadhurst gives a conjectured g.f.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000757 | OEIS-A007484

a(n) = 3*a(n-1) + 2*a(n-2), with a(0)=2, a(1)=7.

8*a(n) = A007482(n+2) + A007483(n+1) (conjectured, see A104934 for related formula). - _Creighton Dement_, Apr 15 2005 Conjecture verified using generating functions. - _Robert Israel_, Jul 12 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000758 | OEIS-A007489

a(n) = Sum_{k=1..n} k!.

Conjectured to be the length of the shortest word over {1,...,n} that contains each of the n! permutations as a factor (cf. A180632) [see Johnston]. - _N. J. A. Sloane_, May 25 2013 The above conjecture has been disproven for n>=6. See A180632 and the Houston 2014 reference. - _Dmitry Kamenetsky_, Mar 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000759 | OEIS-A007504

Sum of the first n primes.

Conjecture: Each integer n >= 5340 can be written as a(i) + a(j) + 2*a(k) for some i,j,k > 0. (This has been verified for n <= 10^5.) - _Zhi-Wei Sun_, Oct 09 2025 Conjecture: Any integer n > 16866 can be written as p + a(k) with p prime and k positive. Also, each integer n > 13392 can be written as q + a(k) with q practical and k positive. (This has been verified for n <= 2*10^7.) - _Zhi-Wei Sun_, Oct 15 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000760 | OEIS-A007526

a(n) = n*(a(n-1) + 1), a(0) = 0.

Conjecture : a(n) = (n+2)*a(n-1) - (2*n-1)*a(n-2) + (n-2)*a(n-3). - _R. J. Mathar_, Dec 04 2012 [Conjecture verified by _Robert FERREOL_, Aug 04 2018]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000761 | OEIS-A007531

a(n) = n*(n-1)*(n-2) (or n!/(n-3)!).

Ed Pegg Jr conjectures that n^3 - n = k! has a solution if and only if n is 2, 3, 5 or 9 (when k is 3, 4, 5 and 6).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000762 | OEIS-A007534

Positive even numbers that cannot be written as p + q, where p and q are primes each belonging to a twin-prime pair (p = q is allowed).

Conjectured to be complete (although if this were proved it would prove the "twin primes conjecture"!).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000763 | OEIS-A007540

Wilson primes: primes p such that (p-1)! == -1 (mod p^2).

Conjecture: Odd primes p such that 1^(p-1) + 2^(p-1) + ... + (p-1)^(p-1) == p-1 (mod p^2). - _Thomas Ordowski_ and _Giovanni Resta_, Jul 25 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000764 | OEIS-A007549

Number of increasing rooted connected graphs where every block is a complete graph.

Conjecture: a(n) = Sum_{i=0..2^(n-2) - 1} b(i) for n > 1 with a(1) = 1 where b(n) = (L(n) + 2)*b(f(n)) + Sum_{k=0..L(n) - 1} (1 - R(n,k))*b(f(n) + 2^k*(1 - R(n,k))) for n > 0 with b(0) = 1, L(n) = A000523(n), f(n) = A053645(n) and where R(n,k) = floor(n/2^k) mod 2. Here R(n,k) is the (k+1)-th bit from the right side in the binary expansion of n. - _Mikhail Kurkov_, Jul 21 2024 Conjecture: a(n) = D^(n-1)(exp(x)) evaluated at x = 0, where D denotes the operator exp(x)*(1 + x)*d/dx. - _Peter Bala_, Feb 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000765 | OEIS-A007581

a(n) = (2^n+1)*(2^n+2)/6.

Dimension of the universal embedding of the symplectic dual polar space DSp(2n,2) (conjectured by A. Brouwer, proved by P. Li). - J. Taylor (jt_cpp(AT)yahoo.com), Apr 02 2004.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ and _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000766 | OEIS-A007598

Squared Fibonacci numbers: a(n) = F(n)^2 where F = A000045.

Conjecture: Sum_{n >= 1, n != 2*k+1} 1/(a(n) + (-1)^n*a(2*k+1)) = 1/a(4*k+2) for k = 0,1,2,.... (End) The conjecture is true (see Fried link). - _Sela Fried_, Nov 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000767 | OEIS-A007618

a(n) = a(n-1) + sum of digits of a(n-1), a(1) = 5.

a(2) = 10 and a(590) = 10000 are the first two powers of 10 in this sequence; there are no others below a(19017393928) = 1000000000093. Conjecture: the sequence contains infinitely many powers of 10. - _Charles R Greathouse IV_, Mar 29 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Robert G. Wilson v_, _Mira Bernstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000768 | OEIS-A007655

Standard deviation of A007654.

4*a(n+1) + A046184(n) = A055793(n+2) + A098301(n+1) 4*a(n+1) + A098301(n+1) + A055793(n+2) = A046184(n+1) (4*a(n+1))^2 = A098301(2n+1) (conjectures). - _Creighton Dement_, Nov 02 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000769 | OEIS-A007675

Numbers m such that m, m+1 and m+2 are squarefree.

Conjecture: for every prime p, the numbers p# - 1, p#, p# + 1 are squarefree, where primorial p# = product of all primes <= p. - _Thomas Ordowski_, Apr 21 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000770 | OEIS-A007691

Multiply-perfect numbers: n divides sigma(n).

Luca's solution of problem 11090, which proves that for k>1 there are an infinite number of n such that n divides sigma_k(n), does not apply to this sequence. However, it is conjectured that this sequence is also infinite. - _T. D. Noe_, Nov 04 2007 Conjecture: If n is such that 2^n-1 is in A066175 then a(n) is a triangular number. - _Ivan N. Ianakiev_, Aug 26 2013 Conjecture: Every multiply-perfect number is practical (A005153). I've verified this conjecture for the first 5261 terms with abundancy > 2 using Achim Flammenkamp's data. The even perfect numbers are easily shown to be practical, but every practical number > 1 is even, so a weak form says every even multiply-perfect number is practical. - _Jaycob Coleman_, Oct 15 2013 The only terms m > 1 of this sequence that are not in A145551 are m for which sigma(m)/m is not a divisor of m. Conjecture: after 1, A323653 lists all such m (and no other numbers). - _Antti Karttunen_, Mar 19 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000771 | OEIS-A007755

Smallest number m such that the trajectory of m under iteration of Euler's totient function phi(n) [A000010] contains exactly n distinct numbers, including m and the fixed point.

Shapiro shows that the smallest number is greater than 2^(n-1). Catlin shows that if a(n) is odd and composite, then its factors are among the a(k), k < n. For example a(12) = a(5) a(8). There is a conjecture that all terms of this sequence are odd. - _T. D. Noe_, Mar 08 2004 Shapiro mentions on page 30 of his paper the conjecture that a(n) is prime for each n > 1, but a(13) is composite and so the conjecture fails. - _Charles R Greathouse IV_, Oct 28 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | Pepijn van Erp [vanerp(AT)sci.kun.nl] | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000772 | OEIS-A007757

Dwork-Kontsevich sequence evaluated at 2*n.

A formula, conditional on a widely believed conjecture, can be found in the Krattenthaler-Rivoal (2007-2009) paper; see Theorem 4 with k = 1 and the remarks on the top of page 8. Since R. E. Borcherds defined a sequence b(n), but then only entered b(2*n) in the OEIS, the formula has to be taken with n replaced by 2*n. - Christian Krattenthaler (Christian.Krattenthaler(AT)univie.ac.at), Sep 12 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | Richard E. Borcherds (reb(AT)math.berkeley.edu) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000773 | OEIS-A007770

Happy numbers: numbers whose trajectory under iteration of sum of squares of digits map (see A003132) includes 1.

7) It is conjectured that the sequence includes an infinite number of primes (see A035497).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, A.R.McKenzie(AT)bnr.co.uk | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000774 | OEIS-A007780

Losing initial configurations in 2-hole Tchuka Ruma.

Conjectures from _Colin Barker_, Jan 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Darrah Chavey_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000775 | OEIS-A007782

Number of factors in the infinite word formed by the Kolakoski sequence A000002.

a(n) = number of different substrings of length n found in Kolakoski sequence A000002. It is conjectured that a(n) grows like n^(log(3)/log(3/2)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Patricia Lamas (lamas(AT)math.uqam.ca) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000776 | OEIS-A007808

Number of directed column-convex polyominoes of height n: a(k+1)=(k+1)*a(k)+(a(1)+...+a(k)).

Conjectures: E.g.f.: 1 + (-(x^2+1)*exp(-x)+1)*exp(x)/(-1+x)^2; a(n) = round(n!*n*(exp(1)-2)) for n >= 1. - _Simon Plouffe_, Dec 08 2009 [Corrected by _Alper Ferudun_, Oct 19 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Zimmermann_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000777 | OEIS-A007828

Largest t such that a spherical t-design with n points exists in 3 dimensions.

Most of these values are just conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _R. H. Hardin_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000778 | OEIS-A007837

Number of partitions of n-set with distinct block sizes.

Conjecture: the Gauss congruences a(n*p^k) == a(n*p^(k-1)) (mod p^k) hold for all primes p and positive integers n and k. Cf. A185895. - _Peter Bala_, Mar 17 2022 Since A261599(n) = (1/n)*Sum_{d|n} mu(n/d)*a(d) is an integer, n divides Sum_{d|n} mu(n/d)*a(d). This proves the conjecture above. - _Vladeta Jovovic_, Sep 26 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arnold Knopfmacher_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000779 | OEIS-A007840

Number of factorizations of permutations of n letters into ordered cycles.

Conjecture: a(n) = Integral_{x=0..oo} Gamma(x + n)/(exp(x)*Gamma(x)) dx. - _Velin Yanev_, Jan 10 2026 Yanev's conjecture is true (see Fried link). - _Sela Fried_, Feb 14 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arnold Knopfmacher_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000780 | OEIS-A007857

Number of independent sets in rooted plane trees on n nodes.

To prove _R. J. Mathar_'s conjecture, let b(n) = A007226(n-1) = (2*/n)*binomial(3*(n-1), n-1) and c(n) = A000108(n-1) = binomial(2*(n-1), n-1)/n. Since a(n) = b(n) - c(n), it is enough to prove that each of the sequences (b(n): n >= 1) and (c(n): n >= 1) satisfies the same recurrence as (a(n): n >= 1). Conjecture: 2*n*(n - 1)*(2*n - 3)*(44*n - 69)*a(n) + (n - 1)*(176*n^3 - 9591*n^2 + 38703*n - 40640)*a(n-1) + (-17479*n^4 + 218005*n^3 - 959616*n^2 + 1797890*n - 1221920)*a(n-2) + 6*(3*n - 10)*(2*n - 7)*(3*n - 11)*(517*n - 1198)*a(n-3) = 0 for n >= 4. - _R. J. Mathar_, Nov 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Martin Klazar_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000781 | OEIS-A007887

a(n) = Fibonacci(n) mod 9.

Period 24 = A001175(9). Proof: F_{n+24} = F_n + 9*(5152 F_{n+1} + 3184 F_n). - Olivier Wittenberg, following a conjecture by _Ralf Stephan_, Sep 28 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000782 | OEIS-A007901

Number of minimal unavoidable n-celled pebbling configurations.

Conjecture: (n-3)*(n-8)*a(n) +(-11*n^2+127*n-324)*a(n-1) +42*(n^2-12*n+34)*a(n-2) +(-65*n^2+799*n-2400)*a(n-3) +18*(n-6)*(2*n-13)*a(n-4)=0. - _R. J. Mathar_, Aug 14 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | A. M. Odlyzko (odlyzko(AT)dtc.umn.edu) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000783 | OEIS-A007917

Version 1 of the "previous prime" function: largest prime <= n.

For n > 10, these values are close to the most negative eigenvalues of A191898 (conjecture). - _Mats Granvik_, Nov 04 2011 Conjecture: a(n) = floor(log(Sum_{k=2..n} exp(A000010(k)+1))). - _Joseph M. Shunia_, Aug 09 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | R. Muller | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000784 | OEIS-A007921

Numbers that are not the difference of two primes.

Conjecturally, odd numbers k such that k+2 is composite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | R. Muller | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000785 | OEIS-A007933

Some permutation of digits is a prime.

Conjecture: a(n) ~ 3n/2. - _Charles R Greathouse IV_, Sep 19 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | R. Muller | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000786 | OEIS-A007934

Some nontrivial permutation of digits is a prime.

Conjecture: a(n) ~ 3n/2. - _Charles R Greathouse IV_, Sep 19 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | R. Muller | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000787 | OEIS-A007935

Composite numbers such that some permutation of digits is a prime.

Conjecture: a(n) ~ 3n/2. - _Charles R Greathouse IV_, Sep 19 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | R. Muller | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000788 | OEIS-A007949

Greatest k such that 3^k divides n. Or, 3-adic valuation of n.

Lexicographically earliest cubefree sequence, which also (conjecturally) appears in the construction of the lexicographically earliest cubefree {0,1}-sequence

A282317, cf. Example section of A286940. - _M. F. Hasler_, May 21 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | R. Muller | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000789 | OEIS-A007950

Binary sieve: delete every 2nd number, then every 4th, 8th, etc.

Conjectures: there are infinitely many primes that belong to this sequence; also infinitely many composite numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | R. Muller | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000790 | OEIS-A007969

Rectangular numbers.

It seems that D(n) = 4*a(n) gives precisely those even discriminants D from 4*A000037 of indefinite binary quadratic forms that have only improper solutions of the Pell equation x^2 - D*y^2 = +4. Conjecture tested for n = 1..66. Alternatively, the conjecture is that this sequence gives the r values for the Pell equation X^2 + r Y^2 = +1 whenever Y is even. See A261249 and A261250. - _Wolfdieter Lang_, Sep 16 2015 The proof of these two versions of the conjecture is given in the W. Lang link. - _Wolfdieter Lang_, Sep 19 2015 (revised Oct 03 2015)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. H. Conway_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000791 | OEIS-A007970

Rhombic numbers.

This sequence gives the values d of the Pell equation x^2 - d*y^2 = +1 that have positive fundamental solutions (x0, y0) with odd y0. This was first conjectured and is proved provided Conway's theorem in the link is assumed and the proof of the conjecture stated in A007869, given there in a W. Lang link, is used. - _Wolfdieter Lang_, Sep 19 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. H. Conway_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000792 | OEIS-A007972

Number of permutations that are 2 "block reversals" away from 12...n.

a(n) = (n^4+6*n^3+11*n^2-12*n+6)/6 (conjectured). - _Giovanni Resta_, Jun 08 2015 Conjectured g.f.: (-2-5x+3x^2+x^3-x^4)/(-1+x)^5. - _Benedict W. J. Irwin_, Feb 20 2016 a(n) = A228396(n) - A000124(n-1). See C. Homberger links from A228396. This proves the above conjectured formulas up to offset. - _Martin Fuller_, Mar 31 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. H. Conway_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000793 | OEIS-A007980

Expansion of (1+x^2)/((1-x)^2*(1-x^3)).

Conjecture: a(n) is the maximum number of pieces a triangle can be cut into by n cevians. - _Anton Zakharov_, Apr 04 2017 Zakharov's conjecture is true (see Fried link). - _Sela Fried_, Sep 25 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000794 | OEIS-A008287

Triangle of quadrinomial coefficients, row n is the sequence of coefficients of (1 + x + x^2 + x^3)^n.

T(n,k) = Sum_{j=0..k} (-2)^j*binomial(n,j)*binomial(3*n-2*j,k-j) for n >= 0 and 0 <= k <= 3*n (conjectured). - _Werner Schulte_, Sep 09 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000795 | OEIS-A008319

Smallest number that is sum of squares of distinct earlier terms.

Resta's conjecture is correct. Let x = floor(sqrt(n) - 12). For n > 1935, x^2 > n/2. For n > 1853, n - x^2 > 892. So n > 1935 can be decomposed into x^2 plus a number greater than 892. Since the other number is smaller than x^2, any decomposition into squares will use only numbers smaller than x. By induction, all numbers greater than 1935 (and hence greater than 892) are in this sequence. - _Charles R Greathouse IV_, Oct 06 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | R. Muller | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000796 | OEIS-A008347

a(n) = Sum_{i=0..n-1} (-1)^i * prime(n-i).

Conjecture: For any m = 1, 2, 3, ... and r = 0, ..., m - 1, there are infinitely many positive integers n with a(n) == r (mod m). - _Zhi-Wei Sun_, Feb 27 2013 Conjectures: Conjecture: For any n > 9 we have a(n+1) < a(n-1)^(1+2/(n+2)). (This yields an upper bound for prime(n+1) - prime(n) in terms of prime(1), ..., prime(n-1). The conjecture has been verified for n up to 10^8.) - _Zhi-Wei Sun_, Jun 09 2013 Conjecture (iv) above is false since a(1379694977463) = 20922789888000 = 16!. - _Giovanni Resta_, Sep 04 2018 Conjecture: We have {a(m)+a(n): m,n>0} = {2,3,...}. Also, {a(m)-a(n): m,n>0} contains all the integers, and {a(m)/a(n): m,n>0} contains all the positive rational numbers. (I have noted that {a(m)/a(n): m,n = 1..60000} contains {a/b: a,b = 1..1000}.) - _Zhi-Wei Sun_, May 23 2019 Let d(n) = a(n) - a(n-1). Since a(n-1) = prime(n) - a(n), d(n) = 2*a(n) - prime(n). If lim inf a(n)/prime(n) = 1/2 as conjectured by Joseph L. Pe above holds, lim inf d(n)/prime(n) = 2*lim inf a(n)/prime(n) - 1 = 0. Numerical analysis of a(n) for n up to 10^9 shows that abs(d(n))/sqrt(prime(n)) < 15, and thus abs(d(n)) = O(sqrt(prime(n))) is conjectured. - _Ya-Ping Lu_, Aug 31 2020 Meštrovic (2018), following Pillai, conjectures that with a similar conjecture for a(2k+1). - _N. J. A. Sloane_, Dec 21 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _J. H. Conway_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000797 | OEIS-A008364

11-rough numbers: not divisible by 2, 3, 5 or 7.

Conjecture: these are numbers n such that (Sum_{k=1..n} k^4) mod n = 0 and (Sum_{k=1..n} k^6) mod n = 0. - _Gary Detlefs_, Dec 20 2011 The above conjecture is true. Let m be even and let the m-th Bernoulli number be written in reduced form as Bernoulli(m) = N(m)/D(m). Apply Ireland and Rosen, Proposition 15.2.2, to show the congruence

D(m)*(Sum_{k = 1..n} k^m)/n = N(m) (mod n) holds for all n >= 1. It follows easily from this congruence that (Sum_{k = 1..n} k^m)/n is integral iff n is coprime to D(m). Now Bernoulli(4) = -1/(2*3*5) and Bernoulli(6) = 1/(2*3*7) so the numbers n such that both (Sum_{k=1..n} k^4) mod n = 0 and (Sum_{k=1..n} k^6) mod n = 0 are exactly those numbers coprime to the primes 2, 3, 5 and 7, that is, the 11-rough numbers. (End) Conjecture: these are numbers n such that (n^6 mod 210 = 1) or (n^6 mod 210 = 169). - _Gary Detlefs_, Dec 30 2011 The second Detlefs conjecture above is true and extremely easy to verify with some basic properties of congruences: take the terms of this sequence up to 209 and compute their sixth powers modulo 210: there should only be 1's and 169's there. Then take the complement of this sequence up to 210, where you will see no instances of 1 or 169. - _Alonso del Arte_, Jan 12 2014 It is well-known that the product of 7 consecutive integers is divisible by 7!. Conjecture: This sequence is exactly the set of positive values of r such that (Product_{k = 0..6} n + k*r)/7! is an integer for all n. - _Peter Bala_, Nov 14 2015 This conjecture is true. The first part of the proof deals with numbers not in A008364, i.e., numbers which are divisible by p (p either 2, 3, 5, 7). Let r = p*s and n = 1, then (Product_{k = 0..6} n + k*r) is not divisible by p, because none of the factors 1 + k*p*s are divisible by p. Hence dividing the product by 7! does not return an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000798 | OEIS-A008365

13-rough numbers: positive integers that have no prime factors less than 13.

Conjecture: Numbers n such that n^24 is congruent to {1,421,631,841} mod 2310. - _Gary Detlefs_, Dec 30 2011 Gary Detlefs' conjecture of Dec 30 2011 is false: the list of residues is incomplete. The smallest counterexample is n = 17 = a(3). Indeed, 17^24 == 1 (mod 210), while 17 == 6 (mod 11) and 6^4 == 9 (mod 11), so the Chinese remainder theorem gives 17^24 == 1681 (mod 2310). The correct statement is: n is a term of A008365 iff n^24 == 1, 421, 631, 841, or 1681 (mod 2310). Proof: Suppose gcd(n,2310) = 1. Since 1, 2, 4, and 6 all divide 24, Fermat's little theorem gives n^24 == 1 (mod p) for p = 2, 3, 5, and 7, hence n^24 == 1 (mod 210). Modulo 11, Fermat's little theorem gives n^24 == n^4. The map x -> x^4 on (Z/11Z)^* has kernel of size gcd(4,10) = 2, hence image of size 5; this image is precisely the set of nonzero quadratic residues {1,3,4,5,9}. Since 210 == 1 (mod 11), every residue congruent to 1 modulo 210 has the form 1 + 210*j, with 1 + 210*j == 1 + j (mod 11). Thus the five possible values modulo 11 correspond to j = 0, 2, 3, 4, 8, giving exactly the residues 1, 421, 631, 841, and 1681 modulo 2310. Conversely, each of these five residues is coprime to 2310, so if n^24 is congruent to one of them modulo 2310, then gcd(n^24,2310) = 1, and therefore gcd(n,2310) = 1. - _Eiver Rodríguez_, Sep 15 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000799 | OEIS-A008366

Smallest prime factor is >= 17.

Numbers n > 1 such that ((Sum_{k=1..n} k^10) mod n = 0) and ((Sum_{k=1..n} k^12) mod n = 0) (conjecture). - _Gary Detlefs_, Dec 27 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000800 | OEIS-A008368

Number of orbits on points that are at n steps from the origin in the f.c.c. lattice.

G.f. is conjectured to be (1-x^5)/((1-x)*(1-x^2)^2*(1-x^3)) (a conjecture of Simon Plouffe, see A023054).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _J. H. Conway_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000801 | OEIS-A008474

If n = Product (p_j^k_j) then a(n) = Sum (p_j + k_j).

Conjecture: for m > 4, by iterating the map m -> A008474(m) one always reaches 5 [tested up to m = 320000]. - _Ivan N. Ianakiev_, Nov 10 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000802 | OEIS-A008484

Number of partitions of n into parts >= 4.

a(n) is also the number of partitions of n for which three times the number of ones is twice the number of parts (conjectured). - _George Beck_, Aug 19 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | T. Forbes (anthony.d.forbes(AT)googlemail.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000803 | OEIS-A008486

Expansion of (1 + x + x^2)/(1 - x)^2.

Conjecture: This is also the maximum number of edges possible in a planar simple graph with n+2 vertices. - _Dmitry Kamenetsky_, Jun 29 2008 The conjecture is correct. Proof: For n=0 the theorem holds, the maximum planar graph has n+2=2 vertices and 1 edge. Now suppose that we have a connected planar graph with at least 3 vertices. If it contains a face that is not a triangle, we can add an edge that divides this face into two without breaking its planarity. Hence all maximum planar graphs are triangulations. Euler's formula for planar graphs states that in any planar simple graph with V vertices, E edges and F faces we have V+F-E=2. If all faces are triangles, then F=2E/3, which gives us E=3V-6. Hence for n>0 each maximum planar simple graph with n+2 vertices has 3n edges. - _Michal Forisek_, Apr 23 2009 a(n) is conjectured to be the number of polygons added after n iterations of the polygon expansions (type A, B, C, D & E) shown in the Ngaokrajang link. The patterns are supposed to become the planar Archimedean net 3.3.3.3.3.3, 3.6.3.6, 3.12.12, 3.3.3.3.6 and 4.6.12 respectively when n - > infinity. - _Kival Ngaokrajang_, Dec 28 2014 Conjecture: let m = n + 2, p is the polyhedron formed by the convex hull of m points, q is the number of quadrilateral faces of p (see the Wikipedia link below), and f(m) = a(n) - q. Then f(m) would be the solution of the Thompson problem for all m in 3-space. - _Sergey Pavlov_, Feb 03 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000804 | OEIS-A008544

Triple factorial numbers: Product_{k=0..n-1} (3*k+2).

The Mathar conjecture is true. Generally from the factorial form, the last term is the "extra" product beyond the prior term, from k=n-1 and 3k+2 evaluates to 3*(n-1)+2 = 3n-1, yielding a(n) = a(n-1)*(3n-1) (eqn1). Similarly, a(n) = a(n-2)*(3n-1)*(3(n-2)+2) = a(n-2)*(3n-1)*(3n-4) (eqn2) and a(n) = a(n-3)*(3n-1)*(3n-4)*(3*(n-2)+2) = a(n-3)*(3n-1)*(3n-4)*(3n-7) (eqn3). We equate (eqn2) and (eqn3) to get a(n-2)*(3n-1)*(3n-4) = a(n-3)*(3n-1)*(3n-4)*(3n-7) or a(n-2)+(7-3n)*a(n-3) = 0 (eqn4). From (eqn1) we have a(n)+(1-3n)*a(n-1) = 0 (eqn5). Combining (eqn4) and (eqn5) yields a(n)+(1-3n)*a(n-1)+a(n-2)+(7-3n)*a(n-3) = 0. - _Bill McEachen_, Jan 01 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | Joe Keane (jgk(AT)jgk.org) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000805 | OEIS-A008577

Crystal ball sequence for planar net 4.8.8.

The above g.f. and formula were originally stated as conjectures, but I now have a proof. This also justifies the b-file. Details will be added later. - _N. J. A. Sloane_, Dec 29 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000806 | OEIS-A008585

a(n) = 3*n.

Moser conjectured, and Newman proved, that the terms of this sequence are more likely to have an even number of 1s in binary than an odd number. The excess is an undulating multiple of n^(log 3/log 4). See also Coquet, who refines this result. - _Charles R Greathouse IV_, Jul 17 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000807 | OEIS-A008590

Multiples of 8.

Conjecture: let gcd_2(b,c) be the second greatest common divisor and lcd_2(b,c) be the second least common divisor of not coprime integers b and c. Consecutive elements of this sequence (for a(n) > 0) are consecutive integers m for which both Sum_{k=1..m, gcd(k,m)<>1} gcd_2(k,m) and Sum_{k=1..m, gcd(k,m) <>1} lcd_2(k,m) are even numbers. The conjecture was checked for 5*10^4 consecutive integers. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000808 | OEIS-A008611

a(n) = a(n-3) + 1, with a(0)=a(2)=1, a(1)=0.

For the proof of the preceding conjecture see my comments under A008615 and A049310. Chebyshev S(n,x) = i^n*F(n+1,-i*x), with i = sqrt(-1). - _Wolfdieter Lang_, May 06 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 15 1996 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000809 | OEIS-A008642

Quarter-squares repeated.

Conjecture: Consider the number of compositions of n>=4*k+8 into odd parts, where the order of the parts 1,3,..,2k+1 does not count. Then, as k approaches infinity, a(n-4*k-8) is equal to the number of these restricted compositions minus A000009(n), the number of strict partitions of n. - _Gregory L. Simay_, Aug 12 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000810 | OEIS-A008676

Expansion of 1/((1-x^3)*(1-x^5)).

Conjecture: a(n) = Floor(2*(n + 3)/3) - Floor(3*(n + 3)/5). - _John W. Layman_, Sep 23 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000811 | OEIS-A008705

Coefficient of x^n in (Product_{m=1..n}(1-x^m))^n.

Conjectures: the supercongruences a(p) == -1 - p (mod p^2) and a(2*p) == p - 1 (mod p^2) hold for all primes p >= 3. - _Peter Bala_, Apr 18 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | T. Forbes (anthony.d.forbes(AT)googlemail.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000812 | OEIS-A008749

Expansion of (1+x^6)/((1-x)*(1-x^2)*(1-x^3)).

Conjecture: For n >= 1, A067628(a(n+2)) appears for the first time in A067628. Equivalently, A067628(a(n+2)) is the first T such that the minimal perimeter of polyiamonds of T triangles is a(n+2). - Winston C. Yang (winston(AT)cs.wisc.edu), Feb 05 2002 Conjecture: Let b(n>=0) = (0, 1, 1, 1, 1, 3, 1, 3, 3, 3, 3, 5, 3, 5, 5, 5, 5, 7, 3, ...). Equivalently, let b(0) = 0, b(n>=1) = 2*floor((n-1)/6) + 1 + (2 if n+1=0 mod 6; 0 else). Then a(0) = 1, a(n>=1) = a(n-1) + b(n-1). - Winston C. Yang (winston(AT)cs.wisc.edu), Feb 05 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000813 | OEIS-A008780

a(n) = (n-dimensional partitions of 6) + C(n,4) + C(n,3).

These are the conjectured numbers of d-dimensional partitions for n=6, coming from a formula proposed by MacMahon in the general case that turned out to be wrong. Still, here for n=6, MacMahon's formula coincides for d < 3 with the first three terms of A042984. - _Michel Marcus_, Aug 16 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000814 | OEIS-A008784

Numbers k such that sqrt(-1) mod k exists; or, numbers that are primitively represented by x^2 + y^2.

In appears that {a(n)} is the set of proper divisors of numbers of the form m^2+1. - Kaloyan Todorov (kaloyan.todorov(AT)gmail.com), Mar 25 2009 [This conjecture is correct. - _Franklin T. Adams-Watters_, Oct 07 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000815 | OEIS-A008793

The problem of the calissons: number of ways to tile a hexagon of edge n with diamonds of side 1. Also number of plane partitions whose Young diagrams fit inside an n X n X n box.

Conjecture: the supercongruences a(n*p^r) == a(n*p^(r-1))^p (mod p^(4*r)) hold for all primes p and positive integers n and r. - _Peter Bala_, Apr 07 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonas Wallgren_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000816 | OEIS-A008846

Hypotenuses of primitive Pythagorean triangles.

Conjecture: numbers p for which sqrt(-1) exists in the p-adic numbering system. For example the 5-adic number ...2431212, when squared, gives ...4444444, which is -1, and 5 is in the sequence. - _Thierry Banel_, Aug 19 2022 The above conjecture was proven true by George Bergman. 3 known facts: (1) prime factors of a(n) are equal to 1 mod 4, (2) modulo such primes, sqrt(-1) exists, (3) if sqrt(m) exists mod r, r being odd, this extends to sqrt(m) in the r-adic ring. - _Thierry Banel_, Jul 04 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Ralph Peterson (RALPHP(AT)LIBRARY.nrl.navy.mil) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000817 | OEIS-A008848

Squares whose sum of divisors is a square.

Conjectures: (1) a(2) = 81 is the only prime power (A246655) in this sequence. (2) 81 and 400 are only terms x for which sigma(x) is in A246655. (3) x = 1 is the only such term that sigma(x) is also a term. See also comments in A074386, A336547 and A350072. - _Antti Karttunen_, Jul 03 2023, (2) corrected in May 11 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000818 | OEIS-A008892

Aliquot sequence starting at 276.

It is an open question whether this sequence ever reaches 0. The trajectory has been calculated to 2145 terms, and is still growing, term 2145 being a 214-digit number (see FactorDB link). - _N. J. A. Sloane_, Jan 11 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000819 | OEIS-A008909

Join 2n points on a line with n arcs above the line; form graph with the arcs as nodes, joining 2 nodes when the arcs cross. a(n) is the number of cases in which the graph is a path.

G.f. (conjecture): 1 - G(0)/(1-x), where G(k)= 1 - 1/(1 - x/(1 - x/(1 - x/(1 - x/(x - 1/G(k+1)))))); (continued fraction). - _Sergei N. Gladkovskii_, Jul 12 2013 G.f. (conjecture): (2*x^3-x^2+2*x-1+sqrt(x^4+2*x^2-4*x+1))/(2*x^2-2*x). - _Michael D. Weiner_, Dec 17 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Martin Klazar_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000820 | OEIS-A008934

Number of tournament sequences: sequences (a_1, a_2, ..., a_n) with a_1 = 1 such that a_i < a_{i+1} <= 2*a_i for all i.

Conjecture: a(n) is given by a series of nested sums as follows:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mauro Torelli_, _Jeffrey Shallit_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000821 | OEIS-A008978

a(n) = (5*n)!/(n!)^5.

For positive integer r, define sequences {u(n) : n >= 0} and {v(n) : n >= 0} by setting u(n) = [x^n] E(x)^(r*n) and v(n) = [x^n] F(x)^(r*n), where F(x) is defined above in the posting dated Jul 12 2016. We conjecture that the supercongruences u(m*p^k) == u(m*p^(k-1)) (mod p^(3*k)) and v(m*p^k) == v(m*p^(k-1)) (mod p^(3*k)) hold for all primes p >= 5 and positive integers m and k. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000822 | OEIS-A008996

Increasing length runs of consecutive composite numbers (records).

Conjecture: a(n) = O(n^2); specifically, a(n) <= n^2. - _Alexei Kourbatov_, Jan 23 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | Mark Cramer (m.cramer(AT)qut.edu.au), Mar 15 1996 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000823 | OEIS-A009003

Hypotenuse numbers (squares are sums of 2 nonzero squares).

Conjecture: these are bases n in which exists an n-adic integer x satisfying x^5 = x, and 5 is the smallest k>1 such that x^k =x (so x^2, x^3 and x^4 are not x). Example: the 10-adic integer x = ...499879186432 (A120817) satisfies x^5 = x, and x^2, x^3, and x^4 are not x, so 10 is in this sequence. See also A120817, A210850 and A331548. - _Patrick A. Thomas_, Mar 01 2020 Conjecture: Numbers m such that the limit: Limit_{s->1} zeta(s)*Sum_{k=1..m} [k|m]*A008683(k)*(i^k)/(k^(s - 1)) exists, which is equivalent to numbers m such that abs(Sum_{k=1..m} [k|m]*A008683(k)*(i^k)) = 0. - _Mats Granvik_, Jul 06 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000824 | OEIS-A009101

Fixed point when iterating the function f on n, where f(x) = x + product of digits of x.

Record values of the number of steps to obtain a(n) are 10 for n = 1, 27 for n = 187, 28 for n = 3237326, 32 for n = 3515987, 33 for n = 22572473. It is conjectured that a(n) is obtained in a finite number of steps for every n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000825 | OEIS-A009190

"2p-twin peaks": a(n) is the least x with lpf(x) = lpf(x + 2p) = p = prime(n) and lpf(y) < p for all x < y < x + 2p, where lpf = least prime factor; a(n) = -1 if no such x exists.

It is (weakly) conjectured that a(n) = -1 for 14 <= n <= 19, which is why this entry has offset 20. - _N. J. A. Sloane_, Aug 19 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000826 | OEIS-A009191

a(n) = gcd(n, d(n)), where d(n) is the number of divisors of n (A000005).

First occurrence of k: 1, 2, 9, 8, 400, 12, 3136, 24, 36, 80, 123904, 60, 692224, 448, 2025, 384, ... Conjecture: each k is present. - _Robert G. Wilson v_, Mar 27 2013 Conjecture is true. See _David A. Corneth_'s comment in A324553. - _Antti Karttunen_, Mar 06 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000827 | OEIS-A009388

If a, b in sequence, so is a*b-1.

All terms are congruent to 0 or 2 mod 3. It follows that no three consecutive integers are in the sequence. - _Franklin T. Adams-Watters_, Aug 31 2016, conjectured by _David W. Wilson_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000828 | OEIS-A009744

Expansion of e.g.f. tan(x)*sin(x) (even powers only).

Conjectural o.g.f.: Sum_{n >= 0} 4*x/2^n * Sum_{k = 0..n} (-1)^k*(k+1)*binomial(n,k)/((1 + x*(2*k + 1)^2)*(1 + x*(2*k + 3)^2)) = 2*x + 4*x^2 + 62*x^3 + 1384*x^4 + ... - _Peter Bala_, Mar 03 2015 Bala's conjectured o.g.f. may be simplified to 2 * Sum_{k >= 0} (-1)^k/(1 + x*(2*k+1)^2) - 1/(1 + x). - _Sela Fried_, Dec 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000829 | OEIS-A009843

Expansion of e.g.f. x/cos(x) (odd powers only).

Conjecturally, a(n) = the unsigned numerator of B(2*n+1,1/4). The conjecture is true (see Fried link). - _Sela Fried_, Dec 22 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000830 | OEIS-A010033

The one-sided one-lie Rényi-Ulam game: a(n) is the maximum size of a set M such that one can determine a chosen element x with at most n questions of the form "Is x in T?" for some set T, when the answer can be a lie once, but only when the true answer is Yes.

a(n+1) = 2*a(n) - b(n) with b = o(a), b(n) > 0 for all n >= 0 except for b(1) = b(3) = 0; b(n+1) = 2*b(n) - c(n) with c = o(b) (conjectured). - _M. F. Hasler_, Jun 26 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | Thorsten Prenzel (prenzel(AT)unitas.or.uni-bonn.de) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000831 | OEIS-A010048

Triangle of Fibonomial coefficients, read by rows.

Conjecture: polynomials with (positive) Fibonomial coefficients are reducible iff n odd > 1. - _Ralf Stephan_, Oct 29 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000832 | OEIS-A010051

Characteristic function of primes: 1 if n is prime, else 0.

The spectral components (excluding the zero frequency) of the Fourier transform of the partial sequences {a(j)} with j=1..n and n an even number, exhibit a remarkable symmetry with respect to the central frequency component at position 1 + n/4. See the Fourier spectrum of the first 2^20 terms in Links, Comments in A289777, and Conjectures in A001223 of Sep 01 2019. It also appears that the symmetry grows with n. - _Andres Cicuttin_, Aug 23 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000833 | OEIS-A010052

Characteristic function of squares: a(n) = 1 if n is a square, otherwise 0.

a(n*m) = a(n/gcd(n,m))*a(m/gcd(n,m)) for all n and m > 0 (conjectured). - _Velin Yanev_, Feb 13 2019 [Proof from _Michael B. Porter_, Feb 16 2019: If nm is a square, nm = product_i (p_i^2), where p_i are prime, not necessarily distinct. Each p_i either appears twice in n, twice in m, or one time in each and therefore in the gcd. So n/gcd(n,m) and m/gcd(n,m) are both squares. If nm is not a square, there is a q_j that appears in one of n or m but not in the gcd. So either n/gcd(n,m) or m/gcd(n,m) is not a square.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000834 | OEIS-A010055

1 if n is a prime power p^k (k >= 0), otherwise 0.

Conjecture: a(n) = (n - A048671(n))/A000010(n) for all n > 1. - _Velin Yanev_, Mar 10 2021 [The conjecture is true. - _Andrey Zabolotskiy_, Mar 11 2021]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000835 | OEIS-A010059

Another version of the Thue-Morse sequence: let A_k denote the first 2^k terms; then A_0 = 1 and for k >= 0, A_{k+1} = A_k B_k, where B_k is obtained from A_k by interchanging 0's and 1's.

Two conjectures: If n is even, the ratio of 1's to 0's = 1:1. Conjecture (verified for the first 280000 entries): this is the characteristic function of A001969. - _R. J. Mathar_, Sep 05 2008 Proof of these three conjectures: the first two follow directly from the third, because the sequence A010059 is the binary complement of the Thue-Morse sequence A010060. For the third conjecture: the odious and evil numbers occur as quadruples EOOE and OEEO, simply by their definition. To obtain the mod 2 version of the interleave of the odious and evil numbers we therefore have to apply a transformation

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000836 | OEIS-A010331

Numbers that are expressible as C(m,5) + C(n,5) in at least 2 ways.

No additional solutions with n <= m <= 25000 exist. Conjecture: no additional solutions exist. - _Jon E. Schoenfield_, Apr 13 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000837 | OEIS-A010466

Decimal expansion of square root of 8.

Conjecture: Equals Sum_{k>=0} A210067(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000838 | OEIS-A010503

Decimal expansion of 1/sqrt(2).

Conjecture: Equals Sum_{k>=0} A139820(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000839 | OEIS-A010527

Decimal expansion of sqrt(3)/2.

Gilbert & Pollak conjectured that this is the Steiner ratio rho_2, the least upper bound of the ratio of the length of the Steiner minimal tree to the length of the minimal tree in dimension 2. (See Ivanov & Tuzhilin for the status of this conjecture as of 2012.) - _Charles R Greathouse IV_, Dec 11 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000840 | OEIS-A010767

Decimal expansion of 4th root of 2.

Conjecture: Equals Sum_{k>=0} A007096(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025 Plouffe's conjecture is correct: Sum_{k >= 0} A007096(k) / exp(k*Pi) = eta(q^2)^6 / (eta(q)^4 * eta(q^4)^2) evaluated at q = exp(-Pi), with the convention for Dedekind's eta function that if q = exp(2*Pi*i*t) then eta(q) = eta(t). At q = exp(-Pi), t = i/2. Hence the sum equals eta(i)^6/(eta(i/2)^4 * eta(2*i)^2), which simplifies to 2^(1/4) using the known values eta(i) = alpha/2, eta(i/2) = alpha/2^(7/8) and eta(2*i) = alpha/2^(11/8), where alpha = Gamma(1/4)/Pi^(3/4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000841 | OEIS-A010786

Floor-factorial numbers: a(n) = Product_{k=1..n} floor(n/k).

Conjecture: c = A085361. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000842 | OEIS-A010791

a(n) = n!*(n+2)!/2.

This sequence has been shown to contain infinitely many squares. From the Hong and Liu abstract: Recently Cilleruelo proved that the product Product_{k=1..n} (k^2 + 1) is a square only for n = 3 which confirms a conjecture of Amdeberhan, Medina and Moll. In this paper, we show that the sequence Product_{k=2..n} (k^2 - 1) contains infinitely many squares. Furthermore, we determine all squares in this sequence. We also give a formula for the p-adic valuation of the terms in this sequence. - _Jonathan Vos Post_, Oct 21 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000843 | OEIS-A010815

From Euler's Pentagonal Theorem: coefficient of q^n in Product_{m>=1} (1 - q^m).

Conjecture: Sum_{k>=0} a(k) / exp(k*Pi) = (1/2) * exp(1/24*Pi) * Pi^(1/4) * 2^(5/8) / Gamma(3/4) = A259148. - _Simon Plouffe_, Sep 09 2025 The conjecture is correct: By Euler's Pentagonal Number Theorem, the sum equals Product_{k >= 1} (1 - exp(-Pi*k)) = eta(i/2)*exp(Pi/24), where eta(t) = q^(1/24) * Product_{n >= 1} (1 - q^n), q = exp(2*Pi*i*t), is Dedekind's eta fumction. The conjecture follows from the standard result that eta(i/2) = Gamma(1/4)/(2^(7/8)*Pi^(3/4)) (see, for example, the Wikipedia entry on the eta function) and the reflection formula for the Gamma function. - _Peter Bala_, May 22 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000844 | OEIS-A010845

a(n) = 3*n*a(n-1) + 1, a(0) = 1.

Conjecture: a(n) +(-3*n-1)*a(n-1) +3*(n-1)*a(n-2)=0. - _R. J. Mathar_, Feb 16 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000845 | OEIS-A010904

Pisot sequence E(4,14): a(n) = floor(a(n-1)^2/a(n-2)+1/2) for n>1, a(0)=4, a(1)=14.

Theorem: a(0)=4, a(1)=14, a(2)=49; for n>2, a(n) = 4*a(n-1)-2*a(n-2)+a(n-3). Proved using the PtoRv program of Ekhad-Sloane-Zeilberger. (Conjectured by _Harvey P. Dale_, May 02 2012.) - _N. J. A. Sloane_, Sep 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000846 | OEIS-A010905

Pisot sequence E(4,15): a(n) = floor(a(n-1)^2/a(n-2)+1/2) for n>1, a(0)=4, a(1)=15.

This was conjectured by _Colin Barker_, Apr 16 2012, and implies the G.f.: (4-x)/(1-4*x+x^2) and the formula a(n) = ((1+sqrt(3))^(2*n+4)-(1-sqrt(3))^(2*n+4))/(2^(n+3)*sqrt(3)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000847 | OEIS-A010908

Pisot sequence E(4,21), a(n) = floor(a(n-1)^2/a(n-2) + 1/2).

Conjectures from _Colin Barker_, Jun 04 2016: (Start) Theorem: a(n) = 6*a(n-1) - 4*a(n-2) for n >= 2. (Proved using the PtoRv program of Ekhad-Sloane-Zeilberger.) This implies the above conjectures. - _N. J. A. Sloane_, Sep 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000848 | OEIS-A010909

Pisot sequence E(4,25): a(n) = floor(a(n-1)^2/a(n-2)+1/2) for n>1, a(0)=4, a(1)=25.

Theorem: a(n) = 6 a(n - 1) + a(n - 2) + 3 a(n - 3). (Conjectured by _Colin Barker_, Jun 05 2016. Proved using the PtoRv program of Ekhad-Sloane-Zeilberger.) - _N. J. A. Sloane_, Sep 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000849 | OEIS-A010910

Pisot sequence E(4,27): a(n) = floor(a(n-1)^2/a(n-2)+1/2) for n>1, a(0)=4, a(1)=27.

Conjectures from _Colin Barker_, Jun 05 2016: (Start) Theorem: a(n) = 6 a(n - 1) + 5 a(n - 2) for n >= 2. Proved using the PtoRv program of Ekhad-Sloane-Zeilberger, and implies the above conjectures. - _N. J. A. Sloane_, Sep 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000850 | OEIS-A010911

Pisot sequence E(3,11), a(n) = floor(a(n-1)^2/a(n-2) + 1/2).

Theorem: a(n) = 3 a(n - 1) + 2 a(n - 2) + a(n - 3) for n>=3. Proved using the PtoRv program of Ekhad-Sloane-Zeilberger, and implies the above conjectures. - _N. J. A. Sloane_, Sep 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000851 | OEIS-A010912

Pisot sequences E(3,7), P(3,7).

Since Pisot (1938) showed that E(3,k) always satisfies a linear recurrence, presumably it would not be difficult to prove that the above conjectures are correct. - _N. J. A. Sloane_, Jul 30 2016 Theorem: a(n) = 3 a(n - 1) - 2 a(n - 2) + a(n - 3) for n>=3. Proved using the PtoRv program of Ekhad-Sloane-Zeilberger, and implies the above conjectures. - _N. J. A. Sloane_, Sep 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000852 | OEIS-A010913

Pisot sequence E(3,17), a(n) = floor(a(n-1)^2/a(n-2)+1/2).

Conjecture: If p[i]=fibonacci(2i+2) and if A is the Hessenberg matrix of order n defined by: A[i,j]=p[j-i+1], (i<=j), A[i,j]=-1, (i=j+1), and A[i,j]=0 otherwise. Then, for n>=1, a(n-1)= det A. - _Milan Janjic_, May 08 2010 Conjectures from _Colin Barker_, Jun 05 2016: (Start) Theorem: a(n) = 6 a(n - 1) - 2 a(n - 2) for n>=2. Proved using the PtoRv program of Ekhad-Sloane-Zeilberger, and implies the above conjectures. - _N. J. A. Sloane_, Sep 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000853 | OEIS-A010915

Pisot sequence E(6,16), a(n) = floor(a(n-1)^2/a(n-2) + 1/2).

Conjectured not to satisfy a linear recurrence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000854 | OEIS-A010920

Pisot sequence T(3,13), a(n) = floor(a(n-1)^2/a(n-2)).

Note the warning in A010925 from Pab Ter (pabrlos(AT)yahoo.com), May 23 2004: [A010925] and other examples show that it is essential to reject conjectured generating functions for Pisot sequences until a proof or reference is provided. - _N. J. A. Sloane_, Jul 26 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000855 | OEIS-A010924

Pisot sequence E(8,55), a(n) = floor(a(n-1)^2/a(n-2) + 1/2).

Conjecture: a(n) = 6*a(n-1) + 6*a(n-2), n > 1; a(0)=8, a(1)=55; g.f.: (8+7x)/(1-6x-6x^2). - _Philippe Deléham_, Nov 19 2008 Theorem: a(n) = 6*a(n-1) + 6*a(n-2) for n >= 2. Proved using the PtoRv program of Ekhad-Sloane-Zeilberger,

and implies the above conjectures. - _N. J. A. Sloane_, Sep 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000856 | OEIS-A010925

Pisot sequence T(5,21), a(n) = floor(a(n-1)^2/a(n-2)).

This and other examples show that it is essential to reject conjectured generating functions for Pisot sequences until a proof or reference is provided. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000857 | OEIS-A011257

Numbers k such that the geometric mean of phi(k) and sigma(k) is an integer.

For these terms the arithmetic mean is also an integer. It is conjectured that sigma(k) for these numbers is never odd. See also A065146, A028982, A028983. - _Labos Elemer_, Oct 18 2001 From a(2633) = 6931232 on, it is no longer true (as was once conjectured) that a(n) > n^2. - _M. F. Hasler_, Feb 07 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000858 | OEIS-A011541

Taxicab, taxi-cab or Hardy-Ramanujan numbers: the smallest number that is the sum of 2 positive integral cubes in n ways.

Conjecture: the number of distinct prime factors of a(n) is strictly increasing as n grows (this is not true if a(7) is equal to the upper bound given above), but never exceeds 2*n. - _Sergey Pavlov_, Mar 01 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000859 | OEIS-A011553

Number of standard Young tableaux of type (n,n,n) whose (2,1) entry is odd.

Conjecture: D-finite with recurrence 6*(n+2)*(n+1)^2*a(n) -(n+1)*(164*n^2-179*n+51) *a(n-1) +(46*n^3-609*n^2+812*n+12) *a(n-2) +12*(3*n-4) *(2*n-5) *(3*n-5)*a(n-3)=0. - _R. J. Mathar_, Nov 22 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | giambruno(AT)ipamat.math.unipa.it | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000860 | OEIS-A011558

Expansion of (x + x^3)/(1 + x + ... + x^4) mod 2.

Conjecture: a(n+1) is the number of ways of partitioning n into distinct parts of A084215. - _R. J. Mathar_, Mar 01 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000861 | OEIS-A011774

Nonprimes k that divide sigma(k) + phi(k).

Conjecture: For n > 1, a(n) is a Zumkeller number (A083207). Verified for all n in [2,63]. - _Ivan N. Ianakiev_, Jan 25 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000862 | OEIS-A011848

a(n) = floor(binomial(n, 2)/2).

Conjecture: For n = 2, a(n) is the greatest k such that A123663(k) < A000217(n - 2). - _Peter Kagey_, Nov 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 11 1996 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000863 | OEIS-A011946

Number of Barlow packings with group P63/mmc(S) that repeat after 4n layers.

Conjecture: Mobius transform of A127804. - _R. J. Mathar_, Sep 14 2011 Conjecture: a(n) = A045683(2n). - _R. J. Mathar_, Apr 15 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000864 | OEIS-A011965

Second differences of Bell numbers.

Conjecture: a(n) = Sum_{k=0..2^n - 1} b(k) for n >= 0 where b(2n+1) = b(n) + b(A025480(n-1)), b(2n) = b(n - 2^f(n)) + b(2n - 2^f(n)) + b(A025480(n-1)) for n > 0 with b(0) = b(1) = 1 and where f(n) = A007814(n). Also b((4^n - 1)/3) = A141154(n+1). - _Mikhail Kurkov_, Jan 27 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000865 | OEIS-A012200

arctanh(arctan(arcsin(x))) = x + 1/3!*x^3 + 17/5!*x^5 + 393/7!*x^7 + 25601/9!*x^9 + ...

Conjecture: a(n) ~ (2*n)! / (sin(tan(1)))^(2*n+1). Because sin(tan(1)) = 0.9999103740052..., is not possible to verify this numerically from 1000 terms. - _Vaclav Kotesovec_, Feb 07 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick Demichel_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000866 | OEIS-A012780

Expansion of e.g.f. arcsin(tan(x)) (odd powers only).

Conjecture: the sequence {a(n)} reduced modulo a positive integer k is eventually periodic with the period dividing phi(k). For example, the sequence taken modulo 7 begins [1, 3, 3, 0, 4, 4, 0, 3, 3, 0, 4, 4, 0, ...], with an apparent period of 6 = phi(7) beginning at a(1). - _Peter Bala_, Feb 15 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick Demichel_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000867 | OEIS-A013302

Expansion of e.g.f. cosh(log(x+1)-arctanh(x)) (even powers only).

Conjectures from _R. J. Mathar_, Oct 05 2014: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick Demichel_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000868 | OEIS-A013588

Smallest positive integer not the determinant of an n X n {0,1}-matrix.

This majorizes the sequence of maximal determinants only up to the 6th term. It is conjectured that the sequence of maximal determinants majorizes this for all later terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Gerhard R. Paseman (paseman(AT)prado.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000869 | OEIS-A013640

f-vector for 20-dimensional simplicial polytope providing counterexample to unimodality conjecture.

f-vector for 20-dimensional simplicial polytope providing counterexample to unimodality conjecture. Fails unimodality conjecture because a(11) > a(12) < a(13). - _Sean A. Irvine_, Aug 17 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000870 | OEIS-A013641

f-vector for simplicial polytope providing counterexample to unimodality conjecture.

f-vector for simplicial polytope providing counterexample to unimodality conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000871 | OEIS-A013663

Decimal expansion of zeta(5).

Conjecture: for k >= 1, there exist rational numbers A(k), B(k) and c(k) such that zeta(5) = A(k) + B(k)*Sum_{n >= 1} (n^2 + 2*k*n + c(k))/(n*(n + 1)*...*(n + 2*k))^5. A similar conjecture can be made for the constant zeta(3). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000872 | OEIS-A013936

a(n) = Sum_{k=1..n} floor(n/k^2).

Conjecture : limit n ->infinity (Pi^2/6*n-a(n))/sqrt(n) = c = 1.45... - _Benoit Cloitre_, Jan 10 2003 In fact we have: a(n) = zeta(2)*n+zeta(1/2)*n^(1/2)+O(n^theta) with theta<1/2 and we conjecture that theta=1/4+epsilon is the best possible choice. Also a(n)=sum_{1<=k<=n}floor(sqrt(n/k)). - _Benoit Cloitre_, Nov 05 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Henri Lifchitz_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000873 | OEIS-A013937

a(n) = Sum_{k=1..n} floor(n/k^3).

a(n) = a(n-1)+A061704(n). a(n) = Sum_{k=1..n} floor((n/k)^(1/3)) with asymptotic formula: a(n) = zeta(3)*n+zeta(1/3)*n^(1/3)+O(n^theta) where theta<1/3 and we conjecture that theta=1/4+epsilon is the best possible choice. - _Benoit Cloitre_, Nov 05 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Henri Lifchitz_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000874 | OEIS-A013999

From applying the "rational mean" to the number e.

Conjecture: this is also the number of acyclic orientations of the complement of the path graph. - _Martin Rubey_, Oct 15 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | Domingo Gomez Morin (Dgomezm(AT)etheron.net) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000875 | OEIS-A014015

Denominators of sign-alternating Egyptian fraction expansion of e - 2.

a(n) >= a(n-1)^2 + a(n-1) for n >= 1, so the ratio log(a(n))/2^n is strictly increasing. But does it approach a limit? Conjecture: lim_{n->infinity} log(a(n))/2^n = 0.9748... - _Jon E. Schoenfield_, Jun 22 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000876 | OEIS-A014068

a(n) = binomial(n*(n+1)/2, n).

Conjectural: a(2*p) == p*(2*p + 1) (mod p^4) for prime p >= 5. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000877 | OEIS-A014076

Odd nonprimes.

Conjecture: There exist infinitely many pairs [a(n), a(n)+6] such that a(n)/3 and (a(n)+6)/3 are twin primes. - _Eric Desbiaux_, Sep 25 2014.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Warut Roonguthai_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000878 | OEIS-A014117

Numbers k such that m^(k+1) == m (mod k) holds for all m.

In 2005, B. C. Kellner proved E. W. Weisstein's conjecture that denom(B_n) = n only if n = 1806. - _Jonathan Sondow_, Oct 14 2013 Conjecture: Numbers n such that gcd(d+1, n) > 1 for every proper divisor d of n. Verified up to 10^696. - _David Radcliffe_, May 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Broadhurst_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000879 | OEIS-A014120

Smallest number of persistence n over product-of-nonzero-digits function.

Conjectured terms follow: It seems a safe conjecture that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000880 | OEIS-A014137

Partial sums of Catalan numbers (A000108).

Mathar's formula reduces to 2*(2*n-1)*C(n-1) = (n+1)*C(n), which is a known recurrence of the Catalan numbers, so the conjecture is true. - _Peter J. Taylor_, Mar 23 2015 G.f. (conjecture): Q(0)/(1-x), where Q(k)= 1 + (4*k + 1)*x/(k + 1 - 2*x*(k + 1)*(4*k + 3)/(2*x*(4*k + 3) + (2*k + 3)/Q(k+1))); (continued fraction). - _Sergei N. Gladkovskii_, May 14 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000881 | OEIS-A014153

Expansion of 1/((1-x)^2*Product_{k>=1} (1-x^k)).

Let t(n_, k_) = Sum_{i = 0..k} Sum_{j = 0..n} s(n, j)*C(i, j)*p(k - n - i), where s(n, j) are Stirling numbers of the first kind, C(i, j) are the number of compositions of i distinct objects into j parts, and p is the integer partition function. Then a(k) = t(2, k+2) (conjectured). The formula for t(n, k) is the same as at A126442 except that there the Stirling numbers are of the second kind. - _George Beck_, May 21 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000882 | OEIS-A014176

Decimal expansion of the silver mean, 1+sqrt(2).

Conjecture: 1+sqrt(2) = lim_{n->oo} A179807(n+1)/A179807(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000883 | OEIS-A014201

Number of solutions to x^2 + x*y + y^2 <= n excluding (0,0).

Conjecture: a(n) = 2*(Pi/sqrt(3))*n + O(n^(1/4 + epsilon)), similar to the Gauss circle or Dirichlet divisor problems. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000884 | OEIS-A014307

Expansion of the e.g.f. sqrt(exp(x) / (2 - exp(x))).

Conjecture from _Mikhail Kurkov_, Jun 24 2025: (Start) More generally, R(n,0,m) gives expansion of the e.g.f. (exp(x) / (m - (m-1)*exp(x)))^(1/m) for any m>0. (End) [Conjecture is true, see MathOverflow link. - _Mikhail Kurkov_, Mar 29 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000885 | OEIS-A014320

The next new gap between successive primes.

Conjecture: a(n) = O(n). See arXiv:2002.02115 for discussion. - _Alexei Kourbatov_, Jun 04 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | Hynek Mlcousek (hynek(AT)dior.ics.muni.cz) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000886 | OEIS-A014442

Largest prime factor of n^2 + 1.

All a(n) except for a(1) = 2 are the Pythagorean primes, i.e., primes of form 4*k + 1. Conjecture: every Pythagorean prime appears in a(n) at least once. Conjecture: every Pythagorean prime p appears exactly two times among the first p integers of the sequence. Further: if a(i) = a(j) = p and both i and j are less than p (and i is not equal to j), then i + j = p and ij == 1 (mod p). [If a(k) = p as well, then k > p; in fact, k is in A256011.] Two examples: a(2) = a(3) = 5, with 2 + 3 = 5 and 2*3 = 6 == 1 (mod 5); a(4) = a(13) = 17, with 4 + 13 = 17 and 4*13 = 52 == 1 (mod 17). (End) The conjecture is true. If p is a Pythagorean prime, -1 is a quadratic residue mod p. Then -1 has exactly two square roots mod p, i.e., there are exactly two integers x,y with 1 <= x,y <= p-1 such that x^2 == y^2 == -1 (mod p), i.e., p divides x^2 + 1 and y^2 + 1, and moreover y == -x (mod p) so x + y = p, and x*y == -x^2 == 1 (mod p). Any other prime factor q of x^2 + 1 must divide (x^2 + 1)/p, and since x^2+1 < p^2 we have q < p, so a(x) = p and similarly a(y) = p. - _Robert Israel_, Jun 11 2015 Conjecture: if n is even and a(n) > n, then n+a(n) is in A256011. Examples: 2 + a(2) = 2 + 5 = 7, 4 + a(4) = 4 + 17 = 21, 6+a(6) = 6+37 = 43, and so on. Note that 18+a(18) is NOT in A256011, but 18 itself is. - _Michael Kaltman_, Jun 13 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | Glen Burch (gburch(AT)erols.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000887 | OEIS-A014484

Expansion of (1+2x)/(1-2x)^4 (E.g.f.).

Conjecture: a(n) +(-2*n-9)*a(n-1) +6*(n+1)*a(n-2)=0. - _R. J. Mathar_, Jun 11 2013 Conjecture verified by _Robert Israel_, Jul 11 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000888 | OEIS-A014495

Central binomial coefficient - 1.

D-finite with recurrence: -(n+1)*(n-2)*a(n) +(n^2+n-4)*a(n-1) +2*(n-1)*(2*n-5)*a(n-2) -4*(n-1)*(n-2)*a(n-3)=0. - Conjectured by _R. J. Mathar_, Jan 04 2017, confirmed by _Robert Israel_, Feb 11 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Denis Pochuev (denis(AT)cdc.Informatik.TH-Darmstadt.de) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000889 | OEIS-A014525

Order of shuffle group for deck of 3n cards.

Conjecture: a(3^m) = (m+1) * 6^(m+1), a(4*m) = (1/2) * (3*4*m)!, otherwise a(n) = (3*n)!. - _Sean A. Irvine_, Nov 07 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000890 | OEIS-A014543

Crossing counts in toroidal drawings of K_n exhibited by Guy, Jenkyns and Schaer (1968).

For 11 <= n <= 16, the exhibited drawings establish only cr_1(K_n) <= a(n); Guy, Jenkyns and Schaer conjectured equality for n=11. - _Eric W. Weisstein_, Aug 29 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000891 | OEIS-A014549

Decimal expansion of 1 / M(1,sqrt(2)) (Gauss's constant).

Conjecture: Equals Sum_{k>=0} A104794(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000892 | OEIS-A014569

Super-3 Numbers (3n^3 contains substring '333' in its decimal expansion).

Conjecture: a(n) ~ n. - _Charles R Greathouse IV_, Dec 04 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000893 | OEIS-A014573

Smallest k such that phi(x) = k has exactly n solutions, n>=0 with Carmichael conjecture.

Smallest k such that phi(x) = k has exactly n solutions, n>=0 with Carmichael conjecture. Carmichael conjectured that no term exists for n=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000894 | OEIS-A014577

The regular paper-folding sequence (or dragon curve sequence). Alphabet {1,0}.

Since the Heighway dragon is composed of right angles, it can be mapped with unit trajectories (Right = 1, Left = (-1), Up = i and Down = -i) on the complex plane where i = sqrt(-1). The initial (0th) iterate is chosen in this case to be the unit line from (0,0) to (-1,0). Then follow the directions below as indicated, resulting in a reflected variant of the dragon curve shown at the Eric Weisstein link. The conjectured system of complex plane trajectories is: The conjecture succeeds through the 4th iterate. It appears that to generate the (n+1)-th row, bring down the n-th row as the left half of row (n+1). For the right half of row (n+1), bring down the n-th row but change the signs of the first half of row n. For example, to get the complex plane instructions for the third iterate of the dragon curve, bring down (-1, i, 1, i) as the left half, and the right half is (1, -i, 1, i). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000895 | OEIS-A014587

Nim function for Take-a-Factorial-Game (a subtraction game).

Conjecture: Appears to be periodic with period of length 25 = [0, 1, 2, 0, 1, 2, 3, 0, 1, 2, 0, 1, 2, 3, 0, 1, 2, 0, 1, 2, 3, 0, 1, 2, 3] starting with the initial term (there is no preamble). - _Michel Dekking_, Jul 26 2019 This conjecture is false, because moving from 10! = 3628800 to 0 is a legal move, and so a(3628800) cannot be zero. A similar argument shows that for no value of P is this sequence periodic with period P starting at term 0 (for a(P!) cannot be zero). - _Nathan Fox_, Jul 28 2019. The first counterexample to the conjecture above is a(5050) = 4. - _Pontus von Brömssen_, Jul 09 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Achim Flammenkamp_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000896 | OEIS-A014595

Conjectured dimensions of spaces of weight systems of chord diagrams.

Conjectured dimensions of spaces of weight systems of chord diagrams.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Broadhurst_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000897 | OEIS-A014596

Conjectured numbers of Vassiliev invariants of knots.

Conjectured numbers of Vassiliev invariants of knots.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Broadhurst_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000898 | OEIS-A014605

Partial sums of A001935; at one time this was conjectured to agree with A007478.

Partial sums of A001935; at one time this was conjectured to agree with A007478.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Broadhurst_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000899 | OEIS-A014617

Antidiagonals of the prime-composite array B(m,n) (see A067681) that are zeros from the first Borve conjecture.

Antidiagonals of the prime-composite array B(m,n) (see A067681) that are zeros from the first Borve conjecture. Let c(m) be the m-th composite and p(n) be the n-th prime. The prime-composite array, B, is defined such that each element B(m,n) is the highest power of p(n) that is contained within c(m). The m-th antidiagonal of the array consists of the m elements B(m,1), B(m-1,2), B(m-2,3), ..., B(1,m). The First Borve Conjecture states that there are infinitely many zero-only antidiagonals.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Feb 04 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000900 | OEIS-A014664

Order of 2 modulo the n-th prime.

Also A002326((p_n-1)/2). Conjecture: If p_n is not a Wieferich prime (1093, 3511, ...) then A002326(((p_n)^k-1)/2) = a(n)*(p_n)^(k-1). - _Vladimir Shevelev_, May 26 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000901 | OEIS-A014701

Number of multiplications to compute n-th power by the Chandah-sutra method.

Conjecture: a(n+1) is the minimal number of steps to go from 0 to n, by choosing before each step, after the first step, whether to keep the same step length or double it. The initial step length is 1. - _Jean-Marc Rebert_, May 15 2025 The above conjecture is correct. - _Neal Patel_, Aug 23 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | James Kilfiger (jamesk(AT)maths.warwick.ac.uk) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000902 | OEIS-A014742

Expansion of (1+x^2)/(1 - 2*x - 2*x^2 + x^3).

The preceding comment is a conjecture using a(n) = floor(LL(n)*L(n)) with LL(n) = sqrt(F(n+2)^2 + F(n)^2) and L(n) = LL(n-1), n >= 1 (its author agreed with this in an email). See also, e.g., the Koshy reference for the dissection fallacy, sect. 6, 100 - 108. The proof of the conjecture uses first the identity (LL(n)*LL(n-1))^2 - a(n)^2 = + 1 with a(n) = F(n-1)*F(n) + F(n+1)*F(n+2) (see the formula section for a(n)). This identity is due to the factorization of the left-hand side which is A(n)^2 with A(n) = F(n)*F(n+1) - F(n-1)*F(n+2). But A(n) = (-1)^(n+1) is a special instance of a well known Fibonacci identity (Koshy, p. 88, Nr. 19 for h=-1, k=2, F(-1) = 1). Now one has (LL(n)*LL(n-1))^2 = 1 + a(n)^2, that is LL(n)*LL(n-1) = sqrt(1 + a(n)^2). Because a(n) < sqrt(1 + a(n)^2) < a(n) + 1 (just square both inequalities using a(n) > 0) one has now proved that floor(LL(n)*LL(n-1)) = a(n), n >= 1. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000903 | OEIS-A014847

Numbers k such that k-th Catalan number C(2k,k)/(k+1) is divisible by k.

If A is the set of all a(k)'s, Pomerance proved that the upper density of A is at most 1 - log 2 = 0.30685... and conjectured that A has positive lower density. I improved Pomerance's result by showing that the upper density of A is at most 1 - log 2 - 0.05551 = 0.25134... Numerically, this upper density seems to be less than 0.11. - _Carlo Sanna_, Jan 28 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000904 | OEIS-A014865

Numbers k that divide s(k), where s(1)=1, s(j)=15*s(j-1)+j.

Conjecture: all terms are odd, all terms greater than 7 are composite, and all terms greater than 1 are divisible by 3 or 5 or 7 some combination therof. - _Harvey P. Dale_, Dec 04 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000905 | OEIS-A014945

Numbers k such that k divides 4^k - 1.

Conjecture: if k divides 4^k - 1, then (4^k - 1)/k is squarefree. - _Thomas Ordowski_, Dec 24 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000906 | OEIS-A014960

Integers n such that n divides 24^n - 1.

All n > 1 in the sequence are multiple of 23. - Conjectured by _Thomas Baruchel_, Oct 10 2003; proved by _Max Alekseyev_, Nov 16 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000907 | OEIS-A014963

Exponential of Mangoldt function M(n): a(n) = 1 unless n is a prime or prime power, in which case a(n) = that prime.

a(n) = exp(Sum_{k>=1} A191898(n,k)/k) for n>1 (conjecture). - _Mats Granvik_, Jun 19 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marc LeBrun_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000908 | OEIS-A014972

Expansion of (theta_3(q) / theta_4(q))^4 in powers of q; also of 1 / (1 - lambda(z)).

Conjecture: Sum_{k>=0} a(k) / exp(k*Pi) = 2. - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000909 | OEIS-A015134

Consider Fibonacci-type sequences b(0)=X, b(1)=Y, b(k)=b(k-1)+b(k-2) mod n; all are periodic; sequence gives number of distinct periods.

Conjectures from _Stephen Bartell_, Aug 20 2023: (Start) Conjectures from _Stephen Bartell_, Aug 16 2024: (Start) Conjecture: a(n) >= (n+1)/2. - _Elliot Nutt_, Dec 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Phil Carmody_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000910 | OEIS-A015713

Numbers m such that phi(m) * sigma(m) + k^2 is not a square for any k.

Conjecture: {4, p^(2*m), 2*p^(2*m), p = 4*k+3 is prime}. - _Sean A. Irvine_, Dec 06 2018 The conjecture is true. It can be proved using the multiplicative property of A062354(n), i.e., A062354(p^e) = p^(e-1)*(p^(e+1)-1), and that if m is a term then A007814(A062354(m)) = 1. - _Amiram Eldar_, Feb 11 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000911 | OEIS-A015762

Numbers n such that phi(n) | sigma_4(n).

sigma_{8j+4}(x)/phi(x) is an integer for j=0..500, x=1,2,3,6,249,498, and this is conjectured to hold for possible larger terms of A015762 and all j. Compare with comments to A015759, A091285, A015770. - _Labos Elemer_, May 27 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000912 | OEIS-A015910

a(n) = 2^n mod n.

This sequence is conjectured to include every integer k >= 0 except k = 1. A036236 includes a proof that k = 1 is not in this sequence, and n = A036236(k) solves a(n) = k for all other 0 <= k <= 1000. - _David W. Wilson_, Oct 11 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000913 | OEIS-A016016

Number of iterations of Reverse and Add which lead to a palindrome, or -1 if no palindrome is ever reached.

It is conjectured that a(196) = -1, see A023108.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000914 | OEIS-A016054

Numbers n such that (13^n - 1)/12 is prime.

For Repunits in bases from -14 to 14, base 13 is a lucky number with the highest relative rate of primes being discovered. Base 7 is the most unlucky base with the lowest rate of primes being discovered. There is a Generalized Repunit Conjecture implying that all bases will eventually converge to the same relative rate of occurrence (ref 1). - _Paul Bourdelais_, Mar 01 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000915 | OEIS-A016069

Numbers k such that k^2 contains exactly 2 distinct digits.

a(n) = ((n-1) mod 3 + 1)*10^(ceiling(n/3)-7) for n >= 34 (conjectured). - _Chai Wah Wu_, Dec 17 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000916 | OEIS-A016090

a(n) = 16^(5^n) mod 10^n: Automorphic numbers ending in digit 6, with repetitions.

Conjecture: For any m coprime to 10 and for any k, the density of n such that a(n) == k (mod m) is 1/m. - _Eric M. Schmidt_, Aug 01 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000917 | OEIS-A016578

Decimal expansion of log(3/2).

Conjectured asymptotic density of Fouvry primes (A073024), see link to mathoverflow discussion in A395819. - _Hugo Pfoertner_, May 08 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000918 | OEIS-A016627

Decimal expansion of log(4).

log(4) = Sum_{k >= 1} A191907(4, k)/k, (conjecture). - _Mats Granvik_, Jun 19 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000919 | OEIS-A016729

Highest minimal Hamming distance of any Type 4^H+ Hermitian additive self-dual code over GF(4) of length n.

Conjectures from _Chai Wah Wu_, Apr 13 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000920 | OEIS-A016742

Even squares: a(n) = (2*n)^2.

It is conjectured (I think) that a regular Hadamard matrix of order n exists iff n is an even square (cf. Seberry and Yamada, Th. 10.11). A Hadamard matrix is regular if the sum of the entries in each row is the same. - _N. J. A. Sloane_, Nov 13 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000921 | OEIS-A016744

a(n) = (2*n)^4.

Sum_{n>=1} 1/a(n) = Pi^4/1440 (conjecturally A258945).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000922 | OEIS-A016777

a(n) = 3*n + 1.

Erdős & Sárközy conjecture that a set of n positive integers with property P must have some element at least a(n-1) = 3n - 2. Property P states that, for x, y, and z in the set and z < x, y, z does not divide x+y. An example of such a set is {2n-1, 2n, ..., 3n-2}. Bedert proves this for large enough n. (This is an upper bound, and is exact for all known n; I have verified it for n up to 12.) - _Charles R Greathouse IV_, Feb 06 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 11 1996 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000923 | OEIS-A016789

a(n) = 3*n + 2.

The above conjecture is true - immediate with the special case of Fermat's little theorem: y^3 == y (mod 3). - _Sela Fried_, Oct 27 2025 Conjecture: a(n) = n XOR A005351(n+1) XOR A005352(n+1). - Gilian Breysens, Jul 21 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000924 | OEIS-A016957

a(n) = 6*n + 4.

Conjecture: this sequence contains some but not all, even numbers with odd abundance A088827. They appear in this sequence at indices A186424(n) - 1. - _John Tyler Rascoe_, Jul 09 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000925 | OEIS-A017101

a(n) = 8n + 3.

Conjecture: 2*a(n) is the half-period of oscillation of a Langton's ant colony that is n basic blocks in length. To construct such blocks use a pair of ants facing north at (x,y) and (x+1,y+2) (using Golly's coordinate system). Each successive block is placed 1 cell away from the previous one, i.e., the x coordinate shifts by 3, so we have (x+3k,y) and (x+3k+1,y+2). Also, because of the symmetry of the oscillation pattern, 4*a(n) is the length of the whole period (see MathOverflow link for details). - _Mikhail Kurkov_, Nov 20 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000926 | OEIS-A018187

Restricted Perrin pseudoprimes.

Some sources have conjectured the restricted Perrin pseudoprimes can be derived from the unrestricted Perrin pseudoprimes by checking if { M=[0,1,0; 0,0,1; 1,1,0]; Mod(M,n) == Mod(M,n)^n }. Counterexamples include 52437986833, 60518537641, 364573433665, and 4094040693601. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000927 | OEIS-A018188

The $620 prime list.

In particular the above conjecture implies that there is a BPSW pseudoprime smaller than 10^25286. - _Charles R Greathouse IV_, Sep 28 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jon Grantham (grantham(AT)math.uga.edu) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000928 | OEIS-A018851

a(n)^2 is smallest square beginning with n.

For a given n, d can range from 0 (if n is a square) to d_max, where it appears that d_max approx. equals 3 + floor(log_10(n/25)). The successive points where d_max increases are given in A272678, and that entry contains more precise conjectures about the values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000929 | OEIS-A018888

Numbers which are not the sum of seven nonnegative cubes.

Old name: Write n = m_1^3 + ... +m_k^3 where the m_i are positive integers and k is minimal; sequence gives conjectured list of numbers for which k = 8 or 9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000930 | OEIS-A018890

Numbers whose smallest expression as a sum of positive cubes requires exactly 7 cubes.

It is conjectured that a(121)=8042 is the last term. - _Jud McCranie_

猜想/问题提及段 | historical_or_current_mention_unreviewed | Anonymous | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000931 | OEIS-A018894

Numbers k such that sigma(k)/phi(k) sets a new record.

The LCM of these found terms is a proper divisor and for all primes p <= 13 the exponent is less than the assumed prime. Conjecture: These 466 terms are the terms <= 10^75. For all 240 terms 1 < t <= 10^40 the following holds: there exists a p|t such that t/p is a term. Conjecture: This holds for all terms t > 1. See attached file with terms assuming these conjectures. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel ten Voorde_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000932 | OEIS-A018917

Define the generalized Pisot sequence T(a(0),a(1)) by: a(n+2) is the greatest integer such that a(n+2)/a(n+1) < a(n+1)/a(n). This is T(3,5).

Conjecture: a(n)=a(n-1)+a(n-2)-a(n-4). G.f.: (3+2*x-x^3)/(1-x)/(1-x^2-x^3). [_Colin Barker_, Feb 16 2012] Conjecture: a(n) = a(n-1) + A000931(n+8). - _Reinhard Zumkeller_, Dec 30 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000933 | OEIS-A018918

Define the generalized Pisot sequence T(a(0),a(1)) by: a(n+2) is the greatest integer such that a(n+2)/a(n+1) < a(n+1)/a(n). This is T(3,6).

Conjectures from _Colin Barker_, Dec 21 2012: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000934 | OEIS-A018925

Number of isotropic subspaces of a 2n-dimensional unitary space over GF(4).

Conjecture: a(n) = Product_{k=1..n} (4^k + 1). - _Vaclav Kotesovec_, Nov 21 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000935 | OEIS-A019319

Number of possible chess diagrams after n plies.

Even though the sequence may be infinite (if none of the rules for draw is ever invoked by any of the players), the sequence becomes constant from a given rank n on, since it is increasing (I conjecture - even though some positions available at the n-th move might not be available on the (1+n)-th move) and bounded, thus it has a limit. The challenge is now to find this limit (or at least nontrivial upper bounds) and the rank from which on the sequence becomes constant. - _M. F. Hasler_, Feb 15 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | Bernd Schwarzkopf (schwarzkopf(AT)uni-duesseldorf.de) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000936 | OEIS-A019334

Primes with primitive root 3.

then by Artin's conjecture, Q(N) ~ C*N/log(N) ~ 2*C*(Pi(N,5) + Pi(N,7)), where C = A005596 is Artin's constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000937 | OEIS-A019335

Primes with primitive root 5.

then by Artin's conjecture, Q(N) ~ (20/19)*C*N/log(N) ~ (40/19)*C*(Pi(N,2) + Pi(N,3)), where C = A005596 is Artin's constant. Conjecture: if we further define

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000938 | OEIS-A019337

Primes with primitive root 7.

All terms apart from the first are == 5, 11, 13, 15, 17, 23 (mod 28) since 7 is a quadratic residue modulo any other prime. By Artin's conjecture, this sequence contains about 37.395% of all primes, that is, about 74.79% of all primes == 5, 11, 13, 15, 17, 23 (mod 28). - _Jianing Song_, Sep 05 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000939 | OEIS-A019338

Primes with primitive root 8.

On Artin's conjecture, a(n) ~ (5/3A) n log n, where A = A005596 is Artin's constant. - _Charles R Greathouse IV_, May 21 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000940 | OEIS-A019353

Primes with primitive root 27.

According to Artin's conjecture, the number of terms <= N is roughly ((3/5)*C)*PrimePi(N), where C is Artin's constant = A005596, PrimePi = A000720. Compare: the number of terms of A001122 that are no greater than N is roughly C*PrimePi(N). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000941 | OEIS-A019429

Continued fraction for tan(1/6).

Conjecture: a(n) = (-1+3*(-1)^n-6*(-1+(-1)^n)*n)/2 for n>1. a(n) = 2*a(n-2)-a(n-4) for n>5. G.f.: x*(x^4-x^3+6*x^2+x+5) / ((x-1)^2*(x+1)^2). - _Colin Barker_, May 28 2013 The above conjectures are correct. The simple continued fraction expansion for tan(1/6) may be derived by setting z = 1/6 in Lambert's continued fraction tan(z) = z/(1 - z^2/(3 - z^2/(5 - ...))) and then, after using an equivalence transformation, making repeated use of the identity 1/(n - 1/m) = 1/((n - 1) + 1/(1 + 1/(m - 1))).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000942 | OEIS-A019434

Fermat primes: primes of the form 2^(2^k) + 1, for some k >= 0.

It is conjectured that there are only 5 terms. Currently it has been shown that 2^(2^k) + 1 is composite for 5 <= k <= 32 (see Eric Weisstein's Fermat Primes link). - _Dmitry Kamenetsky_, Sep 28 2008 Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000943 | OEIS-A019442

Numbers m such that a Hadamard matrix of order m exists.

It is conjectured that this sequence consists of 1, 2 and all multiples of 4. Conjectured g.f.: (2*x^3 + x^2 + 1)/(x - 1)^2. - _Jean-François Alcover_, Oct 03 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 16 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000944 | OEIS-A019445

Form a permutation of the positive integers, p_1, p_2, ..., such that the average of each initial segment is an integer, using the greedy algorithm to define p_n; sequence gives p_1 + ... + p_n.

Somos's conjecture is proved in both Shapovalov (1996) and Venkatachala (2009). - _Jeffrey Shallit_, Jul 18 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_, Tom Halverson (halverson(AT)macalester.edu) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000945 | OEIS-A019446

a(n) = ceiling(n/tau), where tau = (1+sqrt(5))/2.

Here is a proof of the conjecture by Ivan N. Ianakiev. Let b = (b(n)) be the sequence of occurrences of the "singleton terms" in (a(n)). We have to show that b = A276885.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_, Tom Halverson (halverson(AT)macalester.edu) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000946 | OEIS-A019450

Conjectured formula for irreducible 5-fold Euler sums of weight 2n+13.

Conjectured formula for irreducible 5-fold Euler sums of weight 2n+13. The conjectured formula has been verified by _David Broadhurst_ up to a(12) = 165.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Broadhurst_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000947 | OEIS-A019459

Conjectured formula for irreducible 6-fold Euler sums of weight 2n+16.

Conjectured formula for irreducible 6-fold Euler sums of weight 2n+16. The conjectured formula has been verified by _David Broadhurst_ up to a(6) = 45.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Broadhurst_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000948 | OEIS-A019469

Numbers k such that k does not divide binomial(2*k-4, k-2).

Conjecture (confirmed, see links): sequence is union of powers of two > 1 (A000079) and 3 * A096304.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000949 | OEIS-A019505

a(1)=1; for n > 1, a(n) is the smallest number with the same number of divisors as 2*a(n-1).

Conjectures: Subsequence of A002182. [This conjecture is false. The 64th term is 97039187544499200, which has exactly 63360 divisors, but is NOT in A002182; which has the smaller number 74801040398884800, which has 64512 divisors. - _J. Lowell_, Nov 29 2021] The value of c is less than twice b. [This conjecture is false. Terms 121-123 are 93635537220943523586899838720000, 14258138622280036546187020896000, and 26139920807513400334676204976000. - _J. Lowell_, Jan 28 2022] The primes and the powers of 3 are the only numbers that never "come and go" from the lists of divisors of the numbers in the sequence as the terms advance. [This conjecture is false. The 122nd term, 14258138622280036546187020896000, is a multiple of 3^5=243, but the 123rd term, 26139920807513400334676204976000, is not. - _J. Lowell_, Jan 28 2022]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000950 | OEIS-A019565

The squarefree numbers ordered lexicographically by their prime factorization (with factors written in decreasing order). a(n) = Product_{k in I} prime(k+1), where I is the set of indices of nonzero binary digits in n = Sum_{k in I} 2^k.

Conjecture: there are no finite cycles of any length. My grounds for this conjecture: any finite cycle in this sequence, if such cycles exist at all, must have at least one member that occurs somewhere in A285319, the terms that seem already to be quite rare. Moreover, any such a number n should satisfy in addition to A019565(n) < n also that A048675^{k}(n) is squarefree, not just for k=0, 1 but for all k >= 0. As there is on average a probability of only 6/(Pi^2) = 0.6079... that any further term encountered on the trajectory of A048675 is squarefree, the total chance that all of them would be squarefree (which is required from the elements of A019565-cycles) is soon minuscule, especially as A048675 is not very tightly bounded (many trajectories seem to skyrocket, at least initially). I am also assuming that usually there is no significant correlation between the binary expansions of n and A048675(n) (apart from their least significant bits), or, for that matter, between their prime factorizations. See also the slightly stronger conjecture in A285320, which implies that there would neither be any two-way infinite cycles. If either of the conjectures is false (there are cycles), then certainly neither sequence A285332 nor its inverse A285331 can be a permutation of natural numbers. (End) The conjecture made in A087207 (see also A288569) implies the two conjectures mentioned above. A further constraint for cycles is that in any A019565-trajectory which starts from a squarefree number (A005117), every other term is of the form 4k+2, while every other term is of the form 6k+3. - _Antti Karttunen_, Jun 18 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marc LeBrun_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000951 | OEIS-A019609

Decimal expansion of Pi*e.

Not known to be irrational (though of course conjectured transcendental), see e.g. Klee & Wagon. - _Charles R Greathouse IV_, Jul 23 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000952 | OEIS-A019952

Decimal expansion of tangent of 54 degrees.

Conjecture: Product (2/3) * (8/7) * (12/13) * (18/17) * (22/23) * (32/33) * ... * (a_n/b_n) = sqrt(25 + 10*sqrt(5))/5 = tan(3*Pi/10) = A019952, where a_n even, a_n + b_n = a(n), |a_n - b_n| = 1, n >= 0. - _Dimitris Valianatos_, Feb 14 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000953 | OEIS-A020481

Least p with p, q both prime, p+q = 2n.

Conjecture: a(n) ~ O(n^1/2). - _Jon Perry_, Apr 29 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000954 | OEIS-A020483

Least prime p such that p+2n is also prime.

It is conjectured that a(n) always exists. a(n) has been computed for n < 5 * 10^11, with largest value a(248281210271) = 3307. - _Jens Kruse Andersen_, Nov 28 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000955 | OEIS-A020497

Conjecturally, this is the minimal y such that n primes occur infinitely often among (x+1, ..., x+y), that is, pi(x+y) - pi(x) >= n for infinitely many x.

Conjecturally, this is the minimal y such that n primes occur infinitely often among (x+1, ..., x+y), that is, pi(x+y) - pi(x) >= n for infinitely many x.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, _Christopher E. Thompson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000956 | OEIS-A020498

a(n) is the least number > a(n-1) such that there is no prime p for which a(1) through a(n) would contain all residues modulo p.

Conjecturally, a(n) is the smallest number such that n primes occur infinitely often among (x+a(1), ...,x+a(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_ and _Christopher E. Thompson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000957 | OEIS-A020665

a(n) is the (conjectured) maximal exponent k such that n^k does not contain a digit zero in its decimal expansion.

a(n) is the (conjectured) maximal exponent k such that n^k does not contain a digit zero in its decimal expansion. The following values and bounds are for the actual maximal exponents (not conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000958 | OEIS-A020698

a(n) = 5*a(n-1) - 2*a(n-2), with a(0)=2, a(1)=9.

Theorem: E(2,9) satisfies a(n) = 5 a(n - 1) 2 2 a(n - 2) for n>=2. Proved using the PtoRv program of Ekhad-Sloane-Zeilberger, and implies the conjecture. - _N. J. A. Sloane_, Sep 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000959 | OEIS-A020704

Pisot sequences E(3,10), P(3,10).

Conjectures from _Colin Barker_, Jun 05 2016: (Start) Theorem: For E(3,10), a(n) = 3 a(n - 1) + a(n - 2) for n>=2. Proved using the PtoRv program of Ekhad-Sloane-Zeilberger, and implies the above conjectures. - _N. J. A. Sloane_, Sep 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000960 | OEIS-A020708

Pisot sequences E(4,9), P(4,9).

Theorem: E(4,9) satisfies a(n) = 2 a(n - 1) + a(n - 3) for n >= 3. Proved using the PtoRv program of Ekhad-Sloane-Zeilberger, and implies the above conjectures. - _N. J. A. Sloane_, Sep 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000961 | OEIS-A020709

Pisot sequence E(4,10).

Theorem: a(n) = 3 a(n - 1) - 2 a(n - 2) + 2 a(n - 3) for n>=3. Proved using the PtoRv program of Ekhad-Sloane-Zeilberger, and implies the above conjectures. - _N. J. A. Sloane_, Sep 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000962 | OEIS-A020711

Pisot sequences E(5,7), P(5,7).

Theorem: E(5,7) satisfies a(n) = 3 a(n - 1) + 2 a(n - 2) + a(n - 3) - a(n - 4) for n >= 4. Proved using the PtoRv program of Ekhad-Sloane-Zeilberger, and implies the above conjectures. - _N. J. A. Sloane_, Sep 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000963 | OEIS-A020713

Pisot sequences E(5,9), P(5,9).

Theorem: E(5,9) satisfies a(n) = 2 a(n - 1) - a(n - 2) + a(n - 3) for n>=3. Proved using the PtoRv program of Ekhad-Sloane-Zeilberger, and implies the above conjectures. - _N. J. A. Sloane_, Sep 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000964 | OEIS-A020716

Pisot sequences E(6,8), P(6,8).

Theorem: E(6,8) satisfies a(n) = a(n - 1) + a(n - 2) - a(n - 4) for n>=4. Proved using the PtoRv program of Ekhad-Sloane-Zeilberger. This shows that the above conjectures are correct. - _N. J. A. Sloane_, Sep 10 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000965 | OEIS-A020718

Pisot sequences E(6,10), P(6,10).

G.f.: (-4*x^5+x^4+x^3-3*x^2-2*x+6)/((1-x)*(1-x-x^2-x^5)) (conjectured). - _Ralf Stephan_, May 12 2004 a(n) = 2*a(n-1)-a(n-3)+a(n-5)-a(n-6) for n>5 (conjectured). - _Colin Barker_, Jun 05 2016 Theorem: E(6,10) satisfies a(n) = 2 a(n - 1) - a(n - 3) + a(n - 5) - a(n - 6) for n >= 6. Proved using the PtoRv program of Ekhad-Sloane-Zeilberger. This shows that the above conjectures are correct. - _N. J. A. Sloane_, Sep 10 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000966 | OEIS-A020721

Pisot sequences E(7,10), P(7,10).

Theorem: E(7,10) satisfies a(n) = 2 a(n - 1) - a(n - 2) + a(n - 3) - a(n - 4) for n>=4. Proved using the PtoRv program of Ekhad-Sloane-Zeilberger. This shows that the above conjectures are correct. - _N. J. A. Sloane_, Sep 10 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000967 | OEIS-A020735

Odd numbers >= 5.

Sun conjectures that any member of this sequence is of the form m^2 + m + p, where p is prime. Blanco-Chacon, McGuire, & Robinson prove that the primes of this form have density 1. - _Charles R Greathouse IV_, Jun 20 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000968 | OEIS-A020744

Pisot sequences P(8,10), T(8,10).

Conjecturally, even sums of four primes. - _Charles R Greathouse IV_, Feb 16 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000969 | OEIS-A020746

Pisot sequence T(3,7), a(n) = floor(a(n-1)^2/a(n-2)).

Conjectured g.f.: (-x^5+x^4-x^3+x^2-2*x+3)/((1-x)*(1-2*x-x^3-x^5)). - _Ralf Stephan_, May 12 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000970 | OEIS-A020748

Pisot sequence T(4,10), a(n) = floor(a(n-1)^2/a(n-2)).

G.f.: (-3x^5+2x^4+x^3-x^2-2x+4)/[(1-x)(1-2x-x^2-2x^5)] (conjectured). - _Ralf Stephan_, May 12 2004 Note the warning in A010925 from Pab Ter (pabrlos(AT)yahoo.com), May 23 2004: [A010925] and other examples show that it is essential to reject conjectured generating functions for Pisot sequences until a proof or reference is provided. - _N. J. A. Sloane_, Jul 26 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000971 | OEIS-A020753

Sizes of successive increasing gaps between squarefree numbers.

Conjecture: a(n) ~ log(A020754(n))/2. - _Thomas Ordowski_, Jul 23 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000972 | OEIS-A020879

Number of elementary edge-subgraphs in Moebius ladder M_n.

Conjectures from _Colin Barker_, Dec 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000973 | OEIS-A020880

Number of strong elementary edge-subgraphs in Moebius ladder M_n.

Conjectures from _Colin Barker_, Dec 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000974 | OEIS-A020881

Number of strong restricted edge-subgraphs in Moebius ladder M_n.

Conjectures from _Colin Barker_, Dec 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000975 | OEIS-A020882

Ordered hypotenuses (with multiplicity) of primitive Pythagorean triangles.

The density of such points in a circle with radius squared = a(n) is ~ Pi * a(n). Restricting to a > b > 0 reduces this by a factor of 1/8; requiring gcd(a,b)=1 provides a factor of 6/Pi^2; and a, b not both odd is a factor of 2/3. (2/3, not 3/4, because the case a, b both even has already been eliminated.) Multiplying, a(n) * Pi * 1/8 * 6/Pi^2 * 2/3 is a(n) / (2 * Pi). But n is approximately this number of points, so a(n) ~ 2 * Pi * n. Conjectured by _David W. Wilson_, proof by _Franklin T. Adams-Watters_, Mar 15 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000976 | OEIS-A020903

Lim f(f(...f(n))) where f is the fractal sequence given by f(n)=A002260(n+1).

Conjecture: a(n) and a(n+1) are never both 2. - _Robert Israel_, Sep 03 2015 Proof of the conjecture: Let f(n)=A002260(n+1)=1,2,1,2,3,1,2,3,4,... Then (f(n)) is a concatenation of ladders 1,2 followed by 1,2,3 followed by 1,2,3,4 etc. The proof is by induction. Note that the sequence (a(n)) can be seen as map from the positive integers to the positive integers. The induction starts from the observation that a(1) and a(2) are not both 2. We use that f(k)<k for all k>2. Any pair (k,k+1) from a ladder has image (a(k),a(k+1)) = (a(f(k)),a(f(k+1))), which occurs either as image of two adjacent integers (j,j+1) earlier in the sequence, and so will not be equal to (2,2) by the induction hypothesis, or as image of a pair (j,1), whose image is also not equal to (2,2). The same holds for a pair consisting of the end of a ladder and the next entry. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000977 | OEIS-A020942

First column of 3rd-order Zeckendorf array A136189.

Conjecture: a(n) = A202342(n) + n. - _Sean A. Irvine_, May 05 2019 [proved in corrected form in Shallit (2025); it should read a(n) = A202342(n) + n-1]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000978 | OEIS-A020963

a(n) = Sum_{k>=1} floor(2*(1+sqrt(2))^(n-k)).

a(n) = 3*a(n-1) - 4*a(n-3) + a(n-4) + a(n-5) for n > 5 (conjectured). G.f.: x*(x^4 - 2*x^3 + x^2 - 2)/((x - 1)^2*(x + 1)*(x^2 + 2*x - 1)) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000979 | OEIS-A020964

a(n) = Sum_{k>=1} floor(3*(1+sqrt(2))^(n-k)).

Conjectures from _Chai Wah Wu_, Sep 14 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000980 | OEIS-A020999

Conjectured number of irreducible multiple zeta values of depth n and weight 3n (confirmed up to n=7).

Conjectured number of irreducible multiple zeta values of depth n and weight 3n (confirmed up to n=7).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Broadhurst_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000981 | OEIS-A021004

Pisot sequence P(4,10).

Conjecture: a(n) = 2a(n-1) + a(n-2) + a(n-3) - a(n-4) - a(n-6) (checked up to n = 1000) Conjectured G.f.: (4+2 x+x^2-2 x^3-x^4-2 x^5)/(1-2 x-x^2-x^3+ x^4+x^6) - _Harvey P. Dale_, Mar 12 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000982 | OEIS-A021008

Pisot sequence P(5,11), a(0)=5, a(1)=11, a(n+1) is the nearest integer to a(n)^2/a(n-1).

Conjecture: G.f.: (2x^3+x^2-4x+5)/(-x^4+2x^2-3x+1). - _Ralf Stephan_, May 12 2004 Conjecture: a(0)=5, a(1)=11, a(2)=24, a(3)=52, a(n)=3*a(n-1)-2*a(n-2)+a(n-4). - _Harvey P. Dale_, May 19 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000983 | OEIS-A021011

Pisot sequence P(6,11), a(0)=6, a(1)=11, a(n+1) is the nearest integer to a(n)^2/a(n-1).

G.f.: (3x^5+2x^4+x^3+4x^2-x+6)/(-x^6-x^3+x^2-2x+1) (conjectured). - _Ralf Stephan_, May 12 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000984 | OEIS-A022008

Initial member of prime sextuples (p, p+4, p+6, p+10, p+12, p+16).

The prime sextuples starting at a(n) give the highest concentration of primes that can occur on an interval of 17 integers (apart intervals starting at p < 7). It is conjectured that there are infinitely many such sextuples.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Warut Roonguthai_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000985 | OEIS-A022025

Define the sequence S(a(0),a(1)) by a(n+2) is the least integer such that a(n+2)/a(n+1) > a(n+1)/a(n) for n >= 0. This is S(6,102).

The conjectured g.f. (6-5*x^2)/(1-17*x-x^2+14*x^3) yields the same initial terms a(0..271) but from a(272) on a different sequence. - _Bruno Berselli_ and _M. F. Hasler_, Feb 11 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000986 | OEIS-A022027

Define the sequence T(a(0),a(1)) by a(n+2) is the greatest integer such that a(n+2)/a(n+1) < a(n+1)/a(n) for n >= 0. This is T(2,16).

Conjectures: a(n) = 8*a(n-1)-4*a(n-3). G.f.: -(x^2-2) / (4*x^3-8*x+1). - _Colin Barker_, Sep 18 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000987 | OEIS-A022028

Define the sequence T(a(0),a(1)) by a(n+2) is the greatest integer such that a(n+2)/a(n+1) < a(n+1)/a(n) for n >= 0. This is T(2,32).

Conjecture: a(n) = 16*a(n-1)-8*a(n-3). G.f.: -(x^2-2) / (8*x^3-16*x+1). - _Colin Barker_, Sep 18 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000988 | OEIS-A022163

First row of spectral array W(sqrt(3/2)).

Conjectures: a(n) = 6*a(n-2)-3*a(n-4). G.f.: -(3*x^3-5*x-1) / (3*x^4-6*x^2+1). - _Colin Barker_, Oct 23 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000989 | OEIS-A022165

First row of spectral array W(sqrt(5)-1).

Conjectures: a(n) = 6*a(n-2)-4*a(n-4). G.f.: -(x+1)*(4*x^2-4*x-1) / (4*x^4-6*x^2+1). - _Colin Barker_, Oct 23 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000990 | OEIS-A022292

Exactly half of first a(n) terms of Kolakoski sequence A000002 are 1's (not known to be infinite).

Conjecture: a(n) is asymptotic to c*n*log(n) for some constant c <= 1. - _Benoit Cloitre_, Nov 17 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000991 | OEIS-A022344

Allan Wechsler's "J determinant" sequence.

Conjecture: Every pair of Fibonacci sequences, F1 and F2, appear in rows n and m of Wythoff's Array, respectively and have respective characteristics a(n) and a(m). Also, there is

a third Fibonacci sequence F3, defined by F3(i) = F1(i) * F2(j+1) - F1(i+1)*F2(j) where j is held constant. The sequence F3 appears in row p of Wythoff's array and has the characteristic a(p) = a(n)*a(m). - _Kenneth J Ramsey_, Feb 11 2007 A product p of rows n and m of the Wythoff Array, such that a(p) = a(n)*a(m) as described in the conjecture above, is defined by A357097(n, m). - _Peter Munn_, Aug 15 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000992 | OEIS-A022440

a(n) = c(n-1) + c(n-3) where c is the sequence of positive numbers not in a.

From _N. J. A. Sloane_, Nov 24 2004: I'm not sure of the minimal hypotheses needed to generate this sequence, but one method that works is the following:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000993 | OEIS-A022444

Number of self-avoiding closed walks (from (0,0) to (0,0)) of length 2n in strip {-1, 0, 1} X Z.

G.f.: (-12*x^7+16*x^6-36*x^5+37*x^4-24*x^3+14*x^2-4*x+1) / ((1+x^2)^2*(1-2*x)^2) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jacques Labelle (labelle.jacques(AT)uqam.ca) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000994 | OEIS-A022445

Number of self-avoiding closed walks (from 0 to 0) of length 2n in the strip {0, 1, 2} X Z of the square lattice Z X Z.

G.f.: (-6*x^7+10*x^6-18*x^5+27*x^4-14*x^3+10*x^2-4*x+1) / ((1+x^2)^2*(1-2*x)^2) (conjectured). - _Ralf Stephan_, Apr 28 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jacques Labelle (labelle.jacques(AT)uqam.ca) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000995 | OEIS-A022493

Fishburn numbers: number of linearized chord diagrams of degree n; also number of nonisomorphic interval orders on n unlabeled points.

G.f. (conjecture): Q(0), where Q(k) = 1 + (1 - (1-x)^(2*k+1))/(1 - (1-(1-x)^(2*k+2))/(1 -(1-x)^(2*k+2) + 1/Q(k+1))); (continued fraction). - _Sergei N. Gladkovskii_, May 13 2013 Conjectural g.f.s:

猜想/问题提及段 | historical_or_current_mention_unreviewed | Alexander Stoimenow (stoimeno(AT)math.toronto.edu) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000996 | OEIS-A022495

Conjectured number of irreducible multiple zeta values of depth 7 and weight 2n+19.

Conjectured number of irreducible multiple zeta values of depth 7 and weight 2n+19.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Broadhurst_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000997 | OEIS-A022496

Conjectured number of irreducible multiple zeta values of depth 8 and weight 2n+22.

Conjectured number of irreducible multiple zeta values of depth 8 and weight 2n+22.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Broadhurst_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000998 | OEIS-A022497

Conjectured number of irreducible multiple zeta values of depth 9 and weight 2n+25.

Conjectured number of irreducible multiple zeta values of depth 9 and weight 2n+25.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Broadhurst_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-000999 | OEIS-A022498

Conjectured number of irreducible multiple zeta values of depth 10 and weight 2n+28.

Conjectured number of irreducible multiple zeta values of depth 10 and weight 2n+28.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Broadhurst_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001000 | OEIS-A022544

Numbers that are not the sum of 2 squares.

Conjecture: if k is not the sum of 2 squares then sigma(k) == 0 (mod 4) (the converse does not hold, as demonstrated by the entries in A025303). - _Benoit Cloitre_, May 19 2002 Proof of Cloitre's 2002 conjecture: if n is not the sum of 2 squares, then sigma(n) == 0 (mod 4). By the sum-of-two-squares theorem, n is not a sum of two squares iff some prime p == 3 (mod 4) divides n to an odd power e = 2j+1. For such p: sigma(p^e) = Sum_{k=0..e} p^k. Since p==3(mod 4), p^k==1(mod 4) for k even and p^k==3(mod 4) for k odd. As e is odd, e+1 is even, so the terms pair as (p^0+p^1)+(p^2+p^3)+...+(p^(e-1)+p^e), with each pair == 1+3 = 0 (mod 4). Hence sigma(p^e) == 0 (mod 4), and by multiplicativity of sigma, sigma(n) == 0 (mod 4). QED. - _Marc-André Leduc_, Mar 11 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001001 | OEIS-A022552

Numbers that are not the sum of 2 squares and a nonnegative cube.

For n = 1..434, a(n) + 2 is a term of A022551. _Zhi-Wei Sun_ conjectures that Any n can be written as x^2 + y^2 + z^3 + 0(or 2). - _XU Pingya_, Jun 02 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Will Jagy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001002 | OEIS-A022553

Number of binary Lyndon words containing n letters of each type; periodic binary sequences of period 2n with n zeros and n ones in each period.

Conjecturally, number of irreducible alternating Euler sums of depth n and weight 3n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Broadhurst_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001003 | OEIS-A022558

Number of permutations of length n avoiding the pattern 1342.

Conjecture: a(n) is the number of permutations of length n simultaneously avoiding patterns 2143 and 415263. - _Alexander Burstein_, Mar 21 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Miklos Bona_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001004 | OEIS-A022566

Numbers that are not the sum of 4 nonnegative cubes.

Conjecture: The sequence is finite, with 113936676 terms, the largest being 7373170279850 (cf. Deshouillers et al, 2000, Conj. 1 and 2). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001005 | OEIS-A022629

Expansion of Product_{m>=1} (1 + m*q^m).

Conjecture: log(a(n)) ~ sqrt(n/2) * (log(2*n) - 2). - _Vaclav Kotesovec_, May 08 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001006 | OEIS-A022693

Expansion of Product_{m>=1} 1/(1 + m*q^m).

Conjecture: a(n) = Sum_{i_1,i_2,i_3,...}[(-1)^(i_1+i_2+i_3+...)*Product_{n>0} n^i_n], where the sum is over all valid sequences of positive i_k such that i_1+2*i_2+3*i_3+4*i_4+...= n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001007 | OEIS-A022885

Primes p=prime(k) such that prime(k) + prime(k+3) = prime(k+1) + prime(k+2).

These are primes p for which the subsequent alternate prime gaps are equal, so (p(k+3)-p(k+2))/(p(k+1)-p(k)) = 1. It is conjectured that the most frequent alternate prime gaps ratio is one. - _Andres Cicuttin_, Nov 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001008 | OEIS-A022894

Number of solutions to c(1)*prime(1) +...+ c(2n+1)*prime(2n+1) = 0, where c(i) = +-1 for i > 1, c(1) = 1.

Conjecture: limit_{n->oo} a(n)^(1/n) = 4. - _Vaclav Kotesovec_, Jun 05 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001009 | OEIS-A022917

Multinomial coefficient n!/ ([n/4]!, [(n+1)/4]!, [(n+2)/4]!, [(n+3)/4]!).

Conjecture: -(126*n-1)*(n+3)*(n+2)*(n+1)*a(n) +4*(-208*n^4-51*n^3+793*n^2+376*n-6)*a(n-1) +16*(334*n^3-114*n^2-5*n-369)*a(n-2) +64*(n-2)*(334*n^2+264*n-213)*a(n-3) +768*(n-2)*(n-3)*(42*n^2+153*n+113)*a(n-4) +1024*(208*n+177)*(n-2)*(n-3)*(n-4)*a(n-5)=0. - _R. J. Mathar_, Aug 06 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001010 | OEIS-A023037

a(n) = n^0 + n^1 + ... + n^(n-1), or a(n) = (n^n-1)/(n-1) with a(0)=0; a(1)=1.

For prime n, a(n) is conjectured to be the period of Bell numbers (mod n). See A054767. - _T. D. Noe_, Oct 12 2007 For prime n, a(n) is a multiple of the period of Bell numbers mod n (and conjectured to be exactly the period, as mentioned above). - _Charles R Greathouse IV_, Jul 31 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001011 | OEIS-A023048

Smallest prime having least positive primitive root n, or 0 if no such prime exists.

Of course if n is a perfect power then a(n) = 0, but it seems that the other direction is true only assuming the generalized Artin's conjecture. See the link from Tomás Oliveira e Silva below. - _Jianing Song_, Jan 22 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001012 | OEIS-A023054

Simon Plouffe's conjectured extension of sequence A008368.

Simon Plouffe's conjectured extension of sequence A008368.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _J. H. Conway_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001013 | OEIS-A023108

Positive integers which apparently never result in a palindrome under repeated applications of the function A056964(x) = x + (x with digits reversed).

196 is conjectured to be smallest initial term which does not lead to a palindrome. John Walker, Tim Irvin and others have extended this to millions of digits without finding one (see A006960).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001014 | OEIS-A023110

Squares which remain squares when the last digit is removed.

Conjectures from _Colin Barker_, Dec 31 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001015 | OEIS-A023189

Conjecturally, number of infinitely recurring prime patterns of width 2n-1.

Conjecturally, number of infinitely recurring prime patterns of width 2n-1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001016 | OEIS-A023190

Conjecturally, maximum number of primes in an infinitely-recurring prime pattern of width 2*n-1.

Conjecturally, maximum number of primes in an infinitely-recurring prime pattern of width 2*n-1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001017 | OEIS-A023191

Conjecturally, number of maximal infinitely-recurring prime patterns of width 2*n-1.

Conjecturally, number of maximal infinitely-recurring prime patterns of width 2*n-1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001018 | OEIS-A023192

Conjecturally, number of infinitely-recurring prime patterns on n consecutive integers.

Conjecturally, number of infinitely-recurring prime patterns on n consecutive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001019 | OEIS-A023199

a(n) is the least k with sigma(k) >= n*k.

The conjecture a(n)|a(n+1) holds out to n=10. - Devin Kilminster (devin(AT)maths.uwa.edu.au), Mar 10 2003 The conjecture a(n)|a(n+1) fails for n=15. - _T. D. Noe_, Jul 08 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001020 | OEIS-A023200

Primes p such that p + 4 is also prime.

Conjecture: The sequence is infinite and for every n, a(n+1) < a(n)^(1+1/n). Namely a(n)^(1/n) is a strictly decreasing function of n. - _Jahangeer Kholdi_ and _Farideh Firoozbakht_, Nov 24 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001021 | OEIS-A023365

a(n) = 2^(3^(n-1)).

a(n+1) is conjectured to be the reduced denominator of b(n) = b(n-1)*(3/2 - b(n-1)^2); b(0) = 1/2. - _Steven Finch_, Oct 08 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001022 | OEIS-A023393

Maximal number of circles of radius 1 that can be packed in a circle of radius n.

The terms for n>5 are only conjectures supported by extensive computations.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001023 | OEIS-A023408

If any power of 2 ends with k 5's and 6's, they must be the first k terms of this sequence in reverse order.

Conjecture: a(n) + A023402(n) = 9, for n > 1. - _J. Lowell_, Nov 16 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001024 | OEIS-A023503

Greatest prime divisor of prime(n) - 1.

Conjecture: every prime is in the sequence. Cf. A035095 (see my comment). - _Thomas Ordowski_, Aug 06 2017 a(n) is 2 for n in A159611, and is at most 3 for n in A174099. Conjecture: liminf a(n) = 3. - _Jeppe Stig Nielsen_, Jul 04 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001025 | OEIS-A023554

Convolution of natural numbers >= 3 and (Fib(2), Fib(3), Fib(4), ...).

2*(n+5) = A022308(n+4) - a(n+1) (conjectured). Note offset of A022308 is 0. - _Creighton Dement_, Feb 02 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001026 | OEIS-A023634

s(3n)-s(3n-1), where s( ) is sequence A023633.

Conjecture: (a(n)) is fixed point of the morphism

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001027 | OEIS-A023688

Numbers with exactly 6 ones in binary expansion.

Sequence appears to include all numbers m such that 8^5 is the highest power of 2 dividing A005148(m). General conjecture: numbers k such that 8^j is the highest power of 2 dividing A005148(k) is the same sequence as numbers having exactly (j+1) 1's in their binary representation. - _Benoit Cloitre_, Jun 22 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001028 | OEIS-A023811

Largest metadrome (number with digits in strict ascending order) in base n.

lim_{n->infinity} a(n)/a(n-1) - a(n-1)/a(n-2) = exp(1). - Conjectured by _Gerald McGarvey_, Sep 26 2004. Follows from the formula below and lim_{n->infinity} (1+1/n)^n = e. - _Franklin T. Adams-Watters_, Jan 25 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001029 | OEIS-A023861

a(n) = 1*t(n) + 2*t(n-1) + ... + k*t(n+1-k), where k=floor((n+1)/2) and t = A000032 (Lucas numbers).

Conjecture: G.f.: x*(-1-3*x^5+x^4+2*x^3-4*x^2-2*x)/((x^2+x-1)*(x^4+x^2-1)^2). - Maksym Voznyy (voznyy(AT)mail.ru), Jul 27 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001030 | OEIS-A023871

Expansion of Product_{k>=1} (1 - x^k)^(-k^2).

Let A(x) = Product_{k >= 1} (1 - x^k)^(-k^2). The sequence defined by u(n) := [x^n] A(x)^n is conjectured to satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 7 and all positive integers n and r. See A380290. - _Peter Bala_, Feb 02 2025 G.f. (conjectured): 1/Product_{n>=1} E(x^n)^J2(n) where E(x) = Product_{n>=1} 1-x^n and J2(n) = A007434(n) [follows from the identity Sum_{d|n} J2(d) = n^2 - _Peter Bala_, Feb 02 2025]. - _Joerg Arndt_, Jan 25 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001031 | OEIS-A023873

Expansion of Product_{k>=1} (1 - x^k)^(-k^4).

Let A(x) denote the g.f. and let m be an integer. Define a sequence by u(n) = [x^n] A(x)^(m*n). We conjecture that the supercongruence u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) holds for all positive integers n and r and all primes p >= 7. Cf. A380581. - _Peter Bala_, Jan 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001032 | OEIS-A023900

Dirichlet inverse of Euler totient function (A000010).

Conjecture for n>1: Let n = 2^(A007814(n))*m = 2^(ruler(n))*odd_part(n), where m = A000265(n), then a(n) = (-1)^(m=n)*(0+Sum_{i=1..m and gcd(i,m)=1} (4*min(i,m-i)-m)) = (-1)^(m<n)*(1+Sum_{i=1..m and gcd(i,m)>1} (4*min(i,m-i)-m)). - _I. V. Serov_, May 02 2017 The conjecture is true (see Fried link). - _Sela Fried_, Nov 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001033 | OEIS-A024152

a(n) = 12^n - n^12.

Conjecture: satisfies a linear recurrence having signature (25, -234, 1222, -4147, 9867, -17160, 22308, -21879, 16159, -8866, 3510, -949, 157, -12). - _Harvey P. Dale_, Jan 27 2019 The conjecture above is correct. From the general formula for {a(n)} we can see that the roots for the characteristic polynomial are one 12 and thirteen 1's, so the characteristic polynomial is (x - 12)*(x - 1)^13 = x^14 - 25*x^13 + 234*x^12 - ... + 12, with corresponding recurrence coefficients 25, -234, ..., -12. - _Jianing Song_, Jan 28 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001034 | OEIS-A024162

Number of incongruent integer-sided triangles with sides a,b,c, a<b<c, a+b+c=n such that gcd(a,b,c)=1.

Conjecture: a(2*k-1) >= a(2*k). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001035 | OEIS-A024187

n-th elementary symmetric function of 3,4,...,n+3.

Conjecture: essentially the same as A001711. - _Ralf Stephan_, Dec 30 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001036 | OEIS-A024195

Integer part of (4th elementary symmetric function of S(n))/(3rd elementary symmetric of S(n)), where S(n) = {3,4, ..., n+5}.

Conjectured g.f.: (-1-x^14+3*x^13-4*x^12+3*x^11-x^10+x^7-x^6-x^2+x)/((x^2+1)*(x^4+1)*(x-1)^3). - Maksym Voznyy (voznyy(AT)mail.ru), Jul 27 2009, confirmed by _Robert Israel_, May 06 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001037 | OEIS-A024199

a(n) = (2n-1)!! * Sum_{k=0..n-1}(-1)^k/(2k+1).

Conjecture: a(n) = 1/2*Sum_{k = 0..2*n-1} (-1)^(n-k+1)*k!*(2*n - 2*k - 3)!!, where the double factorial of an odd integer (positive or negative) may be defined in terms of the gamma function as (2*N - 1)!! = 2^((N+1)/2)*Gamma(N/2 + 1)/sqrt(Pi).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001038 | OEIS-A024206

Expansion of x^2*(1+x-x^2)/((1-x^2)*(1-x)^2).

Conjecture: if a(n) = p#(primorial)-1 for some prime number p, then q=(n+1) is also a prime number where p#=floor(q^2/4). Tested up to n=10^100000 no counterexamples are found. It seems that the subsequence is very scattered. So far the triples (p,q,a(q-1)) are {(2,3,1), (3,5,5), (5,11,29), (7,29,209), (17,1429,510509)}. - _David Morales Marciel_, Oct 02 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001039 | OEIS-A024219

a(n) = floor((2nd elementary symmetric function of S(n))/(first elementary symmetric function of S(n))), where S(n) = {first n+1 positive integers congruent to 1 mod 3}.

Conjecture: a(n) = 3*a(n-1) - 4*a(n-2) + 4*a(n-3) - 3*a(n-4) + a(n-5); The above conjectures are true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001040 | OEIS-A024220

a(n) = [(3rd elementary symmetric function of S(n))/(first elementary symmetric function of S(n))], where S(n) = {first n+2 positive integers congruent to 1 mod 3}.

Conjecture: a(n) = 4*a(n-1) - 6*a(n-2) + 5*a(n-3) - 5*a(n-4) + 6*a(n-5) - 4*a(n-6) + a(n-7). G.f.: x*(-2-11*x-7*x^2-8*x^3+x^4) / ((1+x+x^2)*(x-1)^5). - _R. J. Mathar_, Oct 08 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001041 | OEIS-A024283

E.g.f. (1/2) * tan(x)^2 (even powers only).

Conjectural o.g.f.: x/(1 + x - 9*x/(1 - 8*x/(1 + x - 25*x/(1 - 24*x/(1 + x - ... - (2*n+1)^2*x/(1 - 4*n*(n+1)*x/(1 + x - ...))))))). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_. This sequence was in the 1973 "Handbook", but was then omitted from the database. Resubmitted by _R. H. Hardin_. Entry revised by _N. J. A. Sloane_, Jun 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001042 | OEIS-A024337

Expansion of sinh(log(1+x))*log(1+x)/2.

Conjecture: a(n) = Sum_{k=0..n/2} Stirling1(n,2*k)*k. - _Benedict W. J. Irwin_, Oct 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001043 | OEIS-A024385

a(n) = [(2nd elementary symmetric function of S(n))/(first elementary symmetric function of S(n))], where S(n) = {first n+1 positive integers congruent to 1 mod 4}.

Conjecture: a(n) = 2*a(n-1) - a(n-2) + a(n-6) - 2*a(n-7) + a(n-8). G.f. x^2*(-3-3*x-x^2-2*x^3-2*x^4-2*x^5+x^6) / ((1+x)*(1+x+x^2)*(x^2-x+1)*(x-1)^3). - _R. J. Mathar_, Oct 08 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001044 | OEIS-A024399

a(n) = [(3rd elementary symmetric function of S(n))/(first elementary symmetric function of S(n))], where S(n) = {first n+2 positive integers congruent to 2 mod 3}.

Conjecture: a(n)= +4*a(n-1) -6*a(n-2) +5*a(n-3) -5*a(n-4) +6*a(n-5) -4*a(n-6) +a(n-7). G.f. x*(-5-11*x-7*x^2-5*x^3+x^4) / ((1+x+x^2)*(x-1)^5). - _R. J. Mathar_, Oct 08 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001045 | OEIS-A024401

a(n) = [(3rd elementary symmetric function of S(n))/(2nd elementary symmetric function of S(n))], where S(n) = {first n+2 positive integers congruent to 2 mod 3}.

Conjecture: a(n) = 2*a(n-1) - a(n-2) + a(n-3) - 2*a(n-4) + a(n-5). G.f. x*(-1-x-x^3-x^2+x^4) / ((1+x+x^2)*(x-1)^3). - _R. J. Mathar_, Oct 08 2011 A024392(n)/A024391(n+1) = (3*n^2 + 5*n - 2)/6 + 8*(2*n + 5)/(3*(3*n + 5)*(3*n + 8)). The fractional part of (3*n^2 + 5*n - 2)/6 is 2/3, 0, or 1/3 according as n is congruent to 0, 1, or 2 modulo 3. Since 0 < 8*(2*n + 5)/(3*(3*n + 5)*(3*n + 8)) < 1/3 for n >= 1, the formula follows. In particular, this proves the recurrence and generating function conjectured above. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001046 | OEIS-A024417

s(n,a(n)) = max{s(n,k): k=1,2,...,n}, n >= 1, where s(n,k) = Stirling numbers of the second kind.

a(n) ~ n/LambertW(n) - 1 (conjecture). - _Mats Granvik_, Oct 16 2013 Granvik's conjecture follows from W(n) = log(n) - log(log(n)) + o(1). (See also saddle-point-based asymptotic (for rows' centres of mass rather than maximums) in A000110, which uses W directly.) - _Natalia L. Skirrow_, Nov 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001047 | OEIS-A024451

a(n) is the numerator of Sum_{i = 1..n} 1/prime(i).

Conjecture: All terms are squarefree numbers. - _Nicolas Bělohoubek_, Apr 13 2022 The above conjecture would imply that for n > 0, gcd(a(n), A369651(n)) = 1. See corollary 2 on the page 4 of Ufnarovski-Åhlander paper. - _Antti Karttunen_, Jan 31 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001048 | OEIS-A024459

a(n) = s(1)*t(n) + s(2)*t(n-1) + ... + s(k)*t(n+1-k), where k = [(n+1)/2], s = (Fibonacci numbers), t = (Lucas numbers).

Conjectured g.f.: x*(-1-2*x-x^2+3*x^3) / ((x^2+1)*(x^2-x-1)*(x^2+x-1)^2). - _R. J. Mathar_, Apr 27 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001049 | OEIS-A024719

a(n) = (1/3)*(2 + Sum_{k=0..n} C(3k,k)).

Conjecture: 2*n*(2*n-1)*a(n) + (-31*n^2 + 29*n - 6)*a(n-1) +3*(3*n-1)*(3*n-2)*a(n-2) = 0. - _R. J. Mathar_, Sep 29 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001050 | OEIS-A024812

Numbers n for which there is exactly one positive integer m such that n = floor(cot(Pi/(2m))).

Conjecture (verified for k <= 10^6 by _M. F. Hasler_): A024812(k) = (A024813(k)+k-1)/2, k=1,2,.... - _L. Edson Jeffery_, Mar 21 2013 Conjecture: a(n) = a(n-1) + a(n-3) - a(n-4); g.f.: x*(x^15-x^14+3*x^2+2*x+2) / ((x-1)^2*(x^2+x+1)). - _Colin Barker_, Jan 03 2014 Barker's 2014 recurrence conjecture fails at n=15: a(14)+a(12)-a(11) = 32+28-25 = 35, but a(15) = 34. - _Michal Buran_, Mar 24 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001051 | OEIS-A024813

Positive integers m for which f(m-1) < f(m) < f(m+1), where f(m) = floor(cot(Pi/(2m))).

Conjecture (verified for m < 10^6 by _M. F. Hasler_): a(n) = 2*A024812(n) - n + 1, n=1,2,.... - _L. Edson Jeffery_, Mar 21 2013 The above conjecture follows from the Laurent series for cot(x) = 1/x - x/3 + O(x^3) and the conjecture n/a(n) ~ 4/Pi-1. - _M. F. Hasler_, Mar 25 2013 Proof of both L. Edson Jeffery's conjecture above and M. F. Hasler's asymptotic conjecture below: Let g(x) = cot(Pi/(2*x)). For x >= 2, g'(x) = (2/Pi)*((Pi/(2*x))/sin(Pi/(2*x)))^2 is strictly decreasing with 2/Pi < g'(x) <= g'(2) = Pi/4. By the Mean Value Theorem, for all m >= 2, g(m+1) - g(m) <= Pi/4 < 1 and g(m+2) - g(m) > 4/Pi > 1. Hence f(m) = floor(g(m)) takes every positive integer value either once or twice. Since f(1) = 0, among the A024812(n) - 1 positive integers below A024812(n), exactly n - 1 are taken once and A024812(n) - n are taken twice, giving a(n) = 1 + (n - 1) + 2*(A024812(n) - n) + 1 = 2*A024812(n) - n + 1 for all n >= 1. Since A024812(n) = floor(cot(Pi/(2*a(n)))) = (2/Pi)*a(n) + O(1), substituting into a(n) = 2*A024812(n) - n + 1 immediately yields lim_{n->oo} n/a(n) = 4/Pi - 1. - _Hriday Rungta_, Oct 04 2026 n/a(n) ~ 4/Pi - 1 (as n -> oo), or a(n) ~ 3.65979 n. (Conjectured.) - _M. F. Hasler_, Mar 25 2013 Conjecture: a(n) = a(n-1) + a(n-3) - a(n-4); g.f.: x*(2*x^15-2*x^14-x^3+5*x^2+3*x+4) / ((x-1)^2*(x^2+x+1)). - _Colin Barker_, Jan 03 2014 Barker's 2014 recurrence conjecture is false: a(14)+a(12)-a(11) = 51+45-40 = 56, but a(15) = 54. - _Michal Buran_, Mar 24 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001052 | OEIS-A024849

a(n) = least m such that if r and s in {|F(h+1)-tau*F(h)|: h = 1,2,...,n} satisfy r < s, then r < k/m < s for some integer k, where F = A000045 (Fibonacci numbers) and tau = (1+sqrt(5))/2 (golden ratio).

Conjectures from _Chai Wah Wu_, May 15 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001053 | OEIS-A024998

a(n) = number of (s(0), s(1), ..., s(n)) such that every s(i) is an integer, s(0) = 0, |s(i) - s(i-1)| = 1 for i = 1,2; |s(i) - s(i-1)| <= 1 for i >= 3, s(n) = 1. Also a(n) = T(n,n-1), where T is the array defined in A024996.

Conjecture: (n+1)*a(n) +(-3*n+2)*a(n-1) +(-n-7)*a(n-2) +3*(n-4)*a(n-3)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001054 | OEIS-A025012

Central heptanomial coefficients: largest coefficient of (1+x+...+x^6)^n.

I checked that Mathar's conjectured formula fits the 200 terms of the b-file. Note that the coefficients are powers of 7: 49=7^2, 343=7^3, 2401=7^4, 16807=7^5, 117649=7^6. - _T. D. Noe_, Nov 15 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001055 | OEIS-A025060

Numbers of the form i*j + j*k + k*i, where 1 <= i < j < k.

Conjecture: If i, j and k are allowed to be negative, but not zero, and are still distinct, then the sequence is all the integers. - _Jon Perry_, Apr 21 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001056 | OEIS-A025064

Position of numbers of form 3*n^2 in A025060 (numbers of form j*k + k*i + i*j, where 1 <=i < j < k).

It is conjectured that A000926 ends at 1848, in which case a(n) = 3*n^2+18*n-38 for all n >= 22. - _Robert Israel_, Sep 06 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001057 | OEIS-A025093

a(n) = s(1)t(n) + s(2)t(n-1) + ... + s(k)t(n-k+1), where k = [n/2], s = (Lucas numbers), t = (odd natural numbers).

Conjectures from _Colin Barker_, Aug 07 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001058 | OEIS-A025142

Fixed point of the square runlength transform, with a(1) = 1 (the runlength transform of a binary sequence is the sequence of the lengths of its runs).

Conjecture: as for the Kolakoski sequence A000002, which is a fixed point of RL, it remains to be shown that the limiting frequency of 1's and 2's in this sequence is 0.5. - _Jean-Christophe Hervé_, Oct 21 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001059 | OEIS-A025163

The value of the associated Legendre polynomial of index n and order 1 evaluated at x=2^(-1/2) multiplied by 2^(3*n/2-1).

Conjecture: (-n+1)*a(n) +2*(2*n-1)*a(n-1) -8*n*a(n-2)=0. - _R. J. Mathar_, Feb 05 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001060 | OEIS-A025165

a(n) = H_n(1) / 2^floor(n/2) where H_n is the n-th Hermite polynomial.

Conjecture: a(n) +a(n-1) +(2*n-5)*a(n-2) +(2*n-7)*a(n-3) +(n-2)*(n-3)*a(n-4) +(n-3)*(n-4)*a(n-5)=0. - _R. J. Mathar_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001061 | OEIS-A025166

Expansion of e.g.f. -exp(-x/(1-2*x))/(1-2*x).

Conjecture: a(n) + (-4*n+3)*a(n-1) + 4*(n-1)^2*a(n-2) = 0. - _R. J. Mathar_, Feb 05 2013 Conjecture verified using the differential equation (4*x - 1)*y(x) + (4*x^2 - 4*x + 1)*y'(x) = 0 satisfied by the e.g.f. - _Robert Israel_, May 17 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001062 | OEIS-A025175

Jacobi polynomial P((1, 1), n, (1/2)).

Conjecture: (n+2)*a(n) + 6*(-n-1)*a(n-1) + 12*(2*n+1)*a(n-2) + 32*(-n+1)*a(n-3) = 0. - _R. J. Mathar_, Mar 03 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001063 | OEIS-A025180

a(n) = number of (s(0), s(1), ..., s(n)) such that s(i) is an integer, s(0) = 0, |s(1)| = 1, |s(i) - s(i-1)| <= 1 for i >= 2, s(n) = 2. Also a(n) = T(n,n-2), where T is the array defined in A025177.

Conjecture: -(n+2)*(n-3)*a(n) +2*(2*n+1)*(n-3)*a(n-1) +(-n^2+10*n-18)*a(n-2) -3*(2*n-3)*(n-3)*a(n-3)=0. - _R. J. Mathar_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001064 | OEIS-A025181

a(n) = number of (s(0), s(1), ..., s(n)) such that s(i) is an integer, s(0) = 0, |s(1)| = 1, |s(i) - s(i-1)| <= 1 for i >= 2, s(n) = 3. Also a(n) = T(n,n-3), where T is the array defined in A025177.

Conjecture: +(n+3)*a(n) +(-5*n-7)*a(n-1) +(3*n-7)*a(n-2) +(11*n-7)*a(n-3) +4*(-n+6)*a(n-4) +6*(-n+5)*a(n-5)=0. - _R. J. Mathar_, Feb 25 2015 Conjecture: -(n+3)*(n-3)*(4*n^2-12*n+17)*a(n) +(n-1)*(8*n^3-20*n^2+30*n-81)*a(n-1) +3*(n-1)*(n-2)*(4*n^2-4*n+9)*a(n-2)=0. - _R. J. Mathar_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001065 | OEIS-A025182

a(n) = number of (s(0), s(1), ..., s(n)) such that s(i) is an integer, s(0) = 0, |s(1)| = 1, |s(i) - s(i-1)| <= 1 for i >= 2, s(n) = 4. Also a(n) = T(n,n-4), where T is the array defined in A025177.

Conjecture: -(n-4)*(n+4)*a(n) +(4*n^2-7*n-29)*a(n-1) +(-2*n^2+17*n-2)*a(n-2) -(4*n+1)*(n-3)*a(n-3) +3*(n-3)*(n-4)*a(n-4)=0. - _R. J. Mathar_, Feb 25 2015 Conjecture: -(n-4)*(n+4)*(n^2-3*n+6)*a(n) +(n-1)*(2*n^3-5*n^2+11*n-36)*a(n-1) +3*(n-1)*(n-2)*(n^2-n+4)*a(n-2)=0. - _R. J. Mathar_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001066 | OEIS-A025183

a(n) = T(2n-1,n), where T is the array defined in A025177.

Conjecture: 9*n*(10833*n-26779)*(3*n-4)*(3*n-2)*a(n) -3*(n-1)*(749757*n^3+5096

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001067 | OEIS-A025184

a(n) = T(2n,n), where T is the array defined in A025177.

Conjecture: 33*n*(3*n-1)*(3*n-2)*a(n) +11*(2047*n^3-10725*n^2+17192*n-8520)*a(n-1) +9*(-4397*n^3-10169*n^2+110500*n-145368)*a(n-2) -54*(2*n-5)*(5353*n^2-33313*n+53904)*a(n-3) -115668*(2*n-5)*(2*n-7)*(n-4)*a(n-4)=0. - _R. J. Mathar_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001068 | OEIS-A025187

a(n) = T(2n,n-1), where T is the array defined in A025177.

Conjecture: +12*n*(1573*n-2085)*(3*n-1)*(3*n+1)*a(n) +(259519*n^4-6550685*n^3+19840883*n^2-20466403*n+6720078)*a(n-1) -54*(2*n-3)*(63141*n^3-343882*n^2+645379*n-388606)*a(n-2) -324*(n-3)*(4693*n-5417)*(2*n-3)*(2*n-5)*a(n-3)=0. - _R. J. Mathar_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001069 | OEIS-A025191

a(n) = Sum_{k=0..n} T(n,k), where T is the array defined in A025177.

Conjectures: a(n) = A027914(n)-A027914(n-1) = (A081673(n)-A081673(n-1))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001070 | OEIS-A025244

a(n) = a(1)*a(n-1) + a(2)*a(n-2) + ... + a(n-3)*a(3) for n >= 4, with initial terms 1, 1, 2, 2.

Conjecture: n*a(n) +(-2*n+3)*a(n-1) +(-n+3)*a(n-2) +3*(-2*n+9)*a(n-3) +(n-6)*a(n-4)=0. - _R. J. Mathar_, Feb 25 2015 Conjecture verified (for n >= 6) using differential equation 2*x^4 - 3*x^2 - x - 1 + (-2*x^3 + 9*x^2 + x + 1)*g(x) + (x^4 - 6*x^3 - x^2 - 2*x + 1)*g'(x) = 0 satisfied by g.f. - _Robert Israel_, Sep 28 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001071 | OEIS-A025245

a(n) = (1/2)*s(n+3), where s = A025244.

Conjecture: (n+3)*a(n) +(-2*n-3)*a(n-1) -n*a(n-2) +3*(-2*n+3)*a(n-3) +(n-3)*a(n-4)=0. - _R. J. Mathar_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001072 | OEIS-A025248

a(n) = a(1)*a(n-1) + a(2)*a(n-2) + ... + a(n-3)*a(3) for n >= 4, with initial terms 1, 0, 2, 2.

Conjecture: n*a(n) +(-2*n+3)*a(n-1) +(n-3)*a(n-2) +4*(-2*n+9)*a(n-3)=0. - _R. J. Mathar_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001073 | OEIS-A025249

a(n) = (1/2)*s(n+3), where s = A025248.

Conjecture: +(n+3)*a(n) +(-2*n-3)*a(n-1) +n*a(n-2) +4*(-2*n+3)*a(n-3)=0. - _R. J. Mathar_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001074 | OEIS-A025251

a(n) = a(1)*a(n-1) + a(2)*a(n-2) + ... + a(n-3)*a(3) for n >= 4, with initial terms 0, 1, 2, 0.

Conjecture: n*(n^2+n+1)*a(n) +(n-1)*(n^2+2*n+3)*a(n-1) -2*(n-3)*(n^2+n+1)*a(n-2) +2*(-5*n^3+16*n^2+19*n+30)*a(n-3) +(-7*n^3+23*n^2+59*n+126)*a(n-4) +(n-7)*(n^2+2*n+3)*a(n-5)=0. - _R. J. Mathar_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001075 | OEIS-A025255

a(n) = a(1)*a(n-1) + a(2)*a(n-2) + ... + a(n-3)*a(3) for n >= 4, with initial terms 1, 3, 1, 1.

Conjecture: n*a(n) +(-2*n+3)*a(n-1) +5*(-n+3)*a(n-2) +(2*n-9)*a(n-3) +9*(n-6)*a(n-4)=0. - _R. J. Mathar_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001076 | OEIS-A025256

a(n) = a(1)*a(n-1) + a(2)*a(n-2) + ... + a(n-3)*a(3) for n >= 4, with initial terms 1, 1, 3, 3.

Conjecture: n*a(n) +(-2*n+3)*a(n-1) +(-n+3)*a(n-2) +5*(-2*n+9)*a(n-3) +(n-6)*a(n-4)=0. - _R. J. Mathar_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001077 | OEIS-A025257

(1/3)*s(n+3), where s = A025256.

Conjecture: (n+3)*a(n) +(-2*n-3)*a(n-1) -n*a(n-2) +5*(-2*n+3)*a(n-3) +(n-3)*a(n-4)=0. - _R. J. Mathar_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001078 | OEIS-A025258

a(n) = a(1)*a(n-1) + a(2)*a(n-2) + ... + a(n-3)*a(3) for n >= 4, with initial terms 1, -1, 1, 1.

Conjecture: -n*a(n) +(2*n-3)*a(n-1) +3*(-n+3)*a(n-2) +3*(2*n-9)*a(n-3) +(-n+6)*a(n-4)=0. - _R. J. Mathar_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001079 | OEIS-A025259

a(n) = a(1)*a(n-1) + a(2)*a(n-2) + ... + a(n-3)*a(3) for n >= 4, with initial terms 2, -1, 1, 2.

Conjecture: n*a(n) +2*(3-2*n)*a(n-1) +6*(n-3)*a(n-2) +4*(9-2*n)*a(n-3) +(n-6)*a(n-4)=0. - _R. J. Mathar_, Sep 29 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001080 | OEIS-A025260

a(n) = a(1)*a(n-1) + a(2)*a(n-2) + ... + a(n-3)*a(3) for n >= 4, with initial terms 2, -2, 1, 2.

Conjecture: n*a(n) +2*(-2*n+3)*a(n-1) +8*(n-3)*a(n-2) +6*(-2*n+9)*a(n-3) +4*(n-6)*a(n-4)=0. - _R. J. Mathar_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001081 | OEIS-A025261

a(n) = a(1)*a(n-1) + a(2)*a(n-2) + ... + a(n-3)*a(3) for n >= 4, with initial terms 3, -2, 1, 3.

Conjecture: n*a(n) +3*(-2*n+3)*a(n-1) +13*(n-3)*a(n-2) +8*(-2*n+9)*a(n-3) +4*(n-6)*a(n-4)=0. - _R. J. Mathar_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001082 | OEIS-A025268

a(n) = a(1)*a(n-1) + a(2)*a(n-2) + ...+ a(n-1)*a(1) for n >= 5, with initial values 1,1,1,1.

Conjecture: n*a(n) +(n+1)*a(n-1) +10*(-2*n+5)*a(n-2) +2*(2*n-9)*a(n-3) +2*(14*n-79)*a(n-4) +40*(n-7)*a(n-5)=0. - _R. J. Mathar_, Jan 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001083 | OEIS-A025269

a(n) = a(1)*a(n-1) + a(2)*a(n-2) + ...+ a(n-1)*a(1) for n >= 5, with initial values 2,1,1,1.

Conjecture: n*a(n) +4*(-2*n+3)*a(n-1) +12*(n-3)*a(n-2) +6*(2*n-9)*a(n-3) +16*(n-6)*a(n-4)=0. - _R. J. Mathar_, Nov 21 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001084 | OEIS-A025270

a(n) = a(1)*a(n-1) + a(2)*a(n-2) + ...+ a(n-1)*a(1) for n >= 5, with initial values 1,2,1,1.

Conjecture: n*a(n) +2*(-2*n+3)*a(n-1) +4*(-n+3)*a(n-2) +6*(2*n-9)*a(n-3) +20*(n-6)*a(n-4)=0. - _R. J. Mathar_, Nov 21 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001085 | OEIS-A025272

a(n) = a(1)*a(n-1) + a(2)*a(n-2) + ...+ a(n-1)*a(1) for n >= 5, with initial values 1,1,1,2.

Conjecture: n*a(n) +(n+1)*a(n-1) +10*(-2*n+5)*a(n-2) +2*(2*n-9)*a(n-3) +2*(12*n-67)*a(n-4) +20*(n-7)*a(n-5)=0. - _R. J. Mathar_, Nov 21 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001086 | OEIS-A025273

a(n) = a(1)*a(n-1) + a(2)*a(n-2) + ...+ a(n-1)*a(1) for n >= 5, starting 1,0,1,1.

Conjecture: (with offset 0 instead of 1) (n+1)*a(n) +2*(-2*n+1)*a(n-1) +4*(n-2)*a(n-2) +2*(-2*n+7)*a(n-3) +4*(n-5)*a(n-4)=0. - _R. J. Mathar_, Nov 24 2012 Conjecture follows from the differential equation 4*x^3-3*x^2+2*x-1+(-4*x^4+2*x^3-2*x+1)*g(x)+(4*x^5-4*x^4+4*x^3-4*x^2+x)*g'(x)=0 satisfied by the g.f. - _Robert Israel_, Nov 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001087 | OEIS-A025275

a(n) = a(1)*a(n-1) + a(2)*a(n-2) + ...+ a(n-1)*a(1) for n >= 5, with initial values 1,0,1,0.

Conjecture: n*a(n) +2*(-2*n+3)*a(n-1) +4*(n-3)*a(n-2) +2*(-2*n+9)*a(n-3) +8*(n-6)*a(n-4)=0. - _R. J. Mathar_, May 30 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001088 | OEIS-A025358

Numbers that are the sum of 4 nonzero squares in exactly 2 ways.

Conjecture: the even members of this sequence are all numbers of the form

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001089 | OEIS-A025416

Least sum of 4 nonzero squares in exactly n ways.

Conjecture: The sequence never becomes monotonic increasing. - _Jon Perry_, Nov 03 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001090 | OEIS-A025426

Number of partitions of n into 2 nonzero squares.

Conjecture: Sum_{k=1..n} a(k) ~ n*Pi/8. - _Vaclav Kotesovec_, Dec 28 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001091 | OEIS-A025482

a(0) = 0; a(n) = a(n/3)/3 if n = a(n/3) = 0 (mod 3); a(n) = a(n-1)+1 otherwise.

Bounded, conjectured maximum is a(257) = 17.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001092 | OEIS-A025567

a(n) = T(n,n+1), where T is the array defined in A025564.

Conjecture: (n+3)*a(n) +4*(-n-2)*a(n-1) +2*a(n-2) +8*(n-1)*a(n-3) +3*(n-3)*a(n-4)=0. - _R. J. Mathar_, Apr 03 2015 Conjecture: (n-1)*(n-2)*(n+3)*a(n) -2*n*(n-2)*(n+2)*a(n-1) -3*n*(n-1)^2*a(n-2)=0. - _R. J. Mathar_, Apr 03 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001093 | OEIS-A025577

Expansion of (x/(1-x))*sqrt((1+x)/(1-3*x)).

Conjecture: (-n+1)*a(n) +3*(n-1)*a(n-1) +(n-7)*a(n-2) +3*(-n+3)*a(n-3)=0. - _R. J. Mathar_, Jul 02 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001094 | OEIS-A025591

Maximal coefficient of Product_{k<=n} (1 + x^k). Number of solutions to +- 1 +- 2 +- 3 +- ... +- n = 0 or 1.

a(n) ~ sqrt(6/Pi) * 2^n / n^(3/2) [conjectured by Andrica and Tomescu (2002) and proved by Sullivan (2013)]. - _Vaclav Kotesovec_, Mar 17 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001095 | OEIS-A025754

9th-order Patalan numbers (generalization of Catalan numbers).

Conjecture: n*a(n) + 9*(-9*n+10)*a(n-1) = 0. - _R. J. Mathar_, Jul 28 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001096 | OEIS-A025755

10th-order Patalan numbers (generalization of Catalan numbers).

Conjecture: n*a(n) + 10*(-10*n+11)*a(n-1) = 0. - _R. J. Mathar_, Jul 28 2014 a(n) = 100^(n-1)*Pochhammer(9/10, n-1)/n! for n >= 1. Maple confirms this satisfies Mathar's conjecture for n >= 2 (it's not true for n=1). - _Robert Israel_, Oct 05 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001097 | OEIS-A025756

3rd-order Vatalan numbers (generalization of Catalan numbers).

Conjecture: n*(n-1)*a(n) -(n-1)*(19*n-36)*a(n-1) +9*(11*n^2-51*n+60)*a(n-2) -9*(3*n-7)*(3*n-8)*a(n-3) = 0. - _R. J. Mathar_, Nov 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001098 | OEIS-A025757

4th-order Vatalan numbers (generalization of Catalan numbers).

Conjecture: 5*n*(n-1)*(n-2)*a(n) -(239*n-600)*(n-1)*(n-2)*a(n-1) +24*(n-2)*(158*n^2-953*n+1445)*a(n-2) +16*(-1232*n^3+13056*n^2-45949*n+53730)*a(n-3) -128*(4*n-15)*(2*n-7)*(4*n-13)*a(n-4)=0. - _R. J. Mathar_, Jul 28 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001099 | OEIS-A025758

5th-order Vatalan numbers (generalization of Catalan numbers).

Conjecture: 41*n*(n-1)*(n-2)*(n-3)*a(n) -3*(1367*n-4100)*(n-1)*(n-2)*(n-3)*a(n-1) +50*(n-2)*(n-3)*(3077*n^2-21533*n+38136)*a(n-2) -250*(n-3)*(10265*n^3-123135*n^2+494446*n-664572)*a(n-3) +1875*(8575*n^4-154250*n^3+1039765*n^2-3112730*n+3491808)*a(n-4) -625*(5*n-22)*(5*n-21)*(5*n-24)*(5*n-23)*a(n-5)=0. - _R. J. Mathar_, Jul 28 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001100 | OEIS-A026002

a(n) = T(n,n+2), where T = Delannoy triangle (A008288).

Conjecture: define an infinite array to have m(n,1) = m(1,n) = n*(n-1)+1 in the first row and column, and m(i,j) = m(i,j-1) + m(i-1,j-1) + m(i-1,j); then m(n,n) = a(n). - _J. M. Bergot_, Apr 24 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001101 | OEIS-A026004

a(n) = T(3n+1,n), where T = Catalan triangle (A008315).

G.f.: ((sqrt(x)*sin(2/3*arcsin((3*sqrt(3)*sqrt(x))/2)))/sqrt(4/3-9*x)-cos(1/3*arccos(1-(27*x)/2))+1)/(3*x). - conjectured by _Harvey P. Dale_, Jun 30 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001102 | OEIS-A026005

a(n) = T(4*n,n), where T = Catalan triangle (A008315).

Conjecture: 9*n*(3*n+2)*(3*n+1)*a(n) +12*(-55*n^3-59*n^2+65*n-11)*a(n-1) -32*(4*n-5)*(4*n-3)*(2*n-3)*a(n-2)=0. - _R. J. Mathar_, May 22 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001103 | OEIS-A026008

a(n) = T(n, floor(n/2)), where T = Catalan triangle (A008315).

Conjecture: (n+5)*a(n) + (n+3)*a(n-1) + (-5*n-9)*a(n-2) - 4*n*a(n-3) +4*(n-2)*a(n-4) = 0. - _R. J. Mathar_, Jun 10 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001104 | OEIS-A026010

a(n) = number of (s(0), s(1), ..., s(n)) such that s(i) is a nonnegative integer and |s(i) - s(i-1)| = 1 for i = 1,2,...,n and s(0) = 2. Also a(n) = sum of numbers in row n+1 of array T defined in A026009.

Conjecture: a(n) is the number of integer compositions of n + 2 in which the even parts appear as often at even positions as at odd positions (confirmed up to n = 19). - _Gus Wiseman_, Mar 17 2018 Conjecture: (n+3)*a(n) - 2*a(n-1) + (-5*n-3)*a(n-2) + 2*a(n-3) + 4*(n-3)*a(n-4) = 0. - _R. J. Mathar_, Jun 20 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001105 | OEIS-A026013

a(n) = number of (s(0), s(1), ..., s(2n)) such that s(i) is a nonnegative integer and |s(i) - s(i-1)| = 1 for i = 1,2,...,n, s(0) = 2, s(2n) = 4. Also a(n) = T(2n,n-1), where T is the array defined in A026009.

Conjecture: (n+4)*(n+1)*a(n) -4*(n^2+4*n+6)*a(n-1) +4*(2*n-3)*a(n-2)=0. - _R. J. Mathar_, Nov 13 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001106 | OEIS-A026017

a(n) = number of (s(0), s(1), ..., s(2*n-1)) such that s(i) is a nonnegative integer and |s(i) - s(i-1)| = 1 for i = 1,2,...,n, s(0) = 2, s(2*n-1) = 5. Also a(n) = T(2*n-1,n-2), where T is the array defined in A026009.

Conjecture: (n+4)*a(n) + (-8*n-17)*a(n-1) + (19*n+1)*a(n-2) + 6*(-2*n+5)*a(n-3) = 0. - _R. J. Mathar_, Jun 20 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001107 | OEIS-A026018

a(n) = number of (s(0), s(1), ..., s(2*n-1)) such that s(i) is a nonnegative integer and |s(i) - s(i-1)| = 1 for i = 1,2,...,n, s(0) = 2, s(2*n-1) = 7. Also a(n) = T(2*n-1,n-3), where T is the array defined in A026009.

Conjecture: -(n+5)*(3*n-37)*a(n) + 3*(-n^2-84*n-173)*a(n-1) + 2*(32*n^2+295*n+254)*a(n-2) - 8*(n+25)*(2*n-5)*a(n-3) = 0. - _R. J. Mathar_, Jun 20 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001108 | OEIS-A026019

a(n) = binomial(3*n,n) - binomial(3*n,n-3).

Conjecture: -2*(2*n+3)*(13*n-9)*(n+1)*a(n) + (499*n^3-7*n^2-120*n-54)*a(n-1) - 3*(3*n-5)*(37*n-24)*(3*n-4)*a(n-2) = 0. - _R. J. Mathar_, Jun 20 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001109 | OEIS-A026021

a(n) = T(n, [n/2]), where T is the array defined in A026009.

Conjecture: -(n+7)*(5*n-34)*a(n) + 8*(-n-14)*a(n-1) + (27*n^2-83*n-428)*a(n-2) + 8*(4*n+5)*a(n-3) 44*(n-3)*(7*n-27)*a(n-4) = 0. - _R. J. Mathar_, Jun 20 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001110 | OEIS-A026023

a(n) = number of (s(0), s(1), ..., s(n)) such that s(i) is a nonnegative integer and |s(i) - s(i-1)| = 1 for i = 1,2,...,n and s(0) = 3. Also a(n) = Sum{T(n,k), k = 0,1,...,[(n+3)/2]}, where T is defined in A026022.

Conjecture: (n+4)*(n-1)*a(n) +(n-1)*(n+1)*a(n-1) -2*(n+1)*(2*n+1)*a(n-2) -4*(n-1)*(n+1)*a(n-3)=0. - _R. J. Mathar_, Sep 29 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001111 | OEIS-A026026

a(n) = number of (s(0), s(1), ..., s(2*n-1)) such that s(i) is a nonnegative integer and |s(i) - s(i-1)| = 1 for i = 1,2,...,n, s(0) = 3, s(2*n-1) = 4. Also a(n) = T(2*n-1,n-1), where T is defined in A026022.

Conjecture: -(n+3)*(11*n-38)*a(n) +2*(35*n^2-86*n-102)*a(n-1) -4*(13*n-30)*(2*n-5)*a(n-2)=0. - _R. J. Mathar_, Jun 22 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001112 | OEIS-A026027

a(n) = number of (s(0), s(1), ..., s(2*n-1)) such that s(i) is a nonnegative integer and |s(i) - s(i-1)| = 1 for i = 1,2,...,n, s(0) = 3, s(2*n-1) = 6. Also a(n) = T(2*n-1,n-2), where T is defined in A026022.

Conjecture: (13*n+118)*(n+5)*a(n) +(13*n^2-969*n-3490)*a(n-1) +2*(-203*n^2+1080*n+758)*a(n-2) +4*(73*n-182)*(2*n-5)*a(n-3)=0. - _R. J. Mathar_, Jun 22 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001113 | OEIS-A026029

Number of (s(0), s(1), ..., s(2n)) such that s(i) is a nonnegative integer and |s(i) - s(i-1)| = 1 for i = 1,2,...,n, s(0) = 3, s(2n) = 3. Also T(2n,n), where T is defined in A026022.

Conjecture: (n+4)*a(n) + 6*(-n-2)*a(n-1) + 4*(2*n-1)*a(n-2) = 0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001114 | OEIS-A026030

a(n) = T(2n,n-1), where T is defined in A026022.

Conjecture: (n+5)*a(n) -2*(5*n+16)*a(n-1) +(35*n+47)*a(n-2) +2*(-25*n+14)*a(n-3) + 12*(2*n-5)*a(n-4) = 0. - _R. J. Mathar_, Jun 15 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001115 | OEIS-A026031

a(n) = number of (s(0), s(1), ..., s(2n)) such that s(i) is a nonnegative integer and |s(i) - s(i-1)| = 1 for i = 1,2,...,n, s(0) = 3, s(2n) = 7. Also a(n) = T(2n,n-2), where T is defined in A026022.

Conjecture: (n+6)*a(n) +10*(-n-4)*a(n-1) +2*(17*n+32)*a(n-2) +4*(-11*n+4)*a(n-3) +8*(2*n-5)*a(n-4)=0. - _R. J. Mathar_, Jun 22 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001116 | OEIS-A026050

a(n) = (d(n)-r(n))/2, where d = A026049 and r is the periodic sequence with fundamental period (1,0,0,1).

Conjecture: G.f. x^7*(59-158*x+203*x^2-202*x^3+145*x^4-45*x^5) / ((x^2+1)*(x-1)^4). - _R. J. Mathar_, Jun 22 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001117 | OEIS-A026070

a(n) = number of (s(0), s(1), ..., s(n)) such that every s(i) is an integer, s(0) = 0, |s(i) - s(i-1)| = 1 for i = 1,2; |s(i) - s(i-1)| <= 1 for i >= 3, s(n) = 3. Also a(n) = T(n,n-3), where T is the array defined in A024996.

Conjecture: (n+3)*a(n) +(-5*n-6)*a(n-1) +(3*n-11)*a(n-2) +(11*n-8)*a(n-3) +2*(-2*n+17)*a(n-4) +6*(-n+6)*a(n-5)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001118 | OEIS-A026071

a(n) = number of (s(0), s(1), ..., s(n)) such that every s(i) is an integer, s(0) = 0, |s(i) - s(i-1)| = 1 for i = 1,2; |s(i) - s(i-1)| <= 1 for i >= 3, s(n) = 4. Also a(n) = T(n,n-4), where T is the array defined in A024996.

Conjecture: -(n-4)*(n+4)*a(n) +(4*n+7)*(n-4)*a(n-1) +(-2*n^2+23*n-12)*a(n-2) -(4*n+3)*(n-4)*a(n-3) +3*(n-4)*(n-5)*a(n-4)=0. - _R. J. Mathar_, Jun 22 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001119 | OEIS-A026083

a(n) = number of (s(0), s(1), ..., s(n)) such that every s(i) is an integer, s(0) = 0 = s(n), |s(i) - s(i-1)| = 1 for i = 1,2,3; |s(i) - s(i-1)| <= 1 for i >= 4. Also a(n) = T(n,n), where T is the array defined in A026082.

Conjecture: n*a(n) +(-3*n+5)*a(n-1) +(-n-6)*a(n-2) +3*(n-5)*a(n-3)=0. - _R. J. Mathar_, Jun 22 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001120 | OEIS-A026084

a(n) = number of (s(0), s(1), ..., s(n)) such that every s(i) is an integer, s(0) = 0, |s(i) - s(i-1)| = 1 for i = 1,2,3; |s(i) - s(i-1)| <= 1 for i >= 4, s(n) = 1. Also a(n) = T(n,n-1), where T is the array defined in A026082.

Conjecture: (n+1)*a(n) +3*(-n+1)*a(n-1) +(-n-9)*a(n-2) +3*(n-5)*a(n-3)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001121 | OEIS-A026085

a(n) = number of (s(0), s(1), ..., s(n)) such that every s(i) is an integer, s(0) = 0, |s(i) - s(i-1)| = 1 for i = 1,2,3; |s(i) - s(i-1)| <= 1 for i >= 4, s(n) = 2. Also a(n) = T(n,n-2), where T is the array defined in A026082.

Conjecture: (n+2)*a(n) +(-5*n-1)*a(n-1) +4*(n-4)*a(n-2) +8*(n-1)*a(n-3) +(-5*n+34)*a(n-4) +3*(-n+7)*a(n-5)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001122 | OEIS-A026086

Number of (s(0), s(1), ..., s(n)) such that s(0) = 0, |s(i) - s(i-1)| = 1 for i = 1,2,3; |s(i) - s(i-1)| <= 1 for i >= 4, s(n) = 3; also a(n) = T(n,n-3), where T is the array defined in A026082.

Conjecture: (n+3)*a(n) +5*(-n-1)*a(n-1) +3*(n-5)*a(n-2) +(11*n-9)*a(n-3) +4*(-n+11)*a(n-4) +6*(-n+7)*a(n-5)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001123 | OEIS-A026087

a(n) = number of (s(0), s(1), ..., s(n)) such that every s(i) is an integer, s(0) = 0, |s(i) - s(i-1)| = 1 for i = 1,2,3; |s(i) - s(i-1)| <= 1 for i >= 4, s(n) = 4. Also a(n) = T(n,n-4), where T is the array defined in A026082.

Conjecture: -(n-4)*(n+4)*a(n) +(4*n^2-11*n-27)*a(n-1) +(-2*n^2+29*n-24)*a(n-2) -(4*n+5)*(n-5)*a(n-3) +3*(n-5)*(n-6)*a(n-4)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001124 | OEIS-A026110

a(n) = number of (s(0), s(1), ..., s(n)) such that every s(i) is a nonnegative integer, s(0) = 0, s(1) = 1, s(n) = 4, |s(i) - s(i-1)| <= 1 for i >= 2. Also a(n) = T(n,n-4), where T is the array defined in A026105.

Conjecture: -(n+6)*(n-4)*a(n) +(4*n^2-n-51)*a(n-1) +(-2*n^2+11*n+18)*a(n-2) -(4*n-1)*(n-3)*a(n-3) +3*(n-3)*(n-4)*a(n-4)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001125 | OEIS-A026122

a(n) is the number of (s(0),s(1),...,s(n)) such that every s(i) is a nonnegative integer, s(0) = 1, s(n) = 1, |s(1) - s(0)| = 1, |s(i) - s(i-1)| <= 1 for i >= 2. Also a(n) = T(n,n), where T is the array in A026120.

Conjecture: (n+4)*a(n) +(-3*n-5)*a(n-1) +(-n-6)*a(n-2) +3*(n-3)*a(n-3)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001126 | OEIS-A026125

a(n) = number of (s(0),s(1),...,s(n)) such that every s(i) is a nonnegative integer, s(0) = 1, s(n) = 4, |s(1) - s(0)| = 1, |s(i) - s(i-1)| <= 1 for i >= 2. Also a(n) = T(n,n-3), where T is the array in A026120.

Conjecture: -(n+7)*(n-3)*a(n) +(4*n+17)*(n-3)*a(n-1) +(-2*n^2+13*n+27)*a(n-2) -(4*n+5)*(n-3)*a(n-3) +3*(n-3)*(n-4)*a(n-4)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001127 | OEIS-A026126

a(n) = number of (s(0),s(1),...,s(n)) such that every s(i) is a nonnegative integer, s(0) = 1, s(n) = 5, |s(1) - s(0)| = 1, |s(i) - s(i-1)| <= 1 for i >= 2. Also a(n) = T(n,n-4), where T is the array in A026120.

Conjecture: -(n+8)*(n-4)*a(n) +(4*n^2+5*n-73)*a(n-1) +(-2*n^2+13*n+38)*a(n-2) -(4*n+5)*(n-3)*a(n-3) +3*(n-3)*(n-4)*a(n-4)=0. - _R. J. Mathar_, Jun 10 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001128 | OEIS-A026135

Number of (s(0),s(1),...,s(n)) such that every s(i) is a nonnegative integer, s(0) = 1, |s(1) - s(0)| = 1, |s(i) - s(i-1)| <= 1 for i >= 2. Also sum of numbers in row n+1 of the array T defined in A026120.

Conjecture: (n+2)*a(n) +3*(-n-1)*a(n-1) +(-n-2)*a(n-2) +3*(n-3)*a(n-3)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001129 | OEIS-A026152

a(n) = T(n,n-1), where T is the array in A026148.

Conjecture: -(n+7)*(31*n-87)*a(n) +(164*n^2+313*n-2115)*a(n-1) +(-182*n^2+229*n+891)*a(n-2) +(-164*n^2+59*n+519)*a(n-3) +3*(71*n-104)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001130 | OEIS-A026153

T(n,n-2), where T is the array in A026148.

Conjecture: -(n+8)*(n-118)*a(n) +(-n^2-561*n-2878)*a(n-1) +2*(38*n^2+203*n-1070)*a(n-2) +2*(-116*n^2+781*n+2493)*a(n-3) +(13*n^2-1012*n+3888)*a(n-4) +3*(107*n-423)*(n-4)*a(n-5)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001131 | OEIS-A026154

a(n) = T(n,n-3), where T is the array in A026148.

Conjecture: (n+9)*a(n) +(-7*n-47)*a(n-1) +(13*n+51)*a(n-2) +(5*n+37)*a(n-3) +2*(-13*n-6)*a(n-4) +2*(n-13)*a(n-5) +12*(n-4)*a(n-6)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001132 | OEIS-A026155

T(n,n-4), where T is the array in A026148.

Conjectures from _Chai Wah Wu_, Mar 23 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001133 | OEIS-A026163

Sum{T(k,k-1)}, k = 1,2,...,n, where T is the array in A026148.

Conjectures from _Mark van Hoeij_, Oct 30 2011: (Start)
Conjecture: -(n+7)*(3*n-31)*a(n) +3*(-n^2-35*n-76)*a(n-1) +2*(32*n^2+27*n-459)*a(n-2) +(-47*n^2+286*n-204)*a(n-3) -3*(37*n-51)*(n-2)*a(n-4)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001134 | OEIS-A026165

Number of (s(0), s(1), ..., s(n)) such that every s(i) is a nonnegative integer, s(0) = 2, |s(1) - s(0)| = 1, |s(i) - s(i-1)| <= 1 for i >= 2. Also sum of numbers in row n+1 of the array T in A026148.

Conjecture: (n+3)*a(n) + (-5*n-9)*a(n-1) + (5*n+1)*a(n-2) + 5*(n-1)*a(n-3) + 6*(-n+3)*a(n-4) = 0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001135 | OEIS-A026179

Numbers k such that A026177(j) < A026177(k) for all j < k.

Old conjecture: (a(n)) = complement of A026225 after removal of the initial 1 here. [Note that following the proof of the form of its terms, A026225 has been renamed accordingly. - _Peter Munn_, Mar 24 2022]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001136 | OEIS-A026201

Position of n in A026200.

Conjectures from _Colin Barker_, Oct 14 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001137 | OEIS-A026219

Position of n in A026218.

Conjectures from _Colin Barker_, Oct 15 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001138 | OEIS-A026225

Numbers of the form 3^i * (3k+1).

Conjectures based on old name: these are numbers of the form (3*i+1)*3^j; see A182828, and they comprise the complement of A026179, except for the initial 1 in A026179.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001139 | OEIS-A026243

a(n) = A000522(n) - 2.

Conjecture: a(n) +(-n-2)*a(n-1) +(2*n-1)*a(n-2) +(-n+2)*a(n-3)=0. - _R. J. Mathar_, Jun 23 2013 [Confirmed by _Altug Alkan_, May 18 2018]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, based on a message from a correspondent who wishes to remain anonymous, Dec 21 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001140 | OEIS-A026270

Number of (s(0), s(1), ..., s(n)) such that every s(i) is a nonnegative integer, s(0) = 0, s(1) = 1 = s(n), |s(i) - s(i-1)| <= 1 for i >= 2, |s(2) - s(1)| = 1, |s(3) - s(2)| = 1 if s(2) = 1. Also T(n,n-1), where T is the array in A026268.

Conjecture: (n+3)*a(n) +3*(-n-1)*a(n-1) +(-n-1)*a(n-2) +3*(n-5)*a(n-3)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001141 | OEIS-A026273

a(n) = least k such that s(k) = n, where s = A026272.

This observation also gives an answer to Lenormand's question, and a simple proof of Mathar's conjecture in A059426.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001142 | OEIS-A026274

Greatest k such that s(k) = n, where s = A026272.

Conjecture: This sequence consists precisely of those numbers without a 1 or 2 in their Zeckendorf representation. [In other words, numbers which are the sum of distinct nonconsecutive Fibonacci numbers greater than 2.] - _Charles R Greathouse IV_, Jan 28 2015 Proof of the conjecture by _Charles R Greathouse IV_. Then the same characterization holds for n with d(1)d(0) = 01, since 11 does not appear in a Zeckendorf expansion. But such an n has predecessor n-1 which always has an expansion with d(1)d(0) = 00. Combined with my comment from March 2018, this proves the conjecture (ignoring n = 0). (End) a(n) = A003622(n+1) - 1 for n>1 (conjectured). - _Michel Marcus_, Oct 30 2020 This conjectured formula follows directly from the formula a(n) = floor((n+1)*r^2)-2 in my Mar 12 2018 comment above. - _Michel Dekking_, Oct 31 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001143 | OEIS-A026299

Number of (s(0), s(1), ..., s(n)) such that every s(i) is a nonnegative integer, s(0) = 0, s(1) = 1, |s(i) - s(i-1)| <= 1 for i >= 2, |s(2) - s(1)| = 1, |s(3) - s(2)| = 1 if s(2) = 1. Also sum of numbers in row n+1 of the array T in A026268.

Conjecture: (n+2)*a(n) +(-3*n-4)*a(n-1) +(-n+2)*a(n-2) +3*(n-4)*a(n-3)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001144 | OEIS-A026327

a(n) = number of (s(0), s(1), ..., s(n)) such that s(i) is a nonnegative integer and |s(i) - s(i-1)| <= 1 for i = 1,2,...,n, s(0) = 2, s(n) = 4. Also a(n) = T(n,n-2), where T is the array in A026323.

Conjecture: -(n+8)*(n-2)*a(n) +3*(2*n^2+7*n-28)*a(n-1) +3*(-3*n^2-n+36)*a(n-2) -4*(n+4)*(n-1)*a(n-3) +12*(n-1)*(n-2)*a(n-4)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001145 | OEIS-A026328

a(n) = number of (s(0), s(1), ..., s(n)) such that s(i) is a nonnegative integer and |s(i) - s(i-1)| <= 1 for i = 1,2,...,n, s(0) = 2, s(n) = 5. Also a(n) = T(n,n-3), where T is the array in A026323.

Conjecture: -(n+9)*(n-3)*a(n) +(6*n^2+21*n-128)*a(n-1) +(-9*n^2-3*n+152)*a(n-2) -4*(n+4)*(n-1)*a(n-3) +12*(n-1)*(n-2)*a(n-4)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001146 | OEIS-A026329

a(n) = number of (s(0), s(1), ..., s(n)) such that s(i) is a nonnegative integer and |s(i) - s(i-1)| <= 1 for i = 1,2,...,n, s(0) = 2, s(n) = 6. Also a(n) = T(n,n-4), where T is the array in A026323.

Conjecture: -(n+10)*(n-4)*a(n) +3*(2*n+15)*(n-4)*a(n-1) +3*(-3*n^2-n+68)*a(n-2) -4*(n+4)*(n-1)*a(n-3) +12*(n-1)*(n-2)*a(n-4)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001147 | OEIS-A026363

a(n) is the least k such that s(k) = n, where s = A026362.

Conjecture: -1 < n*r - a(n) < 2 for n>=1, where r = (1 + sqrt(3))/2. - _Clark Kimberling_, Apr 29 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001148 | OEIS-A026364

a(n) = greatest k such that s(k) = n, where s = A026362.

Complement of A026363. Positions of 0 in the fixed point of the morphism 0->11, 1->101; see A285430. Conjecture: -1 < n*r - a(n) < 4 for n>=1, where r = 2 + sqrt(3). - _Clark Kimberling_, Apr 28 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001149 | OEIS-A026367

a(n) = least k such that s(k) = n, where s = A026366.

Conjecture: -1 < n*r - a(n) < 2 for n>=1, where r = (1 + sqrt(3))/2. - _Clark Kimberling_, Apr 29 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001150 | OEIS-A026368

a(n) = greatest k such that s(k) = n, where s = A026366.

Positions of 0 in the fixed point of the morphism 0->11, 1->110; see A285431. Conjecture: -2 < n*r - a(n) < 4 for n>=1, where r = 2 + sqrt(3). - _Clark Kimberling_, Apr 29 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001151 | OEIS-A026377

a(n) = number of integer strings s(0),...,s(n) counted by array T in A026374 that have s(n)=4; also a(n) = T(2n,n-2).

Conjecture: n*(n+4)*a(n) - 3*(n+2)*(2*n+3)*a(n-1) + 5*(n+2)*(n+1)*a(n-2) = 0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001152 | OEIS-A026389

a(n) = number of integer strings s(0),...,s(n) counted by array T in A026386 that have s(n)=4; also a(n) = T(2n,n-2).

Conjecture: (n+2)*a(n) +4*(-3*n-2)*a(n-1) +2*(24*n-19)*a(n-2) +4*(-18*n+43)*a(n-3) +35*(n-4)*a(n-4)=0. - _R. J. Mathar_, May 29 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001153 | OEIS-A026392

T(n,[n/2]), where T is the array in A026386.

Conjecture: (n+1)*a(n) +2*(n-1)*a(n-1) +2*(-3*n+1)*a(n-2) +4*(-3*n+7)*a(n-3) +5*(n-3)*a(n-4) +10*(n-5)*a(n-5)=0. - _R. J. Mathar_, Feb 10 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001154 | OEIS-A026465

Length of n-th run of identical symbols in the Thue-Morse sequence A010060 (or A001285).

Under tau^2 the runs of length 1 are mapped to the sequence 1,2,1 of run lengths, and the runs of length 2 are mapped to the sequence 1,2,2,2,1 of run lengths. This proves John Layman's conjecture number (1): it follows that (a(n)) is fixed point of the morphism alpha Since alpha(1) and alpha(2) are both palindromes, this also proves Alexander Povolotsky's conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001155 | OEIS-A026477

a(1) = 1, a(2) = 2, a(3) = 3; and for n > 3, a(n) = smallest number > a(n-1) and not of the form a(i)*a(j)*a(k) for 1 <= i < j < k < n.

Once a term with a given prime signature S (i.e., multiset of prime exponents) appears, then all numbers with the same prime signature follow. So either all or no terms with the same prime signature appear (first conjectured by _Charles R Greathouse IV_).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001156 | OEIS-A026488

a(n) is the least positive integer > a(n-1) and not a(i)*a(j)-a(k) for 1 <= i <= j <= k <= n, where a(1) = 1.

Conjecture: This is the increasing sequence of numbers congruent to 1 mod 3, with 3 inserted. Conjecture holds for n <= 1300. - _Sean A. Irvine_, Oct 12 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001157 | OEIS-A026570

a(n) = A026568(n,n-1), also a(n) = number of integer strings s(0),...,s(n) counted by A026568 such that s(n)=1.

Conjecture: (n+1)*a(n) -2*n*a(n-1) +(-3*n-1)*a(n-2) +2*(2*n-3)*a(n-3)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001158 | OEIS-A026571

a(n) = T(n,n-2), T given by A026568. Also a(n) = number of integer strings s(0), ..., s(n) counted by T, such that s(n) = 2.

Conjecture: (n+2)*a(n) + 3*(-n-1)*a(n-1) - 3*n*a(n-2) + 11*(n-1)*a(n-3) + 2*(n-6)*a(n-4) + 4*(-2*n+7)*a(n-5) = 0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001159 | OEIS-A026572

a(n) = T(n,n-3), T given by A026568. Also a(n) = number of integer strings s(0),...,s(n) counted by T, such that s(n)=3.

Conjectured g.f.: (1/2)*((-3*x^2-x+1)*((x-1)/(4*x^2+x-1))^(1/2)-1+x+x^2)/x^3. - _Mark van Hoeij_, Oct 30 2011
Conjecture: (n+3)*a(n) +(-3*n-5)*a(n-1) +4*(-n-1)*a(n-2) +(13*n-9)*a(n-3) +(5*n-9)*a(n-4) +6*(-2*n+7)*a(n-5)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001160 | OEIS-A026573

a(n)=T(n,n-4), T given by A026568. Also a(n) = number of integer strings s(0),...,s(n) counted by T, such that s(n)=4.

Conjecture: -2*(n-4)*(n+4)*a(n) +(5*n^2-5*n-46)*a(n-1) +(4*n^2+n+18)*a(n-2) +(-11*n^2+24*n-16)*a(n-3) +2*(n-2)*(2*n-5)*a(n-4)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001161 | OEIS-A026580

T(n,0) + T(n,1) + ... + T(n,n), T given by A026568.

Conjecture: n*a(n) +(n-2)*a(n-1) +(-17*n+28)*a(n-2) +5*(-n+4)*a(n-3) +2*(34*n-107)*a(n-4) +16*(n-5)*a(n-5) +32*(-2*n+9)*a(n-6)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001162 | OEIS-A026582

a(n) = Sum{T(i,j)}, 0<=j<=i, 0<=i<=n, T given by A026568.

Conjecture: n*a(n) +2*(-2*n+1)*a(n-1) +2*(-n+3)*a(n-2) +20*(n-2)*a(n-3) +(-7*n+6)*a(n-4) +2*(-12*n+41)*a(n-5) +8*(2*n-7)*a(n-6)=0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001163 | OEIS-A026587

a(n) = T(n, n-2), T given by A026584. Also a(n) = number of integer strings s(0),...,s(n) counted by T, such that s(n)=2.

Conjecture: (n+2)*a(n) = (3*n+2)*a(n-1) +(3*n+2)*a(n-2) -(11*n-16)*a(n-3) -2*(n-3)*a(n-4) +4*(2*n-9)*a(n-5). - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001164 | OEIS-A026589

a(n) = T(n,n-4), T given by A026584. Also a(n) = number of integer strings s(0),...,s(n) counted by T, such that s(n)=4.

Conjecture: -(n+4)*(65*n-269)*a(n) +(-65*n^2+140*n+1933)*a(n-1) +(809*n^2-2431*n-4514)*a(n-2) +(-123*n^2+2496*n-205)*a(n-3) +2*(-726*n^2+3737*n-4395)*a(n-4) +8*(56*n-215)*(2*n-9)*a(n-5) = 0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001165 | OEIS-A026596

Row sums of A026584.

Conjecture: n*a(n) -3*(n-1)*a(n-1) -(5*n-6)*a(n-2) +3*(5*n-13)*a(n-3) +2*(4*n-9)*a(n-4) -8*(2*n-9)*a(n-5) = 0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001166 | OEIS-A026598

a(n) = Sum_{i=0..n} Sum_{j=0..i} T(i,j), T given by A026584.

Conjecture: n*a(n) - (4*n-3)*a(n-1) - (2*n-3)*a(n-2) + 5*(4*n-9)*a(n-3) - 7*(n-3)*a(n-4) - 6*(4*n-15)*a(n-5) + 8*(2*n-9)*a(n-6) = 0. - _R. J. Mathar_, Jun 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001167 | OEIS-A026641

Number of nodes of even outdegree (including leaves) in all ordered trees with n edges.

Bala's conjecture is true (see Fried link). - _Sela Fried_, Oct 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001168 | OEIS-A026672

a(n) = T(2n,n-1), T given by A026670. Also T(2n,n-1)=T(2n+1,n+2), T given by A026725; and T(2n,n-1), T given by A026736.

Conjecture: -(n+1)*(n-6)*a(n) +2*(4*n^2-23*n+3)*a(n-1) +3*(-5*n^2+33*n-42)*a(n-2) -2*(2*n-3)*(n-5)*a(n-3)=0. - _R. J. Mathar_, Aug 08 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001169 | OEIS-A026710

T(n,1) + T(n,2) + ... + T(n,n), T given by A026703.

Conjecture: D-finite with recurrence 2*(-n+2)*a(n) +2*(-n+6)*a(n-1) +(21*n-62)*a(n-2) +(19*n-118)*a(n-3) +(-69*n+296)*a(n-4) +2*(-25*n+161)*a(n-5) +3*(25*n-158)*a(n-6) +(25*n-162)*a(n-7) +(-29*n+212)*a(n-8) +2*(-2*n+13)*a(n-9) +4*(n-8)*a(n-10)=0. - _R. J. Mathar_, Jun 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001170 | OEIS-A026732

a(n) = Sum_{k=0..n} T(n,k), T given by A026725.

Conjecture: +(-n+1)*a(n) +2*a(n-1) +3*(3*n-7)*a(n-2) -10*a(n-3) +(-23*n+95)*a(n-4) +6*a(n-5) +(11*n-95)*a(n-6) +2*a(n-7) +4*(n-7)*a(n-8)=0. - _R. J. Mathar_, Oct 26 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001171 | OEIS-A026733

a(n) = Sum_{k=0..floor(n/2)} T(n,k), T given by A026725.

Conjecture: (-n+2)*a(n) +(n-2)*a(n-1) +2*(4*n-13)*a(n-2) +8*(-n+4)*a(n-3) +5*(-3*n+14)*a(n-4) +(15*n-94)*a(n-5) +2*(-2*n+9)*a(n-6) +4*(n-6)*a(n-7)=0. - _R. J. Mathar_, Oct 26 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001172 | OEIS-A026735

a(n) = Sum_{k=0..floor(n/2)} T(n-k,k), T given by A026725.

Conjecture: G.f.:-1/2*(2*x^6-5*x^4+8*x^3+x-2+x*(x-1)*(x^2+x+1)*(1-4*x^3)^(1/2))/(x^6+4*x^3-1)/(x^2+x-1). - Maksym Voznyy (voznyy(AT)mail.ru), Aug 11 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001173 | OEIS-A026741

a(n) = n if n odd, n/2 if n even.

Conjecture: lcm(n, p) = p*a(n, p), gcd(n, p) = n/a(n, p). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | J. Carl Bellinger (carlb(AT)ctron.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001174 | OEIS-A026743

a(n) = Sum_{j=0..n} T(n,j), T given by A026736.

Conjecture D-finite with recurrence n*a(n) -2*a(n-1) +(-11*n+20)*a(n-2) +14*a(n-3) +(39*n-152)*a(n-4) -22*a(n-5) +(-41*n+268)*a(n-6) -6*a(n-7) +12*(-n+6)*a(n-8)=0. - _R. J. Mathar_, Jan 13 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001175 | OEIS-A026765

a(n) = Sum_{k=0..n} T(n,k), T given by A026758.

Conjecture: G.f.: -(1-2*x-5*x^2+10*x^3 - sqrt(1-10*x^2+29*x^4-20*x^6))/(2*x*(1-2*x-5*x^2+10*x^3)). - Maksym Voznyy (voznyy(AT)mail.ru), Aug 11 2009 Conjecture: (n+1)*a(n) -2*a(n-1) +2*(-5*n+3)*a(n-2) +12*a(n-3) +(29*n-71)*a(n-4) -10*a(n-5) +20*(-n+5)*a(n-6)=0. - _R. J. Mathar_, Jun 30 2013 Conjecture: a(n) ~ (2+sqrt(5) + (-1)^n*(2-sqrt(5))) * 5^(n/2) / sqrt(2*Pi*n). - _Vaclav Kotesovec_, Feb 12 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001176 | OEIS-A026767

a(n) = Sum_{i=0..n} Sum_{j=0..n} T(i,j), T given by A026758.

Conjecture: (n+1)*a(n) +(-n-3)*a(n-1) +2*(-5*n+4)*a(n-2) +2*(5*n+3)*a(n-3) +(29*n-83)*a(n-4) +(-29*n+61)*a(n-5) +10*(-2*n+11)*a(n-6) +20*(n-5)*a(n-7)=0. - _R. J. Mathar_, Jun 30 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001177 | OEIS-A026772

a(n) = T(2n, n-2), T given by A026769.

Conjecture D-finite with recurrence +n*(683440564498775*n-3571267774541221)*(n+3)*(n+2)*a(n) -(n+2)*(14680943042715007*n^3 -77306402200099276*n^2 -165100190278545 95*n +32930242824559728)*a(n-1) +(115576812096278739*n^4 -727966310087371570*n^3 +337449371031094809*n^2 +647464064695228030*n -329302428245597280)*a(n-2) +(-3467163524964043 27*n^4 +3103154468191226534*n^3 -7064840562647596109*n^2 +5446737046090126798*n -1580651655578866944)*a(n-3) +(-254358906902769779*n^4 +522032785791792 6494*n^3 -3553950692892363997*n^2 +8709131554139818 6522*n -5991616393029499 0824)*a(n-4) +(4096127756151299723*n^4 -7503744282987220 6910*n^3 +5023015019600095 75069*n^2 -1440579074098159574434*n +1469340925940446228608)*a(n-5) +(-9245640254696084879*n^4 +2005594358095559 13066*n^3 -1613433648622480143829*n^2 +5696178287053438445634*n -7425674215347839990712)*a(n-6) +(752059790053091 2203*n^4 -1914053265088322077 18*n^3 +1814954924843492047041*n^2 -760547392719340998 4358*n

+11895887009351293007160)*a(n-7)
+4*(-461380856200630754*n^4
+13108517874955450215*n^3
-137711835536290769230*n^2
+633868893726787892361*n
-107789691593668009176O)*a(n-8)
+4*(n-10)*(8653064513346403*n
-44432714618958651)*(2*n-13)*(2*n-19)*a(n-9)=0. - _R. J. Mathar_, Jun 30 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001178 | OEIS-A026796

Number of partitions of n in which the least part is 3.

Let b(k) be the number of partitions of k for which twice the number of ones is the number of parts, k = 0, 1, 2, ... Then a(n+4) = b(n), n = 0, 1, 2, ... (conjectured). - _George Beck_, Aug 19 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001179 | OEIS-A026876

T(2n+1, n+1), T given by A026758.

The invert transform of this sequences (shifted) appears to be A165538 (conjectured). - _Michael D. Weiner_, Aug 04 2018 G.f. f satisfies: x^6*f^4 + x^3*(4*x-1)*(x-1)*f^3 + x*(x-1)*(6*x^2-6*x+1)*f^2 + (4*x-1)*(x-1)^2*f + (x-1)^2 = 0 (conjectured). - _Michael D. Weiner_, Aug 04 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001180 | OEIS-A026923

Number of partitions of n into an odd number of parts, the greatest being 3; also, a(n+5) = number of partitions of n+2 into an even number of parts, each <= 3.

Conjectures from _Colin Barker_, Sep 01 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001181 | OEIS-A026927

Number of partitions of n into an even number of parts, the greatest being 3; also, a(n+5) = number of partitions of n+2 into an odd number of parts, each <= 3.

Conjectures from _Colin Barker_, Sep 01 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001182 | OEIS-A026970

a(n) = Sum_{k=0..n} (k+1) * A026637(n,k).

Conjectures from _Colin Barker_, Feb 18 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001183 | OEIS-A026975

a(n) = Sum_{k=0..n} (k+1) * A026648(n,k).

Conjectures from _Colin Barker_, Feb 18 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001184 | OEIS-A027027

a(n) = T(n, 2n-3), T given by A027023.

Conjecture: D-finite with recurrence (n+1)*a(n) +(-8*n-1)*a(n-1) +(19*n-14)*a(n-2) +2*(-3*n-1)*a(n-3) +(-21*n+89)*a(n-4) +(8*n-45)*a(n-5) +(n-4)*a(n-6) +6*(n-4)*a(n-7)=0. - _R. J. Mathar_, Jun 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001185 | OEIS-A027028

a(n) = T(n,2n-4), T given by A027023.

Conjecture D-finite with recurrence -(n+2)*(n-6)*a(n) +3*(n+1)*(2*n-11)*a(n-1) -n*(7*n-31)*a(n-2) -2*(n-3)*(4*n-19)*a(n-3) +(5*n^2-38*n+84)*a(n-4) +(2*n-5)*(n-3)*a(n-5) +3*(n-3)*(n-4)*a(n-6)=0. - _R. J. Mathar_, Jun 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001186 | OEIS-A027050

a(n) = T(n,2n-1), T given by A027023.

Conjecture D-finite with recurrence (-n+1)*a(n) +3*(2*n-3)*a(n-1) +(-7*n+10)*a(n-2) +2*(-4*n+19)*a(n-3) +(5*n-23)*a(n-4) +(2*n-5)*a(n-5) +3*(n-4)*a(n-6)=0. - _R. J. Mathar_, Jun 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001187 | OEIS-A027051

a(n) = T(n,2n-2), T given by A027023.

Conjecture: D-finite with recurrence n*a(n) +(-7*n+5)*a(n-1) +(13*n-18)*a(n-2) +(n-13)*a(n-3) +(-13*n+64)*a(n-4) +(3*n-25)*a(n-5) +(-n+2)*a(n-6) +3*(n-5)*a(n-7)=0. - _R. J. Mathar_, Jun 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001188 | OEIS-A027056

a(n) = A027052(n, 2n-1).

Conjecture:b D-finite with recurrence (-n+1)*a(n) +2*(3*n-4)*a(n-1) +(-7*n+3)*a(n-2) +4*(-2*n+13)*a(n-3) +(5*n-29)*a(n-4) +2*(n-2)*a(n-5) +3*(n-5)*a(n-6)=0. - _R. J. Mathar_, Jun 15 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001189 | OEIS-A027058

a(n) = A027052(n, 2n-2).

Conjecture: D-finite with recurrence n*a(n) +(-7*n+4)*a(n-1) +(13*n-10)*a(n-2) +(n-34)*a(n-3) +(-13*n+84)*a(n-4) +(3*n-32)*a(n-5) +(-n+6)*a(n-6) +3*(n-6)*a(n-7)=0. - _R. J. Mathar_, Jun 15 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001190 | OEIS-A027059

a(n) = A027052(n, 2n-3).

Conjecture D-finite with recurrence (n+1)*a(n) +2*(-4*n-1)*a(n-1) +(19*n-5)*a(n-2) +6*(-n-5)*a(n-3) +3*(-7*n+41)*a(n-4) +2*(4*n-29)*a(n-5) +(n+1)*a(n-6) +6*(n-5)*a(n-7)=0. - _R. J. Mathar_, Jun 15 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001191 | OEIS-A027087

a(n) = A027082(n, 2n).

Conjecture: D-finite with recurrence +3*n*a(n) +3*(-5*n+3)*a(n-1) +(7*n-4)*a(n-2) +(25*n-67)*a(n-3) +(17*n-44)*a(n-4) +(19*n-97)*a(n-5) +(5*n-32)*a(n-6) +3*(n-7)*a(n-7)=0. - _R. J. Mathar_, Jun 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001192 | OEIS-A027088

a(n) = A027082(n, 2n-1).

Conjecture: D-finite with recurrence 3*(n+1)*(18*n-85)*a(n) +(292*n^2-1316*n-255)*a(n-1) +(-218*n^2+1213*n-1056)*a(n-2) +2*(-248*n^2+1666*n-2223)*a(n-3) +(-50*n^2-497*n+3243)*a(n-4) +(-52*n^2+80*n+765)*a(n-5) +3*(22*n-83)*(n-6)*a(n-6)=0. - _R. J. Mathar_, Jun 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001193 | OEIS-A027089

a(n) = A027082(n, 2n-2).

Conjecture: D-finite with recurrence -(n+2)*(244*n-1065)*a(n) +3*(n+1)*(422*n -1703)*a(n-1) +2*(-359*n^2 +1211*n -1420)*a(n-2) +2*(-1174*n^2 +6319*n -7410)*a(n-3) +(-760*n^2 +2173*n +1616)*a(n-4) +(-502*n^2 +2245*n -1123)*a(n-5) +6*(23*n-67) *(n-5)*a(n-6)=0. - _R. J. Mathar_, Jun 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001194 | OEIS-A027116

a(n) = A027113(n, n+3).

Conjectures from _Colin Barker_, Feb 20 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001195 | OEIS-A027117

a(n) = A027113(n, n+4).

Conjectures from _Colin Barker_, Feb 20 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001196 | OEIS-A027154

a(n) = Sum_{k=0..floor(n/2)} A027144(n-k, k).

Conjectures from _Chai Wah Wu_, Jun 10 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001197 | OEIS-A027174

a(n) = A027170(2n, n-2).

Conjecture: D-finite with recurrence -2*(n+3)*(139*n-175)*a(n) +(1279*n^2+1334*n-1089)*a(n-1) -2*(167*n+121)*(2*n-3)*a(n-2) +12*(111*n^2+456*n+229)=0. - _R. J. Mathar_, May 06 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001198 | OEIS-A027175

a(n) = A027170(2n-1, n-1).

Conjecture: D-finite with recurrence 2*(n+1)*a(n) +(-7*n-5)*a(n-1) +2*(-2*n+3)*a(n-2) +12*(-3*n+1)=0. - _R. J. Mathar_, May 06 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001199 | OEIS-A027179

a(n) = T(n,0) + T(n,1) + ... + T(n,[n/2]), T given by A027170.

Conjecture: D-finite with recurrence -(n+2)*(3*n-20)*a(n) -(n+1)*(3*n-11)*a(n-1) +(33*n^2-180*n-100)*a(n-2) +(15*n^2-56*n+1)*a(n-3) +2*(-51*n^2+313*n-208)*a(n-4) -12*(n-1)*(n-4)*a(n-5) +8*(9*n-11)*(n-5)*a(n-6) +4*(-24*n^2+196*n-365)=0. - _R. J. Mathar_, May 06 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001200 | OEIS-A027182

a(n) = self-convolution of row n of array T given by A027170.

Conjecture: D-finite with recurrence 2*(n+2)*a(n) +(-17*n-23)*a(n-1) +(47*n+36)*a(n-2) +2*(-22*n-17)*a(n-3) +4*(-n+20)*a(n-4) +8*(2*n-9)*a(n-5) +24*(12*n-47)=0. - _R. J. Mathar_, May 06 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001201 | OEIS-A027261

a(n) = Sum_{k=0..2n} (k+1) * A025177(n, k).

a(n) = 2(n+1)*3^(n-1), for n>1 (conjectured). - _Ralf Stephan_, Feb 02 2004 Conjecture: a(n) = 6*a(n-1)-9*a(n-2), for n>3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001202 | OEIS-A027302

a(n) = Sum_{k=0..floor((n-1)/2)} T(n,k) * T(n,k+1), with T given by A008315.

Conjecture D-finite with recurrence -(n+2)*(13*n-2)*(3+n)^2*a(n) +10*(8*n^2+3*n-8)*(n+2)^2*a(n-1) +8*(12*n^4+47*n^3+52*n^2+67*n+20)*a(n-2) -160*(8*n^2+3*n-8)*(n-1)^2*a(n-3) +128*(7*n+4)*(2*n-5)*(-2+n)^2*a(n-4)=0. - _R. J. Mathar_,

Nov 22 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001203 | OEIS-A027433

Sum over all 2^(2n) pairs (u,v) of binary sequences of length n of length of maximal common subsequence between them.

The proved bounds for gamma_2 (see asymptotic formula below) are 0.788071 <= gamma_2 <= 0.82628, and conjectured value is around 0.811 [see Dixon].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Zimmermann_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001204 | OEIS-A027481

Second subdiagonal of triangle A027477, constructed from the Stirling numbers of the first kind.

Conjectures from _Chai Wah Wu_, Apr 08 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001205 | OEIS-A027615

Number of 1's when n is written in base -2.

Conjecture: a(n) != A053737(n) if and only if there exists even k >= 4 such that n mod 2^k >= (5*2^(k+1) + 2)/3. If this holds, then the probability of a random chosen number n to satisfy a(n) != A053737(n) is 1/6. (End) a(n) = 3*A072894(n+1) - 2*n - 3. Proof by Nikolaus Meyberg, following a conjecture by _Ralf Stephan_. - _R. J. Mathar_, Jan 11 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pontus von Brömssen_, Nov 14 1997 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001206 | OEIS-A027617

Number of permutations of n elements containing a 3-cycle.

Conjectures from _Stéphane Rézel_, Dec 11 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Joe Keane (jgk(AT)jgk.org) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001207 | OEIS-A027619

Numbers k such that Hofstadter Q-sequence Q(k) (A005185) satisfies Q(k) = k/2.

Conjecture: a(n)/n^2 is bounded. - _Benoit Cloitre_, Oct 26 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | G. R. Bower (fsgrb(AT)aurora.alaska.edu) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001208 | OEIS-A027649

a(n) = 2*(3^n) - 2^n.

Related to the number of even a's in a nontrivial cycle (should one exist) in the 3x+1 Problem, where a <= floor(log_2(2*(3^n) - 2^n)). The value n correlates to the number of odds in such a nontrivial cycle. See page 1288 of Crandall's paper. Also, this relation gives another proof that the number of odds divided by the number of evens in a nontrivial cycle is bounded by log 2 / log 3 (this observation does not resolve the finite cycles conjecture as the value could be arbitrarily close to this bound). However, the same argument gives that log 2 / log 3 is less than or equal to the number of odds divided by the number of evens in a divergent sequence (should one exist), as log 2 / log 3 is the limit value for a cycle of an arbitrarily large length, where the length is given by the value n. - _Jeffrey R. Goodwin_, Aug 04 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001209 | OEIS-A027687

4-perfect (quadruply-perfect or sous-triple) numbers: sum of divisors of n is 4n.

It is conjectured that there are only finitely many terms. - _N. J. A. Sloane_, Jul 22 2012 Conjecture: A010888(a(n)) divides a(n). Tested for n up to 36 incl. - _Ivan N. Ianakiev_, Oct 31 2013 It seems that for all such equations there exist such an infinite set of solutions. So I conjecture that the sequence is infinite! (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jean-Yves Perrier (nperrj(AT)ascom.ch) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001210 | OEIS-A027697

Odious primes: primes with odd number of 1's in binary expansion.

Conjecture: a(n) < A027699(n) except for n = 2; verified up to n=5*10^7. Moreover, I conjecture that A027699(n) - a(n) tends to infinity. - _Vladimir Shevelev_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001211 | OEIS-A027699

Evil primes: primes with even number of 1's in their binary expansion.

Comment from _Vladimir Shevelev_, Jun 01 2007: Conjecture: If pi_1(m) is the number of a(n) not exceeding m and pi_2(m) is the number of A027697(n) not exceeding m then pi_1(m) <= smaller than pi_2(m) for all natural m except m=5 and m=6. I verified this conjecture up to 10^9. Moreover I conjecture that pi_2(m)-pi_1(m) tends to infinity with records at the primes m=2, 13, 41, 61, 67, 79, 109, 131, 137, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001212 | OEIS-A027833

Distances between successive 2's in sequence A001223 of differences between consecutive primes.

Conjecture: a(n) < log(A014574(n))^2. - _Thomas Ordowski_, Jul 21 2012 Conjecture: All positive integers are represented in this sequence. This is verified up to 184, by searching up to prime indexes of ~128000000. The rate of filling-in the smallest remaining gap among the integers, and the growth in the maximum value found, both slow down considerably relative to a fixed quantity of twin prime incidences examined in each pass. The maximum value found was 237. - _Richard R. Forberg_, Jul 28 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jean-Marc MALASOMA (Malasoma(AT)entpe.fr) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001213 | OEIS-A027854

Mutinous numbers: n > 1 such that n/p^k > p, where p is the largest prime dividing n and p^k is the highest power of p dividing n.

If p = A006530(a(n)) then p * a(n) is in the sequence. E.g., as 12 is in the sequence with gpf(12) = A006530(12) = 3, 12*3^k is in the sequence for k > 0. Conjecture: if m is in the sequence then so is A003961(m). - _David A. Corneth_, Jul 13 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001214 | OEIS-A027870

Number of zero digits in 2^n.

I conjecture that any value x = 0, 1, 2, ... occurs only a finite number of times N(x) = 36, 41, 31, 34, 25, 32, 37, 23, 43, 47, 33, ... in this sequence, for the last time at well defined indices i(x) = 86, 229, 231, 359, 283, 357, 475, 476, 649, 733, 648, ... - _M. F. Hasler_, Jul 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001215 | OEIS-A027908

a(n) = T(2*n, n), T given by A027907.

Conjectural: a(n*p^k) == a(n*p^(k-1)) (mod p^(2*k)) for all primes p >= 5 and positive integers n and k. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001216 | OEIS-A027984

a(n) = Sum_{k=0..n} T(n, k)*T(n, n+k), T given by A027960.

Conjectures from _Colin Barker_, Feb 25 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001217 | OEIS-A028231

From hexagons in a circle problem.

a(n) = 15*a(n-1)-15*a(n-2)+a(n-3). G.f.: (1+7*x-2*x^2)/((1-x)*(1-14*x+x^2)). - conjectured by _Colin Barker_, Apr 10 2012; these follow easily from the formula.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001218 | OEIS-A028342

Expansion of Product_{i>=1} (1 - x^i)^(-1/i); also of exp(Sum_{n>=1} (d(n)*x^n/n)) where d is number of divisors function.

It appears that the sequence taken modulo 10 is periodic with period 10. More generally, we conjecture Peter Bala's conjectures are correct (see Kallat, 2026). - _Ahaan Kallat_, Jul 23 2026 Conjecture: log(a(n)/n!) ~ log(2)/2 * log(n)^2. - _Vaclav Kotesovec_, Sep 15 2018 The above conjecture is incorrect:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001219 | OEIS-A028355

How the astronomical clock ("Orloj") in Prague would strike 1,2,3,...,24,25,.. (digits follow 12343212343... (A028356), n-th group adds to n).

Conjectures from _Chai Wah Wu_, Apr 18 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _J. H. Conway_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001220 | OEIS-A028359

Two-bell analog of A028355.

Conjectures from _Chai Wah Wu_, Apr 18 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001221 | OEIS-A028362

Total number of self-dual binary codes of length 2n. Totally isotropic spaces of index n in symplectic geometry of dimension 2n.

Conjectural o.g.f. as a continued fraction of Stieltjes type (S-fraction):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001222 | OEIS-A028388

Good primes (version 2): prime(n) such that prime(n)^2 > prime(n-i)*prime(n+i) for all 1 <= i <= n-1.

Selfridge conjectured, and Pomerance proved, that there are infinitely many numbers in this sequence. Pomerance asks if the sequence has density 0. - _Charles R Greathouse IV_, Apr 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001223 | OEIS-A028393

Iterate the map in A006368 starting at 8.

It is conjectured that this trajectory never repeats, but no proof of this has been found. - _N. J. A. Sloane_, Jul 14 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. H. Conway_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001224 | OEIS-A028495

Expansion of g.f. (1-x^2)/(1-x-2*x^2+x^3).

Conjectures: a(n) ~ A160389 * a(n-1). Also a(n) ~ sqrt(A116425)*a(n-1). - _Bill McEachen_, Feb 11 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001225 | OEIS-A028576

a(n) = (1/(4*n)) * Sum_{d|n} mu(n/d) * binomial(2*d, d)^2.

Conjecture: a(n) is divisible by n and 3*a(n) by n^2. - _F. Chapoton_, Feb 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | Lionel Levine (levine(AT)ultranet.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001226 | OEIS-A028723

a(n) = (1/4)*floor(n/2)*floor((n-1)/2)*floor((n-2)/2)*floor((n-3)/2).

Conjectured to be crossing number of complete graph K_n, see A000241. For n >= 3, a(n-2) is conjectured to be the crossing number of the path complement graph on n vertices. - _Eric W. Weisstein_, Oct 02 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001227 | OEIS-A028836

Iterated sum of digits of n is a power of 2.

Conjectures from _Colin Barker_, Feb 10 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001228 | OEIS-A028841

Iterated sum of digits of n is a Fibonacci number.

Conjectures from _Colin Barker_, Feb 18 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001229 | OEIS-A028870

Numbers k such that k^2 - 2 is prime.

It is conjectured that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001230 | OEIS-A028884

a(n) = (n + 3)^2 - 8.

Conjecture: a(n) = A079896(A265284(n-1)). -

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001231 | OEIS-A029576

Number of permutations of an n-set containing a 9-cycle.

Conjectures from _Stéphane Rézel_, Nov 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rob Pratt_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001232 | OEIS-A029577

Number of permutations of an n-set containing a 10-cycle.

Conjectures from _Stéphane Rézel_, Dec 11 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rob Pratt_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001233 | OEIS-A029766

Unary-binary rooted trees with n nodes.

Conjecture: for n > 0, a(n) = n! * A040039(n-1). - _Vaclav Kotesovec_, Sep 30 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001234 | OEIS-A029790

None of the digits in k is present in k^2 or k^3.

I strongly conjecture that a(28) does not exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001235 | OEIS-A029795

Numbers k such that k and k^3 have the same set of digits.

Conjecture: there exists some m and N for which a(n) = m + n for all n >= N. - _Charles R Greathouse IV_, Jun 28 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001236 | OEIS-A029797

Numbers k such that k^2 and k^3 have the same set of digits.

Conjecture: there exists some m and N for which a(n) = m + n for all n >= N. [_Charles R Greathouse IV_, Jun 28 2011] This conjecture is false. If the conjecture is true then for some N we would have k is in the sequence if k >= n. But 10^e + 1 (A062397) is not in the sequence for any integer e >= 0. - _David A. Corneth_, Nov 13 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001237 | OEIS-A029858

a(n) = (3^n - 3)/2.

Conjectured to be the number of integers from 0 to 10^(n-1) - 1 that lack 0, 1, 2, 3, 4, 5 and 6 as a digit. - _Alexandre Wajnberg_, Apr 25 2005. This is easily verified to be true. - _Renzo Benedetti_, Sep 25 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001238 | OEIS-A029902

h(n), where strategic move on g(n) X 3 rectangle in Chomp is (h(n),2). Conjectured to be complementary to r(n), see A029905.

h(n), where strategic move on g(n) X 3 rectangle in Chomp is (h(n),2). Conjectured to be complementary to r(n), see A029905.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Fred Lunnon_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001239 | OEIS-A029905

r(n), where exists one-parameter family of strategic partitions (k+p(n)+q(n), k+q(n), r(n)) for k = 0,1,2,... in Chomp.

Conjectured to be complementary to h(n), see A029902. Maybe the conjecture is false - see the Brouwer link. - _Andries E. Brouwer_, May 31 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Fred Lunnon_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001240 | OEIS-A030051

Numbers from the 290-theorem.

The 290-theorem, conjectured by Conway and Schneeberger and proved by Bhargava and Hanke, asserts that a positive definite quadratic form represents all numbers iff it represents the numbers in this sequence. - _T. D. Noe_, Mar 30 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001241 | OEIS-A030052

Smallest number whose n-th power is a sum of distinct smaller positive n-th powers.

a(n) >= A332101(n) = A078607(n)+2 (conjectured). - _M. F. Hasler_, May 25 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | Richard C. Schroeppel | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001242 | OEIS-A030057

Least number that is not a sum of distinct divisors of n.

The indices of records occur at the highly abundant numbers, excluding 3 and 10, if _Jaycob Coleman_'s conjecture at A002093 that all these numbers are practical numbers (A005153) is true. - _Amiram Eldar_, Jun 13 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001243 | OEIS-A030063

Fermat's Diophantine m-tuple: 1 + the product of any two distinct terms is a square.

It is conjectured that there do not exist five strictly positive integers with the property that 1 + the product of any two distinct terms is a square. (See Dujella's links.) - _Jonathan Sondow_, Apr 04 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | Graham Lewis (grahaml(AT)levygee.com.uk) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001244 | OEIS-A030077

Take n equally spaced points on circle, connect them by a path with n-1 line segments; sequence gives number of distinct path lengths.

For n points on a circle, there are floor(n/2) distinct line segment lengths. Hence an upper bound for a(n) is the number of compositions of n-1 into floor(n/2) nonnegative parts, which is A099578(n-2). Conjecture: the upper bound is attained if n is prime. There are A052558(n-2) paths to be considered. - _T. D. Noe_, Jan 09 2007 [Edited by _Petros Hadjicostas_, Jul 19 2018]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Lurie Gittelson_, Dec 11 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001245 | OEIS-A030101

a(n) is the number produced when n is converted to binary digits, the binary digits are reversed and then converted back into a decimal number.

Conjecture: a(n) = 2*w(n) - 2*w(A053645(n)) - 1 for n > 0, where w = A264596. - _Velin Yanev_, Sep 12 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001246 | OEIS-A030112

Number of antichains in the product of a fence poset with n elements and a chain with 7 elements; also number of paths with n turns when light is reflected from 8 glass plates.

For a k-glass sequence, say a(n,k), a(n,k) is always asymptotic to z(k)*w(k)^n where w(k)=(1/2)/cos(k*Pi/(2k+1)) and it is conjectured that z(k) is the root 1<x<2 of a polynomial of degree Phi(2k+1)/2. - _Benoit Cloitre_, Oct 16 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jacques Haubrich (jhaubrich(AT)freeler.nl) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001247 | OEIS-A030127

Term at which n-Somos sequence first becomes nonintegral.

Conjecture: for even terms, a(n) = 2*n, else a(n) = 2*n+1 when n is in A101591. - _Paul D. Hanna_, Dec 08 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001248 | OEIS-A030173

Differences p(i)-p(j) between primes, sorted in numerical order.

Conjectured (Polignac 1849) to be union of even numbers and the odd primes minus 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Alexander Grasser [Graesser] (alex(AT)computicket.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001249 | OEIS-A030179

Quarter-squares squared: A002620^2.

Conjectured to be crossing number of complete bipartite graph K_{n,n}. Known to be true for n <= 7. If the Zarankiewicz conjecture is true, then a(n) is also the

rectilinear crossing number of K_{n,n}. - _Eric W. Weisstein_, Apr 24 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001250 | OEIS-A030204

Expansion of q^(-1/8) * eta(q) * eta(q^2) in powers of q.

Conjecture: E = A244920 = 2*log(1+sqrt(2)). - _Vaclav Kotesovec_, Sep 26 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001251 | OEIS-A030210

Expansion of (eta(q) * eta(q^5))^4 in powers of q.

Conjecture: |a(p)| < 2*p^(3/2) for p prime. - _Michael Somos_, Oct 31 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001252 | OEIS-A030229

Numbers that are the product of an even number of distinct primes.

Conjecture: For the matrix M(i,j) = 1 if j|i and 0 otherwise, Inverse(M)(a,1) = -1, for any a in this sequence. - _Benedict W. J. Irwin_, Jul 26 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001253 | OEIS-A030469

Primes which are concatenations of three consecutive primes.

It is conjectured that sequence is infinite. - from Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Nov 09 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001254 | OEIS-A030503

Graham-Sloane-type lower bound on the size of a ternary (n,3,3) constant-weight code.

Conjectures from _Colin Barker_, Aug 02 2019: (Start) Conjectures confirmed.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Mattias Svanstrom (mattias(AT)isy.liu.se) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001255 | OEIS-A030547

Number of terms (including the initial term) needed to reach a palindrome when the Reverse Then Add! map (x -> x + (x-with-digits-reversed)) is repeatedly applied to n, or -1 if a palindrome is never reached.

It is conjectured that a(196) is the smallest term equal to -1. See A023108.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001256 | OEIS-A030701

Decimal expansion of 4^n contains no zeros (probably finite).

Integers in A007377 / 2. Conjectured to be finite, and probably complete. - _M. F. Hasler_, Mar 08 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001257 | OEIS-A030797

Decimal expansion of the constant x such that x^x = e. Inverse of W(1), where W is Lambert's function.

Conjecture: Another series can be defined as follows. Let z = a + b*i <> 0 be complex, and let z = v^v. Then log(z) + v = v*(1 + log(v)), so f(z, v) = (log(z) + v)/(1 + log(v)) = v. Suppose lim_{n -> infinity} (log(z) + v(n))/(1 + Log(v(n))) = v, for some sequence {v(n)}. Then, since v(n) -> v(n+1), similarly f_(n+1)(z, v) = v(n+1) = (log(z) + v(n))/(1 + log(v(n))). If Im(z) <> 0, recall that log(z) is multi-valued, so one might take both log(z) and log(v(n)) modulo 2*Pi*i. If Im(z) = 0 (i.e., if z is real), then one should use the recurrence f_(n+1)(z, v) = v(n+1) = (log(z) + v(n))/(1 + abs(log(v(n)))). For example, when z = e, we have lim_{n -> infinity} (1 + v(n))/(1 + abs(log(v(n)))) = 1.763222..., for v(0) <> 1/e, with apparent quadratic convergence, and most rapidly when v(0) = 1. Pathologies occur when v(0) is in the vicinity of a fixed point of f(z, v); e.g., if z = 2^(1/4), then such a fixed point is c = 0.806693797003867301..., so f_(n)(z, v) -> c, for all n, with a(0) near c. The constant c was calculated to 250 digits by Joerg Arndt. - _L. Edson Jeffery_, Apr 12 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Simon Plouffe_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001258 | OEIS-A031149

Numbers whose square with its last digit deleted is also a square.

a(n) = 38*a(n-7) - a(n-14) for n>15 (conjectured). - _Colin Barker_, Dec 31 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001259 | OEIS-A031172

a(n) = prime(n+10) - prime(n).

In principle, moderate values should appear infinitely many times, by analogy with twin primes hypothesis. For example, a(n) = 44 for n = 9, 17, 206, 1604467, 12905293, 18008874, 26545460, 32655424, 57848470, 58313630, 59022635, 66275281, 81581956, 123780499, 160884754, 167797255, 179786560, 181569324, 239542290, ... - _Zak Seidov_, Sep 14 2014, edited by _M. F. Hasler_, Dec 03 2018 According to the k-tuple conjecture, any admissible k-tuple of primes occurs with calculable nonzero asymptotic density, i.e., in particular, infinitely many times. For k = 11, number of primes in the interval [prime(n), prime(n+10)], the smallest possible diameter of a k-tuple is A008407(11) = 36, and there are A083409(11) = 2 such constellations: {0, 4, 6, 10, 16, 18, 24, 28, 30, 34, 36}, first occurring at A213646(1) = 1418575498573, and {0, 2, 6, 8, 12, 18, 20, 26, 30, 32, 36}, first occurring at A213647(1) = 11. The combined list { prime(n) | a(n) = 36 } is A257129. - _M. F. Hasler_, Dec 03 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeff Burch_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001260 | OEIS-A031217

Number of terms in longest arithmetic progression of consecutive primes starting at n-th prime (conjectured to be unbounded).

Number of terms in longest arithmetic progression of consecutive primes starting at n-th prime (conjectured to be unbounded).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001261 | OEIS-A031740

Numbers k such that the least term in the periodic part of the continued fraction for sqrt(k) is 62.

Conjecture: a(n) = n*(961*n+1). a(n) = 3*a(n-1)-3*a(n-2)+a(n-3). G.f.: 2*x*(481+480*x)/(1-x)^3. - _Colin Barker_, Sep 30 2012 Conjecture is false. m*(961*m+1) for m >= 1 is a subsequence (see comment in A031712) and 3940288 is a term not of the form m*(961*m+1). - _Chai Wah Wu_, Jun 19 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001262 | OEIS-A031924

Primes followed by a gap of 6, i.e., next prime is p + 6.

Conjecture: The sequence is infinite and for every n >= 7746, a(n+1) < a(n)^(1+1/n). Namely for n >= 7746, a(n)^(1/n) is a strictly decreasing function of n (See comment lines of the sequence A248855). - _Jahangeer Kholdi_ and _Farideh Firoozbakht_, Nov 29 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeff Burch_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001263 | OEIS-A031926

Lower prime of a difference of 8 between consecutive primes.

Conjecture: The sequence is infinite and for every n, a(n+1) < a(n)^(1+1/n). Namely a(n)^(1/n) is a strictly decreasing function of n (see comment lines of the sequence A248855). - _Jahangeer Kholdi_ and _Farideh Firoozbakht_, Nov 29 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeff Burch_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001264 | OEIS-A031928

Lower prime of a difference of 10 between consecutive primes.

Conjecture: The sequence is infinite and for every n, a(n+1) < a(n)^(1+1/n). Namely, a(n)^(1/n) is a strictly decreasing function of n (see comments at A248855). - _Jahangeer Kholdi_ and _Farideh Firoozbakht_, Nov 29 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Jul 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001265 | OEIS-A031932

Lower prime of a pair of consecutive primes having a difference of 14.

Conjecture: The sequence is infinite and for every n, a(n+1) < a(n)^(1+1/n); i.e., a(n)^(1/n) is a strictly decreasing function of n (see comment lines of the sequence A248855). - _Jahangeer Kholdi_ and _Farideh Firoozbakht_, Nov 29 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeff Burch_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001266 | OEIS-A032086

Number of reversible strings with n beads of 3 colors. If more than 1 bead, not palindromic.

Conjectures from _Colin Barker_, Apr 02 2012: (Start) Conjectures from _Colin Barker_, Mar 09 2017: (Start) The above conjectures are true: The second set follows from the definition and the first set can be derived from that. - _Andrew Howroyd_, Oct 10 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001267 | OEIS-A032087

Number of reversible strings with n beads of 4 colors. If more than 1 bead, not palindromic.

The formula for a(n) for this sequence was Ralf Stephan's conjecture 72. It was solved by Elizabeth Wilmer (see Proposition 1 in one of the links below). She does not accept Bower's assertion that a string of length 1 is not palindromic. - _Petros Hadjicostas_, Jul 05 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001268 | OEIS-A032088

Number of reversible strings with n beads of 5 colors. If more than 1 bead, not palindromic.

Conjectures from _Colin Barker_, Jul 07 2012: (Start) Conjectures from _Colin Barker_, Mar 09 2017: (Start) The above conjectures are true: The second set follows from the definition and the first set can be derived from that. - _Andrew Howroyd_, Oct 10 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001269 | OEIS-A032091

Number of reversible strings with n-1 beads of 2 colors. 4 beads are black. String is not palindromic.

The formula for a(n) for this sequence was Ralf Stephan's conjecture 73. It was solved by Elizabeth Wilmer (see Proposition 2 in one of the links below). There is a minor typo in the original conjecture. - _Petros Hadjicostas_, Jul 04 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001270 | OEIS-A032096

"BHK" (reversible, identity, unlabeled) transform of 2,2,2,2,...

For the current sequence, c(n) = 2 for all n >= 1, and thus, C(x) = 2*x/(1-x). Substituting into the above formula for A(x), and doing the algebra, we get A(x) =

x*(2-5*x-4*x^2+15*x^3)/((1-x)*(1-3*x)*(1-3*x^2)), which is the formula conjectured by _Colin Barker_ below. Conjecture: a(n) = 4*a(n-1) - 12*a(n-3) + 9*a(n-4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001271 | OEIS-A032098

"BHK" (reversible, identity, unlabeled) transform of 3,3,3,3,...

For the current sequence, c(n) = 3 for all n >= 1, and thus, C(x) = 3*x/(1-x). Substituting into the above formula for A(x), and doing the algebra, we get A(x) = 3*x*(7*x^2 - 5*x + 1)/((1-x)*(1-2*x)*(1-4*x)), which is the formula conjectured by _Colin Barker_ below. Using partial fraction decomposition, we get A(x) = -21/8 + 3/(1-x) - 3/(4*(1-2*x)) + 3/(8*(1-4*x)), from which we get a(n) = 3 * (2^(2*n-3) - 2^(n-2) + 1), which is _Ralf Stephan_'s conjecture below. Conjecture: a(n) = 3 * (2^(2*n-3) - 2^(n-2) + 1). - _Ralf Stephan_, Sep 11 2003 Conjecture: a(n) = 7*a(n-1) - 14*a(n-2) + 8*a(n-3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001272 | OEIS-A032106

Number of reversible strings with n black beads and n-1 white beads. String is not palindromic.

The formula for a(n) for this sequence was Ralf Stephan's conjecture 74. It was solved by Elizabeth Wilmer (see Proposition 3 in one of the links below). She does not accept Bower's assertion that a string of length 1 is not palindromic. - _Petros Hadjicostas_, Jul 04 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001273 | OEIS-A032115

"BIJ" (reversible, indistinct, labeled) transform of 1,3,5,7...

Conjecture: D-finite with recurrence a(n) +2*(-n-1)*a(n-1) +4*(n-1)*a(n-2) +2*a(n-3) +(-4*n+11)*a(n-4) +2*(n-4)*a(n-5)=0. - _R. J. Mathar_, May 08 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001274 | OEIS-A032123

Number of 2n-bead black-white reversible strings with n black beads.

Conjecture: D-finite with recurrence n*(n-1)*a(n) -2*(n-1)*(3*n-4)*a(n-1) +4*(2*n^2-14*n+19)*a(n-2) +8*(n^2+5*n-19)*a(n-3) -16*(n-3)*(3*n-10)*a(n-4) +32*(n-4)*(2*n-9)*a(n-5)=0, n>5. - _R. J. Mathar_, Nov 09 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001275 | OEIS-A032132

Number of series-reduced dyslexic planted planar trees with n leaves.

Note that b(m=0, n) = 0 for n <> 1, but b(m=0, n=1) = 1; b(m,n) = 0 when m >= n >= 1; and b(m=1, n) = 1 for n>=2. Also, b(m,m+1) = A001190(m+1) for m>=1, which are the Wedderburn-Etherington numbers, and apparently b(m=2, n) = A024206(n-1) for n>=2 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001276 | OEIS-A032191

Number of necklaces with 6 black beads and n-6 white beads.

G.f.: (1-x+x^2+4*x^3+2*x^4+3*x^6+x^7+x^8)/((1-x)^6*(1+x)^3*(1+x+x^2)^2*(1-x+x^2)) (conjectured). - _Ralf Stephan_, May 05 2004 G.f.: (x^6)*(1-x+x^2+4*x^3+2*x^4+3*x^6+x^7+x^8)/((1-x)^2*(1-x^2)^2*(1-x^3)*(1-x^6)) . (proving the R. Stephan conjecture (with the correct offset) in a different version; see Comments entry above). - _Wolfdieter Lang_, Feb 15 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001277 | OEIS-A032247

"DHK[6]" (bracelet, identity, unlabeled, 6 parts) transform of 1,1,1,1,...

Here, a(n) is the number of aperiodic bracelets with k = 6 black beads and n-k = n-6 white beads that have no reflection symmetry. We conjecture that we can use Herbert Kociemba's formula from the documentation of sequences A008804 and A032246 to derive the g.f. of (a(n): n >= 1). See below for more details. - _Petros Hadjicostas_, Feb 24 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001278 | OEIS-A032249

"DHK[8]" (bracelet, identity, unlabeled, 8 parts) transform of 1,1,1,1,...

Here, a(n) is the number of aperiodic bracelets with k = 8 black beads and n-k = n-8 white beads that have no reflection symmetry. We conjecture that we can use Herbert Kociemba's formula from the documentation of sequences A008804 and A032246 to derive the g.f. of (a(n): n >= 1). See below for more details. - _Petros Hadjicostas_, Feb 24 2019 We conjecture that g.f. = Sum_{n>=1} a(n)*x^n = gf(8,x) - gf(4, x^2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001279 | OEIS-A032250

"DHK[n](2n)" (bracelet, identity, unlabeled, n parts, evaluated at 2n) transform of 1,1,1,1,...

We conjecture that gg(k, x)= Sum_{d|k} mu(d)*ff(k/d, x^d).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001280 | OEIS-A032312

"EGJ" (unordered, element, labeled) transform of 2,2,2,2...

Conjecture: the Gauss congruences a(n*p^k) == a(n*p^(k-1)) (mod p^k) hold for all primes p and positive integers n and k. Cf. A007837.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001281 | OEIS-A032357

Convolution of Catalan numbers and powers of -1.

Conjecture: (n+1)*a(n) + 3*(-n+1)*a(n-1) + 2*(-2*n+1)*a(n-2) = 0. - _R. J. Mathar_, Nov 30 2012 Conjecture is true since the g.f. satisfies (x - 3*x^2 - 4*x^3)*g'(x) + (1 - 6*x^2)*g(x) = 1. - _Robert Israel_, May 22 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001282 | OEIS-A032446

Number of solutions to phi(k) = 2n.

By Carmichael's conjecture, a(n) <> 1 for any n. See A074987. - _Thomas Ordowski_, Sep 13 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ursula Gagelmann (gagelmann(AT)altavista.net) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001283 | OEIS-A032448

Smallest prime == -1 modulo prime(n).

It appears that a(n) <= prime(n)^2-1, where prime(n) = A000040(n) is the n-th prime; see A035095 for a related conjecture. If correct, this implies A006530(a(n)+1)=prime(n), which in turn implies that there are no duplicated values in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jun 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001284 | OEIS-A032450

Period of finite sequence g(n) related to Poulet's Conjecture.

Period of finite sequence g(n) related to Poulet's Conjecture. Poulet's Conjecture states that for any integer n, the sequence f_0(n) = n, f_2k+1(n)=sigma(f_2k(n)), f_2k(n)=phi(f_2k-1(n)) (where sigma = A000203 and phi = A000010) is eventually periodic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ursula Gagelmann (gagelmann(AT)altavista.net), Apr 07 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001285 | OEIS-A032594

Greater of two consecutive palindromes, both of which are prime.

Conjecture: sequence is finite. - _Charles R Greathouse IV_, Mar 22 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, May 15 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001286 | OEIS-A032743

Super-2 Numbers (2 * n^2 contains substring '22' in its decimal expansion).

Conjecture: a(n) ~ n. - _Charles R Greathouse IV_, Dec 04 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, May 15 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001287 | OEIS-A032744

Super-4 Numbers (4 * n^4 contains substring '4444' in its decimal expansion).

Conjecture: a(n) ~ n. - _Charles R Greathouse IV_, Dec 04 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, May 15 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001288 | OEIS-A032745

Super-5 Numbers (5 * n^5 contains substring '55555' in its decimal expansion).

Conjecture: a(n) ~ n. - _Charles R Greathouse IV_, Dec 04 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, May 15 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001289 | OEIS-A032746

Super-6 Numbers (6 * n^6 contains substring '666666' in its decimal expansion).

Conjecture: a(n) ~ n. - _Charles R Greathouse IV_, Dec 04 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, May 15 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001290 | OEIS-A032747

Super-7 Numbers (7 * n^7 contains substring '7777777' in its decimal expansion).

Conjecture: a(n) ~ n. - _Charles R Greathouse IV_, Dec 04 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, May 15 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001291 | OEIS-A032748

Super-8 Numbers (8 * n^8 contains substring '88888888' in its decimal expansion).

Conjecture: a(n) ~ n. - _Charles R Greathouse IV_, Dec 04 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, May 15 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001292 | OEIS-A032908

One of four 3rd-order recurring sequences for which the first derived sequence and the Galois transformed sequence coincide.

Conjecture: consecutive terms of this sequence and consecutive terms of A101265 provide all the positive integer solutions of (a+b)*(a+b+1) == 0 (mod (a*b)). - _Robert Israel_, Aug 26 2015 Conjecture is true: see Mathematics Stack Exchange link. - _Robert Israel_, Sep 06 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | Michele Elia (elia(AT)polito.it) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001293 | OEIS-A033093

Number of 0's when n is written in base b for 2<=b<=n+1.

Conjecture: lim inf a(n) = infinity. The lim inf grows quite slowly: e.g. a(2047)=7. Weaker conjecture: 2, 3 and 7 are the only n for which a(n) = 1. Note that a(n)=1 implies n=2 or n=2^k-1; probabilistically, the chance that 2^k-1 has no zeros just in base 3 is C^k, where C = (2/3)^(log(2)/log(3)) ~ .774, so the number of exceptions should be finite. It appears that 2^15-1 is the last n with no zeros in either base 2 or 3 (and it does have zeros in base 5). - _Franklin T. Adams-Watters_, Nov 07 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001294 | OEIS-A033188

Minimal difference of any increasing arithmetic progression of n primes.

David W. Wilson conjectures that a(n) = n# (n primorial, A034386) for n >= 8. If this is true then a(22) onwards are: 9699690, 223092870, 223092870, 223092870, 223092870, 223092870, 223092870, 6469693230, 6469693230, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001295 | OEIS-A033212

Primes congruent to 1 or 19 (mod 30).

Three further theorems (these were originally stated as conjectures, but are now known to be theorems, thanks to the work of J. B. Tunnell - see link):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001296 | OEIS-A033235

Primes of the form x^2 + 55*y^2.

Conjecture: consecutive elements of this sequence are consecutive primes satisfying the congruence b(k) == 1 (mod k) for k>0, where b(k) is recursive sequence defined as follows: b(k) = -b(k-1) - b(k-2) + b(k-3) - b(k-4) with b(0)=2, b(1)=1, b(2)=0, b(3)=-1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001297 | OEIS-A033282

Triangle read by rows: T(n, k) is the number of diagonal dissections of a convex n-gon into k+1 regions.

The preceding conjecture by Lang is true. It is implicit in Copeland's 2011 comments in A086810 on the relations among a gf and its compositional inverse for that entry and inversion through A133437 (a differently normalized version of A111785), whose integer partitions are the same as those for A134685. (An inversion pair in Copeland's 2008 formulas below can also be used to prove the conjecture.) In addition, it follows from the relation between the inversion formula of A111785/A133437 and the enumeration of distinct faces of associahedra. See the MathOverflow link concernimg Loday and the Aguiar and Ardila reference in A133437 for proofs of the relations between the partition polynomials for inversion and enumeration of the distinct faces of the A_n associahedra, or Stasheff polytopes. - _Tom Copeland_, Dec 21 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001298 | OEIS-A033451

Initial prime in set of 4 consecutive primes with common difference 6.

It is conjectured that there exist arbitrarily long sequences of consecutive primes in arithmetic progression. As of March 2013 the record is 10 primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeff Burch_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001299 | OEIS-A033485

a(n) = a(n-1) + a(floor(n/2)), a(1) = 1.

Conjecture: lim_{n->oo} a(2n)/a(n)*log(n)/n = c = 1.64.... and a(n)/A(n) is bounded where A(n)=1 if n is a power of 2, otherwise A(n) = sqrt(n)*Product_{k<log_2(n)} n/2^k/log(n/2^k). - _Benoit Cloitre_, Jan 26 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_. This was in the 1973 "Handbook", but was then dropped from the database. Resubmitted by _Philippe Delélham_. Entry revised by _N. J. A. Sloane_, Jun 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001300 | OEIS-A033487

a(n) = n*(n+1)*(n+2)*(n+3)/4.

Crossing number of the (n+3)-cocktail party graph (conjectured). - _Eric W. Weisstein_, Apr 29 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001301 | OEIS-A033493

Sum of the numbers in the trajectory of n for the 3x+1 problem.

Conjecture: a(n) = A006577(n)^2 only at a(3) = 49. Verified for n <= 10^7. - _Luca Santarsiero_, Jul 13 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeff Burch_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001302 | OEIS-A033616

Coordination sequence T1 for Zeolite Code TSC.

The following is a another conjectured recurrence, found by gfun, using the command rec:=gfun[listtorec](t1, a(n)); (where t1 is a list of the initial terms) suggested by _Paul Zimmermann_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf W. Grosse-Kunstleve_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001303 | OEIS-A033617

Coordination sequence T2 for Zeolite Code TSC.

The following is another conjectured recurrence, found by gfun, using the command rec:=gfun[listtorec](t1, a(n)); (where t1 is a list of the initial terms) suggested by _Paul Zimmermann_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf W. Grosse-Kunstleve_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001304 | OEIS-A033627

0-additive sequence: not the sum of any previous pair.

Conjecture: a(n+1) is the number of distinct numbers of steps required for the last n digits of integers to repeat themselves by iterating the map m -> m^2 + 1. - _Ya-Ping Lu_, Oct 19 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001305 | OEIS-A033631

Numbers k such that sigma(phi(k)) = sigma(k) where sigma is the sum of divisors function A000203 and phi is the Euler totient function A000010.

There is another class of [conjecturally] infinite subsets connected to A005385 (safe primes). Examples: Let s,t be safe primes, s<>t, then 3^2*5*251*s, 2^2*61*71*s, 2*61*s*t and 2*19*311*s are in this sequence. So is 3*s*A108510(m). (There are some obvious exceptions for small s, t.) - _Vim Wenders_, Dec 27 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001306 | OEIS-A033662

Possible digital sums of Smith numbers (conjectural).

Possible digital sums of Smith numbers (conjectural).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001307 | OEIS-A033686

One-ninth of theta series of A2[hole]^2.

Conjectural g.f.: A(x) = Sum_{n = -oo..oo} x^n/(1 - x^(3*n+2))^2 = Sum_{n = -oo..oo} x^(5*n+1)/(1 - x^(3*n+1))^2. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001308 | OEIS-A033810

Number of people needed so that probability of at least two sharing a birthday out of n possible days is at least 50%.

a(n) = ceiling(sqrt(2*n*log(2)) + (3 - 2*log(2))/6 + (9 - 4*log(2)^2) / (72*sqrt(2*n*log(2))) - 2*log(2)^2/(135*n)) for all n up to 10^18. It is conjectured that this formula holds for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001309 | OEIS-A033919

Odd k for which k+2^m is composite for all m < k.

In an archived website, Payam Samidoost gives these numbers and other results about the dual Sierpiński problem. It is conjectured that, for each of these k<78557, there is an m such that k+2^m is prime. Then a covering argument would show that 78557 is the least odd number such that 78557+2^m is composite for all m. The impediment in the "dual" problem is that it is currently very difficult to prove the primality of large numbers of the form k+2^m. It is much easier to prove the Proth primes of the form k*2^m+1 which occur in the usual Sierpiński problem. According to the distributed search project "Five or Bust", 40291 is the only value of k < 78557 for which there is currently no m known making k + 2^m a prime or probable prime. - _T. D. Noe_, Jun 14 2007 and Phil Moore (moorep(AT)lanecc.edu), Dec 14 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dan Hoey_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001310 | OEIS-A033932

Least positive m such that n! + m is prime.

Conjecture: No term is a composite number. a(n) is a prime > 3*prime(k), where k is such that prime(k) < n <= prime(k+1). - _Amarnath Murthy_, Apr 07 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeff Burch_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001311 | OEIS-A033933

Least nonnegative m such that n! - m is prime.

Conjecture: for n >= 3, a(n) is 1 or a prime. - _Amarnath Murthy_, Mar 19 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeff Burch_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001312 | OEIS-A033955

Sum of the remainders when the n-th prime is divided by primes up to the (n-1)-th prime.

The sequence does not change if defined to go up to the n-th prime, as that contributes 0 to the sum; it's merely a computational optimization. Conjecture: For all n > 6, a(n) > prime(n). A corollary of this conjecture is that a(6) = 13 is the only instance of a(n) = prime(n) in the sequence. - _Alonso del Arte_, Jul 27 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | Armand Turpel (armandt(AT)unforgettable.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001313 | OEIS-A034007

First differences of A045891.

Conjecture 1: For compositions of n+k-1, a(n) is the number of runs of 1 of length k. Example: Among the compositions of 4+2-1 = 5, there are a(4) = 4 runs of two 1's: 3,[1,1]; [1,1],3; 1,2,[1,1] and [1,1],2,1. - _Gregory L. Simay_, Feb 18 2018 Conjecture 2: More generally, let R(n,m,k) = the number of runs of k m's in all compositions of n. Then R(n,m,k) = A045623(n-m*k) - 2*A045623(n-m*(k+1)) + A045623(n-m*(k+2)). For example, R(7,1,1) = A045623(6) - 2*A045623(5) + A045623(4) = 144 - 2*64 + 28 = 44 = a(7). - _Gregory L. Simay_, Feb 20 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001314 | OEIS-A034015

Small 3-Schroeder numbers: a(n) = Sum_{k=0..n} 2^k*binomial(n, k)*binomial(2*n+2, k)/(k + 1).

a(n) = Sum_{i=0..n} Sum_{j=0..i} (-2)^(n-i)*binomial(i, j)*binomial(2*i+j, n)*binomial(n+1, i)/(n + 1) (conjectured). - _Michael D. Weiner_, May 25 2017 Weiner's conjecture is true - see (Theorem 3.8 of) the linked preprint by Easwar. - _Rohun Easwar_, Aug 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001315 | OEIS-A034097

Fractional part of square root of a(n) starts with digit 1.

For any a(n) in this sequence, letting k = integer part of sqrt(a(n)), a(n) belongs to a run of round(1/25 + 1/5 k) consecutive terms that differ by one (conjectured). E.g. round(1/25 + 1/5 floor(sqrt(330))) =4 and 330 is in the run 328, 329, 330, 331, consecutive terms of the series differing by one. - _Dimitri Papadopoulos_, Oct 28 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Sep 15 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001316 | OEIS-A034108

Fractional part of square root of a(n) starts with 2: first term of runs.

Conjectures from _Chai Wah Wu_, Apr 14 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Sep 15 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001317 | OEIS-A034171

Related to triple factorial numbers A007559(n+1).

Working with an offset of 1, we conjecture a(p*n) = a(n) (mod p^2) for prime p = 1 (mod 3) and all positive integers n except those n of the form n = m*p + k for 0 <= m <= (p-1)/3 and 1 <= k <= (p-1)/3. Cf. A298799, A004981 and A004982. - _Peter Bala_, Dec 23 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001318 | OEIS-A034175

a(n) is minimal such that a(n)+a(n-1) is a square and a(n) is not in {a(0), ..., a(n-1)}.

Conjectured to be a permutation of the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, Oct 01 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001319 | OEIS-A034267

a(n) = f(n,n) where f is given in A034261.

Conjecture D-finite with recurrence -(n+2)*(11*n-7)*a(n) + 2*(23*n^2+44*n+30)*a(n-1) - 4*(n+5)*(2*n-3)*a(n-2) = 0. - _R. J. Mathar_, Feb 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001320 | OEIS-A034296

Number of flat partitions of n: partitions {a_i} with each |a_i - a_{i-1}| <= 1.

Conjecture: Also the sum of the smallest parts in the distinct partitions of n with an odd number of parts. - _George Beck_, May 06 2017 Above conjecture was proved by Shane Chern, see link. - _George Beck_, Aug 12 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Erich Friedman_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001321 | OEIS-A034383

Number of labeled groups.

Also by definition, a(n) >= A000001(n)*n!/A059773(n). If the conjecture A059773(2^r) = A002884(r) is true, then A059773(2^r) <= 2^(r^2), while A000001(2^r) >= 2^((2/27)*r^2*(r-6)) (see the Math Stack Exchange link below), so a(2^r)/A034381(2^r) tends to infinity quickly as r tends to infinity. The sequence is strictly increasing for the first 256 terms (a(256) > A034381(256) > A034381(255) = a(255) since 255 is in A003277). On the other hand, assuming that A059773(2^r) = A002884(r), then a(2^20)/(2^20)! >= A000001(2^20)/A002884(20) > 99798.4, while a(2^20+1)/(2^20)! = A034381(2^20+1)/(2^20)! = (2^20+1)/phi(2^20+1) since 2^20+1 = 17*61681 is in A003277, so we would have a(2^20) > a(2^20+1). It is conjectured a(2^r) > a(2^r+1) for all sufficiently large r. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001322 | OEIS-A034444

a(n) is the number of unitary divisors of n (d such that d divides n, gcd(d, n/d) = 1).

Conjecture: For any nonnegative integer k and positive integer n, the sum of the k-th powers of the unitary divisors of n is divisible by the sum of the k-th powers of the odd unitary divisors of n (note that this sequence lists the sum of the 0th powers of the unitary divisors of n). - _Ivan N. Ianakiev_, Feb 18 2018 The above conjecture is true. If n is odd, the asserted divisibility is equality. Otherwise, let n = m*(2^b), where b > 0 and m is odd. The set of unitary divisors of n is equal to the disjoint union of U(m) and {d*(2^b): d in U(m)}, where U(m) denotes the set of unitary divisors of m (all of which are odd). Hence the sum of the k-th powers of the unitary divisors of n = Sum_{d in U(m)} d^k + Sum_{d in U(m)} (d*2^b)^k = (1 + 2^(b*k)) * Sum_{d in U(m)} d^k = (1 + 2^(b*k)) * the sum of the k-th powers of the odd unitary divisors of n. - _Peter Bala_, May 20 2026 [Corrected by _David Radcliffe_, Jul 20 2026] Conjecture: Let b(i; n), n > 0, be multiplicative sequences for some fixed integer i >= 0 with b(i; p^e) = (Sum_{k=1..i+1} A164652(i, k) * e^(k-1)) * (i+2) / (i!) for prime p and e > 0. Then we have Dirichlet generating functions: Sum_{n > 0} b(i; n) / n^s = (zeta(s))^(i+2) / zeta((i+2) * s). Examples for i=0 this sequence, for i=1 A226602, and for i=2 A286779. - _Werner Schulte_, Feb 17 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001323 | OEIS-A034496

Sum of n-th powers of divisors of 8.

Conjecture: No primes in this sequence (checked for first 10000 terms). - _Artur Jasinski_, Sep 23 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001324 | OEIS-A034602

Wolstenholme quotient W_p = (binomial(2p-1,p) - 1)/p^3 for prime p=A000040(n).

The values of this sequence's terms are replicated by conjectured general formula, given in A223886 (and also added to the formula section here) for k=2, j=1 and n>=3. - _Alexander R. Povolotsky_, Apr 18 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001325 | OEIS-A034693

Smallest k such that k*n+1 is prime.

Conjecture: for every n > 1 there exists a number k < n such that n*k + 1 is a prime. - _Amarnath Murthy_, Apr 17 2001 A stronger conjecture: for every n there exists a number k < 1 + n^(.75) such that n*k + 1 is a prime. I have verified this up to n = 10^6. Also, the expression 1 + n^(.74) does not work as an upper bound (counterexample: n = 19). - _Joseph L. Pe_, Jul 16 2002 Stronger version of the conjecture verified up to 10^9. - _Mauro Fiorentini_, Jul 23 2023 Conjecture: a(n) = O(log(n)*log(log(n))). - _Thomas Ordowski_, Oct 17 2014 I conjecture the opposite: a(n) / (log n log log n) is unbounded. See A194945 for records in this sequence. - _Charles R Greathouse IV_, Mar 21 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001326 | OEIS-A034807

Triangle T(n,k) of coefficients of Lucas (or Cardan) polynomials.

Conjecture: If a(n) = H(a,b,c,d,n) is a second-order linear recurrence with constant coefficients defined as a(0) = a, a(1)= b, a(n) = c*a(n-1) + d*a(n-2) then a(m*n) = H(a, H(a,b,c,d,m), Sum_{k=0..floor(m/2)} T(m,k)*c^(m-2*k)*d^k, (-1)^(m+1)*d^m, n) (Wolfdieter Lang). - _Gary Detlefs_, Feb 06 2023 For the proof of the preceding conjecture see the Detlefs and Lang link. There also proofs for several properties of this table are found. - _Wolfdieter Lang_, Apr 25 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001327 | OEIS-A034836

Number of ways to write n as n = x*y*z with 1 <= x <= y <= z.

Conjecture: a(n) = (np + 3*(nsq - ntp) + 5*ntp)/6 = (np + 3*nsq + 2*ntp)/6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Erich Friedman_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001328 | OEIS-A034863

a(n) = n!*(4*n^3 - 30*n^2 + 40*n + 3)/24.

Conjecture: (-96*n + 27161)*a(n) + (96*n^2 - 99580*n + 199917)*a(n-1) +(72707*n - 61983)*(n-1)*a(n-2) = 0. - _R. J. Mathar_, Apr 03 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001329 | OEIS-A034878

Numbers k such that k! can be written as the product of smaller factorials.

Except for the numbers 2, 9 and 10 this sequence is conjectured to be the same as A001013. By Murthy's trick, if k>2 is a product of factorials then k is a term. So half of the above conjecture is true: A001013 is a subsequence except for the number 2. - _Jonathan Sondow_, Nov 08 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Erich Friedman_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001330 | OEIS-A034917

Minimal determinant (negated) of n X n persymmetric matrix with entries {-1,0,+1}.

Conjectures: These minimal determinant matrices have no zero entries; a(10) is 73728, a(11) = 327680 and a(12) = 2097152. - _Jean-François Alcover_, Dec 15 2017 If Alcover's conjecture above is true, a(13) = 14929920, a(14) = 68853760, a(15) = 390905856. - _Pontus von Brömssen_, Jan 30 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Fred Lunnon_, Dec 11 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001331 | OEIS-A034918

Maximal determinant of n X n persymmetric matrix with entries {-1,0,+1}.

Conjectured: a(10) = 55296, a(11) = 327680, a(12) = 2097152. - _Jean-François Alcover_, Dec 16 2017 Conjecture: For n != 2, the maximal determinant is achieved by a matrix with entries {-1, +1} only. - _Jeffrey Shallit_, Jan 16 2026 If Shallit's conjecture above is true, a(13) = 14929920, a(14) = 53747712, and a(15) = 390905856. - _Pontus von Brömssen_, Jan 30 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Fred Lunnon_, Dec 11 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001332 | OEIS-A034950

Expansion of eta(8z)*eta(16z)*theta_3(2z).

Conjecture: Sum_{k=1..n} a(k)^2 ~ c * n^(3/2), where c = 0.787... - _Vaclav Kotesovec_, Sep 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001333 | OEIS-A035014

a(n) contains n digits (either '3' or '4') and is divisible by 2^n.

Proof of conjecture that a(n) ends with a(n-1): If a(n) is divisible by 2^n, then a(n) is divisible by 2^(n-1), so a(n)-k*10^(n-1) is divisible by 2^(n-1) for integer k, but if k is first digit of a(n) then a(n)-k*10^(n-1) is an (n-1)-digit

number made up of 3s and 4s and divisible by 2^(n-1) and so must be a(n-1). - _Henry Bottomley_, Feb 14 2000

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001334 | OEIS-A035050

a(n) is the smallest k such that k*2^n + 1 is prime.

Conjecture: for n > 0, a(n) = k < 2^n, so k*2^n + 1 is a Proth prime A080076. - _Thomas Ordowski_, Apr 13 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001335 | OEIS-A035062

Numbers n such that 2^n does not contain the digit 7.

It is conjectured that 71 is the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Nov 15 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001336 | OEIS-A035137

Numbers that are not the sum of 2 palindromes (where 0 is considered a palindrome).

A261675(a(n)) >= 3 (and, conjecturally, = 3). - _N. J. A. Sloane_, Sep 03 2015 The conjecture that every number is the sum of 3 palindromes fails iff there is a term a(n) such that for all palindromes P < a(n), the difference a(n) - P is also a term of this sequence. - _M. F. Hasler_, Sep 08 2015 Cilleruelo and Luca (see links) have proved the conjecture that every positive integer is the sum of at most three palindromes (in bases >= 5), and also that the density of those that require three is positive. - _Christopher E. Thompson_, Apr 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Nov 15 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001337 | OEIS-A035263

Trajectory of 1 under the morphism 0 -> 11, 1 -> 10; parity of 2-adic valuation of 2n: a(n) = A000035(A001511(n)).

A conjectured identity relating to the partition sequence, A000041: p(x) = A(x) * A(x^2) when A(x) = the Euler transform of A035263 = polcoeff A174065: (1 + x + x^2 + 2x^3 + 3x^4 + 4x^5 + ...). - _Gary W. Adamson_, Mar 21 2010 A conjectured identity relating to the partition sequence, A000041 as polcoeff p(x); A003159, and its characteristic function A035263: (1, 0, 1, 1, 1, 0, 1, 0, 1, 0, 1, 1, 1, ...); and A036554 indicating n-th terms with zeros in A035263: (2, 6, 8, 10, 14, 18, 22, ...). The conjecture states that p(x) = A(x) = A(x^2) when A(x) = polcoeffA174065 = the Euler transform of A035263 = 1/(1-x)*(1-x^3)*(1-x^4)*(1-x^5)*... = (1 + x + x^2 + 2x^3 + 3x^4 + 4x^5 + ...) and the aerated variant = the Euler transform of the complement of A035263: 1/(1-x^2)*(1-x^6)*(1-x^8)*... = (1 + x^2 + x^4 + 2x^6 + 3x^8 + 4x^10 + ...). The conjecture above was proved by _Jean-Paul Allouche_ on Dec 21 2013.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Karamanos Konstantinos_, _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001338 | OEIS-A035312

Zorach additive triangle, read by rows.

Conjectured to form a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Zorach_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001339 | OEIS-A035313

(Largest) diagonal of the Zorach additive triangle A035312.

Conjecturally, every positive integer occurs in the sequence or one of its n-th differences, which would imply that the sequence and its n-th differences partition the positive integers. - _David W. Wilson_, Feb 26 2012 Conjecture: lim(n->infinity, a(n+1)/a(n)) = 2. - _David W. Wilson_, Feb 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Zorach_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001340 | OEIS-A035319

Number of rooted maps of genus n with one vertex and one face; the maps are considered on orientable surfaces and contain 2n edges.

It appears that this is given by the formula (4n)!/2^{2n}(2n+1)! = (4n-1)!!/(2n+1). (This sequence arose -- conjecturally, but it shouldn't be too hard to make it rigorous -- as the unique nontrivial Betti number of a certain poset associated to the hyperoctahedral group.) - Eric M. Rains (rains(AT)caltech.edu), Jan 24 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001341 | OEIS-A035337

Third column of Wythoff array.

The formula a(n) = 3*A003622(n)-n+1 = 3AA(n)-n+1 conjectured by Layman below is correct, since it is well known that AA(n)+1 = B(n) = A(n)+n, where B = A001950, and so 3AA(n)-n+1 = 3B(n)-n-2 = 3A(n)+2n-2. - _Michel Dekking_, Aug 31 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _J. H. Conway_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001342 | OEIS-A035363

Number of partitions of n into even parts.

The above conjecture follows easily from the known value eta(i) = Gamma(1/4)/(2*Pi^(3/4)), where eta(t) is the Dedekind eta function. - _Peter Bala_, Jun 09 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001343 | OEIS-A035378

Coefficients in expansion of Sum_{k>=0} Product_{j=1..k} (1-x^j) about x = -1.

Conjectures: a(5*n+3) == 0 (mod 5), a(5*n+4) == 0 (mod 5) and a(5*n+2) - 2*a(5*n+1) == (0 mod 5) (all checked up to n = 49). - _Peter Bala_, Jun 19 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bill Gosper_, Aug 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001344 | OEIS-A035485

Card on top of deck at n-th stage of R. K. Guy's shuffling problem.

It is conjectured that eventually every number appears on top of the deck.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001345 | OEIS-A035486

Kimberling's expulsion array read by antidiagonals.

It is conjectured since 1992 that the main diagonal elements (A007063) are a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001346 | OEIS-A035513

Wythoff array read by falling antidiagonals.

Conjecture: If m >= 2, then {(T(n,1), T(n,2)) mod m} has cardinality m^2. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001347 | OEIS-A035615

Number of winning length n strings with a 2-symbol alphabet in "same game".

G.f.: x(2x^6 - 6x^5 + 8x^4 + 2x^3 - 6x^2 + 2x)/[(1 - x^2)(1 - 2x)(1 - x - x^2)^2] (conjectured). - _Ralf Stephan_, May 11 2004. Established by Burns and Purcell - see link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Erich Friedman_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001348 | OEIS-A035617

Number of winning length n strings with a 3-symbol alphabet in "same game".

For binary strings, the formula for the number of winning strings of length n has been conjectured by Ralf Stephan and proved by Burns and Purcell (2005, 2007). For b-ary strings with b >= 3, the same problem seems to be unsolved. - _Petros Hadjicostas_, Dec 27 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Erich Friedman_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001349 | OEIS-A036012

a(n) = smallest number > 1 such that a(1)a(2)...a(n) + 1 is prime.

Conjecture: a(n) = O(n). - _Thomas Ordowski_, Aug 08 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Russell Easterly_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001350 | OEIS-A036038

Triangle T(n,k) read by rows: multinomial coefficients for the partitions of n listed in Abramowitz-Stegun order, 1 <= k <= A000041(n).

The conjecture on the Jack polynomials of zero order is true as evident from equation a) on p. 80 of the Stanley reference, suggested to me by Steve Kass. The conventions for denoting the more general Jack polynomials J(n,alpha) vary. Using Stanley's convention, these Jack polynomials are the umbral extensions of the multinomial expansion of (s_1*x_1 + s_2*x_2 + ... + s_(n+1)*x_(n+1))^n in which the subscripts of the (s_k)^j in the symmetric monomial expansions are finally ignored and the exponent dropped to give s_j(alpha) = j-th row polynomial of A094638 or |A008276| in ascending powers of alpha. (The MOPS table has some inconsistency between n = 3 and n = 4.) - _Tom Copeland_, Nov 26 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001351 | OEIS-A036229

Smallest n-digit prime containing only digits 1 or 2 or -1 if no such prime exists.

It is conjectured that such a prime always exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _G. L. Honaker, Jr._ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001352 | OEIS-A036263

Second differences of primes.

Conjecture: |a(1)| + |a(2)| + ... + |a(n)| ~ prime(n). - _Thomas Ordowski_, Jul 21 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001353 | OEIS-A036284

Periodic vertical binary vectors of Fibonacci numbers.

Conjecture: For n>=1, each term a(n), when considered as a GF(2)[X]-polynomial, is divisible by GF(2)[X] -polynomial (x^3 + 1) ^ A000225(n-1). If this holds, then for n>=1, a(n) = A048720bi(A136380(n),A048723bi(9,A000225(n-1))). Conjecture 2: there is also one extra (x^1 + 1) factor present, see A136384.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 01 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001354 | OEIS-A036447

Double and reverse digits.

Conjecture: a(n)^(1/n) tends to sqrt(10). - _Vaclav Kotesovec_, Jan 03 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rodolfo Kurchan_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001355 | OEIS-A036462

Conjecturally, a power of 2 written in base 3 cannot have this many 0's.

Conjecturally, a power of 2 written in base 3 cannot have this many 0's. Conjecture 1: Sequence A104320 never obtains the values in this sequence, so A104321(a(n)) is undefined. Conjecture 2: This sequence is infinite. - _David W. Wilson_, Oct 21 2016, edited by _Antti Karttunen_, Oct 29 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001356 | OEIS-A036463

Conjecturally, a power of 2 written in base 3 cannot have this many 2's.

Conjecturally, a power of 2 written in base 3 cannot have this many 2's.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001357 | OEIS-A036468

Number of ways to represent 2n+1 as a+b with 0 < a < b and a^2 + b^2 prime.

This open problem is mentioned by Guy at the end of section C1. - _T. D. Noe_, Apr 22 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001358 | OEIS-A036492

Offsets for the Atkin Partition Congruence theorem.

The Atkin Theorem, inspired by a famous conjecture of Ramanujan, gives congruences properties of certain partition numbers, generalizing many previous results.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001359 | OEIS-A036496

Number of lines that intersect the first n points on a spiral on a triangular lattice. The spiral starts at (0,0), goes to (1,0) and (1/2, sqrt(3)/2) and continues counterclockwise.

Conjecture: a(n) is half the minimal perimeter of a polyhex of n hexagons. - Winston C. Yang (winston(AT)cs.wisc.edu), Apr 06 2002. This conjecture follows from the Brunvoll et al. reference. - _Sascha Kurz_, Mar 17 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | Mario VELUCCHI (mathchess(AT)velucchi.it) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001360 | OEIS-A036559

Squared distances that cannot arise in any Barlow packing of spheres of diameter sqrt(3) (the scale being the simplest one that makes this an integer sequence).

Conjectured to consist of 4^n times 1 or 7 or 13 or 16k+10 together with 2, 5, 14, 23, 38, 230.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. H. Conway_, _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001361 | OEIS-A036766

Number of ordered rooted trees with n non-root nodes and all outdegrees <= four.

Conjecture: -3*(3*n+2)*(133*n-347)*(3*n+4)*(n+1)*a(n) +(111457*n^4-364730*n^3+228995*n^2+19310*n-33312)*a(n-1) +5*(-68503*n^4+34661

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001362 | OEIS-A036820

Number of partitions satisfying (cn(2,5) = cn(3,5) = 0).

It appears that this sequence is related to the generalized heptagonal numbers A085787 in the same way as the partition numbers A000041 are related to the generalized pentagonal numbers A001318. (See the table in comments section of A195825.) Conjecture: Column 1 of triangle A195837. Also 1 together with the row sums of triangle A195837. Also column 3 of the square array A195825. - _Omar E. Pol_, Oct 08 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001363 | OEIS-A036837

Schoenheim bound L_1(n,n-5,n-6).

Conjecture: G.f.: x^7*(4-x^9+5*x^8-10*x^7+10*x^6-5*x^5+2*x^4-5*x^3+10*x^2-9*x) / ((x^4+x^3+x^2+x+1)*(x-1)^6). - Maksym Voznyy (voznyy(AT)mail.ru), Jul 28 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001364 | OEIS-A036840

a(n) is the average of the repeating terms of {T(n,k)} rounded to the nearest integer (rounding up if there's a choice), if {T(n,k)} is eventually periodic; = 0 otherwise. Here T(n,k) is the "phi/sigma tug-of-war sequence with seed n" defined by T(n,1) = phi(n), T(n,2) = sigma(phi(n)), T(n,3) = phi(sigma(phi(n))), ..., T(n,k) = phi(T(n,k-1)) if k is odd and = sigma(T(n,k-1)) if k is even.

Conjecture: a(n) is never 0; i.e. the sequence {T(n,k)} is eventually periodic for every n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Jan 09 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001365 | OEIS-A036845

a(n) = min_{k} {T(n,k)} where T(n,k) is the "phi/sigma tug-of-war sequence with seed n" defined by T(n,1) = phi(n), T(n,2) = sigma(phi(n)), T(n,3) = phi(sigma(phi(n))), ..., T(n,k) = phi(T(n,k-1)) if k is odd and = sigma(T(n,k-1)) if k is even.

Conjecture: The sequence {T(n,k)} is eventually periodic for every n, so a(n) can be computed in finite time. Conjecture: a(n) -> infinity as n -> infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Jan 09 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001366 | OEIS-A036937

Smallest n-digit prime containing only digits 2 and 3.

It is conjectured that such a prime exists for every value of n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Jan 04 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001367 | OEIS-A036968

Genocchi numbers (of first kind): expansion of 2*x/(exp(x)+1).

Conjecture: For any positive integer n, -a(n+1) is the permanent of the n X n matrix M with M(j, k) = floor((2*j - k)/n), (j,k=1..n). - _Zhi-Wei Sun_, Sep 07 2021 A corresponding conjecture can also be made for L. Seidel's 'Genocchi numbers of second kind' A005439. - _Peter Luschny_, Sep 08 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001368 | OEIS-A036983

Number of invariants in Hilbert basis for binary forms of degree n.

Extending this is related to a "gorgeous open question of great antiquity" [Towber].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001369 | OEIS-A037000

Positions of the digit '1' in the decimal expansion of Pi.

Conjecturally, a(n) ~ 10n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Nicolau C. Saldanha (nicolau(AT)mat.puc-rio.br) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001370 | OEIS-A037018

Conjecturally, largest k with no prime in [k, k(1+1/n)].

Conjecturally, largest k with no prime in [k, k(1+1/n)].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001371 | OEIS-A037053

Smallest prime containing exactly n 0's.

Certain entries require n+3 digits such as a(13). Those which do not require three digits besides n zeros are in A085824. Conjecture: a prime of this form, A037053, requires at most three judiciously placed nonzero digits, two on the exterior and one inside. - _Robert G. Wilson v_ Conjecture 1: (following _Robert G. Wilson v_'s conjecture above and insight provided by Hans Havermann): terms with two substrings of zeros have first digit of 1. Conjecture 2: as n increases, there are more terms with two substrings of zeros than one consecutive string.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Jan 04 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001372 | OEIS-A037055

Smallest prime containing exactly n 1's.

For n > 1, A037055 is conjectured to be identical to A084673. - _Robert G. Wilson v_, Jul 04 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Jan 04 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001373 | OEIS-A037057

Smallest prime containing exactly n 2's.

For n > 1, the last digit cannot be 2, so a(n) has at least n+1 digits. The probability is big that none of [10^n/9]*20 + {1,3,7,9} is prime, in which case a(n) must have at least n+2 digits. This is the most frequent case. We can even conjecture that for all n > 1, a(n) equals [10^(n+1)/9]*20 + b with 1 <= b <= 9 and one of the (first) digits 2 replaced by 0 or 1. - _M. F. Hasler_, Feb 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Jan 04 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001374 | OEIS-A037059

Smallest prime containing exactly n 3's.

For almost all n >= 0, a(n) equals [10^(n+1)/3] with one of the (first) digits 3 replaced by a digit 1 or 2. We conjecture that in the few other cases (e.g., for n = 12, 119, ...) the statement holds with some digit 3 replaced by a digit among {4, 5, 7, 8}, except for the special case a(1) = 3. - _M. F. Hasler_, Feb 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Jan 04 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001375 | OEIS-A037061

Smallest prime containing exactly n 4's.

The last digit of n cannot be 4, therefore a(n) must have at least n+1 digits. It is probable that none among [10^n/9]*40 + {1,3,7,9} is prime in which case a(n) must have n+2 digits. We conjecture that for all n >= 0, a(n) equals [10^(n+1)/9]*40 + b with 1 <= b <= 9 and one of the (first) digits 4 replaced by a 0, 1, 2 or 3. - _M. F. Hasler_, Feb 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Jan 04 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001376 | OEIS-A037063

Smallest prime containing exactly n 5's.

For n > 1, the last digit of n cannot be 5, therefore a(n) must have at least n+1 digits. It is probable that none among [10^n/9]*50 + {1,3,7,9} is prime in which case a(n) must have n+2 digits. We conjecture that for all n > 1, a(n) equals [10^(n+1)/9]*50 + b with 1 <= b <= 9 and one of the (first) digits 5 replaced by a 0, 1, 2, 3 or 4. - _M. F. Hasler_, Feb 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Jan 04 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001377 | OEIS-A037065

Smallest prime containing exactly n 6's.

The last digit of n cannot be 6, therefore a(n) must have at least n+1 digits. It is probable that none among [10^n/9]*60 + {1,3,7,9} is prime in which case a(n) must have n+2 digits. We conjecture that for all n >= 0, a(n) equals [10^(n+1)/9]*60 + b with 1 <= b <= 9 and one of the (first) digits 6 replaced by a digit among {0, 1, 2, 3, 4, 5}. - _M. F. Hasler_, Feb 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Jan 04 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001378 | OEIS-A037067

Smallest prime containing exactly n 7's.

We conjecture that for all n >= 2, a(n) equals floor(10^(n+1)/9)*7 with one of the (first) digits 7 replaced by a digit among {0, ..., 6}. - _M. F. Hasler_, Feb 22 2016 The conjecture is false: a(668) = 7*(10^669-1)/9 + 10^276. - _Robert Israel_, Jul 13 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Jan 04 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001379 | OEIS-A037069

Smallest prime containing exactly n 8's.

The last digit of n cannot be 8, therefore a(n) must have at least n+1 digits. It is probable (using [] for floor) that none among [10^n/9]*80 + {1,3,7,9} is prime in which case a(n) must have n+2 digits. We conjecture that for all n >= 0, a(n) equals [10^(n+1)/9]*80 + b with 1 <= b <= 9 and one of the (first) digits 8 replaced by a digit among {0, ..., 7}. - _M. F. Hasler_, Feb 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Jan 04 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001380 | OEIS-A037071

Smallest prime containing exactly n 9's.

We conjecture that for all n >= 0, a(n) equals [10^(n+1)/9]*9 with one of the (first) digits 9 replaced by a digit among {1, 2, 4, 5, 7, 8}. - _M. F. Hasler_, Feb 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Jan 04 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001381 | OEIS-A037084

Positive integers not going to 1 under iterations of the map in A001281: n->3n-1 if n odd, n->n/2 if n even.

Conjecture : for any x, the iterated process "x ->3x-1" if x is odd or "x ->x/2" if x is even leads to one of the following three cycles: (1, 2), (5, 14, 7, 20, 10), (41, 122, 61, 182, 91, 272, 136, 68, 34, 17, 50, 25, 74, 37, 110, 55, 164, 82). - _Benoit Cloitre_, May 14 2002 Complement (in N*) of A039500 ; union of A039501 and A039502 (conjectured). - _M. F. Hasler_, Nov 26 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | Robert W. Craigen (craigen(AT)fresno.edu) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001382 | OEIS-A037096

Periodic vertical binary vectors computed for powers of 3: a(n) = Sum_{k=0 .. (2^n)-1} (floor((3^k)/(2^n)) mod 2) * 2^k.

Conjecture: For n >= 3, each term a(n), when considered as a GF(2)[X] polynomial, is divisible by the GF(2)[X] polynomial (x + 1) ^ A055010(n-1). If this holds, then for n >= 3, a(n) = A048720(A136386(n), A048723(3,A055010(n-1))).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 29 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001383 | OEIS-A037097

Periodic vertical binary vectors of powers of 3, starting from bit-column 2 (halved).

Conjecture: For n>=3, each term a(n), when considered as a GF(2)[X]-polynomial, is divisible by GF(2)[X] -polynomial (x + 1) ^ A000225(n-2) (= A051179(n-2)). If this holds, then for n>=3, a(n) = A048720bi(A136386(n), A048723bi(3, A000225(n-2))) = A048720bi(A136386(n), A051179(n-2)). [corrected by _Qiyuan Gu_, Jul 24 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 29 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001384 | OEIS-A037197

Numbers k such that tau(sigma(k)) = tau(k) where tau(k) is the number of divisors of k and sigma(k) their sum.

Conjecture: for n > 10^6, a(n) < n^2. - _Benoit Cloitre_, Aug 24 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001385 | OEIS-A037223

Number of solutions to non-attacking rooks problem on n X n board that are invariant under 180-degree rotation.

Conjecture: a(n) = Product_{i=1..n and phi(i)<=floor(i/2)} i. - _Enrique Pérez Herrero_, May 31 2012. This conjecture is WRONG, counterexample is n=105. - _Vaclav Kotesovec_, Sep 07 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | Miklos SZABO (mike(AT)ludens.elte.hu) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001386 | OEIS-A037225

a(n) = phi(2n+1).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001387 | OEIS-A037250

a(n) = n^2*(n^2 + 1)*(n-1).

Conjecture: satisfies a linear recurrence having signature (6, -15, 20, -15, 6, -1). - _Harvey P. Dale_, Jul 27 2019 This conjecture is true since for any series a(n) = P(n) (P polynomial in n of degree d) there is an o.g.f. Q(x)/(1-x)^(d+1). - _Georg Fischer_, Feb 17 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001388 | OEIS-A037281

Number of iterations of transformation in A037280 needed to reach 1 or a prime, or -1 if no such number exists.

Continuation from a(12): -1, 0, 3, 4, 1, 0, -1, 0, -1, 1, 1, 0, -1, 1, 4, 2, -1, 0, -1, 0, 2, 1, -1, 3, -1, 0, 2, 1, -1, 0, -1, 0, -1, -1, 1, 0, -1, 1, -1, 1, -1, 0, 1, 2, -1, 2, 1, 0, -1; where all the -1's are conjectural. - _Franklin T. Adams-Watters_, Jul 28 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001389 | OEIS-A037289

Number of commutative rings with n elements.

This sequence is multiplicative. See the reference "The Numbers of Small Rings" below, which proves the result for all rings; restricting to commutative rings only makes the proof easier. - Conjecture by _Mitch Harris_, Apr 19 2005, proof found by _Franklin T. Adams-Watters_, Jul 10 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, Jun 15 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001390 | OEIS-A037301

Numbers whose base-2 and base-3 expansions have the same digit sum.

a(5) = 10, a(6) = 11, a(7) = 12 and a(8) = 13 is the first time that four consecutive terms appear in this sequence. Conjecture: There is no occurrence of five or more consecutive terms of a(n). Tested by exhaustive search up to a(n) = 3^29. - _Thomas König_, Aug 15 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001391 | OEIS-A037958

a(n) = binomial(n, floor((n-8)/2)).

Conjecture: -(n+9)*(n-8)*a(n) +2*(n)*a(n-1) + 4*n*(n-1)*a(n-2) = 0. - _R. J. Mathar_, Jul 26 2015, verified by _Robert Israel_, Jun 28 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001392 | OEIS-A037988

Critical values in Conway's game of one-dimensional phutball.

Conjectured partial sums of A272729 (true for at least 8192 terms). - _Sean A. Irvine_, Jun 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Erich Friedman_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001393 | OEIS-A038261

First line of 5-wave sequence A038201, also bisection of A006358.

Conjecture: GF = -(x-1)*(x^3-6*x^2+9*x-1) / (-1+15*x-35*x^2+28*x^3-9*x^4+x^5). - _Wouter Meeussen_, Mar 19 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Floor van Lamoen_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001394 | OEIS-A038348

Expansion of (1/(1-x^2))*Product_{m>=0} 1/(1-x^(2m+1)).

Conjecture: The n-th derivative of Gamma(x+1) at x = 0 has a(n+1) terms. For example, d^4/dx^4_(x = 0) Gamma(x+1) = 8*eulergamma*zeta(3) + eulergamma^4 + eulergamma^2*Pi^2 + 3*Pi^4/20 which has a(5) = 4 terms. - _David Ulgenes_, Dec 05 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001395 | OEIS-A038360

Ranks of certain relations among Euler sums of weight n.

It is conjectured that this is (apart from offset) the same as A216714.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001396 | OEIS-A038396

Concatenate first n even numbers in reverse order.

F. Smarandache conjectures that a(n) is never a perfect power (see link sequence 163 page 80). - _Michel Marcus_, Mar 03 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | M. I. Petrescu (mipetrescu(AT)yahoo.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001397 | OEIS-A038460

Maximal value of difference between successive primes among numbers < 10^n.

For 3 < n < 19, a(n) <= 6 (n - 1)(n - 2). Conjecture: for any n > 3, a(n) <= 6 (n - 1)(n - 2). Let q = 6 (n - 1)(n - 2) and d = (10^n) - (10^(n/2) - 1)^2. Since for any even n, d is the smallest difference between two consecutive squares of the form a^2 - b^2, where a^2 = 10^n, b = a - 1, for any even n > 2, d > 5q (where 3q is, according to the conjecture, not less than the sum of the three largest gaps between 4 consecutive primes p1...p4, or 3 * a(n), and 2q is, respectively, not less than the sum of the two largest gaps (p1 - p0) + (p5 - p4), or 2 * a(n)). In the same way, we can state that for any odd n > 3, if a^2 is the smallest square such that a^2 has (n+1) digits, b = a - 1, and d = a^2 - b^2, then d > 5q. The correctness of the above conjecture would establish the well-known Brocard's and Legendre's conjectures (see the link below for both definitions), since they are proved for the first 10000 primes. - _Sergey Pavlov_, Jan 30 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enoch Haga_, Dec 11 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001398 | OEIS-A038505

Sum of every 4th entry of row n in Pascal's triangle, starting at binomial(n,2).

Conjecture: 2*a(n+2) = A038504(n+2) + A000749(n+2) + 2*A009545(n). - _Creighton Dement_, May 22 2005 Proof of the conjecture by _Creighton Dement_ (May 22 2005): using the first formula of Theorem 1 in [Shevelev link] for n=4, omega=i=sqrt(-1), i:=1,2,3,4, m:=n>=1, we have

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank Ruskey_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001399 | OEIS-A038554

Derivative of n: write n in binary, replace each pair of adjacent bits with their mod 2 sum (a(0)=a(1)=0 by convention). Also n XOR (n shift 1).

a(0) = a(1) = 0, a(4n) = 2*a(2n), a(4n+2) = 2*a(2n+1)+1, a(4n+1) = 2*a(2n)+1, a(4n+3) = 2*a(2n+1). Proof by Nikolaus Meyberg following a conjecture by _Ralf Stephan_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001400 | OEIS-A038663

[n/F_2] + [n/F_3] + [n/F_4] +..., F_n=Fibonacci numbers.

Conjecture: a(n) ~ c * n, where c = A079586 - 1. - _Vaclav Kotesovec_, Aug 30 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | Antreas P. Hatzipolakis (xpolakis(AT)hol.gr) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001401 | OEIS-A038701

Prime powers q for which f(g(m(q))) = m(q), where f = A051703, g = A008475 and m = A003418.

There are no more prime powers in the list <= 199. Conjecture: The sequence is finite, i.e., f(g(m(q))) > m(q) for sufficiently great prime powers q.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, May 01 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001402 | OEIS-A038867

a(n) = (n+5)^3 - n^3.

Conjecture: There are no squares in this sequence, verified up to n = 100000. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeff Burch_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001403 | OEIS-A038991

Number of sublattices of index n in generic 4-dimensional lattice.

Conjectured g.f.: Sum_{k>=1} Sum {l>=1} Sum {m>=1} k*l^2*m^3*x^(k*l*m)/(1 - x^(k*l*m)) (by extension of g.f for A001001). - _Miles Wilson_, Apr 05 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001404 | OEIS-A039004

Numbers whose base-4 representation has the same number of 1's and 2's.

Conjecture: a(C(2n,n)-1) = 4^n - 1. (A000984 is C(2n,n)). - _Gerald McGarvey_, Nov 18 2007 We prove the McGarvey conjecture (A) a(e(n,n)-1) = 4^n-1, with e(n,m) = A034870(n,m) = binomial(2n,m), the even rows of Pascal's triangle. By the comment from Hendel in A034870, we have the function s(n,k) = #{n-digit, base-4 numbers with n-k more 1-digits than 2-digits}. As shown in A034870, (B) #s(n,k)= e(n,k) with # indicating cardinality, that is, e(n,k) = binomial(2n,k) gives the number of n-digit, base-4 numbers with n-k more 1-digits than 2-digits. We now show that (B) implies (A). By definition, s(n,n) contains the e(n,n) = binomial(2n,n) numbers with an equal number of 1-digits and 2-digits. The biggest n-digit, base-4 number is 333...3 (n copies of 3). Since 333...33 has zero 1-digits and zero 2-digits it follows that 333...333 is a member of s(n,n) and hence it is the biggest member of s(n,n). But 333...333 (n copies of 3) in base 4 has value 4^n-1. Since A039004 starts with index 0 (that is, 0 is the 0th member of A039004), it immediately follows that 4^n-1 is the (e(n,n)-1)st member of A039004, proving the McGarvey conjecture. (End) Conjecture: there is a constant c around 5 such that a(n) is asymptotic to c*n. - _Benoit Cloitre_, Nov 24 2002 That conjecture is false. The number of members of the sequence from 0 to 4^d-1 is binomial(2d,d) which by Stirling's formula is asymptotic to 4^d/sqrt(Pi*d). If Cloitre's conjecture were true we would have 4^d-1 asymptotic to c*4^d/sqrt(Pi*d), a contradiction. - _Robert Israel_, Jun 24 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001405 | OEIS-A039508

Conjecturally, number of minimum points of '3x+(2n+1)' problem.

Conjecturally, number of minimum points of '3x+(2n+1)' problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, Feb 15 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001406 | OEIS-A039515

Conjecturally, largest attractor in '3x+(2n+1)' problem.

Conjecturally, largest attractor in '3x+(2n+1)' problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, Feb 15 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001407 | OEIS-A039595

Positive Fibonacci numbers that are also triangular numbers.

Vern Hoggatt conjectured and Luo Ming confirmed that these are all the terms. - _Tomohiro Yamada_, Sep 23 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felice Russo_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001408 | OEIS-A039622

Number of n X n Young tableaux.

This is the factor g_n that appears in a conjectured formula for 2n-th moment of the Riemann zeta function on the critical line. (See Conrey articles.) - _Michael Somos_, Apr 15 2003 [Comment revised by _N. J. A. Sloane_, Jun 21 2016]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Floor van Lamoen_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001409 | OEIS-A039650

Prime reached by iterating f(x) = phi(x)+1 on n.

Sequence A229487 gives the conjectured largest number that converges to prime(n). - _T. D. Noe_, Oct 17 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001410 | OEIS-A039654

a(n) = prime reached by iterating f(x) = sigma(x)-1 starting at n, or -1 if no prime is ever reached.

Guy (2004) attributes this conjecture to Erdos. See Erdos et al. (1990). - _N. J. A. Sloane_, Aug 30 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001411 | OEIS-A039661

Decimal expansion of exp(Pi).

Conjecture: Equals Sum_{k>=0} A132136(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001412 | OEIS-A039678

Smallest number m > 1 such that m^(p-1)-1 is divisible by p^2, where p = n-th prime.

Using Fermat's little theorem twice, it is easy to see that m=p^2-1 solves this problem for all odd primes p. In fact, there appear to be exactly p-1 values of m with 1 <= m <= p^2 for which m^(p-1) == 1 (mod p^2). See A096082 for the related open problem. - _T. D. Noe_, Aug 24 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felice Russo_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001413 | OEIS-A039692

Jabotinsky-triangle related to A039647.

Conjectured row sums: sum_{m=1..n} a(n,m) = A005442(n). - _R. J. Mathar_, Jun 01 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001414 | OEIS-A039799

For S a subset of [n] = {1,2,3,...n}, let B_S = {x+y : x,y in S, x<y}; then a(n) is maximal cardinality of B_S intersect B_{[n]-S}.

Conjecture: a(n) = 2n-11 for n >= 14. - _Rob Pratt_, Jul 14 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Erich Friedman_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001415 | OEIS-A039825

a(n) = floor((n^2 + n + 8) / 4).

Conjecturally, apart from the first term, the sequence terms are the exponents in the expansion of Sum_{n >= 1} q^(3*n) * (Product_{k >= 2*n} 1 - q^k) = q^3 - q^5 - q^7 + q^9 + q^12 - q^16 - q^20 + + - - .... Cf. A054925. - _Peter Bala_, Apr 13 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001416 | OEIS-A039827

Number of different coefficient values in expansion of Product (1+q^i+q^(2i)), i=1 to n.

Conjectures from _Chai Wah Wu_, Jun 22 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001417 | OEIS-A039909

Largest coefficient in expansion of Product_{i=1..n} (1-q^1+q^2-...+(-q)^i).

Conjecture: a(n) ~ 6 * n^n / exp(n). - _Vaclav Kotesovec_, Jan 05 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001418 | OEIS-A039924

G.f.: Sum_{k>=0} x^(k^2)*(-1)^k/(Product_{i=1..k} 1-x^i).

a(n) = A286041(n) - A286316(n) (conjectured). - _George Beck_, May 05 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001419 | OEIS-A039986

Primes such that every distinct permutation of digits is composite (including permutations with leading zeros).

The sequence can be seen as a table with the n-digit terms in row n. Row lengths would then be (4, 13, 34, 45, 68, 67, 47, 36, 40, 46, 33, 45, 35, 38, 32, ...). In these rows there are (0, 0, 0, 6, 9, 3, 0, 1, 0, 0, ...) terms with >= 4 distinct digits: this seems to happen only for terms with 4, 5, 6 or 8 digits. I conjecture that there are no more than these 6 + 9 + 3 + 1 = 19 terms (2861, 4027, 4801, 5209, 5623, 5849, 24889, 26561, 40609, 40883, 66541, 66853, 85087, 85843, 86441, 288689, 442469, 558541, 55555429) with 4, and none with 5 or more distinct digits. - _M. F. Hasler_, Jul 01 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001420 | OEIS-A040027

The Gould numbers.

Conjecture: a(n) = Sum_{k = 0..n} A058006(k)*A048993(n+1, k+1) - _Velin Yanev_, Aug 31 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Gould_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001421 | OEIS-A040051

Parity of partition function A000041.

Now we know that among the first ten million values of p(n) 5002137 of them are odd. It is conjectured (T. R. Parkin and D. Shanks) that p(n) is equally often even and odd. Lower bound estimates for the number of times p(n) is even among the first N values of p(n) for any given N are known (Scott

Ahlgren; and Nicolas, Rusza and Sárközy among others). My 1966 conjecture that in every arithmetic progression r (mod s) for arbitrary integral r and s, there are infinitely many integers n for which p(n) is odd - with a similar statement for p(n) even - was proved for the even case by Ken Ono (1996) and for the odd case for all s up to 10^5 and for all s which are powers of 2 by Bolyan and Ono, 2002.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001422 | OEIS-A040076

Smallest m >= 0 such that n*2^m + 1 is prime, or -1 if no such m exists.

Sierpiński showed that a(n) = -1 infinitely often. John Selfridge showed that a(78557) = -1 and it is conjectured that a(n) >= 0 for all n < 78557.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001423 | OEIS-A041017

Denominators of continued fraction convergents to sqrt(12).

Also, values i where A227790(i)/i reaches a new maximum (conjectured). - _Ralf Stephan_, Sep 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001424 | OEIS-A041042

Numerators of continued fraction convergents to sqrt(27).

Subset of |A002316| (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001425 | OEIS-A041243

Denominators of continued fraction convergents to sqrt(133).

Conjecture: satisfies a linear recurrence having signature (0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 5177198, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, -1). - _Harvey P. Dale_, Dec 03 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001426 | OEIS-A042220

Numerators of continued fraction convergents to sqrt(636).

Conjecture: satisfies a linear recurrence having signature (0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 7011902, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, -1). - _Harvey P. Dale_, May 08 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001427 | OEIS-A042945

Numbers congruent to 2,3,6,11 mod 12 missing from A042944 (conjectured to be finite).

Numbers congruent to 2,3,6,11 mod 12 missing from A042944 (conjectured to be finite).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Allan Wilks_ and _Brian Galebach_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001428 | OEIS-A042963

Numbers congruent to 1 or 2 mod 4.

Equivalent to the following variation on Fermat's Diophantine m-tuple: 1 + the product of any two distinct terms is not a square; this sequence, which we'll call sequence S, is produced by the following algorithm. At the start, S is initially empty. At stage n, starting at n = 1, the algorithm checks whether there exists a number m already in the sequence, such that mn+1 is a perfect square. If such a number m is found, then n is not added to the sequence; otherwise, n is added. Then n is incremented to n + 1, and we repeat the procedure. Proof by Clark R. Lyons: We prove by strong induction that n is in the sequence S if and only if n == 1 (mod 4) or n == 2 (mod 4). Suppose now that this holds for all k < n. In case 1, either n == 1 (mod 4) or n == 2 (mod 4), and we wish to show that n does indeed enter the sequence S. That is, we wish to show that there does not exist m < n, already in the sequence at this point such that mn+1 is a square. By the inductive hypothesis m == 1 (mod 4) or m == 2 (mod 4). This means that both m and n are one of 1, 2, 5, or 6 mod 8. Using a multiplication table mod 8, we see that this implies mn+1 is congruent to one of 2, 3, 5, 6, or 7 mod 8. But we also see that mod 8, a perfect square is congruent to 0, 1, or 4. Thus mn+1 is not a perfect square, so n is added to the sequence. In case 2, n == 0 (mod 4) or n == 3 (mod 4), and we wish to show that n is not added to the sequence. That is, we wish to show that there exists m < n already in the sequence such that mn+1 is a perfect square. For this we let m = n - 2, which is positive since n >= 3. By the inductive hypothesis, since m == 1 (mod 4) or m == 2 (mod 4) and m < n, m is already in the sequence. And we have m*n + 1 = (n - 2)*n + 1 = n^2 - 2*n + 1 = (n - 1)^2, so mn+1 is indeed a perfect square, and so n is not added to the sequence. Thus n is added to the sequence if and only if n == 1 (mod 4) or n == 2 (mod 4). This completes the proof. - _Robert C. Lyons_, Jun 30 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001429 | OEIS-A042977

Triangle T(n,k) read by rows: coefficients of a polynomial sequence occurring when calculating the n-th derivative of Lambert function W.

Conjecture: the polynomials have no real roots greater than or equal to -1. This is equivalent to the statement that the derivatives of the 0th branch of the Lambert W function have no real roots greater than -1/e. - _Colin Linzer_, Jan 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001430 | OEIS-A042978

Stern primes: primes not of the form p + 2b^2 for p prime and b > 0.

Related to a conjecture of Goldbach.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001431 | OEIS-A044140

Numbers k such that substring "02" occurs in the base-7 representation of k but not of k-1.

Conjectures from _Colin Barker_, May 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001432 | OEIS-A044279

Numbers k such that string 3,1 occurs in the base 9 representation of k but not of k-1.

Conjecture: the sum of the digits of the terms of the sequence is always a multiple of 9 or 1 plus a multiple of 9. - _Harvey P. Dale_, May 25 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001433 | OEIS-A045406

A diagonal of A008296.

Conjecture: D-finite with recurrence a(n) +(2*n-7)*a(n-1) +(n-4)^2*a(n-2)=0. - _R. J. Mathar_, Sep 15 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 26 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001434 | OEIS-A045473

Primes congruent to 6 mod 7.

Conjecture: Primes p such that ((x+1)^7-1)/x has 3 irreducible factors of degree 2 over GF(p). - _Federico Provvedi_, Mar 31 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001435 | OEIS-A045575

Nonnegative numbers of the form x^y - y^x, for x,y > 1.

Conjecture: For d > 11, 10^d - d^10 is the largest (base-ten) d-digit term. - _Hans Havermann_, Jun 12 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Erich Friedman_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001436 | OEIS-A045576

Numbers k that divide 3^k + 2^k.

Conjecture: Every member of the sequence except 1 is of the form p*m where p is prime and m is in the sequence. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001437 | OEIS-A045626

Bends in loxodromic sequence of spheres in which each 5 consecutive spheres are in mutual contact.

G.f.: -(1-3*x-2*x^2+x^3+2*x^4)/((1+x)*(1-2*x+x^2-2*x^3+x^4)) (conjectured). - _Ralf Stephan_, May 06 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001438 | OEIS-A045650

Numbers that cannot be expressed as k + floor(log(k)) where k is an integer.

Conjecture: lim n->infinity a(n) = e^n. - _Ron R. King_, Nov 10 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001439 | OEIS-A045671

Extension of Beatty sequence; complement of A045672.

For n>=1, these are the positions of 1 in the fixed point of the morphism 0->11, 1->1110; see A285671. Conjecture: -1 < n*r - a(n) < 2 for n>=0, where r = (1 + sqrt(17))/4. - _Clark Kimberling_, May 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aviezri S. Fraenkel_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001440 | OEIS-A045672

Extension of Beatty sequence; complement of A045671 (apart from the initial 0).

For n>=1, these are the positions of 0 in the fixed point of the morphism 0->11, 1->1110; see A285671 and Mathematica program. Conjecture: -1 < n*r - a(n) < 3 for n>=0, where r = (5 + sqrt(17))/2. - _Clark Kimberling_, May 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aviezri S. Fraenkel_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001441 | OEIS-A045679

Numbers congruent to 0,1,4,9 mod 12 missing from A045673 (conjectured to be finite).

Numbers congruent to 0,1,4,9 mod 12 missing from A045673 (conjectured to be finite).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Allan Wilks_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001442 | OEIS-A045739

Number of edges in all noncrossing forests on n nodes on a circle.

Conjecture D-finite with recurrence -8*(n-1)*(42011*n-237357)*(2*n-1)*a(n) -2*(2*n-3)*(168044*n^2+2298095*n-1588326)*a(n-1) +2*(-8750238*n^3+268256261*n^2-1419101561*n+1999934970)*a(n-2) +4*(137297737*n^3-1755498200*n^2+747339 5243*n-10564858105)*a(n-3) +5*(25384204*n^3-3505044 39*n^2+1587560537*n-2286713022)*a(n-4) -25*(n-4)*(n-7)*(3362075*n-9824604)*a(n-5)=0. - _R. J. Mathar_, Jul 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001443 | OEIS-A045740

Number of components in all forests on nodes on a circle.

Conjecture D-finite with recurrence -2*(n-1)*(2*n-1)*(7912210314*n^2 +24034951267*n -109031255382)*a(n) +2*(-15824420628*n^4 +759853283620*n^3 -1653756416501*n^2 -3170366114943*n +6074871939666) *a(n-1) +2*(1171007126472*n^4 -5580539787848*n^3 -21281457754861*n^2 +151349953543323*n -205322404158756) *a(n-2) +(530118091038*n^4 -3085109917817*n^3 -8408054715093*n^2 +76142928591932*n -101713943817720) *a(n-3)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001444 | OEIS-A045742

Number of interior faces in all noncrossing connected graphs on n nodes on a circle.

Conjecture D-finite with recurrence n*(n-1)*(523*n-1993)*a(n) -6*(n-1)*(759*n^2-6019*n+12790)*a(n-1) -12*(n-2)*(4707*n^2-17937*n+6260)*a(n-2) +72*(3*n-10)*(759*n-2224)*(3*n-11)*a(n-3)=0. - _R. J. Mathar_, Jul 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001445 | OEIS-A045743

Number of noncrossing connected graphs on n nodes on a circle having no triangular faces.

Conjecture D-finite with recurrence 5*n*(n-1)*(n-2)*(1353818*n-3651663)*a(n) +16*(n-1)*(n-2)*(2236406*n^2-6864025*n+2014500)*a(n-1) -16*(n-2)*(46338048*n^3-326486432*n^2+734160494*n-506411475)*a(n-2) +8*(2*n-7)*(26592864*n^3-436441232*n^2+2071128458*n-3030735075)*a(n-3) +128*(94246*n-334675)*(4*n-19)*(2*n-9)*(4*n-17)*a(n-4)=0. - _R. J. Mathar_, Jul 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001446 | OEIS-A045761

Define polynomials Pn by P0 = 0, P1 = x, P2 = P1 + P0, P3 = P2 * P1, P4 = P3 + P2, etc. alternately adding or multiplying. For even n > 2k, then first k coefficients of Pn remain unchanged and their values are the first k terms of the sequence.

Conjecture: 1/d = 0.39952466709679946... = A268107. - _Jean-François Alcover_, Aug 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | James Boudinot (jboudinot(AT)yahoo.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001447 | OEIS-A045764

n + 1 - d(n) - phi(n) never takes these values (conjecture).

n + 1 - d(n) - phi(n) never takes these values (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001448 | OEIS-A045768

Numbers k such that sigma(k) == 2 (mod k).

k | sigma(k) produces the multiperfect numbers (A007691). It is an open question whether k | sigma(k) - 1 iff k is a prime or 1. In particular, it is not known if there exist solutions to sigma(k) = 2k+1, called quasiperfect numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dan Hoey_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001449 | OEIS-A045797

Evenish numbers (prime to 10 and 10's digit is even).

Conjecture a(n) = a(n-1)+a(n-4)-a(n-5). G.f.: x*(1+2*x+4*x^2+2*x^3+11*x^4) / ((1-x)^2*(1+x)*(1+x^2)). - _Colin Barker_, Apr 14 2012 The conjecture above is correct. - _Jianing Song_, Apr 27 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. H. Conway_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001450 | OEIS-A045798

Oddish numbers (prime to 10 and 10's digit is odd).

Conjecture a(n) = a(n-1)+a(n-4)-a(n-5). G.f.: x*(11+2*x+4*x^2+2*x^3+x^4)/((1-x)^2*(1+x)*(1+x^2)). - _Colin Barker_, Apr 14 2012 a(n+4) = a(n) + 20. This confirms Barker's conjecture. - _Robert Israel_, Mar 27 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. H. Conway_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001451 | OEIS-A045831

Number of 4-core partitions of n.

Conjecturally Sum_n a(n)q^(8n+5) equals theta series of sodalite. - _Fred Lunnon_, Mar 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001452 | OEIS-A045994

a(0)=1, a(n) = M(n) + Sum_{k=1..n-1} M(k)*a(n-k-1), where M(n) are the Motzkin numbers (A001006).

Conjecture: (n+1)*a(n) + 2*(-2*n-1)*a(n-1) + 2*(-n+3)*a(n-2) + (11*n-19)*a(n-3) + (11*n-27)*a(n-4) + 3*(n-3)*a(n-5) = 0. - _R. J. Mathar_, Sep 27 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001453 | OEIS-A045999

Length of n-th term of binary Gleichniszahlen-Reihe (BGR) sequence A045998.

Reference gives a conjectured formula.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001454 | OEIS-A046040

Numbers that are the sum of 6 but no fewer positive cubes.

According to the McCurley article, it is conjectured that there are exactly 3922 terms of which the largest is a(3922) = 1290740.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001455 | OEIS-A046041

Numbers (not of the form 9n+-4) that were conjectured not to be the sum of 3 positive or negative cubes. However, further research suggests that this sequence is in fact empty.

Numbers (not of the form 9n+-4) that were conjectured not to be the sum of 3 positive or negative cubes. However, further research suggests that this sequence is in fact empty.

猜想/问题提及段 | historical_or_current_mention_unreviewed | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001456 | OEIS-A046052

Number of prime factors of Fermat number F(n).

John Selfridge apparently conjectured that this sequence is not monotonic, so at some point a(n+1) < a(n). Related sequences such as A275377 and A275379 already exhibit such behavior. - _Jeppe Stig Nielsen_, Jun 08 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001457 | OEIS-A046057

Smallest order m > 0 for which there are n nonisomorphic finite groups of order m, or 0 if no such order exists.

R. Keith Dennis conjectures that there are no 0's in this sequence. See A053403 for details.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001458 | OEIS-A046060

5-multiperfect numbers.

Conjectured finite and probably these are the only terms; cf. Flammenkamp's link. [_Georgi Guninski_, Jul 25 2012]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001459 | OEIS-A046094

Agoh's congruence; a(n) is conjectured to be 1 iff n is prime.

Agoh's congruence; a(n) is conjectured to be 1 iff n is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001460 | OEIS-A046160

Bends of spheres in Soddy's bowl of integers which are tangent to the unit sphere and a single 2-sphere.

Conjecture: the list of bends of all spheres not in the plane is A049636 "Congruent to 0 or 2 mod 3, but not equal to 0 or 3." (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001461 | OEIS-A046232

Cubes which are palindromes in base 4.

Conjectured g.f.: x^2*(1 + 40*x - 4284*x^2 + 30080*x^3 - 28672*x^4)/((1 - x)*(1 - 4*x)*(1 - 16*x)*(1 - 64*x)). - _Stefano Spezia_, Jul 31 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, May 15 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001462 | OEIS-A046385

Smallest palindrome with exactly n palindromic prime factors (counted with multiplicity), and no other prime factors.

There are 3 more known terms: a(15)=405504, a(16)=40955904, a(20)=677707776. Any other terms would have at least 18 decimal digits. Conjecture: The sequence is finite and has no other terms than those shown here. - _Hugo Pfoertner_, Aug 13 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Jun 15 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001463 | OEIS-A046529

Largest number in largest finite set fixed by mapping f(k) = sum of n-th power of digits of k.

The terms a(12) = 1281243062759 and beyond are conjectural, but almost certainly correct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001464 | OEIS-A046675

Expansion of Product_{i>0} (1-x^{p_i}), where p_i are the primes.

Conjecture: (-1)^n * a(n) >= 0 for all n >= 50; equivalently, A024939(n) >= A024939(n-2) for all n >= 50. For n <= 10000, the values with (-1)^n * a(n) < 0 are exactly 2, 9, 25, 33, 38, and 49. - _Alejandro Radisic_, Aug 14 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001465 | OEIS-A046699

a(1) = a(2) = 1, a(n) = a(n - a(n-1)) + a(n-1 - a(n-2)) if n > 2.

Conjecture: a(n+1) = a(n) + A215530(a(n) + n) for all n > 0. - _Velin Yanev_, Oct 17 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001466 | OEIS-A046704

Additive primes: sum of digits is a prime.

Harman 2012 also shows, under a conjecture about primes in short intervals, that there are 3/2 * x/(log x log log x) terms up to x. - _Charles R Greathouse IV_, Nov 17 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felice Russo_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001467 | OEIS-A046747

Number of n X n rational {0,1}-matrices of determinant 0.

The probability that a random n X n {0,1}-matrix is singular is conjectured to be asymptotic to C(n+1, 2)*(1/2)^(n-1). [Corrected by _N. J. A. Sloane_, Jan 02 2007]

猜想/问题提及段 | historical_or_current_mention_unreviewed | Günter M. Ziegler (ziegler(AT)math.tu-berlin.de) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001468 | OEIS-A046869

Good primes (version 1): prime(n)^2 > prime(n-1)*prime(n+1).

Conjecture: primes p(n) such that 2*p(n) >= p(n-1) + p(n+1). - _Thomas Ordowski_, Jul 25 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001469 | OEIS-A046921

Number of ways to express 2n+1 as p+2a^2; p = 1 or prime, a >= 0.

Goldbach conjectured this sequence is never zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001470 | OEIS-A046927

Number of ways to express 2n+1 as p+2q where p and q are primes.

This is related to a conjecture of Lemoine (also sometimes called Levy's conjecture, although Levy was anticipated by Lemoine 69 years earlier). - _Zhi-Wei Sun_, Jun 10 2008 The conjecture states that any odd number greater than 5 can be written as p+2q where p and q are primes. It can be conjectured that 1, 3, 5, 59 and 151 are the only odd integers n such that n + 2p and n + 2q both are composite for all primes p,q with n = p + 2q. (Following an observation from V. Shevelev, cf. link to SeqFan list.) - _M. F. Hasler_, Apr 10 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001471 | OEIS-A046932

a(n) = period of x^n + x + 1 over GF(2), i.e., the smallest integer m>0 such that x^n + x + 1 divides x^m + 1 over GF(2).

Conjecture: a(2^k - 1) = 2^a(k) - 1. [See Bartholdi, 2000] More general conjecture: a((2^(k*m) - 1) / (2^m-1)) = (2^(a(k)*m) - 1) / (2^m-1). For m=1, it would imply Bartholdi conjecture. - _Max Alekseyev_, Oct 21 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Russell Walsmith_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001472 | OEIS-A046960

Bessel function J_0(n) is a monotonically decreasing positive sequence.

Conjectures from _Chai Wah Wu_, Mar 05 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001473 | OEIS-A046961

Bessel function Y_0(n) is a monotonically decreasing positive sequence.

Conjectures from _Chai Wah Wu_, Mar 05 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001474 | OEIS-A046995

Number of Greek-key tours on a 4 X n board; i.e., self-avoiding walks on 4 X n grid starting in top left corner.

a(n) = 3*a(n-1)+3*a(n-2)-9*a(n-3)-6*a(n-4)+5*a(n-5)+a(n-6)-3*a(n-7)+a(n-8) for n>=10. [conjectured by _Dean Hickerson_, Apr 05 2003; proved by _Jay Pantone_, Klotz, and Sullivan, Aug 01 2024] G.f.: x*(-(x-1)*(x^7-x^6-2*x^5+3*x^4-2*x^3-4*x^2-2*x-1))/((x^4-2*x^3+2*x^2+2*x-1)*(x^4-x^3-3*x^2-x+1)). [conjectured by Maksym Voznyy (voznyy(AT)mail.ru), Jul 27 2009; proved by _Jay Pantone_, Klotz, and Sullivan, Aug 01 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | Antreas P. Hatzipolakis (xpolakis(AT)otenet.gr) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001475 | OEIS-A047098

a(n) = 2*binomial(3*n, n) - Sum_{k=0..n} binomial(3*n, k).

Conjecture: 2*n*(2*n-1)*a(n) +(-71*n^2+112*n-48)*a(n-1) +3*(131*n^2-391*n+296)*a(n-2) -72*(3*n-7)*(3*n-8)*a(n-3)=0. - _R. J. Mathar_, Nov 30 2012 Conjecture verified using the differential equation 6*x + 3 + (-144*x^2 + 114*x - 7)*y + (-1296*x^3 + 792*x^2 - 101*x + 2)*y' + (-648*x^4 + 393*x^3 - 71*x^2 + 4*x)*y'' = 0 satisfied by the g.f. y(x). - _Robert Israel_, Mar 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 15 1998 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001476 | OEIS-A047211

Numbers that are congruent to {2, 4} mod 5.

Conjecture: n such that the characteristic polynomial of M(n) is irreducible over the rationals where M(n) is an n X n matrix with ones on the skew diagonal and below it and the skew line two positions above it and otherwise zeros; see example for one such matrix. Tested up to n=177. - _Joerg Arndt_, Aug 10 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 11 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001477 | OEIS-A047212

Numbers that are congruent to {0, 2, 4} mod 5.

Except for a(1) and a(4), also the arboricity of the n X n queen graph (conjectured). - _Eric W. Weisstein_, Jan 30 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001478 | OEIS-A047222

Numbers that are congruent to {0, 2, 3} mod 5.

Conjecture: Apart from 0, the sequence gives the values for c/6, such that an infinite number of primes, p, result in both p^2-c and p^2+c being positive primes, except when c is a square. When c is square solutions exist for c (both within and outside of the a(n) set), but occur at only a single prime p. See A274609. Other c values with only one prime providing a solution occur when p^2-c=3. See A274610. The only remaining c values with single p solutions are: c=2 (with p=3) and c=6 (with p=5). - _Richard R. Forberg_, Jun 26 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001479 | OEIS-A047239

Numbers that are congruent to {1, 2} (mod 6).

Conjecture: a(n) is the least positive integer > a(n-1) that is not equal to a(i) + a(j) + a(k) for any i <= j <= k <= n. - _Clark Kimberling_, Oct 09 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001480 | OEIS-A047241

Numbers that are congruent to {1, 3} mod 6.

12*a(n) is conjectured to be the length of the boundary after n iterations of the hexagon and square expansion shown in the link. The squares and hexagons have side length 1 in some units. The pattern is supposed to become the planar Archimedean net 4.6.12 when n -> infinity. - _Kival Ngaokrajang_, Nov 30 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001481 | OEIS-A047363

Numbers that are congruent to {0, 2, 3, 4, 5} mod 7.

Conjecture: Apart from 0, and the further exclusions noted below, the sequence gives the values of c/6 such that an infinite number of primes, p, result in both p^3+c and p^3-c being positive primes. Taking the complement we say: the excluded c/6 values are {1,6} mod 7. See A005097 for a conjecture on the modulo patterns of excluded c/6 values for the general case of p^q + c and p^q - c both prime, for any q > 0, and see A047222 for q=2. Note that polynomial factorization also excludes a few c/6 values. This occurs here when c is an even cube (A016743), which requires a further exclusion of certain c/6 values in this sequence when (6c)^3/6 == 0 (mod 7), or c/6 = {0, 12348, 98784, ...}. - _Richard R. Forberg_, Jun 28 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001482 | OEIS-A047618

Numbers that are congruent to {0, 2, 5} mod 8.

Conjecture: a(n)+a(n+1)+a(n+2) = 8*n-1; or a(n) = 8*(n-2)-a(n-1)-a(n-2)-1, n>3, with a(1)=0, a(2)=2, a(3)=5. - _L. Edson Jeffery_, Sep 02 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001483 | OEIS-A047658

Numbers k such that the initial k digits in decimal portion of Pi form a prime number.

Conjecture: this sequence is finite. - _Carlos Rivera_ Rivera's conjecture that this sequence is finite conflicts with heuristics; the next entry is almost certainly 6205, since floor((Pi-3)*10^6205) is (very) probably prime, though its proof may take decades. - _David Broadhurst_, Nov 08 2000

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carlos Rivera_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001484 | OEIS-A047704

Numbers that are the sum of 5 but no fewer positive cubes.

This sequence is conjectured to have a finite number of terms. - _T. D. Noe_, Dec 13 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | Arlin Anderson (starship1(AT)gmail.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001485 | OEIS-A047781

a(n) = Sum_{k=0..n-1} binomial(n-1,k)*binomial(n+k,k). Also a(n) = T(n,n), array T as in A049600.

This array is now listed as A142978, where some conjectural congruences for the present sequence are given. - _Peter Bala_, Nov 13 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001486 | OEIS-A047788

Numerators of Glaisher's I-numbers.

Conjecture: L(2n+1, chi3) = a(n)/A047789(n) * (2*Pi)^(2n+1)/((2n)!*3^(2n+3/2)), where L(s, chi3) = Sum_{k>=1} Legendre(k,3)/k^s = Sum_{k>=1} A102283(k)/k^s is the Dirichlet L-function for the non-principal character modulo 3. - _Jianing Song_, Nov 17 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001487 | OEIS-A047802

Smallest abundant number (sigma(x) > 2x) which is not divisible by any of the first n primes.

Conjecture: other than a(1) = 945, all terms are cubefree. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Schimke (ulrschimke(AT)aol.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001488 | OEIS-A047837

Honaker's triangle problem: form a triangle with base of length n, all entries different, all row sums equal; a(n) gives minimal row sum.

Agrees with A047873 at least for n < 365, conjectured to always agree.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001489 | OEIS-A047849

a(n) = (4^n + 2)/3.

A. A. Ivanov conjectures that the dimension of the universal embedding of the unitary dual polar space DSU(2n,4) is a(n). - J. Taylor (jt_cpp(AT)yahoo.com), Apr 02 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001490 | OEIS-A047872

a(n) = floor(abs(B(2*n + 2)/B(2*n))) where B(n) is the n-th Bernoulli number.

a(n) = floor((n+1)*(2*n+1)/(2*Pi^2)) (conjectured). - _Bill McEachen_, Dec 08 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001491 | OEIS-A047873

a(n) = max_{r=1..n-1} ceiling(t(t(n)-t(r-1))/(n-r+1)), where t() = triangular numbers A000217.

Another lower bound for Honaker triangle problem (A047837); conjectured to be exact value.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mike Keith_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001492 | OEIS-A047876

a(n) = Sum{T(i,n-i): i=0,1,...,n}, array T as in A049604.

a(n) = 2*a(n-3) + a(n-4) - a(n-6) - 2*a(n-7) + a(n-10) for n > 14 (conjectured). G.f.: 2*x*(x^13 + x^11 - x^9 - 2*x^8 - 4*x^7 - x^6 + x^5 + 2*x^4 - 4*x^2 - 4*x - 3)/(x^10 - 2*x^7 - x^6 + x^4 + 2*x^3 - 1) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001493 | OEIS-A047877

a(n) = (1/2)*Sum_{i=0..n} T(i,n-i), array T as in A049604.

a(n) = 2*a(n-3) + a(n-4) - a(n-6) - 2*a(n-7) + a(n-10) for n > 14 (conjectured). G.f.: x*(x^13 + x^11 - x^9 - 2*x^8 - 4*x^7 - x^6 + x^5 + 2*x^4 - 4*x^2 - 4*x - 3)/(x^10 - 2*x^7 - x^6 + x^4 + 2*x^3 - 1) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001494 | OEIS-A047881

a(n) = Sum_{h=0..n, k=0..n} T(h,k), array T counting knights' moves as in A049604.

Conjecture: a(n) = a(n-1) + a(n-2) - 2*a(n-5) + a(n-8) + a(n-9) - a(n-10) for n > 11. - _Chai Wah Wu_, May 25 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001495 | OEIS-A047909

Array read by antidiagonals upwards: h(n,k) = number of sequences with n copies each of 1,2,...,k and longest increasing subsequence of length k (n>=1, k>=1).

Conjecture: Column k > 1 is asymptotic to k^(k*n + 1/2) / (2*Pi*n)^((k-1)/2). - _Vaclav Kotesovec_, Feb 21 2016 Conjecture: Row k > 1 is asymptotic to sqrt(k) * (k^k/(k-1)!)^n * n^((k-1)*n) / exp((k-1)*(n+1)). - _Vaclav Kotesovec_, Feb 21 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001496 | OEIS-A047926

a(n) = (3^(n+1) + 2*n + 1)/4.

Conjecture: Number of representations of 3^(2n) as a sum a^2 + b^2 + c^2 with 0 < a <= b <= c. That is, a(1) = 3 because 3^2 = 1^2 + 2^2 + 2^2, a(2) = 3 because 3^4 = 1^2 + 4^2 + 8^2 = 3^2 + 6^2 + 6^2 = 4^2 + 4^2 + 7^2. - _Zak Seidov_, Mar 01 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001497 | OEIS-A047946

a(n) = 5*F(n)^2 + 3*(-1)^n where F(n) are the Fibonacci numbers A000045.

For n>0, the linear recurrence for the sequence F(n*k)^2 has signature (a(n),a(n),-1) for n odd, and (a(n),-a(n), 1) for n even. For example, the linear recurrence for the sequence F(3*k)^2 has signature (17, 17, -1) (conjectured). - _Greg Dresden_, Aug 30 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, May 21 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001498 | OEIS-A047947

a(n) is the number of k values for which A023193(k) = n.

The old name was: "Schinzel's rhobar(n), number of distinct lengths of a block of consecutive integers on which a maximum of n primes occurs infinitely often (under the k-tuple conjecture)." [Note that "rhobar" is A023193.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001499 | OEIS-A047968

a(n) = Sum_{d|n} p(d), where p(d) = A000041 = number of partitions of d.

Conjecture: for n > 6, a(n) is strictly increasing. - _Franklin T. Adams-Watters_, Apr 19 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 11 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001500 | OEIS-A047972

Distance of n-th prime to nearest square.

Conjecture: a(n) < sqrt(prime(n)) and lim_{n->infinity} a(n)/sqrt(prime(n)) = 1, where prime(n) is the n-th prime. - _Ya-Ping Lu_, Nov 29 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enoch Haga_, Dec 11 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001501 | OEIS-A048161

Primes p such that q = (p^2 + 1)/2 is also a prime.

It is conjectured that there are an infinite number of such triangles.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Harvey Dubner (harvey(AT)dubner.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001502 | OEIS-A048193

Number of chordal graphs (or triangulated graphs) on n vertices.

Conjectured partial sums of A079456. - _Sean A. Irvine_, Jun 25 2022 This conjecture is true, because each connected component of a complement of a chordal graph is an isolated vertex, except for at most one. Indeed, otherwise there would exist edges A-B and C-D in two different connected components of the complement, and then A-C-B-D-A is an induced 4-cycle in the original graph. - _Andrei Zabolotskii_, Jan 02 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gordon Royle_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001503 | OEIS-A048200

Minimal length pair-exchange / set-rotate sequence to reverse n distinct ordered elements.

Conjecture: a(n) = (3*n^2/4)-2*n if n is even and a(n) = (3*n^2-10*n+15)/4 if n is odd. See links for more information. - _Sai Satwik Kuppili_ and _Bhadrachalam Chitturi_, Jun 09 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tony Bartoletti_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001504 | OEIS-A048211

Number of distinct resistances that can be produced from a circuit of n equal resistors using only series and parallel combinations.

Conjecture: a(n) ~ K * a(n-1), K approx 2.54. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tony Bartoletti_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001505 | OEIS-A048247

Every prime occurs to this power in some factorial.

There are no primes in the sequence, as the prime p fails the base p test. The set of positive integers failing the base p test for membership has density 1/p. Also, when n is a nonmember of the set, any base p whose test n fails has p<=n. Therefore one conjectural estimate for the number of members of the set <=x would be x*product{primes p<=x}(1-1/p) ~ e^(-gamma)*x/log(x). However, a similar heuristic for the primes fails, as pi(x) ~ x/log(x) and not e^(-gamma)*x/log(x). Here gamma denotes the Euler-Mascheroni constant. - _David L. Harden_, Aug 24 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | Antreas P. Hatzipolakis (xpolakis(AT)otenet.gr) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001506 | OEIS-A048270

Sequence of 2 Pythagorean triangles, each with a leg and hypotenuse prime. The leg of the second triangle is the hypotenuse of the first.

It is conjectured that there are infinitely many such pairs of triangles.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Harvey Dubner (harvey(AT)dubner.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001507 | OEIS-A048285

Number of Dyck paths of length 2n with nondecreasing peaks.

Conjectural g.f.: Sum_{n>=0} (x*(1 - x))^n/(Product_{i=2..n+1} (1 - 2*x + x^i)) (checked up to x^50). - _Peter Bala_, Mar 31 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | Olivier Roques (roques(AT)labri.u-bordeaux.fr) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001508 | OEIS-A048330

Numbers that are repdigits in base 5.

Conjecture: G.f.: x*(1+2*x+3*x^2+4*x^3) / ((x-1)*(1+x)*(x^2+1)*(5*x^4-1)) with a(n) = 6*a(n-4) - 5*a(n-8). - _R. J. Mathar_, Mar 15 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Feb 15 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001509 | OEIS-A048331

Numbers that are repdigits in base 6.

Conjecture: G.f.: x*(1+2*x+3*x^2+4*x^3+5*x^4) / ((x-1)*(x^4+x^3+x^2+x+1)*(6*x^5-1)) with a(n) = 7*a(n-5) - 6*a(n-10). - _R. J. Mathar_, Mar 15 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Feb 15 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001510 | OEIS-A048582

Pisot sequence L(4,9).

a(n) = 3*a(n-1) - 2*a(n-2) + a(n-3) + a(n-4) - 2*a(n-5) + a(n-6) - a(n-7) (conjectured). Recurrence is satisfied for at least 760000 terms. - _Chai Wah Wu_, Jul 25 2016 Note the warning in A010925 from Pab Ter (pabrlos(AT)yahoo.com), May 23 2004: [A010925] and other examples show that it is essential to reject conjectured generating functions for Pisot sequences until a proof or reference is provided. - _N. J. A. Sloane_, Jul 26 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001511 | OEIS-A048590

Pisot sequence L(8,9).

Conjecture: satisfies a linear recurrence having signature (2,-1,0,1,-1). - _Harvey P. Dale_, Dec 31 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001512 | OEIS-A048625

Pisot sequence P(4,6).

Conjecture: satisfies a linear recurrence having signature (1, 0, 1). - _Harvey P. Dale_, Jun 05 2021 G.f.: (conjecture) ((Q(0)-1)/2 -(x+x^2+x^3+2*x^4+3*x^5))/x^6, where Q(k) = 1 + x^3 + (2*k+3)*x - x*(2*k+1 + x^2)/Q(k+1); (continued fraction). - _Sergei N. Gladkovskii_, Oct 05 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001513 | OEIS-A048651

Decimal expansion of Product_{k >= 1} (1 - 1/2^k).

Limit of A177510(n)/A000079(n-1) as n->infinity (conjecture). - _Mats Granvik_, Mar 27 2011 (Sum_{k>0} (4^k-1)/(Product_{i=1..k} ((4^i-1)*(2*4^i-1))))*2 = 2/7 + 2/(3*7*31) + 2/(3*7*15*31*127)+2/(3*7*15*31*63*127*511) + ... (conjecture). - _Werner Schulte_, Dec 22 2016 Constant C = Sum_{n >= 0} (-1)^n/(Product_{k = 1..n} (2^k - 1)). The above conjectural result by Schulte follows by adding terms of this series in pairs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001514 | OEIS-A048702

Binary palindromes of even length divided by 3.

Conjecture: a(n) = 2^floor(log_2(n)) * Sum_{i=1..n} 1/(2^v_2(i)), for n >= 1, where v_2(i) = A007814(i) is the exponent of the highest power of 2 dividing i. Conjecture: a(n) = n*2^floor(log_2(n)) - Sum_{i=1..floor(log_2(n))} 2^(floor(log_2(n)) - i)*floor(n/(2^i)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 07 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001515 | OEIS-A048715

Binary expansion matches (100(0)*)*(0|1|10)?; or, Zeckendorf-like expansion of n using recurrence f(n) = f(n-1) + f(n-3).

The conjecture by Benoit is correct. This is easily proved using the well-known result that the multiplicity with which a prime p divides C(n+m,n) is the number of carries when adding n+m in base p. - _Franklin T. Adams-Watters_, Oct 06 2009 a(n) XOR 6*a(n) = 7*a(n); 3*a(n) XOR 4*a(n) = 7*a(n); 3*a(n) XOR 5*a(n) = 6*a(n); (conjectures). - _Paul D. Hanna_, Jan 22 2006 The conjectures can be verified using the Walnut theorem-prover (see links). - _Sebastian Karlsson_, Dec 31 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 30 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001516 | OEIS-A048718

Binary expansion matches ((0)*0001)*(0*); or, Zeckendorf-like expansion of n using recurrence f(n) = f(n-1) + f(n-4).

a(n) XOR 14*a(n) = 15*a(n); 3*a(n) XOR 9*a(n) = 10*a(n); 3*a(n) XOR 13*a(n) = 14*a(n); 5*a(n) XOR 9*a(n) = 12*a(n); 5*a(n) XOR 11*a(n) = 14*a(n); 6*a(n) XOR 11*a(n) = 13*a(n); 7*a(n) XOR 9*a(n) = 14*a(n); 7*a(n) XOR 10*a(n) = 13*a(n); 7*a(n) XOR 11*a(n) = 12*a(n); 12*a(n) XOR 21*a(n) = 25*a(n); 12*a(n) XOR 37*a(n) = 41*a(n); etc. (conjectures). - _Paul D. Hanna_, Jan 22 2006 The conjectures can be verified using the Walnut theorem-prover (see links). - _Sebastian Karlsson_, Dec 31 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 30 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001517 | OEIS-A048799

Decimal expansion of Sum_{n >= 2} 1/S(n)!, where S(n) is the Kempner number A002034(n).

I conjecture that the constants defined in the present sequence, A048834, A071120, A048835, A048836, A048837, A048838 are irrational. - Sukanto Bhattacharya (susant5au(AT)yahoo.com.au), Apr 28 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | Charles T. Le (charlestle(AT)yahoo.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001518 | OEIS-A048819

Decimal expansion of the Minkowski-Bower constant.

It is conjectured that Minkowski's question mark function has exactly five fixed points on [0,1], or four modulo 1; see Gayfulin and Shulga, Conjecture 1. - _Eric W. Weisstein_, Oct 08 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, Apr 15 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001519 | OEIS-A048868

Numbers whose reduced residue system contains more primes than nonprimes.

Conjecture: all terms are even.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001520 | OEIS-A048881

a(n) = A000120(n+1) - 1 = wt(n+1) - 1.

Conjecture: a(n) is the sum of digits of the n-th word in A076478, for n >= 1; has been confirmed for n up to 20000. - _Clark Kimberling_, Jul 14 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001521 | OEIS-A048888

a(n) = Sum_{m=1..n} T(m,n+1-m), array T as in A048887.

Define a "numbral arithmetic" by replacing addition with binary bitwise inclusive-OR (so that [3] + [5] = [7] etc.) and multiplication becomes shift-&-OR instead of shift-&-add (so that [3] * [3] = [7] etc.). [d] divides [n] means there exists an [e] with [d] * [e] = [n]. For example the six divisors of [14] are [1], [2], [3], [6], [7] and [14]. Then it appears that this sequence gives the number of proper divisors of [2^n-1]. Conjecture confirmed by Richard C. Schroeppel, Dec 14 2001. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001522 | OEIS-A048963

Table in which n-th row lists digits in periodic part of decimal expansion of reciprocal of n-th prime.

Conjecture: regarded as a decimal fraction, this number is normal in base 10. - _Franklin T. Adams-Watters_, Aug 20 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001523 | OEIS-A049048

Composite numbers n such that k! == 1 (mod n) for some k > 2.

Conjecture: If c is composite then n! == 1 (mod c) for at most one n >= 2. - David Wilson, Oct 06 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001524 | OEIS-A049067

LaBar's conjecture (steps to return n to 1 after division by 3 and, if needed, multiplication by 2, addition of 1 or 2).

LaBar's conjecture (steps to return n to 1 after division by 3 and, if needed, multiplication by 2, addition of 1 or 2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enoch Haga_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001525 | OEIS-A049074

Ulam's conjecture (steps to return n to 1 after division by 2 and, if needed, multiplication by 3 with 1 added).

Ulam's conjecture (steps to return n to 1 after division by 2 and, if needed, multiplication by 3 with 1 added).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enoch Haga_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001526 | OEIS-A049094

Numbers m such that 2^m - 1 is divisible by a square > 1.

Conjecture: 2^n-1 is squarefree iff gcd(n,2^n-1)=1. If true, the conjecture would imply that Mersenne numbers (A001348) are squarefree. - _Vladeta Jovovic_, Apr 12 2002. But the conjecture is not true: counterexamples are n = 364 and n = 1755, i.e., gcd(364,2^364-1) = 1 and (2^364-1) mod 1093^2 = 0 and gcd(1755,2^1755-1) = 1 and (2^1755-1) mod 3511^2 = 0, cf. A001220. - _Vladeta Jovovic_, Nov 01 2005. The conjecture is true with assumption that n is not a multiple of A002326((q-1)/2), where q is a Wieferich prime A001220. - _Thomas Ordowski_, Nov 17 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001527 | OEIS-A049096

Numbers k such that 2^k + 1 is divisible by a square > 1.

Conjecture: lim n -> infinity a(n)/n = C exists and 4 < C < 9/2. There seems to be a sequence of primes p such that p^2 never divides numbers of the form 2^x + 1: the first few are 2, 7, 23, 31. - _Benoit Cloitre_, Aug 20 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001528 | OEIS-A049125

Revert transform of (1 + x - x^2) / (1 + x)^2.

Conjecturally, the number of sequences (e(1), ..., e(n-1)), 0 <= e(i) < i, such that there is no triple i < j < k with e(i) >= e(k). [Martinez and Savage, 2.10] - _Eric M. Schmidt_, Jul 17 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001529 | OEIS-A049310

Triangle of coefficients of Chebyshev's S(n,x) := U(n,x/2) polynomials (exponents in increasing order).

The number of zeros Z(n) of the S(n, x) polynomials in the open interval (-1,+1) is 2*b(n) for even n >= 0 and 1 + 2*b(n) for odd n >= 1, where b(n) = floor(n/2) - floor((n+1)/3). This b(n) is the number of integers k in the interval (n+1)/3 < k <= floor(n/2). See a comment on the zeros of S(n, x) above, and b(n) = A008615(n-2), n >= 0. The numbers Z(n) have been proposed (with a conjecture related to A008611) by _Michel Lagneau_, as the number of zeros of Fibonacci polynomials on the imaginary axis (-I,+I), with I=sqrt(-1). They are Z(n) = A008611(n-1), n >= 0, with A008611(-1) = 0. Also Z(n) = A194960(n-4), n >= 0. Proof using the A008611 version. A194960 follows from this.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001530 | OEIS-A049425

Row sums of triangle A049404.

Conjecture: -a(n) +a(n-1) +(2*n-2)*a(n-2) +(2-3*n+n^2)*a(n-3)=0. - _R. J. Mathar_, Nov 14 2011 So, the above conjecture is true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001531 | OEIS-A049486

Maximum length of non-crossing path on n X n square lattice.

Conjecture: a(n) = (9 + (-1)^n - 8*n + 4*n^2)/2 for n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Arlin Anderson (starship1(AT)gmail.com) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001532 | OEIS-A049505

a(n) = Product_{1<=i<=j<=n} (i+j+n-1)/(i+j-1), number of symmetric plane partitions in n-cube.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001533 | OEIS-A049591

Odd primes p such that p+2 is composite.

Conjecture: start from any initial value f(1) >= 2 and define f(n) to be the largest prime factor of f(1) +f(2) + ... +f(n-1); then f(n) = n/2 + O(log(n)) and there are infinitely many primes p such that f(2p)=p. Conjecture: current sequence gives primes satisfying f(2p)=p when f(1)=3. - _Benoit Cloitre_, Jun 04 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001534 | OEIS-A049608

a(n)=T(n,n+2), array T as in A049600.

Conjecture: (n-1)*(2*n-1)*(n+2)*a(n) -12*n*(n-1)*(n+1)*a(n-1) +(n-2)*(2*n+1)*(n+1)*a(n-2)=0. - _R. J. Mathar_, Oct 26 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001535 | OEIS-A049609

a(n)=T(n,n+3), array T as in A049600.

Conjecture: -(n+3)*(n-1)*a(n) +(7*n^2+8*n-9)*a(n-1) +(-7*n^2+8*n+9)*a(n-2) +(n+1)*(n-3)*a(n-3)=0. - _R. J. Mathar_, Oct 26 2015 Conjecture: +n*(n+3)*(n-1)*a(n) -(2*n+1)*(3*n^2+3*n-4)*a(n-1) +(n-2)*(n+2)*(n+1)*a(n-2)=0. - _R. J. Mathar_, Oct 26 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001536 | OEIS-A049638

a(n) = T(n,4), array T as in A049627.

Conjectured g.f.: 2 -x*(-7-9*x-5*x^2+2*x^3+3*x^4) / ((1+x)*(1+x+x^2)*(x-1)^2). - _R. J. Mathar_, Oct 26 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001537 | OEIS-A049652

a(n) = (F(3*n+2) - 1)/4, where F=A000045 (the Fibonacci sequence).

"A beautiful combinatorial proof of Dodgson's result was recently given by Zeilberger and an application is presented by Amdeberhan and Ekhad, where a conjecture of Kuperberg and Propp is proved using Dodgson's formula." (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001538 | OEIS-A049777

Triangular array read by rows: T(m,n) = n + n+1 + ... + m = (m+n)(m-n+1)/2.

Conjecture: There is at least one prime in interval [T(n,k), T(n,k+1)]. Since T(n,k+1)/T(n,k) decreases to (k+1)/k as n increases, this is true for k=1 ("Bertrand's Postulate", first proved by P. Chebyshev), k=2 (proved by El Bachraoui) and k=3 (proved by Loo).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001539 | OEIS-A049803

a(n) = n mod 3 + n mod 9 + ... + n mod 3^k, where 3^k <= n < 3^(k+1).

Conjecture: For b >= 2, consider the function s(n,b) = Sum_{1 <= b^j <= n} (n mod b^j) from p. 8 in Dearden et al. (2011). Then s(b*n + r, b) = b*s(n,b) + r*N(n,b) for 0 <= r <= b-1, where N(n,b) = floor(log_b(n)) + 1 is the number of digits in the base-b representation of n. As initial conditions, we have s(n,b) = 0 for 1 <= n <= b. (This is a generalization of a result by _Robert Israel_ in A049802.) If A_b(x) = Sum_{n >= 1} s(n,b)*x^n is the g.f. of the sequence (s(n,b): n >= 1) and the above conjecture is correct, then it can be proved that A_b(x) = b * A_b(x^b) * (1-x^b)/(1-x) + x * ((b-1)*x^b - b*x^(b-1) + 1)/((1-x)^2 * (1-x^b)) * Sum_{k >= 1} x^(b^k). (End) Conjecture: a(3*n+r) = 3*a(n) + r*A081604(n) = 3*a(n) + r*(floor(log_3(n)) + 1) for n >= 1 and r = 0, 1, 2. If the conjecture above is true, the g.f. A(x) satisfies A(x) = 3*(1+x+x^2)*A(x^3) + x*(2*x+1)/(1-x^3) * Sum_{k >= 1} x^(3^k). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001540 | OEIS-A049804

a(n) = n mod 4 + n mod 16 + ... + n mod 4^k, where 4^k <= n < 4^(k+1).

Conjecture: For b >= 2, consider the function s(n,b) = Sum_{1 <= b^j <= n} (n mod b^j) from p. 8 in Dearden et al. (2011). Then s(b*n + r, b) = b*s(n,b) + r*N(n,b) for 0 <= r <= b-1, where N(n,b) = floor(log_b(n)) + 1 is the number of digits in the base-b representation of n. As initial conditions, we have s(n,b) = 0 for 1 <= n <= b. (This is a generalization of a result by _Robert Israel_ in A049802.) If A_b(x) = Sum_{n >= 1} s(n,b)*x^n is the g.f. of the sequence (s(n,b): n >= 1) and the above conjecture is correct, then it can be proved that A_b(x) = b * A_b(x^b) * (1-x^b)/(1-x) + x * ((b-1)*x^b - b*x^(b-1) + 1)/((1-x)^2 * (1-x^b)) * Sum_{k >= 1} x^(b^k). (End) Conjecture: a(4*n+r) = 4*a(n) + r*A110591(n) = 4*a(n) + r*(floor(log_4(n)) + 1) for n >= 1 and r = 0, 1, 2, 3. If the conjecture above is true, the g.f. A(x) satisfies A(x) = 4*(1 + x + x^2 + x^3)*A(x^4) + x*(1 + 2*x + 3*x^2)/(1 - x^4) * Sum_{k >= 1} x^(4^k). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001541 | OEIS-A049828

Triangular array T given by rows: T(n,k)=sum of remainders when Euclidean algorithm acts on n,k; for k=1,2,...,n; n >= 1.

For a fixed n, {(k,T(n,k)), k=1..n} is conjectured to be fractal in nature (see link). - _Tiberiu Szocs-Mihai_, Aug 17 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001542 | OEIS-A049997

Numbers of the form Fibonacci(i)*Fibonacci(j).

Conjecture: Fibonacci(m)*Fibonacci(n) with 2 < m < n is a perfect power only for (m,n) = (3,6). This has been verified for 2 < m < n <= 900. - _Zhi-Wei Sun_, Jan 02 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001543 | OEIS-A049998

a(n) = b(n)-b(n-1), where b=A049997 are numbers of the form Fibonacci(i)*Fibonacci(j).

David W. Wilson conjectured (Dec 14 2005) that this sequence consists only of Fibonacci numbers. Proofs were found by _Franklin T. Adams-Watters_ and _Don Reble_, Dec 14 2005. The following is Reble's proof:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001544 | OEIS-A049999

a(n) = smallest index k such that Fibonacci(k) = d(n), where d = A049998 (sequence of first differences of ordered products of Fibonacci numbers, i.e., of A049997, with no duplicates).

"_David W. Wilson_ conjectured (Dec 14 2005) that" sequence A049998 "consists only of Fibonacci numbers. Proofs were found by _Franklin T. Adams-Watters_ and _Don Reble_, Dec 14 2005." - _Petros Hadjicostas_, Nov 08 2019 [This comment was copied from A049998, which includes _Don Reble_'s proof of the conjecture.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

B02 A050000 - A099999 | 1,917 条

V3-001545 | OEIS-A050168

a(0) = 1; for n > 0, a(n) = binomial(n, floor(n/2)) + binomial(n-1, floor(n/2)).

Conjecture: (n+1)*a(n) + (n-3)*a(n-1) + 2*(-2*n+1)*a(n-2) + 4*(-n+3)*a(n-3) = 0. - _R. J. Mathar_, Nov 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001546 | OEIS-A050181

T(2n+3, n), array T as in A051168; a count of Lyndon words.

Conjecture: -(n-1)*(n+3)*(n+2)*a(n) + 2*(3*n-4)*(n+2)*(n+1)*a(n-1) - 4*n*(n+1)*(2*n-5)*a(n-2) + 2*(n-1)*(n+2)*(2*n-3)*a(n-3) - 4*(2*n-5)*(3*n-4)*(n+1)*a(n-4) + 8*n*(2*n-5)*(2*n-7)*a(n-5) = 0. - _R. J. Mathar_, Jul 20 2016 Using the above formulae, one can verify _R. J. Mathar_'s conjecture above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001547 | OEIS-A050189

T(n,4), array T as in A050186; a count of aperiodic binary words.

a(n) = 2*a(n-1) + 2*a(n-2) - 6*a(n-3) + 6*a(n-5) - 2*a(n-6) - 2*a(n-7) + a(n-8) for n > 11 (conjectured). G.f.: x^5*(5 + 2*x + x^2)/((1 - x)^5*(1 + x)^3) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001548 | OEIS-A050327

Number of factorizations into distinct squarefree factors indexed by prime signatures. A050326(A025487).

Conjecture: prime signatures corresponding to primorials A002110(i) with i > 1 set records in this sequence. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, Oct 15 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001549 | OEIS-A050385

Reversion of Moebius function A008683.

Unfortunately this conjecture is not correct; it fails at a(13). - _Jeffrey Shallit_, Nov 19 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, Nov 15 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001550 | OEIS-A050412

Riesel problem: start with n; repeatedly double and add 1 until reaching a prime. Sequence gives number of steps to reach a prime or 0 if no prime is ever reached.

It is conjectured that n = 509203 is the smallest Riesel number, i.e., n*2^k - 1 is composite for every k>0. - _Robert G. Wilson v_, Mar 01 2015. [This would imply that a(509203) is the first zero term in this sequence. - _N. J. A. Sloane_, Jul 31 2024] If a(n) = 0, then a(2n+1) is also 0. Conjecture: If a(n) = 1, then a(2n+1) is not 0. - _Jeppe Stig Nielsen_, Feb 12 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Dec 22 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001551 | OEIS-A050446

Table read by ascending antidiagonals: T(n, m) giving total degree of n-th-order elementary symmetric polynomials in m variables.

Conjecture: T(n,m) is the number of words of length n over the alphabet [m] such that any pair of adjacent letters sum to at most m + 1. - _John Tyler Rascoe_, Jun 06 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 23 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001552 | OEIS-A050498

Arithmetic progressions of at least 4 terms with common difference 6 having the same value of phi(x) start at these numbers.

Conjecture: a(n) == 0 (mod 6) for n >= 1. After division by 6 the sequence becomes [12, 36, 12721, 16921, 19441, 24481, 49681, 61561, 104161, 229321, 255361, 259681, 266401, 291721, 298201, 311041, 331921, ...].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_, Dec 27 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001553 | OEIS-A050647

a(n+1)^3 is next smallest cube ending with a(n)^3, initial term is 1.

Frank Ellermann conjectures (Jun 11 2001) that if b(n) = 10^(3^n -1), i.e., 1, 100, 100000000, etc. then a(n) = concatenation b(n-2) || a(n-1) for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Jul 15 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001554 | OEIS-A050686

Number of palindromes of length n and containing the digit 1 (or any other fixed nonzero digit).

The conjectured g.f is correct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Aug 15 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001555 | OEIS-A050811

Partition numbers rounded to nearest integer given by the Hardy-Ramanujan approximate formula.

This conjecture is false, for correct approximation see the formula below. - _Vaclav Kotesovec_, Apr 03 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Oct 15 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001556 | OEIS-A050931

Numbers having a prime factor congruent to 1 mod 6.

Conjecture: Numbers m such that abs(Sum_{k=1..m} [k|m]*A008683(k)*(-1)^(2*k/3)) = 0. - _Mats Granvik_, Jul 06 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | Klaus Strassburger (strass(AT)ddfi.uni-duesseldorf.de), Dec 30 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001557 | OEIS-A050970

Numerator of S(n)/Pi^n, where S(n) = Sum_{k=-inf..+inf} (4k+1)^(-n).

a(2n) = A046976(n), a(2n+1) = A089171(n+1) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001558 | OEIS-A050972

Smaller member of friendly pairs ordered by smallest maximal element.

Conjecture: If a(n) is in A005153, then A050973(n) is in A005153. It is natural to ask if there exists a generalization of the indicator function for A005153, call it m(n), such that m(n) = 1 for n in A005153, 0 < m(n) < 1 otherwise, and m(a(n)) <= m(A050973(n)) for all n. See also A017666. - _Jaycob Coleman_, Sep 27 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001559 | OEIS-A051064

3^a(n) exactly divides 3n. Or, 3-adic valuation of 3n.

Conjecture: a(n) = A007949(A000045(4*n)), all other 3-adic quadrisections A007949(A000045(.))=0. [Lengyel?]. - _R. J. Mathar_, Jun 28 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, _Gary W. Adamson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001560 | OEIS-A051159

Triangle read by rows: T(n, k) = binomial(n mod 2, k mod 2) * binomial(n div 2, k div 2), where 'div' denotes integer division.

T is the infinite lower triangular matrix for this sequence; define two others, U and V; let U(n,k)=e_k(-1,2,-3,...,(-1)^n n), where e_k is the k-th elementary symmetric polynomial, and let V be the diagonal matrix A057077 (periodic sequence 1,1,-1,-1). Clearly V^-1 = V. Conjecture: U = U^-1, T = U . V, T^-1 = V . U, and |T| = |U|. - _George Beck_, Dec 16 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Oct 14 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001561 | OEIS-A051179

a(n) = 2^(2^n) - 1.

I conjecture that, for n > 7, omega(a(n)) > omega(a(n-1)) > n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Alan DeKok (aland(AT)ox.org) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001562 | OEIS-A051195

T(2n+2,n), array T as in A050186; a count of aperiodic binary words.

Conjecture: -(n+2)*(n-1)*a(n) +2*(n+1)*(3*n-4)*a(n-1) +4*(-n^2+5*n-1)*a(n-2) -8*n*(3*n-4)*a(n-3) +16*(n-1)*(2*n-5)*a(n-4)=0. - _R. J. Mathar_, May 20 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001563 | OEIS-A051222

Numbers k such that Bernoulli number B_{k} has denominator 6.

Conjecture: When the expression (p+q^b)/2 is required to be prime, p is prime, and q is a prime >=5, then all p values are prime congruent to 1 (mod 12) (A068228), if and only if the exponent b is a member of this set. - _Richard R. Forberg_, Apr 07 2025 There are additional exponential expressions conjectured for generating each of several known prime subsequences (e.g., Pythagorean primes, A002144) where the sequence is invariant to the exponent, if and only if the exponent is a member of this set. See Forberg link. - _Richard R. Forberg_, Apr 25 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001564 | OEIS-A051250

Numbers whose reduced residue system consists of 1 and prime powers only.

Conjecture: the sequence is finite and 60 is the largest term, empirically verified up to 10^7;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001565 | OEIS-A051252

Number of essentially different ways of arranging numbers 1 through 2n around a circle so that sum of each pair of adjacent numbers is prime.

See A228917 for a similar conjecture on twin primes. - _Zhi-Wei Sun_, Sep 08 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001566 | OEIS-A051283

Numbers k such that if one writes k = Product p_i^e_i (p_i primes) and P = max p_i^e_i, then k/P > P.

Numbers k such that A346418(k) > 1 (conjectured). - _Amiram Eldar_, Jul 16 2021 Numbers k for which A354933(k) > A034699(k), i.e., where A354995(k) > 0 (conjectured).- _Antti Karttunen_, Jun 19 2022 a(n) = A080170(n) + 1 (conjectured). - _Ralf Stephan_, Feb 20 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001567 | OEIS-A051286

Whitney number of level n of the lattice of the ideals of the fence of order 2n.

The g.f. A(x) satisfies the differential equation (1-2*x-x^2-2*x^3+x^4)*A'(x) = (1+x+3*x^2-2*x^3)*A(x), from which the recurrence conjectured by Mathar follows. - _Emanuele Munarini_, Dec 18 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001568 | OEIS-A051397

a(n) = (2*n-2)*(2*n-1)*a(n-1)+1.

Conjecture: a(n) +(-4*n^2+6*n-3)*a(n-1) +2*(2*n-3)*(n-2)*a(n-2)=0. - _R. J. Mathar_, Jan 31 2014 Mathar's conjectured recurrence a(n) = (4*n^2 - 6*n + 3)*a(n-1) - (2*n - 3)*(2*n - 4)*a(n-2) follows easily from the defining recurrence. The sequence b(n) := (2*n - 1)! also satisfies Mathar's recurrence but with b(1) = 1, b(2) = 6. This leads to the continued fraction representation a(n) = (2*n - 1)!*(1 + 1/(6 - 6/(21 - 20/(43 - ... - (2*n - 3)*(2*n - 4)/(4*n^2 - 6*n + 3))))) for n >= 3. Taking the limit gives the continued fraction representation sinh(1) = A073742 = 1 + 1/(6 - 6/(21 - 20/(43 - ... - (2*n - 3)*(2*n - 4)/((4*n^2 - 6*n + 3) - ...)))). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aleksandar Petojevic_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001569 | OEIS-A051451

a(n) = lcm{ 1,2,...,x } where x is the n-th prime power (A000961).

Conjecture: For every n > 2, there exists a twin prime pair [p, p+2] with p < a(n), such that [a(n)+p, a(n)+p+2] is also a twin prime pair. Example: For n=6 we can take p=11, because for a(6) = 420 is [420+11, 420+13] = [431, 433] also a twin prime pair. This has been verified for 3 <= n <= 1000. - _Mike Winkler_, Sep 12 2013, May 19 2026 Conjecture: Every term = A001013(j)*A129912(k) for some j,k. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Dec 11 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001570 | OEIS-A051470

a(n) is least value of m for which the sum of Liouville's function from 1 to m is n.

It was once conjectured that the sum of Liouville's function was never > 0 except for the first term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001571 | OEIS-A051524

Second unsigned column of triangle A051338.

Conjecture: a(n) +(-2*n-9)*a(n-1) +(n+4)^2*a(n-2)=0. - _R. J. Mathar_, Aug 04 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001572 | OEIS-A051560

Second unsigned column of triangle A051379.

Conjecture: a(n) +(-2*n-13)*a(n-1) +(n+6)^2*a(n-2)=0. - _R. J. Mathar_, Aug 04 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001573 | OEIS-A051602

a(n) is the maximal number of squares that can be formed from n points in the plane.

Values for n <= 9 [now 16] are exact and are the same for a and b (see proofs below by Hugo van der Sanden and Benoît Jubin for n <= 7 and Sascha Kurz for n <= 9, which furthermore classify all optimal configurations for these values). Values for n > 9 are conjectural since they were obtained by exhaustive search for grid points within a square of side ceil(sqrt(n)), which is a reasonable assumption. A proof that optimal configurations (with gcd of side lengths equal to 1) have a diameter at most f(n) with f of moderate growth would permit exact computation of values from exhaustive searches. Conjecture: the asymptotic density of the numbers n such that there is no maximal arrangement formed by all the grid points within a suitably chosen circle, is 0. - _Peter Munn_, Sep 30 2021 For the known values of a(n) (n <= 17), there is a maximal arrangement formed using circles as specified by the conjecture above. For n <= 100 no arrangement has yet been found to contain more squares than the best attained using a circle as specified by A348469. - _Peter Munn_, Nov 12 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Erich Friedman_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001574 | OEIS-A051634

Strong primes: prime(k) > (prime(k-1) + prime(k+1))/2.

Conjecture: Limit_{N->oo} Sum_{n=1..N} (NextPrime(a(n))-a(n)) / a(N) = 1/4. [A heuristic proof is available at www.primepuzzles.net - Conjecture 91] - _Alain Rocchelli_, Nov 14 2022 Conjecture: Limit_{n->oo} n / PrimePi(a(n)) = 1/2. - _Alain Rocchelli_, Mar 17 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felice Russo_, Nov 15 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001575 | OEIS-A051635

Weak primes: prime(n) < (prime(n-1) + prime(n+1))/2.

Erdős called a weak prime an "early prime." He conjectured that there are infinitely many consecutive pairs of early primes, and offered $100 for a proof and $25000 for a disproof (Kuperberg 1992). See A229832 for a stronger conjecture. - _Jonathan Sondow_, Oct 13 2013 Conjecture: Limit_{n->oo} n / PrimePi(a(n)) = 1/2. - _Alain Rocchelli_, Mar 17 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felice Russo_, Nov 15 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001576 | OEIS-A051694

Smallest Fibonacci number that is divisible by n-th prime.

It is conjectured that a(n) is not divisible by prime(n)^2. See Remark on p. 528 of Wall and Conjectures in CNRS links. - _Michel Marcus_, Feb 24 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001577 | OEIS-A051699

Distance from n to closest prime.

Conjecture: S(n) = Sum_{k=1..n} a(k) is asymptotic to C*n*log(n) with C=0.29..... - _Benoit Cloitre_, Aug 11 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001578 | OEIS-A051754

Consider problem of placing N queens on an n X n board so that each queen attacks precisely 1 other. Sequence gives maximum number of queens.

Conjecture: a(n) = 2*floor(2*n/3) for n >= 6. - _Alexander D. Healy_, Feb 10 2024 Conjecture is true; see link to proof below. - _Alexander D. Healy_, Aug 23 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | Robert Trent (trentrd(AT)hotmail.com), Aug 23 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001579 | OEIS-A051756

Consider the problem of placing N queens on an n X n board so that each queen attacks precisely 3 others. Sequence gives maximal number of queens.

Conjecture: a(n) = 2[(6n-2)/5] for n >= 2; verified up to n = 100. - _Alexander D. Healy_, Feb 11 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | Robert Trent (trentrd(AT)hotmail.com), Aug 23 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001580 | OEIS-A051783

Numbers k such that 3^k + 2 is prime.

Conjecture: The number n = 3^k + 2 is prime if and only if 2^((n-1)/2) == -1 (mod n). - _Maheswara Rao Valluri_, Jun 01 2020. [Note that this is an if and only if assertion, so it does not follow from Fermat's Little Theorem. - _N. J. A. Sloane_, Sep 07 2020]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_, Dec 09 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001581 | OEIS-A051827

Number of numbers below 10^n with nonzero multiplicative digital root 7.

Conjectured o(10^n)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dan Hoey_, Dec 09 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001582 | OEIS-A051909

Subset of strict Egyptian numbers - there is a unique representation of 1 as the sum of distinct unit fractions with the sum of denominators being these numbers.

McCranie conjectured that the sequence is finite. He was right. Numbers 92-269 all have at least two representations without 15 appearing among the reciprocals (list attached). In the same way that Graham did for strict Egyptian numbers, we can transform each representation (a_n) into an even (2, (2a_n)) and an odd (3, 9, 30, 45, (2a_n)). Because there are no 15's in the original representations, the odd transformation has distinct unit fractions. The same is trivially true for the even transformation because there are no 1's either. So if the original sum is s, the new sums are 2s+2 and 2s+87. 2*92+2 = 186, so the even sums can be obtained from transformations after that. 2*92+87 = 271, so the same holds for odd sums from that point. And as the pairs of original representations were distinct, so are the transformed pairs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_, Dec 16 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001583 | OEIS-A051924

a(n) = binomial(2*n,n) - binomial(2*n-2,n-1); or (3n-2)*C(n-1), where C = Catalan numbers (A000108).

Conjecture: a(n) mod n^2 = n+2 iff n is an odd prime. - _Gary Detlefs_, Feb 19 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Barry E. Williams_, Dec 19 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001584 | OEIS-A051960

a(n) = C(n)*(3n+2) where C(n) = Catalan numbers = A000108.

(n+1)*a(n) - 2*(n+2)*a(n-1) - 4*(2*n-3)*a(n-2) = 0. - conjectured by _R. J. Mathar_, Oct 02 2014, verified by _Robert Israel_, Sep 30 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Barry E. Williams_, Jan 05 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001585 | OEIS-A052040

Numbers whose square is zeroless.

This sequence is infinite, since 33...334^2 = 11...11155...556, for example. This answers an open problem stated in HAKMEM. - _Karl W. Heuer_, Aug 19 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Dec 15 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001586 | OEIS-A052130

a(n) is the number of numbers between 1 and 2^m with m-n prime factors (counted with multiplicity), for m sufficiently large.

m is sufficiently large precisely when 2^m > 3^(m-n), i.e., when m >= floor(n*log(3)/log(1.5)) = A117630(n+1) = A126281(n) for n > 1. (_Robert G. Wilson v_ asks if this conjecture holds in a comment to A126281.) - _David A. Corneth_, Apr 09 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | Bernd-Rainer Lauber (br.lauber(AT)surf1.de), Jan 21 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001587 | OEIS-A052131

Decimal expansion of conjectured limit of ratio of successive terms of A052109.

Decimal expansion of conjectured limit of ratio of successive terms of A052109.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Robert Lozyniak (11(AT)onna.com), Jan 21 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001588 | OEIS-A052183

a(n) = (n + 2) * binomial(3*n, n) / (2*n + 1).

Conjecture: 2*n*(2*n+1)*a(n) + (-47*n^2+50*n-12)*a(n-1) + 15*(3*n-4)*(3*n-5)*a(n-2) = 0. - _R. J. Mathar_, Sep 27 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Barry E. Williams_, Jan 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001589 | OEIS-A052210

Numbers k such that k^3 starts with k itself (in base 10).

Replace the first term with 4, then add 1 to all the others to find numbers k where k^3 starts with k+2. Similar processes can be used for any k+2m. (conjectured) - _Dhilan Lahoti_, Aug 30 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Erich Friedman_, Jan 29 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001590 | OEIS-A052213

Numbers k with prime signature(k) = prime signature(k+1).

Sequence is not the same as A280074, first deviation is at a(212): a(212) = 2041, A280074(212) = 2024. Number 2024 is the smallest number n such that A007425(n) = A007425(n+1) with different prime signatures of numbers n and n+1 (2024 = 2^3 * 11 * 23, 2025 = 3^4 * 5^2; A007425(2024) = A007425(2025) = 90). Conjecture: also numbers n such that Product_{d|n} tau(d) = Product_{d|n+1} tau(d). - _Jaroslav Krizek_, Dec 25 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Erich Friedman_, Jan 29 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001591 | OEIS-A052227

a(n) = (4*n+1)*binomial(3*n,n)/(2*n+1).

Conjecture: 8*n*(2*n+1)*a(n) + 6*(-8*n^2-25*n+13)*a(n-1) - 45*(3*n-4)*(3*n-5)*a(n-2) = 0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Barry E. Williams_, Jan 29 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001592 | OEIS-A052276

Nonnegative numbers of the form n^3 (+/-) 3, n >= 0.

It is conjectured that A003325 and A052276 (the current sequence) have infinitely many numbers in common, although only one example (128) is known.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 05 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001593 | OEIS-A052362

Indices of records in length of English name of n including spaces and dashes (A052360): n such that k < n => A052360(k) < A052360(n).

I conjecture that from 103 on all d-digit terms have a smaller (most often but not always (d-1)-digit) term as suffix, and from 173 on they also have an earlier term as prefix. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Allan C. Wechsler_, Mar 07 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001594 | OEIS-A052379

Number of integers from 1 to 10^(n+1)-1 that lack 0 and 1 as a digit.

Conjecture: a(n) = A023001(n+2)-1. - _R. J. Mathar_, May 18 2007. Comment from _Vim Wenders_, Mar 26 2008: The conjecture is true: the g.f. leads to the closed form a(n) = -(8/7)*(1^n) + (64/7)*(8^n) = (-8 + 64*8^n)/7 = (-8 + 8^(n+2))/7 = (8^(n+2) - 1)/7 - 1 = A023001(n+2) - 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Odimar Fabeny_, Mar 12 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001595 | OEIS-A052427

Baxter-Hickerson numbers.

Pegg (1999) conjectured that the sequence of zeroless cubes (A052045) is finite. On April 19, 1999, Hickerson gave the counterexample: if n == 2 (mod 3) and n >= 5, then the cube of (2*10^(5*n) - 10^(4*n) + 17*10^(3*n-1) + 10^(2*n) + 10^n - 2)/3 is zeroless. Three days later, Baxter gave a simpler variation which is valid for all n>=0 and is given in the Formula section. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001596 | OEIS-A052470

Conjectured orders of sociable numbers A003416.

Conjectured orders of sociable numbers A003416. This is only conjectural, since we don't know that A003416 is complete (cf. A122726). - _N. J. A. Sloane_, Sep 17 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001597 | OEIS-A052542

a(n) = 2*a(n-1) + a(n-2), with a(0) = 1, a(1) = 2, a(2) = 4.

Conjecture: Apart from the initial 1, a(n) is the number of compositions of two types of n having no even parts. - _Gregory L. Simay_, Feb 17 2018 The above conjecture is true, as the g.f. can be written as 1/(1 - (2*x)/(1 - x^2)). - _John Tyler Rascoe_, Jun 01 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _INRIA Encyclopedia of Combinatorial Structures_, Jan 25 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001598 | OEIS-A052731

E.g.f. [1-x -sqrt(1-2x-3x^2)]/(2x) - [1+x-sqrt(1-2x-3x^2)]/2 .

Conjecture: a(n) = n!*A002026(n-2). - _R. J. Mathar_, Oct 16 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _INRIA Encyclopedia of Combinatorial Structures_, Jan 25 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001599 | OEIS-A052736

E.g.f. [1 -3x -sqrt(1-6x+x^2) -x*(1-x-sqrt(1-6x+x^2))]/2.

Conjecture: a(n) = n!*A006319(n-1). - _R. J. Mathar_, Oct 16 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _INRIA Encyclopedia of Combinatorial Structures_, Jan 25 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001600 | OEIS-A052870

First differences of A052829.

Conjectures: G.f. satisfies A(x) = exp(Sum_{k>=1} (-1)^(k+1) * A(x^k) * x^k/(k * (1 - x^k))).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _INRIA Encyclopedia of Combinatorial Structures_, Jan 25 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001601 | OEIS-A052895

Expansion of e.g.f. (1/2)/(exp(x) - 1) * (1 - (5 - 4*exp(x))^(1/2)).

Conjecture: for fixed k = 1,2,..., the sequence a(n) (mod k) is eventually periodic with the exact period dividing phi(k), where phi(k) is the Euler totient function A000010. For example, modulo 10 the sequence becomes (1, 1, 5, 3, 5, 1, 5, 3, 5, ...), with an apparent period 1, 5, 3, 5 of length 4 = phi(10) beginning at a(1). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _INRIA Encyclopedia of Combinatorial Structures_, Jan 25 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001602 | OEIS-A052952

a(n) = Fibonacci(n+2) - (1-(-1)^n)/2.

a(n) = floor(|F((1+i)*(n+2))|), n >= 0, with the complex Fibonacci function F: C -> C, z -> F(z) with F(z) := (exp(log(phi)*z) - exp(i*Pi*z)*exp(-log(phi)*z))/(2*phi-1) with the modulus |z|, the imaginary unit i and the golden section phi:=(1+sqrt(5))/2. A Conjecture: For F(z) see, e.g., the T. Koshy reference. ch. 45, p. 523, where F is called f, given in A000045. - _Wolfdieter Lang_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _INRIA Encyclopedia of Combinatorial Structures_, Jan 25 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001603 | OEIS-A052999

Take n-th prime p, let P(p) = all primes that can be obtained by permuting the digits of p and possibly adding or omitting zeros; a(n) = |p-q| where q in P(p) is the closest to p but different from p (a(n)=0 if no such q exists).

Conjecture: a(n) > 0 for n > 4. - _Sean A. Irvine_, Nov 23 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 16 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001604 | OEIS-A053004

Decimal expansion of AGM(1,sqrt(2)).

Conjecturally equals 1/(Sum_{n = -infinity..infinity} exp(-Pi*(n+1/2)^2))^2. Cf. A096427. - _Peter Bala_, Jun 10 2019 Conjecture: Equals Sum_{k>=0} A001934(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 21 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001605 | OEIS-A053027

Odd primes p with 2 zeros in Fibonacci numbers mod p.

Conjecturely, this sequence has density 1/3 in the primes. (End) [Comment rewritten by _Jianing Song_, Jun 16 2024 and Jun 25 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Feb 23 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001606 | OEIS-A053029

Numbers with 4 zeros in Fibonacci numbers mod m.

Conjecture: m is on this list iff m is an odd number all of whose factors are on this list or m is twice such an odd number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Feb 23 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001607 | OEIS-A053031

Numbers with 1 zero in Fibonacci numbers mod m.

Conjecture: m is on this list iff m is an odd number all of whose factors are on this list or m is 2 or 4 times such an odd number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Feb 23 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001608 | OEIS-A053033

Numbers which are the average of two primes in more ways than any smaller number.

Conjecture 1: This sequence is infinite. Conjecture 2: If this sequence is infinite, then for each prime number p > 2, there exists a minimum sufficiently large number k such that all terms >= k are multiples of p. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Len Smiley_, Feb 23 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001609 | OEIS-A053168

Hamming weights (or nonlinearity) of degree 4 rotation-symmetric functions.

T. W. Cusick and P. Stanica conjectured that the Hamming weight and the nonlinearity are the same for rotation-symmetric functions of degree 3. We conjecture that the same is true for rotation-symmetric functions of any degree. The conjecture is true for all such functions of degree >= 3 and at most 13 variables. - _Charlie Neder_, Feb 05 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | Pantelimon Stanica (pstanica(AT)mail.aum.edu), Feb 29 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001610 | OEIS-A053182

Primes p such that p^2 + p + 1 is prime.

Roger Horn computed the first 776 terms of this sequence around 1961 to test (with Paul Bateman) their conjecture on the density of simultaneous primes in polynomials. - _Charles R Greathouse IV_, Apr 05 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enoch Haga_, Mar 01 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001611 | OEIS-A053253

Coefficients of the '3rd-order' mock theta function omega(q).

By Campbell's conjecture above that a(n) is the number of partitions of 2n+1 with all odd parts and all odd conjugate parts, the a(0) = 1 through a(5) = 8 partitions are (B = 11):

Conjectural g.f.: 1/(1 - x)*(1 + Sum_{n >= 0} x^(3*n+1) /((1 - x)*(1 - x^3)*...*(1 - x^(2*n+1)))). - _Peter Bala_, Nov 18 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, Dec 19 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001612 | OEIS-A053254

Coefficients of the '3rd-order' mock theta function nu(q).

Conjecture: a(n) = (-1)^n * (A376628(n) + A376628(n+1)), or equivalently, (1 + q * nu(-q))/(1 + q) = Sum_{n >= 0} q^(n*(n+1))/((1 - q)*(1 - q^3)*...*(1 - q^(2*n-1))), the g.f. of A376628. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, Dec 19 1999 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001613 | OEIS-A053289

First differences of consecutive perfect powers (A001597).

Michel Waldschmidt writes: Conjecture 1.3 (Pillai). Let k be a positive integer. The equation x^p - y^q = k where the unknowns x, y, p and q take integer values, all >= 2, has only finitely many solutions (x,y,p,q). This means that in the increasing sequence of perfect powers [A001597] the difference between two consecutive terms [the present sequence] tends to infinity. It is not even known whether for, say, k=2, Pillai's equation has only finitely many solutions. A related open question is whether the number 6 occurs as a difference between two perfect powers. See Sierpiński [1970], problem 238a, p. 116. - _Jonathan Vos Post_, Feb 18 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Mar 03 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001614 | OEIS-A053319

Distance between the smaller members of successive twin prime pairs.

Conjecture: a(n) < log(A014574(n))^3 for n > 2. - _Thomas Ordowski_, Jul 21 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Mar 06 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001615 | OEIS-A053392

a(n) is the concatenation of the sums of every pair of consecutive digits of n (with a(n) = 0 for 0 <= n <= 9).

The Blomberg link asks whether a number n can have more than 10 predecessors, i.e. values of p such that a(p) = n. The answer is no, because there can be at most one predecessor ending with any given digit d. That can be proved by induction by observing that d and the last digit of n determine the last 1-or-2-digit sum in the concatenation of sums forming n, and hence the penultimate digit of p. That is either incompatible with the known value of n, or it tells us what the last digit of p' is, where p' = p with its last digit removed. We also know that p' is the predecessor of n' = n with its known last digit sum removed, and so we know there is at most one solution for p' by inductive hypothesis, and hence at most one solution for p. - _David J. Seal_, Nov 06 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 07 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001616 | OEIS-A053403

Consider the set P of pairs (a,b) generated by the rules: (1,1) is in P; if (a,b) is in P then (b,a+b) is in P; if (a,b) and (a',b') are in P then (aa', bb') is in P. Sequence gives numbers not appearing in P.

Sequence has 508 known terms, the largest of which is 55487. Conjecturally it is finite. If it is and 55487 is the largest term, then the function the number of groups of order n takes on all positive integers as values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | R. Keith Dennis (dennis(AT)math.cornell.edu), Jan 07 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001617 | OEIS-A053406

Number of stars brighter than visual magnitude n-1.

a(n) would be proportional to 10^(0.6*n) in the simplifying hypothesis of a uniform distribution of identical stars in an infinite universe. Any empirical formula matching the actual sequence for n=0 to 10 is welcome.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jean Fontaine (jfontain(AT)odyssee.net), Jan 08 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001618 | OEIS-A053482

Binomial transform of A029767.

Conjecture: a(n) -(3*n+1)*a(n-1) +(2*n+3)*(n-1)*a(n-2) -2*(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Sep 29 2012 a(n) = 3*n*a(n-2)-2*n*(n-1)*a(n-2)+1, derived from the array defined in the comment, which proves the previous conjecture. - _Richard Choulet_, Dec 17 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 15 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001619 | OEIS-A053532

Expansion of e.g.f.: (1-x)^(-1/2)*exp(-x/2 -x^2/4 -x^3/6).

Conjecture: D-finite with recurrence 2*a(n) +2*(-n+1)*a(n-1) -(n-1)*(n-2)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Jul 06 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 16 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001620 | OEIS-A053576

Smallest number whose Euler totient is divisible by 2^n.

Adamczewski provides a proof abstract regarding Greathouse's conjecture in linked article. - _Michael De Vlieger_, Aug 20 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jan 18 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001621 | OEIS-A053599

Number of nonempty subsequences {s(k)} of 1..n such that the difference sequence is palindromic.

Conjecture: let b(n) be the number of subsets S of {1,2,...,n} having more than one element such that (sum of least two elements of S) > max(S). Then b(0) = b(1) = 0 and b(n+2) = a(n+1) for n >= 0. - _Clark Kimberling_ Sep 27 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jan 19 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001622 | OEIS-A053606

a(n) = (Fibonacci(6*n+3) - 2)/4.

We prove the two McGarvey-CAMI conjectures mentioned at the beginning of the Comments section. Let, as usual, F(n) = A000045(n), the Fibonacci numbers. In the sequel we indicate equations with upper case letters ((A), (B), (C), (D)) for ease of reference.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _James Sellers_, Jan 20 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001623 | OEIS-A053617

Number of permutations of length n which avoid the patterns 1234 and 1324.

G.f. conjectured to be non-D-finite (see Albert et al. link). - _Jay Pantone_, Oct 01 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Moa Apagodu_, Mar 20 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001624 | OEIS-A053624

Highly composite odd numbers: odd numbers where d(n) increases to a record.

Conjecture: all terms after the first three terms are congruent to 5 mod 10. - _Harvey P. Dale_, Jul 05 2023 Dale's conjecture is correct. a(n) can't be even, since then a(n)/2 would be a smaller number with the same number of odd divisors. The respective powers of the successive odd primes can't increase, since if they did, swapping them would give a smaller number with the same number of divisors, e.g., 3^2 * 5^4 has the same number of divisors as 3^4 * 5^2, and the latter is smaller. As such, every a(n) must be an odd multiple of 5, hence congruent to 5 mod 10, unless it's simply a power of 3. But multiplying a power of 3 by 3 gives just one more divisor while multiplying a power of 3 by 5 doubles the number of divisors, so after a(n) = 9 all a(n) must be congruent to 5 mod 10, i.e., have a rightmost decimal digit of 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Stefano Lanfranco (lastefano(AT)yahoo.it), Mar 21 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001625 | OEIS-A053645

Distance to largest power of 2 less than or equal to n; write n in binary, change the first digit to zero, and convert back to decimal.

Conjecture: a(n) = (1 - A036987(n-1))*(1 + a(n-1)) for n > 1 with a(1) = 0. - _Mikhail Kurkov_, Jul 16 2019 Kurkov's conjecture is true (see Fried link). - _Sela Fried_, Nov 20 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Mar 22 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001626 | OEIS-A053695

Differences between record prime gaps.

Conjecture: a(n) <= A005250(n). Based on the equivalent statement at A005250: A005250(n+1) / A005250(n) <= 2. - _John W. Nicholson_, Dec 30 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeff Burch_, Mar 23 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001627 | OEIS-A053697

a(n+1)=a(n)+a^(n), where the addition is in base 11 and where a^(n) is obtained from a(n) by replacing each digit with its multiplicative inverse modulo 11. Zero digits, if any, are deleted.

Conjecture. For any positive integer a(1), the sequence generated according to the above rule eventually cycles through the forms a(k)=[1][4^a][3][(84)^b],..., a(k+6)=[1][4^a][3][(84)^(b+1)], or through a(k)=[1][5^a][4][(84)^b],..., a(k+6)=[1][5^a][4][(84)^(b+1)], for nonnegative integers a and b. The sequence listed above, with a(1)=1, is an example of the first type. Conjectures from _Colin Barker_, Feb 15 2016: (Start) Colin Barker's conjectures follow from these. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Feb 14 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001628 | OEIS-A053871

a(n) = 2*(n-1)*(a(n-1) + a(n-2)), starting a(0) = 1; a(1) = 0.

Harry Richman's conjecture in the Formula section was proved by Ahaan Kallat (2026, Theorem 59); see the paper in the Links section. - _Ahaan Kallat_, Oct 14 2026 Conjectured g.f.: T(0)/(1+x), where T(k) = 1 - x*(k+1)/(x*(k+1) - (1+x)/T(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 13 2013 Conjecture: if m == n (mod q) for q odd, then (-1)^m*a(m) == (-1)^n*a(n) (mod q). - _Harry Richman_, Aug 29 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | Cris Moore (moore(AT)santafe.edu) and _Christian G. Bower_, Mar 29 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001629 | OEIS-A053922

Numbers k such that k^2 contains only digits {2,4,6}.

Conjecture: every number composed of the numeral six repeated n times and ending in the numeral 8 is a term of this sequence. - _Harvey P. Dale_, Jun 16 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Mar 15 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001630 | OEIS-A054030

Sigma(k)/k for k such that sigma(k) is divisible by k.

The graph supports the conjecture that all numbers except 2 appear only a finite number of times. Sequences A000396, A005820, A027687, A046060 and A046061 give the n for which the abundancy sigma(n)/n is 2, 3, 4, 5 and 6, respectively. See A134639 for the number of n having abundancy greater than 2. - _T. D. Noe_, Nov 04 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Asher Auel_, Jan 19 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001631 | OEIS-A054096

T(n,2), array T as in A054090.

Conjecture: sequence is A006183 shifted right. - _Ralf Stephan_, Jan 15 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001632 | OEIS-A054109

a(n) = T(2*n+1, n), array T as in A054106.

Conjecture: (n+1)*a(n) + (-3*n-1)*a(n-1) + 2*(-2*n-1)*a(n-2) = 0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001633 | OEIS-A054116

a(n) = T(n,n-1), array T as in A054115.

Conjecture: a(n) - (n+1)*a(n-1) + n*a(n-2) = 0. - _R. J. Mathar_, Jun 13 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001634 | OEIS-A054117

T(n,n-2), array T as in A054115.

Conjecture: a(n) -(n+1)*a(n-1) +n*a(n-2)=0. - _R. J. Mathar_, Jun 13 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001635 | OEIS-A054118

Subdiagonal T(n,n-3), array T as in A054115.

Conjecture: a(n) = (n+1)*a(n-1) - n*a(n-2) for n>=6. - _R. J. Mathar_, Jun 13 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001636 | OEIS-A054120

Triangular array T(n,k): start with T(n,0)=T(n,n)=1 for n >= 0; recursively, draw vertical lines through T(n-1,k-1) if present and T(n-1,k) if present; then T(n,k) is the sum of T(i,j) that lie on or between the lines and not below T(n,k).

Conjecture: T(n,k) = T(n-1,k-1) + 2*T(n-2,k-1) + T(n-1,k) (except for T(0,0) = 1 and T(2,1) = 3 and assuming that T(n,k) = 0 for elements outside the triangular array). - _Gerald McGarvey_, Sep 20 2007 Conjecture: T(n,k) = A081577(n,k) - A081577(n-2,k-1). (A081577 is Pascal-(1,2,1) array). - _Gerald McGarvey_, Sep 20 2007 We prove the 1st McGarvey conjecture, (A) T(n,k)=T(n-1,k-1)+2*T(n-2,k-1)+T(n-1,k), with McGarvey's added modifications that n >= 3 and that T(n,k)=0 for n<0, k<0, and k>n. We make explicit that the rows and columns of the triangle start at 0 and furthermore, that both diagonals are at 45 degrees to the perpendicular (in both directions). Note that we use letters (A), (B), (C) to indicate equations.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001637 | OEIS-A054149

T(2n-1,n), array T as in A054144.

Conjectures from _Colin Barker_, Feb 11 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 18 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001638 | OEIS-A054271

Difference between prime(n)^2 and the previous prime.

Conjecture: every other even integer appears in the sequence an infinite number of times. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, May 05 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001639 | OEIS-A054272

Number of primes in the interval [prime(n), prime(n)^2].

These are precisely the primes available for the solution of Aguilar's conjecture or Haga's conjecture in Carlos Rivera's The Prime Puzzles and Problems Connection, (conjecture 26). Aguilar's conjecture states that at least one prime will be available for placement on each row and column of a p X p square array. Haga's conjecture states that just p primes are required for such placement in any p X p array. - _Enoch Haga_, Jan 23 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, May 05 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001640 | OEIS-A054341

Row sums of triangle A054336 (central binomial convolutions).

Conjecture: 2*(n+1)*a(n) +5*(-n-1)*a(n-1) +8*(-n+2)*a(n-2) +20*(n-2)*a(n-3)=0. - _R. J. Mathar_, Nov 30 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Mar 13 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001641 | OEIS-A054390

Number of ways of writing n as a sum of powers of 3, each power being used at most three times.

Conjecture: Number of ways of partitioning n into distinct parts of A038754. - _R. J. Mathar_, Mar 01 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, May 09 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001642 | OEIS-A054391

Number of permutations with certain forbidden subsequences.

Conjecture: Also the number of non-capturing set partitions of {1..n}. A set partition is capturing if it has two blocks of the form {...x...y...} and {...z...t...} where x < z and y > t or x > z and y < t. This is a weaker condition than nesting, so for example {{1,3,5},{2,4}} is capturing but not nesting. The a(0) = 1 through a(4) = 14 non-capturing set partitions are: The above conjecture is true: A partition is

non-capturing iff its representation in canonical sequential form avoids the patterns 1221 and 2112. In the context of these partition representations, the pattern 2112 is equivalent to the pattern 12112. Partitions whose canonical sequence form avoid 1221 and 12112 are one of the classes that are handled in the Mansour/Shattuck "Pattern Avoiding Partitions,..." paper. It shows that they are counted by this sequence. - _Christian Sievers_, Oct 29 2024 G.f.: 1/(1-x-x^2/(1-2x-x^2/(1-x-x^2/(1-x-x^2/(1-x-x^2/(1-x-x^2/(1-... (continued fraction) (conjecture). - _Paul Barry_, Jan 19 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Elisa Pergola (elisa(AT)dsi.unifi.it), May 21 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001643 | OEIS-A054408

a(n) = smallest positive integer not already in sequence such that the partial sum a(1)+...+a(n) is prime.

Conjecture: Every even number appears. - _N. J. A. Sloane_, May 29 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, May 09 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001644 | OEIS-A054414

a(n) = 1 + floor(n/(1-log(2)/log(3))).

Also, numbers n such that the first digit in ternary expansion on 2^n is 2. _N. J. A. Sloane_ conjectured that, for any integer n >=15, 2^n always has a 0 in its ternary expansion. - Mohammed Bouayoun (Mohammed.bouayoun(AT)sanef.com), Apr 24 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | B. Schaaf (m.m.schaaf-visch(AT)wxs.nl), May 20 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001645 | OEIS-A054465

Numbers k such that Sum_{j=1..k} d(j) is an integer where d(j) is the decimal fraction 0.2j (e.g., d(14) = 0.28).

Conjecture: the number of k > 0 such that a(k) <= n is asymptotic to C*log(n), C > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001646 | OEIS-A054531

Triangular array T read by rows: T(n,k) = n/gcd(n,k) (n >= 1, 1 <= k <= n).

Read as a linear sequence, this is conjectured to be the length of the shortest cycle of pebble-moves among the partitions of n (cf. A201144). - _Andrew V. Sutherland_, Nov 27 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 09 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001647 | OEIS-A054639

Queneau numbers: numbers n such that the Queneau-Daniel permutation {1, 2, 3, ..., n} -> {n, 1, n-1, 2, n-2, 3, ...} is of order n.

a(n) >> n log n, and on the Bateman-Horn-Stemmler conjecture a(n) << n log^2 n. I imagine a(n) ≍ n log n, and numerics suggest that perhaps a(n) ~ kn log n for some constant k (which seems to be around 1.122). - _Charles R Greathouse IV_, Aug 02 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | Gilles Esposito-Farese (gef(AT)cpt.univ-mrs.fr), May 17 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001648 | OEIS-A054656

Number of primes <= n which do not appear in any partitions of n into distinct primes.

Conjecture: if n >= 23 then a(n)=2 if both (n-6) and (n-4) are prime, a(n)=1 if one of (n-6), (n-4) or (n-1) is prime, a(n)=0 otherwise

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Apr 17 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001649 | OEIS-A054723

Prime exponents of composite Mersenne numbers.

Assuming a conjecture of Dickson, we can prove that this sequence is infinite. See Ribenboim. - _T. D. Noe_, Jul 30 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeppe Stig Nielsen_, Apr 20 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001650 | OEIS-A054726

Number of graphs with n nodes on a circle without crossing edges.

Conjecture: Also the number of simple graphs with vertices {1..n} not containing any pair of nesting edges. Two edges {a,b}, {c,d} where a < b and c < d are nesting if a < c and b > d or a > c and b < d. For example, the a(0) = 1 through a(3) = 8 non-nesting edge-sets are:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Flajolet_, Apr 20 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001651 | OEIS-A054727

Number of forests of rooted trees with n nodes on a circle without crossing edges.

Conjecture: -2(37*n-80) *(n-1) *(2*n-1) *a(n) +2*(592*n^3-3056*n^2+5045*n-2665) *a(n-1) +2*(148*n^3-986*n^2+2021*n-1255) *a(n-2) -5*(n-5) *(n-2) *(37*n-43) *a(n-3)=0. - _R. J. Mathar_, Apr 30 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Flajolet_, Apr 20 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001652 | OEIS-A054763

Residues of consecutive prime differences modulo 6.

For n>2, only the 0-residues may arise several times after each other, that is, there are no "2,2" and no "4,4". Let nz(k) denote the nonzero values of A054763(n). Then nz(0)=1, nz(1)=2, nz(2)=2, and nz(k+1)=6-nz(k) for k>1. Conjecture: the percentage of zeros in A054763(n) asymptotically runs to 50%. - _Alex Ratushnyak_, Apr 18 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, May 17 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001653 | OEIS-A054767

Period of the sequence of Bell numbers A000110 (mod n).

a(2^s) = 3*2^s for s >= 2 (Theorem 6.4 in the Lunnon article). For an odd prime p, if a(p) = (p^p-1)/(p-1) (which is conjectured to hold for all p), then a(p^s) = p^(s-1)*(p^p-1)/(p-1) (Theorem 6.2 in the Lunnon article). - _Jianing Song_, Jun 18 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Feb 09 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001654 | OEIS-A054770

Numbers that are not the sum of distinct Lucas numbers 1,3,4,7,11, ... (A000204).

Conjecture: this is the sequence of numbers for which the base phi representation includes phi itself, where phi = (1 + sqrt(5))/2 = the golden ratio. Example: let r = phi; then 6 = r^3 + r + r^(-4). - _Clark Kimberling_, Oct 17 2012 This conjecture is proved in my paper 'Base phi representations and golden mean beta-expansions', using the formula by Wilson/Agol/Carlitz et al. - _Michel Dekking_, Jun 25 2019 a(n) = floor(((5+sqrt(5))/2)*n)-1 (conjectured by _David W. Wilson_; proved by Ian Agol (iagol(AT)math.ucdavis.edu), Jun 08 2000)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Antreas P. Hatzipolakis (xpolakis(AT)otenet.gr), May 28 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001655 | OEIS-A054783

(n^2)-th Fibonacci number.

The sequence (5*a(n+1))_{n>=1} = (5, 15, 170, 4935, ...) is realizable in the sense that there is a self-map on a set T:X->X with the property that a(n) = #{x in X:T^nx=x} for all n >= 1. This is the simplest illustrative example of two different phenomena. The Fibonacci sequence sampled along an odd power cannot be made realizable after multiplication by a constant; the Fibonacci sequence sampled along an even power becomes realizable after multiplication by 5 (the discriminant of the sequence). This is now known to be an instance of a more general phenomenon in the following sense. If (a(n)) is a linear recurrence sequence whose characteristic polynomial F has simple zeros then the sequence (Ma(n^s)) satisfies the Dold congruence, where M=|discriminant(F)| and s is an integer multiple of the exponent of the Galois group of the splitting field of F over the rationals. Under an additional hypothesis on the signs of the coefficients of F, the sequence (Ma(n^s)) is realizable. - _Thomas Ward_, May 06 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeff Burch_, May 22 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001656 | OEIS-A054872

Number of (12345, 13245, 21345, 23145, 31245, 32145)-avoiding permutations.

Conjectured to be the number of permutations of length n+1 avoiding the partially ordered pattern (POP) {5>1, 1>2, 1>3, 1>4} of length 5. That is, conjectured to be the number of length n+1 permutations having no subsequences of length 5 in which the fifth element is the largest and the first element is the next largest - _Sergey Kitaev_, Dec 13 2020 This conjecture has been proven. There are six sets of permutations avoiding six size five permutations including the two sets discussed in this sequence that are known to match this sequence. A further two are conjectured to match this sequence. - _Christian Bean_, Jul 23 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | Elisa Pergola (elisa(AT)dsi.unifi.it), May 26 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001657 | OEIS-A054906

Smallest number x such that sigma(x+2n) = sigma(x)+2n (first definition).

The 3 definitions are identical or conjectured to be identical.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, May 23 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001658 | OEIS-A054925

a(n) = ceiling(n*(n-1)/4).

Conjecture: For n>11, there always exists a prime number p such that a(n)<p<=a(n+1). - _Raul Prisacariu_, Sep 01 2024 Conjecturally, apart from the first term, the sequence terms are the exponents in the expansion of Sum_{n >= 0} q^(3*n) * (Product_{k >= 2*n+1} 1 - q^k) = 1 - q - q^2 + q^3 + q^5 - q^8 - q^11 + + - - .... Cf. A039825. - _Peter Bala_, Apr 13 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 24 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001659 | OEIS-A054977

a(0)=2, a(n)=1 for n >= 1.

Arises in Gilbreath-Proth conjecture; see A036262.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Gould_, May 29 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001660 | OEIS-A054978

Obtained from sequence of lucky numbers (A000959) by taking iterated absolute value differences of terms and extracting the leading diagonal.

The classical Gilbreath-Proth Conjecture is that when iterated absolute differences are formed from the sequence of primes, the leading diagonal is 2,1,1,1,1,1,1,1,1,... (see A036262). This is an analog for the lucky numbers sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Gould_, May 29 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001661 | OEIS-A054979

e-perfect numbers: numbers k such that the sum of the e-divisors (exponential divisors) of k equals 2*k.

Conjecture: Every e-perfect number is divisible by 36, see A219016. - _Jon Perry_, Nov 13 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_, May 29 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001662 | OEIS-A054988

Number of prime divisors of 1 + (product of first n primes), with multiplicity.

It is an open question as to whether omega(p#+1) = bigomega(p#+1) = a(n); that is, as to whether the Euclid numbers are squarefree. Any square dividing p#+1 must exceed 2.5*10^15 (see Vardi, p. 87). - _Sean A. Irvine_, Oct 21 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | Arne Ring (arne.ring(AT)epost.de), May 30 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001663 | OEIS-A055008

Numbers k such that gcd(phi(k), sigma(k)) = 1 with phi = A000010, sigma = A000203.

Conjecture: Every term is a square or twice a square. - _Jason Yuen_, May 16 2024 The conjecture is true: If k is neither a square nor twice a square (i.e., in A028983), then sigma(k) is even. Since gcd(phi(k), sigma(k)) = 1, then phi(k) must be odd, but phi(k) is odd only for k = 1 and 2. - _Amiram Eldar_, May 19 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, May 31 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001664 | OEIS-A055023

a(n) = n/A055032(n).

It is conjectured that this is n iff n is 1 or a prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 11 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001665 | OEIS-A055030

(Sum(m^(p-1),m=1..p-1)+1)/p as p runs through the primes.

It is conjectured that (Sum(m^(n-1),m=1..n-1)+1)/n is an integer iff n is 1 or a prime. Always an integer from little Fermat theorem. Converse is conjectured to be true: if p | (1+1^(p-1)+2^(p-1)+3^(p-1)+...+(p-1)^(p-1)) and p > 1, then p is prime. That was checked by Giuga up to p <= 10^1000. [_Benoit Cloitre_, Jun 09 2002]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 11 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001666 | OEIS-A055032

Denominator of (Sum(m^(n-1),m=1..n-1)+1)/n.

It is conjectured that this is 1 iff n is 1 or a prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 11 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001667 | OEIS-A055040

Numbers of the form 3^(2i+1)*(3*j+2).

G.f.: [x(x+2)(x^2+x+1)(x^7+x^3+1)]/(x^11-x^10-x+1) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 01 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001668 | OEIS-A055078

Write n as a sum of terms of the form (p^2-1)/24 where p is a prime > 4; sequence gives those n which require at least 4 terms.

Chowla conjectures that all numbers are the sum of no more than four terms of the form (p^2-1)/24 where p is a prime > 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_, Jun 20 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001669 | OEIS-A055170

n-th distinct number to appear in A055168; also the n-th to appear in A217760.

Conjecture: this sequence is a permutation of the nonnegative integers. The conjecture is true: the number 0 appears in every segment of A055168, and, for n > 0, n appears in the (n+1)-th segment (as the number of occurrences of 0 in the previous segments). - _Rémy Sigrist_, Oct 16 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001670 | OEIS-A055171

Cumulative counting sequence: method B (noun,adjective)-pairs with first term 1.

Conjecture: every positive integer occurs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001671 | OEIS-A055173

n-th distinct number to appear in A055171; also the n-th to appear in A055187.

Conjecture: this sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001672 | OEIS-A055174

Cumulative counting sequence: method B (noun,adjective)-pairs with first term 2.

Conjecture: every positive integer occurs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001673 | OEIS-A055176

n-th distinct number to appear in A055174; also the n-th to appear in A055190.

Conjecture: this sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001674 | OEIS-A055177

Cumulative counting sequence: method B (noun,adjective)-pairs with first term 3.

Conjecture: every positive integer occurs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001675 | OEIS-A055179

n-th distinct number to appear in A055171; also the n-th to appear in A055191.

Conjecture: this sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001676 | OEIS-A055180

Cumulative counting sequence: method B (noun,adjective)-pairs with first term 4.

Conjecture: every positive integer occurs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001677 | OEIS-A055182

n-th distinct number to appear in A055174.; also the n-th to appear in A055192.

Conjecture: this sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001678 | OEIS-A055183

Cumulative counting sequence: method B (noun,adjective)-pairs with first term 5.

Conjecture: every positive integer occurs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001679 | OEIS-A055185

n-th distinct number to appear in A055177; also the n-th to appear in A055194.

Conjecture: this sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001680 | OEIS-A055186

Cumulative counting sequence: method A (adjective-before-noun)-pairs with first term 0.

For the original version, where "increasing order..." is "order of 1st appearance", see A217760. The conjecture formerly placed here applies to A217760. - _Clark Kimberling_, Mar 24 2013 Conjectures: a(n) < 2*sqrt(n); limit as n goes to infinity Max(a(k) : 1<=k<=n)/sqrt(n) exist = 2. - _Benoit Cloitre_, Jan 28 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001681 | OEIS-A055187

Cumulative counting sequence: method A (adjective-before-noun)-pairs with first term 1.

Conjectures: limit as n goes to infinity of max {a(k) : 1<=k<=n}/sqrt(n) = 2;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001682 | OEIS-A055188

Cumulative counting sequence: method A (adjective-before-noun) pairs, starting with 2.

Segments (generated as at A217760): 2; 1,2; 2,2,1,1; 4,2,3,1; 5,2,4,1,1,4,1,3; ... The 4th segment is read "4 2's and 3 1's" as a record of what was previously written; 4,2 are adjectives and 2 1 are nouns. Conjecture: every nonnegative integer occurs. - _Clark Kimberling_, Mar 25 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001683 | OEIS-A055189

Cumulative counting sequence: method A (adjective-before-noun)-pairs, starting with 3.

The 4th segment is read "3 3's and 3 1's and 1 2" as a record of what was previously written; 3 3 1 are adjectives and 3 1 2 are nouns. Conjecture: every nonnegative integer occurs. - _Clark Kimberling_, Mar 25 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001684 | OEIS-A055190

Cumulative counting sequence: method A (adjective-before-noun) pairs, starting with 4.

The 4th segment is read "3 4's and 3 1's and 1 2" as a record of what was previously written; 3,3,1 are adjectives and 4 1 2 are nouns. Conjecture: every positive integer occurs. - _Clark Kimberling_, Mar 25 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001685 | OEIS-A055191

Cumulative counting sequence: method A (adjective-before-noun) pairs, starting with 5.

Segments (generated as at A217760): 5; 1,5; 2,5,1,1; 3,5,3,1,1,2; ... The 4th segment is read "3 5's and 3 1's and 1 2" as a record of what was previously written; 3,3,1 are adjectives and 5 1 2 are nouns. Conjecture: every nonnegative integer occurs. - _Clark Kimberling_, Mar 25 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001686 | OEIS-A055203

Number of different relations between n intervals on a line.

Conjectures: a(n) == 1 (mod 12); for fixed k, the sequence a(n) (mod k) eventually becomes periodic with exact period a divisor of phi(k), where phi(k) is Euler's totient function A000010. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Sylviane R. Schwer (schwer(AT)lipn.univ-paris13.fr), Jun 22 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001687 | OEIS-A055211

Lesser Fortunate numbers.

a(1) is not defined. The first 1000 terms are all prime and it is conjectured that all terms are primes. Conjecture: Limit_{N->oo} (Sum_{n=2..N} a(n)) / (Sum_{n=2..N} prime(n)) = 3/2. - _Alain Rocchelli_, Nov 07 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jul 04 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001688 | OEIS-A055229

Greatest common divisor of largest square dividing n and squarefree part of n.

a(n) = core(n)*rad(n/core(n))/rad(n), where core = A007913 and rad = A007947. - Conjecture by _Velin Yanev_, proof by _David J. Seal_, Sep 19 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jun 21 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001689 | OEIS-A055231

Powerfree part of n: product of primes that divide n only once.

a(n*m) = a(n)*a(m)/(gcd(n,a(m))*gcd(m,a(n))) for all n and m > 0 (conjectured). - _Velin Yanev_, Feb 06 2019. [This follows easily from the comment of _Vladeta Jovovic_. - _N. J. A. Sloane_, Mar 14 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jun 21 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001690 | OEIS-A055234

Smallest x such that sigma(x) = n*phi(x), or -1 if no such x exists.

Conjecture: For each n, a(n) > 0. - _Farideh Firoozbakht_, Sep 12 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_, Jun 21 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001691 | OEIS-A055246

At step number k >= 1 the 2^(k-1) open intervals that are erased from [0,1] in the Cantor middle-third set construction are I(k,n) = (a(n)/3^k, (1+a(n))/3^k), n=1..2^(k-1).

Conjecture: Similarly to A191107, this increasing sequence is generated by the rules: a(1) = 1, and if x is in the sequence, then 3*x-2 and 3*x+4 are also in the sequence. - _L. Edson Jeffery_, Nov 17 2015 Conjecture: a(n) = 2*A191107(n) - 1 = 6*A003278(n) - 5 = (a((2*n-1)*2^(k-1))+2)/3^k, k>0. - _L. Edson Jeffery_, Nov 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, May 23 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001692 | OEIS-A055272

First differences of 7^n (A000420).

Conjecture in "Introduction à la théorie des nombres" by J. M. Deconinck and Armel Mercier: this is the period length of the fraction 1/7^n. For example 1/7^2=0.02040816326530612244897959183673469387755102040....with a period of 42 digits =6*7=a(2). The period of 1/7^3 has exactly 294=a(3) digits. - _Benoit Cloitre_, Feb 02 2002 For n >= 1, the conjecture of De Koninck and Mercier cited in the comment of Feb 02 2002 holds: a(n) is the period length of the decimal expansion of 1/7^n. Indeed, 10 is a primitive root modulo 7 (see A001913), and 10^6 == 8 (mod 49), so 7 is not in A045616. By lifting theorem for primitive roots, 10 is a primitive root modulo 7^n for every n >= 1. Consequently, the multiplicative order of 10 modulo 7^n is phi(7^n) = 6*7^(n-1) = a(n). Since gcd(10, 7^n) = 1, the decimal expansion of 1/7^n is purely periodic with this period length. - _Eiver Rodríguez_, Sep 05 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Barry E. Williams_, May 28 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001693 | OEIS-A055278

Number of rooted trees with n nodes and 3 leaves.

G.f.: x^4*(x^3+x^2+1)/((1-x^2)*(1-x^3)*(1-x)^3) (conjectured). - _Ralf Stephan_, Mar 07 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, May 09 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001694 | OEIS-A055342

Number of mobiles (circular rooted trees) with n nodes and 4 leaves.

Conjectured g.f.: x^5*(-1-x-5*x^2-6*x^3-9*x^4-5*x^5-2*x^6-2*x^7+x^9) / ((x^2+1)*(1+x+x^2) *(1+x)^3 *(x-1)^7). - _R. J. Mathar_, Sep 18 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, May 14 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001695 | OEIS-A055356

Triangle of increasing mobiles (circular rooted trees) with n nodes and k leaves.

Conjecture: row polynomials are R(n-2,0) for n > 1 where R(n,k) = R(n-1,k+1) + x*Sum_{i=0..n-1} Sum_{j=0..k} binomial(n-1, i)*R(n-i-1,j)*R(i,k-j) for n > 0, k >= 0 with R(0,k) = 1 for k >= 0. - _Mikhail Kurkov_, Apr 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, May 15 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001696 | OEIS-A055392

Number of bracketings of 0#0#0#...#0 giving result 0, where 0#0 = 1, 0#1 = 1#0 = 1#1 = 0.

Conjecture: 500*n*(n-1)*a(n) +100*(n-1)*(5*n -12)*a(n-1) +20*(25*n^2 -463*n +846)*a(n-2) +(-140161*n^2 +966559*n -1637508)*a(n-3) +2*(250*n^2 -26509*n +105084)*a(n-4) +98036*(4*n -19)*(4*n -21)*a(n-5) = 0. - _R. J. Mathar_, Nov 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeppe Stig Nielsen_, Jun 24 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001697 | OEIS-A055417

Number of points in N^n of norm <= 2.

a(n) = (n^3 - 3*n^2 + 14*n + 24)*(n+1)/24. Proof: The coordinates of such a point are a permutation of one of the vectors (0, ..., 0), (0, ..., 0, 1), (0, ..., 0, 2), (0, ..., 0, 1, 1), (0, ..., 0, 1, 1, 1), or (0, ..., 0, 1, 1, 1, 1), so the number of points is 1 + n + n + binomial(n,2) + binomial(n,3) + binomial(n,4). - Formula conjectured by _Frank Ellermann_, Mar 16 2002 and explained by _Michael Somos_, Apr 25 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001698 | OEIS-A055427

Number of points in Z^n of norm <= 3.

Above conjectures confirmed by later additions of b-file from Andrew Howroyd and program from _Jean-François Alcover_ with connection to A302997. - _Ray Chandler_, Jun 27 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001699 | OEIS-A055428

Number of points in Z^n of norm <= 4.

Above conjecture confirmed by later additions of b-file from Andrew Howroyd and program from _Jean-François Alcover_ with connection to A302997. - _Ray Chandler_, Jun 27 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001700 | OEIS-A055469

Primes of the form k(k+1)/2+1 (i.e., central polygonal numbers, or one more than triangular numbers).

It is conjectured that this sequence is infinite. - _Daniel Forgues_, Apr 21 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Jun 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001701 | OEIS-A055496

a(1) = 2; a(n) is smallest prime > 2*a(n-1).

Conjecture: lim_{n->infinity} a(n)/A006992(n) = 5.1648264... - _Thomas Ordowski_, Apr 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 07 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001702 | OEIS-A055573

Number of terms in simple continued fraction for n-th harmonic number H_n = Sum_{k=1..n} (1/k).

Conjecture: limit n -> infinity a(n)/n = 12*log(2)/Pi^2 = 0.84..... = A089729 Levy's constant. - _Benoit Cloitre_, Jan 17 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Jul 10 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001703 | OEIS-A055615

a(n) = n * mu(n), where mu is the Möbius function A008683.

Conjectures: lim b->1+ Sum n=1..inf a(n)*b^(-n) = -12 and lim b->1- Sum n=1..inf a(n)*b^n = -12 (+ indicates that b decreases to 1, - indicates it increases to 1), both considering that zeta(-1) = -1/12 and calculations (more generally mu(n)*n^s is Abel summable to zeta(-s)). - _Gerald McGarvey_, Sep 26 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Jun 04 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001704 | OEIS-A055836

T(2n+2, n), where T is the array in A055830.

Conjecture: 5*n*(n+1)*(7*n-5)*a(n) - n*(154*n^2+2*n-77)*a(n-1) - 3*(3*n-4)*(7*n+2)*(3*n-2)*a(n-2) = 0. - _R. J. Mathar_, Mar 13 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 28 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001705 | OEIS-A055837

T(2n+3,n), where T is the array in A055830.

Conjecture: 5*n*(n+2)*(11*n-4)*a(n) +(-242*n^3-330*n^2+29*n+42)*a(n-1) -3*(3*n-1)*(11*n+7)*(3*n-2)*a(n-2)=0. - _R. J. Mathar_, Mar 13 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 28 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001706 | OEIS-A055838

T(2n+4,n), where T is the array in A055830.

Conjecture: 5*n*(n+3)*(n-1)*a(n) -2*(n-1)*(11*n+8)*(n+2)*a(n-1) -3*(3*n-1)*(3*n-2)*(n+1)*a(n-2)=0. - _R. J. Mathar_, Mar 13 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 28 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001707 | OEIS-A055938

Integers not generated by b(n) = b(floor(n/2)) + n (complement of A005187).

Conjecture: a(n)=b(n) defined as b(1)=2, for n>1, b(n+1)=b(n)+1 if n is already in the sequence, b(n+1)=b(n)+3 otherwise. If so, then see Cloitre comment in A080578. - _Ralf Stephan_, Dec 27 2013 a(n) = A080578(n+1) - 2 = A080468(n+1) + 2*n (conjectured). - _Ralf Stephan_, Dec 27 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alford Arnold_, Jul 21 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001708 | OEIS-A056053

a(n) = smallest odd number 2m+1 such that the partial sum of the odd harmonic series Sum_{j=0..m} 1/(2j+1) is > n.

a(n) = A092315(n)*2 + 1 = floor(exp(n*2-Euler)/4+1/8)*2+1 for all n (conjectured). - _M. F. Hasler_, Jan 24 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jul 25 2000 and Jan 11 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001709 | OEIS-A056084

Numbers k such that k^8 == 1 (mod 9^3).

a(n) = (1458*n-729+(-1)^n*725)/4. - Conjectured by _Philipp Emanuel Weidmann_, explained by _Robert Israel_, Apr 13 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jun 08 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001710 | OEIS-A056142

Concatenate n, floor[n/10], floor[n/100] ... (but do not continue if floor[.]=0).

Conjecture: if n has 3 or more digits, a(n) is a palindrome only if all the digits of n are the same. It is easy to see that any palindrome can have at most 2 distinct digits: matching digits from the initial n in the concatenation matches each digit after the second with an earlier digit. - _Franklin T. Adams-Watters_, Sep 07 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Jun 15 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001711 | OEIS-A056243

Third diagonal of triangle A056242.

Conjecture: a(n) = 2^(-6+n)*(32-35*n+9*n^2). G.f.: x^3*(1+3*x-x^2)/(1-2*x)^3. - _Colin Barker_, Mar 20 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Mallows_, Aug 23 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001712 | OEIS-A056541

a(n) = 2*n*a(n-1) + 1 with a(0)=0.

Conjecture: a(n) +(-2*n-1)*a(n-1) +2*(n-1)*a(n-2)=0. - _R. J. Mathar_, May 29 2013 Mathar's conjecture is true. Proof: back the name expression up one step to get a(n-1) = 2*(n-1)*a(n-2) + 1. Substituting a(n-1) - 2*(n-1)*a(n-2) for 1 in the name expression and rearranging yields Mathar's form directly. - _Bill McEachen_, May 09 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Jun 20 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001713 | OEIS-A056608

Least prime factor of the n-th composite number.

Record values are seen when n = A120389(m). Conjecture: at each new record the count of the prior record follows A247509. Records seen are 2, 3, 5, 7, 11, ... and when 3, 5, 7, 11 are first seen, there have been 3, 3, 2, and 4 occurrences of 2, 3, 5, and 7. These are A247509(1) through A247509(4). Thus, the count for prime(60) would be A247509(60). - _Bill McEachen_, Jun 17 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Odimar Fabeny_, Aug 07 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001714 | OEIS-A056737

Minimum nonnegative integer m such that n = k*(k+m) for some positive integer k.

Conjecture: There exists some constant, k, approximately equal to 1.7, such that a(n) is of average order k*n/log(n). See Tooth Link for evidence. - _Clive Tooth_, Mar 18 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Aug 26 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001715 | OEIS-A056761

Odd numbers less than the cube of their number of divisors.

Last term is a(267) = 883575, confirming the author's conjecture. - _Charles R Greathouse IV_, Apr 27 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Aug 16 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001716 | OEIS-A056777

Composite numbers k such that both phi(k+12) = phi(k) + 12 and sigma(k+12) = sigma(k) + 12.

It is easy to show that if p, p+2, p+6 and p+8 are all prime (a prime quadruple as defined in A007530, which lists the values of p) with x=p(p+8), x+12=(p+2)(p+6), then x is in the sequence. I conjecture that all members of the sequence are of this form. - _Jud McCranie_, Oct 11 2000 The conjecture that all terms arise from prime quadruplets, and that all terms are congruent to 65 (mod 72), has been verified up to 10^12. - _Himaghna Roy Choudhury_, Jun 05 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Aug 17 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001717 | OEIS-A056793

Number of divisors of lcm(1..n).

Conjectured partial sums of A101207. - _Sean A. Irvine_, Jun 25 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Aug 28 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001718 | OEIS-A056826

Primes p such that (p^p + 1)/(p + 1) is a prime.

Let b(0) = 1, b(1) = 3; b(n+1) = (b(n)^2 + b(n-1)^2)/2. Conjecture: if b(n) = p is prime, then (p^p+1)/(p+1) is prime. Note that b(1) = 3, b(2) = 5, b(3) = 17, b(4) = 157 and b(9) is also prime. - _Thomas Ordowski_, Jul 29 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Aug 29 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001719 | OEIS-A056851

Integers n such that the number of digits in n! is a cube.

Ed Pegg Jr conjectures that n^3 - n = k! has a solution if and only if n is 2, 3, 5 or 9 (when k is 3, 4, 5 and 6).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Aug 30 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001720 | OEIS-A056856

Triangle of numbers related to rooted trees and unrooted planar trees.

Conjecture: T(n,k) = A130534(n,k)* n^(k-1). - _R. J. Mathar_, Mar 31 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _F. Chapoton_, Aug 31 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001721 | OEIS-A056888

a(n) = number of k such that sum of digits of 9^k is 9n.

All terms except the first are only conjectures. For the theorem that a(n) is always finite, see Senge-Straus and Stewart. - _N. J. A. Sloane_, Jan 06 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 05 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001722 | OEIS-A056921

a(0) = 0, a(1) = 1, a(2*n) = n*a(2*n-1) + a(2*n-2), a(2*n+1) = a(2*n) + a(2*n-1).

Conjecture: for n >= 1, a(n-1) is not divisible by n, except at n=5 (verified for n <= 10^7). - _Davide Oliveri_, Oct 03 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aleksandar Petojevic_, Sep 05 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001723 | OEIS-A056929

Difference between n^2 and average of smallest prime greater than n^2 and largest prime less than n^2.

Conjecture: the most frequent value will be 1 (including sequence variants with any even power n^2k). - _Bill McEachen_, Dec 12 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Jul 12 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001724 | OEIS-A057005

Number of conjugacy classes of subgroups of index n in free group of rank 2.

Response from Timothy R. Walsh, Aug 01 2012: The conjecture in the previous comment is true. A combinatorial map is a connected graph, loops and multiple edges allowed, in which a cyclic order of the incident edge-ends is assigned to every node. The equivalence between combinatorial maps and topological maps was conjectured by several researchers and finally proved by Jones and Singerman. In my 1975 paper "Generating nonisomorphic maps without storing them", I established a genus-preserving bijection between hypermaps with n darts, w vertices and b edges and properly bicolored bipartite maps with n edges, w white vertices and b black vertices. A bipartite map can't have any loops; so a combinatorial bipartite map is a multigraph and it suffices to impose a cyclic order of the edges, rather than the edge-ends, incident to each node. Thus it is just the child's drawing described above by Liskovets.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 09 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001725 | OEIS-A057015

Numbers k such that k^2 + 1, k^2 + 3, k^2 + 7 and k^2 + 9 are all primes.

Conjecture: all terms after the first term are equal to zero mod 10. - _Harvey P. Dale_, Sep 04 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Sep 09 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001726 | OEIS-A057030

Let P(n) of a sequence s(1),s(2),s(3),... be obtained by leaving s(1),...,s(n-1) fixed and reversing every n consecutive terms thereafter; apply P(2) to 1,2,3,... to get PS(2), then apply P(3) to PS(2) to get PS(3), then apply P(4) to PS(3), etc. The limit of PS(n) is A057030.

Conjecture: a(n) = A007062(n+1) - 1 for n > 0. - _Mikhail Kurkov_, Mar 10 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 29 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001727 | OEIS-A057032

Let P(n) of a sequence s(1), s(2), s(3), ... be obtained by leaving s(1), ..., s(n-1) fixed and forward-cyclically permuting every n consecutive terms thereafter; apply P(2) to 1, 2, 3, ... to get PS(2), then apply P(3) to PS(2) to get PS(3), then apply P(4) to PS(3), etc. The limit of PS(n) as n -> oo is this sequence.

Conjecture: a(n) - 1 is prime if and only if a(n) = n + 1. - _Mikhail Kurkov_, Mar 10 2022 Conjecture: a(n) = A057064(n+1) - 1 for n > 0. - _Mikhail Kurkov_, Mar 10 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 29 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001728 | OEIS-A057033

Let P(n) of a sequence s(1),s(2),s(3),... be obtained by leaving s(1),...s(n-1) fixed and reverse-cyclically permuting every n consecutive terms thereafter; apply P(2) to 1,2,3,... to get PS(2), then apply P(3) to PS(2) to get PS(3), then apply P(4) to PS(3), etc. The limit of PS(n) is A057033.

Conjecture: a(n) = A057063(n+1) - 1 for n > 0. - _Mikhail Kurkov_, Mar 08 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 29 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001729 | OEIS-A057059

Let R(i,j) be the rectangle with antidiagonals 1; 2,3; 4,5,6; ... Define i(m) and j(m) by R(i(m),j(m)) = m. Then a(n) = j(A057027(n)).

For the proof of the (slightly extended) conjecture stated in the formula section by _L. Edson Jeffery_ see the W. Lang link. - _Wolfdieter Lang_, Sep 14 2016 (Conjecture) Define the Chebyshev polynomials of the second kind by U_0(t) = 1, U_1(t) = 2*t, and U_r(t) = 2*t*U_(r-1)(t) - U_(r-2)(t) (r>1). Then T(n,k) = Sum_{j=1..n} U_(k-1)(cos((2*j-1)*Pi/(2*n+1))), 1<=k<=n. - _L. Edson Jeffery_, Jan 09 2012 (See the Sep 14 2016 comment above.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 30 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001730 | OEIS-A057063

Let P(n) of a sequence s(1),s(2),s(3),... be obtained by leaving s(1),...,s(n) fixed and reverse-cyclically permuting every n consecutive terms thereafter; apply P(2) to 1,2,3,... to get PS(2), then apply P(3) to PS(2) to get PS(3), then apply P(4) to PS(3), etc. The limit of PS(n) is A057063.

Conjecture: a(n) = A057033(n-1) + 1 for n > 1 with a(1) = 1. - _Mikhail Kurkov_, Mar 10 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 01 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001731 | OEIS-A057081

Even-indexed Chebyshev U-polynomials evaluated at sqrt(11)/2.

Conjecture: for k >= 1, Sum_{n >= 1} (-1)^(n+1)/(a(k*n) - s(k)/a(k*n)) = 1/(1 + a(k)) where s(k) = a(0) + a(1) + ... + a(k-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Aug 04 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001732 | OEIS-A057083

Scaled Chebyshev U-polynomials evaluated at sqrt(3)/2; expansion of 1/(1 - 3*x + 3*x^2).

Conjecture: Let M be any endomorphism on any vector space, such that M^3 = 1 (identity). Then (1-M)^n = A057681(n) - A057682(n)*M + z(n)*M^2, where z(0) = z(1) = 0 and, apparently, z(n+2) = a(n). - _Stanislav Sykora_, Jun 10 2012 By the conjecture: Start with x(0)=1, y(0)=0, z(0)=0 and set x(n+1) = x(n) - z(n), y(n+1) = y(n) - x(n), z(n+1) = z(n) - y(n). Then a(n) = z(n+2). This recurrence indeed ends up in a repetitive cycle of length 6 and multiplicative factor -27, confirming G. McGarvey's observation. - _Stanislav Sykora_, Jun 10 2012 a(n) = Sum_{k = 0..floor(n/3)} (-1)^k*binomial(n+2,3*k+2). Sykora's conjecture in the Comments section follows easily from this. - _Peter Bala_, Nov 21 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Aug 11 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001733 | OEIS-A057096

Saint-Exupéry numbers: ordered products of the three sides of Pythagorean triangles.

It is an open question whether any two distinct Pythagorean Triples can have the same product of their sides.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Aug 01 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001734 | OEIS-A057171

Numbers k such that (5^k + 1)/6 is a prime.

With the discovery of a(15), the best fit line slope G=0.55167 (see link to Generalized Repunit Conjecture). This sequence is converging nicely to the conjectured slope G=0.56145948 (A080130). - _Paul Bourdelais_, Feb 26 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 15 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001735 | OEIS-A057192

Least m such that 1 + prime(n)*2^m is a prime, or -1 if no such m exists.

Aguirre conjectured that, for every n > 1, a(n) is even if and only if prime(n) mod 3 = 1 (see the MathStackExchange link below). - _Lorenzo Sauras Altuzarra_, Feb 12 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jan 10 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001736 | OEIS-A057194

a(1) = 1; a(n+1) = (Product_{k=1..n} a(k)) * Sum_{k=1..n} a(k).

Conjecture: a(n) < A216151(n) for all n > 1. - _Jon Perry_, Sep 12 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Sep 15 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001737 | OEIS-A057199

The first nontrivial (k>n+2) palindromic prime in both bases n and n+2 or -1 if it does not exist.

Conjecture: a(n) <> -1 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Sep 16 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001738 | OEIS-A057300

Binary counter with odd/even bit positions swapped; base-4 counter with 1's replaced by 2's and vice versa.

It is an open question if every sequence of integers with bounded gaps must contain a double 3-term arithmetic progression. This problem is equivalent to the well known additive square problem in infinite words: Is it true that any infinite word with a finite set of integers as its alphabet contains two consecutive blocks of the same length and the same sum? For more details about the additive square problem in infinite words see the following references: Ardal, et al.; Brown and Freedman; Freedman; Grytczuk; Halbeisen and Hungerbuhler, and Pirillo and Varricchio. Conjecture: a(2*n) = -2*a(n) + 5*n, a(2*n+1) = -2*a(n) + 5*n + 2. - _Ralf Stephan_, Oct 11 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marc LeBrun_, Aug 24 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001739 | OEIS-A057475

Number of k, 1 <= k <= n, such that gcd(n,k) = gcd(n+1,k) = 1.

It is conjectured that every positive integer appears. - _Jon Perry_, Dec 12 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Sep 27 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001740 | OEIS-A057566

Number of collinear triples in a 3 X n rectangular grid.

Conjecture: a(n) = 5*floor((2n^3 - 3n^2 - 2n + 3)/24) + floor((2(n-1)^3 - 3(n-1)^2 - 2(n-1) + 3)/24) + n, which fits all of the listed terms. [Corrected by _Alper Ferudun_, Oct 21 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Oct 04 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001741 | OEIS-A057652

Numbers n such that n-2^k is a lucky number for all k such that 1 < 2^k < n.

Lucky numbers have asymptotic properties very similar to prime numbers, so one can conjecture finiteness of this sequence in the same way as Erdős did for A039669, and this should generalize to any sequence created using a similar sieve. - _M. F. Hasler_, Oct 15 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Oct 14 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001742 | OEIS-A057655

The circle problem: number of points (x,y) in square lattice with x^2 + y^2 <= n.

a(n) - Pi*n = O(sqrt(n)) (Gauss). a(n) - Pi*n = O(n^c), c = 23/73 + epsilon ~ 0.3151 (Huxley). If a(n) - Pi*n = O(n^c) then c > 1/4 (Landau, Hardy). It is conjectured that a(n) - Pi*n = O(n^(1/4 + epsilon)) for all epsilon > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 15 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001743 | OEIS-A057659

Prime numbers whose square is composed of just two different decimal digits.

Conjectured to be a finite sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carlos Rivera_, Oct 15 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001744 | OEIS-A057663

Primes p such that p + 2^p is also a prime.

These terms satisfy phi(k + 2^k) = phi(k) + 2^k, where phi is A000010, the Euler totient function. Conjecture: this sequence gives all numbers k that satisfy the condition phi(k + 2^k) = phi(k) + 2^k. - _Juri-Stepan Gerasimov_, May 23 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Oct 16 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001745 | OEIS-A057693

Number of permutations on n letters that have only cycles of length 3 or less.

Conjectural 3-adic values (each one checked up to n = 1000):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dennis P. Walsh_, Oct 20 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001746 | OEIS-A057702

Even numbers not the sum of a pair of twin lucky numbers.

Conjectured to be complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Oct 23 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001747 | OEIS-A057746

Even numbers not the sum of a pair of twin even-lucky-numbers (from A045955, A045956).

Conjectured to be complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Oct 30 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001748 | OEIS-A057748

Number of decompositions of 2n-1 into sum of a twin lucky number and a twin even-lucky-number (from A045955, A045956).

Conjecture: this sequence is always positive (with n>1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Oct 30 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001749 | OEIS-A057768

From Goldbach problem: number of decompositions of 2n-1 into sum of a prime lucky number(from A031157) and a twin even-lucky-number(from A045955, A045956).

Conjecture: this sequence is always positive (with n>2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Nov 01 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001750 | OEIS-A057773

a(n) = Sum_{i=1..n} nu_2(prime(i) - 1) where nu_2(m) = exponent of highest power of 2 dividing m.

Conjecture: a(n) ~ 2n. - _Charles R Greathouse IV_, Jun 02 2015 The conjecture is true. See the comment in A023506 about the asymptotic distribution of nu_2(prime(i) - 1). - _Amiram Eldar_, Mar 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Nov 02 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001751 | OEIS-A057778

a(n) is the least odd k such that k*2^n + 1 is prime.

Conjecture: for n > 0, a(n) = k < 2^n, so k*2^n + 1 is a Proth prime A080076. - _Thomas Ordowski_, Apr 13 2019 Conjecture: a(n) = O(n*log(n)). - _Thomas Ordowski_, Oct 16 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Nov 02 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001752 | OEIS-A057856

Least k such that (n+1)^k + n^k is a prime.

Conjecture: For every pair of relatively prime numbers (x, y) there exist at least one number n=2^m and one prime number p such that p = x^n + y^n. This sequence shows one case of this conjecture where y = x + 1. - _Tomas Xordan_, Jun 02 2007 The conjecture for arbitrary coprime x and y is false. Take x = 8 and y = 27. For every integer r >= 1, 8^r + 27^r = (2^r + 3^r)*(4^r - 6^r + 9^r), where the first factor is at least 5 and the second is (3^r - 2^r)^2 + 6^r >= 7. In particular, 8^(2^m) + 27^(2^m) is composite for every integer m >= 0. This does not refute the special case y = x + 1. - _Eiver Rodríguez_, Oct 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Nov 14 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001753 | OEIS-A057870

Number of singular points on n-th order Chmutov surface.

Conjectures from _Colin Barker_, Jan 02 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001754 | OEIS-A057896

Nonnegative numbers that can be written as m^k - m (with m and k nonnegative) in more than one way.

Conjectured to be finite and complete by Bennett (2001).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Sep 26 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001755 | OEIS-A057905

Positive integers that are not the sum of exactly four positive cubes.

It is conjectured that this sequence is finite, with 7373170279850 as its last member. - _Charles R Greathouse IV_, Jan 14 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001756 | OEIS-A057923

a(0)=0, a(1)=2, a(n) = smallest number such that sequence b(n) = {a(n-1) BITWISE OR a(n)} is strictly monotonically increasing.

Conjecture: a(n+2) > a(n). - _Robert Israel_, Aug 13 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | Larry Reeves (larryr(AT)acm.org), Oct 03 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001757 | OEIS-A057961

Number of points in square lattice covered by a disc centered at (0,0) as its radius increases.

Conjecture: the number of lattice points in a quadrant of the disk is equal to A000592(n-1). - _L. Edson Jeffery_, Feb 10 2014 The conjecture is true because a(n) is the cumulative sum of A105352(n), which is the number of distinct squared hypotenuses that are the radius of the disc whose center is (0,0) as it increases, once that a(A047808(n))=A000328(n) for n>=1; see A000161 and A000925. - _Flávio V. Fernandes_, Jan 09 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ken Takusagawa_, Oct 15 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001758 | OEIS-A058047

Generalized Collatz sequences: primes resulting in a cycle containing 1.

All terms are as yet only conjectures. _Jeff Heleen_ checked the primes < 1000 and start points up to 10000000 (see Prime Puzzle 114 and example below). a(1)=3 is the ordinary Collatz problem. - _Frank Ellermann_, Jan 20 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | Murad A. AlDamen (Divisibility(AT)yahoo.com), Nov 17 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001759 | OEIS-A058048

For each prime P consider the generalized Collatz sequence of each integer N > 1 defined by c(0) = N, c(m+1) = c(m) * P + 1 if F > P, else c(m+1) = c(m) / F, where F is the smallest factor of c(m), until the sequence cycles. If all c(i) > 1 for some starting number N then P belongs to the sequence (and vice versa).

Missing primes are as yet only conjectures. _Jeff Heleen_ checked the primes < 1000 and start points up to 10000000 (see Prime Puzzle 114 and example below). P=3 is the ordinary Collatz problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Murad A. AlDamen (Divisibility(AT)yahoo.com), Nov 17 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001760 | OEIS-A058128

a(1) = 1, a(n) = (n^n-n)/(n-1)^2 for n >= 2.

Conjecture: For even n, a(n) is prime or a weak Fermat pseudoprime to base n. - _Davide Rotondo_, Nov 06 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dennis P. Walsh_, Nov 14 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001761 | OEIS-A058188

Number of primes between prime(n) and prime(n) + sqrt(prime(n)), where prime(n) is the n-th prime.

Conjecture: if prime(n)>=127, there is always at least one prime between prime(n) and prime(n) + sqrt(prime(n)). Easily checked for prime(n)<1.1e15 in existing maximal gap tables

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Adam Kertesz_, Dec 04 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001762 | OEIS-A058211

Numbers k such that there is an arrangement of k points in the plane such that no point is on >= k/2 connecting lines.

Counterexamples to a conjecture of Dirac.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 30 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001763 | OEIS-A058213

Triangular arrangement of solutions of phi(x) = 2^n (n >= 0), where phi=A000010 is Euler's totient function. Each row corresponds to a particular n and its length is n+2 for 0 <= n <= 31, 32 for n >= 32. (This assumes that there are only 5 Fermat primes.)

phi(x) is a power of 2 if and only if x is a power of 2 multiplied by a product of distinct Fermat primes. So if, as is conjectured, there are only 5 Fermat primes, then there are only 32 possibilities for the odd part of x, namely the divisors of 2^32-1, given in A004729.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Nov 30 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001764 | OEIS-A058227

Number of edges in all simple (loopless) paths, connecting any node with all the remaining ones in optimal graphs of degree 4.

Conjectures from _Sean A. Irvine_, Jul 29 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | S. Bujnowski (slawb(AT)atr.bydgoszcz.pl), Feb 13 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001765 | OEIS-A058241

Number of sets of n positive integers that can be placed along a circle such that the set of sums of adjacent integers forms { 1, 2, ..., n^2-n+1 }.

Conjecture: for n > 2, p prime, e > 0, if n-1 is of the form p^e then a(n) > 0, otherwise a(n)=0. Conjecture: for n > 2, p prime, e > 0, if n-1 is of the form p^e then a(n) = A000010(n^2-n+1)/(6e), otherwise a(n)=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Jan 16 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001766 | OEIS-A058249

(Smallest prime >= 2^n) - (largest prime <= 2^n).

This sequence gives the gap between consecutive primes on either side of 2^n. The average gap between primes near 2^n should be about g=n*log(2). Cramer's conjecture would allow gaps to be as large as about g^2. - _T. D. Noe_, Jul 17 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Dec 05 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001767 | OEIS-A058265

Decimal expansion of the tribonacci constant t, the real root of x^3 - x^2 - x - 1.

Conjecture: equals (A(exp(Pi*sqrt(11))*A(exp(Pi/sqrt(11))))/sqrt(2), where A(x) = Product_{n >= 0} (1 + x^(2*n+1))/(1 - x^(2*n+1)) is the g.f. of A080054. - _Peter Bala_, May 30 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Dec 07 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001768 | OEIS-A058277

Number of values of k such that phi(k) = n, where n runs through the values (A002202) taken by phi.

Carmichael (1922) conjectured that the number 1 never appears in this sequence. Sierpiński conjectured and Ford (1998) proved that all integers greater than 1 occur in the sequence. Erdős (1958) proved that if s >= 1 appears in the sequence then it appears infinitely often. - _Nick Hobson_, Nov 04 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | Claude Lenormand (claude.lenormand(AT)free.fr), Jan 05 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001769 | OEIS-A058340

Primes p such that phi(x) = p-1 has only 2 solutions, namely x = p and x = 2p.

Conjecture: if q > 7 is in A005385, then q is in the sequence. - _Thomas Ordowski_, Jan 04 2017 Proof of conjecture: q'=(q-1)/2 is an odd prime > 3. If phi(x)=2q', which has 2-adic order 1 but is not a power of 2, there must be exactly one odd prime r dividing x. We could also have a factor of 2 (but no higher power, which would contribute more 2's to phi(x)). If x = r^e or 2r^e, then phi(x) = (r-1) r^(e-1). For this to be 2q' one possibility is r-1 = 2 and r^(e-1)=q', but then q'=r=3, ruled out by q > 7. The only other possibility is r-1=2q' and e=1, which makes r=q and x=q or 2q. - _Robert Israel_, Jan 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Dec 14 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001770 | OEIS-A058347

Array T(n,k), n,k nonnegative: the total number of checks required by a "double-support" algorithm to find out which rows and columns of each of the n by k zero-one matrices are nonzero.

Conjecture: T(n,k) = T(k,n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | M.R.C. van Dongen (dongen(AT)cs.ucc.ie), Dec 15 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001771 | OEIS-A058404

Coefficient triangle of polynomials (falling powers) related to Pell number convolutions. Companion triangle is A058405.

a(k,0)= A057084(k), k >= 0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Dec 11 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001772 | OEIS-A058764

Smallest number x such that cototient(x) = 2^n.

a(n) = (if n>1 then 3 else 4)*2^(n-1) = A007283(n-1) for n>1. (Conjectured.) - _M. F. Hasler_, Nov 10 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jan 02 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001773 | OEIS-A058799

Column 2 of A007754.

Conjectured partial sums of A093935. - _Sean A. Irvine_, Jun 25 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, Dec 02 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001774 | OEIS-A058814

Numbers k such that k divides the number of digits of k!.

For large m, 10^m*C -> 10^m*e, where e is Euler's or Napier's constant (A001113). Conjecture: There exist at least two contiguous terms for each k > 0, sometimes three contiguous terms, but never four. - _Robert G. Wilson v_, Feb 04 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jan 03 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001775 | OEIS-A058853

Primes p such that x^43 = 2 has a solution mod p.

It is conjectured that this sequence has density 42/43 ~ 0.976744 over all the primes. If the conjecture is correct, then a(n) ~ 43/42 * n log n. The generalized conjecture above is equivalent to: let P(p,1) be the set of primes congruent to 1 modulo p, P(p,1;a) be the set of primes q congruent to 1 modulo p such that x^p == a (mod q) has a solution, where p is a prime, a is not a p-th power, then the density of P(p,1;a) over P(p,1) is 1/p. - _Jianing Song_, Mar 09 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Dec 15 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001776 | OEIS-A058927

Numerators of series related to triangular cacti.

Conjecture: S(x) = Sum_{n>=0} ((2*n+1)^(n-1)/(n!*2^n))*x^(2*n+1) = (A052750(n)/A000165(n))*x^(2*n+1). Letting D_n be the set of divisors of n! and d_n = max(k in D_n : k | (2*n+1)^(n-1)), then a(n)=A052750(n)/d_n. (End) The above conjecture is correct and follows from formula given in A034940 for the number of rooted labeled triangular cacti with 2n+1 nodes. - _Andrew Howroyd_, Aug 30 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 12 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001777 | OEIS-A058928

Denominators of series related to triangular cacti.

Conjecture: S(x) = Sum_{n>=0} ((2*n+1)^(n-1)/(n!*2^n))*x^(2*n+1) = (A052750(n)/A000165(n))*x^(2*n+1). Letting D_n be the set of divisors of n! and d_n = max(k in D_n : k | (2*n+1)^(n-1)), then a(n)=A000165(n)/d_n. (End) The above conjecture is correct and follows from formula given in A034940 for the number of rooted labeled triangular cacti with 2n+1 nodes. - _Andrew Howroyd_, Aug 30 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 12 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001778 | OEIS-A058953

Numbers k of the form k=p*2^x, with p prime and x>=0, such that tau(k)-m = A058933(k) where tau(k) is the number of divisors of k and m is 0 or 1.

(p=2 => m=1); (Conjecture: [for p>2]; if p=A040069 then m=0 else m=1.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Jan 13 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001779 | OEIS-A059099

Expansion of e.g.f. exp(exp(x)-1)/(2-exp(x)).

Conjecture: a(n) = exp(1) * Sum_{k>=0} k^n * Gamma(k + 1, 2)/(2^(k + 1) * k!) for n > 0. - _Velin Yanev_, Aug 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Jan 02 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001780 | OEIS-A059100

a(n) = n^2 + 2.

Let s(n) = Sum_{k>=1} 1/n^(2^k). Then I conjecture that the maximum element in the continued fraction for s(n) is n^2 + 2. - _Benoit Cloitre_, Aug 15 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Feb 13 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001781 | OEIS-A059133

A hierarchical sequence (S(W2{2}c) - see A059126).

Conjectures from _Colin Barker_, Oct 07 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonas Wallgren_, Jan 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001782 | OEIS-A059135

A hierarchical sequence (S(W2{3}c) - see A059126).

Conjectures from _Colin Barker_, Oct 07 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonas Wallgren_, Jan 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001783 | OEIS-A059138

A hierarchical sequence (S(W3{2,2}cc) - see A059126).

Conjectures from _Colin Barker_, Oct 07 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonas Wallgren_, Jan 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001784 | OEIS-A059142

A hierarchical sequence (S(W2{3}*) - see A059126).

Conjectures from _Colin Barker_, Oct 06 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonas Wallgren_, Jan 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001785 | OEIS-A059158

A hierarchical sequence (S(W'3{2,2}cc) - see A059126).

a(n) = (24/7) * (8*16^n - 2^n) (conjectured). - _Ralf Stephan_, May 05 2004 Conjectures from _Colin Barker_, Oct 07 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonas Wallgren_, Feb 02 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001786 | OEIS-A059163

A hierarchical sequence (W'3{2,2}*c - see A059126).

Conjecture: a(n) = 2^(1+A059129(n)). - _R. J. Mathar_, Apr 04 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonas Wallgren_, Feb 02 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001787 | OEIS-A059259

Triangle read by rows giving coefficient T(i,j) of x^i y^j in 1/(1-x-x*y-y^2) = 1/((1+y)(1-x-y)) for (i,j) = (0,0), (1,0), (0,1), (2,0), (1,1), (0,2), ...

Conjecture: every second column of this triangle is identical to a column in the square array A071921. For example, column 4 of A059259 (1, 3, 9, 22, 46, ...) appears to be the same as column 3 of A071921; column 6 of A059259 (1, 4, 16, 50, 130, 296, ...) appears to be the same as column 4 of A071921; and in general column 2k of A059259 appears to be the same as column k+1 of A071921. Furthermore, since A225010 is a transposition of A071921 (ignoring the latter's top row and two leftmost columns), there appears to be a correspondence between column 2k of A059259 and row k of A225010. - _Mathew Englander_, May 17 2014 See the Edwards-Allen 2020 paper, page 14, for proof of Englander's conjecture. - _Michael De Vlieger_, Dec 10 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 23 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001788 | OEIS-A059269

Numbers m for which the number of divisors, tau(m), is divisible by 3.

Conjecture: a(n) ~ k*n where k = 1/(1 - Product(1 - (p-1)/(p^(3*i)))) = 3.743455... where p ranges over the primes and i ranges over the positive integers. - _Charles R Greathouse IV_, Apr 13 2012 The asymptotic density of this sequence is 1 - zeta(3)/zeta(2) = 1 - 6*zeta(3)/Pi^2 = 0.2692370305... (Sathe, 1945). Therefore, the above conjecture, a(n) ~ k*n, is true, but k = 1/(1-6*zeta(3)/Pi^2) = 3.7141993349... - _Amiram Eldar_, Jul 26 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | Avi Peretz (njk(AT)netvision.net.il), Jan 24 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001789 | OEIS-A059279

G.f. is ((1-x)/(1-2*x)) * G(x*(1-x)/(1-2*x)) where G(x) is g.f. for Catalan numbers A000108.

Conjecture: (n+1)*a(n) +2*(1-4*n)*a(n-1) + 4*(4*n-5)*a(n-2) +4*(5-2*n)*a(n-3)=0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 24 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001790 | OEIS-A059297

Triangle of idempotent numbers binomial(n,k)*k^(n-k), version 1.

Conjecture: the matrix inverse is A137452. - _R. J. Mathar_, Mar 12 2013 The above conjecture is correct. This triangle is the exponential Riordan array [1, x*exp(x)]. Thus the inverse array is the exponential Riordan array [1, W(x)], which equals A137452. - _Peter Bala_, Apr 08 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 25 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001791 | OEIS-A059304

a(n) = 2^n * (2*n)! / (n!)^2.

Theorem: For n >= 0, a(n) = Sum_{0 <= j, k <= n} binomial(n, j)*binomial(n, k)*binomial(j+k, n). [Conjectured by _Peter Bala_, Jul 16 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Jan 25 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001792 | OEIS-A059324

Numbers n such that 6n + 5 is composite.

Conjecture: There exists no pair of primes (p, q > p^2) such that q - p^2 = 6*n - 4 (see A138479). - Philippe LALLOUET (philip.lallouet(AT)orange.fr), Mar 20 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anton Joha_, Jan 26 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001793 | OEIS-A059363

Lexicographically earliest increasing sequence of positive integers such that A001222(a(i)+a(j)) = A001222(a(i)) + A001222(a(j)) for all 1<=i<j.

The values of A001222(a(n)) are 0,1,2,2,2,... and are conjectured to be 2 for all n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Jan 28 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001794 | OEIS-A059456

Unsafe primes: primes not in A005385.

Conjecture: From the sequence of prime numbers, let 2 and remove the first data iteration of 2*p+1; leave 3 and remove the prime data by the iteration 2*p+1 and we get the sequence. Example for p=2, remove(5,11,23,47); p=3, remove(7); p=13, p=17, p=19, p=23, remove(47); and so on. - _Vincenzo Librandi_, Aug 07 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Feb 02 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001795 | OEIS-A059571

From Mertens's conjecture (1): floor(sqrt(n)) - |M(n)|, where M is Mertens's function A002321.

From Mertens's conjecture (1): floor(sqrt(n)) - |M(n)|, where M is Mertens's function A002321. Mertens conjectured that |A002321(n)| < sqrt(n) for all n > 1. This is now known to be false. So eventually there will be negative terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 16 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001796 | OEIS-A059572

From Mertens's conjecture (2): floor(sqrt(n)) - Mertens's function A002321(n).

From Mertens's conjecture (2): floor(sqrt(n)) - Mertens's function A002321(n). Mertens conjectured that |A002321(n)| < sqrt(n) for all n > 1. This is now known to be false.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 16 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001797 | OEIS-A059581

From Von Sterneck's conjecture: floor(sqrt(n)) - 2*|Mertens's function A002321(n)|.

From Von Sterneck's conjecture: floor(sqrt(n)) - 2*|Mertens's function A002321(n)|. Von Sterneck conjectured that 2*|A002321(n)| < sqrt(n) for all sufficiently large n. This is now known to be false. This is different from the Mertens conjecture that |A002321(n)| < sqrt(n) for all n > 1 (which is also false).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 16 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001798 | OEIS-A059686

Grimm numbers (1): a(n) = largest k so that for each m in {n+1, n+2, ..., n+k} there corresponds a different prime factor p_m.

Guy (2004) discusses some conjectures of Grimm that could produce related sequences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 06 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001799 | OEIS-A059751

Grimm numbers (2): a(n) = largest k so that for each composite m in {n+1, n+2, ..., n+k} there corresponds a different divisor d_m with 1 < d_m < m.

Erdős and Pomerance conjectured that the number n+a(n)+1, which "blocks" a(n) from becoming larger, is always an odd semiprime. They verified this conjecture up to n=492 and proved it for large n. The numbers n at which n+a(n)+1 increases also appear to be semiprimes. - _T. D. Noe_, Feb 18 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 11 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001800 | OEIS-A059774

Consider the line segment in R^n from the origin to the point P=(1,2,3,...,n); let d = squared distance to this line from the closest point of Z^n (excluding the endpoints). Sequence gives d times P.P.

Officially these are just conjectures so far.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Vinay Vaishampayan_, Feb 21 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001801 | OEIS-A059777

Number of self-conjugate three-quadrant Ferrers graphs that partition n.

G.f.: conjecture: 1/(Q(0) - 1), where Q(k) = 1 + (-x)^k - (-x)^(k+2)/Q(k+1); (continued fraction). - _Sergei N. Gladkovskii_, Nov 25 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 21 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001802 | OEIS-A059786

Smallest prime after 2*(n-th prime).

Conjecture: If k is an even integer greater than 2, then a(n) - A059788(n) = k for infinitely many n. - _Clark Kimberling_, Dec 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Feb 22 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001803 | OEIS-A059799

Primes which when added to their reversals are squares.

Idea from Carlos Rivera's The Prime Puzzles and Problems Connection, Conjecture 23.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enoch Haga_, Feb 23 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001804 | OEIS-A059804

Consider the line segment in R^n from the origin to the point v=(2,3,5,7,11,...) with prime coordinates; let d = squared distance to this line from the closest point of Z^n (excluding the endpoints). Sequence gives d times v.v.

Officially these are just conjectures so far.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Vinay Vaishampayan_, Feb 21 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001805 | OEIS-A059837

Diagonal T(s,s) of triangle A059836.

Conjecture: (n+1)*a(n) - 2*n*a(n-1) - 4*n*(n-1)^2*a(n-2) = 0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 25 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001806 | OEIS-A059850

Decimal expansion of Pi^e.

Pi^e is conjectured to be transcendental.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Hsu, Po-Wei (Benny) (arsene_lupin(AT)intekom.co.za), Jan 13 2000 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001807 | OEIS-A059861

a(n) = Product_{i=2..n} (prime(i) - 2).

Arises in Hardy-Littlewood k-tuple conjecture. Also a(n) is the exact number of d=2 and also d=4 differences in dRRS[modulus=n-th primorial]; see A049296 (dRRS[m]=set of first differences of reduced residue system modulo m).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Feb 28 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001808 | OEIS-A059864

a(n) = Product_{i=4..n} (prime(i) - 5).

Such products arise in Hardy-Littlewood prime k-tuplet conjectural formulas.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Feb 28 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001809 | OEIS-A059865

a(n) = Product_{i=4..n} (prime(i) - 6).

Arises in Hardy-Littlewood prime k-tuplet conjectural formulas. Also the sequence gives the exact numbers of X42424Y difference-pattern in dRRS[m], where m=modulus=A002110(n). See A049296 (=dRRS[210]=list of first differences of reduced residue system modulo 210=4th primorial). A pattern X42424Y corresponds to a residue-sextuple or it is their difference-quintuple, X,Y > 4. Analogous pattern for primes is in A022008.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Feb 28 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001810 | OEIS-A059871

Number of solutions to the equation p_i = (1+mod(i,2))*p_{i-1} +- p_{i-2} +- p_{i-3} +- ... +- 2 +- 1, where p_i is the i-th prime number (where p_1 = 2 and the "zeroth prime" p_0 is defined to be 1).

This is Scherk's theorem, which was conjectured by Scherk in 1833 and proved by Pillai in 1928. [_T. D. Noe_, Oct 03 2008]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 05 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001811 | OEIS-A059894

Complement and reverse the order of all but the most significant bit in binary expansion of n. n = 1ab..yz -> 1ZY..BA = a(n), where A = 1-a, B = 1-b, ...

a(1) = 1, a(2n) = a(n) + 2^(floor(log_2(n))+1), a(2n+1) = a(n) + 2^floor(log_2(n)) (conjectured). - _Ralf Stephan_, Aug 21 2003 A000120(a(n)) = A000120(A054429(n)) = A023416(n) + 1 (conjectured). - _Ralf Stephan_, Oct 05 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marc LeBrun_, Feb 06 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001812 | OEIS-A059906

Index of second half of decomposition of integers into pairs based on A000695.

Conjecture: a(n) = n - (1/2)*Sum_{k=1..n} (sqrt(2)^A007814(k) + (-sqrt(2))^A007814(k)) = -Sum_{k=1..n} (-1)^k * 2^floor(k/2) * floor(n/2^k). - _Velin Yanev_, Dec 01 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marc LeBrun_, Feb 07 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001813 | OEIS-A059917

a(n) = (3^(2^n) + 1)/2 = A059919(n)/2, n >= 0.

a(0), a(1), a(2), a(4), a(5), and a(6) are prime. Conjecture: a(n) is composite for all n > 6. - _Thomas Ordowski_, Dec 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Feb 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001814 | OEIS-A059928

The periodic point counting sequence for the toral automorphism given by the polynomial of (conjectural) smallest Mahler measure. The map is x -> Ax mod 1 for x in [0,1)^10, where A is the companion matrix for the polynomial x^10+x^9-x^7-x^6-x^5-x^4-x^3+x+1.

The periodic point counting sequence for the toral automorphism given by the polynomial of (conjectural) smallest Mahler measure. The map is x -> Ax mod 1 for x in [0,1)^10, where A is the companion matrix for the polynomial x^10+x^9-x^7-x^6-x^5-x^4-x^3+x+1. The terms in this sequence are all squares. The sequence of square roots, A087612, is conjectured to contain an infinite number of primes. - _T. D. Noe_, Sep 15 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | Graham Everest (g.everest(AT)uea.ac.uk), Mar 01 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001815 | OEIS-A059929

a(n) = Fibonacci(n)*Fibonacci(n+2).

The preceding conjecture is true. Proof: For n >= 1 let c(n) := confrac(repeat(1^{n-1}, 2)) where 1^{k} denotes 1 taken k times. This can be computed, e.g. from [Perron, third and fourth eq. on p. 62], as c(n) = (F(n) + sqrt(F(n+1)^2 - (-1)^n)) / F(n+1), which is the conjectured formula because F(n+1)^2 - (-1)^n = a(n). - _Wolfdieter Lang_, Jan 05 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Feb 09 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001816 | OEIS-A059958

Smallest number m such that m*(m+1) has at least n distinct prime factors.

Conjecture: a(n) = A069354(n) - 1. - _Robert G. Wilson v_, Feb 18 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Mar 02 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001817 | OEIS-A059965

For m>0, each n+m > 5 is expressed as Sum_{k = x,y,z}(pk)_m where (pk)_m is a prime with x <= y <= z; a(n) = largest m such that (pk)_1 = (pk)_2 = ... = (pk)_m.

Goldbach conjectured that every integer >5 is the sum of three primes. 6=2+2+2, 7=2+2+3, 8=2+3+3, 9=3+3+3=2+2+5,...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Mar 05 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001818 | OEIS-A059970

Nim-factorials: a(1)=1 and, for n>1, a(n)=n*a(n-1), where * denotes Nim multiplication.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Mar 05 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001819 | OEIS-A059992

Numbers with an increasing number of nonprime divisors.

Conjecture: This sequence includes all highly composite numbers (from A002182) except 2 and 6, but there are other terms in this sequence (e.g., a(3) = 8, a(9) = 72) that are not highly composite. Conjecture: a(n)/A007947(a(n)) is in A301414. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Mar 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001820 | OEIS-A059998

Number of different primes occurring when n is expressed as p1+q1+r1 = ... = pk+qk+rk where pk,qk,rk are primes with pk <= qk <= rk.

Goldbach conjectured that every integer >5 is the sum of three primes. 6=2+2+2, 7=2+2+3, 8=2+3+3, 9=3+3+3=2+2+5,.....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Mar 10 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001821 | OEIS-A060003

Odd numbers not of the form p + 2*k^2, k>0, p prime.

Goldbach conjectured that all odd composites are sum of a prime and twice a square. a(9) = 5777 and a(10) = 5993 are the only known exceptions. Elements a(2) .. a(8) are the odd Stern primes (cf. A042978). The next element of the sequence, if it exists, is larger than 10^9. - _M. F. Hasler_, Nov 16 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Mar 14 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001822 | OEIS-A060014

Sum of orders of all permutations of n letters.

Conjecture: This sequence eventually becomes cyclic mod m for all m. - _Isaac Saffold_, Dec 01 2019 log(a(n)/n!) = O(sqrt(n/log n)) [conjectured in Erdős & Turán, 1968].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 17 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001823 | OEIS-A060035

Least m >= 0 such that 2^m has n 2's in its base-3 expansion.

"Paul Erdős conjectured that for n > 8, 2^n is not a sum of distinct powers of 3. In terms of digits, this states that powers of 2 for n > 8 must always contain a '2' in their base 3 expansion." The value of a(42) is conjectured to be -1 because no power of 2 up to 2^10^7 has exactly 42 2's.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Mar 17 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001824 | OEIS-A060064

First subsequent, disjoint occurrence of n consecutive nonprimes.

Conjecture: a(n) - 1 is always prime. - _Robert Israel_, May 04 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Mar 25 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001825 | OEIS-A060096

Numerator of coefficients of Euler polynomials (rising powers).

The first (m=0) column of the rational triangle is conjectured to be E(n,0) = ((-1)^n)*A198631(n) / A006519(n+1). See also the first column shown in A209308 (different signs). - _Wolfdieter Lang_, Jun 15 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Mar 29 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001826 | OEIS-A060212

Primes q such that 6*q-1 and 6*q+1 are twin primes. Proper subset of A002822.

Conjecture: a(n) ~ n*log(n)*log(n*log(n))*log(log(n)). - _Carl R. White_, Nov 16 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Mar 20 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001827 | OEIS-A060236

If n mod 3 = 0 then a(n) = a(n/3), otherwise a(n) = n mod 3.

Equals A005148(n) reduced mod 3. In "On a sequence Arising in Series for Pi" Morris Newman and Daniel Shanks conjectured that 3 never divides A005148(n) and D. Zagier proved it. - _Benoit Cloitre_, Jun 22 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Mar 21 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001828 | OEIS-A060254

Primes which are the sum of two consecutive composite numbers.

Conjecture: a(n) ~ n log n. - _Charles R Greathouse IV_, Apr 29 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Mar 22 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001829 | OEIS-A060264

First prime after 2n.

Conjecture: for n > 2, this is the least prime p such that 1^2, 2^2, 3^2, ..., n^2 are distinct mod p. Checked to 10^4. - _Charles R Greathouse IV_, Dec 03 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Mar 23 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001830 | OEIS-A060266

Difference between 2n and the following prime.

Conjecture: Limit_{n->oo} (Sum_{k=1..n} a(k)) / (Sum_{k=1..n} log(2*k)) = 1. - _Alain Rocchelli_, Oct 24 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Mar 23 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001831 | OEIS-A060294

Decimal expansion of Buffon's constant 2/Pi.

Equals lim_{n->infinity} (1/n) * Sum_{k=1..n} abs(sin(k * m)) for all nonzero integers m (conjectured). Works with cos also. - _Dimitri Papadopoulos_, Jul 17 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Mar 28 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001832 | OEIS-A060295

Decimal expansion of exp(Pi*sqrt(163)).

It is sometimes called "Ramanujan's constant" due to an April Fool's joke by Gardner (1975) in which he claimed that Ramanujan conjectured that this constant is an integer, and that a fictitious "John Brillo" of the University of Arizona proved it on May 1974.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Mar 24 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001833 | OEIS-A060298

Number of powers x^y (x,y > 1) with n digits.

Conjectures from _Robert G. Wilson v_, Aug 29 2012: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel ten Voorde_, Apr 10 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001834 | OEIS-A060305

Pisano periods for primes: period of Fibonacci numbers mod prime(n).

Assuming Wall's conjecture (which is still open) allows one to calculate A001175(m) when m is a prime power since for any k >= 1: A001175(prime(n)^k) = a(n)*prime(n)^(k-1). For example: A001175(2^k) = 3*2^(k-1) = A007283(k-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Louis Mello (mellols(AT)aol.com), Mar 26 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001835 | OEIS-A060318

Exponent of highest power of 3 dividing the n-th odd Catalan number Catalan(2^n - 1).

Conjecture: all odd Catalan numbers have smallest prime factor 3, except Catalan(3), which has smallest prime factor 5, and Catalan(31) and Catalan(255), which have smallest prime factor 7 (checked up to Catalan(-1 + 2^2048)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_, Mar 28 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001836 | OEIS-A060324

a(n) is the minimal prime q such that n*(q+1)-1 is prime, that is, the smallest prime q so that n = (p+1)/(q+1) with p prime; or a(n) = -1 if no such q exists.

A conjecture of Schinzel, if true, would imply that such a q always exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthew Conroy_, Mar 29 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001837 | OEIS-A060330

Primes that are the sum of five consecutive composite numbers.

Conjecture: all these primes are isolated primes (A007510). - _Davide Rotondo_, Dec 31 2024 Stronger conjecture: all p are 7 or 23 mod 30. - _Charles R Greathouse IV_, Jan 21 2025 Above conjectures are true. Proof sketch: n + n+1 + n+2 + n+3 + n+4 = 5n+10, so there must be at least one prime sandwiched between the five composite numbers. If p and p+4 are prime, then p-1 + p+1 + p+2 + p+3 + p+5 = 5p + 10 is composite. If neither p-2 nor p+2 are prime, the sums p-4 + p-3 + p-2 + p-1 + p+1, etc., are 5p-9, 5p-3, 5p+3, and 5p+9 which are even for p > 2 (and p = 2 does not work). So we must have p and p+2 prime, which yield p-3 + p-2 + p-1 + p+1 + p+3 = 5p-2, p-2 + p-1 + p+1 + p+3 + p+4 = 5p+5, and p-1 + p+1 + p+3 + p+4 + p+5 = 5p+12. 5p+5 is composite, but the others can work. Now note that the first form yields only 5p-2 = 23 mod 30 and the second 5p+12 = 7 mod 30. - _Charles R Greathouse IV_, Jan 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Mar 30 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001838 | OEIS-A060382

In base n, a(n) is the smallest number m that leads to a palindrome-free sequence, using the following process: start with m; reverse the digits and add it to m, repeat. Stop if you reach a palindrome.

Only a(2) is proved, all the others are conjectured. - _Eric Chen_, Apr 20 2015 [corrected by _A.H.M. Smeets_, May 27 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel ten Voorde_, Apr 03 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001839 | OEIS-A060393

Values of k associated with A060380.

Additional terms require finding the minimal values for the primepuzzles.net conjecture for primes >= 269.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 04 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001840 | OEIS-A060432

Partial sums of A002024.

Conjecture: a(n) is the minimum diameter of an (n+1)-element set of integers for which no nonzero d has more than d representations as a difference of elements of the set. Known to be true for n <= 16. - _Aditya A Gupta_ and _Sean A. Irvine_, Apr 26 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | Robert A. Stump (bobess(AT)netzero.net), Apr 06 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001841 | OEIS-A060454

Consider the line segment in R^n from the origin to the point v = (1,4,9,...,n^2); let d = squared distance to this line from the closest point of Z^n (excluding the endpoints). Sequence gives d times v.v.

Officially these are just conjectures so far.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 09 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001842 | OEIS-A060464

Numbers that are not congruent to 4 or 5 mod 9.

Conjecture: n is a sum of three cubes iff n is in this sequence. Heath-Brown conjectures that n is a sum of three cubes in infinitely many ways iff n is in this sequence (and not at all otherwise). See his paper for a conjectural asymptotic. - _Charles R Greathouse IV_, Mar 12 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 10 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001843 | OEIS-A060516

Number of series-reduced (or homeomorphically irreducible) graphs with loops on n labeled nodes.

Conjecture: the number of connected graphs of this type is 1, 1, 4, 13, 203, 7754, ... (n>=0). - _R. J. Mathar_, Apr 29 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Mar 24 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001844 | OEIS-A060521

Number of 3 X n grids of black and white cells, no 3 of same color vertically or horizontally contiguous.

The conjectured recursion is correct: For each n count the solutions separately where the last two rows differ in 0, 1, 2, or 3 places; a linear recursion is then readily found. The corresponding matrix has characteristic polynomial x^4 - 2 x^3 - 5 x^2 + 1, matching the recursion recursion a(n+4) = 2a(n+3) + 5a(n+2) - a(n). [From _Hagen von Eitzen_, Oct 21 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | Tom Womack (tom(AT)womack.net) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001845 | OEIS-A060602

Number of tilings of the d-dimensional zonotope constructed from d+3 vectors.

Conjectures from _Colin Barker_, Feb 20 2013: (Start) The above conjectures are correct; see Proposition 7.1 in Ziegler's article. - _Manfred Scheucher_, Feb 09 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | Matthieu Latapy (latapy(AT)liafa.jussieu.fr), Apr 12 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001846 | OEIS-A060651

Smallest odd prime p such that Q(sqrt(-p)) has class number 2n+1.

Conjecture: a(n) = A002146(n) for all n >= 1. That is to say, A002148(n) > A002146(n) for all n >= 1. - _Jianing Song_, Jul 20 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Apr 17 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001847 | OEIS-A060693

Triangle (0 <= k <= n) read by rows: T(n, k) is the number of Schröder paths from (0,0) to (2n,0) having k peaks.

Conjecture: Sum_{k=0..n} (-1)^k*T(n,k)*(n+1-k)^2 = 1+n+n^2. - _Werner Schulte_, Jan 11 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _F. Chapoton_, Apr 20 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001848 | OEIS-A060699

a(n) = floor(A^(C^n)), where A = 2.084551112207285611..., C = 1.221.

The extension of the sequence is guaranteed by the Cramer conjecture. That is: If the needed change in Y(n) for obtaining the next prime (superior or inferior) is as maximum = (log Y(n))^2/2, then the effect on Y(n-1) is less than K*C^(2n-1)*Y(n-1)/Y(n). K = (1/2)*(log A)^2 = 0.269784 This value diminishes with n. Example: For n = 23, a change in Y(23) by 2630 only changes Y(22) by 0.0043. _Jens Kruse Anderson_ with A = 2.084551112197624209091521123 calculated Y(n) = floor(A^(C^n)) from n = 1 to n = 3, obtaining 22 different primes. - Luis Rodriguez-Torres (ludovicusmagister(AT)yahoo.com), Feb 10 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | Luis Rodriguez-Torres (ludovicusmagister(AT)yahoo.com), Apr 20 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001849 | OEIS-A060715

Number of primes between n and 2n exclusive.

For prime n conjecturally a(n) = A226859(n). - _Vladimir Shevelev_, Jun 27 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Apr 25 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001850 | OEIS-A060774

a(n) = number of lattice paths from (0,0,0) to (n,n,n) along the cracks on the surface of a Rubik-ized n X n X n cube so that no step increases distance from goal.

Conjecture: 2*n*(2*n-1)*(n-1)*a(n) + (n-1)*(13*n^2-209*n+258)*a(n-1) + 2*(-259*n^3+1785*n^2-3728*n+2460)*a(n-2) + 6*(295*n^3-2068*n^2+4833*n-3780)*a(n-3) - 36*(3*n-10)*(2*n-7)*(3*n-11)*a(n-4) = 0. - _R. J. Mathar_, Oct 31 2015 Conjecture: 2*n*(n-1)*(2*n-1)*(11*n^2-33*n+24)*a(n) - (n-1)*(473*n^4-1892*n^3+2561*n^2-1338*n+216)*a(n-1) + 6*(3*n-5)*(3*n-4)*(2*n-3)*(11*n^2-11*n+2)*a(n-2) = 0. - _R. J. Mathar_, Oct 31 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Len Smiley_, Apr 25 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001851 | OEIS-A060782

Degree of the curve C(n) relative to a triangle ABC with side lengths a, b and c, given by (x^n-b^n)(y^n-c^n)(z^n-a^n) = (x^n-c^n)(y^n-a^n)(z^n-b^n) where x, y and z denote the distances from the variable point to vertices A, B and C respectively.

It is known that a(n) = 2n-1 for n even; a(1) = 14; a(3) = 46; and it is conjectured that a(n) = 16n-2 for n odd.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Antreas P. Hatzipolakis (xpolakis(AT)otenet.gr), Apr 28 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001852 | OEIS-A060897

Number of walks of length n on square lattice, starting at origin, staying in first and third quadrants.

Is there a formula analogous to the (conjectured) formula for A060900?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_, May 05 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001853 | OEIS-A060898

Number of walks of length n on square lattice, starting at origin, staying in first, second and third quadrants.

Is there a formula analogous to the (conjectured) formula for A060900?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_, May 05 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001854 | OEIS-A060900

Number of walks of length n on square lattice, starting at origin, staying on points with x >= 0, y <= x.

The following conjectural formula for this sequence is apparently due to _Ira M. Gessel_: a(0) = 1, a(2n) = a(2n-1)*(12n+2)/(3n+1), a(2n+1) = a(2n)*(4n+2)/(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_, May 05 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001855 | OEIS-A060903

a(n) = floor(6*n*sqrt(n)/Pi^2).

Conjecture: the sum of the divisors of n is less than a(n) for n exceeding 12. - _Robert G. Wilson v_, May 14 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, May 05 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001856 | OEIS-A060941

Duchon's numbers: the number of paths of length 5*n from the origin to the line y = 2*x/3 with unit East and North steps that stay below the line or touch it.

The sequence defined by b(n) := [x^n] A(x)^n begins [1, 2, 50, 1415, 42258, 1300727, 40820837, 1298493730, ...] and conjecturally satisfies the congruence b(p) == b(1) (mod p^3) for prime p >= 7 (checked up to p = 101). [Added 23 Oct 2024: More generally, let r be an integer and s a positive integer and define a sequence u(n) by u(n) = [x^(s*n)] A(x)^(r*n). Then we conjecture that the supercongruences u(n*p^k) == u(n*p^(k-1)) (mod p^(3*k)) hold for all primes p >= 7 and positive integers n and k.] - _Peter Bala_, Sep 12 2021 Inductively define a family of sequences {a(i,n) : n >= 0}, i >= 1, by setting a(1,n) = a(n) and, for i >= 2, a(i,n) = [x^n] (exp(Sum_{k >= 1} a(i-1,k)*x^k/k))^n. We conjecture that the sequences {a(i,n) : n >= 0}, i >= 2, also satisfy the supercongruences u(n*p^k) == u(n*p^(k-1)) (mod p^(3*k)) for primes p >= 7 and positive integers n and k. - _Peter Bala_, Oct 24 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Flajolet_, May 12 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001857 | OEIS-A060973

a(2*n+1) = a(n+1)+a(n), a(2*n) = 2*a(n), with a(1)=0 and a(2)=1.

Conjectures from _Peter Bala_, Aug 03 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, May 09 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001858 | OEIS-A061002

As p runs through the primes >= 5, sequence gives { numerator of Sum_{k=1..p-1} 1/k } / p^2.

Conjecture: If p is the n-th prime and H(n) is the n-th harmonic number, then denominator(H(p)/H(p-1))/numerator(H(p-1)/p^2) = p^3. A193758(p)/a(n) = p^3, p > 3. - _Gary Detlefs_, Feb 20 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 15 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001859 | OEIS-A061060

Write product of first n primes as x*y with x<y and x maximal; sequence gives value of y-x.

Conjecture: Limit_{N->oo} (Sum_{n=1..N} log(a(n))) / (Sum_{n=1..N} prime(n)) = 1/e (A068985). - _Alain Rocchelli_, Nov 13 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ed Pegg Jr_, May 28 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001860 | OEIS-A061062

Sum of squared factorials: (0!)^2 + (1!)^2 + (2!)^2 + (3!)^2 +...+ (n!)^2.

There is a Kurepa-like conjecture (see A049782) for this sequence: for primes p>3, p does not divide a(p-1). However, the conjecture fails for p=20879. - _T. D. Noe_, Dec 08 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, May 27 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001861 | OEIS-A061068

Primes which are the sum of a prime and its subscript.

This sequence is the intersection of A014688 with the set of primes. Conjecture: this sequence is infinite, yet derives from arbitrarily long "prime deserts" such as the 11 composites in A014688 between a(6) = 19 and a(18) = 79 and the 17 composites in A014688 between a(48) = 271 and a(66) = 383. - _Jonathan Vos Post_, Nov 22 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, May 28 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001862 | OEIS-A061078

Sum of the products of the digits of the first n positive even numbers.

Conjecture: let a >= 1, b >= 0, where p is not a multiple of 2 nor 5. Then:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 14 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001863 | OEIS-A061100

Squares with digital root 4.

Conjecture: Conjecture is true since x^2 == 4 (mod 9) if and only if x == 2 or 7 (mod 9). The odd-numbered terms are (2+9*k)^2 and the even-numbered terms are (7+9*k)^2. - _Robert Israel_, Jun 13 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001864 | OEIS-A061101

Squares with digital root 7.

Conjecture: a(n)=(9*n-8)^2/4 for n even. a(n)=(9*n-1)^2/4 for n odd. G.f.: x*(16+9*x+112*x^2+9*x^3+16*x^4)/((1-x)^3*(1+x)^2). - _Colin Barker_, Apr 21 2012 Conjecture is true, because x^2 == 7 (mod 9) if and only if x == 4 or 5 (mod 9). - _Robert Israel_, Jan 31 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001865 | OEIS-A061220

Least entry in character table of the symmetric group S_n.

Conjecture: For n >= 2, the least entry in the character table of S_n occurs in the column indexed by the transpositions, of cycle type (2,1^(n-2)). Computationally verified for 2 <= n <= 70. - _Vladeta Jovovic_, Oct 21 2026 a(n) = min_{lambda |- n} chi^lambda(2,1^(n-2)) for n >= 2 (conjectured; computationally verified for 2 <= n <= 70), where chi^lambda is the irreducible character of S_n indexed by

lambda. - _Vladeta Jovovic_, Oct 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ola Veshta (olaveshta(AT)my-deja.com), May 30 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001866 | OEIS-A061232

Number of primes p with n! < p <= (n+1)!.

I conjecture that for n>2 we have n + 1/2 <= a(n)/a(n-1) <= n + 2/3. If this conjecture is true we have floor(a(n+1)/a(n)) = n. - Mohammed Bouayoun (mohammed.bouayoun(AT)sanef.com), Apr 03 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 23 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001867 | OEIS-A061242

Primes of the form 9*k - 1.

Conjecture: if a(n) = 9x - 1, the integer formed by the repeating digits in the decimal fraction x/a(n) is the smallest integer such that rotating the digits to the left produces a number which is (x+1)/x times larger. An additional conjecture is that the values of x from this sequence are the only values where rotating an integer one to the left produces a value (x+1)/x times as large. (End) The last conjecture is false. For example, for x = 3 we have 230769*(4/3) = 307692, but 9*3-1 = 26 is not in the sequence. - _Giovanni Resta_, Jul 28 2015 Conjecture: Primes p such that ((x+1)^9-1)/x has 4 irreducible factors of degree 2 over GF(p). - _Federico Provvedi_, Jun 27 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 23 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001868 | OEIS-A061243

a(n) = n+r where r is the smallest number such that n divides (n+1)(n+2)(n+3)...(n+r).

It is conjectured that a(n) = n + A002034(n). _Dean Hickerson_ remarks (Oct 07 2001) that the above conjecture is easy to prove: (n+1)(n+2)...(n+r) is congruent to r! (mod n), so n divides (n+1)(n+2)...(n+r) if and only if n divides r!.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 23 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001869 | OEIS-A061256

Euler transform of sigma(n), cf. A000203.

This is also the number of ordered triples of permutations f, g, h in Symm(n) which all commute, divided by n!. This was conjectured by _Franklin T. Adams-Watters_, Jan 16 2006, and proved by J. R. Britnell in 2012.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Apr 21 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001870 | OEIS-A061262

Smallest number representable as the sum of 3 triangular numbers in exactly n ways.

Conjecture: for n large enough, 1 < a(n)/n^2 < 2. - _Benoit Cloitre_, May 10 2003 Conjecture: No term a(n) with n > 2 is congruent to 0 or 3 modulo 5. - _Zhi-Wei Sun_, Feb 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ed Pegg Jr_, Apr 24 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001871 | OEIS-A061300

Least number whose number of divisors is n!.

a(n) = A037019(n!) for all n <= 12 except for 4. I conjecture that this remains true for all larger n, i.e., 4! is the only "exceptional" factorial (see A037019). - _David Wasserman_, Jun 13 2002 Conjecture is confirmed for n <= 30. - _Max Alekseyev_, Sep 05 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and _Labos Elemer_, Apr 26 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001872 | OEIS-A061303

Given a prime p, let s(p,0)=p and let s(p,n+1) be the smallest prime == 1 (mod s(p,n)). Let S(p) be the sequence {s(p,n): n=0,1,...}. Let a(0)=2 and let a(n+1) be the smallest prime not in any of the sequences S(a(0)), ..., S(a(n)).

It is conjectured for primes p and q the sequences S(p) and S(q) are disjoint, unless one is contained in the other.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 26 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001873 | OEIS-A061313

Minimal number of steps to get from 1 to n by (a) subtracting 1 or (b) multiplying by 2.

Conjecture: a(n)+2 is the sum of the terms of the Hirzebruch (negative) continued fraction for the Stern-Brocot tree fraction A007305(n)/A007306(n). - _Andrey Zabolotskiy_, Apr 17 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Jun 06 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001874 | OEIS-A061341

Numbers not ending in 0 whose cubes are concatenations of other cubes.

Conjecture: the last digit of a(n) is in {1, 2, 3, 4, 5, 6, 8} and if it's 3 then there is k>=2 such that a(n) = 3*10^k+3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Erich Friedman_, Jun 06 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001875 | OEIS-A061357

Number of 0<k<n such that n-k and n+k are both primes.

Conjecture: for n>=4 a(n)>0. - _Benoit Cloitre_, Apr 29 2003 Conjectures from _Rick L. Shepherd_, Jun 24 2003: (Start) 1) For each integer N>=1 there exists a positive integer m(N) such that for n>=m(N) a(n)>a(N). (After the first m(N)-1 terms, a(N) does not reappear). In particular, for N=1 (or 2 or 3), m(N)=4 and a(N)=0, giving Benoit Cloitre's conjecture. (cont.) (cont.) Conjectures based upon observing a(1),...,a(10000):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 28 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001876 | OEIS-A061401

From solution to a Picard-Fuchs equation.

Conjecture: Let r be an integer and s a positive integer. The sequence defined by u(n) = [x^(s*n)] F(x)^(r*n) satisfies the supercongruence u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and positive integers n and r. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 10 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001877 | OEIS-A061639

Number of planar planted trees with n non-root nodes and every 2-valent node isolated.

Conjecture: n*a(n) +2*(2-n)*a(n-1) +(14-5*n)*a(n-2) +6*(3-n)*a(n-3) +3*(4-n)*a(n-4)=0. - _R. J. Mathar_, Nov 15 2011 G.f. A(x) satisfies the differential equation (3*x^4 + 6*x^3 + 5*x^2 + 2*x - 1)*A(x)' - (4*x + 2)*A(x) + 2*x + 1 = 0 with A(0) = 0. Mathar's conjectural recurrence above follows from this.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Jun 13 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001878 | OEIS-A061652

Even superperfect numbers: 2^(p-1) where 2^p-1 is a Mersenne prime (A000668).

It is conjectured that there are no odd superperfect numbers, in which case this coincides with A019279.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jun 16 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001879 | OEIS-A061712

Smallest prime with Hamming weight n (i.e., with exactly n 1's when written in binary).

Conjecture: a(n) < 2^(n+3). - _T. D. Noe_, Mar 14 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander D. Healy_, Jun 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001880 | OEIS-A061871

|First digit - second digit + third digit - fourth digit ...| = 2.

Conjecture: a(n) ~ k*n sqrt(log n) for some k. Possibly the constant is sqrt(1650*Pi)/(18*sqrt(log(10))). - _Charles R Greathouse IV_, Jan 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, May 10 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001881 | OEIS-A062049

Integer part of geometric mean of first n primes.

For large n, the ratio prime(n)/a(n) tends to e (very slowly). This was conjectured by Anton Vrba in 2010 (see Rivera, 2010) and proved by Sandor and Verroken (2011). Tighter bounds and asymptotics for a(n) are proved in the note "On the geometric mean of the first n primes" (2016) (see links). Better formulas prime(n)/a(n) ~ exp(1 + several terms A233824(k) / log^k(prime(n))) exist for larger n; see examples in the formula section. - _Alexei Kourbatov_, Feb 27 2016.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jun 06 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001882 | OEIS-A062071

a(n) = [n/1] + [n/(2^2)] + [n/(3^3)] + [n/(4^4)] + ... + [n/(k^k)] + ..., up to infinity, where [ ] is the floor function.

Conjecture: a(n) ~ c * n, where c = A073009. - _Vaclav Kotesovec_, Aug 30 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jun 13 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001883 | OEIS-A062091

a(1) = 2, a(n)= smallest even number which does not divide the product of all previous terms.

Conjecture: a(n) = 2*A062090(n-1) for n>=3. - _R. J. Mathar_, Jul 08 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jun 16 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001884 | OEIS-A062205

Number of alignments of n strings of length 4.

Conjectures: a(n) == 1 (mod 80); for fixed k, the sequence a(n) (mod k) eventually becomes periodic. - _Peter Bala_, Dec 19 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Angelo Dalli_, Jun 13 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001885 | OEIS-A062208

a(n) = Sum_{m>=0} binomial(m,3)^n*2^(-m-1).

Conjectures: a(2*n) = 3 (mod 60) and a(2*n+1) = 1 (mod 60); for fixed k, the sequence a(n) (mod k) eventually becomes periodic with exact period a divisor of phi(k), where phi(k) is Euler's totient function A000010. - _Peter Bala_, Feb 04 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Angelo Dalli_, Jun 13 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001886 | OEIS-A062251

Take minimal prime q such that n(q+1)-1 is prime (A060324), that is, the smallest prime q so that n = (p+1)/(q+1) with p prime; sequence gives values of p.

A conjecture of Schinzel, if true, would imply that such a p always exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 01 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001887 | OEIS-A062282

Number of permutations of n elements with an even number of fixed points.

Conjecture: a(n) = Sum_{k=0..n} A008290(n, k)*A059841(k). - _John Keith_, Jun 30 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ahmed Fares (ahmedfares(AT)my-deja.com), Jul 04 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001888 | OEIS-A062368

Multiplicative with a(p^e) = (e+1)*(e+2)*(4*e+3)/6.

Conjecture: this is the third inverse Mobius transform of the sequence 4^A001221(n). - _R. J. Mathar_, Aug 09 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Jul 07 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001889 | OEIS-A062402

a(n) = sigma(phi(n)).

Makowski and Schinzel conjectured in 1964 that a(n) = sigma(phi(n)) >= n/2 for all n (B42 of Guy Unsolved Problems...). This has been verified for various classes of numbers and proved to be true in general if it is true for squarefree integers (see Cohen's paper). - Antonio G. Astudillo (afg_astudillo(AT)hotmail.com), Sep 07 2002 Atanassov proves the above conjecture. - _Charles R Greathouse IV_, Dec 06 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001890 | OEIS-A062518

Conjectural largest exponent k such that n^k does not contain all of the digits 0 through 9 (in decimal notation) or 0 if no such k exists (for example if n is a power of 10).

Conjectural largest exponent k such that n^k does not contain all of the digits 0 through 9 (in decimal notation) or 0 if no such k exists (for example if n is a power of 10). Conjectured first occurrence of k for k >= 0: 1, 156224, 22148, 7342, 3376, 861, 609, 477, 295, 152, 153, 149, 138, 69, 139, 47, 49, 38, 32, 42, 43, 67, 92, 24, 22, 18, 61, 17, 27, 21, 53, 26, 36, 56, 14, 190, 271, 13, 110, 45, ?40?, 11, 16, ?43?, 19, 29, ..., .

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jun 24 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001891 | OEIS-A062549

a(n) is the smallest number m such that m! is divisible by the left factorial !n (A003422).

Conjecture: a(n) is the largest prime factor of !n for all n > 2. - _Sean A. Irvine_, Apr 01 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jonathan Ayres (jonathan.ayres(AT)btinternet.com), Nov 22 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001892 | OEIS-A062647

Numbers k such that 81^k - 80^k is prime.

Factors of these numbers are of the form p = 2*n*k + 1, just like the repunit numbers, but the PRP tests are significantly slower since there is not a fast mod() property like the repunits, where (number mod Rn) can be calculated with a folding of the digits at length n in base b. However, numbers of this form b^p - (b-1)^p seem to be prime with greater relative frequency than the repunits. While the repunits have a linear fit coefficient that approaches 0.56145948 (see link below for a Generalized Repunit Conjecture), this sequence currently has a linear fit coefficient of 0.22559. - _Paul Bourdelais_, Jul 06 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mike Oakes_, May 18 2001, May 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001893 | OEIS-A062693

Squarefree n such that the elliptic curve n*y^2 = x^3 - x arising in the "congruent number" problem has rank 3.

Conjectural, as detailed in the pages from which it is extracted (see the first few links at the web site mentioned for details), but the conjecture is supported by much numerical and theoretical evidence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Noam D. Elkies_, Jul 04 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001894 | OEIS-A062730

Rows of Pascal's triangle which contain 3 terms in arithmetic progression.

G.f.: (-5x^8+3x^7+7x^6-3x^5+5x^4-5x^3-12x^2+5x+7) /[(1-x)(1-x^2)^2] (conjectured). - _Ralf Stephan_, May 08 2004 a(n)=(n^2+8*n+8)/4 for n>4 and even; a(n)=(n^2+10*n+9)/4 for n>4 and odd (conjectured). - _Colin Barker_, Aug 29 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Erich Friedman_, Jul 11 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001895 | OEIS-A062747

Row sums of (unsigned) staircase array A062746.

Conjectural recurrence: n*(2*n+1)*a(n) = (4*n-3)*(13*n-4)*a(n-1) + 6*(3*n-1)*(3*n-2)*a(n-2) with a(0) = 1, a(1) = 7. - _Peter Bala_, Mar 25 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jul 12 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001896 | OEIS-A062771

Order of automorphism group of the group C_n X C_2 (where C_n is the cyclic group with n elements).

Conjecture: a(n) = 6*phi(n) if n mod 4 = 2 and a(n) = 4*phi(n) if n mod 4 = 0. - _Vladeta Jovovic_, Jul 20 2001 Conjecture confirmed. - _Christian G. Bower_, May 20 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ahmed Fares (ahmedfares(AT)my-deja.com), Jul 18 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001897 | OEIS-A062772

Smallest prime larger than square of n-th prime.

Conjecture: If k is an even integer greater than 4, then a(n) - A054270(n) = k for infinitely many n. - _Clark Kimberling_, Dec 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jul 18 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001898 | OEIS-A062854

First differences of A027424.

Conjecture: a(n) > n/log(n) for n > 2. - _Thomas Ordowski_, Jan 28 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ron Lalonde (ronronronlalonde(AT)hotmail.com), Jun 25 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001899 | OEIS-A062866

Triangle of number of permutations by barycenter.

Conjecture: e.g.f.: Sum_{n>=0} Sum_{k} T(n,k) * t^k * z^n / n! = (1-t^2) * exp(z) / (exp(t*z) - t^2 * exp(z/t)). - _Robert S. Maier_, Jan 17 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_, Jun 26 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001900 | OEIS-A062870

Number of permutations of degree n with greatest sum of distances.

Number of possible values is 1,2,3,5,7,10,13,17,21,... which I conjecture to be A033638. Maximum distance divided by 2 is the same minus one, i.e., 0,1,2,4,6,9,12,16,20,... which seems to be A002620. a(n) = (n/2)!^2 if n is even else n*((n-1)/2)!^2, cf. A092186. - Conjectured by _Vladeta Jovovic_, Aug 21 2007; proved (see the link) by _Max Alekseyev_, Aug 21 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_, Jun 26 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001901 | OEIS-A062887

Smallest multiple of 2n+1 with the property that its digits are odd and each digit is two less (mod 10) than the previous digit, or 0 if no such number exists.

Conjecture: The only numbers n for which a(n)=0 are those for which 2*n+1 is divisible by 125. - _Sean A. Irvine_, Apr 14 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jun 28 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001902 | OEIS-A062894

The prime indices of sequence A067836 (that sequence is conjectured to contain only primes).

The prime indices of sequence A067836 (that sequence is conjectured to contain only primes).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Frank Buss (fb(AT)frank-buss.de), Feb 13 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001903 | OEIS-A062964

Pi in hexadecimal.

Bailey and Crandall conjecture that the terms of this sequence, apart from the first, are given by the formula floor(16*(x(n) - floor(x(n)))), where x(n) is determined by the recurrence equation x(n) = 16*x(n-1) + (120*n^2 - 89*n + 16)/(512*n^4 - 1024*n^3 + 712*n^2 - 206*n + 21) with the initial condition x(0) = 0 (see A374334). They have numerically verified the conjecture for the first 100000 terms of the sequence. - _Peter Bala_, Oct 31 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | Robert Lozyniak (11(AT)onna.com), Jul 22 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001904 | OEIS-A063019

Reversion of y - y^2 + y^3 - y^4.

Conjecture: 32*n*(n-1)*(n-2)*a(n)
-8*(n-1)*(n-2)*(16*n-9)*a(n-1)
+2*(n-2)*(71*n^2+46*n-549)*a(n-2)
+(97*n^3-2250*n^2+10859*n-14850)*a(n-3)
-12*(4*n-15)*(4*n-14)*(4*n-17)*a(n-4)=0. - _R. J. Mathar_, Oct 01 2012 Conjecture confirmed for n >= 5 using the fact that the g.f. satisfies Conjecture: a(n+1) = (1/(n+1))*Sum_{k=0..floor(n/4)} binomial(n+1, n-4*k)*binomial(n+k, n) (compare to the formula from _Peter Bala_ in A215340). - _Joerg Arndt_, Apr 01 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_, Jul 05 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001905 | OEIS-A063043

Collatz-2 (A063041) trajectory starting at 17.

a(23) = a(28) = 9 and a(i) = a(j) <==> i = j for i, j < 28; same cycle values as A063042 but the conjecture that this is the only cycle for all starting values is not true: see A063044

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jul 07 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001906 | OEIS-A063081

Dimension of the space of weight 2n cusp forms for Gamma_0(13).

Conjectured: a(n) = a(n-2) + a(n-3) - a(n-5) for n > 6. Conjectured: G.f.: x^2*(x^4 + x^3 + 4*x^2 + 5*x + 3)/(x^5 -x^3 -x^2 +1). (End) Conjecture: a(n) = a(n-6) + 14, n>7. - _Bill McEachen_, Jun 16 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001907 | OEIS-A063087

Dimension of the space of weight 2n cusp forms for Gamma_0(19).

Conjecture: a(n) = 4*(n-1)-2*floor((n-1)/3) for n > 1. - _Wesley Ivan Hurt_, Dec 27 2016 Conjectures from _Colin Barker_, Jun 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001908 | OEIS-A063089

Dimension of the space of weight 2n cusp forms for Gamma_0(21).

Conjectures from _Colin Barker_, Jun 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001909 | OEIS-A063094

Dimension of the space of weight 2n cusp forms for Gamma_0(26).

Conjectures from _Colin Barker_, Jun 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001910 | OEIS-A063098

Dimension of the space of weight 2n cusp forms for Gamma_0(30).

Conjectures from _Colin Barker_, Jun 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001911 | OEIS-A063099

Dimension of the space of weight 2n cusp forms for Gamma_0(31).

Conjectures from _Colin Barker_, Jun 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001912 | OEIS-A063102

Dimension of the space of weight 2n cusp forms for Gamma_0(34).

Conjectures from _Colin Barker_, Jun 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001913 | OEIS-A063105

Dimension of the space of weight 2n cusp forms for Gamma_0(37).

Conjecture: a(n) = a(n-2)+a(n-3)-a(n-5) for n>6. G.f.: x*(2 +9*x +13*x^2 +10*x^3 +3*x^4 +x^5)/((1 -x)^2*(1 +x)*(1 +x +x^2)). [_Colin Barker_, Sep 27 2012]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001914 | OEIS-A063107

Dimension of the space of weight 2n cusp forms for Gamma_0(39).

Conjectures from _Colin Barker_, Jun 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001915 | OEIS-A063109

Dimension of the space of weight 2n cusp forms for Gamma_0(41).

Conjectures from _Colin Barker_, Jun 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001916 | OEIS-A063110

Dimension of the space of weight 2n cusp forms for Gamma_0(42).

Conjecture: a(n) = 16*n-12 for n>1. a(n) = 2*a(n-1)-a(n-2) for n>3. G.f.: x*(5+10*x+x^2)/(1-x)^2. - _Colin Barker_, Sep 23 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001917 | OEIS-A063111

Dimension of the space of weight 2n cusp forms for Gamma_0(43).

Conjectures from _Colin Barker_, Jun 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001918 | OEIS-A063116

Dimension of the space of weight 2n cusp forms for Gamma_0(48).

Conjecture: For n>=2, a(n) is the difference between the greater of successive pairs of odd triangular numbers(A000217), namely A(B(n-1)+1) - A(B(n-2)+1), where A=A000217 and B=A016813. This reduces to a(n)=2*(8*n-7) (see Formula). (1,3),(15,21) are the first consecutive pairs of odd triangular numbers, and 21-3=18, which is a(2). - _David James Sycamore_, Sep 12 2019 Conjectures from _Colin Barker_, Jun 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001919 | OEIS-A063117

Dimension of the space of weight 2n cusp forms for Gamma_0(49).

Conjectures from _Colin Barker_, Jun 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001920 | OEIS-A063118

Dimension of the space of weight 2n cusp forms for Gamma_0(50).

G.f. (conjectured): x*(x^3 + 12*x^2 + 15*x + 2) / ((x - 1)^2*(x + 1)). Recurrences (conjectured): Closed formula (conjectured): a(n) = (30*n+(-1)^n-27)/2 for n > 1. - _Bruno Berselli_, Mar 10 2011 Recurrence (conjectured): a(n) = 2*a(n-1) -a(n-2) +2*(-1)^n, n > 3. - _Vincenzo Librandi_, Mar 24 2011 Conjecture: a(n) = A007775(4*n - 3), n > 1. - _Bill McEachen_, May 15 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001921 | OEIS-A063120

Dimension of the space of weight 2n cusp forms for Gamma_0(52).

Conjectures from _Colin Barker_, Jun 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001922 | OEIS-A063121

Dimension of the space of weight 2n cusp forms for Gamma_0(53).

Conjectures from _Colin Barker_, Jun 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001923 | OEIS-A063122

Dimension of the space of weight 2n cusp forms for Gamma_0(54).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001924 | OEIS-A063125

Dimension of the space of weight 2n cusp forms for Gamma_0(57).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001925 | OEIS-A063126

Dimension of the space of weight 2n cusp forms for Gamma_0(58).

Conjecture: a(n) = (-19+(-1)^n+30*n)/2 for n>1. a(n) = a(n-1)+a(n-2)-a(n-3) for n>4. g.f.: x*(x+2)*(x^2+6*x+3) / ((x-1)^2*(x+1)). [_Colin Barker_, Feb 07 2013]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001926 | OEIS-A063128

Dimension of the space of weight 2n cusp forms for Gamma_0(60).

Conjectures from _Colin Barker_, Jun 08 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001927 | OEIS-A063130

Dimension of the space of weight 2n cusp forms for Gamma_0(62).

Conjectures from _Colin Barker_, Jun 13 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001928 | OEIS-A063133

Dimension of the space of weight 2n cusp forms for Gamma_0(65).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001929 | OEIS-A063134

Dimension of the space of weight 2n cusp forms for Gamma_0(66).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001930 | OEIS-A063135

Dimension of the space of weight 2n cusp forms for Gamma_0(67).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001931 | OEIS-A063136

Dimension of the space of weight 2n cusp forms for Gamma_0(68).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001932 | OEIS-A063140

Dimension of the space of weight 2n cusp forms for Gamma_0(72).

Conjectures from _Colin Barker_, Sep 23 2012: (Start) Conjectured e.g.f.: 20 + x + 4*exp(x)*(6*x - 5). - _Stefano Spezia_, Mar 02 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001933 | OEIS-A063141

Dimension of the space of weight 2n cusp forms for Gamma_0(73).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001934 | OEIS-A063142

Dimension of the space of weight 2n cusp forms for Gamma_0(74).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001935 | OEIS-A063143

Dimension of the space of weight 2n cusp forms for Gamma_0(75).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001936 | OEIS-A063144

Dimension of the space of weight 2n cusp forms for Gamma_0(76).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001937 | OEIS-A063146

Dimension of the space of weight 2n cusp forms for Gamma_0(78).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001938 | OEIS-A063147

Dimension of the space of weight 2n cusp forms for Gamma_0(79).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001939 | OEIS-A063148

Dimension of the space of weight 2n cusp forms for Gamma_0(80).

Conjecture: a(n) = A063128(n). - _R. J. Mathar_, Jul 21 2015
Conjectures from _Colin Barker_, Jun 08 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001940 | OEIS-A063150

Dimension of the space of weight 2n cusp forms for Gamma_0(82).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001941 | OEIS-A063151

Dimension of the space of weight 2n cusp forms for Gamma_0(83).

Conjectures from _Colin Barker_, Jun 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001942 | OEIS-A063152

Dimension of the space of weight 2n cusp forms for Gamma_0(84).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001943 | OEIS-A063153

Dimension of the space of weight 2n cusp forms for Gamma_0(85).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001944 | OEIS-A063154

Dimension of the space of weight 2n cusp forms for Gamma_0(86).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001945 | OEIS-A063155

Dimension of the space of weight 2n cusp forms for Gamma_0(87).

Conjectures from _Colin Barker_, Nov 08 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001946 | OEIS-A063157

Dimension of the space of weight 2n cusp forms for Gamma_0(89).

Conjecture: a(n) = ((-1)^n+30*n-17)/2 for n>1. a(n) = a(n-1)+a(n-2)-a(n-3) for n>4. G.f.: x*(7+15*x+7*x^2+x^3)/((1-x)^2*(1+x)). [_Colin Barker_, Sep 26 2012]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001947 | OEIS-A063158

Dimension of the space of weight 2n cusp forms for Gamma_0(90).

Conjectures from _Colin Barker_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001948 | OEIS-A063159

Dimension of the space of weight 2n cusp forms for Gamma_0(91).

Conjectures from _Colin Barker_, Jun 11 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001949 | OEIS-A063160

Dimension of the space of weight 2n cusp forms for Gamma_0(92).

Conjectures from _Colin Barker_, Jun 11 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001950 | OEIS-A063161

Dimension of the space of weight 2n cusp forms for Gamma_0(93).

Conjecture: a(n) = a(n-1)+a(n-3)-a(n-4) for n>5. G.f.: x*(1+x)*(9+12*x+10*x^2+x^3)/((1-x)^2*(1+x+x^2)). [_Colin Barker_, Sep 26 2012]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001951 | OEIS-A063162

Dimension of the space of weight 2n cusp forms for Gamma_0(94).

Conjectures from _Colin Barker_, Jun 11 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001952 | OEIS-A063164

Dimension of the space of weight 2n cusp forms for Gamma_0(96).

Conjectures from _Colin Barker_, Jun 08 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001953 | OEIS-A063165

Dimension of the space of weight 2n cusp forms for Gamma_0(97).

Conjecture: a(n) = a(n-2)+a(n-3)-a(n-5) for n>6. x*(7 +24*x +33*x^2 +25*x^3 +8*x^4 +x^5)/((1 -x)^2*(1 +x)*(1 +x +x^2)). [_Colin Barker_, Sep 27 2012]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001954 | OEIS-A063166

Dimension of the space of weight 2n cusp forms for Gamma_0(98).

Conjectures from _Colin Barker_, Jun 11 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001955 | OEIS-A063168

Dimension of the space of weight 2n cusp forms for Gamma_0(100).

a(n) = 2*a(n-1) - a(n-2) for n > 3 (conjectured). G.f.: x*(x^2 + 22*x + 7)/(x - 1)^2 (conjectured). (End) Conjecture: a(n) = a(n-1) + 30, n>2. - _Bill McEachen_, Jun 16 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001956 | OEIS-A063170

Schenker sums with n-th term.

Conjectures: The exponent in the power of 2 in the prime factorization of a(n) (its 2-adic valuation) equals 1 if n is odd and equals n - A000120(n) if n is even. - _Gerald McGarvey_, Nov 17 2007, Jun 29 2012 Amdeberhan, Callan, and Moll (2012) have proved McGarvey's conjectures. - _Jonathan Sondow_, Jul 16 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | Marijke van Gans (marijke(AT)maxwellian.demon.co.uk) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001957 | OEIS-A063247

Dimension of the space of weight 2n cuspidal newforms for Gamma_0(100).

Conjectures from _Colin Barker_, Jun 08 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 10 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001958 | OEIS-A063250

Number of binary right-rotations (iterations of A038572) to reach fixed point.

Conjectured g.f.: 1/(1-x) * Sum_{j>=0} x^(2^j) - (1-x^(2^j)) * Sum_{k>=1} x^((2^j)*(2^k-1)). - _Mikhail Kurkov_, Sep 29 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marc LeBrun_, Jul 11 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001959 | OEIS-A063292

Dimension of the space of weight n cuspidal newforms for Gamma_1(19).

Conjectures from _Colin Barker_, Feb 24 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 14 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001960 | OEIS-A063305

Dimension of the space S_n^{new}(Gamma_1(32)) of weight n cuspidal newforms for Gamma_1(32).

Conjecture: a(n) = 18*n - 41 for n > 2. - _Peter Luschny_, Mar 04 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 14 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001961 | OEIS-A063321

Dimension of the space of weight n cuspidal newforms for Gamma_1(48).

Conjecture: a(n) = 13*(n + (n-1) mod 2) + 14*(n + n mod 2) - 72 for n > 2. - _Peter Luschny_, Mar 05 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 14 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001962 | OEIS-A063337

Dimension of the space of weight n cuspidal newforms for Gamma_1(64).

Conjecture: a(n) = 72*n - 155 for n > 2. - Peter Luschny, Mar 05 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 14 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001963 | OEIS-A063395

T(2n,n) with T(n,m) as in A063394.

Conjecture: (-n+1)*a(n) +(11*n-17)*a(n-1) +(-19*n+43)*a(n-2) +9*(n-3)*a(n-3)=0. - _R. J. Mathar_, Jun 10 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Floor van Lamoen_, Jul 16 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001964 | OEIS-A063520

Sum divides product: number of solutions (r,s,t), r>=s>=t>0, to the equation rst = n(r+s+t).

Conjecturally, includes all positive integers except 2, 4, 7 and 11 - _David W. Wilson_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_ and _Vladeta Jovovic_, Aug 01 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001965 | OEIS-A063524

Characteristic function of 1.

a(n) for 1 <= n <= 4 and conjectured for n > 4 is the number of Hamiltonian circuits in a 2n X 2n square lattice of nodes, reduced for symmetry, where the orbits under the symmetry group of the square, D4, have 1 element: When n=1, there is only 1 Hamiltonian circuit in a 2 X 2 square lattice, as illustrated below. The circuit is the same when rotated and/or reflected and so has only 1 orbital element under the symmetry group of the square.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jul 30 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001966 | OEIS-A063534

Numbers k such that C(k) = H(k) + d(k), where C(k) is Chowla's function A048050, H(k) is the half-totient function A023022 and d(k) is the number of divisors function A000005.

Conjecture: a(n) = A001748(n), n <> 2. - _R. J. Mathar_, Dec 15 2008 The conjecture is false. The least counterexample is a(11546) = 368335 = 5 * 11 * 37 * 181. The next counterexample is 4922335, and there are no more below 10^10. - _Amiram Eldar_, Apr 15 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Aug 02 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001967 | OEIS-A063535

Primes prime(n) such that prime(n+1)^2 < prime(n)*prime(n+2).

Conjecture: these are the primes such that prime(n+2) - 2*prime(n+1) + prime(n) > 0. If so, this sequence with A122535 and A147812 partition the primes. - _Clark Kimberling_, May 16 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel ten Voorde_, Aug 02 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001968 | OEIS-A063544

Smallest number of triangulations of n points in the plane.

In a so-called double circle, half of the points are extremal, and for every edge of the convex hull, there is one interior point that is arbitrarily close to it. The double circles are conjectured to minimize the number of triangulations, and in this case the next terms would be 203748, 674949, 2031054, 6807382, 20662980, ... - _Manfred Scheucher_, Aug 22 2016 Conjecture: a(n) = sqrt(12)^(n-Theta(log n)). - _Manfred Scheucher_, Aug 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 14 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001969 | OEIS-A063647

Number of ways to write 1/n as a difference of exactly 2 unit fractions.

If n is prime a(n)=1. Conjecture: (1/n)*Sum_{i=1..n} a(i) = C*log(n)*log(log(n)) + o(log(n)) with C=0.7... [The conjecture is false. See the plot and the asymptotic formula below. - _Amiram Eldar_, Oct 03 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Jul 23 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001970 | OEIS-A063656

Numbers k such that the truncated square root of k is equal to the rounded square root of k.

Conjecture: the following definition produces a(n) for n >= 1: a(1) = 1; for n > 1, smallest number > a(n-1) satisfying the condition that a(n) is a square if and only if n is a triangular number. - _J. Lowell_, May 13 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Floor van Lamoen_, Jul 24 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001971 | OEIS-A063719

Numbers n such that usigma(cototient(n)) is a prime.

If usigma(x) is prime, it must be a Fermat prime. It is conjectured that there are only 5 Fermat primes. If this conjecture is true, this sequence has no more terms. - _David Wasserman_, Jul 09 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Aug 23 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001972 | OEIS-A063726

a(n) = gcd(1 + Fibonacci(n+1), 1 + Fibonacci(n)).

Conjectured by _Colin Barker_, Jan 30 2018; proved by _Max Alekseyev_, Mar 13 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Aug 11 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001973 | OEIS-A063790

a(n) is the smallest prime gap between n^2 and (n+1)^2.

This is a stronger version of the conjecture (on which the definition of this sequence relies!) that there is always a prime between n^2 and (n+1)^2. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Aug 17 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001974 | OEIS-A063793

a(n) = Min{ q prime | nextprime(q) - q - 1 = prime(n)}, or 0 if none exist.

Conjecture: a(n) > 0 for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Aug 18 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001975 | OEIS-A063865

Number of solutions to +- 1 +- 2 +- 3 +- ... +- n = 0.

The asymptotic formula below was stated as a conjecture by Andrica & Tomescu in 2002 and proved by B. D. Sullivan in 2013. See his paper and H.-K. Hwang's review MR 2003j:05005 of the JIS paper. - _Jonathan Sondow_, Nov 11 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, suggested by _J. H. Conway_, Aug 27 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001976 | OEIS-A063909

Primes p such that 2*p - 5 is also prime.

There are several interesting computer generated conjectures for this sequence at _Jon Maiga_'s Sequence Machine site. - _Antti Karttunen_, Dec 07 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 31 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001977 | OEIS-A063956

Sum of the unitary prime divisors of n.

a(n*m) = a(n) + a(m) - a(gcd(n^2, m)) - a(gcd(n, m^2)) for all n and m > 0 (conjecture). - _Velin Yanev_, Feb 17 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Sep 04 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001978 | OEIS-A064097

A quasi-logarithm defined inductively by a(1) = 0 and a(p) = 1 + a(p-1) if p is prime and a(n*m) = a(n) + a(m) if m,n > 1.

Conjectures: for n>1, log(n) < a(n) < (5/2)*log(n); lim n ->infinity sum(k=1, n, a(k))/(n*log(n)-n) = C = 1.8(4)... - _Benoit Cloitre_, Oct 30 2002 Conjecture: for n>1, floor(log_2(n)) <= a(n) < (5/2)*log(n). - _Robert G. Wilson v_, Aug 10 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | Thomas Schulze (jazariel(AT)tiscalenet.it), Sep 16 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001979 | OEIS-A064098

a(n+1) = (a(n)^2 + a(n-1)^2)/a(n-2), with a(1) = a(2) = a(3) = 1.

Any 3 successive terms of the sequence satisfy the Markov equation x^2 + y^2 + z^2 = 3 xyz. Therefore from the 3rd term on this is a subsequence of the Markov numbers, A002559. Also, we conjecture that the limit of log(log(a(n)))/n is log((sqrt(5) + 1)/2). - Martin Giese (martin.giese(AT)oeaw.ac.at), Oct 13 2005 Conjecture: lim_{n -> infinity} log(log(a(n)))/n exists = 0.48.... - _Benoit Cloitre_, Aug 07 2002. This is true - see below. For this subsequence of the Markoff numbers, we have 2^(F(n-1) - 1) < a(n) < 3^(F(n-1) - 1) for n > 4, where F(n) are the Fibonacci numbers with F(0)=0, F(1)=1, F(n+1) = F(n) + F(n-1). Hence log(log(a(n)))/n tends to log((1 + sqrt(5))/2) as previously conjectured. - _Andrew Hone_, Jan 16 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Santi Spadaro_, Sep 16 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001980 | OEIS-A064107

Continued fraction quotients for e^e = 15.15426224... (A073226).

It was conjectured (but remains unproved) that this sequence is infinite and aperiodic, but it is difficult to determine who first posed this problem. - _Vladimir Reshetnikov_, Apr 27 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Sep 17 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001981 | OEIS-A064114

Unitary weird numbers: unitary abundant (A034683) but not unitary pseudoperfect (A293188).

Conjecture: All the terms are divisible by 10 (tested on the first 10^6 terms). - _Amiram Eldar_, Oct 19 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Sep 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001982 | OEIS-A064170

a(1) = 1; a(n+1) = product of numerator and denominator in Sum_{k=1..n} 1/a(k).

The known G.f. of {F(2*n)^2} from A049684 leds then to the conjectured formula by _R. J. Mathar_ below, and this proves also the recurrence given there.. Conjecture: a(n) = 8*a(n-1)-8*a(n-2)+a(n-3), n>4. G.f.: -x*(2*x^2+x^3-7*x+1)/((x-1)*(x^2-7*x+1)). - _R. J. Mathar_, Jul 03 2009 [For a proof see the W. Lang comment above.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Sep 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001983 | OEIS-A064173

Number of partitions of n with positive rank.

G.f.: Product_{k>=1} (1/(1-q^k)) * Sum_{k>=1} ((-1)^k * (-q^(3*k^2/2+k/2))) (conjectured). - _Thomas Baruchel_, May 12 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Sep 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001984 | OEIS-A064224

Numbers having more than one representation as the product of consecutive integers > 1.

MacLeod and Barrodale prove that the equation x(x+1)...(x+m-1) = y(y+1)...(y+n-1) has no solutions x>1 and y>1 for the following pairs of (m,n): (2,4), (2,6), (2,8), (2,12), (4,8), (5,10). They also show that (2,3) has two solutions and (3,6) has one solution. They conjecture that (2,k) has no solution for k>3. - _T. D. Noe_, Jul 29 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Sep 22 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001985 | OEIS-A064234

The least k such that A063994(k) = Product_{primes p dividing k} gcd(p-1, k-1) = n, or 0 if there's no such k.

Conjecture: a(n) = 0 iff n == 2 mod 4 and n+1 is composite, if this conjecture is true, then a(54), a(110), a(294), etc. would be 0. Another conjecture: If A063994(k) = n and n == 2 mod 4, then n+1 is prime and k is a power of n+1. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Sep 22 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001986 | OEIS-A064237

Numbers k such that k! + 1 is divisible by a square.

Contains A007540(n)-1 for all n. That sequence is conjectured to be infinite. - _Robert Israel_, Jul 04 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Sep 22 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001987 | OEIS-A064337

Minimal prime numbers with increasing prime differences.

Conjecture: a(n) ~ K*n^2*log(n). - _Bill McEachen_, Apr 04 2024 a(n) >= A007504(n) >> n^2 log n. On Cramér's conjecture, a(n) << n^2 log^2 n. - _Charles R Greathouse IV_, Apr 10 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lior Manor_, Sep 13 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001988 | OEIS-A064377

Numbers n such that sigma_4(n) > phi(n)^5.

It is conjectured that there are no other solutions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Sep 27 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001989 | OEIS-A064386

Numbers of the form 2^k+1 or 4^k-2^k+1.

Call m exceptional if the binary cyclic code of length 2^k-1 with zeros w and w^m (w primitive in GF(2^k)) is double-error-correcting for infinitely many k. It is conjectured that this sequence (with the initial terms 1 and 2 omitted) gives all odd exceptional m's. Conjectures from _Colin Barker_, Mar 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 28 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001990 | OEIS-A064390

Numbers of the form 2^k*(2^n+1) or 2^k*(4^n-2^n+1).

Call m exceptional if the binary cyclic code of length 2^k-1 with zeros w and w^m (w primitive in GF(2^k)) is double-error-correcting for infinitely many k. It is conjectured that this sequence (with the powers of 2 omitted) gives all exceptional m's.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 28 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001991 | OEIS-A064402

Numbers k such that prime(k) + k is prime, where prime(k) is the k-th prime.

Except for the first one all terms are even. Conjecture: First differences include all even integers. - _Zak Seidov_, Nov 10 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Sep 28 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001992 | OEIS-A064414

Numbers that divide at least one term in every Fibonacci sequence.

Conjecture: these are the numbers j such that j^2 = Sum_{d|j} phi(d)*A001177(d), where phi = Euler's totient function (A000010). See A232656. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Oct 15 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001993 | OEIS-A064415

a(1) = 0, a(n) = iter(n) if n is even, a(n) = iter(n)-1 if n is odd, where iter(n) = A003434(n) = smallest number of iterations of Euler totient function phi needed to reach 1.

a(n) = A007814(A309243(n)). [From _Rémy Sigrist_'s conjecture in the latter sequence. This reduces to a(n) = sum(p|n, a(p-1)) formula above, thus holds also]

猜想/问题提及段 | historical_or_current_mention_unreviewed | Christian WEINSBERG (cweinsbe(AT)fr.packardbell.org), Sep 30 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001994 | OEIS-A064428

Number of partitions of n with nonnegative crank.

Conjecture: Also the number of integer partitions y of n without a fixed point y(i) = i, ranked by A352826. This is stated at A238394, but Resta tells me he may not have had a proof. The a(2) = 1 through a(7) = 8 partitions without a fixed point are: The above conjecture is true. See Section 4 of the Blecher-Knopfmacher paper in the Links section. - _Jeremy Lovejoy_, Sep 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Sep 30 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001995 | OEIS-A064466

a(0) = 6 and a(n) = Min { m > a(n-1) | both a(n-1) - p and m - p are prime for some prime p } for n > 0.

The initially very frequent case a(k+1) = a(k) + 2 means that there is a twin prime (q, q + 2) with a(k+1) = p + (q + 2) and a(k) = p + q. This might illustrate a certain coherence of two famous conjectures: Goldbach and twin primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Oct 02 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001996 | OEIS-A064537

Limit of the recursion B_[k] = P[k](B_[k-1]), where B_[0] = (1,2,3,4,5,...) and P[k] is the permutation that permutes the entries k+j and 2k + j for all j = 1,..,k.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Klaus Strassburger (strass(AT)ddfi.uni-duesseldorf.de), Oct 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001997 | OEIS-A064591

Nonunitary perfect numbers: k is the sum of its nonunitary divisors; i.e., k = sigma(k) - usigma(k).

Conjecture: let s0 be the sum of the inverses of the even divisors of a number n and s1 the sum of the inverses of the odd divisors of n; then n is in the sequence iff 2*s0-s1 = 1. - _Michel Lagneau_, Jan 21 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, Sep 25 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001998 | OEIS-A064618

Stirling transform of (n!)^2.

Conjecture: for fixed k = 1,2,..., the sequence a(n) (mod k) is eventually periodic with the exact period dividing phi(k), where phi(k) is the Euler totient function A000010. For example, modulo 10 the sequence becomes (1, 1, 5, 9, 1, 1, 5, 9, ...), with an apparent period 1, 1, 5, 9 of length 4 = phi(10) beginning at a(0). - _Peter Bala_, Jan 15 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Karol A. Penson_, Sep 26 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-001999 | OEIS-A064653

Integers not expressible as p + q*a^2, a>1 and p, q are primes.

Dean Hickerson (Oct 12 2001) writes: I suspect that there are no more terms in the sequence. In fact, I'll make the stronger conjecture that for all n>1260, n can be written as p + q*a^2 where a is the smallest prime that does not divide n. For example, for n=10080, a=11 and we have the representation 10080 = 7297 + 23 * 11^2. Hickerson's stronger conjecture holds for n <= 10^9. Therefore, there are no other terms up to 10^9. - _David A. Corneth_, Jun 17 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Oct 07 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002000 | OEIS-A064671

Number of n-digit base 4 biquanimous numbers (with leading 0's allowed, but not all-0 string).

Conjectures from _Colin Barker_, Dec 16 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Oct 09 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002001 | OEIS-A064708

Initial term of run of (at least) n consecutive numbers with just 2 distinct prime factors.

Eggleton and MacDougall prove that no terms exist beyond a(9) and conjecture that a(9) does not exist. - _Jason Kimberley_, Jul 08 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Oct 13 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002002 | OEIS-A064709

Initial term of run of (exactly) n consecutive numbers with just 2 distinct prime factors.

The given terms up to a(8) = 141 are the only terms less than 10^18. To speed the search, note that any string of 6 or more consecutive numbers contains a multiple of 6 and hence must contain a number of the form 2^a * 3^b. Conjecture: 141 is the last term, because numbers with only two different prime factors get pretty rare, so having several in a row near a number of the form 2^a * 3^b is pretty unlikely. - _Joshua Zucker_, May 05 2006 Wenders's argument is incomplete because the consecutive even numbers can have the form 2^a p^b. As stated in the paper by Eggleton and MacDougall, it is still a conjecture that 9 consecutive omega-2 numbers do not exist. - _T. D. Noe_, Oct 13 2008 Eggleton and MacDougall prove that no terms exist beyond a(9) and conjecture that a(9) does not exist. - _Jason Kimberley_, Jul 08 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Oct 13 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002003 | OEIS-A064757

a(n) = n*11^n - 1.

Conjecture: satisfies a linear recurrence having signature (23,-143,121). - _Harvey P. Dale_, May 12 2019 This conjecture is true since a(n) - a(n-1) yields the recurrence 1 + 10*n + 11*n*a(n-1) - (n-1)*a(n) = 0 with polynomial coefficients in n. - _Georg Fischer_, Feb 19 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002004 | OEIS-A064796

Largest integer m such that every permutation (p_1, ..., p_n) of (1, ..., n) satisfies p_i * p_{i+1} >= m for some i, 1 <= i <= n, where p_{n+1} = p_1.

Conjecture: a(n) = (n+1)(n+3)/4 for odd n, a(n) = (n)(n+4)/4 for even n. - _Jud McCranie_, Oct 25 2001

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 21 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002005 | OEIS-A064807

Numbers which are divisible by their digital root (A010888).

Conjecture: All k-multiply perfect numbers belong to this sequence. - _Ivan N. Ianakiev_, May 10 2016 The even perfect numbers are a subsequence. It is an open question whether the odd perfect numbers are a subsequence; this would involve ruling out 148 residue classes mod 2520 as OPNs. - _Charles R Greathouse IV_, Jan 03 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Oct 21 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002006 | OEIS-A064821

Number of ways of writing the numbers 1 .. n in a sequence so that the sum of any two adjacent numbers is a prime; reversing the sequence does not count as different.

I conjecture a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 23 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002007 | OEIS-A064873

First of four sequences representing the lexicographical minimal decomposition of n in 4 squares: n = a(n)^2 + A064874(n)^2 + A064875(n)^2 + A064876(n)^2.

Conjecture: a(n) = A064874(n - A064874(n)^2). - _Zhuorui He_, Oct 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Oct 10 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002008 | OEIS-A064915

Positive integers n that are not of the form p + q * a^2, where p and q are primes and a is the smallest prime not dividing n.

There are no other terms up to 10^7. Conjecture: There are no more terms in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, Oct 13 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002009 | OEIS-A064928

Inverse of A034175 considered as a permutation of the nonnegative integers (which it is conjectured to be).

Inverse of A034175 considered as a permutation of the nonnegative integers (which it is conjectured to be).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, Oct 14 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002010 | OEIS-A064929

Square of A034175 considered as a permutation of the nonnegative integers (which it is conjectured to be).

Square of A034175 considered as a permutation of the nonnegative integers (which it is conjectured to be).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, Oct 14 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002011 | OEIS-A064930

Inverse square of A034175 considered as a permutation of the nonnegative integers (which it is conjectured to be).

Inverse square of A034175 considered as a permutation of the nonnegative integers (which it is conjectured to be).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, Oct 14 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002012 | OEIS-A065001

a(n) = (presumed) number of palindromes in the 'Reverse and Add!' trajectory of n, or -1 if this number is not finite.

Presumably a(196) = 0 (see A016016). Conjecture: There is no n > 0 such that the trajectory of n contains an infinite number of palindromes; the trajectory of n eventually leads to a term in the trajectory of some integer k which belongs to sequence A063048, i.e. whose trajectory (presumably) never leads to a palindrome.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 01 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002013 | OEIS-A065026

a(1) = 1; for n>1, a(n) = smallest number that is not a sum or product of any subset of the numbers a(1) to a(n-1).

Conjectures from _Colin Barker_, Jun 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jonathan Ayres (jonathan.ayres(AT)btinternet.com), Nov 02 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002014 | OEIS-A065069

Numbers n such that Fibonacci(n) is not squarefree, but for all proper divisors k of n, Fibonacci(k) is squarefree.

This sequence is closely related to A001602(n), which gives the index of the smallest Fibonacci number divisible by prime(n). It can be shown that the index of the first Fibonacci number divisible by prime(n)^2 is A001602(n)*prime(n). This sequence is the collection of numbers A001602(n)*prime(n) with multiples removed. For example, A001602(2)*prime(2) = 12, but all multiples of 12 will generated by 6, the first number in this sequence. The Mathematica code assumes that Fibonacci numbers do not have any square primitive prime factors -- an assumption whose truth is an open question. - _T. D. Noe_, Jul 24 2003 Terms after a(12) are conjectures until the factorizations of F(1271), F(1273), etc. are completed. - _Charles R Greathouse IV_, Feb 02 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Len Smiley_, Nov 07 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002015 | OEIS-A065075

Sum of digits of the sum of the preceding numbers.

Conjecture: every non-multiple of 3 does appear in the sequence. - _Franklin T. Adams-Watters_, Jun 29 2009. See A230289. - _N. J. A. Sloane_, Oct 17 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bodo Zinser_, Nov 09 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002016 | OEIS-A065106

Smallest Fibonacci index to produce a factor p^2 (for primes p).

Assuming that there are no square primitive factors in the Fibonacci sequence (an open question), then this sequence continues 1431, 1892, 2147, 2701, 2943, 3029, 3422, 3852, 4378, 4556, 4753, 4970, 5050, 5513, 6162, 6394, 6972, 7550, 7868, 8841, 8862, 9453. This is obtained by sorting the sequence prime(n)*A001602(n). - _T. D. Noe_, Apr 15 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Len Smiley_, Nov 21 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002017 | OEIS-A065107

Start of the permutation of the primes in the order in which p^2 first appears as a factor of a number in the Fibonacci sequence.

Assuming that there are no square primitive factors in the Fibonacci sequence (an open question), then this sequence continues 53, 43, 113, 73, 109, 233, 59, 107, 199, 67, 97, 71, 101, 149, 79, 139, 83, 151, 281, 421, 211, 137, 103, 157, 307, 521. This is obtained by sorting the pairs (prime(n)*A001602(n), prime(n)) by the first position and noting the order of the primes in the second position. - _T. D. Noe_, Apr 15 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Len Smiley_, Nov 21 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002018 | OEIS-A065129

a(n) is the least m such that m/A003285(m) = n, or 0 if no such m exists.

Conjecture: A003285(m) = even or A004613, if m is divisible by A003285(m). [This sentence appears to be saying that all odd terms of this sequence are in A004613.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Nov 14 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002019 | OEIS-A065164

Permutation t->t+1 of Z, folded to N.

This is, starting at a(2) = 4, the same as the "increasing oscillating sequence" shown in Proposition 3.1, p.7 and plotted in the right of figure 1, of Vatter. The same paper, p.4, cites Comtet and uses without giving the A-number of A003319. Abstract: We prove that there are permutation classes (hereditary properties of permutations) of every growth rate (Stanley-Wilf limit) at least lambda = approx 2.48187, the unique real root of x^5-2x^4-2x^2-2x-1, thereby establishing a conjecture of Albert and Linton. - _Jonathan Vos Post_, Jul 18 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002020 | OEIS-A065237

Number of winning length n strings with a 4-symbol alphabet in "same game".

For binary strings, the formula for the number of winning strings of length n has been conjectured by Ralf Stephan and proved by Burns and Purcell (2005, 2007). For b-ary strings with b >= 3, the same problem seems to be unsolved. - _Petros Hadjicostas_, Jul 05 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sascha Kurz_, Oct 23 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002021 | OEIS-A065238

Number of winning length n strings with a 5-symbol alphabet in "same game".

For binary strings, the formula for the number of winning strings of length n has been conjectured by Ralf Stephan and proved by Burns and Purcell (2005, 2007). For b-ary strings with b >= 3, the same problem seems to be unsolved. - _Petros Hadjicostas_, Aug 31 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sascha Kurz_, Oct 23 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002022 | OEIS-A065255

Consider the sequence b(k) such that b(k) and sigma(b(k)) end with the same digit in base 10. Sequence gives values of b(k) such that b(k)/k = 10.

Conjecture: lim n -> infinity b(k)/k = 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002023 | OEIS-A065450

Make an infinite chessboard from the squares in the first quadrant; sequence gives number of squares a knight can reach in n moves starting at the origin.

The first conjecture is true: Partial sums of A047356 = b(n) = (14*(n*(n+1)) + 2*n + 5 + 3*(-1)^n)/8, since A047356(n)=(14*n+1+3*(-1)^n)/4. And b(n) has g.f. (4*x^2 + 2*x + 1)/(-x^4 + 2*x^3 - 2*x + 1). The difference a(n) - b(n) = 0,2,2,0,0,0,0,0,0..., which has g.f. 2*x^2 + 2*x. Then (4*x^2 + 2*x + 1)/(-x^4 + 2*x^3 - 2*x + 1) + 2*x^2 + 2*x = (-2*x^6 + 2*x^5 + 4*x^4 - 4*x^3 + 2*x^2 + 4*x + 1)/(-x^4 + 2*x^3 - 2*x + 1). - _Vim Wenders_, Apr 16 2008 Conjectures: G.f.: [1+6x^2+4x^3-4x^4-2x^5+2x^6]/[(1+x)*(1-x)^3]. For n>3, partial sums of A047356. - _Ralf Stephan_, Mar 06 2004 The second conjecture "For n>3, partial sums of A047356" is also true. From the last possible move, we can either move back to the second last possible move or to b(n)=A047883(n) new squares. So a(n) = a(n-2)+b(n). For n>6, b(n)=7(n-1)+4=A017029(n-1). But a number of the form 7n+4 is naturally the sum of two consecutive terms in A047356 (4=1+3,11=3+8,18=8+10, ...). The conjecture follows. - _Vim Wenders_, Apr 12 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bodo Zinser_, Nov 18 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002024 | OEIS-A065486

Decimal expansion of Product_{p prime} (1 + 1/(p+1)^2).

The constant c in the conjectured formula Sum_{a,b <= x} f(a,b) ~ c * x^2, where f(a, b) = gcd(ab, a + b)/gcd(a, b) (Tóth, 2026, p. 5, eq. (11)). - _Amiram Eldar_, Jun 24 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002025 | OEIS-A065558

Let G_n be the group of invertible 2 X 2 matrices mod n (sequence A000252). a(n) is the maximal degree of an irreducible representation of G_n.

Conjecture: a(2n) = 2*A001615(n). - _Ralf Stephan_, Mar 26 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | Dan Fux (dan.fux(AT)OpenGaia.com or danfux(AT)OpenGaia.com), Nov 29 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002026 | OEIS-A065603

Transposition diameter: maximal number of moves in an optimal sorting of n objects by moving blocks.

Eriksson et al. (2001) conjectured that a(n) = ceiling((n+1)/2) for n >= 3 except for n = 13 and 15.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 02 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002027 | OEIS-A065657

Numbers m that minimize | k / EulerPhi(k) - golden ratio phi | when k runs over all the numbers with the same number of digits as m.

Since phi is irrational, n/EulerPhi(n) can only approximate phi. Probably an open question: can |(n/EulerPhi(n)) - phi| be made arbitrarily close to 0? The listed terms have this property: for r = 1,...,5, all r-digit terms share the same set of prime factors. For example, all three 3-digit terms have prime factors 3 and 13. Furthermore, all listed terms are multiples of 3. I conjecture that these properties hold in general.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Dec 03 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002028 | OEIS-A065704

Number of squares or twice squares dividing n.

a(n) = A005361(2*n) for n>0 (conjectured). - _Werner Schulte_, Jan 15 2018 [This is true only for numbers whose odd part (A000265) is cubefree (A004709). Therefore, the least counterexample is n=3^3=27: a(27) = 2 while A005361(2*27) = 3. - _Amiram Eldar_, Sep 25 2022]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Dec 04 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002029 | OEIS-A065706

Least member p1 of prime octuplets (p1, p2, p3, ..., p8 = p1 + 26), the eight p's being consecutive primes.

See A022011, A022012 and A022013 for the three different possible patterns. The sequence is conjectured to be infinite, although it is not even proved that there are infinitely many

twin primes (p1, p2 = p1+2). - _M. F. Hasler_, May 02 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank Ellermann_, Dec 05 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002030 | OEIS-A065710

Number of 2's in the decimal expansion of 2^n.

See A034293 for indices of zeros: It is conjectured that the last 0 appears at index 168 = A094776(2). More generally, I conjecture that any value x = 0, 1, 2, 3, ... occurs only a finite number of times N(x) = 23, 35, 28, 26, 41, 37, 34, 26, 34, 38, 33, 41, ... in this sequence, for the last time at a well defined index i(x) = 168, 176, 186, 268, 423, 361, 472, 555, 470, 562, 563, 735, ... - _M. F. Hasler_, Feb 10 2023, edited by _M. F. Hasler_, Jul 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Dec 04 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002031 | OEIS-A065711

The Alon-Tarsi constants AT(n).

The extended Alon-Tarsi conjecture states AT(n)<>0 for every positive integer n. Several cases have been proved - see references.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jan Siwanowicz (jansiwanowicz(AT)yahoo.com), Dec 04 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002032 | OEIS-A065712

Number of 1's in decimal expansion of 2^n.

I conjecture that any value x = 0, 1, 2, ... occurs only a finite number of times N(x) = 26, 34, 30, 40, 26, 33, 39, 30, 30, 30, 38, ... in this sequence, for the last time at well defined indices i(x) = 91, 152, 185, 412, 245, 505, 346, 422, 499, 565, 529, 575, ... - _M. F. Hasler_, Jul 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Dec 04 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002033 | OEIS-A065713

Sum of digits of 4^n.

a(n) = A001370(2n). Results given there imply a(n) > log_4(n) + 1/2, n > 0, but we can conjecture & expect a(n) ~ 9*log_10(2)*n. - _M. F. Hasler_, May 18 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 11 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002034 | OEIS-A065714

Number of 3's in decimal expansion of 2^n.

I conjecture that any value x = 0, 1, 2, ... occurs only a finite number of times N(x) = 34, 34, 24, 34, 39, 34, 35, 34, 35, 32, 33, 31, ... in this sequence, for the last time at well defined indices i(x) = 153, 139, 226, 237, 308, 386, 413, 506, 461, 578, 644, 732, 857, 657, 743, 768, 784, 848, 906, ... - _M. F. Hasler_, Jul 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Dec 04 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002035 | OEIS-A065744

Number of 9's in the decimal expansion of 2^n.

See A035064 for the indices of zeros. I conjecture that any value x = 0, 1, 2, ... occurs only a finite number of times N(x) = 37, 27, 36, 46, 20, 31, 32, 30, 46, 29, 22, ... in this sequence, for the last time at well defined indices i(x) = 108, 197, 296, 277, 278, 315, 379, 555, 503, 504, 539, 696, 667, ... - _M. F. Hasler_, Jul 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Dec 04 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002036 | OEIS-A065796

Alternating sum of digits of n^2.

Conjecture; there are an infinite number of values which do not appear in this sequence (in the signed version, of course). The first example appears to be 2. _Sean A. Irvine_ has checked this up to 10^9. - _Robert G. Wilson v_, Dec 10 2001 a(n) == n^2 (mod 11). In particular, values == 2, 6, 7, 8, 10 (mod 11) do not appear, and the conjecture is true. - _Robert Israel_, Oct 24 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | Benny Wegner (jaeger(AT)clan-efg.de), Dec 05 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002037 | OEIS-A065806

A pseudo-random sequence.

Conjectures: -1 < a(n) < n+2; If a(n) = n+1 then a(n+1) = 0; All integers >1 occur in the sequence; For each c in [0,1] there exists a subsequence a(i_j) with (a(i_j)+1)/(i_j+3)-> c, j->infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Klaus Strassburger (strass(AT)ddfi.uni-duesseldorf.de), Nov 21 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002038 | OEIS-A065825

Smallest k such that n numbers may be picked in {1,...,k} with no three terms in arithmetic progression.

Ed Pegg Jr conjectures the 2^k term always equals (3^k+1)/2 and calls these "unprogressive" sets. Jaroslaw Wroblewski (jwr(AT)math.uni.wroc.pl), Nov 04 2003, remarks that this conjecture is known to be false.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ed Pegg Jr_, Nov 23 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002039 | OEIS-A065907

First solution mod p of x^4 = 2 for primes p such that only two solutions exist.

Conjecture: no integer occurs more than three times in this sequence. Confirmed for the first 2399 terms of A007522 (primes < 100000). There are integers which do occur thrice, e.g. 221, 1159.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 29 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002040 | OEIS-A065908

Second solution mod p of x^4 = 2 for primes p such that only two solutions exist.

Conjecture: no integer occurs more than three times in this sequence. Confirmed for the first 2399 terms of A007522 (primes < 100000). In this section, there are no integers which do occur thrice.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 29 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002041 | OEIS-A065909

First solution mod p of x^4 = 2 for primes p such that more than two solutions exist.

Conjecture: no integer occurs more than three time in this sequence. Confirmed for the first 1182 terms of A014754 (primes < 100000). There are integers which do occur thrice, e.g. 6624. Moreover, no integer is first, second, third or fourth solution for more than three primes. Confirmed for the first 2399 terms of A007522 and the first 1182 terms of A014754 (primes < 100000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 29 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002042 | OEIS-A065910

Second solution mod p of x^4 = 2 for primes p such that more than two solution exists.

Conjecture: no integer occurs more than three time in this sequence. Confirmed for the first 1182 terms of A014754 (primes < 100000). In this section, there are no integers which do occur thrice. Moreover, no integer is first, second, third or fourth solution for more than three primes. Confirmed for the first 2399 terms of A007522 and the first 1182 terms of A014754 (primes < 100000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 29 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002043 | OEIS-A065919

Bessel polynomial y_n(4).

Conjecture: (Sum_{n >= 0} a(n)*x^n/n!)/(1 + Sum_{n >= 0} a(n)*x^(n+1)/(n+1)!) = Sum_{n >= 0} A059304(n)*x^n/n!. - _Peter Bala_, Dec 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 08 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002044 | OEIS-A065941

T(n,k) = binomial(n-floor((k+1)/2), floor(k/2)). Triangle read by rows, for 0 <= k <= n.

For example, for N = 11, we get: (1, 2, 4, 3, 5), showing that when confronted with two choices after the 4: (8 and -3), pick the smaller (abs) term, = 3. Then for the next row pick 7 (not used) and repeat the algorithm; succeeding only if the trajectory produces new terms. But 7 is also (-4) mod 11 and 4 was used. Therefore what I call the "r-t table" (for roots trajectory) has only one row: (1, 2, 4, 3, 5). Conjecture: The numbers of terms in the first row is equal to A003558 corresponding to N, i.e., 5 in this case with period 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Len Smiley_, Nov 29 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002045 | OEIS-A065942

Central column of triangle A065941.

Conjecture: 8*n*(n-1)*a(n) - 36*(n-1)*(n-3)*a(n-1) + 6*(-9*n^2+18*n-14)*a(n-2) + 27*(3*n-7)*(3*n-8)*a(n-3) = 0. - _R. J. Mathar_, Jun 13 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Len Smiley_, Nov 29 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002046 | OEIS-A066001

Sum of digits of 5^n.

We can expect and conjecture that a(n) ~ 4.5*log_10(5)*n, but for n ~ 10^3..10^4 there are still fluctuations of +- 1%, e.g., a(10^3)/log_10(5) ~ 4538, a(10^4)/log_10(5) ~ 44518. Modulo 9, the sequence is periodic with period (1, 5, 7, 8, 4, 2) of length 6. No term is divisible by 3, a(n) = (-1)^n (mod 3). - _M. F. Hasler_, May 18 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 11 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002047 | OEIS-A066023

(a(n)^7+1)/(n^7+1) is the smallest integer > 1.

Conjecture: a(n) = A000578(n), n>1. - _R. J. Mathar_, Aug 28 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Dec 11 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002048 | OEIS-A066049

Numbers k such that 2*k^2 - 1 is a prime.

It is conjectured that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 09 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002049 | OEIS-A066052

Number of permutations in the symmetric group S_n with order >= 3.

Conjecture: (-n+3)*a(n) +(n^2-2*n-1)*a(n-1) +(-n+1)*a(n-2) -(n-1)*(n-2)^2*a(n-3)=0. - _R. J. Mathar_, Jan 04 2017 Conjecture: a(n) +(-n-3)*a(n-1) +(2*n-1)*a(n-2) +n*(n-2)*a(n-3) -2*(n-2)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Jan 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | Sharon Sela (sharonsela(AT)hotmail.com), Dec 29 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002050 | OEIS-A066108

Sum n^d over all divisors of n.

Conjectured g.f.: Sum_{m>=1} (m^m*Sum_{k=1..m}(x^(k*m)*Sum_{j=0..k}((-1)^j*(k - j)^m*binomial(m + 1, j)))/(1 - x^m)^(m + 1)), where the inner summation is the Triangle of Eulerian numbers A008292. - _Miles Wilson_, Jan 12 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Dec 05 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002051 | OEIS-A066175

Numbers k such that sigma(phi(sigma(k))) = k.

It is conjectured that there are infinitely many Mersenne primes. So this conjecture also supports that this sequence is infinite. Additionally, if n=2^k-1, where either k=1, or n is a Mersenne prime (A000668), or sigma(n)=3*2^(k-1), then A000217(n) divides sigma(A000217(n)). - _Altug Alkan_, Jul 25 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Dec 15 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002052 | OEIS-A066223

Bisection of A000085.

Conjecture: a(n) +2*(-2*n+1)*a(n-1) +2*(n-1)*(2*n-3)*a(n-2)=0. - _R. J. Mathar_, Nov 24 2012 Remark: the above conjectured recurrence is true and can be obtained by the e.g.f. - _Emanuele Munarini_, Aug 31 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002053 | OEIS-A066337

Decimal equivalent of A066327.

Conjectures from _Chai Wah Wu_, Apr 18 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _George E. Antoniou_, Dec 15 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002054 | OEIS-A066351

Bisection of A007059.

Conjecture: a(n) ~ 0.721... * 4^n / n. - _Vaclav Kotesovec_, Aug 25 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002055 | OEIS-A066352

Pillai sequence: a(n) is the smallest term in A007924 requiring n primes.

On Cramer's conjecture, the number of primes required is O(log* n), where log* is the iterated logarithm, so the rate of growth of a(n) is tetrational in n. - _Charles R Greathouse IV_, Aug 28 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | Copied from www.primepuzzles.net by _Frank Ellermann_, Dec 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002056 | OEIS-A066357

Number of ordered (i.e., planar) trees on 2n edges with every subtree at the root having an even number of edges.

These numbers have the same parity as the Catalan numbers C(n), that is, a(n) is even except when n has the form 2^m - 1. This follows immediately from the formula a(n) = C(2*n+1) + 2*C(2*n) - 2^(2*n + 1)*C(n) given below by Callan. We conjecture that a(n) and C(n) have the same 2-adic valuation (checked up to n = 100). - _Peter Bala_, Aug 02 2016 The conjecture above is true. Let C(n) denote the n-th Catalan number and let s_2(n) denote the binary digit sum. From C(n) = binomial(2*n,n)/(n+1) and Kummer theorem, v_2(C(n)) = s_2(n+1)-1. Using Callan formula a(n) = C(2*n+1) + 2*C(2*n) - 2^(2*n+1)*C(n), the three terms have 2-adic valuations s_2(n+1)-1, s_2(n)+1, and 2*n+s_2(n+1), respectively. Since s_2(n+1) <= s_2(n)+1, the first valuation is strictly smaller than the other two. Hence v_2(a(n)) = s_2(n+1)-1 = v_2(C(n)) for all n >= 0, proving the conjecture. - _Clément Garrot_, Sep 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Louis Shapiro_, Feb 01 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002057 | OEIS-A066385

Smallest maximum of sum of 3 consecutive terms in any arrangement of [1..n] in a circle.

Conjecture: for all odd n > 15, a(n) = f(n). (End) Conjectures from _Chai Wah Wu_, Apr 17 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rainer Rosenthal_, Dec 23 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002058 | OEIS-A066400

Smallest values of t arising in R. L. Graham's sequence (A006255).

It is conjectured that 2 never appears in this sequence. a(n) = 2 if and only if A006255(n) = A072905(n). - _Peter Kagey_, Oct 25 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 25 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002059 | OEIS-A066426

Conjectured values for a(n) = least natural number k such that phi(n+k) = phi(n) + phi(k), if k exists; otherwise 0.

Conjectured values for a(n) = least natural number k such that phi(n+k) = phi(n) + phi(k), if k exists; otherwise 0. It would be nice to remove the word "Conjectured" from the description. - _N. J. A. Sloane_ The values of a(3), a(15) and a(21) listed above, namely 0, are conjectural. There is no natural number k < 10^6 satisfying the "homomorphic condition" phi(n+k) = phi(n) + phi(k) for n = 3, 15, 21.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Dec 27 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002060 | OEIS-A066435

Conjectured values for a(n) = least natural number k such that sigma(n+k) = sigma(n)+sigma(k) if it exists; otherwise 0.

Conjectured values for a(n) = least natural number k such that sigma(n+k) = sigma(n)+sigma(k) if it exists; otherwise 0. It would be nice to remove the word "Conjectured" from the description - _N. J. A. Sloane_. The values of a(3), a(9), a(15) and a(21) listed above, namely 0, are conjectural. There is no natural number k < 10^6 satisfying the "homomorphic condition" sigma(n+k)=sigma(n)+sigma(k) for n=3,9,15,21.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Dec 27 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002061 | OEIS-A066437

a(n) = max_{k} {T(n,k)} where T(n,k) is the "phi/sigma tug-of-war sequence with seed n" defined by T(n,1) = phi(n), T(n,2) = sigma(phi(n)), T(n,3) = phi(sigma(phi(n))), ..., T(n,k) = phi(T(n,k-1)) if k is odd and = sigma(T(n,k-1)) if k is even.

Conjecture: a(n) is always finite; i.e. the sequence {T(n,k)} is eventually periodic for every n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Jan 08 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002062 | OEIS-A066450

a(n) is the conjectured value of the minimal number to which repeated application of the "reverse and add!" algorithm in base n does not terminate in a palindrome. If there is no such number in base n, then a(n) := -1.

a(n) is the conjectured value of the minimal number to which repeated application of the "reverse and add!" algorithm in base n does not terminate in a palindrome. If there is no such number in base n, then a(n) := -1. It would be nice to remove the word "Conjectured" from the description. - _N. J. A. Sloane_ All the terms in this sequence except the first are only conjectures. (See Walker, Irvin on a(10)=196 and Brockhaus on a(2)=22.) Conjectures: a(n) shows the same asymptotic behavior as n^2. For infinitely many n, a(n) = n^2 - n - 1. Again, it is an open question, if the values of the sequence really lead to infinitely many "reverse and add!" steps or not. Is the sequence always positive?

猜想/问题提及段 | historical_or_current_mention_unreviewed | Frederick Magata (frederick.magata(AT)uni-muenster.de), Dec 29 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002063 | OEIS-A066488

Composite numbers k which divide A001045(k-1).

Conjecture: these are composites k such that 2^M(k-1) == -1 (mod M(k)^2 + M(k) + 1), where M(k) = 2^k - 1. - _Amiram Eldar_ and _Thomas Ordowski_, Dec 19 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jan 03 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002064 | OEIS-A066510

Conjectured list of positive numbers which are not of the form r^i-s^j, where r,s,i,j are integers with i>1, j>1.

Conjectured list of positive numbers which are not of the form r^i-s^j, where r,s,i,j are integers with i>1, j>1. This is a famous hard problem and the terms shown are only conjectured values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Don Reble_, Oct 12 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002065 | OEIS-A066523

Crowded numbers: for any k in the sequence, d(k)/k is larger than d(m)/m for all m > k.

It was briefly conjectured that this sequence was the same as the highly composite numbers (A002182) larger than 1, but this is false: 30 is crowded but not highly composite and 50400 is highly composite but not crowded. Is every super-abundant number (A004394) crowded?

猜想/问题提及段 | historical_or_current_mention_unreviewed | Roy Maulbogat (maulbogat(AT)gmail.com), Jan 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002066 | OEIS-A066542

Nonnegative integers all of whose anti-divisors are either 2 or odd.

The following conjectures have been proved by Bob Selcoe. - _Michael Somos_, Feb 28 2014 Additional conjectures suggested by computational experiments: Calculations suggest the following conjecture. This sequence consists of all odd primes and nonnegative powers of 2 and no other terms. This has been verified for to n=100000. _Robert G. Wilson v_ extended the conjecture out to 2^20.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jan 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002067 | OEIS-A066661

Numbers m such that floor((1+1/m)^k)=k for an integer k>1.

Conjecture: a(n)=C*n*log(n) asymptotically with C=1.5...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jan 14 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002068 | OEIS-A066674

Least number m such that phi(m) = A000010(m) is divisible by the n-th prime.

2. If ERH holds, then q = O(p log p)^2, see Heath-Brown (1990), (1992). Explicitly, on the same hypothesis, q < 2(p log p)^2, see Bach and Sorenson (1996). 3. By Linnik's theorem, q = O(p^c) for some c > 0. This is unconditional, but the best known value of c, equal to 5.18 -- see Xylouris (2011) -- is nowhere near 2. Heath-Brown (1992) mentions the conjecture (generalized to Linnik's theorem) that q <= p^2. If true, a(n) and b(n) are identical, since p^2 cannot be 1 mod p. (End) Summary: A066674 and A125878 are the same, and A035095 is probably also the same, but this is an open question.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Dec 19 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002069 | OEIS-A066722

Numbers that can be expressed as the sum of two primes in exactly six ways.

No other terms below 10000. I conjecture there are no further terms in this sequence and A067188, A067189, etc. _R. K. Guy_ (Jan 14 2002) remarks: "I believe that these conjectures follow from a more general one by Hardy and Littlewood (probably in Some problems of 'partitio numerorum' III, on the expression of a number as a sum of primes, Acta Math. 44(1922) 1-70)."

猜想/问题提及段 | historical_or_current_mention_unreviewed | Peter Bertok (peter(AT)bertok.com), Jan 13 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002070 | OEIS-A066746

Conjectured values of a(n) defined by a(n) = least number of applications of f(k) = k^2 + 1 to n to yield a prime, if this number exists; = -1 otherwise.

Conjectured values of a(n) defined by a(n) = least number of applications of f(k) = k^2 + 1 to n to yield a prime, if this number exists; = -1 otherwise. a(9) = -1 is conjectural. If

a(9) is positive, then a(9) > 14. (f^15)(9) already has several thousand digits. (f^n denotes f applied n times.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Jan 16 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002071 | OEIS-A066774

A066728(a(n))=3.

Conjecture: the set of k such that a(k+1)-a(k)=2 is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jan 16 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002072 | OEIS-A066803

a(n) = gcd(2^n + 1, 3^n + 1).

Conjecture: a(2^k) = 1 for k != 1. That is to say, there is no prime p > 5 such that ord(2,p) and ord(3,p) is the same power of 2, where ord(a,p) is the multiplicative order of a modulo p. - _Jianing Song_, Nov 20 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jan 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002073 | OEIS-A066817

Conjectured values of first prime in the orbit f(m), f(f(m)), ..., where f(n) = A067599(n) and m = n-th composite number; or 0 if none exists.

Conjectured values of first prime in the orbit f(m), f(f(m)), ..., where f(n) = A067599(n) and m = n-th composite number; or 0 if none exists. The terms with 0 value listed above are conjectural. There are no primes < 10^30.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Feb 01 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002074 | OEIS-A066831

Numbers k such that sigma(k) divides sigma(phi(k)).

But suppose n=2m, m odd: again with phi(2m)=phi(m) and sigma(2)=3, sigma(phi(2m)) / sigma(2m)=3 => sigma(phi(m)) /3sigma(m)=3 => sigma(phi(m)) / sigma(m)=9; and with sigma(4)=7 sigma(phi(4m))/ sigma(4m)=1 => sigma(2phi(m))/7sigma(m)=1 => sigma(2phi(m))/ sigma(m)=7. So we get the condition sigma(phi(m)) / sigma(m)=9 <=> sigma(2phi(m))/ sigma(m)=7 which will fail. So if there is a (very) big odd number n in A066831 (numbers n such that sigma(n) divides sigma(phi(n))) with A066831(n) = 9, the conjecture is wrong. I admit I could not yet find such a number, nor do i really know it exists, i.e., A067385(9) exists. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jan 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002075 | OEIS-A066840

Sum of positive integers k where k <= n/2 and gcd(k,n) = 1.

Conjecture: a(n) = n*phi(n)/8 + O(n). - _Thomas Ordowski_, Nov 12 2014 Above conjecture is true (see Baum article). - _Chai Wah Wu_, Apr 10 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Jan 20 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002076 | OEIS-A066852

Odd values arising in A066820.

Conjecture: There are an infinite number of odd values in A066820.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jan 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002077 | OEIS-A066861

For x > 0 let f(x) = x/2 if x is even, f(x) = (3*x+1)/2 if x is odd (3x+1 or Collatz problem). Sequence gives numbers such that k-1 applications of f lead to k for some k > 0.

Conjecture: a(n) = n*log(n) asymptotically.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jan 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002078 | OEIS-A066868

a(n) is the least k such that n + Sum_{i=1..k} A005385(i) is prime; or 0 if none exists.

Conjecture: a(n) is nonzero for all n, so every n can be represented as the difference between a prime and a partial sum of the safe primes series. See A066753 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Jan 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002079 | OEIS-A066881

Numbers k such that sigma(phi(k))/sigma(k) is an integer >= 4.

Conjecture: all natural numbers 1,2,3,...,n will eventually occur among the integer values of sigma(phi(k))/sigma(k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jan 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002080 | OEIS-A066883

Number of primes in the interval [p(n), p(n)^2] minus p(n), where p(n) is the n-th prime.

Haga's conjecture (see link below) is that if the integers from 1 to p^2 (p prime) are put in a p by p square in standard order, then there's a transversal consisting of primes; i.e., a set of p primes containing exactly one number in each row and column. E.g., for p=5 the primes 5, 7, 11, 19, 23 work. Since p is needed for the p-th column, primes less than p can't be used. a(n) is the number of primes available minus the number needed for the transversal.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enoch Haga_, Jan 26 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002081 | OEIS-A066885

a(n) = (prime(n)^2 + 1)/2.

a(n) is the average of the numbers from 1 to prime(n)^2. It's also the average of the primes in a prime(n) X prime(n) example of Haga's conjecture (see link below).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enoch Haga_, Jan 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002082 | OEIS-A066886

Sum of the elements in any transversal of a prime(n) X prime(n) array containing the numbers from 1 to prime(n)^2 in standard order.

a(n) is the sum of the primes in a prime(n) X prime(n) example of Haga's conjecture (see link below).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enoch Haga_, Jan 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002083 | OEIS-A066888

Number of primes p between triangular numbers T(n) < p <= T(n+1).

It is conjectured that for n > 0, a(n) > 0. See also A190661. - _John W. Nicholson_, May 18 2011 If the above conjecture is true, then for any k > 1 there is a prime p > k such that p <= (n+1)(n+2)/2, where n = floor(sqrt(2k)+1/2). Ignoring the floor function we can obtain a looser (but nicer) lower bound of p <= 1 + k + 2*sqrt(2k). - _Dmitry Kamenetsky_, Nov 26 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002084 | OEIS-A066919

a(n) = least number of applications of f to n to reach 1, where f is defined by f(n) = phi(n) if n is even; = sigma(n) if n is odd.

a(n) is in [0,19] for n < 10^5. Conjecture: a(n) exists for all n, i.e., repeated application of f to n eventually yields 1, for any n. The only way this could fail is if n, f(n), f(f(n)), ... are all odd squares. The conjecture is true, since sequence A055021 (smallest x such that n iterations of sigma() are required for the result to be >= 2x) is complete. - _Vim Wenders_, Apr 07 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Jan 23 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002085 | OEIS-A066999

a(n) = 3^n * Sum_{i=1..n} i^3/3^i.

Conjecture: g.f.:(-1-x^2-4*x)/((3*x-1)*(x-1)^4). - Maksym Voznyy (voznyy(AT)mail.ru), Jul 27 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jan 27 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002086 | OEIS-A067016

Start with a(0)=1, a(1)=4, a(2)=3, a(3)=2; for n>=3, a(n+1) = max_i (a(i)+a(n-i)).

Conjecture: a(n) = (-3+(-1)^n+4*n)/2 for n>3. G.f.: -(2*x^6 -2*x^5 -6*x^4 +4*x^3 +2*x^2 -3*x -1) / ((x -1)^2*(x +1)). - _Colin Barker_, Apr 01 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 17 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002087 | OEIS-A067057

Let A(n) = {1,2,3,...n}. Let B(r) and C(n-r) be two subsets of A(n) having r and n-r elements respectively, such that B(r) U C(n-r) = A(n) and B and C are disjoint; then a(n) = sum of the products of all combination sums of elements of B and C for r =1 to n-1.

Conjecture: G.f.: (-2*x^2*(x+1))/(2*x-1)^5. [From Maksym Voznyy (voznyy(AT)mail.ru), Jul 26 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jan 02 2002, May 31 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002088 | OEIS-A067069

Numbers n such that for every k the following condition holds: (*) The number of nonnegative solutions of the Diophantine equation x^2 + n*y^2 = 1+n*k^2 equals one half of the number of divisors of 1+n*k^2 if 1+n*k^2 is not a square and one half of 1 + the number of divisors of 1+n*k^2 if 1+n*k^2 is a square.

The sequence is conjectural. It depends on a proof of conjecture 1: If condition (*) holds for k and n < u(k), then it holds for k+1 and n, where u(k) is some nondecreasing function of k. Computationally this is confirmed for 1 < k < 300 and n < 100, 2 < k < 250 and n < 238, 3 < k < 250 and n < 282, 5 < k < 200 and n < 652, 8 < k < 200 and n < 2088, 9 < k < 200 and n <= 1000000. No further terms up to 1000000 were found; this suggests conjecture 2: The sequence is finite and 1848 its last term. Perhaps easier to prove is the stronger conjecture 3: For every n > c there is a k < d such that condition (*) is violated for k and n. Computationally this is confirmed for n <=1000000 and c = 1848, d = 11.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Holger Stephan (stephan(AT)wias-berlin.de), Feb 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002089 | OEIS-A067076

Numbers k such that 2*k + 3 is a prime.

Conjecture: a(n) = A008507(n) + n - 1 = A005097(n) - 1 = A102781(n+1) - 1. - _R. J. Mathar_, Jul 07 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Williams_, Aug 17 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002090 | OEIS-A067129

Numbers k such that sigma(k-3) + sigma(k+3) = sigma(2*k).

Conjecture: sequence gives odd values only.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002091 | OEIS-A067151

Number of regions in regular n-gon which are quadrilaterals (4-gons) when all its diagonals are drawn.

Conjecture: a(n) ~ c * n^4. Is c = 1/64 ? - _Bill McEachen_, Mar 03 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sascha Kurz_, Jan 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002092 | OEIS-A067182

Smallest Fibonacci number with digit sum n, or -1 if no such number exists.

Starting at n = 11, the terms a(11), a(12), ... are probably -1, -1, -1, 4181, -1, -1, 89, -1, 2584, 10946, 317811, 1597, 514229, 987, -1, -1, 46368, 28657, 196418, 2178309, 1346269, -1, 701408733, 3524578, 9227465, -1, 5702887, -1, -1, -1, 433494437, -1, 63245986, 39088169, -1, 267914296, -1, ... However, these -1's are only conjectural. It appears that 0.9*n < A004090(n) < n for all but a few small n: In the range [0..10^5] the slope of A004090 is roughly 0.93. I conjecture that A004090(n) - n has 92 as its maximum, at n = 2619. This would prove that the given -1's are correct. - _M. F. Hasler_, Dec 26 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jan 09 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002093 | OEIS-A067191

Numbers that can be expressed as the sum of two primes in exactly five ways.

There are no other terms below 10000 and I conjecture there are no further terms in this sequence and A067188, A067189, etc. - Peter Bertok (peter(AT)bertok.com), Jan 13 2002 I believe that these conjectures follow from a more general one by Hardy and Littlewood (probably in Some problems of 'partitio numerorum' III, on the expression of a number as a sum of primes, Acta Math. 44(1922) 1-70). - _R. K. Guy_, Jan 14 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jan 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002094 | OEIS-A067274

Number of ordered integer pairs (b,c), with -n<=b<=n, -n<=c<=n, such that both roots of x^2+bx+c=0 are integers.

Conjecture: The difference a(n)-a(n-1) is 6 if and only if n is a prime number. This has been checked up to about n=300 and may be easy to prove. Preceding conjecture is a corollary of Jovovic's formula below. - _Eric M. Schmidt_, Aug 19 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Feb 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002095 | OEIS-A067281

Number of permutations of {1,2,3,...,n} where the elements of n are considered indistinguishable if they differ by a power of 2 (for example 3, 12 and 24 are all considered equivalent).

The conjecture a(2n+1)=(2n+1)a(2n) is obviously true from the definition of the sequence and the fact that 2n+1 is the smallest element of its equivalence class. - Brian Rothbach (rothbach(AT)Math.Berkeley.EDU), Sep 15 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | Brian Rothbach (rothbach(AT)math.berkeley.edu), Feb 23 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002096 | OEIS-A067299

Second column of triangle A067298.

Conjecture: n*(17*n-7)*a(n) +2*(-119*n^2+270*n-96)*a(n-1) -16*(17*n+10)*(2*n-3)*a(n-2)=0. - _R. J. Mathar_, Jul 21 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Feb 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002097 | OEIS-A067360

a(n) = 17^n sin(2n arctan(1/4)) or numerator of tan(2n arctan(1/4)).

Conjectures from _Colin Barker_, Jul 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Barbara Haas Margolius (b.margolius(AT)csuohio.edu), Jan 17 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002098 | OEIS-A067368

a(n) is the smallest positive even integer that cannot be expressed as the product of two or three previous terms (not necessarily distinct).

Conjecture: a(n) = a(n-1) + 2 if (n = 2a(k) + k + 1) or (n = 2a(k) + k) for some k, otherwise a(n) = a(n-1) + 4. This has been confirmed for several hundred terms. The above conjecture is correct because there are 2*(a(k+1)-a(k)) terms that are not divisible by 4 in the k-th interval which are determined by terms that are divisible by 4. For example, there are 2*(a(2)-a(1)) = 2*(6-2) = 8 terms between a(5) = 16 and a(14) = 48 because numbers of the form 2*s are always terms where s is an odd number. So first differences of a(n) determine the corresponding intervals and the formula above always holds. - _Altug Alkan_, Sep 24 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jeremiah K. Hower (jhower(AT)vt.edu), Jan 20 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002099 | OEIS-A067392

Sum of numbers <= n which have common prime factors with n.

V. S. Guba conjectured that for any positive n and prime p, a(n) != a(n+p). - _Max Alekseyev_, May 08 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jan 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002100 | OEIS-A067396

a(n) is the position of the n-th occurrence of the pair "2,2" in A067395 (the first difference sequence of A067368).

A comparison of this sequence with A067368 suggests the following conjecture: a(n)=2*A067368(n)+n-1. This has been confirmed for several hundred terms. Above conjecture is true, and it is same as conjecture in Formula section of A067368. - _Altug Alkan_, Sep 26 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jan 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002101 | OEIS-A067523

The smallest prime with a possible given digit sum.

Except for 3 no other prime has a digit sum which is a multiple of 3. Hence the possible digit sums are 2,3,4,5,7,8,10,11,13,14,16,..., etc. Conjecture: For every possible digit sum there exists a prime. For n > 2, this is (conjecturally) the smallest prime with digit sum A001651(n). - _Lekraj Beedassy_, Mar 04 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Feb 14 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002102 | OEIS-A067567

Odd numbers with an odd number of partitions.

We generalize the above conjecture as follows. We further conjecture that A305123(n) is odd iff n is a term of this sequence. (End) The conjectures above and in the Maple section by Somos and Bala are true (see Fried link). - _Sela Fried_, Oct 15 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Jan 30 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002103 | OEIS-A067576

Array T(i,j) read by downward antidiagonals, where T(i,j) is the j-th term whose binary expansion has i 1's.

Row i shows the numbers m, in increasing order, such that 2^i divides binomial(2*m, m) and 2^(i+1) does not. Let T'(i, j) = 1 + T(i, j) for all i and j. Conjecture: every column of T' is eventually geometric with common ratio 2. Assuming the conjecture is true, let d(k) be the number in column k of the row in which this doubling begins; then d(1) = 1, d(2) = 1, d(3) = 2, d(4) = 3, d(5) = 4. - _Clark Kimberling_, Nov 08 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jan 30 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002104 | OEIS-A067579

Conjectured values of a(n) defined by: a(n) = first prime appearing in the orbit n, f(n), f(f(n)),...., if it exists; = 0 otherwise, where f(n) = n + sigma(n).

Conjectured values of a(n) defined by: a(n) = first prime appearing in the orbit n, f(n), f(f(n)),...., if it exists; = 0 otherwise, where f(n) = n + sigma(n). The values n with a(n) = 0 listed here are conjectural. There is no prime < 10^30 appearing in the orbits of these values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Jan 30 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002105 | OEIS-A067589

Numbers k such that A067588(k) is an odd number.

Apparently groups of two odd pentagonal numbers (A000326, A014632) followed by two odd 2nd pentagonal numbers (A005449), which leads to the conjectured generating function x*(x^2+4*x+1)*(x^4-2*x^3+4*x^2-2*x+1)/((x^2+1)^2*(1-x)^3). - _R. J. Mathar_, Jul 26 2009 For positive integer m, define b_m(n) = Sum_{k = 1..n} k^(2*m+1)*A000009(k)*A000009(n-k). We conjecture that

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Jan 31 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002106 | OEIS-A067590

Number of partitions of n into odious numbers (A000069).

Conjecture: a(n) ~ c * exp(Pi*sqrt(n/3)) / n^(5/8), where c = 0.221864833... - _Vaclav Kotesovec_, Jul 06 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Jan 31 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002107 | OEIS-A067603

Least k such that gcd(prime(k)+1, prime(k+1)+1) = 2n.

Conjecture: a(n) = least k such that A001223(k) = 2n and A000040(k) == -1 (mod 2n). - _Zak Seidov_, Aug 16 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jan 31 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002108 | OEIS-A067611

Numbers of the form 6xy +- x +- y, where x, y are positive integers.

Conjecture 1: With u(k) = floor(k(k + 1)/4) one has A071538(a(u(k))*6) = a(u(k)) - u(k) + 1, for k >= 2 (u > 1). Conjecture 2: In the interval [T(k-1)+1, T(k)], with T(k) = A000217(k), k >= 2, there exists at least one number that is not a member of the present sequence. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Feb 01 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002109 | OEIS-A067677

Diagonals of the prime-composite array, B(m,n) which are zeros from the Second Borve Conjecture.

Diagonals of the prime-composite array, B(m,n) which are zeros from the Second Borve Conjecture. The Second Borve Conjecture states that there are infinitely many zero-only diagonals.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Feb 04 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002110 | OEIS-A067681

Diagonals and antidiagonals of the prime-composite array, B(m,n) which are zeros from the Third Borve Conjecture.

Diagonals and antidiagonals of the prime-composite array, B(m,n) which are zeros from the Third Borve Conjecture. The Third Borve Conjecture states that there are infinitely many integers m for which the m-th diagonal and m-th antidiagonal are both zero-only.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Feb 04 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002111 | OEIS-A067723

Nontrivial palindromes k such that phi(k) is also a palindrome.

Conjecture: There are no terms with an even number of digits. - _Klaus Brockhaus_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Feb 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002112 | OEIS-A067730

Numbers n such that sigma(n-1) + sigma(n+1) = sigma(2n).

Conjecture: sequence contains odd values only. - _Benoit Cloitre_, Feb 18 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Feb 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002113 | OEIS-A067742

Number of middle divisors of n, i.e., divisors in the half-open interval [sqrt(n/2), sqrt(n*2)).

The proofs are straightforward. These results were conjectured by _Omar E. Pol_ in 2017. (End) Conjecture 1: This is the central column of the isosceles triangle of A249351. Conjecture 2: a(n) is also the width of the terrace at the n-th level in the main diagonal of the pyramid described in A245092. Conjecture 3: a(n) is also the number of central subparts of the symmetric representation of sigma(n). For more information see A279387. Conjectures 2 and 3 were proposed after _Michel Marcus_'s conjecture in A237593. (End) Conjectures 1, 2, and 3 are all true. - _N. J. A. Sloane_, Feb 11 2021 The above conjecture is essentially the same as _Michel Marcus_'s conjecture in A237593. - _Omar E. Pol_, Dec 20 2020 Conjecture: a(n) = A082647(n) - A131576(n) = A001227(n) - 2*A131576(n). - _Omar E. Pol_, Feb 06 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marc LeBrun_, Jan 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002114 | OEIS-A067745

Numerator of ((3*n - 2)/(n^(2*n - 1)*(2*n - 1)*4^(n - 1))).

Conjecture: Odd part of 3n-2. - _Ralf Stephan_, Nov 18 2010 Conjecture is true. Note that gcd(3n-2,2n-1)=1 (because 2(3n-2)-3(2n-1) = -1) and gcd(3n-2,n) = 1 or 2. If 2^k | (3n-2), then k <= log_2(3n-2) < (n-1)/2 for n >= 11. So only the cases n <= 10 need to be checked individually. - _Robert Israel_, May 16 2017 Assuming the above conjecture, a(n) = a((8+(3*n-2)*4^k)/12), for all k >= 1. - _L. Edson Jeffery_, Feb 15 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marc LeBrun_, Jan 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002115 | OEIS-A067786

Numbers n such that phi(n) = 2*pi(n) where pi(n) = A000720(n) and phi(n) = A000010(n).

There are no more terms < 10^10. Conjectured to be complete. - _Donovan Johnson_, Nov 24 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002116 | OEIS-A067793

Nonprimes n such that phi(n) > 2n/3.

Conjecture: Odd composite n such that (n^2 + 8) mod 3 = 0. (All primes > 3 meet this criterion). - _Gary Detlefs_, May 03 2012 Both conjectures are wrong. The first counterexample is 385. - _Robert Israel_, May 17 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002117 | OEIS-A067843

Least solution k>n of phi(k-n)+phi(k+n)=phi(2k).

Conjecture: a(n) >= n+4 for all n. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Feb 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002118 | OEIS-A067864

Numbers k such that k divides the sum of digits of 6^k.

Conjecture: the sequence is finite, since (sum of the digits of 6^k)/k -> log_10(6)*4.5 ~ 3.50168 as k->infinity (this is also a conjecture). - _Robert Gerbicz_, May 08 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and _Shyam Sunder Gupta_, Feb 16 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002119 | OEIS-A067873

Numbers x satisfying x^2 - D*y^2 = 1 for more than one value of D distinct from x^2 - 1.

Sequence includes mostly odd entries in consecutive pairs. It seems reasonable to conjecture that an even entry e satisfies big omega(E) - omega(E) > 2, where E = e^2 - 1. (See A001222 and A001221.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Feb 25 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002120 | OEIS-A067874

Positive integers x satisfying x^2 - D*y^2 = 1 for a unique integer D.

Conjecture: All terms of A002110 > 1 are a subsequence. - _Griffin N. Macris_, Apr 11 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Feb 25 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002121 | OEIS-A067955

Number of dissections of a convex polygon by nonintersecting diagonals into polygons with even number of sides and having a total number of n edges (sides and diagonals).

Conjecture: Number of lattice paths that do not cross below the x-axis from (1,0) to (n,0) using up-step (1,1) and down-steps {(1,-z): z is a positive even integer}. For example, a(8) = 1: [(1,1)(1,1)(1,1)(1,1)(1,1)(1,1)(1,-6)]. - _Nicholas Ham_, Aug 24 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Mar 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002122 | OEIS-A067992

a(0)=1 and, for n > 0, a(n) is the smallest positive integer such that the ratios min(a(k)/a(k-1), a(k-1)/a(k)) for 0 < k <= n are all distinct.

Every positive rational number appears exactly once as the ratio of adjacent terms (in either order). Conjecture: adjacent terms are always relatively prime. - _Franklin T. Adams-Watters_, Sep 13 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Feb 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002123 | OEIS-A068012

Number of subsets of {1,2,3,...,n} that sum to 0 mod 6.

Conjecture: a(n) = 2*a(n-1) for n > 2 and 3 does not divide n-1. This conjecture holds for 1 <= n <= 2000. If this conjecture is true then a(3*k + 1 + r) = ((2*4^k - 2)/3 + 1)*2^(k-1+r) where k >= 1 and r in {0, 1, 2}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002124 | OEIS-A068073

Period 4 sequence [1, 2, 3, 2, ...].

Conjecture: a(n) = Sum_{k=0..n} e^(i*Pi*(A000120(A001045(n)) - A001045(A000120(n)))), i=sqrt(-1). - _Paul Barry_, Jan 14 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Mar 01 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002125 | OEIS-A068082

a(1) = 1, a(n) = smallest triangular number of the form k*a(n-1) + 1 for some positive integer k.

Conjectures from _Colin Barker_, May 30 2016: (Start) Conjectures confirmed by _Robert Israel_, Jun 05 2018 (see link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Feb 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002126 | OEIS-A068085

Numbers k such that k and 10*k are both triangular numbers.

Conjecture: satisfies a linear recurrence having signature (1, 0, 1442, -1442, 0, -1, 1). - _Harvey P. Dale_, Sep 03 2020 This conjecture is true because of the connection between (generalized) Pell equations and continued fractions of quadratic irrationals. - _Georg Fischer_, Feb 23 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Feb 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002127 | OEIS-A068128

Triangular numbers with sum of digits = 6.

1. The sequence is unbounded, as the (2*10^k +1)-th triangular number is a term. 2. The sum of the digits of triangular numbers in most cases is a triangular number. 3. Conjecture: For every triangular number T there exist infinitely many triangular numbers with sum of digits = T.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Feb 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002128 | OEIS-A068129

Triangular numbers with sum of digits = 10.

1. The sequence is unbounded, as the (2*10^k +2)-th triangular number is a term. 2. The sum of the digits of triangular numbers in most cases is a triangular number. 3. Conjecture: For every triangular number T there exist infinitely many triangular numbers with sum of digits = T.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Feb 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002129 | OEIS-A068130

Triangular numbers with sum of digits = 15.

1. The sequence is unbounded, as the (2*10^k +3)-th triangular number is a term. 2. The sum of the digits of triangular numbers in most cases is a triangular number. 3. Conjecture: For every triangular number T there exist infinitely many triangular numbers with sum of digits = T.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Feb 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002130 | OEIS-A068131

Triangular numbers with sum of digits = 21.

1. The sequence is unbounded, as the (5*10^k + 6)-th triangular number is a term. 2. The sum of the digits of triangular numbers in most cases is a triangular number. 3. Conjecture: For every triangular number T there exist infinitely many triangular numbers with sum of digits = T.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Feb 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002131 | OEIS-A068132

Triangular numbers with sum of digits = 28.

Conjecture: For every triangular number T there exist infinitely many triangular numbers with sum of digits = T.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Feb 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002132 | OEIS-A068133

First triangular number with digit sum = n-th triangular number.

The sum of the digits of triangular numbers in most cases is a triangular number. Conjecture: For every triangular number T there exist infinitely many triangular numbers with sum of digits = T. There exist only finitely many triangular numbers whose digit sum is T(0)=0 or T(1)=1: T(0)=0 is, of course, the only one with digit sum 0, and T(1)=1 and T(4)=10 are the only two with digit sum 1. However, for digit sums equal to each of at least the next several triangular numbers, the conjecture can be easily confirmed by observing that, e.g., T(2), T(20), T(200), T(2000), etc., all have digit sum T(2)=3; T(2+1), T(20+1), T(200+1), T(2000+1), etc., all have digit sum T(3)=6; T(20+2), T(200+2), T(2000+2), T(20000+2), etc., all have digit sum T(4)=10; and, similarly, for all sufficiently large values of j, triangular numbers of the form T(2*10^j+m), where m = 3, 9, 23, 34, 132, 368, 1332, 3943, 19388, 88248, 244948, 1788848, 9838483, 19994343, respectively, will have digit sums T(5)=15, T(6)=21, ..., T(18)=171, respectively. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Feb 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002133 | OEIS-A068192

Let a(1)=2, f(n) = 4*a(1)*a(2)*...*a(n-1) for n >= 1 and a(n) = f(n)-prevprime(f(n)-1) for n >= 2, where prevprime(x) is the largest prime < x.

The terms are easily seen to be distinct. It is conjectured that every element is prime. Do all primes occur in the sequence?

猜想/问题提及段 | historical_or_current_mention_unreviewed | Frank Buss (fb(AT)frank-buss.de), Feb 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002134 | OEIS-A068193

The prime indices of sequence A068192. (That sequence is conjectured to contain only primes.).

The prime indices of sequence A068192. (That sequence is conjectured to contain only primes.).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Frank Buss (fb(AT)frank-buss.de), Feb 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002135 | OEIS-A068206

Let N(2n) denote the numerator of B(2n), the 2n-th Bernoulli number and D(2n) the denominator; sequence gives values of n such that gcd(N(2n),D(2n-2))=7.

Conjecture: All terms are of the form 7 + 21*j. - _Vaclav Kotesovec_, Apr 29 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Mar 23 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002136 | OEIS-A068540

Numbers k such that phi(k) = bigomega(k)*tau(k)^2.

Conjecture: a(116) = 14137200 is the last term of this sequence (checked up to 10^10). - _Amiram Eldar_, Jun 24 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Mar 25 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002137 | OEIS-A068591

Nonessential triangular numbers (conjectured).

Nonessential triangular numbers (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ed Pegg Jr_, Mar 30 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002138 | OEIS-A068595

Number of functions from {1,2,...,n} to {1,2,...,n} such that the sum of the function values is 0 mod 3.

This conjecture is false for n=8, n=14, and n=20. - _Sean A. Irvine_, Feb 26 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Mar 13 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002139 | OEIS-A068598

Number of maximal sets of partitions of n with property that all parts in all partitions in the set are distinct.

Conjecture: a(n) asymptotically grows as a*exp(bx^c), where a≈0.57, b≈0.062, c≈1.54. - _Elijah Beregovsky_, Nov 12 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Mar 28 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002140 | OEIS-A068689

Primes which are a sandwich of numbers made of only one decimal digit between two 9's.

Conjecture: Inner digits 1, 2 and 8 are the only repeating digits for which the resulting numbers can be prime for outer digits 9. I.e., 9444..4449, 9555..5559, 9777..7779 are composite. The cases for inner digits 0, 3, 6 and 9 give composite numbers. - _Cino Hilliard_, Jul 11 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 02 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002141 | OEIS-A068704

a(n) = smallest prime obtained as the concatenation of n^k, n^(k-1), n^(k-2), ..., n^2, n, 1 for some k >= 1; or 0 if no such prime exists.

Conjecture: a(n) > 0, that is, for every n there exists a k which yields a prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 04 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002142 | OEIS-A068828

Geometrically weak primes: primes that are smaller than the geometric mean of their neighbors (2 is included by convention).

For primes from 3 to 99973229 (and perhaps much beyond), primes in this sequence are the same as the primes in A051635. So one may conjecture that this is essentially the same as A051635. - _R. J. Mathar_, Sep 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 08 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002143 | OEIS-A068869

Smallest number k such that n! + k is a square.

According to my conjecture that n! + n^2 != m^2 for n >= 1, m >= 0 (see A004664), for all terms: a(n) != n^2. - _Alexander R. Povolotsky_, Oct 06 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 13 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002144 | OEIS-A068873

Smallest prime which is a sum of n distinct primes.

Conjectured terms a(50)-a(76): 5147, 5623, 5591, 6079, 6101, 6599, 6607, 7151, 7151, 7699, 7699, 8273, 8293, 8893, 8893, 9521, 9547, 10211, 10223, 10889, 10891, 11597, 11617, 12343, 12373, 13099, 13127. - _Jean-François Alcover_, Apr 22 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002145 | OEIS-A068875

Expansion of (1 + x*C)*C, where C = (1 - (1 - 4*x)^(1/2))/(2*x) is the g.f. for Catalan numbers, A000108.

If A(x)^t = 1 + 2*t*x + Sum_{n >= 2} t*P(n,t)*x^n, then we conjecture that all the zeros of the polynomial P(n,t) lie on the vertical line Re(t) = -n/2 in the complex plane. - _Peter Bala_, Oct 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002146 | OEIS-A068890

a(1) = 1; a(n) = smallest nontrivial n-th power with property that digits alternate in parity.

Conjecture: the sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 20 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002147 | OEIS-A068891

a(1) = 1; a(n) = smallest nontrivial n-th power with property that digits alternate in parity individually as well as in concatenation with previous terms.

Conjecture: the sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 20 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002148 | OEIS-A068912

Number of n step walks (each step +/-1 starting from 0) which are never more than 3 or less than -3.

Conjecture: a(n) = floor((1+r)^(n/2)*(r+(2*(1+r))^(1/2)+(-1)^n*(r-(2*(1+r))^(1/2)))/4) where r = 1 + 2^(1/2). - _Peter Luschny_, Sep 20 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Mar 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002149 | OEIS-A068962

Number of successive terms of A028356 that add to n; or length of n-th term of A028355.

Conjectures from _Chai Wah Wu_, Apr 18 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 08 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002150 | OEIS-A068991

Numbers k such that Sum_{d divides k} sigma(d)/phi(d) is an integer.

Conjecture: if k is in the sequence and k is squarefree then the denominator of the 2k-th Bernoulli's number contains k. E.g., 2310 is squarefree, is in the sequence and A002445(2310)=74453515937201616371390013892945833 0 is divisible by 2310.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002151 | OEIS-A069051

Primes p such that p-1 divides 2^p-2.

Conjecture: if n-1 divides 2^n-2, then (2^n-2)/(n-1) is squarefree.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 03 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002152 | OEIS-A069056

Numbers k such that Sum_{d|k} d^2*mu(d) divides k^2.

If n > 1, a(n+1) - a(n) == 0 (mod 12), so a(n+1) - a(n) = 12 for n=2,3,4,5,7,8,...; a(n+1) - a(n) = 24 for n=6,9,.... Conjecture: if c > 2 and n > 1, Sum_{d|n} d^c*mu(d) never divides n^c. Hence A063453(n) never divides n^3 for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002153 | OEIS-A069136

Numbers that are not the sum of 5 nonnegative cubes.

Sequence is conjectured to be finite. Comment from Richard C. Schroeppel, Sep 22 2010: It is conjectured that 7373170279850 is the largest number requiring more than four cubes (see Deshouillers et al.).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 08 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002154 | OEIS-A069137

Numbers which are sums of neither 1, 2, 3, 4, 5 or 6 nonnegative cubes.

Sequence is conjectured to be finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 08 2002; edited Sep 15 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002155 | OEIS-A069160

Number of primes p such that n^2 < p < n^2 + pi(n), where pi(n) is the number of primes less than n.

A more restrictive version of the conjecture that there is always a prime between n^2 and (n+1)^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 09 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002156 | OEIS-A069180

F(n) and n! are relatively prime where F(n) are the Fibonacci numbers.

Conjecture : a(n) = C*n*Log(n) + 0(n*Log(n)) with 0, 6 < C < 0, 7

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002157 | OEIS-A069187

Numbers k such that core(k) = ceiling(sqrt(k)) where core(k) is the squarefree part of k (the smallest integer such that k*core(k) is a square).

Conjecture: sequence is A071253 minus those entries of A071253 that have their index in A049532, i.e., a(n) is of form n^2*(n^2+1) for all n not in A049532. - _Ralf Stephan_, Aug 18 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 14 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002158 | OEIS-A069191

Determinant of n X n matrix defined by m(i,j)=1 if i+j is a prime, m(i,j)=0 otherwise.

Conjecture: Let M(n) be the n X n matrix with (i,j)-entry equal to 1 or 0 according as i + j is prime or not. For each n = 24, 25, ..., the characteristic polynomial of M(n) is irreducible over the field of rational numbers, and the n eigenvalues can be listed as lambda(1), ..., lambda(n) such that lambda(1) > -lambda(2) > lambda(3) > -lambda(4) > .... > (-1)^n*lambda(n) > 0. - _Zhi-Wei Sun_, Aug 25 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Santi Spadaro_, Apr 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002159 | OEIS-A069197

a(1)=1 a(2)=4 a(n+2)=(a(n+1)+a(n))/3 if (a(n+1)+a(n)==0 (mod 3)); a(n+2)=(a(n+1)+a(n))/2 if (a(n+1)+a(n)==0 (mod 2)); a(n+2)=a(n+1)+a(n) otherwise.

A Collatz-Fibonacci mixture. Does this sequence diverge to infinity? Conjecture : if a(1)=1 and a(2)=2 sequence is constant = 1, if a(2)=5 sequence is cyclic = (5,2,7,3) but if a(2)=m, different from 2 or 5, sequence diverges.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002160 | OEIS-A069202

A Collatz-Fibonacci mixture: a(1) = 1, a(2) = 2, a(n+2) = a(n+1)/2+a(n)/2 if a(n+1) and a(n) have the same parity, a(n+2) = a(n+1)+a(n) otherwise.

Conjecture: More generally, let a(1)=x and a(2)=y be two distinct positive integers; then for any x,y > 0, lim n -> infinity log(a(n))/n = 1/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002161 | OEIS-A069227

a(1)=1, a(2)=2; a(n+2) = (a(n+1) + a(n))/b(n) where b(n) = gcd(a(n+1) + a(n), 4).

Conjecture: sequence diverges to infinity (a(500)=151767071).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 12 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002162 | OEIS-A069228

a(1)=1, a(2)=4, a(n+2)=(a(n+1)+a(n))/b(n), where b(n)=gcd(a(n+1)+a(n),4).

A Collatz-Fibonacci mixture. Conjecture : sequence diverges to infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 12 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002163 | OEIS-A069269

Second level generalization of Catalan triangle (0th level is Pascal's triangle A007318; first level is Catalan triangle A009766; 3rd level is A069270).

With offset 1 for n and k, T(n,k) is (conjecturally) the number of permutations of [n] that avoid the patterns 4-2-3-1 and 4-2-5-1-3 and for which the last ascent ends at position k (k=1 if there are no ascents). For example, T(4,1) = 1 counts 4321; T(4,2) = 3 counts 1432, 2431, 3421; T(4,3) = 7 counts 1243, 1342, 2143, 2341, 3142, 3241, 4132. - _David Callan_, Jul 22 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Mar 12 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002164 | OEIS-A069429

Half the number of 3 X n binary arrays with no path of adjacent 1's or adjacent 0's from top row to bottom row.

The above conjectures are true and follow from formulas given in A069361 and A069396.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002165 | OEIS-A069482

a(n) = prime(n+1)^2 - prime(n)^2.

Conjecture: The sum of prime factors with repetition (sopfr) applied to a(n), A001414(a(n)), covers all integers covered by sopfr, except 2,3,4,6,7,10,13,15. See A001414 for the sopfr sequence, which does not cover 0 and 1. - _Richard R. Forberg_, Feb 07 2015 Conjecture: There is no upper bound on the number of repetitions (i.e., size of a tally) that will occur for some a(n) values, because the number of possible ways of producing a value of a(n) grows with increasing n, despite decreasing prime density. This happens because there is increasing range in the size of prime gaps which increases the range of primes that can produce the same a(n) value much faster than the decrease in prime density which is decelerating with larger n. - _Richard R. Forberg_, Feb 17 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Mar 29 2002, Aug 05 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002166 | OEIS-A069517

a(n) = (-1)*Sum_{d|n} (moebius(d)*(-1)^d).

Conjecture: Sum_{d | n} a(d)^2 * phi(d) = A393210(n). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 16 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002167 | OEIS-A069548

Primes of the form sum 6d/(2 + mu(d)) for some k and all d dividing k.

The function is the sum of 3d or -d (mu(d) -1 or 1, resp.) over the squarefree divisors d of n, plus 3*sigma(n). As a result, for n > 1, the squarefree part is even, and thus n generates odd numbers only when 3*sigma(n) is odd. This happens only when n is a square or twice a square, as conjectured by Alonso Del Arte. - _Charles R Greathouse IV_, Feb 18 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002168 | OEIS-A069581

Triangle T(m,n) giving number of unit fractions (with odd denominators) needed to represent m/n, rational (n odd), using the greedy algorithm.

If m/n, a rational number (n odd) is expressed as sum (1/xi), where the xi are successively chosen to be the least possible odd integers which leave a nonnegative remainder, is the sum always finite? My conjecture: odd m needs odd, even m needs even unit fractions. In the triangle: rows are the (odd) denominators, columns are 1<m<n numerators.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Adam Kertesz_, Apr 24 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002169 | OEIS-A069586

Smallest prime power p^k (k >= 2) such that there is a difference of exactly n between p^k and some other prime power q^m (m >= 2); or 0 if no such q^m exists.

The nonzero terms are sure terms, only the zero terms are conjectured. - _Michel Marcus_, Nov 02 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 24 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002170 | OEIS-A069623

Number of perfect powers <= n.

a(n) = O(sqrt(n)) (conjectured). a(n) = A076411(n+1) = Sum_{k=1..n} A075802(k). - _Chayim Lowen_, Jul 24 2015 The conjecture is true: The number of squares < n is n^(1/2) + O(1). The number of higher powers < n is nonnegative and less than n^(1/3) log_2(n). Thus a(n) = n^(1/2) + O(n^(1/3) log n). - _Robert Israel_, Jul 31 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 27 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002171 | OEIS-A069685

Denominators of coefficients in -log(1+x)log(1-x) power series.

a(n) = n*A076100(n) (conjectured). - _F. Chapoton_, Nov 09 2009 Chapoton's 2009 conjecture is false: a(8) = 576576 but 8*A076100(8) = 2882880. - _Michal Buran_, Mar 24 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 03 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002172 | OEIS-A069748

Numbers k such that k and k^3 are both palindromes.

For n >= 5, there are no terms with digit sum 5. Conjecture: all terms belong to one of 3 disjoint classes of the following forms: 10^k+1, 10^(2*t)+10^t+1, t > 0, and (10^u+1)*(10^v+1), u,v > 0, with digit sums 2, 3 and 4 correspondingly. - _Vladimir Shevelev_, May 31 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Apr 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002173 | OEIS-A069755

Frobenius number of the numerical semigroup generated by 3 consecutive triangular numbers.

Conjectures from _Colin Barker_, Nov 22 2012: (Start) Conjectures from _Colin Barker_, Mar 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Victoria A Sapko (vsapko(AT)canes.gsw.edu), Apr 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002174 | OEIS-A069761

Frobenius number of the numerical semigroup generated by four consecutive tetrahedral numbers.

Conjecture: a(n)= +a(n-1) +4*a(n-6) -4*a(n-7) -6*a(n-12) +6*a(n-13) +4*a(n-18) -4*a(n-19) -a(n-24) +a(n-25). - _R. J. Mathar_, Aug 15 2025 Conjectured g.f.: x^2*(-4*x^2 -600*x^3 -390*x^4 -1680*x^9 -282*x^8 -496*x^11 -804*x^10 -208*x -312*x^15 -144*x^14 -768*x^13 -772*x^12-41 -32*x^18 -40*x^17 -102*x^16 -2*x^20 -8*x^19 -1608*x^7 +x^24 -884*x^6 -328*x^5) / ((1+x)^4 *(x^2-x+1)^4 *(1+x+x^2)^4 *(x-1)^5). - _R. J. Mathar_, Aug 15 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | Victoria A Sapko (vsapko(AT)canes.gsw.edu), Apr 09 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002175 | OEIS-A069837

Smallest prime which is a concatenation of n primes.

Conjecture: For every n there exists an n-digit prime which is composed of the digits 2,3,5 and 7. I.e., no prime > 7 is required in this concatenation. I.e., a(n) of A069637 contains exactly n digits. This is a weaker conjecture than the one by _Patrick De Geest_ in A036937. If the conjecture is true then this also gives the smallest n-digit prime with prime digits. - _Amarnath Murthy_, Apr 02 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 16 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002176 | OEIS-A069852

a(n) = Sum_{i=0..2n} B(i)*C(2n+1,i)*5^i where B(i) are the Bernoulli numbers, C(2n,i) the binomial numbers.

Conjecturally, a(n) = the signed numerator of B(2*n + 1,1/5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 01 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002177 | OEIS-A069869

Largest prime that is a concatenation of the parts of a partition of n, or 0 if no such prime exists.

Conjecture: a(n) = 0 only for n = 1 or n = 3k with k>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002178 | OEIS-A069883

Smallest nontrivial palindromic n-th perfect power (a(n) >1 for n>1).

Conjecture: sequence is bounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 30 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002179 | OEIS-A069911

Expansion of Product_{i in A069909} 1/(1 - x^i).

Let b(k) be the convolution inverse of A035457, k=1, 2, 3, ...; then a(n) = -b(4n+3), n = 0, 1, 2, 3, ... (conjectured). - _George Beck_, Aug 19 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002180 | OEIS-A069917

In base 6, the reversal of n equals the sum of the reversals of the proper divisors of n.

A number n is called "picture-perfect" if the reversal of n equals the sum of the reversals of the proper divisors of n. These base-6 picture-perfect numbers were found by Mark Ganson while searching for (base-10) picture-perfect

numbers. He observes that the digital sum of their base-10 representations = 10 and conjectures that this is the case for all base-6 picture-perfect numbers. The only (base-10) picture-perfect numbers not exceeding 1.3 * 10^9 are 6, 10311 and 21661371.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Apr 24 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002181 | OEIS-A069929

Number of k, 1 <= k <= n, such that k^3+1 divides n^3+1.

Conjecture: (1/n)*Sum_{k=1..n} a(k) = C*log(log(n)) + o(log(log(n))) with 1 < C < 3/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002182 | OEIS-A069931

Number of k, 1<=k<=n, such that k divides sigma(n).

Asymptotically (still conjectured): sum(k=1, n, a(k)) = C*n*log(n)^2 + o(n*log(n)^2) with C=0.35...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002183 | OEIS-A069932

Number of k, 1<=k<=n, such that phi(k) divides n.

Asymptotically (still conjectured): Sum_{k=1..n} a(k) = C*n*log(n) + o(n*log(n)) with C = 1.5...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002184 | OEIS-A069941

Number of primes p such that n! <= p <= n! + n^2.

Conjecture: if n>=2 there are at least 3 primes p such that n!<=p<=n!+n^2 (or stronger: for n>1, a(n) > log(n)). This is stronger than the conjecture described in A037151(n). Because if n!+k is prime, k composite, k=A*B, where A and B must have, each one, at least one prime factor>n (if not: A=q*A' q<=n then n!+k is divisible by q), hence k>n^2.

Also stronger (but more restrictive) than the Schinzel conjecture: "for m large enough there's at least one prime p such that m <= p <= m + log(m)^2" since n^2 < log(n!)^2 for n>5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 04 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002185 | OEIS-A069942

Reversal of n equals the sum of the reversals of the proper divisors of n.

These numbers are called picture-perfect numbers (ppn's). If a ppn is placed on one side of an equal sign and its proper divisors on the other side, then the resulting equation read backwards is valid. The first three ppn's were found by _Joseph L. Pe_. The fourth ppn was discovered by Daniel Dockery. Mark Ganson conjectures that every ppn is divisible by 3. (Compare this with the still unresolved conjecture that every perfect number is divisible by 2.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Apr 26 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002186 | OEIS-A069972

a(n) = Sum_{d|2*n,d+1|2*n} d.

G.f.: (conjectured) Sum_{n>=1} n*x^(n*(n+1)/2)/(1-x^(n*(n+1)/2)). - _Joerg Arndt_, Jan 30 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Apr 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002187 | OEIS-A069994

a(n) = Sum_{i=0..2n} B(i)*C(2n+1,i)*6^i where B(i) are the Bernoulli numbers, C(2n,i) the binomial coefficients.

Conjecturally, a(n) = 2 * the unsigned numerator of B(2*n - 1,1/6). If true then this sequence is a bisection of 2*A158073.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 01 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002188 | OEIS-A070000

Let B(n) denote the number of distinct norms <=n in the square lattice and let B_3(n) denote the number of distinct norms <=n in the hexagonal lattice; sequence gives B(n) - B3(n).

It was conjectured by Schmutz Schaller that B(x)>=B_3(x) for every x; this was proved in the reference given.

猜想/问题提及段 | historical_or_current_mention_unreviewed | P. Moree (moree(AT)mpim-bonn.mpg.de), May 03 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002189 | OEIS-A070056

Number of Bottleneck-Monge matrices with 8 rows.

G.f.: x*(256 - 4609*x + 40669*x^2 - 232695*x^3 + 961183*x^4 - 3017869*x^5 + 7388387*x^6 - 14256058*x^7 + 21694227*x^8 - 25838259*x^9 + 23693517*x^10 - 16267523*x^11 + 7986763*x^12 - 2589705*x^13 + 474491*x^14 - 32768*x^15) / ((1 - x)*(1 - 2*x)^15) (conjectured). - _Colin Barker_, Sep 10 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | Pascal Prea (pascal.prea(AT)lim.univ-mrs.fr), Apr 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002190 | OEIS-A070057

Number of Bottleneck-Monge matrices with 9 rows.

G.f.: x*(512 - 9985*x + 96161*x^2 - 607903*x^3 + 2821517*x^4 - 10164757*x^5 + 29260931*x^6 - 68263237*x^7 + 129682732*x^8 - 200278047*x^9 + 249653465*x^10 - 247990203*x^11 + 192418577*x^12 - 113113161*x^13 + 48011519*x^14 - 13562349*x^15 + 2175308*x^16 - 131072*x^17) / ((1 - x)*(1 - 2*x)^17) (conjectured). - _Colin Barker_, Sep 10 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | Pascal Prea (pascal.prea(AT)lim.univ-mrs.fr), Apr 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002191 | OEIS-A070077

Greatest common divisor of n-th squarefree number and n-th cubefree number.

Conjecture: for any squarefree k exists m such that a(m)=k; see A070078.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002192 | OEIS-A070219

Smallest prime obtained as a concatenation of n and a number m greater than n.

Conjectures: There is at least one number k with n<k<2n such that the concatenation n and n+k is a prime. For all m there is at least one number k with m*n < k < (m+1)*n such that the concatenation n and n+k is a prime. As written, both conjectures are false. For n = 2, the only integer k with n < k < 2*n is k = 3, and the decimal concatenation of n and n+k is 25 = 5^2. This also refutes the second conjecture with m = 1. Updated first conjecture: the only positive integers n for which no integer k with n < k < 2*n gives a prime by concatenating n and n+k are 1, 2, 4, 5. - _Eiver Rodríguez_, Oct 07 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 17 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002193 | OEIS-A070226

Numbers k such that the squarefree kernel of k is equal to the number of divisors of k.

Conjecture : If A000005(k) divides A007947(k) for some k, then A007947(k) = A000005(k). (End) The conjecture above is true: For any k >= 1, A001222(A000005(k)) >= A001221(k) (from the multiplicative formula for A000005). If A000005(k) divides A007947(k), then A000005(k) is squarefree, so A001221(A000005(k)) = A001222(A000005(k)) >= A001221(k). However, because A000005(k) divides A007947(k), we also have A001221(A000005(k)) <= A001221(A007947(k)) = A001221(k). Therefore, A001221(A000005(k)) = A001221(A007947(k)), and A000005(k) cannot be a proper divisor of A007947(k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, May 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002194 | OEIS-A070237

Numbers k such that the sign of core(k)-phi(k) is not equal to 2*mu(k)^2-1, where core(k) is the squarefree part of k.

a(n) = C*n + O(n), with C a constant conjectured to be a(2) = 420.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 08 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002195 | OEIS-A070247

Palindromic primes with digit sum 5.

It is conjectured that are just 3 palindromic primes with digit sum 2, namely 2, 11 and 101. If any others exist, they must be of the form 10^(2^k) + 1 with k > 14.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002196 | OEIS-A070255

Numbers which divided into two substrings by breaking at any digit then one of them divides the other (0 is not to be considered a substring, multiples of 10 are excluded).

Conjecture: sequence is finite. The conjecture above is false as terms of the form d00...00d, where d goes from 1 to 9, are always in the sequence. - _FUNG Cheok Yin_, Aug 12 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002197 | OEIS-A070295

Smallest composite number such that there are exactly n primes less than it (and coprime to it) which do not divide it, or 0 if no such number exists.

Conjecture: a(223) = 0. - _Sean A. Irvine_, Jun 07 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002198 | OEIS-A070297

Smallest number with n unrelated numbers (numbers which are neither coprime nor a divisor), or 0 if no such number exists.

The 0's are only conjectures (search extended to 2000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002199 | OEIS-A070318

a(n) = Max_{k=1..n} (sigma(k)-k) where sigma(k)-k is the sum of proper divisors of k.

Conjecture: Limit_{n -> oo} (1/(n^2*log(log(n)))) * Sum_{i=1..n} a(i) = C = 0.7... . (see the second plot in the links section). - _Amiram Eldar_, Aug 04 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002200 | OEIS-A070542

Numbers n such that binomial(2n,n) does not have two prime factors that add to 2n.

No others for n < 10^8. It is conjectured that there are no other n with this property. - _T. D. Noe_, Dec 23 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, May 02 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002201 | OEIS-A070563

a(n) = 0 if 3 divides the Ramanujan number tau(n) (A000594(n)), otherwise 1.

Multiplicative because A000594 is. Conjecture: a(3^k) = 0, if p == 1 mod 3, a(p^2k) = 0 and a(p^(2k+1)) = 1, if p == -1 mod 3, a(p^2k) = 1 and a(p^(2k+1)) = 0. - _Christian G. Bower_, Jun 10 2005 The above conjecture is not correct. The first counterexample occurs at n = 2401 = 7^4. My improved conjecture is that this is actually a characteristic function of nonmultiples of 3 whose sum of divisors is also a nonmultiple of 3, that is, having a following multiplicative formula: a(3^k) = 0, if p == 1 mod 3, a(p^e) = 1 if e != 2 (mod 3), otherwise 0, and if p == -1 mod 3, a(p^2k) = 1 and a(p^(2k+1)) = 0. This conjecture has now been proved

correct by _Seiichi Manyama_. a(n) = A011655(n) * A353815(n), conjectured by _Antti Karttunen_, proved by _Seiichi Manyama_, Jul 03 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002202 | OEIS-A070788

Positive integers n such that the Reverse and Add! trajectory of n (presumably) does not join the trajectory of any m < n.

The conjecture that the trajectories of the terms of this sequence do not join is based on the observation that if the trajectories of two integers below 10000 join, this happens after at most 15 steps, while for any two terms the trajectories do not join within 1200 steps. For pairs from 1, 3, 5, 7, 9, 100, 102, 106 this has even been checked for 10000 steps.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, May 07 2002, revised Oct 15 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002203 | OEIS-A070814

Solutions to phi(gpf(x)) - gpf(phi(x)) = 14 = c are special multiples of 17, x = 17k, where greatest prime factors of factor k were observed from {2, 3, 5}, i.e., it is smaller than 17. See solutions to other even cases of c (=A070813): A007283 for 0, A070004 for 2, A070815 for 254, A070816 for 65534. Gpf = greatest prime factor.

For n > 10, a(n) = 2a(n-4) (conjectured). - _Ralf Stephan_, May 09 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, May 09 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002204 | OEIS-A070831

Palindromic primes with digit sum = 11.

Conjecture: The sequence is unbounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, May 15 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002205 | OEIS-A070864

a(1) = a(2) = 1; a(n) = 2 + a(n - a(n-1)).

Conjecture. Let a(1)=a(2)=1 and for n > 2 let k = floor(sqrt(n+1))-1 and d=n-k(k+2). Then, if (d is 0, 1, or 2) OR (d=0 mod 2), a(n)=2k+1; otherwise a(n)=2k+3. This has been verified for n <= 15000. Thus the asymptotic behavior appears to be a(n) ~ floor(sqrt(n+1)). - _John W. Layman_, May 21 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002206 | OEIS-A070875

Binary expansion is 1x100...0 where x = 0 or 1.

Conjecture: a(n) = -1 + positions of the ones in A309019(n+2) - A002487(n+2). - _George Beck_, Mar 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002207 | OEIS-A070916

a(1)=1, a(n) is the smallest integer >= a(n-1) such that a(n)*a(n-1) - 1 is prime.

Conjecture: lim_{n -> infinity} a(n)/(n*log(n)) = C = 1.075... .

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 20 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002208 | OEIS-A070931

Numbers k such that the smallest integer value >= 0 of the form x^3 - k^2 equals the smallest integer value >= 0 of the form x^2 - k^3.

Conjectures from _Colin Barker_, Jun 29 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 20 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002209 | OEIS-A070945

Number of permutations on n letters that have only cycles of length 4 or less.

Conjecturally, for k >= 1, the terms of the sequence are eventually divisible by 3^k. More precisely, we conjecture the following 3-adic values (each one checked up to n = 1000):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and Sharon Sela, May 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002210 | OEIS-A070965

a(1) = 1; a(n+1) = sum{k|n} a(k) * mu(k), where the sum is over the positive divisors, k, of n; and mu(k) is the Moebius function.

Conjecture: all integers are present. - _W. Edwin Clark_, Aug 20 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, May 16 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002211 | OEIS-A070973

Smallest integer k such that n divides floor((3/2)^k).

a(n)=n for some n = 1,2,3,16,56,283,....Conjectures : (i) Log(n) < a(n) < n*Log(n)^2 for n>6; (ii) the equation a(x)=n always has a solution.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 24 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002212 | OEIS-A071010

Sigma(k)/4 when k is not a sum of 2 squares.

Conjecture : if n is not the sum of 2 squares sigma(n) == 0 (mod 4) (converse is not true : if sigma(n) == 0 (mod 4), n is sometimes the sum of 2 squares : sigma(65) = 84 == 0 (mod 4) but 65 = 49+16 is a sum of 2 squares).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002213 | OEIS-A071011

Numbers n such that n is a sum of 2 squares (i.e., n is in A001481(k)) and sigma(n) == 0 (mod 4).

It is conjectured that if m is not a sum of 2 squares (i.e., m is in A022544(k)) sigma(m) == 0 (mod 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002214 | OEIS-A071033

a(n) = n-th state of cellular automaton generated by "Rule 94" when started with a single ON cell.

Conjecture: a(n) = floor((1099*100^n + 9090)/990) + 1 for odd n > 1; a(n) = floor((1090*100^n + 10)/990) + 1 for even n > 1. - _Karl V. Keller, Jr._, Oct 25 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hans Havermann_, May 26 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002215 | OEIS-A071052

Number of 0's in n-th row of triangle in A071036 (cellular automaton "Rule 150").

a(2n) = a(n) + 2n (conjectured). - _Ralf Stephan_, Mar 03 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hans Havermann_, May 26 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002216 | OEIS-A071055

Number of 0's in n-th row of triangle in A071038.

a(n)=b(n+1), with b(0)=0, b(2n)=b(n), b(2n+1)=2b(n)+2-2[n==0] (conjectured). - _Ralf Stephan_, Mar 05 2004 a(n) = pext(n, n + 1) (conjectured) where pext is the "parallel bits extract" instruction of the x86 CPU; pext(x, mask) extracts bits from x at the bit locations specified by mask to contiguous low bits. - _Falk Hüffner_, Jul 26 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hans Havermann_, May 26 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002217 | OEIS-A071060

Largest n-digit prime with only prime digits.

Conjecture: a(n) ~ floor((7/9) * 10^n). - _Bill McEachen_, Apr 07 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rick L. Shepherd_, May 26 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002218 | OEIS-A071066

Palindromic prime pyramid with a(1)=2 such that (number of digits in a(n+1)) = (number of digits in a(n)) + 6.

G. L. Honaker Jr. and C. Caldwell conjectured that this pyramid has a finite number of terms (about 193 terms).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 26 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002219 | OEIS-A071069

a(n) = min { A070923(k) | n^3 < k < (n+1)^3 }.

Strong conjecture : for n>12, n^2/2<a(n)<n^2. Weak conjecture : a(n)>n for any n. A simple application of the weak conjecture could be to determine if the equation x^3-y^2 = A (A integer) has no solution in integers. For example the equation x^3-y^2 = 5 would have no solution in integers since a(n)>5 for n>5 and from a direct calculus, A070923(k) is different from 5, k = 1^3 to 6^3. The strong conjecture does not hold for n = 23, 26, 28, 30, 36, 42, 46, 47, 48, 49, ... - Lambert Klasen (Lambert.Klasen(AT)gmx.net), Dec 17 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 26 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002220 | OEIS-A071071

Minimal "powers of 2" set in base 10: any power of 2 contains at least one term of this sequence in its decimal expansion.

Conjectured by J. Shallit to be complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 26 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002221 | OEIS-A071089

Remainder when sum of first n primes is divided by n-th prime.

Conjecture: Every nonnegative integer can appear in the sequence at most finitely many times. - _Thomas Ordowski_, Jul 22 2013 I conjecture the opposite. Heuristically a given number should appear log log x times below x. - _Charles R Greathouse IV_, Jul 22 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Randy L. Ekl_, May 26 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002222 | OEIS-A071111

a(n) is the least integer x such that there exists an integer in the open interval (x/(i+1), x/i) for i= n-1, n-2 ..., 3, 2, 1.

"This formula was found by Florian Herzig (then a student at Cambridge, UK), who also proved that the proposer's conjecture that a(n) = cases (1 + (n-m)^2, if m^2 <= n - 2, 1 + (n-m)^2 + (n - m), otherwise) where m = floor ((1 + sqrt (4 n - 7)) / 2) also is true although 'the proof of this fact is quite challenging'.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Fernando Delgado, Paul Monasterios, and Adolfo Rodriguez (misterioso53(AT)hotmail.com), May 27 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002223 | OEIS-A071234

Smallest prime beginning and ending in 2n+1 or 0 if no such prime exists.

Conjecture: a(n) = 0 if and only if n > 2 and n == 2 mod 5 (which makes 2*n+1 == 0 mod 5). - _Robert Israel_, May 26 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002224 | OEIS-A071241

Arithmetic mean of k and R(k) where k is the n-th nonnegative number using only even digits and R(k) is its digit reversal (A004086).

Conjecture: 101 is the largest prime term, the only other primes being 2 and 11. The conjecture is false: for example, 181 and 383 are prime terms. There are 150 prime terms less than 75000. - _Harvey P. Dale_, Sep 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 20 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002225 | OEIS-A071256

Smallest multiple of n sandwiched between twin primes.

Conjecture: lim sup n ->infinity a(n)/n^2 exists = C, where 0<C<1. - _Benoit Cloitre_, May 23 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002226 | OEIS-A071264

Expansion of (1+x*C^4)*C^3, where C = (1-(1-4*x)^(1/2))/(2*x) is g.f. for Catalan numbers, A000108.

Conjecture: D-finite with recurrence (n+6) *(8587*n^2 +62958*n +61667) *a(n) +(8587*n^3 -377960*n^2 -2268967*n -1091596) *a(n-1) -2 *(2*n-1) *(42935*n^2 -14497*n -123334) *a(n-2)=0. - _R. J. Mathar_, Dec 10 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002227 | OEIS-A071272

Palindromes in A066492.

Conjecture: m is a term if and only if 1<=m<=9 or m is a palindrome of odd length 2k+1 such that the sum of the k-th and (k+1)-th decimal digits is less than or equal to 9. - _Chai Wah Wu_, May 30 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jun 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002228 | OEIS-A071281

Numerators of Peirce sequence of order 3.

Conjectures from _Colin Barker_, Mar 29 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002229 | OEIS-A071282

Denominators of Peirce sequence of order 3.

Conjectures from _Colin Barker_, Mar 29 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002230 | OEIS-A071283

Numerators of Peirce sequence of order 4.

Conjectures from _Colin Barker_, Mar 29 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002231 | OEIS-A071284

Denominators of Peirce sequence of order 4.

Conjectures from _Colin Barker_, Mar 29 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002232 | OEIS-A071285

Numerators of Peirce sequence of order 5.

Conjectures from _Colin Barker_, Mar 29 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002233 | OEIS-A071286

Denominators of Peirce sequence of order 5.

Conjectures from _Colin Barker_, Mar 29 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002234 | OEIS-A071287

Numerators of Peirce sequence of order 6.

Conjectures from _Colin Barker_, Mar 29 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002235 | OEIS-A071288

Denominators of Peirce sequence of order 6.

Conjectures from _Colin Barker_, Mar 29 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002236 | OEIS-A071302

a(n) = (1/2) * (number of n X n 0..2 matrices M with MM' mod 3 = I, where M' is the transpose of M and I is the n X n identity matrix).

a(n+1) = a(n) * A318609(n+1) for n >= 1. - conjectured by _Petros Hadjicostas_, Dec 18 2019; proved based on the explicit formula by _Max Alekseyev_, Nov 06 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002237 | OEIS-A071304

a(n) = (1/2) * (number of n X n 0..4 matrices M with MM' mod 5 = I, where M' is the transpose of M and I is the n X n identity matrix).

We conjecture that a(n+1) = a(n)*b(n+1) for n >= 1. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002238 | OEIS-A071306

a(n) = (1/2) * (number of n X n 0..6 matrices M with MM' mod 7 = I, where M' is the transpose of M and I is the n X n identity matrix).

Conjecture: Let b(n) be the number of solutions to the equation Sum_{i = 1..n} x_i^2 = 1 (mod 7) with x_i in 0..6. We conjecture that b(n) = 7*b(n-1) - 7*b(n-2) + 49*b(n-3)

for n >= 4 with b(1) = 2, b(2) = 8, and b(3) = 42. We also conjecture that a(n+1) = a(n)*b(n+1) for n >= 1. - _Petros Hadjicostas_, Dec 19 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002239 | OEIS-A071307

1/2 number of n X n 0..8 matrices with MM' mod 9 = I.

Conjecture: a(n) = 3^(n*(n-1)/2) * A071302(n) for n >= 1. - _Michel Marcus_, Nov 08 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002240 | OEIS-A071356

Expansion of (1 - 2*x - sqrt(1 - 4*x - 4*x^2))/(4*x^2).

Conjecturally, the number of sequences (e(1), ..., e(n+1)), 0 <= e(i) < i, such that there is no triple i < j < k with e(i) > e(j) <= e(k). [Martinez and Savage, 2.19] - _Eric M. Schmidt_, Jul 17 2017 Let s denote West's stack-sorting map, and let Av_n(tau_1, ..., tau_r) denote the set of permutations of [n] that avoid the patterns tau_1, ..., tau_r. It is conjectured that a(n) = |s^{-1}(Av_{n+1}(132, 231))| = |s^{-1}(Av_{n+1}(132, 312))| = |s^{-1}(Av_{n+1}(231, 312))|. Only the last of these equalities is known. - _Colin Defant_, Sep 16 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 12 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002241 | OEIS-A071383

Squared radii of the circles around (0,0) that contain record numbers of lattice points.

lim n ->infinity Log(a(n))/n = 1. [Conjectured by _Benoit Cloitre_, proved by _J. H. Conway_]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, May 23 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002242 | OEIS-A071387

Smallest number k for which the set of solutions to phi(x) = k has 2n-1 entries.

a(n) = Min({x; Card(InvPhi(x)) = 2n-1}); a(1)=0 because of Carmichael conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, May 23 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002243 | OEIS-A071536

Number of extensions to a semigroup of the (categorical) composition of arrows in the complete directed graph on n labeled nodes.

Terms obtained, especially the 147, lend some support to the peculiar conjecture that the composition in any category can be extended to a total, associative operation, that is to a semigroup (not, of course, a monoid).

猜想/问题提及段 | historical_or_current_mention_unreviewed | F. Lockwood Morris (lockwood(AT)ecs.syr.edu), May 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002244 | OEIS-A071537

Least integer m >= n such that there exist integers k_n, k_n+1, ..., k_m with nonzero k_n, for which n^(k_n) * (n+1)^(k_n+1) * ... * m^(k_m) = 1.

Conjecture: this is a permutation of the nonprimes A018252. The conjecture is true. It is easy to see that (i) a(n)=m cannot be prime; (ii) k_m is always nonzero; (iii) for any nonprime s, there exist integers k_p,...,k_s, where p is the smallest prime divisor of s, k_p and k_s are nonzero, for which p^(k_p)*...*s^(k_s) = 1. The bijectivity of a(n) (as a function between positive integers and nonprimes) now follows from the lemma.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Fink_, May 30 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002245 | OEIS-A071558

Smallest k such that n*k + 1 and n*k - 1 are twin primes.

Conjecture: a(n) < sqrt(n)*log(n) for all n > 17261. This has been verified for n up to 3*10^7. It implies the inequality a(n) < n for each n > 127. - _Zhi-Wei Sun_, Jan 07 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 30 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002246 | OEIS-A071561

Numbers with no middle divisors (cf. A071090).

Conjecture: lim_{n->oo} a(n)/n = 4/3. Regarding the above conjecture, numerical calculations suggest that this limit is smaller than 4/3. See A071540. - _Amiram Eldar_, Jul 27 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, May 30 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002247 | OEIS-A071621

Primes that can be written as "a * b + c * d", where a, b, c and d are also primes.

As size in the Mathematica coding is increased, the primes not previously covered will probably be forthcoming. Conjecture: Only the primes 2, 3, 5, 7, 11 and 17 are not representable by this form.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Arnoud Buzing (arnoudb(AT)wolfram.com), Jun 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002248 | OEIS-A071622

a(n) = (-1)*Sum_{k=1..n} (-1)^floor((4/3)^k).

Conjecture: for n>2300, a(n) > log(n)^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jun 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002249 | OEIS-A071642

Numbers n such that x^n + x^(n-1) + x^(n-2) + ... + x + 1 is irreducible over GF(2).

Conjecture: Terms >= 2 are numbers n such that P^n + P^(n-1) + P^(n-2) + ... + P + 1 is irreducible over GF(2), where P=x^2+x+1. - _Luis H. Gallardo_, Dec 23 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002250 | OEIS-A071644

a(n) = A005148(2^n-1)/8^(n-1).

Appears to always be an integer. General conjecture: the numbers k such that 8^a is the highest power of 2 dividing A005148(k) is the same sequence as numbers k such that k has exactly (a+1) 1's in its binary representation. Hence this sequence gives the smallest integer of the form A005148(k) /8^(n-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jun 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002251 | OEIS-A071684

Number of plane trees with n edges and having an odd number of leaves.

Conjecture: n*(n+1)*(10*n-37)*a(n) + 2*n*(5*n^2-42*n+91)*a(n-1) + 4*(-40*n^3+270*n^2-560*n+357)*a(n-2) + 8*(n-3)*(5*n^2-42*n+91)*a(n-3) - 16*(n-4)*(25*n-51)*(2*n-7)*a(n-4) = 0. - _R. J. Mathar_, Jul 05 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sen-peng Eu_, Jun 23 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002252 | OEIS-A071713

a(n) = smallest value of |x^n + y^n + z^n| for x,y,z distinct nonzero integers.

The conjectured minimum of 12 for a(5) can be achieved by (x,y,z) = (13,16,-17). - _Stefan Steinerberger_, Dec 13 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jun 03 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002253 | OEIS-A071715

Expansion of (1+x*C^4)*C^4, where C = (1-(1-4*x)^(1/2))/(2*x) is g.f. for Catalan numbers, A000108.

Conjecture: (891*n+6824)*(n+7)*a(n) + (891*n^2-38453*n-250970)*a(n-1) + 4*(-5040*n^2-4271*n+8095)*a(n-2) + 4*(1170*n+3847)*(2*n-1)*a(n-3) = 0. - _R. J. Mathar_, Aug 25 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002254 | OEIS-A071717

Expansion of (1 + x^2*C)*C^2, where C = (1 - sqrt(1-4*x))/(2*x) is g.f. for Catalan numbers, A000108.

Conjecture: (n+2)*a(n) +(-3*n-2)*a(n-1) +(-5*n+8)*a(n-2) +2*(2*n-7)*a(n-3)=0. - _R. J. Mathar_, Aug 25 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002255 | OEIS-A071719

Expansion of (1+x^2*C)*C^4, where C = (1-(1-4*x)^(1/2))/(2*x) is g.f. for Catalan numbers, A000108.

Conjecture: -3*(n+4)*(3*n-1)*a(n) +5*(n+2)*(7*n+3)*a(n-1) +2*(2*n-3)*(n-7)*a(n-2)=0. - _R. J. Mathar_, Aug 25 2013 Conjecture confirmed using the differential equation (4*x^4+35*x^3-9*x^2)*g'' + (-14*x^3+190*x^2-42*x)*g' + (-10*x^2+150*x+12)*g = 30*x+12 satisfied by the G.f. - _Robert Israel_, Jun 07 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002256 | OEIS-A071722

Expansion of (1+x^2*C^2)*C^3, where C = (1-(1-4*x)^(1/2))/(2*x) is g.f. for Catalan numbers, A000108.

Conjecture: 2*(n+3)*a(n) + 4*(-3*n-4)*a(n-1) + (17*n-9)*a(n-2) + 2*(-2*n+5)*a(n-3) = 0. - _R. J. Mathar_, Aug 25 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002257 | OEIS-A071823

Number of numbers x <= n with largest prime factor of the form 4k+3.

Conjecture : for n > 1000, a(n) - n/2 > sqrt(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jun 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002258 | OEIS-A071824

Number of integers <= n whose largest prime factor is of the form 4*k+1.

Conjecture : for n > 1000 a(n) - n/2 < - sqrt(n) (if b(n) denotes the number of x with largest prime factor of the form 4k+3 less than or equal to n, it is conjectured that if n > 1000 b(n)- n/2 > sqrt(n))

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jun 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002259 | OEIS-A071825

a(n) = (number of x with largest prime factor of the form 4k+3 less than or equal to n) minus (number of x with largest prime factor of the form 4k+1 less than or equal to n).

Weak conjecture: sequence gives positive values only; strong conjecture: for n large enough a(n) > 2*sqrt(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jun 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002260 | OEIS-A071868

Number of integers k (1 <= k <= n) such that k^2+1 is prime.

Hardy and Littlewood conjectured that : a(n) ~ c* sqrt(n)/Log(n) where c = Product_{p prime} (1 - (-1)^((p-1)/2)/(p-1)) = 1.3728... (A199401).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jun 09 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002261 | OEIS-A071880

Number of combinatorial types of n-dimensional parallelohedra.

The papers by Dutuor Sikiric, Garber et al say that actually a(5) = 110244. The claim that every parallelotope is a Voronoi cell of some lattice in R^n up to an affine transformation is a conjecture open for n > 5. - _Andrey Zabolotskiy_, Feb 20 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002262 | OEIS-A071887

Conjectured values of n whose trajectory under map x -> A002487(x)*A002487(x+1) does not go into a cycle.

Conjectured values of n whose trajectory under map x -> A002487(x)*A002487(x+1) does not go into a cycle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. H. Conway_, Jun 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002263 | OEIS-A071900

1/4 times the number of n X n 0..7 matrices with MM' mod 8 = I, where M' is the transpose of M and I is the n X n identity matrix.

Conjecture: a(n) = 2^(n*(n-1)/2) * A071303(n) for n >= 1. - _Michel Marcus_, Nov 08 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 12 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002264 | OEIS-A072031

Row sums of A072030.

This conjecture is equivalent to the following identity for the divisor function: for n >= 2 there holds sum {k = 1..n} tau(k)*tau(n-k) = sum {d divides n} (sum {k = 1..d} tau(k*(d - k))), where we take tau(0) = 0. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Jun 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002265 | OEIS-A072041

a(n) is the smallest number of the form k + reverse(k) for exactly n integers k, or -1 if no such number exists.

The negative terms are conjectural. Moreover they have a rather low degree of confirmation, since due to the time-consuming computations only numbers from 0 to 65000 have been taken into account. Random tests of larger numbers however seem to indicate, that only selected values of n occur. - In the cognate sequence A071266 two numbers a and b are counted only once, if n = a + b, a = reverse(b), b = reverse(a). Then 33 = 12 + 21 = 30 + 03 has a count of 2 and 44 = 13 + 31 = 22 + 22 = 40 + 04 has a count of 3, so 44 appears in A071266 instead of 33.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Jun 08 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002266 | OEIS-A072061

[t], 1+[t], [2t], 2+[2t], [3t], 3+[3t], ..., where t=tau = (1+sqrt(5))/2 and []=floor.

Conjecture: For even n, the ratio a(n)/a(n-1) is asymptotic to (1 + sqrt(5))/2 as n becomes large. (At n=3000, the ratio is 1.61804697, compared to the exact value 1.61803399.) - _John W. Layman_, Jul 08 2004 A more general conjecture may be stated as follows: Define {a(n)} by a(1)=1 and, for n>1, a(n) = a(n-1)+floor(kn) if n is even, else a(n)=smallest positive integer which has not yet appeared in the sequence, where k is a positive real number. Then a(2n)/a(2n-1) is asymptotic to k+sqrt(k^2+1) for large n. - _John W. Layman_, Jul 08 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2002, Aug 17 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002267 | OEIS-A072100

Column 2 of the array m(i,1)=m(1,j)=1 m(i,j)=m(i-1,j-1)+m(i-1,j+1) (a(n)=m(n,2)).

Conjecture: (-n+1)*a(n) +(n+1)*a(n-1) +2*(2*n-7)*a(n-2) +4*(-n+3)*a(n-3)=0. - _R. J. Mathar_, Nov 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jul 30 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002268 | OEIS-A072105

Let c(1)=x, c(n+1) = c(n)/2 + n if c(n) is even, c(n+1)= 2c(n) - n otherwise; then a(n)=c(n) for c(1)=3.

It seems that for any n, 2n <= a(n) <16n. If x=0,1,2,4 or 6 we have c(k+1)-c(k)=2 for k large enough and then lim k -> oo c(k)/k=2. For x=3,5 and for any x >6 there is a conjectured constant 4 < C < 5 such that lim N -> oo (1/N)*sum(k=1,N,c(k)/k) = C. Hence lim N -> oo (1/N)*sum(k=1,N,a(k)/k) should be C=4.6...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jul 30 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002269 | OEIS-A072124

a(n)-th factorial is the smallest factorial containing exactly n 1's, or 0 if no such number exists.

By checking the factorials of all the numbers below 10^6, it is conjectured that up to 10^4 there are 746 values of n for which a(n) = 0: n = 84, 164, 167, 169, 182, ... (see the link for more values). - _Amiram Eldar_, Sep 01 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shyam Sunder Gupta_, Jul 30 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002270 | OEIS-A072134

Second generation sequence in which each number is skipped that can be written as sum of distinct previous entries. To make the first generation we start with all natural numbers: this gives the powers of 2 (A000079). For the second generation we start with the natural numbers from which are removed the numbers of the first generation.

Conjectures from _Colin Barker_, Sep 09 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Floor van Lamoen_, Jun 26 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002271 | OEIS-A072141

Numbers n such that two applications of 'Reverse and Subtract' lead to n, whereas one application does not lead to n.

I conjecture that for some k there are no solutions, while in other cases the minimal solutions will have 20, 24, 28, ... digits (which however are very hard to find).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Jun 24 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002272 | OEIS-A072154

Coordination sequence for the planar net 4.6.12.

Theorem: For n >= 7, a(n) = a(n-1) + a(n-5) - a(n-6), and a(5k) = 12k (k > 0), a(5k+m) = 12k + 2m + 1 (k >= 0, 1 <= m < 5). This also implies the conjectured g.f.'s. - _N. J. A. Sloane_, conjectured Dec 20 2015, proved Jan 20 2018.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 28 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002273 | OEIS-A072190

Indices of primes with primitive root 2.

Artin conjectured that this sequence is infinite (this is the famous Artin Conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Miklos Kristof_, Jul 02 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002274 | OEIS-A072203

(Number of oddly factored numbers <= n) - (number of evenly factored numbers <= n).

Polya conjectured that a(n) >= 0 for all n, but this was disproved by Haselgrove. Lehman gave the first explicit counterexample, a(906180359) = -1; the first counterexample is at 906150257 (Tanaka).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Bill Dubuque (wgd(AT)zurich.ai.mit.edu), Jul 03 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002275 | OEIS-A072216

Consider the Reverse and Add! problem (cf. A001127); of all the n-digit numbers N which eventually reach a palindrome, pick that number N which takes the greatest number of steps to converge (in case of a tie, pick the smallest N); sequence gives values of N.

Since we do not even know if 196 eventually converges (see A006960, A023108) for n >= 3 these values are only conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002276 | OEIS-A072234

Numbers k such that reverse(k) = sum of the proper divisors of k.

Mark Ganson conjectures that all terms are divisible by 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Jul 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002277 | OEIS-A072326

Conjectured values of n such that the sum of proper divisors of x*n is never prime for any x >= 1 (i.e., sigma(xn) - xn is never prime for x >= 1).

Conjectured values of n such that the sum of proper divisors of x*n is never prime for any x >= 1 (i.e., sigma(xn) - xn is never prime for x >= 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jul 31 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002278 | OEIS-A072339

Any number n can be written (in two ways, one with m even and one with m odd) in the form n = 2^k_1 - 2^k_2 + 2^k_3 - ... + 2^k_m where the signs alternate and k_1 > k_2 > k_3 > ... >k_m >= 0; sequence gives minimal value of m.

Conjecture: a(n)=1 if n=2^k, a(n)=a(2^k-i)+1 if 2^k<n+i<2^(k+1). - _John W. Layman_, Jul 18 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jul 15 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002279 | OEIS-A072340

Number of steps to reach an integer starting with n/3 and iterating the map x -> x*ceiling(x), or -1 if no integer is ever reached.

We conjecture that an integer is always reached.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and J. C. Lagarias, Sep 03 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002280 | OEIS-A072341

a(n) = the least natural number k such that k*sigma(n) + 1 is prime.

Conjecture: a(n) is less than or equal to n for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Jul 16 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002281 | OEIS-A072342

a(n) = the least natural number k such that k*reverse(n) + 1 is prime.

Conjecture: a(n) is less than or equal to n for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Jul 16 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002282 | OEIS-A072348

Number of primes whose decimal representation has the form rns where r and s are single digits.

a(n) <= 16 for all n, a(5) = 16, conjecture: a(n)<16 for n>5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jul 17 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002283 | OEIS-A072447

Number of connectedness systems on n vertices that contain all singletons and the set of all the vertices.

Conjecture concerning the original definition: a(n) is also the number of families of subsets of {1, ..., n} that contain both the universe and the empty set, are closed under intersection and contain no sets of cardinality n-1. - _Tian

Vlasic_, Nov 04 2022. [This was false, as pointed out by _Christian Sievers_, Oct 20 2023. It is easy to see that for n>1, a(n) is also the number of families of subsets of {1, ..., n} that contain both the universe and the empty set, are closed under union of nondisjoint sets, and contain no singletons; whereas by duality, the sequence suggested in the conjecture is also the number of those families that are also closed under arbitrary union. For details see the Sievers link. - _N. J. A. Sloane_, Oct 21 2023]

猜想/问题提及段 | historical_or_current_mention_unreviewed | Wim van Dam (vandam(AT)cs.berkeley.edu), Jun 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002284 | OEIS-A072491

Define f(1) = 0. For n>=2, let f(n) = n - p where p is the largest prime <= n. a(n) = number of iterations of f to reach 0, starting from n.

On Cramér's conjecture, a(n) = O(log* n). - _Charles R Greathouse IV_, Feb 04 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 14 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002285 | OEIS-A072493

a(1) = 1 and a(n) = ceiling((Sum_{k=1..n-1} a(k))/3) for n >= 2.

Is this sequence, with its first 8 terms removed, the same as A005427? See also the similar conjecture with A005428/A073941. - _Ralf Stephan_, Apr 04 2003 Conjecture 1: a(n) equals the number of multiples of 3 whose representation in base 4/3 (see A024631) has n-1 digits. For example, a(8) = 4 because there are four multiples of 3 with n-1 = 7 digits in their representation in base 4/3: 33 = 3210201, 36 = 3210230, 39 = 3210233, and 42 = 3213122. Conjecture 2: a(n) equals 1/4 times the number of nonnegative integers with the property that their 4/3-expansion has n digits (assuming that the 4/3-expansion of 0 has 1 digit). For example, a(7)*4 = 12 because the following 12 numbers have 4/3 expansions with n = 7 digits: 32 = 3210200, 33 = 3210201, 34 = 3210202, ..., 42 = 3213122, 43 = 3213123. (End) Both conjectures are correct. We first notice that the maximum number with at most n digits in its base-4/3 expansion is 4*A087192(n) - 1, and so the number of multiples of 3 having n digits is floor((4*A087192(n) - 1)/3) - floor((4*A087192(n-1) - 1)/3) = (ceiling(4*A087192(n)/3) - 1) - (ceiling(4*A087192(n-1)/3) - 1) = A087192(n+1) - A087192(n). Let f_n = A087192(n+1), with alpha = 4/3, a = 0, and b = 2/3, then we see that (A390321/4)*(4/3)^n - 1/2 < A087192(n+1) - A087192(n) < (A390321/4)*(4/3)^n + 1/2, since we have neither e_n = -1/3 for all sufficiently large n nor e_n = 1/3 for all sufficiently large n. We conclude that a(n) = A087192(n) - A087192(n-1) for n > 1 by the first formula below, confirming both conjectures. (End) Conjecture: The constant c above equals (3/16)*K(4), where K(q) = C(q/(q-1)) (q > 1) is described in Odlyzko and Wilf (1991). The conjecture above is correct. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002286 | OEIS-A072523

Sum of remainders when n-th Fibonacci number is divided by all smaller Fibonacci numbers > 1.

Conjecture: lim n->inf F(n)/a(n) = sqrt(5)/2 where F(n) is the n-th Fibonacci number and therefore lim n->inf a(n)/a(n-1) = Phi (i.e. (sqrt(5)+1)/2 or lim n->inf F(n)/F(n-1)) - _Gerald McGarvey_, Jul 14 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 31 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002287 | OEIS-A072532

a(0)=1; for n>0, a(n) = smallest prime of the form k*a(n-1)-1 with k>1.

Conjecture: If a(n) = k*a(n-1)-1 then k < a(n-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 02 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002288 | OEIS-A072540

Numbers that are not differences between successive Ulam numbers (A002858).

Unfortunately, it appears that the entries are all conjectural. I am not aware of any proofs that the listed numbers can never appear. The terms shown are those listed by Pickover. - _N. J. A. Sloane_, Jun 19 2008 The sequence is conjectured to continue: 23, 26, 28, 31, 33. These gaps do not appear in the first 158000000 terms, so are candidates for the sequence. [_Jud McCranie_, Sep 12 2013] The conjectured list of gaps that don't appear holds through the first 28 billion Ulam numbers. - _Jud McCranie_, Jan 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 04 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002289 | OEIS-A072564

Numbers m such that the absolute values of the real and imaginary part of zeta(1/2 + m*i) are both < 1.

Conjecture: lim_{n->infinity} a(n)/n = C exists, with 5 < C < 6. [The conjecture was based on erroneous terms; C is about 2.05 (see graph). - _Vaclav Kotesovec_, Feb 18 2021]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002290 | OEIS-A072567

A variant of Zarankiewicz problem: maximal number of 1s in n X n 01-matrix with no four 1s forming a rectangle.

Conjecture: the same number of 1s is achieved for symmetric n X n matrices (cf. A350189). - _Max Alekseyev_, Apr 03 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | Xuli Le (leshlie(AT)eyou.com), Jun 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002291 | OEIS-A072592

Even numbers with at least one prime factor of form 4*k+1.

Conjecture: this is exactly the sequence whose terms are twice those of A009003. (This has been verified for all terms<=500.) Compare A009003. - _John W. Layman_, Mar 12 2008 The conjecture is true. See comments on A008846 and A004613. - Lambert Herrgesell (zero815(AT)googlemail.com), Apr 24 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jun 23 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002292 | OEIS-A072611

Numbers k such that phi(k) divides sigma(k+1) - sigma(k).

Conjecture : Limit_{n -> infinity} log(a(n))/log(n) exists and = 2.6... .

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002293 | OEIS-A072612

Minimal value of { abs(n-sigma(k)) : k>0 }.

Conjecture : S(n) = sum(k=1, n, a(k)) = n + O(sqrt(n)) and more precisely for n large enough : n - (5/2)*sqrt(n) < S(n) < n - (3/2)*sqrt(n)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002294 | OEIS-A072768

The RASTxx transformation of the sequence A072643.

Differs from A071673 first time at the position n=37, where A072768(37) = 4, while A071673(37) = 5. RASTxx(A072768) differs from A071673 first time at the position n=704, which leads to conjecture that the repeated applications of RASTxx starting from A072643 converge towards A071673, the fixed point of RASTxx transformation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jun 12 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002295 | OEIS-A072808

Smallest m such that sigma(m) mod phi(m) = n or 0 if no solution exists.

Warning: It is only conjectured that there are no solutions for n such that a(n) = 0. The search for solutions tested all m <= 10^10 for these n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jul 12 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002296 | OEIS-A072834

Exponents occurring in expansion of F_8(q^2).

Conjecture: a(n) = 4*n + pa(n mod 4), where pa(k) = 0,3,4,3 for k=0,1,2,3 respectively; lim_{n->infinity} a(n)/n = 4; a(n)/n >= 4; a(n+4) = a(n) + 16. - _Jerzy R Borysowicz_, Jan 16 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 25 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002297 | OEIS-A072842

Largest m such that we can partition the set {1,2,...,m} into n subsets with the property that we never have a+b=c for any distinct elements a, b, c in one subset.

The following coloring demonstrates that a(5) >= 207, confirming this number remains an open problem:

猜想/问题提及段 | historical_or_current_mention_unreviewed | Tor G. J. Myklebust (pi(AT)flyingteapot.bnr.usu.edu), Jul 24 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002298 | OEIS-A072849

Prime(a(n)) + ... + prime(a(n)+3) is a square = A051395(n)^2.

Conjecture: this sequence and A064397 are disjoint. That is to say, prime(n) + prime(n+1) and prime(n) + prime(n+1) + prime(n+2) + prime(n+3) cannot be squares at the same time. - _Jianing Song_, Nov 13 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Jun 21 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002299 | OEIS-A072888

Sum of the coefficients of the Schur function expansion of the square of the Vandermonde determinant in n variables.

I conjecture that a(n) = prod_{x=0..floor(n/2)} (-3x+1) * prod_{x=0..floor((n-1)/2)} (6x+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Brian G Wybourne (bgw(AT)phys.uni.torun.pl), Jul 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002300 | OEIS-A072893

Let c(k) be defined as follows: c(1)=1, c(2)=x, c(k+2) = c(k+1)/2 + c(k)/2 if c(k+1) and c(k) have the same parity; c(k+2) = c(k+1) - c(k) otherwise. Sequence gives values of x such that c(k)=1 for any k large enough.

Conjectures: (1) Lim n ->infinity a(n)/n = C = 3.5.... (2) For any n > 2, a(n+1)-a(n) = 2, 3, 5, 7, 8 or 10 only. First differences a(n+1)-a(n) for n>2 are 3, 2, 3, 2, 5, 3, 2, 5, 3, 2, 10, 3, 5, 2, 3, 2, 8... (3) If x is not in this sequence, for k large enough, c(k)= -1 or c(k) reaches one of the two cycles {3, 1, 2, -1} or {0, -1, 1, 0, 1, -1}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jul 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002301 | OEIS-A072894

Let c(k) be defined as follows: c(1)=1, c(2)=n, c(k+2) = c(k+1)/2 + c(k)/2 if c(k+1) and c(k) have the same parity; c(k+2) = c(k+1)/2 + c(k)/2 + 1/2 otherwise; a(n) = limit_{ k -> infinity} c(k).

Conjectures : (1) a(n+1)-a(n) = 0 or 1; (2) lim n ->infinity a(n)/n = 2/3; (3) 1/2 < (3a(n)-2n)/Log(n) <3/2 for any n > 1000. Does lim n -> infinity (3a(n)-2n)/Log(n) = 1 ? Conjectures (1) and (2) are true while (3) is false and the answer to the question is "no". See Fried link. - _Sela Fried_, Mar 03 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jul 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002302 | OEIS-A072901

Composite numbers n such that the discriminant of the quadratic field Q(sqrt(n)) equals 4n.

Conjecture: All terms are squarefree. Example: 6=2*3; 15=3*5; 30=2*3*5; 154=2*7*11; 195=3*5*13. - _Vincenzo Librandi_, Aug 08 2010 and _Michel Marcus_, Nov 26 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002303 | OEIS-A072907

Decimal expansion of the solution of equation log(2) - X/8 - exp(-3X/7) = 0 (4.251844....).

This constant was conjectured to be the exact value for the 3-clause threshold in the satisfiability problem (Olivier Dubois 1993).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002304 | OEIS-A072908

Decimal expansion of the solution of equation log(2)-X*2^(-r)-exp(-X*r/(2^r-1)) = 0 for r = 4 . Solution is 9.96955802...

This constant was conjectured to be the exact value for the 4-clause threshold in satisfiability Problem (Olivier Dubois 1993).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002305 | OEIS-A072924

Least k such that floor((1+1/k)^n) is prime.

On the contrary, A.L. Whiteman conjectured that the sequence floor(r^n) for non-integer rational r > 1 always contains infinitely many primes. If this conjecture is true for some r=1+1/k, then lim inf_{n -> infinity} a(n) is finite. - _Robert Israel_, Jan 09 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002306 | OEIS-A072926

a(n) = Sum_{k=1..n} A051699(k).

Conjecture: a(n) is asymptotic to C*n*log(n) with C = 0.29... .

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002307 | OEIS-A072933

Numbers m such that the least k such that floor((1+1/k)^m) = floor((1+1/m)^k) is not equal to round(m*91/100).

Conjecture: lim n -> infinity a(n)/n = 10

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 13 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002308 | OEIS-A072939

Define a sequence c depending on n as follows: c(1)=1 and c(2)=n; c(k+2) = (c(k+1) + c(k))/2 if c(k+1) and c(k) have the same parity; otherwise c(k+2) = abs(c(k+1) - 2*c(k)); sequence gives values of n such that lim_{k->oo} c(k) = infinity.

Conjecture: a(n) = A036554(n)+1. - _Vladeta Jovovic_, Apr 01 2003 Conjecture: this sequence gives the positions of 0's in the limiting 0-word of the morphism 0->11, 1->10, A285384. - _Clark Kimberling_, Apr 26 2017 Conjecture: This also gives the positions of the 1's in A328979. - _N. J. A. Sloane_, Nov 05 2019 Conjecture: lim_{n->oo} a(n)/n = 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 12 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002309 | OEIS-A072949

Number of permutations p of {1,2,3,...,n} such that Sum_{k=1..n} abs(k-p(k)) = 2n.

a(n) is always even for n>=1. More generally, A062869(n,k) is even whenever k >= n. - Conjectured by _Franklin T. Adams-Watters_, proved by _Max Alekseyev_. (see link in A062869)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 20 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002310 | OEIS-A072989

Numbers m>0 such that the number of solutions to x^m==1 (mod m), 1<=x<=m, is not equal to gcd(m, phi(m)).

Conjecture: limit of a(n)/n is zero. This conjecture is certainly wrong as stated, because sequences "Numbers such that..." have lim a(n)/n >= 1 and a(n) > n for all indices following the first one for which this holds, as here: a(1) > 1. - _M. F. Hasler_, Feb 24 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002311 | OEIS-A072995

Least k > 0 such that the number of solutions to x^k == 1 (mod k) 1 <= x <= k is equal to n, or 0 if no such k exists.

Conjecture: a(n)=0 iff n/2 is in A061346. - _Robert G. Wilson v_, Feb 23 2014 [n=420 seems to be a counterexample to the above conjecture. - _M. F. Hasler_, Feb 24 2014]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002312 | OEIS-A073008

Decimal expansion of the Traveling Salesman constant.

Conjectured to be equal to (4/153)*(1+2*sqrt(2))*sqrt(51).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Aug 03 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002313 | OEIS-A073117

a(n+1) = a(n) + a(n) mod n; a(1) = 1.

Conjecture (seems provable): More generally let a and b(1) be integers. If b(n+1) = b(n) + b(n) (mod(n+a)) there is an integer x(a,b(1)) such that b(n+1) = b(n) + x(a,b(1)) for n sufficiently large. We have x(0,1) = x(1,1) = x(2,1) = 97, x(3,1) = 1, x(4,1) = 3, x(5,1) = 3, x(6,1) = 6, ..., x(97,1) = 43, x(0,11) = 2, etc. - _Benoit Cloitre_, Aug 20 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Aug 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002314 | OEIS-A073131

a(n) = prime(prime(n+1)) - prime(prime(n)).

The first differences of A006450. Conjecture: a(n) < log^3 A006450(n) for sufficiently large n. - _Thomas Ordowski_, Mar 22 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jul 16 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002315 | OEIS-A073133

Table by antidiagonals of T(n,k) = n*T(n,k-1) + T(n,k-2) starting with T(n,1) = 1.

This results in an infinite array in which sequences above the (1, 0, 1, 0, ...) are reflections of the sequences below, except for the alternate signs. Any sequence in the (+ sign) row starting (1, n, ...) is the (2*n-th) INVERT transform of the same sequence but with alternate signs. Example: (1, 2, 5, 12, ...) is the (2*2) = fourth INVERT transform of (1, -2, 5, -12, ...) by inspection. Conjecture: This "reflection" principle will result from taking successive INVERT transforms of any aerated sequence starting 1, ... and with positive signs. Likewise, the rows above the aerated sequence are successive INVERTi transforms of the aerated sequence. - _Gary W. Adamson_, Jul 14 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Jul 16 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002316 | OEIS-A073155

Leftmost column sequence of triangle A073153.

Conjecture: (n+1)*a(n) + (-3*n+4)*a(n-1) + 2*(-6*n+7)*a(n-2) + 2*(-6*n+11)*a(n-3) + 2*(-2*n+5)*a(n-4)=0. - _R. J. Mathar_, Nov 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002317 | OEIS-A073165

Triangle T(n,k) read by rows: related to David G. Cantor's sigma function.

Row n of T(n+k,k) has g.f. _(floor(n/2)+1)F_(floor(n/2))(1,3/2, 5/2,...,(2*floor(n/2)+1)/2;n,n-1,...,n-floor(n/2)+1;2^n*x) (conjecture). [_Paul Barry_, Jan 23 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Jul 24 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002318 | OEIS-A073271

a(n) = floor(prime(n)*prime(n+2) / prime(n+1)).

Conjecture: a(n) is even except for a(1)=3, a(3)=7 and a(8)=23 (A073272(n)=0). Conjecture: when a(n) = a(n+1) then A001223(n+1) = A001223(n) + A001223(n+2). - _Gionata Neri_, May 31 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jul 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002319 | OEIS-A073272

A000040(n+1) - A073271(n).

Observation/conjecture: a(n)=0 iff A073271(n) in {3, 7, 23}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jul 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002320 | OEIS-A073310

a(n) is the smallest number k such that 2+k and 2n+k are both prime.

Conjecture: a(n) < 2n. See A073316 for a generalization for all positive even numbers less than 2n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Aug 02 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002321 | OEIS-A073316

a(n) = Max d(j), j=1..n-1, where d(j) is the smallest positive number such that 2j+d(j) and 2n+d(j) are both prime. A generalization of A073310.

Conjecture: a(n) < 2n. Note that the truth of this conjecture implies that for any pair of positive even numbers e1 < e2 <= 2n, there is a positive odd number d < 2n such that e1+d and e2+d are primes. Note that this conjecture can also be stated with odd and even swapped: for any pair of positive odd numbers d1 < d2 < 2n, there is a positive even number e <= 2n such that e+d1 and e+d2 are primes. Also note that proving this conjecture would prove the twin primes conjecture. This is equivalent to a conjecture by Erdős mentioned by _R. K. Guy_ at the end of section C1 of his book. The conjecture has been verified for n < 10^5. - _T. D. Noe_, Nov 04 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Aug 02 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002322 | OEIS-A073330

Let c(1)=n and c(k+1)=floor(c(k)/Pi) if c(k) is even, or c(k+1)=floor(Pi*c(k)) otherwise; sequence gives the smallest value a(n) such that c(a(n))=0.

Conjecture: sequence always exists (a(123456789)=61, a(2^30)=70) and there is a constant C=2.8... such that sum(k=1, n, a(k)) is asymptotic to C*n*Log(n)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002323 | OEIS-A073341

Number of steps to reach an integer starting with (2n+1)/4 and iterating the map x -> x*ceiling(x).

We conjecture that an integer is always reached.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and J. C. Lagarias, Sep 04 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002324 | OEIS-A073369

Let c(1)=n, e=2.71... and c(k+1)=floor(c(k)/e) if c(k) is even, c(k+1)=floor(e*c(k)) otherwise; sequence gives the smallest value a(n) such that c(a(n))=0.

Conjecture: sequence always exists and there is a constant C=2.9... such that sum(k=1, n, a(k)) is asymptotic to C*n*Log(n). Is this constant the same as those for the sequence involving Pi instead of e?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 23 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002325 | OEIS-A073409

Largest prime factor of the denominator of the Bernoulli number B(2*n) (A002445).

Conjecture: a(n) is the largest prime p <= 2n+1 such that p * A000367(n) == - A002445(n) (mod p^2) for n > 0. Note that 2^(2n) == 1 (mod a(n)), since a(n) is the largest prime p such that b^(2n)== 1 (mod p) for every b coprime to p; i.e., a(n) is the largest prime p such that p-1 | 2n. - _Thomas Ordowski_, May 17 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 23 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002326 | OEIS-A073503

Numbers n such that the number of solutions to x^4 == 1 (mod n) is twice the number of solutions of x^2 == 1 (mod n).

Conjectures: 2n > a(n) or 2n < a(n) for infinitely many values of n and abs(a(n)-2n) < sqrt(n) for n > 45. a(n)=2n for n = 318, 338, 350, 488, 490, 492, 494,... Numbers divisible by 16 which have no prime factors = 1 mod 4, together with numbers not divisible by 16 which have exactly one prime factor = 1 mod 4. This refutes the conjectures. - _Charles R Greathouse IV_, Apr 16 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002327 | OEIS-A073504

A possible basis for finite fractal sequences: let u(1) = 1, u(2) = n, u(k) = floor(u(k-1)/2) + floor(u(k-2)/2); then a(n) = lim_{k->infinity} u(k).

Conjecture: a(n) = 2*(n - A027615(n))/3. - _Alan Michael Gómez Calderón_, Oct 16 2025 Calderón's conjecture is true (see Fried link in A072894). - _Sela Fried_, Mar 03 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_ and Boris Gourevitch (boris(AT)pi314.net), Aug 16 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002328 | OEIS-A073591

a(n) = A000522(n) + 1.

Conjecture: a(n) +(-n-2)*a(n-1) +(2*n-1)*a(n-2) +(-n+2)*a(n-3)=0. - _R. J. Mathar_, Feb 16 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Aug 28 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002329 | OEIS-A073602

a(n) is the smallest prime different from a(k) for k < n such that sum(a(j), j=1..n) is a multiple of the n-th prime.

Conjecture: Every prime belongs to this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 04 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002330 | OEIS-A073608

a(1) = 1, a(n) = smallest number such that a(n) - a(n-k) is a prime power > 1 for all k.

Differences |a(i)-a(j)| are prime powers for all i,j. Conjecture: sequence is bounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 04 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002331 | OEIS-A073631

Nonprimes k such that k divides 3^(k-1) - 2^(k-1).

All primes p>3 divide 3^(p-1) - 2^(p-1). It appears that a(1) = 1 and a(4) = 529 = 23^2 are the only perfect squares in a(n). Most terms of a(n) are squarefree. First 50 nonsquarefree terms of a(n) are the multiples of 23^2. Conjecture: All nonsquarefree terms of a(n) are the multiples of 23^2. Numbers n such that k=n*23^2 divides 3^(k-1) - 2^(k-1) are listed in A130058 = {1, 67, 89, 133, 199, 331, 617, 793, 881, 5281, 8911, 1419, 13333,...}. - _Alexander Adamchuk_, May 04 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002332 | OEIS-A073641

a(1) = 2; a(n) = smallest prime not included earlier such that concatenation of two successive terms is a prime.

Conjecture: every prime besides 5 is in this list. - Gabriel Cunningham (gcasey(AT)mit.edu), Apr 11 2003 The above conjecture by Cunningham (Apr 11 2003) is false: Since a(2)=3 and a(3)=7 == 1 mod 6, all subsequent terms must also be 1 mod 6 because concatenations of numbers 1 mod 6 with 5 mod 6 are 0 mod 3. - _Bob Selcoe_, Aug 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 09 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002333 | OEIS-A073642

Replace 2^k in the binary representation of n with k (i.e., if n = 2^b + 2^c + 2^d + ... then a(n) = b + c + d + ...).

Conjecture: a(n) = (3*A011371(n) - Sum_{k=1..n} A007814(k)^2)/2 for n > 0. - _Velin Yanev_, Sep 09 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002334 | OEIS-A073645

a(1)=2 and, for all n>=1, a(n) is the length of the n-th run of increasing consecutive integers with each run after the first starting with 1.

Conjecture: Let P(k)=1 + k/3 + k^2/2 + k^3/6. Then a(n)=3 if n=P(k) for some k, a(n)=2 if P(k-1)<n<P(k) for some k and P(k)-n=m(m+1)/2 for some m, else a(n)=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Aug 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002335 | OEIS-A073652

Primes which occur as the difference of consecutive prime powers >1 as and when they occur.

Conjecture: Every prime is a member.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002336 | OEIS-A073653

a(1)=3, a(2)=5; for n > 2, a(n) = smallest prime not included earlier such that a(n-2) + a(n-1) + a(n) is a prime.

Conjecture: Every odd prime eventually appears; a(n) ~ prime(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002337 | OEIS-A073659

a(1) = 1; for n > 1, a(n) is the smallest even number not already in the sequence such that a(1) + ... + a(n) is a prime.

Conjecture: Every even number appears. - _N. J. A. Sloane_, May 29 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002338 | OEIS-A073673

Rearrangement of natural numbers such that every partial product + 1 is a prime.

Conjecture: n/log(n) << a(n) << n*log(n). - _Thomas Ordowski_, Aug 09 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002339 | OEIS-A073681

Smallest of three consecutive primes whose sum is a prime.

Conjecture: for n -> oo, a(n) ~ prime(n) * (log(prime(n)))^C, where C = 8/Pi^2 (cf. A217739). - _Alain Rocchelli_, Sep 04 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002340 | OEIS-A073751

Prime numbers that when multiplied in order yield the sequence of colossally abundant numbers A004490.

This calculation assumes that the ratio of consecutive colossally abundant numbers is always prime, which is implied by a conjecture mentioned in Lagarias's paper.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Aug 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002341 | OEIS-A073759

Largest number < n that is neither a divisor of nor relatively prime to n, or 0 if no such number exists.

Conjecture: a(n) = 2*A046667(n) if n>=5. - _R. J. Mathar_, May 15 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Aug 15 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002342 | OEIS-A073761

Primitive numbers whose decimal expansion of 1/n is equidistributed in base 10.

Conjecture 1: let p be a prime such that ord(10,p) is a multiple of 10, where ord(a,m) denotes the multiplicative order of a modulo m. Then p is a term if and only if 10 is a primitive root modulo p. Conjecture 2: suppose that m is a term with bigomega(m) = 2, then m = p*q, where p == 1 (mod 10), q == 9 (mod 10), gcd(p-1,q-1) = 2, ord(10,p) = (p-1)/2, and ord(10,q) = q-1. Note that the converse is not true, though. There are no counterexamples to the conjectures above below 100000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Donald S. McDonald_, Sep 02 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002343 | OEIS-A073763

Least number of unrelated set belonging to these numbers is odd.

Conjecture: a(n) = 36*n - 18 - 6*(-1)^n = 24 * A001651(n). - _Ralf Stephan_, Oct 19 2013 The conjecture is false, first counterexample being a(1541) = 55440. - _Robert Israel_, Sep 11 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Aug 08 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002344 | OEIS-A073783

First differences of composite numbers.

Conjecture: a(n) = A065310(n+1). - _R. J. Mathar_, Jul 17 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lior Manor_, Aug 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002345 | OEIS-A073825

Numbers n such that Sum_{k=1..n} k^k is prime.

Conjecture: This sequence is infinite. - _Daniel Hoying_, Jul 20 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rick L. Shepherd_, Aug 13 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002346 | OEIS-A073842

a(1) = 1; for n>1, a(n) = the smallest positive integer root of n not included earlier, if such a root exists, otherwise the smallest power of n not included earlier, subject to a(n)<>n.

Conjectured to be a rearrangement of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 13 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002347 | OEIS-A073846

a(1) = 1; thereafter, every even-indexed term is prime and every odd-indexed term is composite.

* R#(34) (A261314) has been checked up to f(34,86) = 1091595086717, R#(42) up to f(42,108) = 106838266736, R#(50) up to f(50,98) = 1078406742163, R#(60) up to f(60,80) = 765456394363, R#(74) up to f(74,78) = 687059343029, R#(82) up to f(82,75) = 682580868743 R#(86) up to f(86,74) = 182831963148, R#(88) up to f(88,66) = 719074799059, and R#(98) up to f(98,88) = 641383978721 without repeated values. Hence, their periods are either extremely large or nonexistent (infinite). I

conjecture that the latter is the case. Note that these sequences are not necessarily all distinct as any two may simply be the same sequence with a large offset. Conjecturally, the integers that belong to one of these cycles are: 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 35, 37, 39, 41, 43, 47, 53, 62, 84, 87, 92, 121, 127, 132, 135, 181, 199, 205, 317. - _Michel Marcus_, Mar 07 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 14 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002348 | OEIS-A073883

a(1)=1. For n > 1, a(n) = the smallest positive integer not included earlier such that the n-th partial sum is n times a prime.

Conjectured not to be a permutation of the natural numbers. The numbers that have not been used before the 2000th term: 4, 8, 9, 10, 12, 14, 15, 16, 18, 20, 22, ... - _Derek Orr_, Jun 26 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 16 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002349 | OEIS-A073918

Smallest prime which is 1 more than a product of n distinct primes: a(n) is a prime and a(n) - 1 is a squarefree number with n prime factors.

Conjecture: For any m > 0 there is K > 0 such that for all k > K, a(k)-1 is divisible by the first m primes. Conjecture 2: Let K(m) be the smallest possible K satisfying the above Conjecture. Then K(m) ~ m, i.e., a(k) ~ A002110(k), only very few of the last factors will be a bit larger. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002350 | OEIS-A073923

a(1)=1, a(n) is the smallest power of 2 not included earlier such that Sum_{k=1..n} a(k) is prime.

Conjecture: a(n) = A059661(n)-A059661(n-1). - _R. J. Mathar_, May 07 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002351 | OEIS-A073942

Numbers k such that A002977(k) - A002977(k-1) = 1.

Conjecture : a(n) is asymptotic to c*n where c is around 3.8...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 20 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002352 | OEIS-A074057

a(n) = 2*phi(n-2)-(n-1).

Conjecture : a(n)=0 if and only if n is a Fermat prime (A019434). The conjecture appears to be false, a(83623937)=0 and 83623937 is not a Fermat number (A000215). See A232720 for sequence of n such that a(n)=0. - _Michel Marcus_, Nov 28 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Sep 15 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002353 | OEIS-A074063

a(n) is the number of essentially different ways in which the integers 1,2,3,...,n can be arranged in a sequence such that (1) adjacent integers sum to a prime number and (2) squares of adjacent numbers sum to a prime number. Rotations and reversals are counted only once.

It is known that a(n) > 0 for 58 <= n <= 200. It is conjectured that a(n) > 0 for all n > 57. A greedy algorithm can be used to quickly find a solution for many n. See the link to puzzle 189 for more details. The Mathematica program uses a backtracking algorithm to count the arrangements. To print the unique arrangements, remove the comments from around the print statement.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Aug 17 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002354 | OEIS-A074100

Cubes using only digits 1, 2, 3, 5 and 7.

Conjecture: the sequence is finite. Opposite conjecture: the sequence is infinite. The frequency of terms with k digits is 4, 3, 5, 5, 9, 11, 12, 13, 22, 29, 33, 37, 49, 49, 78 for k = 1..15 respectively. - _David A. Corneth_, Mar 17 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002355 | OEIS-A074162

a(1) = 1; a(n) = smallest multiple of n (>n) using all nonprime digits (1,4,6,8,9 and 0) if n is a prime else using all prime digits and 1 if n is composite; or 0 if no such number exists.

A pigeonhole argument shows the conjecture to be true (that only a(10k) = 0). - _Joshua Zucker_, May 08 2006 Conjecture (contradicts Zucker statement): a(k*5^6) = 0. - _Sean A. Irvine_, Jan 13 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002356 | OEIS-A074208

Least k > 1 such that n divides sigma(k) - k.

Conjecture: Sum_{k=1..n} a(k) = O(n^{2+epsilon}) for any epsilon > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Sep 17 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002357 | OEIS-A074216

Squares m satisfying sigma(m) == 0 (mod 3).

Conjecture: a(n) = A072864(n)^2. - _R. J. Mathar_, May 19 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Sep 17 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002358 | OEIS-A074270

a(n) = the least positive number X such that Cn(X) is X-smooth, where Cn is the n-th cyclotomic polynomial and "X-smooth" means "has no prime factor greater than X".

Inspired by a conjecture of David W. Wilson: for each nonzero polynomial P with integer coefficients, there is an integer X such that P(X) is X-smooth. Certain cyclotomic polynomials seem to stress the conjecture, but no refutation is yet known. If a(11) exists, it is greater than 1060000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Don Reble_, Sep 20 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002359 | OEIS-A074354

Coefficient of q^3 in nu(n), where nu(0)=1, nu(1)=b and, for n>=2, nu(n)=b*nu(n-1)+lambda*(1+q+q^2+...+q^(n-2))*nu(n-2) with (b,lambda)=(1,2).

Conjectures from _Colin Barker_, Nov 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Y. Kelly Itakura (yitkr(AT)mta.ca), Aug 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002360 | OEIS-A074355

Coefficient of q^1 in nu(n), where nu(0)=1, nu(1)=b and, for n>=2, nu(n)=b*nu(n-1)+lambda*(1+q+q^2+...+q^(n-2))*nu(n-2) with (b,lambda)=(1,3).

G.f.: (9x^4+3x^3)/(1-3x-3x^2)^2 (conjectured). - _Ralf Stephan_, May 09 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | Y. Kelly Itakura (yitkr(AT)mta.ca), Aug 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002361 | OEIS-A074356

Coefficient of q^2 in nu(n), where nu(0)=1, nu(1)=b and, for n>=2, nu(n)=b*nu(n-1)+lambda*(1+q+q^2+...+q^(n-2))*nu(n-2) with (b,lambda)=(1,3).

Conjectures from _Colin Barker_, Nov 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Y. Kelly Itakura (yitkr(AT)mta.ca), Aug 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002362 | OEIS-A074357

Coefficient of q^3 in nu(n), where nu(0)=1, nu(1)=b and, for n>=2, nu(n)=b*nu(n-1)+lambda*(1+q+q^2+...+q^(n-2))*nu(n-2) with (b,lambda)=(1,3).

Conjecture: O.g.f.: 3*x^5*(3*x+1)*(36*x^4+24*x^3-29*x^2-14*x+10)/(3*x^2+x-1)^4. - _R. J. Mathar_, Jul 22 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | Y. Kelly Itakura (yitkr(AT)mta.ca), Aug 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002363 | OEIS-A074358

Coefficient of q^1 in nu(n), where nu(0)=1, nu(1)=b and, for n >= 2, nu(n) = b*nu(n-1) + lambda*(1 + q + q^2 + ... + q^(n-2))*nu(n-2) with (b,lambda)=(2,2).

G.f.: 4*x^3*(x + 1)/(2*x^2 + 2*x - 1)^2 (conjectured). - _Chai Wah Wu_, May 30 2016 a(n) = (1/18)*((1 + sqrt(3))^n*(-9 + 2*sqrt(3)) - (1 - sqrt(3))^n*(9 + 2*sqrt(3)) + 3*((1 - sqrt(3))^n + (1 + sqrt(3))^n)*n) for n > 0 (conjectured). - _Colin Barker_, Nov 17 2017 a(n) = 4*a(n-1) - 8*a(n-3) - 4*a(n-4) for n > 4 (conjectured). - _Colin Barker_, Nov 17 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | Y. Kelly Itakura (yitkr(AT)mta.ca), Aug 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002364 | OEIS-A074359

Coefficient of q^2 in nu(n), where nu(0)=1, nu(1)=b and, for n>=2, nu(n)=b*nu(n-1)+lambda*(1+q+q^2+...+q^(n-2))*nu(n-2) with (b,lambda)=(2,2).

Conjecture: O.g.f: 4*x^4*(-3+2*x+8*x^2+4*x^3)/(2*x^2+2*x-1)^3. - _R. J. Mathar_, Jul 22 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | Y. Kelly Itakura (yitkr(AT)mta.ca), Aug 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002365 | OEIS-A074360

Coefficient of q^3 in nu(n), where nu(0)=1, nu(1)=b and, for n>=2, nu(n)=b*nu(n-1)+lambda*(1+q+q^2+...+q^(n-2))*nu(n-2) with (b,lambda)=(2,2).

Conjecture: O.g.f: 8*x^5*(1+x)*(12*x^4+24*x^3-2*x^2-16*x+5)/(2*x^2+2*x-1)^4. - _R. J. Mathar_, Jul 22 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | Y. Kelly Itakura (yitkr(AT)mta.ca), Aug 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002366 | OEIS-A074482

Consider the recursion b(1,n) = 1, b(k+1,n) = b(k,n) + (b(k,n) reduced mod(k+n)); then there is a number x such that b(k,n) - b(k-1,n) is a constant x depending only on n, for k > y = A074483(n). Sequence gives values of x.

Conjecture: a(n) is defined for all n (as well as A074483);

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_ and _Benoit Cloitre_, Aug 23 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002367 | OEIS-A074483

Consider the recursion b(1,n) = 1, b(k+1,n) = b(k,n) + (b(k,n) reduced mod(k+n)); then there is a number y such that b(k,n)-b(k-1,n) is a constant (= A074482(n)) for k > y. Sequence gives values of y.

Conjecture: a(n) is defined for all n (as well as A074482);

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_ and _Benoit Cloitre_, Aug 23 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002368 | OEIS-A074650

Table T(n,k) read by downward antidiagonals: number of Lyndon words (aperiodic necklaces) with n beads of k colors, n >= 1, k >= 1.

The fact that T(3,4) = 20 coincides with the number of the amino acids encoded by DNA made Francis Crick, John Griffith and Leslie Orgel conjecture in 1957 that the genetic code is a comma-free code, which later turned out to be false. [Hayes] - _Andrey Zabolotskiy_, Mar 24 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, Aug 28 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002369 | OEIS-A074724

Highest power of 3 dividing F(4n) where F(k) is the k-th Fibonacci number.

a(n) = 3^A051064(n) (conjectured). - _Michel Marcus_, May 17 2022 Conjecture: a(n) = (sigma(3*n) - sigma(n))/(sigma(3*n) - 3*sigma(n)), where sigma(n) = A000203(n). Equivalently, a(n) = A088838(n) - A074724(n). - _Peter Bala_, Jun 10 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Sep 04 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002370 | OEIS-A074726

Numbers k such that sigma(F(k)) > 2*F(k) where F(k) is the k-th Fibonacci number.

Conjecture: sigma(F(n)) > 2*F(n) if and only if F(n) is a Zumkeller number except for n = 12. Verified for n <= 371. - _M. Farrokhi D. G._, Aug 16 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Sep 04 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002371 | OEIS-A074757

Numbers n such that tau(n) = (tau(n+1)+tau(n-1))/2.

a(n) seems to be asymptotic to c*n with c around 23. [This conjecture is false. The limit a(n)/n, if it exists, is > 29, see graphs. - _Vaclav Kotesovec_, Feb 14 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Sep 28 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002372 | OEIS-A074772

Numbers k such that tau(k) < tau(k+1) and phi(k) < phi(k+1).

It seems that a(n) is asymptotic to c*n with 14<=c<=16. [This conjecture is false, see plot. - _Vaclav Kotesovec_, Feb 16 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Sep 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002373 | OEIS-A074777

Integers n such that sigma(phi(n))/n = 1/2.

Since 2^(2^n)+1 is prime for n=0,1,...,4 (Fermat primes), 2^(2^(n-1)+1)-2 is in the sequence for n=1,2,...,6. Conjecture: There are no further terms. - _Farideh Firoozbakht_, Sep 14 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | Antonio G. Astudillo (afg_astudillo(AT)hotmail.com), Sep 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002374 | OEIS-A074794

Number of numbers k <= n such that tau(k) == 1 (mod 3) where tau(k) = A000005(k) is the number of divisors of k.

The constant is conjecturally 3*zeta(3)/Pi^2 = 0.3653814847007... (A346602). See A211337 for more details. - _Amiram Eldar_, Feb 01 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Sep 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002375 | OEIS-A074796

Number of numbers k <= n such that tau(k) == 2 (mod 3) where tau(k) = A000005(k) is the number of divisors of k.

The constant is conjecturally 3*zeta(3)/Pi^2 = 0.3653814847007... (A346602). See A211338 for more details. - _Amiram Eldar_, Feb 01 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Sep 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002376 | OEIS-A074882

Number of integers in {1, 2, ..., sigma(n)} that are coprime to n.

a(n) <= n, with equality if n is in A000961. Conjecture: a(n) < n if n is not in A000961. - _Robert Israel_, Aug 10 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Oct 04 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002377 | OEIS-A074915

Largest x such that the number of nonprimes (i.e., 1 and composites) in the reduced residue set (RRS(x)) of x equals n, or 0 if there are no such numbers.

It is conjectured that x is always bounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Oct 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002378 | OEIS-A074943

a(n) = tau(n) mod 3.

Conjecture: Asymptotic mean: Limit_{m->oo} (1/m) * Sum_{k=1..m} a(k) = 9*zeta(3)/Pi^2 = 1.0961... . The conjecture is true if A211337 and A211338 have the same asymptotic density (see also A059269). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 04 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002379 | OEIS-A074944

Number of k with 1 <= k <= n such that gcd(n,k) = tau(k), where tau is A000005, number of divisors function.

The above conjecture seems to be wrong. See the graph in the Links section. - _Amiram Eldar_, Mar 10 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002380 | OEIS-A074964

Numbers k such that Max (sigma(x*y) : 1 <= x <= k, 1 <= y <= k) = sigma(k^2).

Conjecture: subsequence of A066522, implying finiteness. - _Reinhard Zumkeller_, Nov 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002381 | OEIS-A074980

Numbers which are not of the form m^p - n^q where p = 2 or 3, q = 2 or 3.

This is a famous hard problem and the terms shown are only conjectured values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Oct 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002382 | OEIS-A074981

Conjectured list of positive numbers which are not of the form r^i - s^j, where r,s,i,j are integers with r>0, s>0, i>1, j>1.

Conjectured list of positive numbers which are not of the form r^i - s^j, where r,s,i,j are integers with r>0, s>0, i>1, j>1. This is a famous hard problem and the terms shown are only conjectured values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Oct 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002383 | OEIS-A074987

a(n) is the least m not equal to n such that phi(m) = phi(n).

I conjecture that a(n) <= 2n. I have checked this for all n <= 10^4. (It is not possible to do better than the 2n upper bound since a(11) = 2*11.) For odd n the conjecture is true because phi(n)=phi(2n). - _T. D. Noe_, Oct 18 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Oct 02 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002384 | OEIS-A075000

Smallest number such that n*a(n) is a concatenation of n consecutive integers; or 0 if no such number exists.

Conjecture: For every n there exists a nonzero a(n). The conjecture that a(n) > 0 is proved in Wang (2026). - _Qichao Wang_, Sep 23 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 31 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002385 | OEIS-A075001

Smallest k such that the concatenation of n consecutive numbers starting with k (from k to n+k-1) is a multiple of n; or 0 if no such number exists.

Conjecture: For every n there exists a k. The conjecture that a(n) > 0 for all n >= 1 is proved in Wang (2026). - _Qichao Wang_, Sep 20 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 31 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002386 | OEIS-A075036

Smaller of two smallest consecutive numbers with 2n divisors.

Conjecture: if a term k is even, the highest p-adic order of k (the maximum may be attained by several p's) occurs at p=2 and the highest p-adic order of k+1 occurs at p=3. If a term k is odd, the highest p-adic order of k occurs at p=3 and the highest p-adic order of k+1 occurs at p=2. - _Chai Wah Wu_, Mar 12 2019 a(n) <= A215199(n-1) for n > 1. Conjecture: if p is prime, then a(p) = A215199(p-1). This conjecture is true if the conjecture in A215199 is true. The b-file of A215199 thus shows that a(p) = A215199(p-1) for prime p < 1279. - _Chai Wah Wu_, Mar 12 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 03 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002387 | OEIS-A075045

Coefficients A_n for the s=3 tennis ball problem.

Conjecture: D-finite with recurrence 8*(2*n+3)*(7*n+1)*(n+1)*a(n) +6*(-252*n^3-477*n^2-220*n-11)*a(n-1) +81*(7*n+8)*(3*n-1)*(3*n+1)*a(n-2)=0. - _Jean-François Alcover_, Feb 07 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002388 | OEIS-A075075

a(1) = 1, a(2) = 2 and then the smallest number not occurring earlier such that every term divides the product of its neighbors: a(n-1)*a(n+1)/a(n) is an integer.

This is a permutation of natural numbers. [_Leroy Quet_ asks (May 06 2009) if this is a theorem or just a conjecture.] Conjecture: if a(n) divides a(n-1) then a(n+1) is prime. - _Thomas Ordowski_, Jul 11 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 09 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002389 | OEIS-A075077

a(1) = 1 and then the smallest number not occurring earlier such that a(n) + a(n+1) gives a different composite number (not obtained earlier).

Conjecture: for n > 6, n - 2 <= a(n) <= n + 1. - _Robert Israel_, Jun 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 09 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002390 | OEIS-A075082

Numbers n such that n! is a product of distinct factorials k!*l!*m!*... with k, l, m, etc. < n.

Subsequence of A034878 (all n such that n! is a product of smaller factorials). It is conjectured that A034878 and A001013 (Jordan-Polya numbers = products of factorials) are the same sequence (except for the numbers 2, 9 and 10). If this is true, then obviously A075082 (without the number 10) is also a subsequence of A001013. On the other hand, this special case of the conjecture might be easier to prove. (a(n)!)^2 is a member of A058295 (products of distinct factorials); for example, (6!)^2 = 6!*5!*3!. - _Jonathan Sondow_, Dec 21 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002391 | OEIS-A075092

Sum of generalized tribonacci numbers (A001644) and reflected generalized tribonacci numbers (A073145).

Conjecture: a(n) >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Mario Catalani (mario.catalani(AT)unito.it), Aug 31 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002392 | OEIS-A075125

Number of parallelogram polyominoes of site-perimeter n (also called staircase polyominoes, although that term is overused).

Conjecture: -(2*n-11)*(n-2)*(2*n-9)*a(n) +4*(2*n-11)*(n-3)*(n-5)*a(n-1) +(4*n^3-60*n^2+317*n-582)*a(n-2) +2*(2*n-7)*(2*n^2-26*n+81)*a(n-3) -(n-10)*(2*n-7)*(2*n-9)*a(n-4)=0. - _R. J. Mathar_, May 30 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | Andrew Rechnitzer (a.rechnitzer(AT)ms.unimelb.edu.au), Sep 09 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002393 | OEIS-A075127

Safe perfect powers: perfect powers n such that (n-1)/2 is also a perfect power.

Conjectures from _Colin Barker_, Oct 28 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Oct 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002394 | OEIS-A075326

Anti-Fibonacci numbers: start with a(0) = 0, and extend by the rule that the next term is the sum of the two smallest numbers that are not in the sequence nor were used to form an earlier sum.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 16 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002395 | OEIS-A075343

a(1) = 1, a(n+1) is the smallest number such that there are n composite numbers between a(n) and a(n+1) exclusive.

Conjecture: there exists a prime between consecutive terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002396 | OEIS-A075344

Number of primes between consecutive terms of A075343 (excluding endpoints).

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002397 | OEIS-A075432

Primes with no squarefree neighbors.

This sequence is infinite by Dirichlet's theorem, since there are infinitely many primes == 17 or 19 (mod 36) and these have no squarefree neighbors. Ludovicus's conjecture about density is correct. Capsule proof: either p-1 or p+1 is divisible by 4, so it suffices to consider the other number (without loss of generality, p+1). For some fixed bound L, p is not divisible by any prime q < L (with finitely many exceptions) so there are q^2 - q possible residue classes for p. The primes in each are uniformly distributed so the probability that p+1 is divisible by q^2 is 1/(q^2 - q). The product of the complements goes to 2A as L increases without bound, and since 2A is an upper bound the limit is sandwiched between. - _Charles R Greathouse IV_, Aug 27 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Sep 15 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002398 | OEIS-A075568

Rearrangement of noncomposite numbers such that every partial sum is composite.

Conjectured to consist of consecutive primes after 53. (See also comments in A123055.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 25 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002399 | OEIS-A075586

Primes p such that the number of distinct prime divisors of all composite numbers between p and the next prime is 6.

For very large n, the probability of a(n) not being a twin prime is extremely small, unless the twin primes conjecture is false. - _Sam Alexander_, Oct 20 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 26 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002400 | OEIS-A075594

a(1) = 1, a(2n) = composite, a(2n+1) = prime and sum of two successive terms is alternately prime or composite.

Conjecture: a(n+r)= A075593(n) for n > k for some k and r. What is the value of k and r?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 27 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002401 | OEIS-A075786

Palindromic perfect powers.

Conjecture: there are no palindromic perfect powers with prime exponent > 3. - _Chai Wah Wu_, Aug 26 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Oct 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002402 | OEIS-A075788

If n can be written in the form r^i-s^j (see A074981), where r,s,i,j are integers with r>0, s>0, i>1, j>1 choose the representation with smallest r^i (in case of ties, minimize i and j); or if n is not of this form set r=s=i=j=0; sequence gives values of r.

The zeros are only conjectures (cf. A074981).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Oct 13 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002403 | OEIS-A075789

Value of i, when n is written as r^i - s^j with the smallest possible r^i (with minimal i) and r, s > 0, i, j > 1; or 0 if n is not of this form.

The zeros are only conjectures (cf. A074981).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Oct 13 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002404 | OEIS-A075790

If n can be written in the form r^i-s^j (see A074981), where r,s,i,j are integers with r>0, s>0, i>1, j>1 choose the representation with smallest r^i (in case of ties, minimize i and j); or if n is not of this form set r=s=i=j=0; sequence gives values of s.

The zeros are only conjectures (cf. A074981).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Oct 13 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002405 | OEIS-A075791

If n can be written in the form r^i-s^j (see A074981), where r,s,i,j are integers with r>0, s>0, i>1, j>1 choose the representation with smallest r^i (in case of ties, minimize i and j); or if n is not of this form set r=s=i=j=0; sequence gives values of j.

The zeros are only conjectures (cf. A074981).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Oct 13 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002406 | OEIS-A075855

Maximum number of black squares on an n X n chessboard (with a black square in at least one corner) that can be covered by a single path, traveling only to adjacent black squares.

For n odd, a(n)=(n-1)^2/2+1. For n even, it is conjectured that a(n)=(n^2-n+2)/2 (it is easy to show this is a lower bound).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Oct 15 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002407 | OEIS-A075885

a(n) = 1 + n + n*[n/2] + n*[n/2]*[n/3] + n*[n/2]*[n/3]*[n/4] +... where [x]=floor(x).

Conjecture: limit a(n)^(1/n) = L where L = 2.200161058099... is the geometric mean of Luroth expansions, where log(L) = Sum_{n>=1} log(n)/(n*(n+1)) = 0.7885305659115... (cf. A085361).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 16 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002408 | OEIS-A075989

Number of k satisfying 1<=k<=n and {n/k} >= 1/2, where {n/k} is the fractional part of n/k, i.e., {n/k} = n/k - floor(n/k).

a(n) = Sum_{k=1..n} (floor(2*n/(2*k+1)) - floor(2*n/(2*k+2))); a(n) = (2*(log 2)-1)*n + O(n^(1/2)). Conjecture: a(n) = (2*(log 2)-1)*n + O(n^(1/4 + epsilon)) like for the divisor and Circle problems. - _Benoit Cloitre_, Oct 21 2012 Conjecture: Let f(a,b)=1, if (a+b) mod |a-b| != (a mod |a-b|)+(b mod |a-b|), and 0 otherwise. a(n) = Sum_{k=1..n-1} f(n,k). - _Benedict W. J. Irwin_, Sep 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 28 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002409 | OEIS-A075997

a(n) = [n/2] - [n/3] + [n/4] - [n/5] + [n/6] - ..., where [n/k] = floor(n/k).

Conjecture: Let f(a,b)=1, if (a+b) mod |a-b| != (a mod |a-b|)+(b mod |a-b|), and 0 otherwise. a(n) = Sum_{k=1..n-1} f(n+k,n-k). - _Benedict W. J. Irwin_, Sep 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 28 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002410 | OEIS-A076035

Expansion of g.f. 1/(1-4*x*C) where C = (1/2-1/2*(1-4*x)^(1/2))/x = g.f. for Catalan numbers A000108.

Conjecture: a(n) = 4*A076025(n), n>0. - _R. J. Mathar_, Apr 01 2022 Mathar's conjecture is true (follows from the g.f.s of the two sequences). - _Sela Fried_, Dec 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002411 | OEIS-A076043

a(1)=2; for n>1, a(n) is smallest squarefree number (greater than 1) not occurring earlier such that a(n) + a(n-1) is a square.

Conjecture: every squarefree number except 1 is a member.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 03 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002412 | OEIS-A076044

Largest a(n) values with at most n primes between a(n) and a(n)+sqrt(a(n)) inclusive.

Conjecture: for every m greater than a(n), there are more than n primes between m and m+sqrt(m); true if a(n) less than 1000000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Adam Kertesz_, Oct 28 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002413 | OEIS-A076046

Ramanujan-Nagell numbers: the triangular numbers (of the form a*(a+1)/2) which are also of the form 2^b - 1.

Ramanujan conjectured and Nagell proved, that the given numbers are the only ones. This sequence is equivalent to A060728, the list of numbers n such that x^2 + 7 = 2^n is soluble, by changing from n to 2^(n-3)-1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Burt Totaro (b.totaro(AT)dpmms.cam.ac.uk), Oct 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002414 | OEIS-A076070

Smallest multiple of n that can be formed by using the digits of the next n numbers, or 0 if no such number exists.

Conjecture: No entry is zero. (At least one multiple of n can be formed by using the digits of the next n numbers).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Oct 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002415 | OEIS-A076099

Triangle read by rows in which the n-th row contains n numbers not occurring in the previous rows whose product is an n-th power. The first (n-1) numbers of the n-th row are the smallest number not occurring earlier and the n-th term is chosen to satisfy the requirement.

The first n-1 elements of the n-th group are the smallest n-1 numbers that haven't already appeared, say u1, u2, ..., u_(n-1) and let u_n be the unknown final element of the n-th group. Let u1*u2*u3*...*u_(n-1) = (p1^e1)(p2^e2)...(pr^er). Then u_n = product(i=1 to r) p_i^(ei + n*floor(ei/n) - n) ...unless this has already appeared in the sequence (probably this never happens). More simply, I conjecture that u_n = product(i=1 to r) p_i^(ei - n). - _Sam Alexander_, Dec 31 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Oct 08 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002416 | OEIS-A076135

Numbers n such that O(n) = E(n)-1, where O(n), E(n) denote the number of integers k, 1 <= k <= n, such that Omega(n) is even or odd, respectively and Omega(n) denotes the number of prime factors of n, counting multiplicity.

In 1919, George Polya conjectured that O(n) >= E(n) for n >= 2. However, in 1958, C. B. Haselgrove showed that there are infinitely many positive integers n for which O(n) < E(n). In 1966, R.S. Lehman showed that 906180359 is the smallest positive integer for which O(n) = E(n) - 1. (Tattersall, p. 92)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Oct 30 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002417 | OEIS-A076157

Continued fraction expansion for c = Sum_{k>=0} 1/2^(k!).

Conjecture: a(3*2^n) = -1 + 2^[(n+1)((n+2)!)]. - _Ralf Stephan_, May 17 2005 The conjecture follows from the theorem in Shallit's paper. The continued fraction has a "folded" overall structure. - _Georg Fischer_, Aug 29 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 02 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002418 | OEIS-A076217

a(1)=1, a(n) = a(n-1) + n * sign(n-a(n-1)).

Conjecture: a(n) = -a(n-1)+a(n-2)+a(n-3) for n>5. G.f.: -x*(27*x^28 +54*x^27 +27*x^26 +9*x^10 +18*x^9 +9*x^8 +3*x^4 +6*x^3 +5*x^2 +4*x +1) / ((x -1)*(x +1)^2). - _Colin Barker_, Feb 25 2013 Regarding Barker's conjectured recurrence, it seems to fail at n= powers of 3, and the 2 successive terms. So it holds except for n= 9-11, 27-29, 81-83, 243-245, ... - _Bill McEachen_, Mar 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 03 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002419 | OEIS-A076218

Numbers n such that 2*n^2 - 3*n + 1 is a square.

Conjecture: a nonzero number occurs twice in A055524 if and only if it's in this sequence. - _J. Lowell_, Jul 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gregory V. Richardson_, Nov 03 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002420 | OEIS-A076335

Brier numbers: numbers that are both Riesel and Sierpiński [Sierpinski], or odd n such that for all k >= 1 the numbers n*2^k + 1 and n*2^k - 1 are composite.

It is a conjecture that every such number has more than 10 digits. In 2011 I have calculated that for any n < 10^10 there is a k such that either n*2^k + 1 or n*2^k - 1 has all its prime factors greater than 1321. - _Arkadiusz Wesolowski_, Feb 03 2016 [Editor's note: The comment below states that the conjecture is now proved. - _M. F. Hasler_, Oct 06 2021]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_, Nov 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002421 | OEIS-A076337

Riesel numbers: odd numbers n such that for all k >= 1 the numbers n*2^k - 1 are composite.

509203 has been proved to be a member of the sequence, and is conjectured to be the smallest member. However, as of 2009, there are still several smaller numbers which are candidates and have not yet been ruled out (see links). Riesel numbers are proved by exhibiting a periodic sequence p of prime divisors with p(k) | n*2^k-1 and disproved by finding prime n*2^k-1. It is conjectured that numbers that cannot be proved Riesel in this way are non-Riesel. However, some numbers resist both proof and disproof. Others conjecture the opposite: that there are infinitely many Riesel numbers that do not arise from a covering system, see A101036. The word "odd" is needed in the definition because otherwise for any term n, all numbers n*2^m, m >= 1, would also be Riesel numbers, but we don't want them in this sequence (as is manifest from A101036). Since 1 and 3 obviously are not in this sequence, for any n in this sequence n-1 is an even number > 2 and therefore composite, so one could replace "k >= 1" equivalently by "k >= 0". - _M. F. Hasler_, Aug 20 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002422 | OEIS-A076449

Least number whose digits can be used to form exactly n different primes (not necessarily using all digits).

Mike Keith conjectures that a(n) always exists and reports that he has checked this for n <= 66. - _N. J. A. Sloane_, Jan 25 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Nov 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002423 | OEIS-A076502

a(1)=1, a(n)=n-a(n-a(n-a(n-1))).

Conjecture : a(n) - c*n is bounded and a(n)=floor(c*n) + 0 + 1 or +2. The conjecture that a(n) is always one of floor(c*n), floor(c*n) + 1, or floor(c*n) + 2 is false. The first counterexample is at n = 1167, for which the recurrence gives a(1167) = 664. Let P(x) = x^3 - x^2 + 2*x - 1, and let c be its positive root. Then P(665/1167) = -9043/1589324463 < 0 and P(666/1167) = 92227/58863869 > 0. Since P'(x) = 3*(x - 1/3)^2 + 5/3 > 0 for every real x, P is strictly increasing, so 665/1167 < c < 666/1167. Hence floor(1167*c) = 665, whereas a(1167) = 664 = floor(1167*c) - 1. This does not disprove the separate conjecture that a(n) - c*n is bounded. - _Eiver Rodríguez_, Sep 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 08 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002424 | OEIS-A076552

a(n) = (-1)^(n+1)/3/(2n+1) * Sum_{k=0..n} 16^k*B(2k)*C(2n+1,2k) where B(k) denotes the k-th Bernoulli number.

Conjectural e.g.f. with offset 0 (checked up to a(14)): 1/3*(2 - cos(x)^2 + 2*cos(x)^4)/cos(x)^3 = 1 + x^2/2! + 21*x^4/4! + 461*x^6/6! + ... [added 24 Aug 2026: this is correct.] (End) Conjecture: a(n) = -1/3*(-4)^n*E(2*n,-1/2), where E(n,x) is the n-th Euler polynomial. - _Peter Bala_, Sep 25 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002425 | OEIS-A076564

List of numbers k such that the k-th Fibonacci number (A000045) does not contain the digit zero.

Altassan and Luca (2019) conjectured that 184 is the last term. They proved that the asymptotic density of this sequence is 0, and moreover, the number of terms below x is << x^(1-c), where c = 1 - log_10(9) = 0.045757... . - _Amiram Eldar_, Jan 02 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Oct 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002426 | OEIS-A076614

Numbers of the form 2^k + 3^k + ... + p_n^k.

It is conjectured that 160 is the only number to appear twice: 2+3+...+31=160 and 2^3+3^3+5^3=160

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Nov 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002427 | OEIS-A076689

Smallest k such that k*prime(n)# + 1 is prime where prime(n)# is the n-th primorial number A002110(n).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Nov 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002428 | OEIS-A076768

Positive integers not expressible as the sum of a prime and a triangular number.

This sequence contains 216 (and possibly other nontriangular numbers) together with an infinite number of triangular numbers. The indices of the triangular numbers are in A138666. This is related to the Sun's conjecture (see A132399) that every number except 216 is the sum of a triangular number and a prime or 0. - _T. D. Noe_, Mar 26 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Nov 14 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002429 | OEIS-A076771

Even numbers n representable as the sum of two non-coprime odd composites.

It is conjectured that there is no N such that for all n > N, every even number is represented. The conjecture is true since 2p, p prime, is never a member of this sequence. - _Charles R Greathouse IV_, Aug 08 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Nov 14 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002430 | OEIS-A076823

Array of coefficients of 1/det(M_n)*P(M_n) where P(M_n) is the characteristic polynomial of the n-th n X n Hilbert matrix M_n(i,j)=1/(i+j-1).

Montgomery made a conjecture related to the largest eigenvalue of the Hilbert matrix (cf. Matthews link)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 27 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002431 | OEIS-A076825

Let u(1)=1, u(n)=2^u(n-1) (mod n), sequence gives values of n such that u(n)=1.

Conjecture: a(n)/n^2 -> oo.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 24 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002432 | OEIS-A076847

Ramanujan function tau(p) as p runs through the primes.

Note that the product representation of the Dirichlet series Sum_{n >=1} tau(n)/Sum_{n >= 1} tau(n)/n^s = Prod_{n >= 1} 1/(1 - a(n)/prime(n)^s + prime(n)^(11) / prime(n)^(2*s)) (see the Mordell reference, eq. (2)) leads also to this formula for tau(p^k) for primes p after expanding the factors of the product and collecting powers of 1/p^(k*s). If one insists on convergence of the product one can use s >= 7, if one uses Ramanujan's 1916 conjecture (proved by P. Deligne 1974) |tau(p)| <= 2*p^(11/2), i.e., |a(n)| <= 2*sqrt(prime(n)^11).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 23 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002433 | OEIS-A076868

a(n) = smallest m such that a spherical n-design with m points exists in three dimensions.

Most of these values are just conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 25 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002434 | OEIS-A076869

Largest t such that a spherical t-design with n points exists in four dimensions.

Most of these values are just conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 25 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002435 | OEIS-A076870

a(n) = smallest m such that a spherical n-design with m points exists in four dimensions.

Most of these values are just conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 25 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002436 | OEIS-A076874

a(n) = n - floor ((4*n + 1)^(1/2)).

Conjecture: For n>=7, a(n)-2 is the maximum number of steps in a 2D self-avoiding random walk trapped after n steps having only 2 choices for the next step. a(n) >= A077484(n) + 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Nov 24 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002437 | OEIS-A076897

a(1)=1, a(n)=n-a(floor(3n/4)).

Conjecture: a(n) is asymptotic to 4n/7; (a(n)-4n/7)/log(n) is bounded. Conjecture is true: in fact if D(n) = a(n) - 4*n/7, D(n) = (3*n-4*m)/7-D(m) where m = floor(3*n/4), so |D(n)| <= 4/7 + |D(m)|. Thus D(n) <= 3/7 + 4/7 log_(4/3)(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 26 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002438 | OEIS-A076902

a(1)=1, a(n) = floor(n/2) - a(floor(n/2)).

a(n) is asymptotic to n/3; conjecture: abs(a(n)-n/3) < log(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 26 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002439 | OEIS-A076905

a(1)=1, a(n)=n-a(n-1)+a(a(n-1))-a(a(a(n-1))).

Conjecture: let c be the real root of 1-2*x+x^2-x^3, c=0.5698402909980532659113..., then a(n)=floor(cn) +0 /+ 1

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 27 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002440 | OEIS-A076935

a(1)=1, a(n)=n-a(n-1)+a(a(n-1))-a(a(a(n-1)))+a(a(a(a(n-1))))-a(a(a(a(a(n-1))))).

Conjecture: let c be the real root of 1-2*x+x^2-x^3+x^4-x^5, c=0.597497008682517962677071696444 6..., then a(n)=floor(cn) +0 /+ 1

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 27 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002441 | OEIS-A076957

Smallest k such that there are exactly n primes strictly between k^2 and (k+1)^2.

Conjecturally, a(n) is undefined for n = 1 and defined for all n >= 2. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Oct 20 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002442 | OEIS-A076964

a(1) = 3, then a(n) = smallest multiple of the previous term with as many divisors.

If the first conjecture in A262981 from _Vladimir Letsko_ is true, then this is also: a(2) = 9, thereafter a(n+1) = least k with a(n) divisors. - _Jianing Song_, May 31 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Oct 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002443 | OEIS-A076973

Starting with 2, largest prime divisor of the sum of all previous terms.

Conjecture: start from any initial value a(1) = m >= 2 and define a(n) to be the largest prime factor of a(1)+a(2)+...+a(n-1); then a(n) = n/2 + O(log(n)) and there are infinitely many primes p such that a(2p)=p. - _Benoit Cloitre_, Jun 04 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Oct 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002444 | OEIS-A076980

Leyland numbers: 3, together with numbers expressible as n^k + k^n nontrivially, i.e., n,k > 1 (to avoid n = (n-1)^1 + 1^(n-1)).

Conjecture: For d > 11, 10^(d-1)+(d-1)^10 is the smallest (base ten) d-digit term. - _Hans Havermann_, May 21 2018 Conjecture from _Zhi-Wei Sun_, Feb 26 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Oct 23 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002445 | OEIS-A076984

Number of Fibonacci numbers that are divisors of the n-th Fibonacci number.

a(n) = tau(n) - [n is even] = A000005(n) - A059841(n). Proof: gcd(Fib(m), Fib(n)) = Fib(gcd(m, n)) and Fib(2) = 1. - Olivier Wittenberg, following a conjecture of _Ralf Stephan_, Sep 28 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Oct 25 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002446 | OEIS-A077026

a(n) = Sum_{k=1..n} floor(n/k + 1)-floor(n/k + 1/2).

Conjecture: Let f(a,b)=1, if (a+b) mod |a-b| != (a mod |a-b|)+(b mod |a-b|), and 0 otherwise. a(n) = Sum_{k=1..n-1} 1-f(n,k). - _Benedict W. J. Irwin_, Sep 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002447 | OEIS-A077038

Least difference of primes p, q such that p < n^3 < q.

There are numbers k other than 2 such that a(k) = 4. The first few (up to 1000) are 129 189 369 435 549 555 561 819. Conjecture: every even integer greater than 2 occurs infinitely often in this sequence. - _Franklin T. Adams-Watters_, May 13 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Oct 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002448 | OEIS-A077071

Row sums of A077070.

Conjecture: given n balls, all of which are initially in the first of n numbered boxes, a(n-1) is the number of steps of the following process required to move them all to the last box. A step consists of first identifying j, the lowest numbered box which has at least one ball. If it has only one ball then move it to box j+1; otherwise move half its balls rounded down to box j+1 and (unless it's the first box) half its balls rounded down to box j-1. See also A356254. - _Mikhail Kurkov_, Nov 24 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Oct 25 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002449 | OEIS-A077085

Initial values such that if A077080(x)=phi(sigma(x)+phi(x)) is started at these numbers then the sequence does not converge.

These terms are only conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Oct 28 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002450 | OEIS-A077128

Smallest number greater than the previous term which is relatively prime to each of the group of the next n numbers.

Conjecture : every member is a prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Oct 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002451 | OEIS-A077183

Smallest number k such that the reverse concatenation of natural numbers from k to 1 is divisible by prime(n), or 0 if no such number exists.

Conjecture: a(n) > 0 for all n > 3, since prime(1) = 2 and prime(3) = 5 are the only primes whose multiples cannot end in 1. - _Ryan Propper_, Jul 29 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 01 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002452 | OEIS-A077199

Smallest k such that both k and k+n are squarefree.

Conjecture: every term is < 10, i.e. for every n at least one of the numbers n+2, n+3, n+5, n+6 or n+7 is squarefree. The conjecture is false. Here are 9 counterexamples, each of which is less than 10000: 1857, 2522, 3570, 4470, 6169, 6645, 7981, 9553, 9745. There are 16 counterexamples within the first 10000 squarefree numbers. - _Harvey P. Dale_, May 24 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 01 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002453 | OEIS-A077202

a(1) = 1, a(n) = smallest number such that the concatenation of two successive terms gives a prime which has not occurred earlier.

Conjecture: Every odd prime occurs in this sequence infinitely many times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 02 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002454 | OEIS-A077214

a(1) = 1. a(n+1) is the smallest number which is greater than a(n), coprime to it and has a different prime signature.

Subsidiary sequence: First occurrence of a difference n between successive terms. Conjecture : the maximum value of a(k+1) - a(k) = 7 that too occurs only once for k = 2.(i.e. the subsidiary sequence has only 7 terms.) I conjecture that a(n+1)-a(n) is unbounded. The simplest way to get such a gap is to find n divisible by lcm(1,2,...,k) where n+1 has the same prime signature as n. I expect that such numbers exist, although they are likely to be quite large. (Since n-1 lacks small divisors, it is very likely to be in the sequence and so n is also very likely to be in the sequence.) Note that the sequence includes every prime except 3, 5 and 7. - _Franklin T. Adams-Watters_, Jul 18 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 02 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002455 | OEIS-A077220

a(1) = 1; thereafter a(n) is smallest number not occurring earlier such that a(n-1)+a(n) is a triangular number.

Conjectured to be a permutation of the natural numbers (cf. A099130). The first few fixed points are: 1, 2, 19, 92, 220, 467, 556, 616, 690, 842.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 03 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002456 | OEIS-A077223

a(1) = 1 and then alternately even and odd numbers not occurring earlier such that the sum of any two successive terms is a squarefree number.

Conjecture: There are infinitely many strings of 8 terms of the form k, k+1, k+2, ..., k+7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 03 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002457 | OEIS-A077361

Smallest n-digit prime whose external digits as well as internal digits form a prime, or 0 if no such number exists.

Conjecture: no entry is zero for n>2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002458 | OEIS-A077362

Largest n-digit prime whose external digits as well as internal digits form a prime, or 0 if no such number exists.

Conjecture: no entry is zero for n>2. Conjecture: each term after the first two terms ends with 7. - _Harvey P. Dale_, May 26 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002459 | OEIS-A077371

Fibonacci numbers whose internal digits form a Fibonacci number. Equivalently, Fibonacci numbers from which deleting the MSD and LSD leaves a Fibonacci number.

Conjecture: The sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002460 | OEIS-A077373

Fibonacci numbers whose external digits as well as internal digits form a Fibonacci number.

Conjecture: sequence is finite. Is there any term > 89 in this sequence?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002461 | OEIS-A077391

Smallest n-digit prime which leaves a prime at every step if most significant digit and least significant digit are deleted until a one digit or two digit prime is obtained, or 0 if no such prime exists.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 07 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002462 | OEIS-A077395

Lesser of two successive squarefree numbers whose product is not squarefree.

Conjecture: lim_{n->oo} n/a(n) > 0. - _Thomas Ordowski_ and _Robert Israel_, Aug 18 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Nov 04 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002463 | OEIS-A077402

Reverse and Add! carried out in base 3; number of steps needed to reach a palindrome, or -1 if no palindrome is ever reached.

Base-3 analog of A066057 (base 2), A075685 (base 4) and A033665 (base 10). a(103) = -1 is a conjecture (cf. A066450, A077408). For values of n such that presumably a(n) = -1 see A077404.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002464 | OEIS-A077568

a(1,n) as defined in A003148.

Conjecture: a(n) +a(n-1) -2*(2*n+1)*(n-1)*a(n-2)=0. - _R. J. Mathar_, May 26 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jul 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002465 | OEIS-A077594

Smallest number whose Reverse and Add! trajectory (presumably) contains exactly n palindromes, or -1 if there is no such number.

Conjecture 1: For each k > 0 the trajectory of k eventually leads to a term in the trajectory of some j which belongs to A063048, i.e. whose trajectory (presumably) never leads to a palindrome. Conjecture 2: There is no k > 0 such that the trajectory of k contains more than twelve palindromes, i.e. a(n) = -1 for n > 12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 08 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002466 | OEIS-A077602

Decimal expansion of lim_{n->inf} M(n,1)/2^n, where M(n,1) is the sum of the coefficients of the n-th Moebius polynomial (cf. A074587).

Conjecture: M(n,1) ~ A077596(n) * sqrt(Pi*n/2), where A077596(n) is the largest coefficient of the n-th Moebius polynomial, M(n,x).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_ and _Paul D. Hanna_, Nov 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002467 | OEIS-A077627

Largest term in periodic part of continued fraction expansion of square root of -1+3^n.

a(2*m) = 2*(3^m-1); in general a(n) is close to 2*(3^(n/2)-1) and for any n, 0 <= a(n) - 2*(3^(n/2)-1) < 2. Conjecture: a(n)=ceiling(2*(3^(n/2)-1)) except for n=3, 9, 27 and all powers of 3, in this case a(n)=1+ceiling(2*(3^(n/2)-1)). - _Benoit Cloitre_, Nov 24 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Nov 13 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002468 | OEIS-A077653

a(1)=1, a(2)=2, a(3)=2, a(n) = abs(a(n-1) - a(n-2) - a(n-3)).

Conjecture : let z(1)=x; z(2)=y; z(3)= z; z(n)=abs(z(n-1)-z(n-2)-z(n-3)) if z(n) is unbounded (i.e. x,y,z are such that z(n) doesn't reach a cycle of length 2), then there are 2 integers n(x,y,z) and w(x,y,z) such that M(n) = floor(sqrt(n+w(x,y,z))) for n>n(,x,y,z) where M(n) = Max (a(k) : 1<=k<=n). As example : w(1,2,2)=9 n(1,2,2)=4; w(1,2,4)=29 n(1,2,4)=4; w(1,2,8)=157 n(1,2,8)=9

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Dec 02 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002469 | OEIS-A077712

a(1) = 1, a(n) = the smallest squarefree number > a(n-1) which contains all the digits of a(n-1).

Conjecture: Terms contain only two types of digits, i.e., 0 and 1. Beginning with a(3), sequence follows a regular pattern: 10^2+1, 10^2+10, 10^3+1, 10^3+10, etc. until at a(21) the pattern is disrupted by 10^11+1, which is not squarefree (see A086982). 10^12+10 is also absent from the sequence since it is also not squarefree. The pattern resumes after this disruption until the next occurrence of 10^k+1 which is not squarefree, k=21, 33, 39, 55, ... The conjecture that the sequence is composed of terms containing only the digits 0 and 1 is certainly true up to 10^406+1 where both it and 10^407+1 are not squarefree. Indeed beginning with a(3) the terms contain exactly two 1 digits and the rest 0's up to this point. The term following 10^406+10 will introduce a third nonzero digit, perhaps a 1, but the pattern of the sequence changes dramatically at this point. - _Ray Chandler_, Aug 02 2003 Term following a(739)=10^406+10 is a(740)=10^407+11 so the conjecture is still in play. - _Ray Chandler_, Aug 05 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002470 | OEIS-A077713

a(1) = 3; thereafter a(n) = the smallest prime of the form d0...0a(n-1), where d is a single digit, or 0 if no such prime exists.

a(n) is the smallest prime obtained by prefixing a(n-1) with a number of the form d*10^k where d is a single digit, 0 < d < 10, and k >= 0. Conjecture: d*10^k always exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002471 | OEIS-A077739

Smallest multiple of n beginning and ending in n and having a digit sum of n, or 0 if no such number exists.

Conjecture: a(n) = 0 if and only if n is 11, 17 or 19.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 20 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002472 | OEIS-A077754

Smallest multiple of n with two or more digits, none of them zeros, whose digit sum equals n, or 0 if no such multiple exists.

Conjecture: zero occurs only for indices which are multiples of 10 apart from 1, 2, 5 and 11. (a(n) = 0 only for n = 1,2,5,11 or n = 10k.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 20 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002473 | OEIS-A077761

Decimal expansion of Mertens's constant, which is the limit of (Sum_{i=1..k} 1/prime(i)) - log(log(prime(k))) as k goes to infinity, where prime(i) is the i-th prime number.

Equals lim_{k->oo} -k + Sum_{p prime} 1/(p*log(p)^(1/k)) conjectured by Meissel in 1866 and proven by Peter Lindqvist and Jaak Peetre in 1997 see links - _Artur Jasinski_, Mar 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Nov 14 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002474 | OEIS-A077800

List of twin primes {p, p+2}.

Sum_{n>=1} 1/a(n) is in the interval (1.840503, 2.288490) (Platt and Trudgian, 2020). The conjectured value based on assumptions about the distribution of twin primes is A065421. - _Amiram Eldar_, Oct 15 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 03 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002475 | OEIS-A077864

Expansion of (1-x)^(-1)/(1-x-2*x^2-x^3).

Conjecture: a(n) = Sum_{j=0..n/2} A027907(n+1-j,2*j+1), n >= 0. - _Werner Schulte_, Sep 29 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 17 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002476 | OEIS-A077866

Expansion of (1-x)^(-1)/(1 - x - 2*x^2 + 2*x^3).

Conjecture: let b(n) be the number of subsets S of {1,2,...,n} having more than one element such that (sum of least two elements of S) = max(S). Then b(0) = b(1) = b(2) = 0 and b(n+3) = a(n) for n >= 0. - _Clark Kimberling_ Sep 27 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 17 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002477 | OEIS-A078008

Expansion of (1 - x)/((1 + x)*(1 - 2*x)).

Conjecture: a(n) = the number of fractions in the infinite Farey row of 2^n terms with even denominators. Compare the Salamin & Gosper item in the Beeler et al. link. - _Gary W. Adamson_, Oct 27 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 17 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002478 | OEIS-A078111

a(n) = floor((n+2)^(n+2)/n^n).

a(n) - e^2(n+1)^2 is bounded. Conjecture: there's a constant c = 2.462... such that a(n) = floor(e^2*(n+1)^2-c). I believe (n+2)^(n+2)/n^n = e^2(n^2 + 2n + 2/3 + o(1)), so the conjecture should be correct (with perhaps finitely many exceptions) with c = e^2/3 = 2.463.... - _Charles R Greathouse IV_, May 07 2013 Greathouse IV's conjecture is true (see Fried link). - _Sela Fried_, Mar 27 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Dec 03 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002479 | OEIS-A078113

Let u(1)=u(2)=1, u(3)=n, u(k) = (1/2)*abs(2*u(k-1) -u(k-2)-u(k-3)); sequence gives values of n such that Sum_{k>=1} u(k) is an integer.

Conjecture: a(n) = -3+2^(1/2*(-5+n))*(10-10*(-1)^n+9*sqrt(2)+9*(-1)^n*sqrt(2)). a(n) = a(n-1)+2*a(n-2)-2*a(n-3). G.f.: x*(3*x^2-4*x-2) / ((x-1)*(2*x^2-1)). - _Colin Barker_, Aug 14 2013 Conjecture: a(n) = 2*a(n-2) + 3, n odd>2 = A154117((n+1)/2). - _Bill McEachen_, Jun 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Dec 04 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002480 | OEIS-A078121

Infinite lower triangular matrix, M, that satisfies [M^2](i,j) = M(i+1,j+1) for all i,j>=0 where [M^n](i,j) denotes the element at row i, column j, of the n-th power of matrix M, with M(0,k)=1 and M(k,k)=1 for all k>=0.

M(n,k) = the coefficient of x^(2^n - 2^(n-k)) in the power series expansion of 1/Product_{j=0..n-k} (1-x^(2^j)) whenever 0<=k<n for all n>0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002481 | OEIS-A078122

Infinite lower triangular matrix, M, that satisfies [M^3](i,j) = M(i+1,j+1) for all i,j>=0 where [M^n](i,j) denotes the element at row i, column j, of the n-th power of matrix M, with M(0,k)=1 and M(k,k)=1 for all k>=0.

Conjecture: the sum of the n-th row equals the number of partitions of 3^n into powers of 3 (A078125). M(n, k) = the coefficient of x^(3^n - 3^(n-k)) in the power series expansion of 1/Product_{j=0..n-k}(1-x^(3^j)) whenever 0<=k<n for all n>0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002482 | OEIS-A078124

Second column, M(n+1,1) for n>=0, of infinite lower triangular matrix M defined in A078122.

The partitions of 2*3^n into powers of 3, or, the coefficient of x^(2*3^n) in 1/Product_{j=0..inf}(1-x^(3^j)) (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002483 | OEIS-A078128

Number of ways to write n as sum of cubes > 1.

Conjecture (lower bound): for all k exists b(k) such that a(n)>k for n>b(k); see b(0)=A078129(83)=154 and b(1)=A078130(63)=218.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Nov 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002484 | OEIS-A078130

Numbers having exactly one representation as sum of cubes > 1.

Conjecture: the sequence is finite; is a(63)=218 the last entry?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Nov 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002485 | OEIS-A078134

Number of ways to write n as sum of squares > 1.

Conjecture (lower bound): for all k exists b(k) such that a(n)>k for n>b(k); see b(0)=A078135(12)=23 and b(1)=A078136(15)=39. This is true - see comments by _Hieronymus Fischer_. The author's conjecture holds true. Proof by induction: b(0) exists; if b(k) exists, then a(j)>k for all j>b(k). Setting m:=b(k)+1, we find that there are k+1 sums B(0,i) of squares >1, 1<=i<=k+1, with m=B(0,i). Further there are k+1 such sum expressions B(1,i), B(2,i) and B(3,i), 1<=i<=k+1, representing m+1, m+2 and m+3, respectively. For n>b(k) we have n=m+4*floor((n-m)/4)+(n-m) mod 4. A more precise formulation of the author's conjecture is "b(k):=min(n | a(j)>k for all j>n) exists for all k>=0". (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Nov 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002486 | OEIS-A078173

a(n) = max(min(r,s,t)), where the maximum is taken over all positive integer solutions of the Diophantine equation r*s*t = n * (r+s+t).

Conjecture: max(t) = n*(n+2) and max(s) = 2*n, where the maxima are taken over all positive integer solutions r<=s<=t of the above Diophantine equation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Nov 20 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002487 | OEIS-A078181

a(n) = Sum_{d|n, d == 1 (mod 3)} d.

Conjecture. If a(n)=n+1 then n==1 (mod 3). (Is this easy to settle? It has been verified for n=1,2,3,...,2000.) - _John W. Layman_, Apr 03 2006 The conjecture is false. The first and only counterexample below 10^8 is a(6800) = 6801 and 6800 == 2 (mod 3). - Lambert Herrgesell (zero815(AT)googlemail.com), May 06 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Nov 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002488 | OEIS-A078251

a(n) = smallest multiple of the n-th prime whose decimal expansion is nnn...n, or 0 if no such number exists.

Conjecture: No entry is zero for n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 24 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002489 | OEIS-A078257

a(n) = denominator(N) where N = 0.123...n (concatenation of 1 to n after decimal point).

Conjecture: sequence is not equal to the sequence of denominators presented in A172495 and A172506. - _Jaroslav Krizek_, Feb 05 2010 The conjecture is false for both other sequences; see A172495 and A172506 for proofs. - _Michael S. Branicky_, Nov 30 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 24 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002490 | OEIS-A078334

Primes in A005728, which counts the terms in the Farey sequence of order n.

Guy, in his Example 8, citing Leo Moser as his source, noted that the first 9 values of A005728(n) = 1 + Sum_{i=1..n} phi(i) = 1 + Sum_{i=1..n} A000010(i) are all primes, but that the pattern breaks down at A005728(10) = 33 = 3*11. As Guy warns, in several paraphrases of the same law, "Capricious coincidences cause careless conjectures." That is, for 1 <= n <= 9 we have A005728(n) = A078334(n), but for n > 9 we sometimes (n = {11, 12, 13, 15, 17, 18, 22, ...}) have A005728(n) prime, but other times (n = {10, 14, 16, 19, 20, 21, ...}) have A005728(n) composite. [_Jonathan Vos Post_, Sep 06 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Nov 21 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002491 | OEIS-A078345

Numbers k such that F(k) mod k divides F(F(k) mod k) where F(k) denotes the k-th Fibonacci number.

Conjecture: a(n) is asymptotic to c*n*log(n) with c>0.7

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002492 | OEIS-A078414

a(n) = (a(n-1)+a(n-2))/7^k, where 7^k is the highest power of 7 dividing a(n-1)+a(n-2).

An open question: Is this sequence bounded?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yasutoshi Kohmoto_, Dec 28 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002493 | OEIS-A078437

Decimal expansion of sum of alternating series of reciprocals of primes.

Conjectured values a(14)-a(18) = 4,5,9,4,8

猜想/问题提及段 | historical_or_current_mention_unreviewed | _G. L. Honaker, Jr._, Dec 31 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002494 | OEIS-A078441

a(n) begins the first chain of n consecutive positive integers that have equal h-values, where h(k) is the length of the finite sequence k, f(k), f(f(k)), ..., 1 in the Collatz (or 3x + 1) problem. (The earliest "1" is meant.)

Garner (1985) conjectured that there are arbitrarily long runs of consecutive integers with the same height. As recalled by Gao (1993), the runs should have the same height, computed with f as here, and the same number of halving steps (A006666); in that form the conjecture implies that a(n) exists for every n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Dec 31 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002495 | OEIS-A078466

First cycle is reached under a "Collatz-Fibonacci" transform (cf. comment).

Conjecture : for any n x(k) reaches a cycle among 2 cycles : (1;1;0) and (1;2;3;5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jan 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002496 | OEIS-A078474

a(1)=a(2)=a(3)=1, a(n)=n-a(a(n-1))-a(a(n-2))-a(a(n-3)).

Conjecture: partial sums of A284680 (verified for n <= 100000). - _Sean A. Irvine_, Jul 16 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jan 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002497 | OEIS-A078489

a(n)=j such that binomial(n,j)<binomial(n-1,j-2).

Conjecture: a(n) = A003258(n) - n + 1. - _Ralf Stephan_, Feb 24 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Jan 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002498 | OEIS-A078506

Decimal expansion of sum of inverses of unrestricted partition function.

Conjecture: this is a transcendental number. - _Zhi-Wei Sun_, May 24 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Jan 05 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002499 | OEIS-A078508

Number of primes between sqrt(n^3) and sqrt((n+1)^3).

Conjecture 1: for n>1051, a prime always exists between n^1.5 and (n+1)^1.5. Conjecture 2: for n>7295, more than 2 primes always exist between n^1.5 and (n+1)^1.5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jan 05 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002500 | OEIS-A078522

Numbers k such that (k+1)*(2*k+1) is a perfect square.

The sequence of areas k(n)*q(n)/2, of the ordered Pythagorean triples (k(n), q(n) = k(n) + 2, c(n)) with k(1)=0, q(1)=2, c(1)=0, a(1)=0, and k(2)=6, q(2)=8, c(2)=10, a(2)=24 (conjectured). Conjecture: let f(n) be a sequence of form x(n)*y(n)/2, of the ordered Pythagorean triples (x(n), y(n) = x(n) + v, z(n)) with x(1)=0, y(1)=v, z(1)=0, f(1)=0, where v is an even number. Then there exists such subset p(i) that p(1) = 0, p(2) = 24*(v/2)^2, for any i > 2, p(i) = 34*p(i-1) - p(i-2) + 24*(v/2)^2, and any p(i) is a term of the above sequence f(n) (see also the first formula by _Benoit Cloitre_ in the Formula section).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Jan 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002501 | OEIS-A078527

Number of maximally 2-constrained walks on square lattice trapped after n steps.

In a 2D self-avoiding walk there may be steps, where the number of free target positions is less than 3. A step is called k-constrained, if only k<3 neighbors were not visited before. Self-trapping occurs at step n (the next step would have k=0). A maximally 2-constrained n-step walk contains n-floor((4*n+1)^(1/2))-2 steps with k=2 (conjectured). The first step is chosen fixed (0,0)->(1,0), all other steps have k=3. This sequence counts those walks among all possible self-trapping n-step walks A077482(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Nov 27 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002502 | OEIS-A078532

Coefficients of power series that satisfies A(x)^3 - 9*x*A(x)^4 = 1, A(0)=1.

If A(x) = Sum_{k>=1} a(k)x^k satisfies A(x)^n - (n^2)*x*A(x)^(n+1) = 1, then a(n-1) = n^(2n-3) and a(2n-1) = n^(4n-2) (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 28 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002503 | OEIS-A078533

Coefficients of power series that satisfies A(x)^4 - 16x*A(x)^5 = 1, A(0)=1.

If A(x) = Sum_{k>=1} a(k)x^k satisfies A(x)^n - (n^2)*x*A(x)^(n+1) = 1, then a(n-1) = n^(2n-3) and a(2n-1) = n^(4n-2) (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 28 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002504 | OEIS-A078534

Coefficients of power series that satisfies A(x)^5 - 25*x*A(x)^6 = 1, A(0)=1.

If A(x) = Sum_{k>=1} a(k)*x^k satisfies A(x)^n - (n^2)*x*A(x)^(n+1) = 1, then a(n-1) = n^(2*n-3) and a(2*n-1) = n^(4*n-2) (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 28 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002505 | OEIS-A078535

Coefficients of power series that satisfies A(x)^6 - 36x*A(x)^7 = 1, A(0)=1.

If A(x)=sum_{k=1..inf} a(k)x^k satisfies A(x)^n - (n^2)*x*A(x)^(n+1) = 1, then a(n-1) = n^(2n-3) and a(2n-1) = n^(4n-2) (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 28 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002506 | OEIS-A078536

Infinite lower triangular matrix, M, that satisfies [M^4](i,j) = M(i+1,j+1) for all i,j>=0 where [M^n](i,j) denotes the element at row i, column j, of the n-th power of matrix M, with M(0,k)=1 and M(k,k)=1 for all k>=0.

M also satisfies: [M^(4k)](i,j) = [M^k](i+1,j+1) for all i,j,k>=0; thus [M^(4^n)](i,j) = M(i+n,j+n) for all n>=0. Conjecture: sum of the n-th row equals the partitions of 4^n into powers of 4. M(n, k) = the coefficient of x^(4^n - 4^(n-k)) in the power series expansion of 1/Product_{j=0..n-k}(1-x^(4^j)) whenever 0<=k<n for all n>0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002507 | OEIS-A078537

Number of partitions of 4^n into powers of 4 (without regard to order).

Conjecture: a(n) = sum of the n-th row of lower triangular matrix A078536.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002508 | OEIS-A078563

a(n) is the least positive integer k such that g(k) = n*g(k-1), where g(k) = prime(k+1) - prime(k).

Conjecture: The equation g(k) = n*g(k-1), for a fixed positive integer n, is always solvable for k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Jan 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002509 | OEIS-A078605

Sum of square displacements over all self-avoiding n-step walks on the cubic lattice with the first step specified. Numerator of mean square displacement s(n)=a(n)/(A001412(n)/6).

A comparison with the conjectured asymptotic behavior of the mean square displacement s(n) over all n-step self-avoiding walks given in Weisstein's article is shown in "Asymptotic Behavior of Mean Square Displacement" at the Pfoertner link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 09 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002510 | OEIS-A078611

Radius of the shortest interval (of positive length) centered at prime(n) that has prime endpoints.

a(1) and a(2) are undefined. Alternatively, a(n) = least k, 1 < k < n, such that prime(n) + k and prime(n) - k are both prime. I conjecture that a(n) is defined for all n > 2. Equivalently, every prime > 3 is the average of two distinct primes. a(n) embodies the difference between weak and strong Goldbach conjectures, and therefore between A047160 and A082467 which differ only for prime arguments (a(n)=A082467(prime(n)), while A047160(prime(n))=0). - _Stanislav Sykora_, Mar 14 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Dec 09 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002511 | OEIS-A078613

Same numbers of distinct prime factors of forms 4*k+1 and 4*k+3.

Conjecture: a(n) is asymptotic to c*n where c is around 4. - _Benoit Cloitre_, Jan 06 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Dec 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002512 | OEIS-A078615

a(n) = rad(n)^2, where rad is the squarefree kernel of n (A007947).

It is conjectured that only 1 and 1782 satisfy a(k) = sigma(k). See Broughan link. - _Michel Marcus_, Feb 28 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Dec 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002513 | OEIS-A078616

a(n) = Sum_{k=0..n} A010815(k).

a(n) = (Number of ones in the distinct partitions of n with an odd number of parts) - (number of ones in the distinct partitions of n with an even number of parts) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Dec 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002514 | OEIS-A078623

Number of matched parentheses and brackets of length n, where a closing bracket will close any remaining open parentheses back to the matching open bracket (as in some versions of LISP).

Conjecturally, a(n-1) = (-1)^(n-1)*(1/n)*Sum_{k=1..n} binomial(n,k)*binomial(n - 2*k,k - 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Brian T. Howard (bhoward(AT)depauw.edu), Dec 11 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002515 | OEIS-A078642

Numbers with two representations as the sum of two Fibonacci numbers.

A positive integer n has exactly two representations as the sum of two Fibonacci numbers if and only if n is twice a Fibonacci number and n >= 4. Conjectured by _John W. Layman_, Dec 20 2002. Proved by Max Alekseyev, Mar 02 2007.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Dec 12 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002516 | OEIS-A078671

Number of times the n-th prime appears among the decimal digits of 2^(2^n) + 1, the Fermat numbers.

Conjectures: Is a(n) monotonically increasing for n > 4? Does lim{n->inf} a(n)/a(n+1) = 0.5? - _Ryan Propper_, Jan 04 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Dec 16 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002517 | OEIS-A078693

Largest k such that round(1/(sqrt(prime(k+1))-sqrt(prime(k)))) = n where prime(n) denotes the n-th prime (conjectured values).

Largest k such that round(1/(sqrt(prime(k+1))-sqrt(prime(k)))) = n where prime(n) denotes the n-th prime (conjectured values).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Dec 17 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002518 | OEIS-A078698

Number of ways to lace a shoe that has n pairs of eyelets such that each eyelet has at least one direct connection to the opposite side.

Conjecture: a(n) = (n-1)!^2*A051286(n). - _Vladeta Jovovic_, Sep 14 2005 (correct, see the Khrabrov/Kokhas reference, _Joerg Arndt_, May 26 2015)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002519 | OEIS-A078700

Number of symmetric ways to lace a shoe that has n pairs of eyelets such that each eyelet has at least one direct connection to the opposite side.

a(n) = (n-1)!*Fibonacci(n+1) = A000142(n-1)*A000045(n+1). - Conjectured by _Vladeta Jovovic_, Jan 23 2005, proved by _Antti Karttunen_, Jan 06 2007. Proof of Jovovic's conjecture from _Antti Karttunen_, Jan 06 2007: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002520 | OEIS-A078703

Number of ways of subtracting twice a triangular number from a perfect square to obtain the integer n.

G.f.: Sum_{n >= 1} x^(n^2)/(1-x^(2*n-1)) (conjecture). - _Joerg Arndt_, Jan 04 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | R. L. Coffman, K. W. McLaughlin and R. J. Dawson (robert.l.coffman(AT)uwrf.edu), Dec 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002521 | OEIS-A078712

Series expansion of (-3 - 2*x)/(1 + x - x^3) in powers of x.

Similar to the Perrin sequence A001608, I conjecture that if p is a prime then a(p) == 1 (mod p). This implies that A001945(n) == 1 (mod p) and A001608(2*n) == 2 (mod p). - _Michael Somos_, Dec 25 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Dec 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002522 | OEIS-A078720

Integer part of the ratio of even to odd terms among n, f(n), f(f(n)), ..., 1 for the Collatz function (that is, until reaching "1" for the first time), or -1 if 1 is never reached.

1. The Collatz function (related to the "3x+1 problem") is defined by: f(n) = n/2 if n is even; f(n) = 3n + 1 if n is odd. A famous conjecture states that n, f(n), f(f(n)), ... eventually reaches 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Dec 20 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002523 | OEIS-A078754

Numbers k such that for all m>k there are more than n primes between m and 2m (inclusive).

The terms shown are only conjectured values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jan 08 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002524 | OEIS-A078777

a(n) = the least positive integer k such that binomial(2k,k) + k + n is prime, if such k exists; = 0, otherwise.

The value a(1) = 0 is conjectural. There is no k < 8.5 x 10^3 such that binomial(2k,k)+k+1 is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Jan 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002525 | OEIS-A078786

Period of cycle of the inventory sequence (as in A063850) starting with n.

It can be proved that any inventory sequence ends in a cycle all of whose terms are <= 10^20. Conjecture: a(n) <= 4 for all n. It suffices to check this for all inventory sequences starting with n, where n <= 10^20.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Jan 14 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002526 | OEIS-A078797

Sum of square displacements over all self-avoiding n-step walks on a square lattice with the first step specified. Numerator of mean square displacement s(n)=a(n)/A046661(n).

A comparison with the conjectured asymptotic behavior of the mean square displacement s(n) over all n-step self-avoiding walks given in E. Weisstein's MathWorld article is shown in the "Asymptotic Behavior of Mean Square Displacement" link. [I'm not sure this comment is correct. There may be some confusion with A176177. - _N. J. A. Sloane_, Aug 02 2015]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 05 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002527 | OEIS-A078798

Sum of Manhattan distances over all self-avoiding n-step walks on square lattice. Numerator of mean Manhattan displacement s(n) = a(n)/A046661(n).

A conjectured asymptotic behavior for the mean Manhattan displacement lim n-> infinity a(n)/(A046661(n)*n^(3/4)) = constant is illustrated in "Asymptotic Behavior of Mean Manhattan Displacement" at first link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002528 | OEIS-A078816

Erroneous version of A081846.

Conjecture : a(n) is asymptotic to c*n with 6<c<7

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Dec 06 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002529 | OEIS-A078841

Main diagonal of the table of k-almost primes (A078840): a(n) = (n+1)-st integer that is an n-almost prime.

Conjecture: Lim as n->inf. of a(n+1)/a(n) = 2. - _Robert G. Wilson v_, Nov 13 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_ and _Paul D. Hanna_, Dec 10 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002530 | OEIS-A078869

Number of n-tuples with elements in {2,4,6} which can occur as the differences between n+1 consecutive primes > n+1. (Values of a(11), ..., a(18) are conjectured to be correct, but are only known to be upper bounds.)

Number of n-tuples with elements in {2,4,6} which can occur as the differences between n+1 consecutive primes > n+1. (Values of a(11), ..., a(18) are conjectured to be correct, but are only known to be upper bounds.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Dec 19 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002531 | OEIS-A078900

Generalized Fermat numbers of the form (k+1)^2^m + k^2^m, with m>0.

It can be shown that, like the Fermat numbers, two of these generalized Fermat numbers are coprime if they have the same base k. However, unlike the Fermat numbers (which are conjectured to be squarefree), these generalized Fermat numbers are not necessarily squarefree for k > 1. Riesel tabulates some prime factors of generalized Fermat numbers for k <= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Dec 12 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002532 | OEIS-A078902

Generalized Fermat primes of the form (k+1)^2^m + k^2^m, with m>1.

For k=1, these are the Fermat primes A019434. Is the set of generalized Fermat primes infinite? Conjecture that there are only a finite number of generalized Fermat primes for

each value of k. See A077659, which shows that in cases such as k=11, there appear to be no primes. See A078901 for generalized Fermat numbers. See A080131 for the conjectured number of primes for each k. See A080208 for the least k such that (k+1)^2^n + k^2^n is prime. The largest probable prime of this form discovered to date is the 10217-digit 312^2^12 + 311^2^12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Dec 12 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002533 | OEIS-A078941

Flipping burnt pancakes. Maximum number of spatula flips to sort a stack of n pancakes of different sizes, each burnt on one side, so that the smallest ends up on top, ..., the largest at the bottom and each has its burnt side down.

It is conjectured that the initial configuration in which the pancakes are in the correct order but all of the burnt sides are up is a worst case for the problem. If so, then this sequence is identical to A078942. The conjecture above is false, as Cibulka (2009, 2011) proves that a(15) = 25 whereas A078942(15) = 24. - _Pontus von Brömssen_, Aug 04 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, Dec 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002534 | OEIS-A078942

Flipping burnt pancakes. Given a sorted stack of n burnt pancakes of different sizes (smallest on top, ..., largest at the bottom), each with its burnt side up, a(n) is the number of spatula flips needed to restore them to their initial order but with the burnt sides down.

The conjecture that this initial configuration is a worst case for the general problem of burnt pancakes has been disproved for n=15 by Cibulka, 2011.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, Dec 18 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002535 | OEIS-A078943

a(1)=1; a(n+1) is either a(n)-n or a(n)+n, where we choose the smallest positive integer that's not among the values a(1), ..., a(n).

A356080 is targeted to be such a sequence, but starting from 0. Its definition incorporates a limited look-ahead condition that is clearly a necessary condition for the sequence not to encounter a dead end (i.e., be finite) and is conjectured to be a sufficient condition. - _Peter Munn_, Feb 09 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Dec 15 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002536 | OEIS-A078999

Coefficients A_n for the s=4 tennis ball problem.

Conjecture D-finite with recurrence -729*(3*n+2)*(447758283*n-407746117) *(3*n+4) *(n+1)*a(n) +216*(182049960672*n^4 +605681769096*n^3 -358290749358*n^2 -265170598015*n -38328134998)*a(n-1) +1536 *(30350980224*n^4 -947048676672*n^3 +1377152586736*n^2 -569141632910*n +54868443093)*a(n-2) -131072*(4*n-5) *(351198196*n -151260957) *(4*n-7) *(2*n-3)*a(n-3)=0. - _R. J. Mathar_, Mar 31 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002537 | OEIS-A079006

Expansion of q^(-1/4) * (eta(q) * eta(q^4)^2 / eta(q^2)^3)^2 in powers of q.

a(n) = A208933(4*n+1) - A215348(4*n+1) (conjectured). - _Thomas Baruchel_, May 14 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Dec 22 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002538 | OEIS-A079034

Determinant of M(n), the n X n matrix defined by m(i,i) = 1, m(i,j) = i-j.

In 2022, Han Wang and Zhi-Wei Sun provided a proof of the formula a(n) = 1 + n^2*(n^2-1)/12 via eigenvalues. See A355175 for my conjecture on det[(i-j)^2+d(i,j)]_{1<=i,j<=n}, where d(i,j) is 1 or 0 according as i = j or not. - _Zhi-Wei Sun_, Jun 28 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002539 | OEIS-A079044

Numbers k such that Sum_{j=0..k} sin(j/Pi) < 0.

Conjecture: a(n) = floor(2 * Pi^2 * n), checked for n <= 10^4. - _Vincenzo Librandi_, Sep 03 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002540 | OEIS-A079047

Number of primes between prime(n) and prime(n)^2.

I conjecture that 25 and 64 are the only terms that are also square numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jose R. Brox (tautocrona(AT)terra.es), Feb 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002541 | OEIS-A079051

Recamán variation: a(0) = 0; for n >= 1, a(n) = a(n-1) - f(n) if that number is positive and not already in the sequence, otherwise a(n) = a(n-1) + f(n), where f(n) = floor(sqrt(n)) (A000196).

Conjecture: for n>100, 1/2 < a(n)/(n*log(n)) < 1. The conjecture is false. In fact, a(n) = n^(3/2)/6 + O(n). - _N. J. A. Sloane_, Apr 29 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002542 | OEIS-A079104

Number of permutations of length n containing the minimum number of monotone subsequences of length 4.

Conjectured to equal A079105 (and so have period 3) from a(17) onwards.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph Myers_, Dec 23 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002543 | OEIS-A079128

Number of degree-n permutations with (mutually) relatively prime cycle lengths.

a(p) = p!-(p-1)! for prime p. Conjecture: a(n) is divisible by n^2-1 for n>3. Conjecture: gcd(a(n),n)=1. - _Vladeta Jovovic_, Jan 25 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, _Vladimir Baltic_, Dec 27 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002544 | OEIS-A079144

Number of labeled interval orders on n elements: (2+2)-free posets.

We make the following conjectures: Conjectural form for the o.g.f. as a continued fraction: For a proof of above conjectural continued fraction representation of the o.g.f. see the Bala link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Detlef Pauly (dettodet(AT)yahoo.de), Dec 27 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002545 | OEIS-A079156

Sum of end-to-end Manhattan distances over all self-avoiding n-step walks on cubic lattice. Numerator of mean Manhattan displacement s(n)=a(n)/A078717.

A conjectured asymptotic behavior for the mean Manhattan displacement is shown in a diagram lim n-> infinity a(n)/(A078717(n)*n^nu)=c, for some values of nu near 0.59 at Pfoertner link

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 29 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002546 | OEIS-A079213

Triangle read by rows giving T(n,k) = number of sets of k black squares in an n X n checkerboard with the upper left corner colored black, such that the line joining any 2 squares slopes down to the right, 0 <= k <= n.

G.f. (conjectured): Sum_{n>=0, k>=0} T(n, k) x^n y^k = sqrt((1+x)/((1+x-x*y)((1-x)^2 - x*y*(1+x)))).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, Jan 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002547 | OEIS-A079267

d(n,s) = number of perfect matchings on {1, 2, ..., 2*n} with s short pairs.

Conjecture: Asymptotically, the n-th row has a Poisson distribution with mean 1. - _David Callan_, Nov 11 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jeremy Martin (martin(AT)math.umn.edu), Feb 05 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002548 | OEIS-A079279

a(n)=n for n<=3; for n>3, a(n) is next integer sharing common factors with 1 or 2 of previous 3 terms.

Conjectures from _Colin Barker_, Oct 15 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthew Vandermast_, Feb 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002549 | OEIS-A079291

Squares of Pell numbers.

(Conjectured) For any primitive Pythagorean triple of the form (X, Y, Z=Y+1), it appears that Y or Z will always be (and only be) a square Pell number if X = A001333(n), for n > 1. If n is even, Y is always a square Pell number, and if n is odd, then Z is always a square Pell number. For example: (3, 4, 5), (7, 24, 25), (17, 144, 145), (41, 840, 841), (99, 4900, 4901). - _Jules Beauchamp_, Feb 02 2022 Conjectured formula for (X, Y, Z) for primitive Pythagorean triple of the form (X, Y, Z=Y+1) is (A001333(n)^2, A079291(n)^2, A079291(n)^2-1) or (A001333(n)^2, A079291(n)^2-1, A079291(n)^2). As a closed formula (X, Y, Z) = (((1-sqrt(2))^n + (1+sqrt(2))^n)/2, (((1-sqrt(2))^n + (1+sqrt(2))^n)^2 - 4)/8, (((1-sqrt(2))^n + (1+sqrt(2))^n)^2 + 4)/8). - _Jules Beauchamp_, Feb 02 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Feb 08 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002550 | OEIS-A079321

Composite numbers of the form 1^1 * 2^2 * 3^3 * 4^4 * ... * n^n + 1.

No primes other than 2,5,109 found in this sequence for n <= 1000. Conjecture: There are no primes in the sequence 2^2 * 3^3 * 4^4 * ... *n^n + 1 for n > 3. Conjecture: There are no primes in the sequence 2^2 * 3^3 * 4^4 * ... * n^n + 61 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Feb 12 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002551 | OEIS-A079340

Absolute value of determinant of n X n matrix whose entries are the integers from 1 to n^2 spiraling outward, ending in a corner.

a(n) = (2*n^2-3*n+3) (2n-2)!/(2 (n-1)!) = A096376(n-1)*A000407(n-2), n>1. - Conjectured by _Dean Hickerson_, Jan 30 2003. Proved in the article by Bhatnagar and Krattenthaler.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Kit Vongmahadlek (kit119(AT)yahoo.com), Jan 03 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002552 | OEIS-A079390

a(n) = lpf(n) * lpf(n+1) * lpf(n+2), where lpf(n) = A020639(n) is the least prime factor of n.

For n>2, a(n) is a record value iff (n,n+2) is a twin prime pair (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Feb 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002553 | OEIS-A079402

Number of permutations of n^2 distinct integers free of any monotonic increasing or decreasing (n+1)-subsequence.

Conjecture: this is equal to the number of permutations of n^2 distinct integers free of any monotonic increasing or decreasing (n+1)-subsequence. (By the Erdos-Szekeres Theorem, every permutation of n^2+1 distinct integers has such a subsequence.) - _Joseph Myers_, Jan 04 2003 Claude Lenormand (claude.lenormand(AT)free.fr) confirms this conjecture. - Jan 06, 2002.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, Jan 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002554 | OEIS-A079404

Let G(n) be the set of numbers between 2^(n-1) and 2^n-1, inclusive. There is a unique number m(n) in G(n) so that the denominator of the m(n)-th partial sum of the double harmonic series is divisible by smaller 2-power than that of others in G(n). This power is defined to be a(n).

The sequence is conjectured to go to positive infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jianqiang Zhao (jqz(AT)math.upenn.edu), Jan 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002555 | OEIS-A079411

a(1)=1, a(n) = n - a(a(ceiling(n/2))).

a(n) is asymptotic to n*(sqrt(3)-1); conjecture: (a(n)-n*(sqrt(3)-1))/log(n) is bounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 16 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002556 | OEIS-A079429

a(0) = 2, a(1) = 3, a(2) = 5; a(n) = a(n-1) + [a(n-1)-a(n-2)] * [a(n-2)-a(n-3)].

Conjecture: a(n)=A011455(n-1)+5 where defined. - _R. J. Mathar_, Apr 26 2007 Proof of conjecture: if d(n) = log_2(a(n+1)-a(n)), we have d(0)=0, d(1)=1, d(n)=d(n-1)+d(n-2), so d(n) = Fibonacci(n). - _Robert Israel_, Oct 25 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ajay Chhabra (ajay(AT)cantab.net), Jan 08 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002557 | OEIS-A079438

a(0) = a(1) = 1, a(n) = 2*(floor((n+1)/3) + (if n >= 14) (floor((n-10)/4) + floor((n-14)/8))).

However, _David Callan_'s A123050 claims to give more correct version of that count from n=26 onward, so I probably made a little mistake when converting my insights into the formula given here. At that time I reckoned that if the conjecture given in A080070 were true, then it would imply that the formula given here were exact, otherwise it would give only a lower bound.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002558 | OEIS-A079484

a(n) = (2n-1)!! * (2n+1)!!, where the double factorial is A006882.

In 1881, R. F. Scott posed a conjecture that the absolute value of permanent of square matrix with elements a(i,j)= (x_i - y_j)^(-1), where x_1,...,x_n are roots of x^n=1, while y_1,...,y_n are roots of y^n=-1, equals a((n-1)/2)/2^n, if n>=1 is odd, and 0, if n>=2 is even. After a century (in 1979), the conjecture was proved by H. Minc. - _Vladimir Shevelev_, Dec 01 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jan 17 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002559 | OEIS-A079489

a(n) = (2*4^n*binomial(2*n, n) - binomial(4*n + 1, 2*n)) / (n + 1).

For integer r and positive integer s, define a sequence {u(n) : n >= 0} by setting u(n) = [x^(s*n)] A(x)^(r*n). We conjecture that the supercongruence u(n*p^k) == u(n*p^(k-1)) (mod p^(3*k)) holds for all primes p >= 5 and for all positive integers n and k. Let B(x) = 1/x * series_reversion(x*A(x)). Define a sequence {v(n) : n >= 0} by setting v(n) = [x^(s*n)] B(x)^(r*n). We conjecture that the supercongruence v(n*p^k) == v(n*p^(k-1)) (mod p^(3*k)) holds for all primes p >= 5 and for all positive integers n and k. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002560 | OEIS-A079491

Numerator of Sum_{k=0..n} binomial(n,k)/2^(k*(k-1)/2).

Conjecture: Also the number of loop-graphs on n vertices without any non-loop edge having loops at both ends, with formula a(n) = Sum_{k=0..n} binomial(n,k) 2^(k*(n-k) + binomial(k,2)). The unlabeled version is A339832. - _Gus Wiseman_, Jan 25 2024 The above conjecture is true since (n-k)*k + binomial(n-k,2) = binomial(n,2) - binomial(k,2) and A006125 gives the denominators for this sequence. - _Andrew Howroyd_, Feb 20 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002561 | OEIS-A079495

Numbers k such that the sum of the squares of the digits of k in base 3 is 0 (mod 3).

In base 2 this gives the "Evil Numbers" (cf. A001969) and slope 2. One may conjecture that in base b the asymptotic slope will be b and might suspect asymptotic density 1/b for each result (mod b). For nonprime b larger variations occur and "very big" numbers must be considered to believe in the conjecture (1 million or more...). (Related to A006287, here mod b is considered)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carlos Alves_, Jan 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002562 | OEIS-A079498

Numbers whose sum of digits in base b gives 0 (mod b), for b = 3.

In base 2 this gives the "Evil Numbers" (cf. A001969). One may conjecture that in base b the asymptotic slope will be b and asymptotic density 1/b for each result (mod b). Cases b=31 or b=61 gave considerable number of primes on the sequence. Proof of this conjecture: in general, the sequence

d with terms d(n) = sum of digits of n written in base b (mod b) is a fixed point of the generalized Thue-Morse morphism 0->01..b-1, 1->12..0, etc. See A053839 for the case b=4. It follows directly from this that all symbols have asymptotic density 1/b, and therefore that the positional sequences all have asymptotic slope b. - _Michel Dekking_, Apr 18 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carlos Alves_, Jan 21 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002563 | OEIS-A079555

Decimal expansion of Product_{k>=1} (1 + 1/2^k) = 2.384231029031371...

2 + Sum_{k>1} 1/(Product_{i=2..k} (2^i-1)) = 2 + 1/3 + 1/(3*7) + 1/(3*7*15) + 1/(3*7*15*31) + 1/(3*7*15*31*63) + ... (conjecture). - _Werner Schulte_, Dec 22 2016 The above conjecture of Schulte follows by setting x = 1/2 and t = -1 in the identity Product_{k >= 1} (1 - t*x^k) = Sum_{n >= 0} (-1)^n*x^(n*(n+1)/2)*t^n/(Product_{k = 1..n} 1 - x^k), due to Euler.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jan 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002564 | OEIS-A079559

Number of partitions of n into distinct parts of the form 2^j-1, j=1,2,....

A006697(k) gives number of distinct subwords of length k, conjectured to be equal to A094913(k)+1. - _M. F. Hasler_, Dec 19 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Jan 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002565 | OEIS-A079611

Waring's problem: conjectured values for G(n), the smallest number m such that every sufficiently large number is the sum of at most m n-th powers of positive integers.

Waring's problem: conjectured values for G(n), the smallest number m such that every sufficiently large number is the sum of at most m n-th powers of positive integers. See A002804 for the simpler problem of Waring's original conjecture, which does not restrict the bound to "sufficiently large" numbers. - _M. F. Hasler_, Jun 29 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002566 | OEIS-A079641

Matrix product of Stirling2-triangle A008277(n,k) and unsigned Stirling1-triangle |A008275(n,k)|.

Conjecture: o.g.f. as a continued fraction of Stieltjes type: 1/(1 - x*z/(1 - 2*z/(1 - (x + 1)*z/(1 - 4*z/(1 - (x + 2)*z/(1 - 6*z/(1 - (x + 3)*z/(1 - 8*z/(1 - ...))))))))) = 1 + x*z + (2*x + x^2)*z^2 + (6*x + 6*x^2 + x^3)*z^3 + ... - _Peter Bala_, Dec 12 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Jan 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002567 | OEIS-A079665

Triangular array read by rows: row s contains integers of the form (2^s+1)/(2^r+1) in order of increasing r <= s-1.

Conjecture: (b^s+1)/(b^r+1) is an integer if and only if: 1) r<s/2, 2) if s==1 (mod 2) then r is divisor of s, 3) if s=k*2^t with gcd(k,2^t)=1 then r is 2^t*u with u dividing k

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jose R. Brox (tautocrona(AT)terra.es), Jan 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002568 | OEIS-A079670

a(n) = (gcd(2^p-1, Fibonacci(p))-1)/(8*p) > 0 where p = A080050(n).

The members of the sequence seem to be all integers. Conjecture: a(n) > 1 if A080050(n) !== 1 (mod 10). The reverse is not true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Jan 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002569 | OEIS-A079726

a(1)=1, a(n) is the smallest integer > a(n-1) such that a(n) is even if k(n)=1, a(n) is odd if k(n)=2, where k(n) denotes the n-th term of the Kolakoski sequence (A000002).

a(n) is conjectured to be asymptotic to (4/3)*n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 17 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002570 | OEIS-A079729

Kolakoski-(1,2,3) sequence: a(n) is the length of the n-th run.

Conjecture: the frequencies of 1's, 2's and 3's in (a(n)) exist and are all equal to 1/3. This conjecture implies the conjecture of Benoit Cloitre on the partial sum sequence. - _Michel Dekking_, Jan 31 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 17 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002571 | OEIS-A079853

Primes p for which (p-2)! == 1 (mod p^2).

Zhi-Wei SUN conjectures that for n>1, a(n) == 3 (mod 8). (Posting to the Number Theory Mailing List, Nov 02 2009; added by _N. J. A. Sloane_, Nov 02 2009) Conjecture: primes p such that Sum_{k=1..p-1} k^(1-p) == -1 (mod p^2) are the odd terms of this sequence. - _Thomas Ordowski_, Jul 02 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | Pavlos Saridis (pavlos19(AT)yahoo.com), Sep 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002572 | OEIS-A079883

a(1) = 1; a(n) = prime(n) - prime(n-1)* a(n-1) if n > 1.

1. Let s(n) be a sequence such that lim s(n)/s(n+1) = K different from -1. The "oscillator sequence" (or simply "oscillator") of s(n) is the sequence s'(n) defined by the rules: s'(1) = 1; s'(n) = 1 - (s(n-1)/s(n)) s'(n-1). 2. It is an open problem whether the oscillator (prime)' converges to 1/2 or diverges. 3. For s(n) = prime(n), one has s'(n) = 1 - (p(n-1)/p(n)) s'(n-1) = [p(n) - p(n-1) s'(n-1)]/p(n). The

numerator is the expression p(n) - p(n-1) s'(n-1), which motivates the definition of the above sequence a(n). a(n) is called the "integral oscillator" of prime(n). In general the integral oscillator of s(n) can be defined similarly.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Feb 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002573 | OEIS-A079895

a(1) = 1; a(n) = phi(n) - phi(n-1)* a(n-1) if n > 1.

1. Let s(n) be a sequence such that lim s(n)/s(n+1) = K different from -1. The "oscillator sequence" (or simply "oscillator") of s(n) is the sequence s'(n) defined by the rules: s'(1) = 1; s'(n) = 1 - (s(n-1)/s(n)) s'(n-1). 2. It is an open problem whether the oscillator (prime)' converges to 1/2 or diverges. 3. s'(n) = 1 - (s(n-1)/s(n)) s'(n-1) = [s(n) - s(n-1) s'(n-1)]/s(n). The numerator is the expression s(n) - s(n-1) s'(n-1), which motivates the definition of the above sequence a(n). a(n) is called the "integral oscillator" of phi(n). In general the integral oscillator of s(n) can be defined similarly.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Feb 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002574 | OEIS-A079897

a(1) = 1; a(n) = sigma(n) - sigma(n-1)* a(n-1) if n > 1.

1. Let s(n) be a sequence such that lim s(n)/s(n+1) = K different from -1. The "oscillator sequence" (or simply "oscillator") of s(n) is the sequence s'(n) defined by the rules: s'(1) = 1; s'(n) = 1 - (s(n-1)/s(n)) s'(n-1). 2. It is an open problem whether the oscillator (prime)' converges to 1/2 or diverges. 3. s'(n) = 1 - (s(n-1)/s(n)) s'(n-1) = [s(n) - s(n-1) s'(n-1)]/s(n). The numerator is the expression s(n) - s(n-1) s'(n-1), which motivates the definition of the above sequence a(n). a(n) is called the "integral oscillator" of sigma(n). In general the integral oscillator of s(n) can be defined similarly.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Feb 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002575 | OEIS-A079898

a(1) = 1; a(n) = tau(n) - tau(n-1)* a(n-1) if n > 1.

1. Let s(n) be a sequence such that lim s(n)/s(n+1) = K different from -1. The "oscillator sequence" (or simply "oscillator") of s(n) is the sequence s'(n) defined by the rules: s'(1) = 1; s'(n) = 1 - (s(n-1)/s(n)) s'(n-1). 2. It is an open problem whether the oscillator (prime)' converges to 1/2 or diverges. 3. s'(n) = 1 - (s(n-1)/s(n)) s'(n-1) = [s(n) - s(n-1) s'(n-1)]/s(n). The numerator is the expression s(n) - s(n-1) s'(n-1), which motivates the definition of the above sequence a(n). a(n) is called the "integral oscillator" of tau(n). In general the integral oscillator of s(n) can be defined similarly.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Feb 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002576 | OEIS-A079899

a(1) = 1; a(n) = Fibonacci(n) - Fibonacci(n-1)* a(n-1) if n > 1.

1. Let s(n) be a sequence such that lim s(n)/s(n+1) = K different from -1. The "oscillator sequence" (or simply "oscillator") of s(n) is the sequence s'(n) defined by the rules: s'(1) = 1; s'(n) = 1 - (s(n-1)/s(n)) s'(n-1). 2. It is an open problem whether the oscillator (prime)' converges to 1/2 or diverges. 3. s'(n) = 1 - (s(n-1)/s(n)) s'(n-1) = [s(n) - s(n-1) s'(n-1)]/s(n). The numerator is the expression s(n) - s(n-1) s'(n-1), which motivates the definition of the above sequence a(n). a(n) is called the "integral oscillator" of Fibonacci(n). In general the integral oscillator of s(n) can be defined similarly.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Feb 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002577 | OEIS-A080070

Decimal encoding of parenthesizations produced by simple iteration starting from empty parentheses and where each successive parenthesization is obtained from the previous by reflecting it as a general tree/parenthesization, then adding an extra stem below the root and then reflecting the underlying binary tree.

Conjecture: only the terms in positions 0,1,2 and 4 are symmetric; i.e., A057164(A080068(n)) = A080068(n) (equivalently: A036044(A080069(n)) = A080069(n)) only when n is one of {0,1,2,4}. If this is true, then the formula given in A123050 is exact. I have checked this up to n=404631 with no other occurrence of a symmetric (general) tree. (See also A154477).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002578 | OEIS-A080076

Proth primes: primes of the form k*2^m + 1 with odd k < 2^m, m >= 1.

Conjecture: a(n) ~ (n log n)^2 / 2. - _Thomas Ordowski_, Oct 19 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Jan 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002579 | OEIS-A080131

Conjectured number of generalized Fermat primes of the form (n+1)^2^k + n^2^k, with k>1.

Conjectured number of generalized Fermat primes of the form (n+1)^2^k + n^2^k, with k>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jan 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002580 | OEIS-A080133

Conjectured number of generalized Fermat primes of the form (n+1)^2^k + n^2^k, with k>0.

Conjectured number of generalized Fermat primes of the form (n+1)^2^k + n^2^k, with k>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jan 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002581 | OEIS-A080192

Complement of A080191 relative to A000040. Prime p is a term iff there is no prime between 2*p and 2*q, where q is the next prime after p.

Conjecture: a(n)/A000040(n) is asymptotic to 3. Reason: I expect the distribution of terms in A102820 to converge to a geometric distribution with mean value 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Feb 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002582 | OEIS-A080201

Numbers that do not occur as differences between terms of the Mian-Chowla variant A051788.

Terms are only conjectures. Elements might be eliminated by higher terms of A051788, which was checked up to 4*10^6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Feb 05 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002583 | OEIS-A080208

a(n) is the least k such that the generalized Fermat number (k+1)^(2^n) + k^(2^n) is prime.

The first five terms correspond to the five known Fermat primes. The sequence A078902 lists some of the generalized Fermat primes. Bjorn and Riesel examined generalized Fermat numbers for k <= 11 and n <= 999. The sequence A080134 lists the conjectured number of primes for each k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Feb 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002584 | OEIS-A080239

Antidiagonal sums of triangle A035317.

A different formula for a(n) as a function of the Fibonacci numbers f(n) may be conjectured. The pattern is of the form a(n) = f(p)*f(p-q) - 1 if n mod 4 = 3, else f(p)*f(p-q) where p = 2,2,4,4,4,4,6,6,6,6,8,8,8,8... and q = 0,1,3,2,0,1,3,2,0,1,3,2... p(n) = 2 * A002265(n+4) = 2*(floor((n+3)/2) - floor((n+3)/4)) (see comment by _Jonathan Vos Post_ at A002265). A general formula for sequences having period 4 with terms a,b,c,d is given in A121262 (the discrete Fourier transform, as for all periodic sequences) and is a function of t(n)= 1/4*(2*cos(n*Pi/2) + 1 + (-1)^n). r4(a,b,c,d,n) = a*t(n+3) + b*t(n+2) + c*t(n+1) + d*t(n). This same formula may be used to subtract the 1 at n mod 4 = 3. a(n) = f(p(n))*f(p(n) - r4(1,0,3,2,n)) - r4(0,0,1,0,n). - _Gary Detlefs_, Dec 09 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 11 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002585 | OEIS-A080243

Signed super-Catalan or little Schroeder numbers.

Conjecture: (n+1)*a(n) +3*(2*n-1)*a(n-1) +(n-2)*a(n-2)=0. - _R. J. Mathar_, Nov 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002586 | OEIS-A080244

Signed generalized Fibonacci numbers.

Conjecture: (n+1)*a(n) +(7*n-2)*a(n-1) +4*(2*n-1)*a(n-2) +6*(n-1)*a(n-3) +(-5*n+1)*a(n-4) +(-n+2)*a(n-5)=0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002587 | OEIS-A080253

a(n) is the number of elements in the Coxeter complex of type B_n (or C_n).

Conjecture: Let k be a positive integer. The sequence obtained by reducing a(n) modulo k is eventually periodic with the period dividing phi(k) = A000010(k). For example, modulo 7 we obtain the sequence [1, 3, 3, 0, 3, 4, 0, 3, 3, 0, 3, 4, 0, 3, 3, 0, 3, 4, 0, ...], with an apparent period of 6 = phi(7) beginning at a(1). Cf. A000670. - _Peter Bala_, Dec 31 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Boddington_ and _Tim Honeywill_, Feb 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002588 | OEIS-A080277

Partial sums of A038712.

a(n) is conjectured to be asymptotic to n*log(n)/log(2). - _Klaus Brockhaus_, Mar 23 2003 [See Bannister et al., 2013. - _N. J. A. Sloane_, Nov 26 2013]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002589 | OEIS-A080289

Integers n for which the ratio phi(n)/pi(n) is smaller than for any subsequent n. Here phi(n) is Euler's totient function and pi(n) is the number of primes that are at most n.

Discovered while proving that phi(n) > 2pi(n) for all n, to prove a conjecture on www.primepuzzles.net. The conjecture stated that for all sufficiently large even n, n is the sum of two coprime composite numbers. This is in fact true for all even n > 210. For more details email the author.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Luke Pebody (ltp1000(AT)hermes.cam.ac.uk), Feb 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002590 | OEIS-A080333

Partial sums of A080278.

a(n) is conjectured to be asymptotic to n*log(n)/log(3). - _Klaus Brockhaus_, Mar 23 2003 [This follows from the asymptotics of A333979. - _Pontus von Brömssen_, Sep 06 2020]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002591 | OEIS-A080353

a(1)=5; for n>1, a(n)=a(n-1)+1 if n is already in the sequence, a(n)=a(n-1)+2 otherwise.

Conjecture: This sequence is equivalent to N \ A008478. - _Michael De Vlieger_, Jul 02 2025 See Cloitre 2026 for a proof of the above conjecture. - _Michael De Vlieger_, Oct 11 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002592 | OEIS-A080371

a(n) is the smallest x such that the quotient d(x+1)/d(x) equals n, where d = A000005.

Conjecture: a(p) = 2^(p-1) * k - 1 for some k > 0 and prime p. - _David A. Corneth_, Jan 26 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Feb 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002593 | OEIS-A080435

a(1) = 1; a(n) = least k > a(n-1) such that each prime of form a(i)+a(j) occurs for unique i <= j.

Conjecture: There are infinitely many primes not of the form a(i)+a(j). - _David W. Wilson_, Apr 14 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Feb 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002594 | OEIS-A080439

a(1) = 11, a(n) is the smallest prime obtained by inserting digits between every pair of digits of a(n-1).

Conjecture: Only one digit needs to be inserted between each pair of digits of a(n-1) to get a(n); i.e., a(n) contains exactly 2n-1 digits for n > 1. The conjecture above is false: a(5)=10000000000060571 has 17 digits instead of 2*5-1=9. A refined conjecture is: a(n) contains exactly 2^(n-1) + 1 digits for all n>0. This follows trivially by induction from the initial conjecture (above) of only one digit needed between each pair, and the fact that we start with 11, a 2-digit number, and holds true at least till a(12). - _Julio Cesar Hernandez-Castro_, Jul 05 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Feb 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002595 | OEIS-A080440

a(1) = 13, a(n) is the smallest prime obtained by inserting digits between every pair of digits of a(n-1).

Conjecture: Only one digit needs to be inserted between each pair of digits of a(n-1) to get a(n); i.e., a(n) contains exactly 2n-1 digits for n > 1. The conjecture above is false: a(4) = 100000963 has 9 digits instead of 2*4 - 1 = 7. A refined conjecture is: a(n) contains exactly 2^(n-1) + 1 digits for all n > 0. This follows trivially by induction from the initial above conjecture of only one digit needed between each pair, and the fact that we start with 13, a 2-digit number, and holds true at least till a(12). - _Julio Cesar Hernandez-Castro_, Jul 06 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Feb 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002596 | OEIS-A080441

a(1) = 17, a(n) is the smallest prime obtained by inserting digits between every pair of digits of a(n-1).

Conjecture: Only one digit needs to be inserted between each pair of digits of a(n-1) to get a(n); i.e. a(n) contains exactly 2n-1 digits for n > 1. The conjecture above is false: a(5) = 10000000000003037 has 17 digits instead of 2*5 - 1 = 9. A refined conjecture is: a(n) contains exactly 2^(n-1) + 1 digits for all n > 0. This follows trivially by induction from the initial conjecture above of only one digit needed between each pair, and the fact that we start with 17, a 2-digit number, and holds true at least till a(12).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Feb 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002597 | OEIS-A080442

a(1) = 19, a(n) is the smallest prime obtained by inserting digits between every pair of digits of a(n-1).

Conjecture: Only one digit needs to be inserted between each pair of digit of a(n-1) to get a(n); i.e., a(n) contains exactly 2n-1 digits for n > 1. The conjecture above is false: a(5) = 10000000000002359 has 17 digits instead of 2*5 - 1 = 9. A refined conjecture is: a(n) contains exactly 2^(n-1) + 1 digits for all n > 0. This follows trivially from the initial conjecture of only one digit needed between each pair, and the fact that we start with 19, a 2-digit number, and holds true at least till a(12). - _Julio Cesar Hernandez-Castro_, Jul 07 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Feb 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002598 | OEIS-A080469

Composite numbers k such that binomial(3*k, k) == 3^k (mod k).

A subsequence of A109641. The terms a(n) with n=2, 6, 7, 8, 9, 10 are of the form 3^k*p where p is prime and k=1, 3, 2, 5, 6, 7, respectively. It is tempting to conjecture that there are (infinitely many?) more terms of that form. - _M. F. Hasler_, Nov 11 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 15 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002599 | OEIS-A080529

Number of nucleons in longest known radioactive decay series ending with Lead 206 ("uranium series"), reversed.

The numbers are all equivalent to 2 mod 4. The sequence was submitted reversed, as it is open in the past. It is unknown if reversed decay series are finite (the "island of stability" hypothesis).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Feb 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002600 | OEIS-A080530

Number of nucleons in longest known radioactive decay series ending with Lead 207 ("actinium series"), reversed.

The numbers are all equivalent to 3 mod 4. The sequence was submitted reversed, as it is open in the past. It is unknown if reversed decay series are finite (the "island of stability" hypothesis).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Feb 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002601 | OEIS-A080531

Number of nucleons in longest known radioactive decay series ending with Lead 208 ("thorium series"), reversed.

The numbers are all equivalent to 0 mod 4. The sequence was submitted reversed, as it is open in the past. It is unknown if reversed decay series are finite (the "island of stability" hypothesis).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Feb 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002602 | OEIS-A080532

Number of nucleons in longest known radioactive decay series ending with Lead 209, reversed.

The numbers are all == 1 mod 4. The sequence was submitted reversed, as it is open in the past. It is unknown if reversed decay series are finite (the "island of stability" hypothesis).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Feb 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002603 | OEIS-A080572

Number of ordered pairs (i,j), 0 <= i,j < n, for which (i & j) is nonzero, where & is the bitwise AND operator.

Conjectured to be less than or equal to lcs(n) (see sequence A063437). The value of a(2^n) is that given in Stinson and van Rees and the value of a(2^n-1) is that given in Fu, Fu and Liao. This function gives an easy way to generate these two constructions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Bean_, Feb 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002604 | OEIS-A080578

a(1)=1; for n > 1, a(n) = a(n-1) + 1 if n is already in the sequence, a(n) = a(n-1) + 3 otherwise.

a(n) = A055938(n-1) + 2 (conjectured). - _Ralf Stephan_, Dec 27 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Benoit Cloitre_, Mar 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002605 | OEIS-A080612

Numbers m such that (1/p(2*m+1)) * Sum_{k=1..m} p(2*k+1)-p(2*k) >= (1/p(2*m))* Sum_{k=1..m} p(2*k)-p(2*k-1) where p(k) denotes the k-th prime.

Conjectured to be finite with last term = 314. This conjecture is false, 1494 is the first term after 314. - _Sean A. Irvine_, Sep 24 2025 Other conjecture log(n)^2 * (1/p(2*n+1) * Sum_{k=1..n} (p(2*k+1)-p(2*k)) - (1/p(2*n)) * Sum_{k=1..n} (p(2*k)-p(2*k-1))) -> constant. Weaker : previous formula is bounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002606 | OEIS-A080651

Numbers n such that n and Fibonacci(n) have the same number of divisors.

a(n) = A001605(n-8) for n >= 21 (conjectured). - _Chai Wah Wu_, Dec 30 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph L. Pe_, Feb 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002607 | OEIS-A080674

a(n) = (4/3)*(4^n - 1).

Conjectured to be the number of integers from 0 to (10^n)-1 that lack 0, 1, 2, 3, 4 and 5 as a digit. - _Alexandre Wajnberg_, Apr 25 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002608 | OEIS-A080692

a(n)=(-1)^(n+1)*det(M(n)) where M(n) is the n X n matrix M(i,j)=min(abs(i-j),i).

a(n) ~ 2^(n-1) * (c*(log(n) + gamma) - 1), where gamma is the Euler-Mascheroni constant A001620 and 1/2 < c < 1. Conjecture: c = 1/sqrt(2). - _Vaclav Kotesovec_, Aug 23 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Mar 03 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002609 | OEIS-A080693

Numbers of the form p^2*q + r*s where p,q,r,s are (not necessarily distinct) primes.

A conjecture of Goldbach type says every number >= 26 is of this form.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Mario Maqueda Garcia [Garci'a] (israelmira(AT)terra.es), Mar 03 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002610 | OEIS-A080735

a(1)=1, then a(n)=2*a(n-1) if a(n-1) is prime, a(n)=a(n-1)+1 otherwise.

Conjectures: (Strong) Let x,y be 2 positive integers and define a(n) as a(1)=1, a(n)=x*a(n-1) if a(n-1) is prime, a(n)=a(n-1)+y otherwise; then lim_{n->oo} log(a(n))/sqrt(n) = C(x,y) exists. (Weak) log(a(n))/sqrt(n) is bounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Mar 08 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002611 | OEIS-A080783

a(1)=1, a(n)=a(n-1)-1 if n is already in the sequence, a(n)=a(n-1)+5 otherwise.

Conjectured to be asymptotic to 3n as n -> infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Mar 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002612 | OEIS-A080795

Number of minimax trees on n nodes.

a(2*n) = 2^n * A006154(2*n), n>0 (conjectured). - _Ralf Stephan_, Apr 29 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, Mar 14 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002613 | OEIS-A080842

Numbers in the x/3 + 1 conjecture: Repeat until 1 is reached: if x is divisible by 3 then divide by 3, otherwise add 1.

Numbers in the x/3 + 1 conjecture: Repeat until 1 is reached: if x is divisible by 3 then divide by 3, otherwise add 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Mar 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002614 | OEIS-A080865

Order of symmetry groups of n points on 3-dimensional sphere with minimal distance between them maximized, also known as hostile neighbor or Tammes problem.

If more than one best packing exists (this occurs for n = 15, 62, 76, 117, ...; see Buddenhagen, Kottwitz link) for a given n, the one with the largest symmetry group is chosen. A conjectured (except n=24) continuation of the sequence starting with n=15 would be: 3, 16, 4, 2, 2, 12, 1, 1, 1, 24, 3, 2, 4, 1, 1, 6, 5, 6, 3, 2, 1, 4, 2, 24, 1, 3, 1, 10, 1, 2, 1, 2, 1, 24, 2, 12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Feb 21 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002615 | OEIS-A080866

Number of equal shortest edges in the solutions of the Tammes' problem.

A conjectured continuation of the sequence starting with n=15 would be: 30, 32, 34, 34, 34, 39, 40, 42, 43, 60, 48, 46, 52, 52, 54, 63, 60, 66, 66, 68, 66, 72, 66, 72, 76, 78, 81, 85, 82, 88, 84, 91, 89, 120, 96, 102.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Feb 21 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002616 | OEIS-A080894

Expansion of the exponential series exp(x M(x)) = exp((1-sqrt(1-2x-3x^2))/(2x)), where M(x) is the ordinary generating series of the Motzkin numbers A001006.

Conjecture D-finite with recurrence: +(-2*n+3)*a(n) +(-2*n^3+9*n^2-9*n+1)*a(n-1) +(n-1)*(n-2)*(4*n^2-2*n-3)*a(n-2) +3*(n-1)*(n-3)*(2*n-1)*(n-2)^2*a(n-3)=0. - _R. J. Mathar_, Jan 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, Mar 31 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002617 | OEIS-A080902

a(1)=1, a(n)=a(n-1)+2 if (n and n+3 are in the sequence), a(n)=a(n-1)+3 otherwise.

Conjectured to be asymptotic to r*n, where r=2.72... Is r the largest root of q^2-3*q+3/4, which is 2.72474487139...?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002618 | OEIS-A080937

Number of Catalan paths (nonnegative, starting and ending at 0, step +/-1) of 2*n steps with all values <= 5.

Proof: (M_3)^n = b(n-2)*(M_3)^2 - (6*b(n-3) - b(n-4))*M_3 + b(n-3)*1_3, for n >= 0, with b(n) = A005021(n), for n >= -4. For the proof of this see a comment in A005021. Hence (M_3)^n[1,1] = 2*b(n-2) - 5*b(n-3) + b(n-4), for n >= 0. This proves the 3 X 3 part of the conjecture in A332602 by _Gary W. Adamson_. The formula for a(n) given below in terms of r = rho(7) = A160389 proves that a(n)/a(n-1) converges to rho(7)^2 = A116425 = 3.2469796..., because r - 2/r = 0.6920... < 1, and r^2 - 3 = 0.2469... < 1. This limit was conjectured in A332602 by _Gary W. Adamson_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Feb 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002619 | OEIS-A080938

Number of Catalan paths (nonnegative, starting and ending at 0, step +-1) of 2*n steps with all values less than or equal to 7.

For A094256(n+1)/A094256(n), like for A094829(n+1)/A094829(n), the limit for n -> infinity is rho(9)^2 = A332438 = 3.53208888..., with rho(9) = 2*cos(Pi/9) = A332437. Therefore the formula of a(n) in terms of A094256 shows that the same limit is reached for a(n+1)/a(n). See this conjecture by _Gary W. Adamson_ in A332602.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Feb 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002620 | OEIS-A080958

a(n) = n!*(2/1 - 3/2 + 4/3 - ... + s*(n+1)/n), where s = (-1)^(n+1).

Conjecture: -(n+1)*a(n+1) + a(n) + n^2*(n+2)*a(n-1) = 0. - _R. J. Mathar_, Sep 27 2012, corrected for offset 1 by _Robert Israel_, Dec 26 2018 Conjecture verified, using the differential equation (x^3-x)*g''(x) + (5*x^2-1)*g'(x) + (3*x+1)*g(x) + 2 = 0 satisfied by the e.g.f. - _Robert Israel_, Dec 26 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002621 | OEIS-A080968

Orbit size of each branch-reduced tree encoded by A080981(n) under Donaghey's "Map M" Catalan automorphism.

If the conjecture given in A080070 is true, then this sequence contains only six 2's. Questions: are there any (other) values with finite number of occurrences? Which primes will eventually appear?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002622 | OEIS-A080978

a(n) = 2*A006046(n) + 1.

Conjectured partial sums of A131136. - _Sean A. Irvine_, Jun 25 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002623 | OEIS-A081053

Sum of a prime and the square of a prime.

Suggested by a Goldbach-type conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Pilar Guerra Cardenas (pilarguerracardenas(AT)hotmail.com), Mar 03 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002624 | OEIS-A081071

a(n) = Lucas(4*n+2)-2 = Lucas(2*n+1)^2.

Conjecture: a(n) = Fibonacci(4*n+3) + Sum_{k=2..2*n} Fibonacci(2*k). - _Alex Ratushnyak_, May 06 2012 The above conjecture is true for n >= 1. - _Nguyen Tuan Anh_, Aug 02 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_, Mar 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002625 | OEIS-A081086

Smallest partial quotients of an infinite simple continued fraction such that the fractional remainders sum to unity.

Conjecture: log(a(n+1))/log(a(n)) -> 2. The 9th term has 69 digits, the 10th term has 140 digits. The decimal expansion of the continued fraction: [0;2,2,9,91,14201,252238179,...] = 0.404250350307436947086987047594...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hans Havermann_ and _Paul D. Hanna_, Mar 05 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002626 | OEIS-A081113

Number of paths of length n-1 a king can take from one side of an n X n chessboard to the opposite side.

Conjecture: g.f.(x)=z*A064808(z), where z=x*A001006(x) and A...(x) are the corresponding generating functions. - _R. J. Mathar_, Jul 07 2009 Conjecture from WolframAlpha (verified for 1<=n<=180): (n+3)*a(n+4) = 27*n*a(n) +27*a(n+1) -9*(2*n+5)*a(n+2) +(8*n+21)*a(n+3). - _Alexander R. Povolotsky_, Jan 04 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Scambler_, Apr 16 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002627 | OEIS-A081124

Binomial transform of floor(n/2)!.

Conjecture: 2*a(n) -4*a(n-1) +(-n+2)*a(n-2) +(n-1)*a(n-3)=0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002628 | OEIS-A081145

a(1)=1; thereafter, a(n) is the least positive integer which has not already occurred and is such that |a(n)-a(n-1)| is different from any |a(k)-a(k-1)| which has already occurred.

The 1977 paper of Slater and Vélez proves that this sequence is a permutation of positive integers and conjectures that its absolute difference sequence (see A308007) is also a permutation. If we call this the "Slater-Vélez permutation of the first kind", then they also constructed another permutation (the 2nd kind), for which they are able to prove that both the sequence (A129198) and its absolute difference (A129199) are true permutations. - Ferenc Adorjan, Apr 03 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Don Reble_, Mar 08 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002629 | OEIS-A081169

Triangle in which n-th row gives trajectory of n (including n itself) under the map x -> x/2 if x is even, x -> 3*x-1 if x is odd, stopping when reaching 1, 5 or 17.

It is conjectured that the sequence will always end in one of three loops: 1, 2,1,1, ...; 5 14 7 20 10 5...; or 17 50 25 74 37 110 55 164 82 41 122 61 182 91 272 136 68 34 17...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Apr 16 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002630 | OEIS-A081199

5th binomial transform of (0,1,0,1,...), A000035.

Conjecture (verified up to a(9)): Number of collinear 4-tuples of points in a 4 X 4 X 4 X ... n-dimensional cubic grid. - _R. H. Hardin_, May 24 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 11 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002631 | OEIS-A081200

6th binomial transform of (0,1,0,1,0,1,...), A000035.

Conjecture (verified up to a(9)): Number of collinear 5-tuples of points in a 5 X 5 X 5 X ... n-dimensional cubic grid. - _Ron Hardin_, May 24 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 11 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002632 | OEIS-A081201

7th binomial transform of (0,1,0,1,0,1,....), A000035.

Conjecture (verified up to a(8)): Number of collinear 6-tuples of points in a 6 X 6 X 6 X... n-dimensional cubic grid. [_R. H. Hardin_, May 23 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 11 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002633 | OEIS-A081207

Main diagonal of number square A081206.

Conjecture: n*(n-2)*a(n) +(-2*n^2+5*n-1)*a(n-1) +(-n^2+3*n-4)*a(n-2) +(-2*n^2+7*n-4)*a(n-3) +(n-1)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Nov 12 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 11 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002634 | OEIS-A081237

Numbers n with no k, 0<k<n, such that mu(n-k)=mu(n)=mu(n+k), where mu=A008683 (Moebius function).

Conjecture: there are no more terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Mar 11 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002635 | OEIS-A081246

Triangle in which (2^n+1)-st row gives trajectory of x=2^n+1 under the map x -> x/2 if x is even, x -> x+1 if x is odd, stopping when reaching 1.

This is the 2^n+1 conjecture and is easily proved to converge to 1. The number of steps required to reach 1 is always 2n+2. Since (2^(n)+1+1)/2 = 2^(n-1)+1 (2^(n-1)+1+1)/2 = 2^(n-2)+1 .... (2^(n-n+1)+1+1)/2 = 2^(n-n)+1 = 2 2/2 = 1 thus 1 is guaranteed.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Apr 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002636 | OEIS-A081278

Binomial transform of Chebyshev polynomial coefficients A001793.

Conjecture: a(n) is the number of intervals in a poset closely related to Hochschild lattices. The cardinality of the poset is A001792(n). For example, the total order on 3 elements has 6 intervals. - _F. Chapoton_, Feb 06 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 16 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002637 | OEIS-A081314

Order of symmetry groups of n points on 3-dimensional sphere with the volume enclosed by their convex hull maximized.

If more than one configuration with maximal volume exists for a given n, the one with the largest symmetry group is chosen. Berman and Hanes give optimality proofs for n<=8. Higher terms are only conjectures. An independent verification of the results by Hardin, Sloane and Smith has been performed by Pfoertner in 1992 for n<28. An archive of the results with improvements for n=23,24 added in 2003 is available at link. A conjectured continuation of the sequence starting with n=28 is: 12,6,2,6,120,2,4,4,2,20,4,12,24,12,20,4,8,2,2,2,4,1,24

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Mar 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002638 | OEIS-A081319

Smallest squarefree integer k such that Q(sqrt(-k)) has class number n, or 0 if no such k exists.

Conjecture: all numbers at even positions are congruent 2 (mod 3). - _Ralf Stephan_, May 23 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, Mar 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002639 | OEIS-A081320

Largest 3-smooth divisor of n-th Fibonacci number.

Conjecture: for n>12 and n>0 modulo 12: a(n)=a(n-12) and a(12*k)=A065331(k)*144. The first part of the conjecture follows from the fact that the Fibonacci numbers are a strong divisibility sequence. - _Charles R Greathouse IV_, Sep 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, May 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002640 | OEIS-A081324

Twice a square but not the sum of 2 distinct squares.

Conjecture: for n>1 this is A050804. Conjecture is true. Proof : Conjecture: sequence consists of numbers of form 2*k^2 such that sigma(2*k^2)==3 (mod 4) and k is not divisible by 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002641 | OEIS-A081352

Main diagonal of square maze arrangement of natural numbers A081349.

Conjecture: let a and b be integers such that 0 < a < b so that 0 < a/b is a proper fraction. Define the map f(a,b,D) = a/b + gcd(a,b)/D. Of course, all such a/b can be partially ordered by value, i.e., 1/2 = 0.5 < 2/3 = 4/6 = 6/9 = 0.6666... < 3/4 = 6/8 = 0.75 < 4/5 = 0.8 etc. The map f appears to specify a total strict order on the co-domain for all a/b that is consistent with the given partial order of the domain, i.e., the partial order remains intact, while equivalent fractions are given a total strict order themselves. Moreover, equivalent fractions are strictly ordered by numerator (or denominator), e.g., 1/2 < 2/4 < 3/6 etc. The conditions are that for n >= 4 all of the fractions with denominator b <= n are listed and the minimum integer value of D to achieve the total strict order of the co-domain is 2*C(n-1,2) - (-1)^(n-1). So, a(n-3) = D for n >= 4. Example: given n = 4, we have D = 2*(4-1,2) - (-1)^(4-1) = 2*3 + 1 = 7 = a(4-3) = a(1). Partial order of domain. 1/4 < 1/3 < 1/2 = 2/4 < 2/3 < 3/4. Total order of co-domain. f(1,4,7) = 1/4 + 1/7 = 33/84 < f(1,3,7) = 1/3 + 1/7 = 40/84 < f(1,2,7) = 1/2 + 1/7 = 54/84 < f(2,4,7) = 2/4 + 2/7 = 66/84 < f(2,3,7) = 2/3 + 2/7 = 68/84 < f(3,4,7) = 3/4 + 1/7 = 75/84. Observe that if D = 6, then f(2,4,6) = 2/4 + 2/6 = 10/12 = f(2,3,6) = 2/3 + 1/6. Computation shows the same failure to achieve total strict order of the co-domain for D = 2..5. (As a >= 1, then b >=2, from the above). Computation also shows that the conjecture holds for n = 4..17. - _Ross La Haye_, Oct 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002642 | OEIS-A081362

Expansion of q^(1/24) * eta(q) / eta(q^2) in powers of q.

Conjecture: Sum_{k>=0} a(k) / exp(k*Pi) = exp(-1/24*Pi)*2^(1/8) = A292824. - _Simon Plouffe_, Sep 09 2025 Plouffe's conjecture is true (see Fried link) - _Sela Fried_, Nov 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Mar 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002643 | OEIS-A081391

Numbers k such that the central binomial coefficient C(2*k,k) has only one prime divisor whose exponent exceeds one.

No more terms through 10^6; conjecture: no terms after 786. - _Jon E. Schoenfield_, Jul 29 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Mar 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002644 | OEIS-A081477

Complement of A086377.

Conjectures from _Clark Kimberling_, Aug 03 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 12 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002645 | OEIS-A081515

Sum of terms in n-th row of A081517.

a(n) is > the n-th triangular number. Conjecture: There are no composite members.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002646 | OEIS-A081545

Smallest prime which is 1 more than the product of n distinct composite numbers.

Let K(m) be the smallest possible K satisfying the Theorem. Conjecture: K(m) ~ m, i.e., a(k) ~ A002808(1)*...*A002808(k), only very few of the last factors will be insignificantly larger. Let K(b,r) be the smallest possible K satisfying the Corollary, i.e., the index from which on all a(k)-1 are multiples of b^r. With the preceding conjecture, there are asymptotically at least (k+PrimePi(A002808(k)))/b multiples of b among the factors of a(k)-1, so this is an (asymptotic) lower bound on r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002647 | OEIS-A081567

Second binomial transform of F(n+1).

Conjecture: a(n) = 2*a(n-1) + A039717(n). - _Benito van der Zander_, Nov 20 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002648 | OEIS-A081601

Numbers m such that 3 does not divide Sum_{k=0..m} binomial(2k,k) = A006134(m).

G.f.: (x/(1 - x))*Sum_{k>=0} 3^(k+1)*x^(2^k)/(1 + x^(2^k)) (conjecture). - _Ilya Gutkovskiy_, Jul 23 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002649 | OEIS-A081621

Number of n-node triangulations of the sphere with minimal degree 5.

Grace conjectured that all polyhedra inscribed in the unit sphere with maximal volume are "medial" (all faces triangular and vertex degree either m or m+1 where m < 6 - 12/n < m+1). For n = 12 and n > 13 the medial polyhedra have 12 vertices of degree 5 and n-12 vertices of degree 6. The numerical solutions for n<=50 of the maximal volume problem (A081314) have this property, except for n=33. The polyhedron with conjectured maximal volume with 33 vertices is very likely a counterexample to Grace's conjecture, as it has a vertex with degree 7. See the visualization in the Pfoertner link. - _Hugo Pfoertner_, Jan 08 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Mar 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002650 | OEIS-A081658

Triangle read by rows: T(n, k) = (-2)^k*binomial(n, k)*Euler(k, 1/2).

Conjecture: If n is prime then n divides T(n, k) for 1 <= k <= n-1. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002651 | OEIS-A081669

Expansion of exp(2x)+exp(x)BesselI_0(2x).

Conjecture: n*a(n) +(-6*n+5)*a(n-1) +(9*n-17)*a(n-2) +4*(n-1)*a(n-3) +12*(-n+3)*a(n-4)=0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002652 | OEIS-A081672

Expansion of exp(2x) - exp(0) + BesselI_0(2x).

Conjecture: n*a(n) +2*(1-n)*a(n-1) +4*(1-n)*a(n-2) +8*(n-2)*a(n-3)=0. - _R. J. Mathar_, Nov 12 2012 E.g.f. satisfies y''' - (2*x-2)*y'' - (4*x + 2)*y' + (8*x-4)*y + 8x - 4 = 0, which implies Mathar's conjectured recurrence. - _Robert Israel_, Jun 03 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002653 | OEIS-A081673

Expansion of exp(3*x) - exp(x)*(1-BesselI_0(2*x)).

Conjecture: n*(2*n - 7)*a(n) +(-12*n^2 + 50*n - 33)*a(n-1) +(16*n^2 - 76*n + 87)*a(n-2) +3*(4*n^2 - 22*n + 27)*a(n-3) -9*(2*n-5)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Nov 24 2012 This implies Mathar's conjectured recursion. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002654 | OEIS-A081692

Define two sequences by A_n = mex{A_i,B_i : 0 <= i < n}, B_n = B_{n-1} + (A_n-A_{n-1})(A_n-A_{n-1}+1), where the mex of a set is the smallest nonnegative integer not in the set. Sequence gives A_n. B_n is in A081693.

Conjecture: Except for the initial 0, this is the sequence of positions of 0 in the fixed point of the morphism 0->01, 1->0000; see A284683. - _Clark Kimberling_, Apr 13 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002655 | OEIS-A081693

Define two sequences by A_n = mex{A_i,B_i : 0 <= i < n}, B_n = B_{n-1} + (A_n-A_{n-1})(A_n-A_{n-1}+1), where the mex of a set is the smallest nonnegative integer not in the set. Sequence gives B_n. A_n is in A081692.

Conjecture: Except for the initial 0, this is the sequence of positions of 1 in the fixed point of the morphism 0->01, 1->0000; see A284683. - _Clark Kimberling_, Apr 13 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002656 | OEIS-A081732

Number of numbers that differ from n in ternary representation by exactly one edit-operation: deletion, insertion, or substitution.

Conjecture: For m >= 3,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002657 | OEIS-A081739

a(n) = (-1/2^n)*(A000367(2^n)+1).

a(n) = A076547(n)-1 (Conjecture). - _R. J. Mathar_, Apr 22 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002658 | OEIS-A081760

Decimal expansion of Gamma(1/3)*Gamma(5/6)/Gamma(1/6).

Conjectured to be the value of Landau's constant L.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002659 | OEIS-A081767

Numbers k such that k^2 - 1 divides binomial(2k,k).

The data does not appear to support the conjectured asymptote statement (neither the constant nor being linear). - _Bill McEachen_, Feb 26 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002660 | OEIS-A081798

a(n) = Sum_{k = 0..n} C(n,k) * C(n+k,k) * C(n+2*k,k).

Conjecture: n^2*(3*n-4)*a(n) -(3*n-2)*(30*n^2-50*n+13)*a(n-1) +(9*n^3-30*n^2+29*n-6)*a(n-2) -(3*n-1)*(n-2)^2*a(n-3)=0. - _R. J. Mathar_, Apr 15 2016 Conjecture: (n^2)*a(n) +(-28*n^2+24*n-3)*a(n-1) +3*(-19*n^2+78*n-77)*a(n-2) +(5*n-12)*(n-3)*a(n-3) -2*(n-3)^2*a(n-4)=0. - _R. J. Mathar_, Apr 15 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, Apr 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002661 | OEIS-A081829

Numbers k such that the k-th term of Hofstadter Q-sequence is > (k+1)-th term.

Conjecture : a(n)/n>2 and a(n)/n->2

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002662 | OEIS-A081831

a(1)=a(2)=1, a(n) = a(a(n - a(n-1))) + a(n - a(n-2)).

Conjecture: a(n)/n -> C = 0.2....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002663 | OEIS-A081832

a(1)=a(2)=1, a(n) = a(n+1-2*a(n-1)) + a(n-2*a(n-2)).

Conjectures: a(n)/n -> C=1/4; a(n+1)-a(n)=1 or 0, first differences are 0, 1, 0, 0, 1, 0, 0, 1, 0, 0, 0, 1, 0, 0, 1, 0, 0, 0, 1, 0, 0, 1, 0, 0, 1, 0, 0, 0, 0, 1, 0, 0, 1, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002664 | OEIS-A081884

Number of steps needed to reach an integer starting with n+1/8 and iterating the map x-->x*ceiling(x).

Conjecture : let b(n,m) denotes the number of steps needed to reach an integer starting with n+1/2^m and iterating the map x-->x*ceiling(x); then sum(k=1,n,b(k,m)) is asymptotic to (m+1)*n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002665 | OEIS-A081920

Expansion of exp(2x)/sqrt(1-x^2).

Conjecture: a(n) -2*a(n-1) -(n-1)^2*a(n-2) +2*(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Nov 24 2012 Conjecture confirmed using d.e. (x^2-1)*y' + (-2*x^2+x+2)*y = 0 satisfied by the E.g.f. - _Robert Israel_, Feb 19 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002666 | OEIS-A081921

Expansion of exp(3x)/sqrt(1-x^2).

Conjecture: a(n)-3*a(n-1) -(n-1)^2*a(n-2) +3*(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Nov 24 2012 Mathar's conjecture follows from the differential equation (-3*x^2+x+3)*y+(x^2-1)*y'=0 satisfied by the E.g.f. - _Robert Israel_, Mar 14 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002667 | OEIS-A081923

Expansion of e.g.f.: exp(2x)/(1-x)^2.

Conjecture: a(n) +(-n-3)*a(n-1) +2*(n-1)*a(n-2)=0. - _R. J. Mathar_, Nov 24 2012 Applying Maple's ZeilbergerRecurrence command to the above series of Walsh for a(n) results in the first-order recurrence equation (n - 1)*a(n+1) = n*(n + 1)*a(n) - 2^(n+2) with a(0) = 1 and a(2) = 18. Using this it is easy to verify that a(n) satisfies the second-order recurrence a(n) = (n + 3)*a(n-1) - 2*(n - 1)*a(n-2) conjectured above by Mathar.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002668 | OEIS-A081978

Smallest triangular number with exactly n divisors, or 0 if no such number exists.

Conjecture: a(2j) > 0 for all j > 1. (This conjecture holds at least through n = 2j = 1000. The largest a(n) for even n <= 1000 is a(898) = 20599^448 * (2 * 20599^448 - 1) = 3.21...*10^3865.) (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 03 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002669 | OEIS-A082031

Expansion of e.g.f. exp(2*x)/(1-x)^3.

Based on this series the ZeilbergerRecurrence command in Maple 17 produces the first-order recurrence (n^2 - 3*n + 4)*a(n) = 2^(n+2) + n*(n^2 - n + 2)*a(n-1). Using this it is easy to verify the second-order recurrence conjectured above by Mathar.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002670 | OEIS-A082090

Length of iteration sequence if function A056239, a pseudo-logarithm is iterated and started at n. Fixed point equals zero for all initial values.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Apr 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002671 | OEIS-A082143

First subdiagonal of number array A082137.

Conjecture: (n+1)*a(n) +4*(-2*n-1)*a(n-1)=0. - _R. J. Mathar_, Oct 19 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002672 | OEIS-A082184

The a(n)-th triangular number is the sum of the n-th triangular number and the smallest triangular number possible.

a(n) is triangular if n+1 is triangular. Conjectures: partial maxima of sequence are at index i with value from A068195 and also a(i) - A082183(i) = 1, where i is in A068194.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Apr 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002673 | OEIS-A082207

Palindromes whose product of digits is a positive palindrome.

Conjecture: There are infinitely many terms in the sequence (of the type 777) for which the product of digits > 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002674 | OEIS-A082235

Triangular numbers obtained as a concatenation of successive terms of A081847.

Conjecture: Every triangular number other than 3, 6 and 10 is a member. (This is a re-arrangement of triangular numbers barring 3,6 and 10).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002675 | OEIS-A082250

a(n) = A082249(n)/7.

Some pattern is visible which would change when numbers of three or more digits occur in A082249. Conjecture: If the concatenation of k consecutive r-digit numbers is divisible by n then the concatenation of the next k numbers would also be divisible by n provided it contains the same number of digits as the first number. This should be true for both forward and reverse concatenation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 11 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002676 | OEIS-A082274

Palindromes k such that k + 2 is also a palindrome.

The above conjecture is true. Adding two to the least significant digit of a number can result in a carry of at most 1, which only happens if the digit of least significance is 8 or 9. If the least significant digit is 8, adding two results in that digit becoming 0, so the resulting number can't be palindromic. If only the k least and most significant digits are 9, the least significant digit will become 1 and all other adjacent digits 9 will turn into the digit 0 and produce a carry of 1. For the starting number to have been palindromic, the k most significant digits must also be 9's. Any digits that are not 9's between the 9's will not produce a carry on their own when increased by one through the previous carry, resulting in a nonpalindromic number with some 9's as most significant digits and a single 1 and 0's as least significant digits. - _Felix Fröhlich_, Jul 22 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002677 | OEIS-A082300

Numbers relatively prime to the sum of their prime factors (with repetition).

Conjecture: a(n) ~ (Pi^2/6)n. - _Charles R Greathouse IV_, Aug 04 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 08 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002678 | OEIS-A082381

Sequence of the squared digital root of a number until 1 or 4 is reached. The initial numbers 1,2,..n are not output.

Conjecture: The sequence always terminates with 1 or the 4 16 37 58 89 145 42 20 4... loop (cf. A080709).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Apr 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002679 | OEIS-A082382

Table which contains in row n the track of n->A003132(n) until reaching 1 or 4.

Conjecture: Each sequence terminates with 1 or the 4 16 37 58 89 145 42 20 4... loop.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Apr 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002680 | OEIS-A082385

For each n append T(n), T(T(n)), T^3(n), ..., T^r(n), where T(n) = A055012(n) and r is the smallest integer such that T^r(n) is one of the following numbers: 1, 55, 136, 153, 160, 370, 371, 407, 919.

Conjecture: The sequence always terminates with one of the following:(tested to n=1000000) 1,55,136,153,160,370,371,407,919 which eventually loop back to themselves. 1,153,370,371,407 loop back in 1 step and are the sum of the cubes of their digits. The others are 55,250,133,55. 136,244,136. 160,217,352,160. 919,1459,919. A046156, A046157 indicate this as a limit of possibilities of numbers that cubed digital roots roll back to the origional number. Proof? - _Cino Hilliard_, Apr 13 2003

Proof: In A055012 _T. D. Noe_ notes that for n > 1999, A055012(n) < n. This means that by repeatedly applying A055012, we eventually reach a number smaller than 2000. As checked by _Cino Hilliard_, all numbers below 10^6 end in one of the listed cycles. - _Stefan Steinerberger_, Sep 05 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Apr 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002681 | OEIS-A082397

Number of directed aggregates of height <= 2 with n cells.

Conjecture: partial sums of A342912. - _Sean A. Irvine_, Jul 16 2022 Conjecture D-finite with recurrence +(n+2)*a(n) +(-3*n-2)*a(n-1) -n*a(n-2) +3*n*a(n-3)=0. - _R. J. Mathar_, Jun 27 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Fouad IBN MAJDOUB HASSANI_, Apr 14 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002682 | OEIS-A082447

a(n) = the number k such that s(k)=0 where s(0)=n and s(i)=s(i-1)-(s(i-1) modulo (i+1)).

Conjecture: a(n) = sqrt(Pi*n) + O(1)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002683 | OEIS-A082449

Let f(p) = greatest prime divisor of p-1. Sequence gives smallest prime which takes at least n steps to reach 2 when f is iterated.

Conjecture: a(n) is the smallest prime p such that b(p) = n, where f(2) = 0 and for an odd prime p, f(p) = 1 + max{q|(p-1), q prime} f(q). In other words, a(n) is the smallest prime p such that A364332(primepi(p)) = n. Verified for n <= 13. - _Jianing Song_, Apr 28 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002684 | OEIS-A082477

Number of divisors d of n such that d+1 is also a divisor of n+1.

The partial sums of this sequence from k = 1 to 10^m, for m = 1, 2, ..., are 15, 185, 1952, 19852, 199538, 1998538, 19995362, 199985379, 1999953823, ... Conjecture: The asymptotic mean of this sequence is 2. - _Amiram Eldar_, Jun 04 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002685 | OEIS-A082512

a(n) = p is the smallest prime introducing a consecutive prime-difference pattern as follows: [2,2n,2], i.e., [p, p+2, p+2+2n, p+2+2n+2] are consecutive primes. Increasing middle prime gap in the immediate neighborhood of two small gaps(=2); a(n) = 0 if no such pattern exists.

It is conjectured that the twin primes in the neighborhood can be separated by an arbitrarily large gap.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Apr 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002686 | OEIS-A082528

Least k such that x(k)=0 where x(1)=n x(k)=k^3*floor(x(k-1)/k^3).

Conjecture : define sequence a(n,m) m real >0 as the least k such that x(k)=0 where x(1)=n x(k)=k^m*floor(x(k-1)/k^m) then a(n,m) is asymptotic to (c(m)*n)^(1/(m+1)). where c(m) is a constant depending on m.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002687 | OEIS-A082579

Expansion of e.g.f.: exp(x/(1-x)^2).

It appears that the sequence taken modulo 10 is periodic with period 5. More generally, we conjecture that for k = 2,3,4,... the difference a(n+k) - a(n) is divisible by k: if true, then for each k the sequence a(n) taken modulo k would be

periodic with period dividing k. - _Peter Bala_, Nov 14 2017 The above conjecture is true - see the Bala link. - _Peter Bala_, Jan 20 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, May 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002688 | OEIS-A082582

Expansion of (1 + x^2 - sqrt(1 - 4*x + 2*x^2 + x^4)) / (2*x) in powers of x.

Conjecture: Also the number of maximal simple graphs with vertices {1..n} and no weakly nesting edges. Two edges {a,b}, {c,d} are weakly nesting if a <= c < d <= b or c <= a < b <= d. For example, the a(1) = 1 through a(5) = 13 edge-sets are: In arXiv:2102.02777 I conjectured that this sequence counts the *Dyck natural numbers* of semilength n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, May 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002689 | OEIS-A082607

a(0)=1; for n > 0, a(n) = least k not included earlier such that k*a(n-1) - 1 is a square.

Conjecture: this is a permutation of A008784. - _Robert Israel_, Aug 25 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002690 | OEIS-A082613

Smallest palindrome divisible by an n-th power.

Conjecture: the n-th power that divides a(n) is 2^n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Apr 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002691 | OEIS-A082615

Palindromes such that the GCD of any pair of successive terms is a palindromic prime.

I conjecture that every sequence generated in this fashion, beginning with a palindrome that has a palindromic prime factor, is infinite and eventually coincides with this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Apr 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002692 | OEIS-A082630

Limit of the sequence obtained from S(0) = (1,1) and, for n > 0, S(n) = I(S(n-1)), where I consists of inserting, for i = 1, 2, 3..., the term a(i) + a(i+1) between any two terms for which 7*a(i+1) <= 11*a(i).

Conjecturally, a(n) = x + y, where these values solve x^2 - floor(y^2/3) = 1, see related sequences and formula below. - _Richard R. Forberg_, Sep 08 2013 Conjecturally, a(n) = A143643(n-1) + A005246(n), for n => 2, as derived from comment above. - _Richard R. Forberg_, Sep 08 2013 If the above conjectures are true, then a(n) = A001353(n)/A005246(n+1). - _Andrey Zabolotskiy_, Sep 26 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, May 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002693 | OEIS-A082662

Numbers k such that the odd part of k is less than sqrt(2k).

A further equivalent property, (P9), follows at once from (P4). This was conjectured by _Omar E. Pol_, Apr 18 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, May 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002694 | OEIS-A082758

Sum of the squares of the trinomial coefficients (A027907).

Conjecture: a(n) = [x^n] ((1 + x + x^3 + x^4)/(1 - x)^2)^n. If the conjecture is true then the Gauss congruences a(n*p^k) == a(n*p^(k-1)) (mod p^k) hold for all primes p and positive integers n and k. [added Aug 29 2025: The conjecture is true. The conjecture leads to the double sum representation a(n) = Sum_{j, k} binomial(n, k)*binomial(n, j)*binomial(3*n-3*j-k-1, n-3*j-k), which satisfies the second-order recurrence given above by _Peter Luschny_, as can be verified using the MulZeil procedure in Doron Zeilberger's MultiZeilberger Maple package.] The supercongruence a(n*p^k) == a(n*p^(k-1)) (mod p^(2*k)) for primes p >= 5 conjectured above by Bala follows from the theorem of Pan and Sun (see the links): by the Somos formula above, a(n*p^k) - a(n*p^(k-1)) = A002426(p*m) - A002426(m) with m = 2*n*p^(k-1), and their theorem gives p-adic valuation >= 2 + 2*v_p(m) >= 2*k. - _Andrew R. Epstein_, Jul 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, May 21 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002695 | OEIS-A082769

a(n) = smallest palindromic prime that begins with A082768(n), or 0 if no such number exists.

Conjecture: no entry is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002696 | OEIS-A082770

a(n) = smallest palindromic prime that begins with A082768(n) and contains more than twice the number of digits in A082768(n), or 0 if no such number exists.

Conjecture: no entry is zero. In most cases the number of digits required is 2k+1 where k is the number of digits in A082768(n). What is the first entry that requires more (than 2k+1) digits?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002697 | OEIS-A082844

Start with 3,2 and apply the rule a(a(1)+a(2)+...+a(n)) = a(n), fill in any undefined terms with a(t) = 2 if a(t-1) = 3 and a(t) = 3 if a(t-1) = 2.

Conjecture: a(n) = A097509(n+1). - _Benedict W. J. Irwin_, Mar 13 2016. [See the discussion in A097509. - _N. J. A. Sloane_, Mar 09 2021]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 15 2003; revised Jun 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002698 | OEIS-A082871

First component x of pairs (x,y) where x <= y and x!+y! is a square, sorted on x.

Conjecture: the only solutions for x!+y! = n^2 (x <= y) are x=0,0,0,1,1,1,2,4 and y=4,5,7,4,5,7,2,5 respectively.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, May 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002699 | OEIS-A082892

a(n) = floor(2*n/log(A000230(n))).

For these larger and larger gap-initiating primes, integer part of relevant quotient,q, may exceed 27, all values between 1 and 28 occur. Observation supports conjecture that infsup(q) is infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Apr 17 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002700 | OEIS-A082935

Smallest palindrome beginning with n and a digit sum of n at some stage.

Checked the conjecture above to n=100. - _Robert G. Wilson v_

猜想/问题提及段 | historical_or_current_mention_unreviewed | Meenakshi Srikanth (menakan_s(AT)yahoo.com), Apr 16 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002701 | OEIS-A082957

Numbers k such that sigma(2*k) < sigma(2*k+1).

Conjecture : a(n) is asymptotic to c*n where 18 < c < 18.5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002702 | OEIS-A082981

Start with the sequence S(0)={1,1} and for k>0 define S(k) to be I(S(k-1)) where I denotes the operation of inserting, for i=1,2,3..., the term a(i)+a(i+1) between any two terms for which 4a(i+1)<=5a(i). The listed terms are the initial terms of the limit of this process as k goes to infinity.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, May 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002703 | OEIS-A082991

a(1) = 1 and for n > 1, a(n) = 2 * length of the cycle reached for the map x -> A062401(x), starting at n [where A062401(n) = phi(sigma(n))], or -1 if no finite cycle is ever reached.

From the original definition: Define a sequence u_n as follows: u_n(1) = n, thereafter u_n(2k) = sigma(u_n(2k-1)), u_n(2k+1) = phi(u_n(2k)); then a(n) is the length of the ultimate period of u_n(k) (which is conjectured to become ultimately periodic for any n>=1). Conjecture: despite results for small terms, all even number are obtained as values. (For example, 12 occurs since a(12102) = 12).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002704 | OEIS-A082994

Numbers k such that all the following properties hold: (i) k*reverse(k) is a square; (ii) k != reverse(k); (iii) k and reverse(k) are not both squares; and (iv) k and reverse(k) have the same number of digits.

These numbers are counterexamples to the following conjecture given in the Ogilvy-Anderson reference: "When an integer and its reversal are unequal, their product is never a square except when both are squares." This sequence excludes terms like 2200, i.e. 2200*22 = 48400.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, May 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002705 | OEIS-A083005

a(1)=1, a(n)=2a(n-1)+1 if a(n-1) is prime, a(n)=a(n-1)+1 otherwise.

Conjecture : limit n ->oo log(a(n))/sqrt(n)=c= 1.3....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 31 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002706 | OEIS-A083020

Irregular triangle read by row of the prime factors of 3^(3^n) + 2.

Conjecture: These numbers have no repeating factors or 3^(3^n) + 2 is squarefree.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, May 31 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002707 | OEIS-A083125

a(1) = 1, then the smallest number not occurring earlier such that the product of a pair of successive terms is a palindrome.

Conjecture: except for multiples of 10, all natural numbers belong to this sequence. - _Reinhard Zumkeller_, May 02 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Apr 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002708 | OEIS-A083137

Smallest palindromic number relatively prime to all the previous terms.

323 is the first composite entry. Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Apr 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002709 | OEIS-A083139

Primes in A083137.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Apr 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002710 | OEIS-A083143

Smallest n-digit palindromic multiple of prime(n), or 0 if no such number exists.

Conjecture: no entry is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Apr 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002711 | OEIS-A083144

Smallest k such that k*prime(n) is an n-digit palindrome, or 0 if no such number exists.

Conjecture: no entry is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Apr 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002712 | OEIS-A083146

Smallest palindromic multiple of n-th palindrome which is not a concatenation of copies of that palindrome.

Conjecture: (1). The sequence differs from A083145 only for finitely many values of n. (2) Values of n are 5,6,7,8 and 9. Conjecture: The sequence differs from A083145 for infinitely many values of n. - _Franklin T. Adams-Watters_, May 16 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Apr 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002713 | OEIS-A083178

Numbers with a digit sum of n and a maximum product of digits. In case of two identical products choose the largest number.

Conjecture: a(n) = 10*a(n-3)+a(n-6)-10*a(n-9) for n>10. - _Colin Barker_, Oct 14 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002714 | OEIS-A083183

Smallest palindromic prime using only prime digits (2,3,5,7) and having a sum of digits = prime(n), or 0 if no such number exists.

Conjecture: no entry is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Apr 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002715 | OEIS-A083184

Smallest palindromic prime having a sum of digits = prime(n), or 0 if no such number exists.

Conjecture: no entry is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Apr 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002716 | OEIS-A083207

Zumkeller or integer-perfect numbers: numbers n whose divisors can be partitioned into two disjoint sets with equal sum.

The 229026 Zumkeller numbers less than 10^6 have a maximum difference of 12. This leads to the conjecture that any 12 consecutive numbers include at least one Zumkeller number. There are 1989 odd Zumkeller numbers less than 10^6; they are exactly the odd abundant numbers that have even abundance, A174865. - _T. D. Noe_, Mar 31 2010 Conjecture: Any 4 consecutive terms include at least one number k such that sigma(k)/2 is also a Zumkeller number (verified for the first 10^5 Zumkeller numbers). - _Ivan N. Ianakiev_, Apr 03 2017 Conjecture: If d > 1, d|k and tau(d)*sigma(d) = k, then k is a Zumkeller number (cf. A331668). - _Ivan N. Ianakiev_, Apr 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002717 | OEIS-A083214

Numbers k for which 3 | p(k), where p(k) = A000041(k) is the k-th partition number.

Conjecture : a(n) = 3n + o(n). - _Benoit Cloitre_, Oct 06 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Jun 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002718 | OEIS-A083219

a(n) = n - 2*floor(n/4).

Conjecture: number of roots of P(x) = x^n - x^(n-1) - x^(n-2) - ... - x - 1 in the left half-plane. - _Michel Lagneau_, Apr 09 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002719 | OEIS-A083338

Number of partitions of odd numbers into three primes and of even numbers into two primes.

a(n) > 0 for all n iff Goldbach's conjectures hold.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002720 | OEIS-A083339

a(n) is the number of distinct prime factors of n that occur in partitions into two primes when n is even and into three primes when n is odd.

Number of distinct prime factors of n that occur in prime-partitions confirming Goldbach's conjectures. (The original name of this sequence.) Conjecture: Apart from k=2, A070826(k): 1, 3, 15, 105, 1155, 15015, 255255, gives the positions of records (each equal to k-1). This follows from the conjectured formula. - _Antti Karttunen_, Sep 14 2017 If n is even, a(n) = A010051(n/2), if n is an odd prime, a(n) = 0, and for odd composites (conjecturally), a(n) = A001221(n). - _Antti Karttunen_, Sep 14 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002721 | OEIS-A083370

Primes satisfying f(2p)=p when f(1)=5 (see comment).

Conjecture: f(n) = n/2 + O(log(n)) and there are infinitely many primes p such that f(2p)=p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jun 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002722 | OEIS-A083371

Primes p such that q-p >= 8, where q is the next prime after p.

The original definition by Cloitre was: [Start from any initial value F(1) >= 2 and define F(n) as the largest prime factor of F(1)+F(2)+F(3)+...+F(n-1). The sequence contains the primes satisfying F(2*p)=p supposed F(1)=7. Conjecture: F(n)= n/2+O(log n) and the sequence is infinite.] _Don Reble_ showed Jan 22 2022 that these are the same primes p followed by a prime gap of q-p >=8, where q is the next prime after p: [

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jun 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002723 | OEIS-A083392

Alternating partial sums of A000217.

Conjecture: for n > 0, a(n-1) is equal to the determinant of an n X n symmetric Toeplitz matrix M(n) whose first row consists of a single zero followed by successive positive integers repeated (A004526). - _Stefano Spezia_, Jan 10 2020 Spezia's conjecture is true (see Fried link). - _Sela Fried_, Mar 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Jun 11 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002724 | OEIS-A083396

Least n such that n and 2k+n are both brilliant numbers.

Conjecture: for any k >= 1 there will always be a brilliant constellation of the form {n, 2k+n} for some n. (True for all k <= 5000.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jun 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002725 | OEIS-A083414

Write the numbers from 1 to n^2 consecutively in n rows of length n; let c(k) = number of primes in k-th column; a(n) = minimal c(k) for gcd(k,n) = 1.

Conjectured to be always positive for n>1. The conjecture is true for all n <= 40000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002726 | OEIS-A083428

Smallest prime that can be formed using the digits of first 3n+1 numbers, or 0 if no such number exists.

The digits of first 3n and first 3n-1 numbers yield a composite number which is a multiple of 3. Conjecture: No entry is zero. Subsidiary sequence: the number of distinct primes that can be formed using the digits of first 3n + 1 numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Apr 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002727 | OEIS-A083434

a(n) = smallest palindrome not included earlier and with the same prime signature as n, or 0 if no such palindrome exists.

Conjecture: There are infinitely many zeros in this sequence. a(32) = 0 is only a conjecture. If there is a palindromic 5th power, it is greater than 10^40. - _David Wasserman_, Mar 22 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Apr 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002728 | OEIS-A083439

a(1) = 3; then a(n+1) = smallest prime not already in the sequence such that the concatenations a(n)a(n+1) and a(n+1)a(n) are both primes.

Conjecture: every prime == 1 (mod 6) occurs in the sequence. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Apr 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002729 | OEIS-A083451

(n concatenated n times) - n^n.

Amarnath Murthy conjectured that all terms are positive for n>1. This is true. Consider the number of digits. There are n*ceiling(log_10(n+1)) digits in the concatenation, but only log_10(n^n)=n*log_10(n) in n^n. Therefore the terms are never negative. - Hauke Worpel (hw1(AT)email.com), Jun 03 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002730 | OEIS-A083462

Smallest palindrome k such that k - n is a distinct power of 2; i.e., a power of 2 that has not arisen earlier; or 0 if no such k exists.

After a(3) the sequence continues (with 0's conjectural): 0, 0, 22, 0, 0, 0, 0, 0, 44, 77, 0, 0, 272, 0, 4114, 0, 65556, 0, 0, 151, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 545, 0, 0, 8228, 0, 0, 0, 0, ... a(n) > 0 for the following values of n: 0, 1, 2, 3, 6, 12, 13, 16, 18, 20, 23, 33, 36, 55, 59, 64, 77, 87, 118, 137, 230, 545, 610, 750, 825. Conjecturally, there are no other such n <= 1000. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002731 | OEIS-A083470

Smallest prime which is the concatenation of n distinct primes in increasing order.

The 9th, 11th, 12th, 13th, ... terms are other counterexamples to the above "conjecture". On the other hand, it seems almost sure that the n-th term starts with the concatenation of the first n-2 primes. - _M. F. Hasler_, Mar 20 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002732 | OEIS-A083476

Positions of 0's in A083475.

Strong conjecture: There are only finitely many terms in this sequence not divisible by 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 03 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002733 | OEIS-A083480

Compress the triangular array A049597 by suppressing zero entries and reversing the order of each row.

Row sums => A000041. Diagonals are sums of Gaussian polynomials (which then sum to powers of two). The number of entries on each row is conjectured to conform to: 0 1 1 1 2 2 3 3 4 5 5 6 7 7 8 9 10 10 11 12 13 13 14 15 16 17 17 ... a sequence which stutters after values 0,1,2,4,6,9,12,16,...A002620. Regarding the first element of the sequence as T(1,0), it appears that this is the number of partitions of n with k elements not in the first hook; i.e., with n - (max part size) - (number of parts) + 1 = k. If this is correct, we have T(n,0) = n and for k > 0, T(n,k) = sum_{j >= max(0,2k-n+2)} j * T(k,j). This is equivalent to T(n,k) = T(n-1,k) + sum_{j >= max(0,2k-n+2)} T(k,j) and thus to T(n,k) = 2* T(n-1,k) - T(n-2,k) + T(k,2k-n+2) [taking T(n,k) = 0 if k < 0]. It also implies the correctness of the conjecture about the row lengths. - _Franklin T. Adams-Watters_, May 27 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alford Arnold_, Jun 08 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002734 | OEIS-A083502

Smallest k such that n*(n+k) + 1 is an n-th power.

[Since there is actually no comment in A083503: this probably means to say that (conjectural!) A083503(prime(n)) = A008864(n) which leads to a(p) = Sum_{s=2..p} binomial(p,s)*p^(s-1) for primes p, an increasing subsequence. - _R. J. Mathar_, Aug 01 2025]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 03 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002735 | OEIS-A083514

Number of steps for iteration of map x -> (4/3)*ceiling(x) to reach an integer > 3n+1 when started at 3n+1, or -1 if no such integer is ever reached.

It is conjectured that an integer is always reached.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Sep 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002736 | OEIS-A083518

Beginning with 3, a(i)*a(j) + 2 is prime for all i, j, i != j.

Conjecture: sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 05 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002737 | OEIS-A083552

Quotient when LCM of 2 consecutive prime differences is divided by GCD of the same two differences.

Conjecture: Every positive integer appears infinitely many times in this sequence. Example: a(834) = a(909) = ... = a(9901) = ... = 4. - _Jerzy R Borysowicz_, Dec 22 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, May 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002738 | OEIS-A083557

a(n) is the greatest prime factor of 3*a(n-1)+2.

Conjecture: if a(1)=m then the sequence becomes cyclic, for any m. Conjecture verified up to 25000000 by _Jud McCranie_, Jun 11 2003 G.f.: x*(3 + 11*x + 7*x^2 + 23*x^3 + 71*x^4 + 43*x^5 + 131*x^6 + 79*x^7 + 239*x^8 + 719*x^9 + 127*x^10 + 383*x^11 + 1151*x^12 + 691*x^13 + 83*x^14 + 251*x^15 + 151*x^16 + 13*x^17 + 41*x^18 + 2*x^19 + 6*x^20 + 46*x^21) / (1 - x^19) (conjectured). - _Colin Barker_, Jul 15 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yasutoshi Kohmoto_, Jun 05 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002739 | OEIS-A083561

a(1) = 1 and for n>1: a(n) = n + Max{a(k): k<n, gcd(n,a(k))=1}.

Conjecture: a(n) ~ n^2/3 as n -> infinity. - _Robert Israel_, Jun 11 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jun 12 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002740 | OEIS-A083567

Let B(k) be the number of binary digits in k equal to 1. This is the sequence of positive integers k such that 2B(k)=B(k^2).

Conjecture: The counting function p(n) satisfies p(n)=c n/log n + o(n/log n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giuseppe Melfi_, Jun 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002741 | OEIS-A083677

Define f(n, k) to be the concatenation of the first n primes, with n-1 k's inserted between the primes. Then a(n) is the smallest k >= 0 such that f(n, k) is prime, or -1 if no such prime exists.

a(3) = -1 because f(3, k) is always a multiple of 5. For any n such that n = 1 (mod 3) and A007504(n) = 0 (mod 3), a(n) = -1 because f(n, k) is always a multiple of 3. It is my conjecture that for all other n, -1 < a(n) < n*p(n). I've checked for all n < 270.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Jun 15 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002742 | OEIS-A083684

Numbers k such that there is no nonnegative integer m such that m < k*prime(k) and the concatenated decimal number fp(k,m) = prime(1).m.prime(2).m. ... .prime(k-1).m.prime(k) is prime.

is in the sequence. I conjecture that 3 is the only term of the sequence which is not of this form.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Jun 15 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002743 | OEIS-A083754

a(1) = 1 and then smallest odd number not occurring earlier such that the concatenation a(1)a(2)a(3)... is a prime.

Conjecture: all odd numbers not of the type 10k+5 are members.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002744 | OEIS-A083756

Beginning with 5, rearrangement of odd primes such that every concatenation beginning with the first term is a prime.

Conjecture: every odd prime is a member.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002745 | OEIS-A083758

Lexicographically earliest infinite sequence of distinct primes such that the concatenation of the initial n terms is a prime for all n >= 1.

Conjecture: every prime except 5 is a term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002746 | OEIS-A083760

Rearrangement of even numbers such that concatenation of terms beginning with the first with a 1 as the least significant digit is a prime.

Conjecture: every even number is a member.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002747 | OEIS-A083762

Rearrangement of natural numbers such that concatenation of terms beginning with the first one with a 1 as the least significant digit is a prime.

Conjecture: every natural number belongs to this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002748 | OEIS-A083807

Perfect powers using only composite digits 4,6,8,9 and 0.

Call terms trivial if they are of the form 10^k*a(m) (e.g., 400,40000,900,90000). Conjecture: sequence has infinitely many nontrivial terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 08 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002749 | OEIS-A083809

Let f(n) be the smallest prime == 1 mod n (cf. A034694). Sequence gives triangle T(j,k) = f^k(j) for 1 <= k <= j, read by rows.

It has been proved in the reference that for every prime p there exists a prime of the form k*p+1. Conjecture: sequence is infinite, i.e., for every n there exists a prime of the form n*k+1 (cf. A034693).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 08 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002750 | OEIS-A083833

Palindromes p such that 5p + 1 is also a palindrome.

(Conjectures)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002751 | OEIS-A083834

Palindromes of the form 5p + 1 where p is also a palindrome. Palindromes arising in A083833.

(Conjectures)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002752 | OEIS-A083845

a(n)^2 + 1 is largest prime of the form x^2 + 1 <= 10^n.

It is conjectured that the number of primes of the form x^2+1 is infinite and thus this sequence never becomes a constant, but this has not been proved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Harry J. Smith_, May 05 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002753 | OEIS-A083846

a(n) is the largest prime of the form x^2 + 1 <= 10^n.

It is conjectured that the number of primes of the form x^2+1 is infinite and thus this sequence does not become a constant, but this has not been proved. It is easily shown that all terms greater than 5 end in 1 or 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Harry J. Smith_, May 05 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002754 | OEIS-A083848

a(n)^2 + 1 is largest prime of the form x^2 + 1 <= 2^n.

It is conjectured that this sequence is infinite, but this has never been proved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Harry J. Smith_, May 05 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002755 | OEIS-A083849

a(n) is the largest prime of the form x^2 + 1 <= 2^n.

It is conjectured that this sequence is increasing, but this has never been proved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Harry J. Smith_, May 05 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002756 | OEIS-A083863

Consider recurrence b(0) = n/3, b(n) = b(n-1)*floor(b(n-1)); sequence gives first integer reached, or -1 if no integer is ever reached.

It is conjectured that an integer is always reached if the initial value is >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002757 | OEIS-A083942

Positions of breadth-first-wise encodings (A002542) of the complete binary trees (A084107) in A014486.

Conjecture: a(n) = A014138(2^n-2) for n>0. - _Alexander Adamchuk_, Nov 10 2007 Conjecture: a(n) = Sum_{k=1..2^n-1} A000108(k). - _Alexander Adamchuk_, Nov 10 2007 Let h(n) = -((C(2*n,n)*hypergeom([1,1/2+n],[2+n],4))/(1+n)+I*sqrt(3)/2+1/2). Assuming Adamchuk's conjecture a(n) = h(2^n) and A014138(n) = h(n+1). - _Peter Luschny_, Mar 09 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, May 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002758 | OEIS-A083964

Smallest palindromic multiple of 2*n-1 beginning with the digit string of 2*n-1; or 0 if no such number exists.

Conjecture: If n is not of the type (5*k+1)/2 then a(n) is not zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002759 | OEIS-A083965

a(n) = A083964(n)/(2n-1).

Conjecture: If n is not of the form (5k+1)/2 then a(n) is not zero. This conjecture is false (e.g. a(28)=1), see comments in A083964. - _Sean A. Irvine_, Feb 27 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002760 | OEIS-A084013

Smallest n-digit palindromic multiple of n. For n = 10k it is sufficient that the multiple is palindromic with leading zeros ignored. 0 if no such number exists.

Conjecture: no entry is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002761 | OEIS-A084015

Largest n-digit palindromic multiple of n. For n = 10k it is sufficient that the multiple is palindromic with leading zeros ignored. 0 if no such number exists.

Conjecture: no entry is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002762 | OEIS-A084029

Triangle read by rows: n-th row contains the first n n-digit multiples of n with digit sum n. If there are fewer than n such numbers, the rest of the row is filled with 0's.

Conjecture: Only one term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002763 | OEIS-A084030

Main diagonal of triangle A084029.

Conjecture: Only one term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002764 | OEIS-A084043

Smallest palindromic multiple of n in which the digit string of n appears as sandwiched between at least a pair of digits, or 0 if n = 10k or no such number exists.

Conjecture: a(n) = 0 iff n = 10k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002765 | OEIS-A084044

Smallest palindromic multiple of n in which n is a substring (anywhere), or 0 if n = 10k or no such number exists.

Conjecture: a(n) = 0 iff n = 10k. Though many terms match, not the same as A077741.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), May 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002766 | OEIS-A084068

a(1) = 1, a(2) = 2; a(2*k) = 2*a(2*k-1) - a(2*k-2), a(2*k+1) = 4*a(2*k) - a(2*k-1).

Conjecture: a(n) is the earliest occurrence of n in A348295, which is to say, a(n) is the least m such that Sum_{k=1..m} (-1)^(floor(k*(sqrt(2)-1))) = Sum_{k=1..m} (-1)^A097508(k) = n. This has been confirmed for the first 32 terms by _Chai Wah Wu_, Oct 21 2021. - _Jianing Song_, Jul 16 2022 Conjecture: a(n) = gcd(A001108(n), A001109(n)). - _Joseph A. Stocke_, Jul 28 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002767 | OEIS-A084091

Expansion of Sum_{k>=0} x^2^k/(1+x^2^k+x^2^(k+1)).

Dirichlet g.f. (conjectured): L(chi_2(3),s)/(1-2^(-s)), with chi_2(3) the nontrivial Dirichlet character modulo 3. - _Ralf Stephan_, Mar 27 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, May 11 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002768 | OEIS-A084141

a(n) is the index in sequence A084138 when the value of that sequence is 1; i.e., there is exactly one case where there are exactly a(n) primes between m and 2m, exclusively, for m > 0.

This calculation relies on the fact that Pi(2*m) - Pi(m) > m/(3*log(m)) for m >= 5. Conjecture: There are an infinite number of terms in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Harry J. Smith_, May 15 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002769 | OEIS-A084162

a(n) is the length of the gap in sequence A084161.

Conjecture: a(n) < phi(4)*log^2(A268963(n)); A268963 are the end-of-gap primes. Conjecture: a(n) < phi(4)*n^2 for all n > 2. (Note the starting offset 0.) - _Alexei Kourbatov_, Aug 12 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sven Simon_, May 17 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002770 | OEIS-A084189

Minimum digit out of both prime factors of the numbers in A046376.

Conjecture: a(n) is never greater than 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jun 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002771 | OEIS-A084199

Smallest prime q < prime(n) such that q+1 divides prime(n)+1, a(n) = 1 if no such q exists.

Conjecture: A008578 is the range of values (or A000040 = range of values > 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, May 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002772 | OEIS-A084229

Let b(1)=1, b(2)=2, b(n) = sum of digits of b(1)+b(2)+b(3)+...+b(n-1), sequence gives values of n such that b(n)=3.

Conjecture : a(n)/n^3 is bounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jun 21 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002773 | OEIS-A084261

A binomial transform of factorial numbers.

Conjecture: 2*a(n) -4*a(n-1) +(-n+2)*a(n-2) +(n-2)*a(n-3)=0. - _R. J. Mathar_, Nov 30 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002774 | OEIS-A084262

Binomial transform of double factorials.

G.f.: conjecture: T(0)/(1-x), where T(k) = 1 - x*(k+1)/(x*(k+1) - (1-x)/T(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 13 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002775 | OEIS-A084302

Remainder of tau(n) modulo 6.

The sums of the first 10^k terms, for k = 1, 2, ..., are 27, 236, 2275, 22166, 220070, 2195376, 21933228, 219259514, 2192385128, 21923168052, ... Conjecture: the asymptotic mean of this sequence is 3*zeta(3)/zeta(2) = 3 * A253905 = 2.192288... . The conjecture is true if A211337 and A211338 have an equal asymptotic density (see also A059269). - _Amiram Eldar_, Jul 11 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jun 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002776 | OEIS-A084318

Iterate function described in A084317 if started at initial value n until reaching a fixed point.

Conjecture: fixed point always exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jun 16 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002777 | OEIS-A084352

Minimum digit out of both prime factors of palindromic brilliant numbers.

Conjecture: a(n) is never greater than 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jun 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002778 | OEIS-A084391

For n>3, a(n) = smallest number divisible by exactly n-2 previous terms; a(n)=n for n<=3.

G.f. [x * (6x^6-6x^5-4x^3-3x^2-2x-1)]/(1-6x^4) (conjectured). - _Ralf Stephan_, Feb 03 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002779 | OEIS-A084432

Expansion of 2/(1-x) + Sum_{k>=0} t^2(3-t)/(1+t)/(1-t)^2, where t=x^2^k.

Conjecture: a(n) is the number of integer solutions (x,y,z) to the Diophantine equation 2^x*(y+z) = n, where 0 <= x,y,z <= n. - _Joseph M. Shunia_, Aug 27 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Jun 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002780 | OEIS-A084481

Number of fault-free tilings of a 4 X 2n rectangle with L tetrominoes.

If the conjectured G.F. in A183304 is true, then a(n)= 2*A183304(n-1), n>3. - _R. J. Mathar_, Dec 02 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, May 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002781 | OEIS-A084519

Number of indecomposable ground-state 3-ball juggling sequences of period n.

a(n) seems to satisfy the recurrence: a(1) = a(2) = 1, a(3) = 3 and a(n) = 3*a(n-1)+2*a(n-2)+2*a(n-3). If so, a(n) = floor(A*B^n+1/2) where B = 3.6890953... is the real positive root of x^3-3x^2-2x-2 = 0 and A = 0.0687059... is the real positive root of 118*x^3+118*x^2+35*x-3 = 0. - _Benoit Cloitre_, Jun 14 2003 [This conjecture is established in the Chung-Graham paper.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jun 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002782 | OEIS-A084562

Even numbers having no more than two partitions into two odd primes.

Conjecture: there are no more terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, May 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002783 | OEIS-A084569

Partial sums of A084570.

Conjecture: a(n) is the number of perimeter-magic (hollow) squares of order 3 with magic sum n+3. Order 3 means each of the 4 edges has 3 elements >=1; the square has 8 elements. The elements do not need to be distinct, and squares obtained by rotations are counted only once. The square (read ccw) for magic sum 3 has elements 1 1 1 1 1 1 1 1. The 3 squares with magic sum 4 are 1 1 2 1 1 1 2 1, 1 1 2 1 1 2 1 2 and 1 2 1 2 1 2 1 2. - _R. J. Mathar_, Mar 08 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 31 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002784 | OEIS-A084597

Largest k such that there are exactly n primes between k^2 and (k+1)^2.

a(n) is the index of last occurrence of n in A014085. This sequence relies on a heuristic calculation and there is no proof that it is correct. Conjecture: There is no k that has only one prime between k^2 and (k+1)^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Harry J. Smith_, May 31 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002785 | OEIS-A084611

a(n) = sum of absolute values of coefficients of (1+x-x^2)^n.

Limit_{n -> oo} a(n+1)/a(n) does not exist; however, lim_{n -> oo} a(n)^(1/n) = sqrt(5) (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002786 | OEIS-A084616

Maximum number of circles of diameter 1 that can be packed in a square of area n (i.e., with side length n^(1/2)).

Most sequence terms beyond n=20 are only conjectures supported by comprehensive numerical results. No proof is available for the following observations: n=30 is the first case where a square of area < n (29.74921576) is sufficient to cover n circles. The first case where more than n circles can be covered occurs for n=38. The required area to cover 39 circles is 37.76050335. n=59 is the last case where a square of area n does not suffice to cover n+1 circles (60 circles require square area 59.11626524).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002787 | OEIS-A084617

Maximum number of circles with diameter 1 that can be packed in a square of side length n.

For n>5 the sequence terms are only conjectures. For more information see comment given in A084616.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002788 | OEIS-A084618

Maximum number of circles of area 1 that can be packed in a circle of area n.

For most values of n these are only conjectures, supported by numerical results.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002789 | OEIS-A084644

Best packings of m>1 equal circles into a larger circle setting a new density record.

Sequence terms for n>5 are only conjectures. The arrangement of 37 circles consists of one central circle surrounded by 3 rings of 6,12 and 18 circles. For n=7, 49 of the 55 circles are arranged in a rigid hexagonal lattice with 6 "rattlers" inserted in gaps at the circumference.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002790 | OEIS-A084653

Fermat base-2 pseudoprimes whose prime factors do not divide any smaller pseudoprime.

Here pseudoprime means a Fermat base-2 pseudoprime; sequence A001567, a composite number n such that n divides 2^(n-1) - 1. All numbers in this sequence seem to have only two prime factors - a conjecture that has been tested for all pseudoprimes < 10^15. The two prime factors are given in A084654 and A084655. The two prime factors are the same when the pseudoprime is the square of a Wieferich prime (A001220).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jun 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002791 | OEIS-A084663

a(1) = 8; a(n) = a(n-1) + gcd(a(n-1), n).

The proof of the conjecture is identical to the proof in the Rowland link. - _Yifan Xie_, Apr 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | Matthew Frank (mfrank(AT)wopr.wolfram.com) on behalf of the 2003 New Kind of Science Summer School, Jul 15 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002792 | OEIS-A084673

Smallest prime in which a digit appears n times.

For n > 1, conjectured to be equal to A037055(n), the smallest prime in { R-10^n, R-10^(n-1), ..., R-10; R+a*10^b, a = 1, ..., 8, b = 0, 1, 2, ..., n }, where R = (10^(n+1)-1)/9 is the (n+1)-digit repunit. - _M. F. Hasler_, Feb 25 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Harvey P. Dale_, Jun 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002793 | OEIS-A084697

a(1) = 2; for n >= 1, k>=1, a(n+1) = a(n) + k*n is the smallest such prime.

Conjecture: a(n) ~ c n^2 log(n) for some positive constant c. - _Robert Israel_, Oct 26 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jun 05 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002794 | OEIS-A084727

Primes arising in A084726.

Conjecture: All terms exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Jun 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002795 | OEIS-A084743

Smallest prime of the form n^k + k, or 0 if no such prime exists.

a(5) > 5^18 +18 = 3814697265643. Conjecture: No entry is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Jun 15 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002796 | OEIS-A084746

Smallest k such that n^k - k is a prime, or 0 if no such number exists.

Conjecture: no entry is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Jun 15 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002797 | OEIS-A084758

The slowest increasing sequence of primes such that difference of successive terms is unique.

The sequence of successive differences is 1,2,6,8,4,14,10,12,20,18,16,... Conjecture: every even number is a term of this sequence. For every even number e there exists some k such that a(k) - a(k-1) = e.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Jun 17 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002798 | OEIS-A084759

Composite numbers in ascending order such that the difference of successive terms is unique. a(m) - a(m-1) = a(k) - a(k-1) iff m = k.

The sequence of first differences is 2, 3, 1, 4, 6, 5, 7, 8, 9, 11, 10, 12, 13, 15, 14, 16, 18, 17, 19, 20, 21, 23, 22, 24, 25, 26, 27, 28, ... Conjecture: every number is a term of this sequence. For every number r there exists some k such that a(k) - a(k-1) = r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Jun 17 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002799 | OEIS-A084760

Squarefree numbers in ascending order such that the difference of successive terms is unique. a(m) - a(m-1) = a(k) - a(k-1) iff m = k.

The sequence of first differences is 1, 2, 5, 3, 4, 6, 7, 8, 9, 10, 12, 13, 11, 14, 15, 16, 17, 18, 20, 21, 19, ... Conjecture: (1) every number is a term of this sequence. For every number r there exists some k such that a(k) - a(k-1) = r. Question: What is the longest string of consecutive integers in this sequence (of successive differences)?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Jun 17 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002800 | OEIS-A084761

A sequence of primes such that (a(n)-a(n-1)) / (a(n-1)-a(n-2)) is a unique integer.

Conjecture: every number is a term of this sequence, or for every number r there exists some k such that (a(k)-a(k-1)) / (a(k-1)-a(k-2)) = r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Jun 17 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002801 | OEIS-A084824

Maximum number of spheres of diameter one that can be packed in a cube of volume n (i.e., with edge length n^(1/3)).

Higher sequence terms are only conjectures found by numerical experimentation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 12 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002802 | OEIS-A084826

Best packing of m>1 equal spheres in a cube setting a new density record.

The terms >=31 are only conjectures found by numerical experimentation. In the table given at the Pfoertner link, the densities are given relative to the density of the cubic lattice packing (Pi/6). The first known arrangement with higher density than that of the cubic lattice packing was found for m=31 spheres. In the region 8<m<27 increasing packing densities occur for m=9,13,14. m=14 comes very close (0.994949) to the cubic lattice packing density.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 12 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002803 | OEIS-A084827

Maximum number of spheres of volume one that can be packed in a sphere of volume n.

Higher terms of the sequence are only conjectures derived from numerical results. The first 12 arrangements are identical with the solutions of the Tammes problem (see A080865).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002804 | OEIS-A084828

Maximum number of spheres of radius one that can be packed in a sphere of radius n.

a(4) and a(5) are experimental values. Although A121346(5) claims a lower bound of a(5)=68, it is conjectured from extensive numerical search that this value is unachievable and therefore a(5)=67. The conjecture a(5)=67 was proved wrong by Yu Liang, who found an arrangement of 68 spheres of radius 1 fitting into a sphere of radius 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 12 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002805 | OEIS-A084829

Best packing of m>1 equal spheres in a sphere setting a new density record.

All terms beyond m=9 are only conjectures found by numerical experimentation. The density is defined as the fraction of the volume of the large sphere occupied by the small spheres. For 2 spheres the density is 0.25. The first known configuration with density exceeding 0.5 occurs for 31 spheres.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 12 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002806 | OEIS-A084865

Primes of the form 2x^2 + 3y^2.

Conjecture: A084863(a(n))=1? Comment: The truth of the conjecture A084863(a(n))=1 follows from the genus theory of quadratic forms (see Cox, page 61). By comparing enough terms, we see that the conjecture a(n) = A019338(n+1) is false. - _T. D. Noe_, May 02 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jun 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002807 | OEIS-A084923

Consider the sequence b(1) = n, b(k) is the greatest prime factor of 3*b(k-1)+2. It is conjectured that this always becomes cyclic; a(n) = length of cycle (or 0 if no cycle is reached).

Consider the sequence b(1) = n, b(k) is the greatest prime factor of 3*b(k-1)+2. It is conjectured that this always becomes cyclic; a(n) = length of cycle (or 0 if no cycle is reached).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_ & _Jud McCranie_, Jun 11 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002808 | OEIS-A084950

Array of coefficients of denominator polynomials of the n-th approximation of the continued fraction x/(1+x/(2+x/(3+..., related to Laguerre polynomial coefficients.

Conjecture: also coefficient triangle of the denominators of the (n-th convergents to) the continued fraction w/(1+w/(2+w/(3+w/... This continued fraction converges to 0.697774657964... = Bessell(1,2)/Bessell(0,2) for w=1. For instance, the denominator of w/(1 + w/(2 + w/(3 + w/(4 + w/5)))) equals 120 + 96*w + 9*w^2. - _Wouter Meeussen_, Aug 08 2010 The continued fraction 0 + K_{k>=1}(x/k) = x/(1+x/(2+x/(3+... has n-th approximation P(n,x)/Q(n,x). These polynomials satisfy the recurrence q(n,x) = n*q(n-1,x) + x*q(n-2,x), for q replaced by P or Q with inputs P(-1,x) = 1, P(0,x) = 0 and Q(-1,x) = 0 and Q(0,1) = 1. The present array provides the Q-coefficients: Q(n,x) = sum(a(n,m)*x^m, m=0 .. floor(n/2)), n >= 0. For the P(n,x)/x coefficients see the companion array A221913. This proves the first part of W. Meeussen's conjecture given above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Jun 14 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002809 | OEIS-A085044

Smallest number k such that tau(n) +tau(k) =tau(n+k), or 0 if no such number exists.

Conjecture: No entry is zero. If n = p^2 where p is an odd prime then a(n) < p^2 or a(n) = p^2 as tau(2p^2) = 6 = tau(p^2) + tau(p^2). The (n,k) pairs are given below. (1,3),(2,10),(3,1),(4,841),(5,3),(6,66),(7,3),(8,37),(9,9),(10,2),(11,3),... Subsidiary sequence:(1) members of this sequence such that a(n) = n. E.g. a(9) = 9. (2)(harder one) Smallest k such that sigma(n) +sigma(k) = sigma(n+k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Jun 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002810 | OEIS-A085057

a(n) is the smallest integer of the form a*b*c.../p*q*r..., where the numerator and the denominator contain n numbers each and a,b,c,...p,q,r... are all the integers from 1 to 2n.

Conjecture: a(n) always equals that lower bound. - _Don Reble_, Jul 01 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Jun 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002811 | OEIS-A085068

Number of steps >= 1 for iteration of map x -> (4/3)*ceiling(x) to reach an integer when started at n, or -1 if no such integer is ever reached.

It is conjectured that an integer is always reached.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 11 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002812 | OEIS-A085094

Smallest k such that n*k-1 is a palindrome, or 0 if no such number exists.

Conjecture: No entry is zero. For every n there exists a k such that n*k-1 is a palindrome.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Jul 03 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002813 | OEIS-A085096

Index of the first occurrence of n in A082744, or 0 if n does not occur in the sequence A082744.

Conjecture: a(11) = 0, or in other words,if 11k+1 is a palindrome then at least one of the numbers in the sequence k+1,2k+1, 3k+1,...,10k+1 is also a palindrome. E.g. for k = 10 and k = 55, 11k+1 is a palindrome but 10*1 +1 and 55*2 +1 are also palindromes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Jul 03 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002814 | OEIS-A085104

Primes of the form 1 + n + n^2 + n^3 + ... + n^k, n > 1, k > 1.

The conjecture proposed in Quadrature "No Sophie Germain prime is Brazilian (prime)" (see link Bernard Schott, Quadrature, Conjecture 1, page 36) is false. Thanks to _Giovanni Resta_, who found that a(856) = 28792661 = 1 + 73 + 73^2 + 73^3 + 73^4 = (11111)_73 is the 141385th Sophie Germain prime. - _Bernard Schott_, Mar 08 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Jul 03 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002815 | OEIS-A085105

Starting with prime(n) go on adding smaller primes until one gets a prime. a(n) = this prime, or 0 if no such prime exists.

Conjecture: Except a(3) no other term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Jul 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002816 | OEIS-A085106

Starting with composite(n) go on adding smaller composite numbers until one gets a prime. a(n) = this prime, or 0 if no such prime exists.

Conjecture: No entry is zero for n >10. There are only four terms which are zero. The conjecture is false as a(31) is zero. There is, however, no further zero up to a(26754), so the conjecture may be rephrased as: no entry is zero for n>31 and there are only five terms which are zero. - _Harvey P. Dale_, May 04 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Jul 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002817 | OEIS-A085108

Decimal expansion of sum 1/(p*2^p) where p runs through the set of Artin primes (primes with primitive root 2).

David H. Bailey and Richard E. Crandal conjectured that this constant is 2-normal.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002818 | OEIS-A085114

Binary expansion of Sum 1/(p*2^p) where p runs through the set of Artin primes (primes with primitive root 2).

David H. Bailey and Richard E. Crandal conjectured that this constant is 2-normal.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002819 | OEIS-A085123

Smallest member of A002473 whose digital product = A002473(n), or 0 if no such number exists.

Conjecture: No entry is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002820 | OEIS-A085124

Smallest multiple of n whose digital product is also a positive multiple of n, or 0 if no such number exists.

Conjecture: otherwise, a(n) is positive.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002821 | OEIS-A085135

Numbers whose cyclic permutations are all 7-smooth (A002473).

Sequence is infinite with trivial terms of the form k*10^r, k = 1 to 9. Conjecture: There are finitely many nontrivial terms (not of the form k*10^r). All the two-digit terms are the same as those of A085133.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and Meenakshi Srikanth (menakan_s(AT)yahoo.com), Jul 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002822 | OEIS-A085245

Least k such that k*2^n + 1 is a semiprime.

The first few values of n such that 78557*2^n + 1 is a semiprime, where k = 78557 (the conjectured smallest Sierpinski number), are: 2, 3, 7, 15, 17, 18, 24, 60, 71, 89, 92, 107, 140, 143, 163,... Conjecture: there are infinitely many semiprimes of this form.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Aug 11 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002823 | OEIS-A085260

Ratio-determined insertion sequence I(0.0833344) (see the link below).

It appears that the b-file, formulas and programs are based on the conjectured, so far apparently unproved recurrence relation. - _M. F. Hasler_, Nov 05 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jun 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002824 | OEIS-A085263

Number of ways to write n as the sum of a squarefree number (A005117) and a positive square (A000290).

First occurrence of k: 2, 6, 11, 23, 30, 38, 62, 71, 83, 110, 138, 155, 182, 203, 227, 263, 302, 327, 383, 435, 447, 503, 542, 602, 635, ..., . Conjecture: For each k above, there is a finite number of terms; for example, only the two numbers 1 and 13 cannot be represented as the sum of a squarefree number and a square. The number of k terms beginning with 0: 2, 9, 19, 27, 38, 36, 57, 63, 62, 74, 94, ..., . - _Robert G. Wilson v_, May 16 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jun 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002825 | OEIS-A085264

Smallest number with exactly n representations as sum of a squarefree number (A005117) and a square (A000290).

Conjecture: Just as there is a least integer that can be represented in n ways, so is there a greatest integer. Conjecture: The last occurrence of k, beginning with 0, in A085263: 13, 61, 85, 196, 225, 441, 621, 909, 1089, 1125, 1521, 2025, 2700, 2200, 2925, 3969, 3825, 4500, 5625, 4869, 6084, 8100, 11025, 7425, 9900, 9981, 10584, 11925, 12825, 14400, 13500, 14625, 18081, 18225, 17424, 20925, 22500, 27225, 21825, 25425, 27000, 28224, 27900, 38025, 44100, 33300, 35721, 35325, 39825, 37044, 39600, 40725, 44325, 55125, 50625, 53100, 52200, 54000, 60300, 65025, 63900, 60025, 63504, 64125, 74529, 81225, 77400, 99225, 88200, 76500, 79200, 87525, 90000, 108900, 88425, 91800, 95400, 96300, 100125, 107325, 132300, ..., . Conjecture: For each j, there is a finite number of positive integers that can be represented as the sum of a squarefree number and a square in exactly j ways; e.g., for j=0, only the two integers 1 and 13 cannot be represented as the sum of a squarefree number and a square.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jun 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002826 | OEIS-A085265

Numbers that can be written as sum of a positive squarefree number and a positive square.

Conjecture: a(n) = n + 2 for n > 11. That is, only 1 and 13 are missing. - _Charles R Greathouse IV_, Aug 21 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jun 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002827 | OEIS-A085267

Numbers having at least two representations as sum of a squarefree number and a nonzero square.

Conjecture: for n > 50, a(n) = n + 11. - _Charles R Greathouse IV_, Oct 26 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jun 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002828 | OEIS-A085298

a(n) is the smallest exponent x such that prime(n)^x when reversed is a prime.

It is conjectured that for every n such exponent exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jun 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002829 | OEIS-A085331

Numbers k such that phi(rev(k)) = k.

rev(2*(10^k-4)) = 3*(10^k-3). If 10^k-3 is prime, then phi(3*(10^k-3)) = 2*(10^k-4), so 2*(10^k-4) is a term. 10^1-3=7 is prime, so 2*(10^1-4)=12 is a term, a(2). 10^2-3=97 is prime, so 2*(10^2-4)=192 is a term, a(4). 10^3-3=997 is prime, so 2*(10^3-4)=1992 is a term, a(5). 10^17-3 is prime, so 2*(10^17-4)=199999999999999992 is a term. 10^140-3 is prime, so 2*(10^140-4) is a term. 10^990-3 is prime, so 2*(10^990-4) is a term. Conjecture: sequence is infinite. - _Ray Chandler_, Jul 20 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jul 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002830 | OEIS-A085376

Ratio-dependent insertion sequence I(0.36704) (see the link below).

Conjecture: partial sums of A129445. - _Sean A. Irvine_, Jul 14 2022 It is conjectured that a(n) = 10*a(n-2) - a(n-4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jun 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002831 | OEIS-A085387

E.g.f exp(x)*cosh(x+x^2/2).

Conjecture: a(n) -3*a(n-1) +2*a(n-2) +(-2*n+3)*a(n-3) +-(n-3)^2*a(n-4) +(n-3)*(n-4)*a(n-5)=0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jun 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002832 | OEIS-A085420

For each n, let p(n,b) be the smallest prime in the arithmetic progression k*n+b, with k > 0. Then a(n) = max(p(n,b)) with 0 < b < n and gcd(b,n) = 1.

Linnik proved that there are n0 and L such that a(n) < n^L for all n > n0. It has been conjectured that a(n) < n^2. The sequence A034694 has the primes p(n,1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jun 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002833 | OEIS-A085427

Least k such that k*2^n - 1 is prime.

Conjecture: for every n there exists a number k < 3n such that k*2^n - 1 is prime. Comment from _T. D. Noe_: this fails at n=624, where a(n)=2163.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Aug 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002834 | OEIS-A085450

a(n) is the smallest m > 1 such that m divides Sum_{k=1..m} prime(k)^n.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Aug 14 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002835 | OEIS-A085495

Maximal order of a finite subgroup of the group GL(n,Q).

Conjecture: a(n)/2 is the maximal order of a finite subgroup of the group SL(n,Q) (or equivalently, SL(n,Z), since a finite subgroup of SL(n,Q) must be conjugate to a subgroup of SL(n,Z)). This is obviously true for odd n: given a finite subgroup G of SL(n,Q), a subgroup of GL(n,Q) of order 2*|G| can be attained by adjoining -I_n where I_n is the identity matrix. The conjecture is also correct for n = 2, since the finite subgroup of the maximal order of SL(n,Q) is isomorphic to C_6. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Yuval Dekel (dekelyuval(AT)hotmail.com), Aug 15 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002836 | OEIS-A085508

Decimal expansion of conjectured value for the Bloch constant.

Decimal expansion of conjectured value for the Bloch constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Jul 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002837 | OEIS-A085574

Least k such that n*(n+1)*(n+2) + k is palindromic.

Conjecture: For all n > 108, a(n) is > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jul 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002838 | OEIS-A085577

Size of largest subset of the n^2 cells in an n X n grid such that there are at least 3 edges between any pair of chosen cells.

Conjecture: if n == 4 (mod 5), then a(n)=(n^2+4)/5. - _Erich Friedman_, Apr 19 2015 More general conjecture: if n != 5, then a(n) = ceiling(n^2/5). - _Rob Pratt_, Jul 10 2015 Conjecture holds for n <= 70. - _Giovanni Resta_, Jul 29 2015 The conjectures are correct. See Eq. (1) of Fisher and Beel for the 2-packing number of grid graphs. - _Andrei Zabolotskii_, Jul 13 2026 Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2003; entry revised Apr 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002839 | OEIS-A085579

a(n) is the n-th digit after the decimal point of the positive real root of x^2+n*x-2.

Conjecture: this diagonal expressed as a decimal is irrational and transcendental.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jul 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002840 | OEIS-A085640

Resultant of the polynomial x^3-1 and the Chebyshev polynomial of the first kind T_n(x).

G.f.: (1+3*x+3*x^2-x^3)/(1-4*x-6*x^2-4*x^3+x^4) (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Yuval Dekel (dekelyuval(AT)hotmail.com), Aug 15 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002841 | OEIS-A085642

Number of columns in the character table of the symmetric group S_n that have zero sum.

Conjecture: Equals the number of partitions of n with at least one part congruent to 2 mod 4. - _Vladeta Jovovic_, Jul 12 2003. This conjecture was established by _Christine Bessenrodt_ and Jorn B. Olsson (olsson(AT)math.ku.dk), Sep 13 2004.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Yuval Dekel (dekelyuval(AT)hotmail.com), Jul 11 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002842 | OEIS-A085710

Smallest k such that kn+1 is a semiprime.

Conjecture: for every n > 2 there exists a number k < n such that n*k + 1 is a semiprime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jul 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002843 | OEIS-A085765

Partial sums and bisection of A086450.

Conjecture: log(a(n))/log(n) grows unboundedly. Conjecture: a(n) mod 2 repeats the 7-pattern 0,0,1,1,1,0,1. The conjecture concerning the mod 2 pattern follows directly from the corresponding conjecture proved in A086450. - Lambert Herrgesell (zero815(AT)googlemail.com), May 08 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Jul 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002844 | OEIS-A085774

Numbers k such that (k / sum of digits of k) and (k+1 / sum of digits of k+1) are both semiprime.

Note the quadruple: 12255,12256,12257,12258 and the triple: 20174,20175,20176. Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jul 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002845 | OEIS-A085780

Numbers that are a product of 2 triangular numbers.

Conjecture: There are about sqrt(x)*log(x) terms up to x. - _Charles R Greathouse IV_, Jul 11 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Jul 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002846 | OEIS-A085785

Start at (2n+1)/4 and iterate the map x -> x*ceiling(x); sequence gives number of steps for denominator to drop to 1 or 2; or -1 if this never happens.

We conjecture that the denominator always does drop.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 16 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002847 | OEIS-A085794

Lexicographically earliest squarefree infinite ternary word.

Open Problem 1.10.2 in the Allouche & Shallit reference asks for a good alternate characterization of this sequence. (Is it morphic, if so for which morphism?)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Claude Lenormand (claude.lenormand(AT)free.fr), Jul 24 2003; corrected Jul 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002848 | OEIS-A085799

Determinant of the symmetric n X n matrix A defined by A[i,j] = abs(i^2 - j^2) for 1 <= i,j <= n.

For n>=2, a(n) = (-1)^(n-1) * (n+1)/2 * (2*n-1)! / (n-2)!. Furthermore, the conjecture by Lajos is true (see Fried link). - _Sela Fried_, Feb 04 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | Yuval Dekel (dekelyuval(AT)hotmail.com), Jul 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002849 | OEIS-A085800

Maximal number of zeros in a row of the character table of the symmetric group S_n.

Conjecture: For k >= 6 and n = k*(k+1)/2 + 2, a(n) is attained precisely by the two rows of the character table of S_n indexed by (k+2, k-1, k-2, ..., 2, 1) and its conjugate. Verified for 6 <= k <= 11, giving a(23) = 1110, a(30) = 5219, a(38) = 24899, a(47) = 121544, a(57) = 604559, and a(68) = 3058703. - _Vladeta Jovovic_, Oct 17 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | Yuval Dekel (dekelyuval(AT)hotmail.com), Jul 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002850 | OEIS-A085910

a(n) = smallest prime(k) not equal to prime(n) and not included earlier, such that A001223(k) = A001223(n); i.e., prime(k+1) - prime(k) = prime(n+1) - prime(n); or 0 is no such prime exists.

Conjecture: a rearrangement of odd primes for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002851 | OEIS-A085928

a(1) = 1, a(2) = 2, then the absolute difference of the concatenation of two previous terms and its digit reversal.

a(n) is a multiple of 99 for n > 4. Conjecture: The sequence has finitely many nonzero terms. (This would happen when the concatenation of two successive terms becomes a palindrome and the next term is 0.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_ and _Jason Earls_, Jul 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002852 | OEIS-A085929

Smaller of two consecutive numbers with the same prime signature not occurring earlier.

Conjecture: (1) The sequence is infinite. (2) For every prime signature other than perfect powers, there exists a number k such that k+1 also has the same prime signature. Up to at least 3*10^8, all numbers in this sequence have signatures with some prime to power 1 (for example, p^2*q^3 does not appear), casting some doubt on the second conjecture. - _Joseph Myers_, Feb 24 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 14 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002853 | OEIS-A086004

Primes which remain prime after one and after two and after three applications of the rotate-and-add operation of A086002.

Conjecture 1: Rotation and addition of primes with even numbers of digits never yields a prime. Conjecture 2: There are no 5-Rotation Cycle Primes. [Conjecture 1 is true because rotation for even numbers of the form 10^k*a+b yields 10^k*b+a, so rotation-and-add yields (10^k+1)*(a+b), which obviously contains a divisor A000533. RJM, Sep 17 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chuck Seggelin_, Jul 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002854 | OEIS-A086018

Number of cyclic numbers (A001913) <= 10^n.

Conjectured ratio a(n)/A006880(n) as n->infinity is Artin's constant 0.3739558136...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Jul 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002855 | OEIS-A086057

Decimal expansion of a constant related to a conjectured value of the Masser-Gramain constant.

Decimal expansion of a constant related to a conjectured value of the Masser-Gramain constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Jul 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002856 | OEIS-A086058

Decimal expansion of conjectured value of delta related to the Masser-Gramain constant.

Decimal expansion of conjectured value of delta related to the Masser-Gramain constant. Numerical work by Melquiond et al. (see reference) disproves Gramain's conjecture, correct bounds for the Masser-Gramain constant delta are: 1.819776 < delta < 1.819833. - _Vaclav Kotesovec_, Apr 27 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Jul 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002857 | OEIS-A086097

Numbers n such that n and n+1 are semiprimes with a semiprime number of 1's in their binary representation.

3721 is the first square entry. Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jul 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002858 | OEIS-A086140

Primes p such that three (the maximum number) primes occur between p and p+12.

A086140 is the union of A022006 and A022007. By merging the two b-files I have extended the current b-file up to n=10000 (nearly n=20000 would have been possible). I add a comparison (see Links) between the frequency of prime 5-tuples and an asymptotic approximation, which is unproven but likely to be true, and based on a conjecture first published by Hardy and Littlewood in 1923. Twins, triples and quadruplets are treated as well. - _Gerhard Kirchner_, Dec 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jul 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002859 | OEIS-A086144

a(n) = 2*A071640(n) - n.

It is conjectured that A071640(n)/n -> 1/2. - _Benoit Cloitre_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Sep 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002860 | OEIS-A086205

Determinant of n X n matrix M_(i,j) = binomial(i^2, j).

Closed form can be deduced from the conjecture that the count of triangular semi-standard Young tableaux with shape (n,...,1) and max part n equals 2^((n-1)n/2); see Mathematica line. - _Wouter Meeussen_, Nov 26 2017 Conjecture: a(n) = the determinant of the n X n matrix (binomial(i^2 + z, j)) for arbitrary complex z. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002861 | OEIS-A086212

Primes in A071998.

Conjecture: there is an infinity of primes in A071998.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Sep 08 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002862 | OEIS-A086222

a(n) = card{ (x,y,z) | x <= y <= z and lcm(x,y,z) = n }.

A number of conjectures are possible, many of which should be easy to prove. Examples: (1) If n is a product of two primes then a(n)=13. (2) If n is a square of a prime then a(n)=6. - _John W. Layman_, Sep 01 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | Yuval Dekel (dekelyuval(AT)hotmail.com), Aug 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002863 | OEIS-A086377

a(1)=1; a(n)=a(n-1)+2 if n is in the sequence; a(n)=a(n-1)+2 if n and (n-1) are not in the sequence; a(n)=a(n-1)+3 if n is not in the sequence but (n-1) is in the sequence.

The three conjectures by respectively Biberstein, Hanna, and Kimberling have all been proved, see the paper by Bosma et al. in the Links. - _Michel Dekking_, Oct 05 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Sep 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002864 | OEIS-A086398

a(1)=1; a(n)=a(n-1)+2 if n is in the sequence; a(n)=a(n-1)+2 if n and (n-1) are not in the sequence; a(n)=a(n-1)+4 if n is not in the sequence but (n-1) is in the sequence.

Conjecture: the positions of 1 in the fixed point of the morphism 0 -> 10, 1 -> 1000, and -1 < n*(1 + sqrt(3)) - a(n) < 4 for n>=1; see A285301. - _Clark Kimberling_, Apr 25 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Sep 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002865 | OEIS-A086411

Greatest prime factor of 3-smooth numbers.

Conjecture: a(n) = A049237(n+1) for n>1. - _R. J. Mathar_, Jun 06 2024 Conjecture is true. This is due to the facts that A007694(n+1) = 2*A003586(n), A049237(n) = A006530(A007694(n)), and A006530(x) = A006530(2x) for x>1 a term in A003586. - _Chai Wah Wu_, Mar 12 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jul 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002866 | OEIS-A086449

a(0) = 1, a(2n+1) = a(n), a(2n) = a(n) + a(n-1) + ... + a(n-2^m) + ... where a(n) = 0 for n < 0.

Conjecture: all a(n) are even except a(2^k-1) = 1. Also a(2^k-2) = 2^(k-1). [For proof see link.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Jul 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002867 | OEIS-A086450

a(0) = 1, a(2n+1) = a(n), a(2n) = a(n) + a(n-1) + ... + a(n-m) + ... where a(n<0) = 0.

Conjecture: a(n) mod 2 repeats the 7-pattern 1,1,0,1,0,0,1 (A011657). The conjecture is easily proved by induction: a(0) to a(14) = 1, 1, 2, 1, 4, 2, 5, 1, 9, 4, 11, 2, 16, 5 read mod 2 gives 1, 1, 0, 1, 0, 0, 1, 1, 1, 0, 1, 0, 0, 1. Assume the conjecture is true up to n = 14k. Then the next 7 odd entries a(14k+1), a(14k+3), ..., a(14k+13) are read from a(7k) to a(7k+6), which follow the correct mod 2 pattern by assumption. For the even entries a(14k), a(14k+10)... a(14k+12), the sum over the first 7k-1 addends is even, simply because of each consecutive 7 addends exactly 4 are odd. So again a(7k) to a(7k+6) determines the outcome and again gives the desired pattern. a(14k) is odd, since a(7k) is odd, a(14k+2) is even, since a(7k) and a(7k+1) are odd and so on ... - Lambert Herrgesell (zero815(AT)googlemail.com), May 08 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Jul 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002868 | OEIS-A086469

Sum of the distinct (smallest) prime signature divisors of n. In case of two or more divisors with the same prime signature the smallest is considered to evaluate the sum. Let this function be defined as psigma(n).

Define n as a 'psigma perfect number' if psigma(n) = 2n. 18 is a psigma perfect number. The p sigma divisors are 1,2,6,9 and 18 and the sum = 36. Conjecture: 18 is the only psigma perfect number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 21 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002869 | OEIS-A086519

a(n) is the least prime not already used such that (a(n)+a(n-1))/2 is prime.

Second term onwards every prime == 1 (mod 6). Conjecture: every prime of the type 6k+1 is a member. After the first term, every member = 7 (mod 12). Conjecture: every prime of the form 12k+7 is a member. - _David Wasserman_, Mar 11 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002870 | OEIS-A086522

Primes arising as the arithmetic mean of a pair of successive terms of A086519.

Conjecture: every prime of the type 6k+1 is a member. Comment from _Vim Wenders_, May 27 2008: The conjecture is worng. For example 19 is missing..

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002871 | OEIS-A086523

Beginning with 5, distinct odd primes such that the arithmetic mean of every pair of successive terms is prime.

Conjecture: every prime of the form 6k-1 is a member. Comment from _Vim Wenders_, May 27 2008: The conjecture is wrong. For example 11 and 23 are missing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002872 | OEIS-A086524

Primes arising as the arithmetic mean of a pair of successive terms of A086523.

Every term == -1 (mod 6). Conjecture: every prime of the form 6k-1 is a member.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002873 | OEIS-A086528

Smallest prime which occurs exactly n times in the sequence A086527.

Conjecture: The sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 31 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002874 | OEIS-A086534

Smallest prime p sandwiched between two numbers that are divisible by n-th powers.

Conjecture: sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 17 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002875 | OEIS-A086537

Beginning with 1, the smallest number not yet seen such that every partial sum has a distinct prime signature.

Conjecture: this is a rearrangement of natural numbers (i.e., every natural number is a term).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002876 | OEIS-A086538

n-th partial sum of A086537.

Every term has a unique prime signature. Conjecture: Every prime signature occurs in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002877 | OEIS-A086550

Smallest k such that tau(k) - tau(k-1) = n, where tau(k) = number of divisors of k, or 0 if no such number exists.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002878 | OEIS-A086551

a(n) = smallest k such that tau(k) = n*tau(k-1) where tau(k) = number of divisors of k, or 0 if no such number exists.

Conjectures: (1) No term is zero. (2) a(n)-1 is a prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002879 | OEIS-A086552

Numbers x such that tau(x)/tau(x-1) is an integer, where tau() is the number of divisors function.

Conjecture: (1) tau(x)/tau(x-1) = n has solutions for every n. (2) If x is the smallest number for a given n such that tau(x)/tau(x-1) = n > 1, then x-1 is a prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002880 | OEIS-A086553

a(1) = 1, a(n) = smallest number not included earlier such that both a(n) + 1 and a(n)*a(n-1) + 1 are primes.

Conjecture: every number p-1 is a term for all primes p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002881 | OEIS-A086554

a(1) = 1; for n>1, a(n) = smallest number not included earlier such that each P+1 is prime, where P is the product of an (n-1)-set from the previous n terms.

Conjecture: sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002882 | OEIS-A086558

Smallest n-digit number beginning with n and having prime signature of n, or 0 if no such number exists.

Conjecture: no term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002883 | OEIS-A086566

a(1)=2; for n>1, a(n) is the largest prime number m such that a(n-1)^(1/(n-1)) > m^(1/n).

For this subsequence of prime numbers the sequence a(n)^(1/n) is decreasing and with this property a(n)-a(n-1) is maximal. Note that conjecture 30 of www.primepuzzles.net says A000040(n)^(1/n) is decreasing. But A000040(n)-A000040(n-1) is not maximal.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Sep 11 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002884 | OEIS-A086642

Maximal number of zeros in a column of the character table of the symmetric group S_n.

The conjecture has been computationally verified for 3 <= n <= 200. Every column of the character table of S_n was certified to contain at least n-1 nonzero entries, while the column corresponding to cycle type (n-1,1) has exactly n-1 nonzero entries. - _Vladeta Jovovic_, Oct 17 2026 Conjecture: a(n) = A000041(n)-n+1, n>2. - _Vladeta Jovovic_, Jul 28 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | Yuval Dekel (dekelyuval(AT)hotmail.com), Jul 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002885 | OEIS-A086685

Number of 1 <= i < n such that i*n+1 is prime.

Number of primes p < n^2 such that p == 1 (mod n). The standard conjecture here is that a(n) ~ n^2/(2 phi(n)log n), where Euler's totient function phi(n) = A000010(n). - _Thomas Ordowski_, Oct 21 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Jul 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002886 | OEIS-A086711

Primes p such that A001414(p-1) = A001414(p+1), where A001414 = sum of primes dividing n (with repetition).

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jul 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002887 | OEIS-A086744

Decimal expansion of Product_{n>=1} (2n/(2n+1))^((-1)^t(n)), where t(n) = A010060(n) is the Thue-Morse sequence.

It is an open problem to decide if this number is algebraic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 12 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002888 | OEIS-A086753

Number of distinct entries in a slice of A046816.

Conjectured asymptotic: a(n) ~ (n^2 + 6*n - 12)/12. - _Vladimir Reshetnikov_, Dec 28 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Jul 31 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002889 | OEIS-A086811

Average (scaled by a certain explicit factor) over all integers k of a_k(n), the n-th coefficient of the k-th cyclotomic polynomial.

When n is odd the n-th term is an integer. If n is even then twice the n-th term is an integer. Conjecturally (Y. Gallot) the n-th term is always an integer. For n <= 128 this has been verified numerically by Yves Gallot. It is also an unproved conjecture due to H. Möller (1970) that no term of this sequence is negative.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Pieter Moree (moree(AT)mpim-bonn.mpg.de), Aug 05 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002890 | OEIS-A086818

a(1) = 111, a(n) = the smallest squarefree number > a(n-1) which contains all the digits of a(n-1).

Conjecture: Terms contain exactly three 1's digits and the rest 0's digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ray Chandler_, Aug 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002891 | OEIS-A086824

Least positive k such that k! >= n^k.

Conjecture: for n>3 a(n) = round(e*n-(1/2)*log(2*Pi*n)-1/n). - _Benoit Cloitre_, Dec 14 2005 Above conjecture is false: For n = 195 we have: a(n) = 526 < 527 = round(exp(1)*n -(1/2)*log(2*Pi*n)-1/n). - _Alois P. Heinz_, Jan 15 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Aug 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002892 | OEIS-A086844

Odd numbers m such that the sequence defined by b(1) = m; for k>1, b(k) = floor((1+sqrt(3))*b(k-1)) contains only odd numbers.

Conjecture: let r(z)= (1/2) *(z+sqrt(z^2+4*z)), for any integral z>=1. Then the sequence a(n)-4n (where a(n) is the sequence of odd numbers m such that the sequence defined by b(1) = m; for k>1, b(k) = floor(r(z)*b(k-1)) contains only odd numbers) becomes ultimately periodic. _Benoit Cloitre_, Aug 10 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Aug 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002893 | OEIS-A086845

a(1) = 0, a(n) = a(floor(n/2)) + 2*a(ceiling(n/2)) + floor(n/2).

Conjecture: partial sums of A087808. - _Sean A. Irvine_, Jul 14 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Aug 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002894 | OEIS-A086848

First differences are formed by interleaving {2^k - 1, k = 1, 2, 3, ...} and {11*k, k = 5, 4, 3, ...}.

Conjectured g.f. above confirmed. Interleaving of two linear recurrences is another linear recurrence, as is the first difference. - _Ray Chandler_, May 07 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | Michael M. Taylor, Aug 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002895 | OEIS-A086853

Number of permutations of length n with exactly 2 rising or falling successions.

Conjecture: (-514*n+2465)*a(n)
+2*(257*n^2-955*n-1085)*a(n-1)
+(-555*n^2+2483*n-1670)*a(n-2)
+16*(-17*n^2+73*n-75)*a(n-3)
+(354*n^2+528*n-2299)*a(n-4)
+2*(-121*n^2+1045*n-1401)*a(n-5)
+3*(67*n-115)*(n-4)*a(n-6)=0. - _R. J. Mathar_, Jun 06 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002896 | OEIS-A086892

Greatest common divisor of 2^n-1 and 3^n-1.

a(n) is a simple (the simplest?) example of a divisibility sequence associated to a rational point on an algebraic group of dimension larger than two. Specifically, it is the divisibility sequence associated to the point (2,3) on the two-dimensional torus G_m^2. Ailon and Rudnick conjecture that a(n) = 1 for infinitely many n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Joseph H. Silverman (jhs(AT)math.brown.edu), Sep 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002897 | OEIS-A086920

Smallest k such that n times concatenation of k with itself followed by a 1 is a prime, or 0 if no such number exists.

Conjecture: no term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002898 | OEIS-A086922

Number of idempotent n X n (0,1) matrices over the reals.

Conjecture: There is no closed form expression for this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Yuval Dekel (dekelyuval(AT)hotmail.com), Sep 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002899 | OEIS-A087037

Smallest integer x > 0 such that x^x + n is prime, or 0 if no such x exists.

It is conjectured that such x always exists. - _Dean Hickerson_ We can show that for all n=(6k-1)^3, k > 0, there is no such x, which disproves the conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jul 31 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002900 | OEIS-A087038

Smallest integer x > 1 such that x^x + n is prime, or 0 if no such x exists.

It is conjectured that such x always exists. - _Dean Hickerson_ We can show that for all n=(6k-1)^3, k > 0, there is no such x, which disproves the conjecture. See the main entry A087037 for more details. - _Farideh Firoozbakht_ and _M. F. Hasler_, Nov 27 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jul 31 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002901 | OEIS-A087092

Descending squares: which squares can be packed with at least 2 squares so that every square directly above a square is strictly smaller? It is conjectured that the answer is all except those in this sequence.

Descending squares: which squares can be packed with at least 2 squares so that every square directly above a square is strictly smaller? It is conjectured that the answer is all except those in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Erich Friedman_, Oct 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002902 | OEIS-A087094

a(n) = smallest k such that (10^k-1)/9 == 0 mod prime(n)^2, or 0 if no such k exists.

The union of the collection of sequences formed from the nonzero terms of a(n)*A000027, gives the values of k for which (10^k-1)/9 is not squarefree, see A046412. All of terms of the sequence a(n) are integer multiples of prime(n) for primes <1000 except for a(93)=486 where prime(93)=487. Conjecture: there are no 0 terms after a(3). That conjecture is easily proved, for a(n) is just the multiplicative order of 10 modulo (prime(n))^2 for n>3. - _Jeppe Stig Nielsen_, Dec 28 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ray Chandler_, Aug 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002903 | OEIS-A087107

This table shows the coefficients of combinatorial formulas needed for generating the sequential sums of p-th powers of tetrahedral numbers. The p-th row (p>=1) contains a(i,p) for i=1 to 3*p-2, where a(i,p) satisfies Sum_{i=1..n} C(i+2,3)^p = 4 * C(n+3,4) * Sum_{i=1..3*p-2} a(i,p) * C(n-1,i-1)/(i+3).

Conjectural formula for table entries: T(n,k) = Sum_{j = 0..k} (-1)^(k+j)*binomial(k,j)*binomial(j+3,3)^n. Conjecturally, the n-th row polynomial R(n,x) = 1/(1 + x)*Sum_{i >= 0} binomial(i+3,3)^n *(x/(1 + x))^n. (End) The conjectures above are correct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _André F. Labossière_, Aug 11 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002904 | OEIS-A087192

a(n) = ceiling(a(n-1)*4/3), with a(1) = 1.

Conjecture: There is a constant k (with a value of about 1.557296798) such that, for all n, 0<k*(4/3)^n-a(n)<2. This conjecture has been verified for all n up to 10^6. - _Clive Tooth_, Dec 24 2025 The conjecture above is correct. In fact, the constant k is (3/4)*A390321.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002905 | OEIS-A087200

a(n) is the smallest m such that m > A005235(n) and A002110(n)+m is prime.

a(n) is the second m (first m is A005235(n)) such that m > 1 and A002110(n)+m is prime. I guess every term of this sequence (compare the conjecture about A005235) is prime. I checked this conjecture for n < 373.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Aug 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002906 | OEIS-A087201

a(n) is the smallest m such that m > A055211(n) and A002110(n)-m is prime.

a(1) and a(2) are not defined. a(n) is the second m (first m is A055211(n)) such that m > 1 and A002110(n)-m is prime. I guess every term of this sequence (compare the conjecture about A055211) is prime. I checked this conjecture for n < 418.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Aug 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002907 | OEIS-A087202

a(n) is the smallest m such that m > A037153(n) and n!+ m is prime.

Conjecture: For n > 1, a(n) is prime (compare the conjecture about A037153). Conjecture holds through 1200 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Sep 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002908 | OEIS-A087203

a(n) is the smallest m such that m > A037155(n) and n! - m is prime.

a(1) and a(2) are not defined. a(n) is the second m (first m is A037155(n)) such that m > 1 and n!- m is prime. For 3 < n < 643, a(n) is prime. I guess (compare the conjecture about A087202) except for the first term, every term of this sequence is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Sep 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002909 | OEIS-A087212

Number of ways in which n X n cells may be numbered using nonzero digits so that n n-digit squares are formed when read across or down the grids.

Sequence conjectured to be complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Oct 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002910 | OEIS-A087214

Expansion of e.g.f.: exp(x)/(1-x^2/2).

Conjecture: 2*a(n) -2*a(n-1) -n*(n-1)*a(n-2) +(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Aug 19 2014 Conjecture verified using differential equation (-x^2 + 2*x + 2)*y(x) + (x^2 - 2)*y'(x) = 0 satisfied by the e.g.f. - _Robert Israel_, Apr 30 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Oct 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002911 | OEIS-A087262

Integer quotient of largest and initial values in 3x+1 iteration, started at n.

It is conjectured that if peak/initial value is an integer then it equals 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Sep 11 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002912 | OEIS-A087304

Smallest nontrivial multiple of n whose nonzero digit product is the same as that of the nonzero digit product of n. By nontrivial one means a(n) is not equal to n or (10^k)*n. 0 if no such number exists.

Conjecture: no term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002913 | OEIS-A087318

Smallest number of the form n^k +1 with the prime signature of n, or 0 if no such number exists.

If n is an i-th power for i > 1, than a(n) = 0. The other zeros are conjectures. I have checked 12, 20 and 24 for k up to 400; 28, 40, 45, 48, 50, 52, 54, 55, 56, 63 and 68 for k up to 200 and 60 for k up to 100. - _David Wasserman_, May 09 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 03 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002914 | OEIS-A087319

Smallest number of the form n^k +1, with tau(a(n)) = tau(n), or 0 if no such number exists.

If n is a square then a(n) = 0; the other zeros (n = 12, 20, 27, 28, 45, 48, 50, 52 and 55) are conjectures. I have checked them for k up to 255. - _David Wasserman_, May 09 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 03 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002915 | OEIS-A087324

Numbers n such that n and its 9's complement both have the same prime signature.

Conjecture: (1) Sequence is infinite. (2) For every prime signature there corresponds a term in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002916 | OEIS-A087325

Numbers k such that k and its 10's complement both have the same prime signature.

Conjecture: (1) Sequence is infinite. (2) For every prime signature there corresponds a term in this sequence. Conjecture (2) is false: k and its 10's complement can't both have prime signature p^m where m is even.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002917 | OEIS-A087331

Smallest number with all identical digits having n distinct prime divisors.

The conjecture that 'for n > 2, a(n) == 0 (mod 3)' is not true since a(12) = 32 2's which is == 1 (mod 3). - _Robert G. Wilson v_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 05 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002918 | OEIS-A087339

Numbers k such that both the sum of the digits of k and 1 plus the product of its digits are primes.

Sequence is infinite. Proof: (10^p-1)/9 is a term if p is a prime. The sum of the digits = p and the product of digits + 1 = 2. Conjecture: There are infinitely many terms not of the form (10^p-1)/9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 06 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002919 | OEIS-A087347

Numbers k that divide Sum_{j=1..k} Fibonacci(j*k).

Conjecture: a(n) is asymptotic to c*n*log(n) with c around 1.3. [This conjecture is probably false, see plot. - _Vaclav Kotesovec_, Feb 16 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002920 | OEIS-A087357

Index of powers of 2 arising as the difference of a pair of successive terms of A087356.

Conjecture: Every number is a member.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 08 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002921 | OEIS-A087376

Leading diagonal of A087377.

Conjecture: a(n) < n^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002922 | OEIS-A087379

Beginning with 2, primes such that the difference between two successive terms is a distinct composite number.

The sequence of successive differences is given by the following distinct composite numbers 9,6,12,8,4,18,14,10,20,.... And trivially second term onwards only even composite numbers occur. Conjecture: Let a(m+1)-a(m) be composite (k). Then there exists a constant C such that m < C*k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002923 | OEIS-A087403

a(n) = smallest prime of the form 10*K(n) + 1, where K is a number obtained by concatenation of n with itself, or 0 if no such prime exists.

Conjecture: No term is zero. The conjecture is not true. There exist many numbers n such that a(n)=0. Conjecture: If n is not of the form 10^m then a(n) is nonzero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002924 | OEIS-A087444

Numbers that are congruent to {1, 4} mod 9.

3*a(n) is conjectured to be the total number of sides (straight double bonds (long side) and single bond (short side)) of a certain equilateral triangle expansion shown in one of the links. The pattern is supposed to become the planar Archimedean net 3.3.3.3.6 when n -> infinity. 3*a(n) is also conjectured to be the total number of sided (bonds) in another version of an equilateral triangle expansion that is supposed to become the planar Archimedean net 3.6.3.6. See the illustrations in the links. - _Kival Ngaokrajang_, Nov 30 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002925 | OEIS-A087445

Numbers that are congruent to 1 or 5 mod 12.

Conjecture: at least one number in this sequence must appear in all Collatz sequences. The sequence is composed of all numbers in congruence classes T(n,1) mod 2^(n+k) in A259663 (i.e., T"(1) in array T259663(n,k)) not congruent to 0 mod 3. Therefore the conjecture above is true (see A259663 for additional explanation). - _Bob Selcoe_, Jul 15 2017 The above conjecture is true: this is because even numbers and odd numbers divisible by 3 will lead to the set of odd numbers not divisible by 3. Odd numbers of the form 4k - 1 can also be ignored, as this consists of odd numbers that grow between themselves and the next odd term through Collatz iteration. No infinite sequence of growth between consecutive odd terms is possible, so all numbers of the form 4k - 1 will lead to an odd number that shrinks between itself and the next odd number. All numbers 4k - 1 will lead to a number in 4k - 3, the odd numbers that shrink between themselves and the following odd term. What we are left after that elimination is this sequence. - _Aidan Simmons_, Feb 25 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002926 | OEIS-A087446

Numbers that are congruent to {1, 6} mod 15.

3*a(n) is conjectured to be the number of edges (bonds) visited when walking around the boundary of a certain equilateral triangle construction at the n-th iteration. See the illustration in the link. Note that isthmus edges (bridges) are counted twice. The pattern is supposed to become the planar Archimedean net 3.12.12 when n -> infinity. - _Kival Ngaokrajang_, Nov 30 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002927 | OEIS-A087482

Number of binary polynomials of degree n irreducible over the integers.

Kogler proves that a(n) ~ 2^(n-1); this was the Odlyzko-Poonen conjecture. - _Charles R Greathouse IV_, Oct 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Sep 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002928 | OEIS-A087485

Odd numbers m such that 2*m - sigma(m) = 6.

Conjecture: a term where gcd(a(n),5)=1 for n > 1 does exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Oct 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002929 | OEIS-A087547

a(n) = n!*2^(n+1) * (Integral_{x = 0..1} 1/(1+x^2)^(n+1) dx) - Pi*(2*n)!/(2^(n+1)*n!).

Bala's conjecture is true. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Al Hakanson (hawkuu(AT)excite.com), Oct 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002930 | OEIS-A087552

a(1) = 1, then the smallest prime divisor of A065447(n) not included earlier.

Conjecture: Every prime is a member and this is a rearrangement of the noncomposite numbers. Proof of conjecture: primes 2=a(1) and 5=a(3) are terms, while any other prime divides infinitely many numbers of the form A002275([(n+1)/2]) = (10^[(n+1)/2]-1)/9, which in turn divide A065447(n). Thus every prime will sooner or later appear as a(n). - _Max Alekseyev_, Jul 03 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002931 | OEIS-A087553

Squares in which the neighboring digits differ at most by 1. Neighbors of 1 are 0 and 2, neighbors of 0 are 1 and 9.

The sequence is infinite with trivial terms like 10^(2k) and 9*10^(2k). Conjecture: There are infinitely many nontrivial terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002932 | OEIS-A087555

a(n) = largest prime that is the concatenation of two numbers whose sum is n.

a(3k) = 0. Conjecture: No other term is zero. Conjecture is false: first counterexample is 4477. See A375938. - _Robert Israel_, Sep 03 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002933 | OEIS-A087556

a(n) = smallest prime that is the concatenation of two numbers whose sum is n.

a(3k) = 0. Conjecture: No other term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002934 | OEIS-A087562

Primes such that the next n successive differences are identical.

The sequence of successive differences is (1),(2,2),(12,12,12),(60,60,60,60),... Conjecture: Sequence is infinite. The successive differences == 0 (mod 30) for n > 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002935 | OEIS-A087564

a(n) = smallest prime of the form n*(n+1)*(n+2)*...*(n+k) + 1, or 0 if no such prime exists.

a(n) = n+1 iff n+1 is prime. Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 15 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002936 | OEIS-A087582

Primes such that successive differences are distinct palindromes.

Conjecture: Sequence is infinite and every even palindrome is a member.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 17 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002937 | OEIS-A087583

Distinct primes such that the absolute values of successive differences are distinct palindromes. a(n+1) is chosen to be < a(n) if such a prime exists, minimizing a(n)-a(n+1); otherwise the minimal a(n+1) > a(n) is chosen.

The sequence of absolute differences is 1,2,6,4,22,8,66,44.... Conjecture: this sequence is infinite and contains every even palindrome.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 17 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002938 | OEIS-A087593

Define dd(n) = the number formed by concatenating the absolute difference of successive digits. Sequence contains primes p such that dd(p) is also prime. (Primes in which the number formed by successive digit difference is also a prime.).

Conjecture: Sequence is infinite. Subsidiary sequence: number of n-digit members.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002939 | OEIS-A087594

Define dd(n) = the number formed by concatenating the absolute difference of successive digits. Sequence contains primes p such that dd(p)=q is a prime, dd(q) is also a prime = r and so on until a single-digit prime (2,3,5,7) arises. (Primes in which the number formed by successive digit differences are primes at every step until a single-digit prime is obtained.).

Conjecture: Sequence is infinite. Subsidiary sequence: number of n-digit members.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002940 | OEIS-A087595

Smallest n-digit member of A087593. Define dd(k) = the number formed by concatenating the absolute difference of successive digits of k. Sequence contains smallest n-digit prime p such that dd(p) is also prime.

Conjecture: Sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002941 | OEIS-A087596

Largest n-digit member of A087593. Define dd(k) = the number formed by concatenating the absolute difference of successive digits of k. Sequence contains largest n-digit prime p such that dd(p) is also prime.

Conjecture: Sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002942 | OEIS-A087597

Triangular numbers m such that A040115(m) is also triangular.

Conjecture: Sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002943 | OEIS-A087599

Smallest nonzero n-digit term of A087597, or 0 if no such number exists.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002944 | OEIS-A087600

Largest n-digit term of A087597, or 0 if no such number exists.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002945 | OEIS-A087604

Smallest k such that n times concatenation of k with itself followed by a 3 is a prime, or 0 if no such number exists.

Trivially a(3n) = 0. Conjecture: no other term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002946 | OEIS-A087606

Smallest k such that n times concatenation of k with itself followed by a 9 is a prime, or 0 if no such number exists.

Conjecture: a(3n) = 0. No other term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002947 | OEIS-A087612

A divisibility sequence derived from Lehmer's polynomial x^10+x^9-x^7-x^6-x^5-x^4-x^3+x+1. Square root of the terms in A059928.

The sequence is conjectured to contain an infinite number of primes. The first 100 terms contain 33 unique primes. As stated by Everest and Ward, except for a finite number of composite n, a(n) can be prime only if n is prime. For this sequence, n=23*47 is the largest composite for which a(n) is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Sep 15 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002948 | OEIS-A087617

Gabcke sequence: a(0)=1; (n+1) a(n+1) = Sum_{k=0..n} 2^(4k+1) |E(2k+2)| a(n-k), where |E(2k+2)| are Euler numbers (E(2k)=(-1)^k A000364(k)).

Gabcke conjectured and _Juan Arias-de-Reyna_ proved that the terms are integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juan Arias-de-Reyna_, Sep 12 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002949 | OEIS-A087659

a(n) = hypergeom([-n, (n+4)/2, (n+5)/2], [3, 2], -4).

A few weeks ago I conjectured that 2 binomial(n,i) (n+2i+3)! / ((i+1)!(i+2)!(n+3)!) is always an integer (summed on i, this gives the current sequence). which I also conjecture integral. These conjectures are true. Consider the partition p(m,k,n)=(n+m,m,...,m) of n+m*(k+1), where m is repeated k times. It is easy to see that C(m,k,n) equals the dimension of the irreducible representation of S_(n+m*(k+1)) corresponding to p(m,k,n) calculated using hook length formula.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bill Gosper_, Sep 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002950 | OEIS-A087664

Consider recurrence b(0) = n/4, b(k) = b(k-1)*floor(b(k-1)); sequence gives number of steps to reach an integer, or -1 if no integer is ever reached.

It is conjectured that an integer is always reached if the initial value is >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002951 | OEIS-A087665

Consider recurrence b(0) = n/4, b(k) = b(k-1)*floor(b(k-1)); sequence gives first integer reached, or -1 if no integer is ever reached.

It is conjectured that an integer is always reached if the initial value is >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002952 | OEIS-A087666

Consider recurrence b(0) = n/3, b(k+1) = b(k)*floor(b(k)); a(n) is the least k such that b(k) is an integer, or -1 if no integer is ever reached.

It is conjectured that an integer is always reached if the initial value n/3 is >= 2. a(n)=0 iff n == 0 (mod 3), a(n)==1 iff n == 1 or 2 (mod 3^2), a(n)=2 iff n == 14,22,25,26 (mod 3^3). In general a(n)=m iff n == x (mod 3^m) where x pertains to a set of 2^m distinct elements included in {1,2,...,(3^m)-1}. Conjecture: a(6) + a(7) + a(8) + ... + a(n) = 2n + O(sqrt(n)). - _Benoit Cloitre_, Sep 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002953 | OEIS-A087669

The smallest index m such that b(m) is an integer, where b(0)=(2n+1)/n and b(k+1)=b(k)*floor(b(k)) for k>=0.

It is conjectured that an integer is always reached if the initial value is >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002954 | OEIS-A087670

The first integer in the sequence defined by b(0)=(2n+1)/n and b(k+1)=b(k)*floor(b(k)) for k>=0.

It is conjectured that an integer is always reached if the initial value is >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002955 | OEIS-A087676

Consider recurrence b(0) = n/3, b(n) = b(0)*floor(b(n-1)); sequence gives number of steps to reach an integer, or -1 if no integer is ever reached.

It is conjectured that an integer is always reached if the initial value is >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, following a suggestion of Bela Bajnok (bbajnok(AT)gettysburg.edu), Sep 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002956 | OEIS-A087677

Consider recurrence b(0) = n/3, b(n) = b(0)*floor(b(n-1)); sequence gives first integer reached, or -1 if no integer is ever reached.

It is conjectured that an integer is always reached if the initial value is >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, following a suggestion of Bela Bajnok (bbajnok(AT)gettysburg.edu), Sep 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002957 | OEIS-A087704

Number of steps for iteration of map x -> (5/3)*floor(x) to reach an integer > n when started at n, or -1 if no such integer is ever reached.

It is conjectured that an integer is always reached.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002958 | OEIS-A087705

First integer > n reached under iteration of map x -> (5/3)*floor(x) when started at n, or -1 if no such integer is ever reached.

It is conjectured that an integer is always reached.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002959 | OEIS-A087707

Number of steps for iteration of map x -> (5/3)*ceiling(x) to reach an integer > n when started at n, or -1 if no such integer is ever reached.

It is conjectured that an integer is always reached.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002960 | OEIS-A087708

First integer > n reached under iteration of map x -> (5/3)*ceiling(x) when started at n, or -1 if no such integer is ever reached.

It is conjectured that an integer is always reached.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002961 | OEIS-A087711

a(n) = smallest number k such that both k-n and k+n are primes.

Let b(n), c(n) and d(n) be respectively, smallest number m such that phi(m-n) + sigma(m+n) = 2n, smallest number m such that phi(m+n) + sigma(m-n) = 2n and smallest number m such that phi(m-n) + sigma(m+n) = phi(m+n) + sigma(m-n), we conjecture that for each positive integer n, a(n)=b(n)=c(n)=d(n). Namely we conjecture that for each positive integer n, a(n) < A244446(n), a(n) < A244447(n) and a(n) < A244448(n). - _Jahangeer Kholdi_ and _Farideh Firoozbakht_, Sep 05 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Sep 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002962 | OEIS-A087713

Greatest prime factor of the product of the neighbors of the n-th prime.

Conjecture: The record values are A120628 \ {2}. - _Jason Yuen_, Jan 19 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Sep 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002963 | OEIS-A087714

Primes p = prime(i) such that p(i)# - p(i+1) and p(i)# + p(i+1) are both primes, where p# = A002110.

Conjecture: there are only 4 primes in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Sep 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002964 | OEIS-A087716

Base-2 pseudoprimes (see A001567) of the form j*p(i)# - p(k) or j*p(i)# + p(k), p(i) and p(k) primes with p(i) < p(k) < p(i+1)^2 and 0 < j < p(i+1).

Conjecture: sequence has only 5 terms. This has been checked for all i <= 150.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Sep 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002965 | OEIS-A087717

Start with x=n, then iterate the map x -> A322982(x) with A322982(x)=2*x-1 if x is noncomposite, otherwise A322982(x) = A032742(x), the largest proper divisor of x. If this iteration leads to a fixed point then a(n) is the value of that fixed point. If the iteration leads to a cycle, a(n) is the smallest value in the cycle. If the iteration never becomes periodic then a(n)=0.

Conjecture. For n > 1, the iteration given in the definition above always leads to the 3-cycle {3,5,9,3} or the 6-cycle {19,37,73,145,29,57,19}, thus a(n) takes on only the values 3 or 19 for n=2,3,4,.... This has been verified to n=1000000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Sep 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002966 | OEIS-A087725

Maximal number of moves required for the n X n generalization of the sliding block 15-puzzle (or fifteen-puzzle).

Conjecture: a(n) ~ (4/3)*n^3. - _Ben Whitmore_, May 29 2021 The above conjecture is false. Zhixian Zhong proved that a(n) = n^3 + O(n^(11/4)), and ciphrae shows that a(n) = n^3 + O(n^2 * log(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_, Sep 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002967 | OEIS-A087734

a(n) = f(f(n)), where f() = A035327().

Some conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002968 | OEIS-A087790

Partial sums of A085068.

Conjecture : a(n)=3n+O(sqrt(n))

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002969 | OEIS-A087791

Indices n such that S^n(1) is an integer where S(x) = 3/2*ceiling(x).

Conjecture : a(n)=2n+O(sqrt(n))

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002970 | OEIS-A087792

Indices n such that S^n(1) is an integer where S(x) = 4/3*ceiling(x).

Conjecture : a(n)=3n+O(sqrt(n))

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002971 | OEIS-A087809

Number of triangulations (by Euclidean triangles) having 3+3n vertices of a triangle with each side subdivided by n additional points.

Conjecture: 2*n*(2*n-1)*(5*n^2 - 29*n + 30)*a(n) + (-295*n^4 + 1926*n^3 - 3425*n^2 + 2106*n - 360)*a(n-1) + 24*(3*n-4)*(3*n-5)*(5*n^2 - 19*n + 6)*a(n-2) = 0. - _R. J. Mathar_, Apr 23 2015. Proved by Andrei Asinowski, C. Krattenthaler, T. Mansour, Counting triangulations of balanced subdivisions of convex polygons, 2016.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roland Bacher_, Oct 16 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002972 | OEIS-A087817

a(n) = a(a(a(n-1)))+a(n-a(n-1)) with a(1)=a(2)=1.

Conjecture: a(n) is asymptotic to n^c with c=0.677.... - _Benoit Cloitre_, Jun 05 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Oct 05 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002973 | OEIS-A087847

a(n) = a(|n - a(n-1)|) + a(a(a(|n - a(n-4)|))).

The conjecture is that even higher order recursions of the term one and term two type for the original and skip term versions of A005185 Hofstadter Q will exist as well. I have invented this way of naming the larger generalization of Hofstadter Q type sequences as being descriptive of their formation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Oct 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002974 | OEIS-A087942

Number of partitions of n into as many primes as n has prime factors.

Conjecture, for m>1: a(m)=0 iff n is an odd semiprime such that m-2 is not prime, i.e. m=A089268(k) for some k. - _Reinhard Zumkeller_, Oct 28 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Oct 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002975 | OEIS-A087952

Smallest prime == 1 (mod n) and > n^2.

Since A014085(n) ~ n/log(n) one may conjecture that a(n) < 2*n^2 for all n > 1. Numerically we find a(n) = n^2*(1 + O(1/sqrt(n))). - _M. F. Hasler_, Feb 27 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ray Chandler_, Sep 16 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002976 | OEIS-A087979

a(n) = Min {x : sigma(x) = phi(n*x), x is not a prime}, least nonprime solutions to sigma(x) = phi(n*x).

For n >= 5, a(n) = A256527(n). - Conjectured by _Manfred Scheucher_, May 28 2015; proved by _Max Alekseyev_, Sep 29 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Sep 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002977 | OEIS-A087981

Expansion of e.g.f.: exp(-2*x) / (1-x)^2.

It is conjectured by Kräuter and Seifter that for n >= 5 a(n-1) is the maximal possible value for the permanent of a nonsingular n X n (+1, -1)-matrix. I do not know for which values of n this has been confirmed - compare A087982. - _N. J. A. Sloane_ The Kräuter conjecture on permanents is true (see Budrevich and Guterman). - _Sergei Shteiner_, Jan 17 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gordon Royle_, Oct 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002978 | OEIS-A087982

Maximal permanent of a nonsingular n X n (+1,-1)-matrix.

It is conjectured by Kraeuter and Seifter that for n >= 5 the maximal permanent of a nonsingular n X n (+1,-1)-matrix is attained by a matrix with exactly n-1 -1's on the diagonal (compare A087981).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002979 | OEIS-A088021

a(n) = (n^2)!/(n!)^2.

Based on an observation of _Hugo Pfoertner_, _W. Edwin Clark_ conjectured and Xiang-dong Hou proved that (n^2)!/(n!)^2 gives the number of distinct determinants of the generic n X n matrix whose entries are n^2 different indeterminates under all (n^2)! permutations of the entries.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Sep 18 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002980 | OEIS-A088023

Set a(1) = 1. Then take the list of defined initial terms, reverse their order, add 1, 2, 3, ... to the reversed list in succession and append this new list to the right of the previously defined terms. Repeat this process indefinitely.

Conjecture: a(n+1) >= a(n). Comments from _Don Reble_, Nov 13 2005: The conjecture is plainly true. In fact, a(n+1)-a(n) = 0 or 1. Also a(A091072(n)) = n; a(A091072(n)+1) = n+1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Sep 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002981 | OEIS-A088039

Smallest k such that k^3 == 1 (mod some n-th power), k > 1.

Conjecture: a(n) = 2^n+1 for n>2. a(n) = 3*a(n-1)-2*a(n-2) for n>4. G.f.: -x*(2*x^3-x^2+2*x-2) / ((x-1)*(2*x-1)). [_Colin Barker_, Feb 05 2013]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002982 | OEIS-A088041

Smallest k such that k^4 - 1 is divisible by an n-th power, k > 1.

Conjecture: a(n) = 3*a(n-1)-2*a(n-2) for n>5. G.f.: x*(4*x^4-2*x^2-3*x+2) / ((x-1)*(2*x-1)). [_Colin Barker_, Feb 05 2013] Conjectures confirmed (see link). - _Robert Israel_, Aug 19 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002983 | OEIS-A088060

For each Fibonacci number F(m) = A000045(m), m >= 1, look for the smallest Fibonacci number F(k) > F(m) such that F(m) + F(k) is a square. If it exists, append F(k) to the sequence.

Conjecture: sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 21 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002984 | OEIS-A088070

Numbers sandwiched between two numbers having the same number of prime divisors.

Subsidiary sequence: (1) Start of the first run of a string of n successive integers in this sequence. Conjecture: For every k there is a string of k successive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002985 | OEIS-A088072

Numbers sandwiched between two numbers having only one prime divisor (at least) one of which is composite.

Conjecture (1): Sequence is infinite. Conjecture (2): There are only finitely many terms which are sandwiched between composite numbers (26 is the only such term below 1000000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002986 | OEIS-A088090

For n > 1, a(n) is the smallest number such that n-th concatenation is prime and the smallest palindrome beginning with (but not equal to) this concatenation is also prime.

Conjecture: sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002987 | OEIS-A088093

Beginning with 2, smallest palindrome >= the previous term such that every concatenation is a prime.

Conjectures: (1) Sequence is infinite. (2) Two successive terms are not equal for n > 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002988 | OEIS-A088110

Numbers n such that 2^n + n and 2^n - n are noncomposite.

Conjecture: sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002989 | OEIS-A088112

Perfect powers whose digit reversal is prime.

Conjectures: (1) Sequence is infinite. (2) For every n there are infinitely many members of the form k^n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002990 | OEIS-A088113

a(n) = digit reversal of (11^n) divided by 11.

Conjecture: There are finitely many palindromic terms. Question: How many are palindromes? Observation: a(n) is further divisible by 11 for n = 2,3,4,5,6,7,8,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002991 | OEIS-A088120

Smallest prime obtained as a product of n terms of an arithmetic progression + the common difference.

Conjectures: (1) Sequence is infinite. (2) The common difference arising = 1 except for n = 4 and 5 in which case it is 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002992 | OEIS-A088126

Let f(n, x) = x+3x^2+6x^3+...+(n(n+1)/2)x^n; then a(n) = least x such that f(n, x) is a triangular number, or 0 if no such x exists.

Zero values are conjectures. I have searched for a(4) up to x = 10^7, a(5) up to x = 10^6 and the rest up to x = 10^4. (Wasserman)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002993 | OEIS-A088144

Sum of primitive roots of n-th prime.

It is a result that goes back to Mirsky that the set of primes p for which p-1 is squarefree has density A, where A denotes the Artin constant (A = Product_{q prime} (1-1/(q*(q-1)))). Numerically A = 0.3739558136.. = A005596. More precisely, Sum_{p <= x} mu(p-1)^2 = Ax/log x + o(x/log x) as x tends to infinity. Conjecture: sum_{p <= x, mu(p-1)=1} 1 = (A/2)x/log x + o(x/log x) and sum_{p <= x, mu(p-1)=-1} 1 = (A/2)x/log x + o(x/log x). - Pieter Moree (moree(AT)mpim-bonn.mpg.de), Nov 03 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ed Pegg Jr_, Nov 03 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002994 | OEIS-A088164

Wolstenholme primes: primes p such that binomial(2p-1,p-1) == 1 (mod p^4).

Primes p such that p^3 divides A001008(p-1) (Zhao, 2007, p. 18). Also: Primes p such that (p, p-3) is an irregular pair (cf. Buhler, Crandall, Ernvall, Metsänkylä, 1993, p. 152). Keith Conrad observes that for the two known (as of 2015) terms ord_p(H_p-1) = 3 is satisfied, where ord_p(H_p-1) gives the p-adic valuation of H_p-1 (cf. Conrad, p. 5). Romeo Mestrovic conjectures that p is a Wolstenholme prime if and only if S_(p-2)(p) == 0 (mod p^3), where S_k(i) denotes the sum of the k-th powers of the positive integers up to and including (i-1) (cf. Mestrovic, 2012, conjecture 2.10). - _Felix Fröhlich_, May 20 2015 The converse of Wolstenholme's theorem implies that if an integer n satisfies the congruence binomial(2*n-1, n-1) == 1 (mod n^4), then n is a term of this sequence, i.e., then n is necessarily prime, or, equivalently, A298946(i) > 1 for all i > 0. Whether this is true for all such n is an open problem. - _Felix Fröhlich_, Feb 21 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian Schroeder_, Sep 21 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002995 | OEIS-A088178

Sequence of distinct products b(n)*b(n+1), n=1,2,3,..., of the terms b(n) of A088177.

Conjecture: the term a(n) is a prime if and only if every number < a(n) belongs to the set {a(1), a(2), ..., a(n-1)}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Sep 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002996 | OEIS-A088183

Number of ways to write n as a sum of two coprime semiprimes.

Is a(n)>0 for n>210? see conjecture in A072931.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Sep 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002997 | OEIS-A088185

Numbers which are not the sum of two coprime semiprimes.

is this sequence finite/full? See conjecture in A072966.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Sep 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002998 | OEIS-A088195

Distance (A088192) of primes from the largest quadratic residues modulo the primes (A088190), where the latter is non-monotonic.

It is conjectured that if we denote the members of A088194 by D(n) and the member of this sequence by M(n) then if D(n)=-1 then M(n)=7, while if M(n)=3 then D(n)=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ferenc Adorjan (fadorjan(AT)freemail.hu), Sep 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-002999 | OEIS-A088199

Primes where the difference sequence (A088197) of LQnR(p_n) (A088196) is <= 0.

The members of the sequence are always == 1 modulo 8 (conjectured!).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ferenc Adorjan (fadorjan(AT)freemail.hu), Sep 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003000 | OEIS-A088200

Members of the difference sequence (A088197) of LQnR(p_n) (A088196) where it is <= 0.

The members of the sequence are always even (conjectured!). The conjecture above is false. The first odd term is a(137) = -3. - _Qiyuan Gu_, Aug 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ferenc Adorjan (fadorjan(AT)freemail.hu), Sep 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003001 | OEIS-A088201

Distance p_n-LQnR(p_n) (A088198) where the difference sequence (A088197) of LQnR(p_n) (A088196) is <= 0.

The terms are conjectured to be odd primes > 3. It is also conjectured that the i-th member of A088200 is -2 if and only if a(i) is 5. The terms are conjectured to be odd primes > 3 (the primality is provable).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ferenc Adorjan (fadorjan(AT)freemail.hu), Sep 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003002 | OEIS-A088207

a(n) = Sum_{k=0..n} floor(k*phi^2) where phi=(1+sqrt(5))/2.

Conjecture: a(n)/A054347(n) tends to phi. Example: a(28)/A054347(28) = 1049/643 = 1.6314... Proof of Adamson's conjecture. We know that lim_{n->oo} A054347(n)/(n*(n+1)) = phi/2 (see A054347).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Sep 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003003 | OEIS-A088212

Smallest k>1 such that n+k^2 is prime.

Conjecture: all integers >1 eventually appear. Among first 300000 terms, first absent integers are 113, 119, 122, 124, 127, 130, 134, 136, 137, 139, 140, 142, 143, 145, 146, 148, 149, 151, 152, 154, 155, 157, 158, 160, 161, 163, 164, 166, 167, 169, 170, 172, 173, 175, 176, 178, 179, 181, 182, 184, 185, 186, 187, 188, 190, 191, 193, 194, 196, 197, 199, 200. - _Zak Seidov_, Jun 06 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Sep 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003004 | OEIS-A088226

a(1)=0, a(2)=0, a(3)=1; for n>3, a(n) = abs(a(n-1) - a(n-2) - a(n-3)).

Conjecture: a(n) = m/2 where m is the smallest even distance from n to a square. - _Ralf Stephan_, Sep 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003005 | OEIS-A088249

Smallest palindromic prime beginning with the n-th prime, or 0 if no such prime exists.

Conjecture: Except for n=1 and 3, a(n) = 0 iff prime(n) has Most Significant digit one among (2,4,5,6,8).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003006 | OEIS-A088251

A088250(n) + 1.

Conjecture: (1) Sequence is infinite. (2) For every n there are infinitely many arithmetic progressions with n successive primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003007 | OEIS-A088252

n-th row of the following triangle contains smallest set of n primes which form n successive terms of an arithmetic progression from the 2nd to (n+1)th term with the first term 1. Sequence contains the leading diagonal.

Conjectures: (1) Sequence is infinite. (2) For every n there are infinitely many arithmetic progressions with n successive primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003008 | OEIS-A088254

Primes p with property that if p = prime(k), if the product of all composite numbers between prime(k) and prime(k+1) is one less than a prime.

Conjecture: the sequence is infinite. The smaller member of the twin prime pairs is a term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003009 | OEIS-A088256

Primorial numbers k such that both k-1 and k+1 are prime.

Conjecture: sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003010 | OEIS-A088258

Factorials F which have at least one prime neighbor, i.e., F+1 or F-1 or both are primes.

Conjecture: sequence is infinite. But there are finitely many members which are sandwiched between twin primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003011 | OEIS-A088259

Perfect powers which have at least one prime neighbor.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003012 | OEIS-A088260

Primorials which yield palindromes if the trailing zeros are deleted.

Conjecture: sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003013 | OEIS-A088268

Palindromes in A007530.

Conjecture: Sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003014 | OEIS-A088276

Smallest k such that 10^k + r is a prime ending (least significant digits) in r where r is the n-th odd number not divisible by 5. 0 if no such number exists.

Conjecture: (1) The only zero entries are for values of r == 2 (mod 3). (2) There are infinitely many values of k such that 10^k + r is a prime if r is not == 2 (mod 3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003015 | OEIS-A088280

Noncomposites in A088279.

Conjecture: sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003016 | OEIS-A088288

Palindromic primes made of only two digits say a and b such that replacing a with b and vice versa one gets another prime.

Conjecture: Sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Sep 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003017 | OEIS-A088333

A version of Josephus problem: a(n) is the surviving integer under the following elimination process. Arrange 1,2,3,...,n in a circle, increasing clockwise. Starting with i=1, delete the integer 3 places clockwise from i. Repeat, counting 3 places from the next undeleted integer, until only one integer remains.

It is tempting (in view of A054995) to conjecture that a(1)=1 and, for n>1, a(n) = (a(n-1)+4) mod n. The conjecture is false; counterexample: a(21)=21; a(20)=17; (a(20)+4)mod 21=0; corrected formula: a(n)=(a(n-1)+3) mod n +1; The conjecture is true. After removing the 4th number, we are reduced to the n-1 case, but starting with 5 instead of 1. - _David Wasserman_, Aug 08 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003018 | OEIS-A088367

Decimal expansion of Krivine's bound for Grothendieck's constant, Pi/(2*log(1+sqrt(2))).

Krivine (1977) proved that Grothendieck's constant <= Pi/(2*log(1+sqrt(2))), and conjectured that this bound is the exact value of the constant. His conjecture was refuted by Braverman et al. (2013). - _Amiram Eldar_, Jun 24 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Sep 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003019 | OEIS-A088431

Half of the (n+1)-st component of the continued fraction expansion of Sum_{k>=0} 1/2^(2^k).

Hendriks' conjecture is proved in Bunder, Bates, and Arnold (2024). Also see Shallit (2024). - _Jeffrey Shallit_, Mar 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 08 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003020 | OEIS-A088500

Expansion of e.g.f. 1/(1+2*log(1-x)).

Conjecture: a(n) = Integral_{x=0..oo} Pochhammer(2*x, n)/exp(x) dx. - _Velin Yanev_, Jan 11 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Nov 12 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003021 | OEIS-A088508

Smallest m such that n divides floor(Pi*10^m), Pi=3.14...

Conjecture: a(n) always exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Nov 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003022 | OEIS-A088519

2-Golomb's sequence G(n,2): earliest positive increasing sequence starting with (1,2) and satisfying "length of n-th run = n-th partial sum".

conjecture: a(n) = (2+sqrt(2))^(1/(2+sqrt(2)))*n^(sqrt(2)-1) + O(1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003023 | OEIS-A088568

3*n - 2*(partial sums of Kolakoski sequence A000002).

It is conjectured that a(n) = o(n). It is conjectured that the density of 1's and that of 2's in the Kolakoski sequence A000002 are equal to 1/2. The deficit of 2's in the Kolakoski sequence at rank n being defined as n/2 - number of 2's in

the Kolakoski word of length n, a(n) is equal to twice the deficit of 2's (or twice the excess of 1's). Equivalently, the number of 2's up to rank n in the Kolakoski sequence is (n - a(n))/2. - _Jean-Christophe Hervé_, Oct 05 2014 The conjecture about the densities of 1's and 2's is equivalent to a(n) = o(n). The graph shows that a(n) seems to oscillate around 0 with a pseudo-periodic and fractal pattern. - _Jean-Christophe Hervé_, Oct 05 2014 It is conjectured that a(n) = O(log(n)) (see PlanetMath link). Note that for a random sequence of 1's and -1's, we would have O(sqrt(n)). - _Daniel Forgues_, Jul 10 2015 The linked PlanetMath text mentions 0.5*n + O(log(n)) only in respect of an empirical observation, apparently to support the density conjecture (the conjecture described above in the first comment dated Oct 05 2014). - _Peter Munn_, Aug 03 2022 The conjecture that a(n) = O(log(n)) seems incorrect as |a(n)| seems to grow as fast as sqrt(n), see A289323 and note that a(2^n) = -A289323(n), so for example a(2^64) = -A289323(64) = -836086974 which is much larger in absolute value than log(2^64), but about 0.19*2^32. - _Richard P. Brent_, Jul 07 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 17 2003; definition changed Oct 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003024 | OEIS-A088601

Number of steps to reach 0 when iterating A261424(x) = x - (the largest palindrome less than x), starting at n.

While it has been conjectured [proved by Cilleruelo & Luca, 2016 -Ed.] that every number can be represented as a sum of at most 3 palindromes, the terms of this sequence, which correspond to a greedy representation, can be larger than 3 (see A109326). For example, 1022 can be represented as 33 + 989, but a(1022) = 4, because the greedy decomposition gives 1022 = 1001 + 11 + 9 + 1. - _Giovanni Resta_, Aug 20 2015 In A109326 an explicit formula for a smaller (conjectured sharp) upper bound was already given earlier. - _M. F. Hasler_, Sep 09 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Oct 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003025 | OEIS-A088606

Smallest number k such that concatenation of k and prime(n) is a prime, or 0 if no other number exists. a(1) = a(3) = 0.

Subsidiary sequences: (set(1)) Index of the start of the first occurrence of a string of n consecutive 1's or 2's or 3's etc. (set (2)): a(n) = smallest prime such that concatenation of 1 with n successive primes starting from a(n) gives primes in each case. (n primes are obtained.) Similarly for 2, 3, etc. Conjecture: The subsidiary sequences are infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Oct 15 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003026 | OEIS-A088614

Alternate prime and composite numbers not included earlier such that every partial concatenation is a prime: a(2n) is composite and a(2n-1) is prime.

Conjecture: 1 and 5 are the only two odd nonmembers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Oct 16 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003027 | OEIS-A088618

Rearrangement of odd numbers such that every concatenation of at least two terms is composite.

Conjecture: Sequence is infinite. And every odd number is a member.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Oct 17 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003028 | OEIS-A088628

Smallest prime using all the digits of first n numbers. If necessary, extra digits can be used.

Conjecture: There exists a number k such that for all m > k, m not == 0 or 2 (mod 3), a(m) does not use any extra digit.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Oct 21 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003029 | OEIS-A088643

Triangle read by rows: row n >= 1 is obtained as follows. Start with n, next term is always largest number m with 1 <= m < n which has not yet appeared in that row and such that m + previous term in the row is a prime. Stop when no further m can be found.

It is conjectured that row n is always a permutation of {1..n}. This has been verified for n <= 400000. I conjecture that almost all rows have exactly 7 (but not more) trailing terms in common with the initial terms of A055265 = (1, 2, 3, 4, 7, 6, 5, 8, ...): After row 10 whose reversal matches the first 10 terms of A055265, and rows n = 14, 15 and 16 having the last 2 (but not 3) terms equal to A055265(1..2), all rows up to n = 500 have either (about 25%) exactly 1 or (about 73%) exactly 7 trailing terms equal to the first terms of A055265. Between n = 501 and n = 10000 and beyond, all rows end in (..., 9, 14, 5, 6, 7, 4, 3, 2, 1), so they all have exactly m = 7 but not m = 8 trailing terms equal to A055265(1..m). - _M. F. Hasler_, Aug 03 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003030 | OEIS-A088647

a(1) = 1, then the smallest number such that the concatenation a(r), a(s) is prime for all s > r.

Conjecture: sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Oct 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003031 | OEIS-A088648

a(1) = 1, then the smallest odd number not occurring earlier such that the concatenation a(r), a(s) is composite for all s > r.

The sequence is infinite with trivial terms of the form 10k+5. Subsidiary sequence: sequence of nontrivial terms. Conjecture: the subsidiary sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Oct 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003032 | OEIS-A088677

Numbers that can be represented as j^6 + k^6, with 0 < j < k, in exactly one way.

Conjecture: no number can be expressed as such a sum in more than one way.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Nov 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003033 | OEIS-A088719

Numbers that can be represented as a^7 + b^7, with 0 < a < b, in exactly one way.

Conjecture: no number can be expressed as such a sum in more than one way.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Nov 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003034 | OEIS-A088734

First primes in p^p - 2m where p is prime and m = 1,2..

Conjecture: 2,3,5,7,11,13,17,19 are the only 1 and 2 digit primes in the sequence. The smallest 8 prime power less 2n primes PPL2NP?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Nov 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003035 | OEIS-A088753

Numbers k such that the Reverse and Add! trajectory of k (presumably) does not reach a palindrome (with the exception of k itself) and does not join the trajectory of any term m < k.

It is not easy to determine whether the orbit of a given term will never merge into the orbit of an earlier term. It seems that the property of "disjoint orbit" is as conjectural as the property of not reaching a palindrome. One could specify a "search limit" in order to get a well-defined sequence. The given list of terms has been checked and extended by considering the orbits up to members of size <= 10^199 at least. Given that the number increases by a factor 10 roughly every 2.416 iterations, this corresponds to about 500 iterations. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003036 | OEIS-A088783

Numbers n such that 10*n^k + 1 is composite for all k > 0.

The prime 10*173^264234+1 is found to solve the generalized Sierpinski problem in base 173 (with conjectured smallest Sierpinski number k=28, see the Barnes link and A123159 for these problems). - _Eric Chen_, Jun 07 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ray Chandler_, Oct 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003037 | OEIS-A088827

Even numbers with odd abundance: even squares or two times squares.

Conjecture: a(n) = ((2*r) + 1)^2 * 2^(c+1) where r and c are the corresponding row and column of n in the table format of A191432, where the first row and column are 0. - _John Tyler Rascoe_, Jul 12 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Oct 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003038 | OEIS-A088831

Numbers k whose abundance is 2: sigma(k) - 2k = 2.

Conjecture: There are no terms of other forms. So the next terms of this sequence are likely 2^93*(2^94-3), 2^115*(2^116-3), 2^121*(2^122-3), 2^149*(2^150-3), etc. (End) The conjecture in the previous comment is incorrect. The first counterexample is 650, which has factorization 2*5^2*13. - _T. D. Noe_, May 10 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Oct 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003039 | OEIS-A088862

Smallest number with exactly n prime factors (possibly with repetitions) such that in binary representation the divisors >1 form a prefix code.

Conjecture: the sequence is defined for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Nov 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003040 | OEIS-A088883

Primes which when concatenated with their reverse and incremented by 2 yield a new prime.

Conjecture: All these primes are of the form 6*k + 1. - _Davide Rotondo_, Apr 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chuck Seggelin_, Oct 21 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003041 | OEIS-A088884

Primes which when concatenated with their reverse and decremented by 2 yield a new prime.

Conjecture: except for 3, all these primes are of the form 6*k - 1. - _Davide Rotondo_, Apr 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chuck Seggelin_, Oct 21 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003042 | OEIS-A088910

Conjectured minimal required number k of terms in a representation n = Sum_{i=1..k} e_i*(p_i)^2 by distinct primes p_i, where e_i is 1 or -1.

Conjectured minimal required number k of terms in a representation n = Sum_{i=1..k} e_i*(p_i)^2 by distinct primes p_i, where e_i is 1 or -1. It is conjectured that all sequence terms are <= 5. The terms with a(n)=5 were provided by _W. Edwin Clark_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Oct 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003043 | OEIS-A088913

a(n) = (Lucas(2n) + Lucas(2n+1)*phi)/kappa(phi/Fibonacci(4n)) where kappa(x) is the sum of successive remainders in computing the Euclidean algorithm for (1,x).

Conjecture: a(n) = Lucas(6*n) - Lucas(2*n). - _Peter Bala_, Jul 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Baruchel_, Nov 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003044 | OEIS-A088966

Numbers k such that A007947(k) = A007947(m+1) and A007947(m) = A007947(k+1), where k > m.

G.f.: Conjecture: Q(0)/x - 1/x where Q(k)= 1 + 2^k*x/(1 - 2*x/(2*x + 2^k*x/Q(k+1))); (continued fraction). - _Sergei N. Gladkovskii_, Apr 10 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, Oct 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003045 | OEIS-A088982

Primes that are between consecutive prime-indexed primes.

Conjecture: For x > 1 there is at least 1 prime between prime(prime(x)) and prime(prime(x+1)). This conjecture is equivalent to saying that there is at least one prime index between prime(x) and prime(x+1), which is trivially true because both are odd for x > 1; one has prime(prime(x)) < prime(prime(x)+1) < prime(prime(x+1)). Obviously the definition is equivalent to "primes > 2 with nonprime index", i.e., sequence A007821 without the initial 2. - _M. F. Hasler_, Jul 31 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Oct 31 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003046 | OEIS-A088983

Numbers k such that each of the 6 consecutive numbers k through k+5 has exactly two distinct prime factors.

This sequence has no run of four consecutive integers, since Eggleton and MacDougall prove that there are no more than 9 consecutive integers with A001221(k) = 2. They conjecture that A007774 contains no runs of 9 consecutive integers, and has only two runs of size 8 (at 141 and 212) and two maximal runs of size 7 (at 323 and 2302); they add that the maximal run of size 6 at 91 might be the only such run, so A088983 might be complete. - Roger Eggleton via _Jason Kimberley_, Jul 12 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Sep 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003047 | OEIS-A088987

Triples of primes between two consecutive prime-indexed primes.

Conjecture: There cannot be an even number total of primes between two consecutive prime-indexed primes. The conjecture is true except for the first two prime-indexed primes (3 and 5), which have zero primes between them.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Oct 31 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003048 | OEIS-A088995

Least k > 0 such that the first n digits of 2^k and 5^k are identical.

The terms grow like 10^n, oscillating by a few orders of magnitude around this value. I conjecture that log(a(n)/10^n) = O(log log n). - _M. F. Hasler_, Mar 22 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Dec 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003049 | OEIS-A089001

Numbers k such that 2*k^2 + 1 is prime.

Conjecture: these are the positive numbers that cannot be represented as d + t/d where t is a triangular number (A000217) and d is a divisor of t. Note that if k = d + n*(n+1)/(2*d), then 2*k^2 + 1 = (2*d^2 + (n+1)^2)*(2*d^2 + n^2)/(2*d^2) cannot be prime.- _Robert Israel_, Mar 10 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003050 | OEIS-A089039

Number of circular permutations of 2n letters that are free of jealousy.

a(n) = (n-1)!*(A001040(n-1) + A001053(n)) = 2*A276356(n), n > 1. - Conjectured by _Mikhail Kurkov_, Feb 10 2019 and proved by _Max Alekseyev_, Apr 23 2024 (see MO link) a(n+4) = -(n+3)*(n+2)*(n*(n+1)*a(n) + 2*(n+1)^2*a(n+1) + n*(n+3)*a(n+2) - 2*a(n+3)) for all integer n>1. - conjectured by _Michael Somos_, Apr 21 2024. [The conjecture is equivalent to Kurkov's formula and thus is also proved. - _Max Alekseyev_, Apr 23 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | Akemi Nakamura, Michihiro Takahashi, Shogaku Meitantei (naka(AT)sansu.org), Dec 03 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003051 | OEIS-A089046

Least edge-length of a square dissectible into at least n squares in the Mrs. Perkins's quilt problem.

It is not clear which terms have been proved to be correct and which are just conjectures. - _Geoffrey H. Morley_, Aug 29 2012; _N. J. A. Sloane_, Jul 06 2017 n <= 15 (and possibly 16) proved minimal by _J. H. Conway_ (Conway, J. H. "Re: [math-fun] Mrs. Perkins Quilt - Orders 89, 90 improved over UPIG." math-fun mailing list. October 10, 2003.). The conjectures are best currently known values of a(n) for n > 16. - _Stuart E Anderson_, Apr 21 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_, Dec 03 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003052 | OEIS-A089047

Edge length of largest square dissectible into up to n squares in Mrs. Perkins's quilt problem.

It is not clear which terms have been proved to be correct and which are just conjectures. - _Geoffrey H. Morley_, Sep 07 2012; _N. J. A. Sloane_, Jul 06 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_, Dec 03 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003053 | OEIS-A089073

Number of symmetric non-crossing connected graphs on n equidistant nodes on a circle.

Conjecture D-finite with recurrence n*(n+2)*(23*n^2-162*n+199) *a(n) +12*(27*n^2-47*n-10) *a(n-1) +24*(-27*n^2+47*n+10) *a(n-2) +48 *(27*n^2-47*n-10) *a(n-3) -12*(3*n-13)*(3*n-5)*(23*n^2-116*n+60) *a(n-4)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003054 | OEIS-A089270

Positive numbers represented by the integer binary quadratic form x^2 + x*y - y^2 with x and y relatively prime.

(Conjecture) The terms are the products of primes congruent to {0,1,4} mod 5 (using at most a single 5, but repeating other primes is allowed), which is A038872. - _T. D. Noe_, Nov 14 2010 [Comment in brackets from _Shreevatsa R_, Mar 27 2019. For a proof see the W. Lang link, Lemma 1, iii) and the Proposition. - _Wolfdieter Lang_, Jul 04 2019] The number of fundamental solutions to n = |y^2 - x^2 - x*y| with relatively prime x and y is 0 or 2^k, where k is the number of distinct prime factors of n that are congruent to {1,4} mod 5 (conjecture). For example, n=187891=11*19*29*31 has 16=2^4 solutions; the prime n=9999999929 has 2=2^1 solutions; n=84182245951=31^3*41^4 has 4=2^2 solutions. [For a proof of the conjecture see the W. Lang link, Lemma 1, iii) and the Proposition. - _Wolfdieter Lang_, Jul 04 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Nov 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003055 | OEIS-A089305

Smallest prime of the form n*(n+1)*(n+2)...(n+k) + 1, k > 0, i.e., a(n) > n+1, or 0 if no such prime exists.

Conjecture: No entry is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003056 | OEIS-A089324

Number of lattice paths from (0,0) to the line x+y=n that use the step set {(0,1),(1,0),(2,0),(3,0),...} and never pass below y=x.

Conjecture: (n+1)*a(n) +3*(-n-1)*a(n-1) +(-5*n+13)*a(n-2) +18*(n-2)*a(n-3) +(-5*n+7)*a(n-4) +3*(-n+5)*a(n-5) +(n-5)*a(n-6)=0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003057 | OEIS-A089337

a(1) = 1; for n > 1, a(n) is the smallest number that is either a divisor or a multiple, in that priority (order), of a(n-1) such that it is a distinct palindrome not included earlier.

Conjecture: Every palindrome is a member.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003058 | OEIS-A089348

Primes of the form smallest multiple of n followed by a 1.

Conjecture: {(a(n)-1)}/{10n} < n, or a(n) < 10*n^2 + 1. The conjecture holds for 2 < n < 10^8. Heuristically, the chance of its failing for larger n is about 10^-300000 based on the integral of exp(-2.5*x/log(x)^2) using the Poisson model for numbers coprime to 10. - _Charles R Greathouse IV_, Feb 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 05 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003059 | OEIS-A089367

Smallest prime p such that np +1 is a prime, or 0 if no such prime exists.

a(2n+1) = 0 if 4n+3 is composite. Conjecture: There are no other zeros.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 08 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003060 | OEIS-A089381

L-th order palindromes with L > 2.

Let P(m) = m/2 if m is even, m + rev(m) if m is odd, where rev(m) is m's base 10 representation reversed. It is conjectured that any number k eventually cycles when P is repeatedly applied to it. If the cycle has length L, k is called an L-th order palindrome.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Darrell Plank (jar_czar(AT)msn.com), Dec 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003061 | OEIS-A089392

Magnanimous primes: primes with the property that inserting a "+" in any place between two digits yields a sum which is prime.

Partition the digits of n into two groups by placing a '+' sign anywhere inside; the result of the expression is prime in every case. Conjecture: sequence is infinite. 11 is the largest term with all odd digits. 2 is the only member with all even digits. Observation: all two-digit primes with the most significant digit even are members. In contradiction to the above conjecture, it is rather expected that this sequence is finite, cf. the link to C. Rivera's "Puzzle 401", and G. Resta's web page. Concerning the statement about 2 and 11, one can say that all terms except 2, 11 and 101 consist of even digits followed by a final odd digit. - _M. F. Hasler_, Dec 25 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003062 | OEIS-A089395

Prime productive numbers m: Let the digits of m be abcd. Then the numbers bcd*a+1, cd*ab+1, d*abc+1, abcd+1 etc. are all primes. If m is a k-digit number it produces k such primes.

Conjecture: Sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003063 | OEIS-A089494

a(n) = smallest non-palindromic k such that the Reverse and Add! trajectory of k is palindrome-free and joins the trajectory of A070788(n).

a(3), a(14) and a(18) are conjectural; it is not yet ensured that they are minimal. The 1-1 property of the mapping depends on the conjecture that the Reverse and Add! trajectory of each term of A070788 contains only a finite number of palindromes (cf. A077594). - _Klaus Brockhaus_, Dec 09 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 04 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003064 | OEIS-A089522

a(n) = smallest non-palindromic number k such that the Reverse and Add! trajectory of k joins the trajectory of A089521(n).

a(12), a(15) to a(18), a(20), a(21) are conjectural; it is not yet ensured that they are minimal.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003065 | OEIS-A089574

Column 4 of an array closely related to A083480. (Both arrays have shape sequence A083479).

First differences of the conjectured polynomial formula for A006468. [From _R. J. Mathar_, Jun 26 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alford Arnold_, Dec 29 2003; extended May 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003066 | OEIS-A089587

a(n) is the smallest integer k, 0<k<n, that most often satisfies the condition: k^m > k^(m+1) (modulo n) as m varies from 1 to n-1, for n>2, with a(1)=0 and a(2)=1.

The values at a(2^n+1), n>=0, are given by A089588: {1,2,2,7,2,9,38,79,2,220,..} and it is conjectured that a(n)=2 only when n is a Fermat number 2^(2^j)+1. a(2^(2^n)+1)=2; a(2^n-1)=2^(n-1); a(4^n-2)=4^(n-1) (conjectures).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003067 | OEIS-A089588

a(n) = A089587(2^n+1) for n >= 0.

A089587(n) is the smallest integer k, 0 < k < n, that most often satisfies the condition: k^m > k^(m+1) (modulo n) as m varies from 1 to n-1, for n > 2, with a(1)=0 and a(2)=1. It is conjectured that A089587(n)=2 only when n is a Fermat number 2^(2^j) + 1 for j >= 0. Conjecture: a(2^k+1) = 2 for k >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 09 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003068 | OEIS-A089605

Let P(m) = m/2 if m is even, m + rev(m) if m is odd, where rev(m) is m's base 10 representation reversed. It is conjectured that any number k eventually cycles when P is repeatedly applied to it. Sequence gives number of steps before the cycle is reached.

Let P(m) = m/2 if m is even, m + rev(m) if m is odd, where rev(m) is m's base 10 representation reversed. It is conjectured that any number k eventually cycles when P is repeatedly applied to it. Sequence gives number of steps before the cycle is reached.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 01 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003069 | OEIS-A089609

Prime number of primes between squares of consecutive primes; or, primes occurring in A050216.

For small values of n, these numbers exhibit higher and lower values as n increases. Conjecture: There exists an n such that seq(n1) is < seq(n1+1) for all n1 >= n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Dec 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003070 | OEIS-A089610

Number of primes between n^2 and (n+1/2)^2.

For small values of n, these numbers exhibit higher and lower values as n increases. Conjectures: After n=17 a(n) > 1. There exists an n_1 such that a(n) is < a(n+1) for all n >= n_1. Same as the number of primes between n^2 and n^2+n. Oppermann conjectured in 1882 that a(n)>0. - _T. D. Noe_, Sep 16 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Dec 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003071 | OEIS-A089618

Continued fraction elements constructed out of the van der Corput discrepancy sequence.

That conjecture has been proven by Wieting. Moreover, the r-th power mean of the sequence also converges to the corresponding constant K_r for any real r<1. - _Andrei Zabolotskii_, Feb 20 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hans Havermann_, Jan 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003072 | OEIS-A089640

Enumeration of partial sums of 1 + [1,2] + [2,3] + [1,2] + [2,3] + ...

Conjectures from _Chai Wah Wu_, Jan 13 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Jan 01 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003073 | OEIS-A089654

Table T(n,k), read by rows, related to a conjecture of P. Erdos (see A039669).

Table T(n,k), read by rows, related to a conjecture of P. Erdos (see A039669). P. Erdos conjectures that T(n,k) are all primes for n = 3, 7, 10, 22, 37, 52 and these are the only values of n with property . The conjecture has been verified for n up to 2^77. example : n=10; 19, 17, 13, 5 are all primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Jan 04 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003074 | OEIS-A089655

a(1)=1 and for n>=2 a(n) is the denominator of A(n) (see comment for A(n) definition).

For n>=2, A(n) is the least rational value >1 such that A(n)*(n^(2k)-1)*B(2k) is an integer value for k=1 up to 200, where B(2k) is the 2k-th Bernoulli number. It appears that sequence of numerators of A(n) coincide with A007947 (terms were computed by _W. Edwin Clark_). We conjecture : A(n)*(n^(2k)-1)*B(2k) is an integer value for all k>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jan 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003075 | OEIS-A089677

Exponential convolution of A000670(n), with A000670(0)=0, with the sequence of all ones alternating in sign.

Occurs also as first column of a matrix-inversion occurring in a sum-of-like-powers problem. Consider the problem for any fixed natural number m>2 of finding solutions to sum(k=1,n,k^m) = (k+1)^m. Erdos conjectured that there are no solutions for n,m>2. Let D be the matrix of differences of D[m,n] := sum(k=1,n,k^m) - (k+1)^m. Then the generating functions for the rows of this matrix D constitute a set of polynomials in n (for varying n along columns) and the m-th polynomial defining the m-th row. Let GF_D be the matrix of the coefficients of this set of polynomials. Then the present sequence is the (unsigned) second column of GF_D^-1. - _Gottfried Helms_, Apr 01 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | Mario Catalani (mario.catalani(AT)unito.it), Jan 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003076 | OEIS-A089695

Numbers m such that placing as many possible '+' signs anywhere in between the digits yields a prime in every case. Let abcd... be the digits of m; then abcd, a + bcd, ab + cd, abc + d, a + b + cd, a + bc + d, ab + c + d, a + b + c + d, ... are all prime.

Though the first 27 terms match those of A089392, the next term of A089392 (2221) is not a member of this sequence. Conjecture: sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003077 | OEIS-A089696

Numbers k such that the numbers obtained by placing as many '*' signs as possible anywhere between the digits and then adding 1 yields a prime in every case: let abc.. be the digits of k, then abc+1, a*bc+1, ab*c+1, a*b*c+1, ... must all be primes.

Though the first 14 terms match with that of A089395, the next term of A089395 306 is not a member of this sequence. Conjecture: Sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003078 | OEIS-A089705

Smallest n-digit prime in which the unit place digit is 1 and the k-th most significant digit is prime (2,3,5,7) if k is prime else is composite (4,6,8,9,0).

Conjecture: All the composite positions are occupied by a zero except the most significant digit, which in most cases perhaps is 4. Most of the prime positions for p > 5 are occupied by 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 10 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003079 | OEIS-A089708

a(1) = 1, a(2) = 2, a(n) = a(n-1) + d where d is the sum of the absolute differences between all pairs of previous terms.

Conjecture: a(n) = c n! (1+2/n+(5/2)/n^2+(31/6)/n^3+(317/24)/n^4+O(1/n^5)), where c is about 0.1289432494744. - _Dean Hickerson_, Nov 15 2003

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 14 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003080 | OEIS-A089758

Primes arising in A089757.

Conjecture: Sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003081 | OEIS-A089763

a(1) = 1, then the least n-digit prime such that the k-th digit from the Most significant side is prime if k is prime else is composite. The most significant digit is always a 1. 0 if no such prime exists.

If n is composite then the least significant digit is mandatorily a 9. Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 22 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003082 | OEIS-A089771

Largest n-digit prime containing no prime substrings, or 0 if no such prime exists.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003083 | OEIS-A089777

a(n) = smallest prime of the form n followed by a prime.

Open problem(?): show that a(n) always exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003084 | OEIS-A089781

Successive coprime numbers with distinct successive differences: gcd(a(k+1),a(k)) = gcd(a(m+1),a(m)) = 1 and a(k+1)-a(k) = a(m+1)-a(m) <==> m=k.

Conjecture: If a(k+1)-a(k) = n then k < C*n. Can someone find an estimate for the constant C?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 24 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003085 | OEIS-A089893

a(n) = (A001317(2n)-1)/4.

Conjecture: a(2^k) = 2^(2^(k+1)-2). [This conjecture is true. - _Vladimir Shevelev_, Nov 28 2010] Conjectures: lim(n->inf, a(2n+1)/a(2n)) = 5, lim(n->inf, a(4n+2)/a(4n+1)) = 17/5, lim(n->inf, a(8n+4)/a(8n+3)) = 257/85 etc. [This follows from the formula, for n>=0, t>=1: (4*a(2^t*n+2^(t-1))+1)/(4*a(2^t*n+2^(t-1)-1)+1) = 3*F_t/(F_t-2), where F_t= A000215(t) - _Vladimir Shevelev_, Nov 28 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Jan 10 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003086 | OEIS-A089936

Number of 5 X n matrices with entries {0,1} without adjacent 0's in any row or column. 5th row of A089934.

G.f.: x*(13 + 47*x - 37*x^2 - 129*x^3 + 68*x^4 + 49*x^5 - 23*x^6 - 3*x^7 + x^8) / (1 - 4*x - 36*x^2 + 105*x^4 - 15*x^5 - 64*x^6 + 20*x^7 + 4*x^8 - x^9) (conjectured). - _Colin Barker_, Jun 06 2017 The above conjecture is correct since the order of the recurrence is A089935(5) = 9. - _Andrew Howroyd_, Dec 24 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marc LeBrun_, Nov 15 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003087 | OEIS-A089937

Number of 6 X n matrices with entries {0,1} without adjacent 0's in any row or column. 6th row of A089934.

G.f.: x*(21 + 71*x - 215*x^2 - 385*x^3 + 668*x^4 + 234*x^5 - 400*x^6 + 9*x^7 + 49*x^8 - 3*x^9 - x^10) / (1 - 8*x - 62*x^2 + 78*x^3 + 375*x^4 - 300*x^5 - 486*x^6 + 385*x^7 + 30*x^8 - 52*x^9 + 2*x^10 + x^11) (conjectured). - _Colin Barker_, Jun 06 2017 The above conjecture is correct because the order of the recurrence is A089935(6) = 11. - _Andrew Howroyd_, Dec 24 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marc LeBrun_, Nov 15 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003088 | OEIS-A089941

Row sums of triangle A089940.

Conjecture: +n*(n+6)*(n-5)*a(n) -(n+6)*(n^2-3*n-6)*a(n-1) -2*(n+6)*(3*n^2-20*n

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 16 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003089 | OEIS-A089948

Smallest prime factor of the concatenation of the first n odd primes.

3 is the only number < 10000 whose concatenation of odd primes is prime. Conjecture: Beyond 3 the concatenation of odd primes produce only composite numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jan 11 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003090 | OEIS-A090072

Numbers n such that there are (presumably) eleven palindromes in the Reverse and Add! trajectory of n.

Only two numbers are known whose Reverse and Add trajectory contains twelve palindromes: 10000 and 10001. It is conjectured that these are the only such numbers and it has been conjectured before (cf. A077594) that no Reverse and Add trajectory contains more than twelve palindromes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003091 | OEIS-A090075

(Presumed) number of palindromes in the Reverse and Add! trajectory of 10^n.

The absolute maximum 12 at n = 4 and a(n) = 11 for n > 5 support the conjecture (cf. A077594) that there is no positive integer whose trajectory contains more than twelve palindromes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003092 | OEIS-A090079

In binary expansion of n: reduce contiguous blocks of 0's to 0 and contiguous blocks of 1's to 1.

Conjecture: a(n) = (2^(A005811(n)+1) + (1-(-1)^n)/2 - 2)/3. - _Velin Yanev_, Dec 12 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Nov 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003093 | OEIS-A090086

Smallest pseudoprime to base n, not necessarily exceeding n (cf. A007535).

Conjecture: a(n) = A007535(n) for finitely many n. For n > 2; if a(n) > n, then n-1 is prime (find all these primes). - _Thomas Ordowski_, Aug 09 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Nov 25 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003094 | OEIS-A090196

Odd integers with two divisors a, b such that a < b <= 2a.

Clearly all even integers have two such divisors a, b. Consider the set S of all integers satisfying this property. Maier & Tenenbaum proved Erdős' conjecture that S has asymptotic density 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Finch_, Jan 22 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003095 | OEIS-A090238

Triangle T(n, k) read by rows. T(n, k) is the number of lists of k unlabeled permutations whose total length is n.

Conjectural o.g.f.: 1/(1 + t - t*F(x)) = 1 + t*x + (2*t + t^2)*x^2 + (6*t + 4*t^2 + t^3)*x^3 + ..., where F(x) = Sum_{n >= 0} n!*x^n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Jan 23 2004, Jun 14 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003096 | OEIS-A090252

The Two-Up sequence: a(n) is the least positive number not already used that is coprime to the previous floor(n/2) terms.

Conjecture 1. A090252 is a subsequence of A354144 (prime powers and semiprimes). Conjecture 2. The terms of A354144 that are missing from A090252 are 6, 10, 14, 15, 22, 33, 34, 35, 38, 39, 46, 51, 58, 62, 65, 69, 74, 77, 82, 86, 87, 91, 93, 94, 95, 106, 111, 115, 118, 119, 122, 123, 129, 133, 134, 141, 142, 143, 145, 146, 155, 158, 166, 177, 178, 183, 185, 187, 194, 201, 202, 203, 209, 213, 214, 215, 218, 219, 221, ... Conjecture 3. For a prime p, there are constants v_1, v_2, ..., v_K and c such that (This sequence of lambdas does not seem to have any simpler explanation, is not in the OEIS, and cannot be since the terms shown are all conjectural.) Conjecture 2 is a consequence of Conjecture 3. For example, 6 does not appear in A090252, since the sets S_2 and S_3 are disjoint. Assuming the numbers that appear to be missing (see Conjecture 2) really are missing, the numbers that take a record number of steps to appear are 1, 2, 3, 4, 7, 8, 16, 26, 32, 64, 128, 206, 256, 478, 512, 933, ..., and the indices where they appear are 1, 2, 3, 5, 6, 11, 23, 47, 95, 191, 383, 767, 1535, 3071, 6143, 8191, ... These two sequences are not yet in the OEIS, and cannot be added since the terms are all conjectural. If we consider the May 23 2022 comment, note the conjectured indices show near complete overlap with terms of A081026: 1, 2, 3, 5, 6, 11, 23, 47, 95, 191, 383, 767, 1535, 3071, 6143, 8191. - _Bill McEachen_, Aug 09 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003097 | OEIS-A090257

a(1) = 1, then least square such that every partial concatenation is a prime.

a(n) == 0 (mod 9) for n>1. Conjecture: Sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003098 | OEIS-A090260

Beginning with 1, distinct numbers such that concatenation a(n+1), a(n), a(n+1) is a prime.

Conjecture: all positive integers coprime to 10 occur in the sequence. - _Robert Israel_, Jan 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 28 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003099 | OEIS-A090287

Smallest prime obtained by sandwiching n between a number with identical digits, or 0 if no such prime exists. Primes of the form k n k where all the digits of k are identical.

a(n) = 0 if n is a palindrome with even number of digits. Conjecture: No other term is zero. The conjecture is false. a(231) = 0, a(420) = 0, a(n) = 0 if 11 divides n and n has an even number of digits. a(1414) has over 2000 digits. - _Chai Wah Wu_, Mar 31 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003100 | OEIS-A090291

Index of first occurrence of n in A090290, or 0 if n does not occur in A090290.

10^(n) -prime(a(n)) is a prime and least such prime. Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003101 | OEIS-A090303

Begin with n and consider numbers obtained by successively subtracting 2,3,5,7,11,...; a(n) = largest prime that arises, i.e., largest prime of the form n - sum prime(r), or 0 if no such prime exists.

Conjecture: There are finitely many zeros in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003102 | OEIS-A090304

Indices of 0's in A090303.

Conjecture: sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003103 | OEIS-A090315

Least k such that k and digit reversal of k both have n divisors, or 0 if no such number exists.

For a(7) one needs a number of the form p^6 whose digit reversal is q^6, p, q are primes. Hence a(7) perhaps is zero (not sure). Conjecture: There are infinitely many nonzero terms as well as zeros in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003104 | OEIS-A090335

Smallest number with exactly n distinct prime factors such that in binary representation the divisors >1 form a prefix code.

Conjecture: the sequence is defined for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Nov 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003105 | OEIS-A090365

Shifts 1 place left under the INVERT transform of the BINOMIAL transform of this sequence.

Conjecture: a(n) = b(2^(n-1) - 1) for n > 0 with a(0) = 1 where b(n) = b((n - 2^f(n))/2) + b(floor((2n - 2^f(n))/2)) + b(A025480(n-1)) for n > 0 with b(0) = 1 and where f(n) = A007814(n). - _Mikhail Kurkov_, Jan 11 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 26 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003106 | OEIS-A090381

Expansion of (1+4x+7x^2)/((1-x)^2*(1-x^2)).

Conjecture: a(n) is the maximum number of areas that are defined by the 3n angle (n+1)-sectors in a triangle. - _Nicolas Nagel_, Sep 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 30 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003107 | OEIS-A090461

Numbers k for which there exists a permutation of the numbers 1 to k such that the sum of adjacent numbers is a square.

Conjecture: sequence includes all integers k > 24. See A090460 for the number of essentially different solutions. The conjecture has been proved: every k >= 25 is in the sequence, moreover for k >= 32 there is a Hamiltonian cycle; see Mersenneforum topic for a code and deterministic algorithm to find a sequence. - _Robert Gerbicz_, Jan 21 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Dec 01 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003108 | OEIS-A090469

Least (distinct) composite k such that k+n, k+2n, k+3n, ... up to n terms are all composite.

Conjectures: 1. For n > 1, a(n) is the least composite k not already used such that gcd(n, k) > 1. 2. For any prime p > 5, a(p) = 2p. - _David Wasserman_, Nov 08 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003109 | OEIS-A090483

Indices of zero in A090482.

I conjecture there are no more terms; I've checked up to 200. - _David Wasserman_, Nov 17 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 02 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003110 | OEIS-A090494

Product_{j=1..n} Product_{k=1..n} lcm(j,k).

Let p be a prime and let ordp(n,p) denote the exponent of the largest power of p which divides n. For example, ordp(48,2)=4 since 48 = 3*(2^4). Then the prime factorization of a(n) appears to be given by the formula ordp(a(n),p)= sum_{k >= 1} [(2*(p^k)-1)*floor((n/(p^k)))^2] + 2*sum_{k >= 1} [floor(n/(p^k))*mod(n,p^k)], for each prime p. See the comments sections of A092143, A092287, A129365 and A129454 for similar conjectural prime factorizations. - _Peter Bala_, Apr 23 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003111 | OEIS-A090496

Ratio of numerator(Bernoulli(2*n)/(2*n)) to numerator(Bernoulli(2*n)/(2*n*(2*n-1))) for n's for which they are different.

All values yielding 37 are of the form 574+666*k, k=0,1,2,3,4,... and form thus an arithmetic progression with step 666=18*37=((37-1)/2)*37. All values yielding 59 are of the form 1269+1711*k, k=0,1,2,3 and 1711=28*59=((59-1)/2)*59. The two values yielding 67 are at distance 2211=((67-1)/2)*67. Conjecture: all indices yielding a given prime p form an arithmetic progression of step ((p-1)/2)*p. See A092291. - _Roland Bacher_, Feb 04 2004 Roland Bacher conjectures that values of n yielding the same quotient p form an arithmetic progression n0+d*k, where d = p(p-1)/2. Actual and conjectured values of n0 are in the sequence A092291. Composite values do occur. An example is 2n = 272876, which yields a quotient of 37*59. This was found by tdn using the Kummer congruences and CRT: using the irregular pairs (37,32) and (59,44), we know that the following Diophantine equations must be solved for (k,l,m): 32+36*k = 44+58*l = 1+37*59*m. Some quotients are not possible, e.g., 37*67, 37*103. All quotients are the product of irregular primes A000928. Composite quotients imply there are missing terms in the arithmetic progression conjectured by Bacher. - _T. D. Noe_, Feb 12 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003112 | OEIS-A090518

Primes arising in A090517, or 0 if A090517(n) = 0.

Conjecture: No term is zero. The conjecture is true. For every n >= 1, k = 5*10^(n-1) gives floor(10^n/k) = 2, which is prime. Hence the set of admissible k is nonempty, so A090517(n) is well-defined. The corresponding quotient A090518(n) is prime by definition and therefore nonzero. - _Lien-Hung Su_, Oct 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003113 | OEIS-A090519

Smallest prime p such that floor((10^n)/p) is prime, or 0 if no such number exists.

Conjecture: No term is zero. Subsidiary Sequence: Number of primes in floor((10^n)/p), p is a prime. a(1) = 3, the primes are 10/2, floor(10/3) and 10/5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003114 | OEIS-A090520

Primes arising in A090519, or 0 if A090519(n)= 0.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003115 | OEIS-A090521

Least k such that floor(n!/k) is prime.

Conjecture: There exists a number k such that for all n > k, a(n) is prime. Motivation: If p is the least prime >n then all the numbers from n to p-1 divide n!. And most of the numbers from p+1 to q also divide n! where q is the least prime > p, etc. and the dividend is composite in almost all cases. I conjecture the opposite: there are infinitely many composites in this sequence. Indeed, heuristics suggest the composites may be of density 1. - _Charles R Greathouse IV_, Apr 07 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003116 | OEIS-A090523

Smallest prime p such that floor(n!/p) is prime, or 0 if no such prime exists.

Conjecture: There are no zeros for n > 2. This conjecture is correct. For m > 1, there is always a prime between m and 2*m. Taking m = n!/4, this gives us a prime p such that floor(n!/p) = 2 or 3. - _Franklin T. Adams-Watters_, Jul 28 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003117 | OEIS-A090524

Primes arising in A090523, or 0 if no such prime exists.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003118 | OEIS-A090525

Least number k such that floor((n^n)/k) is prime, or 0 if no such number exists.

Conjecture: No term is zero. The conjecture is true: If p is a prime factor of n, k = (n^n)/p gives an upper bound. - _James Rayman_, Mar 01 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003119 | OEIS-A090526

Primes arising in A090525, or 0 if A090525(n) = 0.

Conjecture: No term is zero. Conjecture is true: see A090525. - _Robert Israel_, Mar 02 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003120 | OEIS-A090527

Smallest prime p such that floor((n^n)/p) is prime, or 0 if no such number exists.

Conjecture: No term is zero. As long as p(j+1)/p(j) < 2 for all j, then for any integer n >= 4, there exists at least one p such that p and floor(n/p) are both prime. (I do not know a proof for the premise above; however, it seems quite weak compared to other conjectures and theorems about primes. It may be that it follows from the results in sequence A038458.) In fact, there exists a prime p such that either floor(n/p) = 2 or floor(n/p) = 3. Outline of proof: (1) If p is a prime number, then for all n with 2p <= n < 3p, floor(n/p) = 2, which is prime. (2) In addition, for all n with 3p <= n < 4p, floor(n/p) = 3, which is prime. So for any n >= 4, consider the largest prime, p, with 2p <= n. (3) floor(n/p) can't be less than 2, since 2 <= n/p. (4) If floor(n/p) = 2, then p and floor(n/p) are both prime, so we are done. (5) Similarly, if floor(n/p) = 3, we are done. The only remaining case is that 4p <= n. Let p_1 be the next prime after p. (6) p_n must not meet 2(p_1) <= n, since p is the largest that does. Therefore 2(p_1) > n. (7) 4p <= n < 2(p_1) (8) (p_1 / p) > 2 (9) As long as p(j+1)/p(j) < 2 for all j, the case of 4p <= n is not possible. - Weston Markham (WMarkham(AT)paradigmgenetics.com), Jun 15 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003121 | OEIS-A090528

Primes arising in A090527.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003122 | OEIS-A090530

Least multiple k of prime(n) such that (k-1,k+1) forms a twin prime pair, or 0 if no such number exists.

a(n) is a multiple of 6*prime(n) for n>2. Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 07 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003123 | OEIS-A090534

Least n-digit prime in which every two-digit string is also a prime, or 0 if no such number exists. (n-1 two-digit string primes occur.)

Conjecture: sequence is infinite. Notes: (1) If n==1 (mod 3) then in most cases a(n) is a permutation of (n-1), 1's and a 7, else a(n) is a digit permutation of (n-1) ones and a 3; (2) Obviously all the digits except the most significant one must be odd.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 08 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003124 | OEIS-A090634

Start with the sequence [1, 1/2, 1/3, ..., 1/n]; form new sequence of n-1 terms by taking averages of successive terms; repeat until reach a single number F(n); a(n) = denominator of F(n).

a(n) = denominator(2*(1-1/2^n)/n) (conjectured). - _Michel Marcus_, Sep 12 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003125 | OEIS-A090691

5x - 1 sequence starting at 50 (a(n+1) = a(n)/2 if a(n) is even, or 5*a(n)-1 if a(n) is odd).

There is a general conjecture (see, e.g., the paper of Heppner, Eine Bemerkung zum Hasse-Syracuse-Algorithmus, Arch. Math. (Basel) 31 (1978/79), 317-320, that for the 5x-1 problem there are divergent sequences (since 5 > 2^{2/(2-1)}, although for the Collatz problem 3 < 2^{2/(2-1)}). - _Jean-Paul Allouche_, Sep 14 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | Daniil Khanin (daniil(AT)ispms.tsc.ru), Feb 06 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003126 | OEIS-A090738

Integers n such that the concatenation of n^2 and (n+1)^2 is prime.

I conjecture this sequence to be infinite. Searching through the first 200000 values, I found 7000 primes, of which over 400 were "twins", i.e. both n^2*(n+1)^2 and (n+2)^2*(n+3)^2 were prime, where "*" denotes concatenation. I conjecture there to be an infinitude of such twins and the obvious generalizations.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Alex Kontorovich (alexk(AT)math.columbia.edu), Jan 19 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003127 | OEIS-A090780

a(n) = n*Product_{p prime, p|n} (p - 1)/2.

a(2n+1) is the conjectured value of the length of period of sequence of Genocchi number of first kind read modulo (2n + 1) (cf. A001469).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 12 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003128 | OEIS-A090786

Least nonnegative integer k such that n! + n + k + 1 is prime.

Conjectures: By definition a(n)+n!+1 is prime, but is a(n)+n+1=A037153(n) also a prime number for all n > 2? Is the growth of b(n) := Sum_{k=0..n} a(k) quadratic, that is, is b(n)=O(n^2)?

猜想/问题提及段 | historical_or_current_mention_unreviewed | Frederick Magata (frederick.magata(AT)uni-muenster.de), Feb 09 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003129 | OEIS-A090825

Nonprimes n such that (3/2)*(1/n)*(2*n+1)*(3^n+1)*B(2*n) is an integer, where B(k) denotes the k-th Bernoulli number.

Conjecture: composite numbers with all prime factors in A053176 are in the sequence. For p prime (3/2)*(1/p)*(2*p+1)*(3^p+1)*B(2*p) == 1 (mod p). There are few terms n with (3/2)*(1/n)*(2*n+1)*(3^n+1)*B(2*n) == 1 (mod n): 91,247,....Is this subsequence finite?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 11 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003130 | OEIS-A090826

Convolution of Catalan and Fibonacci numbers.

Conjecture: n*a(n) + (-5*n+6)*a(n-1) + 3*(n-2)*a(n-2) + 2*(2*n-3)*a(n-3)=0. - _R. J. Mathar_, Jul 09 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 20 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003131 | OEIS-A090866

Primes p == 1 (mod 4) such that (p-1)/4 is prime.

Same as Chebyshev's subsequence of the primes with primitive root 2, because Chebyshev showed that 2 is a primitive root of all primes p = 4*q+1 with q prime. If the sequence is infinite, then Artin's conjecture ("every nonsquare positive integer n is a primitive root of infinitely many primes q") is true for n = 2. - _Jonathan Sondow_, Feb 04 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 12 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003132 | OEIS-A090867

Number of partitions of n such that the set of even parts has only one element.

Conjecture: a(n) is also the difference between the number of parts in the odd partitions of n and the number of parts in the distinct partitions of n (offset 0). For example, if n = 5, there are 9 parts in the odd partitions of 5 (5, 311, 11111) and 5 parts in the distinct partitions of 5 (5, 41, 32), with difference 4. - _George Beck_, Apr 22 2017 George E. Andrews has kindly informed me that he has proved this conjecture and the result will be included in his article "Euler's Partition Identity and Two Problems of George Beck" which will appear in The Mathematics Student, 86, Nos. 1-2, January - June (2017). - _George Beck_, Apr 23 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Feb 12 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003133 | OEIS-A090904

Row products of the irregular triangle defined in A090905.

Conjecture: For n > 4 the last term of the n-th group is 2p where p is the largest prime in the (n-1)th group. And these are the Bertrand primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 13 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003134 | OEIS-A090909

Terms a(k) of A073869 for which a(k-1) = a(k), and a(k) and a(k+1) are distinct.

Proof of Mihailovs's conjecture: This follows immediately from the result in my 2023 paper in JIS that A073869 is equal to Hofstadter's G-sequence A005206, and my recent comment in A005206 on the pairs of duplicate values in A005206.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 14 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003135 | OEIS-A090940

a(n) = least odd prime distinct from earlier elements such that average of first n elements is prime.

Conjectured to include all odd prime numbers. - _David W. Wilson_, Nov 23 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Dec 29 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003136 | OEIS-A090975

Least integer k such that n!+1-k is prime.

The (n-1) consecutive numbers n!+2,...,n!+n (for n>=2) are not prime. This fact implies that there are arbitrarily large gaps in the distribution of the prime numbers. n!+1 itself may be a prime number as in the case of n=3, 11, 27 (see A002981 for all such n). Now a(n) measures, when the first prime number previous to n!+2 appears. Thus a(n)=8 means that n!+1-3 is prime and so on. Obviously, the values of a(n) are always even numbers. Conjectures: |a(n)-1| is either 1 or a prime number. Is the growth of b(n) := sum(a(k),k=0..n) quadratic, that is b(n)=O(n^2)?

猜想/问题提及段 | historical_or_current_mention_unreviewed | Frederick Magata (frederick.magata(AT)uni-muenster.de), Feb 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003137 | OEIS-A090981

Triangle read by rows: T(n,k) = number of Schroeder paths of length 2n and having k ascents.

The expected number of ascents in a Schroeder n-path is asymptotically (sqrt(2)-1)*n for large n. (Previously formulated as conjecture by _David Callan_, Jul 25 2008, now proven.) - _Valerie Roitner_, Aug 06 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Feb 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003138 | OEIS-A091067

Numbers whose odd part is of the form 4*k+3.

Alternate definition of the same sequence: a(1)=3 and a(n) is the smallest number > a(n-1) so that no number that is the sum of at most 2 terms in this sequence is a power of 2. Conjectured by _J. Lowell_, Jan 20 2024; proved by _Jaisal Khurana_, Sep 24 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Feb 22 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003139 | OEIS-A091137

The Hirzebruch numbers. a(n) = Product_{2 <= p <= n+1, p prime} p^floor(n / (p - 1)).

Conjecture: The denominators of the series reversion of the sequence with e.g.f. Polylog(2,x). - _Benedict W. J. Irwin_, Jan 05 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henry Bottomley_, Dec 19 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003140 | OEIS-A091155

Numbers m such that m - 2^k is squarefree for all 1 <= 2^k < m.

Erdős conjectures that this sequence is infinite. It appears that m == 3 (mod 12) except for m = 2, 4, 7 and 23.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Dec 23 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003141 | OEIS-A091182

Number of ways to write n = x + y (x >= y > 0) with xy - 1 and xy + 1 both prime.

Conjecture: a(n) > 0 for all n > 3120. This has been verified for n up to 5*10^7. Note that if x >= y > 0 and x+y = n then n-1 = x+y-1 <= xy <= ((x+y)/2)^2 = n^2/4. So the conjecture implies that there are infinitely many twin primes. More generally, we have the following conjecture: Let m be any positive integer. If n is sufficiently large and (m-1)n is even, then we can write n as x+y, where x and y are positive integers with xy-m and xy+m both prime. This general conjecture implies that for any positive even integer d there are infinitely many primes p and q with difference d. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ray Chandler_, Dec 27 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003142 | OEIS-A091323

Minimum number of transversals in a Latin square of order 2n+1.

Ryser conjectured that a(n) >= 1 for all n. For even orders the number is 0, since the group table for Z_2n has no transversals.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Bean_, Feb 17 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003143 | OEIS-A091338

a(n) = (3/n), where (k/n) is the Kronecker symbol.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Dec 30 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003144 | OEIS-A091340

Amicable numbers with property that each member m of the corresponding amicable pair is divisible by sopfr(m) (the sum of prime factors with repetition, A001414).

Conjecture: sequence is finite (even though there are many known amicable numbers - about 8*10^7 currently).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sven Simon_, Dec 31 2003 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003145 | OEIS-A091351

Triangle T, read by rows, such that T(n,k) equals the (n-k)-th row sum of T^k, where T^k is the k-th power of T as a lower triangular matrix.

Conjecture: given this triangle T as a triangular matrix, then we have the relation [T^p](n+q-1,q-1) = [T^q](n+p-1,p-1) for n >= 0, p > 0, q > 0; that is, column q-1 of matrix power T^p equals column p-1 of matrix power T^q, for all positive integers p and q. - _Paul D. Hanna_, Jul 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 02 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003146 | OEIS-A091361

Numbers n such that A001840(n) == 0 (mod n).

Apart from 1 and 2 it is conjectured that the only values present are congruent to 3 mod 6 (all these values are present). Proof of the conjecture that this is 1 and 2 followed by A016945 follows by considering the 6 cases n=6k-1, 6k, 6k+1, 6k+2, 6k+3 or 6k+4, individual evaluation of A001840(n) with their corresponding 3 formulas quoted in A001840 in each case and searching for solutions of the form A001840(n) = t*n for integer t. G.f.: conjecture: 2*(1+x)/(1-x)/G(0) +x, where G(k)= 1 + 1/(1 - x*(3*k+1)/(x*(3*k+4) + 1/G(k+1))); (continued fraction). - _Sergei N. Gladkovskii_, Jun 06 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Mar 01 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003147 | OEIS-A091366

Primes p such that the sum of the cubes of the digits of p is prime.

I conjecture the contrary: the relative density of numbers in this sequence with prime digit sum is 0. - _Charles R Greathouse IV_, Sep 08 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chuck Seggelin_, Jan 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003148 | OEIS-A091443

Multiperfect numbers n which are divisible by sopfr(n) (multiperfect number: sigma(n) = k*n with k integer, sopfr: Sum of prime factors with repetition).

Conjecture: the sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sven Simon_, Jan 10 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003149 | OEIS-A091457

Greatest numerator of the remainder in a reciprocal expansion of 1.

Conjecture: in the "extremal" expansion x_i = A000058(i) for i=1..n-3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Jan 11 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003150 | OEIS-A091508

Least k such that b(k) = 1, where b(1) = n and b(k+1) = b(k)+k+1 if gcd(k+1, b(k)) = 1 or else b(k)/gcd(k+1, b(k)).

I conjecture a(n) always exists. That means sequence (b(k)) becomes ultimately regular for any n. i.e. there is always k0 such that b(k0) = 1, so b(k0+1) = b(k0) + k0 = k0 + 1 since gcd(k0, b(k0)) = 1 and gcd(k0+1, b(k0+1)) = k0+1 implies b(k0+2) = b(k0+1)/(k0+1) = 1 and from that point k0 sequence (b(k)) continues : 1, k0+1, 1, k0+2, 1, k0+3,1,... and is "regular". Benoit's conjecture is true for all n <= 10^15. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Mar 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003151 | OEIS-A091524

a(m) is the multiplier of sqrt(2) in the constant alpha(m) = a(m)*sqrt(2) - b(m), where alpha(m) is the value of the constant determined by the binary bits in the recurrence associated with the Graham-Pollak sequence.

Conjecture: a(n) = sqrt(A028982(n)/A006337(n)). - _Mikhail Kurkov_, Apr 25 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Jan 18 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003152 | OEIS-A091527

a(n) = ((3*n)!/n!^2)*(Gamma(1+n/2)/Gamma(1+3n/2)).

Conjecture: the stronger supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) hold for primes p >= 5 and positive integers n and k. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Jan 18 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003153 | OEIS-A091529

When sum {k=1 to n} H(k) k! (n-k)! (mod {n+1}) (A091528) is not zero.

It is conjectured that the density decreases. In each century the number of members are so far: 41, 30, 26, 28, 23, 24, 24, 21, 24, 22, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Jan 08 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003154 | OEIS-A091539

Second column (k=3) of array A091534 ((5,2)-Stirling2) divided by 10.

Conjecture D-finite with recurrence a(n) -(3*n-1)*(6*n-5)*a(n-1) +3*(3*n-5)*(n-1)*(3*n-1)*(3*n-4)*a(n-2)=0. - _R. J. Mathar_, Jul 15 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Feb 13 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003155 | OEIS-A091540

Rescaled second column A091539 of array A091534 ((5,2)-Stirling2).

The recurrence conjectured by _R. J. Mathar_ on Jul 15 2026 follows from the triple factorial formula. Put q(n) = Product_{j=0..n-1} (3*j+1). Then a(n) = (3^n*n! - 3*q(n))/6, so a(n) - 3*n*a(n-1) = q(n-1) for n >= 3. Since q(n-1) = (3*n-5)*q(n-2), we obtain a(n) - 3*n*a(n-1) = (3*n-5)*(a(n-1) - 3*(n-1)*a(n-2)) for n >= 4. Rearranging gives a(n) = (6*n-5)*a(n-1) - 3*(n-1)*(3*n-5)*a(n-2), as conjectured. - _Eiver Rodríguez_, Sep 20 2026 Conjecture D-finite with recurrence a(n) +(-6*n+5)*a(n-1) +3*(n-1)*(3*n-5)*a(n-2)=0. - _R. J. Mathar_, Jul 15 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Feb 13 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003156 | OEIS-A091588

A smoothed version of A091587.

a(n+1) = a(n) + A357063(n+1) + A091840(n+1). See Conjecture 4.2 of F. J. van de Bult et al., proved p. 54 of Levi van de Pol. - _Levi van de Pol_, Nov 04 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 05 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003157 | OEIS-A091590

Number of terms in the simple continued fraction for the 10^n-th harmonic number, H_n = sum_{k=1 to n} (1/k).

Conjecture: lim n -> infinity, a(n)/10^n -> C = 12*log(2)/Pi^2 = 0.842... - _Benoit Cloitre_, May 04 2002

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jan 22 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003158 | OEIS-A091592

Numbers n such that there are no twin primes between n^2 and (n+1)^2.

The first 7 terms of this sequence were given by Ernst Jung in a discussion in the Newsgroup de.sci.mathematik entitled "Primzahlen zwischen (2x-1)^2 und (2x+1)^2" (primes between ...and...) with other significant contributions from

Hermann Kremer and Rainer Rosenthal. It is conjectured that there are no further terms beyond a(12)=122. This has been tested to 50000 by _Robert G. Wilson v_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jan 25 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003159 | OEIS-A091593

Reversion of Jacobsthal numbers A001045.

Conjecture: (n+2)*a(n) +(2*n+1)*a(n-1) +9*(n-1)*a(n-2)=0. - _R. J. Mathar_, Nov 26 2012 R. J. Mathar's conjecture confirmed by Maple using this hypergeom form. - _Robert Israel_, Sep 22 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 23 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003160 | OEIS-A091666

Difference between prime(n)^2 and the next prime.

Conjecturally, a(n) << log^2 n (with constant around 8/e^gamma in the supremum). [_Charles R Greathouse IV_, Dec 27 2011] Conjecture: Limit_{N->oo} (Sum_{n=1..N} a(n)) / (Sum_{n=1..N} log(prime(n))) = 2. - _Alain Rocchelli_, Oct 04 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Jan 27 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003161 | OEIS-A091675

Positive integers n such that the trajectory of n under the Reverse and Add! operation carried out in base 4 (presumably) does not join the trajectory of any m < n.

The conjecture that the base-4 trajectories of the terms do not join is based on the observation that if the trajectories of two integers below 4120 join, this happens after at most 28 steps, while for any two terms listed above the trajectories do not join within 1000 steps. For pairs from 1, 3, 21, 22, 26, 29, 31, 55 this has even been checked for 5000 steps.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Jan 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003162 | OEIS-A091677

a(n) = smallest non-palindromic k such that the base-4 Reverse and Add! trajectory of k is palindrome-free and joins the trajectory of A091675(n).

a(1), a(5), a(22), a(23) and a(30) are conjectural; it is not yet ensured that they are minimal. The 1-1 property of the mapping depends on the conjecture that the base-4 Reverse and Add! trajectory of each term of A091675 contains only a finite number of palindromes (cf. A091680).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Jan 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003163 | OEIS-A091680

Smallest number whose base-4 Reverse and Add! trajectory (presumably) contains exactly n base-4 palindromes, or -1 if there is no such number.

Conjecture 1: For each k > 0 the trajectory of k eventually leads to a term in the trajectory of some j which belongs to A075421, i.e., whose trajectory (presumably) never leads to a palindrome. Conjecture 2: There is no k > 0 such that the trajectory of k contains more than twelve palindromes, i.e., a(n) = -1 for n > 12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Jan 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003164 | OEIS-A091689

Smallest partition number with n-th prime as factor.

Erdős conjectured that every prime divides at least one value of the partition function, see Ahlgren and Ono link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jan 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003165 | OEIS-A091699

Row sums of triangle A091698.

Conjecture: 2*n*a(n) + (13*n-16)*a(n-1) + 4*(4*n-7)*a(n-2) + 5*(n-2)*a(n-3) = 0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, Jan 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003166 | OEIS-A091713

G.f. A(x) satisfies A(x) = x + x*A(A(A(x))).

Conjecture: all terms are odd. - _Paul D. Hanna_, Dec 01 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 31 2004, Jun 04 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003167 | OEIS-A091752

Generalized Stirling2 array (-1,2)S2. Irregular triangle a(n, m) for n >= 1 and 2 <= m <= 2*n.

For the case of the (irregular) triangles (-1,s)S2, for s >= 1, W. Schulte conjectured that a(s; n, m) = T(s; (s+1)*n - m - 1, m - s), for n > = 1 and m = s, s+1, ..., s*n, with the row polynomials RT(s; n, x) = Sum_{m = 0..s*n} T(s; n, m)*x^m of the triangle T(s), n >= 0, defined by the Rodrigues-type formula exp(x^(s+1)/(s+1)) (d/dx)^n exp(-x^(s+1)/(s+1)) = (-1)^n*RT(s; n, x). Thus the rows of (-1,s)S2 are every (s+1)-th upwards antidiagonals of T(s), but with offset m = s instead of m = 0. The proof of the conjecture follows by showing the Carlitz recurrence for (-1,s)S2, but with offset n = 0 and m = 0; that is, a(s; n+1, m+s) =: aHat(s; n, m) = Sum_{j = 0..s} binomial(s, j)*fallfac(s + m - j - (s+1)*n , s-j)*aHat(s; n-1, m-j), for n >= 0, m = 0, 1, ..., s*n, with aHat(s; 0, 0) = 1 and aHat(s; n, m) = 0 for m < 0 and m > s*n. The falling factorials are fallfac(x, n). With aHat(s; n, m) = T(s; (s+1)*n - m, m) this leads to a recurrence for T(s) which is equivalent to the recurrence for the row polynomials RT(s), namely RT(s; n, x) = Sum_{j=0 .. s} binomial(s, j)*x^j*fallfac(s - j - n, s - j)*RT(s; n - (s+1) + j, x), for n >= 1, and RT(s; 0, x) = 1. This, in turn, can be proved by induction over n >= 1 from the simpler recurrence for RT(s) obtained directly from the Rodrigues-type definition, namely RT(s; n, x) = x^s*RT(s; n-1, x) - (d/dx)RT(s; n-1, x), n >= 1,with RT(s; 0, x) = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Feb 27 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003168 | OEIS-A091849

Maximal size of (n,4,2) optical orthogonal code.

Conjectured to agree with A091848 except for n = 9 and 13.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 13 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003169 | OEIS-A091892

Numbers k having only one partition into parts which are a sum of exactly as many distinct powers of 2 as there are 1's in the binary representation of k.

Conjecture: if the second leftmost bit in the binary expansion of k+1 equals 0, then k is a term if and only if A007814(k+1) >= 2^(f(k)-1) + f(k). Otherwise, k is a term if and only if A007814(k+1) >= 2^f(k). Here f(k) = A086784(k+1). - _Mikhail Kurkov_, Oct 03 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Feb 10 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003170 | OEIS-A091915

Maximum of even products of partitions of n.

For n>6, a(n+3) = 3a(n) (conjectured). - _Ralf Stephan_, Dec 02 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Feb 12 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003171 | OEIS-A091916

Maximum of odd products of partitions of n.

For n>5, a(n+3) = 3*a(n) (conjectured). - _Ralf Stephan_, Dec 02 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Feb 12 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003172 | OEIS-A091984

Number of steps required for initial p = 10^n to reach 1 in the recurrence p = floor(Li(p)).

The number of steps for the recurrence in this sequence stopping at 1 compares closely to the steps in the pi(n) recurrence stopping at 0. If we define Li(1) = 0 and allow that step then A(using Li seq#) - A(using pi seq#) = 1 for n <=13. So one may conjecture the steps in the Li method is always 1 greater than the steps in the pi method. Question is can the difference be greater than 1? For the largest value allowed in the link (3*10^13) A(Li) = 17 (assuming L1(1)= 0) and n=13 A(pi) = 16 from Booker so the difference = 1 as before.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Mar 16 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003173 | OEIS-A092028

a(n) is the smallest m > 1 such that m divides n^m-1.

Each prime factor of n-1 is a solution of the equation n^x-1 == 0 (mod x), so a(n) is not greater than the smallest prime factor of n-1. Conjecture 1: All terms of this sequence are primes. Conjecture 2: a(n) is the smallest prime factor of n-1 or for n>2, A092028(n) = A020639(n-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Mar 26 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003174 | OEIS-A092035

Decimal expansion of (Pi/e)^2.

Conjecture: Equals 12*Sum_{k >= 1} ((-1)^(k+1)/k^2) * cos(sqrt(2*k^2*Pi^2/3 - 4)). Cf. A100074. - _Peter Bala_, Jun 20 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mohammad K. Azarian_, Mar 26 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003175 | OEIS-A092067

a(n) is the smallest number m such that m > 1 and m divides n^m + 1.

Conjecture 1: All terms of this sequence are primes. We know if n is odd a(n) is the smallest prime factor of n+1. Conjecture 2: For each n, a(n) is the smallest prime factor of n+1 or a(n)=A020639(n+1). The theorem plus the conjecture 2 in A092028 imply a(n) = A092028(n+2). - _R. J. Mathar_, Mar 21 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Mar 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003176 | OEIS-A092097

Limit number of (m-n)-almost-primes in range [2^m..2^{m+1}-1].

The value of m at which the number of (m-n)-almost-primes reaches its limit is floor(n/(log_2(3)-1))+n-1: 1,4,7,9,12,15,17,20,23,26,28; not A026356: 2,4,7,9,12,15,17,20,22,25,28 as originally conjectured. - _Franklin T. Adams-Watters_, Jun 20 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew S. Plewe_, Feb 19 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003177 | OEIS-A092101

Harmonic primes.

For p = prime(n), Boyd defines J_p to be the set of numbers k such that p divides A001008(k), the numerator of the harmonic number H(k). For harmonic primes, J_p contains only the three numbers p-1, (p-1)p and (p-1)(p+1). It has been conjectured that there are an infinite number of these primes and that their density in the primes is 1/e.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Feb 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003178 | OEIS-A092103

Number of values of k for which prime(n) divides A001008(k), the numerator of the k-th harmonic number.

Eswarathasan and Levine conjectured that for any prime number p the set J_p is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Feb 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003179 | OEIS-A092119

Euler transform of A001511.

Let M = triangle A173238 as an infinite lower triangular matrix. Then A092119 = lim_{n->infinity} M^n. Let P(x) = polcoeff A000041 = (1 + x + 2x^2 + 3x^3 + ...), and A(x) = polcoeff A092119. Then P(x) = A(x) / A(x^2), an example of a conjectured infinite set of operations (cf. A173238). - _Gary W. Adamson_, Feb 13 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Mar 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003180 | OEIS-A092131

Distance from 2^n to the next prime.

Conjecture: Every odd number which is not a Sierpiński number is a term. In other words, for every odd k which is not a Sierpiński number, there exists some n >= 1 such that 2^n + 1, 2^n + 3, ..., 2^n + (k-2) are all composite while 2^n + k is prime. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Helmut Richter (richter(AT)lrz.de), Mar 30 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003181 | OEIS-A092143

Cumulative product of all divisors of 1..n.

Let p be a prime and let ordp(n,p) denote the exponent of the largest power of p which divides n. For example, ordp(48,2) = 4 since 48 = 3*(2^4). Let b(n) = A006218(n) = Sum_{k = 1..n} floor(n/k). The prime factorization of a(n) appears to be given by the following conjectural formula: ordp(a(n),p) = b(floor(n/p)) + b(floor(n/p^2)) + b(floor(n/p^3)) + ... Compare with the comments in A129365. [added 13 May 2026: for a proof of this conjecture by Grok AI see the Bala link.] - _Peter Bala_, Apr 15 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Mar 31 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003182 | OEIS-A092193

Number of generations for which prime(n) divides A001008(k) for some k.

For any prime p, generation m consists of the numbers p^(m-1) <= k < p^m. The zeroth generation consists of just the number 0. When there is a k in generation m such that p divides A001008(k), then that k may generate solutions in generation m+1. It is conjectured that for all primes there are solutions for only a finite number of generations.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Feb 24 2004; corrected Jul 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003183 | OEIS-A092210

Positive integers n such that the trajectory of n under the Reverse and Add! operation carried out in base 2 (presumably) does not join the trajectory of any m < n.

The conjecture that the base 2 trajectories of the terms do not join is based on the observation that if the trajectories of two integers < 12000 join, this happens after at most 93 steps, while for any two terms listed above the trajectories do not join within 1000 steps. For pairs from 1, 16, 64, 74, 98, 107 this has even been checked for 5000 steps.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Feb 25 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003184 | OEIS-A092212

a(n) = smallest non-palindromic k such that the base-2 Reverse and Add! trajectory of k is palindrome-free and joins the trajectory of A092210(n).

Terms a(9) to a(29) are 205796147 (conjectured), 4402, 16720, 1089448, 442, 537, unknown, 1050177, 1575, 28822, unknown, 40573, 1066, 1587, unknown, unknown, 1081, 1082, 1085, 1115, 4185. The 1-to-1 property of the mapping depends on the conjecture that the base-2 Reverse and Add! trajectory of each term of A092210 contains only a finite number of palindromes (cf. A092215).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Feb 25 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003185 | OEIS-A092215

Smallest number whose base-2 Reverse and Add! trajectory (presumably) contains exactly n base-2 palindromes, or -1 if there is no such number.

Conjecture 1: For each k > 0 the trajectory of k eventually leads to a term in the trajectory of some j which belongs to A075252, i.e., whose trajectory (presumably) never leads to a palindrome. Conjecture 2: There is no k > 0 such that the trajectory of k contains more than eleven base 2 palindromes, i.e., a(n) = -1 for n > 11.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Feb 25 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003186 | OEIS-A092265

Sum of smallest parts of all partitions of n into distinct parts.

Conjecture: a(n) = A034296(n) + A237665(n). - _George Beck_, May 06 2017 The preceding conjecture holds (see Chern, 2017). - _Andrew R. Epstein_, Jul 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Feb 14 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003187 | OEIS-A092287

a(n) = Product_{j=1..n} Product_{k=1..n} gcd(j,k).

Conjecture: Let p be a prime and let ordp(n,p) denote the exponent of the highest power of p that divides n. For example, ordp(48,2)=4, since 48=3*(2^4). Then we conjecture that the prime factorization of a(n) is given by the formula: ordp(a(n),p) = (floor(n/p))^2 + (floor(n/p^2))^2 + (floor(n/p^3))^2 + ... Compare this to the de Polignac-Legendre formula for the prime factorization of n!: ordp(n!,p) = floor(n/p) + floor(n/p^2) + floor(n/p^3) + ... This suggests that a(n) can be considered as generalization of n!. See A129453 for the analog for a(n) of Pascal's triangle. See A129454 for the sequence defined as a triple product of gcd(i,j,k). - _Peter Bala_, Apr 16 2007 The conjecture is correct. - _Charles R Greathouse IV_, Apr 02 2013 Conjecture: For a product over a rectangle, f(n,m) = Product_{j=1..n} Product_{k=1..m} gcd(j,k), a factorization similar to the one given above for the square case takes place: ordp(f(n,m),p) = floor(n/p)*floor(m/p) + floor(n/p^2)*floor(m/p^2) + ... By way of directly computing the values of f(n,m), it can be verified that the conjecture holds at least for all 1 <= m <= n <= 200. - _Andrey Kaydalov_, Mar 11 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, based on a suggestion from _Leroy Quet_, Feb 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003188 | OEIS-A092291

Let p = n-th irregular prime, A000928(n). Then a(n) = smallest value of m such that numerator(Bernoulli(2*m)/(2*m)) / numerator(Bernoulli(2*m)/(2*m*(2*m-1))) equals p.

It was conjectured that a(n) = (1 + A000928(n) * (A035112(n) - 1))/2. However, Bernd Kellner's insightful paper shows that this formula first fails for the irregular prime 6449. - _T. D. Noe_, Feb 10 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, based on a suggestion of _Roland Bacher_, Feb 05 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003189 | OEIS-A092305

Length of period of sequence of Genocchi number of first kind read modulo (2n+1).

Conjecture : a(n)= (2n+1)*prod(p | 2n+1, (p-1)/2)=(2n+1)/(-2)^omega(2n+1)*sum(d | 2n+1, d*mu(d))

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 12 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003190 | OEIS-A092315

a(n) is the smallest m such that the partial sum of the odd harmonic series Sum_{j=0..m} 1/(2j+1) is > n.

a(n) = floor(exp(2*n-gamma)/4+1/8) for all n >= 1 (conjectured; see also comments in A002387). - _M. F. Hasler_, Jan 22 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 16 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003191 | OEIS-A092356

UO-sigma multiperfect numbers: n such that A069184(n)/n is an integer.

The initial values of k are 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2, 2. However, I conjecture that every positive integer >= 2 must appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yasutoshi Kohmoto_, Mar 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003192 | OEIS-A092363

n^(1/a(n)) is the closest to an integer on 2..n with a(n) minimal.

The sequence is conjectured to tend to n, as n^(1/n)->1. Is the density of nonnegative entries 0?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Mar 19 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003193 | OEIS-A092372

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by zero loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by zero loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Saibal Mitra (smitra(AT)zonnet.nl), Mar 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003194 | OEIS-A092373

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by one loop is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by one loop is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Saibal Mitra (smitra(AT)zonnet.nl), Mar 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003195 | OEIS-A092374

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by two loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by two loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number

of n X n half turn symmetric alternating sign matrices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Saibal Mitra (smitra(AT)zonnet.nl), Mar 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003196 | OEIS-A092375

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by three loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by three loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Saibal Mitra (smitra(AT)zonnet.nl), Mar 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003197 | OEIS-A092376

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by four loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by four loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Saibal Mitra (smitra(AT)zonnet.nl), Mar 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003198 | OEIS-A092377

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by five loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by five loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Saibal Mitra (smitra(AT)zonnet.nl), Mar 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003199 | OEIS-A092378

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by six loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by six loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Saibal Mitra (smitra(AT)zonnet.nl), Mar 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003200 | OEIS-A092379

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by seven loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by seven loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number

of n X n half turn symmetric alternating sign matrices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Saibal Mitra (smitra(AT)zonnet.nl), Mar 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003201 | OEIS-A092380

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by eight loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by eight loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Saibal Mitra (smitra(AT)zonnet.nl), Mar 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003202 | OEIS-A092381

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by nine loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by nine loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Saibal Mitra (smitra(AT)zonnet.nl), Mar 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003203 | OEIS-A092382

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by ten loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

The O(1) loop model on the square lattice is defined as follows: At every vertex the loop turns to the left or to the right with equal probability, unless the vertex has been visited before, in which case the loop leaves the vertex via the unused edge. Every vertex is visited twice. The probability that a face of the lattice on an n X infinity cylinder is surrounded by ten loops is conjectured to be given by a(n)/A_{HT}(n)^2, where A_{HT}(n) is the number of n X n half turn symmetric alternating sign matrices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Saibal Mitra (smitra(AT)zonnet.nl), Mar 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003204 | OEIS-A092387

a(n) = Fibonacci(2*n+1) + Fibonacci(2*n-1) + 2.

Consider a circle C of radius 5/4 (in some length units) with a chord of length 2. This has been chosen so that the larger sagitta also has length 2. The smaller sagitta has length 1/2. The input, besides the circle C, is the circle C_0 with radius R_0 = 1/4, touching the chord and circle C. The following sequence of circles C_n with radii R_n, n >= 1, is obtained from the conditions that C_n touches (i) the circle C, (ii) the chord and (iii) the circle C_(n-1). The curvature of the n-th circle, C_n = 1/R_n, n >= 0, is conjectured to be a(n). See an illustration given in the link. As found by _Wolfdieter Lang_ (see part II of the proof given by W. Lang in the link), this circle problem is related to the nonnegative solutions of the Pell equation X^2 - 5*Y^2 = 4: a(n) = 2 + X(n) = 2 + A005248(n). For the larger segment below the chord (with sagitta length 2) the sequence would be A115032, see W. Lang's proof given in part I of the link. Conjectures (proved in the next entry) from Colin Barker, Aug 25 2014 (and Aug 27 2014): (Start) The first of the _Colin Barker_ conjectures above is true because of the Binet-de Moivre formula for L(2*n) (see the Jul 24 2003 _Dennis P. Walsh_ comment on A005248). With phi = (1+sqrt(5))/2, use 1/phi = phi-1, phi^2 = phi+1, (phi-1)^2 = 2 - phi. His second conjecture (recurrence) with input a(-3) = 20, a(-2) = 9 and a(-1) = 5 (from the above given recurrence) leads to his g.f. with the expanded denominator. Thus both conjectures are true. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Mar 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003205 | OEIS-A092429

a(n) = n! * Sum_{i,j,k,l >= 0, i+j+k+l = n} 1/(i!*j!*k!*l!).

Conjecture: a(n) ~ 4^n/4!. - _Vaclav Kotesovec_, Mar 07 2014 Vaclav Kotesovec's conjecture is true: a(n) = 4^n/24 + O(4^n/sqrt(n)); hence a(n) ~ 4^n/4!. - _Xinjun Wang_, Jun 05 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Mar 22 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003206 | OEIS-A092434

Number of words X=x(1)x(2)x(3)...x(n) of length n in three digits {0,1,2} that are invariant under the mapping X -> Y, where y(i)=((AD)^(i-1))x(1) and where (AD) denotes the absolute difference (AD)x(i)=abs(x(i+1)-x(i)) (in other words, y(i) is the i-th element in the diagonal of leading entries in the table of absolute differences of {x(1), x(2),...,x(n)}).

It is conjectured that a(n)=(n+2)*2^((n-1) div 2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Mar 23 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003207 | OEIS-A092468

a(n) = Sum_{i+j+k=n, 0<=i<=j<=k<=n} (n+2k)!/(i! * j! * (3*k)!).

Conjecture: a(n+1)/a(n) tends to (73 + 6*sqrt(87))^(1/3)/3 + 13/(3*(73 + 6*sqrt(87))^(1/3)) + 7/3 = 4.875129794... - _Vaclav Kotesovec_, Oct 31 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Mar 25 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003208 | OEIS-A092482

Sequence contains no 3-term arithmetic progression, other than its initial terms 1, 2, 3.

For n > 2, a(n+2) = 1 + 2^floor(log_2(n)) + Sum_{k=1..n} (3^A007814(k) + 1)/2 = 1 + A053644(n) + A005836(n) (conjectured and checked up to n=512). The above conjecture is correct. - _Neal Patel_, Aug 23 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Apr 04 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003209 | OEIS-A092484

Expansion of Product_{m>=1} (1 + m^2*q^m).

Conjecture: log(a(n)) ~ sqrt(2*n) * (log(2*n) - 2). - _Vaclav Kotesovec_, Dec 27 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Apr 04 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003210 | OEIS-A092506

Prime numbers of the form 2^n + 1.

Conjecture: these are the only solutions to the equation A000010(x)+A000010(x-1)=floor((3x-2)/2). - _Benoit Cloitre_, Mar 02 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jorge Coveiro_, Apr 05 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003211 | OEIS-A092517

Product of tau values for consecutive integers.

a(n) is the maximum number of b > 0, which allows us to write (n+1)^2 as a sum of n+1 parts. Each part is of the form b^c and c is an integer >= 0 independent for each part. For n = 2 this is 3^2 = 2^2 + 2^2 + 2^0 = 3^1 + 3^1 + 3^1 = 4^1 + 4^1 + 4^0 = 7^1 + 7^0 + 7^0, b = 2;3;4;7 and a(2) = 4. It is conjectured that for all n the number of possible b reaches a(n). - _Thomas Scheuerle_, Jan 12 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 06 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003212 | OEIS-A092634

a(n) = 1 - Sum_{k=2..n} k*k!.

Conjecture: a(n) +(-n-3)*a(n-1) +(2*n+1)*a(n-2) +(-n+1)*a(n-3)=0. - _R. J. Mathar_, Sep 27 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | Doug Ray (draymath(AT)iastate.edu), Apr 11 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003213 | OEIS-A092635

Coefficients in asymptotic expansion of (Pi*n/16^n)*binomial(2n,n)^2 in powers of 1/(8n).

Conjecture from _Paul D. Hanna_, Aug 15 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Mar 02 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003214 | OEIS-A092674

Derived from a(n)=binomial(n+1,2) - sum{i=1,n-1,a(i)*floor(n/i)} (see A000010) - this is the value of the constant term.

It is conjectured that a(n) is never less than 0 (tested to n=2000)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Mar 02 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003215 | OEIS-A092687

First column and main diagonal of triangle A092686, in which the convolution of each row with {1,2} produces a triangle that, when flattened, equals the flattened form of A092686.

Conjecture: Limit n->infinity a(n)^(1/n) = 3. - _Vaclav Kotesovec_, Jun 29 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 04 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003216 | OEIS-A092749

a(n) is the least k such that m^2 + m + k is prime for m = 0..n.

An obvious question of course is whether for given n, a(n) exists at all. This is far from obvious. Assuming the generally believed k-tuplets conjecture, the answer is yes as was shown by Andrew Granville. For a proof (which is not very difficult) see the paper by Mollin. George Bright conjectured that a(n) exists for every n (private communication, 1974; see Dudley). - _Charles R Greathouse IV_, Sep 12 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | Gabriel Cunningham (gcasey(AT)mit.edu), Apr 12 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003217 | OEIS-A092765

Consider the 1-D random walk with jumps to next-nearest neighbors. Sequence gives number of paths of length n ending at origin.

Conjecture: the stronger congruences a(n*p^k) == a(n*p^(k-1)) (mod p^(2*k)) hold for any prime p, except p = 3, and positive integers n and k.(End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sergey Perepechko_, Apr 19 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003218 | OEIS-A092842

a(n) is the solution of the equation A073666(x) = n (A073666(a(n)) = n).

For each n number of solutions of the equation A073666(x) =n is less than 2 because the terms of the sequence A073666 are distinct. Conjecture: This sequence is

infinite(see A073666).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Apr 15 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003219 | OEIS-A092843

a(n) = Sum_{k|n, k>1} phi(k-1), where phi() is the Euler phi function.

Conjecture: Sum_{k=1..n} a(k) ~ n^2/2. - _Vaclav Kotesovec_, Jun 25 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Nov 09 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003220 | OEIS-A092878

Number of initial odd numbers in class n of the iterated phi function.

Class n, for n>0, contains all numbers k such that n iterations of the Euler phi function applied to k yields 2; class 0 contains only the numbers 1 and 2. There is a conjecture that the smallest number in class n is always odd. This increasing sequence supports that conjecture. As shown by Shapiro, all the initial odd numbers in class n>0 are between 2^n and 2^(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 10 2004, Nov 30 2007, Nov 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003221 | OEIS-A092905

Triangle, read by rows, such that the partial sums of the n-th row form the n-th diagonal, for n>=0, where each row begins with 1.

The triangle with entry a(n,m) = T(n-1,m-1), n >= 1, m = 1, ..., n, is obtained from the partition array A103921 when in row n all entries belonging to part number m are summed (a conjecture). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 12 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003222 | OEIS-A092925

Smallest prime of the form n(n-1)(n-2)...(n-k)-1, or 0 if no such prime exists.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 18 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003223 | OEIS-A092950

Beginning with n, add the next number, subtract the previous number and so on until one gets a prime, or 0 if no such prime is reached in 2n-1 steps: a(n) = n + (n+1) - (n-1) +(n+2) -(n-2) +(n+3)-(n-3)...+...is the first occurring prime at any step.

a(p) = p, p is a prime. If the process is continued until a 1 is subtracted the result is n^2. Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 24 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003224 | OEIS-A092951

Beginning with n, add the next number, subtract the previous number, and so on; then a(n) is the largest prime arising in this process, or 0 if no prime is reached in 2n-1 steps: a(n) is the largest occurring prime in the sum n + (n+1) - (n-1) + (n+2) - (n-2) + (n+3) - (n-3) + ... +- 1 occurring at any stage.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 24 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003225 | OEIS-A092959

Least square of the form 'product of n successive terms of an arithmetic progression + 1', or 0 if no such square exists.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 25 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003226 | OEIS-A092960

Least square of the form 'sum of n successive terms of a geometric progression + 1', or 0 if no such square exists.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 25 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003227 | OEIS-A092969

a(1) = 2; for n>1, a(n) = largest prime of the form n!/k + 1, where k < n, or 0 if no such prime exists.

Conjecture: There are only finitely many zeros in this sequence. In other words the sequence is identical to A092965 barring a finite set of terms which are zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 26 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003228 | OEIS-A092984

a(n) = the least k >= 1 such that n! + k is squarefree.

Conjecture: There exists a finite k such that a(n) < k for all n. Subsidiary sequence: Index of the first occurrence of n in this sequence. In case the conjecture is true, this sequence would be finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Mar 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003229 | OEIS-A093154

Primes resulting from serial multiplication of even composites, plus 1.

a(12) = 2^118*119!+1, a(13) = 2^142*143!+1. I conjecture that a(13) is the last prime number of this form. - _Jorge Coveiro_, Apr 01 2004 Conjecture that a(13) is the last prime of this form is false:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enoch Haga_, Mar 25 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003230 | OEIS-A093161

Even integers k such that there exists a prime p with p = min{q: q prime and (k - q) prime} and (k - p) < p^3.

It is conjectured that the sequence is finite with last term a(104820) = 5714500178 and it is proven that there are no more terms below 4*10^18. This is an extension of A307542. - _Corinna Regina Böger_, Apr 14 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, May 10 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003231 | OEIS-A093210

Row sums of A092964.

Conjecture: a(n) = A060477(n+4) - 2. - _Ralf Stephan_, Apr 24 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | Thomas O. Hoffbauer (Thomas.Hoffbauer(AT)cibamberg.de), Apr 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003232 | OEIS-A093245

a(n) is the lesser term of the smallest twin prime pair such that if P=(a(n)^2+n)^2+n, then P and P+2 are also twin primes. a(n) is 0 if no such pair exists.

Conjecture: a(n) = 0 only if n is a multiple of 3. Note that this implies the existence of infinitely many twin primes. - _Robert Israel_, Apr 15 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ray G. Opao_, May 11 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003233 | OEIS-A093328

a(n) = 2*n^2 + 3.

Conjecture: no perfect powers. - _Zak Seidov_, Sep 27 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Apr 25 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003234 | OEIS-A093343

Primes of form (prime(n)^2 + prime(n+1)^2)/2.

Conjecture: a(n) ~ A224888(n). - _Thomas Ordowski_, Jul 25 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Teofilatto_, Apr 26 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003235 | OEIS-A093345

a(n) = n! * {1 + Sum[i=1..n, 1/i*Sum(j=0..i-1, 1/j!)]}.

Conjecture: a(n) -2*n*a(n-1) +(n^2-2)*a(n-2) -(n-2)^2*a(n-3)=0. - _R. J. Mathar_, May 30 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Apr 26 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003236 | OEIS-A093348

A 5-fractal "castle" starting with 0.

Conjecture: a(n+1) = (n mod 2) + Sum_{k=0..infinity} (4*5^k*(floor(n/5^(k+1)) mod 2)). - _Charlie Neder_, May 25 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Apr 26 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003237 | OEIS-A093387

a(n) = 2^(n-1) - binomial(n, floor(n/2)).

a(2*n+1) = 2*a(2*n) + A000108(n). Together with the first formula by _Emeric Deutsch_, we have a simple system of recursions. Using them, we can prove Mathar's conjecture. For example, let n be odd, n=2*m+1. By the left hand side of Mathar's conjecture, we have (2*m+2)*a(2*m+1) - 2*(2*m+2)*a(2*m) - 4*(2*m-1)*a(2*m-1) + 8(2*m-1)*a(2*m-2) = (2*m+2)*(2*a(2*m) + A000108(m) - 2*a(2*m)) - 4*(2*m-1)*(2*a(2*m-2) + A000108(m-1) - 2*a(2*m-2)) = (2*m+2)*A000108(m) - 4*(2*m-1)*A000108(m-1) = 0, since A000108(m) = binomial(2*m, m)/(m+1). - _Vladimir Shevelev_, Nov 17 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthijs Coster_, Apr 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003238 | OEIS-A093392

[n/25] + [(n+1)/25] + [(n+2)/25] + [(n+3)/25] + [(n+4)/25].

Conjectured g.f. confirmed with more terms similar to A093390, A093391, A093393.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Mar 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003239 | OEIS-A093483

a(1) = 2; for n>1, a(n) = smallest integer > a(n-1) such that a(n) + a(i) + 1 is prime for all 1 <= i <= n-1.

The Hardy-Littlewood k-tuple conjecture would imply that this sequence is infinite. Note that, for n > 2, a(n)+3 and a(n)+5 are both primes, so a proof that this sequence is infinite would also show that there are infinitely many twin primes. - _N. J. A. Sloane_, Apr 21 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 14 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003240 | OEIS-A093489

Define the total signature symmetry of a number n to be the number of values r takes such that n-r and n+r have the same prime signature. Sequence contains the total signature symmetry of n.

Number of partitions of 2n in two parts with identical prime signatures. Conjecture: (1) No term is zero for n > 3. (2) Every number k appears a finitely many times in the sequence. I.e., for every k there exists a number f(k) such that for all n > f(k), a(n) > k. Subsidiary sequence: the frequency of n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 16 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003241 | OEIS-A093492

Define the divisor symmetry of a number n to be k if n-r and n+r have the same number of divisors for r = 1 to k but not for k+1. Sequence contains the divisor symmetry of n.

I conjecture that the sequence is unbounded. [_Charles R Greathouse IV_, Dec 19 2011]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 16 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003242 | OEIS-A093493

Define the total divisor symmetry of a number n to be the number of values r takes such that n-r and n+r have the same number of divisors. Sequence contains the total divisor symmetry of n.

Number of partitions of 2n in two parts with equal number divisors. Conjecture: (1) No term is zero for n > 3. (2) Every number k appears finitely many times in the sequence. i.e. for every k there exists a number f(k) so that for all n > f(k), a(n) > k. Subsidiary sequences: (1) The frequency of n. (2) The greatest number m so that a(m) = n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 16 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003243 | OEIS-A093504

A093503(k) is a term if A093503(k) and A093503(k+1) are successive primes.

There are no further terms < A093503(1000000); this suggests the conjecture that there are no other terms. - Klaus Brockhaus

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 17 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003244 | OEIS-A093505

a(n) = floor(A001969(n)/2 + 1/2).

Conjecture: Essentially partial sums of A036577 (true for at least 10000 terms). - _Sean A. Irvine_, Jul 17 2022 Proof of the conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Joseph Halm_, May 13 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003245 | OEIS-A093511

Transform of the prime sequence by the Rule45 cellular automaton.

Conjecture: For n > 3, the a(n) correspond to the following construct (in numerical order). a(n) terms include "bookend" values at every prime p + 1 (6,8,12,14,18,20,...). Additionally, the values between the bookends are included, unless adjacent to non-"twin composite" bookends. For example, consider bookends 6 and 8. There is only a single value 7 between these, so it is included. This means terms 6, 7 and 8 are included. Consider bookends 89 + 1, 97 + 1. Ignoring 91 and 97 adjacencies, values 92 through 96 are included. This means terms 90, 92-96 and 98 are included. - _Bill McEachen_, Jun 12 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ferenc Adorjan (fadorjan(AT)freemail.hu) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003246 | OEIS-A093525

Decimal expansion of 13/720 - Pi^2/15015.

Klee (1969) conjectured that the average volume is 1/60 and stated that according to Monte Carlo experiments 1/57 is the integer-reciprocal closest to this value. - _Amiram Eldar_, Apr 09 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Mar 30 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003247 | OEIS-A093538

Number of primes between 10^n and 10^n+100.

Conjecture: a(n) = 1 for infinitely many n, but a(n)>=2 for only finitely many; in fact it may be that a(n)<=1 for n > 775. - _Robert Israel_, Feb 28 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, May 14 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003248 | OEIS-A093585

Numerators of even raw moments in the distribution of a triangle picked at random from points on the circumference of a unit circle.

Conjecture: the moments are sqrt(3)*Gamma(n+2/3)*Gamma(n+1/3)*(27/16)^n/(2*Pi*(n!)^2). - _Robert Israel_, Jan 01 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Apr 01 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003249 | OEIS-A093600

Numerator of Sum_{1<=k<=n, gcd(k,n)=1} 1/k.

Problem: are there numbers n > 1 such that n^4 | a(n)? Let b(n) be the numerator of Sum_{1<=k<=n, gcd(k,n)=1} 1/k^2. Conjecture: if, for some e > 0, n^e | a(n), then n^(e-1) | b(n). It appears that, for any odd number n, n^e | a(n) if and only if n^(e-1) | b(n). - _Thomas Ordowski_, Aug 12 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003250 | OEIS-A093652

Let a(1) = 1, a(2) = 2, a(3) = 7, a(4) = 15 and for n >= 5 set a(n) = (n*b(n) - b(n-2)) / 2, where b(n) = 4*b(n-2) - b(n-4) for n >= 5 and b(1) = 1, b(2) = 2, b(3) = 5, b(4) = 8.

Conjecture: b(n) = A082630(n). If true, we can write a(n) = (n*A082630(n) - A082630(n-2)) / 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Harri Aaltonen_, May 15 2004, Apr 12 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003251 | OEIS-A093681

Sequence contains no 3-term arithmetic progression, starting with 1, 28.

a(n) = (Sum_{k=1..n-1} (3^A007814(k) + 1)/2) + f(n), with f(n) a 16-periodic function with values {1, 27, 26, 27, 23, 27, 26, 27, 14, 28, 28, 28, 28, 28, 28, 28, ...}, n >= 1 (conjectured and checked up to n=1000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Apr 09 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003252 | OEIS-A093682

Array T(m,n) by antidiagonals: nonarithmetic-3-progression sequences with simple closed forms.

T(m, n) = (Sum_{k=1..n-1} (3^A007814(k) + 1)/2) + f(n), with f(n) a P-periodic function, where P <= 2^floor((m+3)/2) (conjectured and checked up to m=13, n=1000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Apr 09 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003253 | OEIS-A093695

Number of one-element transitions among partitions of the integer n for unlabeled parts.

G.f.: 2*x^2 / (x^3-x^2-x+1) * Product_{m>=1} (1/(1-x^m)) (conjectured). - _Thomas Baruchel_, May 12 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Wieder_, Apr 10 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003254 | OEIS-A093714

a(n) = smallest number coprime to a(n-1), not equal to a(n-1)+1, and not occurring earlier; a(1)=1.

Conjecture 1: If p is a prime >= 3, p-1 appears after p. Conjecture 2: If p is a prime, the first term divisible by p is p itself. Conjecture 3: If a(n) = p is a prime >= 5, then n < p. Conjecture 1 is false. The smallest counterexample is p = 53: a(49) = 52 and a(52) = 53, so p-1 appears before p. - _Clément Garrot_, Sep 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 12 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003255 | OEIS-A093726

Given the infinite continued fraction (1+i)+1/((1+i)+1/((1+i)+...)), where i is the square root of (-1), this is the numerator of the imaginary part of the convergents.

Conjecture: a(n)/A093727(n) = A317974(n+1)/A138573(n). - _Zhuorui He_, Jun 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Mar 11 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003256 | OEIS-A093811

Sum of the digital products of the divisors of n.

We conjecture that a(n) < n for all n > 798, and that n = 10^k - 4 and n = 10^k - 2 set new record values of a(n) ~ 94*9^(k-2) resp. 9^k*4/3 for all k (resp. k > 4), hence lim sup a(n)/n = 0. - _M. F. Hasler_, Jul 11 2026 Conjecture: lim sup_{n -> oo} a(n)/9^log_10(n) = lim_{k -> oo} a(10^k - 2)/9^k = 4/3. - _M. F. Hasler_, Jul 11 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, May 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003257 | OEIS-A093875

Denominators in Kepler's tree of harmonic fractions.

Conjecture of the comment in detail: a(2n+1) = a(2n), n > 0; a(2n+1) = A071585(n), n >= 0; a(2n) = A071585(n), n > 0. - _Yosu Yurramendi_, Jun 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Reinhard Zumkeller_, May 24 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003258 | OEIS-A093892

Index of first occurrence of n in A093890, or 0 if no such number exists.

Conjecture: there are infinitely many zeros in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 23 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003259 | OEIS-A093894

Composite members of A093893.

Comment: Most terms of this sequence have four divisors. Some terms (the squares of primes) have three divisors; very few terms have more than four divisors (the first such term is 4753, with six). Conjecture: This sequence is infinite. - Adam M. Kalman (mocha(AT)clarityconnect.com), Nov 11 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 23 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003260 | OEIS-A093954

Decimal expansion of Pi/(2*sqrt(2)).

We conjecture the asymptotic expansion Pi/(2*sqrt(2)) - Sum_{k = 0..n-1} (-1)^floor(k/2)/(2*k + 1) ~ 1/(2*n) - 3/(2*n)^3 + 57/(2*n)^5 - 2763/(2*n)^7 + ..., where n is a multiple of 4 and the sequence of unsigned coefficients [1, 3, 57, 2763, ...] is A000281. An example with n = 5000 is given below. (End) c = 2 * Sum_{k >= 0} (-1)^k * (4*k + 2)/((4*k + 1)*(4*k + 3)) = A181048 + A181049. The asymptotic expansion conjectured above follows from the asymptotic expansions given in A181048 and A181049.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Apr 19 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003261 | OEIS-A093963

Antidiagonal sums of array in A093966.

Conjecture: 2*a(n) -5*a(n-1) -(n+2)*a(n-2) +2*(n+6)*a(n-3) +(n-13)*a(n-4) -4*(n-3)*a(n-5) +2*(n-3)*a(n-6) = 0. - _R. J. Mathar_, Nov 10 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Apr 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003262 | OEIS-A093964

a(n) = Sum_{k=1..n} k*k!*C(n,k).

G.f.: conjecture: T(0)/x - 1/x, where T(k) = 1 - x^2*(k+1)^2/(x^2*(k+1)^2 - (1 - 2*x*(k+1))*(1 - 2*x*(k+2))/T(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 18 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Apr 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003263 | OEIS-A094043

Alternate composite and prime numbers not included earlier such that every partial concatenation is a prime: a(2n) is prime and a(2n-1) is not prime.

Conjecture: 2 and 5 are the only two nonmembers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Apr 23 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003264 | OEIS-A094044

Alternate prime and nonprime numbers not included earlier such that every concatenation of a pair of terms is a prime: a(2n) is nonprime and a(2n-1) is prime.

Conjecture: all members of A045572 are in the sequence. - _Robert Israel_, Oct 24 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Apr 23 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003265 | OEIS-A094045

Alternate composite and prime numbers not included earlier such that every concatenation of a pair of terms is a prime: a(2n) is prime and a(2n-1) is nonprime.

Conjecture: 2 and 5 are the only two nonmembers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Apr 23 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003266 | OEIS-A094047

Number of seating arrangements of n couples around a round table (up to rotations) so that each person sits between two people of the opposite sex and no couple is seated together.

Conjecture: a(n) +(-n^2+2*n-3)*a(n-1) -(n-2)*(n^2-3*n+5)*a(n-2) -3*(n-2)*(n-3)*a(n-3) +(n-2)*(n-3)*(n-4)*a(n-4)=0. - _R. J. Mathar_, Nov 02 2015 Conjecture: (-n+2)*a(n) +(n-1)*(n^2-3*n+3)*a(n-1) +(n-2)*(n-1)*(n^2-3*n+3)*a(n-2) +(n-2)*(n-3)*(n-1)^2*a(n-3)=0. - _R. J. Mathar_, Nov 02 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthijs Coster_, Apr 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003267 | OEIS-A094061

Number of n-moves paths of a king starting and ending at the origin of an infinite chessboard.

Conjecture: the stronger congruences a(n*p^k) == a(n*p^(k-1)) (mod p^(2*k)) hold for all primes p >= 5 and positive integers n and k. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthijs Coster_, Apr 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003268 | OEIS-A094076

Smallest k such that prime(n) + 2^k is prime, or -1 if no such prime exists.

Conjecture: k > 0 for all n. For all primes p < 1000 there exists a k such that p + 2^k is prime. However, for p = prime(321) = 2131, p + 2^k is not prime for all k < 30000. The conjecture may be in question. Similarly, I cannot find k such that p + 2^k is prime for p = 7013, 8543, 10711, 14033 for k < 20000. - _Cino Hilliard_, Jun 27 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003269 | OEIS-A094087

Domination number of the Cartesian product of two n-cycles.

1/5 <= a(n)/n^2 <= 1/4 for n >= 4; it is conjectured that a(5n-1) = 5*n^2 - n and a(5n+1) = 5n^2 + 4n - 1 for n >= 1. - _Richard Bean_, Sep 08 2006 [Assadian proves that the both conjectured formulas give the upper bounds. - _Andrey Zabolotskiy_, Dec 23 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Bean_, May 01 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003270 | OEIS-A094187

Numerator of I(n) = Integral_{x=1..9/8} (sqrt(x^2-1)/x)^(2*n) dx.

Conjecture D-finite with recurrence a(n) + (-196*n+51)*a(n-1) + 2754*(n-1)*(2*n-3)*a(n-2) = 0. - _R. J. Mathar_, Feb 04 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | Al Hakanson (hawkuu(AT)excite.com), May 24 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003271 | OEIS-A094189

Number of primes between n^2-n and n^2 (inclusive).

Conjecture: for n>11, a(n)>1. Oppermann conjectured in 1882 that a(n)>0 for n>1. - _T. D. Noe_, Sep 16 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, May 25 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003272 | OEIS-A094213

a(n) = Sum_{k=0..n} binomial(9*n,9*k).

Conjectures from _Colin Barker_, May 27 2019: (Start) a(n) ~ (1/9)*exp(n*9*log(2)) (conjecture). - _Bill McEachen_, Aug 11 2025 [Proved in Easwar (2026). - _Rohun Easwar_, Aug 31 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 27 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003273 | OEIS-A094262

Triangle read by rows: T(n,k) is the number of rooted trees with k nodes which are disjoint sets of labels with union {1..n}. If a node has an empty set of labels then it must have at least two children.

Consequently, R(n+1,z) = (1 + z)^2*z*d/dz(R(n,z)) for n >= 1 as conjectured above by Copeland.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _André F. Labossière_, Jun 01 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003274 | OEIS-A094287

Number of (s(0), s(1), ..., s(n)) such that 0 < s(i) < 7 and |s(i) - s(i-1)| <= 1 for i = 1,2,...,n, s(0) = 1, s(n) = 1.

Conjecture: a(n)= +6*a(n-1) -10*a(n-2) +9*a(n-4) -2*a(n-5) -a(n-6) with g.f. -x*(-1+4*x-2*x^2-5*x^3+2*x^4+x^5) / ((x^3+3*x^2-4*x+1)*(x^3-x^2-2*x+1)). - _R. J. Mathar_, Dec 20 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Herbert Kociemba_, Jun 02 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003275 | OEIS-A094288

Number of (s(0), s(1), ..., s(n)) such that 0 < s(i) < 8 and |s(i) - s(i-1)| <= 1 for i = 1,2,...,n, s(0) = 1, s(n) = 1.

Conjecture: a(n)= +7*a(n-1) -15*a(n-2) +5*a(n-3) +15*a(n-4) -9*a(n-5) -3*a(n-6) +a(n-7) with g.f. -x*(1-5*x+5*x^2+6*x^3-7*x^4-2*x^5+x^6) / ((x-1)*(x^2+2*x-1)*(x^4-4*x^3-2*x^2+4*x-1)). - _R. J. Mathar_, Dec 20 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Herbert Kociemba_, Jun 02 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003276 | OEIS-A094291

a(n) = maximal value of C(i, j) * C(n-j, n-i) for 0 <= j <= i <= n.

I conjecture that the maximum value is always reached for i = n-k where k=j or k=j+1, cf. FORMULA. The j-values are as follows, where * marks those where k=j+1: j(n=1,2,3...) = (0*, 0*, 1, 1, 1*, 1*, 2, 2, 2*, 3, 3, 3*, 3*, 4, 4, 4*, 4*, 5, 5, 5*, 6, 6, ...). Indices of the starred values appear to be the disjoint union of A094350 and A081009 (for which C(n-j, j)^2 gives the same value). - _M. F. Hasler_, Sep 02 2026 Conjecture: a(n) = C(n-k(n), j(n))*C(n-j(n), k(n)) where [n/4] <= j(n) <= k(n) <= j(n)+1; the j (resp. k) value can increase only when j = k-1 (resp. j = k). - _M. F. Hasler_, Sep 02 2026 Conjecture: Limit_{n->oo} a(n)^(1/n) = (3 + sqrt(5))/2. - _Vaclav Kotesovec_, Sep 03 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo van der Sanden_, Jun 15 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003277 | OEIS-A094294

a(n) = n*a(n-1) - n + 2 for n > 1; a(1)=1.

Conjecture: (-n+3)*a(n) + (n^2-2*n-2)*a(n-1) - (n-1)*(n-2)*a(n-2) = 0. - _R. J. Mathar_, Sep 10 2015 Conjecture confirmed using Maple's gfun package. - _Robert Israel_, Oct 07 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003278 | OEIS-A094303

a(1) = 1, a(2) = 2, and a(n+1) = a(n) * sum of all previous terms up to a(n-1) for n >= 2.

R. J. Mathar's conjecture is correct and this is identical to A064847 starting at n = 3. To see why this is the case, consider the sequences u(n) and v(n) defined by u(1) = v(1) = 1, and u(k+1) = u(k) + v(k), v(k+1) = u(k)*v(k) for k >= 1. Then u(n) = A003686(n) and v(n) = A064847(n) for n >= 1. Conjecture: a(n) = A003686(n+1) - A003686(n) for n >= 3. - _R. J. Mathar_, Apr 24 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Apr 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003279 | OEIS-A094336

Product of next n numbers divided by the largest factorial divisor.

Conjecture: a(n) > 1 for n > 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 17 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003280 | OEIS-A094339

Beginning with 2, least number not occurring earlier that divides the sum of all previous terms.

Conjecture: this is a rearrangement of natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 17 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003281 | OEIS-A094340

a(n) = n-th partial sum of A094339 divided by A094339(n+1).

Conjecture: Every natural number occurs in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 17 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003282 | OEIS-A094358

Squarefree products of factors of Fermat numbers (A023394).

Conjectured (by Munafo, see link) to be the same as: numbers n such that 2^^n == 1 mod n, where 2^^n is A014221(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Munafo_, Apr 26 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003283 | OEIS-A094373

Expansion of (1-x-x^2)/((1-x)*(1-2*x)).

Take A007843 and count the repeated values. The result is 1,1,2,1,3,1,2,1,4,1,2,1,3,1,2,1,5,.... Build a third sequence, where a(1) = 1 and a(n) equals the length (greater than 1) of the shortest palindromic subsequence of consecutive terms of the second sequence starting with a(n) of the second sequence. The third sequence starts 1,3,5,3,9,3,5,3,17,3,5,3,9,3,5,3,33,.... Conjecturally, in the third sequence: (1) the indices of the first occurrence of each value form the present sequence and (2) for n>1, a(n) is in the a(n-1)-th position. - _Ivan N. Ianakiev_, Aug 20 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003284 | OEIS-A094383

Primes p such that d>0 exists and p-d, p-2*d and p-3*d are also primes.

Conjecture: only 25 primes are not in the sequence, namely 2, 3, 5, 7, 11, 13, 17, 19, 31, 37, 47, 61, 67, 71, 73, 89, 109, 137, 179, 211, 277, 337, 379, 499, 557. - _Alex Ratushnyak_, Sep 08 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003285 | OEIS-A094384

Determinant of n X n partial Hadamard matrix with coefficient m(i,j) 1<=i,j<=n (see comment).

Conjecture: a(n) = (-2)^A000788(n-1). - _Chai Wah Wu_, Nov 12 2024 The comment above regarding the signs is false but the conjecture is true (see Fried link). - _Sela Fried_, Feb 05 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jun 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003286 | OEIS-A094470

Numbers k such that phi(k) | sigma_18(k).

sigma_18(n) is the sum of the 18th powers of the divisors of n (A013966). Conjecture: analogous sequences with any 4j+2 exponent instead of 18, includes all terms of A015759 [with exponent=2].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, May 25 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003287 | OEIS-A094504

T(n,m) equals number of solid partitions of n containing m plane partitions.

Finding a G.f. for the solid partitions is an open problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_, Jun 05 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003288 | OEIS-A094508

Triangle read by rows: T[n,m] = number of solid partitions of n with trace m, where the trace of a solid partitions is defined as the sum of the traces of the constituent plane partitions.

Finding a GF for the solid partitions is an open problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_, Jun 05 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003289 | OEIS-A094540

Last digit of the n-th perfect number.

Conjecture: Ratio of 6's to 8's approaches 1.5 as a limit as this sequence approaches infinity. - _J. Lowell_, Jun 26 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, May 08 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003290 | OEIS-A094591

a(0) = 1; a(n) = n + (largest element of {a} <= n).

a(n) = A037988(n-1) + 1. [Conjectured by the original author, apparently proved by Eric Angelini.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Jun 07 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003291 | OEIS-A094603

a(n) is the length of the maximal sequence of rightmost black cells in the n-th row of Rule 30 (begun from an initial black cell).

Conjecture: a(n) = A094604(A007814(n)). - _Alan Michael Gómez Calderón_, Feb 17 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Rowland_, May 13 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003292 | OEIS-A094639

Partial sums of squares of Catalan numbers (A000108).

Conjecture: For any positive integer n, the polynomial P_n(x) = sum_{k = 0}^n(C_k)^2*x^k (with C_k = binomial(2k, k)/(k+1)) is irreducible over the field of rational numbers. [_Zhi-Wei Sun_, Mar 23 2013]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _André F. Labossière_, May 27 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003293 | OEIS-A094739

Numbers m such that 4^k*m, for integer k >= 0, are numbers having a unique partition into three squares.

Lehmer's paper has an erroneous version of this sequence. He omits 163 and includes 162 (which has 4 partitions) and 182 (which has 3 partitions). Lehmer conjectures that there are no more terms. Note that squares are allowed to be zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, May 24 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003294 | OEIS-A094740

Numbers n such that 4^k n, for k >= 0, are numbers having a unique partition into three positive squares.

It is conjectured that this sequence is complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, May 24 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003295 | OEIS-A094750

Rearrangement of composite numbers so that the successive differences are distinct prime numbers. Priority is given to the least successive difference (prime) that has not occurred earlier.

First priority is given to minimize the absolute value |a(n+1)-a(n)| of the difference (a prime), second priority to minimize a(n+1). The term "rearrangement" is conjectural as it remains to be proved that all positive composite numbers A002808 are in the list. - _R. J. Mathar_, Aug 03 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 24 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003296 | OEIS-A094757

Least positive k <= n such that n*pi(k) = k*pi(n), where pi(n) is the number of primes <= n (A000720).

Conjecture: For every n there exists a k different from n (possibly k > n) such that n*pi(k) = k*pi(n). _Amarnath Murthy_'s conjecture above is false. For n = 3 we have pi(n)/n = 2/3. For no other k we have pi(k)/k = 2/3. Therefore the conjecture is false. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 30 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003297 | OEIS-A094759

Least k <= n such that n*sigma(k) = k*sigma(n), where sigma(n) is the sum of divisors of n (A000203).

Conjecture: There are infinitely many terms such that a(n)<n. A050973 has those n, A050972 has the a(n). That conjecture is an easy fact: Since, e.g., (6,28) is a friendly pair, then so is (6k,28k) for any multiplier k with gcd(42,k)=1. So any n=28k, where gcd(42,k)=1, satisfies a(n)<n. This even shows that A095301 does not have asymptotic density zero. - _Jeppe Stig Nielsen_, Jul 09 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, May 30 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003298 | OEIS-A094776

a(n) = largest k such that the decimal representation of 2^k does not contain the digit n.

These values are only conjectural. The sequence could be extended to any nonnegative integer index n defining a(n) to be the largest k such that n does not appear as substring in the decimal expansion of 2^k. I conjecture that for n = 10, 11, 12, ... it continues (2000, 3020, 1942, 1465, 1859, 2507, 1950, 1849, 1850, ...). For example, curiously enough, the largest power of 2 in which the string "10" does not appear seems to be 2^2000. - _M. F. Hasler_, Feb 10 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Taktikos_, Jun 09 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003299 | OEIS-A094777

Number of legal positions in Go played on an n X n grid (each group must have at least one liberty).

Tromp & Farnebäck prove that a(n) = (1 + o(1)) * L^(n^2), and conjecture that a(n) ~ A * B^(2n) * L^(n^2) * (1 + O(n*p^n)) for some constants A, B, L, and p < 1. - _Charles R Greathouse IV_, Feb 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jan Kristian Haugland_, Jun 09 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003300 | OEIS-A094783

Numbers k such that, for all m < k, d_i(k) <= d_i(m) for i=1 to Min(d(k),d(m)), where d_i(k) denotes the i-th smallest divisor of k.

Conjecture: If k is in this sequence, then so is the smallest number with k divisors. (This conjecture is definitely false for A002182 (k=840) and A019505 (k=240).) - _J. Lowell_, Jan 24 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthew Vandermast_, Jun 10 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003301 | OEIS-A094787

a(n) = smallest prime p such that p + n is a perfect power m^k, k >= 2.

Conjecture: every prime is contained in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jun 10 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003302 | OEIS-A094802

a(n) = smallest k such that all of 1 through n divides k!.

It is conjectured that after n=4 the sequence is prime for n prime or the previous prime for n not prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Jun 11 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003303 | OEIS-A094822

Expansion of e.g.f.: exp(3*x)/(1-3*x)^(1/3).

Conjecture: D-finite with recurrence: a(n) -(3*n+1)*a(n-1) +9*(n-1)*a(n-2) = 0. - _R. J. Mathar_, Nov 14 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Jun 12 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003304 | OEIS-A094863

Maximal number of longest common subsequences between any two strings of length n (Version 2).

For even n it seems that the maximum is attained for X = 123412341234..., Y = 432143214321..., giving values : (conjectured) maximum number of maximum-length-common-subsequences of 2 strings of length 2*n over an arbitrary (infinite) alphabet f(2*n) = 2,4,10,28,78,220,624,1780,5100,14668,.. Note that (3*f(2*n)-f(2*n+2))/2 gives 1,1,1,3,7,18,46,120,316,841,2257,6103,16611,45475,125139,.. which is A026107. Is there an explanation for this?

猜想/问题提及段 | historical_or_current_mention_unreviewed | Guenter Stertenbrink (Sterten(AT)aol.com), Jun 14 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003305 | OEIS-A094869

E.g.f.: exp(5x)/(1-5x)^(1/5).

Conjectured to be D-finite with recurrence: a(n) +(-5*n-1)*a(n-1) +25*(n-1)*a(n-2)=0. - _R. J. Mathar_, Nov 15 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Jun 16 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003306 | OEIS-A094870

a(1)=1; for n > 1, a(n) is the minimal positive integer t not equal to a(1), ..., a(n-1) such that t - a(n-i) is not equal to a(n-i) - a(n-2i) for all 1 <= i < n/2.

Conjecture: lim_{n->infinity} a(n)/n = 1 (P. Hegarty).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Alec Mihailovs (alec(AT)mihailovs.com), Jun 16 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003307 | OEIS-A094905

Expansion of e.g.f.: exp(6*x)/(1-6*x)^(1/6).

Conjectured to be D-finite with recurrence: a(n) +(-6*n-1)*a(n-1) +36*(n-1)*a(n-2) = 0. - _R. J. Mathar_, Nov 15 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Jun 16 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003308 | OEIS-A094911

Expansion of e.g.f. exp(7*x)/(1-7*x)^(1/7).

Conjectured to be D-finite with recurrence: a(n) +(-7*n-1)*a(n-1) +49*(n-1)*a(n-2)=0. - _R. J. Mathar_, Nov 15 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Delélham_, Jun 17 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003309 | OEIS-A094913

Maximal number of distinct nonempty substrings of any binary string of length n.

It would be more natural to include the empty substring, which would result in the sequence b(n)=a(n)+1; all values computed so far confirm the conjecture that a(n)+1 = A006697(n). - _M. F. Hasler_, Dec 17 2007 Calculations suggest that, for any n>0, binary strings of length n exist such that the number of distinct substrings of length k, 1<=k<=n, is as large as possible consistent with basic counting principles, i.e., n-k+1 substrings of length k starting at each of the n-k+1 possible starting positions, subject to the condition, however, that there cannot be more than 2^k distinct binary strings of length k. This suggests the following conjecture. Conjecture: The maximum number a(n) of distinct substrings of a binary string of length n is given by a(n) = Sum_{k=1..n} t(k) where t(k) is the smaller of 2^k and n-k+1. Conjecture: a(n) = A006697(n)-1. - _Vladeta Jovovic_, Sep 19 2005 Conjecture: a(n) = 2^(m+1)-2 + (n-m)(n-m+1)/2, where m = floor(n+1-LambertW(2^(n+1) * log(2)) / log(2)) = integer part of the solution to 2^x = n+1-x. This is based on the reasoning that you can construct the word of length n so that it contains the maximal possible number, max(n-k+1, 2^k), of different substrings of length k. - _M. F. Hasler_, Dec 17 2007 Conjecture: column sums of:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jun 17 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003310 | OEIS-A094925

A hexagonal spiral Fibonacci sequence.

a(n) ~ c*phi^n with phi=1.61803... being the golden ratio and c = 0.78529667298898361017570049509486675274402985275383398273772345738007479334754... (conjectured). Cf. A094926. - _Manfred Scheucher_, Jun 03 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yasutoshi Kohmoto_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003311 | OEIS-A094960

Positive integers k such that the derivative of the k-th Bernoulli polynomial B(k,x) contains only integer coefficients.

Conjecture: for every integer k > 78, there exists a prime p >= 3 such that the sum of last two base-p digits of k is at least p. This conjecture would imply that this sequence is finite and 60 is the last term. (End) The conjecture is true for all k such that k+1 is not a prime, a power of 2, or a Giuga number (A007850). In this case, there exists a prime p >= 3 such that the base-p representation of k ends in a,p-1 with a > 0. - _Max Alekseyev_, Feb 16 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jun 19 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003312 | OEIS-A095017

Number of lesser twin primes (A001359) in range]2^n, 2^(n+1)].

Conjecture: a(n) > 0 for all n. This holds for all n <= 100. - _Charles R Greathouse IV_, May 14 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_ and _Labos Elemer_, Jun 01 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003313 | OEIS-A095029

An example of a (v,k,lambda)=(21,5,1) cyclic difference set.

A (v,k,lambda) cyclic difference set is a subset D={d_1,d_2,...,d_k} of the integers modulo v such that {1,2,...,v-1} can each be represented as a difference (d_i-d_j) modulo v in exactly lambda different ways. Difference sets with lambda=1 (planar difference sets) have order n=k-1. The Prime Power Conjecture states that all Abelian planar difference sets have order n a prime power. It is known that no cyclic planar difference sets of nonprime power order n exist with n < 2*10^9 (see Baumert, Gordon link)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, May 27 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003314 | OEIS-A095076

Parity of 1-fibits in Zeckendorf expansion A014417(n).

Let u = A000201 = (lower Wythoff sequence) and v = A001950 = (upper Wythoff sequence). Conjecture: This sequence is the sequence p given by p(1) = 0 and p(u(k)) = p(k); p(v(k)) = 1-p(k). - _Clark Kimberling_, Apr 15 2011 Proof of Kimberling's conjecture, by virtue of the four symbol morphism sigma in the previous comment. If we combine CLAIM 1 and CLAIM 2 with lambda(1) = lambda(3) = 0, we obtain the first half of Kimberling's conjecture, simply because delta = mu rho, and rho(R) = R. The second half of the conjecture is obtained in a similar way. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jun 01 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003315 | OEIS-A095116

a(n) = prime(n) + n - 1.

On March 23 2024, _Davide Rotondo_ sent me an email with the following conjecture. (I've simplified it a bit.) The conjecture is that after some initial terms, b becomes periodic with period length 1 or 2, and the n for which the period is 2 are 3 together with the present sequence, that is, 2, 3, 4, 7, 10, 15, 18, 23, 26, 31, 38, 41, 48, ... (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, following a suggestion of _Leroy Quet_, May 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003316 | OEIS-A095121

Expansion of (1-x+2x^2)/((1-x)*(1-2x)).

Conjecture: a(n) is the least m>0 such that A007814(A000108(m)) = n, where A000108 gives the Catalan numbers and A007814(x) is the 2-adic valuation of x (cf. A048881). - _L. Edson Jeffery_, Nov 26 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003317 | OEIS-A095149

Triangle read by rows: Aitken's array (A011971) but with a leading diagonal before it given by the Bell numbers (A000110), 1, 1, 2, 5, 15, 52, ...

Conjecture: Also the number of divisors d with distinct prime multiplicities of the superprimorial A006939(n) that are of the form d = m * 2^k where m is odd. For example, row n = 4 counts the following divisors:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, May 30 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003318 | OEIS-A095180

Reverse digits of primes, append to sequence if result is a prime.

Conjecture: the Benford law limit is 2=Sum[N[Log[10, 1 + 1/d[[n]]]], {n, 1, Length[d]}]^2/((#totalprimes/#totalPrimes)). At 50000 primes total it is 2.05931. - _Roger L. Bagula_ and _Gary W. Adamson_, Jul 02 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jun 21 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003319 | OEIS-A095188

Smallest prime formed by the digit string after decimal point of n^(1/2), or 0 if no such prime exists.

Conjecture: a(n) is nonzero if n is not a perfect square.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jun 02 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003320 | OEIS-A095189

Smallest prime formed by the digit string after decimal point of n^(1/n), or 0 if no such prime exists.

Conjecture: a(n) is nonzero for all n>1. Generates surprisingly large primes that are easily certified using Elliptic curve techniques (Mathematica's NumberTheory`PrimeQ`). For n=24 no certifiable prime was found with fewer than 1024 digits. - _Wouter Meeussen_, Jun 04 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jun 02 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003321 | OEIS-A095205

Least nonnegative integer not already in the sequence such that the frequency of digits in the entire sequence is as close to a uniform distribution as possible. A difference in frequency of at most 1 across all digits is allowed, and that difference must always be in favor of digits closer to 0.

Obviously no term contains identical digits and eventually for some m, a(m) = 1023456789 = a(n) for all n > m. Let d = number of numbers containing all distinct digits, using digits 0 to 9, then is it conjectured that: m is not equal to d, i.e., m < d. Question: what is m?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jun 06 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003322 | OEIS-A095223

a(n) is the least Fibonacci number such that a(n) - a(n-1) is prime; a(0)=1.

Conjecture: The sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jun 10 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003323 | OEIS-A095224

Least squarefree Fibonacci number with exactly n prime divisors.

Conjecture: The sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jun 10 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003324 | OEIS-A095237

a(1)=1; then for n even, a(n)=(sum of previous terms times n) plus 1, for n odd, a(n)=(sum of previous terms times n) minus 1.

Conjecture: There are infinitely many primes in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jun 13 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003325 | OEIS-A095238

a(1) = 1, a(n) = n*(sum of all previous terms mod n).

An open question is whether the sequence contains zeros except for the 3rd and the 5th number. I checked this up to a(10000), which happens to be 99990000. - _Johan Claes_, Jun 16 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jun 15 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003326 | OEIS-A095258

a(n+1) is the smallest divisor of (2 + sum of first n terms) but not occurring earlier; a(1) = 1.

Conjecture: integer permutation with inverse A095259: a(A095259(n))=A095259(a(n))=n. - Comment revised: _Reinhard Zumkeller_, Dec 31 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, May 31 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003327 | OEIS-A095259

Smallest m such that A095258(m) = n.

Inverse of A095258, if the conjecture is true, that this is a permutation: a(A095258(n)) = A095258(a(n)) = n;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, May 31 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003328 | OEIS-A095276

Length of n-th run of identical symbols in A095076 and A095111.

Conjecture: it appears that the asymptotic frequencies of terms 1, 2 and 3 are 1/2, 1/(2*phi^2) and 1/(2*phi) respectively, where phi = (1+sqrt(5))/2 is the golden ratio. - _Vladimir Reshetnikov_, Mar 17 2022 For a proof of this conjecture see my link to a095276.pdf. - _Michel Dekking_, Jun 25 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jun 01 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003329 | OEIS-A095280

Lower Wythoff primes, i.e., primes in A000201.

For generalizations and conjectures, see A184774.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jun 04 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003330 | OEIS-A095371

Number of distinct prime factors of record setting repunits (A328899).

Conjecture: a(24) = 73, a(25) = 94, a(26) = 99, a(27) >= 107, a(28) >= 127, a(29) >= 136, a(30) >= 140, a(31) >= 151, a(32) >= 159, a(33) >= 163, a(34) >= 178, a(35) >= 184, a(36) >= 213, a(37) >= 214. - _Chai Wah Wu_, Nov 01 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jun 04 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003331 | OEIS-A095380

Number of integers not exceeding 2^n that are impossible as sum-of-divisors of other numbers.

Conjecture: ratio of non-sigma numbers tends to one. Increasing majority of numbers is impossible as a sum of divisors.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_ and _Antti Karttunen_, Jun 07 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003332 | OEIS-A095415

Length of repunits of which the prime factor-digit-excess computed by A095414 equals 0.

541, 701, 857 are also terms. Conjecture: Except for the number 4, A046413 is a subsequence. Conjecture: except for the prime powers 9 and 27, all terms are prime. - _Chai Wah Wu_, Nov 03 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Jun 22 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003333 | OEIS-A095689

Permutation of natural numbers, inverse permutation to A094870.

Conjecture: lim_{n->infinity} a(n)/n = 1 (P. Hegarty).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Alec Mihailovs (alec(AT)mihailovs.com), Jul 05 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003334 | OEIS-A095698

Number of permutations of {1,2,3,...,n} where, for 1 < i <= n, the i-th number has maximized sum of the i-1 absolute differences from all previous numbers of the permutation.

With Lewis's formulas, the addition of the g.f.s for a(2*n) and a(2*n+1) yields the conjectures below: 2*x/(-3*x^2+1) - (-x^2+1)/(-6*x^4+5*x^2-1) = (-4*x^3-x^2+2*x+1)/(6*x^4-5*x^2+1). - _Georg Fischer_, Nov 19 2022 a(1) = 1; Conjectured: For k >= 1, a(2k) = a(2k-1) + 2^(k-1) and a(2k+1) = 2*a(2k-1) + a(2k) (needs proof or a reference). Conjecture: a(n) = 5*a(n-2) - 6*a(n-4); g.f.: x*(1+2*x-x^2-4*x^3)/((1-2*x^2)*(1-3*x^2)). - _Colin Barker_, Jul 27 2012 Conjecture: a(n) = 2^(((-1)^n + 2*n - 5)/4)*((-1)^n-1) - 2*3^(((-1)^n + 2*n - 5)/4)*((-1)^n-2). - _Luce ETIENNE_, Dec 20 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rick L. Shepherd_, Jul 06 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003335 | OEIS-A095720

a(1)=1 and, for n>1, a(n)=a(n-1)+Floor(3n/4) if n is even, else a(n)=smallest positive integer which has not yet appeared in the sequence.

Conjecture. For even n, the ratio a(n)/a(n-1) is asymptotic to 2 as n becomes large. (At n=3000, the ratio is 2.0004446.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jul 08 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003336 | OEIS-A095751

Conjectured list of integers known to be friendly but not known to be primitive friendly.

Conjectured list of integers known to be friendly but not known to be primitive friendly.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Walter Nissen_, Jul 09 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003337 | OEIS-A095769

a(1)=a(2)=1; a(n)=a(a(a(a(n-1))))+a(n-a(a(a(n-1)))).

Conjecture: lim_{n->oo} a(n)/n = 0.682327803828... (the real positive root of x^3 + x = 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jun 05 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003338 | OEIS-A095800

Triangle T(n,k) = abs(k *((2*n+1)*(-1)^(n+k)+2*k-1) /4) read by rows, 1<=k<=n.

2. As a conjecture, each row of A095800 could be a distribution governing distinct subsets of types of triangles having the sum in the "How Many Triangles" series A002717. Thus 1 = 1; 5 = (1 + 4), 13 = (2 + 2 + 9)...etc.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Jun 07 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003339 | OEIS-A095811

Greatest number, not divisible by 4, having exactly n partitions into three squares.

These are conjectured values. The Mathematica program checks numbers up to 10^6. Note that a square can be zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jun 07 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003340 | OEIS-A095812

Greatest number, not divisible by 4, having exactly n partitions into three positive squares.

These are conjectured values. The Mathematica program checks numbers up to 10^6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jun 07 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003341 | OEIS-A095849

Numbers j where sigma_k(j) increases to a record for all real values of k.

Conjecture: a number is in this sequence if and only if it is in both A002182 and A095848. - _J. Lowell_, Jun 21 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthew Vandermast_, Jun 09 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003342 | OEIS-A096010

Number of different cycles computed with the generalized 3x+1 problem using C=2, B=Cn+m, A=C^m.

Generalize the 3x+1-Problem from S:= S / 2 if S is even, S:= (S * 3) + 1 if S is odd to S:= S / C if C | S S:= (S * B) + A otherwise. For B=Cn+A, A=C^m the number of different cycles z are computed. Every S leads to a cycle, so it can be conjectured that the number of cycles is infinite. But the number of different cycles seems to be finite. It is conjectured that the last new cycle occurs at the starting number S = B. This was tested with A=1; B=3; C=2 up to S=100000000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Peter Lengler (PeterLengler(AT)t-online.de), Jul 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003343 | OEIS-A096017

Numbers n such that 4^k*n, for k >= 0, have a unique partition into three distinct positive squares.

It is conjectured that this sequence is complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jun 15 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003344 | OEIS-A096021

Greatest number, not divisible by 4, having exactly n partitions into three distinct positive squares.

These are conjectured values. The Mathematica program checks numbers up to 10^6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jun 15 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003345 | OEIS-A096069

Smallest prime ending in prime(n) and == 1 (mod prime(n)), or 0 if no such prime exists.

a(1) = a(3) = 0. Conjecture: No other term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jun 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003346 | OEIS-A096097

a(1) = 2, a(2) = 1; for n >= 3, a(n) = least prime not included earlier that divides the concatenation of all previous terms.

Conjecture:(1) Every concatenation is squarefree. (2) This is a rearrangement of the noncomposite numbers other than 5. Conjecture (1) is false. 3^2 divides the concatenation for a(22) and a(30). - _Sean A. Irvine_, Nov 25 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jun 24 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003347 | OEIS-A096098

a(1) = 2, a(2) = 1; for n >= 3, a(n) = least number not included earlier that divides the concatenation of all previous terms.

Conjecture (1) Every concatenation is squarefree. Conjecture (2) This is a rearrangement of the squarefree numbers not divisible by 5. False! (The a(n) are not always squarefree, since a(12)=49 and a(14)=9.) Conjecture (3) the sequence for n>=2 is a permutation of the positive integers not divisible by 2 or 5. Assuming Conjecture (3), the smallest number yet to appear is 89. - _Sean A. Irvine_, May 11 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jun 24 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003348 | OEIS-A096121

Number of full spectrum rook's walks on a (2 X n) board.

Conjecture: a(n) = 2*n!*A102038(n). - _Mikhail Kurkov_, Feb 07 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo van der Sanden_, Jul 22 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003349 | OEIS-A096123

Least product n*(n-1)*(n-2)*...*(n-k+1) divisible by (n-k)!.

Conjecture: a(n) = n!/(p-1)! for all sufficiently large n, where p is the least prime such that n <= 2*p (Amarnath Murthy). A096974 gives numbers n such that a(n) = n!/(nextprime(n/2)-1)! and indicates that this conjecture is most probably false.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 01 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003350 | OEIS-A096126

a(n) is the least integer of the form (n^2)!/(n!)^k.

(n^2)!/(n!)^(n+1) is an integer for every n (see A057599). Hence k >= n+1. Conjecture: k=n+1 only when n is prime or a power of a prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003351 | OEIS-A096127

a(n) is the largest k such that (n^2)!/(n!)^k is an integer.

Conjecture: a(n)=n+1 only when n is prime or a power of a prime. [Verified for n=2..5000. - _Amiram Eldar_, Apr 06 2021]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003352 | OEIS-A096135

Triangle read by rows which contains in row n that permutation of the n numbers T(n-1)+1..T(n) which yields a smallest multiple of n after concatenation. T(n) are the triangular numbers. If no such multiple exists, the row contains zeros.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 06 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003353 | OEIS-A096164

Triangle T(n,m) read by rows: matrix product A053121 * A038207.

Conjecture: Sum_{m=0..n} T(n,m) = A126931(n). - _R. J. Mathar_, Mar 25 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Jun 19 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003354 | OEIS-A096222

Number of different rectangles when a piece of paper is folded n times in alternate directions.

Conjectures from _Colin Barker_, Feb 04 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Bill Liebeskind (billlieb(AT)hotmail.com), Jul 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003355 | OEIS-A096231

Number of n-th generation triangles in the tiling of the hyperbolic plane by triangles with angles {Pi/2, Pi/3, 0}.

The g.f. (1+X)^2 * (1+X+X^2) / (1-X^2-X^3) follows from the Cannon-Wagreich paper, Prop. 3.1, so the g.f. and the recurrence are now a theorem, no longer conjectures, and the additional terms and the Mma program are now justified. - _N. J. A. Sloane_, Dec 29 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | Bellovin, Kennedy, Stansifer, Wong (chrkenn(AT)bergen.org), Jul 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003356 | OEIS-A096270

Fixed point of the morphism 0->01, 1->011.

Conjecture: a(n) is given recursively by a(1)=0 and, for n>1, by a(n)=1 if n=F(2k+1) and a(n)=a(n-F(2k+1)) otherwise, where F(2k+1) is the largest odd-indexed Fibonacci number smaller than or equal to n. (This has been confirmed for more than nine million terms.) The odd-indexed bisection of the Fibonacci numbers (A001519) is {1, 2, 5, 13, 34, 89, ...}. So by the conjecture, we would expect that a(30) = a(30-13) = a(17) = a(17-13) = a(4) = a(4-2) = a(2) = 1, which is in fact correct. - _John W. Layman_, Jun 29 2004 Proof of the above conjecture: Let g be the morphism above: g(0)=01, g(1)=011. Then g^n(0) has length F(2n+1), and (a(n)) starts with g^n(0) for all n>0. Obviously g^n(0) ends in 1 for all n, proving the first part of the conjecture. Since v(n) has length F(2n-1), which is the largest odd-indexed Fibonacci number smaller than or equal to m for all m between F(2n-1) and F(2n+1), the claim proves the second part of the conjecture. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 22 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003357 | OEIS-A096293

Number of iterations of n -> n + (sum of squares of digits of n) needed for the trajectory of n to join the trajectory of A033936.

Conjecture: For any positive integer starting value n, iterations of n -> n + (sum of squares of digits of n) will eventually join A033936 (verified for all n up to 20000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jun 24 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003358 | OEIS-A096303

Number of iterations of n -> n + (number of 1's in binary representation of n) needed for the trajectory of n to join the trajectory of A010062.

Conjecture: For any positive integer starting value n, iterations of n -> n + (number of 1's in binary representation of n) will eventually join A010062.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jun 25 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003359 | OEIS-A096304

Numbers k such that 3k does not divide (6k-4)!/((3k-2)!*(3k-1)!).

a(n) = 9 * A005836(floor(n/6)) + (n mod 6) (conjectured) (confirmed, see links).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Aug 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003360 | OEIS-A096306

Number of iterations of n -> n + (largest digit of n) needed for the trajectory of n to join the trajectory of A045844.

Conjecture: For any positive integer starting value n, iterations of n -> n + (largest digit of n) will eventually join A045844.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jun 26 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003361 | OEIS-A096316

Given the number wheel 0,1,2,3,4,5,6,7,8,9 then starting with 2, the next number is a prime p number of positions from the previous number found, for p=2,3,...

Conjecture: This sequence carried to infinity is non-repeating and non-terminating. If we concatenate the numbers and introduce a decimal point somewhere, we will get an irrational number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Aug 02 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003362 | OEIS-A096335

Number of iterations of n -> n + tau(n) needed for the trajectory of n to join the trajectory of A064491, or -1 if the two trajectories never merge.

Conjecture: For any positive integer starting value n, iterations of n -> n + tau(n) will eventually join A064491 (verified for all n up to 50000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jun 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003363 | OEIS-A096340

Decimal expansion of conjectured value of constant C such that, if f(n) is the minimal area of a convex lattice polygon with n vertices, then f(n)/n^3 -> C as n -> infinity.

Decimal expansion of conjectured value of constant C such that, if f(n) is the minimal area of a convex lattice polygon with n vertices, then f(n)/n^3 -> C as n -> infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 30 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003364 | OEIS-A096380

Differences between the sum of the first three primes and the fourth prime in consecutive prime quadruples.

There are occurrences where the next term is less than the current term. Conjecture: The number of occurrences where the current term exceeds the next term is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Aug 04 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003365 | OEIS-A096427

Decimal expansion of 1/(sqrt(2)*G), where G is Gauss's constant A014549.

Bala's 2019 theta-series conjecture follows from the Jacobi imaginary transformation: Sum_{k=-oo..oo} exp(-Pi*(k+1/2)^2) = Sum_{k=-oo..oo} (-1)^k*exp(-Pi*k^2). - _Eric W. Weisstein_, Oct 19 2026 Conjecturally, C = (1/sqrt(2)) * 1/(Sum_{n = -oo..oo} exp(-Pi*(n+1/2)^2))^2. Conjecture: Equals Sum_{k>=0} A004403(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Jul 21 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003366 | OEIS-A096469

Smallest number m such that the concatenation of n+1 numbers m^0, m^1,..., m^(n-1), m^n is a prime.

Conjecture: This sequence is infinite. a(n) = 1 if and only if n + 1 is in the sequence A004023, so a(1) = a(18) = a(22) = a(316) = a(1030) = a(49080) = a(86452) = 1 and there are no other n less than 86453 such that a(n) = 1. Every term of this sequence is odd and for each n, 5 doesn't divide a(n). a(50) is greater than 11111.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Jun 23 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003367 | OEIS-A096471

Number of degree-n permutations with exactly one even cycle.

Conjecture D-finite with recurrence a(n) -3*a(n-1) +(-2*n^2+8*n-3)*a(n-2) +(n-2)*(4*n-15)*a(n-3) +(n-3)*(n^3-9*n^2+22*n-7)*a(n-4) -(n-3)*(n-4)*(n-5)^2*a(n-5)=0. - _R. J. Mathar_, Jul 06 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Aug 13 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003368 | OEIS-A096537

E.g.f.: A(x) = exp(x*exp(2*x*exp(3*x*exp(...exp(n*x*exp(...))...)))), for n>=1.

Conjecture: a(n) ~ 2 * Pi * n^(2*n + 1/2) / (exp(n) * (exp(1) - 1)^n). - _Vaclav Kotesovec_, Dec 16 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 24 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003369 | OEIS-A096597

Triangle read by rows: T[n,m] = number of plane partitions of n whose 3-dimensional Ferrers plot just fits inside an m X m X m box, i.e., with Max[parts, rows, columns] = m.

Conjecture: the last (floor(n/2)) terms of each row read backwards are 3*A091360 (partial sums of A000219).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_, Aug 14 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003370 | OEIS-A096732

a(n)=det(M_n) where M_n is the n X n matrix m(i,j)=i^j*j^i.

Conjecture: lim_{n->oo} a(n)^(1/n^2)/n = 2*exp(-3/2). - _Vaclav Kotesovec_, Apr 19 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 30 2002 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003371 | OEIS-A096858

Triangle read by rows in which row n gives the n-set obtained as the differences {b(n)-b(n-i), 0 <= i <= n-1}, where b() = A005318().

It is conjectured that the triangle has the property that all 2^n subsets of row n have distinct sums. This conjecture was proved by T. Bohman in 1996 - _N. J. A. Sloane_, Feb 09 2012 It is also conjectured that in some sense this triangle is optimal. See A005318 for further information and additional references.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 18 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003372 | OEIS-A096913

Numbers k such that (largest digit of k)^(smallest digit of k) + k is a square.

Conjecture: 0 and 3844 are the only squares in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Aug 17 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003373 | OEIS-A096969

Number of ways to number the cells of an n X n square grid with 1,2,3,...,n^2 so that successive integers are in adjacent cells (horizontally or vertically).

Conjecture: Limit_{n->oo} log_(n+1)!(a(n+1)) - log_n!(a(n)) = c, where 0.09 < c < 0.11. - _Nicolas Bělohoubek_, Jun 12 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jul 16 2004, at the suggestion of _Leroy Quet_, Jul 05 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003374 | OEIS-A096974

Numbers n such that A096123(n) = n!/(nextprime(n/2)-1)!.

Suggested by the apparently false conjecture (A. Murthy) that A096123(n) = n!/(nextprime(n/2)-1)! for all sufficiently large n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Jul 17 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003375 | OEIS-A097042

G.f. = (1 + 4 * g.f. for A096661)/(1 + 2 Sum_{m>=1} (-1)^m*q^(m^2)).

Conjectural g.f.: 1 + 2*Sum_{n >= 1} q^(n*(n+1)/2)/((1 + q^n) * Product_{k = 1..n} 1 - q^k). - _Peter Bala_, Feb 19 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 15 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003376 | OEIS-A097060

Revrepfigits (reverse replicating Fibonacci-like digits): Numbers k whose reversal occurs in a sequence generated by starting with the k digits of a number and then continuing the sequence with a number that is the sum of the previous k terms.

Numbers ending in zero are not permitted since the zeros are dropped upon reversal. However, terms with internal zeros such as 90711 are permitted. Conjectures: 1. Sequence is infinite. 2. Revrepfigits are more rare than repfigits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Sep 15 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003377 | OEIS-A097083

Positive values of k such that there is exactly one permutation p of (1,2,3,...,k) such that i+p(i) is a Fibonacci number for 1<=i<=k.

Conjecture: a(n) = floor(F(n+3)/sqrt(5)), where F(n) = A000045(n) are Fibonacci numbers. - _Vladimir Reshetnikov_, Nov 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jul 23 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003378 | OEIS-A097143

a(n) is the least prime greater than a(n-1) such that n is a quadratic residue mod p.

Conjecture: Lim_n->Inf a(n)/A097144(n) = 1 <==> a(P_2n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jul 26 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003379 | OEIS-A097144

Primes not appearing in A097143.

Conjecture: Lim_n->Inf a(n)/A097143(n) = 1 <==> a(P_2n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jul 26 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003380 | OEIS-A097154

Number of iterations of f(x) = reverse(x) - maxdigit(x) needed to reach zero, or -1 if zero is never reached, where maxdig(n) is the largest digit of n.

a(89)=-1 because 89 is the first number that will not reach zero: 98-9=89. It is conjectured that 0 and 89 are the only fixed points of f. See A097155 for numbers that reach 89.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Aug 01 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003381 | OEIS-A097155

Numbers that reach the fixed point 89 under iteration of f(x) = reverse(x) - maxdigit(x).

89 is the first number that will not reach zero under iteration of f(x) = reverse(x) - maxdigit(x), since 98-9=89. It is conjectured that 0 and 89 are the only fixed points of f.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Aug 01 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003382 | OEIS-A097180

Row sums of triangle A097179, in which the n-th row polynomial R_n(y) is formed from the initial (n+1) terms of g.f. A077860(y)^(n+1), where R_n(1/2) = 4^n for all n>=0.

Conjecture: n*(n-1)*(n+1)*a(n) -12*n*(2*n-1)*(n-1)*a(n-1) +12*(n-1) * (16*n^2-32*n+17)*a(n-2) -16*(4*n-5)*(4*n-7)*(2*n-3)*a(n-3) = 0. - _R. J. Mathar_, Nov 16 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 02 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003383 | OEIS-A097188

G.f. A(x) satisfies A057083(x*A(x)) = A(x) and so equals the ratio of the g.f.s of any two adjacent diagonals of triangle A097186.

Conjecture: (n+1)*a(n) +3*(-3*n+1)*a(n-1)=0. - _R. J. Mathar_, Nov 16 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003384 | OEIS-A097189

Row sums of triangle A097186, in which the n-th row polynomial R_n(y) is formed from the initial (n+1) terms of g.f. A057083(y)^(n+1), where R_n(1/3) = 3^n for all n >= 0.

Conjecture: n*(n-1)*a(n) - (19*n-18)*(n-1)*a(n-1) + 9*(11*n^2-31*n+22)*a(n-2) - 9*(3*n-4)*(3*n-5)*a(n-3) = 0. - _R. J. Mathar_, Nov 16 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003385 | OEIS-A097192

Main diagonal of triangle A097190, in which the n-th row polynomial R_n(y) is formed from the initial (n+1) terms of g.f. A097191(y)^(n+1), where R_n(1/3) = 9^n for all n>=0.

Conjecture: n*a(n) +3*(1-9*n)*a(n-1) = 0. - _R. J. Mathar_, Nov 16 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003386 | OEIS-A097204

Binomial transform of A033312.

Conjecture: a(n) +(-n-5)*a(n-1) +(5*n+3)*a(n-2) +4*(-2*n+3)*a(n-3) +4*(n-3)*a(n-4)=0. - _R. J. Mathar_, Nov 16 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ross La Haye_, Sep 16 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003387 | OEIS-A097243

Expansion of 1 + 32 * (eta(q^4) / eta(q))^8 in powers of q.

Conjecture: Sum_{k>=0} a(k) / exp(k*Pi) = 3. - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Aug 02 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003388 | OEIS-A097267

Least positive integer that is the diagonal of n primitive Pythagorean quadruples, or zero if no such integer exists.

Conjecture: a(15)=a(38)=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ray Chandler_, Aug 19 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003389 | OEIS-A097324

Numbers n such that A067655(n) is different from A049606(n).

We conjecture that the sequence is infinite, the sequence and its complement (cases where the two values are equal) equipartition N and the difference between consecutive members of this sequence never exceeds c=7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Aug 11 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003390 | OEIS-A097331

Expansion of 1 + 2x/(1 + sqrt(1 - 4x^2)).

Conjecture: (n+1)*a(n) + n*a(n-1) + 4*(-n+2)*a(n-2) + 4*(-n+3)*a(n-3)=0. - _R. J. Mathar_, Dec 02 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Aug 05 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003391 | OEIS-A097332

Expansion of (1/(1-x))(1+2x/(1-x+sqrt(1-2x-3x^2))).

Conjecture D-finite with recurrence: (n+1)*a(n) - 3*n*a(n-1) + (-n+5)*a(n-2) + 3*(n-2)*a(n-3) = 0. - _R. J. Mathar_, Nov 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Aug 05 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003392 | OEIS-A097356

Number of partitions of n into parts not greater than sqrt(n).

a(n^2) ~ c * d^n / n^2, where d = A258268 = 9.153370192454122461948530292401354... and c = 0.158208720267250414976631099923 8... [see A206226, constant c(1)]. The upper bound of a(n) is c * d^sqrt(n) / n, see graph. For the lower bound, the constant c = 0.088154883798697116... (conjectured). - _Vaclav Kotesovec_, Jan 08 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Aug 08 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003393 | OEIS-A097386

Numbers n such that (largest digit of n)^(smallest digit of n) + n is prime.

No term ends in 9. Conjecture: Let f(x)=(largest digit of x)^(smallest digit of x) + x. There are infinitely many positive integers n such that f(n) and f(n+1) are both prime; see A097387.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Aug 18 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003394 | OEIS-A097387

Let f(x)=(largest digit of x)^(smallest digit of x) + x (A097385). Sequence gives numbers n such that f(n) and f(n+1) are both prime.

Conjecture: Terms will always be multiples of ten. Aug 21, 2004: _Dean Hickerson_ proved this.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Aug 18 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003395 | OEIS-A097391

The number of hierarchies of n labeled elements with at least one subhierarchy composed of exactly 2 levels and no subhierarchy with more than 2 levels.

Conjectured to be (A378854) /3. - _Wouter Meeussen_, Feb 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Wieder_, Aug 13 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003396 | OEIS-A097399

Maximum of the determinant over all permutations of the entries of a 3 X 3 matrix which are consecutive integers in the range (n-4,n+4).

G.f.: (x^13+12*x^3+50*x^2-68*x+86) / (x-1)^2. [_Colin Barker_, Dec 13 2012] [I suspect this is merely a conjecture. - _N. J. A. Sloane_, Jun 09 2018]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Aug 19 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003397 | OEIS-A097406

Largest primitive prime factor of 2^n-1, or a(n) = 1 if no such prime exists.

Conjectures: (1) For every n the highest unique prime factor is of the form kn+1. The values for k are in A097407. (2) For each composite n many factors of the form kn+1 occur intermittently but always singly in any cofactor pair. (3) For each prime n every factor is of the form kn+1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Matosic_, Aug 16 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003398 | OEIS-A097509

a(n) is the number of times that n occurs as floor(k * sqrt(2)) - k.

Conjecture: a(n+1) = A082844(n). - _Benedict W. J. Irwin_, Mar 13 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Odimar Fabeny_, Aug 26 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003399 | OEIS-A097587

a(n)= floor(P(k)/P(n)) with P(k)=Fortunate prime numbers for P(n)# (A005235).

Conjecture : a(n) is always < 8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Aug 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003400 | OEIS-A097588

a(n)=floor(P(k)/P(n)) with P(k)=lesser Fortunate numbers for P(n)# (A055211).

Conjecture : a(n) is always < 8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Aug 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003401 | OEIS-A097602

a(n+1) = a(n) + number of squares so far; a(1) = 1.

Conjecture: a(n) = m^2 iff m mod 3 > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Aug 30 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003402 | OEIS-A097702

a(n) = (A063880(n) - 108)/216.

Conjecture: k is a term iff 6*k+3 is squarefree. - _Vladeta Jovovic_, Aug 27 2004 It is only a conjecture that all terms are integers (confirmed up to 10^6 by _Robert G. Wilson v_). The first conjecture is true. If m = 216*k + 108 = 108 * (2*k + 1) is a term of A063880, then 2*k+1 is a squarefree number coprime to 6. This is because sigma(n)/usigma(n) is multiplicative, equals 1 if and only if n is squarefree and larger than 1 otherwise, sigma(108)/usigma(108) = 2 and sigma(3^k)/usigma(3^k) increases with k. 6*k+3 = 3*(2*k+1) is squarefree because 2*k+1 is a squarefree coprime to 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Aug 26 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003403 | OEIS-A097703

Numbers j such that m = 216*j + 108 satisfies sigma(m) != 2*usigma(m).

Conjecture: all numbers of form 3k + 1 are here. Other terms are listed in A097704. The conjecture is true. If j = 3*k+1, then m = 324*(2*k+1). Let e = A007949(2*k+1) >= 0, so 2*k+1 = 3^e * i and i coprime to 6. Then sigma(m)/(2 * usigma(m)) = (7/20) * (3^(e+5)-1)/(3^(e+4)+1) * sigma(i)/usigma(i) >= 847/820 > 1, because sigma(i)/usigma(i) >= 1 for all i.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Aug 26 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003404 | OEIS-A097704

Terms of A097703 that are not of the form 3*k + 1.

Conjecture: "most" of the terms also belong to [(A067778-1)/2]. Exceptions are {302, 2117, ...} (A098241). In other words, most terms satisfy: GCD(2*k+1, numerator(B(4*k+2))) is not squarefree, with B(n) the Bernoulli numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Aug 26 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003405 | OEIS-A097724

Triangle read by rows: T(n,k) is the number of left factors of Motzkin paths without peaks, having length n and endpoint height k.

Conjecture: the row reverse triangle is the triangle of connection constants for expressing the polynomial u(n,x+1) as a linear combination of the polynomials u(k,x), 0 <= k <= n, where u(n,x) = U(n,x/2) with U(n,x) the n-th Chebyshev polynomial of the second kind. An example is given below. Cf. A205810. - _Peter Bala_, Jun 26 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Sep 11 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003406 | OEIS-A097785

Counterexamples to the conjecture that an even, prime-indexed triangular plus 1 equals a prime or that an odd, prime-indexed triangular minus 2 equals a prime.

Counterexamples to the conjecture that an even, prime-indexed triangular plus 1 equals a prime or that an odd, prime-indexed triangular minus 2 equals a prime. In the entry for 496, Wells remarks that it is the smallest counterexample to the conjecture that an even, prime-indexed triangular plus 1 equals a prime, since 497 is not prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alonso del Arte_ and _Robert G. Wilson v_, Aug 24 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003407 | OEIS-A097814

Expansion of e.g.f. exp(3*x)/(1-3*x).

Conjecture: a(n) + 3*(-n-1)*a(n-1) + 9*(n-1)*a(n-2) = 0. - _R. J. Mathar_, Dec 21 2014 Mathar's conjecture is true (see Fried link). - _Sela Fried_, Dec 08 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Aug 26 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003408 | OEIS-A097877

Triangle read by rows: T(n,k) is the number of Dyck n-paths with k large components, 0 <= k <= n/2.

Conjecture: This is the statistic "indegree" of elements in Pallo's comb posets. For the statistic "outdegree", see A009766. - _F. Chapoton_, Apr 18 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Callan_, Sep 21 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003409 | OEIS-A097913

G.f.: (1+x^18)/((1-x)*(1-x^8)*(1-x^12)*(1-x^24)).

Conjectured Poincaré series [or Poincare series] for genus 2 Siegel theta series of odd unimodular lattices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 04 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003410 | OEIS-A097933

Primes p that divide 3^((p-1)/2) - 1.

Conjecture: Let r(n) = (a(n) - 1)/(a(n) + 1) if a(n) mod 4 = 1, (a(n) + 1)/(a(n) - 1) otherwise; then Product_{n>=1} r(n) = (6/5) * (6/7) * (12/11) * (18/19) * ... = 2/sqrt(3). - _Dimitris Valianatos_, Mar 27 2017 Conjecture: These are the exclusive prime factors of the set of integers d > 1 such that there exist primitive Heronian triangles with sides {b, b+d, b+2d} for one or more integers b.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Sep 04 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003411 | OEIS-A097945

a(n) = mu(n)*phi(n) where mu(n) is the Mobius function (A008683) and phi(n) is the Euler totient function (A000010).

Conjecture: Sum_{n>=1} mu(n)/phi(n) = Sum_{n>=1} a(n)/phi(n)^2 = 0. It is true that Sum_{n>=1} mu(n)/phi(n)^s = 0 at least for s > 1 since: phi(2)=1, phi is multiplicative, so for n's that are squarefree, the phi(n) values can be partitioned in pairs where phi(m)=phi(2m) and mu(m) = -mu(2m). So Sum_{i=1..n} mu(i)/phi(i)^s < Sum_{j=floor(n/2)..n} 1/phi(j)^s, which approaches 0 as n increases since (1) n^(1-e) < phi(n) < n for any e > 0 and n > N(e) and (2) Sum_{i..n} 1/n^s converges for s > 1. Conjecture: Sum_{n>=1} mu(n)/phi(n)^z = 0 for Re(z) > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerald McGarvey_, Sep 04 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003412 | OEIS-A097967

a(n) = Sum_{k=1..n} (P(n,k) + C(n,k)).

Conjecture: a(n) +(-n-6)*a(n-1) +(6*n+7)*a(n-2) +(-13*n+14)*a(n-3) +4*(3*n-8)*a(n-4) +4*(-n+4)*a(n-5)=0. - _R. J. Mathar_, May 29 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ross La Haye_, Sep 21 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003413 | OEIS-A098009

Numbers k such that the transient part of the aliquot sequence for k is finite and sets a new record.

In order to extend this there is the problem that there are small numbers (276, 552, etc.) for which it is not presently known if they cycle. I propose that we assume these do not cycle, but mark the records beyond where this becomes an issue as conjectural only.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 10 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003414 | OEIS-A098034

Numbers that are divisible both by the sum and by the product of the squares of their digits.

Called "insolite numbers" in the paper by J.-M. De Koninck and N. Doyon, which contains a list of insolite numbers below 10^18. Their list lacks a(63)=112264112111616. They conjectured that 11111111113111113111111111175 is the smallest insolite number containing the digit '5'. I verified this claim and found a further such number, 1111111111111111171111111111911111375, which may not be the second one. - _Giovanni Resta_, Oct 19 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Sep 10 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003415 | OEIS-A098041

Arithmetic mean of successive Mersenne primes.

Sequence is based on the following (false) conjecture Shanks mentions in his paper (p.6), which someone else made: "If M_p1 and M_p2 are successive Mersenne primes, their arithmetic mean is also prime." The first four values in the sequence are prime, while the rest up to (2^756839-1 + 2^216091-1)/2 are not. Conjecture: there are no more primes in this sequence after 4159.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Oct 24 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003416 | OEIS-A098052

T(n,k) counts the solid partitions of n that can be extended to a solid partition of n+1 in exactly (k+3) ways. Equivalently, the number of solid partitions of n that have exactly k+3 partitions of n+1 majoring them.

First column is conjectured to be A007426 = tau_4(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_, Sep 11 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003417 | OEIS-A098058

Prime(n) such that 4 does not divide the difference between prime(n) and prime(n+1).

The sequence is infinite, by Dirichlet's theorem on primes in arithmetic progressions. The sequence contains arbitrarily long gaps, by Daniel Shiu's theorem on strings of congruent primes (see A057619 and A057622). Conjecture: The sequence contains arbitrarily long strings of consecutive primes (see A289118). - _Jonathan Sondow_, Jun 25 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Sep 11 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003418 | OEIS-A098151

Number of partitions of 2*n with no part divisible by 3 and all odd parts occurring with even multiplicities.

Let A(q) denote the g.f. of this sequence. Let m be a nonzero integer. The simple continued fraction expansions of the real numbers A(1/(2*m)) and A(1/(2*m+1)) may be predictable. For a given positive integer n, the sequence of the n-th partial denominators of the continued fractions are conjecturally polynomial or quasi-polynomial in m for m sufficiently large. An example is given below. Cf. A080054. - _Peter Bala_, Jun 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Noureddine Chair_, Aug 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003419 | OEIS-A098169

a(n) = Sum_{i=1..n} A098168(i).

Conjecture: a(n)/triangular(n) -> Log(e*Pi/2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Aug 30 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003420 | OEIS-A098274

Sum_{k = 0..n} C(n, k)^2*C(n+k, n)*C(n+2*k, n).

We conjecture that the present sequence satisfies the same pair of supercongruences. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Sep 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003421 | OEIS-A098301

Member r=16 of the family of Chebyshev sequences S_r(n) defined in A092184.

Conjecture: 4*A007655(n+1) + A046184(n) = A055793(n+2) + a(n+1). - _Creighton Dement_, Nov 01 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 18 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003422 | OEIS-A098348

Triangular array read by rows: T(n, k) = number of ordered factorizations of a "hook-type" number with n total prime factors and k distinct prime factors. "Hook-type" means that only one prime can have multiplicity > 1.

The following conjectural formula for the triangle entries agrees with the values listed above: T(n,k) = Sum_{j=0..n-k} 2^(n-k-j)*binomial(n-k,j)*a(k,j), where a(k,j) = 2^j*Sum_{i=j+1..k} binomial(i,j+1)*(i-1)!*Stirling2(k,i). See A098384 for related conjectures. - _Peter Bala_, Apr 20 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alford Arnold_, Sep 04 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003423 | OEIS-A098375

(1/p)*abs(p*(p^(p-1)-1)*B(p-1)-1) when p runs through the primes and B(k) denotes the k-th Bernoulli's number.

Conjecture: p is an odd prime iff p divides p*(p^(p-1)-1)*B(p-1)-1. Seems to be the equivalent (with integer moduli) to Agoh's conjecture (which involves rational moduli).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 26 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003424 | OEIS-A098384

Triangle read by rows of coefficients used to generate diagonals of ordered factorizations as displayed in A098348.

The following formulas are all conjectural:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alford Arnold_, Sep 06 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003425 | OEIS-A098411

Expansion of 1/(sqrt(1-4x)sqrt(1-12x)).

Nguyen and Taggart (see link) conjecture: det[a(i+j) for i,j=0..n] = b(n)*b(n+1)/2 with b(n) = A139685(n). - _Peter Luschny_, May 19 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 07 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003426 | OEIS-A098464

Numbers k such that lcm(1,2,3,...,k) equals the denominator of the k-th harmonic number H(k).

Shiu (2016) conjectured that this sequence is infinite. - _Amiram Eldar_, Feb 02 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Sep 09 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003427 | OEIS-A098472

Least k such that Mersenne-prime(n)*2^k-1 is prime (A000668(n)*2^k-1), or 0 if no such k exists.

Conjecture: Mersenne-prime(12) is a Riesel prime (that is, all numbers k^2*M(12)-1 are composite for all k) and similarly for M(14) and M(15).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Sep 09 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003428 | OEIS-A098519

E.g.f. exp(x)*BesselI(1,2*sqrt(3)*x)/sqrt(3).

Conjecture: (n-1)*(n+1)*a(n) - n*(2*n-1)*a(n-1) - 11*n*(n-1)*a(n-2) = 0. - _R. J. Mathar_, Nov 23 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 12 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003429 | OEIS-A098520

E.g.f. exp(x)*BesselI(1,4*x)/2.

Conjecture: (n-1)*(n+1)*a(n) - n*(2n-1)*a(n-1) - 15*n*(n-1)*a(n-2) = 0. - _R. J. Mathar_, Dec 11 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 12 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003430 | OEIS-A098521

E.g.f. exp(x)*BesselI(2,2*sqrt(2)*x)/2.

Conjecture: (n+2)*a(n) -(4n+3)*a(n-1) -3*(2n+1)*a(n-2) +(20n-29)*a(n-3) +21*(n-3)*a(n-4)=0. - _R. J. Mathar_, Dec 08 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 12 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003431 | OEIS-A098530

T(n,k) counts solid partitions of n+1 that can be 'shrunk' in k ways to a solid partition of n by removing 1 element from it. Equivalently, it counts how many solid partitions of n+1 have k different solid partitions of n it just covers.

Sequence starts 4; 4,6; 10,12,4; 4,42,12,1; 16,60,60,4; 4,105,164,34; Row sums are A000293= the solid partitions of n+1 apart from offset. First column conjectured to be the (beheaded) A007426.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_, Sep 12 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003432 | OEIS-A098550

The Yellowstone permutation: a(n) = n if n <= 3, otherwise the smallest number not occurring earlier having at least one common factor with a(n-2), but none with a(n-1).

Conjectures: 7. Based on the first 250000 terms, I conjectured on Nov 30 2014 that a(n)/n <= (Pi/2)*log n. 8. The primes in the sequence appear in their natural order.

This conjecture is very plausible but as yet there is no proof. - _N. J. A. Sloane_, Jan 29 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Sep 14 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003433 | OEIS-A098557

Expansion of e.g.f. (1/2)*(1+x)*log((1+x)/(1-x)).

conjecture: -a(n) +a(n-1) +(n-1)*(n-3)*a(n-2)=0. - _R. J. Mathar_, Nov 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 14 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003434 | OEIS-A098660

E.g.f. BesselI(0,2*sqrt(2)*x) + BesselI(1,2*sqrt(2)*x)/sqrt(2).

Conjecture: (n+1)*a(n) + (n-1)*a(n-1) - n*a(n-2) + (2-n)*a(n-3) = 0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003435 | OEIS-A098662

Expansion of e.g.f. BesselI(0,2*sqrt(3)*x) + BesselI(1,2*sqrt(3)*x)/sqrt(3).

Conjecture: (n+1)*a(n) + 6(n-1)*a(n-1) - 12n*a(n-2) + 72*(2-n)*a(n-3) = 0. - _R. J. Mathar_, Dec 08 2011 Conjecture confirmed using the differential equation x*(6x+1)*(12*x^2-1) * g'(x) + (6*x-1)*(12*x^2+6*x+1)*g(x) + 2*x + 1 = 0 satisfied by the g.f. - _Robert Israel_, Aug 23 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003436 | OEIS-A098664

Expansion of e.g.f. BesselI(0,4*x)+BesselI(1,4*x)/2.

Conjecture: (n+1)*a(n) + 8*(n-1)*a(n-1) - 16*n*a(n-2) + 128*(2-n)*a(n-3) = 0. - _R. J. Mathar_, Dec 08 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003437 | OEIS-A098746

Number of permutations of [1..n] which avoid 4231 and 42513.

Conjecture: a(n) is the number of permutations of [1..n] which avoid 1342 and 13254. - _Alexander Burstein_, Oct 19 2017 Conjecture: 2*(2*n-1)*(n-1)*a(n) +3*(11*n^2-67*n+76)*a(n-1) +3*(-155*n^2+931*n-1356)*a(n-2) +(469*n^2-2799*n+4070)*a(n-3) -48*(3*n-8)*(3*n-10)*a(n-4)=0. - _R. J. Mathar_, May 30 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 30 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003438 | OEIS-A098833

Numbers n such that the sum of primes dividing n (with repetition) is a Fibonacci number.

Conjecture: there are infinitely many consecutive terms in this sequence. For example, sopfr(143335) = 377 and sopfr(143336) = 89 are both Fibonacci numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Oct 10 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003439 | OEIS-A098860

Primes that are conjectured to lead to a one-cycle for every natural number x in the following (nontrivial) generalization of the (3x+1) problem.

Primes that are conjectured to lead to a one-cycle for every natural number x in the following (nontrivial) generalization of the (3x+1) problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Herman Roelants_, Oct 11 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003440 | OEIS-A098894

Values of k such that {s(1),...,s(k)} is a palindrome, where {s(1),s(2),...} is the fixed point of the substitutions 0->1 and 1->110.

Conjecture: partial sums of A152113. - _Sean A. Irvine_, Jul 14 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Nov 04 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003441 | OEIS-A098926

Permanent of the (0,1)-matrix with ij-th entry equal to zero iff (i=1,j=1),(i=1,j=2),(i=1,j=3),(i=2,j=3),(i=3,j=3),... In other words, the ij-th entry of the matrix is zero iff it is on the path which start from the entry (i=1,j=1) and moves in the matrix alternating 3 steps to the right to 3 steps down.

Conjectured recurrence of order 8 and degree 7: (3*n^5 + 80*n^4 + 763*n^3 + 3184*n^2 + 5915*n + 4080)*a(n+8) + (-3*n^5 - 83*n^4 - 833*n^3 - 3663*n^2 - 6967*n - 4465)*a(n+7) + (-3*n^7 - 122*n^6 - 2039*n^5 - 18038*n^4 - 90333*n^3 - 252920*n^2 - 364438*n - 211080)*a(n+6) + (17*n^5 + 445*n^4 + 4253*n^3 + 17161*n^2 + 24893*n + 1765)*a(n+5) + (9*n^7 + 318*n^6 + 4409*n^5 + 30672*n^4 + 113879*n^3 + 219268*n^2 + 186788*n + 35600)*a(n+4) + (-n^5 + 103*n^4 + 2125*n^3 + 14395*n^2 + 38283*n + 32845)*a(n+3) + (-9*n^7 - 294*n^6 - 3677*n^5 - 22722*n^4 - 76591*n^3 - 146304*n^2 - 157554*n - 81720)*a(n+2) - (n+1)*(13*n^4 + 388*n^3 + 3717*n^2 + 13424*n + 16865)*a(n+1) + n*(n+1)*(3*n^5 + 95*n^4 + 1113*n^3 + 5983*n^2 + 14907*n + 14025)*a(n) = 0. Conjectured differential equation for the generating function: x^5*(x^2 - 1)^3*(x^8 - 2*x^7 - 12*x^6 + 28*x^5 - 10*x^4 - 22*x^3 + 4*x^2 + 4*x + 1)*f'''(x) + x^4*(x^2 - 1)*(5*x^12 - 12*x^11 - 76*x^10 + 212*x^9 - 143*x^8 - 156*x^7 + 80*x^6 + 36*x^5 + 7*x^4 - 24*x^3 - 4*x^2 + 8*x + 3)*f''(x) + x*(x^2 - 1)*(4*x^14 - 9*x^13 - 95*x^12 + 342*x^11 - 288*x^10 - 29*x^9 - 75*x^8 - 336*x^7 + 252*x^6 + 121*x^5 - 53*x^4 + 10*x^3 - 3*x - 1)*f'(x) + 2*(2*x^14 - 13*x^13 + 12*x^12 - 128*x^11 + 40*x^10 + 513*x^9 - 410*x^8 - 24*x^7 + 218*x^6 - 283*x^5 + 16*x^4 + 104*x^3 - 4*x^2 - 9*x - 2)*f(x) = 0. (End)

Conjecture: a(n) ~ exp(-2) * n!, based on the recurrence by _Manuel Kauers_ and _Christoph Koutschan_. - _Vaclav Kotesovec_, Mar 02 2023 For even n, a(n) = A002464(n); for odd n, a(n) = A002464(n) - A002464(n-1). These formulae allow to easily verify the above conjectures. - _Max Alekseyev_, Jun 01 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simone Severini_, Oct 19 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003442 | OEIS-A099118

Conjectured number of times that S(k+n) = S(k), where S is the Kempner function A002034.

Conjectured number of times that S(k+n) = S(k), where S is the Kempner function A002034. Numbers k up to 10^8 have been tested. Tutescu's conjecture is the case n=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Sep 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003443 | OEIS-A099119

Conjectured greatest k such that S(k) = S(k-n), or 0 if no k is known, where S is the Kempner function A002034.

Conjectured greatest k such that S(k) = S(k-n), or 0 if no k is known, where S is the Kempner function A002034. Numbers k up to 10^8 have been tested. Tutescu's conjecture is the case n=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Sep 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003444 | OEIS-A099120

Least m such that n = S(k) = S(k+m) for some k, where S is the Kempner function A002034.

Consider the set Sn of d(n!)-d((n-1)!) positive integers k with S(k) = n, where d is the divisor counting function A000005. For each n, a(n) gives the least difference of integers in the set Sn. For prime n, a(n) = n. For n a power of a prime, a(n) = A046021(n), the least k in Sn. The Tutescu conjecture, which states that the equation S(k) = S(k+1) has no solutions, is equivalent to a(n) > 1 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Sep 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003445 | OEIS-A099143

Least k such that S(k) = S(k+n), or 0 if there is no k, where S is the Kempner function A002034.

The case of n=1 corresponds to the Tutescu conjecture, which states that the equation S(k) = S(k+1) has no solutions. It is conjectured that S(k) = S(k+6) also has no solutions. For odd prime p, a(p) = p. It appears that a(n) <= n, except for n = 1, 2, 6, 24, 48, 120, 240, 720 (A099144).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Sep 30 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003446 | OEIS-A099155

Maximum length of a simple path with no chords in the n-dimensional hypercube, also known as snake-in-the-box problem.

Some confusion seems to exist in the distinction between n-snakes and n-coils. Earlier papers and also A000937 used "snake" to mean a closed path, which is called n-coil in newer notation, see Harary et al. a(8) is conjectured to be 97 by Rajan and Shende. [The true value, however, is 98. See Ostergard and Ville, 2014. - _N. J. A. Sloane_, Apr 06 2014]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Oct 11 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003447 | OEIS-A099188

a(n) = 2*ceiling(n/sqrt(2)).

Conjecturally, length of shortest polygonal path from (0,0) to (n,n) with integer vertices and edges. This is true for n <= 10000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_, Mar 26 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003448 | OEIS-A099237

a(n) = Sum_{k=0..n} binomial(n*(n-k), k).

Conjectures: a(2^k) == 1 (mod 2^k) and a(3^k) == 1 (mod 3^(k+1)); a(p^k) == 1 (mod p^(k+1)) for all primes p >= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 08 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003449 | OEIS-A099251

Bisection of Motzkin sums (A005043).

Conjecture: a(n) = (2/Pi)*Integral_{t=0..1} sqrt((1 - t)/t)*(1 - 8*t + 16*t^2)^n. - _Benedict W. J. Irwin_, Oct 05 2016 Irwin's conjecture is true (see Fried link). - _Sela Fried_, Mar 15 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 16 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003450 | OEIS-A099302

Number of integer solutions to x' = n, where x' is the arithmetic derivative of x.

It appears that there are a finite number of even numbers having a given number of solutions. This conjecture is explored in A189561 and A189562.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Oct 12 2004, Apr 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003451 | OEIS-A099309

Numbers n whose k-th arithmetic derivative is nonzero for all k. Complement of A099308.

Numbers of the form n = m*p^p (where p is prime), i.e., multiples of some term in A051674, have n' = (m + m')*p^p, which is again of the same form, but strictly larger iff m > 1. Therefore successive derivatives grow to infinity in this case, and they are constant when m = 1. There are other terms in this sequence, but I conjecture that they all eventually lead to a term of this form, e.g., 26 -> 15 -> 8 etc. - _M. F. Hasler_, Apr 09 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Oct 12 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003452 | OEIS-A099363

An inverse Chebyshev transform of 1-x.

Conjecture: (n+3)*a(n) +(-n-1)*a(n-1) -4*n*a(n-2) +4*(n-2)*a(n-3)=0. - _R. J. Mathar_, Nov 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 13 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003453 | OEIS-A099370

Chebyshev polynomial of the first kind, T(n,x), evaluated at x=33.

From a(n) = T(n, 33) (see the formula section) and the de Moivre-Binet formula for T(n,x=33) follows a(n+1)/a(n) = 33 + 8*sqrt(17), which is the conjectured value (4+sqrt(17))^2 given in the previous comment by Rick L. Shepherd. - _Wolfdieter Lang_, Jun 28 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 18 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003454 | OEIS-A099468

Numbers n such that there are no primes < 2n in the sequence m(0)=n, m(k+1)=m(k)+4k.

No others < 10^8. Note that 3 divides all these n > 1. This sequence is conjectured to be complete. Related to a question posed in A036468 by Zhang Ming-Zhi. Let r=2s+1 be an odd number. If n = (s+1)^2+s^2, then the sequence m(0)=n, m(k+1)=m(k)+4k for k=0,1,...s calculates the s+1 distinct sums of two squares (r-i)^2+i^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Oct 17 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003455 | OEIS-A099500

Number of distinct products of subsets of integers in the interval [n^2+1, (n+1)^2-1] which are twice a square.

Granville and Selfridge prove Kaplansky's conjecture that a(n) > 0 for all n. Interestingly, the first 24 terms are powers of 2, but the 25th term is not.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Oct 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003456 | OEIS-A099553

Number of rooted 2-connected loopless 4-regular planar maps with n inner faces.

Conjecture: 3*n *(3*n-1) *(5*n-8) *(3*n-2)*a(n) -(n-2) *(2*n-3) *(355*n^2 -703*n +300)*a(n-1) -98*(n-2) *(5*n-3) *(2*n-3) *(2*n-5) *a(n-2)=0. - _R. J. Mathar_, Aug 28 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 18 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003457 | OEIS-A099576

Row sums of triangle A099575.

Conjecture: 4*n*(n-1)*(3*n+2)*(n+2)*a(n) - 36*(n-1)*(n+1)*a(n-1) - 3*n*(3*n+5)*(3*n-1)*(3*n-2)*a(n-2) = 0. - _R. J. Mathar_, Nov 28 2014 The conjecture follows from this. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 23 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003458 | OEIS-A099612

Numerators of the coefficients in the Taylor expansion of sec(x) + tan(x) around x=0.

Conjecture: For n >= 0, (-1)^n * a(n+1) is the numerator of the n-th term of the Taylor expansion of 1/(1 + sin(x)) around x = 0. [This is based on the fact that (sec(x) + tan(x))' = 1/(1 + sin(-x)). _Clark Kimberling_ in A279107 states my conjecture as a fact, but no proof or reference is given.] - _Petros Hadjicostas_, Oct 06 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 18 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003459 | OEIS-A099617

Denominators of the coefficients in the Taylor expansion of sec(x) + tan(x) around x=0.

Conjecture: For n >= 0, (-1)^n * a(n+1)/(n+1) is the denominator of the n-th term of the Taylor expansion of 1/(1 + sin(x)) around x = 0. [This is based on the fact that (sec(x) + tan(x))' = 1/(1 + sin(-x)). Cf. also the comments in A099612 and A279107.] - _Petros Hadjicostas_, Oct 06 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 19 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003460 | OEIS-A099728

Least number B such that (A001359(n) - B^2)^2 - B is also the lesser of larger twin primes, or 0 if no such B exists.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ray G. Opao_, Nov 07 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003461 | OEIS-A099976

Bisection of A000984.

Conjecture: A000265(a(n)) = A063079(n+1), odd part of a(n). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 19 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

B03 A100000 - A149999 | 1,397 条

V3-003462 | OEIS-A100037

Positions of occurrences of the natural numbers as a second subsequence in A100035.

a(n) = 2*n^2 - n + 3 (conjectured). - _Ralf Stephan_, May 15 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Oct 31 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003463 | OEIS-A100038

Positions of occurrences of the natural numbers as third subsequence in A100035.

a(n) = 2*n^2 + 3*n + 6 (conjectured). - _Ralf Stephan_, May 15 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Oct 31 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003464 | OEIS-A100039

Positions of occurrences of the natural numbers as fourth subsequence in A100035.

2n^2 + 7n + 13 (conjectured). - _Ralf Stephan_, May 15 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Oct 31 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003465 | OEIS-A100056

Sum of absolute differences of p(n) defined in A054065.

Conjecture: a(n) provides the maximum sum of difference terms of any permutation of 1 through n. A054065 (or reversals) give the optimal permutations and a(n) gives the sum.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Oct 31 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003466 | OEIS-A100095

An inverse Chebyshev transform of the Fibonacci numbers.

Conjecture: (-n+2)*a(n) +(-n+3)*a(n-1) +(9*n-22)*a(n-2) +(9*n-31)*a(n-3) +20*(-n+3)*a(n-4) +20*(-n+4)*a(n-5)=0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003467 | OEIS-A100096

An inverse Chebyshev transform of the Jacobsthal numbers.

Conjecture: 2*(-n+1)*a(n) +(n+3)*a(n-1) +18*(n-2)*a(n-2) +4*(-n-2)*a(n-3) +40*(-n+3)*a(n-4)=0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003468 | OEIS-A100097

An inverse Chebyshev transform of the Pell numbers.

Conjecture: (-n+2)*a(n) +(-n+3)*a(n-1) +4*(3*n-7)*a(n-2) +4*(3*n-10)*a(n-3) +32*(-n+3)*a(n-4) +32*(-n+4)*a(n-5)=0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003469 | OEIS-A100098

An inverse Chebyshev transform of (1-x)/(1-2x).

Conjecture: 2*n*a(n) + (-5*n+4)*a(n-1) + 2*(-4*n+3)*a(n-2) + 20*(n-2)*a(n-3) = 0. - _R. J. Mathar_, Nov 22 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 04 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003470 | OEIS-A100099

An inverse Chebyshev transform of x/(1-2x).

Conjecture: 2n*a(n) +(-13n+16)*a(n-1) +4(3n-8)*a(n-2) +4(13n-29)*a(n-3) +80(3-n)*a(n-4)=0. - _R. J. Mathar_, Dec 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 04 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003471 | OEIS-A100113

a(n) = n if n <= 2, otherwise (smallest squarefree number m not occurring earlier such that gcd(m, a(n-1)) > 1).

Conjecture: With the three exceptions p = 2, 5, 13, and 31, when a prime p appears it is preceded by 2*p and followed by 3*p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Nov 07 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003472 | OEIS-A100193

a(n) = Sum_{k=0..n} binomial(2*n,n+k)*3^k.

Conjecture: 9*n*a(n) + 6*(11-18*n)*a(n-1) + 16*(26*n-37)*a(n-2) + 256*(5-2*n)*a(n-3) = 0. - _R. J. Mathar_, Nov 09 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 08 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003473 | OEIS-A100200

Decimal Goedelization of antitheorems from propositional calculus, in Richard C. Schroeppel's metatheory of A101273.

Besides A101273, there can also be the subsequence of theorems that can be proved within the more restricted Intuitionist logic; the sequence of well-formed formulas whose truth value is contingent on the truth values of their variables; and many others. As with A101273, I conjecture that a power law approximates the number of integers in this sequence, where the number with N digits is approximately N to the power of some real number D.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Dec 27 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003474 | OEIS-A100348

Numbers n such that n-4^k is a prime for all k > 0 with 4^k < n.

The largest term appears to be 5833497. No others < 10^9; conjectured to be finite. Similar to A067528, which also contains 5 and 17, but a more direct generalization of A039669, a problem due to Erdos.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Nov 18 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003475 | OEIS-A100350

Primes p such that p-2^k is a prime or semiprime for all k > 0 with 2^k < p.

These are the primes in A100349. No others < 10^9; conjecture that this sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Nov 18 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003476 | OEIS-A100351

Numbers n such that n-2^k is a semiprime for all k > 0 with 2^k < n.

A subset of A100349. No others < 10^9; conjecture that this sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Nov 18 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003477 | OEIS-A100571

Cubes m^3 such that m^3 is the sum of m-1 consecutive primes plus a larger prime.

Conjecture: sequence consists of all the cubes > 1 except 8^3=512. - _Giovanni Teofilatto_, Apr 23 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Teofilatto_, Nov 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003478 | OEIS-A100631

Triangle read by rows: T(n,k) = 2*(T(n-1,k-1) - T(n-2,k-1) + T(n-1,k)) for 0 < k < n, T(n,0) = T(n,n) = 1.

All the conjectures below are true because one has to prove only one of them, and the rest follow from the proved one. Conjecture based on _Peter Luschny_'s formulas in other sequences: T(n,k) = 2^(n-k)*hypergeom([-k + 1, n-k], [1], -1) = 2^k*hypergeom([-(n-k) + 1, k], [1], -1). Conjecture based on _Peter Luschny_'s formulas in other sequences: R(n,k) = 2^n*hypergeom([-k + 1, n], [1], -1) = 2^k*hypergeom([-n + 1, k], [1], -1). (End) The above conjecture is true (see the comments).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Dec 03 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003479 | OEIS-A100665

a(n) = round(sqrt(Fibonacci(n))).

This is a special case of the following conjectured relationship: Given a starting sequence of g(0)=0, g(1)=1, ..., g(m)=1 (i.e., 0 then m-1 1's) and then a recurrence relation g(n) = b_1*g(n-1)^c_1 + b_2*g(n-2)^c_2 + ... + b_m*g(n-m)^c_m where the b_i's and c_i's are real numbers > 0 then (b_1*(g(n-1)^c_1)^x + b_2*(g(n-2)^c_2)^x + ... + b_m*(g(n-m)^c_m)^x)^(1/x) = g(n)^(1/x) where x is a real number > 0. - _Gerald McGarvey_, Dec 12 2004

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerald McGarvey_, Dec 04 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003480 | OEIS-A100666

a(n) = Sum_{k>=0} k^n/Catalan(k) rounded to nearest integer.

The exact values are conjectured to be close to integers, but there is no sound basis for it as yet. The coincidence for the first 10 instances is however intriguing. Conjecture is false: a(15) = round(547489559469.6423...), a(23) = round(1405230835254158841468.5247...), in fact, it seems values arbitrarily close to m+1/2 are possible. - _Sean A. Irvine_, Mar 07 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_, Dec 05 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003481 | OEIS-A100682

Floor of 4th root of pentatope numbers.

Conjecture: a(n) = floor((n - 3/2)/24^(1/4)) for n not in {0, 1, 6, 17, 2403, 5318}. - _Charles R Greathouse IV_, May 01 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Dec 06 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003482 | OEIS-A100691

Number of self-avoiding paths with n steps on a triangular lattice in the strip Z x {0,1}.

Conjecture: a(n) = A193641(n+2)-3, n>0 - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 07 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003483 | OEIS-A100826

a(1) = 1; a(n+1) = gcd(n,a(n))-th smallest positive integer not already in the sequence.

This is a permutation iff, infinitely often, gcd(n, a(n)) = 1. Through n = 1000, gcd(n, a(n)) = 1 612 times, which is consistent with the natural conjecture that this is true with density 6/Pi^2 ~ 0.6079.... - _Franklin T. Adams-Watters_, Nov 28 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Jan 06 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003484 | OEIS-A100833

Smallest positive palindrome-free and squarefree sequence.

Conjectures from _Robert Israel_, Jan 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jan 07 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003485 | OEIS-A100873

Pseudotwinprimes p+2 for primes p such that p+2 divides p^(p+2)+2 and p+2 is composite.

Conjecture 1: If p and p+2 are prime (twin primes), then p+2 divides p^(p+2)+2. Compared to the 1517 twin primes less than 130000, there were 33 pseudoprime occurrences. Conjecture 2: If for a randomly chosen prime p, p+2 divides p^(p+2)+2, then there is a greater than 98% chance that p and p+2 are twin primes. The sequence also contains several Carmichael numbers. In addition, If we relax the condition that p is prime or just odd, we get A001567: 341, 561, 645, 1105, 1387, 1729, 1905, 2047, ... (Sarrus numbers).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jan 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003486 | OEIS-A100949

Number of partitions of n into a prime and a semiprime.

Marnell conjectures that a(n) > 0 for n > 10 after analyzing "many thousands of whole numbers". I find no exceptions below 100 million. - _Charles R Greathouse IV_, May 04 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Nov 23 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003487 | OEIS-A100952

Numbers that cannot be written as p*q+r with three distinct primes p, q and r.

Conjecture: the sequence is complete. A weaker conjecture: every integer greater than 60 (or some larger value based on further search) may be partitioned into a prime p and a semiprime qr, where the prime p is bounded by log(min(q,r)). Chen (1978) showed that all sufficiently large even numbers are the sum of a prime and the product of at most two primes. Zumkeller's conjecture effectively extends this from "even" to both even and odd integers. - _Jonathan Vos Post_, Nov 25 2004 Conjecture: Every positive integer can be represented as p*q-r with distinct primes p, q, r. - _Zak Seidov_, Aug 28 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Nov 23 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003488 | OEIS-A100967

Least k such that binomial(2k + 1, k - n) >= binomial(2k, k).

Open question: Does binomial(2*a(n) + 1, a(n) - n) = binomial(2*a(n), a(n)) for any n? An affirmative answer would settle whether there exists an odd term greater than 3 in A003016. - _Danny Rorabaugh_, Mar 16 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Nov 23 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003489 | OEIS-A100982

Number of admissible sequences of order n; related to 3x+1 problem and Wagon's constant.

Conjecture: a(n) is given for each n > 3 by a formula using a(2)..a(n-1). This allows us to create an iterative algorithm which generates a(n) for each n > 6. This has been proved for each n <= 53. For higher values of n the algorithm must be slightly modified. - _Mike Winkler_, Jan 03 2018 a(n) = (m+n-2)!/(m!*(n-2)!) - Sum_{i=2..n-1} binomial(floor((3*(n-i)+b)/2), n-i)*a(i), where m = floor((n-1)*log_2(3))-(n-1) and b assumes different integer values within the sum at intervals of 5 or 6 terms. (Conjecture)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Finch_, Jan 13 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003490 | OEIS-A101035

Dirichlet inverse of the gcd-sum function (A018804).

Conjecture: Logarithmic g.f. Sum_{n>0,k>0} mu(n)*mu(k)*log(1/(1-x^(n*k))). - _Benedict W. J. Irwin_, Jul 26 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerard P. Michon_, Nov 27 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003491 | OEIS-A101036

Riesel numbers n (n*2^k-1 is composite for all k>0, n odd) that have a covering set.

Conjecture: there are infinitely many Riesel numbers that do not arise from a covering system. See page 16 of the Filaseta et al. reference. - _Arkadiusz Wesolowski_, Nov 17 2014 Conjecture: if R is a Riesel number (that has a covering set), then there exists a prime P such that R^p is also a Riesel number for every prime p > P. - _Thomas Ordowski_, Jul 12 2022 Problem: are there numbers K such that K + 2^m is a Riesel number for every m > 0? If so, then (K + 2^m)*2^n - 1 is composite for every pair of positive integers m,n. Also, by the dual Riesel conjecture, |K + 2^m - 2^n| are always composite. Note that, by the dual Riesel conjecture, if p is an odd prime and n is a positive integer, then there exists n such that (p + 2^m)*2^n - 1 is prime. So if such a number K exists, it must be composite. - _Thomas Ordowski_, Jul 20 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_, Jan 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003492 | OEIS-A101042

a(n) is the smallest positive d such that the n-th prime is the smallest prime p for which p+d is also prime.

Except for n=1, A020483(a(n)/2) is the first appearance of the n-th prime. It is conjectured that a(n) always exists. a(386) is the first number which must be above 10^12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jens Kruse Andersen_, Nov 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003493 | OEIS-A101044

Primes corresponding to A101043 (which is A101042 sorted).

The order in which the primes (except 2) first appear in A020483. It is conjectured that all primes are in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jens Kruse Andersen_, Nov 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003494 | OEIS-A101045

Record size primes in A101044.

Conjecture: lim_{n->infinity} a(n)/n^2 = 1. - _Ya-Ping Lu_, Sep 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jens Kruse Andersen_, Nov 28 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003495 | OEIS-A101183

Number of times (>0) function f must be applied in A100800 to arrive at multiple of n, or 0 if multiple of n is never reached.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ray Chandler_, Dec 19 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003496 | OEIS-A101201

Maximal number of kings in the toroidal king's graph on an n X n board such that each king is attacking no more than four other kings.

a(n) is approximately 3n^2/5 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eugen J. Ionascu_, Aug 12 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003497 | OEIS-A101208

Smallest odd prime p such that n = (p - 1) / ord_p(2).

Conjecture: a(n) is defined for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Leigh Ellison (le(AT)maths.gla.ac.uk), Dec 14 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003498 | OEIS-A101215

Numbers n such that concatenation of n with 54321 is a prime.

Conjecture 1 : for any integer N ending in 1 or 3 or 7 or 9 there is an infinite sequence of numbers n such that the concatenation nN is a prime. Conjecture 2 : for any integer N there is an infinite sequence of numbers n such that the concatenation Nn is a prime

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Dec 14 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003499 | OEIS-A101248

Decimal Goedelization of contingent WFFs (well-formed formulas) from propositional calculus, in Richard C. Schroeppel's metatheory of A101273. Truth value depends on truth value of variables, but is neither always true (theorem) nor always false (antitheorem).

As with A101273, I conjecture that a power law approximates the number of integers in this sequence, where the number with N digits is approximately N to the power of some real number D. The union of A101273, A100200 and this sequence is the set of all WFFs in Richard C. Schroeppel's metatheory of A101273.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jan 23 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003500 | OEIS-A101265

a(1) = 1, a(2) = 2, a(3) = 6; a(n) = 5*a(n-1) - 5*a(n-2) + a(n-3) for n > 3.

Conjecture: consecutive terms of this sequence and consecutive terms of A032908 provide all the positive integer solutions of (a+b)*(a+b+1) == 0 (mod (a*b)). - _Robert Israel_, Aug 26 2015 Conjecture is true: see StackExchange link. - _Robert Israel_, Sep 06 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | Lambert Klasen (lambert.klasen(AT)gmx.net) and _Gary W. Adamson_, Jan 25 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003501 | OEIS-A101282

Triangle read by rows: T(n,k) is the number of Schroeder paths of length 2n and having k valleys.

Conjecture: The n-th row polynomial is R(n+1,0) where

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003502 | OEIS-A101292

a(n) = n! + Sum_{i=1..n} i.

The only Fibonacci numbers in this sequence are 1, 2, 5, 34. [Ruiz and Luca, verifying a conjecture of Castillo] - _Eric M. Schmidt_, Nov 07 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Parthasarathy Nambi_, Dec 21 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003503 | OEIS-A101312

Number of "Friday the 13ths" in year n (starting at 1901).

Conjecture: The same basic repeating pattern results if we seek the number of "Sunday the 22nds" or "Wednesday the 8ths" or anything else similar, with the only difference being that the sequence starts at a different point in the repeating pattern.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Adam M. Kalman (mocha(AT)clarityconnect.com), Dec 22 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003504 | OEIS-A101348

Arises from a particular cyclic transformation of the floretion - .5'i - .5i' - .5'ij' - .5'ik' + .5'ji' + .5'ki'.

Conjectures from _Colin Barker_, Apr 30 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Creighton Dement_, Dec 25 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003505 | OEIS-A101349

Numbers of cubes between prime(n) and prime(n+1).

a(n) <= 1 for all n if and only if there is a prime between k^3 and (k+1)^3 for every k >= 1, which is an open conjecture. It has been verified by computation for k^3 <= 1.649*10^40 (Johnston, Sorenson, Thomas and Webster), so a(n) <= 1 whenever prime(n+1) <= 1.6*10^40; analytically it is known only for k >= exp(exp(32.537)) (Cully-Hugill). - _Pierre Colmant_, Oct 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Teofilatto_, Dec 25 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003506 | OEIS-A101363

In the interior of a regular 2n-gon with all diagonals drawn, the number of points where exactly three diagonals intersect.

I checked the following conjecture up to n=210: "An n-gon with n=30k has 5n points where 6 or 7 diagonals meet and no points where more than 7 diagonals meet; If k is odd, then 6 diagonals meet in each of 4n points and 7 diagonals meet in each of n points; If k is even, then no groups of exactly 6 diagonals meet in a point, while exactly 7 diagonals meet in each of 5n points."

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Graeme McRae_, Dec 26 2004, revised Feb 23 2008, Feb 26 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003507 | OEIS-A101364

In the interior of a regular n-gon with all diagonals drawn, the number of points where exactly four diagonals intersect.

I checked the following conjecture up to n=210: "An n-gon with n=30k has 5n points where 6 or 7 diagonals meet and no interior point other than the center where more than 7 diagonals meet; If k is odd, then 6 diagonals meet in each of 4n points and 7 diagonals meet in each of n points; If k is even, then no groups of exactly 6 diagonals meet in a point, while exactly 7 diagonals meet in each of 5n points (all points interior excluding the center)."

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Graeme McRae_, Dec 26 2004, revised Feb 23 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003508 | OEIS-A101368

The sequence solves the following problem: find all the pairs (i,j) such that i divides 1+j+j^2 and j divides 1+i+i^2. In fact, the pairs (a(n),a(n+1)), n>0, are all the solutions.

G.f.: x(1 - 5x + 3x^2) / [(1-x)(1 - 5x + x^2)]; a(n) = 2 * A089817(n-3) + 1, n>2. - Conjectured by _Ralf Stephan_, Jan 14 2005, proved by _Max Alekseyev_, Aug 03 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | M. Benito, O. Ciaurri and E. Fernandez (oscar.ciaurri(AT)dmc.unirioja.es), Jan 13 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003509 | OEIS-A101424

Number of different cuboids with volume p^4 * q^n, where p,q are distinct prime numbers.

The following is conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anthony C Robin_, Jan 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003510 | OEIS-A101425

Number of different cuboids with volume p^5 X q^n, where p,q are distinct prime numbers.

The following is conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anthony C Robin_, Jan 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003511 | OEIS-A101462

Smallest k such that 2^k-prime(n) is prime.

Conjecture: sequence is defined for all n. First unproved n: 286 Prime(286)=1871, up to date, tested up to k=40959, none 2^k-Prime(286) is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Jan 20 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003512 | OEIS-A101478

G.f. satisfies A(x) = x*(1+A(x))^4/(1+A(x)^2).

Conjecture: n*(n-1)*(n+1)*a(n) -12*n*(n-1)*(2*n-3)*a(n-1) +12*(n-1)*(16*n^2-64*n+65)*a(n-2) -16*(2*n-5)*(4*n-9)*(4*n-11)*a(n-3)=0. - _R. J. Mathar_, Nov 10 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Jan 21 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003513 | OEIS-A101481

Column 0 of triangular matrix T=A101479, in which row n equals row (n-1) of T^(n-1) followed by '1'.

Conjecture: a(n) is described by a series of nested sums,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 21 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003514 | OEIS-A101500

A Chebyshev transform of the central binomial numbers.

Conjecture: n*a(n) +2*(-2*n+1)*a(n-1) +2*(n-1)*a(n-2) +2*(-2*n+3)*a(n-3) +(n-2)*a(n-4)=0. - _R. J. Mathar_, Nov 16 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Dec 04 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003515 | OEIS-A101513

a(1) = 1, a(2) = 2, a(3) = 3; triangle where n-th row has lowest n positive integers not yet in the sequence such that each integer has a prime divisor in common with at least one element of the (n-1)th row.

Conjectures from _N. J. A. Sloane_, Apr 22 2005: (Start) "Call a number "postponed" if it cannot be placed right away, that is, if it is relatively prime to the numbers in the previous row. Then I conjecture that:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Jan 25 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003516 | OEIS-A101591

Indices k >= 8 designating the k-Somos sequences that first become nonintegral at odd-indexed terms (i.e., positions of odd terms in A030127).

When k is in this sequence, the k-Somos sequence first becomes nonintegral at 2*k + 1; otherwise, the k-Somos sequence first becomes nonintegral at 2*k (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 08 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003517 | OEIS-A101596

Expansion of g.f. c(2*x)^4, where c(x) is the g.f. of A000108.

Conjecture: (n+4)*a(n) - 4*(3n+7)*a(n-1) + 16*(2n+1)*a(n-2) = 0. - _R. J. Mathar_, Dec 13 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Dec 08 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003518 | OEIS-A101601

Expansion of g.f. c(3x)^3, where c(x) is the g.f. of A000108.

Conjecture: (n+3)*a(n) - 3*(5*n+7)*a(n-1) + 18*(2*n-1)*a(n-2) = 0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Dec 08 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003519 | OEIS-A101602

Expansion of g.f. c(3x)^4, where c(x) is the g.f. of A000108.

Conjecture: (n+4)*a(n) - 6*(3*n+7)*a(n-1) + 36*(2*n+1)*a(n-2) = 0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Dec 08 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003520 | OEIS-A101695

a(n) = n-th n-almost prime.

Conjecture: lim_{ n->inf.} a(n+1)/a(n) = 2. - _Robert G. Wilson v_, Oct 07 2007, Nov 13 2007 Stronger conjecture: a(n)/(n * 2^n) is polylogarithmic in n. That is, there exist real numbers b < c such that (log n)^b < a(n)/(n * 2^n) < (log n)^c for large enough n. Probably b and c can be chosen close to 0. - _Charles R Greathouse IV_, Aug 28 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Dec 12 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003521 | OEIS-A101900

Column 0 of triangular matrix A101897, in which column k equals column 0 of A101897^(k+1) and where the n-th row sums are zero for n>0.

|a(n)| is the number of n-length permutations avoiding the dotted pattern 23\dot{1}(conjectured). - _Michael D. Weiner_, Jul 26 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003522 | OEIS-A101902

Numbers n that are not of the form ab+bc+cd+de+ea with 1<=a<=b<=c<=d<=e.

Conjecture that this sequence is complete. Note that, except for n=1 and n=2, n-1 is prime. The case of the 3-term binary form is treated in A025052. For the 4-term case, ab+bc+cd+da, no prime is representable because the 4 terms factor as (a+c)(b+d).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Dec 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003523 | OEIS-A101914

G.f. satisfies: A(x) = 1/(1 + x*A(x^4)) and also the continued fraction: 1 + x*A(x^5) = [1; 1/x, 1/x^4, 1/x^16, 1/x^64, ..., 1/x^(4^(n-1)), ...].

The conjecture is wrong. The g.f. is dependent on the number of terms. With twenty terms the g.f. is (1 + x^4)/(1 + x + x^4). - _Johannes W. Meijer_, Aug 08 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003524 | OEIS-A101917

G.f. satisfies: A(x) = 1/(1 + x*A(x^7)) and also the continued fraction: 1 + x*A(x^8) = [1; 1/x, 1/x^7, 1/x^49, 1/x^343, ..., 1/x^(7^(n-1)), ...].

This was conjectured to have g.f. (1+x^7) / (1+x+x^7) by _Ralf Stephan_, May 17 2007, but this is wrong. This g.f. produces a sequence which differs at a(57) = -153367. The g.f. gives a(57) = -153366. - _Johannes W. Meijer_, Aug 08 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003525 | OEIS-A101918

G.f. satisfies: A(x) = 1/(1 + x*A(x^8)) and also the continued fraction: 1 + x*A(x^9) = [1; 1/x, 1/x^8, 1/x^64, 1/x^512, ..., 1/x^(8^(n-1)), ...].

The conjecture that this has g.f. (1+x^8) / (1+x+x^8) by _Ralf Stephan_, May 17 2007 is wrong. The first difference occurs at a(73) = -42106. The g.f. gives a(73) = -42105. - _Johannes W. Meijer_, Aug 08 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 20 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003526 | OEIS-A101992

Numerator of Sum_{i=2..n} (-1)^i/(i*phi(i)).

I conjecture that there exists a limit for Sum_{i>=2} (-1)^i/(i*phi(i)) which is ca. 0.558.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Orges Leka_, Dec 23 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003527 | OEIS-A102086

Triangular matrix, read by rows, that satisfies: T(n,k) = [T^2](n-1,k) when n>k>=0, with T(n,n) = (n+1).

Conjecture: n-th row polynomial is P(n+1, n) where P(n, k) = P(n, k-1) + (k+1) * P(n-1, k) for 0 < k < n with P(n, n) = x^n, P(n, 0) = 1, P(n, k) = 0 for k < 0 or k > n. - _Mikhail Kurkov_, Jul 03 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003528 | OEIS-A102091

Number of perfect matchings in the C_{2n} X P_3 graph (C_{2n} is the cycle graph on 2n vertices and P_3 is the path graph on 3 vertices).

Conjecture: For an integer n > 6, primes of the form either 12k+5 or 12k+7 will never appear in factors of any centered n-gonal number, if and only if n is in A102091. - _Steven Lu_, Nov 15 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 29 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003529 | OEIS-A102167

Lightest finite strictly increasing sequence of integers showing once and only once every pair of digits from [00] to [99]. The "weight" of such a sequence is the sum of all its a(n). Weight of this sequence is 2995.

It seems that it is only a conjecture that this sequence is really the lightest! - _N. J. A. Sloane_, Mar 15 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | Geoff Bailey, Richard Tucker, "Alex" and _Eric Angelini_, Feb 16 2005; revised Mar 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003530 | OEIS-A102188

a(n) = Sum_{m=0..n} (-1)^m * binomial(n,m)*(1*3*5*...*(4m-1)).

Comment from Kevin Buzzard (k.buzzard(AT)imperial.ac.uk), Feb 17 2005: 2^k exactly divides f(k). Applying the theory of Wilf and Zeilberger to this problem gives a one-line proof that 16*(k+1)*(k+2)*f(k) - 32*(k+2)^2*f(k+1) + (16k^2 + 80k + 98)*f(k+2) + f(k+3) = 0 for all k >= 0, from which the conjecture follows easily (check for the first few terms and then easy induction on k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003531 | OEIS-A102237

Smallest number equal to the product of n primes which is also equal to the sum of n distinct primes.

Conjecture: a(n) = 2^n for n>5. - _Ray Chandler_ Feb 25 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Teofilatto_, Feb 18 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003532 | OEIS-A102239

a(n) = (Sum_{i=0..n} 5^i) + 1 - (Sum_{i=0..n} 5^i) mod 2.

a(n) = 4*a(n-1) + 5*a(n-2) - 2 (conjecture). - _Creighton Dement_, Apr 13 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Mar 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003533 | OEIS-A102297

Number of distinct divisors of n+1 where n and n+1 are composite or twin composite numbers.

It is interesting to note that the first such consecutive pair of composite numbers is 8 and 9 which are perfect powers: 2^3 and 3^2. Conjecture: 8 and 9 are the only 2 consecutive composite numbers that are both perfect powers. Or, if x>2, x^m+1 != y^n for all m,n,x,y. Now if we relax the condition that 0 and 1 are not composite, we have 0^m+1 = 1^n for all m,n an infinity of solutions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Feb 19 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003534 | OEIS-A102309

a(n) = Sum_{d divides n} moebius(d) * binomial(n/d,2).

The conjectured relation to A100448 is true (see Fried link). - _Sela Fried_, Mar 15 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Jan 03 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003535 | OEIS-A102318

A mean binomial transform of the Catalan numbers.

Conjecture: -(n-1)*(n+1)*a(n) +2*(5*n^2-9*n+1)*a(n-1) +2*(-15*n^2+58*n-49)*a(n-2) +2*(10*n^2-76*n+123)*a(n-3) +(31*n-55)*(n-3)*a(n-4) -30*(n-3)*(n-4)*a(n-5)=0. - _R. J. Mathar_, Jun 08 2016
Conjecture: +(3*n-10)*(n-1)*(n+1)*a(n) +2*(-12*n^3+58*n^2-67*n+10)*a(n-1) +2*(21*n^3-136*n^2+289*n-196)*a(n-2) +2*(n-2)*(12*n^2-46*n+27)*a(n-3) -15*(n-2)*(n-3)*(3*n-7)*a(n-4)=0. - _R. J. Mathar_, Jun 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003536 | OEIS-A102319

A mean binomial transform of the central binomial numbers.

Conjecture: n*(n-1)*a(n) -4*(n-1)*(3*n-4)*a(n-1) +(53*n^2-221*n+232)*a(n-2) +8*(-13*n^2+85*n-134)*a(n-3) +(51*n^2-563*n+1308)*a(n-4) +4*(29*n-93)*(n-4)*a(n-5) -105*(n-4)*(n-5)*a(n-6)=0. - _R. J. Mathar_, Feb 20 2015
Conjecture:+n*(n-1)*(12*n^2-48*n+41)*a(n) -8*(n-1)*(12*n^3-54*n^2+65*n-17)*a(n-1) +2*(84*n^4-504*n^3+1025*n^2-775*n+131)*a(n-2) +8*(n-2)*(12*n^3-54*n^2+65*n-20)*a(n-3) -15*(n-2)*(n-3)*(12*n^2-24*n+5)*a(n-4)=0. - _R. J. Mathar_, Feb 20 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003537 | OEIS-A102341

Areas of 'close-to-equilateral' integer triangles.

Then, the next four terms are {three sides a<=b<=c and area}: {46816, 46817, 46817, 949077360}, {174725, 174725, 174726, 13219419708}, {652080, 652081, 652081, 184120982760}, {2433601, 2433601, 2433602, 2564481115560}. Also, if we allow degenerate triangles (area 0), the first case would be {1,1,2,0}. We have 12 cases and a weak conjecture is that the total number of the

'close-to-equilateral' triangles is finite. - _Zak Seidov_, Feb 23 2005 (2/3) [A007655(n+2) - (-1)^n*A001353(n+1)] (conjectured). - _Ralf Stephan_, May 17 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | Johannes Koelman (Joc_Kay(AT)hotmail.com), Feb 20 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003538 | OEIS-A102395

A mod 2 related Jacobsthal sequence.

Conjecture: all the terms are in A078008. The conjecture is true since a(n) = A078008(A000120(n)). - _Paul Barry_, Jan 07 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 06 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003539 | OEIS-A102483

Numbers k such that 2^k contains no zeros in base 3.

I conjectured in 1973 that there are no further terms. This question is still open.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 25 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003540 | OEIS-A102612

First nonzero terms in the sequences formed by the unique count of primes between an and (a+1)n.

Care should be taken to choose a sufficiently large n for a given m range for a in the script below. (50,1000) -> 22,11,18,62,59,29,12,47,106,40,17,104,94,44,83,22,.. (50,2000) -> 22,11,18,62,59,29,12,47,106,40,17,104,94,44,83,180,.. Notice the breakdown at the end. While the terms in the sequence tend to oscillate increasing, strange things are possible when more terms are listed. Conjecture: The number of terms in this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jan 30 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003541 | OEIS-A102751

Numbers k such that 1 + (k-1)^2 and ((k-1)/2)^2 + ((k+1)/2)^2 = (1/2)*(k^2+1) are primes.

Conjectured to be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robin Garcia_, Feb 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003542 | OEIS-A102753

Decimal expansion of (Pi^2)/2.

Conjecture: Pi^2/2 = Sum_{n = -oo..oo} (cos(Pi*sqrt(n^2+1)) - cos(Pi*n)) (using the Eisenstein summation convention). - _Peter Bala_, Oct 08 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jun Mizuki (suzuki32(AT)sanken.osaka-u.ac.jp), Feb 10 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003543 | OEIS-A102818

Irregular array a(m,k) = A001035(k) mod m read by rows, 1<=k<=16, 3<=m.

Conjectures (based on mod values up to n=99): the sequence A001035(m) is (pre)periodic modulo n for all n, the lengths of the ending periods mod n (except n=4) being given by A011773 (which is related to Carmichael's lambda function).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerald McGarvey_, Feb 26 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003544 | OEIS-A102820

Number of primes between 2*prime(n) and 2*prime(n+1), where prime(n) is the n-th prime.

Conjecture: the asymptotic proportions of 0's, 1's, ... , k's, ... are 1/3, 2/9, ... , 2^k/3^(k+1), ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ali A. Tanara (tanara(AT)khayam.ut.ac.ir), Feb 27 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003545 | OEIS-A102827

"True already", base 10, start 1: a(n) is the least integer such that the sequence up to a(n-1) written in base 10 contains floor(a(n)/10) copies of the digit a(n) % 10, with a(0) = 1.

Conjecture: this sequence in various bases never includes a term divisible by the base.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo van der Sanden_, Feb 26 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003546 | OEIS-A102842

Insipid numbers: n is defined to be insipid if "G is a primitive subgroup of the symmetric group S_n" implies that "G=A_n or G=S_n".

It seems hard to find long gaps in this sequence. It seems plausible (this is implied by some conjectures in number theory) that there are infinitely many strings of 5 consecutive positive integers not in this sequence; however, I do not know of a construction which should yield infinitely many strings of 6 consecutive positive integers which are in the sequence (this may be just a reflection of my ignorance of the right families of finite groups); the largest example I know of a string of more than 5 consecutive integers not in this sequence has length 7 and first term 2^150-5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David L. Harden_, Feb 27 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003547 | OEIS-A102879

A Chebyshev transform of the first kind of the central binomial numbers.

Conjecture: n*(n-3)*a(n) - 2*(2*n-1)*(n-3)*a(n-1) + 2*(2-4*n+n^2)*a(n-2) - 2*(n-1)*(2*n-7)*a(n-3) + (n-1)*(n-4)*a(n-4) = 0. - _R. J. Mathar_, Nov 09 2012 Conjecture verified using the differential equation x*(x^2+1)*(x^2-4*x+1)*g'' + (4*x^4-10*x^3+2*x^2+2*x-2)*g' + 4*(x^2-x+1)*g = 0 satisfied by the g.f. - _Robert Israel_, Aug 28 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003548 | OEIS-A102880

A Chebyshev transform of the first kind of the Catalan numbers.

Conjecture: (n+1)*(n-3)*a(n) -2*(2*n-1)*(n-3)*a(n-1) +2*(1-4*n+n^2)*a(n-2) -2*(n-1)*(2*n-7)*a(n-3) +(n-1)*(n-5)*a(n-4)=0. - _R. J. Mathar_, Nov 09 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003549 | OEIS-A102882

Expansion of (1+2x)/sqrt((1-3x^2)(1+4x+5x^2)).

Conjecture: n*(n+1)*a(n) +2*(2*n+3)*(n-1)*a(n-1) +2*(n^2-7)*a(n-2) +6*(5+n-2*n^2)*a(n-3) -15*(n+2)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Nov 09 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003550 | OEIS-A103117

Berend Jan van der Zwaag's conjectured complete list of numbers that start different "expanding periodic loops" under the mapping described in A053392 and A060630.

Berend Jan van der Zwaag's conjectured complete list of numbers that start different "expanding periodic loops" under the mapping described in A053392 and A060630.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 06 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003551 | OEIS-A103141

Riordan array (1/(1-x), x*(1 + x + x^2 + x^3)/(1-x)).

T(n,k) = Sum_{i=0..n-k} binomial(n-i,k) * (Sum_{j=0..i} binomial(k,j) * binomial(3*k-2*j,i-j) * (-2)^j) for 0 <= k <= n (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003552 | OEIS-A103152

Smallest odd number which can be written as a sum 2p+q (where p and q are both odd primes, A065091) in exactly n ways, zero if there are no such odd number.

Conjecture: except for the 2nd, 3rd and 4th terms, all other terms are divisible by 3; See also comments in A103151.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Feb 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003553 | OEIS-A103176

Let p = prime(sigma(n)) and q = prime(phi(n)), then p is in the sequence if p-q = 6.

Conjecture: In all cases sigma(n)-phi(n)=2, i.e., n is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Mar 02 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003554 | OEIS-A103225

Number of Gaussian integers z with abs(z) < n and gcd(n,z)=1.

Four of the dominant lines of the plot appear to align to k(i)*Pi*n^2, with k(i) = 1, 8/9, 1/2, and 4/9. Conjecture: a(n) < Pi*n^2. - _Bill McEachen_, Aug 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jan 26 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003555 | OEIS-A103267

Positive integers x such that there exist positive integers y and z satisfying y^3 - x^3 = z^4.

Conjecture: The greatest prime divisor of x divides y and z. Conjecture is false: 126^4 = 639^3 - 207^3, the largest prime divisor of 207 is 23 and neither 126 nor 639 is divisible by 23. For max(x,y) < 4.3*10^9, there are no additional terms < 2440. It is likely this is true for all x,y. - _Chai Wah Wu_, Jan 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Mar 20 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003556 | OEIS-A103274

Number of ways of writing prime(n) in the form 2*prime(i)+prime(j).

Marnell conjectures that a(n) > 0 for n > 3. I find no exceptions below 10^9. - _Charles R Greathouse IV_, May 04 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yasutoshi Kohmoto_, Jan 27 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003557 | OEIS-A103287

Main diagonal of triangle A103286, in which row n+1 is formed by sorting, in ascending order, the result of the convolution of row n with {2,1}.

a(n) ~ 3^n*log_3(n)/(18n) (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 28 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003558 | OEIS-A103288

Numbers k such that sigma(k) >= 2k-1 (union of perfect, abundant and least deficient numbers).

If the only least deficient numbers are the powers of 2 (open problem) then this sequence is the union of A023196 and A000079.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Jan 28 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003559 | OEIS-A103362

Reduced numerators of the fraction of primes < 10^n that are full reptend primes.

A103362/A103363 is conjectured to approach Artin's constant A005596.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Feb 02 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003560 | OEIS-A103363

Reduced denominators of the fraction of primes < 10^n that are full reptend primes.

A103362/A103363 is conjectured to approach Artin's constant A005596.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Feb 02 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003561 | OEIS-A103469

Number of polyominoes consisting of 3 regular unit n-gons.

Conjecture: if n > 3, then a(n + 3) is the number of connected components of the subgraph that is vertex-induced on Collatz's graph by the vertex subset {1, ..., n} (see Problem 3.11 of my article, available from the links). - _Lorenzo Sauras Altuzarra_, Apr 07 2020 [Corrected by _Pontus von Brömssen_, Jan 22 2021]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sascha Kurz_, Feb 07 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003562 | OEIS-A103528

a(n) = Sum_{k = 1..n-1 such that n == k (mod 2^k)} 2^(k-1).

Conjecture: a(n) = (b(n) - b(n-1) - 1)/2 for n > 1 where b(n) = Sum_{k=0..A000523(n)} c(n-k, k) and c(n, m) = n - (n mod 2^m) with a(1) = 0. - _Mikhail Kurkov_, Jun 01 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 22 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003563 | OEIS-A103621

Trajectory of 7 under repeated application of the map n -> A102370(n).

Conjectures from _Chai Wah Wu_, Feb 01 2018: (Start) The above conjectures are incompatible with A102370(2^37-37) = 2^38-3. - _Peter Munn_, Jan 13 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Mar 31 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003564 | OEIS-A103627

Let S(n) = {n,1,n}; sequence gives concatenation S(0), S(1), S(2), ...

Conjecture: a(n) = (2*n+1 + (2*n-8)*cos((2*n+1)*Pi/3) + sqrt(3)*cos((8*n+1)*Pi/6) + sqrt(3)*sin((2*n+1)*Pi/3))/9. - _Wesley Ivan Hurt_, Sep 25 2017 Conjectures from _Colin Barker_, Sep 27 2017: (Start) a(3*k+1) = 1, a(3*k) = a(3*k+2) = k. The conjectures easily follow from that. - _Robert Israel_, Jan 11 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Mar 25 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003565 | OEIS-A103663

Smallest integer base x > 1 such that x^n has no digit 0 in its decimal representation, or 0 if no such x exists.

Zero values are conjectural. a(40) was checked up to 10^9 by _Joshua Zucker_. All other zero values were checked up to 10^5 by _David Wasserman_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Feb 28 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003566 | OEIS-A103667

Primes p such that the largest prime divisor of p-1 is greater than the largest prime divisor of p+1.

Conjecture: this sequence is of positive relative density in the primes, perhaps even 1/2. - _Charles R Greathouse IV_, Dec 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Feb 19 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003567 | OEIS-A103670

Smallest m such that A102730(m) = n.

Reinhard Zumkeller conjectures (at A102730) that this sequence is finite. I conjecture the contrary, that a(n) exists for every n. Further, I expect a(n) << n^n. - _Charles R Greathouse IV_, Aug 21 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Feb 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003568 | OEIS-A103671

Smallest m such that the binary representation of n! does not contain m!.

Reinhard Zumkeller conjectures (at A102730) that this sequence is bounded. I conjecture the contrary, that for every k there is n with a(n) > k. - _Charles R Greathouse IV_, Apr 07 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Feb 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003569 | OEIS-A103673

If in binary representation n! contains 5! then 1 else 0.

Conjecture: a(n) = 1 for n > 65. - _Charles R Greathouse IV_, Apr 07 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Feb 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003570 | OEIS-A103674

If in binary representation n! contains 6! then 1 else 0.

Conjecture: a(n) = 1 for n > 802. - _Charles R Greathouse IV_, Apr 07 2013 Conjecture checked up to n <= 5*10^5. - _Giovanni Resta_, Apr 07 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Feb 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003571 | OEIS-A103677

Numbers m such that in binary representation m! doesn't contain 5!.

Conjecture: there are no further terms;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Feb 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003572 | OEIS-A103683

a(1)=1, a(2)=2, a(3)=3, a(n) = smallest positive integer not occurring earlier in sequence and coprime to a(n-1), a(n-2) and a(n-3).

Conjectured to be not a permutation of the natural numbers. Conjecture: for n >= 67, a(n) is even if and only if n == 2 (mod 4) and divisible by 3 if and only if n == 3 (mod 4). In particular, this implies the last value divisible by 6 is a(66) = 36. - _Robert Israel_, May 12 2015 a(102982) = 42, see A105214. Conjecture above is false. - _Sergio Pimentel_, Apr 18 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Mar 26 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003573 | OEIS-A103769

Trinomial transform of central binomial coefficients A001405.

Conjecture: n*(2n+1)*a(n) +2(-61n^2+57n-20)*a(n-1) +3*(205n^2-523*n+346) * a(n-2) -72*(n-2)*(16n-33)*a(n-3) +567*(n-2)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Dec 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003574 | OEIS-A103777

Numbers k such that f(k), f(k+1) and f(k+2) are all primes, where f(k) = 8*k^2 + 4*k + 1.

All terms are divisible by 5, hence conjecture: there is no such k that f(k), f(k+1),f(k+2) and f(k+3) are primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Feb 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003575 | OEIS-A103784

Minimal n that makes primorial P(k)*n-1 prime, k>=2, n>0.

Minimal n of sequence A103783. Weak conjecture: sequence is defined for all k>=2; strong conjecture: a(k)<=(prime(k))^2;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Feb 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003576 | OEIS-A103786

a(n) is the minimum k that makes primorial P(n)/A019565(k)+A019565(k) prime, k>=0, n>0.

This is the k value of A103785. Conjecture: sequence is defined for all n>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Feb 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003577 | OEIS-A103794

Smallest number b such that b^prime(n) - (b-1)^prime(n) is prime.

Conjecture: sequence is defined for all positive indices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Feb 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003578 | OEIS-A103795

Minimal base b such that (b^prime(n)+1)/(b+1) is prime.

Conjecture: sequence is defined for any n>=2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Feb 23 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003579 | OEIS-A103821

A Whitney transform of the central binomial coefficients A000984.

Conjecture: n*a(n) + (2-5*n)*a(n-1) + 2*a(n-2) + 4*(n-1)*a(n-3) = 0. - _R. J. Mathar_, Dec 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003580 | OEIS-A103828

Sequence of odd numbers defined recursively by: a(1)=1 and a(n) is the first odd number greater than a(n-1) such that a(n) + a(i) + 1 is prime for 1<=i<=n-1.

The Hardy-Littlewood k-tuple conjecture would imply that this sequence is infinite. Note that, for n>2, a(n)+2 and a(n)+4 are both primes, so a proof that this sequence is infinite would also show that there are infinitely many twin primes. - _N. J. A. Sloane_, Apr 21 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Walter Kehowski_, May 29 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003581 | OEIS-A103905

Square array T(n,k) read by antidiagonals: number of tilings of an <n,k,n> hexagon.

Conjectural formula as a sum of squares of Vandermonde determinants: T(n,k) = 1/((1!*2! ... *(n-1)!)^2*n!)* Sum_{1 <= x_1, ..., x_n <= k} (det V(x_1, ..., x_n))^2, where V(x_1, ..., x_n) is the Vandermonde matrix of order n. Compare with A133112. - _Peter Bala_, Sep 18 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Feb 22 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003582 | OEIS-A103910

Triangle T read by rows: inverse of fibonomial triangle (A010048).

Conjecture: all rows except the first add to zero. See below. Conjecture: T(n+k, n) = A010048(n+k-1, k)*T(k, 1), n>1. See below. It follows that the n-th row polynomial is Sum_{k = 0..n} T(k, 0)*A010048(n, k)*x^(n-k) and that for n > 0, the row sums are zero. See the above conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Feb 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003583 | OEIS-A103973

Expansion of (sqrt(1-8*x^2)+8*x^2+2*x-1)/(2*x*sqrt(1-8*x^2)).

Conjecture D-finite with recurrence: 11*n*(n+1)*a(n)+4*n*(4*n+1)*a(n-1) +8*(27-11*n^2)*a(n-2) -32*(4*n+9)*(n-3)*a(n-3)=0. - _R. J. Mathar_, Nov 09 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 23 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003584 | OEIS-A103974

Smaller sides (a) in (a,a,a+1)-integer triangle with integer area.

Equivalently, positive integers a such that 3/16*a^4 + 1/4*a^3 - 1/8*a^2 - 1/4*a - 1/16 is a square (A000290), a direct result of Heron's formula. Conjecture: lim_{n->oo} a(n+1)/a(n) = 7 + 4*sqrt(3) (= 7 + A010502). - _Rick L. Shepherd_, Sep 04 2005 "a(n)^2 = A011922(n)^2 + (4*A007655(n))^2, so that A011922(n) = 1/2 base of triangles, A007655(n) = 1/4 height of triangles (conjectured by Paul Hanna).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Feb 23 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003585 | OEIS-A103990

Reduced numerators of the hypercube line-picking integrand sqrt(Pi)*I(n,0).

Conjecture: Let b(n) be A103991(n), then a(n)/b(n) = (2/3)*(n^3-n^2+2*n+1)/(2*n+1). - _Gerald McGarvey_, Aug 11 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Feb 23 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003586 | OEIS-A103995

First column of triangle A103910.

Recurrence: a(n) = -Sum[i=1..n-1, a(i)*A010048(n-1, i-1)] (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Feb 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003587 | OEIS-A104007

Denominators of coefficients in expansion of x^2*(1-exp(-2*x))^(-2).

It appears that a(2n+2) = A002432(n). As A098087(n)/A104007(n) = x*(csch(x)^4)/(4*(coth(x)-1)^2), then a(2n+2) would represent the sequence of denominators for just the even powers of the full series representation at x=0. A002432 could be conjectured to be the non-hyperbolic, or circle trigonometric, function equivalent where the full series of denominators could be found by the formula x*((csc(x)^2)/4) - cot(x)/2) + 1 for a(n) > 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003588 | OEIS-A104016

Devaraj numbers: squarefree r-prime-factor (r>1) integers N=p1*...*pr such that phi(N)=(p1-1)*...*(pr-1) divides gcd(p1-1,...,pr-1)^2*(N-1)^(r-2).

A.K. Devaraj conjectured that these numbers are exactly Carmichael numbers. It was proved (see Alekseyev link) that every Carmichael number is indeed a Devaraj number, but the converse is not true. Devaraj numbers that are not Carmichael are given by A104017.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Feb 25 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003589 | OEIS-A104041

Triangular matrix T, read by rows, such that column k is equal (in absolute value) to row (k-1) in the matrix inverse T^-1 (with extrapolated zeros) for k>0, with T(n,n)=1 (n>=0) and T(n,n-1)=-n (n>=1).

Conjectures from _Peter Bala_, May 25 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 02 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003590 | OEIS-A104102

Number of primes >= p(n)# and < p(n)# + p(n+1)^2 where p(i)=i-th prime, p(i)#=i-th primorial.

Conjecture: a(n) is always > p(n+1)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Mar 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003591 | OEIS-A104109

Numbers n such that whenever a group G has a solvable subgroup of index n, then G itself is solvable.

As for the primes in the sequence, a theorem proved by Burnside guarantees that any unsolvable permutation group on a prime number of points is doubly transitive. The classification of doubly transitive permutation groups has been completed and it can be used to show that the only primes not in this sequence are 7, 13 and the Fermat primes larger than 3. So assuming the standard conjecture that 65537 is the largest Fermat prime, all primes except for these six (5, 7, 13, 17, 257 and 65537) are in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David L. Harden_, Mar 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003592 | OEIS-A104117

For n=2^k, a(n) = k+1, else 0.

Row sums of A103994 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Apr 15 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003593 | OEIS-A104148

Difference between n^3 and largest m less than n^3, where m > 0 is either a prime or a square.

a(n) < 2n^(3/2) - 1. Conjecturally a(n) << log^2 n. - _Charles R Greathouse IV_, Jun 03 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Teofilatto_, Mar 08 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003594 | OEIS-A104157

Smallest of n^2 consecutive primes that form an n X n magic square with the least magic constant, or 0 if no such magic square exists.

Conjecture: for n > 4, a(n) = prime(s) where s > 1 is the smallest integer such that (Sum_{i=s..s+n^2-1} prime(i))/n is an integer of the same parity as n. - _Max Alekseyev_, Jan 29 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Mar 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003595 | OEIS-A104169

Highly decomposable Smith numbers. A Smith number which sets a record for the number of prime factors (counting multiplicity) starting from first Smith number is called a highly decomposable Smith number.

It is conjectured that except for 27, all highly decomposable Smith numbers are even.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shyam Sunder Gupta_, Mar 10 2005 and May 03 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003596 | OEIS-A104272

Ramanujan primes R_n: a(n) is the smallest number such that if x >= a(n), then pi(x) - pi(x/2) >= n, where pi(x) is the number of primes <= x.

If n tends to infinity, then limsup(a(n) - A080359(n-1)) = infinity; conjecture: also limsup(a(n) - A080359(n)) = infinity (cf. A182366). - _Vladimir Shevelev_, Apr 27 2012 Sondow, Nicholson, and Noe's 2011 conjecture that pi(R_{m*n}) <= m*pi(R_n) for m >= 1 and n >= N_m (see A190413, A190414) was proved for n > 10^300 by Shichun Yang and Alain Togbé in 2015. - _Jonathan Sondow_, Dec 01 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Feb 27 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003597 | OEIS-A104310

Maximum length of perfect rulers that can be made from segments not exceeding n.

We conjecture the extension a(8)=81. For nomenclature related to perfect and optimal rulers see Peter Luschny's "Perfect Rulers" web pages.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Mar 01 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003598 | OEIS-A104385

Number of distinct partitions of triangular numbers n*(n+1)/2 into 3 parts for n>=1.

a(n) = round((3n^4 + 6n^3 - 33n^2 - 36n + 110)/144) (conjectured). - _Ralf Stephan_, Nov 13 2010.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003599 | OEIS-A104624

Expansion of exp(arcsinh(-2*x)) in powers of x.

Conjecture: n*a(n) + (n-1)*a(n-1) + 4*(n-3)*a(n-2) + 4*(n-4)*a(n-3) = 0. - _R. J. Mathar_, Nov 15 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003600 | OEIS-A104625

Expansion of 1/(sqrt(1-4*x) - x^2).

Conjecture: n*a(n) + (n-3)*a(n-1) + 2*(-28*n+51)*a(n-2) + 72*(2*n-5)*a(n-3) - n*a(n-4) + (-5*n+3)*a(n-5) + 18*(2*n-5)*a(n-6) = 0. - _R. J. Mathar_, Feb 20 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003601 | OEIS-A104638

Number of odd digits in n-th prime.

On the other hand, a(n) = 2 if prime(n) is in A159352, which is conjectured to be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Mar 18 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003602 | OEIS-A104722

Self-convolution of repeated Catalan numbers.

Conjecture: (n+4)*a(n) + (n+1)*a(n-1) - 4*(n+1)*a(n-2) + 4*(2-n)*a(n-3) = 0. - _R. J. Mathar_, Nov 09 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 20 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003603 | OEIS-A104794

Expansion of theta_4(q)^2 in powers of q.

Conjecture: Sum_{k>=0} a(k) / exp(k*Pi) = (1/2)*Pi^(1/2)*sqrt(2)/Gamma(3/4)^2 = A014549. - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Mar 26 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003604 | OEIS-A104857

Positive integers that cannot be represented as the sum of distinct Lucas 3-step numbers (A001644).

Similar to A054770 "Numbers that are not the sum of distinct Lucas numbers (A000204)" but with Lucas 3-step numbers (A001644). Wanted: equivalent of _David W. Wilson_ conjecture (A054770) as proved by Ian Agol. Note that all positive integers can be presented as the sum of distinct Fibonacci numbers in A000119 way. Catalani called Lucas 3-step numbers "generalized Lucas numbers" but that is quite ambiguous. These are also called tribonacci-Lucas numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Apr 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003605 | OEIS-A104863

a(n) = floor(sqrt(a(n-1)^2 + a(n-2)^2)), a(1)=10, a(2)=30.

For n>=17, a(n) = a(n-2) + a(n-4) + 1 (conjectured). If true then for m>5, a(2*m+1) = 4*F(m) + 25*F(m+1) + 1 and a(2*m+2) = 8*F(m) + 30*F(m+1) + 1 with F(n) = A000045(n). - _Ralf Stephan_, Nov 15 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Mar 28 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003606 | OEIS-A104895

a(0)=0; thereafter a(2n) = -2*a(n), a(2n+1) = 2*a(n) - 1.

Conjecture: the positions where 0, 1, 2, 3, ... appear are given by A048724; the positions where -1, -2, -3, ... appear are given by A065621.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Apr 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003607 | OEIS-A104896

a(0) = 0; a(n) = 7*a(n-1) + 7.

Conjecture: this is also the number of integers from 0 to 10^n - 1 that lack 0, 1 and 2 as a digit.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexandre Wajnberg_, Apr 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003608 | OEIS-A104934

Expansion of (1-x)/(1 - 3*x - 2*x^2).

A floretion-generated sequence relating A007482, A007483, A007484. Inverse is A046717. Inverse of Fibonacci(3n+1), A033887. Binomial transform is A052984. Inverse binomial transform is A006131. Note: the conjectured relation 2*a(n) = A007482(n) + A007483(n-1) is a result of the FAMP identity dia[I] + dia[J] + dia[K] = jes + fam Define A007483(-1) = 1. Then 2*a(n) = A007482(n) + A007483(n-1) (conjecture);

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Creighton Dement_, Mar 29 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003609 | OEIS-A104977

Defining sequence for an inverse Fredholm-Rueppel triangle.

G.f.: 1 - x / (1 + x / (1 + x / (1 - x / (1 + x / (1 - x / ...))))) = 1 + b(1) * x / (1 + b(2) * x / (1 + b(3) * x / ...)) where b(n) = (-1)^ A090678(n+1) [Conjecture]. - _Michael Somos_, Jan 03 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 30 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003610 | OEIS-A105018

Indices of the irregular primes.

Conjecture (Siegel, 1964): Limit_{n -> oo} a(n)/n = 1 - 1 / sqrt(e) (A290506). Of the 788060 primes < 12000000, 310443 are irregular.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Mar 31 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003611 | OEIS-A105151

Greatest numerator among the n! ratios equal to the continued fractions which have the permutations of (1,2,3,...,n) for terms.

Conjecture: a(n) = n*a(n-1) + (n-1)*a(n-3) + a(n-4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Apr 10 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003612 | OEIS-A105183

a(n) = 1 + a(n-1)*(a(n-1) + 1), with a(0)=2.

Conjecture: a(n) is asymptotic to d - 1/2 -(5/2^3)/d -(65/2^7)/d^3 -(650/2^11)/d^5 -(19045/2^15)/d^7 -(274950/2^19)/d^9 -(6979050/2^23)/d^11 -(130292500/2^27)/d^13 ... where d = c^(2^n) and c is a constant = 1.28820319268448517784561078485170040482944371277007918595955446677757748635242025560391582836183314154 6.... (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Apr 11 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003613 | OEIS-A105214

Companion sequence to A103683: a(n) = k if A103683(k) = n; a(n) = 0 if n is not in {A103683}.

Conjecture: a(42 + 6*k) = 0. This would follow from the conjecture that for n >= 67, A103683(n) is divisible by 3 iff n == 3 mod 4 and by 2 iff n == 2 mod 4. - _Robert Israel_, May 12 2015 The conjecture is false; a(42) = 102982. - _Rémy Sigrist_, Jan 05 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_ and _Robert G. Wilson v_, Apr 13 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003614 | OEIS-A105216

Maximum denominator among the n! ratios equal to the continued fractions which have the permutations of (1,2,3,...,n) for terms.

Conjecture: a(n) = n*a(n-1) + (n-1)*a(n-3) + a(n-4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Apr 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003615 | OEIS-A105233

Conjectured numbers n such that the trajectory of n as defined in A003508 is unique.

Conjectured numbers n such that the trajectory of n as defined in A003508 is unique.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ and _Robert G. Wilson v_, Apr 14 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003616 | OEIS-A105403

Numbers n such that prime(n)-1 and prime(n+1)-1 have the same largest prime factor.

No more terms < 100000000. Conjecture: sequence is full. - _Jon E. Schoenfield_, Jul 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, May 01 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003617 | OEIS-A105524

Diagonal sums of number triangle A105522.

Conjecture: 2*(n+2)*(13*n^2-35*n+6)*a(n) +24*(n-8)*a(n-1) +(91*n^3-375*n^2+152*n+372)*a(n-2) +12*(8-n)*a(n-3) -4*(n-3)*(13*n^2-9*n-16)*a(n-4) = 0. - _R. J. Mathar_, Nov 09 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 13 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003618 | OEIS-A105595

Row sums of number triangle A105594.

Conjecture : all terms are odd.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 14 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003619 | OEIS-A105596

a(n) = Sum_{k=0..n} A105595(k)*(-1)^k*A105595(n-k) (interpolated zeros suppressed).

Conjecture : a(2n)=1 mod 4 for all n, a(2n+1)=0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 14 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003620 | OEIS-A105600

Assume the conjectured terms of A105594 are the correct beginnings of the trajectories described in A003508. a(n) is a record length of b(n) iterations to arrive at the collected trajectories. This sequence cites the a(n)'s.

Assume the conjectured terms of A105594 are the correct beginnings of the trajectories described in A003508. a(n) is a record length of b(n) iterations to arrive at the collected trajectories. This sequence cites the a(n)'s.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ and _Robert G. Wilson v_, Apr 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003621 | OEIS-A105601

Assume the conjectured terms of A105594 are the correct beginnings of the trajectories described in A003508. a(n) is a record length of b(n) iterations to arrive at the collected trajectories. This sequence cites the b(n)'s.

Assume the conjectured terms of A105594 are the correct beginnings of the trajectories described in A003508. a(n) is a record length of b(n) iterations to arrive at the collected trajectories. This sequence cites the b(n)'s.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ and _Robert G. Wilson v_, Apr 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003622 | OEIS-A105675

Highest minimal distance of any Type II doubly-even binary self-dual code of length 8n.

Is a(9) = 12 or 16? This is an open question of long standing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 06 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003623 | OEIS-A105695

Expansion of (1-x)*c(x/(1+x)), where c(x) is the g.f. of the Catalan numbers (A000108).

Conjecture: (n+1)*a(n)+(2-3n)*a(n-1) +(1-n)*a(n-2)+3*(n-4)*a(n-3)=0. - _R. J. Mathar_, Dec 13 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003624 | OEIS-A105696

Expansion of (1-x)/sqrt((1-3*x)/(1+x)).

Conjecture D-finite with recurrence: n*a(n) +(-3*n+2)*a(n-1) +(-n+2)*a(n-2) +3*(n-4)*a(n-3)=0. - _R. J. Mathar_, Jan 22 2020 Conjecture verified using the differential equation (-3*x^2 - 1)*y(x) + (3*x^3 - x^2 - 3*x + 1)*y'(x) = 0 satisfied by the g.f. - _Robert Israel_, Jun 05 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003625 | OEIS-A105730

Number of different initial values for 3x+1 trajectories in which the largest term appearing in the iteration is 2^(6n+4).

Conjecture: All numbers greater than 10 appear in the sequence (see also A033496 and A233293). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Wasserman_, Apr 18 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003626 | OEIS-A105750

Real part of Product_{k = 0..n} (1 + k*i), i = sqrt(-1).

We conjecture that the set of type 2 primes consists of primes p == 1 (mod 4), equivalently, rational primes that split in the field extension Q(sqrt(-1)) of Q, together with the prime p = 2, which ramifies in Q(sqrt(-1)). See A002144. We conjecture that the sets of type 1 and type 3 primes taken together consist of primes p == 3 (mod 4), equivalently, rational primes that remain inert in the field extension Q(sqrt(-1)) of Q. See A002145. (End) Conjecture: a(n) -3*a(n-1) +(n^2-n+3)*a(n-2) +(-n^2+4*n-5)*a(n-3)=0. - _R. J. Mathar_, May 23 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 18 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003627 | OEIS-A105752

Expansion of e.g.f. cos(i*log(1 + x)), i = sqrt(-1).

Conjecture: a(n) = Sum_{k=0..n} Stirling1(n, 2*k). - _Benedict W. J. Irwin_, Oct 19 2016 The conjecture is true (see Fried link). - _Sela Fried_, Dec 19 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 18 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003628 | OEIS-A105794

Inverse of a generalized Stirling number triangle of first kind.

Conjecture 1: Sum_{k=0..n} (-1)^(n-k) * T(n, k) * (k+m)! / m! = (m+2)^n for m >= 0. Conjecture 2: (-1)^n - B(n) = Sum_{k=1..n} (-1)^k * T(n, k) * (k-1)! / (k+1) where B(n) = B(n, 0) is n-th Bernoulli number. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 20 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003629 | OEIS-A105802

Smallest m such that the m-th Fibonacci number has exactly n divisors that are also Fibonacci numbers.

Conjecture: a(2k+1) = 3*2^(Prime[k-1]-1) for k>3. It appears that a(2k+1) = 3*2^k for k = {1,2,3,4,6,10,12,16,18,...} = A068499[n] Numbers n such that n! reduced modulo (n+1) is not zero. - _Alexander Adamchuk_, Sep 15 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 20 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003630 | OEIS-A105849

Row sums of number triangle A105848.

Conjecture: (n+2)*a(n) +2*(-4*n-3)*a(n-1) +4*(5*n-2)*a(n-2) +2*(-10*n+17)*a(n-3) +8*(n-3)*a(n-4)=0. - _R. J. Mathar_, Nov 16 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 22 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003631 | OEIS-A105864

Expansion of (1/(1-x^2))*c(x/(1-x^2)), where c(x) is the g.f. of A000108.

Conjecture: (n+1)*a(n) + 2*(1-2*n)*a(n-1) + 2*(1-n)*a(n-2) + 2*(2*n-3)*a(n-3) + (n-3)*a(n-4) = 0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 23 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003632 | OEIS-A105865

Expansion of (1/(1-2*x^2))*c(x/(1-2*x^2)), where c(x) is the g.f. of A000108.

Conjecture: (n+1)*a(n) +2(1-2n)*a(n-1) +4*(1-n)*a(n-2) +4*(2n-3)*a(n-3) +4*(n-3)*a(n-4)=0. - _R. J. Mathar_, Dec 13 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 23 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003633 | OEIS-A105871

a(n) = sum{k=0..floor(n/2), C(2*n-3*k, n)*C(n-k, k)}.

Conjecture: 5*n*(n-1)*(3*n-10)*a(n)
-3*(n-1)*(21*n^2-63*n-20)*a(n-1)
+3*(-3*n^3+107*n^2-446*n+444)*a(n-2)
+(3*n^3-259*n^2+1279*n-1575)*a(n-3)
+3*(-21*n^3+210*n^2-673*n+694)*a(n-4)
-3*(n-3)*(3*n^2-8*n-7)*a(n-5)
-2*(n-4)*(3*n-7)*(2*n-9)*a(n-6)=0. - _R. J. Mathar_, Feb 20 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 23 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003634 | OEIS-A105875

Primes for which -3 is a primitive root.

On Artin's conjecture, the number of terms <= N is asymptotically equivalent to (6/5)*C*primepi(N), where C = A005596 is Artin's constant. - _Jianing Song_, Sep 12 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003635 | OEIS-A105876

Primes for which -4 is a primitive root.

On Artin's conjecture, the number of terms <= N is asymptotically equivalent to C*primepi(N), where C = A005596 is Artin's constant. - _Jianing Song_, Oct 06 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003636 | OEIS-A105877

Primes for which -5 is a primitive root.

Conjecture: the penultimate digit of a(n) is always odd. This characteristic seems to be proper of primes for which -5*n^2 is a primitive root. - _Davide Rotondo_, Oct 26 2024 For any n > 1, a(n) == 11, 13, 17, or 19 (mod 20), which implies the conjecture above. - _Max Alekseyev_, Nov 01 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003637 | OEIS-A105887

Primes for which -15 is a primitive root.

then by Artin's conjecture, Q(N) ~ (94/95)*C*N/log(N) ~ (188/95)*C*(Pi(N,7) + Pi(N,11) + Pi(N,13) + Pi(N,14)), where C = A005596 is Artin's constant. Conjecture: if we further define Conjectured | 0.21277 | 0.25532 | 0.21277 | 0.31915 | 1.00000

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003638 | OEIS-A105974

(A001035(n) - 1)/2 where A001035 is the number of partially ordered sets ("posets") with n labeled elements.

Conjectures: a(n) is (pre)periodic modulo m for all m, the lengths of the ending periods being given by 2*A000010(m) if m is a power of 2 and A000010(m) otherwise. A000010 is Euler's totient function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerald McGarvey_, Apr 28 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003639 | OEIS-A106181

Expansion of c(-x^2)(1+2x-sqrt(1+4x^2))/2, c(x) the g.f. of A000108.

Conjecture: (n+2)*a(n) + n*a(n-1) + 4*(n-1)*a(n-2) + 4*(n-3)*a(n-3) = 0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003640 | OEIS-A106228

G.f. A(x) satisfies A(x) = 1 + x*A(x)/(1 - x*A(x)^2).

Conjecturally, the number of sequences (e(1), ..., e(n)), 0 <= e(i) < i, such that there is no triple i < j < k with e(i) > e(j) and e(i) <= e(k). [Martinez and Savage, 2.21] - _Eric M. Schmidt_, Jul 17 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 19 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003641 | OEIS-A106249

Expansion of (1-x+x^2+x^3)/((1+x)*(1+x^2)*(1-x)^2).

Conjecture: number of roots of x^n + 1 in the left half-plane for n > 0. - _Michel Lagneau_, Oct 31 2012 a(n) = Sum_{k=0..n} Jacobi(2^k, 2k+1) (Conjecture). - _Paul Barry_, Jul 23 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 27 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003642 | OEIS-A106252

Number of positive integer triples (x,y,z), with x<=y<=z<=n, such that each of x,y and z divides the sum of the other two.

The following conjecture is probably not very difficult: Conjecture. The sequence (A106253) of differences of this sequence is periodic with period 6. That the difference sequence in the above conjecture is periodic follows from a formula in the Formula and Mathematica sections; see A211701 for a discussion. - _Clark Kimberling_, Apr 20 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Apr 27 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003643 | OEIS-A106271

Row sums of number triangle A106270.

To prove _R. J. Mathar_'s conjecture, let A(x) be the g.f. of the current sequence. We note first that which proves the conjecture. (End) Conjecture: (n+1)*a(n) + (1-5*n)*a(n-1) + 2*(2*n-1)*a(n-2) = 0. - _R. J. Mathar_, Nov 09 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 28 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003644 | OEIS-A106272

Antidiagonal sums of number triangle A106270.

To prove _R. J. Mathar_'s conjecture, let A(x) be the g.f. of the current sequence. We note first that which proves the conjecture. - _Petros Hadjicostas_, Jul 15 2019 Conjecture: (n+1)*a(n) + 2*(1-2*n)*a(n-1) - (n+1)*a(n-2) + 2*(2*n-1)*a(n-3) = 0. - _R. J. Mathar_, Nov 09 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 28 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003645 | OEIS-A106297

Period of the Lucas 5-step sequence A074048 mod n.

Conjectures: a(5^k) = 781*5^(k-1) for k > 0. a(2^k) = 3*2^(k-1) for k > 1. If a(p) != a(p^2) for prime p > 2, then a(p^k) = p^(k-1)*a(p) for k > 0. - _Chai Wah Wu_, Feb 25 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, May 02 2005, Nov 19 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003646 | OEIS-A106303

Period of the Fibonacci 5-step sequence A001591 mod n.

Conjectures: a(5^k) = 781*5^(k-1) for k > 0. If a(p) != a(p^2) for p prime, then a(p^k) = p^(k-1)*a(p) for k > 0. - _Chai Wah Wu_, Feb 25 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, May 02 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003647 | OEIS-A106346

Inverse of number triangle A106344.

Rows sums are A036987 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 29 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003648 | OEIS-A106369

Number of circular compositions of n such that no two adjacent parts are equal.

Conjecture: a(n) ~ A241902^n / n. - _Vaclav Kotesovec_, Sep 06 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, Apr 29 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003649 | OEIS-A106391

A "cosh transform" of binomial(2n,n-1).

Conjecture: -(n+1)*(n-2)*a(n) +4*n*(3*n-7)*a(n-1) +(-49*n^2+193*n-148)*a(n-2) +8*(9*n-19)*(n-4)*a(n-3) +(5*n^2+115*n-418)*a(n-4) -12*(7*n-19)*(n-4)*a(n-5) +45*(n-4)*(n-5)*a(n-6)=0. - _R. J. Mathar_, Feb 20 2015 Conjecture: -(n-2)*(n+1)*(4*n^2-16*n+19)*a(n) +8*n*(n-2)*(4*n^2-14*n+13)*a(n-1) +2*(-28*n^4+168*n^3-375*n^2+401*n-180)*a(n-2) -8*(n-2)*(4*n^3-18*n^2+27*n-10)*a(n-3) +15*(n-2)*(n-3)*(4*n^2-8*n+7)*a(n-4)=0. - _R. J. Mathar_, Feb 20 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 01 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003650 | OEIS-A106521

Numbers m such that Sum_{k=0..10} (m+k)^2 is a square.

11*((m+5)^2+10) is a square iff the second factor is divisible by 11 and the quotient is a square, i.e., iff m = 11*k - 4 or m = 11*k - 6 and 11*k^2 +- 2 k + 1 is a square. Thus a(n) == (7,5,5,7,7,5,5,7,...) (mod 11), repeating with period 4 and the values are obtained by solving these Pell-type equations (cf. link to Dario Alpern's quadratic solver). The corresponding recurrence equations (see PARI code) should make it possible to prove the conjectured g.f. - _M. F. Hasler_, Jan 27 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, May 30 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003651 | OEIS-A106564

Perfect squares which are not the difference of two primes.

It is conjectured (see A020483) that every even number is a difference of primes, and this is known to be true for even numbers < 10^11. If so,this sequence consists of the odd squares n such that n+2 is composite. - _Robert Israel_, Feb 28 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexandre Wajnberg_, May 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003652 | OEIS-A106575

Perfect squares which are both the sum and the difference of two primes.

By well-known conjectures, every even integer > 2 is both the sum and the difference of two primes; this would be a special case. - _Franklin T. Adams-Watters_, Sep 13 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexandre Wajnberg_, May 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003653 | OEIS-A106741

Numbers n such that n divides the denominator of 2n-th Bernoulli number.

In 2005, B. C. Kellner proved E. W. Weisstein's conjecture that denom(B_n) = n only if n = 1806. - _Jonathan Sondow_, Oct 14 2013.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003654 | OEIS-A107055

Integer part of Sum_{k>=0} Sum_{j=0..k} n^j*A107045(k,j)/A107046(k,j).

Limit a(n+1)/a(n) exists and is conjectured to equal exp(exp(-1)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003655 | OEIS-A107099

G.f. satisfies A(A(x)) = x + 4*x^3, where A(x) = Sum_{n>=0} a(n)*x^(2*n+1).

Coefficients [x^n] A(x) = 0 (mod 3) except at n = 3^k (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 13 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003656 | OEIS-A107231

a(n) = C(n+2,2)*C(n,floor(n/2)).

Conjecture: n*a(n) +(n-4)*a(n-1) +2*(-2*n-5)*a(n-2) -4*n*a(n-3)=0. - _R. J. Mathar_, Nov 24 2012 Conjecture confirmed using the differential equation (12*x^2 + 18*x + 3)*y(x) + (4*x^3 + 4*x^2 - x - 1)*y'(x) = 0 satisfied by the G.f. - _Robert Israel_, Oct 07 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 13 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003657 | OEIS-A107263

Position where n (presumably) appears the last time in A032531, or 0 if n keeps appearing.

I extended the analysis of A032531(n) to all n<= 10^7. Same comments apply considering the new limit and, of course, the uniqueness of Stephan's sequence remains as always only a conjecture since there's no proof that the sequence should be anything different from the zero sequence for all, most or even any of the terms - Herman Jamke (hermanjamke(AT)fastmail.fm), Nov 08 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, May 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003658 | OEIS-A107267

A square array of Motzkin related transforms, read by antidiagonals.

Rows are transforms of k^n, k>=0, under the matrix A107131. As a number triangle, with T(n,k)=if(k<=n,sum{j=0..n-k, (1/(j+1))C(j+1,n-k-j+1)C(n-k,j)k^j},0), row sums are A107268 and diagonal sums are A107269. Rows are series reversions of x/(1+kx+kx^2), k>=0. Conjecture: rows count weighted Motzkin paths.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003659 | OEIS-A107311

Decimal expansion of the solution to zeta(x) = 2.

Conjecture 1: the real parts of the zeros of the prime zeta function are not greater than this constant. Conjecture 2: the real parts of the zeros of the anyone subset of the prime zeta function are not greater than this constant. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, May 20 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003660 | OEIS-A107342

Semiprimes with semiprime digits (digits 4, 6, 9 only).

Conjecture: almost all terms (asymptotic density 1) end with 9 and have either 3k+1 or 3k+2 occurrences of the digit 4 for some nonnegative k. (Otherwise they'd be divisible by 2 or 3 and these semiprimes would be expected to be rare; the sequence is too thin to prove this directly.) - _Charles R Greathouse IV_, Nov 12 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, May 22 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003661 | OEIS-A107363

Expansion of (1 - x)*(1 + x)^2*(1 + x^2)*(1 - x^2 + 2*x^3 + x^4) / ((1 - x^2 - x^4)*(1 + x^2 + 2*x^4 - x^6 + x^8)).

Conjectures: { Fib(n) | n in naturals } = { a(n) | n in naturals, a(n) >= 0 } = { a(n) | n in naturals, n not of the form 6*n+2 } (naturals include 0). a(6*n+2) = - A048876(n) (Generalized Pellian with second term of 7), conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Creighton Dement_, May 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003662 | OEIS-A107578

Prime index of A000101(n), maximal gap upper end prime index.

Conjecture: log a(n) ~ n/2. That is, record prime gaps occur about twice as often as records in an i.i.d. random sequence of comparable length (see arXiv:1709.05508 for a heuristic explanation). - _Alexei Kourbatov_, Jan 18 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | Alex Beveridge, Apr 25 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003663 | OEIS-A107587

Number of Motzkin n-paths with an even number of up steps.

Conjecture: n*(n+2)*a(n) + (-5*n^2-n+3)*a(n-1) + (10*n^2-16*n+3)*a(n-2) + (-10*n^2+34*n-27)*a(n-3) - (11*n-5)*(n-3)*a(n-4) + 15*(n-3)*(n-4)*a(n-5) = 0. - _R. J. Mathar_, Feb 20 2015 [conjecture is true for n = 0..800; checked by _Gennady Eremin_, Jan 25 2021] Conjecture: n*(n-2)*(n+2)*a(n) - (2*n-1)*(2*n^2-2*n-3)*a(n-1) + 3*(n-1)*(2*n^2-4*n+1)*a(n-2) - 2*(n-1)*(n-2)*(2*n-3)*a(n-3) - 15*(n-1)*(n-2)*(n-3)*a(n-4) = 0. - _R. J. Mathar_, Feb 20 2015 [conjecture is true for n = 0..800; checked by _Gennady Eremin_, Jan 25 2021]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003664 | OEIS-A107651

Numbers n such that phi(sigma(n)) + phi(phi(n)) = n.

If both (3^n-1)/2 and 2*3^n-1 are prime then 48*(2*3^n-1) is in the sequence (the proof is easy). So if n is in the intersection of A028491 and A003307 then 48*(2*3^n-1) is in this sequence. Conjecture: There exist only two such terms, namely 2544 and 209904. If both (3^n*31-1)/2 and

2*3^n*31-1 are prime then 48*(2*3^n*31-1) is in the sequence (the proof is easy). Conjecture: There exist only three such terms, namely 26736, 2169456, and 263761038440858432614846 56.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, May 26 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003665 | OEIS-A107655

a(n) is the smallest number m greater than 1 such that phi(m) = d(m)^n, where d(m) is number of positive divisors of m; if there is no such m, a(n)=1.

Conjecture: A000005(a(n)) <= 12 for all n. [_Max Alekseyev_, May 07 2010] This conjecture holds throughout the first 102 terms. - _David A. Corneth_, Jun 14 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Jun 06 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003666 | OEIS-A107668

Column 0 of triangle A107667.

Conjecture: a(n) = (n+1) * P(n+1, n+1) where P(n, k) = P(n-1, k) + n * P(n, k-1) for 1 < k < n with P(n, 1) = 1, P(n, n) = n * P(n, n-1). - _Mikhail Kurkov_, Jun 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 07 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003667 | OEIS-A107740

Number of numbers m such that prime(n) = m + (digit sum of m).

a(n) <= 2 for n <= 10^5. Conjecture: sequence is bounded. I would rather conjecture the opposite. Of course a(n) >= m implies n >= A006064(m), having more than A230857(m) digits, i.e., 14, 25 and 1111111111125 digits of n, for a(n) = 3, 4, 5. - _M. F. Hasler_, Nov 09 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, May 23 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003668 | OEIS-A107742

G.f.: Product_{j>=1} Product_{i>=1} (1 + x^(i*j)).

Conjecture: log(a(n)) ~ Pi*sqrt(n*log(n)/6). - _Vaclav Kotesovec_, Aug 29 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Jun 11 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003669 | OEIS-A107750

If n=0 then 0, else smallest number greater than its predecessor and having either more or fewer zeros in its binary representation.

Conjectures from _Colin Barker_, Jul 24 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, May 23 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003670 | OEIS-A107793

Differences between successive indices of 1's in the ternary tribonacci sequence A305390.

Conjecture (_N. J. A. Sloane_, Jun 22 2018) This is a disguised form of A275925. More precisely, if we replace the 5's by 6's and the 4's by 5's, and ignore the first two terms, we appear to get a sequence which is a shifted version of A275925.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Jun 11 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003671 | OEIS-A107861

Number of distinct values taken by the sums of all subsets of the n-th roots of unity.

Note that a(6)=19, a(12)=19^2 and a(18)=19^3. Similarly, a(10)=211 and a(20)=211^2. For prime n, a(n)=2^n-1. For powers of 2, we have a(2^n)=3^(2^(n-1)). It appears that _David W. Wilson_'s conjectured formula for A103314 may apply to this sequence also. Observe that due to symmetry, n divides a(n)-1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, May 25 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003672 | OEIS-A107863

Column 1 of triangle A107862; a(n) = binomial(n*(n+1)/2 + n, n).

Conjectural congruences:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003673 | OEIS-A107877

Column 1 of triangle A107876.

Conjecture: a(n) can be expressed with a series of nested sums,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 04 2005, Apr 10 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003674 | OEIS-A107878

Column 2 of triangle A107876.

Conjecture: a(n) is given by a series of nested sums,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003675 | OEIS-A107882

Column 1 of triangle A107880.

Conjecture: a(n) can be expressed with a series of nested sums,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003676 | OEIS-A107886

Column 1 of triangle A107884.

Conjecture: a(n) can be expressed with a series of nested sums,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003677 | OEIS-A107928

a(n) is the numerator of harmonic mean of a(n-1) and a(n-2).

Conjecture: G.f.: x*(2+3*x+12*x^2+8*x^3+8*x^4)/[(1-2*x)*(1+2*x+4*x^2)]. a(3n) = (3/2)*8^n, a(3n+1) = 3*8^n, a(3n+2) = 2*8^n, for n>0. - _Ralf Stephan_, Dec 01 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Jun 10 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003678 | OEIS-A107986

Composite numbers of the form p+2 where p is prime.

This sequence is analogous to the sequence formed by the Goldbach-Euler conjecture that every even number greater than 2 is the sum of 2 primes. If p + 2 is prime then p and p + 2 are twin primes. The number of terms in this sequence is infinite. This follows immediately from the proof that the number of primes p is infinite. Conjecture: The ratio of the number of terms in this sequence to Pi(n) tends to a limit < 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jun 13 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003679 | OEIS-A108137

Primes p such that p + 6^k is composite for all k >= 0.

The PARI code given suggests that these values are merely conjectures. - _N. J. A. Sloane_, Jun 30 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jun 27 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003680 | OEIS-A108200

Least positive k such that k*n + 1 is a golden semiprime (A108540).

Conjecture: for every n > 4 there exists a number k < n^[n/2] such that k*n + 1 is a golden semiprime, where [] is the floor function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jun 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003681 | OEIS-A108219

Numbers n such that A001414(n) is a golden semiprime, where A001414 is the sum of primes dividing n (with repetition).

Numbers n such that A001414(n) and A001414(n+1) are both golden semiprimes: 8, 125, 153759, 247455, 678807, 1243499, 1243500, ... Notice that the last two terms indicate a triple. Conjecture: this subsequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jun 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003682 | OEIS-A108225

a(0) = 0, a(1) = 2; for n >= 2, a(n) = (a(n-1) + a(n-2))*(a(n-1) - a(n-2) + 1)/2.

Conjecture: a(n) = A006894(n) + 1. - _R. J. Mathar_, Apr 23 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003683 | OEIS-A108296

Diagonal sums of the number triangle associated to A086617.

Conjecture: (n+3)*a(n) +(-n-2)*a(n-1) +2*(-n-1)*a(n-2) +2*(-n+3)*a(n-3) +(n+1)*a(n-4) +3*(n-2)*a(n-5)=0. - _R. J. Mathar_, Nov 16 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 31 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003684 | OEIS-A108299

Triangle read by rows, 0 <= k <= n: T(n,k) = binomial(n-[(k+1)/2],[k/2])*(-1)^[(k+1)/2].

Conjecture: Let N=2*n+1, with n > 2. Then T(n,k) (0 <= k <= n) gives the k-th coefficient in the characteristic function p_N(x)=0, of degree n in x, for the n X n tridiagonal unit-primitive matrix G_N (see [Jeffery]) of the form

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jun 01 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003685 | OEIS-A108307

Number of set partitions of {1, ..., n} that avoid enhanced 3-crossings (or enhanced 3-nestings).

Conjecturally, the number of sequences (e(1), ..., e(n)), 0 <= e(i) < i, such that there is no triple i < j < k with e(j) <= e(k) and e(i) >= e(k). [Martinez and Savage, 2.16] Conjecturally, the number of sequences (e(1), ..., e(n)), 0 <= e(i) < i, such that there is no triple i < j < k with e(i) >= e(j) >= e(k). [Martinez and Savage, 2.16] The second of the above-mentioned conjectures is proved in Zhicong Lin's paper. - _Eric M. Schmidt_, Nov 25 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mireille Bousquet-Mélou_, Jun 29 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003686 | OEIS-A108308

Expansion of 1/(1-x^2*c(2*x)), c(x) the g.f. of A000108.

Conjecture: 2*(n-1)*a(n) + (41-17*n)*a(n-1) + 4*(2*n-5)*a(n-2) + (n-1)*a(n-3) + 4*(5-2*n)*a(n-4) = 0. - _R. J. Mathar_, Dec 10 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003687 | OEIS-A108315

Composite numbers n such that a^v + b^v + c^v + ... is prime, where a*b*c* ... is the prime factorization of n and v is the number of primes dividing n (counted with repetition).

5166592 generates a 41 digit prime. Some consecutive terms are (74,75) (152,153) (325,326) ... Conjecture: there are infinitely many consecutive values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jun 30 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003688 | OEIS-A108317

Smallest a(n) such that a(n) n's plus a(n) is prime, or 0 if no such a(n) exists.

Some or all of the zero values are merely conjectures. - _N. J. A. Sloane_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ray G. Opao_, Jun 30 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003689 | OEIS-A108345

The set N such that each positive integer can be written in the form s^2 + n, s>=0, n in N, in an even number of ways.

This set is conjectured to have zero density. The only even values are the numbers of the form 2n^2. See the paper by Cooper, Eichhorn and O'Bryant for more details. - _Franklin T. Adams-Watters_, May 16 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Jul 01 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003690 | OEIS-A108399

Least positive k such that n^2 + k is a golden semiprime (A108540).

Conjecture: for every n > 1 there exists a number k < n^3 such that n^2 + k is a golden semiprime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jul 03 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003691 | OEIS-A108411

a(n) = 3^floor(n/2). Powers of 3 repeated.

Conjecture: a(n+1) is the number of distinct subsets S of {0,1,2,...,n} such that the sumset S+S does not contain n. - _Michael Chu_, Oct 05 2021. _Andrew Howroyd_, Nov 20 2021: The conjecture is true: If there are m pairs of numbers that add to n then inclusion/exclusion gives sum(k=0, m, binomial(m,k)*(-1)^k*2^(2*m-2*k)) as the number of sets that don't contain any of those pairs which equals 3^m. For even n , n/2 cannot be included in any set.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Jun 05 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003692 | OEIS-A108430

Number of d steps in all paths from (0,0) to (3n,0) that stay in the first quadrant (but may touch the horizontal axis), consisting of steps u=(2,1), U=(1,2), or d=(1,-1).

Conjecture D-finite with recurrence n*(2*n+1)*(23982*n-28681)*a(n) +(-640736*n^3+1168048*n^2-901220*n+247035)*a(n-1) +(1196488*n^3-6448608*n^2+10992587*n-5911365)*a(n-2) +2*(2*n-5)*(28283*n-6993)*(n-3)*a(n-3)=0. - _R. J. Mathar_, Jul 24 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jun 03 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003693 | OEIS-A108449

Number of paths from (0,0) to (3n,0) that stay in the first quadrant (but may touch the horizontal axis), consisting of steps u=(2,1),U=(1,2), or d=(1,-1) and having no pyramids (a pyramid is a sequence u^pd^p or U^pd^(2p) for some positive integer p, starting at the x-axis).

Conjecture D-finite with recurrence +n*(2*n+1)*(431*n-2895)*a(n) +2*(-9395*n^3+68622*n^2-64084*n+26109)*a(n-1) +2*(59288*n^3-508196*n^2+1044822*n-574587)*a(n-2) +2*(-94965*n^3+1070605*n^2-3607435*n+3485484)*a(n-3) +4*(29036*n^3-351474*n^2+1402336*n-1970505)*a(n-4) +6*(-6703*n^3+63052*n^2-99178*n-237177)*a(n-5) +6*(1012*n^3-14914*n^2+74580*n-127341)*a(n-6) +6*(1127*n^3-21429*n^2+135199*n-282762

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jun 11 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003694 | OEIS-A108475

Expansion of (1-3*x) / (1-5*x-5*x^2+x^3).

Conjecture: a(n) = A000129(n+1)*A001333(n). - _R. J. Mathar_, Jul 08 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jun 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003695 | OEIS-A108476

Expansion of (1-4*x)/(1-6*x-12*x^2+8*x^3).

Conjecture: a(n) = A002605(n+1)*A026150(n). - _R. J. Mathar_, Jul 08 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jun 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003696 | OEIS-A108524

Number of ordered rooted trees with n generators.

Conjecture: 2*n*a(n) +(24-17*n)*a(n-1) +4*(4*n-9)*a(n-2) +4*(3-n)*a(n-3)=0. - _R. J. Mathar_, Nov 14 2011 The o.g.f. A(x) satisfies the differential equation (2 - 17*x + 16*x^2 - 4*x^3)*A'(x) + (7 - 4*x)*A(x) = 2 - 2*x. Mathar's conjectural recurrence above follows from this.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, Jun 07 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003697 | OEIS-A108600

Number of freely braided permutations of length n; the freely braided permutations are those that avoid 3421, 4231, 4312 and 4321.

Conjecture: (1-n)*a(n) +(7*n-10)*a(n-1) +2*(1-4*n)*a(n-2) +8*(11-2*n)*a(n-3) +(n-1)*a(n-4) +3*(2-n)*a(n-5) +2*(11-2*n)*a(n-6)=0. - _R. J. Mathar_, Aug 24 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincent Vatter_, Jun 11 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003698 | OEIS-A108625

Square array, read by antidiagonals, where row n equals the crystal ball sequence for the A_n lattice.

Conjectural result for superdiagonals: zeta(2) = 1 + 1/2^2 + ... + 1/k^2 + Sum_{n >= 1} (-1)^(n+1) * (5*n^2 + 6*k*n + 2*k^2)/(n^2*(n+k)^2*T(n-1,n+k-1)*T(n,n+k)), k = 0,1,2... . Conjectural result for subdiagonals: zeta(2) = 2*(1 - 1/2^2 + ... + (-1)^(k+1)/k^2) + (-1)^k*Sum_{n >= 1} (-1)^(n+1)*(5*n^2 + 4*k*n + k^2)/(n^2*(n+k)^2*T(n+k-1,n-1)*T(n+k,n)), k = 0,1,2... . Conjectural congruences: the main superdiagonal numbers S(n) := T(n,n+1) appear to satisfy the supercongruences

S(m*p^r - 1) = S(m*p^(r-1) - 1) (mod p^(3*r)) for all primes p >= 5 and all positive integers m and r. If p is prime of the form 4*n + 1 we can write p = a^2 + b^2 with a an odd number. Then calculation suggests the congruence S((p-1)/2) == 2*a^2 (mod p). (End) Conjecture: for positive integer m, Sum_{k = 0..n} (-1)^(n+k) * binomial(n, k) * T(m*n, k) = ((m+1)*n)!/(((m-1)*n)!*n!^2) (verified up to m = 10 using the MulZeil procedure in Doron Zeilberger's MultiZeilberger package). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003699 | OEIS-A108628

n-th term of the crystal ball sequence for A_{n+1} lattice for n >= 0.

Conjectures: a((p-1)/2) == 0 (mod p) and a((p^3-1)/2) == 0 (mod p^3) for primes p of the form 4*m + 1; a((p^3-1)/2) == 0 (mod p^2) for primes p of the form 4*m + 3; a((p^2-1)/2) == 0 (mod p^2) for primes p >= 5. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 14 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003700 | OEIS-A108629

Expansion of g.f.: (1+x - sqrt(1-2*x-3*x^2-4*x^3))/(2+2*x+2*x^2).

Conjecture D-finite with recurrence: n*a(n) = (n-3)*a(n-1) +4*(n-3)*a(n-2) +3*(3*n-10)*a(n-3) +(7*n-27)*a(n-4) +2*(2*n-9)*a(n-5). - _R. J. Mathar_, Jan 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, Jun 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003701 | OEIS-A108630

Expansion of g.f.: (1+x-sqrt(1-2*x-7*x^2-8*x^3-4*x^4))/(2*(1+x+x^2)).

Conjecture D-finite with recurrence: (n-1)*a(n) = (n-4)*a(n-1) +8*(n-4)*a(n-2) +(17*n-77)*a(n-3) +(19*n-100)*a(n-4) +12*(n-6)*a(n-5) +4*(n-7)*a(n-6). - _R. J. Mathar_, Sep 27 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, Jun 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003702 | OEIS-A108668

Self-erasure surviving integers in the concatenation of all nonnegative integers.

Conjecture: the sequence is finite. Last term?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Alexandre Wajnberg_, Jul 07 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003703 | OEIS-A108704

Number of partitions of 112233...nn into n pairs.

Conjecture: a(n)-(2*n-1)*a(n-1)-(n-1)*a(n-2)+2*(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Jun 08 2016 The exponential generating series A(t) satisfies the differential equation (1-2*t)*A'(t) = (1+t-2*t^2)*A(t), which is equivalent to the conjectured recurrence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Miklos Kristof_, Jun 20 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003704 | OEIS-A108716

a(n) = tan(Pi/14)^(-2n) + tan(3*Pi/14)^(-2n) + tan(5*Pi/14)^(-2n).

Conjecture: a(n) = 21*a(n-1)-35*a(n-2)+7*a(n-3). G.f.: -(35*x^2-42*x+3) / (7*x^3-35*x^2+21*x-1). - _Colin Barker_, Jun 01 2013 To verify conjecture, note that the roots of 7*x^3-35*x^2+21*x-1 are tan(Pi/14)^2, tan(3*Pi/14)^2 and tan(5*Pi/14)^2. - _Robert Israel_, Aug 23 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Jun 20 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003705 | OEIS-A108764

Products of exactly two supersingular primes (A002267).

Peter Luschny's link shows how this sequence may be connected to Schinzel-Sierpinski conjecture and the Calkin-Wilf tree.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jun 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003706 | OEIS-A108770

Numbers k such that k^2 + (k+1)^2 is a brilliant number.

Consecutive terms are (254,255) (4099,4100) (11159,11160) (25094,25095) (31754,31755) (40189,40190) ... Conjecture: There are infinitely many consecutive values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jun 25 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003707 | OEIS-A108794

Primes whose 10's complement is a semiprime, i.e., p is prime and 10^L - p is a semiprime, where L is the number of digits in p.

Some twin prime terms are (149,151), (311,313), (419,421), (659,661), ... Conjecture: there are infinitely many twin primes in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jul 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003708 | OEIS-A108795

Conjectured greatest number k such that C(2k,k) is not divisible by any odd prime to the n-th power.

Conjectured greatest number k such that C(2k,k) is not divisible by any odd prime to the n-th power.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_ and _Robert G. Wilson v_, Nov 29 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003709 | OEIS-A108863

Number of Dyck paths containing exactly one UUUD.

Conjecture: this is the Motzkin transform of the sequence of three zeros followed by A001651. - _R. J. Mathar_, Dec 11 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Callan_, Jul 25 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003710 | OEIS-A108897

Numbers k such that 60*k^2 + 30*k - 30 +- 1 is a twin prime pair.

Some consecutive terms in this sequence are (102:103), (105:106), (143:144), ... (1320071:1320072), (1320250:1320251) ... Conjecture: There are infinitely many of these consecutive pairs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jul 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003711 | OEIS-A108946

a(2n) = A001570(n), a(2n+1) = -A007654(n+1).

Define c(n) = a(n) - 4*a(n+1) - a(n+2) and d(n) = -a(n) - 4*a(n+1) - a(n+2); Conjectures: I: c(2n) = 24*A076139(n); (Triangular numbers that are one-third of another triangular number) II: c(2n+1) = -A011943(n+1); (Numbers n such that any group of n consecutive integers has integral standard deviation) III: d(2n) = -2; IV: d(2n+1) = -1

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Creighton Dement_, Jul 21 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003712 | OEIS-A108954

a(n) = pi(2*n) - pi(n). Number of primes in the interval (n,2n].

Conjecture: G.f.: Sum_{i>0} Sum_{j>=i|i+j is prime} x^j. - _Benedict W. J. Irwin_, Mar 31 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jul 22 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003713 | OEIS-A108957

Values of n such that n - 2^k is deficient for all 1 <= 2^k < n.

Conjectures: a. Sequence is infinite. b. There are infinitely many consecutive pairs, such as (5:6), (11:12), (17:18), (53:54), ... (204005:204006).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jul 22 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003714 | OEIS-A109035

Number of irreducible partitions into squares. A partition is irreducible if no subpartition with 2 or more parts sums to a square.

Sequence is unbounded, as can be seen by considering sums of 2 squares (thanks to _David L. Harden_). Obviously it contains infinitely many 1's, at square indices. At nonsquare indices, series appears to go to infinity, but this is conjecture and growth rate is entirely unknown. Also unknown is whether the sequence is onto the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Jun 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003715 | OEIS-A109036

Number of irreducible partitions into smaller squares. A partition is irreducible if no subpartition with 2 or more parts sums to a square smaller than n.

This is the same as A109035 except for the values at squares. Conjecture that lim_{n->oo} a(n) = oo.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Jun 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003716 | OEIS-A109037

Number of irreducible partitions into triangular numbers. A partition is irreducible if no subpartition with 2 or more parts sums to a triangular number.

Sequence is almost certainly unbounded. Obviously it contains infinitely many 1's, at triangular indices. At non-triangular indices, series appears to go to infinity, but this is conjecture and growth rate is entirely unknown. Also unknown is whether the sequence is onto the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Jun 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003717 | OEIS-A109038

Number of irreducible partitions into smaller triangular numbers. A partition is irreducible if no subpartition with 2 or more parts sums to a smaller triangular number.

Same as A109037 except at triangular indices. Conjecture than lim_{n->oo} a(n) = oo.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Jun 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003718 | OEIS-A109089

Decimal expansion of a = f(1) = f(2) with function f(x) from A109087.

Continued fraction expansion of a is in A109090. Conjecture: a is transcendental.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Nikolaus Meyberg (Nikolaus.Meyberg(AT)t-online.de), Jun 20 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003719 | OEIS-A109095

Numbers N such that N! is the product of exactly two smaller factorials (larger than 1).

All terms except a(2) = 10 appear to be trivial solutions. (From Erdős's paper this is known as Surányi's conjecture.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_, Jun 19 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003720 | OEIS-A109132

Numbers n such that n! + 1 is a Chen prime. n! + 1 is then called a factorial Chen prime.

Conjecture: 77! + 1 is the largest factorial Chen prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Aug 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003721 | OEIS-A109134

Decimal expansion of Phi, the real root of the equation 1/x = (x-1)^2.

Suppose that r is a real number in the interval [3/2, 5/3). Let C(r) = (c(k)) be the sequence of coefficients in the Maclaurin series for 1/(Sum_{k>=0} floor((k+1)*r))(-x)^k). Conjectures: the limit L(r) of c(k+1)/c(k) as k -> oo exists, L(r) is discontinuous at 5/3 (cf. A279676), and the left limit of L(r) as r->5/3 is Phi. - _Clark Kimberling_, Jul 11 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Aug 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003722 | OEIS-A109136

Even numbers that are not congruent to 1 mod 3 nor are squares nor are the sum of a prime and a square of a prime.

We conjecture that every sufficiently large even number is congruent to 1 mod 3, or is a square, or is a sum of a prime and a square of a prime. The numbers listed here are believed to be the only exceptions. This has been verified up to 100000000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Dmytro Taranovsky (dmytro(AT)mit.edu), Aug 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003723 | OEIS-A109224

Minimum number of moves to solve the second Panex puzzle of order n of exchanging the two side towers.

Conjecture: for n >= 5, a(n) = ((7/2) * (7 - (4 * sqrt(2))) * (1 + sqrt(2))^n) - (8 * n) - 2 + ((7/2) * (7 + (4 * sqrt(2))) * (1 - sqrt(2))^n). Confirmed to n = 13, and known to be an upper bound for all larger n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _William Rex Marshall_, Aug 19 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003724 | OEIS-A109326

Smallest positive number that requires n steps to be represented as a sum of palindromes using the greedy algorithm.

The greedy algorithm means iteration of A261424 until a palindrome is reached. For n = 3, 4, ... we have a(n+1) = 10^L(n) + a(n) + 1 with L(n) = 2^(n-2) + 1 = length(a(n))*2 - 3 for n > 3. We have a(7) <= 10^17 + 1000101025, a(8) <= 10^33 + 10^17 + 1000101026, a(9) <= 10^65 + 10^33 + 10^17 + 1000101027, a(10) <= 10^129 + 10^65 + 10^33 + 10^17 + 1000101028, etc, with conjectured equality. - _M. F. Hasler_, Sep 08 2015, edited Sep 09 2018 a(n) = Sum_{0 <= k <= n-3} 10^(2^k+1) + n - 82, for n > 2 (conjectured). - _M. F. Hasler_, Sep 08 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Wasserman_, Aug 11 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003725 | OEIS-A109381

Maximum digit of n^2 written in factorial base.

Conjecture: lim_{n->infinity} a(n) = infinity. If true, convergence is very slow, since a(1183893) = 1. Sequence is certainly unbounded, since for n >= 4, there is always a square between n*n! and (n+1)!.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Aug 25 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003726 | OEIS-A109385

Maximum number of prime implicants of a symmetric function of n Boolean variables.

Many people have conjectured that this sequence is equal to A003039. Certainly it is a lower bound. An upper bound is given in A109388.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Don Knuth_, Aug 25 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003727 | OEIS-A109464

Numbers n such that A112131(n) and A112131(n+1) are both Harshad numbers.

Conjecture: Sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Aug 28 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003728 | OEIS-A109474

a(1)=1, a(2)=3; thereafter, a(n) = least positive integer > a(n-1) and not equal to a(i)+a(j)+a(k) for 1<=i<=j<=k<=n-1.

Conjecture: Except for the first term, a(n)=10*n-a(n-1)-23 (with a(2)=3). - _Vincenzo Librandi_, Dec 07 2010 [This is easily proved. - _N. J. A. Sloane_, Aug 07 2017]

猜想/问题提及段 | historical_or_current_mention_unreviewed | Bela Bajnok (bbajnok(AT)gettysburg.edu), Aug 10 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003729 | OEIS-A109516

a(n) is the (1,2)-entry of the n-th power of the 2 X 2 matrix [0,1;n-1,n-1].

a(n+1) = Sum_{k=0..floor(n/2)} binomial(n-k,k)*n^(n-k) for n>=0 (conjectured). - _Werner Schulte_, Oct 21 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Jun 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003730 | OEIS-A109634

Number of 1's that appear among all ternary strings of length n that contain no consecutive 1's.

Conjecture: a(n)=A073388(n-1). [From _R. J. Mathar_, Aug 18 2008]

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ralph P. Grimaldi (ralph.grimaldi(AT)rose-hulman.edu), Aug 03 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003731 | OEIS-A109730

Lesser of twin admirable numbers: k such that k and k+2 are both admirable numbers.

Conjecture: Sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Aug 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003732 | OEIS-A109733

Sequence is its own base-10 ASCII representation.

I conjecture that a(n) = 53 when n is congruent to 1, 3 or 5 (mod 8) and a(n) = 51 when n is congruent to 2 (mod 8). - Jonathan Cross (jcross(AT)juggler.net), Oct 14 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Nadia Heninger_, Aug 10 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003733 | OEIS-A109746

Number of distinct sequences of the type m^k mod n for n fixed, k>=1.

It appears that a(p)= p-1 (for prime p) and a(2p) = p-1 (for odd prime p), but this is only speculation. I (or someone else) need to calculate more values of a(n) to settle this conjecture. These conjectures hold for n <= 10000. (End) a(n) = A002322(n) + A051903(n) - 1 (conjectured). - _Ben Whitmore_, Jul 23 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | Bruce Corrigan (scentman(AT)myfamily.com), Aug 10 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003734 | OEIS-A109799

Primes p such that 2^p - 1 is a Chen prime.

Conjecture: 2^127 - 1 is the largest Mersenne-Chen prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Aug 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003735 | OEIS-A109812

a(1)=1; thereafter a(n) = smallest positive integer not among the earlier terms of the sequence such that a(n) and a(n-1) have no common 1-bits in their binary representations.

It appears that a(n)/n is bounded (it is probably less than 4 for all n), and n/a(n) is unbounded. See A352336, A352359, A352917-A352923 and the conjectures therein. - _David Broadhurst_, Apr 17 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Aug 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003736 | OEIS-A109852

a(1)=1; a(2) = 2; for n >= 1 and 1 <= k < 2^n, a(2^n+k) is the least multiple of a(2^n-k) not included earlier and a(2^n) is the least number not included earlier.

Conjecture: a(2^n) is prime if n is not 0 nor 2. Conjecture: for n>2, every odd prime >4 is encountered in order at a(2^n-1), a(2^n). - _Bill McEachen_, May 06 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 07 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003737 | OEIS-A109860

Triangle whose n-th row contains the next n numbers relatively prime to n! and not occurring in a previous row. a(1) = 1.

2 is the only prime that is not a member. Conjecture: There are finitely many composite terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 08 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003738 | OEIS-A109862

Palindromic primes p such that p's 10's complement is also a prime.

Conjecture: Sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 08 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003739 | OEIS-A109863

Primes arising as the 10's complement of A109862(n).

a(n) - 1 is a palindrome. Obviously 7, 5 and 3 are the only palindromic terms. Conjecture: Sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 08 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003740 | OEIS-A109864

Least palindromic prime beginning with n, or 0 if no such prime exists.

Conjecture: a(n) = 0 if and only if n > 3 and n begins with 2,4,6,8 or 5 (most significant digit).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 08 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003741 | OEIS-A109874

Largest exponent e such that n^e that divides A001142(n).

Conjectures: (1) If n is even and n = 2^r*m, m odd and >1, then a(n)= n-r-1. (2) If n = 2^r then a(n) = n-3. (3) If n is odd and composite then a(n) = n-2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 10 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003742 | OEIS-A109875

Chen primes p such that their p + 2 counterpart is a golden semiprime.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Aug 31 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003743 | OEIS-A109890

a(1)=1; for n>1, a(n) is the smallest number not already present which is a divisor or a multiple of a(1)+...+a(n-1).

Conjectured to be a rearrangement of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 13 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003744 | OEIS-A109904

a(1) = 5. a(n+1) is the greatest prime of the form k*(a(n)-k) + 1. The least prime occurs for k = 1 and a(n+1) = a(n) in that case if no other value of k gives a prime then the sequence terminates.

For the first five terms k = floor(a(n)/2). k can take values from 1 to floor(a(n)/2). It is conjectured that at least one value of k, 2 <= k < floor(a(n)/2) gives a prime and the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003745 | OEIS-A109907

Beginning with 7, a(n+1) = greatest prime of the form k*{a(n)-k}-1. If no prime is obtained the sequence ends at that point.

Conjecture: The sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003746 | OEIS-A109912

Beginning with 1, least multiple of a(n) not divisible by 5 such that no digit is common between a(n) and a(n+1).

Conjecture: Sequence is infinite with terms from a(12) onwards alternating between integers with the four digits 4,5,6,9 and integers with the remaining six digits 0,1,2,3,7,8. - _William Rex Marshall_, Jul 19 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003747 | OEIS-A109926

Least k such that k-2^r is prime for n values of r. Index of the first occurrence of n in A109925.

Conjecture: a(n)==0 (mod 3) for n > 2. Then n-2^k is not == 0 (mod 3) and a prime is more probable. - _Robert G. Wilson v_, Jul 21 2005 Conjecture: a(n+15)==0 (mod 30) for n > 4. - _Robert G. Wilson v_, Jul 21 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003748 | OEIS-A109943

Maximal number of distinct primes in the solution of the n X n generalization of the Gordon Lee puzzle.

The Gordon Lee puzzle asks for an n X n array of single digits such that as many distinct primes as possible are formed by joining consecutive digits in any horizontal, vertical or diagonal direction, forward or backward. a(4)=63 was proved in March 2005 by Mike Oakes. a(5) and a(6) are conjectured best values that resisted any improvement since 1998. a(5), a(6) and a(7) are conjectured best values that have resisted any improvement since 1998, including the joint effort of more than 100 participants in a programming contest in summer 2005. The best currently (September 2005) known lower bounds for the next terms are a(8)>=394 and a(9)>=527. - _Hugo Pfoertner_, Sep 21 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jul 05 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003749 | OEIS-A109969

First differences of A109968.

Conjecture: a(n) is bounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jul 06 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003750 | OEIS-A109994

Chen primes p such that their p + 2 counterpart is a Sarrus number (pseudoprime to base 2).

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Sep 01 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003751 | OEIS-A110000

Minimum number of polygonal pieces in a dissection of a regular n-gon to an equilateral triangle (conjectured).

Minimum number of polygonal pieces in a dissection of a regular n-gon to an equilateral triangle (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 11 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003752 | OEIS-A110034

Row sums of a characteristic triangle for the Fibonacci numbers.

Conjecture: |a(n)| = Sum_{k=1..n-1} A061646(k). - _J. M. Bergot_, Jun 10 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 08 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003753 | OEIS-A110037

Signed version of A090678 and congruent to A088567 mod 2.

Conjecture: a(n) = A073089(n) - A073089(n+1) for n >= 2. - _Alan Michael Gómez Calderón_, Aug 19 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003754 | OEIS-A110074

Numbers n such that sigma(n) = 2n - 3*phi(phi(n)).

Each prime number p of the form 2^k*3^j+1 where k & j are natural numbers is in the sequence because 2p-3*phi(phi(p))=2p-3*phi (2^k*3^j)=2p-3*(1-1/2)*(1-1/3)*2^k*3^j=2p-2^k*3^j=p+1=sigma(p). Conjecture: The sequence is infinite and 369 is the only composite term. I checked the validity of this conjecture up to 1.5*10^9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Jul 25 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003755 | OEIS-A110076

a(n) is the largest number m such that sigma(m)=10^n, or if there is no such m a(n)=0.

Conjecture: For n>3 a(n) is positive.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Jul 31 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003756 | OEIS-A110077

a(n) is the smallest number m such that sigma(m)=10^n and if there is no such m, a(n)=0.

Conjecture: For n>3 a(n) is positive.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Aug 01 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003757 | OEIS-A110081

Integers k such that the digit set D = (0, 1, -k) used in base 3 expansions of the form Sum_{ -N < j < oo} d_j 3^{-j}, all d_j in D, can represent all real numbers.

The sequence is known to be infinite and irregular and is conjectured to have density zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, based on correspondence from _R. K. Guy_ and Jeff Lagarias, Aug 31 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003758 | OEIS-A110108

Conjectured values of n such that sigma(n)+sigma(k)=sigma(n+k) has no solution.

Conjectured values of n such that sigma(n)+sigma(k)=sigma(n+k) has no solution.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003759 | OEIS-A110128

Number of permutations p of 1,2,...,n satisfying |p(i+2)-p(i)| not equal to 2 for all 0<i<n-1.

Note that the conjectured recurrence was based on the 600-term b-file, not the other way round. - _N. J. A. Sloane_, Dec 07 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roberto Tauraso_, A. Nicolosi and G. Minenkov, Jul 13 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003760 | OEIS-A110166

Row sums of Riordan array A110165.

Conjectures: for k >= 2, the number of k-ary words of length n such that the number of 1's <= the number of 0's is equal to the coefficient of x^n in the expansion of (k*x + 1/(1 + x))^n, and satisfies the recurrence u(0) = 1, u(1) = k-1 and n*u(n) = (k-2)*(2*n-1)*u(n-1) - k*(k-4)*(n-1)* u(n-2) + k^(n-1) for n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 14 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003761 | OEIS-A110172

Conjectured numbers j such that phi(j) + phi(k) = phi(j+k) has no solution k, where phi is Euler's totient function.

Conjectured numbers j such that phi(j) + phi(k) = phi(j+k) has no solution k, where phi is Euler's totient function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003762 | OEIS-A110194

First differences of A110095 (least nonprime sequence whose first differences are nonprime).

Conjecture: from a(5) on, are there only substrings <1 1 4> and <1 1 1 1 4 4 >?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexandre Wajnberg_ and _Eric Angelini_, Sep 06 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003763 | OEIS-A110198

Antidiagonal sums of number triangle A110197.

Conjecture: n*a(n) +(-3*n+1)*a(n-1) +n*a(n-2) +(-n+2)*a(n-3) +(3*n-5)*a(n-4) +(-n+2)*a(n-5)=0. - _R. J. Mathar_, Nov 15 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003764 | OEIS-A110199

a(n) = Sum_{k=0..floor(n/2)} Catalan(k).

Conjecture: -(n+2)*a(n) + (n+2)*a(n-1) + 4*(n-1)*a(n-2) + 4*(1-n)*a(n-3) = 0. - _R. J. Mathar_, Nov 09 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003765 | OEIS-A110278

Values of n such that the perfect deficiency (A109883) of n and n+1 are both squares.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Earls_, Jul 18 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003766 | OEIS-A110312

Minimum number of polygonal pieces in a dissection of a regular n-gon into a square (conjectured).

Minimum number of polygonal pieces in a dissection of a regular n-gon into a square (conjectured). The three related sequences A110312, A362938, A362939 are exceptions to the usual OEIS policy of requiring that all terms in sequences must be known exactly. These sequences are included because of their importance and in the hope that someone will establish the truth of some of the conjectured values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 11 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003767 | OEIS-A110320

Number of blocks in all RNA secondary structures with n nodes (an RNA secondary structure can be viewed as a restricted noncrossing partition).

Conjecture: a(n) = (A051292(n+2)-A051286(n+1))/2. - _Gerald McGarvey_, Jan 14 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jul 19 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003768 | OEIS-A110322

Row sums of A110321, a number triangle related to the Jacobsthal numbers.

Conjecture: a(n) +(-n-1)*a(n-1) -(2*n-1)*(n-1)*a(n-2) +2*(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Nov 11 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 20 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003769 | OEIS-A110356

Array read by antidiagonals: T(n,k) (n>=3, k>=3) = minimal number of polygonal pieces in a dissection of a regular n-gon to a regular k-gon (conjectured).

Array read by antidiagonals: T(n,k) (n>=3, k>=3) = minimal number of polygonal pieces in a dissection of a regular n-gon to a regular k-gon (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 11 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003770 | OEIS-A110358

Beginning with 3, the least prime which is the product of one or more previous terms + 2.

Conjecture: The sequence is infinite. Subbarao & Yip prove that if there is an integer m such that the equation Phi_2(x) = m has a unique solution, where Phi_2 is the 2nd Schemmel totient function (A058026), then x == 0 (mod a(n)^2) for each term in this sequence. They conjectured an analog to Carmichael's conjecture, that this equation has no unique solution to any integer m, and prove that any counterexample to this conjecture is > 10^120000, a bound calculated from the first 10000 terms of this sequence. A proof that this sequence is infinite would prove the conjecture. - _Amiram Eldar_, Mar 25 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 23 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003771 | OEIS-A110408

Smallest prime which is the repeated reverse concatenation of n followed by n (ending in n), where n == 1,3,7 or 9 (mod 10), or 0 if no such prime exists.

Conjecture: a(n) is 0 iff n is a palindromic prime == 1,3,7 or 9 (mod 10) and/or n ==0 (mod 3). As rev(n) followed by n is == 0 (mod 11), hence every nonzero term contains at least two reverse concatenations of n followed by n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Jul 30 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003772 | OEIS-A110434

Distinct numbers with least significant digit following the repetitive pattern 1,3,7,9,1,3,7,9,... such that every partial concatenation is a prime.

Conjecture: every number == 1,3,7,9 (mod 10) is a term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 02 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003773 | OEIS-A110454

Largest composite number obtained by concatenation of parts of a distinct partition of n, or 0 if no such number exist.

Conjecture a(n) = 0 only for n = 1, 2 and 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003774 | OEIS-A110456

Largest prime obtained by concatenation of parts of a distinct partition of n. 0 if no such number exist.

Conjecture a(n) = 0 only for n = 1, or n == 0 (mod 3), n is > 3. Subsidiary sequence (hard): Number of primes generated by concatenation of distinct partitions of n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003775 | OEIS-A110464

Oddly colossally abundant numbers.

The usual colossally abundant numbers (CANs) are defined in A004490; none of them is odd. Oddly CANs are defined similarly, except only odd primes are allowed in the factorization. It is conjectured that the ratio of consecutive oddly CANs is always a prime number. The sequence of those prime numbers is given in A110465.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 21 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003776 | OEIS-A110491

Expansion of e.g.f.: sqrt(1+2x)/sqrt(1-2x).

Conjecture: {a(n-1), n>=1} is the T-transform of A093178, where T maps a sequence {b(n), n>=1} to the sequence {c(n)} defined by c(n) = det(M_n), where M_n is the n X n matrix with elements M_n(i,j) = b(2*j) for i>j and M_n(i,j) = b(i+j-1) for i<=j. - _Lechoslaw Ratajczak_, Aug 04 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 22 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003777 | OEIS-A110499

Largest k such that n is the highest power in the factorization of binomial(2k,k).

Checked all k < 10^8. a(1) has been proved. a(2) and a(3) are conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 22 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003778 | OEIS-A110508

Expansion of 1/(1-(x+x^2)c(2x)), c(x) the g.f. of A000108.

Conjecture: n*(3*n-7)*a(n) -4*(3*n-4)*(2*n-5)*a(n-1) +2*n*(3*n-7) +(-45*n^2+177*n-160)*a(n-3) -4*(3*n-4)*(2*n-5)*a(n-4) = 0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003779 | OEIS-A110514

Expansion of (1 - x + x^2 + x^3)/(1 - x^2 - x^4 + x^6).

a(n) = Sum_{k=0..floor(n/2)} Jacobi(2^(n-2k), 2(n-2k)+1) [conjecture].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003780 | OEIS-A110515

Sequence array for (1 - x + x^2 + x^3)/(1 - x^4).

T(n, k) = if(k <= n, Jacobi(2^(n-k), 2(n-k)+1), 0) [conjecture].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003781 | OEIS-A110521

Expansion of 1/(1 - (x + x^2)c(3x)), c(x) the g.f. of A000108.

Conjecture: 2*n*a(n) + 2*(16-11n)*a(n-1) + 2*(42-11n)*a(n-2) + (32-21n)*a(n-3) + 5*(20-7n)*a(n-4) + 6*(7-2n)*a(n-5) = 0. - _R. J. Mathar_, Dec 10 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 24 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003782 | OEIS-A110545

a(n) is smallest positive integer m such that n divides either the numerator or the denominator of the (reduced) fraction H(m) = Sum_{k=1..m} 1/k.

Conjecture: a(n) <= n for all positive n's.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Sep 11 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003783 | OEIS-A110612

a(n) is least number k such that k/P > P^n, with n and P (=P(k)) as in A051283.

a(n) is the least number k such that A346418(k) = n+1 (conjectured). - _Amiram Eldar_, Jul 16 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rick L. Shepherd_, Jul 31 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003784 | OEIS-A110666

Sequence is {a(1,n)}, where a(m,n) is defined at sequence A110665.

Conjecture: G.f. x*(-1+2*x) / ((x-1)*(x^2-x+1)^2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Aug 02 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003785 | OEIS-A110667

Sequence is {a(2,n)}, where a(m,n) is defined at sequence A110665.

Conjecture: g.f.: -x*(-1+2*x) / ((x-1)^2*(x^2-x+1)^2). - _R. J. Mathar_, Oct 09 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Aug 02 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003786 | OEIS-A110668

Sequence is {a(3,n)}, where a(m,n) is defined at sequence A110665.

Conjecture: g.f.: x*(-1+2*x) / ((x^2-x+1)^2*(x-1)^3). - _R. J. Mathar_, Oct 09 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Aug 02 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003787 | OEIS-A110700

Number of zeros in the smallest prime with Hamming weight n (given by A061712).

Observe that a(n)=3 for n=6, 14, 30, 62, 126, 254, 510, 1022, ... which is A000918. Conjecture: a(n) is never greater than 3. - _T. D. Noe_, Mar 14 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Aug 03 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003788 | OEIS-A110707

Number of linear arrangements of n blue, n red and n green items such that there are no adjacent items of the same color (first and last elements considered as adjacent).

Conjecture: n^2*a(n) -3*n*(2*n-1)*a(n-1) -3*(n-1)*(5*n-12)*a(n-2) -8*(n-3)^2*a(n-3)=0. - _R. J. Mathar_, Jul 26 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Aug 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003789 | OEIS-A110711

Number of linear arrangements of n blue, n red and n green items such that first and last elements have the same color but there are no adjacent items of the same color.

Conjecture: -(n+1)*(n-2)*a(n) + (7*n^2 - 13*n + 4)*a(n-1) + 8*(n-2)^2*a(n-2) = 0. - _R. J. Mathar_, Nov 01 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Aug 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003790 | OEIS-A110734

Let n = a_1a_2...a_k, where the a_i are digits. a(n) = least multiple of n of the form a_1b_1a_2b_2...b_{k-1}a_k, obtained by inserting single digits b_i in the gaps; 0 if no such number exists.

Conjecture: for large n, a(n) is nonzero. For a k-digit number there are k-1 gaps and 10^(k-1) candidates, so the chances that one of them is a multiple of n increases with k on the one hand though the probability decreases because n becomes large.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003791 | OEIS-A110735

Let n = a_1a_2...a_k, where the a_i are digits. a(n) = least multiple of n of the type b_1a_1b_2a_2...a_kb_{k+1}, obtained by inserting single digits b_i in the gaps and both ends; 0 if no such number exists.

Conjecture: no term is zero. For a k-digit number there are k+1 spaces and 10^(k+1) candidates, so the chances that one of them is a multiple of n increases with k on the one hand although the probability decreases because n becomes large.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003792 | OEIS-A110772

Beginning with 2, least number not occurring earlier such that every partial concatenation is prime.

Conjecture: every odd number not divisible by 5 is a member.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003793 | OEIS-A110875

Minimum positive integer such that length of the gap described at A109322 is exactly n (in contrast to A109322 where the gap length is >= n).

Conjectures and open problems: 1) It is not known whether the sequence is infinite; 2) It is conjectured that for every n there is corresponding a(n). If Conjecture 2) were proved, Conjecture 1) would follow as a direct consequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Bojan Basic (bbasic(AT)ptt.yu), Sep 18 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003794 | OEIS-A111000

Integer part of zeta(zeta(n)).

Notice the previous term is almost 1/2 the next term. Conjecture: lim_{n -> infinity} zeta(zeta(n))/zeta(zeta(n+1)) = 1/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Sep 30 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003795 | OEIS-A111002

a(n) = gcd(f(n), f(n+1)) where f(n) = n^4 + n^2 + 1.

Conjectures from _Colin Barker_, Oct 06 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pahikkala Jussi_, Sep 30 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003796 | OEIS-A111015

Primes in A002535.

Conjecture: Starting with 1/1, there are infinitely many primes in the numerators of fractions built according to the rule: add top and bottom to get the new bottom, add top and 2k times bottom to get the new top, for k>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Oct 02 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003797 | OEIS-A111018

Indices of Catalan numbers that are divisible by 3.

Conjecture: The sequence contains all numbers n such that n and n+1 are both in the sequence A074940, or, equivalently, such that neither n nor n+1 is in the sequence A005836. - _L. Edson Jeffery_, Dec 02 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Sep 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003798 | OEIS-A111042

Odd terms of A059756.

Dowe (1989) conjectured that all Erdős-Woods numbers (A059756) are even. The first counterexamples were found in 2001 by Marcin Bienkowski, Mirek Korzeniowski and Krysztof Lorys, and independently by Nik Lygeros (Cégielski et al., 2003). - _Amiram Eldar_, Jun 20 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Victor S. Miller_, Oct 08 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003799 | OEIS-A111053

Number of permutations which avoid the patterns 1324 and (2143 with Bruhat restriction {2<->3}). Also the number of permutations whose graphs are acyclic.

Conjecture: n*(n+5)*a(n) +3*(20-13*n-3*n^2)*a(n-1) +2*(11*n^2+40*n-150)*a(n-2) +3*(40-11*n-3*n^2)*a(n-3) +2*(n+6)*(2*n-5)*a(n-4)=0. - _R. J. Mathar_, Aug 14 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steve Butler_, Oct 06 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003800 | OEIS-A111081

Successive generations of an alternating Kolakoski rule.

Conjecture : length of n-th string is asymptotic to c*(3/2)^n for some c.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 11 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003801 | OEIS-A111093

Like sequence A111072 but moving right by the squares of the sequence of positive integers.

Conjectures from _Chai Wah Wu_, Jan 28 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giorgio Balzarotti_ & _Paolo P. Lava_, Oct 13 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003802 | OEIS-A111097

Maximum likelihood of the number of distinguishable marbles in an urn if repeated random sampling of one marble with replacement yields n different marbles before the first repeated marble.

Conjecture: Second order difference is {2, 0, 2, 1, 0, 2, 1, 0, 2, ..., 1, 0, 2}. (End) Conjectured g.f.: x*(1 + 2*x^2 - x^3 + 2*x^4 - x^5)/((1 - x)^3*(1 + x + x^2)). - _Stefano Spezia_, Apr 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | Marc A. Brodie (mbrodie(AT)wju.edu), Oct 13 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003803 | OEIS-A111108

a(n) = A001333(n) - (-2)^(n-1), n > 0.

Conjecture: for odd primes p, p divides a(p). Note that (a(n)) and A001333 have different offsets. The conjecture follows from the formula A001333(n) = ((1-sqrt(2))^n + (1+sqrt(2))^n)/2. - _Max Alekseyev_, Oct 16 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Creighton Dement_, Oct 14 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003804 | OEIS-A111151

Numbers not of the form a^2 + b^3 + c^4 + d^5 for a,b,c,d >= 0.

It is conjectured that this list is complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_, Oct 19 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003805 | OEIS-A111152

Smallest number that is a sum of two n-th powers of positive rationals but not of two n-th powers of positive integers.

a(6) and a(7) are only conjectures; the earlier terms have (apparently) been proved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_, Oct 19 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003806 | OEIS-A111159

Numbers k such that A111075(k) is odd.

Conjecture: A111075(n) is odd whenever: (i) n = m^2 for all m>=1 such that 3 does not divide m, or (ii) n = 3*A028982(m) for all m>=1. Also, A028982 gives "Union of nonzero squares and twice squares." It is not difficult to show that the conjecture is true. - _Max Alekseyev_, Oct 12 2005

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003807 | OEIS-A111166

Let p < q be consecutive primes; p is in the sequence if p/(q-p) is a record.

Conjecture: Except for first term, the sequence coincides with A001359. This is true for all primes < 2*10^7. Conjecture: Except for first term, the sequence coincides with A001359. This is true for all primes < 7*10^16. Let n >= 2 be an integer, N +- 1 and M +- 1 two consecutive twin pairs where M>n*N. Finding a counterexample is the same as finding two consecutive primes P1 and P2 with n*N<P1<M and P2-P1 <= n. However, such gaps are unknown even for n=2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernardo Boncompagni_, Oct 21 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003808 | OEIS-A111184

Triangle T(n,k), 0<=k<=n, read by rows, given by [0, 2, 1, 3, 2, 4, 3, 5, 4, 6, 5, 7, 6, ...] DELTA [1, 0, 1, 0, 1, 0, 1, 0, 1, 0, 1, 0, 1, 0, ...] where DELTA is the operator defined in A084938.

Conjecture 1: Sum_{k=1..n} T(n,k)*q^(k-1) = A111528(q,n) for n > 0, q >= 0. Conjecture 2: Sum_{k=1..n} T(n,k)*(-1/q)^(k-1) = R(n,q)/q^(n-1) for n > 0, q > 0 where log(1 + x + q*x*[Sum_{k>=1} R(k,q)*x^k]) = Sum_{k>=1} R(k,q)/k*x^k. Conjecture 3: n-th row polynomial is x*v_n for n > 0 where we start with vector v of fixed length m with elements v_i = 1, and for i=1..m-1, j=i+1..m apply A := v_i (at the beginning of each cycle for i) and also apply A := A + v_j, v_j := (j-i+x-1)*v_j + A. Here last conjecture provides fast and simple algorithm, that allow to compute sums in the previous conjectures by substituting x = q and x = -1/q respectively. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Delépham_ and _Paul D. Hanna_, Oct 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003809 | OEIS-A111205

Nonsquare composite numbers k such that the difference between the closest squares surrounding k and k have a common divisor greater than 1.

Conjecture: The number of terms in this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Nov 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003810 | OEIS-A111213

Difference between the closest squares surrounding prime p is prime.

Conjecture: The number of terms in this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Nov 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003811 | OEIS-A111273

a(n) is the smallest divisor of triangular number T(n) := n(n+1)/2 not already in the sequence.

Conjecture: For all odd primes p, a(p-1) = p. Equivalently, it appears that if an initial 0 is appended (the smallest divisor of 0, the zeroth triangular number), then the fixed points in this include the odd primes. - _Enrique Navarrete_, Jul 24 2019 [Wording of the equivalent property corrected by _Peter Munn_, Jul 27 2019] The above conjecture is true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 03 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003812 | OEIS-A111338

Conjectured values of Cald's sequence A006509 after sorting and removal of duplicate zeros.

Conjectured values of Cald's sequence A006509 after sorting and removal of duplicate zeros. In other words, this is conjecturally the range of A006509. - _M. F. Hasler_, Mar 06 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 02 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003813 | OEIS-A111339

Conjectured to be the numbers that do not appear in A006509.

Conjectured to be the numbers that do not appear in A006509. Conjectured to be the complement of A111338.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 02 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003814 | OEIS-A111364

Index of first occurrence of n in A033093 (or 0 if not present).

Conjecture that the sequence contains no zeros; that is, A033093 is onto the nonnegative integers. Note that most of the values in the sequence are highly divisible, as would expected. a(7) appears to be the last odd value.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Nov 07 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003815 | OEIS-A111403

a(n) = f(f(n+1)) - f(f(n)), where f(m) = 2^m.

Conjecture: a(n) = A002716(2*n)-1. - _R. J. Mathar_, May 15 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003816 | OEIS-A111441

Numbers k such that the sum of the squares of the first k primes is divisible by k.

Conjecture: There are no terms that are 3 or 9 modulo 12. This seems to hold for all related sequences with even powers of primes, not just squares. Compare "sums of powers of primes divisibility sequences", linked below. - _Daniel Bamberger_, Dec 03 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefan Steinerberger_, Nov 14 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003817 | OEIS-A111488

Primes having only {0, 1, 3, 6} as digits.

Includes all repunit primes (A004022). Conjecture: an infinite sequence. Note twin primes: (11, 13), (101, 103), (311, 313), (1031, 1033), (1061, 1063), (1301, 1303), (6131, 6133), (10301, 10303), (10331, 10333), (13001, 13003).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Nov 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003818 | OEIS-A111495

Floor of 10^n/Li(10^n) - 1.

Conjecture: The difference between successive terms is between 2 and 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Nov 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003819 | OEIS-A111671

Smallest prime p > 3 such that p-1 has a prime factor > (p-1)^(n/(n+1)).

a(1) = 7 = A111668(1). Conjectures: (1) This is a subsequence of the safe primes (A005385), (2) lim{n --> oo}(a(n+1)/a(n)) = 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rick L. Shepherd_, Aug 15 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003820 | OEIS-A111677

Array of primes of the type k concatenated with 2n-1 where k < 2n-1. 1---> no prime 13,23 5---> no prime 17,37,47,67 19,29,59,79,89 211,311,811,911 113,313,613,1013,1213 15---> no prime 317,617,... ... Sequence contains the number of terms in the n-th rows.

Conjecture: a(n)=0 iff n== 3 (mod 5). [Corrected by _R. J. Mathar_, Aug 20 2007]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Aug 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003821 | OEIS-A111779

Sixth column (m=5) of unsigned triangle A111595.

Conjecture: +(-n+5)*a(n) +5*n*a(n-1) +n*(n-1)^2*a(n-2)=0. - _R. J. Mathar_, Jun 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Aug 23 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003822 | OEIS-A111802

a(n) = n^2 - n - 1 for n>3; a(1)=1; a(2)=2; a(3)=3.

Inspired by attempt to determine what incorrectly-named A000519 may really be. Conjecture: Sequence is number of different main-diagonal sums among Latin squares of order n. Confirmed for first five terms. Guaranteed to be an upper bound as the diagonal sum can only be in the range from n to n^2 inclusive and it is impossible for the sum to be n+1 or n^2-1. There is probably an easy proof that all other sums in this interval can be realized as the only restriction seems to be that it is not permissible for exactly n-1 numbers on a diagonal to be identical.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rick L. Shepherd_, Aug 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003823 | OEIS-A111883

Unsigned row sums of triangle A111595 (normalized rescaled squared Hermite polynomials).

Conjecture: a(n) -n*a(n-1) -n*(n-1)*a(n-2) +(n-1)*(n-2)^2*a(n-3)=0. - _R. J. Mathar_, Oct 05 2014 Remark: the above conjectured recurrence is true and can be easily obtained by the e.g.f. - _Emanuele Munarini_, Aug 31 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Aug 23 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003824 | OEIS-A111913

Expansion of x*(-2-3*x-x^2+x^7+x^8+2*x^4) / ((x-1)*(x+1)*(x^8-x^4+1)).

The above conjecture is correct, since x^24 = 1 mod (x-1)*(x+1)*(x^8-x^4+1). - _Charles R Greathouse IV_, Feb 07 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Creighton Dement_, Aug 20 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003825 | OEIS-A111967

Inverse of number triangle A101688.

Row sums are the Fredholm-Rueppel sequence A036987 [conjecture].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Aug 23 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003826 | OEIS-A111984

Expansion of g.f.: A(x) = (Sum_{n>=0} (2*n+1)*8^n*x^(n*(n+1)/2))^(1/3).

Conjecture: a(5*n+4) == 0 (mod 5) (checked up to n = 200). - _Peter Bala_, Feb 26 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 25 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003827 | OEIS-A112000

One half of third column (k=2) of triangle A111999.

Conjecture: +n*(4*n+5)*a(n) +(2*n+3)*(n+2)*(4*n+9)*a(n-1)=0. - _R. J. Mathar_, Jul 09 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Sep 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003828 | OEIS-A112001

One fourth of fourth column (k=3) of triangle A111999.

Conjecture: +n*(20*n^2+75*n+67)*a(n) +(2*n+5)*(n+2)*(20*n^2+115*n+162)*a(n-1)=0. - _R. J. Mathar_, Jul 09 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Sep 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003829 | OEIS-A112032

Denominator of 3/4 + 1/4 - 3/8 - 1/8 + 3/16 + 1/16 - 3/32 - 1/32 + 3/64 ...

Conjectures from _Colin Barker_, Apr 05 2013: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Aug 27 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003830 | OEIS-A112041

Numbers k such that 1*k + 1, 3*k + 1, 9*k + 1, 27*k + 1 are all primes.

Conjecture: all terms except the first are multiples of 10. - _Harvey P. Dale_, Mar 26 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Aug 26 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003831 | OEIS-A112243

Expansion of exp(x*(1+x)/(1-2*x)).

Conjecture: a(n) +(-4*n+3)*a(n-1) +2*(n-1)*(2*n-5)*a(n-2) +2*(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Feb 20 2015 The e.g.f. A(x) satisfies the differential equation (4*x^2 - 4*x + 1)*A'(x) + (2*x^2 - 2*x - 1)*A(x) = 0 with A(0) = 1. Mathar's conjectured recurrence follows easily from this. - _Peter Bala_, Jul 15 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Aug 29 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003832 | OEIS-A112283

Consider the array T(n, m) where the n-th row is the sequence of integer coefficients of A(x), where 1<=a(n)<=n, such that A(x)^(1/n) consists entirely of integer coefficients and where m is the (m+1)-th coefficient. This sequence is the first occurrence of 1 after the first column.

Conjecture: a(n)=n if n is 1, a prime or the square of the odd primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_ and _Robert G. Wilson v_, Aug 20 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003833 | OEIS-A112290

Numerator of sum{k=1 to n} 1/S(n,k), where S(n,k) is a Stirling number of the second kind.

Conjecture: a(n)/A112291(n) tends to 2 as n tends to infinity. - _Vaclav Kotesovec_, Jun 02 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Sep 01 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003834 | OEIS-A112293

Row sums of number triangle A112292.

Conjecture: a(n) - 2*n*a(n-1) + (2*n-3)*a(n-2) = 0. - _R. J. Mathar_, Nov 28 2014 a(n) = a(n-1) * (2*n-1) + 1 for n > 0 and a(0) = 1; that proves the conjecture of R. J. Mathar from Nov 28 2014; G.f. A(x) satisfies the equation: A(x) = 1/(1-x)^2 + A'(x) * 2*x^2/(1-x), where A' is the first derivative of A. - _Werner Schulte_, Oct 18 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 01 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003835 | OEIS-A112302

Decimal expansion of quadratic recurrence constant sqrt(1 * sqrt(2 * sqrt(3 * sqrt(4 * ...)))).

With Phi(z, p, q) the Lerch transcendent, define LP(n) = (1/n) * sum(Phi(1/2, n-k, 1) * LP(k), k=0..n-1), with LP(0) = 1. Conjecture: Lim_{n -> infinity} LP(n) = A112302.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Sep 02 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003836 | OEIS-A112388

a(n) is the smallest prime p such that p^n contains every digit.

Conjecture: a(n)=2 for all n>168. Checked up to n = 20000. - _Robert Israel_, Aug 28 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tanya Khovanova_, Dec 05 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003837 | OEIS-A112399

a(n) = Sum_{k=1..n, gcd(k,n)=1} mu(k), where mu(k) = A008683(k) (the Moebius function).

Conjectures from _Jamie Morken_, Sep 18 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Dec 06 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003838 | OEIS-A112486

Coefficient triangle for polynomials used for e.g.f.s for unsigned Stirling1 diagonals.

Conjecture: row polynomials are R(n,x) = Sum_{i=0..n} Sum_{j=0..i} Sum_{k=0..j} (n+i)!*Stirling2(n+j-k,j-k)*x^k*(x+1)^(j-k)*(-1)^(n+j+k)/((n+j-k)!*(i-j)!*k!). - _Mikhail Kurkov_, Apr 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Sep 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003839 | OEIS-A112647

a(n)=x is the smallest solution to abs(sigma(x+1)-sigma(x))=n, or 0 if no solution exists.

The known values plus conjectured zero values are: 14, 2, 1, 3, 6, 9, 5, 7, 62, 0, 13, 25, 22, 16, 12, 32, 11, 0, 104, 18, 837, 17, 19, 63, 46, 0, 28, 0, 116, 24, 58, 31, 2222, 0, 39, 242, 23, 0, 147, 0, 30, 675, 29, 35, 52, 0, 777, 0, 40, 0, 435, 0, 42, 36, 41, 0, 91, 0, 67, 0, 65, 99, 0, 195, 110, 80, 53, 48, 124, 0, 243, 0, 70, 97.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labos Elemer_, Sep 28 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003840 | OEIS-A112695

Number of steps needed to reach 4,2,1 in Collatz' 3*n+1 conjecture.

Number of steps needed to reach 4,2,1 in Collatz' 3*n+1 conjecture. a(n) = number of iterations of the Collatz 3*x+1 map applied to n until the conjectured 4,2,1 sequence is reached.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 31 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003841 | OEIS-A112700

Partial sum of Catalan numbers A000108 multiplied by powers of 6.

Conjecture: (n+1)*a(n) +(-25*n+11)*a(n-1) +12*(2*n-1)*a(n-2)=0. - _R. J. Mathar_, Jun 08 2016, verified by _Robert Israel_, Jun 28 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 31 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003842 | OEIS-A112701

Partial sum of Catalan numbers (A000108) multiplied by powers of 7.

Conjecture: (n+1)*a(n) +(-29*n+13)*a(n-1) +14*(2*n-1)*a(n-2)=0. - _R. J. Mathar_, Jun 08 2016 Conjecture verified using the d.e. (28*x^3-29*x^2+x)*y' + (42*x^2-16*x+1)*y=1 satisfied by the g.f. - _Robert Israel_, Aug 04 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 31 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003843 | OEIS-A112702

Partial sum of Catalan numbers A000108 multiplied by powers of 8.

Conjecture: (n+1)*a(n) +3*(-11*n+5)*a(n-1) +16*(2*n-1)*a(n-2)=0. - _R. J. Mathar_, Jun 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 31 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003844 | OEIS-A112703

Partial sum of Catalan numbers A000108 multiplied by powers of 9.

Conjecture: +(n+1)*a(n) +(-37*n+17)*a(n-1) +18*(2*n-1)*a(n-2)=0. - _R. J. Mathar_, Jun 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 31 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003845 | OEIS-A112725

Smallest positive palindromic multiple of 3^n.

a(0)=1; a(1)=3 and it is easily shown that for n>1, 10^3^(n-2)-1 is a palindromic multiple of 3^n(see comments line of A062567). So for each n, a(n) exists and for n>1, a(n)<=10^3^(n-2)-1. This sequence is a subsequence of A020485(a(n)=A020485(3^n)) and for all n, A062567(3^n)<=a(n) because for all n, A062567(n)<=A020485(n). Jud McCranie conjectures that for n>1 A062567(3^n) =10^3^(n-2)-1, if his conjecture were true then from the above facts we conclude that for n>1 a(n)=10^3^(n-2)-1, but we see that for 4<n<=18 a(n) is much smaller than 10^3^(n-2)-1, so his conjecture will be rejected.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Nov 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003846 | OEIS-A112726

First positive multiple of 3^n whose reverse is also a multiple of 3^n.

a(0)=1; a(1)=3 and it is easily shown that for n>1, 10^3^(n-2)-1 is a multiple of 3^n whose reverse is also a multiple of 3^n (see comments line of A062567), so for each n, a(n) exists and for n>1, a(n)<=10^3^(n-2)-1. This sequence is a subsequence of A062567, a(n)=A062567(3^n). Jud McCranie conjectures that for n>1, a(n)=10^3^(n-2)-1 (see comments line of A062567), but we see that for n>4, a(n) is much smaller than 10^3^(n-2)-1, so his conjecture is rejected. It seems that only for n=2,3 & 4 we have, a(n)=10^3^(n-2)-1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Nov 13 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003847 | OEIS-A112730

Numbers k such that the equation sigma(x)-phi(x)=k has at least one solution.

It is conjectured that if n>2 then all solutions of the equation sigma(x)-phi(x)=n (*) are less than or equal to (n-1)^2/4 and 2 is the only number n such that the equation (*) has infinitely many solutions. In fact in the case n=2 prime numbers are all solutions of (*). All numbers of the form

2p+1 where p is prime are in the sequence because p^2 is a solution for the equation sigma(x)-phi(x)=2p+1. All numbers of the form 3*2^n-1 are in the sequence because 2^(n+1) is a solution for the equation sigma(x)-phi(x)=3*2^n-1 and etc. The conjecture in the previous comment was established by Luke Pebody, see the Rivera link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Dec 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003848 | OEIS-A112799

Least odd number such that all greater odd numbers can be represented as sum of three integers with n distinct prime factors (conjectured).

Least odd number such that all greater odd numbers can be represented as sum of three integers with n distinct prime factors (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_ and _Ray Chandler_, Sep 19 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003849 | OEIS-A112826

Conjectured values of A112825 which are 0.

Conjectured values of A112825 which are 0. It is conjectured that there does not exist a Goldbach partition yielding a Goldbach "gap" of n as defined, for n=58,62,82,....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Sep 05 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003850 | OEIS-A112859

Primes such that the sum of the predecessor and successor primes is divisible by 29.

Under the Hardy-Littlewood k-tuple conjecture, a(n) ~ 28*n log n. - _Charles R Greathouse IV_, Apr 08 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jan 01 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003851 | OEIS-A112884

Number of bits required to represent binomial(2^n, 2^(n-1)).

G.f.: (-3*x^3 + 2*x^2 + x - 1)/((x - 1)^2*(2*x^2 + x - 1)). - Conjectured by _Harvey P. Dale_, Apr 06 2011 The conjectured formula 2^n - floor(n/2) and consequent g.f. are true (see links). - _Sela Fried_, Oct 03 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | Matt Erbst (matt(AT)erbst.org), Oct 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003852 | OEIS-A112971

Row sums of the matrix ((1,xc(x))^2 mod 2), where c(x) is the g.f. of A000108.

(1,xc(x)) is the Riordan array T(n,k)=[x^n](xc(x))^k. Conjectures: a(2^n)=a(2^(n+1)+1)=A005578(n);a(2^n-1)=a(3*2^n-1)=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 07 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003853 | OEIS-A112992

Numbers k such that ceiling(2^(2^k mod k)/3) is odd.

Conjecture: the odd values of ceiling(2^(2^k mod k)/3) are Jacobsthal numbers (and the even values are 1 plus a Jacobsthal number).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 08 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003854 | OEIS-A112994

Numbers whose cubes are exclusionary: numbers k such that k has no repeated digits and k and k^3 have no digits in common.

A number k with no repeated digits has an exclusionary cube k^3 if the latter is made up of digits not appearing in k. (This is a subsequence of A029785.) For the corresponding exclusionary cubes see A112993. Conjectured to be complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Oct 13 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003855 | OEIS-A113011

Decimal expansion of 1/(sqrt(e) - 1).

Conjecture: 1/(sqrt(e) - 1) = 1 + K_{n>=1} 2*n/(4*n^2-1)/1, where K is the Gauss notation for an infinite continued fraction. In the expanded form, 1/(sqrt(e) - 1) = 1 + 2/3/(1 + 4/15/(1 + 6/35/(1 + ...))) (see Beit-Halachmi and Kaminer). - _Stefano Spezia_, Dec 27 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, following a suggestion of Grover W. Hughes, Oct 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003856 | OEIS-A113023

Number of terms in A095810 which have n digits.

Conjecture: a(n) = 18*5^(n-2) for n>1. - _Mohammed Yaseen_, Jul 19 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_ and _Robert G. Wilson v_, Aug 27 2004 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003857 | OEIS-A113040

Number of solutions to +-p(1)+-p(2)+-...+-p(2n)=1 where p(i) is the i-th prime.

Conjecture: limit_{n->infinity} a(n)^(1/n) = 4. - _Vaclav Kotesovec_, Jun 05 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Floor van Lamoen_, Oct 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003858 | OEIS-A113047

a(n) = C(3n,n)/(2n+1) mod 3.

a(n) differs from 0 only when n=(3^j-1)/2, j>=0. [Conjecture confirmed by _Kevin Ryde_, Jun 23 2021; see links]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 11 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003859 | OEIS-A113049

Triangle of sums of Jacobsthal numbers related to binomial(4n,n)/(3n+1) mod 4.

First column is A007583(n+1). Main diagonal is A072197. Conjecture: A113048(n)=2 only if n=T(i,j).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 11 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003860 | OEIS-A113052

Binomial(5n,n)/(4n+1) mod 5.

Conjecture: for p prime, mod(C(p*n,n)/((p-1)*n+1),p) is the indicator function of the sequence (p^k-1)/(p-1). Conjecture is true: see link. - _Robert Israel_, Jun 11 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 12 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003861 | OEIS-A113191

Difference of two Lucas numbers.

Conjecture: L(m) - L(n) with m > 1 and m > n >= 0 is a perfect power but not a square only for (m,n) = (7,0), (5,2). This has been verified for n < m <= 500. Note that L(7) - L(0) = 29 - 2 = 3^3 and L(5) - L(2) = 11 - 3 = 2^3. - _Zhi-Wei Sun_, Jan 02 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Oct 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003862 | OEIS-A113251

Corresponds to m = 5 in a family of 4th-order linear recurrence sequences given by a(n) = m^4*a(n-4) + (2*m)^2*a(n-3) - 4*a(n-1), a(0) = -1, a(1) = 4, a(2) = -13 + 6*(m-1) + 3*(m-1)^2, a(3) = (-8+m^2)^2.

Conjecture: a(m, 2*n+1) is a perfect square for all m,n (see A113249).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Creighton Dement_, Nov 18 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003863 | OEIS-A113253

Corresponds to m = 7 in a family of 4th-order linear recurrence sequences given by a(n) = m^4*a(n-4) + (2*m)^2*a(n-3) - 4*a(n-1), a(0) = -1, a(1) = 4, a(2) = -13 + 6*(m-1) + 3*(m-1)^2, a(3) = (-8+m^2)^2.

Conjecture: a(m, 2*n+1) is a perfect square for all m,n (see A113249).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Creighton Dement_, Nov 18 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003864 | OEIS-A113256

Corresponds to m = 10 in a family of 4th-order linear recurrence sequences given by a(n) = m^4*a(n-4) + (2*m)^2*a(n-3) - 4*a(n-1), a(0) = -1, a(1) = 4, a(2) = -13 + 6*(m-1) + 3*(m-1)^2, a(3) = (-8+m^2)^2.

Conjecture: a(m, 2*n+1) is a perfect square for all m,n (see A113249).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Creighton Dement_, Nov 18 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003865 | OEIS-A113274

Record gaps between twin primes.

(2) Estimate for the actual size of the maximal gap that ends at p: maximal gap = a(log(p/a)-1.2), where a = 0.76*(log p)^2 is the average gap between twin primes near p, as predicted by the Hardy-Littlewood k-tuple conjecture. Both formulas (1) and (2) are derived from the Hardy-Littlewood k-tuple conjecture via probability-based heuristics relating the expected maximal gap size to the average gap. Neither of the formulas has a rigorous proof (the k-tuple conjecture itself has no formal proof either). In both formulas, the constant ~0.76 is reciprocal to the twin prime constant 1.32032...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernardo Boncompagni_, Oct 21 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003866 | OEIS-A113318

Numbers whose biquadrates (fourth powers) are exclusionary.

The number m with no repeated digits has an exclusionary fourth power m^4 if the latter is made up of digits not appearing in m. Is a subsequence of A111116. Conjectured to be complete. For the corresponding exclusionary biquadrates m^4, see A113317.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Oct 26 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003867 | OEIS-A113404

Record gaps between prime quadruplets.

Prime quadruplets (p, p+2, p+6, p+8) are densest permissible constellations of 4 primes (A007530). By the Hardy-Littlewood k-tuple conjecture, average gaps between prime k-tuples are O(log^k(p)), with k=4 for quadruplets. If a gap is larger than all preceding gaps, we call it a maximal gap, or a record gap. Maximal gaps may be significantly larger than average gaps. This sequence suggests that maximal gaps between prime quadruplets are O(log^5(p)). - _Alexei Kourbatov_, Jan 04 2012 (2) Estimate for the actual size of the maximal gap that ends at p: maximal gap = a*(log(p/a)-0.55), where a = 0.241*(log p)^4 is the average gap between quadruplets near p, as predicted by the Hardy-Littlewood k-tuple conjecture. Both formulas (1) and (2) are derived from the Hardy-Littlewood k-tuple conjecture via probability-based heuristics relating the expected maximal gap size to the average gap. Neither of the formulas has a rigorous proof (the k-tuple conjecture itself has no formal proof either). In both formulas, the constant ~0.241 is reciprocal to the Hardy-Littlewood 4-tuple constant 4.15118... (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernardo Boncompagni_, Oct 28 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003868 | OEIS-A113409

A transform of the central binomial coefficients A001405.

Conjecture: (n+2)*a(n)-2*(n+1)*a(n-1) +(n-4)*a(n-2) +2*a(n-3) +4*(2-n)*a(n-4)=0. - _R. J. Mathar_, Nov 07 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 28 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003869 | OEIS-A113411

Excess of number of divisors of 2n+1 of form 8k+1, 8k+3 over those of form 8k+5, 8k+7.

Conjectural g.f.s: A(x) = Sum_{n >= 0} (-1)^(n*(n-1)/2)*x^n/(1 - x^(2*n+1)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Oct 29 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003870 | OEIS-A113413

A Riordan array of coordination sequences.

T(n, k) = (Sum_{i=2..k+2} A137513(k+2, i) * (n-k)^(i-2)) / (k!) for 0 <= k < n (conjectured). - _Werner Schulte_, Oct 31 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 29 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003871 | OEIS-A113424

a(n) = (6*n)!/((3*n)!*(2*n)!*n!).

a(n) = [x^n] (F(x))^(60*n), where F(x) = 1 + x + 56*x^2 + 7355*x^3 + 1290319*x^4 + 264117464*x^5 + 59508459679*x^6 + ... appears to have integer coefficients. We conjecture that for k >= 1 the sequence defined by b_k(n) := [x^n] F(x)^(k*n) satisfies the above supercongruences for primes p >= 7. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Oct 31 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003872 | OEIS-A113459

Least number that begins an arithmetic progression of n numbers with the same prime signature.

Conjecture: For n > 1, a(n) = A007918(n). - _David Wasserman_, Jan 08 2006 I disagree with that conjecture! Ignoring the initial terms, this will agree with A007918 up to some point and then (presumably) drop below A007918. The initial term in the arithmetic progression (of length n) must be >= n, but it is likely to be less than A007918(n) if n is large. - _N. J. A. Sloane_, Oct 18 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Wasserman_, Jan 08 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003873 | OEIS-A113543

Numbers both squarefree and triangle-free.

The cardinality (count, enumeration) of these through n equals n - card{squarefree numbers <= n} - card{trianglefree numbers <= n} + card{numbers <= n which are both square and triangular} = n - card{numbers <= n in A005117} - card{numbers <=n in A112886} + card{numbers <= n in A001110}. "There is no known polynomial time algorithm for recognizing squarefree integers or for computing the squarefree part of an integer. In fact, this problem may be no easier than the general problem of integer factorization (obviously, if an integer can be factored completely, is squarefree iff it contains no duplicated factors). This problem is an important unsolved problem in number theory" [Weisstein]. Conjecture: there is no polynomial time algorithm for recognizing numbers which are both squarefree and triangle-free.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jan 13 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003874 | OEIS-A113547

Triangle read by rows: number of labeled partitions of n with maximin m.

Conjectures: T(n,3) = A007689(n-3). T(n,4) = 2^(n-4)+3^(n-3)+4^(n-4).- _R. J. Mathar_, Mar 13 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Jan 13 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003875 | OEIS-A113552

Beginning with 1, least divisor of the previous term not included earlier, otherwise the least multiple of the previous term having at least one prime divisor coprime to it and not included earlier.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 03 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003876 | OEIS-A113570

Consider the function f(k) in A110545, i.e., the smallest positive integer j such that k divides either the numerator or the denominator of the reduced Harmonic number H(j). This sequence lists numbers k where f(k)=k.

Conjecture: only prime powers are present (A025475) in addition to 1 and 2. Thus this sequence would be a proper subset of A000961.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_ and _Robert G. Wilson v_, Sep 29 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003877 | OEIS-A113571

Let H(n) be the reduced fraction Sum_{i=1..n} 1/i. a(n) is the least factor of H(n)'s numerator or denominator that doesn't divide either part of any earlier H(m).

Conjectured last occurrence of n: 1,3,11,25,137,49,363,761,7129,7381,83711, [sic] 6617,72072,1117 (is this A001008?)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_ and _Robert G. Wilson v_, Sep 29 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003878 | OEIS-A113572

Least n-digit multiple of n such that the r-th digit is prime if r is a prime else it is composite. The location and value of the most significant digit is 1. 0 if no such number exists.

a(5*t)=0, for all t>1. This is because the last digit of 5*t is always 0 or 5 yet we require this digit to be composite for t>1. There are no other zero terms below a(10000). Conjecture: No other term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 06 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003879 | OEIS-A113573

a(n) = A113572(n)/n.

a(5*t) = 0 for t > 1. Conjecture: No other term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 06 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003880 | OEIS-A113575

Least (2n-1)-digit palindromic prime == 1 (mod n), or 0 if no such prime exists.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 06 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003881 | OEIS-A113576

Least (2n-1)-digit palindromic prime == 1 (mod 2n-1), or 0 if no such prime exists.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 06 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003882 | OEIS-A113593

Minimal differences that appear in arithmetic progressions of primes.

Collapsed version of sequence A033188. If a conjecture about arithmetic prime progressions is correct (see A033188), this sequence is simply the primorial numbers with 150 inserted into the list. Zero is also included (since p,p is an arithmetic progression with difference 0).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Wild_, Jan 26 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003883 | OEIS-A113598

Least n-digit prime whose digit permutations yield at least n distinct primes, or 0 if no such prime exists.

Conjecture: No term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 07 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003884 | OEIS-A113599

Least n-digit multiple of n whose digit permutations yield at least n distinct multiples of n, or 0 if no such number exists.

Conjecture: No term is zero. The proof should be simple. The conjecture is true: There are 9*10^(n-1) n-digit numbers, meaning at least floor(9*10^(n-1)/n) n-digit multiples of n. There are binomial(n+9, 9) multisets of n digits. Thus, by the pigeonhole principle, one of these multisets contains at least ceiling(floor(9*10^(n-1)/n)/binomial(n+9, 9)) multiples of n, and this number is at least n whenever n > 3. - _James Rayman_, Feb 14 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 07 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003885 | OEIS-A113604

Define f(k) = k + sum of digits of n. a(n) is the first prime that results by applying f zero or more times to n, or 0 if no such prime exists.

Conjecture: a(n) = 0 only if n == 0 (mod 3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003886 | OEIS-A113610

Sum of digits of all composite numbers between prime(n) and prime(n+1).

Conjecture: For n > 9 if a(n) > a(n+1) < a(n+2) then prime(n) and prime(n+1) form a twin prime pair.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003887 | OEIS-A113682

Expansion of 2/(sqrt(1-2*x-3*x^2)*(1+x+sqrt(1-2*x-3*x^2))).

Conjectural congruences (working with an offset of 1): a(n*p^k) == a(n*p^(k-1)) (mod p^(2*k)) for prime p >= 5 and positive integers n and k. - _Peter Bala_, Mar 15 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 04 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003888 | OEIS-A113702

Trajectory of 10 under map k -> A111273(k).

Conjecture: Sequence is not periodic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 08 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003889 | OEIS-A113703

Retrograde trajectory of 10 under map k -> A111273(k).

Conjecture: Sequence is not periodic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 08 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003890 | OEIS-A113755

Sequence implicit in Jaroma's corollary to Nagura's theorem on primes.

Tightening the bounds on J. Bertrand's 1845 conjecture that for any integer n > 3 there exists at least one prime between n and 2*n-2 (proved by P. Tchebechev in 1852), Nagura proved that, for n >= 25, there exists at least one prime number between n and (6/5)*n. John H. Jaroma gives an elementary proof by induction of the corollary: prime(n) < (1.2)^n for n > 25. Equivalently, this sequence, implicit in Nagura and Jaroma, is always positive after a(25). The minimum is reached with min((1.2)^n - prime(n)) = (1.2)^17 - 59 = -36.8138889.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jan 16 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003891 | OEIS-A113768

a(1) = 1, a(n+1) = a(n) + floor(a(n)^(1/3)).

Conjecture: a(n) ~ (2/3)*n*sqrt((2/3)*n). - _José María Grau Ribas_, Feb 13 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jan 19 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003892 | OEIS-A113846

Palindromic primes that are the sum of 3 consecutive primes, the middle one palindromic.

Conjecture: For all terms in, the corresponding primes p_1 has the same number of digits and the same first digit as p_2 and p_3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Resta_, Jan 24 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003893 | OEIS-A113880

Variation on Recamán's sequence utilizing the four basic operations (/,-,+,*) in that order.

By a(1000) there are 3 division steps, 928 subtraction steps, 59 addition steps and 10 multiplication steps. It is unlikely that every number belongs to the sequence since there are many "holes". It is an open question if there are any repetitions after a multiplication step. Can anybody expand the series?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sergio Pimentel_, Jan 27 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003894 | OEIS-A113910

Integers of the form (Lucas(i+1) - 2*A006206(i+2))/(A006206(i+2) - A006206(i)), i > 2; Lucas = A000204.

It is conjectured that a(n+4) = A001359(n+2) for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Creighton Dement_, Jan 29 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003895 | OEIS-A113917

Maximal element of Image^inf({ 2 }) under repeated base-n zero-split squaring.

Conjecture: a(n) is finite for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo van der Sanden_ extending a suggestion from _David W. Wilson_, Jan 31 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003896 | OEIS-A113918

Cardinality of Image^inf({ 2 }) under repeated base-n zero-split squaring.

Conjecture: a(n) is finite for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo van der Sanden_ extending a suggestion from _David W. Wilson_, Jan 31 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003897 | OEIS-A113956

Expansion of (1/((1-4x)c(x)))/(1-x^2c(x)/sqrt(1-4x)), c(x) the g.f. of A000108.

Conjecture D-finite with recurrence: n*a(n) +2*(-5*n+4)*a(n-1) +(27*n-46)*a(n-2) +(5*n-2)*a(n-3) +(-57*n+188)*a(n-4) +2*(-21*n+83)*a(n-5) +4*(-2*n+9)*a(n-6)=0. - _R. J. Mathar_, Jan 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003898 | OEIS-A114021

Number of semiprimes between n and n + sqrt(n).

It appears that for n > 37 it is always true that a(n) > 0. The exponent can be reduced further. Since 597 + 597^(0.4129) > 611, leaping the record semiprime gap between 597 and 611, it seems that for n > 597 it is always true that there is a semiprime between n and n^(0.4129). It seems that for n > 2705 it is always true that there is a semiprime between n and n^(0.3509). These conjectures are related to the various sequences about semiprime gaps and the merit of such gaps.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jan 31 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003899 | OEIS-A114111

Where records occur in A111390.

Conjectures from _Chai Wah Wu_, Jul 27 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Nov 13 2005, Jan 01 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003900 | OEIS-A114112

a(1)=1, a(2)=2; thereafter a(n) = n+1 if n odd, n-1 if n even.

The induction hypotheses are: for i<n, a(2i)=2i+1, a(2i)=2i-1, A(2i)=i(2i+1), A(2i) = 2i^2+3i+2. This implies that when looking for a(2i), we have seen all the numbers 1 through 2i except 2i-1, so the two smallest candidates for a(2i) are 2i-1 and 2i+1. Since 2i-1 does not divide 2i^2-i-1, a(n)=2i-1. When looking for a(2i+1), we have seen all the numbers 1 through 2i already, so the two smallest candidates for a(2i+1) are 2i+1 and 2i+2. But 2i+1 does divide A(2i) and 2i+2 does not, so a(2i+1)=2i+2. QED The g.f. for the b(n) sequence is x*(x^3-3*x^2+2*x+2)/((1-x)*(1-x^2)). - Conjectured (correctly) by _Colin Barker_, Mar 04 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Nov 13 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003901 | OEIS-A114121

Expansion of (sqrt(1 - 4*x) + (1 - 2*x))/(2*(1 - 4*x)).

Conjecture: n*a(n) + 2*(3 - 4*n)*a(n-1) + 8*(2*n-3)*a(n-2) = 0. - _R. J. Mathar_, Nov 07 2012 Conjecture verified using the differential equation (16*x^2-8*x+1)*g'(x) + (8*x-2)*g(x)-2*x=0 satisfied by the G.f. - _Robert Israel_, Jul 27 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 13 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003902 | OEIS-A114190

Expansion of 1/(1+x*(1-x)*c(-2*x)), c(x) the g.f. of A000108.

Conjecture: n*a(n) + (7*n-10)*a(n-1) + 2*(14-3*n)*a(n-2) + (13*n-20)*a(n-3) + (66-23*n)*a(n-4) + 4*(2*n-7)*a(n-5) = 0. - _R. J. Mathar_, Dec 10 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003903 | OEIS-A114191

Expansion of 1/(1+x*c(-2*x)), c(x) the g.f. of A000108.

Conjecture: n*a(n) + (7*n-12)*a(n-1) + 4*(3-2*n)*a(n-2) = 0. - _R. J. Mathar_, Nov 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003904 | OEIS-A114194

Expansion of 1/(1+x*(2-x)*c(-2*x)), c(x) the g.f. of A000108.

Conjecture: 6*(n+1)*a(n) + (37n-29)*a(n-1) + 2*(45-41n)*a(n-2) + (47n-87)*a(n-3) + 4*(5-2n)*a(n-4) = 0. - _R. J. Mathar_, Dec 10 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 16 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003905 | OEIS-A114214

Diagonal sums of number triangle A114213.

Conjecture: a(n) = A007306(floor(n/2)+1). - _Georg Fischer_, Nov 28 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003906 | OEIS-A114379

Sums of p-th to the q-th prime where p and q are twin primes.

Conjecture: The number of terms in this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Feb 10 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003907 | OEIS-A114381

Sums of p-th to the q-th prime where p and q are consecutive primes.

The number of terms in this sequence is infinite since there is no largest prime number. Conjecture: There will always be an n and i such that a(n) >= a(n+i) or the sequence will alternate forever. Equality does take place in the small sample shown with the entry 991. Certainly the proof of an infinity many twin primes would be a strong probable proof of this assertion. My guess is the alternation would always occur when a twin prime is encountered and often for other consecutive primes such as those differing by 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Feb 10 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003908 | OEIS-A114464

Number of Dyck paths of semilength n having no ascents of length 2 that start at an even level.

Conjecture: (n+1)*a(n) +(-5*n+3)*a(n-1) +2*(3*n-7)*a(n-2) +2*(-3*n+11)*a(n-3) +(5*n-27)*a(n-4) +(-n+7)*a(n-5)=0. - _R. J. Mathar_, Nov 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Nov 29 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003909 | OEIS-A114467

Number of decimal digits in the numerator of the 10^n-th harmonic number.

Eric Weisstein's link conjectures that for both this sequence and A114468, a(n) ~ (log_10(e) = A002285)*10^n. - _Natalia L. Skirrow_, Jun 22 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Nov 29 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003910 | OEIS-A114534

a(n) is the permanent of the n X n matrix M defined as follows: if we concatenate the rows of M to form a vector v of length n^2, v_i is the i-th Fibonacci number.

Conjecture: The rank of the matrix M is 2 for every n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simone Severini_, Feb 15 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003911 | OEIS-A114584

Number of Motzkin paths of length n having no UHD's (U=(1,1), H=(1,0), D=(1,-1)).

Conjecture: +(n+2)*a(n) +(n+2)*a(n-1) +3*(-3*n+2)*a(n-2) +(-7*n+13)*a(n-3) +(4*n-13)*a(n-4) +6*(-n+5)*a(n-5) +(n-7)*a(n-6) +3*(n-8)*a(n-7)=0. - _R. J. Mathar_, Feb 16 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 09 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003912 | OEIS-A114589

Number of hill-free Dyck paths of semilength n+3 and having no peaks at even levels (a hill in a Dyck path is a peak at level 1).

Conjecture: 2*(n+4)*a(n) +2*(-n-1)*a(n-1) +3*(-3*n-4)*a(n-2) +(-8*n-11)*a(n-3) +3*(-n-1)*a(n-4)=0. - _R. J. Mathar_, Jul 02 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 11 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003913 | OEIS-A114590

Number of peaks at even levels in all hill-free Dyck paths of semilength n+2 (a hill in a Dyck path is a peak at level 1).

Conjecture: 2*(n+2)*a(n) +(-7*n-9)*a(n-1) -18*a(n-2) +2*(-7*n+19)*a(n-3) +4*(-2*n+3)*a(n-4)=0. - _R. J. Mathar_, Nov 15 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 11 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003914 | OEIS-A114646

Let M(n) be the n X n matrix m(i,j)=min(i,j) for 1<=i,j<=n then a(n) is the trace of M(n)^(-4).

(Conjecture) G.f.: F(x)=x*(1+45*x+24*x^2)/(1-x)^2. - _L. Edson Jeffery_, Jan 21 2012 (Conjecture) a(n)=2*a(n-1)-a(n-2), n>1, a(1)=1, a(2)=47. - _L. Edson Jeffery_, Jan 21 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 09 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003915 | OEIS-A114698

Let M(n) be the n X n matrix m(i,j)=min(i,j) for 1<=i,j<=n then a(n) is the trace of M(n)^(-3).

(Conjecture) G.f.: F(x)=(1+16*x+3*x^2)/(1-x)^2. - _L. Edson Jeffery_, Jan 21 2012 (Conjecture) a(n)=2*a(n-1)-a(n-2), n>1, a(0)=1, a(1)=18. - _L. Edson Jeffery_, Jan 21 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 09 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003916 | OEIS-A114726

Fixed points of permutation A114650.

Conjecture: Sequence is infinite and lim n -> infinity a(n+1)/a(n) = e = 2.718281828...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Dec 29 2005, Jan 10 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003917 | OEIS-A114782

Smallest prime of the form 10^k + prime(n), k >= d, the number of digits in prime(n). 0 if no such primes exist. a(n) = 0 if prime(n) + 1 == 0 (mod 3).

Conjecture a(n) = 0 iff prime(n) + 1 == 0 (mod 3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amarnath Murthy_, Nov 17 2005 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003918 | OEIS-A114851

Number of lambda calculus terms of size n, where size(lambda x.M) = 2 + size(M), size(M N) = 2 + size(M) + size(N), and size(V) = 1 + i for a variable V bound by the i-th enclosing lambda (corresponding to a binary encoding).

Conjecture: (n+2)*a(n) + 2*(-n-1)*a(n-1) + (-n+2)*a(n-2) + 4*(n-2)*a(n-3) + (-5*n+18)*a(n-4) + 2*(n-4)*a(n-5) + (n-6)*a(n-6) = 0. - _R. J. Mathar_, Mar 04 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John Tromp_, Feb 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003919 | OEIS-A114887

Multiperfect numbers sigma(n) = k*n, which are divisible by the sum of their prime factors without repetition.

Conjecture: the sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sven Simon_, Feb 19 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003920 | OEIS-A114888

Multiperfect numbers, sigma(n) = k*n, which are divisible by their sums of prime factors with and without repetition.

Conjecture: the sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sven Simon_, Feb 19 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003921 | OEIS-A114889

a(1)=1 and, for n>1, a(n) is the smallest integer greater than a(n-1) such that a(n)+a(i) is not a power of 3, for i=1,..., n-1.

The differences of {a(n)}, together with a conjectured formula for them, is given in A114890.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jan 04 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003922 | OEIS-A114890

First differences of A114889.

Conjecture. a(1)=2 and, for n>1, a(n)=1+3+9+...+3^(k-1) + 2, if n=3^k, a(n)=a(n-3^k), if 3^k < n < 2*3^k - 1, else a(n)=1. (This has been verified for n<=3000.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jan 04 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003923 | OEIS-A114948

a(n) = n^2 + 10.

Conjecture: n^2 + 10 != x^k for all n,x, and k > 1. The conjecture is true: See Cohn. - _James Rayman_, Feb 14 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Feb 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003924 | OEIS-A114965

a(n) = n^2 + 34.

Conjecture: n^2 + 34 != x^k for all n,x and k > 1. The conjecture is true: See Cohn. - _James Rayman_, Feb 14 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Feb 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003925 | OEIS-A114976

Number of subsets of {1,2,....,n} with an arithmetic mean that is an integer and also a divisor of n.

a(n) = 2 iff n is prime, just as for the number of divisors of n and also, at least for the very first terms, a(n)=odd iff n is a square: these observations might suggest conjectures on a deeper relationship with A000005.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Feb 22 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003926 | OEIS-A115016

a(n) is the smallest number k that has a shortest addition chain whose length A003313(k) = A003313(n*k), or 0 if this never happens.

It was conjectured that no shortest addition chains exist such that A003313(m)=A003313(m*2^k) for k>1. This is now known to be false, since a(4) != 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Feb 26 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003927 | OEIS-A115038

a(n) is the smallest x such that x^2 + prime(n) is a perfect power.

Conjecture: There will always be an x,y,n such that x^2 + p = y^n for all primes p. In otherwords, there is a one-one mapping of the prime numbers to y^n - x^2 for some x,y,n. Therefore primes of the form y^n - x^2 are infinite in number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Feb 26 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003928 | OEIS-A115047

a(3)=1; a(n) = Sum_{i=1..n-3} binomial(n-4,i-1)*binomial(n,i+1)*a(i+2)*a(n-i)*i*(n-i-2)/(2*(n-1)) for n > 3.

I conjecture that a(n) is divisible by n if n is a prime. - _Michael Somos_, Sep 20 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, based on an email message from John McKay, Feb 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003929 | OEIS-A115136

Row sums of triangle A113647.

Conjecture: (n+2)*a(n) +(-11*n-4)*a(n-1) +(25*n-42)*a(n-2) +5*(-n+12)*a(n-3) +2*(-13*n+20)*a(n-4) +8*(2*n-7)*a(n-5)=0. - _R. J. Mathar_, Nov 10 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jan 13 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003930 | OEIS-A115156

Smallest number having exactly n ones in binary representation and also exactly n prime factors (counted with multiplicity).

Does a(n) exist for every n? It exists for large enough n due to a result of Drmota, Mauduit, & Rivat, see A061712. _T. D. Noe_'s conjecture there implies that a(n) < 4*4^n. - _Charles R Greathouse IV_, Jul 30 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jan 14 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003931 | OEIS-A115256

Diagonal sums of correlation triangle of central binomial coefficients.

Conjecture: n*a(n) + 2*(-2*n+1)*a(n-1) + 4*(-n+1)*a(n-2) + 3*(5*n-8)*a(n-3) + 2*(2*n-1)*a(n-4) + 4*(n-1)*a(n-5) + 8*(-2*n+3)*a(n-6) = 0. - _R. J. Mathar_, Jun 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 18 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003932 | OEIS-A115362

Row sums of ((1,x) + (x,x^2))^(-1)*((1,x)-(x,x^2))^(-1) (using Riordan array notation).

Dirichlet g.f. (conjectured): zeta(s)/(1-2^(-2s)). - _Ralf Stephan_, Mar 27 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003933 | OEIS-A115399

Expansion of c(x^2-x^3), c(x) the g.f. of A000108.

Conjecture: (n+2)*a(n) +(-n-2)*a(n-1) +4*(n-1)*a(n-2) +2*(-4*n+7)*a(n-3) +2*(2*n-5)*a(n-4)=0. - _R. J. Mathar_, Feb 20 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 14 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003934 | OEIS-A115526

Inverse of number triangle A115524.

Row sums are number of zeros in binary expansion of n+2, that is, A023416(n+2) [conjecture].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 25 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003935 | OEIS-A115770

Numbers k such that 7*k = A048720(11,k), where A048720 is carryless base-2 multiplication.

Conjecture: also numbers k such that 5*k = A048720(13,k). This has been checked for all k in range 0 .. 2^25. - _Antti Karttunen_, Dec 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 30 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003936 | OEIS-A115792

a(n) = ceiling(g(A000073(n))) with g(k) = (k-1)^2/(4k).

Conjectures from _Chai Wah Wu_, Dec 21 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Mar 13 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003937 | OEIS-A115801

Numbers k such that 9*k = A048720(25,k), where A048720 is carryless base-2 multiplication.

Conjecture: also k such that 15*k = A048720(19,k). This has been checked for all k in range 0 .. 2^25. - _Antti Karttunen_, Dec 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 01 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003938 | OEIS-A115967

Expansion of 1/(2*sqrt(1-2*x-3*x^2) - 1).

Conjecture: 3*n*a(n) + (-14*n+9)*a(n-1) + (-5*n+3)*a(n-2) + 12*(4*n-9)* a(n-3) + 36*(n-3)*a(n-4) = 0. - _R. J. Mathar_, Nov 15 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 03 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003939 | OEIS-A116020

Numbers n such that sigma(n) - phi(n) is a repdigit greater than 2.

(I). If both numbers p=4*10^n+1 & q=(4*10^n-13)/9 are primes then m=p*q is in the sequence because sigma(m)-phi(m)=8*(10^(n+1)-1)/9 is a repdigit number. Conjecture: 123, 17243 & 1772443 are all such terms. - _Farideh Firoozbakht_, Aug 24 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Resta_, Feb 13 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003940 | OEIS-A116078

Column 0 of triangle A116077.

Conjecture: a(n) = 2^(2*n)*(Sum_{j=1..n+2-floor((n+3)/2)} (cos(j*Pi/(n+3)))^(2*n)). - _L. Edson Jeffery_, Nov 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 04 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003941 | OEIS-A116083

Numbers k such that phi(sigma(k)) - sigma(phi(k)) = 1.

Also if n is a natural number less than 6 then 3*2^(2^n-1) is in the sequence, the proof is similar to the case m=2^(2^n-1). Note that all known terms of the sequence are the ten numbers m*2^(2^n-1) m=1 & 3 and n=1,2,3,4 & 5. Conjecture: There is no other term. - _Farideh Firoozbakht_, Mar 24 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Mar 12 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003942 | OEIS-A116086

Perfect powers n with no primes between n and the next larger perfect power, which is in A116455.

No other n<10^12. There is a conjecture that this sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003943 | OEIS-A116184

Numbers n such that 37^3 divides the numerator of generalized harmonic number H(36,n) = Sum[1/k^n, {k,1,36}].

Note the pattern in the first differences of a(n): {34,2,34,2,36,36,1,35,2,34,2,34,4,32,4,32,6,30,6,30,8,28,8,28,10,26,10,26,12,24,12,24,14,22,14,22,16,20,16,20,18,18,18,18,20,16,20,16,22,14,22,14,24,...}. Conjecture: All terms of the arithmetic progression 3+36k belong to a(n). Prime terms in a(n) are {3, 37, 73, 443, 739, 887, 1109, ...}. It appears that all primes in a(n) that are greater than 37 are of the form 37k-1. For example, 73 = 37*2-1, 443 = 37*12-1, 739 = 37*20-1, 887 = 37*24-1, 1109 = 37*30-1. Many terms in a(n) are the multiples of 37. There are terms of the form 37*m with m = {1,3,4,5,7,9,11,13,15,17,19,21,23,25,27,29,31,33,37,39,41,...}. Note that 37^4 divides the numerator of generalized harmonic number H(36,n) for n = {111, 147, 1047, 1369, 1443, 1479, ...} = {3*37, 3+4*36, 3+29*36, 37^2, 3+40*36, 3+41*36, ...}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Apr 08 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003944 | OEIS-A116383

Row sums of number triangle A116382.

Conjecture: n*a(n) + (n-2)*a(n-1) + (-13*n+20)*a(n-2) + (-9*n+22)*a(n-3) + 4*(14*n-45)*a(n-4) + 20*(n-3)*a(n-5) + 80*(-n+5)*a(n-6) = 0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 12 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003945 | OEIS-A116385

Expansion of e.g.f. Bessel_I(2,2x) + 2*Bessel_I(3,2x) + Bessel_I(4,2x).

Conjecture: (n+4)*a(n) -2*a(n-1) +(-7*n-8)*a(n-2) +6*a(n-3) +12*(n-2)*a(n-4)=0. - _R. J. Mathar_, Jun 13 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 12 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003946 | OEIS-A116387

Expansion of 1/(sqrt(1-2*x-3*x^2)*(2-M(x))), where M(x) is the g.f. of the Motzkin numbers A001006.

Conjecture: n*(17*n-142)*a(n) + (17*n^2 + 95*n + 138)*a(n-1) + (-391*n^2 + 2488*n - 2908)*a(n-2) + (-17*n^2 - 603*n + 1892)*a(n-3) + 2*(697*n-2021)*(n-4)*a(n-4) + 60*(17*n-47)*(n-4)*a(n-5) = 0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 12 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003947 | OEIS-A116388

Expansion of 1/((1+x*(1-M(x)))*sqrt(1-2*x-3*x^2)), M(x) the g.f. of A001006.

Conjecture: n*a(n) + 7*(-n+1)*a(n-1) + 2*(4*n-9)*a(n-2) + (25*n-58)*a(n-3) + (-18*n+65)*a(n-4) + (-52*n+199)*a(n-5) + (-31*n+135)*a(n-6) + 6*(-n+5)*a(n-7) = 0. - _R. J. Mathar_, Jun 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 12 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003948 | OEIS-A116391

Expansion of 1/((1+x)*(sqrt(1-4*x^2)-x)).

Conjecture D-finite with recurrence: n*a(n) +(n)*a(n-1) +3*(-3*n+4)*a(n-2) +3*(-3*n+4)*a(n-3) +20*(n-3)*a(n-4) +20*(n-3)*a(n-5)=0. - _R. J. Mathar_, Jan 23 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 12 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003949 | OEIS-A116408

E.g.f. exp(x)*(Bessel_I(2,2*x) - Bessel_I(3,2*x) + Bessel_I(4,2*x)).

Conjecture: 2*(n+4)*(n-23)*a(n) + 2*(n^2+64n+79)*a(n-1) +(-59n^2+232n+740)*a(n-2) +(64n^2-585n+1019)*a(n-3) +3(n-3)*(41n-124)*a(n-4)=0. - _R. J. Mathar_, Dec 10 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 13 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003950 | OEIS-A116409

Expansion of (1-x-2x^2+sqrt(1-2x-3x^2))/(2(1-x)(1-2x-3x^2)).

Conjecture: n*a(n) +2*(-3*n+2)*a(n-1) +2*(4*n-5)*a(n-2) +6*(n-2)*a(n-3) +9*(-n+2)*a(n-4)=0. - _R. J. Mathar_, Nov 11 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 13 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003951 | OEIS-A116421

a(n) = 2^(n-1)*binomial(2*n-1,n-1)^2.

n^2*a(n) - 8*(2*n-1)^2*a(n-1) = 0 for n >= 2. - conjectured by _R. J. Mathar_, Nov 16 2011, corrected and proved by _Victor X. Ocampo_, Aug 04 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 14 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003952 | OEIS-A116473

Number of partitions with exactly n subpartitions.

Conjecture that lim_{n->infinity} a(n) = infinity. Definitely a(n) > 2 for n > 8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Mar 17 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003953 | OEIS-A116485

Number of permutations in S_n that avoid the pattern 12453 (or equivalently, 31245).

Conjecture: a(n) + A158423(n) = n!. - _Benedict W. J. Irwin_, Mar 15 2016 The conjecture is true: All that is needed is to show that 23145 is Wilf-equivalent to 31245, but that's obvious since they are inverses. - _Doron Zeilberger_ and _Yonah Biers-Ariel_, Feb 26 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | Zvezdelina Stankova (stankova(AT)mills.edu), Mar 19 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003954 | OEIS-A116492

Number of partitions of n into distinct odious numbers.

Conjecture: a(n) ~ c * exp(Pi*sqrt(n/6)) / n^(3/4), where c = 0.26864248.... - _Vaclav Kotesovec_, Jul 06 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian G. Bower_, Feb 17 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003955 | OEIS-A116518

Odd numbers k such that k and phi(k) have the same number of divisors.

It is an open question whether this sequence contains infinitely many terms; see Bellaouar et al., 2023. - _Allen Stenger_, Feb 16 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Mar 24 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003956 | OEIS-A116628

Positions where A116624 is a power of 2.

Conjecture: all the powers of 2 occur in A116624 (implying that none occur in A116625) and they occur in ascending order, in which case A116624(a(n)) = A000079(n-1). Checked up to n=16.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_ and _Antti Karttunen_, Feb 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003957 | OEIS-A116637

G.f. satisfies: A(x) = x/series_reversion(x/G(x)) where A(x) + A(-x) = 2*G(x^2) and G(x) is the g.f. of A046646.

Conjecture: 4*n*(n+1)*(3*n-1)*a(n) -36*n*a(n-1) -3*(3*n-5)*(3*n+2)*(3*n-4)*a(n-2)=0. - _R. J. Mathar_, Jun 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 19 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003958 | OEIS-A116649

Positions where A116648(i) is greater than A116648(i-1).

Equivalently, from i>1 onward those i where A116626(i) = A116648(i-1). Conjecture: all the terms are odd.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003959 | OEIS-A116674

Triangle read by rows: T(n,k) is the number of partitions of n into odd parts and having exactly k distinct parts (n>=1, k>=1).

Conjecture: Also the number of strict integer partitions of n with k maximal runs of consecutive parts decreasing by 1. - _Gus Wiseman_, Jun 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Feb 22 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003960 | OEIS-A116731

Number of permutations of length n which avoid the patterns 321, 2143, 3124; or avoid the patterns 132, 2314, 4312, etc.

Conjecture: also counts the distinct pairs (flips, iterations) that a bubble sort program generates when sorting all permutations of 1..n. - _Wouter Meeussen_, Dec 13 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lara Pudwell_, Feb 26 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003961 | OEIS-A116861

Triangle read by rows: T(n,k) is the number of partitions of n such that the sum of the parts, counted without multiplicities, is equal to k (n>=1, k>=1).

Conjecture: Reverse the rows of the table to get an infinite lower-triangular matrix b with 1's on the main diagonal. The third diagonal of the inverse of b is minus A137719. - _George Beck_, Oct 26 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Feb 27 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003962 | OEIS-A116863

Array used to find the eigenvalues of the quadratic Casimir operator for the Lie algebras A_n = su(n+1), n>=1.

The row sums are conjectured to be 3*A002415(n+1)= [3, 18, 60, 150, 315, 588, 1008, 1620, 2475, 3630,...]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Mar 24 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003963 | OEIS-A116881

Row sums of triangle A116880 (generalized Catalan C(1,2)).

Conjecture: n*(3*n-4)*a(n) +(-21*n^2+43*n-10)*a(n-1) -4*(3*n-1)*(2*n-1)*a(n-2)=0. - _R. J. Mathar_, Jun 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Mar 24 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003964 | OEIS-A116943

Number of 4s digits plus non-final 3s digits 3 base 5 expansion of 2^n.

In his comment on A038003 Frank Adams-Watters conjectures "that 2^n contains such a base 5 digit for n>=9. This is almost certainly true." That is equivalent to a(n) > 0 for n>=9, which is also equivalent to A094389(n) = 5 where A094389 is last decimal digit of the odd Catalan number A038003(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Mar 23 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003965 | OEIS-A116944

a(1) = 1; for n>1, a(n) = (a(n-1)+1)^n.

There was a conjecture that this recurrence would produce the sequence A068894, but this is not the case.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vim Wenders_, Apr 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003966 | OEIS-A117054

Number of ordered ways of writing n = i + j, where i is a prime and j is of the form k*(k+1), k > 0.

Based on a posting by _Zhi-Wei Sun_ to the Number Theory Mailing List, Mar 23 2008, where he conjectures that a(2n+1) > 0 for n >= 2. _Zhi-Wei Sun_ has offered a monetary reward for settling this conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 15 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003967 | OEIS-A117078

a(n) is the smallest k such that prime(n+1) = prime(n) + (prime(n) mod k), or 0 if no such k exists.

a(n) = 0 only for primes 2, 3 and 7. Conjecture: 2, 3 and 7 are the only primes for which log(A000040(n)) < sqrt(A001223(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémi Eismann_, Apr 18 2006, Dec 10 2006, Feb 14 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003968 | OEIS-A117081

a(n) = 36*n^2 - 810*n + 2753, producing the conjectured record number of 45 primes in a contiguous range of n for quadratic polynomials, i.e., abs(a(n)) is prime for 0 <= n < 44.

a(n) = 36*n^2 - 810*n + 2753, producing the conjectured record number of 45 primes in a contiguous range of n for quadratic polynomials, i.e., abs(a(n)) is prime for 0 <= n < 44.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Apr 17 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003969 | OEIS-A117098

Numbers k such that nextprime(5*k)<5*nextprime(k) (if p is prime then nextprime(p)=p).

denote f(x)=nextprime(x) let A(p)={n/f(p*n)<f(p)*f(n)} and A(q)={n/f(q*n)<f(q)*f(n)} (q,p are prime) if q<p all terms of A(q) are in A(p). if B(p)={n/f(p*n)>f(p)*f(n)} I conjectured that when p --> infinity, A(p) = All composite positive integers when p --> infinity, B(p) = All prime numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Mohammed Bouayoun (mohammed.bouayoun(AT)sanef.com), Apr 18 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003970 | OEIS-A117119

Number of partitions of 2*n into two odd prime powers.

Conjecture: For all n, a(n) > 0; a(n) > A002375(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 15 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003971 | OEIS-A117149

Trajectory of 4 under map k -> A094077(k).

Conjecture: Sequence is not periodic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Feb 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003972 | OEIS-A117150

Retrograde trajectory of 4 under map k -> A094077(k).

Conjecture: Sequence is not periodic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Feb 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003973 | OEIS-A117186

Expansion of (1+x)c(x^2)/((1-xc(x^2))*sqrt(1-4x^2)), c(x) the g.f. of A000108.

Conjecture: a(n) = A058622(n) + A058622(n+1). [_R. J. Mathar_, Nov 21 2008] Conjecture: -(n+1)*a(n) +(n+1)*a(n-1) +2*(3*n-2)*a(n-2) -4*n*a(n-3) +8*(3-n)*a(n-4)=0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 01 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003974 | OEIS-A117187

Expansion of (1+x)c(x^2)/((1-x^2*c(x^2))sqrt(1-4x^2)), c(x) the g.f. of A000108.

Conjecture D-finite with recurrence: (n+4)*(31n-12)*a(n) +(37n^2+24n-156)*a(n-1) -4*(31n^2+81n+24)*a(n-2) -4*(37n+24)*(n-1)*a(n-3) =0. - _R. J. Mathar_, Dec 10 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 01 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003975 | OEIS-A117215

Number of divisors of x^n-1 having the maximal height A114536(n).

Let p be a prime. Then a(p)=4 because the divisors are x^p-1, x^(p-1)+x^(p-2)+...+1, x-1 and 1. Similarly, a(p^k)=2^(k+1). For n=p*2^k, a(n)=2. For odd primes p

and q, a(pq)=1. Conjectures: if n is odd and squarefree, then a(n)=1; if n/2^k is odd and squarefree for k>0, then a(n)=2. All the divisors of x^n-1 are products of cyclotomic polynomials cyclo(d) for various d. When n is the product of distinct odd primes p1..pk, it appears that each cyclotomic index has the form d=p1^e1...pk^ek, where the ei are either 0 or 1 and sum(ei) is odd.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 03 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003976 | OEIS-A117223

Numbers n such that Phi(n,x) is a flat cyclotomic polynomial of order three.

Kaplan shows that the sequence also includes pqr if r = +-1 (mod pq). Sequence A160353 lists the subsequence of all odd numbers of this form, while A160355 lists the elements which are not of this form. More cases are covered by _David Broadhurst_'s conjectures, cf. link. - _M. F. Hasler_, May 15 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 04 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003977 | OEIS-A117231

Decimal expansion of Integral_{x=0..1} cos((log x)/x)/x dx.

Conjecturally, the value of the Borel sum of the divergent series Sum_{n >= 1} (-1)^(n+1)*(2*n)^(2*n-1). See Finch, p. 62. - _Peter Bala_, Dec 28 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Mar 05 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003978 | OEIS-A117301

a(n) = prime(n+3)*prime(n) - prime(n+1)*prime(n+2).

Cino Hilliard conjectures that this ratio converges and that there are infinitely many elements in the sequence whose absolute value is 12. It appears that the positions of negative multiples of 12 are given by A064026(n+1) for n >= 1. If so, then Hilliard's conjecture is true, and a further conjecture is that if k >= 2 then there are infinitely many multiples of -12*k in this sequence. - _Clark Kimberling_, Jan 01 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Apr 24 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003979 | OEIS-A117339

a(n) = a(n-1) + a(n-2); if a(n) is not prime divide a(n) by its largest prime factor.

Sequence repeats cyclically. Conjecture: sequence ends with cycle for any positive a(1) and a(2). This has been checked for a(1)=1 and a(2)=1,...,2000. Some other cases shown below: a(1), a(2), length of s, sequence s 1,1,10,{1,1,2,3,5,4,3,7,2,3}, 1,2,9,{1,2,3,5,4,3,7,2,3}, 1,3,9,{1,3,2,5,7,4,11,3,2}, 1,4,20,{1,4,5,3,4,7,11,6,17,23,8,31,3,2,5,7,4,11,3,2}, 1,5,12,{1,5,2,7,3,2,5,7,4,11,3,2}, 1,6,23,{1,6,7,13,4,17,3,4,7,11,6,17,23,8,31,3,2,5,7,4,11,3,2}, 1,7,9,{1,7,4,11,3,2,5,7,4}, 1,8,30,{1,8,3,11,2,13,3,8,11,19,6,5,11,8,19,9,4,13,17,6,23,29,4,3,7,2,3,5,4,3}, 1,9,34,{1,9,2,11,13,8,3,11,2,13,3,8,11,19,6,5,11,8,19,9,4,13,17,6,23,29,4,3,7,2,3,5,4,3}, 1,10,10,{1,10,11,3,2,5,7,4,11,3}. Even exotic seeds give no long sequences, e.g. 967,3000,24,{967,3000,3967,6967,154,7121,75,28,103,131,18,149,167,4,9,13,2,3,5,4,3,7,2,3}. For a(1)=1, maximal length = 55 is found for a(2)=1207: 1,1207,55,{1,1207,8,243,251,26,277,3,40,43,83,18,101,7,36,43,79,2,27,29,8,37,9,2,11,13,8,3,11,2,13,3,8,11,19,6,5,11,8,19,9,4,13,17,6,23,29,4,3,7,2,3,5,4,3}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Mar 09 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003980 | OEIS-A117351

Where n appears in A082224, provided A082224 is read as a sequence rather than as an array.

Conjecture: Sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Mar 09 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003981 | OEIS-A117436

Triangle related to exp(x)*sec(2*x).

Conjecture: The d-th diagonal (starting with d=0) is proportional to the sequence of generalized binomial coefficients binomial(-x, d) where x is the column index. - _Søren G. Have_, Feb 26 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 16 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003982 | OEIS-A117547

Numbers k such that 2*k^2+1 is a perfect power.

Conjectures from _Colin Barker_, Apr 28 2012: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 29 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003983 | OEIS-A117549

Values of n for which there exist d(1),...,d(n), each in {0,1,...,4} and an r in {1,...,4} such that Sum[d(i)d(i+k),i=1,n-k]=r (mod 5) for all k=0,...,n-1. (Such a sequence is called a very(5,r) sequence. See the link.).

Conjecture. Let A be a very(5,1) (respectively very(5,4)) sequence of length n and let Z be a sequence of n-1 0's.. Then AZ(3A)ZA is a very(5,1) (respectively very(5,4)) sequence of length 5n-2. (Here 3A denotes the result of multiplying each term of A by 3, then reducing modulo 5; and juxtaposition of symbols denotes concatenation of sequences.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Apr 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003984 | OEIS-A117633

Number of self-avoiding walks of n steps on a Manhattan square lattice.

It is proved that a(n)^(1/n) has a limit mu called the "connective constant" of the Manhattan lattice; approximate value of mu: 1.733735. It is only conjectured that a(n + 1) ~ mu * a(n). - _Robert FERREOL_, Dec 08 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Apr 08 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003985 | OEIS-A117729

Orders k of cyclic groups C_k such that the map "G -> Automorphism group of G" eventually reaches the trivial group when started at C_k.

Conjecture: If A003434(k) = A185816(k), then k is a term of this sequence. The conjecture is verified for k <= 10^8. (End) The conjecture is verified for k <= 10^9. - _Stan Wagon_, Dec 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, based on a communication from _J. H. Conway_, Apr 14 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003986 | OEIS-A117769

Lucas numbers for which the product of the digits is a Fibonacci number.

A000204 INTERSECT A011540 is a subsequence. As a consequence of Carmichael's theorem, the product of the digits of terms in the sequence must be in the set {0, 1, 2, 3, 5, 8, 21, 144} and if a term is zeroless (A052382), then at most 6 digits are not equal to 1. Conjecture: all terms > 18 have a 0 digit, i.e. is a member of A011540. - _Chai Wah Wu_, Mar 12 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | Luc Stevens (lms022(AT)yahoo.com), Apr 15 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003987 | OEIS-A117770

Fibonacci numbers for which the product of the digits is also a Fibonacci number.

A000045 INTERSECT A011540 is a subsequence. As a consequence of Carmichael's theorem, the product of the digits of terms in the sequence must be in the set {0, 1, 2, 3, 5, 8, 21, 144} and if a term is zeroless (A052382), then at most 6 digits are not equal to 1. Conjecture: all terms > 21 have a 0 digit, i.e. is a member of A011540. - _Chai Wah Wu_, Mar 12 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | Luc Stevens (lms022(AT)yahoo.com), Apr 15 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003988 | OEIS-A117831

Let S_n be the infinite sequence formed by starting with n and repeatedly reversing the digits and adding 4 to get the next term. Sequence gives number of steps for S_n to reach a cycle, or -1 if no cycle is ever reached.

It is conjectured that S_n always reaches a cycle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, following discussions with Luc Stevens, May 03 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003989 | OEIS-A117846

Given n, define a(n) as follows: let a sequence b(k) be defined by b(k+1)=b(k)+b(k)mod k; b(1)=2n-1. (Here b(k)mod k denotes the least nonnegative residue of b(k) modulo k). Let a(n) be the common value of b(k+1)-b(k) for all large k if such exists; otherwise let a(n) be 0.

Putting b(1)=2n gives essentially the same sequence as putting b(1)=2n-1. It is a plausible conjecture or at least an interesting open problem that a(n) is never zero; that is all the sequences b(k) are arithmetic progressions from some point on. Sequence A073117 is the sequence b(k) with b(1)=1. Do the values a(n) include all positive numbers?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Abercrombie_, Mar 22 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003990 | OEIS-A117896

Number of perfect powers between consecutive squares n^2 and (n+1)^2.

Trivially, a(n) << log n/log log n. Turk gives a(n) << sqrt(log n) and Loxton improves this to a(n) <= exp(40 sqrt(log log n log log log n)). Stewart improves the constant from 40 to 30 and conjectures that a(n) < 3 for all but finitely many n. - _Charles R Greathouse IV_, Dec 11 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 31 2006, Feb 15 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003991 | OEIS-A117922

Inverse of A055265, assuming it is a permutation, as conjectured.

Inverse of A055265, assuming it is a permutation, as conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 03 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003992 | OEIS-A117964

a(n) = A117963(n) mod 2.

a(2*A081601(n)) = a(1+2*A081601(n)) = 1. [Conjectured, also these two formulas together seem to give the positions of all 1's] - _Antti Karttunen_, Jan 01 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 05 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003993 | OEIS-A117969

Start of least run of maximal length of consecutive n-almost primes.

For n>=2 there cannot be more than 2^n - 1 consecutive n-almost primes. Is it known whether there always exists such a run of length 2^n - 1? If not, I conjecture so. This is confirmed to be true for terms through a(4). Terms here equal the last terms of corresponding finite sequences: a(3) = A067813(6). a(4) was computed by _Don Reble_ as A067814(14). a(5) >= A067820(12).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rick L. Shepherd_, Apr 05 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003994 | OEIS-A117971

One-based position of the first 2 from the least significant digit in the ternary expansion of 2^n, or 0 if there are no 2's present.

a(0), a(2) and a(8) are the conjectured to be the only terms equal to 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Apr 06 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003995 | OEIS-A118006

Define a sequence of binary words by w(1) = 01 and w(n+1) = w(n)w(n)Reverse[w(n)]. Sequence gives the limiting word w(infinity).

The article by W. Hebisch and M. Rubey gives a conjectured functional equation for the g.f. for this sequence. - _N. J. A. Sloane_, Sep 08 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Apr 09 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003996 | OEIS-A118076

Numbers n such that n divides sigma_(2^k)(n), the sum of the 2^k powers of the divisors of n, for all k>0.

Although these numbers have been tested up to k=20, it is conjectured that n divides sigma_(2^k)(n) for all k>0. Intersection of A046762 and A066292.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 11 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003997 | OEIS-A118093

Numbers of rooted hypermaps on the torus with n darts (darts are semi-edges in the particular case of ordinary maps).

Conjecture: +n*(5*n-17)*a(n) -15*(n-1)*(5*n-16)*a(n-1) +12*(20*n^2-103*n+140)*a(n-2) +32*(5*n-12)*(2*n-5)*a(n-3)=0. - _R. J. Mathar_, Apr 05 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Valery A. Liskovets_, Apr 13 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003998 | OEIS-A118109

Binary representation of n-th iteration of the Rule 54 elementary cellular automaton starting with a single black cell.

Conjectures from _Colin Barker_, Dec 08 2015 and Apr 16 2019: (Start) Conjecture: a(n) = floor((10000+1100*(n mod 2))*100^n/9999). - _Karl V. Keller, Jr._, Sep 24 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Apr 13 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-003999 | OEIS-A118122

Least prime of level 2n-1 (cf. A117563).

Levels of primes are defined in A117563. Conjecture: there are an infinite number of prime members at each level.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémi Eismann_ and _Robert G. Wilson v_, May 12 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004000 | OEIS-A118201

Smallest difference such that both difference and number do not occur previously.

Similar to Recamán's sequence (A005132), but increases the difference to avoid duplicating values. Conjecture that every nonnegative integer occurs in this sequence. Evaluating through n=20000, the smallest number that has not occurred is 139.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Apr 14 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004001 | OEIS-A118202

Inverse of A118201, or -1 if n does not occur in A118201.

Conjecture that a(n) is never -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Apr 14 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004002 | OEIS-A118278

Conjectured largest number that is not the sum of three n-gonal numbers, or -1 if there is no largest number.

Conjectured largest number that is not the sum of three n-gonal numbers, or -1 if there is no largest number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004003 | OEIS-A118279

Conjectured number of numbers that are not the sum of three n-gonal numbers, or -1 if infinite.

Conjectured number of numbers that are not the sum of three n-gonal numbers, or -1 if infinite. See A118278 for the conjectured largest number that is not the sum of three n-gonal numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004004 | OEIS-A118280

Conjectured largest number that is not the sum of three (2n+1)-gonal numbers; bisection of A118278.

Conjectured largest number that is not the sum of three (2n+1)-gonal numbers; bisection of A118278.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004005 | OEIS-A118281

Conjectured number of numbers that are not the sum of three (2n+1)-gonal numbers; bisection of A118279.

Conjectured number of numbers that are not the sum of three (2n+1)-gonal numbers; bisection of A118279.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004006 | OEIS-A118282

Conjectured largest number that is not the sum of three generalized n-gonal numbers, or -1 if there is no largest number.

Conjectured largest number that is not the sum of three generalized n-gonal numbers, or -1 if there is no largest number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004007 | OEIS-A118283

Conjectured number of numbers that are not the sum of three generalized n-gonal numbers, or -1 if infinite.

Conjectured number of numbers that are not the sum of three generalized n-gonal numbers, or -1 if infinite. See A118282 for the conjectured largest number that is not the sum of three generalized n-gonal numbers. For n=7, only four numbers are not representable: 10, 16, 76 and 307.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004008 | OEIS-A118284

Conjectured largest number that is not the sum of three generalized (2n+1)-gonal numbers; bisection of A118282.

Conjectured largest number that is not the sum of three generalized (2n+1)-gonal numbers; bisection of A118282.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 21 2006; revised Apr 23 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004009 | OEIS-A118285

Conjectured number of numbers that are not the sum of three generalized (2n+1)-gonal numbers; bisection of A118283.

Conjectured number of numbers that are not the sum of three generalized (2n+1)-gonal numbers; bisection of A118283.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004010 | OEIS-A118286

Numbers n such that n == 0 (mod 4) or n == 2 (mod 12).

Except for n=2, conjectured n such that A118278(n)=-1 and A118282(n)=-1; n such that there is an infinite set of numbers not representable as the sum of three n-gonal numbers or three generalized n-gonal numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 23 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004011 | OEIS-A118301

Number of partitions of n into distinct parts with largest part congruent to n modulo 2.

Conjectural g.f.: A(x) = Limit_{N -> oo} (Sum_{n = 0..2*N+1} (-1)^(n+1)/Product_{k = 1..n} 1 - x^(2*k-1)). - _Peter Bala_, Feb 11 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 22 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004012 | OEIS-A118302

Number of partitions of n into distinct parts with largest part not congruent to n modulo 2.

Conjectural g.f.: A(x) = Limit_{N -> oo} (Sum_{n = 1..2*N} (-1)^n/Product_{k = 1..n} 1 - x^(2*k-1)). - _Peter Bala_, Feb 11 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 22 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004013 | OEIS-A118313

Sum of squared end-to-end distances of all n-step self-avoiding walks on the simple cubic lattice.

a(5) is 25556 according to MacDonald et al., but 25566 according to Clisby et al. and is therefore conjectural for now. - _R. J. Mathar_, Aug 31 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, May 14 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004014 | OEIS-A118368

Largest number that is not the sum of four (2n+1)-gonal numbers.

See A118367 for the largest number that is not the sum of five n-gonal numbers. Sequence A118280 has the conjectured bounds for three (2n+1)-gonal numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 25 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004015 | OEIS-A118376

Number of all trees of weight n, where nodes have positive integer weights and the sum of the weights of the children of a node is equal to the weight of the node.

Conjectured to be the number of permutations of length n avoiding the partially ordered pattern (POP) {1>2, 1>3, 3>4, 3>5} of length 5. That is, conjectured to be the number of length n permutations having no subsequences of length 5 in which the first element is the largest, and the third element is larger than the fourth and fifth elements. - _Sergey Kitaev_, Dec 13 2020 This conjecture has been proven. It can be restated as the number of size n permutations avoiding 51423, 51432, 52413, 52431, 53412, 53421, 54312, 54321. There are twelve sets of permutations avoiding eight size five permutations that are known to match this sequence. A further four are conjectured to match this sequence. - _Christian Bean_, Jul 24 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jeremy Johnson (jjohnson(AT)cs.drexel.edu), May 15 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004016 | OEIS-A118447

Number of rooted n-edge one-vertex maps on the Klein bottle (dually: one-face maps).

Conjecture: -(n-2)*(n-1)^2*a(n) +2*n*(4*n-5)*(n-2)*a(n-1) -8*n*(n-1)*(2*n-3)*a(n-2)=0. - _R. J. Mathar_, Jun 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Valery A. Liskovets_, May 04 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004017 | OEIS-A118510

Define sequence S_m by: initial term = m, reverse digits and add 1 to get next term. It is conjectured that S_m always reaches a cycle of length 9, as in A117230. Sequence gives records for number of steps to reach cycle.

Define sequence S_m by: initial term = m, reverse digits and add 1 to get next term. It is conjectured that S_m always reaches a cycle of length 9, as in A117230. Sequence gives records for number of steps to reach cycle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 06 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004018 | OEIS-A118511

Define sequence S_n by: initial term = n, reverse digits and add 1 to get next term. It is conjectured that S_n always reaches a cycle of length 9, as in A117230. Sequence gives number of steps for S_n to reach the cycle.

Define sequence S_n by: initial term = n, reverse digits and add 1 to get next term. It is conjectured that S_n always reaches a cycle of length 9, as in A117230. Sequence gives number of steps for S_n to reach the cycle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 06 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004019 | OEIS-A118514

Define sequence S_n by: initial term = n, reverse digits and add 2 to get next term. It is conjectured that S_n always reaches a cycle. Sequence gives number of steps for S_n to reach a cycle.

Define sequence S_n by: initial term = n, reverse digits and add 2 to get next term. It is conjectured that S_n always reaches a cycle. Sequence gives number of steps for S_n to reach a cycle. There is one cycle of length 81 (least component is 3, all components have at most three digits, cf. A117521), 22 cycles of length 90 with 4-digit components (least components are 1013 + 2*k for k = 0, ..., 21, cf. A120214) and one cycle of length 45 with 4-digit components (least component is 1057, cf. A120215). Furthermore there are 22 cycles of length 1890 (least components are 100013 + 2*k for k = 0, ..., 21, cf. A120216), one cycle of length 945 (least component is 100057, cf. A120217) and 225 cycles of length 900 (least components are 100103 + 2*k for k = 0, ..., 224, cf. A120218), all having 6-digit components. It is conjectured that there are also cycles of increasing length with 8-, 10-, 12-, ... digit components. - _Klaus Brockhaus_, Jun 10 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 06 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004020 | OEIS-A118522

Define sequence S_n by: initial term = n, reverse digits and add 3 to get next term. It is conjectured that S_n always reaches a cycle. Sequence gives number of steps for S_n to reach a cycle.

Define sequence S_n by: initial term = n, reverse digits and add 3 to get next term. It is conjectured that S_n always reaches a cycle. Sequence gives number of steps for S_n to reach a cycle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 06 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004021 | OEIS-A118650

Center column a(2n,n) of A106597.

Conjecture: n*a(n) +3*(-3*n+2)*a(n-1) +4*(5*n-8)*a(n-2) +2*(-8*n+21)*a(n-3) +2*(2*n-7)*a(n-4)=0. - _R. J. Mathar_, Nov 10 2013 This proves the above conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joshua Zucker_, May 10 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004022 | OEIS-A118739

Let L_n be the infinite sequence formed by starting with 1 and repeatedly placing the first digit at the end of the number and adding n to get the next term. If L_n eventually reaches a cycle, sequence gives length of that cycle, otherwise -1.

It is conjectured that L_n always reaches a cycle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Luc Stevens (lms022(AT)yahoo.com), May 24 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004023 | OEIS-A118740

Let L_n be the infinite sequence formed by starting with 1 and repeatedly placing the first digit at the end of the number and adding n to get the next term. Sequence gives number of steps for L_n to enter a cycle, or -1 if no cycle is ever reached.

It is conjectured that L_n always reaches a cycle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Luc Stevens (lms022(AT)yahoo.com), May 24 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004024 | OEIS-A118767

Fixed points of permutations A118763, A118764, A118765 and A118766.

No more terms less than 10^4. Looking at the pattern mod 9, I conjecture that there will be more terms past 10^8. - _Joshua Zucker_, May 14 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, May 01 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004025 | OEIS-A118955

Numbers of the form 2^k + prime.

The lower density is at least 0.09368 (Pintz) and upper density is at most 0.49095 (Habsieger & Roblot). The density, if it exists, is called Romanov's constant. Romani conjectures that it is around 0.434. - _Charles R Greathouse IV_, Mar 12 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, May 07 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004026 | OEIS-A118974

Sum of the lengths of the first descents in all hill-free Dyck paths of semilength n (a hill in a Dyck path is a peak at level 1).

Conjecture: 2*(n+1)*(17*n^2-65*n+60)*a(n) -3*(3*n-4)*(17*n^2-48*n+15)*a(n-1) +3*(17*n^3-82*n^2+121*n-60)*a(n-2) +2*(2*n-5) *(17*n^2-31*n+12) *a(n-3)=0. - _R. J. Mathar_, Jun 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, May 08 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004027 | OEIS-A119012

Number of valleys strictly above the x-axis in all hill-free Dyck paths of semilength n (a hill in a Dyck path is a peak at level 1).

Conjecture: 2*(n-3)*(3*n-2)*(n+2)*a(n) +(-21*n^3+50*n^2-13*n-4)*a(n-1) -2*(n-1) *(2*n-1)*(3*n+1)*a(n-2)=0. - _R. J. Mathar_, Jun 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, May 08 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004028 | OEIS-A119018

a(1) = 1; a(n) is the smallest number not yet used that is not relatively prime to n nor relatively prime to a(n-1).

Conjecture: this sequence includes every integer greater than 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, May 08 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004029 | OEIS-A119259

Central terms of the triangle in A119258.

More generally, we conjecture that a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, May 11 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004030 | OEIS-A119262

Number of B-trees of order infinity with n leaves, where a(n) = Sum_{k=1..floor(n/2)} a(k)*C(n-k-1,n-2*k) for n >= 2, with a(0)=0, a(1)=1.

Conjecture: for n>=1, a(n) is the number of lower sets of size n-1 in the Young-Fibonacci lattice (tested up to n = 28). See A393296 for the number of lower sets in the Young lattice. - _Ludovic Schwob_, Feb 11 2026 Conjecture: Let M_n be an n X n matrix whose elements are m_ij = 0 for i < j - 1, m_ij = -1 for i = j - 1, and m_ij = binomial(i - j, n - i) otherwise. Then a(n + 1) = Det(M_n). - _Benedict W. J. Irwin_, Apr 19 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 11 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004031 | OEIS-A119350

First differences of A118374.

Conjecture. Define g(1)=1 and, for i>1, g(i)=2*g(i-1)+2^(i-2). Also define h(1)=1, h(2)=4 and, for i>2, h(i)=h(i-1)+2^(i-2). Then a(n)=h(i) if n=g(i), a(n)=1 if g(i)-2^(i-2)<=n<g(i) and a(n)=a(n-g(i-1)) if g(i-1)<n<g(i)-2^(i-2). (This has been verified for the first 1000 terms.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, May 23 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004032 | OEIS-A119356

Numbers k such that A000330(k) is squarefree.

The numbers of terms that do not exceed 10^k, for k = 1, 2, ..., are 8, 53, 504, 5029, 50187, 501925, 5019527, 50194688, 501948054, 5019478733, ... Conjecture: The asymptotic density of this sequence is 4 * Product_{p prime} (1 - 3/p^2) = 4 * A206256 = 0.50194792... . - _Amiram Eldar_, Sep 24 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, May 16 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004033 | OEIS-A119366

Number of rooted planar n-trees whose number of leaves is equal to 1 modulo 3.

Conjecture D-finite with recurrence +n*(881*n-4580)*(n-2)*(n+1)*a(n) -3*n*(612*n^3-2827*n^2-2988*n+10135)*a(n-1) +3*(-3088*n^4+42803*n^3-190361*n^2+313702*n-167988)*a(n-2) +(43042*n^4-600920*n^3+2924411*n^2-5860777*n+4115562)*a(n-3) +3*(-38600*n^4+558681*n^3-2904370*n^2+6389913*n-4965528)*a(n-4) +3*(-14776*n^4+162695*n^3-434711*n^2-415064*n+1878084)*a(n-5) -9*(n-6)*(10835*n^3-106831*n^2+290611*n-173519)*a(n-6) +54*(n-6)*(n-7)*(593*n-1429)*(2*n-13)*a(n-7)=0. - _R. J. Mathar_, Feb 03 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 16 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004034 | OEIS-A119370

G.f. satisfies A(x) = 1 + x*A(x)^2 + x^2*(A(x)^2 - A(x)).

Conjecture: A(x) = 1 + x*exp(Sum_{n >= 1} g(n, x)*x^n/n), where g(n, x) = Sum_{k = 0..n} binomial(n, k)^2*(1 + x)^k. Cf. A105633 and A167638. - _Peter Bala_, Sep 10 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 16 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004035 | OEIS-A119414

Number of triangle-free graphs g on n nodes for which the chromatic number chi(g) equals r(g) = ceiling((Delta(g) + 1 + omega(g))/2).

Here Delta(g) is the maximum node degree of g and omega(g) is the clique number of g (=2 for triangle-free graphs). r(g) is conjectured by Reed to be an upper bound for chi(g) for all graphs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Keith Briggs_, Jul 26 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004036 | OEIS-A119449

Primes with even digit sum.

On average, there are as many prime numbers for which the sum of decimal digits is even as prime numbers for which it is odd [A119450]. This hypothesis, first made in 1968, has recently been proved by researchers from the Institut de Mathematiques de Luminy. - _Jonathan Vos Post_, May 13 2010 Also primes such that absolute value of difference between largest digit and the sum of all the other digits is an even integer. This is in accordance with hypothesis of Alexandre Gelfond, proved by C. Mauduit and J. Rivat, see the link below. - _Osama Abuajamieh_, Feb 10 2017.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, May 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004037 | OEIS-A119450

Primes with odd digit sum.

On average, there are as many prime numbers for which the sum of decimal digits is even as prime numbers for which it is odd [A119450]. This hypothesis, first made in 1968, has

recently been proved by researchers from the Institut de Mathematiques de Luminy. Also primes such that absolute value of difference between largest digit and the sum of all the other digits is an odd integer. This is in accordance with hypothesis of Alexandre Gelfond, proved by C. Mauduit and J. Rivat as stated in Links section. - _Osama Abuajamieh_, Feb 10 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, May 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004038 | OEIS-A119451

Let S_n be the infinite sequence formed by starting with n and repeatedly reversing the digits and adding 16 to get the next term. Sequence gives number of steps for S_n to reach a cycle, or -1 if no cycle is ever reached.

It is conjectured that S_n always reaches a cycle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, May 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004039 | OEIS-A119479

Length of longest run of consecutive integers having exactly n divisors.

Schinzel's conjecture H would imply that:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Jul 26 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004040 | OEIS-A119485

Number of children for which any subset can be generated by a counting-out game.

Conjecture (by J. Fricke and G. Woeginger): The sequence contains exactly: powers of 2, primes and doubled primes, A058363 with a 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jan Fricke_, May 23 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004041 | OEIS-A119486

Numbers of children for which there is a subset which cannot be generated by a counting-out game.

Conjecture (by J. Fricke and G. Woeginger): The sequence contains all numbers n with an odd prime divisor p satisfying n/p>2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jan Fricke_, May 23 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004042 | OEIS-A119498

Consider the sign of A067276: determinant of n X n matrix containing the first n^2 primes in increasing order; then a(n) = 0 if negative and 1 if positive.

Conjecture: This sequence never cycles.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, May 26 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004043 | OEIS-A119505

The Pi-th digit of Pi where the digit value of 0 is interpreted as decimal 10.

The numbers formed in this sequence are 1,2,3,4,5,6,9. Conjecture: The terms of this sequence are nonrepeating and nonterminating.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, May 27 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004044 | OEIS-A119506

The e-th digit of e where the digit value of 0 is interpreted as decimal 10.

Conjecture: this sequence is non-repeating and infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, May 27 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004045 | OEIS-A119537

Determinant of n X n matrices of first n^2 denumerants (A000115).

Conjecture: a(n>7)=0. - _Robert G. Wilson v_, Jun 07 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, May 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004046 | OEIS-A119574

a(n) = binomial(2*n,n)*(n+2)^2/(n+1).

Conjectured g.f.: (-1 + 14*x - 36*x^2 + (1 - 4*x)^(3/2))/(2*x*(1 - 4*x)^(3/2)). - _Harvey P. Dale_, Jun 02 2024. The conjecture is true (see links). - _Sela Fried_, Oct 02 2024.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zerinvary Lajos_, May 31 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004047 | OEIS-A119623

Composite numbers for which the second elementary symmetric function of divisors (s2) is prime.

Terms in A119616 are always prime if n is prime p and s2(p)=p, hence it is interesting to find composite numbers for which s2 is also prime. Relative values of s2 are: s2=47,97,163,457,733,2203,3733,7993,10723,11317,21313,22147,26557,33403,57283,61417,67153,79393,101467,149323,160453,162727,174337,272683,296827,318793,358273,432907,440383,486583,551767,639007,832687,843043,911917,961183,1152913,1202017,1277593,1322743,1375303,1462897,1567327,1824997,1878883. Otherwise the sequence s2 gives numbers which appear in A119616 at least twice (and conjecture is that exactly twice).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Jun 08 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004048 | OEIS-A119651

Number of different values of exactly n standard American coins (pennies, nickels, dimes and quarters).

Conjectures from _Colin Barker_, Oct 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tanya Khovanova_, Jul 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004049 | OEIS-A119652

Number of different values of <= n standard American coins (pennies, nickels, dimes and quarters).

Conjectures from _Colin Barker_, Oct 25 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tanya Khovanova_, Jul 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004050 | OEIS-A119659

Floor of the area of consecutive Prime-Indexed Prime triangles.

Conjecture: The triples (3,5,11), (5,11,17), (11,17,31) are the only consecutive PIP triples that cannot form a triangle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jul 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004051 | OEIS-A119690

a(n) = n! mod n*(n+1)/2.

Layman's conjecture above is true for every k >= 0. Proof: if n+1 is not an odd prime, then n(n+1)/2 divides n! (see A069268), so every positive power of n! is 0 modulo n(n+1)/2. If n+1 = p is an odd prime, write n(n+1)/2 = h*p with h = n/2, gcd(h,p) = 1. Then n! == 0 (mod h), and n! == -1 (mod p) by Wilson's theorem, so for odd j, (n!)^j == 0 (mod h) and == -1 (mod p). Since n = 2h == 0 (mod h) and n = p-1 == -1 (mod p), the Chinese remainder theorem gives (n!)^j == n (mod n(n+1)/2), and 0 <= n < n(n+1)/2. - _Scott M. Foltz_, Sep 24 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_ and _Giorgio Balzarotti_, Jun 09 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004052 | OEIS-A119936

Least common multiple (LCM) of denominators of the rows of the triangle of rationals A119935/A119932.

a(n)/n^3 = A027451(n) = A002944(n)^2 (the second equation is a conjecture). a(n)/n^3 = (A099946(n)*(n-1))^2, n >= 2 (from the conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jul 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004053 | OEIS-A119967

A transform of the central binomial coefficients C(n,floor(n/2)).

Conjecture: (n+2)*(n-3)*a(n) +(2+2*n-n^2)*a(n-1) +(3*n-5)*(n-2)*a(n-2) -(n-2)*(n-3)*a(n-3) -(n-2)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Nov 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 31 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004054 | OEIS-A119975

E.g.f. exp(x)*(Bessel_I(0,2*sqrt(2)x) + Bessel_I(1,2*sqrt(2)x)/sqrt(2)).

Conjecture: (n+1)*a(n) +(n-4)*a(n-1) +(9-13*n)*a(n-2) +21*(2-n)*a(n-3) =0. - _R. J. Mathar_, Dec 10 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jun 02 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004055 | OEIS-A120062

Number of triangles with integer sides a <= b <= c having integer inradius n.

It is conjectured that the longest possible side c of a triangle with integer sides and inradius n is given by A057721(n) = n^4 + 3*n^2 + 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 11 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004056 | OEIS-A120069

Denominators of partial sums of a convergent series involving scaled Catalan numbers A000108.

Conjecture: a(n) = denominator(Catalan(n)/4^(n-1)). - _Groux Roland_, Dec 11 2010 Conjecture: a(n) = 2^(A283208(n)-2). - _Alexander Staunton_, Feb 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jul 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004057 | OEIS-A120078

Coefficient triangle of numerator polynomials appearing in certain column o.g.f.s related to the H-atom spectrum.

See the W. Lang link under A120072 for the precise form of the o.g.f.s: G(x,n) = -dilog(1-x) + x*P(n,4)/*(A(n)*(n^2)*(1-x)), with A(n) = [1, 1, 4, 9, 144, 100, 3600, 11025, 78400, 63504, ...] = conjectured to be A027451(n), n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jul 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004058 | OEIS-A120114

a(n) = lcm(1, ..., 2n+4)/lcm(1, ..., 2n+2).

Conjectures from _Davide Rotondo_, Dec 02 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jun 09 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004059 | OEIS-A120121

Numbers m such that m=phi((d_1*d_2*...*d_k)*(d_1+d_2+...+d_k)) where d_1 d_2... d_k is the decimal expansion of m.

Conjecture: 139968 is the largest term. Except for the first term all terms are even. It's interesting that for the number 139968 we have the following relations: 139968=phi((1*3*9*9*6*8)*(1+3+9+9+6+8))=phi(1*3*9*9*6*8) *(1+3+9+9+6+8)=(1*3*9*9*6*8)*phi(1+3+9+9+6+8).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Aug 12 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004060 | OEIS-A120123

Numbers m such that m=(d_1*d_2*...*d_k)*phi(d_1+d_2+...+d_k) where d_1 d_2... d_k is the decimal expansion of m.

Conjecture: 139968 is the largest term. Except for the first term, all terms are even. 139968 is also in A120121 & A120122, so it has a very pleasant property (see the Comments lines of A120121).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Aug 12 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004061 | OEIS-A120160

a(n) = ceiling(Sum_{i=1..n-1} a(i)/4) for n >= 2 starting with a(1) = 1.

Conjecture 1: a(n) equals the number of multiples of 4 whose representation in base 5/4 (see A024634) has n-1 digits. For example, a(8) = 3 because there are three multiples of 4 with n-1 = 7 digits in their representation in base 5/4: 36 = 4321031, 40 = 4321420, and 44 = 4321424. Conjecture 2: a(n) equals 1/5 times the number of nonnegative integers with the property that their 5/4-expansion has n digits (assuming that the 5/4-expansion of 0 has 1 digit). For example, a(7)*5 = 10 because the following 10 numbers have 5/4 expansions with n = 7 digits: 35 = 4321030, 36 = 4321031, 37 = 4321032, 38 = 4321033, 39 = 4321034, 40 = 4321420, 41 = 4321421, 42 = 4321422, 43 = 4321423, and 44 = 4321424. (End) Starting at a(11) = 5, we conjecture that this sequence gives all those numbers m for which when we place m persons on a circle, label them 1 through m, start counting at person number 1, and remove every 5th person, the last survivors have numbers in {1, 2, 3, 4}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Graeme McRae_, Jun 10 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004062 | OEIS-A120275

Smallest prime factor of the odd Catalan number A038003(n).

a(n) <> 3 iff the base-3 representation of 2^n-1 has no 2's. Conjecture: this only occurs for n = 2, 5, 8. I verified it up to n = 10^4. - _Robert Israel_, Nov 18 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Jul 04 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004063 | OEIS-A120292

Absolute value of numerator of determinant of n X n matrix with elements M[i,j] = prime(i)/(1+prime(i)) if i=j and 1 otherwise.

Conjecture: All composite terms are semiprime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Jul 08 2006, Jul 04 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004064 | OEIS-A120299

Largest prime factor of Stirling numbers of first kind s(n,2) = A000254(n).

a(n) = gpf(A096617(n)), where gpf = A006530 is the greatest prime factor, and A096617 is a "reduced" variant of A001008 and thus A000254. [Conjectured; true if this gpf is always > n.] - _M. F. Hasler_, Jul 04 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Jul 11 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004065 | OEIS-A120334

Odd primes of the form p = 2^k +- 1 or p = 4^k +- 3 or such that 2^p - 1 is prime or (2^p + 1)/3 is prime or PRP.

Odd primes satisfying at least one of the criteria considered in the New Mersenne Prime Conjecture (cf. link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jorge Coveiro_, Sep 11 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004066 | OEIS-A120337

Euler-irregular primes p dividing E(2k) for some 2k < p-1.

Conjecture (Ernvall and Metsänkylä, 1978): The asymptotic density of this sequence within the primes is 1 - 1/sqrt(e) = 0.393469... (A290506), the same as the corresponding conjectured density of the irregular primes (A000928). - _Amiram Eldar_, Dec 06 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefan Krämer_, Jun 22 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004067 | OEIS-A120355

a(n) = min{k>0: the n-th convergent to e equals m/k! for some m}.

Conjecture: for n>5, a(n) = A006530(A007677(n)). - _Chai Wah Wu_, Feb 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Aug 16 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004068 | OEIS-A120362

Numerators of bivariate Taylor expansion of the incomplete elliptic integral of the second kind.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jun 26 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004069 | OEIS-A120397

Minimal number of steps needed to represent a prime >=5 as a sum of at most 3 primes such that all the previous odd primes are represented.

This sequence contains probably every natural number, though this should be checked further. Conjecture: Every prime >=5 can be represented as a sum of at most 3 primes, such that all the previous odd primes are represented.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robin Garcia_, Jul 01 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004070 | OEIS-A120427

For each y >= 1 there are only finitely many values of x >= 1 such that x-y and x+y are both squares; list all such pairs (x,y) with gcd(x,y) = 1 ordered by values of y; sequence gives y values.

Conjecture: lim_{n->oo} a(n)/n = 1/Pi. - _Lothar Selle_, Jun 19 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 02 2001 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004071 | OEIS-A120437

Differences of A037314 (sum of base-3 digits of n = sum of base-9 digits of n).

a(n) = A066443(A007949(n)). (This is equivalent to the conjectured recursion above; that recursion is correct.) - _Franklin T. Adams-Watters_, Jul 24 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jul 17 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004072 | OEIS-A120451

Number of ways a prime number p can be expressed as 2*(p1-p2) + 3*p3, where p1, p2, p3 are primes or 1, less than or equal to p.

At least for the first 200 primes, it is true that every prime p > 2 can be expressed as 2*(p1-p2) + 3*p3, where p1, p2, p3 are primes or 1, less than or equal to p (the proof would be straightforward if both (a) Levy's conjecture and (b) the conjecture saying that every prime p > 3 can be expressed as 2*p1 + 3*p2, where p1, p2 are primes, were true). It would be interesting to study how the sequence changes when we remove the restriction for p1, p2, p3 to be less than or equal to p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vassilis Papadimitriou_, Jul 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004073 | OEIS-A120503

Generalized meta-Fibonacci sequence a(n) with parameters s=0 and k=3.

a(n) = A007844(n)/3. - _Michel Marcus_, Aug 19 2013, conjectured. This is true: see the analogous sequence A007843 for a sketch of the proof. - _M. F. Hasler_, Dec 27 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank Ruskey_ and Chris Deugau (deugaucj(AT)uvic.ca), Jun 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004074 | OEIS-A120561

Numbers n such that Lucas(prime(n)) is prime, where Lucas = A000032.

All prime Lucas numbers A000032[n] have indices that are prime, zero or a power of 2. It is a conjecture that all indices of prime Lucas numbers are prime, except n = 0, 4, 8, 16.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Aug 07 2006, Oct 05 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004075 | OEIS-A120629

Numbers k with property that -k is not a perfect power and the squarefree part of -k is not congruent to 1 modulo 4.

According to a famous 1927 conjecture of Emil Artin, modified by Dick Lehmer, these negative numbers are primitive roots modulo each prime of a set whose density among primes equals Artin's constant (see A005596). The positive numbers with the same property are given by A085397.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerard P. Michon_, Jun 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004076 | OEIS-A120643

Table T(n,k) = number of fractal initial sequences (where new values are successive integers) of length n whose last term is k.

A fractal sequence is one where, when the first instance of each integer is removed, the original sequence results. We require also that these first instances occur in order: 1,1,2,3 is OK, but 1,1,3,2 is not. A finite sequence is an initial subsequence of (uncountably many) fractal sequences when the result after removing the first instance of each number is an initial subsequence. The total number of such sequences of length n is 2^{n-1}. At each index after the first, the next value can be either a new value or a uniquely determined repetition of some earlier value. Conjecture: column 1 of this array is A007059.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Aug 17 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004077 | OEIS-A120737

Numbers k whose number of divisors d(k) is divisible by every prime factor of k.

Numbers k such that A000005(k)/A007947(k) is an integer. A070226 is a subsequence of this sequence. Conjecture: If A000005(k) divides A007947(k) for some k, then A007947(k)/A000005(k)=1. - _Ctibor O. Zizka_, Feb 05 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Jun 29 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004078 | OEIS-A120740

Numbers n such that n = Sum_digits[k*abs(n-k)] for some k>=0.

Conjecture: a(n) = (18*n-(-1)^n-35)/4 for n>2. a(n) = a(n-1)+a(n-2)-a(n-3) for n>5. G.f.: x^2*(4+x+4*x^3)/((1-x)^2*(1+x)). [_Colin Barker_, Apr 10 2012] (The comment above on the 2-periodicity of the first difference and comparison with A130877 mean this is correct. - _R. J. Mathar_, Mar 25 2026)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_ and _Giorgio Balzarotti_, Jul 26 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004079 | OEIS-A120778

Numerators of partial sums of Catalan numbers scaled by powers of 1/4.

a(n) = A141244(2n+2) = A141244(2n+3) (conjectural). - _Greg Martin_, Aug 16 2014, corrected by _M. F. Hasler_, Aug 18 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jul 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004080 | OEIS-A120811

Positive integers n such that n+d+1 is prime for all proper divisors d of n. Generalization of twin prime to all integers.

This sequence (A120811) is a generalization of twin prime (A001359), the sequence A120776 is a generalization of Sophie Germain prime (A005384), while A120806 is the generalization of Sophie Germain twin prime (A045536). The same observations apply to A120811 as to A120806: the elements are (a) twin primes, (b) semiprimes pq, (c) 3-almost-primes, (d) 4-almost-primes. Moreover, the sequence includes all twin primes but in (b), (c) and (d) the containments are proper. The first occurrence of (d) is A120811(3980)=3^3*13147. Any others? A120811 CONJECTURE: These are all the elements, that is, no element of A120811 has more than 3 prime factors with no degree (sum of exponents) higher than 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Walter Kehowski_, Jul 07 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004081 | OEIS-A120860

Fixed-j dispersion for Q = 8: Square array D(g,h) (g, h >= 1), read by ascending antidiagonals.

Indeed, _Clark Kimberling_, the author of this sequence and of sequence A187968, conjectured that A187968 is the complement of A098021, and that A098021 is the complement of the first column of array D here. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 09 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004082 | OEIS-A120862

Fixed-j dispersion for Q = 13: array D(g,h) (g, h >= 1), read by ascending antidiagonals.

For arbitrary g >= 1, once row g is defined, define D(g+1,1) = least positive integer not in rows 1,2,...,g; D(g+1,2) = f(D(g+1,1)); and D(g+1,h) = 11*D(g+1,h-1) - D(g+1,h-2) for h >= 3. All rows after row 1 are thus inductively defined. [Corrected using Conjecture 1 in Kimberling (2007) by _Petros Hadjicostas_, Jul 07 2020]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 09 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004083 | OEIS-A120863

Fixed-k dispersion for Q = 13: array D(g,h) (g, h >= 1), read by ascending antidiagonals.

For arbitrary g >= 1, once row g is defined, define D(g+1,1) = least positive integer not in rows 1,2,...,g; D(g+1,2) = f(D(g+1,1)); and D(g+1,h) = 11*D(g+1,h-1) - D(g+1,h-2) for h >= 3. All rows after row 1 are thus inductively defined. [Corrected using Conjecture 2 in Kimberling (2007) by _Petros Hadjicostas_, Jul 07 2020]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 09 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004084 | OEIS-A120890

Ordered odd leg of primitive Pythagorean triangles.

Conjecture: lim_{n->oo} a(n)/n = 1/Pi. Limit is also conjectured to be equal to lim_{n->oo} A120427(n)/n, see Selle reference, chapter 2.3.10. - _Lothar Selle_, Jun 21 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Jul 12 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004085 | OEIS-A121065

a(n) is the smallest number in English which contains n letter 'E's.

a(n) = a(n-1) + 1000*a(n-7) - 1000*a(n-8) for n > 9 (conjectured). G.f.: (-1000*x^9 + 3000*x^8 - 1800*x^7 + 6*x^6 + 94*x^5 + 6*x^4 + 8*x^3 + 3*x^2 - 2*x + 2)/((x - 1)*(1000*x^7 - 1)) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ray G. Opao_, Aug 10 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004086 | OEIS-A121082

Number of primitive Pythagorean-like triples a^2+b^2=c^2+k for k=-1 with c<=10^n.

It is conjectured by the first author that a(n)/10^n as n->inf is 1/8 = 0.125.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tito Piezas III_, Aug 11 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004087 | OEIS-A121083

Number of primitive Pythagorean-like triples a^2+b^2=c^2+k for k=-2 with 0<c<=10^n.

It is conjectured by the first author that a(n)/10^n as n->inf is 1/(4*sqrt(2)) = 0.17677...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tito Piezas III_, Aug 11 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004088 | OEIS-A121085

Number of primitive Pythagorean-like triples a^2+b^2=c^2+k for k=-3 with 0<c<=10^n.

It is conjectured by the first author that a(n)/10^n as n->inf is 1/(2*sqrt(3)) = 0.28867...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tito Piezas III_, Aug 11 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004089 | OEIS-A121087

Number of primitive Pythagorean-like triples a^2+b^2=c^2+k for k=-5 with 0<c<=10^n.

It is conjectured by the first author that a(n)/10^n as n->inf is 1/(2*sqrt(5)) = 0.22360...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tito Piezas III_, Aug 11 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004090 | OEIS-A121111

Trajectory of 277777788888899 under "x -> product of digits of x" map.

This is the smallest number of persistence 11 (cf. A003001). It is conjectured that no number has persistence greater than 11.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 12 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004091 | OEIS-A121177

Catapolyoctagons (see Cyvin et al. for precise definition).

The conjecture by _Philipp Emanuel Weidmann_ (see link below) is correct. In Cyvin et al. (1997), this sequence has a double meaning. See Eqs. (6) and (7) and Table I on p. 58 in that paper. The terms of the sequence are related to the enumeration of unbranched catapolyoctagons.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 15 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004092 | OEIS-A121217

a(1)=1, a(2)=2, a(3)=3; for n > 3, a(n) is the smallest positive integer which does not occur earlier in the sequence and which is not coprime to a(n-2).

Conjecture: this is a permutation of the positive integers, cf. A256618. - _Reinhard Zumkeller_, Apr 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Aug 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004093 | OEIS-A121254

Number of conjugated cycles composed of six carbons in (n,n)-nanotubes in terms of the number of naphthalene units.

Conjecture: a(n)=6a(n-1)-11a(n-2)+6a(n-3)-a(n-4)=4*A117202(n). G.f.: 4*x*(1-2*x+2*x^2)/(x^2-3*x+1)^2. [From _R. J. Mathar_, Mar 18 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Parthasarathy Nambi_, Aug 22 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004094 | OEIS-A121255

Number of conjugated cycles composed of ten carbons in (n,n)-nanotubes in terms of the number of naphthalene units.

Conjecture: a(n)=6a(n-1)-11a(n-2)+6a(n-3)-a(n-4). G.f.: -4*x^2*(-2+3*x)/(x^2-3*x+1)^2. [From _R. J. Mathar_, Mar 18 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Parthasarathy Nambi_, Aug 22 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004095 | OEIS-A121285

a(0) = 1; for n>0, a(n) = (n+3)*2^(n-2)-n*binomial(n-1, floor((n-1)/2))-(n-1)*binomial(n-2,floor((n-2)/2)).

Conjecture: -(n-1)*(3*n^2-27*n+56)*a(n) +2*(n-5)*(3*n^2-12*n+5)*a(n-1) +4*(3*n^3-33*n^2+110*n-118)*a(n-2) -8*(n-3)*(3*n^2-21*n+32)*a(n-3)=0. - _R. J. Mathar_, Jan 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004096 | OEIS-A121320

Number of vertices in all ordered (plane) trees with n edges that are at distance two from all the leaves above them.

Conjecture: (-n+2)*a(n) +(5*n-12)*a(n-1) +(-3*n+8)*a(n-2) +2*(-2*n+5)*a(n-3)=0. - _R. J. Mathar_, Jun 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Louis Shapiro_, Aug 25 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004097 | OEIS-A121346

Conjectured lower bound for the number of spheres of radius 1 that can be packed in a sphere of radius n.

Conjectured lower bound for the number of spheres of radius 1 that can be packed in a sphere of radius n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jul 22 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004098 | OEIS-A121411

Positive integers k for which there are primes of the form a^2+k^n=b^2+k^m with positive integers (a,b,m,n) and a > b.

Conjecturally, a(n) is the n-th positive nonsquare integer that is not congruent to -1 mod 4, nor to -1 mod 5, nor to -7 mod 16.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Broadhurst_, Jul 29 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004099 | OEIS-A121437

Matrix inverse of triangle A122177, where A122177(n,k) = C(k*(k+1)/2 + n-k + 2, n-k) for n>=k>=0.

Conjecture: The sequence (column 2 of triangle) 14, -62, 365, -2790, 26749, ... is described by a series of nested sums:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 27 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004100 | OEIS-A121451

Maximum product over partitions into parts of the form 3k+2.

Conjecture. a(1)=a(3)=0, otherwise a(n)=2^(n/2) if n is even and a(n)=5*2^((n-5)/2) if n is odd. (This has been verified for up to n=40.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Apr 26 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004101 | OEIS-A121507

Conjectured list of numbers whose aliquot sequence eventually reaches a cycle of length two or more.

Conjectured list of numbers whose aliquot sequence eventually reaches a cycle of length two or more.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joshua Zucker_, Aug 04 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004102 | OEIS-A121508

Conjectured list of numbers whose aliquot sequence eventually reaches a cycle of length two or more, but which are not themselves part of the cycle.

Conjectured list of numbers whose aliquot sequence eventually reaches a cycle of length two or more, but which are not themselves part of the cycle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joshua Zucker_, Aug 04 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004103 | OEIS-A121510

Conjectured to be complete list of cycle lengths for 3x+1 (or Collatz) problem.

Conjectured to be complete list of cycle lengths for 3x+1 (or Collatz) problem. The sequence counts cycles over the domain of all integer values: (0), (-1, -2), (4, 2, 1), (-5, -14, -7, -20, -10), and (-17, -50, -25, -74, -37, -110, -55, -164, -82, -41, -122, -61, -182, -91, -272, -136, -68, -34); the generalized cycle conjecture asserts that these are the only such cycles (see Kaneda link). - _Michael S. Branicky_, Nov 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | Daniel Asimov (dasimov(AT)earthlink.net), Sep 17 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004104 | OEIS-A121553

Total area of all deco polyominoes of height n. A deco polyomino is a directed column-convex polyomino in which the height, measured along the diagonal, is attained only in the last column.

Conjecture D-finite with recurrence a(n) +(-2*n-3)*a(n-1) +(n^2+4*n-3)*a(n-2) +2*(-n^2+n+3)*a(n-3) +2*(n-3)^2*a(n-4)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Aug 08 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004105 | OEIS-A121571

Largest number that is not the sum of n-th powers of distinct primes.

Fuller & Nichols prove _T. D. Noe_'s conjecture that a(3) = 1866000. They also prove that 483370 positive numbers cannot be written as the sum of cubes of distinct primes. - _Robert Nichols_, Sep 08 2017 Noe conjectures that a(4) = 340250525752 and that 31332338304 positive numbers cannot be written as the sum of fourth powers of distinct primes. - _Charles R Greathouse IV_, Nov 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Aug 08 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004106 | OEIS-A121580

Number of cells in column 1 of all deco polyominoes of height n. A deco polyomino is a directed column-convex polyomino in which the height, measured along the diagonal, is attained only in the last column.

Conjecture D-finite with recurrence a(n) +(-n-4)*a(n-1) +3*(n+1)*a(n-2) +2*(-2*n+3)*a(n-3) +2*(n-3)*a(n-4)=0. - _R. J. Mathar_, Jul 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Aug 09 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004107 | OEIS-A121584

Number of cells in columns 1 and 2 of all deco polyominoes of height n. A deco polyomino is a directed column-convex polyomino in which the height, measured along the diagonal, is attained only in the last column.

Conjecture D-finite with recurrence 14*(-n+1)*a(n) +(14*n^2+1731*n-6995)*a(n-1) +3*(-577*n^2+480*n+7243)*a(n-2) +2*(2781*n^2-11952*n+6004)*a(n-3) +(-5987*n^2+36181*n-54220)*a(n-4) +2*(1071*n-3433)*(n-4)*a(n-5)=0. - _R. J. Mathar_, Jul 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Aug 11 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004108 | OEIS-A121627

Real part of a complex operation analogous to the factorials.

Conjecture (1): Current nonzero term divides next nonzero term; as an operation, getting a new sequence: 12, 8, 5, 84, 16, 9, 220, 24, 13, ...; using unsigned terms). Example: a(5) = -480 and a(7)= 40320, the next nonzero term in the sequence. Then 40320/480 = 84. Conjecture (2): If the term in (12, 8, 5, ...) is generated from a(n)/a(n-1) or a(n)/a(n-2), then n divides each (12, 8, 5, ...). Example: a(11) = -1277337600 and the previous nonzero term = a(9) = 5806080. Then a(11)/a(9) = 220 and 11 divides 220: 220/11 = 20. Due to the statements (1) and (2) below a(n) divides a(n+k) for all k>=0, only provided both the residues of n (mod 8) and (n+k) (mod 8) are 0, 1, 3, 4, 5 or 7. As a consequence it follows that conjecture (1) plainly holds true (set k=1). From statements (1) and (2) it further follows that n! divides a(n) (for n with a(n)<>0) and therefore conjecture(2) also holds true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Aug 12 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004109 | OEIS-A121636

Number of 2-cell columns starting at level 0 in all of deco polyominoes of height n. A deco polyomino is a directed column-convex polyomino in which the height, measured along the diagonal, is attained only in the last column.

Conjecture D-finite with recurrence a(n) +(-2*n+3)*a(n-1) +(n^2-5*n+7)*a(n-2) +(n-3)^2*a(n-3)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Aug 13 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004110 | OEIS-A121639

Number of 2-cell columns in all deco polyominoes of height n. A deco polyomino is a directed column-convex polyomino in which the height, measured along the diagonal, is attained only in the last column.

Conjecture D-finite with recurrence a(n) +(-2*n+1)*a(n-1) +n*(n-2)*a(n-2) +(2*n-7)*a(n-3) -(n-3)*(n-5)*a(n-4)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Aug 14 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004111 | OEIS-A121669

Numbers with sum of digits = 19, divisible by 19 and containing the string "19".

Conjecture: There are approximately k(n-1)(n-2)^(n-2) terms of this sequence up to 10^n, where k is about e/(19e-19). - _Charles R Greathouse IV_, Oct 13 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | Hassan Taifour (hytaifour(AT)yahoo.co.uk), Sep 10 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004112 | OEIS-A121695

Number of odd-length first columns in all deco polyominoes of height n. A deco polyomino is a directed column-convex polyomino in which the height, measured along the diagonal, is attained only in the last column.

Conjecture D-finite with recurrence a(n) +a(n-1) -n*(n-2)*a(n-2) -(2*n-3)*(n-2)*a(n-3) -(n-2)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Aug 17 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004113 | OEIS-A121696

Number of even-length first columns in all deco polyominoes of height n. A deco polyomino is a directed column-convex polyomino in which the height, measured along the diagonal, is attained only in the last column.

Conjecture D-finite with recurrence a(n) -(n-1)^2*a(n-2) -n*(n-2)*a(n-3)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Aug 17 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004114 | OEIS-A121716

Fixed points of the permutations A121711, A121713, A121714 and A121715.

Conjecture: the sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Aug 18 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004115 | OEIS-A121724

Generalized central binomial coefficients for k=2.

Conjecture: (n+1)*a(n) = 5*(n+1)*a(n-1) + 16*(n-2)*a(n-2) - 80*(n-2)*a(n-3). - _R. J. Mathar_, Nov 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Aug 17 2006, Feb 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004116 | OEIS-A121725

Generalized central coefficients for k=3.

Conjecture: (n+1)*a(n) = 10*(n+1)*a(n-1) + 36*(n-2)*a(n-2) - 360*(n-2)*a(n-3). - _R. J. Mathar_, Nov 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Aug 17 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004117 | OEIS-A121747

Number of columns of odd length in all deco polyominoes of height n. A deco polyomino is a directed column-convex polyomino in which the height, measured along the diagonal, is attained only in the last column.

Conjecture D-finite with recurrence (-9*n+38)*a(n) +3*(3*n^2-19*n+27)*a(n-1) +(9*n^3-37*n^2-24*n+127)*a(n-2) +(-9*n^4+102*n^3-392*n^2+525*n-85)*a(n-3) -(n-3)*(28*n^3-223*n^2+510*n-237)*a(n-4) -(19*n-15)*(n-3)*(n-4)^2*a(n-5)=0. - _R. J. Mathar_, Jul 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Aug 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004118 | OEIS-A121750

Number of columns of even length in all deco polyominoes of height n. A deco polyomino is a directed column-convex polyomino in which the height, measured along the diagonal, is attained only in the last column.

Conjecture D-finite with recurrence a(n) +(-n-2)*a(n-1) +(-n^2+4*n-2)*a(n-2) +(n^3-3*n^2-4*n+11)*a(n-3) -(n-1)*(n^2-11*n+26)*a(n-4) +(-n^3+5*n^2+7*n-43)*a(n-5) +(n-3)*(n-5)^2*a(n-6)=0. - _R. J. Mathar_, Jul 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Aug 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004119 | OEIS-A121751

Number of deco polyominoes of height n in which all columns end at an even level. A deco polyomino is a directed column-convex polyomino in which the height, measured along the diagonal, is attained only in the last column.

Conjecture D-finite with recurrence +256*a(n) -384*a(n-1) +16*(-8*n^2+40*n-67)*a(n-2) +16*(8*n^2-54*n+87)*a(n-3) +4*(4*n^4-56*n^3+242*n^2-304*n-21)*a(n-4) +4*(-2*n^4+38*n^3-249*n^2+647*n-554)*a(n-5) +(n-4)*(n-8)*(n^2-12*n+31)*a(n-6)=0. - _R. J. Mathar_, Jul 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Aug 23 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004120 | OEIS-A121752

Number of columns ending at an odd level in all deco polyominoes of height n. A deco polyomino is a directed column-convex polyomino in which the height, measured along the diagonal, is attained only in the last column.

Conjecture D-finite with recurrence (-460*n+1223)*a(n) +(460*n^2+460*n-5569)*a(n-1) +(460*n^3-3826*n^2+7853*n+1313)*a(n-2) +(-460*n^4+1840*n^3+5501*n^2-31045*n+31726)*a(n-3) +(1223*n^4-13205*n^3+45787*n^2-51389*n+558)*a(n-4) -2*(426*n-1111)*(n-6)*(n-4)^2*a(n-5)=0. - _R. J. Mathar_, Jul 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Aug 23 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004121 | OEIS-A121754

Number of columns ending at an even level in all deco polyominoes of height n. A deco polyomino is a directed column-convex polyomino in which the height, measured along the diagonal, is attained only in the last column.

Conjecture D-finite with recurrence 16*(n+1)*a(n) +(-16*n^2-178*n+531)*a(n-1) +(-16*n^3+178*n^2-393*n-510)*a(n-2) +(16*n^4+98*n^3-1439*n^2+4222*n-3623)*a(n-3) +(-146*n^4+1479*n^3-4483*n^2+3054*n+2841)*a(n-4) +(130*n-311)*(n-6)*(-4+n)^2*a(n-5)=0. - _R. J. Mathar_, Jul 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Aug 23 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004122 | OEIS-A121757

Triangle read by rows: multiply Pascal's triangle by 1,2,6,24,120,720,... = A000142.

a(n,k) = D(n+1,k+1) Array D in A253938 is part of a conjectured formula for F(n,p,r) that relates Dyck path peaks and returns. a(n,k) was discovered prior to array D. - _Roger Ford_, May 19 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alford Arnold_, Aug 19 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004123 | OEIS-A121782

Series expansion for mean-squared radius of gyration of rectangles on square lattice.

Conjectures from _Colin Barker_, Apr 17 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 30 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004124 | OEIS-A121805

The "comma sequence": the lexicographically earliest sequence of positive numbers with the property that the sequence formed by the pairs of digits adjacent to the commas between the terms is the same as the sequence of successive differences between the terms.

Conjecture: This sequence is finite, for any initial term. - _N. J. A. Sloane_, Nov 14 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_, Dec 11 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004125 | OEIS-A121816

Conjectured chromatic number of the square of an outerplanar graph G^2 as a function of the maximum degree of a vertex of G.

Conjectured chromatic number of the square of an outerplanar graph G^2 as a function of the maximum degree of a vertex of G.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Aug 26 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004126 | OEIS-A121860

a(n) = Sum_{d|n} n!/(d!*(n/d)!).

Conjecture: the sequence a(n) taken modulo a positive integer k >= 3 eventually becomes constant equal to 2. For example, the sequence taken modulo 11 is [1, 2, 2, 8, 2, 1, 2, 10, 6, 3, 2, 2, 2, 2, 2, 2, ...]. - _Peter Bala_, Aug 08 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Sep 09 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004127 | OEIS-A121861

Least previously nonoccurring positive integer such that partial sum + 1 is prime.

Conjecture: a(n) = {1,3} UNION {permutation of even positive numbers}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Aug 30 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004128 | OEIS-A121878

a(1)=1. a(n) = the smallest positive integer not occurring earlier in the sequence such that a(n-1)+a(n) is squarefree.

Conjectured to be a permutation of the natural numbers. - _Derek Orr_, Jun 01 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Aug 31 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004129 | OEIS-A121926

a(n) = prime(n) + n!.

It was conjectured by _Alexander R. Povolotsky_ and proved (in SeqFan email exchange) by _David L. Harden_ (with proof improvement from Daniel Berend) that no value of n exists such that (n! + prime(n)) yields an integral square. He also conjectured (see the link) that no value of n exists such that (n! + prime(n)) yields an integral m^k where k>1. - _Alexander R. Povolotsky_, Aug 13 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 13 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004130 | OEIS-A122045

Euler (or secant) numbers E(n).

Odd primes p such that p | E(p-1) are primes p == 1 (mod 4), A002144. Conjecture: odd composites m such that m | E(m-1) are Carmichael numbers m such that p == 1 (mod 4) for every prime p|m, A265237. - _Thomas Ordowski_, Feb 06 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Sep 13 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004131 | OEIS-A122047

Degree of the polynomial P(n,x), defined by a Somos-6 type sequence: P(n,x)=(x^(n-1)*P(n-1,x)*P(n-5,x) + P(n-2,x)*P(n-4,x))/P(n-6,x), initialized with P(n,x)=1 at n<0.

Conjectures from _R. J. Mathar_, Jul 15 2008: (Start) Conjecture: a(n) = (A000292(n+1) - n - 2 - (-1)^floor((n-1)/5)*A099443(n+1))/5. - _R. J. Mathar_, Jul 15 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Sep 13 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004132 | OEIS-A122129

Expansion of 1 + Sum_{k>0} x^k^2/((1-x)(1-x^2)...(1-x^(2k))).

Wiseman's first conjecture above was proved by Gordon, Theorem 7. For two other combinatorial interpretations of this sequence see Connor, Proposition 1. - _Peter Bala_, Dec 22 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Aug 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004133 | OEIS-A122130

Expansion of f(-x^4, -x^16) / psi(-x) in powers of x where psi() is a Ramanujan theta function and f(,) is Ramanujan's general theta function.

Wiseman's first conjecture above was proved by Connor, Proposition 2. - _Peter Bala_, Dec 22 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Aug 21 2006, corrected Aug 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004134 | OEIS-A122134

Expansion of Sum_{k>=0} x^(k^2+k)/((1-x)(1-x^2)...(1-x^(2k))).

Conjecture: Also the number of even-length integer partitions y of n such that y_i != y_{i+1} for all even i. For example, the a(2) = 1 through a(9) = 7 partitions are: For Wiseman's conjecture above and three other partition-theoretic interpretations of this sequence see Connor, Proposition 3. - _Peter Bala_, Jan 02 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Aug 21 2006, Oct 10 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004135 | OEIS-A122135

Expansion of f(x, -x^4) / phi(-x^2) in powers of x where f(,) and phi() are Ramanujan theta functions.

Conjecture: Also the number of integer partitions y of n such that y_i > y_{i+1} for all even i. For example, the a(1) = 1 through a(9) = 12 partitions are: For Wiseman's conjecture above and three other partition-theoretic interpretations of this sequence see Connor, Proposition 4. - _Peter Bala_, Jan 02 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Aug 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004136 | OEIS-A122224

Length of the longest possible self avoiding path on the 2-dimensional square lattice such that the path fits into a circle of diameter n.

It is conjectured that a(n) >= A123690(n)-1, i.e., that it is always possible to find a path visiting all grid points covered by a circle, irrespective of the position of its center. - _Hugo Pfoertner_, Mar 02 2018 This conjecture is false. A counterexample is the configuration with 81 points and a diameter of 10, shown in the pdf with the illustrations for A123690. Not all of the four isolated points on the axes and the circumference of the enclosing circle, each with only one neighbor in the grid, can be part of a connecting Hamiltonian path. - _Hugo Pfoertner_, Jun 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Sep 25 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004137 | OEIS-A122226

Length of the longest possible self-avoiding path on the 2-dimensional triangular lattice such that the path fits into a circle of diameter n.

It is conjectured that the sequence is identical with A125852 for all n>1. That means that it is always possible to find an Hamiltonian cycle on the maximum possible number of lattice points that can be covered by circular disks of diameter >=2. For the given additional terms it was easily possible to construct such closed paths by hand, using the lattice subset found by the exhaustive search for A125852. See the examples at the end of the linked pdf file a122226.pdf that were all generated without using a program. - _Hugo Pfoertner_, Jan 12 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Sep 25 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004138 | OEIS-A122249

Numerators of Hankel transform of 1/(2n+1).

Conjecture - requires proof.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Aug 27 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004139 | OEIS-A122270

Numbers m such that the numerator of the Bernoulli number B(m) is divisible by a cube.

Conjecture: For all regular primes p > 3 and integers k > 0, the numerator of the Bernoulli number B(2*p^k) is divisible by p^k. Moreover, for all regular primes p > 3 and integers k > 0, m = 2*p^k is the smallest index such that the numerator of the Bernoulli number B(m) is divisible by p^k. Also, for all regular primes p > 3 and integers k > 0, all m such that the numerator of the Bernoulli number B(m) is divisible by p^k are of the form m = 2*s*p^k, where s > 0 is an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Aug 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004140 | OEIS-A122276

If b(n-1) + b(n-2) < n then a(n) = 0, otherwise a(n) = 1, where b(i) = A096535(i).

Conjecture: lim {n -> infinity} x_n / y_n = 1, where x_n is the number of j <= n such that A096535(j) = A096535(j-1) + A096535(j-2) and y_n is the number of j <= n such that A096535(j) = A096535(j-1) + A096535(j-2) - j. Computational support: x_n / y_n = 0.9999917 for n = 10^9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Aug 29 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004141 | OEIS-A122378

Numbers m such that m^2 > S(m)!, where S(m)! is the smallest factorial divisible by m.

It is conjectured that m^2 < S(m)! for almost all m.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Sep 03 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004142 | OEIS-A122379

Numbers n such that S(n)! > n^2 > P(n)!, where S(n)! is the smallest factorial divisible by n and P(n) is the greatest prime factor of n.

It is conjectured that n^2 < P(n)! for almost all n. It is known that S(n) = P(n) for almost all n. Clearly, S(n) >= P(n) for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Sep 03 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004143 | OEIS-A122380

Numbers k such that k^2 > P(k)!, where P(k) is the greatest prime factor of k.

It is conjectured that k^2 < P(k)! for almost all k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Sep 03 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004144 | OEIS-A122399

a(n) = Sum_{k=0..n} k^n * k! * Stirling2(n,k).

Conjecture: Let p be prime. The sequence obtained by reducing a(n) modulo p for n >= 1 is purely periodic with period p - 1. For example, modulo 7 the sequence becomes [1, 2, 1, 3, 3, 0, 1, 2, 1, 3, 3, 0, ...], with an apparent period of 6. Cf. A338040. - _Peter Bala_, May 31 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Aug 31 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004145 | OEIS-A122437

Allowable values of the "dropping time" of the Collatz (3x+1) iteration.

Whether only these values occur depends on Terras's conjecture on the coefficient stopping time; see the comment of Olivier Rozier in A126241. - _Ruud H.G. van Tol_, Oct 11 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Sep 06 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004146 | OEIS-A122535

Smallest prime of a triple of successive primes, where the middle one is the arithmetic mean of the other two.

These are primes for which the subsequent prime gaps are equal, so (p(k+2)-p(k+1))/(p(k+1)-p(k)) = 1. It is conjectured that prime gaps ratios equal to one are less frequent than those equal to 1/2, 2, 3/2, 2/3, 1/3 and 3. - _Andres Cicuttin_, Nov 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Miklos Kristof_, Sep 18 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004147 | OEIS-A122536

Number of binary sequences of length n with no initial repeats (or, with no final repeats).

Conjecture: a_n ~ C * 2^n where C is 0.27004339525895354325... [Chaffin, Linderman, Sloane, Wilks, 2012]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sarah Nibs_, Sep 18 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004148 | OEIS-A122537

a(1) = 1; for n>1, a(n) is smallest number greater than a(n-1), divisible by n and not equal to any a(i)+a(j) with i and j <= n-1.

Jonathan Vos Post asks if 1, 2, 4 and 5 are the only values of n for which n^2 divides a(n), Sep 19 2006. _J. Lowell_, Oct 02 2006 remarks that n = 1, 2, 4, 5 and 10 have this property and conjectures that there are no other values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Sep 18 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004149 | OEIS-A122648

Expansion of e.g.f.: exp(x^2)*(exp(2*x)+1)/2.

Conjecture: D-finite with recurrence a(n) -2*a(n-1) +2*(-2*n+3)*a(n-2) +4*(n-2)*a(n-3) +4*(n-2)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Aug 20 2021 Mathar's conjecture is true (see Fried link). - _Sela Fried_, Dec 07 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 22 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004150 | OEIS-A122707

Smallest prime equal to the sum of squares of 8 consecutive primes divided by 2^n, or 1 if such a prime does not exist.

Conjecture: a(n) > 1 exists for all n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Sep 23 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004151 | OEIS-A122722

Number of triangulations of Delta^2 x Delta^(k-1).

Conjectures: a(n) = n! * A273464(n, n*(n+1)/2) for n >= 1; a(n) = A011555(n-1) for n >= 2. [A273464(n,k) is defined for n >= 1 and 0 <= k <= n*(n+1)/2.] - _Petros Hadjicostas_, Sep 12 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Oct 22 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004152 | OEIS-A122835

Number of topologies on n labeled elements in which no element belongs to any pair of noncomparable members of the topology.

Conjectural closed form for n>0: 3*2^(k-3)(LerchPhi[1/4, -k, 1/2] + 2 PolyLog[-k, 1/4]) - 1/2. - _Vladimir Reshetnikov_, Jan 07 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | Nathan K. McGregor (mcgregnk(AT)ese.wustl.edu), Sep 15 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004153 | OEIS-A122858

Expansion of E(k) * K(k) * (2/Pi)^2 in powers of q^2 where E(), K() are complete elliptic integrals and the nome q = exp(-Pi * K(k') / K(k)).

Conjecture: -3 A122858(n) - A229616(n) + 4 A282031(n) = 0 for all n. - _Thomas Baruchel_, Jun 23 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Sep 15 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004154 | OEIS-A122863

Smallest integer not the sum of nonzero powers of previous terms.

It can be shown that a(n) is even for n >= 2 and a(n+1)/a(n) >= 2. Conjecture: lim_{n->infinity} a(n+1)/a(n) = e and lim_{n->infinity} a(n)/e^(n-1) = 1. - _Ya-Ping Lu_, Feb 18 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_, Sep 15 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004155 | OEIS-A122869

Primes p that divide Lucas((p-1)/2), where Lucas is A000032.

Conjecture: This sequence is just the primes congruent to 11 or 19 mod 20. - _Charles R Greathouse IV_, May 25 2011 [The conjecture is correct. - _Jianing Song_, Jun 20 2025]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Sep 16 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004156 | OEIS-A122920

Diagonal sums of number triangle A122919.

Conjecture: 2*n*(n+3)*a(n) - (7*n^2+9*n+4)*a(n-1) - 2*(n+1)*(2*n+1)*a(n-2) = 0. - _R. J. Mathar_, Nov 05 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 19 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004157 | OEIS-A122946

a(0)=a(1)=0, a(2)=2; for n >= 3, a(n) = a(n-1) + 4*a(n-3).

Conjecture: count of even Markov numbers in generation n (with generations 0, 1 and 2 labeled as {5}, {13, 29} and {34, 194, 433, 169}. (Checked up to generation 20.) - _Wouter Meeussen_, Jan 16 2024 Wouter Meeussen's conjecture is true. Proof: label the Markov tree with Markov triples according to the scheme described at A368546. Mod 2, the triples are: row 0: (1,1,0); row 1: (1,1,1), (1,1,0); row 2: (1,0,1), (1,0,1), (1,1,1), (1,1,0); row 3: (1,1,0), (0,1,1), (1,1,0), (0,1,1), (1,0,1), (1,0,1), (1,1,1), (1,1,0); etc. Note that the Markov number labels of the tree (the center numbers of the triples) in rows 0 and 1 include no even numbers, while those in row 2 include two even numbers. Observing that the second triple in row 1 and the first four triples in row 3 are the same or the reverse of the root triple, and noting that every vertex in row 3 and beyond is in a subtree with one of these triples as root, the recurrence follows. - _William P. Orrick_, Mar 05 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 24 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004158 | OEIS-A122955

Numbers k such that although the number of digits in 5^k is larger, the sum of the digits is less than that of the previous power of 5.

Conjecture: Natural density is log 5/log 100. - _Charles R Greathouse IV_, Nov 15 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Oct 25 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004159 | OEIS-A123050

Conjectured number of ordered trees on n edges for which the conjugate and transpose commute.

Conjectured number of ordered trees on n edges for which the conjugate and transpose commute.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Callan_, Sep 25 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004160 | OEIS-A123066

(Number of numbers <= n with an odd number of distinct prime factors) - (number of numbers <= n with an even number of distinct prime factors).

Analog of A072203 for number of distinct factors. Conjecture that sequence changes sign infinitely often, although the next sign change is probably large.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Sep 26 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004161 | OEIS-A123070

Hofstadter Flip-G-sequence.

Conjecture: a(0) = 0, a(1) = 1, a(2) = 1, a(3) = 2; a(n) = (n+1) - a(a(n-1)+1) for n>=4. [Conjecture has been proved; see Letouzey link. - _Pierre Letouzey_, Sep 09 2015] Conjecture: a(0) = 0, a(1) = 1, a(2) = 1, a(3) = 2; for n>=4, let T = n-a(n-1). If T = (n-1) - a((n-1)-1) then a(n) = max{a-inv(T)} otherwise a(n) = min{a-inv(T)}, where a-inv(n) is the inverse of a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Albert Neumueller (albert.neu(AT)gmail.com), Sep 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004162 | OEIS-A123071

Bishops on a 2n+1 X 2n+1 board (see Robinson paper for details).

Conjecture: 2*a(n) +a(n-1) -2*n*a(n-2) +(-n-10)*a(n-3) -2*(n-2)*(n+2)*a(n-4) +(-n^2-2*n+23)*a(n-5) +2*(n-5)*(n^2-7*n+11)*a(n-6) +(n-6)*(n-5)^2*a(n-7)=0. - _R. J. Mathar_, Apr 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004163 | OEIS-A123078

Integers that are the arithmetic mean of 1000 consecutive primes.

Conjecture: a(n) ~ 1000 n log n. - _Charles R Greathouse IV_, Oct 23 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Sep 27 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004164 | OEIS-A123120

Numbers k>0 such that m+k is not the sum of m nonzero squares for any m>5.

Conjecture: All but (n+6) positive numbers are equal to the sum of n>5 nonzero squares. For all n>5 the only (n+6) positive numbers that are not equal to the sum of n nonzero squares are {1,2,3,...,n-3,n-2,n-1,n+1,n+2,n+4,n+5,n+7,n+10,n+13}. The above conjecture appears as Theorem 2 on p. 73 in the Grosswald reference, where it is attributed to E. Dubouis (1911). - _Wolfdieter Lang_, Mar 27 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Sep 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004165 | OEIS-A123145

a(1) = 1, a(n) = a(n-1) if n == 1 (mod 4), otherwise a(n) = n * a(n-1) for n >= 2.

Conjecture: E.g.f.: E(x)=d(G(0))/dx where G(k) = 1 + x/(4*k+1 - x*(4*k+1)/(1 + x - x/(1 + x - x/(x + 1/G(k+1))))), or shift on 1 left G(0); (continued fraction,5-step). - _Sergei N. Gladkovskii_, Nov 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Oct 01 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004166 | OEIS-A123159

Conjectured smallest Sierpiński numbers of the second kind S, base b=2,3,4,5,..., where S*b^n+1 is composite for all n>=1 and gcd(S+1, b-1) = 1.

Conjectured smallest Sierpiński numbers of the second kind S, base b=2,3,4,5,..., where S*b^n+1 is composite for all n>=1 and gcd(S+1, b-1) = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Robert Smith (robert_smith44(AT)hotmail.com), Oct 02 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004167 | OEIS-A123164

Row sums of A123160.

Conjecture: a(n*p^k) == a(n*p^(k-1)) mod(p^(3*k)) for prime p >= 5 and all positive integers n and k. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Oct 02 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004168 | OEIS-A123167

Continued fraction for c=sqrt(2)*(exp(sqrt(2))+1)/(exp(sqrt(2))-1). a(2*n-1) = 8*n-6, a(2*n) = 4*n-1.

Conjecture: Numerator of 4/n - 2/n^2. - _Wesley Ivan Hurt_, Jul 11 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 02 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004169 | OEIS-A123233

Difference between the (10^n)-th prime and the Riemann-Gram approximation of the (10^n)-th prime.

The algorithm in the PARI script below produces the 10^n-th prime accurate to first n/2 places. Conjecture: The sign of the terms in this sequence changes infinitely often. Based on the small sample presented here, it appears the negative terms occur much more often.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Oct 06 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004170 | OEIS-A123329

Let M be the matrix defined in A111490. Sequence gives M(2,1)-M(1,2), M(2,1)+M(3,1)+M(3,2)-M(1,2)-M(1,3)-M(2,3), etc.

Conjectures: In the three conjectures assuming that here the offset is 1. a(n) = A175254(n+1) - A000292(n+1), conjectured by _Omar E. Pol_, Jan 20 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_ and _Giorgio Balzarotti_, Oct 06 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004171 | OEIS-A123348

Kekulé numbers for certain benzenoids (see the Cyvin-Gutman book for details).

Conjectured g.f.: 1/(1 - Q(0)) - 1, where Q(k)= 1 - 1/(4^k - 2*x*16^k/(2*x*4^k - 1/(1 + 1/(2*4^k - 8*x*16^k/(4*x*4^k + 1/Q(k+1)))))); (continued fraction). - _Sergei N. Gladkovskii_, Apr 13 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 10 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004172 | OEIS-A123365

Values of k such that A046530(k) = (k+2)/3, where A046530(k) is the number of distinct residues of cubes mod k.

Conjecture: With the exception of the first term a(1)=1, this is exactly the sequence of primes of the form 6k+1 (A002476). This has been verified up to a(n)=2000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Oct 12 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004173 | OEIS-A123371

Smallest number k>1 such that (1 + Sum[i^(2n - 1), {i,1,k}]) is prime, or 0 if no such number k exists.

Conjecture: a(n)>0 exists for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Nov 09 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004174 | OEIS-A123375

Least k such that the difference between consecutive semiprimes A065516(k) equals n, or 0 if no such k exists.

a(n) equals least k such that A065516(k) = n, or 0 if no such k exists. Conjecture: a(n)>0 exists for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Nov 09 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004175 | OEIS-A123487

Smallest prime q such that (q^p-1)/(q-1) is prime, where p = prime(n); or 0 if no such prime q exists.

Conjecture: Never is a(n) equal to 0. Jones & Zvonkin conjecture (as did Robert G. Wilson v above) that a(n) > 0 for all n. - _Charles R Greathouse IV_, Jul 23 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Sep 30 2006, Oct 02 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004176 | OEIS-A123509

Rohrbach's problem: a(n) is the largest integer such that there exists a set of n integers that is a basis of order 2 for (0, 1, ..., a(n)-1).

According to Zhining Yang, Jul 08 2017, a(13) to a(20) are 65, 70, 79, 90, 101, 112, 123, 134, but there is some doubt about these terms, and they should be confirmed before they are accepted. They do not agree with the conjecture, so perhaps the VBA program is not correct. a(n) = A001212(n-1)+1 (conjecture). - _R. J. Mathar_, Oct 08 2006. Comment from _Martin Fuller_, Mar 18 2009: I agree with this conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Warren D. Smith_, Oct 02 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004177 | OEIS-A123689

Number of points in a square lattice covered by a circle of diameter n if the center of the circle is chosen such that the circle covers the minimum possible number of lattice points.

Using brute force computation and a step size of 1/1000 (though 1/200 suffices), the [conjectured] terms a(21) to a(40) would be: 332, 374, 408, 438, 484, 522, 560, 608, 648, 698, 740, 794, 846, 894, 952, 1006, 1060, 1124, 1184, 1248. - _Jean-François Alcover_, Jan 08 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Oct 09 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004178 | OEIS-A123710

Indices k such that 4 = A123709(k) = number of nonzero terms in row k of triangle A123706.

a(n) = A025475(n) for n>2 (conjectured). - _M. F. Hasler_, Feb 12 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 09 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004179 | OEIS-A123737

Partial sums of (-1)^floor(n*sqrt(2)).

Conjecture: A001652(n) is the index of the first occurrence of n in sequence A123737, A001108(n) is the index of the first occurrence of -n in sequence A123737. - _Vaclav Kotesovec_, Jun 02 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Oct 11 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004180 | OEIS-A123922

Number of 2143-avoiding Dumont paths of the 2nd kind of length 2n.

Conjecture: 16*n*(n+2)*(n+1)^2*a(n)
-108*n*(n+1)*(2*n-1)*a(n-1)
-9*(3*n-5)*(3*n-1)*(3*n-4)*(3*n-2)*a(n-2)=0. - _R. J. Mathar_, Jan 25 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Nov 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004181 | OEIS-A123929

Simili-primes of order 2.

Conjecture : a(n) is asymptotic to c*n*log(n) with c about 1.5. - _Benoit Cloitre_, Feb 11 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Hugo van der Sanden_, Nov 22 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004182 | OEIS-A124000

Semiprimes in A006987(n), or semiprime binomial coefficients: C(n,k), 2 <= k <= n-2.

Conjecture: all a(n) except a(1) = 6 and a(2) = 10 are odd. Conjecture: all a(n) except a(5) = 35 are triangular numbers of the form p*(2p +/- 1) that belong to A068443(n) = {6, 10, 15, 21, 55, 91, 253, 703, 1081, 1711, 1891, 2701, ...} Triangular numbers with two distinct prime factors.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Oct 31 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004183 | OEIS-A124168

Union of all n-Fibonacci sequences, that is, all sequences s(0) = s(1) = ... = s(n-2) = 0, s(n-1) = 1 and for k >= n, s(k) = s(k-1) + ... + s(k-n).

Noe and Post conjectured that the only positive terms that are common to any two distinct n-step Fibonacci sequences are the powers of 2 that begin each sequence and 13 (in 2- and 3-step) and 504 (in 3- and 7-step). Perhaps we should also include 8 (in 2- and 4-step). - _T. D. Noe_, Dec 05 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carlos Alves_, Dec 03 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004184 | OEIS-A124197

Number of subsets S of {1,2,3,...,n}, including the empty subset, such that if x and y are in S with x<y and x+y even, then (x+y)/2 is also in S.

The sequence appeared in Problem B3 on the 2009 Putnam exam, which asked one to find all n for which the second difference equals 1. The second difference is the number of such subsets of {1,2,...,n+1} that contain both 1 and n+1. One such subset is {1,2,...,n+1}, and if n has an odd factor d>1 then the arithmetic progression {1,d+1,2d+1,...,n+1} works as well; hence the second difference is 1 iff n is a power of 2. [Note that the Putnam problem uses n+1 for our n.] This also means that the conjectural formula for the second difference is a lower bound. To prove the conjecture, note that consecutive elements of S alternate in parity (else S contains their average); thus if x,s,y are consecutive elements then x+y is even, so s=(x+y)/2, which means that S is a finite arithmetic progression with odd common difference. Since conversely any such arithmetic progression works, we are done. - _Noam D. Elkies_, Dec 05 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Dec 06 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004185 | OEIS-A124247

Numbers n such that n!!-8 and n!!+8 are two consecutive primes.

n is in the sequence iff n!!-8 and n!!+8 are primes and the four numbers n!!-2,n!!+2, n!!-4 & n!!+4 are composites. It's interesting that related to the term 341 we can say 341!!-(3+4+1) and 341!!+(3+4+1) are two consecutive primes with length 341+prime(3+4+1). Conjecture: 341 is the largest term of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Nov 19 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004186 | OEIS-A124254

Forest-and-trees problem: number of trees visible. (See Comments.)

Conjecture: a(n) is the number of pairs of coprime numbers (x,y) such that x <= y and x^2 + y^2 < n^2. The Mathematica program is based on this assumption. How important is the decreasing function 1/n of the trunk's radius with respect to n? Does the sequence change if this function changes? - _Andres Cicuttin_, Feb 24 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Oct 22 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004187 | OEIS-A124352

Number of directed Hamiltonian paths on the n-antiprism graph.

Conjectures from _Colin Barker_, Dec 21 2015: (Start) Equivalent conjecture: a(n) = 2*a(n-1) +3*a(n-2) -2*a(n-3) -6*a(n-4) -4*a(n-5) -a(n-6) +672*n -1376 -72*n^2 if n>=9. - _R. J. Mathar_, Jan 25 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Oct 27 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004188 | OEIS-A124359

Numbers n for which the sum of the digits of n^k, for some k, is equal to n.

Most of the values are conjectural. So far not much is proved about the function A247889. - _M. F. Hasler_ and _Robert Israel_, May 18 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_ and _Giorgio Balzarotti_, Oct 27 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004189 | OEIS-A124367

Numbers that raised to any exponent do not produce a number whose sum of digits is equal to the initial number.

Most of the values are conjectural, so far not much is really proved about the function A247889. - _M. F. Hasler_ and _Robert Israel_, May 18 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_ and _Giorgio Balzarotti_, Oct 30 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004190 | OEIS-A124380

O.g.f.: A(x) = Sum_{n>=0} x^n*Product_{k=0..n} (1 + k*x).

Conjecture: log(a(n)) ~ n*log(n)/2 - n*(1 + log(2))/2. - _Vaclav Kotesovec_, Sep 18 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004191 | OEIS-A124418

Triangle read by rows: T(n,k) is the number of partitions of the set {1,2,...,n} having exactly k blocks that contain both odd and even entries (0<=k<=floor(n/2)).

Conjecture: T(n,k) = k!*A049020([n/2],k)*A049020([(n+1)/2],k) where A049020(n,k)=Sum_{i=0..n} S2(n,i)*C(i,k) and S2(n,k)=(1/k!)*Sum_{j=0..k} (-1)^(k-j)*C(k,j)*j^n (the Stirling numbers of 2nd kind). - _Paul D. Hanna_, Nov 08 2006

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Oct 31 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004192 | OEIS-A124431

a(n) = Sum_{k=0..n} 2^k*C([(n+k)/2],k)*C([(n+k+1)/2],k) where [x]=floor(x).

Conjecture: (n+1)*a(n) -2*(2*n+1)*a(n-1) +2*(n-1)*a(n-2) +2*(5-2*n)*a(n-3) +(n-3)*a(n-4) =0. - _R. J. Mathar_, Aug 09 2012 Conjecture: +(n+1)*a(n) +2*(-2*n-1)*a(n-1) +2*(n-1)*a(n-2) +2*(-2*n+5)*a(n-3) +(n-3)*a(n-4)=0. - _R. J. Mathar_, Jun 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 31 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004193 | OEIS-A124435

Number of effective multiple alignments of three equal-length sequences.

Conjecture: the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(2*r)) holds for positive integers n and r and all primes p >= 5. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lee A. Newberg_, Dec 15 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004194 | OEIS-A124453

Define Tuba numbers T(k,d,b) (0 <= d < b) by Sum_{j=0..k} binomial(k,j)*floor((k+d)/b)^(k-j)*T(j,d,b) = delta(k,0). Sequence gives T(n,0,2).

This family of sequences appeared when I was solving an open problem presented in volume 3 of the Art of Computer Programming by D. E. Knuth. This problem asked for the expected value of the search cost of a random element in a linear probing hashing table with buckets of size b. In our paper in Algorithmica 21(1): 37-71 (1998) we solve this open problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Alfredo Viola (viola(AT)fing.edu.uy), Dec 16 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004195 | OEIS-A124478

a(n) = prime(n) - floor((2/n)*Sum_{i=1..n} prime(i)).

Robert Mandl conjectured and Rosser and Schoenfeld proved that prime(n)/2 > (Sum_{i=1..n} prime(i))/n for n >= 9 (implying that a(n) > 0 for n >= 9).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 17 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004196 | OEIS-A124496

Triangle read by rows: T(n,k) is the number of set partitions of {1,2,...,n} in which the size of the last block is k, 1<=k<=n; the blocks are ordered with increasing least elements.

Conjectures: T(n,n-3) = A000096(n). T(n,n-4)= A055831(n+1). - _R. J. Mathar_, Mar 13 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Nov 14 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004197 | OEIS-A124526

Triangle, read by rows, where T(n,k) = A049020([n/2],k)*A049020([(n+1)/2],k).

A124418(n,k) = k!*T(n,k) (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 08 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004198 | OEIS-A124529

a(n) = Sum_{k=0..n} k!*A124526(n+k,k) for n>=0.

Equals diagonal sums of triangle A124418: a(n) = Sum_{k=0..n} A124418(n+k,k) for n>=0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 08 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004199 | OEIS-A124571

Expansion of limit b(n)/x^n where b(n) = b(n-1)^2 + b(n-1)*x, b(1) = x^2.

Conjecture: the limit is equal to GoldenRatio. - _Vaclav Kotesovec_, Jul 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Nov 04 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004200 | OEIS-A124588

Primes p such that q - p <= 2, where q is the next prime after p.

"Assuming certain (admittedly difficult) conjectures on the distribution of primes in arithmetic progressions, [Goldston-Pintz-Yildirim] prove the existence of infinitely many prime pairs that differ at most by 16." - Soundararajan

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 19 2006; edited May 15 2008 at the suggestion of _R. J. Mathar_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004201 | OEIS-A124598

Primes p of the form k^2+s where k > 1 and 1 <= s < (k+1)^2, such that q = k^4+s is prime and larger than p.

The terms of this sequence illustrate a special case of the conjecture from A126769.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tomas Xordan_, Mar 02 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004202 | OEIS-A124642

Antidiagonal sums of A096465.

Conjecture: G.f.: -(1/2)*z*(2*z+(1-4*z^2)^(1/2)+1)/(1-4*z^2)^(1/2)/(z^2-1). - Maksym Voznyy (voznyy(AT)mail.ru), Aug 11 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerald McGarvey_, Dec 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004203 | OEIS-A124648

Numbers n such that n^i (i=1..7) are all zeroless.

Subsequence of A253647, the analog with i <= 6 instead of 7. Conjectured to be finite. - _M. F. Hasler_, Jan 07 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Dec 22 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004204 | OEIS-A124689

Position of the first n in the decimal expansion of e^n, or -1 if the digits of n never appear.

Conjecture: -1 only occurs in the first entry of the sequence. 1000-digit precision was used in the calculation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Dec 25 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004205 | OEIS-A124690

Position of the first n in the decimal expansion of Pi^n, or -1 if the digits of n never appear.

Conjecture: -1 only occurs in the first entry of the sequence. 2000 digits of precision was used in the calculation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Dec 25 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004206 | OEIS-A124698

Number of base 5 circular n-digit numbers with adjacent digits differing by 1 or less.

Conjectures from _Colin Barker_, Jun 04 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004207 | OEIS-A124699

Number of base 6 circular n-digit numbers with adjacent digits differing by 1 or less.

Conjectures from _Colin Barker_, Jun 04 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004208 | OEIS-A124753

a(3n+k) = (k+1)*binomial(4n+k, n)/(3n+k+1), where k is n reduced mod 3.

Conjecture: +8019*n*(n-1)*(n+1)*a(n) +17496*n*(n-1)*(n-3)*a(n-1) +2592*(3*n-5)*(n-1)*(3*n-16)*a(n-2) +216*(-224*n^3+48*n^2+3926*n-6331)*a(n-3) +576*(-288*n^3+2448*n^2-6558*n+5443)*a(n-4) +768*(-288*n^3+3600*n^2-14878*n+20375)*a(n-5) -8192*(4*n-23)*(2*n-11)*(4*n-21)*a(n-6)=0. - _R. J. Mathar_, Oct 30 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 06 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004209 | OEIS-A124758

Product of the parts of the compositions in standard order.

Some conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Nov 06 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004210 | OEIS-A124791

Row sums of number triangle A124790.

Conjecture: (n+1)*a(n) +(-n+2)*a(n-1) +(-5*n+7)*a(n-2) +3*(-n+2)*a(n-3) = 0. - _R. J. Mathar_, Dec 02 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 07 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004211 | OEIS-A124828

Number of base 7 circular n-digit numbers with adjacent digits differing by 2 or less.

See A285280 for confirmation of linear recurrence and code to produce sequence; also confirms conjectured g.f. from _Colin Barker_. - _Ray Chandler_, Aug 12 2023.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004212 | OEIS-A124867

Numbers that are the sum of 3 distinct primes.

(Conjecture) Every number n > 17 is the sum of 3 distinct primes. Natural numbers that are not the sum of 3 distinct primes are listed in A124868.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Nov 11 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004213 | OEIS-A124868

Natural numbers that are not the sum of 3 distinct primes.

(Conjectured) Every number n > 17 is the sum of 3 distinct primes. a(n) is the complement of A124867(n) = {10, 12, 14, 15, 16, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, ...}, numbers that are the sum of 3 distinct primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Nov 11 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004214 | OEIS-A124869

Numerator of real part of (3*i - 1)^(-n).

Conjectures from _Colin Barker_, Jul 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Nov 11 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004215 | OEIS-A124870

Denominator of real part of (3*i - 1)^(-n).

Conjectures from _Colin Barker_, Jul 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Nov 11 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004216 | OEIS-A124999

Number of base 5 circular n-digit numbers with adjacent digits differing by 3 or less.

Conjectures from _Colin Barker_, Jun 02 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004217 | OEIS-A125054

Central terms of triangle A125053.

Conjectured g.f.: 1/(1 - x) * (1 + 2*x/(1 - 9*x/(1 - 8*x/(1 - 25*x/(1 - 24*x/(1 - 49*x/(1 - 48*x/(1 - 81*x/(1 - 80*x/(1 - ...))))))))). - _Peter Bala_, Dec 15 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004218 | OEIS-A125148

a(n) = smallest prime p = z + x, where x is the n-th odd composite number not divisible by 5 and z is a multiple of 10.

Conjecture: for every odd (prime or nonprime) number x>=1 that is not a multiple of 5 there exists a prime p such that p = z + x; where z is a multiple of 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tomas Xordan_, Jan 11 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004219 | OEIS-A125172

Triangle T(n,k) with partial column sums of the even-indexed Fibonacci numbers.

Conjecture: T(n,k) = T(n,k-1) - A121460(n+1,k). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Nov 22 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004220 | OEIS-A125189

Number of symmetric bushes with n edges. I.e., number of ordered trees with n edges, no non-root vertices of outdegree 1 and symmetrical with respect to the vertical axis passing through the root.

Conjecture: (n+1)*a(n) -3*a(n-1) +(-4*n+5)*a(n-2) +(-2*n+7)*a(n-3) +3*a(n-4) +(4*n-5)*a(n-5) +(8*n-49)*a(n-6) +3*(2*n-13)*a(n-7) +3*(n-8)*a(n-8)=0. - _R. J. Mathar_, Jun 08 2016 The conjecture follows from the differential equation 3*z^7 + z^6 + 3*z^5 + 5*z^4 + z^3 + 3*z^2 + z - 1 + (3*z^7 - z^6 + 15*z^5 + 3*z^4 + z^3 - 3*z^2 - 3*z + 1)*g(z) + (3*z^9 + 6*z^8 + 8*z^7 + 4*z^6 - 2*z^4 - 4*z^3 + z)*g'(z) = 0 satisfied by the g.f.. - _Robert Israel_, Nov 21 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004221 | OEIS-A125197

Numbers such that pre- and post-concatenating 7, 13 and 31 results in six primes.

It is conjectured that 7, 13 and 31 is the only triple with this property

猜想/问题提及段 | historical_or_current_mention_unreviewed | Anton Vrba (antonvrba(AT)yahoo.com), Jan 13 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004222 | OEIS-A125266

Number of repetitions in A007918.

Except for the first 2 terms, these numbers are always even. Conjecture: the number 2 occurs infinitely often in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jan 15 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004223 | OEIS-A125267

Number of Motzkin paths with no peaks and with level steps at height 0 having three colors except that consecutive level steps at height 0 must have different colors.

Conjecture: -(n+1)*(n-2)*a(n) +2*(n^2-n-3)*a(n-1) +(n^2-3*n+8)*a(n-2) +2*(n^2-5*n+3)*a(n-3) -(n-1)*(n-4)*a(n-4)=0. - _R. J. Mathar_, Jun 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Louis Shapiro_ and Gi-Sang Cheon, Jan 15 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004224 | OEIS-A125312

Moessner triangle based on primes.

Row sums are 2, 8, 44, 262, 1724, 13024, ... Conjecture: log row n-th sum tends to (2n-3) + some unknown fractional part. E.g., log 1724 = 7.45... while log 13024 = 9.43... Right border = A011756.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Dec 10 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004225 | OEIS-A125316

Number of base 6 circular n-digit numbers with adjacent digits differing by 3 or less.

Conjectures from _Colin Barker_, Jun 04 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004226 | OEIS-A125343

Number of base 6 circular n-digit numbers with adjacent digits differing by 4 or less.

Conjectures from _Colin Barker_, Jun 02 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004227 | OEIS-A125370

Number of base 7 circular n-digit numbers with adjacent digits differing by 5 or less.

Conjectures from _Colin Barker_, Jun 02 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004228 | OEIS-A125396

Number of base 8 circular n-digit numbers with adjacent digits differing by 6 or less.

Conjectures from _Colin Barker_, Jun 02 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004229 | OEIS-A125421

Number of base 9 circular n-digit numbers with adjacent digits differing by 7 or less.

Conjectures from _Colin Barker_, Jun 02 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004230 | OEIS-A125445

Number of base 10 circular n-digit numbers with adjacent digits differing by 8 or less.

Conjectures from _Colin Barker_, Jun 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004231 | OEIS-A125468

Number of base 11 circular n-digit numbers with adjacent digits differing by 9 or less.

Conjectures from _Colin Barker_, Jun 04 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004232 | OEIS-A125606

Row sums of A125605.

Conjectures: a(n) acts like u^n, where u = 2 zeta(3) / zeta(2). In increasing order of strength: (1) lim_{n->infinity} a(n)^(1/n) = u; (2) lim_{n->infinity} a(n+1)/a(n) = u; (3) a(n) = c u^n + o(u^n) for some c. (Conjectures 2 and 3 are very speculative.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Nov 27 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004233 | OEIS-A125607

Lesser of the smallest pair of consecutive positive reduced quadratic residues modulo p = prime(n) > 5.

The constant 0.133141413191633911131141331131441391... = sum(a(n)/10^(n-3)) is conjectured to be irrational.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nick Hobson_, Nov 30 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004234 | OEIS-A125641

Square of the (3,1)-entry of the 3 X 3 matrix M^n, where M = [1,0,0; 1,1,0; 1,i,1].

Conjecture [False!]: Draw the segments joining every lattice point on axis X with every lattice point on axis Y for 1 <= x <= n and 1 <= y <= n. The number of regions formed with these segments and axis X and Y is a(n). - _César Eliud Lozada_, Feb 14 2013 The above conjecture appears to be wrong. The number of regions formed by this construction is given in A332953, which differs from this sequence for n > 5. - _Scott R. Shannon_, Mar 04 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Nov 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004235 | OEIS-A125644

Table with T(n,k) = the number of Dyck paths whose ascent lengths, in order, are the k-th composition of n, for the standard composition order.

Conjecture: a(n) = A347205(A035327(n)) for n >= 0 (noticed by Sequence Machine). - _Mikhail Kurkov_, Dec 15 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Nov 29 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004236 | OEIS-A125696

Number of categories with n morphisms.

Conjecture: a(n) = A058129(n)*(1+o(n)). See Cruttwell presentation in Links. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_ and _Christian G. Bower_, Jan 05 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004237 | OEIS-A125731

a(n) = minimal number of steps to get from 1 to n, where a step is x -> 3x+1 if x is odd, or x -> either x/2 or 3x+1 if x is even. Set a(n) = -1 if n cannot be reached from 1.

Clearly a(3k) = -1 for all k; we conjecture that a(n) >= 0 otherwise.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Applegate_ and _N. J. A. Sloane_, Feb 02 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004238 | OEIS-A125758

Numbers congruent to 4 or 7 (mod 9).

For a given integer m, write its binary representation in reverse order, as in A125626, A125754, etc.; let a 0 mean "halving" and a 1 mean "k -> 3k+1". Then m specifies an operation on real numbers given by k -> f_m(k). Suppose the equation f_m(k) = k has a positive integer solution for some m. Then we conjecture that the values of k are precisely the terms of this sequence. In other words, we conjecture that this sequence coincides with A125757 sorted and with duplicates removed.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _David Applegate_, Feb 02 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004239 | OEIS-A125790

Rectangular table where column k equals row sums of matrix power A078121^k, read by antidiagonals.

Conjecture: g.f. for n-th row is (Sum_{i=0..n-1} x^i Sum_{j=0..i} binomial(n+1,j)*T(n,i-j)*(-1)^j)/(1-x)^(n+1) for n > 0. - _Mikhail Kurkov_, May 03 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 10 2006, corrected Dec 12 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004240 | OEIS-A125792

Column 2 of table A125790; also equals row sums of matrix power A078121^2.

Conjecture: a(n+1) = Sum_{i_1=1..3} Sum_{i_2=1..2*i_1-1} ... Sum_{i_n=1..2*i_(n-1)-1} (2*i_n - 1). For example:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 10 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004241 | OEIS-A125878

Duplicate of A066674.

It also seems likely that this is the same as A035095, but this is an open problem. Conjecture: this sequence contains only primes (this would follow if this is indeed the same as A035095).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Dec 13 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004242 | OEIS-A126026

Conjectured upper bound on area of the convex hull of any edge-to-edge connected system of regular unit hexagons (n-polyhexes).

Conjectured upper bound on area of the convex hull of any edge-to-edge connected system of regular unit hexagons (n-polyhexes). Kurz proved the polyomino equivalent of this conjecture as A122133 and abstracts: "In this article we prove a conjecture of Bezdek, Brass and Harborth concerning the maximum volume of the convex hull of any facet-to-facet connected system of n unit hypercubes in the d-dimensional Euclidean space. For d=2 we enumerate the extremal polyominoes and determine the set of possible areas of the convex hull for each n."

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Feb 27 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004243 | OEIS-A126029

The smallest positive k such that (sopfr(k)*tau(k))^n = sigma(k) where sopfr is the sum of prime factors with multiplicity (A001414).

a(3) = 14844221560107739 (conjectured) is most likely minimal but it hasn't been proved. No solutions have been found (minimal or otherwise) where the number was not squarefree.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Fred Schneider_, Dec 14 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004244 | OEIS-A126089

Expansion of e.g.f.: (1-2*x)*sqrt(1-4*x).

Conjecture D-finite with recurrence: (-n+4)*a(n) +2*(2*n-5)*(n-3)*a(n-1)=0. - _R. J. Mathar_, Jan 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 22 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004245 | OEIS-A126127

Inverse square of A061554.

G.f. as triangle (conjectured): (1-x)*(1-x+x^2)/(1-x*y+3*x^2-x^3*y+x^4). - _Robert Israel_, Jul 07 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Dec 17 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004246 | OEIS-A126156

Expansion of e.g.f. sqrt(sec(sqrt(2)*x)), showing coefficients of only the even powers of x.

Conjecture: the sequence {a(n)} reduced modulo a positive integer k is eventually periodic with the period dividing phi(k), where phi(n) = A000010(n). An example is given below. - _Peter Bala_, Mar 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 20 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004247 | OEIS-A126180

Number of branches in all hex trees with n edges (n>=1).

Conjecture: -2*(n+2)*(239*n-1252)*a(n) +21*(220*n^2-997*n-578)*a(n-1) +(-12902*n^2+76676*n-77157)*a(n-2) +15*(n-3)*(584*n-2237)*a(n-3)=0. - _R. J. Mathar_, Jun 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 19 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004248 | OEIS-A126195

Conjectured values for maximal number of solid spheres of radius 1 that can be rolled all in touch with and on the outside surface of a sphere of radius n.

Conjectured values for maximal number of solid spheres of radius 1 that can be rolled all in touch with and on the outside surface of a sphere of radius n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Mar 07 2007, corrected Apr 03 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004249 | OEIS-A126198

Triangle read by rows: T(n,k) (1 <= k <= n) = number of compositions of n into parts of size <= k.

Conjecture: Sum_{k=1..n} T(n,k) = A039671(n), n>0. - _L. Edson Jeffery_, Nov 29 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 09 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004250 | OEIS-A126235

Minimum length of a codeword in Huffman encoding of n symbols, where the k-th symbol has frequency k.

Conjecture: a(n) = A099396(n+1) = floor(log_2(2(n+1)/3)). Equivalently, a(n) = a(n-1) + 1 if n has the form 3*2^k-1, a(n) = a(n-1) otherwise. This is true at least for n up to 1000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, Dec 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004251 | OEIS-A126236

Maximum length of a codeword in Huffman encoding of n symbols, where the k-th symbol has frequency k.

Conjecture: a(n) = floor(log_2(n)) + floor(log_2(2n/3)). Equivalently, a(n) = a(n-1) + 1 if n has the form 2^k or 3*2^k, a(n) = a(n-1) otherwise. This is true at least for n up to 1000. The conjecture is true (see Szewczyk link). - _Kamila Szewczyk_, Jul 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, Dec 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004252 | OEIS-A126237

Length of row n in table A126014.

I conjecture that there are no gaps in the set of codeword lengths; that is, every integer that's between the minimum and maximum codeword lengths occurs as a codeword length. If so, then a(n) = A126236(n) - A126235(n). If, in addition, the conjectured formulas for the min and max lengths are correct, then a(n) = floor(log_2(n)) unless n has the form 3*2^k-1, in which case a(n) = floor(log_2(n)) - 1. This is true at least for n up to 1000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dean Hickerson_, Dec 21 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004253 | OEIS-A126241

Dropping time of n in the 3n+1 problem (or the Collatz problem). Let T(n):=n/2 if n is even, (3n+1)/2 otherwise (A014682). Let a(n) be the smallest integer k such that T^(k)(n)<n, where T^(k) is the k-th iterate, or infinity otherwise; a(n) is called the dropping time of n.

Terras defines a coefficient stopping time (definition 1.5) tau(n) = d which is the smallest d for which 3^u/2^d < 1 where u is the number of tripling steps among the first d steps starting from n. Clearly tau(n) <= a(n), and Terras conjectures (conjecture 2.9) that tau(n) = a(n) for n>=2. - _Olivier Rozier_, May 13 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | Christof Menzel (christof.menzel(AT)hs-niederrhein.de), Mar 08 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004254 | OEIS-A126269

Numbers k such that hcl(k,k) < hcl(k,k-1) where hcl(k,i) is the Huffman code length; see comments.

Conjecture: a(2k) = A033484(k-1) and a(2k-1) = A068156(k-1), k >= 2. Conjectures from _Colin Barker_, Aug 06 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Serhat Sevki Dincer (mesti_mudam(AT)yahoo.com), Dec 22 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004255 | OEIS-A126322

Number of hex trees with n edges and no branches of length 1.

Conjecture: (n+4)*(25*n^2+230*n+1137)*a(n) +3*(-50*n^3-585*n^2-3169*n-4248) *a(n-1) +9*(25*n^3+255*n^2+932*n-1764) *a(n-2) +29484*a(n-3) -36*(n-4) *(25*n^2+280*n+1392) *a(n-4)=0. - _R. J. Mathar_, Jun 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 25 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004256 | OEIS-A126323

Number of branches of length 1 in all hex trees with n edges.

Conjecture: -(n+2)*(1176*n^2+350135*n-2095015)*a(n) +3*(-392*n^3+1110577*n^2-6230546*n-1231580)*a(n-1) +(43512*n^3-9103273*n^2+68264245*n-103131090) *a(n-2) -15*(n-4)*(2744*n^2-398547*n+1917624)*a(n-3)=0. - _R. J. Mathar_, Jun 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 25 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004257 | OEIS-A126324

a(2n) = Cat(n), a(2n+1) = 3*Cat(n), where Cat(n) = binomial(2n,n)/(n+1) are the Catalan numbers (A000108).

Conjecture: (5*n+17)*(n+2)*a(n) - 36*a(n-1) - 4*(5*n+22)*(n-2)*a(n-2) = 0. - _R. J. Mathar_, Jun 17 2016 Conjecture confirmed, because the g.f. satisfies the d.e. (-36*z+34)*g(z)+(-148*z^3+32*z)*g'(z)+(-20*z^4+5*z^2)*g''(z)-162*z-34 = 0. - _Robert Israel_, Dec 29 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 25 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004258 | OEIS-A126326

a(1) = 1; for n>1, a(n) = smallest number which is not a sum or product or power of any subset of the numbers a(1) to a(n-1).

Conjectures from _Colin Barker_, Jun 21 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Mar 11 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004259 | OEIS-A126360

Number of base 6 n-digit numbers with adjacent digits differing by one or less.

Conjecture: a(n) = 4*a(n-1) - 3*a(n-2) - a(n-3) for n > 3. Barker's conjecture follows from the fact that (A^3 - 4*A^2 + 3*A + 1) e = 0. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 26 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004260 | OEIS-A126362

Number of base 8 n-digit numbers with adjacent digits differing by one or less.

Conjecture: a(n) = 5*a(n-1) - 6*a(n-2) - a(n-3) + 2*a(n-4) for n > 4. a(n) = e^T A^(n-1) e for n>=1, where A is the 8 X 8 matrix with 1 on the main diagonal and first super- and subdiagonals, 0 elsewhere, and e the column vector (1,1,1,1,1,1,1,1). Barker's conjecture follows from the fact that (A^4 - 5*A^3 + 6*A^2 + A - 2*I)*e = 0. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 26 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004261 | OEIS-A126364

Number of base 10 n-digit numbers with adjacent digits differing by one or less.

Conjecture: a(n) = 3*a(n-1) - A025565(n), n >= 2. - _Daniel Forgues_, Aug 23 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 26 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004262 | OEIS-A126395

Number of base 8 n-digit numbers with adjacent digits differing by two or less.

Conjectures from _Colin Barker_, Jun 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004263 | OEIS-A126396

Number of base 9 n-digit numbers with adjacent digits differing by two or less.

Conjectures from _Colin Barker_, Jun 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004264 | OEIS-A126397

Number of base 10 n-digit numbers with adjacent digits differing by two or less.

Conjectures from _Colin Barker_, May 31 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004265 | OEIS-A126474

Number of arrays in [1..6]^n with adjacent elements differing by three or less.

Conjectures from _Colin Barker_, Jan 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 27 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004266 | OEIS-A126475

Number of base 7 n-digit numbers with adjacent digits differing by three or less.

Conjectures from _Colin Barker_, Jun 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 27 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004267 | OEIS-A126476

Number of base 8 n-digit numbers with adjacent digits differing by three or less.

Conjectures from _Colin Barker_, Jun 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 27 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004268 | OEIS-A126477

Number of base 9 n-digit numbers with adjacent digits differing by three or less.

Conjectures from _Colin Barker_, Jun 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 27 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004269 | OEIS-A126478

Number of base 10 n-digit numbers with adjacent digits differing by three or less.

Conjectures from _Colin Barker_, May 31 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 27 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004270 | OEIS-A126501

Number of n-tuples of numbers [0..5] (leading zeros allowed) in which adjacent digits differ by 4 or less.

Conjecture: a(n) = (2^(-1-n)*((5-sqrt(41))^n*(-7+sqrt(41)) + (5+sqrt(41))^n*(7+sqrt(41)))) / sqrt(41). - _Colin Barker_, Jan 20 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004271 | OEIS-A126502

Number of base 7 n-digit numbers with adjacent digits differing by four or less.

Conjectures from _Colin Barker_, Jun 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004272 | OEIS-A126503

Number of base 8 n-digit numbers with adjacent digits differing by four or less.

Conjectures from _Colin Barker_, Jun 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004273 | OEIS-A126504

Number of base 9 n-digit numbers with adjacent digits differing by four or less.

Conjectures from _Colin Barker_, Jun 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004274 | OEIS-A126505

Number of base 10 n-digit numbers with adjacent digits differing by four or less.

Conjectures from _Colin Barker_, May 31 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004275 | OEIS-A126529

Number of base 8 n-digit numbers with adjacent digits differing by five or less.

Conjectures from _Colin Barker_, Jun 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004276 | OEIS-A126530

Number of base 9 n-digit numbers with adjacent digits differing by five or less.

Conjectures from _Colin Barker_, Jun 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004277 | OEIS-A126531

Number of base 10 n-digit numbers with adjacent digits differing by five or less.

Conjectures from _Colin Barker_, May 31 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004278 | OEIS-A126568

Binomial transform of A026641.

Conjecture: 4*n*a(n) +2*(2*n-7)*a(n-1) +(-163*n+267)*a(n-2) +10*(23*n-58)*a(n-3) +75*(-n+3)*a(n-4) = 0. - _R. J. Mathar_, Jun 30 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Mar 13 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004279 | OEIS-A126674

a(n) = n!*Sum_{j=0..n-1} 2^j/(j+1).

Conjecture: a(n) + (-3*n+2)*a(n-1) + 2*(n-1)^2*a(n-2) = 0. - _R. J. Mathar_, May 23 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and Carlo Wood (carlo(AT)alinoe.com), Feb 13 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004280 | OEIS-A126699

Lesser of twin simili-primes of order 2.

q > p are "twin simili-primes of order 2" if (p, q) are in A123929 and satisfy q - p = 3. Conjecture : this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 11 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004281 | OEIS-A126717

Least odd k such that k*2^n-1 is prime.

Conjecture: a(n) = O(n log n). - _Thomas Ordowski_, Oct 15 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernardo Boncompagni_, Feb 13 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004282 | OEIS-A126726

Number of squares (of nonnegative integers) that require n binary (base-2) digits.

Conjecture: a(n) = A190568(n)-(-1)^n for n>0. - _R. J. Mathar_, May 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | Andrew G. West (westa(AT)wlu.edu), Mar 13 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004283 | OEIS-A126757

Number of n-node connected graphs with no cycles of length less than 5.

This is the inverse Euler transform of A006787. - Conjectured by _Vladeta Jovovic_, Jun 16 2008, proved by _Max Alekseyev_ and _Brendan McKay_, Jun 17 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 18 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004284 | OEIS-A126762

a(n) is the least k > n such that the remainder when n^k is divided by k is n.

a(n) is the smallest number k > n such that n^k == n (mod k). Conjecture: a(n) is the smallest number k > n such that n^(k-1) == 1 (mod k). Thus a(n) is coprime to n. - _Thomas Ordowski_, Aug 03 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Feb 17 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004285 | OEIS-A126764

Number of L-convex polyominoes with n cells, that is, convex polyominoes where any two cells can be connected by a path internal to the polyomino and which has at most 1 change of direction (i.e., one of the four orientations of the letter L).

Conjecture: a(n) ~ c * exp(Pi*sqrt(13*n/6)) / n^(3/2), where c = 13*sqrt(2) / 768. - _Anthony Guttmann_ and _Vaclav Kotesovec_, Jun 09 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, based on email from Simone Rinaldi (rinaldi(AT)unisi.it), Feb 23 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004286 | OEIS-A126769

Primes p of the form k^4 + s where k > 1, s >= 1 and k^2 + s is also prime.

Conjecture: Every prime q > 3 can be written in a nontrivial way as the sum of two or more squares, q = Sum_{i} (k_i)^2, such that the sum of the fourth powers of the squared numbers is again prime, p = Sum_{i} (k_i)^4. (Tomas Xordan) This sequence illustrates an easy case of the conjecture: For primes q arising in the sequence there exists an integer k > 1, a positive integer s and a prime p such that k^2 < q, s = q - k^2, p = k^4 + s and p > q.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tomas Xordan_, Feb 16 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004287 | OEIS-A126892

a(n) = row of Wythoff's array T(n,j) containing the sequence of values T(n,j-1) + T(n,j+1).

Conjecture: If the Zeckendorf representation of n ends in 1, then a(n) = 15 + H(n-H(n))*29 + (n-H(n) - H(n-H(n)))*18, where H(n) is Hofstadter's G sequence A005206. Otherwise, a(n) = 1 + H(H(n))*7 + (H(n) - H(H(n)))*4 unless the Zeckendorf representation of n has one of the 0-endings listed in the first comment line, in which case a(n) = a(n+1) - 11.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kenneth J Ramsey_, Jan 13 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004288 | OEIS-A126932

Binomial transform of A127358.

Conjecture: +2*n*a(n) -11*n*a(n-1) +4*(2*n+3)*a(n-2) +21*(n-2)*a(n-3)=0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Mar 17 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004289 | OEIS-A127054

Rectangular table, read by antidiagonals, defined by the following rule: start with all 1's in row zero; from then on, row n+1 equals the partial sums of row n excluding terms in columns k = m*(m+1)/2 (m>=1).

Conjecture: A(n,k) = A(n-1,k) + A396981(n,k) with A(0,k) = 1. - _Mikhail Kurkov_, Jun 12 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 04 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004290 | OEIS-A127068

Let d(m, 0) = 1, d(m, 1) = m, and d(m, k) = (m - k + 1)*d(m+1, k-1) - (k-1)*(m+1) d(m+2, k-2). Sequence gives d(3,n).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | Vincent v.d. Noort, Mar 21 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004291 | OEIS-A127069

Number of lines in a Pauli graph of order n.

Conjectures from _Colin Barker_, May 28 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Mar 21 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004292 | OEIS-A127081

One-sided kissing number for spheres in n-dimensional Euclidean space.

a(8) = 183. Musin's conjectures: a(5) = 32, a(24) = 144855.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Mar 21 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004293 | OEIS-A127154

Number of Dyck paths of semilength n and having no UDUD's starting at level 0; here U=(1,1), D=(1,-1).

Conjecture: (n+1)*a(n) +3*a(n-1) +(-11*n+16)*a(n-2) +(-17*n+43)*a(n-3) +(-11*n+28)*a(n-4) +2*(-2*n+7)*a(n-5)=0. - _R. J. Mathar_, Jun 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Feb 27 2007, Dec 13 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004294 | OEIS-A127202

a(1)=1, a(2)=2; a(n) = the smallest positive integer not occurring earlier in the sequence such that gcd(a(n), a(n-1)) does not equal gcd(a(n-1), a(n-2)).

Conjecture: The two lines in the graph are (apart from small local deviations) defined by the same equations as the two lines in the graph of A283312. - _N. J. A. Sloane_, Mar 12 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Jan 08 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004295 | OEIS-A127208

Union of all n-step Lucas sequences, that is, all sequences s(1-n) = s(2-n) = ... = s(-1) = -1, s(0) = n and for k > 0, s(k) = s(k-1) + ... + s(k-n).

Noe and Post conjectured that the only positive terms that are common to any two distinct n-step Lucas sequences are the Mersenne numbers (A001348) that begin each sequence and 7 and 11 (in 2- and 3-step) and 5071 (in 3- and 4-step). The intersection of this sequence with the union of all the n-step Fibonacci sequences (A124168) appears to consist of 4, 21, 29, the Mersenne numbers 2^n-1 for all n and the infinite set of Eulerian numbers in A127232.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jan 09 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004296 | OEIS-A127317

Numbers n such that (256^n + 1)/257 is prime.

We can even conjecture that ((Fj-1)^p + 1)/Fj is always squarefree for any odd prime p and for any Fermat number Fj with j >= 0. Note that this is not true if p is not restricted to be a prime, even if p and Fj are coprime, as shown by the following counterexample relative to the case j = 1, f1 = 5: 4^91 + 1 == 0 mod 1093^2. Remark that any such counterexample has to be a Wieferich prime (A001220), but not every Wieferich prime gives a counterexample, as shown by the second known Wieferich prime (3511), which cannot match here because it belongs to A072936.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Mar 29 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004297 | OEIS-A127361

a(n) = Sum_{k=0..n} binomial(n, floor(k/2))*(-2)^(n-k).

Conjecture: 2*n*a(n) + (5*n-4)*a(n-1) - 2*(4*n-3)*a(n-2) - 20*(n-2)*a(n-3) = 0. - _R. J. Mathar_, Nov 30 2012 [Proved in Easwar (2026). - _Rohun Easwar_, Aug 31 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 11 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004298 | OEIS-A127363

a(n) = Sum_{k=0..n} C(n,floor(k/2))*(-4)^(n-k).

Conjecture: 4*n*a(n) +(17*n-8)*a(n-1) +2*(-8*n-1)*a(n-2) +68*(-n+2)*a(n-3)=0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 11 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004299 | OEIS-A127628

G.f. 1/(1-6*x*c(x)) where c(x) is the g.f. of A000108.

Conjecture: 5*n*a(n) + 2*(15-28*n)*a(n-1) + 72*(2*n-3)*a(n-2) = 0. - _R. J. Mathar_, Nov 14 2011 G.f.: (2+3*sqrt(1-4*x))/(5-36*x). Mathar's conjecture verified using the differential equation

(144*x^2-56*x+5)*y'+(72*x-26)*y = 4 satisfied by the g.f. - _Robert Israel_, Aug 28 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 20 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004300 | OEIS-A127632

Expansion of c(x*c(x)), where c(x) is the g.f. for A000108.

Conjecture: 3*n*(n-1)*(4*n-9)*(n+1)*a(n) - 8*n*(n-1)*(20*n^2-67*n+48)*a(n-1) + 8*(4*n-5)*(n-1)*(22*n^2-94*n+99)*a(n-2) - 8*(4*n-11)*(4*n-5)*(4*n-9)*(2*n-5)*a(n-3) = 0. - _R. J. Mathar_, May 04 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 20 2007, Jan 25 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004301 | OEIS-A127710

Inverse of the triangle A(n,k)=if(k<=n,if(n<=2k,1/(n+1),0),0).

It is conjectured that all elements of this triangle are integers. It appears that the row sums are the ruler function A001511.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 24 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004302 | OEIS-A127711

Inverse of the triangle A(n,k) = 1/F(n+1) if k <= n <= 2k, 0 otherwise.

It is conjectured that all elements of this triangle are integers. Row sums are A127712.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 24 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004303 | OEIS-A127728

Sum of squared coefficients of q in the q-factorials.

Conjecture: a(n) ~ 6 * sqrt(Pi) * n^(2*n - 1/2) / exp(2*n). - _Vaclav Kotesovec_, Oct 22 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 25 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004304 | OEIS-A127749

Inverse of number triangle A(n,k) = 1/(2n+1) if k <= n <= 2k, 0 otherwise.

Conjectures: row sums modulo 2 are the Fredholm-Rueppel sequence A036987; row sums of triangle modulo 2 are A111982. Row sums are A127750. The first conjecture is equivalent to the row sums conjecture in A111967. - _R. J. Mathar_, Apr 21 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004305 | OEIS-A127750

Row sums of inverse of number triangle A(n,k) = 1/(2n+1) if k <= n <= 2k, 0 otherwise.

Conjecture: a(n) mod 2 gives Fredholm-Rueppel sequence A036987. The conjecture is true at least up to n=2048. - _Antti Karttunen_, Sep 29 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004306 | OEIS-A127751

Inverse of number triangle A(n,k)=if(k<=n,if(n<=2k,1/(3n+1),0),0).

Row sums A127752 mod 2 are conjectured to be first Feigenbaum symbolic sequence A035263.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004307 | OEIS-A127752

Row sums of inverse of number triangle A(n,k) = 1/(3n+1) if k <= n <= 2k, 0 otherwise.

a(n) mod 2 is first Feigenbaum symbolic sequence A035263 (conjecture). The conjecture is true at least up to 2048 first terms. (But please note the different indexing, here 0-based.) - _Antti Karttunen_, Sep 29 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004308 | OEIS-A127755

Inverse of number triangle A(n,k) = 1/n! if k <= n <= 2k, 0 otherwise.

It is conjectured that this is an integer triangle. Row sums are A127756.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004309 | OEIS-A127767

Inverse of number triangle A(n,k) = 1/C(n) if k <= n <= 2k, 0 otherwise, where C(n) = A000108(n).

It is conjectured that all elements of the triangle are integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004310 | OEIS-A127771

Inverse of number triangle A(n,k) = 1/Euler_phi(n+1) if k <= n <= 2k, 0 otherwise.

It is conjectured that all elements of the triangle are integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004311 | OEIS-A127789

Record indices of 2^h(n)/(3^t(n)*n), where h and t are the number of halving resp. tripling steps in the '3x + 1' problem.

Eric Roosendaal calls 2^j / (n*3^k) the residue of n and conjectures that 993 yields the highest residue. - _T. D. Noe_, Apr 08 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Boddington_, Apr 04 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004312 | OEIS-A127793

Inverse of number triangle t(n,k) = 1/floor((n+2)/2) if k <= n <= 2*k, 0 otherwise.

It is conjectured that the triangle is an integer triangle. The triangle and its inverse both appear to have row sums equal to the all 1's sequence. The triangle is equivalent to the lower semi-matrix T = e_{1,1} + Sum_{i>=2} Sum_{p>=0} (e_{2^p i, i} ceiling(i/2) - e_{2^p (i+1), i} ceiling(i/2)) , where e_{i,j} is the matrix unit. The conjecture above is true, deduced from the formula of the matrix. - _FUNG Cheok Yin_, Sep 12 2022 All the conjectures above are true (see Fried link). - _Sela Fried_, Nov 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 29 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004313 | OEIS-A127800

Inverse of number triangle A(n,k) = 1/(2 - 0^n), if k <= n <= 2k, 0 otherwise.

Row sums are A069517 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 29 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004314 | OEIS-A127823

Weighted Catalan numbers: number of combinatorial types of plane Morse links of order n.

a(n) has the same number of factors of 2 as does A000108(n) (Catalan numbers). "Given a sequence of integers b = (b_0,b_1,b_2,...) one gives a Dyck path P of length 2n the weight wt(P) = b_{h_1} b_{h_2} ... b_{h_n}, where h_i is the height of the i-th ascent of P. The corresponding weighted Catalan number is C_n^b = sum_P wt(P), where the sum is over all Dyck paths of length 2n. So in particular, the ordinary Catalan numbers C_n correspond to b_i = 1 for all i >= 0. Let xi(n) stand for the base two exponent of n, i.e., the largest power of 2 dividing n. We give a condition on b which implies that xi(C_n^b) = xi(C_n). In the special case b_i=(2i+1)^2, this settles a conjecture of Postnikov about the number of plane Morse links." - Alexander Postnikov, Bruce Sagan.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 30 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004315 | OEIS-A127854

Largest number k such that k^2 divides A007781(6n+1).

Conjecture: a(n) = 12n^2 + 6n + 1. Conjecture: a(n) = 3*a(n-1) - 3*a(n-2) + a(n-3); g.f.: x*(19 + 4*x + x^2)/(1-x)^3. - _Colin Barker_, Mar 16 2012 These conjectures are false. For n=74, 12*n^2 + 6*n + 1 = 66157 but A007781(6*74+1) is divisible by 5491031^2. - _Robert Israel_, Nov 19 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Apr 05 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004316 | OEIS-A127905

Construct triangle in which n-th row is obtained by expanding (1+x+x^3)^n and take the next-to-central column.

Conjecture: 2*(2*n+1)*(n-1)^2*a(n) -2*n*(6*n^2-12*n+5)*a(n-1) +6*n*(n-1)*(2*n-3)*a(n-2) -31*n*(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Feb 23 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 05 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004317 | OEIS-A127923

Difference between squares of legs of primitive Pythagorean triangles, sorted (with multiplicity).

The sequence is conjectural (and may miss entries) because it is generated from a finite list of primitive Pythagorean triangles. The associated lengths in a^2+b^2=c^2 are (a,b)=(3,4), (21,20), (5,12), (15,8), (119,120), (7,24), (55,48), (65,72), (35,12), (45,28), (697,696), (9,40), (33,56), (105,88), (11,60), (63,16), (297,304), (77,36), (91,60), (39,80), (403,396), (133,156), (13,84), (207,224), (4059, 4060), (99,20), (171,140), (85,132), (117,44), (275,252), (15,112), (153,104), (51,140), (555,572), (95,168), (143,24), (17,144), (253, 204), (225,272), (165,52), (1755,1748), (429,460),... with gcd(a, b)=1 and |a^2-b^2| in the sequence. - _R. J. Mathar_, Sep 24 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Feb 06 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004318 | OEIS-A127925

Primes p such that 2p < prime(k-i) + prime(k+i) for i=1..k-1, where p=prime(k).

Conjecture: these are the numbers k > 1 for which 2*pi(k) = pi(k-m) + pi(k+m) has no solution 0 < m < k. The conjecture is true if and only if for every composite number k this equation has such a solution m. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Feb 06 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004319 | OEIS-A127929

a(n) = A127928(n) mod 18.

Aside from "3", all terms of A127928 must be 1 or 7 mod 18 (see A127928 for mod rules); but not all primes mod 1 or 7 are pure hailstone numbers. For example, the prime 61 == 7 mod 18 but 61 is impure. Conjecture: for large n, the numbers of 1 and 7 mod 18 terms are approximately equal.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Feb 07 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004320 | OEIS-A127974

Numerators in expansion of (1-x)^(-2/3).

Conjecture: a(n-1) = the numerator of (1/(2*sqrt(3)*Pi)) * Integral_{x >= 0} 1/(1 + x^3)^n. - _Peter Bala_, Jun 11 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 09 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004321 | OEIS-A128057

Expansion of (1+x)/sqrt(1+4x^2).

Conjecture: n*(5*n-9)*a(n) +4*a(n-1) +4*(5*n-4)*(n-2)*a(n-2)=0. - _R. J. Mathar_, Dec 02 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 13 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004322 | OEIS-A128100

Triangle read by rows: T(n,k) is the number of ways to tile a 2 X n rectangle with k pieces of 2 X 2 tiles and n-2k pieces of 1 X 2 tiles (0 <= k <= floor(n/2)).

Row n gives the coefficients of the polynomial p(n,x) defined as the numerator of the rational function given by f(n,x) = 1 + (x + 1)/f(n-1,x), where f(x,0) = 1. Conjecture: for n > 2, p(n,x) is irreducible if and only if n is a (prime - 2). - _Clark Kimberling_, Oct 22 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Feb 18 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004323 | OEIS-A128103

Number of permutations of [n] with an even number of rises.

Conjecture: a(n) = Sum_{k = 0..n} Sum_{j = 0..k} (-1)^(n+j)*binomial(n,k-j)*j^n. - _Peter Bala_, Jan 22 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, May 09 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004324 | OEIS-A128127

The number of regular pentagons found by constructing n equally-spaced points on each side of the pentagon and drawing lines parallel to the pentagon side.

a(n) = (10*n^2 - 4*n + 5 -(-1)^n)/4 (conjectural).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Noah Priluck (npriluck(AT)gmail.com), May 02 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004325 | OEIS-A128135

Row sums of A128134.

Conjecture: a(n)/a(n-1) tends to sqrt(5). (E.g., a(10)/a(9) = 2.235294....) The conjecture is false. The fraction a(n)/a(n-1) tends to 2 as n grows. - Philipp Zumstein (zuphilip(AT)inf.ethz.ch), Oct 05 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Feb 16 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004326 | OEIS-A128153

The number of regular pentagons found by constructing n equally-spaced points on each side of the pentagon and drawing lines parallel to the pentagon sides, as well as lines connecting vertices.

Conjecture: a(n) = (10*n^2 + 16*n + 9 -(-1)^n)/4 for n > 0. a(n) = 2*a(n-1) - 2*a(n-3) + a(n-4) for n > 4 (conjectured). G.f.: (-x^4 + x^3 - 2*x^2 - 7*x - 1)/((x - 1)^3*(x + 1)) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Noah Priluck (npriluck(AT)gmail.com), May 02 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004327 | OEIS-A128280

a(n) is the least number not occurring earlier such that a(n)+a(n-1) is prime, a(0) = 0.

Without altering the definition nor the existing values, one can as well start with a(0) = 0 and get (conjecturally) a permutation of the nonnegative integers. This sequence is in some sense the "arithmetic" analog of the "digital" variant A231433: Here we add subsequent terms, there the digits are concatenated. - _M. F. Hasler_, Nov 09 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, May 03 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004328 | OEIS-A128318

G.f.: A(x) = 1+x*(1+2x*(1+3x*(...(1+n*x*(...)^2)^2...)^2)^2)^2.

Conjecture: a(n) ~ n! * (8/3)^n / sqrt(n). - _Vaclav Kotesovec_, Mar 19 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 07 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004329 | OEIS-A128378

Partial sums of A128379.

Conjecture: Given the infinite set of sequences generated from using the partial sum operator on A000594, (i.e. A000012^k * A000594, k in succession = 1, 2, 3, ...), k = 23 and k = 24 are the only two sequences in the set with zeros. In A128379, k = 23: (1, -1, -24, 0, 276, 300, -1748, -4300,

...).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Feb 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004330 | OEIS-A128379

A000012^23 * A000594.

Conjecture: given A000012^k * A000594, k=23 and 24 are the only k's generating sequences with zeros. k = 24 in A128378: (1, 0, -24, -24, 252, 552, -1196, -5496, ...).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Feb 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004331 | OEIS-A128380

A097806^24 * A000594.

Conjecture: Given the infinite set of sequences generated from the pairwise operation on A000594 (A097806^k * A000594), k = 24, (A128380) is the only sequence in the set with a zero. The sequence generated from k=23 = (1, -1, -47, 23, 1081, 47, -15939, ...). Analogous conjecture with the partial sum operator: (Cf. A128378, A128379); in which zeros are conjectured to occur only with k=23 and k=24. A128380 mod 24 = 1, 0, 0, 0, 0, 0, -4, 0, 0, 16, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Feb 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004332 | OEIS-A128381

A007318^24 * A000594.

Conjecture: given the bto performed any k times on A000594 (k = 1, 2, 3, ...); k = 6 and k = 24 are the only members of the set with zeros. k = 6 generates (1, -18, 0, 688, 4494, 5508, ...).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Feb 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004333 | OEIS-A128382

Inverse Moebius transform operation performed 24 times on A000594: A051731^24 * A000594.

Conjecture: given the inverse Moebius transform operation performed any k times (k=1,2,3,...); k=24 is the only such sequence with zeros. A weaker conjecture: "zero" occurs an infinite number of times in A128382.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Feb 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004334 | OEIS-A128391

A054523^24 * A000594.

Conjecture: Given A054523^k, k = any positive integer, "zero" appears only in the sequence A018391 (k=24).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Feb 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004335 | OEIS-A128392

A126988^12 * A000594.

Conjecture: Given A126988^k, k any positive integer, A128392 is the only sequence in the infinite set with zeros.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Feb 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004336 | OEIS-A128424

a(n) = floor(sqrt(a(n-1)^2 + a(n-2)^2 + a(n-1)*a(n-2))), a(1)=1, a(2)=3.

Conjectures from _Colin Barker_, Nov 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, May 04 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004337 | OEIS-A128427

Last point where sum of n consecutive n-th powers does not exceed the next n-th power.

Denoting by x(n) the largest real root of (x-n)^n+...+(x-1)^n=x^N (so that by definition a(n)=floor(x(n))), it is conjectured ("Cyprian's Last Theorem") but not fully proved that x(n) is never an integer for n>3. A (very rapidly very good) asymptotic approximation to x(n) is 1.5 + n / ln2 + O(1/n), but this needs proof. This yields floor(1.5 + n / ln2) as an approximation to a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Martin Kochanski (mjk(AT)cardbox.com), May 10 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004338 | OEIS-A128440

Array T(n,k) = floor(k*r^n) where r = golden ratio = (1 + sqrt(5))/2, read by descending antidiagonals.

Conjecture: rows 2n-1 and 2n are disjoint for every positive integer n. - _Clark Kimberling_, Nov 11 2022 Stronger conjecture: for any positive integer n, if the numbers in rows 2n-1 and 2n are jointly arranged in increasing order, and each number is replaced by its position in the ordering, then the resulting two rows are identical to the first two rows. - _Clark Kimberling_, Nov 13 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 03 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004339 | OEIS-A128508

Number of partitions p of n such that max(p) - min(p) = 3.

Conjecture. a(1)=0 and, for n>1, a(n+1)=a(n)+d(n), where d(n) is defined as follows: d=0,0,0,1,0 for n=1,...,5 and, for n>5, d(n)=d(n-2)+1 if n=6k or n=6k+4, d(n)=d(n-2) if n=6k+1 or n=6k+3, d(n)=d(n-2)+2Floor[n/6] if n=6k+2 and d(n)=d(n-5) if n=6k+5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, May 07 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004340 | OEIS-A128605

Smallest number m having exactly n divisors d with sqrt(m/2) <= d < sqrt(2*m).

Conjecture: a(n) = k where p(k) and p(k-1) are the first pair of Dyck paths for the symmetric representation of sigma(k) and sigma(k-1), as described in A237593, having a gap of exactly n units on the diagonal, i.e., it is the sequence of record gaps in sequence A240542; tested through 2000000

with a variant of function A279286. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Mar 14 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004341 | OEIS-A128607

Perfect (or pure) powers pp such that sigma(pp) is also a perfect (pure) power.

Denote by egcd(n) the gcd of all the powers in the prime factorization of n. In our context, a square has egcd=2, a cube has egcd=3 and so on. The only elements n in the sequence for which egcd(n)>2 are 81 and 343. Are there any others? Conjecture I: egcd(A128607(n))=2 for all n>2. Let A128608(n)=sigma(A128607(n)). Note that A128607(11)=1857437604=(2^2)*(3^2)*(11^2)*(653^2) has A128608(11)=5168743489=(7^3)*(13^3)*(19^3). Any other cubes or higher egcd's in A128608? Conjecture II: egcd(A128608(n))=2 for all n ne 11.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Walter Kehowski_, Mar 20 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004342 | OEIS-A128608

a(n)=sigma(A128607(n)), where A128607(n) is the sequence of perfect (or pure) powers such that a(n) is a perfect power.

Denote by egcd(n) the gcd of all the powers in the prime factorization of n. In our context, a square has egcd=2, a cube has egcd=3 and so on. The only elements n in the sequence for which egcd(n)>2 are 81 and 343. Are there any others? Conjecture I: egcd(A128607(n))=2 for all n>2. Let a(n)=sigma(A128607(n)). Note that A128607(11)=1857437604=(2^2)*(3^2)*(11^2)*(653^2) has a(11)=5168743489=(7^3)*(13^3)*(19^3). Any other cubes or higher egcd's in this sequence? Conjecture II: egcd(a(n))=2 for all n ne 11.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Walter Kehowski_, Mar 20 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004343 | OEIS-A128666

Least generalized Wilson prime p such that p^2 divides (n-1)!(p-n)! - (-1)^n; or 0 if no such prime exists.

Conjecture: a(n)>0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Mar 25 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004344 | OEIS-A128699

Highly abundant numbers that are not superabundant, i.e., the complement of A004394 w.r.t. A002093.

In 1944, Alaoglu and Erdős conjectured that this sequence was infinite and this was proved to be true by Nicolas in 1969.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ant King_, Mar 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004345 | OEIS-A128723

Number of skew Dyck paths of semilength n having no peaks at level 1.

Conjecture: +3*(n+1)*a(n) +(-17*n+10)*a(n-1) +9*(n-3)*a(n-2) +5*(n-2)*a(n-3)=0. - _R. J. Mathar_, Jun 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Mar 31 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004346 | OEIS-A128750

Number of skew Dyck paths of semilength n having no ascents of length 1.

Conjecture: (n+1)*a(n) +(-4*n+3)*a(n-1) +(-2*n-1)*a(n-2) +(4*n-11)*a(n-3) +(n-4)*a(n-4)=0. - _R. J. Mathar_, Nov 15 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Mar 31 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004347 | OEIS-A128760

Number of ways to write n as the absolute difference of a power of 2 and a power of 3.

Conjecture: there exists c>=23 such that a(n)<2 for n>c.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Mar 25 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004348 | OEIS-A128818

Examples of integer-encoded mathematical structures in Max Tegmark's "The Mathematical Universe".

Abstract from "The Mathematical Universe" (see link below): "I explore physics implications of the External Reality Hypothesis (ERH) that there exists an external physical reality completely independent of us humans. I argue that with a sufficiently broad definition of mathematics, it implies the Mathematical Universe Hypothesis (MUH) that our physical world is an abstract mathematical structure. I discuss various implications of the ERH and MUH, ranging from standard physics topics like symmetries, irreducible representations, units, free parameters and initial conditions to broader issues like consciousness, parallel universes and Godel incompleteness. I hypothesize that only computable and decidable (in Godel's sense) structures exist, which alleviates the cosmological measure problem and help explain why our physical laws appear so simple. I also comment on the intimate relation between mathematical structures, computations, simulations and physical systems."

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Apr 10 2007, Oct 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004349 | OEIS-A128828

The number of regular hexagons found by constructing n equally-spaced points on each side of the hexagon and drawing lines parallel to the hexagon side.

(5*n^4-46*n^3 + 1515*n^2 - 6046*n + 7940)/108 when n = 2 + 3k (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Noah Priluck (npriluck(AT)gmail.com), May 08 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004350 | OEIS-A128889

a(n) = (2^(n^2) - 1)/(2^n - 1).

a(n) is prime for n in A156585. Conjecture: gpf(a(n)) = gpf(Phi(n,2^n)), where Phi(n,2^n) = A070526(n). - _Thomas Ordowski_, Feb 16 2014 The conjecture fails at n = 26, where 33407622839523953295063270233033 > 21565632938289155092019246266l. Next counterexample for n = 30, but no odd counterexamples found so far. - _Charles R Greathouse IV_, Feb 17 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Apr 19 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004351 | OEIS-A128909

3D version of A005670. The problem is to dissect an n X n X n cube into smaller integer cubes, the gcd of whose sides is 1, using the smallest number of cubes. The gcd condition exclude dissecting a 6 X 6 X 6 cube into eight 3 X 3 X 3 cubes.

As far as I know, no term, (except trivial cases) has been proved optimal. Repeated dissection, as in the above example, shows that if the side is a composite number mn, a(mn) <= a(m) + a(n) - 1. It is an open problem to find a number mn for which a(mn) < a(m) + a(n) - 1. Dissecting a cube with side n into a cube with side n - 1 and several unit cubes gives a trivial bound: a(n) <= 3n^2 - 3n + 2. Dissecting a cube with side n = 2k + 1 into a cube with side k + 1, 7 with side k and several unit cubes gives another trivial bound: a(n) <= (9n^2 - 12n + 31) / 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mauro Fiorentini_, Apr 23 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004352 | OEIS-A128939

Maximal product over partitions of n into parts of the form 3k+1.

Conjecture. a(n)=1 if n<4, else a(n)=4^[n/4] if n<>4m+3, else a(n)=7*4^([n/4]-1), where [...] denotes the Floor function. (This has been verified up to n=40.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Apr 27 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004353 | OEIS-A128957

a(n) is equal to the number of positive integers m less than or equal to 10^n such that m is not divisible by at least one of the primes 2,5 and is not divisible by at least one of the primes 3,7.

Conjectures from _Colin Barker_, Nov 06 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Milan Janjic_, Apr 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004354 | OEIS-A129011

a(n) = floor(n^(4/3)).

Churchhouse (1971), as an early example of the use of computers in number theory, conjectured that every positive integer N is the sum of two elements of this sequence and verified the conjecture up to N = 10,000 using the Atlas 1 computer of the Atlas Computer Laboratory at Chilton, U.K. He was able to prove that every sufficiently large integer, N, can be expressed in the form N = floor(n^s) + floor(m^s), n and m being positive integers and s being any number in the interval (1, 4/3). - _Peter Bala_, Jan 13 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, May 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004355 | OEIS-A129112

Decimal expansion of constant equal to concatenated semiprimes.

Is this, as Copeland and Erdos (1946) showed for the Copeland-Erdos constant, a normal number in base 10? I conjecture that it is, despite the fact that the density of odd semiprimes exceeds the density of even semiprimes. What are the first few digits of the continued fraction of this constant? What are the positions of the first occurrence of n in the continued fraction? What are the incrementally largest terms and at what positions do they occur?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, May 24 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004356 | OEIS-A129147

Expansion of c(x(1+2x)), c(x) the g.f. of A000108.

Conjecture: (n+1)*a(n) +2*(2-n)*a(n-1) +4*(5-4n)*a(n-2) +16*(2-n)*a(n-3)=0. - _R. J. Mathar_, Dec 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004357 | OEIS-A129149

Permutations with exactly 7 fixed points.

Conjecture: (-n+7)*a(n) +n*(n-8)*a(n-1) +n*(n-1)*a(n-2)=0. - _R. J. Mathar_, Nov 02 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zerinvary Lajos_, May 25 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004358 | OEIS-A129150

The n-th arithmetic derivative of 2^3.

Conjecture: a strictly increasing sequence. - _J. Lowell_, Sep 10 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004359 | OEIS-A129179

Triangle read by rows: T(n, k) is the number of Schroeder paths of semilength n such that the area between the x-axis and the path is k (n >= 0; 0 <= k <= n^2).

Conjecture: T(n, k) = [z^k] R(n, 0) for n >= 0, k >= 0 where R(n, q) = Sum_{j=0..q + (q mod 2) + 1} z^j*R(n-1, j) for n > 0, q >= 0 with R(0, q) = 1 for q >= 0. - _Mikhail Kurkov_, Aug 03 2023 Conjecture above can be simplified as follows: T(n, k) = [z^k] P(n+1, n+1) for n >= 0, k >= 0 where P(n, k) = P(n-1, k) + z^(n-k)*P(n + ((n-k) mod 2), k-1) for 0 < k <= n with P(n, k) = 0 for k > n, P(n, 0) = 1 for n >= 0. - _Mikhail Kurkov_, Oct 02 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Apr 08 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004360 | OEIS-A129364

a(n) = Product_{k = 1..n} A066841(k).

Conjecture: a(n) divides A092287(n) for all n - see comments in A129365.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 11 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004361 | OEIS-A129366

a(n) = Sum_{k=0..floor(n/2)} A000108(n-k).

Conjecture: n*(12*n+35)*(n-1)*a(n) + (n-1)*(12*n^2-701*n+1236)*a(n-1) + 2*(6*n^3-385*n^2+2285*n-3432)*a(n-2) + 4*(-405*n^3+5313*n^2-19970*n +23175)*a(n-3) + 8*(156*n^3-1724*n^2+5498*n-5175)*a(n-4) + 16*(393*n^3-4981*n^2+20393*n-26820)*a(n-5) - 32*(n-5)*(93*n-268)*(2*n-9)*a(n-6) = 0. - _R. J. Mathar_, Feb 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 11 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004362 | OEIS-A129393

Row sums of A129392.

Conjectures from _Colin Barker_, Oct 07 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonas Wallgren_, Apr 13 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004363 | OEIS-A129439

An analog of Pascal's triangle: T(n,k) = A092143(n)/(A092143(n-k)*A092143(k)), 0 <= k <= n.

It appears that the T(n,k) are always integers. This would follow from the conjectured prime factorization given in the comments section of A092143.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 15 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004364 | OEIS-A129453

An analog of Pascal's triangle based on A092287. T(n,k) = A092287(n)/(A092287(n-k)*A092287(k)), 0 <= k <= n.

It appears that the T(n,k) are always integers. This would follow from the conjectured prime factorization given in A092287. Calculation suggests that the binomial coefficients C(n,k) divide T(n,k) and indeed that T(n,k)/C(n,k) are perfect squares.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 16 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004365 | OEIS-A129454

a(n) = Product{i=1..n-1} Product{j=1..n-1} Product{k=1..n-1} gcd(i,j,k).

Conjecture: Let p be a prime and let ordp(n,p) denote the largest power of p which divides n. For example, ordp(48,2)=4 since 48 = 3*(2^4). Then we conjecture that the prime factorization of a(n) is given by ordp(a(n),p)=(floor(n/p))^3 + (floor(n/p^2))^3 + (floor(n/p^3))^3 + ... Compare with the comments in A092287.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 16 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004366 | OEIS-A129455

An analog of Pascal's triangle based on A129454. T(n, k) = A129454(n+1)/(A129454(n-k+1)*A129454(k+1)).

It appears that the T(n,k) are always integers. This would follow from the conjectured prime factorization given in A129454. Calculation suggests that the binomial coefficients C(n,k) divide T(n,k) and that T(n,k)/C(n,k) are perfect sixth powers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 16 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004367 | OEIS-A129507

G.f.: (1+x+x^2-sqrt(1+2x+3x^2-2x^3+x^4))*2.

Conjecture: g.f.: 4*q^2*(1 - 1/G(0)) where G(k) = 1 + q/(1 + q^2 / G(k+1)). - _Joerg Arndt_, Jul 17 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 18 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004368 | OEIS-A129527

a(2n) = a(n) + 2n, a(2n+1) = 2n + 1.

Conjecture: a(n) = 2*n-A000265(n) for n > 0. - _Velin Yanev_, Jun 23 2017. [_Joerg Arndt_, Jun 23 2017: For odd n the conjecture holds, for even n induction should work. _Andrey Zabolotskiy_, Aug 03 2017: Confirm: induction works, the conjecture holds for all n.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, May 29 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004369 | OEIS-A129654

Number of different ways to represent n as general polygonal number P(m,r) = 1/2*r*((m-2)*r-(m-4)) = n>1, for m,r>1.

G.f.: x * Sum_{k>=1} x^k / (1 - x^(k*(k + 1)/2)) (conjecture). - _Ilya Gutkovskiy_, Apr 09 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Apr 27 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004370 | OEIS-A129683

Expansion of (1/(1-2x))*exp(2x/(1-2x)).

Conjecture: a(n) -4*n*a(n-1) +4*(n-1)^2*a(n-2)=0. - _R. J. Mathar_, Nov 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004371 | OEIS-A129695

Laguerre transform of the Jacobsthal numbers.

Conjecture: a(n) +(-4*n+3)*a(n-1) +(6*n^2-16*n+9)*a(n-2) -(4*n-7)*(n-2)^2*a(n-3) +(n-2)^2*(n-3)^2*a(n-4)=0. - _R. J. Mathar_, Feb 23 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004372 | OEIS-A129756

Repetitions of odd numbers four times.

Conjecture: number of roots of P(x) = x^n - x^(n-1) - x^(n-2) - ... - x - 1 in the right half-plane. - _Michel Lagneau_, Apr 09 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_ and _Giorgio Balzarotti_, May 15 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004373 | OEIS-A129758

Smallest prime p such that there are primes q and r with the property that p, q and r form an arithmetic progression and their sum is the same as three times the n-th prime number.

Conjecture: Limit_{N->oo} (Sum_{n=1..N} a(n)) / (Sum_{n=1..N} prime(n+2)) = 1. - _Alain Rocchelli_, May 01 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Teofilatto_, May 15 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004374 | OEIS-A129759

For the Landau function L(n), A000793, this sequence gives the largest prime which is a factor of L(n).

Nicolas showed that a(n) ~ sqrt(n log n) and Grantham showed that a(n) <= 1.328 sqrt(n log n) for n > 4. Massias, Nicolas, & Robin conjecture that a(n) <= 1.265... sqrt(n log n) in this range with equality at n = 215. - _Charles R Greathouse IV_, Jun 02 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anthony C Robin_, May 15 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004375 | OEIS-A129762

Sum of all elements of n X n X n cubic array M[i,j,k] = Fibonacci[i+j+k-2].

a(n) is prime for n = {2, 7, 19, 47, 175, 179, ...}. The formula a(n) = F(3n+4) - 3F(2n+4) + 3F(n+4) - 3 and its generalization for k-dimensional hypercubes with elements M(i,j,...) = F(i+j+...-k+1) was stated and proved by the user 1istik_figi in private communication at LiveJournal on Oct 10 2007. The k-dimensional formula is a(n) = Sum[(-1)^i*Binomial[k,i]*Fibonacci[(k-i)*n+k+1],{i,0,k}]. Conjecture: if prime p divides F(p-1) then p^k divides a(n) in k-dimensional case.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, May 15 2007, Oct 11 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004376 | OEIS-A129783

Conjectured numbers n with the property that there exist two consecutive primes p and q such that pq + n is a square.

Conjectured numbers n with the property that there exist two consecutive primes p and q such that pq + n is a square.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, May 18 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004377 | OEIS-A129807

Primes congruent to +-7 mod 18.

Conjecture: Equals (A061241 UNION A061238) MINUS {2}. - _R. J. Mathar_, Jun 10 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 22 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004378 | OEIS-A129811

Lesser of consecutive prime pairs whose last digits differ by 4.

Conjecture: The number of terms in this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, May 19 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004379 | OEIS-A129812

Lesser of consecutive prime pairs whose last digits differ by 6.

Conjecture: The number of terms in this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, May 19 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004380 | OEIS-A129813

Lesser of consecutive prime pairs whose last digits differ by 8.

Conjecture: The number of terms in this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, May 19 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004381 | OEIS-A129816

Conjectured numbers n such that there do not exist two consecutive primes whose product + n is a square.

Conjectured numbers n such that there do not exist two consecutive primes whose product + n is a square.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, May 20 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004382 | OEIS-A129817

Number of alternating fixed-point-free permutations on n letters.

We conjecture that for n >= 3, A000111(2n)/a(2n) < e < A000111(2n)/A129815(2n), so that A000111(2n)/a(2n) increases while A000111(2n)/A129815(2n) decreases (and both quotients tend to e). - _Vladimir Shevelev_, Apr 29 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, May 20 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004383 | OEIS-A129829

Conjectured first occurrence of numbers n with the property that there exist two consecutive primes p and q such that pq + n is a cube.

Conjectured first occurrence of numbers n with the property that there exist two consecutive primes p and q such that pq + n is a cube.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, May 20 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004384 | OEIS-A129830

Conjectured first occurrence of numbers n with the property that there exist two consecutive primes p and q such that pq + n is a fourth power.

Conjectured first occurrence of numbers n with the property that there exist two consecutive primes p and q such that pq + n is a fourth power.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, May 21 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004385 | OEIS-A129833

a(n) = Sum_{k = 0..n } binomial(n + 1, k + 1)*binomial(n, k)*k!.

Conjecture: +(n+1)*a(n) -n*(2*n+5)*a(n-1) +(n-1)*(n^2+6*n+3)*a(n-2) -(n-2)*(3*n^2-2)*a(n-3) +(n-2)*(n-3)*(3*n-4)*a(n-4) -(n-4)*(n-3)^2*a(n-5) = 0. - _R. J. Mathar_, Feb 28 2015 Conjecture: (n+1)*(n^2-4*n+2)*a(n) -n*(2*n^3-5*n^2-6*n+3)*a(n-1) +n*(n-1)*(n^3-2*n^2-2*n-2)*a(n-2) -(n-2)*(n^2-2*n-1)*(n-1)^2*a(n-3) = 0. - _R. J. Mathar_, Feb 28 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, May 21 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004386 | OEIS-A129890

a(n) = (2*n+2)!! - (2*n+1)!!.

Conjecture: a(n) = (2*n + 2)*(2*n + 2)! * Sum_{k >= 1} (-1)^(k+1)/Product_{j = 0..n+1} (k + 2*j) [added Dec 03 2025: this identity can be verified using Maple's sum command]. - _Peter Bala_, Jul 06 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_ and _Giorgio Balzarotti_, Jun 04 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004387 | OEIS-A129924

Primes p such that p divides both A061354(p-3) and A061354(p-1).

Conjecture: a(n) = A064384(n+1). The conjecture is true. It is the case n = p-3 of the relation GCD(A061354(n), A061354(n+2)) = A124779(n), which follows from the Comments in A064384 and A124779. For a proof, see the link "The Taylor series for e ...". - _Jonathan Sondow_, Jun 12 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Jun 06 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004388 | OEIS-A129995

a(n) = (n + 1)*(n^2 + 2)*(n^3 + 3)*(n^4 + 4)/4!.

Following my conjecture, computations by _Peter J. C. Moses_, mediation by _Clark Kimberling_ and helpful comments from George E. Andrews, it is now known that a(n) = (n^1 + 1)*(n^2 + 2)*(n^3 + 3)*...*(n^k + k)/k! is an integer-valued sequence if and only if k belongs to {1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 17, 18, 19, 20, 21}; this is the case for k=4; see generalization in A131685.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander R. Povolotsky_, Aug 19 2007, Aug 25 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004389 | OEIS-A130012

Smallest natural number whose cube is the sum of n cubes of distinct natural numbers, or 0 if no such number exists.

Conjecture: sequence is nondecreasing for n > 4. - _Zhining Yang_, Apr 24 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Jun 15 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004390 | OEIS-A130058

Numbers m such that k = m*23^2 divides 3^(k-1) - 2^(k-1).

Conjecture: All nonsquarefree terms of A073631 are the multiples of 23^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, May 04 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004391 | OEIS-A130095

Inverse Möbius transform of odd-indexed Fibonacci numbers.

Conjecture: a(n)/a(n-1) tends to phi^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, May 06 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004392 | OEIS-A130145

Number of nonisomorphic orthogonal arrays OA(8*n+4,4,2,2).

Conjecture: partial sums of A115266. - _Sean A. Irvine_, Jul 14 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Aug 15 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004393 | OEIS-A130187

Numerators of rationals r(n) related to the z-sequence of the Sheffer matrix A060821 for Hermite polynomials.

Conjecture: a(n) = A001147(n+1)/A000265(n+1), n >= 0. (Motivated to reconsider this sequence by an e-mail of Thomas Olson.) - _Wolfdieter Lang_, Jan 04 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jun 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004394 | OEIS-A130287

a(n) = A121296(n) - A121263(n).

We conjecture that all terms are >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Applegate_ and _N. J. A. Sloane_, Sep 21 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004395 | OEIS-A130519

a(n) = Sum_{k=0..n} floor(k/4). (Partial sums of A002265.)

Conjecture: a(n+2) is the maximum effective weight of a numerical semigroup S of genus n (see Nathan Pflueger). - _Stefano Spezia_, Jan 04 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hieronymus Fischer_, Jun 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004396 | OEIS-A130536

Prime numbers arising from A057856.

Conjecture: For all pairs of relative prime numbers (x, y) there exists at least one number n = 2^m and one prime number p such that p = x^n + y^n. This sequence shows one case of this conjecture where y = x + 1. The conjecture for arbitrary coprime positive integers x and y is false. Take x = 8 and y = 27. For every integer r >= 1, 8^r + 27^r = (2^r + 3^r)*(4^r - 6^r + 9^r), with both factors greater than 1. In particular, 8^(2^m) + 27^(2^m) is composite for every integer m >= 0. This does not refute the special case y = x + 1. See also A057856. - _Eiver Rodríguez_, Oct 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tomas Xordan_, Jun 02 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004397 | OEIS-A130542

The maximum absolute value of the L-series coefficient for an elliptic curve.

The values of a(n) and the multiplicativity are conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Jun 04 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004398 | OEIS-A130568

Generalized Beatty sequence 1+2*floor(n*phi), which contains infinitely many primes.

Conjecture: Sequence gives n of A163873 whose connection to a(n) crosses (in the tree of A163873) another path. Is this generalizable in any way for A163874, A163875? - Daniel Platt (d.platt(AT)web.de), Sep 14 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Aug 09 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004399 | OEIS-A130595

Triangle read by rows: lower triangular matrix which is inverse to Pascal's triangle (A007318) regarded as a lower triangular matrix.

Conjecture from _Dale Gerdemann_, Nov 28 2015:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Delépham_, Jun 17 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004400 | OEIS-A130633

Additive persistence of Fibonacci numbers.

Up to 10^22 the maximum value is 4. Conjecture: for n > 15, a(n) > 1. I checked the conjecture up to 10^10. - _Charles R Greathouse IV_, Feb 12 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_ and _Giorgio Balzarotti_, Jun 19 2007, corrected Jun 22 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004401 | OEIS-A130651

1/16 the number of permutations of 0..n having exactly 3 maxima.

Conjecture: a(n) = 2^(n-8)*(3^(n+1) - 2^(n+2)*n + 2*n^2 - 3). - _Vaclav Kotesovec_, Nov 29 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 11 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004402 | OEIS-A130655

Catalan transform of Catalan numbers C(n+1).

Conjecture: 3*n*(n-2)*(n+2)*a(n)
+4*(-10*n^3+21*n^2+7*n-15)*a(n-1)
+16*(11*n^3-47*n^2+57*n-15)*a(n-2)
-8*(2*n-5)*(4*n-9)*(4*n-7)*a(n-3)=0. - _R. J. Mathar_, Mar 01 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Delépham_, Jun 21 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004403 | OEIS-A130694

Exponents of powers of 2 that contain all ten digits.

a(n) = n + 91 for n >= 78 (conjectured). - _Chai Wah Wu_, Jan 27 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Greg Dresden_, Jul 10 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004404 | OEIS-A130729

Fermat numbers of order 3 or F(n,3) = 2^(2^n) + 3.

This is equivalent to F(n)+2 or 2^(2^n)+ 1 + 2. Conjecture: If n is odd, 7 is a divisor of F(n,3). The conjecture is true: the order of 2 mod 7 is 3, and if n is odd then 2^n == 2 mod 3 so 2^(2^n) + 3 == 2^2 + 3 == 0 mod 7. - _Robert Israel_, Nov 20 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jul 05 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004405 | OEIS-A130740

a(n) = 2^(4^n) + 5.

Numbers of the form 2^(4^n) + 5 are divisible by 7. We prove this by induction making use of the expansion (1) a^m - b^m = (a-b)(a^(m-1)+a^(m-2)b +...+ b^(m-1). For n = 1, we have 2^4 + 5 = 21 = 7*3. So the statement is true for n=1. Now assume the statement is true for some integer k and show that it is also true for k+1. Thus we have 2^(4^k) + 5 = 7h for some h. Let 2^(4^(k+1))+ 5 = h1. Now consider the difference h1 - 7h. If this is a multiple of 7 then so is h1. So we have 2^(4^(k+1))+ 5 - (2^(4^k) + 5)) = 16^(4^k)-2^(4^k). This is of the form (1) where n = 4^k, a = 16 and b = 2. So the difference, a^n-b^n is divisible by (a-b) = (16-2) = 14 = 7*2. This implies 2^(4^(k+1))+5 is divisible by 7. So we have assumed the statement was true for k and have shown it to be true for k+1. Therefore by the induction hypothesis, the statement is true for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jul 07 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004406 | OEIS-A130741

a(n) = 4^(4^n) + 3.

Numbers of the form 4^(4^n) + 3 are divisible by 7. We prove this by induction making use of the expansion (1) a^m - b^m = (a-b)(a^(m-1)+a^(m-2)b+...+b^(m-1). For n = 1, we have 4^4 + 3 = 259 = 7*37. So the statement is true for n=1. Now assume the statement is true for some integer k and show that it is also true for k+1. Thus we have 4^(4^k) + 3 = 7h for some h. Let 4^(4^(k+1))+ 3 = h1. Now consider the difference h1 - 7h. If this is a multiple of 7 then so is h1. So we have 4^(4^(k+1))+ 3 - (4^(4^k) + 3)) = 256^(4^k)-4^(4^k). This is of the form (1) where n = 4^k, a = 256 and b = 4. So the difference, a^n-b^n is divisible by (a-b) = (256-4) = 252 = 7*36. This implies 4^(4^(k+1))+3 is divisible by 7. So we have assumed the statement was true for k and have shown it to be true for k+1. Therefore by the induction hypothesis, the statement is true for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jul 07 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004407 | OEIS-A130770

One third of the least common multiple of 3 and n^2+n+1.

Conjecture: a(n) = A046163(n), n>0. - _R. J. Mathar_, Jun 13 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _W. Neville Holmes_, Jul 14 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004408 | OEIS-A130789

The primes prime(n) sorted according to increasing prime(n)/prime(n+1).

Or: primes sorted according to decreasing ratio A001223(n)/A000040(n). All values are conjectural, derived from the finite list up to prime(200000): large prime gaps at higher indices may still insert numbers above prime(200000) at low positions of the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jul 15 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004409 | OEIS-A130896

For D_n type groups as polyhedra that are pyramid-like: {F,V,E,dimension}->{2*n+1,2*n+1,4*n,(2*n+1)*((2*n+1)-1)/2} such that Euler's equation is true: V=E-F+2.

Conjectures from _Colin Barker_, Jan 02 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Aug 22 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004410 | OEIS-A131011

a(n) = A121295(n) - A121263(n).

We conjecture that all terms are >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 24 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004411 | OEIS-A131012

a(n) = A121295(n) - A121265(n).

We conjecture that from some point on, all terms are >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 24 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004412 | OEIS-A131016

Smallest semiprime == 1 (mod n).

This is to semiprimes A001358 as A034694 is to primes A000040. Conjecture: every semiprime occurs in this sequence. a(21)-a(65) from _Robert G. Wilson v_, Sep 21 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Sep 22 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004413 | OEIS-A131135

Denominator of (ordinary) expansion of log((x/2-1)/(x-1)).

a(n) = denominator((1-1/2^n)/n), for n > 0, conjectured. - _Michel Marcus_, Sep 12 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jun 17 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004414 | OEIS-A131137

Denominator of (exponential) expansion of log((2*x/3-1)/(x-1)).

Conjecture: starting at third term, also equals the denominator of polylog(-n,1/4)/4. - _Wouter Meeussen_, Feb 13 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jun 17 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004415 | OEIS-A131178

Non-plane increasing unary binary (0-1-2) trees where the nodes of outdegree 1 come in 2 colors.

G.f.: conjecture: T(0)/(1-2*x) -1, where T(k) = 1 - x^2*(k+1)*(k+2)/(x^2*(k+1)*(k+2) - 2*(1-2*x*(k+1))*(1-2*x*(k+2))/Q(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 19 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wenjin Woan_, Oct 31 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004416 | OEIS-A131280

Sums of exactly 4 positive octahedral numbers A005900.

Pollock (1850) conjectured that every number is the sum of at most 7 octahedral numbers. Which octahedral numbers are themselves the sum of exactly 4 positive octahedral numbers? To begin with, Oc(3) = Oc(2) + Oc(2) + Oc(2) + Oc(1) = 6 + 6 + 6 + 1 = 19.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Oct 21 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004417 | OEIS-A131357

a(1) = 1, a(2) = 2; for n > 1, a(n) = sum of the next two smallest integers > a(n-1) which are coprime to the sum s = a(1) + ... + a(n-1).

Conjecture: a(n) ~ 2^(n+1) * 0.9873939997556885598467... - _Petros Hadjicostas_, Jun 23 2019 [Constant evaluated using _Michel Marcus_'s bfile]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Sep 30 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004418 | OEIS-A131388

Sequence (a(n)) generated by Rule 1 (in Comments) with a(1) = 1 and d(1) = 0.

Conjecture: if a(1) is an nonnegative integer and d(1) is an integer, then (a(n)) is a permutation of the nonnegative integers (if a(1) = 0) or a permutation of the positive integers (if a(1) > 0). Moreover, (d(n)) is a permutation of the integers if d(1) = 0, or of the nonzero integers if d(1) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 05 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004419 | OEIS-A131389

Sequence (d(n)) generated by Rule 1 (see Comments) with a(1) = 1 and d(1) = 0.

Rule 1 is given at A131388. Conjecture: Every integer occurs exactly once in A131389.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 05 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004420 | OEIS-A131391

Positions of positive integers in A131389, minus 1.

If A131389 is a permutation of the integers, as conjectured, then A131391 is the complement of A131392.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 06 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004421 | OEIS-A131392

Positions of negative integers in A131389, minus 1.

If A131389 is a permutation of the integers, as conjectured, then A131392 is the complement of A131391.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 06 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004422 | OEIS-A131394

Conjectured permutation of the integers using Rule 2 with A131393(1)=1.

Conjectured permutation of the integers using Rule 2 with A131393(1)=1. Rule 2 is given at A131393. Conjecture: A131394 is a permutation of the integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 05 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004423 | OEIS-A131396

Positions of n-th positive integer in A131394, minus 1.

If A131394 is a permutation of the integers, as conjectured, then A131396 is the complement of A131397.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 06 2007, Jul 21 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004424 | OEIS-A131397

Positions of negative integers in A131394, minus 1.

If A131394 is a permutation of the integers, as conjectured, then A131397 is the complement of A131396.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 06 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004425 | OEIS-A131401

Least number dividing Fibonacci(n) but not dividing Fibonacci(m) for m < n, or 0 if there is no such number.

Conjecture: only a(2)=0. I have not found values of a(n) < 2*106 less than 100 for n = 43, 47, 74, 82, 83 & 94. In other words, the conjecture is true. For n > 2, Fibonacci(n) has at least one divisor that does not divide Fibonacci(k) for k < n. The number of such divisors is A120256(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Herbert A. Hauptman (hauptman(AT)hwi.buffalo.edu) & _Robert G. Wilson v_, Jul 07 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004426 | OEIS-A131418

Numbers n such that Sum_digits(n)=Sum_digits[n+Sum_digits(n)], with n>=0.

Conjecture: all terms are divisible by 9. - _Harvey P. Dale_, Mar 28 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_ and _Giorgio Balzarotti_, Jul 09 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004427 | OEIS-A131439

Inverse binomial transform of A131438 (assuming zero offset in both sequences).

Conjecture: The sequence appears to be (1, 7, ...) followed by 4k + 14; k=0,1,2,...; thus: (1, 7, 14, 18, 22, 26, ...).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Jul 11 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004428 | OEIS-A131450

a(n) is the number of integers x that can be written x = (2^c(1) - 2^c(2) - 3*2^c(3) - 3^2*2^c(4) - ... - 3^(m-2)*2^c(m) - 3^(m-1)) / 3^m for integers c(1), c(2), ..., c(m) such that n = c(1) > c(2) > ... > c(m) > 0 and c(1) - c(2) != 2 if m >= 2.

It is conjectured that lim_{n->oo} a(n)/a(n-1) = 4/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Perry Dobbie_, Jul 11 2007, Jul 12 2007, Jul 13 2007, Jul 17 2007, Jul 22 2007, Oct 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004429 | OEIS-A131498

For D_2 type groups as polyhedra: {F,V,E,dimension}->{n+2,2*n,3*n,2*n*(2*n-1)/2} such that Euler's equation is true: V=E-F+2.

G.f.: x*(3 + 2*x + 3*x^2 + x^3 - 5*x^4 - 2*x^5 - 3*x^6 + 3*x^7 + 2*x^8) / ((1 - x)^3*(1 + x)^3*(1 + x^2)^3) (conjectured). - _Colin Barker_, Jan 03 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Aug 12 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004430 | OEIS-A131501

Xm/CV where Xm is a point of maximum error using an approximation method for x^(1/2) which I have found and CV is the population coefficient of variation from my list of error values.

a(n) = A146951(n) for 1 <= n <= 10, but more terms would be needed to justify such a hypothesis. - _Georg Fischer_, Nov 03 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | Anthony J. Browne (tony2theipi(AT)yahoo.com), Aug 13 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004431 | OEIS-A131576

Number of ways to represent n as a sum of an even number of consecutive integers.

Conjecture: a(n) is also the number of pairs of subparts in the symmetric representation of sigma(n) which are mirror images of each other in the main diagonal. (Cf. A279387). - _Omar E. Pol_, Feb 22 2017 [Conjecture clarified by _N. J. A. Sloane_, Dec 16 2020] Conjectures: a(n) = (A001227(n) - A067742(n))/2 = A082647(n) - A067742(n). - _Omar E. Pol_, Feb 22 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Aug 28 2007, Sep 16 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004432 | OEIS-A131607

Pell companion numbers A001333 without last digit.

Conjectures from _Colin Barker_, Jan 03 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Oct 02 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004433 | OEIS-A131613

Numbers k such that the decimal expansion of 3^k contains no 9.

I conjecture that 56 is the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shyam Sunder Gupta_, Sep 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004434 | OEIS-A131614

Numbers k such that the decimal expansion of 3^k contains no 8.

I conjecture that 100 is the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shyam Sunder Gupta_, Sep 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004435 | OEIS-A131615

Numbers k such that the decimal expansion of 3^k contains no 7.

I conjecture that 101 is the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shyam Sunder Gupta_, Sep 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004436 | OEIS-A131616

Numbers k such that the decimal expansion of 3^k contains no 6.

I conjecture that 104 is the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shyam Sunder Gupta_, Sep 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004437 | OEIS-A131617

Numbers k such that the decimal expansion of 3^k contains no 5.

I conjecture that 91 is the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shyam Sunder Gupta_, Sep 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004438 | OEIS-A131618

Numbers k such that the decimal expansion of 3^k contains no 4.

I conjecture that 106 is the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shyam Sunder Gupta_, Sep 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004439 | OEIS-A131625

Numbers k such that decimal expansion of 3^k contains no 2.

I conjecture that 59 is the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shyam Sunder Gupta_, Sep 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004440 | OEIS-A131626

Prime subsequences in the evaluation of the first derivatives of quadratic equations with positive successive prime coefficients.

Conjecture: The number of primes in a row for f'(x,p) = 2*prime(p)*x + prime(p+1) is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Sep 03 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004441 | OEIS-A131627

Numbers k such that the decimal expansion of 3^k contains no 1.

I conjecture that 43 is the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shyam Sunder Gupta_, Sep 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004442 | OEIS-A131629

Numbers k such that the decimal expansion of 3^k contains no 3.

I conjecture that 84 is the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shyam Sunder Gupta_, Sep 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004443 | OEIS-A131646

Numbers that can be written from base 2 to base 18 using only the digits 0 to 9 (conjectured to be complete).

Numbers that can be written from base 2 to base 18 using only the digits 0 to 9 (conjectured to be complete). It is a plausible conjecture that there are no more terms, but this has not been proved. - _N. J. A. Sloane_, Nov 17 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Mondot_, Sep 08 2007, Nov 02 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004444 | OEIS-A131649

Number of distinct improper 2-coloring of edges for odd-order cyclic graphs.

Conjectured g.f.: 4 + (2*x*((x-2)^2*x-4))/(x-1)^3. - _Harvey P. Dale_, Jul 07 2013 For n>4, a(n) = 3*n^2 - 17*n + 26 (conjectured). - _Ralf Stephan_, Jul 08 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | M. Razid Black (mrazidblack(AT)hotmail.com), Sep 10 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004445 | OEIS-A131657

For n >= 1, put A_n(z) = Sum_{j>=0} (n*j)!/(j!^n) * z^j and B_n(z) = Sum_{j>=0} (n*j)!/(j!^n) * z^j * (Sum_{k=1..j*n} (1/k)), and let b(n) be the largest integer for which exp(B_n(z)/(b(n)*A_n(z))) has integral coefficients. The sequence is b(n).

Krattenhaler and Rivoal (2007-2009) conjecture that a(n) = n!*Xi(n), where Xi(1) = 1, Xi(7) = 1/140, and Xi(n) = Product_{p <= n} p^min(2, v_p(H_n)) for n <> 1, 7, where v_p(r) is the p-adic valuation of rational r. (Here p indicates a prime and H_n is the n-th harmonic number.) - _Petros Hadjicostas_, May 24 2020 A formula, conditional on a widely believed conjecture, can be found in Theorem 3 with k = 1 in the article by Krattenhaler and Rivoal (2007-2009) cited in the references; see the remarks before Theorem 4 in that article.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Christian Krattenthaler (Christian.Krattenthaler(AT)univie.ac.at), Sep 12 2007, Sep 30 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004446 | OEIS-A131658

For n >= 1, put A_n(z) = Sum_{j>=0} (n*j)!/(j!^n) * z^j and B_n(z) = Sum_{j>=0} (n*j)!/(j!^n) * z^j * (Sum__{k=j+1..j*n} (1/k)), and let u(n) be the largest integer for which exp(B_n(z)/(u(n)*A_n(z))) has integral coefficients. The sequence is u(n).

A formula, conditional on a widely believed conjecture, can be found in the article by Krattenthaler and Rivoal (2007-2009) cited in the references: see Theorem 4 and the accompanying remarks.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Christian Krattenthaler (Christian.Krattenthaler(AT)univie.ac.at), Sep 12 2007, Sep 30 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004447 | OEIS-A131662

Numbers n where either n or n+1 is divisible by the numbers from 1 to 12.

Conjectures from _Colin Barker_, Jun 15 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tanya Khovanova_, Sep 13 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004448 | OEIS-A131853

Numbers m such that z(m)=(0,0) with z as defined in A131851.

conjecture: a(n) mod 5 = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jul 22 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004449 | OEIS-A131883

a(n) = the minimum value from among (phi(n+1),phi(n+2),phi(n+3),...,phi(2n)), where phi(m) (A000010) is the number of positive integers which are coprime to m and are <= m.

Conjecture: After omitting multiple occurrences we get A036912. - _Vladeta Jovovic_, Oct 31 2007. This conjecture has been established by _Max Alekseyev_ - see link below.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Oct 24 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004450 | OEIS-A131884

Numbers conjectured to have an infinite, aperiodic, aliquot sequence.

Numbers conjectured to have an infinite, aperiodic, aliquot sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Oct 24 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004451 | OEIS-A131935

a(n) is the number of Khalimsky-continuous functions with four-point codomain and an n-point range.

Conjectures from _Colin Barker_, Jan 13 2018: (Start) (End) [Since we have an explicit set of recurrences that produce a(n), it should be straightforward to prove these conjectures. - _N. J. A. Sloane_, Jan 14 2018]

猜想/问题提及段 | historical_or_current_mention_unreviewed | Shiva Samieinia (shiva(AT)math.su.se), Oct 05 2007, Oct 09 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004452 | OEIS-A131937

a(1)=1; a(2)=4, a(n) = a(n-1) + (n-th positive integer which does not occur in sequence A131938).

Conjecture: Starting with any infinite positive integer sequence A1(n) whose complement is also infinite, let B1(n) be the sequence of partial sums of the complement of sequence A1. Let A2(n) be the sequence of partial sums of the complement of sequence B1. Iteratively let B[k](n) be the sequence of partial sums of the complement of sequence A[k], and let A[k+1](n) be the sequence of partial sums of the complement of sequence B[k]. Call the stable iterative limit sequences A(n) and B(n), which are universal and do not depend upon the initial sequence A1(n). These two limiting sequences are conjectured to be A131938 and this A131937. - _Gregory Gerard Wojnar_, Dec 12 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Jul 30 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004453 | OEIS-A131939

Least k such that the difference between consecutive 3-almost primes A014612(k) equals n, or 0 if no such k exists.

Index of smallest 3-almost prime gap equal to n. This is to 3-almost primes A014612 as A123375 is to semiprimes A001358. a(25) = 0 or a(25) > 10000. Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Oct 05 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004454 | OEIS-A131971

a(0) = a(1) = a(2) = 1; a(n) = (a(n-1) + a(n-2) + a(n-3)) mod n.

Tribonacci (A000213) analog of A096535. The analogous 3 _Klaus Brockhaus_ conjectures are applicable: (1) All numbers appear infinitely often, i.e., for every number k >= 0 and every frequency f > 0 there is an index i such that a(i) = k is the f-th occurrence of k in the sequence. (2) a(j) = a(j-1) + a(j-2) + a(j-3) and a(j) = a(j-1) + a(j-2) + a(j-3) - j occur approximately equally often, i.e., lim_{n -> infinity} x_n / y_n = 1, where x_n is the number of j <= n such that a(j) = a(j-1) + a(j-2) + a(j-3) and y_n is the number of j <= n such that a(j) = a(j-1) + a(j-2) + a(j-3) - j (cf. A122276). (3) There are sections a(g+1), ..., a(g+k) of arbitrary length k such that a(g+h) = a(g+h-1) + a(g+h-2) + a(g+h-3) for h = 1,...,k, i.e., the sequence is nondecreasing in these sections (cf. A122277, A122278, A122279).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Oct 05 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004455 | OEIS-A131978

A049614(n)/A131685(n).

Conjectured to be always integral.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander R. Povolotsky_, Sep 21 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004456 | OEIS-A132064

Numbers multiplied by 4 and written backwards.

Conjecture: a(n)^(1/n) tends to 10. - _Vaclav Kotesovec_, Jan 03 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | Rachit Agrawal (rachit_agrawal(AT)daiict.ac.in), Oct 30 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004457 | OEIS-A132075

A conjectured permutation of the positive integers such that for every n, a(n) is the largest number among a(1), a(2), ..., a(n) that when added to a(n+1) gives a prime.

A conjectured permutation of the positive integers such that for every n, a(n) is the largest number among a(1), a(2), ..., a(n) that when added to a(n+1) gives a prime. It is not until the 26th truncated row - [1, 2, 3, 4, 7, 6] - that the initial list is extended at all. It is unclear whether this process can be continued indefinitely, although I have verified by computer that it generates a sequence of at least 2000 terms. Conjecturally: (1) the sequence is infinite, (2) it is the unique sequence containing infinitely many complete rows of table A088643, and (3) for every n > 0 there exists N > 0 such that the first n terms of this sequence are contained in every row of table A088643 from the N-th onwards.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Boddington_, Oct 30 2007, Mar 06 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004458 | OEIS-A132089

Triangle T, read by rows, equal to the matrix square of Losanitsch's triangle (A034851).

Conjectures: (1) The first column equals the row sums of Losanitsch's triangle (which equals A005418). (2) Let L^n denote the n-th power of Losanitsch's triangle. Then the first column of L^(n+1) equals the row sums of L^n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerald McGarvey_, Oct 30 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004459 | OEIS-A132091

Expansion of psi(x^3) * chi(-x^9) / f(-x^2) in powers of x where psi(), chi(), f() are Ramanujan theta functions.

Also number of partitions of n into parts not divisible by 3 with every part repeated at least twice. Conjectured by _R. H. Hardin_, Jun 06 2009, proved by _Max Alekseyev_, Jun 06 2009.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Aug 09 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004460 | OEIS-A132113

Multiply previous term by 8 and reverse.

Conjecture: a(n)^(1/n) tends to 10. - _Vaclav Kotesovec_, Jan 03 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | Rachit Agrawal (rachit_agrawal(AT)daiict.ac.in), Oct 31 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004461 | OEIS-A132163

Triangle read by rows. For row n, start with 1 but from the second term onwards always choose the largest positive integer between 1 and n inclusive that i) has not already appeared in the row ii) gives a prime when added to the previous term. Stop if no such integer can be found.

The following statements are conjectural: 1) The n-th row is always a permutation of 1,...,n. 2) For the even rows, the last term is one less than a prime (so the row gives a solution to the prime circle problem - see A051252). 3) There exists a (unique) sequence b(2), b(3),... with the property that for every n > 1 there is a positive integer N such that every even row of the triangle from the 2N-th onwards ends b(n), ..., b(3), b(2) and every odd row from the (2N - 1)-th onwards ends b(n)+(-1)^n, ..., b(3)-1, b(2)+1. (If the sequence b(n) exists it is probably A132075 without the initial term 1.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Boddington_, Nov 04 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004462 | OEIS-A132293

Maximal number of right angles in an n-gon.

Conjecture from _R. J. Mathar_, Mar 07 2008, Apr 21 2008: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Asher Stuhlman (asherstuhlman(AT)gmail.com), Nov 06 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004463 | OEIS-A132297

Number of distinct Markov type classes of order 2 possible in binary strings of length n.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | Lav R. Varshney (lrv(AT)mit.edu), Aug 17 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004464 | OEIS-A132364

Expansion of 1/(1-x^2*c(x)), c(x) the g.f. of A000108.

Conjecture: +(-n+1)*a(n) +(5*n-11)*a(n-1) +2*(-2*n+5)*a(n-2) +(-n+1)*a(n-3) +2*(2*n-5)*a(n-4)=0. - _R. J. Mathar_, Aug 28 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Nov 08 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004465 | OEIS-A132399

Number of ordered ways of writing n = i + j, where i is 0 or a prime and j is a triangular number (A000217) >= 0.

Based on a posting by _Zhi-Wei Sun_ to the Number Theory Mailing List, Mar 23 2008, where he conjectures that a(n) > 0 except for n = 216. _Zhi-Wei Sun_ has offered a monetary reward for settling this conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 23 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004466 | OEIS-A132411

a(0) = 0, a(1) = 1 and a(n) = n^2 - 1 with n >= 2.

Conjecture: the sequence terms are the exponents in the expansion of the q-series Sum_{n >= 0} q^(2*n) * Product_{k >= 2*n+2} 1 - (-q)^k = 1 + q^3 + q^8 + q^15 + q^24 + ... - _Peter Bala_, May 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mohamed Bouhamida_, Nov 12 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004467 | OEIS-A132415

Integers that can be expressed as the sum of distinct squares in exactly a prime number of ways.

Conjecture: Indices of primes in sequence A033461. - _Vaclav Kotesovec_, Aug 04 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Nov 12 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004468 | OEIS-A132616

Column 0 of triangle A132615.

Conjecture: a(n) is described by a series of nested sums,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 24 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004469 | OEIS-A132634

a(n) = Fibonacci(n) mod n^2.

a(n)=0 for n=1 and n=12 only (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hieronymus Fischer_, Aug 24 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004470 | OEIS-A132794

Numbers n such that sigma(phi(n)) -phi(n) -1 = phi(sigma(n) -n -1).

Conjecture: a(n)=2^(A000043(n)+1), if true the next terms are: 4294967296, 4611686018427387904, 1237940039285380274899124224... - _Jon Maiga_, Dec 14 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_ and _Giorgio Balzarotti_, Aug 31 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004471 | OEIS-A132827

Table based upon insertion points of n into sequence A132828 and having a specific formula.

Conjecture: The array A132827 is the dispersion of the sequence f given by f(n)=floor(n*x+n+1), where x=(golden ratio). Evidence: use f(n_):=Floor[n*x+n+1] in the Mathematica program at A191426. - _Clark Kimberling_, Jun 03 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kenneth J Ramsey_, Sep 03 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004472 | OEIS-A132864

Expansion of 1/(1-4x*c(5x)), where c(x) is the g.f. of A000108.

Conjecture: n*a(n) + 2*(15-2*n)*a(n-1) + 160*(3-2*n)*a(n-2) = 0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Nov 18 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004473 | OEIS-A132900

Colored Motzkin paths where each of the steps has three possible colors.

Conjecture: (n+2)*a(n) - 3*(2*n+1)*a(n-1) + 27*(1-n)*a(n-2) = 0. - _R. J. Mathar_, Nov 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 04 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004474 | OEIS-A133058

a(0) = a(1) = 1; for n > 1, a(n) = a(n-1) + n + 1 if a(n-1) and n are coprime, otherwise a(n) = a(n-1)/gcd(a(n-1),n).

It is conjectured that quasi-periodic sequences exist only for R = 0, 1, 2 or 3 in a(n) = a(n-1) + n + R and that for R >= 4 the recurrence is not quasi-periodic. For R = 0, 1, 2 all starting values give a quasi-periodic sequence. The respective loop is (1, x) for R = 0, (1, x, 2x, 2) for R = 1 (this sequence), (1, x, 2x, x) or (2x, x) for R = 2. For R = 3 only some starting values converge to a 6-loop (4x+2, 2x+1, 3x+6, x+2, 2x+9, 3x+17). - _Ctibor O. Zizka_, Oct 27 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Dec 16 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004475 | OEIS-A133144

Start with n and repeatedly apply the powerback map of A133048. Sequence gives number of steps to the point where the next number would be one that has appeared before.

It is conjectured that every number eventually reaches a fixed point (see A131571) or the cycle of length 2 given by (175 <-> 78125).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. H. Conway_ and _N. J. A. Sloane_, Jan 01 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004476 | OEIS-A133154

a(n) is the smallest m<=p-1 such that p-1 is the only value of j in 1<=j<=2p for which m^j+j==0 (mod p), where p is the n-th prime.

Andrew Granville, based on submitter's analysis of the data in A131685, made the following conjecture: "For some n with 1<=n<=p-1, there does not exist a value of j, with 1 <= j <= 2p, other than j=p-1, for which n^j+j == 0 (mod p)." Max Alekseyev's calculations confirm that the conjecture is true for the primes between 5 and 10^5. The sequence consists of the first such "n" (referred to as "m" in this sequence's definition) for each prime. a(n)=0 means that there is no corresponding m; this occurs at n=1 (p=2), n=2 (p=3), and n=3 (p=5), and at no other primes p<10^5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander R. Povolotsky_, Oct 08 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004477 | OEIS-A133158

Binomial transform of A126568, second binomial transform of A026641.

Conjecture: 2*n*a(n) + (-19*n+12)*a(n-1) + 6*(8*n-11)*a(n-2) + 36*(-n+2)*a(n-3) = 0. - _R. J. Mathar_, Jun 30 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Oct 08 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004478 | OEIS-A133184

Let k'' be the result of applying the powertrain map of A133500 twice to k. Sequence gives conjectured list of all possible values of k''.

Let k'' be the result of applying the powertrain map of A133500 twice to k. Sequence gives conjectured list of all possible values of k''.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. H. Conway_ and _N. J. A. Sloane_, Jan 01 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004479 | OEIS-A133185

Conjectured list of numbers that are 7-smooth (A002473) but are not in A133184.

Conjectured list of numbers that are 7-smooth (A002473) but are not in A133184.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. H. Conway_ and _N. J. A. Sloane_, Jan 01 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004480 | OEIS-A133205

Fully multiplicative with a(p) = p*(p+1)/2 for prime p.

Conjecture: Sum_{k=1..n} a(k) = O(n^3/log(n)). - _Vaclav Kotesovec_, Jan 28 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Oct 10 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004481 | OEIS-A133212

a(n) = 4*a(n-1) - 6*a(n-2) + 4*a(n-3), n > 3; a(0) = 1, a(1) = 4, a(2) = 12, a(3) = 32.

Conjecture: a(n) = 2*A038503(n+3) if n > 0. - _R. J. Mathar_, Oct 23 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Oct 11 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004482 | OEIS-A133219

Smallest composite integer in base n which remains composite after altering any one or two digits.

Changing the most significant digit to 0 is allowed. The problem (base 10) was posed by W. Sierpinski, published in 1977. There are an infinite number of solutions if a certain Erdos conjecture on congruences is true. a(2) through a(9) are proved minimal, a(10) has not yet been proved minimal.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Oct 11 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004483 | OEIS-A133223

Sum of digits of primes (A007605), sorted and with duplicates removed.

Conjecture: for s > 10 and not a multiple of 3, there exists a prime with digit-sum s consisting only of the digits 2 and 3 (cf. A137269). This conjecture has been verified for s <= 2995. Conjecture: for s > 18 and not a multiple of 3, there exists a prime with digit-sum s consisting only of the digits 3 and 4. This conjecture has been verified for s <= 1345. Conjecture: for s > 90 and not a multiple of 3, there exists a prime with digit-sum s consisting only of the digits 8 and 9. This conjecture has been verified for s <= 8995. Conjecture: for 0 < a < b < 10, gcd(a, b) = 1 and ab not a multiple of 10, if s > 90 and s is not a multiple of 3, then there exists a prime with digit-sum s consisting only of the digits a and b. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Dec 19 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004484 | OEIS-A133266

a(1) = 30; for n >= 2, choose smallest a(n) so that no sum of 2 or more terms equals a prime.

a(n+1) = a(n) + 30 for n >= 7 (conjectured). - _Chai Wah Wu_, Feb 15 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jan 01 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004485 | OEIS-A133297

a(n) = n! * Sum_{k=1..n} (-1)^(k + 1) * n^(n - k - 1)/(n - k)!.

Conjecture: a(n) = (n-1)!*(Sum_{k >= 0} (-1)^k * n^(n+k)/(n+k)! - (-1/e)^n) for n >= 1. Cf. A000435. - _Peter Bala_, Jul 23 2021 This conjecture is true. Let "gamma" be the lower incomplete gamma function: gamma(n, x) = (n-1)! (1 - exp(-x)*Sum_{k = 0..n-1} x^k/k!), then we can get the upper incomplete gamma function Gamma(n, x) = gamma(n, oo) - gamma(n, x). By inserting according the formula below, we will obtain the formula from Peter Bala.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Oct 17 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004486 | OEIS-A133305

a(n) = (1/n)*Sum_{i=0..n-1} C(n,i)*C(n,i+1)*4^i*5^(n-i), a(0) = 1.

Conjecture: (n+1)*a(n) + 9*(-2*n+1)*a(n-1) + (n-2)*a(n-2) = 0. - _R. J. Mathar_, May 23 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Oct 18 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004487 | OEIS-A133306

a(n) = (1/n)*Sum_{i=0..n-1} C(n,i)*C(n,i+1)*5^i*6^(n-i), a(0)=1.

Conjecture: (n+1)*a(n) + 11*(-2*n+1)*a(n-1) + (n-2)*a(n-2) = 0. - _R. J. Mathar_, May 23 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Oct 18 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004488 | OEIS-A133307

a(n) = (1/n)*Sum_{i=0..n-1} C(n,i)*C(n,i+1)*6^i*7^(n-i), a(0)=1.

Conjecture: (n+1)*a(n) + 13*(-2*n+1)*a(n-1) + (n-2)*a(n-2) = 0. - _R. J. Mathar_, May 23 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Oct 18 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004489 | OEIS-A133308

a(n) = (1/n)*Sum_{i=0..n-1} C(n,i)*C(n,i+1)*7^i*8^(n-i), a(0)=1.

Conjecture: (n+1)*a(n) + 15*(-2*n+1)*a(n-1) + (n-2)*a(n-2) = 0. - _R. J. Mathar_, May 23 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Oct 18 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004490 | OEIS-A133361

Multiply by 9 and reverse.

Conjecture: a(n)^(1/n) tends to 10. - _Vaclav Kotesovec_, Jan 03 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | Rachit Agrawal (rachit_agrawal(AT)daiict.ac.in), Oct 26 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004491 | OEIS-A133411

Smallest highly composite number of the form k*a(n-1) where k is an integer greater than 1.

Conjecture: subsequence of A019505.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Nov 25 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004492 | OEIS-A133470

Numbers k > 1 for which floor(b(k)) = floor(b(k-1)), where b(m) = Sum_{j=1..m} (j/(j+2)).

I conjecture that lim_{n->infinity} a(n)/a(n-1) = sqrt(e). For integers not in the sequence, b(m) = 1 + b(m-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Philippe LALLOUET (philip.lallouet(AT)orange.fr), Nov 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004493 | OEIS-A133491

Order of the symmetry group of the (in some cases conjectural) minimal-energy configuration of n identical charged particles confined to the surface of a sphere.

Order of the symmetry group of the (in some cases conjectural) minimal-energy configuration of n identical charged particles confined to the surface of a sphere.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Keenan Pepper_, Nov 30 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004494 | OEIS-A133501

Number of steps for "powertrain" operation to converge when started at n.

It is conjectured that every number converges to a fixed-point.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. H. Conway_ and _N. J. A. Sloane_, Dec 03 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004495 | OEIS-A133552

Number of length n binary sequences with at most 4 of every adjacent 6 bits set.

Conjectures from _Colin Barker_, Feb 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 24 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004496 | OEIS-A133568

Triangle: row n is least sequence of n positive integers such that A131655(n) distinct rational numbers are generated only from them using only +, -, * and / and each number in the row no more than once in a given expression.

The list of rational numbers generated by row 5 does not contain a 1 or 0 and therefore it still might be possible to construct row 6 by using the numbers from row 5. Futhermore, one might conjecture that all subsequent rows fully contain the previous one. - _Mikelis Emils Mikelsons_, Sep 01 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rick L. Shepherd_, Sep 16 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004497 | OEIS-A133575

Table, read by rows, giving the number of vertices possible in 2 X n nondegenerate classical transportation polytopes.

This paper discusses properties of the graphs of 2-way and 3-way transportation polytopes, in particular, their possible numbers of vertices and their diameters. Our main results include a quadratic bound on the diameter of axial 3-way transportation polytopes and a catalog of non-degenerate transportation polytopes of small sizes. The catalog disproves five conjectures about these polyhedra stated in the monograph by Yemelichev et al. (1984). It also allowed us to discover some new results. For example, we prove that the number of vertices of an m X n transportation polytope is a multiple of the greatest common divisor of m and n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Sep 17 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004498 | OEIS-A133587

Conjectured order of the symmetry group of the (numerically computed) least-perimeter cluster of n nonoverlapping circles.

Conjectured order of the symmetry group of the (numerically computed) least-perimeter cluster of n nonoverlapping circles.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Keenan Pepper_, Dec 27 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004499 | OEIS-A133602

The matrix-vector product A133080 * A000108.

Conjecture: n*(n-1)*(n-3)*(3*n-4)*a(n) -8*(n-1)*(2*n-5)*a(n-1) -4*(n-2)*(3*n-1)*(2*n-5)*(2*n-7)*a(n-2)=0. - _R. J. Mathar_, Jun 20 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Sep 18 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004500 | OEIS-A133603

The matrix-vector product A133566 * A000108.

Conjecture: n*(n-2)*(3*n-1)*(n+1)*a(n) -8*n*(2*n-3)*a(n-1) -4*(n-1)*(3*n+2)*(

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Sep 18 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004501 | OEIS-A133604

Elements of A005282 that are also the sum of a pair of not necessarily distinct elements of A005282.

Conjecture: 2, 4 and 8 are the only terms n such that n = 2*A005282(k) for some k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Sep 18 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004502 | OEIS-A133697

a(n) = smallest number k such that P(k)/P(k+1) > P(k+1)/P(k+2) > ... > P(k+n+1)/P(k+n+2), where P(k) = k-th prime = A000040(k).

If 113 is, as conjectured, the last term of A124129, then P(a(n)) = A158939(n+2). Proof: Let x and y be the prime gaps following the prime p = P(j) > 113, so that P(j+1) = P(j) + x and P(j+2) = P(j) + x + y. The inequality P(j)/P(j+1) > P(j+1)/P(j+2) can be written as p/(p+x) > (p+x)/(p+x+y), which simplifies to y > x+x^2/p. By assumption, x^2 < p, so this holds if and only if y > x. So the condition P(j)/P(j+1) > P(j+1)/P(j+2) is equivalent to increasing prime gaps, P(j+2) - P(j+1) > P(j+1) - P(j). (In fact, since all prime gaps except the first are even, it is enough to assume the weaker conjecture that 7 is the only prime P(j) such that (P(j+1)-P(j))^2 >= 2*P(j).) - _Pontus von Brömssen_, Nov 19 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | Philippe LALLOUET (philip.lallouet(AT)orange.fr), Jan 04 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004503 | OEIS-A133710

Column l=3 of irregular triangle in A133709.

Conjectures from _Colin Barker_, Apr 03 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 30 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004504 | OEIS-A133722

Column 3 of triangle in A133721.

Conjectures from _Colin Barker_, Apr 03 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 30 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004505 | OEIS-A133745

Numbers n such that A133744(n) = 0.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Sep 24 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004506 | OEIS-A133832

Least number k > n such that the binary trinomial 1 + 2^n + 2^k is prime, or 0 if there is no such k.

Conjecture: a(n) is nonzero for all n. These binary trinomials can also be written as f*2^n+1, where f=2^m+1 for some m, which is reminiscent of the Sierpinski problem (see A076336). The conjecture is equivalent to no Sierpinski numbers of the form 2^m+1 existing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Sep 26 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004507 | OEIS-A133857

Numbers k such that (18^k - 1)/17 is prime.

With the discovery of a(7), G=0.54789, which is very close to the expected constant 0.56145948 mentioned in the Generalized Repunit Conjecture below. - _Paul Bourdelais_, Dec 08 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Sep 28 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004508 | OEIS-A133901

Numbers in A128921 whose square is not a palindrome.

Conjecture: Sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_ and _Zak Seidov_, Sep 29 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004509 | OEIS-A133907

Least prime number p such that binomial(n+p, p) mod p = 1.

Conjecture: a(n) is the smallest prime p such that Sum_{k=1..n} k^(p-1) == n (mod p). Thus a(n) >= A317358(n). - _Thomas Ordowski_, Jul 29 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hieronymus Fischer_, Oct 20 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004510 | OEIS-A133910

Period numbers of A133900 divided by n^2.

I conjecture that a(n) is divisible by n iff n is in {1} U A051283. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hieronymus Fischer_, Oct 20 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004511 | OEIS-A133954

Difference between the numbers of nonnegative evil and odious multiples of p_n less than 2^p_n, where p_n = n-th prime.

A comparison of Gerbicz's calculations up to a(46) with A001122 and A139035 shows that one can conjecture that the converse statements are true as well.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 30 2007, Dec 17 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004512 | OEIS-A134002

Positive integers n such that n(n+5)=a(a+5)+b(b+5) is solvable in positive integers.

Conjecture. If n a positive integer not a term of this sequence, then n^2+(n+5)^2 is prime. (This has been verified up to n=500.) Examples. For n=1,2,3,4,6,7, n^2+(n+5)^2 is 37, 53,73, 97, 157 and 193, each of which is prime. See A134003 for the complement of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Oct 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004513 | OEIS-A134003

Positive integers n for which n^2+(n+5)^2 is prime.

Conjecture. If n is in this sequence then n(n+5)=a(a+5)+b(b+5) is not solvable in integers. (This has been verified up to n=500.) See A134002 for the complement of this sequence (in the positive integers).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Oct 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004514 | OEIS-A134016

Inverse score permutation of an Fibonacci -anti-Fibonacci zero sum game of 2 X 2 matrices.

Conjecture: a(n) = 4*a(n-2)+a(n-4). G.f.: x*(x^3-2*x^2-6*x-2)/(x^4+4*x^2-1). [_Colin Barker_, Nov 01 2012]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Jan 10 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004515 | OEIS-A134119

a(n) = floor(n^2/10) - floor((n-1)^2/10).

The above conjectured g.f. is correct. - _Sela Fried_, Dec 08 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander R. Povolotsky_, Jan 12 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004516 | OEIS-A134143

Let T(n) = (p, p+2) denote the n-th pair of twin primes. Let S(n) = 2p+2 (see A054735). Then a(n) = number of ways of writing S(n) as S(i) + S(j) with i <= j < m.

It is conjectured that a(n) > 0 for n >= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 25 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004517 | OEIS-A134162

Let S(k) be the sequence s() defined by s(1) = k; for i > 1, s(i) = s(i-1) + gcd(s(i-1), i). Start with the list of positive integers and remove any k's for which S(k) merges with an S(m) with m < k. Each value k > 1 is conjectural.

Let S(k) be the sequence s() defined by s(1) = k; for i > 1, s(i) = s(i-1) + gcd(s(i-1), i). Start with the list of positive integers and remove any k's for which S(k) merges with an S(m) with m < k. Each value k > 1 is conjectural. For the given initial values k, it is conjectural that their sequences S(k) never merge. The S(k) have been checked to be distinct for 2^60 terms (see Rowland link), but it is possible that they merge later on.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Rowland_, Jan 29 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004518 | OEIS-A134195

Antidiagonal sums of square array A126885.

Conjecture: partial sums of A104879. - _Sean A. Irvine_, Jul 14 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Oct 12 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004519 | OEIS-A134341

Numbers whose fifth powers have a partition as a sum of fifth powers of four positive integers.

The paper by Lander and Parkin where they just give the first known counterexample to Euler's conjecture, 27^5 + 84^5 + 110^5 + 133^5 = 144^5, found using a CDC6600, is known as one of the shortest published proofs. - _M. F.

Hasler_, Mar 11 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Oct 21 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004520 | OEIS-A134347

A007318^(-1) * A134346.

Conjecture: T(n,k) = 2^(k+1)*binomial(n,k) for k < n; T(n,n) = 2^(n+1)-1. - _Knud Werner_, Jan 05 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Oct 21 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004521 | OEIS-A134357

Denominator of binomial(6*n-2,2*n)/(2*binomial(4*n-1,2*n)).

It is conjectured that binomial(6*n-2,2*n)/(2*binomial(4*n-1,2*n)) = A005156(n+1)/A005156(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 04 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004522 | OEIS-A134376

Numbers whose sum of prime factors (counted with multiplicity) is not prime.

Conjecture: a(n) ~ n. - _Charles R Greathouse IV_, Apr 28 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hieronymus Fischer_, Oct 23 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004523 | OEIS-A134389

A transform of floor((n+2)/2) with Hankel transform floor((n+2)/2)*(cos(Pi*n/2) + sin(Pi*n/2)).

Conjecture: (n-1)*a(n) - 2*n*a(n-1) + 4*(4-n)*a(n-2) + 8*(n-3)*a(n-3) = 0. - _R. J. Mathar_, Oct 25 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 23 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004524 | OEIS-A134424

Area under all paths in the first quadrant from (0,0) to (n,0) using steps U=(1,1), D=(1,-1), h=(1,0) and H=(2,0).

Conjecture D-finite with recurrence -(n+2)*(5*n-7)*a(n) -(n+1)*(5*n-127)*a(n-1) +(135*n^2-655*n-42)*a(n-2) +2*(5*n^2-275*n-108)*a(n-3) +(-725*n^2+4941*n-5734)*a(n-4) +(-235*n^2+1880*n-1173)*a(n-5) +(725*n^2-6659*n+12606)*a(n-6) +2*(5*n^2+195*n-1988)*a(n-7) +(-135*n^2+1505*n-3358)*a(n-8) -(5*n+87)*(n-9)*a(n-9) +(5*n-33)*(n-10)*a(n-10)=0. - _R. J. Mathar_, Jul 24 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Oct 25 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004525 | OEIS-A134449

Sum of even products minus sum of odd products of different pairs of numbers from 1 to n.

Conjectures from _Colin Barker_, Mar 20 2015: (Start) The above conjectures are true. - _Sela Fried_, Dec 08 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_ and _Giorgio Balzarotti_, Jan 31 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004526 | OEIS-A134460

First differences of A067186.

Conjecture: Except for the first term a(1)=2, this sequence contains no terms of the form 4m+2. (This has been confirmed for the first several thousand terms.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Oct 26 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004527 | OEIS-A134466

See A134457.

Note that a(n) = a(n-1) for n = [1,] 3, 6, 11, 20,... = A006127(2^k + k) (conjectured); this corresponds to the case where the string c (see A134457) satisfies c(n)="0".c(n-1) (pre-pended "0"). The next string c(n+1) is obtained by inserting "0" after the first "1" and adding 1 (except for c(4) where both of these operations are equivalent and only one must be performed). - _M. F. Hasler_, Dec 17 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_, Dec 17 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004528 | OEIS-A134543

Triangle read by rows: T(n,k) = Sum_{i=k..n} (i+1-k)*A134541(n,i).

Conjecture: row terms starting from the right tend to the triangular series.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Oct 31 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004529 | OEIS-A134546

Triangle read by rows: T(n, k) = Sum_{j=0..n} floor(j / k).

Conjecture: For T(n, k) n >= k^(3/2), there is at least one prime in the interval [T(n-1, k+1), T(n, k)]. - _Bob Selcoe_, Aug 21 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Oct 31 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004530 | OEIS-A134558

Array read by antidiagonals, a(n,k) = gamma(n+1,k)*e^k, where gamma(n,k) is the upper incomplete gamma function and e is the exponential constant 2.71828...

Conjecture: a(n,k) is the permanent of the n X n matrix with k+1 on the main diagonal and 1 elsewhere.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ross La Haye_, Jan 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004531 | OEIS-A134559

Triangle read by rows: T(n,k) is the sum of the divisors of n that are >= k.

Conjecture: let P(n) be the n-th antidiagonal product (P(n) = Product_{k=0..ceiling(n/2)-1} b_n(k)). Consecutive n satisfying two equations: gcd(P(n),n) = 1 and

gcd(P(n+1),n+1) = 2 are consecutive elements of A005383 (primes p such that (p+1)/2 are also primes, save A005383(1) = 3 and A005383(2) = 5). The conjecture is false if for any prime number p belonging to A005383 gcd(P(p),p) = p. The conjecture was checked for 2000 consecutive integers. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Oct 31 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004532 | OEIS-A134635

Row sums of triangle A134634.

Conjecture: (n+1)*a(n) +(-7*n+1)*a(n-1) +2*(7*n-8)*a(n-2) +4*(-2*n+5)*a(n-3)=0. - _R. J. Mathar_, May 30 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Nov 04 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004533 | OEIS-A134639

Conjectured number of numbers k such that sigma(k)/k = n.

Conjectured number of numbers k such that sigma(k)/k = n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Nov 05 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004534 | OEIS-A134641

Prime permutational numbers A134640.

Conjecture: This sequence is finite and complete. The conjecture is correct: in base 2k permutational numbers are divisible by 2k-1 and in base 2k+1 permutational numbers are divisible by k. Hence it suffices to check bases up to 3. - _Charles R Greathouse IV_, Dec 14 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Nov 05 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004535 | OEIS-A134650

Numbers that are the sum of two consecutive primes (i.e., in A001043) but are not the sum of two sums of two consecutive primes.

Conjectured to be finite, may be complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 25 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004536 | OEIS-A134664

Number of 3-stack sortable permutations on n letters.

Lim_{n->infinity} a(n)^(1/n) >= 9.4854... (a new rigorous lower bound). Lim_{n->infinity} = 9.69963634535... (conjecture). [Defant, Elvey Price, Guttmann, 2020] - _Vaclav Kotesovec_, Jun 12 2021, following a suggestion of _Anthony Guttmann_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Rowland_, Jan 25 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004537 | OEIS-A134675

Row sums of triangle A134674.

a(n) = A007434(n) + A001065(n). - Conjectured by _John Mason_ and proved by _Max Alekseyev_, Jan 07 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Nov 05 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004538 | OEIS-A134680

a(n) = length (or lifetime) of the meta-Fibonacci sequence {f(1) = ... = f(n) = 1; f(k)=f(k-f(k-1))+f(k-f(k-n))} if that sequence is only defined for finitely many terms, or 0 if that sequence is infinite.

The term a(2) = 0 is only conjectural - see A005185. a(4) = 0 is a theorem of Balamohan et al. (2007). - _N. J. A. Sloane_, Nov 07 2007, Apr 18 2014.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Nov 06 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004539 | OEIS-A134799

a(n) = 3^((3^n - 1)/2).

a(n) is conjectured to be one-third the reduced denominator of b(n) = (3/2)*b(n-1)*(1 - b(n-1)^2); b(0) = 1/3. - _Steven Finch_, Oct 08 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yasutoshi Kohmoto_, Jan 09 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004540 | OEIS-A134972

Decimal expansion of 2 divided by golden ratio = 2/phi = 4/(1 + sqrt(5)) = 2*(-1 + phi).

Conjecture: Equals Sum_{k>=0} A225701(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Nov 15 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004541 | OEIS-A135010

Triangle read by rows in which row n lists A000041(n-1) 1's followed by the list of juxtaposed lexicographically ordered partitions of n that do not contain 1 as a part.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Nov 17 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004542 | OEIS-A135051

Pyramid game person numbers that have integer solutions.

Conjecture: a(n) = 9 - 2*(-1)^n + 4*(-8+(-1)^n)*n + 32*n^2. Conjecture: G.f.: x*(7 + 72*x + 98*x^2 + 72*x^3 + 7*x^4)/((1-x)^3*(1+x)^2). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Jan 31 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004543 | OEIS-A135052

Expansion of g.f.: (1-2*x-sqrt(1-4*x+8*x^3-4*x^4))/(2*x^2*(1-x)).

Conjecture: (n+2)*a(n) +(-5*n-4)*a(n-1) +2*(2*n+1)*a(n-2) +4*(2*n-5)*a(n-3) +12*(-n+3)*a(n-4) +4*(n-4)*a(n-5)=0. - _R. J. Mathar_, Apr 19 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 15 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004544 | OEIS-A135060

a(n) is the smallest number m for which none of the first n multiples of m has twice as many divisors as m.

Conjecture: every number in this sequence is also in A002182. [_J. Lowell_ disproved this conjecture at a(24) = 48188059920. - _Ray Chandler_] The conjecture that every term is a multiple of the preceding term is disproved at n = 15; a(15) = 1081080, which is not a multiple of a(14) = 720720. - _J. Lowell_, Jun 06 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Feb 11 2008, Jul 08 2008, Jul 14 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004545 | OEIS-A135064

Numbers n such that the quintic polynomial x^5 - 10*n*x^2 - 24*n has Galois group A_5 over rationals.

If this agreement is provable then of course it provides recurrences, generating functions, etc., for this sequence. - _N. J. A. Sloane_, Nov 24 2007 However, at present this is only a conjecture, and should not be used as the basis for formulas or computer programs. - _N. J. A. Sloane_, Mar 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Nov 15 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004546 | OEIS-A135096

Decimal expansion of certain constant L.

Conjecture: With any real number c in the interval [L,U] there exists for each natural number n>2 a natural number m and a prime number p so that n=floor(m^c)+p. L is the lower bound of c. U is the upper bound of c. L = 0.57812965263056281372405775798 03... U = 1.36521238896851872972933866767 77... (A135097).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner D. Sand_, Nov 25 2007, Dec 01 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004547 | OEIS-A135231

Row sums of triangle A135230.

Conjectures from _Chai Wah Wu_, Aug 31 2023: (Start) Equivalent conjecture: a(n) = A128209(n+1) for n>0. - _R. J. Mathar_, Mar 03 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Nov 23 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004548 | OEIS-A135238

Numbers n such that phi(sigma(n)) = reversal(n).

If both numbers 10^m-3 & 5*10^(m-1)-1 are primes and n=3*(10^m-3) then phi(sigma(n))=reversal(n), namely n is in the sequence (the proof is easy). Conjecture: n=2991 is the only such term of the sequence. there is no further term up to 35*10^7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Dec 26 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004549 | OEIS-A135255

Partial sums of A102659 read as decimal integers.

The initial 5 ones of 11111843 of a(32) leads me to conjecture that some element of this sequence is, base 10, a concatenation of the digits (1,2). Could there be an element which is also one of the Lyndon words in the underlying A102659?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Nov 30 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004550 | OEIS-A135303

a(n) = phi(2*n+1)/(2*A003558(n)), where phi = A000010.

Conjecture relating to primes with more than one coach: The combined set of integers in the top rows of all coaches of these primes is composed of a permutation of the first q odd integers, where prime p = (4q-1) or (4q+1), (q > 0). Example: As shown for 17, this prime has two coaches with the top rows [1] and [3, 7, 5]. 43 has three coaches with q = 11. The top rows are [1, 21, 11], [3, 5, 19], [7, 9, 17, 13, 15]. The comment of Sep 08 2012 in A216371 applies to primes with one coach, in which case "all coaches" is reduced to one and the set of q odd integers is in the top row of the coach. - _Gary W. Adamson_, Sep 10 2012 Conjecture [Carl Schick]: If 2*n+1 is prime, then these are the number of distinct cycles of f(k) = |(2*n+1) - 2*k| beginning at an odd number 0 < k < 2*n. - _Jonathan Skowera_, Aug 03 2013 [See also the Brändli and Beyne link, eq. (2). - _Wolfdieter Lang_, Feb 08 2020] Conjecture of Aug 03 2013 proved by Jean Pedersen. By way of example, take A003558(5) = 11, such that

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 05 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004551 | OEIS-A135313

Triangle of numbers T(n,k) (n>=0, n>=k>=0) of transitive reflexive early confluent binary relations R on n labeled elements where k=max_{x}(|{y : xRy}|), read by rows.

Conjecture: For fixed k>=0, A135313(n+k,n) ~ n! * n^(2*k) / (2^(k+1) * k! * log(2)^(n+k+1)). - _Vaclav Kotesovec_, Nov 20 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Dec 05 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004552 | OEIS-A135334

Number of Dyck paths of semilength n having no UDDU's starting at level 1.

Conjecture: 2*(n+1)*a(n) + 2*(1-5n)*a(n-1) + 3*(3n-1)*a(n-2) + 2*(1-2n)*a(n-3) = 0. - _R. J. Mathar_, Dec 18 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 07 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004553 | OEIS-A135335

Number of Dyck paths of semilength n having no DDUU's starting at level 2.

Conjecture: n*a(n) +(-7*n+6)*a(n-1) +2*(8*n-9)*a(n-2) +6*(-3*n+4)*a(n-3) +3*(3*n-4)*a(n-4) +2*(-2*n+3)*a(n-5)=0. - _R. J. Mathar_, Jun 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 07 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004554 | OEIS-A135339

Number of Dyck paths of semilength n having no DUDU's starting at level 1.

Conjecture: 2*(n + 1)*a(n) + (-5*n + 1)*a(n-1) + (-11*n + 28)*a(n-2) + 2*(-2*n + 7)*a(n-3) = 0 for n >= 4. - _R. J. Mathar_, Oct 20 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 07 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004555 | OEIS-A135401

a(n) = number of permutations (p(1),p(2),p(3),...,p(n)) of (1,2,3,...n) each of which is its own inverse and is such that p(k) = n + 1 - p(n+1-k) for all k in the range 1 <= k <= n.

a(n) = A000898(floor(n/2)). - Conjectured by _Leroy Quet_, Jan 20 2008; proved by _Max Alekseyev_, Jan 21 2008: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Dec 11 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004556 | OEIS-A135404

Gessel sequence: the number of paths of length 2m in the plane, starting and ending at (0,1), with unit steps in the four directions (north, east, south, west) and staying in the region y > 0, x > -y.

According to Ira Gessel's student, Guoce Xin, Ira Gessel made his intriguing conjecture in 2001. On June 25, 2008, the Gessel Conjecture became the Kauers-Koutschan-Zeilberger theorem - see the link. - _Doron Zeilberger_, Jul 01 2008 The Ira Gessel Conjecture is that a(m)=16^m*(5/6)_m*(1/2)_m/ ((2)_m*(5/3)_m), where (a)_m:=a*(a+1)*...*(a+m-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Doron Zeilberger_, Dec 11 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004557 | OEIS-A135406

Sum of squares of gaps between consecutive semiprimes.

This is to semiprimes A001358 as A074741 is to primes A000040. What is the semiprime analog of D. R. Heath-Brown's conjecture: Sum_{prime(n)<=N} (prime(n)-prime(n-1))^2 ~ 2*N*log(N) and Marek Wolf's conjecture: Sum_{prime(n)<N} (prime(n)-prime(n-1))^2 = 2*N^2/pi(N) + error term(N), pi(N)=A000720(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Dec 09 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004558 | OEIS-A135414

a(1)=a(2)=1 and for n>=3, a(n) = n - a(a(n-2)).

Conjecture: A recursively built tree structure can be obtained from the sequence:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 17 2008, Feb 19 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004559 | OEIS-A135422

a(1) = 1, a(n) = Sum(k=1,n-1, a(k)^F(n-k)) where F(m) is the m-th Fibonacci number.

Conjecture: limit n->infinity Phi^(Phi^(n-c))/a(n) = 1 where c is a constant = 3.47348961175710091....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerald McGarvey_, Dec 12 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004560 | OEIS-A135470

Composite numbers n that divide 2 * sigma(n) - d(n) [that is, 2 * sum of divisors - number of divisors].

It is conjectured that there are no odd terms and that the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. K. Guy_, Feb 08 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004561 | OEIS-A135499

Numbers for which Sum_digits(odd positions) = Sum_digits(even positions).

Conjecture: this is a subsequence of A008593 (verified for the first 50 thousand terms). - _R. J. Mathar_, Feb 10 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_ and _Giorgio Balzarotti_, Feb 08 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004562 | OEIS-A135543

Record number of steps under iterations of "map n to n - (largest prime <= n)" (A064722) until reaching the limiting value 0 or 1. Also, places where A121561 reaches a new record.

Conjecture: there exist positive values of n for which a(n) != A175079(n) - 1. - _Jaroslav Krizek_, Feb 05 2010 a(n) = A175079(n) - 1 for n >= 1, i.e., the conjecture in the Comments is false. This follows from the result that A175078(n) = A121561(n-1) for n >= 2. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sergio Pimentel_, Feb 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004563 | OEIS-A135571

Positive integers that are the difference of two positive triangular numbers in an odd number of ways.

Conjecture. This sequence is just the sequence of positive integers that are either square, twice a square, or triangular, but not both square and triangular (A001110). (This has been verified for n up to 100000.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Feb 23 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004564 | OEIS-A135582

Expansion of 1/((1-x^2*c(x))(1-x-2x^2)) where c(x) is the g.f. of A000108.

Conjecture: (-n+1)*a(n) +6*(n-2)*a(n-1) +(-7*n+19)*a(n-2) +(-7*n+13)*a(n-3) +(13*n-31)*a(n-4) +2*(-n+4)*a(n-5) +4*(-2*n+5)*a(n-6)=0. - _R. J. Mathar_, Feb 23 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004565 | OEIS-A135594

a(n) = (1/2^n) * Sum_{i=0..n} (-1)^(n-i) * binomial(n,i) * A000364(i).

Conjecture: e.g.f. as a continued fraction: 2*exp(-t)/(2 - (1-exp(-4*t))/(2 - (1-exp(-8*t))/(2 - (1-exp(-12*t))/(2 - ...)))) = 1 + t^2/2! + 6*t^3/3! + 73*t^4/4! + ... Cf. A000657 and A005799. - _Peter Bala_, Dec 21 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Feb 25 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004566 | OEIS-A135611

Decimal expansion of sqrt(2) + sqrt(3).

Karl Popper considers whether this approximation to Pi might have been known to Plato, or even conjectured to be exact. - _Charles R Greathouse IV_, Apr 26 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 03 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004567 | OEIS-A135627

Perfect numbers minus 1.

Conjecture: every odd prime is a factor of at least one number in this sequence. - _J. Lowell_, Apr 17 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Nov 27 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004568 | OEIS-A135646

a(m, n) is the number of coprime pairs (i, j) with 1 <= i <= m, 1 <= j <= n; table of a(m, n) read by antidiagonals.

Conjecture: a(m, n) ~ mn - sum_1^m{ (i - phi(i)) (n / i) } = n sum_1^m{ phi(i) / i } ~ 6mn / pi^2 as m -> oo.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo van der Sanden_, Nov 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004569 | OEIS-A135659

a(n) = 24*n + 7.

Conjecture: All Mersenne Primes (A000668) > 3 are in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Nov 25 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004570 | OEIS-A135730

Number of steps to reach the minimum of the final cycle under iterations of the map A001281: x->3x-1 if x odd, x/2 otherwise.

It is conjectured that the trajectory of any odd number will eventually reach 1, 9, or 33, and so enter one of the loops (1,3), (9, 27, 13, 39, 19), or (33, 99, 49, 147, 73, 219, 109, 327, 163, 81, 243, 121, 363, 181, 543, 271, 135, 67). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Nov 26 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004571 | OEIS-A135736

Nearest integer to n*Sum_{k=1..n} 1/k = rounded expected coupon collection numbers.

Conjecture: a(n) = round(n*(log(n) + gamma) + 1/2) for n > 0, where gamma = A001620. - _Ilya Gutkovskiy_, Oct 31 2016 The conjecture is false: a(2416101) = 36905656 while round(2416101*(log(2416101) + gamma) + 1/2) = 36905657, with the unrounded numbers being 36905656.499999982... and 36905656.500000016.... Heuristically this should happen infinitely often. - _Charles R Greathouse IV_, Oct 31 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Nov 29 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004572 | OEIS-A135833

Number of Section I primes between 2^n and 2^(n+1). See A135832.

Comparing these numbers with A036378, the number of primes between 2^n and 2^(n+1), leads one to conjecture that the density of Section I primes is 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Nov 30 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004573 | OEIS-A135908

Clique number of commuting graph of symmetric group S_n.

Conjecture: a(n) = a(n-1) + 3*a(n-3) - 3*a(n-4) for n>7; g.f.: -x^3*(3*x^3 - 2*x^2 - x - 2) / ((x-1)*(3*x^3-1)). - _Colin Barker_, Jul 26 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 07 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004574 | OEIS-A135909

Clique number of commuting graph of alternating group A_n.

Conjecture: a(n) = a(n-1)+3*a(n-3)-3*a(n-4) for n>6. G.f.: -x^3*(x^6-x^5+2*x^3-x^2-x-2) / ((x-1)*(3*x^3-1)). - _Colin Barker_, Jul 26 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 07 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004575 | OEIS-A135920

O.g.f.: A(x) = Sum_{n>=0} x^n / Product_{k=0..n} (1 - k^2*x).

Conjecture: a(n) = A369527(n-1, 0) = A369595(n-1, 0) for n > 0 with a(0) = 1. - _Mikhail Kurkov_, Apr 25 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 06 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004576 | OEIS-A135925

G.f.: (x^2+x-1)*(1+2*x+S)*(1-5*x+2*x^2+(3*x-1)*S)^2*(-1+3*x+2*x^2+(1-x)*S), where S=sqrt(1-4*x).

Conjecture D-finite with recurrence: +4*(n-1)*(7678757997599*n-100709801435251)*a(n) +(-291262905844111*n^2+4745710679447853*n-1293539319282124)*a(n-1) +3*(268523622217993*n^2-4978734571371597*n+20251709258114720)*a(n-2) +2*(-196528865827703*n^2+4147266222561252*n-21888137659762186)*a(n-3) -12*(22463548140013*n-247087251730058)*(2*n-25)*a(n-4)=0. - _R. J. Mathar_, Jan 23 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 09 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004577 | OEIS-A135936

Irregular triangle read by rows: row n gives coefficients of Boubaker polynomial B_n(x) in order of decreasing exponents (another version).

Conjectures from _Thomas Baruchel_, Jun 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 09 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004578 | OEIS-A135953

(Nonprime Fibonacci numbers with prime indices) that have exactly 2 prime factors.

Conjecture: All numbers in this sequence are products of two sums of two squares, e.g. 4181 = 37*113 = (1^2+6^2)*(7^2+8^2), 1346269 = 557*2417 = (14^2+19^2)*(4^2+49^2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Dec 08 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004579 | OEIS-A135954

Nonprime Fibonacci numbers with prime indices (A050937) that have exactly 3 prime factors.

Conjecture: All numbers in this sequence are products of three sums of two squares.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Dec 08 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004580 | OEIS-A135955

(Nonprime Fibonacci numbers with prime indices, A050937) which have exactly 4 prime factors.

Conjecture: All numbers in this sequence are products of four sums of two squares.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Dec 08 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004581 | OEIS-A135956

Members of A050937 (nonprime Fibonacci numbers with prime index) with 5 or more distinct prime factors.

Conjecture: all numbers in this sequence are product of 5 or more sum of two squares

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Dec 08 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004582 | OEIS-A136032

Number of prime factors (with multiplicity) of Mersenne composites (A065341).

If the conjecture that all Mersenne composites are squarefree is true, then this sequence is identical to A135975. - _Felix Fröhlich_, Aug 24 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Dec 11 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004583 | OEIS-A136050

Sum of digits of product of twin primes A037074.

Conjecture: except for the initial term, each term is one less than a multiple of 9. - _Harvey P. Dale_, Dec 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Dec 12 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004584 | OEIS-A136092

Bisection of A138540.

a(n) = Sum_{j=0..n+1} 180*(2n)!*(2j+2)!*(2n-3j+2)/(j!*(j+1)!*(j+2)!*(j+4)!*(n-j+1)!*(n-j+3)!) (conjectured). - _Michael D. Weiner_, May 11 2016 36*(n-1)*(2*n-1)*(2*n-3)*a(n-2)-4*(2*n-1)*(5*n^2+22*n+15)*a(n-1)+(n+3)*(n+5)*(n+6)*a(n) = 0 (conjectured). - _Michael D. Weiner_, May 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 25 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004585 | OEIS-A136097

a(n) = A135951(n) /[(2^(n+1)-1) * 2^(n*(n-1)/2)].

Conjecture: the n-th central term of the matrix inverse of the triangle of Gaussian binomial coefficients in q is divisible by [(q^(n+1)-1)/(q-1) * q^(n*(n-1)/2)] for n>=0 and integer q > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 13 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004586 | OEIS-A136117

Pentagonal numbers (A000326) which are the sum of 2 other positive pentagonal numbers.

It is conjectured that every integer and hence every pentagonal number, greater than 33066, hence greater than A000326(149) = 33227, can be represented as the sum of three pentagonal numbers. - _Jonathan Vos Post_, Dec 18 2007

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Dec 15 2007; corrected Dec 25 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004587 | OEIS-A136119

Limiting sequence when we start with the positive integers (A000027) and delete in step n >= 1 the term at position n + a(n).

Apparently a(n) = A001953(n-1)+1 = floor((n-1/2)*sqrt(2))+1 (confirmed for n < 20000) and a(n+1) - a(n) = A001030(n). From the definitions these conjectures are by no means obvious. Can they be proved? - _Klaus Brockhaus_, Apr 15 2008 [For an affirmative answer, see the Cloitre link.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Mar 16 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004588 | OEIS-A136162

List of prime quadruplets {p, p+2, p+6, p+8}.

{11, 13, 17, 19} is the only prime quadruplet {p, p+2, p+6, p+8} of the form {Q-4, Q-2, Q+2, Q+4} where Q is a product of a pair of twin primes {q, q+2} (for prime q = 3) because numbers Q-2 and Q+4 are for q>3 composites of the form 3*(12*k^2-1) and 3*(12*k^2+1) respectively (k is an integer). Conjecture: {11, 13, 17, 19} is the only prime quadruplet {p, p+2, p+6, p+8} of the form {q*(nextprime(q))-4, q*(nextprime(q))-2, q*(nextprime(q))+2, q*(nextprime(q))+4} where q is a prime (for prime q = 3). - _Jaroslav Krizek_, Jul 07 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Harry J. Smith_, Dec 17 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004589 | OEIS-A136191

Primes p such that 2p-3 and 2p+3 are both prime (A092110), with last decimal being 3.

Theorem: If in the triple (2n-3,n,2n+3) all numbers are primes then n=5 or the decimal representation of n ends in 3 or 7. Proof: Consider Q=(2n-3)n(2n+3), by hypothesis factorized into primes. If n is prime, n=10k+r with r=1,3,7 or 9. We want to exclude r=1 and r=9. Case n=10k+1. Then Q=5(-1+6k+240k^2+800k^3) and 5 is a factor; thus 2n-3=5 or n=5 or 2n+1=5 : this means n=4 (not prime); or n=5 (included); or n=2 (impossible, because 2n-3=1). Case n=10k+9. Then Q=5(567+1926k+2160k^2+800k^3) and 5 is a factor; the arguments, for the previous case, also hold.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carlos Alves_, Dec 20 2007 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004590 | OEIS-A136269

Floor of sum of the first 10^n cube roots.

Conjecture: For n = 3k, a(n) = (3/4)*10^(4n/3)+5*10^(n/3-1)-1. The conjecture is true (see links) - _Sela Fried_, Oct 02 2024.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Mar 19 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004591 | OEIS-A136289

Start with three pennies touching each other on a tabletop. In each generation, add pennies subject to the rule that a penny can be placed only when (at least) two pennies are already in position to determine its position; sequence gives number of pennies added at generation n.

The conjectured formulae are correct: Looking at Knecht's diagram one can see that a line of three numbers in any direction must be a permutation of {1,2,3} by applying the triple rule three times to one of the adjacent pairs, like so: Conjecture: a(n) = a(n-3) + 6, implying g.f. 3*(1+x^2)/((1-x)^2*(1+x+x^2)). - _R. J. Mathar_, Apr 15 2008 Conjecture: a(n) = 2n + 1 + ((n+2) mod 3). - _Wesley Ivan Hurt_, Jul 07 2013 Conjecture: a(n) = 3*floor(2*n/3) + 3. - _Jon E. Schoenfield_, Jul 30 2015 Conjecture: a(n) = 2*(n+1+sin(2*(n+1)*Pi/3)/sqrt(3)). - _Wesley Ivan Hurt_, Sep 27 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Mallows_, Apr 13 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004592 | OEIS-A136301

Frequency of occurrence for each possible "probability of derangement" for a Secret Santa drawing in which each person draws a name in sequence and the only person who does not draw someone else's name is the one who draws the final name.

Conjecture: Assume the table represented as in the Example section. Then row 2^n is row n + 1 of A371761. - _Peter Luschny_, Apr 10 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Parsonnet_, Mar 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004593 | OEIS-A136304

Expansion of g.f. (1-z)^2*(1-sqrt(1-4*z))/(2*z*(1 - 3*z + 2*z^2 - z^3)).

Conjecture: (n+1)*a(n) + (-8*n+1)*a(n-1) + 3*(7*n-8)*a(n-2) + (-23*n+49)*a(n-3) + (13*n-32)*a(n-4) + 2*(-2*n+7)*a(n-5) = 0. - _R. J. Mathar_, Feb 29 2016

Conjecture verified using differential equation -z^3 + 3*z^2 - 3*z + 1 + (6*z^5 - 20*z^4 + 20*z^3 - 18*z^2 + 7*z - 1)*g(z) + (4*z^6 - 13*z^5 + 23*z^4 - 21*z^3 + 8*z^2 - z)*g'(z) satisfied by the G.f. - _Robert Israel_, Feb 23 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Mar 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004594 | OEIS-A136404

Square numbers with more divisors than any smaller square number.

Conjecture: square roots of the terms of this sequence are the same terms as A126098.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Mar 30 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004595 | OEIS-A136433

a(n+2) = a(n+1)*(n mod 3 + 1) + (n mod 2 + 1), a(1) = 11.

Conjecture: a(n)=6*a(n-3)+a(n-6)-6*a(n-9). [_R. J. Mathar_, Oct 30 2008]

猜想/问题提及段 | historical_or_current_mention_unreviewed | Herman Jamke (hermanjamke(AT)fastmail.fm), Apr 01 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004596 | OEIS-A136510

Smallest k>1 such that in binary representation n is contained in n^k.

conjecture: a(n) is defined for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jan 03 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004597 | OEIS-A136564

Array read by rows: T(n,k) is the number of directed multigraphs with loops with n arcs, k vertices, and no vertex of degree 0.

T(n,2n-k) = A144047(k) for n large enough (conjecturally, n >= 2k is enough).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Jubin_, Apr 14 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004598 | OEIS-A136576

Series reversion of x*c(x)/(1 - 2*x), c(x) the g.f. of A000108.

Hankel transform of a(n+1) is A136577 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 08 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004599 | OEIS-A136612

a(n) = ((prime(n+3) + prime(n+1)) - (prime(n+2) + prime(n))).

a(n) is the sum of two prime gaps, thus a(n) >= 4 for n > 1. Conjecturally a(n) << log^2 n (probably with constant around 2). - _Charles R Greathouse IV_, Aug 25 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Odimar Fabeny_, Apr 14 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004600 | OEIS-A136746

G.f.: (z^12+1-z^11-z^10+z^8-z^6+z^5-z^3+z)/((z+1)*(z-1)^2).

Simon Plouffe conjectured in 1992 that this is a g.f. for A003003 - it is not.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 07 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004601 | OEIS-A136774

n! never ends in this many 0's in base 16.

Conjecture: a(n) ~ 16n. This holds with probability 1 in a random model. - _Charles R Greathouse IV_, Sep 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carl R. White_, Jan 21 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004602 | OEIS-A136859

Numbers k such that k and k^2 use only the digits 0, 1, 4 and 6.

Are the formulas conjectured or proved? For example, the analogous sequence for {0,1,2,4} contains the sporadic solution 1010000104010000101. Clearly, if a(n) is in the sequence then 10*a(n) is also in the sequence. Is there any term that is not 0, 1, or 4 times a power of 10? - _M. F. Hasler_, Jan 26 2016 Answer: the formulas were merely conjectures. It appears that it is an open question as to whether there is any other type of term. - _N. J. A. Sloane_, Jan 29 2016 Conjectures from _Philippe Deléham_, Mar 11 2014: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jonathan Wellons (wellons(AT)gmail.com), Jan 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004603 | OEIS-A136863

Numbers k such that k and k^2 use only the digits 0, 1, 4 and 7.

Conjecture: the only terms not ending in 0 are 1 and 1071. Thus for n >= 5, a(n) = 10^((n+1)/2) if n is odd, 1071 * 10^((n-6)/2) if n is even.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jonathan Wellons (wellons(AT)gmail.com), Jan 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004604 | OEIS-A136958

Numbers k such that k and k^2 use only the digits 0, 4, 7 and 8.

Conjectures: (1) the digit 4 never appears in the terms of the sequence (although it does appear in the digits of the squares of the terms of the sequence), and (2) all nonzero terms are of the form 88 * 10^m or 878 * 10^m. - _Harvey P. Dale_, Oct 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jonathan Wellons (wellons(AT)gmail.com), Jan 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004605 | OEIS-A136967

Numbers k such that k and k^2 use only the digits 1, 2, 3 and 4.

No further terms up to 444444444444. Conjecture: these are all the terms. - _Robert Israel_, Mar 01 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jonathan Wellons (wellons(AT)gmail.com), Jan 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004606 | OEIS-A137051

Numbers k such that k and k^2 use only the digits 1, 4, 5, 8 and 9.

Conjecture: The only squares present in the sequence are 1, 9, and 441. - _Stefano Spezia_, Apr 18 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jonathan Wellons (wellons(AT)gmail.com), Jan 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004607 | OEIS-A137098

Numbers k such that k and k^2 use only the digits 2, 4, 5, 7 and 9.

Conjecture: the last digit of all terms except the first two is 7. - _Harvey P. Dale_, Jan 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jonathan Wellons (wellons(AT)gmail.com), Jan 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004608 | OEIS-A137225

Triangle T(k,q) of minimal q-Niven numbers: smallest number such that the sum of its digits in base q equals k, 2<=q<=k+1.

T(k,2)=A000225(k). T(k,k+1)=2k-1. Conjecture: T(k,3)=A062318(k), verified up to k=23.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Mar 07 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004609 | OEIS-A137248

The smallest prime with digit sum n and maximum digit product (or 0 if no such prime exists).

Conjecture: If n > 4 and a(n) > 0, then a(n) consists only of the digits 2, 3 and 4. - _Chai Wah Wu_, Dec 17 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Mar 10 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004610 | OEIS-A137264

Prime number gaps read modulo 3.

Conjecture: The only digit that is repeated in the sequence is 0 except for n=2 and n=3 where 2 repeats. So 1 may be followed by 2 or 0; 2 may be followed by 1 or 0; 0 may be followed by 0 or 1 or 2. this has been confirmed for the first million prime gaps. The conjecture is true, because any three numbers whose differences are (1, 1) or (2, 2) will form a complete residue system modulo 3, and hence one of them will be a multiple of 3. - _Karl W. Heuer_, Mar 16 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | Noel H. Patson (n.patson(AT)cqu.edu.au), Mar 12 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004611 | OEIS-A137281

Numbers k such that T(k) is not divisible by T(i), 1 < i < k, where T(k) = k-th triangular number A000217(k).

Conjecture: a(n) ~ C * n * log(n) for some constant C, in analogy with the prime number theorem (see A000040). - _Harry Richman_, Mar 05 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Mar 14 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004612 | OEIS-A137284

a(0)=1 and a(n) for n > 0 equals the minimal positive integer such that addition of 2^(-a(n)) to Sum_{k = 0,1,...,n-1} 2^(-a(k)) changes only trailing zeros in its decimal representation.

a(n+1) = ceiling(a(n)*log_2(10)) = ceiling(a(n)*A020862). - Conjectured by _R. J. Mathar_, proved by _Max Alekseyev_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_ and _Giorgio Balzarotti_, Mar 14 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004613 | OEIS-A137295

P-positions in sparse Atomic Wari, octal game .37.

The sequence is only conjectured to be finite as of the publication in 1996, but there are definitely no more terms under 200000, so the conjecture seems extremely likely to be true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joshua Zucker_, Mar 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004614 | OEIS-A137432

Number of ways to place n^2 nonattacking kings on a 2n X 2n cylindrical chessboard.

Conjecture: limit of a(n+1)/(n*a(n)) as n->infinity is e.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Aug 31 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004615 | OEIS-A137560

Let f(z) = z^2 + c, then row k lists the expansion of the n-fold composition f(f(...f(0)...)) in rising powers of c.

The sequence of polynomials is conjectured to tend to the Catalan numbers (A000108). - _Jon Perry_, Oct 31 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Apr 25 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004616 | OEIS-A137582

Numbers having no inner zeros in decimal representation of their factorial.

Conjecture: the sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jan 27 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004617 | OEIS-A137613

Omit the 1's from Rowland's sequence f(n) - f(n-1) = gcd(n,f(n-1)), where f(1) = 7.

a(n) = A168008(2*n+4) (conjectured). - _Jon Maiga_, May 20 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Jan 29 2008, Jan 30 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004618 | OEIS-A137675

Prime numbers p such that p +- a, p +- b, p +- c and p +- d are prime numbers, where a, b, c and d are distinct positive integers with p > d, d > c, c > b and b > a.

Conjecture: includes all primes > 61. - _Robert Israel_, May 24 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Joseph Stephan Orlovsky_, Apr 27 2008; corrected May 02 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004619 | OEIS-A137715

Prime values of n for which n*2^k + 1 is composite for all positive integers k.

Note that the terms in A076336 are presently conjectural. - _Joerg Arndt_, Jun 29 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ant King_, Feb 09 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004620 | OEIS-A137727

Final digit of prime(n)*prime(n+1).

a(n) is 1, 3, 7, or 9 for n > 3. I conjecture that 1 and 9 appear 17/66 of the time and 3 and 7 appear 8/33 of the time. - _Charles R Greathouse IV_, Jan 03 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Feb 08 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004621 | OEIS-A137738

Coefficients of the polynomial giving the n-th diagonal of A137743 * n!, read as an upper right triangle.

For m>0, T(n,n+m+3) = sum(T(k,k+m+2), k=m+1..n+1) - (1/m!) Q[m](n), where Q[m] is a monic polynomial of degree <= m with integer coefficients (conjectured - see examples).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Mar 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004622 | OEIS-A137741

Number of different strings of length n+4 obtained from "123...n" by iteratively duplicating any substring.

a(n) = n^4 for n=1,2; a(n) = 1/24*(n+3)*(n^3+15*n^2+50*n-96) for n>2 (conjectured). G.f.: x*(x^6+x^5-8*x^4+x^3+16*x^2-11*x-1) / (x-1)^5 (conjectured). - _Colin Barker_, Nov 04 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 10 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004623 | OEIS-A137777

Triangular sequence of coefficients from the expansion of the derivative of the Bernoulli polynomial function: p(x,t) = t*exp(x*t)/(exp(t)-1); q(x,t) = p'(x,t) = dp(x,t)/dt.

Conjecture on row sums: Sum_{k=0..n+1} T(n,k) = 2*A129825(n+2). - _R. J. Mathar_, Jun 03 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_ and _Gary W. Adamson__, Apr 28 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004624 | OEIS-A137864

a(n) = n^4 - 10n^3 + 35n^2 - 48n + 23.

This sequence appears at first to be the sequence of odd numbers but then rapidly becomes something different altogether. It is a good example of why more than a few terms are needed to check a hypothesis.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Christopher Martin (christopher.j.martin(AT)gmail.com), Feb 17 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004625 | OEIS-A137876

a(n) = (nextprime(18n)-previousprime(18n))/2.

Note that a(n) cannot be 2 (because, for arbitrary number m, if (6*m-1) is prime then (6*m+3) is not, and similarly, if (6*m+1) is prime then (6*m-3) is not). I conjecture that all other values are possible and a(n) == 0 (mod 3) are (much) more abundant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Feb 19 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004626 | OEIS-A137996

Numbers which are not the sum of a triangular number and zero or a prime = 1 (mod 4).

Zhi-Wei SUN conjectured that n=216 is the only number not of the form n = p + k(k+1)/2 with p a prime or zero and more precisely that the numbers given in this sequence A137996(1..123) = 2,...,88956 are the only numbers which cannot be written in this form with p=0 or p=1 (mod 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Mar 24 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004627 | OEIS-A137997

Numbers which are not the sum of a triangular number and zero or a prime = 3 (mod 4).

Zhi-Wei SUN conjectured that n=216 is the only number not of the form n = p + k(k+1)/2 with p a prime or zero and more precisely that the numbers given in this sequence A137997(1..112) = 2,...,90441 are the only numbers which cannot be written in this form with p=0 or p=3 (mod 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Mar 24 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004628 | OEIS-A137998

Least k such that floor(16^n / 10^k) = 1, 2, 4 or 8 (mod 10), or zero if no such k exists.

This is related to A071071, i.e., the conjecture by J. Shallit that {1,2,4,8,65536} is a minimal set for powers of 2 written in base 10. This amounts to saying that any power of 2 contains either a digit among {1,2,4,8}, or 65536 as subword (obtained by deleting zero or more digits of the number). The present sequence gives the position of the first digit (starting with units) of these powers which is among {1,2,4,8}. It appears as if 16^4=65536 is the only power of 2 which does not contain any of these four digits. This would imply the conjecture (and be an even stronger statement).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Mar 26 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004629 | OEIS-A138000

a(n) is the least prime such that the subsets of { a(1), ..., a(n) } sum up to 2^n different values.

It can be seen that a(n) is always very close to Sum_{i=1..n-1} a(i). As a consequence, the sequence grows like a(n) ~ 2^(n-0.256...) and thus is not a counterexample to Erdős's conjecture mentioned in the cited paper.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 08 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004630 | OEIS-A138004

Odd numbers k for which all nontrivial cycles of the Ducci map on k-tuples modulo 2 have the same length.

It is conjectured that all terms of this sequence are prime numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Feb 26 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004631 | OEIS-A138099

Irregular triangle read by rows: T(n,k) = k + floor(n/2), 1 <= k <= ceiling(n/2).

Conjecture: a(n) = HookSum(T(n))-HookSum(T(n-1)), where HookSum(A) is the sum of all hooks across a tableau A, and T(n) are the spiraling sequence of tableaux:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, May 03 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004632 | OEIS-A138123

Antidiagonal sums of a triangle of coefficients of recurrences of the Fibonacci sequence.

Conjecture: The Fibonacci sequence F obeys all the recurrences: A000045(n)=F(n)= L(p)*F(n-p)-(-1)^p*F(n-2p), any p>0, L=A000204. [Proof: conjecture is equivalent to the existence of a g.f. of F with denominator 1-L(p)x^p+(-1)^p*x^(2p). Since 1-x-x^2 is known to be a denominator of such a g.f. of A000045, the conjecture is that 1-L(p)*x^p+(-1)^p*x^(2p) can be reduced to 1-x-x^2. One finds: {1-L(p)*x^p+(-1)^p*x^(2p)}/(1-x-x^2) = sum{n=0..p-1}F(n+1)x^n-sum{n=0..p-2} (-1)^(n+p)F(n+1)x^(2p-n-2) is a polynomial with integer coefficients, which is proved by multiplication with 1-x-x^2 and via F(n)+F(n+1)=F(n+2) and L(n)=F(n-1)+F(n+1). - _R. J. Mathar_, Jul 10 2008]. Conjecture: The Lucas sequence L also obeys all the recurrences: L(n)= L(p)*L(n-p)-(-1)^p*L(n-2p), any p>0, L=A000204. Conjectures from _Chai Wah Wu_, Apr 15 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, May 04 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004633 | OEIS-A138155

Triangle read by rows: T(n,k) is the number of Dyck paths with nondecreasing peaks having semilength n and with height of last peak equal to k (1 <= k <= n).

Conjectural g.f.: Sum_{n>=1} (t*x*(1 - x))^n/(Product_{i=2..n+1} (1 - 2*x + x^i)) = t*x + (t + t^2)*x^2 + (t + 2*t^2 + t^3)*x^3 + ... (checked up to x^12). - _Peter Bala_, Mar 31 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Mar 04 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004634 | OEIS-A138159

Triangle read by rows: T(n,k) is the number of permutations of [n] having k occurrences of the pattern 321 (n>=1, 0<=k<=n(n-1)(n-2)/6).

The number of 321-patterns of a given permutation p of [n] is given by Sum(L[i]R[i],i=1..n), where L (R) is the left (right) inversion vector of p. L and R are related by R[i]+i=p[i]+L[i] (the given Maple program makes use of this approach). References contain formulas and generating functions for the first few columns (some are only conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Mar 27 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004635 | OEIS-A138164

Row sums of Riordan array (c(-x^2),xc(-x^2)^2)^(-1) where c(x) is the g.f. of A000108.

Conjecture: -8*n*(n-1)*a(n) + 12*(n+3)*(n-1)*a(n-1) + 2*(89*n^2-512*n+651)*a(n-2) + 3*(-127*n^2+715*n-966)*a(n-3) + 3*(-247*n^2+2431*n-5698)*a(n-4) + 9*(75*n-341)*(3*n-16)*a(n-5) - 72*(3*n-14)*(3*n-16)*a(n-6) = 0. - _R. J. Mathar_, Feb 24 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 03 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004636 | OEIS-A138175

Row sums of the Riordan array (1/(1+x),x(1+2x)/(1+x)^3)^(-1).

Conjecture: 12*n*(n-1)*(39*n-88)*a(n) -8*(n-1)*(156*n^2-352*n+27)*a(n-1) +9*(-351*n^3-2196*n^2-4325*n+2680)*a(n-2) +24*(39*n-49)*(3*n-7)*(3*n-8)*a(n-3)=0.- _R. J. Mathar_, Oct 25 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 03 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004637 | OEIS-A138220

Numbers k such that 900*k^2 + 1 is prime.

Conjecture: (30*k)^(2^m)+1 is prime for some numbers k at all values of m >= 0. - _Richard R. Forberg_, Feb 06 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, May 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004638 | OEIS-A138265

Number of upper triangular zero-one matrices with n ones and no zero rows or columns.

For each m, a(5m+4) mod 5 = 0. Conjectured by Andrews-Sellers, and proved by Garvan (see Remark 1.4(ii) in Garvan's paper). Conjectural g.f.: 1 + Sum_{n >= 0} n/(1+x)^(n+1) * (Product_{i = 1..n} 1 - 1/(1+x)^i). Cf. A194530. - _Peter Bala_, Aug 21 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Mar 10 2008, Mar 11 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004639 | OEIS-A138289

Row sums of A138060.

Conjecture: a(n) = a(n-1) + a(n-4) with g.f. x*(-1 - 2*x - 3*x^2 - 4*x^3) / (-1 + x + x^4). - _R. J. Mathar_, Jul 08 2011 The conjectured recurrence is true. Proof:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, May 02 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004640 | OEIS-A138290

Numbers m such that 2^(m+1) - 2^k - 1 is composite for all 0 <= k < m.

Conjecture: 2^j - 2 are terms for j > 2. - _Chai Wah Wu_, Sep 07 2021 The proof of this conjecture is in A369375. - _Thomas Ordowski_, Mar 20 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 13 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004641 | OEIS-A138312

Decimal expansion of Mertens's constant B_3 minus Euler's constant.

Conjectured to be equivalent to 'kappa' = lim_{n->oo} ((Sum_{k=1..n} mu^2(k)/phi(k)) - H_n), where mu(k) is the Mobius function, phi(k) is Euler's Totient and H_n is the n-th harmonic number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Dick Boland (abstract(AT)imathination.org), Mar 13 2008, Mar 14 2008, Mar 27 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004642 | OEIS-A138313

Decimal expansion of constant 'kappa' = lim_{n -> infinity} (F_n - H_n), where H_n are harmonic numbers, F_n are squarefree totient analogs of H_n.

Conjectured to be equivalent to Mertens's constant B_3 minus Euler's constant (A138312). B_3 - gamma is given by Sum_{i>=1} log p_i/(p_i*(p_i-1)), where p_i is the i^th prime = Sum_{j>=2} mu(j)*zeta'(j)/zeta(j), mu(j) is the Mobius function, zeta'(j) is the derivative of zeta(j).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Dick Boland (abstract(AT)imathination.org), Mar 13 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004643 | OEIS-A138316

Numerators of the squarefree totient analogs of the harmonic numbers F_n.

F_n-H_n approaches a constant, 'kappa', conjectured to be equivalent to the difference of B_3-gamma, where B_3 is Mertens' 3rd constant and gamma is Euler's constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Dick Boland (abstract(AT)imathination.org), Mar 13 2008, Mar 27 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004644 | OEIS-A138350

Moment sequence of tr(A^2) in USp(4).

Conjectured e.g.f. Bessell[1,2x](Bessell[0,2x]-Bessell[1,2x])/x. - _Benjamin Phillabaum_, Feb 25 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew V. Sutherland_, Mar 16 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004645 | OEIS-A138365

a(n) = A006190(n) * A006190(n+2).

For n >= 4, a(n) = 10*a(n-1) + 10*a(n-2) - a(n-3); from this it follows that the conjectured generating function is correct. - _Sela Fried_, Oct 14 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Mar 16 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004646 | OEIS-A138413

A bisection of A000957.

Conjecture: D-finite with recurrence 4*n*(2*n-1)*(7*n-13)*a(n) +(-910*n^3+3489*n^2-4277*n+1680)*a(n-1) +2*(4*n-7)*(7*n-6)*(4*n-5)*a(n-2)=0. Telescoping would provide another recurrence for A000957. - _R. J. Mathar_, Jun 26 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 08 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004647 | OEIS-A138467

a(1)=1, then for n>=2 a(n) = n - floor((1/3)*a(a(n-1))).

Conjecture: a(n) = floor(r(p)*(n+1)) with r(p)=(1/2)*(sqrt(p*(p+4))-p). - _Yalcin Aktar_, Jul 13 2008 Conjecture: b_p(n) = floor(r(p)*(n+1)) where r(p) =(1/2)*(sqrt(p*(p+4))-p).(End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, May 09 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004648 | OEIS-A138572

Numbers k that divide the sum of the digits of k^k in base 2.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Gerbicz_, May 12 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004649 | OEIS-A138670

Indices of 0's in Stewart's choral sequence.

Conjecture: essentially partial sums of A106036 (verified for n <= 10000). - _Sean A. Irvine_, Jul 19 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | Joel Reyes Noche (joel.noche(AT)up.edu.ph), Mar 26 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004650 | OEIS-A138684

First differences of A138683.

Conjecture. Let x(1)=1, x(2)=2, x(n) = 2*x(n-1) + x(n-2), and let y(1)=1, y(2)=3, y(n) = y(n-1) + y(n-2). Then if n=x(k), a(n)=y(k); if x(k) < n < 2*x(k), a(n) = a(n-x(k)); and if 2*x(k) <= n < x(k+1), a(n)=1. (This has been confirmed for n < 500.) - _John W. Layman_, Nov 26 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Mar 26 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004651 | OEIS-A138738

E.g.f. satisfies: A(x) = exp(x*exp(x^2*exp(x^3*exp(...exp(x^n*...))))).

a(n) == 1 (mod 6) (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 27 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004652 | OEIS-A138808

Number of integer pairs (x,y), x > 0, y > 0, such that x <= p, y <= q for any factorization n = p*q.

Conjecture: the row sums of the plane partitions A010766 are upper bounds. - _R. J. Mathar_, Aug 06 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonas Wallgren_, May 16 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004653 | OEIS-A138830

Bisection of imperfect numbers A132999.

The conjectured g.f. (-x^14+x^13-x^3+x^2+2)/(x^2-2*x+1) that a correspondent suggested is wrong and fails after roughly 240 terms [_R. J. Mathar_, Jun 15 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Apr 06 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004654 | OEIS-A139003

Number of operations A000142 (i.e., x!) or A000196 (i.e., floor(sqrt(x))) needed to get n, starting with 3.

Knuth conjectured that any number can be obtained in that way, starting from 4. In fact a single 2 is enough to get any positive integer, if Knuth's conjecture that one 4 is enough is true. From 2, (((-tan(2.))!)!)! = 5.592..., then floor, factorial gets 120, then sqrt, sqrt gives 3.162..., and floor gives 3, or negate, floor, negate gives 4. - _N. J. A. Sloane_, Feb 26 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 09 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004655 | OEIS-A139004

Number of operations A000142 (i.e., x!) or A000196 (i.e., floor(sqrt(x))) needed to get n, starting with 4.

Knuth conjectured that any number can be obtained in this way, starting from 4. If we are allowed to use tan(x) just once, then a single 2 is enough to get any positive integer, if Knuth's conjecture that one 4 is enough is true. From 2, (((-tan(2.))!)!)! = 5.592..., then floor, factorial gets 120, then sqrt, sqrt gives 3.162..., and floor gives 3, or negate, floor, negate gives 4. - _N. J. A. Sloane_, Feb 26 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 09 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004656 | OEIS-A139250

The toothpick sequence (see Comments lines for definition).

Conjecture: Consider the rectangles in the sieve (including the squares). The area of each rectangle (A = b*c) and the edges (b and c) are powers of 2, but at least one of the edges (b or c) is <= 2. Conjectures: after 2^k stages, for k >= 2, and for m = 1 to k - 1, there are 4^(m-1) substructures of size s = k - m, where every substructure has 4*s rectangles. The total number of substructures is equal to (4^(k-1)-1)/3 = A002450(k-1). For example: If k = 5 (after 32 stages) we can see that:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Apr 24 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004657 | OEIS-A139251

First differences of toothpicks numbers A139250.

It appears that there is a relation between this sequence, even superperfect numbers, Mersenne primes and even perfect numbers. Conjecture: The sum of the toothpicks added to the toothpick structure between the stage A061652(k) and the stage A000668(k) is equal to the k-th even perfect number, for k >= 1. For example: A000396(1) = 2+4 = 6. A000396(2) = 4+4+8+12 = 28. A000396(3) = 16+4+8+12+12+16+28+32+20+16+28+36+40+60+88+80 = 496. - _Omar E. Pol_, May 04 2009 Concerning this conjecture, see _David Applegate_'s comments on the conjectures in A153006. - _N. J. A. Sloane_, May 14 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Apr 24 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004658 | OEIS-A139262

Total number of two-element anti-chains over all ordered trees on n edges.

Conjecture: n*(n-2)*a(n) +2*(-4*n^2+9*n-3)*a(n-1) +8*(n-1)*(2*n-3)*a(n-2)=0. - _R. J. Mathar_, Feb 03 2013 The above conjecture follows easily from the formula by J. M. Bergot. - _Robert Israel_, Feb 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lifoma Salaam_, Apr 12 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004659 | OEIS-A139376

Expansion of 1/((1-x^2*c(x))(1-x-x^2)) where c(x) is the g.f. of A000108.

Conjecture: (-n+1)*a(n) +6*(n-2)*a(n-1) +4*(-2*n+5)*a(n-2) +2*(-n+1)*a(n-3) +3*(3*n-7)*a(n-4) +3*(-n+3)*a(n-5) +2*(-2*n+5)*a(n-6)=0. - _R. J. Mathar_, Feb 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004660 | OEIS-A139383

Number of n-level labeled rooted trees with n leaves.

Conjecture: a(n) ~ c * n^(2*n-5/6) / (2^(n-1) * exp(n)), where c = 2.86539...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 16 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004661 | OEIS-A139399

Number of steps to reach a cycle in Collatz problem.

a(n) = A006577(n) - 2 for n > 2 (if the conjecture holds).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004662 | OEIS-A139474

a(n) = ((2*sqrt(2) + 3)^(2^(prime(n) - 1) - 1) - (3 - 2*sqrt(2))^(2^(prime(n) - 1) - 1))/(4*sqrt(2)).

Conjecture of _Kenneth J Ramsey_ from May 16 2006 (see A001109): a(n) is divisible by 2^prime(n)-1 if and only 2^prime(n)-1 is a Mersenne prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Apr 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004663 | OEIS-A139476

Positions of squares in the EKG sequence (A064413).

Conjecture: the squares appear in increasing order. The conjecture is false; in the EKG sequence, 158^2 is at position 24142, but 157^2 is at position 24146. Note that 157 is prime. If we let p=157, then the two terms on either side of p^2 are p(p+2) and p(p+3), which is unusual because for all primes 3 < p < 157, the three terms are p(p+1), p^2, p(p+2). The next unusual prime is 661. There are no others less than 8164.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, May 10 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004664 | OEIS-A139490

Numbers n such that the quadratic form x^2 + n*x*y + y^2 represents exactly the same primes as the quadratic form x^2 + m*y^2 for some m.

Conjecture: This sequence is finite and complete (checked for range n<=200 and m<=500).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Apr 24 2008, Apr 26 2008, Apr 27 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004665 | OEIS-A139544

Numbers which are not the difference of two squares of positive integers.

Conjecture: these numbers do not occur in A139491.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Apr 25 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004666 | OEIS-A139758

a(n) is the smallest prime such that (a(n) - the n-th prime) is a power of 2.

It is only conjectured (see _Cino Hilliard_'s comment at A094076) that this sequence is defined for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, May 19 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004667 | OEIS-A139772

Number of linear trimer coverings of a 3n X 3n square.

Conjecture: a(n) ~ alpha * beta^n * gamma^(n^2), where 0.933 < alpha < 0.979, 0.4619 < beta < 0.4696, 4.1691 < gamma < 4.1739. Notice similarity with A004003. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Finch_, May 20 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004668 | OEIS-A139804

Smallest m such that (3^n)! + m is prime.

Conjecture: There are no composite numbers in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, May 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004669 | OEIS-A140100

Start with Y(0)=0, X(1)=1, Y(1)=2. For n > 1, choose least positive integers Y(n) > X(n) such that neither Y(n) nor X(n) appear in {Y(k), 1 <= k < n} or {X(k), 1 <= k < n} and such that Y(n) - X(n) does not appear in {Y(k) - X(k), 1 <= k < n} or {Y(k) + X(k), 1 <= k < n}; sequence gives X(n) (for Y(n) see A140101).

Conjecture: A140100(n) - A140098(n) = A276404(n) is always 0 or 1; see A276406 for the positions where a difference of 1 occurs. The sequence is "tribonacci-synchronized"; this means there is a finite automaton recognizing the tribonacci representation of (n,a(n)) input in parallel, where a shorter input is padded with leading zeros. This finite automaton has 22 states and was verified with Walnut. In particular this finite automaton and a similar one for A140101 was used to verify that (conjecture of J. Cassaigne) either a(b(n)) = a(n)+b(n) or b(a(n)) = a(n)+b(n) for all n>=1, where b(n) = A140101(n). - _Jeffrey Shallit_, Oct 04 2022 Conjecture: the limit of X(n)/n = 1+1/t and limit of Y(n)/n = 1+t where the limit of Y(n)/X(n) = t = tribonacci constant (A058265), and thus the limit of (Y(n) + X(n))/(Y(n) - X(n)) = t^2 and the limit of (Y(n)^2 + X(n)^2)/(Y(n)^2 - X(n)^2) = t. It is conjectured in A305392 that the first differences of (X(n)) as a word are given by 212121 delta(x), where x is the tribonacci word x = A092782, and delta is the morphism This conjecture implies the frequency conjectures above: let N(i,n) be the number of letters i in x(1)x(2)...x(n). Then simple counting gives

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 04 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004670 | OEIS-A140101

Start with Y(0)=0, X(1)=1, Y(1)=2. For n > 1, choose least positive integers Y(n) > X(n) such that neither Y(n) nor X(n) appear in {Y(k), 1 <= k < n} or {X(k), 1 <= k < n} and such that Y(n)-X(n) does not appear in {Y(k)-X(k), 1 <= k < n} or {Y(k)+X(k), 1 <= k < n}; sequence gives Y(n) (for X(n) see A140100).

Conjecture: a(n) = A003144(n) + n. (This is from my notebook Lattices 115 page 20 from Oct 25 2016. It is now a theorem - see the Dekking et al. paper.) - _N. J. A. Sloane_, Jul 22 2019 The sequence is "tribonacci-synchronized"; this means there is a finite automaton recognizing the tribonacci representation of (n,a(n)) input in parallel, where a shorter input is padded with leading zeros. This finite automaton has 23 states and was verified with Walnut. In particular this finite automaton and a similar one for A140101 was used to verify that (conjecture of J. Cassaigne) either a(b(n)) = a(n)+b(n) or b(a(n)) = a(n)+b(n) for all n>=1, where b(n) = A140100(n). - _Jeffrey Shallit_, Oct 04 2022 CONJECTURE: the limit of a(n)/n = 1+t and limit of X(n)/n = 1+1/t so that limit of a(n)/X(n) = t = tribonacci constant (A058265), and thus the limit of [a(n) + X(n)]/[a(n) - X(n)] = t^2 and the limit of [a(n)^2 + X(n)^2]/[a(n)^2 - X(n)^2] = t. Conjectured recursion: Take first differences: 3, 3, 3, 2, 3, 3, 3, 3, 3, 2, ... (appears to consist of only 3's and 2's); list the run lengths: 3, 1, 6, 1, 5, 1, 6, 1, 3, 1, 6, 1, 5, 1, 6, 1, ... (it appears that every second term is 1 and the other terms are 3, 5, and 6); and bisect, getting 3, 6, 5, 6, 3, 6, 5, 6, 6, 5, 6, 3, 6, ... This is (although I do not have a proof) the recursively defined A275925. Thanks to _Alois P. Heinz_ for providing enough terms of A273059 to enable a (morally) convincing check of this conjecture. - _N. J. A. Sloane_, Aug 30 2016 This conjecture can be reformulated as follows (cf. A140100). This conjecture implies the frequency conjecture above: let N(i,n) be the number of letters i in a(1)a(2)...a(n). Then simple counting gives

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 04 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004671 | OEIS-A140103

Term-by-term sums of A140101 and A140100; also, equals the complement of A140102, which is the term-by-term differences of A140101 and A140100, where A140101 is the complement of A140100.

Conjecture: a(n) = A003145(n) + n. This is the most direct connection between the Greedy Queens sequence and the tribonacci word that I know. - _Michel Dekking_, Mar 19 2019. [My notes show that I made this conjecture on Jul 20 2018. There are many similar conjectures relating the two problems. For example A140100 = A003145(n)-A003144(n), A140101(n) = A003146(n)-A003145(n), a(n) = A003146(n)-A003144(n). - _N. J. A. Sloane_, Mar 19 2019] All these conjectures are now theorems - see the Dekking et al. paper. - _N. J. A. Sloane_, Jul 22 2019 Conjecture: the limit of A140103(n)/A140102(n) = t^2 = 3.38297576... (cf. A276800) where the limit of A140101(n)/A140100(n) = t =

1.839286755.. and t = tribonacci constant satisfies: t^3 = 1 + t + t^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 04 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004672 | OEIS-A140110

Numbers n such that for all divisors of n, ratios of 2 consecutive divisors of n will always reduce to lowest terms as a ratio of consecutive integers.

Proof of John W. Layman's conjecture that every term after the second is a multiple of 4 or 6. The first divisor of any number is always 1. Because the only divisor of 1 is 1, the second divisor of any member of this sequence greater than 1 is 2. Because the divisors of 2 are 1 and 2, the third divisor of any term of this sequence is always either 3 (proving it is a multiple of 6 because 6 is the LCM of 2 and 3) or 4. Therefore every term greater than 2 in this sequence is a multiple of at least one of 4 and 6. - _J. Lowell_, Jul 07 2008 It appears that consecutive elements of this sequence (for n>1) are consecutive solutions of the equation n = 1 + Sum_{k=1..tau(n)-1} gcd(d_k(n), d_(k+1)(n)), where d_k(n) denotes the k-th smallest divisor. The conjecture was checked for 10^8 consecutive integers. - _Lechoslaw Ratajczak_, Jan 09 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Jun 21 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004673 | OEIS-A140124

a(n) = degree in N of the number of orbits under S_N of the set of n-tuples of partitions of {1,...,N} into n subsets.

This formula and its first two values are given in Bergman, p. 18. Abstract: Erdos and Szekeres showed in 1978 that for any four positive integers satisfying m_1+m_2 = n_1+n_2, the two binomial coefficients (m_1+m_2)!/m_1! m_2! and (n_1+n_2)!/n_1! n_2! have a common divisor >1. The analogous statement for families of k k-nomial coefficients (k>1) was conjectured in 1997 by David Wasserman. Erdos and Szekeres remark that if m_1, m_2, n_1, n_2 as above are all >1, there is probably a lower bound on the common divisor in question which goes to infinity as a function of m_1+m_2. Such a bound is here obtained. Results are proved that narrow the class of possible counterexamples to Wasserman's conjecture. On the other hand, several plausible generalizations of that conjecture are shown to be false.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jun 04 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004674 | OEIS-A140141

Positions of second appearances of primes in A039649.

If prime(n) > 7 is in A005385, then a(n) = 2*prime(n). - _Thomas Ordowski_ (conjecture) and _Robert Israel_ (proof), Jan 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 10 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004675 | OEIS-A140320

a(n) = A137576((3^n-1)/2).

Conjecture. a(n) = 2n*3^(n-1)+1. If conjecture is true then limsup(A137576(n)/n)=infinity while liminf(A137576(n)/n)=2 with a realization on primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 26 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004676 | OEIS-A140355

Smallest integer such that the first n multiples of a(n) have a number of divisors that is strictly increasing.

Conjecture: this sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, May 30 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004677 | OEIS-A140356

Triangle T(n,m) read by rows: m! if m <= floor(n/2), and (n-m)! otherwise.

Conjecture: limit_{n->infinity} sum_{m=0..n} (1/binomial(n,m) - T(n,m)) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_ and _Gary W. Adamson_, May 30 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004678 | OEIS-A140357

a(1)=1; a(n)=floor(4*a(n-1)*(n-a(n-1)) / n) for n > 1.

Conjecture: starting with any given n and any 1 <= a(n) <= n and applying the rule for the sequence produces a sequence which eventually joins this one. For example, starting with a(9)=5, the sequence continues 10,3,9,11,9, at which point it has joined.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, May 30 2008, May 31 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004679 | OEIS-A140358

Smallest nonnegative integer k such that n = +-1+-2+-...+-k for some choice of +'s and -'s.

Conjecture when n is greater than 0. Choose k so that t(k)<=n<t(k+1) where t(n) is the n-th triangular number t(n)=n(n+1)/2. If n=t(k), a(n)=k, otherwise if k is odd then a(n)=k+2 if n-t(k) is odd, a(n)=k+1 if n-t(k) is even, else if k is even than a(n)=k+1 if n-t(k) is odd, a(n)=k+3 if n-t(k) is even. (This has been verified for n up to 100.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jun 23 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004680 | OEIS-A140427

Arises in relating doubly-even error-correcting codes, graphs and irreducible representations of N-extended supersymmetry.

Conjecture: essentially partial sums of A169675 (verified for n <= 10000). - _Sean A. Irvine_, Jul 19 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jun 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004681 | OEIS-A140444

Primes congruent to 1 (mod 14).

Conjecture: The monic polynomial P(x) = (x+1)^7/x - 1/x = ((x+1)^7-1)/x is irreducible but factorizable over Galois field (mod a(n)) with exactly 6 distinct irreducible factors of degree 1. Examples:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jun 26 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004682 | OEIS-A140454

Number of primes p less than 10^n such that p^2-2 is prime.

Korevaar gives these values in Table 1, p. 18, attributing the calculation to Fokko van de Bult. Abstract: For any positive integer r, let pi_{2r}(x) denote the number of prime pairs (p, p+2r) with p not exceeding (large) x. According to the prime-pair conjecture of Hardy and Littlewood, pi_{2r}(x) should be asymptotic to 2C_{2r}li_2(x) with an explicit positive constant C_{2r}. A heuristic argument indicates that the remainder e_{2r}(x) in this approximation cannot be of lower order than x^beta, where beta is the supremum of the real parts of zeta's zeros. The argument also suggests an approximation for pi_{2r}(x) similar to one of Riemann for pi(x).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jun 26 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004683 | OEIS-A140472

a(n) = a(n - a(n-1)) + a(floor(n/2)).

a(n) = (n+A006519(n))/2 for n > 0 (conjectured). - _Jon Maiga_, Aug 16 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_ and _Gary W. Adamson_, Jun 28 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004684 | OEIS-A140601

Values of m such that binomial(m, a) + binomial(m, b) divides binomial(m, a + b) for some distinct nonnegative integers a and b with a + b <= m.

This sequence was first suggested by _Stefan Steinerberger_, who conjectures that there are infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew V. Sutherland_, May 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004685 | OEIS-A140608

"Right" odd composite numbers n for which n > A140607((n-1)/2).

Conjecture. The sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004686 | OEIS-A140717

Triangle read by rows: T(n,k) is the number of Dyck paths d of semilength n such that sum of peakheights of d - number of peaks of d = k (n >= 0, 0 <= k <= floor(n^2/4)).

Conjecture: antidiagonal sums equal A227309. - _Mikhail Kurkov_, Aug 30 2024 Conjecture: n-th row polynomial equals t_n for n > 0 where we start with vector v of fixed length m with elements v_i = 1, then set t = v and for i=1..m-1, for j=i+1..m apply [v_i, v_j] := [v_i + z^(j-i)*v_j, z*v_i + v_j] (here square brackets mean that instead of sequentially assigning v_i and then v_j, we reserve their values (for example, as A = v_i, B = v_j) and then assign them in any order) and t_{i+1} := v_{i+1} (after ending each cycle for j). It also seems that if we change z^(j-i) to z^(2*(j-i)) it gives us equivalence of t_n and n-th row polynomial of A227543. - _Mikhail Kurkov_, Aug 30 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jun 08 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004687 | OEIS-A140794

One of the four smallest counterexamples to the conjecture that the cardinality of the sumset is less than or equal to the cardinality of the difference set of every finite set of integers.

One of the four smallest counterexamples to the conjecture that the cardinality of the sumset is less than or equal to the cardinality of the difference set of every finite set of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jul 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004688 | OEIS-A140829

a(0)=1; for n >= 1, a(n) = ceiling(Fibonacci(n)/a(n-1)).

Conjectures from _Colin Barker_, Mar 08 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Jul 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004689 | OEIS-A140831

Numbers in whose canonical prime factorization the powers of the primes do not form an increasing sequence.

Conjecture: There are infinitely many terms k such that k+1 is also a term. - _Mike Jones_, Jun 18 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Jul 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004690 | OEIS-A140869

Triangle read by rows where T(m,n) = floor((2mn+m+n-2)/2), m >= n >= 1.

Conjecture: If h does not belong to the sequence, then 4*h+5 is prime. - _Vincenzo Librandi_, Nov 18 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Jan 16 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004691 | OEIS-A140901

Number of 3 X 5 matrices with elements in 0..n with each row and each column in nondecreasing order. 3,5,n can be permuted, see formula.

(Conjecture) G.f.: (1 + 40*x + 400*x^2 + 1456*x^3 + 2212*x^4 + 1456*x^5 + 400*x^6 + 40*x^7 + x^8)/(1-x)^16. - _Bruno Berselli_, May 08 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004692 | OEIS-A140934

Number of 2 X 11 matrices with elements in 0..n with each row and each column in nondecreasing order. 2,11,n can be permuted, see formula.

Conjecture: 12th column (and diagonal) of the triangle A001263. - _Bruno Berselli_, May 07 2012 G.f. conjectured: (1 + 55*x + 825*x^2 + 4950*x^3 + 13860*x^4 + 19404*x^5 + 13860*x^6 + 4950*x^7 + 825*x^8 + 55*x^9 + x^10)/(1 - x)^23. - _Bruno Berselli_, May 07 2012 Conjecture: a(n) = ((n+12)/(12*n+12))*binomial(n+11,11)^2. - _Bruno Berselli_, May 07 2012 Conjecture: a(n) = Product_{i=1..11} A002378(n+i)/A002378(i). - _Bruno Berselli_, Sep 01 2016 Conjecture: Sum_{n>=0} 1/a(n) = 3538258540001/8820 - 40646320*Pi^2. Conjecture: Sum_{n>=0} (-1)^n/a(n) = 1678950598/2205 - 23068672*log(2)/21. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004693 | OEIS-A140955

Even integers that do not have at least two partitions into 2 distinct primes.

If A056636(3) = 128 (as is conjectured), then 38 is the last term in the sequence. - _Charles R Greathouse IV_, Sep 07 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gil Broussard_, Jul 25 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004694 | OEIS-A140999

Members of A067128 that are the smallest numbers with their prime signatures.

Conjecture (false!): includes all members of A094348. Counterexample to conjecture: 5354228880, the smallest positive multiple of the first 23 positive integers, does not belong to A067128. It is the smallest member of A003418 (a subsequence of A094348) not to be largely composite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Jul 28 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004695 | OEIS-A141116

Smallest n-digit prime with no identical adjacent digits (or 0 if no such prime exists).

For n >= 1, a(n) >= A056830(n), the least n-digit positive integer with no identical adjacent digits (also the least positive integer whose digits occur in n runs). Conjecture: For all n, a(n) <> 0. If the conjecture is true, then this sequence and the following two sequences are equivalent: i) Smallest prime with exactly n runs of digits and ii) Smallest prime with at least n runs of digits. For each n <= 625, a(n) is an n-digit prime (provided that each probable prime shown in the link is indeed a prime -- or at least one of very many (slightly) larger probable prime candidates is prime).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rick L. Shepherd_, Jun 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004696 | OEIS-A141117

G.f.: A(x) = G(G(x)) = G(x)*(1 + A(A(x))) where G(x) = x + x*G(G(G(x))) = g.f. of A091713.

Conjecture: a(n) is divisible by 4 iff n is divisible by 3, otherwise a(n) == 2 (mod 4) for n > 1 with a(1) = 1. - _Paul D. Hanna_, Apr 03 2026 The conjecture is true (see link). - _Yuan Chi_, Aug 20 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 04 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004697 | OEIS-A141135

Minimal number of unit edges required to construct n regular pentagons when allowing edge-sharing.

Conjectures from _Colin Barker_, Apr 05 2019: (Start) Conjecture: if n is a term in A121149, a(n) = a(n-1) + 3, otherwise a(n) = a(n-1) + 4. - _Jinyuan Wang_, Apr 05 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ralph H. Buchholz (teufel_pi(AT)yahoo.com), Jun 08 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004698 | OEIS-A141146

Number of linear arrangements of n blue, n red and n green items such that first and last elements are blue but there are no adjacent items of the same color.

Conjecture: -(n+1)*(n-2)*a(n) +(7*n^2-13*n+4)*a(n-1) +8*(n-2)^2*a(n-2)=0. - _R. J. Mathar_, Jul 23 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Jun 10 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004699 | OEIS-A141204

Let sequences X and Y consist of the least positive integers such that 2X+Y is the complement of X and X+Y is the complement of Y, starting with X(1)=1 and Y(1)=1; then this sequence equals X, while 2X+Y=A141205, Y=A141206 and X+Y=A141207.

CONJECTURES on evaluating limits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 21 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004700 | OEIS-A141205

Let sequences X and Y consist of the least positive integers such that 2X+Y is the complement of X and X+Y is the complement of Y, starting with X(1)=1 and Y(1)=1; then this sequence equals 2X+Y, while X=A141204, Y=A141206 and X+Y=A141207.

CONJECTURES on evaluating limits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 21 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004701 | OEIS-A141206

Let sequences X and Y consist of the least positive integers such that 2X+Y is the complement of X and X+Y is the complement of Y, starting with X(1)=1 and Y(1)=1; then this sequence equals Y, while X=A141204, 2X+Y=A141205 and X+Y=A141207.

CONJECTURES on evaluating limits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 21 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004702 | OEIS-A141207

Let sequences X and Y consist of the least positive integers such that 2X+Y is the complement of X and X+Y is the complement of Y, starting with X(1)=1 and Y(1)=1; then this sequence equals X+Y, while X=A141204, 2X+Y=A141205, Y=A141206.

CONJECTURES on evaluating limits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 21 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004703 | OEIS-A141222

Expansion of -1/(2*x) + (2*x-1)^2/(2*x*(1-4x)^(3/2)).

Conjecture: (n+1)*a(n) + 2*(-3*n-1)*a(n-1) + 4*(2*n-3)*a(n-2) = 0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jun 14 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004704 | OEIS-A141244

Numerators in the expansion of (1-sqrt(1-x^2))/(1-x).

a(2n) = a(2n+1) = A120778(n-1) (conjectural). - _Greg Martin_, Aug 16 2014, corrected by _M. F. Hasler_, Aug 18 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jun 17 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004705 | OEIS-A141273

Conjectured to be complete list of numbers not of the form 4*x^2 + 4*y^2 + triangular number.

Conjectured to be complete list of numbers not of the form 4*x^2 + 4*y^2 + triangular number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004706 | OEIS-A141292

Conjectured values for minimal number of pieces required in a 2n-gon to square dissection that uses translation alone.

Conjectured values for minimal number of pieces required in a 2n-gon to square dissection that uses translation alone.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Pamela Pierce (PPierce(AT)wooster.edu), Jeffrey Willert (jwillert09(AT)wooster.edu) and Wenyuan Wu (wwu11(AT)wooster.edu), Aug 01 2008, Aug 12 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004707 | OEIS-A141342

A transform of the Fibonacci numbers.

Conjecture: (n-1)*a(n) +4*(n-4)*a(n-1) + (65-29*n)*a(n-2) +12*(7-2*n)*a(n-3)+ 4*(4-n)*a(n-4) =0. - _R. J. Mathar_, Nov 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jun 26 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004708 | OEIS-A141344

Expansion of (2-sqrt(1+4x))/(2-x-sqrt(1+4x)).

Conjectured to be D-finite with recurrence: 3*(n-1)*a(n) +2*(2*n-11)*a(n-1) +(79-31*n)*a(n-2) +2*(2*n-5)*a(n-3)=0. - _R. J. Mathar_, Oct 25 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jun 26 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004709 | OEIS-A141386

Conjectured to be complete list of numbers not of the form x^2 + y^2 + 5*triangular number.

Conjectured to be complete list of numbers not of the form x^2 + y^2 + 5*triangular number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004710 | OEIS-A141387

Triangle read by rows: T(n,m) = n + 2*m*(n - m) (0 <= m <= n).

Construct the infinite-dimensional matrix representation of angular momentum operators (J_1,J_2,J_3) in Jordan-Schwinger form (cf. Harter, Klee, Schwinger). The triangle terms T(n,k)=T(2j,j+m) satisfy:(1/4)T(2j,j+m) = <j,m|J_1^2|j,m> = <j,m|J_2^2|j,m>. Matrices for J_1^2 and J_2^2 are sparse. These diagonal equalities and the off-diagonal equalities of A268759 determine the only nonzero entries. Comments on A268759 provide a conjecture for the clear interpretation of these numbers in the context of binomial coefficients and other geometrical sequences. - _Bradley Klee_, Feb 20 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Aug 03 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004711 | OEIS-A141412

Triangle c(n,k) of the denominators of coefficients [x^k] P(n,x) of the polynomials P(n,x) of A129891.

Conjecture: T(n, k) = d(n+1, k+1), with d(n,k) = denominator(A000254(n, k)*k!/n!) where A000254 are the unsigned Stirling numbers of the 1st kind. See d(n,k) in Farhi link. - _Michel Marcus_, Oct 18 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Aug 04 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004712 | OEIS-A141423

Conjectured to be complete list of numbers not of the form 3*x^2 + 4*y^2 + triangular number.

Conjectured to be complete list of numbers not of the form 3*x^2 + 4*y^2 + triangular number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004713 | OEIS-A141428

Conjectured to be complete list of numbers not of the form x^2 + 5*y^2 + 3*triangular number.

Conjectured to be complete list of numbers not of the form x^2 + 5*y^2 + 3*triangular number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004714 | OEIS-A141436

Tisdale's sieve: generators for an infinite set of disjoint prime/nonprime sequences (see Comments for precise definition).

Conjecture: Equals A102615 UNION A006450. - _R. J. Mathar_, Aug 14 2008 Proof of Conjecture, from _David Applegate_, Dec 28 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Tisdale_, Aug 06 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004715 | OEIS-A141443

Conjectured to be complete list of numbers not of the form 2*x^2 + y^2 + 12*triangular number.

Conjectured to be complete list of numbers not of the form 2*x^2 + y^2 + 12*triangular number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004716 | OEIS-A141459

a(n) = Product_{p-1 divides n} p, where p is an odd prime.

Conjecture: a(n) = denominator of integral_{0..1}(log(1-1/x)^n) dx. - _Jean-François Alcover_, Feb 01 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Aug 08 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004717 | OEIS-A141463

Conjectured to be the complete list of numbers not of the form 2*x^2 + 5*y^2 + 4*triangular number.

Conjectured to be the complete list of numbers not of the form 2*x^2 + 5*y^2 + 4*triangular number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004718 | OEIS-A141464

Conjectured to be complete list of numbers not of the form 4*x^2 + 3*triangular number + triangular number.

Conjectured to be complete list of numbers not of the form 4*x^2 + 3*triangular number + triangular number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004719 | OEIS-A141491

Conjectured to be complete list of numbers not of the form 5*x^2 + 3*triangular number + 2*triangular number.

Conjectured to be complete list of numbers not of the form 5*x^2 + 3*triangular number + 2*triangular number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004720 | OEIS-A141497

Conjectured to be complete list of numbers not of the form 2*x^2 + 4*triangular number + 5*triangular number.

Conjectured to be complete list of numbers not of the form 2*x^2 + 4*triangular number + 5*triangular number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004721 | OEIS-A141771

Expansion of (1-sqrt(1-4*x))/(2*x) + 8*x^3/((sqrt(1-4*x))*(1+sqrt(1-4*x))^3).

Conjecture: (n+1) *(n^2+5*n-12) *(n^3-5*n^2+38*n-250)*a(n) -2 *(2*n-3) *(n^2+7*n-6) *(n^3-5*n^2+38*n-250) *a(n-1)=0. - _R. J. Mathar_, Apr 30 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004722 | OEIS-A141779

Numbers k such that A120292(k) is composite.

Conjecture: All composite terms of A120292 are semiprime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Jul 04 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004723 | OEIS-A141821

Least number k < n and coprime to n such that the largest term of the continued fraction of k/n is as small as possible.

See A141822 for the value of the largest term in the continued fraction of a(n)/n. Zaremba conjectured that the largest value is 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 08 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004724 | OEIS-A141822

Maximum term in the continued fraction of A141821(n)/n.

Zaremba conjectured that a(n)<=5, a bound that is attained for n in A195901. It appears that n=150 may be the largest integer with a(n)=5, while n=6234 may be the largest integer with a(n)=4. Conjecture: There exists an integer N0 such that a(n) = 2 for all n > N0. - _Bartlomiej Pawlik_, Mar 04 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 08 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004725 | OEIS-A141823

Integers k>1 such that A141822(k)=4.

Zaremba conjectured that A141832, A141833, A195901, and this sequence form a partition of the integers >1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 08 2008, Jul 09 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004726 | OEIS-A141832

Integers n>1 such that A141822(n)=2.

Zaremba conjectured that A141823, A141833, A195901, and this sequence form a partition of the integers >1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 09 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004727 | OEIS-A141833

Integers n>1 such that A141822(n)=3.

Zaremba conjectured that A141823, A141832, A195901, and this sequence form a partition of the integers >1.Is this sequence finite or infinite?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 09 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004728 | OEIS-A141849

Primes congruent to 1 mod 11.

Conjecture: Also primes p such that ((x+1)^11-1)/x has 10 distinct irreducible factors of degree 1 over GF(p). - _Federico Provvedi_, Apr 17 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 11 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004729 | OEIS-A142240

A triangular sequence from the pattern in row sums of Pascal's triangle A007318, Eulerian numbers A008292 and A060187: Delta_diagonal=m; m={0,1,2,3,...k}.

The conjecture that goes with this triangular sequence is that there on n levels like these for Pascal combinatorial quantum levels.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_ and _Gary W. Adamson_, Sep 17 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004730 | OEIS-A142248

Odd numbers in A138123.

Conjectures from _Chai Wah Wu_, Apr 15 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Sep 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004731 | OEIS-A142591

Composite terms in A143578.

Conjecture: This consists exactly of the semiprimes p*q for which p + q divides p*q + 1. - _Mohamed Bouhamida_, Aug 17 2009 (Comment edited by _N. J. A. Sloane_, Sep 01 2019.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Aug 24 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004732 | OEIS-A142978

Table of figurate numbers for the n-dimensional cross polytopes.

Conjectural congruences for main diagonal entries: Put A(n) = T(n,n). Calculation suggests the following congruences: for prime p > 3 and m, r >= 1, A(m*p^r) == A(m*p^(r-1)) (mod p^(3*r));

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004733 | OEIS-A143007

Square array, read by antidiagonals, where row n equals the crystal ball sequence for the 2*n-dimensional lattice A_n x A_n.

Main diagonal: zeta(3) = 6 * Sum_{n >= 1} 1/(n^3*T(n-1,n-1)*T(n,n)). Conjectural result for other diagonals: zeta(3) = 1 + 1/2^3 + ... + 1/k^3 + Sum_{n >= 1} (2*n+k)*(3*n^2+3*n*k+k^2)/(n^3*(n+k)^3*T(n-1,n+k-1)*T(n,n+k)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004734 | OEIS-A143013

Number of Motzkin n-paths with two kinds of level steps one of which is a final step.

Conjecture: (n+2)*a(n) -(2*n+1)*a(n-1) +3*(1-n)*a(n-2) +2*(5-2*n)*a(n-3)=0. - _R. J. Mathar_, Oct 25 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Jul 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004735 | OEIS-A143017

Number of {2-1-3, 2'^e-31}-avoiding permutations of size n (see definition in the Elizalde paper).

Conjecture: -8*n*(n+1)*a(n) + 4*n*(2*n+5)*a(n-1) + 4*n*(n+7)*a(n-2) + 2*(70*n^2-395*n+564)*a(n-3) + 2*(25*n^2-143*n+222)*a(n-4) + 4*(49*n-228)*(n-5)*a(n-5) - 45*(n-5)*(n-6)*a(n-6) = 0. - _R. J. Mathar_, Mar 14 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jul 17 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004736 | OEIS-A143020

Sum of the distances from a fixed node (root) to the next node in all non-crossing graphs on n nodes on a circle.

Conjecture: -(n-1) *(n+1) *(39*n^2-136*n+60) *a(n) +60 *(-18*n^2+57*n-7) *a(n-1) +12 *(3*n-7) *(3*n-8) *(39*n^2-58*n-37) *a(n-2)=0. - _R. J. Mathar_, May 10 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jul 30 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004737 | OEIS-A143027

Sturdy prime numbers: p such that in binary notation k*p has at least as many 1-bits as p for all k > 0.

Max Alekseyev reports that 6700417, 13264529, 20394401, and 97685839 are not sturdy because each number divides a number having fewer 1-bits: 6700417 divides 2^32 + 1, 13264529 divides 331613225, 20394401 divides 1611157679, and 97685839 divides 18014398643699713. He conjectures that 616318177 is the next term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 17 2008, Jul 21 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004738 | OEIS-A143062

Expansion of false theta series variation of Euler's pentagonal number series in powers of x.

A conjectural g.f: 1 + Sum_{n >= 0} (-1)^n*x^(2*n-1)/Product_{k = 1..n} 1 + x^(2*k-1). Conjectural g.f.: (1/(1 + x)) * (2 - Sum_{n >= 0} (-1)^n * x^(3*n)/Product_{k = 1..n} 1 + x^(2*k)). - _Peter Bala_, Jan 19 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Jul 21 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004739 | OEIS-A143105

Let g_0(x,y)=x, g_1(x,y)=x^3+3xy and g_{n+2}(x,y) = (x^2+2y)g_{n+1}(x+y)-y^2g_n(x,y). The entries of the sequence are those odd d for which g_d(x,y) and cx^jy^kg_m(x,y) have at least two terms in common (same coefficients) for some c > 0 and integers j,k and such that g_d(x,y) + cx^jy^k(1+y^m - g_m(x,y)) has all positive coefficients.

It is unknown if this sequence is infinite (conjectured to be so). Furthermore A143106 is definitely a subsequence of this sequence, but it is unknown if the two are in fact equal.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jiri Lebl (jlebl(AT)math.uiuc.edu), Jul 25 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004740 | OEIS-A143106

Odd degrees for which (up to swapping of variables) there exists a unique polynomial p(x,y), such that p(x,y)=1 when x+y=1, with positive coefficients and such that the number of terms is minimal (equal to (d+3)/2). There always exists a group invariant polynomial (see any of the references), but for many degrees, other such extremal polynomials exist.

This sequence is a subsequence of A143105. It is unknown if this is the same sequence, nor if this sequence is infinite (conjectured to be such). It is not currently computationally feasible to find out if 21 belongs in this sequence or not.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jiri Lebl_, Jul 25 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004741 | OEIS-A143109

Let H(2,d) be the space of polynomials p(x,y) of two variables with nonnegative coefficients such that p(x,y)=1 whenever x + y = 1. a(n) is the number of different polynomials in H(2,d) with exactly n distinct monomials and of maximum degree minus two, i.e., of degree 2n-5.

It is unknown but conjectured that this is a sequence of finite numbers. Note that if we went one degree lower and look at polynomials of degree 2n-6, then there are infinitely many if any exist in H(2,d).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Jiri Lebl (jlebl(AT)math.uiuc.edu), Jul 25 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004742 | OEIS-A143132

Binomial transform of [1, 5, 15, 35, 70, 0, 0, 0, ...].

Conjecture: rightmost digit of terms is cyclic: (1, 6, 6, 6, ... repeat).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Jul 27 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004743 | OEIS-A143138

E.g.f.: A(x) = x + (exp(A(x)) - 1)^2.

Conjecture: the sequence {a(n)} reduced modulo a positive integer k is eventually periodic with the period dividing phi(k), where phi(n) = A000010(n). For example, modulo 24 the sequence becomes [1, 2, 18, 14, 18, 14, 18, 14, ...] with an apparent period of 2 = phi(24)/4 starting at n = 2. - _Peter Bala_, Feb 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 27 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004744 | OEIS-A143139

E.g.f.: A(x) = exp(x + A(x)^2) - 1.

Conjecture: the sequence {a(n)} reduced modulo a positive integer k is eventually periodic with the period dividing phi(k), where phi(n) = A000010(n). For example, modulo 14 the sequence becomes [1, 3, 11, 1, 13, 1, 1, 3, 11, 1, 13, 1, 1, ...] with an apparent period of 6 = phi(14) starting at n = 2. - _Peter Bala_, Feb 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 27 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004745 | OEIS-A143412

Main diagonal of A143410.

a(n) is the expectation of U_{2n}(X) where X is a standard Gaussian random variable and U_n is the n-th Chebyshev polynomial of second kind (conjectured). - _Benjamin Dadoun_, Dec 16 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Aug 14 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004746 | OEIS-A143413

Apéry-like numbers for the constant e: a(n) = 1/(n-1)!*Sum_{k = 0..n+1} (-1)^k*C(n+1,k)*(2*n-k)! for n >= 1.

Conjectural congruences: for r >= 0 and odd prime p, calculation suggests that a(p^r*(p+1)) + a(p^r) == 0 (mod p^(r+1)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Aug 14 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004747 | OEIS-A143415

Another sequence of Apery-like numbers for the constant 1/e: a(n) = 1/(n+1)!*Sum_{k = 0..n-1} C(n-1,k)*(2*n-k)!.

Conjectural congruences: for r >= 0 and prime p, calculation suggests the congruences a(p^r*(p+1)) == a(p^r) (mod p^(r+1)) may hold.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Aug 14 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004748 | OEIS-A143510

Numbers m such that the equation phi(x) = m has even but no odd solutions.

In the unlikely event that Carmichael's conjecture is proved false, the counterexamples will be in this sequence. The number a(1) = 16842752 = 257*2^16 is mentioned in problem E3361. If there are only five Fermat primes, then 2^k is in this sequence for all k>31. It appears that for every product d of Fermat primes (A143512), the number 2^k * d is in this sequence for some k. The link to "Numbers Like 16842752" lists examples for various d. Conjecture: if the least solution to phi(x) = m is even, then m is in this sequence. - _Jianing Song_, Nov 07 2022 The conjecture holds for the first 26 such numbers, see A387221. - _Jud McCranie_, Dec 17 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Aug 21 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004749 | OEIS-A143512

Numbers of the form 3^a * 5^b * 17^c * 257^d * 65537^e; products of Fermat primes.

If the well-known conjecture that there are only five prime Fermat numbers F_k = 2^(2^k) + 1, k=0,1,2,3,4, is true, then we have exactly Sum_{n>=1} 1/a(n) = Product_{k=0..4} F_k/(F_k-1) = 4294967295/2147483648 = 1.99999999953433871269226074218 75. - _Vladimir Shevelev_ and _T. D. Noe_, Dec 01 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Aug 21 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004750 | OEIS-A143513

Numbers of the form 2^a * 3^b * 5^c * 17^d * 257^e * 65537^f; products of 2 and the Fermat primes.

Sum_{a(n) is odd} 1/a(n) = Sum_{a(n) is even} 1/a(n). If there are only five Fermat primes: 3,5,17,257,65537 (this is a well-known conjecture), then we have exactly Sum_{n>=1} 1/a(n) = 4294967295/1073741824 = 3.99999999906867742538452148 4375, which is twice the sum of the reciprocals of A143512. - _Vladimir Shevelev_ and _T. D. Noe_, Dec 01 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Aug 21 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004751 | OEIS-A143514

Array D of denominators of Best Remaining Lower Approximates of x=(1+sqrt(5))/2, by antidiagonals.

(5) Conjecture: Every (N(n,k+1)-N(n,k))/(D(n,k+1)-D(n,k)) is a principal convergent to x. (6) Suppose n>=1 and p/q and s/t are consecutive terms in row n of R. Then (conjecture) q*s-p*t=n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 22 2008, Aug 25 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004752 | OEIS-A143515

Array D of denominators of Best Remaining Upper Approximates of x=(1+sqrt(5))/2, by antidiagonals.

(5) Conjecture: Every (N(n,k+1)-N(n,k))/(D(n,k+1)-D(n,k)) is a principal convergent to x. (6) Suppose n>=1 and p/q and s/t are consecutive terms in row n of R. Then (conjecture) p*t-q*s=n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004753 | OEIS-A143527

Array D of denominators of Best Remaining Lower Approximates of x=sqrt(2), by antidiagonals.

(5) Conjecture: Every (N(n,k+1)-N(n,k))/(D(n,k+1)-D(n,k)) is an upper principal convergent to x. (6) Suppose n>=1 and p/q and s/t are consecutive terms in row n of R. Then (conjecture) q*s-p*t=n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004754 | OEIS-A143528

Array D of denominators of Best Remaining Upper Approximates of x=sqrt(2), by antidiagonals.

(5) Conjecture: Every (N(n,k+1)-N(n,k))/(D(n,k+1)-D(n,k)) is a lower principal convergent to x. (6) Suppose n>=1 and p/q and s/t are consecutive terms in row n of R. Then (conjecture) p*t-q*s=n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 23 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004755 | OEIS-A143548

Irregular triangle of numbers k < p^2 such that p^2 divides k^(p-1)-1, with p=prime(n).

Row n begins with 1 and has prime(n)-1 terms. The first differences of each row are symmetric. For k > p^2, the solutions are just shifted by m*p^2 for m > 0. An open question is whether every integer appears in this sequence. For instance, 2 does not appear until the prime 1093 and 5 does not appear until the prime 20771.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Aug 24 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004756 | OEIS-A143586

Length of row n of the Kolakoski fan A143477.

Conjecture: if L(n)=A143586(n), then lim(L(n+1)/L(n)) = 3/2. For a more general conjecture see A143589.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 25 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004757 | OEIS-A143589

Kolakoski fan based on A000034 with initial row 1.

Conjecture (following Benoit Cloitre's conjecture at A111090): if L(n) is the number (assumed finite) of terms in row n of K, then L(n)*(2/3)^n approaches a constant. (L= A143590.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 25 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004758 | OEIS-A143590

Length of row n of A143589 (a Kolakoski fan).

Conjecture (following Benoit Cloitre's conjecture at A111090): a(n)*(2/3)^n approaches a constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 25 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004759 | OEIS-A143727

Near-prime numbers: numbers n such that n-2=prime, n+2=prime, n-4=prime, n+4=prime, n-8=prime, n+8=prime.

Conjecture: all terms are multiples of 15. - _Harvey P. Dale_, Aug 03 2026 This conjecture is true. Checking all residues mod 15 shows that for all residues but 0 one of the numbers is a multiple of 3 or a multiple of 5 hence not prime. - _David A. Corneth_, Aug 03 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Joseph Stephan Orlovsky_, Aug 30 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004760 | OEIS-A143772

If m is the n-th composite, then a(n) = gcd(k + m/k), where k is over all divisors of m.

Conjecture: All even numbers are terms and the only odd numbers which are terms are 1 and 3. - _Robert G. Wilson v_, Sep 08 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Aug 31 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004761 | OEIS-A143805

Eigensequence of triangle A130534.

Conjecture: a(n) = R(n,0) where R(n,k) = R(n-1,n-1) + Sum_{j=0..k-1} (j+1)*R(n-1,j) for 0 <= k <= n with R(0,0) = 1. - _Mikhail Kurkov_, Jul 18 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Sep 01 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004762 | OEIS-A143836

Triangle read by rows: T(r,c) = (prime(r+2) + prime(c+1))/2.

The number of appearances of m >= 1 in this sequence is A061357(m). Conjecture: Every integer >= 4 appears at least once in this sequence. - _Ya-Ping Lu_, Mar 05 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Sep 02 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004763 | OEIS-A143898

Number of primes between n^K and (n+1)^K, where K = log(1151)/log(95).

This value of K is conjectured to be the least possible such that there is at least one prime in the range n^K to (n+1)^K for n > 0. This value of K was found using exact interval arithmetic. For each n <= 110 and for each prime p in the range n to n^1.7, we computed an interval k(n,p) such that p is between n^k(n,p) and (n+1)^k(n,p). The intersection of all these intervals produces a list of intervals. The least value in those intervals is K, which is log(1151)/log(95). We computed 10^5 terms of this sequence to give us confidence that a(n) > 0 for all n. 1151 counts as a qualifying prime towards both a(94)=1 and a(95)=3, in accordance with use of closed ranges. If prime p were counted only when n^K < p <= (n+1)^K, then term 95 would be 2. If prime p were counted only when n^K <= p < (n+1)^K, then term 94 would be 0. The conjecture in the author's comment implies K is the greatest real value such that for all e <= K there exists n > 0 with no prime p satisfying n^e <= p < (n+1)^e. - _Peter Munn_, Mar 02 2017 The author's description of the calculation of K implies that K is not an isolated qualifying value; equivalently that K is also the least real value for which there is a positive epsilon such that for all exponent e, K <= e <= K+epsilon and integer n > 0 there is a prime p satisfying n^e <= p <= (n+1)^e. This is a necessary precondition for my Mar 02 2017 deduction from the author's conjecture. - _Peter Munn_, Aug 21 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Sep 04 2008, Sep 26 2009, Oct 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004764 | OEIS-A143902

Rectangular array R by antidiagonals: R(m,n) = number of black squares.

Antidiagonal sums: (2,4,10,16,28,40,60,80,...)=A006584 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 04 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004765 | OEIS-A143935

Number of primes between n^K and (n+1)^K, inclusive, where K=log(127)/log(16).

This value of K is conjectured to be the least possible such that there is at least one prime in the range n^k and (n+1)^k for all n>0 and k>=K. This value of K was found using exact interval arithmetic. For each n <= 300 and for each prime p in the range n to n^2, we computed an interval k(n,p) such that p is between n^k(n,p) and (n+1)^k(n,p). The intersection of all these intervals produces a list of 29 intervals. The last interval appears to be semi-infinite beginning with K, which is log(127)/log(16). See A143898 for the smallest number in the first interval.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Sep 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004766 | OEIS-A143955

Sum of the altitudes of the leftmost valleys of all Dyck paths of semilength n (if path has no valley, then this altitude is taken to be 0).

Conjecture: (n+2)*a(n) -4*(2*n+1)*a(n-1) +2*(10*n-9)*a(n-2) +17*(2-n)*a(n-3) +2*(2*n-7)*a(n-4)=0. - _R. J. Mathar_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Oct 14 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004767 | OEIS-A143991

Numerators of numbers with g.f. exp(1-(1-x)^(1/2)).

Conjecture: a(n) = numerator((e/Pi)*Integral_{x=-oo..+oo} cos(x)/(1 + x^2)^n dx) for n > 0. See A381845 for the numerators of this integral. - _Stefano Spezia_, Mar 12 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 02 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004768 | OEIS-A144014

E.g.f. satisfies: A'(x) = 1 + x*A(x)^4 where A(0) = 1.

a(n) ~ n^(n-1/6) * sqrt(2*Pi) / (3^(1/3) * GAMMA(1/3) * exp(n) * r^(n+2/3)), where r = 0.52731343741213... (multiplicative constant is conjectured, holds 14 decimal places). - _Vaclav Kotesovec_, Feb 24 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 10 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004769 | OEIS-A144084

T(n,k) is the number of partial bijections of height k (height(alpha) = |Im(alpha)|) of an n-element set.

Conjecture: T(n, k) = Sum_{j=0..k} |Stirling1(n, n-j)| * Stirling2(n-j+1, n-k+1). - _Mikhail Kurkov_, Jun 23 2026 [The conjecture is true. Both k!*C(n, k)^2 and the claimed sum satisfy the same recurrence relation and the initial condition T(n, 0) = 1, thus they are equal for all 0 <= k <= n by induction. - _Peter Luschny_, Jun 23 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Abdullahi Umar_, Sep 10 2008, Sep 30 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004770 | OEIS-A144135

a(n) = GegenbauerC[n,2,8].

Conjecture: a(n) = 32*a(n-1)-258*a(n-2)+32*a(n-3)-a(n-4). - _Colin Barker_, Oct 27 2014 Conjecture and empirical g.f. verified. - _Robert Israel_, Jul 26 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Joseph Stephan Orlovsky_, Sep 11 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004771 | OEIS-A144247

Prime numbers of the form 11^k +- 10.

It is conjectured that 131 is the only palindrome in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reikku Kulon_, Sep 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004772 | OEIS-A144264

Numbers k such that A014085(k) = A014085(k+1).

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Sep 16 2008, Sep 17 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004773 | OEIS-A144301

a(0) = a(1) = 1; thereafter a(n) = (2*n-3)*a(n-1) + a(n-2).

G.f.: conjecture: 1 + x*(1-x)/(1-3*x+x^2)*Q(0), where Q(k) = 1 - 2*(k+1)*(2*k+1)*x^2/(2*(k+1)*(2*k+1)*x^2 - (1 - 3*x + x^2 - 4*x*k)*(1 - 7*x + x^2 - 4*x*k)/Q(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 19 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Applegate_ and _N. J. A. Sloane_, Dec 07 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004774 | OEIS-A144313

Prime numbers p such that p - 1 is the fourth a-figurate number, seventh b-figurate number and possibly tenth c-figurate number for some a, b and c and not a d-figurate number for any nontrivial d.

It is conjectured:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reikku Kulon_, Sep 17 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004775 | OEIS-A144315

Prime numbers p such that p - 1 is the fourth a-figurate number, eighth b-figurate number and possibly sixteenth c-figurate number for some a, b and c and not a d-figurate number for any nontrivial d.

It is conjectured:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reikku Kulon_, Sep 17 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004776 | OEIS-A144325

Prime numbers p such that p - 1 is the third a-figurate number, sixth b-figurate number and possibly twelfth c-figurate number for some a, b and c and not a d-figurate number for any nontrivial d.

It is conjectured:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reikku Kulon_, Sep 17 2008, Sep 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004777 | OEIS-A144326

Prime numbers that cannot be Mersenne prime exponents, by conjecture of A144325.

Prime numbers that cannot be Mersenne prime exponents, by conjecture of A144325. Conjecture: 191, 1217, 1559 and 1901 are not in fact members of this sequence, noting that they are (4, 19) k-figurate numbers; 19 is a member of A138694. Determining whether a Mersenne prime exponent one greater than a (4, 19) k-figurate number exists is sufficient to determine whether these primes are members.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reikku Kulon_, Sep 17 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004778 | OEIS-A144327

Prime numbers p such that p - 1 is the fourth a-figurate number and nineteenth b-figurate number for some a and b.

Sequence of primes sufficient to determine the conjecture of A144326. Presumed to include many terms that cannot be Mersenne prime exponents, noting A096676: either infinitely many, or all.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reikku Kulon_, Sep 17 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004779 | OEIS-A144406

Rectangular array A read by upward antidiagonals: entry A(n,k) in row n and column k gives the number of compositions of k in which no part exceeds n, n>=1, k>=0.

Conjecture: Since the array row sequences successively tend to A000079, the absolute values of nonzero differences between two successive row sequences tend to A045623 = {1,2,5,12,28,64,144,320,704,1536,...}, as k -> infinity. - _L. Edson Jeffery_, Dec 26 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_ and _Gary W. Adamson_, Sep 29 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004780 | OEIS-A144437

Period 3: repeat [3, 3, 1].

Conjecture: a(n) is the separatix. See A045944.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Oct 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004781 | OEIS-A144503

Sum of n-th antidiagonal of array in A144502.

Conjecture: a(n) = 1 + Integral_{x=0..oo} ChebyshevU(n - 1, x + 1)/exp(x) dx, where ChebyshevU(n, x) is the n-th Chebyshev polynomial of the second kind with variable x. - _Velin Yanev_, Jan 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Applegate_ and _N. J. A. Sloane_, Dec 13 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004782 | OEIS-A144559

a(n) = number of triples [i,j,k] with i+j+k = n, i an odd prime, j an odd Fibonacci number and k a positive Fibonacci number.

Zhi-Wei Sun conjectured on the Number Theory Mailing List that a(n) > 0 for all n > 4. The conjecture has been verified by _D. S. McNeil_ for all n < 10^13.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004783 | OEIS-A144575

E,g.f.: exp(1-sqrt(1-2*x-3*x^2)).

Conjecture D-finite with recurrence: a(n) +(-2*n+3)*a(n-1) +(-3*n^2+12*n-10)*a(n-2) +3*(-2*n+3)*a(n-3) -9*(n-1)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Jan 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004784 | OEIS-A144590

Number of ordered ways of writing 2n+1 = i + j, where i is a prime and j is of the form k*(k+1), k > 0.

Based on a posting by _Zhi-Wei Sun_ to the Number Theory Mailing List, Mar 23 2008, where he conjectures that a(n) > 0 for n >= 2. _Zhi-Wei Sun_ has offered a monetary reward for settling this conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 15 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004785 | OEIS-A144610

Sturmian word of slope e.

Conjecture: a(n) = A085369(n). - _R. J. Mathar_, Jan 27 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004786 | OEIS-A144611

Sturmian word of slope 2-sqrt(2).

Conjecture: a(n) = floor((n+1)*log(3)/log(2)) - floor(n*log(3)/log(2)) - 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004787 | OEIS-A144621

a(0) = a(1) = 1 and a(n+2) = a(n+1) * (3*a(n+1) - 2*a(n)^2), n>=0.

Conjecture: a(n+1) = determinant of 2^n x 2^n matrix M where M(i,j) = 1 + exponent of 2 in the factorization of i+j-1 = A001511(i+j-1), i,j>0, n>2. - _Ralf Stephan_, Sep 27 2013 The conjecture above is true. The matrix M(n+1) is equivalent via an even number of row and column interchanges to the matrix T(n+1) defined by: T(0)=1; T(n) = I(2) X T(n-1) + 1(2^n)(1(2^n))^T, where X is the Kronecker product, I(2) is the 2 X 2 identity matrix, 1(k) is a column vector of k 1s and T is matrix transpose. All rows of this matrix have the same sum (n+1) + sum_{j=1..n}j*2^(n-j) = 2^(n+1)-1 hence this is the eigenvalue of the eigenvector 1(2^n). Now det(T(n)) = (det(T(n-1)))(det(T(n-1)+2*1(2^(n-1))(1(2^(n-1)))^T)) = (det(T(n-1))^2)(1+2*(1(2^(n-1)))^T (T(n-1))^{-1} 1(2^(n-1))) = (det(T(n-1))^2)((2-2^(-n))/(1-2^(-n))). Let v(n)=(2-2^(-(n+1)))/(1-2^(-(n+1))) and notice that v(n) = 3-2/(v(n-1)). Then det(T(n)) = (det(T(n-1))^2)(3-2/v(n-1)) = (det(T(n-1))^2)(3-2(det(T(n-2))^2)/(v(n-1)det(T(n-2))^2))= det(T(n-1))(3det(T(n-1))-2det(T(n-2))^2). In the closed-form product formula we see the eigenvalues of T(n). The matrix T(n) is the matrix of times of most recent common ancestors for a 2^n tips, fully balanced binary phylogenetic tree with unit branch lengths. - _Krzysztof Bartoszek_, Mar 23 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 16 2009, based on email from Lionel Levine (levine(AT)Math.Berkeley.EDU), May 04 2006 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004788 | OEIS-A144636

Column 1 of triangle in A144633.

Conjecture: 6^(n-1)*a(n)/n! is odd iff n is a power of 2. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004789 | OEIS-A144637

Column 2 of triangle in A144633.

Conjecture: (1/18)*6^(n)*a(n)/n! is odd iff n = 2^k for k >= 1. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004790 | OEIS-A144642

Positive numbers that are not the sum of a triangular number, a square and a Fibonacci number.

Prompted by a conjecture of _Zhi-Wei Sun_ (now known not to be true) that every number can be written in the stated form.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Kurt Foster (drsardonicus(AT)earthlink.net), Jan 25 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004791 | OEIS-A144652

Triangle, read by rows, where T(m,n) = floor((2mn+m+n)/2) with m >= n >= 1.

Conjecture: If h does not belong to the sequence, then 4*h+1 is prime. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Jan 27 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004792 | OEIS-A144686

Maximum size of a connected acyclic domain of permutations of n elements with diameter n*(n-1)/2.

a(n) is at most 2.487^n and at least 2.076^n for large enough n (see Felsner & Valtr). Originally conjectured to equal A144685, but in fact a(n) is asymptotically larger and exceeds A144685 at least for n >= 34 (see Karpov & Slinko). - _Clayton Thomas_, Aug 19 2019 [Updated by _Andrei Zabolotskii_, Dec 31 2023]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004793 | OEIS-A144709

Integers m > 1 such that there is not a pair of primes (m^2+m-k, m^2+m+k), 1 <= k < m, between successive squares m^2 and (m+1)^2.

Conjecture: the intervals (n^2,(n+1)^2) that do not contain at least one such pair are sparse as n gets large.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Tisdale_, Sep 19 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004794 | OEIS-A144743

Recurrence sequence a(n)=a(n-1)^2-a(n-1)-1, a(0)=3.

Conjecture: A130282 and this sequence are disjoint. If this is true, for n >= 1, a(n+1) is the smallest m such that (m^2-1) / (a(n)^2-1) + 1 is a square. - _Jianing Song_, Mar 19 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Sep 20 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004795 | OEIS-A144750

A098777 mod 9.

a(n) = a(n-2) + a(n-3) - a(n-5) - a(n-6) + a(n-8) for n > 7 (conjectured). G.f.: (-8*x^7 - 4*x^6 + 3*x^5 + 8*x^4 + 7*x^3 - 6*x^2 - 8*x - 1)/((x - 1)*(x + 1)*(x^6 - x^3 + 1)) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004796 | OEIS-A144751

a(1) = 3, a(n + 1) = 1 + a(n) + least odd prime factor of a(n).

The conjectured g.f. (-4*x^11 + 12*x^10 + 8*x^9 + 24*x^8 + 56*x^7 + 72*x^6 + 43*x^5 + 39*x^4 + 19*x^3 + 15*x^2 + 7*x + 3)/(4*x^12 - 5*x^6 + 1) is wrong; it matches only the first 41 terms of the sequence [_R. J. Mathar_, Jun 16 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | Peter Pein, Feb 09 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004797 | OEIS-A144812

Integers having ideal digital mean up to base 7.

There are no integers less than 2^32 for which this is true to base 8. It is believed there are either infinitely many starting at some larger n, or none. If they exist, it is conjectured that the set of all similar sequences continues at least to base ten, almost certainly to base 16 and likely to arbitrarily large b. Sequences for b at least ten have an intersection with A144777.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reikku Kulon_, Sep 21 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004798 | OEIS-A144841

a(n) = 280*C(n,9)+10*C(n,6)+6*C(n,4)+C(n,3)+1 where C = binomial.

This was formerly conjectured to be "values of maximal number of regions obtained by joining each triple of n points on a 3-dimensional sphere (S_2) by a plane", a generalization of A000127. However, members of Sugaku_day pointed out that there could be 48 (or even 54) regions for n=5 and not 41. Those counterexamples were visualized, so the conjecture was incorrect. - _Taichi Aoki_, Nov 13 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | Sep, Jorik & Noud Thijssen (sep.thijssen(AT)gmail.com), Sep 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004799 | OEIS-A144876

Maximal number of distinct polyominoes into which an n X n square can be divided.

The maximal cardinality of a set of distinct polyominoes with total area n^2 is an upper bound, and I conjecture that this bound is always attainable. - _Charlie Neder_, Mar 06 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ken Duisenberg (Ken.Duisenberg(AT)hp.com), Sep 24 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004800 | OEIS-A144981

Decimal expansion of cos(Pi/8) = cos(22.5 degrees).

Conjecture: Equals Sum_{k>=0} A210065(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Sep 28 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004801 | OEIS-A145015

Run lengths in A037861 having the same sign.

a(n) seems (conjecture) to contain only the numbers 2^n - 1 (for n>=1) and 2^n - 3 (for n>=3)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Denis Boigelot (dboigelo(AT)ulb.ac.be), Sep 29 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004802 | OEIS-A145017

Squarefree positive integers k for which k-(floor(sqrt(k)))^2 is a perfect square.

Conjecture. For every n>=1 there exist infinitely many primes p of the form 4k+1 for which for a(n) > 1 we have s*p-(floor(sqrt(s*p)))^2 is not a perfect square for s=1,...,a(n)-1 while a(n)*p-(floor(sqrt(a(n)*p)))^2 is a perfect square. (See A145016(s=1) and A145022, A145023, A145047, A145048, A145049, A145050 correspondingly for s=2, s=5, s=10, s=13, s=17, s=26.) - _Vladimir Shevelev_, Sep 30 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 29 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004803 | OEIS-A145027

a(n) = a(n-1) + a(n-2) + a(n-3) with a(1) = 2, a(2) = 3, a(3) = 4.

If the conjectured recurrence in A000382 is correct, then a(n) = A000382(n+2) - A000382(n+1), n>=4. - _R. J. Mathar_, Jan 30 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Joseph Stephan Orlovsky_, Sep 30 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004804 | OEIS-A145040

Primes p such that 2^p-1 is prime and congruent to 31 mod 5!.

Conjecture: For n > 1, the Mersenne number M(n) = 2^n - 1 is in this sequence iff 5^M(n-1) == 1 (mod M(n)). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Sep 30 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004805 | OEIS-A145047

Primes p of the form 4k+1 for which s=10 is the least positive integer such that sp-(floor(sqrt(sp)))^2 is a square.

Conjecture: The least positive integer s can take values only from A008784 (see for s=1,2,5,10 sequences A145016, A145022, A145023 and this sequence).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 30 2008, Oct 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004806 | OEIS-A145050

Primes p of the form 4*k+1 for which s=26 is the least positive integer such that s*p-(floor(sqrt(s*p)))^2 is a square.

For all primes of the form 4*k+1 not exceeding 10000 the least integer s takes only values: 1, 2, 5, 10, 13, 17, 26. These values are the first numbers in A145017 (see our conjecture at A145047).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 30 2008, Oct 03 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004807 | OEIS-A145052

One-third of the number of n X n nonnegative integer arrays with every 3 X 3 subblock summing to 1.

Conjectured g.f.: x^3*(3 + 4*x+ 4*x^2 - 5*x^3)/(1 - x - 3*x^3 + 3*x^4). - _Alexander R. Povolotsky_, Apr 15 2015 Conjectured formula: a(n) = 6*(3^(floor(n / 3))/3 - 1) + 3 + 4 * (n mod 3) * 3^(floor(n / 3))/3. - _David A. Corneth_, Apr 15 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 30 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004808 | OEIS-A145053

One-third of the number of n X n nonnegative integer arrays with every 3 X 3 subblock summing to 2.

Second differences of (a(3k), a(3k + 1), a(3k + 1)) and (a(3k + 1), a(3k + 2), a(3k + 3)) appear to be 48 * 9^(k - 1). Based on this conjecture, we would have a(15) = 306435. - _David A. Corneth_, Apr 15 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 30 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004809 | OEIS-A145110

Number of elements in the Redheffer matrix that contribute to the Moebius function.

a(11) to a(30) are conjectured to be 11, 23, 13, 18, 19, 25, 17, 29, 19, 30, 26, 26, 23, 45, 27, 30, 32, 39, 29, 48. [From _Mats Granvik_, Dec 21 2008]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Oct 02 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004810 | OEIS-A145156

Number of Greek-key tours on a 5 X n board; i.e., self-avoiding walks on 5 X n grid starting in top left corner.

G.f.: -x*(3*x^13 -3*x^12 +17*x^11 -11*x^10 +11*x^9 -21*x^8 +67*x^7 -29*x^6 -65*x^5 +45*x^4 +8*x^3 -4*x^2 -x -1) / ((x +1)*(x^6 -x^5 +8*x^4 -8*x^3 -2*x^2 +5*x -1)*(2*x^6 +11*x^2 -1)). [conjectured by _Colin Barker_, Nov 09 2015; proved by _Jay Pantone_, Klotz, and Sullivan, Aug 01 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nathaniel Johnston_, Oct 03 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004811 | OEIS-A145215

a(n) is the minimal prime of the form 4k+1 for which s=A008784(n) is the minimal positive integer such that s*a(n)-floor(sqrt(s*a(n)))^2 is a square.

See the conjecture in the comment at A145047. In addition, I conjecture that for every such s there exist infinitely many primes of the form 4k+1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Oct 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004812 | OEIS-A145236

a(n) is the least positive integer such that if p_n is the n-th prime then (ceiling(sqrt(a(n)*p_n)))^2 - a(n)*p_n is a perfect square.

Conjectures: 1) for n >= 2, the sequence does not decrease; 2) for n > 1, a(n) is odd; 3) a(n) can be equal to a(n+1) only for twins: p_(n+1) - p_n = 2 (although there also exist twins

for which a(n) < a(n+1)). All these conjectures are proved using the formula a(n) = p_n - 2*floor(sqrt(2p_n)) + 2, n > 1. See also A145701 and A145714. - _Vladimir Shevelev_, Oct 18 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Oct 05 2008, Oct 07 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004813 | OEIS-A145300

a(n) is the maximal prime such that if p_n is the n-th prime then ceiling(sqrt(a(n)*p_n))^2 - a(n)*p_n is a perfect square.

Or a(n) is the maximal prime q_n > p_n such that sqrt(q_n)-sqrt(p_n) < sqrt(2) [or (p_n+q_n)/2 < sqrt(p_n*q_n)+1]. I conjecture that lim_{n->infinity} (sqrt(q_n) - sqrt(p_n)) = sqrt(2). Note that in the considered case this conjecture is equivalent to the following: lim_{n->infinity} fract(sqrt(p_n*q_n)) = 0, where fract(x) denotes the fractional part of x. - _Vladimir Shevelev_, Oct 09 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Oct 06 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004814 | OEIS-A145354

It is conjectured that for each m >= 1 there exist primes Q=Q(m) and P=P(m) with (2m)^2 + 1 <= Q <= (2m+1)^2 - 2m <= P <= (2m+1)^2; then set a(2m-1) = Q, a(2m) = P.

It is conjectured that for each m >= 1 there exist primes Q=Q(m) and P=P(m) with (2m)^2 + 1 <= Q <= (2m+1)^2 - 2m <= P <= (2m+1)^2; then set a(2m-1) = Q, a(2m) = P.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Mar 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004815 | OEIS-A145376

If T(n)>=n is the nearest triangular number to n and p_n is the n-th prime, then a(n) is the least prime such that T(a(n)*p_n)-a(n)p_n is a triangular number.

Conjecture. For n>=2 the sequence is nondecreasing. The conjecture holds for the first 10000 terms. [_Charles R Greathouse IV_, Feb 21 2011]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Oct 09 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004816 | OEIS-A145384

The number of terms of A050791 bracketed by successive terms of A141326.

a(107),a(108) are the first occurrence of 2 consecutive 0's and a(119),a(120),a(121) are the first occurrence of 3 consecutive 0's. This leads to the following conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | Lewis Mammel (l_mammel(AT)att.net), Oct 10 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004817 | OEIS-A145552

Square numbers which can be expressed as the sum of 2 positive cubes in exactly 2 different ways.

This is conjectured to be an infinite sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Iain Renfrew (iain.renfrew(AT)btinternet.com), Oct 13 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004818 | OEIS-A145553

Numbers n such that n^2 can be expressed as the sum of 2 positive cubes in exactly 2 different ways.

This is conjectured to be an infinite sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Iain Renfrew (iain.renfrew(AT)btinternet.com), Oct 13 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004819 | OEIS-A145646

Wavelength (in ångströms) of the series limit of the Hydrogen spectrum for main quantum number n.

Conjectures from _Colin Barker_, May 07 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Oct 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004820 | OEIS-A145647

First differences of A145646.

Conjectures from _Colin Barker_, May 08 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Oct 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004821 | OEIS-A145656

a(n) = numerator of polynomial of genus 1 and level n for m = 2.

Conjecture: a(n) = numerator((2^n)*log(2) - 2^(n+1)*Integral_{x = 0..1} x^(2*n-1)/(1 + x^2)^n). - _Peter Bala_, Jun 10 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Oct 16 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004822 | OEIS-A145679

Lower limit of backward value of 2^n and n!.

The sum of constants in this sequence and A023415 is conjectured to be 11 exactly. The equality of the backward sequences for n! omitting the trailing sequence zeros and 2^n holds under the conjecture that for each k and each r not divisible by 5 there are infinitely many n for which n! without trailing zeros = r mod 5^k. Actually this conjecture is true because for m > 0 g(k)^m = ((5^k)(5^(4m) - 1)/(5^4 - 1))! without leading zeros mod 5^k, and g(k) is a generator of the multiplicative group mod 5^k for k=3 mod 4 by induction using g(k+4) = g(k) mod 5^k. The above conjecture about the sum of constants in this sequence and A023415 can be proved using the recursion formulas for the sequences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_, Mar 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004823 | OEIS-A145731

Conjectured integers m such that |A144912(2, p^m)| > 0 for all Mersenne prime exponents p > 7.

Conjectured integers m such that |A144912(2, p^m)| > 0 for all Mersenne prime exponents p > 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reikku Kulon_, Oct 17 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004824 | OEIS-A145819

Union of A145812 and A145818 with double repetition of 1, so that a(1)=1, a(2)=1.

Theorem. For every even integer m there exists a representation of the form m=a(r)+a(s). If A(x) is the counting function of a(n)<=x, then A(x)=O(sqrt(x))and Omega(sqrt(x)). Conjecture. The sequence is minimal in the following sense: if any sequence has the counting function B(x)<=A(x) for all x>=1 and B(x) < A(x) for x>=x_0, then there exists an even integer N which is not expressible as a sum of two terms of such sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Oct 20 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004825 | OEIS-A145824

Lower twin primes p1 such that p1-1 is a square.

Conjecture: There exist an infinite number of primes of this form.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Oct 20 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004826 | OEIS-A145855

Number of n-element subsets of {1,2,...,2n-1} whose elements sum to a multiple of n.

a(n) = (1/(2*n))*Sum_{d|n} (-1)^(n+d)*phi(n/d)*binomial(2*d,d). Conjectured by _Vladeta Jovovic_, Oct 22 2008; proved by _Max Alekseyev_, Oct 23 2008 (see link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Oct 21 2008, Oct 22 2008, Oct 24 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004827 | OEIS-A145875

Repdigit Kaprekar numbers.

Conjectures from _Daniel Mondot_, Mar 28 2025: (Start) The above two conjectures are linked, they are easily proved using modular arithmetic, and correspond to the explicit formula given below. - _M. F. Hasler_, Mar 28 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | Howard Berman (howard_berman(AT)hotmail.com), Oct 22 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004828 | OEIS-A145878

Triangle read by rows: T(n,k) is the number of permutations of [n] having k strong fixed points (0 <= k <= n).

T(3+k,k)=3k+3, T(4+k,k)=(k+1)(k+28)/2, T(5+k,k)=(k+1)(3k+77), T(6+k,k)=(k+1)(k^2+110k+2982)/6, T(7+k,k)=(k+1)(3k^2+235k+7352)/2 (previous conjectures).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Oct 29 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004829 | OEIS-A145879

Triangle read by rows: T(n,k) is the number of permutations of {1,2,...,n} having exactly k entries that are midpoints of 321 patterns (0 <= k <= n-2 for n >= 2; k=0 for n=1).

The following formulas are conjectural and assume different offsets: Conjecture: T(n, k) = [z^k] R_1(n-1, 0) where R_1(n, q) = (q*z + 1)*R_1(n-1, q+1) + Sum_{j=0..q} R_1(n-1, j) for n > 0, q >= 0 with R_1(0, q) = 1 for q >= 0. - _Mikhail Kurkov_, Dec 26 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Oct 30 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004830 | OEIS-A145920

List of numbers that are both pentagonal (A000326) and binomial coefficients C(n,4) (A000332).

Conjecture: a(n) = a(n-1) + 4a(n-2) - 4a(n-3) - 6a(n-4) + 6a(n-5) + 4a(n-6) - 4a(n-7) - a(n-8) + a(n-9). - _R. J. Mathar_, Oct 29 2008 Conjecture: G.f.: x^2(1 + 4x + 26x^2 + 19x^3 + 4x^5 + x^6 + 26x^4)/((1+x)^4(1-x)^5). - _R. J. Mathar_, Oct 29 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthew Vandermast_, Oct 28 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004831 | OEIS-A145996

Numbers k such that quintic polynomial 4*k - k^2 + 5*k^2*x + (20*k - 20*k^2)*x^3 + (16 - 32*k + 16*k^2)*x^5 has a rational root.

Conjecture: This sequence is finite and complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Oct 26 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004832 | OEIS-A146025

Numbers that can be written in bases 2, 3, 4, and 5 using only the digits 0 and 1.

Conjectured to be complete. a(4), if it exists, is greater than 10^15. - _Charles R Greathouse IV_, Apr 06 2012 It is a plausible conjecture that there are no more terms, but this has not been proved. - _N. J. A. Sloane_, Feb 06 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Mondot_, Oct 26 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004833 | OEIS-A146026

Numbers that can be written from base 2 to base 9 using only the digits 0 to 3.

Conjectured to be complete. It is a plausible conjecture that there are no more terms, but this has not been proved. - _N. J. A. Sloane_, Nov 17 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Mondot_, Oct 27 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004834 | OEIS-A146028

Numbers that can be written from base 2 to base 15 using only the digits 0 to 7.

It is a plausible conjecture that there are no more terms, but this has not been proved. - _N. J. A. Sloane_, Nov 17 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Mondot_, Nov 02 2008, Nov 06 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004835 | OEIS-A146029

Numbers that can be written from base 2 to base 17 using only the digits 0 to 8 (conjectured to be complete).

Numbers that can be written from base 2 to base 17 using only the digits 0 to 8 (conjectured to be complete). It is a plausible conjecture that there are no more terms, but this has not been proved. - _N. J. A. Sloane_, Nov 17 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Mondot_, Nov 02 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004836 | OEIS-A146107

Bell numbers (A000110) read mod 17.

Conjecture: a(n+51702516367896047761) = a(n). - _G. C. Greubel_, Feb 01 2016 [Conjecture is correct; see A054767. - _Jianing Song_, Jun 20 2025]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004837 | OEIS-A146109

Bell numbers (A000110) read mod 19.

Conjecture: a(n+1099122030922396438402221) = a(n). - _G. C. Greubel_, Feb 01 2016 [Conjecture is correct; see A054767. - _Jianing Song_, Jun 20 2025]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004838 | OEIS-A146113

Bell numbers (A000110) read mod 23.

Conjecture: a(n+94911218181126872883431967753) = a(n). - _G. C. Greubel_, Feb 02 2016 [Conjecture is correct; see A054767. - _Jianing Song_, Jun 20 2025]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004839 | OEIS-A146119

Bell numbers (A000110) read mod 29.

Conjecture: a(n+91703076898614683377208150526107718802981) = a(n). - _G. C. Greubel_, Feb 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004840 | OEIS-A146121

Bell numbers (A000110) read mod 31.

Conjecture: a(n+5689724710241078652870214343019771 58534824481) = a(n). - _G. C. Greubel_, Feb 02 2016 [Conjecture is correct; see A054767. - _Jianing Song_, Jun 20 2025]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004841 | OEIS-A146123

Number of n X n arrays of squares of integers with every (n-3) X (n-3) subblock summing to 4 and every element equal to at least one neighbor.

Conjectures from _Colin Barker_, Mar 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004842 | OEIS-A146124

Number of n X n arrays of squares of integers summing to 5 with every element equal to at least one neighbor.

Conjectures from _Colin Barker_, Mar 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004843 | OEIS-A146125

Number of n X n arrays of squares of integers with every (n-1) X (n-1) subblock summing to 5 and every element equal to at least one neighbor.

Conjectures from _Colin Barker_, Mar 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004844 | OEIS-A146126

Number of n X n arrays of squares of integers with every (n-2) X (n-2) subblock summing to 5 and every element equal to at least one neighbor.

Conjectures from _Colin Barker_, Mar 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004845 | OEIS-A146295

Integers which are not the sum of a 4-almost prime and a prime.

Largest term is 60060 (see b-file). No more terms < 10^8. Conjectured to be complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Donovan Johnson_, Nov 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004846 | OEIS-A146296

Integers which are not the sum of a 5-almost prime and a prime.

Largest term is 3573570 (see b-file). No more terms < 10^9. Conjectured to be complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Donovan Johnson_, Nov 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004847 | OEIS-A146304

Number of distinct ways to place bishops (up to 2n-2) on an n X n chessboard so that no bishop is attacking another and that it is not possible to add another bishop.

Conjectured to be a(n) = O(n^(n-1)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Bonzini_, Oct 29 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004848 | OEIS-A146557

Number of collinear triples of distinct points in Zn x Zn with no two on the same "horizontal" or "vertical" line.

Conjecture: if 3 does not divide n then n^2 divides a(n). - _Peter Bala_, Jul 24 2025 The conjecture easily follows from the second formula for a(n). - _Max Alekseyev_, Sep 29 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Oct 31 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004849 | OEIS-A146564

a(n) is the number of solutions of the equation k*n/(k-n) = c. k,c integers.

Conjecture: a(n) = A048691(n)+A063647(n). - _R. J. Mathar_, Nov 21 2008 (See Corollary 4 in Cerruti's paper.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Nov 01 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004850 | OEIS-A146567

Numbers k such that k*sigma_0(k)/(k+sigma_0(k)) = c, c an integer.

For a given n let x be the minimal natural number such that n*x/(n+x)=c. I conjecture: from a certain n onward, x>sigma_0(n) for all n. Thus, there is no other solution bigger than 1260, and this sequence is finite. - _Ctibor O. Zizka_, Sep 13 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Nov 01 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004851 | OEIS-A146955

A functionally symmetric Polynomial as a triangle of coefficients: p(x,n)=If[n == 0, 1, (x + 1)^n + 2^(n - 4)*Sum[(2^m + 2*m)*x^m*(1 + x^(n - 2*m)), {m, 1, n - 1}]].

Conjecture: row sums are 2^n*(2^n+6-n+n^2)/8 for n>0. [From _R. J. Mathar_, Nov 30 2008]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Nov 03 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004852 | OEIS-A146961

Numbers k = p*q*r, with odd primes p < q < r, such that Sister Beiter's cyclotomic coefficient conjecture is false.

Numbers k = p*q*r, with odd primes p < q < r, such that Sister Beiter's cyclotomic coefficient conjecture is false. In 1968, Sister Beiter conjectured that for k = p*q*r, with odd primes p < q < r, the maximum coefficient (in absolute value) of the cyclotomic polynomial Phi(k,x) is <= (p+1)/2. Up to 10^6, all counterexamples have p > 7. Gallot and Moree prove the conjecture is false for p > 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Nov 03 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004853 | OEIS-A147296

a(n) = n*(9*n+2).

a(n) = n*(9*n + 2), as conjectured by V. Librandi. - _M. F. Hasler_, Mar 01 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Nov 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004854 | OEIS-A147672

a(n) = a(n-2) + 2^(n-1) + 5 for n>3, a(0..3) = (0,1,2,7).

The name comes from the puzzle "Crossing the bridge" (cf. link): If this puzzle is generalized to a group of n people who need c(0),...,c(n-1) minutes to cross the bridge, then b(n) is an upper bound for the optimal solution, conjectured to be optimal, cf. explanations from _Franklin T. Adams-Watters_ to the SeqFan list, Nov 10 2008.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Nov 10 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004855 | OEIS-A147792

A quadrisection of A061042.

Conjecture: a(n) = 3*a(n-2) -6*a(n-4) +10*a(n-6) -12*a(n-8) +12*a(n-10) -10*a(n-12) +6*a(n-14) - 3*a(n-16) +a(n-18). - _R. J. Mathar_, Nov 14 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Nov 13 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004856 | OEIS-A147796

Number of consistent sets of 3 irreflexive binary order relationships over n objects.

Conjectures from _Colin Barker_, Apr 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004857 | OEIS-A147817

Number of consistent sets of 4 irreflexive binary order relationships over n objects.

Conjectures from _Colin Barker_, Apr 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004858 | OEIS-A147821

Number of consistent sets of 5 irreflexive binary order relationships over n objects.

Conjectures from _Colin Barker_, Apr 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

B04 A150000 - A199999 | 1,627 条

V3-004859 | OEIS-A151282

Number of walks within N^2 (the first quadrant of Z^2) starting at (0,0) and consisting of n steps taken from {(-1, -1), (-1, 0), (0, 1), (1, 1)}.

Conjecture: (n+1)*a(n)-3*(2n+1)*a(n-1) +(n+10)*a(n-2) +28(n-2)*a(n-3)=0. - _R. J. Mathar_, Dec 08 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Manuel Kauers_, Nov 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004860 | OEIS-A151323

Number of walks within N^2 (the first quadrant of Z^2) starting at (0,0) and consisting of n steps taken from {(-1, -1), (-1, 0), (0, -1), (0, 1), (1, 0), (1, 1)}.

Conjecture: (n+1)*a(n) -2*(2*n+1)*a(n-1) -12*(n-1)*a(n-2) = 0. - _R. J. Mathar_, Oct 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Manuel Kauers_, Nov 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004861 | OEIS-A151341

Number of walks within N^2 (the first quadrant of Z^2) starting and ending at (0,0) and consisting of 2 n steps taken from {(-1, -1), (-1, 0), (-1, 1), (1, 0)}.

Conjecture: n*(n+2)*(n+1)*a(n) -2*n*(2*n-1)*(2*n+1)*a(n-1) -12*(n-1)*(2*n-1)*(2*n-3)*a(n-2) = 0. - _R. J. Mathar_, Jul 21 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Manuel Kauers_, Nov 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004862 | OEIS-A151366

Number of walks within N^2 (the first quadrant of Z^2) starting and ending at (0,0) and consisting of n steps taken from {(-1, 0), (-1, 1), (0, -1), (0, 1), (1, -1), (1, 0)}.

Conjecture: (n+4)*(n+3)*a(n) -n*(n-1)*a(n-1) -12*(2*n+1)*(n-1)*a(n-2) -36*(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Jul 21 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Manuel Kauers_, Nov 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004863 | OEIS-A151413

Index at which n occurs in A160256, or -1 if n never occurs.

A160218(n) = a(A000040(n)), conjectured. - _M. F. Hasler_, May 16 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 16 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004864 | OEIS-A151483

Number of walks within N^2 (the first quadrant of Z^2) starting at (0,0), ending on the vertical axis and consisting of n steps taken from {(-1, 0), (-1, 1), (0, -1), (0, 1), (1, -1), (1, 0)}.

Conjecture: +(n+3)*a(n) -4*n*a(n-1) +12*(-n+1)*a(n-2)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Manuel Kauers_, Nov 18 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004865 | OEIS-A151512

The triangle in A151359 read by rows upwards.

Conjectured: The i-th element of row j is the number of different equivalence relationships, within a set of (j-1) element, having (j-i) equivalence classes. For example: row 5 = [1, 6, 7, 1, 0] means that, in a set of 4 elements, there exists 7 equivalence relationships having 3 different equivalence classes. - _Philippe Beaudoin_, Nov 09 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004866 | OEIS-A151800

Least prime > n (version 2 of the "next prime" function).

Conjecture: if n > 3, then a(n) < n^(n^(1/n)). - _Thomas Ordowski_, Feb 23 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 29 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004867 | OEIS-A151819

First component x of pairs (x,y) where x!+y! is a square, sorted on x.

Conjecture: the only solutions for x!+y!=n^2 are x=0,0,0,1,1,1,2,4,4,4 and y=4,5,7,4,5,7,2,0,1,5 respectively.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Jul 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004868 | OEIS-A151987

Number of legal starting configurations in the glass worms (or vers de verres) game with n glasses.

It is conjectured that a(n)~(n+1)^(n+1)/exp(n). - _Dennis Hou_, May 05 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jacques Tramu_, Mar 09 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004869 | OEIS-A152009

(L)-sieve transform of {1,4,7,10,...,3n-2,...} (A016777).

The (L)-sieve transform is quite different from the transform introduced by _N. J. A. Sloane_ in A099361 and used by T. D. Noe in A100424 - A100426 and seems to lead to more interesting results and relationships among sequences. An interesting property of the (L)-sieve transform is that the (L)-sieve transform of the sequence {1,3,6,10,...,n(n+1)/2,...} of triangular numbers is again the triangular numbers. Another (conjectured) connection with the triangular numbers is given in the following. Conjecture. Let x(0) be a random sequence of positive integers and, for n>0, let x(n)=S[x(n-1)], where S is the (L)-sieve transform. Then the limit of {x(n)} as n goes to infinity is the sequence of triangular numbers {1,3,6,10,...,n(n+1)/2,...}. Illustration of the conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Nov 19 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004870 | OEIS-A152020

Denominator of 8/(9*n^2) divided by 9.

Conjecture: a(n) = denominator((n-2)^3/n^2). - _Andres Cicuttin_, Sep 19 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Nov 20 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004871 | OEIS-A152046

a(n) = Product_{k=1..floor((n-1)/2)} (1 + 8*cos(k*Pi/n)^2) for n >= 0.

Apparently the same as A001045 after the first term. - _R. J. Mathar_, Nov 27 2008 [This conjecture is true. - _Peter Luschny_, Nov 28 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Nov 21 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004872 | OEIS-A152079

Primes p such that A000695(p) are also prime.

Numbers a(n) are in A077718. Conjecture: The sequence contains infinitely many terms. Supporting the conjecture, there are 8398 terms up to a million, 56915 up to 10 million, 407875 terms up to 100 million, and 3135962 terms up to a billion. - _Charles R Greathouse IV_, Apr 28 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Nov 23 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004873 | OEIS-A152120

a(n) = 2^n * Product_{k=1..(n-1)/2} (2 + 3*cos(k*Pi/n)^2).

Conjecture: a(n) = 14*a(n-2) - 9*a(n-4) for n > 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_ and _Gary W. Adamson_, Nov 24 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004874 | OEIS-A152139

Correlation classes of pairs of different words.

Given the first terms of the sequence above, an obvious conjecture is that b(m,q) = b(m,4) for all q > 4. Conjecture that b(m,q) = b(m,4) for all q > 4, is now verified for m to and including 8. Also b(9,q)=b(9,4) for q=5,6,7,8 at least. - _Paul Leopardi_, Jun 14 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Leopardi_, Nov 26 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004875 | OEIS-A152165

Largest number which is not the sum of an n-almost prime and a prime.

All the values are conjectural and untrustworthy. - _N. J. A. Sloane_, Oct 05 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Mar 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004876 | OEIS-A152191

a(n)=Product_{k=1..floor((n-1)/2)} (1 + 4*cos(k*Pi/n)^2)*(1 - 4*sin(k*Pi/n)^2).

Conjecture: a(2n) = A108196(n-1), n>=2. a(n) = (-1)^(n+1)*A000045(n) *A101675(n-1), n>0. G.f.: 1 -x*(x-1)*(x^2-x+1)*(1+x)^3 / ((x^4-x^3+2*x^2+x+1)*(x^4+x^3+2*x^2-x+1)). - _R. J. Mathar_, Mar 08 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_ and _Gary W. Adamson_, Nov 28 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004877 | OEIS-A152225

Number of Dyck paths of semilength n with no peaks at height 0 (mod 3) and no valleys at height 2 (mod 3).

Conjecture: (n+2)*a(n) - 2*(2*n+1)*a(n-1) + 4*(n-1)*a(n-2) + 2*(5-2*n)*a(n-3)=0. - _R. J. Mathar_, Aug 14 2012 This conjecture follows from the differential equation (4*x^4-4*x^3+4*x^2-x)*y' + (2*x^3-4*x^2+6*x-2)*y - 2*x^3+2*x^2-3*x+2=0 satisfied by the g.f. - _Robert Israel_, Jan 09 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | Majun (majun(AT)math.sinica.edu.tw), Nov 29 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004878 | OEIS-A152260

Triangle T(n, k) = [x^k] p(n, x), where p(n, x) = (1/n)*(1-x)^(2*n) * Sum_{j >= 0} binomial(n+j-1, j) * j^n * x^(j-1).

T(n, m) = Sum_{k=1..m} binomial(n+k,k)*binomial(n-k,m-k)*k!*(-1)^(m-k) * Stirling2(n,k)*1/(n+k) (conjectured). - _Michael D. Weiner_, Jul 01 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Dec 01 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004879 | OEIS-A152396

Let f(M,k) denote the decimal concatenation of k numbers starting with M: M | M-1 | M-2 | ... | M-k+1, k > 1. Then a(n) is the smallest M such that for all m in {1,..,n} an m-th prime occurs as f(M,k) for the smallest possible k, order prioritized m = 1 through n.

The sequence of k values is 2, 4, 8, 10, 14, 20, 28, 32, 34, 40, 50, etc. a(n) is necessarily congruent to 10 modulo 30 starting with n=4 (and is coincidentally so for n=2 and 3). Each successive unknown term's existence is only conjectural, but is supported by standard heuristics. The values in the lead sentence come through considerations modulo the smallest primes. Concatenation of 22 numbers is ruled out by consideration modulo 11 in concert with the prior need for the stem to be 10 mod 30: A prime concatenation of 22 values necessarily involves passage from one length number to another, with a power of 10 not leading. Beginning with a(11), however, with each time a multiple of 22 for the number of concatenated values is passed there is a measure of uncertainty. Theoretically, it seems a(11) could include for largest prime a concatenation of 44 (rather than 50) values, but with these primes very sparse it is a near certainty heuristically that this is not so. Mathematically, a conjectured a(11) would have a higher bar for strict proof than a(n) for n < 11, and the same holds repeatedly for even less accessible terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James G. Merickel_, Oct 20 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004880 | OEIS-A152458

Numbers k that A064413(k) = k.

Conjecture: This sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Dec 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004881 | OEIS-A152461

Primes p such that there does not exist any positive integer k and prime q with p > q and 3^k = p + 2q or 3^k = q + 2p.

If there exists a sequence N_k = 3^n_k such that N_k has O((N_k)^v), v < 1/2, representations of the considered form, then removing the maximal primes in every such representation, we obtain an analog B of A152461 with the counting function Z(x) = pi(x) - O(x^v). Replacing in B the first N terms with N consecutive primes (with arbitrarily large N), we obtain a sequence which essentially is indistinguishable from the sequence of all primes with the help of the approximation of pi(x) by li(x), since, according to the well-known Littlewood result, the remainder term in the theorem of primes cannot be less than sqrt(x)logloglog(x)/log(x). But for this sequence we have infinitely many odd numbers which are not expressible by sum p + 2q with p, q primes. Thus in this case the Lemoine-Levy conjecture is essentially unprovable. Nevertheless, we conjecture that there does not exist a considered abnormal case of sequence (N_k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Dec 05 2008, Dec 12 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004882 | OEIS-A152466

a(1) = 252, a(n) is a(n-1) multiplied by the smallest prime factor of a(n-1)+1.

Conjecture: this sequence contains no terms k where k+1 is prime. (All similar sequences that start with numbers less than 252 are known to contain terms k where k+1 is prime.) The above conjecture has now been disproved: adding 1 to the 39th term of this sequence gives a prime number. - _Andrea Concaro_, Dec 31 2022 Conjecture: a(n) = A238642(a(n-1)). - _J. Lowell_, Mar 25 2014 [This conjecture fails at n=40; see the above comment from _Andrea Concaro_. - _Jon E. Schoenfield_, Jan 15 2023]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Dec 05 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004883 | OEIS-A152483

Even numbers which are not the sum of 2 odd semiprimes.

No more terms < 10^9. Conjectured to be complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Donovan Johnson_, Dec 06 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004884 | OEIS-A152484

Odd numbers which are not the sum of an even semiprime and an odd semiprime.

No more terms < 10^9. Conjectured to be complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Donovan Johnson_, Dec 06 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004885 | OEIS-A152491

Numbers n such that 1/c = 1/n + 1/S(n). c, n positive integers (A000027(n)), S(n) sum of digits of n (A007953(n)).

For a given n let x be the minimal natural number such that n*x/(n+x)=c. I conjecture: from a certain n onward, x>S(n) for all n. Thus there is no other solution bigger than 72, and this sequence is finite. - _Ctibor O. Zizka_, Sep 13 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Dec 06 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004886 | OEIS-A152519

Numbers k such that 2*A064413(k) = 3*k.

Conjecture: Number of points on 3 critical lines of EKG sequence is finite (and complete?). - _Artur Jasinski_, Dec 08 2008

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Dec 06 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004887 | OEIS-A152522

a(n) is the least even number such that if p_i is the i-th prime then a(n)-p_i, i=1..n, are composite numbers.

Conjecture (A. Granville, H. te Riele and J. van de Lune, 1989): Let, for even N, p=p(N) be the least prime such that N-p is prime as well. Then p(N)=O((log(N))^2log(log(N))). - _Vladimir Shevelev_, Dec 08 2008 It is conjectured that a(n) >= prime(k+1) + 3. See Conjecture C p. 2 in Phong & Dongdong. - _Michel Marcus_, Aug 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Dec 06 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004888 | OEIS-A152601

a(n) = Sum_{k=0..n} C(n+k,2k)*A000108(k)*3^k*2^(n-k).

Conjecture: (n+1)*a(n) +8*(-2*n+1)*a(n-1) +4*(n-2)*a(n-2)=0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Dec 09 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004889 | OEIS-A152650

Triangle of the numerators of coefficients c(n,k) = [x^k] P(n,x) of certain polynomials P(n,x) given below.

Conjecture: All roots of P(n,x) are real, hence negative. - _Jean-François Alcover_, Oct 10 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Dec 10 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004890 | OEIS-A152661

Number of permutations of [n] for which the first two entries have the same parity (n>=2).

Conjecture: (-n+3)*a(n) + 2*(n-2)*a(n-1) + (n-1)*(n-2)*(n-3)*a(n-2) = 0. - _R. J. Mathar_, Apr 20 2015 Conjecture: a(n) = 2*A077613(n-1). - _R. J. Mathar_, Apr 20 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 12 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004891 | OEIS-A152718

a(n) = 2*a(n-1) - a(n-2) + 2*a(n-3) - a(n-4) + a(n-5), n > 5.

Conjecture: this is the inverse Motzkin transform of A054393.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Dec 11 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004892 | OEIS-A152829

Numbers k whose squares can be written in exactly one way as a sum of three squares: k^2 = a^2 + b^2 + c^2 with 1 <= a <= b <= c.

2k is a term iff k is also a term, so the conjecture from _Colin Barker_ (see Formula) is true iff 3, 7, and 13 are the only odd terms. - _Jon E. Schoenfield_, Mar 22 2022 Conjecture: a(n) = 2*a(n-3) for n > 5. - _Colin Barker_, Mar 12 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | Peter Pein (petsie(AT)dordos.net), Dec 13 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004893 | OEIS-A152920

Triangle read by rows: triangle A062111 reversed.

Define the array B(n,p) = (Sum_{k=0..n} binomial(p+k,p) * T(n,k))/(n+p+1) for n >= 0 and p >= 0. Then see the comment of Robert Coquereaux (2014) at A193844. Conjecture: B(n+1,p) = A(n,p). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Dec 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004894 | OEIS-A152929

Number of sets (in the Hausdorff metric geometry) at each location between two sets defining a polygonal configuration consisting of two 4-gonal polygonal components chained with string components of length l as l varies.

a(n) = (163*A000045(n)+63*A000032(n))/2. - Conjectured by _Philipp Emanuel Weidmann_, cf. LINKS.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Schlicker_, Dec 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004895 | OEIS-A152930

Number of sets (in the Hausdorff metric geometry) at each location between two sets defining a polygonal configuration consisting of k 4-gonal polygonal components chained with string components of length 2 as k varies.

Conjectures from _Colin Barker_, Jul 09 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Schlicker_, Dec 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004896 | OEIS-A152932

Number of sets (in the Hausdorff metric geometry) at each location between two sets defining a polygonal configuration consisting of three 6-gonal polygonal components chained with string components of length l as l varies.

Conjectures from _Colin Barker_, Jul 09 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Schlicker_, Dec 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004897 | OEIS-A152933

Number of sets (in the Hausdorff metric geometry) at each location between two sets defining a polygonal configuration consisting of k 6-gonal polygonal components chained with string components of length 2 as k varies.

Conjectures from _Colin Barker_, Jul 09 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Schlicker_, Dec 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004898 | OEIS-A152934

Number of sets (in the Hausdorff metric geometry) at each location between two sets defining a polygonal configuration consisting of two m-gonal polygonal components chained with string components of length 3 as m varies.

Conjectures from _Colin Barker_, Jul 09 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Schlicker_, Dec 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004899 | OEIS-A152939

Number of sets (in the Hausdorff metric geometry) at each location between two sets defining a polygonal configuration consisting of four 4-gonal polygonal components chained with string components of length l as l varies.

Conjectures from _Colin Barker_, Jul 09 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Schlicker_, Dec 15 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004900 | OEIS-A152959

Number of correlation classes for pairs of different words in an alphabet of size 4.

Let b(m,q) be the number of correlation classes of pairs of different q-ary words of length m, per the definition of sequence A152139. Then here a(m)=b(m,4), the number of correlation classes of pairs of different words of length m in an alphabet of size 4. In other words, for m>1, a(m)=c(m*(m-1)+4), where c is given by A152139. A conjecture mentioned in the comments to A152139 translates here to b(m,q) = a(m) for all q > 4. For more details, see the comments in A152139.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Leopardi_, Dec 15 2008, Dec 28 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004901 | OEIS-A153010

Indices of A153007 where the entry equals zero.

Conjectures from _Colin Barker_, Nov 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Dec 21 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004902 | OEIS-A153197

Binomial transform of Bessel numbers A006789.

Conjecture: Given any sequence with Hankel transform of [1,1,1,...], performing alternate operations: binomial transform followed by INVERT transform, then binomial transform of the last result (repeat); or INVERT transform starting first, will converge upon A006789 and A153197 as a two sequence limit cycle. The conjecture can be extended to any Hankel transform (and their accompanying sequence set): analogous operations will converge upon a Bessel-type sequence and its binomial transform.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Dec 20 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004903 | OEIS-A153303

G.f.: cm4(x)^4 = Sum_{n>=0} a(n)*x^(4n)/(4n)!, where cm4(x) is defined by A153300.

Conjecture: a(n)/2^(4n-1) is an odd integer for n>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004904 | OEIS-A153367

Number of zig-zag paths from top to bottom of a rectangle of width 9 with 2n-1 rows whose color is not that of the top right corner.

Conjectures from _Colin Barker_, Feb 11 2018: (Start) Assuming Colin Barker's conjectures, a(2*n) = 2*5^(n-1)*Lucas(2*(n+1)), a(2*n+1) = 2*5^n*Fibonacci(2*n+3). - _Ehren Metcalfe_, Apr 21 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph Myers_, Dec 24 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004905 | OEIS-A153635

Primes of the form 4*x^3 + 27*y^2, with x positive or negative.

Hardy and Wright: If there are an infinite number of these primes, then there are infinitely many cubic polynomials with integer coefficients and prime discriminant. It would also resolve the open conjecture that there are infinitely many non-isomorphic elliptic curves defined over the rationals and having prime conductor. Granville: "The most desired open problem in [asymptotic sieves] is to show that 4a^3 + 27b^2 is prime for infinitely many pairs of integers a, b (this is of interest because if 4a^3 + 27b^2 is prime then it is usually the conductor of the elliptic curve y^2 = x^3 + ax + b)." - _Charles R Greathouse IV_, Jun 06 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Dec 29 2008, Jan 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004906 | OEIS-A153641

Nonzero coefficients of the Swiss-Knife polynomials for the computation of Euler, tangent, and Bernoulli numbers (triangle read by rows).

Conjectural o.g.f.: Sum_{n >= 0} (1/2^((n-1)/2))*cos((n+1)*Pi/4)*(Sum_{k = 0..n} (-1)^k*binomial(n,k)/(1 - (k + x)*t)) = 1 + x*t + (x^2 - 1)*t^2 + (x^3 - 3*x)*t^3 + ... (checked up to O(t^13)), which leads to W_n(x) = Sum_{k = 0..n} 1/2^((k - 1)/2)*cos((k + 1)*Pi/4)*(Sum_{j = 0..k} (-1)^j*binomial(k, j)*(j + x)^n). - _Peter Bala_, Oct 03 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Dec 29 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004907 | OEIS-A153659

Triangle read by rows. A074206 interleaved with k-1 zeros in the k-th column.

By moving the term in the lower right corner to the upper right corner and then calculating the determinant one gets the sequence 1,-1,-1,-1,-1,-1,-1,-1,... The contributing permutation matrices have the same signs as the contributing permutation matrices in the Möbius function determinant. It is conjectured that their number is A074206.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Dec 30 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004908 | OEIS-A153772

a(n) = (2^n + 2*(-1)^n - 6)/3.

Conjecture: For n >= 6, a(n) is the third largest natural number whose Collatz orbit has length n+2. - _Markus Sigg_, Sep 14 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Jan 01 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004909 | OEIS-A153815

Indices of nontrivial zeros of the Riemann zeta function where the real part of zeta'(s) becomes negative.

Conjecture 1: Indices n of nontrivial zeros of the Riemann zeta function such that: abs(floor(im(zetazero(n))/(2*Pi)*log(im(zetazero(n))/(2*Pi*e)) + 7/8) - n + 1) = 1. Conjecture 2: The zeta zeros with these indices are also the locations where the zeta zero counting sequence A135297 disagrees with the zeta zero counting function: (RiemannSiegelTheta(t) + im(log(zeta(1/2 + I*t))))/Pi + 1. The locations where the counting function overcounts are given by A282793, and the locations where the counting function undercounts are given by A282794. Conjecture 3: Union of A282793 and A282794. Floor(im(zetazero(n))/(2*Pi)*log(im(zetazero(n))/(2*Pi*e)) + 7/8) - n + 1 is the branch of the argument of zeta at the n-th zero on the critical line (conjectured). - _Stephen Crowley_, Mar 09 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Reshetnikov_, Jan 02 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004910 | OEIS-A153818

a(n) = Sum_{k=1..n} floor(n^2/k^2).

Conjecture: a(n) = zeta(2)*n^2 + zeta(1/2)*n + O(n^0.5/log(n)) (see link). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Jan 02 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004911 | OEIS-A154138

Indices k such that 3 plus the k-th triangular number is a perfect square.

Conjectures: G.f.: x*(1 + 2*x + 3*x^2 - 2*x^3 - 2*x^4)/((1-x)*(x^2-2*x-1)*(x^2+2*x-1)). [Comment from _Zak Seidov_, Oct 21 2009: I believe both of these conjectures are correct.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004912 | OEIS-A154140

Indices k such that 6 plus the k-th triangular number is a perfect square.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004913 | OEIS-A154141

Indices k such that 8 plus the k-th triangular number is a perfect square.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004914 | OEIS-A154142

Indices k such that 9 plus the k-th triangular number is a perfect square.

Conjectures: (Start) These conjectures follow from the theory of Pell-like equations.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004915 | OEIS-A154143

Indices k such that 10 plus the k-th triangular number is a perfect square.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004916 | OEIS-A154144

Indices k such that 13 plus the k-th triangular number is a perfect square.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004917 | OEIS-A154145

Indices k such that 15 plus the k-th triangular number is a perfect square.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004918 | OEIS-A154146

Numbers k such that 16 plus the k-th triangular number is a perfect square.

Conjectures: (Start) Conjectures confirmed: see link. - _Robert Israel_, Jul 18 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004919 | OEIS-A154148

Numbers k such that 21 plus the k-th triangular number is a perfect square.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004920 | OEIS-A154150

Numbers k such that 24 plus the k-th triangular number is a perfect square.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004921 | OEIS-A154151

Indices k such that 25 plus the k-th triangular number is a perfect square.

Conjectures: (Start) The first conjecture is true for the first 1000 terms of the sequence. - _Harvey P. Dale_, Jun 15 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004922 | OEIS-A154153

Numbers k such that 28 plus the k-th triangular number is a perfect square.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004923 | OEIS-A154154

Numbers k such that 30 plus the k-th triangular number is a perfect square.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004924 | OEIS-A154255

Floor of harmonic energy of the conjectural minimal-energy configurations of n points on the unit sphere in R^4, under the harmonic potential function.

Floor of harmonic energy of the conjectural minimal-energy configurations of n points on the unit sphere in R^4, under the harmonic potential function. Floor of the right-hand column of the subtable for k=4 given by Ballinger et al., in the table of data related to their paper. This table shows the conjectural minimal-energy configurations of n points on the unit sphere in R^k (for k from 2 to 64), under the harmonic potential function: the potential energy between two points at distance r is 1/r^(k-2). In each entry in the table, the column for n links to coordinates for the configuration (given as a plain text file in which each line lists the coordinates for one point).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jan 05 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004925 | OEIS-A154258

Number of triples <p,s,t> such that p+F_s+(F_t)^2=n, where p is an odd prime, s and t are greater than one and the Fibonacci number F_s or F_t is odd.

Zhi-Wei Sun conjectured that a(n)>0 for all n=5,6,... (i.e., any integer n>4 can be written as the sum of an odd prime, a positive Fibonacci number and a square of a positive Fibonacci number, with one of the two Fibonacci numbers odd). He has verified this for n up to 3*10^7. _Zhi-Wei Sun_ has offered a monetary reward for settling this conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 05 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004926 | OEIS-A154263

Number of triples <p, s,t> such that p+F_s+(F_t)^3=n, where p is an odd prime, s and t are greater than one and F_s or F_t is odd.

Zhi-Wei Sun conjectured that a(n)>0 for every n=5,6,...; in other words, any integer n>4 can be written as the sum of an odd prime, a positive Fibonacci number and a cube of a positive Fibonacci number, with one of two Fibonacci numbers odd. He has verified this up to 3*10^7. _Zhi-Wei Sun_ has offered a monetary reward for settling this conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004927 | OEIS-A154285

Number of ordered triples <p,s,t> satisfying p+L_s+L_t=n, where p is an odd prime, s and t are nonnegative and the Lucas number L_s or L_t is odd.

Zhi-Wei Sun conjectured that a(n)>0 for all n=5,6,...; in other words, any integer n>4 can be represented as the sum of an odd prime, an odd Lucas number and a Lucas number. This has been verified up to 1.5*10^8. Sun thought that the constant lim inf_n a(n)/log(n) is greater than 2 and smaller than 3. For k=2,3 Sun also conjectured that any integer n>4 can be written in the form p+L_s+(L_t)^k, where p is an odd prime and L_s or L_t is odd. _Zhi-Wei Sun_ has offered a monetary reward for settling this conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004928 | OEIS-A154290

Number of ordered triples <p,s,t> satisfying p+F_s+L_t = n, where p is an odd prime, s >= 2 and F_s or L_t is odd.

Zhi-Wei Sun conjectured that a(n)>0 for every n=5,6,...; in other words, any integer n>4 can be written as the sum of an odd prime, a positive Fibonacci number and a Lucas number, with the Fibonacci number or the Lucas number odd. Moreover, Sun conjectured that lim inf_n a(n)/log(n) is greater than 3 and smaller than 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 06 2009, Jan 07 2008 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004929 | OEIS-A154293

Integers of the form t/6, where t is a triangular number (A000217).

Conjecture: A193828 divided by 2. - _Omar E. Pol_, Aug 19 2011 The above conjecture is correct. - _Charles R Greathouse IV_, Jan 02 2012 G.f.: x^2*(x^2-x+1)*(x^2+4*x+1)/((1+x^2)^2*(1-x)^3) (conjectured). (End) The conjectured g.f. is correct. - _Charles R Greathouse IV_, Jan 02 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Joseph Stephan Orlovsky_, Jan 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004930 | OEIS-A154364

Number of ways to express n as the sum of an odd prime, a positive Pell number and a companion Pell number.

This is inspired by the sequence A154290 and related conjectures of Sun. On Jan 08 2009, Zhi-Wei Sun and Qing-Hu Hou conjectured that a(n)>0 for n=6,7,...; in other words, any integer n>5 can be written as the sum of an odd prime, a positive Pell number and a companian Pell number. The Pell numbers are defined by P_0=0, P_1=1 and P_{n+1}=2P_n+P_{n-1} (n=1,2,3,...) and the companion Pell numbers are given by Q_0=Q_1=2 and Q_{n+1}=2Q_n+Q_{n-1} (n=1,2,3...). Note that for n>5 both P_n and Q_n are greater than 2^n. _D. S. McNeil_ disproved the conjecture by finding the 4 initial counterexamples: 169421772576, 189661491306, 257744272674, 534268276332. - _Zhi-Wei Sun_, Jan 17 2009 On Feb 01 2009, Zhi-Wei Sun observed that these 4 counterexamples are divisible by 42 and guessed that all counterexamples to the conjecture of Sun and Hou should be multiples of 42. - _Zhi-Wei Sun_, Feb 01 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004931 | OEIS-A154404

Number of ways to express n as the sum of an odd prime, a positive Fibonacci number and a Catalan number.

Motivated by _Zhi-Wei Sun_'s conjecture that each integer n>4 can be expressed as the sum of an odd prime, an odd Fibonacci number and a positive Fibonacci number (cf. A154257), during their visit to Nanjing Univ. Qing-Hu Hou (Nankai Univ.) and Jiang Zeng (Univ. of Lyon-I) conjectured on Jan 09 2009 that a(n)>0 for every n=5,6,.... and verified this up to 5*10^8. _D. S. McNeil_ has verified the conjecture up to 5*10^13 and Hou and Zeng have offered prizes for settling their conjecture (see Sun 2009).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Qing-Hu Hou (hou(AT)nankai.edu.cn), Jan 09 2009, Jan 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004932 | OEIS-A154417

Number of ways to express n as the sum of an odd prime, a positive Fibonacci number and half of a positive Fibonacci number.

On Jan 09 2009, _Zhi-Wei Sun_ conjectured that a(n)>0 for every n=5,6,...; in other words, any integer n>4 can be written in the form p+F_s+F_{3t}/2 with p an odd prime and s,t>0. Sun verified this up to 5*10^6 and Qing-Hu Hou continued the verification (on Sun's request) up to 3*10^8. Note that 932633 cannot be written as p+F_s+F_{3t}/2 with p a prime and (F_s or F_{3t}/2) odd. If we set u_0=0, u_1=1

and u_{n+1}=4u_n+u_{n-1} for n=1,2,3,..., then F_{3t}/2=u_t is at least 4^{t-1} for each t=1,2,3,.... In a recent paper K. J. Wu and Z. W. Sun constructed a residue class which contains no integers of the form p+F_{3t}/2 with p a prime and t nonnegative.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 09 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004933 | OEIS-A154421

Number of ways to express n as the sum of an odd prime, a positive Fibonacci number and an even Lucas number.

On Jan 09 2009, _Zhi-Wei Sun_ conjectured that a(n)>0 for all n=6,7,.... ; in other words, any integer n>5 can be written in the form p+F_s+L_{3t} with p an odd prime, s positive and t nonnegative. [Compare this with the conjecture related to the sequence A154290.] Sun verified the above conjecture up to 5*10^6 and Qing-Hu Hou continued the verification up to 2*10^8. If we set v_0=2, v_1=4 and v_{n+1}=4v_n+v_{n-1} for n=1,2,3,..., then L_{3t}=v_t is at least 4^t for every t=0,1,2,.... On Jan 17 2009, _D. S. McNeil_ found that 36930553345551 cannot be written as the sum of a prime, a Fibonacci number and an even Lucas number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 09 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004934 | OEIS-A154536

Positive integers that can be written as the sum of a positive Pell number and twice a positive Pell number.

On Jan 10 2009, _Zhi-Wei Sun_ conjectured that any integer greater than 5 can be expressed as the sum of an odd prime and a term in the above sequence; in other words, each n=6,7,... can be written in the form p+P_s+2*P_t with p an odd prime and s,t>0. This has been verified up to 5*10^13 by _D. S. McNeil_ (from London Univ.). Motivated by this conjecture, Qing-Hu Hou (from Nankai Univ.) observed and _Zhi-Wei Sun_ proved that each term a(n) in the above sequence can be uniquely written in the form P_s+2P_t with s,t>0. Sun noted that 2176 cannot be written as the sum of a prime and two Pell numbers; _D. S. McNeil_ found that 393185153350 cannot be written in the form p+P_s+3P_t and 872377759846 cannot be written in the form p+P_s+4P_t, where p is a prime and s and t are nonnegative. _Zhi-Wei Sun_ has offered a monetary reward for settling this conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 11 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004935 | OEIS-A154623

Expansion of 1+(x/(1-5*x))*c(x/(1-5*x)) where c(x) is the g.f. of A000108.

Conjecture: n*a(n) + 2*(8-7*n)*a(n-1) + 45*(n-2)*a(n-2) = 0. - _R. J. Mathar_, Dec 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004936 | OEIS-A154752

Least prime p (or 0) such that n = p + T, where T is a triangular number (A000217), or -1 if there is no such representation.

Zhi-Wei Sun conjectures that only n=216 has no such representation. It appears that n = 2, 7, 61 and 211 are the only numbers for which the triangular number 0 is required in the representation (see A065397). When n is a triangular number, then a(n)=0. Sequence A132399 gives the number of representations of n as p+T. As n becomes larger, the largest prime required to verify the conjecture increases slowly. For example, for n<=10^3, the largest prime required is 953; for n<=10^6 it is 373361; for n<=10^9 it is only 36455351. Using primes less than 10^9, all n<243277591560 have been verified.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jan 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004937 | OEIS-A154767

Primes with even largest digit.

Relative density 0 in the primes. It is natural to conjecture that there are infinitely many, but can this be proved? - _Charles R Greathouse IV_, Nov 12 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jan 15 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004938 | OEIS-A154800

List of pairs of primes that are equidistant from and nearest to n, or the pair (0,0) if there is no such pair.

Conjecture: There are only six 0's in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jan 15 2009, Jan 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004939 | OEIS-A154806

Numbers such that every run length in base 2 is 4.

Conjecture: a(n) = 1/17*2^(4*n+4) + 15/34*(-1)^(n+1) - 1/2. - _Vaclav Kotesovec_, Nov 30 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jan 25 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004940 | OEIS-A154808

Numbers such that every run length in base 2 is 5.

Conjecture: a(n) = (-33-31*(-1)^n+2^(6+5*n))/66. g.f.: 31*x / ((x-1)*(x+1)*(32*x-1)). - _Colin Barker_, Sep 16 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jan 25 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004941 | OEIS-A154940

Number of ways to express n as the sum of an odd prime, a Lucas number and a Catalan number.

On Jan 16 2009, _Zhi-Wei Sun_ conjectured that a(n)>0 for n=5,6,... and verified this up to 5*10^6. (Sun also thought that lim inf_n a(n)/log(n) is a positive constant.) _D. S. McNeil_ continued the verification up to 10^13 and found no counterexamples. The conjecture is similar to a conjecture of Qing-Hu Hou and Jiang Zeng related to the sequence A154404; both conjectures were motivated by Sun's recent conjecture on sums of primes and Fibonacci numbers (cf. A154257).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004942 | OEIS-A155005

Smallest number having exactly n divisors that are contained in its decimal representation.

Conjecture: a(5+5n)=1125*10^n for n>0. [_Robert G. Wilson v_, Jan 23 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jan 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004943 | OEIS-A155051

Expansion of c(x^2)*(1+x)/(1-x), c(x) the g.f. of A000108.

Conjecture: A000975(n) = A264784(a(n-1)) for n > 0. - _Reinhard Zumkeller_, Dec 04 2015 Conjecture: (n+2)*a(n) -2*a(n-1) +(-5*n+4)*a(n-2) +8*a(n-3) +4*(n-3)*a(n-4)=0. - _R. J. Mathar_, Feb 05 2015 Conjecture: -(n+2)*(n-3)*a(n) +(n^2-n-10)*a(n-1) +4*(n^2-4*n+5)*a(n-2) -4*(n-2)^2*a(n-3)=0. - _R. J. Mathar_, Feb 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 19 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004944 | OEIS-A155105

Positive numbers appearing in the third column of A155103.

a(1) = 1, a(n) = a(n-1)*(1+3*2^(n-2)) (conjectured). a(n) = Product_{k=1..n-1} 1+3*2^(k-1) = QPochhammer[-3, 2, n-1]. (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Jan 20 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004945 | OEIS-A155114

Number of ways to express n as the sum of an odd prime, a positive Fibonacci number and twice a positive Fibonacci number.

Motivated by his conjecture related to A154257, on Dec 26 2008, _Zhi-Wei Sun_ conjectured that a(n)>0 for n=6,7,... On Jan 15 2009, _D. S. McNeil_ verified this up to 10^12 and found no counterexamples. See the sequence A154536 for another conjecture of this sort. Sun also conjectured that any integer n>7 can be written as the sum of an odd prime, twice a positive Fibonacci number and the square of a positive Fibonacci number; this has been verified up to 2*10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 20 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004946 | OEIS-A155151

Triangle T(n, k) = 4*n*k + 2*n + 2*k + 2, read by rows.

Conjecture: Let p = prime number. If 2^p belongs to the sequence, then 2^p-1 is not a Mersenne prime. - _Vincenzo Librandi_, Dec 12 2012 Conjecture is true because if T(n, k) = 2^p with p prime, then 2^p-1 = 4*n*k + 2*n + 2*k + 1 = (2*n+1)*(2*k+1) hence 2^p-1 is not prime. - _Michel Marcus_, May 31 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Jan 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004947 | OEIS-A155190

Non-triangular numbers having a unique representation as p + t, where p is prime and t is a triangular number.

Conjectured to be finite. The prime terms are in A065397.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jan 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004948 | OEIS-A155200

G.f.: A(x) = exp(Sum_{n>=1} 2^(n^2) * x^n/n), a power series in x with integer coefficients.

Conjecture: highest exponent of 2 dividing a(n) = A000120(n) = number of 1's in binary expansion of n, so that a(n)/2^A000120(n) is odd for n >= 0. - _Paul D. Hanna_, Sep 01 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004949 | OEIS-A155214

Palindromes n with the property that for some prime p, n = p + prime(p).

It is conjectured that the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Jan 22 2009, Mar 02 2009, Mar 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004950 | OEIS-A155456

Write (1+1/x)*log(1+x) = Sum c(n)*x^n; then a(n) = (n+1)!*c(n).

G.f.: conjecture: T(0)*x^2/(1+2*x) - 1 - x, where T(k) = 1 - x^2*(k+1)*(k+2)/(x^2*(k+1)*(k+2) - (1+2*x*(k+1))*(1+2*x*(k+2))/T(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 19 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Jan 22 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004951 | OEIS-A155587

Expansion of (1 + x*c(x))/(1 - x), where c(x) is the g.f. of A000108.

To prove _R. J. Mathar_'s conjecture, note that the o.g.f. of the sequence implies (Sum_{n >= 0} a(n)*x^n)*(1 - x) = 1 + x*c(x); i.e., a(0) + Sum_{n >= 1} (a(n) - a(n-1))*x^n = 1 + Sum_{n >= 1} C(n-1)*x^n, where C(n) = A000108(n) (Catalan numbers). Thus, C(n-1) = a(n) - a(n-1) (for n >= 1), and hence C(n) = a(n+1) - a(n). Since 2*(2*n - 1)*C(n-1) = (n + 1)*C(n), we get (n + 1)*a(n+1) + (-5*n + 1)*a(n) + 2*(2*n - 1)*a(n-1) = 0. The last equation implies _R. J. Mathar_'s conjecture. (End) Conjecture: n*a(n) + (6-5*n)*a(n-1) + 2*(2*n-3)*a(n-2) = 0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004952 | OEIS-A155828

Number of integers k in {1,2,3,..,n} such that kn+1 is a square.

Conjecture: All terms a(n) are of the form 2^m - 1. This has been verified up to n = 1000. The conjecture that a(n) is of the form 2^m - 1 holds trivially. In fact, the conjecture can be restated as follows: For any positive integer n, the solutions of the congruence x^2 = 1 (mod n) with 1 <= x <= n is a power of two. By the Chinese remainder theorem, this reduces to the case when n is a prime power. It is well known that the solution of x^2 = 1 (mod p^a) is two when p is an odd prime (one may obtain this by induction on a). As for the number of solutions of the congruence x^2 = 1 (mod 2^a), it equals 1, 2, 4 according to whether a=1, 2, or a>2. - _Zhi-Wei Sun_, Feb 11 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jan 28 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004953 | OEIS-A155858

Diagonal sums of triangle A155856.

Conjecture: a(n) -(n-1)*a(n-1) -(n-2)*a(n-2) +(n-3)*a(n-3) +(n-10)*a(n-4) -5*a(n-5) +3*a(n-6) +3*a(n-7) = 0. - _R. J. Mathar_, Feb 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 29 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004954 | OEIS-A155860

Number of ways to write 2n-1 as p + 2^x + 3*2^y with p an odd prime and x,y positive integers.

On Jan 21 2009, _Zhi-Wei Sun_ conjectured that a(n)>0 for n=6,7,...; in other words, any odd integer m>10 can be written as the sum of an odd prime, a positive power of 2 and three times a positive power of 2. Sun verified this for odd integers m<10^7. On Sun's request, Qing-Hu Hou and _Charles R Greathouse IV_ continued the verification for odd integers below 2*10^8 and 10^10 respectively and found no counterexamples. As 3*2^y = 2^y + 2^{y+1}, Sun's conjecture implies that each odd integer m>8 can be written as the sum of an odd prime and three positive powers of two. Note that Paul Erdős asked whether there is a positive integer r such that every odd integer m>3 can be written as the sum of a prime and at most r powers of 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 29 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004955 | OEIS-A155867

A 'Morgan Voyce' transform of the large Schroeder numbers A006318.

Conjecture: (n+1)*a(n) + (4-11*n)*a(n-1) + (29*n-43)*a(n-2) +(73-29*n)*a(n-3) + (11*n-40)*a(n-4) + (5-n)*a(n-5) = 0. - _R. J. Mathar_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 29 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004956 | OEIS-A155904

Number of ways to write 2n-1 as p+2^x+5*2^y with p an odd prime and x,y positive integers.

On Jan 21 2009, _Zhi-Wei Sun_ conjectured that a(n)>0 for n=8,9,...; in other words, any odd integer m>=15 can be written as the sum of an odd prime, a positive power of 2 and five times a positive power of 2. Sun has verified this for odd integers m<10^8. As 5*2^y=2^y+2^{y+2}, the conjecture implies that each odd integer m>8 can be written as the sum of an odd prime and three positive powers of two. [It is known that there are infinitely many positive odd integers not of the form p+2^x+2^y (R. Crocker, 1971).] Sun also conjectured that there are infinitely many positive integers n with a(n)=a(n+1); here is the list of such positive integers n: 1, 2, 3, 4, 5, 6, 9, 10, 11, 19, 24, 36, 54, 60, 75, 90, 98, 101, 105, 135, 153, 173, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004957 | OEIS-A155919

Number of squared hypotenuses mod n in three dimensions.

Conjecture: a(2^k) = A171231(k-1) for k > 0; a(p^k) = p^k for odd prime p. - _Andrew Howroyd_, Nov 17 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Finch_, Jan 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004958 | OEIS-A155973

Smallest x such that prime(2n)x^(2n-1) + prime(2n-1)x^(2n-2) + prime(2n-2)x^(2n-3) +...+ prime(2)x^1 + 2x^0 evaluates to an odd prime.

Conjecture: The number of 1's in this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jan 31 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004959 | OEIS-A156164

Decimal expansion of 17 + 12*sqrt(2).

Conjecturally, the fractional part 0.97056 27484 ... of this constant equals ((1 + 2 * Sum_{n >= 1} (-1)^n*exp(-2*Pi*n^2))/(1 + 2 * Sum_{n >= 1} exp(-2*Pi*n^2)))^4. The series are rapidly converging. For example, summing both series from n = 1 to n = 2 approximates the fractional part of the constant as ((1 - 2*exp(-2*Pi) + 2*exp(-8*Pi))/(1 + 2*exp(-2*Pi) + 2*exp(-8*Pi)))^4 = 0.97056 27484 77140 58562 026(89) ..., correct to 23 decimal places. - _Peter Bala_, Jun 05 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Feb 09 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004960 | OEIS-A156170

G.f.: exp(Sum_{n>=1} [Sum_{k>=1} k^n*x^k]^n/n), a power series in x with integer coefficients.

Conjecture: log(a(n)) ~ n^2 * log(2)/4. - _Vaclav Kotesovec_, Sep 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 05 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004961 | OEIS-A156201

Numerator of Euler(n, 1/8).

G.f.: conjecture T(0)/(1+3*x), where T(k) = 1 - 16*x^2*(k+1)^2/(16*x^2*(k+1)^2 + (1+3*x)^2/T(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 12 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004962 | OEIS-A156216

G.f.: A(x) = exp(Sum_{n>=1} A000204(n)^n * x^n/n), a power series in x with integer coefficients.

More generally, if exp(Sum_{n>=1} C(n) * x^n/n) equals a power series in x with integer coefficients, then exp(Sum_{n>=1} C(n)^n * x^n/n) also equals a power series in x with integer coefficients (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004963 | OEIS-A156265

a(n)=3*n-A078649(n).

Conjectured to be o(n)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004964 | OEIS-A156267

a(n)=A054353(2*n)-A078649(n).

Conjectured to be o(n)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004965 | OEIS-A156268

a(n)=2*A054353(n)-A078649(n).

Conjectured to be o(n)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004966 | OEIS-A156276

Denominator of Euler(n, 1/10).

Conjecture: satisfies a linear recurrence having signature (0, 100). - _Harvey P. Dale_, Apr 17 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004967 | OEIS-A156330

Numerator of Euler(n, 1/12).

G.f.: conjecture T(0)/(1+5*x), where T(k) = 1 - 36*x^2*(k+1)^2/(36*x^2*(k+1)^2 + (1+5*x)^2/T(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 12 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004968 | OEIS-A156344

Number of steps to reach a square starting from n and iterating the map: x -> x*ceiling(sqrt(x))/floor(sqrt(x)) or zero if no square is reached.

We conjecture sequence is never zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004969 | OEIS-A156351

a(n) = Sum_{k=1..n} (-1)^K(k+1)*(K(k+1)-K(k)) where K(k) = A000002(k).

a(n) = gcd(a(a(n-1)),2) + a(n-2) (conjectured). - _Jon Maiga_, Dec 07 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004970 | OEIS-A156377

Numerator of Euler(n, 1/16).

G.f.: conjecture T(0)/(1+7*x), where T(k) = 1 - 64*x^2*(k+1)^2/(64*x^2*(k+1)^2 + (1+7*x)^2/T(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 12 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004971 | OEIS-A156552

Unary-encoded compressed factorization of natural numbers.

A037011(a(n)) = A010052(n) [conjectured, depends on the exact definition of A037011].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leonid Broukhis_, Feb 09 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004972 | OEIS-A156554

The number of integer sequences of length d = 2n+1 such that the sum of the terms is 0 and the sum of the absolute values of the terms is d-1.

Conjectural: for any prime p >= 5, a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for all positive integers n and k. More generally, if r and s are positive integers, we conjecture that the same supercongruences hold for the sequence defined by [x^(r*n)] P(s*n,(1 + x)/(1 - x)). (End) Even more generally, we conjecture that the same supercongruences hold for the sequence defined by [x^(r*n)] (1 + x)^(A*n) * (1 - x)^(B*n) * P(s*n,(1 + x)/(1 - x)), where A and B are integers. - _Peter Bala_, Mar 17 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _W. Edwin Clark_, Feb 09 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004973 | OEIS-A156628

Square array, read by antidiagonals, where row n+1 is generated from row n by first removing terms in row n at positions 0 and {(m+1)*(m+2)/2-2, m>0} and then taking partial sums, starting with all 1's in row 0.

Conjecture: A(n,k) = A396981(n,k+2) - Sum_{j=0..n-1} (n-j)! * A(j,k). - _Mikhail Kurkov_, Jun 12 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004974 | OEIS-A156642

Number of decompositions of 4n+2 into unordered sums of two primes of the form 4k+3.

Conjecture. For n >= 1, a(n) > 0. This conjecture does not follow from the validity of the Goldbach binary conjecture because numbers of the form 4n+2, generally speaking, also have decompositions into sums of two primes of the form 4k+1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Feb 12 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004975 | OEIS-A156655

Primes of the form 1000*k + 1.

Let d be any divisors of 1000, then d and -d are quadratic residues modulo these primes. All terms are of the form x^2 - d*y^2 for d = -40, -25, -10, -8, -5, -4, -2, -1, 2, 5, 8, 10. Conjecture: they are also of the form x^2 - d*y^2 for d = 20, 40, 50, 200. - _Jianing Song_, Aug 29 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | Avik Roy (avik_3.1416(AT)yahoo.co.in), Feb 12 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004976 | OEIS-A156695

Odd numbers that are not of the form p + 2^a + 2^b, a, b > 0, p prime.

All members above 5 found so far (up to 2.5 * 10^11) are divisible by 255 = 3 * 5 * 17, and many are divisible by 257. I conjecture that all members of this sequence greater than 5 are divisible by 255. This implies that all odd numbers (greater than 7) are the sum of a prime and at most three positive powers of two. All terms > 5 are numbers k > 3 such that k - 2^n is a de Polignac number (A006285) for every n > 0 with 2^n < k. Are there numbers K such that |K - 2^n| is a Riesel number (A101036) for every n > 0? If so, ||K - 2^n| - 2^m| is composite for every pair m,n > 0, by the dual Riesel conjecture. - _Thomas Ordowski_, Jan 06 2024 In keeping with the example's connection to A000215, the lowest ki for ki * Product_{i=0..11} (F(i)) to belong to A156695 are 1, 433007, 25471, 17047, 1291, 7, 101, 807, 83, 347, 9, 179. So for example, 433007*(3*5) is a term. This implies a variant of the first commented conjecture accordingly. - _Bill McEachen_, Apr 17 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Feb 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004977 | OEIS-A156703

String of digits encountered in decimal expansion of successive ratios k/(k+1), treating only non-repeating expansions, with decimal point and leading and trailing zeros removed.

Conjecture: subsets of the sequence (as it fills out) will correspond to the odd integers by length.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bill McEachen_, Feb 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004978 | OEIS-A156746

Numerator of Euler(n, 1/20).

G.f.: conjecture T(0)/(1+9*x), where T(k) = 1 - 100*x^2*(k+1)^2/(100*x^2*(k+1)^2 + (1+ 9*x)^2/T(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 12 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004979 | OEIS-A156769

a(n) = denominator(2^(2*n-2)/factorial(2*n-1)).

This sequence appears in the Eta and Zeta triangles A160464 and A160474. Its resemblance to the sequence of the denominators of the Taylor series for tan(x) led to the conjecture A156769(n) = A036279(n)*A089170(n-1). - _Johannes W. Meijer_, May 24 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Johannes W. Meijer_, Feb 15 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004980 | OEIS-A156834

A156348 * A000010.

Conjecture: for n>1, a(n) = n iff n is prime. Companion to A156833.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Feb 16 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004981 | OEIS-A156849

Numbers k such that k^2 == 2 (mod 23^2).

Conjecture: a(n) = a(n-1) + a(n-2) - a(n-3) = 529*n/2 - 529/4 - 95*(-1)^n/4. - _R. J. Mathar_, Oct 18 2010 The conjecture is correct. - _Charles R Greathouse IV_, Dec 27 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Feb 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004982 | OEIS-A156887

a(n) = Sum_{k=0..n} C(n,k) * C(4*n+k,k).

Conjecture: 576*n*(32901928701*n-65877527665)*(4*n-3)*(2*n-1)*(4*n-1)*a(n) +(-8795436181229177*n^5 +35251410418024655*n^4 -47934714902592853*n^3 +29414167990853161*n^2 -9060238526902314*n +1466702211905280)*a(n-1) +8*(10299715469615*n^5 -136961193094719*n^4 +872530072905392*n^3 -2699499511785411*n^2 +3902106377543903*n -2123717948975100)*a(n-2) -64*(2*n-5)*(4*n-9)*(n-2)*(27741827*n-2925269736)*(4*n-11)*a(n-3)=0. - _R. J. Mathar_, Feb 25 2015 8*(4*n)*(4*n-1)*(4*n-2)**(4*n-3)*(4633*n^3-19662*n^2+27593*n-12804)*a(n) = (137604733*n^7-859190528*n^6+2179882848*n^5-2890753162*n^4+2144669963*n^3-880916550*n^2+182941416*n-14515200)*a(n-1) -(4*n-4)*(4*n-5)*(4*n-6)*(4*n-7)*(4633*n^3-5763*n^2+2168*n-240)*a(n-2). A proof of Mathar's conjectured third-order recurrence above follows easily using this second-order recurrence. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004983 | OEIS-A156894

a(n) = Sum_{k=0..n} binomial(n,k)*binomial(2*n+k-1,k).

Conjecture: 64*n*(2*n-1)*a(n) -16*(89*n^2 -134*n +63)*a(n-1) +4*(661*n^2 -2619*n +2576)*a(n-2) -3*(119*n^2 -713*n +1092)*a(n-3) +6*(2*n-7)*(n-4)*a(n-4) = 0. - _R. J. Mathar_, Feb 05 2015 Conjecture: 16*n*(782*n +5365)*(2*n-1)*a(n) +8*(3128*n^3 -362053*n^2 +593930*n -290328)*a(n-1) -3*(726869*n^3 -5105981*n^2 +11667946*n -8715544)*a(n-2) +158*(2*n-5)*(n-3)*(391*n -764)*a(n-3) = 0. - _R. J. Mathar_, Feb 05 2015 Conjecture: 4*n*(2*n-1)*(17*n^2 -52*n +39)*a(n) -(1207*n^4 -4899*n^3 +6692*n^2 -3504*n +576)*a(n-1) +2*(n-2)*(2*n-3)*(17*n^2 -18*n +4)*a(n-2) = 0. - _R. J. Mathar_, Feb 05 2015 [the Maple command sumrecursion (binomial(n,k) * binomial(2*n+k-1,k), k, a(n)) verifies this recurrence. - _Peter Bala_, Oct 05 2015] More generally, we conjecture that a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004984 | OEIS-A156903

Abundant numbers (A005101) whose abundance is odd.

If n is in the sequence, so is 100*n (conjectured). - _Sergey Pavlov_, Mar 22 2017. Pavlov's observation trivially follows from the fact that to have odd abundance a number k must be either a square or twice a square. If such a number k is abundant then 100*k = (10^2) * k is abundant as well and has odd abundance. In general, we can say that if k is present, so are t^2*k and 2*t^2*k, for every t>0. - _Giovanni Resta_, Oct 16 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Feb 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004985 | OEIS-A156907

G.f.: A(x) = 1 + x*exp(Sum_{k>=1} [A(2^k*x) - 1]^k/k).

Conjectured to consist entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004986 | OEIS-A156908

G.f.: A(x) = 1 + x*exp(Sum_{k>=1} [A(3^k*x) - 1]^k/k).

Conjectured to consist entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004987 | OEIS-A157002

Transform of Catalan numbers whose Hankel transform gives the Somos-4 sequence.

Conjecture: (n+1)*a(n) +3*(-n+1)*a(n-1) +2*(-2*n+1)*a(n-2) +2*(2*n-7)*a(n-3) +2*(2*n-7)*a(n-4)=0. - _R. J. Mathar_, Nov 19 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 20 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004988 | OEIS-A157021

a(n)=sum{k=0..floor(n/2), C(n,2k)*A000108(floor(k/2))}. Inverse binomial transform is aeration of doubled Catalan numbers.

Conjecture: (n+4)*a(n) +(-7*n-18)*a(n-1) +2*(11*n+12)*a(n-2) +8*(-5*n+2)*a(n-3) +(41*n-76)*a(n-4) +(-19*n+62)*a(n-5) +4*(-n+6)*a(n-6) +6*(n-6)*a(n-7)=0. - _R. J. Mathar_, Feb 05 2015
Conjecture:+(n+4)*(n^2-11*n+32)*a(n) +(-5*n^3+41*n^2-26*n-360)*a(n-1) +2*(5*n^3-47*n^2+112*n+64)*a(n-2) +2*(-5*n^3+53*n^2-182*n+168)*a(n-3) +(n-4)*(n^2-3*n-8)*a(n-4) +3*(n-4)*(n^2-9*n+22)*a(n-5)=0. - _R. J. Mathar_, Feb 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004989 | OEIS-A157100

Transform of Catalan numbers whose Hankel transform satisfies the Somos-4 recurrence.

Conjecture: (n+1)*a(n) +(-5*n+1)*a(n-1) +2*(2*n-1)*a(n-2) +2*(2*n-7)*a(n-3) +2*(-2*n+7)*a(n-4) = 0. - _R. J. Mathar_, Feb 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 22 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004990 | OEIS-A157125

A transform of the Catalan numbers.

Conjecture: (n+3)*(n-2)*a(n) -4*a(n-1) +2*(n^2-n-4)*a(n-2) +2*(2*n^2-7*n+2)*a(n-3) +(n+1)*(n-4)*a(n-4) +2*(n-1)*(2*n-7)*a(n-5)=0. - _R. J. Mathar_, Nov 15 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004991 | OEIS-A157129

An analog of the Kolakoski sequence A000002, only now a(n) = (length of n-th run divided by 2) using 1 and 2 and starting with 1,1.

a(n) = gcd(A284796(ceiling(n/2)), 2) (conjectured). - _Jon Maiga_, Jun 11 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004992 | OEIS-A157182

Square array such that horizontal and vertical neighbors add to a prime, read by antidiagonals, filled the greedy way using positive integers, not more than once each.

Can we conjecture that this is a permutation of the positive integers?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004993 | OEIS-A157196

a(n) = (1/2)*(sum of elements of n-th run) using 1 and 2 starting with 1,1.

We conjecture that the density of 1's in the sequence approaches 2/3 as n -> infinity. This conjecture is proved in the paper of Shallit.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004994 | OEIS-A157200

Numbers k such that k*(2k)^k - 1 is a prime.

Conjecture: there are infinitely many such primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Feb 25 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004995 | OEIS-A157218

Number of ways to write the n-th positive odd integer in the form p+2^x+7*2^y with p a prime congruent to 1 mod 6 and x,y positive integers.

On Feb 24 2009, _Zhi-Wei Sun_ conjectured that a(n)>0 for all n=18,19,...; in other words, any odd integer greater than 34 can be written as the sum of a prime congruent to 1 mod 6, a positive power of 2 and seven times a positive power of 2. Sun verified the conjecture for odd integers below 5*10^7, and Qing-Hu Hou continued the verification for odd integers below 1.5*10^8 (on Sun's request). Compare the conjecture with R. Crocker's result that there are infinitely many positive odd integers not of the form p + 2^x + 2^y with p an odd prime and x,y positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 25 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004996 | OEIS-A157240

a(n) = A128018(n) + 1.

Generating floretion is Y = .5('i + 'j + 'k + i' + j' + k') + ee. ("tes"). Note: A current conjecture is that if X is a floretion for which 4*tes(X^n) is an integer for all n, then X+sigma(X) also has this property. "sigma" is the uniquely defined projection operator which "flips the arrows" of a floretion (i.e. sigma('i) = i', sigma('j) = j', etc.). Taking X = .5('i + 'j + 'k + ee), then tesseq(X) = [-1, -2, -1, 1, 2, 1, -1, -2, -1, 1, 2, 1, ...] is an integer sequence, thus by the conjecture 4*tes(Y^n) = 4*tes((X+sigma)^n) should also be an integer sequence for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Creighton Dement_, Feb 25 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004997 | OEIS-A157242

Number of ways to write the n-th positive odd integer in the form p+2^x+11*2^y with p a prime congruent to 5 mod 6 and x,y positive integers.

On Feb. 24, 2009, Zhi-Wei Sun conjectured that a(n)=0 if and only if n<15 or n=17, 20, 23, 86, 124; in other words, except for 33, 39, 45, 171 and 247, any odd integer greater than 28 can be written as the sum of a prime p=5 (mod 6), a positive power of 2 and eleven times a positive power of 2. Sun verified the conjecture for odd integers below 5*10^7. Knowing the conjecture from Sun, Qing-Hu Hou and D. S. McNeil have continued the verification for odd integers below 1.5*10^8 and 10^12 respectively, and they have found no counterexample. Compare the conjecture with Crocker's result that there are infinitely many positive odd integers not of the form p+2^x+2^y with p an odd prime and x,y positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 25 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004998 | OEIS-A157308

G.f. A(x) satisfies the condition that both A(x) and F(x) = A(x*F(x)) = g.f. of A155585 have zeros for every other coefficient after initial terms; g.f. of dual sequence A157309 satisfies the same condition.

Conjecture: a(m) == 1 (mod 2) iff m is a power of 2 or m=0. [_Paul D. Hanna_, Mar 17 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 11 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-004999 | OEIS-A157328

Expansion of 1/(1-2x*c(4x)) with c(x) g.f. of Catalan numbers (A000108).

Conjecture: n*a(n) +2*(12-7n)*a(n-1) +16*(3-2n)*a(n-2) = 0. - _R. J. Mathar_, Dec 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Feb 27 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005000 | OEIS-A157409

Minimum of { k > 0 : [2^n / 3^k] mod 6 = 3 } if such k exists, 0 otherwise.

It is conjectured that a(n) = 0 only for n in {0,1,2,3,4,7,9,10,14,15,17,26}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Mar 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005001 | OEIS-A157418

a(n) is the number of ways to insert single pairs of parenthesis to completely separate n identical objects in a straight line such that at least one of the objects at the two ends is not enclosed.

Conjecture: n*(7*n-29)*a(n) +(-21*n^2+136*n-192)*a(n-1) +2*(-28*n^2+165*n-210)*a(n-2) -4*(n-3)*(7*n-22)*a(n-3)=0. - _R. J. Mathar_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | William Sit (wyscc(AT)sci.ccny.cuny.edu), Jun 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005002 | OEIS-A157615

On an n X n board, a(n) is the maximal number of squares covered by a self-avoiding path made of alternated vertical and horizontal unitary steps.

David Wilson conjectures that the formula for a(n) is:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Resta_, Mar 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005003 | OEIS-A157616

On an n X n board, a(n) is the maximal number of squares covered by a self-avoiding path that starts from a corner and is made of alternated vertical and horizontal unitary steps.

David Wilson conjectures that the formula for a(n) is:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Resta_, Mar 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005004 | OEIS-A157655

Zeroless primes p such that the next prime after p can be obtained from p by adding the sum and product of the digits of p.

If we allow a zero digit in p, we generate A089824. One could conjecture that the digit 1 must always appear in the entries of this sequence. The idea for this sequence and the description was motivated by A089823.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Mar 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005005 | OEIS-A157656

Maximal possible number of states in a minimal deterministic automaton, equivalent to an n-state nondeterministic automaton over 1-symbol alphabet.

Conjecture: a(n)=A059100(n-1) holds only for all n<20 as well as n=22 and n=23. (Rustem Aidagulov)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Mar 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005006 | OEIS-A157675

G.f.: A(x) = exp(Sum_{n>=1} A(2^n*x)^n * x^n/n).

Conjectured to consist entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005007 | OEIS-A157713

a(n) = (2*n+1)!*(2*n-2)!/((n-1)!*(n!)^2*6).

Conjecture: n*a(n) -4*(2*n+1)*(2*n-3)*a(n-1)=0. - _R. J. Mathar_, Jun 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Karol A. Penson_, Mar 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005008 | OEIS-A157715

Primes sorted on digit sums, then on the primes.

Only correct for n >= 4 if an undiscovered prime of digit sum two (which would have to be a member of A080176) does not exist; this is conjectured but not proved. - _Jeppe Stig Nielsen_, Mar 30 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lekraj Beedassy_, Mar 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005009 | OEIS-A157751

Triangle of coefficients of polynomials F(n,x) in descending powers of x generated by F(n,x)=(x+1)*F(n-1,x)+F(n-1,-x), with initial F(0,x)=1.

Conjecture 1. If n>1 is even then F(n,x) has no real roots. Conjecture 2. If n>0 is odd then F(n,x) has exactly one real root, r, Conjectures 1 and 2 are true. [From Alain Thiery (Alain.Thiery(AT)math.u-bordeaux1.fr), May 14 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 05 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005010 | OEIS-A157792

Least number of lattice points on one side from which every point of a square n X n lattice is visible.

Conjectures from _Chai Wah Wu_, Aug 05 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005011 | OEIS-A157795

Largest subset of the discrete triangular grid { (a,b,c): a+b+c = n, a,b,c >= 0 } that does not contain any upward-pointing triangles (i.e., triples (a+r,b,c), (a,b+r,c), (a,b,c+r) with r positive).

The n=3 case was posed as a problem by Fujimura. The sequence is related to a certain "hyper-optimistic conjecture" regarding the density Hales-Jewett theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Terence Tao_, Mar 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005012 | OEIS-A157950

Primes p such that p^8 + 2^8 is prime.

It is conjectured that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Mar 10 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005013 | OEIS-A157996

Primes which are sum of 1 and two nonconsecutive primes p1 and p2, p2 - p1 > 2.

Conjecture: for n > 1, a(n) = prime(n+5). - _Charles R Greathouse IV_, Mar 12 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Joseph Stephan Orlovsky_, Mar 11 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005014 | OEIS-A158013

100*a(n)+13 and 100*a(n)+27 are consecutive primes, i.e., a prime gap 14.

2) It is conjectured that sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Mar 11 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005015 | OEIS-A158092

Number of solutions to +- 1 +- 2^2 +- 3^2 +- 4^2 +- ... +- n^2 = 0.

The expansion and integral representation formulas below are due to Andrica & Tomescu. The asymptotic formula is a conjecture; see Andrica & Ionascu. - _Jonathan Sondow_, Nov 11 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pietro Majer_, Mar 12 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005016 | OEIS-A158118

Number of solutions of +-1+-2^3+-3^3..+-n^3=0.

The expansion above and the integral representation formula below are due to Andrica & Tomescu. The asymptotic formula is a conjecture; see Andrica & Ionascu. - _Jonathan Sondow_, Nov 06 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pietro Majer_, Mar 12 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005017 | OEIS-A158119

Unsigned bisection of A157308 and A157310.

Conjecture: a(m) == 1 (mod 2) iff m is a power of 2 or m=0. [_Paul D. Hanna_, Mar 16 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 12 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005018 | OEIS-A158196

Expansion of (1-x^2*c(x)^4)/(1-3*x*c(x)^2), c(x) the g.f. of A000108.

Conjecture: +3*(n+1)*a(n) +2*(-26*n+7)*a(n-1) +16*(18*n-25)*a(n-2) +256*(-2*n+5)*a(n-3)=0. - _R. J. Mathar_, Feb 05 2015 Conjecture: 3*(2*n+3)*(n+1)*a(n) +2*(-28*n^2-52*n+21)*a(n-1) +32*(2*n+5)*(2*n-3)*a(n-2)=0. - _R. J. Mathar_, Feb 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005019 | OEIS-A158197

Expansion of (1-x^2*c(x)^4)/(1-4*x*c(x)^2), c(x) the g.f. of A000108.

Conjecture: +4*(n+1)*a(n) +(-81*n+23)*a(n-1) +10*(51*n-70)*a(n-2) +500*(-2*n+5)*a(n-3)=0. - _R. J. Mathar_, Feb 05 2015 Conjecture: +4*(n+1)^2*a(n) +(-41*n^2-58*n+23)*a(n-1) +50*(n+2)*(2*n-3)*a(n-2)=0. - _R. J. Mathar_, Feb 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005020 | OEIS-A158232

Numbers which yield primes when "13" is prefixed or appended: N natural number is a member of the sequence, if P="13N" (prefixed 13) and A="N13" (appended 13) are prime.

It is conjectured and numerically examined that sequences of this type are infinite. It is also conjectured that an infinite number of primes are terms of the sequence; first 20 primes are: 19, 61, 103, 127, 241, 331, 337, 367, 523, 577, 709, 829, 997, 1009, 1129, 1213, 1381, 1489, 1543, 1627.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Mar 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005021 | OEIS-A158233

a(n) = A120070(n+1)*A120070(n+2) mod 9.

Conjecture: this contains only the numbers 0,3,4,6 (verified for the first 5000 terms).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Mar 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005022 | OEIS-A158277

The lesser of twin prime pairs with each prime in a different century.

The sequence is conjecturally infinite; note that those ending in 999 straddle millenia: A158861.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ki Punches_, Mar 15 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005023 | OEIS-A158361

Primes p with property that Q = p^4 + 2^4 is prime.

It is conjectured that sequence a(n) is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Mar 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005024 | OEIS-A158423

Number of permutations of 1..n containing the relative rank sequence { 23145 } at any spacing.

Conjecture: a(n) + A116485(n) = n!. - _Benedict W. J. Irwin_, Mar 15 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005025 | OEIS-A158439

Numbers arising in Kane's conjecture on representing sets with sums of triangular numbers.

Numbers arising in Kane's conjecture on representing sets with sums of triangular numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Mar 19 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005026 | OEIS-A158460

Numbers k such that prime(k-1) + 5 is square and equal to prime(k+1) - 1.

Conjecture: If the condition holds, prime(k-1) and prime(k) are twin primes of the form 10m+1 and 10m+3. I.e., the last digits are 1 and 3. This is so because prime(k-1)+5 square => possible ending digits of 0,1,4,5,6,9. To get ending digits for prime(k-1) we subtract 5 to get ending digits 5,6,9,0,1,4. So 1,9 are the only possible endings since 0,4,5,6 => prime(k-1) not prime, impossible. Now by the condition of equality, prime(k-1)+6 = prime(k+1). So if prime(k-1) ends in 9, prime(k-1)+6 ends in 5 => prime(k+1) not prime, impossible. Therefore prime(k-1) ends in 1 and by the condition of prime(k-1) and prime(k) being twin primes, prime(k) ends in 3. This sequence is a variation of the conjecture provided in the link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Mar 19 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005027 | OEIS-A158465

Number of solutions to +-1+-2^4+-3^4+-4^4...+-n^4=0.

Andrica & Tomescu give a more general integral formula than the one below. The asymptotic formula below is a conjecture by Andrica & Ionascu; it remains unproven. - _Jonathan Sondow_, Nov 11 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pietro Majer_, Mar 19 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005028 | OEIS-A158470

Numbers k such that prime(k-1) + 7 is square and equal to prime(k+1) - 1.

Since prime(k-1) is prime, endings 2,4,5,8 are not allowed. This leaves us with 3,7,9 as possible endings of prime(k-1). Now to get prime(k), which the condition states is 2 greater than prime(k-1), we add 2 and 3+2=5 => prime(k) not prime, impossible. So the possible endings of prime(k-1) are reduced to 7 or 9. Now the condition prime(k-1)+7 = prime(k+1)-1 => prime(k-1)+8 = prime(k+1). Then adding 7 => prime(k+1) ends in 5, impossible. So prime(k-1) must end in 9, and adding 2 makes prime(k) end in 1. This

sequence is a calculation of the conjecture provided in the link. The PARI script provides for the general investigation of numbers of the form prime(k-1)+a and prime(k+1)-b. The values a=5,7; b=1 consistently yield twin primes when the condition holds.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Mar 19 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005029 | OEIS-A158477

Primes p with property that Q(p) = p^32+2^32 is prime.

2) It is conjectured that sequence a(n) is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Mar 20 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005030 | OEIS-A158495

Expansion of ((1-4*x) + sqrt(1-4*x))/(2*(1-2*x)).

Conjecture: n*a(n) +6*(-n+1)*a(n-1) +4*(2*n-3)*a(n-2)=0. - _R. J. Mathar_, Dec 03 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 20 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005031 | OEIS-A158509

Prime(n-1) such that prime(n-1)+5 is square and equal to prime(n+1)-1.

Conjecture: If the condition holds, prime(n-1) and prime(n) are twin primes of the form 10k+1 and 10k+3. I.e., the last digits of the primes are 1 and 3. The 1 ending is evident in the entries here. The conjecture is true: For p = prime(n-1) > 2 in this sequence we know that {p+1, p+3, p+5} are all even, p+6 = prime(n+1), p+5 = r^2 is a square, and p+4 = r^2-1 = (r-1)*(r+1) is composite. So p+2 = prime(n). Then p mod 10 = 1 because no prime > 5 has rightmost digit 5 and no square has rightmost digit 2. - _Alois P. Heinz_, Sep 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Mar 20 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005032 | OEIS-A158521

Primes which yield primes when "13" is prefixed or appended.

It is conjectured that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Mar 20 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005033 | OEIS-A158529

List of primes p with following properties: p = prime(m-1) for some m, p+7 is a square and is equal to prime(m+1)-1.

Conjecture: If the condition holds, prime(m-1) and prime(m) are twin primes of the form 10k+9 and 10k+11, i.e. the last digits of the twin prime pairs are 9 and 1. The 9 ending is obvious in this sequence. The table of the first 101 terms was computed using Zak Seidov's table. Proof of conjecture: Suppose p = prime(m-1) is a term with p + 7 = x^2. Now prime(m) must be p + 2 (it can't be p + 4 because one of p, p+4 and p+8 would be divisible by 3, and it can't be p + 6 = x^2-1 = (x-1)*(x+1)). As an even square, p + 7 ends in 0, 4 or 6, but it can't be 0 because p + 2 would be divisible by 5 and can't be 4 because p + 8 would be divisible by 5. The only possibility is that p + 7 ends in 6, i.e. p ends in 9. - _Robert Israel_, Jun 12 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Mar 20 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005034 | OEIS-A158584

Erroneous version of A079866.

We do not begin with the unit 1 because it has no prime factors. Conjecture: The sequence contains the set of prime numbers more than once.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Mar 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005035 | OEIS-A158594

Numbers which yield a prime whenever a 3 is prefixed, appended or inserted.

1) It is conjectured that sequences of this type are infinite; also that an infinite number of primes is included.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Mar 22 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005036 | OEIS-A158623

Denominator of the reduced fraction A158620(n)/A158621(n).

A158620(n)/A158621(n) = 2(n^2+n+1)/(3n(n+1)). Conjecture: a(n) = 3a(n-3) - 3a(n-6) + a(n-9), so trisections are A152996, A060544 and 3*A081266. - _R. J. Mathar_, Mar 27 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Mar 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005037 | OEIS-A158690

Expansion of the basic hypergeometric series 1 + (1 - exp(-t)) + (1 - exp(-t))*(1 - exp(-3*t)) + (1 - exp(-t))*(1 - exp(-3*t))*(1 - exp(-5*t)) + ... as a series in t.

Conjectures: Peter Bala's conjectures are correct (see Kallat, 2026). - _Ahaan Kallat_, Sep 08 2026 Conjectural g.f.: G(exp(t)) as a formal power series in t, where G(q) := Sum_{n >= 0} q^(2*n+1) * Product_{k = 1..2*n} (1 - q^k). - _Peter Bala_, May 16 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005038 | OEIS-A158691

The number of upper-triangular matrices with at least one nonzero entry in each row and whose entries sum to n.

The above remark, first conjectured by Bala, is a consequence of the identities satisfied by the generating function for a(n). More precisely, the generating function F(x)=Sum_{n>=0} a(n)x^n can be expressed by any of these three formulas: The first two formulas were

conjectured to be equal by Bala. This was confirmed by Andrews and Jelínek, who also derived the third formula. Conjectural g.f.: Sum_{n >= 0} 1/(1 - x)^((n+1)*(2*n+1)) * Product_{i = 1..2*n} ((1 - x)^i - 1). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005039 | OEIS-A158704

Nonnegative integers with an even number of even powers of 2 in their base-2 representation.

Conjecture: take any binary vector of length 4n+3 with n >= 0. We can activate any bits. When a bit is activated, neighboring bits change their values 0 -> 1, 1 -> 0. Our goal is to turn the original binary vector into a vector of only ones by activating the bits. If the value of the binary vector belongs to this sequence, this is possible for a maximum of 4n+3 activations. - _Mikhail Kurkov_, Jun 01 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Mar 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005040 | OEIS-A158721

Primes p such that (p + 1)/3 + p is prime.

Set q = (p + 1)/3 + p, then (p + 1)/(q + 1) = 3/4. If this sequence is proven infinite, that would prove two specific cases of the Schinzel-Sierpiński conjecture regarding rational numbers. - _Alonso del Arte_, Mar 12 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Joseph Stephan Orlovsky_, Mar 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005041 | OEIS-A158793

Triangle read by rows: product of A130595 and A092392 considered as infinite lower triangular arrays.

Conjecture: T(n,1) = A113682(n-1). - _R. J. Mathar_, Oct 06 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_ & _Roger L. Bagula_, Mar 26 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005042 | OEIS-A158794

Multiples of 4 which are not the sum of seven nonnegative cubes.

Boklan and Elkies: It is conjectured that every integer N>454 is the sum of seven nonnegative cubes. We prove the conjecture when N is a multiple of 4. Elkies [2010]: It is conjectured that every integer N>454 is the sum of seven nonnegative cubes. We prove the conjecture when N is congruent to 2 mod 4. This result, together with a recent proof for 4|N, shows that the conjecture is true for all even N. - _Jonathan Vos Post_, Sep 22 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Mar 26 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005043 | OEIS-A158826

Third iteration of x*C(x) where C(x) is the Catalan function (A000108).

Conjecture D-finite with recurrence 1053*n*(n-1)*(n-2)*(n-3)*a(n) -36*(n-1)*(n-2)*(n-3)*(634*n-1367)*a(n-1) +24*(n-2)*(n-3)*(7966*n^2-43500*n+61181)*a(n-2) -8*(n-3)*(96128*n^3-957424*n^2+3221878*n-3665189)*a(n-3) +16*(91904*n^4-1446528*n^3+8575792*n^2-22703688*n+22652013)*a(n-4) -256*(8*n-35)*(8*n-41)*(8*n-39)*(8*n-37)*a(n-5)=0. - _R. J. Mathar_, Aug 30 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 28 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005044 | OEIS-A158861

The lesser of twin prime pairs with each prime in a different millennia.

The sequence is conjecturally infinite; note that terms ending 9999 straddle ten millennia.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ki Punches_, Mar 28 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005045 | OEIS-A158960

Numerator of Hermite(n, 1/5).

The denominators are 5^n = A000351(n) (conjectured). - _M. F. Hasler_, Feb 16 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 12 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005046 | OEIS-A158979

a(n) is the smallest number > n such that n^4 + a(n)^4 is prime.

Conjecture: a(n) exists for all n, i.e., the sequence is well-defined and infinite. Conjecture: a(n)-n = 1 for infinitely many n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Apr 01 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005047 | OEIS-A159021

a(0)=19; a(n) = a(n-1) + floor(sqrt(a(n-1))), n > 0.

Conjecture: The squares are of the form (7*2^k)^2 (see A005009). - _Vincenzo Librandi_, Apr 10 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Delélham_, Apr 02 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005048 | OEIS-A159082

Numbers whose squares added to 7! are prime.

2) Sequence is conjectured to be infinite. 3) It is conjectured that an infinite number of terms are primes. 5) It is conjectured that an infinite number of a(n) are squares, first are 121=11^2, 169=13^2, 529=23^2, 841=29^2, 961=31^2, 1681=41^2, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Apr 05 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005049 | OEIS-A159225

1/4 the number of n X n arrays of squares of integers with every 2X2 subblock summing to 25.

Conjecture: G.f.: x^2*(-7-12*x^4+20*x^3+13*x^2-17*x)/((x-1)*(2*x+1)*(2*x-1)*(2*x^2-1)) [From Maksym Voznyy (voznyy(AT)mail.ru), Jul 26 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005050 | OEIS-A159227

1/4 the number of n X n arrays of squares of integers with every 2X2 subblock summing to 27.

Conjecture: G.f.: x^2*(-7-8*x^4+32*x^3+19*x^2-15*x)/((x-1)*(2*x+1)*(2*x-1)*(2*x^2-1)) [From Maksym Voznyy (voznyy(AT)mail.ru), Jul 26 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005051 | OEIS-A159351

Smallest prime of the form a^2 + b^2 with 0 < a < b such that a + b = 2n+1.

Conjecture: there is always such a prime number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Apr 11 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005052 | OEIS-A159578

Dividend associated with A116536.

a(2)-a(4) are mentioned by Alladi and Erdős (1977). They conjectured that this sequence is infinite. - _Amiram Eldar_, Nov 02 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enoch Haga_, Apr 16 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005053 | OEIS-A159596

G.f.: A(x) = exp(Sum_{n>=1} [D^n x/(1-x)^2]^n/n), where differential operator D = x*d/dx.

Conjecture: limit_{n->oo} a(n)^(1/n^2) = 2^(1/4). - _Vaclav Kotesovec_, Nov 17 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 05 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005054 | OEIS-A159619

Slowest increasing sequence beginning with 4 such that n and a(n) are either both evil or both odious.

There is a conjecture arising in Sequence Machine that a(n) = A026491(2+n)-1. This appears to be true: Here we start from on odious or evil number and apply a minimum number of van-Eck-Transforms (of A171898) to reach a value larger than a(n-1). The Dekking formula in A026491 says that A026491 is essentially a partial sum of the backward van-Eck-Transforms, and in a (vague) manner this seems to match.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 17 2009, Apr 27 2009, May 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005055 | OEIS-A159633

Dimension of Eisenstein subspace of the space of modular forms of weight k/2, level 4*n and trivial character, where k>=5 is odd.

Conjecture: a(n) = 2*chi(n) - if(mod(n+2,4)=0, chi(n)/2, 0) with chi(n) = A001616(n) = Sum_{d|n} phi(gcd(d,n/d)); checked up to n=1024. - _Wouter Meeussen_, Apr 02 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Finch_, Apr 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005056 | OEIS-A159671

Continued fraction expansion of Landau's constant L.

Terms are computed using the conjectured formula. It is conjectured that L = 0.5432589653429767... = Gamma(1/3)*Gamma(5/6)/Gamma(1/6).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Harry J. Smith_, Apr 19 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005057 | OEIS-A159704

Number of different pairs of primes p,q such that : p<q, p is a cousin prime of p-4 or p+4 and q is a cousin prime of q-4 or q+4, 2*n=p+q.

conjecture : for n>5573 there is at least one such pair of primes p+q=2*n

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Apr 20 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005058 | OEIS-A159715

Number of permutations of 2 indistinguishable copies of 1..n arranged in a circle with exactly 1 local maximum.

Conjectures from _Colin Barker_, Mar 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 20 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005059 | OEIS-A159769

Number of n-leaf binary trees that do not contain ((((())())(()(()(())))) as a subtree.

Conjecture: 2*(n+1)*a(n) +3*(-3*n+1)*a(n-1) +2*(2*n-1)*a(n-2) +4*(2*n-7)*a(n-3) +2*(-2*n+7)*a(n-4)=0. - _R. J. Mathar_, May 30 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Rowland_, Apr 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005060 | OEIS-A159802

Number of primes q with (2m)^2+1 <= q < (2m+1)^2-2m.

2) See sequence A145354 and A157884 for more details to this new improved conjecture. 4) It is conjectured that a(m) >= 1. This is a bisection of A089610 and hence related to a conjecture of Oppermann. - _T. D. Noe_, Apr 22 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Apr 22 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005061 | OEIS-A159803

Number of primes p with (2m+1)^2 - 2m <= p < (2m+1)^2.

2) See sequence A145354 and A157884 for more details to this new improved conjecture. 4) It is conjectured that a(m) >= 1. This is a bisection of A094189 and hence related to a conjecture of Oppermann. - _T. D. Noe_, Apr 22 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Apr 22 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005062 | OEIS-A159804

Number of primes q with (2n-1)^2+1 <= q < (2n)^2-(2n-1).

See sequence A145354 and A157884 for more details to this new improved conjecture. It is conjectured that a(n) >= 1 for n >= 1. This is a bisection of A089610 and hence related to a conjecture of Oppermann. [_T. D. Noe_, Apr 22 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Apr 22 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005063 | OEIS-A159805

Number of primes p with (2m)^2-(2m-1) <= p < (2m)^2.

2) See sequence A145354 and A157884 for more details to this new improved conjecture. 4) It is conjectured that a(m) >= 1. Except for a(1), this is a bisection of A094189 and hence related to a conjecture of Oppermann. - _T. D. Noe_, Apr 22 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Apr 22 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005064 | OEIS-A159807

Numerator of Hermite(n, 3/22).

Conjecture: E.g.f.: exp(-x*(121*x-3)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 12 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005065 | OEIS-A159856

Triangle read by rows: T(n,0) = n+1, T(n,k) = 2*T(n-1,k) - T(n-1,k-1), T(n,k) = 0 if k > n and if k < 0.

Conjecture: Sum_{i=0..n} |T(n,k)| = A047926(n). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Apr 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005066 | OEIS-A159864

Difference array of Fibonacci numbers A000045 read by antidiagonals.

Conjecture: row sums are Sum_{k=0..n} T(2n,k)=0. Sum_{k=0..n} T(2n+1,k) = A025169(n). - _R. J. Mathar_, May 29 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Apr 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005067 | OEIS-A159881

Triangle read by rows : T(n,0) = n+1, T(n,k)=0 if k<0 or if k>n, T(n,k) = k*T(n-1,k) - T(n-1,k-1).

Conjecture row sums: Sum_{k=0..n} |T(n,k)| = A029761(n). - _R. J. Mathar_, May 29 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Apr 25 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005068 | OEIS-A159907

Numbers m with half-integral abundancy index, sigma(m)/m = k+1/2 with integer k.

Conjecture: with number 1, multiply-anti-perfect numbers m: m divides antisigma(m) = A024816(m). Sequence of fractions antisigma(m) / m: {0, 0, 10, 2157, 2337, 13101, 4455356, ...}. - _Jaroslav Krizek_, Jul 21 2011 The above conjecture is equivalent to the conjecture that there are no odd multiply perfect numbers (A007691) greater than 1. Proof: (sigma(n)+antisigma(n))/n = (n+1)/2 for all n. If n is even then sigma(n)/n is a half-integer if and only if antisigma(n)/n is an integer. Since all members of this sequence are known to be even, the only way the conjecture can fail is if antisigma(n)/n is an integer, in which case sigma(n)/n is an integer as well. - _Nathaniel Johnston_, Jul 23 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 25 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005069 | OEIS-A159939

Odd solutions of phi(sigma(k)) = sigma(phi(k)).

It is conjectured (see links) that all odd solutions are of this form, for which at least 10130 solutions are known.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Walter Nissen_, Apr 26 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005070 | OEIS-A160002

Number of moves needed to solve 4-peg Tower of Hanoi Puzzle with n disks.

Values for n=21, 22, 23 were computed on SuperMUC-NG by Andreas M. Hinz and Ciril Petr during 2020-2021. The value for n = 23 is the first one that is less than Yang's bound, so the optimal strategy is still an open problem. - _Ciril Petr_, May 03 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yi Yang_, Apr 29 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005071 | OEIS-A160022

Primes p such that p^4 + 5^4 + 3^4 is prime.

It is conjectured that the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Apr 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005072 | OEIS-A160023

Primes p such that p^4 + 7^4 + 3^4 is prime.

It is conjectured that the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Apr 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005073 | OEIS-A160024

Primes p such that p^4 + 11^4 + 3^4 is prime.

It is conjectured that the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Apr 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005074 | OEIS-A160025

Primes p such that p^4 + 13^4 + 3^4 is prime.

It is conjectured that the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Apr 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005075 | OEIS-A160026

Primes p such that p^4 + 17^4 + 3^4 is prime.

It is conjectured that the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Apr 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005076 | OEIS-A160031

Primes p such that p^4 + 2*3^4 is prime.

It is conjectured that the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Apr 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005077 | OEIS-A160077

Numerator of Hermite(n, 19/26).

E.g.f.: exp(-x*(169*x-19)). The conjecture is a consequence. - _Robert Israel_, Jan 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 12 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005078 | OEIS-A160089

The maximum of the absolute value of the coefficients of Pn = (1-x)(1-x^2)(1-x^3)...(1-x^n).

If n is even then a(n) is the absolute value of the coefficient of z^(n(n+1)/4). If n is odd, it is an open question as to which coefficient is a(n). For odd n values, the Berkovich/Uncu reference provides explicit conjectural formulas for a(n). - _Ali Uncu_, Jul 19 2020 Conjecture: a(n)^(1/n) ~ sqrt(A133871(n)^(1/n)) ~ 1.219715476121633689013599 33...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Theodore Kolokolnikov_, May 01 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005079 | OEIS-A160217

Minimal increasing sequence with a(1)=3 and the property that a(n) and n are both in or both not in A003159.

Conjecture: Let m>3 belong to A003159. Define the sequence b(n) to be the minimal increasing sequence with b(1)=m and the property that b(n) and n are both in or both not in A003159. Then a(n)=b(n) for all n larger than some m-dependent minimum index.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005080 | OEIS-A160218

Index at which n-th prime occurs in A160256, or -1 if the prime never occurs.

A160256(a(n))=A000040(n) if and only if both Conjectures 1 and 2 are true: Conjecture 1: Primes occur in A160256 in increasing order. Conjecture 2: All primes occur in A160256. Conjecture 3: Except for A160256(4)=4, the least positive integer which does not occur in A160256 up to a given index is always a prime (and thus of the form A160256(a(k)) for some k). Conjecture 4: A160256(a(n)) is always the least positive integer which did not occur earlier in A160256.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, May 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005081 | OEIS-A160239

Number of "ON" cells in a 2-dimensional cellular automaton ("Fredkin's Replicator") evolving according to the rule that a cell is ON in a given generation if and only if there was an odd number of ON cells among the eight nearest neighbors in the preceding generation, starting with one ON cell.

a(0) = 1; a(2t)=a(t), a(4t+1)=8*a(t), a(4t+3)=2*a(2t+1)+8*a(t) for t >= 0. (Conjectured by _Hrothgar_, Jul 11 2014; proved by _N. J. A. Sloane_, Oct 04 2014.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, May 05 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005082 | OEIS-A160268

Odd numbers k for which A006694((k-1)/2) < A006694((A000265(3k+1)-1)/2).

Conjecture: For every k in the sequence, the number k^2 is in the sequence as well.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005083 | OEIS-A160326

Number of ways to express n=0,1,2,... as the sum of two squares and a pentagonal number.

In April 2009, Zhi-Wei Sun conjectured that a(n)>0 for every n=0,1,2,3,... Note that pentagonal numbers are more sparse than squares. The Gauss-Legendre theorem asserts that n is the sum of three squares if and only if it is not of the form 4^a(8b+7) (a,b=0,1,2,...).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005084 | OEIS-A160389

Decimal expansion of 2*cos(Pi/7).

Conjecture: Equals lim_{n->oo} A060557(n) / A052975(n). - _Bill McEachen_, Feb 17 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Harry J. Smith_, May 31 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005085 | OEIS-A160420

Number of "ON" cells at n-th stage in simple 2-dimensional cellular automaton whose skeleton is the same network as the toothpick structure of A139250 but with toothpicks of length 4.

Conjecture: a(n) = A147614(n)+2*A139250(n). [From _R. J. Mathar_, Jan 22 2010] The above conjecture is true: each toothpick covers exactly two more grid points than the corresponding toothpick in A147614.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, May 13 2009, May 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005086 | OEIS-A160433

a(n) is the least number k such that (k-th prime after n!+1)-n! is not a prime.

The conjectures from A037153 and A087202 can be rephrased using a(n): Also compare this with the conjecture on the fortunate numbers A005235.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Frederick Magata (frederick.magata(AT)web.de), May 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005087 | OEIS-A160434

a(n) is the least number k such that (k-th prime after A002110(n)+1) - A002110(n) is not a prime, where A002110(n) is the n-th primorial.

The conjecture on the fortunate numbers rephrased with a(n) is a(n)>=2 for all n>=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Frederick Magata (frederick.magata(AT)web.de), May 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005088 | OEIS-A160455

Number of triangles that can be built from rods with lengths 1,2,...,n by using and concatenating all rods.

Conjectures from _Colin Barker_, May 12 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hagen von Eitzen_, May 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005089 | OEIS-A160464

The Eta triangle.

Some of our results are conjectures based on numerical evidence, see especially A160466.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Johannes W. Meijer_, May 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005090 | OEIS-A160466

Row sums of the Eta triangle A160464.

It is conjectured that the row sums of the Eta triangle depend on five different sequences. Two Maple algorithms are given. The first one gives the row sums according to the Eta triangle A160464 and the second one gives the row sums according to our conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Johannes W. Meijer_, May 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005091 | OEIS-A160468

Triangle of polynomial coefficients related to the o.g.f.s of the RES1 polynomials.

Some of our results are conjectures based on numerical evidence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Johannes W. Meijer_, May 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005092 | OEIS-A160469

The left hand column of the triangle A160468.

Resembles A002430, the numerators of the Taylor series for tan(x). The first difference occurs at a(12). (Its resemblance to this sequence led to the conjecture A160469(n) = A002430(n)*A089170(n-1).)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Johannes W. Meijer_, May 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005093 | OEIS-A160474

The Zeta triangle.

Some of our results are conjectures based on numerical evidence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Johannes W. Meijer_, May 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005094 | OEIS-A160479

The ZL(n) sequence of the Zeta and Lambda triangles A160474 and A160487.

The first ZL(n) formula is a conjecture. This formula links the ZL(n) to the prime numbers A000040; see A217983, A128060, A130290 and the third Maple program.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Johannes W. Meijer_, May 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005095 | OEIS-A160480

The Beta triangle read by rows.

Some of our results are conjectures based on numerical evidence, see especially A160481.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Johannes W. Meijer_, May 24 2009, Sep 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005096 | OEIS-A160481

Row sums of the Beta triangle A160480.

It is conjectured that the row sums of the Beta triangle depend on three different sequences. Two Maple algorithms are given. The first one gives the row sums according to the Beta triangle A160480 and the second one gives the row sums according to our conjecture. Conjecture: a(n) = (2*n-3)! - 2^(2*n-3)*(n-1)!*(n-2)!, for n >= 2 (gives the first 13 terms). - _Christopher P. Herzog_, Nov 25 2014 Meijer's and Herzog's conjectures can also be written as: a(n) = -A129890(n-2)*A000165(n-2) = A009445(n-2) - A002474(n-2). - _Peter Luschny_, Dec 01 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Johannes W. Meijer_, May 24 2009, Sep 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005097 | OEIS-A160487

The Lambda triangle.

Some of our results are conjectures based on numerical evidence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Johannes W. Meijer_, May 24 2009, Sep 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005098 | OEIS-A160595

Numerator of resilience R(n) = phi(n)/(n-1), with a(1) = 1 by convention.

Conjecture: a(n) > 2 for every composite n > 6. Slightly stronger than Lehmer's totient conjecture (1932). - _Thomas Ordowski_, Mar 13 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, May 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005099 | OEIS-A160627

Numerator of Laguerre(n, 4).

Conjecture: all terms are odd. - _Robert Israel_, Mar 29 2018 Conjecture proven by Wenlin Zhang and Haobo Ma with AI assistance (see links). - _Robert Israel_, Sep 30 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005100 | OEIS-A160648

Second differences of sequence A160644.

Conjecture: a(n) counts unrestricted partitions of even numbers such that

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alford Arnold_, May 25 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005101 | OEIS-A160678

Numbers n whose abundancy is equal to 13/2; sigma(n)/n = 13/2.

This sequence includes many terms but it is conjectured to be finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerard P. Michon_, Jun 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005102 | OEIS-A160686

Numbers n such that n/A000005(A000045(n)) is an integer.

Next term, if it exists, is greater than 3000. Conjecture: the sequence is finite and complete. - _Max Alekseyev_, May 21 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, May 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005103 | OEIS-A160693

Numerator of the 2n-th raw moment for distribution of distances between two points picked at random in the interior of a unit cube.

Conjecture: a(n)/A160694(n) = n!*Sum_{i=0..n,j=0..n-i} 1/((i+1)!(j+1)!(n-i-j+1)!(2i+1)(2j+1)(2(n-i-j)+1)). - _Jérôme Houdayer_, Nov 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, May 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005104 | OEIS-A160702

Sequence such that the Hankel transform of a(n+1) satisfies a generalized Somos-4 recurrence.

In general, we can conjecture that the Hankel transform of h(n) of a(n+1), where a(n)=if(n=0,1,if(n=1,1,if(n=2,1,r*a(n-1)+s*sum{k=0..n-2, a(k)*a(n-1-k)}))), satisfies the generalized Somos-4 recurrence h(n)=(s^2*h(n-1)*h(n-3)+s^3*(2*s+r-2)*h(n-2)^2)/h(n-4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005105 | OEIS-A160703

Generalized Somos-4 Hankel determinant recurrence sequence.

In general, we can conjecture that the Hankel transform a(n) of f(n+1), where f(n)=if(n=0,1,if(n=1,1,if(n=2,1,r*f(n-1)+s*sum{k=0..n-2, f(k)*f(n-1-k)}))) satisfies the generalized Somos-4 recurrence: a(n)=(s^2*a(n-1)*a(n-3)+s^3*(2*s+r-2)*a(n-2)^2)/a(n-4). The case r=s=1 is proved in the Xin reference.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005106 | OEIS-A160720

Number of "ON" cells at n-th stage in 2-dimensional cellular automaton (see Comments for precise definition).

Conjecture: a(n) = 1 + 4*(A266532(n) - 1)/3, n >= 1. - _Omar E. Pol_, Jan 19 2016. This formula is correct! - _N. J. A. Sloane_, Jan 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, May 25 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005107 | OEIS-A160766

Values of n for which A094004(n) > A094004(n-1)+1.

The next three terms are conjectured to be 68, 76 and 77.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benjamin Chaffin_ and _N. J. A. Sloane_, Feb 13 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005108 | OEIS-A160823

A transform of the large Schroeder numbers.

Conjecture: (n+2)*a(n) -3*(n+1)*a(n-1) +3(2-n)*a(n-2) +(11*n-20)*a(n-3) +(11-5*n)*a(n-4) + (4-n)*a(n-5)=0. - _R. J. Mathar_, Nov 16 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 27 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005109 | OEIS-A160845

Number of lines through at least 2 points of a 5 X n grid of points.

Conjectures from _Colin Barker_, May 24 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seppo Mustonen_, May 28 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005110 | OEIS-A160848

Number of lines through at least 2 points of an 8 X n grid of points.

Conjectures from _Colin Barker_, Dec 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seppo Mustonen_, May 28 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005111 | OEIS-A160849

Number of lines through at least 2 points of a 9 X n grid of points.

Conjectures from _Colin Barker_, Dec 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seppo Mustonen_, May 28 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005112 | OEIS-A160850

Number of lines through at least 2 points of a 10 X n grid of points.

Conjectures from _Colin Barker_, Dec 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seppo Mustonen_, May 28 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005113 | OEIS-A160906

Row sums of A159841.

Conjecture: a(2n+1) = A075273(3n). Conjecture: 2*n*(2*n-1)*(5*n-4)*a(n) +(-295*n^3+451*n^2-130*n-24)*a(n-1) +24*(5*n+1)*(3*n-4)*(3*n-2)*a(n-2) = 0. - _R. J. Mathar_, Jul 20 2016 R. J. Mathar's conjecture verified using differential equation (2112*x + 2)*y + (11328*x^2 - 842*x + 2)*y' + (7776*x^3 - 1319*x^2 + 34*x)*y'' + (1080*x^4 - 295*x^3 + 20*x^2)*y''' - 12 satisfied by the g.f. y(x). - _Robert Israel_, Mar 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, May 29 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005114 | OEIS-A160999

Row sums of A027052.

Conjecture: (-n+2)*a(n) +(6*n-11)*a(n-1) +(-7*n+1)*a(n-2) +2*(-4*n+27)*a(n-3) +(5*n-28)*a(n-4) +(2*n-3)*a(n-5) +3*(n-5)*a(n-6)=0. - _R. J. Mathar_, May 26 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jun 01 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005115 | OEIS-A161006

Convolution of A000108 (Catalan numbers) with A126120 (Catalan numbers interpolated with 0's).

Conjecture: -(47*n-42)*(n+3)*(n+2)*(n+1)*a(n) + 2*(n+2)*(n+1)*(270*n^2 - 523*n + 126)*a(n-1) - 8*(n+1)*(211*n^3 - 1022*n^2 + 946*n + 63)*a(n-2) + 8*(-212*n^4 - 326*n^3 + 2489*n^2 - 1636*n + 126)*a(n-3) + 16*(985*n^4 - 8154*n^3 + 23771*n^2 - 28164*n + 11151)*a(n-4) + 32*(-386*n^4 + 5771*n^3 - 29314*n^2 + 60721*n - 43533)*a(n-5) - 64*(n-4)*(2*n-9)*(258*n^2 - 1193*n + 1267)*a(n-6) + 128*(n-5)*(2*n-9)*(2*n-11)*(82*n-203)*a(n-7) = 0. - _R. J. Mathar_, Oct 04 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Jun 02 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005116 | OEIS-A161323

Perfect numbers written in base 8.

All members end in 0, 1, 4, 5, or 6. Conjecturally, apart from the first two terms, all terms end in 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jun 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005117 | OEIS-A161460

Positive integers k such that there is no m different from k where both d(k) = d(m) and d(k+1) = d(m+1), where d(k) is the number of positive divisors of k.

Are these values known to be correct, or are they just conjectures? - _Leroy Quet_, Jun 20 2009 [Answer: all the numbers listed in the Data are known to be correct with the exception of 50175 and 59049, which remain conjectural at this time; see the Mathar link. - _Jon E. Schoenfield_, Feb 08 2021] Conjecture: 2 is the only term that is neither a square nor 1 less than a square. - _Jon E. Schoenfield_, Aug 12 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Jun 10 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005118 | OEIS-A161579

Positions n such that A010060(n) = A010060(n+3).

Conjecture: In every sequence of numbers n such that A010060(n)=A010060(n+k), for fixed odd k, the odious (A000069) and evil (A001969) terms alternate. - _Vladimir Shevelev_, Jul 31 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jun 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005119 | OEIS-A161580

Positions n such that A010060(n) + A010060(n+3) = 1.

Conjecture: In every sequence of numbers n such that A010060(n) + A010060(n+k) = 1, for fixed odd k, the odious (A000069) and evil (A001969) terms alternate. [From _Vladimir Shevelev_, Jul 31 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jun 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005120 | OEIS-A161681

Primes that are the difference between a cube and a square (conjectured values).

Primes that are the difference between a cube and a square (conjectured values). The primes found among the differences are sorted in ascending order and unique primes are then extracted. I call this a "conjectured" sequence since I cannot prove that somewhere on the road to infinity there will never exist an integer pair x,y such that x^3-y^2 = 3,5,17,..., missing prime. For example, testing x^3-y^2 for x,y up to 10000, the count of some duplicates are: Conjecture: The number of primes in x^3-y*2 is infinite. Conjecture: The number of duplicates for a given prime is finite. Then there is the other side - the primes that are not in the sequence 3, 5, 17, 29, 31, 37, 41, 43, 59, 73, 97, 101, 103, ... Looks like a lot of twin components here. Do these

have an analytical form? Is there such a thing as a undecidable sequence?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cino Hilliard_, Jun 16 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005121 | OEIS-A161721

Primes p such that the reversal of p is prime and the product of p with its reversal is a palindrome.

Conjecture: except for a(2) which equals 3, all terms can only be composed of the digits 0, 1 or 2. - _Chai Wah Wu_, Jan 07 2015 Conjecture: the digit 2 can only appear once in each term. - _Robert G. Wilson v_, Jan 07 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tanya Khovanova_, Jun 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005122 | OEIS-A161800

G.f.: A(q) = exp(Sum_{n>=1} A002129(n) * 2*A006519(n) * q^n/n).

[The above statements are conjectures needing proof.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 19 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005123 | OEIS-A161869

Convergent of an infinite product of Pascal's triangles aerated by rows.

The sequence may be the binomial transform of A024493 interleaved with zeros. A024493 = (1, 1, 1, 2, 5, 11, 22,...); so the conjecture succeeds through a(12) = A007318 * [1, 0, 1, 0, 1, 0, 2, 0, 5, 0, 11,...]. Calculating more terms of the sequence shows the above conjecture is incorrect. - _Peter Bala_, Jul 07 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Jun 20 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005124 | OEIS-A161890

Numbers such that A010060(n) = A010060(n+9).

Conjecture. In every sequence of numbers n, such that A010060(n)=A010060(n+k), for fixed odd k, the odious (A000069) and evil (A001969) terms alternate. - _Vladimir Shevelev_, Jul 31 2009 This conjecture was actually proved in a later version of the Shevelev arxiv article cited below, and it can also easily be proved by the Walnut prover. - _Jeffrey Shallit_, Oct 12 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jun 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005125 | OEIS-A161942

Odd part of sum of divisors of n.

It is conjectured that iteration of this function will always reach 1. This implies the nonexistence of odd perfect numbers. This is equivalent to the same question for A000593, which can be expressed as the sum of the divisors of the odd part of n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Jun 22 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005126 | OEIS-A161961

Denominator of ratio in lowest terms of pi(n^2)/(n*pi(n)), where pi(x) = A000720(x).

Conjecture: S(n) = pi(n^2)/(n*pi(n)) ~ 1/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Tisdale_, Jun 22 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005127 | OEIS-A162024

Number of n X n binary arrays with rows and columns, considered as binary numbers, in nondecreasing order, and all but the outermost row or column zero.

Conjectures from _Colin Barker_, Mar 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 25 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005128 | OEIS-A162171

Third column of A162170.

a(n) = n*(n+1)/2 * A000111(n-1) (conjectured). - _Mike Tryczak_, Jun 17 2015 The above conjecture by Tryczak is correct. With an offset of 2, the e.g.f. is x^2/2!*(sec(x) + tan(x)). - _Peter Bala_, Sep 08 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Jun 27 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005129 | OEIS-A162180

Let a(n) = d[m(n),i] be the i-th divisor of the integer m(n) and the index j such that d[m(n),i] = d[m(n+1),j], m(n+1) is the first integer > m(n). The sequence a(n), starting with m(1) = 1, i = 1 is given by the conditions: a(1) = 1 and a(n+1) is the first predecessor of a(n) if n even, or the first successor of a(n) if n odd.

Conjecture: the sequence is the union of an infinity subsets of the form {1, 2}, {1, 5, 2, 3}, ..., {1, p, ..., q}, ... where the number 1 is the first element of each subset.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Sep 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005130 | OEIS-A162281

Values of sigma_0(n) for n in A162279.

Conjecture: every composite number appears in this sequence. - _Max Alekseyev_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Jun 29 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005131 | OEIS-A162290

Let A087788(n) = p*q*r, where p<q<r, be the n-th 3-Carmichael number. Then a(n) = (p-1)*(p*q*r-1)/((q-1)*(r-1)).

A.K. Devaraj conjectured that a(n) is always an integer, and this was proved by Carl Pomerance.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _A.K. Devaraj_, Jul 01 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005132 | OEIS-A162314

Row sums of A162313.

Conjecture: for fixed k = 1,2,..., the sequence obtained by reducing a(n) modulo k is eventually periodic with the exact period dividing phi(k), where phi(k) is the Euler totient function A000010. For example, modulo 24 the sequence becomes [1, 4, 0, 16, 0 16, 0, 16, ...] with an apparent period of 2 beginning at a(2). - _Peter Bala_, Jul 08 2022 Conjectural o.g.f. as a continued fraction of Stieltjes type: 1/(1 - 4*x/(1 - 2*x/(1 - 8*x/(1 - 4*x/(1 - ... - 3*n*x/(1 - 2*n*x/(1 - ...)))))))). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 01 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005133 | OEIS-A162475

Expansion of c(x/(1-x)^4), c(x) the g.f. of A000108.

Conjecture: (n+1)*a(n) +3*(2-3n)*a(n-1) +2*(7n-20)*a(n-2) +2*(22-5n)*a(n-3) +(5n-31)*a(n-4) +(8-n)*a(n-5)=0. - _R. J. Mathar_, Nov 17 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005134 | OEIS-A162477

Expansion of (1/(1-x)^2)*c(x/(1-x)^4) where c(x) is the g.f. of A000108.

Conjecture: (n+1)*a(n) +4*(1-2*n)*a(n-1) +6*(n-2)*a(n-2) +2*(7-2*n)*a(n-3) +(n-5)*a(n-4) = 0. - _R. J. Mathar_, Nov 17 2011 G.f.: (1 - 2*x + x^2 - sqrt(1 - 8*x + 6*x^2 - 4*x^3 + x^4))/(2*x). Remark: using this form of the g.f., it is straightforward to prove the above conjectured recurrence. - _Emanuele Munarini_, Aug 31 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005135 | OEIS-A162479

Expansion of 1/((1-x)*sqrt(1-4x/(1-x)^4)).

Conjecture D-finite with recurrence: n*a(n) +3*(2-3n)*a(n-1) +2*(7n-13)*a(n-2)+2*(12-5n)*a(n-3) +(5n-16)*a(n-4) +(4-n)*a(n-5)=0. - _R. J. Mathar_, Nov 17 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005136 | OEIS-A162481

Expansion of (1/(1-x)^3)*c(x/(1-x)^3), c(x) the g.f. of A000108.

Conjecture: (n+1)*a(n) + 2*(1-4*n)*a(n-1) + 2*(5*n-3)*a(n-2) + 4*(2-n)*a(n-3) + (n-3)*a(n-4) = 0. - _R. J. Mathar_, Dec 11 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005137 | OEIS-A162482

Expansion of (1/(1-x)^3)*M(x/(1-x)^3), M(x) the g.f. of Motzkin numbers A001006.

Conjecture: (n+2)*a(n) +4*(-2*n-1)*a(n-1) +18*(n-1)*a(n-2) +13*(-2*n+5)*a(n-3) +17*(n-4)*a(n-4) +3*(-2*n+11)*a(n-5) +(n-7)*a(n-6)=0. - _R. J. Mathar_, Feb 10 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005138 | OEIS-A162533

a(n) = Sum_{k=0..n} binomial(n,2k)*A002426(k).

Conjecture: n*a(n) + (4-5*n)*a(n-1) + 2*(4*n-7)*a(n-2) + 4*(3-n)*a(n-3) + 4*(2-n)*a(n-4) + 4*(n-4)*a(n-5) = 0. - _R. J. Mathar_, Nov 16 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 05 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005139 | OEIS-A162548

A Chebyshev transform of the little Schroeder numbers A001003.

Conjecture: (n+1)*a(n) +3*(-2*n+1)*a(n-1) +(4*n-5)*a(n-2) +12*(-n+2)*a(n-3) +(4*n-11)*a(n-4) +3*(-2*n+7)*a(n-5) +(n-5)*a(n-6)=0. - _R. J. Mathar_, Nov 15 2012. (Formula verified and used for computations. - _Fung Lam_, Feb 19 2014)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jul 05 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005140 | OEIS-A162634

Numerators of fractions with denominators A000215(n) approximating the Thue-Morse constant.

One can prove that if in the sequence of numbers N for which A010060(N+2^n)= A010060(N) you replace the odious (evil) terms by 1's (0's), then we obtain 2^(n+1)-periodic (0,1)-sequence; if you write it in the form .xx...,i.e., as a binary infinite fraction, then the corresponding fraction has the form a(n)/A000215(n). These fractions very fast converge to the Thue-Morse constant .4124540336401...; e.g a(5)/(2^32+1) approximates this constant up to 10^(-9). These approximations differ from A074072-A074073. Conjecture. For n>=1, the fraction a(n)/A000215(n) is a convergent corresponding to the continued fraction for the Thue-Morse constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jul 08 2009, Jul 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005141 | OEIS-A162657

Least number m such that n is the denominator of sigma_{-1}(m), or zero if no such exists.

Conjecture: a(n) is never zero. Checking up to 1000000, the smallest number not found is 210; and a(210) = 26611200.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Jul 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005142 | OEIS-A162761

Minimal total number of floors an elevator must move to transport n people initially waiting at floors i = 1, ..., n to their destination floors n-i+1 (= n, ..., 1), when the elevator can hold at most one person at a time and starts at floor 1, and no passenger may get off the elevator before reaching his/her destination.

The value a(4) = 9 shows that it is not allowed to unload a passenger before he reaches his destination, cf. examples. This implies that there is no better solution than to transport person 1 to floor n, then person n to floor 1, then person 2 to floor n-1, then person n-1 to floor 2, etc., which yields a(n) = (Sum_{i=1..floor(n/2)} (n+1 - 2*i)*2 + 1) - 1 (for n > 1),

equal to the formula conjectured by C. Barker. It would be interesting to consider the variant of optimal solutions involving temporary drop-off of passengers. - _M. F. Hasler_, Apr 29 2019 a(n) = (n^2 + n + (-1)^n - 3)/2 for n > 1. a(n) = 2*a(n-1) - 2*a(n-3) + a(n-4) for n > 4. G.f.: x^2*(2+x^2-x^3)/((1-x)^3*(1+x)). - Conjectured by _Colin Barker_, Jun 10 2012, edited and proved by _M. F. Hasler_, Apr 29 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | Do Zerg (daidodo(AT)gmail.com), Jul 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005143 | OEIS-A162763

Minimal total number of floors an elevator must travel to get n people waiting, respectively, at floors i = 1, 2, ..., n, to their destinations at floors n+1-i (= n, ..., 1), if the elevator can hold at most C = 3 people and starts at floor 1, and no passenger may get off the elevator before reaching his/her destination.

The simple algorithm of taking the first C (or [n/2] if less) persons up to their destination, then the last C persons down to their destination, and if not finished, starting over with floor C+1 as new ground floor with n' = n - 2C, yields an upper bound a(n) <= 2n - 2 + C + a(n - 2C) for n > 2C + 1. In the example section we show that this is not optimal for n = 8 and 9, i.e., 2C + 2 and 2C + 3. I conjecture, however, that this upper bound is sharp (i.e., yields the exact result) for all n > 2C + 3. More precisely, I think the assumption of strict inequality for some larger value of n (the smallest such index) gives a proof by contradiction. - _M. F. Hasler_, May 15 2019 a(n) = 2n - 2 for n < 2C + 2 = 8, a(n) <= 2n - 2 + C + a(n - 2C) else, see comment for proof. Conjectured to hold with equality for all n except n = 2C + {2, 3} = {8, 9} where a(n) = 4(n - C) - 2 = 4n - 14. - _M. F. Hasler_, May 15 2019 G.f.: x*(2 + 2*x^6 - 2*x^7 + x^8 - x^9)/((1 - x)^3*(1 + x)*(1 - x + x^2)*(1 + x + x^2)) (conjectured). - _M. F. Hasler_, May 15 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | Do Zerg (daidodo(AT)gmail.com), Jul 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005144 | OEIS-A162764

Minimal total number of floors that an elevator must move to get n persons waiting, respectively, on floors i = 1, 2, ..., n, to their destination floors n-i+1 (= n, n-1, ..., 1), if the elevator can hold up to C = 4 persons at a time and starts at floor 1, and no passenger may get off the elevator before reaching his destination.

The simple algorithm of taking the first C (or [n/2] if less) persons up to their destinations, then the last C persons down to their destinations, and if not finished, starting over with floor C+1 as new ground floor with n' = n - 2C, yields an upper bound a(n) <= 2n - 2 + C + a(n - 2C) for n > 2C + 1. In the example section we show that this is not optimal for n = 10 and 11, i.e., 2C + 2 and 2C + 3. I conjecture, however, that this upper bound is sharp (i.e., yields the exact result) for all n > 2C + 3. More precisely, I think the assumption of strict inequality for some larger value of n (the smallest such index) gives a proof by contradiction. - _M. F. Hasler_, May 15 2019 a(n) = 2n - 2 for n < 2C + 2 = 10, a(n) <= 2n - 2 + C + a(n - 2C) for n > 2C + 1 (cf. comments for proof), with equality for all known values except n = 10 & 11 where a(n) = 4n - 18; conjectured to be exact for all n > 2C + 3 = 11. - _M. F. Hasler_, May 15 2019 G.f.: 2*x*(1 + x^8 - x^9 + x^10 - x^11)/((1 - x)^3*(1 + x)*(1 + x^2)*(1 + x^4)) (conjectured). - _M. F. Hasler_, May 15 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | Do Zerg (daidodo(AT)gmail.com), Jul 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005145 | OEIS-A163000

Count of integers x in [0,n] satisfying A000120(x) + A000120(n-x) = A000120(n) + 1.

Conjecture: a(n) = A033264(n)* 2^(A000120(n)-1); from [Davis & Webb]. - _R. J. Mathar_, Jul 28 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jul 20 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005146 | OEIS-A163029

Number of n X 3 binary arrays with all 1's connected and a path of 1's from top row to bottom row.

a(n) = 7*a(n-1) - 11*a(n-2) + 6*a(n-3) + a(n-4) - 7*a(n-5) + a(n-6). [Conjectured by _R. J. Mathar_, Aug 11 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 20 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005147 | OEIS-A163183

Primes dividing 2^j + 1 for some odd j.

Conjecture: this sequence has density 7/24 among the primes (see A014663). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christopher J. Smyth_, Jul 22 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005148 | OEIS-A163198

Sum of the cubes of the first n even-indexed Fibonacci numbers.

a(n) = (F(2*n+1)-1)^2*(F(2*n+1) + 2)/4, n>=0. See the Melham reference for a general conjecture. - _Wolfdieter Lang_, Aug 10 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stuart Clary_, Jul 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005149 | OEIS-A163250

a(n) = A000045(n+6) - (n^2 + 4*n + 8).

a(n) = (n-1)^2 + a(n-1) + a(n-2), n>2 (conjectured). - _Bill McEachen_, Jan 20 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jul 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005150 | OEIS-A163493

Number of binary strings of length n which have the same number of 00 and 01 substrings.

Conjecture: -n*a(n) +(2+n)*a(n-1) +(3n-12)*a(n-2) +(12-n)*a(n-3) +(2n-18)*a(n-4)+(56-12n)*a(n-5) +(8n-40)*a(n-6)=0. - _R. J. Mathar_, Nov 28 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christopher Carl Heckman_, Jul 29 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005151 | OEIS-A163511

a(0)=1. a(n) = p(A000120(n)) * Product_{m=1..A000120(n)} p(m)^A163510(n,m), where p(m) is the m-th prime.

If the conjecture given in A364297 holds, then it implies the above conjecture about A007283. See also A364963. - _Antti Karttunen_, Sep 06 2023 Conjecture: a(n^k) is never of the form x^k, for any integers n > 0, k > 1, x >= 1. This holds at least for squares, cubes, seventh and eleventh powers (see A365808, A365801, A366287 and A366391). - _Antti Karttunen_, Sep 24 2023, Oct 10 2023.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Jul 29 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005152 | OEIS-A163553

First differences of A024816.

It appears that (1) a(n) = A120444(n)+1 if and only if n is a prime, (2) if a(n)<0 then A120444(n)<0, and (3) a(n)<=0 whenever n is of the form 6k-1. Are these conjectures easy to prove/disprove? (A120444 is the first difference of A004125 Sum of remainders of n mod k, for k = 1,2,3,...,n). We have a(n) - A120444(n) = sigma(n) - n. Hence a(n) = A120444(n) + 1 if and only if n is prime, and a(n) < 0 implies A120444(n) < 0. The conjecture that a(n) <= 0 whenever n == 5 (mod 6) is false and the first counterexample is a(65) = 6. - _Qiyuan Gu_, Aug 31 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jul 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005153 | OEIS-A163617

a(2*n) = 2*a(n), a(2*n + 1) = 2*a(n) + 2 + (-1)^n, for all n in Z.

Conjecture: a(n) = A003188(n) + (6*n + 1 - (-1)^n)/4. - _Velin Yanev_, Dec 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Aug 01 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005154 | OEIS-A163704

Number of n X 2 binary arrays with all 1s connected, a path of 1s from left column to lower right corner, and no 1 having more than two 1s adjacent.

Conjectures from _Colin Barker_, Mar 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005155 | OEIS-A163714

Number of n X 2 binary arrays with all 1s connected, a path of 1s from top row to bottom row, and no 1 having more than two 1s adjacent.

Conjectures from _Colin Barker_, Feb 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005156 | OEIS-A163733

Number of n X 2 binary arrays with all 1's connected, all corners 1, and no 1 having more than two 1's adjacent.

G.f.: (1-x^3)/(1-x-x^2) (conjecture). - _Paul Barry_, Aug 10 2009 a(n) = round(phi^(k-1)) - round(phi^(k-1)/sqrt(5)), phi = (1 + sqrt(5))/2 (conjecture). - _Federico Provvedi_, Mar 26 2013 G.f.: 1 + 2*x - x*Q(0), where Q(k) = 1 + x^2 - (2*k+1)*x + x*(2*k-1 - x)/Q(k+1); (conjecture), (continued fraction). - _Sergei N. Gladkovskii_, Oct 05 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005157 | OEIS-A163767

a(n) = tau_{n}(n) = number of ordered n-factorizations of n.

a(n) = n^k when n is the product of k distinct primes (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005158 | OEIS-A163774

Row sums of the central coefficients triangle (A163771).

Conjecture: a(n) = Sum_{k=0..n} (-1)^(n-k)*binomial(n+1,k)*binomial(2*k,k). - _Werner Schulte_, Nov 17 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Aug 05 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005159 | OEIS-A163801

a(n) = n - a(a(n-2)) with a(0)=0, a(1)=1.

Conjecture: Which features a certain structure (Comparable to A005206 or A135414). If the (below) following two constructs (C and D) are added on top of their ends (either marked with C or D) one will (if starting with one instance of D) receive the above tree (x marks a node):

猜想/问题提及段 | historical_or_current_mention_unreviewed | Daniel Platt (d.platt(AT)web.de), Aug 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005160 | OEIS-A163824

Number of permutations of length n in the 2x2 double-chevron permutation grid class.

Conjecture: n*(n^2-6*n+11)*a(n) +(-9*n^3+56*n^2-119*n+60)*a(n-1) +2*(13*n^3-83*n^2+193*n-150)*a(n-2) -12*(2*n-5)*(n^2-4*n+6)*a(n-3) =0 . - _R. J. Mathar_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Bevan_, Jun 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005161 | OEIS-A163846

Starting from a(1)=5, a(n+1) is the smallest prime > a(n) such that 2*a(n)-a(n+1) is also prime.

Conjecture: The graph G constructed above consists of exactly two trees: one containing 7 and all odd primes congruent to 2 modulo 3, and another one containing all primes congruent to 1 modulo 3 except 7. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Joseph Stephan Orlovsky_, Aug 05 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005162 | OEIS-A163873

a(n) = n - a(a(n-2)) with a(0) = a(1) = 0.

Conjecture: This recursive structure exists for every sequence of the above mentioned family. The first node of D always has k+1 child nodes where the first one consists of a new copy of D, the second one consists of another node and then D. The remaining child nodes consist of another node and then C. Between the first and the second leaf there is always space for k-1 nodes of construct C. Construct C, on its way from a(n) to n, always crosses exactly k paths (the right ones from construct D).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Daniel Platt (d.platt(AT)web.de), Aug 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005163 | OEIS-A163874

a(n) = n-a(a(n-3)) with a(0) = a(1) = a(2) = 0.

Conjecture: This features a certain structure (similar to the G-sequence A005206 or other sequences of this family: A163875 and A163873). If the (below) following two constructs (C and D) are added on top of their ends (either marked with C or D) one will (if starting with one instance of D) receive the above tree (x marks a node, o marks spaces for nodes that are not part of the construct but will be filled by the other construct):

猜想/问题提及段 | historical_or_current_mention_unreviewed | Daniel Platt (d.platt(AT)web.de), Aug 08 2009, Sep 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005164 | OEIS-A163875

a(n)=n-a(a(n-4)) with a(0)=a(1)=a(2)=a(3)=0.

Conjecture: This features a certain structure (similar to the G-sequence A005206 or other sequences of this family: A163874 and A163873). If the (below) following two constructs (C and D) are added on top of their ends (either marked with C or D) one will (if starting with one instance of D) receive the above tree (x marks a node, o marks spaces for nodes that are not part of the construct but will be filled by the other construct):

猜想/问题提及段 | historical_or_current_mention_unreviewed | Daniel Platt (d.platt(AT)web.de), Aug 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005165 | OEIS-A163940

Triangle related to the divergent series 1^m*1! - 2^m*2! + 3^m*3! - 4^m*4! + ... for m >= -1.

Conjectural explicit formula: T(n,k) = Stirling2(n,n-k) + (n-k)*Sum_{j = 0..k-1} (-1)^j*Stirling2(n, n+1+j-k)*j! for 0 <= k <= n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Johannes W. Meijer_, Aug 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005166 | OEIS-A163945

Triangle interpolating between (-1)^n (A033999) and the swinging factorial function (A056040) restricted to odd indices (2n+1)$ (A002457), read by rows.

Conjectural g.f.: sqrt(1 + t)/(1 + (1 - 4*x)*t)^(3/2) = 1 + (-1 + 6*x)*t + (1 - 12*x + 30*x^2)*t^2 + ... - _Peter Bala_, Nov 10 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Aug 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005167 | OEIS-A163951

The number of functions in a finite set for which the sequence of composition powers ends in a length 2 cycle.

This is to be seen as a conjecture, and the sequence ending with a length 2 cycle does not seem to have such an easy expression.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carlos Alves_, Aug 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005168 | OEIS-A164004

Zero together with row 4 of the array in A163280.

Conjectures from _Colin Barker_, Apr 07 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005169 | OEIS-A164005

Zero together with row 5 of the array in A163280.

Conjecture: a(n) = A100451(n+2). (See A163280.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005170 | OEIS-A164008

Zero together with row 8 of the array in A163280.

Conjecture: a(n) = A028563(n), n > 9. [_R. J. Mathar_, Jul 31 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005171 | OEIS-A164009

Zero together with row 9 of the array in A163280.

Conjecture: a(n) = A028566(n), n > 12. [_R. J. Mathar_, Jul 31 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005172 | OEIS-A164010

Zero together with row 10 of the array in A163280.

Conjecture: a(n) = A028569(n), n > 16. [_R. J. Mathar_, Jul 31 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005173 | OEIS-A164011

Zero together with row 11 of the array in A163280.

Conjecture: a(n) = A098603(n), n > 20. [_R. J. Mathar_, Jul 31 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005174 | OEIS-A164012

Zero together with row 12 of the array in A163280.

Conjecture: a(n) = A119412(n), n > 36. [_R. J. Mathar_, Jul 31 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005175 | OEIS-A164014

Numbers k for which the Hurwitz equation (x_1)^2 + (x_2)^2 + ... + (x_n)^2 = z * x_1 * x_2 * ... * x_k has solutions only for z = k.

Conjectured finite. Any further terms are greater than 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Jul 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005176 | OEIS-A164032

Number of "ON" cells in a certain 2-dimensional cellular automaton.

This box is full (more precisely, every cell in it has more than one ON neighbor), and at generation G+1 we have just 4 ON cells which are now at the corners of a box of edge 2G+1. Until the next power of 2 there is no interaction between the configurations that grow at the four corners, and so a(2^k+j) = 4a(j), as conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Aug 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005177 | OEIS-A164119

Numbers k that are the smallest number that produces the ordered pair (d(k), d(k+1)), where d(k) is the number of divisors of k.

The set of numbers that produces a given ordered pair (i,j) is either empty, finite, or infinite. The pair (3,3) is produced by no number because d(k)=3 only if k is the square of a prime and no two consecutive numbers are squares of primes. Sequence A161460 lists numbers k that produce a unique ordered pair. It appears that the ordered pair (4,2) is produced by an infinite number of k, which is another way of conjecturing that there are an infinite number of safe primes, A005385. The pair (2,4) is produced by primes in A005383. The numbers in A039832 produce the pair (4,4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Aug 10 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005178 | OEIS-A164137

Number of binary strings of length n with equal numbers of 000 and 001 substrings.

Conjecture: a(n) = ((8*n-72)*a(n-10) + (20*n-160)*a(n-9) + (6*n-26)*a(n-8) + (46-5*n)*a(n-7) - 16*a(n-6) + (56-11*n)*a(n-5) + (12-n)*a(n-4) + (n-18)*a(n-3) + n*a(n-2) + 2*n*a(n-1))/n for n>=10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 11 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005179 | OEIS-A164141

Number of binary strings of length n with equal numbers of 001 and 010 substrings.

Conjecture: (n+1)*a(n) +(-4*n-1)*a(n-1) +(8*n-9)*a(n-2) +(-14*n+27)*a(n-3) +(19*n-40)*a(n-4) +2*(-11*n+35)*a(n-5) +4*(5*n-22)*a(n-6) +8*(-n+5)*a(n-7)=0. - _R. J. Mathar_, Dec 10 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 11 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005180 | OEIS-A164338

Conway's creeper sequence.

John Conway calls this sequence "the creeper" and conjectures that the RATS trajectory of every n >= 1 eventually enters a cycle or the creeper. David Wilson confirms this conjecture for n <= 10^10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_, Aug 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005181 | OEIS-A164512

Prime power pairs of form (p^a, q^b = p^a + 1), a >= 1, b >= 1.

It is conjectured that there exist an infinite number of Mersenne primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Forgues_, Aug 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005182 | OEIS-A164586

Row sums of generalized rhombic triangle A164585.

Conjecture: (n+1)*a(n) +(n-1)*a(n-1) +2*(-2*n-1)*a(n-2) +2*(-7*n+10)*a(n-3) +14*(-n+2)*a(n-4) +2*(-2*n+3)*a(n-5) +(n-7)*a(n-6) +(n-5)*a(n-7)=0. - _R. J. Mathar_, Feb 10 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Aug 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005183 | OEIS-A164651

Number of permutations of length n that avoid both 1243 and 2134.

These formulas were conjectured by _Vaclav Kotesovec_ and proved correct by _David Callan_ (see Link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincent Vatter_, Aug 19 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005184 | OEIS-A164660

Numerators of row sums of triangle of rationals A164658/A164659. Definite integral of Chebyshev polynomials of the first kind: Integral_{x=0..1} T(n,x).

Conjecture for the rationals r(n):= A164660(n)/A164661(n): r(n)= 1 if n=0, if n is even r(n) = -1/((n-1)*(n+1)) and if n is odd r(n) = ((-1)^((n-1)/2))/(2*(2*floor((n-1)/4)+1)). a(n+1) = Product_{k=1..n} (1-2*(floor(k^n/n)-floor((k^n -1)/n))) = (-1)^(A003557(n)) for n>0 (conjecture). - _Anthony Browne_, May 29 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 16 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005185 | OEIS-A164704

A006886 mod 9.

The definition implies that if m is A006886(n) for some n, then m == m^2 mod 9, hence m == 0 or 1 mod 9, as conjectured by Paul Curtz. - _N. J. A. Sloane_, Aug 23 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Aug 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005186 | OEIS-A164733

Number of n-digit fixed points under the Kaprekar map A151949.

Conjectures from _Chai Wah Wu_, Apr 13 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph Myers_, Aug 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005187 | OEIS-A164734

Number of n-digit cycles of length 2 under the Kaprekar map A151949.

Conjectures from _Chai Wah Wu_, Apr 13 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph Myers_, Aug 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005188 | OEIS-A164735

Number of n-digit cycles of length 3 under the Kaprekar map A151949.

Conjectures from _Chai Wah Wu_, Apr 13 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph Myers_, Aug 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005189 | OEIS-A164736

Number of n-digit cycles of length 5 under the Kaprekar map A151949.

Conjectures from _Chai Wah Wu_, Apr 13 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph Myers_, Aug 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005190 | OEIS-A164760

Number of n X 3 1..4 arrays with all 1s connected, all 2s connected, all 3s connected, all 4s connected, 1 in the upper left corner, 2 in the upper right corner, 3 in the lower left corner, 4 in the lower right corner, and with no element having more than 3 neighbors with the same value.

Conjectures from _Colin Barker_, Mar 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005191 | OEIS-A164843

The smallest magic constant of an n X n magic square with distinct prime entries.

a(n) >= m(n), where m(n) is the smallest integer of the same parity as n, which is >= (Sum_{k=1..n^2} prime(k+1))/n. For example, Sum_{k=1..5^2} prime(k+1)/5=231.8, so m(5)=233. Conjecture: for n > 4, a(n)=m(n) or a(n)=m(n)+2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Lelechenko_, Aug 28 2009 and _Natalia Makarova_, Sep 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005192 | OEIS-A164848

a(n) = A026741(n)/A051712(n+1).

Conjecture (checked for the first 3000 entries): periodic with a(n+24)=a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Aug 28 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005193 | OEIS-A164901

a(1)=1, a(2) = 2. For n >=3, a(n) = the smallest integer > a(n-1) that is coprime to every sum of any two distinct earlier terms of this sequence.

Conjecture: No two consecutive primes are absent from this sequence (other than 3 and 5). (Equivalently, if p < q < r are consecutive primes and q > 5 is not a term, then p and r are terms.) See A164980. - _Rick L. Shepherd_, Sep 03 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Aug 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005194 | OEIS-A164934

Number of different ways to select 3 disjoint subsets from {1..n} with equal element sum.

Conjecture: a(n) ~ 4^n / (Pi * sqrt(3) * n^3). - _Vaclav Kotesovec_, Oct 16 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Aug 31 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005195 | OEIS-A164952

Primes p with the property: if p/3 is in the interval (p_m, p_(m+1)), where p_m>=3 and p_k is the k-th prime, then the interval (p, 3p_(m+1)) contains a prime.

For k>1 (not necessarily integer), we call a Ramanujan k-prime R_n^(k) the prime a_k(n) which is the smallest number such that if x >= a_k(n), then pi(x)- pi(x/k) >= n. Note that, the sequence of all primes corresponds to the case of "k=oo". These numbers possess the following properties: R_n^(k)~p_((k/(k-1))n) as n tends to the infinity; if A_k(x) is the counting function of the Ramanujan k-primes not exceeding x, then A_k(x)~(1-1/k)\pi(x) as x tends to the infinity; let p be a Ramanujan k-prime, such that p/k is in the interval (p_m, p_(m+1)), where p_m>=3 and p_n is the n-th prime. Then the interval (p, k*p_(m+1)) contains a prime. Conjecture. For every k>=2 there exist non-Ramanujan k-primes, which possess the latter property. For example, for k=2, the smallest such prime is 109. Problem. For every k>2 to estimate the smallest non-Ramanujan k-prime which possesses the latter property.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 01 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005196 | OEIS-A164958

Primes p with the property that if p/3 is in the interval (p_m, p_(m+1)), where p_m>=3 and p_k is the k-th prime, then the interval (3p_m, p) contains a prime.

For k>1 (not necessarily integer), we call a Labos k-prime L_n^(k) the prime a_k(n) which is the smallest number such that pi(a_k(n)) - pi(a_k(n)/k)= n. Note that, the sequence of all primes corresponds to the case of "k=oo". Let p be a k-Labos prime, such that p/k is in the interval (p_m, p_(m+1)), where p_m>=3 and p_n is the n-th prime. Then the interval (k*p_(m), p) contains a prime. Conjecture. For every k>1 there exist non-k-Labos primes, which possess the latter property. For example, for k=2, the smallest such prime is 131. Problem. For every k>1 to estimate the smallest non-k-Labos prime, which possess the latter property. [From _Vladimir Shevelev_, Sep 02 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 02 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005197 | OEIS-A164980

Number of primes between consecutive terms of A164901.

Conjecture: Each term is 0 or 1. I have confirmed this for the first 3499 terms while calculating this b-file based on A164901's b-file.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rick L. Shepherd_, Sep 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005198 | OEIS-A165025

Number of cycles of n-digit numbers (including fixed points) under the base-4 Kaprekar map A165012.

Conjectures from _Colin Barker_, Jun 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph Myers_, Sep 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005199 | OEIS-A165027

Number of n-digit fixed points under the base-4 Kaprekar map A165012.

Conjectures from _Chai Wah Wu_, Apr 13 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph Myers_, Sep 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005200 | OEIS-A165064

Number of cycles of n-digit numbers (including fixed points) under the base-6 Kaprekar map A165051.

G.f.: x*(1 + x + 2*x^5 - 2*x^7 - 3*x^8 - 3*x^9 - x^10 + 2*x^11 + 4*x^12 + 4*x^13 + 4*x^14 + x^15 - 3*x^16 - 3*x^17 - 2*x^18 - x^19 + x^21 + x^22) / ((1 - x)^4*(1 + x)^3*(1 + x^2)*(1 + x + x^2)*(1 + x + x^2 + x^3 + x^4)) (conjectured). - _Colin Barker_, Jun 01 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph Myers_, Sep 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005201 | OEIS-A165066

Number of n-digit fixed points under the base-6 Kaprekar map A165051.

Conjectures from _Chai Wah Wu_, Apr 13 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph Myers_, Sep 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005202 | OEIS-A165105

Number of n-digit fixed points under the base-8 Kaprekar map A165090.

Conjectures from _Chai Wah Wu_, Apr 13 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph Myers_, Sep 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005203 | OEIS-A165125

Number of n-digit fixed points under the base-9 Kaprekar map A165110.

Conjectures from _Chai Wah Wu_, Apr 13 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph Myers_, Sep 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005204 | OEIS-A165201

Expansion of 1/(1-x*c(x)^3), c(x) the g.f. of A000108.

Conjecture: n*(n-3)*a(n) +2*(-4*n^2+15*n-10)*a(n-1) +(15*n^2-69*n+80)*a(n-2) +2*(n-2)*(2*n-5)*a(n-3) =0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005205 | OEIS-A165203

Expansion of (1+x)*c(x)^3/(1-x*c(x)^3), c(x) the g.f. of A000108.

Conjecture: (n+1)*(5*n-31)*a(n) +(5*n^2+74*n+62)*a(n-1) +(-285*n^2+ 1072*n-757)*a(n-2) +(695*n^2-3674*n+4206)*a(n-3) +2*(45*n-74)*(2*n-7)*a(n-4)=0. - _R. J. Mathar_, Dec 11 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005206 | OEIS-A165205

a(n) = C(2n-1,n) + C(2n+1,n+1) - C(0,n).

This sequence has the following conjectural description using cluster algebras. Consider the cyclic directed graph with n vertices as a cluster seed S of type D_n. Let S_n be the set of seeds that can be obtained from S by iterated mutation of the initial cluster variables in S. Mutations of other cluster variables are not allowed. Then the cardinality of S_n is apparently a(n). This can be checked from n=2 (4) to n=8 (8151). - _F. Chapoton_, Apr 03 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005207 | OEIS-A165217

Count of interior bounded regions in a regular 2n-sided polygon dissected by all diagonals parallel to sides.

Conjecture: a(2n) = (2*n-1)*(4*n^3-4*n^2+6*n-3)/3. - Thomas Young (tyoung(AT)district16.org), Dec 23 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | Chintan (timtamboy63(AT)gmail.com), Sep 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005208 | OEIS-A165242

The larger member of the n-th twin prime pair, modulo 8.

Related to the rank of some elliptic curves by the conjecture on page 2 of [Hatley]:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Sep 09 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005209 | OEIS-A165248

Quintisection A061037(5*n+2).

Conjecture: a(n) = 3*a(n-4) - 3*a(n-8) + a(n-12), n>11. - _R. J. Mathar_, Mar 02 2010 The conjecture is equivalent to a(4n) = 5n*(5n+1), a(4n+1) = 5*(20n+9)*(4n+1), a(4n+2) = 5*(10n+7)*(2n+1) and a(4n+3) = 5*(20n+19)*(4n+3). - _R. J. Mathar_, Feb 13 2011 The conjectures can be proved by taking the closed form of A061037, and writing up the quadrisections case by case. - _Bruno Berselli_, Feb 20 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Sep 10 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005210 | OEIS-A165275

Table read by antidiagonals: T(n, k) is the k-th number with n-1 odd-power summands in its base 2 representation.

Layman's conjecture is true (see Fried link). - _Sela Fried_, Nov 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 12 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005211 | OEIS-A165300

a(n) is the smallest number not already present that permits the cyclic repetition of the path 1,2 of the digits in the sequence.

Conjecture. (1) If n > 1, and a(n) begins and ends with 1, then a(n+1) is obtained by deleting the initial 1 of a(n); (2) if a(n) begins with 1 and ends with 2 then a(n+1) is obtained by adding a final 1 to a(n); (3) if a(n) begins with 2 and ends with 1 then a(n+1) is obtained by adding a final 2 to a(n); (4) if a(n) begins and ends with 2 then a(n+1) is obtained by adding an initial 1 to a(n). This has been confirmed through a(140), which has 71 digits (and should be fairly easy to prove). - _John W. Layman_, Sep 22 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_ and _Giorgio Balzarotti_, Sep 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005212 | OEIS-A165372

Number of slanted n X 3 (i=1..n) X (j=i..3+i-1) 1..4 arrays with all 1s connected, all 2s connected, all 3s connected, all 4s connected, 1 in the upper left corner, 2 in the upper right corner, 3 in the lower left corner, and 4 in the lower right corner.

Conjectures from _Colin Barker_, Mar 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005213 | OEIS-A165373

Number of slanted n X 4 (i=1..n) X (j=i..4+i-1) 1..4 arrays with all 1s connected, all 2s connected, all 3s connected, all 4s connected, 1 in the upper left corner, 2 in the upper right corner, 3 in the lower left corner, and 4 in the lower right corner.

Conjectures from _Colin Barker_, Mar 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005214 | OEIS-A165378

Number of slanted n X 4 (i=1..n) X (j=i..4+i-1) 1..4 arrays with all 1s connected, all 2s connected, all 3s connected, all 4s connected, 1 in the upper left corner, 2 in the upper right corner, 3 in the lower left corner, 4 in the lower right corner, and with no element having more than 2 neighbors with the same value.

Conjectures from _Colin Barker_, Mar 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005215 | OEIS-A165381

Number of slanted n X 4 (i=1..n) X (j=i..4+i-1) 1..4 arrays with all 1s connected, all 2s connected, all 3s connected, all 4s connected, 1 in the upper left corner, 2 in the upper right corner, 3 in the lower left corner, 4 in the lower right corner, and with no element having more than 3 neighbors with the same value.

Conjectures from _Colin Barker_, Feb 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005216 | OEIS-A165386

Number of slanted 3 X n (i=1..3) X (j=i..n+i-1) 1..4 arrays with all 1s connected, all 2s connected, all 3s connected, all 4s connected, 1 in the upper left corner, 2 in the upper right corner, 3 in the lower left corner, and 4 in the lower right corner.

Conjectures from _Colin Barker_, Feb 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005217 | OEIS-A165392

Number of slanted 2 X n (i=1..2) X (j=i..n+i-1) 1..4 arrays with all 1s connected, all 2s connected, all 3s connected, all 4s connected, 1 in the upper left corner, 2 in the upper right corner, 3 in the lower left corner, 4 in the lower right corner, and with no element having more than 2 neighbors with the same value.

Conjectures from _Colin Barker_, Feb 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005218 | OEIS-A165394

Number of slanted 2 X n (i=1..2) X (j=i..n+i-1) 1..4 arrays with all 1s connected, all 2s connected, all 3s connected, all 4s connected, 1 in the upper left corner, 2 in the upper right corner, 3 in the lower left corner, 4 in the lower right corner, and with no element having more than 3 neighbors with the same value.

Conjectures from _Colin Barker_, Mar 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005219 | OEIS-A165433

A transform of the double factorial numbers A001147.

Conjecture: 2*a(n) -3*a(n-1) +(3-2*n)*a(n-2) +(2*n-3)*a(n-3)=0. - _R. J. Mathar_, Nov 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Sep 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005220 | OEIS-A165460

The height at the 1/3 point of Jacobi-bridge, computed for 12n+7. a(n) = Sum_{i=0..(4n+2)} J(i,12n+7), where J(i,m) is the Jacobi symbol.

Conjecture: a(2n) = 2*A165605(2n) and a(2n+1) = (2/3)*A165605(2n+1). - _Antti Karttunen_, Oct 05 2009. (If true, then implies also the truth of conjecture in A165462.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005221 | OEIS-A165461

Positions of zeros in A165460.

Also the positions of zeros in A165605 if the conjecture given in A165462 is true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005222 | OEIS-A165462

a(n) = (A165463(n)-3)/4.

Conjecture: These are all those terms of A165602 which = 1 modulo 3. If this is true, then A165461 gives also the positions of zeros in A165605. - _Antti Karttunen_, Oct 05 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005223 | OEIS-A165463

Numbers of the form 12n+7 for which Sum_{i=0..(4n+2)} J(i,12n+7) = 0, where J(i,m) is the Jacobi symbol.

Conjecture: a subset of A165603. See conjectures in A165460 and A165462.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005224 | OEIS-A165537

Number of permutations of length n which avoid the patterns 4213 and 2143.

Conjecture: n*a(n) + 2*(-9*n+7)*a(n-1) + (121*n-204)*a(n-2) + 28*(-14*n+37)*a(n-3) + 16*(42*n-151)*a(n-4) + 4*(-153*n+694)*a(n-5) + 4*(67*n-364)*a(n-6) + 40*(-n+6)*a(n-7) = 0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincent Vatter_, Sep 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005225 | OEIS-A165540

Number of permutations of length n which avoid the patterns 1234 and 2341.

Conjecture: +(14681*n+187954)*(n+3) *a(n) +(14681*n^2-2696783*n-4897218) *a(n-1) +(-888761*n^2+12771539*n+5490342) *a(n-2) +2*(1635223*n^2-14850835*n+13281012) *a(n-3) +2*(-1908503*n^2+15402653*n-25889820) *a(n-4) +156*(2*n-7)*(3301*n-14374) *a(n-5)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincent Vatter_, Sep 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005226 | OEIS-A165542

Number of permutations of length n which avoid the patterns 4231 and 4123.

G.f. conjectured to be non-D-finite (see Albert et al link). - _Jay Pantone_, Oct 01 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincent Vatter_, Sep 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005227 | OEIS-A165543

Number of permutations of length n which avoid both the patterns 3241 and 4321. Or, equivalently, avoids both 1234 and 1342.

Conjecturally, a(n) is the number of permutations pi of length n such that s(pi) avoids the patterns 231 and 321, where s denotes West's stack-sorting map. - _Colin Defant_, Sep 17 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincent Vatter_, Sep 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005228 | OEIS-A165545

Number of permutations of length n which avoid the patterns 2341 and 3421.

G.f. is conjectured to be non-D-finite (see Albert et al link). - _Jay Pantone_, Oct 01 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincent Vatter_, Sep 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005229 | OEIS-A165556

A symmetric version of the Josephus problem read modulo 2.

Conjecture: a(n) = (1 - (-1)^(n + (n + 1)*floor(log_2(n + 1))))/2. - _Velin Yanev_, Nov 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ryohei Miyadera_ and Masakazu Naito, Sep 22 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005230 | OEIS-A165565

a(n) = n - a(a(a(n-3))) with a(0)=a(1)=a(2)=0.

This will give a (k+1)-ary tree which (Conjecture:) features a certain structure (similar to the G-sequence A005206 or other sequences of the above mentioned family: A163873, A163875 and A163874). If the (below) following two constructs (C and D) are added on top of their ends (either marked with C or D) one will (if starting with one instance of D) receive the above tree (x marks a node, o marks spaces for nodes that are not part of the construct but will be filled by the other construct, constructs apply only to this sequence, comments for the whole family!):

猜想/问题提及段 | historical_or_current_mention_unreviewed | Daniel Platt (d.platt(AT)web.de), Sep 22 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005231 | OEIS-A165605

Trisection 3n+1 of A165601.

Positions of zeros probably given by A165461. See the conjectures in A165460 and A165462.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005232 | OEIS-A165640

Number of distinct multisets of n integers, each of which is -2, +1, or +3, such that the sum of the members of each multiset is 3.

Conjecture: a(n) = floor(4*(n+4)/5) - floor(2*(n+4)/3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Sep 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005233 | OEIS-A165659

Numerators of A007504 divided by A033955, starting from the second term of A033955.

Conjecture: with the exception of the second term, 2 <= a(n)/A165660(n) < 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Creighton Dement_, Sep 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005234 | OEIS-A165660

Denominators of A007504 divided by A033955, starting from the second term of A033955.

Conjecture: with the exception of the second term, 2 <= A165659(n)/a(n) < 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Creighton Dement_, Sep 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005235 | OEIS-A165718

Integers of the form k*(k+7)/6.

Conjecturally, the sequence terms are the exponents in the expansion of x/(1 + x) + Sum_{n >= 1} (-1)^n * x^(2*n-1) / Product_{k = 1..n+1} (1 + x^(2*k-1)) = x^3 - x^5 + x^10 - x^13 + x^20 - x^24 + - .... - _Peter Bala_, Nov 20 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Joseph Stephan Orlovsky_, Sep 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005236 | OEIS-A165719

Integers of the form k*(k+9)/8.

Conjecture: Valid k values are pairs of the form (8*m-1,8*m) for m=1,2,3.... If true, 17 is the only prime term. - _Bill McEachen_, May 06 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Joseph Stephan Orlovsky_, Sep 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005237 | OEIS-A165722

Positive integers k such that the sum of decimal digits of (16^k - 1) equals 6*k.

Conjecture: For k > 223, digsum(16^k - 1) < 6*k. This would mean that no further terms exist in the sequence. - _Iain Fox_, Nov 22 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Sep 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005238 | OEIS-A165742

First differences of A003591.

Conjecturally, the limit infimum is infinite. [_Charles R Greathouse IV_, Jun 28 2011]

猜想/问题提及段 | historical_or_current_mention_unreviewed | Mick Purcell (mickpurcell(AT)gmail.com), Sep 25 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005239 | OEIS-A165885

Minimum sum of a set of positive integers such that every positive integer <= n is the sum of 1 or 2 elements of the set.

By considering sets {1, 2, ..., r, 2r, 3r, ..., (s-1)r}, it is conjectured that the asymptotic behavior is a(n) ~ 3/4 * 2^(1/3) * n^(4/3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Bevan_, Sep 29 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005240 | OEIS-A165910

Wiener indices of Fibonacci trees of order k.

D(T(f),k) = A001629(k+1). Conjecture: G.f. x*(1-x+x^2-2*x^3)/((1-x^2-x) * (1+x)^2 * (x^2-3*x+1)^2). [From _R. J. Mathar_, Apr 19 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _K.V.Iyer_, K. R. Udaya Kumar Reddy, Sep 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005241 | OEIS-A165940

G.f.: Sum_{n>=0} a(n)*x^n/2^(n^2+n) = exp(Sum_{n>=1} x^n/[n*2^(n^2)]).

Conjectured to consist entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 01 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005242 | OEIS-A165943

a(n) = A061037(7*n+2).

a(n) = 3*a(n-4) - 3*a(n-8) + a(n-12), n>12. - Conjectured by _R. J. Mathar_, Mar 02 2010, proved by _Robert Israel_, Apr 20 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Oct 01 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005243 | OEIS-A165968

Number of pairings disjoint to a given pairing, and containing a given pair not in the given pairing.

Conjecture: a(n) +2*(-n+1)*a(n-1) +2*(-n+2)*a(n-2)=0. - _R. J. Mathar_, Nov 15 2012 a(n) * (2n-2) satisfies the recurrence of A053871, so Detlef's conjecture was correct. And if we rewrite Mathar's conjecture as b(n) = 2*(n-1)*b(n-1) +2*(n-2)*b(n-2) it becomes quite clear that Mathar's b(n) = a(n-1). - _Sergey Kirgizov_, Jun 03 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | Lewis Mammel (l_mammel(AT)att.net), Oct 02 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005244 | OEIS-A166040

Number of times Sum_{i=1..u} J(i,2n+1) obtains value zero when u ranges from 1 to (2n+1). Here J(i,k) is the Jacobi symbol.

Four bold conjectures by _Antti Karttunen_, Oct 08 2009: 1) All odd natural numbers occur. 2) Each of them occurs infinitely many times. 3) All even natural numbers occur. 4) Each even number > 0 occurs only finitely many times. (The last can be disputed. For example, 6 occurs four times among the first 400001 terms, at the positions 10, 28, 360, 215832.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005245 | OEIS-A166051

Nonsquare integers of the form 4n+1 for which Sum_{i=1..u} J(i,4n+1) is never negative for any u in range [1,(2n)], where J(i,k) is the Jacobi symbol.

Conjecture: There are no more terms after 165. (Checked up to A016813(290511) = 1162045.) If this is true, then also 5, 13 and 37 are only 4k+1 primes in A080114.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005246 | OEIS-A166076

Expansion of (7+5x-sqrt(1-10x-7x^2))/(2(3+5x+2x^2)).

Conjecture: 3*(n)*a(n) +5*(-5*n+9)*a(n-1) +69*(-n+2)*a(n-2) +5*(-11*n+27)*a(n-3) +14*(-n+3)*a(n-4)=0. - _R. J. Mathar_, Feb 10 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005247 | OEIS-A166078

Expansion of (3(1-x)-sqrt(1+6x-7x^2))/(2(1-x)(1-2x)).

Conjecture: n*a(n) + 2*(2*n-5)*a(n-1) + (34-19*n)*a(n-2) + 14*(n-2)*a(n-3)=0. - _R. J. Mathar_, Nov 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 06 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005248 | OEIS-A166088

Odd integers (that is, of the form 2n+1) for which Sum_{i=1..u} J(i,2n+1) obtains value zero exactly 8 times, when u ranges from 1 to (2n+1). Here J(i,k) is the Jacobi symbol.

Of these eight, all are of the form 4k+1, and all others are primes except 65 (= 5*13) and 177 (= 3*59). Conjecture: There are no more terms after the eight one, 193. (Checked up to the 400000th term of A166040, i.e., up to A005408(400000)=800001.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005249 | OEIS-A166102

Odd numbers k such that A166100((k-1)/2)/k is not an integer.

Conjecture: a(n) = 3*A166103(n). (Checked for terms a(1)-a(92).)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005250 | OEIS-A166103

Squares of A166104.

Conjecture: a(n)=A166102(n)/3. (Checked for terms a(1)-a(92).)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005251 | OEIS-A166135

Number of possible paths to each node that lies along the edge of a cut 4-nomial tree, that is rooted one unit from the cut.

Conjecture: 2*n*(2*n+1)*a(n) + (17*n^2-53*n+24)*a(n-1) + 6*(-13*n^2+43*n-36)*a(n-2) - 108*(2*n-5)*(n-3)*a(n-3) = 0. - _R. J. Mathar_, Oct 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | Rick Jarosh (rick(AT)jarosh.net), Oct 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005252 | OEIS-A166168

G.f.: exp(Sum_{n>=1} Lucas(n^2)*x^n/n) where Lucas(n) = A000204(n).

Conjectured to consist entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005253 | OEIS-A166212

Number of 4 X 4 X 4 triangular nonnegative integer arrays, symmetric under 120 degree rotation, with all sums of an element and its neighbors <= n.

Conjectures from _Colin Barker_, Feb 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 09 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005254 | OEIS-A166213

Number of 5 X 5 X 5 triangular nonnegative integer arrays, symmetric under 120 degree rotation, with all sums of an element and its neighbors <= n.

Conjectures from _Colin Barker_, Feb 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 09 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005255 | OEIS-A166228

Alternating sum of large Schroeder numbers.

Conjecture: (n+1)*a(n) +(4-5n)*a(n-1) +(1-5n)*a(n-2) +(n-2)*a(n-3)=0. - _R. J. Mathar_, Nov 17 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 09 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005256 | OEIS-A166251

Isolated primes: Primes p such that there is no other prime in the interval [2*prevprime(p/2), 2*nextprime(p/2)].

Conjecture: a(n)/A000040(n) is asymptotic to 9. This would follow from my conjectured asymptotic proportion of 1's in A102820 (the first differences of A020900).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Oct 10 2009, Oct 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005257 | OEIS-A166268

Bisection 2n of A166050.

Conjecture: a(n) = A166270(n)/2, and also A165605(A005843(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005258 | OEIS-A166269

Bisection 2n+1 of A166050.

Conjecture: a(n) = -(1/2) * A166271(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005259 | OEIS-A166270

Bisection 2n of A165460.

Conjecture: a(n) = 2*A166268(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005260 | OEIS-A166271

Bisection 2n+1 of A165460.

Conjecture: a(n) = -2 * A166269(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005261 | OEIS-A166273

Bisection 2n+1 of A165605.

Conjecture: a(n) = -3 * A166269(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005262 | OEIS-A166287

Number of peak plateaux in all Dyck paths of semilength n with no UUU's and no DDD's (U=(1,1), D=(1,-1)).

Conjecture: n*a(n) +(-4*n+3)*a(n-1) +3*(n-1)*a(n-2) +(n-9)*a(n-3) +(3*n-5)*a(n-4) +(-3*n+7)*a(n-5) +(-2*n+13)*a(n-6) +(n-6)*a(n-7)=0. - _R. J. Mathar_, Jun 14 2016 Conjecture: n*(2*n-5)*(2*n-7)*a(n) -(2*n-7)*(6*n^2-17*n+8)*a(n-1) +(n-2)*(4*n^2-16*n-1)*a(n-2) +(-4*n^3+32*n^2-71*n+44)*a(n-3) +(2*n-3) (6*n^2-37*n+54)*a(n-4) -(n-4)*(2*n-3)*(2*n-5)*a(n-5)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Oct 12 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005263 | OEIS-A166290

Number of UDUD's in all Dyck paths of semilength n with no UUU's and no DDD's (U=(1,1), D=(1,-1)).

Conjecture: -(n+2)*(60283*n-319055)*a(n) +(-60283*n^2-13090*n+236862)*a(n-1) +(588710*n^2-2858167*n+437486)*a(n-2) +(-32043*n^2+2024290*n-2826488)*a(n-3) +(134686*n^2+585099*n-3417942)*a(n-4) +(-514307*n^2+4366464*n-8945002)*a(n-5) +(166729*n-863288)*(n-6)*a(n-6)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Oct 12 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005264 | OEIS-A166300

Number of Dyck paths of semilength n, having no ascents and no descents of length 1, and having no UUDD's starting at level 0.

Conjecture: (n+1)*a(n) +2*(-n+1)*a(n-1) +(-n+1)*a(n-2) +2*(-n+1)*a(n-3) +(n-3)*a(n-4)=0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Nov 07 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005265 | OEIS-A166363

Number of primes in the half-open interval (n*(log(n))^2..(n+1)*(log(n+1))^2].

The Shanks Conjecture and the Cramer-Granville Conjecture tell us that the intervals of length (log(n))^2 are of very critical length (the constant M > 1 of the Cramer-Granville Conjecture definitely matters!). There seems to be some risk that one such interval does not contain a prime. The Wolf Conjecture (which agrees better with numerical evidence) seems more in favor of each interval's containing at least one prime. The similar sequence A345755 relies on intervals that are slightly more than twice as wide as those in the present sequence. A345755 does not include zero entries for n <= 2772, suggesting that the lengths of prime gaps may be bracketed by the two sequences. We conjecture that prime gaps may be larger than log(p)^2, but are not larger than log_2(p)^2. - _Hal M. Switkay_, Aug 29 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Forgues_, Oct 12 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005266 | OEIS-A166408

a(n) = floor(A166407(n)/3).

See the conjecture in A166407. If true, then a(i) = A166406(i)/A005408(i), whenever i is not in A166101.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005267 | OEIS-A166482

a(n) = Sum_{k=0..n} binomial(n+k,2*k)*Fibonacci(2*k+1).

Conjecture: a(n) is the number of tilings of a 4 X 4n rectangle into L tetrominoes (no reflections, only rotations). - _Nicolas Bělohoubek_, Feb 12 2022 The conjecture above was confirmed by _Nicolas Bělohoubek_ and Antonín Slavík. (See links.) - _Nicolas Bělohoubek_, Jan 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005268 | OEIS-A166483

Number of integers in [1..A112141(n)] that are multiples of the first n semiprimes, where A112141(n) is the product of the first n semiprimes.

Conjecture: sequences for 2-ads, 3-ads (composites of 3 primes), 4-ads, etc., converge to 1/2, 1/4, 1/8, ..., respectively.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Tisdale_, Oct 14 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005269 | OEIS-A166549

The number of halving steps of the Collatz 3x+1 map to reach 1 starting from 2n-1.

Conjecture: a(n) = ceiling(log_2((2*n-1)*3^A075680(n))). - _V. Barbera_, Jan 22 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jimin Park_, Oct 16 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005270 | OEIS-A166574

If p, q are successive primes, and there is a number k with p < k <= q such that r = p+k is a prime, then r is in the sequence.

If A(x) is the counting function of a(n) not exceeding x, then, in view of the symmetry, it is natural to conjecture that A(x)~pi(x)/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Oct 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005271 | OEIS-A166587

A signed variant of the Motzkin numbers.

Conjecture: n*(n+1)*a(n) + n*(n+1)*a(n-1) - (5*n-3)*(n-2)*a(n-2) + 3*(n-2)*(n-3)*a(n-3) = 0. - _R. J. Mathar_, Nov 15 2012 G.f. G(x) satisfies (3*x^3 - 2*x^2 - x)*G'(x) - (x+1)*G(x) + 3*x + 1 = 0, from which follows 3*n*a(n) + (-3-2*n)*a(1+n) + (-3-n)*a(n+2) = 0 as well as Mathar's conjecture. - _Robert Israel_, May 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005272 | OEIS-A166588

Partial sums of A097331; binomial transform of A166587.

Conjecture: (-n-1)*a(n) + (n+1)*a(n-1) + 4*(n-2)*a(n-2) + 4*(-n+2)*a(n-3) = 0. - _R. J. Mathar_, Nov 15 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005273 | OEIS-A166597

Let p = largest prime <= n, with p(0)=p(1)=0, and let q = smallest prime > n; then a(n) = q-p.

Note the large prime gap of 72 between 31397 and 31469. This is the prime gap with the largest merit (cf. A111870), 72/log(31397)=6.95352 for primes less than 100000. Also 72/(log(31397))^2=0.67154 (cf. conjectures of Cramer-Granville, Shanks and Wolf) is largest for primes less than 100000. - _Daniel Forgues_, Oct 23 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Forgues_, Oct 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005274 | OEIS-A166602

Numbers k such that Sum_{i=1..k} i^2 divides Product_{i=1..k} i^2.

The b-files suggest conjecturally that this is 1 plus the terms in A067656. - _R. J. Mathar_, Nov 06 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Adamchuk_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005275 | OEIS-A166678

a(n) = pi((sqrt(P(n))+1)^2) - pi(P(n)), where pi(n) = number of primes <= n and P(n) = n-th primorial.

Conjecture: pi((sqrt(P(n))+1)^2) - pi(P(n)) >= n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Tisdale_, Oct 18 2009, Oct 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005276 | OEIS-A166694

A transform of the large Schroeder numbers, A006318.

Conjecture: (n+1)*a(n) +5*(1-2n)*a(n-1) +19*(n-2)*a(n-2) +5*(7-2*n)*a(n-3) +(n-5)*a(n-4)=0. - _R. J. Mathar_, Nov 16 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005277 | OEIS-A166696

A transform of A103210.

Conjecture: (n+1)*a(n) +7*(-2*n+1)*a(n-1) +27*(n-2)*a(n-2) +7*(-2*n+7)*a(n-3) +(n-5)*a(n-4)=0. - _R. J. Mathar_, Feb 10 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005278 | OEIS-A166697

A "Morgan Voyce" transform of A103210.

Conjecture: (n+1)*a(n) + 3*(-5*n+2)*a(n-1) + (41*n-61)*a(n-2) + (-41*n+103)*a(n-3) + 3*(5*n-18)*a(n-4) + (-n+5)*a(n-5) = 0. - _R. J. Mathar_, Feb 10 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Oct 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005279 | OEIS-A166746

Count of propositional theorems up to 10^n in Richard C. Schroeppel's Goedelization of A101273.

Jonathan Vos Post conjectures that a(n) is approximately D^n for some real number D.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Oct 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005280 | OEIS-A166826

Number of n X 2 1..4 arrays containing at least one of each value, all equal values connected, rows considered as a single number in nondecreasing order, and columns considered as a single number in nonincreasing order.

Conjectures from _Colin Barker_, Mar 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005281 | OEIS-A166872

a(n) = floor(n/2 + 2 - sqrt(17/4 + 2*n)).

Each nonnegative integer k occurs 2 to 6 times in the sequence, with 0 occurring 6 times, 3 occurring 4 times, all others either 2 or 3 times. Conjecture: The sequence of integers k which occur 3 times has the g.f. (1 + 2*x^2 - 2*x^3 + x^6 - x^8 - x^9 + x^11)/(1 - 2*x + x^2 - x^8 + 2*x^9 - x^10). - _Robert Israel_, May 27 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Oct 22 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005282 | OEIS-A166945

Records of first differences of A166944.

Conjecture. Each term of the sequence is the greater of a pair of twin primes (A006512).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Oct 24 2009, Nov 05 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005283 | OEIS-A166968

Minimum k such that for all m >= k there is a prime p with m < p < m * (n+1)/n.

The first term was proved by Chebyshev in 1850: for all m > 1, there is a prime number between m and 2m. It is known by Bertrand's Postulate after Joseph Bertrand, who first conjectured it in 1845, and also by Chebyshev's Theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael B. Porter_, Oct 25 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005284 | OEIS-A167023

Fibonacci numbers where both neighbors are semiprimes.

Next term (if it exists) is larger than 10^10000. I conjecture that this sequence is finite: if neighbors of Fibonacci numbers behave randomly, the expected number of remaining terms is about 0.0103 (or 0.00779 if their behavior mod 6 is taken into account). - _Charles R Greathouse IV_, Nov 09 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Joseph Stephan Orlovsky_, Oct 27 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005285 | OEIS-A167137

E.g.f.: P(exp(x)-1) where P(x) is the g.f. of the partition numbers (A000041).

CONJECTURE: Sum_{n>=0} a(n)^m * log(1+x)^n/n! is an integer series in x for all integer m>0; see A167138 and A167139 for examples. Conjecture: Let k be a positive integer. The sequence obtained by reducing a(n) modulo k is eventually periodic with the period dividing phi(k) = A000010(k). For example, modulo 16 we obtain the sequence [1, 1, 5, 15, 1, 7, 1, 7, 1, 7, ...], with an apparent period of 2 beginning at a(4). More generally, we conjecture that the same property holds for integer sequences having an e.g.f. of the form G(exp(x) - 1), where G(x) is an integral power series. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005286 | OEIS-A167138

G.f.: Sum_{n>=0} A167137(n)^2 * log(1+x)^n/n! where Sum_{n>=0} A167137(n)*log(1+x)^n/n! = g.f. of the partition numbers (A000041).

Conjecture: For all integers m > 0, Sum_{n>=0} L(n)^m * log(1+x)^n/n! is an integer series whenever Sum_{n>=0} L(n)*log(1+x)^n/n! is an integer series.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005287 | OEIS-A167152

Lexicographically earliest injective sequence such that a(n) = A007953(a(a(n))), where A007953 = sum of digits (in base 10).

Otherwise, a(n) = n+1 unless n+1 occurred earlier in the sequence (and therefore is of the form k*10^m - 1), in which case a(n) = n+2 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _M. F. Hasler_, Nov 02 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005288 | OEIS-A167154

Numbers where terms in A167153 change parity: a(n)+1 is in A167153, but a(n)-1 is not.

Sequence A167153 consists of runs of numbers of the same parity. It is conjectured that each time the parity changes, there is a gap of 3 numbers, and the sequence goes on with the successor a(n)+1 (of opposite parity) of the first "missing" term a(n) in the run of terms of given parity (a(n-1)+1, a(n-1)+3, ..., a(n)-2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _M. F. Hasler_, Nov 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005289 | OEIS-A167161

The Catapult Sequence.

This sequence is conjectured to be a permutation of the nonnegative integers, generated by the following process:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Weimholt_, Oct 29 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005290 | OEIS-A167162

Inverse Permutation (conjectured) of Catapult Sequence (A167161).

Inverse Permutation (conjectured) of Catapult Sequence (A167161).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Weimholt_, Oct 29 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005291 | OEIS-A167193

a(n) = (1/3)*(1 - (-2)^n + 3*(-1)^n) = (-1)^(n+1)*A167030(n).

This is the inverse binomial transform of 1, 1, 1, 3, 5, 11,.. (continued as in A001045 and conjectured to be equal to A152046).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Oct 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005292 | OEIS-A167199

First column of A167196.

Conjecture: E.g.f.: 2/(2+4*x+x^2) = G(0)/(1+x) where G(k) = 1 - x/((1+x) - x*(1+x)/(x - (1+x)*2/G(k+1))); (recursively defined continued fraction). - _Sergei N. Gladkovskii_, Dec 28 2012.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Oct 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005293 | OEIS-A167230

The matrix exponential of Sierpiński's triangle (A047999) scaled by exp(-1).

Conjecture: All the nonzero entries in this triangle are Bell numbers (A000110).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerald McGarvey_, Oct 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005294 | OEIS-A167276

Primes p such that p^2=x^2+y^2-1 with x and y also prime.

Since (5*x+13)^2 + 1 = (3*x+7)^2 + (4*x+11)^2, it appears that there are infinitely many members of this sequence of the form 5*x+13 where x is an even number, that is the form of A030431(n). See the solution 78 at page 49 in the given reference (250 Problems in Elementary Number Theory) for the related conjecture. - _Altug Alkan_, Mar 30 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Nov 01 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005295 | OEIS-A167440

5th GegenbauerC polynomial evaluated at powers of 2 (multiplied by 5).

Conjectures from _Colin Barker_, Oct 27 2014: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Joseph Stephan Orlovsky_, Nov 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005296 | OEIS-A167493

a(1) = 2; thereafter a(n) = a(n-1) + gcd(n, a(n-1)) if n is odd, and a(n) = a(n-1) + gcd(n-2, a(n-1)) if n is even.

Conjectures. 1) For n >= 2, every difference a(n) - a(n-1) is 1 or prime; 2) Every record of differences a(n) - a(n-1) greater than 3 belongs to the sequence of the greater of twin primes (A006512). Conjecture #1 above fails at n = 620757, with a(n) = 1241487 and a(n-1) = 1241460, difference = 27. Additionally, the terms of related A167495(m) quickly tend to index n/2. So for example, A167495(14) = 19141 is seen at n = 38284. - _Bill McEachen_, Jan 20 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Nov 05 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005297 | OEIS-A167494

List of first differences of A167493 that are different from 1.

Conjecture. All terms of the sequence are primes. The conjecture is false: a(144)=27, a(146)=25, a(158)=45, etc., which are composite numbers. - _Harvey P. Dale_, Dec 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Nov 05 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005298 | OEIS-A167495

Records in A167494.

Conjecture: each term > 3 of the sequence is the greater member of a twin prime pair (A006512). One can formulate a similar conjecture without verification of the primality of the terms (see Conjecture 4 in my paper). [_Vladimir Shevelev_, Nov 13 2009]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Nov 05 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005299 | OEIS-A167638

Number of Dyck paths of semilength n, having no ascents and no descents of length 1, and having no peaks at even level.

Conjecture: g.f. A(x) = 1 + (x^3)*exp(Sum_{n >= 1} g(n, x)*x^(2*n)/n), where g(n, x) = Sum_{k = 0..n} binomial(n, k)^2*(1 + x)^k. Cf. A129509. - _Peter Bala_, Sep 10 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Nov 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005300 | OEIS-A167675

Least prime p such that p-2 has n divisors, or 0 if no such prime exists.

This sequence is the idea of Alonso Del Arte. For n>2, a(n) is conjectured to be the smallest number that is orderly (see A167408) for n-1 values of k. For example, 11 is orderly for k=3 and 9. See A056899 for other primes p that are orderly for two k. It is a conjecture because it is not known whether there are composite numbers that are orderly for more than one value of k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Nov 09 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005301 | OEIS-A167764

a(n) is the index k of k-th prime prime(k) in the smallest concatenation "2 3 ... prime(k)" where prime(n+1) is a factor.

It is conjectured that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Nov 11 2009, Nov 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005302 | OEIS-A167770

a(n) = prime(n)^2 modulo prime(n+1).

In all other cases, a(n) = A076821(n) = (prime(n+1)-prime(n))^2, is highly probable but not proved conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Nov 11 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005303 | OEIS-A167790

a(n) is the index k of k-th prime prime(k) in the smallest sum s(k)=2+3+...+prime(k)=t*prime(n) of first k primes where t is a true divisor and first occurrence of factor prime(n) (n=1,2,3,...)

It is conjectured that the sequence is infinite

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Nov 12 2009, Nov 13 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005304 | OEIS-A167894

Expansion of g.f.: 1/(Sum_{k >= 0} k!*x^k).

G.f.: conjecture: Q(0), where Q(k) = 1 + k*x - (k+1)*x/Q(k+1). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Nov 15 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005305 | OEIS-A168019

Square array A(n,k) read by antidiagonals, in which row n lists the number of partitions of n into parts divisible by k+1.

Let A(n,k) denote the number of partitions of n into parts divisible by k+1. Let p(n) denote the number of partitions of n. If k+1 is a divisor of n then A(n,k) = p(n/(k+1)) otherwise A(n,k) = 0. [Conjectured by _Omar E. Pol_, Nov 25 2009] - this is trivial, just divide each part size by k - _Franklin T. Adams-Watters_, May 14 2010.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Nov 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005306 | OEIS-A168219

Naturals n for which 1 + 10*n^3 (A168147) is prime.

It is conjectured that sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Nov 20 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005307 | OEIS-A168222

a(n) = A006369(A006369(n)).

Conjectures from _Colin Barker_, Aug 15 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Nov 20 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005308 | OEIS-A168243

Expansion of e.g.f. Product_{i>=1} (1 + x^i)^(1/i).

Conjecture: log(a(n)/n!) ~ (log(2) - 1) * log(n). - _Vaclav Kotesovec_, Sep 10 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladeta Jovovic_, Nov 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005309 | OEIS-A168253

a(n) is the smallest prime q such that, for the previous prime p and the following prime r, the fraction (q-p)/(r-q) has denominator n (or 0, if such a prime does not exist).

Conjecture: a(n)>0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005310 | OEIS-A168260

Triangle read by rows, A168258 * the diagonalized variant of A168259.

Conjecture: Row sum ratios tend to phi^2 = 2.6180339... (cf. A168259).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Nov 21 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005311 | OEIS-A168268

G.f.: exp(Sum_{n>=1} A007837(n)*x^n/n), where A007837(n) = number of partitions of n-set with distinct block sizes.

Conjectured to consist entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005312 | OEIS-A168314

Eigensequence of triangle A168313.

Conjectured convergent of a(n)/a(n-1) = (1+sqrt(2)). E.g.: a(19)/a(18) = 2653699/1099137 = 2.4143478...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Nov 22 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005313 | OEIS-A168317

Eigensequence of triangle A168316.

Conjectured convergent of a(n)/a(n-1) = phi^2 = 2.6180339... E.g.: a(19)/a(18) = 10310703/3938272 = 2.6180779....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Nov 22 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005314 | OEIS-A168327

Primes of concatenated form "1 n^3".

(1) It is conjectured that sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Nov 23 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005315 | OEIS-A168375

Natural numbers n for which the concatenation "1 n^3" (A168327) is prime.

It is conjectured that the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Nov 24 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005316 | OEIS-A168417

Primes q for which 1 concatenated with q^3 (A168327) is prime.

It is conjectured that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Nov 25 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005317 | OEIS-A168452

Self-convolution of A005568, where A005568(n) is the product of two successive Catalan numbers C(n)*C(n+1).

Conjecture D-finite with recurrence 3*(n+4)*(n+3)*(n+2)*(n+1)^2*a(n) -4*n*(n+1) *(32*n^3+164*n^2+233*n+75)*a(n-1) +96*(16*n^5+24*n^4-14*n^3-28*n^2-16*n+3) *a(n-2) +1536*(-8*n^4+22*n^3-32*n+15)*a(n-3) -16384*(2*n-5)*(n-1)*(n-2)*(n-3)*(2*n-3)*a(n-4)=0. - _R. J. Mathar_, Nov 22 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 26 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005318 | OEIS-A168494

Sequence with Hankel transform equal to 3^floor(n^2/3).

Conjecture: (n+1)*a(n) +(4-11*n)*a(n-2) +5*(7*n-11)*a(n-2) +(92-37*n) * a(n-3) +6*(2*n-7)*a(n-4) = 0. - _R. J. Mathar_, Sep 30 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 27 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005319 | OEIS-A168503

Expansion of 1/(1-x/(1-2x/(1-3x/(1-x/(1-2x/(1-3x/(1-... (continued fraction).

Conjecture: (n+1)*a(n) +(-13*n+5)*a(n-1) +6*(8*n-13)*a(n-2) +12*(-5*n+13)*a(n-3) +12*(2*n-7)*a(n-4)=0. - _R. J. Mathar_, Feb 10 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Nov 27 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005320 | OEIS-A168506

Number of rooted plane trees of total weight n whose nodes are themselves planted plane trees whose roots are distinguished as either red or blue, the weight of each such node being equal to the size of the corresponding planted tree.

Conjecture: n*(n-1)*(n-6)*a(n) -2*(6*n-35)*(n-1)*(n-2)*a(n-1) +4*(8*n^3-88*n^2+305*n-330)*a(n-2) +8*(8*n^3-92*n^2+314*n-305)*a(n-3) -16*(n-5)*(4*n-19)*(4*n-17)*a(n-4) =0. - _R. J. Mathar_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | Timothy DeVries (tdevries(AT)math.upenn.edu), Nov 27 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005321 | OEIS-A168540

Natural numbers n for which 100n^3 + 27 is prime.

It is conjectured that sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Nov 29 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005322 | OEIS-A168545

Primes p such that the concatenation of p and 29 is a square number: "p 29" = N = m^2.

(1) It is conjectured that the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Nov 29 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005323 | OEIS-A168568

Primes p such that the concatenation p//29 is a squared prime.

(1) Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Nov 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005324 | OEIS-A168597

Squares of the central trinomial coefficients (A002426).

Conjecture: The stronger congruences a(n*p^k) == a(n*p^(k-1)) (mod p^(2*k)) hold for all primes p >= 5 and positive integers n and k. (End) The supercongruence conjectured above by Bala follows from the theorem of Pan and Sun applied to A002426 (see the links): a(n*p^k) - a(n*p^(k-1)) factors as (A002426(n*p^k) - A002426(n*p^(k-1))) * (A002426(n*p^k) + A002426(n*p^(k-1))), and already the first factor has p-adic valuation >= 2*k for p >= 5 by their theorem. - _Andrew R. Epstein_, Jul 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 01 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005325 | OEIS-A168670

Numbers that are congruent to {1, 8} mod 11.

Conjecture: For no term n>1 in the sequence 36*n^2+72*n+35 is equal to p*(p+2), where p, p+2 are twin primes. This conjecture is evident: in fact, it is sufficient to observe that a(2k) = 11*k-3 and a(2k+1) = 11*k+1, therefore 6*a(2k)+7 = 11*(6*k-1) and 6*a(2k+1)+5 = 11*(6*k+1). [_Bruno Berselli_, Jan 07 2013, modified Jul 07 2015]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Dec 02 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005326 | OEIS-A168671

Numbers that are congruent to {1, 10} mod 13.

Conjecture: For no n>1 in the sequence 36*n^2+72*n+35 is equal to p*(p+2), where p, p+2 are twin primes. The conjecture is evident, it can be proved as in A169599. [_Bruno Berselli_, Jan 07 2013]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Dec 02 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005327 | OEIS-A168672

Numbers that are congruent to {2,13} mod 17.

Conjecture: For no term n>2 in the sequence 36*n^2+72*n+35 is equal to p*(p+2), where p, p+2 are twin primes. The conjecture is evident, it can be proved as in A169599. [_Bruno Berselli_, Jan 07 2013]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Dec 02 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005328 | OEIS-A169586

Primes p in A168540 for which q = 3^3 + 10^2*p^3 (A168487) is prime.

It is conjectured that sequence is infinite

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Dec 02 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005329 | OEIS-A169597

Numbers that are congruent to {2, 15} mod 19.

Conjecture: For no term n>2 of the sequence 36*n^2+72*n+35 = (6*n+5)*(6*n+7) is of the form p*(p+2), where p and p+2 are primes. This conjecture is evident: in fact, it is sufficient to observe that a(2k) = 19k-4 and a(2k+1) = 19*k+2, therefore 6*a(2k)+5 = 19*(6*k-1) and 6*a(2k+1)+7 = 19*(6*k+1). [_Bruno Berselli_, Jan 07 2013, modified Jul 07 2015]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Dec 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005330 | OEIS-A169598

Numbers that are congruent to {3,18} mod 23.

Conjecture: For no n in the sequence 36*n^2+72*n+35 = (6*n+5)*(6*n+7) is of the form p*(p+2), where p and p+2 are primes. The conjecture is evident, it can be proved as in A169599. [_Bruno Berselli_, Jan 07 2013]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Dec 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005331 | OEIS-A169599

Numbers that are congruent to {4, 23} mod 29.

Conjecture: For no number n>4 in the sequence 36*n^2+72*n+35 = (6*n+5)*(6*n+7) is of the form p*(p+2), where p and p+2 are primes. This conjecture is evident: in fact, it is sufficient to observe that a(2k) = 29*k-6 and a(2k+1) = 29*k+4, therefore 6*a(2k)+7 = 29*(6*k-1) and 6*a(2k+1)+5 = 29*(6*k+1). [_Bruno Berselli_, Jan 07 2013, modified Jul 07 2015]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Dec 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005332 | OEIS-A169600

Numbers that are congruent to {4, 25} mod 31.

Conjecture: For no n >4 in the sequence, 36*n^2+72*n+35 = (6*n+5)*(6*n+7) is of the form p*(p+2), where p and p+2 are primes. The conjecture is evident, it can be proved as in A169599. [_Bruno Berselli_, Jan 07 2013]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Dec 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005333 | OEIS-A169610

Numbers that are congruent to {5, 30} mod 37.

The conjecture is evident, it can be proved as in A169599. - _Bruno Berselli_, Jan 07 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Dec 03 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005334 | OEIS-A169626

A conjectured sequence of bud numbers for eucalyptus flowers.

A conjectured sequence of bud numbers for eucalyptus flowers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, based on email from _Omar E. Pol_, Mar 07 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005335 | OEIS-A170941

a(n+1) = a(n) + n*a(n-1) - a(n-2) + a(n-3).

Conjecture: a(n)/A000085(n) -> 1/e as n -> inf. In other words, the asymptotic proportion of involutions that contain no adjacent transpositions is conjecturally = 1/e. - _David Callan_, Nov 11 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Barry Cipra_, Feb 09 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005336 | OEIS-A171091

Digits in the order in which they appear in the fractional part of the decimal expansion of Pi.

3,1,8,0,9,6,7,5,2,4 (see A049541) and 6,1,8,0,3,9,7,4,2,5 (see A094214) are the equivalent sequences for 1/Pi and 1/phi. Conjecture: These sequences are not random but are in ratio of 3/2 between the first six and last four digits and the first six digits and last four are the same.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-Yves BOULAY_, Sep 07 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005337 | OEIS-A171141

Numbers that are congruent to {6,33} mod 41.

Conjecture: Numbers n>6 such that 36*n^2+72*n+35 = (6*n+5)*(6*n+7) is not of the form p*(p+2), where p and p+2 are primes. This conjecture is evident: in fact, it is sufficient to observe that a(2k) = 41*k-8 and a(2k+1) = 41*k+6, therefore 6*a(2k)+7 = 41*(6*k-1) and 6*a(2k+1)+5 = 41*(6*k+1). [_Bruno Berselli_, Jan 07 2013]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Dec 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005338 | OEIS-A171154

Smallest prime whose decimal expansion begins with concatenation of first n primes in descending order.

Sequence is conjectured to be infinite. It is conjectured that R(n)=1 for infinite many n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Dec 04 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005339 | OEIS-A171416

A sequence with Somos-4 Hankel transform.

Conjecture: (n+2)*a(n) +(n+1)*a(n-1) +6*(1-n)*a(n-2) +2*(11-5*n)*a(n-3) +(10-3*n)*a(n-4) +(n-5)*a(n-5)=0. - _R. J. Mathar_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Dec 08 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005340 | OEIS-A171462

Number of hands a bartender needs to have in order to win at the blind bartender's problem with n glasses in a cycle.

Conjecture: n > 1: k=1..n: a(n) = -n*min(A191898(n, k)/k). Verified up to n=10000. - _Mats Granvik_, Apr 19 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Ehrenborg_, Dec 09 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005341 | OEIS-A171481

a(n) = a(a(n-1)) + a(n - a(n-1) - 2) with a(0) = 0, a(1) = a(2) = 1.

Conjecture 1: a(n) - a(n-1) = 0 or 1. Conjecture 2: lim_{n->infinity} a(n)/n = e/10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Dec 09 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005342 | OEIS-A171482

a(n) = a(a(n-1)) + a(n - a(n-1) - 2) with a(0) = a(1) = a(2) = 1.

Hofstadter-Conway-like sequence (see A004001). Conjecture 1: a(n) - a(n-1) = 0 or 1. Conjecture 2: lim n ->infinity a(n)/n = 1/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Dec 09 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005343 | OEIS-A171587

Sequence of the diagonal variant of the Fibonacci word fractal. Sequence of the Fibonacci tile.

Conjecture: A171587=(A001950 mod 2), as suggested for n=1,2,...,500 by Mathematica program below. - _Clark Kimberling_, May 31 2011 Proof of Kimberling's 2011 conjecture, i.e., this sequence is the parity sequence of the Upper Wythoff sequence A001950. The following proposition implies the conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexis Monnerot-Dumaine_, Dec 12 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005344 | OEIS-A171751

Maximum k with 1 <= k <= n such that nk + 1 is prime (or 0 if no such prime exists).

It is conjectured that nonzero terms exist for each n: see A089727, the main entry.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rick L. Shepherd_, Dec 17 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005345 | OEIS-A171775

a(n) = smallest number M such that there exist bases b_2, b_3, ..., b_n with the property that M written in base b_k is a k-digit palindrome for all k=2..n.

There are 135382 numbers M <= 2^72 that are both a 10-digit and a 9-digit palindrome (in some bases); 2^72 is the only one of them that is also an 8-digit palindrome in some base. There are 524157 numbers M <= 2^90 that are both an 11-digit and a 10-digit palindrome; 2^90 is the only one of them that is also a 9-digit palindrome. Hence a(10) = 2^72 and a(11) = 2^90, as conjectured in the Formula section. - _Jeff Sponaugle_, Oct 08 2026 Conjecture: a(n) = 2^((n-1)*(n-2)) for n >= 7. - _Max Alekseyev_, Jun 05 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James G. Merickel_, Dec 18 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005346 | OEIS-A171783

Third smallest divisor of smallest number having exactly n divisors.

Conjecture: a(n) = 4 for all prime numbers >= 3 and 3 for all composites. The conjecture is true since the prime factors of each term in A005179 are consecutive primes starting from 2. - _Jianing Song_, Aug 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Oct 12 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005347 | OEIS-A171792

G.f. A(x) satisfies: A(x) = (x + A(x+x^2))/2 with A(0)=0.

Conjecture: a(n) = Sum_{r=ceiling(n/2)..n-1} binomial(r, n-r)*a(r) with a(1) = 1. See [Aspenberg, Perez]. - _Michel Marcus_, Jun 26 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005348 | OEIS-A171814

Triangle T : T(n,k)= A007318(n,k)*A001700(n-k).

Conjectural g.f.: 1/(2*t)*(sqrt((1 - x*t)/(1 - (4 + x)*t)) - 1) = 1 + (3 + x)*t + (10 + 6*x + x^2)*t^2 + ... - _Peter Bala_, Nov 10 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Delélham_, Dec 19 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005349 | OEIS-A171847

Total area under all peakless Motzkin paths of length n (n>=0).

Conjecture D-finite with recurrence (n+2)*a(n) +5*(-n-1)*a(n-1) +(7*n-4)*a(n-2) +2*(-3*n+7)*a(n-3) +(13*n-18)*a(n-4) +11*(-n+3)*a(n-5) +(13*n-60)*a(n-6) +2*(-3*n+11)*a(n-7) +(7*n-38)*a(n-8) +5*(-n+7)*a(n-9) +(n-8)*a(n-10)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Feb 08 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005350 | OEIS-A171849

Total area under all the level steps in all peakless Motzkin paths of length n (n>=0).

Conjecture D-finite with recurrence (n+1)*a(n) -5*n*a(n-1) +(7*n-11)*a(n-2) +2*(-3*n+10)*a(n-3) +(13*n-31)*a(n-4) +11*(-n+4)*a(n-5) +(13*n-73)*a(n-6) +2*(-3*n+14)*a(n-7) +(7*n-45)*a(n-8) +5*(-n+8)*a(n-9) +(n-9)*a(n-10)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Feb 08 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005351 | OEIS-A171851

The difference between the area under a peakless Motzkin path and the number of its U-steps, summed over all peakless Motzkin paths of length n (n>=0).

Conjecture D-finite with recurrence -2*(n+2)*(1088*n-6417)*a(n) +(10968*n^2-55789*n-59252)*a(n-1) +2*(-6704*n^2+45808*n-27647)*a(n-2) +2*(2264*n^2-18729*n+42256)*a(n-3) +2*(-11968*n^2+82555*n-88362)*a(n-4) +(24904*n^2-243375*n+511724)*a(n-5) +4*(-1088*n^2+9137*n-25680)*a(n-6) +2*(6616*n^2-65333*n+164938)*a(n-7) -2*(5616*n-25895)*(n-7)*a(n-8) +(2264*n-10805)*(n-8)*a(n-9)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Feb 08 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005352 | OEIS-A171853

Sum of the trapezoid weights of all peakless Motzkin paths of length n (n>=0).

Conjecture: (-n+1)*a(n) +(3*n-5)*a(n-1) +(-n+7)*a(n-2) -7*a(n-3) -5*a(n-4) -7*a(n-5) +n*a(n-6) +(-3*n+16)*a(n-7) +(n-6)*a(n-8)=0. - _R. J. Mathar_, Dec 07 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Feb 08 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005353 | OEIS-A171854

Number of ladders in all peakless Motzkin paths of length n (n>=0).

Conjecture D-finite with recurrence -(n+2)*(1390*n-8929)*a(n) +(4884*n^2-25542*n-20107)*a(n-1) +(60*n^2-6110*n-2249)*a(n-2) +(-5080*n^2+49134*n-115735)*a(n-3) +(-8476*n^2+66210*n-98877)*a(n-4) +(-3652*n^2+41338*n-109281)*a(n-5) +(2878*n-9993)*(n-7)*a(n-6)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Feb 08 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005354 | OEIS-A171871

Triangle read by rows: Distinct classifications of N elements containing exactly R binary partitions.

Value is A000055(N) when R=N-1 (last term in each row). (Conjectured by _Robert Munafo_ Dec 28 2009, then proved by _Andrew Weimholt_ and _Franklin T. Adams-Watters_ on Dec 29 2009) Conjecture: In general, in each column, the last 2^(R-1) values are the same as the first 2^(N-1) values from the corresponding row of A039754. - _Robert Munafo_, Dec 30 2009

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Munafo_, Jan 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005355 | OEIS-A171876

Mutual solutions to two classification counting problems: binary block codes of wordlength J with N used words; and classifications of N elements by J partitions.

This connection was conjectured by _Robert Munafo_, then proved by _Andrew Weimholt_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Munafo_, Jan 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005356 | OEIS-A171884

Lexicographically earliest injective nonnegative sequence a(n) satisfying |a(n+1) - a(n)| = n for all n.

A correspondent, Dennis Reichard, conjectures that (i) a(n) <= 3.5*n for all n and (ii) the sequence covers 2/3 of all natural numbers. - _N. J. A. Sloane_, Jun 30 2018 [(i) is true: the indices of records for a(n)/n are n = 1, 2, 3, 4, 6, 7, and 2*3^k+2 for k >= 1, with record values 0, 1/2, 1, 1, 3/2, 7/6, 13/7, and (7*3^k-1)/(2*3^k+2) for k >= 1, so a(n) <= 3.5*n. (ii) needs further justification: the lower natural density is lim_{k->+oo} #{terms <= 7*3^k-2}/(7*3^k-2) = lim_{k->+oo} (4*3^k-1)/(7*3^k-2) = 4/7, and the upper natural density is lim_{k->+oo} #{terms <= 5*3^k-2}/(5*3^k-2) = lim_{k->+oo} (4*3^k-1)/(5*3^k-2) = 4/5. - _Jianing Song_, Oct 07 2022]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Munafo_, Mar 11 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005357 | OEIS-A171889

a(n) = number immediately following n-th 0 in A181391.

This is associated with conjecture (ii) in A181391 (van Eck's sequence). - _Po-chia Chen_, Jun 20 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 18 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005358 | OEIS-A171929

Odd numbers whose abundancy is closer to 2 than any smaller odd number.

So far all known perfect numbers (abundancy = 2) are even, cf. A000396 = (6, 28, 496, 8128, ...). It has been conjectured but not proved that there are no odd perfect numbers. This sequence provides the list of odd numbers that approach perfection (odd numbers which abundancy is closer to two than the abundancy of any smaller odd number).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sergio Pimentel_, Jan 05 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005359 | OEIS-A171949

P-positions for game of Mark-4.

Conjecture: For n>=2, a(n) = A217319(n-1). - _Vladimir Shevelev_, Mar 18 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005360 | OEIS-A172025

Expansion of (2/(3*sqrt(1-4*z)-1+4*z))*((1-sqrt(1-4*z))/(2*z))^k with k=3.

Conjecture: 2*n*(n+3)*a(n) + (-7*n^2 - 17*n - 8)*a(n-1) -2*(n+2)*(2*n+1)*a(n-2) = 0. - _R. J. Mathar_, Feb 19 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Jan 23 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005361 | OEIS-A172062

Expansion of (2/(3*sqrt(1-4*z)-1+4*z))*((1-sqrt(1-4*z))/(2*z))^k with k=5.

Conjecture: 2*n*(n+5)*(3*n+7)*a(n) - (n+3)*(21*n^2+79*n+80)*a(n-1) - 2*(3*n+10)*(2*n+3)*(n+2)*a(n-2) = 0. - _R. J. Mathar_, Feb 19 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Jan 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005362 | OEIS-A172063

Expansion of (2/(3*sqrt(1-4*z)-1+4*z))*((1-sqrt(1-4*z))/(2*z))^k with k=6.

Conjecture: 2*n*(n+6)*(n+3)*a(n) -(7*n^3+59*n^2+166*n+160)*a(n-1) -2*(2*n+5)*(n+4)*(n+2)*a(n-2)=0. - _R. J. Mathar_, Feb 19 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Jan 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005363 | OEIS-A172064

Expansion of (2/(3*sqrt(1-4*z)-1+4*z))*((1-sqrt(1-4*z))/(2*z))^k with k=7.

Conjecture: 2*n*(n+7)*(3*n+11)*a(n) -(21*n^3+212*n^2+719*n+840)*a(n-1) -2*(2*n+5)*(n+3)*(3*n+14)*a(n-2)=0. - _R. J. Mathar_, Feb 19 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Jan 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005364 | OEIS-A172065

Expansion of (2/(3*sqrt(1-4*z)-1+4*z))*((1-sqrt(1-4*z))/(2*z))^k with k=8.

Conjecture: 2*n*(n+8)*(3*n+13)*a(n) -(21*n^3 + 247*n^2 + 980*n + 1344)*a(n-1) - 2*(n+3)*(3*n+16)*(2*n+7)*a(n-2) = 0. - _R. J. Mathar_, Feb 29 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Jan 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005365 | OEIS-A172066

Expansion of (2/(3*sqrt(1-4*z)-1+4*z))*((1-sqrt(1-4*z))/(2*z))^k with k=9.

Conjecture: 2*n*(n+9)*(n+5)*a(n) -(7*n^3+94*n^2+427*n+672)*a(n-1) -2*(2*n+7)*(n+6)*(n+4)*a(n-2)=0. - _R. J. Mathar_, Feb 19 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Jan 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005366 | OEIS-A172067

Expansion of (2/(3*sqrt(1-4*z)-1+4*z))*((1-sqrt(1-4*z))/(2*z))^k with k=10.

Conjecture: 2*n*(n+10)*(3*n+17)*a(n) - (21*n^3 + 317*n^2 + 1622*n + 2880)*a(n-1) - 2*(3*n+20)*(n+4)*(2*n+9)*a(n-2) = 0. - _R. J. Mathar_, Feb 21 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Jan 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005367 | OEIS-A172125

a(n) = smallest prime of the form n*(n+1)*(n+2)...(n+k-1) - 1, or zero if no such a prime exists.

A conjecture is that no term is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Cheng Zhang (cz1(AT)rice.edu), Jan 26 2010, Mar 02 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005368 | OEIS-A172164

Differences between numbers of triangles entirely contained in two consecutive turns of Pythagoras's snail (Theodorus spiral).

Conjecture : The terms are only 18,19,20,21 (From the first thousand turns, there are 2,3% of 18, 36,5% of 19, 46,2% of 20 and 15% of 21). No period found. Probably due to Pi transcendence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sébastien Dumortier_, Jan 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005369 | OEIS-A172236

Array A(n,k) = n*A(n,k-1) + A(n,k-2) read by upward antidiagonals, starting A(n,0) = 0, A(n,1) = 1.

.....1.....1.....1/6 (conjectured)...2/3 (conjectured)...1/6 (conjectured)* .....1.....5.....1/2 (conjectured).....0.....1/2 (conjectured)*

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Jan 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005370 | OEIS-A172251

Arises in the representability of integers as sums of triangular numbers.

Wieb Bosma, p.10: Following the bounds given in the proof of Theorem 1.6, computational evidence suggests that... a proof of the above identity using the techniques of Bhargava and Hanke developed in the proof of the 290-Theorem may require a careful analysis of a possible Siegel zero. The sequence given is thus conjectured to be complete as shown.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jan 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005371 | OEIS-A172253

Numbers k such that the squarefree kernel of 9^k*(9^k - 1) is 3*(9^k - 1)/4.

Conjecture: This sequence is infinite. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jan 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005372 | OEIS-A172271

Smaller member p of a twin prime pair (p,p+2) with a cube sum N^3.

It is conjectured that the number of twin prime pairs is infinite, one of the great open questions in number theory. It is conjectured that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Jan 30 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005373 | OEIS-A172434

G.f.: Sum_{n>=0} a(n)*x^n/n!^4 = [Sum_{n>=0} x^n/n!^4]^3.

Conjecture: the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) hold for primes p >= 5 and positive integers n and k. Extending the sequence to negative n via a(-n) = Sum_{k = 0..n} C(-n,k)^4 * Sum_{j = 0..k} C(k,j)^4 produces the sequence [1, 3, 1491, 1663491, 2742551763, 5400215949753, 12106456954153971, 29631476653021123107, ...], which appears to satisfy the same supercongruences. Cf. A141057. - _Peter Bala_, Aug 05 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005374 | OEIS-A172468

Numbers such that A166986(n)/2 - A167389(n) = 1.

It is conjectured that the successive differences of this sequence, A172470, are limited to three numbers: 9, 52 and 61 where it is noted that 61 - 52 = 9, (52 + 2)/9 = 6, (61 + 2)/9 = 7 and we have lcm(9, 52, 61) = 28548 = 13^4 - 13 and 6 + 7 = 13. The conjectured closure of {9,52,61} then follows from plugging [IV] into [III] to get c = lim_{n->oo} c_n = 7/log(16) - 5/2 and by inspection of (n/log(2)) mod 1 for n <= 61, noting that -log(2)/2 satisfies [V] so that c_n satisfies [I] and [II]. Some basic pigeonholing further restricts the differences to three admissible runs [{52,9}, {61,61,9}, {61,61,61,9}] and issuing the sequence simplifies to evaluating ((n + 2)/log(2)) mod 1 once per run, comparing it to two constants, and measuring a low watermark for c_n as infrequently as possible.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stephen Crowley_, Feb 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005375 | OEIS-A172470

First differences of A172468.

The numbers in the sequence are conjectured to be restricted to 9,52, or 61. Note that 61-52 = 9, (52+2)/(61-52) = (52+2)/9 = 6, (61+2)/(61-52) = (61+2)/9 = 7 and we have lcm(9,52,61) =28548 = 13^4 - 13 and (52+2)/9 + (61+2)/9 = 6+7 = 13.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stephen Crowley_, Feb 04 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005376 | OEIS-A172487

Lesser of twin primes in A172240.

For a(n) > 3, the first differences of the sequence are divisible by 6. (Is this a conjecture or a theorem?)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Teofilatto_, Nov 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005377 | OEIS-A172495

a(n) = numerator of fraction whose decimal representation is (n).(1)(2)(3)...(n-1)(n).

Sequence of denominators: 10, 25, 1000, 5000, 20000, 15625, 10000000, 50000000, 1000000000, 10000000000, ... Conjecture: this sequence is not equal to the sequence A078257. I conjecture that it is the same as A078257. - _Franklin T. Adams-Watters_, Mar 29 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Feb 05 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005378 | OEIS-A172499

a(n) = numerator of fraction whose decimal representation is (n).(1)(2)(3)...(n-1)(n)... with repeating part (1)(2)(3)...(n-1)(n).

Sequence of denominators: 9, 33, 333, 9999, 33333, 333333, 9999999, 11111111, 111111111, 99999999999, 3333333333333, ... Conjecture: this sequence is not equal to the sequence A172498.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Feb 05 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005379 | OEIS-A172505

a(n) = numerator of fraction a/b, where gcd(a, b) = 1, whose decimal representation has the form (1)(2)(3)...(n-1)(n).(1)(2)(3)...(n-1)(n)... with period (1)(2)(3)...(n-1)(n).

Sequence of denominators: 9, 33, 333, 9999, 33333, 333333, ... Conjecture: this sequence is not equal to the sequence A172498.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Feb 05 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005380 | OEIS-A172506

a(n) = numerator of fraction a/b, where gcd(a, b) = 1, whose decimal representation has the form (1)(2)(3)...(n-1)(n).(1)(2)(3)...(n-1)(n).

Sequence of denominators: 10, 25, 1000, 5000, 20000, 15625, 10000000, 50000000, ... Conjecture: this sequence is not equal to the sequence A078257. The conjecture is false: the denominators here are the same as in A078257.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Feb 05 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005381 | OEIS-A172507

a(n) = numerator of fraction a/b, where gcd(a, b) = 1, whose decimal representation has the form (n).(n).

Sequence of denominators: 10, 25, 1000, 5000, 20000, 15625, 10000000, 50000000, ... Conjecture: this sequence is not equal to the sequence A078267.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Feb 05 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005382 | OEIS-A172572

Number of 3*n X 6 0..1 arrays with row sums 2 and column sums n.

Conjectured recurrence of order 3 and degree 9: (n + 2)*(n + 3)^5*(2*n + 5)*(62*n^2 + 217*n + 191)*a(n + 3) - 9*(n + 2)*(3*n + 7)*(3*n + 8)*(5084*n^6 + 68634*n^5 + 383756*n^4 + 1137319*n^3 + 1884032*n^2 + 1653960*n + 601185)*a(n + 2) - 9*(3*n + 4)*(3*n + 5)*(3*n + 7)*(3*n + 8)*(31372*n^5 + 313720*n^4 + 1227805*n^3 + 2354425*n^2 + 2220988*n + 827860)*a(n + 1) + 20250*(n + 1)*(3*n + 1)*(3*n + 2)*(3*n + 4)*(3*n + 5)*(3*n + 7)*(3*n + 8)*(62*n^2 + 341*n + 470)*a(n) = 0. - _Christoph Koutschan_, Feb 26 2023 Conjecture: a(n) ~ 3^(3*n + 1/2) * 5^(3*n + 5/2) / (128 * Pi^(5/2) * n^(5/2)), based on the recurrence by _Christoph Koutschan_. - _Vaclav Kotesovec_, Feb 26 2023 Conjectures: 1) a(p) == 90 (mod p^2) for all primes p (checked up to p = 73).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 06 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005383 | OEIS-A172671

Number of 3*n X 6 0..2 arrays with row sums 2 and column sums n.

Conjectured recurrence of order 4 and degree 13: (n + 2)*(n + 3)^3*(3784*n^4 + 32164*n^3 + 100749*n^2 + 137862*n + 69678)*(n + 4)^5*a(n + 4) - (n + 2)*(n + 3)^3*(3*n + 10)*(3*n + 11)*(3799136*n^7 + 72183584*n^6 + 579689880*n^5 + 2548427912*n^4 + 6617561702*n^3 + 10141503096*n^2 + 8487349821*n + 2991586122)*a(n + 3) - 9*(n + 2)*(3*n + 7)*(3*n + 8)*(3*n + 10)*(3*n + 11)*(10844944*n^8 + 222321352*n^7 + 1973930222*n^6 + 9916013134*n^5 + 30831383530*n^4 + 60768378830*n^3 + 74160044251*n^2 + 51243135187*n + 15352797306)*a(n + 2) + 9*(3*n + 4)*(3*n + 5)*(3*n + 7)*(3*n + 8)*(3*n + 10)*(3*n + 11)*(29681696*n^7 + 504588832*n^6 + 3602458816*n^5 + 14001842392*n^4 + 32010306742*n^3 + 43078657918*n^2 + 31639900193*n + 9799573455)*a(n + 1) + 416745*(n + 1)*(3*n + 1)*(3*n + 2)*(3*n + 4)*(3*n + 5)*(3*n + 7)*(3*n + 8)*(3*n + 10)*(3*n + 11)*(3784*n^4 + 47300*n^3 + 219945*n^2 + 450988*n + 344237)*a(n) = 0. - _Christoph Koutschan_, Feb 26 2023 Conjecture: a(n) ~ 3^(3*n + 1/2) * 7^(3*n + 5/2) / (2^(19/2) * Pi^(5/2) * n^(5/2)), based on the recurrence by _Christoph Koutschan_. - _Vaclav Kotesovec_, Feb 27 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 06 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005384 | OEIS-A173039

Odd numerators of the fractions (1/4-1/n^2), n>= -2.

Conjecture: a(n)= +3*a(n-3) -3*a(n-6) +a(n-9). G.f.: (3+x+3*x^2-14*x^3-6*x^4-30*x^5-21*x^6-3*x^7-5*x^8) / ((x-1)^3*(1+x+x^2)^3). - _R. J. Mathar_, Dec 02 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Nov 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005385 | OEIS-A173102

Number of partitions x + y = z with {x,y,z} in {1,2,3,..,3*n} and z > y >= x.

Conjectures from _Colin Barker_, Sep 04 2013: (Start) Conjecture: a(n) = Sum_{j=1..n} Sum_{i=1..n} ceiling((i+j-n+3)/2). - _Wesley Ivan Hurt_, Mar 12 2015 This also implies that the above conjectures on the recurrence and g.f. are true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Feb 09 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005386 | OEIS-A173195

Values of k such that 4^x + 4^y + 4^z = k^2 with arbitrary integers x <= y <= z.

Conjecture: a(n) = 3*A263132(n). - _George Beck_, May 05 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Feb 12 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005387 | OEIS-A173238

Triangle by columns, A000041 in every column shifted down twice for columns > 0.

Conjectures: Given the infinite set of triangles with A000041 in every column shifted down 0, 1, 2, ... n times, row sums of n-th triangle (where A173238 = 2nd in the set) = the numbers of n's in all partitions of n. E.g., row sums = of A173238 = A024786, the numbers of 2's in all partitions of n. Second conjecture: Given the infinite set of analogous triangles with columns shifted down 2, 3, 4, ..., k times, we let such triangles = T(k) and perform lim_{n->inf} T^n(k), obtaining the left-shifted vectors considered as sequences.

The conjecture states that the infinite set of such left-shifted vectors = the Euler transform of the infinite set of Ruler functions starting with the ruler function for k=2 = A001511: (1, 2, 1, 3, 1, 2, 1, ...) Third conjecture: Let P(x) = polcoeff A000041 = (1 + x + 2x^2 + 3x^3 + ...), and A(k)(x) = the Euler transform of k-th ruler sequence, (k=2,3,...). Then P(x) = A(k)(x) / A(k)(x^k). The conjecture extends the analogous conclusions to all k. Proof of second conjecture received from Helmut Prodinger 02/28/10 with a summary by _R. J. Mathar_:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Feb 13 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005388 | OEIS-A173239

Triangle by columns, A000041 shifted down thrice, k>=0.

Refer to A173238 comments for three conjectures relating A000041 to the infinite set of generalized ruler function sequences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Feb 13 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005389 | OEIS-A173300

a(n) is the denominator of the fraction f = x^n + y^n given that x + y = 1 and x^2 + y^2 = 2.

a(n) = 4*a(n-4), for n >= 7. - conjectured by _Paolo Xausa_, Feb 02 2024; proved by _Max Alekseyev_, Mar 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Feb 15 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005390 | OEIS-A173348

Numbers x such that 0 < |x^7 - y^2| < x^(5/2) for some number y.

Beukers and Stewart conjecture that for coprime integers n and m with n > m >= 2, and for any c > 0, the inequality 0 < |x^n - y^m| < c*X^(1-1/n-1/m) is true for infinitely many positive integers x and y, where X = max(x^n,y^m). They compute such x for 34 pairs (n,m). Given x, it is easy to compute y = round(x^(n/m)). Their tables have been extended to include all terms < 10^7 (or higher to obtain more terms).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Feb 22 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005391 | OEIS-A173419

Length of shortest computation yielding n using addition, subtraction and multiplication (starting from 1).

Conjecture: if n is prime then a(n) >= a(n-1). The conjecture is true for n < 1800. - _Dmitry Kamenetsky_, Dec 26 2019 The conjecture of Dmitry Kamenetsky above holds for every prime p < 3359, but fails at p = 3359, where a(3359) = 7 < 8 = a(3358); a shortest computation for 3359 = 15^3 - 16 is 1, 2, 4, 16, 15, 225, 3375, 3359. There are exactly four counterexamples below 5000: p = 3359; p = 3623 = 29*125 - 2, with a(3623) = 8 < 9 = a(3622), via 1, 2, 4, 5, 25, 125, 29, 3625, 3623; p = 4909 = 17^3 - 4, with a(4909) = 7 < 8 = a(4908), via 1, 2, 4, 16, 17, 289, 4913, 4909; and p = 4943 = 17*291 - 4, with a(4943) = 8 < 9 = a(4942), via 1, 2, 4, 16, 17, 289, 291, 4947, 4943. In each case the prime lies at short additive distance from a product of cheaply reachable numbers. Verified by exhaustive breadth-first enumeration of all computations of length <= 8 with no bound on intermediate values, extended to length 9 for the nine pairs (p, p-1) with p prime < 5000 left undecided at depth 8; all nine satisfy a(p) = a(p-1) = 9. - _Rosario Patanè_, Aug 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Feb 17 2010, Apr 22 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005392 | OEIS-A173515

Consider positive integer solutions to x^3 + y^3 = z^3 - n or 'Fermat near misses' of 1, 2, 3 ... Arrange known solutions by increasing values of n. Sequence gives value of lowest z for a given n.

Author conjectures that no explicit formula or recurrence exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andy Martin_, Feb 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005393 | OEIS-A173518

Solutions z of the Diophantine equation x^3 + y^3 = 6z^3.

A. Nitaj proved Erdős's conjecture (1975) and claimed that there exist infinitely many triples of 3-powerful numbers a,b,c with (a,b) = 1, such that a+b=c, because the equation x^3 + y^3 = 6z^3 admits an infinite number of solutions, and given by the recurrence equations (see formula). It is proved that a=x(k)^3, b=y(k)^3, and c=6c^3, and are 3-powerful numbers for each k >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Feb 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005394 | OEIS-A173524

a(n) = A053737(4^k+n-1) in the limit k->infinity, where k plays the role of a row index in A053737.

It appears that if A053737 is written as a triangle then the rows are initial segments of the present sequence; see the conjecture in A000120. Conjecture: Fixed point of the morphism 1->{1,2,3,...,b}, 2->{2,3,4,...,b+1}, j->{j,j+1,...,j+b-1} for b=4. [_Joerg Arndt_, Dec 08 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Feb 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005395 | OEIS-A173525

a(n) = 1 + A053824(n-1), where A053824 = sum of digits in base 5.

See conjecture in the entry A000120, and the case of base 2 in A063787.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Feb 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005396 | OEIS-A173526

a(n) = 1 + A053827(n-1), where A053827 is the sum-of-digits function in base 6.

See conjecture in the entry A000120. Conjecture: Fixed point of the morphism 1->{1,2,3,...,b}, 2->{2,3,4,...,b+1},

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Feb 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005397 | OEIS-A173527

a(n) = 1 + A053828(n-1), where A053828 is the sum of digits in base 7.

See conjecture in the entry A000120. Conjecture: Fixed point of the morphism 1->{1,2,3,...b}, 2->{2,3,4...,b+1}, j->{j,j+1,...,j+b-1} for b=7. [_Joerg Arndt_, Dec 08 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Feb 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005398 | OEIS-A173528

a(n) = 1 + sum of digits of n-1 written in base 8.

If A053829 is written as a triangle then the rows converge to this sequence; see the conjecture in A000120.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Feb 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005399 | OEIS-A173614

a(n) = lcm_{p is prime and divisor of n} p-1.

Conjecture: a(n) = A002322(A007947(n)). - _Velin Yanev_, Feb 06 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Feb 22 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005400 | OEIS-A173643

Positive numbers of form 2^m - 2^l - 3*2^k (A172233) divisible by 9, divided by 9.

Conjecture: Odd part of a(n) is of form [(1/6)*(8^m-(-1)^m-3)*4^k-1]/3, k,m>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Nov 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005401 | OEIS-A173644

a(n) = smallest positive integer m such that n^2+7m is a square.

Conjecture: a(n)= +2*a(n-7) -a(n-14).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Dec 05 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005402 | OEIS-A173693

a(n) = ceiling(A107293(n)/2).

Conjecture: a(n) = a(n-1) + a(n-2) - a(n-3) + a(n-5) + a(n-31) - a(n-32) - a(n-33) + a(n-34) - a(n-36). - _R. J. Mathar_, Feb 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger L. Bagula_, Nov 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005403 | OEIS-A173774

The arithmetic mean of (21*k + 8)*binomial(2*k,k)^3 (k=0..n-1).

On Feb 10 2010, _Zhi-Wei Sun_ introduced the sequence and conjectured that each term a(n) is an integer divisible by 4*binomial(2*n,n). On Feb 11 2011, _Kasper Andersen_ confirmed this conjecture by noting that the sequence b(n) = a(n)/(4*binomial(2*n,n)), for n > 0, coincides with A112029. It was proved that for every prime p and positive integer a we have a(p^a) == 8 + 16*p^3*B_(p-3) (mod p^4), where B_0, B_1, B_2, ... are Bernoulli numbers. Given a prime p, it has been conjectured that Sum_{k=0..(p-1)/2} (21*k + 8)*binomial(2*k,k)^3 == 8*p + (-1)^((p-1)/2)*32*p^3*E_(p-3) (mod p^4) if p > 3 (where E_0, E_1, E_2, ... are Euler numbers), and that Sum_{k=0..floor(2p^a/3)} (21*k + 8)*binomial(2*k,k)^3 == 8*p^a (mod p^(a + 5 + (-1)^p)) if a is a positive integer with p^a == 1 (mod 3). He also observed that b(n) = a(n)/(4*binomial(2*n,n)) is odd if and only if n is a power of two.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005404 | OEIS-A173777

Infinite sequence gradually builds a triangle plus another more widely spaced triangle on top of it, or overlapping, if you will.

Conjectures from _Colin Barker_, Dec 01 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eric Werner (eric.werner(AT)gmx.net), Feb 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005405 | OEIS-A173781

a(n) is the smallest entry of the n-th column of the matrix of Super Catalan numbers S(m,n).

S(m,n) is a positive integer, but a combinatorial interpretation of S(m,n) is an open problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Joseph Alfano (jalfano(AT)assumption.edu), Feb 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005406 | OEIS-A173948

a(n) = denominator of (Zeta(2, 1/4) - Zeta(2, n+1/4)), where Zeta is the Hurwitz Zeta function.

Presumably conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Mar 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005407 | OEIS-A173951

Positive integers with the property that if the base-3 representation is reversed the result is twice the original number.

(This conjecture is correct - see A223077. - _N. J. A. Sloane_, Mar 19 2013)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Mar 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005408 | OEIS-A173993

Sequence whose Hankel transform is the Somos (4) sequence.

Conjecture: (n+1)*a(n) +2*(-4*n-1)*a(n-1) +(19*n-5)*a(n-2) -36*a(n-3) +8*(-7*n+26)*a(n-4) +2*(34*n-143)*a(n-5) +24*(-n+5)*a(n-6)=0. - _R. J. Mathar_, Oct 10 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 04 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005409 | OEIS-A174013

Sequence whose Hankel transform is a (1,1) Somos-4 sequence.

Conjecture: (n+1)*a(n) +n*a(n-1) +12*(-n+1)*a(n-2) +3*(3*n-8)*a(n-3) +6*(6*n-23)*a(n-4) +11*(-n+2)*a(n-5) +(-49*n+253)*a(n-6) +20*(-n+6)*a(n-7)=0. - _R. J. Mathar_, Jan 12 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 05 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005410 | OEIS-A174015

A generalized Catalan number sequence.

Conjecture: (n+1)*a(n) -3*n*a(n-1) +3*(n-1)*a(n-2) +3*(n-4)*a(n-3) +2*(7-2*n)*a(n-4)=0. _R. J. Mathar_, Nov 13 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 05 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005411 | OEIS-A174016

Row sums of number triangle A174014.

Conjecture: (n+1)*a(n) - 6*n*a(n-1) + 12*(n-1)*a(n-2) - 12*a(n-3) + 8*(7-2*n)*a(n-4) = 0. - _R. J. Mathar_, Nov 13 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 05 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005412 | OEIS-A174031

The smallest integer k>0 such that the double-concatenation prime(n) // prime(n+1) // k is a prime number.

Conjecture: 1 appears infinitely often.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Mar 06 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005413 | OEIS-A174034

The smallest prime p such that the double-concatenation prime(n) // prime(n+1) // p is a prime number.

It is conjectured that a(n) = 3 for infinitely many n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Mar 06 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005414 | OEIS-A174056

Prime sums of three Mersenne primes. Primes in A174055.

No other terms < 10^1000. Conjecture: these are all the terms. - _Robert Israel_, Mar 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Mar 06 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005415 | OEIS-A174065

Convolved with its aerated variant = A000041.

The sequence diverges from A100853 after 16 terms; and is a conjectured Euler transform of A035263: (1, 0, 1, 1, 1, 0, 1, 0, 1, ...).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Mar 06 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005416 | OEIS-A174107

Expansion of (1/(1-x+x^2))c(x/(1-x+x^2)), c(x) the g.f. of A000108.

Conjecture: (n+1)*a(n) +2*(1-3n)*a(n-1) +7*(n-1)*a(n-2) +2*(5-3*n)*a(n-3) +(n-3)*a(n-4)=0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 07 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005417 | OEIS-A174114

Even central polygonal numbers (A193868) divided by 2.

Conjecture: the sequence terms are the exponents in the expansion of Sum_{n >= 1} q^n * (Product_{k >= 2*n} 1 - q^k) = q + q^2 + q^8 + q^11 + q^23 + q^28 + ... Cf. A266883. - _Peter Bala_, May 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Mar 08 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005418 | OEIS-A174161

a(1) = 2. Let k >= 1 be the minimal integer such that 2*k*a(n-1) - 1 has at least one prime divisor which is not already in the sequence. Then a(n) is the smallest such divisor.

Conjectures: 1) The sequence is a permutation of prime numbers; 2) k = k(n) runs all positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Mar 10 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005419 | OEIS-A174169

A generalized Chebyshev transform of the Motzkin numbers A001006.

Conjecture: (n+2)*a(n) -(2*n+1)*a(n-1) +3*(n-1)*a(n-2) +3*(5-2*n)*a(n-3) +9*(n-4)*a(n-4)=0. - _R. J. Mathar_, Sep 30 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 10 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005420 | OEIS-A174193

Expansion of the conjectured generating function for the number of "almost avoiding" permutations in the set Av(321)^{+1}.

Expansion of the conjectured generating function for the number of "almost avoiding" permutations in the set Av(321)^{+1}. Conjecture: -(n+2)*(404489*n-1866368)*a(n) +(3147815*n^2-8680082*n-14098754)*a(n-1) +2*(-2336439*n^2+652022*n+14514849)*a(n-2) +8*(-1044770*n^2+10585023*n-20370553)*a(n-3) +8*(642982*n-2004177)*(2*n-9)*a(n-4)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 26 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005421 | OEIS-A174195

Number of permutations of 1..n that almost avoid 231.

Conjecture: 2*(n+2)*(2013*n^2-10435*n+41550)*a(n) +(4026*n^3 -128025*n^2 +34955*n+59010)*a(n-1) -2 *(2*n-5)*(20130*n^2 -110855*n +93819) *a(n-2)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 26 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005422 | OEIS-A174216

a(1)=15; for n>1, a(n) = the smallest number k >a(n-1) such that 2*A174214(k)= 3*(k-1).

Conjecture. a(n+1)/a(n) tends to 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Mar 12 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005423 | OEIS-A174221

a(n) = n/2 if n is even, otherwise add to n the next three primes > n.

Related to the PrimeLatz conjecture, which states that if this map k -> a(k) is iterated, starting at any n >= 0, then the trajectory will eventually enter a loop.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 26 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005424 | OEIS-A174326

Exactly one of 3^n +- 2^n is prime.

If 3^n + 2^n is prime then n must be a power of 2, and 3^n + 2^n is a generalized Fermat prime. It is conjectured that 3^n + 2^n is prime only for n=1,2,4: see A082101. - _Robert Israel_, Mar 15 2017, edited May 18 2017.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Mar 15 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005425 | OEIS-A174327

Conjecturally, even numbers n such that every even number greater than n has more decompositions as the sum of two primes.

Conjecturally, even numbers n such that every even number greater than n has more decompositions as the sum of two primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 16 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005426 | OEIS-A174341

a(n) = Numerator of Bernoulli(n, 1) + 1/(n+1).

Conjecture: the numerator of (A164555(n)/(n+1) + A027642(n)/(n+1)^2) is a(n) and the denominator of this fraction is equal to 1 if and only if n+1 is prime or 1. Cf. A309132. - _Thomas Ordowski_, Jul 09 2019 The "if" part of the conjecture is true: see the theorems in A309132 and A326690. The values of the numerator when n+1 is prime are A327033. - _Jonathan Sondow_, Aug 15 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Mar 16 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005427 | OEIS-A174386

Smallest possible area for a perfect isosceles right triangled square of order n; or 0 if no such square exists.

These terms are only conjectures, but are thought highly likely to be correct for n<18.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Mar 18 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005428 | OEIS-A174420

Let A(n) = floor((3/2)^n), B(n)=3^n-2^n*A(n); then a(n)=2^n-A(n)-B(n)-2.

a(n) is conjectured to be >= 0 for all n >= 3. If this were proved it would settle Waring's problem (see A002804). It is known that a(n) >= 0 for 3 <= n <= 471600000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005429 | OEIS-A174470

G.F.: exp(Sum_{n>=1} A014578(n)*(3x)^n/n), where A014578 is the binary expansion of Thue constant.

Conjectured to consist entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 28 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005430 | OEIS-A174500

Continued fraction expansion for exp(Sum_{n>=1} 1/(n*A003500(n))), where A003500(n) = (2+sqrt(3))^n + (2-sqrt(3))^n.

a(2n-1) = 1, a(2n) = A003500(n) - 2, for n>=1 [conjecture]. The above conjectures are correct. The real number exp(Sum_{n>=1} 1/(n*A003500(n))) is equal to the infinite product F(x) := Product_{n >= 0} (1 - x^(4*n+3))/(1 - x^(4*n+1)) evaluated at x = 2 - sqrt(3). Ramanujan has given a continued fraction expansion for F(x). Using this we can find the simple continued fraction expansion of the numbers F(1/2*(N - sqrt(N^2 - 4))), N an integer greater than 3. The present case is when N = 4. See the Bala link for details.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005431 | OEIS-A174501

Continued fraction expansion for exp(Sum_{n>=1} 1/(n*A003499(n))), where A003499(n) = (3+sqrt(8))^n + (3-sqrt(8))^n.

a(2n-1) = 1, a(2n) = A003499(n) - 2, for n>=1 [conjecture]. The above conjectures are correct. See the Bala link for details. - _Peter Bala_, Jan 08 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005432 | OEIS-A174502

Continued fraction expansion for exp(Sum_{n>=1} 1/(n*A086903(n))), where A086903(n) = (4+sqrt(15))^n + (4-sqrt(15))^n.

a(2n-2) = 1, a(2n-1) = A086903(n) - 2, for n>=1 [conjecture]. The above conjectures are correct. See the Bala link for details. - _Peter Bala_, Jan 08 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005433 | OEIS-A174503

Continued fraction expansion for exp(Sum_{n>=1} 1/(n*A087799(n))), where A087799(n) = (5+sqrt(24))^n + (5-sqrt(24))^n.

a(2n-2) = 1, a(2n-1) = A087799(n) - 2, for n>=1 [conjecture]. The above conjectures are correct. See the Bala link for details. - _Peter Bala_, Jan 08 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005434 | OEIS-A174504

Continued fraction expansion for exp(Sum_{n>=1} 1/(n*A002203(n))), where A002203(n) = (1+sqrt(2))^n + (1-sqrt(2))^n.

a(3n-3) = 1, a(3n-2) = A002203(2n-1) - 1, a(3n-1) = A002203(2n) - 1, for n>=1 [conjecture]. The above conjectures are correct. The real number exp(Sum {n>=1} 1/(n*A002203(n))) is equal to the infinite product F(x) := product {n >= 0} (1 + x^(4*n+3))/(1 - x^(4*n+1)) evaluated at x = sqrt(2) - 1. Ramanujan has given a continued fraction expansion for the product F(x). Using this we can find the simple continued fraction expansion of the numbers F(1/2*(sqrt(N^2 + 4) - N)), N a positive integer. The present case is when N = 2. See the Bala link for details.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005435 | OEIS-A174505

Continued fraction expansion for exp(Sum_{n>=1} 1/(n*Lucas(n))), where Lucas(n) = A000032(n) = ((1+sqrt(5))/2)^n + ((1-sqrt(5))/2)^n.

a(3n-2) = 1, a(3n-1) = A000032(2n+1) - 1, a(3n) = A000032(2n+2) - 1, for n>=1 [conjecture]. The above conjectures are correct. The real number exp(Sum {n>=1} 1/(n*Lucas(n))) is equal to the infinite product F(x) := product {n >= 0} (1 + x^(4*n+3))/(1 - x^(4*n+1)) evaluated at x = 1/2*(sqrt(5) - 1). Ramanujan has given a continued fraction expansion for the product F(x). Using this we can find the simple continued fraction expansion of the numbers F(1/2*(sqrt(N^2 + 4) - N)), N a positive integer. The present case is when N = 1. See the Bala link for details.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005436 | OEIS-A174506

Continued fraction expansion for exp(Sum_{n>=1} 1/(n*A014448(n))), where A014448(n) = (2+sqrt(5))^n + (2-sqrt(5))^n.

a(3n-3) = 1, a(3n-2) = A014448(2n-1) - 1, a(3n-1) = A014448(2n) - 1, for n>=1 [conjecture]. The above conjectures are correct. The real number exp(Sum {n>=1} 1/(n*A014448(n))) is equal to the infinite product F(x) := product {n >= 0} (1 + x^(4*n+3))/(1 - x^(4*n+1)) evaluated at x = sqrt(5) - 2. Ramanujan has given a continued fraction expansion for the product F(x). Using this we can find the simple continued fraction expansion of the numbers F(1/2*(sqrt(N^2 + 4) - N)), N a positive integer. The present case is when N = 4. See the Bala link for details.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005437 | OEIS-A174507

Continued fraction expansion for exp(Sum_{n>=1} 1/(n*A085447(n))), where A085447(n) = (3+sqrt(10))^n + (3-sqrt(10))^n.

a(3n-3) = 1, a(3n-2) = A085447(2n-1) - 1, a(3n-1) = A085447(2n) - 1, for n>=1 [conjecture]. The above conjectures are correct. The real number exp(Sum {n>=1} 1/(n*A085447(n))) is equal to the infinite product F(x) := product {n >= 0} (1 + x^(4*n+3))/(1 - x^(4*n+1)) evaluated at x = sqrt(10) - 3. Ramanujan has given a continued fraction expansion for the product F(x). Using this we can find the simple continued fraction expansion of the numbers F(1/2*(sqrt(N^2 + 4) - N)), N a positive integer. The present case is when N = 6. See the Bala link for details.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005438 | OEIS-A174508

Continued fraction expansion for exp(Sum_{n>=1} 1/(n*A086594(n))), where A086594(n) = (4+sqrt(17))^n + (4-sqrt(17))^n.

a(3n-3) = 1, a(3n-2) = A086594(2n-1) - 1, a(3n-1) = A086594(2n) - 1, for n>=1 [conjecture]. The above conjectures are correct. The real number exp(Sum {n>=1} 1/(n*A086594(n))) is equal to the infinite product F(x) := product {n >= 0} (1 + x^(4*n+3))/(1 - x^(4*n+1)) evaluated at x = sqrt(17) - 4. Ramanujan has given a continued fraction expansion for the product F(x). Using this we can find the simple continued fraction expansion of the numbers F(1/2*(sqrt(N^2 + 4) - N)), N a positive integer. The present case is when N = 8. See the Bala link for details.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005439 | OEIS-A174509

Continued fraction expansion for exp(Sum_{n>=1} 1/(n*A086927(n))), where A086927(n) = (5+sqrt(26))^n + (5-sqrt(26))^n.

a(3n-3) = 1, a(3n-2) = A086927(2n-1) - 1, a(3n-1) = A086927(2n) - 1, for n>=1 [conjecture]. The above conjectures are correct. The real number exp(Sum {n>=1} 1/(n*A086927(n))) is equal to the infinite product F(x) := product {n >= 0} (1 + x^(4*n+3))/(1 - x^(4*n+1)) evaluated at x = sqrt(26) - 5. Ramanujan has given a continued fraction expansion for the product F(x). Using this we can find the simple continued fraction expansion of the numbers F(1/2*(sqrt(N^2 + 4) - N)), N a positive integer. The present case is when N = 10. See the Bala link for details.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005440 | OEIS-A174510

Continued fraction expansion for exp(Sum_{n>=1} 1/(n*A080040(n))), where A080040(n) = (1+sqrt(3))^n + (1-sqrt(3))^n.

a(3n-3) = 1, a(3n-2) = A080040(2n-1)/2^(n-1) - 1, a(3n-1) = A080040(2n)/2^n - 1, for n>=1 [conjecture].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005441 | OEIS-A174525

Bases N in which ab and ba are different squares, for some a and b.

Conjecture: all odd squares (A016754) except 1 are terms of the sequence. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael R Peake_, Mar 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005442 | OEIS-A174531

Coefficients of polynomials described below.

Conjectures: (1) all terms of the sequence are integers; (2) P_n(0) = (floor((n-1)/2))!*4^(floor((n-1)/2)); (3) if n is even, then the coefficients of P_n do not exceed the corresponding coefficients of P_(n-1) and the equality holds only for the last ones; (4) all coefficients of P_n, except of the last one, are multiples of n iff n is prime. More conjectures: For even x, P[x, 1] = (2^x - 1) (x/2)! / (x+1), and for odd x, P[x, 1] = (2^x - 1) ((x-1) / 2)!. - _Peter J. C. Moses_, Sep 28 2011 The sequence of polynomials R's for k >= 0 begins: 0, 1, n+4, n^2 + 11*n + 32, n^3 + 21*n^2 + 152*n + 384, n^4 + 34*n^3 + 443*n^2 + 2642*n + 6144, ... This formula proves the conjecture of V. Shevelev for P_n(0) and P. Moses for P_n(1). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Mar 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005443 | OEIS-A174583

a(k) is the least n such that the concatenation (n - k)"n is a prime number, for k >= 0.

Conjecture: a(k) = a(k+1) for an infinite number of k's.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Mar 23 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005444 | OEIS-A174604

a(1) = 1, then a(n) is the least integer > a(n-1) such that n is the maximum element in the continued fraction for 1/a(1) + 1/a(2) + ... + 1/a(n).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Mar 23 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005445 | OEIS-A174655

Partial sums of A049486.

Conjecture: a(n) = (3*(-9+(-1)^n)+34*n-12*n^2+8*n^3)/12 for n>1. G.f.: x*(x^5-x^4+3*x^3+2*x^2+2*x+1) / ((x-1)^4*(x+1)). - _Colin Barker_, May 02 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Mar 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005446 | OEIS-A174783

Expansion of (1+2x-sqrt(1-4x^2))/(2(1-x^2)*sqrt(1-4x^2)).

Conjecture: n*a(n) -2*a(n-1) +(8-5*n)*a(n-2) +2*a(n-3) +4*(n-2)*a(n-4)=0. - _R. J. Mathar_, Nov 13 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005447 | OEIS-A174846

E.g.f.: AGM(1, exp(4x)), where AGM(x, y) is the arithmetic-geometric mean of Gauss.

Conjecture: limit |a(n)/n!|^(-1/n) = r exists and is finite with r<0.8...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005448 | OEIS-A174905

Numbers with no pair (d,e) of divisors such that d < e < 2*d.

Conjecture: Numbers m such that A001227(m) equals the number of 2-dense sublists of divisors of m. - _Omar E. Pol_, Feb 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 01 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005449 | OEIS-A174926

Smallest prime which has a decimal representation which shows n^2 embedded in otherwise only decimal square digits 0, 1, 4 and 9.

It is conjectured that sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ulrich Krug_, Apr 02 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005450 | OEIS-A174968

Decimal expansion of (1 + sqrt(2))/2.

Conjecture: Equals Sum_{k>=0} A131126(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Apr 02 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005451 | OEIS-A174979

Smallest prime p which is a concatenation of n^3 and the cubic digits 0, 1, 8.

There are three decimal digits which are cubes: 0 = 0^3, 1 = 1^3, 8 = 2^3. It is conjectured that sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), Apr 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005452 | OEIS-A174989

Partial sums of A003602.

I conjecture that infinitely many terms are prime. For n<=10^5, exactly 5115 terms are prime. For n<=10^7, there are 352704 prime terms. The largest prime for n<10^10 is at n=9999999983, a(n)=16666666618226308891. Below 10^100, n=(10^100)-345. Below 10^500, n=(10^500)-2414. - _Griffin N. Macris_, May 04 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Apr 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005453 | OEIS-A175019

a(0)=a(1)=1; a(n) = a(n-a(n-2)) + 2.

Conjectures from _Chai Wah Wu_, Jun 12 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Apr 04 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005454 | OEIS-A175033

Numbers n such that (ceiling(sqrt(n*n/2)))^2 - n*n/2 = 17/2.

Conjecture: a(n) = 6*a(n-2) - a(n-4). - _Charles R Greathouse IV_, Apr 30 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Nov 09 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005455 | OEIS-A175046

Write n in binary, then increase each run of 0's by one 0, and increase each run of 1's by one 1. a(n) is the decimal equivalent of the result.

a(n) <= (9*n^2 + 12*n)/5, with equality iff n = (2/3)*(4^k-1) = A182512(k) for some k, i.e., n = 10101...10 in binary. - Conjectured by _N. J. A. Sloane_, Sep 09 2018, proved by _M. F. Hasler_, Sep 12 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, Dec 02 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005456 | OEIS-A175073

Primes q with result 1 under iterations of {r mod (max prime p < r)} starting at r = q.

Conjecture: under this iteration procedure, all primes eventually will yield either a 2 or a 1. If a 2 results, all subsequent terms are zeros; if a 1 results, all subsequent terms are -1s. The conjecture is true for the first 2 million primes. - _Harvey P. Dale_, Jan 17 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jan 23 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005457 | OEIS-A175079

The smallest natural numbers m with first occurrence 0, 1, 2, 3, ... for number of steps of iterations of {r mod (max prime p < r)} needed to reach 1 or 2 starting at r = m.

I offer a prize of 100 liters of Pilsner Urquell to the discoverer of a(5). Conjecture: a(n) is not equal A135543(n) + 1 for all n >= 1. See A175071 (natural numbers m with result 1) and A175072 (natural numbers m with result 2). See A175077 (results 1 or 2 under iterations) and A175078 (number of steps of iterations). a(n) = A135543(n) + 1 for n >= 1, i.e., the conjecture in the Comments is false. This follows from the result that A175078(n) = A121561(n-1) for n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jan 23 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005458 | OEIS-A175130

Indices of Fibonacci numbers that are not cubefree.

Conjecture: all terms are multiples of 6 or 125. - _Harvey P. Dale_, Apr 28 2020 The conjecture is false. The counterexamples are 392, 784, 1183, 1210, ... - _Amiram Eldar_, Oct 16 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Feb 16 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005459 | OEIS-A175177

Conjectured number of numbers for which the iteration x -> phi(x) + 1 terminates at prime(n). Cardinality of rooted tree T_p (where p is n-th prime) in Karpenko's book.

Conjectured number of numbers for which the iteration x -> phi(x) + 1 terminates at prime(n). Cardinality of rooted tree T_p (where p is n-th prime) in Karpenko's book.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Mar 01 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005460 | OEIS-A175243

Array read by antidiagonals: total number of spanning trees R_n(m) of the complete prism K_m X C_n.

Each column of the array is a linear divisibility sequence. Conjecturally, the k-th column satisfies a linear recurrence of order 4*k - 2. - _Peter Bala_, May 04 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Mar 13 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005461 | OEIS-A175254

a(n) = Sum_{k<=n} A000203(k)*(n-k+1), where A000203(m) is the sum of divisors of m.

Conjecture: a(n) = Sum_{k=0..n} A006218(n-k). - _R. J. Mathar_, Oct 17 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Mar 14 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005462 | OEIS-A175257

a(n) is the smallest prime p such that 2^(p-1) == 1 (mod a(1)*...*a(n-1)*p).

Conjecture: a(n) is the smallest integer k > 1 such that 2^(k-1) == 1 (mod a(0)*...*a(n-1)*k), with a(0) = 1. - _Thomas Ordowski_, Mar 13 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Manuel Valdivia_, Mar 15 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005463 | OEIS-A175287

Partial sums of ceiling(n^2/4).

Conjecture: a(n) is the number of perimeter-magic (hollow) triangles of order 3 with magic sum n+2. Order 3 means each of the 3 edges has 3 elements >=1; the triangle has 6 elements. The elements do not need to be distinct, and triangles obtained by rotations are counted only once. The triangle (read ccw) for magic sum 3 has elements 1 1 1 1 1 1. The 2 triangles with magic sum 4 are 1 1 2 1 1 2 and 1 2 1 2 1 2. - _R. J. Mathar_, Mar 08 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mircea Merca_, Dec 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005464 | OEIS-A175368

Number of integer 4-tuples (x,y,z,u) satisfying |x|^3 + |y|^3 + |z|^3 + |u|^3 = n, -n <= x,y,z,u <= n.

Conjectured g.f.: (1 + 2*Sum_{j>=1} x^(j^3))^4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Apr 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005465 | OEIS-A175385

a(n) = numerator of Sum_{i=1..n} binomial(2n-i-1,i-1)/i.

We conjecture that Sum_{i=1..n} ((1/i)*C(2n-i-1,i-1)) is not an integer for n>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, _Zak Seidov_, Apr 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005466 | OEIS-A175403

a(n) is the smallest number m requiring n iterations {((((D_1^D_2)^D_3)^D_4)^...)^D_k to reach a one-digit number starting at r = n, where D_k is the k-th digit D of the number r and k is the digit number of the number r in the decimal expansion of r (A055642)}.

Conjecture: sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, May 01 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005467 | OEIS-A175406

The greatest integer k such that (1+1/n)^k <= 2.

a(n) = n log 2 + O(1). Conjecture: a(n) = floor((n + 1/2) log 2). - _Charles R Greathouse IV_, Apr 03 2012 The conjecture by Greathouse fails at n=1121626023352383: a(n)=777451915729367 while floor((n + 1/2) log 2)=777451915729368. - _Per W. Alexandersson_, Oct 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, May 01 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005468 | OEIS-A175424

a(n) is the number of steps of iterations of {((((D_k^D_(k-1))^D_(k-2))^...)^D_1, where D_k is the k-th digit D of number r and k is the number of digits of number r in decimal expansion of r (A055642)} needed to reach a single-digit number starting at r = n, or a(n) = -1 if a single-digit number is never reached.

Conjecture: max(a(n)) = 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, May 09 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005469 | OEIS-A175425

The smallest number m which needs n iterations of the map x->A175420(x) to reach a single-digit number.

Conjecture: sequence a(n) is finite for n = natural number, a(n) = 25 for n = infinity. See A175424.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, May 09 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005470 | OEIS-A175498

a(1)=1. a(n) = the smallest positive integer not occurring earlier such that a(n)-a(n-1) doesn't equal a(k)-a(k-1) for any k with 2 <= k <= n-1.

Conjecture: the lexicographically earliest permutation of {1,2,...n} for which differences of adjacent numbers are all distinct (cf. A131529) has, for n-->infinity, this sequence as its prefix. - _Joerg Arndt_, May 27 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leroy Quet_, May 31 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005471 | OEIS-A175527

Digit sum of 13^n.

a(n) ~ 4.5*log_10(13)*n ~ 5.0127*n (conjectured). - _M. F. Hasler_, May 18 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005472 | OEIS-A175567

a(n) = (n!)^2 modulo n*(n+1)/2.

The conjecture above is true for every even power 2k with k >= 1. Proof: if n+1 is not an odd prime, n(n+1)/2 divides n! (see A069268). If n+1 = p is an odd prime, write n(n+1)/2 = h*p with h = n/2. Then (n!)^(2k) == 0 (mod h), and (n!)^(2k) == 1 (mod p) by Wilson's theorem. Because 2h == -1 (mod p), x = h*(p-2) = n(n-1)/2 satisfies x == 0 (mod h) and x == -2h == 1 (mod p), and 0 <= x < n(n+1)/2, so by the Chinese remainder theorem the residue is n(n-1)/2. - _Scott M. Foltz_, Sep 24 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jul 12 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005473 | OEIS-A175573

Decimal expansion of Pi^(1/4)/Gamma(3/4).

Conjecture: Equals Sum_{k>=0} A000122(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jul 15 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005474 | OEIS-A175574

Decimal expansion of sqrt(Pi) / (Gamma(3/4))^2.

Conjecture: Equals Sum_{k>=0} A004018(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jul 15 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005475 | OEIS-A175582

a(n) = sigma(n-th Zumkeller number)/2.

Conjecture: Any 4 consecutive terms include at least one Zumkeller number (verified for the first 10^5 terms). - _Ivan N. Ianakiev_, Oct 17 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | Vladislav-Stepan Malakhovsky and _Juri-Stepan Gerasimov_, Jul 15 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005476 | OEIS-A175625

Numbers k such that gcd(k, 6) = 1, 2^(k-1) == 1 (mod k), and 2^(k-3) == 1 (mod (k-1)/2).

Conjecture: The prime values belong to A039787. - _Bill McEachen_, Dec 27 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alzhekeyev Ascar M_, Jul 28 2010, Jul 30 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005477 | OEIS-A175669

Triangle of numerators of coefficients of the polynomial Q^(2)_m(n) defined by the recursion Q^(2)_0(n)=1; for m>=1, Q^(2)_m(n) = Sum_{i=1..n} i^2*Q^(2)_(m-1)(i). For m>=0, the denominator for all 3*m+1 terms of the m-th row is A202367(m+1).

Sequence {Q^(r)_m(x)} is defined by the recursion: Q^(r)_0(x)=1; for m>=1 and integer x=n, Q^(r)_m(n)=sum{i=1,...,n}i^rQ^(r)_(m-1)(i). By the induction, we see that polynomial Q^(r)_m(x) has degree (r+1)*m. Note that Q^(0)_m(n) is C(n+m-1,m), Q^(1)_m(n)=S(n+m,n), where S(k,l) are Stirling numbers of the second kind. Thus Q^(r)_m(x) is an r-generalization of binomial coefficients and Stirling numbers of the second kind. Moreover, for every r, LCM of denominators of the coefficients of Q^(r)_m(x) generate sequences of factorial type which possess important arithmetic properties. For r=0, it is n!, for r=1, it is A053657, for r=2,3,4 we obtain A202367, A202368, A202369. Denote the general term of the sequence corresponding to a given r by n!^(r) and, for 0<=m<=n, denote C^(r)(n,m)=n!^(r)/(m!^(r)*(n-m)!^(r). Then, for the "r-Pascal triangle", we have the following conjectural regularity: if a prime p==1 mod r, then the ((p-1)/r)-th row contains two 1's and numbers multiple of p. Cf. triangles A202917, A202941.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Dec 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005478 | OEIS-A175705

Convolution square of A001157 (the sum of squared divisors).

Conjecture: Sum_{k=1..n} a(k) ~ zeta(3)^2 * n^6 / 180. - _Vaclav Kotesovec_, Aug 20 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Aug 10 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005479 | OEIS-A175764

Number of iterations of the mapping k->f(k) to reach one of 2, 5, or 29, starting with k=n, and with f(k)=(k^2+4)/d, where d is the next-to-largest divisor of k^2+4, or -1 if the sequence never reaches one of the required values.

It appears that a(n) is 1 whenever n>29 and n mod 10 is one of {0,1,2,4,6,8,9}. This has been verified to n=5000. Also, it appears that a(n) is 9 whenever n mod 130 is one of {3,23,55,75,107,127}. This has also been verified to n=5000. Are these conjectures easy to prove?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Aug 30 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005480 | OEIS-A175784

Numerators of k/(10+k)+1 for k = 2*n-1.

Conjecture: a(n) = 2*a(n-5) - a(n-10) = 4*A060791(n+2) with g.f. -4*x*(-3 - 4*x - x^2 - 6*x^3 - 7*x^4 - 2*x^5 - x^6 + x^8 + 2*x^9) / ((x-1)^2*(x^4 + x^3 + x^2 + x + 1)^2). [_R. J. Mathar_, Dec 07 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven J. Forsberg_, Dec 04 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005481 | OEIS-A175799

Number of real zeros of the polynomial whose coefficients are the decimal expansion of Pi truncated to n places (A011545).

Conjecture: P(n,x) has nonzero discriminant for all n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Dec 04 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005482 | OEIS-A175809

a(n) is the number of shortest common superstrings of the binary representations of all natural numbers from 1 to n.

a(2^n) = A016031(n) (if conjectured A175808(2^n) = 2^n is true). a(2^n-2) = a(2^n-1) = a(2^n) for n > 1. Conjectured. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Reshetnikov_, Sep 08 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005483 | OEIS-A175815

Numbers k such that the first k decimal digits of Euler's constant gamma contain equal numbers of even and odd decimal digits.

Conjecturally infinite. - _Charles R Greathouse IV_, May 15 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Sep 09 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005484 | OEIS-A175931

Numbers n such that A129363(n-1) = A129363(n) = A129363(n+1).

Conjecture: Sequence is finite and complete with last term 2103.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Oct 23 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005485 | OEIS-A175938

Repeat: Take next nonprime, next prime, next nonprime, then reverse the order.

Conjecture: The permutation has exactly 12 fixed points: 2, 6, 9, 12, 15, 18, 21, 24, 29, 40, 46, 49. The conjecture is true, roughly because the primes are packed far too closely (one every three terms, after all) and as a result the nonprimes are too spread out.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Dec 06 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005486 | OEIS-A176006

The number of branching configurations of RNA (see Sankoff, 1985) with n or fewer hairpins.

Conjecture : n*a(n) +(9-7*n)*a(n-1) +(7*n-12)*a(n-2) +(3-n)*a(n-3)=0. - _R. J. Mathar_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lee A. Newberg_, Apr 05 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005487 | OEIS-A176014

Decimal expansion of (3+sqrt(21))/6.

Also greatest eigenvalue of the 6 X 6 matrix [[3 0 0 3 0 0][0 0 0 0 1 0][0 3 0 0 3 0][0 0 0 0 1 0][0 0 3 0 0 3][0 0 0 0 1 0]]/3. It is conjectured that this is lim_{k->infinity} A005186(k+1)/A005186(k), i.e., the asymptotic growth rate of the number of numbers with the same total stopping time in the Collatz iteration. - _Hugo Pfoertner_, Sep 28 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_, Apr 06 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005488 | OEIS-A176042

For odd n, this is the number of partitions of n*(n-1)/2 with all part sizes between 3 and n, inclusive. For even n, this is the number of partitions of n*(n-2)/2 with all part sizes between 3 and n, inclusive.

Number of different decompositions of complete graphs on n vertices into cycles (for odd n) or into cycles and a 1-factor (for n even) according to a conjecture of Brian Alspach's. This is verified for n <= 14.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Christopher Maitland (c3053540(AT)uon.edu.au), Apr 07 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005489 | OEIS-A176117

Primes p such that T(p) is prime in Juricevic conjecture on classification of Lehmer triples.

Primes p such that T(p) is prime in Juricevic conjecture on classification of Lehmer triples.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Apr 08 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005490 | OEIS-A176187

Least period of sequence {f^(m)(n)},m=1,2,3,..., where f^(m) is the m-th iteration of A002326 (f^(1)=f), after possibly dropping some finite number of iterations.

Conjecture: For every n>=0 the sequence {f^(m)(n)} is eventually periodic. Further calculations may show if there exist terms different from 1 and 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 11 2010, Apr 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005491 | OEIS-A176206

Irregular triangle T(n,k) (n >= 1, k >= 1) read by rows: row n has length A000070(n-1) and every column k gives the positive integers.

Conjecture: all divisors of all terms in row n are also all parts of all partitions of n. For more information see the example and A336811 which contains the most elementary conjecture about the correspondence divisors/partitions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alford Arnold_, Apr 11 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005492 | OEIS-A176268

Primes of a Generalized Cunningham chain of length 9 by the function f(p) = 2 * p + 13.

It is conjectured that arbitrarily long such chains exist

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Apr 13 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005493 | OEIS-A176285

a(1) = 1, and then 4*(2*n + 1)^2*a(n+1) + n^2*a(n) = (205*n^2 + 160*n + 32)*binomial(2*n-1, n)^3 (n = 1, 2, 3, ...).

On Apr 04 2010, _Zhi-Wei Sun_ introduced this sequence and conjectured that each term a(n) is a positive integer. He also guessed that a(n) is odd if and only if n is a power of two. It is easy to see that 8*n^2*binomial(2*n,n)^2*a(n) equals Sum_{k=0..n-1} (205*k^2 + 160*k + 32)*(-1)^(n-1-k)*binomial(2*k,k)^5. Sun also conjectured that for any prime p > 5 we have Sum_{k=0..p-1} (205*k^2 + 160*k + 32)(-1)^k*binomial(2*k,k)^5 == 32*p^2 + 64*p^3*Sum_{k=1..p-1} 1/k (mod p^7) and Sum_{k=0..(p-1)/2} (205*k^2 + 160*k + 32)(-1)^k*binomial(2*k,k)^5 == 32*p^2 + 896/3*p^5*B_{p-3} (mod p^6), where B_0, B_1, B_2, ... are Bernoulli numbers. It is also remarkable that Sum_{n>0} (-1)^n(205*n^2 - 160*n + 32)/(n^5*binomial(2*n,n)^5) = -2*zeta(3) as proved by T. Amdeberhan and D. Zeilberger via the WZ method. Conjecture: the supercongruences a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) hold for all primes p >= 5 and all positive integers n and r. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 14 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005494 | OEIS-A176328

Numerators of the rational sequence with e.g.f. (x/2)*(exp(-x) + 1)/(exp(x) - 1).

Conjecture: a(2*n+1) = -2*n-1. The conjecture is true (follows immediately from Luschny's formula. See Fried link). - _Sela Fried_, Dec 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Apr 15 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005495 | OEIS-A176332

Row sums of triangle A176331.

Conjecture: 6*n*a(n) + 2*(-17*n+10)*a(n-1) + (47*n-60)*a(n-2) + 10*(-3*n+5)*a(n-3) + 4*(2*n-5)*a(n-4) = 0. - _R. J. Mathar_, Nov 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 15 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005496 | OEIS-A176335

Central coefficients T(2n,n) of number triangle A176331.

Conjecture: 224*n^2*(n-1)*a(n) - 48*(n-1)*(65*n^2 - 36*n - 13)*a(n-1) + 4*(-1839*n^3 + 11081*n^2 - 21932*n + 14280)*a(n-2) + 12*(-81*n^3 + 326*n^2 - 591*n + 562)*a(n-3) - (n-3)*(1853*n^2 - 7403*n + 7140)*a(n-4) - 12*(n-4)*(2*n-7)^2*a(n-5) = 0. - _R. J. Mathar_, Feb 10 2015 Conjecture (assuming an offset of 1): the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) holds for all primes p >= 5 and all positive integers n and r [added Nov 29 2024: proved by Coster. See Theorem 4]. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 15 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005497 | OEIS-A176494

Least m >= 1 for which |2^m - prime(n)| is prime.

Conjecture. For every odd prime p, the sequence {|2^n - p|} contains at least one prime. The record values of the sequence appear at n = 2, 10, 31, 68, 341, ... and are 3, 4, 47, 791, ... Note that up to now the value a(341) is not known. _Charles R Greathouse IV_ calculated the following two values: a(815)=16464, a(591)=58091 and noted that a(341) is much larger [private communication, May 27 2010]. - _Vladimir Shevelev_, May 29 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 19 2010, Aug 15 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005498 | OEIS-A176542

Numbers k such that there are only a finite nonzero number of sets of k consecutive triangular numbers that sum to a square.

Conjectures from _Colin Barker_, Sep 24 2015: (Start) These conjectures are true (see Fried link). - _Sela Fried_, Jan 02 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Weimholt_, Apr 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005499 | OEIS-A176545

Numbers k>0 such that 2*k^2+14*k+5 is prime.

Conjecture: If p = 2*k^2 + 14*k + 5 then 2^p - 1 is not prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Apr 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005500 | OEIS-A176547

Numbers n such that 2*n^2 + 6*n + 1 is prime.

Conjecture: If p=2*n^2+6*n+1 is prime then 2^p-1 is not prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Apr 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005501 | OEIS-A176553

Numbers m such that concatenations of divisors of m are noncomposites.

Conjecture: this sequence is a subsequence of A003136 (Loeschian numbers). - _Davide Rotondo_, Jan 02 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Apr 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005502 | OEIS-A176605

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=1, k=0 and l=1.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +(9*n-13)*a(n-2) +8*(-n+3)*a(n-3) +4*(n-4)*a(n-4)=0. - _R. J. Mathar_, Feb 29 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005503 | OEIS-A176606

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=3, k=0 and l=1.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +(n+3)*a(n-2) +8*(n-3)*a(n-3) +4*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Feb 29 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005504 | OEIS-A176607

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=4, k=0 and l=1.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +(-3*n+11)*a(n-2) +16*(n-3)*a(n-3) +8*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Feb 29 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005505 | OEIS-A176609

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=5, k=0 and l=1.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +(-7*n+19)*a(n-2) +24*(n-3)*a(n-3) +12*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Mar 01 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005506 | OEIS-A176610

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=0, k=1 and l=1.

Conjecture: (n+1)*a(n) +(2-7n)*a(n-1) +(19n-29)*a(n-2) +(110-37n)*a(n-3) +36*(n-4)*a(n-4) +12*(5-n)*a(n-5)=0. - _R. J. Mathar_, Nov 17 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005507 | OEIS-A176611

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=1, k=1 and l=1.

Conjecture: (n+1)*a(n) +(-7*n+2)*a(n-1) +3*(5*n-7)*a(n-2) +(-25*n+74)*a(n-3) +24*(n-4)*a(n-4) +8*(-n+5)*a(n-5)=0. - _R. J. Mathar_, Mar 01 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005508 | OEIS-A176612

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=2, k=1 and l=1.

Conjecture: (n+1)*a(n) +(-7*n+2)*a(n-1) +(11*n-13)*a(n-2) +(-13*n+38)*a(n-3) +12*(n-4)*a(n-4) +4*(-n+5)*a(n-5)=0. - _R. J. Mathar_, Feb 29 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005509 | OEIS-A176614

Conjectured least number M such that PrimePi(m)*log(m)/m lies between 1+(1/n) and 1-(1/n) for all m >= M.

Conjectured least number M such that PrimePi(m)*log(m)/m lies between 1+(1/n) and 1-(1/n) for all m >= M.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael B. Porter_, Apr 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005510 | OEIS-A176675

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=0, k=0 and l=-1.

Conjecture: +(n+1)*a(n) +2*(-3*n+1)*a(n-1) +(13*n-21)*a(n-2) +4*(-2*n+5)*a(n-3)=0. - _R. J. Mathar_, Feb 29 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 23 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005511 | OEIS-A176677

Sequence defined by the recurrence formula a(n+1) = Sum_{p=0..n} (a(p)*a(n-p)+k) + l for n>=1, with here a(0)=1, a(1)=1, k=0 and l=-1.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +(9*n-13)*a(n-2) -4*a(n-3) +4*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Mar 01 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 23 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005512 | OEIS-A176678

a(n+1) = Sum_{p=0..n} a(p)*a(n-p)+k+l for n>=1, with here a(0)=1, a(1)=2, k=0 and l=-1.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +5*(n-1)*a(n-2) +4*(2*n-7)*a(n-3) +8*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Feb 18 2016. Proved by _Adarsh Ramamurthy_, Jul 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 23 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005513 | OEIS-A176716

a(1) = 1; a(2*n) = prime(n)*a(n), a(2*n+1) = prime(n)*a(n) + a(n+1), where prime(n) is the n-th prime.

Conjecture: for n > 1, 4 divides a(n) iff n == 0 (mod 3), otherwise A007814(a(n)) = 1. Conjecture is true (proof by mathematical induction). - _Robert Israel_, Feb 25 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Apr 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005514 | OEIS-A176752

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=0, k=0 and l=-2.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +(13*n-21)*a(n-2) +2*(-2*n+3)*a(n-3) +4*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Feb 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005515 | OEIS-A176753

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=1, k=0 and l=-2.

Conjecture: (n+1)*a(n) +2*(1-3*n)*a(n-1) +(9*n-13)*a(n-2) +2*(2*n-9)*a(n-3) +8*(4-n)*a(n-4)=0. - _R. J. Mathar_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005516 | OEIS-A176754

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=2, k=0 and l=-2.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +5*(n-1)*a(n-2) +6*(2*n-7)*a(n-3) +12*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Feb 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005517 | OEIS-A176755

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=3, k=0 and l=-2.

Conjecture: (n+1)*a(n) +2*(1-3*n)*a(n-1) +(n+3)*a(n-2) +2*(10*n-33)*a(n-3) +16*(4-n)*a(n-4) =0. - _R. J. Mathar_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005518 | OEIS-A176756

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=4, k=0 and l=-2.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +(-3*n+11)*a(n-2) +2*(14*n-45)*a(n-3) +20*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Feb 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005519 | OEIS-A176757

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=5, k=0 and l=-2.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +(-7*n+19)*a(n-2) +6*(6*n-19)*a(n-3) +24*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Feb 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005520 | OEIS-A176781

Smallest prime prime(i) such that concatenation 2//0_(n)//prime(i) is prime.

It is conjectured that prime(i) = 3 occurs infinitely often: at n= 0, 2, 4, 5, 6, 11, 15, 16, 21, 23, 34, 114, 119,... (A101951).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Apr 26 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005521 | OEIS-A176806

Consider asymmetric 1-D random walk with set of possible jumps {-1,+1,+2}. Sequence gives number of paths of length n ending at origin.

From this special Lagrange series derivation and the proof that the g.f. g from above coincides with G, the conjecture, given by _Joerg Arndt_ as a comment above, has been proved. This uses [t^n]phi(t)^n = (1/n!)*(d/dt)^n phi(t)^n, evaluated at t = 0, which appears in the considered Lagrange series.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sergey Perepechko_, Apr 26 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005522 | OEIS-A176826

a(n+1) = Sum_{p=0..n} (a(p)*a(n-p)+k)+l for n>=1, where a(0)=1, a(1)=1, k=1 and l=-1.

Conjecture: (n+1)*a(n) +(-7*n+2)*a(n-1) +3*(5*n-7)*a(n-2) +(-17*n+46)*a(n-3) +4*(2*n-7)*a(n-4)=0. - _R. J. Mathar_, Feb 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005523 | OEIS-A176829

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=3, k=1 and l=-1.

Conjecture: (n+1)*a(n) +(-7*n+2)*a(n-1) +(7*n-5)*a(n-2) +(7*n-26)*a(n-3) +4*(-4*n+17)*a(n-4) +8*(n-5)*a(n-5)=0. - _R. J. Mathar_, Feb 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005524 | OEIS-A176830

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=4, k=1 and l=-1.

Conjecture: (n+1)*a(n) +(-7*n+2)*a(n-1) +3*(n+1)*a(n-2) +(19*n-62)*a(n-3) +4*(-7*n+29)*a(n-4) +12*(n-5)*a(n-5)=0. - _R. J. Mathar_, Feb 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005525 | OEIS-A176832

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=5, k=1 and l=-1.

Conjecture: (n+1)*a(n) +(-7*n+2)*a(n-1) +(-n+11)*a(n-2) +(31*n-98)*a(n-3) +4*(-10*n+41)*a(n-4) +16*(n-5)*a(n-5)=0. - _R. J. Mathar_, Feb 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005526 | OEIS-A176848

Number of compositions of n into floor(j/3) kinds of j's for all j>=1.

Conjectural g.f.: (x-1)^2*(x^2+x+1) / (x^4-2*x^3-x+1). - _Colin Barker_, May 15 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Jul 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005527 | OEIS-A176854

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=0, k=-1 and l=0.

Conjecture: (n+1)*a(n) +(-7*n+2)*a(n-1) +(19*n-29)*a(n-2) +(-17*n+40)*a(n-3) +2*(2*n-7)*a(n-4)=0. - _R. J. Mathar_, Mar 01 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005528 | OEIS-A176855

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=1, k=-1 and l=0.

Conjecture: (n+1)*a(n) +(-7*n+2)*a(n-1) +3*(5*n-7)*a(n-2) +(-5*n+4)*a(n-3) +2*(-4*n+17)*a(n-4) +4*(n-5)*a(n-5)=0. - _R. J. Mathar_, Mar 01 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005529 | OEIS-A176856

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=2, k=-1 and l=0.

Conjecture: (n+1)*a(n) +(-7*n+2)*a(n-1) +(11*n-13)*a(n-2) +(7*n-32)*a(n-3) +2*(-10*n+41)*a(n-4) +8*(n-5)*a(n-5)=0. - _R. J. Mathar_, Mar 01 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005530 | OEIS-A176857

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=3, k=-1 and l=0.

Conjecture: (n+1)*a(n) +(-7*n+2)*a(n-1) +(7*n-5)*a(n-2) +(19*n-68)*a(n-3) +2*(-16*n+65)*a(n-4) +12*(n-5)*a(n-5)=0. - _R. J. Mathar_, Mar 01 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005531 | OEIS-A176858

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=4, k=-1 and l=0.

Conjecture: (n+1)*a(n) +(-7*n+2)*a(n-1) +3*(n+1)*a(n-2) +(31*n-104)*a(n-3) +2*(-22*n+89)*a(n-4) +16*(n-5)*a(n-5)=0. - _R. J. Mathar_, Mar 01 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005532 | OEIS-A176859

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=5, k=-1 and l=0.

Conjecture: (n+1)*a(n) +(-7*n+2)*a(n-1) +(-n+11)*a(n-2) +(43*n-140)*a(n-3) +2*(-28*n+113)*a(n-4) +20*(n-5)*a(n-5)=0. - _R. J. Mathar_, Feb 21 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005533 | OEIS-A176898

a(n) = binomial(6*n, 3*n)*binomial(3*n, n)/(2*(2*n+1)*binomial(2*n, n)).

Zhi-Wei Sun formulated two conjectures concerning a(n) (see Conjectures 1.1 and 1.2 in Z.-W. Sun, "Products and sums divisible by central binomial coefficients" and Conjecture A89 in "Open conjectures on congruences"). The first conjecture states that Sum_{n=1..p-1} a(n)/(108^n) is congruent to 0 or -1 modulo a prime p > 3 depending on whether p is congruent to +-1 or +-5 modulo 12, respectively. The second conjecture asks about an exact formula for a companion sequence of a(n). Both conjectures as well as many numerical congruences involving a(n) and (2n+1)a(n) were solved by Kh. Hessami Pilehrood and T. Hessami Pilehrood, see the link below. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 28 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005534 | OEIS-A176948

a(n) is the smallest solution x to A176774(x)=n; a(n)=0 if this equation has no solution.

Conjecture: For every n >= 4, except for n=6, there exists an n-gonal number N which is not k-gonal for 3 <= k < n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005535 | OEIS-A176954

a(n+1) = 1 + Sum_{p=0..n} a(p)*a(n-p)+k for n>=1, with here a(0)=1, a(1)=2, k=-1 and l=-1.

Conjecture: (n+1)*a(n) +(2-7n)*a(n-1) +(11n-13)*a(n-2) +(11n-46)*a(n-3) +4*(29-7n)*a(n-4) +12(n-5)*a(n-5)=0. - _R. J. Mathar_, Nov 21 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, Apr 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005536 | OEIS-A176969

Numbers n such that n^2 + 13^2 is prime.

The n^2 + d conjecture is a famous and still unsolved problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Eva-Maria Zschorn (e-m.zschorn(AT)zaschendorf.km3.de), Apr 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005537 | OEIS-A176997

Integers k such that 2^(k-1) == 1 (mod k).

The union of {1}, the odd primes, and the Fermat pseudoprimes, i.e., {1} U A065091 U A001567. Also, the union of A006005 and A001567 (conjectured by _Alois P. Heinz_, Dec 10 2010). - _Max Alekseyev_, Sep 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Dec 08 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005538 | OEIS-A177025

Number of ways to represent n as a polygonal number.

Conjecture: Every positive integer appears in the sequence. G.f.: x * Sum_{k>=2} x^k / (1 - x^(k*(k + 1)/2)) (conjecture). - _Ilya Gutkovskiy_, Apr 09 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 01 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005539 | OEIS-A177049

Numerator of (3n+1)*(3n+2)/4.

Conjecture: a(n)= +3*a(n-1) -6*a(n-2) +10*a(n-3) -12*a(n-4) +12*a(n-5) -10*a(n-6) +6*a(n-7) -3*a(n-8) +a(n-9) with g.f. -(x^2+4*x+1)*(x^6-2*x^5+12*x^4-13*x^3+12*x^2-2*x+1) / ((x-1)^3*(x^2+1)^3). - _R. J. Mathar_, Dec 12 2010 The conjecture is correct. - _Charles R Greathouse IV_, Feb 08 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Dec 09 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005540 | OEIS-A177111

Sequence defined by the recurrence formula a(n+1) = sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=1, k=-2 and l=0.

Conjecture: +(n+1)*a(n) +(-7*n+2)*a(n-1) +3*(5*n-7)*a(n-2) +3*(n-8)*a(n-3) +2*(-10*n+41)*a(n-4) +8*(n-5)*a(n-5)=0. - _R. J. Mathar_, Mar 02 2016 Conjectured recurrence follows from the differential equation (8*z^6 - 20*z^5 + 3*z^4 + 15*z^3 - 7*z^2 + z) * f'(z) + (2*z^4 - 15*z^3 + 9*z^2 - 5*z + 1) * f(z) + 4*z^4 - 11*z^3 + 9*z^2 + 3*z - 1 = 0 satisfied by the g.f. - _Robert Israel_, Jan 03 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005541 | OEIS-A177113

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=2, k=-2 and l=0.

Conjecture: +(n+1)*a(n) +(-7*n+2)*a(n-1) +(11*n-13)*a(n-2) +15*(n-4)*a(n-3) +2*(-16*n+65)*a(n-4) +12*(n-5)*a(n-5)=0. - _R. J. Mathar_, Mar 02 2016 Conjecture verified, using the differential equation 6*z^4 - 17*z^3 + 15*z^2 + z - 1 + (2*z^4 - 15*z^3 + 9*z^2 - 5*z + 1)*y(z) + (12*z^6 - 32*z^5 + 15*z^4 + 11*z^3 - 7*z^2 + z)*y'(z) = 0 satisfied by the g.f. - _Robert Israel_, Jan 06 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005542 | OEIS-A177115

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=3, k=-2 and l=0.

Conjecture: +(n+1)*a(n) +(-7*n+2)*a(n-1) +(7*n-5)*a(n-2) +3*(9*n-32)*a(n-3) +2*(-22*n+89)*a(n-4) +16*(n-5)*a(n-5)=0. - _R. J. Mathar_, Mar 02 2016 Conjecture verified using the differential equation (8*z^4 - 23*z^3 + 21*z^2 - z - 1 + (2*z^4 - 15*z^3 + 9*z^2 - 5*z + 1)*y(z)) + (16*z^6 - 44*z^5 + 27*z^4 + 7*z^3 - 7*z^2 + z)*y'(z) = 0 satisfied by the g.f. - _Robert Israel_, Jan 06 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005543 | OEIS-A177117

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=4, k=-2 and l=0.

Conjecture: (n+1)*a(n) +(-7*n+2)*a(n-1) +3*(n+1)*a(n-2) +3*(13*n-44)*a(n-3) +2*(-28*n+113)*a(n-4) +20*(n-5)*a(n-5)=0. - _R. J. Mathar_, Mar 02 2016 Conjecture verified using differential equation 10*z^4 - 29*z^3 + 27*z^2 - 3*z - 1 + (2*z^4 - 15*z^3 + 9*z^2 - 5*z + 1)*f(z) + (20*z^6 - 56*z^5 + 39*z^4 + 3*z^3 - 7*z^2 + z)*f'(z) satisfied by the g.f. - _Robert Israel_, Jan 07 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005544 | OEIS-A177118

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=5, k=-2 and l=0.

Conjecture: (n+1)*a(n) -(7*n-2)*a(n-1) -(n-11)*a(n-2) +3*(17*n-56)*a(n-3) -2*(34*n-137)*a(n-4) +24*(n-5)*a(n-5)=0. - _R. J. Mathar_, Jul 24 2012 Conjecture verified using differential equation 12*z^4 - 35*z^3 + 33*z^2 - 5*z - 1 + (2*z^4 - 15*z^3 + 9*z^2 - 5*z + 1)*f(z) + (24*z^6 - 68*z^5 + 51*z^4 - z^3 - 7*z^2 + z)*f'(z) = 0 satisfied by the g.f. - _Robert Israel_, Jan 07 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005545 | OEIS-A177122

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=6, k=1 and l=1.

Conjecture: +(n+1)*a(n) +(-7*n+2)*a(n-1) +(-5*n+19)*a(n-2) +(35*n-106)*a(n-3) +36*(-n+4)*a(n-4) +12*(n-5)*a(n-5)=0. - _R. J. Mathar_, Mar 02 2016 Conjecture verified using differential equation 6*z^4 - 18*z^3 + 18*z^2 - 7*z - 1 + (-z^3 + 9*z^2 - 5*z + 1)*f(z) + (12*z^6 - 36*z^5 + 35*z^4 - 5*z^3 - 7*z^2 + z)*f'(z) = 0 satisfied by the g.f. - _Robert Israel_, Jan 07 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005546 | OEIS-A177123

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=7, k=1 and l=1.

Conjecture: +(n+1)*a(n) +(-7*n+2)*a(n-1) +9*(-n+3)*a(n-2) +(47*n-142)*a(n-3) +48*(-n+4)*a(n-4) +16*(n-5)*a(n-5)=0. - _R. J. Mathar_, Mar 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005547 | OEIS-A177124

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=8, k=1 and l=1.

Conjecture: +(n+1)*a(n) +(-7*n+2)*a(n-1) +(-13*n+35)*a(n-2) +(59*n-178)*a(n-3) +60*(-n+4)*a(n-4) +20*(n-5)*a(n-5)=0. - _R. J. Mathar_, Mar 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005548 | OEIS-A177125

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=9, k=1 and l=1.

Conjecture: +(n+1)*a(n) +(-7*n+2)*a(n-1) +(-17*n+43)*a(n-2) +(71*n-214)*a(n-3) +72*(-n+4)*a(n-4) +24*(n-5)*a(n-5)=0. - _R. J. Mathar_, Mar 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005549 | OEIS-A177126

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=10, k=1 and l=1.

Conjecture: n*(n+1)*a(n) -n*(7*n-2)*a(n-1) -3*n*(7*n-17)*a(n-2) +n*(83*n-250)*a(n-3) -84*n*(n-4)*a(n-4) +28*n*(n-5)*a(n-5) =0. - _R. J. Mathar_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005550 | OEIS-A177127

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=6, k=0 and l=1.

Conjecture: +(n+1)*a(n) +2*(-3*n+1)*a(n-1) +(-11*n+27)*a(n-2) +32*(n-3)*a(n-3) +16*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Mar 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005551 | OEIS-A177128

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=7, k=0 and l=1.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +5*(-3*n+7)*a(n-2) +40*(n-3)*a(n-3) +20*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Mar 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005552 | OEIS-A177129

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=8, k=0 and l=1.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +(-19*n+43)*a(n-2) +48*(n-3)*a(n-3) +24*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005553 | OEIS-A177130

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=9, k=0 and l=1.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +(-23*n+51)*a(n-2) +56*(n-3)*a(n-3) +28*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Mar 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005554 | OEIS-A177131

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=10, k=0 and l=1.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +(-27*n+59)*a(n-2) +64*(n-3)*a(n-3) +32*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005555 | OEIS-A177168

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=6, k=0 and l=-2.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +(-11*n+27)*a(n-2) +2*(22*n-69)*a(n-3) +28*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 04 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005556 | OEIS-A177169

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=7, k=0 and l=-2.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +5*(-3*n+7)*a(n-2) +2*(26*n-81)*a(n-3) +32*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 04 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005557 | OEIS-A177170

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=8, k=0 and l=-2.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +(-19*n+43)*a(n-2) +6*(10*n-31)*a(n-3) +36*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 04 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005558 | OEIS-A177171

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=9, k=0 and l=-2.

Conjecture: +(n+1)*a(n) +2*(-3*n+1)*a(n-1) +(-23*n+51)*a(n-2) +2*(34*n-105)*a(n-3) +40*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Mar 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 04 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005559 | OEIS-A177172

Sequence defined by the recurrence formula a(n+1)=sum(a(p)*a(n-p)+k,p=0..n)+l for n>=1, with here a(0)=1, a(1)=10, k=0 and l=-2.

Conjecture: (n+1)*a(n) +2*(-3*n+1)*a(n-1) +(-27*n+59)*a(n-2) +2*(38*n-117)*a(n-3) +44*(-n+4)*a(n-4)=0. - _R. J. Mathar_, Feb 21 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Choulet_, May 04 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005560 | OEIS-A177258

Number of derangements of {1,2,...,n} having no adjacent transpositions.

Conjecture D-finite with recurrence a(n) = (n-1)*a(n-1) + (n-1)*a(n-2) + (n-1)*a(n-3) + a(n-4). - _R. J. Mathar_, Jul 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, May 08 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005561 | OEIS-A177259

Number of derangements of {1,2,...,n} having no adjacent 3-cycles (an adjacent 3-cycle is a cycle of the form (i,i+1,i+2)).

Conjecture: D-finite with recurrence a(n) = (n-1)*a(n-1) + (n-1)*a(n-2) + a(n-3) + (n-1)*a(n-4) + 2*a(n-6). - _R. J. Mathar_, Jul 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, May 08 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005562 | OEIS-A177260

Number of derangements of {1,2,...,n} having no adjacent 4-cycles (an adjacent 4-cycle is a cycle of the form (i,i+1,i+2,i+3)).

Conjecture: D-finite with recurrence a(n) = (n-1)*a(n-1) + (n-1)*a(n-2) + 2*a(n-4) + (n-1)*a(n-5) + 3*a(n-8). - _R. J. Mathar_, Jul 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, May 08 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005563 | OEIS-A177265

Number of permutations of {1,2,...,n} having exactly one string of consecutive fixed points (including singletons).

Conjecture: D-finite with recurrence a(n) - (n-1)*a(n-1) - (n-1)*a(n-2) +(n-1)*a(n-3) + (n-2)*a(n-4) = 0. - _R. J. Mathar_, Jul 01 2022 The above conjecture is correct. See Ray, 2026. - _Deepnil Ray_, Aug 10 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, May 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005564 | OEIS-A177316

Number of permutations of n copies of 1..4 with all adjacent differences <= 1 in absolute value.

See A103882 and A177317 through A177328 for the number of permutations of n copies of 1..k (for different values of k) with adjacent differences restricted in size. We conjecture that all these sequences satisfy the congruences A(n*p^k) == A(n*p^(k-1)) (mod p^(3*k)) for all positive integers n and k and any prime p >= 5. - _Peter Bala_, Jan 16 2020 Conjecture: a(n) = (1/3)*(A005259(n) + A005259(n-1)). Equivalently, a(n) = Sum_{k = 0..n} binomial(n,k)^2*binomial(n+k-1,k)^2. Cf. A103882. If true, then the sequence satisfies the recurrence a(n) = (2*(102*n^6 - 612*n^5 + 1462*n^4 - 1768*n^3 + 1143*n^2 - 382*n+52) * a(n-1) - (2*n-1)*(3*n^2 - 3*n+1) * (n-2)^3 * a(n-2)) / (n^3*(2*n - 3) * (3*n^2 - 9*n+7)) and the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for all positive integers n and k and any prime p >= 5. [added Apr 18 2022: assuming the recurrence given in the Maple program below is correct then these conjectures are true.] (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 06 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005565 | OEIS-A177322

Number of permutations of n copies of 1..4 with all adjacent differences <= 2 in absolute value.

The following are conjectural:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 06 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005566 | OEIS-A177354

a(n) is the moment of order n for the density measure 24*x^4*exp(-x)/((x^4*exp(-x)*Ei(x) - x^3 - x^2 - 2*x - 6)^2 + Pi^2*x^8*exp(-2*x)) over the interval 0..infinity.

a(n) = 5*A111532(n+1) (conjecture). - _R. J. Mathar_, Dec 14 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Groux Roland_, Dec 10 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005567 | OEIS-A177356

a(n) is the index of the first 0 term in the rumor sequence with initial 0th term 1 and parameters b = 2 and n.

A rumor sequence (running modulus recurrence sequence) is defined as follows: fix integer parameters b > 1 and n > 0. Set z[0] = any integer, and, for k > 0, define z[k] to be the least nonnegative residue of b*z[k-1] modulo (k+n). The rumor sequence conjecture states that all such rumor sequences are eventually 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jerry Metzger_, Dec 10 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005568 | OEIS-A177374

Fibonacci numbers whose decimal expansion does not contain any digit 8.

The above conjecture is true up through Fib(100,000) which has 20,800 digits. - _Harvey P. Dale_, Dec 31 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carmine Suriano_, May 07 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005569 | OEIS-A177390

Triangle T where the g.f. of row n of T^(2n) = (2n^2 + y)^n for n>=0, as read by rows, where T^n denotes the n-th matrix power of T.

Conjecture: for all integer k, there exists an integer triangle J such that the g.f. of row n of J^(k*n) = (k*n^2 + y)^n for n>=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005570 | OEIS-A177430

Least monotonically increasing logarithmic derivative consisting of only squares.

Conjecture: the sequence {a(n)^(m/2), n>=1} forms a logarithmic derivative of an integer sequence only when m is a nonnegative even integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 06 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005571 | OEIS-A177431

a(n) = sqrt(A177430(n)) where A177430 is the least monotonically increasing logarithmic derivative consisting of only squares.

Conjecture: the series exp(Sum_{n>=1} a(n)^m*x^n/n) consists entirely of integer coefficients only when m is a nonnegative even integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 06 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005572 | OEIS-A177680

Numbers n such that 6n and 12n are both the average of twin prime pairs.

Conjecture: all terms but the first term are multiples of 5. - _Harvey P. Dale_, Sep 26 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, May 10 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005573 | OEIS-A177768

Positions of decreases in sequence A061712 (related to Hamming weights of primes).

Conjecture: The numbers 2^k-1 for k > 2 are terms. - _Chai Wah Wu_, Sep 02 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 13 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005574 | OEIS-A177770

Conjectured positive numbers which have more than one representation (m,s) as a difference s^2 - m^5, m >= 1, s > 0.

Conjectured positive numbers which have more than one representation (m,s) as a difference s^2 - m^5, m >= 1, s > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, May 13 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005575 | OEIS-A177775

G.f. A(x) satisfies: [x^n] A^{n}(x) = [x^n] A^{n-1}(x) for n > 2 where A^{n+1}(x) = A^{n}(A(x)) denotes iteration with A^0(x)=x.

Conjecture: a(n) is odd iff n is a power of 2. - _Paul D. Hanna_, May 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 13 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005576 | OEIS-A177780

E.g.f. satisfies: L(x) = x*Sum_{n>=0} (2^n/n!)*Product_{k=0..n-1} L(3^k*x).

More generally, we have the following conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005577 | OEIS-A177781

E.g.f. satisfies: L(x) = x*Sum_{n>=0} 3^n/n!*Product_{k=0..n-1} L(4^k*x).

More generally, we have the following conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005578 | OEIS-A177800

The maximal length of the chain of consecutive primes starting with A177798(n) such that all of them are odious.

In contrast to sequence of all positive integers, where the length of a chain of consecutive odious numbers cannot exceed 2, we conjecture that over primes the length is not bounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Dec 12 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005579 | OEIS-A177801

Record lengths of chains of consecutive evil primes, starting with A177748(n).

In contrast to the sequence of all positive integers, where the length of every chain of consecutive evil numbers cannot exceed 2, we conjecture that for the sequence of primes such length is not bounded with growth of n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Dec 12 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005580 | OEIS-A177960

Numbers of the form A001317(t), excluding those at places of the form t=m*(2^k-1), m>=0, k>=2.

Conjecture: If for every m>=2, to consider triangle of m-nomial coefficients mod 2 converted to decimal, then the sequence lists terms of A001317 which are not in the union of other sequences for m=3 (A038184), 4 (A177882), 5, 6,...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Dec 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005581 | OEIS-A178072

Sequence with Hankel transform equal to the Somos-4 sequence A006769(n+2).

Conjecture: 2*n*(n+1)*a(n) +2*n*(n+1)*a(n-1) -3*(3*n-1)*(3*n-4)*a(n-2) +(61*n^2-191*n+36)*a(n-3) +6*(-2*n^2+2*n-1)*a(n-4) +2*(5*n-1)*(4*n-15)*a(n-5) +n*(n-5)*a(n-6) +(5*n-1)*(n-6)*a(n-7)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 19 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005582 | OEIS-A178113

Transform of C(n+1,floor((n+1)/2)) by A178112.

Conjecture: a(2*n) = A005773(n+1), a(2*n+1) = 2*a(2*n). - _Jason Yuen_, Feb 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, May 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005583 | OEIS-A178127

Lesser of twin Ramanujan primes.

Conjecture: For all n > 570, more than 1/4 of the twin prime pairs < n are both Ramanujan primes. Motivation for the conjecture is in "Ramanujan primes and Bertrand's postulate" Section 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, May 20 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005584 | OEIS-A178169

Numbers in A178168, sorted.

At least the first 12332052 terms of this sequence are unique. It is conjectured that all terms are unique. The sequence is easy to compute by generating the Collatz tree on the fly while keeping track of products.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, May 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005585 | OEIS-A178207

a(n) = 4*n - 2*A000120(n) + 1 where A000120(n) = number of nonzero digits in the binary representation of n.

Baralic provides some evidence for the conjecture that each n-dimensional, compact smooth manifold M^n (n>1), admits a totally skew embedding in the Euclidean space of dimension N = 4n - 2alpha(n) + 1 where alpha(n) = number of nonzero digits in the binary representation of n. The subsequence of primes begins: 3, 7, 23, 31, 37, 47, 53, 71, 79, 83, 101, 107, 109, 113, 127, 149, 157. See also my note on the Whitney immersion theorem in A005187.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, May 22 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005586 | OEIS-A178237

Smallest prime p of the form prime(n)+k^2 such that sum of digits(p) = prime(n).

It is still an open problem if there exist infinitely many primes of form k^2 + d (d integer, no negative square).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), May 23 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005587 | OEIS-A178253

a(n) is the least positive integer m such that (A001969(n))^m is in A000069.

Conjecture: a(n) exists for all n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 24 2010, May 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005588 | OEIS-A178294

Number of collinear point triples in a 4 X 4 X 4 X... n-dimensional cubic grid.

Conjecture: a(n) = 8^n/2-3*4^n/2+6^n. a(n) = +18*a(n-1) -104*a(n-2) +192*a(n-3). G.f.: 4*x*(-1+7*x)/((6*x-1)*(8*x-1)*(4*x-1)). - _R. J. Mathar_, May 24 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, suggested by R. J. Mathar in the Sequence Fans Mailing List, May 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005589 | OEIS-A178295

Number of collinear point 4-tuples in a 5 X 5 X 5 X... n-dimensional cubic grid.

Conjecture: a(n) = 9^n/2+3*7^n/2-2*5^n. a(n)= +21*a(n-1) -143*a(n-2) +315*a(n-3). G.f.: x*(-5+41*x)/((9*x-1)*(7*x-1)*(5*x-1)). [_R. J. Mathar_, May 24 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, suggested by R. J. Mathar in the Sequence Fans Mailing List, May 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005590 | OEIS-A178296

Number of collinear point 5-tuples in a 6 X 6 X 6 X... n-dimensional cubic grid.

Conjecture: a(n) = 10^n/2+2*8^n-5*6^n/2. a(n)= 24*a(n-1) -188*a(n-2) +480*a(n-3). G.f.: 2*x*(-3+28*x)/((6*x-1)*(8*x-1)*(10*x-1)). [_R. J. Mathar_, May 24 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, suggested by R. J. Mathar in the Sequence Fans Mailing List, May 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005591 | OEIS-A178297

Number of collinear point 6-tuples in a 7 X 7 X 7 X... n-dimensional cubic grid.

Conjectures from _R. J. Mathar_, May 24 2010: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, suggested by _R. J. Mathar_ in the Sequence Fans Mailing List, May 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005592 | OEIS-A178299

Number of collinear point 4-tuples in a 6 X 6 X 6 X... n-dimensional cubic grid.

Conjectures from _R. J. Mathar_, May 24 2010: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, suggested by _R. J. Mathar_ in the Sequence Fans Mailing List, May 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005593 | OEIS-A178307

Remove powers of 2 from A000069. Let b(n) be n-th term of the remaining sequence. Then a(n) is the least number m such that (b(n))^m is in A001969.

The sequence {b(n)} coincides with A075930. Conjecture. For every n>=1, a(n) does exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 24 2010, May 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005594 | OEIS-A178319

E.g.f.: (Sum_{n>=0} 3^(n*(n + 1)/2) * x^n/n!)^(1/3).

a(n) = 1 (mod 6) for n >= 0 (conjecture). General conjecture: [x^n/n!] E(q*x, q)^(1/q) = 1 (mod q(q-1)) for n >= 0 and integer q > 1 where E(x, q) = Sum_{n>=0} q^(n*(n - 1)/2)*x^n/n!.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005595 | OEIS-A178351

Number of collinear point 5-tuples in a 7 X 7 X 7 X ... n-dimensional cubic grid.

Conjectures from _R. J. Mathar_, May 25 2010: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, suggested by _R. J. Mathar_ in the Sequence Fans Mailing List, May 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005596 | OEIS-A178352

Number of collinear point triples in a 6 X 6 X 6 X ... n-dimensional cubic grid.

Conjectures from _R. J. Mathar_, May 25 2010: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, suggested by _R. J. Mathar_ in the Sequence Fans Mailing List, May 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005597 | OEIS-A178353

Number of collinear point 4-tuples in a 7 X 7 X 7 X ... n-dimensional cubic grid.

Conjectures from _R. J. Mathar_, May 25 2010: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, suggested by _R. J. Mathar_ in the Sequence Fans Mailing List, May 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005598 | OEIS-A178375

The greatest common prime divisor of A000032(n)-1 and A001608(n), or 1 if no such divisor exists.

Conjecture: Records of the sequence are consecutive primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 26 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005599 | OEIS-A178444

Markov numbers that are prime.

Bourgain, Gamburd, and Sarnak have announced a proof that almost all Markoff numbers are composite--see A256395. Equivalently, the present sequence has density zero among all Markoff numbers. (It is conjectured that the sequence is infinite.) - _Jonathan Sondow_, Apr 30 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Muljadi_, Jan 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005600 | OEIS-A178449

Conjectured expansion of exp(Pi*sqrt(163)) in powers of t, where t = 1/(640320)^3.

Conjectured expansion of exp(Pi*sqrt(163)) in powers of t, where t = 1/(640320)^3. R. W. Gosper asks if the coefficients are well-defined. Until this is answered, the sequence is only conjectural. This sequence is very close to A178451, but presumably different from it.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 22 2010, based on a posting by R. W. Gosper to the Sequence Fans Mailing List, Dec 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005601 | OEIS-A178456

Primes p such that p-1 or p+1 has more than two distinct prime divisors.

Sequence contains many pairs of twin primes. More exactly, denote A(x), t(x),T(x) the counting functions of this sequence, twin primes in this sequence and all twin primes correspondingly. In supposition of the infinitude of twin primes, the very plausible conjectures are: (1) for x tends to infinity, t(x)~T(x) and (2) for x >= 31, t(x)/A(x) > T(x)/pi(x).

Indeed (heuristic arguments), the middles of twin pairs (beginning with the second pair) belong to progression {6*n}. Let us choose randomly n. The probability that n has prime divisors 2,3 only is, as well known, O((log n)^2/n), i.e. it is quite natural to conjecture that almost all twin pairs are in the sequence. Besides, it is natural to conjecture that the inequality is true as well, since A(x)<pi(x).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Dec 23 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005602 | OEIS-A178473

For n>=0, let n!^(4) = A202369(n+1) and, for 0<=m<=n, C^(4)(n,m) = n!^(4)/(m!^(4)*(n-m)!^(4)). The sequence gives triangle of numbers C^(4)(n,m) with rows of length n+1.

Conjecture. If p is prime of the form 4*k+1, then the k-th row contains two 1's and k-1 numbers multiple of p; if p is prime of the form 4*k+3, then the (2*k+1)-th row contains two 1's and 2*k numbers multiple of p. Conjecture. A007814(C^(4)(n,m)) = A007814(C(n,m)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005603 | OEIS-A178481

Number of steps of the map x -> A055566(x), starting at n, before reaching the end of the cycle.

Generalization for the k-th powers and conjecture: For each k >= 1, iteration of taking the k-th powers of digit sums reaches a cycle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, May 28 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005604 | OEIS-A178482

Phi-antipalindromic numbers.

Conjecture: this is the sequence of numbers k for which f(k) is an integer, where f(x) is the change-of-base function defined at A214969 using b=phi and c=b^2. - _Clark Kimberling_, Oct 17 2012 Kimberling's conjecture has been proven by Ingrid Vukusic and myself. Along the way we

prove an alternate characterization of the sequence: they are the positive integers whose base-phi expansion consists only of even exponents of phi. - _Jeffrey Shallit_, Aug 28 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 28 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005605 | OEIS-A178520

Number of ordered trees with n edges and with no vertex of outdegree 2 that have two leaves as their two children.

Conjecture D-finite with recurrence: (n+1)*a(n) +(n+2)*a(n-1) +(n+9)*a(n-2) +(-82*n+203)*a(n-3) +5*(2*n-9)*a(n-4) +21*(2*n-11)*a(n-5) +(n-8)*a(n-6) +5*(n-9)*a(n-7) +21*(n-10)*a(n-8)=0. - _R. J. Mathar_, Jan 28 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, May 31 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005606 | OEIS-A178527

Primes p such that either p - 2 or p + 2 has more than two distinct prime divisors.

Sequence contains "many" pairs of cousin primes. More exactly, our conjectures are: (1) sequence contains almost all cousin primes; (2)for x >= 107, c(x)/A(x) > C(x)/pi(x), where A(x), c(x) and C(x) are the counting functions for this sequence, cousin pairs in this sequence and all cousin pairs respectively. The probability that n has no more than two prime divisors is, as well known, O(log(log(n))/log(n)), i.e., it is natural to conjecture that almost all cousin pairs are in the sequence. Furthermore, it is natural to conjecture that the inequality is true as well, since A(x) < pi(x).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Dec 23 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005607 | OEIS-A178534

Triangle T(n,k) read by rows. T(n,1) = A000045(n+1), k > 1: T(n,k) = (Sum_{i=1..k-1} T(n-i,k-1)) - (Sum_{i=1..k-1} T(n-i,k)).

Conjecture (by extrapolating): G.f. k-th column: x^k*(1-x^2)/((1-x-x^2)*(1-x^k)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, May 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005608 | OEIS-A178536

First column of A178535.

a(n) = Sum_{k|n} A008683(n/k)*([k = 1] - [2|k]) (conjecture). - _Mats Granvik_, Jan 24 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, May 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005609 | OEIS-A178545

Primes p such that q = p^2 + p + 1 is an emirp.

It is conjectured (but still an open problem) that there exist infinitely many primes of the form n^2 + n + 1 = ((2*n+1)^2 + 3)/4. Landau's 4th problem from (1912, 5th Congress of Mathematicians in Cambridge) conjectures that there are infinitely many primes of the form n^2 + 1 (also Euler 1760; Mirsky 1949). Hardy and Littlewood proposed a conjecture about the asymptotic number of primes of the form n^2 + 1. It is conjectured but also unproved that there are infinitely many emirps (see A048054).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Ulrich Krug (leuchtfeuer37(AT)gmx.de), May 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005610 | OEIS-A178612

Positive numbers of the form p^6 - 4*p^4*q + 4*p^2*q^2 + 4*q^3 (and p*q <> 0).

Conjecture: There are no perfect squares in this sequence (in spite of all numbers being congruent to 0 or 1 mod 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, May 30 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005611 | OEIS-A178623

Triangle T(n,m) read by rows: T(n,0)= prime(n); T(n,m)=1 if m>=1.

The sequence reflects a conjecture on the denominator of inverse Bernoulli polynomials in A178340: if the row index is one less than one of the primes in A008578, the row of denominators starts with that prime and contains 1's in the remaining entries.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, May 31 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005612 | OEIS-A178762

Prime numbers that are Fibonacci integers.

A Fibonacci integer is a number that can be written as the product and/or quotient of Fibonacci numbers (A000045). For example, 107 is a Fibonacci integer because Fib(36)/(Fib(18)*Fib(3)*Fib(4)^3) = 107. Observe that the prime Fibonacci numbers (A005478) are a subset of these primes. Luca, Pomerance, and Wagner conjecture that this sequence is infinite. The paper's Remark 2 and sequences A152012, A178763, and A178764 are useful in finding these primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jun 10 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005613 | OEIS-A178768

Decimal expansion of real constant in an explicit counterexample to the Lagarias-Wang finiteness conjecture.

Decimal expansion of real constant in an explicit counterexample to the Lagarias-Wang finiteness conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jun 11 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005614 | OEIS-A178785

a(n) is the smallest n-perfect number of the form 2^(n+1)*L, where L is an odd number with exponents <= n in its prime power factorization, and a(n)=0 if no such n-perfect number exists.

Let k >= 1. In the multiplicative basis Q^(k) = {p^(k+1)^j, p runs through A000040, j=0,1,...} every positive integer m has a unique factorization of the form m = Product_{q is in Q^(k)} q^(m_q), where m_q is in {0,1,...,k}. In particular, in the case of k=1, we have the unique factorization over distinct terms of A050376. Notice that the standard prime basis is the limiting value for k tending to infinity, and, by the definition, Q^(infinity)=A000040. The number d is called a k-divisor of m if the exponents d_q in its factorization in the basis Q^(k) do not exceed m_q. A number m is called k-perfect if it equals to the sum of its proper positive k-divisors. Conjecture: a(11)=0. Note that we also know of n-perfect numbers for n = 12, 14, 15, 16, and 18.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jun 14 2010, Jun 18 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005615 | OEIS-A178790

The arithmetic mean of (2*k+1)*A_k (k=0,...,n-1), where A_0,A_1,... are Apery numbers given by A005259.

Conjecture: the number a(n) = n^(-1)*Sum_{k=0..n-1}(2*k+1)*A_k is always an integer. Conjecture: If p=5,7 (mod 8) is a prime then sum_{k=0}^{p-1}A_k=0 (mod p^2); if p=1,3 (mod 8) is a prime greater than 3 and p=x^2+2y^2 with x,y integers then sum_{k=0}^{p-1}A_k=4x^2-2p (mod p^2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 14 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005616 | OEIS-A178791

The arithmetic mean of (2k+1)(-1)^k*A_k (k=0,...,n-1), where A_0, A_1,... are Apéry numbers given by A005259.

On Jun 14 2010, _Zhi-Wei Sun_ conjectured that the number a(n) = n^{-1}*Sum_{k=0..n-1} (2*k+1)*(-1)^k*A_k is always an integer and that a(p) = p(p/3) (mod p^3) for any prime p>3. He also formulated the following conjecture: If p=1 (mod 3) is a prime and p = x^2 + 3y^2 with x,y integers then Sum_{k=0..p-1} (-1)^k*A_k = 4*x^2 - 2*p (mod p^2); if p=2 (mod 3) is a prime then Sum_{k=0..p-1}(-1)^k*A_k=0 (mod p^2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 14 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005617 | OEIS-A178808

a(n) = (1/n^2) * Sum_{k = 0..n-1} (2*k+1)*(D_k)^2, where D_0, D_1, ... are central Delannoy numbers given by A001850.

On Jun 14 2010, _Zhi-Wei Sun_ conjectured that a(n) = (1/n^2) * Sum_{k = 0..n-1} (2*k+1)*(D_k)^2 is always an integer and that p^2*a(p) = p^2 - 4*p^3*q_p(2) - 2*p^4*q_p(2)^2 (mod p^5) for any prime p > 3, where q_p(2) denotes the Fermat quotient (2^(p-1) - 1)/p (see Sun, Remark 4.3, p. 26, 2014). He also conjectured that Sum_{k = 0..n-1} (2*k+1)*(-1)^k*(D_k)^2 == 0 (mod n*D_n/(3,D_n)) for all n = 1,2,3,....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 16 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005618 | OEIS-A178815

First base of a nonzero Fermat quotient mod the n-th prime.

It is known that a(n) = O((log p_n)^2) as n -> oo. It is conjectured that a(n) = 3 if p_n is a Wieferich prime. See Section 1.1 in Ostafe-Shparlinski (2010).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Jun 17 2010, Jun 24 2010, Jun 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005619 | OEIS-A178841

The number of pure inverting compositions of n.

The pure inverting compositions index a basis for this module, as conjectured by F. Bergeron and C. Reutenauer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Aaron Lauve (lauve(AT)math.luc.edu), Jun 17 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005620 | OEIS-A178942

a(1) = 3; for n >= 2, a(n) is the smallest prime q > a(n-1) such that, for the previous prime p and the following prime r, the fraction (q-p)/(r-q) has denominator equal to A001223(n)/2 (or 0, if no such prime exists).

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005621 | OEIS-A179004

T(n,k)=Maximum 1's in an nXk binary array with no element equal to a strict majority of its horizontal and vertical neighbors.

Conjecture: away from boundaries a sliding 101 pattern would fill with a limiting 2/3 1's

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005622 | OEIS-A179017

Odd numbers c such that c*(c^2 - 1)/4 is squarefree.

This function is minimal orbital in hypothesis (a,b,c).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jun 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005623 | OEIS-A179071

Chapman's "evil" determinants I.

Guy says "Chapman has a number of conjectures which concern the distribution of quadratic residues." One is that if p_n == 3 (mod 4), then a(n) = 1. Chapman also has a conjecture if p_n == 1 (mod 4), involving the fundamental unit and class number of the quadratic field Q(sqrt(p)). (Added Aug 23 2011: Both conjectures have been proved by Vsemirnov.) It appears that a(n) is negative and even, if p_n == 1 (mod 4); see A179073. (Added Aug 28 2011: This conjecture has also been proved by Vsemirnov.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_ and _Wadim Zudilin_, Jun 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005624 | OEIS-A179073

A179071 for p == 1 (mod 4).

It appears that a(n) is negative and even. (Added Aug 28 2011: This conjecture has been proved by Maxim Vsemirnov.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_ and _Wadim Zudilin_, Jun 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005625 | OEIS-A179077

a(n) is the residue ((2^p - 2)/p) mod p, where p is the n-th prime.

Conjecture: a(n) is the residue A036968(p-1) (mod p) for p = prime(n). If the above conjecture is true, then a(n) = 0 if and only if p is a Wieferich prime (A001220) (cf. Hu et al., 2019, section 1.3). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jun 28 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005626 | OEIS-A179089

a(n) = (1/n^2) * Sum_{k=0..n-1} (2k+1)*T_k^2(-3)^(n-1-k), where T_0, T_1, ... are central trinomial coefficients given by A002426.

On Jun 28 2010, _Zhi-Wei Sun_ conjectured that a(n) is an integer for every n=1,2,3,... and that a(p) = (1+p/3)/2 (mod p) for any prime p, where (p/3) is the Legendre symbol. In contrast, he showed that Sum_{k=0..n-1} (2k+1)*T_k^2*3^(n-1-k) = n*T_n*T_{n-1} for all n=1,2,3,...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005627 | OEIS-A179095

Rectified 5-cell numbers: the coefficient of x^{2n-2} in (1+x+x^2+ ... + x^{n-1})^5.

Conjectures: a(n) = n*(11*n^3+6*n^2+n+6)/24. G.f.: x*(1+5*x+5*x^2)/(1-x)^5. - _Colin Barker_, Jan 09 2012 The conjectures in A179095-A179099 are true. Proof: and this agrees with the conjectures for k=5..9 (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael A. Jackson_, Jun 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005628 | OEIS-A179096

Rectified hexateron (5-simplex) numbers: the coefficient of x^(2n-2) in (1+x+x^2+...+x^(n-1))^6.

Conjectures: a(n) = n*(n+1)*(13*n^3+12*n^2-7*n+12)/60. G.f.: x*(1+9*x+x^3+15*x^2)/(x-1)^6. - _R. J. Mathar_, Jul 06 2010 These conjectures are true, see A179095 for proof.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael A. Jackson_, Jun 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005629 | OEIS-A179097

Rectified heptapeton (6-simplex) numbers: the coefficient of x^(2n-2) in (1+x+x^2+...+x^(n-1))^7.

Conjecture: a(n) = n*(40+26*n+5*n^2+75*n^3+75*n^4+19*n^5)/240. G.f.: x*(1+14*x+35*x^2+7*x^3)/(1-x)^7. - _Colin Barker_, Jan 09 2012 These conjectures are true, see A179095 for proof.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael A. Jackson_, Jun 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005630 | OEIS-A179098

Rectified 7-simplex number: the coefficient of x^(2n-2) in (1+x+x^2+...+x^(n-1))^8.

Conjectures: a(n) = n*(90+77*n+140*n^3+210*n^4+98*n^5+15*n^6)/630. G.f.: x*(1+20*x+70*x^2+28*x^3+x^4)/(1-x)^8. - _Colin Barker_, Jan 09 2012 These conjectures are true, see A179095 for proof.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael A. Jackson_, Jun 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005631 | OEIS-A179099

Rectified 8-simplex numbers: the coefficient of x^(2n-2) in (1+x+x^2+...+x^(n-1))^9.

This conjecture is true, see A179095 for proof.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael A. Jackson_, Jun 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005632 | OEIS-A179100

a(n) = (1/n) * Sum_{k=0..n-1} (8k+5) T_k^2, where T_0, T_1, ... are central trinomial coefficients given by A002426.

On Jun 17 2010, _Zhi-Wei Sun_ conjectured that a(n) is an integer for every n=1,2,3,... and that a(p) == 3(p/3) (mod p) for any prime p, where (p/3) is the Legendre symbol. He also observed that Sum_{k=0..n-1} (2k+1) T_k*3^{n-1-k} = n * Sum_{k=0..n-1} C(n-1,k)*(-1)^(n-1-k)*(k+1)*C(2k,k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005633 | OEIS-A179107

Records of minima of positive distance d between a square cubefree integer y and a cube of positive and squarefree integer x and such d = y^2 - x^3.

Next conjectured terms are: 4401169, 87002345, 193234265, 497218657.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jun 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005634 | OEIS-A179108

Values x for records of minima of positive distance d between a square cubefree integer y and a cube of positive and squarefree integer x and such d = y^2 - x^3.

Next conjectured terms are: 53197086958290, 12813608766102800, 810574762403977000, 471477085999389000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jun 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005635 | OEIS-A179113

Odd primes which can never divide 2^a+2^b+1.

Mariusz Skałba conjectures that this sequence has density zero among all primes but contains infinitely many primes based on the following observations. For any prime p in this sequence, the multiplicative order of 2 modulo p is <p^0.8 (Erdős conjectures that the set of such primes must have

density zero among all primes). Moreover, any number of the form 2^m-1 the number of whose prime factors counted with multiplicity is <log m/log 3 has at least one prime factor in this sequence (the infinitude of such numbers may be more tractable than the infinitude of Mersenne primes). - _Tomohiro Yamada_, Aug 08 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Phil Carmody_, Jan 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005636 | OEIS-A179118

Number of Collatz steps to reach 1 starting with 2^n + 1.

There are many long runs of consecutive terms that increase by 1 (see second conjecture in A277109). For n < 40000, the longest run has 1030 terms starting from a(33237) = 244868 and ending with a(34266) = 245897. - _Dmitry Kamenetsky_, Sep 30 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mitch Harris_, Jan 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005637 | OEIS-A179180

Partial sums of A007895.

Total number of summands in Zeckendorf representations of all the numbers 1,2,...,n (for n>0); see the conjecture at A214979. - _Clark Kimberling_, Oct 23 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Walt Rorie-Baety_, Jun 30 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005638 | OEIS-A179194

Bases n in which 1/(n-2) is non-terminating and has period n-3.

Conjecture: a(n) = A001122(n) + 2 (via formula at A217948). - _Bill McEachen_, Jun 11 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Will Nicholes_, Jul 01 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005639 | OEIS-A179210

a(n) is the smallest prime q such that (r-q)/(q-p) = n, where p<q<r are consecutive primes (or 0 if no such prime exists).

Conjecture: a(n) > 0 for all n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005640 | OEIS-A179234

a(n) is the smallest prime q such that, for the previous prime p and the following prime r, the fraction (r-q)/(q-p) has denominator n in lowest terms.

The conjecture that a(n) exists for every n is a weaker conjecture than a related one in the comment to A179210.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005641 | OEIS-A179238

Numerators in convergents to infinitely repeating period 3 palindromic continued fraction [1,2,1,...].

Conjecture relating palindromic continued fractions to the tangents of angles A and 2A: The conjecture applies to any palindromic form such as [a,a,a,...], [a,b,a,a,b,a,...], [a,b,b,a,a,b,b,a,...], [a,b,c,c,b,a,...] and so on. Conjecture proved by the author, Jul 05 2010. - _Gary W. Adamson_, Mar 25 2014 Tan(2) corrected to tan(A), which is the half angle of a right triangle with sides (a, b) and c = sqrt(a^2 + b^2), not an integer. Example: Given a right triangle with sides a = 3, b = 5, c = sqrt(34), we calculate the two acute angles 2A and divide each by two, giving us two angles A; then perform tan(A) on each. These have an algebraic equivalency of two constants: 3/(5 + sqrt(34)) = 0.276983964... and 5/(3 + sqrt(34)) = 0.566190378... The continued fraction representations of these constants are respectively barover[3,1,1,1,1,3] and barover[1,1,3,3,1,1], both palindromic. The conjecture and proof don't apply to Pythagorean triangles, in which case the tangents of the half angles would be rational. - _Gary W. Adamson_, Apr 07 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Jul 04 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005642 | OEIS-A179240

a(n) is the smallest prime q > a(n-1) such that, for the previous prime p and the following prime r, the fraction (q-p)/(r-q) has denominator equal to A006843(n) (or 0, if such a prime does not exist).

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005643 | OEIS-A179256

a(n) is the smallest prime q such that (q-p)/(r-q) = n, where p<q<r are consecutive primes (or 0 if none exist).

Conjecture: a(n)>0 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005644 | OEIS-A179269

Number of partitions of n into distinct parts such that the successive differences of consecutive parts are increasing, and first difference > first part.

G.f.: Sum_{k>=0} x^(k*(k+1)*(k+2)/6) / Product_{j=1..k} (1 - x^(j*(j+1)/2)) (conjecture). - _Ilya Gutkovskiy_, Apr 25 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Jan 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005645 | OEIS-A179289

Smallest index k such that prime(k)*2^n-1 is prime, or zero if there is no prime.

It is conjectured that a(42228) is the first 0 term. This corresponds to the first Riesel number, 509203, which happens to be prime. See A101036. - _T. D. Noe_, Mar 23 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Jul 09 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005646 | OEIS-A179290

Decimal expansion of length of edge of a regular icosahedron with radius of circumscribed sphere = 1.

(10/3)*(this constant)=3.504874080794224... is the volume of the polyhedron with 32 edges with conjectured maximum volume inscribed in a sphere of radius 1. It has 60 congruent triangular faces and the symmetry group of the regular icosahedron. See Pfoertner links for visualizations. - _Hugo Pfoertner_, Aug 02 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Joseph Stephan Orlovsky_, Jul 09 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005647 | OEIS-A179295

a(n) is the least prime number such that prime(n)+a(n)+1 is a prime or -1 if no such prime number exists.

If Maillet's conjecture is true, then a(n) != -1 for all n. - _Chai Wah Wu_, Aug 01 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lars-Erik Svahn_, Jun 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005648 | OEIS-A179328

a(n) is the smallest prime q > a(n-1) such that, for the previous prime p and the following prime r, the fraction (q-p)/(r-q) has denominator prime(n) (or 0, if such a prime does not exist).

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005649 | OEIS-A179383

a(n) = 2*k(n)-1 where k(n) is the sequence of positions of records in A179382.

Some very plausible conjectures: 1) The sequence consists of primes and squares of primes; 2) The set of squares is finite; 3) A prime p>=5 is in the sequence iff it has primitive root 2 (A001122) ; 4) There exists l such that, for n>l, A179383(n) =A139099(n+l) . [From _Vladimir Shevelev_ , Jul 14 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jul 12 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005650 | OEIS-A179406

Conjectured records d such that for all nonsquare x > A179407(n), all positive differences x^5-y^2 exceed d.

Conjectured records d such that for all nonsquare x > A179407(n), all positive differences x^5-y^2 exceed d. Conjecture: For any positive number x >= A179407(n), the distance d between the fifth power of x and the square of any y (such that x != k^2 and y != k^5) can't be less than a(n). - _Artur Jasinski_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jul 13 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005651 | OEIS-A179479

a(n) is the smallest prime q > a(n-1) such that, for the previous prime p and the following prime r, the fraction (q-p)/(r-q) has denominator 2, for odd n and 1 for even n (or 0, if such a prime does not exist).

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005652 | OEIS-A179481

a(n) = 2*t(n)-1 where t(n) is the sequence of records positions of A179480.

In answer to the primality question and pursuant to the Coach Theorem of Hilton and Pedersen: phi(b) = 2 * k * c, with b an odd integer and k in A003558, and c (the numbers of coaches) in A135303; iff phi(b) = (b-1) then b = p, prime. This implies that if b has one coach and k = (b-1)/2, b must be prime since phi(b) = 2 * k * c = 2 * (b-1)/2 * 1 = (b-1). Conjectures: all terms in A179481 have one coach with k = (b-1)/2 and are therefore primes. Next, if A179480(n) is a new record high value, then so is A003558(n-1); but not necessarily the converse (e.g. 13), and the corresponding value of k for b is (b-1)/2. Examples: b = 13 has one coach with k (sum of bottom row terms) = 6 = A003558(6); and r (number of entries in each row) = 3:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jul 16 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005653 | OEIS-A179504

a(n) = sigma(2n,n) for n>0 with a(0)=1.

Conjecture: a(n) = A023887(n)*((2^n - 1)/(2^(n*(A007814(n) + 1)) - 1) + 2^n) for n > 0. - _Velin Yanev_, Jul 08 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 21 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005654 | OEIS-A179508

a(n) is the unique integer such that Sum_{k=0..p-1} b(k)/(-n)^k == a(n) (mod p) for any prime p not dividing n, where b(0), b(1), b(2), ... are Bell numbers given by A000110.

On July 17, 2010 Zhi-Wei Sun conjectured that a(n) exists for every n=1,2,3,... He noted that a(1)=2 since Sum_{k=0..p-1} (-1)^k * b(k) == b(p) (mod p), and conjectured that a(2)=1, a(3)=2, a(4)=-1, a(5)=10, a(6)=-43, a(7)=266, a(8)=-1853, a(9)=14834, a(10)=-133495. It seems that (-1)^(n-1)*a(n) > 0 for all n=3,4,5,...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 17 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005655 | OEIS-A179525

G.f.: A(x) = Sum_{n>=0} Product_{k=1..n} ((1+x)^k - 1).

Conjecturally, a(n) is asymptotically c*n!*(12/Pi^2)^n, where c=6*sqrt(2)*exp(-Pi^2/24)/Pi^2. - _Vít Jelínek_, Feb 12 2012 [This is correct: see Hwang and Jin, Table 3, p. 26. - _Peter Bala_, Jan 31 2021] G.f.: Q(0), where Q(k)= 1 - (1-(1+x)^(2*k+1))/(1 - (1-(1+x)^(2*k+2))/(1 - (1+x)^(2*k+2) - 1/Q(k+1))); (continued fraction). Conjecture. - _Sergei N. Gladkovskii_, May 13 2013 Conjectural g.f.: Sum_{n >= 0} 1/(1 + x)^(n+1)*Product_{k = 1..n} (1 - 1/(1 + x)^(2*k)). Conjectural g.f.: Sum_{n >= 0} (1 + x)^(2*n+1)*Product_{k = 1..2*n} (1 - (1 + x)^k). Cf. A158690, which has e.g.f. A(exp(x) - 1). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 17 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005656 | OEIS-A179529

Number of terms of A083207 in 12 consecutive numbers.

F. Buss and T. D. Noe conjectured a(n) > 0; this is correct (proof by R. Gerbicz);

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jul 19 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005657 | OEIS-A179535

a(n) = Sum_{k=0..n} binomial(n,k)^2 * binomial(n-k,k)^2 * 81^k.

On Jul 17 2010, _Zhi-Wei Sun_ introduced this sequence and made the following conjecture: If p is a prime with p=1,9,11,19 (mod 40) and p = x^2+10y^2 with x,y integers, then Sum_{k=0..p-1} a(k) == 4x^2-2p (mod p^2); if p is a prime with p == 7,13,23,37 (mod 40) and 2p = x^2 + 10y^2 with x,y integers, then Sum_{k=0..p-1} a(k) == 2p - 2x^2 (mod p^2); if p is an odd prime with (-10/p)=-1, then Sum_{k=0..p-1} a(k) == 0 (mod p^2). He also conjectured that Sum_{k=0..n-1} (10k+9)*a(k) == 0 (mod n) for all n=1,2,3,... and that Sum_{k=0..p-1} (10k+9)*a(k) == p(4(-2/p)+5) (mod p^2) for any prime p > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 18 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005658 | OEIS-A179536

a(n) = Sum_{k=0..n} binomial(n,k)^2*binomial(n-k,k)^2*(-324)^k.

On Jul 17 2010, _Zhi-Wei Sun_ introduced this sequence and made the following conjecture: If p is an odd prime with (p/13) = (-1/p) = 1 and p = x^2 + 13y^2 with x,y integers, then Sum_{k=0..p-1} a(k) == 4x^2 - 2p (mod p^2); if p is an odd prime with (p/13) = (-1/p) = -1 and 2p = x^2 + 13y^2 with x,y integers, then Sum_{k=0..p-1} a(k) == 2x^2 - 2p (mod p^2); if p > 3 is a prime with (p/13) = -(-1/p), then Sum_{k=0..p-1} a(k) == 0 (mod p^2). He also conjectured that Sum_{k=0..n-1} (260k+237)*a(k) == 0 (mod n) for all n=1,2,3,... and that Sum_{k=0..p-1} (260k+237)*a(k) == p(130(-1/p)+107) (mod p^2) for any prime p > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 18 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005659 | OEIS-A179538

Numbers 2n+1 for which A003558(n), n>=1, are record values of A003558.

Conjecture. All terms are primes except for a finite set of squares of primes. All terms from a(1) to a(5000) are primes except for a(21) = 121 = 11^2, supporting V. Shevelev's conjecture. [_John W. Layman_, Jul 22 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jul 18 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005660 | OEIS-A179581

Number of permutations of {1, ..., n+6} with deficiencies #{i : p(i) < i} exceeding exceedances #{i : p(i) > i} by n or more.

We have now a proof for the conjectured relations, cf. links. - _M. F. Hasler_, Jul 17 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 19 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005661 | OEIS-A179648

Expansion of (1/(1+4x-2x^2))*c(x/(1+4x-2x^2)), c(x) the g.f. of A000108.

Conjecture: (n+1)*a(n) +2*(2n+1)*a(n-1) +4*(1-n)*a(n-2) +4*(5-2n)*a(n-3) +4*(n-3)*a(n-4)=0. - _R. J. Mathar_, Nov 17 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005662 | OEIS-A179772

Position of ones in A179771.

This seems to give the positions where "L"'s occur in the central column of A122242 (please zoom into the illustration given here). Conjecture, from a(60)=167 onward, all integers >= 167 present.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005663 | OEIS-A179776

Quadrisection of the fourth central column of triangle A122245, a(n) = A179775((4*n)-2).

Conjecture: the last zero (108th) occurs at n=163, after which only ones occur.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005664 | OEIS-A179777

Position of ones in A179776.

This seems to give the positions where "L"'s occur in the central column of A122245 (please zoom into the illustration given here). Conjecture, from a(56)=164 onward, all integers >= 164 present.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005665 | OEIS-A179784

Records for minima of the positive distance d between the seventh power of a positive integer x and the square of an integer y such that d = x^7 - y^2 (x <> k^2 and y <> k^7).

Conjecture (_Artur Jasinski_): For any positive number x >= A179785(n), the distance d between the seventh power of x and the square of any y (such that x <> k^2 and y <> k^7) can't be less than A179784(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jul 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005666 | OEIS-A179790

Records for minima of the positive distance d between the ninth power of a positive integer x and the square of an integer y such that d = x^9 - y^2 (x <> k^2 and y <> k^9).

Conjecture (_Artur Jasinski_): For any positive number x >= A179791(n), the distance d between the ninth power of x and the square of any y (such that x <> k^2 and y <> k^9) can't be less than A179790(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jul 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005667 | OEIS-A179793

Records of minima of the positive distance d between the eleventh power of a positive integer x and the square of an integer y such that d = x^11 - y^2 (x <> k^2 and y <> k^11).

Conjecture (_Artur Jasinski_): For any positive number x >= A179794(n), the distance d between the eleventh power of x and the square of any y (such that x <> k^2 and y <> k^11) can't be less than A179793(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jul 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005668 | OEIS-A179795

Values y for records of the minima of the positive distance d between the eleventh power of a positive integer x and the square of an integer y such that d = x^11 - y^2 (x <> k^2 and y <> k^11).

Conjecture (_Artur Jasinski_):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jul 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005669 | OEIS-A179798

Record minima of the positive distance d between the 11th power of a positive integer x and the square of an integer y such that d = x^13 - y^2 (x <> k^2 and y <> k^13).

Conjecture (_Artur Jasinski_):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jul 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005670 | OEIS-A179799

Values x for records of minima of the positive distance d between an 11th power of a positive integer x and a square of an integer y such that d = x^13 - y^2 (x<>k^2 and y<>k^13).

Conjecture: For any positive number x >= A179799(n) the distance d between the 11th power of x and the square of any y (such that x<>k^2 and y<>k^13) can't be less than A179798(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jul 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005671 | OEIS-A179800

Values y for record minima of the positive distance d between the thirteenth power of a positive integer x and the square of an integer y such that d = x^13 - y^2 (x <> k^2 and y <> k^13).

Conjecture: For any positive number x >= A179799(n), the distance d between the 13th power of x and the square of any y (such that x <> k^2 and y <> k^13) can't be less than A179798(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jul 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005672 | OEIS-A179812

Record minima of the positive distance d between the fifteenth power of a positive integer x and the square of an integer y such that d = x^15 - y^2 (x <> k^2 and y <> k^15).

Conjecture: For any positive number x >= A179813(n), the distance d between the fifteenth power of x and the square of any y (such that x <> k^2 and y <> k^15) can't be less than A179812(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jul 28 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005673 | OEIS-A179831

Quadrisection of the fourth central column of triangle A122242, a(n) = A179830((4*n)+1).

Conjecture: the last zero (119th) occurs at n=164, after which only ones occur.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005674 | OEIS-A179832

Position of ones in A179831.

Look at the central column of A179757. Conjecture, from a(46)=165 onward, all integers >= 165 present.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005675 | OEIS-A179873

Corresponding values of antiharmonic means B(h) to numbers h from A179871 (numbers h such that antiharmonic mean of the numbers k < h such that gcd(k, h) = 1 is an integer).

Conjecture: nondecreasing sequence of odd numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jul 30 2010, Jul 31 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005676 | OEIS-A179874

Possible values of A179873(m) in increasing order, where A179873(m) = corresponding values of antiharmonic means to numbers from A179871 (numbers h such that antiharmonic mean of the numbers k < h such that gcd(k, h) = 1 is an integer).

Conjecture: a(n) = sequence of odd numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jul 30 2010, Jul 31 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005677 | OEIS-A179875

Numbers h such that h and h+1 have same antiharmonic mean of the numbers k < h such that gcd(k, h) = 1.

Conjecture: also numbers k such that mu(k) = 1 and mu(k+1) = -1, where mu is the Möbius function (tested on the first 10^4 terms). - _Amiram Eldar_, Mar 06 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jul 30 2010, Jul 31 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005678 | OEIS-A179892

Numbers which are not the sum of three distinct members of twin primes.

No odd numbers >=19 are in the sequence (conjecture?).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Jul 30 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005679 | OEIS-A179894

Given the series (1, 2, 1, 2, 1, 2, ...), let (1 + 2x + x^2 + 2x^3 + ...) * (1 + 2x^2 + x^3 + 2x^4 + ...) = (1 + 2x + 3x^2 + 7x^3 + ...)

The offset has been selected as "1" to accommodate the conjectured property of the sequence: 3 divides a(n) iff n == 0 mod 3. Example: 3 divides (3, 12, 15, 27, 27, 42, ...) but not other terms through n = 18.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Jul 31 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005680 | OEIS-A179902

Row sums of triangle A179901.

Conjectured mod 2 parity: (1, 0, 1, 1, 1, 0, 1, 1, 1, 0, 1, 1, 1,...).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Jul 31 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005681 | OEIS-A179929

a(n) = 2^n*A(n, -1/2), A(n, x) the Eulerian polynomials.

Conjecture: the sequence {a(n)} reduced modulo a positive integer k is eventually periodic with period a divisor of phi(k), where phi() = A000010(k). For example, modulo 11 the sequence becomes [1, 2, 2, 5, 3, 9, 2, 8, 5, 4, 0, 2, 2, 5, 3, 9, 2, 8, 5, 4, 0, 2, 2, 5, 3, 9, 2, 8, 5, 4, 0, ...], with an apparent period of 10 = phi(11) starting at n = 1. - _Peter Bala_, Feb 20 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Aug 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005682 | OEIS-A179967

Number of permutations p of [2*n+9] such that |p(i)-p(i-1)| > n for i>1.

Conjecture: a(n) = ((154924827300000*n+5704752193340625)/31250000000)*20^n - (3073238635/354294)*18^n - (13406475/235298)*14^n - (105/8192)*8^n for n > 5. Conjecture: G.f.: (362880 - 23733610*x + 574782136*x^2 - 6066574082*x^3 + 23063767496*x^4 + 1306804904*x^5 + 19195078240*x^6 + 99691196928*x^7 + 5798373293056*x^8 + 140546260769280*x^9 - 1838864889858048*x^10)/((1-8*x)*(1-14*x)*(1-18*x)*(1-20*x)^2). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005683 | OEIS-A179988

Smallest m such that n = sum of digits of A108971(m).

conjecture for n>=18: a(n) = A051885(n+1) + 7*(10^floor(n/9) - 1)/9 - 3*(n mod 2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Aug 09 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005684 | OEIS-A180000

a(n) = lcm{1,2,...,n} / swinging_factorial(n) = A003418(n) / A056040(n).

Asymptotics: log a(n) ~ n(1 - log 2). It is conjectured that log a(n) ~ n(1 - log 2) + O(n^{1/2+eps}) for all eps > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Aug 17 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005685 | OEIS-A180021

Second differences of sequence A180005.

It is conjectured that this sequence is bounded between 0 and 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Mario O. Bourgoin (mob(AT)brandeis.edu), Aug 06 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005686 | OEIS-A180044

Let the n-th Carmichael number A002997(n) = p1*p2*...*pr, where p1 < p2 < ... < pr are primes. Then a(n) = (p1-1) * (p1*p2*...*pr - 1)^(r-2) / ((p2-1)*...*(pr-1)).

The conjecture referred to in A162290 was generalized as follows: Let k be an r-factor Carmichael number (p_1 < p_2 < ... < p_r). Then (p_1-1)*(k-1)^(r-2)/((p_2-1)*(p_3-1)*...*(p_r-1)) is an integer. This was proved by _Max Alekseyev_ (see link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _A.K. Devaraj_, Aug 08 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005687 | OEIS-A180048

Coefficient triangle of the denominators of the (n-th convergents to) the continued fraction 1/(w+2/(w+3/(w+4/... . Conjectured to equal unsigned version of A137286.

Coefficient triangle of the denominators of the (n-th convergents to) the continued fraction 1/(w+2/(w+3/(w+4/... . Conjectured to equal unsigned version of A137286. Equivalence to the recurrence formula needs formal proof.

This continued fraction converges to 0.525135276160981... for w=1. A conjecture by Ramanujan puts this equal to -1 + 1/(sqrt(e*Pi/2) - Sum_{k>=1} 1/(2k-1)!!). p(0)=1; p(1)=w; p(n) = w*p(n-1) + n*p(n-2) (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_, Aug 08 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005688 | OEIS-A180049

Coefficient triangle of the numerators of the (n-th convergents to) the continued fraction 1/(w+2/(w+3/(w+4/... .

Equivalence to the recurrence formula needs formal proof. This continued fraction converges to 0.525135276160981... for w=1. A conjecture by Ramanujan puts this equal to -1 + 1/(sqrt(e*Pi/2) - Sum_{k>=1} 1/(2k-1)!!). Row sums equal A059480. b(0)=1; b(1)=w; b(n) = w*b(n-1) + (n+1)*b(n-2) (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_, Aug 08 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005689 | OEIS-A180101

a(0)=0, a(1)=1; thereafter a(n) = largest prime factor of sum of all previous terms.

For the purposes of this paragraph, regard 0 as the (-1)st prime and 1 as the 0th prime. Conjectures: All primes appear; the primes appear in increasing order; the k-th prime p(k) appears p(k+1)-p(k-1) times (cf. A031131); and p(k) appears for the first time at position A164653(k) (sums of two consecutive primes). These assertions are stated as conjectures only because I have not written out a formal proof, but they are surely true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005690 | OEIS-A180195

a(n) = (-1)^n*Sum((-1)^j*b(j), j=1..n), where b(n)=(n-1)!*(n^2 - n + 1) = A001564(n-1) (n>=1).

Conjecture D-finite with recurrence a(n) +(-n-1)*a(n-1) -4*a(n-2) +(n-2)*a(n-3)=0. - _R. J. Mathar_, Jul 24 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Sep 09 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005691 | OEIS-A180247

Prime Brier numbers: primes p such that for all k >= 1 the numbers p*2^k + 1 and p*2^k - 1 are composite.

It is a conjecture that every such number has more than 11 digits. In 2011 I have calculated that for any prime p < 10^11 there is a k such that either p*2^k + 1 or p*2^k - 1 has all its prime factors greater than 1321. - _Arkadiusz Wesolowski_, Feb 03 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Aug 19 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005692 | OEIS-A180248

Odd composite squarefree numbers k such that r = 2*(p - 2 + k/p)/(p-1) is an integer for each prime divisor p of k.

Conjecture: k is a Carmichael number (A002997) if and only if k is a term of this sequence and all r-values of k are even. With this new definition and Korselt's theorem it is easily shown that the proposed conjecture is true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | William F. Sindelar (w_sindelar(AT)juno.com), Aug 19 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005693 | OEIS-A180291

Number of arrangements of n indistinguishable balls in n boxes with the maximum number of balls in any box equal to n-1.

Conjecture: For n > 2, a(n) = 2*A000217(n+1). - _Bruce J. Nicholson_, Apr 05 2017 Clearly a(2)=1. Moreover, for n>2, a(n) = n*(n-1), since one can choose the box with n-1 balls in n ways, and the remaining ball can be put in one of the remaining n-1 boxes. So the conjecture above and the empirical formulas below are all correct. - _Luca Ferrigno_, Jul 13 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, formula from _Robert Gerbicz_ in the Sequence Fans Mailing List, Aug 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005694 | OEIS-A180306

a(n) is the largest integer k for which the Frobenius equation a_1*x_1 + a_2*x_2 + ... + a_n*x_n == k has no nonnegative integer solutions, where the a_i are consecutive primes beginning with the n-th prime.

a(n) <= A007414(n), so conjecturally a(n) ~ 3*prime(n). - _Charles R Greathouse IV_, Apr 03 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Aug 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005695 | OEIS-A180319

Total number of possible standard knight moves on an n X 2n chessboard, if the knight is placed anywhere.

Conjecture: a(n) = 4*(4-9*n+4*n^2) for n>1. a(n) = 3*a(n-1)-3*a(n-2)+a(n-3) for n>4. G.f.: 4*x^2*(2+7*x-x^2)/(1-x)^3. - _Colin Barker_, Mar 11 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | Graziano Aglietti (mg5055(AT)mclink.it), Aug 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005696 | OEIS-A180323

Positive numbers l of the form l = A007913(k^4 - 4*k*m^3), where 1 <= k <= 5*l, 1 <= |m| <= 5*l.

The equation x^3 + y^3 + z^3 = 0 is solvable in numbers of the form N + M*sqrt(a(n)), where M and N are integers. Moreover, it is solvable in numbers of the form N + M*sqrt(l), where l has the form l = A007913(k^4 - 4*k*m^3), where k,|m| >= 1 (without restrictions k,|m| <= 5*l). But in this more general case there could be unknown numbers l having this form; this circumstance does not allow construction of the full sequence of such l. Therefore we restrict ourselves by the condition k,|m| <= 5*l. Note that testing l with respect to this condition is rather simple by sorting all values of k,|m| <= l. One can prove that, at least, if the Fermat numbers (A000215) are squarefree, then the sequence is infinite. Conjecture (necessity of the form of l): If the equation x^3 + y^3 + z^3 = 0 is solvable in numbers of the form N + M*sqrt(l) with integer N,M, then there exist positive integers k,m such that l = A007913(k^4 - 4*k*m^3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Aug 28 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005697 | OEIS-A180336

Positive numbers l of the form l = A007913(4*k*m^3 - k^4), where 1 <= k <= 5*l, 1 <= m <= 5*l.

The equation x^3 + y^3 + z^3 = 0 is solvable in numbers of the form N + M*sqrt(-a(n)), where M and N are integers. Moreover, it is solvable in numbers of the form N + M*sqrt(-l), where l>0 has the form l = A007913(4*k*m^3 - k^4), where k,m >= 1 (without restrictions k,m <= 5*l). But in this more general case there could be unknown numbers l having this form; this circumstance does not allow construction of the full sequence of such l. Therefore we restrict ourselves by the condition k,m <= 5*l. Note that testing l with respect to this condition is rather simple by sorting all values of k,m <= l. One can prove that, at least, in case the Fermat numbers (A000215) are squarefree, the sequence is infinite. Conjecture (necessity of the form of l): If the equation x^3 + y^3 + z^3 = 0 is solvable in numbers of the form N + M*sqrt(-l) with integer N,M, then there exist positive integers k,m such that l = A007913(4*k*m^3 - k^4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Aug 28 2010, Aug 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005698 | OEIS-A180337

Numbers which cannot be expressed as a sum 1 + p1 + p1*p2 + p1*p2*p3 + ... for some collection of primes {p1, p2, p3, ...}.

Conjecture: This sequence is finite with only the terms given above. It can be constructed by taking a1=2, and adjoining all numbers aj*ak + 1, where aj and ak are two prime members of the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jack W Grahl_, Aug 28 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005699 | OEIS-A180346

Primes that divide every circular permutation of the digits of at least one number of the form 123...(k-1)(k) (see A007908), where k is 3 digits long (that is, for some k in the range 99 < k < 1000).

a(28) > prime(2.3316*10^9) if it exists. Conjecture: 18973 is the last term. - _Chai Wah Wu_, Oct 09 2023 Conjecture verified: there are no other terms. - _Robert Israel_, Oct 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Jan 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005700 | OEIS-A180364

a(n) = sum_{k=0..n} C(n,k)*C(n+k,k)*(2*k+1)^2, where C(m,k) denotes the binomial coefficient m!/(k!*(m-k)!).

Conjecture: The sequence a(n+1)/a(n) (n = 0,1,...) is strictly decreasing to the limit 3+2*sqrt(2), and the sequence a(n+1)^(1/(n+1))/a(n)^(1/n) (n = 1,2,3,..) is strictly increasing to the limit 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005701 | OEIS-A180402

a(n) = lcm(1,...,Fibonacci(n)).

Also least period for number of ways of placing k non-attacking queens on an n X n chessboard. [conjectured by Kotesovec; proved for n <= 5. - _Thomas Zaslavsky_, Jun 24 2018]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Sep 02 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005702 | OEIS-A180442

Numbers k such that a sum of two or more consecutive squares beginning with k^2 is a square.

It is conjectured that the sequence continues 103, 104, 106, 112, 115, 117, 119, 121, 124, 128, 129, 131, 132, 137, 140, 142, 146, 158, 168, 170, 172, 175, 178, 181, 183, 192, 193, 197, 199, 200. - _Jean-François Alcover_, Sep 17 2013. Conjecture confirmed (see the Schoenfield link below). - _Jon E. Schoenfield_, Nov 22 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhining Yang_, Jan 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005703 | OEIS-A180460

a(n) is the smallest number m such that sigma(m)=22^n, or 0 if m does not exist.

Conjecture: Given any even integer E not a power of 2 (see A078426) there exists a positive integer N such that for all n>=N the equation sigma(m)=E^n has at least one solution for m.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Walter Kehowski_, Sep 06 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005704 | OEIS-A180476

Smallest k>0 such that q=p+6k, 6kp+q, 6kp-q, 6kq+p and 6kq-p are simultaneously prime, or 0 if no such k exists, where p=A000040(n) is the n-th prime.

It is conjectured that such a k>0 does exist for all primes > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005705 | OEIS-A180481

The smallest prime q > p = prime(n) such that p*(q-p)+q, p*(q-p)-q, q*(q-p)+p and q*(q-p)-p are simultaneously prime, or 0 if no such q exists.

It is conjectured that a(n) > 0 for all n, and for infinitely many terms, a(n) = prime(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005706 | OEIS-A180486

Numbers of the form ceiling(A179896(j)/A018252(j)) where A179896(j) mod A018252(j) <> 0.

Conjectures from _Colin Barker_, Nov 25 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Odimar Fabeny_, Sep 07 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005707 | OEIS-A180632

Minimum length of a string over the alphabet A = {1,2,...,n} that contains every permutation of A as a substring exactly once, also known as length of the minimal super-permutation.

Was conjectured to be equal to A007489, but it is now known that a(n) < A007489(n) for all n > 5. - _Robin Houston_, Aug 22 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Hamm_, Sep 13 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005708 | OEIS-A180638

Phan Thành Nam's upper bound on the number of non-relativistic electrons bound to a nucleus of charge n.

Note that the published version of the paper, as revisions v2 and v3 on the arXiv, use the stronger bound 1.22Z + 3Z^(1/3) instead, where the constant 1.22 can be improved to 1.21684.... In particular this improves the bound for lithium, aluminium, scandium, chromium, cobalt, zinc, arsenic, krypton, yttrium, zirconium, etc. Asymptotically Seco, Sigal, & Solovej show that the maximum ionization is n + O(n^(5/7)) and it is conjectured that the maximum ionization is n + O(1). - _Charles R Greathouse IV_, Sep 13 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Sep 14 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005709 | OEIS-A180710

a(n) = sigma(2^(n-1)*a(n-1)) for n>1 with a(1)=1.

Conjecture: sigma(n) | a(n) for n>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 18 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005710 | OEIS-A180871

Index of term in Sylvester's sequence A000058 divisible by prime A007996(n).

An open problem is to prove that all terms of Sylvester's sequence are squarefree or to find a counterexample. Using the p from A007996 and k found here, it is simple to determine whether A000058(k) = 0 (mod p^2). No p < 10^10 was found to have this property.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Sep 25 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005711 | OEIS-A180921

a(n) is the square root of the sum of the cubes of the b(n) consecutive integers starting from b(n), where b(n) = A180920.

Colin Barker's linear recurrence conjecture confirmed, see A180920. - _Ray Chandler_, Jan 12 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Pletser_, Sep 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005712 | OEIS-A180987

T(n,k)=number of nXk binary matrices with rows in lexicographically nonincreasing order and columns in lexicographically strictly increasing order.

Conjecture: T(n,k) = A180986(n-k+1,k), k<=n . [From _R. J. Mathar_, Oct 18 2010]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 30 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005713 | OEIS-A181018

Maximum number of 1's in an n X n binary matrix with no three 1's adjacent in a line along a row, column or diagonally.

Three or more "1"s may be adjacent in an L-shape or step shape (cf. bottom of first example) or 2 X 2 square (top right of 2nd example) or similar. One possible (not always optimal) solution is therefore to fill the square with 2 X 2 squares of "1"s, separated by rows of "0"s: this yields the lower bound (n - floor(n/3))^2 = ceiling(2n/3)^2 given in FORMULA. I conjecture that this is optimal for n = 2 (mod 3) and that a(n) ~ (2n/3)^2. For n = 3k, the array can be filled with 2k(2k+1) "1"s by repeating the optimal solution for n = 3 on the diagonal, and filling the rest with 2 X 2 blocks separated by rows of "0"s, cf. the 4th example for 6 X 6. - _M. F. Hasler_, Jul 17 2015 [Conjecture proved to be wrong, see below. - _M. F. Hasler_, Jan 19 2016]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 30 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005714 | OEIS-A181030

Duplicate of A006506.

Conjecture: a(n) = A006506(n-1), n>1. - _R. J. Mathar_, Oct 02 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005715 | OEIS-A181104

Dirichlet inverse of Ramanujan's L-series (A000594).

Although it is conjectured that A000594(n) is never 0 here a(n)=0 for infinitely many n. Namely a(n)=0 iff n is not cubefree (n is in A046099).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 03 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005716 | OEIS-A181142

Number of permutations of {1,2,...,2n} , say x(1),x(2), ... , x(2n) , such that x(i) + x(i+1) is not equal to 2n-1 for all i, 1<=i<=2n-1.

Conjecture: D-finite with recurrence (-n+1)*a(n) +2*(2*n-1)*(n^2-n-1)*a(n-1) +4*n^2*(n-2)*a(n-2)=0. - _R. J. Mathar_, Jan 27 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | Arin Chaudhuri (arin.chaudhuri(AT)gmail.com), Oct 06 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005717 | OEIS-A181147

a(n) = Sum_{k=0..n-1} (105k+44)*C(2k,k)^2*T(k)*(-1)^(n-1-k)/(2n*C(2n,n)), where T(k) (k=0,1,2,...) are central trinomial coefficients given by A002426.

On Jan 22 2011, _Zhi-Wei Sun_ conjectured that a(n) is a positive integer for every n=1,2,3,... Let p > 3 be a prime. He also conjectured that a(p) == 5 + 6*(p/3)*(2-3^(p-1)) (mod p^2). Another conjecture of his states that Sum_{k=0..p-1} (-1)^k*binomial(2k,k)^2*T(k) is congruent to b(p) modulo p^2, where b(p)=0 if (p/15)=-1, b(p) = 4x^2-2p if p == 1,4 (mod 15) and p = x^2+15y^2 with x,y integers, and b(p) = 20x^2-2p if p == 2,8 (mod 15) and p=5x^2+3y^2 with x,y integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005718 | OEIS-A181156

Odd Fibonacci numbers F which have a proper Fibonacci divisor G such that F/G is a Lucas number or a product of Lucas numbers.

A conjectural statement that, for odd prime p, the ratio F_{p^2}/F_{p} is never a Lucas number or a product of some Lucas numbers, yields that

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Oct 07 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005719 | OEIS-A181192

Number of n X 5 matrices containing a permutation of 1..n*5 in increasing order rowwise, columnwise, diagonally and (downwards) antidiagonally.

Conjectures from _Colin Barker_, Feb 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 10 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005720 | OEIS-A181198

Number of 4 X n matrices containing a permutation of 1..4*n in increasing order rowwise, columnwise, diagonally and (downwards) antidiagonally.

Conjectured recurrence of order 2 and degree 9: 3*(n + 1)*(2*n + 3)*(3*n + 4)*(3*n + 5)*(7*n^2 - 1)*(n + 2)^3*a(n + 2) - 8*(n + 1)*(2*n + 1)*(4*n + 3)*(4*n + 5)*(364*n^5 + 84*n^4 - 1025*n^3 - 534*n^2 + 157*n + 54)*a(n + 1) - 64*(2*n - 1)^2*(2*n + 1)*(4*n - 1)*(4*n + 1)*(4*n + 3)*(4*n + 5)*(7*n^2 + 14*n + 6)*a(n) = 0. - _Christoph Koutschan_, Feb 26 2023 Conjectured formula, solution to the above recurrence, for n > 1: a(n) = (-64)^n * (n-1) * (-1/2)_{2*n} * (1/2)_{n} / (4*(3*n)!) * (-1 + 3*Sum_{k=2..n-1} (-4)^k * (7*k^2-1) / ((k-1) * k * (k+1)^2 * (2*k-1)^2 * (2*k+1)^3) * binomial(3*k,2*k) * binomial(k+1/2,k)), where (a)_{n} is the Pochhammer symbol.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 10 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005721 | OEIS-A181199

Number of 5 X n matrices containing a permutation of 1..5*n in increasing order rowwise, columnwise, diagonally and (downwards) antidiagonally.

Conjectured recurrence of order 3 and degree 24: 3*(n + 2)^2*(2*n + 3)*(2*n + 5)^2*(3*n + 7)*(3*n + 8)*(4*n + 9)*(4*n + 11)*(137855872*n^11 + 860969696*n^10 + 2047036856*n^9 + 2032587274*n^8 - 24192441*n^7 - 1894061166*n^6 - 1671661480*n^5 - 524330624*n^4 + 36004789*n^3 + 62751860*n^2 + 13865604*n + 927360)*(n + 3)^4*a(n + 3) - (n + 2)^2*(2*n + 3)*(717088749346816*n^21 + 14962008250398720*n^20 + 139665528127686656*n^19 + 760770273998231808*n^18 + 2616700630350746208*n^17 + 5556799141672247640*n^16 + 5462710847171622988*n^15 - 608017104359154861O*n^14 - 3177189218448699433 3*n^13 - 5479130874518334030 9*n^12 - 4663430896095769856 7*n^11 - 1215620047630796049*n^10 + 4883346077919144976 3*n^9 + 6569291812211075457 3*n^8 + 4709707908384811651 1*n^7 + 1962810609828790919 1*n^6 + 3499276436019940446*n^5 - 8502592830253275 24*n^4 - 6858020531017172 88*n^3 - 1800890418886574 40*n^2 - 226039772714112 00*n - 1104708217536000)*a(n + 2) - 15*(2*n + 1)*(5*n + 6)*(5*n + 7)*(5*n + 8)*(5*n + 9)*(9113927409664*n^19 + 166164038596608*n^18 + 1357324488921088*n^17 + 6541981632511232*n^16 + 20586994844739808*n^15 + 44061606220301336*n^14 + 64302800289940820*n^13 + 61042725728660150*n^12 + 30740208891967721*n^11 - 3351043407516635*n^10 - 17410620603191989*n^9 - 12634296322883951*n^8 - 4487751704725767*n^7 - 793918042456325*n^6 - 54713722756179*n^5 + 42448206362469*n^4 + 50327965592874*n^3 + 21777752002716*n^2 + 4014999151560*n + 257403484800)*a(n + 1) + 25*(2*n - 1)^2*(2*n + 1)*(4*n - 1)*(4*n + 1)*(5*n + 1)*(5*n + 2)*(5*n + 3)*(5*n + 4)*(5*n + 6)*(5*n + 7)*(5*n + 8)*(5*n + 9)*(137855872*n^11 + 2377384288*n^10 + 18238806776*n^9 + 81945774178*n^8 + 238738633847*n^7 + 471293180347*n^6 + 638861237719*n^5 + 588363536007*n^4 + 354332674386*n^3 + 128320688700*n^2 + 23068181160*n + 1077753600)*a(n) = 0. - _Christoph Koutschan_, Feb 27 2023 Conjecture: a(n) ~ 9 * 5^(5*n + 1/2) / (2^17 * Pi^2 * n^12), based on the recurrence by _Christoph Koutschan_. - _Vaclav Kotesovec_, Feb 27 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 10 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005722 | OEIS-A181270

Number of 2 X n binary matrices M with rows in strictly increasing order and rows of M*Mtranspose (mod 2) in strictly increasing order.

Conjectures from _Colin Barker_, Mar 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 10 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005723 | OEIS-A181271

Number of 3 X n binary matrices M with rows in strictly increasing order and rows of M*Mtranspose (mod 2) in strictly increasing order.

Conjectures from _Colin Barker_, Mar 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 10 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005724 | OEIS-A181278

Number of 2 X n binary matrices M with rows in strictly increasing order and rows of M*Mtranspose (mod 2) in strictly decreasing order.

Conjectures from _Colin Barker_, Mar 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 10 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005725 | OEIS-A181279

Number of 3 X n binary matrices M with rows in strictly increasing order and rows of M*Mtranspose (mod 2) in strictly decreasing order.

Conjectures from _Colin Barker_, Mar 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 10 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005726 | OEIS-A181280

Number of 4Xn binary matrices M with rows in strictly increasing order and rows of M*Mtranspose (mod 2) in strictly decreasing order.

a(n) = (1/3)*2^(2*n-11)*(6*n^2-219*n+820) - (1/9)*2^(n-5)*(3*n+32) - (113/3)*(-1)^n*2^(3*n-14) + 2^(4*n-9) - (1/3)*(-1)^n*2^(2*n-11)*(13*n-164) +(1/9)*2^(3*n-14)*(288*n-3473) (conjectured). - _Manuel Kauers_ and _Christoph Koutschan_, Mar 02 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 10 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005727 | OEIS-A181311

Highly abundant numbers k whose largest prime factor is less than log(k).

Every highly abundant number (A002093) is either in this sequence or A181312. Although it appears that for every n there is at least one prime p such that n*p is another highly abundant number, this conjecture fails for 46846800, the 227th term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Oct 13 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005728 | OEIS-A181312

Highly abundant numbers k whose largest prime factor is greater than log(k).

Every highly abundant number (A002093) is either in this sequence or A181311. Although it appears that dividing any term by its largest prime factor produces another highly abundant number, this conjecture fails for 2942007868800, the 527th term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Oct 13 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005729 | OEIS-A181373

Least m>0 such that prime(n) divides S(m)=A007908(m)=123...m and all numbers obtained by cyclic permutations of its digits; 0 if no such m exists.

Conjecture: a(n) > 0 for n <> 1 and n <> 3. - _Chai Wah Wu_, Oct 06 2023 The conjecture that a(n) > 0 for all n != 1, 3 is proved in Wang (2026). - _Qichao Wang_, Oct 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_ and _Marco Ripà_, Jan 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005730 | OEIS-A181381

Total number of positive integers below 10^n requiring 5 positive cubes in their representation as sum of cubes.

Deshouillers et al. (2000) conjectured that there are exactly 113936676 positive integers which are not the sum of four nonnegative integral cubes, and that the largest is 7373170279850. Combined with earlier proofs and conjectures, this would imply a(n) = 113932616 for n >= 13. - _Martin Fuller_, Oct 11 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Martin Renner_, Jan 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005731 | OEIS-A181395

Summed lengths of nonintersecting rook paths on a 4 X n board.

Conjectured g.f.: x*(1 - x)*(3 - 29*x + 51*x^2 + 595*x^3 - 3879*x^4 + 9553*x^5 - 8366*x^6 - 8026*x^7 + 22931*x^8 - 13117*x^9 - 5593*x^10 + 7955*x^11 - 6118*x^12 + 6842*x^13 + 1884*x^14 - 6824*x^15 + 519*x^16 + 1991*x^17 - 206*x^18 - 230*x^19 + 13*x^20 + 9*x^21)/((1 - 8*x + 15*x^2 - 5*x^3 - 9*x^4 + 2*x^5 + x^6)^2*(1 - 4*x + 7*x^2 - 3*x^3 - 7*x^4 + 2*x^5 + x^6)^2). - _Andrew Howroyd_, Jan 06 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Scambler_, Oct 17 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005732 | OEIS-A181402

Total number of positive integers below 10^n requiring 7 positive cubes in their representation as sum of cubes.

Conjectured g.f.: x*(1+9*x+63*x^2+48*x^3)/(1-x). - _Colin Barker_, May 04 2012 Conjectured e.g.f.: 121*(exp(x) - 1) - 120*x - 111*x^2/2 - 8*x^3. - _Stefano Spezia_, May 21 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Martin Renner_, Jan 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005733 | OEIS-A181411

a(n) = Sum_{k=0..n} C(n,k)*sigma(n+k) for n>=1.

Conjecture: a(n) ~ c * n * 2^n, where c = Pi^2/4 = A091476. - _Vaclav Kotesovec_, Oct 05 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 19 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005734 | OEIS-A181477

a(n) has generating function 1/((1-x)^k*(1-x^2)^(k*(k-1)/2)) for k=5.

a(n-1,k) is conjectured to also be the count of monomials (or terms) in the Schur polynomials of k variables and degree n, summed over all partitions of n in at most k parts (zero-padded to length k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_, Oct 24 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005735 | OEIS-A181487

Numbers k such that Sum_{d|k, d<k, d not occurring before} d > k.

De Koninck and Ivić conjectured that this sequence has an asymptotic density.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Oct 28 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005736 | OEIS-A181547

a(n) = Sum_{k=0..floor(n/2)} C(n-k,k)^5.

Conjecture: Given F(n,L) = Sum_{k=0..[n/2]} C(n-k,k)^L, then lim_{n->oo} F(n+1,L)/F(n,L) = (Fibonacci(L)*sqrt(5) + Lucas(L))/2 for L>=0 where Fibonacci(n) = A000045(n) and Lucas(n) = A000032(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 29 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005737 | OEIS-A181553

Coefficient of x^n in (x^2 + 98*x + 1)^n.

On Jan 29 2011, _Zhi-Wei Sun_ conjectured that Sum_{k>=0} (40k + 3)*a(k)*binomial(4k, 2k)*binomial(2k, k)/112^(2k) = 70*sqrt(21)/(9*Pi). He also conjectured that 2n(2n + 1)*binomial(2n, n) divides Sum_{k=0..n-1} (40k + 3)*a(k)*binomial(4k, 2k)*binomial(2k, k)112^{2(n - 1 - k)} for each n = 2, 3, 4, ... and that for any prime p different from 2 and 7 we have the congruence Sum_{k=0..p-1} (40k + 3)*a(k)*binomial(4k, 2k)*binomial(2k, k)/112^(2k) == p(-21/p)(5 - 2(-2/p)) (mod p^2). Another conjecture of his states that for any prime p == 1, 3 (mod 8) with p = x^2 + 2y^2 (x, y integers) we have Sum_{k=0..p-1} a(k)*binomial(4k, 2k)*binomial(2k, k)/112^(2k) == (7/p)(4x^2 - 2p) (mod p^2). By a result of Sun, Sum_{k=0..p-1} (40k + 3)*a(k)*binomial(4k, 2k)*binomial(2k, k)/112^(2k) == 0 (mod p^2) for any prime p = 5,7 (mod 8).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005738 | OEIS-A181564

Semiprime convenient numbers.

This is to semiprimes A001358 as A140768 is to primes A000040. Because Euler's "numerus idoneus" (idoneal, or suitable, or convenient numbers) are finite, this subset is finite. Conjecture: this list is complete (see Weinberger's result and comment by _Jud McCranie_ in A140768).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jan 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005739 | OEIS-A181630

Duplicate of A112310.

Conjectured row sums = A023610: (1, 3, 7, 15, 30, 58, ...).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Nov 02 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005740 | OEIS-A181674

Number of partitions of n into a primorial and a semiprime.

Conjecture: Average order is Theta(log log n). - _Charles R Greathouse IV_, May 25 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Nov 04 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005741 | OEIS-A181717

Fibonacci-Collatz sequence: a(1)=0, a(2)=1; for n>2, let fib=a(n-1)+a(n-2); if fib is odd then a(n)=3*fib+1 else a(n)=fib/2.

It is easy to prove that all the terms a(n) with n>=7 are congruent to 7 mod 9. Conjecture: for every k>0 there is an index m such that all the a(n) with n>m have the same residue mod 3^k. - _Giovanni Resta_, Nov 17 2010

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Nov 17 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005742 | OEIS-A181776

a(n) = lambda(lambda(n)), where lambda(n) is the Carmichael lambda function (A002322).

Harland proves the conjecture of Martin & Pomerance that a(n) = n exp ((1 + o(1))(log log n)^2 log log log n) for almost all n, as well as a generalization to k-th iterates. - _Charles R Greathouse IV_, Dec 21 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Nov 12 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005743 | OEIS-A181777

Number of steps to reach 1 in '3x+1' (or Collatz) problem starting with the n-th Mersenne prime.

It is conjectured by Ohira and Watanabe that for large Mersenne primes 2^k-1, the fraction steps/k ~ 2+3*log(3)/log(4/3) or approximately 13.45.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Dec 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005744 | OEIS-A181830

The number of positive integers <= n that are strongly prime to n.

It is conjectured (see Scroggs link) that a(n) is also the number of cardboard braids that work with n slots. - _Matthew Scroggs_, Sep 23 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Nov 17 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005745 | OEIS-A181901

a(n) = least positive integer m such that 2(s_k)^2 for k=1,...,n are pairwise distinct modulo m where s_k = Sum_{j=1..k} (-1)^(k-j)*p_j and p_j is the j-th prime.

On Mar 28 2012, _Zhi-Wei Sun_ conjectured that a(n) is the (n+1)-th prime p_{n+1} with the only exceptions being a(1)=1, a(2)=4, a(4)=9 and a(9)=25. He has shown that 2(s_k)^2 (k=1,...,n) are indeed pairwise distinct modulo p_{n+1} and hence a(n) does not exceed p_{n+1}. The conjecture was verified for n up to 2*10^5 by the author in 2018, and for n up to 3*10^5 by Chang Zhang (a student at Nanjing Univ.) in June 2020. - _Zhi-Wei Sun_, Jun 22 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 31 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005746 | OEIS-A181905

Expansion of (b(q^3)^3 - b(q)^3) / 9 in powers of q where b() is a cubic AGM theta function.

Conjecture: multiplicative with a(3^e) = 0, a(p^e) = ((p^2)^(e+1)-1)/(p^2-1) for p == 1 (mod 3), a(p^e) = (1-(-p^2)^(e+1))/(p^2+1) for p == 2 (mod 3). - _Andrew Howroyd_, Aug 05 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Apr 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005747 | OEIS-A181933

a(n) = Sum_{k=0..n} binomial(n+k,k)*sin(Pi*(n+k)/2).

Conjecture: +2*n*a(n) +8*n*a(n-1) +(-n+20)*a(n-2) +5*(-n+4)*a(n-3) +2*(-2*n+5)*a(n-4)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Apr 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005748 | OEIS-A181996

Triangle of Ward numbers T(n,k) (n>=0, k=0 if n=0, otherwise 0 <= k <= n-1) read by rows.

Conjecture on row sums was proved in the first paragraph of the formula section of the reverse matrix A134991 in 2008 (e.g.f. evaluated at t=1). - _Tom Copeland_, Jan 03 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005749 | OEIS-A182003

a(n) = least prime p_k such that n = p_k - p_{k-1} + ... + (-1)^{k-j}p_j for some 1 <= j <= k, with p_k the k-th prime.

a(n) is obviously at least n. In March 2012, Zhi-Wei Sun conjectured that a(n) always exists and does not exceed 2.2*n for n > 1; he even thought that 2.2*n might be replaced by 2*n + 2.5*sqrt(n) for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005750 | OEIS-A182022

Let p = n-th prime; a(n) = conjectured prime q such that the sequence of primes [p,...,q] has maximal discrepancy.

Let p = n-th prime; a(n) = conjectured prime q such that the sequence of primes [p,...,q] has maximal discrepancy.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 06 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005751 | OEIS-A182025

a(n) = 31*binomial(2*n,n-4) + Sum_{i=1..n-4} binomial(2*n,n-4-i)*(4+i).

Conjecture: 558*(n-4)*(n+4)*a(n) +7*(-631*n^2+777*n+4600)*a(n-1) +4370*(n-1)*(2*n-3)*a(n-2)=0. - _R. J. Mathar_, Aug 08 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 06 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005752 | OEIS-A182049

Numbers m such that m! is not pandigital in decimal representation.

Conjecture: sequence is finite and a(32) = 41 is the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005753 | OEIS-A182079

a(n) = floor(n*floor((n-1)/2)/3).

The conjectured g.f. is correct. See links. - _Sela Fried_, Dec 27 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005754 | OEIS-A182080

a(n) is the maximal depth of an indecomposable exact cover of an n-set.

The values of a(6), a(7), a(8) shown here were only conjectural, but that may have changed since Graver's paper is now nearly 40 years old.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005755 | OEIS-A182084

a(n) = 3*n - n/p, where p is the smallest prime dividing n.

Conjectured to be minimal number of nodes in any non-bipartite regular graph of degree n, diameter 2 and girth 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005756 | OEIS-A182085

Carmichael numbers of the form (30k+7)*(60k+13)*(150k+31).

Conjecture: N = (30k+7)*(60k+13)*(150k+31) is a Carmichael number if (but not only if) 30k+7, 60k+13 and 150k+31 are all three prime numbers. We checked the conjecture up to k = 256; we got Carmichael numbers with three prime divisors for k = 0, 1, 10, 12, 18, 24, 32, 43, 85, 102, 123, 129, 150, 201, 207, 256. The conjecture follows from Korselt's criterion. - _Charles R Greathouse IV_, Oct 02 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Apr 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005757 | OEIS-A182088

Carmichael numbers of the form C = (30n-29)*(60n-59)*(90n-89)*(180n-179), where n is a natural number.

Conjecture: The number C = (30n-29)*(60n-59)*(90n-89)*(180n-179) is a Carmichael number if (but not only if) 30n-29, 60n-59, 90n-89 and 180n-179 are all four prime numbers. The conjecture is checked for 0<n<150; the condition is satisfied just for n = 10, 52, 77, 143.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Apr 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005758 | OEIS-A182089

Numbers of the form (330*k+7)*(660*k+13)*(990*k+19)*(1980*k+37).

Conjecture: C = (330k+7)*(660k+13)*(990k+19)*(1980k+37) is a Carmichael number if 330k+7, 660k+13, 990k+19 and 1980k+37 are all four prime numbers. [The conjecture is true, and can be proved using Korselt's criterion. - _Amiram Eldar_, Apr 24 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Apr 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005759 | OEIS-A182090

Carmichael numbers divisible by 11.

Conjecture: Any Carmichael number C divisible by 11 and not divisible by 3 can be written in one of the following four forms: C = 990k+11; C = 990k+11^2; C = 990k+11*41 or C = 990k+11*71, where k is a natural number. For comparison, we issue the following related conjectures: The first counterexample to the conjecture is 6049681 which is 781 mod 990. 341 and 671 mod 990 also appear. Conjecture 1 is true; the only excluded possibility is 161 mod 210 which is never a 2-pseudoprime (check mod 7). All three parts of conjecture 2 are false; counterexamples include 461502097, 1378483393, and 7044493729. Conjecture 3 is true, consider Korselt's criterion mod 30. - _Charles R Greathouse IV_, Dec 07 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Apr 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005760 | OEIS-A182097

Expansion of 1/(1-x^2-x^3).

Conjectured values of d(n), the dimension of a Z-module in MZV(conv). See the Waldschmidt link. - _Michael Somos_, Mar 14 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005761 | OEIS-A182122

Expansion of exp(arcsinh(2*x)).

Conjecture : n*(n+1)*a(n) + (n+2)*(n-1)*a(n-1) +4*(n+1)*(n-3)*a(n-2) +4*(n+2)*(n-4)*a(n-3) = 0.- _R. J. Mathar_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Apr 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005762 | OEIS-A182129

Number of iterations of the orbit n -> (sum of the decimal digits of n)^n starting with n, needed to stabilize.

Generalization and conjecture: Let k be a positive integer. The number of iterations of the orbit k -> (sum of the decimal digits of k)^n is finite for any exponent n and any starting value k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005763 | OEIS-A182132

Carmichael numbers of the form C = (30n-7)*(90n-23)*(300n-79).

Conjecture: The number C = (30n-7)*(90n-23)*(300n-79) is a Carmichael number if (but not only if) 30n-7, 90n-23 and 300n-79 are all three prime numbers. The conjecture is checked for 0<n<130; we got Carmichael numbers with three prime divisors for n = 2, 9, 15, 32, 43, 48, 58, 64, 67, 78, 97, 128.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Apr 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005764 | OEIS-A182133

Carmichael numbers of the form C = (30n-17)*(90n-53)*(150n-89).

Conjecture: The number C = (30n+13)*(90n+37)*(150n+61) is a Carmichael number if (but not only if) 30n+13, 90n+37 and 150n+61 are all three prime numbers. The conjecture is checked for 0<n<130; the condition is satisfied for n = 0, 1, 5, 12, 14, 12, 27, 28, 49, 55, 56, 71, 83, 121, 125. The conjecture is true (follows from Korselt's criterion). - _Charles R Greathouse IV_, Jul 05 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Apr 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005765 | OEIS-A182142

Abundance d = sigma(N) - 2*N = A033880(N) of numbers N = A153501(n), i.e., N has d > 0 as divisor.

It is conjectured that only powers of 2 can occur more than once.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005766 | OEIS-A182145

XOR(d1,d2), where d1 and d2 are the difference between consecutive primes given in binary.

Conjecture: among values less than 60 the most frequent value is 30 (based on frequencies for n < 2126795281, i.e. primes less than 50176000000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Apr 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005767 | OEIS-A182151

Carmichael numbers divisible by 31.

Conjecture: Any Carmichael number C divisible by 31 can be written in one of the following three forms: C = 2790n+31; C = 2790n+31^2 or C = 2790n+31*61, where n is natural. It follows from Korselt's criterion that such numbers are 31 mod 930, the union of the three residue classes. Thus the conjecture is true. - _Charles R Greathouse IV_, Oct 02 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Apr 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005768 | OEIS-A182160

Number of iterations of the map n -> sum of the n-powers of the decimal digits of n.

Generalization and conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005769 | OEIS-A182172

Number A(n,k) of standard Young tableaux of n cells and height <= k; square array A(n,k), n>=0, k>=0, read by antidiagonals.

Conjecture: A(n,k) ~ k^n/Pi^(k/2) * (k/n)^(k*(k-1)/4) * Product_{j=1..k} Gamma(j/2). - _Vaclav Kotesovec_, Sep 12 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Apr 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005770 | OEIS-A182187

a(n) is the least m >= n such that the Hamming distance D(n,m) = 2.

Using this formula, we can prove the conjecture formulated in comment in A209554 in the case k=2. Moreover, let us show that if N does not have the form 8*t or 8*t+1, then it can be represented in the form n<+>2. Indeed, in the cases N = 8*t+2, 8*t+4, 8*t+6, 8*t+3, 8*t+5 and 8*t+7 it is sufficient to choose n=N-4, n=N-2, n=N-1, n=N-3, n=N-2 and n = N-3 respectively; in the cases 8*t, 8*t+1, for every choice of n <= N, we do not obtain the equality n<+>2 = N.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005771 | OEIS-A182205

Iterate the map in A006368 starting at 40.

Like for iterations with starting points 8 or 14, it is conjectured that also this trajectory never repeats.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005772 | OEIS-A182206

Carmichael numbers of the form C = 37*73*(18n+91).

Conjecture: Any Carmichael number C divisible by 37 and 73 can be written as C = 37*73*(18n+91), where n is natural; checked for the first 22 Carmichael numbers divisible by 37 and 73.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Apr 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005773 | OEIS-A182208

Carmichael numbers divisible by 7.

Conjecture: Any Carmichael number C divisible by 7 can be written in one of two ways: (1) C=7*(6m+1)*(6n+1), where m and n are natural numbers or (2) C=7*(6m-1)*(6n-1), where m and n are natural numbers. In other words, there aren't Carmichael numbers divisible by 7 of the form C=7*(6m+1)*(6n-1). Checked for the first 27 Carmichael numbers divisible by 7. Note: a Carmichael number with more than 3 prime divisors can be written (sometimes) in both ways: 41041 = 7*11*13*41 = 7*13*451 (form 1) = 7*11*533 = 7*41*143 (form 2). The conjecture follows from Korselt's criterion. - _Charles R Greathouse IV_, Oct 02 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Apr 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005774 | OEIS-A182209

a(n) is the least m >= n, such that the Hamming distance D(n,m) = 3.

Using this formula, we can prove conjecture formulated in comment in A209554 in case k=3. Moreover, one can prove that N could be represented in form n<+>2 or n<+>3 iff N is not a number of the forms 32*t, 32*t+1. - _Vladimir Shevelev_, Apr 25 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005775 | OEIS-A182220

Largest number k such that there exists an extensional acyclic digraph on n labeled nodes with k sources.

It appears a(n+1) is the number of distinct possible heights of binary trees with n nodes. The minimum height of an n node binary tree is A000523(n), the maximum height is n-1 and all intermediate heights are possible. This conjecture is therefore equivalent to the conjectured formulas. - _Yuchun Ji_, Mar 22 2021 Conjecture: Partial sums of A347523, thus a(n) is the number of nonpowers of 2 <= n-1, or with offset 0: a(n) is the number of nonpowers of 2 <= n. - _Omar E. Pol_, Sep 30 2021 Conjecture, for all n >= 3: a(n) = A083058(n-1) + 1 = n - 1 - A000523(n-1) = n - 1 - floor(log_2(n-1)). - _Antti Karttunen_, Aug 17 2013 Conjecture: a(1) = 1, a(n) = n - 1 - Sum_{i=1..n} sign(floor((n-1)/ 2^i)), n > 1. - _Wesley Ivan Hurt_, Feb 02 2014 Conjecture: a(n) = n - Sum_{k=0..n-2} A036987(k). - _Paul Barry_, Mar 07 2017 Conjecture: a(1) = 1, a(n) = A001855(n-1) - A061168(n-1) + 1. - _Alexander Krivilev_, Aug 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nathaniel Johnston_, Apr 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005776 | OEIS-A182278

Indices n such that A068869(n)<A068869(n-1).

The sequence A068869 is statistically growing, but I conjecture that there are infinitely many indices n such that A068869(n)<A068869(n-1). However, we don't yet know whether there are two consecutive integers in the sequence, i.e., a term n such that, additionally to the above,

A068869(n+1)<A068869(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_ and _Artur Jasinski_, Apr 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005777 | OEIS-A182319

Triangular array: T(n,k) counts upper triangular matrices with entries from {0,1} having n 1's in total, with k 1's on the main diagonal and at least one nonzero entry in each row.

Let F(x,t) = 1 - (1-t)*(1 - (1-x*t) + (1-x*t)*(1-x^2*t) - (1-x*t)*(1-x^2*t)*(1-x^3*t) + ...). Then F(1+x,1+x*t) = 1 + x*t + (t+t^2)*x^2 + (2*t+4*t^2+t^3)*x^3 + (7*t+16*t^2+9*t^3+t^4)*x^4 + ... is conjecturally a generating function for the triangle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005778 | OEIS-A182336

a(n) is the least m>=n, such that the Hamming distance D(n,m)=4.

Conjecture: for n > 96, n + 1 <= a(n) <= 9n/8 + 1. - _Charles R Greathouse IV_, Apr 25 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005779 | OEIS-A182372

Number of distinct sets of nonnegative integers with perimeter n, as defined in the comments.

Conjecture: number of partitions of n into two sorts of parts such that no two successive parts are the same sort (as the order of sorts in a run of identical parts is immaterial, there can be at most two parts of same size), see example. - _Joerg Arndt_, Jun 01 2013 The conjecture of Arndt [namely that a(n) = A227134(n) + A227135(n)] has been verified for n < 1200. - _Patrick Devlin_, Jul 02 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Apr 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005780 | OEIS-A182381

Carmichael numbers divisible by 17 and 29.

Conjecture: Any Carmichael number C divisible by 17 and 29 can be written as C = 952*n + 561; checked up to the Carmichael number 105823343809. The conjecture follows from Korselt's criterion. More is true: a(n) = 2465 mod 55216. - _Charles R Greathouse IV_, Oct 02 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Apr 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005781 | OEIS-A182398

a(n) = (Sum_{k=1..2n} k^2n) mod 2n.

It appears that a(A007018(n)/2) = 1 and conjecturally a(m*A007018(n)/2) = m for a majority of value m.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005782 | OEIS-A182416

Carmichael numbers of the form (60k+13)*(180k+37)*(300k+61).

Conjecture: N = (60k+13)*(180k+37)*(300k+61) is a Carmichael number if (but not only if) 60k+13, 180k+37 and 300k+61 are all three prime numbers. We checked the conjecture up to k = 175; we got Carmichael numbers with three prime divisors for k = 0, 6, 7, 11, 14, 28, 60, 62, 80, 102, 126, 137, 139, 157, 171. The conjecture follows from Korselt's criterion. - _Charles R Greathouse IV_, Oct 02 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Apr 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005783 | OEIS-A182418

a(0)=0, a(n+1) = (a(n) XOR floor(a(n)/6)) + 1, where XOR is the bitwise exclusive-or operator.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Apr 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005784 | OEIS-A182481

a(n) is the least k such that 6*k*n-1 and 6*k*n+1 are twin primes, and a(n)=0, if such k does not exist.

Conjecture: a(n)>0; equivalently, for every n, the arithmetic progression {6*k*n-1} contains infinitely many lessers of twin primes (A001359).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005785 | OEIS-A182511

a(0)=1, a(1)=2, a(n) = (a(n-2)*a(n-1) mod n) + 1.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, May 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005786 | OEIS-A182515

Carmichael numbers of the form C = 23*67*(66n+23).

Conjecture: Any Carmichael number C divisible by 23 and 67 can be written as C = 23*67*(66n+23). The conjecture follows from Korselt's criterion: 67 | a(n) so a(n) = 1 (mod 66). - _Charles R Greathouse IV_, Oct 02 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, May 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005787 | OEIS-A182531

Extremal graph numbers for a triangle with an edge off it.

Assuming Colin Barker's conjectures this is A002378 and the positive terms of A000290 interleaved, except for a(3) = 3. - _Omar E. Pol_, Jan 25 2018 Colin Barker's conjectures are essentially that for n >= 4, a(n) = floor(n^2/4) = A002620(n). This is true: Conjectures from _Colin Barker_, Jul 04 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John Y. Kim_, May 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005788 | OEIS-A182532

Carmichael numbers divisible by 7 and 17.

Conjecture (1): Any Carmichael number C divisible by 7 and 17 of the form 10k+1 can be written as C = 7*17*(120n+119). We got for the tested numbers the following values for n: 478, 1174, 34348, 38350, 82660, 107110, 158658, 162028, 176746, 179278, 262918, 313570, 377788, 563974, 710950. Conjecture (2): Any Carmichael number C divisible by 7 and 17 of the form 10k+5 can be written as C = 7*17*(120n+95). We got for the tested numbers the following values for n: 57, 3291, 28965, 357567, 451893. Conjecture (3): Any Carmichael number C divisible by 7 and 17 of the form 10k+3 can be written as C = 7*17*(120n+47). We got for the tested numbers the following values for n: 111835, 181157, 222733. Conjecture (4): Any Carmichael number C divisible by 7 and 17 of the form 10k+9 can be written as C = 7*17*(120n+71). We got for the tested number the following value for n: 435446.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, May 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005789 | OEIS-A182554

Composite numbers k that divide Fibonacci(k+1) or Fibonacci(k-1).

Any nonsquarefree term is divisible by the square of a Fibonacci-Wieferich prime (i.e., a prime p such that Fibonacci(j) == 0 (mod p^2) for some j not divisible by p). No Fibonacci-Wieferich primes are known, and there are none < 2*10^14, although it is conjectured that there are infinitely many. - _Robert Israel_, Jul 22 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary Detlefs_, May 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005790 | OEIS-A182580

Position of first occurrence of n in A182576.

Conjecture: for k > 12, a(2k) = a(2k-1) + 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, May 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005791 | OEIS-A182614

Least k such that floor(k/r^n)=n, where r = golden ratio = (1+sqrt(5))/2.

Conjectures from _Chai Wah Wu_, Jan 12 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 22 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005792 | OEIS-A182615

Greatest k such that floor(k/r^n)=n, where r = golden mean = (1+sqrt(5))/2.

Conjectures from _Chai Wah Wu_, Jan 12 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 22 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005793 | OEIS-A182617

Number of toothpicks in a toothpick spiral around n cells on hexagonal net.

Conjecture: a(n) = 2*n + ceiling(sqrt(12*n - 3)), for n > 0. - _Vincenzo Librandi_, Sep 20 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Dec 13 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005794 | OEIS-A182663

Number of ordered ways to write n = p + q with q > 0 such that p and 2*p + prime(q) are both prime.

Conjecture: (i) a(n) > 0 for all n > 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005795 | OEIS-A182749

a(n) = the smallest multiple of n with n digits and exactly n divisors. a(n) = 0 if no such number exists.

Conjecture: a(n) > 0 for all nonsquarefree n except for n = 4. - _Jon E. Schoenfield_, Jun 01 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005796 | OEIS-A182750

a(n) = the largest multiple of n with n digits and exactly n divisors; a(n) = 0 if no such number exists.

Conjecture: a(n) > 0 for all nonsquarefree n except for n = 4. - _Jon E. Schoenfield_, Jun 01 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005797 | OEIS-A182755

Expansion of (1+35*x)/(1-90*x^2).

a(1) = 1, a(2) = 35, for n >= 3; a(n) = the smallest number h > a(n-1) such that [[a(n-2) + a(n-1)] * [a(n-2) + h] * [a(n-1) + h]] / [a(n-2) * a(n-1) * h] is an integer (= 130). (conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005798 | OEIS-A182838

H-toothpick sequence in the first quadrant starting with a D-toothpick placed on the diagonal [(0,1), (1,2)] (see Comments for precise definition).

Conjecture: a(n) = (A182840(n+1) + A267458(n+1) - 2)/4. - _Omar E. Pol_, Feb 10 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Dec 12 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005799 | OEIS-A182839

Number of toothpicks and D-toothpicks added at n-th stage to the H-toothpick structure of A182838.

Conjecture: a(n) = (A182841(n+1) + A010673(n))/4, n >= 2. - _Omar E. Pol_, Feb 10 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Dec 12 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005800 | OEIS-A182879

The sum of the lengths of all weighted lattice paths in L_n.

Conjecture: (n-1)*(310*n-781)*a(n) + (-882*n^2+3845*n-3679)*a(n-1) + (214*n^2-2903*n+3751)*a(n-2) + (-358*n^2-459*n+1955)*a(n-3) + (834*n^2-3631*n+4065)*a(n-4) - (262*n-663)*(n-3)*a(n-5) = 0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 11 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005801 | OEIS-A182881

Number of (1,1)-steps in all weighted lattice paths in L_n.

Conjecture: (-n+3)*a(n) +(2*n-5)*a(n-1) +(n-2)*a(n-2) +(2*n-3)*a(n-3) +(-n+1)*a(n-4)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 11 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005802 | OEIS-A182883

Number of weighted lattice paths of weight n having no (1,0)-steps of weight 1.

D-finite n*a(n) = (2*n - 2)*a(n-2) + (4*n - 6)*a(n-3) - (n - 2)*a(n-4), follows easily by differentiating the o.g.f. Maple's sumrecursion command verifies that Sum_{k = 0..floor(n/2)} binomial(n-k,k)*binomial(k,n-2*k) satisfies the same recurrence with the same initial conditions thus proving the above conjecture. - _Peter Bala_, Feb 07 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 11 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005803 | OEIS-A182887

Number of (1,0)-steps in all weighted lattice paths in L_n.

Conjecture: (n-1)*(182*n-279)*a(n) + (-230*n^2+11*n+643)*a(n-1) + (-450*n^2+1603*n-315)*a(n-2) + (-498*n^2+971*n+57)*a(n-3) + (-86*n^2+959*n-529)*a(n-4) + (134*n-59)*(n-3)*a(n-5) = 0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 11 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005804 | OEIS-A182892

Number of weighted lattice paths in L_n having no (1,0)-steps of weight 2 at level 0.

Conjecture: n*a(n) +(n-2)*a(n-1) +2*(-9*n+16)*a(n-2) +5*(2*n-5)*a(n-3) +(10*n-33) *a(n-4) +2*(26*n-109)*a(n-5) +(13*n-37)*a(n-6) +(13*n-63) *a(n-7) +10*(-n+7) *a(n-8)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 12 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005805 | OEIS-A182894

Number of weighted lattice paths in L_n having no (1,0)-steps at level 0. The members of L_n are paths of weight n that start at (0,0) , end on the horizontal axis and whose steps are of the following four kinds: an (1,0)-step with weight 1, an (1,0)-step with weight 2, a (1,1)-step with weight 2, and a (1,-1)-step with weight 1. The weight of a path is the sum of the weights of its steps.

Conjecture: n*a(n) +(-4*n+3)*a(n-1) +(n-3)*a(n-2) -3*a(n-3) +3*(5*n-14)*a(n-4) +6*(n-3)*a(n-5) +6*(n-4)*a(n-6) +4*(-n+6)*a(n-7)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 12 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005806 | OEIS-A182899

Number of returns to the horizontal axis (both from above and below) in all weighted lattice paths in L_n.

Conjecture D-finite with recurrence n*a(n) +(-4*n+3)*a(n-1) +(2*n-3)*a(n-2) +11*(n-3)*a(n-4) +(2*n-9)*a(n-6) +(-4*n+21)*a(n-7) +(n-6)*a(n-8)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 13 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005807 | OEIS-A182904

Number of weighted lattice paths in L_n having no peaks. The members of L_n are paths of weight n that start at (0,0), end on the horizontal axis and whose steps are of the following four kinds: an (1,0)-step with weight 1, an (1,0)-step with weight 2, a (1,1)-step with weight 2, and a (1,-1)-step with weight 1. The weight of a path is the sum of the weights of its steps. A peak is a (1,1)-step followed by a (1,-1)-step.

Conjecture: n*a(n) +(n-2)*a(n-1) +(-7*n+10)*a(n-2) +3*(-n+2)*a(n-3) +(-n+2)*a(n-4) +(-5*n+14)*a(n-5) +(-5*n+18)*a(n-6) +3*(n-4)*a(n-7)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 16 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005808 | OEIS-A182936

Greatest common divisor of the proper divisors of n, 0 if there are none.

Conjecture: Sum_{k=1..n} a(k) = A072107(n) - A034387(n) - 1. - _Vaclav Kotesovec_, Jan 29 2025 a(n) = A014963(n) - A061397(n) for n > 1. In other words, this sequence is the exponential von Mangoldt function restricted to proper divisors of n. See A380118. This implies the above conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Mar 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005809 | OEIS-A182938

If n = Product (p_j^e_j) then a(n) = Product (binomial(p_j, e_j)).

Conjecture: Sum_{k=1..n} a(k) ~ c * n^2, where c = 0.33754... - _Vaclav Kotesovec_, Mar 28 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jan 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005810 | OEIS-A182959

Expansion of o.g.f. 2*(1+x)^2/(1-2*x+sqrt(1-8*x)).

Conjecture: D-finite with recurrence (n+1)*a(n) +(-7*n+3)*a(n-1) +4*(-2*n+5)*a(n-2)=0. - _R. J. Mathar_, Jan 22 2020 (n + 1)*(3*n - 4)*a(n) = 4*(2*n - 3)*(3*n - 1)*a(n-1) for n >= 3 with a(2) = 20. Mathar's conjectured second order recurrence above follows from this. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 31 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005811 | OEIS-A182960

G.f.: exp(Sum_{n>=1} C(6n-1,2n-1)*x^n/n).

The sequence defined by b(n) := [x^n] A(x)^n begins [1, 5, 215, 10463, 537287, 28435880, 1534398353, 83920389642, ...] and conjecturally satisfies the congruence b(p) == b(1) (mod p^3) for prime p >= 5 (checked up to p = 101). - _Peter Bala_, Sep 14 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 31 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005812 | OEIS-A182963

G.f.: A(x) = exp(Sum_{n>=1} A183235(n)*x^n/n) where A183235 is the sums of the cubes of multinomial coefficients.

Conjectured to consist entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005813 | OEIS-A182987

Least a + b such that a*b = A002110(n), the product of the first n primes, where a, b are positive integers.

Conjecture: Limit_{N->oo} (Sum_{n=1..N} log(a(n)-2*sqrt(prime(n)#))) / (Sum_{n=1..N} prime(n)) = 2/e - 1/2 (i.e., A135002 - 1/2). - _Alain Rocchelli_, Nov 30 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Risto Kauppila_, Feb 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005814 | OEIS-A183003

a(n) = A183002(n)/2.

Conjecture: partial sums of A072670. - _Sean A. Irvine_, Jul 14 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jan 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005815 | OEIS-A183016

Conjectured list of smallest terms of k-sociable cycles of order r.

Conjectured list of smallest terms of k-sociable cycles of order r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _William Rex Marshall_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005816 | OEIS-A183017

Conjectured list of multiplicities of multisociable cycles sorted by smallest term.

Conjectured list of multiplicities of multisociable cycles sorted by smallest term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _William Rex Marshall_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005817 | OEIS-A183018

Conjectured list of orders of multisociable cycles sorted by smallest term.

Conjectured list of orders of multisociable cycles sorted by smallest term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _William Rex Marshall_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005818 | OEIS-A183019

Conjectured list of multisociable numbers.

Conjectured list of multisociable numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _William Rex Marshall_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005819 | OEIS-A183026

Conjectured list of smallest members of fully k-sociable cycles of order r.

Conjectured list of smallest members of fully k-sociable cycles of order r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _William Rex Marshall_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005820 | OEIS-A183027

Conjectured list of multiplicities of fully multisociable cycles sorted by smallest term.

Conjectured list of multiplicities of fully multisociable cycles sorted by smallest term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _William Rex Marshall_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005821 | OEIS-A183028

Conjectured list of orders of fully multisociable cycles sorted by smallest term.

Conjectured list of orders of fully multisociable cycles sorted by smallest term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _William Rex Marshall_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005822 | OEIS-A183029

Conjectured list of fully multisociable numbers.

Conjectured list of fully multisociable numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _William Rex Marshall_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005823 | OEIS-A183068

Central terms of triangle A183065.

Conjecture: the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(2*r)) holds for all primes p and positive integers n and r (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 22 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005824 | OEIS-A183069

L.g.f.: Sum_{n>=1,k>=0} CATALAN(n,k)^2 * x^(n+k)/n = Sum_{n>=1} a(n)*x^n/n, where CATALAN(n,k) = n*C(n+2*k-1,k)/(n+k) is the coefficient of x^k in C(x)^n and C(x) is the g.f. of the Catalan numbers.

We conjecture that the sequence satisfies the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for all primes p >= 5 and positive integers n and k. - _Peter Bala_, Mar 20 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 23 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005825 | OEIS-A183110

Period-length of the ultimate periodic behavior of the orbit of a list [1,1,1,...,1] of n 1's under the mapping defined in the comments.

This is also conjectured to be the length of the longest cycle of pebble-moves among the partitions of n (cf. A201144). - _Andrew V. Sutherland_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Feb 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005826 | OEIS-A183161

Self-convolution equals A183160.

Conjecture: a(n) is never congruent to 3 modulo 4; see A218622. - _Paul D. Hanna_, Nov 03 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 27 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005827 | OEIS-A183204

Central terms of triangle A181544.

Conjecture D-finite with recurrence: n^3*a(n) - (2*n-1)*(13*n^2-13*n+4)*a(n-1) - 3*(n-1)*(3*n-4)*(3*n-2)*a(n-2) = 0. - _R. J. Mathar_, May 15 2016 The conjectured D-finite recurrence can be proved by Zeilberger's algorithm.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 30 2010 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005828 | OEIS-A183214

Least number of squared primes that add up to n; -1 if impossible.

Conjecturally, a(n)<=8 with a(n)=8 infinitely often.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Jan 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005829 | OEIS-A183240

Sums of the squares of multinomial coefficients.

Ignoring initial term, equals the logarithmic derivative of A183241; A183241 is conjectured to consist entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005830 | OEIS-A183241

G.f.: A(x) = exp(Sum_{n>=1} A183240(n)*x^n/n) where A183240 is the sums of the squares of multinomial coefficients.

Conjectured to consist entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005831 | OEIS-A183245

Number of permutations of 1..2*n+2 with each element displaced by at least n.

Conjectures from _Colin Barker_, Mar 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005832 | OEIS-A183246

Number of permutations of 1..2*n+3 with each element displaced by at least n.

Conjectures from _Colin Barker_, Mar 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005833 | OEIS-A183247

Number of permutations of 1..2*n+4 with each element displaced by at least n.

Conjectures from _Colin Barker_, Mar 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005834 | OEIS-A183264

Number of singly defective permutations of 1..n with exactly 1 maximum.

Conjecture: a(n) = n * (3*n-4) * 2^(n-3) for n > 1. - _Andrew Howroyd_, May 12 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005835 | OEIS-A183299

Complement of A005563.

Defining the probability for integer value x in a Poisson distribution of integer mean = q > 0 as P(q,x) = e^(-q)*(q^x)/x! then it is conjectured that P(n, a(n)+1)- P(n, a(n)) <= P(n, k+1)- P(n, k), for every k > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005836 | OEIS-A183356

One quarter the number of n X 4 1..4 arrays with no two neighbors of any element equal to each other.

Conjectures from _Colin Barker_, Mar 28 2018: (Start) Assuming Colin Barker's conjectures, a(n) = (12*Fibonacci(n+1))^2, n>1. - _Ehren Metcalfe_, Apr 21 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005837 | OEIS-A183409

Number of n X 2 binary arrays with each sum of a(1..i,1..j) no greater than i*j/2 and rows and columns in nondecreasing order.

Conjecture: a(n) = (2*n^3+11*n^2+31*n+27+(n^2+n+5)*(-1)^n)/32. - _Luce ETIENNE_, Nov 18 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005838 | OEIS-A183430

One third the number of n X 2 0..3 arrays with no element equal to its row sum plus its column sum mod 4.

Conjectures from _Colin Barker_, Mar 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005839 | OEIS-A183452

Number of n X 5 binary arrays with every 1 having exactly three king-move neighbors equal to 1.

Conjecture: g.f. -x*(1 +5*x +607360*x^20 -11532*x^9 +753306*x^19 +6*x^2 -288*x^26 -57080*x^10 -612804*x^16 +7234*x^8 -600857*x^17 +181998*x^18 -57*x^3 -10592*x^25 -16800*x^24 -182104*x^23 -283*x^4 +5942*x^7 -132231*x^15 -345*x^5 +1200*x^6+ 285093*x^14 +259034*x^13 +52803*x^12 +384*x^29 +768*x^28 +2048*x^27 -67623*x^11 -344072*x^22 -99176*x^21)/(-1 +x +275562*x^20 +15867*x^9 -37876*x^19 +x^2 -8768*x^26 +384*x^30 -7896*x^10 -20075*x^16 +1261*x^8 -184239*x^17 -567185*x^18 +60*x^3 -8096*x^25 -174456*x^24 -168944*x^23 -52*x^4+980*x^7 +311194*x^15-58*x^5 -1393*x^6 +65529*x^14 +24304*x^13 -94858*x^12 +384*x^29 +1664*x^28 -1952*x^27 -12862*x^11 +73800*x^22 +537480*x^21) . - _R. J. Mathar_, Dec 03 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005840 | OEIS-A183527

An Ulam-type sequence: a(n) = n if n<=4; for n>4, a(n) = least number > a(n-1) which is a unique sum of 4 distinct earlier terms.

Conjectured G.f.: (-61*x^124-56*x^118+53*x^117+3*x^116 -x^115+2*x^114-65*x^113-58*x^112+57*x^111 -56*x^110-x^109-57*x^108+54*x^106 +3*x^105+58*x^104-52*x^102+50*x^101-55*x^100 +56*x^95-53*x^94-3*x^93+x^92-2*x^91 +2*x^90+x^87-x^86 +41*x^84-51*x^83-66*x^82-58*x^81 -52*x^79+4*x^78-58*x^77 -x^74-x^73-x^72-54*x^71 -6*x^70-59*x^69-x^68-58*x^67-2*x^66 -x^65-2*x^64-56*x^63-4*x^62-x^61 -58*x^60-2*x^59-59*x^58-x^57-x^56 -2*x^55-x^54-x^53-54*x^52-6*x^51 -59*x^50-x^49-58*x^48-2*x^47-x^46 -2*x^45-56*x^44-4*x^43-x^42-58*x^41 -2*x^40-x^39-58*x^38-2*x^37-x^36 -2*x^35-x^34-55*x^33-5*x^32-59*x^31 -x^30-2*x^29-56*x^28-4*x^27-x^26 -58*x^25-2*x^24-x^23-58*x^22-3*x^21 -x^20-2*x^19-56*x^18-4*x^17-59*x^16 -x^15-2*x^14-x^13-x^12-42*x^11 -3*x^10-x^9-x^8-x^7-6*x^6 -6*x^5-x^4-x^3-x^2-x) / (-x^74+x^73+x-1). (This has been verified for n up to 1000.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_ and _Alois P. Heinz_, Jan 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005841 | OEIS-A183586

1/24 the number of (n+1) X 2 0..3 arrays with every 2 X 2 subblock containing exactly three distinct values.

Conjectures from _Colin Barker_, Mar 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005842 | OEIS-A183615

1/120 the number of (n+1) X 3 0..4 arrays with every 2 X 2 subblock containing four distinct values.

Conjecture: G.f.: 1/(x*(3- G(0))) -1/x, where G(k)= 1 + 2^k/(1 - 2*x/(2*x + 2^k*1/G(k+1))); (continued fraction). - _Sergei N. Gladkovskii_, Jul 21 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005843 | OEIS-A183624

Number of (n+1) X 2 0..2 arrays with every 2 x 2 subblock summing to 4.

Conjectures from _Colin Barker_, Feb 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005844 | OEIS-A183625

Number of (n+1) X 3 0..2 arrays with every 2 X 2 subblock summing to 4.

Conjectures from _Colin Barker_, Mar 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005845 | OEIS-A183626

Number of (n+1) X 4 0..2 arrays with every 2 X 2 subblock summing to 4.

Conjectures from _Colin Barker_, Mar 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005846 | OEIS-A183627

Number of (n+1) X 5 0..2 arrays with every 2 x 2 subblock summing to 4.

Conjectures from _Colin Barker_, Mar 30 2018: (Start) Formula and conjectures confirmed by _Robert Israel_, Mar 30 2018 (see link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005847 | OEIS-A183629

Number of (n+1) X 7 0..2 arrays with every 2 X 2 subblock summing to 4.

Conjectures from _Colin Barker_, Mar 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005848 | OEIS-A183630

Number of (n+1) X 8 0..2 arrays with every 2 X 2 subblock summing to 4.

Conjectures from _Colin Barker_, Mar 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005849 | OEIS-A183631

Number of (n+1) X 9 0..2 arrays with every 2 X 2 subblock summing to 4.

Conjectures from _Colin Barker_, Mar 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005850 | OEIS-A183634

Number of (n+1) X 2 0..3 arrays with every 2 x 2 subblock summing to 6.

Conjecture: Number of (n+1) X (k+1) 0..p arrays with every 2 X 2 subblock summing to 2p is Sum_{i=1..(p+1)} i^(n+1)*(p+2-i)^(k+1) - 2*Sum_{i=1..p} i^(n+1)*(p+1-i)^(k+1) + Sum_{i=1..(p-1)} i^(n+1)*(p-i)^(k+1). - _Zhuorui He_, Jun 20 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005851 | OEIS-A183644

Number of (n+1) X 2 0..4 arrays with every 2 X 2 subblock summing to 8.

Conjectures from _Colin Barker_, Mar 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005852 | OEIS-A183682

Number of (n+1) X 3 binary arrays with every 2 X 2 subblock nonsingular.

Conjectures from _Colin Barker_, Feb 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005853 | OEIS-A183690

Number of (n+1) X 2 0..2 arrays with every 2 X 2 subblock nonsingular.

Conjectures from _Colin Barker_, Apr 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005854 | OEIS-A183702

Number of (n+1) X 2 0..3 arrays with every 2 X 2 subblock nonsingular.

Conjectures from _Colin Barker_, Apr 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005855 | OEIS-A183712

1/20 of the number of (n+1) X 3 0..4 arrays with every 2 X 2 subblock strictly increasing clockwise or counterclockwise with one decrease.

a(n) ~ c*k^n where k = 1.629316... is the largest root of x^3 - 3x^2 - x + 1 and c = 1.6293... is conjecturally the largest root of 148x^3 - 296x^2 + 90x - 1. - _Charles R Greathouse IV_, Sep 15 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005856 | OEIS-A183761

Number of 2 X 2 nonsingular 0..n matrices with rows in increasing order.

This conjecture is true, since half of all 2 X 2 nonsingular 0..n matrices have rows in increasing order, and half of them have positive determinant. - _David Radcliffe_, Aug 13 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005857 | OEIS-A183767

1/32 the number of (n+1) X 5 binary arrays with equal numbers of 2 X 2 subblocks with sum mod two being 0 and 1.

Conjectures from _Peter Bala_, Mar 11 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005858 | OEIS-A183869

a(n) = n + floor(sqrt(4*n + 5)); complement of A004116.

a(n-2) appears to be the minimum number of rectangular tiles to place on an n X n grid of unit squares, possibly of different sizes, such that each side of every tile lies on a grid line, every unit square is covered by at most one tile, and each row and each column of the grid has exactly one unit square that is not covered by any tile. - _Yifan Xie_, Jul 19 2025 [The conjecture is proven. - _Yifan Xie_, Jul 24 2025]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005859 | OEIS-A183875

Triangle T(n,k) for A(x)^k=sum(n>=k T(n,k)*x^n), where o.g.f. A(x) satisfies A(x)=(a+b*x*A(x))/(c-d*x*A(x)), a=1,b=2,c=1,d=2.

Conjecture: T(n,1) = A156017(n-1). - _R. J. Mathar_, Nov 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Feb 12 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005860 | OEIS-A183877

Number of arrangements of n+2 numbers in 0..2 with each number being the sum mod 3 of two others.

Conjectures from _Colin Barker_, Apr 05 2018: (Start) Conjecture is true. The complement consists of arrangements of the forms

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005861 | OEIS-A183878

Number of arrangements of n+2 numbers in 0..3 with each number being the sum mod 4 of two others.

Conjectures from _Colin Barker_, Apr 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005862 | OEIS-A183879

Number of arrangements of n+2 numbers in 0..4 with each number being the sum mod 5 of two others.

Conjectures from _Colin Barker_, Apr 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005863 | OEIS-A183880

Number of arrangements of n+2 numbers in 0..5 with each number being the sum mod 6 of two others.

Conjectures from _Mikhail Smirnov_, Oct 14 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005864 | OEIS-A183885

Number of arrangements of n+3 numbers in 0..2 with each number being the sum mod 3 of three others.

Conjectures from _Colin Barker_, Apr 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005865 | OEIS-A183886

Number of arrangements of n+3 numbers in 0..3 with each number being the sum mod 4 of three others.

Conjectures from _Colin Barker_, Apr 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005866 | OEIS-A183897

Number of nondecreasing arrangements of n+3 numbers in 0..2 with each number being the sum mod 3 of three others.

Conjectures from _Colin Barker_, Apr 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005867 | OEIS-A183898

Number of nondecreasing arrangements of n+3 numbers in 0..3 with each number being the sum mod 4 of three others.

Conjectures from _Colin Barker_, Apr 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005868 | OEIS-A183899

Number of nondecreasing arrangements of n+3 numbers in 0..4 with each number being the sum mod 5 of three others.

Conjectures from _Colin Barker_, Apr 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005869 | OEIS-A183900

Number of nondecreasing arrangements of n+3 numbers in 0..5 with each number being the sum mod 6 of three others.

Conjectures from _Colin Barker_, Apr 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005870 | OEIS-A183901

Number of nondecreasing arrangements of n+3 numbers in 0..6 with each number being the sum mod 7 of three others.

Conjectures from _Colin Barker_, Apr 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005871 | OEIS-A183905

Number of nondecreasing arrangements of n+2 numbers in 0..2 with each number being the sum mod 3 of two others.

Conjectures from _Colin Barker_, Apr 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005872 | OEIS-A183906

Number of nondecreasing arrangements of n+2 numbers in 0..3 with each number being the sum mod 4 of two others.

Conjectures from _Colin Barker_, Apr 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005873 | OEIS-A183907

Number of nondecreasing arrangements of n+2 numbers in 0..4 with each number being the sum mod 5 of two others.

Conjectures from _Colin Barker_, Apr 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005874 | OEIS-A183908

Number of nondecreasing arrangements of n+2 numbers in 0..5 with each number being the sum mod 6 of two others.

Conjectures from _Colin Barker_, Apr 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005875 | OEIS-A183909

Number of nondecreasing arrangements of n+2 numbers in 0..6 with each number being the sum mod 7 of two others.

Conjectures from _Colin Barker_, Apr 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005876 | OEIS-A183910

Number of nondecreasing arrangements of n+2 numbers in 0..7 with each number being the sum mod 8 of two others.

Conjectures from _Colin Barker_, Apr 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005877 | OEIS-A183911

Number of nondecreasing arrangements of n+2 numbers in 0..8 with each number being the sum mod 9 of two others.

Conjectures from _Colin Barker_, Apr 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005878 | OEIS-A183978

1/4 the number of (n+1) X 2 binary arrays with all 2 X 2 subblock sums the same.

Based on the conjectured recursion formula, it is also the number of notches in a sheet of paper when you fold it n times and cut off the four corners (see A274230). - _Philippe Gibone_, Jul 06 2016 Based on the conjectured recursion formula, we may prove (by a tedious induction) that a(n) = (2^ceiling(n/2) + 1) * (2^floor(n/2) + 1) = A051032(n) * A051032(n-1) for n >= 1. - _Philippe Gibone_, Jul 06 2016, corrected by _Robert Israel_, May 21 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005879 | OEIS-A183979

1/4 the number of (n+1) X 3 binary arrays with all 2 X 2 subblock sums the same.

Conjectures from _Colin Barker_, Apr 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005880 | OEIS-A183980

1/4 the number of (n+1) X 4 binary arrays with all 2 X 2 subblock sums the same.

Conjectures from _Colin Barker_, Apr 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005881 | OEIS-A183981

1/4 the number of (n+1) X 5 binary arrays with all 2 X 2 subblock sums the same.

Conjectures from _Colin Barker_, Apr 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005882 | OEIS-A183982

1/4 the number of (n+1) X 6 binary arrays with all 2 X 2 subblock sums the same.

Conjectures from _Colin Barker_, Apr 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005883 | OEIS-A183983

1/4 the number of (n+1) X 7 binary arrays with all 2 X 2 subblock sums the same.

Conjectures from _Colin Barker_, Apr 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005884 | OEIS-A183984

1/4 the number of (n+1) X 8 binary arrays with all 2 X 2 subblock sums the same.

Conjectures from _Colin Barker_, Apr 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005885 | OEIS-A183985

1/4 the number of (n+1) X 9 binary arrays with all 2 X 2 subblock sums the same.

Conjectures from _Colin Barker_, Apr 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005886 | OEIS-A184018

Expansion of c(x/(1-x-x^2)) / (1-x-x^2), c(x) the g.f. of A000108.

Conjecture: (n+1)*a(n) + 2*(1-3*n)*a(n-1) + 3*(n-1)*a(n-2) + 2*(3*n-5)*a(n-3) + (n-3)*a(n-4) = 0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005887 | OEIS-A184031

1/16 the number of (n+1) X 2 0..3 arrays with all 2 X 2 subblocks having the same four values.

Conjectures from _Colin Barker_, Apr 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005888 | OEIS-A184032

1/16 the number of (n+1) X 3 0..3 arrays with all 2 X 2 subblocks having the same four values.

Conjectures from _Colin Barker_, Apr 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005889 | OEIS-A184033

1/16 the number of (n+1) X 4 0..3 arrays with all 2 X 2 subblocks having the same four values.

Conjectures from _Colin Barker_, Apr 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005890 | OEIS-A184034

1/16 the number of (n+1) X 5 0..3 arrays with all 2 X 2 subblocks having the same four values.

Conjectures from _Colin Barker_, Apr 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005891 | OEIS-A184035

1/16 the number of (n+1) X 6 0..3 arrays with all 2 X 2 subblocks having the same four values.

Conjectures from _Colin Barker_, Apr 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005892 | OEIS-A184036

1/16 the number of (n+1) X 7 0..3 arrays with all 2 X 2 subblocks having the same four values.

Conjectures from _Colin Barker_, Apr 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005893 | OEIS-A184037

1/16 the number of (n+1) X 8 0..3 arrays with all 2 X 2 subblocks having the same four values.

Conjectures from _Colin Barker_, Apr 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005894 | OEIS-A184038

1/16 the number of (n+1) X 9 0..3 arrays with all 2 X 2 subblocks having the same four values.

Conjectures from _Colin Barker_, Apr 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005895 | OEIS-A184041

1/9 the number of (n+1) X 3 0..2 arrays with all 2 X 2 subblocks having the same four values.

Conjectures from _Colin Barker_, Apr 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005896 | OEIS-A184042

1/9 the number of (n+1) X 4 0..2 arrays with all 2 X 2 subblocks having the same four values.

Conjectures from _Colin Barker_, Apr 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005897 | OEIS-A184043

1/9 the number of (n+1) X 5 0..2 arrays with all 2 X 2 subblocks having the same four values.

Conjectures from _Colin Barker_, Apr 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005898 | OEIS-A184044

1/9 the number of (n+1) X 6 0..2 arrays with all 2 X 2 subblocks having the same four values.

Conjectures from _Colin Barker_, Apr 10 2018: (Start) Conjectured e.g.f.: 2*cosh(2*x) + 3*cosh(sqrt(2)*x) + 73*sinh(x) + cosh(x)*(73 + 4*sinh(x)) + 2*sqrt(2)*sinh(sqrt(2)*x) - 78. - _Stefano Spezia_, Aug 01 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005899 | OEIS-A184045

1/9 the number of (n+1) X 7 0..2 arrays with all 2 X 2 subblocks having the same four values.

Conjectures from _Colin Barker_, Apr 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005900 | OEIS-A184046

1/9 the number of (n+1) X 8 0..2 arrays with all 2 X 2 subblocks having the same four values.

Conjectures from _Colin Barker_, Apr 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005901 | OEIS-A184047

1/9 the number of (n+1) X 9 0..2 arrays with all 2 X 2 subblocks having the same four values.

Conjectures from _Colin Barker_, Apr 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005902 | OEIS-A184063

Number of (n+1) X 2 binary arrays with rows and columns in nondecreasing order and with no 2 X 2 subblock sum differing from a horizontal or vertical neighbor subblock sum by more than one.

Conjectures from _Colin Barker_, Apr 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005903 | OEIS-A184064

Number of (n+1) X 3 binary arrays with rows and columns in nondecreasing order and with no 2 X 2 subblock sum differing from a horizontal or vertical neighbor subblock sum by more than one.

Conjectures from _Colin Barker_, Apr 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005904 | OEIS-A184065

Number of (n+1) X 4 binary arrays with rows and columns in nondecreasing order and with no 2 X 2 subblock sum differing from a horizontal or vertical neighbor subblock sum by more than one.

Conjectures from _Colin Barker_, Apr 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005905 | OEIS-A184073

Number of (n+1) X 2 0..2 arrays with rows and columns in nondecreasing order and with no 2 X 2 subblock sum differing from a horizontal or vertical neighbor subblock sum by more than one.

Conjectures from _Colin Barker_, Apr 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005906 | OEIS-A184130

Number of n X 2 0..2 arrays with rows and columns in nondecreasing order.

Conjectures from _Colin Barker_, Apr 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005907 | OEIS-A184138

Number of n X 3 binary arrays with rows and columns in nondecreasing order.

Conjectures from _Colin Barker_, Apr 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005908 | OEIS-A184145

1/4 the number of (n+1) X 2 0..3 arrays with every 2 X 2 subblock having at least two equal elements connected horizontally or vertically.

Conjectures from _Colin Barker_, Apr 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005909 | OEIS-A184223

Half the number of (n+1) X 2 0..3 arrays with every 2 X 2 subblock diagonal sum differing from its antidiagonal sum by more than 3.

Conjectures: a(n) = 3*a(n-1)+3*a(n-2)-6*a(n-3); G.f.: -3*x*(2*x+1)*(6*x-5) / (6*x^3-3*x^2-3*x+1). - _Colin Barker_, Sep 26 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005910 | OEIS-A184368

1/3 the number of (n+1) X 2 0..2 arrays with every 2 X 2 subblock having at least two equal elements connected horizontally or vertically.

Conjectures from _Colin Barker_, Feb 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 12 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005911 | OEIS-A184423

a(n) = (2*n)!*(3*n)!/n!^5.

Inductively define a family of sequences {a(i,n) : n >= 0}, i >= 1, by setting a(1,n) = a(n) and, for i >= 2, a(i,n) = [x^n] (exp(Sum_{k >= 1} a(i-1,k)*x^k/k))^n. We conjecture that the sequences {a(i,n) : n >= 0}, i >= 2, also satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for primes p >= 5 and positive integers n and r. Cf. A362730 and A362732. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 13 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005912 | OEIS-A184540

Number of (n+2) X 3 binary arrays with each 3 X 3 subblock having rows and columns in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Apr 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005913 | OEIS-A184541

Number of (n+2) X 4 binary arrays with each 3 X 3 subblock having rows and columns in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Apr 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005914 | OEIS-A184542

Number of (n+2) X 5 binary arrays with each 3 X 3 subblock having rows and columns in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Apr 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005915 | OEIS-A184543

Number of (n+2) X 6 binary arrays with each 3 X 3 subblock having rows and columns in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Apr 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005916 | OEIS-A184544

Number of (n+2) X 7 binary arrays with each 3 X 3 subblock having rows and columns in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Apr 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005917 | OEIS-A184545

Number of (n+2) X 8 binary arrays with each 3 X 3 subblock having rows and columns in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Apr 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005918 | OEIS-A184546

Number of (n+2) X 9 binary arrays with each 3 X 3 subblock having rows and columns in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Apr 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005919 | OEIS-A184547

Number of (n+2) X 10 binary arrays with each 3 X 3 subblock having rows and columns in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Apr 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005920 | OEIS-A184634

a(n) = floor(1/{(10+n^4)^(1/4)}), where {}=fractional part.

This implies the conjectured recurrence for n >= 12. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005921 | OEIS-A184674

a(n) = n+floor((n/2-1/(2*n))^2); complement of A184675.

Conjecture: a(n) = A014616(n-1). - _R. J. Mathar_, Jan 29 2011 The above conjecture is true. - _Stefano Spezia_, Apr 04 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005922 | OEIS-A184679

Number of (n+1) X 3 binary arrays with every 2 X 2 subblock singular.

Conjectures from _Colin Barker_, Apr 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005923 | OEIS-A184688

1/3 the number of n X 3 0..2 arrays with no element equal both to the element above and to the element to its left.

Conjectures from _Colin Barker_, Apr 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005924 | OEIS-A184774

Primes of the form floor(k*sqrt(2)).

The conjecture formerly posted here, that "if r is an irrational number and 1<r<2, then there are infinitely many primes in the set L={floor(n*r)}," is proved in I. M. Vinogradov, The Method of Trigonometrical Sums in the Theory of Numbers, (1954), page 180. The conjecture generalized: if r is a positive irrational number and h is a real number, then each of the sets {floor(n*r+h)}, {round(n*r+h)}, and {ceiling(n*r+h)} contains infinitely many primes. Can the method in Vinogradov be extended to cover these cases? [Update regarding the conjecture from _Clark Kimberling_, Jan 03 2011.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 21 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005925 | OEIS-A184878

Third column (negated) of triangle in A182971.

Conjectures: G.f.: x^5*(225+54*x+x^2)/(x-1)^10 . a(n) = n *(n-1) *(n-2) *(n-3) *(n-4) *(n+2) *(n+1) *(35*n^2+182*n+240) /45360. - _R. J. Mathar_, May 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005926 | OEIS-A184894

a(n) equals the coefficient of x^(2n-1) in the n-th iteration of x+x^3 for n>=1.

Conjecture: a(m) = 0 (mod 3) everywhere except at m = (3^n+1)/2, n>=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005927 | OEIS-A185004

Ramanujan modulo primes R_(3,1)(n): a(n) is the smallest number such that if x >= a(n), then pi_(3,1)(x) - pi_(3,1)(x/2) >= n, where pi_(3,1)(x) is the number of primes==1 (mod 3) <= x.

We conjecture that for all n >= 1 a(n) <= A104272(3*n). This conjecture is based on observation that, if interval (x/2, x] contains >= 3*n primes, then at least n of them are of the form 3*k+1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Dec 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005928 | OEIS-A185005

Ramanujan primes R_(3,2)(n): a(n) is the smallest number such that if x >= a(n), then pi_(3,2)(x) - pi_(3,2)(x/2) >= n, where pi_(3,2)(x) is the number of primes==2 (mod 3) <= x.

We conjecture that for all n >= 1, a(n) <= A104272(3*n). This conjecture is based on observation that, if interval (x/2, x] contains >= 3*n primes, then at least n of them are of the form 3*k+2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Dec 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005929 | OEIS-A185006

Ramanujan primes R_(4,1)(n): a(n) is the smallest number such that if x >= a(n), then pi_(4,1)(x) - pi_(4,1)(x/2) >= n, where pi_(4,1)(x) is the number of primes==1 (mod 4) <= x.

We conjecture that for all n >= 1, a(n) <= A104272(3*n). This conjecture is based on observation that, if interval (x/2, x] contains >= 3*n primes, then at least n of them are of the form 4*k+1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Dec 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005930 | OEIS-A185007

Ramanujan primes R_(4,3)(n): a(n) is the smallest number such that if x >= a(n), then pi_(4,3)(x) - pi_(4,3)(x/2) >= n, where pi_(4,3)(x) is the number of primes==3 (mod 4) <= x.

We conjecture that for all n >= 1, a(n) <= A104272(3*n). This conjecture is based on observation that, if interval (x/2, x] contains >= 3*n primes, then at least n of them are of the form 4*k+3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Dec 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005931 | OEIS-A185010

a(n) = A000108(n)*A015518(n+1), where A000108 are the Catalan numbers and A015518(n) = 2*A015518(n-1) + 3*A015518(n-2).

Conjecture: n*(n+1)*a(n) -4*n*(2*n-1)*a(n-1) -12*(2*n-1)*(2*n-3)*a(n-2)=0. - _R. J. Mathar_, Oct 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005932 | OEIS-A185020

a(n) = A000108(n)*A002605(n+1), where A000108 are the Catalan numbers.

Conjecture: +n*(n+1)*a(n) -4*n*(2*n-1)*a(n-1) -8*(2*n-1)*(2*n-3)*a(n-2)=0. - _R. J. Mathar_, Oct 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005933 | OEIS-A185023

Number of steps to reach the number 1 of the map n -> n/3 if x==0 (mod 3), n -> (4n-1)/3 if x==1 (mod 3), n -> (4n+1)/3 if x==2 (mod 3).

Conjecture: Every positive integer reaches 1 (verified up to 10^11). - _Farhad Banazadeh_, Sep 15 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Feb 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005934 | OEIS-A185046

Smallest prime p such that 2n+1 = p^3 - 2q for some odd prime q, or 0 if no such prime exists.

Conjecture: The odd numbers different from a cube are of the form m = p^3 - 2q where p and q are prime numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005935 | OEIS-A185061

Position of the first occurrence of n in A193358 when it is considered to have the starting offset 1 instead of 0.

Conjecture: Only odd numbers occur after 2. - _Antti Karttunen_, Jun 13 2013 Conjecture is true. - _Chai Wah Wu_, Jun 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Jan 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005936 | OEIS-A185064

Numbers k such that a Golay sequence of length k exists.

It is known that the sequence contains all numbers 2^i 10^j 26^m, and that all terms k > 1 are even and not divisible by any prime == 3 (mod 4). But the full characterization of these numbers is an open problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005937 | OEIS-A185078

Numbers k for which A064380(k) = k/2.

Note that, if there exist infinitely many infinitary perfect numbers (A007357), then, as k tends to infinity over such numbers, A064380(k)/k = 1/2 + o(k^(-1+eps)). We conjecture that here A064380(k)/k = 1/2 infinitely many times, and thus the sequence contains infinitely many infinitary perfect numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Feb 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005938 | OEIS-A185087

a(n) = Sum_{k=0..floor(n/2)} binomial(n-k,k)*A000108(k+1).

Conjecture: (n+4)*a(n) -(2*n+5)*a(n-1) -3*(n+1)*a(n-2) +2*(2*n-1)*a(n-3)=0. - _R. J. Mathar_, Nov 16 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005939 | OEIS-A185089

A transform of the little Schroeder numbers.

conjecture: (n+2)*a(n) -3*(n+1)*a(n-1) +3*(2-n)*a(n-2) +(11*n-20)*a(n-3) +(11-5*n)*a(n-4) +(4-n)*a(n-5) =0. - _R. J. Mathar_, Nov 16 2011 (Formula verified and used for computations. - _Fung Lam_, Feb 24 2014)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005940 | OEIS-A185095

Rectangular array read by antidiagonals: row q has generating function F_q(x) = sum_{r=0,...,q-1} ((q-r)*(-1)^r*binomial(2*q-r,r)*x^r) / sum_{s=0,...,q} ((-1)^s*binomial(2*q-s,s)*x^s), where q=1,2,....

Conjecture. The n-th entry in row q is given by R_q(n) = 2^(2*n)*(sum_{j=1,...,n+1} (cos(j*Pi/(2*q+1)))^(2*n)), q >= 1, n >= 0. Conjecture. G.f. for column n is of the form G_n(x) = H_n(x)/(1-x)^2, where H_n(x) is a polynomial in x, n >= 0. Conjecture. 2*A185095(q,n) = A198632(2*q,n), q >= 1, n >= 0. - _L. Edson Jeffery_, Nov 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Jan 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005941 | OEIS-A185099

Indices k for which A185079(k) > k^2.

Conjecture: The sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Feb 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005942 | OEIS-A185104

Primes with property that each swapping any pair of digits also gives a prime.

Conjecture: next terms are 1111111111111111111 and 11111111111111111111111.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Dec 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005943 | OEIS-A185165

Numbers n such that lambda(n) = lambda(n - lambda(n)).

Lambda is the function in A002322. If there are infinitely many Sophie Germain primes (conjecture), then this sequence is infinite. Proof: The numbers of the form 3p^2 are in a subsequence if p and 2p+1 are both prime with p > 3, because from the property that lambda(3p^2) = p(p-1) and lambda (p(2p+1)) = p(p-1), if m = 3p^2 then lambda(m-phi(m)) = lambda (3p^2 - p(p-1)) = lambda(p(2p+1)) = p(p-1) = lambda(m).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Mar 31 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005944 | OEIS-A185351

Sums of distinct perfect numbers.

The first 131072 terms of this sequence are even. Conjecturally, all terms are even.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Feb 09 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005945 | OEIS-A185371

Product of two consecutive primes divided by the next prime and rounded down.

For n > 3, prime(n)/2 < a(n) < prime(n). prime(n) - a(n) << log(n)^2 under Cramer's conjecture. Weaker estimates follow from the Prime Number Theorem. - _Charles R Greathouse IV_, Jan 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tyler Carrico_, Jan 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005946 | OEIS-A185376

Number of binary necklaces of 2n beads for which a cut exists producing a palindrome.

Conjecture: a(n) = 2^(n-1) + 2^((n-2^t)/(2^(t+1))), where t = number of factors of 2 in n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tony Bartoletti_, Feb 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005947 | OEIS-A185378

Number of binary necklaces of 2n beads that are identical when turned over yet cannot be cut to produce a palindrome.

These necklaces have bilateral symmetry across axes that involve only vertices. a(n) = A029744(2n) - A185333(2n). Conjecture: a(n) = 2^n - 2^((n - 2^t)/(2^(t+1))), where t = number of factors of 2 in n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tony Bartoletti_, Feb 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005948 | OEIS-A185392

Position of g(n) when the numbers f(j) and g(k) are jointly ranked, where f(j) = j + |cos j| and g(k) = k + |sin k|.

For a guide to related sequences and a conjecture, see A206911.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005949 | OEIS-A185410

A decomposition of the double factorials A001147.

Row polynomial R(n,x) = sum {k = 1..n} 2^(n-2*k)*C(2*k,k)*k!*Stirling2(n,k)*(x-1)^(n-k). R(n,4*x)/(1-4*x)^(n+1/2) = sum {k>=0} C(2*k,k)*(2*k+1)^n*x^k. The sequence of rational functions x*R(n,x)/(1-x)^(n+1) conjecturally occurs in the first column of (I - x*A112857)^(-1). (1+x)^(n-1)*R(n,x/(x+1)) gives the n-th row polynomial of A186695.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Jan 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005950 | OEIS-A185450

Define a map of the integers by T_k(i)=i/2 if i even, (3i+k)/2 if i odd; a(n) is the conjectured number of cycles for trajectories under this map in the case k = 5^n.

Define a map of the integers by T_k(i)=i/2 if i even, (3i+k)/2 if i odd; a(n) is the conjectured number of cycles for trajectories under this map in the case k = 5^n. The values are only (strong) conjectures; showing that there are correct will be about as hard as solving the famous "3x+1" problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005951 | OEIS-A185455

Trajectory of 7 under repeated application of the map in A185452.

It is conjectured that this trajectory is unbounded, but this is an open problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005952 | OEIS-A185510

Array of primes in the natural number array A000027, by antidiagonals.

Conjecture: Every row contains infinitely many primes. Every prime occurs exactly once; that is, every prime is uniquely expressible as (1/2)(n^2 + (2k-1)n + (k-2)(k-1)) for some positive integers n and k. We assume as true the conjecture that each row is infinite. - _Clark Kimberling_, Mar 10 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005953 | OEIS-A185511

Array: row n shows numbers m such that prime(m) is in row n of the natural number array A000027, by antidiagonals.

If the conjecture at A185510 is true, then the A185512, as a sequence, is a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005954 | OEIS-A185636

a(n) = |{0 <= k < n: n+k and n+k^2 are both prime}|.

Conjecture: a(n) > 0 for all n > 1. This conjecture has been verified for n up to 10^8. It is stronger than Bertrand's postulate proved by Chebyshev in 1850. Zhi-Wei Sun also guessed the following refinement of the conjecture: For any integer n > 1456 there is an integer k among 0,...,n-1 such that n-k, n+k and n+k^2 are all prime; in other words, there is a prime p <= n such that 2n-p and n+(n-p)^2 are both prime. For other refinements of the conjecture, the reader may consult arXiv:1211.1588. The author also conjectured the following polynomial analogs: In a 2017 paper, the author announced a USD $100 prize for the first solution to his conjecture that for each n = 1,2,3,... there is an integer k among 0,...,n such that n+k and n+k^2 are both prime. - _Zhi-Wei Sun_, Dec 03 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005955 | OEIS-A185645

Number of permutations q_1,...,q_n of the first n primes p_1,...,p_n with q_1 = p_1 = 2 and q_n = p_n, and with |q_1-q_2|, |q_2-q_3|, ..., |q_{n-1}-q_n|, and |q_n-q_1| (if n>2) pairwise distinct.

Conjecture: a(n) > 0 for all n > 0. In general, for any n consecutive primes p_k,...,p_{k+n-1}, there always exists a permutation q_k,...,q_{k+n-1} of p_k,...,p_{k+n-1} with q_{k+n-1} = p_{k+n-1} such that the n-1 numbers |q_k-q_{k+1}|, |q_{k+1}-q_{k+2}|,...,|q_{k+n-2}-q_{k+n-1}| are pairwise distinct. (In the case k = 2, this implies that a(n) > 0.) On Aug 31 2013, Zhi-Wei Sun proved the following extension of the general conjecture: Let a_1 < a_2 < ... < a_n be a sequence of n distinct real numbers in ascending order. Then there is a permutation b_1, ..., b_n of a_1, ..., a_n with b_n = a_n such that |b_1-b_2|, |b_2-b_3|, ..., |b_{n-1}-b_n| are pairwise distinct. In fact, when n = 2*k is even we may take (b_1,...,b_n) = (a_k,a_{k+1},a_{k-1},a_{k+2},...,a_2,a_{2k-1},a_1,a_{2k}); when n = 2*k-1 is odd we may take (b_1,...,b_n) = (a_k,a_{k-1},a_{k+1},a_{k-2},a_{k+2},..., a_2,a_{2k-2},a_1,a_{2k-1}). On Sep 01 2013, Zhi-Wei Sun made the following conjecture: (i) For any n distinct real numbers a_1, a_2, ..., a_n (not necessarily in ascending or descending order), there is a permutation b_1, ..., b_n of a_1, ..., a_n with b_1 = a_1 such that the n-1 distances |b_1-b_2|, |b_2-b_3|, ..., |b_{n-1}-b_n| are pairwise distinct. We believe that part (ii) of the new conjecture holds at least when G is cyclic, and it might also hold when the group G is not abelian.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005956 | OEIS-A185726

Array associated with A049480, by descending antidiagonals.

Conjectures by _Georg Fischer_, Jun 01 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005957 | OEIS-A185761

Number of (n+1) X 2 binary arrays with no 2 X 2 subblock trace equal to any horizontal or vertical neighbor 2 X 2 subblock trace.

Conjectures from _Colin Barker_, Feb 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005958 | OEIS-A185858

1/128 the number of (n+2) X 3 binary arrays with no 3 X 3 subblock trace equal to any horizontal or vertical neighbor 3 X 3 subblock trace.

Conjectures from _Colin Barker_, Apr 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005959 | OEIS-A185897

G.f. A(x) satisfies: A(x - A(x)^2) = x/(1-x).

Conjecture: for n > 1, a(n) is odd when n = 3*2^k through 4*2^k - 1 for k >= 0. - _Paul D. Hanna_, Oct 05 2025 The conjecture is true (see Preil link). Proof sketch: the defining equation determines A(x) uniquely over the integers, so it suffices to solve it mod 2. Over GF(2) put L(y) = Sum_{k>=0} y^(2^k), so that L(y)^2 + L(y) = y. Then B(x) = L(x + x^3)/(1 + x) satisfies B(x + B(x)^2) = x/(1 + x), because with u = x + B^2 one has u = L(x+x^3)/(1+x)^2 and L(u + u^3) = x(1+u)/(1+x), verified by the Artin-Schreier identity. Hence A(x) == L(x + x^3)/(1 + x) (mod 2), whose coefficients are 1 exactly for n = 1 and 3*2^k <= n <= 4*2^k - 1. - _Dovid Preil_, Oct 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005960 | OEIS-A185943

Riordan array ((1/(1-x))^m, x*A000108(x)), m = 2.

Conjecture: R(n,1,2) = A014140(n-1). R(n,2,2) = A014143(n-2). - _R. J. Mathar_, Feb 11 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Feb 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005961 | OEIS-A185956

Fajtlowicz q-primes.

Paul Erdős and C. Larson conjecture that all primes occur in the sequence of q's.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Craig Eric Larson_, Feb 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005962 | OEIS-A185965

Central coefficients of number triangle A185962.

Conjecture: 15*n*(n-1)*a(n) +10*(2*n-1)*(n-1)*a(n-1) +(109*n^2-263*n+180)*a(n-2) +4*(-2*n^2-10*n+57)*a(n-3) +60*(n-4)^2*a(n-4) -6*(2*n-7)*(n-5)*a(n-5)=0. - _R. J. Mathar_, Dec 03 2014 Conjecture: 3*n*(n-1)*(57*n-136)*a(n) +(n-1)*(171*n^2-415*n+14)*a(n-1) +2*(532*n^3-2281*n^2+2755*n-840)*a(n-2) -2*(n-3)*(304*n^2-835*n+216)*a(n-3) +2*(19*n-5)*(n-4)*(2*n-5)*a(n-4)=0. - _R. J. Mathar_, Dec 03 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005963 | OEIS-A185966

Series reversion of A028310.

Conjecture: 6*n*(n+1)*a(n) -n*(n-14)*a(n-1) +2*n*(14*n-19)*a(n-2) -4*(n-2)*(17*n-48)*a(n-3) +6*(2*n-5)*(n-4)*a(n-4)=0. - _R. J. Mathar_, Nov 15 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005964 | OEIS-A186008

Irregular triangle T(n,k) read by rows, in which row n has the pattern of conjectured dropping times in the Collatz iteration.

Irregular triangle T(n,k) read by rows, in which row n has the pattern of conjectured dropping times in the Collatz iteration.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Feb 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005965 | OEIS-A186080

Fourth powers that are palindromic in base 10.

Conjecture: If k^4 is a palindrome > 0, then k begins and ends with digit 1, all other digits of k being 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matevz Markovic_, Feb 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005966 | OEIS-A186185

Expansion of 1/(1 - x*A001764(x/(1-x))/(1-x)).

Conjecture: D-finite with recurrence 2*(2*n-1)*(n-1)*a(n) +(9*n^2-127*n+170)*a(n-1) +10*(-57*n^2+368*n-559)*a(n-2) +(2579*n^2-18451*n+32892)*a(n-3) +3*(-1633*n^2+13195*n-26878)*a(n-4) +6*(712*n-3111)*(n-5)*a(n-5) -1395*(n-5)*(n-6)*a(n-6)=0, n>6. - _R. J. Mathar_, May 07 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Feb 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005967 | OEIS-A186188

Least k such that A156077^(k)(n)=1 where a^(k)=a(a^(k-1)).

conjecture: a(n)=floor(log(n)/log(2))+0 or +1

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005968 | OEIS-A186189

Least k such that A074286^(k)(n) = 1 where a^(k) = a(a^(k-1)).

Conjecture: a(n)=floor(log(n)/log(2))+0 or +1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005969 | OEIS-A186232

Row sums of A185249.

Conjecture: the constant is the same as in A343032. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 15 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005970 | OEIS-A186239

G.f. A(x) satisfies A(x) = 1+x*A(x)+x^2*A(x)^2+2*x^3*A(x)^3.

Conjecture: 4*(2*n+3)*(n+1)*a(n) +(113*n^2-91*n-72)*a(n-1) + 3*(-135*n^2+263*n-108)*a(n-2) -3*(107*n-119)*(n-2)*a(n-3) -1411*(n-2)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Nov 14 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Feb 15 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005971 | OEIS-A186240

G.f. A(x) defined by A(x) = 1 +x*A(x) +x^2*A(x)^2 +3*x^3*A(x)^3.

Conjecture: 6*(2*n+3)*(n+1)*a(n) +(277*n^2-191*n-162)*a(n-1) +6*(-155*n^2+305*n-126)*a(n-2) -4*(181*n-195)*(n-2)*a(n-3) -5304*(n-2)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Sep 27 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Feb 15 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005972 | OEIS-A186253

Indices of zeros of the sequence u(n)=abs(u(n-1)-gcd(u(n-1),n-1)), u(1)=1.

For any fixed integer m>=1 define u(1)=1 and u(n)=abs(u(n-1)-gcd(u(n-1),m*n-1)). Then (b_m(k))_{k>=1} is the sequence of integers such that u(b_m(k))=0 and we conjecture that for k large enough m*b_m(k)+m-1 is a prime number. Here for m=1 it appears a(n) is prime for n>=1. Conjecture: a(n) is asymptotic to c*2^n with c = 1.1861...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005973 | OEIS-A186254

a(n) = 2*b(n)+1, where b(n) lists the zeros of the sequence u(n)=abs(u(n-1)-gcd(u(n-1),2*n-1)), u(1)=1.

For any fixed integer m>=1 define u(1)=1 and u(n)=abs(u(n-1)-gcd(u(n-1),m*n-1)). Then (b_m(k))_{k>=1} is the sequence of integers such that u(b_m(k))=0 and we conjecture that for all k large enough, m*b_m(k)+m-1 is a prime number. Here for m=2 it appears a(n) is prime for n>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005974 | OEIS-A186255

a(n) = 3*b_3(n)+2, where b_3 lists the zeros of the sequence A261303: u(n+1)=abs(u(n)-gcd(u(n),3*n+2)), u(1)=1.

For any fixed integer m>=1 define u(1)=1 and u(n)=abs(u(n-1)-gcd(u(n-1),m*n-1)). Then (b_m(k))_{k>=1} is the sequence of integers such that u(b_m(k))=0 and we conjecture that for k large enough m*b_m(k)+m-1 is a prime number. Here for m=3 it appears a(n) is prime for n>=2. We conjecture that a(n) is asymptotic to c*4^n with c=0.96...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005975 | OEIS-A186256

a(n) = 4*b_4(n)+3, where b_4 lists the indices of zeros of the sequence A261304: u(n) = abs(u(n-1)-gcd(u(n-1),4*n-1)), u(1) = 1.

For any fixed integer m>=1 define u(1)=1 and u(n)=abs(u(n-1)-gcd(u(n-1),m*n-1)). Then (b_m(k))_{k>=1} is the sequence of integers such that u(b_m(k))=0 and we conjecture that for k large enough m*b_m(k)+m-1 is a prime number. Here for m=4 it appears a(n) is prime for n>=1. We conjecture that a(n) is asymptotic to c*5^n with c=1.9408... See the wiki link for a sketch of a proof of this conjecture. We can give more decimals of c = 1.94080675... - _M. F. Hasler_, Aug 22 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005976 | OEIS-A186257

a(n) = 5*b_5(n)+4, where b_5 lists the indices of zeros of the sequence A261305: u(n) = abs(u(n-1)-gcd(u(n-1),5*n-1)), u(1) = 1.

For any fixed integer m>=1 define u(1)=1 and u(n)=abs(u(n-1)-gcd(u(n-1),m*n-1)). Then (b_m(k))_{k>=1} is the sequence of integers such that u(b_m(k))=0 and we conjecture that for k large enough m*b_m(k)+m-1 is a prime number. Here for m=5 it appears a(n) is prime for n>=2. We conjecture that a(n) is asymptotic to c*6^n with c>0. See the wiki link for a sketch of a proof of this conjecture. More precisely we find c = 1.15917467761... - _M. F. Hasler_, Aug 22 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005977 | OEIS-A186258

a(n) = 6*b_6(n)+5, where b_6 lists the indices of zeros of the sequence A261306: u(n) = abs(u(n-1)-gcd(u(n-1),6*n-1)), u(1) = 1.

For any fixed integer m>=1 define u(1)=1 and u(n)=abs(u(n-1)-gcd(u(n-1),m*n-1)). Then (b_m(k))_{k>=1} is the sequence of integers such that u(b_m(k))=0 and we conjecture that for k large enough m*b_m(k)+m-1 is a prime number. Here for m=6 it appears a(n) is prime for n>=1. We

conjecture that a(n) is asymptotic to c*7^n with c>0. See the wiki link for a sketch of a proof of this conjecture. We find c = 1.15917467758687... - _M. F. Hasler_, Aug 22 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005978 | OEIS-A186259

a(n) = 7*b_7(n) + 6, where b_7 lists the indices of zeros of the sequence u(n) = abs(u(n-1) - gcd(u(n-1), 7n-1)), u(1) = 1.

For any fixed integer m>=1 define u(1)=1 and u(n)=abs(u(n-1)-gcd(u(n-1),m*n-1)). Then (b_m(k))_{k>=1} is the sequence of integers such that u(b_m(k))=0 and we conjecture that for k large enough m*b_m(k)+m-1 is a prime number. Here for m=7 it appears a(n) is prime for n>=2. We conjecture that a(n) is asymptotic to c*8^n with c>0. See the wiki link for a sketch of a proof of this conjecture. We find c = 1.48462836... - _M. F. Hasler_, Aug 22 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005979 | OEIS-A186260

a(n) = 8*b_8(n)+7, where b_8 lists the zeros of the sequence A261308: u(n+1)=|u(n)-gcd(u(n), 8n+7)|, u(1)=1.

For any fixed integer m>=1 define u(1)=1 and u(n)=abs(u(n-1)-gcd(u(n-1),m*n-1)). Then (b_m(k))_{k>=1} is the sequence of integers such that u(b_m(k))=0 and we conjecture that for k large enough m*b_m(k)+m-1 is a prime number. Here for m=8 it appears a(n) is prime for n>=1. We conjecture that a(n) is asymptotic to c*9^n with c>0. See the wiki link for a sketch of a proof of this conjecture. We find c = 2.024712577430180... - _M. F. Hasler_, Aug 22 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005980 | OEIS-A186261

a(n) = 9*b_9(n) + 8, where b_9 lists the indices of zeros of the sequence A261309: u(n) = abs(u(n-1) - gcd(u(n-1), 9n-1)), u(1) = 1.

For any fixed integer m>=1 define u(1)=1 and u(n)=abs(u(n-1)-gcd(u(n-1),m*n-1)). Then (b_m(k))_{k>=1} is the sequence of integers such that u(b_m(k))=0 and we conjecture that for k large enough m*b_m(k)+m-1 is a prime number. Here for m=9 it appears a(n) is prime for n>=2. We conjecture that a(n) is asymptotic to c*10^n with c>0. See the wiki link for a sketch of a proof of this conjecture. We find c=2.59490352... - _M. F. Hasler_, Aug 22 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005981 | OEIS-A186263

a(n) = 10*b_10(n) + 9, where b_10 lists the indices of zeros of the sequence A261310: u(n) = abs(u(n-1) - gcd(u(n-1), 10n-1)), u(1) = 1.

For any fixed integer m>=1 define u(1)=1 and u(n)=abs(u(n-1)-gcd(u(n-1),m*n-1)). Then (b_m(k))_{k>=1} is the sequence of integers such that u(b_m(k))=0 and we conjecture that for k large enough m*b_m(k)+m-1 is a prime number. Here for m=10 it appears a(n) is prime for n>=1. We conjecture that a(n) is asymptotic to c*11^n with c>0. See the wiki link for a sketch of a proof of this conjecture. We find c = 2.2163823215... - _M. F. Hasler_, Aug 22 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005982 | OEIS-A186265

a(n) = b_f(n) where f is the 2-periodic sequence f(k) = (-1)^k (see comments).

Let u(1)=1 and u(n)=abs(u(n-1)-gcd(u(n-1),n+f(n))) where f(n) is a periodic sequence with period [f(1),f(2),...,f(beta)]. Then (b_f(k))_{k>=1} is the sequence of integers such that u(b_f(k))=0. We conjecture that for k large enough b_f(k)+1+f(i) is simultaneously prime for i=1,2,...,beta. Here for f(k)=(-1)^k it appears a(n) and a(n)+2 are twin primes for n>=7. If we start with u(1) large enough (such as with u(1)=71) the sequence will produce only twin primes.

Conjecture: a(n) is asymptotic to c*2^n with c>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005983 | OEIS-A186267

a(n) = b_f(n) where f is the 3-periodic sequence [-1,1,5] (see comments).

Let u(1)=1 and u(n)=abs(u(n-1)-gcd(u(n-1),n+f(n))) where f(n) is a periodic sequence with period [f(1),f(2),...,f(beta)]. Then (b_f(k))_{k>=1} is the sequence of integers such that u(b_f(k))=0. We conjecture that for k large enough b_f(k)+1+f(i) is simultaneously prime for i=1,2,...,beta. Here f is a period 3 sequence with period [-1,1,5]. It appears [a(n),a(n)+2,a(n)+6] is a prime triple for n>=14 (a(n)>=2657). Conjecture: a(n) is asymptotic to c*2^n with c>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Feb 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005984 | OEIS-A186335

A transform of the central binomial coefficients.

Conjecture: n*a(n) +(-2*n+1)*a(n-1) +5*(-n+1)*a(n-2) +3*(2*n-3)*a(n-3) +5*(n-2)*a(n-4)=0. - _R. J. Mathar_, Feb 13 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005985 | OEIS-A186338

Expansion of 1/(1-2x/(1-2x/(1-x/(1-2x/(1-2x/(1-x/(1-2x/(1-... (continued fraction).

Conjecture: (n+1)*a(n) +3*(1-4n)*a(n-1) +15*(3n-4)*a(n-2) +6*(26-11n)*a(n-3) +16*(2n-7)*a(n-4)=0. - _R. J. Mathar_, Nov 17 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005986 | OEIS-A186341

a(n)=sum{k=0..floor(n/2), binomial(n-k,k)*A186338(k)}.

Conjecture: (n+2)*a(n) +5*(-n-1)*a(n-1) +2*(-n+3)*a(n-2) +(38*n-59)*a(n-3) +(-22*n+41)*a(n-4) +4*(-22*n+81)*a(n-5) +3*(19*n-79)*a(n-6) +3*(29*n-164)*a(n-7) +2*(-17*n+98)*a(n-8) +16*(-2*n+15)*a(n-9)=0. - _R. J. Mathar_, Oct 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005987 | OEIS-A186364

Number of cycle-up-down permutations of {1,2,...,n} having no fixed points. A permutation is said to be cycle-up-down if it is a product of up-down cycles. A cycle (b(1), b(2), ...) is said to be up-down if, when written with its smallest element in the first position, it satisfies b(1)<b(2)>b(3)<... .

G.f.: conjecture: T(0), where T(k) = 1 - x^2*(k+1)*(k+2)/(x^2*(k+1)*(k+2) - 2*(1-x*k)*(1-x*(k+1))/T(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 21 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Feb 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005988 | OEIS-A186388

Adjusted joint rank sequence of (f(i)) and (g(j)) with f(i) before g(j) when f(i)=g(j), where f(i)=6*i and g(j)=j*(j+1)/2 (triangular number). Complement of A186387.

Both conjectures from _Chai Wah Wu_ hold. Proof: Let T_n = n*(n+1)/2. In the joint ranking, there are n triangular numbers up to T_n. Since multiples of 6 are ranked before T_n in case of equality, the number of positive multiples of 6 counted up to T_n is floor(T_n/6). Hence a(n) = n + floor(T_n/6) = n + floor(n*(n+1)/12). Checking the residue classes of n modulo 12 shows that this formula gives the conjectured generating function; multiplying that generating function by its denominator gives the stated recurrence. - _Kaiming Sun_, Sep 03 2026 Conjectures from _Chai Wah Wu_, May 12 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005989 | OEIS-A186390

Adjusted joint rank sequence of (f(i)) and (g(j)) with f(i) after g(j) when f(i)=g(j), where f(i)=6i and g(j)=j(j+1)/2 (triangular number). Complement of A186389.

Conjectures from _Chai Wah Wu_, May 12 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005990 | OEIS-A186420

a(n) = binomial(2n,n)^4.

We conjecture that for each integer k, the sequence defined by f_k(n) = [x^n] F(x)^(k*n) satisfies the above supercongruences, where F(x) is defined above. We conjecture that for each integer k, the sequence defined by e_k(n) = [x^n] E(x)^(k*n) also satisfies the above supercongruences. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, Feb 21 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005991 | OEIS-A186425

Antidiagonal sums of A179748.

a(n+1)/a(n) tends to the golden ratio. [Note added by _Joerg Arndt_, Mar 16 2013: this is only a conjecture so far!]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Feb 21 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005992 | OEIS-A186433

Matrix inverse of A186432.

Conjectural e.g.f.:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Feb 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005993 | OEIS-A186447

a(n)=a(floor(n/3)+a(n-1)*floor(n/4)) XOR a(floor(n/2)).

Conjecture: All finite binary words appear in the sequence infinitely many times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ben Branman_, Feb 21 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005994 | OEIS-A186522

Smallest prime factor of 2^n - 1 having the form k*n + 1.

Conjecture: a(n) exists for any integer n > 1 (verified for n = 2..300). (End) Proof of the above conjecture: By Bang's theorem, for each n > 1 except 6 there exists an odd prime p such that the multiplicative order of 2 modulo p is n, and therefore n must divide p-1. Note that a(n) <= p. - _Robert Israel_ and _Thomas Ordowski_, Sep 08 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Feb 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005995 | OEIS-A186564

Number of (n+2) X 7 0..2 arrays with every 3 X 3 subblock commuting with each horizontal and vertical neighbor 3 X 3 subblock.

G.f.: x*(12122 - 21627*x + 11695*x^2 - 19684*x^3 + 45954*x^4 - 30458*x^5 - 31744*x^6 + 15662*x^7 + 14520*x^8 - 3552*x^9 - 10944*x^10 + 16928*x^12) / ((1 - x)*(1 - x - 2*x^3 + 2*x^4 - 2*x^6)) (conjectured). - _Colin Barker_, Jan 18 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005996 | OEIS-A186740

Sequence read from antidiagonals of rectangular array with entry in row n and column q given by T(n,q) = 2^(2*n)*(Sum_{j=1..n+1} (cos(j*Pi/(2*q+1)))^(2*n)), n >= 0, q >= 1.

Conjecture: G.f. for column q is F_q(x) = (Sum_{r=0..q-1} ((q-r)*(-1)^r*binomial(2*q-r,r)*x^r)) / (Sum_{s=0..q} ((-1)^s*binomial(2*q-s,s)*x^s)), q >= 1. Conjecture: G.f. for n-th row is of the form G_n(x) = H_n(x)/(1-x)^2, where

H_n(x) is a polynomial in x.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Jan 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005997 | OEIS-A186828

Diagonal sums of number triangle A186826.

Conjecture: (n+1)*a(n) +(-7*n+2)*a(n-1) +(9*n-7)*a(n-2) +(-13*n+36)*a(n-3) +(3*n-35)*a(n-4) +(-5*n+28)*a(n-5) +(-5*n+21)*a(n-6) +(n-6)*a(n-7)=0. - _R. J. Mathar_, Feb 13 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Feb 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005998 | OEIS-A186864

Number of 5-step king's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 19 2018: (Start) The above conjectures are true. - _Stefano Spezia_, Oct 28 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-005999 | OEIS-A186886

Least number k having exactly n distinct prime divisors and the Stern polynomial B(k,x) is irreducible.

Ulas and Ulas tabulate these values and conjecture 6.5, p.20, that a(n) exists for all n. For n > 1, a(n) >= A002110(2+n)/6, conjectured to be equivalence from n=5 onward. - _Antti Karttunen_, Dec 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Feb 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006000 | OEIS-A186891

Numbers m such that the Stern polynomial B(m,x) is irreducible.

Ulas and Ulas conjecture that all primes are here. The nonprime m are in A186892. See A186886 for the least number having m prime factors.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Feb 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006001 | OEIS-A186940

Diagonal sums of number triangle A114709.

Conjecture: 3*(n+1)*a(n) +(10-17*n)*a(n-1) -6*(n+1)*a(n-2) +10*(2*n-1)*a(n-3) +9*(1-n)*a(n-4) +(n-2)*a(n-5) =0. - _R. J. Mathar_, Nov 17 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006002 | OEIS-A186997

G.f. satisfies: A(x) = 1 + x*A(x)^3 + x^2*A(x)^4.

Conjectures (both checked up to n = 100):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006003 | OEIS-A187011

Determinant of the n X n matrix with (i,j)-entry equal to |p_i-p_j|, where p_k denotes the k-th prime.

Conjecture: (-1)^(n-1)*a(n) > 0 for all n > 1, and |a(n)|^{1/n} tends to the infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006004 | OEIS-A187029

Number of 4-step one or two collinear space at a time queen's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006005 | OEIS-A187035

Diagonal sums of number triangle A187034.

Conjectured g.f.: 1/((1-x)*(1-x+x^2)*(1+x)^2) with a(n) = +a(n-2) -a(n-3) +a(n-5). - _R. J. Mathar_, Jun 30 2011 The above conjecture is correct. - _Charles R Greathouse IV_, Dec 28 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006006 | OEIS-A187043

Number of factors in expansion of infinitary perfect numbers (A007357) over distinct terms of A050376.

All terms beyond a(17) should be regarded conjectural, because they reach beyond the known terms in A007357. - _R. J. Mathar_, Mar 30 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Mar 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006007 | OEIS-A187044

Row sums of number triangle A070895.

Conjecture: a(n) -2*a(n-1) +(2-n)*a(n-2) +(n-2)*a(n-3)=0. - _R. J. Mathar_, Sep 29 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006008 | OEIS-A187049

Number of 5-step one or two space at a time bishop's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006009 | OEIS-A187050

Number of 6-step one or two space at a time bishop's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006010 | OEIS-A187051

Number of 7-step one or two space at a time bishop's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006011 | OEIS-A187054

Numbers that are not the sum of three powerful numbers (A001694).

Heath-Brown shows that this sequence is finite, resolving a conjecture of Erdos. Presumably a(12) = 119 is the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Mar 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006012 | OEIS-A187094

Good primes (version 3): p such that p-1 has a prime factor between (log p)^2 and 2 (log p)^2.

Cheng conjectures that there is a positive constant lambda such that the fraction of primes in (n - 2 sqrt(n) + 1, n + 2 sqrt(n) + 1) that are members of this sequence is greater than lambda/log log n for all sufficiently large n. If it exists, lambda <= (log 2)/2 = 0.346....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Mar 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006013 | OEIS-A187142

Smallest number k such that the continued fraction expansion of sqrt(k) contains n distinct numbers.

Conjecture: a(n) is of the form p, 2*p or 4*p, where p is prime. For the first 528 terms, a(n) is of the form 4*p only for n = 10, 11, 12, 14, 81 and 277. - _Chai Wah Wu_, Oct 04 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Joseph Stephan Orlovsky_, Mar 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006014 | OEIS-A187156

Number of 3-step one space at a time bishop's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Feb 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006015 | OEIS-A187157

Number of 4-step one space at a time bishop's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006016 | OEIS-A187158

Number of 5-step one space at a time bishop's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006017 | OEIS-A187159

Number of 6-step one space at a time bishop's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006018 | OEIS-A187160

Number of 7-step one space at a time bishop's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006019 | OEIS-A187161

Number of 8-step one space at a time bishop's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006020 | OEIS-A187164

Number of 3-step self-avoiding walks on an n X n X n cube summed over all starting positions.

Conjectures from _Colin Barker_, Apr 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006021 | OEIS-A187165

Number of 4-step self-avoiding walks on an n X n X n cube summed over all starting positions.

Conjectures from _Colin Barker_, Apr 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006022 | OEIS-A187166

Number of 5-step self-avoiding walks on an n X n X n cube summed over all starting positions.

Conjectures from _Colin Barker_, Apr 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006023 | OEIS-A187167

Number of 6-step self-avoiding walks on an n X n X n cube summed over all starting positions.

Conjectures from _Colin Barker_, Apr 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006024 | OEIS-A187168

Number of 7-step self-avoiding walks on an n X n X n cube summed over all starting positions.

Conjectures from _Colin Barker_, Apr 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006025 | OEIS-A187169

Number of 8-step self-avoiding walks on an n X n X n cube summed over all starting positions.

Conjectures from _Colin Barker_, Apr 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006026 | OEIS-A187170

Number of 9-step self-avoiding walks on an n X n X n cube summed over all starting positions.

Conjectures from _Colin Barker_, Apr 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006027 | OEIS-A187173

Number of 3-step left-handed knight's tours (moves only out two, left one) on an n X n board summed over all starting positions.

G.f.: 4*x^4*(4 - x)*(1 + x) / (1 - x)^3 (conjectured). - _Colin Barker_, Apr 22 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006028 | OEIS-A187174

Number of 4-step left-handed knight's tours (moves only out two, left one) on an n X n board summed over all starting positions.

G.f.: 8*x^4*(1 + 3*x + 7*x^2 - x^3 - x^4) / (1 - x)^3 (conjectured). - _Colin Barker_, Apr 22 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006029 | OEIS-A187175

Number of 5-step left-handed knight's tours (moves only out two, left one) on an n X n board summed over all starting positions.

G.f.: 4*x^5*(2 - x)*(2 + 12*x + 18*x^2 + 15*x^3 + 3*x^4) / (1 - x)^3 (conjectured). - _Colin Barker_, Apr 22 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006030 | OEIS-A187176

Number of 6-step left-handed knight's tours (moves only out two, left one) on an n X n board summed over all starting positions.

G.f.: 8*x^6*(11 + 29*x + 27*x^2 + 18*x^3 - 3*x^4 - 9*x^5 - 2*x^6) / (1 - x)^3 (conjectured). - _Colin Barker_, Apr 22 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006031 | OEIS-A187177

Number of 7-step left-handed knight's tours (moves only out two, left one) on an n X n board summed over all starting positions.

G.f.: 4*x^6*(4 + 80*x + 136*x^2 + 143*x^3 + 85*x^4 + 19*x^5 - 43*x^6 - 29*x^7 - 5*x^8) / (1 - x)^3 (conjectured). - _Colin Barker_, Apr 22 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006032 | OEIS-A187235

Number of ways to place n nonattacking semi-bishops on an n X n board.

Conjecture: Number of parity preserving permutations of the set {1, 2, ..., 2n+1} with exactly n+1 cycles (see A246117). - _Peter Luschny_, Feb 09 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Mar 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006033 | OEIS-A187252

Number of cycles with at least 3 alternating runs in all permutations of [n] (it is assumed that the smallest element of a cycle is in the first position).

Conjecture D-finite with recurrence a(n) +(-2*n-3)*a(n-1) +(n+7)*(n-1)*a(n-2) +4*(-n^2+2*n+1)*a(n-3) +4*(n-3)^2*a(n-4)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Mar 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006034 | OEIS-A187256

Number of peakless Motzkin paths of length n, assuming that the (1,0)-steps come in 2 colors.

Conjecture: (n+2)*a(n) -2*(2*n+1)*a(n-1) +2*(n-1)*a(n-2) +2*(5-2*n)*a(n-3) +(n-4)*a(n-4) = 0. - _R. J. Mathar_, Nov 16 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, May 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006035 | OEIS-A187259

Number of UH^jU's, DH^jD's, and DH^jU's for some j>0, in all peakless Motzkin paths of length n (here U=(1,1), D=(1,-1) and H=(1,0); can be easily expressed using RNA secondary structure terminology).

Conjecture D-finite with recurrence -(n+1)*(42968*n-187991)*a(n) +(-33354*n^2+888062*n+187991)*a(n-1) +(587317*n^2-5596253*n+61483

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, May 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006036 | OEIS-A187277

Let S denote the palindromes in the language {0,1,2,...,n-1}*; a(n) = number of words of length 4 in the language SS.

From _Colin Barker_, Jul 24 2013: (Start) (Conjectured formulas; later proven) The above conjecture is true: A284873(4, n) evaluates to the same polynomial. - _Andrew Howroyd_, Oct 10 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006037 | OEIS-A187288

Number of 3-step one or two space at a time rook's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006038 | OEIS-A187289

Number of 4-step one or two space at a time rook's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006039 | OEIS-A187290

Number of 5-step one or two space at a time rook's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006040 | OEIS-A187291

Number of 6-step one or two space at a time rook's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006041 | OEIS-A187292

Number of 7-step one or two space at a time rook's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006042 | OEIS-A187298

Number of 3-step one space leftwards or up, two space rightwards or down asymmetric rook's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006043 | OEIS-A187299

Number of 4-step one space leftwards or up, two space rightwards or down asymmetric rook's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006044 | OEIS-A187300

Number of 5-step one space leftwards or up, two space rightwards or down asymmetric rook's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006045 | OEIS-A187301

Number of 6-step one space leftwards or up, two space rightwards or down asymmetric rook's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006046 | OEIS-A187302

Number of 7-step one space leftwards or up, two space rightwards or down asymmetric rook's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006047 | OEIS-A187303

Number of 8-step one space leftwards or up, two space rightwards or down asymmetric rook's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006048 | OEIS-A187380

Number of 6-step S, NW and NE-moving king's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006049 | OEIS-A187381

Number of 7-step S, NW and NE-moving king's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006050 | OEIS-A187382

Number of 8-step S, NW and NE-moving king's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006051 | OEIS-A187383

Number of 9-step S, NW and NE-moving king's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006052 | OEIS-A187384

Number of 10-step S, NW and NE-moving king's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006053 | OEIS-A187430

Number of nonnegative walks of n steps with step sizes 1 and 2, starting and ending at 0.

Conjecture: 1 + Sum_{n>=0} a(n)*(-1)^n x^(n+1)/(1-x)^(2*n+2) = C(x), the g.f. for the Catalan numbers A000108. - _Benedict W. J. Irwin_, Jan 13 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Mar 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006054 | OEIS-A187511

Number of 6-step S, E, and NW-moving king's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006055 | OEIS-A187512

Number of 7-step S, E, and NW-moving king's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006056 | OEIS-A187514

Number of 9-step S, E, and NW-moving king's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006057 | OEIS-A187515

Number of 10-step S, E, and NW-moving king's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006058 | OEIS-A187553

Smallest prime p such that the sum of the squares of the digits of p equals n (or 0 if no such prime exists).

a(3) = 0 because the numbers of the form 10..010..01 are divisible by 3. Conjecture : except for the numbers 1 and 3, for every possible square digit sum there exists a prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Mar 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006059 | OEIS-A187563

Least k >=0 and k < prime(n) such that prime(n)*(prime(n)-k)-1 and prime(n)*(prime(n)-k)+1 are twin primes or -1 if no such k exists.

Conjectures: there are only 11 primes (5,11,13,31,37,53,61,73,79,97,127) for which k does not exist and there is always at least a pair of twin primes between prime(n) and prime(n)^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Mar 13 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006060 | OEIS-A187589

Number of 6-step E, S, NW and NE-moving king's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006061 | OEIS-A187590

Number of 7-step E, S, NW and NE-moving king's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006062 | OEIS-A187591

Number of 8-step E, S, NW and NE-moving king's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006063 | OEIS-A187592

Number of 9-step E, S, NW and NE-moving king's tours on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006064 | OEIS-A187607

Number of 3-step one space for components leftwards or up, two space for components rightwards or down asymmetric quasi-bishop's tours (antidiagonal moves become knight moves) on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006065 | OEIS-A187608

Number of 4-step one space for components leftwards or up, two space for components rightwards or down asymmetric quasi-bishop's tours (antidiagonal moves become knight moves) on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006066 | OEIS-A187609

Number of 5-step one space for components leftwards or up, two space for components rightwards or down asymmetric quasi-bishop's tours (antidiagonal moves become knight moves) on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006067 | OEIS-A187610

Number of 6-step one space for components leftwards or up, two space for components rightwards or down asymmetric quasi-bishop's tours (antidiagonal moves become knight moves) on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006068 | OEIS-A187611

Number of 7-step one space for components leftwards or up, two space for components rightwards or down asymmetric quasi-bishop's tours (antidiagonal moves become knight moves) on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006069 | OEIS-A187612

Number of 8-step one space for components leftwards or up, two space for components rightwards or down asymmetric quasi-bishop's tours (antidiagonal moves become knight moves) on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006070 | OEIS-A187679

"Connect the dots": Minimal number of circular arcs needed to connect all points of an n X n grid of points.

a(7)=9 is only a conjecture at present. I include it in the Data line to disambiguate this sequence from its neighbors. For straight line segments the answer is conjectured to be 2n-2 for all n >=3, with a(1)=1, a(2)=3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 12 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006071 | OEIS-A187714

Odd numbers m divisible by 3 such that for every k >= 1, m*2^k - 1 has a divisor in the set {5, 7, 11, 13, 17, 19, 31, 37, 41, 61, 73, 109, 151, 241, 331}.

Conjecture: 7148695169714208807 is the smallest Riesel number that is divisible by 3. - _Arkadiusz Wesolowski_, May 12 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Mar 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006072 | OEIS-A187754

Number of ways of writing the n-th twin prime p as p = q + r + s, where q >= r >= s are twin primes.

The author conjectures that a(n) >= 1 for all n >= 4. By Zhi-Wei Sun's conjecture related to A219157, for any positive integer n not among 1, 10, 430 we can write 6n-1 = p+2q = p+q+q with p,p-2,q,q+2 all prime, also for any integer n>702 we can write 6n+1 = 6(n-1)+7 = p+q+7 with p,p-2,q,q+2 all prime. Thus the author's conjecture is a consequence of Sun's conjecture. - _Zhi-Wei Sun_, Jan 06 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Fabio Mercurio_, Jan 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006073 | OEIS-A187757

Number of ways to write n=x+y (x,y>0) with 6x-1, 6x+1, 6y+1 and 6y+5 all prime.

Conjecture: a(n)>0 for all n>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006074 | OEIS-A187758

Number of ways to write n=x+y (x,y>0) with 2x-3, 2x+3, 6y+1 and 6y+5 all prime.

Conjecture: a(n)>0 for all n>4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006075 | OEIS-A187759

Number of ways to write n=x+y (0<x<y<n) with 6x-1, 6x+1, 6y-1 and 6y+1 all prime.

Conjecture: If n>200 is not among 211, 226, 541, 701, then a(n)>0. This essentially follows from the conjecture related to A219157, since n=x+y for some positive integers x and y with 6x-1,6x+1,6y-1,6y+1 all prime if and only if 6n=p+q for some twin prime pairs {p,p-2} and {q,q+2}. Similarly, the conjecture related to A218867 implies that any integer n>491 can be written as x+y (0<x<=y<n) with 6x+1, 6x+5, 6y+1 and 6y+5 all prime; and the conjecture related to A219055 implies that any integer n>1600 not among 2729 and 4006 can be written as x+y (0<x<=y<n) with 2x-3, 2x+3, 2y-3 and 2y+3 all prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006076 | OEIS-A187761

Number of maps f: [n] -> [n] with f(x)<=x and f(f(x)) = f(f(f(x))).

Conjecture (confirmed below) about the e.g.f. A(x): Let B(x) = Sum_{n>=0} b(n) * x^n/n! = exp(exp(x)-1) the e.g.f. of the Bell numbers (A000110). Then A(x) = Sum_{n>=0} a(n) * x^n/n! = exp(Sum_{n>=0} b(n) * x^(n+1)/(n+1)!), see PARI program. Conjecture (confirmed below): Let C(0,x) = 1 and for k>=1 C(k, x) = exp(integral(C(k-1,x))) then C(k,x) is the e.g.f. for monotonic-labeled forests on n vertices with rooted trees of height less than k (column k of A179455(n,k) for k>=1, n>=k). For k=1 (C(1,x)=exp(x)) and k=2 (C(2,x)=exp(exp(x)-1)) this is known to be true, for k=3 this is the conjecture above. (End) "Which means the conjecture is right. (The integral turns that into 'multisets of things whose sizes plus 1 add up to n'; a tree is a forest together with a new node on top.)"

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Jan 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006077 | OEIS-A187767

Number of bicolored cyclic patterns n X n.

a(1) = 0; a(n) = 2^(n-1)-1 if n is odd, 2^(n-1)+a(n/2) if n is even (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Resta_, Jan 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006078 | OEIS-A187772

T(n,k) = (n-1)*A220555(n,k), n,k = 2,3,....

For n,k >= 2, T(n,k) is equal to the maximal period p of the corresponding sequence (w.r.t. n and k) defined in Conjecture 1 in A220555. T(2,k) = A001175(k) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Jan 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006079 | OEIS-A187779

Numbers n for which the terms of A014320 up to n=A014320(k) are a permutation of {1,2,4,6,...,2*(k-1)}.

An old conjecture states that the equation prime(n+1)-prime(n)=2m is solvable for m=1/2,1,2,... . a(n) indicate the smallest term of A014320 such that among the previous terms there occur 1 and all previous even numbers<=a(n). The primes up to 10^8 do not generate any further terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter J. C. Moses_, Jan 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006080 | OEIS-A187785

Number of ways to write n=x+y (x,y>=0) with {6x-1,6x+1} a twin prime pair and y a triangular number.

Conjecture: a(n)>0 for all n>48624 not equal to 76106. Compare the conjecture with another Sun's conjecture associated with A132399.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006081 | OEIS-A187808

a(n) = |{0<=k<n: 2k+3, n(n-k)-1, n(n+k)-1 are all prime}|.

Conjecture: a(n)>0 for all n>1. Moreover, if n>5 is different from 9, 191, 329, 641, 711, 979, then 2k-3, 2k+3, n(n-k)-1, n(n+k)-1 are all prime for some 0<k<n. Zhi-Wei Sun also made the following conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006082 | OEIS-A187809

a(n) is the smallest prime(m) such that the interval (prime(m)*n, prime(m+1)*n) contains exactly two primes.

Conjecture. In the supposition that there are infinitely many twin primes, every term is 2 or in A001359 (lesser of twin primes).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006083 | OEIS-A187810

a(n) is the smallest prime(m) such that the interval (prime(m)*n, prime(m+1)*n) contains exactly three primes.

Conjecture. In the supposition that there are infinitely many twin primes, every term beginning the fourth is 2 or in A001359 (lesser of twin primes).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006084 | OEIS-A187812

a(n) is the smallest prime(m) such that the interval (prime(m)*n, prime(m+1)*n) contains exactly four primes.

Conjecture. In the supposition that there are infinitely many twin primes, every term beginning with the sixth is 2 or in A001359 (lesser of twin primes).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006085 | OEIS-A187815

Number of permutations q_1, ..., q_7 of the 7 consecutive primes p_n, p_{n+1}, ..., p_{n+6} with q_1 = p_n and q_7 = p_{n+6}, and with |q_1-q_2|, |q_2-q_3|, ..., |q_6-q_7|, |q_7-q_1| pairwise distinct, where p_k denotes the k-th prime.

See also A185645 for a related conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006086 | OEIS-A187822

Smallest k such that the partial sums of the divisors of k (taken in increasing order) contain exactly n primes.

It appears that a(n) is even for n > 0 and nonsquarefree for n > 1. We also conjecture that there is an infinite subsequence of squares 1, 4, 16, 64, 2304, 14400, 57600, 112896, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Dec 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006087 | OEIS-A187851

Number of 3-step king-knight's tours (piece capable of both kinds of moves) on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006088 | OEIS-A187852

Number of 4-step king-knight's tours (piece capable of both kinds of moves) on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006089 | OEIS-A187853

Number of 5-step king-knight's tours (piece capable of both kinds of moves) on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006090 | OEIS-A187854

Number of 6-step king-knight's tours (piece capable of both kinds of moves) on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006091 | OEIS-A187858

Number of 3-step one space for components leftwards or up, two space for components rightwards or down asymmetric quasi-queen's tours (antidiagonal moves become knight moves) on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006092 | OEIS-A187859

Number of 4-step one space for components leftwards or up, two space for components rightwards or down asymmetric quasi-queen's tours (antidiagonal moves become knight moves) on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006093 | OEIS-A187860

Number of 5-step one space for components leftwards or up, two space for components rightwards or down asymmetric quasi-queen's tours (antidiagonal moves become knight moves) on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006094 | OEIS-A187861

Number of 6-step one space for components leftwards or up, two space for components rightwards or down asymmetric quasi-queen's tours (antidiagonal moves become knight moves) on an n X n board summed over all starting positions.

Conjectures from _Colin Barker_, Apr 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006095 | OEIS-A187866

Greatest k such that prime(n)*(prime(n)-k)-1 and prime(n)*(prime(n)-k)+1 are twin primes, k >= 0 and k < prime(n) or -1 if no such k exists.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Mar 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006096 | OEIS-A187916

a(n) = C(n) if n is odd, else C(n) - C(n/2); C(n) are Catalan numbers.

Conjecture: -(n-2)*(n+2)*(n+1)*a(n)
+8*(n+1)*(n^2-3*n+1)*a(n-1)
+4*(-3*n^3+18*n^2-22*n+2)*a(n-2)
-32*(n-2)*(n^2-3*n+1)*a(n-3) +16*

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Mar 15 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006097 | OEIS-A187941

Least number with exactly n even divisors.

Conjecture: a(n) = 2^n only if n is prime or if n = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Mar 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006098 | OEIS-A187968

Positions of 1 in A187967; complement (conjectured) of A098021.

Positions of 1 in A187967; complement (conjectured) of A098021.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006099 | OEIS-A188039

Positions of 0 in A188038; complement of A188040.

There is (conjecturally) a connection a(1+n) = f(n) where f(n) = 3*n +2 +2*floor(n*sqrt 2) is defined in A120861. Tested numerically up to n=40000. - _R. J. Mathar_, Jul 22 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006100 | OEIS-A188070

Positions of 1 in A188068; complement of A188069.

Conjectured partial sums of A245222. - _Sean A. Irvine_, Jun 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006101 | OEIS-A188145

Solutions of the equation n" - n' - n = 0, where n' and n" are the first and second arithmetic derivatives (see A003415).

Solutions of the similar equation n"-n'+n=0 are 30, 858, 1722, etc., apparently Giuga numbers whose derivative is a prime number. In fact the equation can be rewritten as n'=n+n" and if n"=1 it is the conjecture in A007850.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_, Mar 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006102 | OEIS-A188148

Number of 3-step self-avoiding walks on an n X n square summed over all starting positions.

Conjectures from _Colin Barker_, Apr 26 2018: (Start) The above conjectures are correct. See A188147.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006103 | OEIS-A188149

Number of 4-step self-avoiding walks on an n X n square summed over all starting positions.

Conjectures from _Colin Barker_, Apr 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006104 | OEIS-A188150

Number of 5-step self-avoiding walks on an n X n square summed over all starting positions.

Conjectures from _Colin Barker_, Apr 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006105 | OEIS-A188151

Number of 6-step self-avoiding walks on an n X n square summed over all starting positions.

Conjectures from _Colin Barker_, Apr 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006106 | OEIS-A188152

Number of 7-step self-avoiding walks on an n X n square summed over all starting positions.

Conjectures from _Colin Barker_, Apr 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006107 | OEIS-A188153

Number of 8-step self-avoiding walks on an n X n square summed over all starting positions.

Conjectures from _Colin Barker_, Apr 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006108 | OEIS-A188154

Number of 9-step self-avoiding walks on an n X n square summed over all starting positions.

Conjectures from _Colin Barker_, Apr 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006109 | OEIS-A188155

Number of 10-step self-avoiding walks on an n X n square summed over all starting positions.

Conjectures from _Colin Barker_, Apr 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006110 | OEIS-A188266

Coefficient of x^n in the series 1/F(-1/2,1/2;1;16x), where F(a1,a2;b;x) is the hypergeometric series.

Conjecture: a(n) ~ Pi * 2^(4*n-3) / n^2. - _Vaclav Kotesovec_, Apr 12 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, Mar 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006111 | OEIS-A188288

In lunar arithmetic in base 2, the number of divisors of the number 11...1101 (n digits, the binary expansion of 2^n-3).

a(1)=1 by convention. The g.f. is only a conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Adam S. Jobson_ and _N. J. A. Sloane_, Mar 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006112 | OEIS-A188312

Expansion of (1/(1-x^2))*c(x/((1-x)*(1-x^2))) where c(x) is the g.f. of A000108.

Conjecture: (n+1)*a(n) +(3-4*n)*a(n-1) + (7-6*n)*a(n-2) -a(n-3) +(n-4)*a(n-4)=0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006113 | OEIS-A188314

Expansion of (1/(1-x))*c(x/((1-x)*(1-x^2))) where c(x) is the g.f. of A000108.

Conjecture: (n+1)*a(n) +(n+2)*a(n-1) +(42-26*n)*a(n-2) +30*(3-n)*a(n-3) +(n-5)*a(n-4) +5*(n-6)*a(n-5)=0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Mar 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006114 | OEIS-A188342

Smallest odd primitive abundant number (A006038) having n distinct prime factors.

a(15) <= 279258050855730883 2935, a(16) <= 428525983200229616718445, a(17) <= 421623204340052009840800 45. If these a(n) are squarefree and don't have a greatest prime factor more than 3 primes away from that of the preceding term, then these bounds are the actual values of a(n). The PARI code needs only fractions of a second to compute further bounds, which under the given hypotheses are the actual values of a(n). - _M. F. Hasler_, Jul 17 2016 This conjecture is correct up to a(50). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006115 | OEIS-A188350

Number of arrangements of n+1 numbers x(i) in -1..1 with the sum of x(i)*x(i+1) equal to zero.

Conjectures from _Mikhail Smirnov_, Oct 15 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006116 | OEIS-A188458

Expansion of e.g.f. exp(x)/cosh(2*x).

G.f.: conjecture T(0)/(1-x), where T(k) = 1 - 4*x^2*(k+1)^2/(4*x^2*(k+1)^2 + (1-x)^2/T(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 12 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006117 | OEIS-A188460

Diagonal sums of number triangle A119308.

Conjecture: (n+5)*(n+1)*a(n) -(2*n+5)*(n+2)*a(n-1) -(n+2)*(n+1)*a(n-2) -(n+1)*(2*n+1)*a(n-3) +(n-2)*(n+2)*a(n-4)=0. - _R. J. Mathar_, Feb 13 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006118 | OEIS-A188462

Least number of 5th powers needed to represent n.

Vaughan & Wooley (1995) prove that a(n) <= 17 for large enough n; in fact it is conjectured that a(n) <= 6 for large enough n. The maximum value is a(223) = 37. - _Charles R Greathouse IV_, Jul 05 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Apr 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006119 | OEIS-A188464

Diagonal sums of triangle A188463.

Conjecture: (n+4)*a(n)-(4*n+9)*a(n-1) +(2*n-1)*a(n-2) -a(n-3) +(n-3)*a(n-4)=0. - _R. J. Mathar_, Nov 17 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006120 | OEIS-A188474

A generalized Deutsch triangle.

Row sums are A137398(n+1) (conjecture). Diagonal sums are A188476.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006121 | OEIS-A188476

Diagonal sums of number triangle A188474.

Conjecture: (n+3)*a(n) +(-4*n-9)*a(n-1) +n*a(n-2) -3*a(n-3) +3*(5*n+1)*a(n-4) +6*(n)*a(n-5) +6*(n-1)*a(n-6) +4*(-n+3)*a(n-7)=0. - _R. J. Mathar_, Dec 03 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Barry_, Apr 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006122 | OEIS-A188482

Diagonal sums of the Riordan matrix (1/(1-4x),(1-sqrt(1-4x))/(2*sqrt(1-4x))) (A188481).

Conjecture: (-n+1)*a(n) + (7*n-9)*a(n-1) + 2*(-4*n+7)*a(n-2) + (-15*n+23)*a(n-3) + 2*(-2*n+3)*a(n-4) = 0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, Apr 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006123 | OEIS-A188483

Numbers whose squares have an odd square abundance.

Kravitz conjectured that no numbers exist whose abundance is a positive odd square. The squares of this sequence are counterexamples.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ed Pegg Jr_, Apr 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006124 | OEIS-A188484

Numbers n (other than powers of 2) such that abs(abundance(n)) is an odd square.

Superset of Kravitz conjecture counterexamples (A188486). Abundances of these numbers are A188487.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Apr 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006125 | OEIS-A188486

Numbers k such that abundance(k) is an odd square.

Counterexamples to the Kravitz conjecture. Subsequence of A188484 with positive abundances. Abundances are A188488, sigma(k) - 2*k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Apr 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006126 | OEIS-A188516

Number of nX2 binary arrays without the pattern 1 1 0 diagonally, vertically or horizontally.

Conjecture: a(n) = (F(n+3) - 1)^2, where F = A000045 (Fibonacci numbers). - _Clark Kimberling_, Jun 21 2016 Assuming the conjecture, define b(1) = 1 and b(n) = a(n-1) for n > 1. Then b(n) = Sum{F(i,j): (i=n and 1<=j<=n) or (j=n and 1<=i<=n)}, where F is the Fibonacci fusion array, A202453. - _Clark Kimberling_, Jun 21 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006127 | OEIS-A188541

a(n) = 2 * A079500(n) - A079500(n+1).

Arises in studying a conjecture related to lunar divisors in base 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006128 | OEIS-A188553

T(n,k) = Number of n X k binary arrays without the pattern 0 1 diagonally, vertically, antidiagonally or horizontally.

Such a bijection implies that all the empirical/conjectured formulas in A188554, A188555, A188556, A188557, A188558, and A188559 become correct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006129 | OEIS-A188555

Number of 4 X n binary arrays without the pattern 0 1 diagonally, vertically, antidiagonally or horizontally.

The number of such words satisfy the recurrence given below and, hence, the empirical/conjectured formulas become true. (End) Conjectures from _Colin Barker_, Apr 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006130 | OEIS-A188556

Number of 5 X n binary arrays without the pattern 0 1 diagonally, vertically, antidiagonally or horizontally.

Conjectures from _Colin Barker_, Apr 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006131 | OEIS-A188557

Number of 6 X n binary arrays without the pattern 0 1 diagonally, vertically, antidiagonally or horizontally.

Conjectures from _Colin Barker_, Apr 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006132 | OEIS-A188558

Number of 7 X n binary arrays without the pattern 0 1 diagonally, vertically, antidiagonally or horizontally.

Conjectures from _Colin Barker_, Apr 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006133 | OEIS-A188559

Number of 8 X n binary arrays without the pattern 0 1 diagonally, vertically, antidiagonally or horizontally.

Conjectures from _Colin Barker_, Apr 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006134 | OEIS-A188572

a(n) = coefficient of sqrt(3) in the expansion of (1 + sqrt(2) + sqrt(3))^n sequence.

Conjecture: every prime divides a(n) for infinitely many n, and if K(p) = (k(1), k(2),...) is the maximal subsequence of indices n such that p divides a(n), then the difference sequence of K(p) is eventually periodic; indeed, K(p) is purely periodic for the first 6 primes, with respective period lengths 6,8,5,4,4,8 and these periods: Conjectures from _R. J. Mathar_, Jan 09 2013: (Start) The conjectures by Mathar are true. See link. - _Sela Fried_, Jan 01 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mateusz Szymański_, Dec 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006135 | OEIS-A188597

Odd deficient numbers whose abundancy is closer to 2 than any smaller odd deficient number.

a(38) = 1412875743840099 is the next term after 9 not divisible by 15. The number of terms not divisible by 7 is conjectured to be infinite provided no odd perfect numbers exist. - _Alexander Violette_, Sep 26 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006136 | OEIS-A188644

Array of (k^n + k^(-n))/2 where k = (sqrt(x^2-1) + x)^2 for integers x >= 1.

Conjecture: Given the function f(x,y) = (sqrt(x^2+y) + x)^2 and constant k=f(x,y), then for all integers x >= 1 and y=[+-]1, k may be irrational, but (k^n + k^(-n))/2 always produces integer sequences; y=-1 results shown here; y=1 results are A188645.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles L. Hohn_, Apr 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006137 | OEIS-A188646

Array of a(n)=a(n-1)*k-((k-1)/(k^n)) where a(0)=1 and k=(sqrt(x^2-1)+x)^2 for integers x>=1.

Conjecture: Given function f(x, y)=(sqrt(x^2+y)+x)^2; constant k=f(x, y); and initial term a(0)=1; then for all integers x>=1 and y=[+-]1, k may be irrational, but sequence a(n)=a(n-1)*k-((k-1)/(k^n)) always produces integer sequences; y=-1 results shown here; y=1 results are A188647.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles L. Hohn_, Apr 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006138 | OEIS-A188647

Array read by antidiagonals of a(n) = a(n-1)*k-((k-1)/(k^n)) where a(0)=1 and k=(sqrt(x^2+1)+x)^2 for integers x>=1.

Conjecture: Given function f(x, y)=(sqrt(x^2+y)+x)^2; constant k=f(x, y); and initial term a(0)=1; then for all integers x>=1 and y=[+-]1, k may be irrational, but sequence a(n)=a(n-1)*k-((k-1)/(k^n)) always produces integer sequences; y=1 results shown here; y=-1 results are A188646.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles L. Hohn_, Apr 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006139 | OEIS-A188648

Binomial sums a(n) = Sum_{k=0..n} (binomial(2n-k,k))^2.

Conjecture: a(n) = hypergeom([-n,-n,n+1,n+1], [1/2,1/2,1], 1/16). - _Velin Yanev_, Oct 31 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, Apr 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006140 | OEIS-A188672

a(n) is the least r > 1 for which the interval (r*n, r*(n+1)) contains no prime powers (p^k, k >= 1), or a(n) = 0 if no such r exists.

Conjecture: a(n) = 0, iff n = 1, 2, 3, 4, 5, 6, 9, 12, 13, 14, 24, 56.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Aug 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006141 | OEIS-A188785

Number of 2-step self-avoiding walks on an n X n X n X n 4-cube summed over all starting positions.

Conjectures from _Colin Barker_, Apr 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006142 | OEIS-A188786

Number of 3-step self-avoiding walks on an n X n X n X n 4-cube summed over all starting positions.

Conjectures from _Colin Barker_, Apr 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006143 | OEIS-A188787

Number of 4-step self-avoiding walks on an n X n X n X n 4-cube summed over all starting positions.

Conjectures from _Colin Barker_, Apr 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006144 | OEIS-A188788

Number of 5-step self-avoiding walks on an n X n X n X n 4-cube summed over all starting positions.

Conjectures from _Colin Barker_, Apr 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006145 | OEIS-A188789

Number of 6-step self-avoiding walks on an n X n X n X n 4-cube summed over all starting positions.

Conjectures from _Colin Barker_, Apr 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006146 | OEIS-A188793

Start with 1 and 5, then repeatedly adjoin the smallest number that is greater than the last term and not equal to the sum of a subset of the existing terms.

(1) any number from 5 to 10*3^k + 3 (by inductive hypothesis),

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006147 | OEIS-A188819

Number of n X 3 binary arrays without the pattern 0 1 diagonally or antidiagonally.

Conjectures from _Colin Barker_, Apr 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006148 | OEIS-A188820

Number of n X 5 binary arrays without the pattern 0 1 diagonally or antidiagonally.

Conjectures from _Colin Barker_, Apr 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006149 | OEIS-A188821

Number of n X 6 binary arrays without the pattern 0 1 diagonally or antidiagonally.

Conjectures from _Colin Barker_, Apr 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006150 | OEIS-A188822

Number of n X 7 binary arrays without the pattern 0 1 diagonally or antidiagonally.

Conjectures from _Colin Barker_, Apr 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006151 | OEIS-A188861

Number of n X 4 binary arrays without the pattern 0 1 diagonally, vertically or antidiagonally.

Conjectures from _Colin Barker_, Feb 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 12 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006152 | OEIS-A188865

Number of n X 8 binary arrays without the pattern 0 1 diagonally, vertically or antidiagonally.

Conjectures from _Colin Barker_, Feb 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 12 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006153 | OEIS-A188934

Decimal expansion of (1+sqrt(17))/4.

Conjecture: This number is an eigenvalue to infinitely many n*n submatrices of A191898, starting in the upper left corner, divided by the row index. For the first few characteristic polynomials see A260237 and A260238. - _Mats Granvik_, May 12 2016.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 13 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006154 | OEIS-A189007

Number of ON cells after n generations of the 2D cellular automaton described in the comments.

Conjecture: a(1) = 1; a(n) = 2^A056791(n-1) for n > 1. - _Michael De Vlieger_, Nov 02 2022 It is not difficult to show that the preceding conjecture is correct. In fact one can give an explicit formula for the n-th term. At generation n >= 2, the configuration of ON cells consists of a set of concentric diamonds (see the illustration). The sizes of the diamonds are given by the (n-2)nd term of A245191. Let N = A245191(n-2) = Sum_{i>=0} b_i*2^i. Then the ON cells form a set of diamonds with edge-lengths i+2 for each b_i = 1. The i-th diamond contains 4*(i+1) ON cells, and the total number of ON cells is therefore a(n) = 4*Sum_i (i+1)*b_i. The b_i are given explicitly in A245191.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Apr 15 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006155 | OEIS-A189034

Positions of 0 in the zero-one sequence s based on the sequence lower Wythoff sequence p: s(p(k))=s(k); s(q(k))=1-s(k); s(1)=0, q=(upper Wythoff sequence).

It is conjectured at A095076 that the zero-sequence s is A095076.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 15 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006156 | OEIS-A189053

Expansion of d/dx log(1/(1-x/sqrt(1-4*x^2))).

Conjecture: n*a(n) +(n-1)*a(n-1) +(-13*n+12)*a(n-2) +(-13*n+25)*a(n-3) +4*(14*n-27) *a(n-4) +4*(14*n-41)*a(n-5) +80*(-n+3)*a(n-6) +80*(-n+4)*a(n-7)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Apr 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006157 | OEIS-A189123

Positive integers n such that A005245(n-i) + A005245(i) < A005245(n-1) + 1 for some i, 1 < i < n.

Counterexamples to the first conjecture of David Wilson on A005245.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Apr 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006158 | OEIS-A189124

Positive integers n such that A005245(n) < min{ A005245(n-1) + 1, A005245(x) + A005245(y) } where x,y range over positive integers with x*y = n.

Counterexamples to the second conjecture of David Wilson on A005245.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Apr 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006159 | OEIS-A189125

Primes p such that A005245(p) < A005245(p-1)+1.

Counterexamples to the conjecture from section F26 in UPINT.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Apr 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006160 | OEIS-A189176

Row sums of the Riordan matrix (1+x/sqrt(1-4*x),(1-sqrt(1-4*x))/2) (A189175).

Conjecture: (n+1)*a(n) + 3*(1-3*n)*a(n-1) + 10*(2*n-3)*a(n-2) = 0. - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, Apr 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006161 | OEIS-A189202

Let s_k(n) denote the sum of digits of n in base k. Then a(n) is the smallest m>0 such that both s_2(m*(n-1)) and s_n(2*m*(n-1))/(n-1) are even, or a(n)=0, if such m does not exist.

Conjecture: For all n>=2, a(n)>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006162 | OEIS-A189281

Number of permutations p of 1,2,...,n satisfying p(i+2) - p(i) <> 2 for all 1 <= i <= n-2.

The conjectured recurrence would imply the asymptotic expansion for a(n)/n! ~ There are good reasons to believe the conjecture is correct. (It has the expected form.) The problem is one of counting Hamiltonian cycles in the complement of some simple graph. There is a method for counting these efficiently (although I have not implemented in code). Similar to A242522 / A229430. - _Andrew Howroyd_, Mar 06 2022 a(40)-a(300) were computed using an independent solution (dynamic programming, O(N^4) per term), and the conjectured recurrence was further confirmed to be correct up to n=300. Consequently, the keyword "hard" is removed. - _Rintaro Matsuo_, Oct 18 2022

Conjectured recurrence of degree 11 and order 8: (262711*n + 1387742*n^2 - 824875*n^3 - 1855253*n^4 - 111530*n^5 + 680983*n^6 + 364242*n^7 + 84992*n^8 + 10332*n^9 + 640*n^10 + 16*n^11)*a(n) + (-1050844*n - 9705192*n^2 - 7414683*n^3 + 3536494*n^4 + 6459004*n^5 + 3326393*n^6 + 903534*n^7 + 144684*n^8 + 13756*n^9 + 720*n^10 + 16*n^11)*a(n+1) + (3492344 - 2212342*n - 8507169*n^2 - 11544227*n^3 - 12034116*n^4 - 8216995*n^5 - 3442049*n^6 - 890050*n^7 - 142300*n^8 - 13660*n^9 - 720*n^10 - 16*n^11)*a(n+2) + (19817984 + 45323852*n + 825228*n^2 - 57004661*n^3 - 57059306*n^4 - 28077270*n^5 - 8398637*n^6 - 1631510*n^7 - 207980*n^8 - 16828*n^9 - 784*n^10 - 16*n^11)*a(n+3) + (9586160 + 6680237*n - 13772613*n^2 - 27689586*n^3 - 22162455*n^4 - 9855085*n^5 - 2629562*n^6 - 427656*n^7 - 41332*n^8 - 2176*n^9 - 48*n^10)*a(n+4) + (22192864 + 44710768*n - 2924668*n^2 - 52385912*n^3 - 45161616*n^4 - 18784740*n^5 - 4549208*n^6 - 674256*n^7 - 60400*n^8 - 3008*n^9 - 64*n^10)*a(n+5) + (557152 - 2032472*n - 2937392*n^2 - 1594200*n^3 - 517688*n^4 - 122032*n^5 - 19856*n^6 - 1792*n^7 - 64*n^8)*a(n+6) + (3786960 + 7105324*n - 1191064*n^2 - 8059160*n^3 - 5938996*n^4 - 2073752*n^5 - 402736*n^6 - 44528*n^7 - 2624*n^8 - 64*n^9)*a(n+7) + (-598208 - 943004*n + 414196*n^2 + 1213772*n^3 + 728648*n^4 + 203584*n^5 + 29616*n^6 + 2176*n^7 + 64*n^8)*a(n+8) = 0. This recurrence correctly predicted the four new terms in the b-file. - _Christoph Koutschan_, Feb 19 2022 The preceding conjectured recurrence is equivalent to the following, which has degree 3 and order 13, and was obtained by _Doron Zeilberger_ and then reformatted by _Manuel Kauers_ (it uses Mathematica syntax): Conjecture: ((-1 + n)^2*n*a[n])/4 + (n*(-16 + 38*n + 11*n^2)*a[1 + n])/16 +

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Apr 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006163 | OEIS-A189286

a(n):=(Sum_{k=0}^n C(6k,3k)C(3k,k)C(6(n-k),3(n-k))C(3(n-k),n-k))/((2n-1)Binomial[3n,n]).

On Apr 19 2011, _Zhi-Wei Sun_ conjectured that a(n) is an integer for every n=0,1,2,.... He proved that a(p-1)=[(p+1)/6] (mod p) for any prime p, and also made the following conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006164 | OEIS-A189359

Number of homogeneous games for n players.

Conjecture: g.f.: Q(0)*x/(1-x), where Q(k) = 1 + (1-(1-x)^(2*k+2))/(1- (1-(1-x)^(2*k+3))/(1-(1-x)^(2*k+3) + 1/Q(k+1))); (continued fraction). - _Sergei N. Gladkovskii_, May 13 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Fabián Riquelme_, Apr 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006165 | OEIS-A189377

a(n) = n + floor(ns/r) + floor(nt/r) with r=2, s=(-1+sqrt(5))/2, t=(1+sqrt(5))/2.

(Conjecture) These are the numbers n such that (n+1)-sections of the Fibonacci word contain "000" (the commoner bit) but not "111" (the rarer bit). - _Don Reble_, Apr 07 2021 Conjecture proved April 8 2021, using the Walnut theorem prover. - _Jeffrey Shallit_, Apr 09 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006166 | OEIS-A189379

n+[nr/t]+[ns/t]; r=2, s=(-1+sqrt(5))/2, t=(1+sqrt(5))/2.

(Conjecture) These are the numbers n such that (n+1)-sections of the Fibonacci word contain both "000" and "111". - _Don Reble_, Apr 07 2021 Conjecture proved April 8 2021 using the Walnut theorem prover. - _Jeffrey Shallit_, Apr 09 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006167 | OEIS-A189391

The minimum possible value for the apex of a triangle of numbers whose base consists of a permutation of the numbers 0 to n, and each number in a higher row is the sum of the two numbers directly below it.

Conjecture: a(n) is also half the sum of the "cuts-resistance" (see A319416, A319420, A319421) of all binary vectors of length n (see Lenormand, page 4). - _N. J. A. Sloane_, Sep 20 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nathaniel Johnston_, Apr 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006168 | OEIS-A189450

Number of 2 X n array permutations with each element moving zero or one space horizontally or diagonally.

Conjectures from _Colin Barker_, May 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006169 | OEIS-A189457

a(n) = n+[ns/r]+[nt/r]; r=2, s=sqrt(2), t=1+sqrt(2).

With r=2, s=sqrt(2), t=1+sqrt(2), a=A189457, b=A189458, c=A186222 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006170 | OEIS-A189572

Fixed point of the morphism 0->01, 1->001.

Proof of Kimberling's conjecture: one can use Lemma 1 in the Bosma, Dekking, Steiner paper, which gives the positions of 1 in a Sturmian sequence as a Beatty sequence. - _Michel Dekking_, Nov 03 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006171 | OEIS-A189573

Positions of 0 in A189572; complement of A080652 (conjectured).

Positions of 0 in A189572; complement of A080652 (conjectured). The conjecture is proved in A189572. - _Michel Dekking_, Nov 03 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006172 | OEIS-A189576

Fixed point of the morphism 0->01, 1->110.

Some lower-numbered entries are conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006173 | OEIS-A189628

Fixed point of the morphism 0->001, 1->010.

Some lower-numbered entries are conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006174 | OEIS-A189640

Fixed point of the morphism 0->001, 1->101.

Proof of Mathar's third conjecture: Let alpha be the defining morphism for (a(n)): alpha(0)=001, alpha(1)=101. Let beta be the defining morphism for A080846: beta(0)=010, beta(1)=011. To prove the conjecture, it suffices to prove that

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006175 | OEIS-A189665

Positions of 0's in A189664; complement of A189666.

2*a(n) = A284794(n) for n>=1. (Conjectured by Clark Kimberling, see A189664). - _Michel Dekking_, Oct 15 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006176 | OEIS-A189668

Fixed point of the morphism 0->010, 1->100.

Conjecture: a(n) = 2*n - 1 - A285347(n). This conjecture is correct, and proved in A285347. - _Michel Dekking_, Sep 07 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006177 | OEIS-A189759

Numbers pqr such that pq + pr + qr is prime, where p, q, and r are primes.

As mentioned by Ufnarovski and Ahlander, if pq+pr+qr is prime, then the arithmetic derivative (A003415) of pqr is that prime. They conjecture that this sequence and A087054 are infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006178 | OEIS-A189760

Least nonnegative number whose n-th arithmetic derivative (A003415) is zero and lower derivatives are nonzero.

Question: Are there indefinitely long sequences of iterations of A003415 that end with steps ... -> p -> 1 -> 0, with p=5? Are there such sequences for any other prime p? Can we construct a such sequence that is guaranteed to be infinite? See the subtree depicted in A327975 and conjecture #8 in Ufnarovski and Åhlander paper.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006179 | OEIS-A189786

a(n) = n + [nr/t] + [ns/t]; r=Pi/2, s=arcsin(5/13), t=arcsin(12/13).

Conjecture: Sequence consists of all the positive even numbers except numbers of the form 8*x+6, x >= 0. - _Harvey P. Dale_, Dec 07 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006180 | OEIS-A189807

Number of right triangles on an (n+1) X 3 grid.

Conjectures from _Colin Barker_, May 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006181 | OEIS-A189808

Number of right triangles on a (n+1) X 4 grid.

Conjectures from _Colin Barker_, May 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006182 | OEIS-A189809

Number of right triangles on an (n+1) X 5 grid.

Conjectures from _Colin Barker_, May 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006183 | OEIS-A189827

a(n) = d(n-1) + d(n+1), where d(k) is the number of divisors of k.

It is conjectured that every number greater than 3 occurs an infinite number of times. Note that an infinite number of 4's is equivalent to there being an infinite number of twin primes (A001097). An infinite number of 5's is equivalent to there being an infinite number of primes of the form p^2-2 (A028871) or p^2+2 (A056899) for prime p. An infinite number of 6's is equivalent to there being an infinite number of primes of the form p^3-2 (A066878), p^3+2 (A048636), p*q-2 (A063637), or p*q+2 (A063638), where p and q are distinct primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006184 | OEIS-A189830

Pairs of numbers i,j ordered by increasing i, such that 2 <= j < i, gcd(i,j)=1 and gcd(Phi_j(i), Phi_i(j))=2*i*j+1, where Phi_k(t) is the k-th cyclotomic polynomial.

The second pair i=3313, j=17 is the only known counterexample to a conjecture of Stephens that gcd(Phi_p(q), Phi_q(p))=1 for every pair of prime numbers (p,q). This is related to a conjecture of Feit-Thompson. See the corresponding wiki page. Up to i=9400 there are no new terms of the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luis H. Gallardo_, Apr 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006185 | OEIS-A189912

Extended Motzkin numbers, Sum_{k>=0} C(n,k)*C(k), where C(k) is the extended Catalan number A057977(k).

Conjecture: a(n) = Sum_{k=0..floor(n/2)} (n+1-2*k)*A055151(n,k). - _Werner Schulte_, Oct 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, May 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006186 | OEIS-A189925

Expansion of theta_4/theta_3 in powers of q.

Conjecture: Sum_{k>=0} a(k) / exp(k*Pi) = (1/2)*2^(3/4) = A228497. - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, May 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006187 | OEIS-A189974

Numbers m such that d(m-1) = d(m+1) = 4, where d(k) is the number of divisors of k (A000005).

Conjecture: all terms but the first two are even numbers. - _Harvey P. Dale_, Jul 21 2025 Proof of conjecture: if m is odd and > 10 then either m-1 or m+1 is divisible by 4 and > 8 as well. Let t be the number from {m-1, m+1} divisible by 4. Then t is a power of 2 that is > 8 and so has more than two divisors or it has an odd prime divisor such that it has more than 4 divisors. Both exclude the odd m > 8 from the sequence. - _David A. Corneth_, Aug 05 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, May 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006188 | OEIS-A190035

Number of nondecreasing arrangements of n+2 numbers in 0..3 with the last equal to 3 and each after the second equal to the sum of one or two of the preceding three.

Conjectures from _Colin Barker_, May 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006189 | OEIS-A190036

Number of nondecreasing arrangements of n+2 numbers in 0..4 with the last equal to 4 and each after the second equal to the sum of one or two of the preceding three.

Conjectures from _Colin Barker_, May 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006190 | OEIS-A190038

Number of nondecreasing arrangements of n+2 numbers in 0..6 with the last equal to 6 and each after the second equal to the sum of one or two of the preceding three.

Conjectures from _Colin Barker_, May 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006191 | OEIS-A190039

Number of nondecreasing arrangements of n+2 numbers in 0..7 with the last equal to 7 and each after the second equal to the sum of one or two of the preceding three.

Conjectures from _Colin Barker_, May 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006192 | OEIS-A190040

Number of nondecreasing arrangements of n+2 numbers in 0..8 with the last equal to 8 and each after the second equal to the sum of one or two of the preceding three.

Conjectures from _Colin Barker_, May 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006193 | OEIS-A190091

Number of rhombuses on a (n+1) X 3 grid.

Conjectures from _Colin Barker_, May 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006194 | OEIS-A190092

Number of rhombuses on a (n+1) X 4 grid.

Conjectures from _Colin Barker_, May 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006195 | OEIS-A190093

Number of rhombuses on an (n+1) X 5 grid.

Conjectures from _Colin Barker_, May 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006196 | OEIS-A190154

Diagonal sums of the triangle A190152.

Conjecture: G.f. (-1+x+x^3-x^4+2*x^2) / ((x^3-3*x^2+4*x-1)*(x^3+3*x^2+2*x+1)). - _R. J. Mathar_, Mar 15 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, May 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006197 | OEIS-A190160

Number of peakless Motzkin paths of length n containing no subwords of type uh^ju or dh^jd (j>=1), where u=(1,1), h=(1,0), and d=(1,-1) (can be easily expressed using RNA secondary structure terminology).

Conjecture D-finite with recurrence (n+2)*a(n) +(-5*n-6)*a(n-1) +2*(4*n+3)*a(n-2) +(-2*n-9)*a(n-3) +(-8*n+25)*a(n-4) +6*(n-2)*a(n-5) +9*(n-7)*a(n-6) +3*(-5*n+32)*a(n-7) +7*(n-7)*a(n-8) +(-n+8)*a(n-9)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, May 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006198 | OEIS-A190163

Number of subwords of type dh^ju (j>=1), where u=(1,1), h=(1,0), and d=(1,-1), in all peakless Motzkin paths of length n (can be easily expressed using RNA secondary structure terminology).

Conjecture D-finite with recurrence -4*(n+1)*(n-7)*a(n) +(13*n^2-85*n+28)*a(n-1) +(-7*n^2+52*n-41)*a(n-2) +(5*n^2-41*n+67)*a(n-3) +(-13*n^2+103*n-197)*a(n-4) +(7*n-29)*(n-5)*a(n-5) -(n-5)*(n-6)*a(n-6)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, May 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006199 | OEIS-A190171

Number of peakless Motzkin paths of length n having no UHD's starting at level 0; here U=(1,1), H=(1,0), and D=(1,-1).

Conjecture D-finite with recurrence (n+2)*a(n) +(-n+1)*a(n-1) +2*(-2*n-1)*a(n-2) +9*a(n-3) +(-n+1)*a(n-4) -9*a(n-5) +2*(-2*n+5)*a(n-6) +(-n+1)*a(n-7) +(n-4)*a(n-8)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, May 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006200 | OEIS-A190213

Integers m such that m divides (2^m-2)^2 and (m-2)^((k-1)*(1+k*(m-1))) == 1 (mod k), where k = 2^m - 1.

Conjecture: All odd entries are also Mersenne exponents (A000043): primes n such that 2^n-1 is prime. Any exceptions to the conjecture are larger than 10^5. - _Charles R Greathouse IV_, Oct 03 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alzhekeyev Ascar M_, May 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006201 | OEIS-A190273

Numbers n such that n' = m+1, with n and m semiprimes and gcd(m,n)>1, where n' is the arithmetic derivative of n.

The sequence is related to the Rassias Conjecture ("for any prime p there are two primes p1 and p2 such that p*p1=p1+p2+1, p>2, p2>p1", see A190272-A190275), because n = p1*p2, m=p1*p -> p1*p = p1+p2-1. The sequence includes the cases with p=p1 (or p2). Generalization can be achieved by removing semiprimarity condition or accepting gcd(n,m)=1. The differential equation in its general form n'=m+1 includes Giuga Numbers, i.e., n'=b*n+1, or n'=n+1 (A007850).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giorgio Balzarotti_, May 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006202 | OEIS-A190274

Numbers n such that n' = p^2-1, with n = semiprime = p*q, n' is the arithmetic derivative of n. Also: semiprimes of the form p*(p^2-p-1).

The sequence shows similarity with the Rassias Conjecture ("for any prime p there are two primes p1 and p2 such that p*p1=p1+p2+1, p>2, p2>p1") with p1=p we have p*p=p+p2-1 (see A190272). Generalization can be achieved by removing semiprimarity condition and accepting p^e, e>=2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giorgio Balzarotti_, May 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006203 | OEIS-A190302

Smallest number h such that the decimal expansion of n*h starts with 1.

Conjecture: a(n) < 6 for all n (verified to n = 10022141). - _Felix Fröhlich_, Jul 28 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, May 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006204 | OEIS-A190314

The number of cycles in the digraph representation of all endofunctions on {1,2,...,n}.

Conjectures from _Velin Yanev_, Apr 14 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Critzer_, May 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006205 | OEIS-A190363

a(n) = n + [n*r/t] + [n*s/t]; r=1, s=sqrt(5/4), t=sqrt(4/5).

Conjecture: linear recurrence with constant coefficients 0, 0, 0, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 0, 0, 0, -1. - _Harvey P. Dale_, Jan 28 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006206 | OEIS-A190413

primepi(R_{n*m}) <= n*primepi(R_m) for m >= a(n), where R_k is the k-th Ramanujan prime (A104272).

This is Conjecture 1 in the paper by Sondow, Nicholson, and Noe. The conjecture has been verified for n <= 20 and Ramanujan primes less than 10^9. The conjecture has been proven for n > 10^300 by Shichun Yang and Alain Togbé. - _Jonathan Sondow_, Jan 21 2016 The conjecture has been proven for n > 38 and m > 9 by Christian Axler. Complete exception list can be found in remark of paper. - _John W. Nicholson_, Aug 04 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, May 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006207 | OEIS-A190414

primepi(R_m) <= i*primepi(R_j) for any factorization m=i*j if j >= a(i), where R_k is the k-th Ramanujan prime (A104272).

This is another interpretation of Conjecture 1 in the paper by Sondow, Nicholson, and Noe. The conjecture has been verified for i <= 20 and Ramanujan primes less than 10^9. The conjecture has been proven for i > 38 and j > 9 by Christian Axler. Complete exception list can be found in remark of paper. - _John W. Nicholson_, Aug 04 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Nicholson_, May 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006208 | OEIS-A190640

Numbers whose base-3 expansion ends in 2 and does not contain any 1's.

Conjecture: a(n) = A055246(n) + 1. - _Michel Marcus_, Aug 24 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 15 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006209 | OEIS-A190661

a(n) is the least number m such that there are at least n primes in the range (T(k-1), T(k)] for all k >= m, where T(k) is the k-th triangular number.

All values and even whether the sequence is well defined are conjectural. a(n) is the conjectured index of the last occurrence of n in A066888. It is conjectured that for every n >= 0, a(n) > n. With R_n the n-th Ramanujan prime (A104272), it is conjectured that for every n >= 0, (1/2) R_n <= a(n) < (20/13) R_n. These bounds have been verified for all n up to 8000. For most n <= 8000, we have a(n) > R_n, with exceptions listed in A190881.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Nicholson_, May 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006210 | OEIS-A190724

Row sums of Riordan matrix A118384.

Conjecture: n*(2*n+3)*a(n) +2*(-12*n^2-15*n+22)*a(n-1) +(74*n^2+73*n-274)*a(n-2) -6*(2*n+5)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, May 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006211 | OEIS-A190725

Diagonal sums of Riordan matrix A118384.

conjecture: n*(2*n-3)*a(n) +(-6*n^2+15*n-8)*a(n-1) +2*(-16*n^2+32*n-11)*a(n-2) +(-6*n^2+9*n-2)*a(n-3) +(2*n-1)*(n-2)*a(n-4) =0. - _R. J. Mathar_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, May 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006212 | OEIS-A190736

Diagonal sums of the Riordan matrix A121576.

Conjecture: n*(11*n-35)*a(n) + 3*(-33*n^2+149*n-136)*a(n-1) +2*(77*n^2-377*n+396)*a(n-2) +(-209*n^2+1061*n-1200)*a(n-3) +12*(-11*n+30)*a(n-4) +4*(11*n-24)*(n-4)*a(n-5)=0. - _R. J. Mathar_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, May 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006213 | OEIS-A190786

Numbers m such that sigma(2*m-1) = 3*m, where sigma(k) is the sum of the positive divisors of k.

All even perfect numbers are of the form z(2*z-1) with z = 2^(p-1), p prime and 2*z-1 = 2^p-1 prime. It is unknown if there are any odd perfect numbers of this same form. The equation defining the sequence appears while working a special case of the conjecture. It is conjectured that all terms of this sequence are even numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luis H. Gallardo_, May 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006214 | OEIS-A190788

Expansion of ((x-1)*sqrt(1-4*x^2))/((x-1)*sqrt(1-4*x^2)+x).

Conjecture D-finite with recurrence: (-n+1)*a(n) +(3*n-5)*a(n-1) +2*(3*n-7)*a(n-2) +4*(-6*n+19)*a(n-3) +4*(n-3)*a(n-4) +4*(11*n-53)*a(n-5) +16*(-3*n+16)*a(n-6) +16*(n-6)*a(n-7)=0. - _R. J. Mathar_, Nov 28 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, May 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006215 | OEIS-A190803

Increasing sequence generated by these rules: a(1)=1, and if x is in a then 2x-1 and 3x-1 are in a.

Examples, where [A.....] indicates a conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006216 | OEIS-A190898

Least odd prime p>n^2 with (n/p) = 1, where (/) is the Legendre symbol.

Conjecture: a(n)<(n+1)^2 for all n>0. (See also A185150.) This conjecture implies that a(1),a(2),a(3),... are pairwise distinct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006217 | OEIS-A190918

Number of permutations of n copies of 1..4 introduced in order 1..4 with no element equal to another within a distance of 1.

Conjecture: n^3 *(n-1) *(2*n+1) *(5*n-7) *a(n) -2*(n-1) *(415*n^5 -1026*n^4 +727*n^3 -90*n^2 -98*n +36)*a(n-1) +(1630*n^4 -6387*n^3 +7388*n^2 +111*n -3510) *(n-2)^2 *a(n-2) -162*(3+5*n) *(n-2)^2 *(n-3)^3 *a(n-3)=0. - _R. J. Mathar_, Nov 01 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006218 | OEIS-A190940

Number of divisors of LCM(1,2,...,n)/n.

Also, the number of distinct values taken by lcm(a,a+b,a+b+c,...,n), where positive integers a,b,c,... run over the compositions a+b+c+...=n. - Conjectured by _Ridouane Oudra_, Aug 24 2019; proved by _Max Alekseyev_, Nov 22 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, May 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006219 | OEIS-A190952

Largest integer k for which exp(k) < k^n, n>=3.

Conjecture: There are floor((n+1)/log(n+1))-2 terms less than or equal to n. - _Benedict W. J. Irwin_, Jun 15 2016 Conjecture: G.f.: Sum_{ j>=1 } (Sum_{ k>=1 } x^(j+floor((k+1)/log(k+1)))) + x^j. - _Benedict W. J. Irwin_, Jun 15 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shel Kaphan_, May 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006220 | OEIS-A190969

a(n) = 5*a(n-1) - 8*a(n-2), with a(0)=0, a(1)=1.

Let S(p):=Sum_{k=0..p-1} a(4k)*binomial(2k,k)^3/(-4096)^k. Zhi-Wei Sun conjectured that S(p) == 0 (mod p^2) for every odd prime p, and also S(p) == 0 (mod p^3) for any odd prime p == 1,2,4 (mod 7). - _Zhi-Wei Sun_, Mar 13 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Joseph Stephan Orlovsky_, May 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006221 | OEIS-A191017

Primes with Kronecker symbol (p|14) = 1.

Conjecture: primes congruent to {1, 3, 5, 9, 13, 15, 19, 23, 25, 27, 39, 45} mod 56. - _Vincenzo Librandi_, Jun 22 2016 These are primes p such that either one of (a) p == 1, 2, 4 (mod 7), p == 1, 7 (mod 8) or (b) p == 3, 5, 6 (mod 7), p == 3, 5 (mod 8) holds. So the conjecture above is correct. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, May 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006222 | OEIS-A191107

Increasing sequence generated by these rules: a(1)=1, and if x is in a then 3x-2 and 3x+1 are in a.

Conjecture: a(n) = 3*A003278(n) - 2 = (A055246(n) + 1)/2. - _L. Edson Jeffery_, Nov 25 2015 Conjecture: a(n) = A190640(n)/2. - _Michel Marcus_, Aug 24 2016 Conjecture: a(n) = A003278(2n-1). - _Arie Bos_, Aug 07 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006223 | OEIS-A191150

Hypersigma(n), definition 1: sum of the divisors of n plus the recursive sum of the divisors of the restricted divisors.

For n > 1, a(n) = A191161(n) - A074206(n). [Conjectured by Sequence Machine] - _Antti Karttunen_, Nov 22 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alonso del Arte_, May 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006224 | OEIS-A191161

Hypersigma(n), definition 2: sum of the divisors of n plus the recursive sum of the divisors of the proper divisors.

Following formulas were conjectured by Sequence Machine:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alonso del Arte_, May 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006225 | OEIS-A191257

a(n) = A067368(n)/2.

With the previous definition, a(n) = A067368(n)/2 was stated as a conjecture. _Jianing Song_, Sep 21 2018 showed that this is true, and gave us the simpler definition that we have now used. The conjecture is correct because {a(n)} lists the numbers of the form 2^(3t)*s, and {A067368(n)} lists the numbers of the form 2^(3t+1)*s, where s is an odd number. Note also that a(n) = A213258(n)/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006226 | OEIS-A191269

Fixed point of the morphism 0 -> 001, 1 -> 02, 2 -> 01.

Proof of Kimberling's conjecture on the positions of 0 in this sequence: consider the letter to letter projection pi given by pi(0) = 0, pi(1) = 1, pi(2) = 1. Then pi sigma = tau pi, where tau is the morphism on {0,1} given by tau(0) = 001, tau(1) = 01. It follows that pi(a) = x, where x = A188432 is the fixed point of tau. Note that the positions of zero in a = A191269 are equal to the positions of zero in x. Since x is the infinite Fibonacci word with a zero in front, it follows that these positions are given by A026351. - _Michel Dekking_, Aug 24 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006227 | OEIS-A191303

An infinite sequence of 4-digit half-palindromes.

Conjecture. For every k>=2, there exists a polynomial of degree k taking only k-digit half-palindrome values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006228 | OEIS-A191307

Sum of the heights of the first peaks in all dispersed Dyck paths of length n (i.e., in Motzkin paths of length n with no (1,0)-steps at positive heights).

Conjecture: -(n+3)*(n-2)*a(n) +(n^2+3*n-6)*a(n-1) +2*n*(2*n-3)*a(n-2) - 4*n*(n-1)*a(n-3)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, May 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006229 | OEIS-A191309

Number of peaks at height >= 2 in all dispersed Dyck paths of length n (i.e., Motzkin paths of length n with no (1,0) steps at positive heights).

Conjecture: n*(n-4)*a(n) +(n^2-10*n+15)*a(n-1) +2*(-5*n^2+28*n-27)*a(n-2) -4*(n-3)*(n-8) *a(n-3) +24*(n-3)*(n-4)*a(n-4)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, May 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006230 | OEIS-A191313

Sum of the abscissae of the first returns to the horizontal axis (assumed to be 0 if there are no such returns) in all dispersed Dyck paths of length n (i.e., Motzkin paths of length n with no (1,0) steps at positive heights).

Conjecture: n*(3*n-13)*a(n) +2*(-6*n^2+29*n-18)*a(n-1) +(3*n^2-13*n+24)*a(n-2) +2*(21*n^2-124*n+150)*a(n-3) +4*(-15*n^2+92*n-132) *a(n-4) +8*(n-3)*(3*n-10) *a(n-5)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, May 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006231 | OEIS-A191319

Sum of pyramid weights of all dispersed Dyck paths of length n (i.e., of all Motzkin paths of length n with no (1,0) steps at positive heights).

Conjecture: (-n+2)*a(n) +2*a(n-1) +5*(n-2)*a(n-2) -2*a(n-3) +4*(-n+2)*a(n-4)=0. - _R. J. Mathar_, Dec 07 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jun 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006232 | OEIS-A191355

Indices of terms in A069748 with two decimal digits 1 and all others 0.

Conjectural formula: a(n) = 5 + Sum_{i=1..n-1} A105637(i). Equivalent conjecture: a(n) = -(-1)^n/8 + A049347(n)/3 + 115/24 + n/6 + n^2/4. - _R. J. Mathar_, Jul 11 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 31 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006233 | OEIS-A191363

Numbers m whose deficiency is 2: sigma(m) - 2*m = -2.

Conjecture: a(1)=3 is the only odd term of the sequence. Conjecture: All terms of the sequence are of the above form derived from Fermat primes. Conjectures: all terms are second hexagonal numbers (A014105). There are no terms with middle divisors. - _Omar E. Pol_, Oct 31 2018 Conjecture: Numbers m whose two odd divisors are 1 and A033677(m) = 2*A033676(m) + 1. - _Omar E. Pol_, Feb 07 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luis H. Gallardo_, May 31 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006234 | OEIS-A191374

Number of ways (up to rotations and reflections) of arranging numbers 1 through 2n around a circle such that the sum of each pair of adjacent numbers is composite.

One of the obvious analogs of sequence A051252, which has the sums being prime. Presumably it is an open problem as to whether a(n) > 0 always for this problem as well.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bennett Gardiner_, Jun 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006235 | OEIS-A191398

Number of dispersed Dyck paths of length n (i.e., Motzkin paths of length n with no (1,0)-steps at positive heights) having no DHU's (here U=(1,1), H=(1,0), and D=(1,-1)).

Conjecture: -(n+2)*(n-3)*a(n) +(3*n^2-3*n-14)*a(n-1) +2*(n^2-7*n+8)*a(n-2) +4*(-3*n^2+12*n-10)*a(n-3) +(7*n^2-31*n+38)*a(n-4) +4*a(n-5) +4*(n-2)^2*a(n-6)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jun 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006236 | OEIS-A191432

Dispersion of ([n*x+1/x]), where x=sqrt(2) and [ ]=floor, by antidiagonals.

Conjecture: It appears this sequence is related to the even numbers with odd abundance A088827. Looking at the table format if the columns represent the powers of 2 (starting at 2^1) and the rows represent the squares of odd numbers, then taking the product of a term's row and column gives the n-th term in A088827. Example: A088827(67) = (7^2) * (2^6) = 3136. - _John Tyler Rascoe_, Jul 12 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006237 | OEIS-A191450

Dispersion of (3*n-1), read by antidiagonals.

Conjecture: A(n,k) = (3 + (2*A032766(n) - 1)*A000244(k))/6. - _L. Edson Jeffery_, with slight changes by _Antti Karttunen_, Jan 21 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006238 | OEIS-A191526

Number left factors of Dyck paths of length n and having no hills; a hill is a (1,1)-step starting at level 0 and followed by a (1,-1)-step.

Conjecture: -2*(n+1)*(3*n-10)*a(n) +12*(n-5)*a(n-1) +(21*n^2-97*n+122)*a(n-2) +6*(n-5)*a(n-3) +4*(n-2)*(3*n-7)*a(n-4)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jun 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006239 | OEIS-A191531

Sum of lengths of initial and final horizontal segments over all dispersed Dyck paths of semilength n (i.e., over all Motzkin paths of length n with no (1,0)-steps at positive heights).

Conjecture: +n*(n-3)*a(n) -2*(n^2-2*n-2)*a(n-1) -(3*n^2-21*n+32)*a(n-2) +2*(n-2)*(4*n-13)*a(n-3) -4*(n-2)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jun 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006240 | OEIS-A191585

Central coefficients of the Riordan matrix (1/(1-3*x^2),x/(1-x)) (A191582).

Conjecture: 2*n*(n+3)*a(n) +2*(-7*n^2-19*n+24)*a(n-1) +3*(5*n^2+11*n-48)*a(n-2) +18*(n+4)*(2*n-3)*a(n-3)=0. - _R. J. Mathar_, Jun 14 2016 Conjecture: +4*n*a(n) +2*(-23*n+22)*a(n-1) +156*(n-2)*a(n-2) +9*(-7*n+38)*a(n-3) +162*(-2*n+5)*a(n-4)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, Jun 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006241 | OEIS-A191625

Number of n-step two-sided prudent self-avoiding walks ending at the northeast corner of their box.

Conjecture: +(n+1)*a(n) -4*n*a(n-1) +(n-5)*a(n-2) +2*(4*n-3)*a(n-3) +3*(-n+3)*a(n-4) +2*(n-8)*a(n-5) +(-n+13)*a(n-6) +8*(-n+7)*a(n-7) +2*(n-7)*a(n-8) +2*(n-9)*a(n-9)=0. - _R. J. Mathar_, Sep 16 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jun 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006242 | OEIS-A191668

Dispersion of A016825 (4k+2, k>0), by antidiagonals.

Conjecture: a(n,k) = (8 + (3*floor((4*n + 1)/3) - 2)*4^k)/12 = (8 + (3*A042965(n+1) - 2)*A000302(k))/12. - _L. Edson Jeffery_, Feb 14 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006243 | OEIS-A191677

Numbers n such that 1^(n-1)+2^(n-1)+...+n^(n-1) == 0 (mod n).

Fermat's little theorem shows that this sequence contains no primes. Related to Giuga's conjecture that the sum is -1 iff n is prime. - _Charles R Greathouse IV_, Jun 10 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Jun 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006244 | OEIS-A191782

Sum of the lengths of the first ascents in all n-length left factors of Dyck paths.

Conjecture: -(n+3)*(6*n-17)*a(n) +2*(6*n^2-2*n-57)*a(n-1) +3*(6*n^2-17*n+27)*a(n-2) -2*(2*n-3)*(12*n-25)*a(n-3) +4*(6*n-11)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jun 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006245 | OEIS-A191783

Numbers k such that the topological k-sphere has a unique differentiable structure up to diffeomorphism.

Whether 4 is a term is an open question. - _Andrey Zabolotskiy_, Feb 02 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006246 | OEIS-A191786

Number of length n left factors of Dyck paths having no triple-rises (triple-rise = three consecutive (1,1)-steps).

Conjecture: -(n+3)*(13*n-70)*a(n) +(-13*n^2+19*n-102)*a(n-1) +(65*n^2-221*n-516) *a(n-2) +(65*n^2-197*n+288)*a(n-3) -(n+6)*(13*n-97) *a(n-4) +3*(-13*n^2+35*n-70) *a(n-5) +(-169*n^2+1201*n-2208) *a(n-6) -9*(13*n-40)*(n-5) *a(n-7) -6*(13*n-25)*(n-6) *a(n-8)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jun 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006247 | OEIS-A191790

Number of base pyramids in all length n left factors of Dyck paths.

Conjecture: (n+1)*a(n) +(-n-1)*a(n-1) +5*(-n+1)*a(n-2) +(5*n-11)*a(n-3) +2*(2*n-3)*a(n-4) +4*(-n+3)*a(n-5)=0. - _R. J. Mathar_, Jun 14 2016 Conjecture: -(n+1)*(n-2)*a(n)

+2*(n-1)*a(n-1) +(5*n-3)*(n-2)*a(n-2) +2*(-n+1)*a(n-3) -4*(n-1)*(n-2)*a(n-4)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jun 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006248 | OEIS-A191792

Number of length n left factors of Dyck paths having no UDUD's; here U=(1,1) and D=(1,-1).

Conjecture D-finite with recurrence (n+1)*a(n) -2*a(n-1) +2*(-n+1)*a(n-2) +4*(-1)*a(n-3) +5*(-n+3)*a(n-4) +4*a(n-5) +2*(-n+5)*a(n-6) +2*a(n-7) +(n-7)*a(n-8)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jun 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006249 | OEIS-A191796

Number of DUU's in all length n left factors of Dyck paths; here U=(1,1) and D=(1,-1).

Conjecture: -(n+1)*(2527*n^2+15963*n-146560)*a(n) +(-2527*n^3+68000*n^2-231053*n-293120)*a(n-1) +2*(12635*n^3+906*n^2-429395*n+746484)*a(n-2) +4*(2527*n^3-70527*n^2+316742*n-316524)*a(n-3) -24*(n-5)*(2527*n^2-232*n-28664)*a(n-4)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jun 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006250 | OEIS-A191821

a(n) = n*(2^n - n + 1) + 2^(n-1)*(n^2 - 3*n + 2).

Conjecture: generating function = -((2 (-1+6 x-19 x^2+31 x^3-22 x^4+4 x^5))/(1-3 x+2 x^2)^3) - _Harvey P. Dale_, May 10 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Adeniji, Adenike_, Jun 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006251 | OEIS-A191837

Least even number m which can be written as sum of 2n primes p(1) < ... < p(2n) < m/2 such that m-p(i) is also prime for i=1,...,2n.

Conjecture: For all a(n), a(n)-1 can be found in A014092 (numbers not the sum of two primes), and a(n)+1 can be found in A007921. (numbers not the difference of two primes). - _J. Stauduhar_, Aug 28 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_, Jun 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006252 | OEIS-A191873

A problem of Zarankiewicz: maximal number of 1's in an n X n matrix of 0's and 1's with 0's on the main diagonal and no "rectangle" with 1's at the four corners.

Conjecture: for n >= 3, a(n) = A072567(n) = A001197(n) - 1 (per above comment). - _Max Alekseyev_, Jan 29 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_ and _N. J. A. Sloane_, Jun 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006253 | OEIS-A191898

Symmetric square array read by antidiagonals: T(n,1)=1, T(1,k)=1, T(n,k) = -Sum_{i=1..k-1} T(n-i,k) for n >= k, -Sum_{i=1..n-1} T(k-i,n) for n < k.

Rows equal columns and are periodic. No zero elements are found (conjecture). The recurrence is related to the recurrence for the Mahonian numbers. The main diagonal is the Dirichlet inverse of the Euler totient function A023900 (conjecture). [The 2nd and 3rd formulas state that the conjecture is correct. _R. J. Mathar_, Sep 16 2017] The sums from n=1 to infinity of T(n,k)/n converge to the Mangoldt function for column k (conjecture). If gcd(n,k)=1 then T(n,k)=1 and if T(n,k)=1 then gcd(n,k)=1 (conjecture). (conjecture) The signs of the eigenvalues of this matrix are a rearrangement of the Mobius function A008683 (conjecture). The first few eigenvalues are:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Jun 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006254 | OEIS-A191904

Square array read by antidiagonals up: T(n,k) = 1-k if k divides n, else 1.

Conjecture: Sum_{n>=1} T(n,k)/n = log(k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Jun 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006255 | OEIS-A191928

Array read by antidiagonals: T(m,n) = floor(m/2)*floor((m-1)/2)*floor(n/2)*floor((n-1)/2).

T(m,n) is conjectured to be the crossing number of the complete bipartite graph K_{m,n}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006256 | OEIS-A191993

a(n) = 3^(n-1) + C(2*n, n)/2.

Conjecture: n*(n-3)*a(n) - (7*n^2 -23*n +12)*a(n-1) +6*(2*n-3)*(n-2)*a(n-2)=0. - _R. J. Mathar_, Oct 18 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mircea Merca_, Jun 21 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006257 | OEIS-A192004

Alternating row sums of array A187360: minimal polynomial of 2*cos(Pi/n) evaluated for x=-1.

The preceding conjecture can be checked by using for even n Theorem 1A, eq. (41), and for odd n Theorem 2A, eq. (50) of the W. Lang arXiv link given in A187360 putting x = -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jul 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006258 | OEIS-A192023

The Wiener index of the comb-shaped graph |_|_|...|_| with 2n (n>=1) nodes. The Wiener index of a connected graph is the sum of the distances between all unordered pairs of vertices in the graph.

Conjecture: for n>2, A192023(n-2) is the number of 2 X 2 matrices with all terms in {1,2,...,n} and determinant 2n. - _Clark Kimberling_, Mar 31 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jun 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006259 | OEIS-A192031

Rectangular array read by rows: T(n,k) is the number of unordered pairs of nodes at distance k in the helm graph G(n) obtained from a wheel graph with n spokes by adjoining a pendant edge at each node of the cycle (n>=3, k>=1). The entries in row n are the coefficients of the corresponding Wiener polynomial.

Conjectures from _Colin Barker_, Mar 28 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jun 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006260 | OEIS-A192056

a(n) = |{0<k<n/2: k^2+(n-k)^2-3(n-1 mod 2) is prime and JacobiSymbol[k,n+3(n-1 mod 2)]=1}|.

Conjecture: a(n)>0 for all n>2. Moreover, if n>2 is not among 12, 18, 105, 522, then there is 0<k<n/2 such that This conjecture has been verified for n up to 2*10^8. It is stronger than Ming-Zhi Zhang's conjecture that any odd integer n>1 can be written as x+y (x,y>0) with x^2+y^2 prime (see A036468).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006261 | OEIS-A192062

Square Array T(ij) read by antidiagonals (from NE to SW) with columns 2j being the denominators of continued fraction convergents to square root of (j^2 + 2j).

Conjecture re relation of A192062 to the sequence of primes: T(2*n,j) = A(n,j)*T(n,j) where A(n,j) is from the square array A191971. There, A(3*n,j) = A(n,j)*B(n,j) where B(n,j) are integers. It appears further that B(5*n,j)=B(n,j)*C(n,j); C(7*n,j)= C(n,j)*D(n,j); D(11*n,j) = D(n,j)*E(n,j); E(13*n,j) = E(n,j)*F(n,j) and F(17*n,j) = F(n,j)*G(n,j) where C(n,j), D(n,j) etc. are all integers. My conjecture is that this property continues indefinitely and follows the sequence of primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kenneth J Ramsey_, Jun 21 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006262 | OEIS-A192069

(A192249)/5.

Conjecture: G.f.: x^2*(1-2*x+4*x^2-3*x^3+x^4) / ((x-1)*(x^2+x-1)^5). - _R. J. Mathar_, May 04 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006263 | OEIS-A192101

Least number requiring n terms to express it as a sum of signed terms of the form 2^k-1.

Conjecture: a(n) = (4^n)/3 + O(1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank Ruskey_, Jul 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006264 | OEIS-A192189

Polynomial-like numbers: numbers whose sequence of number-derivatives (A003415) is monotonically decreasing.

Conjecture. For every sufficiently large greater q of twin primes, the sequence contains infinite increasing sequence {s_n} of semiprimes beginning with 2*(q-2), such that (s_n)'=s_(n-1). This conjecture is true, if 1) there exist infinitely many twin primes; 2) there exists n_0, such that for every prime p>n_0, number 2*p is sum of two primes r,t, for which r*t-2 is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jun 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006265 | OEIS-A192227

Number of primes in the range (n - 2*sqrt(sqrt(n)), n].

Conjecturally, a(n) ~ 2*n^(1/4)/log n. - _Charles R Greathouse IV_, Jul 01 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jun 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006266 | OEIS-A192229

Indices of records in A087669.

Conjecture: the terms starting with 11 are prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Jun 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006267 | OEIS-A192238

Constant term in the reduction of the polynomial x(x+1)(x+2)...(x+n-1) by x^2 -> x+1.

Conjecture: a(n) +2*(-n+1)*a(n-1) +(n^2-3*n+1)*a(n-2)=0. - _R. J. Mathar_, May 04 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006268 | OEIS-A192240

Constant term in the reduction of the polynomial (x+3)^n by x^2 -> x+1.

a(n) = Sum_{i=0..n} (-1)^i*Fibonacci(i+1)*binomial(n,i)*4^(n-i) (conjecture). - _Rigoberto Florez_, Mar 25 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006269 | OEIS-A192245

1-sequence of reduction of triangular number sequence by x^2 -> x+1.

The above conjectures are true. See link for a proof.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006270 | OEIS-A192248

0-sequence of reduction of binomial coefficient sequence B(n,4)=A000332 by x^2 -> x+1.

Conjecture: G.f.: -x*(-1+5*x-20*x^2+30*x^3-25*x^4+8*x^5) / ((x-1)*(x^2+x-1)^5). - _R. J. Mathar_, May 04 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006271 | OEIS-A192249

1-sequence of reduction of binomial coefficient sequence B(n,4)=A000332 by x^2 -> x+1.

Conjecture: G.f.: 5*x^2*(1-2*x+4*x^2-3*x^3+x^4) / ((x-1)*(x^2+x-1)^5). - _R. J. Mathar_, May 04 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006272 | OEIS-A192250

0-sequence of reduction of central binomial coefficient sequence by x^2 -> x+1.

Conjecture: (n-1)*(n-2)*a(n) -(5*n-7)*(n-2)*a(n-1) -2*(2*n-3)*(3*n-8)*a(n-2) +4*(2*n-3)*(2*n-5)*a(n-3)=0. - _R. J. Mathar_, May 04 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006273 | OEIS-A192251

1-sequence of reduction of central binomial coefficient sequence by x^2 -> x+1.

Conjecture: (n-1)*(n-2)*a(n) -(5*n-7)*(n-2)*a(n-1) -2*(2*n-3)*(3*n-8)*a(n-2) +4*(2*n-3)*(2*n-5)*a(n-3)=0. - _R. J. Mathar_, May 04 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006274 | OEIS-A192252

0-sequence of reduction of (n!) by x^2 -> x+1.

Conjecture: a(n) +(-n-1)*a(n-1) -n*(n-2)*a(n-2) +n*(n-1)*a(n-3)=0. - _R. J. Mathar_, May 04 2014
Conjecture: a(n) = Sum_{k=0..n} A052554(k). - _Sean A. Irvine_, Jul 14 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006275 | OEIS-A192253

1-sequence of reduction of (n!) by x^2 -> x+1.

Conjecture: a(n) -n*a(n-1) -(n-1)*(n-3)*a(n-2) +(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, May 04 2014
Conjecture: partial sums of A005443. - _Sean A. Irvine_, Jul 12 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006276 | OEIS-A192255

1-sequence of reduction of (n^2) by x^2 -> x+1.

a(n) = Sum_{i=1..n+1} i^2*Fibonacci(i-1) (conjectured). - _Greg Dresden_, Oct 10 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006277 | OEIS-A192338

Constant term of the reduction of n-th polynomial at A157751 by x^2->x+2.

Conjecture: a(n) = 4*a(n-1)+a(n-2)-10*a(n-3)-4*a(n-4). G.f.: x*(2*x^3-3*x^2-2*x+1) / ((x^2+2*x-1)*(4*x^2+2*x-1)). [_Colin Barker_, Nov 22 2012]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006278 | OEIS-A192344

Constant term of the reduction of n-th polynomial at A161516 by x^2->x+1.

Conjecture a(n) = 2*a(n-1)+10*a(n-2)-6*a(n-3)-9*a(n-4). G.f.: -(5*x^2+2*x-1) / (9*x^4+6*x^3-10*x^2-2*x+1). [_Colin Barker_, Nov 22 2012]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006279 | OEIS-A192346

Constant term of the reduction (by x^2->x+1) of polynomial p(n,x) identified in Comments.

Conjecture: a(n) = 2*a(n-1)+6*a(n-2)-2*a(n-3)-a(n-4). G.f.: -x*(x+1)*(3*x-1) / (x^4+2*x^3-6*x^2-2*x+1). [_Colin Barker_, Jan 17 2013]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006280 | OEIS-A192347

Coefficient of x in the reduction (by x^2->x+1) of polynomial p(n,x) identified in Comments.

Conjecture: a(n) = 2*a(n-1)+6*a(n-2)-2*a(n-3)-a(n-4). G.f.: x^2*(x^2+1) / (x^4+2*x^3-6*x^2-2*x+1). [_Colin Barker_, Jan 17 2013]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006281 | OEIS-A192357

Constant term of the reduction of the polynomial p(n,x)=(1/2)((x+3)^n+(x-3)^n) by x^2->x+1.

Conjecture: a(n) = 2*a(n-1)+19*a(n-2)-20*a(n-3)-55*a(n-4). G.f.: x*(x^3-9*x^2-2*x+1)/((5*x^2+5*x+1)*(11*x^2-7*x+1)). [_Colin Barker_, Nov 22 2012]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006282 | OEIS-A192358

Coefficients of x in the reduction of the polynomial p(n,x)=(1/2)((x+3)^n+(x-3)^n) by x^2->x+1.

Conjecture: a(n) = 2*a(n-1)+19*a(n-2)-20*a(n-3)-55*a(n-4). G.f.: x^2*(8*x^2-x+1)/((5*x^2+5*x+1)*(11*x^2-7*x+1)). [_Colin Barker_, Nov 22 2012]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006283 | OEIS-A192373

Constant term in the reduction of the polynomial p(n,x) defined at A162517 and below in Comments.

Conjecture: a(n) = 2*a(n-1)+10*a(n-2)-6*a(n-3)-9*a(n-4). G.f.: -x*(x+1)*(3*x-1) / (9*x^4+6*x^3-10*x^2-2*x+1). - _Colin Barker_, May 09 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006284 | OEIS-A192376

Constant term of the reduction by x^2->x+2 of the polynomial p(n,x) defined below in Comments.

Conjecture: a(n) = 2*a(n-1)+6*a(n-2)+2*a(n-3)-a(n-4). G.f.: x*(x-1)^2 / ((x+1)^2*(x^2-4*x+1)). - _Colin Barker_, May 11 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006285 | OEIS-A192379

Constant term of the reduction by x^2->x+1 of the polynomial p(n,x) defined below in Comments.

Conjecture: a(n) = 2*a(n-1)+6*a(n-2)-2*a(n-3)-a(n-4). G.f.: -x*(x^2+2*x-1) / (x^4+2*x^3-6*x^2-2*x+1). - _Colin Barker_, May 11 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006286 | OEIS-A192382

Coefficient of x in the reduction by x^2 -> x+2 of the polynomial p(n,x) defined below in Comments.

Conjectures from _Colin Barker_, May 12 2014: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006287 | OEIS-A192393

Number of primes in the half-open interval [semiprime(n)-sqrt(semiprime(n)), semiprime(n)).

Conjecturally, a(n) ~ sqrt(n log log n / (log n)^3). - _Charles R Greathouse IV_, Jun 29 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jun 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006288 | OEIS-A192416

The sequence obtained from A005117 (the squarefree numbers) by the transformation "forming" as defined in A191612.

Conjecture: a(n) is also a "totally formed" sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Oct 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006289 | OEIS-A192431

Coefficient of x in the reduction by x^2->x+1 of the polynomial p(n,x) defined below in Comments.

Conjectures from _Colin Barker_, Jun 07 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006290 | OEIS-A192466

Coefficient of x in the reduction by x^2->x+2 of the polynomial p(n,x)=1+x^n+x^(2n).

Conjectures from _Colin Barker_, Feb 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006291 | OEIS-A192473

Coefficient of x in the reduction by x^2->x+1 of the polynomial p(n,x)=1+x^n+x^(2n+2).

Conjectures from _Colin Barker_, Jun 07 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006292 | OEIS-A192480

a(n) = n + A000108(n-1) for n > 1; a(0)=0, a(1)=1.

Conjecture: n*(3*n^2-16*n+19)*a(n) +(-15*n^3+95*n^2-188*n+120)*a(n-1) +2*(2*n-5)*(3*n^2-10*n+6)*a(n-2)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Volkan Yildiz_, Jul 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006293 | OEIS-A192628

Nonvanishing exponents of the reciprocal of the modulo 2 generating function for the sum-of-divisor function.

Density upper bound: 1/16. Conjectured density: 1/32. Contains only 0 and positive integers congruent to 1 and 3 (mod 8) and 7 (mod 16). *7 (mod 16): conjectured density 1/32 with upper bound 1/16.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Riasanovsky_, Dec 31 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006294 | OEIS-A192636

Powerful sums of two powerful numbers.

Browning & Valckenborgh conjecture that a(n) ~ kn^2 with k approximately 0.139485255. See their Conjecture 1 and equation (14). Their Theorems 1 and 2 establish upper and lower asymptotic bounds.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Jul 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006295 | OEIS-A192718

Elements of A192628 which are congruent to 7 (mod 8) (equivalently, 7 (mod 16)).

However, the 7 modulo 8 congruence classes appears to have nonzero density, conjectured 1/32. A current upper bound on its density (thus the entire density of A192628) is 1/16.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Riasanovsky_, Dec 31 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006296 | OEIS-A192763

Symmetric square array read by antidiagonals up.

The main diagonal is the Mobius function times the natural numbers A055615 (conjecture). For k>1 the first row is the Mertens function + 2 = A002321 + 2 (conjecture). There is one recurrence for n=1 and k=1, and another recurrence for n>1 and k>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Jul 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006297 | OEIS-A192820

2-Ramanujan primes: the interval (x/2,x] has at least n Ramanujan primes for x >= a(n) but not for x = a(n) - 1.

It is conjectured that primepi(a(n)) <= 7*n for all n. - _T. D. Noe_, Aug 26 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006298 | OEIS-A192821

3-Ramanujan primes; the interval (x/2,x] has at least n 2-Ramanujan primes for x >= a(n) but not for x = a(n)-1.

It is conjectured that primepi(a(n)) < 10*n for large n. - _T. D. Noe_, Aug 26 2011 The sequence is only conjectural without a proof of an upper bound on a(n) (like the bound A104272(n) < prime(3*n) proved by Laishram and used in computing Ramanujan primes). - _Jonathan Sondow_, Aug 26 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006299 | OEIS-A192822

4-Ramanujan primes; the interval (x/2,x] has at least n 3-Ramanujan primes for x >= a(n) but not for x = a(n)-1.

It is conjectured that primepi(a(n)) < 22*n for large n. - _T. D. Noe_, Aug 26 2011 The sequence is only conjectural without a proof of an upper bound on a(n) (like the bound A104272(n) < prime(3*n) proved by Laishram and used in computing Ramanujan primes). - _Jonathan Sondow_, Aug 27 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006300 | OEIS-A192823

5-Ramanujan primes; the interval (x/2,x] has at least n 4-Ramanujan primes for x >= a(n) but not for x = a(n)-1.

It is conjectured that primepi(a(n)) < 50*n for large n. - _T. D. Noe_, Aug 26 2011 The sequence is only conjectural without a proof of an upper bound on a(n) (like the bound A104272(n) < prime(3*n) proved by Laishram and used in computing Ramanujan primes). - _Jonathan Sondow_, Aug 27 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006301 | OEIS-A192824

Least n-Ramanujan prime.

The sequence is only conjectural without a proof of an upper bound on a(n) (like the bound A104272(n) < prime(3*n) proved by Laishram and used in computing Ramanujan primes). - _Jonathan Sondow_, Aug 27 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006302 | OEIS-A192826

Molecular topological indices of the folded cube graphs.

Conjecture: D-finite with recurrence: +(n-1)*(3171347*n-9561097)*a(n) +2*(-22257061*n^2+113080920*n-124259984)*a(n-1) +8*(25480326*n^2-186826067*n+302663821)*a(n-2) +16*(-2142117*n^2+189489004*n-548366652)*a(n-3) +64*(-46156212*n^2+185594099*n-75953597)*a(n-4) +1024*(11001336*n^2-74317052*n+114909451)*a(n-5) +1024*(-17489828*n^2+145166555*n-279390145)*a(n-6) +20480*(530179*n-1731202)*(n-6)*a(n-7)=0. - _R. J. Mathar_, Feb 21 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Jul 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006303 | OEIS-A192869

Thin primes: odd primes p such that p+1 is a prime (or 1) times a power of two.

Broughan & Qizhi conjecture that a(n) << n (log n)^2, matching the lower bound they proved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Jul 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006304 | OEIS-A192924

Constant term in the reduction by (x^2->x+1) of the polynomial p(n,x) defined below at Comments.

Conjecture: a(n) +2*(-n+1)*a(n-1) +(n^2-3*n-1)*a(n-2) +3*(n-2)*a(n-3) +a(n-4)=0. - _R. J. Mathar_, May 08 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 12 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006305 | OEIS-A192931

Numbers n for which A193262(n)=0.

Conjecture: The sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Aug 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006306 | OEIS-A192934

Smallest square number starting with (at least) n identical digits.

Conjecture: total number of digits of a(n) is always 2*n-1. This conjecture is false: length(a(6)) = 10 != 2*6-1 = 11.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Aug 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006307 | OEIS-A192939

Constant term of the reduction by x^2->x+1 of the polynomial p(n,x)=(x+2)(x+4)...(x+2n).

Conjecture: a(n) +(-4*n+1)*a(n-1) +(4*n^2-6*n+1)*a(n-2)=0. - _R. J. Mathar_, May 08 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 13 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006308 | OEIS-A192940

Coefficient of x in the reduction by x^2->x+1 of the polynomial p(n,x)=(x+2)(x+4)...(x+2n).

Conjecture: a(n) +(-4*n+1)*a(n-1) +(4*n^2-6*n+1)*a(n-2)=0. - _R. J. Mathar_, May 08 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 13 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006309 | OEIS-A192984

In repeated iterations of function m->m/3 if m == 0 (mod 3), m->4m+a if 4m+a == 0 (mod 3) where a = 1 or 2, a(n) is maximum value achieved if starting from n.

If a(n) exists (which is the conjecture of A194428) then a(n) must be a multiple of 9, since if a(n) is not divisible by 3 then the next iteration 4*a(n) + 1 or 4*a(n) + 2 would be greater than a(n), while if a(n) was divisible by 3 only once, a(n) = 3x -> x -> 4x+1 or 4x+2 is greater than 3x.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Aug 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006310 | OEIS-A192989

Expansion of e.g.f.: exp((1+x)^3 - 1).

Conjecture: a(n) -3*a(n-1) +6*(-n+1)*a(n-2) -3*(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, May 12 2014 Remark: the above conjectured recurrence is true and can be easily obtained from the e.g.f. - _Emanuele Munarini_, Aug 31 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 13 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006311 | OEIS-A192993

Numbers that are in more than one way the concatenation of the decimal representation of two nonzero squares.

Conjecture: a(n) = A038670(n) for all n >= 1. - _Zhuorui He_, Jun 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Brockhaus_ and _Zak Seidov_, Jul 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006312 | OEIS-A193020

Number of distinct self-shuffles of the word given by the binary representation of n.

An examination of the terms of the sequence leads to the following conjectures (in each case with the caveat that k must exceed a certain lower bound): a(2^k-5)=3k-6, a(2^k-4)=k*(k-1)/2, a(2^k-3)=2k-2, a(2^k-2)=k, a(2^k-1)=1, a(2^k)=k+1, a(2^k+1)=k^2, a(2^k+2)=2*(k-1)^2, a(2^k+3)=k*(k-1)^2/2. To illustrate, consider a(2^k+1); we get, for k=1, 2, 3, ..., a(3)=1, a(5)=4, a(9)=9, a(17)=16, a(33)=25, a(65)=36, a(129)=49, a(257)=64,..., leading to the conjecture that a(2^k+1)=k^2. The other conjectures were arrived at in the same manner.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jul 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006313 | OEIS-A193101

Minimal number of numbers of the form (m^3+5m)/6 (see A004006) needed to sum to n.

It is conjectured that a(n) <= 5 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 15 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006314 | OEIS-A193138

Number of square satins of order n.

This sequence is conjectured to coincide with the multiplicities of the representation of n >= 3 as primitive sums of two squares. Neither the order of the squares nor the signs of the numbers to be squared are taken into account. a(n) = 0 if no such representation exists. Checked for n = 3,4, ..., 1000 (using the program below). The two squares are in each case nonzero and distinct. If one includes also 0 as a square in the primitive sum of two squares one could take a(0) = 0, a(1) = 1, a(2) = 1. If only nonzero squares are considered, then one could take a(0) = 0, a(1) = 0, a(2) = 1. For the numbers n with a(n) > 0 (in this conjectured interpretation of a(n)) see A008784. - _Wolfdieter Lang_, Apr 17 2013 The stated conjecture is true because it follows immediately from Theorem 3.22, p. 165, of the Niven-Zuckerman-Montgomery reference. There r(n) gives the number of primitive solutions of n = x^2 + y^2 with ordered and signed pairs of integers x,y. Because x and y are distinct if n >= 3 one needs here a(n) = r(n)/2^3. This then coincides with the formula for u(n) given in the Grünbaum-Shephard Theorem 5. - _Wolfdieter Lang_, Apr 18 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006315 | OEIS-A193142

Primes which are the sum of 5 distinct positive squares.

Conjecture: a(n) = prime(n+32) for n > 13. [_Charles R Greathouse IV_, Jul 17 2011]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Joseph Stephan Orlovsky_, Jul 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006316 | OEIS-A193168

Number of simple paths from (1, 1) to (n, 3) on an n X 3 grid with king moves allowed.

Conjectures from _Andrew Howroyd_, Nov 05 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matías Benzo_, Jul 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006317 | OEIS-A193229

A double factorial triangle.

It appears that T(n,k) = (2*n-k)!/(2^(n-k)*(n-k)!) with conjectural e.g.f. 1/(x*(1-2*z) + (1-x)*sqrt(1-2*z)) = 1 + (1+x)*z + (3+3*x+2*x^2)*z^2/2! + ... Cf. A102625. - _Peter Bala_, Jul 09 2012 The conjectured formula above is valid. The production matrix gives T(n+1,0) = T(n+1,1), T(n+1,k)-T(n+1,k+1) = k*T(n,k-1) for 1 <= k <= n, and T(n,n) = n!. The displayed formula satisfies these relations and T(0,0)=1, which proves it by induction. - _Philip Weiss_, Jul 09 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Jul 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006318 | OEIS-A193236

Expansion of g.f.: (Sum_{n>=0} (-3)^n*(2*n+1)*x^(n*(n+1)/2))^(-1/3).

Conjectures: a(5*n+4) == 0 (mod 5) and a(7*n+5) == 0 (mod 7) (both checked up to n = 200). - _Peter Bala_, Feb 26 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006319 | OEIS-A193237

Expansion of g.f.: (Sum_{n>=0} (2*n+1)*3^n*(-x)^(n*(n+1)/2))^(-1/3).

Conjectures: a(5*n+4) == 0 (mod 5) and a(7*n+5) == 0 (mod 7) (both checked up to n = 200). - _Peter Bala_, Feb 26 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006320 | OEIS-A193262

Number of representations of 2*p_n as sum of two primes p,q such that p*q-2 is prime (p_n is the n-th prime).

Sequence arising in connection with conjecture in comment to A192189. Conjecture: There exists n_0, such that, for n>n_0, a(n)>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Aug 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006321 | OEIS-A193282

a(n) = (n!/floor(n/2)!)^2.

Conjecture: a(n) + 8*a(n-1) - 4*(n-2)*(n+2)*a(n-2) + 16*(-2*n^2 + 6*n - 3)*a(n-3) - 64*(n-3)^2*a(n-4) = 0. - _R. J. Mathar_, Oct 03 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Sep 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006322 | OEIS-A193351

Numbers k such that A071324(k) is prime.

It is interesting to note that all numbers except the first are squares, or twice a square, hence the conjecture: This conjecture is verified up to k <= 10^8. - _Shreyansh Jaiswal_, Apr 25 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Dec 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006323 | OEIS-A193358

a(0)=1; a(1)=2 and for n>1: a(n)=a(n-a(n-2))+2.

Conjecture: Every positive integer is in this sequence. Conjecture: Odd numbers are not repeated. Both conjectures are true. Sequence follows the pattern: for n = 0,1,2,..., the sequence consists of 2*n terms of value 2*(n+1), followed by the terms 2*n+1 and 2*(n+1). Thus 0 terms of 2 followed by 1 and 2, then 2 terms of 4 followed by 3 and 4, then 4 terms of 6 followed by 5 and 6, etc. In other words, a(n*(3+n)) = 2n+1 and for the rest of the terms, a(n*(n+1)) ,..., a(n+(n+1)^2) = 2*(n+1). It can be shown that this definition satisfies the recurrence relation. - _Chai Wah Wu_, Jun 06 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Jul 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006324 | OEIS-A193418

Expansion of x*(x^2+x-1)/(3*x^6-4*x^5+x^4-3*x^2+4*x-1).

Conjecture: log(A005960-a(n)) ~ (log(2)*(2*n-11)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yalcin Aktar_, Jul 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006325 | OEIS-A193437

Expansion of e.g.f. exp(Sum_{n>=0} x^(3*n+1)/(3*n+1)).

Conjecture: a(n) is divisible by 7^floor(n/7) for n>=0. Conjecture: a(n) is divisible by p^floor(n/p) for prime p == 1 (mod 3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006326 | OEIS-A193438

Expansion of e.g.f. exp(Sum_{n>=0} x^(4*n+1)/(4*n+1)).

Conjecture: a(n) is divisible by 5^floor(n/5) for n>=0. Conjecture: a(n) is divisible by p^floor(n/p) for prime p == 1 (mod 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006327 | OEIS-A193550

O.g.f.: 1/(1 - x/(1 - x/(1 - 9*x/(1 - 9*x/(1 - 25*x/(1 - 25*x/(1 - 49*x/(1 - 49*x/(1-...)))))))))), a continued fraction involving the odd squares.

Conjecture: given o.g.f. A(x), then sqrt((A(x) - 1)/x) is an integer series.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006328 | OEIS-A193555

Numerators of the squared radii of the smallest enclosing circles of n points with integer coordinates and distinct mutual distances, arranged such that the radius of their enclosing circle is minimized. Denominators are given in A193556.

Conjectured next terms are a(17)/A193556(17)=19720/121, a(18)/A193556(18)=5002/25.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jul 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006329 | OEIS-A193558

Differences between consecutive primes of the form k^2+1.

It is conjectured that the sequence of primes of the form k^2+1 is infinite, but this has never been proved. This sequence contains a subset of squares: {64, 144, 100, 1024, 4900, 10816, 11664, 12544, 18496, 102400, 41616, ...}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jul 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006330 | OEIS-A193574

Smallest divisor of sigma(n) that does not divide n.

(4) Conjecture: a(n) > n implies a(n) = sigma(n); tested through n = 20000000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Aug 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006331 | OEIS-A193628

Integer part of A192851 / A191626.

Conjecture: the sequence seems to converge asymptotically to 113.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrea Raffetti_, Aug 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006332 | OEIS-A193649

Q-residue of the (n+1)st Fibonacci polynomial, where Q is the triangular array (t(i,j)) given by t(i,j)=1. (See Comments.)

Conjecture: G.f.: -(1+x)*(2*x-1) / ((x-1)*(4*x^2+x-1)). - _R. J. Mathar_, Feb 19 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006333 | OEIS-A193654

Q-residue of the triangle p(n,k)=floor((n+1)/(n+k+2)/2), 0<=k<=n, where Q is the triangular array (t(i,j)) given by t(i,j)=1. (See Comments.)

Conjecture: G.f.: (-1-2*x+4*x^2+4*x^3-8*x^5) / ((1+x)*(2*x+1)*(x-1)^2*(2*x-1)^3). - _R. J. Mathar_, Feb 19 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006334 | OEIS-A193655

Q-residue of the triangle p(n,k)=floor(1/2+(n+1)/(n+k+2)/2), 0<=k<=n, where Q is the triangular array (t(i,j)) given by t(i,j)=1. (See Comments.)

Conjecture: G.f.: (-1-2*x+3*x^2+9*x^3-8*x^4-4*x^5) / ((1+x)*(2*x+1)*(x-1)^2*(2*x-1)^3). - _R. J. Mathar_, Feb 19 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006335 | OEIS-A193656

Q-residue of the triangle p(n,k)=(2^(n - k))*5^k, 0<=k<=n, where Q is the triangular array (t(i,j)) given by t(i,j)=1. (See Comments.)

Conjecture: a(n) = 3*5^n-2*4^n; G.f.: (1-2*x) / ((5*x-1)*(4*x-1)). - _R. J. Mathar_, Feb 19 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006336 | OEIS-A193658

Q-residue of the triangle A051162, where Q is the triangular array (t(i,j)) given by t(i,j)=1. (See Comments.)

Conjecture: a(n) +(-n-4)*a(n-1) +(4*n-1)*a(n-2) +5*(-n+2)*a(n-3) +2*(n-3)*a(n-4)=0. - _R. J. Mathar_, Feb 19 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006337 | OEIS-A193659

Q-residue of the triangle A094727, where Q=Pascal's triangle. (See Comments.)

Conjecture: a(n) +(-n-5)*a(n-1) +2*(2*n+1)*a(n-2) +(-5*n+8)*a(n-3) +2*(n-3)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006338 | OEIS-A193661

Q-residue of the triangle A193673, where Q is the triangular array (t(i,j)) given by t(i,j)=1. (See Comments.)

Conjecture: G.f.: (-1+8*x-13*x^2) / ((x-1)*(3*x-1)*(7*x-1)). - _R. J. Mathar_, Feb 19 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006339 | OEIS-A193662

Q-residue of the Lucas triangle A114525, where Q is the triangle given by t(i,j)=1 for 0<=i<=j. (See Comments.)

Conjecture: a(n) = 2*a(n-1) +3*a(n-2) -4*a(n-3) if n>3. - _R. J. Mathar_, Feb 19 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006340 | OEIS-A193663

Q-residue of A049310 (triangle of coefficients of Fibonacci polynomials), where Q is the triangle given by t(n,k)=k+1 for 0<=k<=n. (See Comments.)

Conjecture: G.f.: x*(1-x+x^2) / (1-2*x-6*x^2+7*x^3+x^4). - _R. J. Mathar_, Feb 19 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006341 | OEIS-A193665

Q-residue of A075392, where Q=A075392. (See Comments.)

Conjecture: a(n) +(-n-4)*a(n-1) +(n+1)*a(n-2) -a(n-3)=0. - _R. J. Mathar_, Feb 19 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006342 | OEIS-A193679

Sequence related to discriminant of cyclotomic polynomials A004124.

Conjecture: Dirichlet g.f. of log(a(n)): -zeta(s-1)*zeta'(s)/zeta(s)^2, where zeta'(s) is the derivative of zeta(s). This would give a(n) = exp(Sum_{d|n} Lambda(d)*phi(n/d)), with Lambda(n)=log(A014963) and phi(n)=A000010. - _Benedict W. J. Irwin_, Jul 14 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Aug 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006343 | OEIS-A193855

Primes p such that tau(p) is congruent to 1 (mod p), where tau is the Ramanujan tau function.

M. J. Hopkins wrote "It is not known whether or not tau(p) == 1 mod p holds for infinitely many primes". For more information about this open problem see the Sloane comment in A000594.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006344 | OEIS-A193918

Mirror of the triangle A193917.

Here, we extend of the conjecture begun at A193917. Suppose n is an even positive integer and r(n+1,x) is the polynomial matched to row n+1 of A193918 as in the Mathematica program, where the first row is counted as row 0. Conjecture: r(n+1,x) is the product of the following two polynomials whose coefficients are Fibonacci numbers:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006345 | OEIS-A193948

Integers k such that for all i > k the largest prime factor of i(i+1)(i+2)(i+3)(i+4)(i+5)(i+6) exceeds the largest prime factor of k(k+1)(k+2)(k+3)(k+4)(k+5)(k+6).

Currently terms through a(16) = 12874 have been proved to be correct, while remaining terms are conjectural. Stormer's theorem (see link) provides an approach to proving their correctness. Such an approach should be easy for the first few terms. - _Franklin T. Adams-Watters_, Nov 07 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrey V. Kulsha_, Aug 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006346 | OEIS-A194001

Mirror of the triangle A194000.

Here, we extend of the conjecture begun at A194000. Suppose n is an odd positive integer and r(n+1,x) is the polynomial matched to row n+1 of A194001 as in the Mathematica program, where the first row is counted as row 0. Conjecture: r(n+1,x) is the product of the following two polynomials whose coefficients are Fibonacci numbers:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006347 | OEIS-A194263

Numbers that occur more than once in A193873, in order of appearance.

This conjecture only holds for n<62: a(62)=A193873(124)=A193873(2n)=620378449

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Sep 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006348 | OEIS-A194271

Number of toothpicks or D-toothpicks added at n-th stage to the structure of A194270.

Conjecture: a(n) = 4*(b(n-1)-d(n)) + 2*(c(n)-d(n+1)) + 2*(c(n+2)-d(n+1)) + 8, if n >= 3. Conjecture: a(2^k+2) = 32, if k >= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006349 | OEIS-A194362

Decimal expansion of log(1151) / log(95).

This number occurs in A143898. This value K is conjectured to be the least possible such that there is at least one prime in the range n^K to (n+1)^K for n>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Oct 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006350 | OEIS-A194428

Number of iterations of the map n->n/3 if n == 0 (mod 3), n->4*n+a if 4*n+a == 0 (mod 3) where a = 1 or 2, before reaching the end of the cycle.

The problem is as follows: start with any number n. If n is divisible by 3, divide it by 3, otherwise multiply it by 4 and add 1 or 2 in order to find a new integer divisible by 3. Do we always reach the end of a cycle? It is conjectured that the

answer is yes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Aug 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006351 | OEIS-A194433

Number of toothpicks and D-toothpicks added at n-th stage to the structure of A194432.

Conjecture: a(2^k+1) = 16, if k >= 1 .

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Sep 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006352 | OEIS-A194435

Number of toothpicks or D-toothpicks added at n-th stage to the structure of A194434.

Conjecture: a(2^k+1) = 16, if k >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Sep 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006353 | OEIS-A194441

Number of toothpicks or D-toothpicks added at n-th stage to the structure of A194440.

Conjectures for n = 2^k+j, if -1<=j<=3: End of conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006354 | OEIS-A194443

Number of toothpicks or D-toothpicks added at n-th stage to the structure of A194442.

Conjectures for n = 2^k+j, if -6<=j<=6: End of conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006355 | OEIS-A194469

Values of m for which sqrt(m) is curbed by 1/2; see Comments for "curbed by".

Suppose that r and c are real numbers, that 0<c<1, and that s(n)=sum{<c+k*r>-<k*r> : 1<=k<=n}, where < > denotes fractional part. The inequalities s(n)<0, s(n)=0, s(n)>0 yield up to three sequences that partition the set of positive integers, as in the examples cited at A194368. If s(n)>=0 for every n>=1, we say that r is curbed by c. For r=sqrt(m), clearly r is curbed by 1/2 if m is a square. Conjecture: there are infinitely many nonsquare m for which sqrt(m) is curbed by 1/2, and there are infinitely many m for which sqrt(m) is not curbed by 1/2 (see A194470). The terms shown here for A194469 are conjectured, based on examinations of s(n) for 1<=n<=B for various B>100.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006356 | OEIS-A194470

Complement of A194469 (conjectured): numbers m such that sqrt(m) is not curbed by 1/2.

Complement of A194469 (conjectured): numbers m such that sqrt(m) is not curbed by 1/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006357 | OEIS-A194530

Number of unlabeled rigid interval posets with n non-maximal and 2 maximal elements.

Conjectural g.f.:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Aug 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006358 | OEIS-A194558

G.f.: A(x) = exp(Sum_{n>=1} G_n(x)^n/n) where G_n(x) = x + x*G_n(x)^n and A(x) = Sum_{n>=1} a(n)*x^n/floor(n/2)!.

This sequence is conjectured to consist entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006359 | OEIS-A194584

Differences of A035336.

Proof of this conjecture: since A035336(n) = (2*floor(n*phi) + n - 1) (with phi = (1+sqrt(5))/2) is a generalized Beatty sequence, this follows from Lemma 4 in Allouche and Dekking. - _Michel Dekking_, Oct 10 2018 Proof of this conjecture: this follows from the Carlitz-Scoville-Hoggatt theorem: compositions of the Wythoff A and B sequences are generalized Beatty sequences (cf. Theorem 1 in Allouche and Dekking). - _Michel Dekking_, Oct 10 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Aug 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006360 | OEIS-A194595

Triangle by rows T(n,k), showing the number of meanders with length (n+1)*3 and containing (k+1)*3 L's and (n-k)*3 R's, where L's and R's denote arcs of equal length and a central angle of 120 degrees which are positively or negatively oriented.

The conjectured formulas are confirmed by dynamic programming for 0 <= n <= 62. - _Susanne Wienand_, Jun 24 2015 Recursive formula (conjectured): Closed formula (conjectured): T(n,k) = (C(n,k))^3 + C(n,k) * C(n,k+1) * C(n+1,k+1). - _Susanne Wienand_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Susanne Wienand_, Oct 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006361 | OEIS-A194688

First differences of A036554 (numbers whose binary representation ends in an odd number of zeros).

Conjecture. This sequence is self-generated according to the following rule: start with {4} at step 0, then extend by steps, appending {2,2,4} at step n if a(n)=4 or appending {4} if a(n)=2. (This has been verified for several thousand terms.) To illustrate, the first few steps of this process give {4}->{4,2,2,4}, since a(1)=4, ->{4,2,2,4,4}, since a(2)=2, ->{4,2,2,4,4,4}, since a(3)=2, ->{4,2,2,4,4,4,2,2,4}, since

a(4)=4, etc. Equivalently, it appears that {a(n)} is the fixed-point of the morphism 2->4, 4->422, starting with 4. Since A036554 = 2*A003159, this conjecture follows from the paper by Allouche, Shallit and Skordev in 2005, see page 13. - _Michel Dekking_, Jan 06 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Sep 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006362 | OEIS-A194693

Number of toothpicks or D-toothpicks added at n-th stage to the D-toothpick "corner" structure of A194692.

Conjectures for n = 2^k+j, if 0<=j<=3: End conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Sep 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006363 | OEIS-A194701

Number of toothpicks or D-toothpicks added at n-th stage to the structure of A194700.

Essentially the first differences of A194700. First differs from A194271 at a(15). Conjecture: this sequence and A194271 have infinitely many numbers in common.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Sep 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006364 | OEIS-A194724

Number of quaternary words either empty or beginning with the first character of the alphabet, that can be built by inserting n doublets into the initially empty word.

Conjecture: n*a(n) +2*(-14*n+9)*a(n-1) +96*(2*n-3)*a(n-2)=0. - _R. J. Mathar_, Mar 14 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Sep 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006365 | OEIS-A194772

Number of lower triangles of an (n+2) X (n+2) 0..2 array with new values introduced in row major order 0..2 and no element unequal to more than one horizontal or vertical neighbor.

Conjectures from _Colin Barker_, May 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006366 | OEIS-A194894

The number of the ordered triples (A,B,C) satisfying the system of the modular relations {A*B - B*A = C, B*C - C*B = A, C*A - A*C = B}, where A,B,C are distinct 2 X 2 matrices over Z(n).

Remarks for the cases n<=41 (conjectures for n>41):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Erdos Pal_, Sep 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006367 | OEIS-A194943

Greatest d such that d*n+b is the least prime in the arithmetic progression k*n+b for some 0 < b < n with gcd(b, n) = 1.

Heath-Brown's conjecture on Linnik's theorem implies that a(n) < n. Pomerance shows that a(n) > (e^gamma + o(1)) log(n) * phi(n)/n, and Granville & Pomerance conjecture that a(n) >> log(n)^2 * phi(n)/n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Sep 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006368 | OEIS-A194959

Fractalization of (1 + floor(n/2)).

Sum_{k=1..n} A(k,n+1-k)*A209229(k) = 2*n-1. (conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006369 | OEIS-A195000

Number of lower triangles of a 4 X 4 0..n array with each element differing from all of its horizontal and vertical neighbors by one.

Conjectures from _Colin Barker_, May 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006370 | OEIS-A195060

Numbers on the main diagonals and on the axes (x,y) in the square spiral whose vertices are the generalized pentagonal numbers A001318.

Conjectures: a(n) = 2*a(n-1) - 2*a(n-2) + 2*a(n-3) - a(n-4) + a(n-8) - 2*a(n-9) + 2*a(n-10) - 2*a(n-11) + a(n-12); g.f.: -x*(1+x^2+x^3-x^4+x^5+x^7-2*x^8+x^11+2*x^9-2*x^10) / ((1+x)*(x^4+1)*(x^2+1)^2*(x-1)^3). - _R. J. Mathar_, Oct 11 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Sep 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006371 | OEIS-A195100

Numbers n such that there are no primes between (n-1)*sqrt(n-1) and n*sqrt(n).

Cramér's conjecture implies that the sequence is finite. - _Robert Israel_, Aug 11 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Sep 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006372 | OEIS-A195101

Solid numbers. Numbers m such that A005245(m) < A005245(k) + A005245(m-k) for 1<= k < m.

Conjecture that for every solid number m > 1 there is some number n such that A005245(n) = A005245(m)+A005245(n-m) and such that for every representation as a product n = u*v with u, v >= 2 or every 1 < = k < m, we have A005245(n) < A005245(u)+A005245(v) and A005245(n) < A005245(k) + A005245(n-k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juan Arias-de-Reyna_, Sep 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006373 | OEIS-A195194

G.f.: A(x) = INV(x - x*INV(x - 2*x*INV(x - 3*x*INV(x - 4*x*INV(x - ...)))))), where INV(F(x)) = series reversion of F(x).

Conjecture: a(n) is odd iff n is a power of 2. - _Paul D. Hanna_, Apr 07 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006374 | OEIS-A195204

Triangle of coefficients of a sequence of binomial type polynomials.

Conjecture: o.g.f. as a continued fraction of Stieltjes type: 1/(1 - 2*x*z/(1 - z/(1 - 2*(x + 1)*z/(1 - 2*z/(1 - 2*(x + 2)*z/(1 - 3*z/(1 - 2*(x + 3)*z/(1 - 4*z/(1 - ...))))))))). - _Peter Bala_, Dec 12 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 13 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006375 | OEIS-A195233

Number of lower triangles of a 3 X 3 0..n array with each element differing from all of its diagonal, vertical, antidiagonal and horizontal neighbors by one or less.

Conjectures from _Colin Barker_, May 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 13 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006376 | OEIS-A195234

Number of lower triangles of a 4 X 4 0..n array with each element differing from all of its diagonal, vertical, antidiagonal and horizontal neighbors by one or less.

Conjectures from _Colin Barker_, May 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 13 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006377 | OEIS-A195235

Number of lower triangles of a 5 X 5 0..n array with each element differing from all of its diagonal, vertical, antidiagonal and horizontal neighbors by one or less.

Conjectures from _Colin Barker_, May 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 13 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006378 | OEIS-A195236

Number of lower triangles of a 6 X 6 0..n array with each element differing from all of its diagonal, vertical, antidiagonal and horizontal neighbors by one or less.

Conjectures from _Colin Barker_, May 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 13 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006379 | OEIS-A195237

Number of lower triangles of a 7 X 7 0..n array with each element differing from all of its diagonal, vertical, antidiagonal and horizontal neighbors by one or less.

Conjectures from _Colin Barker_, May 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 13 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006380 | OEIS-A195249

Number of lower triangles of a 3 X 3 0..n array with each element differing from all of its diagonal, vertical, antidiagonal and horizontal neighbors by two or less.

Conjectures from _Colin Barker_, May 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 13 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006381 | OEIS-A195264

Iterate x -> A080670(x) (replace x with the concatenation of the primes and exponents in its prime factorization) starting at n until reach 1 or a prime (which is then the value of a(n)); or a(n) = -1 if a prime is never reached.

J. H. Conway offered $1000 for a proof or disproof for his conjecture that every number eventually reaches a 1 or a prime - see OEIS50 link. - _N. J. A. Sloane_, Oct 15 2014 The number D' = 13^532385396179 maps to D and so is a much larger number with a(D') = -1. Repeating this process (by finding a prime prefix of D') should lead to an infinite sequence of counterexamples to Conway's conjecture. - _Hans Havermann_, Jun 09 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 14 2011, based on discussions on the Sequence Fans Mailing List by _Alonso del Arte_, _Franklin T. Adams-Watters_, _D. S. McNeil_, _Charles R Greathouse IV_, _Sean A. Irvine_, and others | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006382 | OEIS-A195279

Number of lower triangles of a 3 X 3 integer array with each element differing from all of its vertical and horizontal neighbors by n or less and triangles differing by a constant counted only once.

Conjectures from _Colin Barker_, Mar 01 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006383 | OEIS-A195307

Where records occur in A129308 and also in A195155.

Conjecture: for each term k > 1 in the sequence there exists prime p such that k/p is in the sequence. In the list of conjectured terms, if actual terms <= 10^16 are 97-smooth and have the following property: a(n+1) = a(n) + k*gcd(a(n), a(n-1), ..., a(n-20)) setting a(n) = 1 for n < 1 then those terms are actual terms. The conjectured terms are 41-smooth and satisfy a(n+1) = a(n) + k*gcd(a(n), a(n-1), ..., a(n-13)). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Oct 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006384 | OEIS-A195334

Numbers the sum of whose even divisors is 2 times a prime.

Conjecture: a(n) = 2*A023194(n). - _R. J. Mathar_, Sep 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Sep 15 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006385 | OEIS-A195350

Expansion of (1 - 3*x - x^2)/(1 - 4*x + 2*x^3 + x^4).

a(n) = A301483(n) - A303647(n-2) + A195339(n-4) (conjectured). - _Gregory Gerard Wojnar_, Apr 27 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bruno Berselli_, Sep 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006386 | OEIS-A195378

Numbers n such that n-th Fibonacci number has all digits 0 to 9.

Brian Trial conjectures that all Fibonacci numbers greater than F285 are pandigital, see the Gupta link. Conjecture: a(n) = n + 131 for n > 154.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kausthub Gudipati_, Sep 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006387 | OEIS-A195422

Permanents of certain n X 2 cyclic (1,-1) matrices.

Conjecture: (-n+2)*a(n) +(n^2-4*n+6)*a(n-1) +2*(n^2-2*n+3)*a(n-2) +8*(n-1)*a(n-3) = 0. - _R. J. Mathar_, Jul 20 2016 Conjecture: a(n) -n*a(n-1) -8*a(n-2) +4*(n-4)*a(n-3) +16*a(n-4) = 0. - _R. J. Mathar_, Jul 20 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Sep 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006388 | OEIS-A195441

a(n) = denominator(Bernoulli_{n+1}(x) - Bernoulli_{n+1}).

The equation a(n-1) = denominator(Bernoulli_n(x) - Bernoulli_n) = rad(n+1) has only finitely many solutions, where rad(n) = A007947(n) is the radical of n. It is conjectured that S = {3, 5, 8, 9, 11, 27, 29, 35, 59} is the full set of all such solutions. Note that (S\{8})+1 joined with {1,2} equals A094960. More precisely, the set S implies the finite sequence of A094960. See Kellner 2023. - _Bernd C. Kellner_, Oct 18 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Sep 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006389 | OEIS-A195442

G.f.: 1 = Sum_{n>=0} a(n)*x^n / Product_{k=1..n+1} (1+k*x)^4.

CONJECTURES. (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006390 | OEIS-A195443

Self-convolution square equals A195442.

CONJECTURES. (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006391 | OEIS-A195465

The first a(n) n-gap primes are lessers of twin primes, a(n) maximal.

Conjecture: a(n)>0 for n>1. This conjecture is equivalent to the conjecture that all terms of A195325 are lessers of twin primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006392 | OEIS-A195507

Smallest integer m greater than n such that m (mod k) == n (mod k) for k = 1..n-1.

Conjecture: The sequence (a(n): n >= 1) satisfies the following recurrence. Write a(n)/n in lowest terms as num/d. Then a(n+1) = d*a(n) - (d-1)*n + 1. Illustration: a(4) = 10 and a(4)/4 = 5/2 in lowest terms. Then a(5) = 2*10 - 1*4 + 1 = 17. (This has been verified up to a(23) = 232792583.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Sep 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006393 | OEIS-A195523

Number of lower triangles of a 5 X 5 -n..n array with all row and column sums zero.

Conjectures from _Colin Barker_, May 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006394 | OEIS-A195538

Denominators a(n) of Pythagorean approximations b(n)/a(n) to sqrt(8).

Conjecture: a(n) = 35*a(n-2) - 35*a(n-4) + a(n-6) with bisections A098602 and A076218. - _R. J. Mathar_, Sep 21 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006395 | OEIS-A195539

Numerators b(n) of Pythagorean approximations b(n)/a(n) to sqrt(8).

Conjecture: a(n) = 34*a(n-2) - a(n-4) with a(2*n+1) = A000129(4*n+4), a(2*n) = A046176(n+1). - _R. J. Mathar_, Sep 21 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006396 | OEIS-A195549

Hypotenuses of primitive Pythagorean triples in A195547 and A195548.

Conjectures from _Colin Barker_, Apr 08 2012: (Start) Conjecture: a(n) is the denominator of the reduced fraction (F(2*n+1)-2)/F(2*n+1), F(n) = Fibonacci(n). - _Sébastien Labbé_, May 06 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006397 | OEIS-A195556

Denominators a(n) of Pythagorean approximations b(n)/a(n) to 1/3.

Conjecture: a(n) = 37*a(n-3) + 37*a(n-6) - a(n-9). - _R. J. Mathar_, Sep 21 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 21 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006398 | OEIS-A195557

Numerators b(n) of Pythagorean approximations b(n)/a(n) to 1/3.

Conjecture: a(n) = +37*a(n-3) +37*a(n-6) -a(n-9). - _R. J. Mathar_, Sep 21 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 21 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006399 | OEIS-A195562

Denominators a(n) of Pythagorean approximations b(n)/a(n) to 1/4.

Conjecture: a(n) = 65*a(n-3) + 65*a(n-6) - a(n-9). - _R. J. Mathar_, Sep 21 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 21 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006400 | OEIS-A195571

Denominators a(n) of Pythagorean approximations b(n)/a(n) to 1/5.

Conjecture: a(n) = 101*a(n-3) + 101*a(n-6) - a(n-9). - _R. J. Mathar_, Sep 21 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 21 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006401 | OEIS-A195572

Numerators b(n) of Pythagorean approximations b(n)/a(n) to 1/5.

Conjecture: satisfies a linear recurrence having signature (0, 0, 101, 0, 0, 101, 0, 0, -1). - _Harvey P. Dale_, Jun 13 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 21 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006402 | OEIS-A195588

a(n) = Sum_{k=0..2*n} (-1)^k * moebius(2*n-k+1) * moebius(k+1).

It is conjectured that all integers appear an infinite number of times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006403 | OEIS-A195608

Numbers n such that Sum_{i=1..n} A(i) = A(n)*A(n+1)/4, where A(n) = A001969(n).

Conjectures: 1) there are only 3 different first differences 1,4,6; 2) the sequence contains either isolated series of terms, e.g., {1},{7},{25},{31},..., or series of 3 consecutive integers, e.g., {11,12,13}, {19,20,21}, etc.; 3)the first terms m of every series satisfy the condition A(m+1)-A(m-1)=5, where A(n)=A001969(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 21 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006404 | OEIS-A195609

Numbers n such that Sum_{i=1..n} A(i) = A(n)*A(n+1)/4, where A(n) = A000069(n).

Conjectures: 1) there are only 3 different first differences 1,4,6; 2) the sequence contains either isolated series of terms, e.g., {9},{23},{33},{39},..., or series of 3 consecutive integers, e.g., {3,4,5}, {15,16,17}, etc.; 3)the first terms m of every series satisfy the condition A(m+1)-A(m-1)=5, where A(n)=A000069(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 21 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006405 | OEIS-A195673

Triangle T(n,k) read by rows: T(0,0)=-2, T(1,0)=3, T(1,1)=0 and T(n,k) = T(n-1,k)-T(n-2,k-2) otherwise.

Conjecture: The polynomials p(n,x) = sum_{k=0..n} T(n,k)*x^(n-k) based on this simple recurrence for other initial constant values of T(0,0)=p and T(1,0)=q are related to the S-polynomials of A053119: p(n,x,p+1,q+1)-p(n,x,p,q) = S(n,x).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Sep 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006406 | OEIS-A195806

Number of triangular of a 5 X 5 X 5 0..n arrays with all rows and diagonals having the same length having the same sum, with corners zero.

Conjectured recurrence: a(n) - 3*a(n+1) + 2*a(n+2) - a(n+3) + 6*a(n+4) - 5*a(n+5) - 3*a(n+6) + 3*a(n+8) + 5*a(n+9) - 6*a(n+10) + a(n+11) - 2*a(n+12) + 3*a(n+13) - a(n+14) = 0. Conjectured closed form as a quasi-polynomial:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006407 | OEIS-A195825

Square array T(n,k) read by antidiagonals, n>=0, k>=1, which arises from a generalization of Euler's Pentagonal Number Theorem.

Conjecture: if k is odd then column k contains (k+1)/2 plateaus whose levels are the first (k+1)/2 terms of A210843 and whose lengths are k+1, k-1, k-3, k-5, ... 2. Otherwise, if k is even then column k contains k/2 plateaus whose levels are the first k/2 terms of A210843 and whose lengths are k+1, k-1, k-3, k-5, ... 3. The sequence A210843 gives the levels of the plateaus of column k, when k -> infinity. For the visualization of the plateaus see the graph of a column, for example see the graph of A210964. - _Omar E. Pol_, Jun 21 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Sep 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006408 | OEIS-A195871

a(n) is the smallest prime(m) such that the interval (prime(m)*n, prime(m+1)*n) contains exactly one prime.

Conjecture: In the supposition that there are infinitely many twin primes, for n>=5 all terms are in A001359 (lesser of twin primes). Note that a unique prime which is contained in an interval of the form (prime(m)*n, prime(m+1)*n) is called n-isolated (see author's link, where a heuristic proof is given that the number of n-isolated primes<=x approaches e^{-2(n-1)}x/log(x) as x goes to infinity (cf. Conjecture 25, Remark 26 and formula (47)). One can easily prove that a(n) is not bounded. This conjecture seems hard, since it's not obvious how to find an upper bound for a(n) (see Conjecture 42 in the Shevelev link). - _Charles R Greathouse IV_, Jan 02 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006409 | OEIS-A195901

Integers k>1 such that A141822(k)=5.

Zaremba conjectured that A141823, A141832, A141833, and this sequence form a partition of the integers >1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Sep 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006410 | OEIS-A195915

Table with T(n,1) = n, otherwise T(n,k) = xor(T(n-1,k-1), T(n-1,k)).

It appears that, if 2^j > n, T(2^j+n, 2^j+k) = T(n,k). This is equivalent to the periodicity conjecture in A195916.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Sep 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006411 | OEIS-A196020

Irregular triangle read by rows: T(n,k), n >= 1, k >= 1, in which column k lists the odd numbers interleaved with k-1 zeros, and the first element of column k is in row k(k+1)/2.

Conjecture: T(n,k) is the difference between the square of the total number of partitions of all positive integers <= n into exactly k consecutive parts, and the square of the total number of partitions of all positive integers < n into exactly k consecutive parts. - _Omar E. Pol_, Feb 14 2018 For the connection with A280851 see also the algorithm of A280850 and the conjecture of A296508. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Feb 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006412 | OEIS-A196079

Difference between the largest and smallest inverse of totient function.

No terms are zero if Carmichael's conjecture is true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franz Vrabec_, Sep 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006413 | OEIS-A196129

A002110(n)-(p[i]+p[i+1]+...+p[i+n-1]), where p[i] is the largest prime such that this is nonnegative.

It is an open problem whether there is n>1 (necessarily even) such that a(n)=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Sep 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006414 | OEIS-A196130

The difference prime(i)+prime(i+1)+...+prime(i+n-1)-A002110(n), where prime(i) is the smallest prime such that the value is nonnegative.

It is an open problem whether there is n>1 (necessarily even) such that a(n)=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Sep 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006415 | OEIS-A196200

Numbers k such that Euler phi(Dedekind psi(k)) > k.

Counterexamples to a conjecture by Sandor.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Dec 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006416 | OEIS-A196226

m such that A054024(m) (sum of divisors of m reduced modulo m) is 3 + m/2.

Conjectures. (1) If m>=14 is a term of this sequence, then sigma(2,m) is congruent to 5 + m/2 modulo m; (2) If m>=22 is a term of this sequence, then sigma(3,m) is congruent to 9 + m/2 modulo m; If m>=38 is a term of this sequence, then sigma(4,m) is congruent to 17 + m/2 modulo m. (sigma(k,m) denotes the sum of the k-th powers of the divisors of m.) Similar conjectures can be made about sigma(k,m) congruent to 2^k+1 + m/2 modulo m, for m a sufficiently large term of this sequence..

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Sep 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006417 | OEIS-A196279

Let r= (7n) mod 10 and x=floor(7n/10) be the last digit and leading part of 7n. Then a(n) = (x-2r)/7.

Conjecture: a(n)= +a(n-1) +a(n-10) -a(n-11) with G.f. x*(-2 +x -2*x^3 +x^4 -2*x^6 +x^7 +x^9 +x^2 +x^5 +x^8) / ((1+x) *(x^4+x^3+x^2+x+1) *(x^4-x^3+x^2-x+1)

*(x-1)^2). - _R. J. Mathar_, Oct 04 2011 The conjecture above is correct. - _Charles R Greathouse IV_, Jan 04 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Sep 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006418 | OEIS-A196523

G.f. A(x) satisfies A(x) = x + x*A(A(A(A(x)))).

Conjecture: a(n) == 1 (mod 3) for n >= 1. - _Paul D. Hanna_, Dec 01 2024 The conjecture is true, and more generally if A(x) = x + x*A^[l](x), where A^[l] is the l-th iterate of A, then a(n) == 1 (mod l-1) for n >= 1. Proof: the equation determines a(n) from a(1..n-1) by integer polynomial recurrences, so A mod l-1 is the unique solution over Z/(l-1). Let B(x) = x/(1-x); by induction B^[k](x) = x/(1-k*x). Since l == 1 (mod l-1), B^[l] = B, and x + x*B = x/(1-x) = B. Hence A == x/(1-x) (mod l-1). - _Amol Bhave_, Oct 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006419 | OEIS-A196579

Length of longest possible 90-degree snake inside a circle of diameter n.

These solutions have been found by experimentation and are in that respect conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kausthub Gudipati_, Oct 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006420 | OEIS-A196660

Smallest k>0 such that k*n+(n-1) is prime.

Conjecture: for every n there exists k < n (apart from a(1)) such that k*n+(n-1) is prime. See A034693.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Oct 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006421 | OEIS-A196750

Numbers k such that 3, 5, 7 and 11 do not divide swing(k) = A056040(k).

It is conjectured that there are no other terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Oct 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006422 | OEIS-A196779

a(n) is the smallest number m such that no prime takes the form of n^m+/-n^k+/-1, while 0 <= k < m and m > 1.

Conjecture: a(n) has finite value when a>4 Hypothesis is that a(2), a(3), and a(4) are infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Oct 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006423 | OEIS-A196837

Coefficient table of numerator polynomials of o.g.f.s for partial sums of powers of positive integers.

This e.g.f. - o.g.f. connection proves some conjectures by _Simon Plouffe_. See the o.g.f. Maple programs under, e.g., A001551(n=4) and A001552 (n=5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006424 | OEIS-A196934

a(n) is the first occurrence of n in sequence A078498.

Conjecture: Any prime number greater than 11 (p) can be the center term of arithmetic progressions prime chain p-6k, p, p+6k, while k>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Oct 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006425 | OEIS-A196935

a(n) is the number of arithmetic progressions prime chains in the form of p(n)-6k, p(n), p(n)+6k, while k > 0 and p(n) > 6k.

Conjecture: a(n) > 0 for all n >= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Oct 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006426 | OEIS-A197171

Values k such that singular quadratic unity of Q(k) have gcd(k,y) = 2.

Conjecture: This sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Oct 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006427 | OEIS-A197184

Triangle of polynomial coefficients of the polynomial factors defined in A074051.

Conjectures on p_n(x) = Sum_{k=0..n-1} T(n,k)*x^k:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Oct 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006428 | OEIS-A197521

Decimal expansion of least x>0 having cos(Pi*x/2)=(cos Pi*x/3)^2.

Conjecture: the constant here, 3.52133782..., is 3 plus the constant in A197383, the latter being the least t>0 satisfying sin(Pi*t/6)=(sin Pi*t/3)^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006429 | OEIS-A197653

Triangle by rows T(n,k), showing the number of meanders with length (n+1)*4 and containing (k+1)*4 Ls and (n-k)*4 Rs, where Ls and Rs denote arcs of equal length and a central angle of 90 degrees which are positively or negatively oriented.

The conjectured formulas are confirmed by dynamic programming for 0 <= n <= 43. - _Susanne Wienand_, Jun 29 2015 Recursive formula (conjectured): closed formula (conjectured): T(n,n) = 1, k = n

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Susanne Wienand_, Oct 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006430 | OEIS-A197654

Triangle by rows T(n,k), showing the number of meanders with length 5(n+1) and containing 5(k+1) L's and 5(n-k) R's, where L's and R's denote arcs of equal length and a central angle of 72 degrees which are positively or negatively oriented.

The conjectured formulas are confirmed by dynamic programming for 0 <= n <= 29. - _Susanne Wienand_, Jul 01 2015 Recursive formula (conjectured): Closed formula (conjectured):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Susanne Wienand_, Oct 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006431 | OEIS-A197655

Triangle by rows T(n,k), showing the number of meanders with length (n+1)*6 and containing (k+1)*6 Ls and (n-k)*6 Rs, where Ls and Rs denote arcs of equal length and a central angle of 60 degrees which are positively or negatively oriented.

The conjectured formulas are confirmed by dynamic programming for 0 <= n <= 19. - _Susanne Wienand_, Jul 04 2015 recursive formula (conjectured): closed formula (conjectured): T(n,n) = 1, k = n

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Susanne Wienand_, Oct 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006432 | OEIS-A197657

Row sums of A194595.

Conjecture: (n+1)^2*a(n) -3*(n+1)*(2*n+1)*a(n-1) -3*n*(5*n-7)*a(n-2) -8*(n-2)^2*a(n-3)=0. - _R. J. Mathar_, Jul 26 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Susanne Wienand_, Oct 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006433 | OEIS-A197702

Smallest positive integer k such that n = +-1 +-3 +-... +-(2k-1) for some choice of +'s and -'s.

Conjecture. Let SO(k) be the sum of the first k odd positive integers. Then a(n)=k if n=SO(k). Otherwise, choose k so that SO(k-1)<n<SO(k). Then if SO(k)-n=4, a(n)=k+2, else if SO(k)-n is odd then a(n)=k+1 else a(n)=k. (This has been verified for n up to 200.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Oct 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006434 | OEIS-A197877

Smallest number m such that A096535(m) = n.

A096535(a(n)) = n and A096535(m) <> n for m < a(n), concerning first conjecture in A096535.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Oct 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006435 | OEIS-A197879

Parity of floor(n*sqrt(8)).

If the fractional part of sqrt(2)*n > 1/2 then a(n) = 1, otherwise a(n) = 0. It is possible to see from this that since 5/4 < sqrt(2) < 7/4, there are no more than two consecutive 0's or 1's (a similar feature is found in A272532 and conjectured in A272170). The sequence looks quasiperiodic and its Fourier spectrum seems to present a maximum component at a frequency which converges to about 0.828 of the maximum frequency. - _Andres Cicuttin_, Jul 09 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Oct 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006436 | OEIS-A197918

Pythagorean primes p such that for all primes q < p, p XOR q is not equal to p - q.

It is conjectured that with the exception of the first three terms (2,5,17) all of the terms are a subset of all primes p such that p XOR 22 = p + 22.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brad Clardy_, Oct 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006437 | OEIS-A197988

Number of isomorphism classes of nanocones with 3 pentagons and a symmetric boundary of length n.

Conjectured g.f.: (x^7 + x^5 + x^4 + 2*x^3 + 2*x + 1)*x/((x - 1)^4*(x + 1)^2*(x^2 + x + 1)). [_D. S. McNeil_, Oct 20 2011]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nico Van Cleemput_, Oct 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006438 | OEIS-A198014

Number of isomorphism classes of nanocones with 3 pentagons and a nearsymmetric boundary of length n.

Conjectured g.f.: (3*x^5 + 2*x^4 + 5*x^3 + 6*x^2 + 6*x + 2)*x/((x - 1)^4*(x + 1)^2*(x^2 + x + 1)). [_D. S. McNeil_, Oct 20 2011]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nico Van Cleemput_, Oct 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006439 | OEIS-A198015

Number of isomorphism classes of nanocones with 4 pentagons and a symmetric boundary of length n.

Conjectured g.f.: x^2*(-x^11 + x^10 - x^9 + 4*x^8 + 5*x^7 + 10*x^6 + 9*x^5 + 11*x^4 + 4*x^3 + 7*x^2 + 2*x + 1) / ((x - 1)^6*(x + 1)^3*(x^2 + 1)*(x^2 + x + 1)). [_D. S. McNeil_, Oct 20 2011]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nico Van Cleemput_, Oct 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006440 | OEIS-A198016

Number of isomorphism classes of nanocones with 4 pentagons and a nearsymmetric boundary of length n.

Conjectured g.f.: x^2*(-2*x^8 + 5*x^7 + x^6 + 13*x^5 + 8*x^4 + 13*x^3 + 9*x^2 + 3*x + 2)/(x^12 - 3*x^11 + 2*x^10 + x^9 - x^8 + 2*x^7 - 4*x^6 + 2*x^5 - x^4 + x^3 + 2*x^2 - 3*x + 1). [_D. S. McNeil_, Oct 20 2011]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nico Van Cleemput_, Oct 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006441 | OEIS-A198060

Array read by antidiagonals, m>=0, n>=0, A(m,n) = Sum_{k=0..n} Sum_{j=0..m} Sum_{i=0..m} (-1)^(j+i)*C(i,j)*C(n,k)^(m+1)*(n+1)^j*(k+1)^(-j).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Nov 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006442 | OEIS-A198195

a(n) is the smallest prime(m) such that the interval (prime(m)*n, prime(m+1)*n) contains exactly five primes.

Conjecture. In the supposition that there are infinitely many twin primes, every term beginning with the 20th is 2 or in A001359 (lesser of twin primes). The sequence is unbounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006443 | OEIS-A198245

Euler primes: Primes p that divide E(p - 3), where E(k) is the k-th Euler number.

Mestrovic (2012) computes that only three primes < 10^7 are in the sequence, but he conjectures that the sequence is infinite. - _Jonathan Sondow_, Dec 18 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Romeo Mestrovic_, Oct 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006444 | OEIS-A198256

Row sums of A197653.

Conjecture 1: a(n) = Sum_{k = 0..n} binomial(n+1,k)^2*binomial(n,k)^2. Conjecture 2: working with offset 1, that is, a(1) = 1, a(2) = 5, ..., then the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) holds for positive integers n and r and all primes p >= 5. (End) Peter Bala's conjecture 1 can equivalently written: a(n) = hypergeom([-n - 1, -n - 1, -n, -n], [1, 1, 1], 1). - _Detlef Meya_, May 28 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Susanne Wienand_, Oct 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006445 | OEIS-A198258

Row sums of A197655.

Conjecture: working with offset 1, that is, a(1) = 1, a(2) = 7, ..., then the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) holds for positive integers n and r and all primes p >= 3. - _Peter Bala_, Mar 21 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Susanne Wienand_, Oct 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006446 | OEIS-A198287

Number of 2n X 2 0..3 arrays with values 0..3 introduced in row major order and each element equal to exactly one horizontal and vertical neighbor.

Conjectures from _Colin Barker_, Mar 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006447 | OEIS-A198319

a(n) is the smallest prime(m) such that the interval (prime(m)*n, prime(m+1)*n) contains exactly six primes.

Conjecture. In the supposition that there are infinitely many twin primes, every term beginning with the 12th is 2 or in A001359 (lesser of twin primes).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006448 | OEIS-A198405

Number of 2n X 2 0..4 arrays with values 0..4 introduced in row major order and each element equal to exactly one horizontal and vertical neighbor.

Conjectures from _Colin Barker_, Mar 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006449 | OEIS-A198443

Conjectured record minima of the positive distance d between the square of an integer y and the fifth power of a positive integer x such that d = y^2 - x^5 (x <> k^2 and y <> k^5).

Conjectured record minima of the positive distance d between the square of an integer y and the fifth power of a positive integer x such that d = y^2 - x^5 (x <> k^2 and y <> k^5). Conjecture: For any positive number x >= A198444(n), the distance d between the square of an integer y and the fifth power of a positive integer x (such that x <> k^2 and y <> k^5) can't be less than A198443(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Oct 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006450 | OEIS-A198444

Values x for record minima of the positive distance d between the square of an integer y and the fifth power of a positive integer x such that d = y^2 - x^5 (x <> k^2 and y <> k^5).

Conjecture: For any positive number x >= A198444(n), the distance d between the square of an integer y and the fifth power of x (such that x <> k^2 and y <> k^5) can't be less than A198443(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Oct 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006451 | OEIS-A198447

Number of 2n X 2 0..2 arrays with values 0..2 introduced in row major order and each element unequal to exactly two horizontal and vertical neighbors.

Conjectures from _Colin Barker_, May 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006452 | OEIS-A198469

a(n) is the smallest prime(m) such that the interval (prime(m)*n, prime(m+1)*n) contains exactly seven primes.

Conjecture. In the supposition that there are infinitely many twin primes, every term beginning with the 15th is 2 or in A001359 (lesser of twin primes). A generalization. Consider sequence A_r: "A_r(n) is the smallest prime(m) such that the interval (prime(m)*n, prime(m+1)*n) contains exactly r>=0 primes". Then the sequence is unbounded and, in the supposition that there are infinitely many twin primes, we conjecture that there exists number N=N(r)>=2, such that for n>=N every term A_r(n) is 2 or in A001359.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006453 | OEIS-A198474

Number of 2n X 2 0..2 arrays with values 0..2 introduced in row major order and each element equal to an odd number of horizontal and vertical neighbors.

Conjectures from _Colin Barker_, Mar 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006454 | OEIS-A198631

Numerators of the rational sequence with e.g.f. 1/(1+exp(-x)).

Conjecture: r(n) = Sum_{k=0..n} A001147(k) * A039755(n, k) * (-1)^k / (k+1) where r(n) = a(n) / A006519(n+1) = (n!) * ([x^n] (2 / (1 + exp(-x)))), for n >= 0. - _Werner Schulte_, Feb 16 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 31 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006455 | OEIS-A198632

Triangle version of the array of the number of closed paths of even length on the graph P_n (n vertices, n-1 edges).

Conjecture: For the array w(n,2*k) in the example below, w(2*q,2*k)/2 = A185095(q,k), q >= 1, k >= 0. - _L. Edson Jeffery_, Nov 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Nov 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006456 | OEIS-A198636

One half of total number of round trips, each of length 2n, on the graph P_6 (o-o-o-o-o-o).

Conjecture: a(n) = 2^(2*n)*(sum_{k=1,2,3} (cos(k*Pi/7))^(2*n)). - _L. Edson Jeffery_, Jan 21 2012 (in fact this conjecture was recently proved in [Barbero, et al.]) Let r,s,t be the roots of x^3 + x^2 - 2x - 1; then apparently a(n) = r^(2n) + s^(2n) + t^(2n). - _James R. Buddenhagen_, Nov 03 2013 [This is equivalent to the conjecture by _L. Edson Jeffery_.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Nov 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006457 | OEIS-A198638

Number of 2n X 2 0..2 arrays with values 0..2 introduced in row major order and each element equal to an even number of horizontal and vertical neighbors.

Conjectures from _Colin Barker_, May 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006458 | OEIS-A198717

Number of n X 2 0..6 arrays with values 0..6 introduced in row major order and no element equal to any horizontal or vertical neighbor.

Conjectures from _Colin Barker_, Feb 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006459 | OEIS-A198900

Number of n X 2 0..4 arrays with values 0..4 introduced in row major order and no element equal to any horizontal or vertical neighbor.

Conjectures from _Colin Barker_, Feb 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 31 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006460 | OEIS-A198908

Number of n X 2 0..7 arrays with values 0..7 introduced in row major order and no element equal to any horizontal or vertical neighbor.

Conjectures from _Colin Barker_, Feb 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 31 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006461 | OEIS-A198956

q-expansion of modular form psi_0^4/t_{3B}.

Conjecture: Multiplicative with a(3^e) = 3^e, a(p^e) = sigma_3(p^e) for prime p <> 3. - _Andrew Howroyd_, Aug 08 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006462 | OEIS-A198976

Number of n X 2 0..5 arrays with values 0..5 introduced in row major order and no element equal to any horizontal or vertical neighbor.

Conjectures from _Colin Barker_, Feb 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 01 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006463 | OEIS-A199085

Number of distinct values taken by 3rd derivative of x^x^...^x (with n x's and parentheses inserted in all possible ways) at x=1.

Conjectured g.f.: (x^6-2*x^4-x^2+x-1)*x/(x^5-2*x^4+x^3-x^2+2*x-1). - _Alois P. Heinz_, Nov 02 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Reshetnikov_, Nov 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006464 | OEIS-A199126

Number of nX1 0..2 arrays with values 0..2 introduced in row major order, the number of instances of each value within one of each other, and no element equal to any horizontal or vertical neighbor.

Conjecture: a(3n) = A190917(n). - _R. J. Mathar_, Nov 01 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006465 | OEIS-A199127

Number of n X 2 0..2 arrays with values 0..2 introduced in row major order, the number of instances of each value within one of each other, and no element equal to any horizontal or vertical neighbor.

Conjecture: a(3n+2) = a(3n+3) = A208881(n+1). - _R. J. Mathar_, Nov 01 2015 Conjecture: -(458*n-1205) *(n+2) *(n+1)*a(n) +(-208*n^3+2578*n^2-4613*n-2410) *a(n-1) +9*(-339*n-638) *a(n-2) +27*(n-2) *(458*n^2-289*n-1146) *a(n-3) +54*(n-2) *(n-3) *(104*n-1081) *a(n-4)=0. - _R. J. Mathar_, Nov 01 2015 Conjecture: (n+2)*(n+1)*a(n) +(5*n^2-2)*a(n-1) +3*(5*n^2-15*n+3) *a(n-2) +3*(n^2 -60*n +81)*a(n-3) +135*(-n^2+3*n-1)*a(n-4) -405*(n-2)*(n-4) *a(n-5) -810*(n-4) *(n-5) *a(n-6)=0. - _R. J. Mathar_, Nov 01 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006466 | OEIS-A199203

Decimal expansion of f(0) where f is the functional square root (half-iterate) of exponent, f(f(x))=exp(x).

The listed digits are a conjecture obtained as a common result of several different numeric approximation algorithms, they haven't been rigorously proved to be correct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Reshetnikov_, Nov 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006467 | OEIS-A199331

Number of ordered ways of writing n as the sum of two semiprimes.

Conjecture: Only the integers 1, 2, 3, 4, 5, 6, 7, 9, 11, 17, 22, 33 (A072966) cannot be partitioned into a set of two semiprimes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Nov 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006468 | OEIS-A199531

Number of -n..n arrays x(0..3) of 4 elements with zero sum and no two consecutive zero elements.

Conjectures from _Colin Barker_, May 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006469 | OEIS-A199532

Number of -n..n arrays x(0..4) of 5 elements with zero sum and no two consecutive zero elements.

Conjectures from _Colin Barker_, May 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006470 | OEIS-A199533

Number of -n..n arrays x(0..5) of 6 elements with zero sum and no two consecutive zero elements.

Conjectures from _Colin Barker_, May 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006471 | OEIS-A199534

Number of -n..n arrays x(0..6) of 7 elements with zero sum and no two consecutive zero elements.

Conjectures from _Colin Barker_, May 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006472 | OEIS-A199550

Decimal expansion of the positive root of x^x^x = 2.

As follows from Gelfond's theorem, the root is irrational, so this sequence is infinite and aperiodic. Its transcendence is, apparently, still an open problem. - _Vladimir Reshetnikov_, Apr 27 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Reshetnikov_, Nov 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006473 | OEIS-A199578

Row sums of coefficient triangle of the monic associated Laguerre polynomials of order 1.

The conjecture on the alternating row sums has been proved by _Wolfdieter Lang_, Dec 12 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Nov 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006474 | OEIS-A199636

Irregular rows of odd numbers that produce n even numbers in their Collatz iteration.

It is conjectured that every odd number greater than 1 eventually appears in this sequence. The smallest and largest terms in row n are A199637(n) and A199638(n). The number of terms in row n is A131450(n) for n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Nov 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006475 | OEIS-A199705

Number of -n..n arrays x(0..2) of 3 elements with zero sum and no two neighbors equal.

Conjectures from _Colin Barker_, May 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 09 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006476 | OEIS-A199812

Number of distinct values taken by w^w^...^w (with n w's and parentheses inserted in all possible ways) where w is the first transfinite ordinal omega.

Conjectured extension of this sequence is given by A255170. Conjecture: a(n) ~ c * d^n * n^(-3/2), where c = 0.664861... and d = A051491 = 2.955765... - _Vladimir Reshetnikov_, Aug 11 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Reshetnikov_, Nov 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006477 | OEIS-A199836

Number of -n..n arrays x(0..6) of 7 elements with zero sum and no two neighbors summing to zero.

Conjectures from _Colin Barker_, May 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006478 | OEIS-A199837

Number of -n..n arrays x(0..7) of 8 elements with zero sum and no two neighbors summing to zero.

Conjectures from _Colin Barker_, May 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006479 | OEIS-A199838

Number of -n..n arrays x(0..8) of 9 elements with zero sum and no two neighbors summing to zero.

Conjectures from _Colin Barker_, Mar 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006480 | OEIS-A199899

Number of -n..n arrays x(0..3) of 4 elements with zero sum, and adjacent elements not both strictly positive and not both strictly negative.

Conjectures from _Colin Barker_, Feb 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006481 | OEIS-A199900

Number of -n..n arrays x(0..4) of 5 elements with zero sum, and adjacent elements not both strictly positive and not both strictly negative.

Conjectures from _Colin Barker_, May 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006482 | OEIS-A199901

Number of -n..n arrays x(0..5) of 6 elements with zero sum, and adjacent elements not both strictly positive and not both strictly negative.

Conjectures from _Colin Barker_, May 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006483 | OEIS-A199902

Number of -n..n arrays x(0..6) of 7 elements with zero sum, and adjacent elements not both strictly positive and not both strictly negative.

Conjectures from _Colin Barker_, May 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006484 | OEIS-A199920

Number of ways to write n = p+k with p, p+6, 6k-1 and 6k+1 all prime.

Conjecture: a(n)>0 for all n>11. This implies that there are infinitely many twin primes and also infinitely many sexy primes. It has been verified for n up to 10^9. See also A199800 for a weaker version of this conjecture. Zhi-Wei Sun also conjectured that any integer n>6 not equal to 319 can be written as p+k with p, p+6, 3k-2+(n mod 2) and 3k+2-(n mod 2) all prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006485 | OEIS-A199944

Number of -n..n arrays x(0..2) of 3 elements with zeroth through 2nd differences all nonzero.

Conjectures from _Colin Barker_, May 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 12 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

B05 A200000 - A249999 | 2,749 条

V3-006486 | OEIS-A200039

Number of -n..n arrays x(0..2) of 3 elements with sum zero and with zeroth through 2nd differences all nonzero.

Conjectures from _Colin Barker_, May 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 12 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006487 | OEIS-A200139

Triangle T(n,k), read by rows, given by (1,1,0,0,0,0,0,0,0,0,...) DELTA (1,0,0,0,0,0,0,0,0,0,...) where DELTA is the operator defined in A084938.

Unsigned A118800, so my conjecture above is true. - _Tom Copeland_, Nov 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Nov 13 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006488 | OEIS-A200166

Number of -n..n arrays x(0..2) of 3 elements with nonzero sum and with zero through 2 differences all nonzero.

Conjectures from _Colin Barker_, May 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 13 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006489 | OEIS-A200216

Danilov's sequence of x satisfying 0 < |x^3-y^2| < sqrt(x) with integer (x,y).

Conjecture: a(n) = 3461450947505*a(n-1) + 6440022564931157490*a(n-2) - 6440022564931157490*a(n-3) - 3461450947505*a(n-4) + a(n-5). - _R. J. Mathar_, Nov 15 2011 Conjecture: g.f. 4092*(1-z)/(5*(1+1860498*z+z^2)) - 7/(10*(z-1)) + 930249*(1-z)/(10*(1-3461452808002*z+z^2)). - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Nov 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006490 | OEIS-A200217

Danilov's sequence of y values satisfying 0 < |x^3 - y^2| < sqrt(x) with integer (x,y).

Conjecture: a(n) = +6440022564929296994*a(n-1) +22291834190970757443015664937985*a(n-2) -41473935220466903245533179036528718020*a(n-3) +22291834190970757443015664937985*a(n-4) +6440022564929296994*a(n-5) -a(n-6). - _R. J. Mathar_, Nov 15 2011 Equivalent conjecture g.f.: -61*(z-1) * (471281*z^4 -396480201682498803123 76*z^3 -41789857533031766983147634306 7314*z^2 -396480201682498803123 76*z +471281) / ((z^2+3461452808002*z+1) *(z^2-6440026026380244498*z+1) *(z^2-1860498*z+1)). - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Nov 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006491 | OEIS-A200218

The differences x^3 - y^2 of Danilov's subsequence of good Hall's examples A078933.

Conjecture: a(n) = -1860497 * a(n-1) + 1860497 * a(n-2) + a(n-3) with g.f. 297 * z * (1 + 14882 * z + z^2) / ((z-1)*(z^2 + 1860498 * z+1)). - _R. J. Mathar_, Nov 15 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Nov 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006492 | OEIS-A200249

Number of 0..5 arrays x(0..n-1) of n elements with each no smaller than the sum of its previous elements modulo 6.

Conjectures from _Colin Barker_, May 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 15 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006493 | OEIS-A200312

a(n) = A000108(n)*A006130(n), where A000108 is the Catalan numbers and A006130(n) = A006130(n-1) + 3*A006130(n-2).

Conjecture: n*(n+1)*a(n) -2*n*(2*n-1)*a(n-1) -12*(2*n-1)*(2*n-3)*a(n-2) = 0. - _R. J. Mathar_, Nov 17 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006494 | OEIS-A200431

Number of -n..n arrays x(0..3) of 4 elements with zero sum and no two or three adjacent elements summing to zero.

Conjectures from _Colin Barker_, May 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006495 | OEIS-A200432

Number of -n..n arrays x(0..4) of 5 elements with zero sum and no two or three adjacent elements summing to zero.

Conjectures from _Colin Barker_, May 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006496 | OEIS-A200433

Number of -n..n arrays x(0..5) of 6 elements with zero sum and no two or three adjacent elements summing to zero.

Conjectures from _Colin Barker_, May 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006497 | OEIS-A200434

Number of -n..n arrays x(0..6) of 7 elements with zero sum and no two or three adjacent elements summing to zero.

Conjectures from _Colin Barker_, May 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006498 | OEIS-A200435

Number of -n..n arrays x(0..7) of 8 elements with zero sum and no two or three adjacent elements summing to zero.

Conjectures from _Colin Barker_, May 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006499 | OEIS-A200503

Record (maximal) gaps between prime sextuplets (p, p+4, p+6, p+10, p+12, p+16).

Prime sextuplets (p, p+4, p+6, p+10, p+12, p+16) are densest permissible constellations of 6 primes. Average gaps between sextuplets (and, more generally, between prime k-tuples) can be deduced from the Hardy-Littlewood k-tuple conjecture and are O(log^6(p)). Maximal gaps may be significantly larger than average gaps; this sequence suggests that maximal gaps are O(log^7(p)). (1) Conjectured upper bound: gaps between prime sextuplets (p, p+4, p+6, p+10, p+12, p+16) are smaller than 0.058*(log p)^7, where p is the prime at the end of the gap. (2) Estimate for the actual size of the maximal gap that ends at p: maximal gap = a(log(p/a)-1/3), where a = 0.058*(log p)^6 is the average gap between sextuplets near p, as predicted by the Hardy-Littlewood k-tuple conjecture. Formulas (1) and (2) are asymptotically equal as p tends to infinity. However, (1) yields values greater than all known gaps, while (2) yields "good guesses" that may be either above or below the actual size of maximal gaps. Both formulas (1) and (2) are derived from the Hardy-Littlewood k-tuple conjecture via probability-based heuristics relating the expected maximal gap size to the average gap. Neither of the formulas has a rigorous proof (the k-tuple conjecture itself has no formal proof either). In both formulas, the constant ~0.058 is reciprocal to the Hardy-Littlewood 6-tuple constant 17.2986...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Nov 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006500 | OEIS-A200526

a(n) = gcd(t(n), t(3n-1)), where t = A200217.

Successive maxima of the GCD in A200217 occur between A200217(n) and A200217(3n-1) terms. Conjecture: All terms have same set of prime divisors, that can be used to complete prime factorization of every term in this sequence by the GCD method. All prime divisors with exception 19 are of the form 4k+1. The integer 19 divides a(3n+1) for n=0,1,2,3,...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Nov 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006501 | OEIS-A200652

a(n) = least k with -3<k<2*n such that n*(n+1)+k and n*(n+1)+k+2 are twin primes, or 0 if no such k exists.

Only 11 values are zero for n < 434. Conjecture: no more 0 values if n>433.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Nov 20 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006502 | OEIS-A200753

G.f. satisfies: A(x) = 1 + (x-x^2)*A(x)^3.

Conjecture: 2n*(2n+1)*a(n) -(13n-7)(3n-2)*a(n-1) +4(29n^2-87n+67)*a(n-2) +9(-15n^2+69n-80)*a(n-3) +6(3n-8)(3n-10)*a(n-4)=0. - _R. J. Mathar_, Nov 22 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 21 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006503 | OEIS-A200770

Number of nX1 0..2 arrays with every row and column running average nondecreasing rightwards and downwards, and the number of instances of each value within one of each other.

Conjecture: (n+2)*(2770*n-4893)*a(n) + 3*(1090*n^2-4919*n+1654)*a(n-1) + 4686*a(n-2) + 2*(-5540*n^2+7016*n+17127)*a(n-3) - 6*(1090*n-2739)*(2*n-7)*a(n-4) = 0. - _R. J. Mathar_, Nov 23 2011 [corrected by _Jason Yuen_, Oct 14 2025]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006504 | OEIS-A200785

T(n,k) is the number of arrays of n+2 elements from {0,1,...,k} with no two consecutive ascents.

All the conjectured formulas are true, and follow from the Burstein-Mansour paper. - _N. J. A. Sloane_, May 21 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006505 | OEIS-A200786

Number of 0..n arrays x(0..3) of 4 elements without any two consecutive increases.

Conjectures from _Colin Barker_, Oct 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006506 | OEIS-A200787

Number of 0..n arrays x(0..4) of 5 elements without any two consecutive increases.

Conjectures from _Colin Barker_, Oct 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006507 | OEIS-A200788

Number of 0..n arrays x(0..5) of 6 elements without any two consecutive increases.

Conjectures from _Colin Barker_, Oct 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006508 | OEIS-A200789

Number of 0..n arrays x(0..6) of 7 elements without any two consecutive increases.

Conjectures from _Colin Barker_, Oct 15 2017: (Start) The formulas below are consistent with the conjectured formula above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006509 | OEIS-A200790

Number of 0..n arrays x(0..7) of 8 elements without any two consecutive increases.

Conjectures from _Colin Barker_, Oct 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006510 | OEIS-A200791

Number of 0..n arrays x(0..8) of 9 elements without any two consecutive increases.

Conjectures from _Colin Barker_, Oct 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006511 | OEIS-A200839

Number of 0..n arrays x(0..3) of 4 elements without any two consecutive increases or two consecutive decreases.

Conjectures from _Colin Barker_, Oct 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006512 | OEIS-A200840

Number of 0..n arrays x(0..4) of 5 elements without any two consecutive increases or two consecutive decreases.

Conjectures from _Colin Barker_, Oct 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006513 | OEIS-A200841

Number of 0..n arrays x(0..5) of 6 elements without any two consecutive increases or two consecutive decreases.

Conjectures from _Colin Barker_, Oct 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006514 | OEIS-A200842

Number of 0..n arrays x(0..6) of 7 elements without any two consecutive increases or two consecutive decreases.

Conjectures from _Colin Barker_, Oct 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006515 | OEIS-A200843

Number of 0..n arrays x(0..7) of 8 elements without any two consecutive increases or two consecutive decreases.

Conjectures from _Colin Barker_, Oct 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006516 | OEIS-A200844

Number of 0..n arrays x(0..8) of 9 elements without any two consecutive increases or two consecutive decreases.

Conjectures from _Colin Barker_, Oct 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006517 | OEIS-A200872

Number of 0..n arrays x(0..3) of 4 elements without any interior element greater than both neighbors or less than both neighbors.

Conjectures from _Colin Barker_, Oct 16 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006518 | OEIS-A200873

Number of 0..n arrays x(0..4) of 5 elements without any interior element greater than both neighbors or less than both neighbors.

Conjectures from _Colin Barker_, Oct 16 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006519 | OEIS-A200874

Number of 0..n arrays x(0..5) of 6 elements without any interior element greater than both neighbors or less than both neighbors.

Conjectures from _Colin Barker_, Oct 16 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006520 | OEIS-A200875

Number of 0..n arrays x(0..6) of 7 elements without any interior element greater than both neighbors or less than both neighbors.

Conjectures from _Colin Barker_, Oct 16 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006521 | OEIS-A200876

Number of 0..n arrays x(0..7) of 8 elements without any interior element greater than both neighbors or less than both neighbors.

Conjectures from _Colin Barker_, Oct 16 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006522 | OEIS-A200877

Number of 0..n arrays x(0..8) of 9 elements without any interior element greater than both neighbors or less than both neighbors.

Conjectures from _Colin Barker_, Oct 16 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006523 | OEIS-A200887

Number of 0..n arrays x(0..3) of 4 elements without any interior element greater than both neighbors.

Conjectures from _Colin Barker_, Oct 16 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006524 | OEIS-A200888

Number of 0..n arrays x(0..4) of 5 elements without any interior element greater than both neighbors.

Conjectures from _Colin Barker_, Oct 16 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006525 | OEIS-A200889

Number of 0..n arrays x(0..5) of 6 elements without any interior element greater than both neighbors.

Conjectures from _Colin Barker_, Oct 16 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006526 | OEIS-A200890

Number of 0..n arrays x(0..6) of 7 elements without any interior element greater than both neighbors.

Conjectures from _Colin Barker_, Oct 16 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006527 | OEIS-A200891

Number of 0..n arrays x(0..7) of 8 elements without any interior element greater than both neighbors.

Conjectures from _Colin Barker_, Oct 16 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006528 | OEIS-A200892

Number of 0..n arrays x(0..8) of 9 elements without any interior element greater than both neighbors.

Conjectures from _Colin Barker_, Oct 16 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006529 | OEIS-A200918

Successive prime factors of (3^1006003 - 3)/1006003^2.

Conjecture (*Artur Jasinski*): If another infinite sequences with good Hall's examples occurred, it would have to contain primes from this sequence as constant divisors of the whole sequence, because parts of Danilov's infinite sequence (A200216, A200217, A200218) contain divisors of (3^A014127(1) - 3)/(A014127(1)^2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Nov 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006530 | OEIS-A200982

Numbers k such that 1^(lambda(k)/2) + 2^(lambda(k)/2) + ... + (k - 1)^(lambda(k)/2) = 0 (mod k), where lambda is the Carmichael reduced totient function.

Conjecture: The asymptotic density is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Nov 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006531 | OEIS-A201043

Number of -n..n arrays of 4 elements with adjacent element differences also in -n..n.

Conjectures from _Colin Barker_, May 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006532 | OEIS-A201044

Number of -n..n arrays of 5 elements with adjacent element differences also in -n..n.

Conjectures from _Colin Barker_, May 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006533 | OEIS-A201045

Number of -n..n arrays of 6 elements with adjacent element differences also in -n..n.

Conjectures from _Colin Barker_, May 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006534 | OEIS-A201046

Number of -n..n arrays of 7 elements with adjacent element differences also in -n..n.

Conjectures from _Colin Barker_, May 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006535 | OEIS-A201051

Record (maximal) gaps between prime septuplets (p, p+2, p+6, p+8, p+12, p+18, p+20).

Prime septuplets (p, p+2, p+6, p+8, p+12, p+18, p+20) are one of the two types of densest permissible constellations of 7 primes (A022009 and A022010). Average gaps between prime k-tuples can be deduced from the Hardy-Littlewood k-tuple conjecture and are O(log^k(p)), with k=7 for septuplets. If a gap is larger than any preceding gap, we call it a maximal gap, or a record gap. Maximal gaps may be significantly larger than average gaps; this sequence suggests that maximal gaps are O(log^8(p)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Nov 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006536 | OEIS-A201052

a(n) is the maximal number c of integers that can be chosen from {1,2,...,n} so that all 2^c subsets have distinct sums.

Conjecture (confirmed through k=7): a(n)=k for all n in the interval A005318(k) <= n < A005318(k+1). - _Jon E. Schoenfield_, Nov 28 2013 [Note: This conjecture is false; see A276661 for a counterexample (n=34808712605260918463) in which n is in the interval A005318(66) <= n < A005318(67), yet a(n)=67, not 66. - _Jon E. Schoenfield_, Nov 05 2016]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006537 | OEIS-A201060

Repdigit numbers with repdigit digital product.

Conjecture: a(n) is repunit for n >= 16.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006538 | OEIS-A201062

Record (maximal) gaps between prime 5-tuples (p, p+4, p+6, p+10, p+12).

Prime quintuplets (p, p+4, p+6, p+10, p+12) are one of the two types of densest permissible constellations of 5 primes (A022006 and A022007). Average gaps between prime k-tuples can be deduced from the Hardy-Littlewood k-tuple conjecture and are O(log^k(p)), with k=5 for quintuplets. If a gap is larger than all preceding gaps, we call it a maximal gap, or a record gap. Maximal gaps may be significantly larger than average gaps; this sequence suggests that maximal gaps between quintuplets are O(log^6(p)). (2) Estimate for the actual size of the maximal gap that ends at p: maximal gap ~ a(log(p/a)-0.4), where a = 0.0987*(log p)^5 is the average gap between quintuplets near p, as predicted by the Hardy-Littlewood k-tuple conjecture. Both formulas (1) and (2) are derived from the Hardy-Littlewood k-tuple conjecture via probability-based heuristics relating the expected maximal gap size to the average gap. Neither of the formulas has a rigorous proof (the k-tuple conjecture itself has no formal proof either). In both formulas, the constant ~0.0987 is reciprocal to the Hardy-Littlewood 5-tuple constant 10.1317...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Nov 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006539 | OEIS-A201073

Record (maximal) gaps between prime 5-tuples (p, p+2, p+6, p+8, p+12).

Prime quintuplets (p, p+2, p+6, p+8, p+12) are one of the two types of densest permissible constellations of 5 primes (A022006 and A022007). Average gaps between prime k-tuples can be deduced from the Hardy-Littlewood k-tuple conjecture and are O(log^k(p)), with k=5 for quintuplets. If a gap is larger than any preceding gap, we call it a maximal gap, or a record gap. Maximal gaps may be significantly larger than average gaps; this sequence suggests that maximal gaps are O(log^6(p)). (2) Estimate for the actual size of the maximal gap that ends at p: maximal gap ~ a(log(p/a)-0.4), where a = 0.0987*(log p)^5 is the average gap between quintuplets near p, as predicted by the Hardy-Littlewood k-tuple conjecture. Both formulas (1) and (2) are derived from the Hardy-Littlewood k-tuple conjecture via probability-based heuristics relating the expected maximal gap size to the average gap. Neither of the formulas has a rigorous proof (the k-tuple conjecture itself has no formal proof either). In both formulas, the constant ~0.0987 is reciprocal to the Hardy-Littlewood 5-tuple constant 10.1317...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Nov 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006540 | OEIS-A201089

Number of -n..n arrays of 4 elements with first and second differences also in -n..n.

Conjectures from _Colin Barker_, Feb 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006541 | OEIS-A201202

Row sums of triangle A201201: first associated monic Laguerre polynomials with parameter alpha=1 evaluated at x=1.

This e.g.f. satisfies the homogeneous second-order differential equation: (1+x)^2*(d^2/dx^2)g(x) + (5+6*x)*(d/dx)g(x) + 6*g(x) = 0, with g(0)=1 and (d/dx)g(x)|_{x=0} = -3. This is equivalent to the recurrence conjectured by _R. J. Mathar_, which is thus proved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Dec 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006542 | OEIS-A201203

Alternating row sums of triangle A201201: first associated monic Laguerre-Sonin(e) polynomials with parameter alpha=1 evaluated at x=-1.

This e.g.f. satisfies the homogeneous ordinary second-order differential equation (1+x)^2*(d^2(g(x))/dx^2) + (7+6*x)*(d(g(x))/dx)+6*g(x), with g(0)=1 and (d(g(x))/dx)_{x=0} = -5. This is equivalent to the recurrence conjectured above by _R. J. Mathar_, which is thus proved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Dec 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006543 | OEIS-A201227

a(n) = (A201225(n))^3 - (A201226(n))^2.

Conjecture: No more infinite families of solutions Mordell curves with extension sqrt(5) than A201227 and A200218.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Nov 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006544 | OEIS-A201251

Record (maximal) gaps between prime septuplets (p, p+2, p+8, p+12, p+14, p+18, p+20).

Prime septuplets (p, p+2, p+8, p+12, p+14, p+18, p+20) are one of the two types of densest permissible constellations of 7 primes (A022009 and A022010). Average gaps between prime k-tuples can be deduced from the Hardy-Littlewood k-tuple conjecture and are O(log^k(p)), with k=7 for septuplets. If a gap is larger than any preceding gap, we call it a maximal gap, or a record gap. Maximal gaps may be significantly larger than average gaps; this sequence suggests that maximal gaps are O(log^8(p)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Nov 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006545 | OEIS-A201267

a(1) = 1 and a(n) is the least integer such that the continued fraction for 1/a(1) + 1/a(2) + ... + 1/a(n) contains exactly n elements.

Conjecture: limsup_{n->infinity} a(n) = infinity. More precisely we claim that log(a(n))/log(n) is bounded and does not converge to zero (see related link). Does a(n) = 2 infinitely many times or does it exist M >= 2 such that a(n) <= M infinitely many times?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jan 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006546 | OEIS-A201268

Distances d=x^3-y^2 for primary extremal points {x,y} of Mordell elliptic curves with quadratic extensions over rationals.

Conjecture (*Artur Jasinski*): Conjecture (*Artur Jasinski*): Extremal points are k-th successive points with maximal coordinates x.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Nov 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006547 | OEIS-A201347

Number of n X 2 0..1 arrays with rows and columns lexicographically nondecreasing and every element equal to at least one horizontal or vertical neighbor.

Conjectures from _Colin Barker_, May 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006548 | OEIS-A201348

Number of n X 3 0..1 arrays with rows and columns lexicographically nondecreasing and every element equal to at least one horizontal or vertical neighbor.

Conjectures from _Colin Barker_, May 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006549 | OEIS-A201371

Number of n X 4 0..1 arrays with rows and columns lexicographically nondecreasing read forwards, and nonincreasing read backwards.

Conjectures from _Colin Barker_, May 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006550 | OEIS-A201372

Number of n X 5 0..1 arrays with rows and columns lexicographically nondecreasing read forwards, and nonincreasing read backwards.

Conjectures from _Colin Barker_, May 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006551 | OEIS-A201373

Number of n X 6 0..1 arrays with rows and columns lexicographically nondecreasing read forwards, and nonincreasing read backwards.

Conjectures from _Colin Barker_, May 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006552 | OEIS-A201488

Decimal expansion of maximal quantum success probability of the CHSH game.

Conjecture: Equals Sum_{k>=0} A210066(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Dec 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006553 | OEIS-A201533

Number of n X 2 0..2 arrays with rows and columns lexicographically nondecreasing and every element equal to at least one horizontal or vertical neighbor.

Conjectures from _Colin Barker_, May 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006554 | OEIS-A201553

Number of arrays of 6 integers in -n..n with sum zero.

Conjectures from _Colin Barker_, May 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006555 | OEIS-A201554

Number of arrays of 7 integers in -n..n with sum zero.

Conjectures from _Colin Barker_, May 23 2018: (Start) Barker conjectures confirmed using technique similar to A201553.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 02 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006556 | OEIS-A201560

a(n) = (Sum(m^(n-1), m=1..n-1) + 1) modulo n.

Equals 0 if n is 1 or a prime, by Fermat's little theorem. It is conjectured that the converse is also true; see A055030, A055032, A204187 and note that a(n) = 0 <==> A055032(n) = 1 <==> A204187(n) = n-1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Jan 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006557 | OEIS-A201596

Record (maximal) gaps between prime triples (p, p+4, p+6).

Prime triples (p, p+4, p+6) are one of the two types of densest permissible constellations of 3 primes (A022004 and A022005). By the Hardy-Littlewood k-tuple conjecture, average gaps between prime k-tuples are O(log^k(p)), with k=3 for triples. If a gap is larger than any preceding gap, we call it a maximal gap, or a record gap. Maximal gaps may be significantly larger than average gaps; this sequence suggests that maximal gaps between triples are O(log^4(p)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Dec 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006558 | OEIS-A201598

Record (maximal) gaps between prime triples (p, p+2, p+6).

Prime triples (p, p+2, p+6) are one of the two types of densest permissible constellations of 3 primes (A022004 and A022005). By the Hardy-Littlewood k-tuple conjecture, average gaps between prime k-tuples are O(log^k(p)), with k=3 for triples. If a gap is larger than any preceding gap, we call it a maximal gap, or a record gap. Maximal gaps may be significantly larger than average gaps; this sequence suggests that maximal gaps between triples are O(log^4(p)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Dec 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006559 | OEIS-A201618

Number of n X 1 0..3 arrays with rows and columns lexicographically nondecreasing and every element equal to at least one horizontal or vertical neighbor.

Conjectures from _Colin Barker_, May 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006560 | OEIS-A201631

a(n) is the number of Fibonacci meanders of length m*n and central angle 360/m degrees where m = 2.

The above conjecture was proved by Baril et al., which also give a formal definition of the Fibonacci meanders and describe a bijection with a certain class of peakless grand Motzkin paths of length n. - _Peter Luschny_, Mar 16 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jan 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006561 | OEIS-A201686

a(n) = binomial(n, [n/2]) - 2.

Conjecture: +(n+1)*a(n) +2*(-n-1)*a(n-1) +(-3*n+7)*a(n-2) +2*(4*n-9)*a(n-3) +4*(-n+3)*a(n-4)=0. - _R. J. Mathar_, Jul 17 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006562 | OEIS-A201695

Number of n X 3 0..2 arrays with rows and columns lexicographically nondecreasing read forwards and nonincreasing read backwards.

Conjectures from _Colin Barker_, May 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006563 | OEIS-A201696

Number of n X 4 0..2 arrays with rows and columns lexicographically nondecreasing read forwards and nonincreasing read backwards.

Conjectures from _Colin Barker_, May 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006564 | OEIS-A201722

Number of n X 1 0..4 arrays with rows and columns lexicographically nondecreasing and no element equal to the number of horizontal and vertical neighbors equal to itself.

Conjectures from _Colin Barker_, May 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 04 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006565 | OEIS-A201812

Number of arrays of 4 integers in -n..n with sum zero and equal numbers of elements greater than zero and less than zero.

Conjectures from _Colin Barker_, May 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006566 | OEIS-A201813

Number of arrays of 5 integers in -n..n with sum zero and equal numbers of elements greater than zero and less than zero.

Conjectures from _Colin Barker_, May 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006567 | OEIS-A201814

Number of arrays of 6 integers in -n..n with sum zero and equal numbers of elements greater than zero and less than zero.

Conjectures from _Colin Barker_, May 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006568 | OEIS-A201815

Number of arrays of 7 integers in -n..n with sum zero and equal numbers of elements greater than zero and less than zero.

Conjectures from _Colin Barker_, May 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 05 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006569 | OEIS-A201828

The smallest A(m) such that the interval (A(m)*n, A(m+1)*n) contains exactly one element of A, where A is the sequence of primes p for which p-2 is not prime.

This is a possible model sequence to understand the role of twin primes in sequences like A195871. In particular, if after a large number N_tw, there are no twin primes, what primes will take their place in A195871? Our observations and expectations are expressed in the following conjecture. Conjecture: For n>=13, every a(n) is the lesser of a pair of cousin primes p and p+4, cf. A023200. Note that it is only conjectured that there are infinitely many pairs of cousin primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006570 | OEIS-A201950

Central coefficients in Product_{k=0..n-1} (1 + k*x + x^2).

Conjecture: a(n) also counts permutations of [n+1] in which the number of left-to-right maxima equals the number of right-to-left maxima. Verified via direct enumeration for n=0..11 and Stirling-number-based calculation for n = 0..29. - _Austen M. Fletcher_, Dec 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 06 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006571 | OEIS-A201975

Number of n X 2 0..3 arrays with rows and columns lexicographically nondecreasing read forwards and nonincreasing read backwards.

Conjectures from _Colin Barker_, May 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006572 | OEIS-A201976

Number of n X 3 0..3 arrays with rows and columns lexicographically nondecreasing read forwards and nonincreasing read backwards.

Conjectures from _Colin Barker_, May 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006573 | OEIS-A201993

Conjectured lower bound for the number of circles of radius 1 that can be packed into a circle of radius n.

Conjectured lower bound for the number of circles of radius 1 that can be packed into a circle of radius n. Bound provided by David W. Cantrell in December 2008. It is conjectured that it is possible to find packings such that A023393(n)>=a(n) for all n. Currently (December 2011) the smallest number of circles, for which the bound is not achieved, is 507.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 07 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006574 | OEIS-A202020

Number of 4-colored Motzkin paths of length n with no peaks at level 1.

Conjecture: 2(n+2)*a(n) -4*(5n+4)*a(n-1) +3*(19n-2)*a(n-2) +4*(11-14n)*a(n-4) +12*(n-1)*a(n-4)=0. - _R. J. Mathar_, Dec 18 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José Luis Ramírez Ramírez_, Dec 08 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006575 | OEIS-A202048

Number of (n+2) X 6 binary arrays avoiding patterns 001 and 110 in rows and columns.

Conjectures from _Colin Barker_, May 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006576 | OEIS-A202049

Number of (n+2) X 7 binary arrays avoiding patterns 001 and 110 in rows and columns.

Conjectures from _Colin Barker_, May 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006577 | OEIS-A202050

Number of (n+2) X 8 binary arrays avoiding patterns 001 and 110 in rows and columns.

Conjectures from _Colin Barker_, May 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006578 | OEIS-A202051

Number of (n+2) X 9 binary arrays avoiding patterns 001 and 110 in rows and columns.

Conjectures from _Colin Barker_, May 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006579 | OEIS-A202054

Smallest x such that Mordell's elliptic curve x^3-y^2=d (for positive d) has an integral point in the quadratic extension sqrt(A202057(n)).

The existence of such x for each A202057(n) is conjectural following the conjecture in A201278.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Dec 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006580 | OEIS-A202055

Smallest y such that Mordell's elliptic curve x^3-y^2=d (for positive d) has an integral point in the quadratic extension sqrt(A202057(n)).

The existence of such y for each A202057(n) is conjectural following the conjecture in A201278.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Dec 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006581 | OEIS-A202056

Smallest d such that Mordell's elliptic curve x^3-y^2=d (for positive d) has an integral point in the quadratic extension sqrt(A202057(n)).

The existence of such d for each A202057(n) is conjectural following the conjecture in A201278.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Dec 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006582 | OEIS-A202057

Numbers which are not perfect squares and such that all prime divisors are congruent to 1 or 2 mod 4.

This sequence follows conjecture from A201278 that Mordell's elliptic curve x^3-y^2 = d can contain points {x,y} with quadratic extension sqrt(k) over rationals if and only k belongs to this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Dec 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006583 | OEIS-A202063

Position of first appearance of n-th prime in A071216.

a(n) = primepi(A188815(n)) (conjectured). - _Michel Marcus_, Jul 05 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Dec 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006584 | OEIS-A202077

Number of arrays of 5 integers in -n..n with sum zero and the sum of every adjacent pair being odd.

Conjectures from _Colin Barker_, May 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 10 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006585 | OEIS-A202137

Numbers k such that 24k + 1 is neither square nor prime.

Conjecture: sequence contains arbitrarily long runs of consecutive integers. Conjecture is easy to prove using the Chinese Remainder Theorem and the fact that the gaps between squares grow. - _Robert Israel_, Jan 25 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Dec 15 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006586 | OEIS-A202146

G.f.: 1/(1-x) + Sum_{n>=1} x^n/(1-x) * Product_{k=1..n} (1 - x^k) / (1 - x^(2*k+1)).

a(k) == 1 (mod 2) iff k = 3*n*(n+1) for n>=0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 12 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006587 | OEIS-A202150

a(n) = A202146(3*n*(n+1)) for n>=0.

Conjecture: this sequence consists of all odd terms in A202146; the g.f. of A202146 is 1/(1-x) + Sum_{n>=1} x^n/(1-x) * Product_{k=1..n} (1 - x^k) / (1 - x^(2*k+1)), which by the conjecture has an odd coefficient of x^m iff m = 3*n*(n+1) for n>=0. The conjecture holds for at least the initial 30300 terms of A202146.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 12 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006588 | OEIS-A202173

Conjectured list of numbers not of the form x^3 + y^3 + 2*z^3.

Conjectured list of numbers not of the form x^3 + y^3 + 2*z^3. Hypothesis: the sequence is empty. See the Cohen book for details. - _Arkadiusz Wesolowski_, Aug 20 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_ and _Arkadiusz Wesolowski_, Dec 22 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006589 | OEIS-A202194

Number of (n+2)X(n+2) binary arrays avoiding patterns 001 and 101 in rows and columns.

Conjecture: a(n) = 4*(n+2)^2 * (2*n+1)!/(n! * (n+1)!). - _Michael Somos_, Sep 11 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006590 | OEIS-A202210

G.f.: A(x) = (Sum_{n>=0} 3^n*(2*n+1) * x^(n*(n+1)/2))^(1/3).

Conjecture: a(5*n+4) == 0 (mod 5) (checked up to n = 200). - _Peter Bala_, Feb 26 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006591 | OEIS-A202254

Number of zero-sum -n..n arrays of 4 elements with adjacent element differences also in -n..n.

Conjectures from _Colin Barker_, Mar 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 14 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006592 | OEIS-A202275

Differences between A074753 (number of integers k such that sigma(k) <= n) and A202276 (number of integers k <= n such that sigma(x) = k has no solution); sigma = A000203.

Conjectures: Max a(n) = 15 for n = 195, 403, 434. For n >= 687, a(n) < 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Dec 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006593 | OEIS-A202277

Numbers m such that number of integers k such sigma(k) <= m (A074753) is equal to number of integers k <= m such that sigma(x) = k has no solution (A202276).

Conjecture: sequence is finite with 13 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Dec 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006594 | OEIS-A202281

Record (maximal) gaps between prime decuplets (p+0,2,6,8,12,18,20,26,30,32).

Average gaps between prime k-tuples are O(log^k(p)), with k=10 for decuplets, by the Hardy-Littlewood k-tuple conjecture. If a gap is larger than any preceding gap, we call it a maximal gap, or a record gap. Maximal gaps may be significantly larger than average gaps; this sequence suggests that maximal gaps are O(log^11(p)). Both formulas (1) and (2) are derived from the Hardy-Littlewood k-tuple conjecture via probability-based heuristics relating the expected maximal gap size to the average gap. Neither of the formulas has a rigorous proof.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Dec 15 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006595 | OEIS-A202319

Lesser of two semiprimes sandwiched each between semiprimes thus forming a twin semiprime-triple.

Conjecture: For every natural number k there are infinitely many semiprime pairs sp and sp' both sandwiched between semiprimes such that sp' - sp = 4k. The number centered between the triples of a twin semiprime-triple must be divisible by 36. Let m be the middle of such a twin semiprime-triple. It is trivial that m is divisible by 4, and that it is congruent to 0, 4, or 5 (mod 9). If it were congruent to 4, then m-1 and m+2 would both be divisible by 3, hence equal to 3 times a prime. But then those two primes would differ by 1, impossible except for primes 2, 3, which can be checked separately. A similar argument eliminates the case m == 5 (mod 9), so m must be divisible by 9. Conjecture by the author, proved by _Franklin T. Adams-Watters_, Dec 18 2011.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V.J. Pohjola_, Dec 16 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006596 | OEIS-A202330

Number of (n+1) X 4 binary arrays with consecutive windows of two bits considered as a binary number nondecreasing in every row and column.

Conjectures from _Colin Barker_, May 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006597 | OEIS-A202331

Number of (n+1) X 5 binary arrays with consecutive windows of two bits considered as a binary number nondecreasing in every row and column.

Conjectures from _Colin Barker_, May 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006598 | OEIS-A202332

Number of (n+1) X 6 binary arrays with consecutive windows of two bits considered as a binary number nondecreasing in every row and column.

Conjectures from _Colin Barker_, May 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006599 | OEIS-A202333

Number of (n+1) X 7 binary arrays with consecutive windows of two bits considered as a binary number nondecreasing in every row and column.

Conjectures from _Colin Barker_, May 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006600 | OEIS-A202334

Number of (n+1) X 8 binary arrays with consecutive windows of two bits considered as a binary number nondecreasing in every row and column.

Conjectures from _Colin Barker_, May 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 17 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006601 | OEIS-A202349

Lexicographically earliest sequence such that the sequence and its first and second differences share no terms, and the 3rd differences are equal to the original sequence.

Is it a theorem that, if x,y,z = 1,3,9, the sequence has the desired properties, or is it just a conjecture? - _N. J. A. Sloane_, Jun 21 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_, Jun 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006602 | OEIS-A202367

LCM of denominators of the coefficients of polynomials Q^(2)_m(n) defined by the recursion Q^(2)_0(n)=1; for m >= 1, Q^(2)_m(n) = Sum_{i=1..n} i^2*Q^(2)_(m-1)(i).

Conjecture: a(n) = Product_{primes p} p^(Sum_{k>=0} floor((n-1)/(ceiling((p-1)/2)*p^k))). If the conjecture is true, then, for n >= 0, A007814(a(n+1)) = A007814(n!) + n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Dec 18 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006603 | OEIS-A202438

G.f.: A(x) = (Sum_{n>=0} 9^n*(2*n+1) * (-x)^(n*(n+1)/2))^(-1/3).

Conjectures: a(25*n+24) == 0 (mod 25) (checked up to n = 50) and a(7*n+5) == 0 (mod 7) (checked up to n = 200). - _Peter Bala_, Feb 26 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006604 | OEIS-A202440

Number of (n+2) X 3 binary arrays avoiding patterns 001 and 110 in rows, columns and nw-to-se diagonals.

Conjectures from _Colin Barker_, May 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006605 | OEIS-A202442

Number of (n+2) X 5 binary arrays avoiding patterns 001 and 110 in rows, columns and nw-to-se diagonals.

Conjectures from _Colin Barker_, May 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006606 | OEIS-A202443

Number of (n+2) X 6 binary arrays avoiding patterns 001 and 110 in rows, columns and nw-to-se diagonals.

Conjectures from _Colin Barker_, May 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006607 | OEIS-A202454

Number of (n+2) X (n+2) binary arrays with consecutive windows of three bits considered as a binary number nondecreasing in every row and column.

Conjecture: a(n) ~ c * 4^n / sqrt(n), where c = 95.3480396... - _Vaclav Kotesovec_, Feb 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006608 | OEIS-A202455

Number of (n+2) X 4 binary arrays with consecutive windows of three bits considered as a binary number nondecreasing in every row and column.

Conjectures from _Colin Barker_, May 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006609 | OEIS-A202456

Number of (n+2) X 5 binary arrays with consecutive windows of three bits considered as a binary number nondecreasing in every row and column.

Conjectures from _Colin Barker_, May 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006610 | OEIS-A202457

Number of (n+2) X 6 binary arrays with consecutive windows of three bits considered as a binary number nondecreasing in every row and column.

Conjectures from _Colin Barker_, May 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006611 | OEIS-A202458

Number of (n+2) X 7 binary arrays with consecutive windows of three bits considered as a binary number nondecreasing in every row and column.

Conjectures from _Colin Barker_, May 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006612 | OEIS-A202459

Number of (n+2) X 8 binary arrays with consecutive windows of three bits considered as a binary number nondecreasing in every row and column.

Conjectures from _Colin Barker_, May 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006613 | OEIS-A202460

Number of (n+2) X 9 binary arrays with consecutive windows of three bits considered as a binary number nondecreasing in every row and column.

Conjectures from _Colin Barker_, Jun 01 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 19 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006614 | OEIS-A202650

Number of ways to write n = p + p(k) + p(m) with 0 < k <= m, where p is a prime and p(.) is the partition function (A000041).

Conjecture: (i) a(n) > 0 for all n > 3. We have verified part (i) of the conjecture for all n = 4, 5, ..., 2*10^7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006615 | OEIS-A202730

Number of n X 3 nonnegative integer arrays each row and column an ascent sequence (interior element no greater than one plus up-steps preceding it) and the lower right element equal to min(n,3)-1.

Conjectures from _Colin Barker_, Jun 01 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006616 | OEIS-A202750

Triangle T(n,k) = binomial(n,k)^4 read by rows, 0<=k<=n.

Zhi-Wei Sun has conjectures related to the arithmetic mean of the polynomials formed from the rows of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Dec 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006617 | OEIS-A202752

Number of n X 4 nonnegative integer arrays with each row and column increasing from zero by 0 or 1.

Conjectures from _Colin Barker_, Jun 01 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006618 | OEIS-A202753

Number of n X 5 nonnegative integer arrays with each row and column increasing from zero by 0 or 1.

Conjectures from _Colin Barker_, Jun 01 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 23 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006619 | OEIS-A202785

Number of 3 X 3 0..n arrays with row and column sums equal.

Conjectures from _Colin Barker_, Jun 01 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006620 | OEIS-A202786

Number of 4 X 4 0..n arrays with row and column sums equal.

Conjecture: a(n) = (29 + 3*(-1)^n)/32 + (34/7)*n + (7202/525)*n^2 + (4658/189)*n^3 + (118873/3780)*n^4 + (5321/180)*n^5 + (36827/1800)*n^6 + (1285/126)*n^7 + (17581/5040)*n^8 + (2789/3780)*n^9 + (2789/37800)*n^10. - _Robert Israel_, May 03 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006621 | OEIS-A202807

Number of n X 3 nonnegative integer arrays with each row and column increasing from zero by 0, 1 or 2.

Conjectures from _Colin Barker_, Jun 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006622 | OEIS-A202808

Number of n X 4 nonnegative integer arrays with each row and column increasing from zero by 0, 1 or 2.

Conjectures from _Colin Barker_, Mar 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 24 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006623 | OEIS-A202856

Moments of the quadratic coefficient of the characteristic polynomial of a random matrix in SU(2) X SU(2) (inside USp(4)).

Conjecture: (n+2)^2*a(n) +2*(-3*n^2-5*n-1)*a(n-1) -4*(n-1)*(n-5)*a(n-2) +24*(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Dec 04 2013 [Maple's sumrecursion command applied to the above formula for a(n) produces this recurrence. - _Peter Bala_, Jul 06 2015]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006624 | OEIS-A202865

Number of 3 X 3 0..n arrays with row and column sums one greater than the previous row and column.

Conjectures from _Colin Barker_, Jun 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 25 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006625 | OEIS-A202917

For n >= 0, let n!^(1) = A053657(n+1) and, for 0 <= m <= n, C^(1)(n,m) = n!^(1)/(m!^(1)*(n-m)!^(1)). The sequence gives a triangle of numbers C^(1)(n,m) with rows of length n+1.

3) If p is an odd prime, then the (p-1)-th row contains two 1's and p-2 numbers that are multiples of p. For a conjectural generalization, see comment in A175669.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Dec 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006626 | OEIS-A202919

Number of n X 3 nonnegative integer arrays with each row and column increasing from zero by 0, 1, 2 or 3.

Conjectures from _Colin Barker_, Jun 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006627 | OEIS-A202933

Number of (n+3) X 5 binary arrays with consecutive windows of four bits considered as a binary number nondecreasing in every row and column.

Conjectures from _Colin Barker_, Jun 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006628 | OEIS-A202934

Number of (n+3) X 6 binary arrays with consecutive windows of four bits considered as a binary number nondecreasing in every row and column.

Conjectures from _Colin Barker_, Jun 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006629 | OEIS-A202935

Number of (n+3) X 7 binary arrays with consecutive windows of four bits considered as a binary number nondecreasing in every row and column.

Conjectures from _Colin Barker_, Jun 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006630 | OEIS-A202936

Number of (n+3) X 8 binary arrays with consecutive windows of four bits considered as a binary number nondecreasing in every row and column.

Conjectures from _Colin Barker_, Jun 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006631 | OEIS-A202937

Number of (n+3) X 9 binary arrays with consecutive windows of four bits considered as a binary number nondecreasing in every row and column.

Conjectures from _Colin Barker_, Jun 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006632 | OEIS-A202940

Positive numbers n such that n is not a multiple of 10 and n^3 has at most three different digits.

Conjecture: no more terms other than those of the form 10^n+1, n=0,1,2,... .

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Dec 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006633 | OEIS-A202941

For n>=0, let n!^(2)=A202367(n+1) and, for 0<=m<=n, C^(2)(n,m)=n!^(2)/(m!^(2)*(n-m)!^(2)). The sequence gives triangle of numbers C^(2)(n,m) with rows of length n+1.

Conjecture. If p is an odd prime, then the ((p-1)/2)-th row contains two 1's and (p-3)/2 numbers multiple of p. If conjectural formula in A202367 is true, then A007814(C^(2)(n,m)) =A007814(C(n,m)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Dec 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006634 | OEIS-A202943

G.f.: [Sum_{n>=0} (n+1)*(n+2)/2 * 2^(n*(n-1)) * x^n]^(1/3).

Conjecture: the characteristic function of a(n) (mod 2) equals (1+x)*(1+x^2)*(1+x^8) * Sum_{n>0} x^(32*A000695(n)), where A000695 is the sums of distinct powers of 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006635 | OEIS-A202955

Decimal expansion of Pi^Pi^Pi^Pi.

Whether this number is an integer or not is an open question. It is also an open question whether Pi^Pi^Pi^...^Pi^Pi n times is an integer for any natural n > 4. - _Eliora Ben-Gurion_, Nov 17 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Dec 26 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006636 | OEIS-A202996

Least prime q such that prime(n)^2 - q^2 - 1 and prime(n)^2 - q^2 + 1 are twin primes or 0 if no solution.

Conjecture: a(n) > 0 for n > 6. Conjecture: With Sq=sum of q for n=1 to N and Sp=sum of p(n) for n=1 to N, lim sup Sq/Sp = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Dec 27 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006637 | OEIS-A202998

Previous integer to m^2/n where m is the next odd prime after n (n excluded).

It is conjectured by Michael B Rees (Dec 2011) that for any n > 0, A110835(n) >= a(n)>=n. The Sierpinski conjecture states that: "For any n >= 2 and any k such that 1 < k <= n there exists a prime number between (k-1)*n and k*n (inclusively)". Rees has conjectured that: "For any n >= 1 and any k such that 1 < k < m^2/n where m is the next odd prime after n (n excluded), there exists a prime number between (k-1)*n and k*n (inclusively)".

猜想/问题提及段 | historical_or_current_mention_unreviewed | Michael B. Rees and _Frank M Jackson_, Jan 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006638 | OEIS-A203069

Lexicographically earliest sequence of distinct positive numbers such that a(n-1)+a(n) is odd and composite.

It is conjectured that this is a permutation of the positive integers. Is there a proof? The terms are distinct, by definition, and the sequence is clearly infinite. But does every number appear?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Dec 28 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006639 | OEIS-A203077

Alternating-parity rearrangement of natural numbers: a(n) is the smallest number such that a(n-1)^2 + a(n)^2 is odd and composite.

The maximum between a(n) and the n-th integer appears to be +-6. In the first 10k terms, the distribution of differences, from -6 to 6 is: 27, 140, 1350, 7002, 1282, 168, 31. Therefore I conjecture that Lim_{n->infinity} a(n) = n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Dec 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006640 | OEIS-A203150

(n-1)-st elementary symmetric function of the first n terms of (1,2,1,2,1,2,1,2,1,2,...)=A000034.

Conjecture: a(n) = (6*r*n+(1+3*(1-r)*n)*(1-(-1)^n))*r^(n-1)/8, where r=sqrt(2). - _Bruno Berselli_, Jan 03 2011

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006641 | OEIS-A203157

(n-1)-st elementary symmetric function of the first n triangular numbers.

Conjecture: 2*(-n+1)*a(n) +n^3*a(n-1)=0. - _R. J. Mathar_, Oct 01 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006642 | OEIS-A203159

(n-1)-st elementary symmetric function of {2,4,6,8,...,2n}.

Conjecture: a(n) +2*(-2*n+1)*a(n-1) +4*(n-1)^2*a(n-2)=0. - _R. J. Mathar_, Oct 01 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 29 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006643 | OEIS-A203165

(n-1)-st elementary symmetric function of the first n terms of the periodic sequence (3,4,1,2,3,4,1,2,...).

Conjecture: G.f. x*(1+7*x+19*x^2+50*x^3+126*x^4+432*x^5+144*x^6) / (-1+24*x^4)^2 with a(n) = 48*a(n-4)-576*a(n-8). - _R. J. Mathar_, Jul 02 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006644 | OEIS-A203166

(n-1)-st elementary symmetric function of the first n terms of (1,2,3,4,5,1,2,3,4,5,...)=A010884.

Conjecture: G.f. x*(1+3*x+11*x^2+50*x^3+274*x^4+154*x^5+188*x^6+324*x^7+576*x^8) / (-1+120*x^5)^2 with a(n) = 240*a(n-5) -14400*a(n-10). - _R. J. Mathar_, Jul 02 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006645 | OEIS-A203167

(n-1)-st elementary symmetric function of the first n terms of (2,2,1,2,2,1,2,2,1,...)=(A130196 for n>0).

Conjecture: a(n) = 8*a(n-3) - 16*a(n-6). Conjecture: G.f.: x*(1+4*x+8*x^2+12*x^3+16*x^4) / (-1+4*x^3)^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006646 | OEIS-A203230

(n-1)-st elementary symmetric function of the first n terms of A010684.

Conjecture: a(n)=6*a(n-2)-9*a(n-4) with G.f. x*(1+4*x+x^2) / (-1+3*x^2)^2 . - _R. J. Mathar_, Oct 15 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006647 | OEIS-A203231

(n-1)-st elementary symmetric function of the first n terms of the periodic sequence (3,1,3,1,3,1,3,1,...).

Conjecture: a(n) = 6*a(n-2)-9*a(n-4) with G.f. x*(1+4*x+9*x^2) / (-1+3*x^2)^2 . - _R. J. Mathar_, Oct 15 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006648 | OEIS-A203232

(n-1)-st elementary symmetric function of the first n terms of the periodic sequence (2,3,2,3,2,3,...).

Conjecture: a(n)=12*a(n-2)-36*a(n-4) with G.f. x*(1+x)*(4*x+1) / (-1+6*x^2)^2 . - _R. J. Mathar_, Oct 15 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006649 | OEIS-A203233

(n-1)-st elementary symmetric function of the first n terms of the periodic sequence (3,2,3,2,3,2,...).

Conjecture: a(n)=12*a(n-2)-36*a(n-4) with G.f. x*(1+5*x+9*x^2) / (-1+6*x^2)^2 . - _R. J. Mathar_, Oct 15 2013, verified by _Robert Israel_, May 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006650 | OEIS-A203234

(n-1)-st elementary symmetric function of the first n terms of the periodic sequence (1,1,1,2,1,1,1,2,...).

Conjecture: a(n) = 4*a(n-4)-4*a(n-8) with G.f. x*(1+2*x+3*x^2+7*x^3+5*x^4+3*x^5+x^6) / (-1+2*x^4)^2 . - _R. J. Mathar_, Oct 15 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006651 | OEIS-A203235

(n-1)-st elementary symmetric function of the first n terms of the periodic sequence (1,1,1,3,1,1,1,3,...).

Conjecture: a(n) = 6*a(n-4)-9*a(n-8) with G.f. x*(1+2*x+3*x^2+10*x^3+7*x^4+4*x^5+x^6) / ((-1+3*x^4)^2). - _R. J. Mathar_, Oct 15 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 30 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006652 | OEIS-A203242

Second elementary symmetric function of the first n terms of (1, 3, 7, 15, 31, ...).

Conjecture: (-103*n+258)*a(n) + (881*n-2116)*a(n-1) + 6*(-427*n+960)*a(n-2) + 4*(766*n-1545)*a(n-3) + 16*(-80*n+121)*a(n-4) = 0. - _R. J. Mathar_, Oct 15 2013 Conjecture and empirical g.f. verified.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 31 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006653 | OEIS-A203246

Second elementary symmetric function of the first n terms of (1,1,2,2,3,3,4,4,...).

Conjectural o.g.f.: x^2*(1 + 3*x + x^2 + x^3)/((1 - x^2)^3*(1 - x)^2). - _Peter Bala_, Aug 15 2014 Conjectural closed form: 64*a(n) = 2*n^2 -16*n/3 -3 +16*n^3/3 +2*n^4 +(-1)^n *(3-2*n^2). - _R. J. Mathar_, Oct 01 2016 Both conjectures are true. See link. - _Sela Fried_, Dec 22 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 31 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006654 | OEIS-A203278

Row sums of triangle A202941.

Conjecture. If p is odd prime, then a((p-1)/2)==2 (mod p)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Dec 31 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006655 | OEIS-A203286

Number of arrays of 2n nondecreasing integers in -3..3 with sum zero and equal numbers greater than zero and less than zero.

Conjectures from _Colin Barker_, Jun 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 31 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006656 | OEIS-A203292

Number of arrays of 4 nondecreasing integers in -n..n with sum zero and equal numbers greater than zero and less than zero.

Conjectures from _Colin Barker_, Jun 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 31 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006657 | OEIS-A203298

Second elementary symmetric function of the first n terms of (1,2,2,3,3,4,4,5,5...).

Conjectures from _Colin Barker_, Jan 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 31 2011 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006658 | OEIS-A203399

T(n, k), a triangular array read by rows, is the number of classes of equivalent 2-color n-bead necklaces (turning over is allowed) that contain k necklaces.

Conjectures: For n >= 1, let b(n) be the number of bracelets of two colors with n beads that are either periodic (period >= 2), or have reflection symmetry (achiral), or both. Then b(n) = A000029(n) - A032239(n) for n >= 3 with b(n) = A000029(n) for n = 1, 2. We have A000029(n) = 2^floor(-3 + n/2) * (7 - (-1)^n) + (1/(2*n)) * Sum_{d|n} phi(d) * 2^(n/d) for n >= 1 and A032239(n) = (1/2) * Sum_{d|n} mu(d) * (-2^floor(-2 + (n/(2*d))) * (7 - (-1)^(n/d)) + 2^(n/d)/n) for n >= 3. For 1 <= k <= n, we conjecture that T(n, k) = Sum_{d|k} mu(d)*b(k/d) for k|n, and = 0 otherwise. Note that, if 3 <= n <= 11, we have T(n, k) = A056493(k) when k|n, but this is not true (for example) for n = 12 and n = 14. We have T(12, 12) = 82 <> 81 = A056493(12) and T(14, 14) = 177 <> 175 = A056493(14). I ran the author's Mathematica program for n = 1..21 and I saw that the conjecture is OK except for n = 18 and n = 21. The program gives T(n=18, k=12) = 1 and T(n=18, k=18) = 742 while my conjecture implies that T(n=18, k=12) = 0 (since k = 12 does not divide n = 18) and T(n=18, k=18) = 743. In addition, the program gives T(n=21, k=14) = 2 and T(n=21, k=21) = 2030, while my conjecture implies that T(n=21, k=14) = 0 (since k = 14 does not divide n = 21) and

T(n=21, k=21) = 2032. Apparently, my conjecture needs to be refined.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Critzer_, Jan 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006659 | OEIS-A203470

a(n) = Product_{2 <= i < j <= n+1} (i + j).

Each term divides its successor, as (conjectured) in A102693. Each term is divisible by the corresponding superfactorial, A000178(n), as in A203471.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006660 | OEIS-A203472

a(n) = Product_{3 <= i < j <= n+2} (i + j).

Each term divides its successor, as in A203470. It is conjectured that each term is divisible by the corresponding superfactorial, A000178(n), as in A203474.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006661 | OEIS-A203484

For n>=0, let n!^(3) = A202368(n+1) and, for 0<=m<=n, C^(3)(n,m) = n!^(3)/(m!^(3)*(n-m)!^(3)). The sequence gives triangle of numbers C^(3)(n,m) with rows of length n+1.

Conjecture. If p is prime of the form 3*k+1, then the k-th row contains two 1's and k-1 numbers multiple of p; if p is prime of the form 3*k+2, then the (2*k+1)-th row contains two 1's and 2*k numbers multiple of p. Conjecture. A007814(C^(3)(n,m)) = A007814(C(n,m)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006662 | OEIS-A203498

Row sums of triangle A203484.

Conjecture. If prime p==1 (mod 3), then a((p-1)/3)==2 (mod p); if prime p==2 mod 3, then a((2*p-1)/3)==2 (mod p).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006663 | OEIS-A203509

Row sums of triangle A178473.

Conjecture. If prime p==1 (mod 4), then a((p-1)/4)==2 (mod p); if prime p==3 (mod 4), then a((p-1)/2)==2 (mod p).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006664 | OEIS-A203510

a(n) = A203482(n) / A000178(n).

It is conjectured that every term of the sequence is an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006665 | OEIS-A203520

v(n)/A000178(n); v=A203518 and A000178=(superfactorials).

It is conjectured that every term of A203520 is an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006666 | OEIS-A203521

a(n) = Product_{1 <= i < j <= n} (prime(i) + prime(j)).

Each term divides its successor, as in A203511. It is conjectured that each term is divisible by the corresponding superfactorial, A000178(n). See A093883 for a guide to related sequences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006667 | OEIS-A203523

v(n)/A000178(n); v=A203521 and A000178=(superfactorials).

It is conjectured that every term of A203523 is an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006668 | OEIS-A203524

a(n) = Product_{2 <= i < j <= n+1} (prime(i) + prime(j)).

Each term divides its successor, as in A203525. It is conjectured that each term is divisible by the corresponding superfactorial, A000178(n); as in A203526. See A093883 for a guide to related sequences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006669 | OEIS-A203526

v(n)/A000178(n); v=A203524 and A000178=(superfactorials).

It is conjectured that every term of A203526 is an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006670 | OEIS-A203527

a(n) = Product_{1 <= i < j <= n} (A018252(i) + A018252(j)); A018252 = nonprime numbers.

Each term divides its successor, as in A203528. It is conjectured that each term is divisible by the corresponding superfactorial, A000178(n); as in A203529. See A093883 for a guide to related sequences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006671 | OEIS-A203529

a(n) = A203527(n)/A000178(n-1); A000178 = (superfactorials).

It is conjectured that every term is an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006672 | OEIS-A203530

a(n) = Product_{1 <= i < j <= n} (c(i) + c(j)); c = A002808 = composite numbers.

It is conjectured that each term is divisible by the corresponding superfactorial, A000178(n); as in A203533.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006673 | OEIS-A203533

a(n) = A203530(n)/A000178(n-1); A000178 = (superfactorials).

It is conjectured that every term is an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006674 | OEIS-A203650

1/25 the number of (n+1) X 3 0..4 arrays with every 2 X 2 subblock having equal diagonal elements or equal antidiagonal elements.

Conjectures from _Colin Barker_, Jun 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006675 | OEIS-A203719

A204521(n)^2 = floor[A055812(n)/5]: Squares which written in base 5, with some digit appended, yield another square.

Conjecture: a(n) = 323*a(n-4)-323*a(n-8)+a(n-12) for n>13. - _Colin Barker_, Sep 20 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006676 | OEIS-A203751

Square root of v(2*n+1)/(3*v(2*n)), where v = A203748.

Conjecture: a(n) = n^2/(n-1)! * Product_{k = 1..n-1} (n^3 - k^3). - _Peter Bala_, Dec 01 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006677 | OEIS-A203774

Square root of v(2n)/v(2n-1), where v=A203773.

a(n) = gamma((1-i)*n)*gamma((1+i)*n)*sinh(n*Pi)/Pi (conjecture). - _Velin Yanev_, Nov 15 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006678 | OEIS-A203775

Square root of v(2*n+1) / (2*v(2*n)), where v=A203773.

a(n) = n*gamma((1-i)*n)*gamma((1+i)*n)*sinh(n*Pi)/Pi (conjecture). - _Velin Yanev_, Nov 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006679 | OEIS-A203829

Number of (n+1) X 3 0..2 arrays with every 2 X 2 subblock having equal diagonal elements or equal antidiagonal elements.

Conjectures from _Colin Barker_, Jun 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 06 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006680 | OEIS-A203985

Lexicographically earliest permutation of the primes such that successive absolute differences yield a permutation of all nonprime numbers.

It is only conjectured that this sequence is a permutation of the primes and that the successive differences yield all composite integers. The definition is rather to be understood as "The next term is chosen as the smallest prime not occurring earlier such that the successive absolute differences only yield 1 or composite numbers, and none of these occurs twice." - _M. F. Hasler_, Jan 09 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_, Jan 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006681 | OEIS-A204021

Triangle read by rows: row n shows the coefficients of the characteristic polynomial of the n-th principal submatrix of min(2i-1,2j-1) (A157454).

Conjectural o.g.f.: t*(1-x-x^2*t)/(1-2*t*(1-x)+t^2*x^2) = (1-x)*t + (2-4*x+x^2)*t^2 + ... (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006682 | OEIS-A204061

G.f.: exp(Sum_{n>=1} A001333(n)^2 * x^n/n) where A001333(n) = A002203(n)/2, one-half the companion Pell numbers.

a(n) == 1 (mod 5) iff n has no 2's in its base 5 expansion (A023729), otherwise a(n) == 0 (mod 5); this is a conjecture needing proof.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006683 | OEIS-A204065

Least nonnegative integer k with n+k and n+k^2 both prime.

Conjecture: For any n > 0 not among 1, 21, 326, 341, 626, we have a(n) < sqrt(n)*log(n). If n > 626 is not equal to 971, then n+k and n+k^2 are both prime for some 0< k <

sqrt(n)*log(n). Also, n+k^2 is prime for some 0< k <= sqrt(n) if n > 43181.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006684 | OEIS-A204070

Number of (n+1) X 3 0..2 arrays with every 2 X 2 subblock having equal diagonal elements or equal antidiagonal elements, and new values 0..2 introduced in row major order.

Conjectures from _Colin Barker_, Jun 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006685 | OEIS-A204137

Triangle T(r+1,c) = |T(r,c) - T(r,c+1)| of positive integers such that T(r,c) is prime iff r=1 and no number occurs twice.

It is conjectured that the first row of the table is a permutation of the primes and the whole table, i.e., this sequence, a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_, Jan 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006686 | OEIS-A204187

a(n) = Sum_{m=1..n-1} m^(n-1) modulo n.

a(n) = n - 1 if n is 1 or a prime, by Fermat's little theorem. It is conjectured that the converse is also true; see A055032 and A201560 and note that a(n) = n-1 <==> A055032(n) = 1 <==> A201560(n) = 0. a(n) = n/2 iff n is of the form 4k+2 (conjectured). - _Ivan Neretin_, Sep 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Jan 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006687 | OEIS-A204214

Number of length 6 nonnegative integer arrays starting and ending with 0 with adjacent elements differing by no more than n.

Conjectures from _Colin Barker_, Jun 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006688 | OEIS-A204215

Number of length 7 nonnegative integer arrays starting and ending with 0 with adjacent elements differing by no more than n.

Conjectures from _Colin Barker_, Jun 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006689 | OEIS-A204216

Number of length 8 nonnegative integer arrays starting and ending with 0 with adjacent elements differing by no more than n.

Conjectures from _Colin Barker_, Jun 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006690 | OEIS-A204235

Permanent of the n-th principal submatrix of A143182.

Conjecture: a(1) and a(2) are the only terms that are odd numbers. - _Stefano Spezia_, Oct 28 2018 The conjecture is true. This is because the permanent and determinant of every integer matrix are congruent modulo 2. The determinant of the matrix here is known to be (-1)^(n-1)*(n+1)*2^(n-2), i.e., even for n >= 3 (see Cloitre's comment in A001792). - _Sela Fried_, Feb 11 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006691 | OEIS-A204238

Determinant of the n-th principal submatrix of A204237.

Conjectures from _Colin Barker_, Feb 21 2015: (Start) a(n) = (-4)^(n-2)*(1-9*n), which confirms Barker's conjectures (see Fried link). - _Sela Fried_, Jan 29 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006692 | OEIS-A204240

Determinant of the n-th principal submatrix of A204158.

Conjectures from _Colin Barker_, Feb 21 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006693 | OEIS-A204262

Permanent of the n-th principal submatrix of A003983.

Conjecture: Limit_{n->oo} (a(n)/n!^2)^(1/n) = 2/Pi. - _Vaclav Kotesovec_, Aug 05 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006694 | OEIS-A204399

Numbers k such that floor(2^k / 3^n) = 1.

The conjecture is false; A195176(n) - a(n) increases without bound (though not monotonically) since log_2(3) < 3 - sqrt(2). - _Charles R Greathouse IV_, Jan 15 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006695 | OEIS-A204503

Squares n^2 such that floor(n^2/9) is again a square.

Conjectures: a(n) = 9*(n-4)^2 for n>5. G.f.: x^2*(7*x^6-12*x^5-11*x^4-x-1) / (x-1)^3. - _Colin Barker_, Sep 15 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006696 | OEIS-A204519

Square root of floor(A055851(n)/6).

Conjecture (for n>=8): a(n) = 10*a(n-3) - a(n-6). - _Vaclav Kotesovec_, Nov 26 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006697 | OEIS-A204558

Row sums of the triangle A045975.

Conjecture: a(n) = (n^2*(-(-1)^n+n^2))/2. G.f.: -x*(x^6 +4*x^5 +31*x^4 +24*x^3 +31*x^2 +4*x +1) / ((x -1)^5*(x +1)^3). - _Colin Barker_, Apr 02 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jan 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006698 | OEIS-A204565

Number of (n+1) X 2 0..3 arrays with every 2 X 2 subblock having equal diagonal elements or equal antidiagonal elements, and new values 0..3 introduced in row major order.

Conjectures from _Colin Barker_, Jun 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006699 | OEIS-A204573

A204519(n)^2 = floor(A055851(n)/6): Squares which written in base 6, with some digit appended, yield another square.

Conjecture: a(n) = 99*a(n-3)-99*a(n-6)+a(n-9) for n>10. - _Colin Barker_, Sep 20 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006700 | OEIS-A204609

Number of (n+1) X 2 0..1 arrays with the determinants of 2 X 2 subblocks nondecreasing rightwards and downwards.

Conjectures from _Colin Barker_, Feb 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006701 | OEIS-A204624

Number of (n+1) X 3 0..2 arrays with every 2 X 2 subblock having unequal diagonal elements or unequal antidiagonal elements, and new values 0..2 introduced in row major order.

Conjectures from _Colin Barker_, Jun 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006702 | OEIS-A204645

Number of (n+1) X 3 0..1 arrays with column and row pair sums b(i,j)=a(i,j)+a(i,j-1) and c(i,j)=a(i,j)+a(i-1,j) nondecreasing in column and row directions, respectively.

Conjectures from _Colin Barker_, Jun 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006703 | OEIS-A204646

Number of (n+1) X 4 0..1 arrays with column and row pair sums b(i,j)=a(i,j)+a(i,j-1) and c(i,j)=a(i,j)+a(i-1,j) nondecreasing in column and row directions, respectively.

Conjectures from _Colin Barker_, Jun 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006704 | OEIS-A204647

Number of (n+1) X 5 0..1 arrays with column and row pair sums b(i,j)=a(i,j)+a(i,j-1) and c(i,j)=a(i,j)+a(i-1,j) nondecreasing in column and row directions, respectively.

Conjectures from _Colin Barker_, Jun 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006705 | OEIS-A204648

Number of (n+1) X 6 0..1 arrays with column and row pair sums b(i,j)=a(i,j)+a(i,j-1) and c(i,j)=a(i,j)+a(i-1,j) nondecreasing in column and row directions, respectively.

Conjectures from _Colin Barker_, Jun 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006706 | OEIS-A204707

Number of (n+1) X 3 0..1 arrays with the permanents of all 2 X 2 subblocks equal and nonzero.

Conjectures from _Colin Barker_, Jun 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006707 | OEIS-A204708

Number of (n+1) X 4 0..1 arrays with the permanents of all 2 X 2 subblocks equal and nonzero.

Conjectures from _Colin Barker_, Jun 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006708 | OEIS-A204770

Expansion of psi(x^3) * f(-x) in powers of x where psi(), f() are Ramanujan theta functions.

Conjecture: density of zero terms slowly approaches 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Jan 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006709 | OEIS-A204842

Triangle by rows relating to A081696.

Conjecture: T(n,1) = 2*A109262(n). T(n,2)=2*A109263(n). - _R. J. Mathar_, Jul 21 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Jan 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006710 | OEIS-A204844

Cyclic primes that are not absolute primes (A003459).

Conjecture: The sequence is finite, with 999331 being the last term (cf. A293142). - _Felix Fröhlich_, Aug 05 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006711 | OEIS-A204983

a(n) = 2^(k-1)-2^(j-1), where (2^(k-1),2^(j-1)) is the least pair of distinct positive powers of 2 for which n divides 2^(k-1)-2^(j-1).

(Conjecture) Equivalently, the solution set of 2^p * (2^q - 1) = x * y, OR 2^q - 1 = 2^p * x * y, for at most one of the naturals x and y being given; unknown p and q in the integers; then a(n) = 2^p * (2^q - 1) where p and q are directly related to n (see formula). - _Andrew T. Porter_, Dec 20 2022 Conjecture: a(n) = 2^A007814(n) * (2^A007733(n) - 1). - _Andrew T. Porter_, Dec 20 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006712 | OEIS-A204988

The index j < k such that n divides 2^k - 2^j, where k is the least index (A204987) for which such j exists.

a(n) = A007814(n) + (1-(-1)^n)/2 (conjecture). - _Velin Yanev_, Nov 14 2016. The above conjecture is true because the definition of this sequence and A204987 requires j to be at least 1 and 2^k - 2^j can be written 2^j*(2^(k-j) - 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006713 | OEIS-A205163

Number of (n+1) X 2 0..3 arrays with every 2 X 2 subblock having unequal diagonal elements or unequal antidiagonal elements, and new values 0..3 introduced in row major order.

Conjectures from _Colin Barker_, Mar 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006714 | OEIS-A205186

Number of (n+1) X (n+1) 0..1 arrays with the number of clockwise edge increases in every 2 X 2 subblock differing from each horizontal or vertical neighbor.

Conjectures from _Colin Barker_, Jan 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006715 | OEIS-A205220

Number of (n+1) X 3 0..1 arrays with the number of equal 2 X 2 subblock diagonal pairs and equal antidiagonal pairs differing from each horizontal or vertical neighbor, and new values 0..1 introduced in row major order.

Conjectures from _Colin Barker_, Jun 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006716 | OEIS-A205322

Smallest k>=0 such that (2^n-k)*2^n-1 and (2^n-k)*2^n+1 are a twin prime pair; or -1 if no such k exists.

Conjecture: there is always at least one k>=0 unless n=3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Jul 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006717 | OEIS-A205329

Number of (n+1) X 3 0..2 arrays with the number of equal 2 X 2 subblock diagonal pairs and equal antidiagonal pairs differing from each horizontal or vertical neighbor, and new values 0..2 introduced in row major order.

Conjectures from _Colin Barker_, Jun 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006718 | OEIS-A205342

Number of length 5 nonnegative integer arrays starting and ending with 0 with adjacent elements unequal but differing by no more than n.

Conjectures from _Colin Barker_, Jun 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006719 | OEIS-A205343

Number of length 6 nonnegative integer arrays starting and ending with 0 with adjacent elements unequal but differing by no more than n.

Conjectures from _Colin Barker_, Jun 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006720 | OEIS-A205344

Number of length 7 nonnegative integer arrays starting and ending with 0 with adjacent elements unequal but differing by no more than n.

Conjectures from _Colin Barker_, Jun 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006721 | OEIS-A205345

Number of length 8 nonnegative integer arrays starting and ending with 0 with adjacent elements unequal but differing by no more than n.

Conjectures from _Colin Barker_, Jun 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006722 | OEIS-A205354

Number of (n+1) X 2 0..2 arrays with every 2 X 2 subblock having the same number of equal diagonal or antidiagonal elements, and new values 0..2 introduced in row major order.

Conjectures from _Colin Barker_, Mar 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006723 | OEIS-A205492

Expansion of (1-x^2-x^3-x^4+x^5)/((1-x)^3*(1-x-x^2)^2*(1-2*x-x^2+x^3)).

CONJECTURE 1. a(n) = M_{n,2} = M_{2,n}, where M = A205497. CONJECTURE 2. lim_{n->oo} a(n+1)/a(n) = (2*cos(Pi/7))^2-1 = A116425-1 = spectral radius of the 3 X 3 unit-primitive matrix (see [Jeffery]) A_{7,2} = [0,0,1; 0,1,1; 1,1,1].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Jan 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006724 | OEIS-A205493

Third row or column of table A205497.

Conjecture 1. a(n) = M_{n,3} = M_{3,n}, where M = A205497. Conjecture 2. Let w=2*cos(Pi/9). Then lim_{n->oo} a(n+1)/a(n) = w^3-2*w = spectral radius of the 4 X 4 unit-primitive matrix (see [Jeffery]) A_{9,3} = [0,0,0,1; 0,0,1,1; 0,1,1,1; 1,1,1,1].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Jan 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006725 | OEIS-A205494

Conjectured row or column n=4 of array A205497.

Conjectured row or column n=4 of array A205497. CONJECTURE 1. a(n) = M_{n,4} = M_{4,n}, where M = A205497. CONJECTURE 2. Let w=2*cos(Pi/11). Then lim_{n->oo} a(n+1)/a(n) = w^4-3*w^2+1 = spectral radius of the 5 X 5 unit-primitive matrix (see [Jeffery]) A_{11,4} = [0,0,0,0,1; 0,0,0,1,1; 0,0,1,1,1; 0,1,1,1,1; 1,1,1,1,1].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Jan 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006726 | OEIS-A205495

Convolution related to array A205497 and to generating functions for the rows of the array form of A050446.

CONJECTURE 1. a(n) = M_{n,5} = M_{5,n}, where M = A205497. CONJECTURE 2. Let w=2*cos(Pi/13). Then lim_{n->oo} a(n+1)/a(n) = w^5-4*w^3+3*w = spectral radius of the 6 X 6 unit-primitive matrix (see [Jeffery]) A_{13,5} = [0,0,0,0,0,1; 0,0,0,0,1,1; 0,0,0,1,1,1; 0,0,1,1,1,1; 0,1,1,1,1,1; 1,1,1,1,1,1].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Jan 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006727 | OEIS-A205497

Triangle read by rows: Zig-zag Eulerian number triangle T(n, k).

(Previous name:) Omitting the first two ones, a rectangular array M read by antidiagonals in which entry M_{n-k, k} in row n-k and column k, 0 <= k <= n, gives the coefficient of x^k in the numerator of the conjectured generating function for row n + 3 of the tabular form of A050446. Conjecture: 1. M_{n, k} = M_{k, n}, for all n and k; that is, M is symmetric about the central terms {1, 3, 31, 623,...}. (This has been verified for the first 100 antidiagonals of M.) Conjecture: 2. For m in {3, 4,...}, row m of array A050446 has generating function of the form H_m(x)/(1 - x)^m, in which the numerator H_m(x) is a polynomial of degree m - 3 in x with coefficients given by the entries of the (m - 3)-th antidiagonal of M containing the sequence of entries {M_{m-3-j,j}}, j=0..m-3 (see the example below). It is known that H_1(x) = H_2(x) = 1. Conjecture: 3. Define the Chebyshev polynomials of the second kind by U_0(t) = 1, U_1(t) = 2*t and U_r(t) = 2*t*U_(r-1)(t) - U_(r-2)(t) (r > 1). Assuming Conjecture 1, lim_{n -> infinity} M_{n+1, k}/M_{n, k} = U_k(cos(Pi/(2*k+3))) = spectral radius of the (k+1) X (k+1) unit-primitive matrix (see [Jeffery]) A_{2*k+3, k} = [0,...,0,1; 0,...,0,1,1; ...; 0,1,...,1; 1,...,1], with identical limits for the columns of the transpose M^T of M. Conjecture: 4. Let S(u, v) denote the entry in row u and column v of triangle S = A187660, 0 <= v <= u. Define the polynomials P_u(x) = Sum[S(u, v)*x^v]. Assuming Conjecture 1, then (i) the generating function for row (or column) n of M is of the form Remarks: The coefficients in the numerators G_n(x) appear to have no pattern. The polynomial P_j(x), j in {1,...,n+1}, of Conjecture 4 is also obtained from the characteristic polynomial of the unit-primitive matrix A_{2*j+3,j} of Conjecture 3 by taking the exponents of the latter in reverse order; and P_j(x) is otherwise identical to the characteristic polynomial of the unit-primitive matrix A_{2*j+3,1}. Conjecture: The Eulerian zig-zag polynomials have only negative and simple real roots and form a Sturm sequence, that is, p(n+1, x) has n real roots separated by the roots of p(n, x). This property was proved by Frobenius for the classical Eulerian polynomials. - _Peter Luschny_, Jun 04 2024 Conjecture: 5.1. G.f. for column 0 of M is 1/(1-x) (A000012). Conjecture: 5.2. G.f. for column 1 of M is 1/((1-x)^2*(1-x-x^2)) (A001924). Conjecture: 5.3. G.f. for column 2 of M is (1 - x^2 - x^3 - x^4 + x^5)/((1-x)^3*(1-x-x^2)^2*(1 - 2*x - x^2 + x^3)) (A205492). Conjecture: 5.4. G.f. for column 3 of M is (1 + x - 6*x^2 - 15*x^3 + 21*x^4 + 35*x^5 - 13*x^6 - 51*x^7 + 3*x^8 + 21*x^9 + 5*x^10 + x^11 - 5*x^12 - x^13 - x^14)/((1-x)^4*(1-x-x^2)^3*(1 - 2*x - x^2 + x^3)^2*(1 - 2*x - 3*x^2 + x^3 + x^4)) (A205493). Conjecture: 5.5. G.f. for column 4 of M is (1 + 4*x - 31*x^2 - 67*x^3 + 348*x^4 + 418*x^5 - 1893*x^6 - 1084*x^7 + 4326*x^8 + 4295*x^9 - 7680*x^10 - 9172*x^11 + 9104*x^12 + 11627*x^13 - 5483*x^14 - 10773*x^15 + 1108*x^16 + 7255*x^17 + 315*x^18 - 3085*x^19 - 228*x^20 + 669*x^21 + 102*x^22 - 23*x^23 - 45*x^24 - 16*x^25 + 11*x^26 + 2*x^27 - x^28)/((1-x)^5*(1-x-x^2)^4*(1 - 2*x - x^2 + x^3)^3*(1 - 2*x - 3*x^2 + x^3 + x^4)^2*(1 - 3*x - 3*x^2 + 4*x^3 + x^4 - x^5)) (A205494).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Jan 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006728 | OEIS-A205510

Binary Hamming distance between prime(n) and prime(n+1).

We call "Hamming's twin primes" the pairs of consecutive primes (p,q) with Hamming distance 1. They are (2,3), (5,7), (17,19), (19,23), (29,31), (41,43), (43,47), (67,71), (97,101), ..., (A205511,A205302). As in Twin Primes Conjecture, we conjecture that there exist infinitely many Hamming's twin pairs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006729 | OEIS-A205572

E.g.f.: 1/(cos(x) - sinh(x)).

a(2^n + k) == a(k) (mod 2^n) for k>=0, n>=1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006730 | OEIS-A205573

Array M read by antidiagonals in which successive rows evidently converge to A001405 (central binomial coefficients).

CONJECTURE 1. Let M(n,k) (n,k >= 0) denote the entry in row n and column k of the array. For all n, M(n,j) = A001405(j), j=0,...,2*n+1; hence row n of M -> A001405 as n -> infinity. CONJECTURE 2 (Rows of M). Let S(n,i) denote term i in row n of A115139, i=0,...,floor(n/2), and let T(n,j) denote term j in row n of A108299, j=0,...,n. The generating function for row n of M is of the form F_n(x) =sum[i=0,...,floor(n/2) S(n,i)*x^(2*i)]/sum[j=0,...,n T(n,j)*x^j]. CONJECTURE 3 (Columns of M). Let D(m,k) denote term m in column k of A191314, m=0,...,floor(k/2). The generating function for column k of M is of the form G_k(x)=sum[m=0,...,floor(k/2) D(m,k)*x^m]/(1-x).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Jan 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006731 | OEIS-A205616

Even numbers that are not the sum of two non-Ramanujan primes (A174635).

No other terms < 2*10^8. Conjectured to be complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Nicholson_ and _Donovan Johnson_, Jan 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006732 | OEIS-A205649

Hamming distance between twin primes.

Conjecture: The sequence is unbounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006733 | OEIS-A205798

List of exceptional parameters u of irreducible two-weight cyclic codes.

The list is conjectured to be complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 31 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006734 | OEIS-A205810

Irregular triangle read by rows: Whitney numbers c_{n,k} (n >= 0, 0 <= k <= 2n) of Lucas lattices.

Conjecture: for n >= 1, t(n, x + 1) = Sum_{k = 0..n} c(n, n-k)*t(k, x) - c(n, n). Cf. A097724. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 31 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006735 | OEIS-A205901

Number of primes between two consecutive primes of the form n^2 + 1.

Conjecture: this sequence is infinite, because it is conjectured that the sequence A002496 is infinite, although this has never been proved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Feb 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006736 | OEIS-A205989

a(n) = smallest prime >= 10^n with primitive root 10.

The decimal expansion of 1/a(n) includes every possible block of n digits. Conjecturally, a(n) is the smallest value with this property. If Artin's conjecture is true, there are an infinite number of primes with primitive root 10, which implies that a(n) exists for all n. Artin's conjecture remains open. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006737 | OEIS-A205999

Inverse Euler transform of A195980.

The sequence is conjectured to be positive, nondecreasing and strictly convex.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 02 2012, Feb 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006738 | OEIS-A206000

Inverse Euler transform of 2 - A195981.

The sequence is conjectured to be nonnegative and convex.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 02 2012, Feb 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006739 | OEIS-A206170

Number of 2 X (n+1) 0..3 arrays with every 2 X 2 subblock in a row having an equal number of equal diagonal or equal antidiagonal elements, adjacent rows differing in this number, and new values 0..3 introduced in row major order.

Conjectures from _Colin Barker_, Mar 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006740 | OEIS-A206198

Number of (n+1) X (n+1) 0..3 arrays with every 2 X 3 or 3 X 2 subblock having exactly one clockwise edge increases.

Conjectures from _Colin Barker_, Jan 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006741 | OEIS-A206246

Numbers k such that the greatest prime divisor p of k^2+1 has the property that (p - k)^2 + 1 = p.

Because this sequence is connected with A002496, it is conjectured that the set of this numbers is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Feb 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006742 | OEIS-A206248

Number of (n+1) X 2 0..3 arrays with every 2 X 2 subblock having zero permanent.

Conjectures from _Colin Barker_, Jun 14 2018: (Start) Barker's conjectures are true (see Fried link). - _Sela Fried_, Feb 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006743 | OEIS-A206261

Number of (n+1) X 3 0..1 arrays with the number of rightwards and downwards edge increases in each 2 X 2 subblock equal to the number in all its horizontal and vertical neighbors.

Conjectures from _Colin Barker_, Jun 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006744 | OEIS-A206262

Number of (n+1) X 4 0..1 arrays with the number of rightwards and downwards edge increases in each 2 X 2 subblock equal to the number in all its horizontal and vertical neighbors.

Conjectures from _Colin Barker_, Jun 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006745 | OEIS-A206263

Number of (n+1) X 5 0..1 arrays with the number of rightwards and downwards edge increases in each 2 X 2 subblock equal to the number in all its horizontal and vertical neighbors.

Conjectures from _Colin Barker_, Jun 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006746 | OEIS-A206264

Number of (n+1) X 6 0..1 arrays with the number of rightwards and downwards edge increases in each 2 X 2 subblock equal to the number in all its horizontal and vertical neighbors.

Conjectures from _Colin Barker_, Jun 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006747 | OEIS-A206332

Complement of A092754.

Conjecture: these are the positions words in A076478 that start with 0 and end with 1. - _Clark Kimberling_, Jul 14 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, May 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006748 | OEIS-A206336

Number of (n+1) X 2 0..3 arrays with every 2 X 2 subblock having nonzero determinant and commuting with every horizontal or vertical neighbor.

Conjectures from _Colin Barker_, Jun 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 06 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006749 | OEIS-A206464

Number of length-n Catalan-RGS (restricted growth strings) such that the RGS is a valid mixed-radix number in falling factorial basis.

Conjecture: a(n) = Sum_{k = 0..floor(n/4)} (-1)^k * C(floor(n/2) + 1 - k, k + 1) * a(n - 1 - k), a(0) = 1. - _Gionata Neri_, Jun 17 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Feb 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006750 | OEIS-A206622

G.f.: Product_{n>0} ((1+x^n)/(1-x^n))^(n^2).

Let A(x) denote the g.f. and let m be an integer. Define a sequence by u(n) = [x^n] A(x)^(m*n). We conjecture that the supercongruence u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) holds for all positive integers n and r and all primes p >= 5. Cf. A380582. - _Peter Bala_, Jan 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006751 | OEIS-A206687

Number of n X 2 0..3 arrays with no element equal to another within two positions in the same row or column, and new values 0..3 introduced in row major order.

Conjectures from _Colin Barker_, Feb 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006752 | OEIS-A206716

(1/5)A206715.

It is conjectured that all the terms of this sequence are integers; this may be equivalent to the conjecture at A131853.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006753 | OEIS-A206815

Position of n+pi(n) in the joint ranking of {j+pi(j)} and {k+(k+1)/log(k+1)}.

The sequences A206815, A206818, A206827, A206828 illustrate the closeness of {j+pi(j)} to {k+(k+1)/log(k+1)}, as suggested by the prime number theorem and the conjecture that all the terms of A206827 and A206828 are in the set {1,2,3}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006754 | OEIS-A206824

Number of solutions (n,k) of s(k) = s(n) (mod n), where 1 <= k < n and s(k) = k(k+1)/2.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006755 | OEIS-A206911

Position of n-th partial sum of the harmonic series when all the partial sums are jointly ranked with the set {log(k+1)}; complement of A206912.

Conjecture: the difference sequence of A206911 consists of 2s and 3s, and the ratio (number of 3s)/(number of 2s) tends to a number between 3.5 and 3.6. Similar conjectures can be stated for difference sequences based on jointly ranked sets, such as A206903, A206906, A206928, A206805, A206812, and A206815. Adamczewski provides a proof abstract regarding the above explicit conjecture in linked article. - _Michael De Vlieger_, Aug 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006756 | OEIS-A206912

Position of log(n+1) when the partial sums of the harmonic series are jointly ranked with the set {log(k+1)}; complement of A206911.

Conjecture: the difference sequence of A206912 consists of 1s and 2s; see Comments at A206911.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006757 | OEIS-A207020

Number of n X 4 0..1 arrays avoiding 0 0 0 and 0 1 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006758 | OEIS-A207064

Number of n X 4 0..1 arrays avoiding 0 0 1 and 0 1 0 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006759 | OEIS-A207065

Number of n X 5 0..1 arrays avoiding 0 0 1 and 0 1 0 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006760 | OEIS-A207066

Number of n X 6 0..1 arrays avoiding 0 0 1 and 0 1 0 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006761 | OEIS-A207067

Number of n X 7 0..1 arrays avoiding 0 0 1 and 0 1 0 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006762 | OEIS-A207080

The smallest Carmichael number k such that phi(k) does not divide (k-1)^n, where phi is the Euler totient function.

Conjecture: phi(a(n)) divides (a(n)-1)^(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Feb 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006763 | OEIS-A207106

Number of n X 3 0..1 arrays avoiding 0 0 0 and 0 1 1 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Feb 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006764 | OEIS-A207107

Number of n X 4 0..1 arrays avoiding 0 0 0 and 0 1 1 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Feb 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006765 | OEIS-A207108

Number of n X 5 0..1 arrays avoiding 0 0 0 and 0 1 1 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006766 | OEIS-A207109

Number of n X 6 0..1 arrays avoiding 0 0 0 and 0 1 1 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006767 | OEIS-A207110

Number of n X 7 0..1 arrays avoiding 0 0 0 and 0 1 1 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006768 | OEIS-A207118

Number of n X 3 0..1 arrays avoiding 0 0 1 and 0 1 0 horizontally and 0 0 1 and 0 1 1 vertically.

Conjectures from _Colin Barker_, Feb 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006769 | OEIS-A207142

Number of (n+1) X 2 0..2 arrays with every 2 X 2 subblock having nonzero determinant and commuting with every horizontal or vertical neighbor.

Conjectures from _Colin Barker_, Jun 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006770 | OEIS-A207165

Number of n X 4 0..1 arrays avoiding 0 0 0 and 1 0 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006771 | OEIS-A207166

Number of n X 5 0..1 arrays avoiding 0 0 0 and 1 0 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006772 | OEIS-A207167

Number of n X 6 0..1 arrays avoiding 0 0 0 and 1 0 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006773 | OEIS-A207168

Number of n X 7 0..1 arrays avoiding 0 0 0 and 1 0 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006774 | OEIS-A207255

Number of 4 X n 0..1 arrays avoiding 0 0 1 and 1 0 1 horizontally and 0 1 0 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006775 | OEIS-A207256

Number of 5 X n 0..1 arrays avoiding 0 0 1 and 1 0 1 horizontally and 0 1 0 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006776 | OEIS-A207257

Number of 6 X n 0..1 arrays avoiding 0 0 1 and 1 0 1 horizontally and 0 1 0 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006777 | OEIS-A207258

Number of 7 X n 0..1 arrays avoiding 0 0 1 and 1 0 1 horizontally and 0 1 0 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006778 | OEIS-A207262

a(n) = 2^(4n - 2) + 1.

Conjecture: Let m = 4n - 2. a(n) equals the sum of the m-th powers of the divisors of m divided by the sum of the m-th powers of the odd divisors of m. - _Ivan N. Ianakiev_, Jan 29 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alonso del Arte_, Feb 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006779 | OEIS-A207302

Number of n X 5 0..1 arrays avoiding 0 0 0 and 1 0 1 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006780 | OEIS-A207303

Number of n X 6 0..1 arrays avoiding 0 0 0 and 1 0 1 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006781 | OEIS-A207304

Number of n X 7 0..1 arrays avoiding 0 0 0 and 1 0 1 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006782 | OEIS-A207318

a(n) = Sum_{k=0..n-1} (-1)^k*A000172(k).

Conjecture: (n-1)^2*a(n) +(2*n-3)*(3*n-5)*a(n-1) +(-15*n^2+53*n-48)*a(n-2) +8*(n-2)^2*a(n-3)=0. - _R. J. Mathar_, Nov 28 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006783 | OEIS-A207319

a(n) = Sum_{k=0..n-1} (-1)^k*k*A000172(k).

Conjecture: (n-1)*(n-2)*a(n) +2*(n^2+3*n-9)*a(n-1) +3*(-13*n^2+63*n-80)*a(n-2) +4*(17*n^2-96*n+136)*a(n-3) -32*(n-3)^2*a(n-4)=0. - _R. J. Mathar_, Nov 28 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006784 | OEIS-A207323

a(n) = Sum_{k=0..n} k*A002893(k).

Conjecture: n*(n-1)*a(n) +(-14*n^2+23*n-12)*a(n-1) +(52*n^2-151*n+120)*a(n-2) +3*(-22*n^2+79*n-72)*a(n-3) +27*(n-2)^2*a(n-4)=0. - _R. J. Mathar_, Nov 28 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006785 | OEIS-A207363

Number of n X 3 0..1 arrays avoiding 0 0 0 and 0 1 1 horizontally and 0 0 1 and 0 1 1 vertically.

Conjectures from _Colin Barker_, Feb 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006786 | OEIS-A207375

Irregular array read by rows in which row n lists the (one or two) central divisors of n in increasing order.

Conjecture 1: It appears that the n-th record in this sequence is the last member of row A008578(n). The conjecture 1 follows from Bertrand's Postulate. - _Charles R Greathouse IV_, Feb 11 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Feb 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006787 | OEIS-A207384

A206815(n+1)-A206815(n).

The sequences A206815, A206818, A207384, A207835 illustrate the closeness of {j+pi(j)} to {k+(k+1)/log(k+1)}, as suggested by the prime number theorem and the conjecture that all the terms of A207384 and A207835 are in the set {1,2,3}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006788 | OEIS-A207386

G.f.: Sum_{n>=0} Product_{k=1..n} (q^k - 1) where q = (1+x)/(1+x^3).

At present Bala's identity is conjectural and needs formal proof. G.f.: Sum_{n>=0} q^(-n^2) * Product_{k=1..n} (q^(2*k-1) - 1) where q = (1+x)/(1+x^3). [Based on Peter Bala's conjecture in A158690]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006789 | OEIS-A207388

Number of n X 5 0..1 arrays avoiding 0 0 1 and 0 1 0 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006790 | OEIS-A207389

Number of n X 6 0..1 arrays avoiding 0 0 1 and 0 1 0 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006791 | OEIS-A207390

Number of n X 7 0..1 arrays avoiding 0 0 1 and 0 1 0 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006792 | OEIS-A207397

G.f.: Sum_{n>=0} Product_{k=1..n} (q^k - 1) where q = (1+x)/(1+x^2).

At present Bala's identity is conjectural and needs formal proof. G.f.: Sum_{n>=0} q^(-n^2) * Product_{k=1..n} (q^(2*k-1) - 1) where q = (1+x)/(1+x^2). [Based on Peter Bala's conjecture in A158690]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006793 | OEIS-A207399

Number of n X 4 0..1 arrays avoiding 0 0 1 and 0 1 1 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Mar 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006794 | OEIS-A207400

Number of n X 5 0..1 arrays avoiding 0 0 1 and 0 1 1 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006795 | OEIS-A207401

Number of n X 6 0..1 arrays avoiding 0 0 1 and 0 1 1 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006796 | OEIS-A207402

Number of n X 7 0..1 arrays avoiding 0 0 1 and 0 1 1 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006797 | OEIS-A207433

G.f.: Sum_{n>=0} Product_{k=1..n} (q^k - 1) where q = (1-x^3)/(1-x).

At present Bala's identity is conjectural and needs formal proof. G.f.: Sum_{n>=0} q^(-n^2) * Product_{k=1..n} (q^(2*k-1) - 1) where q = (1-x^3)/(1-x). [Based on Peter Bala's conjecture in A158690]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006798 | OEIS-A207449

Number of n X 4 0..1 arrays avoiding 0 0 0 and 0 0 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006799 | OEIS-A207450

Number of n X 5 0..1 arrays avoiding 0 0 0 and 0 0 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006800 | OEIS-A207451

Number of n X 6 0..1 arrays avoiding 0 0 0 and 0 0 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006801 | OEIS-A207452

Number of n X 7 0..1 arrays avoiding 0 0 0 and 0 0 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006802 | OEIS-A207569

G.f.: Sum_{n>=0} Product_{k=1..n} ((1+x)^(2*k-1) - 1).

Conjectural g.f.: Sum_{n>=0} (-1)^n*Product_{k = 1..n} 1 + (-1/(1 + x))^k. - _Peter Bala_, Dec 04 2020 Conjectural g.f.s: Sum_{n >= 0} (-1)^n*(1 + x)^(n+1)*Product_{k = 1..n} (1 + (-1)^k*(1 + x)^k)^2. Also

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006803 | OEIS-A207590

Number of 3 X n 0..1 arrays avoiding 0 0 0 and 0 0 1 horizontally and 0 0 1 and 1 1 1 vertically.

Conjectures from _Colin Barker_, Mar 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006804 | OEIS-A207596

Number of n X 5 0..1 arrays avoiding 0 0 0 and 0 1 0 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006805 | OEIS-A207597

Number of n X 6 0..1 arrays avoiding 0 0 0 and 0 1 0 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006806 | OEIS-A207598

Number of n X 7 0..1 arrays avoiding 0 0 0 and 0 1 0 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006807 | OEIS-A207642

Expansion of g.f.: Sum_{n>=0} x^n * Product_{k=0..n-1} (1 + x^(n+k)).

Conjecture: a(n) is the number of partitions p of n into distinct parts such that max(p) <= 1 + 2*min(p), for n >= 1 (as in the Mathematica program at A241061). - _Clark Kimberling_, Apr 16 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006808 | OEIS-A207656

Number of n X 3 0..1 arrays avoiding 0 0 0 and 1 1 1 horizontally and 0 0 1 and 1 0 0 vertically.

Conjectures from _Colin Barker_, Jun 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006809 | OEIS-A207672

n + [ns/r] + [nt/r], where []=floor, r=5, s=(1+sqrt(5))/2, t=1/s.

For a guide to related sequences and a conjecture, see A206911. Adamczewski provides a proof abstract regarding the above conjecture in linked article. - _Michael De Vlieger_, Aug 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006810 | OEIS-A207673

n+[nr/s]+[nt/s], where []=floor, r=5, s=(1+sqrt(5))/2, t=1/s.

For a guide to related sequences and conjectures, see A206911. Adamczewski provides a proof abstract regarding the above conjecture in linked article. - _Michael De Vlieger_, Aug 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006811 | OEIS-A207700

Number of n X 5 0..1 arrays avoiding 0 0 0 and 0 1 0 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006812 | OEIS-A207701

Number of n X 6 0..1 arrays avoiding 0 0 0 and 0 1 0 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006813 | OEIS-A207702

Number of n X 7 0..1 arrays avoiding 0 0 0 and 0 1 0 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006814 | OEIS-A207820

Let A = A025584. a(n) is the smallest A(m) such that the interval (A(m)*n, A(m+1)*n) contains no primes from A.

Conjecture. For n>=3, every a(n) is the lesser of a pair of cousin primes p and p+4, cf. A023200.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006815 | OEIS-A207846

Number of 3 X n 0..1 arrays avoiding 0 0 0 and 0 0 1 horizontally and 0 0 0 and 1 1 1 vertically.

Conjectures from _Colin Barker_, Jun 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006816 | OEIS-A207872

Denominator of Z(n,1/2), where Z(n,x) is the n-th Zeckendorf polynomial.

Conjecture: a(n) is also the reverse binarization of the Zeckendorf representation of n in base Fibonacci. For example, 11 = 1x8 + 0x5 +1x3 +0x2 + 0x1, so 11 =10100 in base Fibonacci. Now read that as binary but in reverse, 00101 = 101 = 5 = A207873(11). - _George Beck_, Sep 02 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006817 | OEIS-A207935

Number of n X 5 0..1 arrays avoiding 0 0 1 and 1 0 0 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006818 | OEIS-A207936

Number of n X 6 0..1 arrays avoiding 0 0 1 and 1 0 0 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006819 | OEIS-A207937

Number of n X 7 0..1 arrays avoiding 0 0 1 and 1 0 0 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006820 | OEIS-A208034

G.f.: exp(Sum_{n>=1} 2*Pell(n)^2 * x^n/n), where Pell(n) = A000129(n).

Conjectures: For all positive integers k,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006821 | OEIS-A208044

Number of n X 3 0..3 arrays with new values 0..3 introduced in row major order and no element equal to any horizontal, vertical or antidiagonal neighbor (colorings ignoring permutations of colors).

Conjectures from _Colin Barker_, Mar 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006822 | OEIS-A208059

Start with n, successively subtract each digit of the resulting sequence (the digits of a negative term being the negatives of that term's digits): a(n) is the number of steps needed to get to the first zero.

As far as I know, it is only a conjecture that for any starting value n we always reach a 0. - _N. J. A. Sloane_, Jun 01 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hans Havermann_, Feb 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006823 | OEIS-A208060

a(n) = 1 + 2*n + 2^2*n*[n/2] + 2^3*n*[n/2]*[n/3] + 2^4*n*[n/2]*[n/3]*[n/4] + ... where [x]=floor(x).

Conjecture: limit a(n)^(1/n) = 2*L where L = 2.200161058099... is the geometric mean of Luroth expansions, where log(L) = Sum_{n>=1} log(n)/(n*(n+1)) = 0.7885305659115... (cf. A085361).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006824 | OEIS-A208064

Number of n X 3 0..1 arrays avoiding 0 0 0 and 1 1 1 horizontally and 0 0 1 and 0 1 1 vertically.

Conjectures from _Colin Barker_, Jun 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006825 | OEIS-A208065

Number of n X 4 0..1 arrays avoiding 0 0 0 and 1 1 1 horizontally and 0 0 1 and 0 1 1 vertically.

Conjectures from _Colin Barker_, Jun 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006826 | OEIS-A208079

Number of 4 X n 0..1 arrays avoiding 0 0 0 and 0 0 1 horizontally and 0 0 0 and 0 1 0 vertically.

Conjectures from _Colin Barker_, Jan 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006827 | OEIS-A208086

Number of 4 X (n+1) 0..1 arrays with every 2 X 2 subblock having the same number of equal edges as its horizontal neighbors and a different number from its vertical neighbors, and new values 0..1 introduced in row major order.

Conjectures from _Colin Barker_, Jun 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006828 | OEIS-A208087

Number of 6 X (n+1) 0..1 arrays with every 2 X 2 subblock having the same number of equal edges as its horizontal neighbors and a different number from its vertical neighbors, and new values 0..1 introduced in row major order.

Conjectures from _Colin Barker_, Jun 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006829 | OEIS-A208088

Number of 7 X (n+1) 0..1 arrays with every 2 X 2 subblock having the same number of equal edges as its horizontal neighbors and a different number from its vertical neighbors, and new values 0..1 introduced in row major order.

Conjectures from _Colin Barker_, Jun 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006830 | OEIS-A208089

Number of 8 X (n+1) 0..1 arrays with every 2 X 2 subblock having the same number of equal edges as its horizontal neighbors and a different number from its vertical neighbors, and new values 0..1 introduced in row major order.

Conjectures from _Colin Barker_, Jun 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006831 | OEIS-A208114

Number of n X 4 0..1 arrays avoiding 0 0 0 and 0 1 0 horizontally and 0 0 1 and 0 1 1 vertically.

Conjectures from _Colin Barker_, Jun 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006832 | OEIS-A208138

Number of n X 4 0..1 arrays avoiding 0 0 1 and 0 1 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006833 | OEIS-A208139

Number of n X 5 0..1 arrays avoiding 0 0 1 and 0 1 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 28 2018: (Start)
Conjectures from _Amiram Eldar_, Jun 16 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006834 | OEIS-A208140

Number of n X 6 0..1 arrays avoiding 0 0 1 and 0 1 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006835 | OEIS-A208141

Number of n X 7 0..1 arrays avoiding 0 0 1 and 0 1 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jun 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006836 | OEIS-A208229

Consider Wolfram's universal 2-state 3-symbol Turing machine on a one-way-infinite tape with all but the first n cells initially blank and the head initially in state 1. a(n) is the maximum number of steps the Turing machine can make before halting.

For initial conditions with n<=11, the machine always halts (the Mathematica code given assumes that the machine halts for all finite configurations). Whether this remains true is an open question.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ben Branman_, Jan 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006837 | OEIS-A208243

Number of ways to write 2n-1 = p+q, where p is a prime, and both q and q+2 are practical numbers (A005153).

Conjecture: a(n)>0 for all n=3,4,... Zhi-Wei Sun also made the following similar conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006838 | OEIS-A208244

Number of ways to write n as the sum of a practical number (A005153) and a triangular number (A000217).

Conjecture: a(n)>0 for all n>0. Somu and Tran (2024) proved the conjecture that a(n)>0 for n>0. - _Duc Van Khanh Tran_, Apr 24 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006839 | OEIS-A208246

Number of ways to write n = p+q with p prime or practical, and q-4, q, q+4 all practical.

Conjecture: a(n)>0 for all n>8. Zhi-Wei Sun also made some similar conjectures, below are few examples.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006840 | OEIS-A208249

Number of primes p with n < p < 2n, such that p-1 and p+1 are both practical.

Conjecture: a(n)>0 for all n>8. Zhi-Wei Sun also made the following conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006841 | OEIS-A208253

Number of n X 3 0..2 arrays with new values 0..2 introduced in row major order and no element equal to more than one of its immediate leftward or upward neighbors.

Conjectures from _Colin Barker_, Jun 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006842 | OEIS-A208277

Smallest number of multiplicative persistence n in factorial base.

a(n) exists for all n, unlike (conjecturally) its decimal equivalent A003001. In particular, with k = a(n-1), a(n) <= k * k! + (k-1)! + ... + 2! + 1! < (a(n-1)+1)! for n > 1. Diamond & Reidpath ask if this upper bound can be improved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Feb 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006843 | OEIS-A208283

Number of n X 4 0..1 arrays avoiding 0 0 0 and 0 0 1 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006844 | OEIS-A208284

Number of n X 5 0..1 arrays avoiding 0 0 0 and 0 0 1 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006845 | OEIS-A208285

Number of n X 6 0..1 arrays avoiding 0 0 0 and 0 0 1 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006846 | OEIS-A208286

Number of n X 7 0..1 arrays avoiding 0 0 0 and 0 0 1 horizontally and 0 0 1 and 1 1 0 vertically.

Conjectures from _Colin Barker_, Jun 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006847 | OEIS-A208296

Smallest positive nontrivial odd solution of the congruence x^2 == 1 (mod A001748(n+2)), n >= 1.

a(n) solves also the congruence x^2 == 1 (Modd A001748(n+2)), n>=1. For Modd n (not to be confused with mod n) see a comment on A203571. This follows from floor(a(n)^2/3*prime(n+2)) being even, in fact it is 8*A024699(n) (see a comment there), hence a(n)^2 (Modd 3*prime(n+2)) = a(n)^2 (mod 3*prime(n+2)) = 1. For those multiplicative groups Modd 3*p with p an odd prime which are cyclic (this is not possible in the mod case, see A033949), a(n) is the representative of the only other nontrivial solution of this congruence. The representative of the trivial solution is 1 (-1 belongs to the same Modd class). (The conjecture stated here earlier is wrong, that is, the multiplicative group Modd (91=7*13) is non-cyclic. It may still be true for 3*p. - _Wolfdieter Lang_, Mar 15 2012)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Mar 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006848 | OEIS-A208309

Number of n X 3 0..1 arrays with new values 0..1 introduced in row major order and no element equal to more than one of its immediate leftward or upward neighbors.

Conjectures from _Colin Barker_, Jun 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006849 | OEIS-A208316

Number of n X 2 0..2 arrays with new values 0..2 introduced in row major order and no element equal to more than two of its immediate leftward or upward or left-upward diagonal neighbors.

Conjectures from _Colin Barker_, Jul 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006850 | OEIS-A208327

Position of f(n) when the numbers f(j) and g(k) are jointly ranked, where f(j)=j + |cos j | and g(k)=k + |sin k|.

For a guide to related sequences and a conjecture, see A206911.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006851 | OEIS-A208341

Triangle read by rows, T(n,k) = hypergeometric_2F1([n-k+1, -k], [1], -1) for n>=0 and k>=0.

Conjecture: T(n, k) * 2^(n-2*k) = T(n, n-k). - _Werner Schulte_, Mar 20 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006852 | OEIS-A208342

Triangle of coefficients of polynomials u(n,x) jointly generated with A208343; see the Formula section.

T(n,k) = Sum_{j=0..floor((k-1)/2)} binomial(k-1-j,j)*binomial(n-k+j,j) for k,n>0 and k<=n (conjectured). - _Werner Schulte_, Mar 07 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006853 | OEIS-A208354

Number of compositions of n with at most one even part.

Conjecture: a(n) is the number of compositions of n if all the 1's are constrained to be in a single run; for example, a(7) counts the compositions 4,1,1,1 and 1,1,1,4 but not the compositions 1,4,1,1 and 1,1,4,1. - _Gregory L. Simay_, Sep 29 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Feb 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006854 | OEIS-A208355

Right edge of the triangle in A208101.

Conjecture: -(n+3)*(n-2)*a(n) - 4*a(n-1) + 4*(n-1)^2*a(n-2) = 0. - _R. J. Mathar_, Aug 04 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Mar 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006855 | OEIS-A208375

Number of n X 4 0..1 arrays avoiding 0 0 0 and 1 1 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jul 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006856 | OEIS-A208376

Number of n X 5 0..1 arrays avoiding 0 0 0 and 1 1 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jul 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006857 | OEIS-A208377

Number of n X 6 0..1 arrays avoiding 0 0 0 and 1 1 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jul 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006858 | OEIS-A208378

Number of n X 7 0..1 arrays avoiding 0 0 0 and 1 1 1 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jul 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006859 | OEIS-A208393

Number of 2 X n 0..2 arrays with new values 0..2 introduced in row major order and no element equal to more than two of its immediate leftward or upward or right-upward antidiagonal neighbors.

Conjectures from _Colin Barker_, Jul 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006860 | OEIS-A208402

Number of n X 2 0..3 arrays with new values 0..3 introduced in row major order and no element equal to more than two of its immediate leftward or upward or right-upward antidiagonal neighbors.

Conjectures from _Colin Barker_, Jul 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006861 | OEIS-A208403

Number of n X 3 0..3 arrays with new values 0..3 introduced in row major order and no element equal to more than two of its immediate leftward or upward or right-upward antidiagonal neighbors.

Conjectures from _Colin Barker_, Jul 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006862 | OEIS-A208425

Expansion of Sum_{n>=0} (3*n)!/n!^3 * x^(2*n)/(1-x)^(3*n+1).

Conjecture: (i) For any prime p > 3 and positive integer n, the number (a(p*n)-a(n))/(p*n)^3 is always a p-adic integer. We have proved part (i) of this conjecture for n = 1. (End) Conjecture: n^2*(3*n-5)*a(n) +(-9*n^3+24*n^2-17*n+4) *a(n-1) -(3*n-4) *(24*n^2-56*n+27)*a(n-2) -(3*n-2)*(n-2)^2*a(n-3)=0. - _R. J. Mathar_, Mar 10 2016 It is easy to show that a(n) = Sum_{k=0..n}C(n,k)*C(n-k,k)*C(n+k,k) = Sum_{k=0..n}C(n+k,k)*C(n,2k)*C(2k,k). By this formula and the Zeilberger algorithm, we confirm the recurrence conjectured by _R. J. Mathar_. - _Zhi-Wei Sun_, Nov 12 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006863 | OEIS-A208428

Number of n X 2 0..2 arrays with new values 0..2 introduced in row major order and no element equal to any knight-move neighbor (colorings ignoring permutations of colors).

Conjectures from _Colin Barker_, Feb 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006864 | OEIS-A208494

Least integer m>1 such that those k! mod m with k=1,...,n are pairwise distinct.

On Feb 27 2012 Zhi-Wei Sun conjectured that a(n) is a prime with the only exception a(5)=10, and that a(n) does not exceed n^2/2 for all n=2,3,4,... He also conjectured that max{n>0: 1!,...,n! are pairwise incongruent mod p} is asymptotically equivalent to sqrt(p), where p is an odd prime. He guessed that if we replace k! in the definition of a(n) by (-1)^k*k! then a(n) is a prime with the only exception a(3)=6. If we replace k! in the definition of a(n) by (2k)! or (-1)^k*(2k)!, then Zhi-Wei Sun conjectured that a(n) will take only prime values. He also has similar conjectures involving (r*k)! or (-1)^k*(r*k)! with r>2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006865 | OEIS-A208502

Number of 4 X n 0..1 arrays avoiding 0 0 0 and 0 0 1 horizontally and 0 1 0 and 1 1 1 vertically.

Conjectures from _Colin Barker_, Jul 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006866 | OEIS-A208531

Number of distinct sums of subsets of the first n squares {1,4,9,...,n^2}.

Conjectures from _Colin Barker_, Feb 15 2016: (Start) For a proof of these conjectures see the Suzuki (2019) link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Feb 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006867 | OEIS-A208548

a(n) = floor((4/3)*(2+prime(n)) - prime(n+1)).

Conjecture: a(n)>0 for all n (cf. A062234, A207480).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Feb 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006868 | OEIS-A208556

Number of 4 X n 0..1 arrays avoiding 0 0 0 and 0 0 1 horizontally and 0 0 1 and 0 1 1 vertically.

Conjectures from _Colin Barker_, Mar 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006869 | OEIS-A208561

Number of n X 2 0..2 arrays with new values 0..2 introduced in row major order and no element equal to more than one of its immediate leftward or upward or left-upward diagonal neighbors.

Conjectures from _Colin Barker_, Jul 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006870 | OEIS-A208598

Number of 4-bead necklaces labeled with numbers -n..n not allowing reversal, with sum zero.

Conjectures from _Colin Barker_, Mar 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006871 | OEIS-A208599

Number of 5-bead necklaces labeled with numbers -n..n not allowing reversal, with sum zero.

Conjectures from _Colin Barker_, Mar 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006872 | OEIS-A208600

Number of 6-bead necklaces labeled with numbers -n..n not allowing reversal, with sum zero.

Conjectures from _Colin Barker_, Mar 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006873 | OEIS-A208601

Number of 7-bead necklaces labeled with numbers -n..n not allowing reversal, with sum zero.

Conjectures from _Colin Barker_, Mar 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006874 | OEIS-A208634

Number of n X 5 0..1 arrays with new values 0..1 introduced in row major order and no element equal to more than one of its immediate leftward or upward or right-upward antidiagonal neighbors.

Conjectures from _Colin Barker_, Jul 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006875 | OEIS-A208638

Number of 3 X n 0..1 arrays with new values 0..1 introduced in row major order and no element equal to more than one of its immediate leftward or upward or right-upward antidiagonal neighbors.

Conjecture: a(n) is equal to half the sum along the edges of (centered, height 2, width n, starting at line n+1) rectangles in Pascal's triangle, as shown here for n=3 (not including the terms inside the rectangles): Conjectures from _Colin Barker_, Mar 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006876 | OEIS-A208639

Number of 4 X n 0..1 arrays with new values 0..1 introduced in row major order and no element equal to more than one of its immediate leftward or upward or right-upward antidiagonal neighbors.

Conjectures from _Colin Barker_, Jul 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006877 | OEIS-A208640

Number of 5 X n 0..1 arrays with new values 0..1 introduced in row major order and no element equal to more than one of its immediate leftward or upward or right-upward antidiagonal neighbors.

Conjectures from _Colin Barker_, Jul 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006878 | OEIS-A208641

Number of 6 X n 0..1 arrays with new values 0..1 introduced in row major order and no element equal to more than one of its immediate leftward or upward or right-upward antidiagonal neighbors.

Conjectures from _Colin Barker_, Jul 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006879 | OEIS-A208642

Number of 7 X n 0..1 arrays with new values 0..1 introduced in row major order and no element equal to more than one of its immediate leftward or upward or right-upward antidiagonal neighbors.

Conjectures from _Colin Barker_, Jul 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006880 | OEIS-A208658

Row sums of A208657.

Conjectures from _Colin Barker_, Mar 20 2012: (Start) The above conjectures are true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006881 | OEIS-A208675

Number of words, either empty or beginning with the first letter of the ternary alphabet, where each letter of the alphabet occurs n times and letters of neighboring word positions are equal or neighbors in the alphabet.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Feb 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006882 | OEIS-A208679

Kashaev's invariant for the (5,2)-torus knot (Solomon's seal knot).

We make the following conjectures: Define F(q) := Sum_{m,n >= 0} (q^(-m*n)*product {i = 1.. m+n} (1-q^i)). For the expansion of F(1-q) and F(exp(-t)) see A208733 and A208730 respectively. Kitami gives the conjectural e.g.f. exp(-9*t)*F(exp(-40*t)) = 1 + 71*t + 14641*t^2/2! + ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006883 | OEIS-A208681

Kashaev's invariant for the (9,2)-torus knot.

We make the following conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006884 | OEIS-A208689

Number of 3 X n 0..1 arrays avoiding 0 0 0 and 0 0 1 horizontally and 0 1 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Mar 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006885 | OEIS-A208705

Number of n X 4 0..1 arrays with new values 0..1 introduced in row major order and no element equal to more than two of its immediate leftward or upward or right-upward antidiagonal neighbors.

Conjectures from _Colin Barker_, Jul 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006886 | OEIS-A208730

Sequence related to Kashaev's invariant for the (5,2)-torus knot.

E.g.f.: F(exp(-t)) = 1 + 2*t + 10*t^2! + 104*t^3/3! + ... For the expansion of F(1-q) see A208733. F(q) also appears in a conjectural e.g.f. for A208679. Conjectural S-fraction for the o.g.f.: 1/(1-2*x/(1-3*x/(1-9*x/(1-11*x/(1-...-1/2*n*(5*n-1)*x/(1-1/2*n*(5*n+1)*x/(1- ....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006887 | OEIS-A208731

Sequence related to Kashaev's invariant for the (7,2)-torus knot.

Conjectural S-fraction for the o.g.f.: 1/(1-3*x/(1-4*x/(1-13*x/(1-15*x/(1-...-1/2*n*(7*n-1)*x/(1-1/2*n*(7*n+1)*x/(1- ....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006888 | OEIS-A208732

Sequence related to Kashaev's invariant for the (9,2)-torus knot.

Conjectural S-fraction for the o.g.f.: 1/(1-4*x/(1-5*x/(1-17*x/(1-19*x/(1-...-1/2*n*(9*n-1)*x/(1-1/2*n*(9*n+1)*x/(1- ....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006889 | OEIS-A208826

Number of 5-bead necklaces labeled with numbers -n..n allowing reversal, with sum zero.

Conjectures from _Colin Barker_, Jul 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006890 | OEIS-A208827

Number of 6-bead necklaces labeled with numbers -n..n allowing reversal, with sum zero.

Conjectures from _Colin Barker_, Jul 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006891 | OEIS-A208850

Expansion of phi(q^2) / phi(-q) in powers of q where phi() is a Ramanujan theta function.

Conjecture: Sum_{k>=0} a(k) / exp(k*Pi) = (1/2)*2^(1/4)*(2+sqrt(2))^(1/2) = A309948. - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Mar 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006892 | OEIS-A208884

a(n) = (a(n-1) + n)/2^k where 2^k is the largest power of 2 dividing a(n-1) + n, for n>1 with a(1)=1.

Conjecture: For any given odd number m, there exists a number n_max such that all odd numbers <= m can be found in the sequence a(n) with n <= n_max. For example: If the conjecture above is true, this sequence contains all odd numbers. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006893 | OEIS-A208951

Twice the maximum of the areas of the convex hulls of permutations {(1,p(1)), (2,p(2)), ..., (n,p(n))} of {1, 2, ..., n}, considered as points in the plane.

Conjecture: a(n) = 2n^2 - 8n + 10 for n >= 4. - _Peter Kagey_, Oct 30 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Mar 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006894 | OEIS-A208971

Number of 4-bead necklaces labeled with numbers -n..n allowing reversal, with sum zero and first and second differences in -n..n.

Conjectures from _Colin Barker_, Jul 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006895 | OEIS-A208995

Number of 4-bead necklaces labeled with numbers -n..n not allowing reversal, with sum zero and first differences in -n..n.

Conjectures from _Colin Barker_, Jul 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006896 | OEIS-A209008

Number of 4-bead necklaces labeled with numbers -n..n not allowing reversal, with sum zero and first and second differences in -n..n.

Conjectures from _Colin Barker_, Jul 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006897 | OEIS-A209078

Conjectured list of values of n such that the Diophantine equation a^4 + b^4 = n*c^2 has infinitely many solutions with 0 < a < b.

Conjectured list of values of n such that the Diophantine equation a^4 + b^4 = n*c^2 has infinitely many solutions with 0 < a < b.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006898 | OEIS-A209094

Number of n X 2 0..2 arrays with new values 0..2 introduced in row major order and no element equal to more than one of its immediate leftward or upward or right-upward antidiagonal neighbors.

Conjectures from _Colin Barker_, Mar 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006899 | OEIS-A209095

Number of n X 3 0..2 arrays with new values 0..2 introduced in row major order and no element equal to more than one of its immediate leftward or upward or right-upward antidiagonal neighbors.

Conjectures from _Colin Barker_, Jul 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006900 | OEIS-A209236

List of integers m>0 with m-1 and m+1 both prime, and m-2, m, m+2 all practical.

Conjecture: a(n) always exists. In other words, there are infinitely many quintuples (m-2, m-1, m, m+1, m+2) with m-1 and m+1 both prime and m-2, m, m+2 all practical.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006901 | OEIS-A209237

Number of practical numbers not exceeding 2^n.

Motivated by Sun's conjecture concerning the asymptotics of A005153.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006902 | OEIS-A209253

Number of ways to write 2n-1 as the sum of a Sophie Germain prime and a practical number.

Conjecture: a(n)>0 for all n>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006903 | OEIS-A209254

Number of ways to write 2n-1 = p+q with q practical, p and p^4+q^4 both prime.

Conjecture: a(n)>0 for all n>1. Zhi-Wei Sun also conjectured that any odd integer greater than one can be written as p+q with q practical, and p and p^2+q^2 both prime. This is a refinement of Ming-Zhi Zhang's problem related to A036468.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006904 | OEIS-A209260

Irregular triangular array read by rows, resulting in a permutation of the natural numbers.

Conjecture 1. 2*n+1 is in the set S_n = {union of first n rows of A209260} if and only if 2*n+1 is composite. (Verified for all n <= 10^6 by _Charles R Greathouse IV_.) This conjecture has been proved (see [Jeffery]). Conjecture 2. Let N be an integer, N>0, and let D_N be the set of all positive integral divisors of N. Let B_N be the set of all integral solutions of d + (h-1)/2 for which d in D_N, with h=N/d and N=d*h (some solutions will not be integers and so are not in B_N). Then N appears in row r = min(B_N) of A209260. Also, N appears in row b of A141419, for each b in B_N. - _L. Edson Jeffery_, Feb 13 2013 For a proof of Conjecture 2 see the link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_ and _Charles R Greathouse IV_, Jan 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006905 | OEIS-A209266

a(n) is the number of 3-prime arithmetic progression prime chains surrounding the n-th prime number with 5-smooth intervals.

Based on the conjecture in A211376, a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Feb 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006906 | OEIS-A209312

Number of practical numbers p<n with n-p and n+p both prime or both practical.

Conjecture: a(n)>0 for all n>2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006907 | OEIS-A209332

a(n) is the minimal positive number k such that n<+>k is prime or 0 if no such number exists (operation <+> defined in A206853).

Conjecture: for even n, a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Mar 06 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006908 | OEIS-A209530

Half the number of (n+1) X 3 0..2 arrays with every 2 X 2 subblock having exactly two distinct clockwise edge differences.

Conjectures from _Colin Barker_, Jul 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006909 | OEIS-A209531

Half the number of (n+1) X 5 0..2 arrays with every 2 X 2 subblock having exactly two distinct clockwise edge differences.

Conjectures from _Colin Barker_, Jul 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006910 | OEIS-A209532

Half the number of (n+1) X 7 0..2 arrays with every 2 X 2 subblock having exactly two distinct clockwise edge differences.

Conjectures from _Colin Barker_, Jul 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006911 | OEIS-A209554

Primes that expressed in none of the forms n<+>2 and n<+>3, where the operation <+> is defined in A206853.

If the formulated below conjecture is true, then for n>=2, A209544 and this sequence coincide with A007519 and A133870 respectively. Conjecture (A209544 and this sequence correspond to cases k=2 and k=3 respectively). For k=2 this conjecture is true (see comment in A182187). - _Vladimir Shevelev_, Apr 18 2012.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Mar 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006912 | OEIS-A209647

Number of n X 5 0..1 arrays avoiding 0 0 1 and 1 0 0 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Mar 07 2018: (Start) Empirical formula (and thus Barker's conjectures) proved by _Robert Israel_, Mar 07 2018: see link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006913 | OEIS-A209648

Number of n X 6 0..1 arrays avoiding 0 0 1 and 1 0 0 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Jul 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006914 | OEIS-A209649

Number of n X 7 0..1 arrays avoiding 0 0 1 and 1 0 0 horizontally and 0 0 1 and 1 0 1 vertically.

Conjectures from _Colin Barker_, Mar 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006915 | OEIS-A209721

1/4 the number of (n+1) X 3 0..2 arrays with every 2 X 2 subblock having distinct clockwise edge differences.

Conjectures: a(n) = A029744(n+1)+1. Also, a(n) = positions of the zeros in A309019(n+2) - A002487(n+2). - _George Beck_, Mar 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006916 | OEIS-A209722

1/4 the number of (n+1) X 4 0..2 arrays with every 2 X 2 subblock having distinct clockwise edge differences.

Conjectures from _Colin Barker_, Jul 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006917 | OEIS-A209723

1/4 the number of (n+1) X 5 0..2 arrays with every 2 X 2 subblock having distinct clockwise edge differences.

Conjectures from _Colin Barker_, Jul 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006918 | OEIS-A209724

1/4 the number of (n+1) X 6 0..2 arrays with every 2 X 2 subblock having distinct clockwise edge differences.

Conjecture: a(1) = 8; for n > 1, a(n) is the smallest integer m such that m = ((2x * a(n-1)) /(x+1)) - x , with x a positive nontrivial divisor of m. (This is true at least for a(1) to a(100).) - _Enric Reverter i Bigas_, Oct 11 2020 Conjectures from _Colin Barker_, Mar 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006919 | OEIS-A209725

1/4 the number of (n+1) X 7 0..2 arrays with every 2 X 2 subblock having distinct clockwise edge differences.

Conjectures from _Colin Barker_, Jul 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006920 | OEIS-A209726

1/4 the number of (n+1) X 8 0..2 arrays with every 2 X 2 subblock having distinct clockwise edge differences.

Conjectures from _Colin Barker_, Jul 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006921 | OEIS-A209789

Half the number of (n+1) X 2 0..2 arrays with every 2 X 2 subblock having exactly one duplicate clockwise edge difference.

Conjectures from _Colin Barker_, Jul 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006922 | OEIS-A209802

Partial sums of exponential Möbius function, A166234.

Analog of Mertens's function, A002321; conjecture: a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Mar 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006923 | OEIS-A209805

Triangle read by rows: T(n,k) is the number of k-block noncrossing partitions of n-set up to rotations.

The above conjecture about the diagonal entries T(2*n-1, n) is true since gcd(2*n-1, n) = gcd(2*n-1, n-1) = 1 and then T(2*n-1, n) simplifies to A001246(n-1) using the formula given below. - _Andrew Howroyd_, Nov 15 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tilman Piesk_, Mar 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006924 | OEIS-A209832

Expansion of the q-series Sum_{n>=0} (-1)^n*q^(n+1)*Product_{k = 1..n} (1 - q^(2*k-1)), q = exp(t), as a formal Taylor series in t.

Conjectural S-fraction expansion for the o.g.f.:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006925 | OEIS-A209860

Fixed points of permutation A209861/A209862.

Conjecture: for every a(n), also A054429(a(n)) is in the sequence. Conversely, if i is not in the sequence, then neither is A054429(i).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006926 | OEIS-A209861

Permutation of nonnegative integers which maps A209641 to A209642.

Conjecture: For all n, A209861(A054429(n)) = A054429(A209861(n)), i.e. A054429 acts as an homomorphism (automorphism) of the cyclic group generated by this permutation. This implies also a weaker conjecture given in A209860.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006927 | OEIS-A209862

Permutation of nonnegative integers which maps A209642 into ascending order (A209641).

Conjecture: For all n, a(A054429(n)) = A054429(a(n)), i.e. A054429 acts as a homomorphism (automorphism) of the cyclic group generated by this permutation. This implies also a weaker conjecture given in A209860.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006928 | OEIS-A209863

a(n) = Number of fixed points of permutation A209861/A209862 in range [2^(n-1),(2^n)-1].

See the conjecture given in A209860. If true, then all the terms from a(2) onward are even. a(0) gives the number of fixed points in range [0,0], i.e. 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006929 | OEIS-A209933

Numbers that are divisible by all digits of their divisors.

There are 24 terms < 10000. Conjecture: next term a(25) = prime repunit with 19 digits 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Apr 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006930 | OEIS-A209934

a(n) is the first value to occur consecutively in the sequence b_n defined by p_2k(b_n(k)) = p_k(n)^2, k=1,2,3,..., where p_k(n) is the n-th k-almost prime.

Conjecture: Each sequence b_n repeats indefinitely. (Example: for n=3, b_n = 9, 8, 8, 8, 8, 8, ... It looks like b_3(k) is 8 for all k > 1.) The conjecture follows from the formula that uses A078843 below (and the strict monotonicity of A078843). However the first repeated value is not for every n the value that repeats indefinitely. For example a(8) = b_8(2) = b_8(3) = 66, but b_8(k) = 64 for k >= 4. - _Peter Munn_, Aug 05 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Tisdale_, Mar 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006931 | OEIS-A209944

a(n) is half the number of (n+1) X 2 0..2 arrays with every 2 X 2 subblock having at most one duplicate clockwise edge difference.

Conjectures from _Colin Barker_, Jul 14 2018: (Start) Conjectured e.g.f.: exp(3*x)*(189*cosh(sqrt(21)*x) + 41*sqrt(21)*sinh(sqrt(21)*x))/42 - 9/2. - _Stefano Spezia_, Aug 02 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006932 | OEIS-A210054

Number of (n+1) X 2 0..3 arrays with every 2 X 2 subblock having one, two or four distinct values, and new values 0..3 introduced in row major order.

Conjectures from _Colin Barker_, Jul 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006933 | OEIS-A210064

Total number of 231 patterns in the set of permutations avoiding 123.

Conjecture: n*(n-3)*a(n) +2*(-4*n^2+11*n-2)*a(n-1) +8*(n-1)*(2*n-3)*a(n-2)=0. - _R. J. Mathar_, Oct 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cheyne Homberger_, Mar 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006934 | OEIS-A210100

Number of (n+1) X 2 0..2 arrays with every 2 X 2 subblock having one or three distinct values, and new values 0..2 introduced in row major order.

Conjectures from _Colin Barker_, Jul 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006935 | OEIS-A210101

Number of (n+1) X 3 0..2 arrays with every 2 X 2 subblock having one or three distinct values, and new values 0..2 introduced in row major order.

Conjectures from _Colin Barker_, Jul 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006936 | OEIS-A210144

a(n) = least integer m>1 such that the product of the first k primes for k=1,...,n are pairwise distinct modulo m.

Conjecture: all the terms are primes and a(n)<n^2 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006937 | OEIS-A210186

a(n) = least integer m>1 such that m divides none of P_i + P_j with 0<i<j<=n where P_k is the product of the first k primes.

Conjecture: all the terms are primes and a(n) < n^2 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006938 | OEIS-A210210

Least number k among 1, ..., n such that pi(p+2*n) - pi(p+n) = pi(p+n) - pi(p) > 0 with p = prime(k), or 0 if such a number k does not exist, where pi(.) is given by A000720.

Conjecture: a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006939 | OEIS-A210245

Signs of the polylogarithm li(-n,-1/2).

Conjecture: a(n) is the same as the sign of cos((n+1)*arctan(Pi/log(2))). - _Mikhail Kurkov_, Apr 13 2021 [Holds for all n<=10000. - _Vaclav Kotesovec_, Mar 02 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Mar 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006940 | OEIS-A210247

a(n) = sign of the polylogarithm li(-n,-1/3) for n > 0, with a(0) = 1.

Conjecture: a(n) = -a(n-28). - _Mikhail Kurkov_, Aug 27 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Mar 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006941 | OEIS-A210374

Number of 2 X 2 matrices with all terms in {0,1,...,n} and (sum of terms) = n+2.

Conjectures from _Colin Barker_, Dec 07 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006942 | OEIS-A210376

Number of 2 X 2 matrices with all terms in {0,1,...,n} and (sum of terms) = n + 4.

Conjectures from _Colin Barker_, Dec 07 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006943 | OEIS-A210377

Number of 2 X 2 matrices with all terms in {0,1,...,n} and (sum of terms) = n + 5.

Conjectures from _Colin Barker_, Dec 07 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006944 | OEIS-A210380

Consider all n-tuples of distinct positive integers for which no two different elements add up to a square. This sequence gives the smallest maximal integer in such tuples.

a(n) <= (32/11)*n - 2. Erdős conjectured that a(n) >= (32/11)*n - k for some fixed k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Mar 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006945 | OEIS-A210393

a(n) = least integer m>1 such that S_k! for k=1,...,n are pairwise distinct modulo m where S_k is the sum of the first k primes.

When n>1, we have S_n!=S_{n-1}!=0 (mod m) for all m=1,...,S_{n-1} and hence a(n)>S_{n-1}. Zhi-Wei Sun conjectured that a(n) is always a prime not exceeding S_n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006946 | OEIS-A210406

Number of (n+1) X 2 0..3 arrays with every 2 X 2 subblock having three or four distinct values, and new values 0..3 introduced in row major order.

Conjectures from _Colin Barker_, Jul 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006947 | OEIS-A210444

a(n) = |{0<k<n: k*n is practical, k*n-1 and k*n+1 are twin primes}|.

Conjecture: a(n)>0 for all n>911. The conjecture has been verified for n up to 10^6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006948 | OEIS-A210445

Least positive integer k with k*n practical.

Conjecture: a(n) < n for all n>1, and a(n) < n/2 for all n>47.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006949 | OEIS-A210456

Period of the sequence of the digital roots of Fibonacci n-step numbers.

Conjecture: All entries >1 are divisible by 3. Conjecture: a(3^n-1)=3^(2*n+1)-3, a(3^n)=3^(2*n+1)+3^(n+1)+3 - Fred W. Helenius (fredh(AT)ix.netcom.com), posting to MathFun, Feb 21 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_, _Robert G. Wilson v_, Jan 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006950 | OEIS-A210465

Let p_(3,1)(m) be the m-th prime == 1(mod 3). Then a(n) is the smallest p_(3,1)(m) such that the interval(p_(3,1)(m)*n, p_(3,1)(m+1)*n) contains exactly one prime == 1(mod 3).

Conjectures: (1) If q is the nearest prime>a(n), then q-a(n)=4 or 6 and both of these cases occur infinitely many times. (2) If q-a(n)=4 then q is the lesser of twin primes. Thus, if the conjectures are true, then there exist infinitely many triples of primes of the form {p,p+4,p+6}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006951 | OEIS-A210467

Let p_(3,2)(m) be the m-th prime == 2(mod 3). Then a(n) is the smallest p_(3,2)(m) such that the interval(p_(3,2)(m)*n, p_(3,2)(m+1)*n) contains exactly one prime == 2 (mod 3).

Conjectures: (1) If q is the nearest prime > a(n), then q-a(n) = 2 or 6 and both of these cases occur infinitely many times. (2) If q-a(n) = 2, then also q is lesser of a pair of cousin primes q and q+4, see A023200. Thus, if the conjectures are true, then there exist infinitely many triples of primes of the form {p,p+2,p+6}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006952 | OEIS-A210468

Collatz (3x+1) problem with rational numbers: number of steps to reach the end of the cycle starting with 1/(2n+1).

This variation of the "3x+1" problem with a class of rational numbers is as follows: start with any number 1/(2n+1). If the numerator is even, divide it by 2, otherwise multiply it by 3 and add 1. Do we always reach the end of a cycle with a rational number? It is conjectured that the answer is yes. If x is of the form x = m/(2n+1) where m is an integer, it is conjectured that the number of steps tends into the end of a finite cycle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006953 | OEIS-A210474

The number of different lattice paths from (0,0) to (2n,0) using steps of S={(i,i) or (i,-i): i=1,2,...,n} with j flaws(j=1,2,...,n-1), where the j flaws is the sum of lengths of down steps below the x-axis. (For down steps that are partly above and partly below the x-axis we just count the part below the x-axis.) This number is independent of the number of flaws.

Conjecture: (n+1)*a(n) +2*(-5*n+3)*a(n-1) +9*(n-3)*a(n-2)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jisun Huh_, Jan 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006954 | OEIS-A210475

Let p_(4,1)(m) be the m-th prime == 1 (mod 4). Then a(n) is the smallest p_(4,1)(m) such that the interval(p_(4,1)(m)*n, p_(4,1)(m+1)*n) contains exactly one prime == 1 (mod 4).

Conjecture: for n >= 12, every a(n) is the lesser of a pair of cousin primes p and p+4, (see A023200).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006955 | OEIS-A210476

Let p_(4,3)(m) be the m-th prime == 3 (mod 4). Then a(n) is the smallest p_(4,3)(m) such that the interval(p_(4,3)(m)*n, p_(4,3)(m+1)*n) contains exactly one prime == 3(mod 4).

Conjecture: every a(n), except for a(7) = 191, is the lesser of a pair of cousin primes p and p+4, (see A023200).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006956 | OEIS-A210484

Ordered areas of primitive integer Soddyian triangles.

A Soddyian triangle is a triangle whose outer Soddy circle has degenerated into a straight line. If it is assumed that the sides a<=b<=c then, 1/Sqrt(s-c) = 1/Sqrt(s-a)+1/Sqrt(s-b) where s is the semiperimeter. All integer Soddyian triangles are Heronian. It is conjectured that a(n) has no multiplicities - checked to a(21886129).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Jan 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006957 | OEIS-A210496

Number of set partitions of [n] avoiding the patterns {1123, 1211}.

Conjecture: (n+2)*a(n) +2*(-3*n-4)*a(n-1) +4*(2*n+3)*a(n-2) +(13*n-40)*a(n-3) +5*(-7*n+18)*a(n-4) +6*(2*n-1)*a(n-5) +22*(n-7)*a(n-6) +(-19*n+134)*a(n-7) +2*(2*n-15)*a(n-8)=0. - _R. J. Mathar_, Oct 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jan 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006958 | OEIS-A210505

Numbers k for which 2*k+7, 4*k+7, 6*k+7, 8*k+7, 10*k+7 and 12*k+7 are primes.

Conjecture. For every odd prime p there exist infinitely many numbers k for which 2*k+p, 4*k+p, ..., 2*(p-1)*k+p are primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006959 | OEIS-A210527

a(n) = 9*n^2 + 39*n + 83.

Conjecture: the total number of primes remains below one-half of the total number of terms generated from and after n = 886. - _Harvey P. Dale_, Mar 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tony Herrys Silva Rabelo_, Jan 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006960 | OEIS-A210528

Number of ways to write 2n = p+q (p<=q) with p, q and p^6+q^6 all practical.

Conjecture: a(n)>0 for all n>0. Moreover, for any positive integers m and n, if m is greater than one or n is not among 17, 181, 211, 251, 313, 337, then 2n can be written as p+q with p, q and p^{3m}+q^{3m} all practical. This conjecture implies that for each m=1,2,3,... there are infinitely many practical numbers of the form p^{3m}+q^{3m} with p and q both practical.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006961 | OEIS-A210531

Number of nonnegative integers k<n with n+k^3 practical.

Conjecture: a(n)>0 for all n>0. Moreover, if n>0 is different from 74, 138, 166, 542, then n+k^3 is practical for some 0<=k<=sqrt(n)*log(n); if n is not equal to 102, then n+k and n+k^3 are both practical for some k=0,...,n-1. Zhi-Wei Sun also conjectured that any integer n>1 can be written as x^3+y (x,y>0) with 2x and 4xy both practical.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006962 | OEIS-A210533

Number of ways to write 2n = x+y (x,y>0) with x-1 and x+1 both prime, and x and x^3+y^3 both practical.

Conjecture: a(n)>0 for all n>2. Moreover, for each m=2,3,4,... any sufficiently large even integer can be written as x+y (x,y>0) with x-1 and x+1 both prime, and x and x^m+y^m both practical.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006963 | OEIS-A210537

a(1)=3; for n>1, a(n)>a(n-1) is the minimal for which the set {a(1),a(2),...,a(n)} lacks at least one residue mod 2, 3, ...

Conjectures: (1) the sequence contains infinitely many "twins" when such differences equal 6; (2) lim a(n)/prime(n)=1 as n goes to infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jan 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006964 | OEIS-A210570

Consider all sequences of n distinct positive integers for which no two different elements have a difference which is a square. This sequence gives the smallest maximal integer in such sequences.

László Lovász conjectured, and Hillel Furstenberg and András Sárközy (1978) independently showed that a(n) is superlinear. Erdős conjectured that a(n) >> n^2/log^k n for some k. Sárközy proved that a(n) = o(n^2/log^k n) for all k, but still conjectured that a(n) >> n^(2-e) for all e > 0. Ruzsa showed that in fact a(n) << n^1.365.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Apr 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006965 | OEIS-A210576

Positive integers that cannot be expressed as sum of one or more nontrivial binomial coefficients.

I conjectured that the sequence is finite, consisting of the terms listed. This conjecture is now proved. - _Douglas Latimer_, Apr 10 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Douglas Latimer_, Mar 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006966 | OEIS-A210627

Constants r_n arising in study of polynomials of least deviation from zero in several variables.

Conjecture: a(n) = n*A038049(n). - _R. J. Mathar_, Mar 27 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006967 | OEIS-A210640

a(n) = least integer m > 1 such that 2S_k^2 (k=1,...,n) are pairwise distinct modulo m, where S_k is the sum of the first k primes.

If a(n) is an odd prime p_k then a(n)|2(S_k^2-S_{k-1}^2) and hence k>n. Zhi-Wei Sun conjectured that a(n) is a prime smaller than n^2 unless n divides 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006968 | OEIS-A210645

Area A of the triangles such that A, the sides and one of the altitudes are four consecutive integers of an arithmetic progression d.

Conjecture: a(n) = 84*n^2. a(n) = 3*a(n-1)-3*a(n-2)+a(n-3). G.f.: 84*x*(1+x)/(1-x)^3. - _Colin Barker_, Apr 19 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Mar 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006969 | OEIS-A210650

Least number k>0 such that at least one of (2^k-1)*2^n-1, (2^k-1)*2^n+1, (2^k+1)*2^n-1, or (2^k+1)*2^n+1 is a prime number.

Conjecture: for n>1 a(n) is always < n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Mar 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006970 | OEIS-A210721

a(n) is the smallest trial division companion for the n-th prime number.

Conjecture: a(n) < prime(n)^2*(Pi^(-2Pi)+log((prime(n)+1)/prime(n)))

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Jan 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006971 | OEIS-A210722

Number of ways to write n = (2-(n mod 2))p+q+2^k with p, q-1, q+1 all prime, and p-1, p+1, q all practical.

Conjecture: a(n)>0 except for n = 1,...,8, 10, 520, 689, 740.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006972 | OEIS-A211008

Triangle read by rows: T(n,k) = number of squares and rectangles of area 2^(k-1) after n-th stage in the toothpick structure of A139250, n>=1, k>=1, assuming the toothpicks have length 2.

It appears that the number of rectangles of area 2 in the toothpick structure of A139250 equals the number of hearts in the Q-toothpick cellular automaton of A187210. See conjecture in formula section.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Sep 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006973 | OEIS-A211163

Numerator of (-1/Pi^n) * integral_{0..1} (log(1-1/x)^n) dx.

Conjecture: sequence of denominators is A141459.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Jan 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006974 | OEIS-A211165

Number of ways to write n as the sum of a prime p with p-1 and p+1 both practical, a prime q with q+2 also prime, and a Fibonacci number.

Conjecture: a(n)>0 for all n>5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006975 | OEIS-A211175

Triangle read by rows: row n gives, in increasing order, the prime divisors of all the composites of the form k^2 + 1 between the two primes A002496(n) and A002496(n+1).

A variety of conjecturally infinite subsequences and starting with {2, 5, ...} can be shown in the graph of the sequence. If the number of primes of the form n^2 + 1 is finite, then the last subsubsequence of the graph abruptly becomes A002144(n) union {2} (odd Pythagorean primes with the number 2). In this case, the discontinued forms of the graph disappear. But this case is highly improbable.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Feb 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006976 | OEIS-A211189

Number of prime divisors formed by {2} and the consecutive Pythagorean primes for all the composites k^2 + 1 between the two primes A002496(n) and A002496(n+1).

These considerations confirm the opinion of the truthfulness of the conjecture about an infinity of primes of the form n^2 + 1. This sequence gives the length of a variety of conjecturally infinite subsequences of consecutive primes starting with {2, 5, ...}. If the number of primes of the form n^2 + 1 were finite, there should exist a last prime p such that this sequence stops abruptly from p because the length of A002144(n) is infinite. In this case, we should observe a contradictory behavior of this sequence between the stability of the slow growth of a(n) and the discontinuity from the prime p. But this case is highly improbable.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Feb 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006977 | OEIS-A211190

Number of ways to write 2n = p+2q+3r with p,q,r terms of A210479.

Conjecture: a(n)>0 for all n>8. Moreover, for positive integers a<=b<=c, all integers n>=3(a+b+c) with n-a-b-c even can be written as a*p+b*q+c*r with p,q,r terms of A210479, if and only if (a,b,c) is among the following 6 triples: (1,2,3), (1,2,4), (1,2,8), (1,2,9), (1,3,5), (1,3,8). The author also conjectured that if n>8 is odd, different from 201 and 447, and not congruent to 1 or -1 modulo 12, then n can be written as a sum of three terms of A210479.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006978 | OEIS-A211193

E.g.f.: exp((1+x)^(1+x)-1).

Conjecture: For n > 1, a(n) == 1 (mod n). - _Mélika Tebni_, Aug 22 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Feb 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006979 | OEIS-A211200

Indices of records in A207709.

Conjecture: the sequence contains only noncomposite numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Apr 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006980 | OEIS-A211276

a(n) = number of n-lettered words in the alphabet {1, 2, 3} with as many occurrences of the substring (consecutive subword) [1, 1] as of [1, 2].

Conjecture: n*a(n) +2*(-3*n+1)*a(n-1) +(7*n+6)*a(n-2) +2*(7*n-37)*a(n-3) +3*(-7*n+40)*a(n-4) +6*(-n-4)*a(n-5) +27*(-n+6)*a(n-6) +54*(n-6)*a(n-7)=0. - _R. J. Mathar_, May 31 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006981 | OEIS-A211278

a(n) is the number of n-lettered words in the alphabet {1, 2, 3} with as many occurrences of the substring (consecutive subword) [1, 1] as of [2, 3].

Conjecture: n*a(n) - 4*n*a(n-1) +2*(n+3)*a(n-2) + 4*(n-3)*a(n-3) + 3*(-n+2)*a(n-4) = 0. - _R. J. Mathar_, Jun 09 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006982 | OEIS-A211309

a(n) = number |fdw(P,(n))| of entangled P-words with s=2.

Conjectural e.g.f.: 2 - 1/A(x), where A(x) = Sum_{n >= 0} (2*n)!/2^n*x^n/n! is the e.g.f. for A000680 (also the o.g.f. for A001147).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006983 | OEIS-A211310

a(n) = number |fdw(P,(n))| of entangled P-words with s=3.

Conjectural e.g.f.: 2 - 1/A(x), where A(x) = Sum_{n >= 0} (3*n)!/6^n*x^n/n! is the e.g.f. for A014606 (also the o.g.f. for A025035).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006984 | OEIS-A211311

a(n) = number |fdw(P,(n))| of entangled P-words with s=4.

Conjectural e.g.f.: 2 - 1/A(x), where A(x) = Sum_{n >= 0} (4*n)!/24^n*x^n/n! is the e.g.f. for A014608 (also the o.g.f. for A025036).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006985 | OEIS-A211322

Number of (n+1) X (n+1) -3..3 symmetric matrices with every 2 X 2 subblock having sum zero and one or two distinct values.

Conjectures from _Colin Barker_, Jul 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006986 | OEIS-A211327

Number of (n+1) X (n+1) -3..3 symmetric matrices with every 2 X 2 subblock having sum zero and one, three or four distinct values.

Conjectures from _Colin Barker_, Jul 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006987 | OEIS-A211337

Numbers k for which the number of divisors, tau(k), is congruent to 1 modulo 3.

The numbers of terms not exceeding 10^k, for k = 1, 2, ..., are 4, 36, 366, 3635, 36499, 365456, 3654240, 36538501, 365382167, 3653804173, ... Conjecture: the asymptotic density of this sequence exists and equals 3*zeta(3)/Pi^2 = 0.3653814847007... (A346602), so, a(n) ~ k*n with k = Pi^2/(3*zeta(3)) = 2.73686555524... . This conjecture is true if this sequence and A211338 have the same density (see A059269). - _Amiram Eldar_, Jan 06 2024 Conjecture: a(n) ~ k*n where k = 2/prod(1 - (p-1)/(p^(3*k))) = 2.7290077... where p ranges over the primes and k ranges over the positive integers. - _Charles R Greathouse IV_, Apr 13 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Douglas Latimer_, Apr 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006988 | OEIS-A211338

Numbers k for which the number of divisors, tau(k), is congruent to 2 modulo 3.

The numbers of terms not exceeding 10^k, for k = 1, 2, ..., are 4, 37, 368, 3681, 36596, 365336, 3653499, 36537962, 365381169, 3653826361, ... Conjecture: the asymptotic density of this sequence exists and equals 3*zeta(3)/Pi^2 = 0.3653814847007... (A346602), so, a(n) ~ k*n with k = Pi^2/(3*zeta(3)) = 2.73686555524... . This conjecture is true if this sequence and A211337 have the same density (see A059269). - _Amiram Eldar_, Jan 06 2024 Conjecture: a(n) ~ k*n where k = 2/prod(1 - (p-1)/(p^(3*k))) = 2.7290077... where p ranges over the primes and k ranges over the positive integers. - _Charles R Greathouse IV_, Apr 13 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Douglas Latimer_, Apr 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006989 | OEIS-A211343

Triangle read by rows: T(n,k), n >= 1, k >= 1, in which column k lists the positive integers interleaved with k-1 zeros, and the first element of column k is in row k(k+1)/2.

Conjecture 1: T(n,k) is the smallest part of the partition of n into k consecutive parts, if T(n,k) > 0. Conjecture 2: the last positive integer in the row n is in the column A109814(n). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Feb 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006990 | OEIS-A211376

a(n) is the smallest 5-smooth number k such that both prime(n) - k and prime(n) + k are prime.

Conjecture: a(n) is defined for all integers n > 2. Conjecture confirmed true up to n = 300000, no exceptions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Feb 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006991 | OEIS-A211419

a(n) = (6*n)!*(2*n)!/((4*n)!*(3*n)!*n!).

It appears that 5*a(n)/(n + 1) and a(n)/(4*n + 1) are integers for all n. More generally, we conjecture that there are constants C(r) > 0 and D(r) > 0, r >= 1, such that a(n) * C(r)/((n + 1)*(n + 2)*...*(n + r)) and a(n) * D(r)/((4*n + 1)*(4*n + 3)*...*(4*n + 2*r - 1)) are integers for all n. More generally, we conjecture that there are constants K(r) > 0, r >= 1, such that a(n) * K(r)/((6*n - 1)*(6*n - 3)*...*(6*n - (2*r - 1))) is an integer for all n. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006992 | OEIS-A211433

Number of ordered triples (w,x,y) with all terms in {-n,...,0,...,n} and w+2x+4y=0.

Also, a(n) is the number of ordered pairs (w,x) with both terms in {-n,...,0,...,n} and w+2x divisible by 4. If (w,x) is such a pair it is easy to see that (-w,x), (-w,-x), and (w,-x) also are such pairs. The number of pairs with both w and x positive is given by A211521(n). If w=0, x must be even, and if x=0, w must be divisible by 4. This means that a(n) = 4*A211521(n) + 2*floor(n/2) + 2*floor(n/4) + 1. Since the sequences A211521(n), floor(n/2), floor(n/4), and the constant sequence all satisfy the recurrence conjectured in the formula section, a(n) must also satisfy the recurrence, so this proves the conjecture. - _Pontus von Brömssen_, Jan 19 2020 Conjectures from _Colin Barker_, May 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006993 | OEIS-A211434

Number of ordered triples (w,x,y) with all terms in {-n,...,0,...,n} and w+2x+5y=0.

Also, a(n) is the number of ordered pairs (w,x) with both terms in {-n,...,0,...,n} and w+2x divisible by 5. If (w,x) is such a pair it is easy to see that (-x,w), (-w,-x), and (x,-w) also are such pairs. If both w and x are nonzero these four pairs lie one in each quadrant. If one of w or x is zero, the other must be a multiple of 5. This means that a(n) equals 4*A211523(n) (the nonzero pairs) plus 4*floor(n/5) + 1 (pairs with w or x equal to zero). Since the sequences A211523(n), floor(n/5), and the constant sequence all satisfy the recurrence conjectured in the formula section, a(n) must also satisfy the recurrence, so this proves the conjecture. - _Pontus von Brömssen_, Jan 17 2020 Conjectures from _Colin Barker_, May 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006994 | OEIS-A211435

Number of ordered triples (w,x,y) with all terms in {-n,...,0,...,n} and w+4x+5y=0.

Conjectures from _Colin Barker_, May 15 2017: (Start) Conjecture: a(n) ~ 4*n^2/5. - _Stefano Spezia_, Oct 17 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006995 | OEIS-A211438

Number of ordered triples (w,x,y) with all terms in {-n,...,0,...,n} and w+2x+2y=0.

Conjecture: a(n) = (5+3*(-1)^n+2*(7+3*(-1)^n)*n+14*n^2)/8. G.f.: (1+2*x+8*x^2+2*x^3+x^4)/((1-x)^3*(1+x)^2). [_Colin Barker_, Apr 18 2012]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006996 | OEIS-A211439

Number of ordered triples (w,x,y) with all terms in {-n,...,0,...,n} and w+3x+3y=0.

Conjectures from _Colin Barker_, May 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006997 | OEIS-A211440

Number of ordered triples (w,x,y) with all terms in {-n,...,0,...,n} and 2w+3x+3y=0.

Conjectures from _Colin Barker_, May 15 2017: (Start) This has the conjectured g.f. and recurrence. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006998 | OEIS-A211476

Number of -1..1 arrays x(i) of n+1 elements i=1..n+1 with x(i)+x(j), x(i+1)+x(j+1), -(x(i)+x(j+1)), and -(x(i+1)+x(j)) having one or three distinct values for every i<=n and j<=n.

Conjectures from _Colin Barker_, Jul 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-006999 | OEIS-A211479

Number of -1..1 arrays x(i) of n+1 elements i=1..n+1 with x(i)+x(j), x(i+1)+x(j+1), -(x(i)+x(j+1)), and -(x(i+1)+x(j)) having two, three or four distinct values for every i<=n and j<=n.

Conjectures from _Colin Barker_, Jul 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007000 | OEIS-A211480

Number of ordered triples (w,x,y) with all terms in {-n,...,0,...,n} and w + 2x + 3y = 1.

Conjecture: Number of partitions of 4n+2 into 3 parts. - _George Beck_, Mar 23 2023 Conjectures from _Colin Barker_, May 15 2017: (Start) Conjecture: a(n) = (8*floor(n/3)*(2*n-3*floor(n/3)-1)+3*(1-(-1)^(n+2-floor((n+2)/3))))/2 = floor((2*n-1)^2/3). - _Luce ETIENNE_, May 25 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007001 | OEIS-A211482

Number of ordered triples (w,x,y) with all terms in {-n,...,0,...,n} and w*x+w*y+x*y=w*x*y.

Conjectures from _Colin Barker_, May 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007002 | OEIS-A211490

Number of (n+1) X (n+1) -4..4 symmetric matrices with every 2 X 2 subblock having sum zero and one or two distinct values.

Conjectures from _Colin Barker_, Jul 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007003 | OEIS-A211516

Number of ordered triples (w,x,y) with all terms in {1,...,n} and w^2=x+y.

Conjecture: a(n) = -(2*n + 5/3)*w(n) + w(n)^2 + (2/3)*w(n)^3 + (2*n + 5/6)*w(2*n) - (1/2)*w(2*n)^2 - (1/3)*w(2*n)^3, where w = A000196. - _Velin Yanev_, Nov 16 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007004 | OEIS-A211526

Number of -1..1 arrays x(i) of n+1 elements i=1..n+1 with set{t,u,v in 0,1}((x[i+t]+x[j+u]+x[k+v])*(-1)^(t+u+v)) having two, four or five distinct values for every i,j,k<=n.

Conjectures from _Colin Barker_, Jul 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007005 | OEIS-A211562

Number of nonnegative integer arrays of length n+2 with new values 0 upwards introduced in order, and containing the value n-1.

Conjectures from _Colin Barker_, Jul 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007006 | OEIS-A211563

Number of nonnegative integer arrays of length n+3 with new values 0 upwards introduced in order, and containing the value n-1.

Conjectures from _Colin Barker_, Jul 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007007 | OEIS-A211564

Number of nonnegative integer arrays of length n+4 with new values 0 upwards introduced in order, and containing the value n-1.

Conjectures from _Colin Barker_, Jul 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007008 | OEIS-A211682

Prime numbers such that all the substrings of length <= 2 are primes.

A211681 restricted to primes. Contains A085823. In contrast to A085823 this sequence seems to be not finite (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hieronymus Fischer_, Apr 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007009 | OEIS-A211715

Number of (n+1) X (n+1) -11..11 symmetric matrices with every 2 X 2 subblock having sum zero and two or four distinct values.

Conjectures from _Colin Barker_, Jul 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007010 | OEIS-A211719

Number of (n+1) X (n+1) -11..11 symmetric matrices with every 2 X 2 subblock having sum zero.

Conjectures from _Colin Barker_, Jul 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007011 | OEIS-A211837

Number of nonnegative integer arrays of length n+4 with new values 0 upwards introduced in order, no three adjacent elements equal, and containing the value n+1.

Conjectures from _Colin Barker_, Jul 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007012 | OEIS-A211838

Number of nonnegative integer arrays of length n+5 with new values 0 upwards introduced in order, no three adjacent elements equal, and containing the value n+1.

Conjectures from _Colin Barker_, Jul 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007013 | OEIS-A211839

Number of nonnegative integer arrays of length n+6 with new values 0 upwards introduced in order, no three adjacent elements equal, and containing the value n+1.

Conjectures from _Colin Barker_, Jul 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007014 | OEIS-A211850

Number of nonnegative integer arrays of length 2n+5 with new values 0 upwards introduced in order, no three adjacent elements all unequal, and containing the value n+1.

Conjectures from _Colin Barker_, Jul 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007015 | OEIS-A211851

Number of nonnegative integer arrays of length 2n+6 with new values 0 upwards introduced in order, no three adjacent elements all unequal, and containing the value n+1.

Conjectures from _Colin Barker_, Jul 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007016 | OEIS-A211852

Number of nonnegative integer arrays of length 2n+7 with new values 0 upwards introduced in order, no three adjacent elements all unequal, and containing the value n+1.

Conjectures from _Colin Barker_, Jul 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007017 | OEIS-A211853

Number of nonnegative integer arrays of length 2n+8 with new values 0 upwards introduced in order, no three adjacent elements all unequal, and containing the value n+1.

Conjectures from _Colin Barker_, Jul 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007018 | OEIS-A211865

Arises in computing maximum information a Boolean function can reveal about noisy inputs.

From Table I: Reduction in number of candidate Boolean functions to be considered for verification of Conjecture 2, Kumar.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Feb 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007019 | OEIS-A211873

Numbers b >= 0 such that 2 b^2 + 3 b + 5 is prime.

An example of Sun's conjecture in A217788: For a(n)>5, also bases b such [2;3;5] seen as a number written in base b, is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007020 | OEIS-A211899

Number of triangular n X n X n arrays colored with integers 0 upwards introduced in row major order, with no element equal to any neighbor, and containing the value n(n+1)/2-2.

Conjectures from _Colin Barker_, Jul 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007021 | OEIS-A211911

Number of lower triangular n X n arrays colored with integers 0 upwards introduced in row major order, with no element equal to any horizontal, vertical, diagonal or antidiagonal neighbor, and containing the value n(n+1)/2-2.

Conjectures from _Colin Barker_, Jul 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007022 | OEIS-A211912

Number of lower triangular n X n arrays colored with integers 0 upwards introduced in row major order, with no element equal to any horizontal, vertical, diagonal or antidiagonal neighbor, and containing the value n(n+1)/2-3.

Conjectures from _Colin Barker_, Jul 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007023 | OEIS-A211924

Number of lower triangular n X n arrays colored with integers 0 upwards introduced in row major order, with no element equal to any horizontal, vertical or antidiagonal neighbor, and containing the value n(n+1)/2-2.

Conjectures from _Colin Barker_, Mar 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007024 | OEIS-A211958

Number of lower triangular n X n arrays colored with integers 0 upwards introduced in row major order, with no element equal to any diagonal or antidiagonal neighbor, and containing the value n(n+1)/2-2.

Conjectures from _Colin Barker_, Jul 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007025 | OEIS-A211995

a(n) = floor(7^n / 2^n) mod 2^n.

Conjectured to be always positive for n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Oct 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007026 | OEIS-A212031

Number of lower triangular n X n arrays colored with integers 0 upwards introduced in row major order, with no element equal to any element at a city block distance of two, and containing the value n(n+1)/2-2.

Conjectures from _Colin Barker_, Jul 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007027 | OEIS-A212039

Number of lower triangular n X n arrays colored with integers 0 upwards introduced in row major order, with no element equal to any element within a city block distance of two, and containing the value n(n+1)/2-2.

Conjectures from _Colin Barker_, Jul 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007028 | OEIS-A212198

Leading diagonal of triangle in A211321.

Conjectured to be the number of permutations of length n avoiding the partially ordered pattern (POP) {1>5, 2>5, 5>3, 5>4} of length 5. That is, conjectured to be the number of length n permutations having no subsequences of length 5 in which the fifth element is larger than the elements in positions 3 and 4, but smaller than the elements in positions 1 and 2. - _Sergey Kitaev_, Dec 13 2020 The conjecture of Kitaev has been proven. It can be restated as the number of size n permutations avoiding 45123, 45213, 54123, 54213. There are nineteen sets of permutations avoiding four size five permutations that are known to match this sequence. A further four are conjectured to match this sequence. - _Christian Bean_, Jul 23 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007029 | OEIS-A212205

G.f.: ((1+2*x)*sqrt(1-6*x^2+x^4)-1+5*x^2-2*x^3)/(2*x*(1-6*x^2)).

Conjecture D-finite with recurrence: (n+1)*a(n) +2*(n-1)*a(n-1) +6*(-2*n+1)*a(n-2) +12*(-2*n+5)*a(n-3) +(37*n-77)*a(n-4) +2*(37*n-151)*a(n-5) +6*(-n+5)*a(n-6) +12*(-n+7)*a(n-7)=0. - _R. J. Mathar_, Aug 20 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007030 | OEIS-A212210

Triangle read by rows: T(n,k) = pi(n) + pi(k) - pi(n+k), n >= 1, 1 <= k <= n, where pi() = A000720().

It is conjectured that pi(x)+pi(y) >= pi(x+y) for 1 < y <= x.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007031 | OEIS-A212211

Triangle read by rows: T(n,k) = pi(n) + pi(k) - pi(n+k), n >= 2, 2 <= k <= n, where pi() = A000720().

It is conjectured that pi(x) + pi(y) >= pi(x+y) for 1 < y <= x.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007032 | OEIS-A212212

Array read by antidiagonals: pi(n) + pi(k) - pi(n+k), where pi() = A000720.

It is conjectured that pi(x) + pi(y) >= pi(x+y) for 1 < y <= x.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007033 | OEIS-A212223

a(n) is the least integer greater than 1 whose expansion in prime bases 2 to prime(n) have equal sum of digits.

Conjecture: the sequence never terminates.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, May 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007034 | OEIS-A212275

Least k such that 4*n*k+1 is a prime of the form m^2+1, or 0 if no such k exists.

Conjecture: a(n)>0. If the conjecture is true, then there exist infinitely many primes of the form m^2+1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007035 | OEIS-A212279

A002144(n+1)^2+1 mod A002144(n), where A002144 are the Pythagorean primes (p=4k+1).

Conjecture: a(n) = A082073(n)^2 + 1 for all n > 159. - _Charles R Greathouse IV_, May 13 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, May 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007036 | OEIS-A212294

Sums of (zero or more) distinct twin primes.

Conjecturally, a(n) = n+4 for n > 5. This holds at least up to 2^500.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Jun 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007037 | OEIS-A212320

Irregular triangle: T(n, k) = k! modulo prime(n), 1<k<prime(n), 1<n.

It is conjectured that only first and second row have all terms distinct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Oct 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007038 | OEIS-A212334

Number of words, either empty or beginning with the first letter of the 4-ary alphabet, where each letter of the alphabet occurs n times and letters of neighboring word positions are equal or neighbors in the alphabet.

Conjecture: for r >= 2, and all primes p >= 5, a(p^r) == a(p^(r-1)) (mod p^(3*r+3)). - _Peter Bala_, Oct 13 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Aug 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007039 | OEIS-A212339

Sequence of coefficients of x in marked mesh pattern generating function Q_{n,132}^(0,0,3,0)(x).

Conjecture: satisfies a linear recurrence having signature (3, 0, 1, -12, -3, 1, 18, 12, 8). - _Harvey P. Dale_, Sep 03 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 09 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007040 | OEIS-A212344

Sequence of coefficients of x^(n-3) in marked mesh pattern generating function Q_{n,132}^(0,3,0,0)(x).

Conjecture: (n-2)*a(n) = 2*(2n-7)*a(n-1). _R. J. Mathar_, Jun 27 2012 G.f.: conjecture: 5*T(0), where T(k) = 1 - x/(x - 1/T(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 26 2013 G.f.: conjecture: 5*(1-sqrt(1-4*x))/(2*x) = 10/T(0), where T(k) = 2*x*(2*k+1) + k+2 - 2*x*(k+2)*(2*k+3)/T(k+1); (continued fraction). - _Sergei N. Gladkovskii_, Nov 26 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 09 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007041 | OEIS-A212346

Sequence of coefficients of x^0 in marked mesh pattern generating function Q_{n,132}^(0,4,0,0)(x).

Conjecture stated in Formula holds through a(35). Conjecture: this appears to equal (n+3)(n^2-4)/6 for n >= 3, see A129936. Conjectures from _Colin Barker_, Feb 11 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 09 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007042 | OEIS-A212392

G.f. satisfies: A(x) = (x + A(A(x)))^2 where g.f. A(x) = Sum_{n>=1} a(n)*x^(3*n-1).

Conjecture: n divides a(n); see A212391. More generally, we have the conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007043 | OEIS-A212425

G.f. satisfies: A(x) = (x + A(A(x)))^3 where g.f. A(x) = Sum_{n>=1} a(n)*x^(8*n-5).

Conjecture: (2*n-1) divides a(n); see A212426. More generally, we have the conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007044 | OEIS-A212488

Smallest k>0 such that (5^n-k)*5^n-1 and (5^n-k)*5^n+1 are a twin prime pair or 0 if no such k exists.

Conjecture: There is always one such k if n>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Jul 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007045 | OEIS-A212490

Least number m > 1 such that A000203(x)*x = m has exactly n solutions.

Conjecture: a(n) <= 2^(n-1) * Product_{k=1..n} A000396(k). - _Daniel Suteu_, Nov 01 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Naohiro Nomoto_, May 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007046 | OEIS-A212495

Numbers all of whose base 11 digits are even.

As n increases, it is most likely that A212495(n) < A033036(n), although exceptions to this rule can be found. I conjecture that A212495(n) > A033036(n) for only finitely many values of n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Douglas Latimer_, May 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007047 | OEIS-A212613

Conjectured largest number having n representations as the sum of two twin primes p + q with p >= q.

Conjectured largest number having n representations as the sum of two twin primes p + q with p >= q.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, May 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007048 | OEIS-A212652

a(n) is the least positive integer M such that n = T(M) - T(k), for k an integer, 0 <= k <= M, where T(r) = A000217(r) is the r-th triangular number.

Conjecture: n appears in row a(n) of A209260. Conjecture: a(n) is the smallest of the largest parts of the partitions of n into consecutive parts. - _Omar E. Pol_, Jan 07 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Feb 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007049 | OEIS-A212696

Central coefficient of the triangle A097609.

Conjecture: 2*n*(n-1)*(2*n+1)*(5*n-8)*a(n) -(n-1)*(115*n^3-344*n^2+299*n-82) *a(n-1) -4*(2*n-3)*(5*n^3+27*n^2-74*n+30)*a(n-2) +36*(n-1)*(5*n-3)*(2*n-3)*(2*n-5) *a(n-3)=0. - _R. J. Mathar_, Oct 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, May 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007050 | OEIS-A212730

Number of 0..2 arrays of length 2*n with sum less than 2*n in any length 2n subsequence (=less than 50% duty cycle).

Conjecture: a(n) ~ 9^n/2. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007051 | OEIS-A212731

Number of 0..2 arrays of length 2*n+1 with sum less than 2*n in any length 2n subsequence (=less than 50% duty cycle).

Conjecture: a(n) ~ 3/2*9^n. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007052 | OEIS-A212732

Number of 0..2 arrays of length 2*n+2 with sum less than 2*n in any length 2n subsequence (=less than 50% duty cycle).

Conjecture: a(n) ~ 9/2*9^n. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007053 | OEIS-A212733

Number of 0..2 arrays of length 2*n+3 with sum less than 2*n in any length 2n subsequence (=less than 50% duty cycle).

Conjecture: a(n) ~ 27/2*9^n. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007054 | OEIS-A212734

Number of 0..2 arrays of length 2*n+4 with sum less than 2*n in any length 2n subsequence (=less than 50% duty cycle).

Conjecture: a(n) ~ 81/2*9^n. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007055 | OEIS-A212797

Number of spanning trees in C_2 X C_n.

Conjecture: a(n) = 16*A001109(n+1) +3*A001109(n) -16*(A144133(n)-6*A144133(n-1)) -n = 3*A144133(n-1) -2*A144133(n-2) +3*A144133(n-3) -n. - _R. J. Mathar_, Apr

16 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007056 | OEIS-A212814

a(n) = number of integers k >= 7 such that A212813(k) = n.

Comment from Hans Havermann, Sequence Fans Mailing List, May 31 2012: The 11 numbers k for which A212813(k)=2 are 9, 11, 14, 20, 24, 27, 28, 40, 45, 48, 54. Empirically, it appears that 2632 is the sum of the number of prime partitions (A000607) of the eleven numbers 8, 10, 13, 19, 23, 26, 27, 39, 44, 47, 53. I hesitate turning this into a conjecture only because the 3 numbers k for which A212813(k)=1 are 7, 10, 12 and the sum of the number of prime partitions of the three numbers 6, 9, 11 is twelve, not eleven (the extra partition being, I think, 2+2+2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 30 2012. I added _Hans Havermann_'s comment May 31 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007057 | OEIS-A212836

Number of 0..n arrays of length 3 with 0 never adjacent to n.

Conjectures from _Colin Barker_, Jul 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007058 | OEIS-A212837

Number of 0..n arrays of length 4 with 0 never adjacent to n.

Conjectures from _Colin Barker_, Jul 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007059 | OEIS-A212838

Number of 0..n arrays of length 5 with 0 never adjacent to n.

Conjectures from _Colin Barker_, Jul 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007060 | OEIS-A212839

Number of 0..n arrays of length 6 with 0 never adjacent to n.

Conjectures from _Colin Barker_, Jul 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007061 | OEIS-A212840

Number of 0..n arrays of length 7 with 0 never adjacent to n.

Conjectures from _Colin Barker_, Jul 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007062 | OEIS-A212841

Number of 0..n arrays of length 8 with 0 never adjacent to n.

Conjectures from _Colin Barker_, Jul 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007063 | OEIS-A212843

Carmichael numbers that have only prime divisors of the form 10k+1.

Conjecture: only Carmichael numbers of the form 10n+1 can have prime divisors of the form 10k+1 (but not all Carmichael numbers of the form 10n+1 have prime divisors of the form 10k+1). Conjecture: all Carmichael numbers C (not only with three prime divisors) of the form 10n+1 that have only prime divisors of the form 10k+1 can be written as C = (30a+1)*(30b+1)*(30c+1), C = (30a+11)*(30b+11)*(30c+11), or C = (30a+1)*(30b+11)*(30c+11). In other words, there are no numbers of the form C = (30a+1)*(30b+1)*(30c+11). The first conjecture is a consequence of Korselt's criterion. - _Charles R Greathouse IV_, Oct 02 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, May 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007064 | OEIS-A212855

T(n,k) = number of n X k arrays with rows being permutations of 0..k-1 and no column j greater than column j-1 in all rows (n, k >= 1).

The integer partitions of k = 1..10 are listed on pp. 831-832 of Abramowitz and Stegun (1964). We see that, for k = 1..6, the corresponding multinomial coefficients k! / ((1!)^a_1 * (2!)^a_2 * ... * (k!)^a_k) are all distinct; that is, A070289(k) = A000041(k) and A309951(k,s) = A325305(k,s) for s = 0..A000041(k). For 7 <= k <= 10, this is not true anymore; i.e., A070289(k) < A000041(k) for 7 <= k <= 10 (and we conjecture that this is the case for all k >= 7).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007065 | OEIS-A212882

Carmichael numbers of the form k*(2*k - 1)*(p*k - p + 1)*(2*p*k - 2*p + 1), where p is odd, p from 3 to 23.

Conjecture: for any odd number p we have an infinite number of Carmichael numbers of the form k*(2*k - 1)*(p*k - p + 1)*(2*p*k - 2*p + 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, May 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007066 | OEIS-A212888

Conjectured number of positive even numbers that can be represented as the sum of two twin primes p + q in exactly n ways, with p >= q.

Conjectured number of positive even numbers that can be represented as the sum of two twin primes p + q in exactly n ways, with p >= q.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Jones_, May 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007067 | OEIS-A212916

Number of standard Young tableaux of n cells and height <= 10.

Conjecture: generally (for tableaux with height <= k), a(n) ~ k^n/Pi^(k/2) * (k/n)^(k*(k-1)/4) * prod(j=1..k,Gamma(j/2)); set k=10 for this sequence. - _Vaclav Kotesovec_, Sep 12 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, May 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007068 | OEIS-A213047

Prime numbers corresponding to cases where A212990(n) is greater than the n-th prime.

Conjecture. On every interval [1,N] the numbers m for which A212990(m) less than or equal to Prime(m) are in majority.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jun 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007069 | OEIS-A213085

First occurrence of n in A212844.

It is conjectured that every integer n>=0 appears in A212844 at least once, and therefore every a(n) is defined.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Jul 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007070 | OEIS-A213202

Number of ways to write n = p + sum_{k=1..m}(-1)^(m-k)*p_k, where p is a Sophie Germain prime and p_k is the k-th prime.

Conjecture: a(n)>0 for all n>2. Let s_n=sum_{k=1}^n(-1)^{n-k}p_k for n=1,2,3,... The author also made the following conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007071 | OEIS-A213203

The sum of the first n! integers, with every n-th integer taken as negative.

Conjecture: a(n) + (-n^2-n-11)*a(n-1) + (n^3+7*n^2-13*n+39)*a(n-2) - 2*(n-2)*(4*n^2-2*n-15)*a(n-3) + 20*(n-2)*(n-3)*(n-4)*a(n-4) = 0. - _R. J. Mathar_, Mar 21 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Cano_, Mar 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007072 | OEIS-A213210

Numbers n such that n and n + 8 are prime and there is a power of two in the interval (n,n+8).

It is a conjecture that this sequence is finite. A search around 2^n was done up to 2^1500.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brad Clardy_, Mar 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007073 | OEIS-A213222

Minimum number of distinct slopes formed by n noncollinear points in the plane.

Scott formulated the problem (on the basis of a similar problem of Erdős), gave bounds, and conjectured the formula which Unger later proved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Mar 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007074 | OEIS-A213243

Number of nonzero elements in GF(2^n) that are cubes.

Conjectures from _Colin Barker_, Aug 23 2014, verified by _Robert Israel_, Apr 22 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Jun 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007075 | OEIS-A213244

Number of nonzero elements in GF(2^n) that are 5th powers.

Conjectures from _Colin Barker_, Aug 23 2014: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Jun 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007076 | OEIS-A213245

Number of nonzero elements in GF(2^n) that are 7th powers.

Conjectures from _Colin Barker_, Aug 23 2014, verified by _Robert Israel_, Nov 20 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Jun 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007077 | OEIS-A213246

Number of nonzero elements in GF(2^n) that are 9th powers.

Conjectures from _Colin Barker_, Aug 23 2014: (Start) Conjectures verified by _Robert Israel_, Jun 27 2018.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Jun 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007078 | OEIS-A213248

Number of nonzero elements in GF(2^n) that are 13th powers.

Conjectures from _Colin Barker_, Aug 24 2014: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Jun 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007079 | OEIS-A213257

a(1) = 1, a(2) = 2 and, for n > 2, a(n) is the smallest integer greater than a(n - 1) such that no three terms of the sequence form a geometric progression of the form {x, 2 x, 4 x}.

Conjecture. The positive integers that are not in this sequence are given by the positions of 2 in the fixed-point of the morphism 0->01, 1->02, 2->03, 3->01 (see A191255). (This has been confirmed for over 5000 terms of A213257.) To illustrate, the fixed-point of the indicated morphism is {0,1,0,2,0,1,0,3,0,1,0,2,0,1,0,1,0,1,0,2,0,...} and 2 occurs at positions {4,12,20,...}, integers that are missing in A213257.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jun 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007080 | OEIS-A213258

Positive integers that are not in A213257.

Conjecture. The terms of this sequence are given by the positions of 2 in the fixed-point of the morphism 0 -> 01, 1 -> 02, 2 -> 03, 3 -> 01 (see A191255). (This has been confirmed for over 5000 terms of A213257.) To illustrate, the fixed-point of the indicated morphism is {0,1,0,2,0,1,0,3,0,1,0,2,0,1,0,1,0,1,0,2,0,...} and 2 occurs at positions {4,12,20,...}, integers in this sequence but missing from A213257. The above two conjectures are correct. This is indeed positions of 2 in A191255, and numbers of the form 2^(3t+2)*s where s is an odd number. - _Jianing Song_, Sep 21 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Jun 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007081 | OEIS-A213325

Number of ways to write n = q + sum_{k=1}^m(-1)^{m-k}p_k, where p_k is the k-th prime, and q is a practical number with q-4 and q+4 also practical.

Conjecture: a(n)>0 for all n>8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007082 | OEIS-A213340

Numbers which are the values of the quadratic polynomial 5+8t+12k+16kt on nonnegative integers.

In 1948 Erdős and Straus conjectured that for any positive integer n >= 2 the equation 4/n = 1/x + 1/y +1/z has a solution with positive integers x, y and z (without the additional requirement 0 < x < y < z).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Mizony_, Jun 09 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007083 | OEIS-A213381

a(n) = n^n mod (n+2).

Conjectures: Conjecture 1) is true: with m = n+2, a(n) = (-2)^(m-2) mod m = 0 iff m divides 2^(m-2), i.e., m = 2^k for some k with k <= m-2 (which is true for k >= 2). Conjecture 2) is true: if n = 3*p-2 where p is prime, then n == 1 (mod 3) so n^n == n (mod 3), and n^(p-1) == 1 (mod p) so n^n == n (mod p), and therefore (if p <> 3) n^n == n (mod 3*p). A separate computation verifies the case p=3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Jun 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007084 | OEIS-A213382

Numbers n such that n^n mod (n + 2) = n.

It was conjectured that A176003 is a subsequence. The conjecture is correct: verify the cases 1 and 3, then it suffices to show that (3p-2)^(3p-2) = 3p-2 mod 3 and mod p. Mod 3 the congruence is 1^(3p-2) = 1, and mod p the congruence is (-2)^(3p-2) = -2 which is true by Fermat's little theorem. - _Charles R Greathouse IV_, Sep 12 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Jun 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007085 | OEIS-A213421

Real part of Q^n, Q being the quaternion 2+i+j+k.

Conjecture: G.f. (1-2x)/(1-4x+7x^2). a(n) = A168175(n)-2*A168175(n-1). - _R. J. Mathar_, Jun 25 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Jun 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007086 | OEIS-A213443

a(0)=5, thereafter a(n) = chromatic number (or Heawood number) Chi(n) of surface of genus n.

This is what A000934 would have looked like if the four-color conjecture had turned out to be false.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007087 | OEIS-A213507

E.g.f.: exp(Sum_{n>=1} A000108(n)*x^n/n), where A000108(n) = binomial(2*n,n)/(n+1) forms the Catalan numbers.

Conjecture: a(n) is not divisible by {2,5,13,29,31,37,41,43,53,71,97}. Conjectural e.g.f.: 1/x * Series_Reversion (x/(1 + F(x))), where F(x) = x + Sum_{n >= 2} (n - 1)^(n - 2)*x^n/n! = x + x^2/2! + 2*x^3/3! + 3^2*x^4/4! + 4^3*x^5/5! + 5^4*x^6/6! + ...; that is, dF/dx = 1 - LambertW(-x) = 1 + Euler's tree function T(x). See A000169. The conjecture is equivalent to the result: Catalan(n) = [x^n] (1 + F(x))^n = [x^n] (2*x + x*T(x) + x/T(x))^n. [added Aug 16 2026: this is correct - see the Bala link.] (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007088 | OEIS-A213510

The sequence N(n) arising in the enumeration of balanced ternary trees.

Conjecture: partial sums of A064099. - _Sean A. Irvine_, Jul 14 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007089 | OEIS-A213514

For composite n, remainder of n - 1 when divided by phi(n), where phi(n) is the totient function (A000010).

D. Lehmer conjectured that a(k) is never 0. He proved that if such k exists, the corresponding composite n must be odd, squarefree, and divisible by at least 7 primes. Cohen and Haggis showed that such n must be larger than 10^20 and have at least 14 prime factors.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Balarka Sen_, Feb 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007090 | OEIS-A213519

Numbers that are the sum of cubes of distinct primes.

The complement of this sequence is conjectured to have 483370 terms, the last one being 1866000 = A121571(3). This conjecture was proved by Fuller and Nichols (see the link). - _Robert Nichols_, Sep 17 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007091 | OEIS-A213523

Numbers not representable as the sum of three heptagonal numbers.

It is conjectured that 1348 positive numbers are not the sum of three heptagonal numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007092 | OEIS-A213525

Numbers not representable as the sum of three 9-gonal numbers.

It is conjectured that 5282 positive numbers are not the sum of three 9-gonal numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007093 | OEIS-A213537

Record first differences of base sequence A213536 (a cousin prime recurrence sequence).

Three conjectures on this sequence are formulated in A213536.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph Benstock_, Jul 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007094 | OEIS-A213648

The minimum number of 1's in the relation n*[n,1,1,...,1,n] = [x,...,x] between simple continued fractions.

Conjecture: the fixed points of this sequence are in A000057. The conjecture above is true; in fact, the fixed points of this sequence are exactly the terms of A000057. See Formula. - _M. Jeremie Lafitte (Levitas)_, Aug 10 2026 Conjecture: a(n)=A001177(n)-1. The conjecture is true. For n>=2, write F(k)=A000045(k) and M(t)={{t,1},{1,0}}. If [n,1,...,1,n] contains m occurrences of 1, then M(n) M(1)^m M(n)={{P,Q},{Q,F(m+1)}}, where P F(m+1)-Q^2=(-1)^m. Applying the Euclidean algorithm to this symmetric continuant matrix, with the canonical convention that the final partial quotient is greater than 1, shows that the canonical simple continued fraction obtained by multiplying [n,1,...,1,n] by n has equal first and last partial quotients if and only if n | F(m+1). Hence a(n)=min{m>=1: n | F(m+1)}=A001177(n)-1. This proves Art DuPre's 2012 conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Jun 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007095 | OEIS-A213653

Least semiprime whose digital sum is n, or 0 if no such integer exists.

I conjecture that all terms > 278, except for 10001, end in the digit "9". What is the next term a(n) > 69 violating monotony, i.e., such that a(n) < a(n-1)? _M. F. Hasler_, Jun 17 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jun 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007096 | OEIS-A213677

Numbers n such that n and n + 12 are prime and there is a power of two in the interval (n, n+12).

It is a conjecture that this is a finite sequence. A search was conducted out to 2^1500.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brad Clardy_, Mar 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007097 | OEIS-A213710

Number of steps to reach 0 when starting from 2^n and iterating the map x -> x - (number of 1's in binary representation of x): a(n) = A071542(2^n) = A218600(n)+1.

Conjecture: A179016(a(n))= 2^n for all n apart from n=2. This is true if all powers of 2 except 2 itself occur in A179016 as in that case they must occur at positions given by this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007098 | OEIS-A213859

a(n) = 2^n mod (n+2).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Jun 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007099 | OEIS-A213883

Least number k such that (10^k-j)*10^n-1 is prime for some single-digit j or 0 if no such prime with 1<=k, 0<=j<=9 exists.

Conjecture: there is always at least one (k,j) solution for each n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Jun 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007100 | OEIS-A213892

Fixed points of a sequence h(n) defined by the minimum number of 3's in the relation n*[n,3,3,...,3,n] = [x,...,x] between simple continued fractions.

We conjecture that this sequence contains prime numbers analogous to the sequence of prime numbers A000057, in the sense that, instead of referring to the Fibonacci sequences (sequences satisfying f(n) = f(n-1) + f(n-2) with arbitrary positive integer values for f(1) and f(2)) it refers to the sequences satisfying f(n) = 3*f(n-1) + f(n-2), A006190, A003688, A052924, etc. This would mean that a prime is in the sequence if and only if it divides some term in each of the sequences satisfying f(n) = 3*f(n-1) + f(n-2). In response to the conjecture above, an exhaustive computation using the rank criterion in the Formula section found no composite term n<=10^6. The proposed characterization of the prime terms by analogy with A000057 is exact, as explained below. For a prime p, these conditions are also equivalent to p dividing a term of every integer sequence satisfying f(k+2)=3*f(k+1)+f(k). Thus the equivalence proposed in the conjecture above holds for primes. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Jun 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007101 | OEIS-A213893

Fixed points of a sequence h(n) defined by the minimum number of 4's in the relation n*[n,4,4,...,4,n] = [x,...,x] between simple continued fractions.

We conjecture that this sequence contains prime numbers analogous to the sequence of prime numbers A000057, in the sense that, instead of referring to the Fibonacci sequences(sequences satisfying f(n) = f(n-1) + f(n-2) with arbitrary positive integer values for f(1) and f(2)) it refers to the sequences satisfying f(n) = 4*f(n-1) + f(n-2), A001076, A001077, A015448, etc. This would mean that a prime is in the sequence if and only if it divides some term in each of the sequences satisfying f(n) = 4*f(n-1) + f(n-2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Jun 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007102 | OEIS-A213894

Fixed points of a sequence h(n) defined by the minimum number of 5's in the relation n*[n,5,5,...,5,n] = [x,...,x] between simple continued fractions.

We conjecture that this sequence contains prime numbers analogous to the sequence of prime numbers A000057, in the sense that, instead of referring to the Fibonacci sequences (sequences satisfying f(n) = f(n-1) + f(n-2) with arbitrary positive integer values for f(1) and f(2)) it refers to the generalized Fibonacci sequences satisfying f(n) = 5*f(n-1) + f(n-2), A052918, A015449, A164581, etc. This would mean that a prime is in the sequence if and only if it divides some term in each of the sequences satisfying f(n) = 5*f(n-1) + f(n-2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Jun 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007103 | OEIS-A213895

Fixed points of a sequence h(n) defined by the minimum number of 6's in the relation n*[n,6,6,...,6,n] = [x,...,x] between simple continued fractions.

We conjecture that this sequence contains numbers is analogous to the sequence of prime numbers A000057, in the sense that, instead of referring to the Fibonacci sequences (sequences satisfying f(n) = f(n-1) + f(n-2) with arbitrary positive integer values for f(1) and f(2)) it refers to the generalized Fibonacci sequences satisfying f(n) = 6*f(n-1) + f(n-2), A005668, A015451, A179237, etc. This would mean that a prime is in the sequence if and only if it divides some term in each of the sequences satisfying f(n) = 6*f(n-1) + f(n-2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Jun 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007104 | OEIS-A213896

Fixed points of a sequence h(n) defined by the minimum number of 7's in the relation n*[n,7,7,...,7,n] = [x,...,x] between simple continued fractions.

We conjecture that this sequence contains prime numbers analogous to the sequence of prime numbers A000057, in the sense that, instead of referring to the Fibonacci sequences (sequences satisfying f(n)=f(n-1)+f(n-2) with arbitrary positive integer values for f(1) and f(2)) it refers to the sequences satisfying f(n)=7*f(n-1)+f(n-2), A054413, A015453, etc. This would mean that a prime is in the sequence A213896 if and only if it divides some term in each of the sequences satisfying f(n)=7*f(n-1)+f(n-2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Jun 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007105 | OEIS-A213897

Fixed points of a sequence h(n) defined by the minimum number of 8's in the relation n*[n,8,8,...,8,n] = [x,...,x] between simple continued fractions.

We conjecture that this sequence contains prime numbers analogous to the sequence of prime numbers A000057, in the sense that, instead of referring to the Fibonacci sequences (sequences satisfying f(n)=f(n-1)+f(n-2) with arbitrary positive integer values for f(1) and f(2)) it refers to the sequences satisfying f(n)=8*f(n-1)+f(n-2), A041025, A015454, etc. This would mean that a prime is in the sequence A213897 if and only if it divides some term in each of the sequences satisfying f(n)=8*f(n-1)+f(n-2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Jun 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007106 | OEIS-A213898

Fixed points of a sequence h(n) defined by the minimum number of 9's in the relation n*[n,9,9,...,9,n] = [x,...,x] between simple continued fractions.

We conjecture that this sequence contains prime numbers analogous to the sequence of prime numbers A000057, in the sense that, instead of referring to the Fibonacci sequence (sequences satisfying f(n)=f(n-1)+f(n-2) with arbitrary positive integer values for f(1) and f(2)) it refers to the sequences satisfying f(n)=9*f(n-1)+f(n-2) like A099371, A015455 etc. This would mean that a prime is in the sequence A213898 if and only if it divides some term in each of the sequences satisfying f(n)=9*f(n-1)+f(n-2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Jun 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007107 | OEIS-A213899

Fixed points of a sequence h(n) defined by the minimum number of 10's in the relation n*[n,10,10,...,10,n] = [x,...,x] between simple continued fractions.

We conjecture that this sequence contains prime numbers analogous to the sequence of prime numbers A000057, in the sense that, instead of referring to the Fibonacci sequences (sequences satisfying f(n) = f(n-1) + f(n-2) with arbitrary positive integer values for f(1) and f(2)) it refers to the sequences satisfying f(n) = 10*f(n-1) + f(n-2), A041041, A015456, etc. This would mean that a prime is in the sequence A213899 if and only if it divides some term in each of the sequences satisfying f(n) = 10*f(n-1) + f(n-2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Jun 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007108 | OEIS-A213900

The minimum number of 11's in the relation n*[n,11,11,...,11,n] = [x,...,x] between simple terminating continued fractions.

If we consider the fixed points such that a(n)=n, we conjecture to obtain the sequence A000057. This sequence consists of prime numbers. We conjecture that this sequence of prime numbers, in addition to its well-known relation to the collection of Fibonacci sequences (sequences satisfying f(n)=f(n-1)+f(n-2) with arbitrary positive integer values for f(1) and f(2)) it also refers to the sequences satisfying f(n)=11*f(n-1)+f(n-2), A049666, A015457, etc. This would mean that a prime is in the sequence A000057 if and only if it divides some term in each of the sequences satisfying f(n)=11*f(n-1)+f(n-2). The conjectures in the preceding comments are true. The fixed points of this sequence are exactly the terms of A000057, and these are precisely the primes p that divide some term of every integer sequence satisfying f(k+2) = 11*f(k+1)+f(k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Jun 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007109 | OEIS-A213904

a(n) is the initial member of the least pair of prime quadruples (of the form p, p+2, p+6, p+8) with a difference of 30*n, with no other prime quadruple between the pair.

a(n) is 0 if no such pair of prime quadruples is conjectured to exist for the indicated difference.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ray G. Opao_, Jun 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007110 | OEIS-A213982

Least k >= 1 such that prime(n) +- k = 2^m * q, m >= 0, where q >= 2 is prime.

What can one say about the average behavior of a(n) for large n? It is interesting in view of the Broughan-Qizhi inequality A192869(n) >> n*(log(n))^2 and their conjecture that A192869(n) = O(n*(log(n))^2). But in the case of A213982 we have, on average, log(n) possible odd positive and negative values of k with |k| < min(prime(n)-prime(n-1), prime(n+1)-prime(n)) which is approximately log(n). Therefore, we conjecture that, on average, a(n) is approximately c*log(n) with c in (0,1). Calculations up to 10^6 (_Peter J. C. Moses_) show that, most likely, c < 0.298.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jun 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007111 | OEIS-A214016

Poulet numbers (2-pseudoprimes) of the form 7200*n^2 + 8820*n + 2701.

Conjecture: There are infinitely many Poulet numbers of the form 7200*n^2 + 8820*n + 2701.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Jul 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007112 | OEIS-A214017

Poulet numbers (2-pseudoprimes) of the form 144*n^2 + 222*n + 85.

Conjecture: There are infinite many Poulet numbers of the form 144*n^2 + 222*n + 85.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Jul 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007113 | OEIS-A214018

Least k >= 1, such that prime(n) + k has the form 2^m * q, m >= 0, where q >= 2 is prime.

By the definition, a(n)<=p_(n+1)-p_n. It is well known that, for large n, p_(n+1)-p_n on average is approximately log(n). What is the average behavior of a(n)? By the Broughan-Qizhi inequality, A192869(n)>>n*(log(n))^2. Besides, they conjecture that A192869(n)=O(n*(log(n))^2). But in the case of this sequence, we have, on average, log(n)/2 possible odd values of k< p_(n+1)-p_n. Therefore, we conjecture that, on average, a(n) is approximately c*log(n) with c in (0,1). Calculations up to 10^6 (_Peter J. C. Moses_) show that, most likely, c < 0.53 (cf. comment in A213892).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jul 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007114 | OEIS-A214027

The number of zeros in the fundamental Pisano period of the sequence A000129 mod n.

a(n) = 2 for n == 3 (mod 8). For primes p, a(p^e) = 1 if p == 7 (mod 8), 4 if p == 5 (mod 8). Conjecture: 1/6 of the primes congruent to 1 mod 8 satisfy a(p^e) = 1, 2/3 of them satisfy a(p^e) = 2 and 1/6 of them satisfy a(p^e) = 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Jul 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007115 | OEIS-A214028

Entry points for the Pell sequence: smallest k such that n divides A000129(k).

Conjecture: A175181(n)/A214027(n) = a(n). This says that the zeros appear somewhat uniformly in a period. The second zero in a period is exactly where n divides the first Lucas number, so this relationship is not really surprising.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Jul 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007116 | OEIS-A214074

Least m>0 such that 2^n-m and 3^n-m are relatively prime.

Conjecture: a(n) = 3 if and only if n = 330*k-209 for some k>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007117 | OEIS-A214160

Number of 0..5 colorings of an n X 3 array circular in the 3 direction with new values 0..5 introduced in row major order.

Conjectures from _Colin Barker_, Jul 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007118 | OEIS-A214181

Number of 0..6 colorings of an n X 3 array circular in the 3 direction with new values 0..6 introduced in row major order.

Conjectures from _Colin Barker_, Jul 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007119 | OEIS-A214196

Unique terms in sequence A210144.

Conjecture: all the terms are prime. See Conjecture 1.5(i) in Sun 2012. - _Jason Yuen_, Feb 25 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007120 | OEIS-A214280

Total number of primes which can be reached via Cunningham chains, starting with prime(n), not counting this starting prime.

It is conjectured there are arbitrarily big terms (see the MathWorld link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Jul 09 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007121 | OEIS-A214305

Fermat pseudoprimes to base 2 with two prime factors.

Conjecture: For any biggest prime factor of a Poulet number p1 with two prime factors, there exists a series with infinitely many Poulet numbers p2 formed this way: p2 mod (p1 - d) = d, where d is the biggest prime factor of p1. Note: it can be seen that the Poulet numbers divisible by 73 bigger than 2701 (7957, 10585, 15841, 31609, etc.) can be written as 1314*n + 73 as well as 2628*m + 73. Conjecture: Any Poulet number p2 divisible by d can be written as (p1 - d)*n + d, where n is a positive integer, if there exists a smaller Poulet number p1 with two prime factors divisible by d. Note: This conjecture can't be extrapolated for Poulet numbers p1 with more than two prime factors; for instance, if p1 = 561 = 3*11*17, there indeed are bigger Poulet numbers divisible by 17 (such as 1105 and 4369) that can be written as 544*n + 17, but there also exist such numbers that can't be written this way, e.g., 2465. But the first conjecture can be extrapolated. Conjecture: For any biggest prime factor of a Poulet number p1 exists a series with infinitely many Poulet numbers p2 formed this way: p2 mod (p1 - d) = d, where d is the biggest prime factor of p1. The conjectured formula p2 = (p1-d)*n+d fails for p1 = 23377 = 97*241, p2 = 34945 = 5*29*241, and d = 241. Here p1 < p2, p1 has exactly two prime factors, and d is its largest prime factor and also divides p2. Both numbers are Poulet numbers. To check this, note that 2^48 == 1 (mod 97) and 2^24 == 1 (mod 241), while both 48 and 24 divide p1-1. Likewise, 2^4 == 1 (mod 5), 2^28 == 1 (mod 29), and 2^24 == 1 (mod 241), while 4, 28, and 24 all divide p2-1. But the proposed formula would require n = (p2-d)/(p1-d) = 34704/23136 = 3/2, which is not an integer. This disproves the representation conjecture, but not the separate infinitude conjectures. - _Eiver Rodríguez_, Sep 22 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Jul 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007122 | OEIS-A214321

A214551 sorted and duplicates removed (conjectured).

A214551 sorted and duplicates removed (conjectured). Based on the 10000-term b-file for A214551. But since little is known about A214551, the values beyond 7 are conjectural. For instance, it is only a conjecture that 8 and 10 do not appear in A214551. Based on the first 10^5 terms of A214551 these conjectured values are correct; it seems that there only 1, 4, 8, 25, 29, 33 and 45 occur more than once. - _Reinhard Zumkeller_, Jul 24 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007123 | OEIS-A214342

Count of the decimal descendants of the n-th prime.

Conjecture: no terms bigger than 35 after a(8)=40.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Jul 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007124 | OEIS-A214353

Number of 3 X 3 X 3 triangular 0..n arrays with no element lying outside the (possibly reversed) range delimited by its sw and se neighbors.

Conjectures from _Colin Barker_, Jul 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007125 | OEIS-A214385

Number of 3 X 3 X 3 triangular 0..n arrays with no element lying outside the (possibly reversed) range delimited by its sw and se neighbors, and no element equal to its horizontal neighbors.

Conjectures from _Colin Barker_, Jul 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007126 | OEIS-A214409

a(n) is the smallest conjectured m such that the irreducible fraction m/n is a known abundancy index.

a(n) is the smallest conjectured m such that the irreducible fraction m/n is a known abundancy index. A known abundancy index is related to a limit. Terms of the sequence have been built with a limit of 10^30. So when n is composite, the values for a(n) are conjectural. A higher limit could provide smaller values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Jul 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007127 | OEIS-A214415

Numbers n such that prevprime(2^n) AND nextprime(2^n) = 1, where AND is the bitwise AND operator.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Aug 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007128 | OEIS-A214419

Numbers not representable as the sum of three 10-gonal numbers.

It is conjectured that 7687 positive numbers are not the sum of three 10-gonal numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007129 | OEIS-A214420

Numbers not representable as the sum of three 11-gonal numbers.

It is conjectured that 12453 positive numbers are not the sum of three 11-gonal numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jul 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007130 | OEIS-A214444

Arithmetic mean of next a(n) successive squares is prime.

Conjecture: for some k and x, a(n)=A214451(n+x) for n>k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Jul 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007131 | OEIS-A214449

Least m>0 such that n! <= m^m.

Conjecture: this sequence results from A118168 by deleting the first two 1s.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007132 | OEIS-A214458

Let S_3(n) denote difference between multiples of 3 in interval [0,n) with even and odd binary digit sums. Then a(n)=(-1)^A000120(n)*(S_3(n)-3*S_3(floor(n/4))).

In 1969, D. J. Newman (see the reference) proved L. Moser's conjecture that difference between numbers of multiples of 3 with even and odd binary digit sums in interval [0,x] is always positive. This fact is known as Moser-Newman phenomenon.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jul 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007133 | OEIS-A214526

Manhattan distances between n and 1 in a square spiral with positive integers and 1 at the center.

For n > 1, a(n) = layer(n) + abs(((n-1) mod (2*layer(n)) - layer(n))) (conjectured) where layer(n) = ceiling(0.5*sqrt(n) - 0.5). - _Karl R. Stephan_, Jan 26 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Aug 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007134 | OEIS-A214551

Reed Kelly's sequence: a(n) = (a(n-1) + a(n-3))/gcd(a(n-1), a(n-3)) with a(0) = a(1) = a(2) = 1.

It is a strong conjecture that 8 and 10 are missing from this sequence, but it would be nice to have a proof! See A214321 for the conjectured values. [I have often referred to this as "Reed Kelly's sequence" in talks.] - _N. J. A. Sloane_, Feb 18 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reed Kelly_, Jul 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007135 | OEIS-A214555

Subsequence of fixed points A099009 of the Kaprekar mapping, with numbers of the form 5{n}49{n+1}4{n}5, where d{n} means n-fold repetition of digit d.

Conjecture: Satisfies a linear recurrence having signature (1111, -112110, 1111000, -1000000). - _Harvey P. Dale_, Nov 23 2022 The given general formula for a(n) confirms the above conjecture. The formula would yield the fixed point 0 for n = -1, which suggests that the nonzero terms could be indexed starting with n = 1, so that the listed terms would be 5{n-1}49{n}4{n-1}5, n >= 1. - _M. F. Hasler_, Jul 17 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Syed Iddi Hasan_, Jul 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007136 | OEIS-A214560

Number of 0's in binary expansion of n^2.

Conjecture: for every x>=0 there is an i such that a(n)>x for n>i. Comment from _N. J. A. Sloane_, Nov 21 2013: See also the conjecture in A231898.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Jul 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007137 | OEIS-A214581

Triangle read by rows: T(n,k) is the number of unordered pairs of nodes at distance k in the circumcoronene H(n) (n=1,2,3,4,5; see definition in the Klavzar papers).

T(n,3) = 3(9n^2-9n+1)= 3*A069131(n) (for n>5 this is a conjecture). T(n,2n) = n(7n^2-1) = 6*A004126(n) (for n>5 this is a conjecture). T(n,4n-2) = 6(n^2+n-1) = 6*A028387(n-1) (for n>5 this is a conjecture). T(n,4n-1) = 3n^2 = A033428(n) (for n>5 this is a conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Aug 31 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007138 | OEIS-A214603

Location of the first gap of exactly n in Ulam numbers, or zero if none is known. The zero terms are conjectural.

Location of the first gap of exactly n in Ulam numbers, or zero if none is known. The zero terms are conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_, Jul 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007139 | OEIS-A214615

Row sums of A060338.

Conjecture: a(n) - a(n-1) +(n-1)^2*a(n-2)=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Mar 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007140 | OEIS-A214619

Numbers k such that at least one other integer m exists with the same smallest prime factor, same largest prime factor, and same multiset of decimal digits as k.

Conjecture: there is an X such that among integers bigger than X more than 50% are in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Jul 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007141 | OEIS-A214621

Numbers k such that at least one other integer m exists with the same smallest and same largest prime factors, and same multisets of decimal and binary digits as k.

Conjecture: there is X such that among integers bigger than X more than 50% are in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Jul 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007142 | OEIS-A214625

Let n=r_1*r_2*...*r_k is Fermi-Dirac factorization of n (see comment). Set g(n) = n + k - 1 and g_i, i>=0 (g_0(n) = n, g_1=g), is i-th iteration of g. a(n) is the minimal i such that g_i(n) is in A050376.

Conjecture: a(n) exists for every n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Feb 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007143 | OEIS-A214724

Expansion of e.g.f.: exp(Sum_{n>=0} x^(n^2+1)/(n^2+1)).

Conjecture: p | a(n) for n>=p when p is a prime of the form m^2+1 (A002496). [Proved in Kallat (2026). - _Ahaan Kallat_, Oct 14 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007144 | OEIS-A214758

Carmichael numbers divisible by a smaller Carmichael number.

Conjecture: A Carmichael number C1 can be written as C1 = (C2 + C3)/2, where C2 and C3 are also Carmichael numbers, only if both C1 and C3 are divisible by C2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Aug 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007145 | OEIS-A214785

Numbers k such that k + 6 = A214783(k).

This set is disjoint from A213784. Conjecture: all terms are prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Aug 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007146 | OEIS-A214786

Union of A214784 and A214785.

The sequences A214784 and A214785 are disjoint and this is their union. It is only a conjecture that these are all primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Aug 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007147 | OEIS-A214790

Numbers n such that A214789(n) = n-6.

It is still a conjecture that 10 is the only nonprime in this sequence. This sequence is disjoint from A214791.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Aug 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007148 | OEIS-A214791

Numbers n such that n+8 = A214789(n).

This sequence is disjoint from A214790. It is only a conjecture that this sequence consists of primes except for 26.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Aug 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007149 | OEIS-A214792

Union of A214790 and A214791.

This is the union of the disjoint sequences A214790 and A214791. It is only a conjecture that 10 and 26 are the only nonprimes in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Aug 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007150 | OEIS-A214794

Numbers n such that n-5=A214793(n).

It is still a conjecture that 28 is the only nonprime in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Aug 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007151 | OEIS-A214796

Numbers n such that n-8=A214795(n).

It is still a conjecture that these are all primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Aug 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007152 | OEIS-A214797

Numbers n such that n+10=A214795(n).

It is still a conjecture that these are all primes. This sequence is disjoint from A214796.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Art DuPre_, Aug 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007153 | OEIS-A214799

Let S be a set of n positive numbers such that all n choose 2 pairwise GCD's are distinct, and let max(S) denote the largest element of S; a(n) is the minimal value of max(S) over all choices for S.

Conjecture: a(n) = 15 * 2^(n-3) for n >= 6 (_Robert Israel_)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007154 | OEIS-A214808

Kostant partition function P(2*rho) for root system A_n, where 2*rho is the sum of the positive roots.

Conjecture: number of lattice points in the reflexive flow polytope of the transitive tournament on n+1 vertices. This holds at least up to n = 5. - _F. Chapoton_, Jan 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007155 | OEIS-A214813

Maximal contact number of a subset of n balls from the face-centered cubic lattice.

If b(n) denotes the maximal contact number of any arrangement of n balls then it is conjectured that a(n) = b(n) for n <= 9. It is also known that b(10)>=25, b(11)>=29, b(12)>=33 and of course b(13) >= a(13) >= 36. [Bezdek 2012]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 31 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007156 | OEIS-A214834

a(n) is the smallest prime number such that both a(n) and (4n + 2) - a(n) are prime numbers of the form 4k + 3.

Any even number of the form of 4k + 2 (k > 0) can be partitioned into 4k + 2 = (4i+3) + (4j+3), where i + j + 1 = k. This sequence implies the conjecture that for any number in the form of 4k + 2 (k > 0), there is a partition for which 4i + 3 and 4j + 3 are both prime. Conjecture tested true up to n=1000000000. In case the conjecture is not true, zero could be used to represent the missing entries.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Mar 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007157 | OEIS-A214852

Indices of Fibonacci numbers with the same number of 1's and 0's in their binary representation.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Mar 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007158 | OEIS-A214853

Fibonacci numbers with only one 0 in the binary representation.

Conjecture: the sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Mar 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007159 | OEIS-A214873

Primes p such that 2*p + 1 is also prime and p + 1 is a highly composite number (definition 1).

Conjecture: for n >= 5, GCD(A000032(a(n)), A000225(a(n))) = 2*a(n) + 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Jul 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007160 | OEIS-A214891

Numbers that are not the sum of two squares and two fourth powers.

Conjecture: The set {a(n): n > 0} coincides with {16^k*m: k = 0, 1, 2, ... and m = 23, 44, 71, 79, 184, 519, 599, 4024}. - _Zhi-Wei Sun_, Jan 27 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Jul 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007161 | OEIS-A214900

Number of ordered ways to represent n as the sum of three squares and one fourth power.

Conjecture: a(n) != 0, that is, all numbers are sums of three squares and one fourth power.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Jul 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007162 | OEIS-A214922

Numbers of the form x^2 + y^2 + z^3 + w^3 (x, y, z, w >= 0).

Conjecture: 23 is the only number not in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Deléham_, Jul 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007163 | OEIS-A214926

Difference A214925(n) - A214924(n), prime count between Ramanujan primes bounding maximal gap primes.

Conjecture: For every n > 0, a(n) > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Nicholson_, Aug 06 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007164 | OEIS-A214944

Number of squarefree words of length 5 in an (n+1)-ary alphabet.

Conjectures from _Colin Barker_, Jul 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007165 | OEIS-A214945

Number of squarefree words of length 6 in an (n+1)-ary alphabet.

Conjectures from _Colin Barker_, Jul 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007166 | OEIS-A214946

Number of squarefree words of length 7 in an (n+1)-ary alphabet.

Conjectures from _Colin Barker_, Jul 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007167 | OEIS-A214948

a(n) is the sum over all proper integer partitions of n of the previous terms.

a(n) ~ exp(k) * a(n-1), k = 0.992632731... (conjecture). - _Bill McEachen_, Dec 26 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_, Jul 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007168 | OEIS-A214970

Integers n that divide f(n), where f is the change-of-base function defined at A214969, using b = (golden ratio) and c=b^2.

Conjecture: L(2k-1) and L(2k)+1 are terms of the sequence, for all positive integers k, where L=A000032 (Lucas numbers).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007169 | OEIS-A214971

Integers k for which the base-phi representation of k includes 1.

Conjecture: L(2k-1) and L(2k)+1 are terms of this sequence for all positive integers k, where L=A000032 (Lucas numbers). Proof of this conjecture: this follows directly from the well known formula L(2k)=phi^{2k}+phi^{-2k}, and the recursion L(2k+1)=L(2k)+L(2k-1). - _Michel Dekking_, Jun 25 2019 Conjecture: If D is the difference sequence, then D-3 is the infinite Fibonacci word A096270. If so, then A214971 can be generated as in Program 3 of the Mathematica section. - _Peter J. C. Moses_, Oct 19 2012 Conjecture: A very simple formula for this sequence seems to be a(n) = ceiling((n-1)*phi) + 2*(n-1) for n>1; thus, see the related sequence A004956. - _Thomas Baruchel_, May 14 2018 Moses' conjecture is equivalent to Baruchel's conjecture: Baruchel's conjecture expresses that this sequence is a generalized Beatty sequence, and since A096270 equals the Fibonacci word A005614 with an initial zero, this follows directly from Lemma 8 in Allouche and Dekking. - _Michel Dekking_, May 04 2019 The conjectures by Baruchel and Moses are proved in my paper 'Base phi representations and golden mean beta-expansions'. - _Michel Dekking_, Jun 25 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007170 | OEIS-A214973

Number of terms in greedy representation of n using Fibonacci and Lucas numbers.

Conjecture: a(A049651(n)) = n and this is the first instance of n in the sequence for all n > 0. In other words, apart from its initial term, A049651 is the RECORDS transform of this sequence. - _Charles R Greathouse IV_, Oct 14 2021 Conjecture: a(n) = A329320(A048679(n)) for n > 0 (noticed by Sequence Machine). - _Mikhail Kurkov_, Oct 13 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007171 | OEIS-A214977

Number of terms in Lucas representations of 1,2,...,n.

See the conjecture at A214979.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007172 | OEIS-A214980

Positions of zeros in A214979.

Let U(n) and V(n) be the number of terms in the Lucas representations and Zeckendorf (Fibonacci) representations, respectively, of all the numbers 1,2,...,n. Then A214980 is the sequence of zeros of the sequence A214979(n) = V(n) - U(n). It is conjectured at A214979 that A214980 is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007173 | OEIS-A214981

Number of terms in the greedy Lucas-and-Fibonacci representations of 1,2,...,n; partial sums of A214973.

For comparison with Zeckendorf (Fibonacci) representations, it is conjectured that the limit of A179180(n)/A214981(n) exists and is between 1.2 and 1.4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007174 | OEIS-A215010

Integer side lengths in arithmetic progression of simple convex hexagons with equal interior angles. Sequence gives the values of m for sides of lengths t+m*d, counterclockwise, for the two primitive solutions.

Conjecture: hexagons are the only simple convex polygons with equal interior angles with such property, due to the fact that cos(Pi/3) = 1/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robin Garcia_, Jul 31 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007175 | OEIS-A215041

a(n) = n^degree(C(n,x))/discriminant(C(n,x)) for the minimal polynomials C(n,x) of 2*cos(Pi/n), given in A187360.

a(1) = 1. Conjectures for a(n), n>=2: i) a(2^k) = 2, k>=1;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Aug 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007176 | OEIS-A215050

Number of primes of the form 1 + b^16 for 1 < b < 10^n.

Primes 1 + b^16 are a form of generalized Fermat primes. It is conjectured that a(n) is asymptotic to 0.229464*li(10^n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henryk Dabrowski_, Aug 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007177 | OEIS-A215056

a(n) = smallest k such that k is both a sum of n nonzero squares and 50-n nonzero squares.

Conjecture: If a vector (x1,...,x50) is a zero to a signature S, then it is only also a zero for S' and no other signature T. {S' is the complement signature of S - all the signs are reversed, so if S=(+,-,-,+,-) then S' is (-,+,+,-,+)}. We therefore modify the conjecture to say that if we define Z_S as the set of integer zeros for S, and Z_t as the collection of integer zeros for all signatures T that are not S or S' (i.e., belong to t), then Z_S is NOT a subset of Z_t. The opposite - for some S, Z_S is a subset of Z_t - implies S has no function. The conjecture is obviously true for k=1, and also for k=1 there are no 'shared zeros'. For k=2, pick one of the 2 squares so that it cannot be equaled by some combination of the other 48 squares, therefore it must be combined with its companion square and so must be a unique integer zero for S(2). For k=3 we require that 2 of the 3 available squares are not matchable, and nor is their sum. This method can be extended for higher k, however the basic existence of such a zero in the first place becomes increasingly less provable.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Aug 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007178 | OEIS-A215058

Number of primes of the form 1 + b^128 for 1 < b < 10^n.

It is conjectured that a(n) is asymptotic to 0.0242888*li(10^n)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henryk Dabrowski_, Aug 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007179 | OEIS-A215066

Expansion of e.g.f.: Sum_{n>=0} Product_{k=1..n} (exp((2*k-1)*x) - 1).

Conjecture: reducing the sequence modulo a positive integer k produces an eventually periodic sequence with period dividing phi(k). (End) Conjectural g.f.: Sum_{n >= 0} (-1)^n*Product_{k = 1..n} (1 + (-1)^k*exp(- k*t)). Cf. A158690. - _Peter Bala_, Jan 28 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007180 | OEIS-A215095

a(0)=0, a(1)=1, a(n) = least k>a(n-1) such that k+a(n-2) is a Jacobsthal number.

Conjecture: G.f. (x+2*x^2)/(1-x-x^2-x^3-2*x^4). - _David Scambler_, Aug 06 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Aug 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007181 | OEIS-A215174

a(n) is the least term of A214723 having n different prime divisors, or 0 if no such term exists.

Conjectures. 1) For every n, a(n)>0; 2) for n>=3, a(n)==0 (mod 10).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, _Zak Seidov_, _Robert G. Wilson v_, Aug 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007182 | OEIS-A215191

Number of arrays of 4 0..n integers with no sum of consecutive elements equal to a disjoint adjacent sum of an equal number of elements.

Conjectures from _Colin Barker_, Jul 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007183 | OEIS-A215199

Smallest number k such that k and k+1 are both of the form p*q^n where p and q are distinct primes.

If k is a term such that k = p*q^n and k+1 = r*s^n, where p,q,r,s are primes, then clearly q != s. Conjecture: q and s are either 2 or 3 for all terms. - _Chai Wah Wu_, Mar 10 2019 Conjecture: Let 1 <= D < 2^n be the denominator of N/D of (3/2)^n. Without loss of generality, if the conjecture above holds that (q, s) = (2, 3) then r = D + k*2^n for some n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Aug 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007184 | OEIS-A215229

Number of length-6 0..k arrays connected end-around, with no sequence of L<n elements immediately followed by itself (periodic "squarefree").

Conjectures from _Colin Barker_, Jul 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 06 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007185 | OEIS-A215230

Number of length-7 0..k arrays connected end-around, with no sequence of L<n elements immediately followed by itself (periodic "squarefree").

Conjectures from _Colin Barker_, Jul 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 06 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007186 | OEIS-A215343

Fermat pseudoprimes to base 2 that can be written as 2*p^2 - p, where p is also a Fermat pseudoprime to base 2.

Conjecture: there are infinitely many Poulet numbers that can be written as (n + 1)*p^2 - n*p, where n is natural, n > 0, and p is another Poulet number. Finally, considering, e.g., that for the Poulet number 645, Poulet numbers were obtained for n = 1, 2, 4 (i.e., 831405, 1246785, 2077545), yet another conjecture: For any Poulet number p, there are infinitely many Poulet numbers that can be written as (n + 1)*p^2 - n*p, where n is natural, n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Aug 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007187 | OEIS-A215416

Numerator of zeta function multiple arising in Zig-Zag Conjecture.

Numerator of zeta function multiple arising in Zig-Zag Conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Aug 09 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007188 | OEIS-A215448

a(0)=1, a(1)=0, a(n) = a(n-1) + a(n-2) + Sum_{i=0...n-1} a(i).

Conjecture: a(n) = +3*a(n-1) -a(n-3) = A076264(n) -3*A076264(n-1) +2*A076264(n-2). G.f. (2*x-1)*(x-1) / (1-3*x+x^3). - _R. J. Mathar_, Aug 11 2012 Proof of the above conjecture: we have a(n) - a(n-1) = a(n-1) + a(n-2) + (a(0) + ... + a(n-1)) - a(n-2) - a(n-3) - (a(0) + ... + a(n-2)), which after simple algebra implies a(n) - a(n-1) = 2*a(n-1) - a(n-3), so the Mathar's formula holds true (see also Witula's comment above). - _Roman Witula_, Aug 27 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Aug 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007189 | OEIS-A215462

Number of decompositions of 2n into ordered sums of two odd nonprimes.

Conjecture: a(n) is never zero for n > 19, verified to 10^9. Conjecture: a(n) > A215461(n) > A002372(n) for sufficiently large n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter A. Lawrence_, Aug 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007190 | OEIS-A215470

Prime intersections in a square spiral with positive integers: primes p such that there are four primes among eight nearest neighbors of p.

Conjecture: the sequence is infinite. - _Alex Ratushnyak_, Sep 19 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Aug 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007191 | OEIS-A215485

Periods of square root continued fractions at which A013646 sets a new record.

Open question: Are there any more even terms after the 2?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick McKinley_, Aug 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007192 | OEIS-A215561

Number A(n,k) of permutations of k indistinguishable copies of 1..n with every partial sum <= the same partial sum averaged over all permutations; square array A(n,k), n>=0, k>=0, read by antidiagonals.

Conjecture: Row r > 1 is asymptotic to c(r) * r^(r*n) / (Pi^((r-1)/2) * n^((r+1)/2)), where c(r) are a constants. - _Vaclav Kotesovec_, Sep 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Aug 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007193 | OEIS-A215570

Number of permutations of n indistinguishable copies of 1..5 with every partial sum <= the same partial sum averaged over all permutations.

Conjectured recurrence of order 3 and degree 15: 3*(n + 1)*(n + 2)^3*(n + 4)^2*(3*n + 8)*(3*n + 10)*(65*n^3 + 398*n^2 + 781*n + 496)*(n + 3)^4*a(n + 3) - 20*(n + 1)*(n + 2)^3*(5*n + 11)*(5*n + 12)*(5*n + 13)*(5*n + 14)*(910*n^5 + 11032*n^4 + 52047*n^3 + 119686*n^2 + 134365*n + 58980)*(n + 3)^2*a(n + 2) + 25*(n + 1)*(n + 2)*(5*n + 6)*(5*n + 7)*(5*n + 8)*(5*n + 9)*(5*n + 11)*(5*n + 12)*(5*n + 13)*(5*n + 14)*(2015*n^5 + 24428*n^4 + 114387*n^3 + 258294*n^2 + 281088*n + 118368)*a(n + 1) - 250*(5*n + 1)*(5*n + 2)*(5*n + 3)*(5*n + 4)*(5*n + 6)*(5*n + 7)*(5*n + 8)*(5*n + 9)*(5*n + 11)*(5*n + 12)*(5*n + 13)*(5*n + 14)*(65*n^3 + 593*n^2 + 1772*n + 1740)*a(n) = 0. - _Manuel Kauers_ and _Christoph Koutschan_, Mar 02 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Aug 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007194 | OEIS-A215594

Expansion of f(-x, -x^4) / f(x, x^4) in powers of x where f(,) is Ramanujan's two-variable theta function.

Let A(q) denote the g.f. of this sequence. Let m be a nonzero integer. The simple continued fraction expansions of the real numbers A(1/(2*m)) and A(1/(2*m+1)) may be predictable. For a given positive integer n, the sequence of the n-th partial denominators of the continued fractions are conjecturally polynomial or quasi-polynomial in m for m

sufficiently large. An example is given below. Cf. A080054 and A098151. - _Peter Bala_, Jun 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Aug 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007195 | OEIS-A215616

From Wendt's determinant compute (-A048954(2*n)/3)^(1/3).

It is known that 3 divides A048954(2*n). It is conjectured that the quotient is a perfect cube.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Aug 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007196 | OEIS-A215641

Smallest prime whose decimal expansion consists of the concatenation of a 1-digit prime, a 2-digit prime, a 3-digit prime, ..., and an n-digit prime.

It is a plausible conjecture that a(n) always exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Aug 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007197 | OEIS-A215647

Smallest semiprime whose decimal expansion consists of the concatenation of a 1-digit semiprime, a 2-digit semiprime, a 3-digit semiprime, ..., and an n-digit semiprime.

This is to A215641 as semiprimes A001358 are to primes A000040. It is a plausible conjecture that a(n) always exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Aug 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007198 | OEIS-A215658

Primes p such that the smallest positive integer k for which p# + k is square satisfies p# + k = k^2, where p# = 2*3*5*7*11*...*p is a primorial.

Conjecture: Not the same sequence as A192579, which is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Sep 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007199 | OEIS-A215672

Fermat pseudoprimes to base 2 with three prime factors.

Conjecture: Any Poulet number P with three or more prime divisors has at least one prime divisor q for that can be written as P = q*((r + 1)*q - r), where r is a natural number. The conjecture fails for a(80) = 617093 = 43 * 113 * 127. - _Charles R Greathouse IV_, Dec 07 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Aug 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007200 | OEIS-A215689

Smallest prime whose decimal expansion consists of the concatenation of a 1-digit square, a 2-digit square, a 3-digit square, ..., and an n-digit square, or 0 if there is no such prime.

The n-th term has n(n+1)/2 digits (cf. A000217). There are (0, 3, 29, 991, 175210, ...) primes of that form, for n = 1, 2, 3, ... We can conjecture that a(n) > 0 for all n, and even that the terms converge to the concatenation of (s(1), s(2), s(3), ...) where s(n) is the smallest n-digit square, cf. formula. - _M. F. Hasler_, Dec 31 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Aug 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007201 | OEIS-A215692

Smallest prime whose decimal expansion consists of the concatenation of a 1-digit cube, a 2-digit cube, a 3-digit cube, ..., and an n-digit cube, or 0 if there is no such prime.

The n-th term has A000217(n) = n(n+1)/2 digits. We can conjecture that a(n) > 0 for all n > 1 and the terms converge to the concatenation of (c(1), c(2), c(3), ...) where c(k) is the smallest k digit cube, cf. formula. The number of such primes between a(n) and A340115(n) (the largest of this form) is (0, 2, 2, 9, 177, 6909, 570166, ...). (In particular, for n = 2 and 3, a(n) and A340115(n) are the only two primes of this form.) This is very close to what we expect, given the number of concatenations of cubes of the respective length (product of 10^(k/3)-10^((k-1)/3), k=1..n) and the density of primes in that range according to the PNT. - _M. F. Hasler_, Dec 31 2020 a(n) ~ 10^(n(n+1)/2)*0.12712510001064810382310000010077696 1... (conjectured) - _M. F. Hasler_, Dec 31 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Aug 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007202 | OEIS-A215698

Number of primes of the form 1 + b^256 for 1 < b < 10^n.

It is conjectured that a(n) is asymptotic to 0.0290422*li(10^n)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henryk Dabrowski_, Aug 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007203 | OEIS-A215699

Number of primes of the form 1 + b^512 for 1 < b < 10^n.

It is conjectured that a(n) is asymptotic to 0.0146271*li(10^n)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henryk Dabrowski_, Aug 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007204 | OEIS-A215700

Number of primes of the form 1 + b^1024 for 1 < b < 10^n.

It is conjectured that a(n) is asymptotic to 0.00783139*li(10^n)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henryk Dabrowski_, Aug 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007205 | OEIS-A215719

The smallest of four consecutive primes with prime gaps {a,b,c} = {10,18,2}.

Conjecture: The terms of any feasible prime gap triple {a,b,c} to form a quadruple of consecutive primes are sums of terms of three consecutive subsequences of the infinite integer sequence with period (4,2,4,2,4,6,2,6). By this token all possible sequences of quadruples of consecutive primes can be generated, including those already in the OEIS.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V.J. Pohjola_, Aug 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007206 | OEIS-A215895

Primes p with property that there exists a number D such that p-3D, p-2D, p-D, p+D, p+2D, p+3D are all primes.

Conjecture: only 130633 primes are not in the sequence: 2, 3, ..., 94532497.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Aug 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007207 | OEIS-A215926

Smallest deficient number k such that the product k*n is non-deficient (perfect or abundant).

Conjecture: a(n) is 1, 3, or a power of 2. Conjecture: The first occurrence of 2^m happens at A014210(m).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Aug 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007208 | OEIS-A215944

Fermat pseudoprimes to base 2 with three prime factors divisible by a smaller Fermat pseudoprime to base 2.

Conjecture: The only Fermat pseudoprimes to base 2 divisible by a smaller Fermat pseudoprime to base 2 that can't be written as p*((m + 1)*p - m)*((n + 1)*p - n), where m, n, p are natural numbers, are multiples of 5461 and can be written as 5461*(42*k - 13). Conjecture is checked for the numbers from the sequence above and for the first 15 Poulet numbers with four prime factors.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Aug 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007209 | OEIS-A215973

a(0) = 1, for n > 0: a(n) = Sum_{k=0..n-1} a(k) * (1 + a(n-1-k)).

Conjecture: (n+1)*a(n) +3*(-3*n+1)*a(n-1) +4*(5*n-7)*a(n-2) +2*(-8*n+19)*a(n-3) +2*(2*n-7)*a(n-4)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Aug 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007210 | OEIS-A216063

a(n) is the conjectured highest power of n which has no two identical digits in succession.

a(n) is the conjectured highest power of n which has no two identical digits in succession. A naive heuristic suggests that there are infinitely many n such that a(n) = 6 but only finitely many a(n) such that a(n) > 6. This suggests a weaker conjecture: this sequence is bounded. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V. Raman_, Sep 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007211 | OEIS-A216064

a(n) is the conjectured highest power of n which has no three identical digits in succession.

a(n) is the conjectured highest power of n which has no three identical digits in succession.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V. Raman_, Sep 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007212 | OEIS-A216065

a(n) is the conjectured highest power of n which has no four identical digits in succession.

a(n) is the conjectured highest power of n which has no four identical digits in succession.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V. Raman_, Sep 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007213 | OEIS-A216094

a(1) = 2; a(n+1) = a(n) + |A036263(n)| + 2.

Conjecture: a(n) ~ prime(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Sep 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007214 | OEIS-A216137

a(n) = conjectured number of integers k such that n^k has no two consecutive identical digits.

a(n) = conjectured number of integers k such that n^k has no two consecutive identical digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V. Raman_, Sep 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007215 | OEIS-A216138

a(n) = conjectured number of integers k such that n^k has no three consecutive identical digits.

a(n) = conjectured number of integers k such that n^k has no three consecutive identical digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V. Raman_, Sep 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007216 | OEIS-A216139

a(n) = conjectured number of integers k such that n^k has no four consecutive identical digits.

a(n) = conjectured number of integers k such that n^k has no four consecutive identical digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V. Raman_, Sep 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007217 | OEIS-A216140

Conjectured number of digits in highest power of n with no two consecutive identical digits.

Conjectured number of digits in highest power of n with no two consecutive identical digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V. Raman_, Sep 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007218 | OEIS-A216141

Conjectured number of digits in highest power of n with no three consecutive identical digits.

Conjectured number of digits in highest power of n with no three consecutive identical digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V. Raman_, Sep 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007219 | OEIS-A216142

Conjectured number of digits in highest power of n with no four consecutive identical digits.

Conjectured number of digits in highest power of n with no four consecutive identical digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V. Raman_, Sep 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007220 | OEIS-A216152

A205957(n) where n is a nonprime number.

a(1),...,a(10) are highly totient numbers (A097942) and products of distinct factorials (A058295). The author conjectures that this is true in general.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Sep 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007221 | OEIS-A216170

Fermat pseudoprimes to base 2 of the form (n^2 + 2*n)/3.

Conjecture: For any m natural, m > 1, there exists a series with infinitely many Fermat pseudoprimes to base 2, P, formed this way: P = (n^m + m*n)/(m+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Sep 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007222 | OEIS-A216196

A diagonalized Sudoku torus.

Conjecture: This is the maximum for a Xoox-free grid.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Mar 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007223 | OEIS-A216211

Number of self-avoiding walks of any length from NW to SW corners of a grid or lattice with n rows and 4 columns.

Conjectures from _Colin Barker_, Nov 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Toby Gottfried_, Mar 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007224 | OEIS-A216217

Smallest k such that 6^n - 2*k*3^n - 1 and 6^n - 2*k*3^n + 1 are twin primes or 0 if no solution, n > 1.

Conjecture: there is only one zero term: a(5) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Mar 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007225 | OEIS-A216231

Fibonacci with priority for primes: a(0)=0, a(1)=1, for n >= 2, a(n) = a(n-1) + a(k), where 0 < k <= n-2 is maximal index such that a(n-1) + a(k) is prime. If there is no such k, then a(n) = a(n-1) + a(n-2).

Conjecture: There exist arbitrarily long chains of consecutive prime terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Mar 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007226 | OEIS-A216240

Composite numbers arising in Eratosthenes sieve with removing the multiples of every other remaining numbers after 2 (see comment).

Conjecture. There exists x_0 such that for every x>=x_0, the number of a(n)<=x is more than pi(x).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Mar 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007227 | OEIS-A216265

Number of primes between n^3 - n and n^3.

Conjecture: a(n) > 0 for n > 13.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Mar 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007228 | OEIS-A216266

Number of primes between n^3 and n^3+n (inclusive).

Conjecture: a(n)>0 for n>23.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Mar 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007229 | OEIS-A216275

Fibonacci + Goldbach: a(1)=6, a(2)=8 and for n>=3, a(n)=g(a(n-1)) + g(a(n-2)), where for m>=3, g(2*m) is the maximal prime p < 2*m such that 2*m - p is prime.

Conjecture. lim a(n+1)/a(n)=phi as n goes to infinity (phi=golden ratio).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Mar 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007230 | OEIS-A216276

Fermat pseudoprimes to base 2 of the form (p^2 + 2*p)/3, where p is also a Fermat pseudoprime to base 2.

Conjecture: For any Fermat pseudoprime to base 2, p1, there exist infinitely many Fermat pseudoprimes to base 2, of the form p2 = (p1^n + n*p1)/(n+1), where n > 1.
Conjecture: For any Carmichael number c there exist infinitely many Carmichael numbers of the form (c^n + n*c)/(n + 1) with n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Sep 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007231 | OEIS-A216359

G.f. satisfies: A(x) = (1 + x*A(x)^2) * (1 + x/A(x)).

For n not of the form 3*m + 1, we conjecture that a(n) = Sum_{k = 0..n} binomial(-n+3*k+1, k)*binomial(-n+3*k+1, n-k)/(-n+3*k+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007232 | OEIS-A216368

Number T(n,k) of distinct values taken by k-th derivative of x^x^...^x (with n x's and parentheses inserted in all possible ways) at x=1; triangle T(n,k), n>=1, 1<=k<=n, read by rows.

Conjecture: T(n,n) = A000081(n) for n>=1. It would be nice to have a proof (or a disproof if the conjecture is wrong).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Sep 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007233 | OEIS-A216371

Odd primes with one coach: primes p such that A135303((p-1)/2) = 1.

Conjecture relating to odd integers: iff an integer is in the set A216371 and is either of the form 4q - 1 or 4q + 1, (q>0); then the top row of its coach (cf. A003558) is composed of a permutation of the first q odd integers. Examples: 11 is of the form 4q - 1, q = 3; with the top row of its coach [1, 5, 3]. 13 is of the form 4q + 1, q = 3; so has a coach of [1, 3, 5]. 37 is of the form 4q + 1, q = 9; so has a coach with the top row composed of a permutation of the first 9 odd integers: [1, 9, 7, 15, 11, 13, 3, 17, 5]. - _Gary W. Adamson_, Sep 08 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Sep 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007234 | OEIS-A216374

Number of ways to express the square of the n-th prime as the sum of four nonzero squares.

The simple counting and the conjectured first formula agree for all the primes from 3 to 997. The counting and the conjectured second formula agree for all the primes from 5 to 997. The author of this sequence would like to know whether the formulas are already known and/or how it could be proved. a(n) = floor((prime(n)^2 + 4*prime(n) + 24)/48) (conjectured for n>1). a(n) = (prime(n)^2 + 4*prime(n) + (19*(5*(prime(n) mod 48)+2)^2) mod 48 - 24)/48 (conjectured for n>2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mark Underwood_, Sep 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007235 | OEIS-A216440

a(n) = smallest m such that 2n-1 | 2^m+1, or 0 if no such m exists.

Conjecture: a(n) is the number of riffle shuffles of 2(n-1) cards required to reverse the order of the deck, or 0 if no such number of shuffles exists. This conjecture is in the context of the in-shuffle variant of the riffle shuffle, e.g., [1, 2, 3, 4] -> [3, 1, 4, 2] -> [4, 3, 2, 1].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V. Raman_, Sep 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007236 | OEIS-A216442

Numbers of the form 2^i + 3^j + 5^k, where i, j, k >= 0.

Conjecture: Unlike A004050, which has a limited set of integers expressible in more than one way, this set has no such limit.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Sep 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007237 | OEIS-A216468

Primes p with property that there exists a number d>0 such that numbers p-k*d, k=1...6, are six primes.

Conjecture: only 312722 primes are not in the sequence: 2, 3, ..., 198702899.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Sep 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007238 | OEIS-A216495

Primes p with property that there exists a number d>0 such that numbers p-d, p-2*d are primes.

Conjecture: only 5 primes are not in the sequence: 2, 3, 5, 13, 37.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Sep 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007239 | OEIS-A216496

Primes that are not in A094383.

Conjecture: a(25)=557 is the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Sep 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007240 | OEIS-A216497

Primes p with property that there exists a number d>0 such that numbers p-k*d, k=1...4, are four primes.

Conjecture: only 653 primes are not in the sequence: 2, 3, ..., 100291.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Sep 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007241 | OEIS-A216498

Primes p with property that there exists a number d>0 such that numbers p-k*d, k=1...5, are five primes.

Conjecture: only 9198 primes are not in the sequence: 2, 3, ..., 2521081.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Sep 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007242 | OEIS-A216539

Largest prime p such that there is no n-term arithmetic progression of primes ending with p.

For n>=3, the value given is only a conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Sep 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007243 | OEIS-A216557

Number of iterations of A216556 until the initial value n appears as a substring of the iterate; 0 if this will never happen.

Can someone prove (and maybe strengthen) the following conjecture: a(n) = 0 whenever A216587(m) = -1 for all m obtained by concatenating any digit to the left and any digit to the right of n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Sep 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007244 | OEIS-A216590

Primes p with property that there exists a number d>0 such that numbers p-k*d, k=1...7, are seven primes.

Conjecture: only 5254157 primes are not in the sequence: 2, 3, ..., 5082095279. Conjecture: for any k>0 there exists p0 such that for any prime p>p0 there exists a k-term arithmetic progression of primes with p at the end.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Sep 09 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007245 | OEIS-A216700

Violations of Rosser's rule: numbers n such that the Gram block [g(n), g(n+k)) contains fewer than k points t such that Z(t) = 0, where Z(t) is the Riemann-Siegel Z-function.

Lehman showed that this sequence is infinite and conjectured (correctly) that a(1) > 10^7. Brent (1979) found a(1)-a(15). Brent, van de Lune, te Riele, & Winter extended this to a(104). Gourdon extended the calculation through a(320624341).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Sep 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007246 | OEIS-A216714

a(n) = 2^(n-5) - A000931(n).

It is conjectured that this sequence (with a different offset) and A038360 are the same.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007247 | OEIS-A216732

Conjectural order of largest subgraph of diameter n of the two-dimensional mesh with maximal degree 3.

Conjectural order of largest subgraph of diameter n of the two-dimensional mesh with maximal degree 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007248 | OEIS-A216814

Given n and a constant C, define a sequence b(m) by the recurrence in the comments; a(n) = smallest positive integer C such that for some prime p the denominators of all b(m) are powers of p (conjectured).

Given n and a constant C, define a sequence b(m) by the recurrence in the comments; a(n) = smallest positive integer C such that for some prime p the denominators of all b(m) are powers of p (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007249 | OEIS-A216835

Fibonacci + Goldbach (dual sequence to A216275). a(1)=5, a(2)=7 and for n>=3, a(n) = g(a(n-1) + a(n-2)), where for m>=3, g(2*m) is the maximal prime p < 2*m such that 2*m - p is prime.

Conjecture. lim a(n+1)/a(n)=phi as n goes to infinity (phi=golden ratio).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Mar 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007250 | OEIS-A216879

G.f. satisfies: A(x) = sqrt(theta_3(x*A(x)) / theta_4(x*A(x))).

Conjectured to be the number of permutations of length n+1 avoiding the partially ordered pattern (POP) {5>1, 1>2, 1>4} of length 5. That is, conjectured to be the number of length n+1 permutations having no subsequences of length 5 in which the fifth element is larger than the first element, which in turn is larger than the second and fourth elements. - _Sergey Kitaev_, Dec 13 2020 The conjecture was disproven. The numbers are actually A366706, which matches the first 9 entries. - _Christian Bean_, Jul 22 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007251 | OEIS-A216938

Number of side-2 hexagonal 0..n arrays with values nondecreasing E, SW and SE.

Conjectures from _Colin Barker_, Jul 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007252 | OEIS-A216940

Number of side-4 hexagonal 0..n arrays with values nondecreasing E, SW and SE.

a(n) = (1+n)*(2+n)*(3+n)*(4+n)*(5+n)*(6+n)^2*(7+n)^3*(8+n)^2*(9+n)*(10+n)*(11+n)*(12+n)*(13+n)*(151184836155757307904000000 + 375573334572799334737920000*n + 45137854540680193956153600*n^2 + 3482984637133501033554048 0*n^3 + 193143943474599207101027 04*n^4 + 816635331585979471929686 4*n^5 + 272690484096441703337652 0*n^6 + 735273283907306553706472*n^7 + 162382123713323392711687*n^8 + 296300153616613712908 44*n^9 + 448755757551481013236 2*n^10 + 56469403484836599633 6*n^11 + 589003614336182448 60*n^12 + 50622267972163529 60*n^13 + 354853893929158096*n^14 + 1996972899878107 2*n^15 + 880856790135603*n^16 + 29345762188932*n^17 + 694580474022*n^18 + 10413780440*n^19 + 74384146*n^20)/22142459927970310563571395723264000000 (conjectured). - _Manuel Kauers_ and _Christoph Koutschan_,

Mar 01 2023 [Proved in Lizardi (2026); see Links. - _Alejandro Lizardi_, Jun 18 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007253 | OEIS-A216944

The least odd prime p for which the n-th prime can be written as 3p + q - 3 for some odd prime q.

Conjecture: Any prime >= 11 can be written this way.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Sep 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007254 | OEIS-A216971

Triangle read by rows: T(n,k) is the number of functions f:{1,2,...,n}->{1,2,...,n} that have exactly k nonrecurrent elements mapped to some (one or more) recurrent element. n >= 1, 0 <= k <= n-1.

Conjecture: every entry in row n is divisible by n. - _Jon Perry_, Sep 21 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Critzer_, Sep 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007255 | OEIS-A216994

Multiples of 7 such that the digit sum is divisible by 7.

Conjecture: Every century has a representation in the sequence. The conjecture is true. See a proof in the links section. - _FUNG Cheok Yin_, Dec 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Sep 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007256 | OEIS-A217016

Least r > 1 without Goldbach partition 2r = p+q such that |p-q| is prime(n)-smooth.

Conjecture [D. Broadhurst]: a(n) is prime(n)-smooth.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Broadhurst_ and _M. F. Hasler_, Sep 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007257 | OEIS-A217147

Prime numbers after which at least four distinct classes modulo 7 are equally represented among the primes to that point.

Only after 13 and 223 are five of the congruence classes modulo 7 equally represented, and it's not unreasonable to conjecture that this holds permanently.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James G. Merickel_, Sep 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007258 | OEIS-A217157

a(n) is the least value of k such that the decimal expansion of n^k contains two consecutive identical digits.

Conjecture: 1 <= a(n) <= 16 for n > 1 and a(n) < 16 for n > 2. - _Chai Wah Wu_, Feb 20 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V. Raman_, Sep 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007259 | OEIS-A217214

G.f.: (P-Q*(sqrt(1-4*x)))/(x*(1-4*x)^3) where P=(1-4*x)^3*(1-x-x^2), Q=1-11*x+39*x^2-40*x^3-22*x^4.

Conjecture D-finite with recurrence: -(n+1)*(423*n-1718)*a(n) +(-423*n^2+4779*n+1718)*a(n-1) +(17801*n^2-84527*n+75390)*a(n-2) -2*(9341*n-20828)*(2*n-7)*a(n-3)=0. - _R. J. Mathar_, Jan 25 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007260 | OEIS-A217280

a(n) = Sum_{k=0..n} binomial(n,k)^3 * 2^k * 3^(n-k).

5 divides a(n) except when a(n) == 1 (mod 5), which occurs only when the base 5 representation of n consists solely of digits '0' and '2' (conjecture). Positions n at which a(n) == 1 (mod 5) begin: [0, 2, 10,12, 50,52,60,62, 250,252,260,262,300,302,310,312, 1250,1252,1260,1262,1300,1302,1310,1312,1500,1502,1510,1512, 1550,1552,1560,1562, ...].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007261 | OEIS-A217317

Number of primes between n^2 and n^2 + log_2(n)^2 (inclusive).

Conjecture: a(n) > 0 for n > 4765516. Conjecture checked up to n = 5 * 10^10. - _Charles R Greathouse IV_, Mar 21 2013 Conjecture checked up to 4 * 10^18. Note that this conjecture is consistent with Granville's conjecture that lim sup (prime(n+1)-prime(n))/log(prime(n))^2 >= 2/e^gamma, where gamma is Euler's constant. - _Charles R Greathouse IV_, Mar 21 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Mar 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007262 | OEIS-A217319

Numbers with binary representation ending in 4*k+2 or 4*k+3 zeros.

Conjecture. For n>=1, a(n) = A171949(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Mar 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007263 | OEIS-A217333

G.f.: exp(Sum_{n>=1} x^n/n * Sum_{k=0..n} binomial(n,k)^2 * x^k/(1-x)^k).

Conjecture: (n+3)*a(n) +2*(-2*n-3)*a(n-1) +4*n*a(n-2) +2*(-2*n+3)*a(n-3) +4*(n-3)*a(n-4)=0. - _R. J. Mathar_, May 17 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007264 | OEIS-A217358

Series reversion of x-x^3-x^4.

Conjecture: 46*n*(n-1)*(n-2)*a(n) -(n-1)*(n-2)*(11*n-74)*a(n-1) -(n-2)*(336*n^2-1359*n+1351)*a(n-2) +(-347*n^3+2190*n^2-3861*n+1330)*a(n-3) + 8*(2*n-7)*(4*n-15)*(4*n-17)*a(n-4) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Oct 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007265 | OEIS-A217376

Numbers n such that n, 2n-1 and 2n+1 all are prime or a prime power.

a(9) = (3^541-1)/2. - Conjectured by _Jack Brennen_, Oct 01 2012; confirmed by _Max Alekseyev_, Nov 10 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, after an idea from Warren Smith, Oct 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007266 | OEIS-A217442

Numbers n such that d(prime(n) - 1) | n, where d(k) is the number of divisors of k.

For n in {1,2,3,4,6}, n = d(prime(n)-1). There are no others with this property, as conjectured by _Raphie Frank_ and proved by _Charles R Greathouse IV_ on Physics Forums (Nov, 2010).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Raphie Frank_, Oct 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007267 | OEIS-A217447

Number of n x n permutation matrices that disconnect their zeros.

Conjecture: 2*a(n) + 2*(-n-1)*a(n-1) + (6*n-11)*a(n-2) + (-5*n+14)*a(n-3) + 3*a(n-4) + (n-6)*a(n-5) = 0. - _R. J. Mathar_, Nov 30 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Hopkins_, Nov 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007268 | OEIS-A217448

Least k > 0 such that 1 + n^2 and 1 + (n+k)^2 have the same smallest prime factor.

Conjecture: a(n) exists for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Oct 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007269 | OEIS-A217471

Partial sum of fifth power of the even-indexed Fibonacci numbers.

The tables of the coefficient of the polynomials which appear in Ozeki's formula and in Melham's conjecture are found in A217472 and A217475, respectively (see References). a(n) =(F(2*n+1)-1)^2*(4*F(2*n+1)^3 + 8*F(2*n+1)^2 - 3*F(2*n+1) - 14)/44 (an example for Melham's conjecture, see the reference, eq. (2.7) for m=2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007270 | OEIS-A217472

Coefficient table for polynomials used for the formula of partial sums of odd powers of even-indexed Fibonacci numbers.

For Melham's conjecture on pL(m)*sum(F(2*k)^(2*m+1),k=0..n) see A217475 where also the reference is given.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007271 | OEIS-A217475

Coefficients of polynomials in a Melham conjecture.

Coefficients of polynomials in a Melham conjecture. A conjecture by Melham (see the reference, eq. 2.7) is: The table a(m,l) lists the coefficients of these polynomials for m=1..6. Thus the conjecture is certainly true for m=1..6. a(m,l) = [x^l]P(2*m-1,x), m>-1, l=0..2*m-1, with the polynomial P appearing in the Melham conjecture stated in the comment section.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007272 | OEIS-A217519

Base-2 state complexity of partitioned deterministic finite automaton (PDFA) for the periodic sequence (123...n)*.

Neder's conjecture was proved by Kreczman, Prigioniero, Rowland, and Stipulanti. - _Eric Rowland_, Feb 02 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007273 | OEIS-A217539

Number of Dyck paths of semilength n which satisfy the condition: number of returns + number of hills < number of peaks.

Conjecture: 2*(n+1)*(n-3)*a(n) +(-15*n^2+53*n-12)*a(n-1) +(28*n^2-157*n+165)*a(n-2) + 3*(3*n^2+2*n-26)*a(n-3) -18*(2*n-7)*(n-4)*a(n-4)=0. - _R. J. Mathar_, Nov 11 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Oct 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007274 | OEIS-A217586

a(1) = 1 and, for all n >= 1, if a(n) = 0 then a(2*n) = 1 and a(2*n+1) = 0 whereas if a(n) = 1 then a(2*n) = 0 and a(2*n+1) = 0.

Conjecture: a(n) = (1 - (1 - 2*A096268(n-1))*(1 - 2*A036987(n-1)))/2. - _Velin Yanev_, Aug 08 2017 Here is a proof of Velin Yanev's conjecture. Note that Yanev's conjecture can be reformulated as From (*) we see now that the conjecture will be proved if one shows that

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Layman_, Oct 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007275 | OEIS-A217596

Expansion of g.f. x / reversion(x - x^2 - x^3).

Conjecture: 25*n*(n-1)*a(n) - 5*(n-1)*(25*n-42)*a(n-1) + 3*(-23*n^2 + 59*n + 4)*a(n-2) + 9*(3*n-10)*(3*n-11)*a(n-3)=0. - _R. J. Mathar_, May 23 2014 Maple's sumrecursion command gives the second-order recurrence equation: 5*n*(n - 1)*(4*n - 9)*a(n) = 2*(n - 1)*(44*n^2 - 165*n + 150)*a(n-1) + 3*(4*n - 5)(3*n - 7)(3*n - 8)*a(n-2) with initial conditions a(1) = -1 and a(2) = -2. Mathar's conjectured third-order recurrence follows from this. - _Peter Bala_, Feb 15 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Mar 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007276 | OEIS-A217606

a(n) is the least unused prime greater than 3 such that (a(n) + a(n-1))/2 is prime, with a(0)=13.

Conjecture: every prime of the form 12k+1 is a member.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pedja Terzic_, Oct 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007277 | OEIS-A217696

Let p = A002145(n) be the n-th prime of the form 4k+3, then a(n) is the smallest number such that p is the smallest prime of the form 4k+3 for which 4*a(n)+2-p is prime.

It is conjectured that a(n) is defined for all positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Mar 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007278 | OEIS-A217703

a(0)=1, a(1)=0, and a(n+1) = 2*n*(n+1)*a(n)-n^4*a(n-1) for n>0.

Conjectures: (i) S_n(x) is irreducible over the field of rational numbers for every n=1,2,3,...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007279 | OEIS-A217711

Total number of 321 patterns in the set of permutations avoiding 123.

Conjecture: -(n+1)*(25*n-3314)*a(n) -5*n*(5*n+9446)*a(n-1) +2*(594*n^2 +128863*n -142613)*a(n-2) +16*(-119*n^2-39230*n+87888)*a(n-3) -32*(2*n-7)*(53*n-8687)*a(n-4)=0. - _R. J. Mathar_, Oct 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cheyne Homberger_, Mar 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007280 | OEIS-A217721

Number of primes between n^2 - log_2(n)^2 and n^2 (inclusive).

Conjecture: a(n) > 0 for n > 1. Conjecture checked up to n = 2^28 - 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Mar 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007281 | OEIS-A217729

Trajectory of 40 under the map n-> A006369(n).

It is conjectured that this trajectory does not close on itself.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007282 | OEIS-A217738

Numbers k such that Fibonacci(k) is divisible by k*(k+1).

Conjectures: the sequence is infinite; all terms are divisible by 12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Mar 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007283 | OEIS-A217772

a(n) = ((p+1)*(3p)!/((2p-1)!*(p+1)!*2p) - 3)/(3p^3), where p is the n-th prime.

This sequence is based on Gary Detlefs's conjecture, which he posted as a comment to A005809. His conjecture is equivalent to the conjecture that the Diophantine equation ((n+1)*(3*n)!/((2*n-1)!*(n+1)!*2*n)-3)/n^3 = m has integer solutions m for all odd primes n. Additionally I conjecture that all m are divisible by 3, therefore terms of this sequence a(n) = m/3. The values of this sequence's terms are replicated by conjectured general formula, given in A223886 (and also added to the formula section here) for k=3, j=1 and n>=2. - _Alexander R. Povolotsky_, Apr 18 2013 a(n) = (binomial(j*k*prime(n), j*prime(n)) - binomial(k*j, j))/(k*prime(n)^3) for k=3, j=1 and n>=2 (conjectured). - _Alexander R. Povolotsky_, Apr 18 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander R. Povolotsky_, Mar 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007284 | OEIS-A217774

Palindromic deviation of terms of A006960.

If 196 ever reaches a palindrome (which is an open problem), a(n) will become 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _W. W. Kokko_, Mar 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007285 | OEIS-A217785

Smallest integer s>n such that 1+2*s+3*s^2+...+n*s^{n-1} is prime.

Conjecture: For each n=2,3,... there are infinitely many primes of the form 1+2*s+...+n*s^{n-1}, where s is a positive integer; moreover, we have a(n)<12*n^2. This is related to the following conjecture of the author: The polynomials s_n(x)=sum_{k=0}^n(k+1)x^k (n=1,2,3,...) are all irreducible over the field of rational numbers; moreover, s_n(x) is reducible modulo every prime if and only if n has the form 8k(k+1), where k is a positive integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007286 | OEIS-A217788

Least integer s > p_n such that sum_{k=1}^n p_k*s^(n-k) (the number [p_1,...,p_n] in base s) is prime, where p_k denotes the k-th prime.

Conjecture: For any integers n >= m > 0, there are infinitely many positive integers s > p_n such that the number sum_{k=m}^n p_k*s^{n-k} (i.e., [p_m,...,p_n] in base s) is prime; moreover the smallest such an integer s (denoted by s(m,n)) does not exceed (n+1)*(m+n+1). A related conjecture of the author states that for each n=2,3,... the polynomial sum_{k=1}^n p_k*x^(n-k) is irreducible modulo some prime. See also the author's comments on A000040. The conjecture can be further extended as follows: If a_1 < ... < a_n are distinct integers with a_n prime, then there are infinitely many integers b > a_n such that [a_1,a_2,...,a_n] in base b is prime. See A224197 for a more general conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007287 | OEIS-A217835

Fermat pseudoprimes to base 2 that can be written as p^2*n - p*n + p, where p is also a Fermat pseudoprime to base 2 and n is a positive integer.

Conjecture: For any Fermat pseudoprime p to base 2 there are infinitely many Fermat pseudoprimes to base 2 equal to p^2*n - p*n + p, where n is a positive integer. Conjecture: For any Fermat pseudoprime p to base 2 and any positive integer k, there are infinitely many Fermat pseudoprimes to base 2 equal to p^2*n - p*n + p^k, where n is a positive integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Oct 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007288 | OEIS-A217853

Fermat pseudoprimes to base 3 of the form (3^(4*k + 2) - 1)/8.

Conjecture: There are infinitely many Fermat pseudoprimes to base 3 of the form (3^(4*k + 2) - 1)/8, where k is a natural number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Oct 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007289 | OEIS-A217866

Last time n appears in the first differences of n*log(n): A217865.

Conjecture: the last time n appears is always before the first time n+2 appears.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Oct 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007290 | OEIS-A217876

Triangle read by rows: distribution of adjacent transpositions in involutions.

Conjecture: The asymptotic distribution of adjacent transpositions in involutions is Poisson with mean 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Callan_, Oct 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007291 | OEIS-A217924

a(n) = n! * [x^n] exp(2*exp(x) - x - 2). Row sums of triangle A217537.

Conjecture: a(n) = Sum_{k=0..2^n-1} A372205(k). - _Mikhail Kurkov_, Nov 21 2021 [Rewritten by _Peter Luschny_, Apr 22 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Oct 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007292 | OEIS-A217948

List of numbers 2n for which the riffle permutation permutes all except the first and last of the 2n cards.

Tiago Januario (email, Jan 12 2015; see also reference) conjectures that these terms are always one more than a prime. - _N. J. A. Sloane_, Mar 02 2015 Conjecture: a(n) = A001122(n) + 1 (via an observation by _Jules Beauchamp_). - _Bill McEachen_, Jun 11 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 07 2012, based on an email message from _Anthony C Robin_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007293 | OEIS-A217976

First time n appears in the first differences of n*log(n): A217865.

Conjecture: the last time n appears is always before the first time n+2 appears.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Oct 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007294 | OEIS-A218010

Primes of the form (24*p + 1)/5, where p is a Fermat pseudoprime to base 2.

Conjecture: There is no absolute Fermat pseudoprime m for which n = (5*m - 1)/24 is a natural number (checked for the first 300 Carmichael numbers; if true, then the formula is a criterion to separate pseudoprimes at least from a subset of primes, because there are 37 primes m from the first 300 primes for which n = (5*m - 1)/24 is a natural number). 3380740301 is a counterexample to the conjecture. - _Charles R Greathouse IV_, Dec 07 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Oct 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007295 | OEIS-A218035

Number of n-digit palindromes with squares that are also palindromes.

The conjectures in the formula section are true. Conjecture: a(n) = n^3/48 - n^2/8 + 2n/3 + 1 if n even, see A011826. Conjecture: a(n) = n^3/24 - 3n^2/8 + 59n/24 - 1/8 if n odd, n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Zvirbulis_, Oct 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007296 | OEIS-A218045

Number of truth tables of bracketed formulas (case 3).

Conjecture D-finite with recurrence n*(n-1)*(2*n-3)*a(n) -2*(n-1)*(11*n^2-39*n+35)*a(n-1) +8*(-3*n^3+42*n^2-159*n+175)*a(n-2) +32*(2*n-7)*(8*n^2-61*n+115)*a(n-3) +128*(n-5)*(2*n-7)*(2*n-9)*a(n-4)=0. - _R. J. Mathar_, Dec 02 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007297 | OEIS-A218046

Primes p such that 8p + 2r is a primorial for some r in A006512.

The empirical evidence suggests that each twin prime pair is associated with at least one p, and each p with a twin prime pair. I conjecture that this sequence (and therefore the sequence of twin primes) is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael G. Kaarhus_, Oct 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007298 | OEIS-A218086

Mersenne prime exponents of prime index equal to 1 or another Mersenne prime exponent.

Conjecture: This sequence is complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Raphie Frank_, Oct 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007299 | OEIS-A218147

Degree of minimal polynomial satisfied by exp(8*Pi*phi_2(1/n,1/n)), where phi_2 is defined in the Comments.

Watson Ladd has proved that the sequence satisfies the following recurrence relations, which were conjectured by _Jason Kimberley_:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Kimberley_, Oct 21 2012 and Apr 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007300 | OEIS-A218185

Number of rows with the value true in the truth tables of all bracketed formulas with n distinct propositions p_1, ..., p_n connected by the binary connective of m-implication (case 3).

Conjecture: n*(n-1)*(n-2)*a(n) -12*(2*n-5)*(n-1)*(n-2)*a(n-1) +4 *(n-2)*(52*n^2-312*n+465) *a(n-2) -48*(2*n-7)*(8*n^2-56*n+95)*a(n-3) +256*(n-5) *(2*n-7) *(2*n-9)*a(n-4)=0. - _R. J. Mathar_, Oct 08 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007301 | OEIS-A218225

G.f. A(x) satisfies: (1 - x*A(x)) / (1 - x^2*A(x)^2)^2 = 1 - x.

Conjecture: a(n) is the number of permutations of [1..n+1] that avoid one of the following sets of patterns: (2134, 42153, 24153), (3124, 42153, 24153), (2143, 42135, 24135). - _Alexander Burstein_, Dec 20 2017 Chern et al. (see links) proves the above conjecture for (3124, 42153, 24153) as well as the following conjecture of Hong and Li:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007302 | OEIS-A218249

Difference sequence of A219096.

Each appearance of 1 represents a prime p for which the next 3 larger primes are p+6, p+12, and p+18. More generally, the sequence gives gap sizes, measured as the number of primes minus 1, between consecutive triples (p, p+6, p+12) and (q, q+6, q+12) of consecutive primes. Conjecture: Every positive integer except 2 occurs infinitely many times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007303 | OEIS-A218263

Number of standard Young tableaux of n cells and height >= 3.

Conjecture: (n-6)*(n-3)*(n+1)*a(n) +(-n^3+6*n^2+11*n-36)*a(n-1) -(n-1)*(n^3-4*n^2-21*n+76)*a(n-2) +2*(n-1)*(n-2)*(3*n-19)*a(n-3) +4*(n-5)*(n-1)*(n-2)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Jan 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Oct 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007304 | OEIS-A218275

a(n) is the smallest n-isolated prime, or a(n)=0 if there are no n-isolated primes.

Conjectures. 1) a(n) > 0; 2) a(n)/n is between 2 and 3 or between the smaller and larger member of a twin prime pair.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Zak Seidov_, Oct 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007305 | OEIS-A218279

Let (p(n), p(n)+2) be the n-th twin prime pair. a(n) is the smallest k, such that there is only one prime in the interval (k*p(n), k*(p(n)+2)), or a(n)=0, if there is no such k.

Conjecture: a(n)>0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Oct 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007306 | OEIS-A218333

The index of the smallest n-isolated prime p such that p/n is not between 2 and 3 and not between the smaller and greater primes of a twin prime pair, or 0 if no such p exists.

Conjecture: all a(n) >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Oct 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007307 | OEIS-A218459

a(n) is the smallest positive integer d such that prime(n) = x^2 + dy^2 has a solution (x,y) in integers.

It appears 75% of values are 1 or 2, with the vast majority of the rest prime, though many are duplicates. Conjecture: Odd composite values belong to A176255. - _Bill McEachen_, Sep 03 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alonso del Arte_, Oct 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007308 | OEIS-A218465

Least integer b>2n+1 such that the numbers written as [1,3,...,2n-1,2n+1] and [2n+1,2n-1,...,3,1] in base b are both prime.

Conjecture: Let n be any positive integer. Then a(n) exists, moreover there are infinitely many integers b > 2n+1 such that [1,3,...,2n-1,2n+1] and [2n+1,2n-1,...,3,1] in base b are both prime. Also, the polynomial S_n(x) = sum_{k=0}^n (2k+1)*x^{n-k} is irreducible modulo some prime p < (n+1)(n+2), and the Galois group of S_n(x) over the field of rational numbers is isomorphic to the symmetric group S_n. This conjecture can be extended by replacing 2k+1 by (2k+1)^m. For example, [1^2,3^2,5^2,...,61^2,63^2] and [63^2,61^2,...,3^2,1^2] in base b=241784 are both prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007309 | OEIS-A218468

a(n+1) is a multiple of at least one digit of a(n); this is the lexicographically earliest injective such sequence.

Conjectured to be a permutation of the positive integers, which is the case if and only if an infinite number of terms have a digit "1". Conjecture: For nonnegative integer x, 2*10^x appears at index 2*10^x (see A226179). _Hans Havermann_, May 29 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _M. F. Hasler_, May 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007310 | OEIS-A218483

Fermat pseudoprimes to base 2 which are congruent to 1 (mod 8).

For p = 5 the formula becomes 40*n + 25. From the first 15 pseudoprimes divisible by 5, 12 are of the form 40*n + 25 (beside 3 of them which are of the form 40*n + 5). Conjecture: there are no pseudoprimes to base 2 of the form 40*n + 15. Conjecture: If a Fermat pseudoprime to base 2 can be written as 8*p*n + p^2, where n is an integer and p one of its prime factors, then it can be written this way for any of its prime factors. Checked for all pseudoprimes from the sequence above. Conjecture: If a Fermat pseudoprime to base 2 with two prime factors can be written as 8*p1*n + p1^2, where n is a natural number and p1 one of its two prime factors, then it can also be written as 8*p2*(-n) + p2^2, where p2 is the other prime factor. Checked for 4033 = 37*109(n = 9), 4369 = 17*257(n = 30), 4681 = 31*151(n = 15), 8321 = 53*157(n = 13), 18721 = 97*193(n = 12), 23377 = 97*241(n = 18), 31417 = 89*353(n = 33), 31609 = 73*433 (n = 45), 65281 = 97*673(n = 72), 85489 = 53*1613 (n = 195). Conjecture: If a Fermat pseudoprime to base 2 cannot be written as 8*p*n + p^2, where n is an integer and p one of its prime factors, then it cannot be written this way for any of its prime factors. Checked for the following pseudoprimes: 341, 645, 1387, 2047, 2701, 2821, 3277, 4371, 5461, 7957, 10261, 13741, 13747, 13981, 14491, 15709, 19951, 29341, 31621, 42799, 49141, 49981, 55245, 60701, 60787, 63973, 65077, 68101, 72885, 80581, 83333. All three conjectures are true (obvious from new characterization). - _Charles R Greathouse IV_, Dec 07 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius Coman_, Oct 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007311 | OEIS-A218505

Decimal expansion of Sum_{k>=1} (H(k)/k)^2, where H(k) = Sum_{j=1..k} 1/j.

This series is called "the quadratic series of Au-Yeung", after Enrico Au-Yeung who discovered numerically the value of the sum on April 1993, when he was an undergraduate student in the Faculty of Mathematics in Waterloo. His conjecture was proved by David Borwein and Jonathan Borwein (1995).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Mar 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007312 | OEIS-A218543

Number of times when an odd number is encountered, when going from 2^(n+1)-1 to (2^n)-1 using the iterative process described in A071542.

What is the reason for that smoothness? (Conjecture: The distribution of "tendrils", i.e. finite subtrees in the beanstalk and its almost fractal nature? Cf: A218787.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007313 | OEIS-A218585

Number of ways to write n as x+y with 0<x<=y and x^2+xy+y^2 prime.

Conjecture: a(n)>0 for all n>1 with the only exception n=8. Zhi-Wei Sun also conjectured that (sum_{p<N, p=1(mod 3)}x(p))/(sum_{p<N, p=1(mod 3)}y(p)) tends to 1+sqrt(3), and (sum_{p<N, p=1(mod 3)}x(p)^2)/(sum_{p<N, p=1(mod 3)}y(p)^2) tends to 52/9. Similar conjectures involving p=x^2+y^2 were recently formulated by Thomas Ordowski.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007314 | OEIS-A218616

The infinite trunk of beanstalk (A179016) with reversed subsections.

This contains the terms in the infinite trunk of beanstalk (A179016) listed in partially reversed manner: after the initial zero each subsequence lists A213709(n) successive terms from A179016, descending from (2^n)-1 downwards, usually down to 2^(n-1) (conjectured to indeed be a power of 2 in each case, apart from 2 itself missing from the beginning of the sequence).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007315 | OEIS-A218621

a(n) = unique divisor d of n such that d + (n/d - 1)/2 is minimal and integral.

Conjecture. For n, p, v, j natural numbers, the conditions on a(n) seem to be the following:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Feb 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007316 | OEIS-A218622

a(n) = A183161(n) (mod 4), n>=0.

Conjecture: a(n) never equals 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007317 | OEIS-A218654

Number of ways to write n as x+y with 0<x<=y and x^2+3xy+y^2 prime.

Conjecture: a(n)>0 for all n=2,3,4,... Zhi-Wei Sun also conjectured that These conjectures are similar to the ones mentioned in the comments in A218585.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007318 | OEIS-A218656

Number of ways to write 2n+1 as x+y with 0 < x < y and x^4 + y^4 prime.

Conjecture: a(n) > 0 for all n >= 1. _Thomas Ordowski_ conjectured on Nov 03 2012 that if x^4 + y^4 in the definition of a(n) is replaced by x^2 + y^2, then a(n) will always be positive. We also have similar conjectures with x^4 + y^4 replaced by x^8 + y^8 or x^16 + y^16. All conjectures verified for 2n+1 up to 10^6: no exceptions for x^2 + y^2 and x^4 + y^4; exceptions 2n+1 = 7, 9, 55, 73, 75 and 105 for x^8 + y^8; exceptions 2n+1 = 5 and 9 for x^16 + y^16. - _Mauro Fiorentini_, Sep 22 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007319 | OEIS-A218755

Denominators of Bernoulli numbers which are == 6 (mod 9).

Conjecture: all first differences 36, 288, 180, 180,... of the sequence are multiples of 36. The conjecture is true, since elements of A090801 are 2 mod 4. - _Charles R Greathouse IV_, Nov 22 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Nov 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007320 | OEIS-A218771

Primes of the form p^2 + 3pq + q^2 with p and q prime.

This sequence is of particular interest due to Zhi-Wei Sun's surprising conjecture related to A218754. That conjecture implies that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 05 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007321 | OEIS-A218830

Largest odd integer not of the form p+2q with p, q, p^2+4(2^n-1)q^2 all prime, or 0 if there would be no such upper bound.

Zhi-Wei Sun has conjectured (Nov 07 2012) that for any n>0, there is only a finite number of positive odd integers not of the given form. See arXiv:1211.1588.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_ and _M. F. Hasler_, Nov 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007322 | OEIS-A218832

Number of positive integer solutions to the Diophantine equation x + y + 2z = n^2.

Conjectures from _Colin Barker_, Apr 01 2013: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dennis P. Walsh_, Mar 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007323 | OEIS-A218851

a(n) = 2*floor((n + sin(n)/(2*sin(1/2))) * log(n)) + 1.

This sequence of odd integers mimics the prime numbers and more specifically it is conjectured a(n) satisfies an analog of the k-tuple conjecture (see link) with similar heuristic asymptotic formulas.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007324 | OEIS-A218867

Number of prime pairs {p,q} with p>q and {p-4,q+4} also prime such that p+(1+(n mod 6))q=n if n is not congruent to 4 (mod 6), and p-q=n and q<n/2 if n=4 (mod 6).

Conjecture: a(n)>0 for all n>50000 with n different from 50627, 61127, 66503. This conjecture implies that there are infinitely many cousin prime pairs. It is similar to the conjectures related to A219157 and A219055.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007325 | OEIS-A218976

a(n) is the smallest positive integer such that 10^(2 + floor(k/a(1)) + floor(k/a(2)) + ... + floor(k/a(n))) divides (k+9)! for all k > 0.

Conjecture: All terms except a(5) are 1 mod 5. - _R. J. Cano_, Nov 11 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Cano_ and _Charles R Greathouse IV_, Nov 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007326 | OEIS-A218980

Integers not in A218852.

Conjecture: This sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Nov 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007327 | OEIS-A218981

Numbers n for which sigma(n) = sigma(w) + sigma(x) + sigma(y) + sigma(z), where n = w + x + y + z, with w, x, y, z all positive.

Conjecture: This sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Nov 08 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007328 | OEIS-A219025

Number of primes p<n such that 6n-p and 6n+p are both prime.

Conjecture: a(n)>0 for all n=6,7,... Zhi-Wei Sun also made the following general conjecture: See also A219023 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007329 | OEIS-A219033

Numbers k such that k = x + y, sigma_1(k) = sigma_1(x) + sigma_1(y) and sigma_2(k) = sigma_2(x) + sigma_2(y).

Conjecture: This sequence is infinite. Conjecture: The sequence only consists of even numbers. Conjecture: The partitions only consist of even numbers. Conjecture: None satisfy sigma_3(n) = sigma_3(x) + sigma_3(y). Conjecture: With the lower partition as 6*A185208(n) and the upper partition 214/3 = 71.3333... of this, then the equalities are satisfied.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Nov 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007330 | OEIS-A219037

Numbers k such that k divides 2^k + 2 and (k-1) divides 2^k + 1.

Conjecture: a(n+1) = 2^a(n) + 2 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Nov 10 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007331 | OEIS-A219167

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal or vertical neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..2 n X 2 array.

Conjectures from _Colin Barker_, Mar 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007332 | OEIS-A219197

Self-convolution equals A199033.

Conjecture: a(n) is never congruent to 3 modulo 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007333 | OEIS-A219211

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal, vertical, diagonal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..2 n X 2 array.

Conjectures from _Colin Barker_, Mar 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007334 | OEIS-A219252

Smallest prime q such that 2*n+1 = p + 4*q for some odd prime p, otherwise 0 if no such q exists.

Conjecture: except m = 77, all odd number > 9 are of the form m = p + 4*q where p and q are prime numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007335 | OEIS-A219254

Number of ways to express 2n+1 as p+4q with p, q primes.

This is related to the conjecture given in A219252.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007336 | OEIS-A219286

Number of n X 3 arrays of the minimum value of corresponding elements and their horizontal or vertical neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..1 n X 3 array.

Conjectures from _Colin Barker_, Jul 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007337 | OEIS-A219287

Number of n X 4 arrays of the minimum value of corresponding elements and their horizontal or vertical neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..1 n X 4 array.

Conjectures from _Colin Barker_, Jul 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007338 | OEIS-A219293

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal, vertical or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..2 n X 2 array.

Conjectures from _Colin Barker_, Jul 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007339 | OEIS-A219303

Numbers which do not reach zero under the repeated iteration x -> floor(sqrt(x)) * (x - floor(sqrt(x))^2).

Conjecture #1: All numbers under the iteration reach 0 or, like the elements of this sequence, reach a finite loop, and none expand indefinitely to infinity. Conjecture #2: There are an infinite number of such finite loops, though there is often significant distance between them. Conjecture #3: There are an infinite number of pairs of consecutive integers in this sequence, e.g. 14 and 15, 197 and 198. This argument is strengthened by the fact there are other

groupings such as triples - The first of these is 11527, 11528 and 11529 - and also that for randomly chosen numbers of hundreds of digits, N, the nearest pair or grouping appears to be within N +/- 1000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carl R. White_, Nov 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007340 | OEIS-A219314

Composition of the inverse binomial transform of Fibonacci numbers and the Catalan transform of Fibonacci numbers.

Conjecture: n*a(n) -2*n*a(n-1) +11*(-n+2)*a(n-2) +4*(2*n-5)*a(n-3) +8*(5*n-17)*a(n-4) +24*(n-4)*a(n-5)=0. - _R. J. Mathar_, Jun 14 2016 Conjecture: n*(5*n-7)*a(n) -4*(5*n^2-12*n+6)*a(n-1) -(15*n^2-11*n-30) *a(n-2) +2*(35*n^2-119*n+66)*a(n-3) +12*(n-3)*(5*n2)*a(n-4)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Nov 17 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007341 | OEIS-A219349

Number of n X 3 arrays of the minimum value of corresponding elements and their horizontal, vertical, diagonal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..1 n X 3 array.

Conjectures from _Colin Barker_, Jul 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007342 | OEIS-A219350

Number of n X 4 arrays of the minimum value of corresponding elements and their horizontal, vertical, diagonal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..1 n X 4 array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007343 | OEIS-A219367

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal or vertical neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..3 n X 2 array.

Conjectures from _Colin Barker_, Mar 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007344 | OEIS-A219382

Number of 2 X n arrays of the minimum value of corresponding elements and their horizontal, diagonal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..2 2 X n array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007345 | OEIS-A219431

Logarithmic derivative of the g.f. of the overpartitions of n^2 (A219430).

Conjecture: a(2*n+1) == 2 (mod 4), a(2*n) == 0 (mod 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007346 | OEIS-A219432

Least 3-smooth number k such that prime(n)*k - 1 is prime.

Conjecture: a(n) < prime(n) for n > 21. Conjecture confirmed up to a(122578) = 55296 < prime(122578) = 1620539.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Nov 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007347 | OEIS-A219433

a(n) is the smallest 3-smooth number such that prime(n)*a(n) + 1 is prime.

Conjecture: a(n) < prime(n) except for a(522720). Conjecture tested up to a(1000000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Nov 19 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007348 | OEIS-A219452

First position of a plateau of length n in the product-of-exponents function A005361.

a(8) is conjectured to be 177001925537328703872748486819210922882 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Nov 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007349 | OEIS-A219454

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal, vertical, diagonal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..3 n X 2 array.

Conjectures from _Colin Barker_, Mar 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007350 | OEIS-A219465

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..3 n X 2 array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007351 | OEIS-A219472

Number of 2 X n arrays of the minimum value of corresponding elements and their horizontal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..3 2 X n array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007352 | OEIS-A219498

Number of n X 4 arrays of the minimum value of corresponding elements and their horizontal or vertical neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..1 n X 4 array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007353 | OEIS-A219499

Number of n X 5 arrays of the minimum value of corresponding elements and their horizontal or vertical neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..1 n X 5 array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007354 | OEIS-A219500

Number of n X 6 arrays of the minimum value of corresponding elements and their horizontal or vertical neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..1 n X 6 array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 20 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007355 | OEIS-A219514

Number of n X 3 arrays of the minimum value of corresponding elements and their horizontal, diagonal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..1 n X 3 array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007356 | OEIS-A219515

Number of n X 4 arrays of the minimum value of corresponding elements and their horizontal, diagonal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..1 n X 4 array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007357 | OEIS-A219520

Number of 3 X n arrays of the minimum value of corresponding elements and their horizontal, diagonal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..1 3 X n array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007358 | OEIS-A219529

Coordination sequence for 3.3.4.3.4 Archimedean tiling.

First few terms provided by _Allan C. Wechsler_; Fred Lunnon and Fred Helenius gave the next few; Fred Lunnon suggested that the recurrence was a(n+3) = a(n) + 16 for n > 1. [This conjecture is true - see the CGS-NJAS link for a proof. - _N. J. A. Sloane_, Dec 31 2017] Conjectured to be a(n) = floor((16n+1)/3) for n>0; a(0) = 1; this is a consequence of the suggested recurrence due to Lunnon (see comments). [This conjecture is true - see the CGS-NJAS link in A296368 for a proof. - _N. J. A. Sloane_, Dec 31 2017]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Allan C. Wechsler_, Nov 21 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007359 | OEIS-A219540

a(n) = (n^2 + 2*(Sum_{j = 1..n} j^n)) (mod n^3).

R. Mestrovic conjectures that, for n >= 3, 2*sum(j = 1, n) j^n == -n^2 (mod n^3) implies that n is prime. See proposition 2.2(ii) - the conjecture is proved to be equivalent to Giuga's conjecture. This proves the first half of Mestrovic's conjecture and hence the first half of Giuga's conjecture : If n>2 is prime, a(n)=0. Proof: [1] [(n-k)^n + k^n] mod n^3 with n odd becomes [2] n^2*k^(n-1) mod n^3. k^(n-1) == 1 mod n if n is prime, so we can write k^(n-1) = 1 + jn. Place this back into [2], we find that [2] is just n^2 mod n^3. From [1] we have (n-1)*n^2 = -n^2 mod n^3. Q.E.D. The second half of Mestrovic's conjecture involves proving that if n is composite then a(n)<>0. It is easy enough to prove a(4k+2)<>0 by noticing that the residue must be even on the RHS, but the residue on the LHS is odd. a(8k+4)<>0 by a similar method. - _Jon Perry_, Dec 01 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Nov 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007360 | OEIS-A219543

Denominators of Bernoulli numbers which are congruent to 3 (mod 9).

Conjecture: all the first differences 36, 72, 144, 72,... of the sequence are multiples of 36. The conjecture is true, since elements of A090801 are 2 mod 4. - _Charles R Greathouse IV_, Nov 22 2012

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Nov 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007361 | OEIS-A219552

First differences of A219303.

Conjecture: 1 appears infinitely often in this sequence (indicating an infinite number of pairs of consecutive integers in the parent sequence).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carl R. White_, Nov 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007362 | OEIS-A219558

Number of odd prime pairs {p,q} (p>q) such that p+(1+(n mod 2))q=n and ((p-1-(n mod 2))/q)=((q+1)/p)=1 where (-) denotes the Legendre symbol.

Zhi-Wei Sun has the following general conjecture: s(m) is always positive. In particular, s(0)=1239, Note that s(1)=1470 means that a(n)>0 for all n=1470,1471,... That s(0)=1239 is related to a conjecture of _Olivier Gérard_ and Zhi-Wei Sun.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007363 | OEIS-A219562

a(n) = Sum_{k=0..n} binomial(n+k,k)^4.

Conjecture: a(p-1) == 1 (mod p^5) for prime p >= 7 (checked up to p = 499). Coster, Theorem 4, proves that a(p-1) == 1 (mod p^3) for primes p >= 5. Conjecture: for r >= 2, the supercongruence a(p^r - 1) == a(p^(r-1) - 1) (mod p^(3*r+3)) may hold for all primes p >= 5. Coster, Theorem 4, proves that a(p^r -1) == a(p^(r-1) - 1) (mod p^(3*r)) for r >= 2 and all primes p >= 5. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Nov 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007364 | OEIS-A219572

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal or vertical neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..3 n X 2 array.

Conjectures from _Colin Barker_, Mar 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007365 | OEIS-A219579

Number of 2 X n arrays of the minimum value of corresponding elements and their horizontal or vertical neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..3 2 X n array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007366 | OEIS-A219589

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal or vertical neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..2 n X 2 array.

Conjectures from _Colin Barker_, Mar 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007367 | OEIS-A219590

Number of n X 3 arrays of the minimum value of corresponding elements and their horizontal or vertical neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..2 n X 3 array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007368 | OEIS-A219596

Number of 3 X n arrays of the minimum value of corresponding elements and their horizontal or vertical neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..2 3 X n array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007369 | OEIS-A219604

Smallest prime p such that 2n+1 = 4q + p for some odd prime q, or 0 if no such prime exists.

Conjecture: except m = 77, all odd numbers > 9 are of the form m = p + 4*q where p and q are prime numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007370 | OEIS-A219621

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..2 n X 2 array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007371 | OEIS-A219628

Number of 2 X n arrays of the minimum value of corresponding elements and their horizontal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..2 2 X n array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 24 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007372 | OEIS-A219665

One more than the partial sums of A219661.

Compare to the conjecture given at A213710.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, May 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007373 | OEIS-A219680

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal, vertical or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..2 n X 2 array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007374 | OEIS-A219687

Number of 3 X n arrays of the minimum value of corresponding elements and their horizontal, vertical or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..2 3 X n array.

Conjectures from _Colin Barker_, Jul 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007375 | OEIS-A219699

Number of n X 3 arrays of the minimum value of corresponding elements and their horizontal, vertical or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..1 n X 3 array.

Conjectures from _Colin Barker_, Jul 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007376 | OEIS-A219700

Number of n X 4 arrays of the minimum value of corresponding elements and their horizontal, vertical or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..1 n X 4 array.

Conjectures from _Colin Barker_, Jul 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007377 | OEIS-A219708

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..3 n X 2 array.

Conjectures from _Colin Barker_, Jul 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007378 | OEIS-A219715

Number of 2 X n arrays of the minimum value of corresponding elements and their horizontal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..3 2 X n array.

Conjectures from _Colin Barker_, Jul 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 26 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007379 | OEIS-A219766

Number of nonsquare simple perfect squared rectangles of order n up to symmetry.

In "A Census of Planar Maps", p. 267, William Tutte gave a conjectured asymptotic formula for the number, a(n) of perfect squared rectangles of order n: Conjectured: a(n) ~ n^(-5/2) * 4^n / (243*sqrt(Pi)). [Corrected by _Stuart E Anderson_, Feb 02 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stuart E Anderson_, Nov 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007380 | OEIS-A219768

Number of n X 3 arrays of the minimum value of corresponding elements and their horizontal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..1 n X 3 array.

Conjectures from _Colin Barker_, Jul 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007381 | OEIS-A219769

Number of n X 4 arrays of the minimum value of corresponding elements and their horizontal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..1 n X 4 array.

Conjectures from _Colin Barker_, Jul 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007382 | OEIS-A219774

Number of 3 X n arrays of the minimum value of corresponding elements and their horizontal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..1 3 X n array.

Conjectures from _Colin Barker_, Jul 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007383 | OEIS-A219782

Number of ways to write n=x+y (0<x<=y) with n^2-xy and n^2+xy both prime.

Conjecture: a(n)>0 if n is not among 1, 8, 10, 18, 20, 41, 46, 58, 78, 116, 440. Zhi-Wei Sun also made the following general conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 27 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007384 | OEIS-A219791

Number of ways to write n=x+y (0<x<=y) with (xy)^2+1 prime.

Conjecture: a(n)>0 if n is different from 1, 6, 16, 24. This conjecture has been verified for n up to 10^7. It implies that there are infinitely many primes of the form x^2+1. Zhi-Wei Sun also made the following general conjecture: For any positive integer k, each sufficiently large integer n cna be written as x+y (x>0, y>0) with (xy)^{2^k}+1 prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007385 | OEIS-A219803

Number of 2 X n arrays of the minimum value of corresponding elements and their horizontal, vertical, diagonal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..3 2 X n array.

Conjectures from _Colin Barker_, Jul 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007386 | OEIS-A219810

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal, vertical or diagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..2 n X 2 array.

Conjectures from _Colin Barker_, Jul 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007387 | OEIS-A219817

Number of 3 X n arrays of the minimum value of corresponding elements and their horizontal, vertical or diagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..2 3 X n array.

Conjectures from _Colin Barker_, Jul 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007388 | OEIS-A219838

Number of ways to write n as x + y with 0 < x <= y and (xy)^2 + xy + 1 prime.

Conjecture: a(n) > 0 for all n > 1. Zhi-Wei Sun has made the following general conjecture: For each prime p, any sufficiently large integer n can be written as x + y, where x and y are positive integers with ((x*y)^p - 1)/(x*y - 1) prime. (For p = 5, 7 it suffices to require n > 28 and n > 46 respectively.) Compare this with Sun's another conjecture related to A219791.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007389 | OEIS-A219846

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..2 n X 2 array.

Conjectures from _Colin Barker_, Jul 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007390 | OEIS-A219853

Number of 2 X n arrays of the minimum value of corresponding elements and their horizontal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..2 2 X n array.

Conjectures from _Colin Barker_, Jul 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 29 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007391 | OEIS-A219864

Number of ways to write n as p+q with p and 2pq+1 both prime.

Conjecture: a(n)>0 for all n>7. Zhi-Wei Sun also made the following general conjecture: For each odd integer m not congruent to 5 modulo 6, any sufficiently large integer n can be written as p+q with p and 2*p*q+m both prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007392 | OEIS-A219884

Number of 3 X n arrays of the minimum value of corresponding elements and their horizontal, diagonal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..2 3 X n array.

Conjectures from _Colin Barker_, Jul 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 30 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007393 | OEIS-A219916

Number of 3 X n arrays of the minimum value of corresponding elements and their horizontal, diagonal, diagonal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..2 3 X n array.

Conjectures from _Colin Barker_, Jul 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007394 | OEIS-A219923

Number of ways to write n=x+y (x>0, y>0) with x-1, x+1 and 2*x*y+1 all prime.

Conjecture: a(n)>0 for all n>623. Zhi-Wei Sun made the following general conjecture: For each nonnegative integer m, any sufficiently large integer n can be written as x+y (x>0, y>0) with x-m, x+m and 2*x*y+1 all prime. Sun also conjectured that for each m=0,1,2,... any sufficiently large integer n with m or n odd can be written as x+y (x>0, y>0) with x-m, x+m and x*y-1 all prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007395 | OEIS-A219940

Number of 2 X n arrays of the minimum value of corresponding elements and their horizontal or diagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..3 2 X n array.

Conjectures from _Colin Barker_, Jul 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 01 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007396 | OEIS-A219960

Numbers which do not reach zero under the repeated iteration x -> ceiling(sqrt(x)) * (ceiling(sqrt(x))^2 - x).

Conjecture #1: All numbers under the iteration reach 0 or, like the elements of this sequence, reach a finite loop, and none expand indefinitely to infinity. Conjecture #2: There are an infinite number of such finite loops, though there is often significant distance between them. Conjecture #3: There are an infinite number of pairs of consecutive integers in this sequence despite being less abundant than in A219303.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carl R. White_, Dec 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007397 | OEIS-A219961

First differences of A219960.

Conjecture: 1 appears infinitely often in this sequence (indicating an infinite number of pairs of consecutive integers in the parent sequence), despite very few appearances compared with A219552, which is derived from a very similar process.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carl R. White_, Dec 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007398 | OEIS-A219963

Numbers which do not reach zero under either of the iterations: x -> floor(sqrt(x)) * (x - floor(sqrt(x))^2) or y -> ceiling(sqrt(y)) * (ceiling(sqrt(y))^2 - y).

It is possible, assuming the infinite-pairs conjectures are true for both parent sequences, that there may also be an infinite number of pairs here, but even then that is not guaranteed.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carl R. White_, Dec 02 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007399 | OEIS-A220004

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal, vertical or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..3 n X 2 array.

Conjectures from _Colin Barker_, Jul 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007400 | OEIS-A220011

Number of 2 X n arrays of the minimum value of corresponding elements and their horizontal, vertical or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..3 2 X n array.

Conjectures from _Colin Barker_, Jul 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007401 | OEIS-A220029

Number of n X 5 arrays of the minimum value of corresponding elements and their horizontal or diagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..1 n X 5 array.

Conjectures from _Colin Barker_, Jul 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007402 | OEIS-A220030

Number of n X 6 arrays of the minimum value of corresponding elements and their horizontal or diagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..1 n X 6 array.

Conjectures from _Colin Barker_, Jul 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007403 | OEIS-A220031

Number of n X 7 arrays of the minimum value of corresponding elements and their horizontal or diagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..1 n X 7 array.

Conjectures from _Colin Barker_, Jul 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007404 | OEIS-A220033

Number of 3 X n arrays of the minimum value of corresponding elements and their horizontal or diagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..1 3 X n array.

Conjectures from _Colin Barker_, Jul 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007405 | OEIS-A220034

Number of 4 X n arrays of the minimum value of corresponding elements and their horizontal or diagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..1 4 X n array.

Conjectures from _Colin Barker_, Jul 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007406 | OEIS-A220035

Number of 5 X n arrays of the minimum value of corresponding elements and their horizontal or diagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..1 5 X n array.

Conjectures from _Colin Barker_, Jul 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007407 | OEIS-A220036

Number of 6 X n arrays of the minimum value of corresponding elements and their horizontal or diagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..1 6 X n array.

Conjectures from _Colin Barker_, Jul 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007408 | OEIS-A220037

Number of 7 X n arrays of the minimum value of corresponding elements and their horizontal or diagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..1 7 X n array.

Conjectures from _Colin Barker_, Jul 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007409 | OEIS-A220045

Number of 2 X n arrays of the minimum value of corresponding elements and their horizontal, diagonal or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..3 2 X n array.

Conjectures from _Colin Barker_, Jul 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007410 | OEIS-A220072

Least prime p such that sum_{k=0}^n A005117(k+1)*x^{n-k} is irreducible modulo p.

Conjecture: For any n>0, we have a(n) <= n*(n+1), and the Galois group of SF_n(x) = sum_{k=0}^n A005117(k+1)*x^{n-k} over the rationals is isomorphic to the symmetric group S_n. For another related conjecture, see the author's comment on A005117.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007411 | OEIS-A220095

Numbers k such that there are no primes between k - sqrt(k) and k.

Conjecture: This sequence is complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Dec 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007412 | OEIS-A220097

Number of words on {1,1,2,2,3,3,...,n,n} avoiding the pattern 123.

Conjecture: a(n) = (2/Pi)*Integral_{t=0..1} sqrt((1 - t)/t)*(16*t^2 - 4*t)^n = Catalan(2*n)*2F1(-1-2*n,-n;1/2-2*n;1/4). - _Benedict W. J. Irwin_, Oct 05 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lara Pudwell_, Dec 04 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007413 | OEIS-A220147

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal or diagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..2 n X 2 array.

Conjectures from _Colin Barker_, Jul 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 06 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007414 | OEIS-A220148

Number of n X 3 arrays of the minimum value of corresponding elements and their horizontal or diagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..2 n X 3 array.

Conjectures from _Colin Barker_, Jul 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 06 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007415 | OEIS-A220154

Number of 2 X n arrays of the minimum value of corresponding elements and their horizontal or diagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..2 2 X n array.

Conjectures from _Colin Barker_, Mar 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 06 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007416 | OEIS-A220167

Number of simple squared rectangles of order n up to symmetry.

Conjecture: a(n) ~ n^(-5/2) * 4^n / (243*sqrt(Pi)), from "A Census of Planar Maps", p. 267, where William Tutte gave a conjectured asymptotic formula for the number of perfect squared rectangles where n is the number of elements in the dissection (the order). [Corrected by _Stuart E Anderson_, Feb 03 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stuart E Anderson_, Dec 06 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007417 | OEIS-A220198

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal, vertical or diagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..3 n X 2 array.

Conjectures from _Colin Barker_, Jul 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007418 | OEIS-A220205

Number of 2 X n arrays of the minimum value of corresponding elements and their horizontal, vertical or diagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and nonincreasing columns, 0..3 2 X n array.

Conjectures from _Colin Barker_, Jul 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 07 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007419 | OEIS-A220272

Number of ways to write n=x^2+y (x>0, y>0) with 2*x*y-1 prime.

Conjecture: a(n)>0 for all n>2. The author guessed that the conjecture can be slightly refined as follows: Any integer n>2 can be written as x^2+y with 2*x*y-1 prime, where x and y are positive integers with x<=y. Zhi-Wei Sun also made the following general conjecture: If m is a positive integer and r is 1 or -1, then any sufficiently large integer n can be written as x^2+y (x>0, y>0) with m*x*y+r prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 09 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007420 | OEIS-A220351

Decimal expansion of (3*sqrt(3)+sqrt(7))/10.

Smith & Smith conjecture that this is the Steiner ratio rho_3, the least upper bound on the ratio of the length of the Steiner minimal tree to the length of the minimal tree in dimension 3. Diaconis & Graham offer $1000 for proof (or disproof) of this conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Dec 11 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007421 | OEIS-A220365

a(n) is conjectured to be the largest power k for which k divides the sum of digits of n^k.

a(n) is conjectured to be the largest power k for which k divides the sum of digits of n^k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Dec 12 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007422 | OEIS-A220413

Number of ways to write n=x+y (x>=0, y>=0) with x^3+2*y^3 prime.

Conjecture: a(n)>0 for every n=1,2,3,... Moreover, any integer n>3 not among 7, 22, 31 can be written as p+q (q>0) with p and p^3+2*q^3 both prime. We have verified this conjecture for n up to 10^8. D. R. Heath-Brown proved in 2001 that there are infinitely many primes in the form x^3+2*y^3, where x and y are positive integers. Zhi-Wei Sun also made the following general conjecture: For each positive odd integer m, any sufficiently large integer n can be written as x+y (x>=0, y>=0) with x^m+2*y^m prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 13 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007423 | OEIS-A220420

Express the Sum_{n>=0} p(n)*x^n, where p(n) is the partition function, as a product Product_{k>=1} (1 + a(k)*x^k).

Alkauskas (2016, Problem 3, p. 3) conjectured that a(8*k+2), a(8*k+4), and a(8*k+6) are all negative, and a(8*k) is positive for k >= 1. [This statement is not wholly true for k = 0.] - _Petros Hadjicostas_, Oct 07 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Dec 14 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007424 | OEIS-A220433

Expansion of e.g.f. for operads for alia algebras.

Conjecture: D-finite with recurrence 4*a(n) +6*(-2*n+3)*a(n-1) -(3*n-5)*(3*n-7)*a(n-2)=0. - _R. J. Mathar_, Feb 27 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007425 | OEIS-A220455

Number of ways to write n=x+y (x>0, y>0) with 3x-2, 3x+2 and 2xy+1 all prime.

Conjecture: a(n)>0 for all n>7. Conjecture verified for n up to 10^9. - _Mauro Fiorentini_, Aug 06 2023 Zhi-Wei Sun also made some other similar conjectures, e.g., he conjectured that any integer n>17 can be written as x+y (x>0, y>0) with 2x-3, 2x+3 and 2xy+1 all prime, and each integer n>28 can be written as x+y (x>0, y>0) with 2x+1, 2y-1 and 2xy+1 all prime. Both conjectures verified for n up to 10^9. - _Mauro Fiorentini_, Aug 06 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007426 | OEIS-A220554

Number of ways to write 2n = p+q (q>0) with p, 2p+1 and (p-1)^2+q^2 all prime.

Conjecture: a(n)>0 for all n>1. Zhi-Wei Sun also made the following related conjectures: All conjectures verified for n up to 10^9. - _Mauro Fiorentini_, Sep 21 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 15 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007427 | OEIS-A220572

Number of ways to write 2n-1=x+y (x,y>=0) with x^18+3*y^18 prime.

Conjecture: a(n)>0 for every n=1,2,3,.... Moreover, any odd integer greater than 2092 can be written as x+y (x,y>0) with x-3, x+3 and x^18+3*y^18 all prime. Zhi-Wei Sun also made the following general conjecture: For each positive integer m, any sufficiently large odd integer n can be written as x+y (x,y>0) with x-3, x+3 and x^m+3*y^m all prime (and hence there are infinitely many primes in the form x^m+3*y^m). In particular, for m = 1, 2, 3, 4, 5, 6, 18 any odd integer greater than one can be written as x+y (x,y>0) with x^m+3*y^m prime, and for m =1, 2, 3 any odd integer n>15 can be written as x+y (x,y>0) with x-3, x+3 and x^m+3*y^m all prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 16 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007428 | OEIS-A220670

Coefficient triangle for powers of x^2 of polynomials appearing in a generalized Melham conjecture on alternating sums of third powers of Chebyshev's S polynomials with odd indices. Coefficients in powers of x^2 of 2 + (-1)^n*S(2*n,x).

Coefficient triangle for powers of x^2 of polynomials appearing in a generalized Melham conjecture on alternating sums of third powers of Chebyshev's S polynomials with odd indices. Coefficients in powers of x^2 of 2 + (-1)^n*S(2*n,x). For the original Melham conjecture on sums of odd powers of even-indexed Fibonacci numbers see the references given in A217475. See especially the Wang and Zhang reference given there. An analog conjecture stated for Chebyshev's S polynomials (see A049310) is (Product_{j=0..m} tau(j,x))*(Sum_{k=0..n} ((-1)^k)*(S(2*k-1,x)/x)^(2*m+1))/(P(n,x^2)^2) = H(m,n,x^2), with P(n,x^2) := (1 - (-1)^n*S(2*n,x))/x^2 and certain integer polynomials H with degree (2*m-1)*n + binomial(m-1,2) in x^2. The coefficients of powers of x^2 of the monic integer polynomials tau(n,x) := 2*T(2*n+1,x/2)/x, with Chebyshev's T polynomials, are given by the signed A111125 triangle (see a comment there from Oct 23 2012). The coefficients of the powers of x^(2*j) of the polynomials P(n,x^2) are found in (-1)^(n-1)*A109954(n-1,j). Here the conjecture is considered for m=1 (third powers): H(1,n,x^2) = Sum_{p=0..n} a(n,p)*x^(2*p), n >= 1. It is conjectured that in fact H(1,n,x^2) = 2 + (-1)^n*S(2*n,x). This has been checked by Maple for n=1..100. Therefore we have added a(0,0) = 3 (in the conjecture above this would be the undetermined 0/0). The original Melham conjecture for m=1 (third powers), appears by putting x = i (the imaginary unit): 1*4*sum(F(2*k)^3)/(1-F(2*n+1))^2 = sum(a(n,p)*(-1)^p) = 2 + F(2*n+1) (the unsigned row sums of the present triangle). This m=1 identity is, of course, proved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jan 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007429 | OEIS-A220671

Coefficient array for powers of x^2 of polynomials appearing in a generalized Melham conjecture on alternating sums of fifth powers of Chebyshev S polynomials with odd indices.

Coefficient array for powers of x^2 of polynomials appearing in a generalized Melham conjecture on alternating sums of fifth powers of Chebyshev S polynomials with odd indices. For the generalized Melham conjecture and links to the references concerned with the Melham conjecture on sums of fifth powers of even-indexed Fibonacci numbers see a comment under A220670. Here the conjecture is considered for m=2 (fifth powers): H(2,n,x^2):= product(tau(j,x), j=0..2) * sum(((-1)^k)*(S(2*k-1,x)/x)^5, k=0..n) / (P(n,x^2)^2), with P(n,x^2):= (1 - (-1)^n*S(2*n,x))/x^2. For tau(j,x):= 2*T(2*j+1,x/2)/x, with Chebyshev's T polynomials see a Oct 23 2012 comment on A111125. For the polynomials P see signed A109954. The conjecture is that H(2,n,x^2) is an integer polynomial of degree 3*n: H(2,n,x^2) = sum(a(n,p)*x^(2*p), p=0..3*n), n >= 1. If one puts x = i (the imaginary unit) one obtains the original Melham conjecture for Fibonacci numbers F = A000045. H(2,n,-1) = +44*sum(F(2*k)^5,k=0..n)/(1+F(2*n+1))^2, n>=1, which is conjectured to be -14 - 3*y(n) + 8*y(n)^2 + 4*y(n)^3, with y(n):=F(2*n+1) (see row m=2 of A217475). It is conjectured that H(2,n,x^2) = h(2,n,x^2) - 3*z(n) + 8*z(n)^2 + 4*z(n)^3, with z(n):= ((-1)^n)*S(2*n,x), with h an integer polynomial of degree 3*n. See A220672 for the coefficients of h(2,n,x^2) for n = 1..5. Because h(2,n,-1) = -14 by the usual Melham conjecture, we put h(2,0,x^2) = -14.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jan 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007430 | OEIS-A220672

Coefficients of powers of x^2 of polynomials, called h(2,n,x^2), appearing in a conjecture on alternating sums of fifth powers of odd-indexed Chebyshev S polynomials stated in A220671.

Coefficients of powers of x^2 of polynomials, called h(2,n,x^2), appearing in a conjecture on alternating sums of fifth powers of odd-indexed Chebyshev S polynomials stated in A220671. A generalized Melham conjecture involving fifth powers (m=2) of odd-indexed Chebyshev S polynomials (see A049310) is H(2,n,x^2):= (x^2-3)*(x^4-5*x^2+5)*sum(((-1)^k)*(S(2*k-1,x)/x)^(2*m+1), k=0..n)/((1 - (-1)^n*S(2*n,x))/x^2)^2 = h(2,n,x^2) - 3*z(n) + 8*z(n)^2 + 4*z(n)^3, with z(n):= ((-1)^n)*S(2*n,x), and h an integer polynomial of degree 3*n. The present array a(n,p) appears as h(2,n,x^2) = sum(a(n,p)*x^(2*p),p=0..3*n), n >= 1. The entry a(0,0) := -14 has been used because, in accordance with the original Melham conjecture (see a comment on A220671), h(2,n,i^2), with the imaginary unit i, is conjectured to be -14, for all n >= 1. a(n,p) = [x^(2p)] h(0,2,n,x^2), with the polynomial h defined above in a comment. The conjecture is that h is an integer polynomial of degree 3n in x^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jan 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007431 | OEIS-A220749

Equals one maps: number of n X 4 binary arrays indicating the locations of corresponding elements equal to exactly one of their horizontal and vertical neighbors in a random 0..3 n X 4 array.

Conjectures from _Colin Barker_, Aug 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 18 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007432 | OEIS-A220752

Terms of A220698 that appear in A224218.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Apr 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007433 | OEIS-A220813

The elements of the set P3 in ascending order.

It is conjectured that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franz Vrabec_, Dec 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007434 | OEIS-A220814

The elements of the set P5 in ascending order.

It is conjectured that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franz Vrabec_, Dec 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007435 | OEIS-A220815

The elements of the set P7 in ascending order.

It is conjectured that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franz Vrabec_, Dec 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007436 | OEIS-A220846

a(n) = sum_(d|n) ((product_(d|n) d) / d).

Conjecture: sequence is injective (all terms of this sequence occur exactly once).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Dec 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007437 | OEIS-A220847

a(n) = numerator of the fraction whose Engel expansion has the positive divisors of n as its terms.

Conjecture: sequence is injective (all terms of this sequence occur only once).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Dec 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007438 | OEIS-A220848

a(n) = sum_(d|n) product_(d_x|n, d_x<=d) d_x.

Conjecture: sequence is injective (all terms of this sequence occur only once). The conjecture is true for the first 100000 terms. - _Harvey P. Dale_, Dec 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Dec 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007439 | OEIS-A220849

a(n) = Product_{d|n} Product_{d_x|n , d_x <= d} d_x.

Conjecture: sequence is injective (all terms of this sequence occur only once).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Dec 22 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007440 | OEIS-A220853

Denominators of the fraction (30*n+7) * binomial(2*n,n)^2 * 2F1([1/2 - n/2, -n/2], [1], 64)/(-256)^n, where 2F1 is the hypergeometric function.

Conjectures from _Alexander R. Povolotsky_, Feb 27 2013: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander R. Povolotsky_, Dec 23 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007441 | OEIS-A220898

Number of primitive maps on n edges.

Conjecture: 2*(n+1)*(2*n+1)*a(n) +(-15*n^2+19*n-8)*a(n-1) -2*(21*n-22)*(n-2)*a(n-2) -23*(n-2)*(n-3)*a(n-3)=0. - _R. J. Mathar_, May 30 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007442 | OEIS-A220899

Number of 2-face-free maps on n edges.

Conjecture: 2*(n+1)*(2*n+1)*a(n) +(-23*n^2+31*n-6)*a(n-1) +2*(-4*n^2-9*n+25)*a(n-2) +2*(23*n^2-142*n+174)*a(n-3) +4*(n+8)*(n-5)*a(n-4) -23*(n-5)*(n-6)*a(n-5)=0. - _R. J. Mathar_, Jul 17 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007443 | OEIS-A220902

a(n) = Catalan(n) - A000245(n-2).

Conjecture: (n+1)*a(n) + (-5*n+3)*a(n-1) + (5*n-13)*a(n-2) + 2*(-2*n+9)*a(n-3) = 0. - _R. J. Mathar_, May 30 2014 Mathar's conjecture can be verified by expressing a in terms of factorials and simplifying.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007444 | OEIS-A220926

Number of n X 2 arrays of the minimum value of corresponding elements and their horizontal, vertical or antidiagonal neighbors in a random, but sorted with lexicographically nondecreasing rows and columns, 0..3 n X 2 array.

Conjectures from _Colin Barker_, Aug 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007445 | OEIS-A220932

Equals two maps: number of n X 3 binary arrays indicating the locations of corresponding elements equal to exactly two of their horizontal and antidiagonal neighbors in a random 0..2 n X 3 array.

Conjectures from _Colin Barker_, Mar 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 25 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007446 | OEIS-A220947

Least prime p such that sum_{k=0}^n F(k+1)*x^{n-k} is irreducible modulo p, where F(j) denotes the Fibonacci number A000045(j).

Conjecture: a(n) <= n^2+12 for all n>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007447 | OEIS-A220949

Least prime p such that sum_{k=0}^n (2k+1)*x^(n-k) is irreducible modulo p.

Conjecture: a(n) <= n^2+22 for all n>0. We have similar conjectures with 2k+1 in the definition replaced by (2k+1)^m (m=2,3,...).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007448 | OEIS-A220951

Primes p such that p+6 is also prime and there is a power of two in the interval (p,p+6).

A search for sexy primes bracketing a power of two was conducted up to 2^1500. It is conjectured that this is a finite sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brad Clardy_, Feb 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007449 | OEIS-A220956

(Binomial(2n, n) - binomial(2n - 2, n - 1)) (mod n^2) - n - 2.

Conjecture: a(n) = 0 iff n is an odd prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary Detlefs_ and _Robert G. Wilson v_, Feb 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007450 | OEIS-A221018

Number k such that Fibonacci(k) is divisible by k, k + 1 and k - 1.

a(n) % 10 == 0, 2, or 8 (conjectured). - _Chai Wah Wu_, Aug 06 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, May 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007451 | OEIS-A221022

Equals one maps: number of n X 4 binary arrays indicating the locations of corresponding elements equal to exactly one of their horizontal, vertical and antidiagonal neighbors in a random 0..3 n X 4 array.

Conjectures from _Colin Barker_, Aug 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 28 2012 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007452 | OEIS-A221082

Number of n X 3 arrays of occupancy after each element moves to some horizontal or antidiagonal neighbor.

Conjectures from _Colin Barker_, Aug 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007453 | OEIS-A221121

Number of n X 3 arrays of occupancy after each element stays put or moves to some horizontal or antidiagonal neighbor.

Conjectures from _Colin Barker_, Aug 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007454 | OEIS-A221211

Numbers n such that n and n + 4 are prime and there is a power of two in the interval (n,n+4).

It is a conjecture that this is a finite sequence. These may be the only known cousin primes with this property.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brad Clardy_, Feb 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007455 | OEIS-A221212

Rounded area of distinct right triangles appearing in the unit golden spiral.

Conjecture: a(n) = 3*a(n-2)-a(n-4) for n>5. G.f.: -x*(x+1)*(x^7-x^6-x^5+x^4+x^3+x^2-2*x-1) / ((x^2-x-1)*(x^2+x-1)). - _Colin Barker_, Mar 12 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Feb 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007456 | OEIS-A221213

Conjectured total number of times that k-n appears in the Collatz (3x+1) sequence of k for k = 1, 2, 3,....

Conjectured total number of times that k-n appears in the Collatz (3x+1) sequence of k for k = 1, 2, 3,....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jayanta Basu_, Feb 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007457 | OEIS-A221218

Let sequence B_n={b_m} be defined by: b_1=prime(n), b_2=prime(n+1); for m>=3, b_m=b_(m-2)+b_(m-1) if b_(m-2)+b_(m-1) is not semiprime, otherwise b_m is the least prime divisor of b_(m-2)+b_(m-1). Then a(n) is the maximal term of sequence B_n, or a(n)=0 if B_n is unbounded.

Conjecture: All a(n)>=570. Conjecture: All sequences B_n are eventually periodic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Feb 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007458 | OEIS-A221280

Numbers m such that lambda(m) = lambda(m+1), where lambda(n) = A008836(n) is the Liouville function.

Sequence is infinite. Density is conjecturally 1/2; it is easy to show that the density is at least 1/4 (one of k, 2k, and 2k+1 is in this sequence).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Jan 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007459 | OEIS-A221374

Number of n X 2 arrays of occupancy after each element stays put or moves to some horizontal, vertical or antidiagonal neighbor, with no occupancy greater than 2.

Conjectures from _Colin Barker_, Aug 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007460 | OEIS-A221464

Number of 0..n arrays of length 5 with each element unequal to at least one neighbor, starting with 0.

Conjectures from _Colin Barker_, Aug 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007461 | OEIS-A221465

Number of 0..n arrays of length 6 with each element unequal to at least one neighbor, starting with 0.

Conjectures from _Colin Barker_, Aug 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007462 | OEIS-A221466

Number of 0..n arrays of length 7 with each element unequal to at least one neighbor, starting with 0.

Conjectures from _Colin Barker_, Aug 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007463 | OEIS-A221471

Integers n such that n^2 is the difference of two Lucas numbers (A000032).

Conjecture: a(n) = 3*a(n-1)-a(n-2) = A002878(n-8) for n>13. G.f.: x^2*(28*x^11-66*x^10-16*x^9-2*x^8-13*x^7-2*x^6-5*x^5-4*x^4-3*x^3-2*x^2-x+1) / (x^2-3*x+1). [_Colin Barker_, Feb 17 2013]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Feb 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007464 | OEIS-A221516

Number of 0..n arrays of length 5 with each element differing from at least one neighbor by 2 or more, starting with 0.

Conjectures from _Colin Barker_, Aug 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007465 | OEIS-A221517

Number of 0..n arrays of length 6 with each element differing from at least one neighbor by 2 or more, starting with 0.

Conjectures from _Colin Barker_, Aug 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007466 | OEIS-A221518

Number of 0..n arrays of length 7 with each element differing from at least one neighbor by 2 or more, starting with 0.

Conjectures from _Colin Barker_, Aug 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007467 | OEIS-A221525

Number of 0..n arrays of length 5 with each element differing from at least one neighbor by 2 or more.

Conjectures from _Colin Barker_, Aug 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007468 | OEIS-A221526

Number of 0..n arrays of length 6 with each element differing from at least one neighbor by 2 or more.

Conjectures from _Colin Barker_, Aug 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007469 | OEIS-A221527

Number of 0..n arrays of length 7 with each element differing from at least one neighbor by 2 or more.

Conjectures from _Colin Barker_, Aug 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007470 | OEIS-A221543

Number of 0..n arrays of length 5 with each element differing from at least one neighbor by something other than 1, starting with 0.

Conjectures from _Colin Barker_, Aug 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007471 | OEIS-A221544

Number of 0..n arrays of length 6 with each element differing from at least one neighbor by something other than 1, starting with 0.

Conjectures from _Colin Barker_, Aug 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007472 | OEIS-A221545

Number of 0..n arrays of length 7 with each element differing from at least one neighbor by something other than 1, starting with 0.

Conjectures from _Colin Barker_, Aug 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007473 | OEIS-A221564

The number of FO4C3 moves required to restore a packet of n playing cards to its original state (order and orientation), where each move Flips Over the top four (4) as a unit and then Cuts three (3) cards from the top to bottom.

Conjecture: a(3k+1) = 2k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colm Mulcahy_, May 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007474 | OEIS-A221574

Number of 0..n arrays of length 3 with each element differing from at least one neighbor by something other than 1.

Conjectures from _Colin Barker_, Aug 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007475 | OEIS-A221575

Number of 0..n arrays of length 5 with each element differing from at least one neighbor by something other than 1.

Conjectures from _Colin Barker_, Aug 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007476 | OEIS-A221576

Number of 0..n arrays of length 6 with each element differing from at least one neighbor by something other than 1.

Conjectures from _Colin Barker_, Aug 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007477 | OEIS-A221577

Number of 0..n arrays of length 7 with each element differing from at least one neighbor by something other than 1.

Conjectures from _Colin Barker_, Aug 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007478 | OEIS-A221588

Equals one maps: number of n X 3 binary arrays indicating the locations of corresponding elements equal to exactly one of their king-move neighbors in a random 0..3 n X 3 array.

Conjectures from _Colin Barker_, Aug 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007479 | OEIS-A221589

Equals one maps: number of n X 4 binary arrays indicating the locations of corresponding elements equal to exactly one of their king-move neighbors in a random 0..3 n X 4 array.

Conjectures from _Colin Barker_, Aug 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007480 | OEIS-A221597

Number of 0..n arrays of length 5 with each element differing from at least one neighbor by 1 or less.

Conjectures from _Colin Barker_, Aug 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007481 | OEIS-A221598

Number of 0..n arrays of length 6 with each element differing from at least one neighbor by 1 or less.

Conjectures from _Colin Barker_, Aug 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007482 | OEIS-A221599

Number of 0..n arrays of length 7 with each element differing from at least one neighbor by 1 or less.

Conjectures from _Colin Barker_, Aug 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007483 | OEIS-A221619

Number of n X 4 arrays with each row a permutation of 1..4 having at least as many downsteps as the preceding row.

Conjectures from _Colin Barker_, Aug 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007484 | OEIS-A221650

Tetrahedron P(n,j,k) = T(j,k)*p(n-j), where T(j,k) = 1 if k divides j otherwise 0.

Conjecture 1: P(n,j,k) is the number of partitions of n that contain at least m parts of size k, where m = j/k, if k divides j otherwise P(n,j,k) = 0. Conjecture 2: P(n,j,k) is the number of parts that are the m-th part of size k in all partitions of n, where m = j/k, if k divides j otherwise P(n,j,k) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jan 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007485 | OEIS-A221671

Maximum number of squares in a non-constant arithmetic progression (AP) of length n.

González-Jiménez and Xarles (2013) compute a(n) up to a(52) = 12 using elliptic curves (see Table 2, where their Q(N) = a(N)). They do not seem to have noticed that a(n) = A193832(n) for n != 5 in the range where they computed a(n). I conjecture that this formula holds for all n != 5. Bombieri & Zannier prove that a(n) << n^(3/5) (log n)^c for some constant c > 0. It is conjectured that a(n) ~ sqrt(8n/3). - _Charles R Greathouse IV_, Jan 21 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Jan 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007486 | OEIS-A221672

Length of shortest non-constant arithmetic progression (AP) containing n squares.

González-Jiménez and Xarles (2013) conjecture that for n >= 5 the sequence a(n)-1 equals the tail 7, 12, 15, 22, 26, 35, 40, 51, ... of A001318 (generalized pentagonal numbers k*(3*k-1)/2 for k = 0, +-1, +-2, ...). They prove it up to a(12)-1 = 51 = 6*(3*6-1)/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Jan 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007487 | OEIS-A221685

Number of 0..n arrays of length 6 with each element differing from at least one neighbor by 1 or less, starting with 0.

Conjectures from _Colin Barker_, Aug 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007488 | OEIS-A221686

Number of 0..n arrays of length 7 with each element differing from at least one neighbor by 1 or less, starting with 0.

Conjectures from _Colin Barker_, Aug 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007489 | OEIS-A221701

Let K be a local ring with a principal maximal ideal J of nilpotent degree 2 with |K/J|>2; a(n) = number of D-invariant ideals in the ring R_n(K,J).

Conjecture: (n+4) *(127*n^2 -82004*n +181245)*a(n) +(127*n^3 +983454*n^2 +469493*n -5725578) *a(n-1) +2*(-3556*n^3 -1976261*n^2 +1650923*n +5023392) *a(n-2) +8*(2*n-3) *(1143*n^2 +332375*n -209308)*a(n-3)=0. - _R. J. Mathar_, Mar 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007490 | OEIS-A221703

Let K be a local ring with a principal maximal ideal J of nilpotent degree 3 with |K/J|>2; a(n) = number of D-invariant ideals in the ring R_n(K,J).

Conjecture: (n+2) *(101496*n^2 -9493711*n +47969511) *a(n) +8 *(12687*n^3 +14002447*n^2 -60550969*n -45758370) *a(n-1) +4*(-1420944*n^3 -111890800*n^2 +622530778*n -414036471) *a(n-2) +48*(2*n-7) *(152244*n^2 +6705204*n -18873683) *a(n-3)=0. - _R. J. Mathar_, Mar 15 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007491 | OEIS-A221731

Number of n X 2 arrays of occupancy after each element stays put or moves to some horizontal or antidiagonal neighbor, without move-in move-out left turns.

Conjectures from _Colin Barker_, Mar 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007492 | OEIS-A221764

Number of n X 3 arrays of occupancy after each element moves to some horizontal or vertical neighbor, without 2-loops or left turns.

Conjectures from _Colin Barker_, Aug 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007493 | OEIS-A221788

Number of 3 X n arrays of occupancy after each element moves to some horizontal or antidiagonal neighbor, without 2-loops or left turns.

Conjectures from _Colin Barker_, Aug 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007494 | OEIS-A221829

Number of 2 X n arrays of occupancy after each element moves to some horizontal, vertical or antidiagonal neighbor, without 2-loops or left turns.

Conjectures from _Colin Barker_, Aug 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007495 | OEIS-A221834

G.f.: Sum_{n>=1} x^n * (1-x^n)^(n-1) / (1-x)^(n-1).

Conjecture: a(n) is the number of compositions of n if all single instances of the part 1 are frozen ([1]). Example: The compositions enumerated by a(5) = 13 are 5; 4,[1]; 3,2; 2,3; 3,1,1; 1,3,1; 1,1,3; 2,2,[1]; 2,1,1,1; 1,2,1,1; 1,1,2,1; 1,1,1,2; 1,1,1,1,1. - _Gregory L. Simay_, Oct 27 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007496 | OEIS-A221858

a(1)=2, a(2)=3, for n>=3, a(n) is the n-th number which is obtained by application of Eratosthenes-like sieve (with removing 1's) to sequence: odd part of digit sum of 2^m, m>=1.

We conjecture that every term is prime; moreover, we conjecture that the sequence is a permutation of the sequence of all primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007497 | OEIS-A221968

Number of -n..n arrays of length 5 with the sum ahead of each element differing from the sum following that element by n or less.

Conjectures from _Colin Barker_, Aug 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007498 | OEIS-A221969

Number of -n..n arrays of length 6 with the sum ahead of each element differing from the sum following that element by n or less.

Conjectures from _Colin Barker_, Aug 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007499 | OEIS-A221970

Number of -n..n arrays of length 7 with the sum ahead of each element differing from the sum following that element by n or less.

Conjectures from _Colin Barker_, Aug 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007500 | OEIS-A222001

Number of n X 3 arrays with each row a permutation of 1..3 having at least as many downsteps as the preceding row, with rows in lexicographically nonincreasing order.

Conjectures from _Colin Barker_, Aug 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007501 | OEIS-A222008

Primes of the form 4^k + 1 for some k > 0.

Chebyshev showed that 3 is a primitive root of all primes of the form 2^(2*k) + 1 with k > 0. If the sequence is infinite, then Artin's conjecture ("every nonsquare integer n != -1 is a primitive root of infinitely many primes q") is true for n = 3. Conjecture: let p a prime number, a(n) is not congruent to p mod (p^2-3)/2. - _Vincenzo Librandi_, Jun 15 2014 This conjecture is false when p = a(n), but may be true for primes p != a(n). - _Jonathan Sondow_, Jun 15 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Feb 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007502 | OEIS-A222013

G.f. satisfies: A(x) = Sum_{n>=0} n! * x^n * A(x)^(n*(n+1)/2) / Product_{k=1..n} (1 + k*x*A(x)^k).

Conjecture: a(n) is odd iff n = 2^k - 1 for some k >= 0. - _Paul D. Hanna_, Dec 06 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007503 | OEIS-A222058

Harmonic-geometric numbers.

Conjecture: a(n) = Sum_{k>=0} k^n * (H(k) - log(2))/2^(k+1), where H(k) is the k-th harmonic number. - _Velin Yanev_, Jul 22 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007504 | OEIS-A222066

Decimal expansion of 1/sqrt(128).

Conjectured to be center density of densest packing of equal spheres in five dimensions (achieved for example by the D_5 lattice).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007505 | OEIS-A222067

Decimal expansion of 1/(8*sqrt(3)).

Conjectured to be center density of densest packing of equal spheres in six dimensions (achieved for example by the E_6 lattice).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007506 | OEIS-A222068

Decimal expansion of (1/16)*Pi^2.

Conjectured to be density of densest packing of equal spheres in four dimensions (achieved for example by the D_4 lattice).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007507 | OEIS-A222069

Decimal expansion of (1/30)*2^(1/2)*Pi^2.

Conjectured to be density of densest packing of equal spheres in five dimensions (achieved for example by the D_5 lattice).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007508 | OEIS-A222070

Decimal expansion of (1/144)*3^(1/2)*Pi^3.

Conjectured to be density of densest packing of equal spheres in six dimensions (achieved for example by the E_6 lattice).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007509 | OEIS-A222071

Decimal expansion of (1/105)*Pi^3.

Conjectured to be density of densest packing of equal spheres in 7 dimensions (achieved for example by the E_7 lattice).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007510 | OEIS-A222072

Decimal expansion of (1/384)*Pi^4.

Conjectured to be density of densest packing of equal spheres in 8 dimensions (achieved for example by the E_8 lattice). The above conjecture is true (cf. Viazovska, 2017). - _Felix Fröhlich_, Jan 08 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007511 | OEIS-A222098

Number of n X 2 0..5 arrays with entries increasing mod 6 by 0, 1, 2 or 3 rightwards and downwards, starting with upper left zero.

Conjectures from _Colin Barker_, Aug 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007512 | OEIS-A222114

Least integer m>1 such that 6*p_k*(p_k-1) (k=1,...,n) are pairwise incongruent modulo m, where p_k denotes the k-th prime.

Conjecture: For each n=3,4,..., a(n) is the first prime p>=p_n dividing none of those p_i+p_j-1 (1<=i<j<=n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007513 | OEIS-A222139

Number of n X 3 0..4 arrays with no entry increasing mod 5 by 4 rightwards or downwards, starting with upper left zero.

Conjectures from _Colin Barker_, Mar 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007514 | OEIS-A222219

Numbers n such that n and n + 18 are prime and there is a power of two in the interval (n,n+18).

It is a conjecture that this sequence is finite. A search around 2^n was done up to 2^1500.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brad Clardy_, Feb 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007515 | OEIS-A222227

Numbers n such that n and n + 16 are prime and there is a power of two in the interval (n,n+16).

It is a conjecture that this is a finite sequence. A search was conducted out to 2^1500.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brad Clardy_, Feb 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007516 | OEIS-A222267

The number of distinct lines defined by an n X n X n grid of points.

Conjecture: a(n) is approximately 0.3639537*n^6, with a relative error of about 10^-5 when n is near 200. - _Clive Tooth_, Mar 03 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clive Tooth_, Feb 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007517 | OEIS-A222276

Number of n X 3 0..5 arrays with no entry increasing mod 6 by 5 rightwards or downwards, starting with upper left zero.

Conjectures from _Colin Barker_, Mar 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007518 | OEIS-A222295

Conjectured number of Fibonacci numbers with exactly n bits set in their binary representation.

Conjectured number of Fibonacci numbers with exactly n bits set in their binary representation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Feb 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007519 | OEIS-A222313

A222311 sorted and duplicates removed (conjectured).

A222311 sorted and duplicates removed (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007520 | OEIS-A222424

Numbers n such that n and n + 14 are prime and there is a power of two in the interval (n,n+14).

It is a conjecture that this is a finite sequence. A search was conducted out to 2^1500.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brad Clardy_, Feb 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007521 | OEIS-A222459

Number of n X 3 0..7 arrays with no entry increasing mod 8 by 7 rightwards or downwards, starting with upper left zero.

Conjectures from _Colin Barker_, Mar 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007522 | OEIS-A222532

a(1)=2; for n >= 1, a(n+1) is the least prime p_m such that a(n)=p_m-p_{m-1}+...+(-1)^{m-k}p_k for some 0<k<m.

Conjecture: For any given prime p, if we define b(1)=p and let b(n+1) be the least prime p_m such that b(n)=p_m-p_{m-1}+...+(-1)^{m-k}p_k for some 0<k<m, then a(n)=b(n') for some positive integers n and n'. In other words, if we take all the primes as vertices of a simple graph T and let two vertices p and q>p adjacent if and only if q is the least prime p_m such that p=p_m-p_{m-1}+...+(-1)^{m-k}p_k for some 0<k<m, then the graph T is a tree!

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007523 | OEIS-A222566

a(1)=2; for n>0, a(n+1) is the least prime p>a(n) such that 2*(a(n)+1)-p is prime.

Conjecture: The graph G constructed above is connected and hence it is a tree!

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007524 | OEIS-A222579

Least prime p_m with p_m+1 practical such that n=p_m-p_{m-1}+...+(-1)^{m-k}p_k for some 0<k<m with p_k-1 practical.

Conjecture: a(n)<=3n for all n>0. Moreover, a(2n-1)/(2n-1) and a(2n)/(2n) have limits 1 and 2 respectively, as n tends to the infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007525 | OEIS-A222580

Number of ways to write n=p_m-p_{m-1}+...+(-1)^{m-k}p_k with k<m and p_m<=3n, and p_m+1 and p_k-1 both practical, where p_j denotes the j-th prime.

Conjecture: All the terms are positive.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007526 | OEIS-A222601

Conjectured number of Fibonacci numbers with exactly n 0-bits in their binary representation.

Conjectured number of Fibonacci numbers with exactly n 0-bits in their binary representation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007527 | OEIS-A222602

Irregular triangle of conjectured Fibonacci numbers with exactly n 0-bits in their binary representation.

Irregular triangle of conjectured Fibonacci numbers with exactly n 0-bits in their binary representation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007528 | OEIS-A222603

a(1)=1; for n>0, a(n+1) is the least practical number q>a(n) such that 2(a(n)+1)-q is practical.

Conjecture: The graph H constructed above is connected and hence it is a tree.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007529 | OEIS-A222757

Irregular table of conjectured indices of Fibonacci numbers with exactly n 0-bits in their binary representation.

Irregular table of conjectured indices of Fibonacci numbers with exactly n 0-bits in their binary representation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007530 | OEIS-A222758

Irregular table of conjectured indices of Fibonacci numbers with exactly n 1-bits in their binary representation.

Irregular table of conjectured indices of Fibonacci numbers with exactly n 1-bits in their binary representation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007531 | OEIS-A222759

Conjectured number of primes p for which binomial(n*p,p) (mod p^3) does not equal n.

Conjectured number of primes p for which binomial(n*p,p) (mod p^3) does not equal n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007532 | OEIS-A222760

Conjectured least prime p for which binomial(n*q,q) (mod q^3) = n for all primes q >= p.

Conjectured least prime p for which binomial(n*q,q) (mod q^3) = n for all primes q >= p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007533 | OEIS-A222763

Number of n X 2 0..1 arrays with exactly floor(nX2/2) elements unequal to at least one horizontal or antidiagonal neighbor, with new values introduced in row major 0..1 order.

Conjecture: Also the number of integer compositions of 2n + 1 with the same length as reverse-alternating sum. Here, the reverse-alternating sum of a sequence (y_1,...,y_k) is Sum_i (-1)^i y_i. For example, the a(4) = 20 compositions are:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007534 | OEIS-A222777

Number of n X 2 0..6 arrays with no element equal to another at a city block distance of exactly two, and new values 0..6 introduced in row major order.

Conjectures from _Colin Barker_, Aug 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007535 | OEIS-A222834

Number of n X 4 0..3 arrays with no element equal to another at a city block distance of exactly two, and new values 0..3 introduced in row major order.

Conjectures from _Colin Barker_, Aug 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007536 | OEIS-A222835

Number of n X 5 0..3 arrays with no element equal to another at a city block distance of exactly two, and new values 0..3 introduced in row major order.

Conjectures from _Colin Barker_, Aug 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007537 | OEIS-A222869

Number of n X 3 0..5 arrays with no element equal to another at a city block distance of exactly two, and new values 0..5 introduced in row major order.

Conjectures from _Colin Barker_, Aug 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007538 | OEIS-A222892

Number of n X 3 0..7 arrays with no element equal to another at a city block distance of exactly two, and new values 0..7 introduced in row major order.

Conjectures from _Colin Barker_, Aug 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007539 | OEIS-A222939

Number of n X 1 0..4 arrays with no element equal to another at a city block distance of exactly two, and new values 0..4 introduced in row major order.

Conjectures from _Colin Barker_, Aug 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007540 | OEIS-A222940

Number of n X 2 0..4 arrays with no element equal to another at a city block distance of exactly two, and new values 0..4 introduced in row major order.

Conjectures from _Colin Barker_, Aug 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007541 | OEIS-A222945

Number of distinct sums i+j+k with |i|, |j|, |k|, |i*j*k| <= n.

Mathar's conjecture is true (see Fried link). - _Sela Fried_, Mar 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007542 | OEIS-A222955

Number of n X 1 0..1 arrays with every row and column least squares fitting to a zero slope straight line, with a single point array taken as having zero slope.

Conjecture: A binary word is counted iff it has the same sum of positions of 1's as its reverse, or, equivalently, the same sum of partial sums as its reverse. - _Gus Wiseman_, Jan 07 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007543 | OEIS-A222993

Number of n X 2 0..2 arrays with successive rows and columns fitting to straight lines with nondecreasing slope, with a single point array taken as having zero slope.

Conjectures from _Colin Barker_, Feb 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007544 | OEIS-A223083

Trajectory of 64 under the map n-> A006369(n).

It is conjectured that this trajectory does not close on itself.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007545 | OEIS-A223084

Trajectory of 80 under the map n-> A006369(n).

It is conjectured that this trajectory does not close on itself.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007546 | OEIS-A223085

Trajectory of 82 under the map n-> A006369(n).

It is conjectured that this trajectory does not close on itself.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007547 | OEIS-A223086

Trajectory of 64 under the map n-> A006368(n).

It is conjectured that this trajectory does not close on itself.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007548 | OEIS-A223087

Trajectory of 80 under the map n-> A006368(n).

It is conjectured that this trajectory does not close on itself.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007549 | OEIS-A223088

Trajectory of 82 under the map n-> A006368(n).

It is conjectured that this trajectory does not close on itself.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007550 | OEIS-A223174

Smallest prime p such that 2n+1 = p + 8*q for some odd prime q, or 0 if no such prime exists.

Conjecture: except m = 25 and 49, all odd numbers > 17 are of the form m = p + 8*q where p and q are prime numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, May 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007551 | OEIS-A223181

Rolling icosahedron footprints: number of n X 3 0..11 arrays starting with 0 where 0..11 label vertices of an icosahedron and every array movement to a horizontal, vertical or antidiagonal neighbor moves along an icosahedral edge.

Conjectures from _Colin Barker_, Aug 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007552 | OEIS-A223197

Rolling cube footprints: number of n X 3 0..5 arrays starting with 0 where 0..5 label faces of a cube and every array movement to a horizontal or vertical neighbor moves across a corresponding cube edge.

Conjectures from _Colin Barker_, Mar 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007553 | OEIS-A223204

Rolling icosahedron face footprints: number of n X 3 0..19 arrays starting with 0 where 0..19 label faces of an icosahedron and every array movement to a horizontal or vertical neighbor moves across an icosahedral edge.

Conjectures from _Colin Barker_, Aug 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007554 | OEIS-A223211

3 X 3 X 3 triangular graph coloring a rectangular array: number of n X 1 0..5 arrays where 0..5 label nodes of a graph with edges 0,1 0,2 1,2 1,3 1,4 2,4 3,4 2,5 4,5 and every array movement to a horizontal or vertical neighbor moves along an edge of this graph.

Conjectures from _Colin Barker_, Aug 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007555 | OEIS-A223228

Rolling icosahedron footprints: number of n X 3 0..11 arrays starting with 0 where 0..11 label vertices of an icosahedron and every array movement to a horizontal, diagonal or antidiagonal neighbor moves along an icosahedral edge.

Conjectures from _Colin Barker_, Aug 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007556 | OEIS-A223234

Rolling icosahedron footprints: number of 2 X n 0..11 arrays starting with 0 where 0..11 label vertices of an icosahedron and every array movement to a horizontal, diagonal or antidiagonal neighbor moves along an icosahedral edge.

Conjectures from _Colin Barker_, Aug 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007557 | OEIS-A223240

3-loop graph coloring a rectangular array: number of n X 1 0..6 arrays where 0..6 label nodes of a graph with edges 0,1 1,2 2,0 0,3 3,4 4,0 0,5 5,6 6,0 and every array movement to a horizontal or vertical neighbor moves along an edge of this graph.

Conjectures from _Colin Barker_, Aug 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007558 | OEIS-A223249

Two-loop graph coloring a rectangular array: number of n X 2 0..4 arrays where 0..4 label nodes of a graph with edges 0,1 1,2 2,0 0,3 3,4 4,0 and every array movement to a horizontal or vertical neighbor moves along an edge of this graph.

Conjectures from _Colin Barker_, Mar 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007559 | OEIS-A223258

Rolling icosahedron footprints: number of n X 3 0..11 arrays starting with 0 where 0..11 label vertices of an icosahedron and every array movement to a horizontal or vertical neighbor moves along an icosahedral edge.

Conjectures from _Colin Barker_, Aug 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007560 | OEIS-A223270

Rolling cube footprints: number of 2 X n 0..5 arrays starting with 0 where 0..5 label faces of a cube and every array movement to a horizontal, diagonal or antidiagonal neighbor moves across a corresponding cube edge.

Conjectures from _Colin Barker_, Aug 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007561 | OEIS-A223277

Rolling icosahedron face footprints: number of n X 3 0..19 arrays starting with 0 where 0..19 label faces of an icosahedron and every array movement to a horizontal, diagonal or antidiagonal neighbor moves across an icosahedral edge.

Conjectures from _Colin Barker_, Aug 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007562 | OEIS-A223278

Rolling icosahedron face footprints: number of n X 4 0..19 arrays starting with 0 where 0..19 label faces of an icosahedron and every array movement to a horizontal, diagonal or antidiagonal neighbor moves across an icosahedral edge.

Conjectures from _Colin Barker_, Aug 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007563 | OEIS-A223291

4-loop graph coloring a rectangular array: number of n X 2 0..8 arrays where 0..8 label nodes of a graph with edges 0,1 1,2 2,0 0,3 3,4 4,0 0,5 5,6 6,0 0,7 7,8 8,0 and every array movement to a horizontal or vertical neighbor moves along an edge of this graph.

Conjectures from _Colin Barker_, Aug 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007564 | OEIS-A223299

4 X 4 X 4 triangular graph coloring a rectangular array: number of n X 2 0..9 arrays where 0..9 label nodes of a graph with edges 0,1 0,2 1,2 1,3 1,4 2,4 3,4 2,5 4,5 3,6 3,7 4,7 6,7 4,8 5,8 7,8 5,9 8,9 and every array movement to a horizontal or vertical neighbor moves along an edge of this graph.

Conjectures from _Colin Barker_, Aug 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007565 | OEIS-A223318

Rolling icosahedron footprints: number of n X 5 0..11 arrays starting with 0 where 0..11 label vertices of an icosahedron and every array movement to a horizontal or antidiagonal neighbor moves along an icosahedral edge.

Conjectures from _Colin Barker_, Aug 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007566 | OEIS-A223322

Rolling icosahedron footprints: number of 2 X n 0..11 arrays starting with 0 where 0..11 label vertices of an icosahedron and every array movement to a horizontal or antidiagonal neighbor moves along an icosahedral edge.

Conjectures from _Colin Barker_, Aug 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007567 | OEIS-A223332

Rolling cube footprints: number of 3 X n 0..7 arrays starting with 0 where 0..7 label vertices of a cube and every array movement to a horizontal or antidiagonal neighbor moves along a corresponding cube edge.

Conjectures from _Colin Barker_, Aug 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007568 | OEIS-A223363

6 X 6 X 6 triangular graph coloring a rectangular array: number of n X 1 0..20 arrays where 0..20 label nodes of the fully triangulated graph and every array movement to a horizontal or vertical neighbor moves along an edge of this graph.

Conjectures from _Colin Barker_, Aug 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007569 | OEIS-A223381

Number of n X 2 0..2 arrays with all horizontally or vertically connected equal values in a straight line, and new values 0..2 introduced in row major order.

Conjectures from _Colin Barker_, Aug 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007570 | OEIS-A223395

4 X 4 square grid graph coloring a rectangular array: number of n X 1 0..15 arrays where 0..15 label nodes of the square grid graph and every array movement to a horizontal or vertical neighbor moves along an edge of this graph.

Conjectures from _Colin Barker_, Aug 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007571 | OEIS-A223443

4-level binary fanout graph coloring a rectangular array: number of n X 2 0..14 arrays where 0..14 label nodes of a graph with edges 0,1 1,3 3,5 3,6 1,4 4,7 4,8 0,2 2,9 9,11 9,12 2,10 10,13 10,14 and every array movement to a horizontal or vertical neighbor moves along an edge of this graph.

Conjectures from _Colin Barker_, Aug 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007572 | OEIS-A223499

Petersen graph (3,1) coloring a rectangular array: number of n X 3 0..5 arrays where 0..5 label nodes of a graph with edges 0,1 0,3 3,5 3,4 1,2 1,4 4,5 2,0 2,5 and every array movement to a horizontal, diagonal or antidiagonal neighbor moves along an edge of this graph, with the array starting at 0.

Conjectures from _Colin Barker_, Aug 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007573 | OEIS-A223541

Array T(m,n) = nim-product(2^m,2^n) (m>=0, n>=0) read by antidiagonals.

Piesk's conjecture above is true: the antidiagonal m + n = N contains exactly A002487(N+1) distinct values, and each of them occurs 2^k times, where k is the number of 1's in the binary expansion of m XOR n; see the Blanco Gómez link. - _Roberto Blanco Gómez_, Oct 05 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tilman Piesk_, Mar 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007574 | OEIS-A223552

Petersen graph (3,1) coloring a rectangular array: number of n X 4 0..5 arrays where 0..5 label nodes of a graph with edges 0,1 0,3 3,5 3,4 1,2 1,4 4,5 2,0 2,5 and every array movement to a horizontal or antidiagonal neighbor moves along an edge of this graph, with the array starting at 0.

Conjectures from _Colin Barker_, Aug 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007575 | OEIS-A223577

Positive integers k for which there is exactly one negative integer m such that -k = floor(cot(Pi/(2*m))).

Conjecture: a(n) = (A223578(n) + n - 1)/2. The conjecture is true (see Fried link). - _Sela Fried_, Oct 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Mar 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007576 | OEIS-A223578

Positive integers k for which f(-k-1) < f(-k) < f(-k+1), where f(m) = floor(cot(Pi/(2m))).

Conjecture: a(n) = 2*A223577(n) - n + 1. The conjecture is true (see Fried link). - _Sela Fried_, Oct 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Mar 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007577 | OEIS-A223615

Number of n X 3 0..1 arrays with rows and columns unimodal.

Conjectures from _Colin Barker_, Aug 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007578 | OEIS-A223616

Number of n X 4 0..1 arrays with rows and columns unimodal.

Conjectures from _Colin Barker_, Aug 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007579 | OEIS-A223632

Number of n X 3 0..1 arrays with rows, antidiagonals and columns unimodal.

Conjectures from _Colin Barker_, Aug 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007580 | OEIS-A223633

Number of n X 4 0..1 arrays with rows, antidiagonals and columns unimodal.

Conjectures from _Colin Barker_, Aug 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007581 | OEIS-A223639

Number of n X 3 0..1 arrays with rows, columns, diagonals and antidiagonals unimodal.

Conjectures from _Colin Barker_, Aug 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007582 | OEIS-A223640

Number of n X 4 0..1 arrays with rows, columns, diagonals and antidiagonals unimodal.

Conjectures from _Colin Barker_, Aug 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007583 | OEIS-A223670

Number of 3 X n 0..1 arrays with rows, diagonals and antidiagonals unimodal.

Conjectures from _Colin Barker_, Mar 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007584 | OEIS-A223681

Number of 3 X n 0..1 arrays with rows and antidiagonals unimodal.

Conjectures from _Colin Barker_, Aug 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007585 | OEIS-A223682

Number of 4 X n 0..1 arrays with rows and antidiagonals unimodal.

Conjectures from _Colin Barker_, Aug 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007586 | OEIS-A223687

Petersen graph (8,2) coloring a rectangular array: number of n X 3 0..15 arrays where 0..15 label nodes of a graph with edges 0,1 0,8 8,14 8,10 1,2 1,9 9,15 9,11 2,3 2,10 10,12 3,4 3,11 11,13 4,5 4,12 12,14 5,6 5,13 13,15 6,7 6,14 7,0 7,15 and every array movement to a horizontal or antidiagonal neighbor moves along an edge of this graph.

Conjectures from _Colin Barker_, Aug 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007587 | OEIS-A223703

Conjectured irregular triangle (with some rows blank) of numbers k such that prime(n) is the largest prime factor of k^3 - 1.

Conjectured irregular triangle (with some rows blank) of numbers k such that prime(n) is the largest prime factor of k^3 - 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007588 | OEIS-A223706

Conjectured least number k such that prime(n) is the largest divisor of k^3 - 1, or 0 if there is no such k.

Conjectured least number k such that prime(n) is the largest divisor of k^3 - 1, or 0 if there is no such k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Apr 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007589 | OEIS-A223711

Number of n X 2 0..1 arrays with row sums and column sums unimodal.

Conjectures from _Colin Barker_, Mar 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007590 | OEIS-A223719

Number of n X 2 0..2 arrays with rows, antidiagonals and columns unimodal.

Conjectures from _Colin Barker_, Feb 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007591 | OEIS-A223730

Multiplicities for representations of positive numbers n as primitive sums of three nonzero squares.

For representations of n as a sum of three nonzero squares see the Grosswald reference, Theorem 7, p. 79. There also the above mentioned set T appears and for the Conjecture it is assumed that the extra eleventh member of T is absent.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Apr 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007592 | OEIS-A223731

All positive numbers that are primitive sums of three nonzero squares.

Conjectured g.f.: (x^77 +2*x^76 -2*x^75 +x^74 -x^73 -x^72 +2*x^50 -x^49 +2*x^47 -2*x^46 -x^45 +x^34 +2*x^33 -2*x^32 +x^31 -x^30 -x^29 +2*x^22 -x^21 +2*x^19 -2*x^18 -x^17 +3*x^15 -2*x^14 +x^13 -x^12 -x^10 +2*x^9 +2*x^7 +2*x^6 -3*x^4 -2*x^3 -3*x^2 -3*x -3)*x / (-x^6 +x^5 +x -1). - _Alois P. Heinz_, Apr 06 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Apr 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007593 | OEIS-A223732

Positive numbers that are the sum of three nonzero squares with no common factor > 1 in exactly one way.

Conjecture: 793 = 6^2 + 9^2 + 26^2 is the largest element of this sequence. - _Alois P. Heinz_, Apr 06 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Apr 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007594 | OEIS-A223733

Positive numbers that are the sum of three nonzero squares with no common factor > 1 in exactly two ways.

Conjecture: a(147) = 1885 = 16^2 + 27^2 + 30^2 = 12^2 + 29^2 + 30^2 is the largest element of this sequence. - _Alois P. Heinz_, Apr 06 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Apr 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007595 | OEIS-A223734

Positive numbers that are the sum of three nonzero squares with no common factor > 1 in exactly three ways.

Conjecture: a(185) = 4075 = 31^2 + 33^2 + 45^2 = 23^2 + 39^2 + 45^2 = 5^2 + 9^2 + 63^2 is the largest element of this sequence. - _Alois P. Heinz_, Apr 06 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Apr 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007596 | OEIS-A223735

Positive numbers that are not representable as a primitive sum of three nonzero squares.

Conjectured g.f.: (2*x^61 -x^60 +2*x^59 -x^58 -2*x^57 +x^43 +3*x^42 -3*x^41 +x^40 -2*x^39 +2*x^32 -x^31 +2*x^30 -x^29 -2*x^28 +x^23 +3*x^22 -3*x^21 +x^20 -2*x^19 +x^18 +2*x^16 -3*x^14 +x^12 +3*x^11 -x^10 +x^6 -x^5 +x^4 +2*x^2 +x +1)*x / (x^4 -x^3 -x +1). - _Alois P. Heinz_, Apr 06 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Apr 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007597 | OEIS-A223756

Number of n X 2 0..3 arrays with rows, antidiagonals and columns unimodal.

Conjectures from _Colin Barker_, Feb 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007598 | OEIS-A223764

Number of n X 2 0..1 arrays with rows, columns and antidiagonals unimodal and diagonals nondecreasing.

Conjectures from _Colin Barker_, Feb 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007599 | OEIS-A223765

Number of n X 3 0..1 arrays with rows, columns and antidiagonals unimodal and diagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007600 | OEIS-A223766

Number of n X 4 0..1 arrays with rows, columns and antidiagonals unimodal and diagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007601 | OEIS-A223772

Number of n X 3 0..1 arrays with rows and columns unimodal and antidiagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007602 | OEIS-A223773

Number of n X 4 0..1 arrays with rows and columns unimodal and antidiagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007603 | OEIS-A223774

Number of n X 5 0..1 arrays with rows and columns unimodal and antidiagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007604 | OEIS-A223833

Number of n X 3 0..1 arrays with rows, diagonals and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007605 | OEIS-A223834

Number of n X 4 0..1 arrays with rows, diagonals and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007606 | OEIS-A223835

Number of n X 5 0..1 arrays with rows, diagonals and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007607 | OEIS-A223836

Number of n X 6 0..1 arrays with rows, diagonals and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007608 | OEIS-A223837

Number of n X 7 0..1 arrays with rows, diagonals and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007609 | OEIS-A223839

Number of 3 X n 0..1 arrays with rows, diagonals and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007610 | OEIS-A223840

Number of 4 X n 0..1 arrays with rows, diagonals and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007611 | OEIS-A223859

Number of n X 3 0..3 arrays with rows and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007612 | OEIS-A223881

Denominators in the expression m!/(prime(m-1)+1) for m > 1 such that this expression is not an integer.

The scatter plot reveals four distinct, well-separated, monotonically increasing curves. It became possible to extract the integers (all conjectured to be primes) corresponding to each of the four subsets.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander R. Povolotsky_, May 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007613 | OEIS-A223886

Numbers (binomial(j*k*prime(n), j*prime(n)) - binomial(k*j, j))/(k*prime(n)^3) for k=4, j=3 and n>=2.

This sequence (together with already present in the OEIS A034602 and A217772) is based on Gary Detlefs' conjecture, which he disclosed to me in a private communication on 3/29/13 and recently he gave me permission to make it public. Specifically he wrote to me the following: "I have a conjecture which is broader than the one I submitted, having to do with binomial(k*n,n) mod n^3. It appears that binomial(j*k*n,j*n) mod n^3 will be binomial(k*j,j) for n sufficiently large." In effect above conjecture further extends Wolstenholme's and Ljunggren's ideas and could also be expressed as follows: starting with some specific (for any given unchanged values of integers k>0 and j>0) sufficiently large value of n=N and further on for n>N it is true that (binomial(j*k*prime(n), j*prime(n)) - binomial(k*j, j))/k/(prime(n))^3 = m(j, k, n), where m(j, k, n) are integer values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander R. Povolotsky_, Mar 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007614 | OEIS-A223911

Number of tiered orders on n nodes (corrected version of A006860).

Conjecture: a(n) ~ c * 2^(n^2/4 + 3*n/2) / sqrt(n), where c = EllipticTheta[3, 0, 1/2^(1/4)] / (sqrt(Pi) * 2^(1/4)) = 2.020039... (based on the numerical analysis of 600 terms). - _Vaclav Kotesovec_, Apr 10 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Mar 29 2013, using information provided by _Joel B. Lewis_, _M. F. Hasler_ and _Michel Marcus_ in A006860 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007615 | OEIS-A223913

Number of n X 3 0..2 arrays with rows and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007616 | OEIS-A223914

Number of n X 4 0..2 arrays with rows and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007617 | OEIS-A223919

Number of 2 X n 0..2 arrays with rows and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Feb 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007618 | OEIS-A223927

Number of n X 2 0..2 arrays with rows, columns and antidiagonals unimodal and diagonals nondecreasing.

Conjectures from _Colin Barker_, Feb 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007619 | OEIS-A223934

Least prime p such that x^n-x-1 is irreducible modulo p.

Conjecture: a(n) < n*(n+3)/2 for all n>1. The conjecture was motivated by E. S. Selmer's result that for any n>1 the polynomial x^n-x-1 is irreducible over the field of rational numbers. We also conjecture that for every n=2,3,... there is a positive integer z not exceeding the (2n-2)-th prime such that z^n-z-1 is prime, and the Galois group of x^n-x-1 over the field of rationals is isomorphic to the symmetric group S_n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007620 | OEIS-A223942

Least prime q such that (x^{p_n}-1)/(x-1) is irreducible modulo q, where p_n is the n-th prime.

Conjecture: a(n) <= sqrt(7*p_n) for all n>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007621 | OEIS-A223950

Number of 3 X n 0..1 arrays with diagonals and antidiagonals unimodal and rows nondecreasing.

Conjectures from _Colin Barker_, Aug 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007622 | OEIS-A223951

Number of 4 X n 0..1 arrays with diagonals and antidiagonals unimodal and rows nondecreasing.

Conjectures from _Colin Barker_, Aug 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007623 | OEIS-A223952

Number of 5 X n 0..1 arrays with diagonals and antidiagonals unimodal and rows nondecreasing.

Conjectures from _Colin Barker_, Aug 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007624 | OEIS-A223953

Number of 6 X n 0..1 arrays with diagonals and antidiagonals unimodal and rows nondecreasing.

Conjectures from _Colin Barker_, Aug 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007625 | OEIS-A223954

Number of 7 X n 0..1 arrays with diagonals and antidiagonals unimodal and rows nondecreasing.

Conjectures from _Colin Barker_, Aug 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007626 | OEIS-A223962

Number of 2 X n 0..3 arrays with antidiagonals unimodal and rows and diagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007627 | OEIS-A223963

Number of 3 X n 0..3 arrays with antidiagonals unimodal and rows and diagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007628 | OEIS-A223982

Number of n X 3 0..3 arrays with rows unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007629 | OEIS-A224000

Number of 2 X n 0..2 arrays with antidiagonals unimodal and rows and diagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007630 | OEIS-A224001

Number of 3 X n 0..2 arrays with antidiagonals unimodal and rows and diagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007631 | OEIS-A224002

Number of 4 X n 0..2 arrays with antidiagonals unimodal and rows and diagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007632 | OEIS-A224013

Number of 3 X n 0..2 arrays with rows nondecreasing and antidiagonals unimodal.

Conjectures from _Colin Barker_, Aug 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007633 | OEIS-A224014

Number of 4 X n 0..2 arrays with rows nondecreasing and antidiagonals unimodal.

Conjectures from _Colin Barker_, Aug 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007634 | OEIS-A224025

Number of 3 X n 0..3 arrays with rows nondecreasing and antidiagonals unimodal.

Conjectures from _Colin Barker_, Aug 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007635 | OEIS-A224030

a(n) = |{0<k<n: 2*n+k and 2*n^3+k^3 are both prime}|.

Conjecture: a(n)>0 for all n>4. We also conjecture that for any integer n>1 there is an integer 0<k<n such that n^2+k^2 is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007636 | OEIS-A224039

Number of 3 X n 0..1 arrays with antidiagonals unimodal and rows and diagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007637 | OEIS-A224040

Number of 4 X n 0..1 arrays with antidiagonals unimodal and rows and diagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007638 | OEIS-A224041

Number of 5 X n 0..1 arrays with antidiagonals unimodal and rows and diagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007639 | OEIS-A224042

Number of 6 X n 0..1 arrays with antidiagonals unimodal and rows and diagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007640 | OEIS-A224043

Number of 7 X n 0..1 arrays with antidiagonals unimodal and rows and diagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007641 | OEIS-A224067

Number of cycles the reduced complex Collatz function using 1 + (2n-1)i takes to hit 1 + i, or 0 if this never happens.

This is conjectured to be infinite and the same as in A224165.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Apr 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007642 | OEIS-A224134

Number of 3 X n 0..1 arrays with rows nondecreasing and antidiagonals unimodal.

Conjectures from _Colin Barker_, Aug 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007643 | OEIS-A224135

Number of 4 X n 0..1 arrays with rows nondecreasing and antidiagonals unimodal.

Conjectures from _Colin Barker_, Aug 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007644 | OEIS-A224136

Number of 5 X n 0..1 arrays with rows nondecreasing and antidiagonals unimodal.

Conjectures from _Colin Barker_, Aug 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007645 | OEIS-A224137

Number of 6 X n 0..1 arrays with rows nondecreasing and antidiagonals unimodal.

Conjectures from _Colin Barker_, Aug 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007646 | OEIS-A224138

Number of 7 X n 0..1 arrays with rows nondecreasing and antidiagonals unimodal.

Conjectures from _Colin Barker_, Aug 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007647 | OEIS-A224141

Number of n X 3 0..1 arrays with rows and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007648 | OEIS-A224142

Number of n X 4 0..1 arrays with rows and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007649 | OEIS-A224143

Number of n X 5 0..1 arrays with rows and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007650 | OEIS-A224144

Number of n X 6 0..1 arrays with rows and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007651 | OEIS-A224145

Number of n X 7 0..1 arrays with rows and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007652 | OEIS-A224147

Number of 3 X n 0..1 arrays with rows and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007653 | OEIS-A224148

Number of 4 X n 0..1 arrays with rows and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007654 | OEIS-A224149

Number of 5 X n 0..1 arrays with rows and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007655 | OEIS-A224150

Number of 6 X n 0..1 arrays with rows and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007656 | OEIS-A224159

Number of 3 X n 0..1 arrays with diagonals and rows unimodal and antidiagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007657 | OEIS-A224160

Number of 4 X n 0..1 arrays with diagonals and rows unimodal and antidiagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007658 | OEIS-A224165

Number of cycles the reduced complex Collatz function using 3 + (2n-1)i takes to hit 1 + i, or 0 if this never happens.

This is conjectured to be infinite and the same as in A224067.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Apr 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007659 | OEIS-A224168

Number of n X 3 0..3 arrays with rows, diagonals and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007660 | OEIS-A224183

Number of halving and tripling steps to reach the last number of the cycle in the Collatz (3x+1) problem for the negative integers (the initial term is not counted).

The Collatz problem with negative numbers is as follows: start with any number n = -1, -2, -3, ... If n is even, divide it by 2, otherwise multiply it by 3 and add 1. Do we always

reach a last number of each cycle? It is conjectured that the answer is yes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007661 | OEIS-A224185

Number of n X 3 0..2 arrays with rows unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007662 | OEIS-A224186

Number of n X 4 0..2 arrays with rows unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007663 | OEIS-A224197

Least b > p_n^2 such that [p_1^2,p_2^2,...,p_n^2] in base b is prime, where p_j denotes the j-th prime.

Conjecture: (i) For any positive integer k and distinct positive integers a_1< a_2 < ... < a_n with a_n prime, there are infinitely many integers b > a_n^k such that [a_1^k,a_2^k,...,a_n^k] in base b is prime. This is the k-th power version of the author's conjecture related to A217788. Note that s(m,n) defined there is identical with s_1(m,n). It seems that s_2(m,n) < p_{n+1}*p_{m+n+1}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007664 | OEIS-A224210

Least prime p such that sum_{k=0}^n (k+1)^2*x^{n-k} is irreducible modulo p.

Conjecture: a(n) does not exceed the (4n-3)-th prime for each n>0. Moreover, for any integers m>1 and n>0 the polynomial sum_{k=0}^n (k+1)^m*x^{n-k} is irreducible modulo some prime, and its Galois group over the rationals is isomorphic to the symmetric group S_n. Also, for m,n=2,3,... there are infinitely many integers b > n^m such that [n^m,...,2^m,1^m] in base b is prime. We have a similar conjecture with the above (k+1)^m replaced by (2k+1)^m.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007665 | OEIS-A224217

Numbers b such that b^2 + 4*b + 9 is prime.

Inspired by Sun's conjecture in A224197: Also, bases b such that the number (with digits) [1;4;9] in base b is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007666 | OEIS-A224241

Numbers k such that k^2 XOR (k+1)^2 is a square, and k^2 XOR (k+2)^2 is also a square, where XOR is the bitwise logical exclusive-or operator.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Apr 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007667 | OEIS-A224242

Numbers k such that k^2 XOR (k+1)^2 is a square, and k^2 XOR (k-1)^2 is a square, where XOR is the bitwise logical XOR operator.

A059153 is a subsequence. Terms that are not in A059153: 0, 44, 263160, 278828, 340028, 25239472, 32490836. Conjecture: the subsequence of non-A059153 terms is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Apr 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007668 | OEIS-A224254

Full cycle lengths in the Collatz (3x+1) problem when the negative integers are used.

There are other cycles of lengths 2, 5 and 18 if negative integers are used. In Z, it is conjectured that the five values of cycle are 1, 2, 3, 5 and 18 (see A121510).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007669 | OEIS-A224257

Number of n X 3 0..2 arrays with rows, diagonals and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007670 | OEIS-A224258

Number of n X 4 0..2 arrays with rows, diagonals and antidiagonals unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Aug 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007671 | OEIS-A224274

a(n) = binomial(4*n,n)/4.

Conjecture: a(n) == 1 (mod n^3) iff n is an odd prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary Detlefs_, Apr 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007672 | OEIS-A224287

Number of multiplicity free permutations in S_n, i.e., permutations whose Stanley symmetric function is multiplicity free.

This family is conjectured to be characterized by a finite set of patterns up to S_11.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sara Billey_, Apr 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007673 | OEIS-A224295

Number of permutations of length n avoiding 12345 and 12354.

Conjectured to be the number of permutations of length n avoiding the partially ordered pattern (POP) {2>1>5>3, 5>4} of length 5. That is, conjectured to be the number of length n permutations having no subsequences of length 5 in which the elements 3 and 4 are the smallest, and the element in position 2 is larger than that in position 1, which in turn is larger than the element in position 5.- _Sergey Kitaev_, Dec 13 2020 Restatement of the comment by Kitaev: Conjectured to be the number of permutations of length n avoiding patterns 45123 and 45213. - _Alexander Burstein_, Feb 05 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Nakamura_, Apr 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007674 | OEIS-A224299

Collatz problem with rational negative numbers: number of steps to reach the end of the cycle starting with 1/(2n+1) where n is negative (the initial term is not counted).

This variation of the "3x+1" problem with a class of rational negative numbers is as follows: start with any number 1/(2n+1), n= -1, -2, -3, ... If the numerator is even, divide it by 2, otherwise multiply it by 3 and add 1. Do we always reach the end of a cycle with a rational number? It is conjectured that the answer is yes. This sequence is an extension of A210468 with n negative.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007675 | OEIS-A224300

Collatz problem with rational negative numbers: number of steps to reach the end of the cycle starting with 1/(2n+1) where n is negative (the initial value is counted).

This variation of the "3x+1" problem with a class of rational negative numbers is as follows: start with any number 1/(2n+1), n= -1, -2, -3, ... If the numerator is even, divide it by 2, otherwise multiply it by 3 and add 1. Do we always reach the end of a cycle with a rational number? It is conjectured that the answer is yes. This sequence is an extension of A210471 with n negative.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007676 | OEIS-A224316

Number of permutations w in S_n whose Kazhdan-Lusztig polynomial P_{id,w}(1) <= 3.

This family of permutations is closed under taking patterns. It is conjectured to be characterized by a finite set of patterns by Billey and Weed.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sara Billey_, Apr 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007677 | OEIS-A224342

Apparently solves the identity: find sequence B that represents the numbers of ordered compositions of n using the terms of A, and vice versa.

Alternatively as a conjecture, A and B solve the unique identity as described in the heading as to ordered compositions with A = A224341 and B = A224342. The INVERT transform of the characteristic function of A = B, and the INVERT transform of the characteristic function of B = A.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary W. Adamson_, Apr 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007678 | OEIS-A224354

Number of 3 X n 0..2 arrays with diagonals and antidiagonals unimodal and rows nondecreasing.

Conjectures from _Colin Barker_, Aug 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007679 | OEIS-A224355

Number of 4 X n 0..2 arrays with diagonals and antidiagonals unimodal and rows nondecreasing.

Conjectures from _Colin Barker_, Aug 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007680 | OEIS-A224392

Number of 3 X n 0..3 arrays with diagonals and antidiagonals unimodal and rows nondecreasing.

Conjectures from _Colin Barker_, Aug 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007681 | OEIS-A224400

Composite numbers that become prime when their digits are put in nondecreasing order.

Conjecture: a(n) ~ 7.75*n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian N. K. Anderson_ and _Kevin L. Schwartz_, Apr 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007682 | OEIS-A224410

Number of 3 X n 0..1 arrays with rows unimodal and antidiagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007683 | OEIS-A224411

Number of 4 X n 0..1 arrays with rows unimodal and antidiagonals nondecreasing.

Conjectures from _Colin Barker_, Aug 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007684 | OEIS-A224416

Least prime p such that the polynomial sum_{k=0}^n C_k*x^{n-k} is irreducible modulo p, where C_k denotes the Catalan number binomial(2k,k)/(k+1).

Conjecture: (i) a(n) does not exceed n^2+n+5 for each n>0, and the Galois group of sum_{k=0}^n C_k*x^{n-k} over the rationals is isomorphic to the symmetric group S_n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007685 | OEIS-A224417

Least prime p such that sum_{k=0}^n B_k*x^{n-k} is irreducible modulo p, where B_k refers to the Bell number A000110(k).

Conjecture: a(n) < 4n^2-1 for all n>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007686 | OEIS-A224418

Least prime q such that sum_{k=0}^n p(k)*x^{n-k} is irreducible modulo q, where p(k) refers to the partition number A000041(k).

Conjecture: a(n) < n^2 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007687 | OEIS-A224480

Least prime q such that x^n + sum_{k=1}^n p_k*x^{n-k} is irreducible modulo q, where p_k denotes the k-th prime.

Conjecture: a(n) <= (n+4)*(n+5)+1 for all n>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007688 | OEIS-A224487

Least integer b > F(n) such that sum_{k=1}^n F(k)*b^{k-1} is prime, where F = A000045.

Conjecture: For any n > 1 we have a(n) < F(n+4); moreover, there are infinitely many integers b > F(n) such that sum_{k=1}^n F(k)*b^{k-1} is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007689 | OEIS-A224492

Smallest k such that k*2*p(n)^2-1=q is prime, k*2*q^2-1=r, k*2*r^2-1=s, k*2*r^2-1=t, r, s, and t are also prime.

conjecture: a(n) exist for all n

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Apr 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007690 | OEIS-A224511

Roots of squares generated in A221643. That is, S = i^2 XOR (i+1)^2; increment i; if S is a square then square root of S is appended to a(n). Initially i=0. XOR is the binary logical exclusive-or operator.

Conjecture: this is a permutation of odd numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Apr 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007691 | OEIS-A224523

Let {f_n(k)} be the n-th sequence of Fibonacci-like numbers defined by recursion f_n(0) = 0, f_n(1) = 1 and, for k>=2, f_n(k) = f_n(k-1) + f_n(k-2) divided by maximal possible powers of primes >= prime(n). a(n) is length of the smallest period of {f_n(k)}, and a(n)=0, if {f_n(k)} is not eventually periodic.

Conjecture. For n>=1, a(n)>0. If Conjecture is true, then we have an answer in affirmative on question in A078414.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007692 | OEIS-A224529

Sequence f_n from a paper by Robert Osburn and Brundaban Sahu.

Conjecture 1.1 of Osburn and Sahu is if p is a prime and JacobiSymbol(p, 23) = 1 and n>0 then a(n * p) == a(n) (mod p). - _Michael Somos_, Sep 21 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Apr 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007693 | OEIS-A224613

a(n) = sigma(6*n).

Conjectures: sigma(6n) > sigma(6n - 1) and sigma(6n) > sigma(6n + 1). Conjectures are false. Try prime 73961483429 for n. One finds sigma(6*73961483429) < sigma(6*73961483429+1). The number n = 105851369791 provides a counterexample for the other case. - _T. D. Noe_, Apr 22 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Apr 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007694 | OEIS-A224666

Number of 4 X 4 0..n matrices with each 2 X 2 subblock idempotent.

Conjectures from _Colin Barker_, Sep 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007695 | OEIS-A224667

Number of 5 X 5 0..n matrices with each 2 X 2 subblock idempotent.

Conjectures from _Colin Barker_, Sep 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007696 | OEIS-A224668

Number of 6 X 6 0..n matrices with each 2 X 2 subblock idempotent.

Conjectures from _Colin Barker_, Sep 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007697 | OEIS-A224669

Number of (n+1) X 2 0..2 matrices with each 2 X 2 subblock idempotent.

Conjectures from _Colin Barker_, Feb 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007698 | OEIS-A224670

Number of (n+1) X 3 0..2 matrices with each 2 X 2 subblock idempotent.

Conjectures from _Colin Barker_, Feb 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007699 | OEIS-A224671

Number of (n+1) X 4 0..2 matrices with each 2 X 2 subblock idempotent.

Conjectures from _Colin Barker_, Feb 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007700 | OEIS-A224673

Number of (n+1) X 6 0..2 matrices with each 2 X 2 subblock idempotent.

Conjectures from _Colin Barker_, Feb 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007701 | OEIS-A224709

The number of unordered partitions {a,b} of the even numbers 2n such that a and b are composite.

Conjecture: a(3n+9) > a(3n+8) and a(3n+10) < a(3n+9) for n>=1. - _Anthony Browne_, Jun 26 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Stauduhar_, Apr 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007702 | OEIS-A224730

Smallest prime q such that 2n+1 = p^3 - 2q for some odd prime p, or 0 if no such prime exists.

See the conjecture and the comments in A185046, and the corresponding primes p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007703 | OEIS-A224748

Given n-th prime p, a(n) = number of primes of the form p + q - 1 where q is any prime < p.

Conjecture: a(n)>0 for all n>2. - _Dmitry Kamenetsky_, Feb 09 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jayanta Basu_, Apr 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007704 | OEIS-A224793

Least prime p which generates exactly n primes of the form p+q-1 where q < p is prime, or 0 if (conjecturally) no such p exists.

Least prime p which generates exactly n primes of the form p+q-1 where q < p is prime, or 0 if (conjecturally) no such p exists. These values of 0 are all conjectural. - _Robert Israel_, Apr 28 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jayanta Basu_, Apr 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007705 | OEIS-A224794

Smallest prime p such that 2n+1 = 2p - q^3 for some odd prime q, or 0 if no such prime exists.

Conjecture: The odd numbers different from a cube are of the form m = 2p - q^3 where p and q are prime numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007706 | OEIS-A224795

Smallest prime q such that 2n+1 = 2p - q^3 for some odd prime p, or 0 if no such prime exists.

See the conjecture and the comments in A224794, and the corresponding primes p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007707 | OEIS-A224832

Numbers k such that the sum of reciprocals of even divisors of k is an integer.

Conjecture: if a(n) is a non-twice even perfect numbers, the sum of reciprocals of even divisors equals 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jul 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007708 | OEIS-A224868

a(1) = greatest k such that H(k) - H(4) < 1/3 + 1/4; a(2) = greatest k such that H(k) - H(a(1)) < H(a(1)) - H(4); and for n > 2, a(n) = greatest k such that H(k) - H(a(n-1)) > H(a(n-1)) - H(a(n-2)), where H = harmonic number.

a(n) = 2*a(n-1) - a(n-2) + a(n-3) - a(n-4) (conjectured). G.f.: (7 - 3 x + 2 x^2 - 4 x^3)/(1 - 2 x + x^2 - x^3 + x^4) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007709 | OEIS-A224887

Honaker trios: Consecutive prime numbers p < q < r such that p | (qr+1).

Caldwell & Cheng show that there are only 3 Honaker trios below 2*10^17 and that these are the only Honaker trios if the Cramér-Granville conjecture is satisfied with a constant M < 199262; they also give other sufficient conditions for the number of Honaker trios to be finite. Strictly speaking, the keywords "fini,full" are thus only conjectured, but given that the Cramér-Granville conjecture is believed to hold with M = 2*exp(-gamma) ~ 1.123, it seems justified to use them.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jul 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007710 | OEIS-A224898

G.f.: Sum_{n>=0} (-1)^n* x^(n*(n+1)) / Product_{k=1..n} (1-x^k).

Conjecture: a(n+1) = A286744(n) - A286745(n). - _George Beck_ May 13 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007711 | OEIS-A224902

O.g.f.: exp(Sum_{n>=1} (sigma(2*n^4) - sigma(n^4)) * x^n/n).

a(n) == 2 (mod 4) for n>0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007712 | OEIS-A224928

Numbers of pairs {x, y} such that x <= y and triangular(x) + triangular(y) = 2^n.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, May 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007713 | OEIS-A224929

Primes such that concatenation of two adjacent terms and two adjacent digits is also prime.

Conjecture: all prime congruent to 1 (mod 3) which only have the odd digits {1, 3, 7, 9} but not 5 eventually appear. So far 139, 193, 199, 331, 337, 397, 733, 739, 919, 937, 991, 997, etc. have not appeared. - _Robert G. Wilson v_, May 06 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lars Blomberg_, inspired by _Eric Angelini_, Apr 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007714 | OEIS-A224935

Numbers n such that a positive number m <= n exists such that n-m, n+m, and n*m are triangular numbers.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Apr 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007715 | OEIS-A224954

Numbers n such that a positive number m < n exists such that n-m, n+m, and n*m are oblong numbers (A002378).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Apr 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007716 | OEIS-A224962

a(n) = number of primes of the form p*q+(p+q) where p = prime(n) and q is any prime < p.

Conjecture: a(n) > 0 for all n > 1. - _Dmitry Kamenetsky_, Jul 18 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jayanta Basu_, Apr 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007717 | OEIS-A224986

a(n) = Product_{k=1..n-4} (n-k-2)!^(k*k!).

Consider words on n symbols that contain every permutation of those n symbols as contiguous substrings. The minimal length of such a string was conjectured to equal A007489(n) (see A180632). This sequence is a lower bound on the number of distinct (up to relabeling the symbols) such strings of the conjectured minimal length. It was conjectured in the Ashlock paper that, for all n, there is only one string of length A007489(n) containing all permutations. This sequence shows that this conjecture fails as n grows. In 2014 Houston has shown that the first conjecture about the minimal length is also false for all n > 5. In particular, A180632(6) <= 872 = A007489(n). - _M. F. Hasler_, Jul 28 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nathaniel Johnston_, Apr 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007718 | OEIS-A225008

Number of n X 6 0..1 arrays with rows unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Sep 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007719 | OEIS-A225009

Number of n X 7 0..1 arrays with rows unimodal and columns nondecreasing.

Conjectures from _Colin Barker_, Sep 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007720 | OEIS-A225034

a(n) is the number of binary words containing n 1's and at most n 0's that do not contain the substring 101.

Conjecture: (n+1)*a(n) - (2*n+3)*a(n-1) - 3*(n-2)*a(n-2) = 0 for n>1. - _Bruno Berselli_, May 02 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Elisabetta Grazzini_, May 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007721 | OEIS-A225040

a(n) is the position of prime(n) in A225039, and a(n)=0, if prime(n) is not in A225039.

Conjecture: All terms are positive, or equivalently, the sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007722 | OEIS-A225053

Second terms of continued fractions for power towers e, e^e, e^e^e, ...

It was conjectured (but remains unproved) that none of the power towers e, e^e, e^e^e, ... are integers. If so, the corresponding continued fractions contain at least 2 terms. If the conjecture fails, let the corresponding a(n) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Reshetnikov_, Apr 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007723 | OEIS-A225060

Discriminants D < 0 such that h(D) > h(D') for D < D' < 0, negated.

Conjecture: except for a(1) = 3, terms are congruent 7 (mod 8). _Ralf Stephan_, May 23 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Apr 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007724 | OEIS-A225064

Decimal expansion of the fractional part of e^e^e^e.

It was conjectured (but remains unproved) that this sequence is infinite and aperiodic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Reshetnikov_, Apr 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007725 | OEIS-A225093

a(1)=2, a(2)=3, for n>=3, a(n) is the n-th number which is obtained by application of Eratosthenes-like sieve to A225091.

We conjecture that every term is prime; moreover, we conjecture that the sequence is a permutation of the sequence of all primes. - _Vladimir Shevelev_, Dec 17 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007726 | OEIS-A225106

Prime powers (A025475) that can be represented as a sum of two distinct prime powers.

Conjecture: the sequence is finite. - _Alex Ratushnyak_, Feb 03 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Apr 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007727 | OEIS-A225114

Number of skew partitions of n whose diagrams have no empty rows and columns.

Conjectured g.f.: 1/(2 - 1/(1 - x/(1 - x/(1 - x^2/(1 - x^2/(1 - x^3/(1 - x^3/(1 - ...)))))))). - _Mikhail Kurkov_, Sep 03 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Apr 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007728 | OEIS-A225127

Convolutory inverse of the nonprimes.

Coefficients in 1/(1+g(x)), where g is the generating functions of the sequence of nonprimes: (1,4,6,8,9,...). For the convolutory inverse of the primes, see A030018. Conjecture: a(n+1)/a(n) has a limit, -2.4633754095588889..., analogous to the Backhouse constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007729 | OEIS-A225128

Numerators of the convolutory inverse of the primes of the form 4m+3.

Coefficients in 1/(1+g(x)), where g is the generating functions of the sequence of primes (3,7,11,19,23,31,...) of primes congruent to 3 mod 4. For the convolutory inverse of the primes, see A030018. Conjecture: a(n+1)/a(n) -> -1.370819405....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007730 | OEIS-A225129

Numerators of the convolutory inverse of the primes of the form 4m+1.

Coefficients in 1/(1+g(x)), where g is the generating functions of the sequence of primes (5,13,17,29,37,...) of primes congruent to 1 mod 4. For the convolutory inverse of the primes, see A030018. Conjecture: a(n+1)/a(n) -> -1.59045463062282....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007731 | OEIS-A225130

Numerators of the convolutory inverse of the primes of the form 6m-1.

Coefficients in 1/(1+g(x)), where g is the generating functions of the sequence of primes (5,11,17,23,29,...) of primes congruent to -1 mod 6. For the convolutory inverse of the primes, see A030018. Conjecture: a(n+1)/a(n) -> -1.24066....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007732 | OEIS-A225131

Numerators of the convolutory inverse of the primes of the form 6m + 1 (A007528).

Coefficients in 1/(1+g(x)), where g is the generating functions of the sequence of primes (7,13,19,31,37,...) of primes congruent to 1 mod 6. For the convolutory inverse of

the primes, see A030018. Conjecture: a(n+1)/a(n) diverges.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007733 | OEIS-A225132

Convolutory inverse of the Thue Morse sequence.

Coefficients in 1/(1+g(x)), where g is the generating function of the Thue-Morse sequence, A010060. Conjecture: a(n+1)/a(n) -> -1.405177106052... .

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007734 | OEIS-A225147

a(n) = Im((1-I)^(1-n)*A_{n, 3}(I)) where A_{n, k}(x) are the generalized Eulerian polynomials.

G.f.: conjecture -T(0)/(1+2*x), where T(k) = 1 - 9*x^2*(k+1)^2/(9*x^2*(k+1)^2 + (1+2*x)^2/T(k+1)); (continued fraction). - _Sergei N. Gladkovskii_, Nov 12 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Apr 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007735 | OEIS-A225179

a(n) = min{2 + c_n + 2(c_1 + c_2 + ... + c_(n-1)) | n = p + q, 1 <= p < q < n, gcd(p, q) = 1, and p/q has continued fraction expansion [0; c_1, c_2, ..., c_n]}.

It is conjectured that a(n) = min{ B(w) | w is unbordered and length(w) = n }, where B(w) is the number of distinct unbordered factors in w. If this conjecture were to be proved, we could use this is the definition, and put the present definition into the FORMULA field. It is known to be true for n <= 20.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 03 2013, based on an email from Kalle Saari, Apr 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007736 | OEIS-A225322

Lucky numbers that are prime powers.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, May 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007737 | OEIS-A225389

Prime powers (A025475) representable as (p+q)/2, where p and q are distinct prime powers, in two or more ways.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, May 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007738 | OEIS-A225400

First occurrence of n in A225399, or -1 if n does not appear in A225399.

Conjecture: a(n) >= 0. A000217(a(n)) = A076983(n+1) for n>0, if the conjecture is true and a(n) >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, May 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007739 | OEIS-A225502

Least m > 0 such that prime(n)*triangular(m) is a triangular number, or 0 if no such m exists.

Conjecture: a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, May 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007740 | OEIS-A225503

Least triangular number t such that t = prime(n)*triangular(m) for some m>0, or 0 if no such t exists.

Conjecture: a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, May 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007741 | OEIS-A225516

Primes corresponding to terms of A225430.

Conjecture: the sequence is a permutation of all primes > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007742 | OEIS-A225518

Number of ways to express 2n+1 as p + 8*q, with p and q primes.

This is related to the conjecture given in A223174 and A223175.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, May 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007743 | OEIS-A225538

Let r(n) denote the reverse of n. For every n, consider the sequence n_1 = n + 1 + r(n+1), and for m >= 2, n_m = n_(m-1) + 1 + r(n_(m-1) + 1). a(n) is the least m for which n_m is a palindrome, or 0 if there is no such m.

Conjecture: the least n's for which a(n) = 0 are 1895, 1985, 2894, 2984, 3893, and 3983. - _Peter J. C. Moses_, May 10 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007744 | OEIS-A225551

Longest checkmate in king and queen versus king endgame on an n X n chessboard.

Conjecture: For n > 8, a(n) = 2*n-3-floor(n/2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, May 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007745 | OEIS-A225552

Longest checkmate in king and rook versus king endgame on an n X n chessboard.

Conjecture: For n > 24, a(n) = 2*n - 3 + floor((n+1)/2) - floor(n/6).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, May 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007746 | OEIS-A225553

Longest checkmate in king and amazon versus king endgame on an n X n chessboard.

Conjecture: for n > 10, a(n) = floor((n+2)/2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, May 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007747 | OEIS-A225554

Longest checkmate in king and empress versus king endgame on an n X n chessboard.

Conjecture: a(n) ~ 7*n/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, May 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007748 | OEIS-A225556

Longest checkmate in king, bishop and knight versus king endgame on an n X n chessboard.

Conjecture: a(n) ~ n^2/6 + 3*n if n is even and a(n) ~ n^2/6 + 2*n if n is odd.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, May 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007749 | OEIS-A225557

Longest checkmate in king and two bishops versus king endgame on an n X n chessboard.

Conjecture: a(n) ~ 11*n/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, May 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007750 | OEIS-A225563

Numbers whose totient-trajectory can be partitioned into two sets with the same sum.

Conjecture: No totient-trajectory can be partitioned into an odd number of sets with the same sum.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kevin L. Schwartz_ and _Christian N. K. Anderson_, May 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007751 | OEIS-A225577

Least integer m>1 such that 1^2,2^2,...,n^2 are pairwise incongruent modulo 2^m-1.

Conjecture: a(n) is the least prime p such that 2^p-1 is a Mersenne prime greater than 2n-1. This conjecture implies that there are infinitely many Mersenne primes. Zhi-Wei Sun also conjectured that for each n>17 the least Fibonacci number modulo which 1^2,2^2,...,n^2 are pairwise incongruent is just the first Fibonacci prime greater than 2n-1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007752 | OEIS-A225595

Conjectured square array T(n,k) read by antidiagonals related to the existence of rectangles of size n*k in the toothpick structure of A139250.

Conjectured square array T(n,k) read by antidiagonals related to the existence of rectangles of size n*k in the toothpick structure of A139250.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, May 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007753 | OEIS-A225603

Palindromic primes whose square is also a palindrome.

Palindromes in A161721. Conjecture: a(n) for n >=3 consists only of the digits 0,1. - _Chai Wah Wu_, Jan 06 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jayanta Basu_, May 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007754 | OEIS-A225605

a(1) = least k such that 1/3 < H(k) - 1/3; a(2) = least k such that H(a(1)) - H(3) < H(k) - H(a(1)), and for n > 2, a(n) = least k such that H(a(n-1)) - H(a(n-2)) > H(k) - H(a(n-1)), where H = harmonic number.

For A225605, (x,y) = (3,3); it appears that H(a(n)) - H(a(n-1)) approaches 0.60937786343... ; it is conjectured a(n)/a(n-1) approaches the constant given at A058265.. a(n) = A192804(n+4) (conjectured). a(n) = 2*a(n-1) - a(n-4) (conjectured). G.f.: (5 - x - 2 x^2 - 3 x^3)/(1 - 2 x + x^4) (conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007755 | OEIS-A225611

Determinant of the (p_n-1) X (p_n-1) matrix with (i,j)-entry equal to the Legendre symbol ((i^2+6*i*j+j^2)/p_n), where p_n is the n-th prime.

Conjecture: We have a(n) = 0 if p_n == 3 (mod 4). For an odd prime p and integers c and d, _Zhi-Wei Sun_ defined (c,d)_p to be the determinant of the (p-1) X (p-1) matrix whose (i,j)-entry is the Legendre symbol ((i^2+c*i*j+d*j^2)/p). It is easy to see that (-c,d)_p = (-1/p)*(c,d)_p. Sun conjectured that for any integer c and nonzero integer d there are infinitely many odd primes p with (c,d)_p = 0, moreover (c,d)_p = 0 if (d/p) = -1. He also formulated some concrete conjectures in the case (d/p) = 1. For example, (3,3)_p = 0 if p == 11 (mod 12), and (10,9)_p = 0 if p == 5 (mod 12); (3,2)_p = (4,2)_p = 0 if p == 7 (mod 8). On August 12 2013, _Zhi-Wei Sun_ conjectured that for any odd prime p and integers c and d with d not divisible by p, if (c,d)_p is nonzero then its p-adic valuation (i.e., p-adic order) must be even.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007756 | OEIS-A225620

Indices of partitions in the table of compositions of A228351.

Conjecture: this sequence is a sorted version of b(n) where b(2^k) = 2^k for k >= 0, b(n) = A080100(n)*(2*b(A053645(n)) + 1) otherwise. - _Mikhail Kurkov_, Oct 21 2023 Conjecture: a(A000070(m) - k) = 2^m - A228354(k) for m > 0, 0 < k <= A000041(m). - _Mikhail Kurkov_, Oct 20 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007757 | OEIS-A225660

Number of nonrepeating vectors in a counting procedure that starts with the digits of (n base 2).

Conjecture: if n>9, then a(n) is odd.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007758 | OEIS-A225661

Number of nonrepeating vectors in a counting procedure that starts with the digits of (n base 3).

The counting procedure and "eventually period 6 theorem" are introduced at A225660. Conjecture: if a(n) > 6, then a(n) is odd.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007759 | OEIS-A225663

Number of nonrepeating vectors in a counting procedure that starts with the digits of (n base 5).

The counting procedure and "eventually period 6 theorem" are introduced at A225660. Conjecture: if a(n) > 8, then a(n) is odd.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007760 | OEIS-A225664

Number of nonrepeating vectors in a counting procedure that starts with the digits of (n base 10).

The counting procedure and "eventually period 6 theorem" are introduced at A225660. Conjecture: if a(n) is odd for only finitely many n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007761 | OEIS-A225684

Nonpalindromic numbers n with property that the sum of the reversed divisors of n is equal to n+1.

Call a number "quasi-perfect" or "slightly excessive" if sigma(n) = 2n+1 (cf. A000203). It is conjectured that no quasi-perfect number exists. The present sequence is a variation that certainly has at least four terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007762 | OEIS-A225692

Number of Dyck paths of semilength n avoiding the pattern U^(n-1) D^(n-1).

Conjecture: -10*(n+1)*(n-4)*a(n) +(51*n^2-196*n+80)*a(n-1) +(-45*n^2+134*n-99)*a(n-2) +2*(2*n-5)*(n-3)*a(n-3)=0. - _R. J. Mathar_, Jul 09 2013 Conjecture: a(n) ~ 4^n / (sqrt(Pi) * n^(3/2)). - _Vaclav Kotesovec_, Nov 30 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007763 | OEIS-A225759

Primes p such that A179382((p+1)/2) = (p-1)/16.

Conjecture 1: If n-1=2^x*c for some x<5, then n is prime. If x > 4, the relative density of primes in such numbers is 1. Conjecture 2: If the period of the decimal expansion of 1/n is n-1 or a divisor of n-1, and if n-1=2^x*c or n+1=2^x*c for some x, then n is prime. The numbers in the sequence are the values of n in the above conjecture when c=4 in case (1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lear Young_, May 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007764 | OEIS-A225775

Numbers k such that antisigma(x) = antisigma(x+1) = k has solution.

Conjecture: also numbers k such that antisigma(n) = antisigma(m) = k has solution for distinct numbers n and m. - _Jaroslav Krizek_, Nov 10 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jul 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007765 | OEIS-A225776

Determinant of the (n+1) X (n+1) matrix with (i,j)-entry equal to f(i+j) for all i,j = 0,...,n, where f(k) = A000172(k) is the k-th Franel number.

Conjecture: a(n)/6^n is always a positive odd integer. Moreover, for any integers r > 1 and n >= 0, the number a(r,n)/2^n is a positive odd integer, where a(r,n) denotes the Hankel determinant |f(r,i+j)|_{i,j=0,...,n} with f(r,k) = Sum_{j=0..k} C(k,j)^r. On Aug 20 2013, _Zhi-Wei Sun_ made the following conjecture: If p is a prime congruent to 1 mod 4 but p is not congruent to 1 mod 24, then p divides a((p-1)/2). Zhu and Sun (2018, Theorem 1.1) proved the conjecture that a(n)/6^n is always a positive odd integer. - _Eiver Rodríguez_, Sep 12 2026 Conjecture: a(n)^(1/n^2) tends to 9/4. - _Vaclav Kotesovec_, Sep 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007766 | OEIS-A225790

Rubik's Square sequence.

Conjecture: a(2*n) = 2^A142463(n) * 3^(n^2) = 2^A142463(n) * 3^A000290(n). - _Eric M. Schmidt_, Nov 05 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sean D Lawton_, May 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007767 | OEIS-A225819

Consider the set of n-tuples such that the sum of cubes of the elements is equal to square of their sum; sequence gives largest element in all such tuples.

Conjecture [Sen]: lim inf log_n a(n) >= 5/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, _Jimmy Zotos_, and _Balarka Sen_, Jul 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007768 | OEIS-A225835

Smallest prime p such that there is a prime q satisfying (2*n + 1)*p^2 - (2*n-1)*q^2 = 2, or 0 if no such p exists.

Conjecture: a(6) = a(7) = 0. _Charles R Greathouse IV_ reports that a(6) must have thousands of digits. - _Michael B. Porter_, May 19 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Irina Gerasimova_, May 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007769 | OEIS-A225855

Difference between the number of nonnegative evil and odious multiples of n less than 2^n.

For prime n and related conjectures, see A133954.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Aug 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007770 | OEIS-A225868

Numbers m for which max_{2 <= k <= (m-2)/2} Sum_{d>1, d|m+k, k|m+d} 1 = 3.

Terms >= 53 are primes p such that p+2 is either prime or semiprime or, relatively rarely, the cube of a prime. However, according to calculations by _Peter J. C. Moses_, up to 4.2*10^13 there are no numbers p in the sequence for which p+2 is cube of a prime. One can prove that if such a prime p exists, then it is necessary (but not sufficient) for all numbers of the quadruple {r, 2*r - 1, 4*r^2 - 6*r + 3, (2*r - 1)^3 - 2} to be primes, where r == 19 (mod 30) is defined by the equality (2r-1)^3 - 2 = p. The first 3 suitable values of r are 229, 3109, and 17449. But the corresponding p's are not in the sequence. We conjecture that all primes of the sequence are Chen primes, that is, all of them are in A109611.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007771 | OEIS-A225887

a(n) = A212205(2*n + 1).

Conjecture: (n+1)*a(n) +3*(-4*n-1)*a(n-1) +(37*n-20)*a(n-2) +6*(-n+2)*a(n-3)=0. - _R. J. Mathar_, May 23 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, May 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007772 | OEIS-A225889

Least prime p_m such that n = p_m-p_{m-1}+...+(-1)^(m-k)*p_k for some 0<k<m, where p_j denotes the j-th prime.

By a conjecture of the author, a(n) <= 2*n+2.2*sqrt(n), and moreover a(n) <= n+4.6*sqrt(n) if n is odd. Clearly a(n)>n. We guess that a(2n)/(2n) --> 2 as n tends to the infinity. Note that this sequence is different from A222579 which involves a stronger conjecture of the author. Zhi-Wei Sun also conjectured that any positive even integer m can be written in the form p_n-p_{n-1}+...+(-1)^{n-k}*p_k with k < n and 2m-3.6*sqrt(m+1) < p_n < 2m+2.2*sqrt(m).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007773 | OEIS-A225894

Number of n X 2 binary arrays whose sum with another n X 2 binary array containing no more than a single 1 has rows and columns in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007774 | OEIS-A225895

Number of n X 3 binary arrays whose sum with another n X 3 binary array containing no more than a single 1 has rows and columns in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007775 | OEIS-A225907

Smallest n-Ramanujan prime that is less than half of the next n-Ramanujan prime, or 0 if none exists.

Only the first three terms 0, 2, 11 are proved (by Chebyshev, Ramanujan, and Paksoy, respectively). The rest are conjectural--see the 2nd comment in A192821.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Jun 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007776 | OEIS-A225913

Composite numbers coprime to 6 such that A179382(n) = A000265(n-1), the odd part of n-1.

Inspired by the Conjecture mentioned in A225759.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, May 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007777 | OEIS-A225918

a(n) is the least k such that f(a(n-1)+1) + ... + f(k) > f(a(n-2)+1) + ... + f(a(n-1)) for n > 1, where f(n) = 1/(n+3) and a(1) = 1.

Suppose that f(n) is a sequence of positive real numbers for which the series f(1) + f(2) + ... diverges. Put a(1) = 1 and a(n) = least k such that f(a(n-1)+1) + ... + f(k) > f(a(n-2)+1) + ... + f(a(n-1)) for n > 1. Conjecture: a(n) is linearly recurrent for the choices of f(n) shown here:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007778 | OEIS-A225919

a(n) is the least k such that f(a(n-1)+1) + ... + f(k) > f(a(n-2)+1) + ... + f(a(n-1)) for n > 1, where f(n) = 1/(n+4) and a(1) = 1.

Conjecture: a(n) is linearly recurrent. See A225918 for details.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007779 | OEIS-A225920

a(n) is the least k such that f(a(n-1)+1) + ... + f(k) > f(a(n-2)+1) + ... + f(a(n-1)) for n > 1, where f(n) = 1/(n+5) and a(1) = 1.

Conjecture: a(n) is linearly recurrent. See A225918 for details.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007780 | OEIS-A225921

a(n) is the least k such that f(a(n-1)+1) + ... + f(k) > f(a(n-2)+1) + ... + f(a(n-1)) for n > 1, where f(n) = 1/(n+6) and a(1) = 1.

Conjecture: a(n) is linearly recurrent. See A225918 for details.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007781 | OEIS-A225922

a(n) is the least k such that f(a(n-1)+1) + ... + f(k) > f(a(n-2)+1) + ... + f(a(n-1)) for n > 1, where f(n) = 1/(n+7) and a(1) = 1.

Conjecture: a(n) is linearly recurrent. See A225918 for details.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007782 | OEIS-A225976

Number of n X 2 binary arrays whose sum with another n X 2 binary array containing no more than two 1s has rows and columns in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007783 | OEIS-A225977

Number of n X 3 binary arrays whose sum with another n X 3 binary array containing no more than two 1s has rows and columns in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007784 | OEIS-A226014

Primes p such that A179382((p+1)/2) = (p-1)/(2^x) for some x>0.

It is conjectured that: The numbers in the sequence are the values of n in the above conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lear Young_, May 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007785 | OEIS-A226036

Let abc... be the decimal expansion of n. a(n) is the number of iterations of the map n -> f(n) needed to reach the last number of the cycle, where f(n) = a^a + b^b + c^c + ...

It is conjectured that the trajectory for every number converges to a single number. The growth of a(n) is very slow; for example, a(457) = 211, a(10337) = 213, a(16669) = 214, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, May 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007786 | OEIS-A226076

Lexicographically least sequence of squares that are sum-free.

Conjecture: a(n) = 4*a(n-3) for n>9. G.f.: -x*(33*x^8 +112*x^7 +80*x^6 +28*x^5 +20*x^4 +12*x^3 +9*x^2 +4*x +1) / (4*x^3 -1). - _Colin Barker_, May 28 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, May 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007787 | OEIS-A226115

Least positive integer not of the form p_m - p_{m-1} + ... +(-1)^(m-k)*p_k with 0 < k < m <= n, where p_j denotes the j-th prime.

Conjecture: sqrt(2*a(n)) > sqrt(p_n)-0.7 for all n > 0, and a(n) is even for any n > 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007788 | OEIS-A226135

Let abcd... be the decimal expansion of n. Number of iterations of the map n -> f(n) needed to reach a number < 10, where f(n) = a^b + c^d + ... which ends in an exponent or a base according as the number of digits is even or odd.

Inspired by the sequence A133501 (Number of steps for "powertrain" operation to converge when started at n). It is conjectured that the trajectory for each number converges to a single number < 10. The conjecture is true, since f(x) < x trivially holds for x > 10^10 and I have verified that for every 10 <= x <= 10^10 there is a k such that f^(k)(x) < x, where f^(k) denotes f applied k times. Both the conventions 0^0 = 1 and 0^0 = 0 have been taken into account. - _Giovanni Resta_, May 28 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, May 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007789 | OEIS-A226136

Positions of the positive integers in the ordering of rational numbers as generated by the rules: 1 is in S, and if nonzero x is in S, then x+1 and -1/x are in S. (See Comments.)

Conjecture: a(n) = a(n-1)+a(n-3) for n>6. G.f.: -x*(x+1) * (x^2+1)^2 / (x^3+x-1). - _Colin Barker_, Jul 03 2013 Conjectures confirmed (see Reid link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007790 | OEIS-A226158

a(n) = 2*n*(2^n - 1)*zeta(1-n) where in the case n=0 the limit is understood, zeta(s) the Riemann zeta function.

G.f.: conjecture: -x/Q(0),where Q(k) = 1 - x*(k+1)/(1 + x*(k+1)/(1 - x*(k+1)/(1 + x*(k+1)/Q(k+1)))); (continued fraction). - _Sergei N. Gladkovskii_, Nov 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jun 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007791 | OEIS-A226160

Least positive integer k such that 1 + 1/2 + ... + 1/k > n/phi, where phi = golden ratio = (1+sqrt(5))/2.

Conjecture: a(n+1)/a(n) converges to 1.8552... Conjecture confirmed: using series expansion of HarmonicNumber(k) one gets a(n+1)/a(n) -> exp(1/phi) = 1.855276958... [_Jean-François Alcover_, Jun 04 2013]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007792 | OEIS-A226161

Least positive integer k such that 1 + 1/2 + ... + 1/k > n/2.

Conjecture: a(n+1)/a(n) converges to 1.64872... The conjecture is correct, a(n+1)/a(n) ~ exp(1/2) (A019774). - _Charles R Greathouse IV_, Jun 03 2013 Conjecture: a(n) = round(exp(n/2-gamma)) for all n, where gamma is the Euler-Mascheroni constant (see A001620). - _Jon E. Schoenfield_, Jul 19 2015 All terms in the b-file are correct. Moreover, the above conjecture that a(n) = round(exp(n/2-gamma)) has been verified for all n in 1..10000. - _Jon E. Schoenfield_, Jul 22 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007793 | OEIS-A226163

Determinant of the (p_n-1)/2-by-(p_n-1)/2 matrix with (i,j)-entry being the Legendre symbol ((i^2-((p_n-1)/2)!*j)/p_n), where p_n is the n-th prime.

Conjecture: a(n) = 0 if and only if p_n == 3 (mod 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007794 | OEIS-A226187

Least positive integer k such that 1 + 1/2 + ... + 1/k > n/3.

Conjecture: a(n+1)/a(n) converges to 1.39...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007795 | OEIS-A226188

Least positive integer k such that 1 + 1/2 + ... + 1/k > 2n/3.

Conjecture: a(n+1)/a(n) converges to 1.94... Conjecture confirmed: series expansion of HarmonicNumber(k) gives a(n+1)/a(n) -> exp(2/3) = 1.947734... - _Jean-François Alcover_, Jun 05 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007796 | OEIS-A226191

Odd numbers n such that n*2^k + 1 is a Carmichael number for some k.

Terms after the first are conjectural. a(2) could be proved by an argument like that on p. 17 on 33 with the addition of the prime 257 (29 and 31 are not in this sequence by 7.1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, May 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007797 | OEIS-A226224

The largest value of k in base n for which the sum of digits of k = sqrt(k).

a(n)=1 iff A226087(n)=1. Conjecture: this occurs exactly once -- in base 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian N. K. Anderson_ and _Kevin L. Schwartz_, May 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007798 | OEIS-A226264

Number of additive Z_2 Z_8 codes of a certain type (see Siap-Aydogdu for precise definition).

Conjectures from _Colin Barker_, Jun 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007799 | OEIS-A226270

Number of primitive permutations with n buds and 3 red or blue elements.

Conjectural recurrence: a(n) = (1/3)*(n+1)*d(n+2) - Sum_{k = 0..n-1} d(n+1-k)*a(k) with a(0)= 1, where d(n) = (2*n-1)!! = A001147(n). If true then the sequence continues [1, 7, 69, 843, 12081, 197127, 3595509, 72393003, 1594224801, 38123341767, 984142175589, ...]. - _Peter Bala_, Dec 26 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007800 | OEIS-A226302

a(n) = P_n(-1), where P_n(x) is a certain polynomial arising in the enumeration of tatami mat coverings.

Conjecture: (-n+2)*a(n) +(-n+2)*a(n-1) +2*(3*n-11)*a(n-2) +2*(3*n-14)*a(n-3) +4*(-2*n+9)*a(n-4) +8*(-n+6)*a(n-5)=0. - _R. J. Mathar_, Nov 06 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007801 | OEIS-A226316

Expansion of g.f. 1/2 + 1/(1+sqrt(1-8*x+8*x^2)).

Conjecture: (n+1)*a(n) +3*(-3*n+1)*a(n-1) +4*(4*n-5)*a(n-2) +8*(-n+2)*a(n-3)=0. - _R. J. Mathar_, Apr 02 2015 Conjecture verified using differential equation -4*x^3 + 6*x^2 - 4*x + 1 + (8*x^3 - 12*x^2 + 6*x - 1)*g(x) + (8*x^4 - 16*x^3 + 9*x^2 - x)*g'(x) = 0 satisfied by the g.f. - _Robert Israel_, Feb 23 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007802 | OEIS-A226318

Positive integers n with p_{n+1}-p_n = 2 and p_{n+3}-p_{n+2} = 2, where p_k denotes the k-th prime.

An old conjecture of de Polignac asserts that for any positive even integer d there are infinitely many n>0 with p_{n+1}-p_n = d. Conjecture: For any positive even integers d_1,...,d_k, there are infinitely many positive integers n such that p_{n+2j-1}-p_{n+2j-2} = d_j for all j=1,...,k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007803 | OEIS-A226334

Number of nondecreasing -2..2 vectors of length n whose dot product with some other -2..2 vector equals n.

Conjectures from _Colin Barker_, Mar 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007804 | OEIS-A226434

The number of sum decomposable permutations which avoid the patterns 3124 and 4312.

Conjecture: +(95*n+537)*(n+2)*a(n) +(95*n^2-16421*n-14748) *a(n-1) +(-6403*n^2+124495*n-60066) *a(n-2) +(21565*n^2-354883*n+596496) *a(n-3) +2*(-5092*n^2+138877*n-395970) *a(n-4) +8*(-2470*n^2+11113*n+12744) *a(n-5) +192*(38*n-67)*(2*n-13)*a(n-6)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jay Pantone_, Sep 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007805 | OEIS-A226435

Number of permutations of {1..n} with fewer than 2 interior elements having values lying between the values of their neighbors.

E.g.f. (conjecture): (sec(x) + tan(x)) - (sec(x) + tan(x))^2 + (sec(x) + tan(x))^3. - _Sergei N. Gladkovskii_, Jun 11 2015 E.g.f. (conjecture): 1 + x + (sec(x) + tan(x) - 1)^2 = x + (1 - sec(x))^2 + (1 - tan(x))^2 + 2*sec(x)*tan(x). - _Peter Bala_, Feb 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007806 | OEIS-A226476

Numbers n with the property that, if tau(n) = k = number of divisors of n, and the d(i) are the divisors [arranged in increasing order], then the sum 1/d(k) + 1/d(k-1) + 1/d(k-2) + ... + 1/d(q) is an integer for some q.

q = 2 if n = 1, 6, 28, 120, 496, 672, 8128, ... is a multiply-perfect number (see A007691 where it is conjectured that this sequence is infinite), which would imply that this sequence is also infinite because A007691 is a subsequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jun 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007807 | OEIS-A226513

Array read by antidiagonals: T(n,k) = number of barred preferential arrangements of k things with n bars (k >=0, n >= 0).

Conjectural g.f. for row n as a continued fraction of Stieltjes type: 1/(1 - (n+1)*x/(1 - 2*x/(1 - (n+2)*x/(1 - 4*x/(1 - (n+3)*x/(1 - 6*x/(1 - ...))))))). Cf. A265609. - _Peter Bala_, Aug 27 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007808 | OEIS-A226515

Row 2 of array in A226513.

Conjectural g.f. as a continued fraction of Stieltjes type: 1/(1 - 3*x/(1 - 2*x/(1 - 4*x/(1 - 4*x/(1 - 5*x/(1 - 6*x/(1 - (n+2)*x/(1 - 2*n*x/(1 - ...))))))))). - _Peter Bala_, Aug 27 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007809 | OEIS-A226518

Irregular triangle read by rows: T(n,k) = Sum_{i=0..k} Legendre(i,prime(n)).

There is a famous open problem concerning upper bounds on |T(n,k)| for small k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007810 | OEIS-A226525

Fortunate semiprimes: least m > 1 such that m + sp(n)# is semiprime, where sp# denotes the product of the semiprimes <= sp.

Just like the Fortunate primes (A005235) we conjecture that all terms are prime!

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sean A. Irvine_, _Jonathan Vos Post_ and _Robert G. Wilson v_, Jun 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007811 | OEIS-A226607

Irregular array read by rows in which row floor(k/3)+1, where gcd(k,6)=1, lists the smallest elements, in ascending order, of conjecturally all primitive cycles of positive integers under iteration by the 3x+k function.

Irregular array read by rows in which row floor(k/3)+1, where gcd(k,6)=1, lists the smallest elements, in ascending order, of conjecturally all primitive cycles of positive integers under iteration by the 3x+k function. Lagarias (1990) called

a T_k cycle primitive if its elements are all relatively prime to k or, equivalently, if its elements are not a common multiple of the elements of another cycle. He conjectured that, for every positive integer k relatively prime to 6, there is at least one primitive cycle of the map T_k and that the number of such cycles is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007812 | OEIS-A226613

a(n) is the conjectured number of primitive cycles of positive integers under iteration by the Collatz-like 3x+k function, where n=floor(k/3)+1.

a(n) is the conjectured number of primitive cycles of positive integers under iteration by the Collatz-like 3x+k function, where n=floor(k/3)+1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007813 | OEIS-A226617

Smallest positive integer k (or 0 if no such k) with a primitive cycle of positive integers, n of which are odd including 1, under iteration by the Collatz-like 3x+k function.

Conjecture: a(n)>0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jul 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007814 | OEIS-A226623

Irregular array read by rows in which row n lists the smallest elements, in ascending order, of conjecturally all primitive cycles of positive integers under iteration by the Collatz-like 3x-k function, where k = A226630(n).

Irregular array read by rows in which row n lists the smallest elements, in ascending order, of conjecturally all primitive cycles of positive integers under iteration by the Collatz-like 3x-k function, where k = A226630(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007815 | OEIS-A226629

a(n) is the conjectured number of primitive cycles of positive integers under iteration by the Collatz-like 3x-k function, where k=A226630(n).

a(n) is the conjectured number of primitive cycles of positive integers under iteration by the Collatz-like 3x-k function, where k=A226630(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007816 | OEIS-A226661

Smallest positive integer k (or 0 if no such k) with a primitive cycle of positive integers, exactly n of which are odd, under iteration by the Collatz-like 3x+k function.

Conjecture: a(n)>0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jul 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007817 | OEIS-A226662

Smallest positive integer k (or 0 if no such k) with conjecturally exactly n primitive cycles of positive integers under iteration by the Collatz-like 3x+k function.

Smallest positive integer k (or 0 if no such k) with conjecturally exactly n primitive cycles of positive integers under iteration by the Collatz-like 3x+k function. Belaga and Mignotte (2000, p.19) conjectured that the number of primitive cycles attains all positive integer values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007818 | OEIS-A226663

Conjectured record-breaking numbers, for ascending positive integers k, of primitive cycles of positive integers under iteration by the Collatz-like 3x+k function.

Conjectured record-breaking numbers, for ascending positive integers k, of primitive cycles of positive integers under iteration by the Collatz-like 3x+k function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007819 | OEIS-A226665

Conjectured record-breaking maximal values, for ascending positive integers k, of the minimal elements of the primitive cycles of positive integers under iteration by the Collatz-like 3x+k function.

Conjectured record-breaking maximal values, for ascending positive integers k, of the minimal elements of the primitive cycles of positive integers under iteration by the Collatz-like 3x+k function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007820 | OEIS-A226666

Integer k associated with the conjectured record-breaking maximal value A226665(n) of the minimal elements of primitive Collatz-like 3x+k cycles.

Integer k associated with the conjectured record-breaking maximal value A226665(n) of the minimal elements of primitive Collatz-like 3x+k cycles.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007821 | OEIS-A226667

Conjectured record-breaking values, for ascending positive integers k, of the maximal element of the primitive cycles of positive integers under iteration by the Collatz-like 3x+k function.

Conjectured record-breaking values, for ascending positive integers k, of the maximal element of the primitive cycles of positive integers under iteration by the Collatz-like 3x+k function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007822 | OEIS-A226668

Integer k associated with the conjectured record-breaking maximal element A226667(n) of primitive Collatz-like 3x+k cycles.

Integer k associated with the conjectured record-breaking maximal element A226667(n) of primitive Collatz-like 3x+k cycles.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007823 | OEIS-A226669

Smallest integer in the primitive Collatz-like 3x+k cycle with the conjectured record-breaking maximal element A226667(n).

Smallest integer in the primitive Collatz-like 3x+k cycle with the conjectured record-breaking maximal element A226667(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007824 | OEIS-A226670

Record-breaking values, for increasing positive integers k == 1 or 5 mod 6, of the conjectured length of the longest primitive cycle(s) of positive integers under iteration by the Collatz-like 3x+k function.

Record-breaking values, for increasing positive integers k == 1 or 5 mod 6, of the conjectured length of the longest primitive cycle(s) of positive integers under iteration by the Collatz-like 3x+k function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007825 | OEIS-A226671

Integer k associated with the conjectured record-breaking length A226670(n) of primitive Collatz-like 3x+k cycles.

Integer k associated with the conjectured record-breaking length A226670(n) of primitive Collatz-like 3x+k cycles.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007826 | OEIS-A226672

Smallest positive integer in the primitive Collatz-like 3x+k cycle(s) with the conjectured record-breaking length A226670(n).

Smallest positive integer in the primitive Collatz-like 3x+k cycle(s) with the conjectured record-breaking length A226670(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007827 | OEIS-A226673

Conjectured record-breaking numbers of odd elements, for ascending positive integers k, in primitive cycles of positive integers under iteration by the Collatz-like 3x+k function.

Conjectured record-breaking numbers of odd elements, for ascending positive integers k, in primitive cycles of positive integers under iteration by the Collatz-like 3x+k function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007828 | OEIS-A226674

Integer k associated with the conjectured record-breaking number A226673(n) of odd elements in a primitive Collatz-like 3x+k cycle.

Integer k associated with the conjectured record-breaking number A226673(n) of odd elements in a primitive Collatz-like 3x+k cycle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007829 | OEIS-A226675

Smallest element in the primitive Collatz-like 3x+k cycle(s) associated with the conjectured record-breaking number A226673(n) of odd elements.

Smallest element in the primitive Collatz-like 3x+k cycle(s) associated with the conjectured record-breaking number A226673(n) of odd elements.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007830 | OEIS-A226676

Smallest positive integer k (or 0 if no such k) with a primitive cycle of n positive integers under iteration by the Collatz-like 3x-k function.

Conjecture: For n>2, a(n)>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jul 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007831 | OEIS-A226678

Smallest positive integer (or 0 if no such k) with conjecturally exactly n primitive cycles of positive integers under iteration by the Collatz-like 3x-k function.

Smallest positive integer (or 0 if no such k) with conjecturally exactly n primitive cycles of positive integers under iteration by the Collatz-like 3x-k function. Conjecture:

a(n)>0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007832 | OEIS-A226679

Record-breaking values, for increasing k = A226630(n), of the conjectured number of primitive cycles of positive integers under iteration by the Collatz-like 3x-k function.

Record-breaking values, for increasing k = A226630(n), of the conjectured number of primitive cycles of positive integers under iteration by the Collatz-like 3x-k function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007833 | OEIS-A226682

Integer k associated with the conjectured record-breaking maximal value A226681(n) of the minimal elements of primitive Collatz-like 3x-k cycles.

Integer k associated with the conjectured record-breaking maximal value A226681(n) of the minimal elements of primitive Collatz-like 3x-k cycles.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007834 | OEIS-A226683

Conjectured record-breaking values, for ascending positive integers k, of the maximal element of the primitive cycles of positive integers under iteration by the 3x-k function.

Conjectured record-breaking values, for ascending positive integers k, of the maximal element of the primitive cycles of positive integers under iteration by the 3x-k function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007835 | OEIS-A226684

Integer k associated with the conjectured record-breaking maximal element A226683(n) of primitive 3x-k cycles.

Integer k associated with the conjectured record-breaking maximal element A226683(n) of primitive 3x-k cycles.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007836 | OEIS-A226685

Minimal element in the primitive 3x-k cycle with the conjectured record-breaking maximal element A226683(n).

Minimal element in the primitive 3x-k cycle with the conjectured record-breaking maximal element A226683(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007837 | OEIS-A226686

Conjectured record-breaking lengths, for ascending positive integers k, of primitive cycles of positive integers under iteration by the Collatz-like 3x-k function.

Conjectured record-breaking lengths, for ascending positive integers k, of primitive cycles of positive integers under iteration by the Collatz-like 3x-k function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007838 | OEIS-A226687

Integer k associated with the conjectured record-breaking length A226686(n) of primitive Collatz-like 3x-k cycles.

Integer k associated with the conjectured record-breaking length A226686(n) of primitive Collatz-like 3x-k cycles.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007839 | OEIS-A226688

Smallest positive integer in the primitive Collatz-like 3x-k cycle(s) with the conjectured record-breaking length A226686(n).

Smallest positive integer in the primitive Collatz-like 3x-k cycle(s) with the conjectured record-breaking length A226686(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey H. Morley_, Jun 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007840 | OEIS-A226727

Places n where A225867(n)=5.

The proportion of prime terms has a tendency to increase. For example, 4 from the first 10 terms; 10 from the first 20 terms; 16 from the first 30 terms; 24 from the first 40 terms; 40 from the first 60 terms, 88 from the first 110 terms. We conjecture that this proportion asymptotically tends to 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jun 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007841 | OEIS-A226729

G.f.: 1 / G(0), where G(k) = 1 - q^(k+1) / (1 - q^(k+1)/G(k+2)).

Conjectures from _John Tyler Rascoe_, Nov 04 2023: (Start) Conjecture: a(n) = Sum_{i=0..floor((n-sqrt(2*n-1))/2)} A129183(n-(2*i),n-i). - _John Tyler Rascoe_, Nov 04 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Jun 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007842 | OEIS-A226738

Row 3 of array in A226513.

Conjectural g.f. as a continued fraction of Stieltjes type: 1/(1 - 4*x/(1 - 2*x/(1 - 5*x/(1 - 4*x/(1 - 6*x/(1 - 6*x/(1 - (n+3)*x/(1 - 2*n*x/(1 - ...)))))))). - _Peter Bala_, Aug 27 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Jun 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007843 | OEIS-A226739

Row 4 of array in A226513.

Conjectural g.f. as a continued fraction of Stieltjes type: 1/(1 - 5*x/(1 - 2*x/(1 - 6*x/(1 - 4*x/(1 - 7*x/(1 - 6*x/(1 - (n+4)*x/(1 - 2*n*x/(1 - ...))))))))). - _Peter Bala_, Aug 27 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Jun 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007844 | OEIS-A226740

Row 5 of array in A226513.

Conjectural g.f. as a continued fraction of Stieltjes type: 1/(1 - 6*x/(1 - 2*x/(1 - 7*x/(1 - 4*x/(1 - 8*x/(1 - 6*x/(1 - (n+5)*x/(1 - 2*n*x/(1 - ...))))))))). - _Peter Bala_, Aug 27 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Jun 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007845 | OEIS-A226751

G.f.: 1 / (1 + 6*x*G(x) - 7*x*G(x)^2), where G(x) = 1 + x*G(x)^3 is the g.f. of A001764.

Conjecture: 18*n*(2*n-1)*(55*n-76)*a(n) +(-11605*n^3+28521*n^2-20870*n+4536)*a(n-1) -24*(55*n-21)*(3*n-4)*(3*n-2)*a(n-2)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007846 | OEIS-A226770

Let n+1 have proper divisors 1 < d_1,...,d_k < n+1; consider all proper divisors of n+d_1,...,n+d_k which did not appear earlier. Let them be d_{1,1}, d_{1,2},..., d_{k,1}, d_{k,2},..., d_{k,t}; then consider proper divisors of n+d_{1,1},...,n+d_{k,t} which did not appear earlier, repeat until no new divisor is introduced. a(n) is the total number of different divisors obtained.

a(n) = 0, iff n = p - 1, where p is prime; we conjecture that a(p) = p - 2 and, more general, for odd prime p and k>=1, a(p^k) = p^k - p^(k-1) - 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jun 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007847 | OEIS-A226806

Numbers of the form 2^j + 4^k, for j and k >= 0.

Conjecture: Any integer n > 1 not equal to 4 can be written as a sum of distinct terms of the current sequence with no summand dividing another. - _Zhi-Wei Sun_, May 01 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jun 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007848 | OEIS-A226807

Numbers of the form 3^j + 4^k, for j and k >= 0.

Conjecture: Each integer n > 8 can be written as a sum of finitely many numbers of the form 3^a + 4^b (a,b >= 0) with no one dividing another. This has been verified for all n <= 1500. - _Zhi-Wei Sun_, Apr 18 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jun 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007849 | OEIS-A226808

Numbers of the form 2^j + 5^k, for j and k >= 0.

Conjecture: Each integer n > 4 can be written as a_1 + ... + a_k, where a_1,...,a_k are numbers of the form 2^a + 5^b (a,b>=0) (i.e., terms of the current sequence) with no one dividing another. This has been verified for n = 5..1200. - _Zhi-Wei Sun_, Apr 14 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jun 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007850 | OEIS-A226812

Numbers of the form 3^j + 6^k, for j and k >= 0.

Conjecture: Any positive integer not among 1, 3, 5, 6, 8, 12, 27 can be written as a sum of distinct terms of the current sequence with no summand dividing another. - _Zhi-Wei Sun_, May 01 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jun 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007851 | OEIS-A226838

E.g.f.: exp(Sum_{n>=1} x^(n^2) / n^n).

a(n) == 1 (mod 6) (conjecture - valid up to n=1024; if true for n>=0, why?).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007852 | OEIS-A226839

E.g.f.: exp(Sum_{n>=1} x^(n*(n+1)/2) / n!).

a(n) == 1 (mod 3) (conjecture - valid up to n=1024; if true for n>=0, why?).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007853 | OEIS-A226857

Numbers that are both the sum of two Fibonacci numbers and the product of two Fibonacci numbers.

Conjecture: a(n) = a(n-3)+a(n-6) for n>12. - _Colin Barker_, Nov 09 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alonso del Arte_, Jun 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007854 | OEIS-A226866

Number of n X 2 (-1,0,1) arrays of determinants of 2 X 2 subblocks of some (n+1) X 3 binary array with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007855 | OEIS-A226867

Number of n X 3 (-1,0,1) arrays of determinants of 2 X 2 subblocks of some (n+1) X 4 binary array with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007856 | OEIS-A226875

Number of n-length words w over a 5-ary alphabet {a1,a2,...,a5} such that #(w,a1) >= #(w,a2) >= ... >= #(w,a5) >= 0, where #(w,x) counts the letters x in word w.

Conjecture: a(n) ~ 5^n/5!. - _Vaclav Kotesovec_, Mar 07 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jun 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007857 | OEIS-A226890

E.g.f.: exp(Sum_{n>=1} sigma(n,n) * x^(n^2) / n^n).

a(n) == 1 (mod 30) (conjecture - valid up to n=4000; if true for n>=0, why?).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007858 | OEIS-A226896

Position of log n in the joint ranking of harmonic numbers H(k) and {log k}, for k >= 1; complement of A226894.

If, in the definition, log k is replaced by g + log k, where g is the Euler-Mascheroni constant, then the position of log n is 2n-1, which leads to a conjecture: limit[1/(H(n) - g - log n) - 2n] = 1/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007859 | OEIS-A226898

Hooley's Delta function: maximum number of divisors of n in [u, e*u] for all u. (Here e is Euler's number 2.718... = A001113.)

Conjecture: For n > 4, a(n) <= d(n)/2 where d(n) is A000005. - _Charles R Greathouse IV_, Jan 24 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Jun 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007860 | OEIS-A226917

Number of non-sphere-homeomorphic crossing optimal 2-page book drawings of complete graph K_{2n+1}.

Conjecture: a(n) = 4*a(n-1) - 2*a(n-2) - 8*a(n-3) + 8*a(n-4); g.f.: -x^2*(5*x^3-5*x^2+1) / ((2*x-1)^2*(2*x^2-1)). - _Colin Barker_, May 08 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007861 | OEIS-A226919

Asymptotic number of completely symmetric polynomials of degree n up to momentum conservation in the limit as the number of particles increases.

The values for n >= 6 are only conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007862 | OEIS-A226955

Number of representations of n! as a sum of 3 positive cubes.

Conjecture: For any t >= 0, there are only finitely many values of n such that a(n) = t. - _Altug Alkan_, Aug 16 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Jun 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007863 | OEIS-A226978

Number of ways to cut an n X n square into squares with integer sides, reduced for symmetry, where the orbits under the symmetry group of the square, D4, have 1 element.

Conjecture: For n>3 the number of dissections is a multiple of 4. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christopher Hunt Gribble_, Jun 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007864 | OEIS-A226988

Number of n X 2 0..4 arrays of sums of 2 X 2 subblocks of some (n+1) X 3 binary array with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007865 | OEIS-A227010

Places n where A225867(n) = 4.

1 from the first 20 terms; 6 from the first 40 terms; 16 from the first 60 terms; 34 from the first 80 terms; 53 from the first 100 terms; 90 from the first 140 terms. We conjecture that this proportion asymptotically tends to 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jun 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007866 | OEIS-A227012

a(n) = floor(M(g(n-1)+1, ..., g(n))), where M = harmonic mean and g(n) = n^3.

Conjecture: if f(k) = u/(v*k + w), where u,v,w are integers, and g(n) is a polynomial, then the sequence with n-th term m(n) = floor(M(n)) is linearly recurrent. The conjecture extends to these cases, in which a,b,c,d are integers and a > 0: a(n+2) = (1/8)*(27 - (-1)^n - 2*cos(n*Pi/2) + 2*sin(n*Pi/2) + 2*n*(4*n^2 + 18 n + 27)) for n >= 1 (conjectured). a(n) = 3*a(n-1) - 3*a(n-2) + a(n-3) + a(n-4) - 3*a(n-5) + 3*a(n-6) - a(n-7) for n >= 10 (conjectured). G.f.: x*(1 + x + 7*x^2 + 6*x^3 + 5*x^4 + 5*x^5 - 2*x^7 + x^8)/(((x - 1)^4)*(1 + x

+ x^2 + x^3)) (conjectured). a(n) = (2*n^3 - 3*n^2 + n + 2)/2 + floor(max(0, n - 3)/4) (conjectured). - _Franck Maminirina Ramaharo_, Apr 12 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007867 | OEIS-A227013

a(n) = floor(M(g(n-1)+1,..,g(n))), where M is the harmonic mean and g(n) = n^4.

a(n) = 47/9 + 14*n + (41*n^2)/3 + 6*n^3 + n^4 - (2/9)Cos(2*n*pi/3) (conjectured). a(n) = 4*a(n-1) - 6*a(n-2) + 5*a(n-3) - 5*a(n-4) + 6*a(n-5) - 4*a(n-6) + a(n-7) for n > 2 (conjectured). G.f.: (-1 - 2*x - 22*x^2 - 23*x^3 - 20*x^4 - 4*x^5 + 2*x^6 - 3*x^7 + x^8)/((-1 + x)^5 (1 + x + x^2)) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007868 | OEIS-A227014

a(n) = floor(M(g(n-1)+1,..,g(n))), where M = harmonic mean and g(n) = n^5.

See A227012. It is conjectured that A227014 is a linear recurrence sequence with signature (5,-10,10,-5,1,...Z...,1,-5,-10,-10,-1,0,0), where ...Z... represents a string of 138 zeros; has been confirmed for a(1), a(2),..., a(150000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007869 | OEIS-A227015

a(n) = floor(M(g(n-1)+1, ..., g(n))), where M = harmonic mean and g(n) = n^3 + n^2 + n + 1.

a(n) = 3*a(n-1) - 3*a(n-2) + a(n-3) + a(n-4) - 3*a(n-5) + 3*a(n-6) - a(n-7) for n > 2 (conjectured). G.f.: x*(2 + 2*x + 8*x^2 + 4*x^3 + 5*x^4 + 4*x^5 - 2*x^6 + 3*x^7 - 3*x^8 + x^9)/((x - 1)^4*(1 + x + x^2 + x^3)) (conjectured). a(n) = (1/2)*((-1)^(n - 1)! + 2*n^3 - n^2 + n + 3 + 2*floor(max(0, n - 4)/4)) (conjectured). E.g.f.: (1/24)*exp(-x)*(exp(x)*(6*sin(x) + 6*cos(x) + 4*x^3 - 24) + exp(2*x)*(24*x^3 + 60*x^2 + 30*x + 15) + 3)

(conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007870 | OEIS-A227016

Floor(M(g(n-1)+1,..,g(n))), where M = harmonic mean and g(n) = n(n + 1)(n + 2)/6.

a(n) + 3*a(n-1) - 3*a(n-2) + a(n-3) + a(n-8) - 3*a(n-9) + 3*a(n-10) - a(n-11) for n > 1 (conjectured). G.f.: (1 - x + 4*x^2 - 2*x^3 + 4*x^4 - x^5 + x^6 + 2*x^7 - x^8 + 3*x^9 - 3*x^10 + x^11)/((x - 1)^4 (1+x) (1+x^2) (1+x^4)) (conjectured). (G.f. found by _Peter J. C. Moses_, Jul 01 2013)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007871 | OEIS-A227017

Floor(M(g(n-1)+1,..,g(n))), where M = harmonic mean and g(n) = n(3n-1)/2 = A000326(n).

a(n) = (1/4)*(1 - (-1)^n + 4*n + 6*n^2) (conjectured). a(n) = 2*a(n-1) - 2*a(n-3) + a(n-4) for n > 1 (conjectured). G.f.: (-1 - x - 2*x^2 - 3*x^3 + x^4)/((-1 + x)^3 (1 + x)). (conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007872 | OEIS-A227018

a(n) = floor(M(g(n-1)+1,..,g(n))), where M = harmonic mean and g(n) = n*(n + 1)*(n + 2)*(n + 3)/24.

Conjectured g.f.: (-1 + x - 3 x^2 - 2 x^3 + 2 x^4 - 2 x^5 - 3 x^6 + 2 x^8 - 5 x^9 - x^12 - x^13 - 4 x^14 + 4 x^15 - 4 x^16 - 2 x^17 + 2 x^18 - 2 x^19 - 4 x^20 + 4 x^21 - 4 x^22 - x^23 - x^24 + x^25 - 4 x^26 + x^27 - x^28 - 3 x^29 + 3 x^30 - 5 x^31 - x^34 - 2 x^35 + x^36 - 3 x^37 + 2 x^38 + 2 x^39 - 2 x^40 - 3 x^41 + 6 x^42 - 3 x^43 - 3 x^44 + 6 x^45 - 4 x^46 + x^47)/((-1 + x)^5 (1 + x) (1 + x^2) (1 - x + x^2) (1 + x + x^2) (1 - x^2 + x^4) (1 - x^3 + x^6) (1 + x^3 + x^6) (1 - x^6 + x^12)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007873 | OEIS-A227021

Number of n X 2 (0,1,2) arrays of permanents of 2 X 2 subblocks of some (n+1) X 3 binary array with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007874 | OEIS-A227036

Expansion of 2*(1+x^2)/((1-x)*(1-x-2*x^3)).

Conjecture: The perimeter of the n-th iteration of the Harter-Heighway dragon is a(n) segments or a(n)/2^(n/2) base units.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roland Kneer_, Jun 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007875 | OEIS-A227051

E.g.f.: 1/(cos(x) - sin(x)*sinh(x)), omitting the zero-valued coefficients of odd powers of x.

a(n) == 3 (mod 10) for n>0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007876 | OEIS-A227054

a(n) = least triangular number t > 0 such that n*t is a triangular number, or 0 if no such t exists.

Indices of conjectured 0's: 4, 9, 16, 25, 49, 64, 81, 121, 144, 169, 225, 256, 289, 361, 400, 441, 529, 576, 625, 729, ... These are squares of 2, 3, 4, 5, 7, 8, 9, 11, 12, 13, 15, 16, 17, 19, 20, 21, 23, 24, 25, 27.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Jun 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007877 | OEIS-A227056

Number of n X 2 -2..2 arrays of 2 X 2 subblock diagonal sums minus antidiagonal sums for some (n+1) X 3 binary array with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007878 | OEIS-A227081

Row sums of A124576.

Conjecture: 3*n*a(n) +2*(-13*n+9)*a(n-1) +4*(13*n-21)*a(n-2) +24*(-n+2)*a(n-3)=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jun 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007879 | OEIS-A227083

Number of ways to write n as a + b/2 with a and b terms of the sequence A008407.

Conjecture: We have a(n) > 0 for all n > 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007880 | OEIS-A227085

Number of n X 2 binary arrays indicating whether each 2 X 2 subblock of a larger binary array has lexicographically increasing rows and columns, for some larger (n+1) X 3 binary array with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007881 | OEIS-A227086

Number of n X 3 binary arrays indicating whether each 2 X 2 subblock of a larger binary array has lexicographically increasing rows and columns, for some larger (n+1) X 4 binary array with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007882 | OEIS-A227099

Number of n X 2 0,1 arrays indicating 2 X 2 subblocks of some larger (n+1) X 3 binary array having a sum of two, with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007883 | OEIS-A227121

Number of n X 2 0,1 arrays indicating 2 X 2 subblocks of some larger (n+1) X 3 binary array having a sum of zero, with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007884 | OEIS-A227122

Number of n X 3 0,1 arrays indicating 2 X 2 subblocks of some larger (n+1) X 4 binary array having a sum of zero, with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007885 | OEIS-A227134

Partitions with parts repeated at most twice and repetition only allowed if first part has an odd index (first index = 1).

Conjecture: A227134(n) + A227135(n) = A182372(n) for n >= 0, see comment in A182372.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Jul 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007886 | OEIS-A227135

Partitions with parts repeated at most twice and repetition only allowed if first part has an even index (first index = 1).

Conjecture: A227134(n) + A227135(n) = A182372(n) for n>=0, see comment in A182372.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Jul 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007887 | OEIS-A227145

Numbers satisfying an infinite nested recurrence relation.

Conjecture: a(F_n) = F_{n-2} for n>1, where F_n is the n-th Fibonacci number. Conjecture: a(n) ~ n*(3-sqrt(5))/2. -_Jeffrey Shallit_, Oct 12 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank Ruskey_, Jul 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007888 | OEIS-A227156

Number of ways to write n as a sum of a square and half of a term of the sequence A008407.

Conjecture: We have a(n) > 0 except for n = 23. We also conjecture that any positive integer can be written as a sum of a triangular number and half of a term of A008407, and each integer n > 4 can be written as x + y (x>0, y>0) with x*y a term of A008407.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007889 | OEIS-A227162

Number of n X 3 0,1 arrays indicating 2 X 2 subblocks of some larger (n+1) X 4 binary array having a sum of one or less, with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007890 | OEIS-A227183

a(n) is the sum of parts of the unique unordered partition encoded in the run lengths of the binary expansion of n; row sums of A227739 for n >= 1.

Each n occurs A000041(n) times in total and occurs for the first time at A227368(n) and for the last time at position A000225(n). See further comments and conjectures at A227368 and A227370.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jul 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007891 | OEIS-A227202

Least prime, q, greater than the previous prime, p, which is a primitive root of p; beginning with 2.

Conjecture: a(n) < Prime[n*E]. Conjecture: All primes will join this sequence eventually.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Sep 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007892 | OEIS-A227244

Smallest prime factor of (2^(2^n) + 1)*2^(2^(2^n)) - 1.

Proof that the 1st conjecture of Chai Wah Wu is right. 5) a(0) = a(5) = 11 and the conjecture is proved. a(n) >= 11 for n >= 0 (conjectured). a(4n+5) = 11 for n >= 0 (conjectured). See the proofs of these 2 conjectures in the comments. (End) Conjecture 1: a(20*n+16) = 101 for n >= 0. Conjecture 2: a(220*n+8) = 47 for n >= 0. Conjecture 3: a(72*n+32) = 149 for n >= 0 and a(72*n+32) is not covered by Conjectures 1 and 2. Conjecture 4: a(168*n+43) = 59 if n>=0 and n mod 55 <> 26. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Oct 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007893 | OEIS-A227249

Number of consecutive composites beginning with the first, to be added to obtain a power.

All powers are squares with the exception of 3^3 for a(2) and 6^9 for a(6). I conjecture these are the only nonsquare powers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robin Garcia_, Jul 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007894 | OEIS-A227250

Number of binary labeled trees with two-colored vertices that have n leaves and avoid the easiest to avoid 6-pattern set.

Conjecture: E.g.f. (for offset 0) satisfies A'(x) = 1 + A(x)^3, with A(0)=1. The next terms are 1203498, 24163110, 549811962, 13982486166, 393026414922, ... - _Vaclav Kotesovec_, Jun 15 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Dotsenko_, Jul 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007895 | OEIS-A227252

Number of n X 2 binary arrays indicating whether each 2 X 2 subblock of a larger binary array has lexicographically nondecreasing rows and columns, for some larger (n+1) X 3 binary array with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007896 | OEIS-A227259

Number of n X 2 0,1 arrays indicating 2 X 2 subblocks of some larger (n+1) X 3 binary array having a sum of two or less, with rows and columns of the latter in lexicographically nondecreasing order.

Conjecture: a(n) is also the number of length n strings of {0,1,2} with digit sum <= 4 (holds if the empirical formula holds). - _Daniel T. Martin_, May 24 2023 G.f.: x*(3 - 6*x + 8*x^2 - 5*x^3 + x^4) / (1 - x)^5. (Conjecture from _Colin Barker_, Sep 07 2018)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007897 | OEIS-A227265

Number of n X 2 0,1 arrays indicating 2 X 2 subblocks of some larger (n+1) X 3 binary array having a sum of three or less, with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007898 | OEIS-A227266

Number of n X 3 0,1 arrays indicating 2 X 2 subblocks of some larger (n+1) X 4 binary array having a sum of three or less, with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007899 | OEIS-A227281

First primes of arithmetic progressions of 5 primes each with the common difference 30.

The minimal possible difference in an AP-k is conjectured to be k# for all k > 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sameen Ahmed Khan_, Jul 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007900 | OEIS-A227282

First primes of arithmetic progressions of 7 primes each with the common difference 210.

The minimal possible difference in an AP-k is conjectured to be k# for all k > 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sameen Ahmed Khan_, Jul 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007901 | OEIS-A227283

First primes of arithmetic progressions of 8 primes each with the common difference 210.

The minimal possible difference in an AP-k is conjectured to be k# for all k > 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sameen Ahmed Khan_, Jul 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007902 | OEIS-A227284

First primes of arithmetic progressions of 9 primes each with the common difference 210.

The minimal possible difference in an AP-k is conjectured to be k# for all k > 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sameen Ahmed Khan_, Jul 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007903 | OEIS-A227285

First primes of arithmetic progressions of 11 primes each with the common difference 2310.

The minimal possible difference in an AP-k is conjectured to be k# for all k > 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sameen Ahmed Khan_, Jul 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007904 | OEIS-A227286

First primes of arithmetic progressions of 13 primes each with the common difference 30030.

The minimal possible difference in an arithmetic progression of k primes is conjectured to be k# = A034386(k) for all k > 7. 13# = 30030.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sameen Ahmed Khan_, Jul 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007905 | OEIS-A227329

Number of n X 2 0,1 arrays indicating 2 X 2 subblocks of some larger (n+1) X 3 binary array having two adjacent 1's and two adjacent 0's, with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007906 | OEIS-A227345

Triangle read by rows, partitions into distinct parts by size of boundary.

Conjecture: there exists a partition (into distinct parts) of n with boundary size k if and only if 0 < k^2 * 3/4 <= n. - _Patrick Devlin_, Jul 13 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Jul 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007907 | OEIS-A227359

Natural numbers that are not of the form (k +- sum of binary digits of k) for any k.

The numbers of terms not exceeding 10^k, for k = 1, 2, ..., are 1, 9, 127, 1362, 12921, 128429, 1261747, 12554142, 125697648, 1257065977, ... Conjecture: This sequence has an asymptotic density (1/2) * A242403 = 0.126330... . - _Amiram Eldar_, Oct 02 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andres M. Torres_, Jul 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007908 | OEIS-A227368

a(n) = Index k where A227183(k) for the first time gets value n; the runlength binary code for minimally runlength-encoded unordered partition of size n.

A000267(n) seems to give the binary width of a(n+1). Compare to the conjecture given at A227370. a(n) = A227369(A227370(n)) [See comments and conjecture at A227370]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jul 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007909 | OEIS-A227369

Positions where A227183 obtains new distinct values for the first time.

a(n) = A227370(A227368(n)). [See comments and conjectures at A227370]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jul 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007910 | OEIS-A227370

Permutation which maps between A227368 and A227369.

Conjecture 1: This is an involution (self-inverse permutation) of nonnegative integers. (Which would imply that both formulas given in A227368 and A227369 involving A227370 are valid). Conjecture 2: (which would automatically imply the conjecture 1): the only transpositions (used to compose the permutation) are of adjacent terms 2k-1 and 2k, where A061887 gives the values of k. This is true at least for the first 35 transpositions (up to k=60).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jul 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007911 | OEIS-A227375

G.f.: 1/(1 - x*(1-x^6)/(1 - x^2*(1-x^7)/(1 - x^3*(1-x^8)/(1 - x^4*(1-x^9)/(1 - x^5*(1-x^10)/(1 - ...))))))), a continued fraction.

Conjecture: G.f. -(x^4+x-1)*(x^5+x^4+x^3-x-1) / ((x-1)*(x^4+x^3+x^2+x+1)*(x^7+x^6+x^5-x^3-x^2-x+1)). - _R. J. Mathar_, Jul 17 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007912 | OEIS-A227382

Number of n X 3 0,1 arrays indicating 2 X 2 subblocks of some larger (n+1) X 4 binary array having a sum of one, with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007913 | OEIS-A227387

Numbers n such that sigma(n)*7 < sigma(n-1) + sigma(n+1).

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Jul 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007914 | OEIS-A227399

Number of permutations i_1, ..., i_n of 1, ..., n with i_1 = 1 and i_n = n such that i_1+2*i_2, i_2+2*i_3, ..., i_{n-1}+2*i_n, i_n+2*i_1 are pairwise distinct modulo n.

Zhi-Wei Sun also made the following general conjecture: Recall that a conjecture of Snevily proved by Arsovski states that for any two n-subsets A and B of an additive abelian group of odd order there is a numbering a_1,...,a_n of all the elements of A and a numbering b_1, ..., b_n of all the elements of B such that the n sums a_1+b_1, ..., a_n+b_n are pairwise distinct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007915 | OEIS-A227405

Order of the symmetry group of the densest possible packing of N circles in a larger circle. (If there are different patterns with the same density pick the largest value of the symmetry group order.)

a(14) is unknown but conjectured to be 2. The conjectured sequence is 4, 6, 8, 10, 12, 12, 14, 16, 2, 2, 6, 2, 2, 10, 2, 2, 12, 12, 2...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Keenan Pepper_, Sep 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007916 | OEIS-A227419

Least k such that the sum of the semiprime divisors equals n times the sum of the prime divisors, or 0 if no such k exists.

Conjecture: a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jul 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007917 | OEIS-A227433

Number of pairs of 2 X 2 matrices over Z/nZ that do not commute.

Conjecturally, a(p) = p^3(p^3-1)(p^2-1), for p prime. - _Giovanni Resta_, Jul 13 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Jul 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007918 | OEIS-A227437

Longest checkmate in king and 3 knights versus king endgame on an n X n chessboard.

With 16 GB of memory it was possible to explore this ending on boards up to 15 X 15. On these boards the ending is a general win. I think that from some greater n this ending is drawn (and the sequence is finite), but this is only conjecture. - _Vaclav Kotesovec_, Jul 19 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jul 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007919 | OEIS-A227456

Number of permutations i_0, i_1, ..., i_n of 0, 1, ..., n with i_0 = 0 and i_n = 1 such that all the n+1 numbers i_0^2+i_1, i_1^2+i_2, ..., i_{n-1}^2+i_n, i_n^2+i_0 are of the form (p+1)/4 with p a prime congruent to 3 modulo 4.

Conjecture: a(n) > 0 for all n > 0. Similarly, for any positive integer n, there is a circular permutation i_0, i_1, ..., i_n of 0, 1, ..., n such that all the n+1 numbers i_0^2+i_1, i_1^2+i_2, ..., i_{n-1}^2+i_n, i_n^2+i_0 are of the form (p-1)/4 with p a prime congruent to 1 modulo 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007920 | OEIS-A227463

E.g.f. equals the series reversion of arcsinh(x) / exp(x).

Conjectural 2-adic values: v_2(a(8*n+1)) = 4*n; v_2(a(8*n+3)) = 4*n+1; v_2(a(8*n+5)) = 4*n+2; v_2(a(8*n+4)) = 8*n+4. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007921 | OEIS-A227500

a(0)=a(1)=0; for n>1, a(n) = numerator(r(n)), where r(n) = r(n-1)+r(n-2)+A027641(n-2)/A027642(n-2) and r(0)=r(1)=a(0).

Conjecture: (b(n)-a(n))/c(n) = 0, 0, 0, 1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, ..., that is two 0 followed by A000045. This conjecture is confirmed up to 100 terms. [_Jean-François Alcover_, Jul 19 2013]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Jul 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007922 | OEIS-A227512

Floor(-1/n + 1/log((2n+1)/(2n-1))).

a(n) = 12*n^3 - floor(9*n/5) - 1. (conjectured, based on computations by _Peter J. C. Moses_, Jul 14 2013) a(n) = 3*a(n-1) - 3*a(n-2) + a(n-3) + a(n-5) - 3*a(n-6) + 3*a(n-7) - a(n-8) (conjectured; verified up to n = 100000). G.f.: (10 + 62 x + 72 x^2 + 72 x^3 + 72 x^4 + 63 x^5 + 8 x^6 + x^7)/((-1 + x)^4 (1 + x + x^2 + x^3 + x^4)) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007923 | OEIS-A227513

Round(-1/n + 1/log((2n+1)/(2n-1))).

a(n) = 12*n^3 - round(9*n/5) (conjectured, based on computations by _Peter J. C. Moses_, Jul 14 2013). a(n) = 3*a(n-1) - 3*a(n-2) + a(n-3) + a(n-5) - 3*a(n-6) + 3*a(n-7) - a(n-8) (conjectured; verified up to n = 100000). G.f.: x*(10 + 62*x + 73*x^2 + 70*x^3 + 73*x^4 + 62*x^5 + 10*x^6)/((-1 + x)^4*(1 + x + x^2 + x^3 + x^4)) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007924 | OEIS-A227533

Smallest e > 1 such that (2n)^e is a totient, or 0 if no such e exists.

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Jul 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007925 | OEIS-A227534

Even numbers n such that the least e with n^e a totient is a new record.

Probably all terms beyond the first are even semiprimes: conjecturally this is a subsequence of A001747.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Jul 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007926 | OEIS-A227543

Triangle defined by g.f. A(x,q) such that: A(x,q) = 1 + x*A(q*x,q)*A(x,q), as read by terms k=0..n*(n-1)/2 in rows n>=0.

Conjecture: b(n) = P(n, n) where b(n) is an integer sequence with g.f. B(x) = 1/(1 - f(0)*x/(1 - f(1)*x/(1 - f(2)*x/(1 - f(3)*x/(1 - f(4)*x/(1 -...)))))), P(n, k) = P(n-1, k) + f(n-k)*P(n, k-1) for 0 < k <= n with P(n, k) = 0 for k > n, P(n, 0) = 1 for n >= 0 and where f(n) is an arbitrary function. In fact for this sequence we have f(n) = q^n. - _Mikhail Kurkov_, Sep 26 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007927 | OEIS-A227554

Number of n X 2 0,1 arrays indicating 2 X 2 subblocks of some larger (n+1) X 3 binary array having nonzero determinant, with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007928 | OEIS-A227555

Number of n X 3 0,1 arrays indicating 2 X 2 subblocks of some larger (n+1) X 4 binary array having nonzero determinant, with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007929 | OEIS-A227576

Numbers k such that F(3*k)/(2*F(k)) is prime, where F(m) is the m-th Fibonacci number.

All terms are primes. Conjecture: this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Jul 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007930 | OEIS-A227578

Number A(n,k) of lattice paths from {n}^k to {0}^k using steps that decrement one component such that for each point (p_1,p_2,...,p_k) we have p_1<=p_2<=...<=p_k; square array A(n,k), n>=0, k>=0, read by antidiagonals.

Conjecture: column k is asymptotic to c(k) * (k+1)^(k*n)/n^((k^2-1)/2), where c(k) is a constant dependent only on k. - _Vaclav Kotesovec_, Jul 21 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007931 | OEIS-A227589

Maximum label within a minimal labeling of n identical 4-sided dice yielding the most possible sums.

The conjectured formula is certainly an upper bound for a(n), using the labeling (1, 2, (n^2+n+4)/2, (n^2+3n+5-(-1)^n)/2). Consider the sets S(y) = {possible sums if exactly y dice roll >2}; these sets have disjoint ranges for y <= n/2, and are interleaved without overlap for y > n/2. Computer search shows this labeling is optimal for n=2..53 at least. - _Martin Fuller_, Jun 08 2026 Conjecture: a(n) = (n^2 + 3n + 5 - (-1)^n)/2 for n > 1. Conjecture: a(n) = 2*a(n-1)-2*a(n-3)+a(n-4) for n>5. G.f.: (x^5-x^4-x^2+2*x+1) / ((x-1)^3*(x+1)). - _Colin Barker_, Aug 01 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jens Voß_, Jul 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007932 | OEIS-A227597

Number of lattice paths from {n}^6 to {0}^6 using steps that decrement one component such that for each point (p_1,p_2,...,p_6) we have p_1<=p_2<=...<=p_6.

Conjecture: a(n) ~ 5 * 7^(6*n+29) / (2^58 * 3^8 * Pi^(5/2) * n^(35/2)). - _Vaclav Kotesovec_, Nov 20 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007933 | OEIS-A227598

Number of lattice paths from {n}^7 to {0}^7 using steps that decrement one component such that for each point (p_1,p_2,...,p_7) we have p_1<=p_2<=...<=p_7.

Conjecture: a(n) ~ 25 * sqrt(7) * 8^(7*n + 44) / (7^17 * 3^43 * Pi^3 * n^24). - _Vaclav Kotesovec_, Nov 21 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007934 | OEIS-A227599

Number of lattice paths from {n}^8 to {0}^8 using steps that decrement one component such that for each point (p_1,p_2,...,p_8) we have p_1<=p_2<=...<=p_8.

Conjecture: a(n) ~ 42 * sqrt(5) * 9^(8*n + 58) / (8^20 * 10^29 * n^(63/2) * Pi^(7/2)). - _Vaclav Kotesovec_, Nov 26 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007935 | OEIS-A227603

Number of lattice paths from {6}^n to {0}^n using steps that decrement one component such that for each point (p_1,p_2,...,p_n) we have p_1<=p_2<=...<=p_n.

Conjecture: a(n) ~ 2^(5/2) * 6^(6*n + 67/2) / (5^29 * Pi^(5/2) * n^(35/2)). - _Vaclav Kotesovec_, Nov 21 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007936 | OEIS-A227604

Number of lattice paths from {7}^n to {0}^n using steps that decrement one component such that for each point (p_1,p_2,...,p_n) we have p_1<=p_2<=...<=p_n.

Conjecture: a(n) ~ 2^4 * 5^2 * 7^(7*n + 85/2) / (6^37 * Pi^3 * n^24). - _Vaclav Kotesovec_, Nov 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007937 | OEIS-A227609

Determinant of the (p_n-1)/2 X (p_n-1)/2 matrix with (i,j)-entry being the Legendre symbol((i^2+j^2)/p_n), where p_n is the n-th prime.

Conjecture: p_n never divides a(n), and moreover -a(n) is a quadratic residue mod p_n. Zhi-Wei Sun also made the following conjecture: These were proved in version 9 of arXiv:1308.2900 (2018). In addition, the author has the following new conjecture. Conjecture: For any prime p == 3 (mod 4), the number -S(1,p) is a positive square divisible by 2^((p-3)/2), i.e., -S(1,p) = (2^((p-3)/4)*m)^2 for some positive integer m. - _Zhi-Wei Sun_, Sep 09 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007938 | OEIS-A227627

Numbers k such that F(5*k)/(5*F(k)) is prime, where F(m) is the m-th Fibonacci number.

Except for k = 25, all other terms k are primes. Conjecture: this sequence is infinite. - _Thomas Ordowski_, Jul 18 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jul 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007939 | OEIS-A227631

Array t(n,k): row n consists of the positive integers m for which the least splitter of H(m) and H(m+1) is n, where H denotes harmonic number.

Suppose that x < y. The least splitter of x and y is introduced here as the least positive integer d such that x <= c/d < y for some integer c; the number c/d is called the least splitting rational of x and y. Conjecture: every row of the array in A227631 is infinite, and every positive integer occurs exactly once. Let r be the limiting ratio of consecutive terms of row 1; is r = e?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007940 | OEIS-A227637

Number of n X 2 0,1 arrays indicating 2 X 2 subblocks of some larger (n+1) X 3 binary array having determinant equal to one, with rows and columns of the latter in nondecreasing lexicographic order.

Conjectures from _Colin Barker_, Sep 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007941 | OEIS-A227638

Number of n X 3 0,1 arrays indicating 2 X 2 subblocks of some larger (n+1) X 4 binary array having determinant equal to one, with rows and columns of the latter in nondecreasing lexicographic order.

Conjectures from _Colin Barker_, Sep 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007942 | OEIS-A227653

a(1) = least k such that 1/2 + 1/3 < H(k) - H(3); a(2) = least k such that H(a(1)) - H(3) < H(k) -H(a(1)), and for n > 2, a(n) = least k such that H(a(n-1)) - H(a(n-2)) > H(k) - H(a(n-1)), where H = harmonic number.

a(n) = A077849(n+1) (conjectured). a(n) = 3*a(n-1) - a(n-2) - a(n-4) (conjectured).G.f.: (8 - 3 x - x^2 - 3 x^3)/(1 - 3 x + x^2 + x^4) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007943 | OEIS-A227675

Number of n X 2 0,1 arrays indicating 2 X 2 subblocks of some larger (n+1) X 3 binary array having an odd sum, with rows and columns of the latter in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007944 | OEIS-A227698

a(n) is the number of ways to form the integer n as (p^2 + q^2 - 2)/24, where p and q are primes > 3 (excluding swaps of p and q).

A persistent tendency for more frequent large values of a(n) for n> 40000 is conjectured, with the likelihood that 8 is NOT the maximum value, and the possibility that ever larger values can always be found at higher n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard R. Forberg_, Sep 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007945 | OEIS-A227719

Floor(1/s(n)), where s(n) = n*log(1+1/n) - (2n-1)/(2n).

That s(n) > 0 for n >=1 follows from the chain 1 < log 2 < 3/4 < 2 log 3/2 < 5/6 < 3 log 4/3 < 7/8 < 4 log 5/4 < ... ; i.e., n*log((n+1)/n) - (2n-1)/(2n) > 0 and (2n+1)/(2n+2) - n*log((n+1)/n) > 0. For the first, closeness to 0 is indicated by A227719 and A227720, and for the second, by A227721 and a sequence which possibly equals A094159. Conjecture: the four sequences are linearly recurrent. a(n) = -2*a(n-1) - a(n-2) + a(n-4) - 2*a(n-5) + a(n-6) (conjectured). G.f.: (-5 - 6 x - 6 x^2 - 6 x^3 - 2 x^4 + x^5)/((-1 + x)^3 (1 + x + x^2 + x^3)) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007946 | OEIS-A227720

Round(1/s(n)), where s(n) = n*log(1+1/n) - (2n-1)/(2n).

That s(n) > 0 for n >=1 follows from the chain 1 < log 2 < 3/4 < 2 log 3/2 < 5/6 < 3 log 4/3 < 7/8 < 4 log 5/4 < ... ; i.e., n log((n+1)/n) - (2n-1)/(2n) > 0 and (2n+1)/(2n+2) - n log((n+1)/n) > 0. For the first, closeness to 0 is indicated by A227719 and A227720, and for the second, by A227721 and a sequence which possibly equals A094159. Conjecture: the four sequences are linearly recurrent. a(n) = -2*a(n-1) - a(n-2) + a(n-4) - 2*a(n-5) + a(n-6) (conjectured). G.f.: (-5 - 6 x - 7 x^2 - 5 x^3 - x^4)/((-1 + x)^3 (1 + x + x^2 + x^3)) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007947 | OEIS-A227721

Floor(1/s(n)), where s(n) = (2n+1)/(2n+2) - n*log((n+1)/n).

That s(n) > 0 for n >=1 follows from the chain 1 < log 2 < 3/4 < 2 log 3/2 < 5/6 < 3 log 4/3 < 7/8 < 4 log 5/4 < ... ; i.e., n log((n+1)/n) - (2n-1)/(2n) > 0 and (2n+1)/(2n+2) - n log((n+1)/n) > 0. For the first, closeness to 0 is indicated by A227719 and A227720, and for the second, by A227721 and a sequence which possibly equals A094159. Conjecture: the four sequences are linearly recurrent. a(n) = 2 + 9*n + 6*n^2 (conjectured). a(n) = 3*a(n-1) - 3*a(n-2) + a(n-3) (conjectured). G.f.: (-17 + 7 x - 2 x^2)/(-1 + x)^3 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007948 | OEIS-A227728

a(1) = greatest k such that H(k) - H(2) < 1/1 + 1/2; a(2) = greatest k such that H(k) - H(a(1)) < H(a(1)) - H(2); and for n>2, a(n) = greatest k such that H(k) - H(a(n-1)) < H(a(n-1)) - H(a(n-2)), where H = harmonic number.

a(n) = 5*a(n-1) - 4*a(n-2) + 2*a(n-3) - 2*a(n-4) + 2*a(n-5) - 2*a(n-6) + 2*a(n-7) - 2*a(n-8) + 2*a(n-9) - 2*a(n-10) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007949 | OEIS-A227761

a(n) is the maximal difference between successive parts in the minimally runlength-encoded unordered partition of n (A227368(n)).

After n=3, only composites may obtain value 0. (But not all of them do; see A227762.) The first nine n for which a(n)=2 are 7, 13, 23, 33, 47, 61, 79, 97, 119, of which all are primes except 33 and 119. Conjecture: these values are given by A227786.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jul 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007950 | OEIS-A227776

a(n) = 6*n^2 + 1.

Least splitter is defined for x < y at A227631 as the least positive integer d such that x <= c/d < y for some integer c; the number c/d is called the least splitting rational of x and y. Conjecture: a(n) is the least splitter of s(n) and s(n+1), where s(n) = n*sin(1/n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007951 | OEIS-A227777

Least splitter of n-th and (n+1)st partial sums of 1/0! + 1/1! + ... + 1/n! + ... = e.

Conjecture: a(n) <= n*sqrt(n!) for all n>0; see scatterplot under Links. - _Jon E. Schoenfield_, Jun 28 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007952 | OEIS-A227786

Take squares larger than 1, subtract 3 from even squares and 2 from odd squares.

Conjecture: from n>=2 onward, a(n) gives the positions of 2's in A227761.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jul 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007953 | OEIS-A227792

Expansion of (1 + 6*x + 17*x^2 - x^3 - 3*x^4)/(1 - 6*x^2 + x^4).

Also, values i where A067060(i)/i reaches a new maximum (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Sep 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007954 | OEIS-A227804

a(1) = greatest k such that H(k) - H(8) < H(8) - H(4); a(2) = greatest k such that H(k) - H(a(1)) < H(a(1)) - H(8), and for n > 2, a(n) = greatest k such that H(k) - H(a(n-1)) > H(a(n-1)) - H(a(n-2)), where H = harmonic number.

a(n) = 3*a(n-1) - 3*a(n-2) + 2*a(n-3) - a(n-4) (conjectured). G.f.: (15 - 18 x + 12 x^2 - 8 x^3)/(1 - 3 x + 3 x^2 - 2 x^3 + x^4) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007955 | OEIS-A227845

G.f.: Sum_{n>=0} x^n / (1-x)^(2*n+1) * [Sum_{k=0..n} C(n,k)^2*x^k]^2.

Conjecture: if p is a prime in A105874 then a(p-1) == 0 (mod p^2) (checked up to p = 24239 = A105874(1000)). - _Peter Bala_, Sep 19 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007956 | OEIS-A227849

a(n) = 2 * floor(3/14 * n^2) if n even, a(n) = 2 * round(3/14 * n^2) -1 if n odd.

The degrees of the sequence of the G_2 polynomials defined by Bruce Westbury is conjectured to be a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Oct 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007957 | OEIS-A227864

Smallest base in which n's digital reversal is prime, or 0 if no such base exists.

Conjecture 1: 0, 1 and 4 are the only values where there is no base in which a digital reversal makes a prime. Conjecture 2: n = 2 is the only prime p which must be represented in base p+1, i.e., trivially, as a single digit, in order for its reversal to be prime. Conjecture 3: With the restriction gcd(base,n) = 1, a(n) = 0 except for n = 2, 3 and 6k+-1, for positive integer k, i.e., terms of A038179.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carl R. White_, Nov 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007958 | OEIS-A227877

Number of ways to write n = x + y + z (x, y, z > 0) such that x*y and x*z are triangular numbers, and 6*y-1 and 6*z+1 are both prime.

Conjecture: a(n) > 0 for all n > 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007959 | OEIS-A227898

Number of primes p < n with p + 6 and n + (n - p)^2 both prime.

Conjecture: (i) a(n) > 0 for all n > 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007960 | OEIS-A227899

Number of primes p < n with 3*p - 4 and n^2 + (n - p)^2 both prime.

Conjecture: a(n) > 0 for all n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007961 | OEIS-A227917

Number of semi-increasing binary plane trees with n vertices.

Conjecture: a(n) = 2^(n + 1)*Integral_{x=0..oo} Pochhammer(x, n)/exp(2*x) dx. - _Velin Yanev_, Jan 11 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brad R. Jones_, Oct 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007962 | OEIS-A227918

Sum over all permutations beginning and ending with ascents, and without double ascents on n elements and each permutation contributes 2 to the power of the number of double descents.

Conjecture: a(n) -n*a(n-1) -a(n-2) +(n-2)*a(n-3) = 0. - _R. J. Mathar_, Jul 17 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Ehrenborg_, Oct 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007963 | OEIS-A227937

Partitions of n labeled elements into subsets of two or three elements.

For n coprime to both 2 and 3 and for k >= 0, we conjecture that n divides a(n+k). - _Peter Bala_, Nov 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Eppstein_, Oct 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007964 | OEIS-A227968

Determinant of the (p_n+1)/2 X (p_n+1)/2 matrix with (i,j)-entry (i,j=0,...,(p_n-1)/2) equal to the Legendre symbol((i^2+j^2)/p_n), where p_n is the n-th prime.

Conjecture: 2*a(n) is always a quadratic residue mod p_n. Zhi-Wei Sun also made the following general conjecture: The general conjecture mentioned above was proved in a recent preprint of Zhi-Wei Sun.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007965 | OEIS-A227971

Determinant of the (p_n+1)/2 X (p_n+1)/2 matrix with (i,j)-entry (i,j=0,...,(p_n-1)/2) being the Legendre symbol((i+j)/p_n), where p_n is the n-th prime.

Conjecture: If p_n == 1 (mod 4), then a(n) == ((p_n-1)/2)! (mod p_n). If p_n == 3 (mod 4), then a(n) == (2/p_n) (mod p_n). Zhi-Wei Sun also made the following general conjecture: On August 19 2013, Zhi-Wei Sun found a formula for a(n). Namely, he made the following conjecture: If p_n == 1 (mod 4) and e(p_n)^{h(p_n)} = (a_n + b_n*sqrt(p_n))/2 with a_n and b_n integers of the same parity (where e(p_n) and h(p_n) are the fundamental unit and the class number of the quadratic field Q(sqrt(p_n)) respectively), then a(n) = - (2/p_n)*2^{(p_n-3)/2}*a_n. If p_n > 3 and p_n == 3 (mod 4), then a(n) = 2^{(p_n-1)/2}. On August 19 2013, Zhi-Wei Sun proved all the conjectured congruences mentioned above by using the identity D(c,d,n) = (-d)^{n*(n+1)/2}*(n!)^{n+1}, where D(c,d,n) is the (n+1) X (n+1) determinant with (i,j)-entry equal to (i+d*j+c)^n for all i,j = 0,...,n. For any prime p == 1 (mod 4) he showed that R(d,p) == (d*(d/p))^{(p-1)/4}*((p-1)/2)! (mod p). Note also that the formula for a(n) found by Sun on August 9, 2013 is actually equivalent to Chapman's result on the evaluation of the determinant |((i+j-1)/p)|_{i,j=1,...,(p+1)/2}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007966 | OEIS-A228005

Determinant of the (p_n-1)/2-by-(p_n-1)/2 matrix with (i,j)-entry being the Legendre symbol ((i^2+j^2+1)/p_n), where p_n is the n-th prime.

Conjecture: We have (a(n)/p_n) = ((2(-1)^{(p_n-1)/2}-4)/p_n). Zhi-Wei Sun also made the following general conjectures for any odd prime p:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007967 | OEIS-A228012

The 2-color Rado number for the equation x_1 + x_2 + ... + x_n = 2*x_0.

Conjecture: For n >= 5, a(n) = (1-(-1)^n+i*(-i)^n-i*i^n+n-(-1)^n*n+2*n^2)/8, where i=sqrt(-1). G.f.: x^2*(x^9-2*x^7-x^6+x^5+x^4-3*x-1) / ((x-1)^3*(x+1)^2*(x^2+1)). - _Colin Barker_, Aug 12 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Donald Vestal_, Aug 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007968 | OEIS-A228016

a(1) = least k such that 1/1+1/2+1/3+1/4+1/5 < H(k) - H(5); a(2) = least k such that H(a(1)) - H(5) < H(k) -H(a(1)), and for n > 2, a(n) = least k such that H(a(n-1)) - H(a(n-2)) > H(k) - H(a(n-1)), where H = harmonic number.

a(n) = 11*a(n-1) - 11*a(n-2) + a(n-3) (conjectured). G.f.: (-54 + 55 x - 5 x^2)/(-1 + 11 x - 11 x^2 + x^3) (conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007969 | OEIS-A228025

a(1) = least k such that 1/2+1/3+1/4+1/5 < H(k) - H(5); a(2) = least k such that H(a(1)) - H(5) < H(k) -H(a(1)), and for n > 2, a(n) = least k such that H(a(n-1)) - H(a(n-2)) > H(k) - H(a(n-1)), where H = harmonic number.

a(n) = A061278(n+1) (conjectured). a(n) = 5*a(n-1) - 5*a(n-2) + a(n-3) (conjectured). G.f.: (-20 + 24 x - 5 x^2)/(-1 + 5 x - 5 x^2 + x^3) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007970 | OEIS-A228036

(10^n)-th primitive antiharmonic number.

We conjecture that lim_{n->oo} a(n)/100^n = lim_{n->oo} A228023(n)/n^2 = 6. This is supported by the values a(n)/(10^n)^2 = 6.05, 5.10, 5.84, 5.99, 6.05 for n = 1..5, as well as by the values of A228023(n)/n^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_ and _Charles R Greathouse IV_, Aug 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007971 | OEIS-A228037

Odd-indexed terms of Agoh's congruence A046094: a(n) is conjectured to be 1 iff 2n+1 is prime.

Odd-indexed terms of Agoh's congruence A046094: a(n) is conjectured to be 1 iff 2n+1 is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Aug 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007972 | OEIS-A228039

Thue-Morse sequence along the squares: A010060(n^2).

(Adapted from Drmota, Mauduit, and Rivat) The Thue-Morse sequence T(n) is a 0-1-sequence that can be defined by T(n) = s2(n) mod 2, where s2(n) denotes the binary sum-of-digits function of n (that is, the number of powers of 2). By definition it is clear that 0 and 1 appear with the same asymptotic frequency 1/2. However, there is no consecutive block of the form 000 or 111, so that the Thue-Morse sequence is not normal. (A 0-1-sequence is normal if every finite 0-1-block appears with the asymptotic frequency 1/2^k, where k denotes the length of the block.) Mauduit and Rivat (2009) showed that the subsequence T(n^2) also has the property that both 0 and 1 appear with the same asymptotic frequency 1/2. This solved a long-standing conjecture by Gelfond (1967/1968). Drmota, Mauduit, and Rivat (2013) proved that the subsequence T(n^2) is actually normal.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Sep 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007973 | OEIS-A228040

Decimal expansion of sum of reciprocals, row 2 of Wythoff array, W = A035513.

Let c be the constant given by A079586, that is, the sum of reciprocals of the Fibonacci numbers F(k) for k>=1. The number c-1, the sum of reciprocals of row 1 of W, is known to be irrational (see A079586). Conjecture: the same is true for all the other rows of W.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007974 | OEIS-A228060

Irregular table of conjectural numbers. Let len(n) be the length (including n) of the Collatz (3x+1) iteration of n. The number n is in row n - len(n).

Irregular table of conjectural numbers. Let len(n) be the length (including n) of the Collatz (3x+1) iteration of n. The number n is in row n - len(n). This sequence is labeled conjectural because (I think) we currently have no way of knowing whether there is some large n that has a long Collatz iteration. Sequence A220139 computes the trajectory for some large numbers, all of which have moderately small length (see A220140). The table below shows numbers in the same row. Notice that each row has only a few numbers. Row 15 is the first with 4 numbers: 26, 37, 103, 124. Row 324 is the first with 5 numbers: 336, 357, 370, 414, 440. Row 16028 has 6 numbers. Row 34937 has 7 numbers. No row in the first 10^7 rows has more than 7 numbers. This sequence is fairly linear; in the first 10^7 terms, no adjacent terms differ by more than 547.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Aug 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007975 | OEIS-A228077

Determinant of the (p_n-1)/2 X (p_n-1)/2 matrix with (i,j)-entry being the Legendre symbol ((j-i)/p_n), where p_n is the n-th prime.

Conjecture: In the case p_n == 1 (mod 4), (2/p_n)*a(n) is a positive odd integer whose prime factors are all congruent to 1 modulo 4, and moreover for some integer b(n) we have b(n) + (2/p_n)*a(n)*sqrt(p_n) = e(p_n)^{(2-(2/p_n))h(p_n)}, where e(p_n) and h(p_n) are the fundamental unit and the class number of the real quadratic field Q(sqrt(p_n)) respectively. M. Vsemirnov has proved _Robin Chapman_'s conjecture on the evaluation of the determinant of the (p+1)/2-by-(p+1)/2 matrix with (i,j)-entry (i,j = 0,...,(p-1)/2) being the Legendre symbol ((j-i)/p), where p is an odd prime. On Aug 14 2013, _Robin Chapman_ informed the author that he first made the conjecture about the exact value of a(n) in a private manuscript dated Aug 05 2003.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007976 | OEIS-A228095

Determinant of the p_n X p_n matrix with (i,j)-entry equal to the Legendre symbol ((i^2 + 3*i*j + 3*j^2)/p_n) for all i,j = 0, 1, ..., p_n-1, where p_n is the n-th prime.

Conjecture: a(n) = 0 if p_n == 5 (mod 6). For an odd prime p and integers c and d, _Zhi-Wei Sun_ defined [c,d]_p to be the determinant of the p X p matrix with (i,j)-entry equal to the Legendre symbol ((i^2+c*i*j+d*j^2)/p). It is easy to see that [-c,d]_p = (-1/p)*[c,d]_p. Sun conjectured that for any integer c and nonzero integer d there are infinitely many odd primes p with [c,d]_p = 0. He also conjectured that if (c,d)_p (defined in the comments in A225611) is nonzero with d not divisible by p, then [c,d]_p/(c,d)_p equals (1-p)/(p-2) or (p-1)/2 according as p divides c^2-4*d or not. He also made some concrete conjectures, for example, [3,2]_p = 0 if p == 3 (mod 4); and [4,2]_p = 0 if p == 5 or 7 (mod 8). On Aug 12 2013, _Zhi-Wei Sun_ conjectured that for any odd prime p and integers c and d with d not divisible by p, if [c,d]_p is nonzero then its p-adic valuation (i.e., p-adic order) must be even.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007977 | OEIS-A228096

Numbers consisting of only odd digits such that no permutation of its digits yields a prime.

Apart from the first term, A061810 is a subsequence. Conjecture: a(n) ~ A061810(n). - _Charles R Greathouse IV_, Feb 15 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jayanta Basu_, Aug 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007978 | OEIS-A228144

Smallest k > n such that j*10^k + m*10^n - 1 is a prime number for at least a pair {j,m} with 0 < j < 10 and 0 < m < 10.

Conjecture: there is always at least one k for each n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Aug 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007979 | OEIS-A228178

The number of boundary edges for all ordered trees with n edges.

Conjecture: 2*(n+3)*a(n) +2*(-7*n-11)*a(n-1) +(29*n+7)*a(n-2) +(-21*n+19)*a(n-3) +2*(2*n-5)*a(n-4)=0. - _R. J. Mathar_, Aug 25 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Louis Shapiro_, Aug 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007980 | OEIS-A228180

The number of single edges on the boundary of ordered trees with n edges.

Conjecture: 2*(n+1)*a(n) +(-13*n+5)*a(n-1) +(23*n-37)*a(n-2) +6*(-2*n+5)*a(n-3)=0. - _R. J. Mathar_, Aug 25 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Louis Shapiro_, Aug 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007981 | OEIS-A228197

Number of n-edge ordered trees with bicolored boundary edges.

Conjecture: (-n+1)*a(n) +2*(5*n-8)*a(n-1) +4*(-8*n+17)*a(n-2) +16*(2*n-5)*a(n-3)=0. - _R. J. Mathar_, Aug 25 2013 a(n) = 2^(2*n)-2^n*JacobiP(n-1,1/2,-n,3) = 2^(2*n)-2*A082590(n-1), which satisfies the above conjecture. - _Benedict W. J. Irwin_, Sep 16 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Louis Shapiro_, Aug 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007982 | OEIS-A228212

Number of second differences of arrays of length n + 2 of numbers in 0..2.

Conjectures from _Colin Barker_, Sep 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007983 | OEIS-A228213

Number of second differences of arrays of length n + 2 of numbers in 0..3.

Conjectures from _Colin Barker_, Sep 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007984 | OEIS-A228214

Number of second differences of arrays of length n + 2 of numbers in 0..4.

Conjectures from _Colin Barker_, Sep 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007985 | OEIS-A228219

Number of second differences of arrays of length 4 of numbers in 0..n.

Conjectures from _Colin Barker_, Mar 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007986 | OEIS-A228220

Number of second differences of arrays of length 5 of numbers in 0..n.

Conjectures from _Colin Barker_, Sep 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007987 | OEIS-A228221

Number of second differences of arrays of length 6 of numbers in 0..n.

Conjectures from _Colin Barker_, Sep 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007988 | OEIS-A228222

Number of second differences of arrays of length 7 of numbers in 0..n.

Conjectures from _Colin Barker_, Sep 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007989 | OEIS-A228223

Number of second differences of arrays of length 8 of numbers in 0..n.

Conjectures from _Colin Barker_, Sep 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007990 | OEIS-A228224

Number of second differences of arrays of length 9 of numbers in 0..n.

Conjectures from _Colin Barker_, Sep 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007991 | OEIS-A228254

Number of third differences of arrays of length n+3 of numbers in 0..2.

Conjectures from _Colin Barker_, Sep 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007992 | OEIS-A228255

Number of third differences of arrays of length n+3 of numbers in 0..3.

Conjectures from _Colin Barker_, Sep 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007993 | OEIS-A228256

Number of third differences of arrays of length n+3 of numbers in 0..4.

Conjectures from _Colin Barker_, Sep 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007994 | OEIS-A228261

Number of third differences of arrays of length 5 of numbers in 0..n.

Conjectures from _Colin Barker_, Sep 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007995 | OEIS-A228262

Number of third differences of arrays of length 6 of numbers in 0..n.

Conjectures from _Colin Barker_, Sep 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007996 | OEIS-A228289

Determinant of the p_n X p_n matrix with (i,j)-entry equal to D(i+j) for all i,j = 0,...,p_n-1, where D(k) = A002895(k) is the k-th Domb number and p_n is the n-th prime.

Conjecture: If p_n == 1 (mod 3) and p_n = x^2 + 3*y^2 with x and y integers, then we have a(n) == (-1)^{(p_n-1)/2}*(4*x^2-2*p_n) (mod p_n^2). In the case p_n == 2 (mod 3), we have a(n) == 0 (mod p_n^2). Zhi-Wei Sun also made the following similar conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007997 | OEIS-A228304

a(n) = Sum_{k=0..n} C(n,k)^4*(-1)^k.

Conjecture: Let p be any odd prime, and let A(p) be the p X p determinant with (i,j)-entry equal to a(i+j) for all i,j = 0,...,p-1. Then A(p) == (-1)^{(p-1)/2} (mod p). Similarly, if c(n) = sum_{k=0}^n (-1)^k*C(n,k)^2*C(2k,k)*C(2(n-k),n-k) and C(p) is the p X p determinant with (i,j)-entry equal to c(i+j) for all i,j = 0,...,p-1, then we have C(p) == 1 (mod p). Adamczewski includes a proof abstract regarding the above conjecture in linked article. - _Michael De Vlieger_, Aug 19 2026 Conjecture: n^3*(n-1)*(12*n^2-63*n+83)*a(n) +(n-2)*(12*n^2-87*n+158)*(n-1)^3*a(n-1) +4*(408*n^6-3774*n^5+13760*n^4-25203*n^3+24465*n^2-11970*n+2340)*a(n-2) +4*(408*n^6-6222*n^5+38750*n^4-126143*n^3+226494*n^2-212867*n+81920)*a(n-3) +16*(n-2)*(12*n^2-15*n+5)*(n-3)^3*a(n-4) +16*(n-3)*(12*n^2-39*n+32)*(n-4)^3*a(n-5)=0. - _R. J. Mathar_, Aug 21 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007998 | OEIS-A228320

The Wiener index of the graph obtained by applying Mycielski's construction to the cycle graph C(n).

The Hosoya-Wiener polynomial is conjectured to be 4nt +(1/2)n(n+9)t^2 + n(n-4)t^3 + (1/2)n(n-7)t^4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Aug 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-007999 | OEIS-A228329

a(n) = Sum_{k=0..n} (k+1)^2*T(n,k)^2 where T(n,k) is the Catalan triangle A039598.

Conjecture: n*(2*n+1)*a(n) + 2*(-26*n^2+25*n-11)*a(n-1) + 20*(4*n-5)*(4*n-7)*a(n-2) = 0. - _R. J. Mathar_, Sep 08 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008000 | OEIS-A228330

Let h(m) denote the sequence whose n-th term is Sum_{k=0..n} (k+1)^m*T(n,k)^2, where T(n,k) is the Catalan triangle A039598. This is h(4).

Conjecture: n*(2*n+1)*(3467*n-4029)*a(n) + 8*(-36721*n^3 + 109040*n^2 - 137926*n + 69822)*a(n-1) + 4*(4*n-9)*(45706*n-7907)*(4*n-7)*a(n-2) = 0. - _R. J. Mathar_, Sep 08 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008001 | OEIS-A228331

Let h(m) denote the sequence whose n-th term is Sum__{k=0..n} (k+1)^m*T(n,k)^2, where T(n,k) is the Catalan triangle A039598. This is h(5).

Conjecture: n^2*a(n) + 4*(2*n^2-22*n+11)*a(n-1) + 16*(-7*n^2-54*n+166)*a(n-2) - 1088*(2*n-3)*(2*n-7)*a(n-3) = 0. - _R. J. Mathar_, Sep 08 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008002 | OEIS-A228333

Let h(m) denote the sequence whose n-th term is Sum_{k=0..n} (k+1)^m*T(n,k)^2, where T(n,k) is the Catalan triangle A039598. This is h(7).

Conjecture: n^2*(304*n-411)*a(n) + 4*(-1814*n^3+2554*n^2-4776*n+7567)*a(n-1) + 32*(2*n-5)*(2*n-1)*(299*n-176)*a(n-2) = 0. - _R. J. Mathar_, Dec 04 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008003 | OEIS-A228379

Determinant of the n X n matrix with (i,j)-entry equal to (i^2+j^2)^n for all i,j = 0,...,n-1.

Conjecture: If n > 2, then (-1)^{n(n-1)/2}*a(n) > 0 and 2*Product_{k=1..n} (k!*(2k-1)!) divides a(n). This conjecture implies that if p = 2*n-1 > 3 is a prime then we have a((p+1)/2) == 0 (mod p). So Sun's Conjecture 3.7 in the reference would follow from the above conjecture. If b(n) denotes the n X n determinant with (i,j)-entry equal to (i+j)^n for all i,j = 0,...,n-1, then we conjecture that b(n)*(-1)^(n*(n-1)/2) / ((n-2)!*n*Product_{k=1..n} (k!)) is a positive integer for any integer n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008004 | OEIS-A228404

The number of complete binary trees with bicolored twigs. A twig is a vertex with one child on the boundary and the other child having no descendants.

Conjecture: -5*(n+3)*(n-2)*a(n) + 5*(-n^2-n+18)*a(n-1) + 5*(-n^2-n+48)*a(n-2) +(-5*n^2+20029*n+720)*a(n-3) + (-5*n^2-104153*n+186654)*a(n-4) + (-5*n^2 +130153*n -508806)*a(n-5) + 13650*(2*n-11)*(n-7)*a(n-6) = 0. - _R. J. Mathar_, Aug 08 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Louis Shapiro_, Aug 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008005 | OEIS-A228407

The digits of a(n) and a(n+1) together can be reordered to form a palindromic integer; the lexicographically earliest injective sequence of nonnegative integers with this property.

It is conjectured that the sequence is a permutation of the nonnegative integers (motivating the choice of offset 0), i.e., that all numbers will eventually occur. (The conjecture is true - see below. - _N. J. A. Sloane_, Nov 12 2013) To test this conjecture, consider the indices n where the smallest integers not yet used occur. If the conjecture is true, this is equivalent to a(m)>a(n) for all m>n; if not, then this list ends at the first missing number. These [n,a(n)] are: [0, 0], [2, 1], [6, 2], [10, 3], [17, 4], [34, 5], [45, 6], [65, 7], [81, 8], [118, 9], [119, 29], [122, 39], [145, 44], [152, 45], [197, 46], [230, 47], [271, 48], [362, 49], [533, 57], [740, 58], [754, 68], [816, 69], [855, 89], [856, 98], [857, 198], [1011, 211], [1012, 222], [1110, 224], [1232, 225], [1385, 234], [1406, 236], [1413, 237], [1767, 238], [1921, 239], [2555, 257], [2590, 259], [2597, 269], [4354, 279], [4355, 297], [4361, 379], [4362, 397], [4368, 479], [4369, 497],...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _M. F. Hasler_, Nov 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008006 | OEIS-A228410

The digits of a(n) and a(n+1) together can be reordered to form a palindrome; lexicographically least injective sequence of positive integers with this property.

It is conjectured that the sequence is a permutation of the positive integers, i.e., that all numbers will eventually occur. To test this conjecture, one can consider the indices n at which occur the numbers equal to the smallest integer not yet used. If the conjecture is true, this is equivalent to a(m)>a(n) for all m>n; if not, then this list ends at the first missing number. These [n,a(n)] are: [1, 1], [5, 2], [12, 3], [19, 4], [35, 5], [45, 6], [62, 7], [78, 8], [88, 9], [89, 29], [92, 39], [118, 44], [149, 45], [187, 46], [314, 47], [432, 49], [477, 59], [506, 67], [507, 76], [521, 78], [531, 79], [572, 89], [573, 98], [574, 198], [954, 211][955, 222], [956, 233], [1602, 234], [1616, 235], [1623, 237], [1924, 238], [1959, 239], [2508, 258], [2515, 278], [2536, 279], [4046, 289], [4047, 298], [4053, 489], [4054, 498], ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Nov 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008007 | OEIS-A228424

Primes that can be written as a sum of a triangular number and a square.

This sequence is interesting because of the conjecture in the comments in A228425.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008008 | OEIS-A228425

Number of ways to write n = x + y (x, y > 0) with x*(x+1)/2 + y^2 prime.

Conjecture: a(n) > 0 for all n > 1. For m = 3, 4, 5, ... the m-gonal numbers are given by p_m(x) = (m-2)*x*(x-1)/2 + x (x = 0, 1, 2, ...). We note that there are many pairs m > k > 2 such that all sufficiently large integers n can be written as x + y (x, y > 0) with p_k(x) + p_m(y) prime. For example, we conjecture that the pair (k, m) works if k is among 3, 4, 6 , and m > k is not congruent to k modulo 2. For k = 5, we guess that the pair (5, m) works if m is congruent to 0 or 4 modulo 6. We conjecture that the only pairs (k,m) with 2 < k <= 10 and k< m <= 100 such that any integer n > 1 can be written as x + y (x, y > 0) with p_k(x) + p_m(y) prime, are as follows: (3,4),(3,6),(3,28),(3,46),(3,52),(3,82),(3,88),(4,7),(4,15),(4,25),(4,27),(4,37),(4,43),(4,63),(4,67),(4,97),(6,25),(6,43),(6,73),(7,10),(7,18),(7,100),(10,15),(10,19),(10,27),(10,37),(10,55),(10,75),(10,79),(10,87),(10,99). We also conjecture that any integer n > 1 can be written as x + y (x, y > 0) with p_k(x) + p_{k+1}(y) prime, if and only if k is among 3, 39, 99.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008009 | OEIS-A228428

Number of ways to write n = x + y (x, y > 0) with p(99, x) + p(100, y) prime, where p(m, k) denotes the m-gonal number (m-2)*k*(k-1)/2 + k.

Conjecture: a(n) > 0 for all n > 1. See also A228425 for other similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008010 | OEIS-A228429

Number of ways to write n = x + y (x, y > 0) with p(39, x) + p(40, y) prime, where p(m, k) denotes the m-gonal number (m-2)*k*(k-1)/2 + k.

By a conjecture in A228425, we should have a(n) > 0 for all n > 1. Conjecture: For each m = 3, 4, ..., any sufficiently large integer n can be written as x + y (x, y > 0) with p(m, x) + p(m+1, y) prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008011 | OEIS-A228430

Number of ways to write n = x + y (x, y > 0) with x^4 + y*(y+1)/2 prime.

Conjecture: (i) a(n) > 0 except for n = 1, 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008012 | OEIS-A228431

Number of ordered ways to write n = x + y (x, y > 0) with p(3, x) + p(6, y) prime, where p(3, k) denotes the triangular number k*(k+1)/2 and p(6, k) denotes the hexagonal number k*(2*k-1) = p(3, 2*k-1).

Conjecture: a(n) > 0 for all n > 1. See also A228425, A228428, A228429 and A228430 for more similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008013 | OEIS-A228441

Expansion of Sum_{k>0} -(-x)^k / (1 + x^k).

Conjecture: a(n) = (7 - 2*(-1)^n)*tau(n) - 4*tau(2*n) = 5*tau(n) - (3 + (-1)^n)*tau(2*n), where tau = A000005. - _Velin Yanev_, Dec 17 2019 The proof of the above conjecture easily follows from the fact that both a(n) and tau(n) are multiplicative arithmetical functions and tau(p^e) = e + 1 for prime p. - _Peter Bala_, Jan 28 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Nov 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008014 | OEIS-A228446

a(n) = smallest prime p such that 2*n+1 = p + x*(x+1) for some positive integer x, or -1 if no such prime exists.

Based on Sun's conjecture 1.4 in the paper referenced below.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bill McEachen_, Oct 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008015 | OEIS-A228462

Number of arrays of maxima of three adjacent elements of some length 7 0..n array.

Conjectures from _Colin Barker_, Sep 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008016 | OEIS-A228463

Number of arrays of maxima of three adjacent elements of some length 8 0..n array.

Conjectures from _Colin Barker_, Sep 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008017 | OEIS-A228464

Number of arrays of maxima of three adjacent elements of some 0..n array of length 9.

Conjectures from _Colin Barker_, Mar 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008018 | OEIS-A228469

a(n) = 6*a(n-2) + a(n-4), where a(0) = 2, a(1) = 8, a(2) = 13, a(3) = 49.

Conjecture: a(n) is the least positive integer c for which there is a positive integer b for which trace(b,c) consists of the first n letters of 01010101010101...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008019 | OEIS-A228471

a(n) = 6*a(n-2) + a(n-4), where a(0) = 3, a(1) = 5, a(2) = 19, a(3) = 31.

Conjecture: a(n) is the least positive integer c for which there is a positive integer b for which trace(b,c) consists of the first n letters of 101010101010101..., where "trace" is as defined at A228469.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008020 | OEIS-A228487

Trace(Pi), where trace is defined in Comments.

Conjectures: (1) trace(pi) is not periodic; (2) if k is a positive integer, then trace(sqrt(k)) is purely periodic; (3) if r is a quadratic irrational, or if r = s*e where s is a positive rational number, or if r = s*e^2 where s is a positive rational number, then trace(r) is eventually periodic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008021 | OEIS-A228488

Period length of trace(sqrt(n)).

It is assumed that trace(sqrt(n)) is purely periodic, as conjectured at A228487 where trace is defined.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008022 | OEIS-A228489

Period of trace(n*phi), where phi = (1+sqrt(5))/2 = golden ratio.

It is assumed that trace(n*phi) is purely periodic, as conjectured at A228487 where trace is defined.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008023 | OEIS-A228497

Decimal expansion of the fourth root of 1/2.

Conjecture: Equals Sum_{k>=0} A189925(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Aug 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008024 | OEIS-A228511

a(n) = sum_{k=0}^n binomial(n,k)^2*4^k*A000108(k).

Conjecture: Let p be any odd prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008025 | OEIS-A228514

a(n) = Sum_{k=0..n} binomial(n,k)^2*binomial(2k,k+1).

Conjecture: Let p > 3 be a prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008026 | OEIS-A228518

a(n) is the least r > 1 for which the interval (r*n, r*(n+1)) contains no terms of A050376 or a(n) = 0 if no such r exists.

Conjecture: a(n) = 0 iff n = 1, 2, 3, 4, 5, 6, 9, 12, 14, 24, 56.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Aug 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008027 | OEIS-A228541

Numbers having at least one prime factor of the form 30*k + 1.

Conjecture: Numbers m such that abs(Sum_{k=1..m} [k|m]*A008683(k)*(-1)^(k/15)) = 0. - _Mats Granvik_, Jul 06 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Aug 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008028 | OEIS-A228549

Determinant of the n X n matrix with (i,j)-entry equal to 1 or 0 according as i + j is squarefree or not.

Conjecture: a(n) is always nonzero. Moreover, |a(n)|^(1/n) tends to infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008029 | OEIS-A228552

Square root of the absolute value of A069191(n).

According to the comments of A069191, a(n) should be always integral. Note that a(2*n) is the absolute value of A228616(n) by the comments of A228591. We conjecture that a(n) > 0 for all n > 15.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008030 | OEIS-A228558

Numbers k such that (17^k + 4^k)/21 is prime.

Conjecture: All terms are prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008031 | OEIS-A228559

Determinant of the n X n matrix with (i,j)-entry equal to 1 or 0 according as i + j is a Sophie Germain prime or not.

Conjecture: a(n) is nonzero if n is greater than 55 and not congruent to 5 modulo 6. Zhi-Wei Sun also had some similar conjectures. For example, if b(n) denotes the n X n determinant with (i,j)-entry equal to 1 or 0 according as i + j and 2*(i + j) - 1 are both prime or not, then b(n) is nonzero when n is greater than 125 and not congruent to 3 modulo 6. (Just like a(6*n-1) = 0, we can show that b(6*n-3) = 0.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008032 | OEIS-A228561

Determinant of the n X n matrix with (i,j)-entry equal to 1 or 0 according as i + j and 4*(i + j)^2 + 1 are both prime or not.

Conjecture: a(n) is nonzero for each n > 28.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008033 | OEIS-A228574

Determinant of the 2*n X 2*n matrix with (i,j)-entry equal to 1 or 0 according as i + j is a prime congruent to 1 mod 4 or not.

Conjecture: a(2*n-1) = 0 for all n > 0, and a(2*n) is nonzero when n > 9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008034 | OEIS-A228591

Determinant of the n X n (0,1)-matrix with (i,j)-entry equal to 1 if and only if i + j is 2 or an odd composite number.

Conjecture: a(n) = 0 for no n > 15.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008035 | OEIS-A228616

Determinant of the n X n matrix with (i,j)-entry equal to 1 or 0 according as 2*(i + j) - 1 is prime or not.

Conjecture: a(n) is nonzero for any n > 7. Clearly this conjecture implies that there are infinitely many primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008036 | OEIS-A228624

Determinant of the n X n matrix with (i,j)-entry equal to 1 or 0 according as i + j is a square or not.

Conjecture: a(n) is nonzero for any n > 21. Zhi-Wei Sun also made the following similar conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008037 | OEIS-A228626

Number of Hamiltonian cycles in the undirected simple graph G_n with vertices 1,...,n which has an edge connecting vertices i and j if and only if |i-j| is prime.

Conjecture: a(n) > 0 for all n > 4. In other words, for each n = 5,6,... there is a permutation i_1,...,i_n of 1,...,n such that |i_1-i_2|, |i_2-i_3|, ..., |i_{n-1}-i_n| and |i_n-i_1| are all prime. Note that this conjecture is different from the prime circle problem in A051252 though they look similar. On August 30 2013, Yong-Gao Chen (from Nanjing Normal University) confirmed the conjecture for n > 12 as follows: If n = 2*k then G_n contains a Hamiltonian cycle (1,3,5,2,7,9,...,2k-5,2k-3,2k,2k-2,2k-4,2k-1,2k-6,2k-8,...,6,4);

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008038 | OEIS-A228638

Determinant of the n X n matrix with (i,j)-entry equal to 1 or 0 according as |i-j| is prime or not.

Conjecture: a(n) is nonzero for any n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008039 | OEIS-A228728

a(1)=1, a(2)=2 and for n > 2, a(n) is the least integer > a(n-1) such that there is a permutation b(1), ..., b(n) of a(1), ..., a(n) with b(1) = a(1) and b(n) = a(n), and with the n numbers |b(1)-b(2)|, |b(2)-b(3)|, ..., |b(n-1)-b(n)|, |b(n)-b(1)| pairwise distinct.

Conjecture: For any n distinct real numbers a_1 < a_2 < ... < a_n, if there is a permutation b_1,b_2,...,b_n of a_1,...,a_n with |b_1-b_2|, |b_2-b_3|, ..., |b_{n-1}-b_n|, |b_n-b_1| pairwise distinct, then there exists a permutation c_1,c_2,...,c_n of a_1,...,a_n with c_1 = a_1 and c_n = a_n such that the n numbers |c_1-c_2|, |c_2-c_3|, ..., |c_{n-1}-c_n|, |c_n-c_1| are pairwise distinct. This conjecture is somewhat curious but we are unable to find a counterexample.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008040 | OEIS-A228730

Lexicographically earliest sequence of distinct nonnegative integers such that the sum of two consecutive terms is a palindrome in base 10.

Conjectured to be a permutation of the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Tek_, Aug 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008041 | OEIS-A228744

Number of arrays of the median of three adjacent elements of some length 9 0..n array.

Conjectures from _Colin Barker_, Mar 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008042 | OEIS-A228767

Second bisection of the inverse binomial transform of the rational sequence with e.g.f. (x/2)*(exp(-x)+1)/(exp(x)-1).

Conjecture: G.f. -x*(2-11*x+21*x^2-2*x^3+8*x^4)/((1-x)^2*(1-4*x)^2). [_Bruno Berselli_, Sep 03 2013] Conjecture: a(n) = (8+4^n)*(1-2*n)/8 for n>1, a(1)=-2. [_Bruno Berselli_, Sep 03 2013]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, following a suggestion of _Paul Curtz_, Sep 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008043 | OEIS-A228770

The number of sum indecomposable permutations which avoid the patterns 3124 and 4312.

Conjecture: -(n+1)*(39961*n-2474598)*a(n) +(-39961*n^2-25975201*n+4949196) *a(n-1) +3*(1460811*n^2+27429105*n-41310802) *a(n-2) +3 *(-8653921*n^2-4750029*n+74360724) *a(n-3) +4*(15005713*n^2-82481258*n+83094771) *a(n-4) +12*(-4937548*n^2+40726604*n-73155719) *a(n-5) +16*(652718*n-2110173)*(2*n-13) *a(n-6)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jay Pantone_, Sep 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008044 | OEIS-A228771

The number of skew sum indecomposable permutations which avoid the patterns 3124 and 4312.

Conjecture: -163*(n+2)*(4*n-413) *a(n) +(-652*n^2-725425*n-452889) *a(n-1) +5*(14473*n^2+512276*n-443094) *a(n-2) +(-410045*n^2-2408964*n+8429009) *a(n-3) +2*(404156*n^2-1297075*n-1518084)*a(n-4) -8*(29333*n-32490)*(2*n-11)*a(n-5)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jay Pantone_, Sep 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008045 | OEIS-A228775

a(n) is the maximal k>=1 such that nextprime(j*n)<=(j+1)*n, j=1,...,k.

Conjectural inequality: for n>=2, a(n) <= log^2(n*a(n)). This essentially corresponds to Cramer's conjecture for prime gaps.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008046 | OEIS-A228781

Irregular triangle read by rows: coefficients of minimal polynomial of a certain algebraic number S2(2*k+1) from Q(2*cos(Pi/n)) related to the regular (2*k+1)-gon, k >= 1.

Conjecture from _Seppo Mustonen_, rewritten for the p(n, x) coefficients for odd primes: p(prime(j), x) = Sum_{i=0..imax(j)} (-1)^(imax(j - i))* binomial(prime(j), 2*i+1)*x^i, with imax(j) = (prime(j)-1)/2. See the adapted eq. (5) of the S. Mustonen paper.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008047 | OEIS-A228791

Number of n X 3 binary arrays with top left element equal to 1 and no two ones adjacent horizontally or nw-se.

Conjectures from _Colin Barker_, Sep 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008048 | OEIS-A228797

Number of 2 X n binary arrays with top left element equal to 1 and no two ones adjacent horizontally or nw-se.

Conjectures from _Colin Barker_, Sep 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008049 | OEIS-A228858

Let A(x,y,z) denote the integer area A of the triangle with integer side lengths (x,y,z). a(n) is the smallest area such that there exist exactly n distinct triangles (x_i, y_i, z_i) satisfying A(x_1, y_1, z_1) = A(x_2, y_2, z_2) = ... = A(x_n, y_n, z_n).

Conjecture: there exists an integer corresponding to the common area of exactly n distinct triangles (x_i, y_i, z_i) for i = 1,2,...,n such that A(x_1, y_1, z_1) = A(x_2, y_2, z_2) = ... = A(x_n, y_n, z_n) for n = 1, 2, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Sep 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008050 | OEIS-A228885

Determinant of the n X n matrix with (i,j)-entry equal to 1 or 0 according as i + j is coprime to n or not.

Conjecture: If n is squarefree, then (-1)^(n*(n-1)/2)*a(n) > 0. Conjectures from _Ridouane Oudra_, Mar 13 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008051 | OEIS-A228886

Number of permutations i_0,i_1,...,i_n of 0,1,...,n with i_0 = 0 and i_n = n, and with i_0+i_1, i_1+i_2, ..., i_{n-1}+i_n, i_n+i_0 all coprime to both n-1 and n+1.

Conjecture: a(n) > 0 except for n = 2, 4. Note that this conjecture is stronger than the one stated in the comments in A228860. Also, there is no permutation i_0, ..., i_7 of 0, ..., 7 with i_0+i_1, i_1+i_2, ..., i_6+i_7, i_7+i_0 all coprime to 7*13 - 1 = 90. We have some other similar conjectures. For example, for any positive integer n there is a permutation

i_0,i_1,...,i_n of 0,1,...,n with i_0 = 0 and i_n = n such that i_0 + i_1, i_1 + i_2, ..., i_{n-1} + i_n, i_n + i_0 are all relatively prime to both 2*n-1 and 2*n+1. On Sep 10 2013, Zhi-Wei Sun proved the conjecture for any positive odd integer n. In fact, if n == 1 or 3 (mod 6) then we may take the permutation 0,n-2,2,n-4,4,...,1,n-1,n which meets the requirement; if n == 3 or 5 (mod 6) then the permutation 0,1,n-1,3,n-3,...,n-2,2,n suffices for our purpose.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008052 | OEIS-A228907

G.f. satisfies: A(x) = 1 + Sum_{n>=0} x^n * (1 - A(x)^(2*n))/(1 - A(x)).

Conjectured to be the number of permutations of length n+1 avoiding the partially ordered pattern (POP) {1>4, 1>5, 2>4, 2>5, 5>3} of length 5. That is, conjectured to be the number of length n+1 permutations having no subsequences of length 5 in which the first and second elements are larger than the elements in positions 4 and 5, and the fifth element is larger than the element in position 3.- _Sergey Kitaev_, Dec 13 2020 This conjecture has been proven. It can be restated as the number of size n permutations avoiding 45123, 45132, 45213, 54123, 54132, 54213. - _Christian Bean_, Jul 24 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008053 | OEIS-A228916

Indices of primes in sequence A108300.

Conjecture: for n>1 a(n) is a multiple of 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Sep 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008054 | OEIS-A228931

Optimal ascending continued fraction expansion of sqrt(2)-1.

Conjecture: The terms from a(3) are all positive and can be generated by the recurrence relation a(k+1) = a(k)^2 - 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Artico_, Sep 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008055 | OEIS-A228933

Optimal ascending continued fraction expansion of phi-1 = 1/phi = (sqrt(5)-1)/2.

Conjecture: The golden ratio (phi) expansion exhibits from the fourth term the recurrence relation a(n) = a(n-1)^2 - 2 described in A228931.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Artico_, Sep 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008056 | OEIS-A228934

Optimal ascending continued fraction expansion of sqrt(44) - 6.

Conjecture: The terms of the expansion of sqrt(x) are all negative starting from a(4) and satisfy these recurrence relations for n>=3: a(2n) = 4*a(2n-1) - 4 and a(2n+1) = -2*a(2n-1)^2 + 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Artico_, Sep 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008057 | OEIS-A228943

Number of decompositions of highly composite numbers (A002182) into unordered sums of two primes.

Conjecture: (a) This sequence is strictly increasing beginning with n=5. (b) For all n>2, if p is the greatest prime with p<A002182(n)-1, then A002182(n)-p is prime. This is a strengthening of a conjecture regarding A117825. - _Jaycob Coleman_, Sep 08 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaycob Coleman_, Sep 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008058 | OEIS-A228945

Number of ways to write highly composite numbers (A002182(n)) as the difference of two primes, both <= 2*A002182(n).

Conjectures: (a) This sequence is strictly increasing beginning with n=7. (b) If p is the smallest prime with p > A002182(n)+1, then p-A002182(n) is prime. This is a strengthening of a conjecture regarding A117825.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaycob Coleman_, Sep 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008059 | OEIS-A228947

a(n) = sigma(n) - phi(n) - n.

a(n) = 0 <=> n = 2 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Oct 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008060 | OEIS-A228993

Decimal expansion of L(Pi), the limit of iterations of continued fraction transforms of Pi.

Conjecture: if x is an irrational number between 3 and 4, then L(x) = L(Pi).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008061 | OEIS-A229004

Indices of Bell numbers divisible by 3.

Wagstaff shows that N(p) = (p^p-1)/(p-1) is the period for all primes p < 102, for p=3 then N(3) = A054767(3) = 13, Bell numbers with indices less than or equal to 13 that are divisible by 3 are those with indices: 4, 8, 9, 11, so the conjecture holds. - _Enrique Pérez Herrero_, Sep 12 2013 Conjectures from _Colin Barker_, Jul 16 2014: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enrique Pérez Herrero_, Sep 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008062 | OEIS-A229005

Number of undirected circular permutations i_0, i_1, ..., i_n of 0, 1, ..., n such that all the n+1 numbers |i_0^2-i_1^2|, |i_1^2-i_2^2|, ..., |i_{n-1}^2-i_n^2|, |i_n^2-i_0^2| are of the form (p-1)/2 with p an odd prime.

Conjecture: a(n) > 0 except for n = 2, 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008063 | OEIS-A229013

Number of arrays of median of three adjacent elements of some length-5 0..n array, with no adjacent equal elements in the latter.

Conjectures from _Colin Barker_, Sep 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008064 | OEIS-A229014

Number of arrays of median of three adjacent elements of some length 6 0..n array, with no adjacent equal elements in the latter.

Conjectures from _Colin Barker_, Sep 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008065 | OEIS-A229015

Number of arrays of median of three adjacent elements of some length 7 0..n array, with no adjacent equal elements in the latter.

Conjectures from _Colin Barker_, Sep 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008066 | OEIS-A229016

Number of arrays of median of three adjacent elements of some length 8 0..n array, with no adjacent equal elements in the latter.

Conjectures from _Colin Barker_, Sep 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008067 | OEIS-A229017

Number of arrays of median of three adjacent elements of some length 9 0..n array, with no adjacent equal elements in the latter.

Conjectures from _Colin Barker_, Sep 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008068 | OEIS-A229019

Minimal position at which the sequence defined in the same way as A159559 but with initial term prime(n) merges with A159559; a(n)=0 if there is no such position.

Conjecture: all terms are positive.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008069 | OEIS-A229038

Number of undirected circular permutations (i_1, ..., i_n) of 1, ..., n such that all the n sums i_1^2+i_2^2, ..., i_{n-1}^2+i_n^2, i_n^2+i_1^2 are among those integers m with the Jacobi symbol (m/(2*n+1)) equal to 1.

Conjecture: a(n) > 0 if 2*n+1 is a prime greater than 13. When p = 2*n+1 > 13 is a prime, the conjecture indicates that there is a numbering a_1, a_2, ..., a_n of all the (p-1)/2 = n quadratic residues modulo p such that all the n adjacent sums a_1+a_2, a_2+a_3, ..., a_{n-1}+a_n, a_n+a_1 are also quadratic residues modulo p. Zhi-Wei Sun also made the following conjecture: Noga Alon has confirmed this conjecture. See also A229878 for a stronger conjecture which remains open.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008070 | OEIS-A229044

G.f. A(x) satisfies: [x^(n+1)] A(x)^(n^2) = 0 for n>=0.

a(n) is odd iff n+1 is a power of 2 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008071 | OEIS-A229046

G.f.: Sum_{n>=0} n! * x^n * (1+x)^n / Product_{k=1..n} (1 + k*x).

Conjecturally, the number of sequences (e(1), ..., e(n)), 0 <= e(i) < i, such that there is no triple i < j < k with e(i) = e(k). [Martinez and Savage, 2.13] - _Eric M. Schmidt_, Jul 17 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008072 | OEIS-A229053

Number of standard Young tableaux of n cells and height <= 11.

Conjecture: a(n) ~ k^n/Pi^(k/2)*(k/n)^(k*(k-1)/4) * prod(j=1,k,Gamma(j/2)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Sep 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008073 | OEIS-A229064

Lesser of Fermi-Dirac twin primes: both a(n)(>=5) and a(n)+2 are in A050376.

There exist conjecturally only 5 Fermat primes F, such that both F-1 and F are in A050376. If we add pair (3,4), then we obtain exactly 6 such pairs as an analog of the unique pair (2,3) in usual arithmetic, which is not considered as a pair of twin primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008074 | OEIS-A229068

Number of standard Young tableaux of n cells and height <= 12.

Conjecture: generally (for tableaux with height <= k), a(n) ~ k^n/Pi^(k/2) * (k/n)^(k*(k-1)/4) * Product_{j=1..k} Gamma(j/2); set k=12 for this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Sep 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008075 | OEIS-A229075

Primes of the form p^2 + q^2 + 21, where p and q are consecutive primes.

Conjecture: the expression p^2+q^2+c with p and q consecutive primes and c=21 generates more primes than any other value of c in the range 1..150. Hence, c=21 is considered for this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _K. D. Bajpai_, Sep 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008076 | OEIS-A229082

Number of circular permutations i_0, i_1, ..., i_n of 0, 1, ..., n such that all the n+1 numbers i_0^2+i_1, i_1^2+i_2, ..., i_{n-1}^2+i_n, i_n^2+i_0 are of the form (p-1)/2 with p an odd prime.

Conjecture: a(n) > 0 except for n = 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008077 | OEIS-A229083

Numbers k such that the distance between the k-th triangular number and the nearest square is at most 1.

G.f.: (x^7 - 2*x^6 + x^5 - 3*x^4 + x^3 + 2*x^2 + x + 1)/((1-2*x^2+x^4)*(1-2*x^2-x^4)*(1-x)) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Sep 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008078 | OEIS-A229086

Numbers k such that the concatenation of the divisors of k written in base 2 is a palindromic number.

Conjecture: a(n) for n > 1 is not equal to A161387, sequence also contains composites.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Sep 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008079 | OEIS-A229088

Numbers k such that sigma(k) mod k = antisigma(k) mod k, where sigma(k) = A000203(k) = sum of divisors of k, antisigma(k) = A024816(k) = sum of non-divisors of k.

Conjecture: For k>1, numbers k such that GCD(sigma(k), k) = n/4. - _Jaroslav Krizek_, Sep 23 2014 This conjecture is equivalent to the absence of odd multiperfect numbers > 1. - _Max Alekseyev_, Jul 31 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Oct 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008080 | OEIS-A229110

Sum of non-divisors of n reduced modulo n.

Numbers n such that a(n)=0 are: 1, 2, 24, 4320, 4680, ... (see A159907, conjecture by _Jaroslav Krizek_ and further comments). - _Michel Marcus_, Sep 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Sep 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008081 | OEIS-A229118

Distance from the n-th triangular number to the nearest square.

Conjecture: running average of a(n) ~ sqrt(s)/2 where s is the associated nearest square. - _Bill McEachen_, Jun 08 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Sep 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008082 | OEIS-A229130

Number of permutations i_0, i_1, ..., i_n of 0, 1, ..., n with i_0 = 0 and i_n = n such that the n+1 numbers i_0^2+i_1, i_1^2+i_2, ..., i_{n-1}^2+i_n, i_n^2+i_0 are all relatively prime to both n-1 and n+1.

Conjecture: a(n) > 0 except for n = 2, 5, 11. Similarly, for any positive integer n not equal to 4, there is a permutation i_0, i_1, ..., i_n of 0, 1, ..., n with i_0 = 0 and i_n = n such that the n+1 numbers i_0^2-i_1, i_1^2-i_2, ..., i_{n-1}^2-i_n, i_n^2-i_0 are all coprime to both n-1 and n+1. Zhi-Wei Sun also made the following general conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008083 | OEIS-A229131

Numbers k such that the distance between the k-th triangular number and the nearest square is exactly 1.

G.f.: (-x^7 + 2*x^6 - 2*x^5 + 4*x^4 - 5*x^3 + 2*x^2 + x + 1)/((1-6*x^3+x^6)*(1-x)) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Sep 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008084 | OEIS-A229136

Number of solutions to Sum_{i=1..n} x_i^2 == 1 (mod 4) with x_i in 0..3.

Conjecture: a(n) = A131885(n)*2^(n-1) for n >= 1. [Corrected by _Petros Hadjicostas_, Dec 20 2019] It follows that 2*b(n-1) = 4*b(n-2) - 4*b(n-3) + 2^(n-2) for n >= 4. Subtracting the last equation from the previous one, we get (after some algebra) b(n) = 4*b(n-1) - 6*b(n-2) + 4*b(n-3) for n >= 4. We can easily verify that b(1) = 2, b(2) = 4, and b(3) = 6, and this proves that b(n) = A131885(n) for n >= 1. This proves the above conjecture. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Sep 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008085 | OEIS-A229140

Smallest k such that k^2 + l^2 = n-th number expressible as sum of two squares (A001481).

Conjecture: the values between two zeros are always distinct from each other.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Sep 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008086 | OEIS-A229141

Number of circular permutations i_1, ..., i_n of 1, ..., n such that all the n sums i_1^2+i_2, ..., i_{n-1}^2+i_n, i_n^2+i_1 are among those integers m with the Jacobi symbol (m/(2n+1)) equal to 1.

Conjecture: a(n) > 0 if 2*n+1 is a prime greater than 11. Zhi-Wei Sun also made the following conjectures: See also the linked arXiv paper of Sun for more conjectures involving primitive roots modulo primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008087 | OEIS-A229166

Number of ordered ways to write n = x*(x+1)/2 + y with y*(y+1)/2 + 1 prime, where x and y are nonnegative integers.

Conjecture: a(n) > 0 for all n > 0. Moreover, if n > 0 is not among 1, 3, 60, then there are positive integers x and y with x*(x+1)/2 + y = n such that y*(y+1)/2 + 1 is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008088 | OEIS-A229232

Number of undirected circular permutations pi(1), ..., pi(n) of 1, ..., n with the n numbers pi(1)*pi(2)-1, pi(2)*pi(3)-1, ..., pi(n-1)*pi(n)-1, pi(n)*pi(1)-1 all prime.

Conjecture: a(n) > 0 for all n > 5 with n not equal to 13. Zhi-Wei Sun also made the following conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008089 | OEIS-A229244

Number of n-permutations such that at least one cycle has size ceiling(n/2).

Conjecture: (n+1)*a(n) +(-3*n+1)*a(n-1) -(n-2)*(n^2-2*n-1)*a(n-2) +(n-2)*(n-3)^2*a(n-3)=0. - _R. J. Mathar_, May 23 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Critzer_, Sep 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008090 | OEIS-A229289

Primes p of the form p = 2^k * m + 1, where (i) m is squarefree and odd, (ii) all primes that divide m are in the sequence, and (iii) k is 0, 1, or 2.

Also prime factors of terms in A341858. It is conjectured that this sequence is infinite. - _Jianing Song_, Feb 22 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Oct 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008091 | OEIS-A229422

Number of n X 2 0..2 arrays with horizontal differences mod 3 never 1, vertical differences mod 3 never -1, rows lexicographically nondecreasing, and columns lexicographically nonincreasing.

Conjectures from _Colin Barker_, Sep 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008092 | OEIS-A229423

Number of n X 3 0..2 arrays with horizontal differences mod 3 never 1, vertical differences mod 3 never -1, rows lexicographically nondecreasing, and columns lexicographically nonincreasing.

Conjectures from _Colin Barker_, Sep 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008093 | OEIS-A229424

Number of n X 4 0..2 arrays with horizontal differences mod 3 never 1, vertical differences mod 3 never -1, rows lexicographically nondecreasing, and columns lexicographically nonincreasing.

Conjectures from _Colin Barker_, Sep 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008094 | OEIS-A229425

Number of n X 5 0..2 arrays with horizontal differences mod 3 never 1, vertical differences mod 3 never -1, rows lexicographically nondecreasing, and columns lexicographically nonincreasing.

Conjectures from _Colin Barker_, Sep 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008095 | OEIS-A229439

Number of n X 2 0..2 arrays with horizontal differences mod 3 never 1, vertical differences mod 3 never -1, and rows and columns lexicographically nondecreasing.

Conjectures from _Colin Barker_, Sep 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008096 | OEIS-A229440

Number of n X 3 0..2 arrays with horizontal differences mod 3 never 1, vertical differences mod 3 never -1, and rows and columns lexicographically nondecreasing.

Conjectures from _Colin Barker_, Sep 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008097 | OEIS-A229441

Number of n X 4 0..2 arrays with horizontal differences mod 3 never 1, vertical differences mod 3 never -1, and rows and columns lexicographically nondecreasing.

Conjectures from _Colin Barker_, Sep 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008098 | OEIS-A229442

Number of n X 5 0..2 arrays with horizontal differences mod 3 never 1, vertical differences mod 3 never -1, and rows and columns lexicographically nondecreasing.

Conjectures from _Colin Barker_, Sep 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008099 | OEIS-A229446

Number of 3 X n 0..2 arrays with horizontal differences mod 3 never 1, vertical differences mod 3 never -1, and rows and columns lexicographically nondecreasing.

Conjectures from _Colin Barker_, Sep 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008100 | OEIS-A229447

Number of 4 X n 0..2 arrays with horizontal differences mod 3 never 1, vertical differences mod 3 never -1, and rows and columns lexicographically nondecreasing.

Conjectures from _Colin Barker_, Sep 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008101 | OEIS-A229448

Number of 5 X n 0..2 arrays with horizontal differences mod 3 never 1, vertical differences mod 3 never -1, and rows and columns lexicographically nondecreasing.

Conjectures from _Colin Barker_, Sep 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008102 | OEIS-A229449

Number of 6 X n 0..2 arrays with horizontal differences mod 3 never 1, vertical differences mod 3 never -1, and rows and columns lexicographically nondecreasing.

Conjectures from _Colin Barker_, Sep 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008103 | OEIS-A229450

Number of 7 X n 0..2 arrays with horizontal differences mod 3 never 1, vertical differences mod 3 never -1, and rows and columns lexicographically nondecreasing.

Conjectures from _Colin Barker_, Sep 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008104 | OEIS-A229451

G.f.: exp(Sum_{n>=1} (3*n)!/n!^3 * x^n/n).

The sixth root of the o.g.f. A(x)^(1/6) = 1 + x + 8*x^2 + 101*x^3 + 1569*x^4 + 27445*x^5 + ... appears to have integer coefficients. See A229452. More generally, if A(m,x) := exp(Sum_{n >= 1} (m*n)!/n!^m * x^n/n), m = 1,2,3,..., then it can be shown that the expansion of A(m,x) has integer coefficients. We conjecture that the expansion of A(m,x)^(1/m!) also has integer coefficients. - _Peter Bala_, Feb 16 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008105 | OEIS-A229452

G.f.: exp(Sum_{n>=1} (3*n)!/(3!*n!^3) * x^n/n).

The sequence defined by b(n) = [x^n] A(x)^n for n >= 1 begins [1, 17, 352, 7969, 189876, 4676768, 117905565, 3024222753, 78607893934, 2064924478892, 54710782664836, ...]. We conjecture that b(n) satisfies the supercongruences b(n*p^k) == b(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and all positive integers n and k [added 20 Oct 2024: more generally, for r a positive integer and s an integer we conjecture that the sequence {b(r,s;n) : n >= 1} defined by b(r,s; n) = [x^(r*n)] A(x)^(s*n) satisfies the same supercongruences]. More generally, for a positive integer m, set A_m(x) = exp(Sum_{n >= 1} (m*n)!/(m!*n!^m) * x^n/n) and define a sequence b_m(n) := [x^n] A_m(x)^n for n >=1. Then we conjecture that b_m(n) is an integer sequence satisfying the same congruences. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008106 | OEIS-A229461

Numbers k such that there is no convex pentagon that can be decomposed into k pairwise congruent regular equilateral triangles.

Conjecture: These 59 numbers are all such exceptions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Suggested by Eike Hertel, _Hugo Pfoertner_, Sep 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008107 | OEIS-A229472

Number of defective 4-colorings of an n X 1 0..3 array connected horizontally, antidiagonally and vertically with exactly one mistake, and colors introduced in row-major 0..3 order.

Conjectures from _Colin Barker_, Mar 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008108 | OEIS-A229473

Number of defective 4-colorings of an n X 2 0..3 array connected horizontally, antidiagonally and vertically with exactly one mistake, and colors introduced in row-major 0..3 order.

Conjectures from _Colin Barker_, Sep 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008109 | OEIS-A229483

Squarefree numbers whose indices are lucky numbers.

Originally arose as "Lucky numbers generated from squarefree numbers" under the hypothesis that Ulam's sieve (the one used to produce lucky numbers) ignores the values of the terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_ and _M. F. Hasler_, Sep 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008110 | OEIS-A229484

Pairs of numbers (m,k) such that 2^m/(m+1) < binomial(m,k) <= 2^m/m and k < m/2.

Sun conjectures that there are an infinite number of pairs (m,k) that satisfy this inequality.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Oct 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008111 | OEIS-A229488

Conjecturally, possible differences between prime(k)^2 and the previous prime for some k.

Conjecturally, possible differences between prime(k)^2 and the previous prime for some k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Oct 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008112 | OEIS-A229489

Conjecturally, possible differences between prime(k)^2 and the next prime for some k.

Conjecturally, possible differences between prime(k)^2 and the next prime for some k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Oct 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008113 | OEIS-A229500

Smaller of Fermi-Dirac twin primes (A229064) which are not the smaller of twin primes (A001359).

The sequence conjecturally infinite. All squares of the sequence are 9, 81, 6561,... and have the form 3^(2^k), k>=1. However, we conjecture that there is only a finite number of them, or, the same, there is only a finite number of primes of the form 3^(2^k) + 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008114 | OEIS-A229505

Number of defective 3-colorings of an n X 3 0..2 array connected diagonally and antidiagonally with exactly one mistake, and colors introduced in row-major 0..2 order.

Conjectures from _Colin Barker_, Jun 16 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008115 | OEIS-A229506

Number of defective 3-colorings of an n X 4 0..2 array connected diagonally and antidiagonally with exactly one mistake, and colors introduced in row-major 0..2 order.

Conjectures from _Colin Barker_, Jun 16 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008116 | OEIS-A229570

Primes of form p*q + 30, where p and q are consecutive primes.

Conjecture: The expression p*q + c with p and q consecutive primes and c = 30 generates more primes than any other value of c in the range 1 < c < 100 and p = 48611 which is 5000th prime. Hence, c = 30 is considered for this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _K. D. Bajpai_, Sep 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008117 | OEIS-A229572

Number of defective 4-colorings of an n X 2 0..3 array connected horizontally, vertically, diagonally and antidiagonally with exactly one mistake, and colors introduced in row-major 0..3 order.

Conjectures from _Colin Barker_, Sep 19 2018: (Start)
Conjectured e.g.f.: 2 + (3*x + exp(2*x)*(7*x - 4))/2. - _Stefano Spezia_, Oct 21 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008118 | OEIS-A229581

Number of defective 3-colorings of an n X 3 0..2 array connected horizontally and antidiagonally with exactly one mistake, and colors introduced in row-major 0..2 order.

Conjectures from _Colin Barker_, Sep 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008119 | OEIS-A229582

Number of defective 3-colorings of an n X 4 0..2 array connected horizontally and antidiagonally with exactly one mistake, and colors introduced in row-major 0..2 order.

Conjectures from _Colin Barker_, Sep 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008120 | OEIS-A229597

Decimal expansion of L(e), the limit of iterations of continued fraction transforms of e.

Conjecture: if x is an irrational number between 2 and 3, then L(x) = L(e).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008121 | OEIS-A229600

Number of defective 3-colorings of an n X 2 0..2 array connected horizontally and vertically with exactly two mistakes, and colors introduced in row-major 0..2 order.

Conjectures from _Colin Barker_, Mar 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008122 | OEIS-A229607

Square array read by antidiagonals downwards in which each row starts with the least prime not in a previous row, and each prime p in a row is followed by the greatest prime < 2*p.

Conjectures: (row 1) = A006992, (column 1) = A104272, and for each row r(k), the limit of r(k)/2^k exists. For rows 1 to 4, the respective limits are 0.303976..., 4.249137..., 6.857407..., 12.235210... . Regarding the conjectures above:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008123 | OEIS-A229608

Square array read by antidiagonals downwards: each row starts with the least prime not in a previous row, and each prime p in a row is followed by the least prime > 2*p.

Conjectures: (row 1) = A055496, (column 1) = A193507, and for each row r(k), the limit of r(k)/2^k exists. For rows 1 to 4, the respective limits are 1.569985..., 2.677285..., 8.230592..., 10.709142...; see _Franklin T. Adams-Watters_'s comment at A055496. The above conjecture row 1 = A055496 is true; additionally, row 2 = A065545; row 3 = A065546; the first 5 terms of row 6 are a contiguous subsequence of A064934; and column 1 = A194598. - _Bob Selcoe_, Oct 27 2015; corrected by _Peter Munn_, Jul 30 2017 The conjecture for column 1 is true iff A194598 and A193507 are equivalent. Is this the case? - _Bob Selcoe_, Oct 29 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008124 | OEIS-A229609

Array: each row starts with the least prime not in a previous row, and each prime p in a row is followed by the greatest prime < 3*p.

Conjectures: (row 1) = A126031, (column 1) = A164952, and for each row r(k), the limit of r(k)/3^k exists. For rows 1 to 4, the respective limits are 0.431270..., 0.636059..., 3.229697..., 5.015914... .

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008125 | OEIS-A229610

Array: each row starts with the least prime not in a previous row, and each prime p in a row is followed by the least prime > 3*p.

Conjectures: (row 1) = A076656, (column 1) = A164958, and for each row r(k), the limit of r(k)/3^k exists. For rows 1 to 4, the respective limits are 0.928655..., 1.447047..., 2.038260..., 4.753271... .

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008126 | OEIS-A229616

Expansion of (phi(-q)^3 / phi(-q^3))^2 in powers of q where phi() is a Ramanujan theta function.

Conjecture: -3 A122858(n) - A229616(n) + 4 A282031(n) = 0 for all n. - _Thomas Baruchel_, Jun 23 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Sep 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008127 | OEIS-A229623

Palindromes m such that m - sum_of_digits(m) is also a palindrome.

It is conjectured that a(n) = 10^(n-18) + 1 for all n > 20. - _Derek Orr_, Apr 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Sep 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008128 | OEIS-A229666

Number of defective 4-colorings of an n X 2 0..3 array connected horizontally, antidiagonally and vertically with exactly two mistakes, and colors introduced in row-major 0..3 order.

Conjectures from _Colin Barker_, Sep 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008129 | OEIS-A229681

Number of defective 3-colorings of an n X 4 0..2 array connected diagonally and antidiagonally with exactly two mistakes, and colors introduced in row-major 0..2 order.

Conjectures from _Colin Barker_, Jun 16 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008130 | OEIS-A229749

Number of defective 4-colorings of an n X 2 0..3 array connected horizontally, vertically, diagonally and antidiagonally with exactly two mistakes, and colors introduced in row-major 0..3 order.

Conjectures from _Colin Barker_, Sep 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008131 | OEIS-A229750

Number of defective 4-colorings of an n X 3 0..3 array connected horizontally, vertically, diagonally and antidiagonally with exactly two mistakes, and colors introduced in row-major 0..3 order.

Conjectures from _Colin Barker_, Sep 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008132 | OEIS-A229769

Not of the form x^2 + P*y^2 for integers x > 0, y > 1, P a prime.

There are no terms <= 10^9 greater than a(3036) = 69808035, leading to the conjecture that the sequence is finite. If true, every sufficiently large number is expressible as x^2 + P*y^2 with x > 0, y > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chris Boyd_, Sep 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008133 | OEIS-A229775

Number of 3 X 3 0..n arrays with rows in lexicographically nondecreasing order and columns in nonincreasing order.

Conjectures from _Colin Barker_, Sep 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008134 | OEIS-A229795

Number of 2 X 2 0..n arrays with rows and columns in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008135 | OEIS-A229796

Number of 3 X 3 0..n arrays with rows and columns in lexicographically nondecreasing order.

Conjectures from _Colin Barker_, Sep 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008136 | OEIS-A229816

Number of partitions of n such that if the length is k then k is not a part.

Conjecture: Also, for n>=1, a(n-1) is the total number of distinct parts of each partition of 2n with partition rank n. - _George Beck_, Jun 23 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Sep 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008137 | OEIS-A229826

Evil (A001969) numbers divisible by 7 but not divisible by 3.

By the Moser-Newman phenomenon, among the first N positive integers divisible by 3, the evil numbers are always in the majority. But what happens if we remove from the positive numbers the multiples of 3? We conjecture that in this case we obtain another phenomenon: among the first N such positive integers divisible by 7, the odious numbers (A000069) are always in the majority.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008138 | OEIS-A229832

First term of smallest sequence of n consecutive weak primes.

Erdős called a weak prime A051635 an "early prime," defined to be one which is less than the arithmetic mean of the prime before it and the prime after it. He conjectured that there are infinitely many consecutive pairs of early primes, and offered $100 for a proof and $25000 for a disproof. See Kuperberg 1992. I make the stronger conjecture that the sequence a(n) is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Oct 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008139 | OEIS-A229835

Number of ways to write n = (p - 1)/6 + q, where p is a prime, and q is a term of the sequence A000009.

Conjecture: a(n) > 0 for all n > 1. Also, any integer n > 1 can be written as (p + 1)/6 + q, where p is a prime and q is a term of A000009.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008140 | OEIS-A229878

a(n) = number of undirected circular permutations tau of 1, ..., (p_n-1)/2 such that tau(1)^2 + tau(2)^2, tau(2)^2 + tau(3)^2, ..., tau((p_n-3)/2)^2 + tau((p_n-1)/2)^2, tau((p_n-1)/2)^2 + tau(1)^2 give all the (p_n-1)/2 quadratic residues modulo p_n, where p_n is the n-th prime.

Conjecture: a(n) > 0 for all n > 6. In other words, for any prime p = 2*n+1 > 13, there is a circular permutation a_1, ..., a_n of the n quadratic residues modulo p such that a_1+a_2, a_2+a_3, ..., a_{n-1}+a_n, a_n+a_1 give all the n quadratic residues modulo p. Zhi-Wei Sun also made the following general conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008141 | OEIS-A229899

a(n) = |{0<g<p_n: g-1, g, g+1 are primitive roots mod p_n}|, where p_n denotes the n-th prime.

Conjecture: a(n) > 0 except for n = 1, 2, 3, 4, 6, 10, 18. In other words, for any prime p > 7 not equal to 13 or 29 or 61, there are three consecutive integers which are primitive roots modulo p. Zhi-Wei Sun also made the following conjectures: The conjecture that a(n) > 0 except for n = 1, 2, 3, 4, 6, 10, 18 follows from Theorem 1 of Cohen, Oliveira e Silva and Trudgian (2015). For an odd prime power q, their theorem states that F_q contains three consecutive primitive elements if and only if q is not one of 3, 5, 7, 9, 13, 25, 29,

61, 81, 121, 169. The primes in this list are 3, 5, 7, 13, 29 and 61, namely p_2, p_3, p_4, p_6, p_10 and p_18. The remaining exception n = 1 (p = 2) is immediate, since at least one of any three consecutive integers is divisible by 2. - _Eiver Rodríguez_, Sep 05 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008142 | OEIS-A229927

Smallest k >= n such that (4^n-1)*2^k - 1 is prime.

Conjecture: there is no "Riesel" number of the form 4^n-1; that is, a(n) exists for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Oct 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008143 | OEIS-A229989

Number of primes in the interval [floor(n/2), floor(3n/2)].

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008144 | OEIS-A230000

Triangular array read by rows: row n shows the coefficients of the polynomial u(n) = c(0) + c(1)*x + ... + c(k)*x^k which is the numerator of the n-th convergent of the continued fraction [1, 1/x, 1/x^2, ... ,1/x^n].

In the Name section, k = n(n+1)/2. For the denominator polynomials, see A230001. Conjecture: every nonnegative integer occurs infinitely many times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008145 | OEIS-A230001

Triangular array read by rows: row n shows the coefficients of the polynomial v(n) = d(0) + d(1)*x + ... + d(k)*x^k which is the denominator of the n-th convergent of the continued fraction [1, 1/x, 1/x^2, ... ,1/x^n].

In the Name section, k = n(n+1)/2 if n is even, and k = (n-1)(n+2)/2 if n is odd. For the numerator polynomials, see A230000. Conjecture: every nonnegative integer occurs infinitely many times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008146 | OEIS-A230022

a(n) = |{the number of primes in the interval (k*n, (k+1)*n]: k = 0, 1, ..., n-1}|.

Conjecture: (i) a(n) is at least sqrt(n-1) for each n > 0, and equality holds only when n is 2 or 26. We have verified part (i) of the conjecture for n up to 10000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008147 | OEIS-A230030

Numbers k not divisible by 5 such that k^2 written in base 5 has no digit > 1.

Conjecture: a(k) = P(5^(k-2)) for every k > 5. - _David Radcliffe_, Sep 14 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Radcliffe_, May 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008148 | OEIS-A230044

Nonnegative numbers k such that k plus a perfect square is a triangular number.

Conjecture: these are the sorted distinct terms of A064784.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Oct 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008149 | OEIS-A230073

Coefficients of the minimal polynomials of the algebraic numbers sqLhat(2*l) from A230072, l >= 1, related to the square of all length in a regular (2*l)-gon inscribed in a circle of radius 1 length unit.

Mustonen's polynomials from his eq. (6) for the even n case coincide with PsqL(l, x) precisely for l = 2^k, k>=1, and for l = 1 (k=0) one has to take the negative. In all other cases the degrees do not fit (Mustonen's polynomials become reducible over the integers). His conjecture for the coefficients can then be rewritten as a conjecture for the present polynomials psqLhat(2^k, x), k >= 0 (see the formula section). S. Mustonen also conjectured about the other zeros of his polynomials. Mustonen's conjecture (adapted, see a comment above) is: a(2^k,m) = ((-1)^m)*binomial(2*2^k,2*m), k >= 1, and for k=0 a(1,0) = -1 and a(1,1) = 1 is trivial.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008150 | OEIS-A230107

Define a sequence by b(1)=n, b(k+1)=b(k)+(sum of digits of b(k)); a(n) is the number of steps needed to reach a term in A004207, or a(n) = -1 if the sequence never joins A004207.

Conjecture: the b sequence, for any starting value n, will eventually merge with one of A000004 (the zero sequence), A004207, A016052 or A016096.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Reinhard Zumkeller_, Oct 15 2013; corrected Oct 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008151 | OEIS-A230113

Digital root of summed Fibonacci and Lucas digital roots indexed by numbers not divisible by 2, 3 or 5.

Conjectures from _Colin Barker_, Sep 22 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary Croft_, Dec 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008152 | OEIS-A230121

Number of ways to write n = x + y + z (0 < x <= y <= z) such that x*(x+1)/2 + y*(y+1)/2 + z*(z+1)/2 is a triangular number.

Conjecture: (i) a(n) > 0 except for n = 1, 2, 4, 5, 7, 12. Moreover, for each n = 20, 21, ... there are three distinct positive integers x, y and z with x + y + z = n such that x*(x+1)/2 + y*(y+1)/2 + z*(z+1)/2 is a triangular number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008153 | OEIS-A230138

List of those primes p with p + 2 and 2*p - 5 both prime.

This sequence is interesting because of the conjectures in the comments in A230140 and A230141.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008154 | OEIS-A230140

Number of ways to write n = x + y + z with 0 < x <= y <= z such that 6*x-1, 6*y-1, 6*z-1 are among those primes p (terms of A230138) with p + 2 and 2*p - 5 also prime.

Conjecture: (i) a(n) > 0 for all n > 2, i.e., 6*n-3 with n > 2 can be expressed as a sum of three terms from A230138. Moreover, for any integer n > 12, there are three distinct positive integers x, y, z with x + y + z = n such that 6*x-1, 6*y-1, 6*z-1 are primes in A230138. Note that part (i) of this conjecture implies that there are infinitely many primes in A230138.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008155 | OEIS-A230147

Record values in A165959.

Because of the bounds on a(n), A182873(n), A192820, and A190661, old conjectures about prime numbers may be proved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John W. Nicholson_, Nov 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008156 | OEIS-A230148

Numbers whose abundancy, sigma(n)/n, is 1024/513.

This list of 7 numbers is conjectured to be complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Oct 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008157 | OEIS-A230179

Number of n X 3 0..2 arrays x(i,j) with each element horizontally or antidiagonally next to at least one element with value 2-x(i,j).

Conjectures from _Colin Barker_, Sep 21 2018: (Start) Conjectured e.g.f.: 6*exp(9*x/2)*sinh(sqrt(65)*x/2)/sqrt(65). - _Stefano Spezia_, Feb 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008158 | OEIS-A230217

List of those primes p with p + 6 and 3*p + 8 also prime.

This sequence is interesting because of the conjecture in the comments in A230219.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008159 | OEIS-A230218

G.f. A(x) satisfies: [x^n] A(x)^(n^2+n+1) = 0 for n>1.

for n>0, a(n) is odd iff n is a power of 2 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008160 | OEIS-A230223

Primes p such that 3*p-4, 3*p-10, and 3*p-14 are all prime.

Conjecture: Any even number greater than 35 can be written as a sum of four terms of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008161 | OEIS-A230224

Number of ways to write 2*n = p + q + r + s with p <= q <= r <= s such that p, q, r, s are primes in A230223.

Conjecture: a(n) > 0 for all n > 17.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008162 | OEIS-A230241

Number of ways to write n = p + q with p, 3*p - 10 and (p-1)*q - 1 all prime, where q is a positive integer.

Conjecture: a(n) > 0 for all n > 5. This implies A. Murthy's conjecture mentioned in A109909. We have verified the conjecture for n up to 10^8. Conjecture verified for n up to 10^9. - _Mauro Fiorentini_, Jul 29 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008163 | OEIS-A230243

Number of primes p < n with 3*p + 8 and (p-1)*n + 1 both prime.

Conjecture: a(n) > 0 for all n > 4. This implies A. Murthy's conjecture (cf. A034693) that for any integer n > 1, there is a positive integer k < n such that k*n + 1 is prime. Conjecture verified for n up to 10^9. - _Mauro Fiorentini_, Sep 21 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008164 | OEIS-A230252

Number of ways to write n = x + y (x, y > 0) with 2*x + 1, x^2 + x + 1 and y^2 + y + 1 all prime.

Conjecture: (i) a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008165 | OEIS-A230254

Number of ways to write n = p + q with p and (p+1)*q/2 + 1 both prime.

Conjecture: a(n) > 0 for all n > 3. We also have some similar conjectures, for example, any integer n > 3 not equal to 17 or 66 can be written as p + q with p and (p+1)*q/2 - 1 both prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008166 | OEIS-A230261

Number of ways to write 2*n - 1 = p + q with p, p + 6 and q^4 + 1 all prime, where q is a positive integer.

Conjecture: (i) a(n) > 0 for all n > 3. Also, any odd number greater than 6 can be written as p + q (q > 0) with p, p + 6 and q^2 + 1 all prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008167 | OEIS-A230267

Coins left after packing 5 curves coins patterns into fountain of coins base n.

G.f.: x*(x^3 - 2*x^2 + 2*x + 1)/((1-x)*(1-x^2)*(1-x^3)) (conjectured). - _Ralf Stephan_, Oct 17 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Oct 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008168 | OEIS-A230269

Number of n X 2 0..2 arrays x(i,j) with each element horizontally or vertically next to at least one element with value (x(i,j)+1) mod 3, and upper left element zero.

Conjectures from _Colin Barker_, Feb 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008169 | OEIS-A230282

Largest k such that (k*n)! >= (k!)^(n+1).

For n > 1, a(n) = floor(e*(n^n) - ((n^2-1)*log(n) + n*(1+log(2*Pi)))/2) [conjectural, but verified for all n in 2..5000]. - _Jon E. Schoenfield_, Oct 22 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Oct 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008170 | OEIS-A230299

Define a sequence b_s by b_s(1)=s, b_s(k+1)=b_s(k)+(sum of digits of b_s(k)); a(n) is the number of steps needed for b_n to reach a term in one of b_0, b_1, b_3 or b_9, or a(n) = -1 if b_n never joins one of these four sequences.

We conjecture that a(n) is never -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_ and _Reinhard Zumkeller_, Oct 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008171 | OEIS-A230312

Squares which cannot be written as the sum of a smaller nonzero square and twice a triangular number.

The conjecture a(n) = A001912(n)^2 (mentioned in the formula part) is easy. In fact, any prime divisor of 4*n^2 + 1 is congruent to 1 modulo 4 and hence it can be written as a sum of two squares. Thus 4*n^2 + 1 = (2*n)^2 + 1^2 is composite if and only if it can be written as a sum of two squares in at least two ways. So the conjecture follows immediately. - _Zhi-Wei Sun_, Feb 23 2014 Conjecture: a(n) = A001912(n)^2, that is, squares of numbers n such that 4n^2 + 1 is prime. - _Alonso del Arte_, Dec 20 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kieren MacMillan_, Dec 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008172 | OEIS-A230351

Number of ordered ways to write n = p + q (q > 0) with p, 2*p^2 - 1 and 2*q^2 - 1 all prime.

Conjecture: a(n) > 0 for all n > 3. Conjecture verified for n up to 10^9. - _Mauro Fiorentini_, Aug 07 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008173 | OEIS-A230360

a(n) is the number of base-4 n-digit numbers requiring only binary digits in bases 3 and 4.

0 is included to mesh with the earlier A146025 ({0, 1, 82000}: the values in decimal with top base 4 here replaced by 5, conjectured complete). It appears unlikely, empirically, that this sequence has a last positive term, and a heuristic approximation of terms is likely not difficult.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James G. Merickel_, Oct 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008174 | OEIS-A230362

Least prime p with 2*p^2 - 1 and 2*(n-p)^2 -1 both prime, or 0 if such a prime p does not exist.

Conjecture: 0 < a(n) < sqrt(2n)*(log n) except for n = 1, 2, 3, 232, 1478, 6457. By the conjecture in the comments in A230351, 0 < a(n) < n for all n > 3. Conjecture verified for n up to 10^9. - _Mauro Fiorentini_, Sep 22 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008175 | OEIS-A230366

a(n) = Sum_{k=1..floor(n/2)} (k^2 mod n).

a(26) and a(27) are both squares. Conjecture: the number of n such that a(n) and a(n+1) are both squares is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Oct 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008176 | OEIS-A230368

A strong divisibility sequence associated with the algebraic integer 1 + i.

Let alpha be an algebraic integer and define a sequence of integers a(n) by the condition a(n) = max { integer d : alpha^n == 1 (mod d)}. Silverman shows that a(n) is a strong divisibility sequence, that is gcd(a(n), a(m)) = a(gcd(n, m)) for all n and m in N; in particular, if n divides m then a(n) divides a(m). For the present sequence we take alpha = 1 + i. For other examples see A230369, A235450 and (conjecturally) A082630.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jan 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008177 | OEIS-A230369

A strong divisibility sequence associated with the algebraic integer 2 + i.

Let alpha be an algebraic integer and define a sequence of integers a(n) by the condition a(n) = max {integer d : alpha^n == 1 (mod d)}. Silverman shows that a(n) is a strong divisibility sequence, that is, gcd(a(n), a(m)) = a(gcd(n, m)) for all n and m in N; in particular, if n divides m then a(n) divides a(m). For the present sequence we take alpha = 2 + i. For other examples see A230368, A235450 and (conjecturally) A082630.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jan 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008178 | OEIS-A230370

Voids left after packing 3 curves coins patterns (3c3s type) into fountain of coins base n.

G.f.: x^3*(11*x^8 - 5*x^7 - 21*x^6 + 6*x^5 + 9*x^4 + x^2 + 3*x + 3)/((1-x)*(1-x^2)^2) (conjectured). _Ralf Stephan_, Oct 19 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Oct 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008179 | OEIS-A230494

Number of ways to write n = x^2 + y (x, y >= 0) with 2*y^2 - 1 prime.

Conjecture: (i) a(n) > 0 for all n > 1. Moreover, if n > 1 is not among 2, 69, 76, then there are positive integers x and y such that x^2 + y is equal to n and 2*y^2 - 1 is prime. We have verified the conjecture for n up to 2*10^7. Both conjectures verified for n up to 10^9. - _Mauro Fiorentini_, Aug 08 2023 See also A230351 and A230493 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008180 | OEIS-A230495

a(n) is the minimal odd evil k, such that k^i, i=1,2,...,n, all are evil, and a(n)=0, if there is no such k.

We conjecture that a(n)>0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Oct 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008181 | OEIS-A230496

a(n) is the minimal odd odious k > 1, such that k^i, i=1,2,...,n, all are odious, or a(n)=0, if there is no such k.

Conjecture: for all n, a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Oct 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008182 | OEIS-A230497

a(n), n>=2, is the minimal odd evil k, such that k^i, i=2,3,...,n, all are odious, and a(n)=0, if there is no such k.

Conjecture: For all n, a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Oct 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008183 | OEIS-A230498

a(n) is the minimal odd odious k>1, such that k^i, i=2,...,n, all are evil, and a(n)=0, if there is no such k.

Conjecture: For all n, a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Oct 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008184 | OEIS-A230506

Numbers n with 2*n + 1 and 2*n^3 + 1 both prime.

This sequence is interesting because of the conjecture in the comments in A230507.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008185 | OEIS-A230516

Number of ways to write n = a + b + c with 0 < a <= b <= c such that {a^2+a-1, a^2+a+1}, {b^2+b-1, b^2+b+1}, {c^2+c-1, c^2+c+1} are twin prime pairs.

Conjecture: a(n) > 0 for all n > 5. Conjecture verified for n up to 10^9. - _Mauro Fiorentini_, Sep 22 2023 See also A230514 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008186 | OEIS-A230546

Least positive integer k <= n such that 2*k^2-1 is a prime and n - k is a square, or 0 if such an integer k does not exist.

By the conjecture in A230494, we should have a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008187 | OEIS-A230548

Twin hearts patterns packing into n X n coins.

G.f.: x^2 * (x^10 + x^8 + 2*x^5 + 3*x^4 + 2*x^3 + 2*x^2 + x)/((1+x^3) * (1-x^3)^2 * (1-x^2)) (conjectured). - _Ralf Stephan_, Oct 30 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Oct 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008188 | OEIS-A230549

Coins left after packing twin hearts patterns into n X n coins.

G.f.: x^2 * (-3*x^10 - 4*x^8 + 3*x^7 + 8*x^6 + 4*x^5 - x^4 + 4*x^3 + 4*x^2 + 5*x + 4)/(1+x^3)*(1-x^3)^2*(1-x^2). (conjectured). - _Ralf Stephan_, Oct 30 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Oct 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008189 | OEIS-A230550

Voids left after packing twin hearts patterns into n X n coins.

G.f.: x^2 * (-2*x^10 + x^9 + 2*x^8 + 8*x^7 + 11*x^6 + 8*x^5 + 6*x^4 + 7*x^3 + 4*x^2 + 2*x + 1)/((1+x^3)*(1-x^3)^2*(1-x^2)) (conjectured). - _Ralf Stephan_, Oct 30 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Oct 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008190 | OEIS-A230578

Smallest number larger than 1 such that multiplying n-th highly composite number by a(n) will give a highly composite number.

Conjecture: all a(n) are prime numbers. This conjecture is false: a(643) = 6 is not prime. - _Alois P. Heinz_, Oct 23 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Oct 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008191 | OEIS-A230579

a(n) = 2^n mod 341.

Granted that without a computer, the task of calculating 2^340 - 1 and dividing by 341 is tedious and error-prone, thus discouraging the discovery of that number as a counterexample to the so-called Chinese hypothesis.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alonso del Arte_, Oct 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008192 | OEIS-A230596

Number of ways to write n = x + y + z with 0 < x <= y <= z such that x*y*z is a triangular number, and that x is a triangular number of the form (p^2 - 1)/8 with p an odd prime.

Conjecture: (i) a(n) > 0 except for n = 1, 2, 4, 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008193 | OEIS-A230667

Smallest starting number for n consecutive odd numbers that are primes or pseudoprime (base 2).

It is conjectured that the sequence is full, that is, a(6) does not exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shyam Sunder Gupta_, Oct 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008194 | OEIS-A230668

Smallest of 5 consecutive odd numbers that are primes or pseudoprime (base 2).

It is conjectured that all numbers in this sequence ends in 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shyam Sunder Gupta_, Oct 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008195 | OEIS-A230718

Smallest n-th power equal to a sum of some consecutive, immediately preceding, positive n-th powers, or 0 if none.

The Erdos-Moser equation is the case k = 1. They conjecture that the only solution is m = n = 1. Any counterexample would be a case of a(n) > 0 with n > 3. And such a case with k = 1 would be a counterexample to the Erdos-Moser conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Oct 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008196 | OEIS-A230747

Number of ways to write n = x + y + 2*z with 0 < x <= y and z > 0 such that x^2 + y^2 + 2*z^2 is a square.

Conjecture: a(n) > 0 for all n > 17. See also A230121 for a related conjecture. The conjecture was confirmed by Chao Haung and Zhi-Wei Sun in 2021. - _Zhi-Wei Sun_, May 09 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008197 | OEIS-A230762

List of commonest number of decompositions of 2k into an unordered sum of two odd primes in range 3 <= k <= m, integer m >= 3, where m is explained below.

Hypothesis: With the increase of m, the commonest number of decompositions of 2n into an unordered sum of two odd primes in the range of 3 <= k <= m ascends. This hypothesis derives that the corresponding smallest m to the terms of this sequence makes an ascending sequence. Or say, when testing with m ascending, once a number a(n) enters this sequence, no number smaller than a(n) will be able to enter this sequence if they had not enter previous.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Oct 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008198 | OEIS-A230811

Smallest starting number for n consecutive odd numbers that are primes or strong pseudoprime (base 2).

It is conjectured that there can be at most one more term, a(5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shyam Sunder Gupta_, Oct 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008199 | OEIS-A230822

a(n) is the smallest integer that makes A230762(n) the commonest number of decompositions of 2k into an unordered sum of two odd primes, where 3 <= k <= a(n).

Hypothesis: With the increase of m, the commonest number of decompositions of 2k into an unordered sum of two odd primes in the range of 3 <= k <= m ascends. This hypothesis derives that a(n) is an ascending sequence. Or say, a(n+1) > a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Oct 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008200 | OEIS-A230823

Number of modified skew Dyck paths of semilength n.

G.f.: 1/(1 - x/(1 - (x + x^2)/(1 - (x + x^2 + x^3)/(1 - (x + x^2 + x^3 + x^4)/(1 - ...))))), a continued fraction (conjecture). - _Ilya Gutkovskiy_, Jun 08 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Scambler_ and _Alois P. Heinz_, Oct 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008201 | OEIS-A230961

Boustrophedon transform of factorials beginning with 1, cf. A000142.

E.g.f.: conjecture: (tan(x)+sec(x))/(1-2*x+x^2) = (1- 12*x/(Q(0)+6*x+3*x^2))/(1-x)^2, where Q(k) = 2*(4*k+1)*(32*k^2+16*k - x^2-6) - x^4*(4*k-1)*(4*k+7)/Q(k+1) ; (continued fraction). - _Sergei N. Gladkovskii_, Nov 18 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Nov 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008202 | OEIS-A231103

Number of n X 3 0..3 arrays x(i,j) with each element horizontally or antidiagonally next to at least one element with value (x(i,j)+1) mod 4, no adjacent elements equal, and upper left element zero.

Conjectures from _Colin Barker_, Sep 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008203 | OEIS-A231104

Number of n X 4 0..3 arrays x(i,j) with each element horizontally or antidiagonally next to at least one element with value (x(i,j)+1) mod 4, no adjacent elements equal, and upper left element zero.

Conjectures from _Colin Barker_, Sep 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008204 | OEIS-A231118

Numbers m such that m*91^k is palindromic for k=0,...,4.

It is an open question whether there exists a,r>1 such that a*r^k is palindromic for k=0,...,5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Nov 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008205 | OEIS-A231119

Least positive k such that n * k^k + 1 is a prime, or 0 if no such k exists.

The number a(5) is conjectured to be zero. Four days of computation have shown that all numbers 5*k^k+1 are composite for k = 1..22733. - _T. D. Noe_, Nov 11 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Nov 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008206 | OEIS-A231123

Array T(n,k) read by antidiagonals: T(n,k) = sum(i=0...n, (-1)^(n+i) * C(n+i,2i) * n/(2i+1) * k^(2i+1)), n>0, k>1.

The polynomial x^(4n+2) - T(n,k)*x^(2n+1) + 1 is reducible. Example: x^10-123x^5+1=(x^2-3x+1)(x^8+3x^7+8x^6+21x^5+55x^4+21x^3+8x^2+3x+1). It is conjectured that for prime p=2n+1, these are the only values where this holds.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Nov 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008207 | OEIS-A231168

Number of ways to write n = x + y + z (x, y, z > 0) such that x^2 + y^2 + z^2 + z is a square, and 6*x + 1, 6*y - 1, 6*z -1 are all prime.

Conjecture: a(n) > 0 for all n > 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008208 | OEIS-A231270

Irregular table read by rows r=0,1,2..., which contain the list of numbers whose (American) English name has no letter in common with that of r.

I conjecture that the table is finite and ends with row 60000000000000000000006006000006 having the entry [3], see examples and the (supposed) complete list given in the links section.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Nov 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008209 | OEIS-A231274

G.f.: Sum_{n>=0} x^n * Product_{k=1..n} (k + x) / (1 - k*x - x^2).

G.f. (conjecture): 1/2 + (1/2)*Sum_{n >= 0} (2*x)^n * Product_{k = 1..n} (k + x)/(1 + k*x). - _Peter Bala_, Jul 06 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008210 | OEIS-A231280

Number of n X 3 0..3 arrays x(i,j) with each element horizontally or antidiagonally next to at least one element with value (x(i,j)+1) mod 4, and upper left element zero.

Conjectures from _Colin Barker_, Sep 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008211 | OEIS-A231317

Number of (n+1) X (1+1) 0..2 arrays with no element equal to a strict majority of its diagonal and antidiagonal neighbors, with values 0..2 introduced in row major order.

Conjectures from _Colin Barker_, Mar 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008212 | OEIS-A231329

Numbers n such that (19^n + 4^n)/23 is prime.

Conjecture: All terms are prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008213 | OEIS-A231364

Numbers n such that antisigma(n) = antisigma(k) has solution for distinct numbers n and k.

Antisigma(n) = A024816(n) = the sum of the non-divisors of n that are between 1 and n. a(9) > 10^5. Conjecture: a(9) = 29886159, a(10) = 29886160.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008214 | OEIS-A231366

Number of numbers whose sum of non-divisors (A024816) is equal to n.

I checked this, and thus Krizek's conjecture below, up to 4*10^19. (End) Conjecture: max a(n) = 2. Conjecture: a(n) = 2 iff n is a number from A225775 (0, 9, 36844389, 129894940, 446591224981504, ...).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008215 | OEIS-A231431

Evil squares.

Conjecture: a(n) ~ 4n^2. - _Charles R Greathouse IV_, Nov 20 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Nov 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008216 | OEIS-A231516

Number of ways to write n = x + y with 0 < x <= y such that x!*y + 1 is prime.

Conjecture: a(n) > 0 for all n > 1. Also, any integer n > 3 can be written as x + y with 0 < x <= y such that x!*y -1 is prime. We have verified the conjecture for n up to 10^6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008217 | OEIS-A231531

Imaginary part of Product_{k = 1..n} (k + i), i = sqrt(-1).

We conjecture that the set of type 2 primes consists of p = 2 and all primes of the form p == 1 (mod 4). See A002144. Moll's conjecture 5.5 about type 2 primes extends to this sequence and takes the form: We conjecture that the set of type 3 primes is A002145, primes of the form 4*k + 3. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Nov 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008218 | OEIS-A231548

Numbers n such that 2*n - 1 < sigma(n) - sigma(n-2).

Conjecture: there are no numbers n such that antisigma(n) < antisigma(n-3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008219 | OEIS-A231555

Number of ways to write n = x + y (x, y > 0) with x*(x + 1) + F(y) prime, where F(y) denotes the y-th Fibonacci number (A000045).

Conjecture: (i) a(n) > 0 for all n > 1. Also, any integer n > 1 can be written as x + y (x, y > 0) with x + F(y) prime. We also have similar conjectures involving some second-order recurrences other than the Fibonacci sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 10 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008220 | OEIS-A231557

Least positive integer k <= n such that 2^k + (n - k) is prime, or 0 if such an integer k does not exist.

Conjecture: a(n) > 0 for all n > 0. See also part (i) of the conjecture in A231201.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008221 | OEIS-A231561

Number of ways to write n = x + y with 0 < x <= y such that 2^x * y + 1 is prime.

Conjecture: (i) a(n) > 0 for all n > 1. Also, any integer n > 1 can be written as x + y (x, y > 0) with 2^x * y^2 + 1 prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008222 | OEIS-A231569

Composite numbers n such that lambda(n) divides 2n-2, where lambda is the Carmichael lambda function (A002322).

Conjecture: the relative asymptotic density of the Carmichael numbers in this sequence exists, is positive and smaller than 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Nov 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008223 | OEIS-A231573

Composite numbers n such that lambda(n) divides 6n-6, where lambda is the Carmichael lambda function (A002322).

Conjecture: the relative asymptotic density of Carmichael numbers in this sequence exists, is positive and smaller than 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Nov 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008224 | OEIS-A231577

Number of ways to write n = x + y (x, y > 0) with 2^x + y*(y+1)/2 prime.

Conjecture: a(n) > 0 for all n > 1. See also A231201, A231555 and A231561 for other similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008225 | OEIS-A231619

E.g.f. A(x) satisfies: A'(x) = A(x*A'(x)) with A(0)=1.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008226 | OEIS-A231631

Least positive integer k < n with k!*(n-k) + 1 prime, or 0 if such an integer k does not exist.

Conjecture: 0 < a(n) < sqrt(n)*(log n) for all n > 2. See also the conjecture in A231516.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008227 | OEIS-A231633

Number of ways to write n = x + y (x, y > 0) with x^2 * y - 1 prime.

Conjectures: Both parts of conjecture (i) verified for n up to 10^9. Conjecture (ii) and both parts of conjecture (iii) verified for n up to 10^7. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008228 | OEIS-A231635

Number of ways to write n = x + y with 0 < x <= y such that lcm(x, y) + 1 is prime.

Conjecture: (i) a(n) > 0 for all n > 1. Also, any integer n > 3 can be written as x + y (x, y > 0) with lcm(x, y) - 1 prime. Both parts of conjecture (i) verified for n up to 10^9. Both parts of conjecture (ii) verified for n up to 10^6. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008229 | OEIS-A231692

Define a sequence of rationals by f(0)=0, thereafter f(n)=f(n-1)-1/n if that is >= 0, otherwise f(n)=f(n-1)+1/n; a(n) = numerator of f(n).

It is conjectured that the terms of the {f(n)} sequence are distinct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 15 2013, Nov 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008230 | OEIS-A231725

Least positive integer k < n such that n + k + 2^k is prime, or 0 if such an integer k does not exist.

Conjecture: a(n) > 0 for all n > 3. We have verified this for n up to 2*10^6; for example, we find the following relatively large values of a(n): a(65958) = 37055, a(299591) = 51116, a(295975) = 13128, a(657671) = 25724, a(797083) = 44940, a(1278071) = 24146, a(1299037) = 34502, a(1351668) = 25121, a(1607237) = 34606, a(1710792) = 11187, a(1712889) = 18438. I conjecture the opposite. In particular I expect that a(n) = 0 for infinitely many values of n. - _Charles R Greathouse IV_, Nov 13 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008231 | OEIS-A231817

Multiplicative digital root of concatenation of all divisors of n (A037278).

Conjecture: a(n) = 0 for almost all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008232 | OEIS-A231883

Number of ways to write n = x + y (x, y > 0) with x^2 + (n-2)*y^2 prime.

Conjecture: (i) a(n) > 0 for all n > 2, and a(n) = 1 only for n = 8, 12, 16. Moreover, if m and n are positive integers with m >= max{2, n-1} and gcd(m, n+1) = 1, then x^2 + n*y^2 is prime for some positive integers x and y with x + y = m, except for the case m = n + 3 = 29.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008233 | OEIS-A231897

a(n) = smallest m such that wt(m^2) = n (where wt(i) = A000120(i)), or -1 if no such m exists.

Conjecture: a(n) is never -1. (It seems likely that the arguments of Lindström (1997) could be modified to establish this conjecture.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008234 | OEIS-A231898

a(n) = smallest k with property that for all m >= k, there is a square N^2 whose binary expansion contains exactly n 1's and m 0's; or -1 if no such k exists.

Conjecture: Apart from n=1, 2 and 4, no other a(n) is -1. See A214560 for a related conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008235 | OEIS-A231899

E.g.f. A(x) satisfies: A'(x) = A(x*A'(x)^3) with A(0)=1.

CONJECTURES.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008236 | OEIS-A231903

Even numbers which are the sum of two different primes > 3 in exactly one way.

Conjecture: there are no more numbers > 62 in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Nov 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008237 | OEIS-A231978

Numbers the squares of which are in A231558.

Prime p is in the sequence, if and only if p, p^2, p^3, p^4, p^2+1 and p^2+p all are odious (A000069). We conjecture that the sequence contains also composite numbers, but the first one should be very large.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Nov 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008238 | OEIS-A232091

Smallest square or promic (oblong) number greater than or equal to n.

It appears that a(n) = A216607(n) + n. (Verified for all n<10^9 by _Lars Blomberg_, Jan 09 2014.) This conjecture now follows from a proof given by _David Applegate_, Jan 10 2014 (see [Applegate]).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Nov 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008239 | OEIS-A232109

Least prime p < n + 5 with n + (p-1)*(p-3)/8 prime, or 0 if such a prime p does not exist.

Conjecture: a(n) > 0 for all n > 0. Moreover, for any integer n > 1 there exists a prime p < 2*sqrt(n)*log(7n) such that n + (p-1)*(p-3)/8 is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008240 | OEIS-A232167

Number of composite integers k less than 10^n such that lambda(k) divides 2k-2, where lambda is the Carmichael lambda function (A002322).

Conjecture: A055553(n)/a(n) has a limit strictly smaller than 1 as n tends to infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Mar 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008241 | OEIS-A232186

Number of ways to write n = p + q (q > 0) with p and p^3 + n*q^2 both prime.

Conjecture: a(n) > 0 for all n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008242 | OEIS-A232193

Numerators of the expected value of the length of a random cycle in a random n-permutation.

a(n) = numerator(Sum_{k=0..n} A002657(k)/A091137(k)) (conjectured). - _Michel Marcus_, Jul 19 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Critzer_, Nov 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008243 | OEIS-A232194

Number of ways to write n = x + y (x, y > 0) with n*x + y and n*y - x both prime.

Conjecture: (i) a(n) > 0 for all n > 2. Also, a(n) = 1 only for n = 3, 4, 6, 20, 24. See also A232174 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008244 | OEIS-A232210

Let b_k=3...3 consist of k>=1 3's. Then a(n) is the smallest k such that the concatenation prime(n)b_k is prime, or a(n)=0 if there is no such prime.

Conjecture: for n>=3, a(n)>0. Counter-conjecture: a(2889)=0. - _Hans Havermann_, Oct 15 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008245 | OEIS-A232239

Lesser of twin-bin primes: primes p such that p+2, x and y are primes, where x is concatenation of binary representations of p and p+2, and y is concatenation of binary representations of p+2 and p: x = p * 2^A070939(p+2) + p+2, y = (p+2) * 2^A070939(p) + p.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Nov 20 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008246 | OEIS-A232248

Denominators of the expected length of a random cycle in a random permutation.

a(n) = denominator(Sum_{k=0..n} A002657(k)/A091137(k)) (conjectured). - _Michel Marcus_, Jul 19 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Critzer_, Nov 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008247 | OEIS-A232269

Number of ways to write 2*n + 1 = x + y (x, y > 0) with x^3 + y^2 and x^2 + y^2 both prime.

Conjecture: a(n) > 0 for all n > 0. Also, any odd integer greater than one can be written as x + y (0 < x < y) with x^3 + y^2 prime. The conjecture implies that there are infinitely many primes of the form x^3 + y^2 (x, y > 0) with x^2 + y^2 also prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008248 | OEIS-A232270

Number of ways to write n = x + y (0 < x <= y) with sigma(x) + phi(y) prime, where sigma(x) is the sum of all (positive) divisors of x and phi is Euler's totient function.

Conjecture: (i) a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 3, 4, 29, 70, 105.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008249 | OEIS-A232271

Start with positive integers (A000027), and at each step n >= 1 subtract from the term at position n + a(n) the value a(n).

The first 71 terms are correct if the following conjecture is true: n+a(n)<=0 or n+a(n)>71 for n >= 2^30.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Nov 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008250 | OEIS-A232353

Number of ways to write n = k + m with k > 0 and m > 0 such that p = prime(k) + phi(m) and p*(p+1) - prime(p) are both prime, where phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for all n > 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008251 | OEIS-A232398

Number of ways to write n = p + (2^k - k) + (2^m - m) with p prime and 0 < k <= m.

Conjecture: a(n) > 0 for all n > 3. This was motivated by A231201. We have verified the conjecture for n up to 2*10^8. It seems that a(n) = 1 for no odd n. On Dec 08 2013, Qing-Hu Hou finished checking the conjecture for n up to 10^10 and found no counterexamples. - _Zhi-Wei Sun_, Dec 08 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008252 | OEIS-A232434

Limit of rows in triangle A232433 when read in reverse order.

Conjecture: a(n) equals sum of f(lambda) over all partitions of n, where f is defined recursively as f({})=1; f(lambda)=binomial(i+j,j) f(mu)f(nu); with i and j the row and column of the box in the Young-Ferrers diagram of lambda such that i+j is maximized, and mu is lambda with the first i rows removed, and nu is lambda with the first j columns removed. See Math Overflow link. - _Wouter Meeussen_, Apr 07 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008253 | OEIS-A232441

Sequence read from antidiagonals of rectangular array given by A(n,k) = 2^(2*k)*(Sum_{j=1..n-floor(n/2)-1} (cos(j*Pi/n))^(2*k)), rows n >= 3, columns k >= 0.

Conjecture: A(n,k) = floor(A198632(n-1,k)/2), n >= 3, k >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Nov 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008254 | OEIS-A232442

a(n) = |{0 < k < n: m*prime(m) - 1 and m*prime(m) + 1 are both prime with m = sigma(k) + phi(n-k)}|, where sigma(k) is the sum of all positive divisors of k and phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for all n > 214. We have verified the conjecture for n up to 10^5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008255 | OEIS-A232443

Number of ways to write n = p + q - pi(q), where p and q are odd primes, and pi(q) is the number of primes not exceeding q.

Conjecture: (i) a(n) > 0 for all n > 3. Moreover, every n = 6, 7, ... can be written as p + q - pi(q) with p, p + 6 and q all prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008256 | OEIS-A232463

Number of ways to write n = p + q - pi(q), where p and q are odd primes not exceeding n, and pi(q) is the number of primes not exceeding q.

Conjecture: a(n) > 0 for all n > 3. Also, a(n) = 1 only for n = 4, 5, 6, 7, 9, 10, 11, 12, 15, 16, 28, 35.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008257 | OEIS-A232502

Number of ways to write n = k + m (0 < k < m) with 2*prime(m) - prime(k) prime.

Conjecture: (i) a(n) > 0 for all n > 4, and a(n) = 1 only for n = 5, 6, 7, 9, 11, 14, 17, 21, 34, 43.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008258 | OEIS-A232504

Number of ways to write n = k + m (k, m > 0) with p(k) + q(m) prime, where p(.) is the partition function (A000041) and q(.) is the strict partition function (A000009).

Conjecture: a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008259 | OEIS-A232540

Numbers n such that (n(n+1)/2) modulo sigma(n) = 1.

Conjecture: supersequence of A112774 (semiprimes of the form 6n+4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008260 | OEIS-A232548

Number of ways to write n = p - pi(p) + 2^k + 2^m with 0 < k <= m, where p is an odd prime and pi(p) is the number of primes not exceeding p.

Conjecture: a(n) > 0 for all n > 4. Qing-Hu Hou has checked the conjecture for n up to 10^7, and found one counterexample: n = 1897048.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008261 | OEIS-A232558

a(1)=5, q=a(n) is the smallest prime > a(n-1) such that q-2*n = p prime.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Nov 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008262 | OEIS-A232559

Sequence (or tree) generated by these rules: 1 is in S, and if x is in S, then x + 1 and 2*x are in S, and duplicates are deleted as they occur.

Conjecture: a(n) = A059894(A348366(n)) for n > 0. - _Mikhail Kurkov_, Jun 14 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008263 | OEIS-A232565

a(n) is the smallest k such that 2^(2^n) - 2^k - 1 is prime, or -1 if no such k exists.

If n > 2, then the (largest) prime P(n) = 2^(2^n)-2^a(n)-1 is a de Polignac number (A065381); i.e., P(n)-2^m is not prime. It seems that if n > 6, then |P(n)-2^m| is composite for every natural m and P(n)*2^m-1 is composite (by the dual Riesel conjecture). So if n > 6, then the prime P(n) may be a Riesel number (A182296). For example, the prime P(7) = 2^(2^7)-(2^18+1) is the first candidate (note that 2^18+1 is the smallest de Polignac number of form 2^k+1). Also, by Crocker's theorem, the smallest number of form 2^(2^n)-2^m-1, namely 2^(2^n-1)-1 is a de Polignac number (A006285) and for n > 6 may be a dual Riesel number (A101036). For example, the double Mersenne prime 2^(2^7-1)-1 probably is a dual Riesel number. It is not known whether these are Riesel numbers with a covering set. - _Thomas Ordowski_, Jan 24 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Nov 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008264 | OEIS-A232616

Least positive integer m such that {2^k - k: k = 1,...,m} contains a complete system of residues modulo n.

Conjectures: Adamczewski disproves Conjecture (i) in linked article, employing a counterexample for n = 550172. - _Michael De Vlieger_, Aug 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008265 | OEIS-A232622

Number of incongruent domino tilings of the 3 X (2n) board.

Conjecture: G.f.: (-1+3*x+4*x^2-10*x^3+4*x^5-x^6) / ((x-1)*(x^2-4*x+1)*(x^4-4*x^2+1)). a(n) = 5a(n-1)-a(n-2)-19a(n-3)+19a(n-4)+a(n-5)-5a(n-6)+a(n-7) for n > 6. - Conjectured by _Jean-François Alcover_, Jan 21 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Nov 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008266 | OEIS-A232624

Coefficient array for the minimal polynomials of 2*cos(2*Pi/n) for n >= 1.

Conjecture: For odd n > 1, the term of the highest degree of (MPR2(2n, x) - MPR2(n, x))/2 is (-1)^omega(n) * x^(phi(n)/2-n/rad(n)) = A076479(n) * x^(A023022(n)-A003557(n)). For example, for n = 15, (MPR2(30, x) - MPR2(15, x))/2 = x^3 - 4x; for n = 105, (MPR2(210, x) - MPR2(105, x))/2 = -x^23 + ...; for n = 225, (MPR2(450, x) - MPR2(225, x))/2 = x^45 + ... If this is true, then for odd n > 1, a(n,A023022(n)-k) = a(2n,A023022(n)-k) = 0 for k = 1, 3, ..., A003557(n)-2; a(n,A023022(n)-A003557(n)) = -A076479(n) and a(2n,A023022(n)-A003557(n)) = A076479(n). - _Jianing Song_, Jul 11 2019 Conjecture: Let MPR2(n, x) equal the odd indexed (n) monic polynomial. If the number of roots with negative signs is even, then n is a term in A014659. Example: n = 7 for x^3 + x^2 - 2x - 1, having two negative roots, (-445041..., and -1.801937...). Two is even so the integer 7 is in A014659. n = 9 for the polynomial x^3 - 3x + 1, with one negative root, (-1.87938). The term 9 is in A014657. - _Gary W. Adamson_, Oct 20 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Nov 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008267 | OEIS-A232636

The second largest value of permanent of (0,1) square matrices of order n with row and column sums equal to 3.

We conjecture that formula below also holds for other values. All terms for n>30 not divisible by 3 are conjectural since our method is based on the following very plausible but yet not proved conjecture with confirms in many cases and without counterexample since 1992 (when it was posed, see ref. [11] in Shevelev link). Let L_n be the set of considered n X n (0,1)-matrices with all row and column sums equal to 3. Let S_n be the subset of completely indecomposable such matrices, i.e., not containing submatrices from L_m (m<n). Let M(n) be the maximal value of permanent on S_n. Then M(n_1+n_2) <= M(n_1)*M(n_2). V. I. Bolshakov proved the result for n divisible by 3, using quite different arguments which, generally speaking, do not work for other n and for third, fourth, etc. largest values of permanent, where our (yet conditional) method does work successfully, at least for sufficiently large n. Conjectures from _Colin Barker_, May 27 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Nov 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008268 | OEIS-A232640

Sequence (or tree) generated by these rules: 1 is in S, and if x is in S, then x + 1 and 2*x + 1 are in S, and duplicates are deleted as they occur.

Conjecture: a(n) = A135533(A003754(n+1)) for n > 0. - _Mikhail Kurkov_, Feb 26 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008269 | OEIS-A232656

The number of pairs of numbers below n that, when generating a Fibonacci-like sequence modulo n, contain zeros.

Conjecture: a(n) = Sum_{d|n} phi(d)*A001177(d), where phi = Euler's totient function (A000010). - _Logan J. Kleinwaks_, Oct 28 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brandon Avila_ and _Tanya Khovanova_, Nov 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008270 | OEIS-A232686

E.g.f. A(x) satisfies: A'(x) = A(x*A'(x))^2.

CONJECTURES.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008271 | OEIS-A232692

E.g.f. satisfies: A(x) = exp(1/A(x)^3 * Integral A(x)^8 dx).

CONJECTURE:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008272 | OEIS-A232694

E.g.f. A(x) satisfies: A'(x) = A(x*A'(x)^4) with A(0)=1.

CONJECTURES.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008273 | OEIS-A232695

E.g.f. A(x) satisfies: A'(x) = A(x*A'(x)^5) with A(0)=1.

CONJECTURES.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008274 | OEIS-A232711

Conjectured list of numbers whose trajectory under the '5x+1' map eventually reaches 1.

Conjectured list of numbers whose trajectory under the '5x+1' map eventually reaches 1. This is conjectural in that there is no known proof that 7, 9, 11, etc. (see A267970) do not eventually cycle. - _N. J. A. Sloane_, Jan 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Perry_, Nov 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008275 | OEIS-A232724

Numbers k satisfying g(k - g(k)) > g(k) = greatest prime divisor of k.

Conjecture: for every positive integer d, there exist infinitely many n for which a(n + 1) - a(n) + d; for d = 1, the first 4 such n are 40, 67, 76, 79.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 11 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008276 | OEIS-A232768

Numbers n with the property that n^2+(n+1)^2 and n^2+(n+1)^2+(n+2)^2 are both prime.

It is an open question whether either of these polynomials produces an infinite number of primes. This sequence lists the values of x that produce a prime in both polynomials. x must be congruent to 0 or 2 (mod 4) and all the generated primes are of the form 4k+1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chris Fry_, Nov 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008277 | OEIS-A232805

a(n) is the length of the n-th chunk of nonprimes in this sequence, the chunks being separated by primes; lexicographically least such sequence starting with a prime.

Another self-describing sequence invented by E. Angelini. Conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_ and _Eric Angelini_, Nov 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008278 | OEIS-A232820

Imaginary part of the sum over the n-th powers of all Gaussian integers in the n X n base square in the first quadrant.

Conjecture: a(4n) = 0. - _Michel Marcus_, Nov 09 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Nov 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008279 | OEIS-A232861

Numbers k with k - 1, k + 1, prime(k) - k, prime(k) + k, k*prime(k) - 1, k*prime(k) + 1 all prime.

Conjecture: (i) This sequence contains infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008280 | OEIS-A232862

Least positive integer m <= n^2/2 + 3 such that the set {prime(k) - k: k = 1,...,m} contains a complete system of residues modulo n, or 0 if such a number m does not exist.

Conjecture: a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008281 | OEIS-A232866

Positions of the nonnegative integers in the sequence (or tree) of complex numbers generated by these rules: 0 is in S, and if x is in S, then x + 1 and i*x are in S, where duplicates are deleted as they occur.

a(n+4) = 2*n^2 + n + 6 for n >= 1 (conjectured). G.f.: x*(-1 + x - x^3 - x^4 - x^5 - x^6)/(x - 1)^3 (conjectured). a(n) = 3*a(n-1) - 3*a(n-2) + a(n-3) for n >= 8 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008282 | OEIS-A232867

Positions of the negative integers in the sequence (or tree) of complex numbers generated by these rules: 0 is in S, and if x is in S, then x + 1 and i*x are in S, where duplicates are deleted as they occur.

a(n+1) = 2*n^2 + n + 9 for n >= 1 (conjectured). G.f.: x*(-8 + 12*x - 7*x^2 - x^3)/(x - 1)^3 (conjectured). a(n) = 3*a(n-1) - 3*a(n-2) + a(n-3) for n >= 5 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008283 | OEIS-A232868

Positions of the integers in the sequence (or tree) of complex numbers generated by these rules: 0 is in S, and if x is in S, then x + 1 and i*x are in S, where duplicates are deleted as they occur.

a(n) = a(n-1) + 2*a(n-2) - 2*a(n-3) - a(n-4) + a(n-5) for n > 10 (conjectured). G.f.: x*(-x^9 - 4*x^7 + 2*x^6 + 2*x^5 - 2*x^4 + x^2 - x - 1)/((x - 1)^3*(x + 1)^2) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 01 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008284 | OEIS-A232890

Denominator of n-th term of sequence (or tree) S of all rational numbers generated by these rules: 0 is in S; if x is in S then x + 1 is in S, and if x + 1 is nonzero, then -1/(x + 1) is in S; duplicates are deleted as they occur.

Conjectures: If b/c is a positive rational number, the position of n + b/c for n >= 0 forms a linear recurrence sequence with signature (1,1), and the position of -n - b/c forms a linear recurrence sequence with signature (4, -4, 1). For n>=1, the numbers -(1 + 1/n) are terminal nodes in the tree, and their positions are linearly recurrent with signature (2,0,-1). For n >=3, the n-th generation g(n) consists of F(n-1) positive numbers and F(n-1) negative numbers, where F = A000045, the Fibonacci numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008285 | OEIS-A232891

Least positive integer m <= n^2/2 + 3 such that {k*prime(k): k = 1,...,m} contains a complete system of residues modulo n, or 0 if such a number m does not exist.

Conjecture: (i) a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008286 | OEIS-A232894

Least positive integer m such that {Catalan(k) - k: k = 1, ..., m} contains a complete system of residues modulo n, or 0 if such a number m does not exist.

Conjecture: (i) Let n be any positive integer. Then 0 < a(n) <= n^2/2 + 7. Also, {Catalan(k) + k: k = 1, ..., [n^2/2] + 23} contains a complete system of residues modulo n, where [.] is the floor function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008287 | OEIS-A232895

Sequence (or tree) S of all positive integers in the order generated by these rules: 1 and 2 are in S; if x is in S then x + 2 and 2*x are in S, where duplicates are deleted as they occur.

Let S be the sequence (or tree) of numbers generated by these rules: 1 and 2 are in S; if x is in S then x + 2 and 2*x are in S, where duplicates are deleted as they occur. Every positive integer occurs exactly once in S, so that S is a permutation of the natural numbers. Deleting duplicates as they occur, the generations of S are given by g(1) = (1,2), g(2) = (3,4), g(3) = (5,6,8), g(4) = (7,10,12,16), ... Concatenating gives 1,2,3,4,5,6,8,... Conjecture: the position of the n-th odd positive integer in S is the linearly recurrent sequence given by A232896(n) for n>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008288 | OEIS-A232896

a(n) = 3*a(n-1) - 2*a(n-2) - a(n-3) + a(n-4), where a(1) = 1, a(2) = 3, a(3) = 5, a(4) = 8.

Conjecture: a(n) is the position of 2*n-1, for n >= 1, in the sequence S = A232895 of positive integers generated by these rules: 1 and 2 are in S; if x is in S then x + 2 and 2*x are in S, where duplicates are deleted as they occur.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008289 | OEIS-A232898

Least positive integer m such that {C(2k,k) + k: k = 1,...,m} contains a complete system of residues modulo n, or 0 if such a number m does not exist.

Conjecture: (i) Let n be any positive integer. Then 0 < a(n) <= n^2/2 + 3. Also, {C(2k,k) - k: k = 1, ..., [n^2/2] + 15} contains a complete system of residues modulo n, where [.] is the floor function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008290 | OEIS-A232921

Number of 2 X n 0..2 arrays with no element x(i,j) adjacent to value 2-x(i,j) horizontally, diagonally or antidiagonally.

Conjectures from _Colin Barker_, Oct 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008291 | OEIS-A232931

The least positive integer k such that Kronecker(D/k) = -1 where D runs through all positive fundamental discriminants (A003658).

With D = A003658(n): Mollin conjectured, and Granville, Mollin, & Williams proved, that for n > 1128, a(n) <= D^0.5 / 2. Treviño proves that for n > 484, a(n) <= D^0.45. Asymptotically the best known upper bound for the exponent is less than 0.16 when D is prime and 1/4 + epsilon (for any epsilon > 0) for general D. - _Charles R Greathouse IV_, Apr 23 2014 (corrected by _Enrique Treviño_, Mar 18 2022)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Finch_, Dec 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008292 | OEIS-A232950

Number of n X 3 0..3 arrays with no element x(i,j) adjacent to value 3-x(i,j) horizontally, diagonally or antidiagonally, and top left element zero.

Conjectures from _Colin Barker_, Oct 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008293 | OEIS-A232951

Number of n X 4 0..3 arrays with no element x(i,j) adjacent to value 3-x(i,j) horizontally, diagonally or antidiagonally, and top left element zero.

Conjectures from _Colin Barker_, Oct 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008294 | OEIS-A232956

Number of 2 X n 0..3 arrays with no element x(i,j) adjacent to value 3-x(i,j) horizontally, diagonally or antidiagonally, and top left element zero.

Conjectures from _Colin Barker_, Oct 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 02 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008295 | OEIS-A232996

a(n) = the constant f(2n+1) appearing in Conjecture 2 of Ahmed et al., 2013.

a(n) = the constant f(2n+1) appearing in Conjecture 2 of Ahmed et al., 2013.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 17 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008296 | OEIS-A233020

Number of n X 2 0..3 arrays with no element x(i,j) adjacent to value 3-x(i,j) horizontally, vertically, diagonally or antidiagonally, and top left element zero.

Conjectures from _Colin Barker_, Oct 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008297 | OEIS-A233044

Pairs p, q for those partial sums p/q of the series e = sum_{n>=0} 1/n! that are not convergents to e.

Sondow (2006) conjectured that 2/1 and 8/3 are the only partial sums of the Taylor series for e that are also convergents to the simple continued fraction for e. Sondow and Schalm (2008, 2010) proved partial results toward the conjecture. Berndt, Kim, and Zaharescu (2012) proved it in full.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Dec 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008298 | OEIS-A233077

Number of n X 3 0..3 arrays with no element x(i,j) adjacent to value 3-x(i,j) horizontally or antidiagonally, top left element zero, and 1 appearing before 2 in row major order.

Conjectures from _Colin Barker_, Oct 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008299 | OEIS-A233078

Number of n X 4 0..3 arrays with no element x(i,j) adjacent to value 3-x(i,j) horizontally or antidiagonally, top left element zero, and 1 appearing before 2 in row major order.

Conjectures from _Colin Barker_, Oct 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008300 | OEIS-A233083

Number of 2 X n 0..3 arrays with no element x(i,j) adjacent to value 3-x(i,j) horizontally or antidiagonally, top left element zero, and 1 appearing before 2 in row major order.

Conjectures from _Colin Barker_, Oct 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 03 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008301 | OEIS-A233094

Number of n X 4 0..3 arrays with no element x(i,j) adjacent to value 3-x(i,j) horizontally, diagonally or antidiagonally, top left element zero, and 1 appearing before 2 in row major order.

Conjectures from _Colin Barker_, Oct 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008302 | OEIS-A233099

Number of 2 X n 0..3 arrays with no element x(i,j) adjacent to value 3-x(i,j) horizontally, diagonally or antidiagonally, top left element zero, and 1 appearing before 2 in row major order.

Conjectures from _Colin Barker_, Oct 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008303 | OEIS-A233106

Number of n X 1 0..5 arrays with no element x(i,j) adjacent to value 5-x(i,j) horizontally or vertically, top left element zero, and 1 appearing before 2 3 and 4, and 2 appearing before 3 in row major order.

Conjectures from _Colin Barker_, Feb 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008304 | OEIS-A233107

Number of n X 2 0..5 arrays with no element x(i,j) adjacent to value 5-x(i,j) horizontally or vertically, top left element zero, and 1 appearing before 2 3 and 4, and 2 appearing before 3 in row major order.

Conjectures from _Colin Barker_, Mar 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008305 | OEIS-A233123

Number of n X 2 0..5 arrays with no element x(i,j) adjacent to itself or value 5-x(i,j) horizontally or vertically, top left element zero, and 1 appearing before 2 3 and 4, and 2 appearing before 3 in row major order (unlabelled 6-colorings with no clashing color pairs).

Conjectures from _Colin Barker_, Mar 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008306 | OEIS-A233124

Number of n X 3 0..5 arrays with no element x(i,j) adjacent to itself or value 5-x(i,j) horizontally or vertically, top left element zero, and 1 appearing before 2 3 and 4, and 2 appearing before 3 in row major order (unlabelled 6-colorings with no clashing color pairs).

Conjectures from _Colin Barker_, Mar 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 04 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008307 | OEIS-A233150

Number of ways to write n = k + m (k, m > 0) with 2^k + prime(m) prime.

Conjecture: a(n) > 0 except for n = 1, 2, 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008308 | OEIS-A233162

Number of n X 1 0..7 arrays with no element x(i,j) adjacent to itself or value 7-x(i,j) horizontally, diagonally, antidiagonally or vertically, top left element zero, and 1 appearing before 2 3 4 5 and 6, 2 appearing before 3 4 and 5, and 3 appearing before 4 in row major order (unlabeled 8-colorings with no clashing color pairs).

Conjectures from _Colin Barker_, Feb 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008309 | OEIS-A233163

Number of n X 3 0..7 arrays with no element x(i,j) adjacent to itself or value 7-x(i,j) horizontally, diagonally, antidiagonally or vertically, top left element zero, and 1 appearing before 2 3 4 5 and 6, 2 appearing before 3 4 and 5, and 3 appearing before 4 in row major order (unlabelled 8-colorings with no clashing color pairs).

Conjectures from _Colin Barker_, Oct 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008310 | OEIS-A233164

Number of n X 4 0..7 arrays with no element x(i,j) adjacent to itself or value 7-x(i,j) horizontally, diagonally, antidiagonally or vertically, top left element zero, and 1 appearing before 2 3 4 5 and 6, 2 appearing before 3 4 and 5, and 3 appearing before 4 in row major order (unlabelled 8-colorings with no clashing color pairs).

Conjectures from _Colin Barker_, Oct 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008311 | OEIS-A233165

Number of n X 5 0..7 arrays with no element x(i,j) adjacent to itself or value 7-x(i,j) horizontally, diagonally, antidiagonally or vertically, top left element zero, and 1 appearing before 2 3 4 5 and 6, 2 appearing before 3 4 and 5, and 3 appearing before 4 in row major order (unlabelled 8-colorings with no clashing color pairs).

Conjectures from _Colin Barker_, Oct 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008312 | OEIS-A233166

Number of n X 6 0..7 arrays with no element x(i,j) adjacent to itself or value 7-x(i,j) horizontally, diagonally, antidiagonally or vertically, top left element zero, and 1 appearing before 2 3 4 5 and 6, 2 appearing before 3 4 and 5, and 3 appearing before 4 in row major order (unlabelled 8-colorings with no clashing color pairs).

Conjectures from _Colin Barker_, Oct 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008313 | OEIS-A233167

Number of n X 7 0..7 arrays with no element x(i,j) adjacent to itself or value 7-x(i,j) horizontally, diagonally, antidiagonally or vertically, top left element zero, and 1 appearing before 2 3 4 5 and 6, 2 appearing before 3 4 and 5, and 3 appearing before 4 in row major order (unlabeled 8-colorings with no clashing color pairs).

Conjectures from _Colin Barker_, Oct 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008314 | OEIS-A233175

Number of 2 X n 0..5 arrays with no element x(i,j) adjacent to itself or value 5-x(i,j) horizontally, diagonally or antidiagonally, top left element zero, and 1 appearing before 2 3 and 4, and 2 appearing before 3 in row major order (unlabelled 6-colorings with no clashing color pairs).

Conjectures from _Colin Barker_, Oct 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008315 | OEIS-A233183

Number of ways to write n = k + m with 0 < k < m such that C(2*k, k) + prime(m) is prime.

Conjecture: a(n) > 0 for all n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008316 | OEIS-A233196

Number of n X 2 0..7 arrays with no element x(i,j) adjacent to itself or value 7-x(i,j) horizontally, antidiagonally or vertically, top left element zero, and 1 appearing before 2 3 4 5 and 6, 2 appearing before 3 4 and 5, and 3 appearing before 4 in row major order (unlabeled 8-colorings with no clashing color pairs).

Conjectures from _Colin Barker_, Oct 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008317 | OEIS-A233204

Number of ways to write n = k + m with 0 < k < m such that 2^k * prime(m) + 3 is prime.

Conjecture: (i) a(n) > 0 for all n > 3. Also, any integer n > 2 can be written as k + m (0 < k <= m) with 2^k*prime(m) - 3 prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008318 | OEIS-A233206

Number of ways to write n = k + m (0 < k <= m) with k! + prime(m) prime.

Conjecture: a(n) > 0 for all n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 05 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008319 | OEIS-A233211

Number of n X 2 0..5 arrays with no element x(i,j) adjacent to value 5-x(i,j) horizontally or antidiagonally, top left element zero, and 1 appearing before 2 3 and 4, and 2 appearing before 3 in row major order.

Conjectures from _Colin Barker_, Oct 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008320 | OEIS-A233212

Number of n X 3 0..5 arrays with no element x(i,j) adjacent to value 5-x(i,j) horizontally or antidiagonally, top left element zero, and 1 appearing before 2 3 and 4, and 2 appearing before 3 in row major order.

Conjectures from _Colin Barker_, Oct 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008321 | OEIS-A233213

Number of n X 4 0..5 arrays with no element x(i,j) adjacent to value 5-x(i,j) horizontally or antidiagonally, top left element zero, and 1 appearing before 2 3 and 4, and 2 appearing before 3 in row major order.

Conjectures from _Colin Barker_, Oct 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008322 | OEIS-A233218

Number of 2 X n 0..5 arrays with no element x(i,j) adjacent to value 5-x(i,j) horizontally or antidiagonally, top left element zero, and 1 appearing before 2 3 and 4, and 2 appearing before 3 in row major order.

Conjectures from _Colin Barker_, Oct 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008323 | OEIS-A233251

Number of n X 3 0..5 arrays with no element x(i,j) adjacent to itself or value 5-x(i,j) horizontally or antidiagonally, top left element zero, and 1 appearing before 2 3 and 4, and 2 appearing before 3 in row major order (unlabelled 6-colorings with no clashing color pairs).

Conjectures from _Colin Barker_, Oct 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008324 | OEIS-A233252

Number of n X 4 0..5 arrays with no element x(i,j) adjacent to itself or value 5-x(i,j) horizontally or antidiagonally, top left element zero, and 1 appearing before 2 3 and 4, and 2 appearing before 3 in row major order (unlabelled 6-colorings with no clashing color pairs).

Conjectures from _Colin Barker_, Oct 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008325 | OEIS-A233257

Number of 2 X n 0..5 arrays with no element x(i,j) adjacent to itself or value 5-x(i,j) horizontally or antidiagonally, top left element zero, and 1 appearing before 2 3 and 4, and 2 appearing before 3 in row major order (unlabelled 6-colorings with no clashing color pairs).

Conjectures from _Colin Barker_, Oct 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008326 | OEIS-A233283

Record values in partial LCM-products (A233287) of Fibonacci entry points (A001177).

Conjecture: sequence gives the positions of records in A233284.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008327 | OEIS-A233293

Smallest number that is the largest value in the Collatz (3x + 1) trajectories of exactly n initial values. (a(n)=0 if no such number exists.)

No number that is the largest value in the 3x + 1 trajectories of exactly 2, 4, 5, 7, or 10 initial values exists, so a(n) = 0 at n = 2, 4, 5, 7, and 10; for all other values of n up to 3000, a(n) > 0. Conjecture: a(n) > 0 for all n > 10. - _Jon E. Schoenfield_, Dec 14 2013

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Dec 06 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008328 | OEIS-A233307

a(n) = |{0 < k < n: p(k)^2 + q(n-k)^2 is prime}|, where p(.) is the partition function (A000041) and q(.) is the strict partition function (A000009).

Conjecture: (i) a(n) > 0 for all n > 1. Also, for any integer n > 4, p(k)*q(n-k) - 1 is prime for some 0 < k < n/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008329 | OEIS-A233319

E.g.f. A(x) satisfies: A'(x) = A(x*A'(x))^3.

CONJECTURES:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008330 | OEIS-A233329

Expansion of (1+4*x+x^2)/((1+x)^2*(1-x)^5).

Conjecture: sequence gives column 1 of A233331 (up to an offset).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Jan 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008331 | OEIS-A233330

Irregular array read by rows: T(n,k) = number of r_{n,k}-cores associated with A233332(n,k), for n>=2, 1<=k<=floor(n/2), explained below.

Conjecture: The entries of column k satisfy sum{h=0..5} ((-1)^h*binomial(5,h)*A(n+5-h,k)) = 0, for appropriate initial conditions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Dec 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008332 | OEIS-A233331

Irregular array read by rows: T(n,k) = number of r_{n,k}-cores associated with A233332(n,k), reduced for symmetry, for n>=2, 1<=k<=floor(n/2), explained below.

Conjecture: G.f. for column 1 is F_1(x) = x^2*(1+4x+x^2)/((1+x)^2*(1-x)^5). Conjecture: Column 1 = A233329, up to an offset. Conjecture: Column k has generating function of the form F_k(x) = G_k(x)/((1+x)^2*(1-x)^5), where G_k(x) is a polynomial in x. Conjecture: A233331(2*M,1) = A076454(M), M>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Jan 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008333 | OEIS-A233332

Irregular array read by rows: A(n,k) = number of first coronas of a fixed rhombus r_{n,k} with characteristics of n-fold rotational symmetry in the Euclidean plane, n>=2, 1<=k<=floor(n/2), as explained below.

Conjecture: For all n and for all k, A(n,k) == n-2 (mod 2^(n-3)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Dec 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008334 | OEIS-A233341

Least number m for which n applications of the mapping r(k) = k - (greatest prime divisor of k) map m to 0.

r(m) = 1 if and only if m = 1 or m is a prime. Conjecture: Every positive integer divides infinitely many terms of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008335 | OEIS-A233346

Primes of the form p(k)^2 + q(m)^2 with k > 0 and m > 0, where p(.) is the partition function (A000041), and q(.) is the strict partition function (A000009).

Conjecture: The sequence contains infinitely many terms. This follows from part (i) of the conjecture in A233307. Similarly, the conjecture in A232504 implies that there are infinitely many primes of the form p(k) + q(m) with k and m positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 07 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008336 | OEIS-A233359

a(n) = |{0 < k < n: L(k) + q(n-k) is prime}|, where L(k) is the k-th Lucas number (A000204), and q(.) is the strict partition function (A000009).

Conjecture: a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008337 | OEIS-A233360

Primes of the form L(k) + q(m) with k > 0 and m > 0, where L(k) is the k-th Lucas number (A000204), and q(.) is the strict partition function (A000009).

Conjecture: The sequence has infinitely many terms. This follows from the conjecture in A233359.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008338 | OEIS-A233386

Number of ways to write n = i + j + k with 0 < i <= j <= k and i, j, k not all equal such that phi(i)*phi(j)*phi(k) is a cube.

Conjecture: For each k = 3, 4, ..., any integer n >= 3*k can be written as n_1 + n_2 + ... + n_k with n_1, n_2, ..., n_k positive and not all equal such that the product phi(n_1)*phi(n_2)*...*phi(n_k) is a k-th power. We have verified this conjecture with k = 3 for n up to 10^5 and with k = 4, 5, 6 for n up to 30000. See also A236998 for a similar conjecture with k = 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008339 | OEIS-A233390

a(n) = |{0 < k < n: 2^k - 1 + q(n-k) is prime}|, where q(.) is the strict partition function (A000009).

Conjecture: a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 08 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008340 | OEIS-A233402

Number of (n+1) X (1+1) 0..2 arrays with row and column sums nondecreasing, and no adjacent elements equal.

Conjectures from _Colin Barker_, Oct 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008341 | OEIS-A233417

a(n) = |{0 < k <= n/2: q(k)*q(n-k) + 1 is prime}|, where q(.) is the strict partition function (A000009).

Conjecture: (i) a(n) > 0 for all n > 1. Similarly, for any integer n > 5, there is a positive integer k < n with q(k)*q(n-k) - 1 prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008342 | OEIS-A233419

Least m such that A232637(m) is divisible by k, where k is the n-th number not divisible by 4; or 0 if no such m exists.

Conjecture: a(n) > 0 for all n. Equivalently, any number not divisible by 4 divides infinitely many members of A232637.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Dec 09 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008343 | OEIS-A233466

Numbers k such that phi(k) = (k-5)/2.

Conjecture (i): There is no solution to the equation phi(x) = (x-1)/2. Conjecture (ii): The sequence has only three terms and a(3) = (2^2^5-5) * (2^2^5-1) = 18446744047939747845. Conjecture (i) is a part of Lehmer's totient problem. Conjecture (ii) is disproved with the term a(3) = 6992962170388485 = 3 * 5 * 17 * 353 * 929 * 83623931. - _Max Alekseyev_, Oct 28 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Dec 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008344 | OEIS-A233513

Triangle read by antidiagonals of the conjectured least index k > 2 of Fibonacci-like sequence f(i+2) = f(i+1) + f(i), with f(1)=m and f(2)=n, such that f(k) is a square, or k=0 if squares do not exist in the corresponding sequence.

Triangle read by antidiagonals of the conjectured least index k > 2 of Fibonacci-like sequence f(i+2) = f(i+1) + f(i), with f(1)=m and f(2)=n, such that f(k) is a square, or k=0 if

squares do not exist in the corresponding sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jan 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008345 | OEIS-A233514

Numbers n such that all numbers of the form Fib(n)/d + d are prime (or nonexistent), where Fib(n) is the n-th Fibonacci number and d is a nontrivial divisor of Fib(n).

Damir et al. conjecture that this sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jan 31 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008346 | OEIS-A233515

Fibonacci numbers whose index is in A233514.

Damir et al. conjecture that this sequence is finite. Note that it is A005478 (prime Fibonacci numbers) and two additional terms (34 and 610) so far.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Jan 31 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008347 | OEIS-A233542

Number of ways to write n = k^2 + m with k > 0 and m > 0 such that phi(k^2)*phi(m) - 1 is prime, where phi(.) is Euler's totient function (A000010).

Conjecture: (i) a(n) > 0 for all n > 5. We have verified part (i) of the conjecture for n up to 2*10^7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008348 | OEIS-A233544

Number of ways to write n = k^2 + m with k > 0 and m >= k^2 such that sigma(k^2) + phi(m) is prime, where sigma(k^2) is the sum of all (positive) divisors of k^2, and phi(.) is Euler's totient function (A000010).

Conjectures: Part (i) of the conjecture is stronger than the conjecture in A232270. We have verified it for n up to 10^8. I verified the conjecture to 3*10^9. The conjecture is almost surely true. - _Charles R Greathouse IV_, Dec 13 2013 There are no counterexamples to conjecture (i) < 5.12 * 10^10. - _Jud McCranie_, Jul 23 2017 The conjectures appeared as Conjecture 3.31 in the linked 2017 paper. - _Zhi-Wei Sun_, Nov 30 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008349 | OEIS-A233549

Number of ways to write n = p + q (q > 0) with p and (phi(p)*phi(q))^4 + 1 prime, where phi(.) is Euler's totient function (A000010).

Conjecture: (i) a(n) > 0 for all n > 2. Part (i) of the conjecture implies that there are infinitely many primes of the form x^4 + 1. We have verified it for n up to 10^7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 12 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008350 | OEIS-A233566

a(n) = |{0 < p < n: p and p*phi(n-p) - 1 are both prime}|, where phi(.) is Euler's totient function (A000010).

Conjecture: a(n) > 0 for all n > 3. Also, for any n > 2 there is a prime p < n with p^2*phi(n-p) - 1 prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008351 | OEIS-A233567

Number of ways to write n = p + q (q > 0) with p and p^4 + phi(q)^4 both prime, where phi(.) is Euler's totient function (A000010).

Conjecture: If n > 2 is not equal to 5, then we have a(n) > 0, also there is a prime p < n with p^2 + phi(n-p)^2 prime. We have verified this for n up to 10^7. The first assertion in the conjecture implies that there are infinitely many primes of the form p^4 + q^4, where p is a prime and q is a positive integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008352 | OEIS-A233574

a(n) is the smallest term of either A233010 or A233572 such that |n-a(n)|<a(n) is a term of one of the two sequences.

It is conjectured that a(n) exists for all n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Dec 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008353 | OEIS-A233578

n >= 2 such that the denominator/6 of Bernoulli(n) is congruent to {1, 5, 7, 13 or 19} modulo 30.

Conjecture: for these and only these n, the absolute value of the numerator of Bernoulli(n) is congruent 1 modulo 6. If my conjecture is true, then you can obtain the residue modulo 6 of the abs. value of Bernoulli numerators by calculating their denominators/6 modulo 30. Program uses the von Staudt-Clausen Theorem. None of these n are in the complementary sequence, A233579 (n such that the denominator/6 of Bernoulli(n) is congruent to {11, 17, 23, 25 or 29} modulo 30). I have checked and verified that, up to n = 50446, the union of A233578 and A233579 is all even numbers >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael G. Kaarhus_, Dec 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008354 | OEIS-A233579

Numbers n such that the denominator/6 of Bernoulli(n) is congruent to {11, 17, 23, 25 or 29} modulo 30.

Conjecture: for these and only these n, the absolute value of the numerator of Bernoulli(n) is congruent 5 modulo 6. If this is true, then you can obtain the residue modulo 6 of the absolute value of Bernoulli numerators by calculating their denominators/6 modulo 30. The program uses the von Staudt-Clausen Theorem. None of these n are in the complementary sequence, A233578 (n >= 2 such that the denominator/6 of Bernoulli_n is congruent to {1, 5, 7, 13 or 19} modulo 30). I have checked and verified that, up to n = 50446, the union of A233578 and A233579 is all even numbers >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael G. Kaarhus_, Dec 13 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008355 | OEIS-A233592

Positive integers k such that the continued fraction expansion sqrt(k) = c(1) + c(1)/(c(2) + c(2)/(c(3) + c(3)/...)) is periodic.

All the listed cases become periodic after just two leading terms (it is a conjecture that this behavior is general); the validity of their expansions was explicitly tested.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Jan 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008356 | OEIS-A233593

Positive integers k such that the continued fraction expansion sqrt(k) = c(1) + c(1)/(c(2) + c(2)/(c(3) + c(3)/....)) is aperiodic.

The cases with known periodic expansions, listed in A233592, all become periodic after just two leading terms. In contrast, the Blazys's expansion of sqrt(a(k)) for every member a(k) of this list remains aperiodic up to at least 1000 terms. It is therefore conjectured, though not proved, that these expansions are indeed aperiodic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Jan 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008357 | OEIS-A233637

Number of (n+1) X (1+1) 0..2 arrays with every 2 X 2 subblock having the sum of the squares of the edge differences equal to 2.

Conjectures from _Colin Barker_, Oct 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008358 | OEIS-A233656

a(n) = 3*a(n-1) - 2*(n-1), with a(0) = 1.

Conjecture: Last digit of a(n) has repeating pattern of 20 digits as follows: {1, 3, 7, 7, 5, 7, 1, 1, 9, 1, 5, 5, 3, 5, 9, 9, 7, 9, 3, 3}, with an equal frequency of the five odd digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard R. Forberg_, Dec 14 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008359 | OEIS-A233766

Digital root of Lucas numbers (A030133) indexed by natural numbers not divisible by 2, 3 or 5 (A007775).

Conjectures from _Colin Barker_, Sep 22 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary Croft_, Dec 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008360 | OEIS-A233793

Least odd prime p such that 2*n - p = sigma(k) for some k > 0, or 0 if such an odd prime p does not exist, where sigma(k) is the sum of all (positive) divisors of k.

Conjecture: a(n) > 0 for all n > 1. Moreover, if n > 180 is not among 284, 293, 371, 542, 788, 1274, then 2*n can be written as p + sigma(m^2), where p is an odd prime and m is a positive integer. See also part (i) of the conjecture in A233654. We have verified part (i) of the conjecture for n up to 10^9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 15 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008361 | OEIS-A233860

E.g.f. satisfies: A'(x) = Product_{n>=1} 1/(1 - A(x)^n), where A(0) = 0.

CONJECTURES.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 16 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008362 | OEIS-A234016

Partial sums of the characteristic function of A055938.

Conjecture: partial sums of A308187 (i.e, A308187 is the characteristic function of A055938). - _Sean A. Irvine_, Jul 16 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 18 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008363 | OEIS-A234133

Number of (n+1) X (1+1) 0..3 arrays with every 2 X 2 subblock having the sum of the absolute values of all six edge and diagonal differences equal to 9.

Conjectures from _Colin Barker_, Oct 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 19 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008364 | OEIS-A234220

Number of (n+1) X (1+1) 0..4 arrays with every 2 X 2 subblock having its diagonal sum differing from its antidiagonal sum by 3 (constant-stress 1 X 1 tilings).

Conjectures from _Colin Barker_, Oct 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008365 | OEIS-A234246

a(n) = |{0 < k < n: k*phi(n-k) + 1 is a square}|, where phi(.) is Euler's totient function.

Conjecture: (i) a(n) > 0 if n is not a divisor of 6. The only values of n with a(n) = 1 are 4, 5, 8, 9, 12, 13, 24, 33, 49.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 21 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008366 | OEIS-A234259

Number of (n+1) X (1+1) 0..2 arrays with every 2 X 2 subblock having its diagonal sum differing from its antidiagonal sum by 2 (constant-stress 1 X 1 tilings).

Conjectures from _Colin Barker_, Oct 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008367 | OEIS-A234269

Expansion of (1-2*x^2-sqrt(1-4*x^2-4*x^3))/(2*x*sqrt(1-4*x^2-4*x^3)).

Conjecture D-finite with recurrence: -(n+1)*(2*n-5)*a(n) -2*n*(n-4)*a(n-1) +4*(n-1)*(2*n-3)*a(n-2) +2*(8*n^2-32*n+29)*a(n-3) +4*(n-2)*(2*n-7)*a(n-4)=0. - _R. J. Mathar_, Jan 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008368 | OEIS-A234285

Positive odd numbers n such that sigma(m) - 2m is never equal to n, where sigma(.) is the sum of divisors function A000203. Conjectural.

Positive odd numbers n such that sigma(m) - 2m is never equal to n, where sigma(.) is the sum of divisors function A000203. Conjectural. The terms shown here are conjectural, based on a search up to 10^20 made by Davis et al. (2013). Moreover I conjecture that, for each even number r, there exist infinitely many primes p of the form 2^k - r - 1, or equivalently, I conjecture that: For each odd number s, there exists infinitely many primes p of the form 2^k - s.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008369 | OEIS-A234286

Positive odd numbers n such that 2m - sigma(m) is never equal to n, where sigma(.) is the sum of divisors function A000203. Conjectural.

Positive odd numbers n such that 2m - sigma(m) is never equal to n, where sigma(.) is the sum of divisors function A000203. Conjectural. The terms shown here are conjectural, based on a search up to 10^20 made by Davis et al. (2013).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008370 | OEIS-A234307

a(n) = Sum_{i=1..n} gcd(2*n-i, i).

Conjecture: a(n) = (1/4)*Sum_{k = 1..4*n} (-1)^k *gcd(k, 8*n). - _Peter Bala_, Jan 01 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Dec 22 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008371 | OEIS-A234309

a(n) = |{2 < k <= n/2: 2^{phi(k)} + 2^{phi(n-k)} - 1 is prime}|, where phi(.) is Euler's totient function.

Conjecture: (i) a(n) > 0 for all n > 5. As phi(k) is even for any k > 2, part (i) of the conjecture implies that there are infinitely many primes of the form 4^a + 4^b - 1 with a and b positive integers (cf. A234310). Note that any Mersenne prime greater than 3 has the form 2^{2a+1} - 1 = 4^a + 4^a - 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008372 | OEIS-A234337

a(n) = |{0 < k < n - 2: 4^k + 2^{phi(n-k)} - 1 is prime}|, where phi(.) is Euler's totient function.

Conjecture: Let a be 2 or 3 or 4. If n > 3, then a^k + a^{phi(n-k)/2} - 1 is prime for some 0 < k < n - 2. This conjecture for a = 4 implies that there are infinitely many terms of the sequence A234310. The conjecture for a = 3 implies that there are infinitely many primes of the form 3^k + 3^m - 1 (cf. A234346), where k and m are positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008373 | OEIS-A234346

Primes of the form 3^k + 3^m - 1, where k and m are positive integers.

By a conjecture in A234337 or A234347, this sequence should have infinitely many terms. Conjecture: For any integer a > 1, there are infinitely many primes of the form a^k + a^m - 1, where k and m are positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 23 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008374 | OEIS-A234360

a(n) = |{0 < k < n: (k+1)^{phi(n-k)} + k is prime}|, where phi(.) is Euler's totient function.

Conjecture: (i) a(n) > 0 for all n > 1. Also, for any n > 5 there is a positive integer k < n with (k+1)^{phi(n-k)/2} - k prime. Adamczewski disproves the above conjecture in linked article, using case n = 1408. - _Michael De Vlieger_, Aug 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 24 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008375 | OEIS-A234366

Primes of the form q(p) + 1, where p is a prime and q(.) is the strict partition function (A000009).

Though the primes in this sequence are very rare, by the conjecture in A234514 there should be infinitely many such primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008376 | OEIS-A234388

Primes of the form 2^k*(2^{phi(m)} - 1) + 1, where k and m are positive integers, and phi(.) is Euler's totient function.

Conjecture: (i) Any integer n > 1 can be written as k + m with k > 0 and m > 0 such that 2^k*(2^{phi(m)} - 1) + 1 is prime. Part (i) of the conjecture implies that this sequence has infinitely many terms. See also A234399.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008377 | OEIS-A234399

a(n) = |{0 < k < n: 2^k*(2^phi(n-k) - 1) + 1 is prime}|, where phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for all n > 1. See also the conjecture in A234388. The conjecture is false. a(5962) = 0. - _Jason Yuen_, Nov 04 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 25 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008378 | OEIS-A234429

Numbers which are the digital sum of the square of some prime.

Conjectures from _Chai Wah Wu_, Apr 16 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antonio Graciá Llorente_, Dec 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008379 | OEIS-A234451

Number of ways to write n = k + m with k > 0 and m > 0 such that 2^(phi(k)/2 + phi(m)/6) + 3 is prime, where phi(.) is Euler's totient function.

Conjecture: (i) a(n) > 0 for all n > 9. Also, any integer n > 13 can be written as k + m with k > 0 and m > 0 such that 2^(phi(k)/2 + phi(m)/6) - 3 is prime. This conjecture implies that there are infinitely many primes in any of the four forms 2^n + 3, 2^n - 3, 3*2^n + 1, 3*2^n - 1. We have verified the conjecture for n up to 50000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008380 | OEIS-A234470

Number of ways to write n = k + m with k > 0 and m > 2 such that p(k + phi(m)/2) is prime, where p(.) is the partition function (A000041) and phi(.) is Euler's totient function.

Conjecture: a(n) > 0 if n > 3 is not among 27, 34, 50, 61, 74, 78, 115, 120, 123, 127.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008381 | OEIS-A234475

Number of ways to write n = k + m with 2 < k <= m such that q(phi(k)*phi(m)/4) + 1 is prime, where phi(.) is Euler's totient function and q(.) is the strict partition function (A000009).

Conjecture: a(n) > 0 for all n > 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008382 | OEIS-A234504

Number of ways to write n = k + m with k > 0 and m > 0 such that 2^(phi(k) + phi(m)/4) - 5 is prime, where phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for all n > 10. We have verified this for n up to 50000. The conjecture implies that there are infinitely many primes of the form 2^n - 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 26 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008383 | OEIS-A234514

Number of ways to write n = k + m with k > 0 and m > 0 such that p = k + phi(m)/2 and q(p) + 1 are both prime, where phi(.) is Euler's totient function, and q(.) is the strict partition function (A000009).

Conjecture: (i) a(n) > 0 for all n > 12. Clearly, part (i) of the conjecture implies that there are infinitely many primes p with q(p) + 1 prime (cf. A234530).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008384 | OEIS-A234530

Primes p with q(p) + 1 also prime, where q(.) is the strict partition function (A000009).

By the conjecture in A234514, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008385 | OEIS-A234535

Numbers n such that n-1 is a divisor of 3^n + 5^n.

It is an open problem to find all numbers n such that (n+1)(n-1) is a divisor of 3^n + 5^n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Siad Daboul_, Dec 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008386 | OEIS-A234536

Numbers k such that k+1 is a divisor of 3^k + 5^k.

It is an open problem to find all numbers k such that (k+1)*(k-1) is a divisor of 3^k + 5^k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Siad Daboul_, Dec 27 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008387 | OEIS-A234566

1/a(n) is the area of the smallest triangle delimited by 3 lines each passing through at least 2 points of an n X n unitary spaced grid.

Conjecture: sequence is well-defined, i.e., a(n) is an integer for every n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Resta_, Dec 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008388 | OEIS-A234567

Number of ways to write n = k + m with k > 0 and m > 0 such that p = phi(k) + phi(m)/2 + 1 and P(p-1) are both prime, where phi(.) is Euler's totient function and P(.) is the partition function (A000041).

Conjecture: (i) a(n) > 0 for all n > 727. Part (i) of the conjecture implies that there are infinitely many primes p with P(p-1) prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008389 | OEIS-A234569

Primes p with P(p-1) also prime, where P(.) is the partition function (A000041).

By the conjecture in A234567, this sequence should have infinitely many terms. It seems that a(n+1) < a(n) + a(n-1) for all n > 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008390 | OEIS-A234572

Primes of the form P(p-1), where p is a prime and P(.) is the partition function (A000041).

Though the primes in this sequence are very rare, by the conjecture in A234567 there should be infinitely many such primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008391 | OEIS-A234604

Floor of the solutions to c = exp(1 + n/c) for n >= 0, using recursion.

Conjecture: a(n) = floor(e^(-e^(t^2/e^t - t)*t^2 + t + 1)), where t = n/e, for all n > 13. - _Jon E. Schoenfield_, Jan 11 2014 [clarified by _Qiyuan Gu_, Sep 14 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard R. Forberg_, Dec 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008392 | OEIS-A234615

Number of ways to write n = k + m with k > 0 and m > 0 such that p = prime(k) + phi(m) and q(p) - 1 are both prime, where phi(.) is Euler's totient function and q(.) is the strict partition function (A000009).

Conjecture: (i) a(n) > 0 for all n > 7. Clearly, part (i) of this conjecture implies that there are infinitely many primes p with q(p) - 1 also prime (cf. A234644).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008393 | OEIS-A234642

Smallest x such that x mod phi(x) = n, or 0 if no such x exists.

Conjecture: a(n) > 0 for all n. This would follow from a form of Goldbach's (binary) conjecture. Checked up to 10^7; largest term in that range is a(9972987) = 4178506411. Pomerance proves that x = n (mod phi(x)) has at least two solutions for each n, but this allows x < n and so does not prove the conjecture above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Dec 28 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008394 | OEIS-A234695

Primes p with prime(p) - p + 1 also prime.

By the conjecture in A234694, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 29 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008395 | OEIS-A234779

Number of (n+1) X (1+1) 0..3 arrays with no adjacent elements equal and with each 2 X 2 subblock having the number of clockwise edge increases equal to the number of counterclockwise edge increases.

Conjectures from _Colin Barker_, Oct 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008396 | OEIS-A234788

Solutions to numerator(Bernoulli(k)) == denominator/6 (Bernoulli(k)) (mod 30).

Conjecture: the residues mod 30 of the numerator and the denominator/6 of Bernoulli(20(n-1) + 2) are equal. Conjecture: the only solutions to the above equation are {1, 7, 11, 13, 17, 19, 23 or 29}. Observation: the differences between these solutions are (6, 4, 2, 4, 2, 4, 6), a sequence with bilateral symmetry. Program checks all nonzero Bernoulli numbers except B(1), but if the above conjecture is true, then it needs check only every 20th Bernoulli Number starting with B(2). This formula is conjectural, but the program verified it for each of the first 300 numbers in this sequence: to obtain k from the n of this sequence, k = 20(n-1) + 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael G. Kaarhus_, Dec 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008397 | OEIS-A234789

Number of (n+1) X (1+1) 0..3 arrays with each 2 X 2 subblock having the number of clockwise edge increases less than or equal to the number of counterclockwise edge increases.

Conjectures from _Colin Barker_, Oct 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008398 | OEIS-A234800

First occurrence of n in A234323: Number of nontrivial zeros of the Riemann zeta function in the interval 1/2 + i[n,n+1).

The occurrence of <0> is probably finite since the average height of a(n) is A234323 is log(n)/(2*Pi). This is a conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_ and _Robert G. Wilson v_, Dec 30 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008399 | OEIS-A234839

a(n) = Sum_{k = 0..n} (-1)^k * binomial(n,k) * binomial(2*n,k).

Conjecture: the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) hold for all primes p >= 5 and positive integers n and k. [added Apr 12 2022: this was proved in 2014 by Osburn et al.; see Example 3.3]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Dec 31 2013 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008400 | OEIS-A234875

Number of (n+1) X (1+1) 0..4 arrays with every 2 X 2 subblock having its diagonal sum differing from its antidiagonal sum by 1, with no adjacent elements equal (constant-stress tilted 1 X 1 tilings).

Conjectures from _Colin Barker_, Oct 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008401 | OEIS-A234903

a(n)*Pi is the total length of irregular spiral (center points: 1, 3, 2) after n rotations.

G.f.: x*(4*x^4 + 7*x^3 + 2*x^2 + 7*x + 4)/((1-x)(1-x^5)) (conjectured). - _Ralf Stephan_, Jan 13 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jan 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008402 | OEIS-A234960

Sequence (or tree) T of primes generated by these rules: 2 is in T; if p is in T, then the greatest prime < 2*p is in T; if p is in T, then the least prime > 2*p is in T; duplicates are deleted as they occur.

The rules generate successive generations g(n) as follows: g(1) = (2), which begets 3 and 5, so that g(2) = (3,5); then g(3) = (7,11); g(4) = (13,17,19,23); etc. The number of primes in g(n) is given by A234961, and primes not generated, beginning with 71, are given by A234962. Conjecture: the limiting relative density of generated primes is 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008403 | OEIS-A234962

Primes missing from the tree generated at A234960.

According to the conjecture at A234961, the limiting relative density of these primes is 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008404 | OEIS-A234963

Number of ways to write n = k + m with k > 0 and m > 2 such that C(2*sigma(k) + phi(m), sigma(k) + phi(m)/2) - 1 is prime, where sigma(k) is the sum of all positive divisors of k and phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for all n >= 180. Clearly, this implies that there are infinitely many primes of the form C(2*n,n) - 1. We have verified the conjecture for n up to 10000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008405 | OEIS-A235061

Number of ways to write n = k*(k+1)/2 + m with k > 0 and m > 0 such that prime(k*(k+1)/2) + phi(m) is prime, where phi(.) is Euler's totient function.

Conjecture: (i) a(n) > 0 for all n > 8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008406 | OEIS-A235063

Continued fraction expansion of Sum(i=1..inf, 1/2^(2^i+1)).

Recurrence: a(8n)=1, a(8n+4)=a(16n+14)=a(32n+26)=2, a(16n+6)=a(32n+10)=3, a(8n+3)=4, a(8n+7)=a(16n+5)=a(32n+9)=8, a(16n+13)=a(32n+25)=12, a(8n+1)=a(4n+1), a(8n+2)=a(4n+2), starting 2,2,4 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Jan 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008407 | OEIS-A235088

a(n)*Pi is the total length of irregular spiral (center points: 1, 2, 3, 4) after n rotations.

a(n) = 2*floor((n-1)^2/4) + 3*ceiling(n^2/2) (conjectured). - _Ralf Stephan_, Jan 13 2014 Conjecture: a(n) = 1-(-1)^n-n+2*n^2. a(n) = 2*a(n-1)-2*a(n-3)+a(n-4). G.f.: -x*(5*x^2+3)/((x-1)^3*(x+1)). - _Colin Barker_, Jan 16 2014 a(n) = 2*n^2 + n - 4*floor(n/2). In particular, Stephan's and Barker's conjectures are true (see Fried link). - _Sela Fried_, May 08 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jan 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008408 | OEIS-A235089

a(n)*Pi is the total length of irregular spiral (center points: 2, 1, 3, 4) after n rotations.

Conjecture: Numbers equivalent 0 or 3 modulo 10. - _Ralf Stephan_, Jan 13 2014 Conjectures from _Colin Barker_, Jan 16 2014: (Start) a(n) = 5*n - 2*floor((n+1)/2) + 2*floor(n/2). In particular, Barker's conjectures are true (see Fried link). - _Sela Fried_, May 08 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jan 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008409 | OEIS-A235112

a(n) = the largest of the M-indices of the trees with n vertices.

It is conjectured that for n>=7 one has a(n) = A235120(n-6).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jan 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008410 | OEIS-A235146

a(n) = Least integer k such that it takes n iterations of "factor and reverse bits of odd prime divisors" (A235027) before a fixed point or cycle of 2 is reached; records in A235145.

Conjecture: additional property is that the primes here belong to that subset of p in A204219 for which A056539(p) > p. The list of such primes begins as: 19, 79, 103, 137, 139, 149, 157, 179, 191, 239, 271, 281, 293, 311, 317, 347, 367, 379, 439, 523, 541, 547, 557, 563, 569, 587, 607, 613, 647, 659, 719, 733, 743, 751, 787, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008411 | OEIS-A235202

Numbers written in an alternating binary-then-quaternary base.

Conjecture: This sequence interpreted as quaternary (base 4) numbers gives A126001 (hence a simplified scheme for computing that sequence).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeremy Gardiner_, Jan 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008412 | OEIS-A235230

Integer radii of circles tiled by square tiles such that the ratio of uncut tiles to cut tiles is an integer and four square tiles meet at the center of the circle.

It is my conjecture that there are an infinite number of solutions and that they occur by chance, accounting for the widening gaps between valid answers as the number of digits for the sums of tiles increases.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gregory V. Richardson_, Jan 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008413 | OEIS-A235264

Tileable numbers: base-2 representation, considered as a fixed disconnected polyomino, tiles all places >= 0.

Conjecturally, a(n) = (2^k-1)/m where k, m >= 1, and base-2 product m*a(n) involves no carries. Confirmed for a(n) <= 2^20. Conjecturally, a(n) is of the form Product (2^(d_i*b_i)-1)/(2^b_i-1) where d_i >= 1, b_i >= 2, and d_i*b_i | d_(i+1). Confirmed for a(n) <= 2^20. First conjecture is equivalent to the 2-adic definition. - _Charlie Neder_, Nov 04 2018 Second conjecture is true, see Neder link. - _Charlie Neder_, Dec 04 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_, Jan 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008414 | OEIS-A235271

Number of (n+1) X (1+1) 0..4 arrays with every 2 X 2 subblock having its diagonal sum differing from its antidiagonal sum by 5 (constant-stress 1 X 1 tilings).

Conjectures from _Colin Barker_, Oct 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008415 | OEIS-A235282

Number of (n+1) X (1+1) 0..3 arrays with every 2 X 2 subblock having its diagonal sum differing from its antidiagonal sum by 4 (constant-stress 1 X 1 tilings).

Conjectures from _Colin Barker_, Oct 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008416 | OEIS-A235334

Numbers n such that for any positive integers (a, b), if a * b = n then a + b is a square.

Conjecture: the numbers n such that for any positive integers (a, b), a * b = n and a + b is a square are the product of twin primes, and a*b+1 is also a perfect square.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008417 | OEIS-A235344

Numbers m with m - 1, m + 1 and q(m) + 1 all prime, where q(.) is the strict partition function (A000009).

Clearly, any term after the first term 4 is a multiple of 6. By part (i) of the conjecture in A235343, this sequence should have infinitely many terms. The prime q(a(51)) + 1 = q(3235368) + 1 has 1412 decimal digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008418 | OEIS-A235356

Primes of the form q(m) + 1 with m - 1 and m + 1 both prime, where q(.) is the strict partition function (A000009).

Though the primes in this sequence are very rare, by part (i) of the conjecture in A235343 there should be infinitely many such primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008419 | OEIS-A235383

Fibonacci numbers that are the product of other Fibonacci numbers.

Saha and Karthik conjectured (without reference to Carmichael's theorem) that the only positive integers k for which A001175(k^2) = A001175(k) are 6 and 12. (A000045(6) = 8 and A000045(12) = 144.) - _L. Edson Jeffery_, Feb 13 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert C. Lyons_, Jan 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008420 | OEIS-A235450

A strong divisibility sequence associated with the algebraic integer 2 + 3*sqrt(3).

Let alpha be an algebraic integer and define a sequence of integers a(n) by the condition a(n) = max {integer d : alpha^n == 1 (mod d)}. Silverman shows that a(n) is a strong divisibility sequence, that is, gcd(a(n), a(m)) = a(gcd(n, m)) for all n and m in N; in particular, if n divides m then a(n) divides a(m). For the present sequence we take alpha = 2 + 3*sqrt(3). For other examples see A230368, A230369 and (conjecturally) A082630.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jan 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008421 | OEIS-A235453

Triangle T(n, k) = Number of non-equivalent (mod D_4) ways to arrange k indistinguishable points on an n X n square grid so that no three of them are collinear. Triangle read by rows.

2n is an upper bound on the number of points that can be placed on the grid. For large n, it is conjectured that this bound is not reached (see MathWorld link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Heinrich Ludwig_, Jan 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008422 | OEIS-A235459

Number of facets of the correlation polytope of degree n.

According to the SMAPO database it is conjectured that a(8) = 12246651158320. This database also says that the above value of a(7) is conjectural, but Ziegler lists it as known.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Victor S. Miller_, Jan 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008423 | OEIS-A235467

Primes whose base-4 representation also is the base-3 representation of a prime.

This is a subsequence of A002144, A002313, A003655, A050150, A062090, A141293, A175768, A192592, A226181 (conjectural).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008424 | OEIS-A235477

Primes whose base-2 representation also is the base-7 representation of a prime.

A subsequence of A027697, A015919, A197636 (conjectural).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008425 | OEIS-A235508

Number of ways to write 2*n = p + q with q > 0 such that p, p*(p+1) - prime(p) and prime(q) - q + 1 are all prime.

Conjecture: a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008426 | OEIS-A235510

Number of (n+1) X (1+1) 0..1 arrays with the sum of each 2 X 2 subblock maximum and minimum lexicographically nondecreasing rowwise and columnwise.

Conjectures from _Colin Barker_, Mar 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008427 | OEIS-A235592

Numbers k such that k*(k+1) - prime(k) is prime.

It is known that prime(k) <= k*(k+1) for any positive integer k. The conjecture in A235613 implies that the sequence has infinitely many terms. Conjecture: This sequence contains infinitely many primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008428 | OEIS-A235608

Triangle read by rows: a non-Riordan array serving as a counterexample to a conjecture about Riordan arrays.

Triangle read by rows: a non-Riordan array serving as a counterexample to a conjecture about Riordan arrays.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008429 | OEIS-A235613

Number of ways to write n = k + m (0 < k <= m) with k and m terms of A235592.

Conjecture: a(n) > 0 for all n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008430 | OEIS-A235614

Number of ordered ways to write n = k + m with k a term of A235592 and m a positive triangular number.

Conjecture: a(n) > 0 for all n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008431 | OEIS-A235643

Total number of sides of a tetraflake-like fractal after n iterations, a(1) = 16 (see comments).

Conjecture from _Colin Barker_, Apr 21 2014: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Apr 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008432 | OEIS-A235644

Number of decompositions of 12*n into the sum of two (not necessarily distinct) twin prime pairs.

In the 1980's, Liang conjectured that (6n)^2 = p_1 + p_2 + p_3 + p_4, where (p_1, p_2) and (p_3, p_4) are twin prime pairs. See reference for more details.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lear Young_, Jun 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008433 | OEIS-A235661

Primes p with p*(p+1) - prime(p) prime.

By the conjecture in A232353, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008434 | OEIS-A235681

Primes p with prime(p) - p + 1 and p*(p+1) - prime(p) both prime.

By the conjecture in A235682, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008435 | OEIS-A235682

Number of ways to write n = k + m with k > 0 and m > 2 such that p = phi(k) + phi(m)/2 + 1, prime(p) - p + 1 and p*(p+1) - prime(p) are all prime, where phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for all n > 84.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008436 | OEIS-A235703

Number of ordered ways to write n = p + q with p a term of A234695 and q a term of A235592.

Conjecture: a(n) > 0 for all n > 3, and a(n) = 1 only for n = 4, 30, 83.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008437 | OEIS-A235727

Odd primes p with (p^2 - 1)/4 - prime((p - 1)/2) and (p^2 - 1)/4 + prime((p - 1)/2) both prime.

By the conjecture in A235728, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008438 | OEIS-A235728

a(n) = |{0 < k < n - 2: 2*m + 1, m*(m+1) - prime(m) and m*(m+1) + prime(m) are all prime with m = phi(k) + phi(n-k)/2}|, where phi(.) is Euler's totient function.

Conjecture: (i) a(n) > 0 for all n > 5, and a(n) = 1 only for n = 191. Clearly, part (i) of the conjecture implies that there are infinitely many odd primes p = 2*m + 1 with m*(m+1) - prime(m) = (p^2-1)/4 - prime((p-1)/2) and m*(m+1) + prime(m) = (p^2-1)/4 + prime((p-1)/2) both prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008439 | OEIS-A235791

Irregular triangle read by rows: T(n,k), n >= 1, k >= 1, in which column k lists k copies of every positive integer in nondecreasing order, and the first element of column k is in row k(k+1)/2.

Conjecture: T(n,k) is also the total number of partitions of all positive integers <= n into exactly k consecutive parts, i.e., the partial column sum of A285898, or in accordance with the triangles of the same family: the partial column sum of A237048. - _Omar E. Pol_, Apr 28 2017, Nov 24 2020 The above conjecture is true. The proof will be added soon (it uses the generating function for the columns). - _N. J. A. Sloane_, Nov 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jan 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008440 | OEIS-A235804

Rectangular array read by upward antidiagonals: A(n,k) = n-2+k*2^(n-3), n>=3, k>=0.

Conjectured equivalence classes associated with the row entries of A233332. Conjecture: G.f. for row n is F_n(x) = ((n-2)+(2^(n-3)-(n-2))*x)/(1-x)^2 = ((n-2)+(2^(n-3)-(n-3)-1)*x)/(1-x)^2 = ((n-2)+A000295(n-3)*x)/(1-x)^2, n>=3. Conjecture: G.f. for column k is G_k(x) = (k+1-2*(k+1)*x+k*x^2)/((1-2*x)*(1-x)^2), k>=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008441 | OEIS-A235877

Number of (n+1) X (1+1) 0..2 arrays with the minimum plus the upper median equal to the lower median plus the maximum in every 2 X 2 subblock.

Conjectures from _Colin Barker_, Oct 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008442 | OEIS-A235878

Number of (n+1) X (2+1) 0..2 arrays with the minimum plus the upper median equal to the lower median plus the maximum in every 2 X 2 subblock.

Conjectures from _Colin Barker_, Oct 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008443 | OEIS-A235879

Number of (n+1) X (3+1) 0..2 arrays with the minimum plus the upper median equal to the lower median plus the maximum in every 2 X 2 subblock.

Conjectures from _Colin Barker_, Oct 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008444 | OEIS-A235880

Number of (n+1) X (4+1) 0..2 arrays with the minimum plus the upper median equal to the lower median plus the maximum in every 2 X 2 subblock.

Conjectures from _Colin Barker_, Oct 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008445 | OEIS-A235881

Number of (n+1) X (5+1) 0..2 arrays with the minimum plus the upper median equal to the lower median plus the maximum in every 2 X 2 subblock.

Conjectures from _Colin Barker_, Oct 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008446 | OEIS-A235882

Number of (n+1) X (6+1) 0..2 arrays with the minimum plus the upper median equal to the lower median plus the maximum in every 2 X 2 subblock.

Conjectures from _Colin Barker_, Oct 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008447 | OEIS-A235883

Number of (n+1) X (7+1) 0..2 arrays with the minimum plus the upper median equal to the lower median plus the maximum in every 2 X 2 subblock.

Conjectures from _Colin Barker_, Oct 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008448 | OEIS-A235886

Number of (n+1) X (1+1) 0..3 arrays with the minimum plus the upper median equal to the lower median plus the maximum in every 2 X 2 subblock.

Conjectures from _Colin Barker_, Oct 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008449 | OEIS-A235887

Number of (n+1) X (2+1) 0..3 arrays with the minimum plus the upper median equal to the lower median plus the maximum in every 2 X 2 subblock.

Conjectures from _Colin Barker_, Oct 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008450 | OEIS-A235888

Number of (n+1) X (3+1) 0..3 arrays with the minimum plus the upper median equal to the lower median plus the maximum in every 2 X 2 subblock.

Conjectures from _Colin Barker_, Oct 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008451 | OEIS-A235889

Number of (n+1) X (4+1) 0..3 arrays with the minimum plus the upper median equal to the lower median plus the maximum in every 2 X 2 subblock.

Conjectures from _Colin Barker_, Oct 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008452 | OEIS-A235890

Number of (n+1) X (5+1) 0..3 arrays with the minimum plus the upper median equal to the lower median plus the maximum in every 2 X 2 subblock.

Conjectures from _Colin Barker_, Oct 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008453 | OEIS-A235891

Number of (n+1) X (6+1) 0..3 arrays with the minimum plus the upper median equal to the lower median plus the maximum in every 2 X 2 subblock.

Conjectures from _Colin Barker_, Oct 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008454 | OEIS-A235892

Number of (n+1) X (7+1) 0..3 arrays with the minimum plus the upper median equal to the lower median plus the maximum in every 2 X 2 subblock.

Conjectures from _Colin Barker_, Oct 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008455 | OEIS-A235895

Number of (n+1) X (1+1) 0..2 arrays with the minimum plus the maximum equal to the lower median plus the upper median in every 2 X 2 subblock.

Conjectures from _Colin Barker_, Oct 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008456 | OEIS-A235914

Odd primes p = 2*m + 1 with m*(m-1) - prime(m) and m*(m+1) - prime(m) both prime.

By the conjecture in A235912, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008457 | OEIS-A235917

a(n) = |{0 < k < n - 2: p = prime(k) + phi(n-k)/2, p^2 - 1 - prime(p) and (p^2 - 1)/2 - prime(p) are all prime}|, where phi(.) is Euler's totient function.

Conjecture: (i) a(n) > 0 for all n > 14. Clearly, part (i) of the conjecture implies that there are infinitely many primes p with p^2 - 1 - prime(p) and (p^2 - 1)/2 - prime(p) both prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008458 | OEIS-A235919

a(n) = |{0 < k < n - 2: p = prime(k) + phi(n-k)/2, prime(p) - p + 1 and (p^2 - 1)/4 - prime(p) are all prime}|, where phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for all n > 24.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008459 | OEIS-A235920

Primes p with prime(p) - p + 1 and (p^2 - 1)/4 - prime(p) both prime.

By the conjecture in A235919, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008460 | OEIS-A235922

Minimal k > 1 such that the base-k representation of the n-th palindrome in A002113, read in decimal, is also palindrome.

We conjecture that limit of average (a(1)+...+a(n))/n exists and equals 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008461 | OEIS-A235923

Minimal k > 1 such that the base-k representation of the n-th Ramanujan prime (A104272), read in decimal, is also a Ramanujan prime.

Conjecture 1. Every number 2,...,10 occurs infinitely many times. Conjecture 2. There exists limit of average (a(1)+...+a(n))/n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008462 | OEIS-A235925

Primes p with q = prime(p) - p + 1 and r = prime(q) - q + 1 both prime.

By the conjecture in A235924, this sequence should have infinitely many terms. Conjecture: For any integer m > 1, there are infinitely many chains p(1) < p(2) < ... < p(m) of m primes with p(k+1) = prime(p(k)) - p(k) + 1 for all 0 < k < m.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008463 | OEIS-A235934

Primes p with f(p), f(f(p)) and f(f(f(p))) all prime, where f(n) = prime(n) - n + 1.

By the general conjecture in A235925, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008464 | OEIS-A235935

Primes p with f(p), f(f(p)), f(f(f(p))), f(f(f(f(p)))) all prime, where f(n) = prime(n) - n + 1.

By the general conjecture in A235925, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008465 | OEIS-A235984

Primes p with f(p), f(f(p)), f(f(f(p))), f(f(f(f(p)))), f(f(f(f(f(p))))) all prime, where f(n) = prime(n) - n + 1.

By the general conjecture in A235925, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008466 | OEIS-A236021

Record values in A236020.

Conjecture: subsequence of A094348.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jan 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008467 | OEIS-A236024

Record values in A236022.

Conjecture: union of 10 and squares of primes A001248.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jan 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008468 | OEIS-A236043

Number of triangular numbers <= 10^n.

Except for 5, all numbers begin with either a 4 or a 1. If strictly less than, the 5 would become a 4, satisfying this conjecture. This is not a conjecture, it is a fact and it is the result from the square root of 2 and 20 times powers of ten. - _Robert G. Wilson v_, Jan 11 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Jan 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008469 | OEIS-A236066

Primes p with g(p), g(g(p)), g(g(g(p))), g(g(g(g(p)))), g(g(g(g(g(p))))) all prime, where g(n) = prime(n) - n - 1.

Conjecture: For any integer m > 1, there are infinitely many chains p(1) < ... < p(m) of m primes with p(k+1) = prime(p(k)) - p(k) - 1 for all 0 < k < m. This is similar to the conjecture in A235925.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008470 | OEIS-A236074

a(n) = |{0 < k < n: p = phi(k) + phi(n-k)/6 + 1, prime(2*p) - 2*prime(p) and prime(p) - 2*prime((p-1)/2) are all prime}|, where phi(.) is Euler's totient function.

Conjecture: (i) a(n) > 0 for all n > 116.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008471 | OEIS-A236075

Odd primes p with prime(2*p) - 2*prime(p) and prime(p) - 2*prime((p-1)/2) both prime.

By the conjecture in A236074, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008472 | OEIS-A236097

a(n) = |{0 < k < n-2: p = phi(k) + phi(n-k)/2 + 1, prime(p) - p - 1 and prime(p) - p + 1 are all prime}|, where phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for all n > 31.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008473 | OEIS-A236102

Numbers whose divisors are partition numbers.

Conjecture: the only composite numbers in this sequence are 15, 22, and 77. - _Jon E. Schoenfield_, Feb 05 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jan 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008474 | OEIS-A236107

Nonprimes whose divisors are partition numbers.

Conjecture: no terms exist beyond 77. - _Jon E. Schoenfield_, Feb 05 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jan 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008475 | OEIS-A236108

Nonprimes whose proper divisors are partition numbers.

This conjecture holds for terms <= 10^16. - _David A. Corneth_, Jan 25 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jan 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008476 | OEIS-A236119

Primes p with prime(p) - p - 1 and prime(p) - p + 1 both prime.

By the conjecture in A236097, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008477 | OEIS-A236128

Primes p such that gpf(gpf(2^p-1)-1) = p.

Conjecture: gpf(gpf(2^p-1)-1) = p for finitely many p. Conjecture: gpf(lpf(2^p-1)-1) = p for infinitely many p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Jan 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008478 | OEIS-A236138

a(n) = |{0 < k < n: p = phi(k) + phi(n-k)/3 - 1, prime(p-1) - (p-1) and prime(p-1) - 2*prime((p-1)/2) are all prime}|, where phi(.) is Euler's totient function.

Conjecture: (i) a(n) > 0 for all n > 73. Clearly, part (i) of the conjecture implies that there are infinitely many odd primes p with prime(p-1) - (p-1) and prime(p-1) - 2*prime((p-1)/2) both prime, and part (ii) implies that there are infinitely many odd primes p with prime(p) - 2*prime((p-1)/2) and prime(p) - 2*prime((p+1)/2) both prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008479 | OEIS-A236143

Odd primes p with prime(p-1) - (p-1) and prime(p-1) - 2*prime((p-1)/2) both prime.

By part (i) of the conjecture in A236138, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008480 | OEIS-A236145

Records in the continued fraction expansion of Sum_{k >= 2} 2^(-Fibonacci(k)) (A006518).

Conjecture: For n>4, a(n) = 2^L(n) - 2^F(n) - 1, with L(n) the Lucas numbers A000032, and F(n) the Fibonacci numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Jan 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008481 | OEIS-A236183

Primes p such that for all primes q dividing p-1 every residue mod p is the sum of two q-th powers mod p.

Related to the conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Esteban Arreaga Ambéliz_, Jan 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008482 | OEIS-A236192

a(n) = |{0 < k < n: p = phi(k) + phi(n-k)/4 + 1, prime(p)^2 + (2*p)^2 and p^2 + (2*prime(p))^2 are all prime}|, where phi(.) is Euler's totient function.

Conjecture: (i) a(n) > 0 for all n > 370. Clearly, part (i) of this conjecture implies that there are infinitely many primes p with prime(p)^2 + (2*p)^2 and p^2 + (2*prime(p))^2 both prime, and part (ii) implies that there are infinitely many primes p with prime(p)^2 + (p-1)^2 and p^2 + (prime(p)-1)^2 both prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008483 | OEIS-A236193

Primes p with prime(p)^2 + (2*p)^2 and p^2 + (2*prime(p))^2 both prime.

By part (i) of the conjecture in A236192, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008484 | OEIS-A236211

Numbers c > 0 for which there exist integers a > 1 and b > 1 such that the equation a^x - b^y = c has two solutions in positive integers x, y.

Bennett conjectured that if a, b, c are positive integers with a > 1 and b > 1, then the equation a^x - b^y = c has at most one solution in positive integers x and y, except for the triples (a,b,c) = (3,2,1), (2,5,3), (6,2,4), (2,3,5), (15,6,9), (13,3,10), (2,3,13), (91,2,89), (280,5,275), (6,3,1215), (4930,30,4900). If this is true, then the present sequence is complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Jan 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008485 | OEIS-A236228

Number of solutions (x,y,z) to the Diophantine equation 2^x + p^y = z^2 where p = prime(n).

The maximum x should be at least 30 because of the identity 2^30 + (2^16+1)^1 = (2^15+1)^2 involving the conjectured greatest prime Fermat number 2^16+1 = 65537. - _Jean-François Alcover_, Jan 24 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008486 | OEIS-A236241

a(n) = |{0 < k < n: m = phi(k) + phi(n-k)/8 is an integer with C(2*m, m) + prime(m) prime}|, where C(2*m, m) = (2*m)!/(m!)^2, and phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for every n = 20, 21, ... The conjecture implies that there are infinitely many primes of the form C(2*m, m) + prime(m).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008487 | OEIS-A236242

Numbers m with C(2*m, m) + prime(m) prime, where C(2*m, m) = (2*m)!/(m!)^2.

According to the conjecture in A236241, this sequence should have infinitely many terms. The prime C(2*a(52),a(52)) + prime(a(52)) = C(61468, 30734) + prime(30734) has 18502 decimal digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008488 | OEIS-A236243

Position (or inverse of) n in A235598.

The terms of this sequence are conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jan 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008489 | OEIS-A236256

a(n) = |{0 < k < n: m = phi(k) + phi(n-k)/4 is an integer with C(2*m, m) - prime(m) prime}|, where C(2*m, m) = (2*m)!/(m!)^2.

Conjecture: a(n) > 0 for all n > 410. This implies that there are infinitely many positive integers m with C(2*m, m) - prime(m) prime. We have verified the conjecture for n up to 51000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008490 | OEIS-A236269

First differences of Stanley sequence S(0,4) (A005487).

While there are conjectures about formulas for S(0,m), m=1,2,3,6,9... (see A093682), m=4 is the first case where the first differences look almost random.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Stephan_, Jan 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008491 | OEIS-A236283

The number of orbits of triples of {1,2,...,n} under the action of the dihedral group of order 2n.

Conjectures from _Colin Barker_, Jan 21 2014: (Start) The above conjectures are true and can be derived from the following formulas for even and odd n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _W. Edwin Clark_, Jan 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008492 | OEIS-A236305

The number of P-positions in the game of Nim with up to 3 piles, allowing for piles of zero, such that the number of objects in each pile does not exceed n.

a(n) = 4^floor(log(n)/log(2)) + 3*(n mod 2^floor(log(n)/log(2)))^2 (conjectured). - _Thomas Baruchel_, May 15 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tanya Khovanova_ and _Joshua Xiong_, Apr 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008493 | OEIS-A236306

Least prime p < prime(n) for which both p and p! are primitive roots modulo prime(n), or 0 if such a prime does not exist.

Conjecture: a(n) > 0 for all n > 4. In other words, for any prime p > 7, there exists a prime q < p such that both q and q! are primitive roots modulo p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008494 | OEIS-A236320

Conjectured number of Sierpiński numbers less than 10^n.

Conjectured number of Sierpiński numbers less than 10^n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Jan 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008495 | OEIS-A236321

Conjectured number of Riesel numbers less than 10^n.

Conjectured number of Riesel numbers less than 10^n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Jan 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008496 | OEIS-A236326

a(n)*Pi is the total length of irregular spiral (center points: 1, 2, 3, 4, 5; pattern 1) after n rotations.

Conjecture: All forms of 120 permutations of 5 center points are non-expanded loops. Conjectures from _Colin Barker_, Jul 12 2014: (Start) a(n) = 3*n + 4*floor(n/5) + 4*floor((n+1)/5) + floor((n+2)/5). In particular, Barker's conjectures are true (see Fried link). - _Sela Fried_, May 08 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jan 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008497 | OEIS-A236327

a(n)*Pi is the total length of irregular spiral (center points: 1, 2, 3, 4, 5; pattern 2) after n rotations.

Conjecture: All forms of 120 permutations 5 center points are non-expanded loops. Conjectures from _Colin Barker_, Jul 12 2014: (Start) a(n) = 2*n + 9*floor(n/4) + 9*floor((n+1)/4) + 4*floor((n+2)/4). In particular, Barker's conjectures are true (see Fried link). - _Sela Fried_, May 08 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jan 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008498 | OEIS-A236332

The number of orbits of 4-tuples of the dihedral group of order 2n acting on {1,2,...,n}.

Conjectures from _Colin Barker_, Jan 22 2014: (Start) The above conjectures are true and can be derived from the following formulas for even and odd n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _W. Edwin Clark_, Jan 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008499 | OEIS-A236375

Positive integers m with 2^(m-1)*phi(m) - 1 prime, where phi(.) is Euler's totient function.

According to the conjecture in A236374, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008500 | OEIS-A236383

Smallest k such that k^2 is a concatenation of two numbers x and y where y = x + n^2 and x and y have the same number of digits.

Conjecture: a(n) exists for all numbers n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008501 | OEIS-A236389

a(n) = |{0 < k < n: m = phi(k)/2 + phi(n-k)/12 is an integer with 2^m*p(m) + 1 prime}|, where p(.) is the partition function (A000041).

Conjecture: a(n) > 0 for all n > 56. The conjecture implies that there are infinitely many positive integers m with 2^m*p(m) + 1 prime. See A236390 for a list of such numbers m.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008502 | OEIS-A236390

Positive integers m with 2^m*p(m) + 1 prime, where p(.) is the partition function (A000041).

According to the conjecture in A236389, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008503 | OEIS-A236397

Weight of the largest-weight sunflower-free set of width n.

Peebles conjectures that if n is even, a(n+1) = 2*A090245(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008504 | OEIS-A236401

Kurepa determinant K_n.

Conjecture: a(n) ~ -(-1)^n * n! * exp(-1) / n^4. - _Vaclav Kotesovec_, Nov 30 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008505 | OEIS-A236408

Number of increasing planar st-graphs with n edges.

Conjecture: (n+2)*(n+3)*a(n) = 2*n*(2*n+1)*a(n-1) + (n-2)*(19*n-9)*a(n-2) + 14*(n-3)*(n-2)*a(n-3). - _Vaclav Kotesovec_, Feb 14 2014 [For a proof of the conjecture see Corollary 4.14 in Bostan et al. arxiv.org/abs/2110.13753. - _Mark van Hoeij_, Nov 05 2023]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008506 | OEIS-A236412

a(n) = |{0 < k < n: m = phi(k)/2 + phi(n-k)/8 is an integer with p(m)^2 + q(m)^2 prime}|, where phi(.) is Euler's totient, p(.) is the partition function (A000041) and q(.) is the strict partition function (A000009).

Conjecture: a(n) > 0 for all n > 17. The conjecture implies that there are infinitely positive integers m with p(m)^2 + q(m)^2 prime. See A236413 for a list of such numbers m. See also A236414 for primes of the form p(m)^2 + q(m)^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008507 | OEIS-A236413

Positive integers m with p(m)^2 + q(m)^2 prime, where p(.) is the partition function (A000041) and q(.) is the strict partition function (A000009).

According to the conjecture in A236412, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008508 | OEIS-A236414

Primes of the form p(m)^2 + q(m)^2 with m > 0, where p(.) is the partition function (A000041) and q(.) is the strict partition function (A000009).

According to the conjecture in A236412, this sequence should have infinitely many terms. See A236413 for positive integers m with p(m)^2 + q(m)^2 prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008509 | OEIS-A236417

a(n) = |{0 < k < n: p = phi(k)/2 + phi(n-k)/12 + 1 and A047967(p) are both prime}|.

Conjecture: a(n) > 0 for all n > 98. The conjecture implies that there are infinitely many primes p with A047967(p) prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008510 | OEIS-A236418

Primes p with A047967(p) also prime.

According to the conjecture in A236417, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008511 | OEIS-A236419

a(n) = |{0 < k < n: r = phi(k) + phi(n-k)/6 + 1 and p(r) + q(r) are both prime}|, where phi(.) is Euler's totient function, p(.) is the partition function (A000041) and q(.) is the strict partition function (A000009).

Conjecture: a(n) > 0 for all n > 127. The conjecture implies that there are infinitely many primes r with p(r) + q(r) also prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008512 | OEIS-A236440

Positive integers m with A000009(m)^2 + A047967(m)^2 prime.

According to the conjecture in A236439, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008513 | OEIS-A236442

Number of ordered ways to write n = k + m with k > 0 and m > 0 such that A000009(k) + A047967(m) is prime.

Conjecture: (i) a(n) > 0 for all n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008514 | OEIS-A236457

Primes p with q = p + 2 and prime(q) + 2 both prime.

According to the conjecture in A236456, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008515 | OEIS-A236460

Number of ordered ways to write n = k + m with k > 0 and m > 0 such that p = phi(k) + phi(m)/2 - 1, prime(p) + 4 and prime(p) + 6 are all prime, where phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for all n > 211.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008516 | OEIS-A236462

Primes p with prime(p) + 4 and prime(p) + 6 both prime.

According to the conjecture in A236460, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008517 | OEIS-A236464

Primes p with prime(p) + 2 and prime(p) + 6 both prime.

According to the conjecture in A236472, this sequence contains infinitely many terms, i.e., there are infinitely many prime triples of the form {prime(p), prime(p) + 2, prime(p) + 6} with p prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008518 | OEIS-A236467

Primes p with p + 2 and prime(p) - 2 both prime.

According to the conjecture in A236468, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008519 | OEIS-A236470

a(n) = |{0 < k < n: p = prime(k) + phi(n-k), p + 2 and prime(p) + 2 are all prime}|, where phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for all n > 948. The conjecture implies that there are infinitely many primes p with p + 2 and prime(p) + 2 both prime. See A236458 for such primes p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008520 | OEIS-A236472

a(n) = |{0 < k < n: p = prime(k) + phi(n-k), prime(p) + 2 and prime(p) + 6 are all prime}|, where phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for every n = 330, 331, ... The conjecture implies that there are infinitely many prime triples of the form {prime(p), prime(p) + 2, prime(p) + 6} with p prime. See A236464 for such primes p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008521 | OEIS-A236480

a(n) = |{0 < k < n-2: p = 2*phi(k) + phi(n-k)/2 + 1, prime(p) + 2 and prime(prime(p)) + 2 are all prime}|, where phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for every n = 640, 641, ... The conjecture implies that there are infinitely many primes p with prime(p) + 2 and prime(prime(p)) + 2 both prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008522 | OEIS-A236481

Primes p with p + 2, prime(p) + 2 and prime(prime(p)) + 2 all prime.

Conjecture: For any positive integer m, there are infinitely many chains p(1) < p(2) < ... < p(m) of m primes with p(k) + 2 prime for all k = 1,...,m such that p(k + 1) = prime(p(k)) for every 0 < k < m.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008523 | OEIS-A236483

a(n) = |{0 < k < n-2: p(k) + 2^(phi(n-k)/2) is prime}|, where p(.) is the partition function (A000041) and phi(.) is Euler's totient function.

Conjecture: (i) a(n) > 0 for all n > 3. Clearly, part (i) of the conjecture implies that there are infinitely many primes of the form p(k) + 2^m, where k and m are positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008524 | OEIS-A236485

Primes p such that gpf(lpf(m)-1) = gpf(gpf(m)-1), where m = 2^p-1 is Mersenne composite.

Conjecture: finitely many such p and gpf(lpf(m)-1) = gpf(gpf(m)-1) = p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Jan 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008525 | OEIS-A236506

Conjectured least index k > 2 of Fibonacci-like sequence f(i+2) = f(i+1) + f(i), with f(1)=1 and f(2)=n, such that f(k) is a square, or k=0 if squares do not exist in the corresponding sequence.

Conjectured least index k > 2 of Fibonacci-like sequence f(i+2) = f(i+1) + f(i), with f(1)=1 and f(2)=n, such that f(k) is a square, or k=0 if squares do not exist in the corresponding sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carmine Suriano_, Jan 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008526 | OEIS-A236508

a(n) = |{0 < k < n-2: p = 2*phi(k) + phi(n-k)/2 - 1, p + 2, p + 6 and prime(p) + 6 are all prime}|, where phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for all n > 146. The conjecture implies that there are infinitely many prime triples {p, p + 2, p + 6} with {prime(p), prime(p) + 6} a sexy prime pair. See A236509 for such primes p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008527 | OEIS-A236509

Primes p with p + 2, p + 6 and prime(p) + 6 all prime.

According to the conjecture in A236508, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008528 | OEIS-A236511

a(n) = |{0 < k < n: p = 3*phi(k) + phi(n-k) - 1, p + 2, p + 6 and p + 8 are all prime}|, where phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for all n > 1075. The above conjecture implies the well-known conjecture that there are infinitely many prime quadruplets (p, p + 2, p + 6, p + 8).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008529 | OEIS-A236535

a(n)*Pi is the total length of irregular spiral (center points: 2, 3, 1; pattern 1) after n rotations.

Conjecture: All forms of 3 center points are non-expanded loops. Conjecture from _Colin Barker_, Jul 12 2014: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jan 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008530 | OEIS-A236539

Integer areas of integer-sided triangles such that the length of two sides are Fibonacci numbers.

Conjecture: there exists only one triangle where three sides are of Fibonacci number length. This triangle is (5, 5, 8), and the area is A = 12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008531 | OEIS-A236541

Number of ways to write 2*n = k + m with 0 < k <= m such that prime(k) + m and k + prime(m) are both prime.

Conjecture: (i) a(n) > 0 for all n > 31. Part (ii) of the conjecture in A232443 implies that any integer n > 7 can be written as k + m (k > 0, m > 0) with prime(k) + m = n + prime(k) - k prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008532 | OEIS-A236548

Number of ways to write n = k^2 + m with k > 0 and m > 0 such that phi(k^2) + prime(m) is prime, where phi(.) is Euler's totient function.

Conjecture: (i) a(n) > 0 for all n > 100.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008533 | OEIS-A236559

Number of partitions of 2n of type EO (see Comments).

Conjecture: call C the character table of the symmetric group S(n), C_2 the matrix C with each entry squared, and w its class sizes, then matrix A = w/n! transpose(C).(C_2) has rows that are all zero according to partitions of type EO if n is even, and of type OO if n is odd. - _Wouter Meeussen_, Sep 03 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008534 | OEIS-A236568

Primes p with prime(p + 2) + 2 prime.

This sequence is interesting because of the conjecture in A236566.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008535 | OEIS-A236569

Least term p of A236568 with 2*n - p prime, or 0 if such a prime p does not exist.

The conjecture in A236566 implies that a(n) > 0 for all n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008536 | OEIS-A236573

Number of ordered ways to write n = k + m (k > 0, m > 0) such that p = 2*k + phi(m) - 1, prime(p + 2) + 2 and 2*n - p are all prime, where phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for all n > 712. This implies the conjecture in A236566.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008537 | OEIS-A236574

Primes p with prime(p)^3 + 2*p^3 and p^3 + 2*prime(p)^3 both prime.

Conjecture: This sequence has infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008538 | OEIS-A236673

Exponents of powers of 3 that contain all ten decimal digits.

It is conjectured that after a(43), a(n) = n + 63 (i.e., natural numbers beginning with 107).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Jan 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008539 | OEIS-A236674

Exponents of powers of 3 that do not contain all ten decimal digits.

It is conjectured that 106 is the final number in this sequence. 3^106 contains all digits except for 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Jan 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008540 | OEIS-A236691

Number of totally symmetric solid partitions which fit in an n X n X n X n box.

There is no conjectured formula for a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Graham H. Hawkes_, Jan 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008541 | OEIS-A236754

Number of irregular spirals (non-expanded loop) appearing with n center points.

Conjecture: (1) For odd n all forms are non-expanded loops and equal to n!. (2) For n mod 4 = 0, there are both non-expanded loops and expanded spirals.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jan 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008542 | OEIS-A236755

Number of irregular spiral (expanded spiral) appearing with n center points.

Conjecture: (1) For even n; except n mod 4 = 0; all forms are expanded spirals and equal to n!. (2) For n mod 4 = 0, there are both non-expanded loops and expanded spirals.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jan 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008543 | OEIS-A236765

Number of ways to write n = k^2 + m with k > 1 and m > 1 such that sigma(k^2) + prime(m) - 1 is prime, where sigma(j) denotes the sum of all positive divisors of j.

Conjecture: (i) If n > 6 is not equal to 18, then a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008544 | OEIS-A236831

Number of ordered ways to write n = p + q with q > 0 such that p, p + 2 and p + prime(q) + 1 are all prime.

Conjecture: a(n) > 0 for all n > 13.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 31 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008545 | OEIS-A236954

E.g.f. satisfies: A'(x) = A(x)^7 * A(-x) with A(0) = 1.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008546 | OEIS-A236968

Number of ordered ways to write n = k + m with k > 0 and m > 0 such that 6*k - 1, 6*k + 1 and k + phi(m) are all prime, where phi(.) is Euler's totient function.

Conjecture: (i) a(n) > 0 for all n > 1. Also, any n > 12 can be written as k + m (k > 0 and m > 2) with 6*k - 1, 6*k + 1 and k + phi(m)/2 all prime. Clearly, part (i) of the conjecture implies that any integer n > 1 can be written as p + m - phi(m), where p is a prime and m is a positive integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008547 | OEIS-A237016

a(n) = |{0 < k < n: phi(k)*sigma(n-k) is a square}|, where phi(.) is Euler's totient function and sigma(j) is the sum of all positive divisors of j.

Conjecture: (i) a(n) > 0 except for n = 1, 7, 17. See also A236998 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008548 | OEIS-A237046

Numbers k such that sigma(k) < 2*k-1.

If the only least-deficient numbers are the powers of 2 (which is an open problem) then the union of A000079 and A023196 and this sequence gives A000027 (see also A103288).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Feb 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008549 | OEIS-A237049

Number of ordered ways to write n = i + j + k (0 < i <= j <= k) with i,j,k not all equal such that sigma(i)*sigma(j)*sigma(k) is a cube, where sigma(m) denotes the sum of all positive divisors of m.

Conjecture: For every k = 2, 3, ... there is a positive integer N(k) such that any integer n > N(k) can be written as n_1 + n_2 + ... + n_k with n_1, n_2, ..., n_k positive and distinct such that the product sigma(n_1)*sigma(n_2)*...*sigma(n_k) is a k-th power. In particular, we may take N(2) = 309, N(3) = 42, N(4) = 25, N(5) = 24, N(6) = 27 and N(7) = 32. This is similar to the conjecture in A233386.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008550 | OEIS-A237050

Number of ways to write n = i_1 + i_2 + i_3 + i_4 + i_5 (0 < i_1 <= i_2 <= i_3 <= i_4 <= i_5) with i_1, i_2, ..., i_5 not all equal such that the product i_1*i_2*i_3*i_4*i_5 is a fifth power.

Conjecture: For each k = 3, 4, ... there is a positive integer M(k) such that any integer n > M(k) can be written as i_1 + i_2 + ... + i_k with i_1, i_2, ..., i_k positive and not all equal such that the product i_1*i_2*...*i_k is a k-th power. In particular, we may take M(3) = 486, M(4) = 23, M(5) = 26, M(6) = 36 and M(7) = 31. This is motivated by the conjectures in A233386 and A237049.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008551 | OEIS-A237053

Smallest number k such that some subset of n+1..n+k can be summed and added to n to produce a prime.

Conjecture: every nonnegative integer except 2 occurs infinitely often in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Feb 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008552 | OEIS-A237123

Number of ways to write n = i + j + k with 0 < i < j < k such that phi(i), phi(j) and phi(k) are all cubes, where phi(.) is Euler's totient function.

Conjecture: For each k = 2, 3, ... there is a positive integer s(k) such that any integer n >= s(k) can be written as i_1 + i_2 + ... + i_k with 0 < i_1 < i_2 < ... < i_k such that all those phi(i_1), phi(i_2), ..., phi(i_k) are k-th powers. In particular, we may take s(2) = 70640, s(3) = 935 and s(4) = 3273.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008553 | OEIS-A237203

Least positive integer k < n/2 with phi(k)*phi(n-k) a square, or 0 if such a number k does not exist.

Conjecture: a(n) < sqrt(n)*log(2*n) for all n > 0. According to the conjecture in A236998, a(n) should be positive for all n > 8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008554 | OEIS-A237259

Least positive integer k < n such that prime(k) + 2 and prime(prime(n-k)) + 2 are both prime, or 0 if such a number k does not exist.

Conjecture: a(n) < 2*sqrt(n)*log(3*n) for all n > 0. According to the conjecture in A218829, a(n) should be positive for all n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008555 | OEIS-A237260

Least positive integer m < n with prime(prime(m)) + 2 and prime(n-m) + 2 both prime, or 0 if such a number m does not exist.

Conjecture: a(n) < sqrt(6*n)*log(3*n) for all n > 0. According to the conjecture in A218829, a(n) should be positive for all n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008556 | OEIS-A237274

a(n) = A236283(n) mod 9.

(Conjecture) This has period 18: repeat 2, 1, 4, 5, 1, 4, 2, 7, 7, 5, 7, 7, 2, 4, 1, 5, 4, 1. The sorted terms in the conjectured period are 1, 1, 1, 1, 2, 2, 2, 4, 4, 4, 4, 5, 5, 5, 7, 7, 7, 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Feb 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008557 | OEIS-A237275

Smallest k divisible by the n-th power of its last decimal digit > 1.

Conjecture: a(n) == 2 (mod 10).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008558 | OEIS-A237283

Primes p with prime(prime(p)) + 2 also prime.

This sequence is interesting because of the conjecture in A237253.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008559 | OEIS-A237367

Number of ordered ways to write n = k + m with k > 0 and m > 0 such that 2*k - 1, prime(k)^2 - 2 and prime(m)^2 - 2 are all prime.

Conjecture: (i) a(n) > 0 for all n > 2, and a(n) = 1 only for n = 3, 73, 81, 534. Clearly, part (i) of this conjecture implies that there are infinitely many primes p with p^2 - 2 also prime. Similar comments apply to parts (ii) and (iii).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008560 | OEIS-A237414

Primes p with p^2 - 2 and prime(p)^2 - 2 both prime.

According to the conjecture in A237413, this sequence should have infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008561 | OEIS-A237424

Numbers of the form (10^a + 10^b + 1)/3.

Conjecture: This sequence is in a sense the maximally dense sequence with this nondecreasing products property. That is, it appears that every maximal sequence is either (i) A237424, (ii) a finite set of "extra" terms plus A237424 restricted to b=0 (which is A093137), or (iii) a finite set of "extra" terms plus A237424 restricted to a=b (which is A067275). (There might be one more case, not yet identified.) - _David Applegate_, Feb 12 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ahmad J. Masad_, Feb 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008562 | OEIS-A237453

Number of primes p < n with p*n + pi(p) prime, where pi(.) is given by A000720.

Conjecture: (i) a(n) > 0 for all n > 4, and a(n) = 1 for no n > 144. Moreover, for any positive integer n, there is a prime p < sqrt(2*n)*log(5n) with p*n + pi(p) prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008563 | OEIS-A237496

Number of ordered ways to write n = k + m (0 < k <= m) with pi(k) + pi(m) - 2 prime, where pi(.) is given by A000720.

Conjecture: (i) a(n) > 0 for all n > 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008564 | OEIS-A237497

a(n) = |{0 < k <= n/2: pi(k*(n-k)) is prime}|, where pi(.) is given by A000720.

Conjecture: a(n) > 0 for all n > 10, and a(n) = 1 for no n > 51. Moreover, for any integer n > 10, there is a positive integer k < n with 2*k + 1 and pi(k*(n-k)) both prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008565 | OEIS-A237515

Decimal expansion of the sum of reciprocals of the strict partition function (the function giving the number of partitions of an integer into distinct parts).

Conjecture: This number is transcendental. - _Zhi-Wei Sun_, May 24 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Feb 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008566 | OEIS-A237518

Least primes that together with prime(n) forms a Heronian triangle, starting at n = 2.

It has been proved that for every integer i > 2 there exists an infinite series of side pairs (j, k) that together with i form a Heronian triangle. It is conjectured that for every prime(n) where n > 1 there exists an infinite series of side pairs (p, q) that together with prime(n) form a Heronian triangle such that either p or q is also prime. See A230666 and A233232 for prime(2) and prime(3). a(n) is the sequence of least such primes for prime(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Feb 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008567 | OEIS-A237523

a(n) = |{0 < k < n/2: phi(k*(n-k)) + 1 is a square}|, where phi(.) is Euler's totient function.

Conjecture: (i) a(n) > 0 for all n > 7, and a(n) = 1 only for n = 8, 9, 13, 15, 20, 132. See also A237524 for a similar conjecture involving higher powers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008568 | OEIS-A237524

Number of ordered ways to write n = i + j + k with 0 < i <= j <= k such that phi(i*j*k) is a cube, where phi(.) is Euler's totient function.

Conjecture: For each k = 3, 4, ..., any integer n > 2*k + 1 can be written as a sum of k positive integers n_1, n_2, ..., n_k such that phi(n_1*n_2*...*n_k) is a k-th power. See also A237523 for a similar conjecture with k = 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008569 | OEIS-A237578

a(n) = |{0 < k < n: pi(k*n) is prime}|, where pi(.) is given by A000720.

Conjecture: a(n) > 0 for all n > 2, and a(n) = 1 only for n = 5, 8, 13. Moreover, for each n = 1, 2, 3, ..., there is a positive integer k < 3*sqrt(n) + 3 with pi(k*n) prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008570 | OEIS-A237582

a(n) = |{0 < k < n: pi(n + k^2) is prime}|, where pi(.) is given by A000720.

Conjecture: (i) For each a = 2, 3, ... there is a positive integer N(a) such that for any integer n > N(a) there is a positive integer k < n with pi(n + k^a) prime. In particular, we may take (N(2), N(3), ..., N(9)) = (1, 1, 9, 26, 8, 9, 18, 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008571 | OEIS-A237588

a(n) = sigma(n) - 2n + 1.

Conjecture: a(n) = 0 iff n is a power of 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Feb 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008572 | OEIS-A237590

a(n) is the total number of regions (or parts) after n-th stage in the diagram of the symmetries of sigma described in A236104.

Conjecture: a(n) is the total number of 2-dense sublists of divisors of all positive integers <= n. - _Omar E. Pol_, Oct 07 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Mar 31 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008573 | OEIS-A237593

Triangle read by rows in which row n lists the elements of the n-th row of A237591 followed by the same elements in reverse order.

Conjecture 1: the area under the n-th Dyck path equals A024916(n), the sum of all divisors of all positive integers <= n. If the conjecture is true then the n-th Dyck path represents the boundary segments after the alternating sum of the elements of the n-th row of A236104. Conjecture 2: two adjacent Dyck paths never cross (checked by hand up to n = 128), hence the total area between the n-th Dyck path and the (n-1)-st Dyck path is equal to sigma(n) = A000203(n), the sum of divisors of n. Checked _Michel Marcus_'s conjecture with two Mathematica functions up to n=100000, for more information see A240542. - _Hartmut F. W. Hoft_, Jul 17 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Feb 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008574 | OEIS-A237595

a(n) = |{1 <= k <= n: n + pi(k^2) is prime}|, where pi(.) is given by A000720.

Conjecture: (i) a(n) > 0 for all n > 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008575 | OEIS-A237597

a(n) = |{0 < k < prime(n): n divides pi(k*n)}|, where pi(.) is given by A000720.

Conjecture: a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008576 | OEIS-A237598

a(n) = |{0 < k < prime(n): pi(k*n) is a square}|, where pi(.) is given by A000720.

Conjecture: (i) a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008577 | OEIS-A237606

Positive integers k such that x^2 - 8xy + y^2 + k = 0 has integer solutions.

Even powers of terms as well as products of an even number of terms belong to A243188. This can be proved with respect to the forms [a,0,-c] and [a, 0, +c] by the following identities: (au^2 - cv^2)(ax^2 - cy^2) = (aux + cvy)^2 - ac(uy + vx)^2 and (au^2 + cv^2)(ax^2 + cy^2) = (aux - cvy)^2 + ac(uy + vx)^2 for all a, c, u, v, x, y in R. This can be verified by expanding both sides of the equations. Generalization (conjecture): This multiplication rule applies to all sequences represented by any binary quadratic form [a, b, c]. Odd powers of terms as well as products of an odd number of terms belong to the sequence. This can be proved with respect to the forms [a,0,-c] and [a, 0, +c] by the following identities: (as^2 - ct^2)(au^2 - cv^2)(ax^2 - cy^2) = a[s(aux + cvy) + ct(uy + vx)]^2 - c[as(uy + vx) + t(aux + cvy)]^2 and (as^2 + ct^2)(au^2 + cv^2)(ax^2 + cy^2) = a[s(aux - cvy) - ct(uy + vx)]^2 + c[as(uy + vx) + t(aux - cvy)]^2 for all a, c, s, t, u, v, x, y in R. This can be verified by expanding both sides of the equations. Generalization (conjecture): This multiplication rule applies to all sequences represented by any binary quadratic form [a, b, c]. Generalization (conjecture): If there are three sequences of a given positive discriminant that are represented by the forms [a1, b1, c1], [a2, b2, c2] and [a1*a2, b3, c3] for a1, a2 != 1, then the BCD rules apply to these sequences. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Barker_, Feb 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008578 | OEIS-A237612

Least positive integer k such that A000720(k*n) is a square, or 0 if such a number k does not exist.

According to the conjecture in A237598, we should always have 0 < a(n) < prime(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008579 | OEIS-A237614

Least positive integer k with A000720(k*n) divisible by n, or 0 if such a number k does not exist.

According to the conjecture in A237597, we should always have 0 < a(n) < prime(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008580 | OEIS-A237615

a(n) = |{0 < k < n: k^2 + k - 1 and pi(k*n) are both prime}|, where pi(.) is given by A000720.

Conjecture: (i) a(n) > 0 for all n > 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008581 | OEIS-A237643

Least positive integer m such that {A000720(k*n): k = 1, ..., m} contains a complete system of residues modulo n, or 0 if such a number m does not exist.

Conjecture: a(n) is always positive. Moreover, a(n) <= 2*prime(n) for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008582 | OEIS-A237656

Least positive integer m such that {A000720(k^2): k = 1, ..., m} contains a complete system of residues modulo n, or 0 if such a number m does not exist.

Conjecture: a(n) is always positive. Moreover, a(n) < 2*prime(n+1) - 2 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008583 | OEIS-A237657

a(n) = |{n < m < 2*n: pi(m) and pi(m^2) are both prime}|, where pi(.) is given by A000720.

Conjecture: (i) a(n) > 0 for all n > 8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008584 | OEIS-A237658

Positive integers m with pi(m) and pi(m^2) both prime, where pi(.) is given by A000720.

The conjecture in A237657 implies that this sequence has infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008585 | OEIS-A237659

Primes p with pi(p) and pi(p^2) both prime, where pi(.) is given by A000720.

Conjecture: The sequence has infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008586 | OEIS-A237665

Number of partitions of n such that the distinct terms arranged in increasing order form a string of two or more consecutive integers.

Conjecture: a(n) = sum of smallest parts in the distinct partitions of n with an even number of parts. - _George Beck_, May 06 2017 The preceding conjecture is correct (see Chern, 2019). - _Andrew R. Epstein_, Jul 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008587 | OEIS-A237666

Number of partitions of n that include a pair of consecutive integers.

Conjecture: for n > 0, a(n) = A000041(n) - A116931(n). - _Vaclav Kotesovec_, Jan 28 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008588 | OEIS-A237671

Let m_n denote the number which is obtained from n-base representation of m if its digits are written in nondecreasing order; then a(n) is the smallest period of the sequence which is defined by the recurrence b(0)=0, b(1)=1, b(k)=(b(k-1) + b(k-2))_n, for k>=2, or a(n)=0, if there is no such period.

We conjecture that the sequence b is always eventually periodic, and so a(n)>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Feb 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008589 | OEIS-A237684

a(n) = floor(n*prime(n) / Sum_{i<=n} prime(i)).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Feb 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008590 | OEIS-A237685

Number of partitions of n having depth 1; see Comments.

Suppose that P is a partition of n. Let x(1), x(2),...,x(k) be the distinct parts of P, and let m(i) be the multiplicity of x(i) in P. Let f(P) be the partition [m(1)*x(1), m(2)*x(2),...,x(k)*m(k)] of n. Define c(0,P) = P, c(1,P) = f(P), ..., c(n,P) = f(c(n-1,P)), and define d(P) = least n such that c(n,P) has no repeated parts; d(P) is introduced here as the depth of P. Clearly d(P) = 0 if and only if P is a strict partition, as in A000009. Conjecture: if d >= 0, then 2^d is the least n that has a partition of depth d.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008591 | OEIS-A237687

Primes p with pi(p), pi(pi(p)) and pi(p^2) all prime, where pi(.) is given by A000720.

Conjecture: The sequence has infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008592 | OEIS-A237705

Number of primes p < n with pi(n-p) prime, where pi(.) is given by A000720.

Conjecture: (i) a(n) > 0 for all n > 4, and a(n) = 1 only for n = 5, 12, 26, 27, 32, 68.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008593 | OEIS-A237706

Number of primes p < n with pi(n-p) a square, where pi(.) is given by A000720.

Conjecture: (i) a(n) > 0 for all n > 2, and a(n) = 1 only for n = 3, 5, 6, 7, 8, 16, 17, 22, 23, 148, 149. See also A237705, A237720 and A237721 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008594 | OEIS-A237710

Least prime p < n with pi(n-p) a square, or 0 if such a prime p does not exist.

According to the conjecture in A237706, a(n) should be positive for all n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008595 | OEIS-A237712

a(n) = |{0 < k < n: k*n + pi(k*n) is prime}|, where pi(.) is given by A000720.

Conjecture: a(n) > 0 for all n > 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008596 | OEIS-A237715

Number of ordered ways to write n = p + q (q > 0) with p, prime(p) - p + 1 and prime(prime(q)) - prime(q) + 1 all prime.

Conjecture: a(n) > 0 for all n > 2, and a(n) = 1 only for n = 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008597 | OEIS-A237720

Number of primes p <= (n+1)/2 with floor(sqrt(n-p)) prime.

Conjecture: (i) a(n) > 0 for all n > 5, and a(n) = 1 only for n = 6, 23, 24, 111, 112, ..., 120. See also A237705, A237706 and A237721 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008598 | OEIS-A237721

Number of primes p <= n with floor(sqrt(n-p)) a square.

Conjecture: a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 9, 10, 11, 12, 15, 16, 17. See also A237705, A237706 and A237720 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008599 | OEIS-A237757

Number of partitions of n such that 2*(least part) = (number of parts).

Conjectural g.f.: Sum_{n >= 0} q^(2*(n+1)^2)/Product_{k = 1..2*n+1} 1 - q^k. - _Peter Bala_, Feb 02 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008600 | OEIS-A237768

Number of primes p < n with pi(n-p) a Sophie Germain prime, where pi(.) is given by A000720.

Conjecture: a(n) > 0 for all n > 4, and a(n) = 1 only for n = 5, 12, 20, 21, 26, 27, 30, 31, 32, 60, 61, 68, 69, 80, 81. This is stronger than part (i) of the conjecture in A237705.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008601 | OEIS-A237769

Number of primes p < n with pi(n-p) - 1 and pi(n-p) + 1 both prime, where pi(.) is given by A000720.

Conjecture: (i) a(n) > 0 for all n > 8, and a(n) = 1 only for n = 9, 34, 35.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008602 | OEIS-A237840

a(n) = |{0 < k <= n: the number of twin prime pairs not exceeding k*n is a square}|.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 for no n > 159.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008603 | OEIS-A237879

Least positive integer k <= n such that the number of twin prime pairs not exceeding k*n is a square, or 0 if such a number k does not exist.

According to the conjecture in A237840, a(n) should be always positive.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008604 | OEIS-A237880

Conjectured number of distinct integers < 10^n that are Sierpiński or Riesel or simultaneously Sierpiński and Riesel numbers.

Conjectured number of distinct integers < 10^n that are Sierpiński or Riesel or simultaneously Sierpiński and Riesel numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Feb 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008605 | OEIS-A237930

a(n) = 3^(n+1) + (3^n-1)/2.

First differences of A108765. The digital root of a(n) for n > 1 is always 4. a(n) is never divisible by 7 or by 12. a(n) == 10 (mod 84) for odd n. a(n) == 31 (mod 84) for even n > 0. Conjecture: This sequence contains no prime factors p == {11, 13, 23, 61 71, 73} (mod 84). - _Klaus Purath_, Apr 13 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Delélham_, Feb 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008606 | OEIS-A237975

Least nonnegative integer m such that for some k = 1, ..., n there are exactly m^2 twin prime pairs not exceeding k*n.

The conjecture in A237840 implies that a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008607 | OEIS-A238003

Number of partitions of n not having depth 1; see Comments.

The depth of a partition is defined at A237685 as follows. Suppose that P is a partition of n. Let x(1), x(2), ..., x(k) be the distinct parts of P, and let m(i) be the multiplicity of x(i) in P. Let f(P) be the partition [m(1)*x(1), m(2)*x(2), ..., x(k)*m(k)] of n. Define c(0,P) = P, c(1,P) = f(P), ..., c(n,P) = f(c(n-1,P)), and define d(P) = least n such that c(n,P) has no repeated parts; d(P) is the depth of P. Conjecture: lim_{n->infinity} a(n)/A000041(n) = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008608 | OEIS-A238005

Number of partitions of n into distinct parts such that (greatest part) - (least part) = (number of parts).

This theorem was conjectured by _Omar E. Pol_ in February 2018, and proved independently by _William J. Keith_ and _Roland Bacher_ on Sep 05 2021. The elegant proof given in the link below is due to _Don Reble_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008609 | OEIS-A238016

Number A(n,k) of partitions of n^k into parts that are at most n; square array A(n,k), n>=0, k>=0, read by antidiagonals.

Conjecture: If f(n) >= O(n^4) then "number of partitions of f(n) into parts that are at most n" is asymptotic to f(n)^(n-1) / (n!*(n-1)!). See also A237998, A238000, A236810 or A258668-A258672. - _Vaclav Kotesovec_, Jun 07 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Feb 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008610 | OEIS-A238165

Number of pairs {j, k} with 0 < j < k <= n such that pi(j*n) divides pi(k*n), where pi(.) is given by A000720.

Conjecture: (i) a(n) > 0 for all n > 1. See also A238224 for a refinement of part (i) of this conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008611 | OEIS-A238194

Conjectured numbers n for which n^n + (-1)^n (n-1)^(n-1) is not squarefree.

Conjectured numbers n for which n^n + (-1)^n (n-1)^(n-1) is not squarefree. The first case (130) yields a number divisible by 83^2. The next 5 terms yield numbers divisible by 59^2. Boyd et al. are not completely certain about the other 994 numbers up to 1000. They conjecture that 0.9934466... of numbers n^n + (-1)^n (n-1)^(n-1) are squarefree.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Feb 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008612 | OEIS-A238208

The total number of 1's in all partitions of n into an odd number of distinct parts.

Faster converging g.f. (conjecture): Sum_{n >= 0} x^((n+1)*(2*n+1))/ Product_{k = 1..2*n} 1 - x^k. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mircea Merca_, Feb 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008613 | OEIS-A238224

Number of pairs {j, k} with 0 < j < k <= n and k == 1 (mod j) such that pi(j*n) divides pi(k*n), where pi(.) is given by A000720.

Conjecture: a(n) > 0 for all n > 1. This is a refinement of part (i) of the conjecture in A238165. We have verified the conjecture for n up to 21500.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008614 | OEIS-A238256

A060308 begins with one 2, one 3, one 5, two 7's, one 11, two 13's, i.e., d(n) = 1, 1, 1, 2, 1, 2, 1, 2, 3, 1,... times the primes (A000040). a(n) uses this distribution with noncomposites (A008578).

Conjecture: a(n) is the greatest noncomposite (A008578) dividing A238691(n-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Mar 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008615 | OEIS-A238266

a(n) is the largest number that can be written in the form a(n) = 2^k1 * p1^k2 + 2^k3 * p2^k4 in n distinct ways, where p1 and p2 are odd prime numbers and k1, k2, k3, and k4 are nonnegative integers.

It is conjectured that, for any positive integer n, there exist only a finite number of positive integers that can be written in the form 2^k1 * p1^k2 + 2^k3 * p2^k4 in n distinct ways.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Feb 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008616 | OEIS-A238267

The number of integers that can be written in the form 2^k1 * p1^k2 + 2^k3 * p2^k4 in n distinct ways, where p1 and p2 are odd prime numbers and k1, k2, k3, and k4 are nonnegative integers.

It is conjectured that a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Feb 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008617 | OEIS-A238268

The number of unordered ways that n can be written as the sum of two numbers of the form p or 2p, where p is prime.

It is conjectured that all terms of this sequence are greater than zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Feb 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008618 | OEIS-A238277

a(n) = |{0 <= k < n: the number of primes in the interval (k*n, (k+1)*n] is a square}|.

Conjecture: a(n) > 0 for all n > 0. See also A238278 and A238281 for related conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008619 | OEIS-A238278

a(n) = |{0 < k < n: the number of primes in the interval ((k-1)*n, k*n] and the number of primes in the interval (k*n, (k+1)*n] are both prime}|.

Conjecture: (i) a(n) > 0 for all n > 3. See also A238277 and A238281 for related conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008620 | OEIS-A238281

a(n) = |{0 < k < n: the two intervals (k*n, (k+1)*n) and ((k+1)*n, (k+2)*n) contain the same number of primes}|.

Conjecture: (i) a(n) > 0 for all n > 1. Moreover, if n > 1 is not equal to 8, then there is a positive integer k < n with 2*k + 1 prime such that the two intervals ((k-1)*n, k*n) and (k*n, (k+1)*n) contain the same number of primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008621 | OEIS-A238289

Least positive integer k such that prime(k*n), prime((k+1)*n) and prime((k+2)*n) form an arithmetic progression, or 0 if such a number k does not exist.

Conjecture: (i) We always have 0 < a(n) < 3*prime(n) + 9. The conjecture is a refinement of the Green-Tao theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008622 | OEIS-A238373

The number of "topped inward" shuffles to reorder a stack of n cards to its original order.

Conjecture (equivalent to the Pfister comment in A216066): if the shuffle were defined again in inward order, but with the odd numbers to the left and the even numbers to the right, A003558 becomes the number of shuffles to return to original order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Feb 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008623 | OEIS-A238386

a(n) = |{0 < k < n-1: p = prime(k) + pi(n-k) and p + 2 are both prime}|, where pi(.) is given by A000720.

Conjecture: (i) a(n) > 0 for all n > 10. We have verified part (i) of the conjecture for n up to 10^7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008624 | OEIS-A238393

a(n) = |{0 < k <= n: 2*p(k)*p(n) + 1 is prime}|, where p(.) is the partition function (A000041).

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 2, 13.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008625 | OEIS-A238405

a(n) = |{0 < k < n: prime(k) + pi(n-k) is a triangular number}|, where pi(.) is given by A000720.

Conjecture: a(n) > 0 for all n > 6, and a(n) = 1 only for n = 5, 7, 13, 16, 18, 24, 29, 71, 72, 158.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008626 | OEIS-A238457

a(n) = |{0 < k <= n: p(n) + k is prime}|, where p(.) is the partition function (A000041).

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 2, 3, 4, 5, 30, 109. Also, for each n > 2 there is a positive integer k <= n+3 such that p(n) - k is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008627 | OEIS-A238458

Number of primes p < n with 2*P(n-p) + 1 prime, where P(.) is the partition function (A000041).

Conjecture: a(n) > 0 for all n > 2. Also, for each n > 3 there is a prime p < n with 2*P(n-p) - 1 prime. We have verified the conjecture for n up to 10^5. See also A238459 for a similar conjecture involving the strict partition function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008628 | OEIS-A238459

Number of primes p < n with q(n-p) + 1 prime, where q(.) is the strict partition function (A000009).

Conjecture: a(n) > 0 for all n > 2. Also, for each n > 6 there is a prime p < n with q(n-p) - 1 prime. We have verified the conjecture for n up to 10^5. See also A238458 for a similar conjecture involving the partition function p(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008629 | OEIS-A238504

Number of primes p <= n with pi(pi((p-1)*n)) prime, where pi(x) denotes the number of primes not exceeding x.

Conjecture: (i) a(n) > 0 for all n > 2. See also A238902 for another conjecture involving pi(pi(x)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008630 | OEIS-A238506

Number of primes of the form: |P_i +- P_n / P_i| or |P_i/2 +- 2P_n / P_i|, where 2 <= i <= n, P_i = the i-th primorial number (A002110).

It is conjectured that all terms of this sequence are greater than zero, but this conjecture might have exceptions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Feb 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008631 | OEIS-A238509

a(n) = |{0 < k < n: p(n) + p(k) - 1 is prime}|, where p(.) is the partition function (A000041).

Conjecture: (i) a(n) > 0 for all n > 1. Also, if n > 2 is different from 8 and 25, then p(n) + p(k) + 1 is prime for some 0 < k < n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008632 | OEIS-A238516

a(n) = |{0 < k < n: (p(k)+1)*p(n) + 1 is prime}|, where p(.) is the partition function (A000041).

Conjecture: (i) a(n) > 0 for all n > 1. Also, for any integer n > 4 there is a positive integer k < n with (p(k)-1)*p(n) - 1 prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008633 | OEIS-A238549

a(n) is one fourth of the total number of free ends of 4 line segments expansion at n iterations (see Comments lines for definition).

We take the official definition to be that provided by the PARI program. From this the assertions in the Formula section follow (they were formerly stated as conjectures). - _N. J. A. Sloane_, Feb 24 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, May 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008634 | OEIS-A238568

a(n) = |{0 < k < n: n^2 - pi(k*n) is prime}|, where pi(x) denotes the number of primes not exceeding x.

Conjecture: (i) a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 3, 4, 8, 10, 24, 41.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008635 | OEIS-A238570

a(n) = |{0 < k < n: pi((k+1)^2) - pi(k^2) and pi(n^2) - pi(k^2) are both prime}|, where pi(x) denotes the number of primes not exceeding x.

Conjecture: (i) a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008636 | OEIS-A238576

Number of odd primes p < 2*n with prime(n*(p-1)/2)^2 - 2 prime.

Conjecture: (i) a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 7, 23, 61, 73. Clearly, either part of the conjecture implies that there are infinitely many primes of the form p^2 - 2 with p prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008637 | OEIS-A238577

a(n) = |{0 < k <= n: p(n)*q(k)*r(k) + 1 is prime}|, where p(.), q(.) and r(.) are given by A000041, A000009 and A047967 respectively.

Conjecture: (i) a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 3, 5, 20. If n > 2, then p(n)*q(k)*r(k) - 1 is prime for some k = 1, ..., n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008638 | OEIS-A238580

a(n) = |{0 < k <= n: 2*k + 1 and prime(k)*prime(n) - 2 are both prime}|.

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 2, 4, 7, 8, 10, 28, 34,37, 77.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008639 | OEIS-A238585

Number of primes p < n with prime(p)^2 + (prime(n)-1)^2 prime.

Conjecture: (i) a(n) > 0 unless n divides 6, and a(n) = 1 only for n = 4, 5, 7, 10, 11, 12, 19, 21, 22, 31, 42, 44.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008640 | OEIS-A238597

Number of primes p < 2*n with 2*pi(p) + 1 and p*(2n-1) - 2 both prime, where pi(.) is given by A000720.

Conjecture: (i) a(n) > 0 for all n > 1, and a(n) = 1 for no n > 195. Part (i) of this conjecture is an extension of the conjecture in A238580.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008641 | OEIS-A238643

Number of primes p <= n such that 2*pi(p) - (-1)^n and p*n +((-1)^n - 3)/2 are both prime, where pi(x) is the number of primes not exceeding x.

Conjecture: (i) a(n) > 0 for all n > 2. Part (i) of the conjecture is a further extension of the conjecture in A238597 to cover the even case.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008642 | OEIS-A238645

Number of odd primes p < 2*n such that the number of squarefree numbers among 1, ..., ((p-1)/2)*n is prime.

Conjecture: a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 4, 7, 14, 19, 29. This is an analog of the conjecture in A237578 for squarefree numbers. We have verified it for n up to 20000. See also A238646 for a similar conjecture involving squarefree numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008643 | OEIS-A238646

Number of primes p < n such that the number of squarefree numbers among 1, ..., n-p is prime.

Conjecture: a(n) > 0 for all n > 3, and a(n) = 1 only for n = 4, 10, 12, 14, 16, 24, 36. This is analog of the conjecture in A237705 for squarefree numbers. We have verified the conjecture for n up to 60000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008644 | OEIS-A238695

Decimal expansion of Product_{k>=0} (1+1/k!).

Conjectured to be irrational, transcendental and normal, none have been shown. Product is sometimes taken from n=1, leading to half the stated value.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frederick Reckless_, Mar 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008645 | OEIS-A238703

a(n) = |{0 < k < n: floor(k*n/3) is prime}|.

Conjecture: If n > m > 0 with n not divisible by m, then floor(k*n/m) is prime for some 0 < k < n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008646 | OEIS-A238732

Number of primes p < n with floor((n-p)/3) a square.

Conjecture: (i) For any integers m > 2 and n > 2, there is a prime p < n such that floor((n-p)/m) is a square.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008647 | OEIS-A238733

Number of primes p < n such that floor((n-p)/3) = (q-1)*(q-3)/8 for some prime q.

Conjecture: (i) For any integers m > 2 and n > 2, there is a prime p < n such that floor((n-p)/m) has the form (q-1)*(q-3)/8 with q an odd prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008648 | OEIS-A238756

Number of ordered ways to write n = k + m (k > 0 and m > 0) such that 2*k + 1, prime(prime(k)) - prime(k) + 1 and prime(prime(m)) - prime(m) + 1 are all prime.

Conjecture: a(n) > 0 for all n > 1. The conjecture suggests that there are infinitely many primes p with 2*pi(p) + 1 and prime(p) - p + 1 both prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008649 | OEIS-A238766

Number of ordered ways to write n = k + m (k > 0 and m > 0) such that prime(prime(k)) - prime(k) + 1, prime(prime(2*k+1)) - prime(2*k+1) + 1 and prime(prime(m)) - prime(m) + 1 are all prime.

Conjecture: a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 3, 11, 44, 55, 149, 371.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008650 | OEIS-A238776

Primes p with prime(p) - p + 1 and prime(q) - q + 1 both prime, where q = prime(2*pi(p)+1) with pi(.) given by A000720.

Conjecture: The sequence has infinitely many terms. This is motivated by the conjecture in A238766. Note that the sequence is a subsequence of A234695.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008651 | OEIS-A238802

Number T(n,k) of standard Young tableaux with n cells where k is the length of the maximal consecutive sequence 1,2,...,k in the first column; triangle T(n,k), n>=0, 0<=k<=n, read by rows.

Conjecture: Generally, column k is asymptotic to sqrt(2)/(2*(k+1)*(k-1)!) * exp(sqrt(n)-n/2-1/4) * n^(n/2) * (1 + 7/(24*sqrt(n))), holds for all k<=10. - _Vaclav Kotesovec_, Mar 08 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_ and _Alois P. Heinz_, Mar 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008652 | OEIS-A238806

Number of n X 2 0..2 arrays with no element equal to one plus the sum of elements to its left or one plus the sum of the sum of elements above it, modulo 3.

Conjectures from _Colin Barker_, Oct 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008653 | OEIS-A238807

Number of n X 3 0..2 arrays with no element equal to one plus the sum of elements to its left or one plus the sum of the sum of elements above it, modulo 3.

Conjectures from _Colin Barker_, Oct 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008654 | OEIS-A238814

Primes p with prime(p) - p + 1 and prime(q) - q + 1 both prime, where q is the first prime after p.

Conjecture: The sequence is infinite, in other words, A234695 contains infinitely many consecutive prime pairs prime(k) and prime(k+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008655 | OEIS-A238822

Number of equilateral triangles bounded by the sides and diagonals of a regular 3n-gon.

Conjecture: For odd n, a(n) = n*(n+1)^3/2 - 3*n^2. - _Vilim Lendvaj_, Jul 24 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008656 | OEIS-A238856

Number of digits of the largest right-truncatable reversible prime in base n.

For definitions and more comments, see A238854 and A238850. A weak conjecture: this sequence might be bounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Mar 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008657 | OEIS-A238873

Number of superdiagonal partitions: partitions (p1, p2, p3, ...) of n such that pi >= i.

Conjecture: Also the number of integer partitions y of n with choosable initial intervals, meaning it is possible to choose distinct positive integers (z_1, z_2, ...) such that z_i <= y_i for all i. - _Gus Wiseman_, Sep 26 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Mar 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008658 | OEIS-A238879

Row sums of the triangle of generalized ballot numbers A238762.

a(2n) = A000108(n), a(2n+1) = A002054(n) (conjectured). - _Ralf Stephan_, Mar 14 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Mar 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008659 | OEIS-A238881

Number of odd primes p < 2*n with prime(n*(p+1)/2) + n*(p+1)/2 prime.

Conjecture: (i) a(n) > 0 for all n > 5, and a(n) = 1 only for n = 2, 3, 7, 9, 23. Moreover, for any r = 1,-1 and n > 5*(2+r) there is a positive integer k < n such that 2*k+r and prime(k*n)+k*n are both prime. This conjecture implies that there are infinitely many positive integers m with prime(m) + m (or prime(m) - m) prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008660 | OEIS-A238890

a(n) = |{0 < k <= n: prime(k*n) - pi(k*n) is prime}|, where pi(x) denotes the number of primes not exceeding x.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 for no n > 28. The conjecture implies that there are infinitely many primes p with p - pi(pi(p)) (or p + pi(pi(p))) prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008661 | OEIS-A238902

a(n) = |{0 < k <= n: pi(pi(k*n)) is a square}|, where pi(x) denotes the number of primes not exceeding x.

Conjecture: (i) a(n) > 0 for all n > 0. See A239884 for a sequence related to part (i) of the conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008662 | OEIS-A238923

Number of (n+1) X (1+1) 0..3 arrays with no element greater than all horizontal neighbors or equal to all vertical neighbors.

Conjectures from _Colin Barker_, Oct 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008663 | OEIS-A238936

Powers of 6 without the digit '0' in their decimal expansion.

Conjectured to be finite and complete. See the OEIS wiki page for further information, references and links.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Mar 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008664 | OEIS-A238977

Number of ballot sequences of length n with exactly 2 fixed points.

Conjecture: a(n+3) == 1 (mod n) iff n is coprime to 2 and 3, that is, iff n is a term of A007310. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_ and _Alois P. Heinz_, Mar 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008665 | OEIS-A238979

Number of ballot sequences of length n with exactly 4 fixed points.

Conjecture: Generally, for fixed k is column k of A238802 asymptotic to sqrt(2)/(2*(k+1)*(k-1)!) * exp(sqrt(n)-n/2-1/4) * n^(n/2) * (1 + 7/(24*sqrt(n))), holds for all k<=10. - _Vaclav Kotesovec_, Mar 08 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_ and _Alois P. Heinz_, Mar 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008666 | OEIS-A239024

Number of n X 2 0..2 arrays with no element equal to the sum of elements to its left or one plus the sum of elements above it, modulo 3.

Conjectures from _Colin Barker_, Oct 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008667 | OEIS-A239031

Number of 4 X n 0..2 arrays with no element equal to the sum of elements to its left or one plus the sum of the elements above it, modulo 3.

Conjectures from _Colin Barker_, Oct 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008668 | OEIS-A239032

Number of 5 X n 0..2 arrays with no element equal to the sum of elements to its left or one plus the sum of the elements above it, modulo 3.

Conjectures from _Colin Barker_, Oct 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008669 | OEIS-A239033

Number of 6 X n 0..2 arrays with no element equal to the sum of elements to its left or one plus the sum of the elements above it, modulo 3.

Conjectures from _Colin Barker_, Oct 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008670 | OEIS-A239066

Triangle read by rows: row n lists the smallest positive ideal multigrade of degree n, or 2n+2 zeros if none.

Ideal multigrades are known only for degrees < 11 and degree 12. The ideal multigrades of degrees 5,6,7,8,9,10 are only conjecturally the smallest ones.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Mar 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008671 | OEIS-A239067

Triangle read by rows: row n lists the smallest positive ideal symmetric multigrade of degree n, or 2n+2 zeros if none.

The ideal symmetric multigrades of degrees 5,6,7,8,9,10 are only conjecturally the smallest ones.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Mar 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008672 | OEIS-A239068

Triangle read by rows: row n lists the smallest positive ideal non-symmetric multigrade of degree n, or 2n+2 zeros if none.

Any ideal multigrade x1,x2;y1,y2 of degree 1 is symmetric, since x1+x2 = y1+y2. Ideal non-symmetric multigrades are known only for degrees 2,3,4,5,6,7. The ones for degrees 5,6,7 are only conjecturally the smallest ones.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Mar 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008673 | OEIS-A239119

Number of ballot sequences of length n with exactly 8 fixed points.

Conjecture: Generally, for fixed k is column k of A238802 asymptotic to sqrt(2)/(2*(k+1)*(k-1)!) * exp(sqrt(n)-n/2-1/4) * n^(n/2) * (1 + 7/(24*sqrt(n))), (holds for all k<=10). - _Vaclav Kotesovec_, Mar 11 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_ and _Alois P. Heinz_, Mar 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008674 | OEIS-A239171

Number of (n+1) X (1+1) 0..2 arrays with no element greater than all horizontal neighbors or equal to all vertical neighbors.

Conjectures from _Colin Barker_, Oct 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008675 | OEIS-A239201

Expansion of -(x * sqrt(5*x^2 -6*x +1) -2*x^3 +3*x^2 -x) / ((3*x^2 -4*x +1) * sqrt(5*x^2 -6*x +1) +5*x^3 -11*x^2 +7*x -1).

Conjecture D-finite with recurrence: (-n+1)*a(n) +(7*n-11)*a(n-1) +(-11*n+25)*a(n-2) +5*(n-3)*a(n-3)=0. - _R. J. Mathar_, Oct 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Mar 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008676 | OEIS-A239204

Expansion of ((x-1)*sqrt(x^2-6*x+1)-x^2-4*x+1)/(8*x^3).

Conjectured to be D-finite with recurrence: (n+3)*a(n) + (-7*n-10)*a(n-1) + (7*n-3)*a(n-2) + (-n+2)*a(n-3) = 0. - _R. J. Mathar_, Nov 02 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Mar 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008677 | OEIS-A239207

a(n) = |{0 < k <= n: k*p(n)*q(n)*r(n) - 1 is prime}|, where p(.), q(.) and r(.) are given by A000041, A000009 and A047967 respectively.

Conjecture: (i) a(n) > 0 for all n > 1. If n > 1 is not equal to 25, then k*p(n)*q(n)*r(n) + 1 is prime for some k = 1, ..., n. See also A239209 and A239214 for related conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008678 | OEIS-A239209

a(n) = |{0 < k < n: k*p(n)*(p(n)-1) + 1 is prime}|, where p(.) is the partition function (A000041).

Conjecture: (i) a(n) > 0 for all n > 1. If n > 1 is not equal to 25, then k*p(n)*(p(n)+1) - 1 is prime for some 0 < k < n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008679 | OEIS-A239214

a(n) = |{0 < k < n: p(k)*p(n)*(p(n)+1) - 1 is prime}|, where p(.) is the partition function (A000041).

Conjecture: (i) a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008680 | OEIS-A239230

Expansion of -x*log'(-sqrt(12*x+2*sqrt(1-4*x)+2)/4+sqrt(1-4*x)/4+5/4).

Conjecture D-finite with recurrence +24*(365025561*n-1672569283)*(n-1)*(n-2)*(2*n-1)*a(n) -4*(n-2)*(27129169947*n^3-209577621466*n^2+463278020461*n-314084557758)*a(n-1) +2*(-28823853823*n^4+487259692534*n^3-3105214937957*n^2+8814274338098*n-9143920331436)*a(n-2) +4*(276083065830*n^4-4172118623320*n^3+24824880820695*n^2-69263721795041*n+75832154222148)*a(n-3) +(-587491214125*n^4+9941738070620*n^3-75070680472775*n^2+281912285021344*n-413197788157152)*a(n-4) +2*(-1186924847911*n^4+26108844767699*n^3-211936472383904*n^2+757584729548632*n-1009721693733312)*a(n-5) +4*(2*n-11)*(328530544924*n^3-5280431217363*n^2+28334632524947*n-50473913356356)*a(n-6) -72*(4143100547*n-18456753180)*(n-6)*(2*n-11)*(2*n-13)*a(n-7)=0. - _R. J. Mathar_, Jul 27 2022 Conjecture: a(n) = Sum_{i=0..n-1} A059260(2*(n-1),n+i-1)*A009766(n+i-1,i)*(-1)^(n+i-1) = n*Sum_{i=0..n-1} binomial(n+2*i, i)/(n+2*i)*(-1)^(n+i-1)*[x^(n+i-1)] (1+(x+1)^(2*n-1))/(x+2) for n >= 0. - _Mikhail Kurkov_, Feb 18 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, May 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008681 | OEIS-A239232

a(n) = |{0 < k <= n: p(n+k) + 1 is prime}|, where p(.) is the partition function (A000041).

Conjecture: (i) a(n) > 0 for all n > 3. The conjecture implies that there are infinitely many positive integers m with p(m) + 1 (or p(m) - 1, or p(m)) prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008682 | OEIS-A239241

Number of partitions of n into distinct parts for which (number of odd parts) = (number of even parts).

Conjecture: c = 1/(2^(5/2)*3^(1/2)). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008683 | OEIS-A239275

a(n) = numerator(2^n * Bernoulli(n, 1)).

Numerators of the binomial transform of A157779(n)/(interleave A001897(n), 1)(conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Mar 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008684 | OEIS-A239293

Smallest composite c > n such that n^c == n (mod c).

Conjecture: a(n) = n+1 if and only if n+1 is an odd composite number. - _Thomas Ordowski_, Aug 03 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert FERREOL_, Mar 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008685 | OEIS-A239328

Number of primes p < n with pi(p*n) - pi((p-1)n) prime, where pi(x) denotes the number of primes not exceeding x.

Conjecture: a(n) > 0 for all n > 3. Also, for any integer n > 4, there is a prime p < n with pi((p+1)*n) - pi(p*n) prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008686 | OEIS-A239330

Number of odd primes p <= n with pi(n*(p+1)/2) - pi(n*(p-1)/2) prime, where pi(x) denotes the number of primes not exceeding x.

Conjecture: a(n) > 0 for all n > 3, and a(n) = 1 only for n = 4, 5, 17.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008687 | OEIS-A239351

Integer solutions x, y, z of x^3 + y^3 + z^3 = 3 with |x| <= |y| <= |z|.

It is conjectured that no other solution exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Apr 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008688 | OEIS-A239352

van Heijst's upper bound on the number of squares inscribed by a real algebraic curve in R^2 of degree n, if the number is finite.

In 1911 Toeplitz conjectured the Square Peg (or Inscribed Square) Problem: Every continuous simple closed curve in the plane contains 4 points that are the vertices of a square. The conjecture is still open. Many special cases have been proved; see Matschke's beautiful 2014 survey. Recently van Heijst proved that any real algebraic curve in R^2 of degree d inscribes either at most (d^4 - 5d^2 + 4d)/4 or infinitely many squares. He conjectured that a generic complex algebraic plane curve inscribes exactly (d^4 - 5d^2 + 4d)/4 squares.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Mar 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008689 | OEIS-A239358

Number of n X 5 0..2 arrays with no element equal to one plus the sum of elements to its left or one plus the sum of elements above it or one plus the sum of the elements diagonally to its northwest, modulo 3.

Conjectures from _Colin Barker_, Oct 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008690 | OEIS-A239359

Number of n X 6 0..2 arrays with no element equal to one plus the sum of elements to its left or one plus the sum of elements above it or one plus the sum of the elements diagonally to its northwest, modulo 3.

Conjectures from _Colin Barker_, Oct 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008691 | OEIS-A239367

The bisection of A238315 that remains constant with changes in the offset of the exponent of the second term.

Conjectures from _Colin Barker_, Dec 30 2014: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard R. Forberg_, Mar 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008692 | OEIS-A239382

Decimal expansion of the probability of a normal-error variable exceeding the mean by more than one standard deviation.

In experimental sciences (hypothesis testing), a measured excursion exceeding background "noise" by just one standard deviation is not significant, unless corroborated by strong additional indications.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Mar 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008693 | OEIS-A239383

Decimal expansion of the probability of a normal-error variable exceeding the mean by more than two standard deviations.

The probability P{(x-m)/s > 2} for a normally distributed random variable x with mean m and standard deviation s. In experimental sciences (hypothesis testing), a measured excursion exceeding background "noise" by more than two standard deviations is considered mildly significant if corroborated by additional indications.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Mar 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008694 | OEIS-A239384

Decimal expansion of the probability of a normal-error variable exceeding the mean by more than three standard deviations.

In experimental sciences (hypothesis testing), a measured excursion exceeding background "noise" by more than three standard deviations is considered fairly significant, unless it is an isolated case among hundreds of measurements.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Mar 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008695 | OEIS-A239385

Decimal expansion of the probability of a normal-error variable exceeding the mean by more than four standard deviations.

In experimental sciences (hypothesis testing), a measured excursion exceeding background "noise" by more than four standard deviations is considered significant, unless it is an isolated case among thousands of iterated measurements.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Mar 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008696 | OEIS-A239386

Decimal expansion of the probability of a normal-error variable exceeding the mean by more than five standard deviations.

In experimental sciences (hypothesis testing), a measured excursion exceeding background "noise" by more than five standard deviations is considered practically certain, unless it is an isolated case among tens of thousands of iterated measurements.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Mar 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008697 | OEIS-A239387

Decimal expansion of the probability of a normal-error variable exceeding the mean by more than six standard deviations.

In experimental sciences (hypothesis testing), a measured excursion exceeding background "noise" by more than six standard deviations is considered certain and "experimentally confirmed".

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Mar 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008698 | OEIS-A239425

Expansion of -16/(sqrt(12*x+2*sqrt(1-4*x)+2)-sqrt(1-4*x)-1)^2+1/x^2-1.

Conjecture: 2*(2*n+1)*(n+2)*(n+1)*a(n) +(n+1)*(n^2-27*n+2)*a(n-1) +2*(-73*n^3+204*n^2-167*n+6)*a(n-2) +12*(n-3)*(2*n-3)*(4*n-7)*a(n-3) +216*(2*n-5)*(n-3)*(2*n-3)*a(n-4)=0. - _R. J. Mathar_, Apr 02 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Mar 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008699 | OEIS-A239428

Number of ordered ways to write n = k + m with 0 < k <= m such that pi(2*k) - pi(k) and pi(2*m) - pi(m) are both prime, where pi(x) denotes the number of primes not exceeding x.

Conjecture: a(n) > 0 for all n > 9, and a(n) = 1 only for n = 8, 10, 11, 26, 33, 50, 52.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008700 | OEIS-A239430

Number of ways to write n = k + m with k > 0 and m > 0 such that pi(2*k) - pi(k) is prime and pi(2*m) - pi(m) is a square, where pi(x) denotes the number of primes not exceeding x.

Conjecture: a(n) > 0 for all n > 4, and a(n) = 1 only for n = 5, 6, 20.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008701 | OEIS-A239451

a(n) = |{1 < k < sqrt(n)*log(n): prime(n) + C(prime(k)-1, (prime(k)-1)/2) is prime}|, where C(2j,j) = (2j)!/(j!)^2.

Conjecture: (i) a(n) > 0 for all n > 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008702 | OEIS-A239455

Number of Look-and-Say partitions of n; see Comments.

Conjecture: a partition is Look-and-Say iff it has a permutation with all distinct run-lengths. For example, the partition y = (2,2,2,1,1,1) has the permutation (2,2,1,1,1,2), with run-lengths (2,3,1), which are all distinct, so y is counted under a(9). - _Gus Wiseman_, Aug 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ and _Peter J. C. Moses_, Mar 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008703 | OEIS-A239530

Number of (n+1) X (1+1) 0..2 arrays with no element equal to all horizontal neighbors or unequal to all vertical neighbors, and new values 0..2 introduced in row major order.

Conjectures from _Colin Barker_, Oct 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008704 | OEIS-A239546

Sum of the prime factors of n, where n runs through perfect numbers.

a(n) = A000668(n) + 2 (conjectured). - _Michel Marcus_, Mar 21 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Madaline McCluskey_, Mar 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008705 | OEIS-A239566

(Round(c^prime(n)) - 1)/prime(n), where c is the heptanacci constant (A118428).

Conjecture. Absolute values of all roots of the equation x^n - x^(n-1) - ... -x - 1 = 0, except for n-bonacci constant c_n, are less than 1. If the conjecture is valid, then for sufficiently large k=k(n), for all m>=k, we have round(c_n^prime(m)) == 1 (mod 2*prime(m)) (cf. Shevelev link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter J. C. Moses_ and _Vladimir Shevelev_, Mar 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008706 | OEIS-A239614

a(n) = A239611(n) / A079458(n).

Conjectures from _Ridouane Oudra_, Jul 22 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Jun 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008707 | OEIS-A239622

Conjecturally, the irregular triangle of numbers k such that prime(n)^2 is the largest squared prime divisor of binomial(2k,k).

Conjecturally, the irregular triangle of numbers k such that prime(n)^2 is the largest squared prime divisor of binomial(2k,k). Row 0 lists the numbers k such that binomial(2k,k) is squarefree. Sequence A110494 lists the first term of each row; A239623 lists the conjectured last term; A239624 lists the conjectured length of each row.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008708 | OEIS-A239623

Conjecturally, the largest k such that prime(n)^2 is the largest squared prime divisor of binomial(2k,k).

Conjecturally, the largest k such that prime(n)^2 is the largest squared prime divisor of binomial(2k,k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008709 | OEIS-A239624

Conjecturally, the number of numbers k such that prime(n)^2 is the largest squared prime divisor of binomial(2k,k).

Conjecturally, the number of numbers k such that prime(n)^2 is the largest squared prime divisor of binomial(2k,k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _T. D. Noe_, Mar 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008710 | OEIS-A239657

Number of odd divisors m of n such that there is a divisor d of n with d < m < 2*d.

Conjecture: number of partitions of n into consecutive parts minus the number of 2-dense sublists of divisors of n. - _Omar E. Pol_, Nov 22 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Mar 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008711 | OEIS-A239669

Total number of prime factors counted with multiplicity of prime(n)-1 and prime(n)+1, where prime(n) is the n-th prime.

If a(n) <= 6, n is a distinguished prime (A106639). According to a conjecture in A106639, 6 is the least term appearing infinitely often. Numerical evidence suggests the conjecture that other - perhaps all - integers > 6 appear infinitely often. The strong conjecture implies that all terms but one in A155800 are positive.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chris Boyd_, Mar 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008712 | OEIS-A239800

First appearance of n in A016014, or 0 if n never occurs.

Conjecture: a(n) > 0 for all n. - _Jianing Song_, Mar 09 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Mar 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008713 | OEIS-A239844

Number of n X 2 0..3 arrays with no element equal to one plus the sum of elements to its left or one plus the sum of elements above it or two plus the sum of the elements diagonally to its northwest, modulo 4.

Conjectures from _Colin Barker_, Oct 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008714 | OEIS-A239884

Least positive integer k <= n with pi(pi(k*n)) a square, or 0 if such a number k does not exist, where pi(x) denotes the number of primes not exceeding x.

According to part (i) of the conjecture in A238902, a(n) should be always positive. We have verified this for all n = 1, ..., 2*10^5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008715 | OEIS-A239903

List of Restricted-Growth Strings a_{k-1}a_{k-2}...a_{2}a_{1}, with k=2 and a_1 in {0,1} or k>2, a_{k-1}=1 and a_{j+1}>=1+a_j, for k-1>j>0.

Please see Dejter's preprint for a more formal mathematical definition and how this number system is applied in relation to Havel's Conjecture on the existence of Hamiltonian cycles in the middle-levels graphs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008716 | OEIS-A239907

Let cn(n,k) denote the number of times 11..1 (k 1's) appears in the binary representation of n; a(n) = n - cn(n,1) + cn(n,2) - cn(n,3) + cn(n,4) - ... .

Conjecture: a(n) = n - A329320(n) for n >= 0 (noticed by Sequence Machine). - _Mikhail Kurkov_, Oct 13 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008717 | OEIS-A239987

Number of 3 X n 0..3 arrays with no element equal to zero plus the sum of elements to its left or two plus the sum of the elements above it or one plus the sum of the elements diagonally to its northwest, modulo 4.

Conjectures from _Colin Barker_, Oct 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008718 | OEIS-A240001

Number of 2 X n 0..3 arrays with no element equal to one plus the sum of elements to its left or two plus the sum of the elements above it or one plus the sum of the elements diagonally to its northwest, modulo 4.

Conjectures from _Colin Barker_, Oct 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008719 | OEIS-A240002

Number of 3 X n 0..3 arrays with no element equal to one plus the sum of elements to its left or two plus the sum of the elements above it or one plus the sum of the elements diagonally to its northwest, modulo 4.

Conjectures from _Colin Barker_, Oct 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008720 | OEIS-A240062

Square array read by antidiagonals in which T(n,k) is the n-th number j with the property that the symmetric representation of sigma(j) has k parts, with j >= 1, n >= 1, k >= 1.

Conjecture: T(n,k) is the n-th positive integer with k 2-dense sublists of divisors. - _Omar E. Pol_, Aug 25 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Apr 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008721 | OEIS-A240087

Smallest difference to start a prime arithmetic progression of three or more terms with the n-th prime number.

It is conjectured that this sequence is defined for all odd prime numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Mar 31 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008722 | OEIS-A240088

The number of ways of writing n as an ordered sum of a triangular number (A000217), a square (A000290) and a pentagonal number (A000326).

It is conjectured that a(n) is always positive - this is one of the conjectures in Conjecture 1.1 of Sun (2009). - _N. J. A. Sloane_, Apr 01 2014 Note that both the conjecture in A160325 and the conjecture in A160324 imply that a(n) is always positive. - _Zhi-Wei Sun_, Apr 01 2014 Greatest number (conjectured) to be represented k ways, k >= 1: 0, 18, 168, 78, 243, 130, 553, 455, 515, 658, 865, 945, 633, 1918, 2258, 1385, 1583, 2828, 2135, 2335, 2785, 4533, 3168, 3478, 2790, 3868, 4193, 7328, 4953, 5278, 6390, 8148, 8015, 4585, 9160, 10485, 7613, 12333, 12025, 10178, 9923, 9720, 12558, 11340, 17420, 11753, 14893, 16155, 16415, 14343, ..., . Conjectured lists of numbers that are represented in k >= 1 ways:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Mar 31 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008723 | OEIS-A240114

Maximal number of points that can be placed on a triangular grid of side n so that no three of them are vertices of an equilateral triangle in any orientation.

Conjecture: for n >= 11, a(n) = 3n-11. (End) The conjecture a(n) = 3*n - 11 (for n >= 11) is false; a(n) grows superlinearly. In triangular-lattice coordinates, a set with no equilateral triangle contains no triple (x,y), (x+r,y), (x,y+r) with r > 0 (these three points are mutually equidistant), so it is corner-free; hence a(n) = o(n^2) by the corners theorem of Ajtai and Szemeredi. Conversely a(n) >> n^(log_7(12)) = n^(1.2769...) by an explicit triangular-lattice construction (Polansky 2019). Hence a(n)/n -> infinity. - _Josh Arnold_, May 30 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Heinrich Ludwig_, Apr 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008724 | OEIS-A240131

Least k such that prime(n)^2 + k^2 is prime, or 0 if none.

k and p_n must be of opposite parity. Conjecture, there is always a k for any p_n. Tested for all primes < 15*10^10. - _Robert G. Wilson v_, Nov 04 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Apr 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008725 | OEIS-A240136

Prime numbers in the Antikythera mechanism.

The Antikythera Mechanism is a unique Greek geared device, constructed around the end of the 2nd Century B.C. This device is a sophisticated arrangement of bronze gears, which can calculate the relative positions of the Sun and Moon, lunar phases and even eclipses. It is conjectured that it could also calculate the positions of the inner planets, but this is unproven. It was recovered from a Roman shipwreck in 1900 as a corroded mass of gear work. Detailed analysis could only really begin once X-ray equipment became available. Scans performed over the last 20 years have added significant clues as to the function and operation of the device. Ratios of gear tooth counts in the device indicate a good match to ratios used by the ancients to calculate the dates of eclipses using the Saros and Metonic cycles. Fragments of text recovered from the device mention the sun and moon and use month names used by the Greek city of Corinth and its colonies.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008726 | OEIS-A240167

Lesser of the first cousin prime pair with n digits.

It is only a (plausible) conjecture that this sequence is well-defined. See A152052. - _N. J. A. Sloane_, Aug 22 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Abhiram R Devesh_, Aug 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008727 | OEIS-A240301

a(n) is the number of sets of three positive integers p_1 < p_2 < p_3 such that 2*p_2 = p_1 + p_3, where p_i (i=1,2,3) is either 1 or a prime number and p_3 = prime(n).

It is conjectured that every positive integer appears a positive finite number of times in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Apr 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008728 | OEIS-A240333

Number of n X 2 0..3 arrays with no element equal to two plus the sum of elements to its left or two plus the sum of elements above it or one plus the sum of the elements diagonally to its northwest, modulo 4.

Conjectures from _Colin Barker_, Oct 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008729 | OEIS-A240359

Number of n X 2 0..3 arrays with no element equal to one plus the sum of elements to its left or two plus the sum of elements above it or two plus the sum of the elements diagonally to its northwest, modulo 4.

Conjectures from _Colin Barker_, Oct 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008730 | OEIS-A240395

Number of 3 X n 0..3 arrays with no element equal to zero plus the sum of elements to its left or one plus the sum of the elements above it or two plus the sum of the elements diagonally to its northwest, modulo 4.

Conjectures from _Colin Barker_, Oct 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008731 | OEIS-A240434

Binomial transform of the sum of the first n even squares (A002492).

Conjecture: a(n) = (2^(n-1)*n*(5+6*n+n^2))/3. G.f.: -4*x*(x-1) / (2*x-1)^4. - _Colin Barker_, Apr 06 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Apr 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008732 | OEIS-A240537

Let a(n) be the least k such that in the prime power factorization of k! the exponents of primes p_1, ...,p_n are even, while the exponent of p_(n+1) is odd.

The sequence is connected with a 1980-Erdős-Graham conjecture that, for every N, there exists an n such that in prime power factorization of n! at least N first exponents are even. In 1997, this conjecture was proved by D. Berend. A generalization was given by Y.-G. Chen (2003).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Apr 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008733 | OEIS-A240542

The midpoint of the (rotated) Dyck path from (0, n) to (n, 0) defined by A237593 has coordinates (a(n), a(n)). Also a(n) is the alternating sum of the n-th row of A235791.

Conjecture 1: the sequence of first positions of the alternating runs of odd and even numbers in a(k) equals A028982. Example: the positions in (1),(2),2,(3),3,5,5,(6),(7),7,7,9,9,9,11,(12),12,(13),13,15,... are 1,2,4,8,9,16,18,... Checked with a Mathematica function through a(1000000). Conjecture 2: The sequence of differences a(n) - a(n-1), for n>=1, appears to be equal to A067742(n), the sequence of middle divisors of n; as an empty sum, a(0) = 0, (which was conjectured by _Michel Marcus_ in the entry A237593). Checked with the two respective Mathematica functions up to n=100000. - _Hartmut F. W. Hoft_, Jul 17 2014 Conjecture 3: a(n) is also the difference between the total number of partitions of all positive integers <= n into an odd number of consecutive parts, and the total number of partitions of all positive integers <= n into an even number of consecutive parts. - _Omar E. Pol_, Apr 28 2017 Conjecture 4: a(n) is also the total number of central subparts of all symmetric representations of sigma of all positive integers <= n. - _Omar E. Pol_, Apr 29 2017 a(n) is the total number of middle divisors of all positive integers <= n (after _Michel Marcus_'s conjecture in A237593). - _Omar E. Pol_, Aug 18 2021 a(n) = A285901(n) - A285902(n), assuming the conjecture 3. - _Omar E. Pol_, Feb 15 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Apr 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008734 | OEIS-A240545

Least k > 0 such that n + p(k) is prime, where p(k) is the number of partitions of k.

Conjecture of Zhi-Wei Sun: a(n) < n for n > 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sean A. Irvine_, Apr 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008735 | OEIS-A240558

a(n) = 2^n*n!/((floor(n/2)+1)*floor(n/2)!^2).

Conjecture: -(n+2)*(n^2-5)*a(n) +8*(-2*n-1)*a(n-1) +16*(n-1)*(n^2+2*n-4)*a(n-2)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Apr 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008736 | OEIS-A240562

Integers whose squares are in A000073 (tribonacci numbers).

No more terms < 10^19300. I conjecture that the sequence is finite. - _Manfred Scheucher_, Aug 17 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Apr 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008737 | OEIS-A240567

a(n) = optimal number of tricks to throw in the game of One Round War (with n cards) in order to maximize the expected number of tricks won.

The formula is conjectured to hold for all n between 60 and 10 million, as well.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dana Mackenzie_, Apr 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008738 | OEIS-A240568

Difference between n times the n-th prime and twice the sum of the first n primes.

Rosser & Schoenfeld prove Robert Mandl's conjecture that a(n) > 0 for n > 8. - _Charles R Greathouse IV_, Jun 11 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Apr 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008739 | OEIS-A240583

Odd primes satisfying a specific condition (see comments).

Consider the even integers E > a(1), and constrain distance <= E. The absolute distance from E to one of the sequence terms is conjectured to nearly always be prime (or adjacent). For example, 1488768 = a(40046)-1488623. There are four early exceptions thru E = 1488694, {104, 128, 134, 212}. Where is the next exception? - _Bill McEachen_, May 31 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Apr 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008740 | OEIS-A240604

Primes p with pi(p), pi(2*p), pi(3*p) and pi(4*p) all prime, where pi(x) denotes the number of primes not exceeding x.

Conjecture: For any positive integer n, there are infinitely many primes p with pi(k*p) (k = 1,...,n) all prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008741 | OEIS-A240669

Number of the first odious exponents (A000069) in the prime power factorization of (2n)!.

Conjecture: The sequence is unbounded. (This conjecture does not follow from Chen's theorem.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008742 | OEIS-A240672

Number of the first evil exponents (A001969) in the prime power factorization of (2n)!.

Conjecture: The sequence is unbounded. (This conjecture does not follow from Chen's theorem.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008743 | OEIS-A240688

Expansion of -(x*sqrt(-4*x^2-4*x+1)-2*x^2-3*x) / ((x+1)*sqrt(-4*x^2-4*x+1)+ 4*x^3+8*x^2+3*x-1).

Conjecture: n*(n-1)*a(n) -(3*n-2)*(n-1)*a(n-1) +2*(-4*n^2+7*n-1)*a(n-2) -4*n*(n-2)*a(n-3)=0. - _R. J. Mathar_, Jun 14 2016 Mathar's conjectural third-order recurrence above can be verified using Maple's gfun:-rectodiffeq command. Conjecture: the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(2*k)) hold for primes p >= 5 and positive integers n and k. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Apr 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008744 | OEIS-A240714

Even numbers whose unordered two primes decomposition set does not contain two groups of n = p1+p2 = p3+p4 such that |p1-p3| = 6 or 12.

Conjecture: this sequence is finite and all elements are listed.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Apr 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008745 | OEIS-A240722

Expansion of log'((-1-sqrt(1-8*x)+sqrt(2+2*sqrt(1-8*x)+8*x))/(4*x)).

Conjecture: -(3*n-1)*(2*n+1)*(n+1)*a(n) +2*(21*n^3+14*n^2-2*n-1)*a(n-1) +8*n*(3*n+2)*(2*n-1)*a(n-2)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Apr 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008746 | OEIS-A240764

Least k such that prime power factorization of A240751(k)! contains p^k when the smallest such p equals prime(n), or a(n)=0 if there is no such k.

Conjecture: all a(n)>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008747 | OEIS-A240775

The six values n in each interval [i*840, (i+1)*840), where i >= 0, for which Mordell's formulas do NOT provide a three-Egyptian-fraction solution for 4/n.

Erdős and Straus conjectured that for all integers n >= 2, the rational number 4/n can be expressed as an Egyptian fraction with exactly three unit fractions -- that is, 4/n = 1/x + 1/y + 1/z where x, y and z are positive integers. The conjecture has been verified to high values of n, and Mordell has provided formulas to compute x, y and z for many n. The values of n NOT included in Mordell's formulas are those for which n modulo 840 = {an element of this sequence}. Each element is the square of a prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kenneth Vollmar_, Apr 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008748 | OEIS-A240809

a(n) = n for 1 <= n <= 4; thereafter a(n) = a(n - a(n-2)) + a(n - a(n-4)).

Conjectured to be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 15 2014, Apr 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008749 | OEIS-A240810

a(n) = index of first nonexisting term of the meta-Fibonacci sequence {f(1) = ... = f(n) = 1; f(k)=f(k-f(k-1))+f(k-f(k-n))} if that sequence is only defined for finitely many terms, or 0 if that sequence is infinite.

a(2)=0 is only a conjecture (see A005185), whereas a(4)=0 is a theorem of Balamohan et al. (2007).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008750 | OEIS-A240811

a(n) = length (or lifetime) of the meta-Fibonacci sequence f(1) = ... = f(n) = 1; f(k)=f(k-f(k-2))+f(k-f(k-n)) if that sequence is only defined for finitely many terms, or 0 if that sequence is infinite.

The term a(4) = 0 is only conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008751 | OEIS-A240813

Triangle read by rows: T(n,k) (n >= 1, 1 <= k <= n) = length (or lifetime) of the meta-Fibonacci sequence f(1) = ... = f(n) = 1; f(i)=f(i-f(i-k))+f(i-f(i-n)) if that sequence is only defined for finitely many terms, or 0 if that sequence is infinite.

The zero entries (except T(4,1)) are only conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008752 | OEIS-A240814

a(n) = index of first nonexisting term of the meta-Fibonacci sequence {f(1) = ... = f(n) = 1; f(k)=f(k-f(k-2))+f(k-f(k-n))} if that sequence is only defined for finitely many terms, or 0 if that sequence is infinite.

a(4)=0 is only a conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008753 | OEIS-A240816

Triangle read by rows: T(n,k) (n >= 1, 1 <= k <= n) = index of first nonexisting term of the meta-Fibonacci sequence {f(1) = ... = f(n) = 1; f(i)=f(i-f(i-k))+f(i-f(i-n))} if that sequence is only defined for finitely many terms, or 0 if that sequence is infinite.

The zero entries (except T(4,1)) are only conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008754 | OEIS-A240818

a(n) = length (or lifetime) of the meta-Fibonacci sequence f(k) = k for k <= n; f(k)=f(k-f(k-1))+f(k-f(k-n)) if that sequence is only defined for finitely many terms, or 0 if that sequence is infinite.

The terms a(2) = 0 and a(4) = 0 are only conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008755 | OEIS-A240819

a(n) = length (or lifetime) of the meta-Fibonacci sequence f(k) = k for k <= n; f(k)=f(k-f(k-2))+f(k-f(k-n)) if that sequence is only defined for finitely many terms, or 0 if that sequence is infinite.

The term a(4) = 0 is only conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008756 | OEIS-A240820

a(n) = length (or lifetime) of the meta-Fibonacci sequence f(k) = k for k <= n; f(k) = f(k-f(k-3)) + f(k-f(k-n)) if that sequence is only defined for finitely many terms, or 0 if that sequence is infinite.

The term a(6) = 0 is only conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008757 | OEIS-A240821

Triangle read by rows: T(n,k) (n >= 1, 1 <= k <= n) = length (or lifetime) of the meta-Fibonacci sequence {f(i) = i for i <= n; f(i)=f(i-f(i-k))+f(i-f(i-n))} if that sequence is only defined for finitely many terms, or 0 if that sequence is infinite.

The zero entries are only conjectural. More precisely, Hofstadter conjectures that T(n,k) = 0 (i.e. the sequence is immortal) iff n = 2k or n = 4k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008758 | OEIS-A240822

a(n) = index of first nonexisting term of the meta-Fibonacci sequence {f(k) = k for k <= n; f(k)=f(k-f(k-1))+f(k-f(k-n))} if that sequence is only defined for finitely many terms, or 0 if that sequence is infinite.

a(2)=0 and a(4)=0 are conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008759 | OEIS-A240823

a(n) = index of first nonexisting term of the meta-Fibonacci sequence {f(k) = k for k <= n; f(k)=f(k-f(k-2))+f(k-f(k-n))} if that sequence is only defined for finitely many terms, or 0 if that sequence is infinite.

a(4)=0 and a(8)=0 are only conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008760 | OEIS-A240824

a(n) = index of first nonexisting term of the meta-Fibonacci sequence {f(k) = k for k <= n; f(k)=f(k-f(k-3))+f(k-f(k-n))} if that sequence is only defined for finitely many terms, or 0 if that sequence is infinite.

a(6)=0 and a(12)=0 are only conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008761 | OEIS-A240827

a(n) = n for 1<=n<=6; thereafter a(n) = a(n-a(n-3))+a(n-a(n-6)).

Conjectured to be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008762 | OEIS-A240886

Primes p equal to the sum in base-3 of the digits of all primes < p.

Conjecture: this sequence is finite and all terms are shown. - _Robert G. Wilson v_, Jul 27 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anthony Sand_, Apr 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008763 | OEIS-A240905

Smallest k such that the minimal factor in factorization of k! over distinct terms of A050376 is A050376(n), or a(n) = 0 if there is no such k.

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008764 | OEIS-A240911

Smallest number greater than a(n-1) divisible by smallest number that is not a divisor of a(n-1).

Conjecture: Numbers that are not in this sequence below any finite limit will never outnumber numbers that are. The expected value of the first differences of this sequence is Sum_{q>=2} q*(p-1)/(2*lcm(2,...,q)) as q = p^k ranges over the prime powers, or approximately 1.39389022808623327303..., so the conjecture is very likely to be true. - _Charlie Neder_, Apr 26 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Apr 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008765 | OEIS-A240912

Numbers not in A240911.

Conjecture: for all N, the number of terms less than N in A240911 exceeds the number of terms less than N in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Apr 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008766 | OEIS-A240920

Prime numbers that occur as divisors of numbers of the form m^2 + 5.

Conjecture: a prime number is in this sequence if and only if its next-to-last digit is even. The law of quadratic reciprocity shows an odd prime is in the sequence if and only if it is 1, 3, 5, 7 or 9 (mod 20). This proves the above conjecture, so the sequence is the union of {2, 5} and A139513. - _Jens Kruse Andersen_, Aug 09 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Aug 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008767 | OEIS-A240945

Powers of 9 without the digit '0' in their decimal expansion.

Conjectured to be finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Aug 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008768 | OEIS-A240998

G.f. satisfies: A(x)^2 = x + A(x + 2*x^2).

For n>0, a(n) == 1 (mod 2) iff n=2^k for k>=0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008769 | OEIS-A241017

Decimal expansion of Sierpiński's S constant, which appears in a series involving the function r(n), defined as the number of representations of the positive integer n as a sum of two squares. This S constant is the usual Sierpiński K constant divided by Pi.

S = A086058 - 1, where A086058 is the conjectured (but erroneous!) value of Masser-Gramain 'delta' constant. [updated by _Vaclav Kotesovec_, Apr 27 2015]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Aug 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008770 | OEIS-A241023

Central terms of the triangle in A102413.

Conjecture: the supercongruences a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) hold for all primes p >= 5 and positive integers n and r. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008771 | OEIS-A241100

Smallest prime with length n having at least n-1 identical digits.

Conjecture: each term consists of at least n-1 digits 1. - _Chai Wah Wu_, Dec 10 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008772 | OEIS-A241151

Number of distinct degrees in the partition graph G(n) defined at A241150.

Conjecture: partial sums of A097806. - _Sean A. Irvine_, Jul 14 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ and _Peter J. C. Moses_, Apr 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008773 | OEIS-A241193

a(n) = Sum_{k=1..n} ((3*n-k-1)/(2*n-k))*(3*n-k-2)!/((n-1)!*(n-1)!*(n-k)!).

Conjecture: -(2*n-1)*(n-1)^2*a(n) +2*(32*n^3-131*n^2+187*n-94)*a(n-1) +3*(-86*n^3+721*n^2-1896*n+1617)*a(n-2) -18*(2*n-5)*(3*n-8)*(3*n-7)*a(n-3)=0. - _R. J. Mathar_, Aug 26 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008774 | OEIS-A241206

Greatest n-digit prime having at least n-1 identical digits.

Conjecture: each term consists of at least n-1 digits 9 and no digit 0. - _Chai Wah Wu_, Dec 10 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008775 | OEIS-A241299

Initial digit of the decimal expansion of n^(n^n) or n^^3 (in _Don Knuth_'s up-arrow notation).

Conjecture: the distribution of the initial digits obey Zipf's law.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Munafo_ and _Robert G. Wilson v_, Apr 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008776 | OEIS-A241427

Smallest prime of the form n^k - k^n for some k, or 0 if no such prime exists.

Conjecture: a(n) > 0 for all n not in A097764.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Aug 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008777 | OEIS-A241476

Least positive primitive root g < prime(n) modulo prime(n) with g - 1 a square, or 0 if such a number g does not exist.

According to the conjecture in A239957, a(n) should be always positive.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008778 | OEIS-A241492

a(n) = |{0 < g < prime(n): g is a primitive root modulo prime(n) and g is a product of two consecutive integers}|.

Conjecture: a(n) > 0 for all n > 4. In other words, any prime p > 7 has a primitive root g < p of the form k*(k+1). See also A239957 and A239963 for similar conjectures. Clearly, for any prime p > 3, one of the three numbers 1*2, 2*3, 3*4 is a quadratic residue modulo p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008779 | OEIS-A241516

Least positive primitive root g < prime(n) modulo prime(n) which is also a partition number given by A000041, or 0 if such a number g does not exist.

According to the conjecture in A241504, a(n) should be always positive.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008780 | OEIS-A241601

Largest divisor of A246006(n) whose prime factors are all >= n+2.

Conjectures: All terms are squarefree, and there are infinitely many n such that a(n) is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, Dec 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008781 | OEIS-A241604

Least Fibonacci number smaller than prime(n)/2 which is a quadratic nonresidue modulo prime(n), or 0 if such a Fibonacci number does not exist.

According to the conjecture in A241568, a(n) should be positive for all n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008782 | OEIS-A241605

a(n) = |{0 < k < sqrt(prime(n))*log(prime(n)) : k is not only a quadratic nonresidue modulo prime(n) but also a Fibonacci number}|.

Conjecture: For any odd prime p, let f(p) be the least Fibonacci number which is a quadratic nonresidue modulo p. Then f(p) = o(p^(0.7)) as p tends to infinity. Moreover,f(p) = O(p^c) for any constant c > c_0 = log_2((1+sqrt(5))/2). This refinement of the conjecture in A241568 seems reasonable in view of the heuristic arguments described below. Let c be any constant greater than c_0 (which has the approximating value 0.694). Roughly speaking, there are about (log_2(p^c))/c_0 positive Fibonacci numbers below p^c. Suppose that a positive Fibonacci number is a quadratic residue modulo a fixed odd prime p with "probability" 1/2. Then we might expect that the "probability" for all positive Fibonacci numbers below p^c being quadratic residues modulo p is about (1/2)^(c*(log_2 p)/c_0) = 1/p^(c/c_0). As sum_p 1/p^(c/c_0) converges, it seems reasonabale to think that there are only finitely many odd primes p for which all positive Fibonacci numbers below p^c are quadratic residues mod p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008783 | OEIS-A241620

Number of length 5+2 0..n arrays with no consecutive three elements summing to more than n.

Conjectures from _Colin Barker_, Oct 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008784 | OEIS-A241621

Number of length 6+2 0..n arrays with no consecutive three elements summing to more than n.

Conjectures from _Colin Barker_, Oct 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008785 | OEIS-A241622

Number of length 7+2 0..n arrays with no consecutive three elements summing to more than n.

Conjectures from _Colin Barker_, Oct 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Apr 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008786 | OEIS-A241656

Smallest semiprime, sp, such that 2n - sp is a semiprime, or a(n)=0 if there is no such sp.

Conjecture: every even number greater than 22 is a sum of two semiprimes. Only 2, 4, 6 & 22 cannot be so represented.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Robert G. Wilson v_, Apr 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008787 | OEIS-A241658

Smallest semiprime, sp, such that n - sp is a semiprime, or a(n)=0 if there is no such sp.

Conjecture: every number greater than 33 is a sum of two semiprimes. Only 1, 2, 3, 4, 5, 6, 7, 9, 11, 17, 22 & 33 cannot be so represented.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Robert G. Wilson v_, Apr 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008788 | OEIS-A241664

Number of iterations of A058026 needed to reach either 0 or 1.

I conjecture that, for n>3 and n odd, we have a(n)>=log((49/15)n)/log(7).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Defant_, Apr 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008789 | OEIS-A241671

Where zeros occur in A065806.

A065806(a(n)-1) = a(n), conjectured in A065806.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Aug 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008790 | OEIS-A241675

a(n) = |{0 < k < n/2: k is a Fibonacci number with x^2 == k (mod n) for no integer x}|.

a(n) > 0 for all n > 4 if and only if the conjecture in A241568 holds. In fact, if n > 0 is a multiple of 4, then x^2 == F(3) = 2 (mod n) for no integer x. If n > 4 is composite with an odd prime divisor p, then by the conjecture in A241568 there should exist a Fibonacci number k < p <= n/2 such that x^2 == k (mod p) for no integer x and hence x^2 == k (mod n) for no integer x.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008791 | OEIS-A241756

A finite sum of products of binomial coefficients: Sum_(m=0..n) binomial(-1/4, m)^2*binomial(-1/4, n-m)^2 (C. C. Grosjean's problem, denominators).

Conjecture (from sequencedb.net): a(n) = 8^A005187(n). - _R. J. Mathar_, Jun 30 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Apr 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008792 | OEIS-A241807

Numerators of c(n) = (n^2+n+2)/((n+1)*(n+2)*(n+3)) as defined in A241269.

a(n) = A014206(n)/period 16: repeat 2, 4, 4, 2, 2, 16, 4, 2, 2, 4, 4, 2, 2, 8, 4, 2 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_ and _Paul Curtz_, Apr 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008793 | OEIS-A241844

Number of ordered ways to write n = k + m with k > 0 and m > 0 such that L(k) + m is prime, where L(k) is the k-th Lucas number.

Conjecture: (i) a(n) > 0 for all n > 1. Moreover, any integer n > 5 can be written as k + m with k > 1 and m > 0 such that L(k) + m is prime. This is similar to the conjecture in A231201. We have verified parts (i) and (ii) for n up to 7*10^6 and 3.7*10^6 respectively.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008794 | OEIS-A241855

Array t(n,k) of sum of successive even powers of primes, where t(n,k) = sum_(j=0..k-1) prime(n)^(2j), with n>=1 and k>=0, read by ascending antidiagonals.

Conjecture: any term, except 0 and 1, is never a square.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Apr 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008795 | OEIS-A241870

Number of compositions of n such that the smallest part has multiplicity ten.

Conjecture: Generally, for k > 0 is column k of A238342 asymptotic to n^k * ((1+sqrt(5))/2)^(n-2*k-1) / (5^((k+1)/2) * k!). - _Vaclav Kotesovec_, May 02 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_ and _Alois P. Heinz_, Apr 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008796 | OEIS-A241887

Smallest m such that A065806(m) = n.

All integers > 1 occur in A065806, see conjecture there;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Aug 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008797 | OEIS-A241937

Number of length 1+4 0..n arrays with no consecutive five elements summing to more than 2*n.

Conjectures from _Colin Barker_, Oct 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008798 | OEIS-A241938

Number of length 2+4 0..n arrays with no consecutive five elements summing to more than 2*n.

Conjectures from _Colin Barker_, Oct 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008799 | OEIS-A241939

Number of length 3+4 0..n arrays with no consecutive five elements summing to more than 2*n.

Conjectures from _Colin Barker_, Oct 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008800 | OEIS-A241940

Number of length 4+4 0..n arrays with no consecutive five elements summing to more than 2*n.

Conjectures from _Colin Barker_, Oct 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008801 | OEIS-A241965

Number of length 2+3 0..n arrays with no consecutive four elements summing to more than 2*n.

Conjectures from _Colin Barker_, Oct 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008802 | OEIS-A241966

Number of length 3+3 0..n arrays with no consecutive four elements summing to more than 2*n.

Conjectures from _Colin Barker_, Oct 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008803 | OEIS-A241967

Number of length 4+3 0..n arrays with no consecutive four elements summing to more than 2*n.

Conjectures from _Colin Barker_, Oct 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008804 | OEIS-A241968

Number of length 5+3 0..n arrays with no consecutive four elements summing to more than 2*n.

Conjectures from _Colin Barker_, Oct 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008805 | OEIS-A242030

Irregular triangle read by rows of a variation of the Collatz iteration with signature (1,1).

It is conjectured that the trajectory of this Collatz-like iteration arrives at 8 in a finite number of steps for any initial value x, (x>2). The iterative step is divide by 2 and add 1 if even, or multiply by 3 and subtract 1 if odd. Two variations belong to the same subset if their (a) parameters are the same and their (b) parameters have the same parity. It is

conjectured that any variations belonging to the same subset have equal row lengths.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _K. Spage_, Aug 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008806 | OEIS-A242033

a(n) = lpf(A245024(n)-1), where lpf = least prime factor (A020639).

Conjecture. The sequence contains all odd primes. The conjecture is true. Consider n-1 = p*q where p is an odd prime and q is a prime > p such that q == p^(-1) mod r for every odd prime r < p. Such primes q exist by Dirichlet's theorem on primes in arithmetic progressions. - _Robert Israel_, Aug 13 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Aug 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008807 | OEIS-A242076

Numbers k for which (2^k + 1)/F is prime where F is a Fermat number.

Conjecture: 6 is the only term whose prime factorization contains a single 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, May 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008808 | OEIS-A242085

Smallest k such that (2*k*3^n-1)*2*k*3^n-1 is prime, with k not divisible by 3.

Conjectures: the ratio a(n)/n is always <10 and sum(a(n)/n)/N for n=1 to N tends to 1 as N tends to infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, May 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008809 | OEIS-A242103

Numbers m such that m^2 - 1 is the product of three distinct Fibonacci numbers > 1.

Conjecture : except the numbers 9, 14, 121, 329, 659, 4523, 7307 and 212533, a(n) is a Lucas number (A000204).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Aug 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008810 | OEIS-A242105

Number of sequences (x(k))_{k=1..n}, of n strictly increasing terms of nonnegative integers {x(1)<x(2)<...<x(n)}, satisfying x(k) <= k^2 for all k.

Conjecture: lim n->infinity a(n)^(1/n)/n = 2. - _Vaclav Kotesovec_, Feb 26 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles Helou_, Aug 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008811 | OEIS-A242118

Number of unit squares that intersect the circumference of a circle of radius n centered at (0,0).

a(n) = 8*n - A046109(n) for n > 0. - conjectured by _Orson R. L. Peters_, Jan 30 2017, proved by _Andrey Zabolotskiy_, Jan 31 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, May 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008812 | OEIS-A242137

Numbers n such that there are no triangular numbers strictly between n^2 and n^2 + n.

For n>2, a(n+1) - a(n) = either 2 or 3 or 5 (conjecture; checked up to a(n) = 2^32).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Aug 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008813 | OEIS-A242145

Number of length 1+5 0..n arrays with no consecutive six elements summing to more than 3*n.

Conjectures from _Colin Barker_, Oct 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008814 | OEIS-A242146

Number of length 2+5 0..n arrays with no consecutive six elements summing to more than 3*n.

Conjectures from _Colin Barker_, Oct 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008815 | OEIS-A242147

Number of length 3+5 0..n arrays with no consecutive six elements summing to more than 3*n.

Conjectures from _Colin Barker_, Oct 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008816 | OEIS-A242169

Least prime divisor of Fr(n) which does not divide any Fr(k) with k < n, or 1 if such a primitive prime divisor of Fr(n) does not exist, where Fr(n) denotes the n-th Franel number given by A000172.

Conjecture: a(n) > 1 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008817 | OEIS-A242170

Least prime divisor of T(n) which does not divide any T(k) with k < n, or 1 if such a primitive prime divisor of T(n) does not exist, where T(n) is the n-th central trinomial coefficient given by A002426.

Conjecture: (i) a(n) > 1 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008818 | OEIS-A242171

Least prime divisor of B(n) which does not divide any B(k) with k < n, or 1 if such a primitive prime divisor of B(n) does not exist, where B(n) is the n-th Bell number given by A000110.

Conjecture: (i) a(n) > 1 for all n > 1. Conjecture: (ii) For any integer n > 2, the derangement number D(n) given by A000166 has a prime divisor dividing none of those D(k) with 1 < k < n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008819 | OEIS-A242172

a(n) = 2^n*binomial((n + 1 + (n mod 2))/2, 1/2).

Conjecture: (n+1)*a(n) -2*a(n-1) +4*(-n-1)*a(n-2)=0. - _R. J. Mathar_, May 11 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, May 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008820 | OEIS-A242173

Least prime divisor of the n-th central Delannoy number D(n) which does not divide any D(k) with k < n, or 1 if such a primitive prime divisor of D(n) does not exist.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008821 | OEIS-A242174

Least prime divisor of A005260(n) which does not divide any previous term A005260(k) with k < n, or 1 if such a primitive prime divisor of A005260(n) does not exist.

Conjecture: a(n) is prime for any n > 0. In general, for any r > 2, if n is large enough then f_r(n) = sum_{k=0..n}C(n,k)^r has a prime divisor which does not divide any previous terms f_r(k) with k < n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008822 | OEIS-A242180

Least prime divisor of q(n) which does not divide any q(k) with k < n, or 1 if such a primitive prime divisor does not exist, where q(.) is the strict partition function given by A000009.

Conjecture: a(n) > 1 for all n > 203.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008823 | OEIS-A242193

Least prime p such that B_{2*n} == 0 (mod p) but there is no k < n with B_{2k} == 0 (mod p), or 1 if such a prime p does not exist, where B_m denotes the m-th Bernoulli number.

Conjecture: a(n) is prime for any n > 4. See also A242194 for a similar conjecture involving Euler numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008824 | OEIS-A242194

Least prime divisor of E_{2*n} which does not divide any E_{2*k} with k < n, or 1 if such a primitive prime divisor of E_{2*n} does not exist, where E_m denotes the m-th Euler number given by A122045.

Conjecture: a(n) is prime for any n > 1. See also A242193 for a similar conjecture involving Bernoulli numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008825 | OEIS-A242195

Least prime divisor of the n-th tangent number T_n which does not divide any T_k with k < n, or 1 if such a primitive prime divisor of T_n does not exist.

Conjecture: (i) a(n) is prime for any n > 3. See also A242193 and A242194 for similar conjectures involving Bernoulli numbers and Euler numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008826 | OEIS-A242207

Least prime divisor of the n-th Domb number D(n) which does not divide any D(k) with k < n, or 1 if such a primitive prime divisor of D(n) does not exist.

Conjecture: a(n) is prime except for n = 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008827 | OEIS-A242222

Number of primes p <= (prime(n)+1)/2 such that the harmonic number H(p-1) = sum_{0<k<p} 1/k is a primitive root modulo prime(n), or 0 if such a prime p does not exist.

Conjecture: a(n) > 0 for all n > 3. In other words, for any prime p > 5, there exists a prime q <= (p+1)/2 such that the harmonic number H(q-1) = sum_{0<k<q}1/k is a primitive root modulo p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008828 | OEIS-A242223

Least prime p such that H(n) == 0 (mod p) but H(k) == 0 (mod p) for no 0 < k < n, or 1 if such a prime p does not exist, where H(n) denotes the n-th harmonic number sum_{k=1..n}1/k.

Conjecture: a(n) is prime except for n = 1, 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008829 | OEIS-A242234

Number of length n+3+1 0..3 arrays with every value 0..3 appearing at least once in every consecutive 3+2 elements, and new values 0..3 introduced in order.

Conjecture: a(n) = 9*A145112(n-1) + A016777(n-1). - _R. J. Mathar_, Aug 16 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008830 | OEIS-A242235

Number of length n+4+1 0..4 arrays with every value 0..4 appearing at least once in every consecutive 4+2 elements, and new values 0..4 introduced in order.

Conjecture: a(n) = 16*A145113(n-1) + A004767(n-2), n > 1. - _R. J. Mathar_, Aug 16 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, May 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008831 | OEIS-A242241

Least prime p such that H(2,n) = sum_{k=1..n}1/k^2 == 0 (mod p) but there is no 0 < k < n with H(2,k) == 0 (mod p), or 1 if such a prime p does not exist.

Conjecture: (i) a(n) is prime for any n > 1. See also A242222 and A242223 for similar conjectures involving harmonic numbers H(n) = sum_{k=1..n}1/k (n > 0).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008832 | OEIS-A242249

Number A(n,k) of rooted trees with n nodes and k-colored non-root nodes; square array A(n,k), n>=0, k>=0, read by antidiagonals.

Column k > 0 is asymptotic to c(k) * d(k)^n / n^(3/2), where constants c(k) and d(k) are dependent only on k. Conjecture: d(k) ~ k * exp(1). Numerically:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, May 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008833 | OEIS-A242250

Least positive integer g < prime(n) such that g, 2^g - 1 and (g-1)! are all primitive roots modulo prime(n), or 0 if such a number g does not exist.

According to the conjecture in A242248, a(n) should be positive for all n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008834 | OEIS-A242258

Number of summands in s-greedy sum of s(n), where s(n) = A000041(n), the partitions numbers.

Conjecture: A000041(n) is A000041-greedy summable for n >= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008835 | OEIS-A242259

Number of summands in s-greedy sum of s(n), where s(n) = A000009(n) (strict partition numbers).

See A242252 for the definitions of greedy sum and summability. Conjecture: A000009(n) is A000009-greedy summable for n >= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008836 | OEIS-A242277

Least positive primitive root g < prime(n) mod prime(n) such that g is the sum of the first k primes for some k > 0, or 0 if such a number g does not exist.

According to the conjecture in A242266, a(n) should be positive for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008837 | OEIS-A242292

Least prime divisor of 2^n - n which does not divide any 2^k - k with 0 < k < n, or 1 if such a primitive prime divisor of 2^n - n does not exist.

Conjecture: a(n) = 1 only for n = 1, 5, 16.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008838 | OEIS-A242364

Irregular triangular array of all the integers ordered as in Comments.

Let f1(x) = 2x, f2(x) = 1-x, f3(x) = 2-x, g(1) = (1), and g(n) = union(f1(g(n-1)), f2(g(n-1)),f3(g(n-1))) for n >1. Let T be the array whose n-th row consists of the numbers in g(n) arranged in increasing order. It is easy to prove that every integer occurs exactly once in T. Conjectures: (1) |g(n)| = 2*F(n-1) for n >=2, where F = A000045 (the Fibonacci numbers), and exactly half of the numbers in g(n) are positive; (2) the number of even numbers in g(n) is 2*F(n-2) and the number of odd numbers is 2*F(n-3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008839 | OEIS-A242365

Irregular triangular array of the positive integers ordered as in Comments.

Conjectures: (1) |g(n)| = F(n-1) for n >=2, where F = A000045 (the Fibonacci numbers); (2) the number of even numbers in g(n) is F(n-2) and the number of odd numbers is F(n-3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008840 | OEIS-A242396

Number of rows of equilateral triangles (sides length = 1) that intersect the circumference of a circle of radius n centered at (0,0) or (1/2,0).

G.f., conjectured: (-2*x^14 + 4*x^13 + 2*x^12 + 2*x^11 + 2*x^10 + 2*x^9 + 2*x^8 + 4*x^7 + 2*x^6 + 2*x^5 + 2*x^4 + 2*x^3 + 2*x^2 + 4*x)/(x^14 - x^13 - x + 1). - _Ralf Stephan_, May 18 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, May 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008841 | OEIS-A242429

Length of longest chain of nonempty proper subsemigroups of the monoid of partial injective order-preserving functions of a chain with n elements.

Conjecture: n*(131*n-376)*a(n) +2*(-563*n^2+1993*n-1185)*a(n-1) +3*(1099*n^2-4678*n+4684)*a(n-2) +2*(-1987*n^2+9803*n-12021)*a(n-3) +4*(209*n-387)*(2*n-7)*a(n-4)=0. - _R. J. Mathar_, Oct 20 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James Mitchell_, May 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008842 | OEIS-A242433

Decimal expansion of one of the Pell-Stevenhagen constants.

P. Stevenhagen conjectured that the asymptotic counting function of the squarefree integers for which the negative Pell equation x^2 - n*y^2 = -1 has an integer solution, was f(n) ~ (6/Pi^2)*P*K*n/sqrt(log(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, May 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008843 | OEIS-A242442

Number of ways of writing n, a positive integer, as an unordered sum of a triangular number (A000217), an odd square (A016754) and a pentagonal number (A000326).

It is conjectured that only 18 cannot be so represented. See Sun, p. 4, Remark 1.2 (b).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, May 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008844 | OEIS-A242444

Number of squares k^2 < n with k^2*p == 1 (mod n) for some prime p < n.

According to the conjecture in A242441, a(n) > 0 for all n > 1848.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008845 | OEIS-A242448

Number of distinct linear polynomials b+c*x in row n of array generated as in Comments.

Conjecture: a(n) = 2*a(n-1) - a(n-3) for n>= 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008846 | OEIS-A242453

(Conjectured) infinite nondecreasing sequence with a(1) = 1 such that the divisors of n appear a total of a(n) times in the sequence.

(Conjectured) infinite nondecreasing sequence with a(1) = 1 such that the divisors of n appear a total of a(n) times in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthew Vandermast_, Jul 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008847 | OEIS-A242462

a(1) = 10; for n>1, a(n) = (a(n-1)-1) * (smallest odd prime factor of a(n-1)) + 1.

Conjecture: this sequence is infinite (so it contains no powers of 2). The similarly defined sequence starting with a(1) = 6 terminates after 2 terms: 6 and 16. Conjecture is true if it turns out that one number in this sequence is one more than a multiple of 41. Any number of the form 2^n-1 that is divisible by 41 is also divisible by 25, which is the square of a prime number greater than 3. Because numbers one less than this sequence's terms are always 9 times a squarefree number, this proves that if a number one less than this sequence is divisible by 41, then this sequence is infinite. - _J. Lowell_, Jul 17 2017 Conjecture is also true if it turns out that one number in this sequence is a multiple of either 19 or 73. Any number of the form 2^n-1 that is divisible by 19 is divisible by 27, and any number of the form 2^n-1 that is divisible by both 73 and 9 is divisible by 27. Numbers one less than this sequence are always divisible by 9 but not by 27. - _J. Lowell_, Feb 27 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, May 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008848 | OEIS-A242463

a(n) = A152466(n) + 1.

Conjecture: this sequence has no prime terms. This conjecture has now been disproved: a(39) is a prime number. - _Andrea Concaro_, Dec 31 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, May 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008849 | OEIS-A242465

Minimum number of mathematical symbols and operators in common use required to write n using exactly three 4's. This version allows only addition, subtraction, multiplication, division, square root, exponentiation, factorial, subfactorial, decimal point, overline (an infinitely repeated digit), parentheses, as well as concatenation.

My web page gives examples of Four Nines up to 774,840,978 = (9-to-the-9) + (9-to-the-9); tablified a Four Pi's; and cites Knuth's Conjecture:"Representing Numbers Using Only One 4", Donald Knuth, [Mathematics Magazine, Vol. 37, Nov/Dec 1964, pp. 308-310]. Knuth shows how (using a computer program he wrote) all integers from 1 through 207 may be represented with only one 4, varying numbers of square roots, varying numbers of factorials, and the floor function. For example: Knuth shows how to make the number 64 using only one 4: |_ sqrt sqrt sqrt sqrt sqrt sqrt sqrt sqrt sqrt |_ sqrt sqrt sqrt sqrt sqrt sqrt sqrt sqrt sqrt |_ sqrt sqrt sqrt sqrt sqrt sqrt sqrt sqrt sqrt sqrt sqrt sqrt sqrt |_ sqrt sqrt sqrt sqrt sqrt sqrt sqrt sqrt sqrt |_ sqrt sqrt sqrt sqrt sqrt sqrt sqrt sqrt sqrt sqrt |_ sqrt |_ sqrt |_ sqrt sqrt sqrt sqrt sqrt (4!)! _| ! _| ! _| ! _| ! _| ! _| ! _| ! _|. - _Jonathan Vos Post_, May 25 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, May 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008850 | OEIS-A242482

Numbers n such that A242481(n) = ((n*(n+1)/2) mod n + sigma(n) mod n + antisigma(n) mod n) / n = 1.

Conjecture: with number 1 complement of A242483.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, May 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008851 | OEIS-A242483

Numbers n such that A242481(n) = ((n*(n+1)/2) mod n + sigma(n) mod n + antisigma(n) mod n) / n = 2.

Conjecture: with number 1 complement of A242482.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, May 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008852 | OEIS-A242513

a(n) = maximal number of shortest knight's move paths, from origin at center of an infinite open chessboard, to any square within n moves.

For n>4 this sequence is conjectured to be identical to A242511.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Fred Lunnon_, May 16 2014 and May 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008853 | OEIS-A242514

a(n) is the maximal number of shortest knight's move paths, from origin at center of an infinite open chessboard, to square with coordinates <= n.

For n > 5 the distinct terms of this sequence are conjectured to be identical to A242512: precisely, A242514(n) = A242512(ceiling(2*(n+1)/3)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Fred Lunnon_, May 16 2014 and May 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008854 | OEIS-A242533

Number of cyclic arrangements of S={1,2,...,2n} such that the difference of any two neighbors is coprime to their sum.

Conjecture: in this case it seems that NPC(n;S;P)=0 for all odd n, so only the even ones are listed. This is definitely not the case when the property P is replaced by its negation (see A242534). The cycle must alternate odd/even, so that 2 is not a common factor of sum and difference. Therefore only even lengths are possible. This proves the above conjecture. - _Martin Fuller_, Mar 12 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, May 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008855 | OEIS-A242560

Least positive number k such that (n!-k)/(n-k) is an integer.

It is conjectured that a(2n) = n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, May 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008856 | OEIS-A242566

Expansion of (1-sqrt(1-(2*(1-sqrt(1-4*x^2)))/x))/2.

Conjecture D-finite with recurrence: 2*n*(2*n+1)*a(n) +(-49*n^2+97*n-36)*a(n-1) +12*(10*n^2-42*n+41)*a(n-2) +4*(49*n-97)*(n-3)*a(n-3) -544*(n-3)*(n-4)*a(n-4)=0. - _R. J. Mathar_, Jan 25 2020 Conjecture verified using the differential equation -18*x + 6 + (36*x - 12)*y(x) + (1088*x^4 - 396*x^3 - 96*x^2 + 50*x - 6)*y'(x) + (544*x^5 - 196*x^4 - 120*x^3 + 49*x^2 - 4*x)*y''(x) satisfied by the G.f. - _Robert Israel_, Oct 06 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Jun 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008857 | OEIS-A242585

Number of divisors of the n-th positive number that is both triangular and square.

Conjectures: Conjecture: a(n) == 0 mod 9 for n different from 1 and 7 [tested up to n = 300]. - _Ivan N. Ianakiev_, May 29 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, May 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008858 | OEIS-A242647

Sign of (the n-th continued fraction convergent to 2*Pi divided by the n-th convergent to Pi minus 2).

Zeros appear in clusters (see A242857). Conjecture: Remove all zeros and the remaining terms will alternate signs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008859 | OEIS-A242719

Smallest even k such that lpf(k-3) > lpf(k-1) >= prime(n), where lpf=least prime factor (A020639).

a(n) <= A242489(n); a(n) >= prime(n)^2+1. Conjecture: a(n) <= prime(n)^4. - _Vladimir Shevelev_, Jun 01 2014 Conjecture: all numbers a(n)-3 are primes. _Peter J. C. Moses_ verified this conjecture up to a(2001) (cf. with conjecture in A242720). - _Vladimir Shevelev_, Jun 09 2014 Conjecturally, a(n) ~ (prime(n))^2, as n goes to infinity (cf. A246748, A246819). - _Vladimir Shevelev_, Sep 02 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008860 | OEIS-A242720

Smallest even k such that the pair {k-3,k-1} is not a twin prime pair and lpf(k-1) > lpf(k-3) >= prime(n), where lpf = least prime factor (A020639).

a(n) >= prime(n)^2+3. Conjecture: a(n) <= prime(n)^4. - _Vladimir Shevelev_, Jun 01 2014 Conjecture. There are only a finite number of composite numbers of the form a(n)-1. _Peter J. C. Moses_ found only two: a(16)-1 = 4189 = 59*71 and a(20)-1 = 6889 = 83^2 and no others up to a(2501). Most likely, there are no others. - _Vladimir Shevelev_, Jun 09 2014 Conjecturally, a(n) ~ (prime(n))^2, as n goes to infinity (cf. A246748, A246821). - _Vladimir Shevelev_, Sep 02 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008861 | OEIS-A242736

Number of solutions of a^2 + b^2 congruent to -1 modulo the n-th prime.

Is this sequence nondecreasing? The data for the first thousand terms supports this conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David S. Newman_, May 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008862 | OEIS-A242747

Largest number k such that (k!+n!)/(k+n) is an integer.

For n > 2, the conjectured second-to-last value of k that makes (k!+n!)/(k+n) an integer is k = n!-n. It is believed that for n > 2, there is a formula for the last value, a(n). Also, for n > 2, k+n is prime. Thus (k-n+1)!+1 (by Wilson's Theorem) and k!+n! are both multiples of k+n. Further, it is conjectured that the smallest prime factor of k!/n!+1 is k+n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, May 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008863 | OEIS-A242752

Primes p such that pi(p) is a primitive root modulo p, where pi(p) is the number of primes not exceeding p.

According to the conjecture in A232748, this sequence should contain infinitely many primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008864 | OEIS-A242753

Number of ordered ways to write n = k + m with 0 < k <= m such that the inverse of k mod prime(k) among 1, ..., prime(k) - 1 is prime and the inverse of m mod prime(m) among 1, ..., prime(m) - 1 is also prime.

Conjecture: a(n) > 0 for all n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008865 | OEIS-A242754

Positive integers k such that k*p == 1 (mod prime(k)) for some prime p < prime(k).

According to the conjecture in A242753, this sequence should have infinitely many terms. Conjecture: The number of terms not exceeding x > 1 has the main term x/(log x) as x tends to infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008866 | OEIS-A242755

Primes p such that pi(p)*q == 1 (mod p) for some prime q < p, where pi(p) is the number of primes not exceeding p.

According to the conjecture in A242753, this sequence should contain infinitely many primes. Conjecture: The number of such primes not exceeding x > 1 has the main term x/(log x)^2 as x tends to infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008867 | OEIS-A242758

Smallest even k such that lpf(k-1) > lpf(k-3) >= prime(n), where lpf=least prime factor (A020639).

It is easy to prove that min(A242719(n), A242720(n)) >= prime(n)^2+1, while we conjecture that max(A242719(n), A242720(n)) <= prime(n)^4. Thus this conjecture implies there are infinitely many twin primes. - _Vladimir Shevelev_, Jun 01 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008868 | OEIS-A242775

Let b_k=3...3 consist of k>=1 3's. Then a(n) is the smallest k such that the concatenation b_k and prime(n) is prime, or a(n)=0 if there is no such prime.

Conjecture: for n>=4, a(n)>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008869 | OEIS-A242781

Expansion of (1 - 2*x - sqrt(1-4*x))/(4*x^2 + sqrt(1-4*x)*(3*x+1) - 5*x + 1).

Conjecture D-finite with recurrence: 3*(n+1)*a(n) -18*n*a(n-1) +2*(11*n-13)*a(n-2) +6*(n-7)*a(n-3) +(7*n-9)*a(n-4) +2*(2*n-7)*a(n-5)=0. - _R. J. Mathar_, Jan 25 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, May 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008870 | OEIS-A242802

Number of primes p < n such that n - p is a term of A242754.

Conjecture: a(n) > 0 for all n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008871 | OEIS-A242817

a(n) = B(n,n), where B(n,x) = Sum_{k=0..n} Stirling2(n,k)*x^k are the Bell polynomials (also known as exponential polynomials or Touchard polynomials).

Conjecture: It appears that the equation a(n)*e^n = Sum_{k>=0} k^n*n^k/k! is true for every positive integer n. - _Nicolas Nagel_, Apr 20 2016 [This is just the special case x=n of the formula B(n,x) = e^(-x) * Sum_{k>=0} k^n*x^k/k!; see for example the World of Mathematics link. - _Pontus von Brömssen_, Dec 05 2020]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, May 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008872 | OEIS-A242819

Number T(n,k) of permutations of [n] with exactly k occurrences of the consecutive step pattern up, down, down; triangle T(n,k), n>=0, 0<=k<=max(0,floor((n-1)/3)), read by rows.

Conjecture: c(k) = c(0) * (c(0)-1)^k / (3^k * k!).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, May 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008873 | OEIS-A242857

Position of zeros in A242647.

Conjecture: The sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hans Havermann_, May 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008874 | OEIS-A242879

Least positive integer k < n such that k*p == 1 (mod prime(k)) for some prime p < prime(k) and (n-k)*q == 1 (mod prime(n-k)) for some prime q < prime(n-k), or 0 if such a number k does not exist.

According to the conjecture in A242753, a(n) should be positive for all n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008875 | OEIS-A242884

Least number k such that (k^k-n^n)/(k-n) is prime or 0 if no such number exists.

a(n) = 0 is confirmed for k <= 5000. (Which means that the zeros are at present only conjectural. - _N. J. A. Sloane_, May 26 2014)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, May 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008876 | OEIS-A242922

Least number k such that (k^n+n^k)/(k+n) is prime or 0 if no such k exists.

a(n) = 0 is definite for n=1. For other n-values where a(n) = 0, it is confirmed for k <= 5000. These are only conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, May 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008877 | OEIS-A242928

Least number k such that (k^k+n^n)/(k+n) is prime or 0 if no such k exists.

a(n) = 0 is confirmed for k <= 2500. These are just conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, May 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008878 | OEIS-A242932

Numbers n such that k*n/(k+n) is prime for some k.

A number n belongs to this sequence iff n belongs to the set {p+1, 2*p, p*(p+1)}, for some prime p. In particular, the conjecture regarding A016742 is true (see Fried link). - _Sela Fried_, Mar 27 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, May 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008879 | OEIS-A242950

Number of ordered ways to write n = k + m with k > 1 and m > 1 such that the least nonnegative residue of prime(k) modulo k is a square and the least nonnegative residue of prime(m) modulo m is a prime.

Conjecture: (i) a(n) > 0 for all n > 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008880 | OEIS-A242970

Decimal expansion of the constant rho = lim f(n)^(1/n), where f(n) = A214833(n) is the number of arithmetic formulas for n (cf. comments).

This is the constant rho, given on page 2 of E. K. Gnang and others. From the abstract: "An arithmetic formula is an expression involving only the constant 1, and the binary operations of addition and multiplication, with multiplication by 1 not allowed. We obtain an asymptotic formula for the number of arithmetic formulas evaluating to n as n goes to infinity, solving a conjecture of E. K. Gnang and D. Zeilberger."

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Vos Post_, Jun 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008881 | OEIS-A242976

Number of ways to write n = i + j + k with 0 < i <= j <= k such that prime(i) mod i, prime(j) mod j and prime(k) mod k are all triangular numbers, where prime(m) mod m denotes the least nonnegative residue of prime(m) modulo m.

Conjecture: (i) a(n) > 0 for all n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008882 | OEIS-A242992

Least k>n/2, k<n, such that 2^(n-k)-1 divides 2^k-2, or 0 if no such k exists.

Conjecture: a(n) = n - A032742(n-1), for n > 2. - _Ridouane Oudra_, Mar 17 2024 The conjecture is true (see Fried link). - _Sela Fried_, Feb 10 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Aug 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008883 | OEIS-A243022

Expansion of -(2*x)/(1-sqrt(1-(2*(1-sqrt(1-4*x^2))))/x)).

Conjecture D-finite with recurrence:
+2*n*(n-1)*(2*n-3)*(5*n-12)*a(n)
-(n-1)*(85*n^3-459*n^2+776*n-400)*a(n-1)

+4*(-40*n^4+396*n^3-1455*n^2+2324*n-1360)*a(n-2) +4*(n-4)*(170*n^3-1258*n^2+3003*n-2185)*a(n-3) +16*(n-4)*(n-5)*(20*n^2-118*n+143)*a(n-4) -272*(n-4)*(n-5)*(n-6)*(5*n-7)*a(n-5)=0. - _R. J. Mathar_, Jul 15 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Jun 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008884 | OEIS-A243050

Composite integers n such that n-1 divided by the binary period of 1/n (=A007733(n)) equals an integral power of 2.

Contains all composite Fermat numbers A000215(k) = 2^(2^k)+1 (which are composite for 5<=k<=32 and conjecturally for any k>=5). In particular, a(5) = A000215(5), a(12) = A000215(6), and a(13) <= A000215(7) = 2^128+1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, May 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008885 | OEIS-A243101

a(n) = (sum_{k=0}^{n-1}(4*k^3-1)*C(n-1,k)*C(n+k,k))/n^2, where C(m,k) denotes the binomial coefficient m!/(k!*(m-k)!).

Conjecture: Let n be any positive integer. The author proved the conjecture in the latest version of arXiv:1408.5381, thus a(n) is always an integer. - _Zhi-Wei Sun_, Sep 01 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008886 | OEIS-A243155

Larger of two consecutive primes whose positive difference is a cube.

The observation is false (unless it is just meant to refer to the Data or the b-file): it is conjectured that every positive even number occurs infinitely often as a prime gap. According to the b-file for A001632 the first case where the gap is 6^3 is where the larger prime is 202551883. - _Robert Israel_, Apr 15 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _K. D. Bajpai_, May 31 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008887 | OEIS-A243160

Least number k such that k^n - k^(n-1) + k^(n-2) - ... + (-1)^n is prime or 0 if no such k exists.

That is only a conjecture. - _Robert Israel_, Apr 15 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, May 31 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008888 | OEIS-A243228

Number of isoscent sequences of length n with exactly two ascents.

G.f.: x^4*(3 - 5*x + x^2)*(1 - x - x^2) / ((1 - x)^3*(1 - 2*x)^2*(1 - 4*x + 3*x^2 + x^3)) (conjectured). - _Colin Barker_, May 05 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_ and _Alois P. Heinz_, Jun 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008889 | OEIS-A243229

Number of isoscent sequences of length n with exactly three ascents.

G.f.: x^6*(17 - 213*x + 1118*x^2 - 3135*x^3 + 4851*x^4 - 3492*x^5 - 262*x^6 + 1707*x^7 + 82*x^8 - 1050*x^9 + 189*x^10 + 297*x^11 - 122*x^12 + 11*x^13 + 3*x^14) / ((1 - x)^4*(1 - 2*x)^3*(1 - 4*x + 3*x^2 + x^3)^2*(1 - 8*x + 21*x^2 - 18*x^3 - 3*x^4 + 5*x^5 + 3*x^6)) (conjectured). - _Colin Barker_, May 04 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_ and _Alois P. Heinz_, Jun 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008890 | OEIS-A243230

Number of isoscent sequences of length n with exactly four ascents.

G.f.: x^7*(2 + 58*x - 3648*x^2 + 72673*x^3 - 842380*x^4 + 6646402*x^5 - 38326607*x^6 + 167767231*x^7 - 568689800*x^8 + 1503499267*x^9 - 3082877524*x^10 + 4785455208*x^11 - 5266898088*x^12 + 3276181912*x^13 + 633509453*x^14 - 3771906622*x^15 + 3792580950*x^16 - 1396200816*x^17 - 686750857*x^18 + 1195721330*x^19 - 1109678200*x^20 + 1163509247*x^21 - 851661184*x^22 + 47666957*x^23 + 460007148*x^24 - 312042368*x^25 - 2183667*x^26 + 84057866*x^27 - 24458487*x^28 - 9560501*x^29 + 5700026*x^30 + 307683*x^31 - 694453*x^32 + 45182*x^33 + 51672*x^34 - 3534*x^35 - 1395*x^36 - 135*x^37) / ((1 - x)^5*(1 - 2*x)^4*(1 - 4*x + 3*x^2 + x^3)^3*(1 - 8*x + 21*x^2 - 18*x^3 - 3*x^4 + 5*x^5 + 3*x^6)^2*(1 - 16*x + 105*x^2 - 361*x^3 + 676*x^4 - 606*x^5 + 68*x^6 + 192*x^7 + 83*x^8 - 157*x^9 - 44*x^10 + 45*x^11 + 15*x^12)) (conjectured). - _Colin Barker_, May 05 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_ and _Alois P. Heinz_, Jun 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008891 | OEIS-A243255

Expansion of 1/2 - (1/2)*(sqrt((2*sqrt(-3*x^2 - 2*x + 1) + 3*x - 2)/x)).

Conjecture D-finite with recurrence: 2*n*(2*n+1)*a(n) + (-65*n^2+119*n-42)*a(n-1) + 3*(97*n^2-373*n+342)*a(n-2) - 9*(23*n-68)*(n-3)*a(n-3) - 567*(n-3)*(n-4)*a(n-4) = 0. - _R. J. Mathar_, Jul 15 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Jun 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008892 | OEIS-A243362

Numbers k such that A243361(k) = 123456789.

Conjecture 1: sequence is infinite. Conjecture 2: a(1), a(2) and a(3) are composites; there are no other numbers n > 3 such that a(n) = composite number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jun 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008893 | OEIS-A243364

Primes whose reverse concatenation of divisors (A176558) contains all the digits 1-9 exactly once; the number of digits 0 is arbitrary (in base 10).

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jun 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008894 | OEIS-A243366

Number T(n,k) of Dyck paths of semilength n having exactly k (possibly overlapping) occurrences of the consecutive steps UDUUDU (with U=(1,1), D=(1,-1)); triangle T(n,k), n>=0, 0<=k<=max(0,floor(n/2)-1), read by rows.

Conjecture: Generally, column k is asymptotic to c(k) * d^n * n^(k-3/2), where d = 3.882159026862850674719436890964338 4... is the root of the equation d^8 - 2*d^7 - 10*d^6 + 12*d^5 - 5*d^4 - 2*d^3 - 5*d^2 - 8*d - 3 = 0, and c(k) are specific constants (independent on n). - _Vaclav Kotesovec_, Jun 05 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jun 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008895 | OEIS-A243374

Least number k such that n//k and k//n are both prime where // is the concatenation function, or 0 if no such k exists.

If n ends in a 2, 4, 5, 6, 8, or 0, then a(n) = 0. It is conjectured that the converse is true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Jun 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008896 | OEIS-A243512

Least index i for which A243473(i)=n, or 0 if no such index exists.

Motivated by the observation that some small numbers (2,12,14,18,...) occur only very late in the recently added sequence A243473, but all numbers seem to appear sooner or later. (The definition is completed by "0 if no such index exists" to guarantee well-definedness in absence of a proof, but I conjecture that no such 0 will ever occur.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008897 | OEIS-A243513

Number of length n+2 0..4 arrays with no three elements in a row with pattern aba or abb (with a!=b) and new values 0..4 introduced in 0..4 order.

Conjectures from _Colin Barker_, Nov 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008898 | OEIS-A243514

Number of length n+2 0..5 arrays with no three elements in a row with pattern aba or abb (with a!=b) and new values 0..5 introduced in 0..5 order.

Conjectures from _Colin Barker_, Nov 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008899 | OEIS-A243515

Number of length n+2 0..6 arrays with no three elements in a row with pattern aba or abb (with a!=b) and new values 0..6 introduced in 0..6 order.

Conjectures from _Colin Barker_, Nov 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008900 | OEIS-A243516

Number of length n+2 0..7 arrays with no three elements in a row with pattern aba or abb (with a!=b) and new values 0..7 introduced in 0..7 order.

Conjectures from _Colin Barker_, Nov 02 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008901 | OEIS-A243585

Expansion of x*log'(C(C(x)-1)-1), C(x) = (1-sqrt(1-4*x))/(2*x).

Conjecture D-finite with recurrence: 2*n*(2*n-1)*(3*n-5)*a(n) +(-123*n^3+328*n^2-249*n+60)*a(n-1) +50*(n-1)*(2*n-3)*(3*n-2)*a(n-2)=0. - _R. J. Mathar_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Jun 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008902 | OEIS-A243610

Irregular triangular array of all the integers, each exactly once, ordered as in Comments.

Let F = A000045 (the Fibonacci numbers). To construct the array, decree the first 4 rows as in the Example. Thereafter, row n consists of F(n) numbers in increasing order, generated as follows: the F(n-1) numbers 2*x from x in row n-1, together with the F(n-2) numbers 1 - 2*x from numbers x in row n-2. For n >= 3, row n consists of F(n-1) negative integers and F(n-2) positive integers; also, row n consists of F(n-1) even integers and F(n-2) odd integers. Conjecture: Every row contains F(k) or -F(k) for some k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008903 | OEIS-A243642

Expansion of -(4*x)/((1-sqrt(1-4*x))/2-sqrt(-3*(1-sqrt(1-4*x))+(1-sqrt(1-4*x))^2/4+1)+1).

Conjecture D-finite with recurrence: +16*n*(n-1)*(n-2)*a(n) -4*(n-1)*(n-2)*(43*n-54)*a(n-1) +2*(n-2)*(52*n^2+695*n-2439)*a(n-2) +(3479*n^3-42714*n^2+169789*n-219234)*a(n-3) +2*(-4633*n^3+64020*n^2-293825*n+447594)*a(n-4) +4*(2*n-11)*(296*n^2-3643*n+11187)*a(n-5) +24*(n-7)*(2*n-11)*(2*n-13)*a(n-6)=0. - _R. J. Mathar_, Jan 25 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Jun 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008904 | OEIS-A243662

Triangle read by rows: the reversed x = 1+q Narayana triangle at m=2.

T(n,k) = (binomial(3*n+1,n) * binomial(n,k-1) * binomial(n-1,k-1)) / (binomial(3*n,k-1) * (3*n+1)) = (A001764(n) * A001263(n,k) * k) / binomial(3*n,k-1) for 1 <= k <= n (conjectured). - _Werner Schulte_, Nov 22 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008905 | OEIS-A243675

Number of hypoplactic classes of 3-parking functions of length n.

This is almost certainly the sequence of small 4-Schroeder numbers as defined by Yang-Jiang (2021). It would be nice to have a proof. Then we could confirm Weiner's conjectured formula, and extend the sequence. Yang & Jiang (2021) give an explicit formula for the small m-Schroeder numbers in Theorems 2.4 and 2.9. - _N. J. A. Sloane_, Mar 28 2021 a(n) = Sum_{i=0..n} Sum_{j=0..i} (-2)^(n-i)*binomial(i,j)*binomial(3i+j, n)*binomial(n+1,i)/(n+1) (conjectured). - _Michael D. Weiner_, May 25 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008906 | OEIS-A243676

Number of hypoplactic classes of 4-parking functions of length n.

This is almost certainly the sequence of small 5-Schroeder numbers as defined by Yang-Jiang (2021). It would be nice to have a proof. Then we could confirm Weiner's conjectured formulas, and extend the sequence. Yang & Jiang (2021) give an explicit formula for the small m-Schroeder numbers in Theorems 2.4 and 2.9. - _N. J. A. Sloane_, Mar 28 2021 a(n+1) = Sum_{i=0..n} Sum_{j=0..i} (-2)^(n-i)*binomial(i,j)*binomial(4*i+j, n)*binomial(n+1,i)/(n+1) (conjectured). - _Michael D. Weiner_, May 25 2017 a(n) = Sum_{i=1..n} binomial(4*n, i-1)*binomial(n, i)*2^(i-1)/n (conjectured). - _Michael D. Weiner_, Jul 24 2019 [This is correct for n>0 - _Jun Yan_, Apr 13 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008907 | OEIS-A243677

Number of hypoplactic classes of 5-parking functions of length n.

a(n) = Sum_{i=0..n} Sum_{j=0..i} (-2)^(n-i)*binomial(i,j)*binomial(5i+j, n)*binomial(n+1,i)/(n+1) (conjectured). - _Michael D. Weiner_, May 25 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008908 | OEIS-A243700

The lexicographically earliest sequence of distinct terms with a(1) = 1 such that a(n) divides the sum of the first a(n) terms.

Conjecture: For n > 1000, a(n) falls within 1% of one of the following six values. a(n) = n, 1.576385*n, 1.788185*n, 2.576385*n, 2.788185*n, or 3.576285*n, using floor at the low bound and ceiling at the high bound, inclusive. Conjectured asymptotic slopes of the 6 lines in the function graph are, from low to high: 1, sqrt(3)/3+1, sqrt(3)/6+3/2, sqrt(3)/3+2, sqrt(3)/6+5/2, sqrt(3)/3+3. - _Hugo Pfoertner_, Dec 19 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008909 | OEIS-A243750

Indices of records in A112801 = number of partitions of 2n-1 into three numbers with two distinct prime factors each.

However, 998 is larger than 103 and congruent to 2 (mod 3), but is not a term, because A112801(995) = 41603 > 41594 = A112801(998). The conjecture holds through a(361) = 995 and first fails at a(362) = 1001 (not 998). - _Qiyuan Gu_, Jul 24 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008910 | OEIS-A243760

Expansion of (sqrt(8*x+4*sqrt(1-4*x)-3)-1)/(2*sqrt(1-4*x)-2).

Conjecture D-finite with recurrence: 245*n*(n-1)*(n+1)*a(n) -140*n*(n-1)*(28*n-53)*a(n-1) +4*(n-1)*(5812*n^2-29864*n+36585)*a(n-2) +16*(-3368*n^3+35040*n^2-111784*n+110667)*a(n-3) +128*(-152*n^3-1224*n^2+14756*n-29655)*a(n-4) +2048*(2*n-9)*(74*n^2-567*n+988)*a(n-5) -98304*(n-6)*(2*n-9)*(2*n-11)*a(n-6)=0. - _R. J. Mathar_, Jun 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Jun 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008911 | OEIS-A243764

Expansion of -((2*sqrt(1-4*x)-2)*x)/(sqrt(8*x+4*sqrt(1-4*x)-3)-1).

Conjecture: 245*n*(n-1)*(n-2)*a(n) -140*(n-1)*(n-2)*(28*n-81)*a(n-1) +4*(n-2)*(5812*n^2-41488*n+72261)*a(n-2) +16*(-3368*n^3+45144*n^2-191968*n+260859)*a(n-3) +128*(-152*n^3-768*n^2+16748*n-45483)*a(n-4) +2048*(2*n-11)*(74*n^2-715*n+1629)*a(n-5) -98304*(n-7)*(2*n-11)*(2*n-13)*a(n-6)=0. - _R. J. Mathar_, Jun 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Jun 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008912 | OEIS-A243814

Expansion of -(x*(1-sqrt((2*(1-sqrt(4*x^2+1)))/x+1)))/(1-sqrt(4*x^2+1)) - 1.

Conjecture D-finite with recurrence: 2*n*(n+1)*(2*n-1)*a(n) -n*(47*n^2-127*n+86)*a(n-1) +4*(38*n^3-186*n^2+291*n-145)*a(n-2) +4*(-94*n^3+724*n^2-1887*n+1650)*a(n-3) +16*(n-4)*(64*n^2-382*n+595)*a(n-4) -16*(n-5)*(47*n-174)*(n-4)*a(n-5) +1920*(n-5)*(n-6)*(n-4)*a(n-6)=0. - _R. J. Mathar_, Jan 25 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Jun 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008913 | OEIS-A243816

Expansion of (x*sqrt(4*x^2+1)-x)/(x*sqrt(-(2*sqrt(4*x^2+1)-x-2)/x) + sqrt(4*x^2+1)-x-1).

Conjecture D-finite with recurrence: 2*n*(n-1)*(2*n-3)*a(n) -(n-1)*(47*n^2-221*n+260)*a(n-1) +4*(38*n^3-300*n^2+777*n-660)*a(n-2) +4*(-94*n^3+1006*n^2-3617*n+4355)*a(n-3) +16*(n-5)*(64*n^2-510*n+1041)*a(n-4) -16*(n-5)*(n-6)*(47*n-221)*a(n-5) +1920*(n-5)*(n-6)*(n-7)*a(n-6)=0. - _R. J. Mathar_, Jan 25 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Jun 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008914 | OEIS-A243826

Maximum number of clues in a certain class of n X n crossword puzzles.

Except for n = 7, 11, and 19, conjectured recursive formula is a(n) = a(n-4) + 4(n-3) - [2 if mod(n,8) in {1,7}]. In particular, conjectured explicit formula is a(n) = 2n floor((n+1)/4) if mod(n,4) = 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rob Pratt_, Jun 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008915 | OEIS-A243924

Irregular triangular array of taxicab norms of Gaussian integers in array G generated as at Comments.

An array G of Gaussian integers is generated as follows: (row 1) = (0), and for n >=2, row n consists of the numbers x+1 and then i*x, where duplicates are deleted as they occur. Every Gaussian integer occurs exactly once in G. The taxicab norm of a Gaussian integer b+c*i is the taxicab distance (also known as Manhattan distance) from 0 to b+c*i, given by |b|+|c|. The norms of numbers in row n are given here in nondecreasing order. Conjecture: the number of numbers in row n is 4n-13 for n >= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008916 | OEIS-A243937

Even numbers n>=6 for which lpf(n-1) > lpf(n-3), where lpf = least prime factor.

Conjecture: All differences are 2, 4 or 6 such that there are no two consecutive terms 2 (..., 2, 2, ...), no two consecutive terms 4, while consecutive terms 6 occur 1, 2, 3 or 4 times; also consecutive pairs of terms 2, 4 appear 1, 2, 3 or 4 times. The conjecture is verified up to n = 2.5*10^7. - _Vladimir Shevelev_ and _Peter J. C. Moses_, Jul 11 2014 Divisibility by 3 means 6m is in the sequence for all m > 0, and 6m + 4 never is, while 6m + 2 is undetermined. Divisibility by 5 means 30m + 8 is always in the sequence, and 30m + 26 never is. This proves the above conjecture. - _Jens Kruse Andersen_, Aug 19 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jul 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008917 | OEIS-A243956

Positive numbers n without a decomposition into a sum n = i+j such that 6i-1, 6i+1, 6j-1, 6j+1 are twin primes.

Conjecture: any integer n > 701 has a decomposition into a sum n = i+j such that 6i-1, 6i+1, 6j-1, 6j+1 are twin primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lear Young_, Jun 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008918 | OEIS-A243977

a(n) is the largest run of identical digits that n^k can end with for some k, or 0 if there is no limit to such runs.

a(53), a(71) and a(77) are conjectured to be infinite. See A244187.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Jun 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008919 | OEIS-A243999

Decimal expansion of B (negated), a constant related to Glaisher's constant A and the Gaussian unitary ensemble hypothesis.

Decimal expansion of B (negated), a constant related to Glaisher's constant A and the Gaussian unitary ensemble hypothesis.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Jun 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008920 | OEIS-A244009

Decimal expansion of 1 - log(2).

Equals 1/(4 - 4/(7 - 12/(10 - ... - 2*n*(n-1)/((3*n+1) - ...)))) (an equivalent continued fraction for 1 - log(2) was conjectured by the Ramanujan machine). - _Peter Bala_, Mar 04 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Jun 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008921 | OEIS-A244041

Sum of digits of n written in fractional base 4/3.

Conjecture: a(n) >> log(n), hence a(n) ≍ log(n). - _Charles R Greathouse IV_, Nov 03 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hailey R. Olafson_, Jun 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008922 | OEIS-A244059

Initial digit of the decimal expansion of n^(n^(n^n)) or n^^4 (in Don Knuth's up-arrow notation).

Conjecture: the distribution of the initial digits obey G. K. Zipf's law.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Munafo_ and _Robert G. Wilson v_, Jun 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008923 | OEIS-A244240

Number of partitions of n into 4 parts such that every i-th smallest part (counted with multiplicity) is different from i.

Conjectures from _Chai Wah Wu_, Apr 18 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jun 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008924 | OEIS-A244241

Number of partitions of n into 5 parts such that every i-th smallest part (counted with multiplicity) is different from i.

Conjectures from _Chai Wah Wu_, Apr 18 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jun 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008925 | OEIS-A244242

Number of partitions of n into 6 parts such that every i-th smallest part (counted with multiplicity) is different from i.

Conjectures from _Chai Wah Wu_, Apr 18 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jun 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008926 | OEIS-A244243

Number of partitions of n into 7 parts such that every i-th smallest part (counted with multiplicity) is different from i.

Conjectures from _Chai Wah Wu_, Apr 18 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jun 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008927 | OEIS-A244250

Triangle read by rows in which row n lists the widths in the first octant of the symmetric representation of sigma(n).

It appears that the leading diagonal is also A067742 (which was conjectured by _Michel Marcus_ in the entry A237593 and checked with two Mathematica functions up to n = 100000 by _Hartmut F. W. Hoft_).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Oct 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008928 | OEIS-A244254

Positive integers n such that all the gaps prime(n+i+1) - prime(n+i) (i = 0..5) are triangular numbers.

Conjecture: For any integer m > 0, there are infinitely many positive integers n such that all the gaps prime(n+i+1) - prime(n+i) (i = 0, ..., m-1) are triangular numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008929 | OEIS-A244264

Positive integers n such that all the gaps prime(n+i+1) - prime(n+i) (i = 0..7) are powers of two.

Conjecture: For any integer m > 0, there are infinitely many positive integers n such that all the gaps prime(n+i+1) - prime(n+i) (i = 0, ..., m-1) are powers of two.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008930 | OEIS-A244266

Positive integers n such that all those prime(n+i) + prime(n+j) (i,j = 0..7) are squarefree.

Conjecture: For any integer m > 0, there are infinitely many positive integers n such that all those prime(n+i) + prime(n+j) (i,j = 0, ..., m) are squarefree.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008931 | OEIS-A244272

Decimal expansion of exp(gamma)/log(2), a conjectural constant related to the asymptotic counting of Mersenne primes, where gamma is Euler's constant.

Decimal expansion of exp(gamma)/log(2), a conjectural constant related to the asymptotic counting of Mersenne primes, where gamma is Euler's constant. Let M(n) denote the number of primes p <= n, for which 2^p-1 is prime. It is conjectured that M(n) -> infinity and that M(n) ~ exp(gamma)/log(2)*log(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Jun 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008932 | OEIS-A244290

Smallest prime a(n) = x^2 + y^2 such that c^2 + d^2 = A002313(n) and c*x + d*y = 1, where c,d,x,y are integers.

Conjecture: a(n) < A002313(n)^2 for n > 1. - _Thomas Ordowski_, Dec 28 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Jun 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008933 | OEIS-A244294

Positive integers n such that (prime(n+i+1) - prime(n+i))/2 is prime for every i = 0, ..., 12.

Conjecture: For any integer m > 0, there are infinitely many positive integers n such that (prime(n+i+1) - prime(n+i))/2 is prime for every i = 0, ..., m-1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008934 | OEIS-A244311

Primes (with d digits, say) that generate another prime when acted on by the "standard" superpermutation of length A007489(d) of d elements (cf. comment).

For primes with more than four digits, the sequence is based on the current information about the conjectured minimal length.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Abhiram R Devesh_, Jun 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008935 | OEIS-A244331

Number of binary digits in the high-water marks of the terms of the continued fraction of the base-2 Champernowne constant.

Conjecture: partial sums of A296965 (equivalent to observation about A183155 below). - _Sean A. Irvine_, Jul 16 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John K. Sikora_, Jun 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008936 | OEIS-A244349

Positive integers n such that all the differences prime(n+j) - prime(n+i) with 0 <= i < j <= 9 are practical numbers.

Conjecture: For any integer m > 0, there are infinitely many positive integers n such that all the differences prime(n+j) - prime(n+i) with 0 <= i < j <= m are practical numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008937 | OEIS-A244364

a(n) is the least integer m > 1 such that n is the largest number of identical digits that can end m^k for positive integer k.

Conjecture for n>5: a(n) = 10*2^(n-1)-1, k = (2r+1)*5^(n-2), r>=0. - _Lars Blomberg_, Jul 08 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Jun 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008938 | OEIS-A244424

Least number k > 0 such that concatenating k consecutive natural numbers beginning with n is prime, or 0 if no such number exists.

The current zero values are only conjectural: a(n) > 5000 - n for all a(n) = 0 shown. [Edited by _M. F. Hasler_, Apr 27 2017]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Jun 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008939 | OEIS-A244432

E.g.f.: exp(Sum_{n>=1} Pell(n)*x^n/n), where Pell(n) = A000129(n).

a(n) == 0 (mod 7^k) for n >= 7*k, for k >= 1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008940 | OEIS-A244474

4th-largest term in n-th row of Stern's diatomic triangle A002487.

G.f.: (-2-2*x-4*x^2-3*x^3-2*x^4-x^5-3*x^6-2*x^7-x^8-x^9-x^10)/(-1+x+x^2) (conjectured) - _Jean-François Alcover_, Mar 12 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008941 | OEIS-A244475

5th-largest term in the n-th row of Stern's diatomic triangle A002487.

Conjectured g.f.: -x^3*(x^14+ x^13+ x^12+ 2*x^11 +3*x^10 +5*x^9 +8*x^8 +x^7 +3*x^6 +3*x^5 +2*x^4 +4*x^3 +5*x^2 +2*x +1) / (x^2+x-1). - _Alois P. Heinz_, Jun 20 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008942 | OEIS-A244626

Composite numbers k congruent to 5 (mod 8) such that 2^((k-1)/2) mod k = k-1.

Conjecture 1: if p is an odd prime congruent to {3,5} (mod 8) then 2^((p-1)/2) mod p = p-1. This conjecture has been tested to 10^8. Conjecture 1 is true. With p = 2k+1 then 2^k mod (2k+1) == 2k. So 2k+1 | 2k-2^k. Prime numbers 2k+1 == +-3 (mod 8) are the prime numbers such that 2k+1 | 2^k+1 (Comments A007520). A reflection across the x-axis and +1 translation across the y-axis of the graph (2k-2^k) / (2k+1) gives the graph (2^k+1) / (2k+1). So the k values of both 2k+1 | 2k-2^k and 2k+1 | 2^k+1 are identical. - _Hilko Koning_, Feb 04 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary Detlefs_, Jul 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008943 | OEIS-A244628

Composite numbers k such that k == 3 (mod 8) and 2^((k-1)/2) == -1 (mod k).

Conjecture 1: if p is a prime congruent to {3,5} mod 8 then 2^((p-1)/2) mod p = p-1. This conjecture has been tested to 10^8. Conjecture 1 is true. With p = 2k+1 then 2^k mod (2k+1) == 2k. So 2k+1 | 2k-2^k . Prime numbers 2k+1 == +-3 (mod 8) are the prime numbers such that 2k+1 | 2^k+1 (Comments A007520). A reflection across the x-axis and +1 translation across the y-axis of the graph (2k-2^k) / (2k+1) gives the graph (2^k+1) / (2k+1). So the k values of both 2k+1 | 2k-2^k and 2k+1 | 2^k+1 are identical. - _Hilko Koning_, Feb 04 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary Detlefs_, Jul 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008944 | OEIS-A244644

Consider the method used by Archimedes to determine the value of Pi (A000796). This sequence denotes the number of iterations of his algorithm which would result in a difference of less than 1/10^n from that of Pi.

Conjecture: There exists a c such that a(n) = floor(n*log(10)/log(4)+c); where c is in the range [0.08554,0.10264]. Critical values to narrow the range are believed to be at a(74), a(133), a(192), a(251), a(310), a(366), a(425), a(484). - _A.H.M. Smeets_, Jul 23 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _William H. Richardson_ and _Robert G. Wilson v_, Jul 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008945 | OEIS-A244713

Positive numbers primitively represented by the binary quadratic form (1, 1, -2).

Conjectures from _Colin Barker_, Oct 31 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jul 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008946 | OEIS-A244743

Smallest number n with ||n-1||-||n|| = k where ||n||=A005245(n) denotes the complexity of n.

It is conjectured that ||n-1||-||n|| is not bounded. But there is no proof that the sequence is infinite or is well defined.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juan Arias-de-Reyna_, Jul 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008947 | OEIS-A244884

Expansion of (-2 +x^2 +x -x*sqrt(1-2*x-3*x^2))/(-1 +x -sqrt(1-2*x-3*x^2)).

Conjecture D-finite with recurrence: (n+2)*a(n) +(-3*n-1)*a(n-1) -n*a(n-2) +3*(n-3)*a(n-3)=0. - _R. J. Mathar_, Jan 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008948 | OEIS-A244886

G.f.: (1-x+sqrt(1-2*x-3*x^2))/(1-3*x+x^2+x^3+(1-x^2)*sqrt(1-2*x-3*x^2)).

Conjecture D-finite with recurrence: n*a(n) +(-5*n+3)*a(n-1) +2*(n)*a(n-2) +(13*n-30)*a(n-3) +3*(-1)*a(n-4) +(-8*n+21)*a(n-5) +3*(-n+3)*a(n-6)=0. - _R. J. Mathar_, Jan 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008949 | OEIS-A244973

a(n) = Sum_{k=0..n} (-1)^k*C(n, k)^2*C(2*k, k), where C(n, k) denotes the binomial coefficient n!/(k!*(n-k)!).

Conjecture: For any prime p > 5 and positive integer n, the number (a(p*n) - a(n))/(p*n)^3 is always a p-adic integer. Conjecture: a(n) = Sum_{k = 0..n} binomial(n, k)*b(k), where b(n) = Sum_{k = 0..n} (-1)^k*binomial(n, k)^2*binomial(2*k, n). [Added Mar 16 2025: this conjecture can be verified using the MulZeil procedure in Doron Zeilberger's MultiZeilberger package for Maple]. - _Peter Bala_, Jul 19 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008950 | OEIS-A245024

Even numbers n for which lpf(n-1) < lpf(n-3), where lpf = least prime factor.

Conjecture: All differences are 2,4 or 6 such that no two consecutive terms 2 (...,2,2,...), no two consecutive terms 4, while consecutive terms 6 occur 1,2,3 or 4 times; also consecutive pairs of terms 4,2 appear 1,2,3 or 4 times. Conjecture is verified up to n = 2.5*10^7. - _Vladimir Shevelev_ and _Peter J. C. Moses_, Jul 11 2014 The first comment is wrong as stated. This would fix it: for every pair of primes (p,q), p>q>=3, if a(n)==1 (mod p) then a(n) not==3 (mod q). Divisibility by 3 means 6m+4 is in the sequence for all m>0, and 6m never is, while 6m+2 is undetermined. Divisibility by 5 means 30m+26 is always in the sequence, and 30m+8 never is. This proves the above conjecture. - _Jens Kruse Andersen_, Jul 13 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jul 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008951 | OEIS-A245046

Smallest non-Fibonacci number k such that k^2 + F(n)^2 = f1*f2 where F(n) = A000045(n) and f1, f2 are distinct Fibonacci numbers.

Conjecture : if k exists, the number of solutions non-Fibonacci k of the equation k^2 + F(n)^2 = f1*f2 is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jul 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008952 | OEIS-A245080

Numbers such that omega(a(n)) is a proper divisor of bigomega(a(n)).

The density of these numbers, i.e., the ratio n/a(n), apparently decreases with n, reaching 0.04420... for n = 10000000. Conjecture: n/a(n) might have a nonzero limit below 0.0427 (the density found in the interval 9500000 < n <= 10000000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Jul 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008953 | OEIS-A245088

a(n) = (Sum_{k=0..n-1} (8*k^2+12*k+5)*A244973(k))/n^2.

Conjecture: All the terms are odd integers. Moreover, a(p) == 3 (mod p) for any prime p. In 2015 Guo, Mao and Pan confirmed the conjecture partially by proving that n*a(n) is always an integer. - _Zhi-Wei Sun_, Nov 14 2015 Conjecture: +n^2*(23656*n-2233)*(n-1)*a(n) +(n-1)*(23656*n^3+655645*n^2-2273556*n+1915600)*a(n-1) -2*(n-2) *(177048*n^3-3748011*n^2+13870329*n-15407140)*a(n-2) -2*(n-3) *(2020728*n^3-18138829*n^2+63885382*n-73802216)*a(n-3) -5*(n-4) *(782552*n^3-11993599*n^2+55520403*n-80051816)*a(n-4) +125*(66088*n-203507)*(n-5)*(n-4)^2*a(n-5)=0. - _R. J. Mathar_, Jul 12 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008954 | OEIS-A245089

The unique integer r with |r| < prime(n)/2 such that B_{prime(n)-2}(1/3) == r (mod prime(n)), where B_m(x) denotes the Bernoulli polynomial of degree m.

Conjecture: a(n) = 0 infinitely often. In other words, there are infinitely many primes p > 3 such that B_{p-2}(1/3) == 0 (mod p). Zhi-Wei Sun made many conjectures on congruences involving B_{p-2}(1/3), see the Sci. China Math. paper and arXiv:1407.0967.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008955 | OEIS-A245094

Total squares count in n-th generation of Pythagoras tree variation which is rhombitrihexagonal tiling.

Conjecture: In the limit n -> infinity this construction produces one of the eight planar semiregular tessellations (one of the 11 Archimedean tessellations, the other three being regular). This is the tessellation (3,4,6,4) because of the sequence of regular 3-, 4- and 6-gons around each vertex. See the Eric Weisstein link. - _Wolfdieter Lang_, Nov 23 2014 Conjectures from _Colin Barker_, Nov 12 2014: (Start) It follows from the above conjecture that this sequence consists of interlaced polynomials for n > 5: a(2n) = 3*(3n-1) and a(2n+1) = 9*n. - _Avi Friedlich_, May 09 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Nov 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008956 | OEIS-A245158

Number of length n 0..3 arrays with new values introduced in order from both ends.

Conjectures from _Colin Barker_, Nov 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008957 | OEIS-A245159

Number of length n 0..4 arrays with new values introduced in order from both ends.

Conjectures from _Colin Barker_, Nov 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008958 | OEIS-A245203

Least k>=0 such that A004767(n) = 4n+3 is the mean of primes 4(n-k)+3 and 4(n+k)+3; or -1 if no such k exists.

It has been conjectured that A004767(n) = 4n+3 is either prime or the mean of two primes in sequence A004767.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jul 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008959 | OEIS-A245204

The unique integer r with |r| < prime(n)/2 such that E_{prime(n)-3}(1/4) == r (mod prime(n)), where E_m(x) denotes the Euler polynomial of degree m.

Conjecture: a(n) = 0 infinitely often. In other words, there are infinitely many odd primes p such that E_{p-3}(1/4) == 0 (mod p) (equivalently, p divides A001586(p-3)). Zhi-Wei Sun made many conjectures on congruences involving E_{p-3}(1/4), see the reference.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008960 | OEIS-A245206

Odd primes p with E_{p-3}(1/4) == 0 (mod p), where E_n(x) denotes the Euler polynomial of degree n.

The conjecture in A245204 asserts that the current sequence contains infinitely many primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008961 | OEIS-A245211

a(n) = Sum_{(d<n) | n} (d * tau(d)).

Conjecture: 21 is only number such that a(n) = n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jul 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008962 | OEIS-A245212

a(n) = n * tau(n) - Sum_{(d<n) | n} (d * tau(d)).

Conjecture: a(n) = sigma(n) iff n is a power of 2 (A000079).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jul 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008963 | OEIS-A245219

Continued fraction expansion of the constant c in A245218; c = sup{f(n,1)}, where f(1,x) = x + 1 and thereafter f(n,x) = x + 1 if n is in A001951, else f(n,x) = 1/x.

Theorem: Referring to Problem B6 in the 81st William Lowell Putnam Mathematical Competition (see link), in the notation of the first solution, the sequence {c_i} equals A245219. This proves the conjecture in the previous comment. - Manjul Bhargava, Kiran Kedlaya, and Lenny Ng, Sep 09 2021.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008964 | OEIS-A245278

Decimal expansion of k3, a Diophantine approximation constant such that the conjectured volume of the "critical parallelepiped" is 2^3*k3 (the 3-D analog of A242671).

Decimal expansion of k3, a Diophantine approximation constant such that the conjectured volume of the "critical parallelepiped" is 2^3*k3 (the 3-D analog of A242671).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Jul 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008965 | OEIS-A245279

Decimal expansion of a1, the first of two constants associated with Djokovic's conjecture on an integral inequality.

Decimal expansion of a1, the first of two constants associated with Djokovic's conjecture on an integral inequality.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Jul 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008966 | OEIS-A245280

Decimal expansion of a2, the second of two constants associated with Djokovic's conjecture on an integral inequality.

Decimal expansion of a2, the second of two constants associated with Djokovic's conjecture on an integral inequality.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Jul 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008967 | OEIS-A245305

Numbers k such that 4k+1, 4k+3, and 6k+5 are all primes.

Infinitude of the sequence would follow from Dickson's (unproved) conjecture. - _Jens Kruse Andersen_, Jul 18 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jul 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008968 | OEIS-A245322

E.g.f.: Sum_{n>=0} sin(n*x)^n.

It appears that for n >= 1, a(2*n) is even and a(2*n-1) is odd. Conjecture: Let p be prime. The sequence obtained by reducing a(n) modulo p for n >= 1 is purely periodic. If p = 4*m + 1 the period appears to be p - 1, while if p = 4*m + 3 the period appears to be 2*(p - 1). Cf. A224899 and A221078. - _Peter Bala_, May 31 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Nov 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008969 | OEIS-A245329

a(n) = sum_{k=0..n}C(n,k)^3*(-8)^k with C(n,k) = n!/(k!(n-k)!).

Conjecture: (i) For any prime p > 3, we have sum_{k=0}^{p-1}(-1)^k*a(k) == (p/3) - p^2/12*B_{p-2}(1/3) (mod p^3) and sum_{k=0}^{p-1}(-1)^k*a(k)*H(k,2) == B_{p-2}(1/3) (mod p), where B_m(x) denotes the Bernoulli polynomial of degree m and H(k,2) stands for sum_{j=1..k} 1/j^2. Also, sum_{k=1}^{p-1}(-1)^k*a(k)/k == - 3*(2^p-2)/p (mod p) for any prime p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008970 | OEIS-A245340

Smallest m such that A125717(m) = n, or -1 if n never appears.

Conjecture: a(n) is never -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jul 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008971 | OEIS-A245372

Prime-partitionable numbers a(n) for which there exists a 2-partition of the set of primes < a(n) that has a smallest subset containing four primes only.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christopher Hunt Gribble_, Nov 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008972 | OEIS-A245474

a(n) = smallest positive integer s such that s*n - floor(sqrt(s*n))^2 is a square.

Conjecture: a(p) < p for prime p == 1 (mod 4). Outline of proof of conjecture: write p = x^2 + y^2. Since gcd(x,y) = 1, there are u,v with x*u + y*v = 1, u^2 + v^2 < y^2 + x^2 = p. Taking s = u^2 + v^2, s*p = (u*y+v*x)^2 + 1^2, and |u*y+v*x| = floor(sqrt(s*p)). - _Robert Israel_, Aug 04 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Jul 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008973 | OEIS-A245525

Unique integer r with -prime(n)/2 < r <= prime(n)/2 such that p(n) == r (mod prime(n)), where p(.) is the partition function given by A000041.

Conjecture: a(n) is always nonzero, i.e., prime(n) never divides the partition number p(n). This conjecture does not hold with the smallest counterexample being n=1119414 (cf. A245662). - _Max Alekseyev_, Jul 27 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008974 | OEIS-A245526

Unique integer r with -prime(n)/2 < r <= prime(n)/2 such that L(2*n) == r (mod prime(n)), where L(k) denotes the Lucas number A000032(k).

Conjecture: a(n) is always nonzero, i.e., prime(n) never divides the Lucas number L(2*n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008975 | OEIS-A245569

Numbers n such that binomial(2^n-1,n) is squarefree.

Conjectured to be finite. Next term, if it exists, is larger than 500.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Nov 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008976 | OEIS-A245615

Sum_{lpf(j)=11, 11<j<=11*prime(n)}(-1)^(A000120(j)+1), where lpf=least prime factor.

Conjecture: All terms are nonnegative. A similar conjecture arises by changing 11 in the definition of the sequence by some other primes: 19,41,67,107,173,...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jul 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008977 | OEIS-A245621

Sequence of distinct least nonnegative numbers such that the average of the first n terms is a cube.

Conjecture: a(2*n+1) = n^3, a(2*n) = 7*n^3 - 9*n^2 + 5*n - 1. Conjecture: a(n) = 4*a(n-2)-6*a(n-4)+4*a(n-6)-a(n-8) for n>8. - _Colin Barker_, Nov 05 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Nov 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008978 | OEIS-A245626

a(n)= 1 (respectively, a(n)= 3) if up to 2^n the number of A245622-terms is more (respectively, less) than the number of A245623-terms; or a(n)=0 if these numbers are equal.

Since 10^2,10^3,... appear earlier than 3*10^2,3*10^3,..., then it is natural to conjecture, that, beginning with n=4, a(n)=0 or 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jul 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008979 | OEIS-A245637

Decimal expansion of Integral_{x = 1..infinity} 1/x^x dx.

Conjecturally, equals 1 - Integral_{x = 1..oo, y = 1..oo} 1/(x*y)^(x*y) dx dy. [added Dec 21 2022: follows from Glasser's Theorem 1.] (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Jul 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008980 | OEIS-A245658

G.f. A(x) satisfies: A(x)^3 = Sum_{n>=0} A247031(n)*x^n.

a(n) == 1 (mod 2) iff n = (4^k - 1)/3 for k>=0 (conjecture). a(n) == 2 (mod 3) iff A247031(3*n) == 2 for n>=0 (mod 3) (conjecture), where A247031 is the self-convolution cube of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008981 | OEIS-A245664

Prime-partitionable numbers a(n) for which there exists a 2-partition of the set of primes < a(n) that has one subset containing two primes only.

Conjecture: If P1 = {p1a, p1b} with p1a and p1b odd primes, p1a < p1b and p1b = 2*k*p1a + 1 for some natural k such that 2*k <= p1a - 3 and if m = p1a + p1b then m is prime-partitionable and belongs to {a(n)}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christopher Hunt Gribble_, Jul 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008982 | OEIS-A245688

Numbers n such that n^2 + 1 is the product of three distinct Fibonacci numbers > 1.

Conjecture: except the numbers 99 and 15075, all the terms belong to A005248 (bisection of Lucas numbers). The above conjecture holds for the first 1000 terms. - _Jens Kruse Andersen_, Aug 10 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jul 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008983 | OEIS-A245691

Irregular triangle of Collatz like iteration, x -> 3x, then repeat (x -> ceiling(x/2) if divisible by 3, otherwise x -> 3x) while x != 6.

It is conjectured that the number of steps for the trajectory to arrive at 6 is equal to the number of steps for the Collatz trajectory to arrive at 1 for the same starting value n (n>1), suggesting the length of the n-th row of the irregular array is given by A008908(n). Note that if the starting value of a trajectory in the Collatz sequence is not treated as a potential stopping value, then the conjecture would also be valid for n = 1. Starting with x the first step in this sequence is always to multiply by 3. Thereafter if x <> 6, divide by 2 (rounding up) if x mod 3 = 0, otherwise multiply by 3. If the initial multiply-by-3 step is omitted the sequence still arrives at 6 for any starting value (conjecturally), but the length of the trajectory would no longer be the same as the length of the Collatz trajectory for starting values (n>1) that are divisible by 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _K. Spage_, Aug 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008984 | OEIS-A245709

Fixed points of A245705 and A245706.

Contains also all such numbers k that A245608(k) = A245708(k), because that condition implies that A245607(A245708(k)) = k = A245707(A245608(k)). Conjecture: contains no numbers outside of that set, that is, for all n, A245608(a(n)) = A245708(a(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jul 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008985 | OEIS-A245760

Maximal multiplicative persistence of n in any base.

It has been conjectured that there is a maximum multiplicative persistence in a given base, but it is not known if this sequence is bounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sergio Pimentel_, Jul 31 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008986 | OEIS-A245769

a(n) = Sum_{k=0..n} C(n, k)*C(n+k, k)/(2k-1), where C(n, k) denotes the binomial coefficient n!/(k!*(n-k)!).

Conjecture: The sequence a(n+1)/a(n) (n = 3,4,...) is strictly increasing to the limit 3+2*sqrt(2), and the sequence a(n+1)^(1/(n+1))/a(n)^(1/n) (n = 5,6,...) is strictly decreasing to the limit 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008987 | OEIS-A245778

Numbers m such that m/tau(m) - sigma(m)/m is an integer.

Conjecture: subsequence of A216793. This conjecture does not hold as A009194(a(19)) < A009194(a(18)). - _Max Alekseyev_, Jul 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Aug 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008988 | OEIS-A245786

Numbers n such that k(n) = (n/tau(n) + sigma(n)/n) is an integer.

Conjecture: Subsequence of A216793.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Aug 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008989 | OEIS-A245800

a(n) is the least number k such that Sum_{j=S(n)+1..S(n)+k} 1/j >= 1/2, where S(n) = Sum_{i=1..n-1} a(i) and S(1) = 0.

Conjecture 1: If the summation threshold (which, for this sequence, is defined as 1/2) is set to any positive number R, then the limiting ratio of consecutive terms in the resulting sequence is e^R. Conjecture 2: If the summation threshold is set to log(phi) = 0.4812118250..., then the Fibonacci sequence is generated.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rob van den Tillaart_, Aug 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008990 | OEIS-A245852

Powers of 8 without the digit '0' in their decimal expansion.

Conjectured to be finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Aug 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008991 | OEIS-A245853

Powers of 12 without the digit '0' in their decimal expansion.

Conjectured to be finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Aug 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008992 | OEIS-A245871

Number of length 2+2 0..n arrays with some pair in every consecutive three terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008993 | OEIS-A245872

Number of length 3+2 0..n arrays with some pair in every consecutive three terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008994 | OEIS-A245873

Number of length 4+2 0..n arrays with some pair in every consecutive three terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008995 | OEIS-A245874

Number of length 5+2 0..n arrays with some pair in every consecutive three terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008996 | OEIS-A245875

Number of length 6+2 0..n arrays with some pair in every consecutive three terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 04 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008997 | OEIS-A245876

Number of length 7+2 0..n arrays with some pair in every consecutive three terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008998 | OEIS-A245926

Expansion of g.f. sqrt((1-x + sqrt(1-14*x+x^2)) / (2*(1-14*x+x^2))).

Conjecture: a(n) = [x^n] ((1 + x + x^2)*(1 + x)^2/(1 - x)^2)^n. Equivalently, a(n) = Sum_{k = 0..n} Sum_{i = 0..k} Sum_{j = 0..n-k-i} C(n, k)*C(k, i)*C(2*n, j)*C(3*n-k-i-j-1, n-k-i-j). If the conjecture is true then the Gauss congruences a(n*p^k) == a(n*p^(k-1)) (mod p^k) hold for all primes p and positive integers n and k. Calculation suggests that the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(2*k)) hold for primes p >= 5 and positive integers n and k. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-008999 | OEIS-A245942

Irregular triangle read by rows of a variation of the Collatz iteration with signature (2,3).

It is conjectured that the trajectory of this Collatz-like iteration arrives at 16 in a finite number of steps for any initial value x, (x>4). The iterative step is divide by 2 and add 2 if even, or multiply by 3 and subtract 5 if odd. For any odd initial value (x>6) the number of steps in the trajectory is the same as the number of steps in the Collatz trajectory starting with floor(x-3)/2. Two variations belong to the same subset if their (a) parameters are the same and their (b) parameters have the same parity. It is conjectured that any variations belonging to the same subset have equal row lengths. Members of the same subset share other properties. For example, if a trajectory of subset type (a=2, b:odd) starts with an odd number, then the function floor((x-b)/2) maps the element values (x) of this trajectory to a trajectory of A245691.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _K. Spage_, Aug 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009000 | OEIS-A245943

Irregular triangle read by rows of a variation of the Collatz iteration with signature (1,2).

It is conjectured that the trajectory of this Collatz-like iteration arrives at 3 in a finite number of steps for any initial value x, (x>2). The iterative step is divide by 2 and add 1 if even, or multiply by 3 and subtract 3 if odd. For any initial value the number of steps in the trajectory is the same as the number of steps in the Collatz trajectory starting with (x-2). Two variations belong to the same subset if their (a) parameters are the same and their (b) parameters have the same parity. It is conjectured that any variations belonging to the same subset have equal row lengths. Members of the same subset share other properties. For example the trajectory of any variation of subset type (a=1, b:even) can be mapped to a Collatz trajectory by b from each element of the trajectory.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _K. Spage_, Aug 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009001 | OEIS-A245944

Irregular triangle read by rows of a variation of the Collatz iteration with signature (2,4).

It is conjectured that the trajectory of this Collatz-like iteration arrives at 6 in a finite number of steps for any initial value x, (x>4). The iterative step is divide by 2 and add 2 if even, or multiply by 3 and subtract 7 if odd. For any even initial value the number of steps in the trajectory is the same as the number of steps in the Collatz trajectory starting with floor(x-4)/2. Two variations belong to the same subset if their (a) parameters are the same and their (b) parameters have the same parity. It is conjectured that any variations belonging to the same subset have equal row lengths.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _K. Spage_, Aug 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009002 | OEIS-A245951

Number of length 1+3 0..n arrays with some pair in every consecutive four terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009003 | OEIS-A245952

Number of length 2+3 0..n arrays with some pair in every consecutive four terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009004 | OEIS-A245953

Number of length 3+3 0..n arrays with some pair in every consecutive four terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009005 | OEIS-A245954

Number of length 4+3 0..n arrays with some pair in every consecutive four terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009006 | OEIS-A245990

Number of length n+2 0..3 arrays with no pair in any consecutive three terms totalling exactly 3.

Conjectures from _Colin Barker_, Nov 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009007 | OEIS-A246048

Numbers for which (n^2)! is divisible by n!^n*(2n)!.

Farideh Firoozbakht observes that all terms are composite numbers. The comment in A057599 and conjecture in A096126 seem to confirm that there are no primes nor powers of primes in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Aug 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009008 | OEIS-A246054

Numerators of the first rational corners of the mean-median map limit value m(x) for x >= 1/2.

The function m(x) is the limit value of the mean-median map. Start with 0<x<1 and find a new number x_4 so that the median of {0,x,1} equals the mean of {0,x,1,x_4}. Then find a new number x_5 so that the median of {0,x,1,x_4} equals the mean of {0,x,1,x_4,x_5}, etc. It is conjectured by Shultz and Shiflett that for all real x this process ends after finitely many steps. Cellarosi and Munday find a range of x for which this conjecture is true (improving results of Chamberland and Martelli), and an explicit piecewise affine formula for m(x). It is enough to study the case of 1/2 <= x <= 2/3. The first rational corners of the map m(x) to the right of 1/2 are found by Cellarosi and Munday and these are the rationals whose numerators are listed above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Francesco Cellarosi_, Aug 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009009 | OEIS-A246055

Denominators of the first rational corners of the function m(x) for x>=1/2.

The function m(x) is the limit value of the mean-median map. Start with 0<x<1 and find a new number x_4 so that the median of {0,x,1} equals the mean of {0,x,1,x_4}. Then find a new number x_5 so that the median of {0,x,1,x_4} equals the mean of {0,x,1,x_4,x_5}, etc. It is conjectured by Shultz and Shiflett that for all real x this process ends after finitely many steps. A range of x for which this conjecture is true found by Cellarosi and Munday (improving the results of Chamberland and Martelli). An explicit piecewise affine formula for m(x) is found. It is enough to study the case of

1/2<=x<=2/3. The first rational corners of the map m(x) to the right of 1/2 are found by Cellarosi and Munday and these are the rationals whose denominators are listed above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Francesco Cellarosi_, Oct 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009010 | OEIS-A246056

G.f.: Sum_{n>=0} x^n / (1-2*x)^(2*n+1) * [Sum_{k=0..n} C(n,k)^2 * 2^k * x^k] * [Sum_{k=0..n} C(n,k)^2 * 3^k * x^k].

Conjecture: a(n) == 1 (mod 3) when n = 2*A005836(k) for k >= 0, and a(n) == 0 (mod 3) otherwise, where A005836 gives numbers whose base-3 representation contains no 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009011 | OEIS-A246065

a(n) = Sum_{k=0..n}C(n,k)^2*C(2k,k)/(2k-1), where C(n,k) denotes the binomial coefficient n!/(k!*(n-k)!).

Conjecture: (i) The sequence a(n+1)/a(n) (n = 2,3,...) is strictly increasing to the limit 9, and the sequence a(n+1)^(1/(n+1))/a(n)^(1/n) (n = 1,2,3,...) is strictly decreasing to the limit 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009012 | OEIS-A246133

a(n) = (binomial(2n, n) - 2) mod n^3.

When e=3, the numbers binomial(2n, n) - 2 mod n^e are 0 whenever n is a prime greater than 3 (Wolstenholme's theorem; see A246130 for introductory comments). No composite number n for which a(n)=0 was found up to n=431500 (conjecture: there are none, and a(n)=0 for n>3 is a deterministic primality test).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Aug 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009013 | OEIS-A246138

a(n) = (Sum_{k=0..n-1} A246065(k)) / n^2.

Part (ii) of the conjecture in A246065 implies that all the terms in the current sequence are integers. Conjecture: The sequence a(n+1)/a(n) (n = 4,5,...) is strictly increasing to the limit 9, and the sequence a(n+1)^(1/(n+1))/a(n)^(1/n) (n = 3,4,...) is strictly decreasing to the limit 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009014 | OEIS-A246317

A246316(2^n-1).

Conjecture: Rules 332, 356, 364, 452, 460, 484, 492, 836, 844, 868, 876, 964, 972, 996 and 1004 also generate this sequence. - _Lars Blomberg_, Jun 14 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009015 | OEIS-A246396

Nonnegative integers k satisfying cos(k) <= 0 and cos(k+1) <= 0.

Conjecture: every term t has at least one neighbor which is equal to t plus or minus one. - _Harvey P. Dale_, Jul 11 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009016 | OEIS-A246402

a(n) = smallest number k such that 2*(k*n)^n-1 is prime, or 0 if no such k exists.

I conjecture that k exists for every n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Aug 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009017 | OEIS-A246423

G.f.: Sum_{n>=0} x^n / (1-3*x)^(2*n+1) * [Sum_{k=0..n} C(n,k)^2 * 2^k * x^k] * [Sum_{k=0..n} C(n,k)^2 * 3^k * x^k].

a(n) == 2 (mod 4) iff n = 2^k for k>=2, and a(n) == 0 (mod 4) elsewhere except at a(0)=1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009018 | OEIS-A246437

Expansion of (1/2)*(1/(x+1)+1/(sqrt(-3*x^2-2*x+1))).

Conjecture: the stronger supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(2*k)) hold for primes p >= 5 and positive integers n and k. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Nov 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009019 | OEIS-A246452

Let A be the set of all numbers m for which m^2+1 is prime (see A005574). The sequence lists the pairs of numbers (a, b) such that every m > 1 included in A can be written in the form m = a + b for a <= b both in A and a minimum.

Conjecture: for n>1, A005574(n) = a+b where a and b are integers in A005574. By definition, a(2n-1) + a(2n) = A005574(n+1), and a(2n-1) is the minimal solution where the pair (a(2n-1), a(2n)) are both in A005574. The above conjecture says such a pair always exists. - _Jens Kruse Andersen_, Aug 27 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Aug 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009020 | OEIS-A246460

a(n) = (sum_{k=0}^{n-1} (2k+1)*C(n-1,k)^2*C(n+k,k)^2)/n^2, where C(n,k) denotes the binomial coefficient n!/(k!(n-k)!).

Conjecture: a(n) is always an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009021 | OEIS-A246471

First of the least n primes that are also consecutive terms of A005117 (squarefree numbers).

If the k-tuple conjecture is true then a(n) always exists. Weak upper bounds are

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hans Havermann_, Aug 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009022 | OEIS-A246480

Number of length 1+3 0..n arrays with no pair in any consecutive four terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009023 | OEIS-A246481

Number of length 2+3 0..n arrays with no pair in any consecutive four terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009024 | OEIS-A246482

Number of length 3+3 0..n arrays with no pair in any consecutive four terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009025 | OEIS-A246488

Gridgeman pairs in increasing order: pairs of palindromic primes which differ only in their middle digits whose difference is equal to 1.

It has been conjectured by Norman T. Gridgeman that infinitely many pairs of such primes exist (see second ref.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_, Aug 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009026 | OEIS-A246501

a(n) is the smallest term of A242719 that is repeated exactly n times, or 0 if there is no such term.

We conjecture that, for all n, a(n)>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Aug 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009027 | OEIS-A246502

a(n) is the smallest term of A242720 that is repeated exactly n times, or 0 if there is no such term.

We conjecture that a(n)>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Aug 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009028 | OEIS-A246512

a(n) = (sum_{k=0}^{n-1}(3k^2+3k+1)*C(n-1,k)^2*C(n+k,k)^2)/n^3, where C(n,k) denotes the binomial coefficient n!/(k!*(n-k)!).

Conjecture: The sequence a(n+1)/a(n) (n = 1,2,3,...) is strictly increasing to the limit 17+12*sqrt(2), and the sequence a(n+1)^(1/(n+1))/a(n)^(1/n) (n = 1,2,3,...) is strictly decreasing to the limit 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009029 | OEIS-A246513

a(n) = (4/n^2)*(Sum_{k=0..n-1} k*A246459(k)).

Conjecture: a(n) is always an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009030 | OEIS-A246535

Largest number with at most n distinct digits in any base b >= 2 (written in decimal).

Generalizing... (Conjecture) a(6) = 2610787117 written in base 7 is 121461216151 (5 distinct digits), and 2610787118 written in base 6^2+1 = 37 is (1)(0)(24)(1)(22)(2)(0) (5 distinct digits). Therefore, Derek Orr's conjecture seems to be wrong.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joonas Pohjonen_, Aug 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009031 | OEIS-A246541

Take the squares of all P_(n+2)-rough numbers less than the (n+1)-th primorial and mod each by the (n+1)-th primorial. There will be a(n) different results.

The conjectured formula from Jon E. Schoenfield is true. This can be seen by considering that each odd prime p has exactly (p+1)/2 quadratic residues (mod p), of which (p-1)/2 are nonzero. The P_(n+2)-rough numbers less than the (n+1)-th primorial comprise all combinations of nonzero residues modulo the first n+1 primes. So for each odd prime p, the p-1 nonzero residues map to (p-1)/2 (nonzero) residues after squaring. - _Bert Dobbelaere_, Aug 09 2023 Conjecture: a(n) = (1/2^n)*Product_{j=1..n} (prime(j+1)-1) = A005867(n+1)/2^n. - _Jon E. Schoenfield_, Feb 20 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John B. Yin_, Aug 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009032 | OEIS-A246542

a(n) = 2/n^2*(sum_{k=0..n-1} (2*k+1)*C(n-1,k)^2*C(n+k,k)), where C(m,k) denotes the binomial coefficient m!/(k!*(m-k)!).

The following general conjecture implies that a(n) is always an integer. Conjecture: Let a and b be nonnegative integers, and let n be a positive integer. If a == b (mod 2), then sum_{k=0}^{n-1}(2k+1)*C(n-1,k)^a*C(-n-1,k)^b == 0 (mod n^2); if a - b is odd, then 2*sum_{k=0}^{n-1}(-1)^k*(2k+1)C(n-1,k)^a*C(-n-1,k)^b == 0 (mod n^2). An extension of the conjecture was proved in the latest version of arXiv:1408.5381. - _Zhi-Wei Sun_, Sep 14 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009033 | OEIS-A246543

a(n) = (2/n^3)*(Sum_{k=0..n-1} (-1)^k*(3*k^2+3*k+1)*binomial(n-1,k)^3*binomial(n+k,k)^3).

Conjecture: Let n be any positive integer. For m = 0, 2, 4, ..., we have Sum_{k=0..n-1} (3k^2+3k+1)*(binomial(n-1,k)*binomial(n+k,k))^m == 0 (mod n^3); for m = 1, 3, 5, ... we have 2*Sum_{k=0..n-1} (-1)^k*(3k^2+3k+1)*(binomial(n-1,k)*binomial(n+k,k))^m == 0 (mod n^3). The author proved the conjecture in the latest version of arXiv:1408.5381. - _Zhi-Wei Sun_, Sep 14 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009034 | OEIS-A246567

a(n) = (sum_{k=0}^{n-1}C(n-1,k)^2*C(-n-1,k)^2/(4*k^2-1))/n, where C(x,k) refers to binomial(x,k).

The following conjecture implies that a(n) is an integer. Conjecture: (i) For any positive integers m and n, the sum sum_{k=0}^{n-1}C(n-1,k)^m*C(-n-1,k)^m/(4k^2-1) is always an integer divisible by n. In the latest version of arXiv:1408.5381, the author proved part (i) of the conjecture, thus a(n) is indeed integral. - _Zhi-Wei Sun_, Sep 04 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009035 | OEIS-A246600

Number of divisors d of n with property that the binary representation of d can be obtained from the binary representation of n by changing any number of 1's to 0's.

The sums of the first 10^k terms for k = 1, 2, ..., are 16, 224, 2580, 26920, 273407, 2745100, 27440305, 274127749, 2738936912, 27373288534, 273631055291, 2735755647065, ... Conjecture: The asymptotic mean of this sequence is 1 + Sum_{k>=1} 1/(k*2^A000120(k)) = 2.7351180693... . - _Amiram Eldar_, Apr 07 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009036 | OEIS-A246606

Central terms of the triangle A116853.

Conjecture: D-finite with recurrence +(-2*n+3)*a(n) +4*(n-1)*(2*n^2-4*n+1)*a(n-1) +(n-1)*(n-2)*(2*n-1)*a(n-2)=0. - _R. J. Mathar_, Jul 01 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Aug 31 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009037 | OEIS-A246655

Prime powers: numbers of the form p^k where p is a prime and k >= 1.

Blagojević & Ziegler prove that for these n and for any convex polygon in the plane, the polygon can be partitioned into n polygons with equal area and equal perimeter. The result is conjectured (by Nandakumar & Rao, who proved the case n = 2) to hold for all n. - _Charles R Greathouse IV_, Apr 24 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_ and _Franklin T. Adams-Watters_, Sep 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009038 | OEIS-A246695

Row sums of the triangular array A246694.

Conjectured linear recurrence: a(n) = 2*a(n-1) + a(n-2) - 4*a(n-3) + a(n-4) + 2*a(n-5) - a(n-6), with a(0) = 1, a(1) = 3, a(2) = 9, a(3) = 18, a(4) = 35, a(5) = 57, a(6) = 91. Conjectured g.f.: (1 + x + 2*x^2 + x^3 + x^4)/((x - 1)^4*(x + 1)^2). Conjecture: a(n) = (1/8)*(n + 1)*((-1)^n + 2*n^2 + 4*n + 7). - _Eric Simon Jacob_, Jul 19 2023 [This conjecture is correct; compare A377802, note offset 1. - _Werner Schulte_, Nov 22 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009039 | OEIS-A246697

Row sums of the triangular array A246696.

Conjectured linear recurrence: a(n) = 2*a(n-1) + a(n-2) - 4*a(n-3) + a(n-4) + 2*a(n-5) - a(n-6), with a(0) = 1, a(1) = 5, a(2) = 16, a(3) = 34, a(4) = 67, a(5) = 111, a(6) = 178. Conjectured g.f.: (1 + 3*x + 5*x^2 + x^3 + 2*x^4)/((x - 1)^4*(x + 1)^2). Conjecture: a(n) = (2*n^3+6*n^2+9*n+4+n*(-1)^n)/4. - _Luce ETIENNE_, Oct 16 2016 Conjectured e.g.f.: ((2 + 8*x + 6*x^2 + x^3)*cosh(x) + (2 + 9*x + 6*x^2 + x^3)*sinh(x))/2. - _Stefano Spezia_, May 10 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009040 | OEIS-A246699

Squarefree n such that C(2^n - 1, n) is also squarefree, where C is the binomial coefficient.

Conjectured to be finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Nov 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009041 | OEIS-A246702

The number of positive k < (2n-1)^2 such that (2^k - 1)/(2n - 1)^2 is an integer.

Conjecture: the positions of 1's, a(k)=1, are exactly given by the 2k-1 which are elements of A167791. - _Antti Karttunen_, Nov 15 2014 Conjecture: Let B be the first value such that (2k-1)^2 divides 2^B - 1. Then either 2k-1 divides B, or 2k-1 is a Wieferich prime (A001220). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Nov 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009042 | OEIS-A246738

Number of length 1+4 0..n arrays with no pair in any consecutive five terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009043 | OEIS-A246739

Number of length 2+4 0..n arrays with no pair in any consecutive five terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009044 | OEIS-A246748

Numbers n such that A242719(n) = (prime(n))^2+1 and A242720(n) - A242719(n) = 2*(prime(n)+1).

In connection with this, one can conjecture that A242719(k) ~ A242720(k) ~ (prime(k))^2, as k goes to infinity (cf. A246819, A246821).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009045 | OEIS-A246779

Strictly increasing terms of the sequence A246776: a(1)= A246776(1) and for n>0 a(n+1) is the next term greater than a(n) after that a(n) appears in A246776 for the first time.

I conjecture that, a(n)=n+17 for all n, n>22.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Sep 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009046 | OEIS-A246780

Strictly increasing terms of the sequence A246778: a(1)= A246778(1) and for n>0 a(n+1) is next term greater than a(n) after that a(n) appears in A246778 for the first time.

I conjecture that, a(n)=n+5 for all n, n>15.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Sep 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009047 | OEIS-A246783

Smallest number m such that all the n numbers np(m+k-1), 0 < k < n+1 are equal, where np(t) is number of primes p with prime(t) < p < prime(t)^(1+1/t).

Conjecture: For every n, a(n) exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_, Oct 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009048 | OEIS-A246786

Record values of A182134.

Also: indices n for which A246810(k) > A246810(n) for all k > n. The missing numbers are {9, 16, 20, 22, 24, 31, 34, ...}, I conjecture that these are exactly the indices for which A246810(n+1) < A246810(n). - _M. F. Hasler_, Oct 16 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Farideh Firoozbakht_ and _Jahangeer Kholdi_, Oct 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009049 | OEIS-A246826

Numbers n such that there is no prime of a prime twin pair between n^2 + n and n^2 + 3*n + 2.

Conjecture: the sequence is finite and given in full.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Sep 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009050 | OEIS-A246844

Decimal expansion of t_0, the lower bound of the conjectured first interval [t_0, t_1] where the real part of zeta(1+i*t) is negative.

Decimal expansion of t_0, the lower bound of the conjectured first interval [t_0, t_1] where the real part of zeta(1+i*t) is negative.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Sep 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009051 | OEIS-A246845

Decimal expansion of t_1, the upper bound of the conjectured first interval [t_0, t_1] where the real part of zeta(1+i*t) is negative.

Decimal expansion of t_1, the upper bound of the conjectured first interval [t_0, t_1] where the real part of zeta(1+i*t) is negative.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Sep 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009052 | OEIS-A246860

Expected value of trace(O)^(2n), where O is a 4 X 4 orthogonal matrix randomly selected according to Haar measure.

In the MathOverflow link, Nathaniel Johnston conjectures a(n) = A000108(n)*(A000108(n)+1)/2. - _Robert Israel_, Jan 17 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nathaniel Johnston_, Sep 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009053 | OEIS-A246893

Number of length 1+4 0..n arrays with some pair in every consecutive five terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009054 | OEIS-A246894

Number of length 2+4 0..n arrays with some pair in every consecutive five terms totalling exactly n.

Conjectures from _Colin Barker_, Nov 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009055 | OEIS-A246907

Numbers n such that sigma(n + sigma(n)) = 3n.

Conjecture: for n >= 2; numbers n of the form 2^k such that 3*(2^k) - 1 is prime. The next terms: 18446744073709551616, 75557863725914323419136, 19807040628566084398385987584, ... Sequence of numbers k: 1, 2, 3, 4, 6, 7, 11, 18, 34, 38, 43, 55, 64, 76, ... Subsequence of A087370 (numbers n such that 3n - 1 is a prime).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Sep 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009056 | OEIS-A246914

Primes p such that sigma(2p+1) = 3*(p+1).

Conjecture: Also primes p such that sigma(2p+1) mod p = 3. - _Jaroslav Krizek_, Sep 28 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Sep 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009057 | OEIS-A246915

Numbers k such that sigma(k + sigma(k)) = sigma((k+1) + sigma(k+1)).

Conjecture: sequence of numbers A246456(a(n)): 12, 24, 48, 168, 2160, 17280, 54720, 77280, 221184, 202176, 185328, 249984, 312480, 599040, 725760, 967680, 864864, 1327104, 1489488, 2048256, 1958400, 2439360, 3110400, 3902976, 4852224, 4713984, ... is sequence of any multiples of 12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Sep 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009058 | OEIS-A246919

The length of the longest nontrivial integral cevian of an equilateral triangle of side n that divides an edge into two integral parts, or 0 if no such cevian exists.

Conjectures from _Colin Barker_, Jun 06 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Barker_, Sep 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009059 | OEIS-A247000

Maximal number of palindromes in a circular binary word of length n.

Conjectures from _Colin Barker_, Nov 14 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009060 | OEIS-A247010

Primes p such that (p-3)/2 and 2*p+3 are both prime.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Sep 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009061 | OEIS-A247031

G.f.: 1 = Sum_{n>=0} a(n)*x^n * [Sum_{k=0..n+1} binomial(n+1, k)^3 * (-x)^k]^3.

a(3*n) == 2 (mod 3) iff A245658(n) == 2 (mod 3), where A245658 is the self-convolution cube root of this sequence (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009062 | OEIS-A247047

Numbers k such that k^2 contains exactly 2 distinct digits and k^3 contains exactly 3 distinct digits.

The numbers k such that k^2 contains 2 distinct digits, k^3 contains 3 distinct digits, and k^4 contains 4 distinct digits are conjectured to only be 6, 8, and 15. (Intersection of A016069, A155146, and A155150.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Sep 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009063 | OEIS-A247077

Composite numbers for which the harmonic mean of proper divisors is an integer.

Conjecture: all terms are of the form m*(sigma(m)-1) where sigma(m)-1 is prime. - _Chai Wah Wu_, Dec 15 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Lignon_, Nov 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009064 | OEIS-A247090

Eric Rowland's generalization of A132199 as a rectangular array A read by upward antidiagonals.

Conjecture [Rowland] (paraphrased): Let A be the above array with entry A(n,k) in row n and column k. For each n, there exists an index N(n) >= 1 such that A(n,j) is either 1 or prime for all j > N(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Nov 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009065 | OEIS-A247102

Expansion of g.f. (6*x+2)/(sqrt(-3*x^2-6*x+1)*(4*x^2+4*x)) - (2*x+1)/(2*x^2+2*x).

Conjecture D-finite with recurrence: (n+1)*a(n) +(-2*n-5)*a(n-1) +3*(-8*n+7)*a(n-2) +15*(-2*n+3)*a(n-3) +9*(-n+2)*a(n-4)=0. - _R. J. Mathar_, Jan 25 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Nov 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009066 | OEIS-A247112

Floor of sums of the cubes of the non-integer square roots of n, as partitioned by the integer roots: floor(Sum_{j=n^2+1..(n+1)^2-1} j^(3/2)).

Conjecture: the corresponding sums for all fractional (rational) powers of n (e.g., 5/2, 7/2, 9/2, ..., 1/3, 2/3, 4/3, ..., 1/4, 3/4, 5/4, ..., 1/5, 2/5, 3/5, ...) will have polynomial integer formulas or recursive integer formulas for their floor, ceiling and/or rounded values, with convergence to a rational fractional portion, with possibly multiple fractional values in a repeating pattern as they converge. This was clear for some additional examples, less so for higher-order examples.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard R. Forberg_, Dec 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009067 | OEIS-A247170

Expansion of (-3/2+(x^3+3*x)/(sqrt(x^4-4*x^3-2*x^2+1)*2*x)).

Conjectured: +n*(2*n-7)*a(n) +(n-1)*(2*n-9)*a(n-1) +2*(-2*n^2+9*n-6)*a(n-2) +2*(-6*n^2+33*n-38)*a(n-3) +3*(-2*n^2+15*n-26)*a(n-4) +(2*n-5)*(n-5)*a(n-5) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Nov 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009068 | OEIS-A247173

Expansion of g.f. (-1)/(2*x^2+2*x) +(-2*x^3-3*x^2+1) / (sqrt(x^4+4*x^3-2*x^2-4*x+1)*(2*x^2+2*x)).

Conjecture D-finite with recurrence: -(n+1)*(3*n-4)*a(n) +(9*n^2-3*n-4)*a(n-1) +2*(9*n^2-21*n+8)*a(n-2) +2*(-3*n^2+n+8)*a(n-3) +(-15*n^2+53*n-12)*a(n-4) -(3*n-1)*(n-4)*a(n-5)=0. - _R. J. Mathar_, Jan 25 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Nov 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009069 | OEIS-A247190

Calculate the smallest positive residues of 2!,3!,...,(prime(n)-1)! modulo prime(n) and stop the calculation at the moment m when for the first time there appear two equal numbers. If m!==k!, then a(n)=k; but a(n)=0 if no such m exists.

There is a conjecture that answer to the Erdős question is no, which is equivalent to having a(n)>0 for n>=4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Nov 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009070 | OEIS-A247202

Smallest odd k > 1 such that k*2^n - 1 is a prime number.

Limit_{N->oo} (Sum_{n=1..N} a(n))/(Sum_{n=1..N} n) = log(2). [[Is there a proof or is this a conjecture? - _Peter Luschny_, Feb 06 2015]]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Nov 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009071 | OEIS-A247225

a(n) = n if n <= 3, a(4)=5, otherwise the smallest number not occurring earlier having at least one common factor with a(n-3), but none with a(n-1)*a(n-2).

Conjecturally the sequence is a permutation of the positive integers. However, to prove this we need more subtle arguments than were used to prove the corresponding property for A098550. - _Vladimir Shevelev_, Jan 14 2015 For n <= 2000, a(3n-1) is even and both a(3n) and a(3n-2) are odd numbers. I conjecture that this is true for all positive integers n. This conjecture is true iff for all positive integers n, a(3n-1) is even. - _Farideh Firoozbakht_, Jan 14 2015 Conjecturally for every k>=2, as in the case k=1, the sequence (*) is a permutation of the positive integers. For k>=3, at first glance, already the appearance of the number 6 seems problematic. However, at the author's request, _Peter J. C. Moses_ found that the positions of 6 are 83, 157, 1190, 206, ... in cases k=3,4,5,6,... respectively (A254003).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009072 | OEIS-A247233

Smallest m such that A075323(m) = n-th odd prime, or zero, if no such m exists.

Conjecture: a(388) = 0, i.e., A065091(388) = 2683 doesn't occur in A075323;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Nov 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009073 | OEIS-A247239

Array a(n,m) = ((n+2)/2)^m*Sum_{k=1..n+1} 1/sin(k*Pi/(n+2))^(2m), n>=0, k>=0, read by ascending antidiagonals.

Unexpectedly, it is conjectured (proof wanted) that the expression ((n+2)/2)^m * Sum_{k=1..n+1} 1/sin(k*Pi/(n+2))^(2m), n>=0, k>=0, always gives an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009074 | OEIS-A247250

Indices of Pell numbers having exactly one primitive prime factor.

Conjecture: The n-th Pell number A000129(n) has a primitive prime factor for all n > 1. (The n-th Fibonacci number A000045(n) has a primitive prime factor for all n except n = 0, 1, 2, 6, and 12.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, Nov 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009075 | OEIS-A247253

First differences of A251239.

Conjecture 1: a(n) > 0, since presumably primes occur in A098550 in natural order; Conjecture 2: it seems that a(n) = 1 only for n = 1, 5, 10 and 13; Conjecture 3: a(n) - 1 = number of composite terms between prime(n) and prime(n+1) in A098550; Conjecture 4: a(n) = A251417(n+5) for n>7. (The first four conjectures are due to _Reinhard Zumkeller_.) Conjecture 5: Apart from first term, this is equal to the sequence of run lengths in A251549. These run lengths begin 2, 6, 6, 7, 1, 7, 13, 8, 10, 1, 17, 8, 1, 13, 13, 13, 5, 10, 11, 5, 9, ... - _N. J. A. Sloane_, Dec 18 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Dec 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009076 | OEIS-A247278

Least integer k > 0 such that k*n - prime(k) is a square.

Conjecture: a(n) exists for any n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009077 | OEIS-A247283

Positions of subrecords in A048673.

Conjectures from _Chai Wah Wu_, Jul 30 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009078 | OEIS-A247284

Subrecords in A048673: maximum value between two consecutive records in A048673.

Conjectures from _Chai Wah Wu_, Jul 30 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009079 | OEIS-A247308

Layer counting sequence in the order-5 cubic honeycomb.

a(d+17) = 3*a(d+16) + 2*a(d+15) + 7*a(d+14) + a(d+13) - 5*a(d+12) + 3*a(d+11) - 2*a(d+10) - 18*a(d+9) + 18*a(d+8) + 2*a(d+7) - 3*a(d+6) + 5*a(d+5) - a(d+4) - 7*a(d+3) - 2*a(d+2) - 3*a(d+1) + a(d) (conjectured, found experimentally and tested from 19 to 135). - _Eryk Kopczynski_, Jul 04 2020 Conjectured G.f.: (1+x) * (1+2*x+8*x^2+9*x^3+8*x^4+17*x^5+10*x^6+10*x^8+10*x^10+17*x^11+8*x^12+9*x^13+8*x^14+2*x^15+x^16) / ((1-x)^2 * (1-2*x-4*x^2-11*x^3-12*x^4-7*x^5-10*x^6-8*x^7+10*x^8-8*x^9-10*x^10-7*x^11-12*x^12-11*x^13-4*x^14-2*x^15+x^16)). - _Natalia L. Skirrow_, Apr 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tim Hutton_, Sep 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009080 | OEIS-A247335

The curvature of touching circles inscribed in a special way in the larger segment of circle of radius 10/9 divided by a chord of length 4/3.

Refer to comment of A092387. Consider a circle C of radius 10/9 (in some length units) with a chord of length 4/3. This has been chosen such that the larger sagitta has length 2. The smaller sagitta has length 2/9. The input, besides the circle C is the circle C_0 with radius R_0 = 1, touching the chord and circle C. The following sequence of circles C_n with radii R_n, n >= 1, is obtained from the conditions that C_n touches (i) the circle C, (ii) the chord and (iii) the circle C_(n-1). The curvature of the n-th circle, C_n = 1/R_n, n >= 0, is conjectured to be a(n). If one considers the curvature of touching circles inscribed in the smaller segment, the sequence would be A247512. See an illustration given in the link. Conjectures from _Colin Barker_, Sep 18 2014: (Start)

The G.f. conjectured by Colin Barker above follows from the one for Chebyshev's T(n, 19) = A078986(n): (1/(1-x) + (1-19*x)/(1-38*x+x^2))/2 = (1-29*x+10*x^2)/((1-x)*(1-38*x+x^2)). The four term recurrence conjectured by Colin Barker above follows from the expanded g.f. denominator: (1-x)* (1-38*x+x^2) = 1- 39*x + 39*x^2 - x^3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Sep 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009081 | OEIS-A247339

a(n) is the least number k such that the greatest prime divisor of k^2+1 is the smallest prime divisor of n^2+1.

Conjecture: for all integer n, there exists at least an integer m <= n such that the smallest prime factor of n^2+1 is also the greatest prime factor of m^2+1. - _Michel Lagneau_, Sep 27 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Sep 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009082 | OEIS-A247341

Let b_k=3...3 consist of k>=1 3's. Then a(n) is the smallest k such that the concatenation 2^n b_k is prime, or a(n)=0 if there is no such prime.

Conjecture: for all n, a(n)>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009083 | OEIS-A247342

Let b_k=3...3 consist of k>=1 3's. Then a(n) is the smallest k such that the odd part (A000265) of concatenation b_k 2^n is prime, or a(n)=0 if there is no such prime.

Conjecture: for all n, a(n)>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009084 | OEIS-A247374

Number of button presses required to try every combination of a binary combination lock with n number buttons.

Conjectured to be D-finite with recurrence: n*a(n) +2*(-2*n+1)*a(n-1) +(n-2)*a(n-2) +2*(7*n-10)*a(n-3) +4*(-5*n+11)*a(n-4) +8*(n-3)*a(n-5)=0. - _R. J. Mathar_, Nov 19 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Elliott Line_, Sep 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009085 | OEIS-A247471

Number of paths in B(n) that start with a u step and end with a d step.

Conjecture D-finite with recurrence (n+3)*a(n) +3*(-n-1)*a(n-1) +(n-9)*a(n-3) +2*(2*n-9)*a(n-4) +(n-9)*a(n-5) +(-n+9)*a(n-6)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Sep 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009086 | OEIS-A247477

Largest m such that n/m + n divides (n/m)^(n/m) + n, (n/m)^n + n/m and n^(n/m) + n/m, or 0 if no such m exists.

Conjecture: smallest k such that k/m = n and k/m + k divides (k/m)^(k/m) + k, (k/m)^k + k/m, k^(k/m) + k/m, or 0 if no such k exists: 2, 1, 10, 0, 36, 0, 78, 0, 136, 0, 210, 0, 312, 0, 406, 0, ... Conjecture: Conjecture: the superdivisor constant is equal to 1/2 + sum_{n >= 1} 1/(4*A000217(2n)) - Sum_{n >= 1} 1/b(n) - Sum_{n >= 1} 1/c(n) - Sum_{n >= 1} 1/d(n), ... = 0.64.., where b(n) = numbers with 2 superdivisors {or 406, 430, 646, 666, 826, 1090, 1236, 1246, 1378, 1596, 1666, 1750, 2002, 2028, 2346, 2410, 2506, 2782, 2796, 2850, ...), c(n) = numbers with 3 superdivisors {or 2926, ...), d(n) = numbers with 4 superdivisors, ... - _Juri-Stepan Gerasimov_, Feb 18 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jan 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009087 | OEIS-A247489

Square array read by antidiagonals: A(k, n) = hypergeometric(P, Q, -k^k/(k-1)^(k-1)) rounded to the nearest integer, P = [(j-n)/k, j=0..k-1] and Q = [(j-n)/(k-1), j=0..k-2], k>=1, n>=0.

Conjecture: hypergeometric(P, Q, -k^k/(k-1)^(k-1)) = sum_{j=0.. floor(n/k)} binomial(n-(k-1)*j, j) for n>=(k-1)^2, P and Q as above. (This means for n>=(k-1)^2 the representation is exact without rounding.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Sep 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009088 | OEIS-A247556

Exact differential base (a B_2 sequence) constructed as follows: Start with a(0)=0. For n>=1, let S be the set of all differences a(j)-a(i) for 0 <= i < j <= n-1, and let d be the smallest positive integer not in S. If, for every i in 1..n-1, a(n-1) + d - a(i) is not in S, then a(n) = a(n-1) + d. Otherwise, let r be the smallest positive integer such that, for every i in 1..n-1, neither a(n-1) + r - a(i) nor a(n-1) + r + d - a(i) is in S; then a(n) = a(n-1) + r and a(n+1) = a(n) + d.

Conjecture: lim inf_{n->oo} (a(n+1) - a(n))/n = 1/2. Conjecture: a(n) ~ log(log(n))*A025582(n+1), where A025582(m)+1 = A005282(m) is the Mian-Chowla sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Sep 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009089 | OEIS-A247601

Least positive integer m with pi(m*n) = phi(m), where pi(.) is the prime-counting function and phi(.) is Euler's totient function.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009090 | OEIS-A247602

Least positive integer m with pi(m*n) = phi(m+n), where pi(.) is the prime-counting function and phi(.) is Euler's totient function.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009091 | OEIS-A247603

Least integer m > 0 with pi(m*n) = sigma(m), where sigma(m) is the sum of all positive divisors of m.

Conjecture: a(n) exists for any n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009092 | OEIS-A247604

Least integer m > 0 with pi(m*n) = sigma(m+n), where pi(.) and sigma(.) are given by A000720 and A000203.

Conjecture: a(n) exists for every n = 5,6,...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009093 | OEIS-A247623

Number of paths from (0,0) to the line x = n, each segment given by a vector (1,1), (1,-1), or (2,0), not crossing the x-axis, and including no horizontal segment on the x-axis.

Conjecture: (n+1)*a(n) +(n-3)*a(n-1) +2*(-3*n+2)*a(n-2) +2*(-3*n+8)*a(n-3) +(n-5)*a(n-4) +(n-5)*a(n-5)=0. - _R. J. Mathar_, Sep 23 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009094 | OEIS-A247672

Least integer m > 0 with pi(m*n) = phi(m) + phi(n), where pi(.) is the prime-counting function and phi(.) is Euler's totient function.

Conjecture: a(n) exists for every n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009095 | OEIS-A247673

Least integer m > 0 with pi(m*n) = sigma(m) + sigma(n), where pi(.) and sigma(.) are given by A000720 and A000203 respectively.

Conjecture: a(n) exists for every n = 5, 6, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009096 | OEIS-A247727

Number of length 1+3 0..n arrays with no disjoint pairs in any consecutive four terms having the same sum.

Conjectures from _Colin Barker_, Nov 07 2018: (Start) Conjectured e.g.f.: exp(x)*x*(8 + 16*x + 8*x^2 + x^3). - _Stefano Spezia_, Mar 12 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009097 | OEIS-A247790

Primes p such that sigma(sigma(2p-1)) is a prime.

Primes p such that A247954(p) = A000203(A000203(2p-1)) = A000203(A008438(p-1)) = A051027(2p-1) is a prime q. The corresponding values of the primes q are: 7, 131071, 524287, ... (A247791). Conjecture: the primes q are Mersenne primes (A000668). Conjecture: the next term is 500461553802019261 (see comment from _Hiroaki Yamanouchi_ in A247821). - _Jaroslav Krizek_, Oct 08 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Sep 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009098 | OEIS-A247791

Primes p such that there is a prime q for which sigma(sigma(2*q-1)) = p.

Conjecture: Subsequence of Mersenne primes. Conjecture: the next term is 2305843009213693951 when 2305843009213693951 = sigma(sigma(2*500461553802019261-1)) where 500461553802019261 is prime (see comment of Hiroaki Yamanouchi in A247821). - _Jaroslav Krizek_, Oct 08 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Sep 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009099 | OEIS-A247793

Least integer m > 0 such that pi(m*n) divides prime(m) + prime(n), where pi(x) denotes the number of primes not exceeding x.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009100 | OEIS-A247821

Numbers k such that sigma(sigma(2k-1)) is a prime p.

Corresponding values of primes p are 7, 8191, 8191, 131071, 524287, 524287, ... (= A247822). Conjecture: The primes p are Mersenne primes (A000668). If the above conjecture is true, the next terms are 366479720500691270, 375344017599431990, 500461553802019261, 554079264075351985, 983755880192409499916700 86, ... - _Hiroaki Yamanouchi_, Oct 01 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Sep 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009101 | OEIS-A247822

Corresponding values of primes p from A247821 and A247838.

Conjecture: all terms are Mersenne primes (A000668). Conjecture: next terms are 2305843009213693951, 2305843009213693951, 2305843009213693951, 2305843009213693951 and 6189700196426901374495621 11. - _Jaroslav Krizek_, Mar 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Sep 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009102 | OEIS-A247834

Maximal non-semiprime number which is a "preprime" of the n-th kind (defined in comment in A247395).

Conjecture: the sequence contains all cubes of primes, except for 3^3 (cf. A030078). Conjecture: every term has the form a(n)= p*q*r, where p<=q<=r are primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009103 | OEIS-A247835

Indices of prime(n)^3 in A247834, or a(n)=0 if prime(n)^3 is not in A247834.

Conjecture: all a(n)>0, except for n=2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009104 | OEIS-A247838

Numbers k such that sigma(sigma(k)) is prime.

Conjecture: Subsequence of A046528 (numbers that are a product of distinct Mersenne primes). Corresponding values of primes p: 7, 8191, 8191, 131071, 524287, 524287, ... (A247822). Conjecture: values of primes p is equal to Mersenne primes (A000668).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Sep 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009105 | OEIS-A247867

a(n) is the smallest prime in the interval [k*sqrt(k), k*sqrt(k+2)], where k = A001359(n), or a(n)=0 if there is no prime in this interval.

The sequence is partly connected with conjecture in A247834. In turn, we conjecture that all terms a(n)>0 for n>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009106 | OEIS-A247893

Least integer k > 0 such that prime(k) - k*n is a square.

Conjecture: a(n) exists for any n > 0. See also A247278 for a related conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009107 | OEIS-A247895

Least integer k > 0 such that prime(k) - k*n is prime.

Conjecture: (i) a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009108 | OEIS-A247937

Least integer m > n such that m + n divides F(m) + F(n), where F(k) refers to the Fibonacci number A000045(k).

Conjecture: Let A be any integer not congruent to 3 modulo 6. Define u(0) = 0, u(1) = 1, and u(n+1) = A*u(n) + u(n-1) for n > 0. Then, for any integer n > 0, there are infinitely many positive integers m such that m + n divides u(m) + u(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009109 | OEIS-A247938

Sum of divisors of 2^prime(n)-1.

Conjecture: a(n)/2^prime(n) reaches its maximum value 135/128 at n = 5. - _Jianing Song_, Dec 31 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Sep 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009110 | OEIS-A247940

Least integer m > n such that m + n divides L(m) + L(n), where L(k) refers to the Lucas number A000032(k).

Conjecture: Let A be any integer not congruent to 3 modulo 6. Define v(0) = 2, v(1) = A, and v(n+1) = A*v(n) + v(n-1) for n > 0. Then, for any integer n > 0, there are infinitely many positive integers m such that m + n divides v(m) + v(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009111 | OEIS-A247975

Least positive integer m such that m + n divides prime(m)^2 + prime(n)^2.

Conjecture: a(n) exists for any n > 0. - _Zhi-Wei Sun_, Sep 28 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009112 | OEIS-A247996

Number of length 1+5 0..n arrays with no disjoint triples in any consecutive six terms having the same sum.

Conjectures from _Colin Barker_, Nov 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009113 | OEIS-A248006

Least positive integer m such that m + n divides phi(m*n), where phi(.) is Euler's totient function.

Conjecture: For any n > 2, a(n) exists and a(n) <= n. See also A248007 and A248008 for similar conjectures. - _Zhi-Wei Sun_, Sep 29 2014 The conjecture is true: One can show that 2*n divides phi(n^2) for all n > 2. So, a(n) is at most n. - _Derek Orr_, Sep 29 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009114 | OEIS-A248007

Least positive integer m such that m + n divides phi(m)*phi(n), where phi(.) is Euler's totient function.

Conjecture: a(n) exists for any n > 6. - _Zhi-Wei Sun_, Sep 29 2014 Numbers n for which a(n) > n: 10, 12, 22, 26, 42, 78, 166, 266, 290. The next term in this mini-sequence, if it exists, is greater than 3*10^4. I conjecture this list is finite. - _Derek Orr_, Sep 29 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009115 | OEIS-A248008

Least positive integer m such that m + n divides sigma(m*n), where sigma(k) denotes the sum of all positive divisors of k.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009116 | OEIS-A248029

Least positive integer m such that m + n divides phi(m)*sigma(n), where phi(.) and sigma(.) are given by A000010 and A000203.

Conjecture: For any n > 1, we have a(n) <= n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009117 | OEIS-A248030

Least positive integer m such that m + n divides sigma(m)*phi(n), where sigma(.) and phi(.) are given by A000203 and A000010.

Conjecture: a(n) < 2*n for any n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009118 | OEIS-A248036

Least positive integer m such that m + n divides sigma(m)^2 + sigma(n)^2, where sigma(k) denotes the number of positive divisors of k.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009119 | OEIS-A248044

Least positive integer m such that m + n divides pi(m)^2 + pi(n)^2, where pi(x) denotes the number of primes not exceeding x.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009120 | OEIS-A248049

a(n) = (a(n-1) + a(n-2)) * (a(n-2) + a(n-3)) / a(n-4) with a(0) = 2, a(1) = a(2) = a(3) = 1.

Somos conjectures that log(a(n)) ~ 1.25255*c^n, where c = A060006. - _Bill McEachen_, Oct 11 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Sep 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009121 | OEIS-A248054

Least positive integer m such that m + n divides sigma(m^2) + sigma(n^2), where sigma(k) is the sum of all positive divisors of k.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009122 | OEIS-A248108

Areas of primitive Heronian triangles K which are perfect squares.

It is known that every positive integer is the area of some rational triangle. Hence for every n > 0 there exists at least one primitive Heronian triangle with area K such that n*k^2 = K for some positive integer k. Therefore for the integer 1 there exists primitive Heronian triangles with area K = k^2. This sequence identifies all such occurrences of square areas from lists of primitive Heronian triangles generated by _Sascha Kurz_ (see link). The sequence excludes repetitive terms and is exhaustive as the Kurz lists searched include all primitive Heronian triangles up to a maximum side length of 6000000 and this sequence only includes areas that do not exceed 6000000 (see _T. D. Noe_ comments at A083875). All 46 terms found are displayed. It is conjectured that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Oct 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009123 | OEIS-A248121

a(n) = floor(1 / (1/n - Pi^2/6 + Sum_{h=1..n} 1/h^2)).

Conjectures from _Chai Wah Wu_, Aug 03 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009124 | OEIS-A248123

Least integer m > 0 such that gcd(m,n) = 1 and m*n | C(m+n), where C(k) refers to the k-th Catalan number binomial(2k,k)/(k+1).

Conjecture: a(n) exists for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009125 | OEIS-A248124

Least integer m > 0 such that gcd(m,n) = 1 and (m+n) | (C(m)+C(n)), where C(k) refers to the k-th Catalan number, binomial(2k,k)/(k+1).

Conjecture: a(n) exists for all n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009126 | OEIS-A248125

Least positive integer m such that m + n divides C(2m,m) + C(2n,n), where C(2k,k) = (2k)!/(k!)^2.

Conjecture: a(n) exists for all n > 0. Moreover, for n > 66 we have a(n) < n except for n = 364, 408.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009127 | OEIS-A248133

Least positive integer m such that m + n divides T(m) + T(n), where T(.) is given by A002426.

Conjecture: a(n) exists for any n > 0. Moreover, a(n) <= n^2 - n + 1 except for n = 274.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009128 | OEIS-A248136

Least positive integer m such that m + n divides D(m) + D(n), where D(.) is given by A001850.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009129 | OEIS-A248137

Least positive integer m such that m + n divides M(m) + M(n), where M(.) is given by A001006.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009130 | OEIS-A248139

Least positive integer m such that m + n divides f(m) + f(n), where f(.) is given by A000172.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009131 | OEIS-A248142

Least positive integer m such that m + n divides A(m) + A(n), where A(.) is given by A005259.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009132 | OEIS-A248143

Least integer m > 0 such that m + n divides p(m) + p(n), where p(.) is the partition function given by A000041.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009133 | OEIS-A248144

Least positive integer m such that m + n divides p(m*n), where p(.) is the partition function given by A000041.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009134 | OEIS-A248175

Least positive integer m such that m + n divides q(m*n), where q(.) is the strict partition function given by A000009.

Conjecture: (i) a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009135 | OEIS-A248183

Least k such that 1/4 - Sum_{h = 1..k} 1/(h*(h+1)*(h+2)) < 1/n^2.

Conjecture: a(n) = floor(sqrt(n^2/2 + 1) - 1/2), for n>1. - _Ridouane Oudra_, Sep 06 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009136 | OEIS-A248186

Least k such that 1/18 - Sum_{h=1..k} 1/(h*(h+1)*(h+2)*(h+3)) < 1/n^3.

Conjecture: a(n) = floor(r*n + r^2/n - 1), where r = 3^(-1/3), for n > 2. - _Ridouane Oudra_, Oct 06 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009137 | OEIS-A248189

Least positive integer m such that m*n divides sigma(m^2+n^2), where sigma(k) is the sum of all positive divisors of k.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009138 | OEIS-A248219

Indices where A248218(n) = 1.

Conjecture: a(n) ~ n^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Oct 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009139 | OEIS-A248230

a(n) = floor(1/(zeta(4) - Sum_{h=1..n} 1/h^4)).

Conjecture: a(n) = 1 + 7*n/2 + 9*n^2/2 + 3*n^3 + floor(n/4), holds for all n <= 10000. - _Vaclav Kotesovec_, Oct 09 2014 [Proved in the article by Sahu, 2018]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009140 | OEIS-A248231

Least k such that zeta(5) - Sum_{h = 1..k} 1/h^5 < 1/n^4.

Conjecture: a(n) = floor(sqrt(n^2/2 - 1) + 1/2), for n>1. - _Ridouane Oudra_, Sep 06 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009141 | OEIS-A248234

a(n) = floor(1/(zeta(5) - Sum_{h=1..n} 1/h^5)).

Conjecture (for n >= 3): a(n) = (12*n*(1+n)*(4+3*n+3*n^2) - 8 - cos(2*n*Pi/3) + sqrt(3)*sin(2*n*Pi/3))/9. - _Vaclav Kotesovec_, Oct 09 2014 [Proved in the article by Sahu, 2018] Conjecture (for n >= 2): a(n) = (2*n+1)*(n^3+(n+1)^3) + A183861(n-1). - _Rui Ferreira_, Aug 29 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009142 | OEIS-A248369

Least positive integer m such that prime(m+n) - prime(m) divides m.

Conjecture: a(n) exists for any n > 0. Moreover, for n > 9 we have a(n) < n^2 except for n = 12, 19, 36, 38.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009143 | OEIS-A248370

a(n+1) is the smallest prime beyond a(n) such that a(n+1) modulo a(n) is (congruent to) a prime; start with a(0)=1.

Shevelev cites El Bachraoui, Lemoine-Levy's conjecture and A046927, cf. links.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _M. F. Hasler_, Oct 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009144 | OEIS-A248381

Multiples of 3 in A247665 in order of appearance.

The initial multiples of 3 occur at positions 6,13,28,57,115,231, ... - conjecturally, for n > 13, A247665(n) is a multiple of 3 iff n = 29*2^k-1, k >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Russ Cox_ and _N. J. A. Sloane_, Oct 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009145 | OEIS-A248427

Circumference of the (n,n)-knight graph.

Colin Barker's conjectures confirmed using first Mathematica program. - _Ray Chandler_, Jan 14 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Oct 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009146 | OEIS-A248516

n^2+1 divided by its largest prime factor.

Conjecture: The Fibonacci numbers in the sequence are 1, 2, 5, 13, 34, 89, 233, 377, 610, 1597, 4181, 10946, 28657, 75025, 121393, 196418, ... including all of A001519.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jun 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009147 | OEIS-A248520

Numbers k such that floor(cot(Pi/(2k+2))) = floor(cot(Pi/(2k))).

Conjecture: a(n) ~ Pi/(Pi-2) * n. - _Vaclav Kotesovec_, Oct 09 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009148 | OEIS-A248521

Numbers k such that floor(cot(Pi/(2k+2))) = floor(cot(Pi/(2k))) + 1.

Conjecture: a(n) ~ Pi*n/2. - _Vaclav Kotesovec_, Oct 09 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009149 | OEIS-A248557

Decimal expansion of (Pi/2)^(1/4)/Gamma(3/4).

Conjecture: Equals Sum_{k>=0} A002448(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Dec 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009150 | OEIS-A248575

Rounded sums of the non-integer cube roots of n, as partitioned by the integer roots: round(Sum_{j=n^3+1..(n+1)^3-1} j^(1/3)).

See A247112 for additional references to similar sequences and a conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard R. Forberg_, Dec 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009151 | OEIS-A248586

a(n) = Sum_{i=0..n} C(n,i)*C(2i,i)^2.

Conjecture: n^2*a(n) +(-19*n^2+19*n-5)*a(n-1) +35*(n-1)^2*a(n-2) -17*(n-1)*(n-2)*a(n-3)=0. Remark: the conjectured recurrence for the coefficients a(n) comes from this diff. eq. for A(t).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Oct 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009152 | OEIS-A248588

Least positive integer m such that m + n divides sigma(m), where sigma(m) is the sum of all positive divisors of m.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009153 | OEIS-A248593

Least positive integer m such that m + n divides F(m), where F(m) is the m-th Fibonacci number given by A000045.

Conjecture: a(n) exists for any n > 0. Moreover, a(n) <= n*(n-1) except for n = 1, 2, 3, 9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009154 | OEIS-A248621

Floor of sums of the squares of the non-integer cube roots of n, as partitioned by the integer roots: floor(Sum_{j=n^3+1..(n+1)^3-1} j^(2/3)).

See A247112 for the cube of the square roots, other references and a conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard R. Forberg_, Dec 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009155 | OEIS-A248626

Least positive integer m such that prime(m+n) divides 2^m - 1.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009156 | OEIS-A248667

Numbers k for which coefficients of the polynomial p(k,x) defined in Comments are relatively prime.

Conjecture: any composite term can be represented as a product a(i)*a(j) (i > 1, j > 1) in at least one way. - _Mikhail Kurkov_, Apr 09 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009157 | OEIS-A248698

Floor of sums of the non-integer fourth roots of n, as partitioned by the integer roots: floor(Sum_{j=n^4+1..(n+1)^4-1} j^(1/4)).

See A247112 for references to other related sequences and a conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard R. Forberg_, Dec 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009158 | OEIS-A248727

A046802(x,y) --> A046802(x,y+1), transform of e.g.f. for the graded number of positroids of the totally nonnegative Grassmannians G+(k,n); enumerates faces of the stellahedra.

Second diagonal is A052944. (Changed from conjecture to fact on Nov 08 2016.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tom Copeland_, Oct 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009159 | OEIS-A248736

Array, read by antidiagonals, of the numbers of digits in the decimal expansion of the number of partitions of b^n employing a conjectured formula. See both the Comments and the Mathematica coding.

Array, read by antidiagonals, of the numbers of digits in the decimal expansion of the number of partitions of b^n employing a conjectured formula. See both the Comments and the Mathematica coding. This array is based upon the conjectured formula by _Charles R Greathouse IV_ in A077644, adapted to other bases.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Oct 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009160 | OEIS-A248833

The curvature of touching circles inscribed in a special way in the larger segment of circle of radius 1/6 divided by a chord of length sqrt(8/75).

Refer to comment of A092387. Consider a circle C of radius 1/6 (in some length units) with a chord of length sqrt(8/75). This has been chosen such that the larger sagitta has length 1/5. The input, besides the circle C, is the circle C_0 with radius R_0 = 1/10, touching the chord and circle C. The following sequence of circles C_n with radii R_n, n >= 1, is obtained from the conditions that C_n touches (i) the circle C, (ii) the chord and (iii) the circle C_(n-1). The curvature of the n-th circle, C_n = 1/R_n, n >= 0, is conjectured to be a(n). If one considers the curvature of touching circles inscribed in the smaller segment (sagitta length 2/15), the sequence would be A248834. See an illustration given in the link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Oct 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009161 | OEIS-A248834

The numerator of curvature of touching circles inscribed in a special way in the smaller segment of circle of radius 1/6 divided by a chord of length sqrt(8/75).

Refer to comment of A092387. Consider a circle C of radius 1/6 (in some length units) with a chord of length sqrt(8/75). This has been chosen such that the smaller sagitta has length 2/15. The input, besides the circle C, is the circle C_0 with radius R_0 = 1/15, touching the chord and circle C. The following sequence of circles C_n with radii R_n, n >= 1, is obtained from the conditions that C_n touches (i) the circle C, (ii) the chord and (iii) the circle C_(n-1). The curvature of the n-th circle is C_n = 1/R_n, n >= 0, and its numerator is conjectured to be a(n). The denominator is A000244 for n > 0. If one considers the curvature of touching circles inscribed in the larger segment (sagitta length 1/5), the sequence would be A248833. See an illustration given in the link. Conjecture: a(n) = 17*a(n-1) - 51*a(n-2) + 27*a(n-3) for n > 3. - _Colin Barker_, Oct 15 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Oct 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009162 | OEIS-A248836

Number of length n arrays x(i), i=1..n with x(i) in 0..i and no value appearing more than 2 times.

Conjecture: E.g.f.: 1/(1 - sin(x))^2. If the above conjecture is correct, the following general term is obtained: Conjecture from _Mikhail Kurkov_, Jun 26 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009163 | OEIS-A248897

Decimal expansion of Sum_{i >= 0} (i!)^2/(2*i+1)!.

Conjecture: for k >= 0, 2*sqrt(3)*Pi/9 = (3/2)^k * k!*Sum_{n = -oo..oo} (-1)^n/ Product_{j = 0..k} (3*n + 3*j + 1). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bruno Berselli_, Mar 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009164 | OEIS-A248919

"Stubborn primes" (see comments in A232210).

However, there are two conjectures: 1) for every n, prime a(n) exists (Shevelev); 2) already prime a(8) does not exist (Havermann).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Oct 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009165 | OEIS-A248923

a(n) is the smallest k >= n such that prime(n)*prime(k) - prime(n+k) is a perfect square.

Conjecture: a(n) exists for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Oct 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009166 | OEIS-A248958

Maximum number of distinct nonempty squares in a binary string of length n.

a(n) < n (proved by Brlek and Li, answering a longstanding conjecture). - _Jeffrey Shallit_, Dec 17 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _W. Edwin Clark_, Oct 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009167 | OEIS-A248982

Sequence of distinct least positive numbers such that the average of the first n terms is a Fibonacci number.

Conjecture: a(n) = 2*a(n-2)+a(n-4)-2*a(n-6)-a(n-8) for n > 17. - _Colin Barker_, Oct 19 2014 Conjecture: For n > 4, a(2*n+1) = A000045(n+3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Oct 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009168 | OEIS-A249056

a(n) = Product_{i=1..floor(n/2)} (2n-i)*(2n+i).

Conjecture: D-finite with recurrence +36*n*(3*n-1)*(3*n-2)*(879441252621897*n^2 -3487718196355549*n +3398030540730836)*a(n) -12*(n-1)*(17789803949566350*n^4 -31081184080278385 5*n^3 +97741257241074 2017*n^2 -12282029746841307 38*n +638585163822021576)*a(n-1) -15*(n-2)*(549650782888685625*n^6 -2729474655610903750*n^5 +4816600358041763875*n^4 -3395195710682405750*n^3 +230889009555801948*n^2 +1176562067066757228*n -715918902620114592)*a(n-2) +5*(5*n-11)*(n-3)*(5*n-14)*(5*n-13)*(5*n-12)*(1976644883285150*n^2 -4064135701440945*n

+2004921500186538)*a(n-3)=0. - _R. J. Mathar_, Jul 27 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Oct 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009169 | OEIS-A249067

Smallest k such that {first digit of j*n, 0<=j<=k} = {0,1,...,9}.

Conjecture: a(n) <= N for all n. Perhaps N can be taken as 81. - _Charles R Greathouse IV_, Oct 20 2014 The conjecture above is true. In fact, a(n) takes on precisely 37 values: 9, 11, 12, 13, 14, 15, 16, 17, 18, 21, 22, 23, 24, 25, 26, 27, 31, 32, 33, 34, 35, 36, 41, 42, 43, 44, 45, 51, 52, 53, 54, 61, 62, 63, 71, 72, and 81. The last of these to occur is a(291) = 31. - _Charles R Greathouse IV_, Mar 17 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian Perfect_, Oct 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009170 | OEIS-A249093

Length of self-iteration of the Kolakoski sequence A000002 starting at A000002(n): a(n) = max { k | A000002(n+i-1) = A000002(i), 0 < i <= k }.

Conjecture: for any k, there is an iteration of length A054352(k) in A000002.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-Christophe Hervé_, Oct 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009171 | OEIS-A249120

Triangle read by rows: T(n,k), n>=1, k>=1, in which column k lists the numbers of A210843 multiplied by A000330(k), and the first element of column k is in row A000217(k).

Conjecture: gives an identity for the sum of all divisors of all positive integers <= n. Alternating sum of row n equals A024916(n), i.e., Sum_{k=1..A003056(n)} (-1)^(k-1)*T(n,k) = A024916(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Dec 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009172 | OEIS-A249129

Lexicographically earliest sequence of distinct nonnegative integers such that a(2n) = a(n) + a(n+1) for all n >= 0.

I conjecture that a(n) ~ n. The differences d(n) = a(n)-n are in A249161. The first occurrences of the small differences are at d(0)=1, d(1)=-1, d(2)=0, d(6)=2, d(9)=-2, d(122)=3, d(127)=-3, d(922)=4, d(929)=-4, d(1994)=5, d(2003)=-5, d(3986)=6, d(3997)=-6,... It seems that if d(n) is the first occurrence of k>0, then n is even and d(n+2k-1) = -k is the first occurrence of the opposite value. - _M. F. Hasler_, Oct 22 2014 Is there a proof of the plausible conjecture that every number appears? - _N. J. A. Sloane_, Oct 17 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _M. F. Hasler_, Oct 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009173 | OEIS-A249167

a(n) = n if n <= 3, otherwise the smallest number not occurring earlier having at least one common Fermi-Dirac factor with a(n-2), but none with a(n-1).

Conjecture: the sequence is a permutation of the positive integers. Conjecture is true. The proof is similar to that for A098550 with minor changes. - _Vladimir Shevelev_, Jan 26 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Dec 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009174 | OEIS-A249223

Triangle read by rows: row n gives partial alternating sums of row n of A237048.

For a proof that T(n, k) = |{ d : d|n and k/2 < d <= k }|, for 1 <= k <= row(n), where row(n) is the length of row n, an identity suggested by _Peter Munn_, see the link. A corollary to it is that the number of divisors of n in the half-open interval (row(n)/2, row(n)] equals the width of the symmetric representation of n at the diagonal: T(n, row(n)) = | { d : d|n and row(n)/2 < d <= row(n) } |. See also the comments and conjectures of _Michel Marcus_ in A067742 and A237593. - _Hartmut F. W. Hoft_, Jun 24 2024 Conjecture 1: Every column is a periodic sequence. Conjecture 2: The periods of the columns 1..8 are respectively: 1, 2, 6, 12, 60, 60, 420, 840. Conjecture 3: Column 3 gives [2, 0] together with A115357, hence column 3 gives 2 together with A171182. Conjectures 1 and 2 are true and the answer to question 1 is affirmative. Conjecture 3 and Question 2 are true. Since T237048(n, 1) = 1, T237208(n, 2) = 1 if n odd and 0 if n even, T237048(n, 3) = 1 if 3|n and 0 otherwise, and T237048(n, 4) = 1 if 4|(n-2) and 0 otherwise, equations T249223(n, 3) = 1 - (n mod 2) + delta(n mod 3) and T249223(n, 4) = 1 - (n mod 2) + delta(n mod 3) - delta((n-2) mod 4) hold where delta(k) = 1 if k = 0 and 0 otherwise. With the 3rd column starting at n = A000217(3) = 6, each period starting in a row that is a multiple of 6 is [2 0 1 1 1 0], and appropriate shifts yield A115357 and A171182. With the 4th column starting at n = A000217(4) = 10, each period starting in a row n with 12|(n+2) is [0 0 2 0 0 1 1 0 1 0 1 1], and with a shift of 9 yields the apparently periodic A337976(10), A337976(11), ... (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Oct 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009175 | OEIS-A249264

Sequence of distinct least nonnegative numbers such that the average of the first n terms is a triangular number.

Conjectured: For n > 6, a(2*n-1) = A000566(n) and a(2*n) = A000217(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Oct 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009176 | OEIS-A249265

Sequence of distinct least nonnegative numbers such that the average of the first n terms is a Fibonacci number.

Conjecture: a(n) = 2*a(n-2) + a(n-4) - 2*a(n-6) - a(n-8) for n > 15. - _Colin Barker_, Oct 24 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Oct 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009177 | OEIS-A249302

Prime-partitionable numbers a(n) for which there exists a 2-partition of the set of primes < a(n) that has a smallest subset containing three primes only.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christopher Hunt Gribble_, Oct 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009178 | OEIS-A249303

Triangular array: row n gives the coefficients of the polynomial p(n,x) defined in Comments.

Conjecture: for n > 2, p(n,x) is irreducible if and only if n is a (prime - 2). More generally, if c is arbitrary and f(n,x) = 1 + (x + c)/f(n-1,x), where f(x,0) = 1, then p(n,x) is irreducible if and only if n is a (prime - 2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009179 | OEIS-A249351

Triangle read by rows in which row n lists the widths of the symmetric representation of sigma(n).

It appears that the middle diagonal is also A067742 (which was conjectured by _Michel Marcus_ in the entry A237593 and checked with two Mathematica functions up to n = 100000 by _Hartmut F. W. Hoft_).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Oct 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009180 | OEIS-A249394

Numbers k such that the k-th ménage number is prime.

This sequence is heuristically conjectured to grow doubly-exponentially.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Adam P. Goucher_, Oct 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009181 | OEIS-A249457

The numerator of curvatures of touching circles inscribed in a special way in the larger segment of a unit circle divided by a chord of length sqrt(84)/5.

The denominators are conjectured to be A005032. Refer to comments and links of A092387. Consider a unit circle with a chord of length sqrt(84)/5. This has been chosen such that the larger sagitta has length 7/5. The input, besides the unit circle C, is the circle C_0 with radius R_0 = 7/10, touching the chord and circle C. The following sequence of circles C_n with radii R_n, n >= 1, is obtained from the conditions that C_n touches (i) the circle C, (ii) the chord and (iii) the circle C_(n-1). The curvature of the n-th circle is C_n = 1/R_n, n >= 0, and its numerator is conjectured to be a(n). O.g.f.: 5*((1 - 17*x)/(1 - 34*x + 9*x^2) + 1/(1-3*x)) = 10*(1 - 27*x + 30*x^2)/((1 - 34*x + 9*x^2)*(1 - 3*x)) proving the conjecture of Colin Barker above. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Oct 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009182 | OEIS-A249458

The numerators of curvatures of touching circles inscribed in a special way in the smaller segment of unit circle divided by a chord of length sqrt(84)/5.

The denominators are conjectured to be A169634. Refer to comments and links of A092387. Consider a unit circle with a chord of length sqrt(84)/5. This has been chosen such that the smaller sagitta has length 3/5. The input, besides the circle C, is the circle C_0 with radius R_0 = 3/10, touching the chord and circle C. The following sequence of circles C_n with radii R_n, n >= 1, is obtained from the conditions that C_n touches (i) the circle C, (ii) the chord and (iii) the circle C_(n-1). The curvature of the n-th circle is C_n = 1/R_n, n >= 0, and its numerator is conjectured to be a(n). If one considers the curvature of touching circles inscribed in the larger segment (sagitta length 7/5), the sequence would be A249457/A005032. See an illustration given in the link. O.g.f.: 5*((1 - 13*x)/(1 - 26*x + (7*x)^2) + 1/(1-7*x)) = 10*(1 - 23*x + 70*x^2)/((1 - 26*x + (7*x)^2)*(1 - 7*x)) proving the conjecture of Colin Barker above. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Oct 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009183 | OEIS-A249482

Numbers n such that the summatory Liouville function L(n) (A002819) is zero and L(n-1)*L(n+1) = -1.

G. Polya (1919) conjectured that L(n)<=0, for n>=2. But this was disproved in 1958 by B. Haselgrove, and in 1980 M. Tanaka found the smallest counterexample, a(2)+1 = 906150257.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 13 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009184 | OEIS-A249507

Length of full iterated word (direct branch and reverse branch) of the Kolakoski sequence A000002 initiated at A000002(n). If the reverse branch is longer than the direct branch, the total length is counted negatively: if A000002(n) = 2, a(n) = 0; if A000002(n) = 1, a(n) = (ld+lr-1)*sign(ld-lr) with ld = max { k | A000002(n-i+1) = A000002(i), 0<i<=k } and lr = max { k | A000002(n-i+1) = A000002(i), 0<i<=k }.

The iterated words are initiated by the 1's in the OK sequence, but the 1's in pairs 11 can only lead to very short iterated words: (1)-1-2(1) or (1)-1-221(2), or the same in reverse order, with values in this sequence of +/-2 or +/-4. On the contrary, each of the single 1's (212) in the OK sequence (except the first) is associated with an iterated word of length at least five, and (conjecture) it is likely that there is no bound to the length of iterated words associated with a single 1. The number of iterated words with the right branch longer than the left seems to be well balanced, so that (conjecture) the limit of partial sums of this sequence could be o(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-Christophe Hervé_, Oct 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009185 | OEIS-A249541

Numbers m such that phi(m-2) divides m-1 where phi is Euler's totient function (A000010).

Conjecture: 4 is the only number m such that 3*phi(m-2) = m-1. (See comment in A203966.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Feb 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009186 | OEIS-A249549

Conjectured largest number k not divisible by 10 such that k^k has exactly n distinct decimal digits.

Conjectured largest number k not divisible by 10 such that k^k has exactly n distinct decimal digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Oct 31 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009187 | OEIS-A249565

Number of self-avoiding walks on the truncated square tiling with n steps.

It is conjectured that a(n) is approximately mu^n*n^{11/32} for large n where mu is the connective constant and mu is approximately 1.80883001(6).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mike Zabrocki_, Nov 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009188 | OEIS-A249566

Numbers n such that A182134(n) = 4, i.e., there exist exactly four primes p with prime(n) < p < prime(n)^(1+1/n).

It is interesting that three consecutive integers n = 20004097201301075, n + 1 and n + 2 are in the sequence. Conjecture: The sequence is infinite. - _Farideh Firoozbakht_, Nov 01 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009189 | OEIS-A249570

Conjectured values of least number k such that k^k + n^n is prime, or 0 if no such k exists.

Conjectured values of least number k such that k^k + n^n is prime, or 0 if no such k exists. The zero entries are only conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Nov 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009190 | OEIS-A249596

Analog of A097717 in base 2.

Conjecture: a(n) = n*A165781(n). - _R. J. Mathar_, Nov 11 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Mar 30 2009 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009191 | OEIS-A249609

a(n) is the smallest m, 1<=m<=n, such that binomial(n,m) is evil (A001969); a(n)=0 if there is no such m.

Conjecture: there are only five n: 0,1,2,7,8, for which all entries of the n-th Pascal row (A007318) are odious (A000069). _Peter J. C. Moses_ verified the conjecture up to n = 10^6. Conjecture verified up to n = 5*10^10. - _Michael S. Branicky_, Jul 13 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Nov 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009192 | OEIS-A249621

Squares that remain squares when prepended with 1 and appended with 25 in base 10.

Conjecture: all the numbers in the sequence end in either 00 or 56.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Nov 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009193 | OEIS-A249623

Conjectured least number k such that k^k - n^n is prime.

Conjectured least number k such that k^k - n^n is prime. The zero entries are only conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Nov 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009194 | OEIS-A249679

Terms of A007504 divisible by 3.

Conjecture: a(n) ~ 4.5 n^2 log n. - _Charles R Greathouse IV_, Nov 03 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Nov 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009195 | OEIS-A249708

Number of length 2+3 0..n arrays with every four consecutive terms having the maximum of some two terms equal to the minimum of the remaining two terms.

Conjectures from _Colin Barker_, Nov 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009196 | OEIS-A249709

Number of length 3+3 0..n arrays with every four consecutive terms having the maximum of some two terms equal to the minimum of the remaining two terms.

Conjectures from _Colin Barker_, Nov 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009197 | OEIS-A249710

Number of length 4+3 0..n arrays with every four consecutive terms having the maximum of some two terms equal to the minimum of the remaining two terms.

Conjectures from _Colin Barker_, Nov 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009198 | OEIS-A249711

Number of length 5+3 0..n arrays with every four consecutive terms having the maximum of some two terms equal to the minimum of the remaining two terms.

Conjectures from _Colin Barker_, Nov 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009199 | OEIS-A249712

Number of length 6+3 0..n arrays with every four consecutive terms having the maximum of some two terms equal to the minimum of the remaining two terms.

Conjectures from _Colin Barker_, Nov 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009200 | OEIS-A249713

Number of length 7+3 0..n arrays with every four consecutive terms having the maximum of some two terms equal to the minimum of the remaining two terms.

Conjectures from _Colin Barker_, Nov 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009201 | OEIS-A249759

Primes p such that sigma(p-1) is a prime q.

Conjectures: 1) sequence is finite; 2) sequence is a subsequence of A019434 (Fermat primes); 3) sequence consists of Fermat primes p such that sigma(p-1) is a Mersenne prime; 4) a(n) = (A249761(n)+3)/2. Conjecture: 3 is the only number n such that n and sigma(n+1) are both prime. Conjecture: subsequence of A256438 and A278741. Conjecture: also primes p such that tau(p-1) is a prime q; corresponding values of primes q are 2, 3, 5, 17. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009202 | OEIS-A249760

Numbers k such that k+1 and sigma(k) are both primes.

Conjectures: (1) sequence is finite; (2) a(n) + 1 is a Fermat prime (A019434); (3) sigma(a(n)) is a Mersenne prime (A000668). Subsequence of A023194, and by a comment in that entry it follows that each term is a prime power. From that conjectures (2) and (3) above easily follow. - _Jeppe Stig Nielsen_, Jan 13 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009203 | OEIS-A249761

Primes p such that there is prime q with sigma(q-1) = p.

Conjectures: 1) sequence is finite; 2) (a(n)+3)/2 is a Fermat prime (A019434); 3) a(n) is subsequence of Mersenne primes (A000668); 4) a(n) = 2*A249759(n)-3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009204 | OEIS-A249833

E.g.f. satisfies: A(x) = 1 + Integral A(x) + A(x)^2*log(A(x)) dx.

Conjecture: a(n) = A370382(n-1, 0) for n > 0 with a(0) = 1. - _Mikhail Kurkov_, Apr 25 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009205 | OEIS-A249836

Numbers n for which tan(n) > n.

It is an open problem whether tan(n) > n for infinitely many integer n. log(a(n)) / n ~ Pi, conjectured. - _M. F. Hasler_, Sep 10 2020 [corrected thanks to _Vaclav Kotesovec_, Feb 22 2021]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jacob Vecht_, Nov 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009206 | OEIS-A249845

Number of length 1+4 0..n arrays with no five consecutive terms having the maximum of any two terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Aug 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009207 | OEIS-A249846

Number of length 2+4 0..n arrays with no five consecutive terms having the maximum of any two terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Aug 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009208 | OEIS-A249847

Number of length 3+4 0..n arrays with no five consecutive terms having the maximum of any two terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Aug 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009209 | OEIS-A249848

Number of length 4+4 0..n arrays with no five consecutive terms having the maximum of any two terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Aug 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009210 | OEIS-A249849

Number of length 5+4 0..n arrays with no five consecutive terms having the maximum of any two terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Aug 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009211 | OEIS-A249850

Number of length 6+4 0..n arrays with no five consecutive terms having the maximum of any two terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Aug 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009212 | OEIS-A249851

Number of length 7+4 0..n arrays with no five consecutive terms having the maximum of any two terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Aug 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009213 | OEIS-A249852

a(n) is the total number of pentagons on the left or the right of the vertical symmetry axis of a pentagon expansion (vertex to vertex) after n iterations.

Conjectures from _Colin Barker_, Mar 07 2015: (Start) Conjectured e.g.f.: (exp(x)*(1 + 8*x + 5*x^2) - cos(x) - sin(x))/4. - _Stefano Spezia_, Feb 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Mar 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009214 | OEIS-A249861

a(n) are values of k that yield "record-breaking" integer sequence lengths for the recursion: b(i) = 3*(b(i-1) + b(i-2))/2, with b(0) = 1 and b(1) = k.

Conjecture: Rules 1 & 2 apply for all such b(i) sequences of the form: b(i) = m*(b(i-1) + b(i-2))/2, b(0) = j, b(1) = k, for j >= 0 and odd integers m > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard R. Forberg_, Nov 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009215 | OEIS-A249884

Number of length 1+6 0..n arrays with no seven consecutive terms having the maximum of any three terms equal to the minimum of the remaining four terms.

Conjectures from _Colin Barker_, Nov 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009216 | OEIS-A249885

Number of length 2+6 0..n arrays with no seven consecutive terms having the maximum of any three terms equal to the minimum of the remaining four terms.

Conjectures from _Colin Barker_, Nov 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009217 | OEIS-A249902

Numbers n such that 2n-1 and sigma(n) are both primes.

Conjecture: 2 and 9 are the only numbers n such that 2n - 1, 2n + 1 and sigma(n) are all primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009218 | OEIS-A249903

Numbers n such that 2n+1 and sigma(n) are both noncomposite numbers.

Conjecture: 2 and 9 are the only numbers n such that 2n - 1, 2n + 1 and sigma(n) are all primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 14 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009219 | OEIS-A249916

a(n) = 4*(n - 1) - a(n-3), n >= 3, a(0) = a(1) = 1, a(2) = 5.

Conjecture: These are the natural numbers of the form 6*j +- 1 in which those of the form 12*k +- 1 are repeated. The second conjecture in Formula section is correct. This also confirms the conjecture in Comment section and the conjecture on the g.f. in Formula section. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Jan 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009220 | OEIS-A249938

E.g.f.: Sum_{n>=0} exp(n^2*x) / 2^(n+1).

Conjectures from _Federico Provvedi_, Nov 07 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009221 | OEIS-A249943

a(n) = smallest k such that the numbers 1..n appear among A098550(1), ..., A098550(k), or a(n) = 0 if there is no such k.

The conjecture that all terms are positive is equivalent to the known conjecture that A098550 is a permutation of the positive integers. The author conjectures that a(n)/n <= a(19)/19 = 43/19. _Peter J. C. Moses_ verified that the strict inequality holds for 19 < n <= 1.1*10^5. - _Vladimir Shevelev_, Dec 06 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Dec 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009222 | OEIS-A249947

Number of available orbitals at increasing subshells in multi-electron atoms.

Theoretically, the integers follow the pattern infinitely, but (conjecturally) the subshells stop after 7p, with 8s not existing as predicted.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alweiss_, Nov 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009223 | OEIS-A249949

Number of generations between each single 1 in the Kolakoski sequence A000002 and its nearest double 1 ancestor.

Conjecture: this sequence takes all integers k >= 1 as values, so there is no bound to the length of the iterated words that appear in the Kolakoski sequence; the limiting frequency of k is 2/3^k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-Christophe Hervé_, Nov 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009224 | OEIS-A249961

Number of length 1+5 0..n arrays with no six consecutive terms having the maximum of any two terms equal to the minimum of the remaining four terms.

Conjectures from _Colin Barker_, Aug 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009225 | OEIS-A249962

Number of length 2+5 0..n arrays with no six consecutive terms having the maximum of any two terms equal to the minimum of the remaining four terms.

Conjectures from _Colin Barker_, Aug 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009226 | OEIS-A249963

Number of length 3+5 0..n arrays with no six consecutive terms having the maximum of any two terms equal to the minimum of the remaining four terms.

Conjectures from _Colin Barker_, Aug 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009227 | OEIS-A249964

Number of length 4+5 0..n arrays with no six consecutive terms having the maximum of any two terms equal to the minimum of the remaining four terms.

Conjectures from _Colin Barker_, Aug 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009228 | OEIS-A249965

Number of length 5+5 0..n arrays with no six consecutive terms having the maximum of any two terms equal to the minimum of the remaining four terms.

Conjectures from _Colin Barker_, Aug 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009229 | OEIS-A249966

Number of length 6+5 0..n arrays with no six consecutive terms having the maximum of any two terms equal to the minimum of the remaining four terms.

Conjectures from _Colin Barker_, Aug 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009230 | OEIS-A249967

Number of length 7+5 0..n arrays with no six consecutive terms having the maximum of any two terms equal to the minimum of the remaining four terms.

Conjectures from _Colin Barker_, Aug 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009231 | OEIS-A249968

Numbers whose squares remain squares when prepended with 2 and appended with 25 in base 10.

Conjecture: squares of numbers in the sequence ends in either 00 or 56.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Nov 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009232 | OEIS-A249969

Numbers whose squares remain squares when prepended with 3 and appended with 25 in base 10.

Conjecture: squares of numbers in the sequence ends in either 00 or 56.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Nov 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009233 | OEIS-A249984

Number of length 4+1 0..2*n arrays with the sum of the absolute values of adjacent differences equal to 4*n.

Conjectures from _Colin Barker_, Nov 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009234 | OEIS-A249986

Number of length 6+1 0..2*n arrays with the sum of the absolute values of adjacent differences equal to 6*n.

Conjectures from _Colin Barker_, Nov 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

B06 A250000 - A299999 | 3,804 条

V3-009235 | OEIS-A250015

Number of length 1+5 0..n arrays with no six consecutive terms having the maximum of any three terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Nov 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009236 | OEIS-A250016

Number of length 2+5 0..n arrays with no six consecutive terms having the maximum of any three terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Nov 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009237 | OEIS-A250017

Number of length 3+5 0..n arrays with no six consecutive terms having the maximum of any three terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Nov 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 10 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009238 | OEIS-A250030

a(n) is the number of iterations of A008474 needed to reach 5, starting at n, where A008474(Product p_i^e_i) = Sum(p_i+e_i).

Conjecture 1: for m > 4, by iterating the map m -> A008474(m) one always reaches 5. Tested for m up to 320000. Conjecture 2: Let S(n) be the sum of the first a(n) numbers in the n-to-5 orbit, i.e., the sum of all the numbers in the orbit without 5. Except for n in [8,9], S(n) is not divisible by n. Verified for n up to 10^6. - _Ivan N. Ianakiev_, May 06 2015 Proof outline for Conjecture 1: 3. The only thing needed to complete the proof is to exhaustively confirm the conjecture for all numbers in [5, 29] that were not handled in items 1 and 2 above, i.e., 5, 6, 8, 9, 10, 11, 13, 14, 15, 21, 22, 25, 26. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ivan N. Ianakiev_, Nov 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009239 | OEIS-A250060

Number of length 1+6 0..n arrays with no seven consecutive terms having the maximum of any two terms equal to the minimum of the remaining five terms.

Conjectures from _Colin Barker_, Aug 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009240 | OEIS-A250061

Number of length 2+6 0..n arrays with no seven consecutive terms having the maximum of any two terms equal to the minimum of the remaining five terms.

Conjectures from _Colin Barker_, Aug 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009241 | OEIS-A250062

Number of length 3+6 0..n arrays with no seven consecutive terms having the maximum of any two terms equal to the minimum of the remaining five terms.

Conjectures from _Colin Barker_, Aug 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009242 | OEIS-A250063

Number of length 5+6 0..n arrays with no seven consecutive terms having the maximum of any two terms equal to the minimum of the remaining five terms.

Conjectures from _Colin Barker_, Aug 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009243 | OEIS-A250064

Number of length 4+6 0..n arrays with no seven consecutive terms having the maximum of any two terms equal to the minimum of the remaining five terms.

Conjectures from _Colin Barker_, Aug 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009244 | OEIS-A250065

Number of length 6+6 0..n arrays with no seven consecutive terms having the maximum of any two terms equal to the minimum of the remaining five terms.

Conjectures from _Colin Barker_, Aug 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009245 | OEIS-A250082

Number of length 1+5 0..n arrays with every six consecutive terms having the maximum of some two terms equal to the minimum of the remaining four terms.

Conjectures from _Colin Barker_, Nov 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009246 | OEIS-A250083

Number of length 2+5 0..n arrays with every six consecutive terms having the maximum of some two terms equal to the minimum of the remaining four terms.

Conjectures from _Colin Barker_, Nov 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009247 | OEIS-A250084

Number of length 3+5 0..n arrays with every six consecutive terms having the maximum of some two terms equal to the minimum of the remaining four terms.

Conjectures from _Colin Barker_, Nov 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009248 | OEIS-A250085

Number of length 4+5 0..n arrays with every six consecutive terms having the maximum of some two terms equal to the minimum of the remaining four terms.

Conjectures from _Colin Barker_, Nov 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009249 | OEIS-A250093

Squarefree part of Fibonacci(n^2).

Conjecture: the only primes in this sequence are 3 and 3001.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Nov 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009250 | OEIS-A250120

Coordination sequence for planar net 3.3.3.3.6 (also called the fsz net).

The Mathematica notebook can compute 30 or 40 iterations, and colors them with period 5. You could also change out images if you want to. These graphs are better for analyzing 5-iteration chunks of the pattern. You can see that under iteration all fragments of the circumferences are preserved in shape and translated outwards a distance approximately sqrt(21) (relative to small triangle edge), the length of a long diagonal of larger rhombus unit cell. The conjectured recurrence should follow from an analysis of how new pieces occur in between the translated pieces. - _Bradley Klee_, Nov 26 2014 Based on computations of Darrah Chavey, Bradley Klee, and Maurizio Paolini, it is conjectured that the first differences of this sequence are 4, 4, 6, 4, 5, 5, 4, 6, 4, 5, 5, 4, 6, 4, 5, 5, ..., that is, 4 followed by (4,6,4,5,5) repeated. All the above conjectures are true - for proof see link to my article with Chaim Goodman-Strauss. - _N. J. A. Sloane_, Jan 14 2018; link added Mar 26 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009251 | OEIS-A250122

Coordination sequence for planar net 3.12.12.

All the above conjectures are true. - _N. J. A. Sloane_, Dec 31 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Darrah Chavey_, Nov 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009252 | OEIS-A250128

Number of triforces generated at iteration n in a Koch-Sierpiński Ninja Star.

Conjecture: a(n) = 2*a(n-1) + 3*a(n-2) + 3*a(n-3). - _Colin Barker_, Nov 13 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kyle M. Zimmer_, Nov 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009253 | OEIS-A250131

a(n) is the odd part of the digital sum of 3^n divided by the maximal possible power of 3.

Consider the sequence {b(n)}, such that b(1)=2, b(2)=3, and for n>=3, b(n)=a(n-2). We conjecture that, if we apply the Eratosthenes-like sieve to b(n) and remove 1's, then we obtain a sequence of primes. _Peter J. C. Moses_ noted that these primes follow with some perturbation of order. For example, 73 appears before 71. Similarly, 101 and 103 appear before 97.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Dec 12 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009254 | OEIS-A250141

Number of length 2+2 0..n arrays with the medians of every three consecutive terms nondecreasing.

Conjectures from _Colin Barker_, Nov 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009255 | OEIS-A250142

Number of length 3+2 0..n arrays with the medians of every three consecutive terms nondecreasing.

Conjectures from _Colin Barker_, Nov 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009256 | OEIS-A250143

Number of length 4+2 0..n arrays with the medians of every three consecutive terms nondecreasing.

Conjectures from _Colin Barker_, Nov 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009257 | OEIS-A250144

Number of length 5+2 0..n arrays with the medians of every three consecutive terms nondecreasing.

Conjectures from _Colin Barker_, Nov 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009258 | OEIS-A250145

Number of length 6+2 0..n arrays with the medians of every three consecutive terms nondecreasing.

Conjectures from _Colin Barker_, Nov 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009259 | OEIS-A250155

Number of length 1+6 0..n arrays with every seven consecutive terms having the maximum of some two terms equal to the minimum of the remaining five terms.

Conjectures from _Colin Barker_, Nov 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009260 | OEIS-A250156

Number of length 2+6 0..n arrays with every seven consecutive terms having the maximum of some two terms equal to the minimum of the remaining five terms.

Conjectures from _Colin Barker_, Nov 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009261 | OEIS-A250157

Number of length 3+6 0..n arrays with every seven consecutive terms having the maximum of some two terms equal to the minimum of the remaining five terms.

Conjectures from _Colin Barker_, Nov 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009262 | OEIS-A250162

Number of length n+1 0..3 arrays with the sum of adjacent differences multiplied by some arrangement of +-1 equal to zero.

Conjectures from _Colin Barker_, Nov 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009263 | OEIS-A250201

Least b such that Phi_n(b, b-1) is prime.

This sequence is not defined at n = 1 since Phi_1(b, b-1) = 1 for all b, and 1 is not prime. Conjecture: a(n) is defined for all n>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, Mar 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009264 | OEIS-A250204

Sierpiński problem in base 6: Least k > 0 such that n*6^k+1 is prime, or 0 if no such k exists.

a(5k+4) = 0, since (5k+4)*6^n+1 is always divisible by 5, but there are infinitely many numbers not in the form 5k+4 such that a(n) = 0. For example, a(174308) = 0 since 174308*6^n+1 is always divisible by 7, 13, 31, 37, or 97 (See A123159). Conjecture: if n is not in the form 5k+4 and n < 174308, then a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, Mar 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009265 | OEIS-A250205

Riesel problem in base 6: Least k > 0 such that n*6^k-1 is prime, or 0 if no such k exists.

Conjecture: if n is not in the form 5j+1 and n < 84687, then a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, Mar 13 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009266 | OEIS-A250209

a(n) = least k such that k * n is in A072226, or 0 if no such k exists.

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, Jan 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009267 | OEIS-A250271

Number of length n+1 0..2 arrays with the sum of the squares of adjacent differences multiplied by some arrangement of +-1 equal to zero.

Conjectures from _Colin Barker_, Nov 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009268 | OEIS-A250307

Number of 2-colored Schroeder paths of semilength n in which there are no (2,0)-steps at level 1.

Conjecture: 2*n*a(n) +3*(-9*n+8)*a(n-1) +4*(28*n-39)*a(n-2) +4*(-43*n+81)*a(n-3) +64*(n-3)*a(n-4)=0. - _R. J. Mathar_, Sep 24 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José Luis Ramírez Ramírez_, Apr 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009269 | OEIS-A250337

Number of length 1+5 0..n arrays with every six consecutive terms having the maximum of some three terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Nov 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009270 | OEIS-A250338

Number of length 2+5 0..n arrays with every six consecutive terms having the maximum of some three terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Nov 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009271 | OEIS-A250339

Number of length 3+5 0..n arrays with every six consecutive terms having the maximum of some three terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Nov 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009272 | OEIS-A250340

Number of length 4+5 0..n arrays with every six consecutive terms having the maximum of some three terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Nov 12 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009273 | OEIS-A250354

Number of length 5 arrays x(i), i=1..5 with x(i) in i..i+n and no value appearing more than 2 times.

Conjectures from _Colin Barker_, Nov 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009274 | OEIS-A250355

Number of length 6 arrays x(i), i=1..6 with x(i) in i..i+n and no value appearing more than 2 times.

Conjectures from _Colin Barker_, Nov 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009275 | OEIS-A250362

Number of length 4 arrays x(i), i=1..4 with x(i) in i..i+n and no value appearing more than 3 times.

Conjectures from _Colin Barker_, Nov 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009276 | OEIS-A250363

Number of length 5 arrays x(i), i=1..5 with x(i) in i..i+n and no value appearing more than 3 times.

Conjectures from _Colin Barker_, Nov 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009277 | OEIS-A250364

Number of length 6 arrays x(i), i=1..6 with x(i) in i..i+n and no value appearing more than 3 times.

Conjectures from _Colin Barker_, Nov 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009278 | OEIS-A250374

Number of length 1+6 0..n arrays with every seven consecutive terms having the maximum of some three terms equal to the minimum of the remaining four terms.

Conjectures from _Colin Barker_, Nov 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009279 | OEIS-A250375

Number of length 2+6 0..n arrays with every seven consecutive terms having the maximum of some three terms equal to the minimum of the remaining four terms.

Conjectures from _Colin Barker_, Nov 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009280 | OEIS-A250376

Number of length 3+6 0..n arrays with every seven consecutive terms having the maximum of some three terms equal to the minimum of the remaining four terms.

Conjectures from _Colin Barker_, Nov 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 19 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009281 | OEIS-A250388

Number of length 2+3 0..n arrays with no four consecutive terms having the maximum of any two terms equal to the minimum of the remaining two terms.

Conjectures from _Colin Barker_, Aug 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009282 | OEIS-A250389

Number of length 3+3 0..n arrays with no four consecutive terms having the maximum of any two terms equal to the minimum of the remaining two terms.

Conjectures from _Colin Barker_, Aug 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009283 | OEIS-A250390

Number of length 4+3 0..n arrays with no four consecutive terms having the maximum of any two terms equal to the minimum of the remaining two terms.

Conjectures from _Colin Barker_, Aug 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009284 | OEIS-A250391

Number of length 5+3 0..n arrays with no four consecutive terms having the maximum of any two terms equal to the minimum of the remaining two terms.

Conjectures from _Colin Barker_, Aug 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009285 | OEIS-A250392

Number of length 6+3 0..n arrays with no four consecutive terms having the maximum of any two terms equal to the minimum of the remaining two terms.

Conjectures from _Colin Barker_, Aug 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009286 | OEIS-A250393

Number of length 7+3 0..n arrays with no four consecutive terms having the maximum of any two terms equal to the minimum of the remaining two terms.

Conjectures from _Colin Barker_, Aug 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009287 | OEIS-A250404

Numbers k such that the set of all distinct values of phi of all divisors of k equals the set of all proper divisors of k+1 where phi is the Euler totient function (A000010).

Conjecture: last term is 4294967295.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009288 | OEIS-A250405

Numbers k such that all values of Euler phi (A000010) of all divisors of k are pairwise distinct and represent all proper divisors of k+1.

Conjecture: last term is 4294967295.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009289 | OEIS-A250461

Number of (n+1)X(1+1) 0..1 arrays with nondecreasing min(x(i,j),x(i,j-1)) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009290 | OEIS-A250610

Number of (n+1) X (7+1) 0..1 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009291 | OEIS-A250658

Number of (4+1) X (n+1) 0..1 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009292 | OEIS-A250659

Number of (5+1) X (n+1) 0..1 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009293 | OEIS-A250660

Number of (6+1) X (n+1) 0..1 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009294 | OEIS-A250661

Number of (7+1) X (n+1) 0..1 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009295 | OEIS-A250756

Number of (1+1) X (n+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing x(i,j)+x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009296 | OEIS-A250757

Number of (2+1) X (n+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing x(i,j)+x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009297 | OEIS-A250758

Number of (3+1) X (n+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing x(i,j)+x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009298 | OEIS-A250759

Number of (4+1) X (n+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing x(i,j)+x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009299 | OEIS-A250760

Number of (5+1) X (n+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing x(i,j)+x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009300 | OEIS-A250761

Number of (6+1) X (n+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing x(i,j)+x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009301 | OEIS-A250762

Number of (7+1) X (n+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing x(i,j)+x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 18 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009302 | OEIS-A250778

Number of (n+1) X (3+1) 0..1 arrays with nondecreasing x(i,j)+x(i,j-1) in the i direction and nondecreasing absolute value of x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009303 | OEIS-A250779

Number of (n+1) X (4+1) 0..1 arrays with nondecreasing x(i,j)+x(i,j-1) in the i direction and nondecreasing absolute value of x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009304 | OEIS-A250780

Number of (n+1) X (5+1) 0..1 arrays with nondecreasing x(i,j)+x(i,j-1) in the i direction and nondecreasing absolute value of x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009305 | OEIS-A250781

Number of (n+1) X (6+1) 0..1 arrays with nondecreasing x(i,j)+x(i,j-1) in the i direction and nondecreasing absolute value of x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009306 | OEIS-A250782

Number of (n+1) X (7+1) 0..1 arrays with nondecreasing x(i,j)+x(i,j-1) in the i direction and nondecreasing absolute value of x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009307 | OEIS-A250784

Number of (2+1) X (n+1) 0..1 arrays with nondecreasing x(i,j)+x(i,j-1) in the i direction and nondecreasing absolute value of x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009308 | OEIS-A250791

Number of (n+1) X (2+1) 0..1 arrays with nondecreasing min(x(i,j),x(i,j-1)) in the i direction and nondecreasing absolute value of x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009309 | OEIS-A250792

Number of (n+1) X (3+1) 0..1 arrays with nondecreasing min(x(i,j),x(i,j-1)) in the i direction and nondecreasing absolute value of x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009310 | OEIS-A250813

Number of (1+1) X (n+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009311 | OEIS-A250814

Number of (2+1) X (n+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009312 | OEIS-A250815

Number of (3+1) X (n+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009313 | OEIS-A250816

Number of (4+1) X (n+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009314 | OEIS-A250817

Number of (5+1) X (n+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009315 | OEIS-A250818

Number of (6+1) X (n+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009316 | OEIS-A250819

Number of (7+1) X (n+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009317 | OEIS-A250846

Number of (n+1) X (1+1) 0..3 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

G.f.: x*(100 - 457*x + 740*x^2 - 384*x^3) / ((1 - x)*(1 - 2*x)*(1 - 3*x)*(1 - 4*x)) (conjectured). - _Colin Barker_, Jan 18 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009318 | OEIS-A250854

Number of (2+1) X (n+1) 0..3 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009319 | OEIS-A250855

Number of (3+1) X (n+1) 0..3 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009320 | OEIS-A250856

Number of (4+1) X (n+1) 0..3 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009321 | OEIS-A250857

Number of (5+1) X (n+1) 0..3 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009322 | OEIS-A250858

Number of (6+1) X (n+1) 0..3 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009323 | OEIS-A250859

Number of (7+1) X (n+1) 0..3 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing min(x(i,j),x(i-1,j)) in the j direction.

Conjectures from _Colin Barker_, Nov 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009324 | OEIS-A250878

Number of (1+1) X (n+1) 0..3 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing x(i,j)+x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009325 | OEIS-A250879

Number of (2+1) X (n+1) 0..3 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing x(i,j)+x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009326 | OEIS-A250880

Number of (3+1) X (n+1) 0..3 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing x(i,j)+x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009327 | OEIS-A250881

Number of (4+1) X (n+1) 0..3 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing x(i,j)+x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009328 | OEIS-A250882

Number of (5+1) X (n+1) 0..3 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing x(i,j)+x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009329 | OEIS-A250883

Number of (6+1) X (n+1) 0..3 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing x(i,j)+x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009330 | OEIS-A250892

Number of (n+1) X (2+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing absolute value of x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009331 | OEIS-A250893

Number of (n+1) X (3+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing absolute value of x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009332 | OEIS-A250894

Number of (n+1) X (4+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing absolute value of x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009333 | OEIS-A250895

Number of (n+1) X (5+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing absolute value of x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009334 | OEIS-A250896

Number of (n+1) X (6+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing absolute value of x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009335 | OEIS-A250897

Number of (n+1) X (7+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing absolute value of x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009336 | OEIS-A250899

Number of (1+1) X (n+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing absolute value of x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009337 | OEIS-A250900

Number of (2+1) X (n+1) 0..2 arrays with nondecreasing x(i,j)-x(i,j-1) in the i direction and nondecreasing absolute value of x(i,j)-x(i-1,j) in the j direction.

Conjectures from _Colin Barker_, Nov 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009338 | OEIS-A250917

Expansion of e.g.f. exp(x*C(x)^3) where C(x) = (1 - sqrt(1-4*x))/(2*x) is the g.f. of the Catalan numbers, A000108.

Conjecture D-finite with recurrence: +2*a(n) +(-11*n+20)*a(n-1) +(n^3+9*n^2-116*n+164)*a(n-2) +(-4*n^4+35*n^3+n^2-317*n+342)*a(n-3) -6*(n-3)*(6*n^3-50*n^2+147*n-176)*a(n-4) +12*(n-5)*(2*n-9)*(n-3)*(n-4)*a(n-5)=0. - _R. J. Mathar_, Jan 25 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009339 | OEIS-A251130

Number of (n+1) X (1+1) 0..2 arrays with no 2 X 2 subblock having the sum of its diagonal elements greater than the maximum of its antidiagonal elements.

Conjectures from _Colin Barker_, Nov 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009340 | OEIS-A251131

Number of (n+1) X (2+1) 0..2 arrays with no 2 X 2 subblock having the sum of its diagonal elements greater than the maximum of its antidiagonal elements.

Conjectures from _Colin Barker_, Nov 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009341 | OEIS-A251132

Number of (n+1) X (3+1) 0..2 arrays with no 2 X 2 subblock having the sum of its diagonal elements greater than the maximum of its antidiagonal elements.

Conjectures from _Colin Barker_, Nov 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009342 | OEIS-A251133

Number of (n+1) X (4+1) 0..2 arrays with no 2 X 2 subblock having the sum of its diagonal elements greater than the maximum of its antidiagonal elements.

Conjectures from _Colin Barker_, Nov 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009343 | OEIS-A251134

Number of (n+1) X (5+1) 0..2 arrays with no 2 X 2 subblock having the sum of its diagonal elements greater than the maximum of its antidiagonal elements.

Conjectures from _Colin Barker_, Nov 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009344 | OEIS-A251135

Number of (n+1) X (6+1) 0..2 arrays with no 2 X 2 subblock having the sum of its diagonal elements greater than the maximum of its antidiagonal elements.

Conjectures from _Colin Barker_, Nov 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009345 | OEIS-A251136

Number of (n+1) X (7+1) 0..2 arrays with no 2 X 2 subblock having the sum of its diagonal elements greater than the maximum of its antidiagonal elements.

Conjectures from _Colin Barker_, Nov 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 30 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009346 | OEIS-A251239

Indices of prime numbers in A098550.

It is conjectured that every prime appears in A098550, and if so then A098550(a(n)) = A000040(n). [Comment edited by _N. J. A. Sloane_, Dec 15 2014] [It is now known that every prime appears in A098550, although it is not known that they appear in their right order. - _N. J. A. Sloane_, Dec 25 2014] Conjecture: a(n) = A251541(n) + 2 for n > 4. - _Reinhard Zumkeller_, Dec 16 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Dec 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009347 | OEIS-A251269

Number of (2+1) X (n+1) 0..1 arrays with no 2 X 2 subblock having x11-x00 less than x10-x01.

Conjectures from _Colin Barker_, Nov 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009348 | OEIS-A251270

Number of (3+1) X (n+1) 0..1 arrays with no 2 X 2 subblock having x11-x00 less than x10-x01.

Conjectures from _Colin Barker_, Nov 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009349 | OEIS-A251271

Number of (4+1) X (n+1) 0..1 arrays with no 2 X 2 subblock having x11-x00 less than x10-x01.

Conjectures from _Colin Barker_, Nov 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009350 | OEIS-A251276

Number of (n+1) X (1+1) 0..3 arrays with no 2 X 2 subblock having its maximum diagonal element less than its minimum antidiagonal element.

Conjectures from _Colin Barker_, Nov 27 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009351 | OEIS-A251293

Number of (n+1) X (1+1) 0..3 arrays with every 2 X 2 subblock summing to 3 6 or 9.

Conjectures from _Colin Barker_, Nov 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009352 | OEIS-A251328

Number of (n+1) X (1+1) 0..2 arrays with every 2 X 2 subblock summing to 2 4 or 6.

Conjectures from _Colin Barker_, Nov 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009353 | OEIS-A251360

Numbers k such that k is the concatenation of prime factors of pi(k), in increasing order.

Conjecture: numbers k such that k = A084317(A000720(k)). - _Chai Wah Wu_, Apr 04 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jahangeer Kholdi_, Dec 01 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009354 | OEIS-A251411

Numbers k such that A098550(k) = k.

There is a strong conjecture that there are no further terms. See the discussion in the comments in A098550.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009355 | OEIS-A251413

a(n) = 2n-1 if n <= 3, otherwise the smallest odd number not occurring earlier having at least one common factor with a(n-2), but none with a(n-1).

Conjecture 1: Sequence is a permutation of the odd numbers. [True, see p. 4 in Applegate et al. link. - _Sela Fried_, Oct 09 2025] Conjecture 2: The odd primes occur in the sequence in their natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009356 | OEIS-A251414

a(n) = (A251413(n) + 1)/2.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009357 | OEIS-A251416

a(n) = smallest number not in {A098550(1), A098550(2), ..., A098550(n)}.

Bradley Klee conjectures that after a(30)=18, all further terms are primes, that every prime appears, and the primes appear in increasing order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009358 | OEIS-A251417

Lengths of runs of identical terms in A251416.

It appears (conjecture) that a(n)>1 for n>18. - _Alexander R. Povolotsky_, Dec 07 2014 Conjecture: a(n) = A247253(n-5) for n>12. - _Reinhard Zumkeller_, Dec 07 2014 The previous conjecture is equivalent to the statement that A251416(n) lists all primes and only primes after a(30)=18. - _M. F. Hasler_, Dec 08 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 03 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009359 | OEIS-A251418

Floor((n^2+7n-23)/14).

This is conjectured to be the value of the dominance number of the triangle grid graph for n >= 14 - see A251419.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009360 | OEIS-A251419

Domination number of the n-triangle grid graph TG_n having n vertices along each side.

Conjectured to equal floor((n^2 + 7n - 23)/14) for n >= 14. See A251418. G.f.: (x^22 - x^21 - x^19 + 2*x^18 - x^17 - x^14 + 2*x^13 - 2*x^11 + 2*x^10 - 2*x^9 + x^8 + x^7 - 2*x^6 + x^5 - x^3 + x^2 - x)/(x^9 - 2*x^8 + x^7 - x^2 + 2*x - 1) (conjectured, equivalent to Wagon's conjectural formula from comments). - _Andy Huchala_, Mar 15 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 04 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009361 | OEIS-A251421

Number of length n+2 0..1 arrays with the sum of the maximum minus twice the median plus the minimum of adjacent triples multiplied by some arrangement of +-1 equal to zero.

Conjectures from _Colin Barker_, Mar 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 02 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009362 | OEIS-A251541

Index of first term in A098550 that is divisible by the n-th prime.

Conjectures. 1: Equals first column of A251716. 2: For n > 4, a(n) = A251239(n) - 2. - _Reinhard Zumkeller_, Dec 15 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009363 | OEIS-A251556

Conjectured values of n which are the minimal cycle representatives of finite cycles in the permutation of the natural numbers defined by A098550.

Conjectured values of n which are the minimal cycle representatives of finite cycles in the permutation of the natural numbers defined by A098550.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009364 | OEIS-A251581

G.f. satisfies: A(x) = A(x^2 + 2*x^3)/(1-x).

Conjecture: limit of a(n+1)/a(n) = 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 31 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009365 | OEIS-A251582

G.f. satisfies: A(x) = A(2*x^2 + x^3)/(1-x).

Conjecture: limit of a(n+1)/a(n) = sqrt(2) + 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 31 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009366 | OEIS-A251595

Distinct terms in A251416.

a(n) = prime(n-4) for n > 11 according to Bradley Klee's conjecture, empirically confirmed for the first 10000 primes;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Dec 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009367 | OEIS-A251598

Dimension of space of invariant tensors in 2n-th tensor power of natural representation of Sp(8).

0=256*(2*n-1)*(n-1)*(n+3)*(2*n-3)*a(n-2)-4*(2*n-1)*(10*n^3+137*n^2+521*n+432)*a(n-1)+(n+4)*(n+7)*(n+9)*(n+10)*a(n), (conjectured). - _Michael D. Weiner_, Apr 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bruce Westbury_, Dec 05 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009368 | OEIS-A251604

A Zumkeller-type sequence (cf. A098550): a(n) = n if n <= 3, otherwise the smallest number not occurring earlier having at least one common factor with a(n-2)+a(n-1), but none with a(n-1).

Conjectured to be a permutation of the positive integers. See also A255972 for this conjecture. - _Reinhard Zumkeller_, Mar 12 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Dec 13 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009369 | OEIS-A251619

Smallest prime factor of smallest multiple of prime(n) in A098550.

conjecture: for n > 2: a(n) = 2 or a(n) = 3; conjecture: for n > 25: a(n) = 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Dec 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009370 | OEIS-A251621

Run lengths in A249943.

Connection with prime gaps: conjecturally, for n>=13, we have a(n) = A001223(n-5). - _Vladimir Shevelev_, Dec 07 2014 _Bradley Klee_ noted that this conjecture and his conjectures in A251416 are equivalent. At least to one side, our conjecture could be deduced from Klee's conjectures by a simple induction. - _Vladimir Shevelev_, Dec 10 2014 As a corollary, we have an explicit conjectural formula for prime(n), n>=8, essentially based on A098550: prime(n) = 19 + sum{i=9,...,n}a(i+4). - _Vladimir Shevelev_, Dec 11 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Dec 06 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009371 | OEIS-A251637

Square array read by antidiagonals containing in row n the multiples of prime(n) in A098550 in order of appearance.

conjecture: A098550 is a permutation of the positive integers iff A001221(n) = number of rows containing n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Dec 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009372 | OEIS-A251716

Array of positions of A251637(n,k) in A098550.

conjecture: A098550 is a permutation of the positive integers iff all integers > 1 occur exactly once.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Dec 09 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009373 | OEIS-A251743

Pairs of nodes in a complete binary tree that are at an absolute height difference of less than 2 from each other.

Conjectures from _Colin Barker_, Dec 09 2014: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dhruv Matani_, Dec 07 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009374 | OEIS-A251758

Let n>=2 be a positive integer with divisors 1 = d_1 < d_2 < ... < d_k = n, and s = d_1*d_2 + d_2*d_3 + ... + d_(k-1)*d_k. The sequence lists the values a(n) = floor(n^2/s).

Conjecture: Terms x, where a(x)=n, x=p#k/p#j, p#i is the i-th primorial, k>j is suitable large k and j is the number of primes less than n. As an example, n=9, x = p#7/p#4 = 2431. For n=10, x = p#6/p#4 = 143 although 121 = 11^2 is the least x where a(x)=10 (see formula section). For n=8, x = p#12/p#4, p#13/p#4, p#14/p#4, p#15/p#4, p#16/p#4, etc. But is p#12/p#4 the least such x? - _Robert G. Wilson v_, Dec 18 2014 Adamczewski includes proof abstract regarding Wilson's conjecture in linked article. - _Michael De Vlieger_, Aug 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Dec 08 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009375 | OEIS-A251936

Number of length 2+2 0..n arrays with the sum of the maximum minus the median of adjacent triples multiplied by some arrangement of +-1 equal to zero.

Conjectures from _Colin Barker_, Dec 01 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009376 | OEIS-A251964

For a prime p, denote by s(p,k) the odd part of the digital sum of p^k. Let k_1 be the smallest k such that s(p,k) is divisible by 5. Sequence lists primes p for which s(p,k_1)=5.

Let p be a prime other than 3. If p is not in the sequence, then either s(p,k_1) >= 25 or k_1 does not exist. We conjecture that k_1=k_1(p) exists for every prime p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Dec 11 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009377 | OEIS-A252178

Number of length 2+2 0..n arrays with the sum of the maximum minus the minimum of adjacent triples multiplied by some arrangement of +-1 equal to zero.

Conjectures from _Colin Barker_, Dec 01 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 15 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009378 | OEIS-A252283

Intersection of A251964, A252280 and A252281.

A weaker conjecture: {2,5} is the intersection of all such sequences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Dec 16 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009379 | OEIS-A252354

Number of Motzkin paths of length n with no level steps at height 2.

Conjecture: (-n+3)*a(n) +3*(2*n-7)*a(n-1) +(-7*n+24)*a(n-2) +2*(-7*n+36)*a(n-3) +2*(11*n-51)*a(n-4) +3*(3*n-23)*a(n-5) +(-10*n+63)*a(n-6) +3*(n-6)*a(n-7)=0. - _R. J. Mathar_, Sep 24 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José Luis Ramírez Ramírez_, Apr 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009380 | OEIS-A252482

Exponents n such that the decimal expansion of the power 12^n contains no zeros.

Conjectured to be finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Dec 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009381 | OEIS-A252486

Smallest k such that n^6 = a_1^6+...+a_k^6 where all the a_i are positive integers less than n.

I conjecture that the sequence is bounded by the initial term. Probably even a(3)=36, a(5)=29, a(6)=22 and some more are followed only by smaller terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Dec 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009382 | OEIS-A252487

Smallest k such that n^7 = a_1^7 + ... + a_k^7 and all a_i are positive integers less than n.

I conjecture that the sequence is bounded by the initial term a(2)=128. Probably even a(4)=66, a(5)=39, a(6)=28 and some more are followed only by smaller terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Dec 17 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009383 | OEIS-A252491

a(n) = (10^(n^2) - 1)/(10^n - 1).

This sequence is relevant for counterexamples to a conjecture in A086766: If p is prime and a(p) is not prime, then A086766(10^(p-1)) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009384 | OEIS-A252503

Smallest prime p such that Phi_n(p) is also prime, where Phi is the cyclotomic polynomial, or 0 if no such p exists.

Conjecture: a(n) > 0 for all n != 2^k (k>5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, Dec 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009385 | OEIS-A252505

Number of biquadratefree (4th power free) divisors of n.

a(n) = Sum_{d|n} A018892(d)*A378214(n/d). [Conjectured]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Critzer_, Mar 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009386 | OEIS-A252593

Number of ways to place 8 nonattacking queens on an n X n board.

Conjectured recurrence order is 477 (see "Non-attacking chess pieces", p. 19). - _Vaclav Kotesovec_, Dec 19 2014

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antal Pinter_, Dec 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009387 | OEIS-A252594

Records in A072994.

Conjecture: f(x)/x -> 5/12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Dec 18 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009388 | OEIS-A252666

Let k be the smallest number such that s(k) = odd part of digital sum of 2^k is multiple of prime(n); then a(n)=k, if s(k) = prime(n). Otherwise, or if there is no such k, a(n)=0.

We conjecture that k in the definition exists for every n>=3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Dec 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009389 | OEIS-A252668

Let k be the smallest number such that s(k) = odd part of digital sum of 5^k is a multiple of prime(n); then a(n)=k, if s(k) = prime(n). Otherwise, or if there is no such k, a(n)=0.

We conjecture that k in the definition exists for every n>=3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Dec 20 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009390 | OEIS-A252739

a(n) = A252738(n) / n.

Note how 6, 720 and 612360000 occur in A244743 as its 0th, 4th and 8th term, from which my bold conjecture that A244743(12) or A244743(16) = 1697781042840960000000000. According to preliminary results from _Janis Iraids_, the value of A005245(a(5)) = ||1697781042840960000000000|| = 160, while ||1697781042840960000000000 - 1|| = 169, which lays to rest my naive conjecture above, as 169 - 160 is neither 12 nor 16. Note also how 5, 719 and 612359999 are all primes, while a(5)-1 factorizes as 1697781042840959999999999 = 13 * 89443 * 908669 * 1606890407869. - _Antti Karttunen_, Dec 20 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009391 | OEIS-A252782

a(n) = n-th term of Euler transform of n-th powers.

Conjecture: limit n->infinity a(n)^(1/n^2) = exp(exp(-1)) = 1.444667861... . - _Vaclav Kotesovec_, Mar 25 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Dec 21 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009392 | OEIS-A252812

Primes whose trajectories under the map x -> A039951(x) enter the cycle {83, 4871} (conjectured).

Primes whose trajectories under the map x -> A039951(x) enter the cycle {83, 4871} (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Dec 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009393 | OEIS-A252814

Number of n X 2 nonnegative integer arrays with upper left 0 and every value within 2 of its city block distance from the upper left and every value increasing by 0 or 1 with every step right or down.

Conjectures from _Colin Barker_, Dec 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009394 | OEIS-A252815

Number of n X 3 nonnegative integer arrays with upper left 0 and every value within 2 of its city block distance from the upper left and every value increasing by 0 or 1 with every step right or down.

Conjectures from _Colin Barker_, Dec 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009395 | OEIS-A252816

Number of n X 4 nonnegative integer arrays with upper left 0 and every value within 2 of its city block distance from the upper left and every value increasing by 0 or 1 with every step right or down.

Conjectures from _Colin Barker_, Dec 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009396 | OEIS-A252817

Number of n X 5 nonnegative integer arrays with upper left 0 and every value within 2 of its city block distance from the upper left and every value increasing by 0 or 1 with every step right or down.

Conjectures from _Colin Barker_, Dec 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009397 | OEIS-A252822

Number of n X 2 nonnegative integer arrays with upper left 0 and every value within 3 of its city block distance from the upper left and every value increasing by 0 or 1 with every step right or down.

Conjectures from _Colin Barker_, Dec 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009398 | OEIS-A252823

Number of n X 3 nonnegative integer arrays with upper left 0 and every value within 3 of its city block distance from the upper left and every value increasing by 0 or 1 with every step right or down.

Conjectures from _Colin Barker_, Dec 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009399 | OEIS-A252831

Number of n X 3 nonnegative integer arrays with upper left 0 and every value within 2 of its king move distance from the upper left and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009400 | OEIS-A252832

Number of n X 4 nonnegative integer arrays with upper left 0 and every value within 2 of its king move distance from the upper left and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009401 | OEIS-A252833

Number of n X 5 nonnegative integer arrays with upper left 0 and every value within 2 of its king move distance from the upper left and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009402 | OEIS-A252834

Number of n X 6 nonnegative integer arrays with upper left 0 and every value within 2 of its king move distance from the upper left and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009403 | OEIS-A252837

Let c = A098550. Let c(n) be prime. If c(n+1) is even, then a(n) is the maximal number r of consecutive even numbers of the form c(n+1), c(n+3), ..., c(n+(2r-1)); if c(n+1) is odd, then a(n)=0.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Dec 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009404 | OEIS-A252838

Let A098550(n) be prime. If A098550(n+1) is even, then suppose A098550(n+3), A098550(n+5), ..., A098550(n+(2*r-1)) are also even, while A098550(n+(2*r+1)) is odd. Then a(n)= A098550(n+(2*r-1))/2; if A098550(n+1) is odd, then a(n)=0.

If conjecture in A252837 is true, then for n>=25, a(n) = prime(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Dec 22 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009405 | OEIS-A252864

Number of pairs in generation n of the tree T defined in Comments.

Conjecture: |g(n)| = |g(n-1)| + |g(n-3)| for n >= 12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 31 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009406 | OEIS-A252865

a(n) = n if n <= 3, otherwise the smallest squarefree number not occurring earlier having at least one common factor with a(n-2), but none with a(n-1).

The sequence has consecutive primes at indices 2 (2 & 3), 10 (7 & 11), 38 (29 & 31), 50 (41 & 43), and 56 (47 & 53). We conjecture that there are no further such pairs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Dec 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009407 | OEIS-A252867

a(n) = n if n <= 2, otherwise the smallest number not occurring earlier having in its binary representation at least one bit in common with a(n-2), but none with a(n-1).

Conjectured to be a permutation of the nonnegative integers. [Comment modified by _N. J. A. Sloane_, Jan 10 2015]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Dec 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009408 | OEIS-A252868

Squarefree version of A252867.

This is a permutation of the squarefree numbers. [I believe this is at present only a conjecture. - _N. J. A. Sloane_, Jan 10 2015]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Dec 23 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009409 | OEIS-A252932

Number of n X 2 nonnegative integer arrays with upper left 0 and every value within 3 of its king move distance from the upper left and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009410 | OEIS-A252933

Number of n X 3 nonnegative integer arrays with upper left 0 and every value within 3 of its king move distance from the upper left and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009411 | OEIS-A252934

Number of n X 4 nonnegative integer arrays with upper left 0 and every value within 3 of its king move distance from the upper left and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009412 | OEIS-A252935

Number of n X 5 nonnegative integer arrays with upper left 0 and every value within 3 of its king move distance from the upper left and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 24 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009413 | OEIS-A252978

Number of n X 3 nonnegative integer arrays with upper left 0 and lower right its king-move distance away minus 2 and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009414 | OEIS-A252979

Number of n X 4 nonnegative integer arrays with upper left 0 and lower right its king-move distance away minus 2 and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009415 | OEIS-A252980

Number of n X 5 nonnegative integer arrays with upper left 0 and lower right its king-move distance away minus 2 and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Mar 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009416 | OEIS-A252984

a(0)=4. a(n) is the smallest number not in the sequence such that Sum_{k=1..n} a(k) divides Product_{k=1..n} a(k).

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Jan 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009417 | OEIS-A252999

Number of n X 3 nonnegative integer arrays with upper left 0 and lower right its king-move distance away minus 3 and every value within 3 of its king move distance from the upper left and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009418 | OEIS-A253000

Number of n X 4 nonnegative integer arrays with upper left 0 and lower right its king-move distance away minus 3 and every value within 3 of its king move distance from the upper left and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009419 | OEIS-A253001

Number of n X 5 nonnegative integer arrays with upper left 0 and lower right its king-move distance away minus 3 and every value within 3 of its king move distance from the upper left and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009420 | OEIS-A253006

Number of n X 3 nonnegative integer arrays with upper left 0 and lower right its king-move distance away minus 3 and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009421 | OEIS-A253007

Number of n X 4 nonnegative integer arrays with upper left 0 and lower right its king-move distance away minus 3 and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 25 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009422 | OEIS-A253029

Number of (n+2) X (1+2) 0..2 arrays with every consecutive three elements in every row and column having exactly two distinct values, and new values 0 upwards introduced in row major order.

Conjectures from _Colin Barker_, Dec 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 26 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009423 | OEIS-A253049

Numbers in A098550 which appear immediately after a prime.

It is conjectured that this is the same as A253048 except for the addition of 3, 13, 31, and 43 (the upper members of the twin primes in A098550).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009424 | OEIS-A253062

Largest order of a rooted tree that does not contain a rooted caterpillar subtree of order n.

Conjectures from _Colin Barker_, Feb 21 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009425 | OEIS-A253073

Lexicographically earliest sequence of distinct numbers such that neither a(n) nor a(n-1)+a(n) is prime.

Conjecture: this is a permutation of the nonprimes. [Proof outline given below by Semeon Artamonov and Pat Devlin.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 01 2015, based on a suggestion from _Patrick Devlin_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009426 | OEIS-A253074

Lexicographically earliest sequence of distinct numbers such that a(n-1)+a(n) is not prime.

Conjecture: this is a permutation of the nonnegative numbers. [Proof outline below due to Patrick Devlin and Semeon Artamonov]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 01 2015, based on a suggestion from _Patrick Devlin_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009427 | OEIS-A253122

Number of length n+2 0..1 arrays with the sum of medians of adjacent triples multiplied by some arrangement of +-1 equal to zero.

Conjectures from _Colin Barker_, Dec 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009428 | OEIS-A253130

Number of length 2+2 0..n arrays with the sum of medians of adjacent triples multiplied by some arrangement of +-1 equal to zero.

Conjectures from _Colin Barker_, Dec 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 27 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009429 | OEIS-A253138

Number of ways to represent the n-th prime as the arithmetic mean of two semiprimes.

Conjecture: a(n)>0 for n>5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Mar 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009430 | OEIS-A253174

For a permutation of the positive integers B={b(k)}, denote L_B(n) = Sum_{k=1..n} lambda(b(k)), where lambda(n) is Liouville's function (A008836). In this sequence, a(n) = L_B(n) in case B = A098550.

In case B=A000027 (the natural numbers), G. Polya (1919) conjectured that L_B(n)<=0, for n>=2. But this was disproved in 1958 by B. Haselgrove, and in 1980 M. Tanaka found the smallest counterexample, 906150257. However, for this sequence we conjecture that a(n)>=0 for all n other than 3. A reason for our conjecture is the later appearance of primes in A098550 than in A000027. By our conjecture, among the first N terms of A098550, the terms with odd s are never in the majority, if N is other than 3. _Peter J. C. Moses_ verified the conjecture up to 2.5*10^5 and, moreover, in this range a(n)>0 for n>6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009431 | OEIS-A253176

Least k>=0 such that both 3n*2^k+1 and 3n*2^k-1 are primes, or -1 if no such k exists.

Conjecture: if n < 79, then a(n) >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, Mar 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009432 | OEIS-A253179

a(n) = mu(n)*Sum_{k=1..n} (n/k) where mu(n) is Möbius (or Moebius) function and (x/y) is Kronecker's symbol.

Conjecture: the partial sum of this sequence changes sign infinitely often.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Mar 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009433 | OEIS-A253187

Number of ordered ways to write n as the sum of a pentagonal number, a second pentagonal number and a generalized decagonal number.

Conjecture: a(n) > 0 for all n. Also, for any ordered pair (k,m) among (5,7), (5,9), (5,13), (6,5), (6,7), (7,5), each nonnegative integer n can be written as the sum of a k-gonal number, a second k-gonal number and a generalized m-gonal number. See also the author's similar conjectures in A254574, A254631, A255916 and the two linked papers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009434 | OEIS-A253202

Number of essentially different ways of arranging the numbers 1 through 2n around a circle so that the sum of each pair of adjacent numbers is semiprime.

Conjecture: a(n) > 0 for all n > 3. This implies the semiprime conjecture, and it is similar to the prime circle problem mentioned in A051252.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Mar 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009435 | OEIS-A253217

Number of n X n nonnegative integer arrays with upper left 0 and lower right its king-move distance away minus 2 and every value within 2 of its king move distance from the upper left and every value increasing by 0 or 1 with every step right, diagonally se or down.

Recurrence: 32*(1 + n)*(1 + 2*n)^2*(161046 + 465785*n + 551943*n^2 + 343020*n^3 + 117954*n^4 + 21285*n^5 + 1575*n^6)*a(n) - 8*(4443102 + 33718283*n + 105734340*n^2 + 180574335*n^3 + 186866686*n^4 + 122556360*n^5 + 51280818*n^6 + 13267683*n^7 + 1933470*n^8 + 121275*n^9)*a(n+1) + 2*(12137328 + 91378536*n + 283626704*n^2 + 478464380*n^3 + 488415476*n^4 + 315713355*n^5 + 130145646*n^6 + 33170868*n^7 + 4763070*n^8 + 294525*n^9)*a(n+2) + (10688508 + 80866406*n + 252913504*n^2 + 431097970*n^3 + 445804136*n^4 + 292620525*n^5 + 122735586*n^6 + 31877118*n^7 + 4668570*n^8 + 294525*n^9)*a(n+3) - (4877748 + 36871922*n + 114948300*n^2 + 194784258*n^3 + 199650088*n^4 + 129484209*n^5 + 53503836*n^6 + 13655808*n^7 + 1961820*n^8 + 121275*n^9)*a(n+4) + 2*(3 + n)^2*(7 + 2*n)*(2428 + 16118*n + 41382*n^2 + 52554*n^3 + 35154*n^4 + 11835*n^5 + 1575*n^6)*a(n+5) = 0. - conjectured by _Manuel Kauers_ and _Christoph Koutschan_, Mar 02 2023; proved by _Robert Dougherty-Bliss_ and _Manuel Kauers_ Conjecture: a(n) ~ 2^(4*n - 2) / (81 * Pi * n), based on the above recurrence - _Vaclav Kotesovec_, Mar 02 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009436 | OEIS-A253218

Number of n X 3 nonnegative integer arrays with upper left 0 and lower right its king-move distance away minus 2 and every value within 2 of its king move distance from the upper left and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009437 | OEIS-A253219

Number of n X 4 nonnegative integer arrays with upper left 0 and lower right its king-move distance away minus 2 and every value within 2 of its king move distance from the upper left and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009438 | OEIS-A253220

Number of n X 5 nonnegative integer arrays with upper left 0 and lower right its king-move distance away minus 2 and every value within 2 of its king move distance from the upper left and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009439 | OEIS-A253221

Number of n X 6 nonnegative integer arrays with upper left 0 and lower right its king-move distance away minus 2 and every value within 2 of its king move distance from the upper left and every value increasing by 0 or 1 with every step right, diagonally se or down.

Conjectures from _Colin Barker_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009440 | OEIS-A253225

Number of (n+1) X (2+1) 0..1 arrays with every 2 X 2 subblock diagonal minimum minus antidiagonal minimum nondecreasing horizontally and diagonal maximum minus antidiagonal maximum nondecreasing vertically.

Conjectures from _Colin Barker_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009441 | OEIS-A253226

Number of (n+1) X (3+1) 0..1 arrays with every 2 X 2 subblock diagonal minimum minus antidiagonal minimum nondecreasing horizontally and diagonal maximum minus antidiagonal maximum nondecreasing vertically.

Conjectures from _Colin Barker_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009442 | OEIS-A253227

Number of (n+1) X (4+1) 0..1 arrays with every 2 X 2 subblock diagonal minimum minus antidiagonal minimum nondecreasing horizontally and diagonal maximum minus antidiagonal maximum nondecreasing vertically.

Conjectures from _Colin Barker_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009443 | OEIS-A253228

Number of (n+1) X (5+1) 0..1 arrays with every 2 X 2 subblock diagonal minimum minus antidiagonal minimum nondecreasing horizontally and diagonal maximum minus antidiagonal maximum nondecreasing vertically.

Conjectures from _Colin Barker_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009444 | OEIS-A253229

Number of (n+1) X (6+1) 0..1 arrays with every 2 X 2 subblock diagonal minimum minus antidiagonal minimum nondecreasing horizontally and diagonal maximum minus antidiagonal maximum nondecreasing vertically.

Conjectures from _Colin Barker_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009445 | OEIS-A253230

Number of (n+1) X (7+1) 0..1 arrays with every 2 X 2 subblock diagonal minimum minus antidiagonal minimum nondecreasing horizontally and diagonal maximum minus antidiagonal maximum nondecreasing vertically.

Conjectures from _Colin Barker_, Dec 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009446 | OEIS-A253233

Smallest even pseudoprime (>2n+1) in base 2n+1.

Conjecture: There are infinitely many even pseudoprimes in every odd base.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, May 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009447 | OEIS-A253237

Conjectured largest perfect power k such that k+n is also a perfect power, or 0 if no such k exists.

Conjectured largest perfect power k such that k+n is also a perfect power, or 0 if no such k exists. Only a(1) is proven, all other terms (even including a(2)) are only conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, Apr 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009448 | OEIS-A253238

Number of ways to write n as a sum of a perfect power (>1) and a prime.

In the past, I conjectured that a(n) > 0 for all n>24, but this is not true. My PARI program found that a(1549) = 0. I also asked which a(n) are 1. For example, 331 is a de Polignac number (A006285), so it cannot be written as 2^n+p with p prime, and 331-6^n must divisible by 5, 331-10^n must divisible by 3, ..., 331-18^2 = 331-324 = 7 is prime (and it is the only prime of the form 331-m^n, with m, n natural numbers, m>1, n>1), so a(331) = 1. Similarly, a(3319) = 1. Conjecture: a(n) > 1 for all n > 3319. This conjecture is not true: a(1771561) = 0. (See A119748) Another conjecture:

For every number m>=0, there is a number k such that a(n)>=m for all n>=k. Another conjecture: Except for k=2, first occurrence of k must be earlier then first occurrence of k+1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, May 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009449 | OEIS-A253241

The "Reverse and Add!" problem in base 12: sequence lists the final palindrome number for n, or -1 if no palindrome is ever reached. (Written in base 10.)

a(237) is conjectured to be -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, Apr 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009450 | OEIS-A253246

Pisano period of A006190 to mod prime(n).

If the generalized Wall's conjecture to A006190 is true, then we can calculate A175182(m) when m is a prime power since for any k>=1 : A175182(prime(n)^k)=a(n)*prime(n)^(k-1). For example: A175182(2^k)=3*2^(k-1)=A007283(k-1). In fact, the conjecture fails on p=241, and this is the only counterexample below 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, Apr 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009451 | OEIS-A253257

Least positive integer k such that prime(k*n) has the form p^2 - 2 with p prime, or 0 if no such k exists.

Conjecture: a(n) > 0 for all n > 0. This is stronger than the conjecture that there are infinitely many primes of the form p^2-2 with p prime. I also conjecture that for any positive integer n there is a positive integer k such that prime(k*n) has the form 2*p^2-1 (or 4*p^2+1, or p^2+p+1) with p prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009452 | OEIS-A253258

Square array read by antidiagonals, j>=1, k>=1: T(j,k) is the j-th number n such that the symmetric representation of sigma(n) has at least a part with maximum width k.

Column T(j,1), j>=1, forms A174905 and is a permutation of A357581. Numbers T(j,k), j>=1 and k>1, form A005279. Conjecture: Every column of the square array contains odd numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jul 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009453 | OEIS-A253260

Brazilian squares.

Conjecture: Let r(n) = (a(n) - 1)/(a(n) + 1); then Product_{n>=1} r(n) = (15/17) * (35/37) * (63/65) * (40/41) * (99/101) * (60/61) * (143/145) * (195/197) * ... = (150 * Pi) / (61 * sinh(Pi)) = 0.668923905.... - _Dimitris Valianatos_, Feb 27 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Apr 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009454 | OEIS-A253261

Odd Brazilian squares.

The previous comment conjectures the 1 and the 121 are the only difference with respect to A062532. - _R. J. Mathar_, Jul 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Apr 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009455 | OEIS-A253279

Arrange numbers in a clockwise spiral with initial terms a(1)=1, a(2)=2, a(4)=3, a(6)=4, a(8)=5; thereafter each number shares a factor with each of its four (N,S,E,W) neighbors.

Conjecture: The sequence is a permutation of the positive integers. - _Vladimir Shevelev_, May 06 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009456 | OEIS-A253288

Each term a(n) satisfies four properties: 1, divisible by all prime factors of n; 2, divisible by only the prime factors of n; 3, not equal to any of the terms a(1), a(2), ... a(n-1); 4, smallest number satisfying 1-3 if A005361(n) is even, or second smallest number satisfying 1-3 if A005361(n) is odd.

Conjecture: this permutation of positive integers is self-inverse. Compare with A358971. The principal distinction between this sequence and A358971 is that fixed points aside from A358971(1) = 1 are explicitly ruled out in the latter. - _Michael De Vlieger_, Dec 10 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009457 | OEIS-A253297

Primes p for which some multiple k*p > 2*p appears before p does in A098550.

It is conjectured (see A098550) that there are no more terms of the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Dec 29 2014 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009458 | OEIS-A253490

Number of (n+1) X (3+1) 0..2 arrays with every 2 X 2 subblock diagonal minus antidiagonal sum nondecreasing horizontally, vertically and ne-to-sw antidiagonally.

Conjectures from _Colin Barker_, Dec 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009459 | OEIS-A253491

Number of (n+1) X (4+1) 0..2 arrays with every 2 X 2 subblock diagonal minus antidiagonal sum nondecreasing horizontally, vertically and ne-to-sw antidiagonally.

Conjectures from _Colin Barker_, Dec 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009460 | OEIS-A253492

Number of (n+1) X (5+1) 0..2 arrays with every 2 X 2 subblock diagonal minus antidiagonal sum nondecreasing horizontally, vertically and ne-to-sw antidiagonally.

Conjectures from _Colin Barker_, Dec 15 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009461 | OEIS-A253493

Number of (n+1) X (6+1) 0..2 arrays with every 2 X 2 subblock diagonal minus antidiagonal sum nondecreasing horizontally, vertically and ne-to-sw antidiagonally.

Conjectures from _Colin Barker_, Dec 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009462 | OEIS-A253503

Number of (n+2) X (1+2) 0..1 arrays with every 2 X 2 and 3 X 3 subblock diagonal maximum minus antidiagonal minimum nondecreasing horizontally and vertically.

Conjectures from _Colin Barker_, Dec 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009463 | OEIS-A253504

Number of (n+2) X (2+2) 0..1 arrays with every 2 X 2 and 3X3 subblock diagonal maximum minus antidiagonal minimum nondecreasing horizontally and vertically.

Conjectures from _Colin Barker_, Dec 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009464 | OEIS-A253505

Number of (n+2) X (3+2) 0..1 arrays with every 2 X 2 and 3 X 3 subblock diagonal maximum minus antidiagonal minimum nondecreasing horizontally and vertically.

Conjectures from _Colin Barker_, Dec 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009465 | OEIS-A253506

Number of (n+2) X (4+2) 0..1 arrays with every 2 X 2 and 3 X 3 subblock diagonal maximum minus antidiagonal minimum nondecreasing horizontally and vertically.

Conjectures from _Colin Barker_, Dec 16 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009466 | OEIS-A253507

Number of (n+2) X (5+2) 0..1 arrays with every 2 X 2 and 3 X 3 subblock diagonal maximum minus antidiagonal minimum nondecreasing horizontally and vertically.

Conjectures from _Colin Barker_, Dec 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009467 | OEIS-A253547

Total number of star-shaped dodecagons appearing in a variant of hexagon expansion after n iterations.

Conjectures from _Colin Barker_, Jan 03 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jan 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009468 | OEIS-A253570

Maximum number of circles of radius 1 that can be packed into a regular n-gon with side length 2 (conjectured).

Maximum number of circles of radius 1 that can be packed into a regular n-gon with side length 2 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Jan 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009469 | OEIS-A253573

Rectangular array a(n,k) read by upwards antidiagonals: row A(n) is the result of applying the function defined in A098550 to the set comprising row n of A253572, for n >= 2.

Conjecture 1: Successive rows tend to A098550. Conjecture 2: The main diagonal is A098550. Conjecture 3: For each n, the primes prime(1), ..., prime(n) in row n appear in their natural order. Conjecture 4: a(n,A251239(j)) = A000040(j), for j = 1,...,n inclusive.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Jan 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009470 | OEIS-A253581

a(n) = A252867(n) AND A252867(n+2).

conjecture: all natural numbers occur, cf. A253603, giving smallest m such that A253581(m) = n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jan 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009471 | OEIS-A253609

Array read by upwards antidiagonals: A(n, k) = index of prime(k)^n in A098550.

Conjecture: For all natural numbers i, j, k, prime(k)^i precedes prime(k)^(i+1) and prime(k)^j precedes prime(k+1)^j in A098550.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Jan 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009472 | OEIS-A253628

a(n) = psi(n) mod n, where psi is the Dedekind psi function (A001615).

Indices of 1 appear to be given by primes A000040 (see conjecture in A068494). The (weaker) statement that a(prime(i)) = 1 is a direct consequence of the multiplicity of A001615.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tom Edgar_, Jan 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009473 | OEIS-A253631

Palindromic primes containing only the digits 0 and 1 such that their squares are palindromes.

Conjecture: a(n) = A225603(n+2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Jan 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009474 | OEIS-A253632

Primes p such that p+d, p+2d, p+4d, p+8d, p+16d, p+32d and p+64d are also primes for d = 30.

Conjecture: In the expression p + d*2^k for k = 0...6; 1 < p < 10^6, d = 30 is the smallest value of d to yield such sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _K. D. Bajpai_, Jan 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009475 | OEIS-A253644

Numbers n such that n^k is zeroless for k=0,...,5.

Conjectured to be finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009476 | OEIS-A253646

Primes p such that p^k is zeroless for k=1,...,6.

Primes in A253647; both sequences are conjectured to be finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_ and _M. F. Hasler_, Jan 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009477 | OEIS-A253647

Numbers n such that n^k is zeroless for k=0,...,6.

There are 55 terms below 10^7. Conjectured to be finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009478 | OEIS-A253686

a(n) is the smallest integer area of the triangle having the sides in the commutative ring Z[sqrt(q)] where q = A005117(n) is a squarefree number.

Conjecture: the set of squarefree numbers q such that the integer area A of the triangles with sides in the commutative ring Z[sqrt(q)] is finite if A < 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009479 | OEIS-A253687

The total number of pentagons in a variant of pentagon expansion (side-to-side, two consecutive sides and one isolated side) after n iterations.

Conjecture: a(n) = 2*a(n-1)-a(n-2)+a(n-4)-2*a(n-5)+a(n-6) for n>7. - _Colin Barker_, Jan 09 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jan 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009480 | OEIS-A253688

The total number of pentagons in a variant of pentagon expansion (vertex-to-vertex, two consecutive vertices and one isolated vertex) after n iterations.

Conjectures from _Colin Barker_, Jan 09 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jan 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009481 | OEIS-A253778

Numbers b that are the first of m^3 consecutive cubes whose sum is a cube c^3, where m^3 is not divisible by 3 (A118719).

Conjectures from _Colin Barker_, Jan 13 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Pletser_, Jan 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009482 | OEIS-A253779

Numbers c whose cubes are equal to the sum of m^3 consecutive cubes for m^3 not divisible by 3 (A118719).

Conjectures from _Colin Barker_, Jan 13 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Pletser_, Jan 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009483 | OEIS-A253809

Array of pairs (x,y) of Markoff triples (x,y,z) with x <= y <= z, for z given in A002559.

Frobenius' conjecture on Markoff triples is that the maximal member z of the triple of positive integers (x,y,z), satisfying x^2 + y^2 + z^2 - 3*x*y*z = 0, with x <= y <= z, determines x and y uniquely. Also, each entry from A002559 (Markoff numbers) is conjectured to appear as a maximal member z. If an entry A002559(n) should not appear as z then one puts z(n) = 0 and row n will be 0, 0. If this Frobenius conjecture is true then the row length of this array is always 2, and only positive numbers appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jan 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009484 | OEIS-A253895

Total number of octagons in two variants of an octagon expansion after n iterations: either "side-to-side" or "vertex-to-vertex", respectively.

Conjectures from _Colin Barker_, Jan 17 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jan 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009485 | OEIS-A253897

Numbers n such that the sequence A127644 with a(1) = n is conjectured to give a permutation of the natural numbers.

Numbers n such that the sequence A127644 with a(1) = n is conjectured to give a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Jan 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009486 | OEIS-A253900

a(n) is the number of squares of the form x^2 + x + n^2 for 0 <= x <= n^2.

Conjecture: the probability that a(n) is a power of 2 is such that 0.703 < p(a(n)=2^p) < 0.705.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009487 | OEIS-A253974

Number of 2n-move closed giraffe paths on an unbounded chessboard from a given square to the same square.

Conjecture: Number of 2n-move closed paths of leaper [r,s] on an unbounded chessboard, where 0 < r < s and gcd(r,s)=1, is asymptotic to 2^(6*n+1) / ((r^2+s^2)*Pi*n) if r+s is even, and 2^(6*n) / ((r^2+s^2)*Pi*n) if r+s is odd.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jan 31 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009488 | OEIS-A253975

a(n) is the smallest number k representable as x*y+x+y, where x>=y>1, in exactly n ways, or -1 if no such k exists.

Conjecture: no term is -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Jan 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009489 | OEIS-A254007

Cardinality of the set of equivalence classes of the set X_n of finite integer sequences {x_1 = 0, x_2, ..., x_n} satisfying |x_k - x_{k+1}| = 1, where two such sequences are deemed equivalent if they are permutations of each other.

G.f.: ((1-x)^3 * (1+x)^2 * (1+x-x^2)) / ((1-x-x^2) * (1-2*x^2)^2) (conjectured). - _David Radcliffe_, May 03 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John J. Harrison_, May 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009490 | OEIS-A254067

Rectangular array A read by upward antidiagonals in which the entry in row n and column k is defined by A(n,k) = S(4*A257499(n,k) - 3), n,k >= 1, where the function S is as defined in A257480.

Conjecture: Arranging the sequence in ascending order gives A189707 (positions of 0 in A189706).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, May 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009491 | OEIS-A254074

Numbers k such that the decimal expansions of both k and k^2 have 5 as the digit with the smallest value and 9 as the digit with the largest value.

The first digit of a term is either 7, 8 or 9 and the last digit is either 5, 6 or 7. Conjecture: no term is a multiple of 5. - _Chai Wah Wu_, Sep 10 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, May 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009492 | OEIS-A254077

a(n) = n if n <= 3, otherwise the smallest number not occurring earlier such that gcd(a(n),a(n-2)) > gcd(a(n),a(n-1)).

Conjecture: The sequence is infinite (that is, a(n) always exists). Conjecture: The sequence is a permutation of the natural numbers. Conjectures: 1) For k>=3, except for k=5, if a(n) = prime(k), then a(n-2) = 2*prime(k) and a(n+2) = 3*prime(k). This conjecture was verified by _Peter J. C. Moses_ for n <= 5000. - _Vladimir Shevelev_, Feb 09 2015. This conjecture verified for n <= 10^7. - _Ray Chandler_, Apr 02 2015. Extended to n <= 10^9. - _John Mason_, Jun 08 2016 2) For k>=3, except for k=4, if a(n)=prime(k)^2, then a(n-2) = prime(k)^2 + prime(k). This conjecture was verified by _Peter J. C. Moses_ for n <= 35000. - _Vladimir Shevelev_, Feb 12 2015. This conjecture verified for n <= 394349. - _Ray Chandler_, Mar 07 2015. This conjecture is false - for n = 4488245, a(n) = 2137^2, but a(n-2) = 2137^2 + 2*2137. - _Ray Chandler_, Mar 30 2015. Next are n = 30655601, a(n) = 5581^2, but a(n-2) = 5581^2 + 2*5581, and n = 922447261, a(n) = 30577^2, but a(n-2) = 30577^2 + 2*30577. - _John Mason_, Sep 15 2016 Theorem: If a(n-2) is prime AND a(n) EXISTS then a(n) is a multiple of a(n-2). For: by sequence definition, assuming a(n-2) = p, prime, then gcd(a(n),p) > gcd(a(n),a(n-1)); hence gcd(a(n),p) > 1; but p is prime and has only 1 and itself as divisors; so gcd(a(n),p)=p, and so a(n) is a multiple of p. (Weaker than the similar conjecture above.) - _John Mason_, Apr 15 2015 [WORDS IN CAPS ADDED BY _N. J. A. Sloane_, Apr 16 2015] Conjecture. The primes exist as elementary terms of the sequence in ascending order. - _John Mason_, Apr 17 2015 Conjecture. For any n > 4, the lowest value x missing from a(1) thru a(n) is prime. - _John Mason_, Apr 29 2015. In fact, taking into account that, apparently, prime p appears in the sequence at position circa 2p, we may conjecture that the lowest k values missing from a(1) thru a(n) are prime, where k = pi(n)-pi(n/2) - see A000720. - _John Mason_, Jun 03 2015 Conjecture. For n > 778, if a(n) < n, then a(n) is prime. This has been confirmed for n through 10^9. - _John Mason_, Jun 03 2015 [Corrected, following suggestion by _John P. Linderman_, by _John Mason_, May 28 2017] Conjecture. As for "Conjecture 1" above, which is its mirror image, except for n=2,3,21, corresponding to primes 2,3,11, if a(n-2)=mp

is the first occurrence of prime p as a factor in the sequence, then m=2 and a(n)=p. Also, if a(n-2)=mp is the first occurrence of prime p as a factor in the sequence, then p does not divide a(n-1). - _John Mason_, May 31 2016 Theorem 4. If a(n) = 2p is the first term having p (prime) as a factor, then a(n+2), if it exists, is either p or 2u for some integer u such that 2u < p. (Note that it is conjectured to be always p, and observation confirms the conjecture.) Theorem 5, generalization of Theorem 4. If a(n) = cp is the first term having p (prime) as a factor (n >3), and as a consequence c=k^i for prime k and i>0, then a(n+2), if it exists, is either p or ku for some integer u such that ku < p. (Note that it is conjectured to be always p, and observation confirms the conjecture.) Conjecture: After 2 and 3, no two primes are consecutive terms. This conjecture is derivable from the previous conjecture : "For k>=3, except for k=5, if a(n) = prime(k), then a(n-2) = 2*prime(k) ...". For, if sequence has terms z,2p,2q,p,q for primes p & q, then gcd(2q,z)>gcd(2q,2p)=2. Hence q divides z. So terms are mq,2p,2q,p,q. So we could have used q in place of 2q. - _John Mason_, May 28 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jan 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009493 | OEIS-A254129

Number of 2n-move closed knight paths on an unbounded chessboard from a given square to the same square.

Recurrence (conjectured): 3*n^2*(3*n-2)*(3*n-1)*(4*n-3)*(4*n-1)*(58625*n^6 - 574525*n^5 + 2317575*n^4 - 4929815*n^3 + 5836090*n^2 - 3647730*n + 940788)*a(n) = 4*(563444875*n^12 - 7212094400*n^11 + 40894216825*n^10 - 135653664390*n^9 + 292742658975*n^8 - 432166599360*n^7 + 446527351283*n^6 - 324481592710*n^5 + 164046706898*n^4 - 56035458036*n^3 + 12203976528*n^2 - 1507156200*n + 78246000)*a(n-1) - 64*(n-1)*(2*n-3)^2*(167726125*n^9 - 1643716025*n^8 + 6735239425*n^7 - 15048594215*n^6 + 20072439970*n^5 - 16473493280*n^4 + 8273936628*n^3 - 2437948332*n^2 + 377982648*n - 22556880)*a(n-2) + 165888*(n-2)*(n-1)*(2*n - 5)^2*(2*n - 3)^2*(58625*n^6 - 222775*n^5 + 324325*n^4 - 232265*n^3 + 86220*n^2 - 15570*n + 1008)*a(n-3). - _Vaclav Kotesovec_, Jan 30 2015 The conjectured recurrence of _Vaclav Kotesovec_ is true. Running the input file inSMAZ6 (see Links) on the Maple program SMAZ gives the recurrence followed by the certificate shown in the output file oSMAZ6. - _Doron Zeilberger_, Mar 29 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David A. Corneth_, Jan 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009494 | OEIS-A254130

Numbers whose factorials are exclusionary: numbers n such that n and n! have no digits in common.

Conjecture: The sequence is finite, with 16 being the last term. If A182049 is finite, then this sequence is finite. If 41 is the largest term in A182049 (as is conjectured), then 16 is the largest term of this sequence. - _M. F. Hasler_, May 04 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, May 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009495 | OEIS-A254211

Number of length n 1..3 arrays with no leading or trailing partial sum equal to a prime and no consecutive values equal.

Conjecture: There are infinitely many n > 0 with a(n) > 0. - _Alois P. Heinz_, Jan 27 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009496 | OEIS-A254311

Set of all natural numbers m such that m < S(m), where the function S is as defined in A257480.

Conjecture: A254312 is a permutation of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, May 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009497 | OEIS-A254312

Rectangular array A read by upward antidiagonals in which the entry in row n and column k is defined by A(n,k) = (2^a(n)*(6*k - (3 - (-1)^a(n))*(1 - (-1)^n)/2) - 2^n + 4)/6, n,k >= 1, where {a(n)} is the Beatty sequence A117630 defined by a(n) = floor(n*log(3)/log(3/2)).

Conjecture: The array A contains without duplication all natural numbers m such that m < S(m), where the function S is as defined in A257480; i.e., the sequence is a permutation of A254311.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, May 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009498 | OEIS-A254337

Lexicographically earliest sequence of distinct numbers such that no sum of consecutive terms is prime.

I conjecture that the sequence contains all even numbers > 2 and no odd number beyond 1. If so, we must simply ensure that the sum a(1)+...+a(n) is not prime, which is always possible for one of the three consecutive even numbers {2n, 2n+2, 2n+4}. As a consequence, it would follow that a(n) ~ 2n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009499 | OEIS-A254439

Median of terms of A254296 in the range (3^(n-1)+1)/2 to (3^n-1)/2.

From conjectured formula, it appears that next terms are 1107102779611719118, 11090084422457163934046, 3020025292945963031585836 42. - _Benedict W. J. Irwin_, Nov 16 2016 Conjecture: for n>3, a(n+3) = Sum_{i_1=1..2} Sum_{i_2=1..3*i_1-1} ... Sum_{i_n..3*i_(n-1)-1} (3*i_n - 1). - _Benedict W. J. Irwin_, Nov 16 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Md. Towhidul Islam_, Mar 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009500 | OEIS-A254460

a(n) is the number of predecessors of the all-ones state of the binary cellular automaton on the n X n grid graph with edges joining diagonal neighbors as well as vertical and horizontal neighbors, whose local rule is f(i,j) = sum of the state at vertex (i,j) and the states at all of its neighbors mod 2.

Later we discovered the relationship to Sutner's paper. A corollary of that paper is that a(n) > 0 for all n. An obvious conjecture is that a(n) = 1 for n mod 3 = 0 or 1 and if n mod 3 = 2 then a(n) = 2^(2n-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _W. Edwin Clark_, Jan 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009501 | OEIS-A254573

Number of ways to write n = x*(x+1) + y*(3*y+1)/2 + z*(3*z-1)/2 with x,y,z nonnegative integers.

Conjecture: a(n) > 0 for all n. Also, a(n) = 1 only for n = 0, 1, 4, 6, 10, 11, 23.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009502 | OEIS-A254574

Number of ways to write n = x*(x+1)/2 + y*(3*y+1)/2 + z*(3*z-1)/2 with x,y,z nonnegative integers.

Conjecture: (i) a(n) > 0 for all n. Also, a(n) = 1 only for n = 0, 49. See also the comments of A254573 for a similar conjecture. We have proved that any nonnegative integer can be written as x*(x+1)/2 + y*(3*y+1)/2 + z*(3*z-1)/2 with x,y,z integers. Note that Zhi-Wei Sun conjectured in 2009 (cf. Conjecture 1.10 of arXiv:0905.0635) that every n = 0,1,... can be expressed as the sum of a triangular number and two pentagonal numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009503 | OEIS-A254595

Least positive integer m such that m can be written as x*(x+1) + y*(3*y+1)/2 + z*(3*z-1)/2 in exactly n ways, where x,y,z are nonnegative integers.

Conjecture: (i) a(n) exists for any n > 0. Moreover, n^2 is the main term of a(n) as n tends to the infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009504 | OEIS-A254623

Number of ways to write n as x^2 + y*(3*y+1)/2 + z*(5*z+3)/2 with x,y,z nonnegative integers.

Conjecture: (i) a(n) > 0 for all n, and a(n) > 1 for all n > 118. Note that k*(3*k+1)/2 (k = 0,1,...) are second pentagonal numbers and k*(5*k+3)/2 (k = 0,1,...) are second heptagonal numbers. The conjecture has been verified for all n = 0.. 2*10^6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009505 | OEIS-A254629

Number of ways to write n as x^2 + y*(y+1) + z*(4*z+1) with x,y,z nonnegative integers.

Conjecture: (i) a(n) > 0 for all n, and a(n) > 1 for all n > 338.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009506 | OEIS-A254631

Number of ways to write n as x*(x+1)/2 + y*(3*y+2) + z*(3*z-2) with x,y,z nonnegative integers.

Conjecture: a(n) > 0 for all n, and a(n) > 1 for all n > 35.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009507 | OEIS-A254661

Number of ways to write n as the sum of a triangular number, an even square and a second pentagonal number.

Conjecture: (i) a(n) > 0 for all n. Also, a(n) = 1 only for n = 0, 1, 2, 4, 9, 11, 13, 20, 35, 69, 75, 188. Compare part (i) of this conjecture with the conjecture in A160325.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009508 | OEIS-A254668

Number of ways to write n as the sum of a square, a second pentagonal number, and a hexagonal number.

Conjecture: a(n) > 0 for all n. Also, a(n) = 1 only for n = 0, 5, 13, 14, 20, 112, 125. Compare this conjecture with the conjecture in A160324. The conjecture that a(n) > 0 for all n = 0,1,2,... appeared in Conjecture 1.2(ii) of the author's JNT paper in the links. - _Zhi-Wei Sun_, Oct 03 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009509 | OEIS-A254669

Consider A098550(x_n)=2*prime(n). Let a(n) be number of even numbers<=2*prime(n) in A098550 in positions <=x_n.

Conjecture. lim a(n)/prime(n)=1, as n goes to infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Feb 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009510 | OEIS-A254687

Number of decompositions of 2n into sums of two primes p1 < p2 such that p2-p1-1 is also a prime.

a(n)=0 for n = 1, 2, 3, 4, 6, 18, 79. It is conjectured that there is not any other n for which a(n)=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Feb 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009511 | OEIS-A254690

Number of decompositions of 2n into a sum of two primes p1 < p2 such that p2-p1 is between a pair of sexy primes.

Conjecture: when n>=7, a(n)>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Feb 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009512 | OEIS-A254699

Number of length 1+4 0..n arrays with every five consecutive terms having the maximum of some two terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Dec 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009513 | OEIS-A254700

Number of length 2+4 0..n arrays with every five consecutive terms having the maximum of some two terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Dec 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009514 | OEIS-A254701

Number of length 3+4 0..n arrays with every five consecutive terms having the maximum of some two terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Dec 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009515 | OEIS-A254702

Number of length 4+4 0..n arrays with every five consecutive terms having the maximum of some two terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Dec 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009516 | OEIS-A254703

Number of length 5+4 0..n arrays with every five consecutive terms having the maximum of some two terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Dec 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009517 | OEIS-A254704

Number of length 6+4 0..n arrays with every five consecutive terms having the maximum of some two terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Dec 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009518 | OEIS-A254705

Number of length 7+4 0..n arrays with every five consecutive terms having the maximum of some two terms equal to the minimum of the remaining three terms.

Conjectures from _Colin Barker_, Dec 17 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009519 | OEIS-A254713

All numbers k such that the number of distinct parts of all A045917(k) Goldbach partitions of 2k is prime.

Conjecture: a(k) is prime for k > 3. Tested for k up to 3*10^4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ivan N. Ianakiev_, Feb 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009520 | OEIS-A254835

Total number of nonagons in a variant of a nonagon expansion ("side-to-side", two consecutive sides) after n iterations.

Conjectures from _Colin Barker_, Feb 08 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Feb 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009521 | OEIS-A255134

First differences of A097764.

conjecture: a(n) > 1; empirically: Min{a(n): 1 <= n <= 10^7} = 9;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Feb 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009522 | OEIS-A255162

Rational part of circle radii in nested circles and hexagons (see comment).

Conjectures from _Colin Barker_, Feb 15 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Feb 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009523 | OEIS-A255163

Irrational parts of circle radii in nested circles and hexagons (see comment).

Conjectures from _Colin Barker_, Feb 15 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Feb 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009524 | OEIS-A255170

a(n) = A087803(n) - n + 1.

Conjectured extension of A199812: number of distinct values taken by w^w^...^w (with n w's and parentheses inserted in all possible ways) where w is the first transfinite ordinal omega. So far all known terms of A199812 (that is, 20 of them) coincide with this sequence. It is conjectured that A199812 is actually identical to this sequence, but it remains unproved, and is computationally difficult to check for n > 20.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Reshetnikov_, Feb 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009525 | OEIS-A255242

Calculate the aliquot parts of a number n and take their sum. Then repeat the process calculating the aliquot parts of all the previous aliquot parts and add their sum to the previous one. Repeat the process until the sum to be added is zero. Sequence lists these sums.

Also, following formulas were conjectured by Sequence Machine:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_, Feb 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009526 | OEIS-A255251

Decimal expansion of d(n) = log(n) + log(log(n)) - prime(n)/n at n = 2688, a (local?) maximum.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard R. Forberg_, Feb 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009527 | OEIS-A255257

Expansion of psi(x) * phi(-x^2)^2 in powers of x where phi(), psi() are Ramanujan theta functions.

Conjecture: Sum_{k=1..n} a(k)^2 ~ c * n^(3/2), where c = 3.147... - _Vaclav Kotesovec_, Sep 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Feb 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009528 | OEIS-A255348

a(1)=1, a(2) = 2; for n > 2, a(n) is the smallest number not already used such that the set of prime divisors of one of the numbers {a(n-1), a(n)} is a subset of the set of prime divisors of the other.

The sequence is an analog of EKG-sequence A064413 with more hard condition on neighbor terms. However, we conjecture that, as A064413, it is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009529 | OEIS-A255350

Number of ways to write n as a*(2a-1)+ b*(2b-1) + c*(2c+1) + d*(2d+1), where a,b,c,d are nonnegative integers with a <= b and c <= d.

Conjecture: (i) a(n) > 0 for all n. In other words, any nonnegative integer can be expressed as the sum of two hexagonal numbers and two second hexagonal numbers. We have verified parts (i) and (ii) of the conjecture for n up to 10^7 and 10^6 respectively.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009530 | OEIS-A255352

List of quadruples (a,b,c,d) with a^4 + b^4 = c^4 + d^4, a < c < d < b, listed in order of the largest term b.

As far as is known, the existence of solutions to the equation with exponent n=5 remains an open question.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009531 | OEIS-A255370

Position of the n-th prime in A257112.

We conjecture that the sequence is infinite or, equivalently, A257112 contains all primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, May 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009532 | OEIS-A255401

Numbers m with the property that its k-th smallest divisor, for all 1 <= k <= tau(m), contains exactly k "1" digits in its binary representation.

Conjectures: 1) Sequence is infinite. 2) 8 is the maximal value of k for numbers with this property. Sequence of the smallest numbers n with k divisors having these properties for k >= 1 or 0 if no solution exists or has been found: 1, 3, 25, 39, 0, 1083, 0, 7955, ...; a(5) = a(7) = 0 if there are only 5 Fermat primes. Conjecture: a(k) = 0 for k > 8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Feb 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009533 | OEIS-A255437

In positive integers: replace k^2 with the first k odd numbers.

conjecture: a(A068722(n)) = (2*n+1)^2, i.e. A068722(n) = gives the position of the first occurrence of n-th odd square;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Mar 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009534 | OEIS-A255472

Number of decompositions of 2n into sums of two primes p <= q such that one or both p and q are elements of A023201.

Conjecture: for all n > 3, a(n) > 0. The conjecture is weaker than a conjecture of Sun posed in 2012 (see A219055). - _Zhi-Wei Sun_, Mar 18 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Feb 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009535 | OEIS-A255517

Number A(n,k) of rooted identity trees with n nodes and k-colored non-root nodes; square array A(n,k), n>=0, k>=0, read by antidiagonals.

Conjecture: For big k the limit asymptotically approaches k*exp(1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Feb 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009536 | OEIS-A255577

Numbers n not coprime to 10 such that there exists an integer k > 1 where n^k contains n as its last digits in base 10.

For n > 1, a(n) <= a(n-1) + 2^(ceiling(log_10(a(n))) + 1) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seyed Ali Tavakoli-Nabavi_, May 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009537 | OEIS-A255617

a(n) is the smallest even term in A098550 less than 2*prime(n) but occurring after 2*prime(n); a(n)=0 if no such term exists.

Conjecture: for n>=5, positive a(n) > prime(n). Together with the known conjecture that, for n>=26, 2*prime(n) is the first multiple of prime(n) appearing in A098550, this conjecture would easily implies that the primes in A098550 occur in the natural order. (Indeed, if 101<=P<Q, then, as known, the first multiple of P, 2*P, appears earlier than the first multiple of Q, 2*Q. By the conjecture, after 2*P there cannot appear any even number <P; thus if a(n)=2*P, then a(n+2)=P appears before 2*Q.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Feb 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009538 | OEIS-A255673

Coefficients of A(x), which satisfies: A(x) = 1 + x*A(x)^3 + x^2*A(x)^6.

a(n+1) = Sum_{j=0..3*n+4} binomial(j,2*j-5*n-7) * binomial(3*n+4,j) / (3*n+4). (conjectured). [_Vladimir Kruchinin_, Mar 09 2013] a(n) = 1/(3*n+1) * Sum_{k=0..n} (-1)^k * binomial(3*n+1, k) * binomial(6*n+2-2*k, n-k). (conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Jul 10 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009539 | OEIS-A255677

Least integer k > 1 such that pi(k)^2 + pi(k*n)^2 is a square, where pi(.) is the prime-counting function given by A000720.

Conjecture: Each positive rational number r < 1 can be written as m/n with 1 < m < n such that pi(m)^2 + pi(n)^2 is a square. Also, any rational number r > 1 can be written as m/n with m > n > 1 such that pi(m)^2 - pi(n)^2 is a square.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 10 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009540 | OEIS-A255679

Least number k such that pi(k*n)^2 = pi(i*n)^2 + pi(j*n)^2 for some 0 < i < j < k.

Conjecture: a(n) exists for any n > 0. In other words, for each n = 1,2,3,... the sequence pi(k*n) (k = 1,2,3,...) contains a Pythagorean triple.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009541 | OEIS-A255683

Sum of the binary numbers whose digits are cyclic permutations of the binary expansion of n.

For n >= 0 and 0 <= i <= 2^n - 1 we conjecture a(2^n + i) = (2^(n+1) - 1)*A063787(i+1). An example is given below. - _Peter Bala_, Mar 02 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_, Mar 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009542 | OEIS-A255746

Squarefree semiprimes n such that n+4 is also a squarefree semiprime.

Conjecturally, the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jul 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009543 | OEIS-A255807

Expansion of e.g.f.: exp(Sum_{k>=1} k^2 * x^k).

It appears that the sequence a(n) taken modulo 10 is periodic with period 5. More generally, we conjecture that for k = 2,3,4,... the difference a(n+k) - a(n) is divisible by k: if true, then the sequence a(n) taken modulo k would be periodic with period dividing k. - _Peter Bala_, Nov 14 2017 The above conjecture is true - see the Bala link. - _Peter Bala_, Jan 20 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Mar 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009544 | OEIS-A255833

Index i such that A166133(i)=n, or 0 if no such i exists.

If A166133 is a permutation, as conjectured, then this is the inverse permutation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009545 | OEIS-A255874

Triangular array T: T(n,k) = number of subset S of {1,2,...,n+1} such that |S| > 1 and max(S*) = k, where S* is the set {x(2)-x(1), x(3)-x(2), ..., x(h+1)-x(h)} when the elements of S are written as x(1) < x(2) < ... < x(h+1).

Column 1: A000217. Conjectures: Column 2 = A014162, and the rows have a limiting tail (1,4,12,32,...) = A001787.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009546 | OEIS-A255895

O.g.f.: exp(Sum_{n>=1} A000364(n+1)*x^n/n), where A000364 is the Euler numbers.

Inspired by Peter Bala's sequence A255881, which suggests the following conjecture: exp(Sum_{n>=1} A000364(n+m)*x^n/n) is an integer series for integer m>=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009547 | OEIS-A255905

Expansion of exp(Sum_{n >= 1} R(n,u)*x^n/n), where R(n,u) denotes the n-th row polynomial of A086646.

The above conjecture can be extensively generalized. The elements of A000364 can be expressed in terms of the Euler polynomial E(n,x) as A000364(n) = (-1)^n*2^(2*n)*E(2*n,1/2). This suggests considering polynomials of the form P(n,u) = Sum_{k = 0..n} binomial(2*n,2*k)*A(n-k)*u^k, where the sequence A(n) is defined in terms of the Euler polynomials. Calculation suggests that in the expansion of exp(Sum_{n >= 1} P(n,u)*x^n/n), the coefficient polynomials in u are always integer polynomials for the following choices of A(n): The present triangle (up to signs) is simply the case of conjecture 1 with the choices h = 1 and k = 2. Similar conjectures can be made if, in the above definition of the polynomial P(n,u), the factor equal to binomial(2*n,2*k) is replaced by binomial(m*n,m*k) for some fixed m = 1,2,3,....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 10 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009548 | OEIS-A255912

O.g.f.: exp(Sum_{n>=1} A000364(2*n)*x^n/n), where A000364 is the Euler numbers.

a(n) == 5 (mod 100) for n>=1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 10 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009549 | OEIS-A255916

Number of ways to write n as the sum of a generalized heptagonal number, an octagonal number and a nonagonal number.

Conjecture: (i) a(n) > 0 for all n. Moreover, for k >= j >=3, every nonnegative integer can be written as the sum of a generalized heptagonal number, a j-gonal number and a k-gonal number, if and only if (j,k) is among the following ordered pairs:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009550 | OEIS-A255926

Expansion of exp(Sum_{n >= 1} A210676(n)*x^n/n).

This sequence is the particular case m = -3 of the following general conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009551 | OEIS-A255928

Expansion of exp(Sum_{n >= 1} A094088(n)*x^n/n).

This sequence is the particular case m = 1 of the following general conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009552 | OEIS-A255929

Expansion of exp(Sum_{n >= 1} A210672(n)*x^n/n).

This sequence is the particular case m = 2 of the following general conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009553 | OEIS-A255930

Expansion of exp(Sum_{n >= 1} A210674(n)*x^n/n).

This sequence is the particular case m = 3 of the following general conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009554 | OEIS-A255932

a(n) is the denominator of Gamma(n+1/2)^2/(2*n*Pi), the value of an integral with sinh in the denominator.

Conjecture: a(n) <= 2^(3*n). - _Vaclav Kotesovec_, Mar 11 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Mar 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009555 | OEIS-A255934

Number of ways to write n as the sum of four unordered generalized octagonal numbers.

I have proved that a(n) > 0 for all n, i.e., any nonnegative integer can be expressed as the sum of four generalized octagonal numbers. I can show that a(n) = 1 if 3*n+4 is among 7, 13, 19, 31, 43, 2^(2k), 5*2^(2k+1), 11*2^(2k+1), 23*2^(2k+1) (k = 0,1,2,...), and conjecture the converse. I also conjecture that each nonnegative integer can be written as the sum of two heptagonal numbers, a second heptagonal number and a generalized heptagonal number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009556 | OEIS-A255972

Smallest number m such that A251604(m) = n.

If the conjecture in A251604 is true, this sequence is the inverse permutation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Mar 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009557 | OEIS-A255992

T(n,k)=Number of length n+k 0..1 arrays with at most one downstep in every k consecutive neighbor pairs.

T(n,k) = n+k+1 + Sum_{r=1..floor((n-2)/k)+2} Sum_{j=0..r-1} (-1)^j*C(r-1,j)*(k-1)^(r-1-j)*(k-2)^j*C(n+(2-r)*k+2*r-1-j,2*r+1). This confirms all conjectures here and in the related sequences A255993-A255998 (see Fried link). - _Sela Fried_, Mar 20 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 13 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009558 | OEIS-A255999

Position of the n-th prime in A253279.

We conjecture that the sequence is infinite or, equivalently, A253279 contains all primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, May 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009559 | OEIS-A256001

Minimum sum of a nonnegative integer triple that takes n moves to reach a 0 component, where a move picks two components, subtracts the smaller from the larger, and doubles the smaller.

Conjecture: last diameters n are for S = 5*2^(n-1) when n>=1, i.e. 5 for diameter 1, 10 for diameter 2, 20 for diameter 3, ... This formula works for all diameters calculated up to S=880K. The diameter of S = 5*2^(n-1) is n which is verified up to n=19; i.e., S=1310720. However, a proof is needed to confirm this formula (as we cannot exclude the possibility that higher S leads to a lower diameter). - _Karl Desfontaines_, Dec 15 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John Tromp_, May 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009560 | OEIS-A256037

Triangle read by rows: number of R-class idempotents of rank k in Brauer monoid B_n.

Conjecture: T(n,k) = (n+k-1)!!/(2k-1)!! for n+k even, T(n,k) = 0 otherwise. - _Jean-François Alcover_, Feb 11 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009561 | OEIS-A256046

Main diagonal of A256045: order of all-2s configuration on the n X n sandpile grid graph.

Conjecture (due to Daniel Tabares): for odd n >= 3, v(a(n)) = v(n+1) + 1, where v(n) = A007814(n) is the 2-adic valuation of n. - _Nikita Kalinin_, Jul 08 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009562 | OEIS-A256060

Queen Dido's puzzle (the founding of Carthage): a(n) is twice the maximal area of a polygon with 1) vertices on integral Cartesian coordinates, 2) no two edges parallel, and 3) all edge lengths less than or equal to n^2.

Are these values known to be optimal or are they conjectures? - _N. J. A. Sloane_, Mar 13 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gordon Hamilton_, Mar 13 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009563 | OEIS-A256071

Number of ordered ways to write n = p + x*(3*x-1)/2, where p is prime or zero, and x is an integer.

Conjecture: a(n) > 0 for all n. Moreover, each nonnegative integer n is either an odd prime, or a generalized pentagonal number, or the sum of an odd prime and a generalized pentagonal number. This is similar to the author's earlier conjecture on sums of primes and triangular numbers (see the reference and also A132399). The conjecture has been verified for all n = 0..10^9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 13 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009564 | OEIS-A256106

Number of ways to write n as w + x + 2*y + 4*z, where w,x,y,z are hexagonal numbers with w <= x.

Conjecture: (i) a(n) > 0 for all n. I have shown the following related result: For m > 4 and 0 < a <= b <= c <= d, if every nonnegative integer can be written as a*w + b*x + c*y + d*z with w,x,y,z m-gonal numbers, then either m = 6 and (a,b,c,d) = (1,1,2,4), or m = 5 and a = 1 and (b,c,d) is among the 15 triples listed in part (ii) of the conjecture. In the preprint arXiv:1608.02022, Xiang-Zi Meng and Zhi-Wei Sun confirmed part (i) of the conjecture, and they also proved that for each triple (b,c,d) = (1,2,2),(1,2,4) any natural number can be written as w + b*x + c*y + d*z with w,x,y,z pentagonal numbers. _Zhi-Wei Sun_, Aug 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009565 | OEIS-A256111

a(n) = squared distance to the origin of the n-th vertex on a Babylonian Spiral.

The name is chosen to mislead school students into making an incorrect hypothesis about the Babylonian Spiral's long-term behavior.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gordon Hamilton_, Mar 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009566 | OEIS-A256119

Least number p that is zero or an odd prime, such that n - p is a generalized pentagonal number.

By the conjecture in A256071, a(n) always exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009567 | OEIS-A256132

Number of ways to write n as w*(3w+1)/2 + x*(3x-1)/2 + y*(3y-1)/2 + z*(3z-1)/2, where w,x,y,z are nonnegative integers with x <= y <= z.

Conjecture: a(n) > 0 for all n. In other words, any nonnegative integer n can be expressed as the sum of three pentagonal numbers and a second pentagonal number. See also A255350 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009568 | OEIS-A256171

Number of ways to write n as the sum of three unordered generalized heptagonal numbers.

Conjecture: (i) a(n) > 0 except for n = 10, 16, 76, 307.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009569 | OEIS-A256174

Boomerang Fractions: Starting with 1, on the first step add 1/n, and on subsequent steps either add 1/n or take the reciprocal. a(n) = minimal number of steps needed to return to 1.

At that event, the sequence was calculated with conjectures for the 11th and 13th terms: 4, 9, 7, 20, 6, 33, 13, 23, 16, 62?, 8, 75?, 18, 17, 25. - _Gordon Hamilton_, Apr 02 2015 Additional values: a(35) = 25, a(48) = 29. Upper bounds: a(n) <= n^2. Add n(n-1) times, take the reciprocal and add n-1 times. a(bc) <= c(b^2-1) + 1. Add n(b-1) times, take the reciprocal and add (b-1)c times. For c > b, a(bc) <= c^2-b^2 + 1. Add c(c-b) times, take the reciprocal and add (c-b)b times. In particular, for n = b(b+1), a(n) <= 2(b+1). The minimum of these bounds give a(n) for n = 2, 3, 4, 6, 8, 10, 12, 15, 20, 30, 35, 42, 48, 56, 72, 90, 110, 132. Exhaustive search show that a(b*(b+1)) = 2(b+1) for b <= 12 which seems to support the conjecture that this is true for all b. - _Chai Wah Wu_, Apr 02 2015 Minimizing the total number of steps under the additional constraint that the number of consecutive "add 1/n" steps immediately before each "take the reciprocal" step must be a multiple of n gives the following upper bound on a(n) for n in [2..13]: 4, 9, 7, 20, 10, 33, 13, 23, 16, 62, 19, 75. Note that these values match the known exact values of a(n) at n = 2, 3, 4, 5, 7, 8, 9, 10, and 11, as well as the conjectured value at n = 13. See the Links for a file giving the value of this upper bound, and a solution at that number of steps, for all n in [2..10000]. Conjecture: For all prime values of n, a(n) is exactly the minimal number of steps to return to 1 under the added constraint that the number of consecutive "add 1/n" steps immediately before each "take the reciprocal" step must be a multiple of n. A stronger conjecture: The above conjecture applies not only when n is prime, but also when n is a prime power. If this is true, then it appears that If the above conjectures about a(n) where n is a prime or prime power are correct, then a(17)..a(49) are 111, 16, 126, 10, 29, 34, 171, 15, 88, 40, 65, 27, 217, 12, 235, 49, 66, 52, 25, 22, 310, 58, 31, 19, 345, 14, 362, 66, 49, 70, 396, 29, 213; a(50) seems very likely to be 73. - _Jon E. Schoenfield_, Apr 20 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gordon Hamilton_, Mar 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009570 | OEIS-A256189

a(n) = n if n <= 3, otherwise a(n) is the smallest number not occurring earlier such that gcd(a(n-2), a(n)) is a prime and gcd(a(n-1), a(n)) = 1.

Moreover, this sequence conjecturally is a permutation of the positive integers: the proof i-v for A098550 applies with essentially no changes, while vi still has a difficulty.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Mar 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009571 | OEIS-A256224

a(n) = n if n <= 3, otherwise a(n) is the smallest number not occurring earlier such that gcd(a(n-2), a(n)) is a prime or a power of a prime (but not 1) and gcd(a(n-1), a(n))=1.

Conjecture: This is a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Mar 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009572 | OEIS-A256260

Total number of ON states after n generations of a cellular automaton-like on the square grid.

See also the conjecture in the Example section.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Mar 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009573 | OEIS-A256271

a(1) = 1; for n > 1, a(n) is the smallest number not already in the sequence such that a(n)^2 + a(n-1)^2 is squarefree.

This is conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Jun 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009574 | OEIS-A256283

Smallest m such that A257905(m) = n.

Conjectured inverse of A257905.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jun 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009575 | OEIS-A256330

Number of H&S Family matchings on n edges.

Jay Pantone has computed the first 1500 terms and has a conjectured g.f. - _N. J. A. Sloane_, Oct 06 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aziza Jefferson_, Mar 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009576 | OEIS-A256366

Primes arising in A=A256189 as gcd(A(n-2), A(n)), n > 3.

Conjecture: the sequence contains all primes such that the first appearances of them are in the natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Mar 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009577 | OEIS-A256368

Primes arising from A098550(n), n >= 2, as smallest prime factor of gcd(A098550(n), A098550(n+2)).

Conjecture: the sequence contains all primes such that the first appearances of them are in the nature order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Mar 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009578 | OEIS-A256386

Numbers m such that m-2, m-1, m+1, m+2 cannot all be represented in the form x*y + x + y for values x, y with x >= y > 1.

Conjectures: Conjecture 2 would follow if we establish the equivalence "t is in sequence" <=> "t is a term of b(n): lesser of twin primes pair p and q such that (p - 1)/2 and (q + 1)/2 are also a pair of twin primes (A077800)". a(n) = A158870(n-5) - 2, n > 5 (conjectured). - _Lamine Ngom_, Feb 12 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Mar 31 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009579 | OEIS-A256387

Numbers n such that no prime can be the arithmetic mean of 2 semiprimes whose difference is 2*n.

There any no even members up to 10^5. Conjecture: all members are odd. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Mar 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009580 | OEIS-A256395

Composite Markoff numbers.

Bourgain, Gamburd, and Sarnak have announced a proof that almost all Markoff numbers A002559 are composite. Equivalently, the prime Markoff numbers A178444 have density zero among all Markoff numbers. (It is conjectured that infinitely many Markoff numbers are prime.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Apr 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009581 | OEIS-A256410

Subtract 1 from the terms of A256407.

Conjectured to be always a prime. But which primes are these? It would be nice to know of some other property that distinguishes these primes. See also the "blog" in A166133.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009582 | OEIS-A256414

Indices of prime terms in A121217.

A010051(A121217(a(n))) = 1; conjecture: sequence is strictly increasing. - _Reinhard Zumkeller_, Apr 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009583 | OEIS-A256429

a(n) is the total number of pentagrams in a variant of pentagram fractal after n iterations.

Conjectures from _Colin Barker_, Mar 29 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Mar 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009584 | OEIS-A256438

Numbers m such that sigma(sigma(m-1)) = 2*(m-1).

Conjecture: numbers of the form 2^k+1 such that sigma(2^k) = prime p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Mar 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009585 | OEIS-A256447

Number of integers in range (prime(n)^2)+1 .. (prime(n)*prime(n+1)) whose smallest prime factor is at least prime(n): a(n) = A250477(n) - A250474(n).

See the conjectures in A256448.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009586 | OEIS-A256461

Smallest k such that A098550(n) + A098550(n+1) + ... + A098550(n+k) is prime, or -1 if there is no such k.

Conjecture: a(n)>=0 for all n. In contrast to this conjecture, for the sequence of the positive integers (A=A000027) there are infinitely many n for which A(n) + A(n+1) + ... + A(n+k) is composite for every k>=0 (cf. A089306, A077654).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Mar 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009587 | OEIS-A256481

Smallest prime obtained by appending a number with identical digits to n or 0 if no such prime exists.

For n <= 15392, a(n) = 0 if and only if n = 6930. Conjecture: if a(n) = 0, then n is divisible by 3. Conjecture verified for n <= 10^6. a(n) = 0 for n = 6930, 50358, 56574, 72975.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Mar 31 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009588 | OEIS-A256527

a(n) is the least number k > 0 such that sigma(k) = phi(n*k).

For n >= 5, a(n) = A087979(n). - Conjectured by _Manfred Scheucher_, May 28 2015; proved by _Max Alekseyev_, Sep 29 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_, Apr 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009589 | OEIS-A256535

The largest number of T-tetrominoes that fit within an n X n square.

Shuxin Zhan proved that it is not possible to tile a square of side 4m+1 or 4m+3 with T-tetrominos and a single monomino. Thus there must be at least 5 empty squares in any partial tiling by T-tetrominos. This bound is achieved for tilings in 5 X 5, 7 X 7, 9 X 9 and 11 X 11 squares. Robert Hochberg proved that for n > 11, there must be either 5 or 9 empty squares. He conjectured that 5 is always enough.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jack W Grahl_, Sep 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009590 | OEIS-A256544

Number of ways to write n as the sum of three unordered elements of the set {floor(T(x)/3): x = 1,2,3,...}, where T(x) denotes the triangular number x*(x+1)/2.

Conjecture: For any positive integer m, every nonnegative integer n can be written as floor(T(x)/m) + floor(T(y)/m) + floor(T(z)/m) with x,y,z nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009591 | OEIS-A256557

a(n) = A166133(n)^2 - 1.

Since it is conjectured that A166133 is a permutation of the natural numbers, it is therefore conjectured that this sequence is a permutation of all numbers of the form n^2-1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bob Selcoe_, Apr 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009592 | OEIS-A256558

Number of ways to write n = p + floor(k*(k+1)/4), where p is a prime and k is a positive integer.

Conjecture: (i) a(n) > 0 for all n > 1. We also have some other conjectures on representations n = p + floor(k*(k+1)/m) with m > 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009593 | OEIS-A256587

Number of ways to write n = r + s + t, where r,s,t are elements of the set {floor(k*(k+1)/4): k = 1,2,3,...} with s odd and r <= s <= t.

Conjecture: (i) a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009594 | OEIS-A256598

Irregular triangle where row n contains the odd terms in the Collatz sequence beginning with 2n+1.

The Collatz function is an integer-valued function given by n/2 if n is even and 3n+1 if n is odd. We build a Collatz sequence by beginning with a natural number and iterating the function indefinitely. It is conjectured that all such sequences terminate at 1. In this triangle, row n is made up of the odd terms of the Collatz sequence beginning with 2n+1. Therefore, it is conjectured that this sequence is well-defined, i.e., that all rows terminate at 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bob Selcoe_, Apr 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009595 | OEIS-A256618

Index i such that A121217(i) = n, or 0 if no such i exists.

If A121217 is a permutation, as conjectured, then this is the inverse permutation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009596 | OEIS-A256619

Numbers n such that there are no primes in the interval [b(n), b(n+1) - 1], where b(n) = 1 + Sum_{k=1..n} floor(1/2 + sqrt(2*k - 2)).

Conjecture: the sequence is finite and complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Apr 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009597 | OEIS-A256628

Index i such that A251622(i) = n, or 0 if no such i exists.

If A251622 is a permutation, as conjectured, then this is the inverse permutation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009598 | OEIS-A256661

Rectangular array by antidiagonals: row n shows the numbers k such that R(k) consists of n terms, where R(k) is the minimal alternating Fibonacci representation of k.

(col 2): A127546 (conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009599 | OEIS-A256750

Start with n, and repeatedly apply the arithmetic derivative A003415. |a(n)| = the number of iterations to reach 0 (then a(n) is taken nonnegative) or a number having a factor of the form p^p with prime p, in which case a(n) = -|a(n)|.

Under iterations of the arithmetic derivative, the orbit of some numbers ends in zero, and the orbit of all others (I conjecture) reaches a number of the form m*p^p with prime p, from where on it keeps this form and grows to infinity iff m>1, or remains at this fixed point if m=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009600 | OEIS-A256852

Number of ways to write prime(n) = a^2 + b^4.

Conjecture: a(n) <= 2, empirically checked for the first 10^6 primes. The conjecture is true, because by the uniqueness part of Fermat's two-squares theorem, at most one duplicate of a^2 + b^4 can exist. Namely, if a is a square, say a = B^2, then a^2 + b^4 = A^2 + B^4 where A = b^2. -_Jonathan Sondow_, Oct 03 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Apr 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009601 | OEIS-A256855

Number of ordered ways to write n as x*(3*x-1)/2 + y*(3*y+1)/2 + z*(3*z+1), where x and y are nonnegative integers and z is an integer.

Conjecture: a(n) > 0 for all n. Also, any nonnegative integer can be written as x*(3*x-1)/2 + y*(3*y-1) + z*(3*z+1)/2 with x,y nonnegative integers and z an integer, and each n = 0,1,2,... can be expressed as x*(3*x+1)/2 + 3*y*(3*y+1)/2 + z*(3*z+1)/2 with x,y nonnegative integers and z an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009602 | OEIS-A256862

a(1)=1, then a(n) = least number > a(n-1) such that 2*a(n-1)+a(n) is prime.

I conjecture that any initial term a(1) eventually merges with this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Apr 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009603 | OEIS-A256908

Initialization a(n)=0 for all n; if a(n)=0, a(n+kp)=n where p=prime(n) and k=0,1,2,...

Conjecture: there is an infinity of numbers n such that a(n)=n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009604 | OEIS-A256963

Partial sums of A005210.

It is conjectured that a(n) grows like n^2/6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009605 | OEIS-A256965

Decimal expansion of sqrt(2) + sqrt(3/2).

Conjectured minimal length of fence ensuring privacy of a square garden.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009606 | OEIS-A257072

Number of 3-colored Schroeder paths of semilength n in which there are no (2,0)-steps at level 1.

Conjecture: 3*n*a(n) +(-53*n+45)*a(n-1) +2*(151*n-213)*a(n-2) +9*(-73*n+144)*a(n-3) +405*(n-3)*a(n-4)=0. - _R. J. Mathar_, Sep 24 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José Luis Ramírez Ramírez_, Apr 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009607 | OEIS-A257104

Number of Motzkin paths of length n with no peaks at level 3.

Conjecture: D-finite with recurrence (-n+2)*a(n) +(7*n-17)*a(n-1) +2*(-7*n+17)*a(n-2) +(n+22)*a(n-3) +(16*n-89)*a(n-4) +(-4*n+23)*a(n-5) +3*(n-5)*a(n-6)=0. - _R. J. Mathar_, Sep 24 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José Luis Ramírez Ramírez_, Apr 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009608 | OEIS-A257109

Number of ways to write n = 3*x^2 + y^2 + z^2, where x,y,z are nonnegative integers with gcd(3,x) = 1 and y <= z.

Conjecture: (i) If n is not divisible by 3, then a(n) > 0 except for n = 1, 2, 10, 26, 31, 34, 47, 59, 166, 278, 451.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009609 | OEIS-A257162

Number of ordered ways to write n as floor(x/5) + floor(y/6) + floor(z/7), where x is a pentagonal number, y is a hexagonal number and z is a heptagonal number.

Conjecture: a(n) > 0 for all n. Moreover, for any i,j,k = 3,4,5,... all sufficiently large integers can be written as floor(x/i) + floor(y/j) + floor(z/k), where x is an i-gonal number, y is a j-gonal number and z is a k-gonal number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009610 | OEIS-A257178

Number of 3-Motzkin paths of length n with no level steps at odd level.

Conjecture: (n+2)*a(n) +6*(-n-1)*a(n-1) +(5*n+4)*a(n-2) +6*(2*n-3)*a(n-3)=0. - _R. J. Mathar_, Sep 24 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José Luis Ramírez Ramírez_, Apr 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009611 | OEIS-A257230

Floor(sqrt(q)-(q-p)), where p and q are consecutive primes.

Conjecture: a(n) is always positive for n > 30, and is negative only for n = 4, 9 and 30, corresponding to prime pairs (7, 11), (23, 29) and (113, 127). Related to prime gap conjectures by (e.g.) Legendre, Oppermann, Andrica and Brocard.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chris Boyd_, Apr 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009612 | OEIS-A257275

Numbers equal to some partial product of the sequence of their digits, repeated over and over again.

The sequence is conjectured to be finite (D. Wilson, Reply on the SeqFan list).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009613 | OEIS-A257280

Inverse permutation of A139080 (a(n+1) = least unused m such that floor(max(m,a(n))/min(m,a(n))) = 2).

A139080 was only conjectured to be a permutation, until a proof has been proposed recently.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009614 | OEIS-A257290

Number of 3-Motzkin paths of length n with no level steps at even level.

Conjecture: (n+2)*a(n) +3*(-2*n-1)*a(n-1) +5*(n-1)*a(n-2) +6*(2*n-5)*a(n-3)=0. - _R. J. Mathar_, Sep 24 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José Luis Ramírez Ramírez_, Apr 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009615 | OEIS-A257300

Number of Motzkin paths of length n with no peaks at level 2.

Conjecture: n*a(n) +(-5*n+3)*a(n-1) +6*(n)*a(n-2) +2*(n-9)*a(n-3) +6*(-n+4)*a(n-4) +(n-6)*a(n-5) +3*(-n+3)*a(n-6)=0. - _R. J. Mathar_, Sep 24 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José Luis Ramírez Ramírez_, Apr 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009616 | OEIS-A257310

Numbers n such that the decimal expansions of both n and n^2 have 2 as smallest digit and 5 as largest digit.

Conjecture: a(n) = A137066(n+2) for all n, i.e., this is A137066 without the initial two terms. (10^(k+1)-1)/3 - 10^k + 2 are terms for k > 1. Conjecture: except for 2354235, 2333524235, and 23333524235, all terms are of this form. - _Chai Wah Wu_, Sep 10 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Apr 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009617 | OEIS-A257311

a(1) = 4; a(2) = 5; for n > 2, a(n) is the smallest number of the form prime + 2 not already used which shares a factor with a(n-1).

Conjecture: the sequence {a(n)-2} is a permutation of the primes (A000040). Conjecture: For every k>=1, the sequence A_k - 2^k is a permutation of the primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009618 | OEIS-A257312

a(1) = 6; a(2) = 7; for n > 2, a(n) is the smallest number of the form prime + 4 not already used which shares a factor with a(n-1).

Conjecture: the sequence {a(n)-4} is a permutation of primes (A000040).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009619 | OEIS-A257313

a(1) = 10; a(2) = 11; for n > 2, a(n) is the smallest number of the form prime + 8 not already used which shares a factor with a(n-1).

Conjecture: the sequence {a(n)-8} is a permutation of primes (A000040).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009620 | OEIS-A257314

a(1) = 18; a(2) = 19; for n > 2, a(n) is the smallest number of the form prime + 16 not already used which shares a factor with a(n-1).

Conjecture: the sequence {a(n)-16} is a permutation of primes (A000040).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009621 | OEIS-A257315

a(1) = 34; a(2) = 35; for n > 2, a(n) is the smallest number of the form prime + 32 not already used which shares a factor with a(n-1).

Conjecture: the sequence {a(n)-32} is a permutation of primes (A000040).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009622 | OEIS-A257329

Construct spiral of numbers on square grid as in Comments; sequence gives terms along the "2" arm.

Conjecture: every number > 1 appears in one of the four arms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009623 | OEIS-A257330

Construct spiral of numbers on square grid as in Comments; sequence gives terms along the "3" arm.

Conjecture: every number > 1 appears in one of the four arms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009624 | OEIS-A257331

Construct spiral of numbers on square grid as in Comments; sequence gives terms along the "5" arm.

Conjecture: every number > 1 appears in one of the four arms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009625 | OEIS-A257332

Construct spiral of numbers on square grid as in Comments; sequence gives terms along the "7" arm.

Conjecture: every number > 1 appears in one of the four arms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009626 | OEIS-A257337

Construct spiral of numbers on square grid as in Comments; sequence gives terms along the "4" arm.

Conjecture: every number > 1 appears in one of the four arms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009627 | OEIS-A257339

Arrange numbers in a single clockwise spiral so that each number is relatively prime to its eight neighbors.

It is conjectured that every number appears.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009628 | OEIS-A257340

Arrange numbers in a single clockwise spiral so that each number is relatively prime to its four (N,S,E,W) neighbors.

It is conjectured that every number appears.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009629 | OEIS-A257347

1 followed by the numbers successively added to the four arms of the spirals in A257329, A257330, A257331, A257332.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009630 | OEIS-A257348

Repeatedly applying the map x -> sigma(x) partitions the natural numbers into a number of disjoint trees; sequence gives the (conjectural) list of minimal representatives of these trees.

Repeatedly applying the map x -> sigma(x) partitions the natural numbers into a number of disjoint trees; sequence gives the (conjectural) list of minimal representatives of these trees. Very little is known for certain. Even the trajectories of 2 (A007497) and 5 (A051572) under repeated application of the map x -> sigma(x) (cf. A000203) are only conjectured to be disjoint.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 01 2015, following a suggestion from _Kerry Mitchell_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009631 | OEIS-A257364

Least prime p such that pi(p*n)^2 = pi(q*n)^2 + pi(r*n)^2 for some primes q and r, where pi(x) denotes the number of primes not exceeding x.

Conjecture: a(n) exists for any n > 0. In other words, for each fixed positive integer n the sequence pi(p*n) with p prime contains a Pythagorean triple. This is stronger than the conjecture in A255679.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009632 | OEIS-A257378

Smallest odd number k such that k*n*2^n+1 is a prime number.

Conjecture: a(n) exists for every n. The conjecture is a corollary of Dirichlet's theorem on primes in arithmetic progressions. - _Robert Israel_, Jan 05 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Apr 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009633 | OEIS-A257388

Number of 4-Motzkin paths of length n with no level steps at odd level.

Conjecture: (n+2)*a(n) +8*(-n-1)*a(n-1) +4*(3*n+1)*a(n-2) +8*(2*n-3)*a(n-3)=0. - _R. J. Mathar_, Sep 24 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José Luis Ramírez Ramírez_, Apr 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009634 | OEIS-A257401

God's number for a Rubik's cube of size n X n X n (using the half turn metric).

Conjecture: a(n) ~ (1/4)*log(24!/4!^6) * n^2/log(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Woodward_, Apr 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009635 | OEIS-A257405

For n=1 or prime, a(n)=n; otherwise, a(n) is the smallest number not already used which shares a factor with a(n-1).

Since limsup(prime(n)) is infinity, it is easy to see that the sequence contains all even numbers. So it is natural to conjecture that the sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009636 | OEIS-A257470

a(n+1) = least number not occurring earlier such that max{a(n), a(n+1)} >= 2 min{a(n), a(n+1)}; a(0) = 0.

In spite of the simplicity of the pattern, it seems not easy to give an explicit formula of a(n), i.e., for a(3k-1), cf. formulas. I conjecture the following properties:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009637 | OEIS-A257499

Array A read by upward antidiagonals in which the entry in row n and column k is defined by A(n,k) = (1 + 2^n*(6*k-3+2*(-1)^n))/3, n,k >= 1.

Conjecture (now Lemma 1): The sequence is a permutation of the odd natural numbers. Reformulating the conjecture, we need to prove that for any integer m >= 0, the equation (1 + 2^n*(6*k - 3 + 2*(-1)^n))/3 = 2*m + 1 has a unique solution in integers n,k >= 1. Simplifying a bit, we have

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Apr 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009638 | OEIS-A257515

Number of 3-generalized 2-Motzkin paths of length n with no level steps H=(3,0) at odd level.

Conjecture: (n+2)*a(n) +(n+1)*a(n-1) +(n+4)*a(n-2) +4*(-2*n+3)*a(n-3) +4*(-6*n+17)*a(n-4) +4*(-3*n+10)*a(n-5) +4*(3*n-11)*a(n-6) +4*(11*n-50)*a(n-7) +20*(n-6)*a(n-8)=0. - _R. J. Mathar_,

Jun 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José Luis Ramírez Ramírez_, Apr 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009639 | OEIS-A257534

Numbers n such that the decimal expansions of both n and n^2 have 4 as the digit with the smallest value and 7 as the digit with the largest value.

Conjecture: a(4) does not exist. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Apr 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009640 | OEIS-A257562

Number of permutations of length n that avoid the patterns 4123, 4231, and 4312.

G.f. conjectured to be non-D-finite (see Albert et al. link). - _Jay Pantone_, Oct 01 2015 Unlike A061552, whose g.f. is also conjectured to be non-D-finite, thousands of terms of the counting sequence are known. - _David Callan_, Aug 29 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jay Pantone_, Apr 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009641 | OEIS-A257594

Consider the hexagonal lattice packing of circles; a(n) is the maximal number of circles that can be enclosed by a closed chain of n circles.

Conjecture (derived from Euler's F+V=E+1 formula): a(n) = 1+(A069813(n)-n)/2 = A001399(n-6), which means g.f. is x^6 / ((1+x)*(1+x+x^2)*(1-x)^3). - _R. J. Mathar_, Jul 14 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009642 | OEIS-A257639

a(n) is the minimal position at which the maximal value of row n appears in row n of triangle A008289.

Conjecture: for n >= 22 there is a unique maximal value in row n of triangle A008289.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Nov 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009643 | OEIS-A257662

Least prime q such that p(q*n) is prime, where p(.) is the partition function given by A000041.

Conjecture: a(n) exists for any n > 0. This implies the conjecture that the sequence p(n) (n = 1,2,3,...) contains infinitely many primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009644 | OEIS-A257663

Least positive integer k such that prime(k*n) - prime(k+n) is a square.

Conjecture: a(n) exists for any n > 1. Also, for any n > 0 there is a number k > 0 such that prime(k*n) + prime(k+n) is a square.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009645 | OEIS-A257716

a(n) = smallest prime of even index not included earlier such that a(n) + a(n-1) + a(n-2) is a prime, beginning with a(1) = 3 and a(2) = 7.

Conjecture: The union of this sequence and A257717 is A065091.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, May 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009646 | OEIS-A257717

a(n) is the smallest odd-indexed prime not included earlier such that a(n) + a(n-1) + a(n-2) is a prime, beginning with a(1) = 5 and a(2) = 11.

Conjecture: The union of this sequence and A257716 is A065091.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, May 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009647 | OEIS-A257743

Let n'= odd part of n. If n' is 1 or prime, then a(n)=1; otherwise a(n) is 1+ the number of times the map x -> (3*x+1)' must be applied to n' in order to reach 1 or a prime.

Conjecture: for every N>0 there exists n=n(N) such that a(n)>N.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009648 | OEIS-A257744

Smallest k such that A257743(k)=n.

If conjecture in comment in A257743 is true, then the sequence is infinite. For example, if this conjecture is true, then A257743 contains 10^9. Indeed, let n_0 requires N>10^9 times of the map in the name. Then there exists n_1 which requires N-1 times of the map, there exists n_2 which requires N-2 times of the map,..., there exists n_(N-10^9+1) which requires 10^9-1 times of the map such that A257743(n_(N-10^9+1)) = 10^9. Therefore, a(10^9) exists. - _Vladimir Shevelev_, May 15 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009649 | OEIS-A257750

Quasi-Carmichael numbers.

If b is negative, then it is always larger than 0 minus the square root of the corresponding Quasi-Carmichael number. But if b is positive, how large can it be in relation to its corresponding Quasi-Carmichael number? Conjecture: It is

always smaller than the square root of the corresponding Quasi-Carmichael number. The conjecture that b < sqrt(n) is false. Look at n = 87061 = 13*37*181, 87365 = 5*101*173, and 96473 = 13*41*181. Their b values are 299, 331, and 351, while the corresponding sqrt(n) values are 295, 295, and 310, respectively.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tim Johannes Ohrtmann_, May 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009650 | OEIS-A257759

Quasi-Carmichael numbers to at least one negative base and at least one positive base.

It is a open question whether any Carmichael number exists that is also a Lucas-Carmichael number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tim Johannes Ohrtmann_, May 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009651 | OEIS-A257778

Smallest k, such that n is a Belgian-k number.

conjecture: a(n) < 9;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, May 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009652 | OEIS-A257810

Smallest number of the cycle in which n ends under iteration of sum-of-the-square-of-two-digits.

The following statements, densities and conjectures are based on calculations for n = 1..10000. It is conjectured that iterations of s_2 always end in cycles of finite period length and besides the 1-cycles for fixed points and the bihappy numbers there are ten different cycles of length > 1. The period lengths are: 2, 2, 4, 5, 5, 6, 10, 14, 35 or 56.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pieter Post_, May 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009653 | OEIS-A257838

Main diagonal of iterated partial sums array of Fibonacci numbers (starting with the first partial sums).

Conjecture: n*(n-3)*a(n) +2*(-4*n^2+13*n-6)*a(n-1) +(15*n^2-53*n+48)*a(n-2) +2*(2*n-3)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Dec 10 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luciano Ancora_, May 10 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009654 | OEIS-A257856

Least positive integer k such that prime(k*n) - prime(k) is a square.

Conjecture: a(n) exists for any n > 0. In general, every rational number r > 1 can be written as m/n with m > n > 0 such that prime(m) - prime(n) is a square. This conjecture is a supplement to the conjecture in A259712.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009655 | OEIS-A257862

a(n) = n if n <= 3, otherwise a(n) is the smallest number not already used and relatively prime to a(n-1) such that the set of prime divisors of one of the numbers {a(n-2), a(n)} is a subset of the set of prime divisors of the other.

The sequence is an analog of the Yellowstone sequence A098550 with a more stringent condition on neighboriong terms at distance 2. However, we conjecture that, as with A098550, this is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009656 | OEIS-A257883

Sequence (a(n)) generated by Algorithm (in Comments) with a(1) = 0 and d(1) = 0.

Conjecture: if a(1) is a nonnegative integer and d(1) is an integer, then (a(n)) is a permutation of the nonnegative integers (if a(1) = 0) or a permutation of the positive integers (if a(1) > 0). Moreover, (d(n)) is a permutation of the integers if d(1) = 0, or of the nonzero integers if d(1) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 13 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009657 | OEIS-A257886

Least positive m such that floor(n! / (2*(floor(n/2)!))) + m is prime.

Conjecture: No term is composite (similar conjecture to A033932 for a different expression).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Morales Marciel_, May 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009658 | OEIS-A257889

Terms satisfy: a(2*n) = a(n)*b(n) and a(2*n+1) = a(n)*b(n+1) for n>=0 with a(0)=1, where A(x)^2 = Sum_{n>=0} b(n)*x^n and g.f. A(x) = Sum_{n>=0} a(n)*x^n.

5 divides a(3*2^n) for n>=0 (conjecture). 5 does not divide a(3*2^n - 1) for n>=0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 13 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009659 | OEIS-A257905

Sequence (a(n)) generated by Rule 3 (in Comments) with a(1) = 0 and d(1) = 0.

Conjecture: suppose that a(1) is an nonnegative integer and d(1) is an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009660 | OEIS-A257906

Sequence (a(n)) generated by Rule 3 (in Comments) with a(1) = 0 and d(1) = 1.

See A257905 for a guide to related sequences and conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009661 | OEIS-A257907

After 1, the first differences of A257906 (its d-sequence).

See A257905 for a guide to related sequences and conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009662 | OEIS-A257908

Sequence (a(n)) generated by Rule 3 (in Comments) with a(1) = 0 and d(1) = 2.

See A257905 for a guide to related sequences and conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009663 | OEIS-A257909

Sequence (d(n)) generated by Rule 3 (in Comments) with a(1) = 0 and d(1) = 2.

See A257905 for a guide to related sequences and conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009664 | OEIS-A257910

Sequence (a(n)) generated by Rule 3 (in Comments) with a(1) = 0 and d(1) = 3.

See A257905 for a guide to related sequences and conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009665 | OEIS-A257922

Practical numbers m with m-1 and m+1 both prime, and prime(m)-1 and prime(m)+1 both practical.

Conjecture: The sequence contains infinitely many terms. In other words, there are infinitely many positive integers n such that {prime(n)-1, prime(n), prime(n)+1} is a "sandwich of the first kind" (A210479) and {n-1, n, n+1} is a "sandwich of the second kind" (A258838).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009666 | OEIS-A257924

Primes p with p-1, p+1, prime(p)-1 and prime(p)+1 all practical.

Conjecture: The sequence contains infinitely many terms. In other words, there are infinitely many prime numbers p such that {p-1, p, p+1} and {prime(p)-1, prime(p), prime(p)+1} are both "sandwiches of the first kind" (A210479).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 13 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009667 | OEIS-A257926

Least positive integer k such that prime(k*n)+2 = prime(i*n)*prime(j*n) for some 0 < i < j.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009668 | OEIS-A257928

Least prime p such that pi(p*n) = pi(q*n)*pi(r*n) for some primes q and r with p > q > r, where pi(x) denotes the number of primes not exceeding x.

Conjecture: a(n) exists for any n > 0. Also, for each positive integer n there are distinct primes p, q and r such that pi(p*n) = pi(q*n) + pi(r*n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 13 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009669 | OEIS-A257938

Least positive integer k such that prime(k*n) - 1 = (prime(i*n)-1)*(prime(j*n)-1) for some integers 0 < i < j < k.

Conjecture: a(n) exists for any n > 0. In general, for any nonzero integer m and positive integer n, the set {prime(k*n)+m: k = 1,2,3,...} always contains three distinct elements x, y and z with x*y = z.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 13 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009670 | OEIS-A257948

Length of cycle in which n ends under iteration of sum-of-squares-of-two-digits map s_2.

The following statements, densities and conjectures are based on calculations for n = 1...10000. It is conjectured that iterations of s_2 always end in cycles of finite period length and besides the 1-cycles there are ten different cycles of length > 1. The period lengths are 2, 2, 4, 5, 5, 6, 10, 14, 35 or 56.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pieter Post_, May 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009671 | OEIS-A257949

The q-series expansion of an expression related to fermionic systems.

Conjecture: a linear combination of modular forms, possibly of different weights.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David McGady_, May 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009672 | OEIS-A257956

Row sums of A232642, when seen as a triangle read by rows.

Conjectures from _Colin Barker_, May 29 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, May 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009673 | OEIS-A257979

Smallest prime p for which exactly n primes k with k < p exist such that F_p-(p/k) == 0 (mod p), where F_i = A000045(i) and (a/b) denotes the Legendre symbol, or 0 if no such p exists.

Conjecture: a(9) = 0 (based on observation of the asymptotic behavior of A257978).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, May 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009674 | OEIS-A257980

Sequence (d(n)) generated by Rule 3 (in Comments) with a(1) = 0 and d(1) = 3.

See A257905 for a guide to related sequences and conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009675 | OEIS-A257981

Sequence (a(n)) generated by Rule 3 (in Comments) with a(1) = 1 and d(1) = 1.

See A257905 for a guide to related sequences and conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009676 | OEIS-A257982

Sequence (d(n)) generated by Rule 3 (in Comments) with a(1) = 1 and d(1) = 1.

See A257905 for a guide to related sequences and conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009677 | OEIS-A257983

Sequence (a(n)) generated by Rule 3 (in Comments) with a(1) = 1 and d(1) = 2.

See A257905 for a guide to related sequences and conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009678 | OEIS-A257985

Sequence (a(n)) generated by Rule 3 (in Comments) with a(1) = 2 and d(1) = 0.

See A257905 for a guide to related sequences and conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009679 | OEIS-A257986

Sequence (a(n)) generated by Rule 3 (in Comments) with a(1) = 2 and d(1) = 1.

See A257905 for a guide to related sequences and conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009680 | OEIS-A257987

Sequence (a(n)) generated by Rule 3 (in Comments) with a(1) = 2 and d(1) = 2.

See A257905 for a guide to related sequences and conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009681 | OEIS-A258021

Eventual fixed point of map x -> floor(tan(x)) when starting the iteration with the initial value x = n.

Pohjola conjectures that no other numbers than 0 and 1 will ever occur in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V.J. Pohjola_ & _Antti Karttunen_, May 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009682 | OEIS-A258022

Nonnegative integers n with property that when starting from x=n, the map x -> floor(tan(x)) reaches [the fixed point] 0 (or any other integer less than 1 if such negative fixed points exist).

V.J. Pohjola conjectures that the only fixed points of function k -> floor(tan(k)) are 0 and 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V.J. Pohjola_ & _Antti Karttunen_, May 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009683 | OEIS-A258024

Natural numbers n such that the iteration of the function floor(tan(k)) applied to n eventually reaches [the fixed point] 1 (or any larger integer if such fixed points exist), where k is interpreted as k radians.

It is conjectured that applying the function floor(tan) k times, with k sufficiently large, on the finite sequence floor(tan(n)), n=0..N, the result is a sequence (cf. A258021) composed only of 0's and 1's for all values of N. The original definition was: "Numbers n with property that floor(tan(n)) reduces to 1 (instead of 0) when the function is applied repeatedly to n with deep enough nesting level." If the conjecture above is true, then the new, in theory more inclusive definition produces exactly the same sequence. It has been checked that for at least up to A249836(13) = 1108341089274117551 there are no other strictly positive fixed points beside 1. - _Antti Karttunen_, May 26 2015 According to _Jan Kristian Haugland_ (cf. link): It is an open problem whether (tan n) > n for infinitely many n, although it has been proved that |tan n| > n for infinitely many n. - _Daniel Forgues_, May 27 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V.J. Pohjola_, May 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009684 | OEIS-A258037

Numbers k such that D(prime(k), k-1) > 0, where D(* , k-1) = (k-1)-st difference.

Conjecture: all the terms of the difference sequence of A258037 belong to {1,2,3}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009685 | OEIS-A258046

Sequence (a(n)) generated by Rule 3 (in Comments) with a(1) = 1 and d(1) = 0.

See A257905 for a guide to related sequences and conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009686 | OEIS-A258047

Sequence (d(n)) generated by Rule 3 (in Comments) with a(1) = 1 and d(1) = 0.

See A257905 for a guide to related sequences and conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009687 | OEIS-A258076

a(0)=1, a(1)=2; for n>=2, a(n) = smallest number not already used which shares a factor with a(n-1) if T(n-1)=T(n), and otherwise is relatively prime to a(n-1), where {T(n)} is Thue-Morse sequence (A010060).

Conjecture: the sequence is a permutation of the positive integers. Conjecture: the primes appear in the natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009688 | OEIS-A258105

Inverse to sequence defined as {b(1) = 1, and for n>1: b(n) = A257906(n)}, which is conjectured to be a permutation of natural numbers.

Inverse to sequence defined as {b(1) = 1, and for n>1: b(n) = A257906(n)}, which is conjectured to be a permutation of natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, May 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009689 | OEIS-A258107

Smallest number > 1 whose representation in all bases up to n consists only of 0's and 1's.

As with A146025, it is a plausible conjecture that there are no more terms, but this has not been proved. - _Daniel Mondot_, Dec 16 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernardo Boncompagni_, May 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009690 | OEIS-A258139

Number of ways to write n as p^2 + q with p and q both prime.

Conjecture: For any integer n > 0, we have a(n+r) > 0 for some r = 0,1,2,3,4,5. Moreover, if n = 6*k + 2, then a(n) > 0 except for k = 0, 1, 12, 28, 102, 117, 168, 4079. We have verified the conjecture for n up to 10^9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009691 | OEIS-A258140

Number of ways to write 6*n + 2 as p^2 + q with p and q both prime.

Conjecture: a(n) > 0 except for n = 0, 1, 12, 28, 102, 117, 168, 4079.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009692 | OEIS-A258153

Numbers of the form p^2 + q with p, q and 2*p + 3 all prime.

The conjecture in A258141 asserts that any six consecutive positive integers contain at least a term of the current sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009693 | OEIS-A258157

Indices of the start of 10 successive distinct digits in the decimal expansion of Pi.

It is natural to conjecture that a(n) ~ 1562500n/567. - _Charles R Greathouse IV_, May 22 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Barker_, May 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009694 | OEIS-A258158

Indices of the start of 9 successive distinct digits in the decimal expansion of Pi.

It is natural to conjecture that a(n) ~ 156250n/567. - _Charles R Greathouse IV_, May 22 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Barker_, May 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009695 | OEIS-A258168

Number of ways to write n as floor((p^2+q)/5) with p and q both prime.

Conjecture: Let n be any positive integer. Then a(n) > 0. Moreover, one of the four consecutive numbers 5*n, 5*n+1, 5*n+2, 5*n+3 can be written as p^2+q with p and q both prime. See also A258141 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009696 | OEIS-A258170

T(n,k) = (1/k!) * Sum_{i=0..k} (-1)^(k-i) * C(k,i) * A185651(n,i); triangle T(n,k), n >= 0, 0 <= k <= n, read by rows.

Conjecture 1: T(n,k) = Stirling2(n,k) for k >= 1 and k <= n <= 2*k - 1. Conjecture 2: T(n,k) = Stirling2(n,k) for k >= 2 and n prime >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, May 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009697 | OEIS-A258173

Sum over all Dyck paths of semilength n of products over all peaks p of y_p, where y_p is the y-coordinate of peak p.

Conjecture: a(n) = A371567(n-1,0) for n > 0 with a(0) = 1. - _Mikhail Kurkov_, Nov 07 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, May 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009698 | OEIS-A258195

Define a<+>b = odd part(odd part(a) + odd part(b)), where odd part(n) = A000265(n); a(n) is not already appeared prime of the form <+>_{1<=i<=k}i with the smallest k, or a(n)=0 if there is no such k (see comment).

Conjecture: the sequence is a permutation of the odd primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009699 | OEIS-A258219

A(n,k) is the sum over all Dyck paths of semilength n of products over all peaks p of (x_p+k*y_p)/y_p, where x_p and y_p are the coordinates of peak p; square array A(n,k), n>=0, k>=0, read by antidiagonals.

Conjecture: the g.f. G(k,x) for the k-th column satisfies the Riccati differential equation 2*x^2*d/dx(G(k,x)) + 1 + (k*x - 1)*G(k,x) + x*G^2(k,x) = 0 and hence, by Stokes 1982, has the continued fraction representation G(k,x) = 1/(1 - (k+1)*x/(1 - 3*x/(1 - (k+3)*x/(1 - 5*x/(1 - (k+5)*x/(1 - 7*x/(1 - ...)))))) of Stieltjes type. - _Peter Bala_, Jul 28 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, May 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009700 | OEIS-A258328

L.g.f.: log(1 + Sum_{n>=1} x^(n^2) + x^(3*n^2)).

a(n) = -1 iff n = 2^k for k>=1 [conjecture]. a(p) = +1 for primes p such that 3 is not a square mod p (A003630), and a(n) = +1 nowhere else except at n=0 [conjecture].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009701 | OEIS-A258333

Number of (primitive) weird numbers of the form 2^n*p*q, with odd primes p < q.

However, p is usually not unique, e.g., for k=3, p=19 we have two pwn (with q=61 and q=71), and for k=5, p=71 yields two pwn (for q=523 and q=541) and p=67 yields three pwn (for q=887, 971 and 1021). I conjecture that there is an increasing number of pwn with, e.g., p=nextprime(2^(k+1)). Also, if 2^k p q and 2^k p' q are both weird, then usually 2^k p" q is weird for all p" between p and p'. There is one exception [p, p', q] = [2713, 2729, 8191] for k=10, five exceptions [6197, 6203, 12049], [6113, 6131, 12289], [6113, 6131, 12301], [6121, 6133, 12323], [5441, 5449, 16411] for k=11, and seven exceptions for k=12. These exceptions occur when q/p is close to an integer, (p, q) ~ (3/4, 3/2)*2^(k+2) or (2/3, 2)*2^(k+2). - _M. F. Hasler_, Jul 16 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Douglas E. Iannucci_ and _Robert G. Wilson v_, May 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009702 | OEIS-A258338

Ternary ménage problem: number of seating arrangements for n opposite-sex couples around a circular table such that no spouses and no triples of the same sex seat next to each other. Seats are labeled.

Conjecture: (a(n)/n!^2)^(1/n) ~ (3+sqrt(5))/2. - _Vaclav Kotesovec_, May 29 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, May 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009703 | OEIS-A258369

Stirling-Bernoulli transform of A027656.

Conjecture: inverse binomial transform of A005649. - _Mikhail Kurkov_, Mar 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philippe Delélham_, May 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009704 | OEIS-A258429

Primes p such that p - 1 = (tau(p - 1) - 1)^k for some k >= 0, where tau(n) is the number of divisors of n (A000005).

Conjecture: the sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, May 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009705 | OEIS-A258490

Number of words of length 2n such that all letters of the ternary alphabet occur at least once and are introduced in ascending order and which can be built by repeatedly inserting doublets into the initially empty word.

Conjecture: 4*n*(n-1)*(46829*n-161203)*a(n) -(n-1)*(4865671*n^2-22433759*n+19821114)*a(n-1) +6*(7756949*n^3-53792553*n^2+117956226*n-84118712)*a(n-2) +(-200071007*n^3+1677158106*n^2-4623144589*n+4201946850)*a(n-3) +2*(2*n-7)*(93171685*n^2-585009841*n+881711802)*a(n-4) -72*(2*n-7)*(2*n-9)*(744719*n-1901876)*a(n-5)=0. - _R. J. Mathar_, Aug 07 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, May 31 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009706 | OEIS-A258580

Least positive integer k such that (prime(j*n)+prime(k*n))/2 = prime(i*n)^2 for some integers i > 0 and 0 < j < k.

Conjecture: a(n) exists for any n > 0. In general, for any positive integers a, m and n, there are integers i,j,k > 0 with i > j such that (prime(i*n)+prime(j*n))/2 (or (prime(i*n)-prime(j*n))/2) is equal to a*prime(k*n)^m.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009707 | OEIS-A258655

a(n) = A256357(n^2), where exp(Sum_{n>=1} A256357(n)*x^n/n) = 1 + Sum_{n>=1} x^(n^2) + x^(2*n^2).

a(4*n) < 0 for n>=1, and a(n) is positive if n is not divisible by 4 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009708 | OEIS-A258661

Positive integers m such that none of the four consecutive numbers m, m+1, m+2, m+3 can be written as p^2 + q with p and q both prime.

Conjecture: Any term not among 1, 2, 1009, 15051, 15052, 21772 has the form 36*k^2-2 or the form 36*k^2-3, where k is a positive integer. Note that this conjecture implies the conjecture in A258168 since neither 36*k^2-2 nor 36*k^2-3 can be a multiple of 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009709 | OEIS-A258663

Numbers n such that 9n-1 is prime.

It is my conjecture that the integer formed by the repeating digits in the decimal fraction a(n)/(a(n)*9-1) is the smallest integer such that rotating the digits to the left produces a number which is ((a(n)+1)/a(n)) times larger. An additional conjecture is that the values x in this sequence are the only values where rotating an integer one to the left produces a value (x+1)/x times as large. For example, the conjecture is that there are integers i that when rotated one to the left produce the value 3i/2, 7i/6 and 9i/8, but none that produce the value 2i/1, 4i/3, 5i/4, 6i/5 or 8i/7. The second conjecture is false. For example, 225806451612903*(8/7) = 258064516129032, or 45 * (6/5) = 54 or 230769*(4/3)=307692. - _Giovanni Resta_, Jul 28 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Doug Bell_, Jun 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009710 | OEIS-A258664

A total of n married couples, including a mathematician M and his wife, are to be seated at the 2n chairs around a circular table, with no man seated next to his wife. After the ladies are seated at every other chair, M is the first man allowed to choose one of the remaining chairs. The sequence gives the number of ways of seating the other men, with no man seated next to his wife, if M chooses the chair that is 3 seats clockwise from his wife's chair.

Therefore, it is natural to conjecture that a(n) ~ e^(-2)*n!/(n-2)*(1 + Sum_{k>=1}(-1)^k/(k!(n-1)_k)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jun 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009711 | OEIS-A258665

A total of n married couples, including a mathematician M and his wife, are to be seated at the 2n chairs around a circular table, with no man seated next to his wife. After the ladies are seated at every other chair, M is the first man allowed to choose one of the remaining chairs. The sequence gives the number of ways of seating the other men, with no man seated next to his wife, if M chooses the chair that is 5 seats clockwise from his wife's chair.

Therefore, it is natural to conjecture that a(n) ~ e^(-2)*n!/(n-2)*(1 + Sum_{k>=1} (-1)^k/(k!(n-1)_k)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jun 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009712 | OEIS-A258666

A total of n married couples, including a mathematician M and his wife, are to be seated at the 2n chairs around a circular table, with no man seated next to his wife. After the ladies are seated at every other chair, M is the first man allowed to choose one of the remaining chairs. The sequence gives the number of ways of seating the other men, with no man seated next to his wife, if M chooses the chair that is 7 seats clockwise from his wife's chair.

Therefore, it is natural to conjecture that a(n) ~ e^(-2)*n!/(n-2)*(1 + Sum_{k>=1} (-1)^k/(k!(n-1)_k)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jun 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009713 | OEIS-A258673

A total of n married couples, including a mathematician M and his wife, are to be seated at the 2n chairs around a circular table, with no man seated next to his wife. After the ladies are seated at every other chair, M is the first man allowed to choose one of the remaining chairs. The sequence gives the number of ways of seating the other men, with no man seated next to his wife, if M chooses the chair that is 11 seats clockwise from his wife's chair.

Therefore, it is natural to conjecture that a(n) ~ e^(-2)*n!/(n-2)*(1 + Sum_{k>=1} (-1)^k/(k!(n-1)_k)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Jun 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009714 | OEIS-A258778

Least base b >= 2 such that prime(n) is an absolute prime in base b.

Conjecture: a(n) < prime(n)-1 for n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Jun 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009715 | OEIS-A258802

Least base b >= 2 such that prime(n) is an absolute prime in base b with at least 2 distinct digits or 0 if no such base exists.

Conjecture: a(n) = 0 if and only if n=1 or n=2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Jun 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009716 | OEIS-A258803

Least prime p such that n = (prime(q)-1)/(prime(p)-1) for some prime q.

Conjecture: a(n) exists for any n > 0. Moreover, for any integers s and t with |s| = |t| = 1, each positive rational number r can be written as (prime(p) + s)/(prime(q) + t) with p and q both prime. - Sun I have verified the conjecture for all those rational numbers r = a/b with a, b = 1, ..., 500. - _Zhi-Wei Sun_, Jun 13 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 10 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009717 | OEIS-A258838

Practical numbers q with q-1 and q+1 twin primes: "Sandwiches of the second kind".

The author introduced two kinds of "sandwiches" in 2013. The conjecture in A258836 essentially says that {a(m)/a(n): m,n = 1,2,3,...} coincides with the set of all positive rational numbers. This implies that the sequence contains infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009718 | OEIS-A258876

Integers k such that both k and prime(k) have the same digital root.

Conjecture: a(n) ~ 9n. - _Charles R Greathouse IV_, Jun 17 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Jun 13 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009719 | OEIS-A258900

E.g.f.: S(x) = Series_Reversion(Integral 1/(1-x^4)^(1/4) dx), where the constant of integration is zero.

Conjecture: c = Gamma(1/3) / (2^(8/3) * 3^(1/6) * Pi^(13/6)) = 0.029412898727575426922981691... (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009720 | OEIS-A258902

Expansion of e.g.f. Series_Reversion(x - x^2/2 - x^3/3).

Conjecture: +19*a(n) +21*(-2*n+3)*a(n-1) -4*(3*n-5)*(3*n-7)*a(n-2)=0. - _R. J. Mathar_, Jun 07 2016 a(n) = ((n-1)! / 2^(n-1)) * sum_{k=0..floor((n-1)/2)} binomial(2n-2-k, n-1-k) * binomial(n-1-k, k) * (4/3)^k. Furthermore, Mathar's conjecture is true (see Fried link). - _Sela Fried_, Dec 07 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009721 | OEIS-A258924

E.g.f.: S(x) = Series_Reversion(Integral 1/(1-x^5)^(1/5) dx), where the constant of integration is zero.

Conjecture: c = Gamma(1/4) / (2^(29/8) * (5 - sqrt(5))^(1/8) * Pi^(11/4)) = 0.011112052155593726141351 19... (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009722 | OEIS-A258945

Decimal expansion of Dickman's constant C_4.

Also (conjecturally) equals Pi^4/1440. Conjecture: Equals (1/10) * Sum_{i,j >= 1} (-1)^(i+j) / (i*j)^2 (which equals Pi^4/1440). - _Amiram Eldar_, Jul 04 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Jun 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009723 | OEIS-A258969

E.g.f.: A'(x) = 1 + A(x)^3, with A(0)=1.

Conjecture: A227250(n+1) = a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jun 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009724 | OEIS-A259032

Numbers k such that digital root of k is 2*(digital root of k-th prime).

Conjecture: a(n) ~ 18n. - _Charles R Greathouse IV_, Jun 18 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Jun 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009725 | OEIS-A259040

Numbers k such that digital root of k is 3*(digital root of k-th prime).

Conjecture: a(n) ~ 27n. - _Charles R Greathouse IV_, Jun 18 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Jun 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009726 | OEIS-A259041

Numbers k such that digital root of k is 4*(digital root of k-th prime).

Conjecture: a(n) ~ 27n. - _Charles R Greathouse IV_, Jun 18 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Jun 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009727 | OEIS-A259056

a(n) gives the determinant of a bisymmetric n X n matrix involving the entries 1, 2, ..., A002620(n+1).

Conjecture: a(2*n-1) = (-1)^(n-1)*A034976(n), n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jul 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009728 | OEIS-A259057

One-third of the even-indexed entries of A259056.

It is conjectured that this is an integer sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jul 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009729 | OEIS-A259081

a(n) = largest k such that the decimal representation of 3^k does not contain the digit n.

These values are only conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009730 | OEIS-A259082

a(n) = largest k such that the decimal representation of 5^k does not contain the digit n, or -1 if every power of 5 contains the digit n.

These values are only conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009731 | OEIS-A259083

a(n) = largest k such that the decimal representation of 7^k does not contain the digit n.

These values are only conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009732 | OEIS-A259084

a(n) = largest k such that the decimal representation of prime(n)^k does not contain the digit 0.

These values are only conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009733 | OEIS-A259085

a(n) = largest k such that the decimal representation of prime(n)^k does not contain the digit 1, or -1 if no such k exists.

These values are only conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009734 | OEIS-A259086

a(n) = largest k such that the decimal representation of prime(n)^k does not contain the digit 2.

These values are only conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009735 | OEIS-A259145

Numbers k such that k^2 - phi(k) is prime, where phi() is A000010.

Conjecture: a(n) is a cyclic number (see A003277) for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carlos Eduardo Olivieri_, Jun 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009736 | OEIS-A259148

Decimal expansion of phi(exp(-Pi)), where phi(q) = Product_{n>=1} (1-q^n) is the Euler modular function.

Conjecture: Equals Sum_{k>=0} A010815(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Jun 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009737 | OEIS-A259207

5x + 1 sequence beginning at 5.

However, there are still unanswered questions about the 5x + 1 problem. Kontorovich and Lagarias (2009) say that it is conjectured that there are very few periodic orbits, one of which is the one exhibited by this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alonso del Arte_, Jun 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009738 | OEIS-A259215

Number of (n+1) X (1+1) 0..1 arrays with each 2 X 2 subblock having clockwise pattern 0000 0011 or 0101.

Conjectures from _Colin Barker_, Dec 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009739 | OEIS-A259226

Numbers n such that A157480(n) = A259232(n).

It is a strong conjecture that the sequence is complete, and the last term is a(99)=9867.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Jun 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009740 | OEIS-A259231

Primitive numbers whose abundance is positive and odd.

See A156903 for the properties of those numbers. Pavlov's 2017 comment below makes observations of some interest, but its conjecture is false -- see _Amiram Eldar_'s comment in A156942 for the first of its terms that is not divisible by 3. - _Peter Munn_, Oct 01 2025 Conjecture: let m == a(n) mod 2. Then a(n) can be written as (2+m)^t * d^2 where t is an integer, t > 0, d is odd, d > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jun 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009741 | OEIS-A259243

Number of (n+1) X (1+1) 0..1 arrays with each 2 X 2 subblock having clockwise pattern 0000 0011 or 0111.

Conjectures from _Colin Barker_, Dec 24 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009742 | OEIS-A259260

With a(1) = 1, a(n) is the smallest number not already in the sequence such that the arithmetic mean of two neighboring terms is a square.

Conjectured to be a permutation of the odd numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Jun 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009743 | OEIS-A259268

G.f. satisfies: A(x) = Series_Reversion(x - 3*A(x)*B(x)^2), where B'(x) = 2*A(x).

Conjecture: (2*n-1) | a(n) for n>=1 and therefore all the terms of this sequence are integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009744 | OEIS-A259269

G.f. A(x) satisfies: A'(x)/2 = Series_Reversion(x - 3*A(x)^2 * A'(x)/2).

a(n) == 1 (mod 3) iff (2*n-1) is a power of 3, and a(n) == 0 (mod 3) elsewhere (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009745 | OEIS-A259270

G.f. satisfies: A(x) = Series_Reversion(x - 2*A(x)*B(x)), where B'(x) = 2*A(x).

Conjecture: n | a(n) for n>=1 and therefore all the terms of this sequence are integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009746 | OEIS-A259271

G.f. A(x) satisfies: A'(x) = 2 * Series_Reversion(x - A(x)*A'(x)).

a(n) == 1 (mod 2) iff n is a power of 2 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009747 | OEIS-A259290

Number of (n+1) X (1+1) 0..1 arrays with each 2 X 2 subblock having clockwise pattern 0001 0101 or 0111.

Conjectures from _Colin Barker_, Dec 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jun 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009748 | OEIS-A259301

Taken over all those prime-partitionable numbers m for which there exists a 2-partition of the set of primes < m that has one subset containing two primes only, a(n) is the frequency with which the smaller prime occurs, where n is the prime index.

Conjecture: If P1 = {p1a, p1b} with p1a and p1b odd primes, p1a < p1b and p1b = 2*k*p1a + 1 for some positive integer k such that 2*k <= p1a - 3 and if m = p1a + p1b then m is prime-partitionable.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christopher Hunt Gribble_, Jun 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009749 | OEIS-A259440

Field discriminant of n-th composite, f(f(...f(r)...)), where r = 2 and f(x) = [x,x,x, ...] (continued fraction).

Conjecture: a(n+1) is divisible by a(n)^2, for n>=1; see Example.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009750 | OEIS-A259457

From higher-order arithmetic progressions.

Conjecture: 3*n*a(n) +(-3*n^2-19*n-44)*a(n-1) -2*(n+2)^2*a(n-2)=0. - _R. J. Mathar_, Jul 15 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009751 | OEIS-A259484

Smallest nonprime number having least positive primitive root n, or 0 if no such root exists.

Conjecture: each odd-indexed value will be populated before either of its even-indexed neighbors.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jun 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009752 | OEIS-A259487

Least positive integer m with prime(m)+2 and prime(prime(m))+2 both prime such that prime(m*n)+2 and prime(prime(m*n))+2 are both prime.

Conjecture: Any positive rational number r can be written as m/n with m and n terms of A259488. I have verified the conjecture for all those r = a/b with a,b = 1,...,400. - _Zhi-Wei Sun_, Jun 29 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009753 | OEIS-A259488

Positive integers k with prime(k)+2 and prime(prime(k))+2 both prime.

The conjecture in A259487 essentially says that {a(m)/a(n): m,n = 1,2,3,...} coincides with the set of all positive rational numbers. This implies that the current sequence has infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009754 | OEIS-A259503

Numbers k such that k^2+1 is the product of a Fibonacci number and a Lucas number.

Conjecture: the sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jun 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009755 | OEIS-A259504

Numbers k such that k and k+1 are the product of exactly three (not necessarily distinct) primes.

Conjecture: this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Nov 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009756 | OEIS-A259531

Least positive integer k such that p(k)^2 + p(k*n)^2 is prime, where p(.) is the partition function given by A000041, or 0 if no such k exists.

Conjecture: Any positive rational number r can be written as m/n, where m and n are positive integers with p(m)^2 + p(n)^2 prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009757 | OEIS-A259540

Least positive integer k such that k and k*n are terms of A259539.

Conjecture: Any positive rational number r can be written as m/n with m and n terms of A259539.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009758 | OEIS-A259542

a(1) = 1, for n > 1 a(n) = smallest number not already in the sequence such that the arithmetic mean of two neighboring terms is a triangular number.

conjecture: sequence is a permutation of the odd numbers, see also A259260, A259429;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jun 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009759 | OEIS-A259544

Minimum greatest integer in a set of n positive integers whose nonempty subsets all have distinct arithmetic means.

Conjecture: The only term that is < 2^(n-1) is a(4)=7. Conjecture: lim a(n)/2^n = infinity. (Note that this is weaker than lim_inf a(n)^(1/n) > 2.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Javier Múgica_, Jun 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009760 | OEIS-A259560

Primes p such that p = 2kq + 1 for k a positive integer, q an odd prime and 2k <= q - 3.

This sequence is associated with the conjecture in A245664 that p + q is prime-partitionable.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christopher Hunt Gribble_, Jun 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009761 | OEIS-A259565

a(1) = 1, for n > 1 a(n) = smallest number not already in the sequence such that the arithmetic mean of two neighboring terms is a squarefree number.

conjecture: sequence is a permutation of the odd numbers;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Jun 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009762 | OEIS-A259569

Triangle T(n,k) read by rows, where T(n,k) is the number of k-dimensional faces of the polytope that is the convex hull of all permutations of the list (0,1,...,1,2), where there are n - 1 ones, for n > 0. T(0,0) is 1.

Conjecture: Sum_{k=0..n-1} T(n,k)*x^(n-k-1) = x^(n+1) - 2(x+1)^(n+1) + (x+2)^(n+1). - _Kevin J. Gomez_, Jul 25 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincent J. Matsko_, Jun 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009763 | OEIS-A259598

Number of representations of n as u(h) + v(k), where u = A000201 (lower Wythoff numbers), v = A001950 (upper Wythoff numbers), h>=1, k>=1.

Three conjectures. The numbers that are not a sum u(h) + v(k) are (1,2,4,7,12, ...) = A000071 = -1 + Fibonacci numbers. The numbers that have exactly one such representation are (3, 5, 9, 15, 25, 41, ...) = A001595. The numbers that have exactly two such representations are (6, 10, 17, 28, 46, ...) = A001610.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009764 | OEIS-A259628

Numbers m with m-1, m+1, prime(m)+2, prime(m)-m, prime(m)+m, m*prime(m)-1 and m*prime(m)+1 all prime.

Conjecture: The sequence contains infinitely many terms. This is stronger than the conjectures in A232861 and A259539.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009765 | OEIS-A259661

Binary representation of the middle column of the "Rule 54" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 08 2015 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009766 | OEIS-A259667

Catalan numbers mod 6.

It is conjectured that the only k which yield a(2^k-1) = 1 are k = 0, 1 and 5. Are there other k than 2 and 8 that yield a(2^k-1) = 5? Otherwise said, is a(2^k-1) = 3 for all k > 8? The question is equivalent to: do 2^k - 1 always contain a digit 2 when converted into base 3 for all k > 8? A similar conjecture has been proposed for 2^k; see A004642. - _Jianing Song_, Sep 04 2018 [Typo corrected by _Jianing Song_, Apr 15 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Nov 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009767 | OEIS-A259672

Number k such that the greatest prime divisor of k+1 is the smallest prime that is larger than the greatest prime divisor of k.

Conjecturally, this subsequence is probably infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jul 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009768 | OEIS-A259678

Primes of the form p(k)^2 + p(m)^2, where k and m are positive integers, and p(.) is the partition function given by A000041.

The conjecture in A259531 implies that the current sequence has infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009769 | OEIS-A259712

Least positive integer k such that prime(k) + prime(k*n) is a square.

Conjecture: a(n) exists for any n > 0. In general, every positive rational number r can be written as m/n, where m and n are positive integers with prime(m) + prime(n) a square. I have verified this conjecture for all those r = a/b with a,b = 1,...,700.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009770 | OEIS-A259714

a(n) = Sum_{k=1..n-1}((k mod 5)*a(n-k)), a(1) = 1.

Conjectures from _Colin Barker_, Jul 04 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anders Hellström_, Jul 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009771 | OEIS-A259724

Numbers k such that [r[s*k]] < [s[r*k]], where r = sqrt(2), s=sqrt(3), and [ ] = floor.

Conjecture: the limits g = lim G(n)/n, e = lim E(n)/n, el = lim L(n)/n exist; if so, then 1/g + 1/e + 1/el = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009772 | OEIS-A259731

Least positive integer k such that prime(k*n)-1 is a square, or 0 if no such k exists.

Conjecture: a(n) > 0 for all n > 0. This is stronger than the well-known conjecture that there are infinitely many primes of the form x^2+1 with x an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009773 | OEIS-A259733

The magic constants of most-perfect magic squares of order 8 composed of distinct prime numbers.

It seems that only the first term, or possibly the first two terms, have been proved to be correct. The other terms are conjectural (that is, there may be missing terms). - _N. J. A. Sloane_, Jul 28 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Natalia Makarova_ and Sergey Zorkin, Jul 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009774 | OEIS-A259762

Smallest integer k_1 such that there exist n positive integers k_1 > k_2 > ... > k_n having the property that k_j * k_n > k_(j+1)^2 for j=1..n-1.

This is one of the orderings of the pairwise products of real numbers in A237749. Conjecture: if we constrain those real numbers to take integer values, then all A237749(n) orderings of pairwise products can be obtained with k_1 = a(n), but this ordering cannot be obtained with k_1 < a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Jul 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009775 | OEIS-A259764

Least prime p such that prime(p*n)-1 is a square, or 0 if no such p exists.

Conjecture: a(n) > 0 for all n > 0. This is stronger than the conjecture in A259731. It implies the well-known conjecture that there are infinitely many primes of the form x^2-1 with x an integer. I also conjecture that for any positive integer n there exists a prime p such that prime(p*n)+2 is a square.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009776 | OEIS-A259775

Stepped path in P(k,n) array of k-th partial sums of squares (A000290).

Conjecture: -(n+5)*(13*n-11)*a(n) +(8*n^2+39*n-35)*a(n-1) +2*(26*n^2+48*n+25)*a(n-2) -4*(8*n+5)*(n-1)*a(n-3)=0. - _R. J. Mathar_, Jul 16 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luciano Ancora_, Jul 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009777 | OEIS-A259776

Number A(n,k) of permutations p of [n] with no fixed points and displacement of elements restricted by k: 1 <= |p(i)-i| <= k, square array A(n,k), n>=0, k>=0, read by antidiagonals.

Conjecture: Column k > 0 has a linear recurrence (with constant coefficients) of order = A005317(k) = (2^k + C(2*k,k))/2. - _Vaclav Kotesovec_, Jul 07 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009778 | OEIS-A259780

Number of permutations p of [n] with no fixed points and displacement of elements restricted by seven: 1 <= |p(i)-i| <= 7.

Conjectured recurrence order is 1780. - _Vaclav Kotesovec_, Jul 07 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009779 | OEIS-A259781

Number of permutations p of [n] with no fixed points and displacement of elements restricted by eight: 1 <= |p(i)-i| <= 8.

Conjectured recurrence order is 6563. - _Vaclav Kotesovec_, Jul 07 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009780 | OEIS-A259782

Number of permutations p of [n] with no fixed points and displacement of elements restricted by nine: 1 <= |p(i)-i| <= 9.

Conjectured recurrence order is 24566. - _Vaclav Kotesovec_, Jul 07 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009781 | OEIS-A259783

Number of permutations p of [n] with no fixed points and displacement of elements restricted by ten: 1 <= |p(i)-i| <= 10.

Conjecture: Column k > 0 of A259776 has a linear recurrence (with constant coefficients) of order = A005317(k) = (2^k + C(2*k,k))/2. For k=10 is conjectured recurrence order 92890. - _Vaclav Kotesovec_, Jul 07 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009782 | OEIS-A259788

Greatest prime factor of phi(binomial(2*n,n)).

Conjectures: All these conjectures follow from the following strengthening of the Bertrand postulate for n>=24: the interval (n, 2*n) contains a safe prime (A005385).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jul 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009783 | OEIS-A259789

Least integer k > 1 such that pi(k)*pi(k*n) is a square, where pi(.) is the prime-counting function given by A000720.

Conjecture: a(n) exists for any n > 0. In general, every positive rational number r can be written as m/n, where m and n are positive integers with pi(m)*pi(n) a positive square.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009784 | OEIS-A259899

a(n) is the minimal position at which the maximal value of row n appears in row n of triangle A080936.

Conjecture: a(n) = floor(sqrt(p*n+q)+r) for all n>=1, where p = 2.67996... = A265179^2 and q,r are some constants (best values found: q=3.6, r=-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009785 | OEIS-A259914

Staircase path through the array P(n,k) of the k-th partial sums of cubes (A000578).

Conjecture: 2*(n+7)*(145672*n^2-236343*n+123525)*a(n) +(-78613*n^3-794662*n^2+327391*n+20220)*a(n-1) +2*(-582688*n^3-1889455*n^2-2148719*n-832650)*a(n-2) +4*(n-1)*(78613*n^2+133361*n+64050)*a(n-3) = 0. - _R. J. Mathar_, Jul 16 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luciano Ancora_, Jul 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009786 | OEIS-A259915

Least positive integer k such that phi(k) and sigma(k*n) are both squares, where phi(.) is Euler's totient function and sigma(m) is the sum of all positive divisors of m.

Conjecture: a(n) exists for any n > 0. In general, every positive rational number r can be written as m/n, where m and n are positive integers with phi(m) and sigma(n) both squares of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009787 | OEIS-A259916

Least positive integer k such that sigma(k) and phi(k*n) are both squares, where sigma(k) is the sum of all positive divisors of k, and phi(.) is Euler's totient function.

The conjecture in A259915 implies that a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009788 | OEIS-A259922

a(n)= Sum_{2 < prime p <= n} c_p - Sum_{n < prime p < 2*n} c_p, where 2^c_p is the greatest power of 2 dividing p-1.

In connection with this, it is natural to conjecture that there exists a number N such that a(n)>0 for all n >= N.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Jul 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009789 | OEIS-A259934

Infinite sequence starting with a(0)=0 such that A049820(a(k)) = a(k-1) for all k>=1, where A049820(n) = n - (number of divisors of n).

Therefore, d(a(k+1) + d(a(k+1))) = d(a(k+2))= d(a(k+1)), i.e., a(k+1) is in A175304. Thus, if there are infinitely many pairs of the same consecutive terms of A259935, then A175304 is infinite (see there my conjecture). (End) If multiple apparently infinite branches would occur at some point of computing, then even if the "lexicographically earliest" condition were then added to the definition, it would not help us much (when computing the sequence), as we would still not know which of the said branches were truly infinite. [See also _Max Alekseyev_'s latter Jul 9 2015 posting on SeqFan-list, where he notes the same thing.] Note that many of the derived sequences tacitly assume that the uniqueness-conjecture is true. See also comments at A262693 and A262896.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Jul 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009790 | OEIS-A259940

Let A={A005574(n)}, the set of all numbers n for which n^2+1 is prime. The sequence lists the number of decompositions A005574(n) = A005574(n1) + A005574(n2) for some n1, n2 and every A005574(n)>1.

We use a little-known conjecture by Goldbach on the primes of form n^2+1: let A be the set of all numbers a for which a^2+1 is prime (A={1, 2, 4, 6, 10, ...}). Then every a in A (a>1) can be written in the form a=b+c for b,c in A.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jul 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009791 | OEIS-A259964

A specially constructed B_2 sequence with sum of reciprocals greater than that of the Mian-Chowla sequence A005282.

It had been conjectured that the sum of reciprocals for B_2 sequences was maximized by A005282. This sequence gives a counterexample.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009792 | OEIS-A260039

Triangle read by rows giving numbers B(n,k) arising in the enumeration of doubly rooted tree maps.

Conjecture: T(n,0) = n*A168452(n-1). - _R. J. Mathar_, Jul 22 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009793 | OEIS-A260040

Triangle read by rows giving numbers H(n,k), number of classes of twin-tree-rooted maps with n edges whose root bond contains k edges.

Conjecture: T(n,n-2) = A005563(n) = 8, 15, 24,.... for n>=2. - _R. J. Mathar_, Jul 22 2015 Conjecture: T(n,n-3)= (n+1)*n*(5*n^2+7*n+6)/12 = 72, 190,.... for n>=3. - _R. J. Mathar_, Jul 22 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009794 | OEIS-A260078

Least positive integer k such that prime(k*n)-1+(prime(h*n)-1) = prime(i*n)-1 and prime(k*n)-1-(prime(h*n)-1) = prime(j*n)-1 for some positive integers h,i,j.

Conjecture: a(n) exists for any n > 0. In general, if m and n > 0 are integers with gcd(6,m) = 1, then the set {prime(k*n)+m: k = 1,2,3,...} contains two distinct elements x and y with x+y and x-y also in the set.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009795 | OEIS-A260080

Least positive integer k such that prime(k*n)^2 - 2 = prime(i*n)*prime(j*n) for some integers 0 < i < j.

Conjecture: a(n) exists for any n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009796 | OEIS-A260082

Least positive integer k such that (prime(k*n)-1)^2 = (prime(i*n)-1)*(prime(j*n)-1) for some integers 0 < i < j.

Conjecture: a(n) exists for any n > 0. In general, for any nonzero integer m and positive integer n there are distinct positive integers i,j,k such that (prime(i*n)+m)*(prime(j*n)+m) = (prime(k*n)+m)^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009797 | OEIS-A260090

Maximum number of kings on an n X n chessboard such that no king attacks more than one other king.

Conjecture: For n != 3, a(n) = n(n+2)/3 + [n mod 3 = 2]/3 - [n mod 6 = 2] Equivalent conjecture for n >= 5: a(n) = a(n-1) + n - A103469(n-2). - _Bob Selcoe_, Jul 17 2015 See A381749 for proofs of the conjectures. - _Yifan Xie_, Mar 22 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rob Pratt_, Jul 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009798 | OEIS-A260113

Maximum number of queens on an n X n chessboard such that no queen attacks more than one other queen.

Taking into account known values, it is reasonable to conjecture that a(n) = floor(4*n/3) for n > 5. - _Giovanni Resta_, Aug 07 2015.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rob Pratt_, Jul 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009799 | OEIS-A260119

a(n) is the least positive integer k such that 2^n-1 and k^n-1 are relatively prime.

According to the conjecture given in A086892, a(n) = 3 infinitely often. Conjecture 1: a(n) is never divisible by 2. Conjecture 2: a(n) is of the form q * m where q is the product of all odd primes that are one more than a divisor of n. Both conjectures are true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tom Edgar_, Jul 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009800 | OEIS-A260120

Least integer k > 0 such that (prime(k*n)-1)^2 = prime(j*n)-1 for some j > 0.

Conjecture: a(n) exists for any n > 0. In general, if a,b,c and m are integers with a > 0, gcd(a,b,c-m) = 1 and c == (a+b+1)*(m+1) (mod 2) such that b^2-4a*(c-m) is not a square and gcd(a*m-b,b^2+b-a*c-1) is not divisible by 3, then for any positive integer n there are two elements x and y of the set {prime(k*n)+m: k = 1,2,3,...} with a*x^2+b*x+c = y. This implies the conjecture in A259731.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009801 | OEIS-A260121

Least positive integer k such that prime(k*n)^2 - 2 = prime(j*n) for some j > 0.

The conjecture in A260120 implies that a(n) exists for any n > 0, which is stronger than the conjecture in A253257.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009802 | OEIS-A260140

Least prime p such that pi(p*n) = pi(q*n)^2 for some prime q, where pi(x) denotes the number of primes not exceeding x.

Conjecture: a(n) exists for any n > 0. In general, for any integers a,b,c,n with a > 0 and n > 0, there are two elements x and y of the set {pi(p*n): p is prime} with a*x^2+b*x+c = y. A supplement to the conjecture: For any integers b,c,n with b > 0 and n > 0, we have b*x+c = y for some elements x and y of the set {pi(p*n): p is prime}. - _Zhi-Wei Sun_, Aug 02 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009803 | OEIS-A260147

G.f. A(x) = Sum_{n=-oo..+oo} x^n * (1 + x^n)^(2*n).

a(2^n) = 1 for n > 0 (conjecture). a(p) = p+1 for primes p > 3 (conjecture). The following are conjectural: Hence, a(p) = p + 1 for primes p > 3 and a(2^n) = 1 for n > 0 as conjectured above. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009804 | OEIS-A260160

a(n) = a(n-2) + a(n-6) - a(n-8) with n>8, the first eight terms are 0 except that for a(5) = a(7) = 1.

a(n) = A264041(n) - n*(n+1)/2, 0<n<=26 (conjectured for n>26).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_ and _Paul Curtz_, Nov 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009805 | OEIS-A260197

Least prime p such that pi(p*n) = prime(q*n) for some prime q, where pi(x) denotes the number of primes not exceeding x.

Conjecture: a(n) exists for any n > 0. Also, for any n > 0, there are primes p and q such that pi(p*n) = q*n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009806 | OEIS-A260208

Least prime p such that 2p*n+1 = prime(q*n) for some prime q.

Conjecture: a(n) exists for any n > 0. In general, if a > 0 is even and b is 1 or -1, then for any positive integer n there are primes p and q such that a*p*n+b = prime(q*n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009807 | OEIS-A260219

Least prime p such that pi(p*n)^2 + 1 = prime(q*n) for some prime q.

Conjecture: a(n) exists for any n > 0. In general, if a,b,c are integers with a > 0 and gcd(a,b,c) = 1, and a+b or c is odd, and b^2 - 4*a*c is not a square, then there are primes p and q such that a*pi(p*n)^2 + b*pi(p*n) + c = prime(q*n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009808 | OEIS-A260252

Least prime p such that n = (prime(6*q)-1)/(prime(6*p)-1) for some prime q.

Conjecture: Let n be any positive integer, and let s and t belong to the set {1,-1}. Then each positive rational number r can be written as (prime(p*n)+s)/(prime(q*n)+t) with p and q both prime, unless n > r = 1 and {s,t} = {1,-1}. This extends the conjecture in A258803.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009809 | OEIS-A260256

Numbers m such that tau(m + 2) = tau(m - 2) where tau(k) = A000005(k).

Pinner proves that this sequence is infinite, and in particular a(n) << n (log n)^7. The correct order is conjectured to be around n sqrt(log n). - _Charles R Greathouse IV_, Jul 21 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jul 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009810 | OEIS-A260310

Pairs with balanced sums of prime divisors (A008472) and inverse prime divisors (A069359), ordered by larger members.

Conjecture: a(2n) < a(2n+2) for all n>0, but there are many times (1/10.84) that a(2n) + 1 = a(2n+2). Conjecture: if a(2n-1) is prime then a(2n) is composite and vice versa. And when a(2n-1) is composite, it is congruent to 0 (mod 6). - _Robert G. Wilson v_, Jul 22 2015 The first conjecture appears to be satisfied because if both x and y were prime then the sum of the A008472 were the sum of the two primes and the sum of the A069359 were two. - _R. J. Mathar_, Aug 03 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jul 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009811 | OEIS-A260311

Difference sequence of A260317.

Conjecture: a(n) is a Fibonacci number (A000045) for every n. Conjectures: n_{2k-1} = n_{2k} and c_{2k-1, j} = c_{2k, j} for all positive k and 0<j<n_{2k}+1. Furthermore c_{2k, j} = c_{2k+1, j} for all positive k and 0<j<n_{2k}. Furthermore c_{i,j} = c_{i,n_i - j} for all positive i and 0<j<n_i + 1. Furthermore (n_{2k})_{k > 0} is the Fibonacci sequence. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 31 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009812 | OEIS-A260332

Labelings of n diamond-shaped posets with 4 vertices per diamond where the labels follow the poset relations whose associated reading permutation avoids 231 in the classical sense.

According to Yang-Jiang (2021) these are the 5-Schroeder numbers. If confirmed, this will prove Michael Weiner's conjectures and enable us to extend the sequence. Yang & Jiang (2021) give an explicit formula for the m-Schroeder numbers in Theorem 2.4. - _N. J. A. Sloane_, Mar 28 2021 Conjecture: Also the number of lattice paths (Schroeder paths) from (0,0) to (n,4n) with unit steps N=(0,1), E=(1,0) and D=(1,1) staying weakly above the line y = 4x. - _Michael D. Weiner_, Jul 24 2019 Conjecture: a(n) = Sum_{k=1..n} binomial(n,k)*binomial(4*n,k-1)*2^k/n for n > 0. - _Michael D. Weiner_, Jul 23 2019 Conjectures: 1) the g.f. A(x) = 1 + 2*x + 18*x^2 + 226*x^3 + ... satisfies A(x)^4 = (1/x) * the series reversion of ((1 - x)/(1 + x))^4. 3) a(n) = 2 * hypergeom([1 - n, -4*n], [2], 2) for n >= 1 (equivalent to Weiner's conjecture above).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Manda Riehl_, Jul 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009813 | OEIS-A260333

Irregular triangle read by rows: T(n,k) = number of ways k brooks (0 <= k <= 2n+1) can be placed on the grid points of an n triboard so that no two brooks lie in the same straight line.

Bennett and Potts give formulas for the first two nontrivial diagonals on the left (A003215 and A047786), and conjectural formulas for the next two diagonals.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009814 | OEIS-A260360

The absolute difference between the largest prime factors of prime(n)-1 and prime(n+1)-1.

It is an open question whether there are infinitely many zeros in this sequence. Are there infinitely many terms below some fixed upper bound?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pratik Koirala_, _Otis Tweneboah_, _Nathan Fox_, _Eugene Fiorini_, Jul 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009815 | OEIS-A260361

G.f.: Sum_{n=-oo..+oo} x^n * (1 + x^n)^n, an even function.

a(2^n) = 2 for n > 0 (conjecture). a(p) = 2*p+2 for primes p > 3 (conjecture). The two conjectures were proved by Bala in A260147. - _Sela Fried_, Dec 08 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009816 | OEIS-A260390

Infinite palindromic word (a(1),a(2),a(3),...) with initial word w(1) = (1,0) and midword sequence (a(n)); see Comments.

w(1) = 0 (see A260455) A260455 A081254 (conjectured)
w(1) = 1 (see A260456) A260456 A081254 (conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 31 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009817 | OEIS-A260406

Numbers n such that (n-1)^2-1 divides 2^(n-1)-1.

The Fermat numbers 2^2^k+1 = A000215(k) with k>1 are a subsequence of this sequence. I conjecture that they are equal to the intersection of this and A260407 (apart from the conventional 1), i.e., the numbers such that (n-1)^4-1 divides 2^(n-1)-1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jul 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009818 | OEIS-A260408

Bisection of A260310.

Conjecture: when a(n) is prime, A260409(n) is composite and vice versa. No contradictions in the first 10000 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jul 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009819 | OEIS-A260409

Bisection of A260310.

Conjecture: when a(n) is composite, A260408(n) is prime and vice versa. No contradictions in the first 10000 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jul 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009820 | OEIS-A260411

Number of ways n can be represented as a sum of a positive cube, a positive square, and a positive triangular number.

It appears that there are 14 zeros and 33 ones. Conjecture: every integer appears in the sequence finitely many times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Jul 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009821 | OEIS-A260418

Number of ways to write 12*n+5 as 4*x^4 + 4*y^2 + z^2, where x is a nonnegative integer, and y and z are positive integers.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 4, 54, 159, 289, 999, 1175, 1404, 16391, 39688. See A290491 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009822 | OEIS-A260457

Field discriminant of n-th composite, f(f(...f(r)...)), where r = 3 and f(x) = [x,x,x, ...] (continued fraction).

Conjecture: a(n+1) is divisible by a(n)^2, for n>=1; see Example.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009823 | OEIS-A260463

a(n) is the smallest number not already in the sequence such that a(n)^2 begins with n.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Jul 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009824 | OEIS-A260465

a(n) is the smallest number not already in the sequence such that a(n)^3 begins with n.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Orr_, Jul 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009825 | OEIS-A260481

Field discriminant of n-th composite, f(f(...f(r)...)), where r = 1 and f(x) is the continued fraction [x,x,x, ...].

Divisibility conjecture: a(n)^2 | a(n+1), for n>=1; see Example.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 13 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009826 | OEIS-A260489

a(n) = 3n - A260488(n).

The sequence is nonnegative for at least the first 10000 terms, so the nonn keyword was used. That it is always nonnegative is a conjecture. That it is unbounded is also a conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Jul 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009827 | OEIS-A260492

Pascal's triangle aerated with columns of zeros.

Conjecturally, the limit of R^n * C^n as n -> oo has as its first column an aerated version of A027826, with zeros elsewhere in the array.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Aug 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009828 | OEIS-A260554

Primes p such that p = q^2 + 4*r^2 where q and r are also primes.

Green and Sawhney prove that this sequence is infinite, verifying a conjecture of Friedlander and Iwaniec. - _Charles R Greathouse IV_, Oct 08 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Barker_, Jul 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009829 | OEIS-A260619

Arithmetic derivative of hyperfactorial(n).

a(n) = A002109(n)*A190121(n) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthew Campbell_, Sep 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009830 | OEIS-A260625

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with (x+3*y+13*z)*x*y*z a square, where x is a positive integer, and y,z,w are nonnegative integers with y >= z.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 7, 39, 47, 95, 191, 239, 327, 439, 871, 1167, 1199, 1367, 1487, 1727, 1751, 2063, 2351, 2471, 4647, 4^k*m (k = 0,1,2,... and m = 1, 3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009831 | OEIS-A260647

Numbers that are the sum of two distinct nonzero triangular numbers.

Conjecture: the sequence contains infinitely many primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Dec 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009832 | OEIS-A260667

a(n) = (1/n^2) * Sum_{k=0..n-1} (2k+1)*S(k,n)^2, where S(k,x) denotes the polynomial Sum_{j=0..k} binomial(k,j)*binomial(x,j)*binomial(x+j,j).

Conjecture: For k = 0,1,2,... define S(k,x):= Sum_{j=0..k} binomial(k,j)*binomial(x,j)*binomial(x+j,j). Our conjecture is motivated by a conjecture of Kimoto and Wakayama which states that Sum_{k=0..p-1} S(k,-1/2)^2 == (-1/p) (mod p^3) for any odd prime p. The Kimoto-Wakayama conjecture was confirmed by Long, Osburn and Swisher in 2014. For more related conjectures, see Sun's paper arXiv.1512.00712. - _Zhi-Wei Sun_, Dec 03 2015 Conjecture: a(p-1) == 1 (mod p^3) for all primes p >= 5. - _Peter Bala_, Aug 15 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009833 | OEIS-A260683

Number of 2's in the expansion of 2^n in base 3.

Erdős conjectures that a(n) > 0 for n > 8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emmanuel Vantieghem_, Nov 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009834 | OEIS-A260708

a(2n) = n*(2*n+1), a(2n+7) = a(2n+1) + 12*n + 28, with a(1)=1, a(3)=6, a(5)=16.

Conjecture: this sequence is 0 followed by A264041.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Nov 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009835 | OEIS-A260753

Least positive integer k such that both k and k*n belong to the set {m>0: prime(prime(m))-prime(m)+1 = prime(p) for some prime p}.

Conjecture: For any s and t in the set {1,-1}, every positive rational number r can be written as m/n with m and n in the set {k>0: prime(prime(k))+s*prime(k)+t = prime(p) for some prime p}. In the case s = -1 and t = 1, the conjecture implies that A261136 has infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009836 | OEIS-A260791

Type II primes (conjectured).

Type II primes (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009837 | OEIS-A260843

Field discriminant of n-th composite, f(f(...f(r)...)), where r = 4 and f(x) = [x,x,x, ...] (continued fraction).

Conjecture: a(n+1) is divisible by a(n)^2, for n>=1; see Example.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009838 | OEIS-A260844

Field discriminant of n-th composite, f(f(...f(r)...)), where r = 5 and f(x) = [x,x,x, ...] (continued fraction).

Conjecture: a(n+1) is divisible by a(n)^2, for n>=1; see Example.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009839 | OEIS-A260876

Number of m-shape set partitions, square array read by ascending antidiagonals, A(m,n) for m, n >= 0.

This result can definitely be used to prove some special cases of my conjecture below. (End) Conjecture: For n >= 0, let P be the set of all possible lists (a_1,...,a_n) of nonnegative integers such that a_1*1 + a_2*2 + ... + a_n*n = n. Consider terms of the form multinomial(n*m, m*[1,..., 1,2,..., 2,..., n,..., n])/(a_1! * a_2! * ... * a_n!), where in the list [1,...,1,2,...,2,...,n,...,n] the number 1 occurs a_1 times, 2 occurs a_2 times, ..., and n occurs a_n times. (Here a_n = 0 or 1.) Summing these terms over P we get A(m, n) provided m >= 1. (End) Conjecture for a recurrence: A(m, n) = Sum_{k = 0..n-1} binomial(m*n - 1, m*k) * A(m, k) with A(m, 0) = 1 for m >= 1 and n >= 0. (Unfortunately, the recurrence does not hold for m = 0.) - _Petros Hadjicostas_, Aug 12 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Aug 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009840 | OEIS-A260878

Number of set partitions of {1, 2, ..., 2*n} with sizes in {[n, n], [2n]}.

Conjecture: p divides a(p+5) - b(6) if p is a prime > b(6).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Aug 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009841 | OEIS-A260879

Coefficients in asymptotic expansion of sequence A229741.

Conjecture: a(k) ~ (k-1)! / (log(2))^k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Aug 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009842 | OEIS-A260883

Number of m-shape ordered set partitions, square array read by ascending antidiagonals, A(m, n) for m, n >= 0.

Conjecture: For n >= 0, let P be the set of all possible lists (a_1, ..., a_n) of nonnegative integers such that a_1*1 + a_2*2 + ... + a_n*n = n. Consider terms of the form multinomial(n*m, m*[1,..., 1, 2,..., 2,..., n,..., n]) * multinomial(a_1 + ... + a_n, [a_1,..., a_n]), where in the list [1,..., 1, 2,..., 2,..., n,..., n] the number 1 occurs a_1 times, 2 occurs a_2 times, ..., and n occurs a_n times. (Here a_n = 0 or 1.) Summing these terms over P we get A(m, n) provided m >= 1. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Aug 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009843 | OEIS-A260884

Number of set partitions of a 2n-set into even blocks which have even length minus the number of partitions into even blocks which have odd length.

Conjecture: a(n) = 2 * exp(1) * Sum_{k=1..oo} k^(2*n) * Bessell(k, -1) for n > 0. - _Velin Yanev_, Feb 20 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Sep 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009844 | OEIS-A260886

Least prime p such that 3 + 4*prime(p*n) = 5*prime(q*n) for some prime q.

Conjecture: Let a,b,c be pairwise relatively prime positive integers with a+b+c even and a not equal to b. Then, for any positive integer n, there are primes p and q such that a*prime(p*n) - b*prime(q*n) = c. This includes the conjectures in A260252 and A260882 as special cases.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009845 | OEIS-A260888

Least prime p such that 2 + 3*pi(p*n) = 4*pi(q*n) for some prime q, where pi(x) denotes the number of primes not exceeding x.

Conjecture: Let a and b be relatively prime positive integers, and let c be any integer. For any positive integer n, there are primes p and q such that a*pi(p*n) - b*pi(q*n) = c. In the case c = 0, this reduces to the conjecture in A260232.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009846 | OEIS-A260896

a(n) gives the number of integers m such that there exist k and h with 2n^2 < mk^2 < 2(n+1)^2 and 2n^2 < 2mh^2 < 2(n+1)^2.

Conjecture: a(n) > 0 for all n > 14.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Aug 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009847 | OEIS-A260911

Least positive integer k < prime(n) such that there are 0 < i < j < k for which i^2 + j^2 = k^2 and i,j,k are all quadratic residues modulo prime(n), or 0 if no such k exists.

Conjecture: (i) a(n) > 0 for all n > 25. In other words, for any prime p > 100, we have a^2 + b^2 = c^2 for some a,b,c in the set R(p) = {0<r<p: r is a quadratic residue mod p}. I have verified the conjecture for primes p < 1.5*10^7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009848 | OEIS-A260913

Coefficients in asymptotic expansion of sequence A059439.

Conjecture: a(k) ~ k * k! / (4 * (log(2))^(k+2)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Aug 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009849 | OEIS-A260929

Least positive integer k < prime(n) such that k + k' is a triangular number, where k' is the unique integer among 1,...,prime(n)-1 with k*k' == 1 (mod prime(n)), or 0 if no such k exists.

Conjecture: a(n) > 0 for all n > 7. In other words, for any prime p > 17 there are two integers k and k' among 1,...,p-1 with k*k' == 1 (mod p) such that k + k' is a triangular number. See also A260928 for a similar conjecture involving squares.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009850 | OEIS-A260947

Least positive integer k < prime(n) such that k = i + j for some 0 < i < j for which i,j,k and i*j*k are all primitive roots modulo prime(n), or 0 if no such k exists.

Conjecture: a(n) > 0 for all n > 6. In other words, for any prime p > 13, there are three distinct elements a,b,c of {1,...,p-1} with a+b = c such that a,b,c and a*b*c are all primitive roots modulo p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009851 | OEIS-A260952

Coefficients in asymptotic expansion of the sequences A109253 and A112225.

Conjecture: a(k) ~ -k! * 2^(k+1) / (9 * (log(3))^(k+1)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Aug 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009852 | OEIS-A260960

Least positive integer k < prime(n) such that there are 0 < i < j with i^2 + j^2 = k^2 for which (i*j)/2 is a primitive root modulo prime(n), or 0 if no such k exists.

Conjecture: a(n) > 0 for any n > 4. In other words, for any prime p > 7, there exists a right triangle whose three sides are among 1,...,p-1 and whose area is a primitive root modulo p. We also conjecture that for any prime p > 31, there exists a right triangle whose three sides are among 1,...,p-1, and whose perimeter and area are quadratic residues modulo p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009853 | OEIS-A260965

Smallest number r>=3 such that for k>=r the number k - 3 + prime(n) can be represented in the form x_1*...*x_k + x_1 + ... + x_k, where 1 <= x_1 <= ... <= x_k, or a(n)=0 if there is no such r.

Conjecture. The sequence contains only a finite number of zero terms. About the conjecture, for n <= 10000, 23 values a(n) are 0, of which 19 have n <= 100. The highest such n is 450. a(n) is at most five for n <= 10000 and mostly 3 (9747 times). - _David A. Corneth_, Aug 16 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Aug 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009854 | OEIS-A260970

Number of hereditarily transitive normal play partisan games born on or before day n.

The hereditarily transitive games born by day n form a distributive lattice whose Hasse diagram is planar. It is conjectured (known for n<=3) that the number of antichains in this lattice is 2^A000372(n)-2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christopher E. Thompson_, Aug 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009855 | OEIS-A261041

Number of partitions of subsets of {1,...,n}, where consecutive integers are required to be in different parts.

Conjecture: Also the number of set partitions of {1, ..., n + 1} where, if x and x + 2 belong to the same block, then so does x + 1. For example, the a(0) = 1 through a(3) = 10 set partitions are:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Aug 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009856 | OEIS-A261058

Column k=1 of A213221.

Conjecture: n*a(n) +(n-3)*a(n-1) +3*(-7*n+10)*a(n-2) +3*(n-1)*a(n-3) +40*(n-2)*a(n-4) +2*(-2*n+3)*a(n-5) +10*(-2*n+5)*a(n-6)=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Aug 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009857 | OEIS-A261061

Number of solutions to c(1)*prime(1)+...+c(2n)*prime(2n) = -1, where c(i) = +-1 for i > 1, c(1) = 1.

Conjecture: limit_{n->infinity} a(n)^(1/n) = 4. - _Vaclav Kotesovec_, Jun 05 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Aug 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009858 | OEIS-A261116

Pairs of integers (a,b) such a^2 + (a+1)^2 = b^2.

The conjectured recurrence formula and g.f. are proved by the formulas for A001652. - _M. F. Hasler_, Aug 11 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Aug 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009859 | OEIS-A261131

Number of ways to write n as the sum of 3 positive palindromes.

Conjecture: a(n) > 0 for n > 2, i.e., every number greater than 2 can be written as the sum of 3 positive palindromes. The conjecture is true (see links). - _Giorgos Kalogeropoulos_, May 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Resta_, Aug 10 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009860 | OEIS-A261133

a(n) = Max{k from {1..n} | T(n,k) = A049831(n)}, where T(n,k) is the triangle defined at A049828.

Conjecture: a(n)/n -> (sqrt(5)-1)/2, see Szocs-Mihai link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tiberiu Szocs-Mihai_, Aug 10 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009861 | OEIS-A261136

Primes p such that prime(p)-p+1 = prime(q) for some prime q.

The conjecture in A260753 implies that the current sequence has infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009862 | OEIS-A261176

Minimum value of (1/2)*Sum_{i=1..n} Sum_{j=1..n} Sum_{k=1..n} Sum_{l=1..n} gcd(b(i,j),b(k,l)) * ((i-k)^2+(j-l)^2) for an n X n matrix b filled with the integers 1 to n^2.

All terms beyond a(5) are conjectured based on numerical results. Terms up to a(17) have at least 5 independent verifications.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Aug 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009863 | OEIS-A261177

Maximum value of (1/2)*Sum_{i=1..n} Sum_{j=1..n} Sum_{k=1..n} Sum_{l=1..n} gcd(b(i,j),b(k,l)) * ((i-k)^2+(j-l)^2) for an n X n matrix b filled with the integers 1 to n^2.

Best results found in Al Zimmermann's Programming Contest "Delacorte Numbers". For more information see A261176. All terms beyond a(5) are conjectured based on numerical results. Terms up to a(11) have at least 5 independent verifications. Lower bounds for the next terms are a(14)>=5189492, a(15)>=8110781, a(16)>=12239616, a(17)>=18073562, a(18)>=26055061, a(19)>=36769303, a(20)>=51095165.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Aug 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009864 | OEIS-A261196

Expansion of sqrt(8*x + sqrt(1 + 64*x^2)).

Conjecture D-finite with recurrence: n*(n-1)*a(n) +(n-1)*(n-2)*a(n-1) +16*(2*n-3)*(2*n-5)*a(n-2) +16*(2*n-5)*(2*n-7)*a(n-3)=0. - _R. J. Mathar_, Jun 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Aug 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009865 | OEIS-A261246

Positive integers D such that the generalized Pell equation X^2 - D Y^2 = 2 is soluble.

Conjecture: The sequence consists of all numbers D not a square and even D = 2*d has odd d with prime factors of the form 1 or 7 (mod 8). Odd D has prime factors of the form 1 or 7 (mod 8) but there is an odd number of primes of the form 7 (mod 8). The following will prove that these conditions for D are necessary in order to have solutions. This conjecture is false. For the odd D case see the counterexamples in A263010, and for the even D in A264352. - _Wolfdieter Lang_, Nov 12 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Sep 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009866 | OEIS-A261249

Number of classes of proper solutions of the Pell equation x^2 - D(n) y^2 = +4 for D(n) = A079896(n), n >= 1.

Conjecture: For even D(n), i.e., D from 4*A000037, and a(n) = 0 one finds for r(n) = D(n)/4 coincidence with Conway's so-called rectangular numbers A007969. The first D values are 8, 20, 24, 40, 48, 52, 56, 68, 72, 80, ... This is equivalent to the conjecture that X^2 - r*y^2 = +1 has an even fundamental positive solution y = y0 precisely for the numbers A007969 (because x has to be even, x = 2*X, and whenever y0 is even all y solutions are even). See A261250 and A262024 for the y0 and x0 values, respectively.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Sep 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009867 | OEIS-A261280

Number of ways to start with set {1,2,...,n} and then repeat n times: partition each set into subsets.

Conjecture: a(n) ~ c * n^(2*n-5/6) / (2^(n-1) * exp(n)), where c = 7.7889...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Aug 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009868 | OEIS-A261282

Least positive integer k such that prime(k)*prime(k*n) = prime(p)+2 for some prime p.

Conjecture: a(n) exists for any n > 0. In general, any positive rational number r can be written as m/n, where m and n are positive integers such that prime(m)*prime(n) = prime(p)+2 for some prime p. The conjecture implies that there are infinitely many primes p such that prime(p)+2 is a product of two primes. Recall that a prime p is called a Chen prime if p+2 is a product of at most two primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009869 | OEIS-A261295

Least positive integer k such that both k and k*n belong to the set {m>0: prime(m) = prime(p)+2 for some prime p}.

Conjecture: (i) Any positive rational number r can be written as m/n with m and n in the set S = {k>0: prime(k) = prime(p)+2 for some prime p} = {p+1: p and prime(p)+2 are both prime}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009870 | OEIS-A261301

a(n+1) = abs(a(n) - gcd(a(n), n)), a(1) = 1.

It is conjectured that a(n) = 0 implies that n is prime, see A186253. (This is the sequence {u(n)} mentioned there.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Aug 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009871 | OEIS-A261302

a(n+1) = abs(a(n) - gcd(a(n), 2n+1)), a(1) = 1.

It is conjectured that for all n, a(n) = 0 implies that 2n+1 = a(n+1) is prime, cf. A186254. (This is the sequence {u(n)} mentioned there.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Aug 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009872 | OEIS-A261303

a(n+1) = abs(a(n) - gcd(a(n), 3n+2)), a(1) = 1.

It is conjectured that a(n) = 0 implies that 3n+2 = a(n+1) is prime, for all n > 2, cf. A186255. (This is the sequence {u(n)} mentioned there.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Aug 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009873 | OEIS-A261304

a(n+1) = abs(a(n) - gcd(a(n), 4n+3)), a(1) = 1.

It is conjectured that a(n) = 0 implies that 4n+3 = a(n+1) is prime, cf. A186256. (This is the sequence {u(n)} mentioned there.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Aug 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009874 | OEIS-A261305

a(n+1) = abs(a(n) - gcd(a(n), 5*n+4)), u(1) = 1.

It is conjectured that a(n) = 0 implies that 5n+4 = a(n+1) is prime for n > 2, cf. A186256. (This is the sequence {u(n)} mentioned there.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Aug 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009875 | OEIS-A261306

a(n+1) = abs((n) - gcd(a(n), 6*n+5)), a(1) = 1.

It is conjectured that for all n, a(n) = 0 implies that a(n+1) = 6n+5 is prime, cf. A186258. (This is the sequence {u(n)} mentioned there.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Aug 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009876 | OEIS-A261307

a(n+1) = abs(a(n) - gcd(a(n), 7*n+6)), a(1) = 1.

It is conjectured that for all n > 2, a(n) = 0 implies that 7n+6 = a(n+1) is prime, cf. A186259. (This is the sequence {u(n)} mentioned there.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Aug 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009877 | OEIS-A261308

a(n+1) = abs(a(n) - gcd(a(n), 8n+7)), a(1) = 1.

It is conjectured that for all n, a(n) = 0 implies that 8n+7 = a(n+1) is prime, cf. A186260. (This is the sequence {u(n)} mentioned there.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Aug 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009878 | OEIS-A261309

a(n+1) = abs(a(n) - gcd(a(n), 9n+8)), u(1) = 1.

It is conjectured that for all n > 2, u(n) = 0 implies that u(n+1) = 9n+8 is prime, cf. A186261. (This is the sequence {u(n)} mentioned there.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Aug 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009879 | OEIS-A261310

a(n+1) = abs(a(n) - gcd(a(n), 10n+9)), a(1) = 1.

It is conjectured that for all n, a(n) = 0 implies that a(n+1) = 10n+9 is prime, cf. A186263.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Aug 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009880 | OEIS-A261312

Numbers n such that the n-th Fibonacci number is not divisible by any prime Fibonacci number.

Conjecture: this sequence has natural density around 0.229. (The sequence has a natural density unless Fibonacci primes are much denser than expected.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Aug 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009881 | OEIS-A261314

Rocket Sequence 34: a(0)=34, a(n) = A073846(a(n-1)).

A given sequence c(n) can be one of two kinds. It can either be periodic with c(m) = c(0) for some m, or it can include infinitely many distinct values. R#(n) is finite for all n<34. However, this sequence has been checked up to a(86) = 1091595086717 without reaching 34. Instead it seems to be slowly climbing in value in both the negative and positive directions. Hence, its period is either extremely large or nonexistent (infinite). I conjecture that the latter is the case. Thus I dubbed the sequence "Rocket" because, as opposed to the "Hailstone" sequences, it never seems to "fall".

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chayim Lowen_, Aug 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009882 | OEIS-A261339

Least positive integer k such that both k and k*n belong to the set {m>0: m+1, m^2+1 and m^2+prime(m)^2 are all prime}.

Conjecture: a(n) exists for any n > 0. In general, any positive rational number r can be written as m/n with m and n in the set {k>0: k+1, k^2+1 and k^2+prime(k)^2 are all prime}. The conjecture implies that there are infinitely many primes p with (p-1)^2+1 and (p-1)^2+prime(p-1)^2 both prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009883 | OEIS-A261352

Primes p such that prime(p)+2 = prime(q)*prime(r) for distinct primes q and r.

Conjecture: The sequence has infinitely many terms. See also A261354 for a similar conjecture, and A261353 for a stronger conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009884 | OEIS-A261353

Least positive integer k such that prime(prime(k))*prime(prime(k*n)) = prime(p)+2 for some prime p.

Conjecture: a(n) exists for any n > 0. In general, any positive rational number r can be written as m/n, where m and n are positive integers such that prime(prime(m))*prime(prime(n)) = prime(p)+2 for some prime p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009885 | OEIS-A261354

Primes p such that prime(p)^2 - 2 = prime(q) for some prime q.

Conjecture: The sequence has infinitely many terms. In general, for any integers a,b,c with a>0 and gcd(a,b,c)=1, if b^2-4*a*c is not a square, a+b+c is odd, and gcd(b,a+c) is not divisible by 3, then there are infinitely many prime pairs {p,q} such that a*prime(p)^2+b*prime(p)+c = prime(q).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009886 | OEIS-A261362

Least positive integer k such that both k and k*n belong to the set {m>0: 2*prime(prime(m))+1 = prime(p) for some prime p}.

Conjecture: Let a,b,c be positive integers with gcd(a,b) = gcd(a,c) = gcd(b,c) = 1. If a+b+c is even and a is not equal to b, then any positive rational number r can be written as m/n with m and n in the set {k>0: a*prime(p) - b*prime(prime(k)) = c for some prime p}. This implies the conjecture in A261361.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009887 | OEIS-A261382

Least positive integer k such that (k-1)^2+(k*n)^2, k^2+(k*n-1)^2, (k+1)^2+(k*n)^2 and k^2+(k*n+1)^2 are all prime.

Conjecture: a(n) exists for any n > 0. In general, any positive rational number r can be written as m/n, where m and n are positive integers with (m-1)^2+n^2, m^2+(n-1)^2, (m+1)^2+n^2 and m^2+(n+1)^2 all prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009888 | OEIS-A261385

Least positive integer k such that (prime(prime(k))-1)*(prime(prime(k*n))-1) = prime(p)-1 for some prime p.

Conjecture: Let d be any nonzero integer. Then each positive rational number r can be written as m/n, where m and n are positive integers with (prime(prime(m))+d)*(prime(prime(n))+d) = prime(p)+d for some prime p. This conjecture implies that for any nonzero integer d the equation x*y = z with x,y,z in the set {prime(p)+d: p is prime} has infinitely many solutions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009889 | OEIS-A261395

Primes p such that (prime(p)-1)^2 = (prime(q)-1)*(prime(r)-1) for some pair of distinct primes q and r.

Conjecture: Let d be any nonzero integer. Then there are infinitely many prime triples (p,q,r) with p,q,r distinct such that (prime(p)+d)^2 = (prime(q)+d)*(prime(r)+d). In other words, the set {prime(p)+d: p is prime} contains infinitely many nontrivial three-term geometric progressions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009890 | OEIS-A261414

2^n+1 appears in A109732 at position a(n).

This assumes van der Poorten's conjecture that every odd number does appear in A109732.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009891 | OEIS-A261437

Least positive integer k such that k*n+1 = prime(p) and k^2*n+1 = prime(q) for some pair of primes p and q.

Conjecture: (i) If n > 0 and r are relatively prime integers, then there are infinitely many positive integers k such that k*n+r = prime(p) for some prime p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009892 | OEIS-A261440

Array of coefficients A(n,k) of the formal power series P(n,x) read by upwards antidiagonals, where P(n,x) = Sum_{k>=0} A(n,k)*x^k = 1+x*P(n,x)^(1*n)+x^2*P(n,x)^(2*n) for n >= 0.

Conjecture: The A(n,k), here n > 0, are the number of lattice paths, if Conjecture: The coefficients B(m,n,k) of the P(n,x)^m (see the formula below), m > 0 and n > 0, are the number of lattice paths, if A(n,k) = 1/(n*k+1)*Sum_{j=0..k} (-1)^j*binomial(n*k+1, j)*binomial(2*n*k+2-2*j, k-j) (conjectured). B(m,n,k) = m/(n*k + m)*(Sum_{j=0..k} (-1)^j*binomial(n*k+m, j)* binomial(2*n*k + 2*m - 2*j, k - j)), if m > 0, and n > 0 (conjectured). P(n,x) = exp(Sum_{k>=1} 1/(n*k)*(Sum{j=0..k} (-1)^j*binomial(n*k,j)* binomial(2*n*k-2*j,k-j))) for n > 0 (conjectured). - _Werner Schulte_, Sep 20 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Aug 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009893 | OEIS-A261462

Least positive integer k such that prime(prime(k)), prime(prime(k*n)), prime(p) and prime(q) form a 4-term arithmetic progression for some pair of primes p and q.

Conjecture: (i) Any positive rational number r can be written as m/n, where m and n are positive integers such that (prime(prime(m))+prime(prime(n)))/2 = prime(p) for some prime p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009894 | OEIS-A261491

a(n) = ceiling(2 + sqrt(8*n-4)).

Conjecture: a(n) = minimal number of stones needed to surround area n in the middle of a Go board (infinite if needed).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juhani Heino_, Aug 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009895 | OEIS-A261513

Least positive integer k with p(prime(k))+p(prime(k*n)) prime, where p(.) is the partition function given by A000041.

Conjecture: Any positive rational number r not equal to one can be written as m/n, where m and n are positive integers with p(prime(m)) + p(prime(n)) prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009896 | OEIS-A261515

Primes of the form p(q) + p(r) with q and r both prime, where p(.) is the partition function given by A000041.

The conjecture in A261513 implies that the current sequence has infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009897 | OEIS-A261524

Odd numbers n such that degree(gcd(1 + x^n, 1 + (1+x)^n)) > 1, where the polynomials are over GF(2).

Conjecture 1: Zs(d,2,1) is a term if and only if d is odd and d != 1, 15, 21. Verified for odd numbers up to 91. Conjecture 2: for a prime p >= 3, gcd(x^(Zs(d,p,1))-1, (x+1)^(Zs(d,p,1))-1) != 1 if and only if d is an odd prime power > 1 other than (d,p) = (7,3), (7,9), (13,3), (37,3), (43,3), (79,3). Note that the comment above shows that both conjectures are true for powers > 1 of a prime r >= 5. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Aug 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009898 | OEIS-A261528

Least positive integer k such that both k and k*n belong to the set {m>0: prime(m)+2 is prime with prime(prime(m)+2) = prime(prime(m))+6}.

Conjecture: Any positive rational number r can be written as m/n with m and n in the set {k>0: prime(k)+2 is prime with prime(prime(k)+2) = prime(prime(k))+6}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009899 | OEIS-A261533

Primes p such that p+2 is prime with prime(p+2)-prime(p)=6.

The conjecture in A261528 implies that the current sequence has infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009900 | OEIS-A261541

Least positive integer m such that both m and m*n belong to the set {k>0: prime(k)+2, prime(k)+6, prime(k)+8 are all prime}.

Conjecture: (i) Each positive rational number r can be written as m/n with m and n in the set {k>0: prime(k)+2, prime(k)+6 and prime(k)+8 are all prime}. Part (i) of the conjecture implies that there are infinitely many primes p with p+2, p+6 and p+8 all prime, while part (ii) implies that there are infinitely many primes p with p+4, p+6 and p+10 all prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009901 | OEIS-A261577

Numbers m such that (4^m + 11) / 3 is prime.

12702 is the first counterexample to Dhilan Lahoti's conjecture: 12702 = 3*4234. - _Bruno Berselli_, Feb 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Aug 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009902 | OEIS-A261580

Primes p such that A214028(p) is odd.

Conjecturely, since (k+2)/2 = 4 is a square, this sequence has density 7/24 in the primes; see the end of Section 1 of my link. (End) [Comment rewritten by _Jianing Song_, Jun 16 2024 and Jun 20 2024] The conjecture above is an analog of Hasse's result that the set {p prime : ord(2,p) is odd} has density 7/24 in the primes, where ord(a,m) is the multiplicative order of a modulo m; see A014663. - _Jianing Song_, Jun 26 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Aug 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009903 | OEIS-A261593

Odd numbers n such that the sum of the binary digits of n and n^2 both equal 12.

12 is the least constant for which the associated sequence is known to be infinite. 1 through 8 make finite sequences, the sequences with 9 and 10 are conjectured finite, and the sequence with 11 is unknown. (See Hare-Laishram-Stoll Theorem 1.3 and 1.4 plus section 5.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Aug 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009904 | OEIS-A261594

G.f.: sqrt(Sum_{n>=0} x^n * Sum_{k=0..n} C(n,k) * 2^(k*(k-1))).

a(k) = 1 (mod 3) iff k = 9*A005836(n) + [0,1,3,4] for n>=0, with a(k) = 0 (mod 3) elsewhere, where A005836 lists numbers n whose base 3 representation contains no 2 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009905 | OEIS-A261598

Product of sexagesimal digits of Fibonacci numbers in base-60 representation.

Conjecture: a(n) = 0 for n > 3329 (empirically checked up to 36000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Sep 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009906 | OEIS-A261602

Triangular array of A(n,k) for n>=1 and 0<=k<=n^2 equal the number of permutations of the set {1,2,...,n}^2 such that first coordinates of first k elements are nondecreasing and second coordinates of the remaining n^2-k elements are nondecreasing.

It is conjectured that A(n,k)>A(n,k-1) for k<=floor(n^2/2) (see Mathoverflow link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Aug 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009907 | OEIS-A261625

Number of primes p <= n such that (p-1)*n+1 is prime.

Conjecture: a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009908 | OEIS-A261628

Least prime p such that n-(p*n'-1) and n+(p*n'-1) are both prime where n' = (3+(-1)^n)/2, or 0 if no such prime p exists.

Conjecture: 0 < a(n) < sqrt(2*n)*log(5*n) for all n > 6. Conjecture verified for n < 1.2 * 10^12. Also, the 5 inside the log function can probably be replaced by 4.26. - _Jud McCranie_, Aug 26 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009909 | OEIS-A261641

Number of practical numbers q such that n+(n mod 2)-q and n-(n mod 2)+q are both practical numbers.

Conjecture: a(n) > 0 except for n = 2. Also, for any integer n > 3, there is a practical number q such that n-(n mod 2)-q and n+(n mod 2)+q are both practical numbers. This is an analog of the author's conjecture in A261627, and it is stronger than Margenstern's conjecture proved by Melfi in 1996.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009910 | OEIS-A261653

Number of primes p < n such that n-p-1 and n+p+1 are both prime or both practical.

Conjecture: a(n) > 0 for all n > 6. Also, for any integer n > 2, there is a prime p < n such that n-(p-1) and n+(p-1) are both prime or both practical.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009911 | OEIS-A261671

If n even, a(n) = 6n+3, otherwise a(n) = n.

It is conjectured that starting with any positive odd number d and iterating f, we always eventually reach either 1, 9, or 33 (see Comments in A135730).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009912 | OEIS-A261678

Even numbers that are not the sum of two binary palindromes.

A subsequence of the numbers that are not the sum of three terms from A006995. The two sequences are equal if every odd number is the sum of three terms from A006995 (which is equivalent to the conjecture in A261680). - _Chai Wah Wu_, Sep 14 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009913 | OEIS-A261681

a(n) = 2^n + binomial(n, floor(n/2)) - 1.

Conjecture: -(n+1)*(n-4)*a(n) +(3*n^2-9*n-8)*a(n-1) +2*(n^2-9*n+16)*a(n-2) +4*(-3*n^2+18*n-25)*a(n-3) +8*(n-3)^2*a(n-4)=0. - _R. J. Mathar_, Jan 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009914 | OEIS-A261682

a(n) = 2^n+(1+(n mod 2)/2)*C(n+1,floor((n+(n mod 2))/2))-1.

Conjecture: -(n+2)*(15*n^2-29*n-56)*a(n) +9*(5*n^3-3*n^2-24*n-20)*a(n-1) +2*(15*n^3-14*n^2-219*n+158)*a(n-2) +36*(-5*n^3+8*n^2+31*n-38)*a(n-3) +8*(n-2)*(15*n^2+n-70)*a(n-4)=0. - _R. J. Mathar_, Jan 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009915 | OEIS-A261699

Triangle read by rows: T(n,k), n >= 1, k >= 1, in which column k lists positive terms interleaved with k-1 zeros, starting in row k(k+1)/2. If k is odd the positive terms of column k are k's, otherwise if k is even the positive terms of column k are the odd numbers greater than k in increasing order.

Conjecture: the positive terms in row n are the odd divisors of n. Proof of the conjecture: let n = 2^m*s*t with s and t odd. The property stated in A237048 verifies the conjecture with odd divisor k <= A003056(n) of n in position k and odd divisor t > A003056(n) in position 2^(m+1)*s. Therefore reading in row n the nonzero odd positions from left to right and then the nonzero even positions from right to left gives all odd divisors of n in increasing order. - _Hartmut F. W. Hoft_, Oct 25 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Sep 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009916 | OEIS-A261715

Putative inverse of conjectured permutation in A261702.

Putative inverse of conjectured permutation in A261702.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Tek_, Aug 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009917 | OEIS-A261725

Lexicographically earliest sequence of distinct terms such that the absolute difference of two successive terms is a power of 10, and can be computed without carry.

Conjectured to be a permutation of the nonnegative integers. See A261729 for putative inverse.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Tek_, Aug 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009918 | OEIS-A261728

a(1)=1; a(2*n) = 3*n; for odd n>1, a(n) is the smallest number not already present which is entailed by the rules (i) k present => 3*k+1 present; (ii) 2*k present => k present.

Proof. We use induction. The induction hypothesis (IH) is that the lemma is true for every even c<=2*k and odd c<=2*k+1. On the other hand, the statement 'the sequence is a permutation of the positive integers' conjecturally is equivalent to the (3*n+1)-conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Aug 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009919 | OEIS-A261729

Putative inverse of conjectured permutation in A261725.

Putative inverse of conjectured permutation in A261725.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Tek_, Aug 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009920 | OEIS-A261781

Number T(n,k) of compositions of n where each part i is marked with a word of length i over a k-ary alphabet whose letters appear in alphabetical order and all k letters occur at least once in the composition; triangle T(n,k), n >= 0, 0 <= k <= n, read by rows.

Conjecture: For k > 0 the recurrence order for column k is equal to k*(k+1)/2. Conjecture: d(k+1) - d(k) tends to 1/log(2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Aug 31 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009921 | OEIS-A261862

Terms in A261524 that are not multiples of earlier terms.

In A261524 it is conjectured that degree(gcd(1 + x^(Zs(d,2,1)), 1 + (1+x)^(Zs(d,2,1))) > 0 for every odd number d != 1, 15, 21, where Zs(d,2,1) is the d-th Zsigmondy number with parameters (2,1) (A064078). Since Zsigmondy numbers with different indices are coprime, if this conjecture is true, then there exists a term of this sequence k with ord(2,k) = d, and k must be a divisor of Zs(d,2,1) for every odd number d != 1, 15, 21. Here ord(a,k) is the multiplicative order of 2 modulo k. In A261524 we show that this conjecture is true for powers > 1 of a prime r >= 5, so there are infinitely many terms in this sequence. One may conjecture that, if k is a term with ord(2,k) = d for even d, then k is a divisor of Zs(d,2,1)*Zs(d/2,2,1). This fails for (d,k) = (20,11275), (40,16962275), (44,165965585), ... Conjecture: a term with ord(2,k) = d for even d exists if and only if d != 12 or 2*p, where p is any Mersonne exponent. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Sep 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009922 | OEIS-A261876

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with (5*x^2+7*y^2+9*z^2)*y*z a square, where x,y,z,w are nonnegative integers with z > 0.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 4^k*m (k = 0,1,2,... and m = 1, 7, 23, 647, 863). Adamczewski includes an abstract of disproof regarding the above conjecture in linked article. - _Michael De Vlieger_, Aug 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009923 | OEIS-A261878

Number of distinct fractional parts of the sums 1/j+...+1/k with 1 <= j <= k <= n, where the fractional part of x is given by x - floor(x).

Conjecture: (i) If 1/j+..+1/k and 1/s+...+1/t have the same fractional part with 0 < min{2,k} <= j <= k, 0 < min{2,t} <= s <= t and j <= s, but the ordered pairs (j,k) and (s,t) are different, then we have 1/j+...+1/k = 1+1/s+...+1/t; moreover, either (j,k) = (2,6) and (s,t) = (4,5), or (j,k) = (2,4) and (s,t) = (12,12), or (j,k) = (2,11) and (s,t) =(5,12), or (j,k) = (3,20) and (s,t) = (7,19). Clearly, part (i) of the conjecture implies that a(n) = n*(n-1)/2 - 3 for all n > 20. See also A261993 for a similar conjecture involving primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009924 | OEIS-A261918

Numbers which in base 5 are neither palindromes nor the sum of two palindromes.

The terms less than 20000, and conjecturally all terms, are the sum of three base 5 palindromes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 13 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009925 | OEIS-A261931

a(0)=1, a(1)=2, for n>=1, a(2*n) = 6*n - 2; a(2*n+1) = min(((a(2*n)-1)/3)^, (2*a(2*n))^, (2*a(2*n-1))^), where, instead of t, we write t^, if t has not appeared earlier in the sequence and is neither of the form 3*k nor of the form 6*m-2.

This sequence is connected with Collatz's (3*n+1)-conjecture. For example, if n=29, then, by the formulas, 29 = a(31) => 88 = a(30) => 11 = a(13) => 34 = a(12) => 17 = a(19) => 52 = a(18) => 13 = a(15) => 40 = a(14) => 5 = a(7) => 16 = a(6) => 1 = a(0).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009926 | OEIS-A261950

Start with a single equilateral triangle for n=0; for the odd n-th generation add a triangle at each expandable vertex of the triangles of the (n-1)-th generation (this is the "side to vertex" version); for the even n-th generation use the "vertex to vertex" version; a(n) is the number of triangles added in the n-th generation.

Conjectures from _Colin Barker_, Sep 10 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Sep 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009927 | OEIS-A261951

Start with a single equilateral triangle for n=0; for the odd n-th generation add a triangle at each expandable vertex of the triangles of the (n-1)-th generation (this is the "vertex to vertex" version); for the even n-th generation use the "vertex to side" version; a(n) is the number of triangles added in the n-th generation.

Conjectures from _Colin Barker_, Sep 10 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Sep 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009928 | OEIS-A261952

Start with a single equilateral triangle for n=0; for the odd n-th generation add a triangle at each expandable vertex of the triangles of the (n-1)-th generation (this is the "vertex to vertex" version); for the even n-th generation use the "side to side" version; a(n) is the number of triangles added in the n-th generation.

Conjectures from _Colin Barker_, Sep 10 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Sep 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009929 | OEIS-A261953

Start with a single equilateral triangle for n=0; for the odd n-th generation add a triangle at each expandable side of the triangles of the (n-1)-th generation (this is the "side to side" version); for the even n-th generation use the "vertex to vertex" version; a(n) is the number of triangles added in the n-th generation.

For finite sets of random points in the real plane with exactly n nearest neighbors, a(n) for n >= 2 is a lower bound for the maximal number of points. Conjecturally, a(n) equals this number. We conjecture that a(n) is optimal. This claim is true if the following assumptions hold:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Sep 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009930 | OEIS-A261954

Start with a single equilateral triangle for n=0; for the odd n-th generation add a triangle at each expandable side of the triangles of the (n-1)-th generation (this is the "side to side" version); for the even n-th generation use the "side to vertex" version; a(n) is the number of triangles added in the n-th generation.

Conjectures from _Colin Barker_, Sep 10 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Sep 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009931 | OEIS-A261955

Start with a single equilateral triangle for n=0; for the odd n-th generation add a triangle at each expandable side of the triangles of the (n-1)-th generation (this is the "side to side" version); for the even n-th generation use the "vertex to side" version; a(n) is the number of triangles added in the n-th generation.

Conjectures from _Colin Barker_, Sep 10 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Sep 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009932 | OEIS-A261956

Start with a single equilateral triangle for n=0; for the odd n-th generation add a triangle at each expandable vertex of the triangles of the (n-1)-th generation (this is the "side to vertex" version); for the even n-th generation use the "side to side" version; a(n) is the number of triangles added in the n-th generation.

Conjectures from _Colin Barker_, Sep 10 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Sep 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009933 | OEIS-A261957

Start with a single equilateral triangle for n=0; for the odd n-th generation add a triangle at each expandable side of the triangles of the (n-1)-th generation (this is the "vertex to side" version); for the even n-th generation use the "side to side" version; a(n) is the number of triangles added in the n-th generation.

Conjectures from _Colin Barker_, Sep 10 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Sep 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009934 | OEIS-A261958

Start with a single square for n=0; for the odd n-th generation add a square at each expandable vertex of the squares of the (n-1)-th generation (this is the "vertex to vertex" version); for the even n-th generation use the "side to vertex" version; a(n) is the number of squares added in the n-th generation.

Conjectures from _Colin Barker_, Sep 10 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Sep 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009935 | OEIS-A261986

Numbers k such that A003313(k) = A003313(4*k).

Terms are counterexamples to the conjecture that no shortest addition chains exist such that A003313(m)=A003313(m*2^k) for k>1, with a(1)=30958077 provided in 2008 by _Neill M. Clift_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Sep 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009936 | OEIS-A261993

Number of distinct fractional parts of the numbers 1/(prime(j)-1)+...+1/(prime(k)-1) with 1 <= j <= k <= n, where the fractional part of x is given by x - floor(x).

Conjecture: Let m be any positive integer. Clearly, part (i) of the conjecture implies that a(n) = n*(n-1)/2 - 2 for all n > 18. See also A261878 for a similar conjecture not involving primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009937 | OEIS-A262005

L.g.f.: log(Sum_{n>=0} x^n/n! * Product_{k=1..n} (k^5 + 1)).

a(n) == 2 (mod 60) for n>=1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009938 | OEIS-A262009

Sum_{d|n} 2^(d^2) * n^2/d^2.

Conjecture: A037227(a(n)) = 2*A037227(n) + 1. Conjecture: a(n) = 2^A037227(n) * d for some odd d, where A037227(n) = 2*m + 1 such that n = 2^m * k for some odd k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009939 | OEIS-A262030

Partition array in Abramowitz-Stegun order: Schur functions evaluated at 1.

Conjecture: The sum of the squares of row numbers give A054688: 1, 10, 165, 3876, ... = binomial(n^2+n-1, n). -_Wouter Meeussen_, Sep 25 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009940 | OEIS-A262082

Array of coefficients A(n,k) of the formal power series P(n,x) read by upwards antidiagonals, where P(n,x) = Sum_{k>=0} A(n,k)*x^k = 1 + x*P(n,x)^(1*n) + x^2*P(n,x)^(2*n) + x^3*P(n,x)^(3*n) for n >= 0.

Conjecture 1: The A(n,k), here n > 0, are the number of lattice paths, if Conjecture 2: The coefficients B(m,n,k) of the P(n,x)^m (see the formula below), m > 0 and n > 0, are the number of lattice paths, if A(n,k) = 1/(n*k+1) * Sum_{j=0..k} (-2)^j*binomial(n*k+1,j)* binomial(3*n*k+3-2*j,k-j) for n >= 0, and k >= 0. (conjectured) B(m,n,k) = m/(n*k+m) * Sum_{j=0..k} (-2)^j*binomial(n*k+m,j)* binomial(3*n*k+3*m-2*j,k-j) for m > 0, n > 0, and k >= 0. (conjectured) P(n,x) = exp(Sum_{k>=1} 1/(n*k)*(Sum_{j=0..k} (-2)^j*binomial(n*k,j)* binomial(3*n*k-2*j,k-j))) for n > 0 (conjectured); (see for n=1: A036765, for n=2: A186241, and for n=3: A200731).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Sep 10 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009941 | OEIS-A262114

Irregular triangle read by rows: row b (b >= 2) gives periodic part of digits of the base-b expansion of 1/5.

Conjectures from _Colin Barker_, Oct 09 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Sep 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009942 | OEIS-A262115

Irregular triangle read by rows: row b (b >= 2) gives periodic part of digits of the base-b expansion of 1/7.

Conjectures from _Colin Barker_, Oct 09 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Sep 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009943 | OEIS-A262136

Number of distinct fractional parts of the numbers Sum_{i=j..k} (-1)^i/i with 1 <= j <= k <= n, where the fractional part of x is given by x - floor(x).

Conjecture: (i) Suppose that Sum_{i=j..k} (-1)^i/i and Sum_{r=s..t} (-1)^r/r with 0 < min{2,k} <= j <= k, 0 < min{2,t} <= s <= t and j <= s have the same fractional part, but the ordered pairs (j,k) and (s,t) are different. Then Sum_{i=j..k} (-1)^i/i = Sum_{r=s..t} (-1)^r/r. Moreover, if j is odd, then j > 1, k = j*(j+1) and (s,t) = (j+2,j*(j+1)-1); if j is even, then either (k = j+1 and s = t = j*(j+1)), or (k = j*(j+1)-1 and (s,t) = (j+2,j*(j+1))). This is an analog of the conjecture in A261878. Part (i) of the conjecture implies that a(n) = n*(n-1)/2 + 2 - floor((sqrt(4n+1)-1)/2) - floor((sqrt(4n+1)-1)/4) for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009944 | OEIS-A262143

Square array read by antidiagonals upwards: the n-th row o.g.f. is exp(Sum_{i >= 1} c(n,i)*x^i/i) for n >= 1, where c(n,k) is Shanks' array of generalized Euler and class numbers.

We conjecture that the entries of the present array are all integers. More generally, we conjecture that for r = 0,1,2,... and for each n >= 1, the expansion of exp(Sum_{i >= 1} c(n,i + r)*x^i/i) has integer coefficients. The case n = 1 was conjectured by Hanna in A255895.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 13 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009945 | OEIS-A262144

Square array read by antidiagonals upwards: the n-th row o.g.f. is exp(Sum_{i >= 1} d(n,i+1)*x^i/i) for n >= 1, where d(n,k) is Shanks's array of generalized Euler and class numbers.

We conjecture that the entries of the present array are all integers. More generally, we conjecture that for r = 1, 2, ... and for each n >= 1, the expansion of exp(Sum_{i >= 1} d(n,i + r)*x^i/i) has integer coefficients. This is the case r = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009946 | OEIS-A262145

Expansion of o.g.f. exp(Sum_{n >= 1} A000182(n+1)*x^n/n), where A000182 is the sequence of tangent numbers.

It appears that the sequence has integer entries. Calculation suggests the following conjecture: the expansion of exp(Sum_{n >= 1} A000182(n + m)*x^n/n) has integer coefficients for m = 1, 2, 3, ... This is the case m = 1. Cf. A255881 and A255895.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 13 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009947 | OEIS-A262147

Numerators of domain values whose images via the flowsnake Q-function follow a spiral trajectory (see comments).

Conjectures from _Colin Barker_, Jan 24 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bradley Klee_, Sep 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009948 | OEIS-A262148

Denominators of domain values whose images via the flowsnake Q-function follow a spiral trajectory.

Conjectures from _Colin Barker_, Jan 24 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bradley Klee_, Sep 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009949 | OEIS-A262149

Rocket sequence 50: a(0)=50, a(n)=A073846(a(n-1)).

This sequence has been checked up to a(98) = 1078406742163 without reaching 50. It seems to be slowly climbing in value in both the negative and positive directions. Hence, its period is either extremely large or, as I conjecture, infinite. Thus I dubbed the sequence "Rocket" because, as opposed the "Hailstone" sequences, it never seems to "fall".

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chayim Lowen_, Sep 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009950 | OEIS-A262177

Decimal expansion of Q_5 = zeta(5) / (Sum_{k>=1} (-1)^(k+1) / (k^5 * binomial(2k, k))), a conjecturally irrational constant defined by an Apéry-like formula.

Decimal expansion of Q_5 = zeta(5) / (Sum_{k>=1} (-1)^(k+1) / (k^5 * binomial(2k, k))), a conjecturally irrational constant defined by an Apéry-like formula.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Sep 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009951 | OEIS-A262179

Signed denominators of the reduced form of the coefficients of degree 2n terms of the Maclaurin series of (t/sinh(t))^x in t.

Conjecture: this is also the integer sequence A202367 up to sign. These numbers show up in the formula for eigenvectors of Adams operations on the K-theory of unitary groups.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chi-Kwong Fok_, Sep 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009952 | OEIS-A262228

Deficiency sequence: a(0) = 1, a(n) is the smallest prime p > a(n-1) such that the product of a(i), 1 <= i < n, is deficient (belongs to A005100).

Conjecture: log(a(n)) ~ e^(an+b) where a and b are approximately 0.6 and -1.6 respectively.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chayim Lowen_, Sep 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009953 | OEIS-A262242

Triangular numbers representable as 2^x + 2^y.

Conjectures from _Colin Barker_, Sep 16 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Sep 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009954 | OEIS-A262258

a(n) = the number of hills (arch length of 1 with no covering arches) for semi-meander solutions with n arches and floor((n+2)/2) arch group returns to the x axis.

Conjectures from _Colin Barker_, May 06 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger Ford_, Sep 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009955 | OEIS-A262259

Numbers k such that the symmetric representation of sigma(k) has only two parts and they meet at the center of the Dyck path.

Conjecture: Numbers k with two 2-dense sublists of divisors and 2*A033676(k) = A033677(k) - 1. - _Omar E. Pol_, Feb 08 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Sep 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009956 | OEIS-A262267

Number of (n+2) X (1+2) 0..1 arrays with each row and column divisible by 5, read as a binary number with top and left being the most significant bits.

Conjectures from _Colin Barker_, Feb 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009957 | OEIS-A262311

Number of ordered ways to write n = x^2 + y^2 + phi(z^2) (0 <= x <= y and z > 0) with y or z prime, where phi(.) is Euler's totient function given by A000010.

Conjecture: a(n) > 0 for all n > 1. See also A262781 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009958 | OEIS-A262326

Number of (n+1) X (2+1) 0..1 arrays with each row and column divisible by 3, read as a binary number with top and left being the most significant bits.

Conjectures from _Colin Barker_, Mar 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009959 | OEIS-A262327

Number of (n+1) X (3+1) 0..1 arrays with each row and column divisible by 3, read as a binary number with top and left being the most significant bits.

Conjectures from _Colin Barker_, Dec 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009960 | OEIS-A262332

T(n,k) = Number of (n+1) X (k+1) 0..1 arrays with each row and column divisible by 3, read as a binary number with top and left being the most significant bits.

It appears that T(m,k) = c1^(mk) * c2^(m+k) * e^(o(m+k)), where c1 = 2.0000(0), c2 = 0.3333(3) based on numerical analysis. We conjecture that c1 = 2, c2 = 1/3; In other words, for sufficiently large m, k, the effect of restrictions on T(m,k) is almost "independent" of each other in the horizontal and vertical directions. - _Liang Kai_, Oct 31 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009961 | OEIS-A262350

a(1) = 2. For n>1, let s denote the binary string of a(n-1) with the leftmost 1 and following consecutive 0's removed. Then a(n) is the smallest prime not yet present whose binary representation begins with s.

This sequence is infinite. The number of primes that are not in this sequence is conjectured to be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Sep 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009962 | OEIS-A262357

Number of ordered ways to write n as w^2 + x^2 + y^2 + z^2 with w^2*x^2 + 5*x^2*y^2 + 80*y^2*z^2 + 20*z^2*w^2 a square, where w is a positive integer and x,y,z are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 4^k*m (k = 0,1,2,... and m = 1, 3, 11, 43, 547, 763, 1739, 6783). See also A268507, A269400, A271510, A271513, A271518, A271665, A271714, A271721, A271724, A271775, A271778 and A271824 for other conjectures refining Lagrange's four-square theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009963 | OEIS-A262375

Smallest prime factor of 4*n! - 1.

The larger terms (conjectured: records) are those whose indices are listed in A099350.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Sep 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009964 | OEIS-A262395

Difference between the numbers of trees on n vertices with an even number and an odd number of leaves.

The sequence could be prepended with a(0)=1 and a(1)=-1. However, it is conjectured that for all n>=2, we have a(n)>=0 (cf. MathOverflow link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Sep 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009965 | OEIS-A262408

Positive integers m such that pi(m^2) = pi(j^2) + pi(k^2) for no 0 < j <= k < m.

Conjecture: (i) There are infinitely many positive integers m such that pi(m^2) = pi(x^2) + pi(y^2) for some 0 < x <= y < m. Also, the current sequence has infinitely many terms. It is interesting to compare this conjecture with Fermat's Last Theorem. See also A262409 for the equation pi(x^3) + pi(y^3) = pi(z^3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009966 | OEIS-A262409

Positive integers m such that pi(m^3) = pi(j^3) + pi(k^3) for some 0 < j <= k < m.

Conjecture: The Diophantine equation pi(x^3) + pi(y^3) = pi(z^3) with 0 < x <= y < z has infinitely many solutions. See also the conjecture in A262408 involving the n-th powers with n = 2,4,5,....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009967 | OEIS-A262415

Number of (n+1) X (3+1) 0..1 arrays with each row divisible by 3 and column not divisible by 3, read as a binary number with top and left being the most significant bits.

Conjectures from _Colin Barker_, Dec 31 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Sep 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009968 | OEIS-A262439

Number of primes not exceeding 1+n*(n+1)/2.

Conjecture: (i) The sequence is strictly increasing, and also a(n)^(1/n) > a(n+1)^(1/(n+1)) for all n = 3,4,.... See also A262446 related to part (ii) of this conjecture. Concerning part (ii) of the conjecture, Neill Clift verified in 2024 that for all 1 < n <= 2^24 = 16777216 we have a(n) = a(k) + a(m) for some 0 < k <= m < n. - _Zhi-Wei Sun_, Jan 29 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009969 | OEIS-A262443

Positive integers m such that pi(m^2) = pi(j^2)*pi(k^2) for some 0 < j < k < m, where pi(x) denotes the number of primes not exceeding x.

Conjecture: (i) The sequence has infinitely many terms. Also, there are infinitely many positive integers m such that pi(m^2) = pi(j^2)*pi(k^2) for no 0 < j <= k < m.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009970 | OEIS-A262447

Primes p such that pi(p^2) = pi(q^2) + pi(r^2) for some distinct primes q and r.

Conjecture: The sequence has infinitely many terms. See also A262408 and A262443 for related conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009971 | OEIS-A262462

Positive integers k with pi(k^3) a square, where pi(x) denotes the number of primes not exceeding x.

Conjecture: (i) The Diophantine equation pi(x^2) = y^2 with x > 0 and y > 0 has infinitely many solutions. The conjecture seems reasonable in view of the heuristic arguments. Part (ii) of the conjecture implies that the only terms of the current sequence are 1, 2, 3, 14 and 1122.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009972 | OEIS-A262526

Any number greater than a(n) can conjecturally be represented in more ways by sums of three base-10 palindromes than a(n).

Any number greater than a(n) can conjecturally be represented in more ways by sums of three base-10 palindromes than a(n). The corresponding representation counts are provided in A262527. Positions of latest occurrence of increasing minima of representation counts in A261132. The sequence provides numerical evidence for the validity of the conjecture that every number is the sum of three palindromes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Sep 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009973 | OEIS-A262527

Conjectured minimum number of ways to represent a number >= A262526(i) by sums of three base-10 palindromes.

Conjectured minimum number of ways to represent a number >= A262526(i) by sums of three base-10 palindromes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Sep 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009974 | OEIS-A262535

Triangle read by rows T(n,k) in which column k lists the partial sums of the k-th column of triangle A261699.

Conjecture: the sum of row n gives A078471(n), the sum of all odd divisors of all positive integers <= n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Sep 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009975 | OEIS-A262536

Positive integers z such that pi(x^3+y^3) = pi(z^3) for some 0 < x <= y < z, where pi(m) denotes the number of primes not exceeding m.

Conjecture: (i) For any integers n > 3 and x,y,z > 0 with {x,y} not equal to {1,z}, we have |x^n+y^n-z^n| >= 2^n-2, unless n = 5, {x,y} = {13,16} and z = 17. We have verified part (i) of the conjecture for n = 4..10 and 0 < x,y,z <= 1700.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009976 | OEIS-A262597

Lüroth expansion of the fractional part of e.

It is conjectured that e is normal with respect to the Lüroth expansion.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthias Paulsen_, Sep 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009977 | OEIS-A262669

Consider the Farey sequence of order n, F_n, and that the average distance between any two adjacent pairs in F_n is 1/A002088(n). Then a(n) is the number of adjacent pairs whose difference is less than the average.

Conjecture: this is a monotonic sequence. For n = 0, 1, 3, 4, 8, 12, 17, 23, 41 & 59, a(n) = a(n+1). Conjecture: Twice the number of pairs less than the average (2*A262669) plus the number of pairs which equal the average (A262670) never exceed the number of pairs which are greater than the average for n greater than 245.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Sep 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009978 | OEIS-A262698

Positive integers m such that pi(k^3) + pi(m^3) is a cube for some k = 1,...,m, where pi(x) denotes the number of primes not exceeding x.

Conjecture: (i) There are infinitely many distinct primes p,q,r such that pi(p^2) + pi(q^2) = r^2. a(26) > 10^5, if it exists. Conjecture (ii) above is false since these further solutions exist: (1339, 7918, 2682), (3360, 8474, 2922), (8443, 13264, 4764), (15590, 16664, 6696), (15883, 27415, 9431), (9719, 39514, 12689), (22265, 48606, 15933), (38606, 51145, 18297). - _Giovanni Resta_, Jun 14 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009979 | OEIS-A262700

Primes p such that pi(p^2)*pi(q^2) is a square for some prime q < p, where pi(x) denotes the number of primes not exceeding x.

Conjecture: (i) The sequence has infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009980 | OEIS-A262707

Positive integers m such that pi(k^2)*pi(m^2) is a square for some 1 < k < m, where pi(x) denotes the number of primes not exceeding x.

Conjecture: The sequence has infinitely many terms. See also A262700 for a related conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009981 | OEIS-A262708

a(n) = p-(p/5) where p = prime(n) and (p/5) is a Legendre symbol.

The sequence lists Fibonacci indices q that are conjectured to produce Fibonacci numbers divisible by p^2, where p is a Fibonacci-Wieferich prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shane Findley_, Sep 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009982 | OEIS-A262722

Positive integers m such that pi(k^3+m^3) is a cube for some k = 1..m, where pi(x) denotes the number of primes not exceeding x.

Conjecture: (i) There are infinitely many distinct primes p,q,r such that pi(p^2+q^2) = r^2. Part (ii) of the conjecture implies that the current sequence only has 12 terms as shown here. Conjecture (ii) is false as there are more terms beyond 1710. It is likely the sequence has an infinite number of terms. (x,y,z) for 1710 < y <= 7564: (1429, 4322, 1514), (1974, 4523, 1604), (3361, 4877, 1840), (3992, 4942, 1949), (3253, 5147, 1902), (971, 5407, 1859), (935, 7564, 2563). -_Chai Wah Wu_, Apr 12 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009983 | OEIS-A262725

The unique function f with f(1)=1 and f(jD!+k)=(-1)^j f(k) for all D, j=1..D, and k=1..D!.

sup_n |Sum_{j=1..n} f(jd)| is finite (but not bounded) for all d, thus giving a counterexample to a strong form of the Erdős discrepancy conjecture (see Remark 1.14 on p. 7 of Tao paper).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Terence Tao_, Sep 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009984 | OEIS-A262730

Primes p such that p^2 = pi(x^3) + pi(y^3) for some positive integers x and y, where pi(m) denotes the number of primes not exceeding m.

Conjecture: (i) The sequence has infinitely many terms. See also A262731 for a related conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009985 | OEIS-A262731

Primes p in the form pi(q^2)+pi(r^2) with q and r both prime, where pi(x) denotes the number of primes not exceeding x.

Conjecture: The sequence has infinitely many terms. In general, for each n = 2,3,4,... there are infinitely many primes p in the form pi(q^n)+pi(r^n) with q and r both prime. Compare this conjecture with the well-known result that there are infinitely many primes p in the form x^2+y^2 with x and y positive integers (such a prime p is either 2 or congruent to 1 modulo 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009986 | OEIS-A262735

Expansion of x*(1-x-x^2)/((1-x)*(1-x-2*x^2-x^3)).

Conjecture: a(n+1) = Sum_{j=0..n/2} A027907(n-j,2*j), n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Sep 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009987 | OEIS-A262746

Number of ordered ways to write n as x^2 + y^2 + pi(z^2) with 0 <= x <= y and z > 0 such that 2*x*y + 3 is prime, where pi(m) denotes the number of primes not exceeding m.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009988 | OEIS-A262747

Number of ordered ways to write n as x^2 + y^2 + phi(z^2) with 0 <= x <= y, z > 0, 2 | x*y*z, and phi(k^2) < n for all 0 < k < z, where phi(.) is Euler's totient function given by A000010.

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 3, 4, 5, 8, 13, 16, 23, 32, 35, 39, 68, 75, 96, 215, 219, 243, 363, 471, 723, 759, 923, 1443, 1551, 1839, 2739, 2883. See also A262311 for a related conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009989 | OEIS-A262750

Least positive integer z such that n - phi(z^2) = x^2 + y^2 for some integers x and y with x*y*z even and phi(k^2) < n for all 0 < k < z, or 0 if no such z exists, where phi(.) is Euler's totient function given by A000010.

Conjecture: a(n) <= sqrt(n) except for n = 3, 8, 13, 32. The conjecture in A262747 implies that a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009990 | OEIS-A262765

a(n) = the number of ways that at least two distinct primes <= prime(n) sum to a prime.

Conjectures: The plot does, in fact, suggest that a(n)/a(n-1) "generally increases" (i.e., generally a(z*n)/a(z*n-1) > a(n)/a(n-1), when z is sufficiently large). In other words, the peaks and the troughs tend to increase, with peaks tending to be higher than next trough. This behavior is consistent with the observation about "general increase" in conjecture iii, as n increases to infinity. However, the plot does not show that generally a(n)/a(n-1) > a(n+1)/a(n), or that there can't be a maximum value of a(n)/a(n-1) after which all other values decrease (two completely different issues). Conjecture ii is clearly false since a(n)/(n-1) is slightly > 2 for a few terms 11 <= n <= 400 (n = {333..340}), therefore weakening the observation in conjecture iii that the limiting ratio must be < 2. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bob Selcoe_, Sep 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009991 | OEIS-A262766

Number of positive integers z such that n - phi(z^2) = x^2 + y^2 for some integers x and y with x*y*z even and phi(k^2) < n for all 0 < k < z.

The conjecture in A262747 implies that a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009992 | OEIS-A262779

Binary representation of the n-th iteration of the "Rule 175" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 20 2016 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009993 | OEIS-A262781

Number of ordered ways to write n as x^2 + phi(y^2) + phi(z^2) (x >= 0 and 0 < y <= z) with y or z prime, where phi(.) is Euler's totient function given by A000010.

Conjecture: (i) a(n) > 0 for all n > 6, and a(n) = 1 only for n = 3, 5, 9, 10, 17, 20, 24, 25, 31, 36, 45, 73, 80, 101, 136, 145, 388, 649. See also A262311 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009994 | OEIS-A262785

Number of ordered ways to write n as x^2 + y^2 + p*(p+d)/2, where 0 <= x <= y, d is 1 or -1, and p is prime.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 2, 9, 34, 45, 49, 72, 97, 241, 337, 538.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009995 | OEIS-A262813

Number of ordered ways to write n as x^3 + y^2 + z*(z+1)/2 with x >= 0, y >=0 and z > 0.

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 9, 21, 35, 98, 152, 306. Conjecture verified up to 10^11. - _Mauro Fiorentini_, Jul 18 2023 In contrast with the conjecture, in 2015 the author refined a result of Euler by proving that any positive integer can be written as the sum of two squares and a positive triangular number. See also A262815, A262816 and A262941 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009996 | OEIS-A262814

Numbers k dividing every cyclic permutation of k^k.

Conjecture: 10^n-1 is a term of the sequence for all n > 0. - _Chai Wah Wu_, Nov 03 2015 Wu's conjecture is true. See a proof in the links section. - _FUNG Cheok Yin_, Jun 04 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Oct 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009997 | OEIS-A262815

Number of ordered ways to write n as x^3 + y*(y+1)/2 + z*(3*z+1)/2, where x, y and z are nonnegative integers.

Conjecture: a(n) > 0 for all n >= 0, and a(n) = 1 only for n = 0, 5, 12, 19, 20, 75, 83, 97, 117. Conjecture verified up to 10^11. - _Mauro Fiorentini_, Jul 20 2023 See also A262813 and A262816 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009998 | OEIS-A262816

Number of ordered ways to write n as x^3 + y^2 + z*(3*z-1)/2, where x and y are nonnegative integers, and z is a nonzero integer.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 4, 216. Conjectures (i) and (ii) verified for n up to 10^9. - _Mauro Fiorentini_, Jul 21 2023 See also A262813 and A262815 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-009999 | OEIS-A262824

Number of ordered ways to write n as w^2 + x^3 + 2*y^3 + 3*z^3, where w, x, y and z are nonnegative integers.

Conjecture: (i) For any m = 3, 4, 5, 6 and n >= 0, there are nonnegative integers w, x, y, z such that n = w^2 + x^3 + 2*y^3 + m*z^3. Conjectures (i) and (ii) verified up to 10^11. - _Mauro Fiorentini_, Jul 22 2023 See also A262827 and A262857 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010000 | OEIS-A262826

a(n) = Sum_{d|n} -(-1)^d * 2^(n^2/d) * d.

Conjecture: a(n) = 2^A007814(n) * 2^n * d for some odd d, where A007814(n) equals the exponent of highest power of 2 dividing n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010001 | OEIS-A262827

Number of ordered ways to write n as w^2 + x^3 + y^4 + 2*z^4, where w, x, y and z are nonnegative integers.

Conjecture: a(n) > 0 for all n >= 0. Also, a(n) = 1 only for the following 41 values of n: 0, 14, 15, 21, 22, 23, 62, 71, 78, 87, 136, 216, 405, 437, 448, 477, 535, 583, 591, 623, 671, 696, 885, 950, 1046, 1135, 1206, 1208, 1248, 1317, 2288, 2383, 2543, 3167, 3717, 3974, 6847, 7918, 8328, 9096, 21935. Conjecture verified up to 10^11. - _Mauro Fiorentini_, Jul 07 2023 We also conjecture that if f(w,x,y,z) is one of the 8 polynomials 2w^2+x^3+4y^3+z^4, w^2+x^3+2y^3+c*z^3 (c = 3,4,5,6) and w^2+x^3+2y^3+d*z^4 (d = 1,3,6) then each n = 0,1,2,... can be written as f(w,x,y,z) with w,x,y,z nonnegative integers. - _Zhi-Wei Sun_, Dec 30 2017 Conjecture verified up to 10^11 for all 8 polynomials. - _Mauro Fiorentini_, Jul 07 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010002 | OEIS-A262857

Number of ordered ways to write n as w^3 + 2*x^3 + y^2 + 2*z^2, where w, x, y and z are nonnegative integers.

Conjecture: We have {a*w^3+b*x^3+c*y^2+d*z^2: w,x,y,z = 0,1,2,...} = {0,1,2,...} if (a,b,c,d) is among the following 63 quadruples: Conjecture verified up to 10^11 for all quadruples. - _Mauro Fiorentini_, Jul 18 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010003 | OEIS-A262880

Number of ordered ways to write n as w*(w+1)/2 + x^3 + y^3 + 2*z^3 with w > 0, 0 <= x <= y and z >= 0.

Conjecture: (i) Any positive integer can be written as w*(w+1)/2 + x^3 + b*y^3 + c*z^3 with w > 0 and x,y,z >= 0, provided that (b,c) is among the following ordered pairs: (1,2),(1,3),(1,4),(1,6),(2,2),(2,3),(2,4),(2,5),(2,6),(2,7),(2,20),(2,21),(2,34),(3,3),(3,4),(3,5),(3,6),(4,10). See also A262813, A262824 and A262857 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010004 | OEIS-A262887

Number of ordered ways to write n as x^3 + y^2 + pi(z^2) (x >= 0, y >= 0 and z > 0) with z-1 or z+1 prime, where pi(m) denotes the number of primes not exceeding m.

Conjectures: Compare these conjectures with the conjectures in A262746.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010005 | OEIS-A262896

If n is in A262892, a(n) = A259934(n), otherwise the largest term in A045765 from which A259934(n) can be reached by iterating A049820, without visiting any other (larger) term of A259934.

Note that without (so far undetected) regularity in A262509, there is no a priori upper bound for the value of a(n), and for some n this might not even be finite, if it happens that contrary to its conjectured nature, A259934 is not the unique infinite component, but just the lexicographically earliest instance of multiple infinite branches of the tree. In that case we might consider this sequence to be well-defined only up to the least such node branching to multiple infinite components, or alternatively, we might mark the nonfinite values at those points with -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010006 | OEIS-A262922

a(1)=1; for n>1, a(n) = a(n-1) + n + 2 if a(n-1) and n are coprime, otherwise a(n) = a(n-1)/gcd(a(n-1),n).

Quasi-periodic sequences exist only for R=0,1,2 or 3 in a(n) = a(n-1) + n + R. For R=0,1,2 all starting values give a quasi-periodic sequence. The respective loop is (1,x) for R=0, (1,x,2x,2) for R=1, (1,x,2x,x) or (x,2x) for R=2. For R=3 only some starting values converge to a 6-loop (4x+2,2x+1,3x+6,x+2,2x+9,3x+17). Conjecture: For R>=4 the recurrence is not quasi-periodic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Oct 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010007 | OEIS-A262941

Number of ordered pairs (x,y) with x >= 0 and y > 0 such that n - x^4 - y*(y+1)/2 is an even square or twice a square.

Conjecture: a(n) > 0 for all n > 0. In other words, any positive integer n can be written as x^4 + 2^k*y^2 + z*(z+1)/2, where k is 1 or 2, and x,y,z are integers with z > 0. See also A262944, A262945, A262954, A262955, A262956 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010008 | OEIS-A262942

Sequence of positive integers where each is chosen to be as small as possible subject to the conditions that no three terms a(j), a(j+k), a(j+2*k) (for any j and k) form an arithmetic progression (in any order) and that no term repeats.

Conjectured permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Barrentine_, Oct 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010009 | OEIS-A262943

Inverse of conjectured permutation A262942.

Inverse of conjectured permutation A262942.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Barrentine_, Oct 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010010 | OEIS-A262944

Number of ordered pairs (x,y) with x >= 0 and y > 0 such that n - x^4 - y*(y+1)/2 is a square or a pentagonal number.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 9, 63, 69, 489, 714, 1089. See also A262941 and A262945 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010011 | OEIS-A262945

Number of ordered pairs (x,y) with x >= 0 and y >= 0 such that n - x^4 - 2*y^2 is a triangular number or a pentagonal number.

Conjecture: a(n) > 0 for every n = 0,1,2,..., and a(n) = 1 only for the following 55 values of n: 0, 26, 27, 32, 41, 42, 50, 65, 80, 97, 112, 122, 130, 160, 196, 227, 239, 272, 322, 371, 612, 647, 736, 967, 995, 1007, 1106, 1127, 1205, 1237, 1240, 1262, 1637, 1657, 1757, 2912, 2987, 3062, 3107, 3524, 3647, 3902, 5387, 5587, 5657, 6047, 6107, 11462, 13427, 14717, 15002, 17132, 20462, 30082, 35750. See also A262941, A262944, A262954 and A262955 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010012 | OEIS-A262954

Number of ordered pairs (x,y) with x >= 0 and y > 0 such that n - x^4 - y*(y+1)/2 has the form z^2 or 8*z^2.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 9, 13, 89, 449, 524, 1049, 2164, 14969, 51334. See also A262941, A262944, A262945, A262954, A262955 and A262956 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010013 | OEIS-A262955

Number of ordered pairs (x,y) with x >= 0 and y > 0 such that n - x^4 - y*(y+1)/2 is a pentagonal number (A000326) or twice a pentagonal number.

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 10, 35, 75, 134, 415, 515, 1465, 2365, 3515, 4140.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010014 | OEIS-A262956

Number of ordered pairs (x,y) with x >= 0 and y > 0 such that n - x^4 - y*(y+1)/2 is a square or a square minus 1.

Conjecture: a(n) > 0 for all n > 0. In other words, for any positive integer n, either n or n + 1 can be written as the sum of a fourth power, a square and a positive triangular number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010015 | OEIS-A262957

Numerators of the n-th iteration of the alternating continued fraction formed from the positive integers, starting with (1 - ...).

Conjecture: b(n) and c(n) are coprime for all n, so that a(n) = b(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mohamed Sabba_, Nov 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010016 | OEIS-A262959

Positive integers that cannot be written as the sum of a fourth power, a square and a positive triangular number.

Conjecture: (i) Each term is congruent to one of 3, 4, 5, 9 modulo 10. Part (ii) of this conjecture is stronger than the conjecture in A262956. Note that a(139) - a(138) = 18089 - 18084 = 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010017 | OEIS-A262961

Crandall numbers: (2/Pi)^4 Integral_{t>=0} ([Pi I_0(t)]^2 - [K_0(t)]^2) I_0(t) [K_0(t)]^5 (2t)^(2n-1) dt.

Anton Mellit and David Broadhurst define the sequence to be the "round" of the integral, with the conjecture that this rounding is exact. No one seems to know how to prove that any of the integrals gives a rational number, let alone an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew R. Reiter_, Oct 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010018 | OEIS-A262965

Least number k such that k mod s = prime(n) where s is the sum of the distinct primes dividing k.

Conjecture: a(n) exists for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Oct 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010019 | OEIS-A262976

Number of ordered ways to write n as 2^x + y^2 + pi(z^2) with x >= 0, y >= 0 and z > 0, where pi(m) denotes the number of primes not exceeding m.

Conjecture: (i) a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010020 | OEIS-A262979

Number of ordered ways to write n as x^4 + phi(y^2) + z*(3*z-1)/2 with x >= 0 and y > 0, where phi(.) is Euler's totient function given by A000010.

Conjecture: (i) a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010021 | OEIS-A262980

Number of ordered ways to write n as p + 2^k + pi(2^m), where p is prime, and k and m are nonnegative integers, and pi(x) denotes the number of primes not exceeding x.

Conjecture: a(n) > 0 for all n > 2. In contrast with this conjecture, in 1971 R. Crocker proved that there are infinitely many positive odd numbers not of the form p + 2^k + 2^m, where p is prime, and k and m are positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010022 | OEIS-A262981

Numbers k such that the least positive integer having exactly k divisors is divisible by k.

Some conjectures for a(n):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Letsko_, Oct 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010023 | OEIS-A262982

Number of ordered ways to write n as x^4 + phi(y^2) + z*(z+1)/2 with x >= 0, y > 0 and z > 0, where phi(.) is Euler's totient function given by A000010.

Conjecture: a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 6, 20, 26, 40, 82, 89, 105, 305, 416, 470, 725, 6135, 25430, 90285. Compare this with the conjectures in A262311, A262785 and A262813.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010024 | OEIS-A262985

Number of ordered ways to write n as 2^x + phi(y^2) + z*(z+1)/2 with x, y and z positive integers, where phi(.) is Euler's totient function given by A000010.

Conjecture: a(n) > 0 for all n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010025 | OEIS-A262995

Number of ordered pairs (k,m) with k > 0 and m > 0 such that n = pi(k*(k+1)/2) + pi(1+m*(m+1)/2), where pi(x) denotes the number of primes not exceeding x.

Conjecture: a(n) > 0 for all n > 0. This has been verified for n up to 10^5. See also A262999 for a similar conjecture. In A262403 and A262439, the author conjectured that the sequences pi(n*(n+1)/2) (n = 1,2,3,...) and pi(1+n*(n+1)/2) (n = 1,2,3,...) are both strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010026 | OEIS-A262999

Number of ordered pairs (k, m) with k > 0 and m > 0 such that n = pi(k*(k+1)/2) + pi(m^2), where pi(x) denotes the number of primes not exceeding x.

Conjecture: a(n) > 0 for all n > 1, and a(n) = 1 only for n = 3, 5, 7, 35, 37, 217, 7439, 10381. See also A262995, A263001 and A263020 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010027 | OEIS-A263001

Number of ordered pairs (k, m) with k > 0 and m > 0 such that n = pi(k*(k+1)) + pi(m*(m+1)/2), where pi(x) denotes the number of primes not exceeding x.

Conjecture: a(n) > 0 for all n > 2, and a(n) = 1 only for n = 1, 4, 6. See also A262995, A262999 and A263020 for similar conjectures. Adamczewski disproves the above conjecture in linked article, using the case a(51156) = 1. - _Michael De Vlieger_, Aug 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010028 | OEIS-A263010

Exceptional odd numbers D that do not admit a solution to the Pell equation X^2 - D Y^2 = +2.

Necessary conditions for nonsquare odd D were shown there to be D == 7 (mod 8), without prime factors 3 or 5 (mod 8) in the composite case. Thus only prime factors +1 (mod 8) and -1 (mod 8) can appear, and the number of the latter is odd. It has been conjectured that all such numbers D appear in A261246, but this conjecture is false as the present sequence shows. For counterexamples to the conjecture in A261246 for even D see A264352.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Nov 10 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010029 | OEIS-A263020

Number of ordered pairs (k, m) with k > 0 and m > 0 such that n = pi(k*(k+1)/2) + pi(m*(3*m-1)/2), where pi(x) denotes the number of primes not exceeding x.

Conjecture: a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010030 | OEIS-A263100

Number of ordered pairs (k, m) with k > 0 and m > 0 such that n = pi(k^2) + pi(m^2/2), where pi(x) denotes the number of primes not exceeding x.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 3, 28. See also A262995, A262999, A263001 and A263020 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010031 | OEIS-A263104

a(3k), a(3k+1), a(3k+2) are the numbers of edges between each vertex-pair in a multigraph with 3 vertices and k edges (with at least one edge between each vertex-pair) which has the minimum number of distinct cycles, if that arrangement of edges is unique.

Viewing the sequence in rows of three (see links) suggests that after a(64) its behavior becomes regular, with each value of a(n) appearing 9 times in 3 contiguous columns of 3 (see conjectured formula). a(n) = floor((n +4*(n%3)-1)/9) for n >= 65 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon R. Donnelly_, Oct 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010032 | OEIS-A263107

Number of ordered pairs (k, m) with k > 0 and m > 0 such that n = pi(k^2/2) + pi(3*m^2/2), where pi(x) denotes the number of primes not exceeding x.

Conjecture: a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010033 | OEIS-A263135

The maximum number of penny-to-penny connections when n pennies are placed on the vertices of a hexagonal tiling.

Conjecture: a(2*n) - A047932(n) = A216256(n) for n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Oct 10 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010034 | OEIS-A263171

Smallest prime starting a sequence of 4 consecutive odd primes such that the center of the symmetrical gaps is 2n.

Conjecture: a(n) exists for all n>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Oct 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010035 | OEIS-A263190

G.f. satisfies: [x^(n-1)] A(x)^(n^2) = (n!)^2 for n>=1.

CONJECTURES. Conjectures hold to at least a(1000). - _Sean A. Irvine_, Oct 21 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010036 | OEIS-A263206

Number of primes p with n^2 < prime(p) < (n+2)^2.

Conjecture: a(n) > 0 for all n > 0. In other words, for each n = 1,2,3,... the interval (n^2, (n+2)^2) contains a prime with prime subscript.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010037 | OEIS-A263207

Number of integer solutions for Product_{k=1..n}(c(k) + 1) = 2 * Product_{k=1..n}(c(k) - 1) with 1 < c(k) <= c(k+1).

For any n>=1 the solution given by c(k)=A204321(k) for k=1..n-1 and c(n)=A204321(n)-1 always exists. Conjecture: there is no solution with a greater c(n). - _Martin Janecke_, Dec 01 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Martin Janecke_, Oct 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010038 | OEIS-A263244

Binary representation of the n-th iteration of the "Rule 155" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 20 2016 and Apr 17 2019: (Start) Conjecture: a(n) = floor(91*100^((n-1)/2)*10^n/9) for odd n; a(n) = floor(901*100^(n/2 - 1)*10^n/9) for even n. - _Karl V. Keller, Jr._, May 01 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010039 | OEIS-A263245

Decimal representation of the n-th iteration of the "Rule 155" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 20 2016 and Apr 17 2019: (Start) Conjecture: a(n) = 3*4^((n-1)/2)*2^n - 1 for odd n; a(n) = 5*4^(n/2 - 1)*2^n - 1 for even n > 0. - _Karl V. Keller, Jr._, May 01 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010040 | OEIS-A263267

Breadth-first traversal of the tree defined by the edge-relation A049820(child) = parent.

It is conjectured that the terms of A259934 trace the only infinite path in this tree.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010041 | OEIS-A263317

Least prime p > n such that the numbers sigma(k^2)/k^2 (k = 1,...,n) are pairwise incongruent modulo p, where sigma(m) is the sum of the divisors of m.

Conjecture: a(n) exists for any n > 0, and a(n) < n^2 for all n > 2. We guess that for each k = 2,3,... all the numbers sigma(n^k)/n^k = Sum_{d|n^k} 1/d (n = 1,2,3,...) are pairwise distinct. See also A001157 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010042 | OEIS-A263319

a(n) = pi(n^2)*phi(n)/2, where pi(x) denotes the number of primes not exceeding x, and phi(.) is Euler's totient function given by A000010.

Conjecture: (i) All the terms of this sequence are pairwise distinct. We have checked this conjecture via Mathematica. For example, we have verified that a(n) (n = 1..4*10^5) are indeed pairwise distinct. See also A263325 for a similar

conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010043 | OEIS-A263321

Least positive integer m such that the numbers phi(k)*pi(k^2) (k = 1..n) are pairwise incongruent modulo m.

Part (i) of the conjecture in A263319 implies that a(n) exists for any n > 0. Conjecture: a(n) <= n^2 for all n > 0, and the only even term is a(7) = 16.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010044 | OEIS-A263325

a(n) = sigma(n)*pi(n^2), where sigma(n) is the sum of all (positive) divisors of n, and pi(x) is the number of primes not exceeding x.

Conjecture: (i) All the terms of this sequence are pairwise distinct. See also A263319 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010045 | OEIS-A263326

Denominator of the rational number Sum_{d|n}1/(d+1).

Conjecture: For any positive integers k and s, all the numbers Sum_{d|n}1/(d+k)^s (n = 1,2,3,...) have pairwise distinct fractional parts, and none of them is an integer. See also A001157 for a similar conjecture involving Sum_{d|n}1/d^s. Adamczewski disproves the above conjecture in linked article, using the case k = 31, s = 1. -_Michael De Vlieger_, Aug 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010046 | OEIS-A263399

Number of ordered pairs (k, m) with 0 < m < n and |k| < prime(m)^2 such that (k+prime(m+1)^2)/(k+prime(m)^2) = (k+prime(n+1)^2)/(k+prime(n)^2).

Conjecture: Let k, m and n be integers with 0 < m < n and -2*prime(m) < k <= 2*prime(m). If (k+prime(m+1)^2)/(k+prime(m)^2) = (k+prime(n+1)^2)/(k+prime(n)^2), then we must have k = -9, m = 5 and n = 11. The conjecture essentially implies that for each k = -3..10 all the ratios (prime(n+1)^2+k)/(prime(n)^2+k) (n = 1,2,...) are pairwise distinct. We have verified that for any k = -1, 1 the ratios (prime(n+1)^2+k)/(prime(n)^2+k) (n = 1..110000) are indeed pairwise distinct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010047 | OEIS-A263427

Denominator of (prime(prime(n))+2)/(prime(n)+2).

Conjecture: For each k = 2,3,6,7,8,9 all the rational numbers (prime(p)+k)/(p+k) with p prime are pairwise distinct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010048 | OEIS-A263429

Smallest prime p such that binomial(2*p-1, p-1) == 1 (mod p^n), or 0 if no such p exists.

Conjecture: a(n) = 0 for n > 4 (McIntosh, 1995, p. 387).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Oct 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010049 | OEIS-A263450

Smallest integer k>0 such that there is at least one zero in the decimal representation of prime(n)^k.

Conjecture: there are an infinite number of ones in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Oct 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010050 | OEIS-A263466

Least k such that prime(n) is the smallest prime p for which k^2 + p^2 is also prime, or 0 if none.

The data support the conjecture in A263977 that if k > 0 is even, then k^2 + p^2 is prime for some prime p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_ and _Robert G. Wilson v_, Nov 02 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010051 | OEIS-A263563

A self-describing sequence: when the sequence is read as a string of decimal digits, a(n) can be read from position n (ignoring leading zeros). This sequence is the lexicographically earliest sequence of distinct terms with this property.

This sequence is conjectured to be a permutation of natural numbers, with putative inverse A263564.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Tek_, Oct 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010052 | OEIS-A263573

Intersection of A024365 and A129912.

The two sequences involve areas of primitive Pythagorean triples and primorial products. Intersections are only considered once (no repeats). Conjecture: the sequence is infinite. Conjecture: The next two entries are a(12) = 200560490130, a(13) = 401120980260.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bill McEachen_, Oct 21 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010053 | OEIS-A263613

Palindromic numbers in base 4 that are cubes.

Conjectures from _Chai Wah Wu_, Dec 25 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010054 | OEIS-A263634

Irregular triangle read by rows: row n gives coefficients of n-th logarithmic polynomial L_n(x_1, x_2, ...) with monomials sorted into standard order.

Conjecture: row polynomials are R(n,1) for n > 0 where R(n,k) = R(n-1,k+1) - Sum_{j=1..n-1} binomial(n-2,j-1)*R(j,k)*R(n-j,1) for n > 1, k > 0 with R(1,k) = x_k for k > 0. - _Mikhail Kurkov_, Mar 30 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010055 | OEIS-A263660

Number of length n arrays of permutations of 0..n-1 with each element moved by -2 to 2 places and with no two consecutive increases or two consecutive decreases.

Conjectures from _Colin Barker_, Jan 02 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010056 | OEIS-A263687

b(n) in (sqrt(2))_n = b(n) + c(n)*sqrt(2), where (x)_n is the Pochhammer symbol, b(n) and c(n) are integers.

Conjecture: a(n) = (-1)^n * Sum_{k=0..n/2} Stirling1(n,2*k)*2^k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Reshetnikov_, Oct 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010057 | OEIS-A263695

Even numbers such that the sum of the even divisors and the sum of the odd divisors are a square or a cube.

Conjecture [False!]: All squares belonging to a pair are associated with a unique cube. Conversely, all cubes are associated with a unique square. The conjecture above is false. Consider for example the triples of numbers {69576, 938184, 7505472} or {958528, 952520, 12382760}. For the first one the (even, odd) sum of divisors pairs are (56^3, 112^2), (1568^2, 56^3), and (4704^2, 56^3). - _Giovanni Resta_, May 28 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, May 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010058 | OEIS-A263722

Integers k > 0 such that k^2 + p^2 is composite for all primes p.

Conjecture: All terms are odd. An equivalent conjecture in A263977 is that if k > 0 is even, then k^2 + p^2 is prime for some prime p. We have checked this for all k <= 12*10^7. An odd number k is a term if and only if k^2 + 2^2 is composite, since k^2 + p^2 is even and > 2 (hence composite) for all odd primes p. Thus if the Conjecture is true, then the sequence is simply odd k such that k^2 + 4 is composite. The sequence of odd k, such that k^2 + 4 is prime, is A007591.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_ and _Robert G. Wilson v_, Oct 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010059 | OEIS-A263789

Triangle read by rows: T(n,k) (n>=0, 0<=k<=floor(n/2)) is the number of permutations of n and k valleys (considered cyclically).

Conjecture: column k > 0 is asymptotic to n * 2^(n-2*k) * k^(n-1). - _Vaclav Kotesovec_, Oct 26 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian Stump_, Oct 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010060 | OEIS-A263794

Number of (n+1) X (3+1) 0..1 arrays with each row and column divisible by 3, read as a binary number with top and left being the most significant bits, and rows and columns lexicographically nonincreasing.

Conjectures from _Colin Barker_, Dec 14 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010061 | OEIS-A263805

Binary representation of the n-th iteration of the "Rule 157" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 20 2016 and Apr 17 2019: (Start) Conjecture: a(n) = (10*10^n*(10^n + 1)/11 - 1)/9 for odd n; a(n) = (10^n*(10^n + 10)/11 - 1)/9 for even n > 0. - _Karl V. Keller, Jr._, May 05 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010062 | OEIS-A263806

Decimal representation of the n-th iteration of the "Rule 157" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 20 2016 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010063 | OEIS-A263810

Numbers k such that k = tau(k) * phi(k-2) + 1.

Conjecture: primes from this sequence are in A254576.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Oct 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010064 | OEIS-A263822

Conjectured number of terms in A263644 less than 10^n.

Conjectured number of terms in A263644 less than 10^n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Oct 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010065 | OEIS-A263843

Reversion of g.f. for A162395 (squares with signs).

Let A(x) = 1 + 4*x + 23*x^2 + ... denote the o.g.f. of this sequence taken with an offset of 0. The sequence defined by b(n) := [x^n] A(x)^n for n >= 0 begins [1, 4, 62, 1084, 19982, 379504, 7347410, 144168392, 2856907662, 57044977168, 1145905776312, 23131265652092, ...]. We conjecture that the supercongruences b(n*p^k) == b(n*p^(k-1)) (mod p^(3*k)) hold for prime p >= 3 and all positive integers n and k. More generally, for a positive integer r and integer s, the sequence {b(r,s;n) : n >= 0} defined by b(r,s;n) := [x^(r*n)] A(x)^(s*n) is conjectured to satisfy the same supercongruences. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010066 | OEIS-A263869

Number of (n+1) X (4+1) 0..1 arrays with each row and column divisible by 3, read as a binary number with top and left being the most significant bits, and rows and columns lexicographically nondecreasing.

Conjectures from _Colin Barker_, Jan 03 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010067 | OEIS-A263907

Number of (2n+2) X (2+2) 0..1 arrays with each row and column modulo 3 equal to 1, read as a binary number with top and left being the most significant bits.

Conjectures from _Colin Barker_, Jan 03 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010068 | OEIS-A263908

Number of (2n+1) X (3+2) 0..1 arrays with each row and column modulo 3 equal to 1, read as a binary number with top and left being the most significant bits.

Conjectures from _Colin Barker_, Jan 03 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010069 | OEIS-A263920

A positive integer n is in this sequence iff arctan(n)^2 can be represented as Sum_{0<k<n} c(k)*arctan(k)^2 with rational c(k). The terms are in increasing order.

It is an open question if the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Reshetnikov_, Oct 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010070 | OEIS-A263931

a(n) = binomial(2*n, n) / Product(p prime | n < p <= 2*n).

By the Erdős squarefree conjecture, proved in 1996, no a(n) with n >= 5 is squarefree. - _Robert FERREOL_, Sep 06 2022 Conjecture: n=5 is the only n for which a(n)^2 > binomial(2*n,n). - _Richard Peterson_, Dec 18 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Oct 31 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010071 | OEIS-A263977

Integers k > 0 such that k^2 + p^2 is prime for some prime p.

Conjecture: Every even number k is a member. (This is equivalent to the Conjecture in A263722.) We have checked this for all k <= 12*10^7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_ and _Robert G. Wilson v_, Oct 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010072 | OEIS-A263992

Number of ordered ways to write n as x^2 + 2*y^2 + phi(z^2) (x >= 0, y >= 0 and z > 0) such that y or z has the form p-1 with p prime, where phi(.) is Euler's totient function.

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 6, 7, 22, 3447. This is similar to the conjecture in A262311, and we have verified it for n up to 10^6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 31 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010073 | OEIS-A263998

Number of ordered ways to write n as x^2 + 2*y^2 + p*(p+d)/2, where x and y are nonnegative integers, d is 1 or -1, and p is prime.

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 2, 8. This is similar to the conjecture in A262785.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 31 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010074 | OEIS-A264010

Number of ways to write n as x^2 + y*(y+1) + z*(z+1)/2, where x, y and z are nonnegative integers such that y or y+1 is prime, and z or z+1 is prime.

Conjectures: (i) a(n) > 0 for all n > 2, and a(n) = 1 only for n = 3, 4, 5, 6, 10, 11, 15, 20, 29, 1125. See also A264025 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 31 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010075 | OEIS-A264025

Number of ways to write n as x^2 + y*(2*y+1) + z*(z+1)/2 where x, y and z are nonnegative integers with z or z+1 prime.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 2, 3, 8, 9, 23, 30, 44, 48, 198, 219, 1344. See also A262785, A263998 and A264010 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010076 | OEIS-A264038

Convolution of Lucas and Jacobsthal numbers.

Referring to the simplicity of definition (II), the authors formulate the following open question: "Since the convolution takes such a simple form, we ask whether it is possible to obtain a purely combinatorial proof of this result." I would suggest another open question: Are there convolutions of other linear homogeneous recurrences with constant coefficients which are equivalent to very simple forms?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Russell Jay Hendel_, Nov 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010077 | OEIS-A264041

a(n) is the maximum number of diagonals that can be placed in an n X n grid made up of 1 X 1 unit squares when diagonals are placed in the unit squares in such a way that no two diagonals may cross or intersect at an endpoint.

Conjecture: partial sums of A260307. - _Sean A. Irvine_, Jul 15 2022 a(n) = a(n-1) + a(n-2) - a(n-3) + a(n-6) - a(n-7) - a(n-8) + a(n-9) for n>9 (conjectured). - _Colin Barker_, Jan 31 2018 a(n) = n*(n+1)/2 for n even, floor(n*(n/2+2/3)+1/6) for n odd (conjecture). - _Bill McEachen_, Aug 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gabriella Pinter_, Stephen Wasielewski, Peter Boyland, Ivan Roth, G. Christopher Hruska, Jeb Willenbring, Oct 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010078 | OEIS-A264083

Number of orthogonal 3 X 3 matrices over the ring Z/nZ.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles Repizo_, Nov 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010079 | OEIS-A264100

Sum of the lengths of the arithmetic progressions in {1,2,3,...,n}, including trivial arithmetic progressions of lengths 1 and 2.

Conjecture: the second differences give A007503(n+1), the sum of the divisors (A000203) plus the number of divisors (A000005) of n+1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gionata Neri_, Nov 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010080 | OEIS-A264104

Numbers k with the property that the symmetric representation of sigma(k) has four parts, each of width one and two regions meet at the center of the Dyck path.

Conjecture: Numbers k with four 2-dense sublists of divisors and four odd divisors such that 2*A033676(k) = A033677(k) - 1. - _Omar E. Pol_, Feb 12 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Nov 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010081 | OEIS-A264120

Values of k such that A001163(k) is positive.

If the conjecture is true, this entry will be removed as a duplicate of A042963. - _R. J. Mathar_, Nov 06 2015 Conjecture: for n > 1, a(n) = ((-1)^n + 4*n - 7)/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Reshetnikov_, Nov 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010082 | OEIS-A264123

Number of (n+1) X (3+1) arrays of permutations of 0..n*4+3 with each element having index change +-(.,.) 0,0 1,-2 or 2,-1.

Conjectures from _Colin Barker_, Jan 04 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010083 | OEIS-A264155

a(n) is the smallest integer m such that n is the least exponent k satisfying sigma(m)^k divides m.

Conjecture: for n > 1, a(n) is of the form 2^n * m generally, sometimes of the form 3^n * m, and sometimes of the form 2^(n-1) * m, depending on sigma(m). Upper bounds are mostly of the form 2^n * m for odd m. For example, a(27) <= 2^27 * 5. - _David A. Corneth_, Feb 14 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Nov 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010084 | OEIS-A264185

Number of (n+1) X (3+1) arrays of permutations of 0..n*4+3 with each element having index change +-(.,.) 0,0 1,-1 or 2,2.

Conjectures from _Colin Barker_, Jan 05 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010085 | OEIS-A264200

Numerator of sum of numbers in set g(n) generated as in Comments.

Conjecture: a(n) = 3*a(n-1) + 4*a(n-2) - 8*a(n-3) - 8*a(n-4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010086 | OEIS-A264273

Number of (2+1) X (n+1) arrays of permutations of 0..n*3+2 with each element having index change +-(.,.) 0,0 1,0 or 1,2.

Conjectures from _Colin Barker_, Jan 07 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 10 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010087 | OEIS-A264337

Number of (n+1) X (4+1) arrays of permutations of 0..n*5+4 with each element having index change +-(.,.) 0,0 0,1 or 1,2.

Conjectures from _Colin Barker_, Jan 07 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 11 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010088 | OEIS-A264354

Even nonsquare D values which admit proper solutions to the Pell equation x^2 - D*y^2 = +8.

The proper positive fundamental solution (x1(n), y1(n)) of the first class is given by (2*A261247(n), A261248(n)) for D(n) = a(n), n >= 1. If there are two classes the proper

positive fundamental solution (x2(n), y2(n)) for the second class is given by (A264357(n), A264386(n)) for D(n). If the fundamental solutions of the two classes coincide then there is only one class (the ambiguous case) for these D(n) values. It is conjectured that there are no more than two classes. For the computation of (x2(n), y2(n)) from (x1(n), -y1(n)) by application of the matrix M(n) for D(n) see a comment under A263012.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Nov 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010089 | OEIS-A264515

Number of (n+1)X(3+1) arrays of permutations of 0..n*4+3 with each element having directed index change -1,0 0,2 -1,-2 or 1,0.

Conjecture: a(n) = A264507(n)^2. - _Bruce Nye_, Feb 11 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010090 | OEIS-A264591

Let G[1](q) denote the g.f. for A003114 and G[2](q) the g.f. for A003106 (the two Rogers-Ramanujan identities). For i>=3, let G[i](q) = (G[i-1](q)-G[i-2](q))/q^(i-2). Sequence gives coefficients of G[4](q).

It is conjectured that G[i](q) = 1 + O(q^i) for all i.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 18 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010091 | OEIS-A264592

Let G[1](q) denote the g.f. for A003114 and G[2](q) the g.f. for A003106 (the two Rogers-Ramanujan identities). For i>=3, let G[i](q) = (G[i-1](q)-G[i-2](q))/q^(i-2). Sequence gives coefficients of G[5](q).

It is conjectured that G[i](q) = 1 + O(q^i) for all i.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010092 | OEIS-A264593

Let G[1](q) denote the g.f. for A003114 and G[2](q) the g.f. for A003106 (the two Rogers-Ramanujan identities). For i>=3, let G[i](q) = (G[i-1](q)-G[i-2](q))/q^(i-2). Sequence gives coefficients of G[6](q).

It is conjectured that G[i](q) = 1 + O(q^i) for all i.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010093 | OEIS-A264594

Let G[1](q) denote the g.f. for A003114 and G[2](q) the g.f. for A003106 (the two Rogers-Ramanujan identities). For i>=3, let G[i](q) = (G[i-1](q)-G[i-2](q))/q^(i-2). Sequence gives coefficients of G[7](q).

It is conjectured that G[i](q) = 1 + O(q^i) for all i.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010094 | OEIS-A264595

Let G[1](q) denote the g.f. for A003114 and G[2](q) the g.f. for A003106 (the two Rogers-Ramanujan identities). For i>=3, let G[i](q) = (G[i-1](q)-G[i-2](q))/q^(i-2). Sequence gives coefficients of G[8](q).

It is conjectured that G[i](q) = 1 + O(q^i) for all i.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010095 | OEIS-A264596

Let S_n be the list of the first n nonnegative numbers written in binary, with least significant bits on the left, and sorted into lexicographic order; a(n) = position of n in S_n, starting indexing at 0.

Conjecture: a(n) = n*(n + 3)/2 - A007814(A293290(n)) for n > 0. - _Velin Yanev_, Sep 12 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010096 | OEIS-A264613

Numbers n such that the Shevelev polynomial {m, n} has a root at m = -1.

Conjectured g.f.: (2 + x*(5 + x*(8 + x*(1 + (-2 - 8*x)* x)))) / (1 + x^3*(-5 + 4*x^3)). - _Peter J. C. Moses_, Dec 12 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010097 | OEIS-A264745

Rectangular array A read by upward antidiagonals in which the entry in row n and column k is defined by A(n,k) = Fibonacci(2^(n-1)*(2*k-1) + 1), n,k >= 1.

Conjectured g.f. for row n: x*(A000045(2^(n-1)+1) - A000045(2^(n-1)-1)*x)/(1 - A001566(n)*x + x^2), n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Nov 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010098 | OEIS-A264752

First differences of A106543.

It is conjectured that, for n > 1, a(n) < 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gionata Neri_, Jul 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010099 | OEIS-A264784

Run lengths of zeros in A265158.

Conjecture: A000975(n) = a(A155051(n-1)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Dec 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010100 | OEIS-A264789

Number of steps needed to reach 1 or to enter the cycle in the "sqrt(3)*x+1" problem.

(iii) The trajectory is divergent trajectory (I conjecture that this cannot occur).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Nov 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010101 | OEIS-A264797

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 342", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 03 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010102 | OEIS-A264803

Numbers with largest ratio A003313(k)/log_2(k) in the range 2^n < k < 2^(n+1).

The quotient A003313(k)/log_2(k) has its conjectured maximum of 1.46347481 for k=71. Values of A003313 up to 2^31-1 are obtained from Achim Flammenkamp's web page, which provides a table computed by _Neill M. Clift_. In the paper by Wattel & Jensen, the conjectured maximum is proved to hold for all k > 71, too. - _Achim Flammenkamp_, Nov 01 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 17 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010103 | OEIS-A264811

Number of steps needed to reach 1 or to enter the cycle in the double '3x+1' and '5x+1' problem.

It is conjectured that the process always stabilize. (iii) Divergent trajectory (it is conjectured that this behavior cannot occur).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Nov 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010104 | OEIS-A264865

Primes of the form 2^x + y (x >= 0 and 0 <= y < 2^x) such that all the numbers 2^(x-a) + (y+a) (0 < a <= x) are composite.

Conjecture: The sequence has infinitely many terms. This is motivated by part (i) of the conjecture in A231201. See also A264866 for a related conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010105 | OEIS-A264904

Primes of the form x^2 + y^2 with 0 < x < y such that all the numbers (x-a)^2 + (y+a)^2 (a = 1,...,x) are composite.

Note that the sequence contains all primes of the form n^2 + 1 with n > 1. A conjecture of Landau states that there are infinitely many primes of the form n^2 + 1. Conjecture: For any prime p > 5 of the form x^2 + y^2 (0 < x < y), there is a prime q not equal to p of the form u^2 + v^2 (0 < u < v) with u + v = x + y. Conjecture: each odd number m > 1 is a unique sum m = x + y with 0 < x < y, where x^2 + y^2 is in the sequence. - _Thomas Ordowski_, Jan 16 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010106 | OEIS-A264906

a(n) is the denominator of the 2nd term of the power series which is the loop length in a regular n-gon. (See comment.)

The values of the loop length L(n) appear to form a power series. Conjectures: the coefficient of the first term is 2*A060819; the numerator and denominator of the coefficient of the 2nd term are -1*A000265 and a(n), respectively; and the numerator of the coefficient of the 3rd term is A109375.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Nov 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010107 | OEIS-A264923

Expansion of g.f. 1 / Product_{n>=0} (1 - x^(n+3))^((n+1)*(n+2)/2!).

Number of partitions of n objects of 3 colors, where each part must contain at least one of each color. [Conjecture - see comment by _Franklin T. Adams-Watters_ in A052847].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010108 | OEIS-A264924

G.f.: 1 / Product_{n>=0} (1 - x^(n+4))^((n+1)*(n+2)*(n+3)/3!).

Number of partitions of n objects of 4 colors, where each part must contain at least one of each color. [Conjecture - see comment by Franklin T. Adams-Watters in A052847].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010109 | OEIS-A264925

G.f.: 1 / Product_{n>=0} (1 - x^(n+5))^((n+1)*(n+2)*(n+3)*(n+4)/4!).

Number of partitions of n objects of 5 colors, where each part must contain at least one of each color. [Conjecture - see comment by Franklin T. Adams-Watters in A052847].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010110 | OEIS-A264926

G.f.: 1 / Product_{n>=0} (1 - x^(n+6))^((n+1)*(n+2)*(n+3)*(n+4)*(n+5)/5!).

Number of partitions of n objects of 6 colors, where each part must contain at least one of each color. [Conjecture - see comment by Franklin T. Adams-Watters in A052847].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010111 | OEIS-A264938

a(n) = n*(2*n-1) + floor(n/3).

Conjecture: after 0, a(n) provides the first bisection of A264041. Conjecture: 2, 9, 20, 34, 53, 76, 102, 133, ... is A248121.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Nov 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010112 | OEIS-A264946

Number of 3 X n arrays containing n copies of 0..3-1 with no equal horizontal neighbors and new values introduced sequentially from 0.

Conjectured recurrence of order 9 and degree 13: (n + 7)*(n + 9)*(n + 11)*(n + 12)*(1125*n^9 + 71100*n^8 + 1849290*n^7 + 25782456*n^6 + 208721101*n^5 + 972463852*n^4 + 2219593700*n^3 + 50298752*n^2 - 10311481536*n - 14857032960)*a(n + 9) - (n + 8)*(n + 11)*(1125*n^11 + 105975*n^10 + 4287165*n^9 + 97635351*n^8 + 1380358747*n^7 + 12566316081*n^6 + 73253605103*n^5 + 255547606333*n^4 + 398433897060*n^3 - 409020958588*n^2 - 2573789550288*n - 2893020641280)*a(n + 8) - (n + 7)*(55125*n^12 + 5043150*n^11 + 203593785*n^10 + 4783109874*n^9 + 72494563771*n^8 + 740695604282*n^7 + 5151899227595*n^6 + 23797245731102*n^5 + 66502991649164*n^4 + 73821686951912*n^3 - 148787634331808*n^2 - 594255830565888*n - 585045220193280)*a(n + 7) + (-111375*n^13 - 10629900*n^12 - 452242260*n^11 - 11349359364*n^10 - 187042020462*n^9 - 2127929201500*n^8 - 17043729555112*n^7 - 95651673757276*n^6 - 362118971182795*n^5 - 819923758737640*n^4 - 569404973885116*n^3 + 2298397321892016*n^2 + 6690359386550016*n + 5779315376271360)*a(n + 6) + 4*(68625*n^13 + 6350850*n^12 + 262268790*n^11 + 6397257006*n^10 + 102643250092*n^9 + 1138907266426*n^8 + 8907524302014*n^7 + 48769985579898*n^6 + 178866084976275*n^5 + 380439835948764*n^4 + 162302253918524*n^3 - 1409022793603840*n^2 - 3584273436805056*n - 2901502849512960)*a(n + 5) + 8*(113625*n^13 + 10109475*n^12 + 400010115*n^11 + 9340598391*n^10 + 143726469497*n^9 + 1536021852345*n^8 + 11649644036681*n^7 + 62442624371309*n^6 + 227373943553018*n^5 + 494618567601008*n^4 + 295071232164264*n^3 - 1523160592469296*n^2 - 4224581839405248*n - 3597712690682880)*a(n + 4) + 16*(10125*n^13 + 791775*n^12 + 27734085*n^11 + 599490219*n^10 + 9285055875*n^9 + 110260636645*n^8 + 1005826848007*n^7 + 6794028883657*n^6 + 32142152171668*n^5 + 97251455731576*n^4 + 145564079336272*n^3 - 68830984119216*n^2 - 616978454170176*n - 699609347458560)*a(n + 3) - 16*(102375*n^13 + 8429850*n^12 + 302519490*n^11 + 6252315786*n^10 + 82728789292*n^9 + 735119632054*n^8 + 4454619907050*n^7 + 18120079094814*n^6 + 45886145527965*n^5 + 51009713975544*n^4 - 7811809365449 2*n^3 - 396941915101552*n^2 - 632960801821824*n - 397766739363840)*a(n + 2) - 64*(n + 1)*(28125*n^11 + 2145375*n^10 + 69983925*n^9 + 1286257695*n^8 + 14737434403*n^7 + 109491713225*n^6 + 526473720215*n^5 + 1544097805013*n^4 + 2140227079172*n^3 - 1382380121116*n^2 - 9705594256208*n - 11040556116480)*n*a(n + 1) - 512*(n - 1)*(n + 1)*(1125*n^9 + 81225*n^8 + 2458590*n^7 + 40812786*n^6 + 406374277*n^5 + 2472650097*n^4 + 8781110488*n^3 + 15058677204*n^2 + 1549576672*n - 21689733120)*n^2*a(n) = 0. - _Manuel Kauers_ and _Christoph Koutschan_, Mar 06 2023 Conjecture: a(n) ~ 3^(7/2) * 2^(3*n - 5) / (Pi*n), based on the recurrence by _Manuel Kauers_ and _Christoph Koutschan_. - _Vaclav Kotesovec_, Mar 07 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010113 | OEIS-A265035

Coordination sequence of 2-uniform tiling {3.4.6.4, 4.6.12} with respect to a point of type 4.6.12.

Conjectured g.f. is equivalent to a(n) = 3*n - A010892(n+1) for n >= 5. - _R. J. Mathar_, Oct 09 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 12 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010114 | OEIS-A265100

a(n) = 9*A005836(n) + 5, n >= 1.

Let C(m) denote the m-th Catalan number (A000108). Let == denote congruence and =!= its negation. _Vladimir Reshetnikov_ asked (see link) how many n exist such that C(n) == 1 (mod 6). It was pointed out by _Robert Israel_ that the only known n are in {1,3,31,255}. Since C(n) is odd if and only if n = 2^m - 1, for some m, _Emmanuel Vantieghem_ (see links) stated the stronger conjecture that C(2^n-1) == 0 (mod 3), for all n>8. This is the motivation for the following. If m is an integer such that the congruences C(m) =!= 0 (mod 3) and C(m-1) == 0 (mod 3) hold simultaneously, then we call m a "gap number." A sequence {m, m+1, ..., m+j-1} of consecutive numbers is called a "gap" (of order j), if C(m+i) =!= 0 (mod 3), for all i such that 0 <= i < j, and if C(m-1) == 0 (mod 3) (i.e., if m is a gap number) and C(m+j) == 0 (mod 3). (The sequence A265104 is conjectured to contain all possible gap numbers.) If C(n) == 0 (mod 3), then we say that n is "gap-avoiding." Conjecture 1: The sequence contains all possible block numbers. Conjecture 2: If m is a block number, then 3*m - 1 is a block number. Conjecture 3: If C(n) == 0 (mod 3), then C(3*n-1) == 0 (mod 3) or, what is the same thing, if n lies in a block, then 3*n - 1 lies in a block. Conjecture 4: Assuming that A265104 contains all possible gap numbers, let B(n) denote the block with block number a(n), n >= 1, so that B(n) = {a(n), a(n)+1, ..., A265104(n)-1}. The (flattened) sequence {B(1), B(2), ...} of blocks contains all numbers m such that the base 3 representations of m and m+1 both contain at least one 2 and is identical to A111018. Conjecture 5: C(n) == 0 (mod 3) if and only if the base 3 representations of n and n + 1 both contain at least one 2. [This conjecture has been proved by _Robert Israel_ (see link for the proof)]. Theorem 1: The following statements are equivalent to Vantieghem's conjecture stated above: (i) For all m>8, 2^m-1 is gap-avoiding; (ii) C(2^n-1) == 0 (mod 3) if and only if the base 3 representations of 2^n - 1 and 2^n both contain at least one 2. Proof: For (i), the statement obviously follows from the definitions, and (ii) follows from the proof of Conjecture 5. Conjecture: a(n) = A265104(n) - A085296(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Dec 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010115 | OEIS-A265108

The number of n-digit numbers in A264847 (pluritriangular numbers).

Conjecture: lim_{n -> infinity} a(n)/a(n-1) = sqrt(10).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Francesco Di Matteo_, Dec 01 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010116 | OEIS-A265159

Rectangular array A read by upward antidiagonals in which the entry in row n and column k is defined by A(n,k) = 5 + 9*A005836(2^(k - 1)*(2 n - 1)), n,k >= 1.

Conjecture 1: The array contains without duplication all possible "block numbers" defined in A265100. Conjecture 2: A(n,k) = (A055246(n)*3^k + 1)/2, so the array and A265100 are related to Cantor's ternary set. G.f. for row n (conjectured): f(n,x) = x*(A265100(n)-(A265100(n)+1)*x)/((1-x)*(1-3*x)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Dec 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010117 | OEIS-A265161

Array A read by upward antidiagonals in which the entry in row n and column k is defined by A(n,k) = (3/2)*(3^k - 1) + A265159(n,k), n,k >= 1.

Conjecture 1: The array contains without duplication all possible "gap numbers" as defined in A265100. Conjecture 2: A(n,k) = A191107(n)*3^k - 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Dec 03 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010118 | OEIS-A265196

Irregular triangle read by rows, where T(n, k) is the coefficient of degree k of the polynomial Product_{j=0..n} (1-x^(3*j+1))*(1-x^(3*j+2)).

Borwein conjectures that T(n,k) >= 0 when k is a multiple of 3, and T(n,k) <= 0 is not a multiple of 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Dec 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010119 | OEIS-A265207

Draw a square and follow these steps: Take a square and place at its edges isosceles right triangles with the edge as hypotenuse. Draw a square at every new edge of the triangles. Repeat for all the new squares of the same size. New figures are only placed on empty space. The structure is symmetric about the first square. The sequence gives the numbers of squares of equal size in successive rings around the center.

Conjectured recurrence: Conjectured formula: ("[]" is the floor function) Conjectures from _Colin Barker_, Dec 07 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marian Kraus_, Dec 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010120 | OEIS-A265210

Irregular triangle read by rows in which row n lists the base 3 digits of 2^n in reverse order, n >= 0.

Conjecture 1: The sequence in column k is periodic, with period p(k) = 2*3^(k-1) = A008776(k-1), k >= 1, and in which the numbers 0,1,2 appear with equal frequency, for each k>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Dec 04 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010121 | OEIS-A265225

Total number of ON (black) cells after n iterations of the "Rule 54" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 08 2015 and Apr 20 2019: (Start) a(n) = n + 1 + (n+1) * floor((n+1)/2), conjectured. - _Wesley Ivan Hurt_, Dec 25 2016 a(n) = A093353(n) + n + 1, conjectured. - _Matej Veselovac_, Jan 21 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 05 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010122 | OEIS-A265233

Number of 3 X n arrays containing n copies of 0..2 with no equal vertical neighbors and new values introduced sequentially from 0.

Conjecture: n^2*a(n) +(-19*n^2+19*n-6)*a(n-1) +96*(n-1)^2*a(n-2) -144*(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Dec 08 2015 Conjecture: a(n) ~ 2^(2*n - 1) * 3^(n - 1/2) / (Pi*n). - _Vaclav Kotesovec_, Mar 08 2023 If conjectured recurrence is true then ogf = (hypergeom([1/3,2/3],[1],27*x*(4*x-1)^2)+5)/6. - _Mark van Hoeij_, Nov 28 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010123 | OEIS-A265251

Number of partitions of n such that there is exactly one part which occurs three times, while all other parts occur only once.

Conjecture: a(n) is also the difference between the number of parts in the distinct partitions of n and the number of distinct parts in the odd partitions of n (offset 0). For example, if n = 5, there are 5 parts in the distinct partitions of 5 (5, 41, 32) and 4 distinct parts in the odd partitions of 5 (namely, 5,3,1,1 in 5,311,11111) with difference 1. - _George Beck_, Apr 22 2017 George E. Andrews has kindly informed me that he has proved this conjecture and the result will be included in his article "Euler's Partition Identity and Two Problems of George Beck" which will appear in The Mathematics Student, 86, Nos. 1-2, January - June (2017). - _George Beck_, Apr 23 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Dec 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010124 | OEIS-A265262

The tree of hemitropic sequences read by rows, arising from an Erdős-Turán conjecture.

The tree of hemitropic sequences read by rows, arising from an Erdős-Turán conjecture. An [order 2 additive] basis of N is a subset A whose profile has no 0's. Erdős and Turán conjectured that the profile of a basis is always unbounded (see the Erdős and Turán link). The conjecture is, up till now,

still undecided. The tree below displays the infinite sequences [1, pA(2), ...], associated to the profiles pA = [1, 1, pA(2), ...] of all the subsets A of N to which 0 and 1 belong. Those are the so-called hemitropic sequences. The Erdős-Turán conjecture would not hold if (and only if) the tree contained an infinite bounded branch with no 0's.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Labib Haddad_ and _Michel Marcus_, Dec 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010125 | OEIS-A265283

Number of ON (black) cells in the n-th iteration of the "Rule 94" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 07 2015 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010126 | OEIS-A265284

Total number of ON (black) cells after n iterations of the "Rule 94" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 07 2015 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010127 | OEIS-A265287

Number of steps needed to reach 1 in the "sqrt(2)*x + 1" problem.

Conjecture: the trajectory reaches 1 for all n. (iii) The trajectory is divergent (I conjecture that this cannot occur).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Dec 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010128 | OEIS-A265310

Least positive k such that the product of divisors of n (A007955) divides k!.

Conjecture: a(n) = n if and only if n is prime, 2*prime, 1, 8 or 9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gionata Neri_, Dec 06 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010129 | OEIS-A265380

Binary representation of the middle column of the "Rule 158" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 14 2015 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010130 | OEIS-A265382

Total number of ON (black) cells after n iterations of the "Rule 158" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 07 2015 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010131 | OEIS-A265385

Sequence defined by a(1)=a(2)=1 and a(n) = gray(a(n-1) + a(n-2)), with gray(m) = A003188(m).

Conjecture: The mean growth rate r(n) = (a(2n)/a(n))^(1/n) appears to converge exactly to 2, with the consecutive-terms ratio s(n) = a(n)/a(n-1) exhibiting relatively small (~1%) but persistent fluctuations around the mean value. This contrasts what one might first expect, that sequence's growth rate were similar to that of the Fibonacci sequence, i.e., the golden ratio, since gray(m) just permutes every block of numbers ranging from 2^k to 2^l-1, for any 0<k<l.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Dec 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010132 | OEIS-A265386

Sequence defined by a(1)=a(2)=1 and a(n) = gray(gray(a(n-1)) + gray(a(n-2))), with gray(m) = A003188(m).

Conjecture: the mean growth rate r(n) = (a(2n)/a(n))^(1/n) appears to converge to sqrt(2), with the consecutive-terms ratio s(n) = a(n)/a(n-1) exhibiting large and persistent fluctuations around the mean value.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Dec 07 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010133 | OEIS-A265416

Number of steps needed to reach 1 or to enter the cycle in the "sqrt(Pi)*x+1" problem.

(iii) The trajectory is divergent (I conjecture that this cannot occur).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Dec 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010134 | OEIS-A265427

Binary representation of the n-th iteration of the "Rule 188" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 09 2015 and Apr 18 2019: (Start) Conjecture: a(n) = floor(11100*100^n/9999) + 10^n - floor(11100*10^n/9999)*10^n. - _Karl V. Keller, Jr._, Dec 15 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010135 | OEIS-A265428

Number of ON (black) cells in the n-th iteration of the "Rule 188" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 09 2015 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010136 | OEIS-A265429

Total number of ON (black) cells after n iterations of the "Rule 188" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 09 2015 and Apr 16 2019: (Start) Conjecture: the sequence consists of all numbers k > 0 such that floor(sqrt(8*(k+1)/3)) != floor(sqrt(8*k/3)). - _Gevorg Hmayakyan_, Sep 01 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010137 | OEIS-A265430

Number of OFF (white) cells in the n-th iteration of the "Rule 188" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 09 2015 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010138 | OEIS-A265431

Total number of OFF (white) cells after n iterations of the "Rule 188" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 09 2015 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 08 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010139 | OEIS-A265436

a(n) is the least m (1 <= m <= n) such that the set of pairs (x, y) of distinct terms from [m, n] can be ordered in such a way that the corresponding sums (x+y) and products (x*y) are monotonic.

Conjecture (derived from the assumed relationship with A035106): for n>5, if sqrt(4n+1) is an odd integer or sqrt(n+1) is an integer, then a(n) = a (n-1), otherwise a(n) = a(n-1)+1. - _Jean-François Alcover_, Dec 21 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bill McEachen_ and _Michel Marcus_, Dec 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010140 | OEIS-A265437

Numbers n such that A265433(n) = 0 and n is not a multiple of 3.

Conjecture: if n > 4 is in the sequence, then A137269(n) > 0 due to primes with only digits 2, 3, 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Dec 10 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010141 | OEIS-A265481

Numbers k such that 9^k - 3^k - 1 is prime.

Conjectures: a(n) is not of the form 7*k+4 or 8*k. a(17) = 11071 mod 7 is 4, so the first half of the conjecture above is not true. - _Robert Price_, Sep 25 2019 a(23) = 78008 is divisible by 8, so the second half of the conjecture above is not true. - _Robert Price_, Sep 25 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Dec 09 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010142 | OEIS-A265482

Numbers k such that 16^k - 4^k - 1 is prime.

Conjecture: the odd terms are not of the form 8*k+7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Dec 10 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010143 | OEIS-A265687

Numbers k such that the concave polygon formed by the trajectory of k in the "3x+1" problem and the straight line between the coordinate points (0,k) and (r,1) where r is the number of iterations needed to reach 1 is not a self-intersecting polygon.

Conjecture: the sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Dec 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010144 | OEIS-A265716

a(n) = n IMPL (2*n), where IMPL is the bitwise logical implication.

a(n)= bitor(A003817(n)-n, 2*n) (conjectured). - _Bill McEachen_, Dec 13 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reinhard Zumkeller_, Dec 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010145 | OEIS-A265722

Number of ON (black) cells in the n-th iteration of the "Rule 1" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 14 2015 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010146 | OEIS-A265723

Number of OFF (white) cells in the n-th iteration of the "Rule 1" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 15 2015 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010147 | OEIS-A265724

Total number of OFF (white) cells after n iterations of the "Rule 1" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 16 2015 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010148 | OEIS-A265733

Number of n-digit primes whose digits include at least n-1 digits "1".

Inspired by A241100 and its conjecture by _Chai Wah Wu_ of Dec 10 2015.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_ and _Robert G. Wilson v_, Dec 14 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010149 | OEIS-A265735

Integers in the interval [Pi*k - 1/k, Pi*k + 1/k] for some k > 0.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Dec 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010150 | OEIS-A265739

Numbers k such that there exists at least one integer in the interval [Pi*k - 1/k, Pi*k + 1/k].

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Dec 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010151 | OEIS-A265740

a(1)=1; a(n+1) is the smallest positive integer not yet used such that all the digits of a(n) and a(n+1) are present in the decimal expansion (including any leading and trailing zeros) of a(n)/a(n+1).

Conjecture: a(n) is a permutation of the natural numbers.

which seems to support the conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_, submitted by _Lars Blomberg_, Dec 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010152 | OEIS-A265741

Integers in the interval [e*k - 1/k, e*k + 1/k] for some k > 0 , where e = 2.71828... is Euler's number.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Dec 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010153 | OEIS-A265742

Numbers k such that there exists at least an integer in the interval [e*k - 1/k, e*k + 1/k] where e = 2.71828... is Euler's number.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Dec 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010154 | OEIS-A265756

a(1)=1; a(n+1) is the smallest positive integer not yet used where all the digits of a(n) and a(n+1) are present in the decimal expansion (excluding any leading or trailing zeros) of a(n)/a(n+1).

Conjecture: a(n) is a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hans Havermann_, Dec 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010155 | OEIS-A265759

Numerators of primes-only best approximates (POBAs) to 1; see Comments.

See A265772 and A265774 for definitions of lower POBA and upper POBA. In the following guide, for example, A001359/A006512 represents (conjecturally in some cases) the Lower POBAs p(n)/q(n) to 1, where p = A001359 and q = A006512 except for first terms in some cases. Every POBA is either a lower POBA or an upper POBA.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 15 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010156 | OEIS-A265840

Expansion of Product_{k>=1} (1 + k^3*x^k).

Conjecture: log(a(n)) ~ 3*sqrt(n/2) * (log(2*n) - 2). - _Vaclav Kotesovec_, Dec 27 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Dec 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010157 | OEIS-A265841

Expansion of Product_{k>=1} (1 + k^4*x^k).

Conjecture: log(a(n)) ~ 4*sqrt(n/2) * (log(2*n) - 2). - _Vaclav Kotesovec_, Dec 27 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Dec 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010158 | OEIS-A265842

Expansion of Product_{k>=1} (1 + k^5*x^k).

Conjecture: log(a(n)) ~ 5*sqrt(n/2) * (log(2*n) - 2). - _Vaclav Kotesovec_, Dec 27 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Dec 16 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010159 | OEIS-A265942

G.f. A(x) satisfies: x = Sum_{n>=1} x^n * A(-n*x).

a(n) = 2 (mod 4) for n>2 [conjecture].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010160 | OEIS-A265946

Sums of positive differences and negated sums of negative differences in the Collatz trajectory of n.

I present an interesting conjecture because of the linearity of the result. Conjecture: a(2n) - a(2n-1) = n - 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Dec 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010161 | OEIS-A265987

Number of n X 3 integer arrays with each element equal to the number of horizontal and antidiagonal neighbors not equal to itself.

Conjectures from _Colin Barker_, Jan 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010162 | OEIS-A265989

Number of n X 6 integer arrays with each element equal to the number of horizontal and antidiagonal neighbors not equal to itself.

Conjectures from _Colin Barker_, Jan 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010163 | OEIS-A266003

Least nonnegative integer y such that n = x^4 - y^3 + z^2 for some nonnegative integers x and z, or -1 if no such y exists.

Conjecture: Any integer m can be written as x^4 - y^3 + z^2, where x, y and z are nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 19 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010164 | OEIS-A266072

Number of ON (black) cells in the n-th iteration of the "Rule 3" elementary cellular automaton starting with a single ON (black) cell.

Conjectured g.f.: (1 + x - x^2 + 3*x^3)/(-1 + x^2)^2. - _Michael De Vlieger_, Dec 21 2015 Conjectures from _Colin Barker_, Dec 21 2015: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010165 | OEIS-A266073

Number of OFF (white) cells in the n-th iteration of the "Rule 3" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 21 2015 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 20 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010166 | OEIS-A266118

a(n) = index of n in A266117 or 0 if n is not present in that sequence.

It is conjectured that A266117 is not just injective, but also surjective on N, i.e., that it is a true permutation of natural numbers. In that case also this sequence is a true permutation, and no hypothetical zero-values are ever needed.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010167 | OEIS-A266121

Lexicographically first injection of natural numbers beginning with a(1)=1 such that 1+(a(n)*a(n+1)) is a fibbinary number (A003714), i.e., has no adjacent 1's in its base-2 representation.

It is conjectured that this sequence is not only injective, but also surjective on N, i.e., that it is a true permutation of natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010168 | OEIS-A266152

Least positive integer y such that n = x^4 - y^3 + z^2 for some positive integers x and z, or 0 if no such y exists.

Conjecture: Any integer m can be written as x^4 - y^3 + z^2, where x, y and z are positive integers. This is slightly stronger than the conjecture in A266003. See also A266153 for a related sequence, and A266212 for a stronger conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 22 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010169 | OEIS-A266181

Numbers k such that k == d_1 (mod 2), k == d_2 (mod 3), k == d_3 (mod 4) etc., where d_1 d_2 d_3 ... is the decimal expansion of k.

If it exists, a(14) >= 10^45 (cf. A284814). Conjecturally the sequence is finite and complete. - _Max Alekseyev_, Jul 26 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pieter Post_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010170 | OEIS-A266195

Match-making permutation: start with a(1) = 1, then always choose for a(n) the least unused number such that multiplying a(n) by a(n-1) does not produce any carries when performed in base 2.

Sequence A266197 gives the positions of these good matches, and A265748 & A265749 give their first and second members respectively. It is an open question whether the algorithm generates an infinite number of good matches or not.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010171 | OEIS-A266212

Positive integers x such that x^3 = y^4 + z^2 for some positive integers y and z.

Conjecture: For any integer m, there are infinitely many triples (x,y,z) of positive integers with x^4 - y^3 + z^2 = m. This is stronger than the conjecture in A266152.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 23 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010172 | OEIS-A266215

Positive integers x such that x^3 - 1 = y^4 + z^2 for some positive integers y and z.

The conjecture in A266212 implies that this sequence has infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010173 | OEIS-A266219

Binary representation of the middle column of the "Rule 7" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 25 2015 and Apr 13 2019: (Start) Conjecture: a(n) = floor(109*10^n/99). - _Karl V. Keller, Jr._, Mar 16 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010174 | OEIS-A266220

Number of ON (black) cells in the n-th iteration of the "Rule 7" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 25 2015 and Apr 13 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010175 | OEIS-A266221

Total number of ON (black) cells after n iterations of the "Rule 7" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 25 2015 and Apr 13 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010176 | OEIS-A266222

Number of OFF (white) cells in the n-th iteration of the "Rule 7" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 26 2015 and Apr 13 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010177 | OEIS-A266223

Total number of OFF (white) cells after n iterations of the "Rule 7" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 26 2015 and Apr 14 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010178 | OEIS-A266224

Least x such that prime(n)*x+x+1 is a prime, or -1 if no such x exists.

Conjecture: a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Dec 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010179 | OEIS-A266225

Least x>1 such that prime(n)*x+x-1 is a prime, or -1 if no such x exists.

Conjecture: a(n) > 1. This conjecture follows from Dirichlet's prime number theorem. - _David Radcliffe_, May 22 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Dec 24 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010180 | OEIS-A266239

a(n) is the smallest k such that the following are four primes: prime(n)*k-1, prime(n)*k+1, prime(n)*k^2-1, prime(n)*k^2+1. Or -1 if no such k exists.

Conjecture: a(n)>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Dec 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010181 | OEIS-A266249

Number of ON (black) cells in the n-th iteration of the "Rule 9" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 28 2015 and Apr 14 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010182 | OEIS-A266250

Total number of ON (black) cells after n iterations of the "Rule 9" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 28 2015 and Apr 14 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010183 | OEIS-A266251

Number of OFF (white) cells in the n-th iteration of the "Rule 9" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 28 2015 and Apr 14 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010184 | OEIS-A266252

Total number of OFF (white) cells after n iterations of the "Rule 9" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 28 2015 and Apr 14 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010185 | OEIS-A266256

Number of ON (black) cells in the n-th iteration of the "Rule 11" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 27 2015 and Apr 14 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010186 | OEIS-A266257

Total number of ON (black) cells after n iterations of the "Rule 11" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 27 2015 and Apr 14 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010187 | OEIS-A266258

Number of OFF (white) cells in the n-th iteration of the "Rule 11" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 27 2015 and Apr 14 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010188 | OEIS-A266259

Total number of OFF (white) cells after n iterations of the "Rule 11" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 27 2015 and Apr 14 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 25 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010189 | OEIS-A266277

Least positive integer x such that n + x^2 = y^3 + z^5 for some positive integers y and z, or 0 if no such x exists.

Conjecture: If {a,b,c} is among the multisets {2,2,p} (p is an odd prime or a product of primes congruent to 1 modulo 4) and {2,3,k} (k = 3,4,5), then for any integer m there are (infinitely many) triples (x,y,z) of positive integers such that m = x^a + y^b - z^c. This implies that a(n) > 0 for all n. Also, it includes the conjectures in A266152, A266212 and A266230 as special cases.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010190 | OEIS-A266285

Number of ON (black) cells in the n-th iteration of the "Rule 13" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 28 2015 and Apr 14 2019: (Start) Conjecture from _Ctibor O. Zizka_, Mar 11 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010191 | OEIS-A266286

Number of OFF (white) cells in the n-th iteration of the "Rule 13" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 28 2015 and Apr 14 2019: (Start) Conjecture: a(n) = (n^2)mod(2n-3) for n>6. - _Andres Cicuttin_, Aug 29 2019 Conjecture: E.g.f.: (1/4)*exp(-x)*(- 3 - 2*x + exp(2*x)*(3 + 4*x)). - _Stefano Spezia_, Aug 31 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010192 | OEIS-A266287

Total number of OFF (white) cells after n iterations of the "Rule 13" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 28 2015 and Apr 14 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010193 | OEIS-A266303

Number of ON (black) cells in the n-th iteration of the "Rule 15" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 28 2015 and Apr 15 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010194 | OEIS-A266304

Total number of OFF (white) cells after n iterations of the "Rule 15" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 28 2015 and Apr 15 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 26 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010195 | OEIS-A266327

Binary representation of the n-th iteration of the "Rule 20" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 28 2015 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 27 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010196 | OEIS-A266378

Recursive 2-parameter sequence allowing calculation of the Möbius function.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gevorg Hmayakyan_, Dec 28 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010197 | OEIS-A266412

Inverse permutation to A266411.

Conjecture: a(2^(k-1)+1) = 2^k, for k > 0. - _Ruud H.G. van Tol_, Apr 03 2026 (see A004001)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010198 | OEIS-A266429

Number of 3 X n binary arrays with rows and columns lexicographically nondecreasing and column sums nondecreasing.

Conjectures from _Colin Barker_, Jan 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010199 | OEIS-A266430

Number of 4 X n binary arrays with rows and columns lexicographically nondecreasing and column sums nondecreasing.

Conjectures from _Colin Barker_, Jan 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010200 | OEIS-A266437

Number of ON (black) cells in the n-th iteration of the "Rule 23" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 30 2015 and Apr 15 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010201 | OEIS-A266438

Total number of ON (black) cells after n iterations of the "Rule 23" elementary cellular automaton starting with a single ON (black) cell.

Conjectured g.f.: (-(1 + x)^(-2) + (-3 + 3*x - 2*x^2)/(-1 + x)^3)/2. - _Michael De Vlieger_, Dec 29 2015 Conjectures from _Colin Barker_, Dec 30 2015 and Apr 15 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010202 | OEIS-A266439

Number of OFF (white) cells in the n-th iteration of the "Rule 23" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 30 2015 and Apr 15 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010203 | OEIS-A266440

Total number of OFF (white) cells after n iterations of the "Rule 23" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 30 2015 and Apr 15 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010204 | OEIS-A266444

Middle column of the "Rule 25" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 31 2015 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010205 | OEIS-A266445

Binary representation of the middle column of the "Rule 25" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 31 2015 and Apr 16 2019: (Start) Conjecture: a(n) = floor(9901000008901*10^n/9900000000000). - _Karl V. Keller, Jr._, Jan 30 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010206 | OEIS-A266446

Decimal representation of the middle column of the "Rule 25" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 30 2015 and Apr 16 2019: (Start) Conjecture: a(n) = floor(6661*2^n/6144). - _Karl V. Keller, Jr._, Dec 26 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010207 | OEIS-A266447

Number of ON (black) cells in the n-th iteration of the "Rule 25" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 30 2015 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010208 | OEIS-A266448

Total number of ON (black) cells after n iterations of the "Rule 25" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 30 2015 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010209 | OEIS-A266449

Number of OFF (white) cells in the n-th iteration of the "Rule 25" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 30 2015 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010210 | OEIS-A266450

Total number of OFF (white) cells after n iterations of the "Rule 25" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 31 2015 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010211 | OEIS-A266460

Binary representation of the n-th iteration of the "Rule 27" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 31 2015 and Apr 16 2019: (Start) Conjecture: a(n) = (10*(100^n - 9*10^floor(n/2)) - 1)/9 for odd n; a(n) = 10^(n/2) for even n. - _Karl V. Keller, Jr._, Sep 29 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010212 | OEIS-A266461

Decimal representation of the n-th iteration of the "Rule 27" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 31 2015 and Apr 16 2019: (Start) Conjecture: a(n) = 2*(4^n - 2^floor(n/2)) - 1 for odd n; a(n) = 2^(n/2) for even n. - _Karl V. Keller, Jr._, Sep 29 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010213 | OEIS-A266471

Number of 4 X n binary arrays with rows and columns lexicographically nondecreasing and column sums nonincreasing.

Conjectures from _Colin Barker_, Jan 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010214 | OEIS-A266472

Number of 5 X n binary arrays with rows and columns lexicographically nondecreasing and column sums nonincreasing.

Conjectures from _Colin Barker_, Jan 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 29 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010215 | OEIS-A266489

G.f. A(x) satisfies: [x^n] A(x/A(x)^n) = 0 for n>1.

CONJECTURES:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010216 | OEIS-A266498

Index of the smallest triangular number greater than 3^n.

a(n) gives a theoretical upper bound for the number of coins such that two fake coins (of equal weight lighter than the other coins) among them can be identified in n weightings on a balance scale. It was shown that the bound is achievable for all n<=10, but it remains an open question if the bound is achievable for n>10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Dec 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010217 | OEIS-A266515

Binary representation of the n-th iteration of the "Rule 29" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Dec 31 2015 and Apr 16 2019: (Start) Conjecture: a(n) = (10*(100^n - 9*10^n) - 1)/9 for odd n; a(n) = 10^n for even n. - _Karl V. Keller, Jr._, Oct 03 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010218 | OEIS-A266516

Decimal representation of the n-th iteration of the "Rule 29" elementary cellular automaton starting with a single ON (black) cell.

Conjecture: a(n) = 2*(4^n - 2^n) - 1 for odd n; a(n) = 2^n for even n. - _Karl V. Keller, Jr._, Oct 03 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 30 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010219 | OEIS-A266532

Total number of Y-toothpicks after n-th stage in the "outward" version of the cellular automaton of A160120.

Conjecture: a(n) = 1 + 3*(A160720(n) - 1)/4 = 1 + 3*A267700(n-1), n >= 1. This formula is correct! - _N. J. A. Sloane_, Jan 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Applegate_ and _Omar E. Pol_, Jan 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010220 | OEIS-A266548

Least prime p such that n + p^3 = x^2 + y^3 for some positive integers x and y, or 0 if no such prime p exists.

Conjecture: (i) Any integer can be written as x^2 + y^3 - p^3, where x and y are positive integers, and p is a prime. See also A266230 and A266277 for related conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 31 2015 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010221 | OEIS-A266577

Square array read by descending antidiagonals: T(n,k) = ((2^(n+1) + 1)^(k-1) + 1)/2.

Conjecture: For integers n and m > 1, let b(n,m)=n^m+1, S(n,m) = set of numbers of the form (b(n,m)^k+...+b(n,m)^((n-1)*k)+1)/n, where k is any nonnegative integer. Then for each positive integer s, (b(n,m)^s-1)*(product of any m not necessarily distinct terms of S(n,m)) is palindromic in base b(n,m). - _Ahmad J. Masad_, Jan 11 2025 The conjecture is true. See my proof in MathOverflow (2025) link. - _Max Alekseyev_, May 15 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ahmad J. Masad_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010222 | OEIS-A266587

Smallest index of a Lucas number (A000032) that is divisible by prime(n), if it exists, or 0 if it does not exist (for n > 1).

Conjecture: Infinite Lucas subsequences divisible by the powers of odd primes, p(n), for which a(n) > 0, are given by:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard R. Forberg_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010223 | OEIS-A266589

Binary representation of the n-th iteration of the "Rule 37" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 01 2016 and Apr 18 2019: (Start) Conjecture: a(n) = (10*100^n - 99999*10^(n-2) - 1)/9 for odd n > 1; a(n) = 111*10^(n-1) for even n > 1. - _Karl V. Keller, Jr._, Oct 06 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010224 | OEIS-A266590

Decimal representation of the n-th iteration of the "Rule 37" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 18 2019: (Start) Conjecture: a(n) = 2*4^n - 31*2^(n-2) - 1 for odd n > 1; a(n) = 7*2^(n-1) for even n > 1. - _Karl V. Keller, Jr._, Oct 06 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010225 | OEIS-A266592

Binary representation of the middle column of the "Rule 37" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010226 | OEIS-A266593

Number of ON (black) cells in the n-th iteration of the "Rule 37" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010227 | OEIS-A266594

Total number of ON (black) cells after n iterations of the "Rule 37" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010228 | OEIS-A266595

Number of OFF (white) cells in the n-th iteration of the "Rule 37" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010229 | OEIS-A266596

Total number of OFF (white) cells after n iterations of the "Rule 37" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010230 | OEIS-A266606

Binary representation of the n-th iteration of the "Rule 39" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010231 | OEIS-A266607

Decimal representation of the n-th iteration of the "Rule 39" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010232 | OEIS-A266609

Binary representation of the n-th iteration of the "Rule 41" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 18 2019: (Start) Conjecture: a(n) = floor((10*100^n + 10 - 1010*(n mod 4))/9) - 10675*0^((n+1) mod 4) for odd n; a(n) = 101^((n mod 4)/2) for even n. - _Karl V. Keller, Jr._, Oct 07 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010233 | OEIS-A266610

Decimal representation of the n-th iteration of the "Rule 41" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 18 2019: (Start) Conjecture: a(n) = 2*4^n + 2 - 10*(n mod 4) for odd n; a(n) = 5^((n mod 4)/2) for even n. - _Karl V. Keller, Jr._, Oct 07 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010234 | OEIS-A266612

Binary representation of the middle column of the "Rule 41" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010235 | OEIS-A266613

Decimal representation of the middle column of the "Rule 41" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 16 2019: (Start) Conjecture: a(n) = A000975(n) + 20*2^(n-5), for n>2. - _Andres Cicuttin_, Mar 31 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010236 | OEIS-A266614

Number of ON (black) cells in the n-th iteration of the "Rule 41" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010237 | OEIS-A266615

Total number of ON (black) cells after n iterations of the "Rule 41" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010238 | OEIS-A266616

Number of OFF (white) cells in the n-th iteration of the "Rule 41" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010239 | OEIS-A266617

Total number of OFF (white) cells after n iterations of the "Rule 41" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 02 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010240 | OEIS-A266651

Nonnegative integers x such that x^3 + 6^3 is a sum of two squares.

Conjecture: For any integer x with gcd(x,6) = 1, the number x^3 + 6^3 is never a sum of two squares. Hao Pan at Nanjing Univ. confirmed the conjecture on Jan. 3, 2016. - _Zhi-Wei Sun_, Jan 06 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010241 | OEIS-A266660

Binary representation of the n-th iteration of the "Rule 47" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 03 2016 and Apr 18 2019: (Start) Conjecture: a(n) = (10*100^n - 100)/9 for odd n > 1; a(n) = 11 for even n > 1. - _Karl V. Keller, Jr._, Oct 10 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010242 | OEIS-A266661

Decimal representation of the n-th iteration of the "Rule 47" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 03 2016 and Apr 18 2019: (Start) Conjecture: a(n) = 2*4^n - 4 for odd n > 1; a(n) = 3 for even n > 1. - _Karl V. Keller, Jr._, Oct 10 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010243 | OEIS-A266662

Number of ON (black) cells in the n-th iteration of the "Rule 47" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 03 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010244 | OEIS-A266663

Total number of ON (black) cells after n iterations of the "Rule 47" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 03 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010245 | OEIS-A266664

Number of OFF (white) cells in the n-th iteration of the "Rule 47" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 03 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010246 | OEIS-A266665

Total number of OFF (white) cells after n iterations of the "Rule 47" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 03 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010247 | OEIS-A266667

Binary representation of the n-th iteration of the "Rule 51" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 03 2016 and Apr 18 2019: (Start) Conjecture: a(n) = (10*100^n - 9*10^n - 1)/9 for odd n; a(n) = 10^n for even n. - _Karl V. Keller, Jr._, Oct 12 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010248 | OEIS-A266668

Decimal representation of the n-th iteration of the "Rule 51" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 03 2016 and Apr 18 2019: (Start) Conjecture: a(n) = 2*4^n - 2^n - 1 for odd n; a(n) = 2^n for even n. - _Karl V. Keller, Jr._, Oct 12 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010249 | OEIS-A266670

Binary representation of the n-th iteration of the "Rule 53" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 03 2016 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010250 | OEIS-A266671

Decimal representation of the n-th iteration of the "Rule 53" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 03 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010251 | OEIS-A266680

Binary representation of the middle column of the "Rule 175" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 19 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010252 | OEIS-A266685

T(n,k) is the number of loops appearing in pattern of circular arc connecting two vertices of regular polygons. (See Comments.)

T(n,k) is conjectured to be even rows of A109004 (excluding the first column).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jan 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010253 | OEIS-A266717

Binary representation of the n-th iteration of the "Rule 59" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 04 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010254 | OEIS-A266718

Decimal representation of the n-th iteration of the "Rule 59" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 04 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010255 | OEIS-A266723

Total number of ON (black) cells after n iterations of the "Rule 59" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 05 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010256 | OEIS-A266724

Number of OFF (white) cells in the n-th iteration of the "Rule 59" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 05 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010257 | OEIS-A266725

Total number of OFF (white) cells after n iterations of the "Rule 59" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 05 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010258 | OEIS-A266729

Numbers m, written in base 5, such that the Stanley sequence S_5(0,m) is well-behaved (or "modular").

It is not clear from Moy and Rolnick which of these terms are known to be correct and which are conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010259 | OEIS-A266734

Number of words on {1,1,2,2,3,3,...,n,n} avoiding the pattern 1234.

Conjecture: +3*n*(620202643096396011773 -608794959941727250938*n +1469492907122431180000*n^2) *(n+1)^2 *(2*n+1)^2 *a(n) -n*(943891175123956180605 44*n^6 -419724075420172456531120*n^5 +442263508538458916585360*n^4 +229131363207555256548194*n^3 -477880029525553894746823*n^2 +160086316440678171209939*n -11163647575735128211914) *a(n-1) -3*(n-1) *(238205220773229085875 84*n^6 -144630446008620178048037 6*n^5 +110804091174537748461455 40*n^4 -35494287160655892321199502*n^3 +57163416479212379649118767*n^2 -45988763994280198223305139*n +14778623468656583258390502) *a(n-2) +36*(n-2) *(419022927350372582170 56*n^6 -78325486543373387621947 2*n^5 +52359701363408117773325 52*n^4 -17094365117036393449118734*n^3 +29518557363755878023892305*n^2 -25895204716899392803468055*n +90757526337816081629440 50) *a(n-3) -8748*(n-2) *(125877543736438014048*n^2 -450267700517870762570*n +37094954161920926847 5) *(n-3)^2 *(2*n-7)^2 *a(n-4)=0. - _R. J. Mathar_, Apr 15 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010260 | OEIS-A266736

Number of words on {1,1,1,2,2,2,3,3,3,4,4,4,...,n,n,n} avoiding the pattern 123.

Conjecture: a(n) = (2/Pi)*Integral_{t=0..1} sqrt((1 - t)/t)*(64*t^3 - 32*t^2)^n = Catalan(3*n)*2F1(-1-3*n,-n;1/2-3*n;1/2). - _Benedict W. J. Irwin_, Oct 05 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010261 | OEIS-A266752

Binary representation of the n-th iteration of the "Rule 163" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 20 2016 and Apr 18 2019: (Start) Conjectured e.g.f.: (99 - 100*exp(x) + 991*exp(100*x))/990. - _Stefano Spezia_, Aug 05 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010262 | OEIS-A266753

Decimal representation of the n-th iteration of the "Rule 163" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 20 2016 and Apr 20 2019: (Start) a(n) = 4*a(n-1) + 2, n>1, conjectured. - _Yosu Yurramendi_, Jan 22 2017 a(n) = 2*A020988(n) - A020988(n-1) = A020988(n) + 2^(2n-1) for n > 0, conjectured. - _Yosu Yurramendi_, Jan 24 2017 [n range correction - _Karl V. Keller, Jr._, May 07 2022] a(n) = A072197(n-1) + A002450(n), n > 0, conjectured. - _Yosu

Yurramendi_, Mar 03 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010263 | OEIS-A266787

Binary representation of the n-th iteration of the "Rule 61" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 03 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010264 | OEIS-A266788

Decimal representation of the n-th iteration of the "Rule 61" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 03 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010265 | OEIS-A266790

Binary representation of the middle column of the "Rule 61" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 04 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010266 | OEIS-A266791

Decimal representation of the middle column of the "Rule 61" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 04 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010267 | OEIS-A266792

Number of ON (black) cells in the n-th iteration of the "Rule 61" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 04 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010268 | OEIS-A266793

Total number of ON (black) cells after n iterations of the "Rule 61" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 04 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010269 | OEIS-A266794

Number of OFF (white) cells in the n-th iteration of the "Rule 61" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 04 2016 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010270 | OEIS-A266795

Total number of OFF (white) cells after n iterations of the "Rule 61" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 04 2016 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010271 | OEIS-A266838

Binary representation of the n-th iteration of the "Rule 67" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 05 2016 and Apr 18 2019: (Start) Conjecture: a(n) = (10*100^n - 999909*1000^floor(n/2)/10000 - 1)/9 for odd n > 3; a(n) = 10111*1000^(n/2)/10000 for even n > 3. - _Karl V. Keller, Jr._, Dec 16 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010272 | OEIS-A266839

Decimal representation of the n-th iteration of the "Rule 67" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 05 2016 and Apr 18 2019: (Start) Conjecture: a(n) = 2*4^n - 61*8^floor(n/2)/16 - 1 for odd n > 2; a(n) = 23*8^(n/2)/16 for even n > 3. - _Karl V. Keller, Jr._, Dec 16 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010273 | OEIS-A266841

Binary representation of the n-th iteration of the "Rule 69" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 05 2016 and Apr 18 2019: (Start) Conjecture: a(n) = (10*100^n + 100^floor(n/2) - 11)/99 for odd n; a(n) = (100*100^n - 10^n)/99 for even n. - _Karl V. Keller, Jr._, Feb 28 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010274 | OEIS-A266842

Decimal representation of the n-th iteration of the "Rule 69" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 05 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010275 | OEIS-A266844

Binary representation of the n-th iteration of the "Rule 70" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 05 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010276 | OEIS-A266846

Decimal representation of the n-th iteration of the "Rule 70" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 05 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010277 | OEIS-A266849

Binary representation of the n-th iteration of the "Rule 71" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 05 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010278 | OEIS-A266850

Decimal representation of the n-th iteration of the "Rule 71" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 05 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010279 | OEIS-A266872

Binary representation of the n-th iteration of the "Rule 77" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 05 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010280 | OEIS-A266873

Decimal representation of the n-th iteration of the "Rule 77" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 05 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010281 | OEIS-A266883

Numbers of the form m*(4*m+1)+1, where m = 0,-1,1,-2,2,-3,3,...

Conjecture: the sequence terms are the exponents in the expansion of Sum_{n >= 1} q^n * (Product_{k >= 2*n-1} 1 - q^k) = q + q^4 + q^6 + q^15 + q^19 + q^34 + ... Cf. A174114. - _Peter Bala_, May 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bruno Berselli_, Jan 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010282 | OEIS-A266891

Expansion of Product_{k>=1} (1 + k*x^k)^k.

Conjecture: log(a(n)) ~ n^(2/3) * (2*log(3*n) - 3) / (4*3^(1/3)). - _Vaclav Kotesovec_, May 08 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jan 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010283 | OEIS-A266928

Number of steps needed to reach 1 or 2 where a step is x -> 3x+1 if x is odd, or x -> x/2p if x is even, 2p being the smallest semiprime dividing x.

Conjecture: all positive integers n not of the form 2^(2m+1) eventually reach 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010284 | OEIS-A266949

a(n) is defined by Product_{i>=1} (1-a(i)*x^i) = Sum_{i>=0} möbius(i+1)*x^i.

Conjecture: All coefficients are positive and strictly increasing starting from 5th term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gevorg Hmayakyan_, Jan 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010285 | OEIS-A266952

Least prime p such that p-2 and 6n-p and 6n+2-p are also prime, or 0 if no such prime exists.

Up to 10^5, the only indices for which a(n)=0 are {0, 1, 16, 67, 86, 131, 151, 186, 191, 211, 226, 541, 701}. I conjecture that this list is finite, and probably complete. Is it a coincidence that all odd numbers > 1 in this list are primes? (See also A144094.) This seems equivalent to a conjecture Zwillinger made in 1978, see reference in LINKS.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010286 | OEIS-A266953

Least prime p such that p-2 and 6n-p and either 6n+2-p or 6n+4-p is also prime, or 0 if no such prime exists.

Up to 10^5, the only indices for which a(n)=0 are {0, 1, 16, 86, 131, 151, 186, 191, 211, 226, 541, 701}. (Only 2 and 67 require the alternative primality of 6n+4-p and have thus A266952(n)=0.) I conjecture that this list is finite, and probably complete. Is it a coincidence that all odd numbers in this list are primes?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010287 | OEIS-A266968

Number of ordered ways to write n as x^5+y^4+z^3+w*(w+1)/2, where x, y, z and w are nonnegative integers with z > 0 and w > 0.

Conjecture: (i) a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 6, 7, 14, 21, 22, 26, 41, 51, 184, 189, 206, 225, 229, 526, 708. Compare this conjecture with the conjectures in A262813, A262827, A270559 and A271026.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010288 | OEIS-A266975

Binary representation of the n-th iteration of the "Rule 78" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 08 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010289 | OEIS-A266976

Decimal representation of the n-th iteration of the "Rule 78" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 08 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010290 | OEIS-A266979

Binary representation of the n-th iteration of the "Rule 79" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 08 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010291 | OEIS-A266980

Decimal representation of the n-th iteration of the "Rule 79" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 08 2016 and Apr 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010292 | OEIS-A266981

Number of ON (black) cells in the n-th iteration of the "Rule 79" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 08 2016 and Apr 19 2019: (Start) Conjecture from _Ctibor O. Zizka_, Mar 11 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010293 | OEIS-A266982

Triangle read by rows giving successive states of cellular automaton generated by "Rule 81" initiated with a single ON (black) cell.

Conjecture: a(0)=1; for n>0, a(n) = (Sum_{i=1..n-2} floor((n-2)/i)) mod 2. - _Wesley Ivan Hurt_, Mar 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010294 | OEIS-A266984

Decimal representation of the n-th iteration of the "Rule 81" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 08 2016 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010295 | OEIS-A266988

The indices of primes for which the average of the primitive roots is < p/2.

These primes are all of the form p==3 (mod 4). (conjecture)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimitri Papadopoulos_, Jan 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010296 | OEIS-A266998

Number of iterations of the map "n -> sum of the triangular numbers whose indices are the digits of n" needed to reach 1.

Conjecture: 1 is reachable for every n. Verified for n up to 10^6. - _Ivan N. Ianakiev_, Jan 10 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ivan N. Ianakiev_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010297 | OEIS-A267002

Binary representation of the n-th iteration of the "Rule 83" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 08 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010298 | OEIS-A267003

Decimal representation of the n-th iteration of the "Rule 83" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 08 2016 and Apr 19 2019: (Start) Conjectured e.g.f.: cosh(2*sqrt(2)*x) - sinh(x) + 2*sinh(4*x) - sinh(2*sqrt(2)*x)/sqrt(2). - _Stefano Spezia_, Mar 16 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010299 | OEIS-A267028

P(n,k) is an array read by rows, with n > 0 and k=1..5, where row n gives the chain of 5 consecutive primes {p(i), p(i+1), p(i+2), p(i+3), p(i+4)} having the symmetrical property p(i) + p(i+4) = p(i+1) + p(i+3) = 2*p(i+2) for some index i.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Feb 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010300 | OEIS-A267035

Binary representation of the n-th iteration of the "Rule 85" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 11 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010301 | OEIS-A267036

Decimal representation of the n-th iteration of the "Rule 85" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 11 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010302 | OEIS-A267041

Binary representation of the n-th iteration of the "Rule 91" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 10 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010303 | OEIS-A267042

Decimal representation of the n-th iteration of the "Rule 91" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 10 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010304 | OEIS-A267044

Binary representation of the middle column of the "Rule 91" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 10 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010305 | OEIS-A267045

Decimal representation of the middle column of the "Rule 91" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 10 2016 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010306 | OEIS-A267046

Number of ON (black) cells in the n-th iteration of the "Rule 91" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 10 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010307 | OEIS-A267047

Total number of ON (black) cells after n iterations of the "Rule 91" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 10 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010308 | OEIS-A267048

Number of OFF (white) cells in the n-th iteration of the "Rule 91" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 11 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010309 | OEIS-A267049

Total number of OFF (white) cells after n iterations of the "Rule 91" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 11 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010310 | OEIS-A267051

Binary representation of the n-th iteration of the "Rule 92" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 10 2016 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010311 | OEIS-A267052

Decimal representation of the n-th iteration of the "Rule 92" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 10 2016 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010312 | OEIS-A267054

Binary representation of the n-th iteration of the "Rule 93" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 10 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010313 | OEIS-A267055

Decimal representation of the n-th iteration of the "Rule 93" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 10 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010314 | OEIS-A267057

Binary representation of the n-th iteration of the "Rule 97" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 10 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010315 | OEIS-A267058

Decimal representation of the n-th iteration of the "Rule 97" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 10 2016 and Apr 19 2019: (Start) Conjecture: a(n) = 256^floor(n/4)*(((n-1)/2 + 1) mod 2) + 20*256^floor(n/4)*(((n-1)/2) mod 2) - 1 for odd n; a(n) = 256^floor(n/4)*((n/2 + 1) mod 2) + 20*256^floor(n/4)*((n/2) mod 2) for even n. - _Karl V. Keller, Jr._, Jul 03 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010316 | OEIS-A267068

a(n) = (n+1) / A189733(n).

Conjectures from _Colin Barker_, Jan 10 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Jan 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010317 | OEIS-A267084

a(n) = ceiling(A007504(n)/n) - floor(A007504(n)/n); a(n) is 0 if n divides the sum of first n primes, 1 otherwise.

It is conjectured that there are infinitely many zeros, but that their density is zero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Jan 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010318 | OEIS-A267089

T(n,k) is decimal conversion of 1's in an n X n table that lie on its principal diagonals.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jan 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010319 | OEIS-A267090

Triangle read by rows: Fill an n X n square with 1's, except for 0's on the two main diagonals. Then T(n,k) is decimal equivalent of the k-th row (0<=k<=n).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jan 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010320 | OEIS-A267120

Triangle of coefficients of Gaussian polynomials [2n+3,3]_q represented as finite sum of terms (1+q^2)^k*q^(g-k), where k = 0,1,...,g with g=3n.

Conjecture: More generally, the Gaussian polynomial [2*n+m+1-(m mod 2),m]_q = Sum_{k=0..g(m;n)} a(m;n,k)*(1+q^2)^k*q^(g(m;n)-k), for m >= 0, n >= 0, where g(m;n) = m*n if m is odd and (2*n+1)*m/2 if m is even, and the tabf array entries a(m;n,k) are the coefficients of the g.f. for the row n polynomials G(m;n,x) = (d^m/dt^m)G(m;n,t,x)/m!|_{t=0}, with G(m;n,t,x) = (1+t)*Product_{k=1..n+(m - m (mod 2))/2}(1 + t^2 + 2*t*T(k,x/2) (Chebyshev's T-polynomials). Hence a(m;n,k) = [x^k]G(m;n,x), for k=0..g(m;n). The present entry is the instance m = 3. (Thanks to _Wolfdieter Lang_ for clarifying the text on the general prescription of a(m;n,k).)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stephen O'Sullivan_, Jan 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010321 | OEIS-A267121

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with x*y*z*(x+9*y+11*z+10*w) a square, where x is a positive integer and y,z,w are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 4^k*m (k = 0,1,2,... and m = 1, 7, 15, 39, 119, 127, 159, 239, 359, 391, 527, 543, 863, 5791).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010322 | OEIS-A267124

Primitive practical numbers: practical numbers that are squarefree or practical numbers that when divided by any of its prime factors whose factorization exponent is greater than 1 is no longer practical.

Conjecture: every odd number, beginning with 3, is the sum of a prime number and a primitive practical number. This is a tighter conjecture to the conjecture posed by _Hal M. Switkay_ (see A005153).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Jan 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010323 | OEIS-A267138

Binary representation of the n-th iteration of the "Rule 103" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 11 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010324 | OEIS-A267139

Decimal representation of the n-th iteration of the "Rule 103" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 11 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010325 | OEIS-A267153

Binary representation of the n-th iteration of the "Rule 107" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 11 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010326 | OEIS-A267154

Decimal representation of the n-th iteration of the "Rule 107" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 11 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010327 | OEIS-A267156

Binary representation of the middle column of the "Rule 107" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 11 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010328 | OEIS-A267157

Decimal representation of the middle column of the "Rule 107" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 11 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010329 | OEIS-A267158

Number of ON (black) cells in the n-th iteration of the "Rule 107" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 11 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010330 | OEIS-A267159

Total number of ON (black) cells after n iterations of the "Rule 107" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 11 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010331 | OEIS-A267160

Number of OFF (white) cells in the n-th iteration of the "Rule 107" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 11 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010332 | OEIS-A267161

Total number of OFF (white) cells after n iterations of the "Rule 107" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 12 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010333 | OEIS-A267182

Row 2 of the square array in A267181.

Colin Barker's conjectures below are true. Proof: A267181(ka,kb) = A267181(a,b) since both operations preserve the greatest common factor of the two coordinates, so A267181(2k,2) = A267181(k,1) = k for k > 1, the second conjecture. For odd coordinates, we have the forced chain (2k+1,2) -> (2,2k+1) -> (2,2k-1) -> ... -> (2,1) -> (1,2) -> (1,1) with k+3 operations, the third conjecture. The rest follow from combining these. (End) Conjectures from _Colin Barker_, Jan 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010334 | OEIS-A267183

Row 3 of the square array in A267181.

Conjectures from _Colin Barker_, Jan 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010335 | OEIS-A267184

Row 4 of the square array in A267181.

Conjectures from _Colin Barker_, Jan 29 2016: (Start) Conjectured e.g.f.: (8 - 5*cos(x) + (1 + x)*cosh(x) - sin(x) + (16 + x)*sinh(x))/4. - _Stefano Spezia_, Nov 29 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010336 | OEIS-A267185

Column 2 of the square array in A267181.

Conjectures from _Colin Barker_, Jan 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010337 | OEIS-A267186

Column 3 of the square array in A267181.

Conjectures from _Colin Barker_, Jan 30 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010338 | OEIS-A267187

Column 4 of the square array in A267181.

Conjectures from _Colin Barker_, Jan 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010339 | OEIS-A267192

Column 3 of triangle in A059317 (the Pascal "Rhombus").

Conjecture: +(n-2)*(n-3)*(n+3)*a(n) -n*(2*n-1)*(n-2)*a(n-1) -(n-1)*(5*n^2-10*n+18)*a(n-2) +n*(2*n-3)*(n-2)*a(n-3) +n*(n+1)*(n-5)*a(n-4)=0. - _R. J. Mathar_, Jul 23 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010340 | OEIS-A267209

Binary representation of the middle column of the "Rule 109" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 13 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010341 | OEIS-A267210

Decimal representation of the middle column of the "Rule 109" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 13 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010342 | OEIS-A267211

Number of ON (black) cells in the n-th iteration of the "Rule 109" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 13 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010343 | OEIS-A267212

Total number of ON (black) cells after n iterations of the "Rule 109" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 13 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010344 | OEIS-A267213

Number of OFF (white) cells in the n-th iteration of the "Rule 109" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 13 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010345 | OEIS-A267214

Total number of OFF (white) cells after n iterations of the "Rule 109" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 13 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010346 | OEIS-A267219

Expansion of exp(Sum_{n >= 1} A002895(n)*x^n/n).

For integer m, define a sequence {u_m(n) : n >= 0} by u_m(n) = [x^n] A(x)^(m*n/4). Conjecture: the supercongruences u_m(n*p^r) == u_m(n*p^(r-1)) (mod p^(2*r)) hold for all primes p >= 5 and positive integers n and r. - _Peter Bala_, Oct 17 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010347 | OEIS-A267226

Number of length-n 0..2 arrays with no following elements greater than or equal to the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010348 | OEIS-A267227

Number of length-n 0..3 arrays with no following elements greater than or equal to the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010349 | OEIS-A267228

Number of length-n 0..4 arrays with no following elements greater than or equal to the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010350 | OEIS-A267229

Number of length-n 0..5 arrays with no following elements greater than or equal to the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010351 | OEIS-A267230

Number of length-n 0..6 arrays with no following elements greater than or equal to the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010352 | OEIS-A267231

Number of length-n 0..7 arrays with no following elements greater than or equal to the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010353 | OEIS-A267233

Number of length-4 0..n arrays with no following elements greater than or equal to the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010354 | OEIS-A267234

Number of length-5 0..n arrays with no following elements greater than or equal to the first repeated value.

Conjectures from _Colin Barker_, Jan 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010355 | OEIS-A267235

Number of length-6 0..n arrays with no following elements greater than or equal to the first repeated value.

Conjectures from _Colin Barker_, Jan 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010356 | OEIS-A267236

Number of length-7 0..n arrays with no following elements greater than or equal to the first repeated value.

Conjectures from _Colin Barker_, Jan 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010357 | OEIS-A267240

Number of n X 3 binary arrays with row sums nondecreasing and columns lexicographically nondecreasing.

Conjectures from _Colin Barker_, Jan 11 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010358 | OEIS-A267257

Binary representation of the middle column of the "Rule 111" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 13 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010359 | OEIS-A267258

Decimal representation of the middle column of the "Rule 111" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 13 2016 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010360 | OEIS-A267259

Number of ON (black) cells in the n-th iteration of the "Rule 111" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 13 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010361 | OEIS-A267260

Total number of ON (black) cells after n iterations of the "Rule 111" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 13 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010362 | OEIS-A267261

Number of OFF (white) cells in the n-th iteration of the "Rule 111" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 14 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010363 | OEIS-A267262

Total number of OFF (white) cells after n iterations of the "Rule 111" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 14 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010364 | OEIS-A267270

Binary representation of the n-th iteration of the "Rule 115" elementary cellular automaton starting with a single ON (black) cell.

Conjecture: a(n) = (10^(2*n + 1) - 9*10^(3*n/2 - 1/2) - 1)/9 for odd n; a(n) = 1100*10^(3*n/2 - 3) for even n > 0. - _Karl V. Keller, Jr._, Apr 02 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010365 | OEIS-A267271

Decimal representation of the n-th iteration of the "Rule 115" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 14 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010366 | OEIS-A267274

Decimal representation of the n-th iteration of the "Rule 117" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 14 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010367 | OEIS-A267293

Binary representation of the n-th iteration of the "Rule 121" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 15 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010368 | OEIS-A267294

Decimal representation of the n-th iteration of the "Rule 121" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 15 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010369 | OEIS-A267297

Square triangular numbers that are the sum of 2 nonzero nonconsecutive triangular numbers.

Subsequence of A001110. Conjecture: except for the terms 0, 1, 36, 48024900, the sequence is identical to A001110. Conjectures from _Colin Barker_, Mar 03 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Feb 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010370 | OEIS-A267350

Binary representation of the n-th iteration of the "Rule 123" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 14 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010371 | OEIS-A267351

Decimal representation of the n-th iteration of the "Rule 123" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 14 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010372 | OEIS-A267352

Number of ON (black) cells in the n-th iteration of the "Rule 123" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 14 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010373 | OEIS-A267353

Total number of ON (black) cells after n iterations of the "Rule 123" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 14 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010374 | OEIS-A267354

Number of OFF (white) cells in the n-th iteration of the "Rule 123" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 14 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010375 | OEIS-A267359

Binary representation of the n-th iteration of the "Rule 125" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 14 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010376 | OEIS-A267360

Decimal representation of the n-th iteration of the "Rule 125" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 14 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010377 | OEIS-A267383

Number A(n,k) of acyclic orientations of the Turán graph T(n,k); square array A(n,k), n>=0, k>=1, read by antidiagonals.

Conjecture: In general, column k > 1 is asymptotic to n! / ((k-1) * (1 - log(k/(k-1)))^((k-1)/2) * k^n * (log(k/(k-1)))^(n+1)). - _Vaclav Kotesovec_, Feb 18 2017 [Proved by Zhiyang Sun, 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jan 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010378 | OEIS-A267414

Integers k such that there exist nonnegative integers x,y,z with k! = x^3 + y^3 + z^3.

Conjecture I: The natural density of this sequence is 1. Conjecture II: All integers > 13 are terms. The decomposition is not necessarily unique; for instance, 12! = 35^3 + 309^3 + 766^3 = 240^3 + 504^3 + 696^3. Deshouillers, Hennecart, & Landreau conjecture (the DHL conjecture) that the sequence of numbers that are a sum of at most three cubes has density 0.0999425... (see links). So roughly, if the DHL conjecture holds and if that density can be used as a probability that holds independently for candidates then we have the probability that 34! is the sum of at most 3 cubes to be 1 - (1-0.0999425)^660 ~= 1 - 6.6*10^-31. For larger k this probability doesn't tend to decrease. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Jan 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010379 | OEIS-A267435

Numbers n such that each reduced Collatz trajectory (mod p): (n, T(n), T(T(n)), ..., 4, 2, 1) / pZ, where the odd prime p is the number of iterations needed to reach 1, contains exactly the p-1 values {1, 2, 3, ..., p-1}.

This sequence provides a link with Artin's conjecture on primitive roots. Conjecture: the sequence is infinite (corollary of Artin's conjecture because the sequence contains the numbers 2^A001122(k) where A001122 are the primes with primitive root 2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010380 | OEIS-A267442

Middle column of the "Rule 129" elementary cellular automaton starting with a single ON (black) cell.

Conjecture: a(n) = A043545(n), n>0. - _R. J. Mathar_, Jan 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010381 | OEIS-A267445

Number of ON (black) cells in the n-th iteration of the "Rule 129" elementary cellular automaton starting with a single ON (black) cell.

Conjecture: a(n) = A071050(n), n>0. - _R. J. Mathar_, Jan 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010382 | OEIS-A267447

Number of OFF (white) cells in the n-th iteration of the "Rule 129" elementary cellular automaton starting with a single ON (black) cell.

Conjecture: a(n) = A071051(n), n>0. - _R. J. Mathar_, Jan 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010383 | OEIS-A267451

Number of ON (black) cells in the n-th iteration of the "Rule 131" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 15 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010384 | OEIS-A267452

Total number of ON (black) cells after n iterations of the "Rule 131" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 15 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010385 | OEIS-A267453

Number of OFF (white) cells in the n-th iteration of the "Rule 131" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 15 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010386 | OEIS-A267454

Total number of OFF (white) cells after n iterations of the "Rule 131" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 15 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010387 | OEIS-A267456

Binary representation of the n-th iteration of the "Rule 133" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 15 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010388 | OEIS-A267457

Decimal representation of the n-th iteration of the "Rule 133" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 15 2016 and Apr 19 2019: (Start) Conjectures from _Stefano Spezia_, Mar 27 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010389 | OEIS-A267458

Number of ON (black) cells in the n-th iteration of the "Rule 133" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 15 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010390 | OEIS-A267459

Total number of ON (black) cells after n iterations of the "Rule 133" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 16 2016 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010391 | OEIS-A267460

Number of OFF (white) cells in the n-th iteration of the "Rule 133" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 16 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010392 | OEIS-A267461

Total number of OFF (white) cells after n iterations of the "Rule 133" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 16 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010393 | OEIS-A267465

Number of length-n 0..2 arrays with no following elements larger than the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010394 | OEIS-A267466

Number of length-n 0..3 arrays with no following elements larger than the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010395 | OEIS-A267467

Number of length-n 0..4 arrays with no following elements larger than the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010396 | OEIS-A267468

Number of length-n 0..5 arrays with no following elements larger than the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010397 | OEIS-A267469

Number of length-n 0..6 arrays with no following elements larger than the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010398 | OEIS-A267470

Number of length-n 0..7 arrays with no following elements larger than the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010399 | OEIS-A267472

Number of length-4 0..n arrays with no following elements larger than the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010400 | OEIS-A267473

Number of length-5 0..n arrays with no following elements larger than the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010401 | OEIS-A267474

Number of length-6 0..n arrays with no following elements larger than the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010402 | OEIS-A267475

Number of length-7 0..n arrays with no following elements larger than the first repeated value.

Conjectures from _Colin Barker_, Feb 05 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010403 | OEIS-A267483

Triangle of coefficients of Gaussian polynomials [2n+3,2]_q represented as finite sum of terms (1+q^2)^k*q^(g-k), where k = 0,1,...,g with g=2n+1.

Conjecture: More generally, the Gaussian polynomial [2*n+m+1-(m mod 2),m]_q = Sum_{k=0..g(m;n)} a(m;n,k)*(1+q^2)^k*q^(g(m;n)-k), for m >= 0, n >= 0, where g(m;n) = m*n if m is odd and (2*n+1)*m/2 if m is even, and the tabf array entries a(m;n,k) are the coefficients of the g.f. for the row n polynomials G(m;n,x) = (d^m/dt^m)G(m;n,t,x)/m!|_{t=0}, with G(m;n,t,x) = (1+t)*Product_{k=1..n+(m - m (mod 2))/2}(1 + t^2 + 2*t*T(k,x/2) (Chebyshev's T-polynomials). Hence a(m;n,k) = [x^k]G(m;n,x), for k=0..g(m;n). The present entry is the instance m = 2. (Thanks to _Wolfdieter Lang_ for clarifying the text on the general prescription of a(m;n,k).)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stephen O'Sullivan_, Jan 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010404 | OEIS-A267489

a(n) = n^2 - 4*floor(n^2/6).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jan 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010405 | OEIS-A267514

Binary representation of the middle column of the "Rule 137" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 16 2016: (Start) This conjectured linear recurrence fails to predict the correct values from a(62) on. - _R. J. Mathar_, Apr 12 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010406 | OEIS-A267515

Decimal representation of the middle column of the "Rule 137" elementary cellular automaton starting with a single ON (black) cell.

[Incorrect] conjectures from _Colin Barker_, Jan 16 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010407 | OEIS-A267524

Binary representation of the middle column of the "Rule 139" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 16 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010408 | OEIS-A267526

Binary representation of the n-th iteration of the "Rule 141" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 16 2016: (Start) Conjecture: a(n) = 10^n*(100^floor(n/2) - 1)/99 + (10^(n-1) - 1)/9 for n > 1. - _Karl V. Keller, Jr._, Sep 22 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010409 | OEIS-A267528

Number of ON (black) cells in the n-th iteration of the "Rule 141" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 16 2016 and Apr 17 2019: (Start) Conjectured e.g.f.: 2 + x + (3*x/2 - 1)*cosh(x) + 3*(x - 1)*sinh(x)/2. - _Stefano Spezia_, Feb 20 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010410 | OEIS-A267529

Total number of ON (black) cells after n iterations of the "Rule 141" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 16 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010411 | OEIS-A267530

Number of OFF (white) cells in the n-th iteration of the "Rule 141" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 17 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010412 | OEIS-A267531

Total number of OFF (white) cells after n iterations of the "Rule 141" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 17 2016: (Start) Conjecture: a(n) = a(n-1) + A004526(n+5) for n > 1. - _J. Stauduhar_, Oct 21 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010413 | OEIS-A267534

Indices of Lucas numbers that are not of the form x^2 + y^2 + z^2 where x, y and z are integers.

Conjectures from _Colin Barker_, Jan 30 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Jan 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010414 | OEIS-A267535

Binary representation of the n-th iteration of the "Rule 143" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 17 2016 and Apr 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010415 | OEIS-A267536

Decimal representation of the n-th iteration of the "Rule 143" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 17 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010416 | OEIS-A267538

Binary representation of the middle column of the "Rule 143" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 17 2016 and Apr 20 2019: (Start) Conjecture: a(n) = floor(9901*10^n/9000). - _Karl V. Keller, Jr._, Apr 24 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010417 | OEIS-A267539

Decimal representation of the middle column of the "Rule 143" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 17 2016 and Apr 20 2019: (Start) {1,3,6} followed by A198274 (conjectured). - _Robert Price_, Jan 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010418 | OEIS-A267588

Binary representation of the middle column of the "Rule 169" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 18 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010419 | OEIS-A267589

Decimal representation of the middle column of the "Rule 169" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 18 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010420 | OEIS-A267595

Binary representation of the n-th iteration of the "Rule 173" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 18 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010421 | OEIS-A267596

Decimal representation of the n-th iteration of the "Rule 173" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 18 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010422 | OEIS-A267598

Triangle read by rows giving successive states of cellular automaton generated by "Rule 177" initiated with a single ON (black) cell.

a(n) = (1 - (-1)^(n + floor(sqrt(n + 2))))/2 (conjecture). - _Velin Yanev_, Nov 15 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010423 | OEIS-A267599

Binary representation of the n-th iteration of the "Rule 177" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 18 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010424 | OEIS-A267604

Decimal representation of the middle column of the "Rule 175" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 18 2016 and Apr 20 2019: (Start) Conjectures from _Altug Alkan_, Jan 18 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010425 | OEIS-A267610

Total number of OFF (white) cells after n iterations of the "Rule 182" elementary cellular automaton starting with a single ON (black) cell.

Conjecture: a(n) = A267700(n) - n. - _Gus Wiseman_, Mar 30 2019 G.f.: (1/x)*(A(x)/x - (x+1)/(1-x)^2) where A(x) is the g.f. for A006046 (conjectured). - _John Tyler Rascoe_, Jul 08 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010426 | OEIS-A267613

Binary representation of the n-th iteration of the "Rule 185" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 18 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010427 | OEIS-A267614

Decimal representation of the n-th iteration of the "Rule 185" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 18 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010428 | OEIS-A267622

Binary representation of the n-th iteration of the "Rule 187" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 19 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010429 | OEIS-A267623

Binary representation of the middle column of the "Rule 187" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 19 2016 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010430 | OEIS-A267632

Triangle T(n, k) read by rows: the k-th column of the n-th row lists the number of ways to select k distinct numbers (k >= 1) from [1..n] so that their sum is divisible by n.

Row less the last element is palindrome for n=odd or n=power of 2 where n is the row number (observation-conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimitri Papadopoulos_, Jan 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010431 | OEIS-A267677

Binary representation of the n-th iteration of the "Rule 197" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 19 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010432 | OEIS-A267678

Decimal representation of the n-th iteration of the "Rule 197" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 19 2016: (Start) Conjecture: a(n) = 2*(-3 + (-1)^n - 2^(2+n) + 3*4^n)/3 for n>1. - _Colin Barker_, Feb 14 2017 Conjecture: a(n) = floor((6*4^n - 8*2^n)/3) - 2^(n mod 2) for n > 1. - _Karl V. Keller, Jr._, Jun 01 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010433 | OEIS-A267680

Binary representation of the n-th iteration of the "Rule 201" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 19 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010434 | OEIS-A267681

Decimal representation of the n-th iteration of the "Rule 201" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 19 2016: (Start) Conjecture: a(n) = 2*4^n - (n%2*2 + [n]*5)*2^(n-1) - 1, where [n] = 1 iff n > 0; n%2 = 1 iff n is odd. - _M. F. Hasler_, Jul 28 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010435 | OEIS-A267684

Binary representation of the n-th iteration of the "Rule 203" elementary cellular automaton starting with a single ON (black) cell.

The above conjectures are correct. Also a(n) = (10*100^n - 9*10^n - 1)/9 for n > 1. - _Karl V. Keller, Jr._, Jun 07 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010436 | OEIS-A267685

Decimal representation of the n-th iteration of the "Rule 203" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from Barker confirmed by later formulas. - _Ray Chandler_, Aug 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010437 | OEIS-A267688

Binary representation of the n-th iteration of the "Rule 199" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 19 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010438 | OEIS-A267689

Decimal representation of the n-th iteration of the "Rule 199" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 19 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010439 | OEIS-A267700

"Tree" sequence in a 90-degree sector of the cellular automaton of A160720.

Conjecture: this is also the "tree" sequence in a 120-degree sector of the cellular automaton of A266532. Conjecture: a(n) is also the total number of Y-toothpicks after n-th stage in the tree that arises from one of the three spokes in a 120-degree sector of the cellular automaton of A266532 on the triangular grid. Comment from _N. J. A. Sloane_, Jan 23 2016: All the above conjectures are true! Conjecture 1: a(n) = (A266532(n+1) - 1)/3. Conjecture 2: a(n) = A160720(n+1) - A266532(n+1). All of the above conjectures are true. - _N. J. A. Sloane_, Jan 23 2016 (Conjecture) a(n) = A267610(n) + n. - _Gus Wiseman_, Mar 31 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jan 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010440 | OEIS-A267703

Conjectured list of numbers whose trajectory under the '7x+1' map eventually reaches 1.

Conjectured list of numbers whose trajectory under the '7x+1' map eventually reaches 1. This is conjectural in that there is no known proof that the missing numbers 3, 6, 7, ... are really missing. It may be that after a very large number of iterations they will cycle. - _N. J. A. Sloane_, Jan 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010441 | OEIS-A267705

Binary representation of the n-th iteration of the "Rule 205" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 20 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010442 | OEIS-A267708

Triangle read by rows giving successive states of cellular automaton generated by "Rule 206" initiated with a single ON (black) cell.

Conjecture: a(n) = (1 + (-1)^floor(2*sqrt(n)))/2. - _Velin Yanev_, Nov 30 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010443 | OEIS-A267774

Decimal representation of the n-th iteration of the "Rule 207" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 20 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010444 | OEIS-A267775

Binary representation of the n-th iteration of the "Rule 207" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 20 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010445 | OEIS-A267777

Binary representation of the n-th iteration of the "Rule 209" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 20 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010446 | OEIS-A267779

Binary representation of the n-th iteration of the "Rule 211" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 20 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010447 | OEIS-A267780

Decimal representation of the n-th iteration of the "Rule 211" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 20 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010448 | OEIS-A267794

a(n) = n-2-i, where i (0 <= i < n-2) is the largest number such that [a(n-2)-a(i); a(n-1)-a(i+1)] = [0; 0], or a(n) = 0 if no such i exists.

Maximal number of successive zeros in A181391 is 1, in this sequence 2. Conjecture : maximal number of successive zeros is k: a(0); ...; a(j)>0; a(j+1)=0; ...; a(j+k)=0; a(j+k+1)>0; ... The conjecture is true since (k+1)-th zero can occur only on the first occurrence of k successive zeros, by definition of such sequences. - _Andrey Zabolotskiy_, Oct 31 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Jan 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010449 | OEIS-A267801

Binary representation of the n-th iteration of the "Rule 213" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 21 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010450 | OEIS-A267802

Decimal representation of the n-th iteration of the "Rule 213" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 21 2016: (Start) Conjecture: a(n) = 2^(2*n+1) - 13 for n>1. - _Colin Barker_, Nov 25 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010451 | OEIS-A267804

Binary representation of the n-th iteration of the "Rule 214" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010452 | OEIS-A267805

Decimal representation of the n-th iteration of the "Rule 214" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010453 | OEIS-A267811

Binary representation of the n-th iteration of the "Rule 217" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from Barker confirmed by later formula. - _Ray Chandler_, Aug 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010454 | OEIS-A267812

Decimal representation of the n-th iteration of the "Rule 217" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010455 | OEIS-A267815

Binary representation of the n-th iteration of the "Rule 221" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010456 | OEIS-A267816

Decimal representation of the n-th iteration of the "Rule 221" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010457 | OEIS-A267824

Composite numbers n such that binomial(2n-1, n-1) == 1 (mod n^2).

Conjecture: n is a term if and only if n = A088164(i)^2 for some i >= 1 (cf. McIntosh, 1995, p. 385). - _Felix Fröhlich_, Jan 27 2016 The "if" part of the conjecture is true: see the McIntosh reference. - _Jonathan Sondow_, Jan 28 2016 The above conjecture implies that this sequence and A228562 are disjoint. - _Felix Fröhlich_, Jan 27 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Jan 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010458 | OEIS-A267826

Numbers not of the form w^3 + 2*x^3 + 3*y^3 + 4*z^3, where w, x, y and z are nonnegative integers.

Conjecture: The sequence has exactly 122 terms the last of which is a(122) = 41405. The conjecture implies that {P(v)+w^3+2*x^3+3*y^3+4*z^3: w,x,y,z = 0,1,2,...} = {0,1,2,...} whenever P(v) is among the polynomials a*v^3 (a = 1,5,6,7,9,10,12,15,18), b*v^4 (b = 1,2,3,5,6,12,18), c*v^5 (c = 1,2,5,12) and d*v^k (d = 5,12; k = 6,7). Moreover, it also implies that {8*t+w^3+2*x^3+3*y^3+4*z^3: t = 0,1; w,x,y,z = 0,1,2,...} = {0,1,2,...}. If a,b,c,d and m are positive integers with {m*t+a*w^3+b*x^3+c*y^3+d*z^3: t = 0,1; w,x,y,z = 0,1,2,...} = {0,1,2,...}, then we must have m = 8 and {a,b,c,d} = {1,2,3,4}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010459 | OEIS-A267840

Number of symmetric difference-closed 4-sets consisting of the empty set and sets consisting of pairwise disjoint 2-subsets of {1,2,...,n}.

Conjectures from _Vaclav Kotesovec_, Apr 10 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John M. Campbell_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010460 | OEIS-A267846

Binary representation of the n-th iteration of the "Rule 227" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 21 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010461 | OEIS-A267847

Decimal representation of the n-th iteration of the "Rule 227" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 21 2016: (Start) Conjecture: a(n) = 2*(4^n - 3) for n>1. - _Colin Barker_, Nov 26 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010462 | OEIS-A267850

Binary representation of the n-th iteration of the "Rule 229" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 21 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010463 | OEIS-A267851

Decimal representation of the n-th iteration of the "Rule 229" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 21 2016: (Start) Conjecture: a(n) = 2*(4^n - 1) = A002446(n) for n>1. - _Colin Barker_, Nov 26 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010464 | OEIS-A267854

Binary representation of the n-th iteration of the "Rule 230" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 21 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010465 | OEIS-A267855

Decimal representation of the n-th iteration of the "Rule 230" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 21 2016 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010466 | OEIS-A267861

Number of ways to write n as 2*t + u^4 + v^4 + 2*w^4 + 3*x^4 + 4*y^4 + 6*z^4, where t is 0 or 1, and u, v, w, x, y, z are nonnegative integers with u <= v and v > 0.

Conjecture: a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 1, 2, 111, 127, 143, 158, 221, 223, 240, 460, 463, 480, 545, 560, 561, 1455, 1695, 1776, 2175. Moreover, any integer n > 10^4 not among 10543, 17935, 37583, 40383, 78543 can be written as u^4 + v^4 + 2*w^4 + 3*x^4 + 4*y^4 + 6*z^4 with u,v,w,x,y,z nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010467 | OEIS-A267867

Binary representation of the n-th iteration of the "Rule 231" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 21 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010468 | OEIS-A267872

Number of ON (black) cells in the n-th iteration of the "Rule 237" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010469 | OEIS-A267873

Number of ON (black) cells in the n-th iteration of the "Rule 235" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010470 | OEIS-A267876

Binary representation of the n-th iteration of the "Rule 233" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010471 | OEIS-A267877

Decimal representation of the n-th iteration of the "Rule 233" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010472 | OEIS-A267879

Binary representation of the middle column of the "Rule 233" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 20 2019: (Start) Conjecture: a(n) = (9091*10^(n-3) - 1)/9 for n > 2. - _Stefano Spezia_, Dec 25 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010473 | OEIS-A267880

Decimal representation of the middle column of the "Rule 233" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010474 | OEIS-A267881

Number of ON (black) cells in the n-th iteration of the "Rule 233" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010475 | OEIS-A267882

Total number of ON (black) cells after n iterations of the "Rule 233" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 20 2019: (Start) Conjectured e.g.f.: exp(x)*(x^2 + 3*x - 5) + x^2/2 + 3*x + 6. - _Stefano Spezia_, Sep 08 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010476 | OEIS-A267884

Total number of OFF (white) cells after n iterations of the "Rule 233" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 24 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010477 | OEIS-A267885

Binary representation of the n-th iteration of the "Rule 235" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010478 | OEIS-A267886

Decimal representation of the n-th iteration of the "Rule 235" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010479 | OEIS-A267887

Binary representation of the n-th iteration of the "Rule 237" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010480 | OEIS-A267888

Decimal representation of the n-th iteration of the "Rule 237" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010481 | OEIS-A267889

Binary representation of the n-th iteration of the "Rule 239" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010482 | OEIS-A267890

Decimal representation of the n-th iteration of the "Rule 239" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010483 | OEIS-A267905

Number of n X 1 0..2 arrays with every repeated value in every row unequal to the previous repeated value, and in every column equal to the previous repeated value, and new values introduced in row-major sequential order.

Conjectures from _Colin Barker_, Jan 11 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010484 | OEIS-A267912

Number of 1 X n 0..2 arrays with every repeated value in every row unequal to the previous repeated value, and in every column equal to the previous repeated value, and new values introduced in row-major sequential order.

Conjectures from _Colin Barker_, Feb 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010485 | OEIS-A267920

Binary representation of the n-th iteration of the "Rule 243" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010486 | OEIS-A267921

Decimal representation of the n-th iteration of the "Rule 243" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 22 2016 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010487 | OEIS-A267923

Binary representation of the n-th iteration of the "Rule 245" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 23 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010488 | OEIS-A267924

Decimal representation of the n-th iteration of the "Rule 245" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 23 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010489 | OEIS-A267925

Binary representation of the n-th iteration of the "Rule 246" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 23 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010490 | OEIS-A267926

Decimal representation of the n-th iteration of the "Rule 246" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 23 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010491 | OEIS-A267934

Binary representation of the n-th iteration of the "Rule 249" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 23 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010492 | OEIS-A267935

Decimal representation of the n-th iteration of the "Rule 249" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 23 2016 and Apr 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010493 | OEIS-A267937

Binary representation of the n-th iteration of the "Rule 251" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 23 2016 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010494 | OEIS-A267938

Decimal representation of the n-th iteration of the "Rule 251" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 23 2016 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010495 | OEIS-A267940

Binary representation of the n-th iteration of the "Rule 253" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 23 2016 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010496 | OEIS-A267941

Decimal representation of the n-th iteration of the "Rule 253" elementary cellular automaton starting with a single ON (black) cell.

Conjectures from _Colin Barker_, Jan 23 2016 and Apr 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010497 | OEIS-A267946

Number of n X 1 0..2 arrays with every repeated value in every row and column one larger mod 3 than the previous repeated value, and upper left element zero.

Conjectures from _Colin Barker_, Mar 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010498 | OEIS-A267960

Number of n X 1 0..2 arrays with every repeated value in every row and column greater than the previous repeated value.

Conjectures from _Colin Barker_, Feb 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010499 | OEIS-A267969

Conjectured list of positive numbers k such that the Collatz-like map T: if x > 1 and x odd, x -> 5*x+1 and, if x even, x -> x/2, when started at k, eventually reaches a cycle that does not contain 1 (cf. A232711).

Conjectured list of positive numbers k such that the Collatz-like map T: if x > 1 and x odd, x -> 5*x+1 and, if x even, x -> x/2, when started at k, eventually reaches a cycle that does not contain 1 (cf. A232711). A companion to A232711 (which conjecturally lists the numbers whose trajectory reaches 1), and A267970 (which conjecturally lists the numbers whose trajectory diverges). This is conjectural in that there is no proof that the list is complete as far as it goes. Some of the terms in A267970 could belong to

A232711 or this sequence if the trajectory is extended far enough. - _N. J. A. Sloane_, Jan 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010500 | OEIS-A267970

Conjectured list of positive numbers n such that the Collatz-like map T: if x > 1 and x odd, x -> 5*x+1 and, if x even, x -> x/2, when started at n, never repeats.

Conjectured list of positive numbers n such that the Collatz-like map T: if x > 1 and x odd, x -> 5*x+1 and, if x even, x -> x/2, when started at n, never repeats. A companion to A232711 (which conjecturally lists the numbers whose trajectory reaches 1), and A267969 (which conjecturally lists the numbers whose trajectory cycles but does not reach 1). The present sequences conjecturally lists the numbers whose trajectory diverges.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010501 | OEIS-A267999

Numbers n > 1 such that gcd(n, 2^n - 2) = 1.

Conjecture: this is a subsequence of A121707. Tested for all terms <= 10^5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_ and _Robert Israel_, Jan 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010502 | OEIS-A268052

Number of n X 4 0..2 arrays with new values introduced in each row and column in sequential order starting with zero.

Conjectures from _Colin Barker_, Jan 11 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010503 | OEIS-A268053

Number of n X 5 0..2 arrays with new values introduced in each row and column in sequential order starting with zero.

Conjectures from _Colin Barker_, Jan 11 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010504 | OEIS-A268081

Least positive integer k such that 3^n-1 and k^n-1 are relatively prime.

According to the conjecture given in A086892, a(n) = 2 infinitely often.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tom Edgar_, Jan 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010505 | OEIS-A268093

Number of 1 X n 0..2 arrays with every repeated value in every row not one larger and in every column one larger mod 3 than the previous repeated value, and upper left element zero.

Conjectures from _Colin Barker_, Jan 11 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010506 | OEIS-A268107

Decimal expansion of 'lambda', a Somos quadratic recurrence constant mentioned by Steven Finch.

Conjecture: lambda is the radius of convergence of the function Sum_{n>=0} A045761(n)*x^n, that is the constant 1/d computed by _Vaclav Kotesovec_ in A045761.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Jan 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010507 | OEIS-A268112

Numbers k for which the numerator of the k-th harmonic number H_k is divisible by the third power of a prime less than k.

It is a widely believed conjecture that there is no pair of an integer k and a prime p for which v_p(H_k) >= 4. If variations of this conjecture hold, then Krattenhaler and Rivoal (2007-2009) would be able to establish some formulas for their theory. See also A007757, A131657, and A131658. - _Petros Hadjicostas_, May 25 2020 Terms a(4) and a(5) are conjectural as possible existence of smaller terms is not eliminated. Carofiglio et al. (2025) computed some further terms (see links). - _Max Alekseyev_, Apr 01 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Jan 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010508 | OEIS-A268132

If a(n) is a square (and n > 1), then a(n+1) = a(n) + a(n-1), else a(n+1) is the smallest positive integer not occurring earlier.

Conjectured to be a permutation of the positive integers (which could be enforced by definition). In case there would occur a duplicate, it must be of the form a(n+1) = a(n) + a(n-1) and equal to an earlier term a(m+1) of the same form, where furthermore the predecessor a(m-1) also is of that form, since otherwise a(m+1) would be smaller than this a(n+1). This seems extremely unlikely to happen, and maybe provably impossible.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010509 | OEIS-A268136

a(n) = (3/n)*Sum_{k=0..n-1} A245769(k)^2.

Conjecture: (i) All the terms are odd integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010510 | OEIS-A268137

a(n) = (1/n)*Sum_{k=0..n-1} A001850(k)*A245769(k).

Conjecture: (i) All the terms are odd integers. For any prime p, if p == 3 (mod 4) then a(p) == -5 (mod p^2), otherwise a(p) == -1 (mod p).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010511 | OEIS-A268149

A double binomial sum involving absolute values.

Conjecture D-finite with recurrence (2*n-1)*(n-1)*a(n) +2*(-22*n^2+27*n-36)*a(n-1) +12*(4*n-5)*(4*n-7)*a(n-2)=0. - _R. J. Mathar_, Feb 27 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard P. Brent_, Jan 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010512 | OEIS-A268164

Number of n X 1 0..3 arrays with every repeated value in every row and column one larger mod 4 than the previous repeated value, and upper left element zero.

Conjectures from _Colin Barker_, Jan 11 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010513 | OEIS-A268197

Number of ordered ways to write n as w^2 + x^2 + y^2 + z^2 with w*(25*w + 24*x + 48*y + 96*z) a square, where w is a positive integer and x,y,z are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 3, 7, 15, 23, 43, 55, 463, 4^k*m (k = 0,1,2,... and m = 1, 31, 34).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010514 | OEIS-A268212

Numbers n of the form 2^k + 1 such that n^2 - n - 1 is a prime q (for k >= 0).

Conjecture: subsequence of prime terms (3, 5, 17, 65537, ...) is not the same as A249759.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jan 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010515 | OEIS-A268251

Expansion of x*(1 + 2*x - 96*x^2 - 148*x^3 + 45*x^4 + 50*x^5 + 2*x^6)/(1 - 99*x^2 + 99*x^4 - x^6).

Conjecture: The sequence lists all nonnegative m, in increasing order, such that floor(m/2)*floor(m/3) is a square. This conjecture has been proved by _Robert Israel_ (see paper in Links section).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bruno Berselli_, Jan 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010516 | OEIS-A268255

Number of length-(n+1) 0..2 arrays with new repeated values introduced in sequential order starting with zero.

Conjectures from _Colin Barker_, Jan 11 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010517 | OEIS-A268262

Number of length-(3+1) 0..n arrays with new repeated values introduced in sequential order starting with zero.

Conjectures from _Colin Barker_, Jan 13 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010518 | OEIS-A268263

Number of length-(4+1) 0..n arrays with new repeated values introduced in sequential order starting with zero.

Conjectures from _Colin Barker_, Jan 13 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010519 | OEIS-A268264

Number of length-(5+1) 0..n arrays with new repeated values introduced in sequential order starting with zero.

Conjectures from _Colin Barker_, Jan 13 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010520 | OEIS-A268265

Number of length-(6+1) 0..n arrays with new repeated values introduced in sequential order starting with zero.

Conjectures from _Colin Barker_, Jan 13 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010521 | OEIS-A268267

Orbit periods of the iteration s(1)=0, s(m+1)=(2*i*Pi*n)/log(3)+log(1/(-1-2^-s(m)))/log(3).

The iteration converges to sequences with periods of orbit equal to the entries in this sequence a(n). The entries in the converged period are then zeros to equations as in the examples in the conjecture below: Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Jan 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010522 | OEIS-A268293

a(n) = A266489(n)/n, for n>=1.

Conjectured to consist entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010523 | OEIS-A268298

G.f.: Sum_{n=-oo..+oo} x^n * (1 - x^n)^(3*n).

For n>0, a(n) = 1 iff n = 3^k for k>=0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010524 | OEIS-A268300

G.f. satisfies: -1 = Product_{n>=1} (1-x^n) * (1 - x^n/A(x)) * (1 - x^(n-1)*A(x)), where g.f. A(x) = Sum_{n>=0} a(n)*2*(x/4)^n.

a(n) is odd iff n = 2^k for k>=0 or n=0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010525 | OEIS-A268317

Irregular triangle read by rows: T(n,k) gives the columns sum in the table Fib(n+1) X Fib(n), where k = 1..Fib(n), and 1's are assigned to cells on the longest diagonal path.

(iii) The row sum of each of these irregular triangles is conjectured to be A000071. (iv) The first differences of the sequence of half of the voids (0's) are conjectured to give A191797.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Feb 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010526 | OEIS-A268318

Irregular triangle read by rows: T(n,k) gives the row sums in the table Fib(n+1) X Fib(n), where k = 1..Fib(n+1), and 1's are assigned to cells on the longest diagonal path.

(iii) The row sums of each of these irregular triangles is conjectured to be A000071. (iv) The first differences of the sequence of half of the voids (0's) are conjectured to give A191797.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Feb 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010527 | OEIS-A268343

Hermit primes: primes which are not a nearest neighbor of another prime.

Conjecture: the asymptotic probability of a prime being in this list is 1/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Karl W. Heuer_, Feb 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010528 | OEIS-A268369

Number of 3 X n arrays containing n copies of 0..3-1 with row sums equal.

Conjecture: n^4*(1057*n^2 -3473*n +2902)*a(n) -(n-1)*(14798*n^5 -63420*n^4 +104199*n^3 -81821*n^2 +31698*n -4968)*a(n-1) +(-235711*n^6 +1717323*n^5 -5084885*n^4 +7773081*n^3 -6422772*n^2 +2716308*n -472176)*a(n-2) -96*(6*n -11) *(6*n-13) *(1057*n^2-1359*n+486) *(n-2)^2*a(n-3)=0. - _R. J. Mathar_, Feb 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010529 | OEIS-A268399

Number of North-East lattice paths from (0,0) to (n,n) that have exactly four east steps below the subdiagonal y = x-1.

Conjecture: -(n+2)*(641*n-5292)*a(n) +(-641*n^2-1681*n+22692)*a(n-1) +(-4067*n^2+10502*n+80184) *a(n-2) +6*(5629*n-11668)*(2*n-11) *a(n-3)=0. - _R. J. Mathar_, Jun 07 2016 Conjecture: +(n+2)*(1023*n^2-11311*n+23730) *a(n) +(1023*n^3+15038*n^2-85751*n+44490)*a(n-1) -10*(2*n-9) *(1023*n^2-2563*n+1450)*a(n-2)=0. - _R. J. Mathar_, Jun 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ran Pan_, Feb 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010530 | OEIS-A268402

Number of North-East lattice paths from (0,0) to (n,n) that bounce off the diagonal y = x to the right exactly four times.

Conjecture: -(n-3)*(n-5)*(55*n-618)*a(n) +(-55*n^3-840*n^2+6969*n-5742)*a(n-1)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ran Pan_, Feb 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010531 | OEIS-A268447

Number of North-East lattice paths from (0,0) to (n,n) that cross the diagonal y = x horizontally exactly four times.

Conjecture: -(n+10)*(n-8)*a(n) +2*n*(2*n+1)*a(n-1)=0. - _R. J. Mathar_, Jun 07 2016 Conjecture follows from the differential equation (6*x + 80)*y(x) + (14*x^2 - 3*x)*y'(x) + (4*x^3 - x^2)*y''(x) = 0 satisfied by the G.f. - _Robert Israel_, May 15 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ran Pan_, Feb 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010532 | OEIS-A268459

Number of length-5 0..n arrays with no adjacent pair x,x+1 followed at any distance by x+1,x.

Conjectures from _Colin Barker_, Jan 13 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010533 | OEIS-A268460

Number of length-6 0..n arrays with no adjacent pair x,x+1 followed at any distance by x+1,x.

Conjectures from _Colin Barker_, Jan 13 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010534 | OEIS-A268461

Number of length-7 0..n arrays with no adjacent pair x,x+1 followed at any distance by x+1,x.

Conjectures from _Colin Barker_, Jan 14 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010535 | OEIS-A268477

Balanced odious primes: primes with an odd number of runs of 1's in their binary expansion.

According to our 2007-conjecture, if pi_1(m) is the number of evil primes (A027699) not exceeding m and pi_2(m) is the number of odious primes (A027697) not exceeding m, then pi_1(m)<=pi_2(m) for all natural m except m=5 and m=6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Feb 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010536 | OEIS-A268483

Primes p such that the numbers of primes not exceeding p in A268476 and A268477 are equal.

In contrast to the analogous sequence for odious and evil primes (A027697, A027699), which, as we conjecture, consists of only primes 3,7,29 (see also our 2007-conjecture in A027697, A027699), here we conjecture that the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Feb 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010537 | OEIS-A268487

Numbers of equal electric charges for which the minimum-potential dislocation on a sphere has nonzero sum of position vectors.

Probably most of these terms are merely conjectural. - _N. J. A. Sloane_, Mar 31 2016 Numeric investigations were carried out by various authors for m = 1 to 204, and then for a number of selected cases (see references in the Wikipedia link). Among the studied cases, 312 is also known to belong to this sequence. All these cases have at most some type of C-symmetry (C_2,C_2v,C_s,C_3,C_3v). So far, 10 cases with no symmetry at all (C_1) were found, namely m = 61, 140, 149, 176, 179, 183, 186, 191, 194, 199. No simple algorithm to handle this open problem, nor a general formula, are known.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Feb 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010538 | OEIS-A268494

Indices n of "late birds" (a(n)<a(k) for all k>n) in A268630 (= a(n)^2+a(n+1) is prime).

Assuming that A268630 is a permutation of the nonnegative integers N (as conjectured), the characterization given in the name is equivalent to say that a(n) equals the least number not occurring earlier. The sequence defined that way is finite if and only if A268630 is not a permutation of N.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010539 | OEIS-A268507

Number of ordered ways to write n as w^2 + x^2 + y^2 + z^2 with w > 0, w >= x <= y <= z such that x^2*y^2 + y^2*z^2 + z^2*x^2 is a square, where w,x,y,z are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 2^k, 4^k*m (k = 0,1,2,... and m = 3, 7, 23, 31, 39, 47, 55, 71, 79, 151, 191, 551). See also A269400, A271510, A271513, A271518, A271665, A271714, A271721, A271724, A271775, A271778 and A271824 for other conjectures refining Lagrange's four-square theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010540 | OEIS-A268539

Numbers k such that 48*k + 25 is a perfect square.

Conjecture: the sequence terms are also the exponents in the expansion of Sum_{n >= 0} q^n/(Product_{k = 1..2*n+1} 1 + q^k) = 1 + q^2 - q^3 - q^7 + q^17 + q^25 - - + + ... - _Peter Bala_, Jan 15 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010541 | OEIS-A268542

The diagonal of the rational function 1/(1 - x - y - x y - x z - y z).

Conjecture: 2*n^2*(21*n-37)*a(n) -32*(7*n-3)*(3*n^2-7*n+3)*a(n-1) +(-1281*n^3+4819*n^2-5610*n+1920)*a(n-2) -3*(3*n-5)*(21*n-16)*(3*n-7)*a(n-3) = 0. - _R. J. Mathar_, Mar 11 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010542 | OEIS-A268543

The diagonal of 1/(1 - (y + z + x z + x w + x y w)).

Conjecture: 2*n^2*(17*n-23)*a(n) +(-1207*n^3+2840*n^2-1897*n+360)*a(n-1) + 4*(17*n-6)*(-3+2*n)^2*a(n-2) = 0. - _R. J. Mathar_, Mar 11 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010543 | OEIS-A268550

Diagonal of 1/(1 - x - y - z - x y - x z - y z).

Conjecture: 2*n^2*(n-1)*a(n) -(n-1)*(107*n^2-107*n+24)*a(n-1) +12*(-9*n^3+27*n^2-22*n+2)*a(n-2) -3*n*(3*n-5)*(3*n-7)*a(n-3)=0. - _R. J. Mathar_, Mar 10 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010544 | OEIS-A268551

Diagonal of 1/(1 - x + y + z + x y + x z - y z + x y z).

Conjecture: n^2*(n-2)*a(n) +(-50*n^3+149*n^2-109*n+21)*a(n-1) -(2*n-3) *(51*n^2-153*n+91)*a(n-2) +(-50*n^3+301*n^2-565*n+315)*a(n-3) +(n-1)*(n-3)^2*a(n-4)=0. - _R. J. Mathar_, Mar 10 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010545 | OEIS-A268552

Diagonal of the rational function 1/((1 - u v - u w - v w - u v w) * (1 - x y - x z - y z)).

Conjecture: the supercongruences a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) hold for all primes p >= 5 and all positive integers n and r. - _Peter Bala_, Jun 22 2023 Conjecture 1. Let m be an integer. The sequence defined by u(n) = [x^n] F(x)^(m*n/6) satisfies the supercongruences u(n*p^r) === u(n*p^(r-1)) (mod p^r) for all primes p >= 5 and all positive integers n and r. Conjecture 2. Let m be an integer. The sequence defined by v(n) = [x^n] E(x)^(m*n/6) satisfies the supercongruences v(n*p^r) === v(n*p^(r-1)) (mod p^r) for all primes p >= 5 and all positive integers n and r. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010546 | OEIS-A268553

Diagonal of the rational function 1/((1 - u v - u w - v w) * (1 - x y - x z - y z)).

Conjecture: n^4*a(n) -9*(3*n-1)^2*(3*n-2)^2*a(n-1) = 0. - _R. J. Mathar_, Mar 11 2016 G.f.: 4F3(1/3,1/3,2/3,2/3,2/3; 1,1,1; 729*x). Mathar's conjecture above is true. - _Benedict W. J. Irwin_, Oct 20 2016 Conjecture 1. Let m be an integer. The sequence defined by u(n) = [x^n] F(x)^(m*n/36) satisfies the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and all positive integers n and r. Conjecture 2. Let m be an integer. The sequence defined by v(n) = [x^n] E(x)^(m*n/36) satisfies the supercongruences v(n*p^r) == v(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and all positive integers n and r. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010547 | OEIS-A268554

Diagonal of the rational function 1/((1 - w - u v) * (1 - x y - x z - y z)).

Conjecture: n^3*(2*n-1)*a(n) -6*(4*n-1)*(3*n-1)*(3*n-2)*(4*n-3)*a(n-1)=0. - _R. J. Mathar_, Mar 10 2016 For integer r and positive integer s, define sequences {u(n) : n >= 0} and {v(n) : n >= 0} by setting u(n) = [x^(s*n)] F(x)^(r*n) and v(n) = [x^(s*n)] E(x)^(r*n). We conjecture that both u(n) and v(n) satisfy the above supercongruences. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010548 | OEIS-A268593

The largest of three distinct primes, a > b > c > 0, such that each of the following four numbers are also prime: a+b+c, a+b-c, a-b+c, a-b-c, regardless of sign.

Once found, the following three properties of these septets are observed without exception for "a" values up to prime(800), resulting in these conjectures: From the first conjecture it follows that no values for "a", "b" or "c" belong to A172256. The first three conjectures can be used to accelerate the search for prime septets. Further observations below are from accelerated searches.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard R. Forberg_, Feb 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010549 | OEIS-A268597

Smallest x such that x-1 mod phi(x) = n, or 0 if no such x exists.

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christina Steffan_, Feb 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010550 | OEIS-A268600

Expansion of 1/(2*f(x)) + 1/(4 - 2*g(x)), where f(x) = sqrt(1 - 4*x) and g(x) = sqrt(1 + 4*x).

Conjecture D-finite with recurrence: -3*n*(n-1)*a(n) +8*(n-1)*(5*n-12)*a(n-1) +4*(-28*n+73)*a(n-2) -160*(2*n-5)*(2*n-7)*a(n-3) +192*(2*n-5)*(2*n-7)*a(n-4)=0. - _R. J. Mathar_, Jan 25 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ran Pan_, Feb 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010551 | OEIS-A268622

Number of n X 2 0..2 arrays with some element plus some horizontally or vertically adjacent neighbor totalling two no more than once.

Conjectures from _Colin Barker_, Mar 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010552 | OEIS-A268630

a(n)^2 + a(n+1) is prime; lexicographically earliest sequence of nonnegative integers with this property and containing no duplicates.

Conjectured to be a permutation of the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _M. F. Hasler_, Feb 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010553 | OEIS-A268633

Number of n X 2 0..2 arrays with some element plus some horizontally or vertically adjacent neighbor totalling two exactly once.

Conjectures from _Colin Barker_, Mar 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010554 | OEIS-A268642

Seelmann's sequence: a(1) = 1; thereafter a(n + 1) = ceiling(a(n)/2) unless this is already in the sequence, in which case a(n + 1) = 3*a(n).

It is conjectured that this is a permutation of the positive integers, along with any Seelmann sequence in which a(n+1) = M*a(n) if the divide by 2 rule cannot be applied, for any integer M>1 and not of the form M = 2^N. [Corrected by _Charlie Neder_, Feb 06 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Feb 09 2016, based on a posting by _David Seelmann_ to the Reddit web site | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010555 | OEIS-A268690

A268642 with a(n+1) = 5*n if ceiling(a(n)/2) is already in the sequence.

It is conjectured that this is a permutation of the integers, along with any Seelmann sequence in which a(n+1) = M*n if the divide by 2 rule is not permitted, for any integer M>1 and not of the form M = 2^N.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Seelmann_, Feb 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010556 | OEIS-A268694

Terms of A268693 that are not divisible by 255.

The conjecture in A156695 implies that each term greater than 1 can be written in the form p + 2^x, where p is a prime. For example: 144465 - 2^2 = 144461 (a prime), 4330455 - 2^8 = 4330199 (a prime).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Feb 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010557 | OEIS-A268755

Variant of A181391: For n >= 1, if there exists an m < n such that a(m) = a(n), take the largest such m and set a(n+1) = #{a(m), a(m+1), ..., a(n)}; otherwise, a(n+1) = 0. Start with a(1) = 0.

Conjecture: every pair of nonnegative integers (x,y) other than (1,1) appear as consecutive entries (a(i) = x, a(i+1) = y, for some i). - _László Kozma_, Aug 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nathaniel Shar_, Feb 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010558 | OEIS-A268757

Unhappycab numbers: the smallest unhappy number that is the sum of two cubes of unhappy numbers in n different ways.

a(n) >= A011541(n) for n > 0. It is conjectured that there is no equality.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Corinna Regina Böger_, Feb 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010559 | OEIS-A268759

Triangle T(n,k) read by rows: T(n,k) = (1/4)*(1 + k)*(2 + k)*(k - n)*(1 + k - n).

Notice that a(n) = T(n,k) is always a multiple of the triangular numbers, up to an offset. Conjecture: the triangle tabulating matrix elements <n(n+1)/2+k+1|J_1^p|n(n+1)/2+k+p+1> is determined entirely by the coefficients: binomial(n,p) (cf. A094053). Various sequences along the diagonals of matrix J_1^p lead to other numbers with geometric interpretations (Cf. A000567, A100165).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bradley Klee_, Feb 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010560 | OEIS-A268768

Number of n X 2 0..2 arrays with some element plus some horizontally, vertically, diagonally or antidiagonally adjacent neighbor totalling two exactly once.

Conjectures from _Colin Barker_, Jan 14 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010561 | OEIS-A268775

Number of n X 2 binary arrays with some element plus some horizontally, vertically, diagonally or antidiagonally adjacent neighbor totalling two no more than once.

Conjectures from _Colin Barker_, Jan 15 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010562 | OEIS-A268790

Magic sums of 3 X 3 magic squares composed of primes.

Conjecture: 3*p is a term for every prime > 859. If conjecture is true, a(n) = 3*prime(n+40) for n >= 110. - _Robert Israel_, Feb 16 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Feb 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010563 | OEIS-A268799

Record (maximal) gaps between primes of the form 4k + 3.

Conjecture: a(n) < phi(4)*log^2(A268801(n)) almost always. Conjecture: a(n) < phi(4)*n^2 for all n>2. - _Alexei Kourbatov_, Aug 12 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Feb 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010564 | OEIS-A268866

Records in A268865.

Conjecture 1: Let m(n) be the position in A268865 corresponding to a(n). Then m(n) = (2/3)*(4^n-1), if n is even, and m(n) = (2/3)*(4^(n-1)-1) + 3*4^(n-1), if n is odd. Conjecture 2: a(n) = 2*m(n) + 2, if n is even, and a(n) = (7*m+12)/11, if n is odd. If Conjectures 1 and 2 are true, then we easily have for even n >= 0, m(n) == 0 (mod 10), a(n) == 2 (mod 10), while a(1) = m(1) = 3 and for odd n >= 3, m(n), a(n) == 8 (mod 10). The sequence {m(n)} begins: 0, 3, 10, 58, 170, 938, 2730, 15018, 43690, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Feb 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010565 | OEIS-A268867

Number of integers of the form m^2+1 between two consecutive pairs of primes of the same form.

Conjecture 1: the sequence is infinite. Conjecture 2: for n>1, the sequence sorted in ascending order with distinct values generates the set B = {b(k)} = {7, 17, 27, 37, ...} = {7 + 10*k}, k >= 0, and {a(n)}/qZ = B/qZ with q = 2^m, m >= 1, is a multiplicative group. However, this concept is based on a conjecture if it is impossible to prove the infinity character of a sequence. This requires, for the calculations, a large number of elements in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Feb 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010566 | OEIS-A268898

Number of n X 2 0..2 arrays with some element plus some horizontally or antidiagonally adjacent neighbor totalling two exactly once.

Conjectures from _Colin Barker_, Jan 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010567 | OEIS-A268899

Number of n X 3 0..2 arrays with some element plus some horizontally or antidiagonally adjacent neighbor totalling two exactly once.

Conjectures from _Colin Barker_, Jan 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010568 | OEIS-A268900

Number of n X 4 0..2 arrays with some element plus some horizontally or antidiagonally adjacent neighbor totalling two exactly once.

Conjectures from _Colin Barker_, Jan 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010569 | OEIS-A268905

Number of 2 X n 0..2 arrays with some element plus some horizontally or antidiagonally adjacent neighbor totalling two exactly once.

Conjectures from _Colin Barker_, Jan 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010570 | OEIS-A268913

Magic sums of 3 X 3 semimagic squares composed of primes.

I conjecture that every odd number greater than 111 belongs to this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Feb 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010571 | OEIS-A268914

Minimum difference between two distinct primes whose sum is 2*prime(n), n>4.

Extrapolating from computational evidence for all primes up to 10^9, we conjecture: For each multiple of 12 there are infinitely many primes p such that p-6k and p+6k are prime and 12k is the minimal difference for two distinct primes whose sum is 2p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Barry Cherkas_, Feb 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010572 | OEIS-A268925

Record (maximal) gaps between primes of the form 6k + 1.

Conjecture: a(n) < phi(6)*log^2(A268927(n)) almost always. Conjecture: phi(6)*n^2/6 < a(n) < phi(6)*n^2 almost always. - _Alexei Kourbatov_, Nov 27 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Feb 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010573 | OEIS-A268939

Number of length-n 0..3 arrays with no repeated value unequal to the previous repeated value plus one mod 3+1.

Conjectures from _Colin Barker_, Jan 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010574 | OEIS-A268941

Number of length-n 0..5 arrays with no repeated value unequal to the previous repeated value plus one mod 5+1.

Conjectures from _Colin Barker_, Jan 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010575 | OEIS-A268943

Number of length-n 0..7 arrays with no repeated value unequal to the previous repeated value plus one mod 7+1.

Conjectures from _Colin Barker_, Jan 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010576 | OEIS-A268945

Number of length-4 0..n arrays with no repeated value unequal to the previous repeated value plus one mod n+1.

Conjectures from _Colin Barker_, Jan 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010577 | OEIS-A268946

Number of length-5 0..n arrays with no repeated value unequal to the previous repeated value plus one mod n+1.

Conjectures from _Colin Barker_, Jan 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010578 | OEIS-A268947

Number of length-6 0..n arrays with no repeated value unequal to the previous repeated value plus one mod n+1.

Conjectures from _Colin Barker_, Jan 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010579 | OEIS-A268948

Number of length-7 0..n arrays with no repeated value unequal to the previous repeated value plus one mod n+1.

Conjectures from _Colin Barker_, Jan 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010580 | OEIS-A268965

Number of n X 2 0..2 arrays with some element plus some horizontally or antidiagonally adjacent neighbor totalling two not more than once.

Conjectures from _Colin Barker_, Jan 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010581 | OEIS-A268966

Number of n X 3 0..2 arrays with some element plus some horizontally or antidiagonally adjacent neighbor totalling two not more than once.

Conjectures from _Colin Barker_, Jan 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010582 | OEIS-A268967

Number of n X 4 0..2 arrays with some element plus some horizontally or antidiagonally adjacent neighbor totalling two not more than once.

Conjectures from _Colin Barker_, Jan 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010583 | OEIS-A268972

Number of 2 X n 0..2 arrays with some element plus some horizontally or antidiagonally adjacent neighbor totalling two not more than once.

Conjectures from _Colin Barker_, Jan 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010584 | OEIS-A268983

Numbers n such that Sum_{d|n} sigma(n/d)^d/d is an integer, where sigma is A000203.

Conjecture: This sequence contains only the three numbers listed. This conjecture is disproven. The number 24083216081502612530611745 53 belongs to this sequence, but it may not be a(4). - _Gevorg Hmayakyan_, May 11 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gevorg Hmayakyan_, Feb 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010585 | OEIS-A268984

Record (maximal) gaps between primes of the form 10k + 1.

Conjecture: a(n) < phi(10)*log^2(A268986(n)) almost always.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Feb 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010586 | OEIS-A269012

Number of 2 X n binary arrays with some element plus some horizontally, diagonally or antidiagonally adjacent neighbor totalling two exactly once.

Conjectures from _Colin Barker_, Jan 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010587 | OEIS-A269021

Number of permutations of [2n] containing an increasing subsequence of length n.

Conjectured recurrence: -64*(1 + n)^2*(2 + n)^2*(3 + n)*(1 + 2*n)^2*(3 + 2*n)^2*(5 + 2*n)^2*(549760 + 3266000*n + 7264534*n^2 + 8663374*n^3 + 6333869*n^4 + 3012795*n^5 + 952323*n^6 + 198469*n^7 + 26156*n^8 + 1968*n^9 + 64*n^10)*a(n) + 16*(2 + n)^2*(3 + n)*(3 + 2*n)^2*(5 + 2*n)^2*(-5543040 - 487964*n + 78563984*n^2 + 229526554*n^3 + 325846005*n^4 + 284054698*n^5 + 165789363*n^6 + 67385886*n^7 + 19359535*n^8 + 3917758*n^9 + 545913*n^10 + 49788*n^11 + 2672*n^12 + 64*n^13)*a(n+1) - 4*(3 + n)*(5 + 2*n)^2*(-250467360 - 946158512*n - 562569136*n^2 + 3271711596*n^3 + 9313604242*n^4 + 12741784568*n^5 + 11118771121*n^6 + 6753094929*n^7 + 2966908118*n^8 + 959068672*n^9 + 228837227*n^10 + 39928763*n^11 + 4969164*n^12 + 419248*n^13 + 21568*n^14 + 512*n^15)*a(n+2) + 2*(-1300242000 - 6360099840*n - 10812498240*n^2 - 778719870*n^3 + 27672902302*n^4 + 55047935941*n^5 + 60004107039*n^6 + 43766004538*n^7 + 22820793074*n^8 + 8747566435*n^9 + 2488583381*n^10 + 523718876*n^11 + 80260596*n^12 + 8677944*n^13 + 624800*n^14 + 26752*n^15 + 512*n^16)*a(n+3) - 3*(4 + n)*(8 + 3*n)*(10 + 3*n)*(-15900 - 61278*n - 68928*n^2 + 48699*n^3 + 204716*n^4 + 233810*n^5 + 143536*n^6 + 52389*n^7 + 11324*n^8 + 1328*n^9 + 64*n^10)*a(n+4) = 0. - _Manuel Kauers_ and _Christoph Koutschan_, Mar 01 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Feb 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010588 | OEIS-A269043

a(n) is the number of distinct values that can be expressed as prime(n+k) + prime(n-k) in at least 2 different ways.

Conjecture: a(n) > 0 for n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Feb 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010589 | OEIS-A269091

Number of n X 2 0..3 arrays with some element plus some horizontally or vertically adjacent neighbor totalling three exactly once.

Conjectures from _Colin Barker_, Mar 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010590 | OEIS-A269103

Number of n X 2 0..3 arrays with some element plus some horizontally or vertically adjacent neighbor totalling three no more than once.

Conjectures from _Colin Barker_, Mar 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010591 | OEIS-A269202

Number of 2 X n 0..3 arrays with some element plus some horizontally, diagonally or antidiagonally adjacent neighbor totalling three no more than once.

Conjectures from _Colin Barker_, Jan 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010592 | OEIS-A269230

Indices for which A037053(n), the smallest prime with n digits '0', does not have n consecutive digits '0'.

It is conjectured that, apart from A037053(0) = 2, all other terms have three nonzero digits and are therefore of the form A037053(n) = a*10^(n+2) + b*10^k + c, where 1 <= a,b,c <= 9 and 1 <= k <= n+1. I conjecture that k < n+1 for all n (where k is function of n, of course).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010593 | OEIS-A269234

Record (maximal) gaps between primes of the form 10k + 3.

Conjecture: a(n) < phi(10)*log^2(A269236(n)) almost always.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Feb 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010594 | OEIS-A269238

Record (maximal) gaps between primes of the form 10k + 7.

Conjecture: a(n) < phi(10)*log^2(A269240(n)) almost always.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Feb 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010595 | OEIS-A269261

Record (maximal) gaps between primes of the form 10k + 9.

Conjecture: a(n) < phi(10)*log^2(A269263(n)) almost always.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Feb 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010596 | OEIS-A269270

Number of n X 2 0..3 arrays with some element plus some horizontally or antidiagonally adjacent neighbor totalling three exactly once.

Conjectures from _Colin Barker_, Jan 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010597 | OEIS-A269271

Number of n X 3 0..3 arrays with some element plus some horizontally or antidiagonally adjacent neighbor totalling three exactly once.

Conjectures from _Colin Barker_, Jan 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010598 | OEIS-A269272

Number of n X 4 0..3 arrays with some element plus some horizontally or antidiagonally adjacent neighbor totalling three exactly once.

Conjectures from _Colin Barker_, Jan 20 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010599 | OEIS-A269277

Number of 2 X n 0..3 arrays with some element plus some horizontally or antidiagonally adjacent neighbor totalling three exactly once.

Conjectures from _Colin Barker_, Jan 21 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010600 | OEIS-A269284

Number of n X 3 0..3 arrays with some element plus some horizontally or antidiagonally adjacent neighbor totalling three no more than once.

Conjectures from _Colin Barker_, Jan 21 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010601 | OEIS-A269285

Number of n X 4 0..3 arrays with some element plus some horizontally or antidiagonally adjacent neighbor totalling three no more than once.

Conjectures from _Colin Barker_, Jan 21 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010602 | OEIS-A269290

Number of 2 X n 0..3 arrays with some element plus some horizontally or antidiagonally adjacent neighbor totalling three no more than once.

Conjectures from _Colin Barker_, Jan 21 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010603 | OEIS-A269304

a(n) = n + n/gpf(n) + 1, where gpf(n) is the greatest prime factor of n or 1 if n = 1.

It is conjectured that any two sequences generated with a(n)=a(n-1)+a(n-1)/gpf(a(n-1))+1 and any initial value >=1 will have a finite number of non-shared terms and an infinite number of shared terms after one initial shared term (see A270807). Example: For a(1)=314, the sequence generated is 314, 317, 319, 331, 333, 343, 393, 397, 399, 421, 423, 433, ...; for a(1)=97, the sequence generated is 97, 99, 109, 111, 115, 121, 133, 141, 145, 151, 153, 163, 165, 181, 183, 187, 199, 201, 205, 211, 213, 217, 225, 271, 273, 295, 301, 309, 313, 315, 361, 381, 385, 421, 423, 433, ...; these sequences have respectively 9 and 33 terms not shared with the other until both reach 421; the following terms of both sequences are identical.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cody M. Haderlie_, Feb 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010604 | OEIS-A269341

Records in A269340.

Conjecture 1. For even n>=2, m(n)=(2/3)*(4^(n-1)-1); for odd n>=3, m(n)=(2/3)*(4^(n-2)-1)+3*4^(n-2). Conjecture 2. For even n>=2, a(n)=2*m(n)+2; for odd n>=3, a(n)=(7*m(n)+12)/11. (Cf. with conjectures in A268866).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Feb 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010605 | OEIS-A269346

Perfect cubes that are not the difference of two primes.

It is still an open conjecture that every even number is the difference of 2 primes. On the other hand, a computer test shows that all even cubes <= 10^21 can be written as the difference of 2 primes. The computer program generating the sequence needs an additional part to test for even cubes besides checking that for odd m^3, m^3+2 is composite. - _Chai Wah Wu_, Mar 03 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Waldemar Puszkarz_, Feb 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010606 | OEIS-A269361

Lexicographically first injection of natural numbers beginning with a(1)=1 such that a(n)*a(n+1) is a fibbinary number (A003714), i.e., has no adjacent 1's in its base-2 representation.

The sequence is conjectured to be a permutation of A091072.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010607 | OEIS-A269363

Lexicographically first injection of natural numbers beginning with a(1)=3 such that for all n >= 1, a(n)*a(n+1) is a fibbinary number (A003714), i.e., has no adjacent 1's in its base-2 representation.

The sequence is conjectured to be a permutation of A091067.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010608 | OEIS-A269400

Number of ordered ways to write n as w^2 + x^2 + y^2 + z^2 with 6*w^2*x^2 + 12*x^2*y^2 + 52*y^2*z^2 + 27*z^2*w^2 a square, where w,x,y are nonnegative integers and z is a positive integer.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 3, 15, 31, 39, 71, 79, 195, 311, 319, 403, 559, 591, 683, 719, 1031, 1439, 1643, 2519, 6879, 2^k, 2^(2k+1)*39 (k = 0,1,2,...). Also, any positive integer can be written as w^2 + x^2 + y^2 + z^2 with x a positive integer and w,y,z nonnegative integer such that 6*w^2*x^2 + 12*x^2*y^2 + 52*y^2*z^2 + 27*z^2*w^2 is a square. See also A268507, A271510, A271513, A271518, A271665, A271714, A271721, A271724, A271775, A271778 and A271824 for other conjectures refining Lagrange's four-square theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010609 | OEIS-A269410

Number of length-4 0..n arrays with no repeated value greater than or equal to the previous repeated value.

Conjectures from _Colin Barker_, Jan 21 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010610 | OEIS-A269411

Number of length-5 0..n arrays with no repeated value greater than or equal to the previous repeated value.

Conjectures from _Colin Barker_, Jan 21 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010611 | OEIS-A269412

Number of length-6 0..n arrays with no repeated value greater than or equal to the previous repeated value.

Conjectures from _Colin Barker_, Jan 21 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010612 | OEIS-A269413

Number of length-7 0..n arrays with no repeated value greater than or equal to the previous repeated value.

Conjectures from _Colin Barker_, Jan 21 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010613 | OEIS-A269420

Record (maximal) gaps between primes of the form 8k + 3.

Conjecture: a(n) < phi(8)*log^2(A269422(n)) almost always.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Feb 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010614 | OEIS-A269424

Record (maximal) gaps between primes of the form 8k + 1.

Conjecture: a(n) < phi(8)*log^2(A269426(n)) almost always.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Feb 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010615 | OEIS-A269427

a(1) = 1, a(n) counts m < n for which n == a(m) (mod m).

I conjecture that this sequence is unbounded. Consider the first k terms of this sequence, and let L be the floor of log(k). If we count the times that each number 1,2,...,2L appears among the first k terms of this sequence, it appears that these sums form a normal distribution centered at L, so that L appears approximately k/10 times among the first k terms of this sequence. (For instance, in the first k = 10000 terms of the sequence, L = log(10000) = 9 appears 1174 times, a maximal count among any value that appears at all.) Thus the sequence appears to be unbounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alec Jones_, Feb 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010616 | OEIS-A269436

Number of length-4 0..n arrays with no repeated value greater than the previous repeated value.

Conjectures from _Colin Barker_, Jan 22 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010617 | OEIS-A269437

Number of length-5 0..n arrays with no repeated value greater than the previous repeated value.

Conjectures from _Colin Barker_, Jan 22 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010618 | OEIS-A269438

Number of length-6 0..n arrays with no repeated value greater than the previous repeated value.

Conjectures from _Colin Barker_, Jan 22 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010619 | OEIS-A269439

Number of length-7 0..n arrays with no repeated value greater than the previous repeated value.

Conjectures from _Colin Barker_, Jan 22 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010620 | OEIS-A269454

Safe primes that are not congruent to -1 mod 8.

Conjecture: If p and q are two distinct safe primes not congruent to -1 mod 8 then the order of 2 mod p*q is phi(p*q)/2. For phi see A000010.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marina Ibrishimova_, Feb 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010621 | OEIS-A269461

Number of length-n 0..2 arrays with no repeated value equal to the previous repeated value.

Conjectures from _Colin Barker_, Mar 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010622 | OEIS-A269462

Number of length-n 0..3 arrays with no repeated value equal to the previous repeated value.

Conjectures from _Colin Barker_, Jan 22 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010623 | OEIS-A269464

Number of length-n 0..5 arrays with no repeated value equal to the previous repeated value.

Conjectures from _Colin Barker_, Jan 22 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010624 | OEIS-A269465

Number of length-n 0..6 arrays with no repeated value equal to the previous repeated value.

Conjectures from _Colin Barker_, Jan 22 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010625 | OEIS-A269466

Number of length-n 0..7 arrays with no repeated value equal to the previous repeated value.

Conjectures from _Colin Barker_, Jan 22 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010626 | OEIS-A269467

T(n,k)=Number of length-n 0..k arrays with no repeated value equal to the previous repeated value.

The conjectures regarding the recursions for column k are correct (see links) - _Sela Fried_, Oct 29 2024.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010627 | OEIS-A269468

Number of length-4 0..n arrays with no repeated value equal to the previous repeated value.

Conjectures from _Colin Barker_, Jan 23 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010628 | OEIS-A269469

Number of length-5 0..n arrays with no repeated value equal to the previous repeated value.

Conjectures from _Colin Barker_, Jan 23 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010629 | OEIS-A269470

Number of length-6 0..n arrays with no repeated value equal to the previous repeated value.

Conjectures from _Colin Barker_, Jan 23 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010630 | OEIS-A269471

Number of length-7 0..n arrays with no repeated value equal to the previous repeated value.

Conjectures from _Colin Barker_, Jan 23 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010631 | OEIS-A269487

Number of length-n 0..n arrays with no repeated value differing from the previous repeated value by one.

Conjecture: a(n) ~ exp(1) * n^n. - _Vaclav Kotesovec_, Feb 28 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010632 | OEIS-A269495

Number of length-4 0..n arrays with no repeated value differing from the previous repeated value by one.

Conjectures from _Colin Barker_, Jan 23 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010633 | OEIS-A269496

Number of length-5 0..n arrays with no repeated value differing from the previous repeated value by one.

Conjectures from _Colin Barker_, Jan 23 2019: (Start)
Conjectured e.g.f.: exp(x)*(1 + 23*x+ 84*x^2 + 65*x^3 + 15*x^4 + x^5) - 1. - _Stefano Spezia_, Feb 20 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010634 | OEIS-A269511

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 342", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 03 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010635 | OEIS-A269512

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 342", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 03 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010636 | OEIS-A269513

Record (maximal) gaps between primes of the form 8k + 5.

Conjecture: a(n) < phi(8)*log^2(A269515(n)) almost always.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Feb 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010637 | OEIS-A269519

Record (maximal) gaps between primes of the form 8k + 7.

Conjecture: a(n) < phi(8)*log^2(A269521(n)) almost always.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Feb 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010638 | OEIS-A269526

Square array T(n,k) (n>=1, k>=1) read by antidiagonals upwards in which each term is the least positive integer satisfying the condition that no row, column, diagonal, or antidiagonal contains a repeated term.

I conjecture that other similar cyclical difference patterns should hold for any column k (i.e., terms a((j^2+j+2*k)/2), j>=k-1), so that each column is a permutation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alec Jones_, Apr 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010639 | OEIS-A269538

Number of length-4 0..n arrays with no repeated value differing from the previous repeated value by other than one.

Conjectures from _Colin Barker_, Jan 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010640 | OEIS-A269539

Number of length-5 0..n arrays with no repeated value differing from the previous repeated value by other than one.

Conjectures from _Colin Barker_, Jan 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010641 | OEIS-A269540

Number of length-6 0..n arrays with no repeated value differing from the previous repeated value by other than one.

Conjectures from _Colin Barker_, Jan 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010642 | OEIS-A269541

Number of length-7 0..n arrays with no repeated value differing from the previous repeated value by other than one.

Conjectures from _Colin Barker_, Jan 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010643 | OEIS-A269565

Array read by antidiagonals: T(n,m) is the number of (directed) Hamiltonian paths in K_n X K_m.

Conjecture: T(n,m) mod n!*m! = 0. - _Mikhail Kurkov_, Feb 08 2019 The above conjecture is true since a path defines an ordering on the rows and columns by the order in which they are first visited by the path. Every permutation of rows and columns therefore gives a different path. - _Andrew Howroyd_, Feb 08 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Howroyd_, Feb 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010644 | OEIS-A269584

Number of length-4 0..n arrays with no repeated value differing from the previous repeated value by more than one.

Conjectures from _Colin Barker_, Jan 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010645 | OEIS-A269585

Number of length-5 0..n arrays with no repeated value differing from the previous repeated value by more than one.

Conjectures from _Colin Barker_, Jan 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010646 | OEIS-A269586

Number of length-6 0..n arrays with no repeated value differing from the previous repeated value by more than one.

Conjectures from _Colin Barker_, Jan 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010647 | OEIS-A269587

Number of length-7 0..n arrays with no repeated value differing from the previous repeated value by more than one.

Conjectures from _Colin Barker_, Jan 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010648 | OEIS-A269607

Number of length-4 0..n arrays with no repeated value differing from the previous repeated value by one or less.

Conjectures from _Colin Barker_, Jan 25 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010649 | OEIS-A269608

Number of length-5 0..n arrays with no repeated value differing from the previous repeated value by one or less.

Conjectures from _Colin Barker_, Jan 25 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010650 | OEIS-A269609

Number of length-6 0..n arrays with no repeated value differing from the previous repeated value by one or less.

Conjectures from _Colin Barker_, Jan 25 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010651 | OEIS-A269610

Number of length-7 0..n arrays with no repeated value differing from the previous repeated value by one or less.

Conjectures from _Colin Barker_, Jan 25 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010652 | OEIS-A269613

Number of length-n 0..2 arrays with no repeated value differing from the previous repeated value by other than plus two, zero or minus 1.

Conjectures from _Colin Barker_, Mar 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010653 | OEIS-A269620

Number of length-4 0..n arrays with no repeated value differing from the previous repeated value by other than plus two, zero or minus 1.

Conjectures from _Colin Barker_, Jan 25 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010654 | OEIS-A269621

Number of length-5 0..n arrays with no repeated value differing from the previous repeated value by other than plus two, zero or minus 1.

Conjectures from _Colin Barker_, Jan 25 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010655 | OEIS-A269622

Number of length-6 0..n arrays with no repeated value differing from the previous repeated value by other than plus two, zero or minus 1.

Conjectures from _Colin Barker_, Jan 25 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010656 | OEIS-A269623

Number of length-7 0..n arrays with no repeated value differing from the previous repeated value by other than plus two, zero or minus 1.

Conjectures from _Colin Barker_, Jan 25 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010657 | OEIS-A269631

a(1) = 0; a(n+1) is the smallest integer not yet used that contains the number of decimal digits of a(n) as a substring.

Conjecture: a(n) is a permutation of the nonnegative integers. and the growth of percentage seems to support the conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Francesco Di Matteo_, Mar 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010658 | OEIS-A269641

Number of length-4 0..n arrays with no repeated value differing from the previous repeated value by other than plus two or minus 1.

Conjectures from _Colin Barker_, Jan 25 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010659 | OEIS-A269642

Number of length-5 0..n arrays with no repeated value differing from the previous repeated value by other than plus two or minus 1.

Conjectures from _Colin Barker_, Jan 25 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010660 | OEIS-A269643

Number of length-6 0..n arrays with no repeated value differing from the previous repeated value by other than plus two or minus 1.

Conjectures from _Colin Barker_, Jan 25 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010661 | OEIS-A269644

Number of length-7 0..n arrays with no repeated value differing from the previous repeated value by other than plus two or minus 1.

Conjectures from _Colin Barker_, Jan 25 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010662 | OEIS-A269657

Number of length-4 0..n arrays with no adjacent pair x,x+1 repeated.

I.e., a(n) = # {x in {0..n}^4 | x[1] != x[0]+1 or x[2] != x[0] or x[3] != x[1]}. The only possibility to have an adjacent x,x+1 pair repeated in a length-4 array is to have the array (x,x+1,x,x+1), with 0 <= x <= n-1 given the restriction on the domain of coefficients. This implies a(n) = (n+1)^4 - n and previously conjectured formulas. - _M. F. Hasler_, Feb 29 2020 Conjectures from _Colin Barker_, Jan 25 2019: (Start) a(n) = (n+1)^4 - n, cf. comment, confirming the above conjectured formulas. - _M. F. Hasler_, Feb 29 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010663 | OEIS-A269658

Number of length-5 0..n arrays with no adjacent pair x,x+1 repeated.

Conjectures from _Colin Barker_, Jan 25 2019: (Start) a(n) = (n+1)^5 - 3*n*(n+1) = A000584(n+1) - A028896(n), cf. comment, which confirms the above conjectured formulas. - _M. F. Hasler_, Feb 29 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010664 | OEIS-A269659

Number of length-6 0..n arrays with no adjacent pair x,x+1 repeated.

Conjectures from _Colin Barker_, Jan 25 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010665 | OEIS-A269660

Number of length-7 0..n arrays with no adjacent pair x,x+1 repeated.

Conjectures from _Colin Barker_, Jan 26 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010666 | OEIS-A269666

Prime sums of five Mersenne primes.

There are 368 terms up to 10^1000, 13 more up to 10^1332, none between 10^1332 and 10^2916, and 9 between 10^2916 and 10^3000. Conjecture: the sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Mar 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010667 | OEIS-A269673

Number of length-n 0..3 arrays with no repeated value differing from the previous repeated value by other than plus or minus one modulo 3+1.

Conjectures from _Colin Barker_, Jan 26 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010668 | OEIS-A269675

Number of length-n 0..5 arrays with no repeated value differing from the previous repeated value by other than plus or minus one modulo 5+1.

Conjectures from _Colin Barker_, Jan 26 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010669 | OEIS-A269677

Number of length-n 0..7 arrays with no repeated value differing from the previous repeated value by other than plus or minus one modulo 7+1.

Conjectures from _Colin Barker_, Jan 26 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010670 | OEIS-A269679

Number of length-4 0..n arrays with no repeated value differing from the previous repeated value by other than plus or minus one modulo n+1.

Conjectures from _Colin Barker_, Jan 26 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010671 | OEIS-A269680

Number of length-5 0..n arrays with no repeated value differing from the previous repeated value by other than plus or minus one modulo n+1.

Conjectures from _Colin Barker_, Jan 26 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010672 | OEIS-A269681

Number of length-6 0..n arrays with no repeated value differing from the previous repeated value by other than plus or minus one modulo n+1.

Conjectures from _Colin Barker_, Jan 26 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010673 | OEIS-A269682

Number of length-7 0..n arrays with no repeated value differing from the previous repeated value by other than plus or minus one modulo n+1.

Conjectures from _Colin Barker_, Jan 26 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010674 | OEIS-A269684

Number of length-n 0..2 arrays with no repeated value differing from the previous repeated value by plus or minus one modulo 2+1.

Conjectures from _Colin Barker_, Jan 27 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010675 | OEIS-A269685

Number of length-n 0..3 arrays with no repeated value differing from the previous repeated value by plus or minus one modulo 3+1.

Conjectures from _Colin Barker_, Jan 27 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010676 | OEIS-A269686

Number of length-n 0..4 arrays with no repeated value differing from the previous repeated value by plus or minus one modulo 4+1.

Conjectures from _Colin Barker_, Jan 27 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010677 | OEIS-A269687

Number of length-n 0..5 arrays with no repeated value differing from the previous repeated value by plus or minus one modulo 5+1.

Conjectures from _Colin Barker_, Jan 27 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010678 | OEIS-A269688

Number of length-n 0..6 arrays with no repeated value differing from the previous repeated value by plus or minus one modulo 6+1.

Conjectures from _Colin Barker_, Jan 27 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010679 | OEIS-A269689

Number of length-n 0..7 arrays with no repeated value differing from the previous repeated value by plus or minus one modulo 7+1.

Conjectures from _Colin Barker_, Jan 27 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010680 | OEIS-A269691

Number of length-5 0..n arrays with no repeated value differing from the previous repeated value by plus or minus one modulo n+1.

Conjectures from _Colin Barker_, Jan 28 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010681 | OEIS-A269692

Number of length-6 0..n arrays with no repeated value differing from the previous repeated value by plus or minus one modulo n+1.

Conjectures from _Colin Barker_, Jan 28 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010682 | OEIS-A269693

Number of length-7 0..n arrays with no repeated value differing from the previous repeated value by plus or minus one modulo n+1.

Conjectures from _Colin Barker_, Jan 28 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010683 | OEIS-A269696

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 6", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010684 | OEIS-A269708

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 14", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010685 | OEIS-A269712

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 20", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010686 | OEIS-A269716

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 22", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010687 | OEIS-A269728

Let k be a number not a power of 2 (see A057716), and define r by 2^(r-1) < k < 2^r; a(n) is smallest prime of the form 2^r*m+1 such that the exponential sum S(sigma_{m,k}) avoids p.

It is only a conjecture that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010688 | OEIS-A269729

a(n) = row number of extended Wythoff array (see A035513) which contains the sequence obtained by reading the n-th row backwards (and adjusting signs).

Conjecture: sequence is its own inverse. - _R. J. Mathar_, May 08 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010689 | OEIS-A269760

Number of n X 1 0..5 arrays with some element plus some horizontally or vertically adjacent neighbor totalling five exactly once.

Conjectures from _Colin Barker_, Mar 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010690 | OEIS-A269761

Number of n X 2 0..5 arrays with some element plus some horizontally or vertically adjacent neighbor totalling five exactly once.

Conjectures from _Colin Barker_, Jan 28 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010691 | OEIS-A269777

Number of length-5 0..n arrays with every repeated value unequal to the previous repeated value plus one mod n+1.

Conjectures from _Colin Barker_, Jan 29 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010692 | OEIS-A269778

Number of length-6 0..n arrays with every repeated value unequal to the previous repeated value plus one mod n+1.

Conjectures from _Colin Barker_, Jan 29 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010693 | OEIS-A269779

Number of length-7 0..n arrays with every repeated value unequal to the previous repeated value plus one mod n+1.

Conjectures from _Colin Barker_, Jan 29 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010694 | OEIS-A269781

a(n) is the smallest k different from n such that (n, k) is an amicably refactorable pair (see the comments).

Conjecture: the sequence contains no zeros.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Waldemar Puszkarz_, May 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010695 | OEIS-A269782

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 65", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 03 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010696 | OEIS-A269811

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 33", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 15 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010697 | OEIS-A269815

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 35", based on the 5-celled von Neumann neighborhood.

Lars Blomberg conjectured that Rules 43 and 59 also produce this sequence. It would be nice to have a proof. Conjecture: a(n) = 14*4^(n-1) - 5*2^n + 1, n>0. - _Lars Blomberg_, Apr 18 2016 Conjectures from _Colin Barker_, Apr 18 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010698 | OEIS-A269822

Number of n X 1 0..4 arrays with some element plus some horizontally, diagonally, antidiagonally or vertically adjacent neighbor totalling four exactly once.

Conjectures from _Colin Barker_, Mar 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010699 | OEIS-A269844

Primes equal to the sum of a pair of consecutive integers which are both squarefree.

Conjecture: sequence has density A271780/2 = A005597*4/Pi^2 = 0.2675535... in the primes. - _Charles R Greathouse IV_, Jan 24 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bill McEachen_, Mar 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010700 | OEIS-A269845

Irregular triangle read by rows: T(n,k) = (k/2+1/2)^2 if odd-k otherwise T(n,k) = (n-k/2)^2 where n >= 1, k = 0..2*n-1.

There are triangles appearing along main diagonal. If the area of the smallest triangles are defined as 1, then the areas of all other triangles seem to be square numbers. Conjectures: (i) Odd terms of row sum/2 is A100157. (ii) Even terms of row sum/2 is A258582. See illustration in links.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Mar 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010701 | OEIS-A269870

Numbers coprime to the number of their odd divisors.

Conjecture: the asymptotic density of this sequence is the same as that of A046642.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Waldemar Puszkarz_, Mar 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010702 | OEIS-A269876

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 43", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010703 | OEIS-A269878

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 43", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010704 | OEIS-A269879

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 43", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010705 | OEIS-A269895

Number of n X 1 0..6 arrays with some element plus some horizontally or vertically adjacent neighbor totalling six exactly once.

Conjectures from _Colin Barker_, Mar 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010706 | OEIS-A269896

Number of n X 2 0..6 arrays with some element plus some horizontally or vertically adjacent neighbor totalling six exactly once.

Conjectures from _Colin Barker_, Jan 29 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010707 | OEIS-A269906

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 1", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010708 | OEIS-A269907

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 1", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010709 | OEIS-A269908

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 1", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010710 | OEIS-A269909

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 1", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010711 | OEIS-A269910

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 3", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010712 | OEIS-A269911

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 3", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010713 | OEIS-A269912

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 3", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010714 | OEIS-A269913

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 3", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010715 | OEIS-A270006

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 5", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 09 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010716 | OEIS-A270008

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 5", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 09 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010717 | OEIS-A270009

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 5", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 09 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010718 | OEIS-A270010

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 7", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 09 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010719 | OEIS-A270012

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 7", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 09 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010720 | OEIS-A270013

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 7", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 09 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010721 | OEIS-A270026

a(n) is the smallest b for which the base-b representation of n contains at least one 0 (or 0 if no such base exists).

It is conjectured that the only terms that are neither 2 nor 3 in this sequence are a(7) and a(32767), which are 7 and 5 respectively.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nathan Fox_, Mar 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010722 | OEIS-A270072

Decimal representation of the n-th iteration of the Conway's game of life starting with four adjacent rows of lines turned on to positive and negative infinity.

It is conjectured that every term has a prime factor that does not divide any previous term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brook Estifanos_, Mar 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010723 | OEIS-A270073

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with x*y + 2*z*w a square, where x,y,z are nonnegative integers and w is an integer with x <= y and z >= |w|.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 3, 11, 15, 23, 43, 67, 79, 155, 211, 331, 347, 403, 427, 659, 899, 1443, 1955, 2^k*m (k = 0,1,2,... and m = 14, 35, 62, 158, 382).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010724 | OEIS-A270079

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 62", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 10 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010725 | OEIS-A270080

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 62", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = 4^n + 2*2^n - 8, n>2. - _Lars Blomberg_, Apr 20 2016 Conjectures from _Colin Barker_, Jun 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010726 | OEIS-A270082

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 62", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 10 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010727 | OEIS-A270084

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 65", based on the 5-celled von Neumann neighborhood.

Lars Blomberg conjectured that Rule 321 also produces this sequence. It would be nice to have a proof. Conjecture: a(n) = 4*4^n - 10*2^n + 12, n>1. - _Lars Blomberg_, Apr 21 2016 Conjectures from _Colin Barker_, Apr 21 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010728 | OEIS-A270085

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 65", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 11 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010729 | OEIS-A270086

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 65", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 11 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010730 | OEIS-A270087

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 73", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 11 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010731 | OEIS-A270088

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 73", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = 4*4^n - 4*2^n + 1, n>2. - _Lars Blomberg_, Apr 21 2016 Conjectures from _Colin Barker_, Apr 21 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010732 | OEIS-A270089

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 73", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 11 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010733 | OEIS-A270090

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 73", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 11 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010734 | OEIS-A270111

Number of n X 1 0..7 arrays with some element plus some horizontally or vertically adjacent neighbor totalling seven exactly once.

Conjectures from _Colin Barker_, Mar 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010735 | OEIS-A270112

Number of n X 2 0..7 arrays with some element plus some horizontally or vertically adjacent neighbor totalling seven exactly once.

Conjectures from _Colin Barker_, Jan 29 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010736 | OEIS-A270126

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 86", based on the 5-celled von Neumann neighborhood.

Lars Blomberg conjectured that Rule 342 also produces this sequence. It would be nice to have a proof. Conjectures from _Colin Barker_, Mar 21 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010737 | OEIS-A270129

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 89", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 12 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010738 | OEIS-A270130

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 89", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = 4*4^n - 4*2^n + 1, n>2. - _Lars Blomberg_, Apr 23 2016 Conjectures from _Colin Barker_, Jun 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010739 | OEIS-A270131

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 89", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 12 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010740 | OEIS-A270132

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 89", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 12 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010741 | OEIS-A270145

Number of n X 2 0..7 arrays with some element plus some horizontally, diagonally, antidiagonally or vertically adjacent neighbor totalling seven exactly once.

Conjectures from _Colin Barker_, Jan 30 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010742 | OEIS-A270164

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 107", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 12 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010743 | OEIS-A270165

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 107", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = 14*4^(n-1) - 3*2^n + 1, n>3. - _Lars Blomberg_, Apr 25 2016 Conjectures from _Colin Barker_, Apr 25 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010744 | OEIS-A270166

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 107", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 13 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010745 | OEIS-A270167

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 107", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 13 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010746 | OEIS-A270211

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 123", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = (14*4^(n-1) + 124*3^(n-3) - 159*2^(n-2) + 615)/3, n>5. - _Lars Blomberg_, Apr 30 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010747 | OEIS-A270218

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 129", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 13 2016: (Start) a(n) = 4*A006095(n+1) (conjectured). - _Michal Stajszczak_, May 20 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010748 | OEIS-A270222

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 131", based on the 5-celled von Neumann neighborhood.

It is conjectured that Rules 163, 171, 227, 235, 771, 787, 803, 811, 819, 827, 835, 851, 867, 875, 883 and 891 also generate this sequence. - _Lars Blomberg_, Apr 30 2016 Conjectures from _Colin Barker_, Mar 13 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010749 | OEIS-A270226

a(n) is the number of terms in the n-th block of consecutive integers of A136119.

Conjecture: Partial average of the sequence converges to 1+sqrt(2). Proof of the conjecture: since A136119(n+1) - A136119(n) = A001030(n), the sequence (a(n+1)) is the fixed point of the substitution sigma: 2->32, 3->322. Here one uses that since sigma(a)=a, the length of the n-th block is coded by the n-th letter. Since the frequencies of 2 and 3 in this fixed point are respectively sqrt(2)/(1+sqrt(2)) and 1/(1+sqrt(2)), the conjecture follows. (Alternatively: (a(n+1)-2) is a Sturmian sequence with density sqrt(2)-1). - _Michel Dekking_, Jan 22 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benedict W. J. Irwin_, Mar 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010750 | OEIS-A270270

The number of n-digit numbers in A270048.

Conjecture: lim_{n -> infinity} a(n)/a(n-1) = sqrt(10).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Francesco Di Matteo_, Mar 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010751 | OEIS-A270287

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 145", based on the 5-celled von Neumann neighborhood.

Lars Blomberg conjectured that Rule 153 also produces this sequence. It would be nice to have a proof. Conjecture: a(n) = 47*4^(n-2) - 4*2^n + 1, n>2. - _Lars Blomberg_, May 02 2016 Conjectures from _Colin Barker_, Jun 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010752 | OEIS-A270309

Irregular triangle read by rows: T(n,k) = ((n-k)+1)^2 if odd-n and odd-k; T(n,k) = k^2 if odd-n and even-k; T(n,k) = (n/2-(k/2-1/2))^2 if even-n and odd-k; T(n,k) = (k/2+1)^2 if even-n and even-k; where n >= 1, k = 1..2*n.

There are triangles appearing along main diagonal. If the area of the smallest triangles are defined as 1, then the areas of all other triangles seem to be square numbers. Conjectures: (i) Even terms of row sum is A002492. (ii) Odd terms of row sum/2 is A100157. See illustration in links.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Mar 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010753 | OEIS-A270360

Least positive integer k such that 5^n-1 and k^n-1 are relatively prime.

Based on conjecture given in A270390, a(n) = 2 infinitely often.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tom Edgar_, Mar 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010754 | OEIS-A270361

Odd primes p for which there exists an odd prime q < p such that p*q - 1 is a square.

Conjecture: For any odd prime p there is at most one odd prime q, with q < p, for which p*q-1 is square.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard R. Forberg_, Mar 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010755 | OEIS-A270363

a(n) = (n+1)*Sum_{k=0..(n-1)/2}((binomial(2*n-3*k-2,n-k-1))/(n-k)).

Conjecture: n*(7*n^2-17*n-2) *a(n) +(-35*n^3+99*n^2+20*n-120) *a(n-1) +2* (2*n-5) *(7*n^2-3*n-12)*a(n-2) +n*(7*n^2-17*n-2) *a(n-3) +2*-(2*n-5) *(7*n^2-3*n-12) *a(n-4)=0. - _R. J. Mathar_, Mar 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Mar 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010756 | OEIS-A270390

Greatest common divisor of 2^n-1 and 5^n-1.

Ailon and Rudnick conjecture that a(n) = 1 infinitely often.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tom Edgar_, Mar 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010757 | OEIS-A270413

Numbers m such that sigma(m-1) is a prime.

Conjecture: supersequence of A256438. Conjecture: 31 is the only prime p such that p = sigma(x-1) = sigma(y-1) for distinct numbers x and y; 31 = sigma(17-1) = sigma(26-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Mar 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010758 | OEIS-A270414

Numbers m such that sigma(m-1) and sigma(phi(m)) are both primes.

Conjecture: union of number 10 and A249759.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Mar 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010759 | OEIS-A270440

Least k such that binomial(k, 2) >= binomial(2*n, n).

Open question: Does binomial(a(n), 2) = binomial(2*n, n) for any n > 2? An affirmative answer would settle whether there exists an odd term greater than 3 in A003016. Conjecture: a(n) ~ 2^(n + 1/2) / (Pi*n)^(1/4). - _Vaclav Kotesovec_, Mar

23 2016 a(n) = ceiling(((8*binomial(2*n,n)+1)^(1/2)+1)/2). The above conjecture is true asymptotically. Using Stirling's formula for the approximation of n!, we get binomial(2*n,n) ~ 2^(2*n)/(Pi*n)^(1/2) and inserting this in the formula for a(n) results in the above approximation for a(n). - _Chai Wah Wu_, Mar 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Danny Rorabaugh_, Mar 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010760 | OEIS-A270454

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 163", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 17 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010761 | OEIS-A270455

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 163", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 17 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010762 | OEIS-A270456

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 163", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 17 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010763 | OEIS-A270469

Number of ordered ways to write n = x^3 + y*(y+1) + z*(3*z+2), where x and y are nonnegative integers and z is a nonzero integer.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 2, 3, 4, 5, 6, 10, 12, 20, 27, 56, 101. See also A262813 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010764 | OEIS-A270488

Number of ordered ways to write n = x^2 + y*(y+1) + z*(z^2+1), where x, y and z are nonnegative integers.

Conjecture: a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 1, 5, 7, 19, 35, 54, 62, 63, 197, 285, 339, 479, 505, 917. Moreover, any integer n > 2 can be written as x^2 + y*(y+1) + z*(z^2+1), where x is a positive integer, and y and z are nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010765 | OEIS-A270490

a(n) = Sum_{i=0..(n+1)/2} binomial(2*i+1,i)*binomial(2*n-2*i,n)/(2*i+1).

Conjecture: 3*n*(n-2)*(n+2)*a(n) -4*(n+1)*(8*n^2-23*n+12)*a(n-1) +16*n *(3*n-4)*(2*n-5)*a(n-2) +8*(2*n-3)*(4*n-7)*a(n-3) -64*(2*n-5)*(n-3)*(2*n-3)*a(n-4)=0. - _R. J. Mathar_, Jun 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Mar 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010766 | OEIS-A270516

Number of ordered ways to write n = x^3*(x+1) + y*(y+1)/2 + z*(3z+2), where x and y are nonnegative integers, and z is an integer.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and the only values of n > 1428 with a(n) = 1 are 2205, 2259, 3556, 4107, 4337, 5387, 9331, 16561, 22237, 27569, 63947, 78610. See also A270533 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010767 | OEIS-A270530

a(n) = Sum_{k=0..n}((binomial(2*k,k)/(k+1)*binomial(2*n+2,n-k))).

Conjecture: 2*n*(2*n+3)*(n+1)*a(n) -n*(77*n^2+27*n-4)*a(n-1) +(549*n^3-987*n^2+686*n-168)*a(n-2) -20*(2*n-3)*(43*n^2-104*n+70)*a(n-3) +500*(2*n-5)*(n-2)*(2*n-3)*a(n-4)=0. - _R. J. Mathar_, Jun 07 2016 Conjecture: 2*n*(2*n+3)*(n+3)*(n+1)*a(n) -n*(57*n^3+228*n^2+107*n+8)*a(n-1) +4*(2*n-1) *(33*n^3+99*n^2-88*n+36)*a(n-2) -100*(n-1)*(2*n-1)*(2*n-3)*(n+4)*a(n-3)=0. - _R. J. Mathar_, Jun 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Mar 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010768 | OEIS-A270533

Number of ordered ways to write n = x^4 + x^3 + y^2 + z*(3z-1)/2, where x and y are nonnegative integers, and z is an integer.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 20, 22, 70, 87, 167, 252, 388, 562, 636, 658, 873, 2598, 14979, 18892, 20824. See also A270516 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010769 | OEIS-A270559

Number of ordered ways to write n as x^4 + x^3 + y^2 + z*(z+1)/2, where x, y and z are integers with x nonzero, y nonnegative and z positive.

Conjecture: (i) a(n) > 0 for all n > 0. In other words, for each n = 1,2,3,... there are integers x and y such that n-(x^4+x^3+y^2) is a positive triangular number. In contrast, the author conjectured in A262813 that any positive integer can be expressed as the sum of a

nonnegative cube, a square and a positive triangular number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010770 | OEIS-A270566

Number of ordered ways to write n as x^4 + y*(3y+1)/2 + z*(7z+1)/2, where x, y and z are integers with x nonnegative.

Conjecture: a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 24, 47, 63, 78, 79, 143, 153, 325, 494, 949, 1079, 3328, 4335, 5609, 7949, 7967, 8888, 9665. Conjecture verified for n up to 10^11. - _Mauro Fiorentini_, Jul 24 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010771 | OEIS-A270592

List of pairs: smallest pair of successive primes p & q such that p+1 & q+1 have n distinct prime divisors in common.

The difference between the pairs is conjectured to be the primorials sequence: A002110. q(n)-p(n)=prime(n)# (conjectured) prime(n)#|(p(n)+1) (conjectured) prime(n)#|(q(n)+1) (conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alon Kellner_, Mar 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010772 | OEIS-A270594

Number of ordered ways to write n as the sum of a triangular number, a positive square and the square of a generalized pentagonal number (A001318).

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 3, 21, 24, 48, 90, 138, 213, 283, 462, 468, 567, 573, 1998, 2068, 2488, 2687, 5208, 5547, 5638, 6093, 6492, 6548, 6717, 7538, 7731, 8522, 14763, 16222, 17143, 24958, 26148. See also A270566 for a similar conjecture involving four powers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010773 | OEIS-A270616

Number of ordered ways to write n as the sum of a positive square, the square of a triangular number, and a generalized pentagonal number (A001318).

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 8, 29, 34, 5949, 10913. See also A270566 and A270594 for more similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010774 | OEIS-A270643

The sequence a of 1's and 2's starting with (1,2,2,1) such that a(n) is the length of the (n+3)rd run of a.

Conjecture: a(n) = A270642(n+1). - _R. J. Mathar_, Jun 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010775 | OEIS-A270647

The sequence a of 1's and 2's starting with (2,2,1,2) such that a(n) is the length of the (n+3)rd run of a.

Conjecture: a(n) = A270643(n+1). - _R. J. Mathar_, Jun 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010776 | OEIS-A270661

a(n) = Sum_{k=0..n}((binomial(2*k,k)*Sum_{i=0..n-k}(binomial(k+1,n-k-i)*binomial(k+i,k)))/(k+1)^2)*(n+1).

Conjecture: (n+1)*a(n) -4* n*a(n-1) +(-13*n+33)*a(n-2) +2*(8*n-17)*a(n-3) +(19*n-69)*a(n-4) +4*(-4*n+15)*a(n-5) +7*(-n+5)*a(n-6) +2*(2*n-13)*a(n-7)=0. - _R. J. Mathar_, Jun 07 2016 Conjecture: (n+1)*(n^2-5*n+3)*a(n) -2*n*(3*n^2-16*n+14)*a(n-1) +2*(n+4)*(n-3)*a(n-2) +2*(5*n^3-31*n^2+39*n+9)*a(n-3) +(-n^3+2*n^2+21)*a(n-4) -2*(2*n-9)*(n^2-3*n-1)*a(n-5)=0. - _R. J. Mathar_, Jun 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Mar 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010777 | OEIS-A270668

Triangle read by rows: The number of domino tilings of the (2n+1) X (2m+1) board with a central free square.

Conjectured g.f. for column 3: (-1 -4*x +543*x^2 -6238*x^3 +17032*x^4 -6238*x^5 +543*x^6 -4*x^7 -x^8) / ((x-1) *(x^2-7*x+1) *(x^2-23*x+1) *(x^4 -161*x^3 +576*x^2 -161*x +1)). - _R. J. Mathar_, Mar 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Mar 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010778 | OEIS-A270681

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 190", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 21 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010779 | OEIS-A270682

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 190", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 254 also generates this sequence. - _Lars Blomberg_, Jun 03 2016 Conjecture: a(n) = 4^n + 4*2^n - 8, n>1. - _Lars Blomberg_, Jun 03 2016 Conjectures from _Colin Barker_, Jun 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010780 | OEIS-A270683

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 190", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 21 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010781 | OEIS-A270684

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 190", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 21 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010782 | OEIS-A270705

Number of ordered ways to write n as x^2*pen(x) + pen(y) + pen(z) with pen(x) = x*(3x+1)/2 and pen(y) <= pen(z), where x, y and z are integers ("pen" stands for "pentagonal").

Conjecture: (i) Any natural number can be written as a*x^2*pen(x) + b*pen(y) + c*pen(z) with x, y and z integers, provided that (a,b,c) is among the following ordered triples: (j,1,k) (j = 1,2; k = 1,2,3,4), (1,2,3), (3,1,4) and (4,1,3). See also A270594 and A270706 for other similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010783 | OEIS-A270706

Number of ordered ways to write n as x^2*T(x) + y^2 + T(z), where x, y and z are integers with x nonzero, y positive and z nonnegative, and T(m) denotes the triangular number m*(m+1)/2.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 3, 90, 438, 480, 7108. See also A270594 and A270705 for some other similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010784 | OEIS-A270724

a(n) = ((n+2)/2)*Sum_{k=0..n/2} (Sum_{i=0..n-2*k} (binomial(k+1,n-2*k-i)*binomial(k+i,k))/(k+1)*C(k)), where C(k) is Catalan numbers.

Conjecture: (n+2)*a(n) +(-n-2)*a(n-1) +(-7*n+6)*a(n-2) +10*a(n-3) +(13*n-32)*a(n-4) +(5*n-32)*a(n-5) +(-11*n+52)*a(n-6) +4*(-n+6)*a(n-7) +4*(n-7)*a(n-8)=0. - _R. J. Mathar_, Oct 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Mar 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010785 | OEIS-A270751

(r,1)-greedy sequence, where r(k) = 1/(k*phi) and phi = golden ratio.

Conjecture: a(n) = A270584(n-1) for n>1. - _R. J. Mathar_, Jun 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010786 | OEIS-A270776

Smallest non-Wieferich prime to base n, i.e., smallest prime p such that n^(p-1) != 1 (mod p^2).

The ABC-(k, Epsilon) conjecture with k >= 2 and Epsilon > 0 such that 1/(1/Epsilon + 1) + 1/k <= log(2)/(24*log(a)) implies that a(n) exists for all n (cf. Broughan, 2006; theorem 5.6).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Mar 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010787 | OEIS-A270803

Formal inverse of Thue-Morse sequence A010060.

Conjecture: a(n) = A151666(floor((n+1)/2)) for n > 0. - _Georg Fischer_, Nov 29 2022 Conjecture confirmed. See the attached file. The idea is to guess the automaton for this sequence (in base 2) and then verify that it satisfies the identities in the Gawron-Ulas paper. Next we make a (base 2) automaton for A151666. This is easy because it is just the numbers consisting only of 0 and 1 in base 4. Then finally we assert Fischer's identity and Walnut returns TRUE. - _Jeffrey Shallit_, Nov 30 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010788 | OEIS-A270807

Trajectory of 1 under the map n -> n + n/gpf(n) + 1 (see A269304).

Cody M. Haderlie (see A269304) conjectures that the trajectory of any initial value will eventually merge with this sequence. The trajectory of 2, for example, begins 2, 4, 7, 9, 13, 15, 19, 21, 25, ... and from 7 on coincides with this sequence. See A271418.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010789 | OEIS-A270916

(r,1)-greedy sequence, where r(k) = 1/(k*sqrt(2)).

Conjecture: a(n) = A270582(n-1). - _R. J. Mathar_, Jun 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010790 | OEIS-A270928

Number of ways to write n = x*(x-1)/2 + y*(y-1)/2 + z*(z-1)/2, where 0 < x <= y <= z, and one of x, y, z is prime.

Conjecture: (i) a(n) > 0 for n > 0 with the only exception n = 15^2 = 225. Also, a(n) = 1 only for n = 1, 2, 4, 5, 6, 8, 11, 14, 15, 18, 20, 29, 36, 50, 53, 60, 62, 96, 117, 119, 218, 411, 540, 645, 1125, 1590, 2346, 4068. See also A270966 for a similar conjecture involving (p-1)^2 with p prime. The conjecture that a(n) > 0 except for n = 225 appeared as Conjecture 1.2(i) of the author's JNT paper in the links.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010791 | OEIS-A270934

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 221", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 26 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010792 | OEIS-A270935

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 221", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 253 also generates this sequence. - _Lars Blomberg_, Jun 06 2016 Conjecture: a(n) = 4*4^n - 4*2^n - 4 = A269907(n), n>1. - _Lars Blomberg_, Jun 06 2016 Conjectures from _Colin Barker_, Jun 07 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010793 | OEIS-A270936

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 221", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 26 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010794 | OEIS-A270937

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 221", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 26 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010795 | OEIS-A270966

Number of ways to write n as x^2 + y^2 + z*(3z+1)/2, where x, y and z are integers with 0 <= x <= y such that x or y has the form p-1 with p prime.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 49, 608. See also A270928 for a similar conjecture involving T(p-1) = p*(p-1)/2 with p prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010796 | OEIS-A270969

Number of ways to write n as w^4 + x^2 + y^2 + z^2, where w, x, y and z are nonnegative integers with x <= y <= z.

Conjecture: For each triple (a,b,c) = (1,2,11),(1,2,12),(1,2,13),(2,3,5), any natural number can be written as w^4+a*x^2+b*y^2+c*z^2 with w,x,y,z integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010797 | OEIS-A270985

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 238", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jun 07 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010798 | OEIS-A270994

a(n) = 9454129 + 11184810*n.

Adamczewski disproves the above conjecture in linked article, using the case for n = 162178349052694. - _Michael De Vlieger_, Aug 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Mar 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010799 | OEIS-A271004

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 245", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 28 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010800 | OEIS-A271005

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 245", based on the 5-celled von Neumann neighborhood.

It is conjectured that Rule 501 also produces this sequence. It would be nice to have a proof. Conjecture: a(n) = 4*4^n - 6*2^n, n>2. - _Lars Blomberg_, Jun 08 2016 Conjectures from _Colin Barker_, Jun 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010801 | OEIS-A271006

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 245", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 28 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010802 | OEIS-A271007

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 245", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 28 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010803 | OEIS-A271017

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 251", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 28 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010804 | OEIS-A271026

Number of ordered ways to write n as x^7 + y^4 + z^3 + w*(3w+1)/2, where x, y, z are nonnegative integers, and w is an integer.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 47, 61, 62, 112, 175, 448, 573, 714, 1073, 1175, 1839, 2167, 8043, 13844. See also A266968 for a related conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010805 | OEIS-A271035

Number of 3 X 3 X 3 triangular 0..n arrays with some element less than a w, nw or ne neighbor exactly once.

Conjectures from _Colin Barker_, Jan 30 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010806 | OEIS-A271036

Number of 4 X 4 X 4 triangular 0..n arrays with some element less than a w, nw or ne neighbor exactly once.

Conjectures from _Colin Barker_, Jan 30 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Mar 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010807 | OEIS-A271051

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 253", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010808 | OEIS-A271052

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 253", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010809 | OEIS-A271053

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 253", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010810 | OEIS-A271055

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 259", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = 4*4^n - 4*2^n - 12, n>1. - _Lars Blomberg_, Jun 09 2016 Conjectures from _Colin Barker_, Dec 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010811 | OEIS-A271060

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 261", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Dec 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010812 | OEIS-A271061

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 261", based on the 5-celled von Neumann neighborhood.

It is conjectured that Rules 269, 277, 285, 293, 301, 309, 317, 325, 333, 341, 349, 357, 365, 373, 381, 413, 445, 477, 509, 645, 653, 661, 669, 677, 685, 693, 701, 709, 717, 725, 733, 741, 749, 757 and 765 also produces this sequence. It would be nice to have a proof. Conjecture: a(n) = 4*4^n - 4*2^n, n>0. - _Lars Blomberg_, Jun 09 2016 Conjectures from _Colin Barker_, Dec 01 2016: (Start) Conjectured e.g.f.: (1 - 2*exp(2*x))^2. - _Stefano Spezia_, Mar 12 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010813 | OEIS-A271062

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 261", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Dec 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010814 | OEIS-A271064

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 261", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Dec 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010815 | OEIS-A271076

Number of ordered ways to write n as u^5 + v^4 + x^3 + 2*y^3 + 3*z^3, where u, v , x, y and z are nonnegative integers with v > 0.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 8, 9, 10, 11, 13, 14, 15, 16, 23, 24, 38, 39, 61, 76, 77, 104, 118, 188, 214, 229. Conjectures (i), (ii) and (iii) verified for n up to 10^11 for all polynomials. - _Mauro Fiorentini_, Sep 20 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010816 | OEIS-A271091

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 275", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 30 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010817 | OEIS-A271092

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 275", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = 4*4^n - 4*2^n - 8, n>1. - _Lars Blomberg_, Jun 10 2016 Conjectures from _Colin Barker_, Jun 10 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010818 | OEIS-A271093

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 275", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 30 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010819 | OEIS-A271094

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 275", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 30 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010820 | OEIS-A271099

Number of ordered ways to write n as u^3 + v^3 + 2*x^3 + 2*y^3 + 3*z^3, where u, v, x, y and z are nonnegative integers with u <= v and x <= y.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 1, 10, 14, 15, 17, 22, 38, 39, 45, 47, 50, 52, 76, 102, 103, 188, 295, 366, 534. This conjecture is stronger than the classical Waring problem on sums of k-th powers. Concerning parts (i) and (ii) of the conjecture, we note that 1+1+2+2+3 = 9 = g(3), 1+1+2+2+3+3+7 = 19 = g(4) and 1+1+1+1+2+4+6+9+12 = 37 = g(5). We have verified that a(n) > 0 for all n = 0..10^6, and that part (ii) of the conjecture holds for n up to 10^5. Concerning part (iii) for k = 6, we conjecture that any natural number can be written as x(1)^6+x(2)^6+x(3)^6+x(4)^6+x(5)^6+3*x(6)^6+5*x(7)^6+6*x(8)^6+10*x(9)^6+18*x(10)^6+26*x(11)^6 with x(1),x(2),...,x(11) nonnegative integers. Note that 1+1+1+1+1+3+5+6+10+18+26 = 73 = g(6). - _Zhi-Wei Sun_, Mar 31 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010821 | OEIS-A271106

Number of ordered ways to write n as x^6 + 3*y^3 + z^3 + w*(w+1)/2, where x and y are nonnegative integers, and z and w are positive integers.

Conjecture: a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 3, 4, 6, 9, 13, 16, 20, 21, 24, 25, 44, 50, 51, 65, 84, 189, 290, 484, 616, 664, 680, 917, 1501, 1639, 3013. Based on our computation, we also formulate the following general conjecture. General Conjecture: Let T(w) = w*(w+1)/2. We have {P(x,y,z,w): x,y,z,w = 0,1,2,...} = {0,1,2,...} for any of the following polynomials P(x,y,z,w): x^3+y^3+c*z^3+T(w) (c = 2,3,4,6), x^3+y^3+c*z^3+2*T(w) (c = 2,3), x^3+b*y^3+3z^3+3*T(w) (b = 1,2), x^3+2y^3+3z^3+w(5w-1)/2, x^3+2y^3+3z^3+w(5w-3)/2, x^3+2y^3+c*z^3+T(w) (c = 2,3,4,5,6,7,12,20,21,34,35,40), x^3+2y^3+c*z^3+2*T(w) (c = 3,4,5,6,11), x^3+2y^3+c*z^3+w^2 (c = 3,4,5,6), x^3+2y^3+4z^3+w(3w-1)/2, x^3+2y^3+4z^3+w(3w+1)/2, x^3+2y^3+4z^3+w(2w-1), x^3+2y^3+6z^3+w(3w-1)/2, x^3+3y^3+c*z^3+T(w) (c = 3,4,5,6,10,11,13,15,16,18,20), x^3+3y^3+c*z^3+2*T(w) (c

= 5,6,11), x^3+4y^3+c*z^3+T(w) (c = 5,10,12,16), x^3+4y^3+5z^3+2*T(w), x^3+5y^3+10z^3+T(w), 2x^3+3y^3+c*z^3+T(w) (c = 4,6), 2x^3+4y^3+8z^3+T(w), x^4+y^3+3z^3+w(3w-1)/2, x^4+y^3+c*z^3+T(w) (c = 2,3,4,5,7,12,13), x^4+y^3+c*z^3+2*T(w) (c = 2,3,4,5), x^4+y^3+2z^3+w^2, x^4+y^3+4z^3+2w^2, x^4+2y^3+c*z^3+T(w) (c = 4,5,12), x^4+2y^3+3z^3+2*T(w), 2x^4+y^3+2z^3+w(3w-1)/2, 2x^4+y^3+c*z^3+T(w) (c = 1,2,3,4,5,6,10,11), 2x^4+y^3+c*z^3+2*T(w) (c = 2,3,4), 2x^4+2y^3+c*z^3+T(w) (c = 3,5), 3x^4+y^3+c*z^3+T(w) (c = 1,2,3,4,5,11), 3x^4+y^3+2z^3+2*T(w), 3x^4+y^3+2z^3+w^2, 3x^4+y^3+2z^3+w(3w-1)/2, 4x^4+y^3+c*z^3+T(w) (c = 2,3,4,6), 4x^4+y^3+2z^3+2*T(w), 5x^4+y^3+c*z^3+T(w) (c = 2,4), a*x^4+y^3+2z^3+T(w) (a = 6,20,28,40), 6x^4+y^3+2z^3+2*T(w), 6x^4+y^3+2z^3+w^2, a*x^4+y^3+3z^3+T(w) (a = 6,8,11), 8x^4+2y^3+4z^3+T(w), x^5+y^3+c*z^3+T(w) (c = 2,3,4), x^5+2y^3+c*z^3+T(w) (c = 3,6,8), 2x^5+y^3+4z^3+T(w), 3x^5+y^3+2z^3+T(w), 5x^5+y^3+c*z^3+T(w) (c = 2,4), x^6+y^3+3z^3+T(w), x^7+y^3+4z^3+T(w), x^4+2y^4+z^3+w^2, x^4+2y^4+2z^3+T(w), x^4+b*y^4+z^3+T(w) (b = 2,3,4), 2x^4+3y^4+z^3+T(w), a*x^5+y^4+z^3+T(w) (a = 1,2), x^5+2y^4+z^3+T(w). The polynomials listed in the general conjecture should exhaust all those polynomials P(x,y,z,w) = a*x^i+b*y^j+c*z^k+w*(s*w+/-t)/2 with {P(x,y,z,w): x,y,z,w = 0,1,2,...} = {0,1,2,...}, where a,b,c,s > 0, 0 <= t <= s, s == t (mod 2), i >= j >= k >= 3, a <= b if i = j, and b <= c if j = k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010822 | OEIS-A271126

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 289", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = (117*4^(n-1) - 76*3^(n-4) - 171*2^(n-2) - 432)/9, n>5. - _Lars Blomberg_, Jun 10 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010823 | OEIS-A271169

Number of ordered ways to write n as s^5 + t^5 + 2*u^5 + 3*v^5 + 4*w^5 + 5*x^5 + 7*y^5 + 14*z^5, where s,t,u,v,w,x,y,z are nonnegative integers with s <= t.

Conjecture: a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 1, 2602. For k = 2,3,4,... define s(k) as the smallest positive integer s such that {a(1)*x(1)^k+...+a(s)*x(s)^k: x(1),...,x(s) = 0,1,2,...} = {0,1,2,...} for some positive integers a(1), ..., a(s), and t(k) as the least positive integer t such that {a(1)*x(1)^k+...+a(t)*x(t)^k: x(1),...,x(t) = 0,1,2,...} = {0,1,2,...} for some positive integers a(1), ..., a(t) with a(1)+...+a(t) = g(k), where g(.) is given by A002804. Then s(k) <= t(k) <= g(k). Part (iii) of the conjecture in A271099 implies that t(k) <= 2k-1 for k > 2. It is easy to see that s(2) = t(2) = 4. Our computation suggests that s(3) = t(3) = 5, s(4) = t(4) = 7, s(5) = t(5) = 8 (which is smaller than 2*5-1), and s(6) = t(6) = 10. We conjecture that s(k) = t(k) for any integer k > 1, and that each natural number can be written as x(1)^6+x(2)^6+x(3)^6+2*x(4)^6+3*x(5)^6+5*x(6)^6+6*x(7)^6+10*x(8)^6+18*x(9)^6+26*x(10)^6, where x(1),x(2),...,x(10) are nonnegative integers. Note that 1+1+1+2+3+5+6+10+18+26 = 73 = g(6). We also conjecture that any natural number can be written as s^5+t^5+2*u^5+3*v^5+4*w^5+6*x^5+8*y^5+12*z^5, with s,t,u,v,w,x,y,z nonnegative integers. Note that 1+1+2+3+4+6+8+12 = 37 = g(5). - _Zhi-Wei Sun_, Apr 04 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010824 | OEIS-A271217

Number of symmetric reduced rearrangement maps.

Conjecture: (-2*n+9)*a(n) -4*a(n-1) +(2*n-3)*(2*n-7)*a(n-2) -4*a(n-3) +2*(2*n-5)*(n-4)*a(n-4)=0. - _R. J. Mathar_, Jan 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Burns_, Apr 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010825 | OEIS-A271218

Number of symmetric linked diagrams with n links and no simple link.

a(n) ~ 2^(-1/2) * e^(-5/8) * (2n/e)^(n/2) * e^(sqrt(n/2)) (conjectured). a(n)/a(n-1) ~ sqrt(2n) (conjectured). a(n)/A047974(n) ~ 1/sqrt(e) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Burns_, Apr 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010826 | OEIS-A271227

Number of solutions to y^2 == x^3 + 17 (mod p) as p runs through the primes.

Conjecture [WL]:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Apr 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010827 | OEIS-A271228

P-defects p - N(p) of the elliptic curve y^2 = x^3 + 17 for primes p, where N(p) is the number of solutions modulo prime p.

See A271227 for details and the conjecture for a(n) if prime(n) == 1 (mod 3). a(n) is negative for the 1 (mod 3) primes 7, 13, 19, 31, 37, 43, 97, 103, 109, 127, 151, 157, 163, 193, 223, 229, 241, 271, 277, 307, 313, 331, ... and positive for the primes 61, 67, 73, 79, 139, 181, 199, 211, 283, 337, 349, ... See A271227 for a comment on the conjectured three types I, II, and III of 1 (mod 3) primes. All three types appear for primes with negative as well as positive a(n) values. Conjecture [WL]: For prime(n) = A002476(m) (a prime == 1 (mod 3)) one has a(n) = + or - sqrt(4*prime(n)) - 3*q(m)^2), with three alternative cases for q(m)^2, namely (2*B(m))^2, (A(m) - B(m))^2 and (A(m) + B(m))^2, where A(m) = A001479(m+1) and B(m) = A001480(m+1), m >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Apr 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010828 | OEIS-A271237

Number of ordered ways to write n as u^3 + 2*v^3 + 3*x^3 + 4*y^3 + 5*z^3, where u, v, x, y and z are nonnegative integers.

Conjecture: We have {u^3+a*v^3+b*x^3+c*y^3+d*z^3: u,v,x,y,z = 0,1,2,...} = {0,1,2,...} whenever (a,b,c,d) is among the following 32 quadruples: (1,2,2,3), (1,2,2,4), (1,2,3,4), (1,2,4,5), (1,2,4,6), (1,2,4,9), (1,2,4,10), (1,2,4,11), (1,2,4,18), (1,3,4,6), (1,3,4,9), (1,3,4,10), (2,2,4,5), (2,2,6,9), (2,3,4,5), (2,3,4,6), (2,3,4,7), (2,3,4,8), (2,3,4,9), (2,3,4,10), (2,3,4,12), (2,3,4,15), (2,3,4,18), (2,3,5,6), (2,3,6,12), (2,3,6,15), (2,4,5,6), (2,4,5,8), (2,4,5,9), (2,4,5,10), (2,4,6,7), (2,4,7,10). If {m*u^3+a*v^3+b*x^3+c*y^3+d*z^3: u,v,x,y,z = 0,1,2,...} = {0,1,2,...} with 1 <= m <= a <= b <= c <= d, then m = 1, and we can show that (a,b,c,d) must be among the 32 quadruples listed in the conjecture (cf. Theorem 1.2 of the linked 2017 paper). Conjecture verified for all the 32 quadruples up to 10^11. - _Mauro Fiorentini_, Jul 09 2023 See also A271099 and A271169 for conjectures refining Waring's problem. We also conjecture that if P(u,v,x,y,z) is one of the four polynomials u^6+v^3+2*x^3+4*y^3+5*z^3 and a*u^6+v^3+2*x^3+3*y^3+4*z^3 (a = 5,8,12) then any natural number can be written as P(u,v,x,y,z) with u,v,x,y,z nonnegative integers. - _Zhi-Wei Sun_, Apr 06 2016 Conjecture verified for all the 4 polynomials up to 10^11. - _Mauro Fiorentini_, Jul 09 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010829 | OEIS-A271254

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 323", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = 4*4^n - 10*2^n + 12, n>1. - _Lars Blomberg_, Jun 14 2016 Conjectures from _Colin Barker_, Dec 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010830 | OEIS-A271256

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 323", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 03 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010831 | OEIS-A271286

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 337", based on the 5-celled von Neumann neighborhood.

It is conjectured that Rule 505 also produces this sequence. It would be nice to have a proof. Conjectures from _Colin Barker_, May 04 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010832 | OEIS-A271325

Number of ordered ways to write n as x^3 + y^2 + z*(3z+1), where x, y and z are integers with x positive and y nonnegative.

Based on our computation, we propose the following general conjecture (which extends the conjectures in A262813 and A270469). Conjecture: Let T(x) = x*(x+1)/2 and pen(x) = x*(3x+1)/2. Every natural number can be written as P(x,y,z) with x a nonnegative integer and y and z integers, where P(x,y,z) is any of the following cubic polynomials: x^3+T(y)+z^2, a*x^3+T(y)+pen(z) (a = 1,2,3,4), x^3+T(y)+z*(5z+1)/2, x^3+T(y)+z*(3z+r) (r = 1,2), x^3+T(y)+z*(7z+3)/2, x^3+T(y)+z*(9z+j)/2 (j = 5,7), x^3+T(y)+z*(5z+r) (r = 2,3), x^3+T(y)+2z*(3z+r) (r = 1,2), x^3+T(y)+z*(6z+5), x^3+T(y)+z*(13z+j)/2 (j = 3,7,9), x^3+T(y)+z*(7z+k) (k = 2,6), a*x^3+y^2+pen(z) (a = 1,2,3,4), x^3+y^2+z*(5z+3)/2, x^3+y^2+2*pen(z), x^3+2*T(y)+pen(z), x^3+2*T(y)+z(5z+j)/2 (j = 1,3), a*x^3+2*T(y)+z*(3z+2) (a = 1,2,3), x^3+2*T(y)+z*(7z+3)/2, x^3+4*T(y)+pen(z), x^3+2y^2+pen(z), x^3+pen(y)+c*pen(z) (c = 1,2,3,4), x^3+b*pen(y)+z*(5z+j)/2 (b = 1,2; j = 1,3), x^3+pen(y)+z*(7z+k)/2 (k = 1,3,5), x^3+pen(y)+z*(4z+j) (j = 1,3), x^3+pen(y)+z*(9z+5)/2, a*x^3+pen(y)+z*(9z+r)/2 (a = 1,2; r = 1,7), x^3+pen(y)+z*(5z+r) (r = 1,2,3,4), a*x^3+pen(y)+z*(11z+9)/2 (a = 1,2), x^3+pen(y)+2z*(3z+2),x^3+pen(y)+z*(13z+11)/2, x^3+pen(y)+z*(7z+k) (k = 4,5,6), x^3+pen(y)+3z*(5z+3)/2, x^3+pen(y)+z*(15z+11)/2, x^3+pen(y)+z*(8z+7), x^3+pen(y)+z*(11z+7), x^3+2*pen(y)+z*(7z+j)/2 (j = 1,5), x^3+2*pen(y)+3*pen(z), x^3+2*pen(y)+z*(4z+1), x^3+2*pen(y)+z*(7z+2), x^3+y*(5y+j)/2+z*(7z+k)/2 (j = 1,3; k = 3,5), x^3+y*(5y+3)/2+z*(9z+7)/2, x^3+y*(3y+2)+z*(4z+1), x^3+y*(3y+2)+z*(5z+1)/2, x^3+y*(7y+3)/2+z*(7z+5)/2, 2x^3+T(y)+z*(5z+3)/2, 2x^3+T(y)+z*(3z+r) (r = 1,2), 2x^3+T(y)+z*(5z+4), 2x^3+2*T(y)+z*(5z+3)/2, 2x^3+3*T(y)+pen(z), 2x^3+y^2+2*pen(z), 2x^3+pen(y)+pen(z), a*x^3+pen(y)+3*pen(z) (a = 2,3,4), a*x^3+pen(y)+z*(7z+5)/2 (a = 2,3,4), 2x^3+pen(y)+z*(5z+k) (k = 1,3), 2x^3+y*(5y+3)/2+z*(7z+5)/2, 2x^3+2*pen(y)+z*(3z+2), 2x^3+2*pen(y)+z*(7z+5)/2, 2x^3+y*(3y+2)+z*(4z+3), 3x^3+pen(y)+z*(7z+3)/2, 4x^3+y^2+z*(5z+1)/2, 4x^3+pen(y)+z*(4z+3). The listed ternary polynomials in the conjecture should exhaust all those P(x,y,z) = a*x^3+y*(s*y+t)/2+z*(u*z+v)/2 with a,s,u > 0, 0 <= t <= s, 0 <= v <= u, s == t (mod 2), u == v (mod 2), and (s-2t)*(u-2v) nonzero, such that any natural number can be written as P(x,y,z) with x a nonnegative integer and y and z integers. Note that those numbers y*(2y+1) with y integral are just triangular numbers. Conjecture verified for all polynomials up to 10^11. - _Mauro Fiorentini_, Aug 03 2023 See also A271106 for another general conjecture on universal sums.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010833 | OEIS-A271356

Difference sequence of the sequence A271354 of the increasing sequence of products of two distinct Fibonacci numbers greater than 1.

Conjecture: every term except the first is a Fibonacci number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010834 | OEIS-A271365

Number of ordered ways to write n as u^2 + v^3 + x^4 + y^5 + z^6, where u is a positive integer, and v, x, y, z are nonnegative integers.

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 8, 15, 16, 23, 24, 56. Moreover, the only positive integers not represented by u^2+v^3+x^4+y^5 (u > 0 and v,x,y >= 0) are 8, 15, 23, 55, 62, 71, 471, 478, 510, 646, 806, 839, 879, 939, 1023, 1063, 1287, 2127, 5135, 6811, 7499, 9191, 26471.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010835 | OEIS-A271381

Number of ordered ways to write n as u^2 + v^2 + x^3 + y^3, where u, v, x, y are nonnegative integers with 2 | u*v, u <= v and x <= y.

Conjecture: (i) a(n) > 0 except for n = 23, and a(n) = 1 only for n = 0, 3, 4, 7, 11, 12, 15, 19, 23, 30, 78, 203, 219.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010836 | OEIS-A271480

Dimension of n-qubit subspace H^{MPS}_{2,n} for bond dimension 2.

Conjectures from _Colin Barker_, Apr 26 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010837 | OEIS-A271485

Maximal term of TRIP-Stern sequence of level n corresponding to permutation triple (e,13,e).

Conjectures from _Colin Barker_, Apr 16 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010838 | OEIS-A271487

Maximal term of TRIP-Stern sequence of level n corresponding to permutation triple (e,13,132).

Conjectures from _Colin Barker_, Apr 16 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010839 | OEIS-A271489

Maximal term of TRIP-Stern sequence of level n corresponding to permutation triple (e,132,e).

Conjectures from _Lars Blomberg_, Jan 08 2018: (Start)
Conjectures from _Colin Barker_, Jan 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010840 | OEIS-A271510

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with x >= y >= 0, z >= 0 and w >= 0 such that x^2 + 8*y^2 + 16*z^2 is a square.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 7, 23, 71, 77, 105, 191, 215, 311, 335, 2903, 4^k*q (k = 0,1,2,... and q = 6, 15, 47, 138). This conjecture is much stronger than Lagrange's four-square theorem. It is apparent that a(m^2*n) >= a(n) for all m,n = 1,2,3,.... See also A271513 and A271518 for related conjectures. Conjectures (i), including the "a(n) = 1" part, (ii), (iii), and (iv) have been verified for n <= 10^9. - _Mauro Fiorentini_, Jun 19 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010841 | OEIS-A271513

Number of ordered ways to write n as w^2 + x^2 + y^2 + z^2 with 3*x^2 + 4*y^2 + 9*z^2 a square, where w, x, y and z are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 3, 11, 23, 43, 47, 67, 83, 107, 155, 323, 683, 803, 4^k*m (k = 0,1,2,... and m = 22, 38). [Conjecture verified for all natural numbers up to 10^9. - _Mauro Fiorentini_, Jul 04 2024] (ii) Any natural number can be written as w^2 + x^2 + y^2 + z^2 with x, y, z integers and a*x^2 + b*y^2 + c*z^2 a square, whenever (a,b,c) is among the following triples: (1,3,12), (1,3,18), (1,3,21), (1,3,60), (1,5,15), (1,8,24), (1,12,15), (1,24,56), (1,24,72), (1,48,72), (1,48,168), (1,120,180), (1,192,288), (1,280,560), (3,9,13), (4,5,12), (4,5,60), (4,9,60), (4,12,21), (4,12,45), (4,12,69), (4,12,93), (4,12,237), (4,21,24), (4,21,36), (4,21,504), (4,24,93), (4,28,77), (4,45,120), (4,45,540), (4,45,600), (5,36,40), (7,9,126), (7,9,588), (8,16,73), (8,16,97), (8,49,112), (9,13,27), (9,16,24), (9,19,36), (9,21,91), (9,24,232), (9,28,63), (9,40,45), (9,40,56), (9,40,120), (9,45,115),(9,45,235), (12,13,24), (12,13,36), (12,36,37), (12,36,133), (13,36,72), (13,36,108), (15,24,25), (15,49,105), (16,17,48), (16,20,45), (16,21,84), (16,33,72), (16,33,176), (16,45,180), (16,48,57), (16,48,105), (16,48,233), (16,48,249), (19,45,57), (19,45,180), (21,25,35), (21,25,75), (21,28,36), (21,28,60), (21,43,105), (21,100,105),(24,25,72), (24,25,120), (24,48,97), (24,81,184), (24,120,145), (25,36,75), (25,40,56), (25,45,51), (25,45,99), (25,48,96), (25,48,144), (25,54,90), (25,75,81), (25,80,184), (25,96,120), (25,200,216), (28,33,36), (28,36,77), (28,72,189), (32,64,73), (33,36,220), (33,48,144), (33,72,256), (33,88,144), (36,45,100), (36,45,172), (37,81,243), (40,81,120), (40,81,240), (41,64,256), (45,48,76), (48,144,177), (49,56,64), (49,63,72), (55,141,165), (57,64,192), (60,105,196), (64,65,160), (72,73,144), (81,160,240), (85,140,196), (105,112,144), (112,144,153), (136,144,153), (144,145,240), (144,160,225),(148,189,252), (175,189,225). [Conjecture verified for all triples and all natural numbers up to 10^9. - _Mauro Fiorentini_, Jul 04 2024] This conjecture is stronger than Lagrange's four-square theorem. Moreover, there are many other suitable triples (a,b,c) for our purpose not listed in part (ii) of the conjecture. If a, b and c are positive integers such that any natural number can be written as w^2 + x^2 + y^2 + z^2 with x, y, z integers and

a*x^2 + b*y^2 + c*z^2 a square, then one of a+b+c, 4*a+b+c, a+4*b+c and a+b+4*c must be a square since 2^2 + 1^2 + 1^2 + 1^2 is the unique way to express 7 as a sum of four squares. See also A271510 and A271518 for related conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010842 | OEIS-A271518

Number of ordered ways to write n as w^2 + x^2 + y^2 + z^2 with x + 3*y + 5*z a square, where w, x, y and z are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 4^k*6 (k = 0,1,2,...), 16^k*m (k = 0,1,2,... and m = 5, 7, 8, 31, 43, 61, 116). See also A271510 and A271513 for related conjectures. Qing-Hu Hou at Tianjin Univ. has verified a(n) > 0 and parts (ii) and (iii) of the above conjecture for n up to 10^9. - _Zhi-Wei Sun_, Dec 04 2016 The conjecture that a(n) > 0 for all n = 0,1,2,... is called the 1-3-5-Conjecture and the author has announced a prize of 1350 US dollars for its solution. - _Zhi-Wei Sun_, Jan 17 2017 The 1-3-5 conjecture was finally proved by António Machiavelo and Nikolaos Tsopanidis in a JNT paper published in 2021. This is a great achivement! - _Zhi-Wei Sun_, Mar 31 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010843 | OEIS-A271578

Magic sums of 4 X 4 magic squares composed of primes.

I conjecture that every even number greater than 152 belongs to this sequence. If conjecture is true, a(n) = A005843(n+65) for n >= 12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Apr 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010844 | OEIS-A271608

Number of ordered ways to write n as pen(u) + pen(v) + pen(x) + pen(y) + pen(z) with u,v,x,y,z nonnegative integers such that u + 2*v + 4*x + 5*y + 6*z is a pentagonal number, where pen(k) denotes the pentagonal number k*(3k-1)/2.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 2, 4, 5, 7, 9, 21, 22, 43. Also, every n = 0,1,2,... can be written as pen(u) + pen(v) + pen(x) + pen(y) + pen(z) with u,v,x,y,z nonnegative integers such that 3*u + 5*v + 11*x + 16*y + 19*z is also a pentagonal number. As conjectured by Fermat and proved by Cauchy, each natural number can be written as the sum of five pentagonal numbers. See also A271510, A271513, A271518 and A271644 for some similar conjectures refining Lagrange's four-square theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010845 | OEIS-A271620

First row (row 0) of the Sprague-Grundy values of "3-pile Sharing Nim".

Ho conjectures that the sequence may be bounded. The highest value in the first thousand entries is a(24)=12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Howard A. Landman_, Apr 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010846 | OEIS-A271623

a(0)=7; a(n) = 7*a(n-1) + 1 if a(n-1) is odd, a(n) = a(n-1)/2 otherwise.

Conjectured: No term of the sequence is equal to a previous term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Apr 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010847 | OEIS-A271644

Number of ordered ways to write n as w^2 + x^2 + y^2 + z^2 such that w*x + 2*x*y + 2*y*z is a square, where w is a positive integer and x,y,z are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 3, 7, 15, 47, 71, 379, 4^k (k = 0,1,2,...). Clearly, part (i) of this conjecture is stronger than Lagrange's four-square theorem. See also A271510, A271513, A271518 and A271608 for related conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010848 | OEIS-A271648

Number of permutations in S_{2*n+3} containing the pattern 2143...(2n)(2n-1).

For n >= 2, a(n) = (32/3)*n^6 + 32*n^5 + (80/3)*n^4 + (16/3)*n^3 + 38/3*n^2 + 59/3*n + 13 (conjectured). Conjectures from _Colin Barker_, Apr 11 2016: (Start) Remark by _Nathaniel Shar_, Apr 13 2016: The preceding three conjectures are equivalent (provided appropriate initial conditions are specified for the recurrence relation).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nathaniel Shar_, Apr 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010849 | OEIS-A271665

Number of ordered ways to write n as w^2 + x^2 + y^2 + z^2 such that w^2 + 4*x*y + 8*y*z + 32*z*x is a square, where w is a positive integer and x,y,z are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 4^k*3^m, 4^k*3^m*43, 4^k*9^m*q (k,m = 0, 1, 2, ... and q = 7, 15, 79, 95, 141, 159, 183). See also A271510, A271513, A271518 and A271644 for other conjectures refining Lagrange's four-square theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010850 | OEIS-A271670

Number of n-step excursions on the 7-dimensional f.c.c. lattice.

a(n) conjecturally satisfies a linear recurrence equation of order 15 with polynomial coefficients of degree 56 (see link above). The probability generating function P(z) = Sum_{n>=0} a(n)*(z/84)^n is given by the 7-fold integral (1/Pi)^7 Int_{0..Pi} ... Int_{0..Pi} 1/(1-z*lambda_7) dk_1 ... dk_7, where the structure function is defined as lambda_7 = (1/binomial(7,2)) Sum_{i=1..7} Sum_{j=(i+1)..7} cos(k_i)*cos(k_j). The function P(z) conjecturally satisfies an eleventh-order linear ODE with polynomial coefficients of degree 68 (see link above).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christoph Koutschan_, Apr 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010851 | OEIS-A271671

Number of n-step excursions on the 8-dimensional f.c.c. lattice.

a(n) conjecturally satisfies a linear recurrence equation of order 20 with polynomial coefficients of degree 109 (see link above). The probability generating function P(z) = Sum_{n>=0} a(n)*(z/112)^n is given by the 8-fold integral (1/Pi)^8 Int_{0..Pi} ... Int_{0..Pi} 1/(1-z*lambda_8) dk_1 ... dk_8, where the structure function is defined as lambda_8 = (1/binomial(8,2)) Sum_{i=1..8} Sum_{j=(i+1)..8} cos(k_i)*cos(k_j). The function P(z) conjecturally satisfies a linear ODE of order 14 with polynomial coefficients of degree 126 (see link above).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christoph Koutschan_, Apr 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010852 | OEIS-A271672

Number of n-step excursions on the 9-dimensional f.c.c. lattice.

a(n) conjecturally satisfies a linear recurrence equation of order 22 with polynomial coefficients of degree 151 (see link above). The probability generating function P(z) = Sum_{n>=0} a(n)*(z/144)^n is given by the 9-fold integral (1/Pi)^9 Int_{0..Pi} ... Int_{0..Pi} 1/(1-z*lambda_9) dk_1 ... dk_9, where the structure function is defined as lambda_9 = (1/binomial(9,2)) Sum_{i=1..9} Sum_{j=(i+1)..9} cos(k_i)*cos(k_j). The function P(z) conjecturally satisfies a linear ODE of order 18 with polynomial coefficients of degree 169 (see link above). Hence a(n) conjecturally satisfies a linear recurrence equation with polynomial coefficients.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christoph Koutschan_, Apr 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010853 | OEIS-A271673

Number of n-step excursions on the 10-dimensional f.c.c. lattice.

a(n) conjecturally satisfies a linear recurrence equation of order 30 with polynomial coefficients of degree 274 (see link above). The probability generating function P(z) = Sum_{n>=0} a(n)*(z/180)^n is given by the 10-fold integral (1/Pi)^10 Int_{0..Pi} ... Int_{0..Pi} 1/(1-z*lambda_10) dk_1 ... dk_10, where the structure function is defined as lambda_10 = (1/binomial(10,2)) Sum_{i=1..10} Sum_{j=(i+1)..10} cos(k_i)*cos(k_j). The function P(z) conjecturally satisfies a linear ODE of order 22 with polynomial coefficients of degree 300 (see link above).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christoph Koutschan_, Apr 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010854 | OEIS-A271674

Number of n-step excursions on the 11-dimensional f.c.c. lattice.

The probability generating function P(z) = Sum_{n>=0} a(n)*(z/220)^n is given by the 11-fold integral (1/Pi)^11 Int_{0..Pi} ... Int_{0..Pi} 1/(1-z*lambda_11) dk_1 ... dk_11, where the structure function is defined as lambda_11 = (1/binomial(11,2)) Sum_{i=1..11} Sum_{j=(i+1)..11} cos(k_i)*cos(k_j). The function P(z) conjecturally satisfies a linear ODE of order 27 with polynomial coefficients of degree 409 (see link above). Hence a(n) conjecturally satisfies a linear recurrence equation with polynomial coefficients.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christoph Koutschan_, Apr 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010855 | OEIS-A271714

Number of ordered ways to write n as w^2 + x^2 + y^2 + z^2 such that (10*w+5*x)^2 + (12*y+36*z)^2 is a square, where w is a positive integer and x,y,z are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 7, 9, 19, 49, 133, 589, 2^k, 2^k*3, 4^k*q (k = 0,1,2,... and q = 14, 67, 71, 199). See also A271510, A271513, A271518, A271644, A271665, A271721 and A271724 for other conjectures refining Lagrange's four-square theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010856 | OEIS-A271719

Number of ordered ways to write n as x + y + z with x >= y > 0, z > 0 and gcd(x,y,z) = 1 such that x^2 + (2*y+z)^2 is a square.

Conjecture: a(n) > 0 for all n > 10, and a(n) = 1 only for n = 11, 12, 14, 18, 24, 54. See also A271714 for a similar conjecture refining Lagrange's four-square theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010857 | OEIS-A271721

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with x >= y >= z >= 0, x > 0 and w >= z such that (x-y)*(w-z) is a square.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 3, 5, 11, 15, 23, 35, 95, 4^k*190 (k = 0,1,2,...). This is stronger than Lagrange's four-square theorem. Note that for k = 2 or 3, any natural number n can be written as w^2 + x^2 + y^2 + z^2 with w,x,y,z nonnegative integers and (x-y)*(w-k*z) = 0, for, if n cannot be represented by x^2 + y^2 + 2*z^2 then it has the form 4^k*(16*m+14) (k,m = 0,1,2,...) and hence it can be represented by x^2 + y^2 + (k^2+1)*z^2. It is known that natural numbers not represented by x^2 + y^2 + 5*z^2 have the form 4^k*(8*m+3), and that positive even numbers not represented by x^2 + y^2 + 10*z^2 have the form 4^k*(16*m+6) (as conjectured by S. Ramanujan and proved by L. E. Dickson). See also A271510, A271513, A271518,

A271644, A271714 and A271724 for other conjectures refining Lagrange's theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010858 | OEIS-A271724

Number of ordered ways to write n as w^2 + x^2 + y^2 + z^2 with w*(x+2*y+3*z) a square, where w,x,y,z are nonnegative integers with x > 0.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 7, 15, 47, 151, 4^k*q (k = 0,1,2,... and q = 1, 23, 71). See also A271510, A271513, A271518, A271644, A271665, A271714 and A271721 for other conjectures refining Lagrange's four-square theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010859 | OEIS-A271725

T(n,k) is an array read by rows, with n > 0 and k=1..4, where row n gives four prime numbers in increasing order with locations in right angles of each concentric square drawn on a distorted version of the Ulam spiral.

Conjecture: there is an infinity of concentric squares having a prime number in each right angle. The number 5 is the center of all the squares.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010860 | OEIS-A271775

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 (x >= y >= z <= w) with x - y a square, where w,x,y,z are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 3, 11, 47, 2^{4k+3}*m (k = 0,1,2,... and m = 1, 3, 7, 15, 79). See also A271510, A271513, A271518, A271644, A271665, A271714, A271721 and A271724 for other conjectures refining Lagrange's four-square theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010861 | OEIS-A271778

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers and x^2 + 3*y^2 + 5*z^2 - 8*w^2 a square.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 1, 15, 29, 33, 47, 53, 65, 89, 129, 689, 1553, 2^(2k+1)*m (k = 0,1,2,... and m = 1, 29). See also A271510, A271513, A271518, A271665, A271714, A271721, A271724 and A271775 for other conjectures refining Lagrange's four-square theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010862 | OEIS-A271780

Decimal expansion of Product_{p odd prime} 1-2/(p*(p-1)), a constant related to Artin's conjecture in the context of quadratic fields.

Decimal expansion of Product_{p odd prime} 1-2/(p*(p-1)), a constant related to Artin's conjecture in the context of quadratic fields.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Apr 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010863 | OEIS-A271804

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 401", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 409 also generates this sequence. - _Lars Blomberg_, Jun 22 2016 Conjectures from _Colin Barker_, Jun 22 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010864 | OEIS-A271824

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with (x+2*y)^2 + 8*z^2 + 40*w^2 a square, where x is a positive integer and y,z,w are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 9, 11, 15, 23, 33, 71, 129, 167, 187, 473, 4^k*m (k = 0,1,2,... and m = 1, 22, 38, 278). Also, any positive integer can be written as x^2 + y^2 + z^2 + w^2 with 9*(x+2*y)^2 + 16*z^2 + 24*w^2 a square, where x is a positive integer and y,z,w are nonnegative integers. See also A271510, A271513, A271518, A271665, A271714, A271721, A271724, A271775 and A271778 for other conjectures refining Lagrange's four-square theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010865 | OEIS-A271834

a(n) = 2^n - Sum_{m=0..n} binomial(n/gcd(n,m), m/gcd(n,m)) = 2^n - A082906.

Compared to A082906, this sequence shows better the drop from 2^n upon replacing every binomial(n,m) in the Newton's expansion of (1+1)^n by the 'reduced' binomial(n/gcd(n,m), m/gcd(n,m)). For n > 1, a(n) is zero if and only if n is prime (no reduction, no drop). The ratio r(n) = a(n)/2^n is always smaller than 1 and presents considerable excursions. For composite n up to 5000, the minimum of 0.01471... occurs for n = 4489, and the maximum of 0.80849... occurs for n = 2310. This apparently large relative difference is actually surprisingly small: on log_2 scale it amounts to just about 5.78; a tiny fraction compared to the full scale, given by the values of n for the extrema. This insight suggests the following conjecture: there exists an average ratio r, defined as r = lim_{n->infinity} Sum_{m=1..n} r(m)/n. Its value appears to be approximately 0.3915+-0.0010, which can be interpreted as the average drop in a binomial value upon the 'reduction' of its arguments.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Apr 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010866 | OEIS-A271861

Recursive sequence based on the central polygonal numbers (A000124) and A002260.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Barrentine_, Apr 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010867 | OEIS-A271863

Recursive sequence based on the central polygonal numbers (A000124) and A004736.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Barrentine_, Apr 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010868 | OEIS-A271865

Recursive sequence based on the central polygonal numbers (A000124) and A004738.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Barrentine_, Apr 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010869 | OEIS-A271886

Decimal expansion of the constant D related to the conjectured asymptotic expression of the counting function of prime triples as D*n/log(n)^3.

Decimal expansion of the constant D related to the conjectured asymptotic expression of the counting function of prime triples as D*n/log(n)^3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Apr 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010870 | OEIS-A271910

Array read by antidiagonals: T(n,k) = number of ways to choose 3 distinct points from an n X k rectangular grid so that they form an isosceles triangle.

It appears that for each n >= 2, there is a number K(n) such that row n satisfies the recurrence a(k) = 2*a(k-1)-2*a(k-3)+a(k-4) for k >= K(n). This is based on the fact that the conjectured generating functions for rows 2, 3, 4, 5 have the same denominator, and on _Colin Barker_'s conjectured recurrence for A271911. K(n) is determined by the degree of the numerator of the g.f. Above conjecture about the recurrence is true for K(n) = (n-1)^2+4 if n is even and K(n) = (n-1)^2+3 if n is odd and not true for smaller K(n). See paper in links. - _Chai Wah Wu_, May 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010871 | OEIS-A271911

Number of ways to choose three distinct points from a 2 X n grid so that they form an isosceles triangle.

Conjectured g.f.: 2*x*(2*x^2-x-2)/((x+1)*(x-1)^3). It would be nice to have a proof! Conjectures from _Colin Barker_, Apr 24 2016: (Start) The conjectured g.f. and recurrence are true. See paper in links. - _Chai Wah Wu_, May 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010872 | OEIS-A271912

Number of ways to choose three distinct points from a 3 X n grid so that they form an isosceles triangle.

Conjectured g.f.: 2*x*(2*x^4+4*x^3+2*x^2-8*x-5)/((x+1)*(x-1)^3). Conjectured recurrence: a(n) = 2*a(n-1)-2*a(n-3)+a(n-4) for n > 6. Conjectures from _Colin Barker_, Apr 25 2016: (Start) The conjectured g.f. and recurrence are true. See paper in links. - _Chai Wah Wu_, May 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010873 | OEIS-A271913

Number of ways to choose three distinct points from a 4 X n grid so that they form an isosceles triangle.

Conjectured g.f.: 4*x*(x^10-x^8+2*x^6+x^5+4*x^4+4*x^3-3*x^2-9*x-4)/((x+1)*(x-1)^3). Conjectured recurrence: a(n) = 2*a(n-1)-2*a(n-3)+a(n-4) for n > 12. Conjectures from _Colin Barker_, Apr 25 2016: (Start) The conjectured g.f. and recurrence are true. See paper in links. - _Chai Wah Wu_, May 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010874 | OEIS-A271914

Symmetric array read by antidiagonals: T(n,k) (n>=1, k>=1) = maximal number of points that can be chosen in an n X k rectangular grid such that no three distinct points form an isosceles triangle.

It is conjectured that T(n,k) <= n+k-1. Conjecture: For n > 4, T(n,3) = n+1 if n is odd and T(n,3) = n+2 if n is even. Conjecture: If n is even, then T(n,k) <= n+k-2 for k >= 2*n. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rob Pratt_ and _N. J. A. Sloane_, Apr 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010875 | OEIS-A271915

Number of ways to choose three distinct points from a 5 X n grid so that they form an isosceles triangle.

Conjectured g.f.: 4*x*(x^16-x^14+2*x^10+2*x^9-x^8-x^7 + 5*x^6+6*x^5+6*x^4+x^3-8*x^2-15*x-6) /((x+1)*(x-1)^3). Conjectured recurrence: a(n) = 2*a(n-1)-2*a(n-3)+a(n-4) for n > 18. The conjectured g.f. and recurrence are true. See paper in links. - _Chai Wah Wu_, May 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010876 | OEIS-A271931

G.f. A(x) satisfies: A(x) = A(x^5 + 5*x*A(x)^5)^(1/5), with A(0)=0, A'(0)=1.

More generally, for prime p there exists an integer series G(x) that satisfies: G(x) = G(x^p + p*x*G(x)^p)^(1/p) with G(0)=0, G'(0)=1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010877 | OEIS-A271960

G.f. A(x) satisfies: A(x)^2 = A((x + 2*A(x)^2)^2).

Conjecture: [x^(2^n-2^k)] (x/A(x))^(2^n) = 0 for k = 0..n-2 and n >= 2. - _Paul D. Hanna_, Jan 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010878 | OEIS-A271983

The smaller of a pair n, m such that phi(n) = phi(m) and there is no other k such that phi(n) = phi(k).

If phi(x) = N has exactly two solutions, x = n and x = m, say (see A007366), it is conjectured that one of n and m is odd and the other even.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Critzer_, Apr 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010879 | OEIS-A272007

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 413", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Jun 16 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010880 | OEIS-A272009

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 413", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Jun 16 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010881 | OEIS-A272010

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 413", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = A269909(n), n>0. - _R. J. Mathar_, Apr 19 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010882 | OEIS-A272032

Positive integers n such that (q^n + r^n)/(q+r) is prime, where q is the number of consecutive composite numbers smaller than n and r is the number of consecutive composite numbers greater than n.

All terms are prime (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marc Morgenegg_, Apr 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010883 | OEIS-A272084

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with 4*x^2 + 5*y^2 + 20*z*w a square, where x,y,z,w are nonnegative integers with z < w.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 2^k*q (k = 0,1,2,... and q = 1, 3, 7), 4^k*m (k = 0,1,2,... and m = 21, 30, 38, 62, 70, 77, 142, 217, 237, 302, 382, 406, 453, 670). See also A262357, A268507, A269400, A271510, A271513, A271518, A271665, A271714, A271721, A271724, A271775, A271778 and A271824 for other conjectures refining Lagrange's four-square theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010884 | OEIS-A272101

Square root of largest square dividing A069482(n).

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benedict W. J. Irwin_, Apr 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010885 | OEIS-A272102

Numerators of bivariate Taylor expansion of the incomplete elliptic integral of the first kind.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Francesco Franco_, Apr 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010886 | OEIS-A272139

Numbers n such that Bernoulli number B_{n} has denominator 1806.

In 2005, B. C. Kellner proved E. W. Weisstein's conjecture that denom(B_n) = n only if n = 1806.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_, Apr 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010887 | OEIS-A272170

Second most significant bit of Fibonacci numbers > 1 written in base 2.

It is conjectured that there are no more than two consecutive "0's" or "1's" (tested up to n=10^5). The sequence looks quasiperiodic and its Fourier spectrum seems to have a fractal structure.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andres Cicuttin_, Apr 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010888 | OEIS-A272175

Least number k such that (k^2+1) mod s = prime(n) where s is the sum of the distinct primes dividing k^2+1, or 0 if no such k exists.

Conjecture: a(n)> 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Apr 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010889 | OEIS-A272198

The p-defect p - N(p) of the congruence y^2 == x^3 + 1 (mod p) for primes p, where N(p) is the number of solutions given by A272197(n).

If prime(n) == 1 (mod 3) = A002476(m) (for a unique m = m(n)) then prime(n) = A(m)^2 + 3*B(m)^2 with A(m) = A001479(m+1) and B(m) = A001480(m+1), m >= 1. In this case (4*prime(n) - a(n)^2)/12, seems to be a square, q(m)^2. In fact is seems that (the positive) q(m) = B(m). This is true at least for the first 80 primes 1 (mod 3), i.e. for such primes <= 997. (In the Silverman reference, in hint c) for Exercise 4.5, on p. 405, a more complicated way is suggested: 4*p is decomposed there non-uniquely instead of p uniquely.) If this conjecture is true then a(n) = 2*(+/-sqrt(prime(n) - 3*B(m)^2)) = +- 2*A(m) for prime(n) = A002476(m). This leads to a bisection of the primes 1 (mod 3) into two types: type I if the + sign applies, and type II for the - sign. Primes of type I are given in A272200: 13, 19, 43, 61, 97, ... and those of type II in A272201: 7, 31, 37, 67, 73, ... The above given conjecture for primes 1 (mod 3) is true because Mordell proved the Ramanujan conjecture on the expansion coefficients of eta^4(6*z), and with the present a(n) the result of Ramanujan follows. See the references and a comment on A000727.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, May 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010890 | OEIS-A272203

P-defects p - N(p) of the congruence y^2 == x^3 - 1 (mod p) for primes p, where N(p) is the number of solutions given by A272202(n).

If prime(n) == 1 (mod 3) = A002476(m) (for a unique m = m(n)) then prime(n) = A(m)^2 + 3*B(m)^2 with A(m) = A001479(m+1) and B(m) = A001480(m+1), m >= 1. In this case (4*prime(n) - a(n)^2)/12, seems to be a square, q(m)^2. In fact is seems that (the positive) q(m) = B(m). If this conjecture is true then a(n) = 2*(+-sqrt(prime(n) - 3*B(m)^2)) = +- 2*A(m) for prime(n) = A002476(m). This leads to a bisection of the primes 1 (mod 3) into two types: type I if the + sign applies, and type II for the - sign. Primes of type I are given in A272204: 7,13,31,61,67, ... and those of type II in A272205: 19,37,43,73,103, ... The above given conjecture for primes 1 (mod 3) is expected to be true by analogy to the case A272198 where only the signs differ.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, May 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010891 | OEIS-A272242

a(n) is the least number c such that there are exactly n abc-hits with third member c, or 0 if no such c exists.

Conjecture: a(n) > 0 for all n. - _Jianing Song_, Sep 21 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Letsko_, Apr 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010892 | OEIS-A272261

Number of one-to-one functions f from [n] to [2n] where f(x) may not be equal to x or to 2n+1-x.

Conjecture: Alternating sign g.f. is Sum_{k>=0} HermiteH[k,sqrt(x)]x^(k/2). - _Benedict W. J. Irwin_, Nov 30 2016 Conjecture D-finite with recurrence: a(n) + 2*(-2*n+1)*a(n-1) - 4*a(n-2)=0. - _R. J. Mathar_, Jan 27 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marko Riedel_, Apr 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010893 | OEIS-A272266

The union of squares (A000290) and 10-gonal numbers (A001107).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Barker_, Apr 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010894 | OEIS-A272267

The union of centered square numbers (A001844) and centered 9-gonal numbers (A060544).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Barker_, Apr 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010895 | OEIS-A272295

a(n) = v(2*n+1), where v is a completely multiplicative sequence defined by v(2)=0 and for prime p >= 3: v(p^e) = ((p+1)/2)^e when p == 1 mod 4, and v(p^e) = ((1-p)/2)^e when p == 3 mod 4.

Conjecture: Sum_{n >= 1, n not divisible by 2 or 3} (moebius(n)/v(n))^2 = 7/5 = 1.4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimitris Valianatos_, Apr 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010896 | OEIS-A272303

Magic sums of 4 X 4 semimagic squares composed of primes.

Conjecture: every even number greater than 134 belongs to this sequence. If conjecture is true, a(n) = A005843(n+61) for n >= 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Apr 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010897 | OEIS-A272322

Numbers n such that n^n does not contain all ten decimal digits.

Conjecture: the only values > 40 are the powers of 10. - _Franklin T. Adams-Watters_, Apr 29 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Apr 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010898 | OEIS-A272332

Number of ordered ways to write n as w^2 + x^2 + y^2 + z^2 with 36*x^2*y + 12*y^2*z + z^2*x a square, where w is a positive integer and x,y,z are nonnegative integers.

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 16^k*m (k = 0,1,2,... and m = 1, 79, 591, 599, 1752, 1839, 10264).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010899 | OEIS-A272336

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with x^4 + y^3*z a square, where x,y,z,w are nonnegative integers with z > 0.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 23, 31, 71, 191, 271, 391, 503, 943, 1591, 2351, 2791, 4791, 8863, 9983, 4^k*m (k = 0,1,2,... and m = 1, 7, 79). See also A280831 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010900 | OEIS-A272342

a(n) = 27*8^n.

a(n) are cubes that can be expressed as sum of exactly four distinct powers of two: a(n)=2^3n + 2^(3n+1) + 2^(3n+3) + 2^(3n+4). For example a(0) = 2^0 + 2^1 + 2^3 + 2^4 = 1 + 2 + 8 + 16 = 27. It is conjectured the a(n) are the only cubes that can be expressed as sum of exactly four distinct nonnegative powers of two (tested on cubes up to (10^7)^3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andres Cicuttin_, Apr 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010901 | OEIS-A272351

Number of ordered ways to write n as w^2 + x^2 + y^2 + z^2 with x^4 + 8*y*z*(y^2+z^2) a fourth power, where w,x,y,z are nonnegative integers with w >= x and y > z.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 2^k, 4^k*m (k = 0,1,2,... and m = 3, 7, 31, 55, 71, 79, 151, 191). The author has proved part (ii) of the conjecture in arXiv:1604.06723. - _Zhi-Wei Sun_, May 09 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010902 | OEIS-A272397

Number of partitions of n into parts congruent to 1, 3, 6, 8 (mod 9).

"Sum side" conjecture: also equals number of partitions pi = (pi_1, pi_2, ...) of n (with pi_1 >= pi_2 >= ...) such that pi(i)-pi(i+2) >= 3 and, if pi(i) - pi(i+1) <= 1, then pi(i) + pi(i+1) is congruent to 0 (mod 3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthew C. Russell_, Apr 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010903 | OEIS-A272398

The union of hexagonal numbers (A000384) and centered 9-gonal numbers (A060544).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Barker_, Apr 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010904 | OEIS-A272417

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 469", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010905 | OEIS-A272459

The total number of different isosceles trapezoids, excluding squares, that can be drawn on an n X n square grid where the corners of each individual trapezoid lie on a lattice point.

a(n) = Sum_{k=0..n} A032438(k) + A000292(n-1). (conjectured) a(n) = A143785(n-2) + A000292(n-1). (conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christopher J. Shore_, Apr 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010906 | OEIS-A272462

E.g.f.: (sin(3^2*x) + sin(4^2*x)) / sin(5^2*x).

Conjecture: Given positive integers a and b, then the coefficient of x^(2*n)/(2*n)! is integral for n>=0 in the power series expansion of ((sin(a*x) + sin(b*x)) / sin((a+b)*x))^(1/(a*b)). See the Bala link for a proof of the conjecture. - _Peter Bala_, Sep 03 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010907 | OEIS-A272468

E.g.f.: ((sin(2*x) + sin(3*x)) / sin(5*x))^(1/6).

Conjecture: Given positive integers a and b, then the coefficient of x^(2*n)/(2*n)! is integral for n>=0 in the power series expansion of ((sin(a*x) + sin(b*x)) / sin((a+b)*x))^(1/(a*b)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010908 | OEIS-A272479

a(n) is the smallest k different from n such that (n, k) is a Harshad amicable pair (see the comments).

Conjecture: the sequence contains no zeros. Adamczewski includes proof abstract regarding the above conjecture in linked article. - _Michael De Vlieger_, Aug 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Waldemar Puszkarz_, May 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010909 | OEIS-A272515

Number of set partitions of [n] into three blocks with distinct sizes.

Conjecture: a(n) ~ 3^n / 6. - _Vaclav Kotesovec_, Dec 11 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, May 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010910 | OEIS-A272544

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 493", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = (2142*4^(n-3) + 196*3^(n-4) - 531*2^(n-3) + 144)/9, n>5. - _Lars Blomberg_, Jul 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010911 | OEIS-A272564

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 501", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 03 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010912 | OEIS-A272566

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 501", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 03 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010913 | OEIS-A272567

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 501", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 03 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010914 | OEIS-A272575

Perfect powers that are the sum of two Fibonacci numbers.

Conjecture: there are no more terms in this sequence. Any remaining terms must have over 10000 digits. - _Charles R Greathouse IV_, May 04 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, May 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010915 | OEIS-A272586

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 507", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = (11*4^n-12*2^n-8)/3 , n>1. - _Lars Blomberg_, Jul 08 2016 Conjectures from _Colin Barker_, Jul 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010916 | OEIS-A272588

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 507", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = A272584(n), n>1. - _R. J. Mathar_, May 20 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010917 | OEIS-A272606

Numbers k such that there is no prime p and index j < k such that A002182(k) = p * A002182(j).

This is a sequence of counterexamples to the second conjecture by Alaoglu & Erdős that such highly composite numbers do not exist (they did not consider 1 to be highly composite number). Robin found the first 3 counterexamples: A002182(815) = 3.622... * 10^65, A002182(4372) = 6.043... * 10^220, and A002182(4996) = 1.115 * 10^244. - _Amiram Eldar_, Jul 09 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Nov 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010918 | OEIS-A272615

Numbers with digits in descending numerical order in base 2, 3 and 4 expansions.

a(10), if it exists, has more than 1000 decimal digits. Conjecture: there are no more terms in this sequence. - _Charles R Greathouse IV_, May 03 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robin Powell_, May 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010919 | OEIS-A272620

Number of ordered ways to write n as w^2 + x^2 + y^2 + z^2 with w + x + y - z a square, where w is an integer and x,y,z are nonnegative integers with |w| <= x >= y <= z < x + y.

Conjecture: a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010920 | OEIS-A272651

The no-3-in-line problem: maximal number of points from an n X n square grid so that no three lie on a line.

It is conjectured that a(n) < 2*n for all sufficiently large n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010921 | OEIS-A272678

Smallest number m such that A272677(m) = n.

a(n) >= 25*10^(n-3). Conjecture: a(n)/(25*10^(n-3)) -> 1 as n -> oo. - _Chai Wah Wu_, May 21 2016 Except for n = 1 and 13, a(n) appears to be approximately equal to either 25*10^(n-3)+sqrt(10^(n-1)) (for n = 0, 2, 3, 5, 6, 9, 11, 12, 15, 18, ...) or 25*10^(n-3)+sqrt(2*10^(n-1)) (for n = 4, 7, 8, 14, 16, 17, ...). For n = 1, a(n) is approximately 25*10^(n-3)+sqrt(3*10^(n-1)) and for n = 13, a(n) is about equal to 25*10^(n-3)+sqrt(4*10^(n-1)). Conjecture: a(n) is always approximately to 25*10^(n-3)+sqrt(k*10^(n-1)) for some small integer k > 0. - _Chai Wah Wu_, May 22 2016 Using the above conjecture as a guide, upper bounds for a(n) can be computed (see file in links) which coincide with a(n) for n <= 19. - _Chai Wah Wu_, May 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nathan Fox_, _Brooke Logan_, and _N. J. A. Sloane_, May 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010922 | OEIS-A272685

a(n) = smallest m such that A265432(m) = A272684(n), or -1 if no such m exists.

It is conjectured that all terms of A272684 eventually appear in A265432.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nathan Fox_, _Brooke Logan_, and _N. J. A. Sloane_, May 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010923 | OEIS-A272688

The antibracket constants {x_n}^n.

There is a conjectured formula - see Conjecture 2.4 of Manetti-Ricciardi.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010924 | OEIS-A272706

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 515", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 523 also generates this sequence. - _Lars Blomberg_, Jul 08 2016 Conjecture: a(n) = 3*4^n - 8*2^n + 9. - _Lars Blomberg_, Jul 08 2016 Conjectures from _Colin Barker_, Jul 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010925 | OEIS-A272719

Maximal number of steps in a nontouching path on an n X n grid such that each step has a different length.

a(10)>=30, a(11)>=32, a(12)>=36, a(13)>=40, a(14)>=44, a(15)>=49, a(16)>=53, a(17)>=58, a(18)>=63. Conjecture: floor(1.19*A226595(n)) <= a(n) <= ceiling(1.5*A226595(n)). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Resta_, May 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010926 | OEIS-A272729

a(n) is the number of repetitions of 2n-1 in A272727.

Conjectured first differences of A037988 (true for at least 8192 terms). - _Sean A. Irvine_, Jun 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ivan Neretin_, May 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010927 | OEIS-A272743

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 526", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = (13*4^(n-1) - 1)/3, n>1. - _Lars Blomberg_, Jul 08 2016 Conjectures from _Colin Barker_, Jul 08 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010928 | OEIS-A272762

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 822", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 886 also generates this sequence. - _Lars Blomberg_, Jul 22 2016 a(n) = 12*a(n-1) - 55*a(n-2) + 120*a(n-3) - 124*a(n-4) + 48*a(n-5) for n > 4 (conjectured). G.f.: (-56*x^4 + 40*x^3 - 20*x^2 + 7*x - 1)/((x - 1)*(2*x - 1)^2*(3*x - 1)*(4*x - 1)) (conjectured). (End) a(n) = 5 - 2^(1+n) + 2^(1+2*n) - 4*3^n + 2^(2+n)*n (conjectured). - _Colin Barker_, Apr 28 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010929 | OEIS-A272766

Numbers n such that A281363(n) = n-1.

Conjecture: primorial numbers (A002110) are terms of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, May 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010930 | OEIS-A272771

Smallest k in the interval [prime(n), 2*prime(n)], such that k has the maximal number of divisors in this interval.

Conjecturally the different values of the sequence are highly composite numbers (A002182, n>=3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010931 | OEIS-A272817

Distance from n-th highly composite number (cf. A002182) to nearest prime or square.

Conjecture: Terms are either 1 or prime, n>7. - _Bill McEachen_, Jun 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, May 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010932 | OEIS-A272832

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 694", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = 2*4^n - 16*3^(n-2) + 2^(n-1) + 1, n>2. - _Lars Blomberg_, Jul 19 2016 Conjectured g.f.: (1 - 5*x + 6*x^2 + 36*x^4 - 92*x^5 + 48*x^6)/((1 - x)*(1 - 2*x)*(1 - 3*x)*(1 - 4*x)). - _Stefano Spezia_, Jun 07 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010933 | OEIS-A272860

Sums of two primes (in increasing order) when equal to the product of their prime-counting functions.

Conjecture: the sequence is infinite and each term has only one decomposition into a sum of suitable primes p,q.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giuseppe Coppoletta_, Jun 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010934 | OEIS-A272882

Discriminator of the Cantor numbers (A005823).

The linked paper by Haque and Shallit gives a conjectured recursive description of the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, May 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010935 | OEIS-A272887

Number of ways to write prime(n) as (4*x + 2)*y + 4*x + 1 where x and y are nonnegative integers.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, May 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010936 | OEIS-A272888

Number of ordered ways to write n as w^2 + x^2 + y^2 + z^2 with w*(x^2 + 8*y^2 - z^2) a square, where w,x,y are nonnegative integers and z is a positive integer.

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 7, 39, 63, 87, 5116, 2^(4k+2)*m (k = 0,1,2,... and m = 1, 7).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010937 | OEIS-A272901

Smallest k>=1 such that A067128(n+k) - A067128(n) is in A067128 or a(n)=0 if there is no such k.

For n=151, the required k, if it exists, should be >9800. So one can conjecture that a(151) is the first zero term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, May 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010938 | OEIS-A272904

Rectangular array, by antidiagonals: row n gives the positions of n in the Fibonacci-products fractal sequence, A272900.

Column 1: A267682 (conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010939 | OEIS-A272909

Numbers that are the product of two Lucas numbers L(i), for i >= 1, using the Lucas numbers as defined in A000204.

Conjecture: if c and d are consecutive terms, then d - c is a product of two Lucas numbers or a product of two Fibonacci numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010940 | OEIS-A272911

Difference sequence of the sequence the increasing sequence of products of two Lucas numbers A000032.

Conjecture: every term is a product of two Lucas numbers or a product of two Fibonacci numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010941 | OEIS-A272916

Numbers that are a product of two Fibonacci (A000045) numbers or a product of two Lucas (A000032) numbers.

Conjecture: if c and d are consecutive terms, then d - c is a term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010942 | OEIS-A272917

Difference sequence of A272916.

Conjecture: every term belongs to A272916.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010943 | OEIS-A272921

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 555", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 571 also generates this sequence. - _Lars Blomberg_, Jul 10 2016 Conjecture: a(n) = 3*4^n - 4*3^(n-2) - 4*2^n + 1, n>1. - _Lars Blomberg_, Jul 10 2016 Conjectures from _Colin Barker_, Dec 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010944 | OEIS-A272930

a(n) is the least k such that sigma(sigma(k)) = n*k, where sigma(n) is the sum of the divisors of n, or 0 if no such k exists.

Is a(n) in fact nonzero for every positive n? - _Franklin T. Adams-Watters_, Jan 22 2019 [who previously conjectured that it is]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, May 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010945 | OEIS-A272977

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with 3*x^2*y + z^2*w a square, where w is a nonzero integer and x,y,z are nonnegative integers with x >= z.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 3, 7, 23, 47, 71, 147, 199, 263, 439, 16^k*m (k = 0,1,2,... and m = 6, 12, 24, 44, 56, 140, 156, 174, 204, 284, 4652). Compare this conjecture with the one in A270073.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010946 | OEIS-A272979

Number of ways to write n as x^2 + 2*y^2 + 3*z^3 + 4*w^4 with x,y,z,w nonnegative integers.

Conjecture: For positive integers a,b,c,d, any natural number can be written as a*x^2 + b*y^2 + c*z^3 + d*w^4 with x,y,z,w nonnegative integers, if and only if (a,b,c,d) is among the following 49 quadruples: (1,2,1,1), (1,3,1,1), (1,6,1,1), (2,3,1,1), (2,4,1,1), (1,1,2,1), (1,4,2,1), (1,2,3,1), (1,2,4,1), (1,2,12,1), (1,1,1,2), (1,2,1,2), (1,3,1,2), (1,4,1,2), (1,5,1,2), (1,11,1,2), (1,12,1,2), (2,4,1,2), (3,5,1,2), (1,1,4,2), (1,1,1,3), (1,2,1,3), (1,3,1,3), (1,2,4,3), (1,2,1,4), (1,3,1,4), (2,3,1,4), (1,1,2,4), (1,2,2,4), (1,8,2,4), (1,2,3,4), (1,1,1,5), (1,2,1,5), (2,3,1,5), (2,4,1,5), (1,3,2,5), (1,1,1,6), (1,3,1,6), (1,1,2,6), (1,2,1,8), (1,2,4,8), (1,2,1,10), (1,1,2,10), (1,2,1,11), (2,4,1,11), (1,2,1,12), (1,1,2,13), (1,2,1,14),(1,2,1,15). See also A262824, A262827, A262857 and A273917 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010947 | OEIS-A273003

Arrange the base 10 digits of the n-th Fibonacci number in descending order.

Conjecture: the largest Fibonacci number F(n) with its base 10 digits already sorted in descending order is F(16) = 987.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alonso del Arte_, May 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010948 | OEIS-A273019

a(n) = hypergeom([-2*n-1, 1/2], [2], 4) + (2*n+1)*hypergeom([-n+1/2, -n], [2], 4).

Conjecture: +(241*n+56) *(2*n+1) *(n+1)*a(n) +(482*n^3-25561*n^2+13831*n-56) *a(n-1) +(-48682*n^3+225897*n^2-300131*n+125310) *a(n-2) +9*(n-2) *(2651*n-3183) *(2*n-3) *a(n-3)=0. - _R. J. Mathar_, May 16 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, May 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010949 | OEIS-A273021

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with 2*x*y + y*z - z*w - w*x a square, where w is a positive integer and x,y,z are nonnegative integers with x <= y.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 11, 31, 47, 55, 71, 105, 115, 119, 253, 383, 385, 4^k*m (k = 0,1,2,... and m = 2, 22, 23, 30, 330). See arXiv:1604.06723 for more conjectural refinements of Lagrange's four-square theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010950 | OEIS-A273038

Least k such that for all m >= k, A067128(m) is divisible by n.

Since by hypothesis k_1 is bounded, for large n we can choose the required x, which gives a contradiction. So k_1 is unbounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David A. Corneth_, May 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010951 | OEIS-A273045

Fibonacci numbers with digits in nondecreasing order.

Conjecture: The sequence is finite, with 377 being the last term. I have verified the conjecture up to A000045(10^5). - _Felix Fröhlich_, May 16 2016 Conjecture verified up to Fibonacci(10^8). - _Charles R Greathouse IV_, May 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, May 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010952 | OEIS-A273059

Positions of 1's in A274640: Greedy Queens on a spiral. Equivalently, positions of 0's in A274641.

This is the Greedy Queens construction on a spiral (cf. A065188). Follow a counterclockwise spiral starting at the origin, and place a queen iff it is not attacked by any existing queen. This same problem is described in a different but equivalent way in A140100 and A140101. See A140101 for a conjectured recurrence which underlies all these sequences. - _N. J. A. Sloane_, Aug 28-30, 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Jul 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010953 | OEIS-A273107

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with (8*x+12*y)^2 + (15*z)^2 a square, where x,y,z,w are nonnegative integers with x+y > 0 and z > 0.

Conjecture: a(n) > 0 for all n > 5, and a(n) = 1 only for n = 7, 9, 23, 25, 31, 55, 2^k*m (k = 1,2,... and m = 1, 5), 2^(2k+1)*m (k = 0,1,2,... and m = 3, 13, 21). This conjecture implies that any integer n > 5 can be written as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that 8*x+12*y and 15*z are the two legs of a right triangle with positive integer sides. See also A271714, A273108, A273110 and A273134 for similar conjectures related to Pythagorean triples. For more conjectural refinements of Lagrange's four-square theorem, one may consult arXiv:1604.06723.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010954 | OEIS-A273108

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with (x + y)^2 + (4z)^2 a square, where x,y,z,w are nonnegative integers with x <= y > z.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 39, 47, 95, 543, 4^k*m (k = 0,1,2,... and m = 1, 3, 7, 15, 23, 135, 183). See also A271714, A273107, A273110 and A273134 for similar conjectures related to Pythagorean triples. For more conjectural refinements of Lagrange's four-square theorem, one may consult arXiv:1604.06723.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010955 | OEIS-A273110

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with (x+4*y+4*z)^2 + (9*x+3*y+3*z)^2 a square, where x,y,z,w are nonnegative integers with y > 0 and y >= z <= w.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 4^k*m (k = 0,1,2,... and m = 1, 7, 23, 31, 39, 47, 55, 71, 79, 119, 151, 191, 311, 671). See also A271714, A273107, A273108 and A273134 for similar conjectures related to Pythagorean triples. For more conjectural refinements of Lagrange's four-square theorem, one may consult arXiv:1604.06723.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010956 | OEIS-A273134

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with (x+8*y+8*z+15*w)^2+(6*(x+y+z+w))^2 a square, where x,y,z,w are nonnegative integers with y < z.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 3, 7, 11, 15, 21, 23, 35, 39, 71, 95, 4^k*m (k = 0,1,2,... and m = 1, 2, 5, 6, 10, 14, 29, 30, 46, 62, 94, 110, 142, 190, 238, 334, 446). Part (i) of this conjecture implies that any positive integer can be written as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that x+8*y+8*z+15*w and 6*(x+y+z+w) are the two legs of a right triangle with positive integer sides. If a nonnegative integer n is not of the form 4^k*(16m+14) (k,m = 0,1,2,...), then n can be written as w^2+x^2+y^2+z^2 with w = x and hence (x+8y+8z+15w)^2 + (6(x+y+z+w))^2 = (8(2x+y+z))^2 + (6(2x+y+z))^2 = (10(2x+y+z))^2. Similar remarks apply to part (ii) of the conjecture. See also A271714, A273107, A273108 and A273110 for similar conjectures related to Pythagorean triples. For more conjectural refinements of Lagrange's four-square theorem, one may consult arXiv:1604.06723.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010957 | OEIS-A273144

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 597", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 16 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010958 | OEIS-A273145

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 597", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = 14*4^(n-1) - 4*2^n + 1, n>1. - _Lars Blomberg_, Jul 13 2016 Conjectures from _Colin Barker_, Jul 13 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010959 | OEIS-A273146

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 597", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 16 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010960 | OEIS-A273147

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 597", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 16 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010961 | OEIS-A273251

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 619", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 635 also generates this sequence. - _Lars Blomberg_, Jul 16 2016 Conjecture: a(n) = (10*4^n - 12*2^n - 1)/3, n>0. - _Lars Blomberg_, Jul 16 2016 Conjectures from _Colin Barker_, Jul 16 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010962 | OEIS-A273278

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that x^2*y^2 + 3*y^2*z^2 + 2*z^2*w^2 is a square.

Conjecture: a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 4^k*m (k = 0,1,2,... and m = 7, 39, 87, 183, 231, 807, 879, 959, 1479, 2391, 2519, 2759, 4359, 10887). See part (ii) of the conjecture in A269400 for similar conjectures. For more conjectural refinements of Lagrange's four-square theorem, one may consult arXiv:1604.06723.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010963 | OEIS-A273294

Least nonnegative integer m such that there are nonnegative integers x,y,z,w for which x^2 + y^2 + z^2 + w^2 = n and x + 3*y + 5*z = m^2.

Clearly, a(n) = 0 if n is a square. Part (i) of the conjecture in A271518 implies that a(n) always exists. For more conjectural refinements of Lagrange's four-square theorem, one may consult arXiv:1604.06723.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010964 | OEIS-A273309

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 641", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 19 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010965 | OEIS-A273310

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 641", based on the 5-celled von Neumann neighborhood.

Conjecture: Rules 649, 897 and 905 also generate this sequence. - _Lars Blomberg_, Jul 17 2016 Conjecture: a(n) = 4*4^n - 4*2^n - 8, n>1. - _Lars Blomberg_, Jul 17 2016 Conjectures from _Colin Barker_, Jul 17 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010966 | OEIS-A273311

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 641", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 19 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010967 | OEIS-A273312

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 641", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 19 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010968 | OEIS-A273313

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 643", based on the 5-celled von Neumann neighborhood.

Conjecture: Rules 651, 707 and 715 also generate this sequence. - _Lars Blomberg_, Jul 17 2016 Conjecture: a(n) = 4*4^n - 4*2^n - 7, n>1. - _Lars Blomberg_, Jul 17 2016 Conjectures from _Colin Barker_, Jul 17 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010969 | OEIS-A273314

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 643", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 19 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010970 | OEIS-A273315

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 643", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 19 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010971 | OEIS-A273316

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 645", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 19 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010972 | OEIS-A273327

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 646", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = 19*4^(n-2) - 16*3^(n-3) + 1, n>2. - _Lars Blomberg_, Jul 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010973 | OEIS-A273331

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 654", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = (53*4^(n-2) - 5)/3, n>2. - _Lars Blomberg_, Jul 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010974 | OEIS-A273334

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 657", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 20 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010975 | OEIS-A273335

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 657", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 665 also generates this sequence. - _Lars Blomberg_, Jul 18 2016 Conjectures from _Colin Barker_, May 20 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010976 | OEIS-A273336

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 657", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 20 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010977 | OEIS-A273337

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 657", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 20 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010978 | OEIS-A273343

The sum of the first-column lengths of all bargraphs of semiperimeter n (n>=2).

Conjecture: (n+2)*a(n) +(-5*n-3)*a(n-1) +(5*n-2)*a(n-2) +(2*n-9)*a(n-3) +(-n+3)*a(n-4) +(-n+4)*a(n-5) +(-n+7)*a(n-6)=0. - _R. J. Mathar_, Jun 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, May 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010979 | OEIS-A273351

Number of up steps in all bargraphs of semiperimeter n (n>=2).

Conjecture: n*(13*n-40)*a(n) +(-55*n^2+211*n-129)*a(n-1) +(38*n^2-212*n+255)*a(n-2) +(-6*n^2+56*n-75)*a(n-3) +(13*n^2-66*n+3)*a(n-4) -(3*n-13)*(n-6)*a(n-5)=0. - _R. J. Mathar_, Jun 06 2016 Conjecture: n*(n-3)*(n-2)^2*a(n) -(n-3)*(2*n-3)*(2*n^2-6*n+3) *a(n-1) +(2*n^4-16*n^3+41*n^2-36*n+5) *a(n-2) +2*(n-1)*(2*n-5) *a(n-3) +(n-2)*(n-5)*(n-1)^2 *a(n-4)=0. - _R. J. Mathar_, Jun 06 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jun 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010980 | OEIS-A273379

Ramanujan's largely composite numbers n (A067128) which are not divisible by all the primes < p, where p is the greatest prime divisor of n.

Moreover, the numerical results of _David A. Corneth_ (cf. A273415) allow us to conjecture that indeed 0 < c < 1. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, May 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010981 | OEIS-A273384

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 659", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 21 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010982 | OEIS-A273385

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 659", based on the 5-celled von Neumann neighborhood.

Conjecture: Rules 667, 723, 731, 931, 939, 947, 955, 995, 1003, 1011 and 1019 also generate this sequence. - _Lars Blomberg_, Jul 18 2016 Conjecture: a(n) = 4*4^n - 4*2^n + 1 = A060867(n+1), n>1. - _Lars Blomberg_, Jul 18 2016 Conjectures from _Colin Barker_, Dec 01 2016: (Start) Conjecture: a(n) = A270007(n) for n>=2. - _R. J. Mathar_, Jun 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010983 | OEIS-A273386

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 659", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 21 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010984 | OEIS-A273387

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 659", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 21 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010985 | OEIS-A273389

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 662", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = 5*4^(n-1) - 2^(n-1) - 51, n>5. - _Lars Blomberg_, Jul 18 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010986 | OEIS-A273404

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with x + 24*y a square, where x,y,z,w are nonnegative integers with z <= w.

Conjecture: a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 16^k*m (k = 0,1,2,... and m = 8, 12, 23, 24, 47, 71, 168, 344, 632). For more conjectural refinements of Lagrange's four-square theorem, one may consult arXiv:1604.06723.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010987 | OEIS-A273405

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 673", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 22 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010988 | OEIS-A273406

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 673", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 22 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010989 | OEIS-A273407

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 673", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 22 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010990 | OEIS-A273408

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 675", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 22 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010991 | OEIS-A273417

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 705", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 22 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010992 | OEIS-A273418

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 705", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 713 also generates this sequence. - _Lars Blomberg_, Jul 19 2016 Conjecture: a(n) = A273313(n), n>1. - _R. J. Mathar_, May 27 2016 Conjecture: a(n) = 4*4^n - 4*2^n - 7, n>1. - _Lars Blomberg_, Jul 19 2016 Conjectures from _Colin Barker_, Dec 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010993 | OEIS-A273419

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 705", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 22 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010994 | OEIS-A273420

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 705", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 22 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010995 | OEIS-A273429

Number of ordered ways to write n as x^6 + y^2 + z^2 + w^2, where x,y,z,w are nonnegative integers with y <= z <= w.

For more conjectural refinements of Lagrange's four-square theorem, one may consult the author's preprint arXiv:1604.06723.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010996 | OEIS-A273432

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with 2*x + y - z a nonnegative cube, where x,y,z,w are nonnegative integers with y <= z.

Conjecture: (i) For each c = 1, 2, 4 and n = 0,1,2,..., we can write n as x^2 + y^2 + z^2 + w^2 with c*(2x+y-z) a nonnegative cube, where x,y,z,w are nonnegative integers with y <= z. See also A273458 for a similar conjecture. For more conjectural refinements of Lagrange's four-square theorem, see the author's preprint arXiv:1604.06723.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010997 | OEIS-A273443

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 721", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 23 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010998 | OEIS-A273446

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 721", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 729 also generates this sequence. - _Lars Blomberg_, Jul 19 2016 Conjecture: a(n) = 4*4^n - 4*2^n, n>2. - _Lars Blomberg_, Jul 19 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-010999 | OEIS-A273447

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 721", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 23 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011000 | OEIS-A273448

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 721", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 23 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011001 | OEIS-A273457

Even numbers 2n that do not have a Goldbach partition 2n = p + q (p < q; p, q prime) satisfying sqrt(n) < p <= sqrt(2n).

It is conjectured that a(12831) = 15702604 is the last term. There are no more terms below 4*10^10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Corinna Regina Böger_, Dec 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011002 | OEIS-A273458

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with x-y+z+w a nonnegative cube, where x,y,z,w are integers with x >= y >= 0 and x >= |z| <= |w|.

Conjecture: a(n) > 0 for all n = 0,1,2,.... For more conjectural refinements of Lagrange's four-square theorem, see the author's preprint arXiv:1604.06723.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011003 | OEIS-A273464

The number of tilings of an equilateral triangle of side length n with k lozenges and n^2 - 2*k unit triangles. Triangle T(n, k) with n >= 1 and 0 <= k <= n*(n + 1)/2, read by rows.

Conjecture: T(n,4) = 3*(n-2)*(n-3)*(9*n^6+9*n^5-135*n^4-81*n^3+670*n^2+104*n-1216)/128. - _Greg Dresden_, Jul 03 2019 Conjecture: T(n,5) = 3*(n-3)*(n+3)* (27*n^8 -135*n^7 -387*n^6 +2835*n^5 -168*n^4 -18732*n^3 +19568*n^2 +36992*n -56320)/1280. - _R. J. Mathar_, Jul 07 2019 Conjecture for rightmost terms: A122722(n) = n! * T(n, n*(n+1)/2) for n >= 1. Conjectures for column k >= 0: Sum_{0 <= s <= 2*k + 1} (-1)^s * binomial(2*k+1, s) * T(n-s, k) = 0 for n >= 2*k+2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, May 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011004 | OEIS-A273480

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 737", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 23 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011005 | OEIS-A273481

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 737", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 23 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011006 | OEIS-A273489

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 758", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = 2*4^n - 4*3^(n-1) + 2*2^n - 23, n>2. - _Lars Blomberg_, Jul 20 2016 Conjectures from _Colin Barker_, Jul 20 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011007 | OEIS-A273539

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 779", based on the 5-celled von Neumann neighborhood.

Conjecture: Rules 795, 843 and 859 also generate this sequence. - _Lars Blomberg_, Jul 20 2016 Conjectures from _Colin Barker_, May 25 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011008 | OEIS-A273565

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 793", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = 43*4^(n-2) - 4*2^n + 1, n>1. - _Lars Blomberg_, Jul 21 2016 Conjectures from _Colin Barker_, Dec 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011009 | OEIS-A273570

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 798", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 862 also generates this sequence. - _Lars Blomberg_, Jul 21 2016 Conjecture: a(n) = 2*4^n - 6*2^n + 13, n>1. - _Lars Blomberg_, Jul 21 2016 Conjectures from _Colin Barker_, Dec 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011010 | OEIS-A273611

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 809", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = (2150*4^(n-4) - 4*3^(n-4) - 12*2^n - 8)/3, n>4. - _Lars Blomberg_, Jul 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011011 | OEIS-A273616

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with (3*x^2+13*y^2)*z a square, where x,y,z,w are nonnegative integers.

Conjecture: For each ordered pair (a,b) = (3,13), (5,11), (15,57), (15,165), (138,150), any natural number can be written as x^2 + y^2 + z^2 + w^2 with (a*x^2+b*y^2)*z a square, where x,y,z,w are nonnegative integers. For more conjectural refinements of Lagrange's four-square theorem, see the author's preprint arXiv:1604.06723.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011012 | OEIS-A273675

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 833", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011013 | OEIS-A273676

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 833", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 849 also generates this sequence. - _Lars Blomberg_, Jul 23 2016 Conjecture: a(n) = A270222(n) for n>1. - _R. J. Mathar_, May 30 2016 Conjecture: a(n) = 3*4^n - 4*2^n + 1, n>1. - _Lars Blomberg_, Jul 23 2016 Conjectures from _Colin Barker_, Dec 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011014 | OEIS-A273677

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 833", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011015 | OEIS-A273678

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 833", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011016 | OEIS-A273684

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 841", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 857 also generates this sequence. - _Lars Blomberg_, Jul 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011017 | OEIS-A273700

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 865", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 881 also generates this sequence. - _Lars Blomberg_, Jul 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011018 | OEIS-A273708

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 873", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 889 also generates this sequence. - _Lars Blomberg_, Jul 24 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011019 | OEIS-A273722

The number of valleys of width 1 (i.e., DHU configurations, where U=(0,1), H(1,0), D=(0,-1)) in all bargraphs of semiperimeter n (n>=2).

Conjecture: -(n-6) *(2*n-7) *(2*n-9) *(n+1)*a(n) +2*(n-3) *(2*n-9) *(4*n^2-24*n+21)*a(n-1) +2*(-4*n^4+56*n^3-289*n^2+651*n-547) *a(n-2) +4*(2*n-5) *(n-4)*a(n-3) -(n-4) *(n-5) *(2*n-5) *(2*n-7) *a(n-4)=0. - _R. J. Mathar_, Jun 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jun 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011020 | OEIS-A273724

Place n equally-spaced points around a circle, labeled 0,1,2,...,n-1. For each i = 0..n-1 such that 3i != i mod n, draw an (undirected) chord from i to (3i mod n). Then a(n) is the total number of distinct chords.

a(n) = n-1 if n>0 is odd, n-2 if n == +-2 (mod 8), n-3 if n == 4 (mod 8), and n-5 if n == 0 (mod 8). These formulas are easily established by observing that the chord at i is missing if 2i == 0 mod n, and the chords starting at i and at 3i coincide if 8i == 0 mod n. The formulas then imply that the g.f. is 4+x^2/(1-x)^2-(4+x^2+2*x^4+x^6)/(1-x^8), which can be rewritten as (5*x^63*x^5+2*x^4+3*x^2-x+2)*x^3/((1-x)*(1-x^8)). (This g.f. was conjectured by _Colin Barker_.) - _Brooke Logan_ and _N. J. A. Sloane_, Jun 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, May 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011021 | OEIS-A273743

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 901", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011022 | OEIS-A273744

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 901", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011023 | OEIS-A273745

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 901", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011024 | OEIS-A273747

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 918", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 982 also generates this sequence. - _Lars Blomberg_, Jul 25 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011025 | OEIS-A273766

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 913", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011026 | OEIS-A273767

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 913", based on the 5-celled von Neumann neighborhood.

Conjecture: Rules 921, 929, 937, 945 and 953 also generate this sequence. - _Lars Blomberg_, Jul 24 2016 Conjecture: a(n) = (2^(n+1) - 1)^2, n>2. - _Lars Blomberg_, Jul 24 2016 Conjectures from _Colin Barker_, Jul 24 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011027 | OEIS-A273768

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 913", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011028 | OEIS-A273769

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 913", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 29 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011029 | OEIS-A273780

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 929", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 30 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011030 | OEIS-A273781

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 929", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 30 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011031 | OEIS-A273782

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 929", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 30 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011032 | OEIS-A273789

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 931", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 31 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011033 | OEIS-A273790

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 931", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 31 2016: (Start) Conjectured e.g.f.: 4 + exp(x)*(4*x^3/3 + 8*x^2 + 9*x - 3). - _Stefano Spezia_, Dec 30 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011034 | OEIS-A273791

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 931", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 31 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011035 | OEIS-A273796

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 942", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = 2*4^n - 16*3^(n-2) + 2^n + 1, n>1. - _Lars Blomberg_, Jul 25 2016 Conjectures from _Colin Barker_, Dec 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011036 | OEIS-A273798

Numbers of the form 2^m*p where p is prime and p < 2^m.

Conjecture: a subsequence of A116882;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, May 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011037 | OEIS-A273821

Triangle read by rows: T(n,k) is the number of 123-avoiding permutations p of [n] (A000108) such that k is maximal with the property that the k largest entries of p, taken in order, avoid 132.

G.f.: Sum_{n>=1, 1<=k<=n} T(n,k) x^n y^k = C(x) - 1 + ((1 - y) (1 - x y) (1 - (1 - x y)C(x)))/((1 - 2 x y) (1 - y + x y^2)) where C(x) = 1 + x + 2x^2 + 5x^3 + ... is the g.f. for the Catalan numbers A000108 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Callan_, May 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011038 | OEIS-A273826

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^2 with x*y + y*z + z*w a fourth power, where x is a positive integer, y is a nonnegative integer, and z and w are integers.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 16^k*m (k = 0,1,2,... and m = 1, 14, 56, 91, 184, 329, 355, 1016). For more conjectural refinements of Lagrange's four-square theorem, see the author's preprint arXiv:1604.06723.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011039 | OEIS-A273828

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 950", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 1014 also generates this sequence. - _Lars Blomberg_, Jul 25 2016 Conjecture: a(n) = 2*4^n - 2*2^n - 3, n>2. - _Lars Blomberg_, Jul 25 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011040 | OEIS-A273831

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 961", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jun 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011041 | OEIS-A273832

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 961", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 977 also generates this sequence. - _Lars Blomberg_, Jul 25 2016 Conjecture: a(n) = A269911(n) for n>1. - _R. J. Mathar_, Jun 02 2016 Conjecture: a(n) = 4*4^n - 4*2^n - 3, n>1. - _Lars Blomberg_, Jul 25 2016 Conjectures

from _Colin Barker_, Dec 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011042 | OEIS-A273833

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 961", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jun 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011043 | OEIS-A273834

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 961", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jun 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011044 | OEIS-A273847

Number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 969", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jun 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011045 | OEIS-A273848

Number of active (ON, black) cells at stage 2^n-1 of the two-dimensional cellular automaton defined by "Rule 969", based on the 5-celled von Neumann neighborhood.

Conjecture: Rule 985 also generates this sequence. - _Lars Blomberg_, Jul 25 2016 Conjecture: a(n) = 4*4^n - 4*2^n + 1, n>2. - _Lars Blomberg_, Jul 25 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011046 | OEIS-A273849

Partial sums of the number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 969", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jun 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011047 | OEIS-A273850

First differences of number of active (ON, black) cells in n-th stage of growth of two-dimensional cellular automaton defined by "Rule 969", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jun 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011048 | OEIS-A273871

Primes p such that (4^(p-1)-1) == 0 mod ((p-1)^2+1).

Conjecture 1: also primes p such that ((4^k)^(p-1)-1) == 0 mod ((p-1)^2+1) for all k >= 0. Conjecture 2: supersequence of Fermat primes (A019434).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jun 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011049 | OEIS-A273875

Number of ordered ways to write n as w^2 + x^2 + y^2 + z^2 with x*y + 2*y*z + 4*z*x a nonnegative cube, where w,x,y,z are integers with w >= 0 and x > 0.

Conjecture: (i) a(n) > 0 for all n > 0. For more conjectural refinements of Lagrange's four-square theorem, see the author's preprint arXiv:1604.06723.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011050 | OEIS-A273905

Number of symmetric bargraphs having semiperimeter n (n>=2).

Conjecture D-finite with recurrence (n+1)*a(n) +2*(-1)*a(n-1) +2*(-2*n+3)*a(n-2) +2*(n-6)*a(n-4) +6*(1)*a(n-5) -2*a(n-6) +(n-9)*a(n-8)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_ and _Sergi Elizalde_, Jun 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011051 | OEIS-A273915

Number of ordered ways to write n as w^5 + x^2 + y^2 + z^2, where w,x,y,z are nonnegative integers with x <= y <= z.

Let c be 1 or 4. Then any nonnegative integer n can be written as c*w^5 + x^2 + y^2 + z^2 with w,x,y,z nonnegative integers. We now prove this by induction on n. For n < 2^(10) this can be verified directly via a computer. If 2^(10) divides n, then by the induction hypothesis we can write n/2^(10) as c*w^5 + x^2 + y^2 + z^2 with w,x,y,z, nonnegative integers, and hence n = c*(2^2*w)^5 + (2^5*x)^2 + (2^5*y)^2 + (2^5*z)^2. If n is not of the form 4^k*(8m+7) with k and m nonnegative integers, then n is the sum of three squares and hence n = c*0^5 + x^2 + y^2 + z^2 for some integers x,y,z. When n = 4^k*(8m+7) > 2^(10) with k < 5, it is easy to see that n - c*1^5 or n - c*2^5 is the sum of three squares. For some conjectural refinements of Lagrange's four-square theorem, one may consult the author's preprint arXiv:1604.06723

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011052 | OEIS-A273917

Number of ordered ways to write n as w^2 + 3*x^2 + y^4 + z^5, where w is a positive integer and x,y,z are nonnegative integers.

Conjectures: Conjectures a(n) > 0 and (ii) verified up to 10^11. - _Mauro Fiorentini_, Jul 19 2023 See also A262813, A262857, A270566, A271106 and A271325 for some other conjectures on representations.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011053 | OEIS-A273926

Given G(x) such that G(G(x)^2 - G(x)^3) = x^2, then G(x) = Sum_{n>=1} A273925(n)*x^n / 2^a(n).

a(2*n-1) = A120738(n-1) = 4*(n-1) - A000120(n-1), for n>=0 (conjecture). a(2*n) = A101925(n-1) = A005187(n-1) + 1, for n>=0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011054 | OEIS-A273927

Absolute difference between A000290(n) and the nearest term of A000578.

Marshall Hall, Jr. conjectured that there exists a constant c > 0 such that for any integers x and y with y^2 != x^3 the relation abs(y^2 - x^3) > c*sqrt(abs(x)) holds.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Jun 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011055 | OEIS-A273959

Decimal expansion of 'C', an auxiliary constant defined by D. Broadhurst and related to Bessel moments (see the referenced paper about the elliptic integral evaluations of Bessel moments).

Also equals s(5,1)/Pi^2 (where the Bessel moment s(5,1) is Integral_{0..inf} x I_0(x) K_0(x)^4 dx), a conjectural equality checked by the authors to 1200 decimal places.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Jun 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011056 | OEIS-A273980

Numbers k such that the number of digits in the numerator of the k-th convergent of the continued fraction 1+1/(2+1/(2+1/(2+...))) = sqrt(2) exceeds the number of digits in the denominator.

Conjecture: an integer k is in the sequence iff s(k)/10^floor(log_10(s(k))) is in the interval [1, sqrt(2)] where s(k) = ((1+sqrt(2))^(k+1))/2. (In other words, k is in the sequence if and only if ((1+sqrt(2))^(k+1))/2, expressed in the form x*10^m where m is the integer such that 1 <= x < 10, satisfies x < sqrt(2).) (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ogetay Kayali_, Nov 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011057 | OEIS-A273984

Decimal expansion of the odd Bessel moment s(5,1) (see the referenced paper about the elliptic integral evaluations of Bessel moments).

Equals Pi^2 C (conjectural, where C is A273959).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Jun 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011058 | OEIS-A273985

Decimal expansion of the odd Bessel moment s(5,3) (see the referenced paper about the elliptic integral evaluations of Bessel moments).

Equals Pi^2 (2/15)^2 (13 C - 1/(10 C)) (conjectural, where C is A273959).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Jun 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011059 | OEIS-A273986

Decimal expansion of the odd Bessel moment s(5,5) (see the referenced paper about the elliptic integral evaluations of Bessel moments).

Equals Pi^2 (4/15)^3 (43 C - 19/(40 C)) (conjectural, where C is A273959).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Jun 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011060 | OEIS-A273987

Smallest Riesel number to base n.

a(2), a(3), a(5), a(6), a(7), a(10), a(15), a(22), a(23), a(30), ... are only conjectural (see links).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tim Johannes Ohrtmann_, Jun 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011061 | OEIS-A273995

Even numbers with a unique representation as the difference of two primes, each of which is a member of a pair of twin primes, and one of which is smaller than the even number under consideration.

Can any even number n > 2 be so written (perhaps not uniquely) as the difference of two (unrelated) twins, one of which is smaller than n? (T. S. Van Kortryk conjectures there are, if any, only a finite number of even integers such that this is not true.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Curtright_, Jun 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011062 | OEIS-A274002

Primes p of the form (q-1)^2+1 that are a divisor of 4^(q-1)-1 where q is prime.

Conjecture: also primes p of the form (q-1)^2+1, where q = prime, that are a divisor of (4^k)^(q-1)-1 for all k>=0. Example: 17 = (5-1)^2+1 is a term because 5 is prime and divides (4^k)^(5-1)-1 for all k>=0: 0/17 = 0 (k=0); 255/17 = 15 (k=1); 65535/17 = 3855 (k=2); 16777215/17 = 986895 (k=3); 4294967295/17 = 252645135 (k=4); 1099511627775/17 = 64677154575 (k=5); ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jun 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011063 | OEIS-A274004

First differences of A002960.

Conjecture: a(n) = A274089(n+2). - _Omar E. Pol_, Aug 10 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alisa Ediger_, Jun 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011064 | OEIS-A274007

Number of ordered ways to write n as x^5 + 2*y^5 + z*(3*z-1)/2 + w*(3*w+1)/2, where x,y,z,w are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 11, 57, 198, 229, 232, 1168, 2624.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011065 | OEIS-A274010

Boris Stechkin function: a(n) is the number of m with 2 <= m <= n and floor(n(m-1)/m) divisible by m-1.

Conjecture: a(n) = tau(n) + tau(n-1) - 2, for n>=2. - _Ridouane Oudra_, Feb 28 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Jun 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011066 | OEIS-A274039

Expansion of (x^4 + x^10) / (1 - 2*x + x^2).

This g.f. was incorrectly conjectured by Plouffe in his 1992 disseration to be the g.f. for A005377.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sean A. Irvine_, Jun 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011067 | OEIS-A274052

Number of factor-free Dyck words with slope 5/2 and length 7n.

Conjectural o.g.f.: Let E(x) = exp(Sum_{n >= 1} binomial(7*n, 2*n)*x^n/n). Then A(x) = (x/series reversion of x*E(x))^(1/7) = 1 + 3*x + 13*x^2 + 94*x^3 + ... Equivalently, [x^n](A(x)^(7*n)) = binomial(7*n, 2*n) for n = 0,1,2,... . - _Peter Bala_, Jan 01 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael D. Weiner_, Jun 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011068 | OEIS-A274055

Relative of Hofstadter Q-sequence: a(n) = n for 1 <= n <= 42; a(n) = a(n-a(n-1)) + a(n-a(n-2)) for n > 42.

If the index is at least 24863, then a(5n) = 24860*A272613(n-4972), a(5n+1) = 4, a(5n+2) = 5*A272611(n-4972), a(5n+3) = 5*A272611(n-4971), a(5n+4) = 5*A272612(n-4971). This pattern lasts as long as A272611 exists (which is conjectured to be forever).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nathan Fox_, Nov 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011069 | OEIS-A274100

Number of partitions of 2^n into at most four parts.

Conjectures from _Colin Barker_, Jun 12 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011070 | OEIS-A274138

Triangle read by rows: Domination number for rectangular queens' graph Q(n,m), 1 <= n <= m.

Chessboard 8 X 11 is of special interest, because it cannot be dominated by 5 queens, although the larger boards 9 X 11, 10 X 11 and 11 X 11 are. It is conjectured that 8 X 11 is the only counterexample of this kind of monotonicity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sandor Bozoki_, Jun 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011071 | OEIS-A274139

a(n) = 2^A000265(n) = 2^numerator(n/2^n), a sequence related to Oresme numbers.

a(n) = 2*A082392(n)^2 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_ and _Paul Curtz_, Jun 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011072 | OEIS-A274163

Number of real integers in n-th generation of tree T(4i) defined in Comments.

For each integer k > 0, let s(k,n) be the number of integers in the n-th generation of T(k*i). Conjecture: there is a limiting sequence S(n) as k increases, and S(n) = F(n) for n >= 1, where F = A000045 (Fibonacci numbers).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011073 | OEIS-A274212

The factorization of n contains only lesser of twin primes.

Arithmetic conjecture: the equation a(n+1) - a(n) = 2r has infinitely many solutions for any fixed integer value r >= 1. Analytic conjecture: a(n) is asymptotic to C*n*log(n)^2 for a constant C > 0.2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jun 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011074 | OEIS-A274232

Number of partitions of 2^n into at most three parts.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Barker_, Jun 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011075 | OEIS-A274242

Numbers whose factorization contains only the greater members of twin prime pairs.

Arithmetic conjecture: the equation a(n+1) - a(n) = 2r has infinitely many solutions for any fixed integer value r >= 1. Analytic conjecture: a(n) is asymptotic to D*n*log(n)^2 for a constant D > 0.4 hence lim_{n->oo} a(n)/A274212(n) exists (see link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jun 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011076 | OEIS-A274244

Number of factor-free Dyck words with slope 7/2 and length 9n.

Conjectural o.g.f.: Let E(x) = exp(Sum_{n >= 1} binomial(9*n, 2*n)*x^n/n). Then A(x) = (x/series reversion of x*E(x))^(1/9) = 1 + 4*x + 34*x^2 + 494*x^3 + ... Equivalently, [x^n](A(x)^(9*n)) = binomial(9*n, 2*n) for n = 0,1,2,... . - _Peter Bala_, Jan 01 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael D. Weiner_, Jun 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011077 | OEIS-A274256

Number of factor-free Dyck words with slope 9/2 and length 11n.

Conjectural o.g.f.: Let E(x) = exp(Sum_{n >= 1} binomial(11*n,2*n)*x^n/n). Then A(x) = (x/series reversion of x*E(x))^(1/11) = 1 + 5*x + 70*x^2 + 1696*x^3 + ... Equivalently, [x^n](A(x)^(11*n)) = binomial(11*n, 2*n) for n = 0,1,2,... . - _Peter Bala_, Jan 01 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael D. Weiner_, Jun 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011078 | OEIS-A274257

Number of factor-free Dyck words with slope 4/3 and length 7n.

Conjectural o.g.f.: Let E(x) = exp(Sum_{n >= 1} binomial(7*n, 3*n)*x^n/n). Then A(x) = (x/series reversion of x*E(x))^(1/7) = 1 + 5*x + 52*x^2 + 880*x^3 + ... Equivalently, [x^n](A(x)^(7*n)) = binomial(7*n, 3*n) for n = 0,1,2,.... - _Peter Bala_, Jan 01 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael D. Weiner_, Jun 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011079 | OEIS-A274258

Number of factor-free Dyck words with slope 5/3 and length 8n.

Conjectural o.g.f.: Let E(x) = exp(Sum_{n >= 1} binomial(8*n, 3*n)*x^n/n). Then A(x) = (x/series reversion of x*E(x))^(1/8) = 1 + 7*x + 133*x^2 + 4140*x^3 + ...

Equivalently, [x^n](A(x)^(8*n)) = binomial(8*n, 3*n) for n = 0,1,2,... . - _Peter Bala_, Jan 01 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael D. Weiner_, Jun 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011080 | OEIS-A274259

Number of factor-free Dyck words with slope 7/3 and length 10n.

Conjectural o.g.f.: Let E(x) = exp(Sum_{n >= 1} binomial(10*n, 3*n)*x^n/n). Then A(x) = (x/series reversion of x*E(x))^(1/10) = 1 + 12*x + 570*x^2 + 44689*x^3 + ... Equivalently, [x^n](A(x)^(10*n)) = binomial(10*n, 3*n) for n = 0,1,2,... . - _Peter Bala_, Jan 03 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael D. Weiner_, Jun 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011081 | OEIS-A274260

Forbidden residues of the greedy prime offset sequence.

The greedy prime offset sequence, A135311, is the closepacked integer sequence, starting with 0, such that for no prime p does the sequence form a complete system of residues modulo p. Instead, at least one residue must be missing for p, this is the (conjectured to be unique) "forbidden residue" for p. The first few terms of the greedy sequence are 0, 2, 6, 8, 12, 18. For the first three primes: 2, 3, 5, the forbidden residues are, respectively: 1, 1, 4. More generally, a(n) gives the forbidden residue for the n-th prime number. Every prime, it appears, has a unique forbidden residue, but this is unproven as far as I know. If this is true then every prime has an "exhaustion number" which is the number of terms of the greedy sequence needed to exhaust all the other residues and determine which one is forbidden; see A274261.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. Michael Perry_, Jun 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011082 | OEIS-A274261

Exhaustion numbers for the greedy prime offset sequence A135311.

The greedy prime offset sequence, A135311, is the close-packed integer sequence, starting with 0, such that for no prime p does the sequence form a complete system of residues modulo p. Instead, at least one residue must be missing for p, this is the (conjectured to be unique) "forbidden residue" for p. Every prime, it appears, has a unique forbidden residue. If this is true then every prime has an "exhaustion number" which is the number of terms of the greedy sequence needed to exhaust all the other residues and determine which one is forbidden. The uniqueness of the forbidden residue for any individual prime can be verified by calculation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. Michael Perry_, Jun 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011083 | OEIS-A274271

Number of partitions of 3^n into at most four parts.

Conjectures (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Barker_, Jun 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011084 | OEIS-A274272

Number of partitions of 5^n into at most four parts.

Conjectures (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Barker_, Jun 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011085 | OEIS-A274274

Number of ordered ways to write n as x^3 + y^2 + z^2, where x,y,z are nonnegative integers with y <= z.

Conjecture: Let n be any nonnegative integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011086 | OEIS-A274295

a(n) = n+binomial(2*n-6,n-3)+binomial(2*n-5,n-3)+binomial(n-1,n-3)+Sum_{i=1..n-3} (binomial(n+i-3,n-3)+2*n-i-5).

Conjecture: (-n+2)*a(n) +(7*n-18)*a(n-1) +14*(-n+3)*a(n-2) +2*(3*n-2)*a(n-3) +(11*n-90)*a(n-4) +(-13*n+102)*a(n-5) +2*(2*n-17)*a(n-6)=0. - _R. J. Mathar_, Oct 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011087 | OEIS-A274315

First row of infinite Sudoku-type array A269526.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011088 | OEIS-A274318

Main diagonal of infinite Sudoku-type array A269526.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011089 | OEIS-A274332

Team size n for which there exists a balanced tournament for 2n+1 players so that in 2n+1 matches each player plays exactly n-1 times with and n times against each other player.

It is conjectured that this sequence is identical to A005097 (ref. Kohen link). Conjectured design scheme for Team 1 (N:= 2n+1; here players count from 0..2n): X, X+1 (mod N), X+1+2 (mod N), X+1+2+3 (mod N), ...; X = 0..2n (match number). Resting player: (X + (n*(n+1)/2) (mod N).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Steyer_, Jun 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011090 | OEIS-A274334

Numbers whose cube is the sum of 4 positive cubes.

Conjecture: a(n) = n + 21 for all n > 50. - _Charles R Greathouse IV_, Jan 14 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Jun 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011091 | OEIS-A274377

E.g.f. satisfies: A(x)^A(x) = exp(2*x) * A(-x)^A(-x).

a(2*n+1) = 6 (mod 10) for n>1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011092 | OEIS-A274397

Positive integers m such that sigma(m) is divisible by 5.

lim_{n->oo} a(k)/k = 2 (conjectured; cf. Examples).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jul 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011093 | OEIS-A274479

G.f. satisfies: A(x)^2 = A(x^2/(1 - 2*x - 4*x^2)).

a(n) = 1 (mod 3) for n>=1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011094 | OEIS-A274521

Number of odd partitions in the multiset of intersections of the set of partitions of n with itself; also number of distinct partitions in that multiset.

By numerical experimentation, it seems a(n) is the convolution of A000009 (with offset 1) and A054440. (conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _George Beck_, Jun 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011095 | OEIS-A274600

Coefficients in an asymptotic expansion of sequence A002893.

Conjecture: a(n) ~ (2/log(3))^n * (n-1)! / (Pi*sqrt(3)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Nov 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011096 | OEIS-A274605

Numbers k such that there is no prime between the k-th perfect power and the (k+1)-st perfect power.

It has been conjectured that 23268 is the last term of the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Jun 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011097 | OEIS-A274650

Triangle read by rows: T(n,k), (0 <= k <= n), in which each term is the least nonnegative integer such that no row, column, diagonal, or antidiagonal contains a repeated term.

Omar E. Pol's conjecture that every column and every diagonal of the triangle is a permutation of the nonnegative integers is true: see the link. - _N. J. A. Sloane_, Jun 07 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jul 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011098 | OEIS-A274651

Triangle read by rows: T(n,k), (1<=k<=n), in which each term is the least positive integer such that no row, column, diagonal, or antidiagonal contains a repeated term.

Omar E. Pol's conjecture that every column and every diagonal of the triangle is a permutation of the positive integers is true: see the link in A274650 (duplicated below). - _N. J. A. Sloane_, Jun 07 2017 Conjecture: as n gets sufficiently large, all coefficients j>4 will appear in a repeating pattern, populating all rows and diagonals around smaller j's near the "middle diagonal" (while I can offer no formal proof, it appears very likely that this is the case).

(End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jul 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011099 | OEIS-A274719

Expansion of Product_{k >= 1} (1-q^(2*k)).

Conjecture: |a(n)| = A089806(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _George Beck_, Jul 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011100 | OEIS-A274720

Odd numbers n such that n is not coprime to the multiplicative order of 2 mod n.

Let G(i, j) = gcd(2^j - 1, j^(2^i) - 1). I conjectured that an odd positive integer n is a term of this sequence if and only if n is not of the form G(i, j). _Jinyuan Wang_ (pers. comm.) proved the direct implication and the fact that, if n is not a term of this sequence, then n divides G(i, j) for some i and j. - _Lorenzo Sauras Altuzarra_, Sep 04 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Jul 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011101 | OEIS-A274733

Number of odd partitions in the multiset of intersections of the set of partitions of n with itself three times; also number of distinct partitions in that multiset.

By numerical experimentation, it seems a(n) is the convolution of A000009 (with offset 1) and A260664. (conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _George Beck_, Jul 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011102 | OEIS-A274786

Diagonal of the rational function 1/(1 - (wxz + wy + wz + xy + xz + y + z)).

Conjecture: a(n) = [x^n] (1 + x)^(2*n) * P(n,(1 + x)/(1 - x))^2, where P(n,x) denotes the n-th Legendre polynomial. Cf. A005260(n) = [x^n] (1 - x)^(2*n) * P(n,(1 + x)/(1 - x))^2, due to Carlitz. - _Peter Bala_, Sep 21 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011103 | OEIS-A274882

a(n) is the largest coefficient of q-binomial(2*n, n) / q-binomial(n+1, 1), which are the q-Catalan polynomials.

Conjecture: a(n) ~ sqrt(3) * 2^(2*n) / (Pi * n^3). - _Vaclav Kotesovec_, Jan 06 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jul 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011104 | OEIS-A274946

Boyd's Pisot-like sequence F(0,5,11).

Conjectures from _Colin Barker_, Jul 26 2016: (Start) Note the warning in A010925 from Pab Ter (pabrlos(AT)yahoo.com), May 23 2004: [A010925] and other examples show that it is essential to reject conjectured generating functions for Pisot sequences until a proof or reference is provided. - _N. J. A. Sloane_, Jul 26 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011105 | OEIS-A274987

Primes p such that A274601(p) is a prime.

It is conjectured that the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Nov 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011106 | OEIS-A275001

Expansion of 1/(1 - Sum_{k>=1} x^(prime(k)^2)).

Conjecture(1): every number > 23 is the sum of at most 8 squares of primes. Conjecture(2): every number > 131 can be represented as a sum of 13 squares of primes. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Dec 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011107 | OEIS-A275011

a(1) = 2; for n > 1, a(n) is the least prime p > a(n-1) such that none of p-a(1), ..., p-a(n-1) is a square.

I conjecture that the sequence is infinite (in fact for any initial term).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Nov 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011108 | OEIS-A275022

Primes not in A001097 and not in A275021 that occur in pairs of the form {p, p+6}.

Conjecture: The sequence has infinitely many terms and the sum of their reciprocals converges.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimitris Valianatos_, Nov 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011109 | OEIS-A275027

a(n) = Sum_{k=0..n} C(n,k)^2*C(n-k,k), where C(n,k) denotes the binomial coefficient n!/(k!*(n-k)!).

Conjecture: For any prime p > 5 and positive integer n, the number (a(p*n)-a(n))/(p*n)^3 is always a p-adic integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011110 | OEIS-A275047

Diagonal of the rational function 1/(1-(1+w)(xy + xz + yz)) [even-indexed terms only].

We conjecture that the sequences {a(i,n) : n >= 0}, i >= 2, also satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5, and positive integers n and r. Cf. A362725 and A362732. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011111 | OEIS-A275051

Expansion of 3F2([1/9, 4/9, 5/9], [1/3,1], 729*x).

"One may consider the following conjecture: all the irreducible factors of the minimal order linear differential operator annihilating a diagonal of a rational function should be homomorphic to their adjoint (possibly on an algebraic extension). [...] "If our conjecture above was correct, this would be a way to show that the series cannot be the diagonal of a rational function." (See Boukraa link.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011112 | OEIS-A275052

G.f.: 3F2([1/7, 2/7, 4/7], [1/2, 1], 2401 x).

"Other hypergeometric 'blind spots' for Christol’s conjecture" - (see Bostan link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011113 | OEIS-A275053

G.f.: 3F2([1/9, 2/9, 7/9], [2/3,1], 729x).

"Other hypergeometric 'blind spots' for Christol’s conjecture" - (see Bostan link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011114 | OEIS-A275054

G.f.: 3F2([1/9, 2/9, 8/9], [2/3,1], 729 x).

"Other hypergeometric 'blind spots' for Christol’s conjecture" - (see Bostan link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011115 | OEIS-A275083

Positive integers congruent to 0 or 1 modulo 4 that cannot be written as x^3 + y^2 + z^2 with x,y,z nonnegative integers.

Conjecture: (i) The sequence has totally 18 terms as listed. Let d be 2 or 6. Clearly, n-d is congruent to 0 or 1 modulo 4 if n is congruent to 2 or 3 modulo 4. So part (i) of the conjecture essentially implies that for each n = 0,1,2,... either n or n-d can be written as x^3 + y^2 + z^2 with x,y,z nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011116 | OEIS-A275113

a(n) is the minimal number of squares needed to enclose n squares with a wall so that there is a gap of at least one cell between the wall and the enclosed cells.

Inspired by beehive construction in which wax is used in the most efficient way. This problem is likened to construction of a fence around a house with minimum materials and maximum enclosed area. I conjectured that a specific house pattern shall be selected. See illustration in links. If the conjecture in A261491 is true (i.e., A261491(n) is the number of squares required to enclose n squares without a gap), then a(n) = A261491(n) + 8. - _Charlie Neder_, Jul 11 2018 The conjecture in A261491 holds through a(16). - _David Consiglio, Jr._, Nov 10 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kival Ngaokrajang_, Jul 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011117 | OEIS-A275115

Least prime of the form x^2 + n*y^2 with x>0 and y>0.

Not all 'd's are squares, such as 24, 132, 258, 1032, 1167, 1518, 2103, 2472, 2652, 2706, 5793. It is conjectured that this list is complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_ and _Robert G. Wilson v_, Jul 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011118 | OEIS-A275122

Pascal's hexagonal pyramid, read by slices, with each slice read by rows.

Odd terms in layer x, where x is 1 less than a power of 2, form a hexaflake (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aresh Pourkavoos_, Jul 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011119 | OEIS-A275138

Number of n X 4 0..2 arrays with no element equal to any value at offset (-2,-2) (-1,-2) or (0,-1) and new values introduced in order 0..2.

Conjectures from _Colin Barker_, Jan 31 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011120 | OEIS-A275139

Number of n X 5 0..2 arrays with no element equal to any value at offset (-2,-2) (-1,-2) or (0,-1) and new values introduced in order 0..2.

Conjectures from _Colin Barker_, Jan 31 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011121 | OEIS-A275150

Number of ordered ways to write n as x^3 + 2*y^2 + k*z^2, where x,y,z are nonnegative integers, k is 1 or 5, and k = 1 if z = 0.

Conjecture 1: a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 15, 79, 120, 218, 399, 454, 622, 725, 3240. Conjecture 2: For any positive integers a, b, c and integers i, j, k greater than one, there are infinitely many positive integers not in the set {a*x^i + b*y^j + c*z^k: x,y,z = 0,1,2,...}. - _Zhi-Wei Sun_, May 24 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011122 | OEIS-A275151

a(1) = 8; a(n) = 3*a(n-1) + 2*sqrt(2*a(n-1)*(a(n-1)-7)) - 7 for n > 1.

Conjectures from _Colin Barker_, Jul 19 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Manuel López Holgueras_, Jul 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011123 | OEIS-A275168

Positive integers not of the form x^3 + 3*y^2 + z^2 with x,y,z nonnegative integers.

Conjecture: The sequence has totally 150 terms as listed in the b-file the largest of which is 182842. Thus any integer n > 182842 can be written as x^3 + 3*y^2 + z^2 with x,y,z nonnegative integers. See also A275169 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011124 | OEIS-A275169

Positive integers not in the form x^3 + 2*y^2 + z^2 with x,y,z nonnegative integers.

Conjecture: The sequence has totally 174 terms as listed in the b-file the largest of which is 375565. This implies the conjecture in A275150. We note that the sequence contains no term greater than 375565 and not exceeding 10^6. See also A275168 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011125 | OEIS-A275206

Expansion of (A(x)^2 - A(x^2))/2 where A(x) = A000108(x) - 1.

Substituting A(x) = Sum_{n >= 1} c(n)*x^n in the above expression, we get (after some algebra) that a(n) = (-2*c(n) + c(n+1) - [(n mod 2) == 0]*c(n/2))/2 for n >= 1. It is then easy to prove that a(n) = 2*A050182(n-2) = (1/n)*(binomial(2*n, n-2) - [(n mod 2) == 0]*binomial(n, (n/2) - 1)) for n >= 2, thus proving the conjecture below. (End) Conjecture: a(n) = 2*A050182(n-2) for n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jul 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011126 | OEIS-A275207

Expansion of (A(x)^2+A(x^2))/2 where A(x) = A001006(x).

Conjecture: a(2n+1) = A026940(n+1). Conjecture D-finite with recurrence -3*(n+4)*(n+3)*(29*n-32)*a(n) +10*(29*n-40)*(n+3)*(n+2)*a(n-1) +2*(n+1)*(149*n^2 +208*n-450)*a(n-2) -2*n*(559*n^2 -381*n-1630)*a(n-3) +4*(-68*n^3 +531*n^2 -904*n+351)*a(n-4) +2*(103*n^3-1701*n^2+5330*n -3600)*a(n-5) +18*(11*n^3 -209*n^2 +834*n -778)*a(n-6) +6*n*(269*n-830)*(n-5)*a(n-7) +9*(n-5)*(n-6)*(95*n-134)*a(n-8)=0. - _R. J. Mathar_, Mar 07 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jul 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011127 | OEIS-A275229

Number of 3 X n 0..2 arrays with no element equal to any value at offset (-2,-1) (-2,1) or (-1,0) and new values introduced in order 0..2.

Conjectures from _Colin Barker_, Feb 02 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011128 | OEIS-A275235

Number of primes between n and n+log(n)^2.

According to Schinzel's formulation of Cramer's conjecture, a(n) > 0 for n > 7; see Guy.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011129 | OEIS-A275288

Least k such that there exists a sequence b_1 < b_2 < ... < b_t = k that includes n and has a reciprocal sum of 1.

Conjectures: Conjecture (confirmed for all n <= 40000): for all n > 1, the lower bound given above is tight, i.e., a(n) = m*P where m and P are as defined above. (See Example section.) (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Aug 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011130 | OEIS-A275297

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^3 with x + 2*y a square, where x,y,z,w are nonnegative integers with z >= w.

Conjecture: a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 3, 7, 8, 11, 12, 15, 23, 24, 32, 39, 47, 71, 103, 120, 136, 159, 176, 183, 218, 359, 463. Compare this conjecture with Conjecture 5.1 of the author's preprint arXiv:1604.06723. See also A275298 and A275299 for similar conjectures. Let c be 1 or 2. By the conjecture in A272979, any n = 0,1,2,... can be written as x^2 + 2*y^2 + z^3 + 2*c^2*w^4 with x,y,z,w nonnegative integers, and hence n = x^2 + (y+c*w^2)^2 + (y-c*w^2)^2 + z^3 with (y+c*w^2)-(y-c*w^2) = 2*c*w^2. If n > 0 is not among the 174 terms in the b-file of A275169, then the conjecture in A275169 implies that n can be written as x^2 + y^2 + z^2 + w^3 with x - y = 0^2, where x,y,z,w are nonnegative integers. If n is among the 174 terms in the b-file of A275169, then we may use a computer to verify that n can be written as x^2 + y^2 + z^2 + w^3 with c*(x-y) a square, where x,y,z,w are nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011131 | OEIS-A275298

Number of ordered ways to write n as w^3 + x^2 + y^2 + z^2 with x - w a square, where x,y,z,w are nonnegative integers with y <= z > w.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 3, 4, 7, 8, 12, 16, 23, 24, 40, 47, 71, 167, 311, 599. See also A275297 and A275299 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011132 | OEIS-A275299

Number of ordered ways to write n as x^3 + y^2 + z^2 + w^2 with x + y - z a square, where x,y,z,w are nonnegative integers with y <= z.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 5, 8, 20, 24, 59, 71, 79, 119, 184, 575, 743, 764, 1471, 2759. See also A275297 and A275298 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011133 | OEIS-A275300

Number of ordered ways to write n as x^2 + y^2 + z^2 + w^3 with x + y + z a square, where x,y,z are integers with x >= |y| <= |z|, and w is a nonnegative integer.

Conjecture: a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 5, 12, 20, 23, 31, 47, 71, 103, 148, 164. See also A275297, A275298, A275299 and A272620 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011134 | OEIS-A275301

Number of ordered ways to write n as x^2 + y^2 + z^2 + 2*w^2 with x + 2*y a cube, where x,y,z,w are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 5, 6, 7, 8, 11, 14, 15, 30, 78, 90, 93, 106, 111, 117, 125, 223, 335. See also A275344 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011135 | OEIS-A275319

Numbers n such that n concatenated with n+1 is not a prime.

Conjecture: a(n) ~ n. - _Charles R Greathouse IV_, Jul 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Jul 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011136 | OEIS-A275322

Decimal expansion of AGM(1, sqrt(2))^2/Pi.

Conjecture: Equals Product_{n odd} (n/(n+2) if n == 1 (mod 4), (n+2)/n otherwise) = (1/3) * (5/3) * (5/7) * (9/7) * (9/11) * (13/11) * (13/15) * (17/15) * (17/19) * (21/19) * (21/23) * (25/23) * (25/27) * ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimitris Valianatos_, Jul 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011137 | OEIS-A275344

Number of ordered ways to write n as x^2 + y^2 + z^2 + 2*w^2 with x + 2*y + 3*z a square, where x,y,z,w are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 1, 3, 5, 7, 14, 15, 16, 25, 30, 33, 84, 169, 225. This is similar to the conjecture in A271518. It is known that any natural number can be written as x^2 + y^2 + z^2 + 2*w^2 with x,y,z,w nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011138 | OEIS-A275345

Characteristic polynomials of a square matrix based on A051731 where A051731(1,N)=1 and A051731(N,N)=0 and where N=size of matrix, analogous to the Redheffer matrix.

Conjecture: The largest absolute value of the eigenvalues of these characteristic polynomials appear to have the same prime signature in the factorization of the matrix sizes N.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Jul 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011139 | OEIS-A275386

The concatenation of |a(n)-a(n+1)| and a(n+1) is prime. Terms are distinct.

The sequence starts with a(1) = 1 and is always extended with the smallest integer not yet present that does not lead to a contradiction. This sequence is conjectured to be a permutation of the odd integers not ending with the digit "5".

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Nov 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011140 | OEIS-A275391

Least k such that n divides sigma(k^k) (k > 0).

Conjecture: a(n) <= n-2 for all n > 15, but a(n) = n-2 for infinitely many n. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Aug 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011141 | OEIS-A275409

Number of ordered ways to write n as 2*w^2 + x^2 + y^2 + z^2 with w + x + 2*y + 4*z a square, where w,x,y,z are nonnegative integers.

Conjecture: (i) a(n) > 0 except for n = 3, 10, and a(n) = 1 only for n = 0, 2, 7, 8, 9, 12, 14, 15, 22, 23, 24, 25, 36, 39, 44, 45, 60, 87, 98, 106, 110, 111, 183. See also A275344 and A275301 for related conjectures. We are able to show that each natural number can be written as x^2 + y^2 + z^2 + 2*w^2 with x,y,z,w integers such that x + y + z = t^2 for some t = 0, 1, 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011142 | OEIS-A275412

Least k such that phi(k*n) = phi(k*n+1), or 0 if no such k exists.

If a(n) = 0, then a(m*n) = 0 for all m > 0. If c is a divisor of a(n), then a(c*n) = a(n)/c. Conjecture: If A275337(n) = 0, then a(n) = 0. - _Chai Wah Wu_, Jul 27 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Jul 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011143 | OEIS-A275418

Numbers n such that n - 1 has exactly as many odd divisors as n + 1.

Conjectures: I don't find any more for conjecture #3 up to 10^10. - _Charles R Greathouse IV_, Aug 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jul 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011144 | OEIS-A275452

G.f.: 3F2([1/9, 4/9, 7/9], [1/3, 1], 729 x).

"Other hypergeometric 'blind spots' for Christol’s conjecture" - (see Bostan link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011145 | OEIS-A275453

G.f.: 3F2([1/9, 4/9, 7/9], [2/3, 1], 729 x).

"Other hypergeometric 'blind spots' for Christol’s conjecture" - (see Bostan link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011146 | OEIS-A275454

G.f.: 3F2([1/9, 4/9, 8/9], [2/3, 1], 729 x).

"Other hypergeometric 'blind spots' for Christol’s conjecture" - (see Bostan link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011147 | OEIS-A275455

G.f.: 3F2([1/9, 5/9, 8/9], [1/3, 1], 729 x).

"Other hypergeometric 'blind spots' for Christol’s conjecture" - (see Bostan link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011148 | OEIS-A275456

G.f.: 3F2([1/9, 7/9, 8/9], [1/3, 1], 729 x).

"Other hypergeometric 'blind spots' for Christol’s conjecture" - (see Bostan link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011149 | OEIS-A275457

G.f.: 3F2([2/9, 4/9, 5/9], [1/3, 1], 729 x).

"Other hypergeometric 'blind spots' for Christol’s conjecture" - (see Bostan link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011150 | OEIS-A275458

Expansion of g.f. hypergeom([4/9, 5/9, 7/9], [2/3, 1], 729 x).

"Other hypergeometric 'blind spots' for Christol’s conjecture" - (see Bostan link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011151 | OEIS-A275459

G.f.: 3F2([4/9, 5/9, 8/9], [2/3, 1], 729 x).

"Other hypergeometric 'blind spots' for Christol’s conjecture" - (see Bostan link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011152 | OEIS-A275460

G.f.: 3F2([2/9, 4/9, 7/9], [1/3, 1], 729 x).

"Other hypergeometric 'blind spots' for Christol’s conjecture" - (see Bostan link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011153 | OEIS-A275461

G.f.: 3F2([2/9, 5/9, 7/9], [2/3, 1], 729 x).

"Other hypergeometric 'blind spots' for Christol’s conjecture" - (see Bostan link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Jul 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011154 | OEIS-A275471

Number of ordered ways to write n as 4^k*(1+x^2+y^2)+z^2, where k,x,y,z are nonnegative integers with x <= y and x == y (mod 2).

Conjecture: a(n) > 0 except for n = 449. See also A275656, A275678 and A275738 for related conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011155 | OEIS-A275482

The sum of the last digit of a(n) and the first digit of a(n+1) is a divisor of a(n) and a(n+1) concatenated.

It is conjectured that this sequence is a permutation of the nonnegative numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Jul 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011156 | OEIS-A275492

Number of primitive weird numbers (A002975) between 10^n and 10^(n+1).

It is not known unconditionally whether there are infinitely many primitive weird numbers (PWN, A002975), although numerical data provides strong evidence: even the number of weird numbers of the form 2^k*p*q (A258882, A258333) seems to increase rapidly as k increases. Melfi has shown that Cramer's conjecture implies the infiniteness of PWN.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jul 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011157 | OEIS-A275505

Number of 5 X n 0..2 arrays with no element equal to any value at offset (-2,-1) (-2,0) or (-1,-1) and new values introduced in order 0..2.

Conjectures from _Colin Barker_, Feb 04 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jul 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011158 | OEIS-A275527

Number of distinct classes of permutations of length n under reversal and complement to n+1.

Conjecture: If n odd a(n) = (n - 1)!/2. If n even a(n) = (n - 2)!!*(1 + (n - 1)!!)/2. Conjectured formula is correct for n>1, see Fuller link. - _Martin Fuller_, Dec 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_, Jul 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011159 | OEIS-A275598

Primes p such that the number of odd divisors of p-1 is a prime q which is equal to the number of odd divisors of p+1.

Conjecture: this sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Aug 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011160 | OEIS-A275602

The concatenation of |a(n)-a(n+1)| and a(n+1) is prime and all such concatenations are different one from another. Terms are distinct.

This sequence is conjectured to be a permutation of the odd integers not ending with the digit "5".

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Nov 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011161 | OEIS-A275603

The concatenation of a(n) and |a(n)-a(n+1)| is prime. Terms are distinct. This is the lexicographically first sequence with this property.

This sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Nov 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011162 | OEIS-A275604

The concatenation of a(n) and |a(n)-a(n+1)| is prime and all such concatenations are different one from another. Terms are distinct.

This sequence is conjectured to be a permutation of the positive integers and is a variant of A275603 that shares with it the first 174 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Nov 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011163 | OEIS-A275648

Number of ordered ways to write n as x^2*(1+y^2+z^2)+w^2, where x is a positive integer and y,z,w are nonnegative integers with y <= z <= w.

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 3, 6, 7, 11, 14, 15, 23, 38, 47, 71, 77, 143, 152, 191, 608, 2^(2k+1) (k = 0,1,2,...). See A275656 for a stronger conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011164 | OEIS-A275652

a(n) = binomial(3*n,3*n/2)*binomial(2*n,n)*binomial(5*n/2,n/2)/binomial(n,n/2).

Conjecture: the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) hold for all primes p >= 5 and all positive integers n and k. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Aug 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011165 | OEIS-A275656

Number of ordered ways to write n as 4^k*(1+x^2+y^2)+z^2, where k,x,y,z are nonnegative integers with x <= y <= z.

Conjecture: a(n) > 0 for all n > 0. This is stronger than Lagrange's four-square theorem and the conjecture in A275648.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011166 | OEIS-A275667

Number of ON cells after n generations in a 2-dimensional "Odd-Rule" cellular automaton on triangular tiling.

If the first conjecture from the Formula section is true then the fact that the right border of the triangle (see Example) gives A000244 follows directly from it. If the second conjecture is true then the numbers just before the right border give A102900. a(0) = 1. Conjecture: a(2*t+1) = 3*a(t). Conjectures: a(8*t+6) = 3*a(4*t+2) + 4*a(2*t), a(8*t+2) = 3*a(4*t) + 4*a(2*t), a(4*t) = a(2*t). These conjectured formulas together give recurrent relations for a(n) for any n. Also, obviously a(2*n) = A247666(n). - _Andrey Zabolotskiy_, Aug 04 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kovba Alexey_, Aug 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011167 | OEIS-A275668

First occurrence of a value in A268755: a(i) = j iff A268755(j) = i-1 and A268755(j+1) = 0.

Conjectured to be infinite (this is equivalent to the conjecture that every positive integer eventually appears in A268755). (Proof given in comments of A268755).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _László Kozma_, Aug 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011168 | OEIS-A275673

List of numbers that are in a spoke of a hexagonal spiral.

Conjectures from _Colin Barker_, Aug 05 2016: (Start) Conjecture: a(n) = (n+4-3*floor((n+4)/6)-2)*floor((n+4)/6)+1. - _Luce ETIENNE_, May 25 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Aug 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011169 | OEIS-A275675

Number of ordered ways to write n as 4^k*(1+x^2+y^2)+5*z^2, where k,x,y,z are nonnegative integers with x <= y.

Conjecture: a(n) > 0 for all n > 0. See also A275656, A275676 and A275678 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011170 | OEIS-A275676

Number of ordered ways to write n as 4^k*(1+5*x^2+y^2) + z^2, where k,x,y,z are nonnegative integers with x <= y.

Conjecture: (i) a(n) > 0 for all n > 0. See also A275656, A275675 and A275678 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011171 | OEIS-A275678

Number of ordered ways to write n as 4^k*(1+4*x^2+y^2) + z^2, where k,x,y,z are nonnegative integers with x <= y.

Conjecture: (i) a(n) > 0 for all n > 0. See also A275656, A275675 and A275676 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011172 | OEIS-A275687

Difference between the smallest 10^n-digit member of a sexy prime quadruple and 10^(10^n - 1), or 0 if no such 10^n-digit number exists.

Conjecture: a(n) = A133449(10^(n-1)) for n >= 1. - _Jason Yuen_, Mar 07 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Aug 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011173 | OEIS-A275738

Number of ordered ways to write n as w^2 + x^2*(1+y^2+z^2), where w,x,y,z are nonnegative integers with x > 0, y <= z and y == z (mod 2).

Conjecture: For any n > 0, we have a(n) > 0, i.e., n can be written as w^2 + x^2*(1+(z-y)^2+(y+z)^2) = w^2 + x^2*(1+2*y^2+2*z^2), where w,x,y,z are nonnegative integers with x > 0 and y <= z. Moreover, any positive integer n not equal to 449 can be written as 4^k*(1+x^2+y^2) + z^2, where k,x,y,z are nonnegative integers with x == y (mod 2). See also A275648, A275656, A275675, A275676 and A275678 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011174 | OEIS-A275786

a(n) = Product_{d|n} T(d) where T(x) = x*(x+1)/2 = A000217(x) = x-th triangular number.

Conjecture: the sequence is injective (all terms of this sequence occur only once).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Aug 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011175 | OEIS-A275815

Maximum total number of possible moves that any number of queens of the same color can make on an n X n chessboard.

Conjecture: a(n) = 8(n-2)^2 for n >= 6. - _Alec Jones_, Nov 16 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Nov 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011176 | OEIS-A275824

Number of ways to write n as pen(x) + pen(y) + pen (z) + 2*pen(w), where x,y,z,w are nonnegative integers with x <= y <= z, and pen(k) denotes the pentagonal number k*(3*k-1)/2.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,...., and a(n) = 1 only for n = 0, 1, 4, 6, 8, 9, 18, 19, 21, 33, 40, 43. Part (i) of the conjecture is stronger than the classical result that any natural number is the sum of five pentagonal numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011177 | OEIS-A275877

Numbers on the East-West axis of A274640 in order of appearance.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011178 | OEIS-A275885

Lengths of runs of successive terms in A199134.

Weak conjecture: all terms are 1, 2 or 3. The first 3 appears at n = 1390. There is no 4 in the first 14000 terms. Does a 4 ever appear? The weak conjecture is true: all terms are 1, 2 or 3, so a 4 never appears (Ho, 2026). - _Boon Suan Ho_, Sep 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011179 | OEIS-A275888

First differences of A275884.

Weak conjecture: a(n) <= 4. Does a 5 ever appear? The weak conjecture is true: all terms are 1, 2, 3 or 4, so a 5 never appears. Moreover, the observations in the previous comment hold for all n: there are exactly 156 such words, of

lengths 2 to 26, five of them contain a 4, and 63 of them are faithful (Ho, 2026). - _Boon Suan Ho_, Sep 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011180 | OEIS-A275904

Order of homogeneous linear recurrence satisfied by the Pisot sequence T(n, n^2-n+1).

Conjecture: a(6) = 6852224. The conjectured generating function for T(6,31) is A(x)/(1+x - x*A(x)) where A(x) = 6 + x - x^2 - x^4 - x^22 - x^1130 - x^6852224 (and as usual there is a common factor of (1+x) in numerator and denominator). - _David Boyd_, Aug 12 2016.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011181 | OEIS-A275916

a(n) = A273059(4n).

A275916-A275919 are the four rays in A273059. There is a conjectured recursion for these sequences: see A140101.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011182 | OEIS-A275925

Trajectory of 3 under repeated application of the morphism sigma: 3 -> 3656, 5 -> 365656, 6 -> 3656656.

Conjecture 1: This sequence is a compressed version of A140101 (see that entry for details). [This was formerly stated as a theorem, but I am no longer sure I have a proof. - _N. J. A. Sloane_, Sep 29 2018. It is true: see the Dekking et al. paper. - _N. J. A. Sloane_, Jul 22 2019] Conjecture 2: For all k=0,1,2,..., the following two finite words are identical: The conjecture would imply that if we omit the initial 3 here, and change 6 to 1, 5 to 2, and leave 3 fixed, we get A092782. Alternatively, if we omit the initial 3 here, and change 6 to 0, 5 to 1, and 3 to 2, we get A080843. Proof of Conjecture 2. where we used the induction hypothesis in the second line. Removing the 656's at the end completes the induction step.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011183 | OEIS-A275974

Partial sums of the Jeffreys binary sequence A275973.

The ratio d(n) = a(n)/n, while obviously bounded, has no limit. Rather, it kind of 'oscillates', at an exponentially decreasing rate, between about 1/3 and 2/3. As mentioned by Jeffreys, the values of liminf and limsup of the set {d(n)} are 1/3 and 2/3, respectively. A proof of this fact by elementary means is relatively easy, for example, using the first formula below, but the following statement is a conjecture: Any real value c in the interval [1/3, 2/3] is an accumulation point of {d(n)}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Aug 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011184 | OEIS-A275975

Decimal expansion of Sum_{k>=0}((-1)^k/2^(2^k)).

Conjecture: Numbers of the type Sum_{k>=0}(x^(2^k)) with algebraic x and |x|<1 are known to be transcendental (Mahler 1930, Adamczewski 2013). It is likely that the alternating sign does not invalidate this property.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Aug 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011185 | OEIS-A276036

Numbers on the North-South axis of A274640 in order of appearance.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Aug 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011186 | OEIS-A276045

Primes p such that d(p*(2p+1)) = 8 where d(n) is the number of divisors of n (A000005).

Conjecture: this sequence is infinite. Conjecture: this sequence contains infinitely many twin primes. The first few twin primes in this sequence are 17,19,59,61,107,109,521,523,599,601,... (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anthony Hernandez_, Aug 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011187 | OEIS-A276051

a(n) is the number of top arches with length =1 for all semi meander solutions with n top arches.

Conjectured formula for n>=3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger Ford_, Aug 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011188 | OEIS-A276068

Sum of the lengths of the first descents in all bargraphs having semiperimeter n (n>=2). A descent is a maximal sequence of consecutive down steps.

Conjecture D-finite with recurrence (n+1)*a(n) +(-6*n+1)*a(n-1) +(10*n-13)*a(n-2) +(-4*n+13)*a(n-3) +(n-4)*a(n-4) +2*(-n+6)*a(n-5)=0. - _R. J. Mathar_, Jul 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Aug 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011189 | OEIS-A276069

Zero terms of A276034.

It is conjectured that this sequence is finite and all terms are found.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Nov 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011190 | OEIS-A276086

Primorial base exp-function: digits in primorial base representation of n become the exponents of successive prime factors whose product a(n) is.

The conjecture given in A327969 asks whether applying this function together with the arithmetic derivative (A003415) in some combination or another can eventually transform every positive integer into zero. Another related open question asks whether there are any other numbers than n=6 such that when starting from that n and by iterating with A003415, one eventually reaches a(n). See comments in A351088.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011191 | OEIS-A276091

Numbers obtained by reinterpreting base-2 representation of n in A001563-base (A276326): a(n) = Sum_{k>=0} A030308(n,k)*A001563(k+1).

Conjectured also to give all numbers n for which A255411(n) = A276340(n) (thus zeros of A276339).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011192 | OEIS-A276096

a(n) is the least number of empty convex pentagons ("convex 5-holes") spanned by a set of n points in the Euclidean plane in general position (i.e., no three points on a line).

Conjecture: a(n) = Theta(n^2). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Manfred Scheucher_, Aug 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011193 | OEIS-A276100

a(n) = (15*n)!*(n/2)!/((15*n/2)!*(5*n)!*(3*n)!).

This is only conjecturally an integer sequence. The similarly defined sequence (15*n)!*floor(n/2)!/(floor(15*n/2)!*(5*n)!*(3*n)!) = A211418(15*n) is integral.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Aug 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011194 | OEIS-A276101

a(n) = (10*n)!*(n/3)!/((5*n)!*(10*n/3)!*(2*n)!).

This is only conjecturally an integer sequence. The similarly defined sequence (10*n)!*floor(n/3)!/((5*n)!*floor(10*n/3)!*(2*n)!) = A211418(10*n) is integral.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Aug 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011195 | OEIS-A276102

a(n) = (6*n)!*(n/5)!/((3*n)!*(2*n)!*(6*n/5)!).

This is only conjecturally an integer sequence. The similarly defined sequence (6*n)!*floor(n/5)!/((3*n)!*(2*n)!*floor(6*n/5)!) = A211418(6*n) is integral.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Aug 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011196 | OEIS-A276136

Numbers m > 1 such that the largest odd divisors of m-1, m, and m+1 are prime.

Conjecture: this sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Aug 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011197 | OEIS-A276140

Replacing every "mixed pair" of integers with the absolute difference of the said pair of integers rebuilds the sequence itself (see "Comments" for the definition of a "mixed pair").

To elaborate reasons for the above conjecture about nonappearance, in the current b-file (8002 terms), only two terms, a(7) = 3 and a(11) = 5, have a value less than half their index. The plot2 link shows an apparently growing propensity for numbers not to appear later than the matching index. - _Peter Munn_, Aug 12 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Aug 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011198 | OEIS-A276141

a(1)=2; thereafter a(n) is the least prime > a(n-1) such that 2*a(n-1)+a(n) is a prime.

Conjecture: starting with any other prime, the sequence will eventually merge with this one. - _Zak Seidov_, Aug 06 2023.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Aug 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011199 | OEIS-A276175

a(n) = (a(n-1)+1)*(a(n-2)+1)*(a(n-3)+1)/a(n-4) with a(0) = a(1) = a(2) = a(3) = 1.

Conjecture: a(n) is an integer for all n >= 0. It has been checked by computer for n <= 40. A proof was proposed by 'mercio' as an answer to the MSE question, which however lacks details and heavily relies on computation. [Updated by _Max Alekseyev_, May 07 2023]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bruno Langlois_, Aug 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011200 | OEIS-A276188

Numbers k > 1 such that the number of odd divisors of k-1 is odd and is equal to the number of odd divisors of k+1.

Conjecture: this sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Aug 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011201 | OEIS-A276202

Smallest integer k such that the number of semiprimes in {k, f(k), f(f(k)), ..., 1} is equal to n, where f is the Collatz function.

Conjecture: a(n) exists for any n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Aug 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011202 | OEIS-A276290

Products of odd primes p and q such that either p or q is in the trajectory of (p*q)+1 under the Collatz 3x+1 map (A014682).

Conjecture: If n is the product of two odd primes p and q and p is equal to 3, then neither p nor q is in the trajectory of (p*q)+1 under the Collatz 3x+1 map (A014682). - _Marina Ibrishimova_, Aug 29 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marina Ibrishimova_, Aug 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011203 | OEIS-A276300

Number of 3 X n 0..2 arrays with no element equal to any value at offset (-2,-1) (-1,1) or (0,-1) and new values introduced in order 0..2.

Conjectures from _Colin Barker_, Feb 05 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Aug 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011204 | OEIS-A276305

Primes p such that d(p*(2p+1)) = 12 where d(n) is the number of divisors of n (A000005).

Conjecture: this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anthony Hernandez_, Aug 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011205 | OEIS-A276307

Primes p such that d(p*(2p+1)) = 16 where d(n) is the number of divisors of n (A000005).

Conjecture: this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anthony Hernandez_, Aug 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011206 | OEIS-A276309

Integer part of the ratio of alternate consecutive prime gaps.

Conjectures: The most frequent ratio among alternate prime gaps is 1, while the most frequent ratios among consecutive prime gaps seems to be 2 and 1/2, both with nearly the same frequency (see links). It also appears that next four most frequent ratios are 2, 1/2, 3 and 1/3, all four with nearly the same frequency (see links).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andres Cicuttin_, Aug 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011207 | OEIS-A276310

G.f. A(x) satisfies: x = A(x)-2*A(x)^2-2*A(x)^3.

Conjecture: 3*n*(n-1)*a(n) -13*(n-1)*(2*n-3)*a(n-1) -3*(3*n-5)*(3*n-7)*a(n-2)=0. - _R. J. Mathar_, Sep 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tom Richardson_, Aug 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011208 | OEIS-A276314

G.f. A(x) satisfies: x = A(x)-A(x)^2-3*A(x)^3.

Conjecture: +169*n*(n+2)*(n-1)*a(n) +13*(n-1) *(13*n^2+26*n-220) *a(n-1) +(-7277*n^3+13423*n^2+43814*n-81700) *a(n-2) -27*(3*n-10) *(3*n-8) *(71*n+197)*a(n-3)=0. - _R. J. Mathar_, Sep 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tom Richardson_, Aug 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011209 | OEIS-A276356

Number of Hamiltonian cycles in the Cartesian product graph K_2 times K_n.

For n > 1, a(n) = ((n-1)!/2)*(A001040(n-1) + A001053(n)). - Conjectured by _Mikhail Kurkov_, Feb 10 2019; proved (see MO link) by _Max Alekseyev_, Apr 23 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joel B. Lewis_, Aug 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011210 | OEIS-A276371

Expansion of e.g.f. exp(x/2)/(2 - exp(2*x))^(1/4).

Conjecture: the sequence a(n) taken modulo a positive integer k is eventually periodic with the period dividing phi(k). For example, the sequence taken modulo 21 begins [1, 1, 3, 19, 9, 19, 9, 1, 3, 19, 9, 19, 9, 1, 3, 19, 9, 19, 9, ...] with an apparent period of 6 (= phi(21)/2) starting at n = 1. - _Peter Bala_, Oct 23 2025 Conjectural o.g.f. as a continued fraction of Stieltjes type: 1/(1 - x/(1 - 2*x/(1 - 5*x/(1 - 4*x/(1 - 9*x/(1 - 6*x/(1 - 13*x/(1 - 8*x/(1 - ...)))))))). - _Peter Bala_, Oct 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011211 | OEIS-A276404

a(n) = A140100(n) - A140098(n).

Conjectured to be always 0 or 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011212 | OEIS-A276415

Number of ways to write n as p + x^3 + y^4, where p is a prime, and x and y are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 9, 10, 16, 17, 36, 52, 502.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011213 | OEIS-A276436

Generating function f(x)=(x+(x+(x+(x+(x+...)^5)^4)^3)^2)^1 is the limit as n->infinity of (f_1(x)=x, f_2(x)=x+x^2, f_3(x)=x+(x+x^3)^2, f_4(x)=x+(x+(x+x^4)^3)^2, ...).

Conjectures: the sequence contains no zeros except for a(0), a(3), a(5) and a(8); the sequence is asymptotically monotonic, the ratios of successive terms a(n+1)/a(n) converge to a constant c, 1<c<1.4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yuriy Sibirmovsky_, Sep 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011214 | OEIS-A276461

Prime numbers whose digits are k+1 1's and k 2's for some k >= 1.

The sequence is conjectured to be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bob Selcoe_, Sep 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011215 | OEIS-A276462

Prime numbers that consist of k 2's digits followed by k+1 1's digits for some k >= 1.

The sequence is conjectured to be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bob Selcoe_, Sep 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011216 | OEIS-A276463

Position at which n first appears in A276457, with a(0)=0.

The conjecture is that every number will appear in A276457, so this sequence should be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yuriy Sibirmovsky_, Sep 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011217 | OEIS-A276464

Fibbinary Fibonacci numbers.

Conjecture: 75025 is the last term of this sequence. - _Altug Alkan_, Jul 07 2020 This conjecture is similar to Erdős's conjecture that 256 is the largest power of 2 whose ternary representation does not contain the digit 2. - _Geoffrey Caveney_, Aug 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Sep 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011218 | OEIS-A276465

Divisors of 16450.

Conjecture: 16450 is the only number such that Sum_{d | n} 0.d is an integer, where 0.d means the decimal fraction of divisors d of n obtained by writing d after the decimal point:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Sep 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011219 | OEIS-A276469

Triangle read by rows: T(n,k) = n-th cyclotomic polynomial evaluated at x = k and then reduced mod n.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter A. Lawrence_, Sep 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011220 | OEIS-A276472

Modified Pascal's triangle read by rows: T(n,k) = T(n-1,k) + T(n-1,k-1), 1<k<n. T(n,1) = T(n-1,1) + T(n,2), n>2. T(n,n) = T(n,n-1) + T(n-1,n-1), n>1. T(1,1) = 1, T(2,1) = 1. n>=1.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yuriy Sibirmovsky_, Sep 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011221 | OEIS-A276493

Perfect numbers whose sum of prime factors is prime.

Conjectures (defining x = 170141183460469231731687303715884105727 = A007013(4)):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Sep 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011222 | OEIS-A276495

Odd numbers not of the form p + 2^m with p prime and m >= 0 for which the smallest k in A067760 such that n + 2^k is prime increases.

There exist de Polignac numbers n such that for all k >= 1 the numbers n + 2^k are composite. It is conjectured that 30666137 is the smallest such number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Sep 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011223 | OEIS-A276511

Primes that are equal to the sum of the prime factors of some perfect number.

Conjectures (defining x = 170141183460469231731687303715884105727 = A007013(4)):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Sep 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011224 | OEIS-A276513

a(n) = the smallest number k>1 such that Sum_{p|k} 0.p = n where p runs through the prime divisors of k.

Conjecture: a(4) = 730610790; Sum_{p|730610790} 0.p = 0.2 + 0.3 + 0.5 + 0.7 + 0.13 + 0.31 + 0.89 + 0.97 = 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Sep 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011225 | OEIS-A276517

Indices k such that A276516(k) = 0.

Conjecture: for k > 7169 there are no more terms in this sequence (tested for k < 10000000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Dec 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011226 | OEIS-A276522

Zero terms of A276520.

It is conjectured that this sequence is finite and all terms are found.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Nov 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011227 | OEIS-A276531

a(n) = (a(n-1) * a(n-5) + a(n-2) * a(n-3) * a(n-4)) / a(n-6), with a(0) = a(1) = a(2) = a(3) = a(4) = a(5) = 1.

Conjecture: a(n) is an integer for all n. It has been checked by computer for 0 <= n <= 50.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Nov 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011228 | OEIS-A276646

a(n) = floor(Sum_{k=1..n} 0.k).

Conjecture: function ((Sum_{k=1..n} 0.k) / n) is bounded above by values 0.55.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Sep 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011229 | OEIS-A276656

Nearest integer to the inverse function of the factorial where the exponential of n is the argument.

Conjecture: Limit_{n->oo} (Im(zetazero(n))/(2*Pi))/(the nearest integer to the inverse function of the factorial where the exponential of n is the argument) = 1. In other words, the non-rounded version of this sequence is asymptotic to imaginary part of the Riemann zeta zeros divided by 2*Pi.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Sep 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011230 | OEIS-A276674

Numbers n such that x - lambda(x) = n has no solution, where lambda(x) = A002322(x).

Conjecture: if x - lambda(x) = 2*m > 0, then x <= 4*m.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Oct 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011231 | OEIS-A276704

Records in A249859.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Barker_, Sep 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011232 | OEIS-A276705

Records in A249860.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Barker_, Sep 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011233 | OEIS-A276706

Indices of records in A249860.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Barker_, Sep 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011234 | OEIS-A276710

Composite numbers m such that Product_{k=0..m} binomial(m,k) is divisible by m^(m-1).

Conjecture, tested so far up to m = 3828: "When m belongs to this list, Product_{k=0..m} binomial(m,k) is divisible also by m^m". Since this is impossible for prime m (see, e.g., A109874), the conjecture is equivalent to the statement "m is prime if and only if Product_{k=0..m} binomial(m,k) is divisible by m^(m-1) but not by m^m". This conjecture has been proved, see corollary 3 in PDF link below.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Sep 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011235 | OEIS-A276711

Number of primes p <= n such that n - p is a perfect power (including 0 and 1).

Conjecture: For n > 1, we have a(n) > 0 except for n = 11^6 = 1771561. In other words, any integer n > 1 other than 11^6 can be written as p + x^k, where p is a prime, x is a nonnegative integer and k is an integer greater than one. Moreover, any integer n > 5 not among 8, 24, 1549, 1771561 can be written as p + x^k, where p is a prime, and x and k are integers greater than one. This has been verified for n up to 10^10. On the author's request Prof. Qing-Hu Hou at Tianjin Univ. verified it for n up to 6*10^9, and then the author used Hou's program to check the conjecture for n from 6*10^9 to 10^10. The author has also finished the verification of the conjecture for squares not exceeding 10^14 and sixth powers not exceeding 10^18, for example, 991^6 - 230^5 is a prime. - _Zhi-Wei Sun_, Sep 22 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011236 | OEIS-A276717

Least prime p < n^2 such that n^2 - p = x^k for some integers x > 1 and k > 1, or 1 if such a prime p does not exist.

The conjecture in A276711 implies that a(n) > 1 for all n > 2 except for n = 11^3 = 1331.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011237 | OEIS-A276751

G.f.: exp(Sum_{n>=1} [Sum_{k>=1} k^(2*n-1) * x^k]^n / n), a power series in x with integer coefficients.

Conjecture: a(n)^(1/n^2) tends to sqrt(2). - _Vaclav Kotesovec_, Oct 17 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011238 | OEIS-A276752

G.f.: exp(Sum_{n>=1} [Sum_{k>=1} k^(2*n) * x^k]^n / n), a power series in x with integer coefficients.

Conjecture: a(n)^(1/n^2) tends to sqrt(2). - _Vaclav Kotesovec_, Oct 17 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011239 | OEIS-A276770

a(n) is the number of runs of an algorithm. Set b_0 = n, if prime, stop; else, set c_0 = largest divisor of n (!=n); set b_1 = c_0 + b_0/c_0. Run with b_1.

For large n: Sum_{k=5..n} a(k) ~ n*log(n)/2 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yuriy Sibirmovsky_, Sep 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011240 | OEIS-A276771

a(n) is the number of runs of an algorithm. Set b_0 = n, if prime or 1 or 0, stop; else, set c_0 = largest divisor of n (!=n); set b_1 = c_0 - b_0/c_0. Run with b_1.

For large n: Sum_{k=1..n} a(k) ~ n*log(n)/2 - n/2 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yuriy Sibirmovsky_, Sep 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011241 | OEIS-A276788

First differences of A003144.

This relation between A003144 and A140101 is a conjecture - _Michel Dekking_, Mar 18 2019 [It has been a theorem since Mar 22 2019. - _N. J. A. Sloane_, Jun 25 2019. (See the Dekking et al. paper)]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011242 | OEIS-A276796

Partial sums of A276793.

Conjecture: A140100(n) - n = a(n-1). - _N. J. A. Sloane_, Oct 26 2016 (added Mar 21 2019). This is true: see the Dekking et al. paper. - _N. J. A. Sloane_, Jul 22 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011243 | OEIS-A276797

Partial sums of A276794.

Conjecture: A140102(n) - n = a(n-1). - _N. J. A. Sloane_, Oct 26 2016 (added Mar 21 2019). This is true - see the Dekking et al. paper. - _N. J. A. Sloane_, Jul 22 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011244 | OEIS-A276798

Partial sums of A276791.

Conjecture: 3*n - A140101(n) = a(n-1). - _N. J. A. Sloane_, Oct 26 2016 (added Mar 21 2019). This is true - see the Dekking et al. paper. - _N. J. A. Sloane_, Jul 22 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011245 | OEIS-A276830

Number of ways to write n as ((p-1)/2)^2 + P_2, where p is an odd prime and P_2 is a product of at most two primes.

Conjecture: a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 3, 4, 9, 17, 21, 25, 33, 49, 109, 169, 189, 361, 841, 961, 12769, 19321. See also A276825 for a similar conjecture involving cubes, and some comments on x^2 + P_2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011246 | OEIS-A276835

Numerator of a modified exponentiated von Mangoldt function defined recursively.

Conjecture: For n>3: If and only if the ratio A276835(n)/A276836(n) is equal to n then n is equal to the greater of the twin primes A006512.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Sep 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011247 | OEIS-A276836

Denominator of modified von Mangoldt function defined recursively.

See A276835 for the conjecture about the ratio A276835(n)/A276836(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Sep 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011248 | OEIS-A276862

First differences of the Beatty sequence A003151 for 1 + sqrt(2).

Conjectures: Equals both A245219 and A097509. - _Michel Dekking_, Feb 18 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011249 | OEIS-A276885

Sums-complement of the Beatty sequence for 1 + phi.

Mathar's conjecture is proved in the paper 'The Frobenius problem for homomorphic embeddings of languages into the integers'. See Example 1 in that paper. - _Michel Dekking_, Dec 21 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011250 | OEIS-A276976

Smallest m such that b^m == b^n (mod n) for every integer b.

Conjecture: a(n) <= n/3 for every n >= 9. Professor Andrzej Schinzel proved this conjecture (in a letter to the author). - _Thomas Ordowski_, Sep 30 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Sep 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011251 | OEIS-A276982

a(n) = number of primes p whose balanced ternary representation is compatible with the binary representation of A276194(n).

It is conjectured that all a(n) > 0. This has been checked for n <= 100000. But it is possible that there is a counterexample for very large n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Oct 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011252 | OEIS-A277028

Numbers that are never pandigital for any base b > 1.

This is a subsequence of A000225, conjectured to be finite (no other term below 2^1000000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Sep 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011253 | OEIS-A277060

a(n) = (1/2) * Sum_{k=0..n} (binomial(n,k) * binomial(n+k,k+1))^2 for n >= 0.

Conjecture: the supercongruences a(p-1) == 1 (mod p^4) holds for all primes p >= 5 and a(p^2-1) == 1 (mod p^5) holds for all primes p >= 3. - _Peter Bala_, Mar 22 2023 The conjecture is true (see Epstein link). - _Andrew R. Epstein_, Sep 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Nov 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011254 | OEIS-A277088

Pisot sequences L(5,12), S(5,12).

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Sep 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011255 | OEIS-A277089

Pisot sequences L(6,15), S(6,15).

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Sep 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011256 | OEIS-A277109

Starting from 2^n+1, the length of the longest sequence of consecutive numbers which all take the same number of steps to reach 1 in the Collatz (or '3x+1') problem.

Conjecture: If a(n) = 2^k - 1 for some k > 1, then a(n-1) = 2^(k-1) - 1. Conjecture holds for n <= 1812. The conjecture is true. Let the lengths of the Collatz runs equal q for all numbers 2^n + 1, 2^n + 2, 2^n + 3, 2^n + 4, ..., 2^n + 2^k - 2, 2^n + 2^k - 1. Then dividing the 2^(k-1) - 1 even numbers by two gives rise to the sequence 2^(n-1) + 1, 2^(n-1) + 2, ..., 2^(n-1) + 2^(k-1) - 1 of numbers for which the lengths of the Collatz runs equals q-1. Furthermore, let the length of the Collatz run of 2^n + 2^k be r != q then the length of the Collatz run of 2^(n-1) + 2^(k-1) is r-1 != q-1, i.e., a(n-1) = 2^(k-1) - 1. Conjecture: Let a(k), ..., a(k+m), m >= 0, be a subsequence of this sequence such that a(k)=a(k+m+1)=1 and a(k+i) > 1, 1 <= i <= m. Then the lengths of the Collatz runs of a(k+i), 0 <= i <= m, increase by 1. In addition, there is an initial segment of increasing numbers a(k), ..., a(k+j), for some 0 <= j <= m, in each such subsequence having the form 2^i - 1, 0 < i <= j. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Sep 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011257 | OEIS-A277120

Number of branching factorizations of n.

The value of a(n) depends only on the prime signature of n. In other words, for all i, j >= 1, it holds that A101296(i) = A101296(j) => a(i) = a(j). Moreover, it seems that the converse proposition also holds, that for all i, j >= 1, a(i) = a(j) => A101296(i) = A101296(j), i.e., for each distinct prime signature there exists a distinct value of a(n). This has been empirically checked up to the first 21001 prime signatures in A025487 (see A366884), and can be proved if one can show that the latter sequence (equally: A366377) is injective. If this conjecture holds, it would imply an unlimited number of statements like those given in the previous comment (see the formula section of A101296).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Oct 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011258 | OEIS-A277141

Lexicographically least sequence of nonnegative integers that avoids a/b-powers for all a/b >= 4/3.

Conjecture: a(336*n + 1666) = A277142(56*n + 17) + 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Rowland_, Oct 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011259 | OEIS-A277147

Lexicographically least sequence of nonnegative integers that avoids 7/5-powers.

This sequence is conjecturally 80874-regular. Conjecture: a(80874 n + 173978) = a(n) + 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Rowland_, Oct 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011260 | OEIS-A277150

Lexicographically least sequence of nonnegative integers that avoids 7/6-powers.

This sequence is conjecturally 41190-regular. Conjecture: a(41190 n + 41201) = a(n) + 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Rowland_, Oct 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011261 | OEIS-A277152

Lexicographically least sequence of nonnegative integers that avoids 8/7-powers.

This sequence is conjecturally 340-regular. Conjecture: a(340 n + 52670) = a(n) + 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Rowland_, Oct 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011262 | OEIS-A277153

Lexicographically least sequence of nonnegative integers that avoids 9/7-powers.

This sequence is conjecturally 44-regular. Conjecture: a(44 n + 2701) = a(n) + 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Rowland_, Oct 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011263 | OEIS-A277154

Lexicographically least sequence of nonnegative integers that avoids 10/7-powers.

This sequence is conjecturally 26-regular. Conjecture: a(26 n + 428) = a(n) + 1 for all n >= 1 (but not for n = 0).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Rowland_, Oct 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011264 | OEIS-A277161

Lexicographically least sequence of nonnegative integers that avoids 15/8-powers.

This sequence is conjecturally 22763-regular. Conjecture: a(22763 n + 22850) = a(n) + 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Rowland_, Oct 01 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011265 | OEIS-A277190

Numbers n such that A277013(n) differs from A001222(n).

If the conjecture by Ulas and Ulas is true, then there are no primes in this sequence. See A186891.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011266 | OEIS-A277201

Primes of the form (p^4 + 1)/2, where p is prime.

Conjecture: the sequence consists of the numbers k such that tau(2k) = 4 and tau(2k-1) = 5. tau(82) = 4 and tau(81) = 5, 82/2 = 41 = a(1). tau(626) = 4 and tau(625) = 5, 626/2 = 313 = a(2). tau(2402) = 4 and tau(2401) = 5, 2402/2 = 1201 = a(3). The conjecture was checked for 10^9 consecutive integers. The above conjecture is true: since tau(2k-1) = 5, 2k-1 must be the 4th power of some prime p, i.e., k = (p^4 + 1)/2 (so p is odd, so p^4 == 1 (mod 16), so k is odd), and since tau(2k) = 4, 2k must be the product of two distinct primes, so k is an odd prime. Thus, the set of numbers k such that tau(2k) = 4 and tau(2k-1) = 5 is the set of primes of the form (p^4 + 1)/2, where p is prime. - _Jon E. Schoenfield_, Mar 17 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lechoslaw Ratajczak_, Oct 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011267 | OEIS-A277215

a(n) is the smallest even number not congruent to 1 modulo 3 that starts a (2n+1)-element alternating sequence of x/2 and (3x+1) iterations ending in the maximum of its Collatz trajectory.

Conjecture: For every n there is an odd number q such that the alternating sequence ends in v_(2n), the maximum of the Collatz trajectory of a(n)=v_0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Nov 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011268 | OEIS-A277223

a(n) = A052489(n)/n.

a(n) is never 1, 2, 3, 4, 5 or 6. Conjecture: if a(n) < 12 then a(n) = 0 or 9. - _Robert Israel_, Oct 06 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Oct 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011269 | OEIS-A277225

Initial values of runs of 5 consecutive numbers all of which are squares, primes, or products of one prime and one square.

The existence of runs of 6 or 7 numbers with the same property is an open question; a run of length 8 is impossible. - _Andrey Zabolotskiy_, Nov 07 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles Bowyer_, Oct 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011270 | OEIS-A277272

a(n+1) is the smallest number not already in the sequence whose sum of prime factors (with repetition) shares a factor with the sum of the prime factors of a(n); a(1)=2.

Inspired by A064413 (the EKG sequence). Conjecture: The sequence is a permutation of the natural numbers >1, in which composite terms appear once only (in irregular fashion) and the primes arise in their natural order. Immediately following the appearance of any prime p>3 the next term is q*(2^r) where q is the largest prime less than p, and r is the unique integer such that p=q+(2*r). The above conjecture fails at a(95)=23, which appears before a(162)=19. (Also, the term appearing immediately after 37 is 218=2*109; the term immediately after 97 is 1335=3*5*89.) - _Jon E. Schoenfield_, Nov 05 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Oct 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011271 | OEIS-A277312

Smallest k such that k - lambda(k) = prime(n), where lambda(k) = A002322(k).

Conjecture: a(n) = prime(n)^2 for infinitely many n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Oct 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011272 | OEIS-A277316

Prime-factorization representation of the prime-th Stern-polynomial: a(n) = A260443(A000040(n)).

If the conjecture by Ulas and Ulas is true, then all these terms can be found from A206284 and then this is also a subsequence of A277318.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 10 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011273 | OEIS-A277338

Reverse and Add! sequence starting with 295.

295 is conjectured to be the second smallest initial term which does not lead to a palindrome. Also, 196 is possibly the smallest initial term which does not lead to a palindrome. a(0) = 196 is described in A006960.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matt C. Anderson_, Oct 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011274 | OEIS-A277346

Maximal coefficient among squares of the polynomials in row n of the triangle of q-binomial coefficients.

Conjecture: a(n) ~ sqrt(3) * 2^(2*n+2) / (Pi^(3/2) * n^(5/2)). - _Vaclav Kotesovec_, Jan 07 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Reshetnikov_, Oct 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011275 | OEIS-A277403

E.g.f. satisfies: A(x - Integral A(x) dx) = x + Integral A(x) dx.

a(n) is divisible by 10 for n>2 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011276 | OEIS-A277411

Column 1 of triangle A277410.

Conjectures from _Colin Barker_, Nov 04 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011277 | OEIS-A277439

Numerators of a sequence defined by a modified recurrence for the exponential of the von Mangoldt function.

Conjecture for n > 2: The ratio is equal to n if and only if n is the average of twin prime pairs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Oct 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011278 | OEIS-A277440

Denominators of a sequence defined by a modified recurrence for the exponential of the von Mangoldt function.

See A277439 for the conjecture about the average of twin primes A014574 and the ratio A277439(n)/A277440(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Oct 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011279 | OEIS-A277506

Expansion of e.g.f. 1/((1+LambertW(-x))*(1-x)).

Conjecture: a(n) is the number of maximal chains in the Hochschild lattice of dimension n+1. For example, there are a(1)=2 maximal chains in the pentagon poset, which has dimension 2. - _F. Chapoton_, Aug 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Oct 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011280 | OEIS-A277514

Irregular triangle read by rows: T(n,k) is the number of primes with n balanced ternary digits of which 2k+1 (3 <= 2k+1 <= n) are nonzero.

It is conjectured that all terms of this sequence are greater than zero, or else there are infinitely many zero terms. The first zero term might appear beyond 10^6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Oct 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011281 | OEIS-A277548

Numbers k such that k/5^m == 4 (mod 5), where 5^m is the greatest power of 5 that divides k.

Conjecture: a(n) = 4*n if and only if n is in A033042. - _David A. Corneth_, Oct 23 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011282 | OEIS-A277560

Binary representation of the x-axis, from the left edge to the origin, or from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 7", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 06 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011283 | OEIS-A277581

Goldbach's problem extended to squares of nonnegative differences of primes: smallest integer >= ((A112823(n) - A234345(n))^2)/n for n >= 2.

Conjecture: 1 <= a(n) <= m for all n, where m is largest value of a(n), i.e., the sequence of records in a(n) {1, 5, 15, 20, ..., m} is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Oct 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011284 | OEIS-A277602

List of indices n such that A092488(n) < A068527(n).

Numbers in this sequence are numbers of the form k^2 + m where m is fairly small compared to k^2 and less than 2k + 1. For m < 2k + 1, if k^2 + m isn't in the sequence then k^2 + m + 1 isn't either. Conjecture: 1 <= m <= t for some t < 2k+1. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Oct 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011285 | OEIS-A277604

Array of coefficients T(k,n) of the formal power series A(k,x) read by upwards antidiagonals, where A(k,x) = sqrt(1 + 2*x*A(k,x) + (4*k+1)*x^2*(A(k,x))^2), k >= 0.

Conjecture: If B(k,n) satisfy B(k,0) = B(k,1) = 1 and B(k,n+2) = B(k,n+1) + k*B(k,n) for k >= 0 and n >= 0 (generalized Fibonacci numbers, see A015441) and G(k,x) = Sum_{n>=0} A000108(n)*B(k,n)*x^n for k >= 0, then you will have (1): A(k,x*G(k,x)) = G(k,x) and (2): G(k,x/A(k,x)) = A(k,x) for k >= 0. Especially for k = 1 see A098615 and for k = 2 see A200376. Conjecture: T(k,2*n) = Sum_{i=0..n} A046521(n,i)*k^(n-i) for k, n >= 0. - _Werner Schulte_, Aug 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Oct 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011286 | OEIS-A277623

Each odd term "t" of the sequence is followed by t even terms and each odd digit "k" of the sequence is followed by k even digits.

The sequence could be started with the even term a(0) = 0, which would not impose any restriction and thus be followed by the next smallest possible integer, a(1) = 1. The sequence is not a permutation of the positive integers because some numbers (11, 13, 15, 17, 19, 31, ..., 97, 99, 100, 102,..., 110, 111, ..., 119, 130, ...) = A277694 can never occur. (The odd digit 1 must be followed by 1 even digit, etc.) The previous comment shows why odd terms occur late, but probably (I conjecture) any odd term which is not excluded a priori will occur sooner or later. - _M. F. Hasler_, Oct 27 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Lars Blomberg_, Oct 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011287 | OEIS-A277624

Composite numbers which have a dominant prime factor. A prime factor p of n is dominant if floor(sqrt(p)) > (n/p).

Conjecture: A092487(a(n)) = A006530(a(n)). - _David A. Corneth_, Oct 27 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Oct 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011288 | OEIS-A277684

Least k such that A277109(k) = 2^n - 1.

Are the terms always increasing? Note, if the conjecture in A277109 is true then the terms in this sequence are guaranteed to be increasing. Since the conjecture in A277109 is true, this sequence is strictly increasing. - _Hartmut F. W. Hoft_, Aug 16 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Oct 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011289 | OEIS-A277688

Odd numbers k such that there is no prime p < k/2 with k - 2*p and k + 2*p both prime.

Or, odd integers k such that k + 2*p is composite for all primes p, q with 2*p + q = k. By the Lemoine-Levy conjecture, for every odd k>5, there are primes p and q such that k=2*p+q. Numbers 1,3,5 formally satisfy the condition. Conjecture: k=59 and k=151 are the only terms k>5 satisfying the additional condition that k + 2*q is composite for every prime p,q such that 2*p+q=k. This conjecture arose from the calculations up to 500001 by _Peter J. C. Moses_ and confirmed by _M. F. Hasler_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011290 | OEIS-A277691

Smallest k such that (k+i)*prime(n)# - 1 is prime for i = 0, 1, 2, 3, 4 with prime(n)# = A002110(n) the n-th primorial, n>1.

Conjecture: for all n>1, there exists an integer k to get 5 primes in arithmetic progression starting with k*prime(n)# - 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Oct 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011291 | OEIS-A277735

Unique fixed point of the morphism 0 -> 01, 1 -> 20, 2 -> 0.

Although I believe the assertions in Kimberling's comment above to be correct, these results are quite tricky to prove, and unless a formal proof is supplied at present these assertions must be regarded as conjectures. - _N. J. A. Sloane_, Aug 20 2018 Here is a proof of Clark Kimberling's conjectures (and more).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011292 | OEIS-A277761

Number of n X 2 0..2 arrays with every element equal to some element at offset (-1,-1) (-1,0) (-1,1) (0,-1) (0,1) or (1,0) both plus 1 mod 3 and minus 1 mod 3, with new values introduced in order 0..2.

Conjectures from _Colin Barker_, Feb 05 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Oct 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011293 | OEIS-A277797

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 1", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Oct 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011294 | OEIS-A277798

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 1", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Oct 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011295 | OEIS-A277799

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 1", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Oct 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011296 | OEIS-A277800

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 1", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 01 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Oct 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011297 | OEIS-A277829

First Series of Hankel determinants based on squares of Catalan numbers.

Conjecture: lim n->infinity log(a(n))/n^2 = 2*log(2). - _Vaclav Kotesovec_, Nov 29 2016 [Proved in the paper by Wenlin Zhang and Haobo Ma, Oct 01 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Karol A. Penson_, Nov 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011298 | OEIS-A277848

Index at which n occurs in A217727, or 0 if there is no such index.

It is conjectured that all integers >= 2 appear in A217727.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Nov 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011299 | OEIS-A277860

a(n) = Sum_{k=0..n-1} binomial(4k, 2k+1)*binomial(2k, k)*48^(n-1-k).

Conjecture: For any prime p > 3 and positive integer n, we have This conjecture with n = 1 gives the congruence a(p) == (5/12)*p^2*B_{p-2}(1/3) (mod p^3) for any prime p > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011300 | OEIS-A277864

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 3", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 03 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011301 | OEIS-A277865

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 3", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 03 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011302 | OEIS-A277866

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 3", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 03 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011303 | OEIS-A277867

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 3", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 03 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011304 | OEIS-A277878

Smallest prime q with q > r such that p = (q + r + 1)/r, where p = prime(n) and r = A277879(n).

Rassias's conjecture is the claim that q and r exist for every p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Nov 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011305 | OEIS-A277879

Smallest prime r such that p = (q + r + 1)/r, where p = prime(n) and q = A277878(n).

Rassias's conjecture is the claim that q and r exist for every p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Nov 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011306 | OEIS-A277913

Nonsquare numbers n for which the smallest y>0 solution of the Pellian equation x^2 - n*y^2 = 1 divides n.

Lim_{n->infinity} a(n+1)/a(n) = 1 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Salvador Cerdá_, Nov 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011307 | OEIS-A277926

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 5", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 04 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011308 | OEIS-A277927

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 5", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 04 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011309 | OEIS-A277928

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 5", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 05 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011310 | OEIS-A277929

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 5", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 05 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011311 | OEIS-A277930

Array of coefficients a(k,n) of the formal power series A(k,x) read by upwards antidiagonals, where A(k,x) = ((2*k+1)*x+sqrt(1+4*k*(k+1)*x^2))/(1-x^2), k>=0.

Conjecture: a(k,2*n+2) = 1+2*k+2*(-k)^(n+2)*(Sum_{i=0..n} A234950(n,i)*k^i) for k>=0 and n>=0. - _Werner Schulte_, Aug 03 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Nov 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011312 | OEIS-A277936

Decimal representation of the x-axis, from the left edge to the origin, or from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 7", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 06 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011313 | OEIS-A277952

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 14", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 06 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011314 | OEIS-A277953

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 14", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 06 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011315 | OEIS-A277954

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 14", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 06 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011316 | OEIS-A277997

Lexicographically first sequence of nonprimes (with no duplicates) whose absolute first differences are prime (with no duplicates).

The sequence starts with a(1) and was always extended with the smallest integer not yet present that does not lead to a contradiction. It is conjectured that the sequence together with its first absolute differences is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Nov 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011317 | OEIS-A278008

Number of n X 2 0..2 arrays with every element plus 1 mod 3 equal to some element at offset (-1,-1) (-1,0) (-1,1) (0,-1) (0,1) or (1,0), with upper left element zero.

Conjectures from _Colin Barker_, Feb 07 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011318 | OEIS-A278023

G.f.: 2*x*(1-x*sqrt(1-4*x))/((1+2*x^2+sqrt(1-4*x))*sqrt(1-4*x)).

Conjecture: +n*(3*n^2-12*n+11) *a(n) -(3*n-5) *(3*n^2-9*n+4) *a(n-1) -2*(2*n-5) *(3*n^2-6*n+2) *a(n-2) +n *(3*n^2-12*n+11) *a(n-3) -2 *(2*n-5) *(3*n^2-6*n+2) *a(n-4)=0. - _R. J. Mathar_, Jun 24 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 09 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011319 | OEIS-A278081

a(n) is 1/12 of the number of primitive quadruples with sum = 0 and sum of squares = 2*m^2, where m = 2*n - 1.

Conjecture: b(m) is multiplicative: for k >= 1, b(2^k) = 0; for p an odd prime, b(p*k) = p^(k-1)*b(p); b(p)= p + 1 for p == (5, 7, 13, 23) (mod 24); b(p) = p-1 for p == (1, 11, 17, 19) (mod 24); and b(3) = 3. It would be nice to have a proof of this.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Mallows_, Nov 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011320 | OEIS-A278082

1/12 of the number of primitive quadruples with sum = n and sum of squares = 3*n^2.

Conjecture: a(n) is multiplicative, with a(2) = 1, a(2^k) = 0 (k >= 2); a(p^k) = p^(k-1)*a(p); a(p) = p + 1 for p == (2, 6, 7, 8, 10)(mod 11), a(p) = p - 1 for p == (1, 3, 4, 5, 9)(mod 11); and p(11) = 11. It would be nice to have a proof of this.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Mallows_, Nov 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011321 | OEIS-A278083

a(n) is 1/6 of the number of primitive integral quadruples with sum = 2*m and sum of squares = 2*m^2, where m = 2*n-1.

Conjecture: b(m) is multiplicative: for k >= 1, b(2^k) = 0; b(p^k) = p^(k-1)*b(p) for p an odd prime; b(p) = p+1 for p == 3 (mod 4); b(p) = p-1 for p == 1 (mod 4). It would be nice to have a proof of this. Conjecture: a(n) = (A000010(2*n - 1)/6) * lim_{k->oo} A005875((2*n - 1)^(2*k))/(2*n - 1)^k. - _Velin Yanev_, Oct 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Mallows_, Nov 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011322 | OEIS-A278084

a(n) is 1/24 of the number of primitive integral quadruples with sum = 2*m and sum of squares = 6*m^2, where m = 2*n-1.

Conjecture: b(m) is multiplicative: for k >= 1, b(2^k) = 0, and for p an odd prime, b(p^k) = p^(k-1)*b(p), with b(p) = p + 1 for p == (11, 13, 17, 19) (mod 20), b(p) = p - 1 for p == (1, 3, 7, 9) (mod 20), b(5) = 5. It would be nice to have a proof of this.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Mallows_, Nov 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011323 | OEIS-A278085

1/4 of the number of primitive integral quadruples with sum = 3*n and sum of squares = 3*n^2.

Conjecture: a(n) is multiplicative, with a(2) = 1, a(2^k) = 0 for k>=2, and for k >= 1 and p an odd prime, a(p^k) = p^(k-1)*a(p), with a(p) = p+1 for p == 5 (mod 6), a(p) = p-1 for p=1 (mod 6), and a(3) = 3. It would be nice to have a proof of this. [See A354766 for additional conjectures. - _N. J. A. Sloane_, Jun 19 2022]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Mallows_, Nov 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011324 | OEIS-A278086

1/12 of the number of integer quadruples with sum = 3*n and sum of squares = 7*n^2.

Conjecture: a(n) is multiplicative with a(2) = 1, a(2^k) = 0 for k >= 2, and for k >= 1 and p an odd prime, a(p^k) = p^(k-1)*a(p) with a(p) = p + 1 for p == (2, 3, 8, 10, 12, 13, 14, 15, 18) (mod 19), a(p) = p - 1 for p == (1, 4, 5, 6, 7, 9, 11, 16, 17) (mod 19), and p(19) = 19. It would be nice to have a proof of this.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Mallows_, Nov 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011325 | OEIS-A278209

a(n) is the smallest number k such that {p(1), p(2), ..., p(k)} contains each residue class mod n, where p(m) is the m-th partition number A000041(m).

It has been conjectured that for each r and n, there is some k such that p(k) = r mod n; on that conjecture, this sequence is defined for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Nov 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011326 | OEIS-A278212

Maximum total number of possible moves that any number of bishops of the same color can make on an n X n chessboard.

Conjecture: a(n) = floor(A275815(n)/2). - _Peter Kagey_, Nov 24 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Nov 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011327 | OEIS-A278319

Number of n X 2 0..1 arrays with rows and columns in lexicographic nondecreasing order but with exactly two mistakes.

Conjectures from _Colin Barker_, Feb 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011328 | OEIS-A278341

a(n) is the number of decompositions of n into unordered form p + c*q, where p, q are terms of A274987 and the difference of trits for p and q is no more than 1, c=1 for even n-s and c=2 for odd n-s.

It is conjectured that this sequence has only 208 zero terms listed in sequence A278342.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Nov 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011329 | OEIS-A278342

Zero terms of A278341.

It is conjectured that this sequence is finite and all 208 terms are found.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Nov 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011330 | OEIS-A278357

Number of n X 2 0..1 arrays with rows and columns in lexicographic nondecreasing order but with exactly one mistake.

Conjectures from _Colin Barker_, Feb 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 19 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011331 | OEIS-A278415

a(n) = Sum_{k=0..n} binomial(n, 2k)*binomial(n-k, k)*(-1)^k.

Conjecture: For any prime p > 3 and positive integer n, the number (a(p*n)-a(n))/(p*n)^2 is always a p-adic integer. See also A275027 and A278405 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011332 | OEIS-A278425

Largest k such that there are no primes between kn and k(n+1); -1 if no such k exists.

This sequence deals with the question of whether there is always a prime between nk and n(k+1). For n<=3 the answer has been proven to be yes (see links and examples). For n>3 the problem remains open, however we can conjecture the values of a(n) by checking the first few hundred k. Conjecture: For every n, there exists a finite m such that for every k>m there is at least one prime between kn and k(n+1). In other words, a(n) is never -1. Conjecture follows from the Prime Number Theorem: for fixed n, the number of primes between kn and k(n+1) is asymptotic to k/log(k) as k -> infinity, and in particular is nonzero for all sufficiently large k. - _Robert Israel_, Nov 28 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Nov 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011333 | OEIS-A278472

a(n) = Sum_{i=0..n} Fibonacci(i+1)*binomial(2*n-i+2, n+2).

Conjecture: +(n+2)*(n^2-4*n+1)*a(n) +2*(-4*n^3+9*n^2+16*n-6)*a(n-1) +(15*n^3-38*n^2-9*n+14)*a(n-2) +2*(2*n-1)*(n^2-2*n-2)*a(n-3)=0. - _R. J. Mathar_, Mar 06 2017 Conjecture: +(n+2)*a(n) +(-11*n-14)*a(n-1) +(37*n+20)*a(n-2) +(-25*n-6)*a(n-3) +2*(-21*n+41)*a(n-4) +4*(-2*n+5)*a(n-5)=0. - _R. J. Mathar_, Mar 06 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Nov 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011334 | OEIS-A278484

Main diagonal of A278482.

a(n) = 2*A073359(n). [Observed by _Arie Groeneveld_ on SeqFan list, Nov 22 2016.] [Conjectural, true for n <= 2000 _David W. Wilson_, Nov 23 2016]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 23 2016 after _David W. Wilson_'s posting on SeqFan list Nov 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011335 | OEIS-A278528

a(n) = number of the round in which n is removed in the Flavius sieve, 0 if it is never removed (when n is one of the terms of A000960).

Conjecture: a(n) = [C > 0] * C where we start with A := n, B := n - 1, C, m := 0 and until A = B consecutively apply m := m + 1, C := A - B, A := abs(A - m - (A mod m)), B := abs(B - m - (B mod m)). - _Mikhail Kurkov_, May 19 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011336 | OEIS-A278546

Even numbers that cannot be expressed as a sum of 3 or fewer terms of A035928.

This is conjectured to be the complete list. There are no other examples smaller than 16773120.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Nov 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011337 | OEIS-A278554

Number of distinct blocks of length n (a.k.a. subword complexity) of the characteristic sequence of the squarefree numbers A008966.

Conjecture 1: this is the number of binary sequences S of length n such that, for all primes p such that p^2 <= n, at least one of the p^2 linearly indexed subsequences of S with gap p^2 starting at the 1st, 2nd, ..., p^2-th position of S, is the all-zeros sequence. In other words, every block that is not explicitly ruled out by congruence conditions for the primes p with p^2 <= n should occur. Conjecture 2: the last new block to actually occur is always 0^n (n copies of 0). Cf. A020754.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Jan 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011338 | OEIS-A278560

Numbers of the form x^2 + y^2 + z^2 with x + 3*y + 5*z a square, where x, y and z are nonnegative integers.

This is motivated by the author's 1-3-5-Conjecture which states that any nonnegative integer can be expressed as the sum of a square and a term of the current sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011339 | OEIS-A278613

Sum of terms in level n of TRIP - Stern sequence associated with permutation triple (e,e,132).

Conjecture: G.f.: (-3-5*x-7*x^2) / (-1+x+2*x^2+6*x^3). - _R. J. Mathar_, Dec 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Amburg_, Nov 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011340 | OEIS-A278614

Sum of terms in level n of TRIP - Stern sequence associated with permutation triple (e,12,12).

Conjecture: G.f.: (3-x-5*x^2) / (1-3*x-x^2+4*x^3). - _R. J. Mathar_, Dec 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Amburg_, Nov 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011341 | OEIS-A278616

Sum of terms in level n of TRIP - Stern sequence associated with permutation triple (e,13,132).

Conjecture: G.f.: (-3-5*x-x^2) / (-1+x+4*x^2+x^3). - _R. J. Mathar_, Dec 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Amburg_, Nov 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011342 | OEIS-A278618

a(n) = Sum_{j=0..n/2} binomial(n-j-1,n-2*j)*binomial(2*n+1,j).

Conjecture: 2*n*(5*n-8)*(2*n-1)*(n+1)*a(n) -n*(115*n^3-344*n^2+299*n-82)*a(n-1) -4*(2*n-1)*(5*n^3+27*n^2-74*n+30)*a(n-2) +36*(n-2)*(5*n-3)*(2*n-1)*(2*n-3)*a(n-3)=0. - _R. J. Mathar_, Dec 02 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Nov 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011343 | OEIS-A278664

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 62", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 26 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011344 | OEIS-A278665

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 62", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 26 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011345 | OEIS-A278666

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 62", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 26 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011346 | OEIS-A278667

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 62", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 26 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011347 | OEIS-A278670

Number of n X 2 0..1 arrays with rows in nondecreasing lexicographic order and columns in nonincreasing lexicographic order, but with exactly one mistake.

Conjectures from _Colin Barker_, Feb 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011348 | OEIS-A278681

Pisot sequence T(3,16).

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Nov 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011349 | OEIS-A278692

Pisot sequence T(4,14).

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Nov 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011350 | OEIS-A278720

The p-defect p - N(p) of the congruence y^2 == x^3 + 4*x (mod p) for primes p, where N(p) is the number of solutions given by A276730.

a(n) = 0 if prime(n) == 2 or 3 (mod 4). (see the conjecture by _Robert Israel_, Sep 28 2016 in A276730). a(n) = 0 if prime(n) == 2 or 3 (mod 4) (this is conjecture II from above).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Dec 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011351 | OEIS-A278741

Odd numbers k such that tau(k-1) is a prime.

Conjecture: prime terms are in A249759: 3, 5, 17, 65537, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011352 | OEIS-A278743

For a base n>1: consider the lexicographically least strictly increasing sequence c_n such that, for any m>0, Sum_{k=1..m} c_n(k) can be computed without carries in base n; the sequence c_n is (conjecturally) eventually linear, and a(n) gives its order.

For a base n>1: consider the lexicographically least strictly increasing sequence c_n such that, for any m>0, Sum_{k=1..m} c_n(k) can be computed without carries in base n; the sequence c_n is (conjecturally) eventually linear, and a(n) gives its order. More precisely, we conjecture that, for any n>1, there are two constants k0 and b such that c_n(k + a(n)) = c_n(k)*n^b for any k>k0. [Corrected by _Rémy Sigrist_, Dec 24 2016]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Nov 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011353 | OEIS-A278753

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 65", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Jun 15 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011354 | OEIS-A278754

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 65", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Jun 15 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011355 | OEIS-A278755

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 65", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Jun 15 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011356 | OEIS-A278756

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 65", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Jun 15 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011357 | OEIS-A278757

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 73", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Jun 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011358 | OEIS-A278758

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 73", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Jun 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011359 | OEIS-A278759

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 73", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Jun 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011360 | OEIS-A278760

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 73", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Jun 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011361 | OEIS-A278764

Pisot sequence T(5,13).

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Nov 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011362 | OEIS-A278770

Second series of Hankel determinants based on squares of Catalan numbers.

Conjecture: lim n->infinity log(a(n))/n^2 = 2*log(2). - _Vaclav Kotesovec_, Nov 28 2016 [Proved in the paper by Wenlin Zhang and Haobo Ma, Oct 01 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Karol A. Penson_, Nov 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011363 | OEIS-A278772

Number of n X 2 0..1 arrays with rows in nondecreasing lexicographic order and columns in nonincreasing lexicographic order, but with exactly two mistakes.

Conjectures from _Colin Barker_, Feb 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Nov 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011364 | OEIS-A278790

Number of real cubic fields with discriminant <= 10^n.

The number of real cubic fields with discriminant <= X is asymptotic to X/(12*zeta(3)) = (0.069325...)*X. The second order term was conjectured by Roberts to be a known constant times X^{5/6}, and this was subsequently proved by Bhargava et al.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christopher E. Thompson_, Nov 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011365 | OEIS-A278791

Number of complex cubic fields with discriminant >= -10^n.

The number of complex cubic fields with discriminant >= -X is asymptotic to X/(4*zeta(3)) = (0.207976...)*X. The second order term was conjectured by Roberts to be a known constant times X^{5/6}, and this was subsequently proved by Bhargava et al.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christopher E. Thompson_, Nov 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011366 | OEIS-A278832

Maximal material difference at the end of the n-th ply of a chess game.

I conjecture that, unless Black is forced to capture a white piece in all of the maximizing positions, every other term will be equal to the preceding one, a(2n) = a(2n-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Nov 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011367 | OEIS-A278833

Indices of records of A004090(n) - n, where A004090 is the sum of digits of the Fibonacci numbers A000045.

Conjectured to be finite and complete. Indeed it appears that 0.9*n < A004090(n) < n for all sufficiently large n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Dec 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011368 | OEIS-A278847

a(n) = permanent M_n where M_n is the n X n matrix m(i,j) = i^2 + j^2.

Conjecture 1: If 2n+1 is composite, then a(n) == 0 (mod 2n+1). Conjecture 2: If p = 4n+1 is prime, then the sum of those Product_{j=1..2n}(j^2-f(j)^2)^{-1} with f over all the derangements of {1,...,2n} is congruent to 1/(n!)^2 modulo p. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Nov 29 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011369 | OEIS-A278897

First series of Hankel determinants based on Bell numbers of argument k^2, Bell(k^2).

If we regard Bell(k^2) as the k-th Stieltjes moment for k>=0, then the solution of the Stieltjes moment problem is given in the P. Blasiak et al. reference, see below. We conjecture that a(n)>0 for n>=0. The positivity of these Hankel determinants a(n), n>=0 is one of the conditions for the existence of a positive solution. Apparently this solution is not unique.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Karol A. Penson_, Nov 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011370 | OEIS-A278898

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 107", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 30 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011371 | OEIS-A278899

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 107", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 30 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011372 | OEIS-A278900

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 107", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 30 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011373 | OEIS-A278901

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 107", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 30 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011374 | OEIS-A278919

Numbers k such that phi(k-2) divides sigma(k-1)+1.

Conjecture: this sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Nov 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011375 | OEIS-A278937

Numbers k such that 3 is the largest decimal digit of k^3.

Conjecture: sequence is equal to A038444 plus terms of the form 684917*10^k for k >= 0. - _Chai Wah Wu_, Sep 02 2017 Conjecture is true up to 4.8*10^18. - _Giovanni Resta_, Sep 03 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Barker_, Dec 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011376 | OEIS-A278962

Each triple of consecutive terms contains a term that divides the product of the other two terms.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Dec 02 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011377 | OEIS-A278988

a(n) is the number of words of length n over an alphabet of size 3 that are in standard order and which have the property that every letter that appears in the word is repeated.

Conjectures from _Colin Barker_, Nov 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011378 | OEIS-A278989

a(n) is the number of words of length n over an alphabet of size 4 that are in standard order and which have the property that every letter that appears in the word is repeated.

Conjectures from _Colin Barker_, Nov 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011379 | OEIS-A278992

Number of simple chord-labeled chord diagrams with n chords.

Conjecture D-finite with recurrence: +(-n+2)*a(n) +(2*n^2-8*n+7)*a(n-1) +(6*n^2-18*n+11)*a(n-2) +(n-1)*(6*n-11)*a(n-3) +2*(n-1)*(n-2)*a(n-4)=0. - _R. J. Mathar_, Jan 27 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 07 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011380 | OEIS-A279013

a(n) = Sum_{k=0..n} binomial(2*k,k)/(k+1)*binomial(2*n-1,n-k).

Conjecture D-finite with recurrence: 2*n*(n+1)*(2*n-3)*a(n) -n*(101*n^2-312*n+203)*a(n-1) +(995*n^3-5570*n^2+9567*n-4984)*a(n-2) +2*(-2393*n^3+19300*n^2-50494*n+42835)*a(n-3) +4*(2*n7)*(1408*n^2-9889*n+16690)*a(n-4) -2600*(n-5)*(2*n-7)*(2*n-9)*a(n-5)=0. - _R. J. Mathar_, Jan 27 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Dec 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011381 | OEIS-A279014

a(n) = Sum_{k=0..n} fibonacci(k+1)*binomial(2*n-1,n-k).

Conjecture: +n*(3*n-62)*a(n) +(3*n^2+362*n-247)*a(n-1) +(-171*n^2+220*n+162)*a(n-2) +(417*n^2-2570*n+3551)*a(n-3) +2*(27*n-59)*(2*n-7)*a(n-4)=0. - _R. J. Mathar_, Mar 12 2017 Conjecture: +n*(n^2-11*n+22)*a(n) +2*(-4*n^3+45*n^2-113*n+60)*a(n-1) +(15*n^3-173*n^2+530*n-480)*a(n-2) +2*(2*n-5)*(n^2-9*n+12)*a(n-3)=0. - _R. J. Mathar_, Mar 12 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Dec 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011382 | OEIS-A279027

a(n) = number of primes of the form 4k + 3 such that 2n - 4k - 3 is prime.

Conjecture: a(n) > 0 for n >= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Dec 31 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011383 | OEIS-A279028

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 129", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 04 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011384 | OEIS-A279030

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 129", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 04 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 03 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011385 | OEIS-A279040

Even numbers 2k such that the smallest prime p satisfying p+q=2k (q prime) is greater than or equal to sqrt(k).

It is conjectured that a(230) = 503222 is the last term. Oliveira e Silva's work shows that there are no more terms below 4*10^18.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Corinna Regina Böger_, Dec 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011386 | OEIS-A279048

a(n) = 0 whenever n is a practical number (A005153) otherwise least powers of 2 that when multiplied by n becomes practical.

A conjecture by _Zhi-Wei Sun_ states that any rational number can be expressed as the sum of distinct unit fractions whose denominators are practical numbers. To prove this conjecture, _David Eppstein_ (see link) used the fact that every natural number when repeatedly multiplied by 2 will eventually become practical.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Dec 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011387 | OEIS-A279049

A 3-dimensional variant of A269526 "Infinite Sudoku": expansion (read first by layer, then by row) of square pyramid P(n,j,k). (See A269526 and "Comments" below for definition).

Conjecture: all infinite lines (i.e., all vertical columns and some multi-layer diagonals) are permutations of the natural numbers (while this has been proven for rows and columns in A269526, proofs here will require more subtle analysis).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bob Selcoe_, Dec 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011388 | OEIS-A279053

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 131", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 05 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 04 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011389 | OEIS-A279056

Number of ways to write n as w^2 + x^2 + y^2 + z^2 with w a positive integer and x,y,z nonnegative integers such that x^3 + 4*y*z*(y-z) is a square.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 16^k*q (k = 0,1,2,... and q = 1, 79, 143, 184, 575). We have verified a(n) > 0 and part (ii) of the conjecture for n up to 3*10^5. For more conjectural refinements of Lagrange's four-square theorem, see Section 4 of arXiv:1604.06723.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 05 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011390 | OEIS-A279087

Smallest number k such that k^j is (at least conjecturally) pandigital in base n for every j > 0.

Smallest number k such that k^j is (at least conjecturally) pandigital in base n for every j > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Jan 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011391 | OEIS-A279118

Binary representation of the x-axis, from the left edge to the origin, (and also from the origin to the right edge) of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 209", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Aug 02 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011392 | OEIS-A279141

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 137", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011393 | OEIS-A279142

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 137", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011394 | OEIS-A279143

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 137", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011395 | OEIS-A279144

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 137", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 06 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011396 | OEIS-A279212

Fill an array by antidiagonals upwards; in the top left cell enter a(0)=1; thereafter, in the n-th cell, enter the sum of the entries of those earlier cells that can be seen from that cell.

Conjecture: Every column has a finite number of odd entries, and every row and diagonal have an infinite number of odd entries. - _Peter Kagey_, Mar 28 2020. The conjecture about columns is true, see that attached pdf file from _Alec Jones_. Column 0 gives A011782. In the column 1, the only powers of 2 occur at positions A233328(k) with value a(k(k+1)/2 + 1), k >=1 (see A335903). Conjecture: Those are the only multiple occurrences of numbers greater than 1 in this sequence (checked through the first 2000 antidiagonals). - _Hartmut F. W. Hoft_, Jun 29 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011397 | OEIS-A279241

Let f(n) = 4*n^2 + 2*n + 41. The values |f(n)| are primes for all n in the range -20 to 19 (but not for n=-21 or 20). The sequence lists this maximal run of primes in the order in which they appear.

The substitution 2n = m changes this quadratic form into Euler's famous quadratic form m^2+m+41 (see A005846). Concerning the conjectured extremal properties of these forms, one should note the comment from _T. D. Noe_ in A005846. For another quadratic form similar to this one, see A145096. - _N. J. A. Sloane_, Dec 17 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles Kusniec_, Dec 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011398 | OEIS-A279262

Number of n X 2 0..1 arrays with no element equal to a strict majority of its horizontal and vertical neighbors, with the exception of exactly one element, and with new values introduced in order 0 sequentially upwards.

Conjectures from _Colin Barker_, Feb 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 08 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011399 | OEIS-A279387

Irregular triangle read by rows: suppose the symmetric representation of sigma(n) consists of m = A250068(n) layers of width 1, arranged in increasing order; then T(n,k) (n >= 1, 1 <= k <= m) is the number of subparts in the k-th layer.

Conjecture: Each row of the irregular triangle of A262045 can be interpreted as a step function of step sizes 1, 0, and -1. The numbers in row n are the widths of the segments in the parts of the symmetric representation of sigma(n). Each new subpart in a segment (in the left half) of row n starts at the same odd index that represents an odd divisor d of n in the irregular triangle of A237048. Either a subpart ends at an even index e, representing a second odd divisor, which satisfies d * e = oddpart(n), and thus the entire subpart is duplicated in the symmetric portion of the representation, or a subpart runs through the center and continues contiguously into the right half of the symmetric portion of the representation. In other words, the number of subparts in row n equals the number of odd divisors of n, i.e., the conjecture is true. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Dec 12 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011400 | OEIS-A279392

Bisection of primes congruent to 1 modulo 4 (A002144), depending on the corresponding sum of the A002972 and 2*A002973 entries being congruent to 1 modulo 4 or not. Here we give the first case.

This bisection appears in the formula for the p-defects of the congruence y^2 == x^3 + 4*x (mod p) for primes p == 1 (mod 4). See A278720 where for nonvanishing entries the sign is conjectured to be + for these part I primes, and it is - for the part II primes from A279393.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Dec 11 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011401 | OEIS-A279419

Numbers k from A284919 such that either k-1 or k+1 is in A277688.

We conjecture that the sequence is infinite. This sequence shows the proximity of A284919 and A277688 based on Goldbach and Lemoine-Levy representations of even and odd numbers respectively.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011402 | OEIS-A279486

Indices k such that A279484(k) = 0.

Conjecture: for k > 353684 there are no more terms in this sequence (tested for k < 1000000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Dec 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011403 | OEIS-A279487

Indices k such that A279485(k) = 0.

Conjecture: Last term is a(1040799) = 64674419. For k > 64674419 there are no more terms in this sequence (tested for k < 150000000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Dec 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011404 | OEIS-A279498

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 161", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 14 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011405 | OEIS-A279499

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 161", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 14 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011406 | OEIS-A279500

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 161", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 14 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011407 | OEIS-A279501

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 161", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 14 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 13 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011408 | OEIS-A279522

Number of ways to write n as w^2 + x^2 + y^2 + z^2 with w + 2*x + 3*y + 5*z a square, where w,x,y,z are nonnegative integers.

Conjecture: (i) a(n) = 0 if and only if n = 16^k*28 for some k = 0,1,2,....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 14 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011409 | OEIS-A279557

Number of length n inversion sequences avoiding the patterns 110, 120, and 021.

Conjecture: a(n) = C_{n+1}-Sum_{i=1..n} C_i where C_i is the i-th Catalan number, binomial(2i,i)/(i+1). Assuming the conjecture a(n) ~ (64/3)*4^n/((4*n+7)^(3/2)*sqrt(Pi)). - _Peter Luschny_, Feb 24 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Megan A. Martinez_, Jan 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011410 | OEIS-A279561

Number of length n inversion sequences avoiding the patterns 101, 102, 201, and 210.

It is conjectured that a_n also counts those length n inversion sequences with no entries e_i, e_j, e_k (where i<j<k) such that e_i < e_j > e_k and e_i <> e_k. This is the same as the set of length n inversion sequences avoiding 021 and 120.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Megan A. Martinez_, Jan 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011411 | OEIS-A279586

Coefficients in the expansion of ([r^2] + [2r^2]x + [3r^2]x^2 + ...)/([r] + [2r]x + [3r]x^2 + ...); [ ] = floor, r = golden ratio = (1 + sqrt(5))/2.

G.f.: 1 + 1/(1 + x/(1 + x/(1 + x^2/(1 + x^3/(1 + x^5/(1 + x^8/(1 + ... + x^Fibonacci(k)/(1 + ...)))))))), a continued fraction (conjecture). - _Ilya Gutkovskiy_, Apr 18 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011412 | OEIS-A279612

Number of ways to write n = x^2 + y^2 + z^2 + w^2 with x + 2*y - 2*z a power of 4 (including 4^0 = 1), where x,y,z,w are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 16^k*q (k = 0,1,2,... and q = 1, 2, 3, 6, 7, 8, 12, 15, 27, 31, 47, 72, 76, 92, 111, 127). We have verified a(n) > 0 for all n = 1..2*10^7. The conjecture that a(n) > 0 for all n > 0 appeared in arXiv:1701.05868.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011413 | OEIS-A279613

Expansion of the g.f. of A160534 in powers of A121593.

asymptotic conjecture: a(n) ~ C n^(-4/3) 7^n cos(n(arctan((3*sqrt 3)/13) +Pi -1.083913253)), where C = 6.502807770...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lynette O'Brien_, Dec 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011414 | OEIS-A279616

Numbers of the form x^2 + y^2 + z^2 with x + 2*y - 2*z a power of four (including 4^0 = 1), where x,y,z are nonnegative integers.

Part (i) of the conjecture in A279612 implies that any positive integer can be written as the sum of a square and a term of the current sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 15 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011415 | OEIS-A279683

Number of move operations required to sort all permutations of [n] by MTF sort.

Conjecture: primes p such that p^4 divides a(p) are the Wolstenholme primes A088164. - _Amiram Eldar_ and _Thomas Ordowski_, Jan 15 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Dec 16 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011416 | OEIS-A279698

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 185", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011417 | OEIS-A279699

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 185", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011418 | OEIS-A279700

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 185", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011419 | OEIS-A279701

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 185", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 17 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011420 | OEIS-A279732

Lexicographically least strictly increasing sequence such that, for any n>0, Sum_{k=1..n} a(k) can be computed without carries in factorial base.

- Conjecture: this happens infinitely often.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Dec 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011421 | OEIS-A279735

Number of n X 2 0..1 arrays with no element equal to a strict majority of its horizontal and antidiagonal neighbors, with the exception of exactly one element, and with new values introduced in order 0 sequentially upwards.

Conjectures from _Colin Barker_, Feb 11 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 18 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011422 | OEIS-A279840

E.g.f.: exp(Integral exp(x^2) dx).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011423 | OEIS-A279859

a(n) is the integer r with |r| < prime(n)/2 such that (prime(n)*Sum_{k=1..prime(n)-1} (3*H(k-1)^2 + 4*H(k-1)/k)/(k^2*binomial(2k,k)) + 3*H(prime(n)-1)/prime(n)^2)/prime(n)^2 == r (mod prime(n)), where H(m) denotes the harmonic number Sum_{k=1..m} 1/k.

Conjecture: (i) We have the new identity It is known that H(p-1) == -(p^2/3)*B_{p-3} (mod p^3) for any prime p > 3. The first congruence in part (ii) of the conjecture implies that a(n) == -B_{prime(n)-5}/5 (mod prime(n)) for all n = 4,5,....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 20 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011424 | OEIS-A279865

Number of n X 1 0..2 arrays with no element unequal to a strict majority of its horizontal and vertical neighbors, with the exception of exactly two elements, and with new values introduced in order 0 sequentially upwards.

Conjectures from _Colin Barker_, Feb 26 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011425 | OEIS-A279872

Decimal representation of the x-axis, from the left edge to the origin, (and also from the origin to the right edge) of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 209", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Aug 02 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011426 | OEIS-A279873

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 211", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 22 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011427 | OEIS-A279874

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 211", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 22 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011428 | OEIS-A279875

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 211", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 21 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011429 | OEIS-A279876

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 211", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 21 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011430 | OEIS-A279877

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 213", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 21 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011431 | OEIS-A279878

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 213", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 21 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011432 | OEIS-A279879

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 213", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 22 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011433 | OEIS-A279880

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 213", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 22 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 21 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011434 | OEIS-A279897

Number of n X 3 0..2 arrays with no element unequal to a strict majority of its king-move neighbors, with the exception of exactly one element, and with new values introduced in order 0 sequentially upwards.

Conjectures from _Colin Barker_, Feb 12 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Dec 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011435 | OEIS-A279915

Numbers m that can be written as x*y with phi(x)*sigma(y) = 2*x*y, where x and y are positive integers, phi(.) is Euler's totient function and sigma(y) is the sum of all positive divisors of y.

Conjecture: (i) All the terms are even. Moreover, if x and y are positive integers with phi(x)*sigma(y) = 2*x*y, then y must be even. As phi(1) = 1, the sequence contains all perfect numbers, and part (i) of the above conjecture implies the well-known conjecture that there are no odd perfect numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 22 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011436 | OEIS-A279959

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 221", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 24 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011437 | OEIS-A279960

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 221", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 24 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011438 | OEIS-A279961

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 221", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 24 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011439 | OEIS-A279962

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 221", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 24 2016: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Dec 23 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011440 | OEIS-A279967

Square array read by antidiagonals upwards in which each term is the sum of prior elements in the same row, column, diagonal, or antidiagonal that divide n; the array is seeded with an initial value a(1)=1.

Conjecture: This conjecture together with Bouniakowsky's conjecture that certain quadratic integer polynomials generate infinitely many primes (e.g. see A002496 for n^2+1 and A188382 for 2*n^2+n+1) implies that in every column in the triangle infinitely many prime sequence indices occur and therefore infinitely many 0's whenever the column contains no 1's. The proof is based on the fact that for a large enough prime sequence index p in whose prior column no 1 occurs then a(p)=0; therefore infinitely many 0's occur in that column. Obviously, once value 1 occurs in a column no 0 value can occur in a subsequent row. Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alec Jones_, Dec 24 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011441 | OEIS-A280051

Let c_n(k) be the sequence defined in A278743; here we give the associated values of k0.

Let c_n(k) be the sequence defined in A278743, for n >= 2. It is conjectured that there are numbers k0 and b such that c_n(k) satisfies the recurrence c_n(k + A278743(n)) = c_n(k)*n^b for k > k0. Here we give the values of k0. The values of b are given in A280052.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011442 | OEIS-A280052

Let c_n(k) be the sequence defined in A278743; here we give the associated values of b.

Let c_n(k) be the sequence defined in A278743, for n >= 2. It is conjectured that there are numbers k0 and b such that c_n(k) satisfies the recurrence c_n(k + A278743(n)) = c_n(k)*n^b for k > k0. Here we give the values of b. The values of k0 are given in A280051.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011443 | OEIS-A280074

Numbers k such that Sum_{d|k} tau(d) = Sum_{d|k+1} tau(d).

Sequence of the smallest numbers k such that Sum_{d|k} tau(d) = Sum_{d|k+1} tau(d) = ... = Sum_{d|k+n-1} tau(d) for n>=1: 1, 2, 33, 19940, 204323, 380480345, 440738966073, ...; conjecture: this sequence is different from A034173.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Dec 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011444 | OEIS-A280098

The sum of the divisors of 24*n - 1, divided by 24.

Conjecture: only the integers k in {1, 3, 4, 6, 8, 12, 24} have the property that the sum of the divisors of (k*n-1)/k is always an integer. - _Robert G. Wilson v_, Dec 25 2016 The finite sequence mentioned in the above conjecture gives the sum of the divisors of the partition numbers of the first seven positive integers (cf. A139041). - _Omar E. Pol_, Dec 25 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Dec 25 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011445 | OEIS-A280134

Number of primes of the form 4(n - k - 1) + 3 where 0 <= k and prime of the form 4k + 3 <= n.

(1) Conjecture: a(n) > 0 for n >= 11. (2) Conjecture: A278287(n) > 0 for n >= 11. Conjecture (1) and conjecture (2) are first and second parts of Goldbach-like conjecture in A001031.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Dec 26 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011446 | OEIS-A280153

Number of ways to write n as x^3 + 2*y^3 + z^2 + 4^k, where x is a positive integer and y,z,k are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 3, 4, 7, 10, 17, 24, 36, 38, 52, 62, 115, 136, 185, 990. We have verified that a(n) > 0 for all n = 2..10^6, and that part (ii) of the conjecture holds for all n = 1..10^6. See also A280356 for a similar conjecture involving powers of two.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011447 | OEIS-A280172

Lexicographically earliest table of positive integers read by antidiagonals such that no row or column contains a repeated term.

Conjecture: The (2^k)-th antidiagonal consists entirely of 2^k. Conjecture: T(2n, k) = 2 * T(n, ceiling(k/2)), n >= 1, 1 <= k <= 2n. E.g. Conjectured algorithm for equilateral triangle (1-indexed rows and row terms), whose concatenated rows give this sequence: T(1, 1) = 1;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Dec 27 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011448 | OEIS-A280183

Positions of pandigital 10-digit numbers after the decimal point in the decimal expansion of Pi.

It is conjectured that Pi is a normal number in base 10, which would imply that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bobby Jacobs_, Dec 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011449 | OEIS-A280185

a(n) = n - A004090(n), where A004090 is the sum of digits of the Fibonacci numbers A000045.

Conjectured to increase to infinity. It appears that the slope of A004090(n) is roughly 0.93, at least in the range 0..10^5. I conjecture that this sequence takes its minimum at a(2619) = -92. - _M. F. Hasler_, Dec 30 2016

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Dec 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011450 | OEIS-A280199

Numbers n such that a^(n-1) == 1 (mod n^2) has solutions with 1 < a < n^2-1.

Conjecture: composite terms are A181780.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_ and _Thomas Ordowski_, Dec 28 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011451 | OEIS-A280246

a(n) = Product_{d|n} A023896(d), where A023896(m) is the sum of totatives of m.

Conjecture: a(n) is odd iff the sum of totatives of n (A023896) is odd. [Proved in Li (2026) - _Wentao Li_, Sep 10 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Dec 30 2016 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011452 | OEIS-A280258

a(n) = Sum_{d|n} pxi(d), where pxi(m) is the product of totatives of m (A001783).

Conjecture: a(n) is odd for numbers in A183300; a(n) is even for numbers in A001105 (2*n^2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jan 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011453 | OEIS-A280321

Number of 2 X 2 matrices with all elements in {0,..,n} having determinant = n*permanent.

Conjectures from _Colin Barker_, Jan 01 2017: (Start) Barker's conjectures are true. The recursion and g.f. follow from the formula for a(n). To see why the formula is true, write the condition as (a*d-b*c)=n*(a*d+b*c) and assume that n>=2. Rearranging, (1-n)*a*d=(1+n)*b*c. Since 1-n<0 and 1+n>0, necessarily a*d=b*c=0. Thus, at least one of a,d and, independently, at least one of b,c must be 0. This gives 2*n+1 possibilities for each of the two pairs, proving that a(n)=(2*n+1)^2. The cases n=0,1 are verified directly. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Indranil Ghosh_, Jan 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011454 | OEIS-A280352

Expansion of Sum_{k>=1} (x/(1 - x))^(k*(k+1)/2).

Conjecture: a(n+1)/a(n) ~ 2. - _Jerzy R Borysowicz_, Jan 14 2023 Conjecture: abs(b(n)-1) < 0.015, where b(n) = a(n)*sqrt(n)/2^(n-1), for n > 781; b(n) does not have a limit. - _Jerzy R Borysowicz_, Feb 17 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Jan 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011455 | OEIS-A280356

Number of ways to write n as x^4 + y^3 + z^2 + 2^k, where x,y,z are nonnegative integers and k is a positive integer.

Conjecture: (i) a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 23, 1135, 6415, 6471. We have verified that a(n) > 0 for all n = 2..2*10^7, and that part (ii) of the conjecture holds for all n = 1..10^7. See also A280153 for a similar conjecture involving powers of 4 or 8. Qing-Hu Hou at Tianjin Univ. has verified that a(n) > 0 for all n = 2..10^9. In 2017, the author announced to offer US $234 as the prize for the first correct solution to his conjecture that a(n) > 0 for all n > 1. - _Zhi-Wei Sun_, Dec 30 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011456 | OEIS-A280364

Number of 2 X 2 matrices with all elements in {0,...,n} with permanent = determinant^n.

Conjectures from _Colin Barker_, Jan 01 2017: (Start) The conjectures are true (see Fried link). - _Sela Fried_, Feb 02 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Indranil Ghosh_, Jan 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011457 | OEIS-A280367

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 259", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jan 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011458 | OEIS-A280368

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 259", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jan 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011459 | OEIS-A280369

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 259", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jan 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011460 | OEIS-A280370

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 259", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jan 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011461 | OEIS-A280371

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 260", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jan 02 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011462 | OEIS-A280373

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 260", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jan 02 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011463 | OEIS-A280386

Number of ways to write n as the sum of two squares and a term of A000009.

Conjecture: a(n) > 0 for all n > 0. The conjecture was verified by the author for n up to 4*10^6. After learning this conjecture from the author, Prof. Qing-Hu Hou at Tianjin Univ. finished his verification of the above conjecture for n up to 10^9. - _Zhi-Wei Sun_, Jan 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011464 | OEIS-A280399

Number of 1 X n 0..2 arrays with no element unequal to a strict majority of its horizontal and antidiagonal neighbors, with the exception of exactly one element, and with new values introduced in order 0 sequentially upwards.

Conjectures from _Colin Barker_, Feb 13 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011465 | OEIS-A280406

Odd semiprimes that can be represented as 2p+3q, where p and q are primes, in an increasing number of ways.

It is conjectured that all terms of this sequence above 1963 are divisible by 5. Initial search to 1000000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Randy L. Ekl_, Jan 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011466 | OEIS-A280410

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 261", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jan 03 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011467 | OEIS-A280411

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 261", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jan 03 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011468 | OEIS-A280412

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 261", based on the 5-celled von Neumann neighborhood.

Conjectures from _Ilya Gutkovskiy_, Jan 02 2017: (Start)
Conjectures from _Colin Barker_, Jan 03 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011469 | OEIS-A280413

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 262", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011470 | OEIS-A280414

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 262", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011471 | OEIS-A280415

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 262", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011472 | OEIS-A280416

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 262", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011473 | OEIS-A280417

Number of distinct length-n blocks (a.k.a. subword complexity) of the characteristic sequence of the prime numbers A010051.

Unlike A023192, this sequence also counts blocks that occur finitely often in A010051. And unlike A023192, the correctness of the numbers provided here depend on no conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Jan 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011474 | OEIS-A280430

Longest word T from 2 equal length strings S using no breakpoint reuse.

Conjecture: (2 +2*x +3*x^2 -7*x^3 -9*x^4 -6*x^5 +14*x^6 +12*x^7 +7*x^8 -7*x^9 -6*x^10 -3*x^11 +x^13+x^14)/ (1+x+x^2)/ (x-1)^2/ (x^6-3*x^3+1)^2 . - _R. J. Mathar_, Mar 06 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nicole Anderson_, Jan 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011475 | OEIS-A280444

Least positive integer m such that n - p(m) = x*(3x-1)/2 + y*(3y+1)/2 for some nonnegative integers x and y, or 0 if no such m exists, where p(.) is the partition function given by A000041.

The conjecture in A280455 asserts that a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011476 | OEIS-A280455

Number of ways to write n as x*(3x-1)/2 + y*(3y+1)/2 + p(z), where x,y,z are nonnegative integers with z > 0, and p(.) is the partition function given by A000041.

Conjecture: (i) a(n) > 0 for all n > 0. This is similar to the author's conjecture in A280386. At the author's request, Prof. Qing-Hu Hou at Tianjin Univ. has verified part (i) of the above conjecture for n up to 10^9. We also have some other similar conjectures. For example, we conjecture that any positive integer can be expressed as the sum of two triangular numbers and a partition number. See also A280472 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011477 | OEIS-A280472

Number of ways to write n as the sum of an octagonal number (A000567), a second octagonal number (A045944), and a strict partition number (A000009).

Conjecture: (i) a(n) > 0 for all n > 0. On the author's request, Prof. Qing-Hu Hou at Tianjin Univ. has verified part (i) of the above conjecture for n up to 10^9. See also A280455 for a similar conjecture of the author involving the partition function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011478 | OEIS-A280474

Number of n X 2 0..2 arrays with no element unequal to a strict majority of its horizontal and antidiagonal neighbors, with the exception of exactly two elements, and with new values introduced in order 0 sequentially upwards.

Conjectures from _Colin Barker_, Feb 13 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011479 | OEIS-A280541

G.f.: Product_{i>=1, j>=1} (1 + x^(i*j))^(i*j).

Conjecture: log(a(n)) ~ 3 * Zeta(3)^(1/3) * log(n)^(1/3) * n^(2/3) / 2^(4/3). - _Vaclav Kotesovec_, Aug 29 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jan 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011480 | OEIS-A280547

Smallest number k such that (k+1)^n - k^n is divisible by a square greater than 1.

Conjecture: for n <= 100, there is no square greater than (10^6)^2 = 10^12 that divides (k+1)^n - k^n at any value of k lower than the upper bound described above; i.e., the upper bound described above is equal to a(n) for each n <= 100. (End) Confirmed Jon E. Schoenfield's conjecture through a(58). - _Robert Price_, Feb 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jan 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011481 | OEIS-A280580

Triangle read by rows: T(n,k) = binomial(2*n,2*k)*binomial(2*n-2*k,n-k)/(n+1-k) for 0<=k<=n.

Conjectures: (1) Sum_{k=0..n} (-1)^k*T(n,k)*A238390(k) = A000007(n);

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Jan 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011482 | OEIS-A280606

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 294", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jan 07 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011483 | OEIS-A280607

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 294", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jan 07 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011484 | OEIS-A280608

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 294", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jan 07 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011485 | OEIS-A280610

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 294", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jan 07 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011486 | OEIS-A280659

Lexicographically earliest sequence of distinct positive terms such that the sum of two consecutive terms has at least 5 distinct prime factors.

Conjecturally: this sequence is a permutation of the natural numbers, and a(n) ~ n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011487 | OEIS-A280668

Number of n X 3 0..2 arrays with no element equal to more than one of its horizontal and antidiagonal neighbors and with new values introduced in order 0 sequentially upwards.

Conjectures from _Colin Barker_, Feb 14 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011488 | OEIS-A280681

Numbers k such that Fibonacci(k) is a totient.

Conjecture: Sequence is infinite. - _Altug Alkan_, Jul 05 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Jan 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011489 | OEIS-A280683

Number of ways to write n as an ordered sum of two positive squarefree semiprimes (A006881).

Conjecture: a(n) > 0 for n > 82 (see comment in A006881 from _Richard R. Forberg_).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Jan 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011490 | OEIS-A280693

Numbers n such that A003961(n) = A250469(n).

Positions of zeros in A280692. Conjectured to be also the positions of ones in A280703.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011491 | OEIS-A280717

Given a prime number p, let b = -p and c = p^2. Assuming that the polynomial P(x) := x^2+b*x+c takes at least one prime value for some positive integer x<p, we define the next term of the sequence to be the maximal element in the set S := {P(x) : x is an integer, 0 < x < p, and P(x) is a prime number}. The first term of the present sequence is p=3.

Conjecture: the sequence is well defined.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luis H. Gallardo_, Jan 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011492 | OEIS-A280738

After S(n)=A280864(n) has been computed, let p(n) = product of distinct primes shared by S(n-1) and S(n); let q(n) = product of distinct primes in S(n) but not in S(n-1); and let r(n) = smallest number not yet in S. Sequence gives p(n).

By definition, all terms are squarefree. Let {i,j,k} be distinct fixed positive numbers. Conjecture: All squarefree numbers appear infinitely often, and all terms a(n) = j are immediately preceded and followed infinitely often by all terms a(n-1) = i and a(n+1) = k. If so, then A280864 is a permutation of the natural numbers. - _Bob Selcoe_, Apr 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011493 | OEIS-A280745

Primes that end a block of terms in A280864.

It is not difficult to show (see A280864) that when a prime appears "naked" in A280864, it either begins a block, in which case the next term is 2*p, or it ends a block. In the latter case we conjecture that the previous term in A280864 is 2*p. This is true for the first 10^7 terms of A280864.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011494 | OEIS-A280801

Least k > 0 such that (2*n)^k is in A002202, or 0 if no such k exists.

Conjecture: a(n) > 0 for all values of n. - _Altug Alkan_, Jan 11 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Jan 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011495 | OEIS-A280849

Square array T(j,k) read by antidiagonals upwards, in which column k lists the numbers n having k odd divisors greater than sqrt(2*n), with j >= 1, k >= 0.

Conjecture: column k lists also the numbers n having k pairs of equidistant subparts in the symmetric representation of sigma(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Feb 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011496 | OEIS-A280850

Irregular triangle read by rows in which row n is constructed with an algorithm using the n-th row of triangle A196020 (see Comments for precise definition).

Conjecture: the odd-indexed terms of the n-th row together with the even-indexed terms of the same row but listed in reverse order give the n-th row of triangle A296508 (this is the same conjecture from A296508). - _Omar E. Pol_, Apr 20 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jan 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011497 | OEIS-A280866

Lexicographically earliest sequence of distinct terms such that, for any prime p, any run of consecutive multiples of p has at least length 2.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jan 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011498 | OEIS-A281000

Triangle read by rows: T(n,k) = binomial(2*n+1, 2*k+1)*binomial(2*n-2*k, n-k)/(n+1-k) for 0 <= k <= n.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Jan 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011499 | OEIS-A281008

Least positive integer k with exactly n odd divisors greater than sqrt(2*k).

Conjecture: a(n) is also the smallest number k having n pairs of equidistant subparts in the symmetric representation of sigma(k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Feb 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011500 | OEIS-A281012

Zig-zag path that we can find descending by the main diagonal of the pyramid described in A245092.

Conjecture: the odd-indexed terms give A281007, the positive terms in A067742.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Feb 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011501 | OEIS-A281022

Single (or isolated or non-twin) primes that are also safe primes.

It is conjectured that there are infinitely many safe primes, but this is still unproved, so it is not known whether this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Jan 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011502 | OEIS-A281112

Decimal expansion of (conjectured) limit c(n+1)/c(n), where c = A078140.

Decimal expansion of (conjectured) limit c(n+1)/c(n), where c = A078140.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011503 | OEIS-A281131

First appearance of 2^n in A281130.

Conjecture: a(n) ~ 2^n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rok Cestnik_, Jan 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011504 | OEIS-A281146

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 515", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011505 | OEIS-A281180

a(n) = (2*n-1)! * [x^(2*n-1)] S(x), where S(x) = Integral (1 + S(x)^2)^2 dx.

Conjecture: the sequence {a(n)} reduced modulo a positive integer k is eventually periodic with the period dividing phi(k)^2, where phi(n) = A000010(n). An example is given below. - _Peter Bala_, Feb 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011506 | OEIS-A281181

E.g.f. C(x) satisfies C(x) = cosh(Integral C(x)^3 dx).

More generally, we conjecture that the sequence {a(n)} reduced modulo a positive integer k is eventually periodic with the period dividing phi(k)^2, where phi(n) = A000010(n). An example is given below. - _Peter Bala_, Feb 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011507 | OEIS-A281183

E.g.f. C(x)^2 = cosh(Integral C(x)^3 dx)^2 where C(x) is described by A281181.

Conjecture: the sequence {a(n)} taken modulo a positive integer k is eventually periodic with the period dividing phi(k)^2, where phi(n) = A000010(n). An example is given below. - _Peter Bala_, Feb 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011508 | OEIS-A281186

E.g.f. satisfies: A(x) = exp(1/A(x) * Integral A(x)^(8/3) dx).

Conjecture: if G(x) = exp(1/G(x) * Integral G(x)^t dx) for some fixed real t, then G(x) will consist entirely of nonnegative coefficients iff t >= 8/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 31 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011509 | OEIS-A281188

Number of refactorable numbers m such that tau(m) = n, or 0 if there are infinitely many such numbers.

Conjecture: a(n) is nonzero if and only if n is squarefree or n = 4. [This conjecture is true; see Links for a proof. - _Jon E. Schoenfield_ and _Altug Alkan_, Jan 17 2017] See also Theorem 5 for the proof of conjecture in Colton link. - _Altug Alkan_, Jan 20 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franklin T. Adams-Watters_, Jan 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011510 | OEIS-A281189

a(n) is the first composite number having the same base-(2n) digits as its prime factors (with multiplicity), excluding zero digits (or 0 if no such composite number exists).

Conjecture: a(n) always exceeds 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ely Golden_ and _Robert G. Wilson v_, Jan 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011511 | OEIS-A281196

Number of n such that A247600(n) = 5.

Conjecture: This sequence has just 22 terms shown.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Jan 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011512 | OEIS-A281199

Number of n X 2 0..1 arrays with no element equal to more than one of its horizontal and antidiagonal neighbors, with the exception of exactly one element, and with new values introduced in order 0 sequentially upwards.

Conjectures from _Colin Barker_, Feb 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011513 | OEIS-A281200

Number of n X 3 0..1 arrays with no element equal to more than one of its horizontal and antidiagonal neighbors, with the exception of exactly one element, and with new values introduced in order 0 sequentially upwards.

Conjectures from _Colin Barker_, Feb 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Jan 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011514 | OEIS-A281214

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 342", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jan 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011515 | OEIS-A281215

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 342", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jan 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011516 | OEIS-A281216

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 342", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jan 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011517 | OEIS-A281217

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 342", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = floor(16^(floor(n/4)+1)/15)*((n+1) mod 2) + 52*floor(16^(floor(n/4))/15)*0^((n+3) mod 4) + 7*floor(16^(floor(n/4)+1)/15)*0^((n+1) mod 4). - _Karl V. Keller, Jr._, Sep 27 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011518 | OEIS-A281229

Smallest number k of the form x^2 + y^2 with 0 <= x <= y such that gcd(x, y) = 1, x + y = n, and k has no other decompositions into a sum of two squares.

Conjecture: for each n there exists such a number k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Jan 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011519 | OEIS-A281256

Runs of consecutive integers in A270877, which is produced by a decaying trapezoidal modification of the sieve of Eratosthenes.

The author conjectures that T(n,m) exists for all n>=1, m>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Munn_, Jan 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011520 | OEIS-A281260

Triangular array of generalized Narayana numbers T(n,k) = 2*binomial(n+1,k)* binomial(n-2,k-1)/(n+1) for n >= 1 and 0 <= k <= n-1, read by rows.

Conjecture: Sum_{k=1..n-m} binomial(n+1-k,m) * N_m(n,k) * A103365(n+1-m-k) = (m+1)^2 * A000007(n-m-1) for n > m >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Jan 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011521 | OEIS-A281267

Main diagonal of A276554.

Conjecture: the stronger supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(2*k)) hold for all primes p >= 3 and all positive integers n and k. (End) Adamczewski includes proof abstract regarding the above conjecture in linked article. - _Michael De Vlieger_, Aug 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Apr 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011522 | OEIS-A281281

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 350", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011523 | OEIS-A281282

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 350", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011524 | OEIS-A281283

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 350", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011525 | OEIS-A281284

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 350", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011526 | OEIS-A281291

Numbers k such that 2*k! is not a refactorable number.

See Conjecture 47 and Theorem 51 in Zelinsky's paper for related points.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Jan 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011527 | OEIS-A281292

Squarefree numbers that are the sum of two squares in exactly one way.

Conjecture: each positive integer is a sum |x| + |y| such that x^2 + y^2 is in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Jan 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011528 | OEIS-A281312

Numbers n such that sigma(4*(n-1)) is prime.

Conjecture: the next terms are: 288230376151711745, 773712524553362671811952 65, 202824096036516704239472512860 17, 212676479325586539664609129644855 13217. Conjecture: prime terms are in A258429: 2, 5, 17, 65537. Conjecture: corresponding primes p are Mersenne primes (A000668) > 3. Sigma is multiplicative, and sigma(m) > 1 for all m > 1, so sigma(m) can be prime only if m is a prime power. Hence all n in this sequence are of the form 2^m + 1 for some m >= 0. This proves the above conjectures and leads to an explicit formula (q.v.) in terms of the Mersenne numbers. - _Charles R Greathouse IV_, Mar 01 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jan 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011529 | OEIS-A281369

Orbit of 25 under repeated applications of the map m -> A280864(m).

It is a strong conjecture that A280864 is a permutation of the positive integers. The orbit containing 25 appears to be the first infinite orbit. See A281370 for the orbit of 25 under the inverse permutation, and see A280864 for other orbits (there are several small cycles).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011530 | OEIS-A281387

Pairs (x, y) of relatively prime positive integers such that (x^2 - 5)/y and (y^2 - 5)/x are both positive integers.

Conjectures from _Chai Wah Wu_, Jan 28 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011531 | OEIS-A281409

Lexicographically first sequence of distinct terms, beginning with a(1)=5, with the property that each triple of consecutive terms contains a term that divides the sum of the other two terms.

Conjecturally, all other variants are not ultimately linear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jan 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011532 | OEIS-A281428

E.g.f. C(x) satisfies C(x) = cosh(Integral C(x)^4 dx).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011533 | OEIS-A281440

E.g.f. C(x) satisfies: C(x) = cosh(Integral C(x)^7 dx).

Conjecture: the sequence {a(n)} reduced modulo a positive integer k is eventually periodic with the period dividing phi(k)^2, where phi(n) = A000010(n). An example is given below. - _Peter Bala_, Feb 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011534 | OEIS-A281505

Numbers of the form y^2 - x^2 such that x^2 + y^2 is a prime and 0 < x < y.

a(n) = n(log n)^c /(log log n)^O(1), where c = 1 - (1 + log log 2)/log 2 = 0.086... Cf. A027424. - Conjectured by Carl Pomerance, Jan 25 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Jan 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011535 | OEIS-A281508

Numbers requiring exactly 261 'Reverse and Add' steps to reach a palindrome.

It is only a conjecture that there are no further terms. - _N. J. A. Sloane_, Jan 24 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | Andrey S. Shchebetov and _Sergei D. Shchebetov_, Jan 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011536 | OEIS-A281532

a(n) = least numerator X such that the proper fraction X/A281529(n) needs n or more terms to be written as a signed sum of distinct unit fractions.

Sierpiński conjectured that for any integer n > 3 there exist positive integers a, b, c so that 5/n = 1/a + 1/b + 1/c. If the conjecture is true, then 5 appears only once in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Jan 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011537 | OEIS-A281543

Number of partitions n = x + y with y >= x > 0 such that x^2 + y^2 or (x^2 + y^2)/2 is prime.

Conjecture: a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_ and _Altug Alkan_, Mar 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011538 | OEIS-A281577

Irregular triangle read by rows: T(n, k) = A281576(n) modulo p^2, where p is the k-th prime factor of A281576(n) with p < sqrt(A281576(n)).

The question whether T(n, k) = 0 for any values of n and k is an open problem (see Ribenboim p. 64, open problem (3)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Jan 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011539 | OEIS-A281579

Lexicographically earliest sequence such that, for any n>0, a(n)=length of the n-th run of consecutive terms in arithmetic progression in this sequence.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jan 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011540 | OEIS-A281622

Numbers k such that sigma(k-1) is a Mersenne prime (A000668).

Conjecture 1: Union of 26 and A256438. Conjecture 2: Mersenne prime 31 is the only prime p such that p = sigma(x-1) = sigma(y-1) for distinct numbers x and y; 31 = sigma(17-1) = sigma(26-1). Conjecture: a(n) = 2^A090748(n) + 1. - _Daniel Suteu_, Feb 08 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jan 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011541 | OEIS-A281659

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with 9*x^2 + 16*y^2 + 24*z^2 + 48*w^2 a square, where x,y,z,w are nonnegative integers.

Conjecture: (i) a(n) > 0 except for n = 71, 85. We have verified part (i) of the conjecture for n up to 1.2*10^5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011542 | OEIS-A281708

a(n) = (Sum_{k=1..n} k^3 * p(k) * p(n-k)) * 2/n where p = A000041.

See A067567 for a conjecture about this sequence from _Peter Bala_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Jan 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011543 | OEIS-A281733

Positive integers T_i such that Sum_{k >= 0} (S_k * x^(2*k+1)) + 1/24 - Sum_{k >= 1} (T_k * x^(2*k)) = (cos((2/3) * arccos(6 * sqrt(3) * x)))/12 for all real x with |x| <= 1/(6 * sqrt(3)), where S_k = A176898(k).

Conjecture: T_p == -2 (mod p) for any prime p (cf. Sun (2013), Conjecture 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Jan 31 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011544 | OEIS-A281735

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 390", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011545 | OEIS-A281736

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 390", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011546 | OEIS-A281737

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 390", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011547 | OEIS-A281738

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 390", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011548 | OEIS-A281739

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 393", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011549 | OEIS-A281740

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 393", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011550 | OEIS-A281741

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 393", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011551 | OEIS-A281742

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 393", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011552 | OEIS-A281748

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 395", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011553 | OEIS-A281749

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 395", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011554 | OEIS-A281750

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 395", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011555 | OEIS-A281751

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 395", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jan 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011556 | OEIS-A281817

a(n) = 2*Sum_{k odd} k!*Stirling2(n,k)/(k + 1).

The numbers a(n) are derived from a formula for the numbers Bernoulli(n). Surprisingly, there also appears to be a connection between a(2*n) and Bernoulli(2*n - 2): we conjecture a(2*n) - 1 = integer * the denominator of Bernoulli(2*n - 2) = integer * (Product_{p prime, p - 1 | 2*n - 2} p) (checked up to n = 200). For example, a(14) - 1 = 956922611190 is divisible by 2*3*5*7*13 where 2, 3, 5, 7 and 13 are the primes p such that p - 1 divides 12, while a(18) - 1 = 241805428075379310 is divisible by 2*3*5*17 where 2, 3, 5 and 17 are the primes p such that p - 1 divides 16. For prime p, a(p) = 1 (mod p). Conjecture: for prime p, a(2*p) = 1 (mod p).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jan 31 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011557 | OEIS-A281820

Numerator of Sum_{k=1..n} (30k-11)/(4*(2k-1)*k^3*binomial(2k,k)^2).

Conjecture: Sum_{n >= k+1} 1/(n^3*(n^2 - 1)^2*(n^2 - 4)^2*...*(n^2 - k^2)^2) = Sum_{n >= k+1} 1/(n*binomial(n,k)^2*binomial(n+k,k)^2*(n-k)^2) = zeta(3) - A281820(k)/A281821(k). - _Peter Bala_, Jan 17 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Jan 31 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011558 | OEIS-A281826

Number of ways to write n as x^3 + 2*y^3 + 3*z^3 + p(k), where x,y,z are nonnegative integers, k is a positive integer, and p(.) is the partition function given by A000041.

Conjecture: (i) a(n) > 0 for all n > 0. We have verified the conjecture for n up to 1.3*10^6. On the author's request, Prof. Qing-Hu Hou at Tianjin Univ. has verified all the three parts of the conjecture for n up to 10^9. - _Zhi-Wei Sun_, Feb 06 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 31 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011559 | OEIS-A281882

Numbers n such that the number of divisors of sum of divisors of n is prime.

Conjecture: 2 and 3 are the only prime numbers in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Feb 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011560 | OEIS-A281900

Index of first occurrence of n in A281579, or -1 if n does not appear there..

It is conjectured that every number >= 2 does appear in A281579.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Feb 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011561 | OEIS-A281910

Minimum possible absolute value over all coefficients of p(x)/(1-x)^n, where p is a power series with +-1 coefficients.

It is known that a(7) >= 2124, and that is conjectured to be the true value.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Feb 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011562 | OEIS-A281912

Number of sequences of balls colored with at most n colors such that exactly one ball is of a color seen earlier in the sequence.

Conjecture: a(n) = 2*A394975(1,n)/(n+1) = n*(a(n-1) + A000522(n) - A000522(n-1)). - _Mikhail Kurkov_, Apr 10 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeremy Dover_, Feb 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011563 | OEIS-A281939

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x - y and 3*z + w both squares, where x,y,z are nonnegative integers and w is an integer.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011564 | OEIS-A281941

Number of ways to write n as w^2 + x^2 + y^2 + z^2 with w and w + x + y + z both squares, where w,x,y,z are integers with |x| <= |y| <= |z|.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011565 | OEIS-A281946

Number of sequences of balls colored with at most n colors such that exactly two balls are of a color seen earlier in the sequence.

Conjecture: a(n) = 3*A394975(2,n)/binomial(n+2,2) = n*(a(n-1) + A281912(n) - A281912(n-1)). - _Mikhail Kurkov_, Apr 10 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeremy Dover_, Feb 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011566 | OEIS-A281975

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers and z <= w such that both x and |x-y| are squares.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,.... See also A281976 and A281977 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011567 | OEIS-A281977

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that both x and -7*x - 8*y + 8*z + 16*w are squares.

Conjecture: a(n) > 0 for all n = 0,1,2,.... We have verified the conjecture for all n = 0..10^6. See also A281976, A282013 and A282014 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011568 | OEIS-A281978

Lexicographically earliest sequence of distinct terms such that, for any n>0, a(2n) is divisible by a(2n-1) and by a(2n+1).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Feb 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011569 | OEIS-A281980

Numbers of the form x^4 + y^2 with x^2 + 24*y a square, where x and y are nonnegative integers.

The conjecture in A281976 implies that any nonnegative integer can be written as the sum of two squares and a term of the current sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011570 | OEIS-A282002

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 413", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 05 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011571 | OEIS-A282003

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 413", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 05 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011572 | OEIS-A282004

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 413", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 05 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011573 | OEIS-A282005

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 413", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 05 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011574 | OEIS-A282013

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that both x and 49*x + 48*(y-z) are squares.

Conjecture: a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 16^k*m (k = 0,1,2,... and m = 5, 8, 14, 31, 47, 71, 79, 143, 248, 463, 1039). See also A281976, A281977 and A282014 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011575 | OEIS-A282014

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that both x and 121*x + 48*(y-z) are squares.

Conjecture: a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 16^k*m (k = 0,1,2,... and m = 8, 29, 31, 40, 94, 104, 143, 319, 671). See also A281976, A281977 and A282013 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011576 | OEIS-A282031

Coefficients in q-expansion of (9*E_2(q^3) - E_2(q))/8.

Conjecture: -3*A122858(n) - A229616(n) + 4*a(n) = 0 for all n. - _Thomas Baruchel_, Jun 23 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011577 | OEIS-A282044

Reduced Kronecker coefficients for the case a=2, b=3, i=4.

Conjectured g.f.: x^4*(1+x^2)/(1-2*x+x^3+3*x^5-4*x^6). - _Jean-François Alcover_, Feb 18 2019.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011578 | OEIS-A282058

Numbers that are not triangular numbers and cannot be written as the sum of a triangular number and two odd squares.

The sequence is conjectured to be complete (cf. Sun, 2009, Conjecture 1.2 (ii)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Feb 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011579 | OEIS-A282088

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 553", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 27 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011580 | OEIS-A282089

Decimal expansion of constant 1.287194... related to a conjectural Viète-like formula for Pi.

Decimal expansion of constant 1.287194... related to a conjectural Viète-like formula for Pi. Conjecture: Pi = lim_{k -> infinity} 2^{k + 1}*(1 - c_k), where the variable c_k is defined by a set of the Viète-like recurrence relations {a_1 = sqrt(2), a_k = sqrt(2 + a_{k - 1}), b_k = sqrt(2 - a_k)/a_{k + 1}, c_1 = b_1, c_k = (c_{k - 1} + b_k)/(1 - c_{k - 1}*b_k)}. From this conjecture it follows that Sum_{k >= 1} (1 - c_k) is convergent [Abrarov and Quine].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sanjar Abrarov_, Feb 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011581 | OEIS-A282091

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x + y - z a cube of an integer, where x,y,z,w are nonnegative integers with x >= y <= z and x == y (mod 2).

Conjecture: (i) a(n) > 0 for all n = 0,1,2,.... Also, any nonnegative integer can be written as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers and x <= y <= z such that x + y - z is a cube of an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011582 | OEIS-A282095

Larger member of a coprime pair (x,y) which solves x^2 + y^2 = z^3 with positive x, y and z.

The ordered values of the bases of the cubes, z, are a subsequence of (and conjecturally the same as) A008846.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Feb 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011583 | OEIS-A282100

a(n) is the least number of successive prime numbers modulo 100 required such that their sum is at least 100.

I conjecture that the limit (a(1) + ... + a(n))/n exists and is equal to e = 2.71828182845904523536028747l.... On the contrary, I conjecture that the limit is smaller, around 2.704, and that furthermore the limit is rational. - _Charles R Greathouse IV_, Feb 06 2017 If the first conjecture is true, then the prime numbers are distributed randomly. If the second is true, we conclude that the prime numbers are not so random. - _Dimitris Valianatos_, Feb 08 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimitris Valianatos_, Feb 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011584 | OEIS-A282121

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 430", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 07 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011585 | OEIS-A282122

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 430", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 07 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011586 | OEIS-A282123

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 430", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 07 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011587 | OEIS-A282124

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 430", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 07 2017: (Start)
Conjectures from _Paul Curtz_, Jun 10 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011588 | OEIS-A282135

Expansion of (Sum_{k = p*q, p prime, q prime} x^k)^3.

Conjecture: a(n) > 0 for n > 15.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Feb 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011589 | OEIS-A282142

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 553", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 27 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011590 | OEIS-A282161

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x and (12*x)^2 + (5*y-10*z)^2 both squares, where x,y,z are nonnegative integers and w is a positive integer.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 16^k*m (k = 0,1,2,... and m = 1, 23, 28, 79, 119, 191, 223, 263, 463, 703, 860, 1052). See also A281976, A281977, A282013 and A282014 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011591 | OEIS-A282166

a(n) is the minimal sum of a positive integer sequence of length n with no duplicate substrings of length greater than 1, and every number different from its neighbors.

For n>=4, we seem to have a(n) = a(n-1) + a(n-2) - a(n-3) + d(n), where d(n) is either 0 or 1 (with a clear formula). This observation leads to the conjecture: for n>=4, a(n) = -3/2 + 2*n + n*m/2 - m*(2*m^2+15*m+46)/24 + (-1)^n*(m%2+2)/4 + (m%2)*3/8, where m is the largest integer such that (2*m^2 + 8*m + 1 + 3*(-1)^m)/4 <= n. - _Max Alekseyev_, May 28 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Feb 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011592 | OEIS-A282172

Expansion of (Sum_{k>=0} x^(k*(k+1)*(k+2)/6))^5.

a(n) > 0 for all n ("Pollock's Conjecture").

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Feb 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011593 | OEIS-A282173

Expansion of (Sum_{k>=0} x^(k*(k+1)*(2*k+1)/6))^6.

Conjecture: a(n) > 0 for all n. Extended conjecture: every number is the sum of at most 6 square pyramidal numbers. Generalized conjecture: every number is the sum of at most k+2 k-gonal pyramidal numbers (except k = 5). - _Ilya Gutkovskiy_, Feb 10 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Feb 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011594 | OEIS-A282175

a(n) is the smallest product M=p_1*p_2*...*p_n with distinct primes p_i such that M+2^n=B, where B=q_1*q_2*...*q_n with distinct primes q_i, or a(n)=0 if there is no such M.

Conjecturally all a(n)>0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Feb 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011595 | OEIS-A282192

Number of ways of writing n as a sum of a prime and a squarefree semiprime.

Conjecture: a(n) > 0 for all n > 30.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Feb 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011596 | OEIS-A282226

Least nonnegative integer m which can be written in exactly n ways as x^2 + y^2 + z^2 + w^2 with both x and x + 24*y squares, where x,y,z,w are nonnegative integers with z <= w.

Conjecture: a(n) exists for any n > 0. See also A281976 for a related conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011597 | OEIS-A282231

First term of A175304 with a given prime signature.

Conjecturally the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Feb 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011598 | OEIS-A282251

Even integers not of the form p + prime(prime(q)) with distinct summands, where p and q are prime.

Conjecture 1: This sequence is finite and its largest term is smaller than 2*e^8. Conjecture 2: For any m > 0, all even numbers greater than 2*e^(4m) are the sum of a prime and an m-(prime-index-prime). See links. Conjecture 3: For any m >= 0 and n >= 0, all large enough even numbers are the sum of an m-(prime-index-prime) and an n-(prime-index-prime). See links.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrei-Lucian Dragoi_, Feb 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011599 | OEIS-A282261

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 446", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011600 | OEIS-A282262

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 446", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011601 | OEIS-A282263

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 446", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011602 | OEIS-A282264

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 446", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011603 | OEIS-A282289

Expansion of (Sum_{p prime, k>=2} x^(p^k))^4.

Conjecture: a(n) > 0 for all n > 27.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Feb 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011604 | OEIS-A282290

Expansion of (Sum_{p prime, i>=2} x^(p^i))*(Sum_{j>=2} mu(j)^2*x^j), where mu() is the Moebius function (A008683).

Conjecture: a(n) > 0 for all n > 8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Feb 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011605 | OEIS-A282291

Lexicographically earliest sequence of distinct terms such that every pair of consecutive terms contains a term that is a unitary divisor of the other term.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Feb 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011606 | OEIS-A282295

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 451", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = A279720(n) for n>=2. - _R. J. Mathar_, Jun 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011607 | OEIS-A282297

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 451", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = A279721(n) for n>=2. - _R. J. Mathar_, Jun 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011608 | OEIS-A282317

Lexicographically earliest infinite cubefree word over alphabet {0,1}.

However, at present it is only a conjecture that the terms from 1000 to 10000 are correct. - _N. J. A. Sloane_, May 22 2017 I reproduced these 10000 terms with an algorithm that did not need any backtracking in the strict sense (cf. PARI code), as follows: (1) Start with S := 0. (2) Let W := S. (3) While SW is not cubefree, "increment" W (see below for details) at the point where the first cube in SW ends. (4) Let S := SW and goto (2). By "increment W at the point k" we mean: if W[k]=1 then decrease k until W[k]=0; now set W[k]=1 and delete all further digits (i.e., W := concat(W[1..k-1],1)). If this never results in a maximal cubefree word S (cf. A282133), then the algorithm yields the correct sequence (minimal and cubefree by construction and infinite by hypothesis). - _M. F. Hasler_, May 20 2017 See A286940 for the lengths of the words W successively appended by this algorithm. These lengths L are sufficient to reconstruct the words W = concat(S[0..L-2],1) (except for the first two 1's which correspond to W = 0). There we give a pattern for that sequence of lengths conjectured to be valid for about 65*3^8 terms, corresponding to more than 10^7 terms of A282317. Beyond that point the behavior is unclear. - _M. F. Hasler_, May 21 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_, Feb 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011609 | OEIS-A282318

Number of ways of writing n as a sum of a prime and a nonprime squarefree number.

Conjecture: a(n) > 0 for all n > 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Feb 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011610 | OEIS-A282322

Greater of twin primes congruent to 13 (mod 30).

Conjecture: The sequence is equivalent to A000040 less primes of A135508 less primes of A356247. - _Bill McEachen_, Sep 25 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Martin Renner_, Feb 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011611 | OEIS-A282342

a(n) is the smallest prime number, with sum of digits equals n and a(n) is greater than previous nonzero terms, except if this is not possible in which case a(n)=0.

I conjecture that there are prime numbers for every n, if n is not divisible by 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimitris Valianatos_, Feb 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011612 | OEIS-A282349

Expansion of (Sum_{k>=0} x^(k*(2*k^2+1)/3))^7.

Pollock (1850) conjectured that every number is the sum of at most 7 octahedral numbers (a(n) > 0 for all n >= 0).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Feb 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011613 | OEIS-A282350

Expansion of (Sum_{k>=0} x^(k*(5*k^2-5*k+2)/2))^15.

Pollock conjectured that every number is the sum of at most 5 tetrahedral numbers and that every number is the sum of at most 7 octahedral numbers. Conjecture: a(n) > 0 for all n >= 0. Extended conjecture: every number is the sum of at most 15 icosahedral numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Feb 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011614 | OEIS-A282355

Expansion of (Sum_{i>=1} x^prime(prime(i)))*(Sum_{j = p*q, p prime, q prime} x^j).

Conjecture: a(n) > 0 for all n > 527 (addition: only 18 positive integers cannot be represented as a sum of a prime number with prime subscript and a semiprime).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Feb 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011615 | OEIS-A282411

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 467", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 15 2017: (Start) Equivalent conjecture: a(n) = A279118(n) for n>=2. - _R. J. Mathar_, Jun 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011616 | OEIS-A282412

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 467", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011617 | OEIS-A282413

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 467", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011618 | OEIS-A282414

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 467", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 15 2017: (Start) Equivalent conjecture: a(n) = A282413(n) for n>=2. - _R. J. Mathar_, Jun 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011619 | OEIS-A282415

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 469", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011620 | OEIS-A282416

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 469", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011621 | OEIS-A282417

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 469", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011622 | OEIS-A282418

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 469", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011623 | OEIS-A282426

Number of ways to write n as x^4 + 4*y^2 + z^2 + 3^k, where x,y,z are nonnegative integers and k is among 0,1,2,3,4.

Conjecture: (i) a(n) > 0 for all n > 0. See arXiv:1701.05868 for more such conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011624 | OEIS-A282451

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 475", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Apr 18 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011625 | OEIS-A282452

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 475", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Apr 18 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011626 | OEIS-A282453

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 475", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Apr 18 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011627 | OEIS-A282454

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 475", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Apr 18 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011628 | OEIS-A282459

Number of composite numbers of the form 2*n - 2^k + 1 (k > 0, 2^k < 2*n + 1).

It is conjectured that a(n) > 0 for all n > 52. See related conjecture and findings in A039669. Also see the graph of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Feb 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011629 | OEIS-A282463

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers, x == y (mod 2) and z <= w such that both x and x^2 + 62*x*y + y^2 are squares.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 1, 3, 43, 723, 1723, 3571, 3911 and 16^k*m (k = 0,1,2,... and m = 7, 23, 31, 71, 79, 143, 303, 1591).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011630 | OEIS-A282482

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 478", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011631 | OEIS-A282483

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 478", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011632 | OEIS-A282484

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 478", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011633 | OEIS-A282485

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 478", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 18 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011634 | OEIS-A282494

Number of ways to write n as x^4 + y^2 + z^2 + w^2 with y*(y+240*z) a positive square, where x,y,z,w are nonnegative integers.

Conjecture: a(n) > 0 for all n > 0. See also A282463 and A282495 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011635 | OEIS-A282495

Number of ways to write n as x^4 + y^2 + z^2 + w^2 with y^2 + 228*y*z + 60*z^2 a square, where x,y,z are nonnegative integers and w is a positive integer.

Conjecture: a(n) > 0 for all n > 0. See also A282463 and A282494 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011636 | OEIS-A282502

Expansion of 1/(1 - Sum_{k>=0} x^(3*k*(k+1)/2+1)).

Conjecture: every number > 1 is the sum of at most 5 centered triangular numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Feb 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011637 | OEIS-A282504

Expansion of 1/(1 - Sum_{k>=0} x^(2*k*(k+1)+1)).

Conjecture: every number > 1 is the sum of at most 6 centered square numbers. Extended conjecture: every number > 1 is the sum of at most k+2 centered k-gonal numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Feb 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011638 | OEIS-A282507

Triangular array read by rows. T(n,k) is the number of chain topologies on an n-set with exactly k open sets where one of the open sets is a single point set, n >= 2, 3 <= k <= n+1.

A conjecture I made to Loic Foissy, who replied: sequence T(n,k) counts surjective maps [n]->> [k] such that k is obtained exactly once, whereas sequence A019538 b(n,k) counts surjective maps [n]->> [k]. To construct an element for T(n,k), you may choose the element of [n] giving k (n choices), then a surjection from the n-1 remaining elements to [k-1] (b(n-1,k-1) choices). This gives T(n,k) = n * b(n-1,k-1), if k,n>1. - _Tom Copeland_, Nov 10 2023 [So it is now a theorem, not a conjecture, right? - _N. J. A. Sloane_, Dec 23 2023]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Critzer_, Feb 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011639 | OEIS-A282510

Irregular triangle T(n,k) read by rows: Each term is the least positive integer such that no row, column, diagonal, or antidiagonal contains a repeated term; and each row terminates at k when it contains all numbers <= k.

Conjecture: All columns and diagonals are permutations of the natural numbers; a proof will be more involved than for A274651.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bob Selcoe_, Feb 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011640 | OEIS-A282523

Number of n X 3 0..1 arrays with no 1 equal to more than two of its king-move neighbors.

Conjectured g.f.: -x*(8+3*x-25*x^2-11*x^3+13*x^4-2*x^5+4*x^6+6*x^7)/(-1+5*x+2*x^2-x^3-24*x^4+13*x^5-2*x^7+6*x^8). - _R. J. Mathar_, Feb 28 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. H. Hardin_, Feb 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011641 | OEIS-A282529

Decimal expansion of Product_{n>=2} (1-1/n!).

Conjectured to be irrational, transcendental and normal, none have been shown.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lucian Craciun_, Feb 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011642 | OEIS-A282535

a(n) is the maximum number of "describing"-steps for an n-chain before entering a loop.

Conjecture: The size of terms of this sequence is unbounded (cf. Marichal, 2007, Corollary 5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Feb 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011643 | OEIS-A282539

a(1)=1, a(n) is the smallest integer not included earlier such that a(n)*a(n-1)-1 and a(n)*a(n-1)+1 are twin primes.

It is only a conjecture that a(n) always exists. - Editors, OEIS, Mar 07 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Feb 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011644 | OEIS-A282542

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that x + 3*y + 5*z and (at least) one of y,z,w are squares.

Conjecture: a(n) > 0 for all n = 0,1,2,.... This is stronger than the 1-3-5 conjecture (cf. A271518).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011645 | OEIS-A282545

Number of ways to write n as x^4 + y^2 + z^2 + w^2 with y^2 + 64*z^2 + 1024*y*z a square, where x,y,z,w are nonnegative integers with y > 0.

Conjecture: (i) a(n) > 0 for all n > 0. See also A282463, A282494 and A282495 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011646 | OEIS-A282561

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that both 2*x - y and 4*z^2 + 724*z*w + w^2 are squares.

Conjecture: a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 16^k*m (k = 0,1,2,... and m = 7, 19, 67, 191, 235, 265, 347, 888, 2559). See also A282562 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011647 | OEIS-A282562

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that both 2*x - y and 9*z^2 + 666*z*w + w^2 are squares.

Conjecture: a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 16^k*m (k = 0,1,2,... and m = 12, 19, 23, 44, 60, 67, 139, 140, 248, 264, 347, 427, 499, 636, 1388, 1867, 1964, 4843). See also A282561 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011648 | OEIS-A282570

Number of ways to write n as an ordered sum of two multiplicatively perfect numbers (A007422).

Conjecture: a(n) > 0 for all n > 51.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Feb 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011649 | OEIS-A282577

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 553", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 27 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011650 | OEIS-A282579

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 553", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 27 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011651 | OEIS-A282585

Number of ways to write n as an ordered sum of 3 squarefree palindromes (A071251).

Conjecture: a(n) > 0 for any sufficiently large n. Additional conjecture: every number > 3 can be written as the sum of 4 squarefree palindromes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Feb 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011652 | OEIS-A282606

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 489", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011653 | OEIS-A282607

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 489", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011654 | OEIS-A282608

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 489", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011655 | OEIS-A282609

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 489", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011656 | OEIS-A282652

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 491", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011657 | OEIS-A282653

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 491", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011658 | OEIS-A282654

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 491", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011659 | OEIS-A282655

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 491", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011660 | OEIS-A282678

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 499", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = A282674(n) for n>=2. - _R. J. Mathar_, Jun 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011661 | OEIS-A282679

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 499", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = A282675(n) for n>=2. - _R. J. Mathar_, Jun 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011662 | OEIS-A282680

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 499", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = A282676(n) for n>=2. - _R. J. Mathar_, Jun 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011663 | OEIS-A282681

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 499", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = A282677(n) for n>=2. - _R. J. Mathar_, Jun 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011664 | OEIS-A282695

A055265(n) - n.

There is no proof at present for the conjecture that A055265 is a permutation of the positive integers. This sequence gives the remarkably small deviations from the identity sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011665 | OEIS-A282715

Number of nonzero entries in row n of the base-3 generalized Pascal triangle P_3.

Leroy et al. (2017) state some conjectured recurrences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011666 | OEIS-A282793

Indices k of nontrivial Riemann zeta zeros such that floor(Im(zetazero(k))/(2*Pi)*log(Im(zetazero(k))/(2*Pi*e)) + 7/8) - k + 1 = 1.

Conjecture 1: The union of this sequence and A282794 is A153815. Conjecture 2: The zeta zeros whose indices are terms of this sequence are the locations where the zeta zero counting function, (RiemannSiegelTheta(t) + Im(log(zeta(1/2 + i*t))))/Pi + 1 overcounts the number of zeta zeros on the critical line. Conjecture 3: This sequence consists of the numbers k such that sign(Im(zetazero(k)) - 2*Pi*e*exp(LambertW((k - 7/8)/e))) = 1. Verified for the first 100000 zeta zeros.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Feb 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011667 | OEIS-A282796

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 505", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011668 | OEIS-A282797

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 505", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011669 | OEIS-A282798

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 505", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011670 | OEIS-A282799

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 505", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011671 | OEIS-A282800

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 507", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011672 | OEIS-A282801

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 507", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011673 | OEIS-A282802

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 507", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011674 | OEIS-A282803

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 507", based on the 5-celled von Neumann neighborhood.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011675 | OEIS-A282814

Table read by antidiagonals (n > 0, k > 0): T(n, k) is the length of the loop in Conway's subprime Fibonacci sequence with starting values of n and k.

All terms are in A214897, which is conjectured to be finite. -_Peter Kagey_, Nov 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Feb 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011676 | OEIS-A282826

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 521", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011677 | OEIS-A282827

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 521", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011678 | OEIS-A282828

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 521", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011679 | OEIS-A282829

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 521", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011680 | OEIS-A282841

Lexicographically earliest sequence of distinct terms such that a(n)=length of first run of multiples of prime(n) in this sequence.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Feb 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011681 | OEIS-A282863

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that 72*x^3 + (y-z)^3 is an even square.

Conjecture: (i) a(n) > 0 for any nonnegative integer n. Also, each n = 0,1,2,... can be written as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that x^3 + ((y-z)/2)^3 is a square (or twice a square).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011682 | OEIS-A282871

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 617", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 06 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011683 | OEIS-A282906

Expansion of 1/(1 - Sum_{j>=1} x^(Sum_{i=1..j} prime(i))).

Conjecture: every number > 3 is the sum of at most 5 partial sums of primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Feb 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011684 | OEIS-A282911

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 526", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011685 | OEIS-A282912

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 526", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011686 | OEIS-A282913

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 526", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011687 | OEIS-A282914

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 526", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011688 | OEIS-A282915

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 529", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011689 | OEIS-A282916

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 529", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011690 | OEIS-A282917

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 529", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011691 | OEIS-A282918

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 529", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Feb 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011692 | OEIS-A282933

Number of ways to write n as x^4 + y^2 + z^2 + w^2 with x,y,z nonnegative integers and w a positive integer such that 8*y^2 - 8*y*z + 9*z^2 is a square.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 16^k*m (k = 0,1,2,... and m = 1, 8, 15, 23, 39, 47, 71, 93, 239, 287, 311, 319, 383, 391, 591, 632, 1663, 2639, 5591, 6236).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011693 | OEIS-A282939

Maximum number of straight lines required to draw the boundary of any polyomino with n squares.

The conjecture that a(n) = 2*n+2 for n (mod 4) == 1, otherwise a(n) = 2*n, is proven. See link. - _John Mason_, Jun 08 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Feb 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011694 | OEIS-A282947

Number of ways of writing n as a sum of a perfect power and a squarefree semiprime.

Conjecture: a(n) > 0 for all n > 108. Conjecture: a(n) > 1 for all n > 604, Conjecture: a(n) > 2 for all n > 1008, etc.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Feb 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011695 | OEIS-A282954

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 533", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011696 | OEIS-A282955

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 533", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011697 | OEIS-A282956

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 533", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011698 | OEIS-A282957

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 533", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011699 | OEIS-A282972

Number of ways to write n as 4*x^4 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that 79*y^2 - 220*y*z + 205*z^2 is a square.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,.... This is much stronger than Lagrange's four-square theorem, and we have verified parts (i) and (ii) of the conjecture for n up to 10^7 and 10^6 respectively. See also A282933 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011700 | OEIS-A282984

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 541", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011701 | OEIS-A282985

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 541", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011702 | OEIS-A282986

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 541", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011703 | OEIS-A282987

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 541", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Feb 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011704 | OEIS-A282999

a(n) = index of first term of A282841 belonging to n runs (see A282841 for precise definition).

This sequence is conjectured to be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Feb 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011705 | OEIS-A283050

Integers that are divisible by the square of their least prime factor.

Conjecture: 1 <= a(n+1) - a(n) <= 4. - _R. J. Cano_, Feb 27 2017 The conjecture is true since all multiples of 4 are in this sequence. - _Charles R Greathouse IV_, Feb 28 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Feb 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011706 | OEIS-A283051

Positive integers n such that none of the primes of the form k*2^n + 1 (with k odd) divide any Fermat number F(m) = 2^(2^m) + 1, m >= 0.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Feb 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011707 | OEIS-A283056

Size of the smallest polyomino that admits a hole of size n.

a(7)-a(49) are now known; every term a(1)-a(49) equals A027709(n) + 3, the additive shift conjectured by Dmitry Kamenetsky in 2017. Each value is established as minimal over holes lying within a bounded rectangular region of the square lattice; that the region is large enough to exclude a smaller polyomino arising from a larger, more elongated hole is supported by a perimeter (isoperimetric) argument but not formally proved in general, so global minimality is conjectured beyond the exhaustively-enumerated small terms. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Feb 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011708 | OEIS-A283107

Numbers k such that tau(4*(k - 1)) is prime.

Conjecturally, a supersequence of A281312. The conjecture is true: the formula at A281312 implies that the number of divisors of 4*A281312(n) - 4 is A000043(n+1). - _Charles R Greathouse IV_, Mar 01 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Feb 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011709 | OEIS-A283152

2-untouchable numbers.

Let sigma(n) denote the sum of divisors of n, and s(n) := sigma(n) - n. Untouchable numbers are those numbers that do not lie in the image of s(n), and they were studied extensively (see the references). In 2016, Pollack and Pomerance conjectured that the set of untouchable numbers has a natural asymptotic density.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anton Mosunov_, Mar 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011710 | OEIS-A283170

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x + 2*y and z + 2*w both squares, where x and w are nonnegative integers, and y and z are integers.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011711 | OEIS-A283196

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with 2*x + y and 2*x + z both squares, where x,y,z are integers with |y| <= |z|, and w is a positive integer.

Conjecture: (i) a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011712 | OEIS-A283204

Numbers of the form x^2 + y^2 with x and y integers such that x + 2*y is a square.

This sequence is interesting since part (i) of the conjecture in A283170 implies that each n = 0,1,2,... can be expressed as the sum of two terms of the current sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011713 | OEIS-A283205

Numbers of the form x^2 + y^2 with x and y integers such that x + 3*y is a square.

This sequence is interesting since part (ii) of the conjecture in A283170 implies that each n = 0,1,2,... can be expressed as the sum of two terms of the current sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011714 | OEIS-A283214

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 597", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 04 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011715 | OEIS-A283215

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 597", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 04 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011716 | OEIS-A283216

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 597", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 04 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011717 | OEIS-A283217

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 597", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 04 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011718 | OEIS-A283225

Primes prime(k) such that prime(k)^2 mod prime(k+2) is different from prime(k+2)^2 mod prime(k).

I conjecture that there are no other terms in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arnaud Vernier_, Mar 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011719 | OEIS-A283239

Number of ways to write n as x^2 + y^2 + z*(3*z-1)/2 with x,y,z integers such that x + 2*y is a square.

Conjecture: (i) a(n) > 0 for all n >= 0, and a(n) = 1 only for n = 0, 8, 43, 84, 133, 253, 399, 488, 523, 803, 7369.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011720 | OEIS-A283269

Number of ways to write n as x^2 + y^2 + z^2 with x,y,z integers such that x + 3*y + 5*z is a square.

Conjecture: (i) a(2^k*m) > 0 for any positive odd integers k and m. Also, a(4*n+1) = 0 only for n = 63. See also A283273 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011721 | OEIS-A283273

Number of ways to write n as x^2 + y^2 + z^2 with x,y integers and z a nonnegative integer such that x + 2*y + 3*z is a square or twice a square.

Conjecture: (i) For any nonnegative integer n not of the form 4^k*(8*m+7) (k,m = 0,1,2,...), we have a(n) > 0. The Gauss-Legendre theorem states that a nonnegative integer can be written as the sum of three squares if and only if it is not of the form 4^k*(8*m+7) with k and m nonnegative integers. Thus a(4^k*(8*m+7)) = 0 for all k,m = 0,1,2,..., and part (i) of our conjecture (verified for n up to 1.6*10^6) is stronger than the Gauss-Legendre theorem. See also A283269 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011722 | OEIS-A283297

The smallest cardinality of a difference-basis in the cyclic group of order n.

It is an open problem whether a(n) = (1+o(1))*sqrt(n). See the MathOverflow link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Taras Banakh_, Mar 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011723 | OEIS-A283299

Number of ways to write 2*n + 1 as x^2 + 2*y^2 + 3*z^2 with x,y,z integers such that x + y + z is a square or twice a square.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 3, 6, 8, 17, 23, 41, 128, 197, 372, 764, 1143, 1893, 3761, 4307, 6408, 6918. See also A283366 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011724 | OEIS-A283348

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 617", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 06 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011725 | OEIS-A283349

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 617", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 06 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011726 | OEIS-A283350

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 617", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 06 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011727 | OEIS-A283351

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 619", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 06 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011728 | OEIS-A283352

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 619", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 06 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011729 | OEIS-A283353

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 619", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 06 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011730 | OEIS-A283355

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 621", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 06 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011731 | OEIS-A283356

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 621", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 06 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011732 | OEIS-A283357

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 621", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 07 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011733 | OEIS-A283358

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 621", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 07 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011734 | OEIS-A283366

Number of ways to write 2*n + 1 as x^2 + y^2 + 2*z^2 with x,y,z integers such that 2*x + y + z is a square or a power of two.

Conjecture: a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 19, 32, 47, 115, 200, 974, 1271, 2240, 2549, 3185, 4865, 9254, 15881. The conjecture implies that any positive integer with even 2-adic order can be written as x^2 + y^2 + 2*z^2 with x,y,z integers such that 2*x + y + z is a square or twice a square. See also A283299 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011735 | OEIS-A283400

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 633", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011736 | OEIS-A283401

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 633", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011737 | OEIS-A283402

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 633", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011738 | OEIS-A283403

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 633", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011739 | OEIS-A283404

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 637", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011740 | OEIS-A283405

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 637", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011741 | OEIS-A283406

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 637", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011742 | OEIS-A283407

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 637", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011743 | OEIS-A283431

a(n) is the number of zeros of the Hermite H(n, x) polynomial in the open interval (-1, +1).

Conjecture: a(n) = A257564(n+2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, May 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011744 | OEIS-A283504

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 641", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 10 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011745 | OEIS-A283505

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 641", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 10 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011746 | OEIS-A283506

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 641", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 10 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011747 | OEIS-A283507

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 641", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 10 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011748 | OEIS-A283508

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 643", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 10 2017: (Start)
Equivalent conjecture: a(n) = A267623(n). - _R. J. Mathar_, Mar 17 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011749 | OEIS-A283523

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 913", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 26 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011750 | OEIS-A283581

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 646", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 12 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011751 | OEIS-A283582

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 646", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 12 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011752 | OEIS-A283583

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 646", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 12 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011753 | OEIS-A283584

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 646", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 12 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011754 | OEIS-A283585

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 654", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 12 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011755 | OEIS-A283586

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 654", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 12 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011756 | OEIS-A283587

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 654", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 12 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011757 | OEIS-A283588

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 654", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 12 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011758 | OEIS-A283589

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 657", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 13 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011759 | OEIS-A283590

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 657", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 13 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011760 | OEIS-A283591

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 657", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 13 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011761 | OEIS-A283592

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 657", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 13 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011762 | OEIS-A283604

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 670", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011763 | OEIS-A283605

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 670", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011764 | OEIS-A283606

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 670", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011765 | OEIS-A283607

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 670", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011766 | OEIS-A283617

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w integers and 0 <= z <= w such that x^3 + 2*y^3 is a square.

Conjecture: (i) Let a and b >= a be positive integers with gcd(a,b) squarefree. Then, every n = 0,1,2,... can be written as x^2 + y^2 + z^2 + w^2 with x,y,z,w integers such that a*x^3 + b*y^3 is a square, if and only if (a,b) is among the ordered pairs (1,2), (1,8), (2,16), (4,23), (4,31), (5,9), (8,9), (8,225), (9,47), (25,88), (50,54).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011767 | OEIS-A283619

a(n) = (conjectured) smallest positive integer k which is neither of the form p + n^x nor of the form p - n^x with x >= 0 and p prime, where gcd(k, n) = 1 and gcd(k^2-1, n-1) = 1.

a(n) = (conjectured) smallest positive integer k which is neither of the form p + n^x nor of the form p - n^x with x >= 0 and p prime, where gcd(k, n) = 1 and gcd(k^2-1, n-1) = 1. What does "conjectured" mean? A positive integer k is a candidate if:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Mar 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011768 | OEIS-A283641

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 678", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011769 | OEIS-A283644

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 694", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011770 | OEIS-A283645

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 694", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011771 | OEIS-A283646

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 694", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011772 | OEIS-A283647

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 694", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011773 | OEIS-A283648

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 705", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011774 | OEIS-A283649

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 705", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011775 | OEIS-A283650

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 705", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011776 | OEIS-A283651

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 705", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011777 | OEIS-A283680

Odd numbers of the form (p^2 - q^2) / 24 with primes p > q > 3.

a(n) ~ cn(loglog n)^(3/2)/(log n)^(1+eta), where eta = 1-(1+loglog 2)/log 2 = 0.086..., perhaps with c about 4/3. - Conjectured by Carl Pomerance, Mar 15 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_ and _Altug Alkan_, Mar 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011778 | OEIS-A283712

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 758", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 10 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011779 | OEIS-A283713

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 758", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 10 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011780 | OEIS-A283752

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 758", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 10 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011781 | OEIS-A283754

The smallest number k such that k*2^n mod 3^n = 1.

Conjecture: 2*a(n+1)-a(n) = 3^n * A055620(n). - _Robert Israel_, Mar 28 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joe Slater_, Mar 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011782 | OEIS-A283794

Positions of 1 in A288375; complement of A288625.

Conjecture: a(n)/n -> 2, and -1/4 < 2 - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011783 | OEIS-A283798

Triangle T(n,k) read by rows: the number of symmetric q-circulant n X n {0,1}-matrices where each column sum, each row sum and the trace equal k.

Conjecture: T(n,k) = binomial(n,k) for odd n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Mar 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011784 | OEIS-A283806

Odd numbers which are uniquely decomposable into the sum of a prime and a power of two.

It is conjectured that none of these numbers is in A101036.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Mar 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011785 | OEIS-A283814

Irregular triangle read by rows in which n-th row lists the numbers m such that 2*prime(m) can be represented as the sum of two primes in exactly n ways.

From b116619.txt it seems that the sequence is correct at least for first 677 terms (first 100 rows of triangle). But as it is usual in number theory better consider this sequence as conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Mar 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011786 | OEIS-A283849

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 734", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011787 | OEIS-A283850

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 734", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011788 | OEIS-A283864

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 734", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011789 | OEIS-A283865

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 734", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011790 | OEIS-A283882

Relative of Hofstadter Q-sequence: a(n) = max(0, n+67) for n <= 0; a(n) = a(n-a(n-1)) + a(n-a(n-2)) for n > 0.

If the index is at least 19047817371, then a(5n) = 5*A272611(n-3809563474), a(5n+1) = 5*A272611(n-3809563473), a(5n+2) = 5*A272612(n-3809563473), a(5n+3) = 19047817435*A272613(n-3809563473), a(5n+4) = 4. This pattern lasts as long as A272611 exists (which is conjectured to be forever).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nathan Fox_, Mar 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011791 | OEIS-A283905

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 758", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Dec 10 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011792 | OEIS-A283929

Number of ways of writing n as a sum of a twin prime (A001097) and a squarefree semiprime (A006881).

Conjecture: a(n) > 0 for all n > 30.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Mar 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011793 | OEIS-A283963

Fixed point of the morphism 0 -> 1, 1 -> 1010.

Let u(n) = # 0's <= n and v(n) = # 1's <= n. Let r = (3+sqrt(3))/2 and s = sqrt(3), so that 1/r + 1/s = 1. Conjecture: -1 < n*r - u(n) < 2 and -1 < n*s - v(n) < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011794 | OEIS-A283964

Positions of 0 in A283963; complement of A283965.

Conjecture: -1 < n*r - a(n) < 2 for n >= 1, where r = (3+sqrt(3))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011795 | OEIS-A283965

Positions of 1 in A283963; complement of A283964.

Conjecture: -1 < n*sqrt(3) - a(n) < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011796 | OEIS-A283966

Fixed point of the morphism 0 -> 1, 1 -> 10101.

Let u(n) = # 0's <= n and v(n) = # 1's <= n. Let r = (7+sqrt(17))/4 and s = (-1+sqrt(17))/2, so that 1/r + 1/s = 1. Conjecture: -1 < n*r - u(n) < 3 and -1 < n*s - v(n) < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011797 | OEIS-A283967

Positions of 0 in A283966; complement of A284015.

Conjecture: -1 < n*r - a(n) < 3 for n >= 1, where r = (7+sqrt(17))/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011798 | OEIS-A283991

a(1) = 0 and for n > 1, a(n) = 1 if Stern polynomial B(n,x) is irreducible, 0 otherwise.

a(n) = A010051(n) + A305892(n). (Conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011799 | OEIS-A284004

a(n) = A046523(A284003(n)).

a(n) = A002110(A001222(A284003(n))) = A002110(A209281(n+1)). [Latter so far only conjectured.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011800 | OEIS-A284015

Positions of 1 in A283966; complement of A283967.

Conjecture: -1 < n*s - a(n) < 2 for n >= 1, where s = (-1+sqrt(17))/2..

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011801 | OEIS-A284024

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 782", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011802 | OEIS-A284025

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 782", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011803 | OEIS-A284026

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 782", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011804 | OEIS-A284027

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 782", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011805 | OEIS-A284028

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 790", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = A283906(n+1). - _R. J. Mathar_, Jun 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011806 | OEIS-A284038

Lexicographically earliest sequence of distinct positive terms such that A000196(n) divides a(n).

Conjecturally:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, May 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011807 | OEIS-A284047

a(n) is the smallest positive integer not already in the sequence such that a(n) + a(n-1) and a(n)^2 + a(n-1)^2 are primes, with a(1) = 1.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Mar 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011808 | OEIS-A284049

a(n) is the smallest positive integer not already in the sequence such that a(n) + a(n-1) is a prime power, with a(1) = 1.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Mar 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011809 | OEIS-A284116

a(n) = largest number of distinct words arising in Post's tag system {00, 1101} applied to a binary word w, over all starting words w of length n, or a(n) = -1 if there is a word w with an unbounded trajectory.

It is an important open question to decide if there is any word whose orbit grows without limit. - _N. J. A. Sloane_, Jul 30 2017, based on an email from _Allan C. Wechsler_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Mar 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011810 | OEIS-A284125

Number of binary strings w of length n for which s, the longest proper suffix of w that appears at least twice in w, is of length 2.

G.f.: 2*x^3*(1 + x^2)*(1 - 4*x + 6*x^2 - 5*x^3 + x^4 + 2*x^5 + 2*x^6 - 2*x^7) / ((1 - x)^2*(1 - x - x^2)*(1 - 2*x + x^2 - x^3)*(1 - x - x^2 - x^3)) (conjectured). - _Colin Barker_, Jun 07 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Mar 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011811 | OEIS-A284145

Triangle read by rows T(n,k) in which each term is the least positive integer not yet appearing in the triangle that is coprime to all the terms in its associated row, column, diagonal and antidiagonal.

Conjecture 1: The triangle is a permutation of the natural numbers. Conjecture 2: Each F(k) and G(n) is a permutation of the prime numbers (except F(1) and G(1), which obviously also contain 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bob Selcoe_, Mar 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011812 | OEIS-A284147

3-untouchable numbers.

Let sigma(n) denote the sum of divisors of n, and s(n) := sigma(n) - n, with the convention that s(0)=0. Untouchable numbers are those numbers that do not lie in the image of s(n), and they were studied extensively (see the references). In 2016, Pollack and Pomerance conjectured that the set of untouchable numbers has a natural asymptotic density.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anton Mosunov_, Mar 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011813 | OEIS-A284170

Array read by antidiagonals: T(i,j) is the largest prime in the sequence defined by a(1) = prime(i), a(2) = prime(j), a(n) = A006530(a(n-1)+a(n-2)+1) for n>=3, or 0 if that sequence contains arbitrarily large primes.

Conjecture: the sequence always eventually repeats, so T(i,j) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Mar 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011814 | OEIS-A284175

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 809", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011815 | OEIS-A284176

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 809", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011816 | OEIS-A284177

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 809", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011817 | OEIS-A284178

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 809", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011818 | OEIS-A284179

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 813", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011819 | OEIS-A284180

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 813", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011820 | OEIS-A284181

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 813", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011821 | OEIS-A284182

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 813", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011822 | OEIS-A284189

Square array T(n,k) read by upward antidiagonals: each term is the least positive integer not yet appearing in the array that is coprime to all the terms in its associated row, column, diagonal and antidiagonal.

Conjecture 1: The array is a permutation of the natural numbers. Conjecture 2: The prime factors of all the terms in each individual row, column and diagonal are permutations of the prime numbers (except the middle diagonal and the first row and column, which obviously also contain 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bob Selcoe_, Mar 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011823 | OEIS-A284230

Number of self-avoiding planar walks starting at (0,0), ending at (n,0), remaining in the first quadrant and using steps (0,1), (1,0), (1,1), (-1,1), and (1,-1) with the restriction that (0,1) is never used below the diagonal and (1,0) is never used above the diagonal.

Conjecture: a(n) -a(n-1) +(-n^2-n+3)*a(n-2) +(-n+2)*a(n-3) +(n-2)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Apr 09 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Mar 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011824 | OEIS-A284235

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 833", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011825 | OEIS-A284236

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 833", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011826 | OEIS-A284237

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 833", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011827 | OEIS-A284238

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 833", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011828 | OEIS-A284243

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 841", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011829 | OEIS-A284244

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 841", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011830 | OEIS-A284245

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 841", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011831 | OEIS-A284246

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 841", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011832 | OEIS-A284278

a(1)=2; for n >= 1, if n+2 is prime then a(2*n+1) = 3*n + 2 and a(2*n) = n + 2, otherwise all terms are 2.

The sequence is motivated by A284172, by the message from B. Jubin dated Mar 23 2017 and by the classic open problem of showing that there are infinitely many primes p for which 2*p-1 is also prime. If there were only finitely many such primes, then there would be a place where this sequence is generated by the same rule as A284172.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Mar 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011833 | OEIS-A284299

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 865", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011834 | OEIS-A284300

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 865", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011835 | OEIS-A284301

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 865", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011836 | OEIS-A284302

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 865", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011837 | OEIS-A284343

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers and y <= z such that 2*x + y - z is either zero or a power of 8 (including 8^0 = 1).

Conjecture: (i) For any c = 1,2,4, each n = 0,1,2,... can be written as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers and y <= z such that c*(2*x+y-z) is either zero or a power of eight (including 8^0 = 1). Part (i) of the conjecture is stronger than the first part of Conjecture 4.4 in the linked JNT paper (see also A273432).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011838 | OEIS-A284347

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 873", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011839 | OEIS-A284348

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 873", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011840 | OEIS-A284349

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 873", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011841 | OEIS-A284350

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 873", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011842 | OEIS-A284351

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 899", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 26 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011843 | OEIS-A284352

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 899", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 26 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011844 | OEIS-A284353

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 899", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 26 2017: (Start)
Conjecture: For n > 0, a(n) = Sum_{k=0..n-1} C(n,k) * (2-(-1)^k). - _Wesley Ivan Hurt_, Sep 23 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011845 | OEIS-A284354

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 899", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 26 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011846 | OEIS-A284364

Fixed point of the morphism 0->1, 1->101010.

Let u(n) = # 0's <= n and v(n) = # 1's <= n. Let r = (9+sqrt(21))/6 and s = (-1+sqrt(21))/2, so that 1/r + 1/s = 1. Conjecture: -2 < n*r - u(n) < 2 and -1 < n*s - v(n) < 2 for n >= 1. Conjecture: The asymptotic mean of this sequence is lim_{m->oo} (1/m) * Sum_{k=1..m} a(k) = (sqrt(21) + 1)/10 = 0.558257... . - _Amiram Eldar_, Apr 13 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011847 | OEIS-A284365

Positions of 0 in A284364; complement of A284366.

Conjecture: -2 < n*r - a(n) < 2 for n >= 1, where r = (9+sqrt(21))/6. Conjecture: The asymptotic density of this sequence is (9-sqrt(21))/10 = 0.441742... . - _Amiram Eldar_, Apr 13 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011848 | OEIS-A284366

Positions of 1 in A284364; complement of A284365.

Conjecture: -1 < n*s - a(n) < 2 for n >= 1, where s = (-1+sqrt(21))/2. Conjecture: The asymptotic density of this sequence is (sqrt(21) + 1)/10 = 0.558257... . - _Amiram Eldar_, Apr 13 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011849 | OEIS-A284368

Fixed point of the morphism 0->1, 1->1011.

Let u(n) = # 0's <= n and v(n) = # 1's <= n. Let r = (5+sqrt(13))/2 and s = (-1+sqrt(13))/2, so that 1/r + 1/s = 1. Conjecture: 1 < n*r - u(n) < 3 and -1 < n*s - v(n) < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011850 | OEIS-A284369

Fixed point of the morphism 0->1, 1->1001.

Let u(n) = # 0's <= n and v(n) = # 1's <= n. Let r = (3+sqrt(3))/2 and s = sqrt(3), so that 1/r + 1/s = 1. Conjecture: -3 < n*r - u(n) < 4 and -2 < n*s - v(n) < 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011851 | OEIS-A284370

Positions of 0 in A284369; complement of A284371.

Conjecture: -3 < n*r - a(n) < 4 for n >= 1, where r = (3+sqrt(3))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011852 | OEIS-A284371

Positions of 1 in A284369; complement of A284370.

Conjecture: -2 < n*sqrt(3) < 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011853 | OEIS-A284386

Fixed point of the morphism 0->1, 1->1101.

Let u(n) = # 0's <= n and v(n) = # 1's <= n. Let r = (5+sqrt(3))/2 and s = (-1+sqrt(13))/2, so that 1/r + 1/s = 1. Conjecture: 0 < n*r - u(n) < 2 and -1 < n*s - v(n) < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011854 | OEIS-A284388

0-limiting word of the morphism 0 -> 1, 1 -> 001.

Conjecture: the limiting frequency of 0's in both limiting words is 1/2. Conjecture: after the first term, same as one less than the run-lengths of the Thue-Morse sequence A010060. - _George Beck_, Mar 24 2021 Conjecture: Excluding the first two terms, if runs of 1s of length one are replaced by 0, and runs of 1s of length three are replaced by 1, we get the same sequence. Conjecture: Index distance between two consecutive 1s is either one or three. Excluding the first two terms, if distances of one are replaced by 0, and distances of three are replaced by 1, we get the same sequence. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011855 | OEIS-A284391

1-limiting word of the morphism 0 -> 1, 1 -> 001.

Conjecture: the limiting frequency of 0's in both limiting words is 1/2. Conjecture: Excluding the first term, if runs of 1s of length one are replaced by 0, and runs of 1s of length three are replaced by 1, we get the same sequence. Conjecture: Index distance between two consecutive 1s is either one or three. Excluding the first term, if distances of one are replaced by 0, and distances of three are replaced by 1, we get the same sequence. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011856 | OEIS-A284403

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 913", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 26 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011857 | OEIS-A284404

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 913", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 26 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011858 | OEIS-A284405

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 913", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 26 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011859 | OEIS-A284469

Positions of 0 in A284468; complement of A284470.

Conjecture: a(n)/n -> (7 + sqrt(13))/6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011860 | OEIS-A284470

Positions of 1 in A284468; complement of A284469.

Conjecture: a(n)/n -> (1 + sqrt(13))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011861 | OEIS-A284472

Positions of 0 in A284471; complement of A284473.

Conjecture: a(n)/n -> (7 + sqrt(13))/6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011862 | OEIS-A284473

Positions of 1 in A284471; complement of A284472.

Conjecture: a(n)/n -> (1 + sqrt(13))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011863 | OEIS-A284480

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 950", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 28 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011864 | OEIS-A284481

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 950", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 28 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011865 | OEIS-A284482

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 950", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 28 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011866 | OEIS-A284483

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 961", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 28 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011867 | OEIS-A284484

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 961", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 28 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011868 | OEIS-A284485

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 961", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 28 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011869 | OEIS-A284486

The lexicographically earliest table read by antidiagonals such that any value appears at most twice on any straight line (of any slope).

Conjecture: Every value appears twice on every line of rational slope that contains an infinite number of points.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Mar 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011870 | OEIS-A284488

Positions of 0 in A284487; complement of A284489.

Conjecture: a(n)/n -> (7 + sqrt(13))/6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011871 | OEIS-A284489

Positions of 1 in A284487; complement of A284488.

Conjecture: a(n)/n -> (1 + sqrt(13))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011872 | OEIS-A284491

Positions of 0 in A284490; complement of A284492.

Conjecture: a(n)/n -> (7 + sqrt(13))/6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011873 | OEIS-A284492

Positions of 1 in A284490; complement of A284491.

Conjecture: a(n)/n -> (1 + sqrt(13))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011874 | OEIS-A284505

Fixed point of the morphism 0->1, 1->1100.

Let u(n) = # 0's <= n and v(n) = # 1's <= n. Let r = (3+sqrt(3))/2 and s = sqrt(3), so that 1/r + 1/s = 1. Conjecture: -3 < n*r - u(n) < 4 and -2 < n*s - v(n) < 4 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011875 | OEIS-A284506

Positions of 0 in A284505; complement of A284507.

Conjecture: -3 < n*r - a(n) < 4 for n >= 1, where r = (3+sqrt(3))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011876 | OEIS-A284507

Positions of 1 in A284505; complement of A284506.

Conjecture: -2 < n*sqrt(3) - a(n) < 4 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011877 | OEIS-A284509

Positions of 0 in A284508; complement of A284510.

Conjecture: a(n)/n -> (7 + sqrt(13))/6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011878 | OEIS-A284510

Positions of 1 in A284508; complement of A284509.

Conjecture: a(n)/n -> (1 + sqrt(13))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011879 | OEIS-A284513

Positions of 0 in A284512; complement of A284514.

Conjecture: a(n)/n -> (7 + sqrt(13))/6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011880 | OEIS-A284514

Positions of 1 in A284512; complement of A284513.

Conjecture: a(n)/n -> (1 + sqrt(13))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011881 | OEIS-A284516

Lexicographically earliest sequence of unique numbers such that for each digit "d" the gap to the nearest other digit "d" is equal to d.

It can be conjectured that any number which is not an intrinsically impossible term will eventually appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Lars Blomberg_, Mar 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011882 | OEIS-A284525

Positions of 0 in A284524; complement of A284526.

Conjecture: a(n)/n -> (7 + sqrt(13))/6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011883 | OEIS-A284526

Positions of 1 in A284524; complement of A284525.

Conjecture: a(n)/n -> (1 + sqrt(13))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011884 | OEIS-A284528

Positions of 0 in A284527; complement of A284529.

Conjecture: a(n)/n -> (7 + sqrt(13))/6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011885 | OEIS-A284529

Positions of 1 in A284527; complement of A284528.

Conjecture: a(n)/n -> (1 + sqrt(13))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011886 | OEIS-A284534

Positions of 0 in A284533; complement of A284535.

Conjecture: a(n)/n -> (3 + sqrt(3))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011887 | OEIS-A284535

Positions of 1 in A284533; complement of A284534.

Conjecture: a(n)/n -> sqrt(3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011888 | OEIS-A284540

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 974", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 29 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011889 | OEIS-A284541

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 974", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 29 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011890 | OEIS-A284542

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 974", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 29 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011891 | OEIS-A284543

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 974", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Mar 29 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011892 | OEIS-A284544

Binary representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 998", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011893 | OEIS-A284545

Binary representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 998", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011894 | OEIS-A284546

Decimal representation of the x-axis, from the left edge to the origin, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 998", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011895 | OEIS-A284547

Decimal representation of the x-axis, from the origin to the right edge, of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 998", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, May 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Mar 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011896 | OEIS-A284589

Positions of 0 in A284588; complement of A284590.

Conjecture: a(n)/n -> (3 + sqrt(3))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011897 | OEIS-A284590

Positions of 1 in A284588; complement of A284589.

Conjecture: a(n)/n -> sqrt(3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011898 | OEIS-A284594

Numbers whose square has a prime number of partitions.

Because asymptotically A072213(n) = A000041(n^2) ~ exp(Pi*sqrt(2/3)*n) / (4*sqrt(3)*n^2), the sum of the prime probabilities ~ 1/log(A072213(n)) is diverging and there are no obvious restrictions on primality; therefore, this sequence may be conjectured to be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Serge Batalov_, Mar 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011899 | OEIS-A284620

{00->2}-transform of the infinite Fibonacci word A003849.

We next prove the first conjecture by Kimberling. We easily see from the form of mu that the letters B and D occur, and only occur, at even positions in the fixed point x of mu. Since lambda(B)=lambda(D)=1, and A and C are not mapped to 1, it follows immediately that the positions of 1 in (a(n)) are given by A005843 = (2n). For a proof of Kimberling's second conjecture, note that a(n)=2 iff x(n)=C, where x is the fixed point of mu. The return words of C in x are CD and CDAB. Coding these two return words by their lengths, mu induces a descendant morphism tau given by This proves Kimberling's conjecture, provided we give the sequence A130568 offset 1, as we should.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011900 | OEIS-A284623

Positions of 0 in A284622; complement of A284626.

a(n) - a(n-1) is in {1,2,3} for n>=2. Conjecture: a(n)/n -> 9/5. This conjecture follows directly from the corresponding result for the positions of 1, proved in A284626. - _Michel Dekking_, Mar 31 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011901 | OEIS-A284624

Positions of 1 in A284749.

Conjectured: a(n) = 1 + A214971(n). Proof of this conjecture: it is easy to see that the infinite Fibonacci word A003849 is a concatenation of the words 01 and 001. So if we replace all 001 by 2, only the 01 remain, i.e., every 0 is directly followed by 1. - _Michel Dekking_, Aug 19 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011902 | OEIS-A284626

Positions of 1 in A284622; complement of A284623.

Conjecture: a(n)/n -> 9/4. Proof of the conjecture. Let x = A010060 be the Thue-Morse sequence, and let y = A284622 be the [0011->0]-transform of x. Let a = A284626 be the positions of 1 in y. There are 3 steps in the proof.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011903 | OEIS-A284641

Expansion of (Sum_{k>=0} x^(k^2*(k+1)^2/4))^12.

Conjecture: a(n) > 0 for all n. Extended conjecture: every number is the sum of at most 12 squares of triangular numbers (or partial sums of cubes).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, May 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011904 | OEIS-A284654

Positions of 0 in A284653; complement of A284655.

Conjecture: -1 < n*r - a(n) < 3 for n >= 1, where r = (3+sqrt(3))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011905 | OEIS-A284655

Positions of 1 in A284653; complement of A284654.

Conjecture: -1 < n*sqrt(3) - a(n) < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011906 | OEIS-A284657

Positions of 0 in A284656; complement of A284658.

Conjecture: -1 < n*r - a(n) < 3 for n >= 1, where r = (3+sqrt(3))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011907 | OEIS-A284658

Positions of 1 in A284656; complement of A284657.

Conjecture: -1 < n*sqrt(3) - a(n) < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011908 | OEIS-A284670

Exponents of powers of 5 that contain all ten decimal digits.

It is conjectured that a(n) = n + 40 for n > 25.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Apr 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011909 | OEIS-A284672

Exponents of powers of 7 that contain all ten decimal digits.

It is conjectured that a(n) = n + 34 for n > 27. G.f.: x*(-x^28 + x^27 - x^7 + x^6 - x^4 - 5*x^3 - 2*x^2 - 9*x + 18)/(x - 1)^2 (as conjectured in comments). - _Chai Wah Wu_, Jan 26 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Apr 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011910 | OEIS-A284675

Positions of 0 in A284674; complement of A284676.

Conjecture: 2 < n*r - a(n) < 4 for n >= 1, where r = (5+sqrt(13))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011911 | OEIS-A284676

Positions of 1 in A284674; complement of A284675.

Conjecture: -1 < n*r - a(n) < 4 for n >= 1, where r = (-1+sqrt(13))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011912 | OEIS-A284677

1-limiting word of the morphism 0->1, 1-> 0111.

Conjecture: a(n) = A284674(n-1). - _R. J. Mathar_, Jun 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011913 | OEIS-A284678

Positions of 0 in A284677; complement of A284679.

Conjecture: 2 < n*r - a(n) < 4 for n >= 1, where r = (5+sqrt(13))/2. If the conjectures here and in A284675 are true, then a(n) = A284675(n) for n>1. - _R. J. Mathar_, May 18 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011914 | OEIS-A284679

Positions of 1 in A284677; complement of A284678.

Conjecture: -1 < n*r - a(n) < 1 for n >= 1, where r = (-1+sqrt(13))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011915 | OEIS-A284681

Positions of 0 in A284680; complement of A284682.

Conjecture: a(n)/n -> (7 + sqrt(13))/6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011916 | OEIS-A284682

Positions of 1 in A284680; complement of A284681.

Conjecture: a(n)/n -> (1 + sqrt(13))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011917 | OEIS-A284684

Positions of 0 in A284627; complement of A284685.

a(n) - a(n-1) is in {1,2,3,4} for n>=2. Conjecture: a(n)/n -> 9/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011918 | OEIS-A284685

Positions of 1 in A284627; complement of A284684.

a(n) - a(n-1) is in {1,2,3} for n>=2. Conjecture: a(n)/n -> 9/5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011919 | OEIS-A284691

Numbers of the form (10^c-1)*the product any two (not necessarily distinct) terms of A074992.

Conjecture 1: all terms are palindromic in base 10. Conjecture 2: the sequence A074992 is the maximally dense sequence with this palindromic products property.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ahmad J. Masad_, Apr 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011920 | OEIS-A284708

Smallest initial prime p for at least n primes in increasing arithmetic progression with a common difference less than p.

Conjecture: a(n) > A034386(n) for every n >= 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Jan 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011921 | OEIS-A284746

Positions of 0 in A284745; complement of A191263.

Conjecture: a(n)/n -> (5 + sqrt(13))/6. From _Andrey Zabolotskiy_, Apr 13 2017: The conjecture is true since it states that the fraction of 0's is equal to 6/(5 + sqrt(13)), which is the invariant value of the fraction of 0's under the morphism.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011922 | OEIS-A284752

Positions of 0 in A284751; complement of A284753.

Conjecture: -1 < n*r - a(n) < 3 for n >= 1, where r = (3+sqrt(3))/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011923 | OEIS-A284753

Positions of 1 in A284751; complement of A284752.

Conjecture: -2 < n*r - a(n) < 3 for n >= 1, where r = 1 + sqrt(3). Interestingly, the conjecture: -2 < n*(1+sqrt(3))-a(n) < 3 , implies a conjecture: -1 < n*(1 + sqrt(3))/2 - A026363(n) < 3/2, which is stronger than the conjecture -1 < n*(1 + sqrt(3))/2 - a(n) < 2 given in the comments of A026363. See Fokkink-Joshi p. 16, Theorem 19, for proof of Kimberling's conjecture, listed first in these comments. - _Michael De Vlieger_, Feb 02 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011924 | OEIS-A284759

a(n) = (Sum_{i=1..n-1} i^(n-2)) mod n^3.

Conjecture: For n > 1, a(n) = 0 if and only if n is a term of A088164, i.e., n is a Wolstenholme prime (cf. Mestrovic, 2012, Conjecture 2.10).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Apr 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011925 | OEIS-A284760

a(n) = Sum_{i=1..n-1}(i^(n-2)) mod n^4.

Mestrovic conjectures that a(n) > 0 for all n > 1 (Conjecture 2.11).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Apr 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011926 | OEIS-A284773

Positions of 0 in A284772; complement of A284774.

Conjecture: -1 < n*r - a(n) < 2 for n >= 1, where r = (3+sqrt(3))/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011927 | OEIS-A284774

Positions of 1 in A284772; complement of A284773.

Conjecture: -1 < n*r - a(n) < 2 for n >= 1, where r = 1 + sqrt(3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011928 | OEIS-A284778

Number of self-avoiding planar walks of length n+1 starting at (0,0), ending at (n,0), remaining in the first quadrant and using steps (0,1), (1,0), (1,1), (-1,1), and (1,-1) with the restriction that (0,1) is never used below the diagonal and (1,0) is never used above the diagonal.

Conjecture: Also the number of topologically series-reduced ordered rooted trees with n + 3 vertices and more than one branch of the root. This would imply a(n) = A187306(n+1) - A005043(n+1). For example, the a(1) = 1 through a(5) = 22 trees are:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Apr 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011929 | OEIS-A284790

Positions of 0 in A284789; complement of A284791.

a(n) - a(n-1) is in {1,2,3} for n>=2. Conjecture: a(n)/n -> 3/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011930 | OEIS-A284791

Positions of 1 in A284789; complement of A284790.

a(n) - a(n-1) is in {1,2,4,5} for n>=2. Conjecture: a(n)/n -> 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011931 | OEIS-A284795

Positions of 0's in A284793.

This sequence and A284795 and A284796 form a partition of the positive integers. Conjecture: for n>=1, we have a(n)-3n+3 in {0,1}, 3*n+2-A284795(n) in {0,1,2,3}, and 3*n-2-A284795(n) in {0,1}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011932 | OEIS-A284796

Positions of 1's in A284793.

This sequence and A284794 and A284795 form a partition of the positive integers. Conjecture: for n>=1, we have a(n)-3n+3 in {0,1}, 3n+2-A284795(n) in {0,1,2,3}, and 3n-2-A284795(n) in {0,1}. Here is a proof of Kimberling's conjecture, i.e., the sequence y defined by y(n) := a(n)-3n+3 takes only values in the alphabet {0,1}. We know that A284793 = 1,-1,0,1,0,-1,... is a morphic sequence(see A284793). Let tau on the alphabet {A,B,C,D} be given by

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011933 | OEIS-A284814

Least number k such that k mod (2, 3, 4, ..., n+1) = (d_1, d_2, ..., d_n), where d_1, d_2, ..., d_n are the digits of n, with MSD(n) = d_1 and LSD(n) = d_n. 0 if such a number does not exist.

Conjecture: a(n) = 0 for all n >= 15. - _Max Alekseyev_, Jul 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_, Apr 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011934 | OEIS-A284815

Least number k such that k mod (2, 3, 4, ... , n+1) = (d_n, d_n-1, ..., d_1), where d_1 , d_2, ..., d_n are the digits of k, with MSD(k) = d_1 and LSD(k) = d_n. 0 if such a number does not exist.

Conjecture: a(n) = 0 for all n >= 15. - _Max Alekseyev_, Nov 10 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_, Apr 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011935 | OEIS-A284820

Positions of 0 in the {1100->0}-transform of A010060; complement of A284847.

a(n) - a(n-1) is in {1,2,3} for n>=2. Conjecture: a(n)/n -> 9/5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011936 | OEIS-A284844

Number of permutations on [n+3] with no circular 3-successions.

Conjecture: a(n) = (n+3)*A055790(n+1). - _R. J. Mathar_, Jul 15 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enrique Navarrete_, Apr 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011937 | OEIS-A284845

Number of permutations on [n+4] with no circular 4-successions.

Conjecture: a(n) = (n+4)*A277609(n+3). - _R. J. Mathar_, Jul 15 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enrique Navarrete_, Apr 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011938 | OEIS-A284847

Positions of 1 in the {1100->0}-transform of A010060; complement of A284820.

a(n) - a(n-1) is in {1,2,3,4} for n>=2. Conjecture: a(n)/n -> 9/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011939 | OEIS-A284849

Number of zeros strictly inside the unit circle of the Bernoulli polynomial B(n,x).

Conjecture: for n > 63, a(n) = 3 for n == 1 (mod 4) and a(n) = 4 otherwise. - _Charles R Greathouse IV_, May 09 2017 [Corrected by _Alper Ferudun_, Oct 18 2026] Conjectures from _Colin Barker_, Jan 22 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, May 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011940 | OEIS-A284852

Positions of 0 in A284851; complement of A284853.

Conjecture: -2 < n*r - a(n) < 2 for n >= 1, where r = (3+sqrt(3))/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011941 | OEIS-A284853

Positions of 1 in A284851; complement of A284852.

Conjecture: -2 < n*r - a(n) < 4 for n >= 1, where r = 1+sqrt(3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011942 | OEIS-A284884

Positions of 1's in A284881.

This sequence and A284882 and A284883 form a partition of the positive integers. Conjecture: for n>=1, we have a(n)-3n-3 in {0,1,2}, 3*n+1-A284883(n) in {0,1,2,3}, and 3*n-1-A284884(n) in {0,1,2}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011943 | OEIS-A284894

Positions of 0 in A284893; complement of A284895.

Conjecture: 2 < n*r - a(n) < 6 for n >= 1, where r = 2 + sqrt(2). Since the positions of 0 in x are partial sums of the terms of d, one can prove, using the Perron Frobenius theorem, that a weak form of the conjecture above holds: the sequence (n*r - a(n)) is bounded. This result can also be derived from the simple part (the second eigenvalue 2-sqrt(2) of the incidence matrix of the morphism tau: 2->24, 4->2444 is smaller than 1) of Theorem 1 in the paper "Symbolic discrepancy and self-similar dynamics" by Boris Adamczewski. - _Michel Dekking_, Jan 16 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011944 | OEIS-A284895

Positions of 1 in A284893; complement of A284894.

Conjecture: 0 < a(n) - n*sqrt(2) < 1 for n >= 1. The conjecture is false, since a(2) - 2*sqrt(2) = 4-2.828... > 1.17. Presumably the new conjecture is 0 < a(n) - n*sqrt(2) < 2 for n >= 1. - _Michel Dekking_, Jan 16 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011945 | OEIS-A284902

Positions of 0 in A284901; complement of A284903.

Conjecture: -3 < n*r - a(n) < 3 for n >= 1, where r = (3+sqrt(3))/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011946 | OEIS-A284903

Positions of 1 in A284901; complement of A284902.

Conjecture: -4 < n*r - a(n) < 6 for n >= 1, where r = 1 + sqrt(3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011947 | OEIS-A284906

Positions of 0 in A284905; complement of A284907.

See the comments of A285250 for a proof of this conjecture. - _Michel Dekking_, Oct 07 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011948 | OEIS-A284917

Lexicographically earliest infinite sequence of positive integers such that the Taxicab distance is unique to each pair of distinct points ((n, a(n)), (k, a(k))).

Conjecture: This sequence is strictly increasing for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Apr 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011949 | OEIS-A284930

Positions of 0 in A284929; complement of A284931.

Conjecture: 1 < n*r - a(n) < 4 for n >= 1, where r = 2 + sqrt(2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011950 | OEIS-A284931

Positions of 1 in A284929; complement of A284930.

Conjecture: -1 < n*sqrt(2) - a(n) < 1, for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011951 | OEIS-A284933

Positions of 0 in A284792; complement of A284934.

a(n) - a(n-1) is in {2,3,4} for n>=2. Conjecture: a(n)/n -> 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011952 | OEIS-A284934

Positions of 1 in A284792; complement of A284933.

a(n) - a(n-1) is in {1,2} for n>=2. Conjecture: a(n)/n -> 3/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011953 | OEIS-A284936

Positions of 0 in A284935; complement of A284937.

Conjecture: a(n) - 2n + 1 is in {0,1,2} for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011954 | OEIS-A284937

Positions of 1 in A284935; complement of A284936.

Conjecture: 2n - a(n) is in {0,1,2} for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011955 | OEIS-A284940

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 107", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 28 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011956 | OEIS-A284941

Positions of 1 in A284939; complement of A080580.

Conjecture: -1 < n*sqrt(2) - a(n) < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011957 | OEIS-A284945

Positions of 0 in A284944; complement of A284946.

Conjecture: -4 < n*r - a(n) < 3 for n >= 1, where r = 2 + sqrt(2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011958 | OEIS-A284946

Positions of 1 in A284944; complement of A284945.

Conjecture: -1 < n*r - a(n) < 3 for n >= 1, where r = 2 + sqrt(2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011959 | OEIS-A284948

1-limiting word of the morphism 0 -> 10, 1 -> 00.

Conjecture: a(n) = A096268(n-2), n>1. - _R. J. Mathar_, Jun 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011960 | OEIS-A284955

Positions of 0 in A284954; complement of A284956.

Conjecture: a(n)/n -> (5 + sqrt(13))/6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011961 | OEIS-A284956

Positions of 1 in A284954; complement of A284955.

Conjecture: a(n)/n -> (3 + sqrt(13))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011962 | OEIS-A284958

Positions of 0 in A284957; complement of A284959.

Conjecture: a(n)/n -> (5 + sqrt(13))/6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011963 | OEIS-A284959

Positions of 1 in A284957; complement of A284958.

Conjecture: a(n)/n -> (3 + sqrt(13))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011964 | OEIS-A285004

Numbers n that enter a cycle of greater length than that for any k < n in the iteration sequence s(0)=n, s(k+1) = s(k) + (-1)^k*d(s(k)), where d(n) is the number of divisors of n (A000005).

Claudia Spiro conjectured that for every initial number n the iteration sequence is eventually periodic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Apr 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011965 | OEIS-A285022

Numbers n such that A002088(n) < 3n^2/Pi^2.

James Joseph Sylvester conjectured in 1883 that A002088(n) > 3n^2/Pi^2 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Apr 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011966 | OEIS-A285032

Positions of 0 in A285031; complement of A285033.

Conjecture: -1 < n*r - a(n) < 2 for n>=1, where r = 1 + sqrt(1/2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011967 | OEIS-A285033

Positions of 1 in A285031; complement of A285032.

Conjecture: -1 < n*r - a(n) < 2 for n>=1, where r = 1 + sqrt(2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011968 | OEIS-A285035

Positions of 0 in A285034; complement of A285036.

Conjecture: -1 < n*r - a(n) < 2 for n>=1, where r = 1 + sqrt(1/2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011969 | OEIS-A285036

Positions of 1 in A285034; complement of A285035.

Conjecture: -1 < n*r - a(n) < 2 for n>=1, where r = 1 + sqrt(2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011970 | OEIS-A285052

Number of idempotent equivalence classes for multiplication in Zn.

Conjecture: a(n) = 4^A001221(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chad Brewbaker_, Apr 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011971 | OEIS-A285074

Positions of 0 in A285073; complement of A285075.

Conjecture: -1 < n*r - a(n) < 1 for n>=1, where r = 1 + sqrt(1/2). Proof of a stronger form of the conjecture: the sequence d:=A285073 can be written as d=01c, where c is the homogeneous Sturmian sequence with slope alpha = sqrt(2)-1 (see comments of A285073). Changing from alpha to 1-alpha = 2-sqrt(2) turns c into its mirror image, so we have to find the positions of 1 in this new sequence. In our case s = 1/(2-sqrt(2)) = 1+sqrt(1/2). It follows that for n=0,1,... one has a(n+2) = floor((n+1)*(1+sqrt(1/2))) + 2, which directly implies that sqrt(1/2)-1 < n*r - a(n) < sqrt(1/2), which is a strengthening of the conjecture (actually there is no strict inequality for n=1: r-a(1) = sqrt(1/2)). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011972 | OEIS-A285075

Positions of 1 in A285073; complement of A285074.

Conjecture: -1 + sqrt(2) < n*r - a(n) < sqrt(2) for n>=1, where r = 1 + sqrt(2). Proof of the conjecture: the sequence d:=A285073 can be written as d=01c, where c is the homogeneous Sturmian sequence with slope sqrt(2)-1 (see comments of A285073). In our case s=1+sqrt(2). It follows that for n=0,1,... one has a(n+2) = floor((n+1)*(1+sqrt(2))) + 2, which directly implies the conjecture. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011973 | OEIS-A285076

1-limiting word of the morphism 0->10, 1-> 010.

Conjecture: a(n) = A285073(n), n>=3. - _R. J. Mathar_, Jun 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011974 | OEIS-A285077

Positions of 0 in A285076; complement of A285078.

Conjecture: -1 < n*r - a(n) < 1 for n>=1, where r = 1 + sqrt(1/2). This conjecture can be proved in the same way as the conjecture on A285074. - _Michel Dekking_, May 30 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011975 | OEIS-A285078

Positions of 1 in A285076; complement of A285077.

Conjecture: -1 + sqrt(2) < n*r - a(n) < sqrt(2) for n>=1, where r = 1 + sqrt(2). Both these conjectures can be proved (for n>=2), see the Comments of A285073 and A285075 - _Michel Dekking_, Sep 24 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011976 | OEIS-A285081

Positions of 0 in A285080; complement of A285082.

Conjecture: 0 < n*r - a(n) < r for n>=1, where r = (3+sqrt(5))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011977 | OEIS-A285082

Positions of 1 in A285080; complement of A285081.

Conjecture: -1 < n*r - a(n) < 1 for n>=1, where r = (1+sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011978 | OEIS-A285084

Positions of 0 in A285083; complement of A285085.

Conjecture: 0 < n*r - a(n) < r for n>=1, where r = (3+sqrt(5))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011979 | OEIS-A285085

Positions of 1 in A285083; complement of A285084.

Conjecture: -1 < n*r - a(n) < 1 for n>=1, where r = (1+sqrt(5))/2, the golden ratio (A001622).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011980 | OEIS-A285086

Numbers n such that the number of partitions of n^2+1 (=A000041(n^2+1)) is prime.

Because asymptotically A000041(n^2+1) ~ exp(Pi*sqrt(2/3*(n^2+1))) / (4*sqrt(3)*(n^2+1)), the sum of the prime probabilities ~ 1/log(A000041(n^2+1)) is diverging and there are no obvious restrictions on primality; therefore, this sequence may be conjectured to be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Serge Batalov_, Apr 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011981 | OEIS-A285088

Numbers n such that the number of partitions of n(n+1)/2 (=A000041(A000217(n))) is prime.

Because asymptotically A000041(n*(n+1)/2) ~ exp(Pi*sqrt(2/3*(n*(n+1)/2))) / (4*sqrt(3)*(n*(n+1)/2)), the sum of the prime probabilities ~1/log(A000041(n*(n+1)/2)) is diverging and there are no obvious restrictions on primality; therefore, this sequence may be conjectured to be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Serge Batalov_, Apr 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011982 | OEIS-A285091

Positions of 0 in A284848; complement of A285092.

a(n) - a(n-1) is in {1,2,3,4} for n>=2. Conjecture: a(n)/n -> 9/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011983 | OEIS-A285092

Positions of 1 in A284848; complement of A285091.

a(n) - a(n-1) is in {1,2,3} for n>=2. Conjecture: a(n)/n -> 9/5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011984 | OEIS-A285126

Positions of 0 in A285125; complement of A285127.

Conjecture: 0 < n*r - a(n) < 1 for n>=1, where r = 1.3903...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011985 | OEIS-A285127

Positions of 1 in A285125; complement of A285126.

Conjecture: -3 < n*r - a(n) < 0 for n>=1, where r = 3.5621...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011986 | OEIS-A285129

Positions of 0 in A285128; complement of A285130.

Conjecture: -1 < n*r - a(n) < 1 for n>=1, where r = 1.3903...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011987 | OEIS-A285130

Positions of 1 in A285128; complement of A285139.

Conjecture: -1 < n*r - a(n) < 3 for n>=1, where r = 3.5621...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011988 | OEIS-A285134

Positions of 0 in A285133; complement of A285135.

Conjecture: -1 < n*r - a(n) < 1 for n>=1, where r = 1 + sqrt(1/3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011989 | OEIS-A285135

Positions of 1 in A285133; complement of A285134.

Conjecture: -1 < n*r - a(n) < 1 for n>=1, where r = 1 + sqrt(3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011990 | OEIS-A285137

Positions of 0 in A285136; complement of A285138.

Conjecture: -1 < n*r - a(n) < 1 for n>=1, where r = 1 + sqrt(1/3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011991 | OEIS-A285138

Positions of 1 in A285136; complement of A285137.

Conjecture: -1 < n*r - a(n) < sqrt(3) for n>=1, where r = 1 + sqrt(3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011992 | OEIS-A285140

Positions of 0 in A285139; complement of A285141.

Conjecture: -1 < n*r - a(n) < 1 for n>=1, where r = (3 + sqrt(3))/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011993 | OEIS-A285141

Positions of 1 in A285139; complement of A285140.

Conjecture: -1 < n*r - a(n) < 1 for n>=1, where r = 1 + sqrt(3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011994 | OEIS-A285143

Positions of 0 in A285142; complement of A285144.

Conjecture: -1 < n*r - a(n) < 1 for n>=1, where r = (3 + sqrt(3))/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011995 | OEIS-A285144

Positions of 1 in A285142; complement of A285143.

Conjecture: -1 < n*r - a(n) < sqrt(3) for n>=1, where r = 1 + sqrt(3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011996 | OEIS-A285160

Positions of 0 in A285159; complement of A285161.

Conjecture: a(n)/n -> 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011997 | OEIS-A285161

Positions of 1 in A285159; complement of A285160.

Conjecture: a(n)/n -> 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011998 | OEIS-A285163

Positions of 0 in A285162; complement of A285164.

Conjecture: a(n)/n -> 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-011999 | OEIS-A285164

Positions of 1 in A285162; complement of A285163.

Conjecture: a(n)/n -> 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012000 | OEIS-A285195

G.f.: 2C(x)/(3-sqrt(1-4xC(x))) where C = g.f. for A000108.

Conjecture D-finite with recurrence: +126*n*(n-1)*(n+1)*a(n) -3*n*(n-1)*(691*n-1037)*a(n-1) +(n-1)*(12023*n^2-48049*n+47736)*a(n-2) +2*(-12889*n^3+93970*n^2-221427*n+169506)*a(n-3) +4*(-248*n^3+12718*n^2-88433*n+160113)*a(n-4) +16*(2918*n^3-41574*n^2+198532*n-317121)*a(n-5) +240*(4*n-19)*(2*n-9)*(4*n-21)*a(n-6)=0. - _R. J. Mathar_, Jan 25 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012001 | OEIS-A285206

Positions of 0 in A285205; complement of A285207.

Conjecture: -1 < n*r - a(n) < 1, where r = 1 + sqrt(1/3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012002 | OEIS-A285207

Positions of 1 in A285205; complement of A285206.

Conjecture: 0 < n*r - a(n) < 3, where r = 1 + sqrt(3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012003 | OEIS-A285209

Positions of 0 in A285208; complement of A285210.

Conjecture: -1 < n*r - a(n) < 1, where r = 1 + sqrt(1/3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012004 | OEIS-A285210

Positions of 1 in A285208; complement of A285209.

Conjecture: 0 < n*r - a(n) < 3, where r = 1 + sqrt(3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012005 | OEIS-A285250

Positions of 0 in A285249; complement of A285251.

Conjecture: 2n - a(n) is in {0,1} for n>=1. Proof of this conjecture: Let (b(n))=A285249. Then clearly b(2n-1)b(2n)=01 or 10 for n=1,2,...So at time 2n you have seen n zeroes in the sequence b. The last of these must have occurred at time 2n or at time 2n-1, so either 2n-a(n)=0 or 2n-a(n)=1. - _Michel Dekking_, Jul 31 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012006 | OEIS-A285251

Positions of 1 in A285249; complement of A285250.

Conjecture: 2n - a(n) is in {0,1} for n>=1. See the comments of A285250 for a proof of this conjecture. - _Michel Dekking_, Oct 07 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012007 | OEIS-A285275

Positions of 0 in A285274; complement of A285276.

Conjecture: 0 < n*r - a(n) < 6 for n>=1, where r = 2+sqrt(2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012008 | OEIS-A285276

Positions of 1 in A285274; complement of A285275.

Conjecture: -2 < n*sqrt(2) - a(n) < 1 for n>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012009 | OEIS-A285278

Positions of 0 in A285277; complement of A285279.

Conjecture: 0 < n*r - a(n) < 6 for n>=1, where r = 2+sqrt(2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012010 | OEIS-A285279

Positions of 1 in A285277; complement of A285278.

Conjecture: 0 < n*r - a(n) < 6 for n>=1, where r = 2+sqrt(2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012011 | OEIS-A285296

Lexicographically earliest sequence of distinct positive terms such that the product of two consecutive terms is divisible by p^2 for some prime p.

Conjecturally, a(n) ~ n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Apr 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012012 | OEIS-A285301

Fixed point of the morphism 0 -> 10, 1 -> 1000.

Conjecture: a(n) = A284751(n+1). - _R. J. Mathar_, May 08 2017 Proof of Mathar's conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012013 | OEIS-A285302

Positions of 0 in A285301, complement of A086398.

Conjecture: -2 < n*r - a(n) < 2 for n>=1, where r = 1 + sqrt(1/3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012014 | OEIS-A285303

Positions of 1 in A284932; complement of A285303.

Conjecture: lim_{n->oo} a(n)/n = 0.78..., and if m denotes this number, then -1 < a(n)/n - m < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012015 | OEIS-A285305

Fixed point of the morphism 0 -> 10, 1 -> 1001.

Conjecture: a(n) = A284775(n+1). - _R. J. Mathar_, May 08 2017 Proof of this conjecture. Let mu: 0 -> 10, 1 -> 1001 be the defining morphism of (a(n)), and let tau: 0 -> 01, 1 -> 0011 be the defining morphism of A284775. Since mu^n(0) starts with 1 for n>1, mu^n(0) tends to (a(n)) as n tends to infinity. So the conjecture will follow directly from the following claim.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012016 | OEIS-A285320

If n == 0 or A008683(n) == 0, then a(n) = 0, otherwise a(n) = 1+a(A048675(n)); number of iterations of A048675 needed before the result is either zero or nonsquarefree number (A013929).

Conjecture: all terms are well-defined (finite). This implies also the conjecture I have made in A019565.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Apr 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012017 | OEIS-A285331

Inverse for A285332: a(1) = 0, a(2) = 1, a(A019565(n)) = 2*a(n), a(A065642(n)) = 1 + 2*a(n).

For the question whether this sequence and A285332 are permutations of natural numbers, see comments in A285332 and the conjecture stated in A019565.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Apr 17 2017, comments edited Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012018 | OEIS-A285332

a(0) = 1, a(1) = 2, a(2n) = A019565(a(n)), a(2n+1) = A065642(a(n)).

As long as it remains conjecture that A019565 has no cycles, it is certainly also an open question whether this is a permutation of the natural numbers: If A019565 has any cycles, then neither any of the terms in those cycles nor any A065642-trajectories starting from those terms (that is, numbers sharing same prime factors) may occur in this tree.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Apr 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012019 | OEIS-A285341

Fixed point of the morphism 0 -> 10, 1 -> 1011.

A short proof of Mathar's conjecture can be obtained by using the fact that sigma and tau are conjugate morphisms. For all words w one has Conjecture: a(n) = A284893(n+1). - _R. J. Mathar_, May 08 2017 Proof of this conjecture: let sigma be the morphism 0 -> 10, 1 -> 1011, and let tau be the morphism 0 -> 01, 1 -> 0111, which has A284893 as a fixed point. It clearly suffices to prove the relation, for all n=1,2,3,...: Using the induction hypothesis with (A) and (B) in the second line, one obtains

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012020 | OEIS-A285342

Positions of 0 in A285341; complement of A285343.

Conjecture: 0 < n*r - a(n) < 4 for n>=1, where r = 2 + sqrt(2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012021 | OEIS-A285343

Positions of 1 in A285341; complement of A285342.

Conjecture: -1 < n*sqrt(2) - a(n) < 1 for n>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012022 | OEIS-A285345

Fixed point of the morphism 0 -> 10, 1 -> 1100.

Conjecture: a(n) = A284905(n+1). - _R. J. Mathar_, May 08 2017 Proof of Mathar's conjecture: let alpha be the morphism 0 -> 10, 1 -> 1100, and let beta be the morphism 0 -> 01, 1 -> 1001, which has A284905 as its fixed point starting with 0. Note that alpha^n(0) tends to (a(n)) as n tends to infinity because alpha(0) starts with 1. It therefore suffices to prove the relation Here we used (B) with w = beta^n(0) in line three, and the induction hypothesis in line four. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012023 | OEIS-A285346

Positions of 0 in A285345; complement of A285347.

Conjecture: 2n + 1 - a(n) is in {0,1} for n>=1; see A285351.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012024 | OEIS-A285347

Positions of 1 in A285345; complement of A285346.

Conjecture: 2n - 1 - a(n) is in {0,1} for n >= 1. Proof of the conjecture by Kimberling. I will show that

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012025 | OEIS-A285352

Positions of 0 in A284932; complement of A285303.

Conjecture: lim_{n->oo} a(n)/n = 0.55..., and if m denotes this number, then -1 < a(n)/n - m < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012026 | OEIS-A285353

Positions of 0 in A285351; complement of A285354.

Conjecture: 3n - 2 - a(n) is in {0,1} for n>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012027 | OEIS-A285354

Positions of 1 in A285351; complement of A285353.

Conjecture: 3n/2 +1/2 - a(n) is in {0,1/2,1} for n>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012028 | OEIS-A285356

Numbers n such that the entries in the n-th row of the irregular triangle A237591 are in nonincreasing order.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Apr 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012029 | OEIS-A285358

Fixed point of the morphism 0 -> 10, 1 -> 1101.

Conjecture: a(n) = A284929(n+1). - _R. J. Mathar_, May 08 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012030 | OEIS-A285359

Positions of 0 in A285358; complement of A285360.

Conjecture: -1 < n*r - a(n) < 1 for n>=1, where r = 2 + sqrt(2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012031 | OEIS-A285360

Positions of 1 in A285358; complement of A285359.

Conjecture: 0 < n*sqrt(2) - a(n) < 2 for n>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012032 | OEIS-A285373

Fixed point of the morphism 0 -> 10, 1 -> 1110.

Proof of Mathar's conjecture of May 08 2017: Furthermore, using the induction hypothesis with (B) and (C) in the first line, and again (C) in the third line, one obtains Conjecture: a(n) = A284939(n+1). - _R. J. Mathar_, May 08 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012033 | OEIS-A285374

Positions of 0 in A285373; complement of A285375.

Conjecture: -4 < n*sqrt(2) - a(n) < 1 for n>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012034 | OEIS-A285375

Positions of 1 in A285373; complement of A285374.

Conjecture: 0 < n*sqrt(2) - a(n) < 3 for n>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012035 | OEIS-A285383

Limiting 0-word of the morphism 0 -> 11, 1 -> 01.

Conjecture: a(n) = A035263(n-1). - _R. J. Mathar_, May 08 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012036 | OEIS-A285385

Positions of 1 in A285384; complement of A072939.

a(n + 1) = a(n) + A096268(a(n) - 1) + 1 for all n > 0. (conjecture) - _Velin Yanev_, Mar 23 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012037 | OEIS-A285401

Positions of 0 in A285177; complement of A285402.

Conjecture: a(n)/n -> (61-sqrt(3))/26 = 2.279... This conjecture is false. In fact,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012038 | OEIS-A285402

Positions of 1 in A285177; complement of A285401.

Conjecture: a(n)/n -> (82 + sqrt(3))/47 = 1.781... This conjecture is false. In fact, a(n)/n --> (5+sqrt(17))/(1+sqrt(17)) = 1.7807764... = A188485. See A285401. It follows in the same way as there, that a(n)/n --> 1/f1, where f1 is the frequency of 1's in A285177, and f1 can be computed using the Perron Frobenius theorem. - _Michel Dekking_, Feb 10 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012039 | OEIS-A285412

Positions of 0 in A285411; complement of A285413.

Conjecture: a(n)/n -> (61-sqrt(3))/26 = 2.279... This conjecture is false. In fact, a(n)/n --> (5+sqrt(17))/ 4 = 2.28077..., the same limit as for A285401(n)/n. The reason is that the generating morphism for (a(n)) has the same incidence matrix as the generating morphism for A285177. - _Michel Dekking_, Feb 10 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012040 | OEIS-A285413

Positions of 1 in A285411; complement of A285412.

Conjecture: a(n)/n -> (61-sqrt(3))/26 = 2.279...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012041 | OEIS-A285419

Positions of 0 in A285418; complement of A285420.

Conjecture: -1 < n*r - a(n) < 5 for n>=1, where r = 2 + sqrt(3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012042 | OEIS-A285420

Positions of 1 in A285418; complement of A285419.

Conjecture: -2 < n*r - a(n) < 2 for n>=1, where r = (1 + sqrt(3))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012043 | OEIS-A285422

Positions of 0 in A285421; complement of A285423.

Conjecture: -1 < n*r - a(n) < 5 for n>=1, where r = 2 + sqrt(3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012044 | OEIS-A285423

Positions of 1 in A285421; complement of A285422.

Conjecture: -2 < n*r - a(n) < 2 for n>=1, where r = (1 + sqrt(3))2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012045 | OEIS-A285428

Positions of 0 in A285427; complement of A285429.

Conjecture: a(n)/n -> (61 - sqrt(3))/26.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012046 | OEIS-A285429

Positions of 1 in A285427; complement of A285428.

Conjecture: a(n)/n -> (82 + sqrt(3))/47.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012047 | OEIS-A285432

Positions of 0 in A286939; complement of A285564.

Conjecture: lim_{n->oo} a(n)/n = 1.25..., and if m denotes this number, then -1 < a(n)/n - m < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012048 | OEIS-A285473

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 3", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 19 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012049 | OEIS-A285474

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 3", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 19 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012050 | OEIS-A285487

Lexicographically earliest sequence of distinct positive terms such that the product of two consecutive terms has at least 5 distinct prime factors.

Conjecturally, a(n) ~ n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012051 | OEIS-A285493

a(n) is the least positive integer not already appearing such that no two distinct unordered pairs of points ((n, a(n)), (m, a(m))) and ((k, a(k)), (j, a(j))) have the same midpoint.

Conjecture: This is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Apr 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012052 | OEIS-A285502

Positions of 0 in A285501; complement of A285503.

Conjecture: a(n)/n -> (3+sqrt(5))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012053 | OEIS-A285503

Positions of 1 in A285501; complement of A285502.

Conjecture: a(n)/n -> (1+sqrt(5))/2 = golden ratio (A001622).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012054 | OEIS-A285505

Positions of 0 in A285504; complement of A285506.

Conjecture: a(n)/n -> (3+sqrt(5))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012055 | OEIS-A285506

Positions of 1 in A285504; complement of A285505.

Conjecture: a(n)/n -> (1+sqrt(5))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012056 | OEIS-A285515

{00->0, 11->1}-transform of A285501.

As a word, A285501 = 0011001100110011111100110011111100110011..., so that the transform of A285501 that results from the substitutions 00-> and 11->1 is 0101010111010111010101010101111010111010101... The sequence is also the 0-limiting word of the morphism 0->11, 1->0101. See A285517 and A285518 for conjectured connections to the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012057 | OEIS-A285516

Positions of 0 in A285515; complement of A285517.

Conjecture: a(n)/n -> (3+sqrt(5))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012058 | OEIS-A285517

Positions of 1 in A285515; complement of A285516.

Conjecture: a(n)/n -> (1+sqrt(5))/2 = golden ratio (A001622).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012059 | OEIS-A285518

{00->0, 11->1}-transform of A285504.

See A285519 and A285520 for conjectured connections to the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012060 | OEIS-A285519

Positions of 0 in A285518; complement of A285520.

Conjecture: a(n)/n -> (3+sqrt(5))/2. Here is a proof of this conjecture. The limit of a(n)/n as n tends to infinity is equal to the inverse 1/f0 of the frequency f0 of the letter 0 in the sequence (a(n)) (see also the argument in A285401).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012061 | OEIS-A285520

Positions of 1 in A285518; complement of A285519.

Conjecture: a(n)/n -> (1+sqrt(5))/2 = golden ratio (A001622). Proof of this conjecture: the limit equals 1/f1, where f1 is the frequency of 1 in (a(n)), and 1/f1 = phi = golden ratio. See A285519 for more details. - _Michel Dekking_, Feb 22 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012062 | OEIS-A285521

Table read by rows: the n-th row gives the lexicographically earliest sequence of length n such that the convex hull of (1, a(1)), ..., (n, a(n)) is an n-gon with minimum height.

Conjecture: Even-length rows have rotational symmetry. Conjecture: The maximum value in any even-length row is the same as the maximum value in the preceding row. Conjecture by Peter Kagey: 24th row is 12, 14, 8, 17, 5, 4, 21, 22, 2, 23, 1, 1, 24, 24, 2, 23, 3, 4, 21, 20, 8, 17, 11, 13. - _Lars Blomberg_, May 06 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Apr 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012063 | OEIS-A285531

Positions of 0 in A285530; complement of A285532.

Conjecture: a(n)/n -> 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012064 | OEIS-A285532

Positions of 1 in A285530; complement of A285533.

Conjecture: a(n)/n -> 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012065 | OEIS-A285534

Positions of 0 in A285533; complement of A285535.

Conjecture: a(n)/n -> 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012066 | OEIS-A285535

Positions of 1 in A285533; complement of A285534.

Conjecture: a(n)/n -> 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012067 | OEIS-A285540

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 35", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012068 | OEIS-A285541

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 35", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012069 | OEIS-A285542

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 35", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012070 | OEIS-A285543

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 35", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 21 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012071 | OEIS-A285549

Smallest weak pseudoprime to all natural bases up to prime(n) that is not a Carmichael number.

Conjecture: lpf(a(n)) > prime(n), where lpf = A020639. - _Thomas Ordowski_, May 13 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Apr 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012072 | OEIS-A285564

Positions of 1 in A286939; complement of A285432.

Conjecture: lim_{n->oo} a(n)/n = 4.89..., and if m denotes this number, then -1 < a(n)/n - m < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012073 | OEIS-A285566

Positions of 0 in A285565; complement of A285567.

Conjecture: a(n)/n -> (3+sqrt(5))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012074 | OEIS-A285567

Positions of 1 in A285565; complement of A285566.

Conjecture: a(n)/n -> (1+sqrt(5))/2 = golden ratio (A001622).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012075 | OEIS-A285569

Positions of 0 in A285568; complement of A286031.

Conjecture: a(n)/n -> (3+sqrt(5))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012076 | OEIS-A285590

Positions of 0 in A285589; complement of A285591.

Conjecture: a(n)/n -> 1+sqrt(5/4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012077 | OEIS-A285591

Positions of 1 in A285589; complement of A285590.

Conjecture: a(n)/n -> 1+sqrt(4/5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012078 | OEIS-A285593

Positions of 0 in A285592; complement of A285594.

Conjecture: a(n)/n -> 1+sqrt(5/4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012079 | OEIS-A285594

Positions of 1 in A285592; complement of A285593.

Conjecture: a(n)/n -> 1+sqrt(4/5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012080 | OEIS-A285597

Positions of 0 in A285596; complement of A285598.

Conjecture: a(n)/n -> (5+sqrt(17))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012081 | OEIS-A285598

Positions of 1 in A285596; complement of A285597.

Conjecture: a(n)/n -> (1+sqrt(17))/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012082 | OEIS-A285600

Positions of 0 in A285599; complement of A285601.

Conjecture: a(n)/n -> (5+sqrt(17))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012083 | OEIS-A285601

Positions of 1 in A285599; complement of A285600.

Conjecture: a(n)/n -> (1+sqrt(17))/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012084 | OEIS-A285612

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 62", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 23 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012085 | OEIS-A285613

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 62", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 23 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012086 | OEIS-A285618

Positions of 0 in A285617; complement of A285619.

Conjecture: a(n)/n -> 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012087 | OEIS-A285619

Positions of 1 in A285617; complement of A285618.

Conjecture: a(n)/n -> 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012088 | OEIS-A285622

Positions of 0 in A285621; complement of A285623.

Conjecture: a(n)/n -> (3 + sqrt(5))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012089 | OEIS-A285623

Positions of 1 in A285621; complement of A285622.

Conjecture: a(n)/n -> (1 + sqrt(5))/2 = golden ratio (A001622).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012090 | OEIS-A285626

Positions of 0 in A285625; complement of A285627.

Conjecture: a(n)/n -> (3 + sqrt(5))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012091 | OEIS-A285627

Positions of 1 in A285625; complement of A285626.

Conjecture: a(n)/n -> (1 + sqrt(5))/2 = golden ratio (A001622).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012092 | OEIS-A285643

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 65", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 23 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012093 | OEIS-A285644

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 65", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 24 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012094 | OEIS-A285645

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 65", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 24 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012095 | OEIS-A285646

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 65", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 24 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012096 | OEIS-A285647

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 73", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 24 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012097 | OEIS-A285648

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 73", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 24 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012098 | OEIS-A285649

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 73", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 24 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012099 | OEIS-A285650

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 73", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 24 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012100 | OEIS-A285655

Lexicographically earliest sequence of distinct positive terms such that the product of two consecutive terms has at least 6 distinct prime factors.

Conjecturally, a(n) ~ n. - conjecturally: d_k(n) ~ n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Apr 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012101 | OEIS-A285658

Positions of 0 in A285657; complement of A285659.

Conjecture: a(n)/n -> (5 + sqrt(17))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012102 | OEIS-A285659

Positions of 1 in A285657; complement of A285658.

Conjecture: a(n)/n -> (1 + sqrt(17))/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012103 | OEIS-A285662

Positions of 0 in A285661; complement of A285663.

Conjecture: a(n)/n -> (3 + sqrt(5))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012104 | OEIS-A285663

Positions of 1 in A285661; complement of A285662.

Conjecture: a(n)/n -> (1 + sqrt(5))/2 = golden ratio (A001622).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012105 | OEIS-A285665

a(n) = a(n-1) + 2*a(n-2) - a(n-3) - a(n-4) + 2*a(n-5) - 2*a(n-7) for n>7 and where a(0)=2, a(1)=3, a(2)=6, a(3)=10, a(4)=17, a(5)=29, a(6)=51.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->001, 1->000, starting with 00; see A286939.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012106 | OEIS-A285666

Fixed point of the mapping 00->001, 1->010, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate of the mapping is given by A001580.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012107 | OEIS-A285667

Positions of 0 in A285666; complement of A286058.

Conjecture: lim_{n->oo} a(n)/n = 1.41..., and if m denotes this number, then -1 < a(n)/n - m < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012108 | OEIS-A285669

Positions of 0 in A285668; complement of A285670.

Conjecture: -1 < n*r - a(n) < 4 for n>=1, where r = (5 + sqrt(17))/2. Here is a proof of a weak form of the conjecture by Kimberling. which identifies the number r in the conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012109 | OEIS-A285670

Positions of 1 in A285668; complement of A285669.

Conjecture: -1 < n*r - a(n) < 4 for n>=1, where r = (1 + sqrt(17))/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012110 | OEIS-A285678

Positions of 0 in A285677.

where 1/t + 1/u + 1/v = 1. Conjecture: a(n) - a(n-1) is in {2,3,4,5} for n>=2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012111 | OEIS-A285679

Positions of 2 in A285677.

Conjecture: a(n) - a(n-1) is in {2,5} for n>=2. See A285683 for a proof of this conjecture. - _Michel Dekking_, Oct 09 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012112 | OEIS-A285681

Positions of 0 in A285680.

where 1/t + 1/u + 1/v = 1. Conjecture: a(n) - a(n-1) is in {1,2} for n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012113 | OEIS-A285682

Positions of 1 in A285680.

where 1/t + 1/u + 1/v = 1. Conjecture: a(n) - a(n-1) is in {5,7} for n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012114 | OEIS-A285683

Positions of 2 in A285680.

Conjecture: a(n) - a(n-1) is in {5,7} for n >= 2. Proof of the correct version of Kimberling's conjecture: This conjecture is false; it should be replaced by a(n) - a(n-1) is in {2,5} for n >= 2. The truth of that conjecture follows from Lemma 8 in Allouche and Dekking, since (a(n)) is a generalized Beatty sequence (see Formula below). - _Michel Dekking_, Oct 09 2018 Proof of Mathar's conjecture: Let xF:=A003849 = 01001010010010100... be the Fibonacci sequence. The two return words of 1010 are r1=10100100, and r2=10100. [See Justin & Vuillon (2000) for definition of return word. - _N. J. A. Sloane_, Sep 23 2019] If these occur at index i, then in BOTH cases there is an occurrence of 0010 in xF at index i+3 in xF. These will be called type-A occurrences. Since r1 has suffix 00, there will also be occurrences of 0010 at index i+6, if r1 occurs at index i. These will be called type-B occurrences. It is clear that there are no other occurrences of 0010 in xF. Now if we perform the [0010->2] transform T on xF, then the type-B occurrences of 0010 will not yield the occurrence of a 2 in T(xF), since the StringReplace procedure acts from left to right, destroying the 0010 occurrences of type-B. The conclusion is that the positions of 2 in the [1010->2] transform and the [0010->2] transform are coupled, occurring at a fixed shift of each other, except that there is an extra first occurrence of 0010 at index 3. By coincidence, the first occurrence of 1010 in xF is at index 5, which equals the index of the second occurrence of 0010, with 3 subtracted (that has to be done because of the shrinking of a block of length 4 to the letter 2). It follows that the two occurrence sequences are equal, except for the extra 3. - _Michel Dekking_, Oct 09 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012115 | OEIS-A285697

Positions of 1 in A288741; complement of A288742.

Conjecture: a(n)/n -> 2, and -1/2 < 2 - a(n)/n < 1/3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012116 | OEIS-A285744

Lexicographically earliest sequence of distinct positive terms such that, for any n>0, n*a(n) has at least 5 distinct prime factors.

Conjecturally, a(n) ~ n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012117 | OEIS-A285766

Maximum spillway height for a zero or one bend minimal area lake in a number square.

Conjectures from _Colin Barker_, May 07 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Craig Knecht_, May 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012118 | OEIS-A285770

Let q_1<q_2<... be primes not dividing 2n+1 such that 2n-q_i + 1 is twice a prime. a(n) is smallest i such that 2n+2q_i+1 is prime or a(n)=0 if there is no such i.

By the Lemoine-Levy conjecture, for every n>=3, there are primes p and q such that 2*n+1=2*p+q. In A277688 were considered the numbers of the form 2n+1+2*p, in this sequence we consider the numbers of the form 2*n+1+2*q.

Then to the same condition as in A277688 satisfies an extremely rare set of numbers that contains numbers {2*n+1} for which a(n)=0: {11, 59, 151,...}. Comparing this with our conjecture in A277688, we conjecture here that this set is finite. For the explanation of this conjecture we need not refer to the minimal number of the representation 2*n+1 for large n (most likely, it is, as in the Goldbach presentations, for 2*n, more than c*n/(log n)^2 with some constant c) since we have a prime only among the first several of these representations, as in this sequence. This leaves us with an important question: why does A277688 contain much more terms than the zeros in this sequence?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012119 | OEIS-A285775

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 89", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 26 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012120 | OEIS-A285776

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 89", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 26 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012121 | OEIS-A285777

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 89", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 26 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012122 | OEIS-A285778

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 89", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 26 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012123 | OEIS-A285786

Number of primes p with 2(n-1)^2 < p <= 2n^2.

The author of the sequence conjectures that a(n) >= 1 for all n. This conjecture is similar to the famous conjecture made by Adrien-Marie Legendre that there is always a prime between n^2 and (n+1)^2, see A014085. - _Antti Karttunen_, May 01 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Steiner_, Apr 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012124 | OEIS-A285796

Number of ways to write n as an ordered sum of two numbers that are the product of an even number of distinct primes (including 1).

Conjecture: a(n) > 0 for all n > 82.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Apr 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012125 | OEIS-A285797

Number of ways to write n as an ordered sum of two numbers that are the product of an odd number of distinct primes.

Conjecture: a(n) > 0 for all n > 57.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Apr 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012126 | OEIS-A285802

Smallest number that when multiplied by n contains the digit 7.

Conjecture: 35 is the highest value to occur in this sequence. Conjecture is true. First of all we notice that a(10*k) = a(k), so we can consider only numbers not ending in 0. Then, it is easy to verify that for any number k between 1 and 99 not ending in 0, there is a multiple of k not larger than 35*k which has a 7 in one of its last two digits. Since the last two digits of the products only depend on the last two digits of k, this extends immediately to larger numbers. - _Giovanni Resta_, Apr 27 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Apr 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012127 | OEIS-A285804

Composite numbers of the form 12*k+5 or 12*k+7 for some k.

Conjecture: Let r(n) = (a(n)-1)/(a(n)+1) if a(n) mod 4 = 1, (a(n)+1)/(a(n)-1) otherwise; then Product_{n>=1} r(n) = (28/27) * (32/33) * (38/39) * (46/45) * ... = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimitris Valianatos_, Apr 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012128 | OEIS-A285833

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 107", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 28 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012129 | OEIS-A285834

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 107", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 28 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012130 | OEIS-A285835

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 107", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 28 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012131 | OEIS-A285842

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 115", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Apr 28 2017: (Start) The conjectured generating function, closed forms, and equivalent linear recurrence are false; the first counterexample occurs at n = 28, where a(28) = 1100110000001100000000000000, whereas the conjectured formulas give 1100000000000000. - _Valentin Haase_, Jul 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012132 | OEIS-A285848

Lexicographically earliest sequence of distinct positive terms such that for any n>0, i > j >=0, gcd(a^i(n), a^j(n)) = 1 (where a^k denotes the k-th iteration of a).

Conjecturally, a(n) ~ 2*n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Apr 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012133 | OEIS-A285869

a(n) is the number of zeros of the Chebyshev S(n, x) polynomial in the open interval (-sqrt(2), +sqrt(2)).

See a May 06 2017 comment on A049310 where these problems are considered which originated in a conjecture by _Michel Lagneau_ (see A008611) on Fibonacci polynomials.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, May 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012134 | OEIS-A285891

Triangle read by rows: T(n,k) = n*A237048(n,k).

Conjecture: T(n,k) = n, is also the sum of the parts of the partition of n into k consecutive parts, if such a partition exists, otherwise T(n,k) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, May 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012135 | OEIS-A285901

Total number of partitions of all positive integers <= n into an odd number of consecutive parts.

Conjecture: a(n) is also the total number of subparts present (totally or partially) in an octant of the symmetric representations of sigma of all positive integers <= n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, May 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012136 | OEIS-A285902

Total number of partitions of all positive integers <= n into an even number of consecutive parts.

Conjecture: a(n) is also the total number of pairs of equidistant subparts of all symmetric representations of sigma of all positive integers <= n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, May 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012137 | OEIS-A285914

Irregular triangle read by rows: T(n,k), n>=1, k>=1, in which column k lists k's interleaved with k-1 zeros, and the first element of column k is in row k(k+1)/2.

Conjecture 1: T(n,k) is the number of parts in the partition of n into k consecutive parts, if T(n,k) > 0. Conjecture 2: row sums give A204217, which should be also the total number of parts in all partitions of n into consecutive parts. (The conjectures are true. See _Joerg Arndt_'s proof in the Links section.) - _Omar E. Pol_, Jun 14 2017 About the above theorem, this is the case for m = 1. For m = 0 see the triangle A127093, in which row sums give A000203. For m = 2 see the triangle A330466, in which row sums give A066839 (conjectured). For m = 3 see the triangle A330888, in which row sums give A330889.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Apr 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012138 | OEIS-A285929

Numbers m such that 2^m + (-1)^m is prime.

Conjecture: if (1 + y)^x + (-y)^x is a prime number then x is zero, or an even power of two, or an odd prime number. The above conjecture can be proved by considering algebraic factorizations of the polynomials involved. - _Jeppe Stig Nielsen_, Feb 19 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Apr 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012139 | OEIS-A285933

a(n) = smallest k such that (6*k-3)*2^n-1 and (6*k-3)*2^n+1 are twin primes.

Conjecture: a(n) is ~ (n*log(2))^2/9 as n increases.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Apr 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012140 | OEIS-A285945

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 123", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 02 2017: (Start) The conjectured generating function, closed forms, and equivalent linear recurrence are false; the first counterexample occurs at n = 28, where a(28) = 110000000011110, whereas the conjectured formulas give 110000000000000. - _Valentin Haase_, Jul 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012141 | OEIS-A285946

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 123", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 02 2017: (Start) The conjectured generating function, closed forms, and equivalent linear recurrence are false; the first counterexample occurs at n = 28, where a(28) = 111100000000110000000000000000, whereas the conjectured formulas give 1100000000000000. - _Valentin Haase_, Jul 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Apr 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012142 | OEIS-A285950

Positions of 0's in A285949; complement of A285951.

Conjecture: 3n/2 - a(n) is in {0, 1/2, 1} for all n >= 1. Proof of the conjecture by Kimberling: more is true. Here follows a proof of the formula below. Let T be the transform T(01)=0, T(1)=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012143 | OEIS-A285951

Positions of 1's in A285949; complement of A285950.

Conjecture: 3n - a(n) is in {0, 1} for all n >= 1. Proof of the conjecture by Kimberling: more is true. Here follows a proof of the formula below. Let T be the transform T(01)=0, T(1)=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012144 | OEIS-A285953

Positions of 0 in A285952; complement of A285954.

Conjecture: 3n - a(n) is in {0, 1} for n >= 1. Proof of the conjecture: Let t = A010060 be the Thue-Morse sequence. Every pair t(2n-1),t(2n) is either 01 or 10. Since 01 and 10 map to 110 and 101 under the transform, which both have length 3, it follows that a(n) = 3n-1+t(2n) for n=1,2,..., and so certainly 3n - a(n) is 0 or 1. - _Michel Dekking_, Jan 05 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012145 | OEIS-A285954

Positions of 1 in A285952; complement of A285953.

Conjecture: 3n/2 - a(n) is in {0, 1/2, 1} for n >= 1. Proof of the conjecture: Let t=A010060 be the Thue-Morse sequence. Every pair t(2n-1),t(2n) is either 01 or 10. Since 01 and 10 map to 110 and 101 under the transform, which both have length 3, it follows that a(2n+1) = 3n+1, and a(2n) = 3n if t(2n)=0, a(2n) = 3n-1 if t(2n)=1 for n=1,2,..., and so certainly 3n/2 - a(n) is 0, 1/2 or 1. - _Michel Dekking_, Jan 05 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012146 | OEIS-A285956

Number of orthogonal rectangles with vertices on an n X n square grid of points but with no vertices on the grid's diagonals.

Conjectures from _Colin Barker_, May 03 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rick L. Shepherd_, Apr 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012147 | OEIS-A285958

Positions of 0 in A285957; complement of A285959.

Conjecture: a(n)/n -> 4/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012148 | OEIS-A285959

Positions of 1 in A285957; complement of A285958.

Conjecture: a(n)/n -> 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012149 | OEIS-A285961

Positions of 0 in A285960; complement of A285962.

Conjecture: lim_{n->oo} a(n)/n = 4. Kimberling's conjecture is equivalent to the property that the frequency of 1's in A285960 is equal to 1/4. This can be computed via the comments of A285960. Another way to prove Kimberling's conjecture: a(n)/n = (1/n)*Sum_{k=1..n} d(k), the average value M of the differences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012150 | OEIS-A285962

Positions of 1 in A285960; complement of A285961.

Conjecture: a(n)/n -> 4/3. Kimberling's conjecture is equivalent to the property that the frequency of 1's in A285960 is equal to 3/4. This follows directly from the corresponding result in A285961.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012151 | OEIS-A285964

Positions of 0 in A285963; complement of A285965.

Conjecture: a(n)/n -> 4/5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012152 | OEIS-A285965

Positions of 1 in A285963; complement of A285964.

Conjecture: a(n)/n -> 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012153 | OEIS-A285967

Positions of 0 in A285966; complement of A285968.

Conjecture: a(n)/n -> 5/3. For a proof of this conjecture see A285968. - _Michel Dekking_, Mar 15 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012154 | OEIS-A285968

Positions of 1 in A285966; complement of A285967.

Conjecture: a(n)/n -> 5/2. Proof of this conjecture: it is equivalent to prove that the frequency of 1 in A285966 exists and equals 2/5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012155 | OEIS-A285970

Positions of 0 in A285969; complement of A285971.

Conjecture: a(n)/n -> 3/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012156 | OEIS-A285971

Positions of 0 in A285969; complement of A285970.

Conjecture: a(n)/n -> 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012157 | OEIS-A285972

{10->1}-transform of the Thue-Morse word A010060.

Conjecture: a(n) = A285960(n-1). - _R. J. Mathar_, May 08 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012158 | OEIS-A285973

Positions of 0 in A285972; complement of A285974.

Conjecture: a(n)/n -> 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012159 | OEIS-A285974

Positions of 1 in A285972; complement of A285973.

Conjecture: a(n)/n -> 4/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012160 | OEIS-A285976

Positions of 0 in A285975; complement of A285977.

Conjecture: a(n)/n -> 5/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012161 | OEIS-A285977

Positions of 1 in A285975; complement of A285976.

Conjecture: a(n)/n -> 5/3. This conjecture can be proved in the same way as for A284626, by showing that the frequency of the letter 1 in A285975 is equal to 3/5 - _Michel Dekking_, Sep 22 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012162 | OEIS-A285979

Positions of 0 in A285978; complement of A285980.

Conjecture: a(n)/n -> 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012163 | OEIS-A285980

Positions of 1 in A285978; complement of A285979.

Conjecture: a(n)/n -> 5/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012164 | OEIS-A286013

Irregular triangle read by rows: T(n,k), n>=1, k>=1, in which column k lists the positive integers starting with k, interleaved with k-1 zeros, and the first element of column k is in row k(k+1)/2.

Conjecture 1: T(n,k) is the largest part of the partition of n into k consecutive parts, if T(n,k) > 0. Conjecture 2: row sums give A286015. The truth of conjecture 1 follows, since each nonzero T(n,k) = k+m corresponds to a trapezoidal pattern of n dots having k rows with lengths (1+m)..(k+m).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012165 | OEIS-A286014

Sum of smallest parts of all partitions of n into consecutive parts.

Conjecture: this is also the row sums of A211343.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012166 | OEIS-A286015

Sum of largest parts of all partitions of n into consecutive parts.

Conjecture: this is also the row sums of A286013.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012167 | OEIS-A286031

Positions of 1 in A285568; complement of A285569.

Conjecture: a(n)/n -> (1+sqrt(5))/2 = golden ratio (A001622). Proof of this conjecture: the limit is 1/f0 where f0 is the frequency of 0 in (a(n)). Since the generating morphism of (a(n)) has the same incidence matrix as the generating morphism 0->11, 1->0101 of the sequence A285518, it follows that 1/f0 = phi = golden ratio. - _Michel Dekking_, Feb 22 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012168 | OEIS-A286040

Matching number of the n X n antelope graph.

Proof of the formula in the preceding comment and of the conjectured formulas below: We prove the more general result that the m X n antelope graph has a matching of size floor(m*n/2) when 20 <= m <= n, i.e., that all these graphs have a matching with at most one unmatched vertex. The proof is by induction on m+n, with the base cases 20 <= m <= n <= 39 checked with a computer program (I used Python with the function matching_number in the networkx module). Assume now that 20 <= m <= n, n >= 40 and that the statement is true for smaller values of m+n. We then know that the m X 20 antelope graph has a perfect matching (since m*20 is even, m >= 20, 20 >= 20, and m+20 < m+n). We also know that the m X (n-20) antelope graph has a matching with at most one unmatched vertex (since m >= 20, n-20 >= 20, and m+n-20 < m+n). Since these two graphs can be seen as disjoint induced subgraphs whose vertex sets partition the m X n antelope graph, these matchings can be combined into a matching of the m X n antelope graph (no edge going between the two subgraphs is included in the matching). Since the resulting matching has at most one unmatched vertex, we are done. - _Pontus von Brömssen_, Jul 16 2026 Conjectures from _Colin Barker_, May 04 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012169 | OEIS-A286043

(1/2)*A285658.

Conjecture: a(n)/n -> (5 + sqrt(17))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012170 | OEIS-A286044

{011->0}-transform of the Thue-Morse word A010060.

Conjecture: a(n) = b(b(n - 2)) - b(b(n - 3)) for n > 3, where b is A003160. - _Velin Yanev_, Feb 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012171 | OEIS-A286045

Positions of 0 in A286044; complement of A003157.

Conjecture: a(n)/n -> 4/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012172 | OEIS-A286047

Positions of 0 in A286046; complement of A286048.

Conjecture: 2n - a(n) is in {0,1} for n >= 1. Obviously Kimberling's conjecture is equivalent to the property that A286046 is a concatenation of the two 2-blocks 01 and 10. This can be read off immediately from the {A, B, C, D} composition of A286046 given in the comments of that sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012173 | OEIS-A286050

Positions of 0 in A286049; complement of A286051.

Conjecture: 2n - a(n) is in {0,1} for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012174 | OEIS-A286051

Positions of 1 in A286049; complement of A286050.

Conjecture: 2n - a(n) is in {0,1} for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012175 | OEIS-A286053

Positions of 0 in A286052; complement of A286054.

Conjecture: a(n)/n -> 4/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012176 | OEIS-A286054

Positions of 1 in A286052; complement of A286053.

Conjecture: a(n)/n -> 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012177 | OEIS-A286056

Positions of 0 in A286055; complement of A286057.

Conjecture: a(n)/n -> 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012178 | OEIS-A286057

Positions of 1 in A286055; complement of A286056.

Conjecture: a(n)/n -> 4/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012179 | OEIS-A286058

Positions of 1 in A285666; complement of A285667.

Conjecture: lim_{n->oo} a(n)/n = 3.38..., and if m denotes this number, then -1 < a(n)/n - m < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012180 | OEIS-A286059

Fixed point of the mapping 00->001, 1->011, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate of the mapping is given by A286062.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012181 | OEIS-A286060

Positions of 0 in A286059; complement of A286061.

Conjecture: lim_{n->oo} a(n)/n = 2.24..., and if m denotes this number, then -1 < a(n)/n - m < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012182 | OEIS-A286061

Positions of 1 in A286059; complement of A286060.

Conjecture: lim_{n->oo} a(n)/n = 1.80..., and if m denotes this number, then -2 < a(n)/n - m < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012183 | OEIS-A286062

a(n) = 2*a(n-1) + a(n-2) - a(n-3), where a(0) = 2, a(1) = 3, a(2) = 6.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->001, 1->011, starting with 00; see A286939.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012184 | OEIS-A286065

Positions of 0 in A286064; complement of A286066.

Conjecture: a(n)/n -> 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012185 | OEIS-A286066

Positions of 1 in A286064; complement of A286065.

Conjecture: a(n)/n -> 3/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012186 | OEIS-A286178

Denominator of Sum_{i=1..n}(A285388(i)*A285389(i+1))/(A285388(i+1)*A285389(i)).

Conjecture: the factorization of a(n) contains all primes 5 < p < 2*(n+1)^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Steiner_, May 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012187 | OEIS-A286182

Number of connected induced (non-null) subgraphs of the prism graph with 2n nodes.

Cases n=1 and n=2 correspond to degenerate prism graphs, but they fit the same (conjectured) linear recurrence as the other terms. a(n) = 6*a(n-1) - 11*a(n-2) + 4*a(n-3) + 5*a(n-4) - 2*a(n-5) - a(n-6), for n > 6 (conjectured). a(n) = A002203(n) + 3*n*A000129(n) - 3*n + 1 (conjectured). - _Eric W. Weisstein_, May 08 2017 G.f.: x*(3 - 5*x + 6*x^2 - 8*x^3 - 5*x^4 - 3*x^5) / ((1 - x)^2*(1 - 2*x - x^2)^2) (conjectured). - _Colin Barker_, May 31 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Resta_, May 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012188 | OEIS-A286183

Number of connected induced (non-null) subgraphs of the antiprism graph with 2n nodes.

a(n) = 8*a(n-1) - 24*a(n-2) + 34*a(n-3) - 24*a(n-4) + 8*a(n-5) - a(n-6), for n > 6 (conjectured). a(n) = A005248(n) - 2*n + 2*n*A001906(n) (conjectured). - _Eric W. Weisstein_, May 08 2017 G.f.: x*(3 - 9*x + 12*x^2 - 15*x^3 + 9*x^4 - 2*x^5) / ((1 - x)^2*(1 - 3*x + x^2)^2) (conjectured). - _Colin Barker_, May 30 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Resta_, May 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012189 | OEIS-A286185

Number of connected induced (non-null) subgraphs of the Möbius ladder graph with 2n nodes.

a(n) = 6*a(n-1) - 11*a(n-2) + 4*a(n-3) + 5*a(n-4) - 2*a(n-5) - a(n-6), for n>6 (conjectured). a(n) = 1/4*((1-sqrt(2))^n*(4-3*sqrt(2)*n) + (1+sqrt(2))^n*(4+3*sqrt(2)*n)) - 1 - n (conjectured). - _Eric W. Weisstein_, May 08 2017 a(n) = Lucas(n, 2) + 3*n*Fibonacci(n, 2) - n - 1, where Lucas(n, 2) = A002203(n) and Fibonacci(n, 2) = A000129(n) (conjectured). - _Eric W. Weisstein_, May 08 2017 G.f. (subject to the above conjectures. In fact all three conjectures are equivalent): (3*x-3*x^2-2*x^3-4*x^4+3*x^5-x^6)/(1-3*x+x^2+x^3)^2. - _Robert Israel_, May 08 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Resta_, May 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012190 | OEIS-A286282

Stage at which Ken Knowlton's elevator (version 2) reaches floor n for the first time.

This implies the following recurrence (conjectured by _N. J. A. Sloane_ on May 09 2017): a(1)=1, and for n>=1, a(n+1) = n*a(n) + n^2 - 3*n + 3. From the asymptotic expansion of b(n) (see A000522), we have a(n) ~ 2*(e-1)*(n-1)!. Conjecture: a(n) +(-n-2)*a(n-1) +3*(n-1)*a(n-2) +(-3*n+8)*a(n-3) +(n-4)*a(n-4)=0. - _R. J. Mathar_, May 21 2017 Conjecture: (n+1)*a(n) +(-n^2+3*n-27)*a(n-1) +3*(-n^2+10*n-13)*a(n-2) +(n-3)*(4*n-17)*a(n-3)=0. - _R. J. Mathar_, May 21 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012191 | OEIS-A286290

A bisection of A064736.

a(n) ~ n^2*(1 + 1.5/n^c) with c=1/2. (Conjectured, although for small n around 10^5 a smaller c ~ 0.478 is a better fit to the data.) - _M. F. Hasler_, May 23 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012192 | OEIS-A286297

Irregular triangle read by rows: successive rows have lengths 1,3,5,7,..., and are filled in across rows with the smallest nonnegative number such that there is no repeat in any row, column, or diagonal of slope +-1.

Conjecture: every column or diagonal of slope +-1 is a permutation of the nonnegative numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012193 | OEIS-A286302

Numbers n such that A133364(n) <= 1.

Conjecture: 10404 is the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, May 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012194 | OEIS-A286337

Positions of 0 in A286336; complement of A286338.

Conjecture: a(n)/n -> 3/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012195 | OEIS-A286338

Positions of 1 in A286336; complement of A286337.

Conjecture: a(n)/n -> 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012196 | OEIS-A286340

Positions of 0 in A286339; complement of A286341.

a(n) - a(n-1) is in {1,2,3} for n>=2. Conjecture: a(n)/n -> 9/5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012197 | OEIS-A286341

Positions of 1 in A286339; complement of A286340.

a(n) - a(n-1) is in {1,2,3} for n>=2. Conjecture: a(n)/n -> 9/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012198 | OEIS-A286343

For n>0, let b(n) = greatest index of n in any Fibonacci-like sequence containing n. This sequence is the ordinal transform of b.

Conjecturally, for any n>1, b(n) = A199088(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, May 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012199 | OEIS-A286348

Numbers m such that 4^m + (-3)^m is prime.

Conjecture: if (1 + y)^x + (-y)^x is a prime number then x is zero, or an even power of two, or an odd prime number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, May 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012200 | OEIS-A286387

a(n) = A002487(n^2).

Conjecture: For all i >= 0, j >= 0: A103391(1+i) = A103391(1+j) => a(i) = a(j). This would be an implication of observation made at A286377, which has been checked up to n=2048. See also A286378.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, May 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012201 | OEIS-A286389

a(0) = 0; a(n) = n - a(floor(a(n-1)/2)).

Conjecture: partial sums of A285431 (verified for n <= 400). - _Sean A. Irvine_, Jul 20 2022 The conjecture has been verified for n <= 50000. - _Michel Dekking_, Jul 06 2023 Irvine's conjecture is now proven using the Walnut theorem prover. - _Jeffrey Shallit_, Oct 21 2023 Conjecture: a(n) ~ c*n, where c = sqrt(3) - 1 = 0.732050807... This conjecture is implied by the conjecture in the COMMENTS, by a simple application of the Perron-Frobenius Theorem. The vector (1, 1 + sqrt(3)) is a right eigenvector of the incidence matrix of the morphism 0->11, 1->110. Therefore the frequency of 1 in A285431 is equal to sqrt(3) - 1. So if the conjecture in the COMMENTS is true, then this implies that a(n)/n converges to sqrt(3) - 1. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, May 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012202 | OEIS-A286401

Positions of 0 in A286400; complement of A286402.

a(n) - a(n-1) is in {1,2,3} for n>=2. Conjecture: a(n)/n -> 9/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012203 | OEIS-A286402

Positions of 1 in A286400; complement of A286401.

a(n) - a(n-1) is in {1,2,3} for n>=2. Conjecture: a(n)/n -> 9/5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012204 | OEIS-A286420

Positions of 0 in A286419; complement of A286421.

a(n) - a(n-1) is in {1,2,3,5,6} for n>=2. Conjecture: a(n)/n -> 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012205 | OEIS-A286421

Positions of 1 in A286419; complement of A286420.

a(n) - a(n-1) is in {1,2,3} for n>=2. Conjecture: a(n)/n -> 3/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012206 | OEIS-A286429

Highest elevation of an island above sea level in a number square.

Conjectures from _Colin Barker_, May 09 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Craig Knecht_, May 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012207 | OEIS-A286430

Least volume of water to surround the largest possible island in a number square.

Conjectures from _Colin Barker_, Jan 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Craig Knecht_, May 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012208 | OEIS-A286485

Positions of 0 in A286484; complement of A286486.

a(n) - a(n-1) is in {1,2,3} for n>=2. Conjecture: a(n)/n -> 9/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012209 | OEIS-A286486

Positions of 1 in A286484; complement of A286485.

a(n) - a(n-1) is in {1,2,3} for n>=2. Conjecture: a(n)/n -> 9/5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012210 | OEIS-A286488

Positions of 0 in A286487; complement of A286489.

a(n) - a(n-1) is in {1,2,3,5,6} for n>=2. Conjecture: a(n)/n -> 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012211 | OEIS-A286489

Positions of 1 in A286487; complement of A286488.

a(n) - a(n-1) is in {1,2,3} for n>=2. Conjecture: a(n)/n -> 3/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012212 | OEIS-A286491

Positions of 0 in A286490; complement of A286492.

a(n) - a(n-1) is in {1,2,3} for n>=2. Conjecture: a(n)/n -> 9/5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012213 | OEIS-A286492

Positions of 1 in A286490; complement of A286491.

a(n) - a(n-1) is in {1,2,3} for n>=2. Conjecture: a(n)/n -> 9/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012214 | OEIS-A286494

Positions of 0 in A286493; complement of A286495.

a(n) - a(n-1) is in {1,2,3} for n>=2. Conjecture: a(n)/n -> 3/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012215 | OEIS-A286495

Positions of 1 in A286493; complement of A286494.

a(n) - a(n-1) is in {1,2,3,5,6} for n>=2. Conjecture: a(n)/n -> 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012216 | OEIS-A286519

Binary representation of the diagonal from the origin to the corner (or of the corner to the origin) of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 659", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012217 | OEIS-A286521

Decimal representation of the diagonal from the origin to the corner (or of the corner to the origin) of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 659", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012218 | OEIS-A286529

a(n) = d(n+d(n)), where d(n) is the number of divisors of n (A000005).

Sum_{k=1..n} a(k) ~ D*n*log(n) + O(n*log(n)/log(log(n))), where D > 0 is a constant (conjectured with an error O(n) by Ivić, 1992; proven by Kátai, 2007). - _Amiram Eldar_, Jul 08 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, May 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012219 | OEIS-A286636

Even numbers that are a sum of two squares plus 1.

If the conjecture in A000952 is true, the two sequences are the same. - _R. J. Mathar_, May 18 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, May 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012220 | OEIS-A286666

Positions of 0 in A286665; complement of A286667.

a(n) - a(n-1) is in {2,3} for n>=2. Conjecture: a(n)/n -> 1 + sqrt(2). This conjecture follows easily from the fact that (a(n)) is a Beatty sequence, see my comments in A286665. - _Michel Dekking_, Mar 11 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012221 | OEIS-A286667

Positions of 1 in A286665; complement of A286666.

a(n) - a(n-1) is in {2,3} for n>=2. Conjecture: a(n)/n -> 1 + sqrt(1/2). This conjecture follows easily from the fact that (a(n)) is a Beatty sequence, see my comments in A286665. - _Michel Dekking_, Mar 11 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012222 | OEIS-A286746

{00->null}-transform of the infinite Fibonacci word A003849.

As a word, A003849 = 0100101001001010010101001001010010..., and deleting each occurrence of 00 gives 0110111011011101110110111011011011011..., in which, it is conjectured, the positions of 0 are given by A214971, and of 1, by A195121. Proof of the two conjectures by Kimberling: first note that the [00->null]-transform is the same as the [001->1]-transform, since 000 does not occur in the infinite Fibonacci word. Next, we do the following trick: replace the [001->1]-transform by the [001->2]-transform. Then the Fibonacci word is mapped to b:= A284749 = 0120122012... Note that the positions of 0 in b are the same as the positions of 0 in a. By Theorem 31 in the Allouche-Dekking paper, the positions of 0 in b are given by the sequence with terms floor(n*phi)+2*n+1, for n=0,1,2.... Transforming to offset 1, this is the sequence (ceiling((n-1)*phi) + 2*(n-1)), conjectured by Baruchel for A214971, and proved in my paper on base-phi representations. Proof of the conjecture by Mathar: this follows directly from Lemma 9.1.3 in the book by Allouche and Shallit. - _Michel Dekking_, Aug 16 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012223 | OEIS-A286758

Numbers k such that sigma(k) divides sigma(k!).

Conjecture: If p is Fermat prime > 3 from A019434 both values sigma((p-1)!) mod sigma(p-1) and sigma(T(p-1)) mod sigma(p-1) are not 0, where T(n) is the n-th triangular number A000217(n) and n! is the factorial number A000142(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, May 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012224 | OEIS-A286770

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 221", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012225 | OEIS-A286771

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 221", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012226 | OEIS-A286772

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 221", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012227 | OEIS-A286773

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 221", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012228 | OEIS-A286807

Fixed point of the mapping 00->001, 1->101, starting with 00.

Conjecture: The number of letters (0's and 1's) in the n-th iterate of the mapping is given by A132045.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012229 | OEIS-A286808

Positions of 0 in A286807; complement of A286809.

Conjecture: a(n)/n-> 2, and -1 < 2 - a(n)/n < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012230 | OEIS-A286809

Positions of 1 in A286807; complement of A286808.

Conjecture: a(n)/n -> 2, and -2 < 2 - a(n)/n < 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012231 | OEIS-A286843

Even k such that k - A001065(k) = 2^m (for some m > 0).

Tianxin Cai conjectured that the sequence is infinite. When p is prime, A001065(2p) = 1 + 2 + p = 3 + p. 2p - A001065(2p) = 2^m iff 2^m + 3 = p. Therefore if A057733 is infinite, Cai's conjecture is correct. In general, for j = 2, 3, ..., if the number of primes of the form 2^m + 2^j - 1 is infinite, then Cai's conjecture is correct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _XU Pingya_, Aug 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012232 | OEIS-A286848

Number of minimal dominating sets in the grid graph P_3 X P_n.

G.f.: x*(2 + 3*x - 2*x^2 - x^3 + 4*x^4 + 10*x^5 - 14*x^6 - 32*x^7 - 29*x^8 - 2*x^9 - 21*x^10 - 140*x^11 - 140*x^12 + 126*x^13 + 80*x^14 + 127*x^15 + 143*x^16 + 695*x^17 + 401*x^18 + 462*x^19 + 582*x^20 - 8*x^21 - 490*x^22 - 660*x^23 - 721*x^24 - 960*x^25 - 714*x^26 - 925*x^27 + 25*x^28 + 206*x^29 + 255*x^30 + 494*x^31 + 155*x^32 + 443*x^33 - 118*x^34 + 80*x^35 - 71*x^36 - 172*x^37 + 78*x^38 - 105*x^39 + 79*x^40 + 7*x^41 - 28*x^42 + 33*x^43 - 7*x^44 - 4*x^45 + 3*x^46 - x^47) / (1 - 2*x - 2*x^2 - 4*x^3 + 8*x^4 - 2*x^5 - 2*x^6 - 23*x^7 + 14*x^8 + 31*x^9 + 31*x^10 - 45*x^11 + 50*x^12 + 83*x^13 + 122*x^14 - 141*x^15 - 54*x^16 - 105*x^17 + 36*x^18 - 85*x^19 - 275*x^20 - 222*x^21 + 63*x^22 + 90*x^23 - 140*x^24 + 253*x^25 + 399*x^26 + 234*x^27 + 190*x^28 - 87*x^29 + 59*x^30 - 219*x^31 - 222*x^32 - 189*x^33 - 270*x^34 + 152*x^35 + 56*x^36 + 123*x^37 + 158*x^38 - 41*x^39 + 75*x^40 - 62*x^41 - 12*x^42 + 21*x^43 - 30*x^44 + 7*x^45 + 4*x^46 - 3*x^47 + x^48) (conjectured). - _Colin Barker_, Aug 02 2017 a(n) = Sum_{k=1..48} c(k)*a(n-k), where c = (2, 2, 4, -8, 2, 2, 23, -14, -31, -31, 45, -50, -83, -122, 141, 54, 105, -36, 85, 275, 222, -63, -90, 140, -253, -399, -234, -190, 87, -59, 219, 222, 189, 270, -152, -56, -123, -158, 41, -75, 62, 12, -21, 30, -7, -4, 3, -1) (conjectured). - _Eric W. Weisstein_, Aug 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Howroyd_, Aug 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012233 | OEIS-A286885

Number of ways to write 6*n+1 as x^2 + 3*y^2 + 54*z^2 with x,y,z nonnegative integers.

Conjecture: a(n) > 0 for all n = 0,1,2,....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012234 | OEIS-A286909

Positions of 1 in A286907; complement of A286908.

Conjectures: a(n) - a(n-1) is in {2,4} for n >= 2, and a(n)/n -> sqrt(8). Proof of the conjectures and the formula in FORMULA: a(n) = 2*A001951(n) for n >= 1, conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012235 | OEIS-A286938

Length of n-th iterate of the mapping 00->001, 1->10, as in A284932.

Conjecture: a(n) = 2*a(n-1) - a(n-2) + 2*a(n-3) -3*a(n-4) + a(n-5) for 5 >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012236 | OEIS-A286944

Number of ways to write n as x^2 + 15*y^2 + z*(3z+1)/2, where x and y are nonnegative integers and z is an integer.

Conjecture: a(n) > 0 for any nonnegative integer n, and a(n) = 1 only for n = 0, 3, 4, 7, 8, 10, 12, 13, 14, 29, 33, 34, 48, 68, 113, 129, 220. Let a,b,c,d,e,f be nonnegative integers with a > b, c > d, e > f, a == b (mod 2), c == d (mod 2), e == f (mod 2), a >= c >= e >= 2, b >= d if a = c, and d >= f if c = e. We have shown in arXiv:1502.03056 that if the ordered tuple (a,b,c,d,e,f) is universal, i.e., each n = 0,1,2,... can be written as x(ax+b)/2 + y(cy+d)/2 + z(ez+f)/2 with x,y,z integers, then (a,b,c,d,e,f) must be among the 12082 tuples listed in the linked a-file. We also conjecture that all the listed tuples not yet proved to be universal are indeed universal. Note that those numbers x(4x+2)/2 with x integral coincide with triangular numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012237 | OEIS-A286971

Number of ways to write n as a sum of two numbers, one of which is the product of an even number of distinct primes (including 1) (A030229) and another is the product of an odd number of distinct primes (A030059).

Conjecture: a(n) > 0 for all n > 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, May 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012238 | OEIS-A286981

Binomial coefficients binomial(n,k) = UV such that n>=2k and U > V, where gpf(U) <= k, gpf(V) > k (gpf(n)= is the greatest prime factor of n).

Ecklund et al. proved that the sequence is finite, and that these are the only terms, except for the cases k = 3, 5 and 7, but they strongly conjectured that the list is complete. They also give the near miss binomial(514,3)=22500864=UV, with U=2^9*3^2=4608, V=19*257=4883, and V/U < 1.06.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, May 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012239 | OEIS-A287076

a(n) = least k > n with the same sum of digits as n in some base b > 1.

Conjecturally, a(n) ~ n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, May 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012240 | OEIS-A287103

Positions of 0 in A288710; complement of A288712.

Conjecture: a(n)/n -> 1.39..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012241 | OEIS-A287104

Start with 0 and repeatedly substitute 0->10, 1->12, 2->0.

Conjecture: if n >=2, then u(n) - u(n-1) is in {2,3,4}, v(n) - v(n-1) is in {2,3}, and w(n) - w(n-1) is in {3,4,5}. See A287105, A287106, and A287107 for proofs of these conjectures, with explicit expressions for u, v, and w. - _Michel Dekking_, Sep 15 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012242 | OEIS-A287126

Positions of 0 in A287125; complement of A287127.

Conjecture: a(n)/n-> 2.22..., and if m denotes this number, then -2 < m - a(n)/n < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012243 | OEIS-A287127

Positions of 1 in A287125; complement of A287126.

Conjecture: a(n)/n-> 1.81..., and if m denotes this number, then -2 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012244 | OEIS-A287128

a(n) = 3*a(n-1) - 2*a(n-2) + 2*a(n-3) - 4*a(n-4) + 2*a(n-5), where a(0) = 2, a(1) =3, a(2) = 6, a(3)=13, a(4) = 29.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->001, 1->110, starting with 00; see A287125.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012245 | OEIS-A287129

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 245", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012246 | OEIS-A287130

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 245", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012247 | OEIS-A287131

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 245", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012248 | OEIS-A287132

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 245", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012249 | OEIS-A287164

Primes having a unique partition into three squares.

D. H. Lehmer conjectures that there are no more terms (see A094739 and A094942).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, May 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012250 | OEIS-A287183

Positions of 0 in A288106; complement of A287184.

Conjecture: a(n)/n-> 1.22..., and if m denotes this number, then -2 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012251 | OEIS-A287184

Positions of 1 in A288106; complement of A287183.

Conjecture: a(n)/n-> 5.42..., and if m denotes this number, then -1 < m - a(n)/n < 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012252 | OEIS-A287185

a(n) = 3*a(n-2) + a(n-3) + a(n-5) - 3*a(n-6) - a(n-7), where a(0) = 2, a(1) = 4, a(2) = 7, a(3) = 14, a(4) = 25, a(5) = 47, a(6) = 88, a(7) = 166.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0010, 1->00, starting with 00; see A288106.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012253 | OEIS-A287186

Positions of 1 in A287457; complement of A287519.

Conjecture: a(n)/n -> 5.90..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012254 | OEIS-A287190

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 253", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012255 | OEIS-A287191

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 253", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012256 | OEIS-A287192

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 253", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012257 | OEIS-A287193

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 253", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Nov 22 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012258 | OEIS-A287199

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 259", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, May 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012259 | OEIS-A287217

Primes p where A196202 reaches a new local minimum, i.e., the value of A196202 is strictly smaller than the value for both the previous prime and the next prime.

Conjecture: There are no consecutive Wieferich primes, i.e., A001220 is a subsequence of this sequence. The truth of the previous conjecture would follow from the truth of the heuristic assumption that there are about log(log(x)) Wieferich primes below x for large x.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, May 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012260 | OEIS-A287233

Numbers whose sum of proper divisors is equal to 88978489594.

Conjecture: there exists a positive real number k such that the number of preimages of an even number n is O((log n)^k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anton Mosunov_, May 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012261 | OEIS-A287238

Numbers whose sum of proper divisors is equal to 95186291194.

Conjecture: there exists a positive real number k > 1 such that the number of preimages of an even number n is O((log n)^k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anton Mosunov_, May 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012262 | OEIS-A287271

a(n) is the number of zeros of the Bernoulli B(n, x) polynomial in the open interval (-1, +1).

Conjecture 1: for n > 32, a(n) = 3 for n odd and a(n) = 4 otherwise. Conjecture 2: A284849(n) - a(n) > 0 if n = 6, 7,...,15 and n = 17. Conjecture 1 is false. It appears that for n => 13, a(n) = 3 for n == 1 (mod 4) and a(n) = 4 otherwise. - _Robert Israel_, May 29 2017 Conjectures from _Colin Barker_, Jan 22 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, May 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012263 | OEIS-A287299

Number of ways of writing n as a sum of a proper prime power (A246547) and a nonprime squarefree number (A000469).

Conjecture: a(n) > 0 for all n > 108.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, May 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012264 | OEIS-A287319

Smallest number k which becomes a power of 2 after being transformed by the reduced Collatz function k=(3*k+1)/2 precisely n times.

I conjecture that all members of a(n) are members of A054646 and A010120, "Smallest number to give 2^(2n) in a hailstone (3x + 1) sequence" and "Smallest start for a `3x+1' sequence containing 2^n".

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joe Slater_, May 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012265 | OEIS-A287372

0-limiting word of the mapping 00->1000, 10->000, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A287439(n), for n >= 0. Here is a proof of the conjecture. We note first that the mapping SR: 00->1000, 10->000, is an algorithmic procedure given by StringReplace in Mathematica. This makes it hard to describe iterates of it. However, in this particular case there is a way to deal with this. Let Thus the conjecture is proved by an application of the Cayley-Hamilton theorem:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012266 | OEIS-A287373

Positions of 0 in A101666.

Proof of these conjectures: Let b(n):=4n-a(n). Note first that a simple way to express the r,s,t property above is to say that the sequence b is obtained from the sequence x := A101666 = 0,1,1,2,1,2,1,0,1,2,1,0,1,..., by changing the alphabet from {0,1,2} to {3,0,1}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012267 | OEIS-A287379

Positions of 0 in A287931; complement of A287380.

Conjecture: a(n)/n-> 1.44..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012268 | OEIS-A287380

Positions of 1 in A287931; complement of A287379.

Conjecture: a(n)/n-> 3.24..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012269 | OEIS-A287381

a(n) = a(n-1) + 2*a(n-2) - a(n-3), where a(0) = 2, a(1) = 4, a(2) = 7.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0010, 1->10, starting with 00; see A287931.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012270 | OEIS-A287382

0-limiting word of the mapping 00->1000, 10->001, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A287525(n), for n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012271 | OEIS-A287402

Positions of 1 in A287372; complement of A287527.

Conjecture: a(n)/n -> 5.89..., as n -> infinity, and if m denotes this number, then -1 < m - a(n)/n <= m - 4 < 2 for n >= 1. Here is a proof of part of this conjecture. We recall from the comments of A287372 that A287372 = delta(x), where x is the fixed point of sigma^2 with x(1)=3. Here sigma is the morphism on {1,2,3} given by

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012272 | OEIS-A287439

a(n) = 2*a(n-2) + 2*a(n-3) for n >= 3, where a(0) = 2, a(2) = 4, a(3) = 7.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iterate of the mapping 00->1000, 10->000, starting with 00; see A288729.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012273 | OEIS-A287440

Positions of 0 in A287382; complement of A287448.

Conjecture: a(n)/n -> 1.45..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012274 | OEIS-A287448

Positions of 1 in A287382; complement of A287440.

Conjecture: a(n)/n -> 3.19..., and if m denotes this number, then -1 < m - a(n)/n < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012275 | OEIS-A287457

1-limiting word of the mapping 00->1000, 10->000, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A287439(n), for n >= 0. See comments of A287372 for a proof of this conjecture. - _Michel Dekking_, Mar 19 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012276 | OEIS-A287468

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 276", based on the 5-celled von Neumann neighborhood.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012277 | OEIS-A287469

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 276", based on the 5-celled von Neumann neighborhood.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012278 | OEIS-A287470

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 276", based on the 5-celled von Neumann neighborhood.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, May 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012279 | OEIS-A287519

Positions of 0 in A287457; complement of A287186.

Conjecture: a(n)/n -> 1.20..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012280 | OEIS-A287521

Positions of 0's in A287520.

a(n) - a(n-1) is in {1, 2, 3, 4, 5} for n >= 1; also, 3n - a(n) is in {0, 1,2} for n >= 1. The first 20 numbers 3n - a(n) are 2, 1, 0, 1, 2, 0, 1, 0, 2, 1, 2, 0, 2, 1, 0, 1, 0, 2, 1, 2, with 0's in positions given by A189630 (conjectured), 1's in positions given by A287522 (conjectured), and 2's in positions given by A287521 (conjectured). [Corrected by _Michel Dekking_, Sep 30 2019] Proof of the conjecture about the positions of 2's in A287520.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012281 | OEIS-A287525

a(n) = a(n-1) + a(n-2) - a(n-3) +a(n-4) + a(n-5) for n >= 6, where a(0) = 2, a(1) = 4, a(2) = 7, a(3) = 12, a(4) = 19, a(5) = 31.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iterate of the mapping 00->1000, 10->001, starting with 00; see A287372.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012282 | OEIS-A287527

Positions of 0 in A287372; complement of A287402.

Conjecture: a(n)/n -> 1.20..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012283 | OEIS-A287530

1-limiting word of the mapping 00->1000, 10->001, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A287525(n), for n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012284 | OEIS-A287531

Positions of 0 in A287530; complement of A287551.

Conjecture: a(n)/n -> 1.45..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012285 | OEIS-A287548

Triangle read by rows: T(n,k), where each row begins with the Catalan number for n nonintersecting arches and transitions through k generations of eliminating and reducing arch configurations to an end row entry equal to number of semi-meander solutions for n arches.

Conjectured: Conjecture: T(n,5) = C(n) - C(n-1) - 3*C(n-2) - 3*C(n-4). - _Hunter Hogan_, Oct 01 2026 Conjecture: T(n,6) = C(n) - C(n-1) - 4*C(n-2) + C(n-3) - 5*C(n-4) - 5*C(n-5). - _Hunter Hogan_, Oct 05 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger Ford_, May 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012286 | OEIS-A287551

Positions of 1 in A287530; complement of A287531.

Conjecture: a(n)/n -> 3.19..., and if m denotes this number, then -1 < m - a(n)/n < 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012287 | OEIS-A287581

Largest squarefree odd primitive abundant number with n prime factors.

a(n+1) = (a(n)/p(n))*p'(n)*q(n), where p(n) = gpf(a(n)), p'(n) = nextprime(p(n)+1), q(n) = precprime(1/(2/sigma[-1](a(n)/p(n)*p'(n))-1)), sigma[-1](x) = sigma(x)/x; conjectured to hold for all n >= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, May 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012288 | OEIS-A287614

Primes of the form (1 + x)^y + (-x)^y for some positive x, y.

Conjecture: If x is a positive number and (1 + x)^y + (-x)^y is an odd prime number, then y is other odd prime number or even power of two.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, May 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012289 | OEIS-A287639

Lexicographically earliest sequence of distinct positive terms such that the sum of two consecutive terms has Hamming weight >= 10.

Conjecturally, this is a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jun 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012290 | OEIS-A287664

Positions of 0's in A287663; complement of A287665.

Conjecture: -2 < n*r - a(n) < 1 for n >= 1, where r = (7 + sqrt(5))/6. [Corrected by _Clark Kimberling_, Aug 19 2019] Proof of Kimberling's conjecture. {} denotes fractional part. Since ({n*phi}) is equidistributed on (0,1), this implies that a(3*n+j-3) takes values in the interval (-(2/3)*phi, -(2/3)*phi+1) for j=1, in the interval (-(1/3)*phi, -(1/3)*phi+1) for j=2, and in the interval (0,1) for j=3. Combining these, we see that a(n) takes values in (-(2/3)*phi, 1), and that this is best possible. Since -(2/3)*phi = -1.078689..., this improves on the lower bound conjectured by Kimberling.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012291 | OEIS-A287665

Positions of 1's in A287663; complement of A287664.

Conjecture: 0 < n*r - a(n) < 3 for n >= 1, where r = (-1 + 3*sqrt(5))/2. [Corrected by _Clark Kimberling_, Aug 19 2019] Proof of Maiga's conjecture: let T denote the morphism {0->1, 1->000}. Proof of Kimberling's conjecture. a(n) = Sum_{k=0..n-1} [3*A001468(k)-2] (conjectured). - _Jon Maiga_, Dec 30 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012292 | OEIS-A287674

{0->1, 1->001}-transform of the infinite Fibonacci word A003849.

This will make it possible to prove the remarkable conjecture above by _R. J. Mathar_. Note that A286749 is a letter-to-letter projection of the SAME sequence x = 123411234... but by a different letter-to-letter map f:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012293 | OEIS-A287675

Positions of 0 in A287674; complement of A287676.

Conjecture: -1 < n*r - a(n) < 2 for n >= 1, where r = (7 + sqrt(5))/4. From this we derive directly Kimberling's conjecture. Note that The two intervals (1/2 - phi/2, 3/2 - phi/2) = (-0.3090..., 0.6090...) and (1,2) are subintervals of the interval (-1,2) in Kimberling's conjecture, and there are no better bounds, since the fractional part of n*phi is equidistributed modulo 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012294 | OEIS-A287676

Positions of 1 in A287674; complement of A287675.

Conjecture: -1 < n*r - a(n) < 2 for n >= 1, where r = 4 - sqrt(5). Conjecture: a(n) = 5*n + 2 - 2 * floor((n+1)*phi), where phi is the golden mean. - _Chunqing Liu_, Sep 27 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012295 | OEIS-A287683

5-tuples of practical numbers: numbers n such that n-6, n-2, n, n+2, n+6 are all practical numbers.

Melfi conjectured that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, May 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012296 | OEIS-A287723

Positions of 0 in A287722; complement of A287724.

Conjecture: -1 < n*r - a(n) < 1 for n >= 1, where r = (7 + sqrt(5))/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012297 | OEIS-A287724

Positions of 1 in A287722; complement of A287723.

Conjecture: 0 < n*r - a(n) < 3 for n >= 1, where r = 4 - sqrt(5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012298 | OEIS-A287726

Positions of 0 in A287725; complement of A287727.

Conjecture: 2 < n*r - a(n) < 3 for n >= 1, where r = (7 + sqrt(5))/2. This immediately implies Kimberling's conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012299 | OEIS-A287727

Positions of 1 in A287725; complement of A287726.

Conjecture: -1 < n*r - a(n) < 1 for n >= 1, where r = (15 - sqrt(5))/10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012300 | OEIS-A287770

Positions of 1 in A287769; complement of A276855.

Conjecture: 0 < n*r - a(n) < 1 for n >= 1, where r = (15 - sqrt(5))/10. It has been verified by computer that a(n) = floor(n*r) for n=1..3*10^6. This conjecture can be proved from the result in the Comments of A287769, where it is shown that A287769 is a Sturmian sequence with slope s := 1 - 1/(3+phi) = (15+sqrt(5))/22. The conjecture then follows from Lemma 9.1.3 in "Automatic sequences", since r = 1/s. - _Michel Dekking_, Oct 11 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012301 | OEIS-A287772

{0->1, 1->00}-transform of the infinite Fibonacci word A003849.

Conjecture: the positions of 1 are given by A050140. This has been checked for the first million terms. Proof of the conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012302 | OEIS-A287775

Positions of 0 in A287772; complement of A050140 (conjectured and proved).

Positions of 0 in A287772; complement of A050140 (conjectured and proved).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012303 | OEIS-A287877

Start with n and repeatedly apply the powertrain map x -> A133500(x); a(n) is the fixed point that is eventually reached, or -1 if the trajectory never reaches a fixed point.

It is conjectured that a fixed point is always reached. It is also conjectured that the only fixed points are the numbers 0 through 9, 2592, and 24547284284866560000000000 (see A135385).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012304 | OEIS-A287945

a(n) = largest prime q such that q | 2^p - 2 and p - 1 | q - 1, where p = prime(n).

First conjecture: a(n) > prime(n) for all n > 6. _Robert Israel_ tested the author's conjecture up to prime(95) = 499. The prime factorizations of the numbers 2^(p-1)-1 for larger p can be checked in available tables, see A005420. Second conjecture: a(n) = gpf(2^prime(n) - 2) for almost all n, in the sense that the set of exceptions {10, 16, 37, 40, ...} has zero natural density.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Sep 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012305 | OEIS-A288020

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 501", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jun 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012306 | OEIS-A288118

Lexicographically earliest sequence of distinct positive terms such that, for any i and j > 0, a(i*j) != a(i) * a(j).

Conjecturally:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jun 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012307 | OEIS-A288131

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 427", based on the 5-celled von Neumann neighborhood.

G.f.: (1 + 11*x - 89*x^2 + 9*x^3 - 1100*x^4 - 900*x^5 - 1100*x^6) / ((1 - x)*(1 - 10*x)*(1 + 10*x)*(1 + x + x^2)*(1 - 10*x^2)*(1 + 10*x^2)) (conjectured). - _Colin Barker_, Jun 06 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012308 | OEIS-A288132

Fixed point of the mapping 00->0010, 1->11, starting with 00.

This proves Kimberling's (corrected) conjecture in the MATHEMATICA program.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012309 | OEIS-A288133

Positions of 0 in A288132; complement of A288134.

This conjecture by Kimberling is proved in A288132. - _Michel Dekking_, Feb 18 2021 Conjectures from _Colin Barker_, Jun 09 2017: (Start) Colin Barker's conjectures are a consequence of

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012310 | OEIS-A288135

Coefficients of 1/(Sum_{k>=0} [(k+1)*r]*(-x)^k), where r = sqrt(7/3) and [ ] = floor.

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012311 | OEIS-A288136

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 427", based on the 5-celled von Neumann neighborhood.

G.f.: (1 + 11*x + 109*x^2 - 900*x^3 - 110*x^4 + 900*x^5 - 11000*x^6) / ((1 - x)*(1 + x)*(1 - 10*x)*(1 - 10*x^2)*(1 + 10*x^2)*(1 + 10*x + 100*x^2)) (conjectured). - _Colin Barker_, Jun 06 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012312 | OEIS-A288137

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 427", based on the 5-celled von Neumann neighborhood.

G.f.: (1 + 3*x - x^2 + x^3 - 12*x^4 - 4*x^5 - 12*x^6) / ((1 - x)*(1 - 2*x)*(1 + 2*x)*(1 + x + x^2)*(1 - 2*x^2)*(1 + 2*x^2)) (conjectured). - _Colin Barker_, Jun 06 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012313 | OEIS-A288138

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 427", based on the 5-celled von Neumann neighborhood.

G.f.: (1 + 3*x + 5*x^2 - 4*x^3 - 6*x^4 + 4*x^5 - 24*x^6) / ((1 - x)*(1 + x)*(1 - 2*x)*(1 - 2*x^2)*(1 + 2*x^2)*(1 + 2*x + 4*x^2)) (conjectured). - _Colin Barker_, Jun 06 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012314 | OEIS-A288139

Lexicographically earliest sequence of distinct positive terms such that the sum of two consecutive terms has Hamming weight <= 4.

Conjecturally, this is a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jun 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012315 | OEIS-A288164

Lexicographically earliest sequence of distinct positive terms such that, for any n > 0, a(n)*a(n+2) has at least 5 distinct prime factors.

Conjecturally, a(n) ~ n. - conjecturally, f_k(n) ~ n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012316 | OEIS-A288168

Positions of 0 in A288167; complement of A288169.

Conjecture: -1 < 6/5 - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012317 | OEIS-A288169

Positions of 1 in A288167; complement of A288168.

Conjecture: -1 < 6 - a(n)/n < 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012318 | OEIS-A288170

a(n) = 3*a(n-1) - a(n-2) - 4*a(n-3) + 2*a(n-4) for n >= 4, where a(0) = 2, a(1) = 4, a(2) = 8, a(3) = 16, a(4) = 34, a(5) = 70 .

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0010, 1->000, starting with 00; see A288167.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012319 | OEIS-A288171

Lexicographically earliest sequence of distinct positive terms such that, for any n > 0, a(n)*a(n+3) has at least 5 distinct prime factors.

Conjecturally, a(n) ~ n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012320 | OEIS-A288173

Fixed point of the mapping 00->0010, 1->001, starting with 00.

See A288176 for a proof of Kimberling's conjecture on the number of letters (0's and 1's) in the n-th iteration of the mapping SR. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012321 | OEIS-A288174

Positions of 0 in A288173; complement of A288175.

Conjecture: lim_{n->infinity} a(n)/n = 1.36..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1. Proof of the first part of this conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012322 | OEIS-A288175

Positions of 1 in A288173; complement of A288174.

Conjecture: lim_{n->infinity} a(n)/n = 3.70..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1. Proof of the first part of this conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012323 | OEIS-A288176

a(n) = 2*a(n-1) + 2*a(n-2) - 4*a(n-3) + a(n-4), where a(0) = 2, a(1) = 4, a(2) = 8, a(3) = 16.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0010, 1->001, starting with 00; see A288173. Proof of this conjecture. This is not the conjectured recursion a(n+4) = 2*a(n+3) +2* a(n+2) - 4*a(n+1) + a(n) for A288176. then we obtain the conjectured recursion. This proves the conjecture (where one uses that the constant sequence (1,1,1,...) satisfies the conjectured recursion). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012324 | OEIS-A288203

Fixed point of the mapping 00->0010, 1->010, starting with 00.

Here is a proof of the conjecture on the sequence of lengths 2, 4, 8, 18, 38, 80, ... of the iterates of the mapping (see also A288206). We note first that the mapping SR: 00->0010, 1->010, is an algorithmic procedure given by StringReplace in Mathematica. This makes it hard in general to describe iterates of it. However, here is an analysis of what happens in this particular case. Let This is not the conjectured recursion a(n+4) = 3*a(n+3) - a(n+2) - 3*a(n+1) + 2*a(n) from A288206. However, if we add this recursion to the shifted recursion a(n+5) = 3*a(n+4) - a(n+3) - 3*a(n+2) + 2*a(n+1), then we obtain the recursion above. This proves the conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012325 | OEIS-A288205

Positions of 1 in A288203; complement of A288204.

a(n) = Sum_{k=1..n} 4 - A036987(k) (conjecture). - _Jon Maiga_, Nov 25 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012326 | OEIS-A288206

a(n) = 3*a(n-1) - a(n-2) - 3*a(n-3) + 2*a(n-4), where a(0) = 2, a(1) = 4, a(2) = 8, a(3) = 18.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0010, 1->010, starting with 00; see A288203. See the Comments of A288203 for a proof of this conjecture. - _Michel Dekking_, Oct 12 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012327 | OEIS-A288209

Numbers k such that prime(k) * prime(k+1) mod prime(k+2) is odd.

Cramér's conjecture implies the sequence is finite. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Jun 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012328 | OEIS-A288213

Fixed point of the mapping 00->0010, 1->011, starting with 00.

Now that we know that this sequence is a morphic sequence, we may prove Kimberling's conjecture that the lengths of the words SR^n(00) are given by A182780. Thus the conjecture is proved by an application of the Cayley-Hamilton theorem: M^3 = 3M^2 - M - Id. a(n) = A001953(n-1) mod 2 (conjectured). - _Jon Maiga_, Sep 27 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012329 | OEIS-A288217

Positions of 0 in A288216; complement of A288218.

Conjecture: a(n)/n -> 1.38..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012330 | OEIS-A288218

Positions of 1 in A288216; complement of A288217.

Conjecture: a(n)/n -> 3.61..., and if m denotes this number, then 0 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012331 | OEIS-A288221

Positions of 0 in A288220; complement of A288222.

Conjecture: a(n)/n -> 1.38..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012332 | OEIS-A288222

Positions of 1 in A288220; complement of A288221.

Conjecture: a(n)/n -> 3.61..., and if m denotes this number, then 0 < m - a(n)/n < 0 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012333 | OEIS-A288224

Positions of 0 in A288223; complement of A288225.

Conjecture: a(n)/n -> 1.38..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012334 | OEIS-A288225

Positions of 1 in A288223; complement of A288224.

Conjecture: a(n)/n -> 3.61..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012335 | OEIS-A288229

Coefficients of 1/(Sum_{k>=0} [(k+1)*r]*(-x)^k), where r = Pi/2 and [ ] = floor.

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012336 | OEIS-A288230

Coefficients of 1/(Sum_{k>=0} [(k+1)*r]*(-x)^k), where r = sqrt(5/2) and [ ] = floor.

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012337 | OEIS-A288231

Coefficients of 1/(Sum_{k>=0} [(k+1)*r]*(-x)^k), where r = 4^(1/3) and [ ] = floor.

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012338 | OEIS-A288232

Coefficients in the expansion of 1/([r]-[2*r]*x+[3*r]*x^2-...); []=floor, r=3*e/5.

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012339 | OEIS-A288233

Coefficients in the expansion of 1/([r]-[2*r]*x+[3*r]*x^2-...); [ ]=floor, r=sqrt(8/3).

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012340 | OEIS-A288234

Coefficients in the expansion of 1/([r]-[2*r]*x+[3*r]*x^2-...); [ ]=floor, r=-1+sqrt(7).

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012341 | OEIS-A288235

Coefficients in the expansion of 1/([r]-[2*r]*x+[3*r]*x^2-...); [ ]=floor, r=sqrt(e).

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012342 | OEIS-A288236

Coefficients in the expansion of 1/([r]-[2*r]*x+[3*r]*x^2-...); [ ]=floor, r=-4/5+sqrt(6).

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012343 | OEIS-A288237

Coefficients in the expansion of 1/([r]-[2*r]*x+[3*r]*x^2-...); [ ]=floor, r=sqrt(11/4).

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012344 | OEIS-A288244

Numbers k such that prime(k)*prime(k+1)*prime(k+2) mod prime(k+3) is even.

Almost surely the above conjecture is true, but it is currently hopeless to prove. It would follow from prime gaps being less than about the cube root, but this is not known even under RH. - _Charles R Greathouse IV_, Jun 10 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Jun 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012345 | OEIS-A288247

2 * smallest possible area of a simple n-sided lattice polygon whose vertex coordinates x and y are both independent permutations of the integers 1 ... n, subject to the condition that none of its edges are mutually parallel.

It is conjectured that a(n) = n-2 for all n > 15, i.e. the bound of Pick's theorem is achievable for all larger n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012346 | OEIS-A288249

2 * largest possible area of a simple n-sided lattice polygon whose vertex coordinates x and y are both independent permutations of the integers 1 ... n, subject to the condition that none of its edges are mutually parallel.

It is conjectured that a(n) = 2*n^2 - 9*(n-1) for all n>22.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012347 | OEIS-A288250

Number of essentially different simple n-sided lattice polygons with maximum area subject to the conditions of A288247.

A conjectured continuation of the sequence starting at a(22) is 6, 3, 3, 6, 19, 36, 61.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012348 | OEIS-A288258

Positions of 0 in A288257; complement of A288259.

Conjecture: a(n)/n-> 1.36..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012349 | OEIS-A288259

Positions of 1 in A288257; complement of A288258.

Conjecture: a(n)/n-> 3.76..., and if m denotes this number, then -1 < m - a(n)/n < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012350 | OEIS-A288260

a(n) = 2*a(n-1) + 2*a(n-3) - 3*a(n-4), where a(0) = 2, a(1) = 4, a(2) = 8, a(3) = 16.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0010, 1->100, starting with 00; see A288257. [Mapping corrected by _Michel Dekking_, Mar 05 2020.] Second Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0101, 1->`010, starting with 00; see A288466. - _Michel Dekking_, Mar 05 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012351 | OEIS-A288307

Positions of 0 in A288306; complement of A288308.

Conjecture: -1 < 2 - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012352 | OEIS-A288308

Positions of 1 in A288306; complement of A288307.

Conjecture: -1 < 2 - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012353 | OEIS-A288309

a(n) = 2*a(n-1) + 2*a(n-2) - 3*a(n-3), where a(0) = 2, a(1) = 4, a(2) = 8.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0010, 1->110, starting with 00; see A288306.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012354 | OEIS-A288315

Positions of 0 in A288314; complement of A288316.

Conjecture: a(n)/n-> 1.22..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012355 | OEIS-A288316

Positions of 1 in A288314; complement of A288315.

Conjecture: a(n)/n-> 5.40..., and if m denotes this number, then -1 < m - a(n)/n < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012356 | OEIS-A288317

a(n) = a(n-1) + 3*a(n-2) - 2*a(n-3) - a(n-4) for n >= 4, where a(0) = 2, a(1) = 4, a(2) = 7, a(3) = 12.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0001, 1->00, starting with 00; see A288314.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012357 | OEIS-A288350

Lexically smallest string of n digits from 1...9, such that no formula using the single digits in the given order exists that evaluates to 0.

The last term a(7)=8985898, which is the only string of 7 digits for which it is not possible to construct an expression with result 0, is conjectured to be the last sequence term, i.e. for all strings of 8 or more digits expressions with result 0 can be found. The conjecture is almost trivially true. Since 8985898 is the only exception for strings of 7 digits, an 8 digit string containing 8985898 can only be of the form x8985898 or 8985898x. But since any of x898589 and 985898x /= 8985898 leads to a 7 digit string that can represent 0, one can always find a substring of length 7 representing 0 in a string of 8 or more digits. Therefore the sequence ends at a(7).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012358 | OEIS-A288351

Number of strings of n digits from 1...9 such that no formula using the single digits in the given order exists that evaluates to 0.

It is conjectured that a(n)=0 for n>=8. The conjecture is true, as shown in the corresponding comment in A288350. - _Hugo Pfoertner_, Jun 09 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012359 | OEIS-A288373

Positions of 0 in A288372; complement of A288316.

Conjecture: a(n)/n-> 1.44..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012360 | OEIS-A288374

Positions of 1 in A288372; complement of A288316.

Conjecture: a(n)/n-> 3.24..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012361 | OEIS-A288378

Positions of 0 in A288377; complement of A288379.

Conjecture: a(n)/n-> 1.46..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012362 | OEIS-A288379

Positions of 1 in A288377; complement of A288378.

Conjecture: a(n)/n-> 3.13..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012363 | OEIS-A288380

a(n) = 3*a(n-1) - 3*a(n-2) + 3*a(n-3) - 3*a(n-4) + a(n-5) for n >= 1, where a(0) = 2, a(1) = 4, a(2) = 7. a(3) = 11, a(4) = 20.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0001, 1->10, starting with 00; see A288377.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012364 | OEIS-A288381

Fixed point of the mapping 00->0001, 1->11, starting with 00.

Proof of Mathar's conjecture: obviously, a(2) = A288132(1), and we know that (a(n): n>1) is generated by a morphism mu (see below), and (A288132(n): n>1) is generated by a morphism nu (see A288132), given by nu(0)=0, nu(1)=11, nu(2)=210, both followed by replacing the letter 2 with the letter 0. Mathar's conjecture will therefore be a consequence of the following claim. Note that the shifting by one place occurs via the prefix 20 2^{-1} on the right side. This proves Kimberling's conjecture in the MATHEMATICA program. This proves Mathar's June 14, 2017 conjecture in A288382.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012365 | OEIS-A288382

Positions of 0 in A288381; complement of A288383.

This conjecture is proved in A288381. - _Michel Dekking_, Feb 18 2021 Conjectures from _Colin Barker_, Jun 10 2017: (Start) Barker's conjectures are implied by Mathar's conjecture. - _Michel Dekking_, Feb 18 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012366 | OEIS-A288424

Partial sums of A288384.

Conjecture: a(n) >= 0 for all n >= 0, and a(n) is unbounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jun 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012367 | OEIS-A288427

Positions of 0 in A288426; complement of A288428.

Conjecture: a(n)/n -> 1.93..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012368 | OEIS-A288428

Positions of 1 in A288426; complement of A288427.

Conjecture: a(n)/n->2.07..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012369 | OEIS-A288429

a(n) = 2*a(n-1) - a(n-2) + a(n-3) - a(n-4), where a(0) = 2, a(1) = 4, a(2) = 5, a(3) = 6.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0101, 1->100, starting with 00; see A288426.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012370 | OEIS-A288462

Fixed point of the mapping 00->0101, 1->10, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288465(n). This conjecture can be proved using the 17-state tribonacci DFAO generating the sequence. - _Jeffrey Shallit_, Oct 02 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012371 | OEIS-A288463

Positions of 0 in A288462; complement of A288464.

Conjecture: a(n)/n->1.83..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1. The constant 1.83... in the conjecture, psi, is the real zero of the polynomial x^3-x^2-x-1, and the conjectured inequality follows from the more precise statement -2 <= floor(psi*n) - a(n) <= 2; this can be proved with the Walnut theorem-prover. - _Jeffrey Shallit_, Oct 03 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012372 | OEIS-A288464

Positions of 1 in A288462; complement of A288463.

Conjecture: a(n)/n->2.19..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1. The constant 2.19... is gamma := (psi^2+1)/2, where psi is the real zero of x^3-x^2-x-1, and the conjectured inequality follows from the stronger statement that -1 <= a(n) - floor(gamma*n) <= 3; it can be proved with the Walnut theorem-prover. -_Jeffrey Shallit_, Oct 03 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012373 | OEIS-A288465

a(n) = 2*a(n-1) - a(n-4), where a(0) = 2, a(1) = 4, a(2) = 6, a(3) = 10.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0101, 1->10, starting with 00; see A288462. This conjecture has now been proved. -_Jeffrey Shallit_, Oct 03 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012374 | OEIS-A288466

Fixed point of the mapping 00->0101, 1->010, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288260(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012375 | OEIS-A288467

Positions of 0 in A288466; complement of A288468.

Conjecture: a(n)/n->1.56..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012376 | OEIS-A288468

Positions of 1 in A288466; complement of A288467.

Conjecture: lim_{n->inf} a(n)/n = 1.76..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012377 | OEIS-A288470

a(n) = Sum_{k=0..n} binomial(n,k)*binomial(2*n,2*k).

Conjecture: the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(2*k)) hold for primes p >= 5 and positive integers n and k. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Jun 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012378 | OEIS-A288473

1-limiting word of the mapping 00->0101, 1->011, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288476(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012379 | OEIS-A288474

Positions of 0 in A288473; complement of A288475.

Conjecture: a(n)/n - >2.34..., and if m denotes this number, then -1 < m - a(n)/n < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012380 | OEIS-A288475

Positions of 1 in A288473; complement of A288474.

Conjecture: a(n)/n -> 1.74..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012381 | OEIS-A288476

a(n) = a(n-1) + 4*a(n-2) - 2*a(n-3), where a(0) = 2, a(1) = 4, a(2) = 8.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0101, 1->011, starting with 00; see A288473.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012382 | OEIS-A288477

a(n) = (2^49 - 2)*n/3 + 247371098957.

Conjecture: a(0) + 1 = 247371098958 is the smallest nonnegative even number m such that for all k >= 1 the numbers m + 2^k + 1 and m + 2^k - 1 are composite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Jun 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012383 | OEIS-A288478

Fixed point of the mapping 00->0101, 1->110, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A182780(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012384 | OEIS-A288479

Positions of 0 in A288478; complement of A288480.

Conjecture: a(n)/n -> 1 + sqrt(2), and -1 < 1 + sqrt(2) - a(n)/n < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012385 | OEIS-A288480

Positions of 1 in A288478; complement of A288479.

Conjecture: a(n)/n -> 1 + sqrt(1/2), and -1 < 1 + sqrt(1/2) - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012386 | OEIS-A288508

0-limiting word of the mapping 00->0101, 10->001, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288511(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012387 | OEIS-A288509

Positions of 0 in A288508; complement of A288510.

Conjecture: a(n)/n -> 1.52..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012388 | OEIS-A288510

Positions of 1 in A288508; complement of A288509.

Conjecture: a(n)/n -> 2.91..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012389 | OEIS-A288511

a(n) = 2*a(n-1) - a(n-2) + 2*a(n-3) - 4*a(n-4) + 4*a(n-5) - 4*a(n-6) + 4*a(n-7) - 4*a(n-8) + 4*a(n-9) - 3*a(n-10) + 2*a(n-11) - 3*a(n-12) + 2*a(n-13) for n >= 16, with initial values as shown.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0101, 10->001, starting with 00; see A288508.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012390 | OEIS-A288512

1-limiting word of the mapping 00->0101, 10->001, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288511(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012391 | OEIS-A288513

Positions of 0 in A288512; complement of A288514.

Conjecture: a(n)/n -> 1.52..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012392 | OEIS-A288514

Positions of 1 in A288512; complement of A288513.

Conjecture: a(n)/n -> 2.91, and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012393 | OEIS-A288520

0-limiting word of the mapping 00->0101, 10->010, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288523(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012394 | OEIS-A288521

Positions of 0 in A288520; complement of A288522.

Conjecture: a(n)/n -> 1.52..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012395 | OEIS-A288522

Positions of 1 in A288520; complement of A288521.

Conjecture: a(n)/n -> 2.91..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012396 | OEIS-A288523

a(n) = a(n-1) + a(n-2) + a(n-3) - 2*a(n-4) for n >= 5, where a(0) = 2, a(1) = 4, a(2) = 5, a(3) = 8, a(4) = 11.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0101, 10->010, starting with 00; see A288520.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012397 | OEIS-A288524

1-limiting word of the mapping 00->0101, 10->010, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288523(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012398 | OEIS-A288525

Positions of 0 in A288524; complement of A288526.

Conjecture: a(n)/n -> 1.52..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012399 | OEIS-A288526

Positions of 1 in A288524; complement of A288525.

Conjecture: a(n)/n -> 2.91..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012400 | OEIS-A288530

Triangle read by rows in reverse order: T(n,k), (0 <= k <= n), in which each term is the least nonnegative integer such that no row, column, diagonal, or antidiagonal contains a repeated term.

Conjecture 3: every column is a permutation of the nonnegative integers. Conjecture 4: every diagonal is a permutation of the right border which gives the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jun 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012401 | OEIS-A288531

Triangle read by rows in reverse order: T(n,k), (1<=k<=n), in which each term is the least positive integer such that no row, column, diagonal, or antidiagonal contains a repeated term.

Conjecture 3: every column is a permutation of the positive integers. Conjecture 4: every diagonal is a permutation of the right border which gives the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jun 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012402 | OEIS-A288537

Array A(b,n) by upward antidiagonals (b>1, n>0): the eventual period of the RATS sequence in base b starting from n; 0 is for infinity.

Conway conjectured that in row (base) 10, all 0's correspond to the same divergent RATS sequence called the Creeper (A164338). In Thiel's terms, it is quasiperiodic with quasiperiod 2, i.e., after every 2 steps the number of one of the digits (in this case, 3 or 6) increases by 1 while other digits stay unchanged. In other bases, 0's may correspond to different divergent RATS sequences. Thiel conjectured that the divergent RATS sequences are always quasiperiodic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrey Zabolotskiy_, Jun 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012403 | OEIS-A288551

0-limiting word of the mapping 00->0101, 1->011, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288476(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012404 | OEIS-A288552

Positions of 0 in A288551; complement of A288553.

Conjecture: a(n)/n -> 2.34..., and if m denotes this number, then -1 < m - a(n)/n < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012405 | OEIS-A288553

Positions of 1 in A288551; complement of A288552.

Conjecture: a(n)/n -> 1.74..., and if m denotes this number, then -2 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012406 | OEIS-A288569

Start with n and repeatedly apply the map x -> A087207(x) until we reach 0; a(n) is the number of steps needed, or -1 if 0 is never reached.

There is a conjecture that 0 is always reached - see A087207. Conjecture 2: a(n) = 2 where n is 2^k, k>0. Conjecture 3: a(n) = 3 where n is an odd prime. Conjecture 4: a(n) = odd where n is an odd composite. (End) Conjecture 2 above follows from the facts that A087207(2^k) = 1 when k >= 1 and A087207(1) = 0. Conjecture 3 follows from Conjecture 2 and A087207(p) = 2^(primepi(p)-1). Provided that a(n) exists, Conjecture 4 follows from the fact that A087207(k) has opposite parity to k. - _Pontus von Brömssen_, Sep 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012407 | OEIS-A288597

Positions of 0 in A288596; complement of A288598.

Conjecture: a(n)/n -> 1.54..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012408 | OEIS-A288598

Positions of 1 in A288596; complement of A288597.

Conjecture: a(n)/n -> 2.83..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012409 | OEIS-A288599

a(n) = 2*a(n-1) - a(n-4) for n >= 4, where a(0) = 2, a(1) = 4, a(2) = 6, a(3) = 10, a(4) = 16.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0101, 10->1000, starting with 00; see A288596.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012410 | OEIS-A288600

Fixed point of the mapping 00->0101, 10->1001, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288603(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012411 | OEIS-A288601

Positions of 0 in A288600; complement of A288602.

Conjecture: a(n)/n -> 2, and -1 < 2 - a(n)/n < 1 for n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012412 | OEIS-A288602

Positions of 1 in A288600; complement of A288601.

Conjecture: a(n)/n -> 2, and -1/2 < 2 - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012413 | OEIS-A288603

a(n) = 2*a(n-1) - a(n-3) for n >= 3, where a(0) = 2, a(1) = 4, a(2) = 6, a(3) = 8.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iteration of the mapping 00->0101, 1->1001, starting with 00; see A288600.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012414 | OEIS-A288622

0-limiting word of the mapping 00->0110, 1->01, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288465(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012415 | OEIS-A288623

Positions of 0 in A288622; complement of A288624.

Conjecture: a(n)/n -> 1.83..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012416 | OEIS-A288624

Positions of 1 in A288622; complement of A288623.

Conjecture: a(n)/n -> 2.19..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012417 | OEIS-A288625

Positions of 0 in A288375; complement of A283794.

Conjecture: a(n)/n -> 2, and 0 < 2 - a(n)/n < 1/2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012418 | OEIS-A288633

Fixed point of the mapping 00->0110, 1->10, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288465(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012419 | OEIS-A288634

Positions of 0 in A288633; complement of A288635.

Conjecture: a(n)/n -> 1.83..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012420 | OEIS-A288635

Positions of 1 in A288633; complement of A288634.

Conjecture: a(n)/n -> 2.19..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012421 | OEIS-A288640

a(n) = (1 + sign(Im(ZetaZero(n)) - 2*Pi*e*exp(LambertW((n - 11/8)/e))))/2.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Jun 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012422 | OEIS-A288661

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 493", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jun 13 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012423 | OEIS-A288662

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 493", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jun 13 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012424 | OEIS-A288663

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 493", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jun 13 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012425 | OEIS-A288664

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 493", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jun 13 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012426 | OEIS-A288665

0-limiting word of the mapping 00->0110, 10->000, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288668(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012427 | OEIS-A288666

Positions of 0 in A288665; complement of A288667.

Conjecture: a(n)/n -> 1.41..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012428 | OEIS-A288667

Positions of 1 in A288665; complement of A288666.

Conjecture: a(n)/n -> 3.39..., and if m denotes this number, then -3 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012429 | OEIS-A288668

a(n) = a(n-2) + 2*a(n-3) for n >= 3, where a(0) = 2, a(2) = 4, a(3) = 5.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iterate of the mapping 00->0110, 10->000, starting with 00; see A288665.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012430 | OEIS-A288670

1-limiting word of the mapping 00->0110, 10->000, starting with 00.

The 1-limiting word is the limit of the n-th iterates for n == 1 mod 3. Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288468(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012431 | OEIS-A288671

Positions of 0 in A288670; complement of A288672.

Conjecture: a(n)/n -> 1.41..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012432 | OEIS-A288672

Positions of 1 in A288670; complement of A288671.

Conjecture: a(n)/n -> 3.39..., and if m denotes this number, then -1 < m - a(n)/n < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012433 | OEIS-A288673

2-limiting word of the mapping 00->0110, 10->000, starting with 00.

The 2-limiting word is the limit of the n-th iterates for n == 2 mod 3. Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288468(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012434 | OEIS-A288674

Positions of 0 in A288673; complement of A288675.

Conjecture: a(n)/n -> 1.41..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012435 | OEIS-A288675

Positions of 1 in A288673; complement of A288674.

Conjecture: a(n)/n -> 3.39..., and if m denotes this number, then -2 < m - a(n)/n < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012436 | OEIS-A288694

Fixed point of the mapping 00->0110, 10->100, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288429(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012437 | OEIS-A288695

Positions of 0 in A288694; complement of A288696.

Conjecture: a(n)/n -> 1.93..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012438 | OEIS-A288696

Positions of 1 in A288694; complement of A288695.

Conjecture: a(n)/n -> 2.07..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012439 | OEIS-A288708

Positions of 0 in A288707; complement of A288709.

Conjecture: a(n)/n -> -1 + sqrt(5), and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1. Proof of this conjecture. One follows the same strategy as in the proof of the conjecture in A288709. We have a(n)/n -> -1 + sqrt(5) if and only if n/a(n) -> (1+sqrt(5))/4, which is the frequency of 0's in A288707. If alpha = (3-sqrt(5))/2 is the frequency of 1's in y = A189661 = 0, 1, 0, 1, 0, 0, 1, 0, 1, 0, ... then

(2*(1-alpha)+alpha)/2 is the frequency of 0's in A288707, since every 0 in y produces 2 zeros in A288707, and every 1 just 1 zero (and one 1). One computes (2*(1-alpha)+alpha)/2 = (1+sqrt(5))/4. The bounds follow from the exponential convergence of the frequencies of 0 and 1 in y. - _Michel Dekking_, Oct 19 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012440 | OEIS-A288709

Positions of 1's in A288707; complement of A288708.

Conjecture: a(n)/n -> 3 + sqrt(5), and if m denotes this number, then -1 < m - a(n)/n < 3 for n >= 1. Here is a proof of this conjecture. Note that

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012441 | OEIS-A288710

0-limiting word of the mapping 00->1000, 10->0001, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288925(n), for n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012442 | OEIS-A288712

Positions of 0 in A288711; complement of A288713.

Conjecture: a(n)/n -> -1 + sqrt(5), and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012443 | OEIS-A288713

Positions of 1 in A288711; complement of A288712.

Conjecture: a(n)/n -> 3 + sqrt(5), and if m denotes this number, then -1 < m - a(n)/n < 5 for n >= 1. The truth of this conjecture follows directly from the same statement for A288709, in combination with Mathar's Jun 19 2017 conjecture. - _Michel Dekking_, Oct 20 2018 Here is a proof of this conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012444 | OEIS-A288729

0-limiting word of the mapping 00->1000, 10->01, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288732(n), for n >= 0. Here is a proof of the conjecture. We note first that the mapping

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012445 | OEIS-A288730

Positions of 0 in A288729; complement of A288731.

Conjecture: a(n)/n -> 2, and 0 < 2 - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012446 | OEIS-A288731

Positions of 1 in A288729; complement of A288730.

Conjecture: a(n)/n -> 2, and 0 < 2 - a(n)/n < 1/4 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012447 | OEIS-A288732

a(n) = a(n-1) + 2*a(n-4) - 2*a(n-5) for n >= 5, where a(0) = 2, a(1) = 4, a(2) = 6, a(3) = 8, a(4) = 10.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iterate of the mapping 00->1000, 10->01, starting with 00; see A288729.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012448 | OEIS-A288733

1-limiting word of the mapping 00->1000, 10->01, starting with 00.

The 1-limiting word is the limit of the n-th iterates for n == 1 mod 4. Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288732(n), for n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012449 | OEIS-A288734

Positions of 0 in A288733; complement of A288735.

Conjecture: a(n)/n -> 2, and 0 < 2 - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012450 | OEIS-A288735

Positions of 1 in A288733; complement of A288734.

Conjecture: a(n)/n -> 2, and 0 < 2 - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012451 | OEIS-A288736

2-limiting word of the mapping 00->1000, 10->01, starting with 00.

The 2-limiting word is the limit of the n-th iterates for n == 2 mod 4. Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288732(n), for n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012452 | OEIS-A288737

Positions of 0 in A288736; complement of A288740.

Conjecture: a(n)/n -> 2, and 0 < 2 - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012453 | OEIS-A288740

Positions of 1 in A288736; complement of A288737.

Conjecture: a(n)/n -> 2, and -2/3 < 2 - a(n)/n < 1/2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012454 | OEIS-A288741

3-limiting word of the mapping 00->1000, 10->01, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288732(n), for n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012455 | OEIS-A288742

Positions of 0 in A288741; complement of A285697.

Conjecture: a(n)/n -> 2, and 0 < 2 - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012456 | OEIS-A288753

Positions of 0 in A288752; complement of A288754.

Conjecture: a(n)/n -> 1.83..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012457 | OEIS-A288754

Positions of 1 in A288752; complement of A288753.

Conjecture: a(n)/n -> 2.19..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012458 | OEIS-A288755

Positions of 1 in A288710; complement of A287103.

Conjecture: a(n)/n -> 3.53..., and if m denotes this number, then -1 < m - a(n)/n < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012459 | OEIS-A288758

Floor of the volume of the "monic slice" of the d-th Chern-Vaaler star body.

Unimodal: increases from 2 to a maximum of 8, then decreases to 0 = a(d) for d >= 15 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012460 | OEIS-A288760

Number of distinct nonnegative rational numbers that can be obtained in n steps by applying addition, subtraction, multiplication and division to any two potentially identical numbers from the complete set of numbers created in n-1 steps, starting with the set {1}.

The conjectured value of a(6)=451572162 needs independent verification.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012461 | OEIS-A288761

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 501", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jun 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012462 | OEIS-A288762

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 501", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jun 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012463 | OEIS-A288763

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 501", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jun 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012464 | OEIS-A288798

Lexicographically earliest sequence of distinct positive terms such that the absolute difference of two consecutive terms has at least 5 distinct prime factors.

Conjecturally: this sequence is a permutation of the natural numbers, and a(n) ~ n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jun 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012465 | OEIS-A288828

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 515", based on the 5-celled von Neumann neighborhood.

a(n) = 3*a(n-1) - 6*a(n-3) + 4*a(n-4) for n > 5 (conjectured). G.f.: (8*x^5 - 8*x^4 - 2*x^3 + 4*x^2 - 1)/((x - 1)*(2*x - 1)*(2*x^2 - 1)) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012466 | OEIS-A288907

Primes p whose distance from the next prime and from the previous prime is less than log(p).

Conjecture: Limit_{n->oo} n / PrimePi(a(n)) = (1-1/e)^2 (A068996). - _Alain Rocchelli_, May 07 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giuseppe Coppoletta_, Jun 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012467 | OEIS-A288921

Lexicographically earliest sequence of distinct positive terms such that the product of two consecutive terms has at least 34 divisors.

Conjecturally, a(n) ~ n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jun 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012468 | OEIS-A288923

Lexicographically earliest sequence of distinct positive terms such that the product of two consecutive terms has at least 6 prime factors (counted with multiplicity).

Conjecturally, a(n) ~ n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jun 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012469 | OEIS-A288925

a(n) = a(n-1) + a(n-2) + 3*a(n-4) - 2*a(n-5) for n >= 5, where a(0) = 2, a(1) = 4, a(2) = 8, a(3) = 13, a(4) = 26.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iterate of the mapping 00->1000, 10->0001, starting with 00; see A288226.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012470 | OEIS-A288926

1-limiting word of the mapping 00->1000, 10->0001, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate is given by A288925(n), for n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012471 | OEIS-A288927

Positions of 0 in A288926; complement of A288928.

Conjecture: a(n)/n -> 1.39..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012472 | OEIS-A288928

Positions of 1 in A288926; complement of A288927.

Conjecture: a(n)/n -> 3.52..., and if m denotes this number, then -1 < m - a(n)/n < 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012473 | OEIS-A288929

Fixed point of the mapping 00->1000, 10->10011, starting with 00.

Conjecture: The number of letters (0's and 1's) in the n-th iterate of the mapping is given by A192967(n+2) for n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012474 | OEIS-A288930

Positions of 0 in A288929; complement of A288931.

Conjecture: a(n)/n -> 3, and 0 < m - a(n)/n < 3/2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012475 | OEIS-A288931

Positions of 1 in A288929; complement of A288930.

Conjecture: a(n)/n -> 3/2, and -1/2 < m - a(n)/n < 3/2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012476 | OEIS-A288932

Fixed point of the mapping 00->1000, 10->10101, starting with 00.

Conjecture: The number of letters (0's and 1's) in the n-th iterate of the mapping is given by A123720.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012477 | OEIS-A288933

Positions of 0 in A288932; complement of A288934.

Conjecture: a(n)/n -> 3, and 0 < 3 - a(n)/n < 1 for n >= 1. a(n) = 3*n-1-HammingWeight(n-1) = 3*n-1-A000120(n-1) (conjectured). - _Gleb Koloskov_, May 06 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012478 | OEIS-A288934

Positions of 1 in A288932; complement of A288933.

Conjecture: a(n)/n -> 3/2, and -2/3 < 3/2 - a(n)/n < 1/2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012479 | OEIS-A288936

Fixed point of the mapping 00->0010, 01->011, 10->011, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate of the mapping is given by A000124. Proof of the conjecture: let N0(n) be the number of 0's, and N1(n) be the number of 1's in the n-th iterate of the StringReplace mapping defining (a(n)). This proves the conjecture:

N0(n)+N1(n) = (n^2 +3n+4)/2 = A000124(n+1) for n=1,2,.... (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012480 | OEIS-A288992

Positions of 0 in A288991; complement of A288993.

Conjecture: a(n)/n -> 1.28... and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012481 | OEIS-A288993

Positions of 1 in A288991; complement of A288992.

Conjecture: a(n)/n -> 4.56... and if m denotes this number, then -1 < m - a(n)/n < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012482 | OEIS-A288997

Fixed point of the mapping 00->0010, 01->001, 10->001, starting with 00.

Conjecture: a(n) = A171588(n+3) for all n>0. - _Michel Dekking_, Mar 21 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012483 | OEIS-A288998

Positions of 0 in A288997; complement of A288999.

Conjecture: a(n)/n -> sqrt(2), and 0 < sqrt(2) - a(n)/n < -1 + sqrt(2) for n >= 1. a(n) = round((n-1/2)*sqrt(2)) (conjectured). - _Chai Wah Wu_, Feb 06 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012484 | OEIS-A288999

Positions of 1 in A288997; complement of A288998.

Conjecture: a(n)/n -> 2 + sqrt(2), and -1 < 2 + sqrt(2) - a(n)/n < -1 + sqrt(2) for n >= 1. a(n) = 2n+round(n*sqrt(2)) (conjectured). - _Chai Wah Wu_, Feb 06 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012485 | OEIS-A289001

Fixed point of the mapping 00->0010, 01->001, 10->010, starting with 00.

This coincides with the Pell word A171588, the fixed point of the morphism 0->001, 1->0. This was conjectured by _R. J. Mathar_, Jul 07 2017 and proved by Dekking and Keane in 2022. As observed by _Michel Dekking_, Mar 09 2022, this also proves the two conjectures about the positions of 0's and 1's stated in the MATHEMATICA section. - _N. J. A. Sloane_, Mar 09 2022 Conjecture: the number of letters (0's and 1's) in the n-th iterate of the mapping is given by A289004.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012486 | OEIS-A289002

Positions of 0 in A289235; complement of A289151.

Conjecture: a(n)/n -> 1.28..., and if m denotes this number, then -1 < m - a(n)/n < m - 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012487 | OEIS-A289004

a(n) = 2*a(n-1) - a(n-2) + 2*a(n-4) - a(n-5) + a(n-7) - a(n-8) - a(n-10) for n >= 10, where a(0) = 2, a(1) = 4, a(2) = 7, a(3) = 11, a(4) = 18, a(5) = 31, a(6) = 52, a(7) = 89, a(9) = 151, a(9) = 257.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iterate of the mapping 00->0010, 01->001, 10->010, starting with 00; see A289001.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012488 | OEIS-A289008

Positions of 0 in A289007; complement of A289009.

Conjecture: a(n)/n -> 2.37..., and if m denotes this number, then 0 < m - a(n)/n < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012489 | OEIS-A289009

Positions of 1 in A289007; complement of A289008.

Conjecture: a(n)/n -> 1.75..., and if m denotes this number, then -2 < m - a(n)/n < 0 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012490 | OEIS-A289012

Positions of 0 in A289011; complement of A289013.

Conjecture: a(n)/n -> sqrt(2), and -1 < sqrt(2) - a(n)/n < -1 + sqrt(2) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012491 | OEIS-A289013

Positions of 1 in A289011; complement of A289012.

Conjecture: a(n)/n -> 2 + sqrt(2), and 0 < 2 + sqrt(2) - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012492 | OEIS-A289017

Positions of 0 in A289016; complement of A289018.

Conjecture: a(n)/n -> 1.28..., and if m denotes this number, then -1 < m - a(n)/n < -1 + m for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012493 | OEIS-A289018

Positions of 1 in A289016; complement of A289017.

Conjecture: a(n)/n -> 4.55..., and if m denotes this numbrer, then -1 < m - a(n)/n < -3 + m for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012494 | OEIS-A289026

Positions of 0 in A289025; complement of A289027.

Conjecture: a(n)/n->sqrt(2), and -1 < sqrt(2)-a(n)/n < -1 + sqrt(2) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012495 | OEIS-A289027

Positions of 1 in A289025; complement of A289026.

Conjecture: a(n)/n -> 2 + sqrt(2), and -1 < 2 + sqrt(2) - a(n)/n < -1 + sqrt(2) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012496 | OEIS-A289035

Fixed point of the mapping 00->0010, 01->010, 10->010, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate of the mapping is given by A289004. Conjecture: a(n) = A171588(n-3) for n>3. - _Michel Dekking_, Mar 20 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012497 | OEIS-A289036

Positions of 0 in A289035; complement of A289037.

Conjecture: a(n)/n->sqrt(2), and -1 < sqrt(2) - a(n)/n < -1 + sqrt(2) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012498 | OEIS-A289037

Positions of 1 in A289035; complement of A289036.

Conjecture: a(n)/n -> 2 + sqrt(2), and 0 < 2 + sqrt(2) - a(n)/n < -1 + sqrt(2) for n >= 1. Proof of (a part of) Kimberling's conjecture. This is based on the formula for the recursion in the FORMULA section of A293077, where we use that the final-letter-removed version of the String-Replace map has the same fixed point A289035. Note: The full conjecture of Kimberling would follow from the conjecture A289035(n) = A171588(n-3) for n>3. (End) Conjecture: a(n+1) - a(n) = A276864(n) for all n. - _Michel Dekking_, Mar 20 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012499 | OEIS-A289057

Fixed point of the mapping 00->0010, 01->010, 10->011, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate of the mapping is given by A289060.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012500 | OEIS-A289060

a(n) = 3*a(n-1) - 3*a(n-2) + *a(n-3) for n >= 8, where a(0) = 2, a(1) = 4, a(2) = 7, a(3) = 11, a(4) = 17, a(5) = 25, a(6) = 36, a(7) = 51.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iterate of the mapping 00->0010, 01->010, 10->011, starting with 00; see A289057.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012501 | OEIS-A289072

Positions of 0 in A289071; complement of A289073.

Conjecture: a(n)/n->sqrt(2), and -1 < sqrt(2) - a(n)/n < -1 + sqrt(2) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012502 | OEIS-A289073

Positions of 1 in A289071; complement of A289072.

Conjecture: a(n)/n -> 2 + sqrt(2), and 0 < 2 + sqrt(2) - a(n)/n < 1 for n >= 1. .

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012503 | OEIS-A289075

Positions of 0 in A289074; complement of A289076.

Conjecture: a(n)/n -> 1.94..., and if m denotes this number, then -1 < m - a(n)/n < -1 + m for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012504 | OEIS-A289076

Positions of 1 in A289074; complement of A289075.

Conjecture: a(n)/n -> 2.05..., and if m denotes this number, then -1 < m - a(n)/n < 0 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012505 | OEIS-A289077

a(n) = 2*a(n-1) + a(n-2) - 4*a(n-3) + 2*a(n-4) + a(n-5) -2*a(n-6) + a(n-7) for n >= 7, a(0) = 2, a(1) = 4, a(2) = 7, a(3) = 11, a(4) = 18, a(5) = 30, a(6) = 47.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iterate of the mapping 00->0010, 01->011, 10->100, starting with 00; see A289057.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012506 | OEIS-A289096

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 545", based on the 5-celled von Neumann neighborhood.

a(n) = 10*a(n-1) + a(n-2) - 10*a(n-3) for n > 5 (conjectured). G.f.: (100*x^5 - 10*x^4 + 10*x^2 + 1)/((x - 1)*(x + 1)*(10*x - 1)) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012507 | OEIS-A289097

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 545", based on the 5-celled von Neumann neighborhood.

a(n) = a(n-1) + 100*a(n-2) - 100*a(n-3) for n > 5 (conjectured). G.f.: (1000*x^5 - 1000*x^4 + 10*x^2 + 1)/((x - 1)*(10*x - 1)*(10*x + 1)) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012508 | OEIS-A289098

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 545", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Jun 13 2025: (Start) Conjectured e.g.f.: 2 + x^2 - 3*cosh(x) + 2*cosh(2*x) - 2*sinh(x) + 2*sinh(2*x). - _Stefano Spezia_, Jun 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012509 | OEIS-A289099

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 545", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Jun 13 2025: (Start) Conjectured e.g.f.: x^2 + (1 + 3*cosh(2*x))/2 - cosh(x) - sinh(x) + sinh(2*x). - _Stefano Spezia_, Jun 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jun 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012510 | OEIS-A289104

Fixed point of the mapping 00->0010, 01->011, 10->000, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate of the mapping is given by A289107. For connections to the golden ratio, see A289105 and A289106.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012511 | OEIS-A289105

Positions of 0 in A289104; complement of A289106.

Conjecture: a(n)/n -> (1 + sqrt(5))/2 = golden ratio = phi, and 1 < a(n)/n < phi for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012512 | OEIS-A289106

Positions of 1 in A289104; complement of A289105.

Conjecture: a(n)/n -> (3 + sqrt(5))/2 = 1 + golden ratio = 1 + phi, and 1 + phi < a(n)/n < 4 for n >= 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012513 | OEIS-A289107

a(n) = 2*a(n-1) - a(n-3) for n >= 5, a(0) = 2, a(1) = 4, a(2) = 7, a(3) = 12, a(4) = 22.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iterate of the mapping 00->0010, 01->011, 10->000, starting with 00; see A289104.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012514 | OEIS-A289111

a(n) = (2^49 - 2)*n/3 + 444813635231.

Conjecture: a(0) + 1 = 444813635232 is the smallest nonnegative even number m such that for all k >= 1 the absolute values of the numbers m - 2^k + 1 and m - 2^k - 1 are composite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Jun 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012515 | OEIS-A289112

Fixed point of the mapping 00->0010, 01->011, 10->001, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate of the mapping is given by A289115.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012516 | OEIS-A289113

Positions of 0 in A289112; complement of A289114.

Conjecture: a(n)/n -> 2.14..., and if m denotes this number, then 0 < a(n)/n < m - 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012517 | OEIS-A289114

Positions of 1 in A289112; complement of A289113.

Conjecture: a(n)/n -> 1.87..., and if m denotes this number, then -2 < a(n)/n < 0 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012518 | OEIS-A289115

a(n) = 3*a(n-1) - 3*a(n-2) + 2*a(n-3) - 2*a(n-4) + a(n-5) for n >= 10, a(0) = 2, a(1) = 4, a(2) = 7, a(3) = 12, a(4) = 20, a(5) = 32, a(6) = 50, a(7) = 77, a(8) = 116, a(9) = 174.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iterate of the mapping 00->0010, 01->011, 10->001, starting with 00; see A289112.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012519 | OEIS-A289118

Least prime beginning a string, of length at least n, of consecutive primes which alternate between types 4*k+1 and 4*k+3 or 4*k+3 and 4*k+1.

Conjecture: the sequence is infinite. (Motivation: the string HTHTHT. . of length n eventually occurs in any sufficiently long sequence of coin tosses.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Jun 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012520 | OEIS-A289128

Fixed point of the mapping 00->0010, 01->011, 10->010, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate of the mapping is given by A289131.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012521 | OEIS-A289129

Positions of 0 in A289128; complement of A289130.

Conjecture: a(n)/n -> 5/2..., and 0 < 5/2 - a(n)/n < 3/2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012522 | OEIS-A289130

Positions of 1 in A289128; complement of A289129.

Conjecture: a(n)/n -> 5/3..., and -4/3 < 5/3 - a(n)/n < 0 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012523 | OEIS-A289131

a(n) = a(n-1) + 3*a(n-2) - 3*a(n-3) - 2*a(n-4) + 2*a(n-5) for n >= 6, a(0) = 2, a(1) = 4, a(2) = 7, a(3) = 11, a(4) = 18, a(5) = 28.

Conjecture: a(n) is the number of letters (0's and 1's) in the n-th iterate of the mapping 00->0010, 01->011, 10->010, starting with 00; see A289128.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012524 | OEIS-A289150

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 597", based on the 5-celled von Neumann neighborhood.

G.f.: (1 - x + x^2 - 10*x^4) / ((1 - x)*(1 - 10*x)*(1 + x^2)*(1 - 10*x^2)) (conjectured). - _Colin Barker_, Jul 14 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012525 | OEIS-A289151

Positions of 1 in A289235; complement of A289002.

Conjecture: a(n)/n -> 1.56..., and if m denotes this number, then -1 < m - a(n)/n < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012526 | OEIS-A289166

Positions of 0 in A289165; complement of A289167.

Conjecture: a(n)/n -> 1.42..., and if m denotes this number, then -1 < m - a(n)/n < m - 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012527 | OEIS-A289167

Positions of 1 in A289165; complement of A289166.

Conjecture: a(n)/n -> 3.35..., and if m denotes this number, then -1 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012528 | OEIS-A289174

Fixed point of the mapping 00->0010, 01->100, 10->011, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate of the mapping is given by A289177.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012529 | OEIS-A289175

Positions of 0 in A289174; complement of A289176.

Conjecture: a(n)/n -> 2.2..., and if m denotes this number, then -1 < m - a(n)/n < m - 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012530 | OEIS-A289176

Positions of 1 in A289174; complement of A289175.

Conjecture: a(n)/n -> 1.8..., and if m denotes this number, then -2 < m - a(n)/n < 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012531 | OEIS-A289194

Lexicographically earliest sequence of distinct positive terms such that the product of two consecutive terms has no isolated 1 in its base-2 representation.

This sequence is conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jun 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012532 | OEIS-A289239

Fixed point of the mapping 00->0010, 01->100, 10->010, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate of the mapping is given by A289004.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012533 | OEIS-A289240

Positions of 0 in A289239; complement of A289241.

Conjecture: a(n)/n -> sqrt(2)..., and -1 < sqrt(2)-a(n)/n < = -1+sqrt(2) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012534 | OEIS-A289241

Positions of 1 in A289239; complement of A289240.

Conjecture: a(n)/n -> 2+sqrt(2)..., and 0 < 2+sqrt(2)-a(n)/n < = 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012535 | OEIS-A289242

Fixed point of the mapping 00->0010, 01->100, 10->100, starting with 00.

Conjecture: the number of letters (0's and 1's) in the n-th iterate of the mapping is given by A289004.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012536 | OEIS-A289243

Positions of 0 in A289242; complement of A289244.

Conjecture: a(n)/n -> sqrt(2), and -1 < sqrt(2) - a(n)/n < = -1 + sqrt(2) n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012537 | OEIS-A289244

Positions of 1 in A289242; complement of A289243.

Conjecture: a(n)/n -> a(n)/n -> 2+sqrt(2), and 0 < 2+sqrt(2)-a(n)/n < = 1 n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012538 | OEIS-A289245

Coefficients of 1/(Sum_{k>=0} [-1 + (k+1)*r](-x)^k), where r = (3 + sqrt(5))/2 = 1 + golden ratio and [ ] = floor.

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012539 | OEIS-A289246

Coefficients in the expansion of 1/Sum_{k >= 0} ([r*(k + 1)] + [s*(k + 1)]) * (-x)^k, where [ ] = floor, r = (1+sqrt(5))/2, s = 1/r.

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012540 | OEIS-A289258

Prime factors of numbers in A289257.

Kalmynin conjectures that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Jun 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012541 | OEIS-A289260

Coefficients in the expansion of 1/([r]-[2r]x+[3r]x^2-...); [ ]=floor, r=8/5.

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012542 | OEIS-A289261

Coefficients in the expansion of 1/([r]-[2r]x+[3r]x^2-...); [ ]=floor, r=13/8.

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012543 | OEIS-A289262

Coefficients in the expansion of 1/([r]-[2r]x+[3r]x^2-...); [ ]=floor, r=11/7.

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012544 | OEIS-A289273

Complement of A242410.

Conjecture: there will never be a point where this sequence has more terms than A242410. (80 is only the 21st term of this sequence.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Jun 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012545 | OEIS-A289313

The number of upper-triangular matrices with integer entries whose absolute sum is equal to n and such that each row contains a nonzero entry.

Conjectural g.f.: Sum_{n >= 0} ((1 + x)/(1 - x))^((n+1)*(2*n+1)) * Product_{i = 1..2*n} (((1 - x)/(1 + x))^i - 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012546 | OEIS-A289314

Number of n X n Fishburn matrices with entries in the set {0,1,2}.

Two conjectural continued fractions for the o.g.f.:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012547 | OEIS-A289315

Number of n X n Fishburn matrices with entries in the set {0,1,2,3}.

Two conjectural continued fractions for the o.g.f.:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012548 | OEIS-A289329

Triangle of coefficients of polynomials p_n(x) (exponents in increasing order) where 2^(x-n-1)p_n(x)/n! counts the ways a game of Nim with two piles of sizes x and n can be played out (x > 0, n >= 0).

The coefficients are conjectured to be positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jane Wang_, Jul 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012549 | OEIS-A289352

Irregular triangle read by rows: T(n,k) = the number of Dyck paths of semilength n with floor((n+2)/2) up movements in odd numbered positions and k returns to the x axis.

T(1,1)=1, T(2,1)=0, T(2,2)=1, For n >= 3, T(n,k) = (1/floor((n-1)/2))*C(n-1,floor((n-3)/2))*C(n-1-k,floor((n-3)/2))*k (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger Ford_, Jul 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012550 | OEIS-A289398

Least integer m > n such that (n^2 + m^2)/2 is a square.

Conjecture: this is "if and only if".

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Jul 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012551 | OEIS-A289404

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 566", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 05 2017: (Start) Conjectures from _Federico Provvedi_, Nov 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012552 | OEIS-A289405

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 566", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 05 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012553 | OEIS-A289428

Sum of factorial-Catalan numbers over a certain family of decorations.

The following are conjectural (Cf. A145879):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012554 | OEIS-A289559

Number of modulo n residues among sums of two fourth powers.

Conjecture: the only primes p for which a(p) < p are 5, 13, 17, 29. - _Robert Israel_, Jul 09 2017 Conjecture is true: see Math Overflow link. - _Robert Israel_, Apr 01 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Jul 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012555 | OEIS-A289587

a(n) is the number of permutations of length n that avoid the pattern 321 and the mesh pattern (12, 174) or the same sequence for the mesh pattern (12, 234).

Conjectured g.f.: (x^4 + 4*x^3 + 11*x^2 - (x^2 + 5*x + 3)*sqrt(x^4 - 2*x^3 - 5*x^2 - 2*x + 1) + 10*x + 3)/(8*x*(x + 1)^2). - _Thomas Scheuerle_, Dec 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012556 | OEIS-A289629

Smallest positive k such that (k+1)^n + (-k)^n is divisible by a square greater than 1.

Conjecture: (k+1)^n + (-k)^n is not squarefree for the following (n, k) patterns, with m >= 1: (22m-11, 2), (20m-6, 3), (20m-2, 3), (3^m, 7), (15m, 7), (20m-15, 14), (16m-8, 19), (42m-35, 23), and (8m-4, 113). In each case, the value of a(n) in this sequence is usually equal to the value specified for k, but sometimes this value is not the smallest such k. For example, in the (n, k) = (20m-2, 3) case, a(20m-2) = 3 for m = 1..3, but at m = 4, a(20*4-2) = a(78) = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Sep 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012557 | OEIS-A289631

Prime powers P for which the number of modulo P residues among sums of two sixth powers is less than P.

Conjecture: 223 is the final prime in this sequence. Conjecture: the terms in this sequence that are the squares of primes are the squares of 13, 37, 61, 73, 109, and every prime not congruent to 1 mod 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Jul 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012558 | OEIS-A289654

Related to number of mesh patterns of length 2 that avoid the pattern 321.

Conjecture: Catalan numbers - 2 (A120304) with first five terms changed to 1,1,1,3,13. _Georg Fischer_, Jan 07 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012559 | OEIS-A289674

Consider the Post tag system described in A284116 (but adapted to the alphabet {1,2}); sequence lists the words that belong to cycles.

It is an important open question to decide if there is any word whose orbit grows without limit.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012560 | OEIS-A289682

Catalan numbers read modulo 16.

Conjecture: a(2^n-1) = 13 and a(2^n) = 6 for n >= 3. - _Robert Israel_, Jul 09 2017 The conjecture is correct. For the first part, see A178854 for more details; for the second part, simply note that Catalan(n+1)/Catalan(n) = 2*(2*n+1)/(n+2), and so in particular Catalan(2^k)/Catalan(2^k-1) == -2 (mod 2^k). - _Jianing Song_, Apr 15 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jul 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012561 | OEIS-A289683

Mixing moments of the busy period of mean steady-state 1/2 in an M/M/1 waiting process.

Conjecture: a(n) +2*(-2*n+3)*a(n-1) +(n-1)*(n-3)*a(n-2)=0. - _R. J. Mathar_, Jul 09 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jul 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012562 | OEIS-A289684

Mixing moments for the waiting time in an M/G/1 waiting queue.

Conjecture: n*a(n) + 2*(-4*n+3)*a(n-1) + 12*(n-2)*a(n-2) + 8*(2*n-3)*a(n-3) = 0. Conjecture verified (for n >= 4) using the differential equation (16*x^3 + 12*x^2 - 8*x + 1)*y' + (24*x^2 - 2)*y -12*x^2 + 2*x = 0 satisfied by the g.f.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jul 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012563 | OEIS-A289740

Prime powers P for which the number of modulo P residues among sums of three sixth powers is less than P.

Conjecture: the largest prime in the sequence is 31. (If this is true, then the next terms after 32768 are 49729, 59049, and 65536.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Jul 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012564 | OEIS-A289748

Thue-Morse constant converted to base -2.

M = Thue-Morse + 4/3 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | Gilian Breysens, Jul 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012565 | OEIS-A289749

Number of ways not ending in 011 to write n in base phi.

Conjecture: a(A005248(k)) = 2k+1 for k=1,2,...(cf. Theorem 2 in the paper by Carlitz.) - _Michel Dekking_, Nov 14 2021 This conjecture is proved in the paper "Counting base phi representations". - _Michel Dekking_, Jul 15 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | Gilian Breysens, Jul 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012566 | OEIS-A289760

Prime powers P for which the number of modulo P residues among sums of four sixth powers is less than P.

Conjecture: the largest prime in the sequence is 13.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Jul 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012567 | OEIS-A289761

Maximum length of a perfect Wichmann ruler with n segments.

Conjectures from _Colin Barker_, Jul 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jul 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012568 | OEIS-A289763

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 597", based on the 5-celled von Neumann neighborhood.

G.f.: (1 - 10*x + 100*x^2 - 1000*x^4) / ((1 - x)*(1 - 10*x)*(1 - 10*x^2)*(1 + 100*x^2)) (conjectured). - _Colin Barker_, Jul 14 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012569 | OEIS-A289764

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 597", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 14 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012570 | OEIS-A289765

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 597", based on the 5-celled von Neumann neighborhood.

G.f.: (1 - 2*x + 4*x^2 - 8*x^4) / ((1 - x)*(1 - 2*x)*(1 - 2*x^2)*(1 + 4*x^2)) (conjectured). - _Colin Barker_, Jul 14 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012571 | OEIS-A289777

Frequency of the largest spectral component of the prime characteristic function of the first n numbers, for n>3, excluding the smallest and largest frequencies. In case of a tie, use the smallest frequency.

Conjecture: lim_{n->inf} abs(4a(n)/n - 1) = 1/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andres Cicuttin_, Jul 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012572 | OEIS-A289873

Related to perfect Wichmann rulers: a(n) = (n^2 - (mod(n, 6) - 3)^2) / 3.

G.f.: x^2*(1 + x - x^2)*(1 + x^2 - x^3 + 2*x^4 + x^5) / ((1 - x)^3*(1 + x)*(1 - x + x^2)*(1 + x + x^2)) (conjectured). - _Colin Barker_, Jul 14 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jul 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012573 | OEIS-A289912

Coefficients of 1/(Sum_{k>=0} round((k+1)*r)(-x)^k), where r = sqrt(2).

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012574 | OEIS-A289914

Coefficients of 1/(Sum_{k>=0} round((k+1)*r)(-x)^k), where r = 7/5.

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012575 | OEIS-A289916

Coefficients of 1/(Sum_{k>=0} round((k+1)*r)(-x)^k), where r = 9/7.

Conjecture: the sequence is strictly increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012576 | OEIS-A289921

Coefficients of 1/([1+r] - [1+2r]x + [1+3r]x^2 - ...), where [ ] = floor and r = 9/10.

Conjecture: all the terms are nonnegative.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012577 | OEIS-A289923

Limiting sequence of coefficients of 1/([1+r] - [1+2r]x + [1+3r]x^2 - ...), where [ ] = floor and r approaches 19/21 from the left.

Conjecture: all the terms are nonnegative.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012578 | OEIS-A289927

p-INVERT of A014217 (starting at n=1), where p(S) = 1 - S - S^2.

Conjectures from _Colin Barker_, Aug 15 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012579 | OEIS-A289997

Numbers n whose trajectory under iteration of the map k -> (sigma(k)+phi(k))/2 never reaches a fraction (that is, either the trajectory reaches a prime, which is a fixed point, or diverges to infinity).

Conjecturally, all terms of A291790 are in the sequence, because their trajectories (see example in A291789 for starting value 270) grow indefinitely. - _Hugo Pfoertner_, Sep 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Sep 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012580 | OEIS-A290003

G.f.: A(x) = Sum_{n=-oo..+oo} (x - x^n)^n.

a(p+1) = 1 for primes p > 3 (conjecture). The above conjecture follows from the following formula: for n >= 2,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012581 | OEIS-A290012

a(n) is the smallest prime number p satisfying p^2 >= Sum_{1 <= k <= n} prime(k)^2.

Conjecture: The only twin prime pair in the sequence is (5, 7).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimitris Valianatos_, Jul 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012582 | OEIS-A290070

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 641", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012583 | OEIS-A290072

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 641", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012584 | OEIS-A290073

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 641", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012585 | OEIS-A290074

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 641", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012586 | OEIS-A290078

Where the ratio A235027(n)/n obtains record values.

Because A056539(n)/n < 2 for all n, and already for the tenth term of this sequence 1349 we have A235027(1349)/1349 = 2.094... it follows that the only primes present are terms a(2) .. a(7): 11, 19, 67, 71, 263, 271. Conjecture: every term after that is a product of some of those six primes. For example: 781 = 11*71, 1273 = 19*67, 1349 = 19*71, 2981 = 11*271, 4757 = 67*71, 5041 = 71*71.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012587 | OEIS-A290136

Positive numbers that are not the sum of two nonprime squarefree numbers (A000469).

The sequence is conjectured to be complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Jul 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012588 | OEIS-A290169

a(n) = least k such that both the sum of the smallest n divisors of k and the sum of its greatest n divisors are prime numbers.

The sequence contains a subsequence of numbers having the property that the sum of the first n divisors is equal to the sum of the last n divisors; for instance, for a(n) = 2, 4, 16 and 64 with n = 2, 3, 5 and 7. Is it possible to conjecture that this subsequence contains all the superperfect numbers (A019279)? The answer is no: for instance, A019279(5) = 4096 = 2^12 => the sum of the 13 terms 1 + 2 + 4 + 8 + ... + 4096 = 8191 is a Mersenne prime, but a(13) = 800 instead 4096 > 800, and we obtain the corresponding pair of primes (293, 1933) instead (8191, 8191).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jul 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012589 | OEIS-A290190

Minimum sum of mutual Manhattan distances of n distinct grid points in a cubic lattice.

Conjecture: a(k^3) = A292045(k), i.e., fully populated cubes are optimal 3-dimensional n-towns. - _Hugo Pfoertner_, Jan 08 2026 The conjecture is false. Optimal 3-dimensional k^3-towns exist for k >= 4 with distance sums < A292045(k). For example, there are 64-towns (see linked illustration) with distance sum = 7539 < A292045(4) = 7680 and 125-towns with distance sum 36510 < A292045(5) = 37500. - _Hugo Pfoertner_, Jan 26 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jul 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012590 | OEIS-A290191

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 673", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) is the binary representation of 2^(n - 1) - 1 - (n mod 2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012591 | OEIS-A290192

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 705", based on the 5-celled von Neumann neighborhood.

a(n) = 10*a(n-1) + a(n-2) - 10*a(n-3) for n > 5 (conjectured). G.f.: (10*x^5 + 89*x^4 + 91*x^3 + 1)/((x - 1)*(x + 1)*(10*x - 1)) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012592 | OEIS-A290193

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 705", based on the 5-celled von Neumann neighborhood.

Conjecture: For even n > 3, binary representation of 3 * 2^(n - 1) - 1. For odd n > 3, binary representation of 2^(n - 1) - 1. - _David A. Corneth_, Jul 23 2017 a(n) = a(n-1) + 100*a(n-2) - 100*a(n-3) for n > 5 (conjectured). G.f.: (10000*x^5 - 10900*x^4 + 910*x^3 + 1)/((x - 1)*(10*x - 1)*(10*x + 1)) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012593 | OEIS-A290194

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 705", based on the 5-celled von Neumann neighborhood.

Conjecture: a(n) = Fibonacci(2*n+1) if n <= 3, for n > 3, a(n) = 2*a(n-1) + 2 if n is even, a(n) = 2*a(n-1) + 5 if n is odd. It would follow that a(n) = 2^(n+1) - 4 + (n mod 2) for n >= 3. - _David A. Corneth_, Jul 23 2017 a(n) = 2*a(n-1) + a(n-2) - 2*a(n-3) for n > 5 (conjectured). G.f.: (2*x^5 + x^4 + 3*x^3 + 1)/((x - 1)*(x + 1)*(2*x - 1)) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012594 | OEIS-A290195

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 705", based on the 5-celled von Neumann neighborhood.

Conjecture: For odd n > 3, a(n) = 2^(n-1) - 1, for even n > 3, a(n) = 3*2^(n-1) - 1. - _David A. Corneth_, Jul 23 2017 a(n) = a(n-1) + 4*a(n-2) - 4*a(n-3) for n > 5 (conjectured). G.f.: (16*x^5 - 20*x^4 + 6*x^3 + 1)/((x - 1)*(2*x - 1)*(2*x + 1)) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012595 | OEIS-A290196

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 721", based on the 5-celled von Neumann neighborhood.

Conjecture: for n>=4, a(n) = 10*(10^n - 1) / 9 (in decimal). - _David A. Corneth_, Jul 23 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012596 | OEIS-A290197

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 721", based on the 5-celled von Neumann neighborhood.

Conjecture: For n>=4, a(n) = (10^n-1) / 9 (in decimal). _David A. Corneth_, Jul 23 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012597 | OEIS-A290198

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 721", based on the 5-celled von Neumann neighborhood.

Conjecture: For n<=3, a(n) = Fibonacci(2*n + 1), for n >= 4, a(n) = 2^(n+1) - 2. _David A. Corneth_, Jul 23 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012598 | OEIS-A290199

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 721", based on the 5-celled von Neumann neighborhood.

Conjecture: if n>=4, a(n) = 2^n - 1. - _David A. Corneth_, Jul 23 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012599 | OEIS-A290203

Numbers m having greatest prime power divisor d such that d is smaller than the difference between m and the largest prime smaller than m.

It is conjectured that for all integers m there exist two primes p and r such that all the binomial coefficients (m,k) with 1 <= k <= m-1 are divisible by either p or r. Using Lucas's Theorem we can prove that the conjecture is true for integers m such that the difference between m and the largest prime smaller than m is smaller than the greatest prime power divisor of m. Therefore this list examines the numbers m that do not satisfy this property.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sílvia Casacuberta Puig_, Jul 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012600 | OEIS-A290232

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 773", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012601 | OEIS-A290233

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 773", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012602 | OEIS-A290234

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 773", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012603 | OEIS-A290235

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 773", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012604 | OEIS-A290251

a(n) is the number of parts in the integer partition having viabin number n.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Jul 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012605 | OEIS-A290268

Number of terms in the fully expanded n-th derivative of x^(x^2).

Conjectured g.f.: (1 + x + 2*x^2 + 2*x^3 + 2*x^4 + 2*x^5 + 2*x^6 + x^7 + 2*x^8 + x^9)/((1 - x)*(1 - x^2)*(1 - x^8)). Conjecture: a(n) = (8*n^2 + 15*n + 14 + (n + 2)*(-1)^n + (2 - 4*sqrt(2)*sin(Pi*n/4))*sin(Pi*n/2))/16. From _Peter Luschny_, Oct 09 2017: (Start) Assuming the conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Reshetnikov_, Oct 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012606 | OEIS-A290277

Inverse Euler Transform of the Motzkin Numbers.

Conjecture: n*a(n) = Sum_{d|n} mobius(d)*A002426(n/d) where mobius=A008683. - _R. J. Mathar_, Nov 05 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jul 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012607 | OEIS-A290278

a(n) = Sum_{k=0..n} (A007953(5*k) - A007953(k)).

- if i and j divides b and k > 0 and 0 <= n <= b^k-1, then d_b(i * n) - d_b(j * n) = d_b(j * (b^k-1 - n)) - d_b(i * (b^k-1 - n)) (this implies the conjecture about the symmetry of A289411),

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jul 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012608 | OEIS-A290281

Numbers k such that (k-1) mod phi(k) = lambda(k), where phi = A000010 and lambda = A002322.

Conjecture: these are numbers k such that phi(k) + lambda(k) = k - 1. Checked up to 2^64. - _Amiram Eldar_ and _Thomas Ordowski_, Dec 06 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_ and _Thomas Ordowski_, Jul 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012609 | OEIS-A290283

Primes p such that A215458(p) is prime.

It is conjectured that there are infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul S. Vanderveen_, Jul 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012610 | OEIS-A290290

Numbers m having greatest prime power divisor d such that d is smaller than the difference between m and the largest prime smaller than m and d is smaller than the difference between m and twice the largest prime smaller than m/2.

It is conjectured that for all integers m there exist two primes p and r such that all the binomial coefficients (m,k) with 1 <= k <= m-1 are divisible by either p or r. Using Lucas's Theorem we can prove that the conjecture is true for integers m such that the difference between m and the largest prime smaller than m is smaller than the greatest prime power divisor of m. In A290203 we examine the integers that do not satisfy this property. However, when this property is not satisfied, if the difference between m and twice the largest prime smaller than m/2 is smaller than the greatest prime power divisor of m we can prove the conjecture for odd numbers and some cases of even numbers. Note that we take the difference between m and twice the largest prime smaller than m/2 because this is the last digit of the base-p representation of m, where p refers to the largest prime smaller than m/2. Therefore, this list examines the numbers that satisfy neither of the two properties and hence are the difficult cases for the proof of the conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sílvia Casacuberta Puig_, Jul 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012611 | OEIS-A290322

Sum modulo n of all units u in Z/nZ such that Phi(5,u) is a unit, where Phi is the cyclotomic polynomial.

Conjecture: If n is divisible by 5 then a(n) > 0. - _Robert Israel_, Jan 23 2024 Conjecture proven by Wenlin Zhang and Haobo Ma with AI assistance (see links). - _Robert Israel_, Sep 30 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joshua Harrington_, _Michael Mueller_, _Jordan Lenchitz_, _Tristan Phillips_, _Madison Wellen_, _Eric Jovinelly_, Jul 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012612 | OEIS-A290334

Numbers that are not the sum of three or fewer terms from A020330.

Not currently proved that there are infinitely many terms. It is conjectured that all integers > 686 are the sum of four binary squares.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Jul 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012613 | OEIS-A290342

Number of ways to write n as x^2 + 2*y^2 + z*(z+1)/2, where x is a nonnegative integer, and y and z are positive integers.

Conjecture: a(n) > 0 for all n > 2. In other words, each n = 0,1,2,... can be written as x^2 + 2y*(y+2) + z*(z+3)/2 with x,y,z nonnegative integers. Let a,c,e be positive integers, and let b,d,f be nonnegative integers with a-b, c-d, e-f all even. Suppose that a|b, c|d and e|f. The author studied in arXiv:1502.03056 when each nonnegative integer can be written as x*(a*x+b)/2 + y*(c*y+d)/2 + z*(e*z+f)/2 with x,y,z nonnegative integers, and conjectured that the answer is positive if (a,b,c,d,e,f) is among the following ten tuples (4,0,2,0,1,3), (4,0,2,0,1,5), (4,0,2,6,1,1), (4,0,2,6,2,0), (4,4,2,0,1,3), (4,8,2,0,1,1), (4,8,2,0,1,3), (4,12,2,0,1,1), (6,0,2,0,1,3), (6,6,2,0,1,3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012614 | OEIS-A290347

Numerators of the Harary index for the n-halved cube graph.

For p > 3, if p is prime, then p^2 divides a(p). Conjecture: for n > 3, if n^2 divides a(n), then n is prime. Primes p such that p^3 divides a(p) are probably A088164. - _Thomas Ordowski_, Mar 30 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Jul 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012615 | OEIS-A290354

a(n) is the n-th term of the n-th Euler transform of the sequence with g.f. 1+x.

Conjecture: a(n) ~ c * 2^n * n^(n-4/3) / Pi^n, where c = 4.4923... - _Vaclav Kotesovec_, Aug 14 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012616 | OEIS-A290397

Least number of nonprime squarefree numbers (A000469) that add up to n.

It is conjectured that a(n) <= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Jul 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012617 | OEIS-A290408

Decimal expansion of the real part of the solution of z = (i+z)^(-i) in C (i is the imaginary unit).

In C, the unique invariant point of the mapping M(z) = (i+z)^(-i) is not the attractor of the mapping (unstable behavior), but it is an attractor of the modified mapping M'(z) = (z+M(z))/2. For M', it takes 5000 iterations to reduce the value of |z - M'(z)| below 10^(-3400). Interestingly, the imaginary part of z seems to be equal to -1/2 (verified to 5000 digits). If this conjecture holds, and considering the definition, one can symbolically write (i+(i+(i+...)^(-i))^(-i))^(-i) = a - i/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Jul 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012618 | OEIS-A290412

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 785", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 31 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012619 | OEIS-A290413

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 785", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 31 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012620 | OEIS-A290414

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 785", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 31 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012621 | OEIS-A290415

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 785", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Jul 31 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Jul 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012622 | OEIS-A290443

a(n) = Catalan(n)*Motzkin(n-1).

Conjecture: n*(n+1)^2*a(n) -2*-n*(2*n-1)^2*a(n-1) -12*(n-2)*(2*n-1)*(2*n-3)*a(n-2)=0. - _R. J. Mathar_, Aug 05 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012623 | OEIS-A290467

Unitary half-Zumkeller numbers: numbers k whose unitary proper divisors can be partitioned into two disjoint sets whose sums are equal.

The conjecture above is false, since 72, 3600 and 19600 do not belong to A246198. - _Ivan N. Ianakiev_, Jan 08 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ivan N. Ianakiev_, Aug 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012624 | OEIS-A290472

Number of ways to write 6*n+1 as x^2 + 3*y^2 + 7*z^2, where x is a positive integer, and y and z are nonnegative integers.

Conjecture: a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 1, 2, 4, 5, 9, 10, 12, 14, 16, 17, 24, 30, 34, 66, 84, 86, 116, 124, 152, 286. We also conjecture that {6n+5: n = 0,1,2,...} is a subset of {2x^2+3y^2+5z^2: x,y,z are nonnegative integers with y > 0}. See A286885 for more similar conjectures. In support of the first conjecture, a(n) > 1 for 286 < n <= 10^7. - _Charles R Greathouse IV_, Aug 04 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012625 | OEIS-A290491

Number of ways to write 12*n+1 as x^2 + 4*y^2 + 8*z^4, where x and y are positive integers and z is a nonnegative integer.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 4, 10, 18, 28, 39, 49, 98, 142, 163, 184, 208, 320, 382, 408, 814, 910, 1414, 2139, 2674, 3188, 3213, 4230, 6279, 25482.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012626 | OEIS-A290515

a(n) = smallest number that is the start of a gap of size n between successive prime powers (A000961), or 0 if no such number exists.

Conjecture: a(n) always exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Aug 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012627 | OEIS-A290525

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 833", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Aug 05 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012628 | OEIS-A290526

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 833", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Aug 05 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012629 | OEIS-A290527

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 833", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Aug 05 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012630 | OEIS-A290528

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 833", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Aug 05 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012631 | OEIS-A290540

Determinant of circulant matrix of order 10 with entries in the first row that are (-1)^(j-1)*Sum_{k>=0} (-1)^k*binomial(n, 10*k+j-1), for j=1..10.

A generalization. For an even N >= 2, consider the determinant of circulant matrix of order N with entries in the first row (-1)^(j-1)K_j(n), j=1..N, where K_j(n) = Sum_{k>=0} (-1)^k*binomial(n, N*k+j-1). Then it is 0 for n == N-1 (mod N). This statement follows from an easily proved identity K_j(N*t + N - 1) = (-1)^t*K_(N - j + 1)(N*t + N - 1) and a known calculation formula for the determinant of circulant matrix [Wikipedia]. Besides, it is 0 for n=1..N-2. We also conjecture that every such sequence contains infinitely many blocks of N-1 negative and N-1 positive terms separated by 0's.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_ and _Peter J. C. Moses_, Aug 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012632 | OEIS-A290575

Apéry-like numbers Sum_{k=0..n} (C(n,k) * C(2*k,n))^2.

Conjecture: The g.f. has a square root with integer coefficients 1, 2, 18, 236, ... - _F. Chapoton_, Apr 23 2026 Chapoton's conjecture is true - see Theorem 6.2 of the linked preprint by Easwar. - _Rohun Easwar_, Aug 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Aug 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012633 | OEIS-A290601

Irregular triangle read by rows: T(n, k) gives the least positive integer imin such that the sequence {A290600(n, k)^i}_{i >= imin} (mod A002808(n)) is periodic.

Conjecture:There exists always an imin >= 1 such that the power sequence A290600(n, k)^i (mod A002808(n)) is periodic for i >= imin. Otherwise T(n, k) is undefined, and one could use T(n, k) = -1. See below for a proof. The above conjecture is trivially true because {K^i}_{i >= 0} (mod N) becomes (or is) always periodic (K and N positive integer). This follows from the fact that at most N values are possible, namely {0, 1, ..., N-1} and for i >= N one of these values has to appear for the second time, leading to periodicity. Thus the above sequence where gcd(N, K) is not 1 is also defined. - _Wolfdieter Lang_, Sep 05 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Aug 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012634 | OEIS-A290615

Number of maximal independent vertex sets (and minimal vertex covers) in the n-triangular honeycomb bishop graph.

Conjecture: a(n) = Sum_{k=0..2^(n-1) - 1} b(k) for n > 0 where b(2n+1) = b(n - 2^f(n)), b(2n) = b(n) + b(2n - 2^f(n)) for n > 0 with b(0) = b(1) = 1 and where f(n) = A007814(n). Also b((4^n - 1)/3) = (floor((n+1)/2)!)^3. - _Mikhail Kurkov_, Sep 18 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Aug 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012635 | OEIS-A290633

Lexicographically earliest sequence of positive integers such that, for any m and n > 0, gcd(a(n), a(n+1)) > 1 and a(n) != a(n+2), and if m < n then a(m) != a(n) or a(m+1) != a(n+1).

Conjecturally, (a(n), a(n+1)) runs over all pairs of noncoprime positive integers; in this sense, this sequence is opposite to sequences like Stern's diatomic series (A002487). This sequence has connections with A067992: here we avoid duplicate ordered pairs of consecutive terms, there unordered pairs, here we deal with noncoprime consecutive terms, there we (conjecturally) have coprime consecutive terms; also, the scatterplots of these sequences have similarities.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Aug 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012636 | OEIS-A290639

a(n) = largest number <= prime(n) such that 1 + a(1)*a(2)*...*a(n) is prime.

Conjecture: a(n) > 1 for every n. - _Thomas Ordowski_, Aug 08 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Aug 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012637 | OEIS-A290642

a(n) is the maximum diameter for an n-threaded binary program. In other words, a(n) is the maximal finite distance in the transition graph of an n-threaded binary program.

Conjecture: a(n) = 3*n for all n >= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Malkis_, Aug 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012638 | OEIS-A290648

a(n) is the smallest number of faces of the triangular lattice required to enclose an area consisting of exactly n faces.

Conjecture: a(n) = 2*A067628(n) + 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian Majenz_, Aug 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012639 | OEIS-A290650

a(1) = 1. For n > 1, a(n) = a(n-1)/2 if a(n-1) is even, a(n) = a(n-1)*n otherwise.

Conjecture: The sequence is unbounded. - _Felix Fröhlich_, Aug 08 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John Bailey_, Aug 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012640 | OEIS-A290653

Number of perfect dissections of equilateral triangles into n equilateral triangles with integer sides.

A perfect dissection has no two triangles of the same side. Triangles of different orientation are counted separately. The computation by A. Drápal and C. Hämäläinen proves Tutte's conjecture that the smallest perfect dissection has size 15.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Aug 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012641 | OEIS-A290660

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 899", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Aug 03 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012642 | OEIS-A290661

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 899", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Aug 03 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012643 | OEIS-A290662

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 899", based on the 5-celled von Neumann neighborhood.

Conjectures from _Chai Wah Wu_, Aug 03 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012644 | OEIS-A290680

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 929", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Aug 09 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012645 | OEIS-A290681

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 929", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Aug 09 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012646 | OEIS-A290682

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 929", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Aug 09 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012647 | OEIS-A290683

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 929", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Aug 09 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012648 | OEIS-A290694

Triangle read by rows, numerators of coefficients (in rising powers) of rational polynomials P(n, x) such that Integral_{x=0..1} P'(n, x) = Bernoulli(n, 1).

Let C_k(n) = [x^k] P_n(x), k > 0 and n even. Conjecture: k is a prime factor of Clausen(n) <=> k = denominator(C_k(n)) <=> k does not divide Stirling2(n, k-1)*(k-1)!. (Note that by a comment in A019538 Stirling2(n, k-1)*(k-1)! is the number of chain topologies on an n-set having k open sets.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Aug 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012649 | OEIS-A290735

a(n) = weighted sum over all the self-conjugate partitions of 4n + 1 into odd parts, with respect to a certain weight.

Andrews conjectures that a(n) > 0 for all n. The conjecture is known to be true for n <= 1000. Andrews also conjectures that a(n) > |A290737(n) + A290739(n)| for n >= 2 (see A290740).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012650 | OEIS-A290740

a(n) = A290735(n) - |A290737(n) + A290739(n)|.

Andrews conjectures that a(n) >= 1 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012651 | OEIS-A290744

Maximum number of distinct Lyndon factors that can appear in words of length n over an alphabet of size 5.

Conjectures from _Colin Barker_, Oct 03 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012652 | OEIS-A290745

Maximum number of distinct Lyndon factors that can appear in words of length n over an alphabet of size 10.

G.f.: x*(10 - 9*x + x^2 + x^3 + x^4 + x^5 + x^6 + x^7 + x^8 + x^9 - 10*x^10 + 10*x^11) / ((1 - x)^3*(1 + x)*(1 - x + x^2 - x^3 + x^4)*(1 + x + x^2 + x^3 + x^4)) (conjectured). - _Colin Barker_, Oct 03 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012653 | OEIS-A290813

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 961", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Aug 12 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012654 | OEIS-A290815

Numbers m such that the numerator of Sum_{k=1..m, gcd(k,m) = 1} 1/k is divisible by m^3.

A general conjecture: if, for some e > 0, m^e | Numerator(Sum_{k=1..m, gcd(k,m)=1} 1/k), then m^(e-1) | Numerator(Sum_{k=1..m, gcd(k,m)=1} 1/k^2). Note: in this case, the exponent e = 3. Problem: are there numbers m > 1 such that m^4 | Numerator(Sum_{k=1..m, gcd(k,m)=1} 1/k)? - _Thomas Ordowski_, Aug 10 2019 This general conjecture was checked up to 10^4. This problem has no solution up to 10^5. - _Amiram Eldar_, Aug 10 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Aug 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012655 | OEIS-A290821

Side length of largest equilateral triangle that can be made from n or fewer equilateral triangles with integer sides s_k, subject to gcd(s_1,s_2,...,s_n) = 1.

No construction from 2, 3 or 5 equilateral triangles exists. The first difference from the Padovan numbers occurs for a(15)=39, where the corresponding term A000931(19)=37. a(16)=A000931(20)=49. a(n) >= A000931(n+3). From the growth behavior of A290697 it is conjectured that a(k) > A000931(k+3) for all k > 20.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Aug 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012656 | OEIS-A290828

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 961", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Aug 12 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012657 | OEIS-A290829

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 961", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Aug 12 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012658 | OEIS-A290834

Binary representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 969", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Aug 12 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012659 | OEIS-A290835

Binary representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 969", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Aug 12 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012660 | OEIS-A290836

Decimal representation of the diagonal from the corner to the origin of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 969", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Aug 12 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012661 | OEIS-A290837

Decimal representation of the diagonal from the origin to the corner of the n-th stage of growth of the two-dimensional cellular automaton defined by "Rule 969", based on the 5-celled von Neumann neighborhood.

Conjectures from _Colin Barker_, Aug 12 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012662 | OEIS-A290844

Square array read by antidiagonals downwards: A(n, k) = (Sum_{i=1..n} i^k) - (n+1)^k for n >= 1, k >= 1.

Paul Erdős conjectured that A(n, k) = 0 only for (n, k) = (2, 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Aug 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012663 | OEIS-A290883

E.g.f. A(x) = sqrt(C(x)^2 + S(x)^2) such that C(x)^2 - S(x)^2 = 1 and C'(x)^2 + S'(x)^2 = 1, where C(x) is the e.g.f. of A290880 and S(x) is the e.g.f. of A290881.

Conjecture: If p is a prime congruent to 1 mod 4 and k is a positive integer, then p^k divides a(n) for n >= (k/2)*p-1. Furthermore, if p^k divides (2n+2)!, then p^k divides a(n). - _Andrew Slattery_, Aug 21 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012664 | OEIS-A290943

Number of ways to write n as an ordered sum of 3 generalized pentagonal numbers (A001318).

Conjecture: every number is the sum of at most k - 4 generalized k-gonal numbers (for k >= 8).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Aug 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012665 | OEIS-A291047

Number of ways to write 4*n+1 as p^2 + q^2 + 8*r^2, where p is prime, and q and r are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n > 0, and the only values of n >= 1000 with a(n) = 1 are 1000, 1052, 1472, 1675, 1967, 4787, 4822, 11962.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012666 | OEIS-A291051

a(n) is the smallest number k such that psi(k) = n*phi(k) where psi(k) is Dedekind psi function (A001615) and phi(k) is Euler totient function (A000010), or 0 if no such k exists.

It is conjectured that A055234(n) > 0 for each n. Is a(n) > 0 for all values of n?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Aug 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012667 | OEIS-A291056

a(n) = k if the first appearance of n in A077618 is at index k, or 0 if k does not appear in A077618.

Conjecture: all positive integers appear in A077618, so a(n) is never 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Oct 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012668 | OEIS-A291123

Number of ways to write 4*n+1 as (p-1)^2 + q^2 + r^2, where p is prime and q and r are nonnegative integers with q <= r.

Conjecture: (i) a(n) > 0 for all n, and a(n) = 1 only for n = 0, 3, 6, 23, 44.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012669 | OEIS-A291127

Consider the zeros of the polynomial P(m,x) whose coefficients are the divisors of a number m. The sequence lists the numbers m such that P(m,x) contains at least two zeros that are purely imaginary numbers.

Conjecture: A032741(a(n))= p where p == 3 (mod 4), p prime. Examples: The conjecture above is false. The first counterexample is a(867) = 1806, for which A032741(1806) = 15 is composite. Indeed, P(1806,x) = (1 + 2*x)*(1 + 3*x^2)*(1 + 7*x^4)*(1 + 43*x^8), so 1806 is in the sequence since 1 + 3*x^2 has two purely imaginary zeros. - _Qiyuan Gu_, Aug 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Aug 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012670 | OEIS-A291139

Primes of the form floor(k^(3/2)) for some integer k.

Conjecturally, a(n) ~ (1.5 n log n)^1.5 and there are ~ x^(2/3)/log x members of this sequence up to x. - _Charles R Greathouse IV_, Oct 14 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Aug 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012671 | OEIS-A291150

Number of ways to write 2*n+1 as x^2 + y^2 + z^2 + 4*w^2, where x,y,z,w are nonnegative integers with x <= y <= z such that 2^x + 2^y + 2^z + 1 is prime.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 1, 2, 3, 5, 7, 11, 14, 15, 21, 23, 35, 41, 43, 59, 71, 309, 435.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012672 | OEIS-A291191

Number of ways to write 2*n+1 as x^2 + y^2 + z^2 + w^2, where x,y,z,w are nonnegative integers with x <= y, z <= w and x + y < z + w such that 2^(x+y) + 2^(z+w) + 1 is prime.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 2, 3, 6, 7, 11, 12, 16, 19, 20, 23, 26, 63, 75, 98. See also A291150 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 20 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012673 | OEIS-A291271

The arithmetic function v_4(n,2).

Conjecture: a(n) = (n-2-cos(n*Pi)-cos(n*Pi/2))/2. - _Wesley Ivan Hurt_, Oct 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012674 | OEIS-A291293

Sequence mod 5 defined by Baldini-Eschgfäller coupled dynamical system (f,lambda,alpha) with f(k) = A000203(k)-1, lambda(y) = 3y+2 mod 5 for y in Y = {0,1,2,3,4}, and alpha(k) = k mod 5 for k in Omega = {primes}.

This sequence assumes that the Erdos conjecture is true, that iterating k -> sigma(k)-1 always reaches a prime (cf. A039654).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012675 | OEIS-A291437

Smallest m >= 0 such that (2*n)*3^m + 1 is prime, or -1 if no such value exists.

It is conjectured that the smallest one is 125050976086 = A123159(3), therefore a(62525488043) = -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Martin Renner_, Aug 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012676 | OEIS-A291438

Smallest prime of the form (2*n)*3^m + 1 for some m >= 0, or -1 if no such prime exists.

It is conjectured that the smallest one is 125050976086 = A123159(3), therefore a(62525488043) = -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Martin Renner_, Aug 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012677 | OEIS-A291455

Number of ways to write 2*n+1 as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that x + 3*y + 5*z + 7*w, x^3 + 3*y^3 + 5*z^3 + 7*w^3 and x^7 + 3*y^7 + 5*z^7 + 7*w^7 are all prime.

Conjecture: (i) a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 3, 4, 7, 17, 23, 34, 39, 51, 66, 69, 99, 109, 115, 171, 191. Also, any integer n > 1 with gcd(n,42) = 1 can be written as x + 3*y + 5*z + 7*w with x,y,z,w nonnegative

integers such that x^3 + 3*y^3 + 5*z^3 + 7*w^3 and x^7 + 3*y^7 + 5*z^7 + 7*w^7 are both prime. See also A290935 for a similar conjecture involving twin primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012678 | OEIS-A291499

The arithmetic function uhat(n,1,5).

Conjectures from _Chai Wah Wu_, Jun 10 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012679 | OEIS-A291502

The arithmetic function uhat(n,1,8).

Conjectures from _Chai Wah Wu_, Jun 10 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012680 | OEIS-A291514

The arithmetic function uhat(n,3,5).

Conjectures from _Chai Wah Wu_, Jun 10 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012681 | OEIS-A291515

The arithmetic function uhat(n,3,6).

Conjectures from _Chai Wah Wu_, Jun 10 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012682 | OEIS-A291520

The arithmetic function uhat(n,4,2).

Conjectures from _Chai Wah Wu_, Jun 10 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012683 | OEIS-A291545

a(n) is the smallest integer k > n such that (k+1)(k+2)...(2k-n)/(k(k-1)...(k-n+1)) is an integer.

A lower bound is given by a(n) >= A104138(j) + j where j = floor((n+1)/2) and A104138(j) is the smallest prime that is followed by j or more nonprimes. Conjecture: this bound is tight for all n > 6. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Aug 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012684 | OEIS-A291561

Diagonal in triangle A291560: a(n) = -A291560(n+1, n) for n>=1.

Conjecture: a(n) = 4^n*gamma(-1/2 + n)*gamma(3/2 + n)*n/(3*Pi). - _Thomas Scheuerle_, Jan 27 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012685 | OEIS-A291568

The arithmetic function uhat(n,5,5).

Conjectures from _Chai Wah Wu_, Jun 10 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012686 | OEIS-A291574

The arithmetic function uhat(n,6,6).

Conjectures from _Chai Wah Wu_, Jun 10 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012687 | OEIS-A291584

The arithmetic function uhat(n,8,8).

Conjectures from _Chai Wah Wu_, Jun 10 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Aug 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012688 | OEIS-A291596

Column 6 of A060244.

Conjecture: column k of A060244 is asymptotic to (6*n)^((k-1)/2) * A000041(n) / (Pi^(k-1) * k!) ~ 2^((k-5)/2) * 3^((k-2)/2) * n^((k-3)/2) * exp(Pi*sqrt(2*n/3)) / (Pi^(k-1) * k!).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Aug 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012689 | OEIS-A291620

Branch term s_n(b), b > 1 of equivalence classes of prime sequences {s_n(k)} for k > 0 derived by records of first differences of Rowland-like recurrences with increasing even start values >= 4.

If the conjecture of an inverse tree of primes with the leaves in A291528 using the same index n holds, except a(1)=0 all terms a(n) == 0 indicates that the branch point is not yet found.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Steiner_, Aug 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012690 | OEIS-A291624

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that p = x + 2*y + 5*z, p - 2 and p + 4 are all prime.

Conjecture: a(n) > 0 for all n > 1 not divisible by 4. See also A291635 for a stronger conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012691 | OEIS-A291635

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that p = x + 2*y + 5*z, p - 2, p + 4 and p + 10 are all prime.

Conjecture: a(n) > 0 for all n > 1 not divisible by 4. This is stronger than the conjecture in A291624. Obviously, it implies that there are infinitely many prime quadruples (p-2, p, p+4, p+10).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012692 | OEIS-A291657

Primes p such that p is a primitive root modulo prime(p).

The conjecture in A291615 implies that the current sequence has infinitely many terms. In fact, if there are only finitely many primes p with p a primitive root modulo prime(p) and we let P denote the product of all such primes, then by Dirichlet's theorem there is a prime q == 1 (mod 4*P) and hence any prime p with p a primitive root modulo prime(p) is a quadratic residue modulo q and hence not a primitive root modulo q. Conjecture: a(n)/(n*log(n)) has a positive limit as n tends to the infinity. Equivalently, all the terms in this sequence form a subset of the set of all primes with positive asymptotic density.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012693 | OEIS-A291690

Least positive integer g which is a primitive root modulo prime(n) and also a primitive root modulo prime(n+1).

Conjecture: (i) For any distinct primes p and q, there is a positive integer g not exceeding sqrt(4*p*q+1) such that g is a primitive root modulo p and also a primitive root modulo q. We may require further that g < sqrt(p*q) if {p,q} is not among the 15 pairs {2,3}, {2,11}, {2,13}, {2,59}, {2,131}, {2,181}, {3,7}, {3,31}, {3,79}, {3,191}, {3,199}, {5,271}, {7,11}, {7,13} and {7,71}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012694 | OEIS-A291698

a(n) = [x^n] Product_{k>=1} (1 + n*x^k).

Conjecture: a(n) ~ exp(sqrt(2*(log(n)^2 + Pi^2/3)*n)) * (log(n)^2 + Pi^2/3)^(1/4) / (sqrt(Pi) * (2*n)^(5/4)). - _Vaclav Kotesovec_, Sep 15 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Aug 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012695 | OEIS-A291712

Lexicographically earliest sequence of positive terms such that, for any m and n > 0, if m < n then a(m) != a(n) or a(m+1) != a(n+1), and if n = least k > m such that a(k) = a(m) then m and n have a different parity.

Conjecturally, (a(n), a(n+1)) uniquely runs over all pairs of positive integers (this is the motivation for this sequence). - A290633 whose pairs of consecutive terms (conjecturally) run over all pairs of noncoprime positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Aug 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012696 | OEIS-A291775

Domination number of the n-Sierpinski carpet graph.

Conjecture: For n>1, a(n) = 2^(3n-2) + 2. There is an independent dominating set of this size consisting of the vertices on every third diagonal and two corner vertices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Aug 31 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012697 | OEIS-A291790

Numbers whose trajectory under iteration of the map k -> (sigma(k)+phi(k))/2 consists only of integers and is unbounded.

It would be nice to have a proof that these trajectories are integral and unbounded, or, of course, that they eventually reach a fractional value (and die), or reach a prime (which is then a fixed point). (Cf. A291787.) If either of the last two things happen, then that value of n will be removed from the sequence. AT PRESENT ALL TERMS ARE CONJECTURAL.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012698 | OEIS-A291834

First positions of records in A252665.

Conjectures based on a(n) < 10^7:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Sep 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012699 | OEIS-A291855

Numbers k such that gpf(2^k - 1) - 1 is not divisible by k.

Conjecture: 4 * A001153(m) for m > 3 is a subsequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_ and _Altug Alkan_, Sep 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012700 | OEIS-A291876

Consider the graph with one central vertex connected to three outer vertices (a star graph). Then, a(n) is the minimum number of moves required to transfer a stack of n pegs from one outer vertex to another outer vertex, moving pegs to adjacent vertices, following the rules of the Towers of Hanoi.

Conjecturally, a(n) = 2*A259823(n). This conjecture was proved by Thierry Bousch, see link. - _Paul Zimmermann_, Oct 05 2015

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric M. Schmidt_, Sep 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012701 | OEIS-A291897

Numerator of E(2*n-1,n), where E(n,x) is the Euler polynomial.

Conjecture: a(n) is divisible by (2*n-1)^2. _Robert G. Wilson v_ verified this conjecture up to 5000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012702 | OEIS-A291934

Largest k achievable by the iteration k->(sigma(k)+phi(k))/2 started at k=n, with termination when sigma(k) becomes odd.

See A291914 for types of termination. The first term with unknown conjecturally unbounded behavior is a(270). More values of n with this type of behavior are given in A291790.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Sep 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012703 | OEIS-A291982

a(n) = Euler(n, n+1) * 2^valuation(n+1, 2), where Euler(n, x) denotes the Euler polynomial.

Conjecture: If n >= 2 is even then n*(n+1) divides a(n). This conjecture was inspired by _Vladimir Shevelev_'s conjecture in A291897.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Sep 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012704 | OEIS-A292034

Linear divisibility sequence based on Salem number of order 4 (case t=6, see formula).

Conjectures from _Colin Barker_, Dec 17 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Sep 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012705 | OEIS-A292035

Linear divisibility sequence based on Salem number of order 4 (case t=7, see formula).

Conjectures from _Colin Barker_, Dec 17 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Sep 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012706 | OEIS-A292042

G.f.: Re((i*x; x)_inf), where (a; q)_inf is the q-Pochhammer symbol, i = sqrt(-1).

Conjectural g.f.: A(x) = (1/2)*Sum_{n >= 0} (-x)^(n*(n-1)/2)/Product_{k = 1..n} (1 - x^k). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Sep 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012707 | OEIS-A292082

Primes p such that (p^2 - 1) / 2 is a square (A000290).

Conjecture: sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Sep 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012708 | OEIS-A292083

Odd numbers k that can be uniquely expressed as (2*p + 1)/q for some prime p and some prime q < 2*k.

Conjecture: the sequence is finite and complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Sep 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012709 | OEIS-A292084

a(n) = least prime p such that 2*p + 1 equals (2*n - 1)*q where q is a prime, or 0 if no such p exists.

Conjecture: 7 <= a(n) <= 4*n^2 - 5*n + 1 for n > 1. This conjecture implies that for every odd k > 1 there exist two primes p and q < 2*k such that k = (2*p + 1)/q.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Sep 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012710 | OEIS-A292114

List of numbers n such that A039655(n) reaches a new record high.

Increasingly many of the values are of the form m*p with a (large) prime p and a smooth m, often m = 2^k (for a(n), n = 12, 14, 21, 23, 26, 28, 29, ...) or m = 2^k*3^k' (n = 7, 9, 19, 22, 30, ...) or m = 2^k*5^k' (n = 20, 25, ...). I conjecture that almost all terms are even. Also, for most terms (n = 1, 2, 3, 4, 5, 7, 10, 12, 14, 15, 16, 17, 19, 20, 21, 23, 24, 25, 26, 29, 30, 31, ...), either a(n)-1 or a(n)+1 has at most 2 prime divisors. - _M. F. Hasler_, Sep 25 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012711 | OEIS-A292177

G.f.: Limit_{K->oo} Sum_{n=-oo..+oo} x^(n-K) * (1 - x^n + n*(n+1)/6 * x^(n+K))^n.

a(p) = 0 for odd prime p (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012712 | OEIS-A292180

G.f.: Sum_{n=-oo..+oo} (1 + x^n)^n / (1 - x^n)^n, ignoring the constant term.

a((2*n-1)^2)/4 is odd for n>=1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012713 | OEIS-A292217

Conjectured list of numbers in increasing order that belong to sociable cycles of length greater than 2 in which the sum of the cycle is divisible by 10.

Conjectured list of numbers in increasing order that belong to sociable cycles of length greater than 2 in which the sum of the cycle is divisible by 10. Up to the known 1593 sociable number cycles, 96.1% of the sociable number cycles satisfy this condition (up to the first 10 sociable number cycles: 40%; up to the first 100 sociable number cycles: 77%; up to the first 500 sociable number cycles: 92%, and up to the first 1000 sociable number cycles: 94.9%). So the conjecture here is that as the number of sociable number cycles increases, the percentage of the sums of the sociable number cycles divisible by 10 approaches 100%. Notice that the sums of amicable pairs are similarly often divisible by 10, but are not included here (see A291422).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zoltan Galantai_, Sep 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012714 | OEIS-A292273

For odd n: a(n) = 0, and for even n: a(n) = -mu(n), where mu is Moebius function (A008683).

Conjecture: a(n) = Re(A008683(n)*(i^n)). - _Mats Granvik_, Jul 06 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012715 | OEIS-A292304

a(n) = [x^n] Product_{k>=1} (1 + n^2*x^k).

Conjecture: a(n) ~ exp(2*sqrt((Pi^2/6 + 2*log(n)^2)*n)) * (Pi^2/6 + 2*log(n)^2)^(1/4) / (2 * sqrt(Pi) * n^(7/4)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Sep 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012716 | OEIS-A292306

a(n) = [x^n] Product_{k>=1} (1 + n^n*x^k).

Conjecture: log(a(n)) ~ (sqrt(2)*n^(3/2) - n/2)*log(n). - _Vaclav Kotesovec_, Aug 22 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Sep 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012717 | OEIS-A292348

"Pri-most" numbers: the majority of bits in the binary representation of these numbers satisfy the following: complementing this bit produces a prime number.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Dec 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012718 | OEIS-A292349

Pri-most primes: primes p such that the majority of bits in the binary representation of p satisfy the following: complementing this bit produces a prime number.

Conjecture: the sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Dec 07 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012719 | OEIS-A292359

E.g.f.: Product_{k>=1} (1 - x^k/k).

Conjecture: a(n) ~ -(-1)^n * n! / n^2. - _Vaclav Kotesovec_, Sep 15 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Sep 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012720 | OEIS-A292364

Composites m such that each prime factor p > m of 2^m - 1 is a primitive prime factor of 2^m - 1.

Let S represent the set of numbers such that the greatest prime factor of 2^n-1 is less than n^2. S begins {2,3,4,6,8,9,10,11,12,14,15,18,20,21,24,28,30,36,48,60} (obtained from A005420), and I conjecture that there are no further terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Sep 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012721 | OEIS-A292389

a(n) = SumXOR_{k=1..n} A292388(k) (where SumXOR is the analog of summation under the binary XOR operation).

Conjecturally, this sequence contains infinitely many 2's.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Sep 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012722 | OEIS-A292394

G.f. A(x) satisfies: [x^n] A(x/A(x)^(n^2)) = 0 for n>1.

a(n) is divisible by n^2 for n>=1 (conjecture): A292395(n) = a(n)/n^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012723 | OEIS-A292395

a(n) = A292394(n)/n^2 for n>=1.

It is conjectured that this sequence consists entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012724 | OEIS-A292414

a(n) = [x^n] Product_{k>=1} (1 + 2^n*x^k).

Conjecture: log(a(n)) ~ sqrt(2)*log(2)*n^(3/2). - _Vaclav Kotesovec_, Aug 22 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Sep 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012725 | OEIS-A292422

Numbers of the form x = 2^k*p*q, p,q odd primes, such that sigma(x)/phi(x) = 4.

A subset of A068390 and A020492 (balanced numbers). Conjectured to be infinite by Broughan and Zhou.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_, Sep 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012726 | OEIS-A292461

Expansion of (1 - x - x^2 + sqrt((1 - x - x^2)^2 - 4*x^3))/2 in powers of x.

Conjecture D-finite with recurrence: n*a(n) +(-2*n+3)*a(n-1) +(-n+3)*a(n-2) +(-2*n+9)*a(n-3) +(n-6)*a(n-4)=0. - _R. J. Mathar_, Jan 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Sep 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012727 | OEIS-A292507

Number of partitions of n with up to n distinct kinds of 1.

Conjecture: log(a(n)) ~ log(2)*n + Pi*sqrt(n/3) - 3*log(n)/2. - _Vaclav Kotesovec_, May 11 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Sep 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012728 | OEIS-A292510

a(n) = smallest k >= 1 such that {1, p(n,2), p(n,3), ..., p(n,k)} can be partitioned into two sets with equal sums, where p(n,m) is m-th n-gonal number.

Conjecture: a(n) = 7 for n > 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Sep 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012729 | OEIS-A292531

a(n) = 0 if n is a power of 2. Otherwise, product of 2 numbers nearest n that have more 2's in their prime factorization than n.

Proposition 2) is true. Proof: 0 mod 2 = 0, so the conjecture trivially holds when n is a power of 2. For n not a power of 2, a(n) has by definition the repeated prime factor 2^2 and so is divisible by 8 when a(n) > 4. - _Felix Fröhlich_, Sep 19 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Sep 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012730 | OEIS-A292559

Composite numbers m such that lpf(2^m - 1) == 1 (mod m).

Conjecture: there are no numbers k in this sequence such that, for each prime factor q of 2^k - 1, q == 1 (mod k). - _Thomas Ordowski_, Sep 20 2017 Note: if all prime factors q of 2^k - 1 are q == 1 (mod k), then 2^k - 1 == 1 (mod k), thus 2^k == 2 (mod k), so k is a pseudoprime. The pseudoprime k = a(42) = 4369 = 17*257 is not a counterexample to this conjecture. A pseudoprime k = P*Q such that both 2^P - 1 and 2^Q - 1 are primes would be a counterexample, but the known Mersenne primes do not give such k. - _Thomas Ordowski_, Oct 02 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_ and _Thomas Ordowski_, Sep 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012731 | OEIS-A292569

Least pandigital number which sums up with the n-th pandigital number A171102(n) to another pandigital number.

For all n, a(n) is well defined, because to any pandigital number N = A171102(n) we can add the number M(N) = 123456789*10^k with k = # digits of N, which is pandigital (in the above extended sense) as well as is the sum N + M(N) (equal to the concatenation of 123456789 and N). In practice, there are much smaller solutions. We conjecture that there is always a 10-digit solution a(n) < 10^10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Sep 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012732 | OEIS-A292570

Least k > 0 such that A171102(n) + A171102(k) is again a term of A171102, the pandigital numbers (having each digit from '0' to '9' at least once).

For all n, a(n) is well defined, because to any pandigital number N = A171102(n) we can add the number M(N) = 123456789*10^k with k = # digits of N, which is pandigital (in the above extended sense) as well as is the sum N + M(N). In practice, there are much smaller solutions. We conjecture that there is always a 10-digit solution a(n) < 10^10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Sep 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012733 | OEIS-A292689

Decimal values of the antidiagonals of the Sierpinski carpet considered as binary numbers.

a(k+1) = 2*a(k)+1 for all k in A003462 = (1, 4, 13, 40, 121, 364, ...). (Conjectured.) - _R. J. Cano_, Oct 25 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Oct 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012734 | OEIS-A292726

a(n) is the number of states that can be achieved when starting from n piles each containing one stone, where any number of stones can be transferred between piles that start with the same number of stones.

Conjecture: A000041(n) - a(n) = 0 if and only if n is a power of 2, and Conjecture: A000041(n) - a(n) = 1 if and only if n is an odd prime. Consider the inverse operation: either divide an even stack in two, or replace two stacks of equal parity with their averages. The states which have no inverse moves are precisely those with all parts equal and odd. If n is a power of two, its only odd divisor is 1 and so every state can be inverted to the all-ones state, and if n is an odd prime then the only states which cannot be inverted are the all-ones state and [n]. If n has an odd divisor d with 1 < d < n, then unreachable states include the all-d's state and any state obtainable from it by combining d's in pairs, of which there is at least one. Therefore, both conjectures are true. - _Charlie Neder_, Jan 28 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Sep 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012735 | OEIS-A292727

a(n) is the number of states that cannot be achieved when starting from n piles each containing one stone, where stones can be transferred between piles only when they start with the same number of stones.

Conjecture: a(n) = 0 if and only if n is a power of 2. Conjecture: a(n) = 1 if and only if n is an odd prime. Conjecture: a(4p) = p+4, a(8p) = 2p+20. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Sep 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012736 | OEIS-A292728

a(n) is the number of terminal states that can be achieved via the following algorithm: start with n piles each containing one stone; stones can be transferred between piles only when the piles start with the same number of stones.

Conjecture: a(n) = A000009(n) if and only if n is a power of 2. Conjecture: A000009(n) - a(n) = 1 if and only if n is an odd prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Sep 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012737 | OEIS-A292740

Indices k such that A292547(k) = 0.

Conjecture: for k > 212594 there are no more terms in this sequence (tested for k < 63000000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Sep 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012738 | OEIS-A292764

Minimal number of moves for the cyclic variant of Hanoi's tower for 4 pegs and n disks, with the final peg two steps away.

Conjecture: for n >= 9, a(n) = a(n-1)+2*a(n-3)+c(n), where c(n) = 18 for odd n and c(n) = 14 for even n. - _Paul Zimmermann_, Oct 23 2017 Conjectures from _Colin Barker_, Oct 25 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 27 2017, following a suggestion from _Paul Zimmermann_ who computed the terms through a(16) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012739 | OEIS-A292766

Numbers n whose trajectory under iteration of the map k -> (sigma(k)+phi(k))/2 consists only of integers and is unbounded, excluding numbers n whose trajectory merges with the trajectory of a smaller number.

These are the "seeds" in A291790, that is, every number which blows up under iteration of the map k -> (sigma(k)+phi(k))/2 belongs to one of these trajectories. AT PRESENT ALL TERMS ARE CONJECTURAL.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 27 2017, based on emails from _Sean A. Irvine_, Sep 14 2017, who computed a(1)-a(9), and _Hans Havermann_, same date, who computed a(10)-a(29). _Hugo Pfoertner_ also computed many of these terms | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012740 | OEIS-A292782

a(n) = E(2n,n)/2, where E(n,x) is the Euler polynomial.

Conjecture. For n >= 2, a(n) is divisible by n(n-1)/2, moreover, for odd n, a(n) is divisible by n^2(n-1)/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012741 | OEIS-A292794

Numbers not congruent to A000045(k) mod A000045(k+1) for all k > 1.

It is conjectured that A035105(n) is always a member of this sequence for n >= 4 but this remains unproved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ely Golden_, Sep 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012742 | OEIS-A292814

G.f.: A(x) satisfies: A(6*x - 5*A(x)) = x - 16*x^2.

Conjecture: In general, if k > 0 and g.f. A(x) satisfies A((k+2)*x - (k+1)*A(x)) = x - (k*x)^2, then a(n) ~ (-1)^n * c(k) * (k*(k+1)/log(k+1))^n * n! / n^((k-1)/(k+1) + (k-2)*log(k+1)/k), where c(k) is a constant. - _Vaclav Kotesovec_, Oct 09 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012743 | OEIS-A292821

Decimal expansion of Product_{k>=1} (1 + exp(-2*Pi*k)).

Conjecture: Equals Sum_{k>=0} A035457(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Sep 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012744 | OEIS-A292824

Decimal expansion of Product_{k>=1} (1 - exp(-Pi*(2*k-1))).

Conjecture: Equals Sum_{k>=0} A081362(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Sep 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012745 | OEIS-A292867

Indices of records in A243823.

Some conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Oct 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012746 | OEIS-A292868

Records in A243823.

See A292867 for comments, linked tables, and conjectures. - _Michael De Vlieger_, Nov 17 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Oct 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012747 | OEIS-A292876

Irregular table whose n-th row lists all k such that A039654(k) = prime(n).

Indeed, A039653(x) >= x with equality iff x is prime, and all orbits of A039653 are conjectured to end in such a fixed point prime p = A039654(k) for any k in this orbit.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Sep 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012748 | OEIS-A292877

G.f. A(x) satisfies: [x^(n-1)] 1/A(x)^(n^2) = 0 for n>2.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012749 | OEIS-A292911

Numbers n such that A291897(n) is divisible by (2*n-1)^3.

Conjecture: Every prime of the form 4*k+1 (A002144) is contained in the sequence {2*a(n)-1}. The author's former conjecture that, for n>=2 the numbers {2*a(n)-1} are consecutive primes of the form 4*k+1, was disproved at n = 553 by _Peter J. C. Moses_. (553*2 - 1 = 1105 is the smallest term which is a product of three distinct (4*k+1)-primes). - _Vladimir Shevelev_, Sep 27 2017 If the conjecture is true, then for n>=2, a(n) <= (A002144(n-1) + 1)/2 (the equality holds up to 90).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Shevelev_, Sep 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012750 | OEIS-A292990

Numbers whose absolute difference from a triangular number is never a prime.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Dec 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012751 | OEIS-A292997

Irregular triangle read by rows: lexicographically earliest, minimum size sets, S_n, such that {1, 2, ... n} is a subset of the union of S_n with the sums of distinct pairs in S_n and the differences of pairs in S_n.

Conjecture: A225687 gives the row lengths.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Sep 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012752 | OEIS-A293010

a(n) is the smallest x > 2 to satisfy pi(x-1)/(x-1)^n < pi(x)/x^n, where pi(x) is the prime counting function (A000720).

Conjecture: a(n) exists for all n, that is, for all n, there exists at least one integer which satisfies the inequality. Conjecture above is true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Josh Marza_, Sep 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012753 | OEIS-A293011

a(n) is the smallest positive k such that f(k) = n*g(k) where f = A007953 and g = A000120, or 0 if no such k exists.

Conjecture: a(n) > 0 for all n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Sep 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012754 | OEIS-A293057

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-2) + 2, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012755 | OEIS-A293058

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-2) + 3, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012756 | OEIS-A293076

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-2), where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012757 | OEIS-A293129

L.g.f.: Sum_{n=-oo..+oo} (x - x^(2*n-1))^(2*n-1) / (2*n-1).

a(2^n + 1) = 1 for n >= 1 (conjecture). The conjecture is true (see Fried link). - _Sela Fried_, Dec 18 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012758 | OEIS-A293136

Irregular triangle T(n,k) read by rows: T(n,k) is the number of strongly unimodal compositions of n (A059618) into k parts.

Conjecture: index k of last nonzero entry in row n of is A293137(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Oct 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012759 | OEIS-A293137

a(0) = 0, and a(n) = floor(2*sqrt(n)) - 1 for n >= 1.

Conjecture: a(n) is index k of last nonzero entry in row n of A293136.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Oct 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012760 | OEIS-A293142

Largest nonrepunit base-n circular prime (conjectured).

Largest nonrepunit base-n circular prime (conjectured). a(3)-a(6) were found via a brute-force approach searching from the largest prime with 12 base-n digits backwards. The number of base-n digits in a(n) for n = 3, 4, 5, 6 is 2, 5, 5, 3, respectively. Since this is much shorter than 12 digits, it is conjectured that the terms are the maximal circular primes for those bases. This also verifies that no circular primes with a length between A055642(a(n)) and 13 digits exist in bases 3, 4, 5 and 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Oct 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012761 | OEIS-A293181

Irregular triangle read by rows: T(n,k) is the number of k-partitions of {1..2n} that are invariant under a permutation consisting of n 2-cycles (1 <= k <= 2n).

Conjecture: T(n,k) = R(n,k)-R(n,k-1), with R(n,k) = Sum_{m=0..k} m^n*A000085(m)*A038205(k-m)/(m!*(k-m)!). - _Mikhail Kurkov_, Jun 26 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Howroyd_, Oct 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012762 | OEIS-A293190

a(n) = |{A001597(n) <= k <= A001597(n+1): 2*k^2-1 is prime}|.

Conjecture: (i) a(n) > 0 for all n > 0. In other words, for any perfect powers x^m and y^n with 0 < x^m < y^n, there is an integer z with x^m <= z <= y^n such that 2*z^2 - 1 is prime. Compare this with the Redmond-Sun conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012763 | OEIS-A293203

Numbers k such that A090086(k), the smallest pseudoprime to base k (not necessarily exceeding k), is a Carmichael number.

Andrzej Schinzel proved that this sequence is infinite. Conjecture: if A090086(n) is a Carmichael number k, then k < n. - _Thomas Ordowski_, Aug 08 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Oct 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012764 | OEIS-A293213

Primes p with phi(p-1) a primitive root modulo p, where phi(.) is Euler's totient function (A000010).

Conjecture: The sequence contains infinitely many terms. Moreover, the number of primes p <= x with phi(p-1) a primitive root modulo p is asymptotically equivalent to c*x/(log x) as x tends to the infinity, where c is a constant with 0.36 < c < 0.37.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012765 | OEIS-A293238

Decimal expansion of the escape probability for a random walk on the 3D bcc lattice.

Conjecture: Equals Sum_{k>=0} A004405(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrei Zabolotskii_, Oct 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012766 | OEIS-A293239

Number of terms in the fully expanded n-th derivative of x^x.

Conjecture: the 2nd differences are eventually periodic: 1, 1, 1, 0, 2, 1, 0, 1, [2, 1, 1, 0]. Conjecture: a(n) ~ n^2/2. - _Vaclav Kotesovec_, Oct 05 2017 Conjectures from _Colin Barker_, Oct 05 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Reshetnikov_, Oct 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012767 | OEIS-A293273

a(n) is the smallest positive k <> n such that f(k) is divisible by f(n) where f = A005132, or 0 if no such k exists.

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Oct 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012768 | OEIS-A293312

Rectangular array read by antidiagonals: A(n,k) = tr((M_n)^k), k >= 0, where M_n is the n X n matrix M_1 = {{1}}, M_n = {{0,...,0,1},{0,...,0,1,1},...,{0,1,...,1},{1,...,1}}, n > 1, and tr(.) is the trace.

Conjecture: For all n >= 1, for all k >= 2, A(n, k) = A293311(k, n); i.e., A(n, k) = number of magic labelings of the graph LOOP X C_k with magic sum n - 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _L. Edson Jeffery_, Oct 10 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012769 | OEIS-A293316

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-2) + 1, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012770 | OEIS-A293349

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-2) + n, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012771 | OEIS-A293350

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-2) + 2n, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012772 | OEIS-A293351

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-2) + n -1, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012773 | OEIS-A293358

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-1), where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012774 | OEIS-A293379

G.f.: exp(Sum_{n>=1} A052886(n) * x^n/n).

This sequence is motivated by the following conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012775 | OEIS-A293406

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-1) + 1, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio. See A293358 for a guide to related sequences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012776 | OEIS-A293452

Triangle T(n,k) read by rows: T(n,k) is the number of iterations to reach a final state for an n X k lattice of sandpiles on a torus according to rules specified in A249872.

Conjecture: T(n,1) = A023855(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Oct 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012777 | OEIS-A293456

G.f.: A(x) satisfies: A(9*x - 8*A(x)) = x - 49*x^2.

Conjecture: In general, if k > 0 and g.f. A(x) satisfies A((k+2)*x - (k+1)*A(x)) = x - (k*x)^2, then a(n) ~ (-1)^n * c(k) * (k*(k+1)/log(k+1))^n * n! / n^((k-1)/(k+1) + (k-2)*log(k+1)/k), where c(k) is a constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Oct 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012778 | OEIS-A293462

Let A_n be a square n X n matrix with entries A_n(i,j)=1 if i+j is a perfect power, and A_n(i,j)=0 otherwise. Then a(n) counts the 1's in A_n.

Conjecture: The golden ratio/golden conjugate are eigenvalues of A_n if and only if n=6, 8 or 9. This has been verified up to n=500. Conjecture: the sequence increases monotonically. - _Robert G. Wilson v_, Oct 09 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anthony Hernandez_, Oct 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012779 | OEIS-A293555

Indices of records in A243822.

Conjecture: all terms of a(n) with n > 92 also appear in A244052, and all terms in A244052 greater than a(92) = 6276270 appear in a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Oct 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012780 | OEIS-A293556

Records in A243822.

See A293555 for comments, linked tables, and conjecture. - _Michael De Vlieger_, Nov 17 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Oct 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012781 | OEIS-A293636

a(1) = a(2) = 1; a(n) = (Sum_{i|(n-1)} a(i)) + Sum_{j|(n-2)} a(j).

a(n) ~ c*phi^n, where phi is the golden ratio (A001622) and c = 1.83226227102725... (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Oct 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012782 | OEIS-A293652

a(n) is the smallest prime number whose a056240-type is n (see Comments).

a(20), a(21) > 14 * 10^9. Conjecture: a(k) > 14 * 10^9 for k > 22. - _David A. Corneth_, Mar 25 2018 a(20), a(21) computed on the basis of the above conjecture. Note that A321983 records the smallest composite number whose sum of prime divisors (with repetition) is a(n). - _David James Sycamore_, Nov 30 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Feb 06 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012783 | OEIS-A293657

Base-4 circular primes that are not base-4 repunits.

Conjecture: The sequence is finite, with 1013 being the last term (see A293142).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Oct 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012784 | OEIS-A293658

Base-5 circular primes that are not base-5 repunits.

Conjecture: The sequence is finite, with 3121 being the last term (see A293142).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Oct 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012785 | OEIS-A293659

Base-6 circular primes that are not base-6 repunits.

Conjecture: The sequence is finite, with 211 being the last term (see A293142).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Oct 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012786 | OEIS-A293660

Base-7 circular primes that are not base-7 repunits.

Conjecture: The sequence is finite, with 13143449029 being the last term. - [Comment extended by _Felix Fröhlich_, May 30 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Dec 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012787 | OEIS-A293661

Base-8 circular primes that are not base-8 repunits.

Conjecture: The sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Dec 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012788 | OEIS-A293662

Base-9 circular primes that are not base-9 repunits.

Conjecture: The sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Dec 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012789 | OEIS-A293663

Circular primes that are not repunits.

Conjecture: The sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Dec 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012790 | OEIS-A293688

Partial sums of A002251.

a(n) = (n+1)*A005206(n) where A005206 is the Hofstadter G-sequence. [Apparently this is a conjecture, rather than a theorem. - _N. J. A. Sloane_, Mar 03 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enrique Navarrete_, Oct 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012791 | OEIS-A293700

First differences of A293698.

Despite its apparent simplicity, it has interesting palindromic and periodic features and may be conjectured not to be represented in a closed form.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V.J. Pohjola_, Oct 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012792 | OEIS-A293715

Numbers k such that A007755(k) is prime.

Shapiro conjectured that A007755(n) is prime for all n > 1, and verified it up to n = 10. Mills showed that A007755(34)=(2^16+1)^2 is composite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Oct 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012793 | OEIS-A293756

a(n) = smallest number k with n prime factors such that d + k/d is prime for every d | k.

If the k-tuple conjecture is true, then this sequence is infinite. - Carl Pomerance, Nov 12 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Nov 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012794 | OEIS-A293765

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-1) + 2, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio. See A293358 for a guide to related sequences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012795 | OEIS-A293766

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-1) + 3, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio. See A293358 for a guide to related sequences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012796 | OEIS-A293771

Minimum number of steps needed to compute n using a machine that can read, write and add starting with the number 1.

Conjecture: reading from memory (operation 2) is never needed to get to a number in the minimal number of steps. Additional conjectures and comments by _Glen Whitney_, Oct 12 2021: (Start) Conjecture II: For each n, there is a minimal-length program for n that stores numbers in memory in increasing order. Conjecture III: For each n, there is a minimal-length program such that the difference between successive numbers stored in memory is strictly increasing. All three conjectures are empirically verified for all programs of length 23 or less, and all values of n up to 2326. However, note that if you are allowed to specify the set of numbers that must be stored in memory, then the first conjecture fails for memory {1,3,9,27,30,54} and conjecture II fails for {1,3,9,27,30,54,60}. (These sequences of numbers in memory are similar to addition chains, see A003313.) Hence, a proof of the first conjecture might need to involve showing that every n has a "good" sequence of numbers that can be stored in memory to produce n, avoiding the need to invoke the read operation. Conjecture III supplies one speculative possibility of a property that might fill the role of "good." A proof of any of these conjectures would tremendously speed up computation of a(n) as compared to brute force. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Floris P. van Doorn_, Oct 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012797 | OEIS-A293821

Number of integer-sided quadrilaterals having perimeter n, modulo rotations but not reflections.

Conjectures from _Colin Barker_, Nov 01 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James East_, Oct 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012798 | OEIS-A293822

Number of integer-sided pentagons having perimeter n, modulo rotations but not reflections.

G.f.: x^5*(1 + x - x^2 + 2*x^3 + 7*x^4 + 2*x^5 - 2*x^6 + x^8) / ((1 - x)^5*(1 + x)^4*(1 + x + x^2 + x^3 + x^4)) (conjectured). - _Colin Barker_, Nov 01 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James East_, Oct 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012799 | OEIS-A293823

Number of integer-sided hexagons having perimeter n, modulo rotations but not reflections.

G.f.: x^6*(1 + x + 5*x^3 + 10*x^4 + 7*x^5 + 3*x^6 + 6*x^7 + 4*x^8 + 2*x^9) / ((1 - x)^6*(1 + x)^5*(1 - x + x^2)*(1 + x + x^2)^2) (conjectured). - _Colin Barker_, Nov 01 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James East_, Oct 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012800 | OEIS-A293881

Number T(n,k) of linear chord diagrams having n chords and minimal chord length k (or k=0 if n=0); triangle T(n,k), n>=0, 0<=k<=n, read by rows.

Conjecture: column k>0 is asymptotic to (exp(-k+1) - exp(-k)) * 2^(n + 1/2) * n^n / exp(n). - _Vaclav Kotesovec_, Oct 25 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Oct 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012801 | OEIS-A293929

Smallest number in base 10 that cannot be collapsed to a single digit using fewer than n plus signs.

Butler et al. (2016) conjectured that collapsing each term x by simply inserting a plus sign in the "middle" of the decimal expansion and adding would require in the order of log (log x) plus signs. Conjecture: Digital root for terms a(n) > 0 is 1. - _Simon Demers_ and _J. Stauduhar_, Nov 16 2017 a(n) <= ((a(n-1)-1)^2)/3 + a(n-2) for n > 1 (conjectured). This would provide a relatively tight upper bound on a(n). If the Demers-Stauduhar conjecture in the Comments turns out to be true, this upper bound will always be an integer. - _Simon Demers_, Nov 29 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Demers_, Oct 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012802 | OEIS-A293982

Length (= size) of the orbit of n under iterations of A293975: x -> x/2 if even, x + nextprime(x) if odd; or -1 if the orbit is infinite.

It is conjectured that for f = A293975, the trajectory (f^k(x); k >= 0) ends in the cycle 1 -> 3 -> 8 -> 4 -> 2 -> 1 for any starting value x.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Nov 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012803 | OEIS-A294064

Numbers k such that 2*k - 3, 2*k + 3, 3*k - 2, 3*k + 2 are primes.

Conjecture: The Sum_{n>=1} 1/a(n) = 0.57... converges.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimitris Valianatos_, Oct 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012804 | OEIS-A294074

Primes of the form p*2^k + 1, where p is an odd prime and k is odd.

I conjecture that a number of the form p*2^k + 1 (with odd prime p and odd k) belongs to this sequence if and only if p*2^k + 1 divides (p + 2)^(p*2^k) - 1. This conjecture has been verified for n up to 10^10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Feb 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012805 | OEIS-A294089

a(n) = b(n) if b(n) := (Sum_{k>=1}(k/5)/k^(2n))*sqrt(5)^(floor((n+1)/2)*floor((n+11)/2)-1)/(24*zeta(2n)) is an integer, and a(n) = 0 otherwise, where (k/5) is the Legendre symbol.

Conjecture: For each n = 1,2,3,... the number (Sum_{k>=1}(k/5)/k^(2n))*sqrt(5)/Pi^(2n) is rational, where (k/5) is the Legendre symbol. Moreover, a(n) is a nonzero integer for every positive integer n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012806 | OEIS-A294112

Practical numbers z such that z^2 = x^2 + y^2 for some practical numbers x and y with gcd(x,y,z) = 4.

Conjecture: The sequence has infinitely many terms. Also, there are infinitely many Pythagorean triples (x,y,z) with x,y,z all practical and gcd(x,y,z) = 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012807 | OEIS-A294115

G.f.: exp(Sum_{n>=1} L(n) * x^n/n), where Sum_{n>=1} L(n) * x^n/n! = Series_Reversion(log(1 + x/A(x))).

This sequence is motivated by the following conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012808 | OEIS-A294139

Sum of the areas of the distinct rectangles (and the areas of the squares on their sides) with positive integer sides such that L + W = n, W < L.

Conjectures from _Colin Barker_, Nov 01 2017: (Start) The first three conjectures of Barker are true. See links. - _Sela Fried_, Aug 11 2024.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Oct 31 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012809 | OEIS-A294159

Alternating row sums of triangle A291844.

Conjecture: D-finite with recurrence n*a(n) +(-3*n+1)*a(n-1) +2*(-n+3)*a(n-2) +2*(n-5)*a(n-3) +(n-4)*a(n-4) +(n-5)*a(n-5)=0. - _R. J. Mathar_, Jun 17 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Nov 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012810 | OEIS-A294172

Maximum value of the cyclic convolution of the first n positive integers with themselves.

Conjecture: a(n) = (n*(13 + 3*(-1)^n + 24*n + 14*n^2))/48, and then lim_{n -> infinity} a(n)/n^3 = 7/24. The conjectured formula is true. See links. - _Sela Fried_, Aug 13 2024. Conjectures from _Colin Barker_, Feb 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andres Cicuttin_, Feb 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012811 | OEIS-A294179

a(n) is the smallest k with n prime factors such that p^k == p (mod k) for every prime p dividing k.

Conjecture: the sequence contains only finitely many Carmichael numbers, A006931. What is the smallest n >= 3 for which a(n) is not a Carmichael number? For n >= 3, a(n) <= A006931(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Feb 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012812 | OEIS-A294185

Number of distinct lesser twin primes which are in Goldbach partitions of 2n.

Tomas Oliveira e Silva in 2012 experimentally confirmed that all even numbers <= 4*10^18 have at least one Goldbach partition (GP) with a prime 9781 or less. Detailed examination of all even numbers < 10^6 showed that the most popular prime in all GPs is 3 (78497 occurrences), then 5 (70328), then 7 (62185), then 11 (48582), then 13 (40916), then 17 (31091), then 19 (29791) -- all these primes are twin primes. These results gave rise to a hypothesis that twin primes should be rather frequent in GP, especially those relatively small.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marcin Barylski_, Feb 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012813 | OEIS-A294200

Primes p such that 2^p - 2 is a practical number.

Conjecture: The sequence has infinitely many terms. In other words, there are infinitely many practical numbers of the form 2^p - 2 with p prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012814 | OEIS-A294203

Number of partitions of n into distinct Lucas parts (A000204) greater than 1.

Positions of 0: 1, 2, 5, 6, 8, 9, 12, 13, ... = A287775(n) - 1 (conjecture). Proof of the 'positions of 0' conjecture: let (z(n))=1,2,5,6,8,9,12,... be the positions of 0. The crucial observation is that if a number n is the sum of distinct Lucas parts greater than 1, then n+1 is a sum of Lucas parts. This implies that (z(2n))=2,6,9,13,... is the sequence of numbers A054770 that are not a sum of Lucas numbers. We see there that Ian Agol proved that b(n):=A054770(n)=floor(phi*n)+2n-1. But then the sequence of first differences (b(n+1)-b(n)) equals the Fibonacci word on the alphabet {4,3}, yielding that (z(2n)-z(2n-1)) equals the Fibonacci word on {3,2}, and we already know that z(2n+1)-z(2n)=1 for all n. On the other hand, A287775 has the same first difference sequence given by A108103. Since A287775(1)=2, the conjecture follows. (End) Positions of 1: 0, 3, 4, 10, 15, 28, 44, 75, ... = A001350(n+1) - 1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Oct 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012815 | OEIS-A294205

a(1) = 2; for n > 1, a(n) is the least prime p not already in the sequence such that the Hamming distance between p and a(n-1) is 1.

Conjecture: this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Oct 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012816 | OEIS-A294224

Triangle read by rows: T(n,k) is the number of graphs with n vertices and skewness k (n >= 1 and k >= 0).

Length of the n-th row kmax(n) is A000124(n-4) for n > 3 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Oct 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012817 | OEIS-A294225

Practical numbers q with q + 2 and q^2 + 2 both practical.

Conjecture: The sequence has infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012818 | OEIS-A294307

Positive integers m with m^k - 1 (k = 1,...,13) all practical.

Conjecture: For any positive integer n, there are infinitely many positive integers m with m^k - 1 (k = 1,...,n) all practical.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012819 | OEIS-A294359

a(n) = [x^n] F(x)^(-(n+1)^2) such that F(x) = F(x^2) + x*F(x^4), where F(x) = Sum_{n>=0} x^A003714(n) and A003714 is the Fibbinary numbers.

It is conjectured that all terms are even after the initial '1'.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012820 | OEIS-A294361

E.g.f.: exp(Sum_{n>=1} sigma(n) * x^n).

It appears that the sequence taken modulo 10 is periodic with period 5. More generally, we conjecture that for k = 2,3,4,... the sequence a(n+k) - a(n) is divisible by k: if true, then for each k the sequence a(n) taken modulo k would be periodic with the exact period dividing k. (End) The above conjectures are true. See the Bala link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Oct 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012821 | OEIS-A294362

E.g.f.: exp(Sum_{n>=1} sigma_2(n) * x^n).

More generally, we conjecture that for k = 2,3,4,... the sequence a(n+k) - a(n) is divisible by k: if true, then for each k the sequence a(n) taken modulo k would be periodic with the exact period dividing k. (End) The above conjectures are true. See the Bala link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Oct 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012822 | OEIS-A294363

E.g.f.: exp(Sum_{n>=1} d(n) * x^n), where d(n) is the number of divisors of n.

It appears that the sequence taken modulo 10 is periodic with period 5. More generally, we conjecture that for k = 2,3,4,... the sequence a(n+k) - a(n) is divisible by k: if true, then for each k the sequence a(n) taken modulo k would be periodic with the exact period dividing k. (End) The above conjectures are true. See the Bala link. Conjecture: log(a(n)/n!) ~ sqrt(2*n*log(n)). - _Vaclav Kotesovec_, Sep 07 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Oct 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012823 | OEIS-A294365

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-1) + n, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio. See A293358 for a guide to related sequences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012824 | OEIS-A294366

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-1) + 2n, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio. See A293358 for a guide to related sequences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012825 | OEIS-A294367

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-1) + n - 1, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio. See A293358 for a guide to related sequences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012826 | OEIS-A294406

Positive odd numbers k such that both (sigma(m) - 2*m) and (2*m - sigma(m)) are never equal to k, where sigma(.) is the sum of divisors function A000203 (conjectured).

Positive odd numbers k such that both (sigma(m) - 2*m) and (2*m - sigma(m)) are never equal to k, where sigma(.) is the sum of divisors function A000203 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Oct 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012827 | OEIS-A294413

Solution of the complementary equation a(n) = a(n-1) + a(n-2) - b(n-1) + 6, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 31 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012828 | OEIS-A294414

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-1) + b(n-2), where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 31 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012829 | OEIS-A294415

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-1) + b(n-2) + 1, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 31 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012830 | OEIS-A294416

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-1) + b(n-2) + n, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 31 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012831 | OEIS-A294418

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-1) + 2*b(n-2), where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 31 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012832 | OEIS-A294419

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + 2*b(n-1) + 2*b(n-2), where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 31 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012833 | OEIS-A294420

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + 2*b(n-1) + b(n-2), where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 31 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012834 | OEIS-A294425

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + 2*b(n-1) - b(n-2) - 1, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012835 | OEIS-A294426

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + 2*b(n-1) - b(n-2) - 2, where a(0) = 1, a(1) = 3, b(0) = 2, b(1) = 4.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2, the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012836 | OEIS-A294472

Squarefree numbers whose odd prime factors are all consecutive primes.

Conjecture: if z(n) is the smallest y such that n*k - k^2 is a squarefree number with no gaps in their prime factors >= prime(y) for some k < n, then z(n) >= 1 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Oct 31 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012837 | OEIS-A294475

a(n) = (-1)^n * A294359(n) / (n+1)^2.

It is conjectured that all terms are even after the initial '1,1'.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012838 | OEIS-A294492

Numbers m that set records for the ratio A045763(n)/n.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Nov 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012839 | OEIS-A294512

Denominators of partial sums of the reciprocals of octagonal numbers.

a(n) = A250400(n+1)/(n+1), n >= 0. [conjecture].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Nov 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012840 | OEIS-A294513

Denominators of the partial sums of the reciprocals of twice the pentagonal numbers.

a(n) = 2*A250327(n+1)/(n+1) [conjecture].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Nov 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012841 | OEIS-A294515

Denominators of partial sums of the reciprocals of the decagonal numbers.

a(n) = A250550(n+1)/(n+1) [conjecture].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Nov 02 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012842 | OEIS-A294527

Number of Dyck paths of length 2n with exactly 2 hills.

Conjecture: 2*(3*n-5) *(n-2) *(3*n+11) *(n+1) *a(n) -(3*n+11) *(n-3) *(21*n^2-35*n+10) *a(n-1) -2*(3*n+11) *(n-1) *(2*n-1) *(3*n-2) *a(n-2)= 0. - _R. J. Mathar_, Jun 24 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric M. Schmidt_, Nov 01 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012843 | OEIS-A294532

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-2), where a(0) = 1, a(1) = 2, b(0) = 3.

Conjecture: for every sequence listed here, a(n)/a(n-1) -> (1 + sqrt(5))/2 = golden ratio (A001622).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012844 | OEIS-A294533

Solution of the complementary equation a(n) = a(n-1) + a(n-2) + b(n-2) + 1, where a(0) = 1, a(1) = 2, b(0) = 3.

The increasing complementary sequences a() and b() are uniquely determined by the titular equation and initial values. See A294532 for a guide to related sequences.

Conjecture: a(n)/a(n-1) -> (1 + sqrt(5))/2 = golden ratio (A001622)..

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012845 | OEIS-A294646

a(n) = (1/2)^(2*n) mod (2*n+1).

Conjecture: a(n) is never 2 or 2*n or 2*n-2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_ and _Thomas Ordowski_, Nov 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012846 | OEIS-A294647

Differential variant of the Kolakoski sequence, with initial terms 1, 1.

Conjecture: the density of 1's is equal to 1/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Nov 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012847 | OEIS-A294652

Positive integers k such that the sum of decimal digits of (4^k - 1) equals 3*k.

Conjecture: For k > 446, digsum(4^k - 1) < 3*k. This would mean that no further terms exist in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Iain Fox_, Nov 27 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012848 | OEIS-A294730

Smallest average >= 6 of a twin prime pair that has exactly 2*n divisors, 0 if no such pair exists.

It is conjectured that a(n)=0 for prime n with the only exceptions given by n=A181490(k)+1, i.e. a(2)=6, a(3)=12, a(7)=192 and a(19)=786432 are the only currently known exceptions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Nov 08 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012849 | OEIS-A294745

Lexicographically first sequence of distinct positive integers in which successive terms differ by distinct triangular numbers.

Conjecture: this sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Dec 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012850 | OEIS-A294925

a(n) is the smallest number k with n prime factors such that p + k/p is prime for every prime p | k.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Nov 11 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012851 | OEIS-A294973

Denominators of the continued fraction convergents to sqrt(7)/2.

The g.f. is correct: denominator recurrence a(n) = b(n)*a(n-1) + a(n-2), a(-1) = 0, a(0) = 1, (a(-2) = a(0) = 1) with b(n) modulo 4 from the continued fraction given above: b(0) = 1, b(4*(k+1)) = 2, b(4*k+1) = 3 = b(4*k+3) and b(4*k+2) = 10, for k >= 0. The 4-section is G(x) = Sum_{k>=0} a(k)*x^k = G_0(x^4) + x*G_1(x^4) + x^2*G_2(x^4) + x^3*G_3(x^4) with G_j(x) = Sum_{k>=0} a(4*k+j)*x^k, for j=0..3. The recurrence leads to four equations (omit the x here): G_1 = 3*G_0 + x*G_3, G_2 = 10*G_1 + G_4, G_3 = 3*G_2 + G_1, G_0 = 2*x*G_3 +1 + x*G_2 (using a(-2) = 1). This can be solved to obtain for G(x) = (1 + 3*x + 31*x^2 + 96*x^3 - 31*x^4 + 3*x^5 - x^6)/(1 - 254*x^4 + x^8), and the numerator and denominator factorize like given in the above conjecture. - _Wolfdieter Lang_, Nov 19 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Nov 18 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012852 | OEIS-A294994

Begin with 2; thereafter a(n) is the least prime not already in the sequence such that the Hamming distance between it and the preceding prime is at most 2.

Conjecture: this sequence is a permutation of the primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Nov 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012853 | OEIS-A295070

Solution of the complementary equation a(n) = a(n-2) + b(n-1) + b(n-2), where a(0) = 1, a(1) = 2, b(0) = 3, and (a(n)) and (b(n)) are increasing complementary sequences.

The increasing complementary sequences a() and b() are uniquely determined by the titular equation and initial values. See A295053 for a guide to related sequences. Conjecture: a(n)/a(n-1) -> 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012854 | OEIS-A295078

Numbers n > 1 such that n and sigma(n) have the same smallest and simultaneously largest prime factors.

Conjecture: A007691 (multiply-perfect numbers) is a subsequence. Note that an odd perfect number (if it exists) would be a counterexample to the conjecture. - _Robert Israel_, Jan 08 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Nov 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012855 | OEIS-A295089

a(n) = 3*n^2 + n + 3.

a(n) = A131649(n+3) + 1, n >= 2 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ron Knott_, Nov 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012856 | OEIS-A295112

a(n) = Sum_{k=0..n} binomial(n,2*k)*binomial(2*k,k)/(2*k-1).

Conjecture: The sequence a(n+1)/a(n) (n = 5,6,...) is strictly increasing with limit 3, and the sequence a(n+1)^(1/(n+1))/a(n)^(1/n) (n = 9,10,...) is strictly decreasing to the limit 1. See also A295113 for a conjecture involving the current sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012857 | OEIS-A295113

a(n) = (1/n)*Sum_{k=0..n-1} (8*k + 9)*A295112(k)^2.

Conjecture: a(n) is always integral for every n = 1,2,3,.... Moreover, for any odd prime p we have a(p) == 24 + 10*Leg(-1,p) - 18*Leg(3,p) - 9*Leg(p,3) (mod p^2), where Leg(m,p) denotes the Legendre symbol (m/p).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012858 | OEIS-A295124

a(n) = smallest number k with n prime factors such that 2d + k/d is prime for every d | k.

Conjecture: the sequence is infinite. It is hard to believe!

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Nov 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012859 | OEIS-A295132

a(n) = (2/n)*Sum_{k=1..n} (2k+1)*M(k)^2 where M(k) is the Motzkin number A001006(k).

Sun (2014) conjectures that for any prime p > 3 we have Sum_{k = 0..p-1} M(k)^2 == (2 - 6*p)(p/3) (mod p^2) and Sum_{k = 0..p - 1} k*M(k)^2 == (9*p - 1)(p/3) (mod p^2), where (p/3) is the Legendre symbol.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012860 | OEIS-A295159

Smallest number with exactly n representations as a sum of five nonnegative squares.

Conjecture: a(448) does not exist, i.e., there is no number with exactly 448 such representations. - _Robert Israel_, Nov 15 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012861 | OEIS-A295169

a(n) = smallest number k with n prime factors such that (2d)^2 + (k/d)^2 is prime for every d | k.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Nov 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012862 | OEIS-A295185

a(n) is the smallest composite number whose prime divisors (with multiplicity) sum to prime(n); n >= 3.

Conjecture: For all n >= 3, a(n) >= A288189(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Nov 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012863 | OEIS-A295196

Numbers n > 1 such that 2^(n-1) and (2*n-m)*2^(((n-1)/2) - floor(log_2(n))) are congruent to 1 (mod n) for at least one of m = 3, m = 7 and m = 15.

This definition arises from the conjecture that pseudoprime numbers (A001567) occur only at certain distances m from the next smaller number of the form 2^n. So, if we know that a certain distance does not appear with pseudoprime numbers, we are able to calculate these numbers using Fermat's little theorem and we know that it has to be prime. To "plot" the distance of pseudoprime numbers to 2^n use m = A001567(n) - 2^floor(log_2(A001567(n))). So, the first values of m which do not have a "safe prime number distance" (values with "safe prime number distance" are those values for m which pseudoprime numbers never have) should be m = 1, 49, 81, 85, 129, 133, 273, 275, 289, 321, ... Conjecture 1: There are no composite numbers in this sequence and perhaps infinitely many primes. Conjecture 2: For m = 7 this definition generates A104066 and for m = 15 this definition generates A144487 (A057197). Conjecture 3: There are (infinitely many?) m for which this definition generates nothing but (infinitely many?) primes of the form p = 2^k + m.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonas Kaiser_, Nov 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012864 | OEIS-A295286

Sum of the products of the smaller and larger parts of the partitions of n into two parts with the smaller part odd.

Conjectures from _Colin Barker_, Nov 20 2017: (Start) Conjectures verified by _Robert Israel_, Dec 05 2017.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Nov 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012865 | OEIS-A295292

Sum of the products of the smaller and larger parts of the partitions of n into two parts with the larger part odd.

Conjectures from _Colin Barker_, Nov 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Nov 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012866 | OEIS-A295317

Sum of the products of the smaller and larger parts of the partitions of n into two distinct parts with the smaller part odd.

Conjectures from _Colin Barker_, Nov 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Nov 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012867 | OEIS-A295318

Sum of the products of the smaller and larger parts of the partitions of n into two distinct parts with the smaller part even.

Conjectures from _Colin Barker_, Nov 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Nov 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012868 | OEIS-A295319

a(n) is the sum of all n-digit palindromes.

Conjecture: All terms after a(1) are of the form 495*10^x where x is any nonnegative integer k such that k != 1 (mod 3). - _Harvey P. Dale_, Mar 05 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Melvin Peralta_, Nov 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012869 | OEIS-A295320

Sum of the products of the smaller and larger parts of the partitions of n into two distinct parts with the larger part odd.

Conjectures from _Colin Barker_, Nov 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Nov 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012870 | OEIS-A295321

Sum of the products of the smaller and larger parts of the partitions of n into two distinct parts with the larger part even.

Conjectures from _Colin Barker_, Nov 20 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Nov 19 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012871 | OEIS-A295339

Least k for the inner Theodorus spiral to complete n revolutions.

Conjecture: a(n) = A072895(n) - 2, n >= 1. This follows from the conjecture that the sequences K := {floor(phi(k)/(2*Pi)}_{k >= 1} with phi given above, and Khat:= {floor(phihat(k)/(2*Pi)}_{k >= 1} with phihat given above satisfy Khat(k-2) = K(k), for k >= 3. Note that phihat(k-2) - phi(k) = alpha(k-2) - alpha(k-1) =: delta(k) = arctan((sqrt(k-1) - sqrt(k-2))/(1 + sqrt((k-1)*(k-2)))) > 0, for k >= 3. Therefore the conjecture is that delta(k) < 2*Pi*(1 - frac(phi(k)/(2*Pi))), for k >= 3, or, equivalently, phihat(k-2) < 2*Pi*(K(k) + 1), for k >= 3. Conjecture: a(n) = A072895(n) - 2, n >= 1 (see the comment above).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Dec 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012872 | OEIS-A295405

a(n) = 1 if n^2+1 is prime, 0 otherwise.

It is conjectured that n^2+1 is prime infinitely often.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Plouffe_, Nov 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012873 | OEIS-A295423

a(n) = size of the part containing the diagonal of the symmetric representation of sigma(k) when k is the n-th number with an odd number of parts.

Conjecture: Every even number, except 2, 4, 6, 10, and 14, occurs as the size of the middle part of the symmetric representation with an odd number of parts. Size 6 is impossible if Conjecture (2) in A285356 is true since the middle area requires legs of length 3 in the Dyck path at the diagonal; and so are sizes 10 and 14 unless the middle part touches the previous part at a point.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Feb 12 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012874 | OEIS-A295424

Number of distinct twin primes which are in Goldbach partitions of 2n.

Tomas Oliveira e Silva in 2012 experimentally confirmed that all even numbers 4 <= n <= 4 * 10^18 have at least one Goldbach partition (GP) with a prime 9781 or less.

Detailed examination of all even numbers less than 10^6 showed that the most popular prime in all GPs is 3 (78497 occurrences), then 5 (70328), then 7 (62185), then 11 (48582), then 13 (40916), then 17 (31091), then 19 (29791) -- all these primes are twin primes. These results gave rise to a hypothesis that twin primes should be rather frequent in GP, especially those relatively small. Conjecture. Further empirical examinations lead to a hypothesis that all even numbers n > 4 have at least 1 twin prime in GP(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marcin Barylski_, Feb 12 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012875 | OEIS-A295431

a(n) = (12*n)!*n! / ((6*n)!*(4*n)!*(3*n)!).

Conjectures: 5*7*11*a(n)/(n + 1), 5*a(n)/(2*n + 1), a(n)/(3*n + 1) and 2*5*7*11*a(n)/((n + 1)*(2*n + 1)*(3*n + 1)) are integers for all n (checked up to n = 1000). It appears that a(n)/(12*n - 1) is integral for all n (checked up to n = 1000). More generally, for r >= 1, we conjecture that there exists a constant D(r) > 0 such that D(r)*a(n)/Product_{i = 1..r, i coprime to 12} (12*n - i) is integral for all n. For n >= 0, we conjecture that both a(n/2) = (6*n)!*(n/2)! / ((3*n)!*(2*n)!*(3*n/2)!) = A347854(n) and a(n/3) = (4*n)!*(n/3)! / ((2*n)!*(4*n/3)!*n!) = A347855(n) are integers, where fractional factorials are defined using the Gamma function; for example, (n/2)! := Gamma(n/2 + 1). - _Peter Bala_, Jun 09 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Nov 22 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012876 | OEIS-A295437

a(n) = (18*n)!*n!/((9*n)!*(6*n)!*(4*n)!).

Conjectures: for n >= 0, both a(n/2) = (9*n)!*(n/2)! / ((9*n/2)!*(3*n)!*(2*n)!) and a(n/3) = (6*n)!*(n/3)! / ((3*n)!*(2*n)!*(4*n/3)!) are integers, where fractional factorials are defined using the Gamma function; for example (n/2)! := Gamma(n/2 + 1). - _Peter Bala_, Jun 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Nov 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012877 | OEIS-A295477

a(n) = (24*n)!*n!/((12*n)!*(8*n)!*(5*n)!).

Conjectures: for n >= 0, both a(n/2) = (12*n)!*(n/2)! / ((6*n)!*(4*n)!*(5*n/2)!) and a(n/4) = (6*n)!*(n/4)! / ((3*n)!*(2*n)!*(5*n/4)!) are integers, where fractional factorials are defined using the Gamma function - for example, (n/2)! := Gamma(n/2 + 1). - _Peter Bala_, Jul 31 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, Nov 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012878 | OEIS-A295622

Number of nonequivalent dissections of an n-gon into 3 polygons by nonintersecting diagonals rooted at a cell up to rotation.

Conjectures from _Colin Barker_, Nov 25 2017: (Start) a(n) = Sum_{k=0..n-5} f(k), where f(n) = Sum_{k=0..n} (3 + lcm(k, 2)) (conjecture). - _Jon Maiga_, Nov 28 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Howroyd_, Nov 24 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012879 | OEIS-A295687

a(n) = a(n-1) + a(n-3) + a(n-4), where a(0) = 1, a(1) = 2, a(2) = 2, a(3) = 1.

Conjecture: a(2*n+7) = Fibonacci(n)^3*Sum_{k >= 1} k^2 * Fibonacci(n*k)/ Fibonacci(n+2)^(k+1), n >= 1. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012880 | OEIS-A295738

a(1)=2, a(2)=3; for n > 2, a(n) is the smallest prime number greater than a(n-1) which has the form a(n-2)*2^k - a(n-1) for some integer k.

Conjecture 1: Conjecture 2:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Nov 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012881 | OEIS-A295741

a(n) is smallest divisor d of the n-th primorial such that d + prime(n)#/d is prime.

Conjecture: a(n) < prime(n)^2 for n > 0. Reo Fortune conjectured that the smallest integer q such that primorial(n) + q is prime, is again a prime: see A005235. This conjecture is still unproven. For this sequence here as well, it is unproven that a(n) always exists. Of course it is trivial that a(n)^2 < prime(n)#, and a(n) must be a squarefree product of primes less than prime(n), if it exists. - _M. F. Hasler_, Apr 11 2026 Conjecture: a(n) < prime(n) for n > 0. - _Alain Rocchelli_, Aug 31 2026 Conjecture: Limit_{N->oo} (Sum_{n=1..N} a(n)) / (Sum_{n=1..N} prime(n)) = 0. - _Alain Rocchelli_, Aug 30 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_ and _Robert G. Wilson v_, Nov 26 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012882 | OEIS-A295762

G.f. A(x) satisfies: A(x - 2*A(x^2)) = x + A(x^2).

Odd terms seem to occur only at a(2^n) for n>=0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012883 | OEIS-A295766

G.f. A(x) satisfies: [x^(n-1)] A(x)^(n^2)/n^2 = [x^(n-2)] A(x)^(n^2) for n>=2 with A'(0) = 1.

a(A075427(k) - 1) is odd for n>=0 and a(n) is even elsewhere (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 31 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012884 | OEIS-A295811

G.f. A(x) satisfies: [x^(n-1)] A(x)^(n^2) = 2*n * [x^(n-2)] A(x)^(n^2) for n>=2, with A(0) = 1.

a(2^k - 1) is odd for k>=0 and a(n) is even elsewhere (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 02 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012885 | OEIS-A295838

Largest value corresponding to a string of n printable ASCII characters.

Conjecture: For every a(n) there exists a sequence of primes (p(1), p(2), p(3), ...) such that for each term a(n) there exists a set of primes that when added to the term result in another prime. For example, a(2)=126 and 126 + {5,11,13,23,37,41,47,...} all are primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Abilene Sukin_, Nov 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012886 | OEIS-A295867

Numbers of the form A000217(n)*A007494(n) that are divisible by 3.

Conjectures from _Colin Barker_, Feb 15 2018: (Start) Colin Barker's conjecture is true. This is a cubic quasipolynomial of order 5: a(n) = 162/125*n^3 + 27/25*n^2 if n is 0 mod 5, 162/125*n^3 + 261/125*n^2 + 111/125*n + 12/125 if n is 4 mod 5, 162/125*n^3 + 297/125*n^2 + 144/125*n + 21/125 if n is 3 mod 5, 162/125*n^3 + 423/125*n^2 + 339/125*n + 84/125 if n is 2 mod 5, and 162/125*n^3 + 459/125*n^2 + 396/125*n + 108/125 if n is 1 mod 5. Generally a(n) = 162/125*n^3 + O(n^2). - _Charles R Greathouse IV_, Feb 20 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Justin Gaetano_, Feb 13 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012887 | OEIS-A295928

Number of triangular matrices T(n,i,k), k <= i <= n, with entries "0" or "1" with the property that each triple {T(n,i,k), T(n,i,k+1), T(n,i-1,k)} containing a single "0" can be successively replaced by {1, 1, 1} until finally no "0" entry remains.

The algorithm for the sequence is rather slow because each start matrix is constructed separately. There exists a faster recursive algorithm which produces the same terms and therefore is likely to be correct, but it is based on a conjecture. For the theory of the recurrence, see "Recursive aspects of fill matrices". Probable extension a(10)-a(14):

821096828, 15804092592, 324709899276, 7081361097108, 163179784397820.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerhard Kirchner_, Nov 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012888 | OEIS-A296056

Determinant of the inverse of the matrix A_n, where A_n is the n X n matrix defined by A_n[i,j] = 1/C(i+j-2) for 1 <= i,j <= n, and C(k) is the k-th Catalan number (A000108).

It is conjectured that a(n) is an integer for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tom Richardson_, Dec 03 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012889 | OEIS-A296075

Sum of deficiencies of divisors of n.

No, Adamczewski gives a counterexample (a(33085221781504) = 1) to the above conjecture in linked paper. - _Vaclav Kotesovec_, Aug 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 04 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012890 | OEIS-A296141

Sum of the smaller parts of the partitions of n into two distinct parts with the larger part even.

Conjectures from _Colin Barker_, Dec 06 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Dec 05 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012891 | OEIS-A296159

Sum of the smaller parts in the partitions of n into two distinct parts with the larger part odd.

Conjectures from _Colin Barker_, Dec 06 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Dec 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012892 | OEIS-A296161

Sum of the larger parts of the partitions of n into two parts such that the smaller part is odd.

Conjectures from _Colin Barker_, Dec 06 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Dec 06 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012893 | OEIS-A296179

Number of points of the inner discrete Theodorus spiral on sheet S_n, n >= 1. First differences of A295339.

The conjecture stated in A295339 implies that a(n) = A295338(n), for n >= 2. Conjecture: a(n) = A295338(n), for n >= 2 (see a comment above).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Dec 13 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012894 | OEIS-A296181

First point of the discrete Theodorus spiral in the fourth quadrant for the n-th revolution, for n >= 1.

This sequence is used in a conjecture on points z_k of the discrete (outer) Theodorus spiral living on quadrant IV of the complex plane of sheet S_n, where S_n := {r*exp(i*phi), r > 0, 2*Pi*(n-1) <= phi < 2*Pi*n}. This corresponds to the n-th revolution, for n >= 1. This conjecture is 2*Pi - varphi(A072895(n)) > arctan(a(n)), n >= 1, with varphi(k) = phi(k) - 2*Pi*floor(phi(k)/(2*Pi)) where z_k = sqrt(k)*exp(i*phi(k)). This conjecture implies a conjecture relating points of the discrete inner spiral to those of the outer ones, namely Khat(k-2) := floor(phihat(k-2)/(2*Pi)) = K(k) =: floor(phi(k)/(2*Pi)) for k >= 3, where zhat_k = sqrt(k)*exp(i*phihat(k)) is a point of the discrete inner Theodorus spiral, given in terms of z_k by zhat(k) = ((k-1 + 2*sqrt(k)*i)/(k+1))*z_k. This implies phihat(k) = phi(k) + arctan((sqrt(k-1) - sqrt(k-2))/(1 + sqrt((k-1)*(k-2)))). The implied conjecture Khat(k-2) = K(k), k >= 3, for the other three quadrants of each sheet S_n can be proved. For the inner spiral see the Waldvogel link. If the implied conjecture is true then A295339(n) = A072895(n) - 2, for n >= 1, hence A296179(n) = A295338(n), for n >= 2. For the conjecture and the proof for the first three quadrants for each sheet S_n see the W. Lang link. - _Wolfdieter Lang_, Jan 24 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jan 05 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012895 | OEIS-A296299

Dimension of the n-th component of a certain graded Lie algebra.

Conjecture: Let b = 122112112221... be the sequence (a(n+1)), written as a word. Let mu be the morphism given by mu(1)=121, mu(2)=12221. Then b = 1221 mu(b). - _Michel Dekking_, Sep 29 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric M. Schmidt_, Dec 09 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012896 | OEIS-A296343

Numbers in cells in spiral A274820 on spoke through center with slope 0 degrees.

It is conjectured that this and A296346 together are a permutation of the nonnegative numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012897 | OEIS-A296344

Numbers in cells in spiral A274820 on spoke through center with slope 60 degrees.

It is conjectured that this and A296347 together are a permutation of the nonnegative numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012898 | OEIS-A296345

Numbers in cells in spiral A274820 on spoke through center with slope 120 degrees.

It is conjectured that this and A296348 together are a permutation of the nonnegative numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012899 | OEIS-A296346

Numbers in cells in spiral A274820 on spoke through center with slope 180 degrees.

It is conjectured that this and A296343 together are a permutation of the nonnegative numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012900 | OEIS-A296347

Numbers in cells in spiral A274820 on spoke through center with slope 240 degrees.

It is conjectured that this and A296344 together are a permutation of the nonnegative numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012901 | OEIS-A296348

Numbers in cells in spiral A274820 on spoke through center with slope 300 degrees.

It is conjectured that this and A296345 together are a permutation of the nonnegative numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 16 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012902 | OEIS-A296362

Number of monohedral disk tilings of type C^t_{5,n}.

G.f.: 2*x*(1 + 763*x + 905*x^2 + 1871*x^3 + 2142*x^4 + 2318*x^5 + 2333*x^6 + 1022*x^7 + 602*x^8 - 348*x^9 - 1422*x^10 - 1599*x^11 - 2949*x^12 - 3041*x^13 - 2413*x^14 - 2329*x^15 - 316*x^16 - 538*x^17 + 175*x^18 + 703*x^19 + 562*x^20 + 1446*x^21 + 852*x^22 + 147*x^23 + 48*x^24 - 646*x^25 - 6*x^26 + 224*x^27 + 16*x^28 + 184*x^29 - 310*x^30 + 107*x^31) / ((1 - x)^9*(1 + x)^2*(1 + x^2)*(1 + x + x^2)^3*(1 + x^3 + x^6)*(1 + x + x^2 + x^3 + x^4 + x^5 + x^6)) (conjectured). - _Colin Barker_, Jan 09 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 15 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012903 | OEIS-A296440

a(1) = 2, a(2) = a(3) = 3, a(4) = 4, a(5) = a(6) = 6; a(n) = a(n-a(n-1)) + a(n-a(n-2)) + a(n-a(n-3)) for n > 6.

Conjecture: Sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Dec 12 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012904 | OEIS-A296508

Irregular triangle read by rows: T(n,k) is the size of the subpart that is adjacent to the k-th peak of the largest Dyck path of the symmetric representation of sigma(n), or T(n,k) = 0 if the mentioned subpart is already associated to a previous peak or if there is no subpart adjacent to the k-th peak, with n >= 1, k >= 1.

Conjecture: row n is formed by the odd-indexed terms of the n-th row of triangle A280850 together with the even-indexed terms of the same row but listed in reverse order. Examples: the 15th row of A280850 is [8, 8, 7, 0, 1] so the 15th row of this triangle is [8, 7, 1, 0, 8]. The 75th row of A280850 is [38, 38, 21, 0, 3, 3, 0, 0, 0, 21, 0] so the 75h row of this triangle is [38, 21, 3, 0, 0, 0, 21, 0, 3, 0, 38].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Feb 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012905 | OEIS-A296516

a(n) is the number of terms in expanded form of bivariate polynomial Q_n, where (P_n, Q_n) is defined by: P_0 = x, Q_0 = y, P_(n+1) = P_n + Q_n, Q_(n+1) = P_n * Q_n.

Programs based on the direct application of the definition quickly reach a limitation by combinatorial explosion, hence this short list of values in section Data. The first conjectured formula (see Formulas) obtained by the observation of a pattern in the 2D shape of Q_n (as drawn in the Examples) is more computationally efficient and makes it possible to produce a significantly longer list of (nonguaranteed) values: see attached a-file in b-file format, section Links. Conjectured: Conjectures from _Colin Barker_, Mar 28 2018: (Start) Derived from the conjectured order-7 linear recurrence above, for n > 0,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luc Rousseau_, Feb 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012906 | OEIS-A296523

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x > 0, y >= 0 and 0 <= z <= w such that 64*x^2 + 65*y^2 = q^2 for some practical number q.

Conjecture: (i) a(n) > 0 for all n > 0. Also, a(n) = 1 only for those numbers 2^k (k = 0,1,2,...), 4^k*m (k = 0,1,...; m = 3, 7), 4^k*47 (k = 0,1,2,3), 4^k*s (k = 0,1,2; s = 15, 23, 31, 39, 71); 4^k*t (k =0,1; t = 87, 111, 119, 159, 191, 311).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012907 | OEIS-A296524

Number of connected (2*k)-regular graphs on 2*n+1 nodes with maximal diameter D(n,k) A294733 written as triangular array T(n,k), 1 <= k <= n.

The next term a(24) corresponding to the 6-regular graphs on 15 nodes is conjectured to be 1. It seems that there exists only one graph with diameter A294733(24)=4. Its adjacency matrix is The conjecture is confirmed using Markus Meringer's GenReg program. Aside from the 1 shown 6-regular graph on 15 nodes with diameter 4 there are 870618932 graphs with diameter 2 and 599674742 graphs with diameter 3. - _Hugo Pfoertner_, Dec 19 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 14 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012908 | OEIS-A296604

Number of Johnson solids with n faces.

Sum_{n>0} a(n) = 92, the number of Johnson solids, as conjectured by Johnson and proved by Zalgaller.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Jan 28 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012909 | OEIS-A296615

Lexicographically earliest sequence of distinct positive terms such that, for any n > 0, a(n) XOR a(n+1) is a cube (where XOR denotes the XOR binary operator).

This sequence is conjectured to the a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Dec 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012910 | OEIS-A296618

Expansion of the e.g.f. exp(-x)/sqrt(1-4*x).

Conjectures: Conjecture 1) above follows immediately from the stated recurrence equation. In fact, a(n+1) == a(n) (mod 8*n) for n >= 1. For a proof of Conjecture 2) see the Bala link (corollary to Theorem 1, p. 5). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emanuele Munarini_, Dec 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012911 | OEIS-A296806

Take a prime, convert it to base 2, remove its most significant digit and its least significant digit and convert it back to base 10. Sequence lists primes that generate another prime by this process.

Conjecture: for any positive integer p, there are infinitely many primes q for which there exists an integer n such that q-(2p+1) = 2^n. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_, _Paolo Iachia_, Dec 21 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012912 | OEIS-A296998

Number of ways to place 4 points on an n X n point grid so that no point is equally distant from two other points on the same row or the same column.

Conjectures from _Colin Barker_, Dec 23 2017: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Heinrich Ludwig_, Dec 23 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012913 | OEIS-A296999

Number of nonequivalent (mod D_8) ways to place 4 points on an n X n point grid so that no point is equally distant from two other points on the same row or the same column.

Conjectures from _Colin Barker_, Jan 21 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Heinrich Ludwig_, Jan 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012914 | OEIS-A297105

Smallest square k > 1 such that iterating the map (k -> number of divisors(k)) n times yields a square on each iteration.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Dec 25 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012915 | OEIS-A297180

a(n) is the smallest positive integer of length (distance from origin) n in the Cayley graph of the integers generated by all powers of 7.

Conjectures from _Colin Barker_, Dec 28 2017: (Start) The second conjecture by Colin Barker is true up to n=1000. - _Lars Blomberg_, Dec 29 2017

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 28 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012916 | OEIS-A297195

Number of bitriangular permutations (row sums of A272644 if that triangle is prefixed with two rows for n=0,1).

Define c(n) = Sum_{m=2..n-1} C(n-1, m-1)^(-2). Define b(1) = x and b(n+1) = b(n) + (Sum_{m=2..n-1} b(m)*b(n+1-m)*C(n-1, m-1)^(-2))/n^2 for n>0. Then b(n) is a polynomial in x and so is (b(n+1)-b(n))/x^2 whose constant term is c(n)/n^2. The Hone et al. (2002) link denotes x with alpha_2 and alpha_k = (k-1)!^2*b(k). Conjecture: Asymptotic expansion of c(n) = 2*Sum_{i>1} a(i)/n^i. - _Michael Somos_, Oct 17 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012917 | OEIS-A297291

Solution (a(n)) of the system of 3 complementary equations in Comments.

Conjectures: for n >=0,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 24 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012918 | OEIS-A297292

Solution (b(n)) of the system of 3 complementary equations in Comments.

Conjectures: for n >= 0,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 24 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012919 | OEIS-A297293

Solution (c(n)) of the system of 3 complementary equations in Comments.

Conjectures: for n >= 0,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 24 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012920 | OEIS-A297402

a(n) = gcd_{k=1..n} (prime(k+1)^n-1)/2.

Conjecture: a(n) equals the GCD of the inradius terms (p^n-1)/2 for all odd primes, i.e. a(n) = GCD_{k=1..oo} (prime(k+1)^n-1)/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Dec 29 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012921 | OEIS-A297440

Rectangular array R by antidiagonals: row n shows the positive integers whose base-3 digits have total variation n, for n>=0. See Comments.

Conjecture: each column, after some number of initial terms, satisfies the linear recurrence relation c(n) = c(n-1) + 2*c(n-2) - 2*c(n-3) + 3*c(n-4) - 3*c(n-5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012922 | OEIS-A297441

Rectangular array R by antidiagonals: row n shows the positive integers whose base-3 digits have down-variation n, for n>=0. See Comments.

Conjecture: each column, after some number of initial terms, satisfies the linear recurrence relation c(n) = c(n-1) + 9*c(n-2) - 9*c(n-3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012923 | OEIS-A297443

a(n) = a(n-1) + 2*a(n-2) - 2*a(n-3) + 3*a(n-4) - 3*a(n-5), where a(0) = 1, a(1) = 3, a(2) = 6, a(3) = 11, a(4) = 20, a(5) = 33.

Conjecture: a(n) = least positive whose base-3 total variation is n; see A297440.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012924 | OEIS-A297444

a(n) = a(n-1) + 9*a(n-2) - 9*a(n-3), where a(0) = 1, a(1) = 3, a(2) = 6, a(3) = 33.

Conjecture: a(n) = least positive whose base-3 down-variation is n; see A297441.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012925 | OEIS-A297445

a(n) = a(n-1) + 9*a(n-2) - 9*a(n-3), where a(0) = 1, a(1) = 5, a(2) = 11.

Conjecture: a(n) = least positive whose base-3 up-variation is n; see A297441.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012926 | OEIS-A297464

Solution (a(n)) of the system of 4 complementary equations in Comments.

Conjecture: for all n >= 0, a(n) = a(n-1) + a(n-3) - a(n-4) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012927 | OEIS-A297465

Solution (b(n)) of the system of 4 complementary equations in Comments.

Conjecture: for all n >= 0,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012928 | OEIS-A297466

Solution (c(n)) of the system of 4 complementary equations in Comments.

Conjecture: for all n >= 0,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012929 | OEIS-A297469

Solution (bb(n)) of the system of 3 complementary equations in Comments.

Conjecture: For t >= 0, bb(2t) = 10t + 1 (+1 if binary expansion of t ends in an odd number of 0's), bb(2t+1) = 10t + 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012930 | OEIS-A297475

Numbers k such that phi(x) = k for more than one value of x, and the smallest such x divides the largest.

Conjecture 1: The number of solutions, excluding endpoints is always 0, or an odd number. (Known to k = 2 * 10^5) Conjecture 2: If both endpoints are divisible by 5, then the number of solutions (excluding terms of A007366) is of the form 4m + 1. (Known to k = 2 * 10^5)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Torlach Rush_, Dec 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012931 | OEIS-A297477

Triangle read by rows: T(n, k) gives the coefficients of x^k of the characteristic polynomial P(n, x) of the n X n matrix M with entries M(i, j) = 1 if i = 1 or j = 1, -1 if i = j > 1, and 0 otherwise. T(0, 0) := 0.

Therefore the spectral radius (absolute value of the maximal eigenvalue) is rho_n = sqrt(n), and the spectral norm of M_n (square root of the maximal eigenvalue of (M_n)^+ M_n) is also sqrt(n), for n >= 1. See the conjecture in the first comment above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Dec 30 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012932 | OEIS-A297551

Rectangular array R by antidiagonals: row n shows the positive integers whose base-4 digits have total variation n, for n>=0. See Comments.

Conjecture: each column, after some number of initial terms, satisfies a homogeneous linear recurrence relation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012933 | OEIS-A297554

a(n) = a(n-2) + 4*a(n-3) - 4*a(n-5), where a(0) = 1, a(1) = 4, a(2) = 7, a(3) = 12, a(4) = 19, a(5) = 28.

Conjecture: a(n) = least positive whose base-4 total variation is n; see A297551.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012934 | OEIS-A297555

a(n) = a(n-1) + 16*a(n-3) - 16*a(n-4), where a(0) = 1, a(1) = 4, a(2) = 8, a(3) = 12, a(4) = 76.

Conjecture: a(n) = least positive whose base-4 down-variation is n; see column 1 of A297552.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012935 | OEIS-A297556

a(n) = a(n-1) + 16*a(n-3) - 16*a(n-4), where a(0) = 1, a(1) = 4, a(2) = 7, a(3) = 19.

Conjecture: a(n) = least positive whose base-4 up-variation is n; see column 1 of A297553.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012936 | OEIS-A297557

Number of maximum independent vertex sets in the n-triangular grid graph.

Conjecture: a(n) = a(n-3) for n >= 12. - _Andrew Howroyd_, Jan 01 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Dec 31 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012937 | OEIS-A297573

Least positive integer m such that m*n divides F(m+n), where F(k) denotes the k-th Fibonacci number A000045(k).

Conjecture: a(n) exists for any n > 0. See also A297574 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012938 | OEIS-A297575

Numbers whose sum of divisors is divisible by 10.

Of the first 10^2, 10^6, and 10^10 positive integers, 17%, 40.8%, and 48.4%, respectively, are in the sequence (See Robert G. Wilson's table below). So the conjecture here is that if this trend continues then as the numbers approach infinity, the percentage of those numbers whose sum of divisors is divisible by 10 approaches 100%.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zoltan Galantai_, Jan 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012939 | OEIS-A297705

a(n) = Sum_{k=0..n} binomial(n, k)*hypergeom([k - n, n + 1], [k + 2], -4).

(End) [Conjectures verified with Maple's FormalPowerSeries Module. - _Peter Luschny_, Nov 10 2022]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jan 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012940 | OEIS-A297741

The number of distinct positions on an infinite chessboard reachable by the (3,4)-leaper in <= n moves.

Conjecture: a(n) = 62*n^2 + 30*n - 55 for n >= 10. Conjectures from _Colin Barker_, Jan 06 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jan 05 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012941 | OEIS-A297789

The number of length 2n - 1 strings over the alphabet {0, 1} such that the first half of any initial odd length substring is a permutation of the second half.

Conjecture: lim_{n->infinity} a(n+1)/a(n) exists and is a value in [1, 2]. [The following comment suggests that on the contrary, this limit may not exist. - _N. J. A. Sloane_, Jan 30 2018, following a comment from _Peter Kagey_, Jan 29 2017] Conjecture: a(n) is odd for all n > 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Jan 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012942 | OEIS-A297826

Solution (a(n)) of the near-complementary equation a(n) = a(1)*b(n-1) - a(0)*b(n-2) + n, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

The sequence (a(n)) generated by the equation a(n) = a(1)*b(n-1) - a(0)*b(n-2) + n, with initial values as shown, includes duplicates; e.g. a(18) = a(19) = 51. If the duplicates are removed from (a(n)), the resulting sequence and (b(n)) are complementary. Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012943 | OEIS-A297827

Difference sequence of A297826.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012944 | OEIS-A297828

Difference sequence of A297997.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012945 | OEIS-A297830

Solution of the complementary equation a(n) = a(1)*b(n-1) - a(0)*b(n-2) + 2*n, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

The increasing complementary sequences a() and b() are uniquely determined by the titular equation and initial values. Conjecture: a(n) - (2 +sqrt(2))*n < 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012946 | OEIS-A297831

Solution of the complementary equation a(n) = a(1)*b(n-1) - a(0)*b(n-2) + 2*n - 1, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjecture: a(n) - (2 +sqrt(2))*n < 5/2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012947 | OEIS-A297833

Solution of the complementary equation a(n) = a(1)*b(n-1) - a(0)*b(n-2) + 2*n - 3, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjecture: -2 < a(n) - (2 +sqrt(2))*n <= 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012948 | OEIS-A297834

Solution of the complementary equation a(n) = a(1)*b(n-1) - a(0)*b(n-2) + 2*n - 4, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjecture: -3 < a(n) - (2 +sqrt(2))*n <= 1 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012949 | OEIS-A297836

Solution of the complementary equation a(n) = a(1)*b(n-1) - a(0)*b(n-2) + 3*n, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjectures: a(n) - (5 + sqrt(13))*n/2 < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012950 | OEIS-A297837

Solution of the complementary equation a(n) = a(1)*b(n-1) - a(0)*b(n-2) + 4*n, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjecture: a(n) - (3 + sqrt(5))*n < 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012951 | OEIS-A297844

a(n) = Sum_{d|n} max(d, n/d)^5.

If p is a prime, then 2*p^5 belongs to this sequence. Conjecture: The converse is true. - _Alexandra Hercilia Pereira Silva_, Oct 04 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Jan 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012952 | OEIS-A297850

Least common prime factor of the members of n-th amicable pair, or 0 if the two members of the pair are coprime.

The question whether a(n) = 0 for any n is an open problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Jan 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012953 | OEIS-A297964

a(n) = A296220(n) - 4n.

It has been checked that -9 <= a(n) <= 8 for n=1..6*10^6. Conjectures: every integer appears infinitely many times, and -2 < a(n) - a(n-1) < 2 for n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012954 | OEIS-A297995

Number of ways to write n as 4*u + v^2 + x^3 + y^4 + 2*z^8, where u is 0 or 1, v is a positive integer and x,y,z are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 398, 496, 498, 500, 1507, 6419, 7843, 7983, 8688, 8947, 9175, 9251, 12923, 12976, 48381.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012955 | OEIS-A297999

Solution (a(n)) of the near-complementary equation a(n) = a(1)*b(n) - a(0)*b(n-1) + n, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, , b(2) = 5, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

The sequence (a(n)) generated by the equation a(n) = a(1)*b(n) - a(0)*b(n-1) + n, with initial values as shown, includes duplicates; e.g. a(18) = a(19) = 52. If the duplicates are removed from (a(n)), the resulting sequence and (b(n)) are complementary. Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012956 | OEIS-A298000

Solution of the complementary equation a(n) = a(1)*b(n) - a(0)*b(n-1) + 2*n, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, b(2) = 5, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjectures: a(n) - (2 +sqrt(2))*n < 4 for n >= 1. Guide to related sequences having initial values a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, b(2) = 5, where (b(n)) is the increasing sequence of positive integers not in (a(n)):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012957 | OEIS-A298001

Solution of the complementary equation a(n) = a(1)*b(n) - a(0)*b(n-1) + 3*n, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, b(2) = 5, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjecture: a(n) - n*L < 4 for n >= 1, where L = (5 + sqrt(13))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012958 | OEIS-A298002

Solution of the complementary equation a(n) = a(1)*b(n) - a(0)*b(n-1) + 4*n, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, b(2) = 5, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjecture: a(n) - n*L < 4 for n >= 1, where L = 3 + sqrt(5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012959 | OEIS-A298003

Solution b( ) of the complementary equation a(n) = a(1)*b(n-1) - a(0)*b(n-2) + 2*n, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjecture: 3/5 < a(n) - n*sqrt(2)*n < 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012960 | OEIS-A298004

Solution b( ) of the complementary equation a(n) = a(1)*b(n-1) - a(0)*b(n-2) + 3*n, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjecture: 7/10 < a(n) - n*L < 3 for n >= 1, where L = (-1 + sqrt(13))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012961 | OEIS-A298005

Solution b( ) of the complementary equation a(n) = a(1)*b(n-1) - a(0)*b(n-2) + 4*n, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjecture: 7/10 < a(n) - n*L < 3 for n >= 1, where L = -1 + sqrt(5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012962 | OEIS-A298006

Solution b( ) of the complementary equation a(n) = a(1)*b(n-1) - a(0)*b(n-2) + 2*n - 1, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjecture: 9/10 < a(n) - n*sqrt(2) < 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012963 | OEIS-A298007

Solution b( ) of the complementary equation a(n) = a(1)*b(n-1) - a(0)*b(n-2) + 2*n - 2, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjecture: 1 < a(n) - n*sqrt(2) < 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012964 | OEIS-A298022

Coordination sequence for Dual(3^3.4^2) tiling with respect to a trivalent node.

Conjectures from _Colin Barker_, Jan 22 2018: (Start) Conjecture: a(n) = 2*(21*n + 3*A049347(n+2)/2)/9 for n > 4. - _Stefano Spezia_, Nov 24 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012965 | OEIS-A298023

Partial sums of A298022.

Conjectures from _Colin Barker_, Jan 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012966 | OEIS-A298024

Expansion of (x^4+3*x^3+6*x^2+3*x+1)/((1-x)*(1-x^3)).

Coordination sequence for Dual(3^3.4^2) tiling with respect to a tetravalent node. This tiling is also called the prismatic pentagonal tiling, or the cem-d net. It is one of the 11 Laves tilings. (The identification of this coordination sequence with the g.f. in the definition was first conjectured by _Colin Barker_, Jan 22 2018.) a(n) = a(n-1) + a(n-3) - a(n-4) for n>4. (Conjectured, correctly, by _Colin Barker_, Jan 22 2018.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012967 | OEIS-A298036

Coordination sequence of Dual(4.6.12) tiling with respect to a 12-valent node.

Conjecture: For n>0, a(n)=6n if n even, otherwise 12n. The conjecture can easily be shown to be true: The vertices at distance 2k consist of 3k 12-valent and 3k 4-alent vertices, and the vertices at distance 2k+1 consist of 6(k+1) 6-valent and 6(k+1) 4-valent vertices. - _Charlie Neder_, Apr 22 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012968 | OEIS-A298038

Coordination sequence of Dual(4.6.12) tiling with respect to a hexavalent node.

Conjecture: For n > 0, a(n)=12n if n even, otherwise 6n. Conjectures from _Colin Barker_, Apr 03 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012969 | OEIS-A298040

Coordination sequence of Dual(4.6.12) tiling with respect to a tetravalent node.

Conjecture: For n>0, a(n)=10n if n even, otherwise 8n. Conjectures from _Colin Barker_, Apr 01 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012970 | OEIS-A298108

Solution b( ) of the complementary equation a(n) = a(1)*b(n-1) - a(0)*b(n-2) + 2*n - 3, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjecture: 3/2 < a(n) - n*sqrt(2) < 4 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012971 | OEIS-A298109

Solution b( ) of the complementary equation a(n) = a(1)*b(n-1) - a(0)*b(n-2) + 2*n - 4, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjecture: 3/2 < a(n) - n*sqrt(2) < 4 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012972 | OEIS-A298111

Solution b( ) of the complementary equation a(n) = a(1)*b(n) - a(0)*b(n-1) + 2*n, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, b(2) = 5, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjecture: 1/5 < a(n) - n*sqrt(2) < 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012973 | OEIS-A298112

Solution b( ) of the complementary equation a(n) = a(1)*b(n) - a(0)*b(n-1) + 3*n, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, b(2) = 5, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjecture: 2/5 < a(n) - n*sqrt(2) < 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012974 | OEIS-A298113

Solution b( ) of the complementary equation a(n) = a(1)*b(n) - a(0)*b(n-1) + 4*n, where a(0) = 1, a(1) = 2, b(0) = 3, b(1) = 4, b(2) = 5, and (b(n)) is the increasing sequence of positive integers not in (a(n)). See Comments.

Conjecture: 1/2 < a(n) - n*sqrt(2) < 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012975 | OEIS-A298206

a(n) is the smallest b >= 2 such that b^(6*2^n) - b^(3*2^n) + 1 is prime.

a(13) = 165394 is a significant outlier from the generally expected trend, which can be conjectured to be 6*2^n*gamma, where gamma is the Euler-Mascheroni constant A001620. Additionally, the next b > a(13) such that b^(6*2^n) - b^(3*2^n) + 1 is prime is 165836, which is remarkably close to a(13). - _Serge Batalov_, Jan 24 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Serge Batalov_, Jan 14 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012976 | OEIS-A298210

Smallest n such that A001542(a(n)) == 0 (mod n), i.e., x=A001541(a(n)) and y=A001542(a(n)) is the fundamental solution of the Pell equation x^2 - 2*(n*y)^2 = 1.

Stronger, but conjectured:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _A.H.M. Smeets_, Jan 15 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012977 | OEIS-A298211

Smallest n such that A001353(a(n)) == 0 (mod n), i.e., x=A001075(a(n)) and y=A001353(a(n)) is the fundamental solution of the Pell equation x^2 - 3*(n*y)^2 = 1.

a(k*A005385(n)) = a(k)*A005384(n) for n>2 and k > 0 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _A.H.M. Smeets_, Jan 15 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012978 | OEIS-A298267

a(n) is the maximum number of heptiamonds in a hexagon of order n.

Conjectures from _Colin Barker_, Jan 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Craig Knecht_, Jan 15 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012979 | OEIS-A298300

Analog of Motzkin numbers for Coxeter type D.

Conjectural algebraic equation: 3*t+2+(3*t^2+5*t-2)*f(t)+(3*t^3-t^2)*f(t)^2 = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _F. Chapoton_, Jan 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012980 | OEIS-A298368

Triangle read by rows: T(n, k) = floor((n-1)/2)*floor(n/2)*floor((k-1)/2)*floor(k/2).

T(n, k) is conjectured by Zarankiewicz's conjecture to be the crossing number of the complete bipartite graph K_{k,n}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Jan 17 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012981 | OEIS-A298468

Solution (aa(n)) of the system of 3 complementary equations in Comments.

Conjecture: For t >= 1, aa(2t) = 5t+1(+1 if binary expansion of t ends in an odd number of 0's), and for t >= 0, aa(2t+1) = 5t+4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012982 | OEIS-A298472

Numbers n such that n and n-1 are both nontrivial binomial coefficients.

Blokhuis et al. show that the values given are complete up to 10^30, and conjecture that there are no more.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christopher E. Thompson_, Jan 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012983 | OEIS-A298607

Powers of 2 with the digit '0' in their decimal expansion.

The complement, A238938, is conjectured to be finite. Furthermore, Khovanova (see link) believes 2^86 = 773712524553362671811952 64 is the largest power of 2 not in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alonso del Arte_, Jan 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012984 | OEIS-A298613

Primes formed by the concatenation of 2^k-1 and 2^(k-1)-1.

Conjectures: The first conjecture has been disproved. In fact, a(36)+1 is divisible by 23^2. - _Paolo Galliani_, Aug 27 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Galliani_, Jan 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012985 | OEIS-A298643

Array A(n, k) read by antidiagonals downwards: k-th base-n non-repunit prime p such that all numbers resulting from switching any two adjacent digits in the base-n representation of p are prime, where k runs over the positive integers, i.e., the offset of k is 1.

Conjecture: All rows of the array are infinite. If the above conjecture is false, then this should have keyword "tabf" rather than "tabl".

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Jan 24 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012986 | OEIS-A298689

G.f. A(x) satisfies: A(x) = Sum_{n>=0} binomial(n^2, n) * x^n / A(x)^(n^2).

a(2^k) is odd for k>=0, and a(n) is even elsewhere except at n=0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 24 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012987 | OEIS-A298788

Coordination sequence for bey tiling (or net) with respect to a trivalent node.

Conjectures from _Colin Barker_, Jan 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012988 | OEIS-A298789

Coordination sequence for bey tiling (or net) with respect to a tetravalent node.

Conjectures from _Colin Barker_, Jan 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012989 | OEIS-A298790

Partial sums of A298788.

Conjectures from _Colin Barker_, Jan 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012990 | OEIS-A298791

Partial sums of A298789.

Conjectures from _Colin Barker_, Jan 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012991 | OEIS-A298792

Coordination sequence for the bil tiling (or net) with respect to a trivalent node of the first type.

Conjectures from _Colin Barker_, Mar 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012992 | OEIS-A298793

Coordination sequence for the bil tiling (or net) with respect to a trivalent node of the second type.

Conjectures from _Colin Barker_, Mar 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012993 | OEIS-A298794

Coordination sequence for the bil tiling (or net) with respect to a trivalent node of the third type.

Conjectures from _Colin Barker_, Mar 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012994 | OEIS-A298795

Coordination sequence for the bil tiling (or net) with respect to a trivalent node of the fourth type.

Conjectures from _Colin Barker_, Mar 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012995 | OEIS-A298799

Expansion of (1-27*x)^(-1/9).

Conjecture: a(p*n) == a(n) (mod p^2) for prime p == 1 (mod 9) and all positive integers n except those n of the form n = m*p + k for 0 <= m <= (p-1)/9 and 1 <= k <= (p-1)/9. Cf. A034171, A004981 and A004982. - _Peter Bala_, Dec 23 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Jun 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012996 | OEIS-A298801

Fourth column of triangular array in A296339.

Conjectures from _Colin Barker_, Feb 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 02 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012997 | OEIS-A298819

a(n) is the period of the oscillating pattern formed by a 1 X n line of cells in the Life-like cellular automaton B2c3-cekq4ikt5i8/S2-in3-acky4aijry5eiky6i, or 0 if the pattern vanishes.

a(2k) = A268754(k); a(2k+1) = lcm(A268754(k), A268754(k+1)) (conjectured). Note that, by an arbitrary/cosmetic convention, A268754 uses 1 for empty patterns and A298819 uses 0. - _WG Zeist_ and _Charlie Neder_, Jan 11 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _WG Zeist_, Jan 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012998 | OEIS-A298856

Triangular numbers k for which A240542(k) = A240542(k-1).

Conjecture 1: Numbers m such that 2*A033676(m) = A033677(m) - 1. - _Omar E. Pol_, Feb 07 2026 Conjecture 2: The positive terms of A014105 are the numbers m whose symmetric representation of sigma(m) has two subparts that meet at the center of the largest Dyck path. - _Omar E. Pol_, Feb 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Jan 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-012999 | OEIS-A298865

The primes p and products 4*p in increasing order.

Conjecture: except for the first term, these are the nonsquares n for which there is a unique pair (x,y) such that x^2 - y^2 = n and x > y >= 0; see A257408.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 13 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013000 | OEIS-A298904

a(n) is the number of distinct primes produced by starting with the n-th prime p and repeatedly looking at all the prime factors of 2p-1, and then performing the same process (double, subtract 1, find all prime factors) with those primes; a(n) = -1 if this produces infinitely many primes.

Just as for A305382, it is conjectured that a(n) is finite for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013001 | OEIS-A298943

Lower of two consecutive Mersenne prime exponents with record first difference.

Conjecture: The sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Jan 30 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013002 | OEIS-A299017

Intersection of A264041 and A000217.

Conjectures (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Barker_, Jan 31 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013003 | OEIS-A299037

a(n) is the number of rooted binary trees with minimal Sackin index and n leaves.

Conjecture 1: column 3 gives the nonzero terms of A000217. Conjecture 2: column 4 gives the nonzero terms of A000330. (End) Conjecture 3: column 5 gives the nonzero terms of A212415. Conjecture 4: row sums give 1 together with A006893. Conjecture 5: T(i,j) has i >= 0 where "i" is the height of the rooted trees. Conjecture 6: For i >= 1, j is the number of leaves minus 2^(i-1). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mareike Fischer_, Feb 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013004 | OEIS-A299104

Number of SNPN-equivalence classes of Boolean functions of n or fewer variables.

Number of Boolean functions distinct under simultaneous complementation of the inputs and/or permutation of the inputs, complementation of the output. Algorithm includes elements that are conjectural in nature. Documentation of the algorithm is at web link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marko Riedel_, Feb 15 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013005 | OEIS-A299110

Prime(r) for r such that prime(r) - prime(r-1) = 12 and prime(r-1) - prime(r-2) = 2.

Conjecture: Sequence has infinitely many terms. Note: p~2(12,2) is just one particular form of a prime of A056240-type k=2; there are others, e.g., 2(18,2), 2(18,4), 2(28,12), 2(24,10). All such prime sequences are also conjectured to produce infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Feb 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013006 | OEIS-A299116

The number of sparse union-closed sets. That is, the number of union-closed sets on n elements containing the empty set and the universe, such that in average each set (not counting the empty set) has at most n/2 elements.

If there is a counterexample to the union-closed set conjecture, it is a sparse union-closed set.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gunnar Brinkmann_, Feb 05 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013007 | OEIS-A299154

a(n) is the distance from prime(n) to the next prime that has the same digital root, or a(n) = 0 if there is no greater prime with the same digital root.

I conjecture that there are infinitely many prime numbers with each of the digital roots from the set {1,2,4,5,7,8}. This would imply that the only prime number with a(n) = 0 is 3. The conjecture is true; for any d in {1,2,4,5,7,8}, gcd(d, 9) = 1, hence, according to Dirichlet's theorem on arithmetic progressions, there are infinitely many primes of the form d + 9*k, and these primes all have digital root d. - _Rémy Sigrist_, Mar 25 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimitris Valianatos_, Feb 03 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013008 | OEIS-A299272

Coordination sequence for "flu" 3D uniform tiling formed from tetrahedra, rhombicuboctahedra, and cubes.

Conjectures from _Colin Barker_, Feb 11 2018: (Start) G.f.: (x^2+1)*(x^4+3*x^3+5*x^2+3*x+1)*(x+1)^3 / (1-x^3)^3. - _N. J. A. Sloane_, Feb 12 2018 (This confirms my conjecture from Feb 10 2018 and the above conjecture from _Colin Barker_.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013009 | OEIS-A299273

Partial sums of A299272.

Conjectures from _Colin Barker_, Feb 11 2018: (Start) These conjectures are correct. - _N. J. A. Sloane_, Feb 12 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013010 | OEIS-A299274

Coordination sequence for "hal" 3D uniform tiling.

However, this g.f. is so much more complicated than the g.f.s for any of the other 27 3D uniform tilings, at present I am only willing to state it as a conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013011 | OEIS-A299275

Partial sums of A299274.

There is a conjectured g.f., see the g.f. for A299274 and divide by 1-x. Note: this should not be used to generate a b-file. - _N. J. A. Sloane_, Feb 13 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013012 | OEIS-A299285

Coordination sequence for "tea" 3D uniform tiling.

[I suspect Barker's formulas only conjectures. - _N. J. A. Sloane_, Jun 12 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013013 | OEIS-A299286

Partial sums of A299285.

Conjectures from _Colin Barker_, Feb 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013014 | OEIS-A299289

Coordination sequence for "tsi" 3D uniform tiling.

Conjectures from _Colin Barker_, Feb 11 2018: (Start) Conjectured e.g.f.: ((4 + 13*x + 13*x^2)*cosh(x) + (3 + 13*x + 13*x^2)*sinh(x) - 2)/2. - _Stefano Spezia_, Jun 08 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013015 | OEIS-A299290

Partial sums of A299289.

Conjectures from _Colin Barker_, Feb 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013016 | OEIS-A299322

Ranks of {2,3}-power towers with neither consecutive 2's nor consecutive 3's; see Comments.

Conjectures from _Colin Barker_, Feb 09 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013017 | OEIS-A299405

Solution (a(n)) of the system of 5 complementary equations in Comments.

Conjecture: for all n >= 0,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013018 | OEIS-A299406

Dirichlet g.f.: Sum_{n>0} a(n)/n^s = (zeta(s)*zeta(6*s))/(zeta(2*s)*zeta(3*s)).

Conjecture: a(n) = A008836(n) * A210826(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Feb 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013019 | OEIS-A299409

Solution (e(n)) of the system of 5 complementary equations in Comments.

Conjecture: for all n >= 0,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013020 | OEIS-A299438

a(n) is the smallest number k such that there do not exist k numbers in the range [1..n] whose product is equal to their sum or -1 if no such number exists.

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Feb 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013021 | OEIS-A299439

a(n) is the smallest number k such that there exists n numbers in the range [1..k] whose product is equal to its sum.

Conjecture: the next number for which a(n) = n is 4354 (see A299438).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Feb 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013022 | OEIS-A299537

Number of ways to write n^2 as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers and z <= w such that x or y is a power of 4 (including 4^0 = 1) and x + 3*y is also a power of 4.

Conjecture (i): a(n) > 0 for all n > 0, and a(n) = 1 only for n = 4^k*m with k = 0,1,2,... and m = 1, 2, 3, 5, 7, 11, 15, 19, 43, 47, 135, 1103. Conjecture (ii): For any integer n > 1, we can write n^2 as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that 2*x or 2*y is a power of 4 and 2*(x+3*y) is also a power of 4. We have verified that the conjecture for n up to 10^7. See also the related comments in A300219 and A300360, and a similar conjecture in A299794.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013023 | OEIS-A299637

Solution (b(n)) of the system of 5 complementary equations in Comments.

Conjecture: for all n >= 0,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013024 | OEIS-A299638

Solution (c(n)) of the system of 5 complementary equations in Comments.

Conjecture: for all n >= 0,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013025 | OEIS-A299639

a(0) = 0, and for n > 0, (a(n)) gives the indices n for which d(n) > d(k) for k < n, where d is the difference sequence of (cos k + sin k).

Conjecture: d(n) -> 1.356...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013026 | OEIS-A299641

Solution (d(n)) of the system of 5 complementary equations in Comments.

Conjecture: for all n >= 0, Conjectures from _Chai Wah Wu_, Sep 07 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013027 | OEIS-A299704

List of primes prime(r) such that prime(r)-prime(r-1)=30, prime(r-1)-prime(r-2)=8 and prime(r-2)-prime(r-3)=6.

Conjecture: Sequence has infinitely many terms. Note: p~3(30,8,6) is one particular form of a prime of a056240-type 3; there are others, e.g., 3(30,12,2), 3(24,6,2), 3(36,6,4), 3(38,10,2), etc. All such prime

sequences are also conjectured to produce infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Feb 17 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013028 | OEIS-A299741

Array read by antidiagonals upwards: a(i,0) = 2, i >= 0; a(i,1) = i+2, i >= 0; a(i,j) = (i+2) * a(i,j-1) - a(i,j-2), for i >= 0, j > 1.

p[j](n) = p[j-1](n) * p[1](n) - p[j-2](n), for j > 1, n >= 0. The coefficients of these polynomials occur as the even numbered, upward diagonals in the OEIS Wiki link. Conjecture: a(i,j) = p[j](i), i,j >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _William W. Collier_, Feb 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013029 | OEIS-A299754

Number of distinct sums of n complex n-th roots of 1.

For prime p, a(p) = binomial(2*p-1,p). - Conjectured by _Robert Israel_, Feb 18 2018; proved by _Max Alekseyev_, Feb 20 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David W. Wilson_, Feb 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013030 | OEIS-A299794

Number of ways to write n^2 as x^2 + y^2 + z^2 + w^2 with x >= y >= 0 <= z <= w such that x or 2*y is a power of 4 (including 4^0 = 1) and x + 15*y is also a power of 4.

Conjecture 1: a(n) > 0 for all n > 0. Also, for any integer n > 1 we can write n^2 as x^2 + y^2 + z^2 + w^2 with x >= y >= 0 <= z <= w such that 2*x or y is a power of 4 and also x + 15*y = 2^(2k+1) for some k = 0,1,2,.... Conjecture 2: Let d be 2 or 8, and let r be 0 or 1. Then any positive square n^2 can be written as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that x or y is a power of 2 and x + d*y = 2^(2k+r) for some k = 0,1,2,.... We have verified Conjecture 1 for n up to 10^7. See also A299537, A300219 and A300396 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013031 | OEIS-A299807

Rectangular array read by antidiagonals: T(n,k) is the number of distinct sums of k complex n-th roots of 1, n >= 1, k >= 0.

The following are all conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Feb 24 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013032 | OEIS-A299825

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that x <= y, x == y (mod 2), and |x+y-z| is a power of 4 (including 4^0 = 1).

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 2, 6, 13, 15, 18, 27, 43, 53, 63, 93, 107, 109, 123, 141, 159, 165, 173, 477, 493, 653, 1005, 16^k*m (k = 0,1,2,... and m = 3, 4, 7, 8). A weaker version of the conjecture was proved by the author in arXiv:1701.05868.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013033 | OEIS-A299896

Coordination sequence for "bar" 3D tiling with respect to second type of node.

For this sequence there is not even a conjectured g.f.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013034 | OEIS-A299898

Coordination sequence for "ftu" 3D tiling with respect to first type of node.

The following is a conjectured recurrence, found by gfun, using the command rec:=gfun[listtorec](t1, a(n)); (where t1 is a list of the initial terms) suggested by _Paul Zimmermann_. Conjectured g.f.: (-x^21 - 3*x^20 - 5*x^19 - 8*x^18 - 13*x^17 - 18*x^16 - 22*x^15 - 27*x^14 - 32*x^13 - 35*x^12 - 36*x^11 - 36*x^10 - 35*x^9 - 32*x^8 - 27*x^7 - 22*x^6 - 17*x^5 - 11*x^4 - 7*x^3 - 5*x^2 - 3*x - 1) / ((x - 1)^3*(1 + x + x^2)*(1 + x^4)*(1 + x^3 + x^6)*(1 + x + x^2 + x^3 + x^4 + x^5 + x^6)). Conjecture: a(n) = 11/7*n^2 - 13/63*n + 31/42 + (((3*n^2+3*n+1) mod 7) + (n mod 7) + ((n+1) mod 7) - ((n+2) mod 7) - ((n+4) mod 7))/7 + ((n^2+n+2) mod 4) + (4*((n+5) mod 8) + 3*((n+4) mod 8) - 4*((n+7) mod 8) - 3*(n mod 8))/8 + (((7*n+4) mod 9) + 2*(n mod 9) + ((n+1) mod 9) + ((n+2) mod 9) - ((n+3) mod 9) - 2*((n+4) mod 9) - 2*((n+5) mod 9) + ((n+8) mod 9))/9 - [n = 0]. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013035 | OEIS-A299900

Coordination sequence for "ftu" 3D tiling with respect to second type of node.

I have to say that I am not very confident about the conjectured g.f. below. We only have 128 terms, and the recurrence has 57 or so coefficients and 57 or so initial terms, and 114 is pretty close to 128. So not much of a safety margin. If the same g.f. still holds when we get 500 terms I will believe it. - _N. J. A. Sloane_, Jan 03 2021 This empirically corroborates the conjectured generating function/recurrence above to the 500-term margin requested in the 2021 comment by _N. J. A. Sloane_. The submitted 501-term table supplies the requested margin. The terms were computed independently from the RCSR CGD description of the ftu net: expanding the Fm-3m symmetry gives a translation-labeled quotient graph with 288 vertices and 576 undirected edges, every vertex of degree 4. The same quotient graph gives A299898 from node type 1 (root vertex 0), agreeing with its existing b-file, and gives this sequence from node type 2 (root vertex 192). The first 128 A299900 terms agree with the existing b-file, and the extension through n = 500 agrees with the proposed rational generating function P(x)/Q(x). The leading growth term in the conjectured formula is (11/7)*n^2. Although Q has quasi-polynomial period 1008, only the constant coefficient has full period 1008; the n^2 coefficient is constant and the n coefficient has period 12. The following is a conjectured recurrence, found by gfun, using the command rec:=gfun[listtorec](t1, a(n)); (where t1 is a list of

the initial terms) suggested by _Paul Zimmermann_. Conjectured g.f., empirically corroborated by the 501-term table below: A(x) = P(x)/Q(x), where P and Q are given in the linked PARI and Python programs as coefficient lists in ascending powers of x. The corresponding degree-53 recurrence from Q is satisfied by the 501-term table for n = 58..500. For this conjectured g.f. the quasi-polynomial form is a(n) = (11/7)*n^2 + b(n mod 12)*n + c(n mod 1008), for n >= 5, where b = [-58/63, -389/504, -52/63, -443/504, -46/63, -437/504, -58/63, -347/504, -52/63, -485/504, -46/63, -395/504]. The constant coefficient c has period 1008.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013036 | OEIS-A299910

Coordination sequence of point of type 3.3.4.3.4 in 3-uniform tiling #3.54 in the Galebach listing.

Conjectures from _Chai Wah Wu_, Jul 10 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013037 | OEIS-A299911

Coordination sequence of point of type 3.3.3.4.4 in 3-uniform tiling #3.54 in the Galebach listing.

Conjectures from _Chai Wah Wu_, Jul 10 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013038 | OEIS-A299924

Number of ways to write n^2 as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that x + 2*y + 3*z is a power of 4 (including 4^0 = 1).

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 3, 7, 13, 49, 61, 4^k*m (k = 0,1,2,... and m = 1, 2, 11, 14, 17).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

B07 A300000 - A349999 | 2,414 条

V3-013039 | OEIS-A300000

The sum of the first n terms of the sequence is the concatenation of the first n digits of the sequence, with a(1) = 1.

Conjecture: a(n) contains all ten digits for n >= 4280. (End) Conjecture holds through a(169000). - _Michael S. Branicky_, Jul 08 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_, Feb 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013040 | OEIS-A300001

Side length of the smallest equilateral triangle that can be dissected into n equilateral triangles with integer sides, or 0 if no such triangle exists.

a(n) = A290820(n) for n <= 8. It is conjectured that a(n) < A290820(n) for all n > 12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Feb 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013041 | OEIS-A300002

Lexicographically earliest sequence of positive integers such that no k+2 points fall on any polynomial of degree k.

Conjecture: This sequence is a permutation of the natural numbers. Conjecture: a(27)-a(32) are 11, 32, 21, 25, 13, 47. If all previous data are correct, no polynomial of degree ceiling(n/2.5) - 1 goes through any set of points. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Apr 17 2017 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013042 | OEIS-A300004

First differences of the list A292794 of numbers not congruent to F(k) modulo F(k+1) for any k > 1, where F are the Fibonacci numbers A000045.

It has been conjectured that every even number occurs in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013043 | OEIS-A300005

Index of first occurrence of 2n in A300004 (or -1 of 2n does not occur), where A300004 are first differences of numbers not congruent to A000045(k) modulo A000045(k+1) for any k > 1.

It is only conjectured that every even number occurs in sequence A300004.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Mar 05 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013044 | OEIS-A300012

A permutation of the positive integers in which adjacent terms differ by a single prime factor and a(n) has no prime factor exceeding prime(n), having the lexicographically earliest inverse.

This permutation can be viewed as a tour of the positive integers, multiplying or dividing by a prime at each step. The "lexicographically earliest inverse" condition means no other such tour visits an integer k earlier, unless it visits an integer less than k later than this tour does (but see the proviso addressed by conjectures 1 and 2 below). The condition "a(n) has no prime factor exceeding prime(n)" is included to ensure the sequence is well-defined. The author anticipates the condition may be redundant, in the sense that if it is removed: (conjecture 1) the lexicographically earliest inverse remains defined; and (conjecture 2) the terms remain the same. Conjecture: A006530(a(n)) <= n; lim sup A006530(a(n))/n < 0.4. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Munn_, Jun 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013045 | OEIS-A300059

Numbers k such that k and k+1 are terms of A279767.

Conjecture: If a run of consecutive terms that are also consecutive numbers of length m exists in this sequence, then there also exists a run of length m + 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Torlach Rush_, Feb 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013046 | OEIS-A300139

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with 4*x - 3*y a square, where x,y,z,w are nonnegative integers with z <= w such that 10*x or y is a square.

Conjecture 1: a(n) > 0 for all n >= 0, and a(n) = 1 only for n = 16^k*m with k = 0,1,2,... and m = 0, 7, 8, 13, 22, 23, 24, 31, 44, 56, 71, 79, 88, 109, 120, 152, 184, 472, 1912, 6008, 9080. Conjecture 2: Each n = 0,1,2,... can be written as x^2 + y^2 + z^2 + w^2 with 3*x - y twice a square, where x,y,z,w are nonnegative integers such that 5*x or y is a square. See also A281976, A300666, A300667, A300708 and A300712 for similar conjectures. a(n) > 0 for all n = 0..10^8. Also, Conjecture 2 holds for all n = 0..10^8. - _Zhi-Wei Sun_, Oct 05 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 12 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013047 | OEIS-A300219

Number of ways to write n^2 as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers and z <= w such that both x and 4*x - 3*y are powers of 4 (including 4^0 = 1).

Conjecture: (i) a(n) > 0 for all n > 0, and a(n) = 1 only for n = 4^k*m (k = 0,1,2,... and m = 1, 2, 3, 5, 15, 37, 83, 263). Also, for each n = 2,3,... we can write n^2 as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that both x and 4*x - 3*y lie in the set {2^(2k+1): k = 0,1,...}. We have verified both parts of the conjecture for n up to 10^7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 28 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013048 | OEIS-A300286

Define a set of generalized Syracuse sequences starting with x(1)=2*n+1 a positive odd integer, if x(i) is odd prime set x(i+1)=67*x(i)+1, if x(i) is odd not prime set x(i+1)=3*x(i)+1 and if x(i) is even then set x(i+1)=x(i)/2. Then a(n) is the first index i > 1 at which x(i) reaches 1.

All the sequences for p < 423 eventually enter a loop (not tested above, but I conjecture that it is the case for any prime, although with different end cycles).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre CAMI_, Mar 02 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013049 | OEIS-A300307

Number of solutions to 1 +- 3 +- 6 +- ... +- n*(n+1)/2 == 0 mod n.

a(2^n) = 2^A000325(n) for n>0 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Mar 02 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013050 | OEIS-A300326

Sum of the largest possible permutations that can be written without repetition of digits in each base from binary to n+1.

For primes 5 <= p <= 23, a(n) (mod p) is p(p-1) periodic. I conjecture this to hold for all p >= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Cano_, Mar 03 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013051 | OEIS-A300356

Number of ways to write n^2 as x^2 + y^2 + z^2 + w^2 with x >= y >= 0 <= z <= w such that x + 63*y = 2^(2k+1) for some nonnegative integer k.

Conjecture: (i) a(n) > 0 for all n > 1, and a(n) = 1 only for n = 5, 13, 25, 29, 59, 61, 91, 95, 101, 103, 211, 247, 2^k (k = 1,2,...), 4^k*79 (k = 0,1,2,...), 2^(2k+1)*m (k = 0,1,2,... and m = 3, 5, 7, 11, 15, 19, 23). See also A299924 and A300219 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 03 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013052 | OEIS-A300360

Number of ways to write n^2 as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers and z <= w such that x or y is a power of 2 (including 1) and x + 63*y = 2^(2k+1) for some k = 0,1,2,....

Conjecture: a(n) > 0 for all n > 1, and a(n) = 1 only for n = 5, 13, 25, 29, 59, 61, 79, 91, 95, 101, 103, 1315, 2^k (k = 1,2,3,...), 2^(2k+1)*m (k = 0,1,2,... and m = 3, 5, 7, 11, 15, 19, 23). In the comments of A300219, the author conjectured that a positive square n^2 can be written as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that both x and x + 3*y are powers of 4 unless n has the form 4^k*81503 with k a nonnegative integer. Since 81503^2 = 208^2 + 16^2 + 51167^2 + 63440^2 with 16 = 4^2 and 208 + 3*16 = 4^4, this implies that any positive square can be written as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that x or y is a power of 4 and x + 3y is also a power of 4. We also conjecture that for any positive integer n not of the form 4^k*m (k =0,1,... and m = 2, 7) we can write n^2 as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that x or y is a power of 4 and x + 2*y is also a power of 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 03 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013053 | OEIS-A300362

Number of ways to write n^2 as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that x + 2*y and (z + 2*w)/3 are squares and w is even.

Conjecture: a(n) > 0 for all n = 0,1,2,..., and a(n) = 1 only for n = 0, 7, 9, 14, 19, 22, 26, 34, 41, 4^k*m (k = 0,1,... and m = 1, 2, 3, 5, 10, 11, 13, 15).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013054 | OEIS-A300386

The number of paths of length 7*n from the origin to the line y = 2*x/5 with unit East and North steps that stay below the line or touch it.

The sequence defined by b(n) := [x^n] A(x)^n begins [1, 3, 161, 9804, 630401, 41789278, 2824792568, 193553976353, ...] and conjecturally satisfies the congruence b(p) == b(1) (mod p^3) for prime p >= 11 (checked up to p = 101). - _Peter Bala_, Sep 14 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bryan T. Ek_, Mar 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013055 | OEIS-A300387

The number of paths of length 9*n from the origin to the line y = 2*x/7 with unit East and North steps that stay below the line or touch it.

The sequence defined by b(n) := [x^n] A(x)^n begins [1, 4, 372, 39298, 4384884, 504464254, 59183637186, 7038517648243, ...] and conjecturally satisfies the congruence b(p) == b(1) (mod p^3) for prime p >= 11 (checked up to p = 101). - _Peter Bala_, Sep 14 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bryan T. Ek_, Mar 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013056 | OEIS-A300388

The number of paths of length 11*n from the origin to the line y = 2*x/9 with unit East and North steps that stay below the line or touch it.

The sequence defined by b(n) := [x^n] A(x)^n begins [1, 5, 715, 116213, 19954187, 3532860880, 637870220023, 116749388814357, ...] and conjecturally satisfies the congruence b(p) == b(1) (mod p^3) for prime p >= 5 except for p = 11 (checked up to p = 101). - _Peter Bala_, Sep 14 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bryan T. Ek_, Mar 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013057 | OEIS-A300389

The number of paths of length 13*n from the origin to the line y = 2*x/11 with unit East and North steps that stay below the line or touch it.

The sequence defined by b(n) := [x^n] A(x)^n begins [1, 6, 1222, 282993, 69239846, 17468997381, 4494716943847, 1172353182893367, ...] and conjecturally satisfies the congruence b(p) == b(1) (mod p^3) for prime p >= 5 except for p = 11 and p = 13 (checked up to p = 101). - _Peter Bala_, Sep 14 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bryan T. Ek_, Mar 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013058 | OEIS-A300392

a(1)=1, a(n) = max(sum0,sum1) mod n, where sum0 is the sum of previous terms with even indices, sum1 with odd indices.

Conjecture: every integer appears infinitely many times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Mar 05 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013059 | OEIS-A300396

Number of ways to write n^2 as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers and z <= w such that 2*x or y is a power of 4 (including 4^0 = 1) and x + 63*y = 2^(2k+1) for some k = 0,1,2,....

Conjecture: a(n) > 0 for all n > 1, and a(n) = 1 only for n = 5, 13, 25, 29, 59, 61, 79, 91, 95, 101, 103, 1315, 2^k (k = 1,2,3,...), 2^(2k+1)*m (k = 0,1,2,... and m = 3, 5, 7, 11, 15, 19, 23, 887). This is stronger than the conjecture that A300360(n) > 0 for all n > 1. Note that a(387) = 3 < A300360(387) = 4 and a(1774) = 1 < A300360(1774) = 2. See also A299537, A299794 and A300219 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 05 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013060 | OEIS-A300399

a(n) = number of primes that end in 7 among the first 10^n primes.

Conjecture: lim_{n->oo} a(n+1)/a(n) = 10. - _Stefano Spezia_, Sep 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _G. L. Honaker, Jr._, Mar 05 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013061 | OEIS-A300448

Number of the integers 4^k*m with k >= 0 and m = 1, 2, 5 such that n^2 - 4^k*m can be written as the sum of two squares.

Conjecture 1: a(n) > 0 for all n > 0. Conjecture 2: Each positive square n^2 can be written as 4^k*m + x^2 + y^2 with k,x,y nonnegative integers and m among 1, 5 17. Conjecture 3: For each n = 1,2,3,... we can write n^2 as 4^k*m + x^2 + y^2 with k,x,y nonnegative integers and m among 1, 5, 65. Conjecture 3 implies that A300441(n) > 0 for all n > 0 since 4*A001353(0)^2 + 1 = 1, 4*A001353(1)^2 + 1 = 5 and 4*A001353(2)^2 + 1 = 65. We have verified Conjectures 1-3 for all n = 1..2*10^7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 06 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013062 | OEIS-A300510

Number of ways to write n^2 as 4^k*(m^2+1) + x^2 + y^2, where m is 1 or 2, and k,x,y are nonnegative integers with x <= y.

Conjecture: a(n) > 0 for all n > 1; in other words, for any integer n > 1 there is a nonnegative integer k such that either n^2 - 2*4^k or n^2 - 5*4^k can be written as the sum of two squares. Moreover, a(n) = 1 only for n = 2^k with k > 0. This conjecture is stronger than the first conjecture in A300448. We have verified that a(n) > 0 for all n = 2..5*10^7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013063 | OEIS-A300531

Matching number of the n-polygon diagonal intersection graph.

Conjecturally, a matching exists for every n with at most one unmatched vertex. This would imply a(n) = floor(A007569(n)/2). A computer search, using a simple nonbacktracking algorithm, has shown the existence of such matchings up to n = 22 and indeed for small n there are large numbers of maximum matchings (A292921). Such a matching could also be constructed from a Hamiltonian path (A300551) by selecting every other edge, so a proof that these graphs are Hamiltonian would also suffice. - _Andrew Howroyd_, Mar 12 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Mar 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013064 | OEIS-A300589

a(n) = A300617(n) / (n*(n+1)/2) for n>=1.

It is conjectured that this sequence consists entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013065 | OEIS-A300593

O.g.f. A(x) satisfies [x^n] exp(n^2 * A(x)) = n^3 * [x^(n-1)] exp(n^2 * A(x)) for n >= 1.

It is conjectured that this sequence consists entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013066 | OEIS-A300595

O.g.f. A(x) satisfies [x^n] exp(n^3 * A(x)) = n^3 * [x^(n-1)] exp(n^3 * A(x)) for n >= 1.

It is conjectured that this sequence consists entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013067 | OEIS-A300617

O.g.f. A(x) satisfies [x^n] exp(n * A(x)) = n^2 * [x^(n-1)] exp(n * A(x)) for n >= 1.

It is conjectured that this sequence consists entirely of integers. a(n) is divisible by n*(n+1)/2 (conjecture); A300589(n) = a(n) / (n*(n+1)/2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013068 | OEIS-A300666

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with nonnegative integers x,y,z,w and z <= w such that x or 2*y is a square and x + 3*y is also a square.

Conjecture 1: a(n) > 0 for all n = 0,1,2,.... Conjecture 2: Any nonnegative integer n not equal to 3 can be written as x^2 + y^2 + z^2 + w^2 with nonnegative integers x,y,z,w such that x or 2*y is a square and 3*x - y is also a square. See also A281976, A300667, A300708 and A300712 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013069 | OEIS-A300667

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers and z <= w such that 3*x or y is a square and x + 2*y is also a square.

Conjecture 1: a(n) > 0 for all n >= 0, and a(n) = 1 only for n = 16^k*m with k = 0,1,2,... and m = 0, 3, 7, 11, 12, 15, 28, 39, 47, 60, 71, 92, 119, 172, 232, 253, 263, 316, 347, 515. Conjecture 2: Each n = 0,1,2,... can be written as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that 3*x or y is a square and 2*x - y is also a square. See also A281976, A300666, A300708 and A300712 for similar conjectures. a(n) > 0 for all n = 0..10^8. Also, Conjecture 2 holds for all n = 0..10^8. In a 2018 paper Y.-C. Sun and Z.-W. Sun proved that any nonnegative integer can be written as x^2 + y^2 + z^2 + w^2 with x + 2*y a square, where x,y,z,w are nonnegative integers. - _Zhi-Wei Sun_, Oct 04 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013070 | OEIS-A300708

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers and z <= w such that x or y is a square and x - y is also a square.

Conjecture: a(n) > 0 for all n >= 0, and a(n) = 1 only for n = 16^k*m with k = 0,1,2,... and m = 0, 8, 12, 23, 24, 44, 47, 56, 71, 79, 92, 95, 140, 168, 184, 248, 344, 428, 568, 632, 1144, 1544. See also A281976, A300666, A300667 and A300712 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013071 | OEIS-A300712

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that 3*x or y is a square and x - y is twice a square.

Conjecture: a(n) > 0 for all n >= 0, and a(n) = 1 only for n = 16^k*m with k = 0,1,2,... and m = 0, 1, 3, 7, 21, 24, 46, 79, 88, 94, 142, 151, 184, 190, 193, 280, 286, 1336. See also A281976, A300666, A300667 and A300708 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013072 | OEIS-A300732

G.f. A(x) satisfies: [x^n] A(x/A(x)^(2*n)) = 0 for n>=1.

Conjecture: n divides a(n) for n>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013073 | OEIS-A300733

G.f. A(x) satisfies: [x^n] A(x/A(x)^(3*n)) = 0 for n>=1.

Conjecture: n divides a(n) for n>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013074 | OEIS-A300734

G.f. A(x) satisfies: [x^n] A(x/A(x)^(2*n^2)) = 0 for n>1.

Conjecture: n^2 divides a(n) for n>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013075 | OEIS-A300751

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x + 3*y + 5*z a positive square, where x,y,z,w are nonnegative integers such that 2*x or y or z is a square.

Conjecture: a(n) > 0 for all n = 1,2,3,..., and a(n) = 1 only for n = 16^k*m with k = 0,1,2,... and m = 0, 1, 4, 5, 6, 7, 8, 20, 24, 28, 31, 36, 43, 61, 71, 79, 100, 116, 157, 188, 200, 344, 351, 388, 632. This is stronger than the author's 1-3-5 conjecture in A271518. See also A300752 for a similar conjecture stronger than the 1-3-5 conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013076 | OEIS-A300752

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x + 3*y + 5*z a positive square, where x,y,z,w are nonnegative integers such that 3*x or y or z is a square.

Conjecture: a(n) > 0 for all n = 1,2,3,.... This is stronger than the author's 1-3-5 conjecture in A271518. See also A300751 for a similar conjecture stronger than the 1-3-5 conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013077 | OEIS-A300789

Heinz numbers of integer partitions whose Young diagram can be tiled by dominos.

This sequence is conjectured to be the Heinz numbers of integer partitions in which the odd parts appear as many times in even as in odd positions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Mar 12 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013078 | OEIS-A300791

Number of ways to write n as x^2 + y^2 + z^2 + w^2, where w is a positive integer and x,y,z are nonnegative integers for which x or y or z is a square and (12*x)^2 + (15*y)^2 + (20*z)^2 is also a square.

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 16^k*m with k = 0,1,2,... and m = 1, 3, 4, 6, 7, 21, 23, 24, 39, 47, 86, 95, 344, 651, 764. See also A300792 for two similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 12 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013079 | OEIS-A300792

Number of ways to write n as x^2 + y^2 + z^2 + w^2, where w is a positive integer and x,y,z are nonnegative integers such that x or y or z is a square and 9*x^2 + 16*y^2 + 24*z^2 is also a square.

Conjecture 1: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 16^k*m with k = 0,1,2,... and m = 1, 3, 6, 14, 21, 24, 56, 79, 119, 143, 248, 301, 383, 591, 728, 959, 1223, 1751, 2311, 6119. Conjecture 2: Any positive integer n can be written as x^2 + y^2 + z^2 + w^2, where w is a positive integer and x,y,z are nonnegative integers such that x or y or z is a square and 36*x^2 + 40*y^2 + 45*z^2 is also a square. See also A300791 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 12 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013080 | OEIS-A300844

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that x or 2*y or z is a square and (12*x)^2 + (21*y)^2 + (28*z)^2 is also a square.

Conjecture 1: a(n) > 0 for all n = 0,1,2,.... Conjecture 2: Any positive integer can be written as x^2 + y^2 + z^2 + w^2 with w a positive integer and x,y,z nonnegative integers such that x or y or z is a square and 144*x^2 + 505*y^2 + 720*z^2 is also a square. In 2016, the author conjectured in A271510 that each n = 0,1,2,... can be written as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers and y >= z such that (3*x)^2 + (4*y)^2 + (12*z)^2 is a square. See also A300791 and A300792 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 13 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013081 | OEIS-A300859

Where records occur in A045763.

Conjectures based on 1255 terms of a(n) < 36,000,000:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Mar 15 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013082 | OEIS-A300871

O.g.f. A(x) satisfies: [x^n] exp(n*(n+1) * A(x)) = n*(n+1) * [x^(n-1)] exp(n*(n+1) * A(x)) for n>=1.

It is conjectured that this sequence consists entirely of integers. a(n) is divisible by n*(n+1)/2 (conjecture); a(n) = n*(n+1)/2 * A300872(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 14 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013083 | OEIS-A300872

a(n) = A300871(n) / (n*(n+1)/2).

It is conjectured that this sequence consists entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 14 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013084 | OEIS-A300874

O.g.f. A(x) satisfies: [x^n] exp(n*(n+1) * A(x)) = 2*n * [x^(n-1)] exp(n*(n+1) * A(x)) for n>=1.

It is conjectured that this sequence consists entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 14 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013085 | OEIS-A300891

Left-truncatable nonzero triangular numbers.

This sequence is finite and full. - Conjectured by _Stanislav Sykora_, Mar 14 2018; proved by _Max Alekseyev_, Jun 03 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, Mar 14 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013086 | OEIS-A300896

Lexicographically earliest sequence of distinct positive terms such that, for any n > 0, a(n) + a(n+1) + a(n+2) is a Fibbinary number (A003714).

This sequence is conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Mar 14 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013087 | OEIS-A300908

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with (3*x)^2 + (4*y)^2 + (12*z)^2 a square , where w is a positive integer and x,y,z are nonnegative integers such that z or 2*z or 3*z is a square.

Conjecture 1: a(n) > 0 for all n > 0. Moreover, any positive integer can be written as x^2 + y^2 + z^2 + w^2 with (3*x)^2 + (4*y)^2 + (12*z)^2 a square, where w is a positive integer and x,y,z are nonnegative integers for which one of z, z/2, z/3 is a square and z/2 (or z/3) is a power of 4 (including 1). Conjecture 2: Each n = 0,1,2,... can be written as x^2 + y^2 + z^2 + w^2 with (3*x)^2 + (4*y)^2 + (12*z)^2 a square, where x,y,z,w are nonnegative integers for which x or z is a square or y = 2^k for some k = 0,1,2,.... Conjecture 3. Let a,b,c be positive integers with a <= b <= c and gcd(a,b,c) = 1. If each n = 0,1,2,... can be written as x^2 + y^2 + z^2 + w^2 with x,y,z,w integers such that (a*x)^2 + (b*y)^2 + (c*z)^2 is a square, then (a,b,c) must be among the three triples (3,4,12), (12,15,20) and (12,21,28). We have verified Conjectures 1 and 2 for n up to 5*10^5 and 10^6 respectively. See also A300791 and A300844 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 15 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013088 | OEIS-A300949

Carmichael numbers whose prime factors form an arithmetic progression.

Conjecture: if n is a Carmichael number and lpf(n)gpf(n)(lpf(n)+gpf(n))/2 = n, then (lpf(n)+gpf(n))/2 is prime; and thus n has exactly three prime factors. Such numbers n form a proper subsequence of this sequence, also subsequence of A262723. - _Charles R Greathouse IV_ and _Thomas Ordowski_, Mar 17 2018. Edited by _Max Alekseyev_, Mar 17 2018 Proof of the above conjecture: Say n = paq with 2 < p < q being primes and a = (p+q)/2, with (a,p!)=1. If n is a Carmichael number, then pa == 1 (mod q-1), so p^2 + pq == 2 (mod q-1), so p^2 + p == 2 (mod q-1). In particular, p^2 + p - 2 >= q-1, which implies that (p+1)^2 > q. Say a has k prime factors, so that a >= (p+2)^k. But a < q, so q > (p+2)^k. Thus, (p+1)^2 > q > (p+2)^k. This implies k=1. - Carl Pomerance (in a letter to the second author), Mar 18 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_ and _Thomas Ordowski_, Mar 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013089 | OEIS-A300984

Numbers whose sum of squarefree divisors and sum of nonsquarefree divisors are both squarefree numbers.

Conjecture: a(n) is of the form a(n) = 2^i*p^j with i, j integers and p prime. This has been verified for n up to 10^7. Generalization: For n <= 10^m with m > 7, it is conjectured that a majority of primes p where a(n) = 2^i*p^j are in A082539. For example, with m = 7, 84% of the primes p are in A082539.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Mar 17 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013090 | OEIS-A301283

Coordination sequence for node of type V1 in "car" 2-D tiling (or net).

Conjectures from _Colin Barker_, Mar 30 2018: (Start) Equivalent conjecture: 12*a(n) = 37*n+4*b(n)+6*(-1)^(n/2)*A142150(n+2)+3*c(n) for n>2, where b(n)=0,-1,1 (3-periodic, n>=0) and c(n) = -6,5,6,-5 (4-periodic, n>=0). - _R. J. Mathar_, Mar 31 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013091 | OEIS-A301285

Coordination sequence for node of type V2 in "car" 2-D tiling (or net).

Conjectures from _Colin Barker_, Mar 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013092 | OEIS-A301286

Partial sums of A301285.

Conjectures from _Chai Wah Wu_, Sep 05 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013093 | OEIS-A301287

Coordination sequence for node of type 3.12.12 in "cph" 2-D tiling (or net).

Equivalent conjecture: 3*a(n) = 8*n+2*A057078(n+1)+3*A228826(n+2). - _R. J. Mathar_, Mar 31 2018 (This is now a theorem. - _N. J. A. Sloane_, Apr 05 2018)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013094 | OEIS-A301310

G.f.: Sum_{n>=0} 2^n * (1+x)^(n^2) / 3^(n+1).

a(n) is divisible by 10 for n>0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013095 | OEIS-A301314

Number of ways to write n as x^2 + y^2 + z^2 + w^2, where w is a positive integer and x,y,z are nonnegative integers such that x + 3*y + 9*z = 2^k*m^3 for some k,m = 0,1,2,....

Conjecture 1: a(n) > 0 for all n > 0. Conjecture 2: Any positive integer can be written as x^2 + y^2 + z^2 + w^2, where x is a positive integer and y,z,w are nonnegative integers such that 2*x + 7*y = 2^k*m^3 for some k = 0,1,2 and m = 1,2,3,.... See also A301303 and A301304 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013096 | OEIS-A301370

Maximum determinant of an n X n (0,1)-matrix that has exactly 2*n ones.

A proved upper bound is abs(a(n)) <= 6^(n/6), provided by Bruhn and Rautenbach. A conjectured sharper bound is abs(a(n)) <= 2^(n/3), provided by the same authors. For n=3*k, the bound is achieved by diagonally concatenating blocks ((1 1 0)(0 1 1)(1 0 1)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Mar 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013097 | OEIS-A301372

Conjectured best solution for the problem stated in A098072.

Conjectured best solution for the problem stated in A098072.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Mar 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013098 | OEIS-A301375

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x,y,z nonnegative integers and w a positive integer such that x*(y+3*z) is a cube or half a cube and y <= z <= w if x = 0.

Conjecture: a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013099 | OEIS-A301376

Number of ways to write n^2 as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers and z <= w such that x^2-(3*y)^2 = 4^k for some k = 0,1,2,....

Conjecture: a(n) > 0 for all n > 0. Moreover, any positive square n^2 can be written as x^2 + y^2 + z^2 + w^2 with x,y,z,w integers and y even such that x^2 - (3*y)^2 = 4^k for some k = 0,1,2,.... Compare this conjecture with the conjectures in A299537. As 3*A001353(n)^2 + 1 = A001075(n)^2, the conjecture in A300441 implies that any positive square can be written as x^2 + y^2 + z^2 + w^2 with x,y,z,w integers such that x^2 - 3*y^2 = 4^k for some k = 0,1,2,.... See also A301391 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013100 | OEIS-A301391

Number of ways to write n^2 as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers and y even such that x^2 - (6*y)^2 = 4^k for some k = 0,1,2,....

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 11, 13, 19, 2^k*m (k = 0,1,2,... and m = 1, 5, 7, 31). See also A301376 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013101 | OEIS-A301420

Sums of positive coefficients in generalized Chebyshev polynomials of the first kind, for a family of 5 data.

G.f.: (-x*(x+1)^4+1)/(x^6+4*x^5+5*x^4-5*x^2-6*x+1); this denominator equals (1-x)*(2-(x+1)^5) (conjectured). a(n+14) = 7*a(n+13) - a(n+12) - 6*a(n+11) + 2*a(n+10) - a(n+9) + 4*a(n+8) + a(n+7) + 4*a(n+5) + 2*a(n+4) - a(n+3) - 5*a(n+2) - 4*a(n+1) - a(n) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gregory Gerard Wojnar_, Mar 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013102 | OEIS-A301421

Sums of positive coefficients of generalized Chebyshev polynomials of the first kind, for a family of 6 data.

G.f.: (-x*(x+1)^5+1)/(x^7+5*x^6+9*x^5+5*x^4-5*x^3-9*x^2-7*x+1); this denominator equals (1-x)*(2-(1+x)^6) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gregory Gerard Wojnar_, Mar 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013103 | OEIS-A301424

Sums of positive coefficients of generalized Chebyshev polynomials of the first kind, for a family of 7 data.

G.f.: (-x*(x+1)^6+1)/(x^2*(x^6+6*x^5+14*x^4+14*x^3-14*x-14)-8*x+1); this denominator equals (1-x)*(2-(1+x)^7) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gregory Gerard Wojnar_, Mar 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013104 | OEIS-A301426

Number of steps required in the worst case for three knights to find the princess in a castle with n rooms arranged in a line (Castle and princess puzzle).

This conjecture is a(n) = A194222(n) for n>2. - _R. J. Mathar_, Mar 27 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Mar 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013105 | OEIS-A301432

G.f.: Sum_{n>=0} 3^n * (1+x)^(n^2) / 4^(n+1).

a(n) is divisible by 21 for n>0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013106 | OEIS-A301452

Number of ways to write n^2 as m*4^k + x^2 + 2*y^2 with m in the set {2, 3} and k,x,y nonnegative integers.

Conjecture: a(n) > 0 for all n > 1. We call this the 2-3 conjecture. It is simialr to the author's 2-5 conjecture which states that A300510(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013107 | OEIS-A301471

Number of ways to write n^2 as x^2 + 2*y^2 + 3*2^z with x,y,z nonnegative integers.

Square Conjecture: a(n) > 0 for all n > 1. Moreover, for any integer n > 3 we can write n^2 as x^2 + 2*y^2 + 3*2^z, where x,y,z are nonnegative integers with y even and z > 1. The Square Conjecture implies that for each n = 1,2,3,... we can write 3*n^2 as x^2 + 2*y^2 + 2^z with x,y,z nonnegative integers. In fact, if (3*n)^2 = u^2 + 2*v^2 + 3*2^z with u,v,z integers and z >= 0, then u^2 == v^2 (mod 3) and thus we may assume u == v (mod 3) without loss of generality, hence 3*n^2 = (u^2+2*v^2)/3 + 2^z = x^2 + 2*y^2 + 2^z with x = (u+2*v)/3 and y = (u-v)/3 integers. On March 25, 2018 Qing-Hu Hou at Tianjin Univ. finished his verification of the Square Conjecture for n <= 4*10^8. Then I used Hou's program to verify the conjecture for n <= 5*10^9. - _Zhi-Wei Sun_, Apr 10 2018 I have found a counterexample to the Square Conjecture, namely a(5884015571) = 0. Note that 5884015571 is the product of the three primes 7, 17 and 49445509. - _Zhi-Wei Sun_, Apr 15 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013108 | OEIS-A301479

Positive integers m such that m^3 cannot be written in the form x^2 + 2*y^2 + 3*2^z with x,y,z nonnegative integers.

It seems that this sequence has infinitely many terms. In contrast, the author conjectured in A301471 and A301472 that any square greater than one can be written as x^2 + 2*y^2 + 3*2^z with x,y,z nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013109 | OEIS-A301483

a(n) = floor(a(n-1)/(2^(1/3)-1)) with a(1)=1.

Conjectures from _Colin Barker_, Apr 01 2018: (Start) a(n) = A195350(n) + A303647(n-2) - A195339(n-4) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gregory Gerard Wojnar_, Mar 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013110 | OEIS-A301510

Smallest positive number b such that ((b+1)^prime(n) + b^prime(n))/(2*b + 1) is prime, or 0 if no such b exists.

Conjecture: a(n) > 0 for every n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tim Johannes Ohrtmann_, Mar 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013111 | OEIS-A301517

Numbers whose ratio (sum of nonsquarefree divisors)/(sum of squarefree divisors) is a positive integer.

Conjecture: if r = A162296(a(n)) / A048250(a(n)) is a perfect square, r belongs to A001248.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Mar 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013112 | OEIS-A301534

Number of ways to write the n-th prime congruent to 7 modulo 12 as x^2 + 3*y^2 + 15*2^z with x,y,z nonnegative integers.

Conjecture: a(n) > 0 for all n > 1. In other words, any prime p > 7 with p == 7 (mod 12) can be written as x^2 + 3*y^2 + 15*2^z with x,y,z nonnegative integers. We have verified the conjecture for all primes p == 7 (mod 12) with 7 < p < 8*10^9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013113 | OEIS-A301554

Expansion of Product_{k>=1} ((1 + x^k)/(1 - x^k))^(sigma_0(k)).

Conjecture: log(a(n)) ~ Pi * sqrt(n*log(n)/2). - _Vaclav Kotesovec_, Sep 03 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Mar 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013114 | OEIS-A301579

Least nonnegative integer k such that n^2 - 3*2^k can be written as x^2 + 2*y^2 with x and y integers, or -1 if no such k exists.

The Square Conjecture in A301471 implies that a(n) >= 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013115 | OEIS-A301601

Numbers k such that k^6 can be written as a sum of 11 positive 6th powers.

Conjecture: 35 is the largest integer not in the sequence. - _Jon E. Schoenfield_, Mar 24 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Mar 24 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013116 | OEIS-A301640

Largest integer k such that n^2 - 3*2^k can be written as x^2 + 2*y^2 with x and y integers, or -1 if no such k exists.

Conjecture: a(n) > 0.6*log_2(log_2 n) for all n > 2, and also lim inf_{n->infinity} a(n)/(log n) = 0. The author's Square Conjecture in A301471 would imply that a(n) >= 0 for all n > 1. We have verified that a(n) > 0.6*log_2(log_2 n) for all n = 3..4*10^9. For n = 2857932461, we have a(n) = 3 and 0.603 < a(n)/log_2(log_2 n) < 0.604.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013117 | OEIS-A301653

Expansion of x*(1 + 2*x)/((1 - x)*(1 + x)*(1 - x - x^2)).

a(n) = Sum_{k>=0} A051601(n-k,k) (conjectured). - _Greg Dresden_, May 18 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Mar 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013118 | OEIS-A301674

Coordination sequence for node of type V1 in "krs" 2-D tiling (or net).

Equivalent conjecture: 9*a(n) = 40*n -18*(-1)^n -6*(-1)^n*A076118(n+1) +6*A049347(n) -4*A049347(n-1). - _R. J. Mathar_, Apr 01 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013119 | OEIS-A301680

Coordination sequence for node of type V2 in "krn" 2-D tiling (or net).

Equivalent conjecture: 7*a(n) = 36*n-2*b(n) for n>3, where b(n>=1) = 4, 1, -2, 2, -1, -4, 0 (continued 7-periodic). - _R. J. Mathar_, Mar 30 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013120 | OEIS-A301697

Coordination sequence for node of type V2 in "krj" 2-D tiling (or net).

Equivalent conjecture: 6*a(n) = 32*n -3*A010892(n-1) + A049347(n-1) for n>0. - _R. J. Mathar_, Nov 26 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013121 | OEIS-A301701

a(n) is the smallest positive integer m, with the property that n appears as a coefficient in the polynomial P_m(x) = (x-1)(x^2-1)...(x^m-1).

We conjecture that all integers appear as a coefficient of a polynomial P_m(x). The conjecture holds for the first 10^5 positive integers, with a maximum on those integers of a(99852) = 1921. - _David A. Corneth_, Apr 08 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ovidiu Bagdasar_, Mar 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013122 | OEIS-A301708

Coordination sequence for node of type V1 in "krc" 2-D tiling (or net).

Conjecture: a(n) = (i*((-i)^n - i^n) + 22*n) / 4 where i=sqrt(-1). - _Colin Barker_, Apr 07 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013123 | OEIS-A301710

Coordination sequence for node of type V2 in "krc" 2-D tiling (or net).

a(n) = ((-i)^(1+n) + i^(1+n) + 22*n) / 4 for n>0, where i=sqrt(-1) (conjectured). - _Colin Barker_, Apr 07 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013124 | OEIS-A301712

Coordination sequence for node of type V1 in "usm" 2-D tiling (or net).

Equivalent conjecture: 5*a(n) = 27*n -b(n) -5*A014017(n-2) for n>0, where b(n) = 2,-1,1,-2,0 (5-periodic) for n>=1. - _R. J. Mathar_, Mar 30 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013125 | OEIS-A301718

Coordination sequence for node of type V2 in "kre" 2-D tiling (or net).

Conjecture: a(n) ~ 28*n/5. - _Stefano Spezia_, Mar 29 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013126 | OEIS-A301719

Partial sums of A301718.

Conjecture: a(n) ~ 14*n^2/5. - _Stefano Spezia_, Mar 28 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013127 | OEIS-A301743

Lexicographically first sequence with no duplicate term whose digits' concatenation is the same as the digits' concatenation of all sums of adjacent terms lined up one by one.

This sequence is conjectured to be a permutation of A000027 (the positive numbers).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Mar 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013128 | OEIS-A301746

Expansion of Product_{k>=1} (1 + x^k)^(sigma_0(k)^2).

Conjecture: log(a(n)) ~ sqrt(n) * log(n)^(3/2) / (2*sqrt(6)). - _Vaclav Kotesovec_, Aug 29 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Mar 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013129 | OEIS-A301757

Positive determinant values assumed by performing all permutations of entries in the 3 X 3 matrix of A301372.

A 3 X 3 matrix with given 9 matrix entries can produce A088021(3)=10080 distinct determinants if all positional permutations are performed. The current sequence provides the 5040 positive determinants of a conjectured optimal matrix minimizing its greatest matrix entry.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Mar 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013130 | OEIS-A301771

Number of odd chordless cycles in the n-Andrásfai graph.

a(11) - a(35) were computed from the definition, not from the conjectured formula. - _Pontus von Brömssen_, Oct 09 2022 Conjectured g.f.: x*(x - 3*x^3 + 4*x^4 + x^5)/(1 - x)^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Mar 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013131 | OEIS-A301776

Prime numbers p with the property that all even numbers n (2 < n <= 2p) are the sum of two primes <= p.

Conjecture: this sequence is finite - it has 7 terms only. Conjecture verified up to first 10^5 primes. This sequence is related to the Goldbach Strong Conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marcin Barylski_, Mar 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013132 | OEIS-A301807

Lexicographically first sequence of distinct integers whose concatenation of digits is the same as the concatenation of the digits of the absolute differences between consecutive terms.

In fact, a(18420) = 42; however that this sequence is a permutation of the natural numbers remains an open question.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Mar 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013133 | OEIS-A301858

Positive integers which can be written as the sum of two squares but cannot be written as x^2 + y^2 + 2*z^2 with x and y integers and z a nonzero integer.

Conjecture 1: The sequence only has the four terms 1, 5, 29 and 65. Conjecture 2: For any integer n > 1 which is neither 17 nor a power of 2, if n = u^2 + 2*v^2 for some integers u and v, then n = x^2 + 2*y^2 + 3*z^2 for some integers x,y,z with z nonzero. Conjecture 3: For any positive integer n not of the form 4^k*m (k = 0,1,2,... and m = 1, 7, 13), if n = u^2 + 3*v^2 for some integers u and v, then n = x^2 + 2*y^2 + 3*z^2 for some integers x,y,z with y nonzero.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013134 | OEIS-A301877

Group the natural numbers into groups (1),(2),(3),(4),(5,6),(7,8,9),... so that the n-th group contains N(n) terms, where N(n) is the Narayana's cows sequence (A000930). Sequence contains the sum of the terms in the n-th group.

Conjectures from _Colin Barker_, Mar 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Mar 28 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013135 | OEIS-A301891

Number of ways to write 2*n^2 as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that x + 3*y + 5*z + 15*w is a power of 4.

Conjecture 1: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 4^k*m with k = 0,1,2,... and m = 2, 3, 6, 10, 38. Conjecture 2: For any positive integer n, we can write 2*n^2 as x^2 + y^2 + z^2 + w^2 with x,y,z,w nonnegative integers such that x + 3*y + 5*z + 15*w is twice a power of 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 28 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013136 | OEIS-A301892

a(n) = A010846(A002182(n)).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Mar 28 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013137 | OEIS-A301893

Numbers m that set records for the ratio A010846(m)/A000005(m).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Mar 28 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013138 | OEIS-A301921

Expansion of e.g.f. 1/(1 - (exp(x) - 1)/(1 - (exp(x) - 1)^2/(1 - (exp(x) - 1)^3/(1 - ...)))), a continued fraction.

Conjecture: Let k be a positive integer. The sequence obtained by reducing a(n) modulo k is eventually periodic with the period dividing phi(k) = A000010(k). For example, modulo 9 we obtain the sequence [0, 1, 3, 1, 6, 4, 0, 7, 3, 1, 6, 4, 0, 7, 3, 1, 6, 4, 0, 7, ...] with an apparent period of 6 = phi(9) beginning at n = 2. Cf. A004123. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Jun 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013139 | OEIS-A301925

a(n) is the smallest prime p such that the gap between p and the next prime is 4*n.

It is a plausible conjecture that a(n) always exists. - _N. J. A. Sloane_, Jul 21 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_ and _Robert G. Wilson v_, Jun 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013140 | OEIS-A301933

G.f. A(x) satisfies: A(x) = x*(1 + 4*A(x)*A'(x)) / (1 + A(x)*A'(x)).

a(n = 2^k) is odd for k>=0, and a(n) is even elsewhere (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 28 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013141 | OEIS-A301976

Number of no-leaf subgraphs of the 3 X n grid.

Conjecture: a(n) mod 10 = 3 for n > 2. Conjectures from _Colin Barker_, Mar 30 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Mar 29 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013142 | OEIS-A301977

a(n) is the number of distinct positive numbers whose binary digits appear in order but not necessarily as consecutive digits in the binary representation of n.

a(n) = Sum_{k=1..n} (Stirling2(n+1,k) mod 2) (conjecture). - _Ilya Gutkovskiy_, Jul 04 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Mar 30 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013143 | OEIS-A301981

Euler transform of A034448.

Conjecture: a(n) ~ exp((3*Pi*n)^(2/3)/2 - 1/2) * A^6 / (2 * 3^(5/6) * Pi^(1/3) * n^(5/6)), where A is the Glaisher-Kinkelin constant A074962.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Mar 30 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013144 | OEIS-A301982

Expansion of Product_{k>=1} (1 + x^k)^A034448(k).

Conjecture: a(n) ~ exp(3 * Pi^(2/3) * n^(2/3) / 2^(5/3)) / (2^(4/3) * sqrt(3) * Pi^(1/6) * n^(2/3)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Mar 30 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013145 | OEIS-A302092

Product of n-th Bell number and n-th Bell number written backwards.

Conjecture: in this sequence only two semiprimes (4,25).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Apr 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013146 | OEIS-A302109

Smallest integer N such that there are exactly n cyclic groups C_2 in the multiplicative group of integers modulo N when decomposed as a product of cyclic groups C_{k_1} x C_{k_2} x ... x C_{k_m}, and k_i divides k_j for i < j.

Conjecture: (a) a(j) is divisible by a(i) if i < j; (b) a(n)/4 is squarefree for all n for n >= 2. [Corrected by _Alper Ferudun_ and _Jianing Song_, Oct 18 2026] Conjecture (b) above is obvious according to this description, while conjecture (a) is false: The smallest multiple N of a(11) such that (Z/NZ)* contains 12 copies of C_2 is N = 179*a(11) = 156451974672904680, but a(12) is at most 120025194365695080. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Apr 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013147 | OEIS-A302112

Number of forests with 2n nodes and n labeled trees. Also number of forests with exactly n edges on 2n labeled nodes.

a(n) ~ c * 2^n * exp(n) * n^(n - 2/3), where c = 0.2305818... = 1 / (2^(1/6) * 3^(1/3) * Gamma(1/3)) [symbolic expression for c is conjectural]. - _Vaclav Kotesovec_, Jul 20 2019, updated Feb 20 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Apr 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013148 | OEIS-A302174

Smallest solution x of x^n + y^(n+1) = z^(n+2), x, y, z >= 1.

We may conjecture that, for n > 4, a(n) is given by these upper limits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jacques Tramu_, Apr 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013149 | OEIS-A302178

The number of 3D walks of semilength n in a quadrant returning to the origin.

Conjecture D-finite with recurrence (n+1)*(n+2)^2*a(n) +2*(-29*n^3-39*n^2+8*n+6)*a(n-1) +36*(4*n-1)*(2*n-3)*(3*n-2)*a(n-2) -648*(2*n-5)*(n-2)*(2*n-3)*a(n-3)=0. - _R. J. Mathar_, Oct 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013150 | OEIS-A302179

The number of 3D walks of length n in an octant returning to axis of origin.

Conjecture D-finite with recurrence 114*(n+5)*(n+4)*(n+2)*a(n) +(-1669*n^3-12592*n^2-26619*n-13644)*a(n-1) +6*(19*n^3+2125*n^2+4411*n-1131)*a(n-2) +8*(7895*n^3-9549*n^2-22925*n+14082)*a(n-3) +48*(-3553*n^3+14958*n^2-6986*n-16383)*a(n-4) -432*(n-4)*(223*n^2-1110*n+1898)*a(n-5) +864*(n-5)*(779*n-1962)*(n-4)*a(n-6) -518400*(n-5)*(n-6)*(n-4)*a(n-7)=0. - _R. J. Mathar_, Oct 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013151 | OEIS-A302180

Number of 3D walks of type aad.

Conjecture D-finite with recurrence (n+4)*(n+2)*a(n) +3*(-n^2-3*n-1)*a(n-1) -13*(n-1)*(n+1)*a(n-2) +15*(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Oct 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013152 | OEIS-A302181

Number of 3D walks of type abb.

Conjecture D-finite with recurrence (n+1)^3*a(n) -2*(2*n-1)*(10*n^2+10*n+3)*a(n-1) +36*n*(2*n-1)*(2*n-3)*a(n-2)=0. - _R. J. Mathar_, Oct 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013153 | OEIS-A302183

Number of 3D n-step walks of type abd.

Conjecture D-finite with recurrence (n+2)^2*a(n) +(-3*n^2-5*n-1)*a(n-1) -(13*n+15)*(n-1)*a(n-2) +15*(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Oct 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013154 | OEIS-A302184

Number of 3D walks of type abe.

Conjecture D-finite with recurrence (n+2)^2*a(n) +2*(-3*n^2-5*n-1)*a(n-1) -4*(n+3)*(n-1)*a(n-2) +24*(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Oct 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013155 | OEIS-A302186

Number of 3D walks of type ace.

Conjecture D-finite with recurrence (n+3)*(n+2)*a(n) +4*(-3*n^2-8*n-3)*a(n-1) +8*(4*n^2+3*n-3)*a(n-2) +48*(n-2)^2*a(n-3) +144*-(n-2)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Oct 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013156 | OEIS-A302187

Number of 3D walks of type bcc.

Conjecture D-finite with recurrence (n+1)*(n+2)^2*a(n) +2*(7*n^3-17*n-6)*a(n-1) -4*(4*n-1)*(4*n^2+9*n-12)*a(n-2) +8*(-70*n^3+330*n^2-461*n+225)*a(n-3) +48*(n-3)*(23*n^2-66*n+64)*a(n-4) +288*(7*n-17)*(n-3)*(n-4)*a(n-5) -3456*(n-5)*(n-3)*(n-4)*a(n-6)=0. - _R. J. Mathar_, Oct 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013157 | OEIS-A302188

Number of 3D walks of type bce.

Conjecture D-finite with recurrence (n+1)^2*a(n) -2*(n+2)*(2*n+1)*a(n-1) +4*(-4*n^2+9*n+1)*a(n-2) +8*(2*n+5)*(n-2)*a(n-3) +48*(n-2)*(n-3)*a(n-4)=0. - _R. J. Mathar_, Oct 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013158 | OEIS-A302249

a(n) is the number of divisors of A279254(n) in Gaussian integers.

Conjecture: a(14) = 63 is the largest odd term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Apr 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013159 | OEIS-A302298

Wiener index of the graph of nodes (i,j) of the square lattice such that abs(i) + abs(j) <= n.

Conjectures from _Colin Barker_, Apr 08 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andres Cicuttin_, Apr 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013160 | OEIS-A302334

A weighted smoothing applied to the primes as a data set: a(n) = floor(A007443(2n-1)/2^(2n-2)), where A007443 is binomial transform of primes.

Conjecture: the second differences are in [-2,2].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Munn_, Apr 05 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013161 | OEIS-A302392

Number of odd parts in the partitions of 3n into 3 parts.

Conjectures from _Colin Barker_, Apr 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Apr 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013162 | OEIS-A302393

Number of even parts in the partitions of 3n into 3 parts.

Conjectures from _Colin Barker_, Apr 07 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Apr 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013163 | OEIS-A302495

a(n) is the least k such that k * A005277(n) is a term of A002202.

Conjecture: Every k > 1 eventually appears in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Torlach Rush_, Jun 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013164 | OEIS-A302599

Numbers k such that digit_sum(k) > digit_sum(2k).

Conjecture: a(n) ~ 2n. - _Charles R Greathouse IV_, Apr 10 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Consiglio, Jr._, Apr 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013165 | OEIS-A302607

a(n) is the greatest possible least prime in any partition of prime(n) into three primes; n >= 4.

Goldbach's weak (ternary) conjecture states that every odd number > 5 can be expressed as the sum of three primes (see link). This sequence applies the conjecture (now proved) to primes > 5. From all possible partitions of prime(n) = p+q+r for primes p,q,r (p <= q <= r), a(n) is chosen to be the maximum possible value of the least prime p. The sequence is not strictly increasing, and although many primes are repeated, some do not appear at all (e.g., 37 and 47 are not included).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Apr 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013166 | OEIS-A302613

Total domination number of the n-folded cube graph.

Conjecture: a(n) = 2^(n-4) for n >= 7. - _Eric W. Weisstein_, Aug 27 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Apr 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013167 | OEIS-A302641

Number of nonnegative integers k such that n^2 - 3*2^k can be written as x^2 + 2*y^2 with x and y integers.

The author's Square Conjecture in A301471 implies that a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013168 | OEIS-A302647

a(n) = (2*n^2*(n^2 - 3) - (2*n^2 + 1)*(-1)^n + 1)/64.

Also the graph crossing number of the (n+2)-barbell graph (assuming Guy's conjecture). - _Eric W. Weisstein_, May 17 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Apr 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013169 | OEIS-A302659

Lexicographically earliest sequence of distinct positive terms such that, for any n > 0, A065368(a(n) * a(n+1)) = 0.

- f_b is conjectured to be a permutation of the natural numbers,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Apr 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013170 | OEIS-A302689

a(n) = 4 + 2^n - 4*n.

a(n) = A258547(n-5) for n > 5 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Apr 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013171 | OEIS-A302694

a(n) is the smallest integer k such that A002828(k*n) = 3.

Conjecture: a(n) <= 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Apr 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013172 | OEIS-A302756

a(n) is the least possible greatest prime in any partition of prime(n) into three primes; n >= 4.

Goldbach's weak (ternary) conjecture states that every odd number > 5 can be expressed as the sum of three primes (see link). This sequence applies the conjecture (now proven) to primes > 5. From all possible partitions of prime(n) = p+q+r for primes p,q,r (p <= q <= r), a(n) is chosen as the least possible value of the greatest prime r (with lower prime p not constrained to be A302607(n)). The sequence is not strictly increasing, and although many primes appear repeatedly, some do not appear at all (e.g. 59 is not included).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Apr 12 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013173 | OEIS-A302864

a(n) is the smallest positive palindromic number whose binary expansion ends in exactly n zeros, or 0 if no such number exists.

Conjecture: a(n) > 0 for all n >= 0. If A082613(n)/A082614(n) = 2^n (see conjecture at A082613) and a(n) > 0, then a(n) >= A082613(n). - _Altug Alkan_, Apr 15 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Apr 14 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013174 | OEIS-A302920

Number of ways to write prime(n)^2 as x^2 + 2*y^2 + 3*2^z with x,y,z nonnegative integers.

Conjecture: a(n) > 0 for all n > 0. In other words, for any prime p there are nonnegative integers x, y and z such that x^2 + 2*y^2 + 3*2^z = p^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 15 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013175 | OEIS-A302930

Maximum number of 6's possible in an infinite Minesweeper grid with n mines.

Conjecture: a(n) = Max_{i=2..floor(n/2)} floor((n-2*i)*(i-1)/i) = n+1-floor(sqrt(8*n-1)) for n >= 6. - _Zhuorui He_, Jun 27 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Apr 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013176 | OEIS-A302982

Number of ways to write n as x^2 + 5*y^2 + 2^z + 3*2^w with x,y,z,w nonnegative integers.

Conjecture: a(n) > 0 for all n > 3. See also A302983 and A302984 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013177 | OEIS-A302983

Number of ways to write n as x^2 + 2*y^2 + 2^z + 3*2^w with x,y,z,w nonnegative integers.

Conjecture: a(n) > 0 for all n > 3. See also A302982 and A302984 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013178 | OEIS-A302984

Number of ways to write n as x^2 + 2*y^2 + 2^z + 5*2^w with x,y,z,w nonnegative integers.

Conjecture: a(n) > 0 for all n > 5. See also A302982 and A302983 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013179 | OEIS-A302985

Number of ordered pairs (x, y) of nonnegative integers such that n - 2^x - 3*2^y has the form u^2 + 2*v^2 with u and v integers.

Conjecture: a(n) > 0 for all n > 3. This is equivalent to the author's conjecture in A302983. We have verified a(n) > 0 for all n = 4...6*10^9. See also A302982 and A302984 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013180 | OEIS-A302987

Primes p such that p+3 divides 3^p+3.

Conjecture: the sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Apr 17 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013181 | OEIS-A303112

Primes p such that (r-q)/(q-p) = 2 or 1/2, and p < q < r are three consecutive primes.

Conjecture: The two most frequent ratios between consecutive prime gaps are 2 and 1/2, and both ratios occur with about the same frequency. Conjecture: lim_{n->inf} n/primepi(a(n)) > k > 0 for some k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andres Cicuttin_, Apr 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013182 | OEIS-A303120

Total area of all rectangles of size p X q such that p + q = n^2 and p <= q.

Conjectures from _Colin Barker_, Apr 19 2018 and Mar 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Apr 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013183 | OEIS-A303233

Number of ways to write n as a*(a+1)/2 + b*(b+1)/2 + 2^c + 2^d, where a,b,c,d are nonnegative integers with a <= b and c <= d.

Conjecture: a(n) > 0 for all n > 1. In other words, any integer n > 1 can be written as the sum of two triangular numbers and two powers of 2. a(n) > 0 for all n = 2..10^9. See A303234 for numbers of the form x*(x+1)/2 + 2^y with x and y nonnegative integers. See also A303363 for a stronger conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013184 | OEIS-A303234

Numbers of the form x*(x+1)/2 + 2^y with x and y nonnegative integers.

Conjecture: Any integer n > 1 can be written as the sum of two terms of the current sequence. This is equivalent to the author's conjecture in A303233.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013185 | OEIS-A303235

Number of ordered pairs (x, y) with 0 <= x <= y such that n - 2^x - 2^y can be written as the sum of two triangular numbers.

Conjecture: a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013186 | OEIS-A303273

Array T(n,k) = binomial(n, 2) + k*n + 1 read by antidiagonals.

T(2*n,n+4) = A186353(4*n+1) (conjectured). T(2*n,n+5) = A184103(8*n+1) (conjectured). T(n+2,2*n) = A249013(5*n+4) (conjectured). T(n+3,2*n) = A186384(5*n+3) = A186386(5*n+3) (conjectured). T(2*n+1,2*n+1) = A143689(2*n+1) (= A030503(3*n+3) (conjectured)). T(2*n,2*n+1) = A104249(2*n) = A093918(2*n+2) = A131355(4*n+1) (= A030503(3*n+5) (conjectured)). T(3*n,n) = A025742(3*n-1), n > 1 (conjectured). T(n*(n+1),n) = A211905(n+1), n > 0 (conjectured). floor(T(n,n/2)) = A267682(n) (conjectured). floor(T(n,n/3)) = A025742(n-1), n > 0 (conjectured). floor(T(n,n/4)) = A263807(n-1), n > 0 (conjectured). ceiling(T(n,2^n)/n) = A134522(n), n > 0 (conjectured). ceiling(T(n,n/2+n)/n) = A051755(n+1) (conjectured). floor(T(n,n)/n) = A133223(n), n > 0 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franck Maminirina Ramaharo_, Apr 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013187 | OEIS-A303338

Number of ways to write n as x^2 + 2*y^2 + 3*2^z + 4^w with x,y,z,w nonnegative integers.

Conjecture: a(n) > 0 for all n > 3. This is stronger than the author's previous conjecture in A302983. It has been verified that a(n) > 0 for all n = 4..10^9. Jiao-Min Lin (a student at Nanjing University) has found a counterexample to the conjecture: a(12558941213) = 0. - _Zhi-Wei Sun_, Jul 30 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013188 | OEIS-A303363

Number of ways to write n as a*(a+1)/2 + b*(b+1)/2 + 2^c + 2^d, where a,b,c,d are nonnegative integers with a <= b, c <= d and 2|c.

Conjecture: a(n) > 0 for all n > 1. This is stronger than the author's conjecture in A303233. I have verified a(n) > 0 for all n = 2..10^9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013189 | OEIS-A303389

Number of ways to write n as a*(a+1)/2 + b*(b+1)/2 + 5^c + 5^d, where a,b,c,d are nonnegative integers with a <= b and c <= d.

Conjecture: a(n) > 0 for all n > 1. In other words, any integers n > 1 can be written as the sum of two triangular numbers and two powers of 5. See also A303401, A303432 and A303540 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013190 | OEIS-A303393

Numbers of the form x*(x+1)/2 + 5^y with x and y nonnegative integers.

The author's conjecture in A303389 has the following equivalent version: Each integer n > 1 can be expressed as the sum of two terms of the current sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013191 | OEIS-A303399

Number of ordered pairs (a, b) with 0 <= a <= b such that n - 5^a - 5^b can be written as the sum of two triangular numbers.

Conjecture: a(n) > 0 for all n > 1. This is equivalent to the author's conjecture in A303389. It has been verified that a(n) > 0 for all n = 2..6*10^9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013192 | OEIS-A303401

Number of ways to write n as a*(3*a-1)/2 + b*(3*b-1)/2 + 3^c + 3^d with a,b,c,d nonnegative integers.

Conjecture: a(n) > 0 for all n > 1. In other words, any integer n > 1 can be written as the sum of two pentagonal numbers and two powers of 3. a(n) > 0 for all n = 2..7*10^6. See A303434 for the numbers of the form x*(3*x-1)/2 + 3^y with x and y nonnegative integers. See also A303389 and A303432 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013193 | OEIS-A303403

Even numbers that are not the sum of two prime-indexed primes.

Bayless et al. conjectured that every even number larger than 80612 is the sum of two prime-indexed primes. If the conjecture is true then this sequence is finite with 733 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, May 13 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013194 | OEIS-A303428

Number of ways to write n as x*(3*x-2) + y*(3*y-2) + 3^u + 3^v, where x,y,u,v are integers with x <= y and 0 <= u <= v.

Conjecture: a(n) > 0 for all n > 1. Moreover, any integer n > 1 can be written as x*(3*x+2) + y*(3*y+2) + 3^z + 3^w, where x is an integer and y,z,w are nonnegative integers. See also A303389, A303401 and A303432 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013195 | OEIS-A303429

Number of ordered pairs (k, m) of nonnegative integers such that n - 3^k - 5^m can be written as the sum of two squares.

Conjecture: a(n) > 0 for all n > 1. This is equivalent to the author's conjecture in A303656.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 28 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013196 | OEIS-A303432

Number of ways to write n as a*(2*a-1) + b*(2*b-1) + 2^c + 2^d, where a,b,c,d are nonnegative integers with a <= b and c <= d.

Conjecture 1: a(n) > 0 for all n > 1. In other words, any integer n > 1 can be written as the sum of two hexagonal numbers and two powers of 2. Conjecture 2: Any integer n > 1 can be written as a*(2*a+1) + b*(2*b+1) + 2^c + 2^d with a,b,c,d nonnegative integers. Conjecture 3: Each integer n > 1 can be written as a*(2*a-1) + b*(2*b+1) + 2^c + 2^d with a,b,c,d nonnegative integers. All the three conjectures hold for n = 2..2*10^6. Note that either of them is stronger than the conjecture in A303233. See also A303363, A303389 and A303401 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013197 | OEIS-A303434

Numbers of the form x*(3*x-1)/2 + 3^y with x and y nonnegative integers.

The author's conjecture in A303401 has the following equivalent version: Each integer n > 1 can be written as the sum of two terms of the current sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013198 | OEIS-A303462

Lexicographically earliest sequence of distinct terms such that the digits succession stays the same if every second digit among all sequence terms is erased.

The sequence starts with a(1) = 1 and is always extended with the smallest integer not used so far that doesn't lead to a contradiction. It is conjectured that this sequence is a permutation of the integers > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Lars Blomberg_, Apr 24 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013199 | OEIS-A303539

Number of ordered pairs (k, m) with 0 <= k <= m such that n - binomial(2*k,k) - binomial(2*m,m) can be written as the sum of two squares.

Conjecture: a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013200 | OEIS-A303540

Number of ways to write n as a^2 + b^2 + binomial(2*c,c) + binomial(2*d,d), where a,b,c,d are nonnegative integers with a <= b and c <= d.

Conjecture: a(n) > 0 for all n > 1. In other words, any integer n > 1 can be written as the sum of two squares and two central binomial coefficients.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013201 | OEIS-A303541

Numbers of the form k^2 + binomial(2*m,m) with k and m nonnegative integers.

The conjecture in A303540 has the following equivalent version: Each integer n > 1 can be written as the sum of two terms of the current sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013202 | OEIS-A303543

Number of ways to write n as a^2 + b^2 + C(k) + C(m) with 0 <= a <= b and 0 < k <= m, where C(k) denotes the Catalan number binomial(2k,k)/(k+1).

Conjecture: a(n) > 0 for all n > 1. In other words, any integer n > 1 can be written as the sum of two squares and two Catalan numbers. This is similar to the author's conjecture in A303540. It has been verified that a(n) > 0 for all n = 2..10^9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013203 | OEIS-A303562

G.f. A(x) satisfies: 0 = [x^(n-1)] 1 / A(x)^(n^2+n-1) for n>2.

a(n) is odd iff n = (4^k - 1)/3 for k >= 0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013204 | OEIS-A303595

Index of n-th prime in A294994, or -1 if that prime is missing.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 28 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013205 | OEIS-A303601

Number of ways to write n as a*(a+1)/2 + b*(b+1)/2 + Bell(k) + Bell(m) with 0 <= a <= b and 0 < k <= m, where Bell(k) denotes the k-th Bell number A000110(k).

Conjecture: a(n) > 0 for all n > 1. In other words, any integer n > 1 can be expressed as the sum of two triangular numbers and two Bell numbers. See also A303389, A303540, A303543 and A303637 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013206 | OEIS-A303637

Number of ways to write n as x^2 + y^2 + 2^z + 5*2^w, where x,y,z,w are nonnegative integers with x <= y.

Conjecture: a(n) > 0 for all n > 5. Jiao-Min Lin (a student at Nanjing University) found a counterexample to the conjecture: a(18836421387) = 0. - _Zhi-Wei Sun_, Jul 21 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013207 | OEIS-A303647

a(n) = ceiling(a(n-1)/(2^(1/3)-1)+1), a(1)=1.

a(n) = A301483(n+2) - A195350(n+2) + A195339(n-3) for n > 2 (conjectured). [Corrected by _Qiyuan Gu_, Sep 18 2026] Conjectures from _Colin Barker_, Apr 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gregory Gerard Wojnar_, Apr 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013208 | OEIS-A303691

a(n) is the number of 3-smooth numbers k such that prime(n)-k is also a prime number, where prime(n) stands for the n-th prime.

Conjecture: a(n)>0 for all n>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Jun 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013209 | OEIS-A303702

Number of ways to write 2*n as p + 2^k + 3^m, where p is a prime, and k and m are nonnegative integers.

Conjecture: a(n) > 0 for all n > 1. In other words, any even number greater than 2 can be written as the sum of a prime, a power of 2 and a power of 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 29 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013210 | OEIS-A303712

a(n) is the smallest number such that there are exactly n numbers k (including a(n) itself) such that U(k) is isomorphic to U(a(n)) (or 0 if no such number exists). Here U(k) is the multiplicative group of integers modulo k.

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Apr 29 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013211 | OEIS-A303762

a(0) = 1, and for n >= 1, a(n) is either the largest divisor of a(n-1) not already present in the sequence, or (if all divisors already used), a(n-1) * {the least prime p such that p does not divide a(n-1) and p*a(n-1) is not already present}.

a(n) = A019565(A303769(n)). [Conjectured]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, May 03 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013212 | OEIS-A303769

a(0) = 0, a(n+1) is either the largest number obtained from a(n) by toggling a single 1-bit off (to 0) if no such number is yet in the sequence, otherwise the least number not yet in sequence that can be obtained from a(n) by toggling a single 0-bit on (to 1). In both cases the bit to be toggled is the rightmost possible that results yet an unencountered number.

The original, but now conjectural, alternative definition is: The equivalence of these definitions is still conjectural. The official definition of this sequence follows the latter one. - _Antti Karttunen_, Jun 08 2018 a(n) = A048675(A303762(n)). [The original definition, now conjectured]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, May 03 2018, with more direct definition from _David A. Corneth_, May 05 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013213 | OEIS-A303821

Number of ways to write 2*n as p + 2^x + 5^y, where p is a prime, and x and y are nonnegative integers.

Conjecture: a(n) > 0 for all n > 1. Moreover, for any integer n > 4, we can write 2*n as p + 2^x + 5^y, where p is an odd prime, and x and y are positive integers. See also A303934 and A304081 for further refinements, and A303932 and A304034 for similar conjectures. The conjecture holds for n up to 2.5*10^12. - _Jud McCranie_, Oct 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013214 | OEIS-A303930

Number of no-leaf subgraphs of the 2 X n grid up to horizontal and vertical reflection.

Conjectures from _Colin Barker_, May 03 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, May 02 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013215 | OEIS-A303932

Number of ways to write 2*n as p + 2^k + 3^m, where p is a prime with 11 a quadratic residue modulo p, and k and m are nonnegative integers.

Conjecture 1. a(n) > 0 for all n > 1, i.e., any even number greater than two can be written as the sum of a power 2, a power of 3 and a prime p with 11 a quadratic residue modulo p. Conjecture 2. For any integer n > 2, we can write 2*n as p + 2^k + 3^m, where p is a prime with 11 a quadratic nonresidue modulo p, and k and m are nonnegative integers. We have verified Conjectures 1 and 2 for n up to 5*10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 02 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013216 | OEIS-A303934

Number of ways to write 2*n as p + 2^k + 5^m with p prime and 2^k + 5^m squarefree, where k and m are nonnegative integers.

Conjecture: a(n) > 0 for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 03 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013217 | OEIS-A303949

Number of ways to write 2*n+1 as p + 2*(2^k+5^m) with p prime and 2^k+5^m a product of at most three distinct primes, where k and m are nonnegative integers.

See also A303934 and A304081 for related conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 05 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013218 | OEIS-A303998

Number of ways to write 2*n+1 as p + 2^k + binomial(2*m,m), where p is a prime, and k and m are positive integers.

Conjecture: a(n) > 0 for all n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013219 | OEIS-A304032

Number of ways to write 2*n as p + 2^k + 3^m with p prime and 2^k + 3^m a product of at most two distinct primes, where k and m are nonnegative integers.

See also A304034 for a related conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013220 | OEIS-A304034

Number of ways to write n as p + 2^k + (1+(n mod 2))*3^m with p prime, where k and m are positive integers with 2^k + (1+(n mod 2))*3^m squarefree.

Conjecture: a(n) > 0 for all n > 11. See also A304081 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 06 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013221 | OEIS-A304081

Number of ways to write n as p + 2^k + (1+(n mod 2))*5^m, where p is an odd prime, and k and m are nonnegative integers with 2^k + (1+(n mod 2))*5^m squarefree.

Conjecture: a(n) > 0 for all n > 7. See also A303821, A303934, A303949, A304031 and A304122 for related information, and A304034 for a similar conjecture. The author would like to offer 2500 US dollars as the prize to the first proof of the conjecture, and 250 US dollars as the prize to the first explicit counterexample. - _Zhi-Wei Sun_, May 08 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 06 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013222 | OEIS-A304122

Squarefree numbers of the form 2^k + 5^m, where k is a positive integer and m is a nonnegative integer.

The conjecture in A304081 has the following equivalent version: Any even number greater than 4 can be written as the sum of a prime and a term of the current sequence, and also any odd number greater than 8 can be written as the sum of a prime and twice a term of the current sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013223 | OEIS-A304174

Decimal expansion of 2^61/(3^32*993), the conjectured maximal residue in the Collatz 3x+1 problem.

Decimal expansion of 2^61/(3^32*993), the conjectured maximal residue in the Collatz 3x+1 problem. The residue of n in the 3x+1 problem is defined as the ratio 2^h(n)/(3^t(n)*n), where h = A006666 is the number of halving steps, and t = A006667 is the number of tripling steps. It is conjectured that n = 993 yields the highest possible residue. See e.g. the Roosendaal page, and A127789 for indices of record residues.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_ (following an idea of _Michel Lagneau_), May 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013224 | OEIS-A304177

Union of sequences b and c defined by: b(1)=8, b(2)=488, b(n)=62*b(n-1) - b(n-2) and c(1)=22, c(2)=10582, c(n)=482*c(n-1) - c(n-2).

Conjecture: Each member of this sequence can be used as an initial value for Inkeri's primality test for Fermat numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pedja Terzic_, May 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013225 | OEIS-A304178

Number of distinct sets of palindrome prefix lengths, over all binary palindromes of length n.

Claesson's conjecture: for n>=3, a(n+1) equals the number of Wilf-classes of Hertzsprung patterns of length n. These patterns correspond to "rigid patterns" from Myers's work. Bóna studied them under the name "very tight patterns". The number of Wilf-classes of Hertzsprung patterns can also be interpreted as the number of different autocorrelation polynomials of self-overlapping permutations. For more details, definitions and examples, see works by Claesson, Bóna, Kirgizov-Nurligareev and Myers in the links section. - _Sergey Kirgizov_, Nov 28 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Jan 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013226 | OEIS-A304239

Indices for which the Mertens function A002321 reaches its extremum between subsequent zeros for the first time.

This is related to the Mertens conjecture, more precisely to record values of Mertens function A002321 in the following sense: Due to the short-scale and long-scale oscillations of A002321, it is less appealing to consider record values in the usual sense (cf. A051402), which yields many slowly growing records and record indices lying closely together, during the approach of a "long-scale" record. Therefore this sequence considers maxima or minima between two subsequent zeros, ignoring the empty intervals between immediately adjacent zeros A002321(k) = A002321(k+1) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, May 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013227 | OEIS-A304275

a(n) = Sum_{k = 1..n} gcd(k,n) / cos(Pi*k/n)^2 for odd n.

a(n) = (1/3)*Sum_{k = 1..4*n-2} (-1)^k*gcd(k,4*n-2)^2. - Conjectured by _Peter Bala_, Dec 26 2023; proved by _Nikolay Osipov_, Oct 05 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, May 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013228 | OEIS-A304312

Logarithmic derivative of F(x) that satisfies: [x^n] exp(n^2 * x) / F(x) = 0 for n>0.

Conjecture: a(n) is the number of connected n-state finite automata with 2 inputs (A006691). [I believe the name of A006691 should be changed to read "(n+1)-state". See my comments in A006691. - _Petros Hadjicostas_, Feb 26 2021]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013229 | OEIS-A304313

Logarithmic derivative of F(x) that satisfies: [x^n] exp(n^3 * x) / F(x) = 0 for n>0.

Conjecture: a(n) is the number of connected n-state finite automata with 3 inputs (A006692).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013230 | OEIS-A304314

Logarithmic derivative of F(x) that satisfies: [x^n] exp(n^4 * x) / F(x) = 0 for n>0.

Conjecture: a(n) is the number of connected n-state finite automata with 4 inputs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013231 | OEIS-A304315

Logarithmic derivative of F(x) that satisfies: [x^n] exp(n^5 * x) / F(x) = 0 for n>0.

Conjecture: a(n) is the number of connected n-state finite automata with 5 inputs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013232 | OEIS-A304321

Table of coefficients in row functions F'(n,x)/F(n,x) such that [x^k] exp(k^n * x) / F(n,x) = 0 for k>=1 and n>=1.

Conjecture: T(n,k) in row n and column k gives the number of connected k-state finite automata with n inputs, for k>=0, for n>=1. For example, row 2 agrees with A006691, the number of connected n-state finite automata with 2 inputs; also, row 3 agrees with A006692, the number of connected n-state finite automata with 3 inputs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013233 | OEIS-A304322

O.g.f. A(x) satisfies: [x^n] exp(n^2 * x) / A(x) = 0 for n>0.

It is conjectured that the coefficients of o.g.f. A(x) consist entirely of integers. Conjecture: given o.g.f. A(x), the coefficient of x^n in A'(x)/A(x) is the number of connected n-state finite automata with 2 inputs (A006691).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013234 | OEIS-A304323

O.g.f. A(x) satisfies: [x^n] exp(n^3 * x) / A(x) = 0 for n>0.

It is conjectured that the coefficients of o.g.f. A(x) consist entirely of integers. Conjecture: given o.g.f. A(x), the coefficient of x^n in A'(x)/A(x) is the number of connected n-state finite automata with 3 inputs (A006692).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013235 | OEIS-A304324

O.g.f. A(x) satisfies: [x^n] exp(n^4 * x) / A(x) = 0 for n>0.

It is conjectured that the coefficients of o.g.f. A(x) consist entirely of integers. Conjecture: given o.g.f. A(x), the coefficient of x^n in A'(x)/A(x) is the number of connected n-state finite automata with 4 inputs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013236 | OEIS-A304325

O.g.f. A(x) satisfies: [x^n] exp(n^5 * x) / A(x) = 0 for n>0.

It is conjectured that the coefficients of o.g.f. A(x) consist entirely of integers. Conjecture: given o.g.f. A(x), the coefficient of x^n in A'(x)/A(x) enumerates the connected n-state finite automata with 5 inputs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013237 | OEIS-A304331

Number of integers k > 1 such that n - F(k) is a positive squarefree number, where F(k) denotes the k-th Fibonacci number A000045(k).

Conjecture: a(n) > 0 for all n > 1. In other words, every n = 2,3,... can be written as the sum of a positive Fibonacci number and a positive squarefree number. See also A304333 for a similar conjecture involving Lucas numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013238 | OEIS-A304333

Number of positive integers k such that n - L(k) is a positive squarefree number, where L(k) denotes the k-th Lucas number A000204(k).

Conjecture: a(n) > 0 for all n > 1. See also A304331 for a similar conjecture involving Fibonacci numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013239 | OEIS-A304362

a(n) = Sum_{d|n, d = 1 or not a perfect power} mu(n/d).

The conjecture that a(n) is always in {-2,-1,0,1,2} is false. In fact, a(30233088000000) = 3. More generally, let N_t = 2^(15+30*t)*3^(10+30*t)*5^(6+30*t), where t >= 0. Then a(N_t) = 3. - _Gerard Gomez_, Jul 23 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, May 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013240 | OEIS-A304451

Solution (a(n)) of the complementary equation in Comments.

Conjectures: a(n)/n -> 5/2 and -1 <= 5/2 - a(n)/n <= 2 for n >= 1;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013241 | OEIS-A304452

Solution (b(n)) of the complementary equation in Comments.

Conjectures: a(n)/n -> 5/2 and -1 <= 5/2 - a(n)/n <= 2 for n >= 1;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013242 | OEIS-A304490

a(1) = a(2) = a(3) = 1, a(4) = 5, a(5) = 6, a(6) = 4; a(n) = a(n-a(n-2)) + a(n-a(n-4)) for n > 6.

Conjectures from _Colin Barker_, May 14 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, May 13 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013243 | OEIS-A304522

Number of ordered ways to write n as the sum of a Fibonacci number and a positive odd squarefree number.

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 2, 27, 83, 31509. This conjecture implies that any integer n > 1 not equal to 83 can be written as the sum of a positive Fibonacci number and a positive odd squarefree number, which has been verified for n up to 10^10. Note that 83 = 0 + 83 = 1 + 2*41, where 0 and 1 are Fibonacci numbers, and 83 and 2*41 are squarefree. The author would like to offer 1000 US dollars as the prize for the first complete solution to his conjecture that any positive integer is the sum of a Fibonacci number and a positive odd squarefree number. See also A304331, A304333 and A304523 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 13 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013244 | OEIS-A304523

Number of ordered ways to write n as the sum of a Lucas number (A000032) and a positive odd squarefree number.

Conjecture: a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 3, 11, 13, 27, 29, 67, 139, 193, 247, 851. See also A304331, A304333 and A304522 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 13 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013245 | OEIS-A304524

Consider the ratio res(p) = 2^A006666(p) / (p*3^A006667(p)) where p is prime. The prime numbers in this sequence are those for which res(p) sets a new record.

Conjecture: for all prime p, res(p) < res(993) = 2^61/(3^32*993) = 1.253142... (see A304174).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, May 14 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013246 | OEIS-A304689

Number of nonnegative integers k such that n - F(k)*F(k+1) is positive and squarefree, where F(k) denotes the k-th Fibonacci number A000045(k).

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 10, 90, 66690, 67452, 22756020.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 17 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013247 | OEIS-A304720

Number of nonnegative integers k such that n - (4^k - k) is positive and squarefree.

Conjecture: a(n) > 0 for all n > 1. See A281192 for N such that none of N - 1 or N + 1 is squarefree: then n = N + 2 is such that n - 1 and n - 3 are not squarefree, i.e., one cannot take k = 0 or k = 1 in the present definition, and k > 1 is required to satisfy the conjecture. - _M. F. Hasler_, May 23 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 17 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013248 | OEIS-A304721

Numbers m with A304720(m) = 1.

Conjecture: The sequence only has 112 terms as listed in the b-file.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 17 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013249 | OEIS-A304752

Beginning with a(1) = 1, for n > 1, a(n) = the least divisor of a(n-1) not included earlier, otherwise a(n) = the least multiple m*a(n-1) such that m is not a divisor of a(n-1) and m*a(n-1) is not included earlier.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, _Michael De Vlieger_, and _Robert G. Wilson v_, May 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013250 | OEIS-A304822

a(n) = A304821(n+3)/A304821(n) - 1.

Conjecture: This sequence consists of 1's and primes only.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pedja Terzic_, May 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013251 | OEIS-A304916

Numbers k such that the number of divisors of the k-th central binomial coefficient is a power of 2.

Conjecture: there are no terms beyond 784.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, May 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013252 | OEIS-A304943

Number of ways to write n as the sum of a positive tribonacci number (A000073) and a positive odd squarefree number.

Conjecture: a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 3, 4, 6, 10, 11, 13, 29, 76, 1332, 25249.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013253 | OEIS-A304945

Number of nonnegative integers k such that n - k*L(k) is positive and squarefree, where L(k) denotes the k-th Lucas number A000032(k).

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 4, 5, 9, 10, 24, 28, 64, 100, 104, 136, 153, 172, 176, 344, 496, 856, 928, 1036, 1084, 1216, 1860.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013254 | OEIS-A305024

Minimal number of squares, not all equal to 1, having as sum prime(n), such that their squares also sum to a prime; 0 if no such decomposition exists.

It has been conjectured (cf. A126769) that any prime p >= 5 can be written in a nontrivial way as p = Sum (b_i)^2 such that Sum (b_i)^4 is also prime. This sequence lists the number of required terms b_i for each prime. The two initial zeros say that this decomposition is not possible for prime(1) = 2 and prime(2) = 3, and are thus conjectured to be the only zeros of the sequence. Since we are interested in the minimal number of terms, we can consider only nonzero b_i >= 1 and min{b_i} >= 2 to avoid the trivial solution b_i = 1 for all i <= k = prime(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, May 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013255 | OEIS-A305058

Totients t such that the number of divisors of t equals the number of solutions of phi(x) = t.

Conjecture: Every term divides one or more subsequent terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Torlach Rush_, May 24 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013256 | OEIS-A305080

Number of connected strict integer partitions of n with pairwise indivisible and squarefree parts.

Conjecture: This sequence is "eventually increasing," meaning that for all k >= 0 there exists an m >= 0 such that a(n) > k for all n > m. For k = 0 it appears we can take m = 18, for example.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, May 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013257 | OEIS-A305123

G.f.: Sum_{k>=1} x^(2*k-1)/(1+x^(2*k-1)) * Product_{k>=1} 1/(1-x^k).

Conjecture: a(n) is odd iff n is a term of A067567. - _Peter Bala_, Jan 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, May 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013258 | OEIS-A305211

a(n) is the number of possible values of (x^3 + y^3) mod n, where x and y are any integers.

Conjecture: keyword mult applies. Furthermore a procedure to find a(n) is as follows: if n = 7k then n -> 5*n/7. if n = 9k then n-> 5*n/9. return(n). - _David A. Corneth_, May 22 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jack Zhang_, May 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013259 | OEIS-A305214

Numbers k such that A305212(k) is not zero.

Conjecture: These are simply the positive integers that are divisible by 7 or 9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jack Zhang_, May 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013260 | OEIS-A305310

Numbers k(n) used for Cassels's Markoff forms MF(n) corresponding to the conjectured unique Markoff triples MT(n) with maximal entry m(n) = A002559(n), for n >= 1.

Numbers k(n) used for Cassels's Markoff forms MF(n) corresponding to the conjectured unique Markoff triples MT(n) with maximal entry m(n) = A002559(n), for n >= 1. Every m(n) is proved to appear as largest member of a Markoff triple MT(n) = (m_1(n), m_2(n), m(n)), with positive integers m_1(n) < m_2(n) < m(n) for n >= 3 (MT(1) = (1, 1, 1) and MT(2) = (1, 1, 2)) satisfying the Markoff equation m_1(n)^2 + m_2(n)^2 + m(n)^2 = 3*m_1(n)*m_2(n)*m(n). The famous Markoff uniqueness conjecture is that m(n) as largest member determines exactly one ordered triple MT(n). See, e.g., the Aigner reference, pp. 38-39, and Corollary 3.5, p. 48. [In numerating the sequence with n related to A002559(n) this conjecture is assumed to be true. - _Wolfdieter Lang_, Jul 29 2018] a(n) = k(n) has been defined in terms of the (conjectured unique) ordered Markoff triple MT(n) = (m_1(n), m_2(n), m(n)) with m(n) = A002559(n) in the comment above as k(n) = min{k1(n), k2(n)}, where m_1(n)*k1(n) - m_2(n) == 0 (mod m(n)), with 0 <= k1(n) < m(n), and m_2(n)*k2(n) - m_1(n) == 0 (mod m(n)), with 0 <= k2(n) < m(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jun 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013261 | OEIS-A305311

Numbers k(n) used for Markoff forms determining quadratic irrationals with purely periodic continued fractions.

The uniqueness conjecture (see A305310, also for the Aigner reference) is here assumed to be true. - _Wolfdieter Lang_, Jul 29 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jun 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013262 | OEIS-A305312

Discriminant a(n) of the indefinite binary quadratic Markoff form m(n)*F_{m(n)}(x, y) with m(n) = A002559(n), for n >= 1.

a(n) is also the discriminant D(n) = a(n) of the indefinite binary quadratic form determining the Markoff triple MT(n) = (x(n), y(n), m(n)) if the largest member is m(n) = A002559(n) and x(n) <= y(n) <= m(n). This is the form x^2 - 3*m*x*y + y^2 = -m^2 (with dropped argument n), or in reduced version X^2 + b*X*Y - b*Y^2 = -m^2, with b = b(n) = 3*m(n) - 2, where X = X(n) = y(n) - x(n) and Y = Y(n) = y(n). The uniqueness of such Markoff triples MT(n) with given largest members m(n) is a conjecture. If the forms for a Markoff triple with largest member m are numerated with n giving m as m(n) = A002559(n)as in the present entry then the uniqueness conjecture is assumed to be true. Otherwise certain m(n) will lead to several different forms. - _Wolfdieter Lang_, Jul 30 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jun 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013263 | OEIS-A305313

Smallest member m_1(n) of the ordered Markoff triple MT(n) with largest member m(n) = A002559(n), n >= 1. These triples are conjectured to be unique.

Smallest member m_1(n) of the ordered Markoff triple MT(n) with largest member m(n) = A002559(n), n >= 1. These triples are conjectured to be unique. The second member m_2 of the Markoff (Markov) triple MT(n) = (m_1(n), m_2(n), m(n)) with m_1(n) <= m_2(n) <= m(n), for n >= 1, with m(n) = A002559(n) is given in A305314(n). For n>=3 the inequalities are strict. The existence of MT(n) with largest number m(n) is proved. The uniqueness is conjectured. The Markoff equation is (the argument n is dropped) m_1^2 + m_2^2 + m^2 = 3*m_1*m_2*m. See the references under A002559. a(n) = m_1(n) is the fundamental proper solution x of the indefinite binary quadratic form x^2 - 3*m(n)*x*y + y^2, of discriminant D(n) = 9*m(n)^2 - 4 = A305312(n), representing -m(n)^2, for n >= 1, with x <= y. The uniqueness conjecture means that there are no other such fundamental solutions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jun 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013264 | OEIS-A305314

Second member m_2(n) of the Markoff triple MT(n) with largest member m(n) = A002559(n), and smallest member m_1(n) = A305313(n), for n >= 1. These triples are conjectured to be unique.

Second member m_2(n) of the Markoff triple MT(n) with largest member m(n) = A002559(n), and smallest member m_1(n) = A305313(n), for n >= 1. These triples are conjectured to be unique. a(n) = m_2(n) is the fundamental proper solution y of the indefinite binary quadratic form x^2 - 3*m(n)*x*y + y^2, of discriminant D(n) = 9*m(n)^2 - 4 = A305312(n), representing -m(n)^2, for n >= 1, with x <= y. The uniqueness conjecture means that there are no other such fundamental solutions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jun 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013265 | OEIS-A305315

a(n) = sqrt(5*b(n)^2 - 4), with b(n) = A134493(n) = Fibonacci(6*n+1), n >= 0.

This shows that ordered Markoff (Markov) triples [1, y, m], with 1 <= y <= m, have for m from the union of sets {m1(k)}_{k>=0} U {m5(k)}_{k>=0) U {m3(k)}_{k>=0)}, with mj(k) = F(6*k+j), for j = 1, 5, and 3, the unique solutions yj(k) = (3*F(6*k+j) - aj(k))/2 < mj(k), namely y1(k) = F(6*k-1) = A134497(k-1) with F(-1) = 1, y5(k) = F(6*k+3) = A134495(k) and y3(k) = F(6*k+1) = A134493. The solutions with the + sign are excluded because they are > mj(k). This trisection of the odd-indexed Fibonacci numbers as m numbers shows again the well known fact that each of them appears as largest member in a Markoff triple if the smallest member is x = 1. The positions of the odd-indexed Fibonacci numbers in the Markoff sequence A002559 are given in A158381. The conjecture in this case is that the odd-indexed Fibonacci numbers appear as largest numbers only in the ordered Markov triples with x = 1. See, e.g., the Aigner reference for the general Frobenius-Markoff conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jul 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013266 | OEIS-A305317

a(n) gives the length of the period of the regular continued fraction of the quadratic irrational of any Markoff form representative Mf(n), n >= 1 (assuming the uniqueness conjecture).

a(n) gives the length of the period of the regular continued fraction of the quadratic irrational of any Markoff form representative Mf(n), n >= 1 (assuming the uniqueness conjecture). The index n enumerates the Markoff triples with largest member m from A002559 in increasing order. If the Markoff-Frobenius uniqueness conjecture (see, e.g. the book of Aigner) is true then the triples can be numbered by n if the largest member is m(n) = A002559(n). In the other (unlikely) case there may be more than one triple (hence forms) for some Markoff numbers m from A002559, and then one orders these triples lexicographically. The indefinite binary quadratic Markoff form Mf(n) = Mf(n;x,y) for the given Markoff number m(n) = A002559(n), n >= 1, (assuming that the mentioned uniqueness conjecture is true) is m(n)*x^2 + (3*m(n) - 2*k(n))*x*y + (l(n) - 3*k(n))y^2 with l(n) = (k(n)^2 +1)/m(n), and k(n) is defined for the representative form (of the unimodualar equvivalence class), e.g., in Cassels as k(n) = k_C(n) = A305310(n). The qudadratic irrational xi(n) is the solution of Mf(n;x,1) = 0 with the positive root. For the representative forms used by Cassels the regular continued fractions for xi(n) = xi_C(n) are not purely periodic. The smallest preperiod is -1 for n = 1 and 0 for n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jul 30 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013267 | OEIS-A305319

Triangle T(n,k) read by rows: coefficients in order of decreasing exponents of characteristic polynomial P_n(t) of the matrix M(i,j) = [(i+j>n) or (i+j)=n-1], 1 <= i,j <= n.

Related to conjecture from entry A047211.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gheorghe Coserea_, May 30 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013268 | OEIS-A305374

First differences of A140101.

This relation between A003144 and A140101 is a conjecture (Daniel Forgues remarks would trivially follow from this relation). - _Michel Dekking_, Mar 18 2019 Conjecture: This sequence is 2.TTW(3,3,2) where TTW is the ternary tribonacci word defined in A080843, or equally it is THETA(3,3,2), where THETA is defined in A275925. - _N. J. A. Sloane_, Mar 19 2019 All these conjectures are now theorems - see the Dekking et al. paper. - _N. J. A. Sloane_, Jul 22 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013269 | OEIS-A305392

First differences of A140100.

Conjecture: (a(n)) = 212121 delta(x). Conjecture: This sequence (prefixed by 1 since A140100 should really begin with 0) is 1.TTW(2,1,1) where TTW is the ternary tribonacci word defined in A080843, or equally it is THETA(2,1,1), where THETA is defined in A275925. - _N. J. A. Sloane_, Mar 19 2019 All these conjectures are now theorems - see the Dekking et al. paper. - _N. J. A. Sloane_, Jul 22 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013270 | OEIS-A305393

First differences of A140102.

Conjecture: (a(n)) = 12 delta(x). Conjecture: This sequence (prefixed by 1 since A140102 should really begin with 0) is 1.TTW(1,2,1) where TTW is the ternary tribonacci word defined in A080843, or equally it is THETA(1,2,1), where THETA is defined in A275925. - _N. J. A. Sloane_, Mar 19 2019 All these conjectures are now theorems - see the Dekking et al. paper. - _N. J. A. Sloane_, Jul 22 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013271 | OEIS-A305394

First differences of A140103.

Conjecture: this sequence is the ternary tribonacci word on the alphabet {5,4,3}, i.e., (a(n)) is the unique fixed point of the morphism 5 -> 54, 4 -> 53, 3 -> 5; see A092782. - _Michel Dekking_, Mar 13 2019 An equivalent conjecture: This sequence (prefixed by 3 since A140103 should really begin with 0) is 3.TTW(5,4,3) where TTW is the ternary tribonacci word defined in A080843, or equally it is THETA(5,4,3), where THETA is defined in A275925. There are similar conjectures for the first differences of A140100, A140101, A140102. - _N. J. A. Sloane_, Mar 14 2019 and Mar 19 2019 All these conjectures are now theorems - see the Dekking et al. paper. - _N. J. A. Sloane_, Jul 22 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013272 | OEIS-A305398

Index of the least prime not dividing p-1, where p = A073918(n) is the smallest prime such that p-1 has n distinct prime factors.

This is related to the conjecture formulated in A073918, that for any m there is K(m) such that prime(m)# | A073918(k)-1 for all k >= K(m): This conjecture is equivalent to lim inf a(n) = oo.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, May 31 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013273 | OEIS-A305400

a(n) = round(1/(A073918(n)/prime(n)# - 1)), where A073918(n) = min { prime p | omega(p-1) = n } and p# = product of primes <= p.

We conjecture that lim inf A073918(n)/A002110(n) = 1 but the value of the lim sup is unknown. Therefore we consider x defined as A073918(n)/A002110(n) = 1 + 1/x, and a(n) = round(x).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, May 31 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013274 | OEIS-A305404

Expansion of Sum_{k>=0} (2*k - 1)!!*x^k/Product_{j=1..k} (1 - j*x).

Conjecture: a(n) = Sum_{k>=0} k^n * binomial(2*k,k) / (2^k * 3^(k + 1/2)). - _Diego Rattaggi_, Oct 11 2020 O.g.f. conjectural: 1/(1 - x/(1 - 3*x/(1 - 3*x/(1 - 6*x/(1 - 5*x/(1 - 9*x/(1 - 7*x/(1 - ... - (2*n-1)*x/(1 - 3*n*x/(1 - ...))))))))) - a continued fraction of Stieltjes-type. - _Peter Bala_, Dec 06 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, May 31 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013275 | OEIS-A305440

Indices k such that A292518(k) = 0.

Conjecture: for k > 3957 there are no more terms in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Jun 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013276 | OEIS-A305441

Indices k such that A305355(k) = 0.

Conjecture: for k > 10316 there are no more terms in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Jun 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013277 | OEIS-A305444

a(n) = Product_{p is odd and prime and divisor of n} (p - 2).

This ratio, multiplied by the twin prime constant, occurs in the asymptotic behavior of prime gaps of size 2*n as decribed by the Hardy-Littlewood asymptotic conjecture for the number of prime pairs. See A005597 for more information. - _Hugo Pfoertner_, Dec 25 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Markus Sigg_, Aug 12 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013278 | OEIS-A305499

Square array A(n,k), n > 0 and k > 0, read by antidiagonals, with initial values A(1,k) = k and recurrence equations A(n+1,k) = A(n,k) for 0 < k <= n and A(n+1,k) = A(n,k) - A000035(n+k) for 0 < n < k.

Conjecture: Sum_{i=1..n} A(n+1-i,i) = A211538(n+3) for n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Jun 03 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013279 | OEIS-A305502

Number of ways to write n as x + y with 0 < x <= y such that x^3 + n*y^2 is prime.

Conjecture: a(n) > 0 for all n > 1, and a(n) = 1 only for n = 2, 3, 4, 5, 6, 7, 8, 10, 11, 12, 16, 20, 22, 28, 42, 50. See also A232174 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 03 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013280 | OEIS-A305503

Largest cardinality of subsets A of {0,1,...,n-1} with |A + A| > |A - A|.

a(n) = n - 7 (conjectured) for all n > 15. Conjectures from _Colin Barker_, Jun 01 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tanuj Mathur_, Jun 03 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013281 | OEIS-A305550

Expansion of e.g.f. Product_{k>=1} (1 + (exp(x) - 1)^k).

Conjecture: Let k be a positive integer. The sequence obtained by reducing a(n) modulo k is eventually periodic with the period dividing phi(k) = A000010(k). For example, modulo 16 we obtain the sequence [1, 1, 3, 3, 7, 3, 15, 11, 7, 3, 15, 11, 7, 3, 15, 11, ...], with an apparent period of 4 beginning at a(4). Cf. A167137. More generally, we conjecture that the same property holds for integer sequences having an e.g.f. of the form G(exp(x) - 1), where G(x) is an integral power series. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Jun 15 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013282 | OEIS-A305562

Coefficients associated with power series solution of g'(x) = g(x/2) e^x at x=0.

Conjecture: a(n) = A386393(n,n) for n > 0. - _Mikhail Kurkov_, Jul 19 2025 [Conjecture is true, see MathOverflow link. - _Mikhail Kurkov_, Apr 01 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Jun 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013283 | OEIS-A305579

Square array read by antidiagonals upwards in which row k has k as its first term and each subsequent term is the least possible value such that the sum of any 2 or more terms does not equal a prime.

Conjecture: these row terms are a proper subset of A005843.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Randy L. Ekl_ and _Robert G. Wilson v_, Jun 05 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013284 | OEIS-A305709

Least k such that there exists a three-term sequence n = b_1 < b_2 < b_3 = k such that b_1 * b_2 * b_3 is square.

Conjecture: a(n) < A072905(n) with finitely many nonsquare exceptions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Jun 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013285 | OEIS-A305756

Coefficients of (q*(j(q)-720))^(1/24) where j(q) is the elliptic modular invariant.

(Conjecture)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Jun 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013286 | OEIS-A305757

Inverse Euler transform of q*(j-720) where j is j-function (A000521).

(Conjecture) Let {b_n} = inverse Euler transform of (q*(j+144*k)). b_n is a multiple of 24.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Jun 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013287 | OEIS-A305827

Beastly primes with beastly prime indices, i.e., primes p such that both the decimal representation of p and the decimal representation of the index of p in A000040 contain the substring "666".

Conjecture 1: The sequence is infinite. An argument in favor of the conjecture goes as follows: Choose any sufficiently large integer i containing the substring "666". Then it seems very likely, assuming the digits of prime(i) are distributed uniformly, that prime(i) contains the substring "666" at least once. Based on that argument, one can make the following stronger conjecture: Conjecture 2: There exists an integer h such that for each i >= h, prime(A051003(i)) is a term of the sequence. This conjecture may be false if some subsequences of the primes are infinite. For example, assume A004022 is infinite. Then it seems likely that there exists a large term of that sequence whose index in A000040 contains the substring "666".

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Jun 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013288 | OEIS-A305875

a(n) is the maximum number m such that 2n = prime(i+k) + prime(j-k), k=1,2,...,m, where i and j are a pair of numbers such that 2n = prime(i+1) + prime(j+1).

Conjecture: there are only 392 n's such that a(n) = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Jun 12 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013289 | OEIS-A305884

Lexicographically first sequence of positive squares, no two or more of which sum to a square.

Conjecture: the only values that appear more than once are 1 and 25.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Jun 13 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013290 | OEIS-A305892

a(n) = 1 if n is composite and the n-th Stern polynomial B(n,x) is irreducible, 0 otherwise.

See also A186891 about related Ulas & Ulas conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jun 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013291 | OEIS-A305924

Irregular table: row n >= 0 lists all k >= 0 such that the decimal representation of 4^k has n digits '0' (conjectured).

Irregular table: row n >= 0 lists all k >= 0 such that the decimal representation of 4^k has n digits '0' (conjectured). It remains an open problem to provide a proof that the rows are complete, in the same way as each of the terms of A020665 is unproved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 14 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013292 | OEIS-A305925

Irregular table read by rows in which row n >= 0 lists all k >= 0 such that the decimal representation of 5^k has n digits '0' (conjectured).

Irregular table read by rows in which row n >= 0 lists all k >= 0 such that the decimal representation of 5^k has n digits '0' (conjectured). It remains an open problem to provide a proof that the rows are complete, in the same way as each of the terms of A020665 is unproved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013293 | OEIS-A305926

Irregular table: row n >= 0 lists all k >= 0 such that the decimal representation of 6^k has n digits '0' (conjectured).

Irregular table: row n >= 0 lists all k >= 0 such that the decimal representation of 6^k has n digits '0' (conjectured). It remains an open problem to provide a proof that the rows are complete, in the same way as each of the terms of A020665 is unproved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013294 | OEIS-A305927

Irregular table: row n >= 0 lists all k >= 0 such that the decimal representation of 7^k has n digits '0' (conjectured).

Irregular table: row n >= 0 lists all k >= 0 such that the decimal representation of 7^k has n digits '0' (conjectured). It remains an open problem to provide a proof that the rows are complete, in the same way as each of the terms of A020665 is unproved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013295 | OEIS-A305928

Irregular table: row n >= 0 lists all k >= 0 such that the decimal representation of 8^k has n digits '0' (conjectured).

Irregular table: row n >= 0 lists all k >= 0 such that the decimal representation of 8^k has n digits '0' (conjectured). It remains an open problem to provide a proof that the rows are complete, in the same way as each of the terms of A020665 is unproved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013296 | OEIS-A305929

Irregular table: row n >= 0 lists all k >= 0 such that the decimal representation of 9^k has n digits '0' (conjectured).

Irregular table: row n >= 0 lists all k >= 0 such that the decimal representation of 9^k has n digits '0' (conjectured). It remains an open problem to provide a proof that the rows are complete, in the same way as each of the terms of A020665 is unproved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013297 | OEIS-A305931

Powers of 3 having at least one digit '0' in their decimal representation.

The complement A238939 is conjectured to have only 23 elements, the largest being 3^68. Thus, all larger powers of 3 are (conjectured to be) in this sequence. Each of the subsequences "powers of 3 with exactly n digits 0" is conjectured to be finite. Provided there is at least one such element for each n >= 0, this leads to a partition of the integers, given in A305933.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 15 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013298 | OEIS-A305932

Irregular table: row n >= 0 lists all k >= 0 such that the decimal representation of 2^k has n digits '0' (conjectured).

Irregular table: row n >= 0 lists all k >= 0 such that the decimal representation of 2^k has n digits '0' (conjectured). Although it remains an open problem to provide a proof that the rows are complete (as are all terms of A020665), we can assume it for the purpose of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 14 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013299 | OEIS-A305933

Irregular table read by rows: row n >= 0 lists all k >= 0 such that the decimal representation of 3^k has n digits '0' (conjectured).

Irregular table read by rows: row n >= 0 lists all k >= 0 such that the decimal representation of 3^k has n digits '0' (conjectured). It remains an open problem to provide a proof that the rows are complete, just as each of the terms of A020665 is unproved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 14 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013300 | OEIS-A305934

Powers of 3 that have exactly one digit '0' in base 10.

The first 6 terms coincide with A305931: powers of 3 having at least one digit 0, with complement A238939 (within A000244: powers of 3) conjectured to be finite, too. Then, a(7..8) = A305931(9..10), etc.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_ (following a suggestion by _Zak Seidov_), Jun 14 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013301 | OEIS-A305938

Number of powers of 8 having exactly n digits '0' (in base 10), conjectured.

Number of powers of 8 having exactly n digits '0' (in base 10), conjectured. These are the row lengths of A305928. It remains an open problem to provide a proof that these rows are complete (as are all terms of A020665), but the search has been pushed to many orders of magnitude beyond the largest known term, and the probability of finding an additional term is vanishing, cf. Khovanova link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013302 | OEIS-A305939

Number of powers of 9 having exactly n digits '0' (in base 10), conjectured.

Number of powers of 9 having exactly n digits '0' (in base 10), conjectured. These are the row lengths of A305929. It remains an open problem to provide a proof that these rows are complete (as for all terms of A020665), but the search has been pushed to many orders of magnitude beyond the largest known term, and the probability of finding an additional term is vanishing, cf. Khovanova link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013303 | OEIS-A305942

Number of powers of 2 having exactly n digits '0' (in base 10), conjectured.

Number of powers of 2 having exactly n digits '0' (in base 10), conjectured. These are the row lengths of A305932. It remains an open problem to provide a proof that these rows are complete (as for all terms of A020665), but the search has been pushed to many orders of magnitude beyond the largest known term, and the probability of finding an additional term is vanishing, cf. Khovanova link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013304 | OEIS-A305943

Number of powers of 3 having exactly n digits '0' (in base 10), conjectured.

Number of powers of 3 having exactly n digits '0' (in base 10), conjectured. These are the row lengths of A305933. It remains an open problem to provide a proof that these rows are complete (as for all terms of A020665), but the search has been pushed to many orders of magnitude beyond the largest known term, and the probability of finding an additional term is vanishingly small, cf. Khovanova link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013305 | OEIS-A305944

Number of powers of 4 having exactly n digits '0' (in base 10), conjectured.

Number of powers of 4 having exactly n digits '0' (in base 10), conjectured. These are the row lengths of A305924. It remains an open problem to provide a proof that these rows are complete (as are all terms of A020665), but the search has been pushed to many orders of magnitude beyond the largest known term, and the probability of finding an additional term is vanishing, cf. Khovanova link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013306 | OEIS-A305945

Number of powers of 5 having exactly n digits '0' (in base 10), conjectured.

Number of powers of 5 having exactly n digits '0' (in base 10), conjectured. These are the row lengths of A305925. It remains an open problem to provide a proof that these rows are complete (as are all terms of A020665), but the search has been pushed to many orders of magnitude beyond the largest known term, and the probability of finding an additional term is vanishing, cf. Khovanova link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013307 | OEIS-A305946

Number of powers of 6 having exactly n digits '0' (in base 10), conjectured.

Number of powers of 6 having exactly n digits '0' (in base 10), conjectured. These are the row lengths of A305926. It remains an open problem to provide a proof that these rows are complete (as for all terms of A020665), but the search has been pushed to many orders of magnitude beyond the largest known term, and the probability of finding an additional term is vanishing, cf. Khovanova link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013308 | OEIS-A305947

Number of powers of 7 having exactly n digits '0' (in base 10), conjectured.

Number of powers of 7 having exactly n digits '0' (in base 10), conjectured. These are the row lengths of A305927. It remains an open problem to provide a proof that these rows are complete (as for all terms of A020665), but the search has been pushed to many orders of magnitude beyond the largest known term, and the probability of finding an additional term is vanishing, cf. Khovanova link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013309 | OEIS-A305992

The sequence whose indicator function is I in conjectured formula A300997(n) = 2*n - Sum_{k=1..n} I(k), as long as the conjecture holds.

The sequence whose indicator function is I in conjectured formula A300997(n) = 2*n - Sum_{k=1..n} I(k), as long as the conjecture holds. A300997(n) is believed to be equal to 2*n - Sum_{k=1..n} I(k), where I is the indicator function of some other sequence -- let it be this sequence. This sequence is finite if the conjecture is false.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luc Rousseau_, Jun 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013310 | OEIS-A306082

Expansion of e.g.f. Product_{k>=1} 1/(1 - (exp(x) - 1)^(k^2)).

Conjecture: for positive integer k, reducing the sequence modulo k produces an eventually periodic sequence with period dividing phi(k) = A000010(k). For example, modulo 7 we obtain the sequence [1, 1, 3, 6, 1, 5, 6, 1, 3, 6, 1, 5, 6, 1, 3, 6, 1, 5, 6, ...], with an apparent period of 6 beginning at a(1). - _Peter Bala_, Feb 22 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jun 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013311 | OEIS-A306112

Largest k such that 2^k has exactly n digits 0 (in base 10), conjectured.

Largest k such that 2^k has exactly n digits 0 (in base 10), conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013312 | OEIS-A306113

Largest k such that 3^k has exactly n digits 0 (in base 10), conjectured.

Largest k such that 3^k has exactly n digits 0 (in base 10), conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013313 | OEIS-A306114

Largest k such that 4^k has exactly n digits 0 (in base 10), conjectured.

Largest k such that 4^k has exactly n digits 0 (in base 10), conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013314 | OEIS-A306115

Largest k such that 5^k has exactly n digits 0 (in base 10), conjectured.

Largest k such that 5^k has exactly n digits 0 (in base 10), conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013315 | OEIS-A306116

Largest k such that 6^k has exactly n digits 0 (in base 10), conjectured.

Largest k such that 6^k has exactly n digits 0 (in base 10), conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013316 | OEIS-A306117

Largest k such that 7^k has exactly n digits 0 (in base 10), conjectured.

Largest k such that 7^k has exactly n digits 0 (in base 10), conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013317 | OEIS-A306118

Largest k such that 8^k has exactly n digits 0 (in base 10), conjectured.

Largest k such that 8^k has exactly n digits 0 (in base 10), conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013318 | OEIS-A306119

Largest k such that 9^k has exactly n digits 0 (in base 10), conjectured.

Largest k such that 9^k has exactly n digits 0 (in base 10), conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013319 | OEIS-A306196

Irregular triangle read by rows where row n lists the primes 2n - k, with 1 < k < 2n-1, and if k is composite also 2n - p has to be prime for some prime divisor p of k.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jan 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013320 | OEIS-A306198

Multiplicative with a(p^e) = p^(e-1)*(p^2 - p - 1).

For any positive integer n and any m coprime to n, define R(n,m) = Product_{primes p divides n} (p - [m == 1 (mod p)]), where [] is an Iverson branket. (Note that Sum_{m=1..n, gcd(m,n)=1} R(n,m) = a(n)). Then we have the following conjecture: (Start) where C is Artin's constant = A005596, PrimePi = A000720. (End) [Conjecture simplified by _Jianing Song_, Sep 13 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Jan 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013321 | OEIS-A306211

Concatenation of the current sequence with the lengths of the runs in the sequence, with a(1) = 1.

Conjecture: All terms are less than or equal to 5. - _Peter Kagey_, Jan 29 2019 Conjecture: Every number appears! (Based on the analogy with the somewhat similar sequence A090822, where the first 5 appeared at around 10^(10^23) steps). - _N. J. A. Sloane_, Jan 29 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _A. D. Skovgaard_, Jan 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013322 | OEIS-A306220

a(n) is the smallest prime p such that Kronecker(-n,p) = -1.

Conjecture: lim sup log(a(n))/log(n) = 0. For example, it seems that log(a(n))/log(n) < 0.5 for all n > 1364.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Jan 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013323 | OEIS-A306224

a(n) is the smallest prime p such that -n is not a square mod p.

Conjecture: lim sup log(a(n))/log(n) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Jan 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013324 | OEIS-A306225

Number of ways to write n as w + x^5 + pen(y) + pen(z), where w is 0 or 1, and x,y,z are integers with x >= w and pen(y) < pen(z), and where pen(m) denotes the pentagonal number m*(3*m-1)/2.

Conjecture: a(n) > 0 for all n > 0. In other words, any positive integer can be written as the sum of two fifth powers of nonnegative integers one of which is 0 or 1, and two distinct generalized pentagonal numbers. We have verified a(n) > 0 for all n = 1..2*10^6. The conjecture implies that the set A = {x^5 + pen(y): x = 0,1,2,... and y is an integer} is an additive basis of order two (i.e., the sumset A + A coincides with {0,1,2,...}). See also A306227 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013325 | OEIS-A306227

Number of ways to write n as w + x^4 + y*(y+1)/2 + z*(z+1)/2, where w is 0 or 1, and x, y, z are nonnegative integers with x >= w and y < z.

Conjecture: a(n) > 0 for all n > 0. In other words, any positive integer can be written as the sum of two fourth powers one of which is 0 or 1, and two distinct triangular numbers. We have verified a(n) > 0 for all n = 1..10^6. The conjecture implies that the set S = {x^4 + y*(y+1)/2: x,y = 0,1,2,...} is an additive basis of order two (i.e., the sumset S + S coincides with {0,1,2,...}). See also A306225 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013326 | OEIS-A306236

a(n) is the smallest integer m > n with integer j > m makes n^2, m^2 and j^2 an arithmetic progression.

Conjecture: if gcd(k, a(k)) = 1, then k is a term in A058529.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jinyuan Wang_, Feb 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013327 | OEIS-A306238

The smallest integer k such that floor(n!/k) is an odd number.

Conjecture: for n > 1, a(n) is a prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jinyuan Wang_, Mar 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013328 | OEIS-A306239

Number of ways to write n as x^3 + y^3 + pen(z) + pen(w), where x, y, z, w are nonnegative integers with x <= y and z <= w, and pen(k) denotes the pentagonal number k*(3*k-1)/2.

Conjecture: a(n) > 0 for all n >= 0, and a(n) = 1 only for n = 0, 4, 5, 16, 20, 46. Also, any nonnegative integer not equal to 16 can be written as x^6 + y^3 + pen(z) + pen(w) with x, y, z, w nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 31 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013329 | OEIS-A306240

Number of ways to write n as x^9 + y^3 + z*(z+1) + w*(w+1), where x,y,z,w are nonnegative integers with x <= 2 and z <= w.

Conjecture: a(n) > 0 for all n >= 0, and a(n) = 1 only for n = 0, 11, 17, 18, 47, 108, 109, 234, 359. Also, any nonnegative integer can be written as x^6 + y^3 + z*(z+1) + w*(w+1), where x,y,z,w are nonnegative integers with x <= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 31 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013330 | OEIS-A306242

Number of ways to write n as x*(3x+1) + y*(3y-1) + z*(3z+2) + w*(3w-2), where x,y,z,w are nonnegative integers.

Conjecture: a(n) > 0 for all n >= 0, and a(n) = 1 only for n = 0, 1, 2, 3, 4, 9, 13. Moreover, any nonnegative integer n can be written as x*(3x+1) + y*(3y-1) + z*(3z+2) + w*(3w-2), where x,y,z,w are nonnegative integers with x or y even. The conjecture has been verified for n up to 10^6. We also have some other similar conjectures. For example, we conjecture that every n = 0,1,2,... can be written as x*(5x+1)/2 + y*(5y-1)/2 + z*(5z+3)/2 + w*(5w-3)/2 with x,y,z,w nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 31 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013331 | OEIS-A306249

Number of ways to write n as x*(2x-1) + y*(3y-1) + z*(4z-1) + w*(5w-1), where x,y,z are nonnegative integers and w is 0 or 1.

Conjecture: a(n) > 0 for any nonnegative integer n. We have some other similar conjectures. For example, we conjecture that each n = 0,1,2,... can be written as x*(3x-1)/2 + y*(5y-1)/2 + z*(7z-1)/2 + w*(9w-1)/2) (or x*(x-1) + y*(2y-1) + z*(3z-1) + w*(4w-1)) with x,y,z,w nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 31 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013332 | OEIS-A306250

Number of ways to write n as x*(3x+1) + y*(3y-1) + z*(3z+2) + w*(3w-2), where x,y,z,w are nonnegative integers with x*y*z = 0.

Conjecture: a(n) > 0 for any nonnegative integer n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013333 | OEIS-A306260

Number of ways to write n as w*(4w+1) + x*(4x-1) + y*(4y-2) + z*(4z-3) with w,x,y,z nonnegative integers.

Conjecture 1: a(n) > 0 for all n >= 0, and a(n) = 1 only for n = 0, 1, 2, 4, 7, 9, 11, 14, 23, 25, 28, 37. Conjecture 2: Each n = 0,1,2,... can be written as w*(4w+2) + x*(4x-1) + y*(4y-2) + z*(4z-3) with w,x,y,z nonnegative integers. Conjecture 3: Each n = 0,1,2,... can be written as 4*w^2 + x*(4x+1) + y*(4y-2) + z*(4z-3) with w,x,y,z nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013334 | OEIS-A306261

Least k > 0 such that 2n - p is prime where p is some prime divisor of 4n^2 - (2k-1)^2 for n >= 4.

Conjecture: a(n) exists for n >= 4. The conjecture holds up to 10^6. Records: a(4) = 1, a(6) = 2, a(34) = 3, a(75) = 4, a(154) = 9, a(1027) = 10, a(1097) = 11, a(1477) = 14, a(1552) = 17, a(5179) = 18, a(10684) = 29, a(70201) = 32, a(79861) = 43, a(519632) = 45, a(1018804) = 46, a(1713031) = 47, ... - _Charles R Greathouse IV_, Feb 17 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Feb 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013335 | OEIS-A306270

Composite numbers k such that b^(k(k-1)) == 1 (mod k^2) for every b coprime to k.

Conjecture: all semiprimes > 4 in this sequence are in A190275. - _Thomas Ordowski_, Jul 19 2020 The conjecture was verified up to 1063290841. - _Amiram Eldar_, Jul 19 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_ and _Thomas Ordowski_, Feb 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013336 | OEIS-A306289

The smallest prime factor of numbers greater than 1 and coprime to 6.

Conjecture: This pattern continues indefinitely. a(n) = A007310(m + 1) when n is congruent to {m, A306277(m + 1)} (mod A091999(m + 1)) but not congruent to {k, A306277(k + 1)} (mod A091999(k + 1)), m > k >= 1. The indices of the first appearance of a number in this sequence supports this conjecture in that they are never, for m > 0, congruent to A306277(m + 1) mod A091999(m + 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Davis Smith_, Feb 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013337 | OEIS-A306302

Number of regions into which a figure made up of a row of n adjacent congruent rectangles is divided upon drawing diagonals of all possible rectangles (a(0)=0 by convention).

a(n) = n + (A114043(n+1) - 1)/2, conjectured by _N. J. A. Sloane_, Feb 07 2019; proved by _Max Alekseyev_, Apr 10 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paarth Jain_, Feb 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013338 | OEIS-A306305

Smallest number m such that 2^m*n has 2 or more identical adjacent decimal digits or -1 if no such m exists.

Conjecture 1: a(n) != -1 for all n > 0. Conjecture 2: there exists K > 0 such that a(n) <= K for all n. Evidence suggests K = 21. The above conjectures are true. In particular, 0 <= a(n) <= 21 for all n > 0. This is proved by showing that for each 0 < n < 1000, there is a number m <= 21 such that 2^m*n mod 1000 has adjacent identical digits. If n > 0 and n == 0 mod 1000, then clearly a(n) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Feb 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013339 | OEIS-A306334

a(n) is the number of different linear hydrocarbon molecules with n carbon atoms.

We prove _Vaclav Kotesovec_'s conjectures from the Formula section. Let M = [[0,0,1], [0,1,1], [1,1,1]], X(n) = M^(n-2), and Y(n) = M^(floor(n/2)-2) = X(floor(n/2)) (with negative powers indicating matrix inverses). Let also, t_1 = [1,1,1]^T, t_2 = [1,2,2]^T, and t_3 = [1,2,3]^T. In addition, define b(n) = (1/2)*(t_1^T X(n) t_1) and c(n) = (1/2)*(t_3^T Y(n) t_1) if n is even and = (1/2)*(t_2^T Y(n) t_1) if n is odd. We have a(n) = b(n) + c(n) for n >= 1. Since the characteristic polynomial of _Vaclav Kotesovec_'s recurrence is x^9 - 2*x^8 - 3*x^7 + 5*x^6 + x^5 + 2*x^3 - 3*x^2 - x + 1 = g(x)*g(x^2), where g(x) = x^3 - 2*x^2 - x + 1, to prove his first conjecture, it suffices to show that b(n) - 2*b(n-1) - b(n-2) + b(n-3) = 0 (whose characteristic polynomial is g(x)) and c(n) - 2*c(n-2) - c(n-4) + c(n-6) = 0 (whose characteristic polynomial is g(x^2)). Since the characteristic polynomial of _Vaclav Kotesovec_'s recurrence is g(x)*g(x^2), which is a polynomial of degree 9, the denominator of the g.f. of the sequence (a(n): n >= 1) should be x^9*g(1/x)*g(1/x^2) = (1 - 2*x - x^2 + x^3)*(1 - 2*x^2 - x^4 + x^6), as _Vaclav Kotesovec_ conjectured below. The numerator of _Vaclav Kotesovec_'s g.f. can be easily derived using the initial conditions (from a(1) = 1 to a(9) = 409). (End) Conjectures from _Vaclav Kotesovec_, Feb 12 2019: (Start) G.f.: (1 - x - 2*x^2 - x^4 + 2*x^5 + x^6 - x^7) / ((1 - 2*x - x^2 + x^3)*(1 - 2*x^2 - x^4 + x^6)) - 1. (End) [These conjectures are true. See my comments above. - _Petros Hadjicostas_, Nov 17 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincent Champain_, Feb 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013340 | OEIS-A306346

Start with the sequence S(1) = [0], and for n >= 1 define S(n+1) to equal the concatenation of S(n) with the RUNS transform of S(n) when read in reverse order. This sequence is the limit of that process as n goes to infinity.

Conjecture 1: a(n) <= 7. Conjecture 2: a(n) is odd for n > 1 and when a(n) = 3, 5 or 7, n is odd. If row 1 = [1], the sequence a(n) becomes b(n) = 1, 1, 2, 1, 2, 1, 1, 1, 2, 1, 3, 1, 1, 1, 2, 1, 3, 1, 1, 1, 3, 1, 1, 1, 2, 1, 3, 1, 3, 1, 1, 1, ... and it is conjectured that b(n) <= 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Feb 09 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013341 | OEIS-A306362

Prime numbers in A317298.

Conjecture: Except the first term a(1) = 3, all the other terms do not end with 3. It is easy to prove that the numbers A317298(n) end with 3 only when n ends with 1. In this case A317298(10*n+1) = (10*n + 1)*(20*n + 3), which is composite for n > 0. Therefore the conjecture is true. - _Bruno Berselli_, Feb 11 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Feb 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013342 | OEIS-A306366

For any sequence s of positive integers without infinitely many consecutive equal terms, let T(s) be the sequence obtained by replacing each run, say of k consecutive t's, with a run of t consecutive k's; this sequence corresponds to T(K) (where K denotes the Kolakoski sequence A000002).

a(n) = A000002(ceiling(2*n/3)) (conjectured). - _Jon Maiga_, Jan 24 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Feb 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013343 | OEIS-A306382

Positive integers not representable as Pen(x) + Pen(y) + 2*Pen(z), where x, y, z are nonnegative integers, and Pen(k) denotes the pentagonal number k*(3k-1)/2.

Conjecture 1: This sequence only has 49 terms as listed. Conjecture 2: Any integer n > 6471 can be written as Pen(x) + Pen(y) + 3*Pen(z) with x,y,z nonnegative integers. Any integer n > 7727 can be written as Pen(x) + 2*Pen(y) + 2*Pen(z) with x,y,z nonnegative integers. Also, any integer n > 4451 can be written as Pen(x) + 2*Pen(y) + 4*Pen(z) with x,y,z nonnegative integers. Conjecture 3: Let N(1) = 5928, N(4) = 2761, N(8) = 8868 and N(11) = 9929. For each c among 1, 4, 8, 11, any integer n > N(c) can be written as x*(3x+1)/2 + y*(3y+1) + c*z*(3z+1)/2 with x,y,z nonnegative integers. Conjecture 4: Any integer n > 5544 can be written as x*(3x+1)/2 + y*(3y+1)/2 + 3z*(3z+1)/2 with x,y,z nonnegative integers. Any integer n > 7093 can be written as x*(3x+1)/2 + 3*y*(3y+1)/2 + 2z*(3z+1) with x,y,z nonnegative integers. Also, any integer n > 8181 can be written as x*(3x+1)/2 + 2y*(3y+1) + 3z*(3z+1) with x,y,z nonnegative integers. Conjecture 5: For each positive integer m, there are only finitely many positive integers not representable as x*(x+1)/2 + y*(y+1)/2 + z*(z+1)/2 with x,y,z in the set {m, m+1, ...}. See also A306383 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013344 | OEIS-A306383

Number of ways to write n as x*(2x+1) + y*(2y+1) + z*(2z+1), where x,y,z are nonnegative integers with x <= y <= z.

Conjecture 1: a(n) > 0 for any integer n > 138158. Conjecture 2: Any integer n > 146858 can be written as the sum of three hexagonal numbers (A000384). Conjecture 3: Any integer n > 33066 can be written as the sum of three pentagonal numbers (A000326). Conjecture 4: Any integer n > 24036 can be written as the sum of three second pentagonal numbers (A005449). Conjecture 5: Let N(1) = 114862, N(-1) = 166897, N(3) = 196987 and N(-3) = 273118. Then, for any r among 1, -1, 3 and -3, each integer n > N(r) can be written as x*(5x+r)/2 + y*(5y+r)/2 + z*(5z+r)/2 with x,y,z nonnegative integers. We have verified Conjectures 2-5 for n up to 10^6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013345 | OEIS-A306384

Start with n and repeatedly double it and apply the "delete any run of identical digits" operation described in A321801; a(n) is the number of steps needed to reach one of 0, 1, or 5, or -1 if none of these three numbers is ever reached.

Conjecture: every number will eventually reach one of 0, 1, or 5. Conjecture is true for n < 10^10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Feb 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013346 | OEIS-A306415

Numbers k such that A179682(k) <> A033996(k).

Conjecture: every member of the sequence is a member of A033996 or is A179682(k) for some k in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Feb 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013347 | OEIS-A306427

Least integer m such that there are exactly n quadruples of distinct divisors (d_i, d_j, d_k, d_l) among the divisors of m having the property d_i * d_j - d_k * d_l = 1, for some i, j, k, l.

Conjecture: for each integer q > 1, there exists a subsequence E(q) of {a(n)} such that q*E(q) is also a subsequence of {a(n)}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Feb 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013348 | OEIS-A306439

Number of ways to write n as x*(3x+1)/2 + y*(3y+1)/2 + z*(3z+1) + 3w*(3w+1)/2, where x,y,z,w are nonnegative integers with x <= y.

Conjecture 1: a(n) > 0 for all n > 5, and a(n) = 1 only for n = 0, 2, 7, 9, 11, 12, 16, 31, 33, 41. Conjecture 2: Let n be any integer greater than 9, and let p(x) denote x*(3x+1)/2. For each c = 2, 4, 9, we can write n as p(x) + 2*p(y) + 3*p(z) + c*p(w) with x,y,z,w nonnegative integers. See also Conjecture 5.2 of the linked 2016 paper.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013349 | OEIS-A306459

Number of ways to write n as w^3 + C(x+2,3) + C(y+2,3) + C(z+2,3), where w,x,y,z are nonnegative integers with x <= y <= z, and C(m,k) denotes the binomial coefficient m!/(k!*(m-k)!).

Conjecture: a(n) > 0 for all n >= 0. In other words, each nonnegative integer can be written as the sum of a nonnegative cube and three tetrahedral numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013350 | OEIS-A306460

Number of ways to write n as x*(2x-1) + y*(y+1)/2 + z*(z+1)*(z+2)/6, where x,y,z are nonnegative integers with x > 0.

Conjecture 1: a(n) > 0 for all n > 0. In other words, any positive integer n can be written as the sum of a positive hexagonal number, a triangular number and a tetrahedral number. Conjecture 2: Let c be 1 or 3. Then each n = 0,1,... can be written as c*x(x+1) + y*(y+1)/2 + z*(z+1)*(z+2)/6 with x,y,z nonnegative integers. Conjecture 3: Let t(x) = x*(x+1)*(x+2)/6. Then each n = 0,1,... can be written as 2*t(w) + t(x) + t(y) + t(z) with w,x,y,z nonnegative integers. We have verified Conjecture 3 for all n = 0..2*10^5. Clearly, Conjecture 3 implies Pollock's conjecture which states that any natural number is the sum of five tetrahedral numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 17 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013351 | OEIS-A306462

Number of ways to write n as C(2w,2) + C(x+2,3) + C(y+3,4) + C(z+4,5), where C(n,k) denotes the binomial coefficient n!/(k!*(n-k)!), w is a positive integer and x,y,z are nonnegative integers.

Conjecture: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 4, 5, 14, 19. See also A306471 and A306477 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 17 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013352 | OEIS-A306471

Number of ways to write n as C(2w+1,2) + C(x+2,3) + C(y+3,4) + C(z+4,5) with w,x,y,z nonnegative integers, where C(n,k) denotes the binomial coefficient n!/(k!*(n-k)!).

Conjecture 1: a(n) > 1 for all n > 0. Conjecture 2: For each r = 0, 1, any positive integer can be written as w^2 + C(x,3) + C(y,4) + C(z,5), where w,x,y,z are nonnegative integers with w - r even. See also A306462 and A306477 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 17 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013353 | OEIS-A306473

a(n) is the maximum number of distinct palindromic not necessarily contiguous subwords over all binary words of length n.

Conjectures from _Colin Barker_, Feb 18 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lars Blomberg_, Feb 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013354 | OEIS-A306516

a(n) is the decimal representation of the binary number with string structure 10s00, s in {0,1}*, such that it results in a non-palindromic cycle of length 4 in the Reverse and Add! procedure in base 2 and is not in the trajectory of any m < a(n).

It is conjectured that all terms in this sequence are derived from S -> S_a -> 10 T_a 00 -> 10 T_a0 00 (in decimal representation here) when we represent the cyclic patterns by the context-free grammar with production rules: It is observed that numbers of the form 81*2^k + 2^floor((k+6)/2) + 2^6*(2^floor((k-8)/2) - 1) + 4 for k >= 11 (i.e. numbers lower(k) as defined in A306514) are all included in this sequence. By assuming this conjecture is true, it is easily to see that the number of seeds for the Reverse and Add! are infinite in base 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _A.H.M. Smeets_, Feb 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013355 | OEIS-A306517

a(n) is the binary number with string structure 10s00, s in {0,1}*, such that it results in a non-palindromic cycle of length 4 in the Reverse and Add! procedure in base 2 and is not in the trajectory of any m < a(n).

It is conjectured that all terms in this sequence are derived from S -> S_a -> 10 T_a 00 -> 10 T_a0 00 when we represent the cyclic patterns by the context-free grammar with production rules: It is observed that all strings 10 T_a0 00, with T_a0 produced by the extended right regular grammar with starting symbol T_a0 and production rules T_a0 -> 1010010 T_a0 | 101, are included in this sequence; i.e. the set of seeds in base 2 is infinite if this conjecture is true. Similar observations can be made for any base b = 2^a, a > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _A.H.M. Smeets_, Feb 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013356 | OEIS-A306521

Least integer N > 2 such that the number of primes (<=N) <= the number of base-n-zero containing numbers (<=N).

a(n) <= A306526(n), equality holds for n = 2, 14, 15, 16, 17, 18, 19, 20, 40, 41, 42, 44, 45, 46, 47, 48, 49, 50, 51, 52 (but a(n) < A306526(n) for all other indices n up to 82). For sufficiently large n, equality holds true for those bases n which satisfy 1/2 <= fract(sqrt(n/log(n)) + O(sqrt(log(n)/n))) < 3/4. This is true for infinitely many further indices, at least for all bases n = ceiling(x), where x is a solution of x/log(x) = k-th triangular number + 1/4, k > 1. For k = 2..10 the corresponding bases are n = 19, 48, 92, 152, 230, 326, 440, 574, 727. Let e(n) be the number of bases m <= n for which a(m) = A306526(m), then lim_{n->infinity} e(n)/n >= 1/4. Conjecture: lim_{n->infinity} e(n)/n = 1/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hieronymus Fischer_, Mar 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013357 | OEIS-A306526

a(n) = greatest integer N such that (number of primes <= N) >= (number of numbers <= N that contain a zero in base n).

a(n) >= A306521(n), equality holds for n = 2, 14, 15, 16, 17, 18, 19, 20, 40, 41, 42, 44, 45, 46, 47, 48, 49, 50, 51, 52 (but a(n) > A306521(n) for all other indices n up to 82). For sufficiently large n, equality holds true for those bases n which satisfy 1/2 <= fract(sqrt(n/log(n)) + O(sqrt(log(n)/n))) < 3/4. This is true for infinitely many indices, at least for all bases n = ceiling(x), where x is a solution of x/log(x) = k-th triangular number + 1/4, k > 1. For k = 2..10 the corresponding bases are n = 19, 48, 92, 152, 230, 326, 440, 574, 727. Let e(n) be the number of bases m <= n for which a(m) = A306521(m), then lim_{n->infinity} e(n)/n >= 1/4. Conjecture: lim_{n->infinity} e(n)/n = 1/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hieronymus Fischer_, Mar 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013358 | OEIS-A306540

Least k >= 1 such that the decimal expansion of n^k starts with 99 or 100.

Conjecture: a(n) <= 231.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Feb 22 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013359 | OEIS-A306559

A self-describing sequence mostly made of 1's and 0's emerging when written in English words (see the Comments section for an explanation).

Jean-Marc Falcoz has computed 100000 terms and found no place where the sequence enters into a loop. It is conjectured that the sequence will grow infinitely without ever entering into such a loop. The proportions of ONEs and ZEROs are conjectured to be 2/3 and 1/3. _Alan C. Wechsler_ (Feb 23 2019) observes that if the first four terms are deleted, this appears to coincide with A039982. Indeed, the resulting pair of sequences agree for at least the first 28442 terms, so this is a very plausible conjecture. - _N. J. A. Sloane_, Feb 23 2019 After deleting the first three terms, this sequence is 0*W(0)*W^2(0)*W^3(0)*... where W is the morphism 0 -> 11, 1 -> 01. It can be shown that this is equivalent to the above conjecture (see link for proof). - _Charlie Neder_, Mar 04 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Feb 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013360 | OEIS-A306562

a(n) = 1 + 2 - 3 - 4 + 5 + 6 + 7 - 8 - 9 - 10 - 11 + 12 + 13 + 14 + 15 + ... + (+-1)*n, where, after the 1st summand there is one plus, two minuses, three plusses, etc.

F(n) = ((-1)^(n+1)(2n+1)(2n^2+2n+5)+21)/16 gives local extrema 3, -4, 14, -24, 46, -71, 111, -157, ... (conjectured). - _Jean-François Alcover_, Jun 01 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brandon J. Butierres_, Feb 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013361 | OEIS-A306570

Values of n such that 5^n ends in n, or expomorphic numbers relative to "base" 5.

Conjecture: if k(n) is expomorphic relative to "base" b, then, the next one in the sequence, k(n+1), consists of the last n+1 digits of b^k(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Feb 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013362 | OEIS-A306571

Number of ways to write n as w*(w+1) + C(x+3,4) + C(y+5,6) + C(z+7,8) with w,x,y,z nonnegative integers, where C(m,k) denotes the binomial coefficient m!/(k!*(m-k)!).

Conjecture: a(n) > 0 for all n >= 0. In other words, each n = 0,1,2,... can be written as 2*C(w,2) + C(x,4) + C(y,6) + C(z,8) with w,x,y,z positive integers. See also A306477 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013363 | OEIS-A306589

a(n) is the smallest positive integer that can be written as the sum or difference of exactly n but no fewer powers of 2 and 3.

Conjecture: The next terms are a(5) = 2581 and a(6) = 27438. - _Eric Rowland_, Jul 22 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Feb 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013364 | OEIS-A306592

Number of ways to write n as u^2 + 2*v^2 + 3*x^2 + 4*y^2 + 5*z^2, where u,v,x,y,z are generalized pentagonal numbers (A001318).

Conjecture: a(n) > 0 for any nonnegative integer n. I'd like to call this the 1-2-3-4-5 conjecture. I have verified it for all n = 0..4*10^5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013365 | OEIS-A306606

Number of ways to write n as w^2 + T(x)^2 + Pen(y)^2 + 2*Pen(z)^2, where w,x,y,z are integers with w > 0 and x >= 0, and T(m) = m*(m+1)/2 and Pen(m) = m*(3m-1)/2.

Conjecture: a(n) > 0 for all n > 0. In other words, any positive integer n can be written as s^2 + t^2 + u^2 + 2*v^2, where s is a positive integer, t is a triangular number, u and v are generalized pentagonal numbers. See also A306614 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013366 | OEIS-A306614

Number of ways to write n as w^2 + (x*(x+1)/2)^2 + (y*(3y+1)/2)^2 + (z*(5z+1)/2)^2, where w,x,y,z are integers with w > 0 and x >= 0.

Conjecture 1 (Big 1-3-5 Conjecture): a(n) > 0 for any positive integer n not divisible by 8. This is stronger than Lagrange's four-square theorem, but different from the 1-3-5 conjecture in A271518. I'd like to call it "the Big 1-3-5 Conjecture". I have verified a(n) > 0 for any positive integer n < 3*10^6 not divisible by 8. Conjecture 2: If c is 5 or 7, then any positive integer n not divisible by 8 can be written as w^2 + (x*(x+1)/2)^2 + (y(3*y+1)/2)^2 + (z*(c*z+3)/2)^2, where w,x,y,z are integers with w > 0. Conjecture 3: If a positive integer n is neither 76 nor divisible by 8, then it can be written as w^2 + (x*(3x+1)/2)^2 + (y*(5y+1)/2)^2 + (z*(5z+3)/2)^2, where w,x,y,z are integers with w > 0. Conjecture 4: Each n = 0,1,2,... can be written as 2*w^2 + (x*(3x+1)/2)^2 + (y*(5y+1)/2)^2 + (z*(5z+3)/2)^2 with w,x,y,z integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013367 | OEIS-A306615

Least positive k such that 2n - p is prime where p is some prime divisor of n^k - 1, or 0 if no such k exists.

Conjecture: a(n) >= 1 for n >= 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Feb 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013368 | OEIS-A306631

Inverse of the Hardy-Ramanujan asymptotic partition function.

Conjecture: a(A000041(n)) = n for all n > 9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Mar 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013369 | OEIS-A306653

a(n) = Sum_{m=1..n} Sum_{k=1..n} [k divides n]*[n/k divides m]*A008683(n/k)*n/k*[k divides m + 2^p]*A008683(k)*k, where p can be a positive integer: 1,2,3,4,5,...

a(n) = A000005(n)*(A008683(n) + Sum_{j=1..n} [4*A056911(j)=n]*A008836(n)*2/3) (conjectured formula verified for n=1..2500). The conjectured Dirichlet generating function simplifies to: Sum_{n>=1} a(n)/n^s = (Product_{j>=1} (1 - 2*prime(j)^(-s)))*(1 + 2^(1 - s)/(2^s - 2)). - _Steven Foster Clark_, Sep 23 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Mar 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013370 | OEIS-A306657

Least primary Carmichael number (A324316) with n prime factors, or -1 if no such number exists.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernd C. Kellner_ and _Jonathan Sondow_, Mar 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013371 | OEIS-A306658

Decimal expansion of the constant S_1 = Sum_{j>=1} prime(2*j - 1)!/prime(2*j)!.

Derived from an open problem published in 2012, the constant S_1 is connected to the gap between the j-th and (j+1)-th primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Mar 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013372 | OEIS-A306683

Integers k for which the base-phi representation of k does not include 1 or phi.

Conjecture (Moses 2012/Baruchel 2018): b is the generalized Beatty sequence b(n) = floor(n*phi) + 2*n + 1. Conjecture (Kimberling 2012): c = A054770 = A000201(n) + 2*n - 1 = floor(n*phi) + 2*n - 1. One can prove that the Moses/Baruchel conjecture and the Kimberling conjecture are equivalent. Conjecture: (a(n)) is a union of two generalized Beatty sequences v and w, given by v(n) = floor(n*phi) + 2*n = A003231(n), and w(n) = 3*floor(n*phi) + n + 1 = A190509(n) + 1. This conjecture is compatible with the Moses/Baruchel/Kimberling conjecture. These three conjectures are proved in my paper 'Base phi representations and golden mean beta-expansions'. - _Michel Dekking_, Jun 26 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Dekking_, May 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013373 | OEIS-A306761

Let digsum(k) = A007953(k) denote the digital sum of k. The sequence lists the smallest integer k such that digsum(k) = digsum (k/d(1)) = digsum (k/d(2)) = ... = digsum (k/d(n)) where d(i) are the n distinct prime factors of k.

Conjecture 1: a(n) == 0 (mod 54) for n > 1. Conjecture 2: a(n)/27 is a squarefree number for n > 1. The smallest multiple of 54 that can be a(10) is 1069808930190, which is also a multiple of 7^2, so the two conjectures above cannot be both true. - _Giovanni Resta_, Mar 08 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Mar 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013374 | OEIS-A306790

Number of ways to write n as x*(3*x+1)/2 + y*(y+1)*(y+2)/2 + z*(z+1)*(z+2)/6, where x is a nonzero integer, and y and z are nonnegative integers.

Conjecture: a(n) > 0 for all n > 0. In other words, each positive integer n can be written as the sum of a positive generalized pentagonal number, a tetrahedral number and a tetrahedral number times three.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013375 | OEIS-A306809

Binomial transform of the continued fraction expansion of e.

Conjectures from _Colin Barker_, Mar 12 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sarah Arpin_, Mar 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013376 | OEIS-A306810

Inverse binomial transform of the continued fraction expansion of e.

Conjectures from _Colin Barker_, Mar 12 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sarah Arpin_, Mar 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013377 | OEIS-A306821

Inverse binomial transform of the "original" Bernoulli numbers [A164555(n)/A027642(n)] with 1 and 1/2 swapped. Denominators.

a(2n) same as denominators of cosecant numbers A001897 for n>0 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Jun 04 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013378 | OEIS-A306825

Primitive part of A001353(n).

Conjecturally there must be infinitely many primes p such that a(p) is prime, but no such p is known. [By the formula below, there is no such p. - _Jianing Song_, Oct 31 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Mar 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013379 | OEIS-A306826

a(0) = 1; a(n) is the smallest integer k > a(n-1) such that 2^(k-1) == 1 (mod a(n-1)*k).

Conjecture: a(n) is prime for every n > 0, namely a(n) is the smallest odd prime p > a(n-1) such that 2^(p-1) == 1 (mod a(n-1)), with a(0) = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Mar 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013380 | OEIS-A306828

Carmichael numbers whose index is a power of 2 (see A174590).

By Lehmer's totient conjecture, there are no composite numbers m such that Od(phi(m)) = Od(m-1), where phi(m) = A000010(m) and Od(m) as above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Mar 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013381 | OEIS-A306861

The concatenation kpk is the number obtained by placing k leading and trailing 1's around the prime p; a(n) is the smallest k such that kpk is prime, where p = prime(n), or -1 if no such k exists.

Conjecture: There are infinitely many -1 terms in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Mar 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013382 | OEIS-A306921

Number of ways of breaking the binary expansion of n into consecutive blocks with no leading zeros.

Conjecture: n appears A067824(n) times for n > 1. Proof: Each partition of n is uniquely determined by a partition of k (call it K) and a choice of some number, from 0 to e, of trailing digits to append to the final part in K, since any remaining digits must appear as singletons. The conjecture follows, since each ordered factorization of a number m produces two numbers n such that a(n) = m, one of each parity, and A067824(n) = 2*A074206(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Mar 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013383 | OEIS-A306948

Expansion of e.g.f. (1 + x)*log(1 + x)*exp(x).

Conjecture: D-finite with recurrence a(n) +(n-5)*a(n-1) +(-3*n+10)*a(n-2) +3*(n-3)*a(n-3) +(-n+3)*a(n-4)=0. - _R. J. Mathar_, Aug 20 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Mar 17 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013384 | OEIS-A306967

a(n) is the first Zagreb index of the Fibonacci cube Gamma(n).

Conjectures from _Colin Barker_, Mar 28 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Mar 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013385 | OEIS-A306971

The number of initial values <= 10^n whose trajectory under the iteration x -> A306938(x) reaches 1.

te Riele noted that of the first initial values <= 10^5 only 459 have a trajectory that reaches 1 and conjectured that almost all trajectories tend to infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Mar 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013386 | OEIS-A306978

Numbers with record number of iterations of x -> A306938(x) required to reach 1 (A306944).

Conjecture: sqrt(3) < a(n)/a(n-1) <= 3. - _Daniel Suteu_, Mar 20 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Mar 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013387 | OEIS-A306995

Number of primes q < prime(n), such that 2*prime(n) + q is prime.

Conjecture: a(n) > 0 for all n >= 3. - _Dmitry Kamenetsky_, Mar 18 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Mar 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013388 | OEIS-A306996

Number of primes q < prime(n), such that 4*prime(n) + q is prime.

Conjecture: a(n) > 0 for all n >= 3. - _Dmitry Kamenetsky_, Mar 18 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Mar 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013389 | OEIS-A306997

Number of primes q < prime(n), such that 6*prime(n) + q is prime.

Conjecture: a(n) > 0 for all n >= 4. - _Dmitry Kamenetsky_, Mar 18 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Mar 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013390 | OEIS-A307087

a(n) is the number of steps it takes for the sequence f(0)=f(1)=n, f(x)=(a*b) mod (a+b+1), where a=f(x-1) and b=f(x-2), to reach a cycle.

Conjecture: for every n, a(n) is finite (that is, the sequence ends up in a cycle).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Costea_, Mar 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013391 | OEIS-A307091

a(n) = Sum_{k=0..floor(n/2)} (-1)^k * binomial(n,2*k)^2.

Conjecture: the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(2*r)) holds for positive integers n and r and all primes p >= 3. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Mar 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013392 | OEIS-A307105

Number of rational numbers which can be constructed from the set of integers between 1 and n, through a combination of multiplication and division.

Conjecture: Let p <= n be prime. If m and p^a*m are two such rationals, then so is p^k*m for all 0 < k < a. - _Yan Sheng Ang_, Feb 13 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Barsotti_, Jul 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013393 | OEIS-A307107

a(n) = A025487(n)/A247451(n).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Mar 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013394 | OEIS-A307110

Index of matching grid point in the bijection between two infinite quadratic grids with one grid rotated by Pi/4 around the common point (0,0), using an enumeration of the grid points by A305575 and A305576.

It is conjectured that the least possible maximum distance within pairs can be reduced to sqrt(5)*sin(Pi/8)=0.855706... (A386241), but not further, and that this can be achieved by "local repairs" of the result of the strip bijection, i.e. by reassigning the connections in groups of 4 pairs, one of which being the pair with d>0.8557... and 3 pairs in the vicinity of the violating pair, but potentially addressing points in neighbor strips. The conjecture is supported by extensive numerical results, but an announced proof by Klaus Nagel remained unpublished.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, following a proposal by _Rainer Rosenthal_, Mar 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013395 | OEIS-A307181

a(n) is the first Zagreb index of the Lucas cube Lambda(n).

Conjectures from _Colin Barker_, Mar 28 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Mar 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013396 | OEIS-A307208

a(n) is the forgotten index of the Fibonacci cube Gamma(n).

Conjectures from _Colin Barker_, Mar 29 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Mar 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013397 | OEIS-A307213

Divide the natural numbers into sets of successive sizes 3,4,5,6,7,...,, starting with {1,2,3}. Cycle through each set until you reach a prime; if the prime was the n-th element in its set, jump to the n-th element of the next set.

If a set does not contain a prime, the sequence goes into an infinite loop, but it is conjectured that this does not happen since the sets are of increasing length.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christopher Cormier_, Mar 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013398 | OEIS-A307244

a(0) = 1; a(n) is the smallest integer k > a(n-1) such that 3^(k-1) == 1 (mod a(n-1)*k).

Conjecture: a(n) is prime for every n > 0, namely a(n) is the smallest prime p > a(n-1) different from 3 such that 3^(p-1) == 1 (mod a(n-1)), with a(0) = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Mar 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013399 | OEIS-A307303

Triangle T(n, k) read as upwards antidiagonals of array A, where A(n, k) is the number of families (also called classes) of proper solutions of the Pell equation x^2 - D(n)*y^2 = -k, for k >= 1.

The position list for nonzero entries in row n = 1 is A057126 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Apr 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013400 | OEIS-A307307

a(n) is the forgotten index of the Lucas cube Lambda(n).

Conjectures from _Colin Barker_, Apr 02 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emeric Deutsch_, Apr 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013401 | OEIS-A307316

Number of unlabeled leafless loopless multigraphs with n edges.

Conjecturally, the asymptotic number of completely symmetric polynomials of degree n up to momentum conservation in the limit as the number of particles increases.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick T. Komiske_, Apr 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013402 | OEIS-A307347

Number of 2n-move closed antelope paths on an unbounded chessboard from a given square to the same square.

Conjecture: a(n) ~ 64^n / (25*Pi*n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Apr 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013403 | OEIS-A307368

a(n) is the minimal positive integer such that 2*a(n)*prime(n)-1 equals another prime.

A more general form of Rassias's conjecture states that for every positive integer a there are two primes p and q such that 2*a*p = q+1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ivan N. Ianakiev_, Apr 17 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013404 | OEIS-A307403

Base-5 based twisted permutation of the nonnegative integers - variant "Hs".

Conjecture: As in A220952, it can be proved (a) that every positive integer is adjacent to exactly two nonnegative integers, and (b) that with this definition of adjacency, the nonnegative integers form a path starting with 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Georg Fischer_, Apr 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013405 | OEIS-A307406

Number of twisted permutations of the nonnegative integers in base 2*n+1 with an adjacency diagram as defined by Knuth in A220952.

Conjecture: a(4) = 673.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Georg Fischer_, Apr 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013406 | OEIS-A307407

Irregular table read by rows: rows list terms that map to the nodes in the graph of the "3x+1" (or Collatz) problem.

Conjecture: The sequence is a permutation of the positive numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Georg Fischer_, Apr 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013407 | OEIS-A307446

First coordinate of a contiguous path on the odd-numbered points of a square lattice leading to a distance-limited bijection with the even-numbered points of a square lattice rotated by Pi/6 around the common point (0,0). A307447 gives the second coordinate.

In a private communication in 2009, _Klaus Nagel_ described a conjectured bijection between the grid points of two quadratic lattices G and H such that the second lattice H has an offset of 1/2 in both coordinate directions and is rotated by an angle of Pi/6 against the first lattice G. If all straight-line connections between points of G and H not exceeding the Euclidean distance d=sqrt(1/2) are drawn (see Nagel link), several types of connected components in the infinite bipartite graph of the vertices of G U H containing equal numbers of grid points from G and H arise.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Apr 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013408 | OEIS-A307486

a(0) = 3; a(n) = smallest k > 1 such that 1 + a(0)*a(1)*...*a(n-1)*k is composite.

Conjecture: for any initial term a(0) > 0, a(n) > 3 only for finitely many n >= 0. Carl Pomerance (in a letter to the author) wrote: Since the sequence a(0)a(1)...a(n-1) grows geometrically, the chance that any given k will give a prime is about 1/n, so the chance that both k = 2 and k = 3 will give primes is about 1/n^2, heuristically, which has a convergent sum. Thus, for this reason I would agree it's reasonable to conjecture that for any starting a(0), the sequence will eventually be made up of 2's and 3's, with 2's predominating. The number of 3's among the first n terms should be proportional to log n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Apr 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013409 | OEIS-A307502

Self-convolution of the Dedekind psi function (A001615).

Conjecture: Sum_{k=1..n} a(k) ~ 75 * n^4 / (8 * Pi^4). - _Vaclav Kotesovec_, Aug 20 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Apr 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013410 | OEIS-A307512

a(n) is the smallest k > 2^n such that 2^(k-1) == 1 (mod (2^n-1)*k).

Conjecture: a(n) is composite if and only if n = 2^j and 2^(2^j) + 1 is composite (presumably for all j > 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Apr 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013411 | OEIS-A307532

a(n) is the smallest k > 2^(2^n)+1 such that 2^(k-1) == 1 (mod (2^(2^n)-1)*k).

Conjecture: a(n) is prime for all n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_ and _Thomas Ordowski_, Apr 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013412 | OEIS-A307535

a(n) is the smallest k >= 0 such that 2^(2^n) + k*2^n + 1 is prime.

Conjecture: 2^(2^n) + a(n)*2^n + 1 = A307532(n) for all n > 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_ and _Thomas Ordowski_, Apr 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013413 | OEIS-A307542

Even integers k such that there exists a prime p with p = min{q: q prime and (k - q) prime} and (k - p) < p^2.

This sequence is an extension of A244408. It is equivalent to "Even numbers 2n such that the smallest prime p satisfying p+q=2n (p, q prime, p<=q) also satisfies p^2+p>2n." If p satisfies additionally p^2 < 2n the corresponding even numbers do not belong to A244408. These numbers are 10, 28, 54, 124, 368, 968, 3526. It is conjectured that a(81)=63274 is the last term. There are no more terms below 4*10^18.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Corinna Regina Böger_, Apr 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013414 | OEIS-A307554

Longest path length in the n X n fiveleaper graph.

Conjectures from _Colin Barker_, Jun 14 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Apr 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013415 | OEIS-A307618

A Calabi-Yau period integral: a(n) = C(4*n,2*n)*C(2*n,n)^3.

We conjecture that for each integer k, the sequence defined by f_k(n) = [x^n] F(x)^(k*n) satisfies the above supercongruences. We conjecture that for each integer k, the sequence defined by e_k(n) = [x^n] E(x)^(k*n) also satisfies the above supercongruences. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bradley Klee_, Jun 04 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013416 | OEIS-A307627

Primes p such that 2 is a primitive root modulo p while 8 is not.

According to Artin's conjecture, the number of terms <= N is roughly ((2/5)*C)*PrimePi(N), where C is Artin's constant = A005596, PrimePi = A000720. Compare: the number of terms of A001122 that are no greater than N is roughly C*PrimePi(N).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Apr 19 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013417 | OEIS-A307628

Primes p such that 2 is a primitive root modulo p while 32 is not.

By Artin's conjecture, the number of terms <= N is roughly ((4/19)*C)*PrimePi(N), where C is Artin's constant = A005596, PrimePi = A000720. Compare: the number of terms of A001122 that are no greater than N is roughly C*PrimePi(N).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Apr 19 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013418 | OEIS-A307663

a(n) = (n-1)!*(Sum_{i=1..n} Sum_{j=1..i} binomial(i,j)*i/j).

Conjectures from _Robert Israel_, Oct 26 2020: (Start) The conjecture regarding the e.g.f. is true. See links. - _Sela Fried_, Jul 30 2024.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pedro Caceres_, Apr 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013419 | OEIS-A307684

Total volume of all rectangular prisms with dimensions r X s X t where r, s and t are the smallest, middle and largest parts in each partition of n into 3 parts.

Conjectures from _Colin Barker_, Apr 22 2019: (Start) These conjectures are true. See links. - _Sela Fried_, Aug 06 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Apr 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013420 | OEIS-A307686

Sum of the smallest side lengths of all integer-sided triangles with perimeter n.

Conjectures from _Colin Barker_, May 12 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013421 | OEIS-A307706

Number of unitary divisors of n that are smaller than sqrt((sqrt(2) - 1)*n).

For k >= 2, the earliest occurrence of k is at n = A132404(k)/2 if A132404(k) is even (and thus being a multiple of 4). Conjecture: this is always the case.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Apr 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013422 | OEIS-A307734

Smallest k such that the adjusted frequency depth of k! is n, and 0 if there is no such k.

Conjecture: this sequence has infinitely many nonzero terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Apr 25 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013423 | OEIS-A307740

Numbers k such that k divides lcm(tau(k), sigma(k)).

Conjecture: multiply-perfect numbers (A007691) are terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Apr 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013424 | OEIS-A307743

a(n) = Sum_{k=1..n} A307742(k).

Conjecture: There is a constant c such that abs(a(n) - 2*n*(c+1)/c) = O(sqrt(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _I. V. Serov_, Apr 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013425 | OEIS-A307744

A fractal function, related to ruler functions and the Cantor set. a(1) = 0; for m >= 0, a(3m) = 1; for m >= 1, a(3m-1) = a(m-1) + sign(a(m-1)), a(3m+1) = a(m+1) + sign(a(m+1)).

Take a Sierpinski arrowhead curve formed of unit edges numbered consecutively from 0 at its axis of symmetry and aligned with an infinite Sierpinski gasket so that each edge is contained in the boundary of either the plane sector occupied by the gasket or a triangular region of the gasket's complement. If a(n) = 0, the n-th edge is contained in the sector boundary, otherwise the relevant triangular region seems to have side 2^(a(n)-1). See A307672 for a fuller description. The conjectured formulas below (that use A094373) derive from summing areas of regions within the gasket. - Corrected by _Peter Munn_, Aug 09 2019 Sum_{n=-3^k..3^k-1} A094373(a(n)) = 3 * 4^k (conjectured). Sum_{n=-3m..3m-1} A094373(a(n)) = 4 * Sum_{n=-m..m-1} A094373(a(n)) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Munn_, Apr 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013426 | OEIS-A307764

Numbers m whose distinct prime factors are exactly the same as the distinct prime factors of each of the numbers obtained by deleting any single digit in the decimal expansion of m.

Conjecture: a(3n-2) = 25*10^(n+1), a(3n-1) = 36*10^(n+1) and a(3n) = 96*10^(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jul 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013427 | OEIS-A307792

Triangular numbers t(n) such that n / (product of decimal digits of t(n)) is an integer.

The name implies that t(n) must be zeroless. I conjecture 1 and 15 are the only two terms with the property n = product of decimal digits of t(n). Are all terms bigger than 1 divisible by 3?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Apr 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013428 | OEIS-A307804

Triangle T(n,k) read by rows: number of labeled 2-regular digraphs (multiple arcs and loops allowed) on n nodes with k components.

Conjectures from _Mikhail Kurkov_, Mar 22 2025: (Start) Note that e.g.f. is exp(t*F(x)) where F(x) = Sum_{n>=1} b(n)*x^n/n! and the same conjectures seems to work if we change b(n) to any sequence with b(1) = 1. (End) Conjectures are true, see MathOverflow links. - _Mikhail Kurkov_, Apr 17 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Apr 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013429 | OEIS-A307812

Triangular numbers t(n) with a zeroless decimal representation such that (product of decimal digits of t(n)) / n is an integer.

Are all terms of the sequence bigger than 1 divisible by 3? I conjecture that 1 and 15 are the only terms for which (product of decimal digits of t(n)) = n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Apr 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013430 | OEIS-A307813

a(n) = (5/32)*4^n - floor((n^2 + 1)/2)*2^(n - 2).

This was formerly conjectured (by Erdos and Guy, 1973) to be the crossing number of the n-hypercube graph for n >= 3; cr(Q_n) = 0 for n = 0, 1, 2, 3. Embeddings of Q_n with cr(Q_n) = a(n) are known for n = 5 and 6. However, the Erdos-Guy conjecture is now refuted since cr(Q_7) <= 1744 < a(7) = 1760 (Clancy et al. 2019).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Apr 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013431 | OEIS-A307818

Numbers k such that A240718(k) = 0.

Conjecture: 31637 is the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Apr 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013432 | OEIS-A307864

a(n) is the number of natural bases b < 2n+1 such that b^n == 1 (mod 2n+1).

Conjecture: if 2n+1 is an absolute Euler pseudoprime, then a(n) = phi(2n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, May 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013433 | OEIS-A307872

Sum of the smallest parts in the partitions of n into 3 parts.

Conjectures from _Colin Barker_, May 02 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013434 | OEIS-A307873

The concatenation pkp is the number obtained by placing prime p either side of R_k, the k-th repunit (1, k times); a(n) is the smallest k such that pkp is prime, where p=prime(n), or -1 if no such k exists.

Conjecture: There are an infinite number of -1 terms in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, May 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013435 | OEIS-A307882

Records of the offset k in A307881, divided by 3.

The divisibility of k by 3 of all terms after a(1) is conjectural. Offsets not divisible by 3 are given as -k instead of k/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, May 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013436 | OEIS-A307933

Number of (undirected) Hamiltonian paths in the n-antiprism graph.

Conjectures from _Colin Barker_, Mar 29 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, May 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013437 | OEIS-A307966

Sum of the largest side lengths of all integer-sided triangles with perimeter n.

Conjectures from _Colin Barker_, May 12 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013438 | OEIS-A307974

Inverse binomial transform of the "original" Bernoulli numbers [A164555(n)/A027642(n)] with 1 and 1/2 swapped. Numerators.

Denominators 3, 15, 21, 15, 33, 1365, 3, 255, ... coincide with cosecant numbers A001897, except 1 (conjectured). a(2*n+1) = n+1 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, May 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013439 | OEIS-A307977

Expansion of g.f.: f'(t)/f(t), where f(t) = Sum_{p prime} t^p.

Conjecture 1: lim_{n->infinity} a(n)/a(n+1) = -0.629233... = gamma (A078756). Conjecture 2: -0.629233... is a root of f(t), where f(t) = Sum_{p prime} t^p. Conjecture 3: From conjecture 2 it follows that, -0.629233... is also a root of H(t) = Sum_{n>=1} pi(n)*t^n, where pi = A000720 is the prime counting function. This follows since H(t) = f(t)/(1-t).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Orges Leka_, May 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013440 | OEIS-A307981

Number of ways to write n as x^3 + 2*y^3 + 3*z^3 + w*(w+1)*(w+2)/6, where x,y,z,w are nonnegative integers.

Conjecture: a(n) > 0 for every nonnegative integer n. In other words, we have {x^3 + 2*y^3 + 3*z^3 + w*(w+1)*(w+2)/6: x,y,z,w = 0,1,2,...} = {0,1,2,...}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013441 | OEIS-A307983

Consider a pentagonal bipyramid with its seven vertices P_1, ..., P_7 lying on a unit sphere; sequence gives decimal expansion of Sum_{i < j} 1/dist(P_i, P_j).

Conjecturally, this is the solution to the Thompson problem with 7 electrons: given 7 points on the surface of a unit sphere, minimize the sum of the inverse distances between pairs of points. (This models electrostatic potential energy in the plum pudding model of the atom. N = 7 electrons is the first unresolved case.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, May 09 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013442 | OEIS-A307985

Number of integer-sided triangles with perimeter n and sides a, b, and c such that a <= b <= c and b|n.

Conjectures from _Colin Barker_, May 15 2019: (Start) (End) [The conjectures are correct. - _Jianing Song_, Oct 23 2022]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013443 | OEIS-A308009

The terms of A081145 appear to lie on three lines; set a(n) = 0 if A081145(n) is on the lower line, 1 if it is on the middle line, and 2 if it is on the upper line.

Certainly the first 10^6 terms of A081145 lie on three roughly straight lines, of slopes roughly 0.56, 1.40, 2.24. See the graph of A081145 and the data file of _Allan Wilks_ attached to that entry. It is a strong conjecture that all terms

of A081145 lie on three roughly straight lines.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013444 | OEIS-A308016

The terms of A308007 appear to lie on two lines; set a(n) = 0 if A308007(n) is on the lower line (defined by A308009(n) <= n), a(n) = 1 if it is on the upper line (defined by a(n) >= n).

It is only a (strong) conjecture that the terms of A308007 line on two lines. Certainly this is true for the first million terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013445 | OEIS-A308021

Start at n=1. Fill in a(n) with a value d > 0 not used earlier such that n-d or n+d is the smallest possible index not visited earlier, then continue with that index.

Conjectured to be a permutation of the positive integers. The conjectured inverse permutation is given in A308049. - _M. F. Hasler_, May 10 2019 The conjecture that {a(n)} is a permutation of the positive integers is equivalent to Slater and Velez's conjecture (see references) that the absolute values of the first differences of A081145 are also a permutation of the positive integers. This problem appears to be still unsolved. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, May 09 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013446 | OEIS-A308034

Number of partitions of n into 3 parts with at least 1 part that is a nondivisor of n.

Conjectures from _Colin Barker_, May 11 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013447 | OEIS-A308047

Sum of subgroup indices of dihedral group, Sum_{H <= D(n)} [D(n):H].

Relation to Lagarias inequality: Let H(G) be the harmonic numbers of D(n) relative to a generating set S <= D(n). Then the conjecture (Lagarias inequality) is:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Orges Leka_, May 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013448 | OEIS-A308049

Inverse permutation to A308021 (Recamán variant).

More precisely, A308021 is by definition injective but presently only conjectured to be a permutation. Until that property is proved, we define this sequence as a(n) = index of the n-th smallest term of A308021.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, May 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013449 | OEIS-A308066

Number of triangles with perimeter n whose side lengths are even.

Conjectures from _Colin Barker_, May 11 2019: (Start) The above conjectures are correct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013450 | OEIS-A308067

Number of integer-sided triangles with perimeter n whose longest side length is odd.

Conjectures from _Colin Barker_, May 11 2019: (Start) Conjecture: a(4*n) = a(4*n+1) = a(4*n+6) = a(4*n-5) = A001840(n-1). - _Marc Bofill Janer_, May 15 2019 The above conjectures are all correct. - _Andrew Howroyd_, Nov 06 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013451 | OEIS-A308068

Number of integer-sided triangles with perimeter n whose longest side length is even.

Conjectures from _Colin Barker_, May 11 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013452 | OEIS-A308090

a(n) = gcd(2^n + n!, 3^n + n!, n+1).

Conjecture: Conversely, if gcd(2^n + n!, 3^n + n!, n+1) = n+1, then n+1 is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pedro Caceres_, May 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013453 | OEIS-A308092

The sum of the first n terms of the sequence is the concatenation of the first n bits of the sequence read as binary, with a(1) = 1.

Conjecture: The number of 1's in the binary representation of each term is weakly increasing, i.e., A000120(a(n)) >= A000120(a(n-1)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, May 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013454 | OEIS-A308096

Take all the integer-sided triangles with perimeter n and sides a, b, and c such that a <= b <= c. a(n) is the sum of all the b's.

Conjectures from _Colin Barker_, May 17 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013455 | OEIS-A308102

Sum of the perimeters of all integer-sided scalene triangles with perimeter n.

Conjectures from _Colin Barker_, May 13 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013456 | OEIS-A308107

Sum of the smallest side lengths of all integer-sided scalene triangles with perimeter n.

Conjectures from _Colin Barker_, May 13 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013457 | OEIS-A308108

Sum of the largest side lengths of all integer-sided scalene triangles with perimeter n.

Conjectures from _Colin Barker_, May 13 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013458 | OEIS-A308121

Irregular triangle read by rows: T(n,k) = A109395(n)*k-A076512(n)*A038566(n,k).

Conjecture: If n is odd, then row 2n can be obtained from row n by interchanging the first and second halves. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jamie Morken_, May 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013459 | OEIS-A308158

Sum of the smallest side lengths of all integer-sided isosceles triangles with perimeter n.

Conjectures from _Colin Barker_, May 15 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013460 | OEIS-A308159

Sum of the largest sides of all integer-sided isosceles triangles with perimeter n.

Conjectures from _Colin Barker_, May 15 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013461 | OEIS-A308160

Take all the integer-sided isosceles triangles with perimeter n and sides a, b, and c such that a <= b <= c. a(n) is the sum of all the b's.

Conjectures from _Colin Barker_, May 15 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013462 | OEIS-A308180

Main diagonal of triangle defined in A308178.

Conjecture: There is a unique fixed point at n=0, and the positive integers form one "cycle", namely (... 59 27 15 9 12 13 8 4 1 3 2 5 6 7 11 14 17 23 30 16 22 19 29 24 38 ...).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013463 | OEIS-A308187

Fixed point (beginning with a) of the morphism a -> aab, b -> b, over the alphabet {a,b} = {0,1}.

Conjecture: characteristic function of A055938. - _Sean A. Irvine_, Jul 16 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013464 | OEIS-A308190

Number of steps to reach 5 when iterating x -> A111234(x) starting at x=n.

It is easy to show that every number n >= 5 eventually reaches 5. This was conjectured by Ali Sada. For A111234 sends a composite n > 5 to a smaller number, and sends a prime > 5 to a smaller number in two steps. Furthermore no number >= 5 can reach a number less than 5. So all numbers >= 5 eventually reach 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013465 | OEIS-A308195

a(n) = smallest m such that A308194(m) = n, or -1 if no such m exists.

Conjecture: all terms, except for a(2), are either primes (A000040) or squarefree semiprimes (A006881). - _Chai Wah Wu_, Jun 18 2019 Conjecture: a(n+1) = (a(n)-prevprime(a(n)))*prevprime(a(n)) infinitely often.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013466 | OEIS-A308215

a(n) is the multiplicative inverse of A001844(n+1) modulo A001844(n); where A001844 is the sequence of centered square numbers.

Conjectures from _Colin Barker_, May 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Hoyt_, May 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013467 | OEIS-A308217

a(n) is the multiplicative inverse of A001844(n) modulo A001844(n+1); where A001844 is the sequence of centered square numbers.

Conjectures from _Colin Barker_, May 16 2019: (Start) Colin Barker's conjectures easily follow. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Hoyt_, May 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013468 | OEIS-A308222

Numbers that are the perimeter of a primitive Heronian isosceles triangle.

a(n) = 2*A096033(n+2) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, May 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013469 | OEIS-A308235

Take apart the sides of each of the integer-sided scalene triangles with perimeter n (at their vertices) and rearrange them orthogonally in 3-space so that their endpoints coincide at a single point. a(n) is the total surface area of all rectangular prisms enclosed in this way.

Conjectures from _Colin Barker_, May 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013470 | OEIS-A308236

Take apart the sides of each of the integer-sided triangles with perimeter n (at their vertices) and rearrange them orthogonally in 3-space so that their endpoints coincide at a single point. a(n) is the total surface area of all rectangular prisms enclosed in this way.

Conjectures from _Colin Barker_, May 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013471 | OEIS-A308247

a(n) is the least integer not the difference of two prime(n)-smooth numbers.

The next few terms are conjectured to be 158857, 681179, 2516509, 10772123, 51292187, 186323681; if they were not, they would provide examples of ABC-triples with quality greater than 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Esteban Crespi de Valldaura_, May 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013472 | OEIS-A308265

Sum of the largest parts in the partitions of n into 3 parts.

Conjectures from _Colin Barker_, Jul 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, May 17 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013473 | OEIS-A308324

Numbers which can be written in the form m^k - m with m an odd prime and k a positive integer.

Besides the trivial example a(1)=0, the only known term which has two representations is a(24) = 2184 = 3^7 - 3 = 13^3 - 13. It is conjectured by Bennett to be the only term with this property.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Craig J. Beisel_, May 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013474 | OEIS-A308342

Number of ways to write 2*n as phi(x^2) + phi(y^2) + phi(z^2), where x,y,z are positive integers with x <= y <= z, and phi(.) is Euler's totient function (A000010).

Conjecture 1: a(n) > 0 for all n > 1. In other words, the set {phi(x^2) + phi(y^2) + phi(z^2): x,y,z = 1,2,3,...} contains all even numbers greater than two. Conjecture 2: For any integer n > 3, we can write 2*n+1 as phi(x^2) + phi(y^2) + sigma(z^2) with x,y,z positive integers, where the function sigma(.) is given by A000203.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013475 | OEIS-A308355

Limiting row sequence of the array A128628.

Conjecture: The length of maximal initial segment of this sequence that is identical to row n of A128628 is A025065(n+1), for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013476 | OEIS-A308359

Triangle T(n,w) read by rows: the number of fixed polyominoes with n cells and width w of the convex hull.

Conjecture: T(n,n-2) = 8*n^2 - 51*n + 86 for n >= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, May 22 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013477 | OEIS-A308376

Least positive integer k < prime(n) such that k + k' is a square, where k' is the unique integer with 0 < k < prime(n) and k*k' == 1 (mod prime(n)), or 0 if such a number k does not exist.

Conjecture 1: a(n) = 0 only for n = 1, 3, 4, 5, 14, 22. In other words, if p is a prime not among 2, 5, 7, 11, 43, 79, then there are k and m among 1,...,p-1 with k*m == 1 (mod p) such that k + m is a square. Conjecture 2: For any prime p > 18, there are k and m among 1,...,p-1 with k*m == 1 (mod p) such that k + m is a triangular number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 22 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013478 | OEIS-A308394

Numbers which can be written in the form m^k - m with m prime and k a positive integer.

The only known terms which have two representations where m is prime are 6 and 2184. It is conjectured by Bennett these are the only terms with this property.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Craig J. Beisel_, May 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013479 | OEIS-A308403

Number of ways to write n as 6^i + 3^j + A008347(k), where i, j and k > 0 are nonnegative integers.

Conjecture 1: a(n) > 0 for all n > 2. In other words, each n = 3,4,... can be written as 6^i + 3^j + prime(k) - prime(k-1) + ... + (-1)^(k-1)*prime(1), where i, j and k > 0 are nonnegative integers. Conjecture 2: If {a,b} is among {2,m} (m = 3..14), {3,4}, {3,5}, then any integer n > 2 can be written as a^i + b^j + A008347(k) with i, j and k > 0

nonnegative integers. Using Qing-Hu Hou's program, we have verified Conjectures 1 and 2 for n up to 10^9 and 10^7 respectively. - _Zhi-Wei Sun_, May 28 2019 Conjecture 1 verified up to 10^10. Conjecture 2 holds up to 10^10 for all cases except {2, 12} since 4551086841 cannot be written as 2^i + 12^j + A008347(k). - _Giovanni Resta_, May 28 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 25 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013480 | OEIS-A308411

Number of ways to write n as 2^i*3^j + A008347(k), where i, j and k > 0 are nonnegative integers.

Conjecture: a(n) > 0 for all n > 1. Conjecture verified for n up to 10^10. - _Giovanni Resta_, May 28 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 25 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013481 | OEIS-A308416

Values of m for which 2*p + m cannot be a square when p is a prime.

Conjecture: for k > 0 and 1 <= j <= k, a(2k^2-2j+1) = 4k^2+4k-4j-3, a(2k^2-2j+2) = 4k^2+4k-4j, a(2k^2+2k-2j+1) = 4k^2+8k-4j, a(2k^2+2k-2j+2) = 4k^2+8k-4j+1. - _Jinyuan Wang_, Jul 23 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bob Andriesse_, May 25 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013482 | OEIS-A308471

Lowest outliers for A057660.

Motivated by _Daniel Forgues_'s conjecture that this sequence consists of 4 and A051451.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charlie Neder_, May 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013483 | OEIS-A308482

Composites c such that T_{c-1} == (c/3)*3^(c-1) (mod c), where T_i denotes the i-th central trinomial coefficient (A002426) and (/) denotes the Kronecker symbol.

Composites satisfying a weaker version of an analog to a congruence satisfied by all primes > 3 (cf. Cao, Sun, 2015, Theorem 1.1 (i); cf. Sun, 2011, Remark to Conjecture A69).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, May 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013484 | OEIS-A308566

Number of ways to write n as w^2 + x*(x+1) + 4^y*5^z with w,x,y,z nonnegative integers.

Conjecture: a(n) > 0 for all n > 0. Equivalently, each n = 1,2,3,... can be written as a*(a+1)/2 + b*(b+1)/2 + 4^c*5^d with a,b,c,d nonnegative integers. See also A308584 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013485 | OEIS-A308584

Number of ways to write n as a*(a+1)/2 + b*(b+1)/2 + 5^c*8^d, where a,b,c,d are nonnegative integers with a <= b.

Conjecture: a(n) > 0 for all n > 0. Equivalently, each n = 1,2,3,... can be written as w^2 + x*(x+1) + 5^y*8^z with w,x,y,z nonnegative integers. See also A308566 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013486 | OEIS-A308597

Number of ways to write n as a*(a+1)/2 + b*(b+1)/2 + 2^c*5^d, where a,b,c,d are nonnegative integers with a <= b and d <= 1.

Conjecture: a(n) > 0 for all n > 0. Moreover, any integer n > 1 can be written as a*(a+1)/2 + b*(b+1)/2 + 2^c*5^d, where a,b,c,d are nonnegative integers with c > 0 and d < 2. Since 2^(k+1) = 2^k + 2^k and 5 = 2^2 + 2^0. the above conjecture implies the conjecture in A303233. On my request, _Giovanni Resta_ found that a(n) = 0 for n = 1217712376, 4371119377. Thus the conjecture fails. - _Zhi-Wei Sun_, Jun 10 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 09 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013487 | OEIS-A308605

Number of distinct subset sums of the subsets of the set of divisors of n. Here 0 (as the sum of an empty subset) is included in the count.

Conjecture: When the terms are sorted and the duplicates deleted a supersequence of A030058 is obtained. Note that A030058 is a result of the same operation applied to A030057.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ivan N. Ianakiev_, Jun 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013488 | OEIS-A308617

Integers i such that the equation A088387(i) = p has N > 1 solutions in the interval prevprime(i)..nextprime(i).

Conjecture: N = 2. Checked up to 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _I. V. Serov_ & _Michel Marcus_, Jun 25 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013489 | OEIS-A308620

Number of states in the evolutionary spatial prisoner's dilemma with n players.

Conjecture: satisfies a linear recurrence having signature (2, 0, -2, 1, 2, -2, 0, 0, 0, -1). - _Harvey P. Dale_, Aug 21 2021 Conjecture: g.f. 2*x -x^2*(-2 +2*x +x^2 -2*x^3 +3*x^5 +x^7 +x^9)/ (x^5+x^2-1)/ (x^5-x^2+2*x-1) .

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jun 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013490 | OEIS-A308621

Number of ways to write n as a*(a+1)/2 + b*(b+1)/2 + 2^c*5^(3d) with a,b,c,d nonnegative integers.

Conjecture: a(n) > 0 for all n > 0. Equivalently, any positive integer n can be written as a^2 + b*(b+1) + 2^c*5^(3d) with a,b,c,d nonnegative integers. This was motivated by A308584, and we verified a(n) > 0 for all n = 1..2*10^8. Then, on the author's request, _Giovanni Resta_ verified the above conjecture for n up to 10^10. G. Resta also noted that 7272774228 cannot be written as a*(a+1)/2 + b*(b+1)/2 + 2^c*250^d with a,b,c,d nonnegative integers. See also A308566, A308584 and A308623 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013491 | OEIS-A308623

Number of ways to write n as a*(a+1)/2 + b*(b+1)/2 + 2^c*10^(2d) with a,b,c,d nonnegative integers.

Conjecture: a(n) > 0 for all n > 0. Equivalently, any positive integer n can be written as a*(a+1)/2 + b*(b+1)/2 + 2^c*10^(2d) with a,b,c,d nonnegative integers. This was motivated by A308566, and we verified a(n) > 0 for all n = 1..2*10^8. Then, on the author's request, _Giovanni Resta_ verified the above conjecture for n up to 10^10. G. Resta also noted that 729546026 cannot be written as a*(a+1)/2 + b*(b+1)/2 + 2^c*3^d with a,b,c,d nonnegative integers. See also A308566, A308594 and A308621 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013492 | OEIS-A308629

Lengths between nodes on a square spiral, such that the n-th node is closer to the (n-2)-th node than to the (n-1)-th node.

Based on an empirical observation of the first several terms, I conjecture that lim_{n->infinity} a(n)/n = 8/9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Barsotti_, Jun 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013493 | OEIS-A308632

Largest aggressor for the maximum number of peaceable coexisting queens as given in A250000.

Conjecture: a(n) - A250000(n) <= 2 for all n. - _Dmitry Kamenetsky_, Oct 14 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roy van Rijn_, Jun 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013494 | OEIS-A308640

Number of ways to write n as (2^a*3^b)^2 + c*(2*c+1) + d*(3*d+1)/2, where a,b,c are nonnegative integers and d is an integer.

Conjecture 1: a(n) > 0 for all n > 0. Conjecture 2: Let k be 1 or 2. Then, any positive integer n can be written as (2^a*3^b)^2 + k*c^2 + d*(3*d+1)/2, where a,b,c are nonnegative integers and d is an integer. Conjecture 3: Let k be 1 or -1. Then, any positive integer n can be written as (2^a*3^b)^2 + c*(5*c+3*k)/2 + d*(3*d+1)/2, where a,b,c are nonnegative integers and d is an integer. We have verified Conjectures 1-3 for all n = 1..10^6. See also A308641 for similar conjectures. Conjecture 1 is false: a(1764630) = 0. Verified by exhaustive enumeration of 1764630 = (2^a*3^b)^2 + c*(2*c+1) + d*(3*d+1)/2, with a,b,c nonnegative integers and d an integer. A bitset scan found this as the only missing positive integer up to 20000000. - _Scott Moore_, May 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013495 | OEIS-A308641

Number of ways to write n as (2^a*5^b)^2 + c*(3*c+1)/2 + d*(3*d+1)/2, where a and b are nonnegative integers, and c and d are integers with c*(3*c+1)/2 <= d*(3*d+1)/2.

Conjecture 1: a(n) > 0 for all n > 0. Conjecture 2: If f(x) is one of the polynomials x^2, x*(5*x+1), x*(5*x+1)/2, x*(7*x+1)/2, x*(7*x+5)/2, then any positive integers n can be written as (2^a*5^b)^2 + f(c) + d*(3*d+1)/2, where a and b are nonnegative integers, and c and d are integers. Conjecture 3: Any positive integers n can be written as (2^a*7^b)^2 + c*(7*c+5)/2 + d*(3*d+1)/2, where a and b are nonnegative integers, and c and d are integers. See also A308640 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013496 | OEIS-A308644

Number of ways to write n as (3^a*5^b)^2 + c*(3c+1)/2 + d*(7d+1)/2, where a and b are nonnegative integers, and c and d are integers.

Conjecture 1: a(n) > 0 for all n > 0. Also, any positive integer n can be written as (3^a*5^b)^2 + c*(3c+1)/2 + d*(7d+3)/2, where a and b are nonnegative integers, and c and d are integers. Conjecture 2: Let r be 1 or 3. Then, any positive integer n can be written as (3^a*4^b)^2 + c*(3c+1)/2 + d*(7d+r)/2, where a and b are nonnegative integers, and c and d are integers. We have verified Conjectures 1-2 for all n = 1..10^6. See also A308640, A308641 and A308656 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013497 | OEIS-A308656

Number of ways to write n as (2^a*9^b)^2 + c*(2c+1) + d*(3d+1), where a and b are nonnegative integers, and c and d are integers.

Conjecture 1: a(n) > 0 for all n > 0. Conjecture 2: If f(x) is one of the polynomials x*(4x+1), x*(5x+2), x*(5x+4), x*(7x+3)/2 and x(7x+5)/2, then any positive integer n can be written as (2^a*9^b)^2 + f(c) + d*(3d+1)/2, where a and b are nonnegative integers, and c and d are integers. Conjecture 3: Let r be 1 or 2. Then any positive integer n can be written as (2^a*7^b)^2 + c*(2c+1) + d*(3d+r), where a and b are nonnegative integers, and c and d are integers. Conjecture 4: If g(x) is one of the polynomials x*(x+1), x*(4x+3), x*(7x+1)/2, x*(7x+3)/2 and x*(7x+5)/2, then any positive integer n can be written as (2^a*7^b)^2 + g(c) + d*(3d+1)/2, where a and b are nonnegative integers, and c and d are integers. We have verified a(n) > 0 for all n = 1..10^8, and Conjectures 2-4 for all n = 1..10^6. See also A308640, A308641, and A308644 for similar conjectures. Jiao-Min Lin (a student at Nanjing University) has found a counterexample to Conjecture 1: a(2109982225) = 0. -

Zhi-Wei Sun, Jul 30 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013498 | OEIS-A308658

Square array read by downward antidiagonals: A(n, k) is the number of primes between the n-th and (n+k)-th perfect powers with exponent > 1, k > 0.

The Redmond-Sun conjecture implies that A(n, 1) is 0 for only finitely many values of n and A(n, k) > 0 for all n and k when k > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Nov 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013499 | OEIS-A308661

Number of ways to write 12*n+5 as (2^a*5^b)^2 + c^2 + d^2, where a,b,c,d are nonnegative integers with a > 0 and c <= d.

Conjecture 1: a(n) > 0 for all n = 0,1,2,.... Equivalently, for each nonnegative integer n we can write 3*n+1 as a*(a+1)/2 + b*(b+1)/2 + (2^c*5^d)^2 with a,b,c,d nonnegative integers. Conjecture 2: For each n = 0,1,2,... we can write 24*n+10 as a^2 + b^2 + (2^c*3^d)^2 with a,b,c,d nonnegative integers and d > 0. We have verified Conjectures 1 and 2 for n up to 2*10^8 and 10^8 respectively. Conjecture 1 holds for n < 8.33*10^9. - _Giovanni Resta_, Jun 19 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013500 | OEIS-A308662

Number of ways to write n as (2^a*5^b)^2 + c*(3c+1) + d*(3d+2), where a and b are nonnegative integers, and c and d are integers.

Conjecture 1: a(n) > 0 for all n > 0. Conjecture 2: Let r be 1 or 2. Then, any positive integer n can be written as (2^a*5^b)^2 + c*(2c+1) + d*(3d+r), where a and b are nonnegative integers, and c and d are integers. We have verified Conjectures 1 and 2 for all n = 1..10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013501 | OEIS-A308669

a(1) = 1, a(2) = 3; for n > 1, a(n) = sum of the next two smallest integers > a(n-1) which are coprime to the sum s = a(1) + ... + a(n-1).

Conjecture 1: a(n) ~ 2^(n+1) * 1.0136719224974731811... Conjecture 2: There are infinitely many n >= 1 such that the ratio (a(n+1) - A131357(n+1))/(a(n) - A131357(n)) equals exactly 2 (for example, for n = 10, 11, 12, 13, 18, 19, 20, 21, 24, 26, etc.).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, Jun 25 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013502 | OEIS-A308691

Numbers k in A320601 such that the fraction of the number of zeros in the decimal expansion of 2^k reaches a record minimum.

Conjecture: there are no more terms beyond 229.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Feb 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013503 | OEIS-A308707

a(n) = gcd(n, phi(n) + sigma(n)), where phi is A000010 and sigma is A000203.

Conjecture: ALL m are primes of the form i^2 + 1 (see A002496); (iv) conjecture: k = m^2 with m >= 1: A000290 \ {0}; if m is prime, it's not a conjecture, see (i). This conjecture is stronger than the conjecture of the 1st comment. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jun 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013504 | OEIS-A308722

Number of edges in the smallest possible regular graceful graph of valence n.

Pegg conjectured the continuation (from n = 13): 130, 147, 180, 192, 221, 243, 285. According to D. E. Knuth (personal communication, Oct. 5, 2026), Shisheng Li proved that no 16-regular graceful graph with 192 edges exists, ruling out the proposed value a(16) = 192. Knuth reports that the existence cases (e,k) = (200,16) and (243,18), where e is the number of edges and k the valence, remain open. - _Eric W. Weisstein_, Oct 06 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Allan C. Wechsler_, Jun 19 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013505 | OEIS-A308726

The number of permutations of length n and tier at most 1, that is, the number of permutations of length n sortable by two passes through a stack where outputting the longest prefix matching the identity permutation is prioritized.

Conjecture: D-finite with recurrence: 3*n*(n-1)*(n+1)*a(n) -n*(n-1)*(67*n-101)*a(n-1) +2*(n-1)*(286*n^2-1112*n+1089)*a(n-2) +4*(-580*n^3+4200*n^2-10106*n+8049)*a(n-3) +24*(184*n^3-1784*n^2+5770*n-6221)*a(n-4) -96*(4*n-15)*(2*n-9)*(4*n-17)*a(n-5)=0. - _R. J. Mathar_, Jan 27 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rebecca Smith_, Jun 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013506 | OEIS-A308727

Lexicographically earliest sequence of distinct terms such that the digits of two contiguous terms sum up to a square.

It is conjectured that this sequence is a permutation of the integers > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-Marc Falcoz_ and _Eric Angelini_, Jun 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013507 | OEIS-A308728

Lexicographically earliest sequence of distinct terms such that the digits of two contiguous terms sum up to a prime.

It is conjectured that this sequence is a permutation of the integers > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Jun 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013508 | OEIS-A308733

Sum of the smallest parts of the partitions of n into 4 parts.

Conjectures from _Colin Barker_, Jun 23 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Jun 22 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013509 | OEIS-A308750

Number of Dumont permutations of the first kind of length 2n avoiding pattern 2143 (or pattern 3421).

Conjecture: The number of Dumont permutations of the first kind avoiding pattern 2143 equals the number of Dumont permutations of the first kind avoiding pattern 3421 for all n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Burstein_ and _Opel Jones_, Jun 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013510 | OEIS-A308758

Sum of the third largest parts of the partitions of n into 4 parts.

Conjectures from _Colin Barker_, Jun 23 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Jun 22 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013511 | OEIS-A308759

Sum of the second largest parts of the partitions of n into 4 parts.

Conjectures from _Colin Barker_, Jun 23 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Jun 22 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013512 | OEIS-A308760

Sum of the largest parts of the partitions of n into 4 parts.

Conjectures from _Colin Barker_, Jun 23 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Jun 22 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013513 | OEIS-A308766

Numbers k such that the minimal mark in a length k sparse ruler is round(sqrt(9 + 12*k)/2) + 1.

Conjecture: Minimal marks k - round(sqrt(9 + 12*k)/2) is always 0 or 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ed Pegg Jr_, Jun 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013514 | OEIS-A308775

Sum of all the parts in the partitions of n into 4 parts.

Conjectures from _Colin Barker_, Jun 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Jun 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013515 | OEIS-A308807

a(n) = 4*5^(n-1) + n.

Conjectures confirmed by _Robert Israel_, Jun 28 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clive Tooth_, Jun 25 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013516 | OEIS-A308815

a(n) = 16^n*Sum_{i=0..n} 2^i*binomial(n, i)*Catalan(2*n-i).

Conjecture: D-finite with recurrence: n*(2*n+1)*a(n) -32*(9*n-4)*(n-1)*a(n-1) +1024*(3*n^2-23*n+29)*a(n-2) +65536*(n-2)*(2*n-5)*a(n-3)=0. - _R. J. Mathar_, Jan 27 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Jun 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013517 | OEIS-A308818

a(n) = a(a(n-1) mod n) + a(a(n-2) mod n) with a(0)=2 and a(1)=3.

Conjectures: (i) This sequence doesn't enter a cycle. (ii) There is an integer greater than 1 that can never appear in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arran Ireland_, Jun 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013518 | OEIS-A308823

Sum of the smallest parts of the partitions of n into 5 parts.

Conjectures from _Colin Barker_, Jun 30 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Jun 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013519 | OEIS-A308824

Sum of the fourth largest parts in the partitions of n into 5 parts.

Conjectures from _Colin Barker_, Jun 30 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Jun 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013520 | OEIS-A308833

Numbers r such that the r-th tetrahedral number A000292(r) divides r!.

Conjecture: for every odd integer r > 1, the following statements are equivalent: a) r is a term of this sequence, b) r + 1 is a term of this sequence, c) r + 2 is composite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lorenzo Sauras Altuzarra_, Jun 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013521 | OEIS-A308838

Orders of Parker finite fields of odd characteristic.

A field or ring is called "Parker" if no 3 X 3 magic square of 9 distinct squared elements can be formed. Conjecture: the sequence is complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Onno M. Cain_, Jun 27 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013522 | OEIS-A308843

Sum of the third largest parts in the partitions of n into 5 squarefree parts.

Conjecture: a(4*k + 3) < a(4*k + 4) for 4*k + 3 >= 195. This conjecture holds for all terms in the b-file. - _David A. Corneth_, Sep 16 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Jun 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013523 | OEIS-A308852

Minimum number k such that the k-th tetrahedral number is not smaller than n!.

Open problem: what is the cardinality of the set of numbers that are simultaneously a tetrahedral number and a factorial number? For example, 1 and 120 belong to this set.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lorenzo Sauras Altuzarra_, Jun 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013524 | OEIS-A308934

Number of ways to write n as (2^a*3^b)^2 + (2^c*3^d)^2 + x^2 + 2*y^2, where a,b,c,d,x,y are nonnegative integers with 2^a*3^b >= 2^c*3^d.

Conjecture 1: a(n) > 0 for all n > 1. Conjecture 2: Any integer n > 1 can be written as (2^a*3^b)^2 + (2^c*5^d)^2 + x^2 + 2*y^2 with a,b,c,d,x,y nonnegative integers. These two conjectures are similar to the Four-square Conjecture in A308734. We have verified Conjectures 1 and 2 for n up to 2*10^9 and 10^9 respectively.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013525 | OEIS-A308950

Number of ways to write n as (p-1)/6 + 2^a*3^b, where p is a prime, and a and b are nonnegative integers.

Conjecture: Let r be 1 or -1. Then, any integer n > 1 can be written as (p-r)/6 + 2^a*3^b, where p is a prime, and a and b are nonnegative integers; in other words, 6*n+r can be written as p + 2^k*3^m, where p is a prime, and k and m are positive integers. Conjecture verified up to n = 10^11. - _Giovanni Resta_, Jul 03 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013526 | OEIS-A309027

Prime powers of the form 12*c^2 + 4*c + 3, where c is an arbitrary integer.

It is conjectured that all terms are prime. See Leung et al. p. 12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Jul 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013527 | OEIS-A309037

Exponential Demlo sequence, like 12345...54321, but for powers of 2 instead.

Conjectures from _Colin Barker_, Jul 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eliora Ben-Gurion_, Jul 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013528 | OEIS-A309042

Numbers k such that there is no prime p and index j > k such that A002182(j) = p * A002182(k).

This is a sequence of counterexamples to the first conjecture by Alaoglu & Erdős that such highly composite numbers do not exist. Robin found the first counterexample: A002182(2557) = 3.000... * 10^153. The sequence of counterexamples to their second conjecture is A272606.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Jul 09 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013529 | OEIS-A309054

a(1) = 0; for m >= 0, a(3m) = 1; for m >= 1, a(3m-1) = 2*a(m-1), a(3m+1) = 2*a(m+1).

Sum_{n=-3m..3m-1} (a(n) + 1) = 4 * Sum_{n=-m..m-1} (a(n) + 1) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Munn_, Jul 09 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013530 | OEIS-A309097

Number of partitions of n avoiding the partition (4,2,1).

Conjecture: for n > 0, a(n) is the number of ordered pairs (r, l) such that there exists a nilpotent matrix of order n whose rank is r and nilpotent index is l. Actually, such a matrix exists if and only if ceiling(n/(n-r)) <= l <= r+1, see my proof below. If this conjecture is true, then a(n) = (n^2 + 3n)/2 - A006590(n) for n > 0. - _Jianing Song_, Nov 04 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan S. Bloom_, Jul 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013531 | OEIS-A309121

a(n) gives the number of primes in the interval I_j = [(j^2 + 3*j - 2)/2, j*(j + 5)/2] = [A034856(j), A095998(j)], for j >= 1.

Conjecture: a(n) >= 1. The length of interval I_n is n+2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jul 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013532 | OEIS-A309129

Numbers k such that -k is a quadratic nonresidue modulo all odd primes p <= sqrt(k) which do not divide k.

Conjecture: 1848 is the last term of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, Jul 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013533 | OEIS-A309130

Smallest prime factor of A077586(n).

a(n) = A077586(n) iff A077586(n) is prime, A077586(n) is prime for 1 <= n <= 4, but composite for 5 <= n <= 17. The status of A077586(18) = 2^(2^61-1)-1 is unknown. It is conjectured that A077586(n) is composite for all n >= 5. Conjecture: All terms are in A122094 (all terms in A263686 are in A122094). For examples related to that conjecture, see A322568. - _Jeppe Stig Nielsen_, Aug 29 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, Jul 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013534 | OEIS-A309156

Rectangular array by antidiagonals: row c is the solution sequence (a(n)) of the complementary equation a(n) = b(n) + b(2n) + c, for c >= 0. See Comments.

Regarding row 1, it is easy to prove that the solution (a(n)) of a(n) = b(n) + b(2n) + 1 is given by a(n) = 4n+3. Conjecture: this is the only linearly recurrent row.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013535 | OEIS-A309158

The smallest prime, a(n), larger than prime(n) for which every even difference from 2 to prime(n) - 1 occurs at least once for some pair of primes from prime(n) to a(n) inclusive.

Conjecture: The sequence is infinite. Conjecture: There are finitely many values of n with a(n) = 2 * prime(n) - 1. Conjecture: There are infinitely many values of n with a(n) = a(n-1). Conjecture: For all n, a(n) <= 3 * prime(n). (This is true for n <= 101.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sally Myers Moite_, Jul 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013536 | OEIS-A309161

a(n) = (x(n)^2 + 1)/m(n), with m(n) = A002559(n) (Markoff numbers) and x(n)= A324601(n), for n >= 3. The Markoff uniqueness conjecture is assumed to be true.

a(n) = (x(n)^2 + 1)/m(n), with m(n) = A002559(n) (Markoff numbers) and x(n)= A324601(n), for n >= 3. The Markoff uniqueness conjecture is assumed to be true. If the equation x^2 + 1 = a(n)*m(n), with m(n) = A002559(n) holds for just one integral x = x(n) in the interval [1, floor(m(n)/2)] then the Markoff uniqueness conjecture is true. x(n) = A324601(n) (if the Markoff conjecture holds).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jul 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013537 | OEIS-A309200

a(n) is the smallest divisor of the Catalan number C(n) = A000108(n) not already in the sequence.

Conjecture: This is a permutation of the positive integers. [The conjecture is true, see A309364. - _Rémy Sigrist_, Jul 25 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 25 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013538 | OEIS-A309221

Expected number of distinct squares visited by a knight's random walk on an infinite chessboard after n steps, normalized to give an integer.

a(0)=1; for n>0, a(n) = (A326954(n)/A326955(n))*2^(3*n-3). (It is only a conjecture that this is always an integer).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013539 | OEIS-A309225

Indices m such that A008348(m)=0.

Probably infinite, on probabilistic grounds, but this is an open question. (Comment modified by _N. J. A. Sloane_, Oct 03 2019.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 31 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013540 | OEIS-A309231

Column 3 of the array at A326662 see Comments.

This is the sequence (c(n)) defined by the complementary equation c(n) = a(2n) + b(2n), with initial value a(1) = 1. See A326662. Conjecture: 9n-c(n) is in {1,2} for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013541 | OEIS-A309263

Terms of A140110 that are not divisible by 6.

Conjecture: The sequence includes all numbers of the form 4*5^n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Jul 19 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013542 | OEIS-A309341

a(n) = Sum_{j=1..n*(n-1)} (n*j mod (n+j)).

Conjecture: a(n) ~ c*n^4 where c = 0.32246....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Jul 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013543 | OEIS-A309353

Let d(1), d(2), ..., d(q) be the q divisors of a number m. The sequence lists the numbers m such that {phi(d(1)), phi(d(2)), ..., phi(d(q))} is the set of the divisors of a number k, where phi is the Euler totient function.

Conjecture: the corresponding numbers k of the sequence is a sequence b(n) of powers of 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Aug 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013544 | OEIS-A309358

Numbers k such that 10^k + 1 is a semiprime.

10^k + 1 is composite unless k is a power of 2, and it can be conjectured that it is composite for all k > 2, cf. A038371 and A185121. - _M. F. Hasler_, Jul 30 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jul 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013545 | OEIS-A309374

Numbers k such that k+j is prime for every j, where 1 <= j < k and gcd(j,k) = 1.

It was conjectured by Recamán Santos in 1976 and proved by Hausman and Shapiro in 1978 that 12 is the largest k possible. Pomerance & Penney (1977) reported in a letter that they have proved that the conjecture is true. - _Amiram Eldar_, May 15 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ivan N. Ianakiev_, Jul 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013546 | OEIS-A309387

a(n) is the smallest positive divisor not yet in the sequence of 7*A000217(n-1); n >= 1.

Conjecture: for large p prime, the odd primes (except p) appear as fixed points of b(n), where b(n) is the smallest positive divisor not yet in the sequence of p*A000217(n-1); n >= 1 (see link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enrique Navarrete_, Jul 27 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013547 | OEIS-A309391

a(n) = gcd(n, A064169(n-2)) for n > 2.

Conjecture: for n > 2, if a(n) = n, then n is a prime. Above conjecture disproved by Adamczewski in linked paper. - _Michael De Vlieger_, Aug 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_ and _Thomas Ordowski_, Jul 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013548 | OEIS-A309393

Let f(n) be equal to n + S(n) + S(S(n)) ... + S(S(S..(n))), where the last term is less than 10 and S(n) is the sum of digits. This is the sequence of numbers k such that the equation f(x) = k has a record number of solutions.

Conjecture from _Daniel Starodubtsev_ and _Dmitry Petukhov_, Nov 19 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Starodubtsev_, Jul 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013549 | OEIS-A309397

a(n) = gcd(n^2, A001008(n-1)) for n > 1.

Conjecture: for n > 3, if a(n) = n^2, then n is a prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_ and _Thomas Ordowski_, Jul 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013550 | OEIS-A309417

Number of steps needed to reduce 10^n to zero by subtracting its digital sum.

Conjecture: lim_{n->infinity} a(n+1)/a(n) = 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Reiner Moewald_, Jul 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013551 | OEIS-A309420

Decimal expansion of 4/(3*Pi-8).

Conjecturally, this can be computed using a recursion formula discovered by an algorithm called "The Ramanujan Machine":

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013552 | OEIS-A309460

a(n) is the smallest number k such that there are n pairs (d,e) of (not necessarily distinct) divisors of k such that the concatenation d_e is a prime.

Conjecture 1: a(n) always exists. Conjecture 2: For any n there are infinitely many choices for m.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius A. Burtea_, Aug 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013553 | OEIS-A309487

Positive integers represented by the quadratic form (the discriminant form) Δ = b^2 - 4ac, where a,b,c are consecutive palindromic primes.

Based on a study of Δ for the case when a,b,c are consecutive primes I conjecture (but have no proof) that now Δ is always negative. The conjecture in the previous comment is true. It says p(n)^2 <= 4*p(n-1)*p(n+1), and this follows from p(n)^2 <= 4*p(n-1)*p(n), i.e. p(n) <= 4*p(n-1), which is true (see A327447, also Mitrinovic, Sect. VII.18 (b)). - _N. J. A. Sloane_, Sep 10 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philip Mizzi_, Sep 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013554 | OEIS-A309514

Number of winning Tak paths on an n X n board.

Conjectured terms above n=5, generated by computer program, are 2012, 22574, 534312, 24816752, and 1908479044.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jakob Gowell_, Aug 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013555 | OEIS-A309515

Sum of the even parts of the partitions of n into 3 parts.

Conjectures from _Colin Barker_, Aug 06 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Aug 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013556 | OEIS-A309516

Number of odd parts in the partitions of n into 4 parts.

Conjectures from _Colin Barker_, Aug 06 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Aug 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013557 | OEIS-A309518

Number of even parts in the partitions of n into 4 parts.

Conjectures from _Colin Barker_, Aug 06 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Aug 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013558 | OEIS-A309580

Primes p with 1 zero in a fundamental period of A000129 mod p.

Conjecturely, since (k+2)/2 = 4 is a square, this sequence has density 7/24 in the primes; see the end of Section 1 of my link. [Comment rewritten by _Jianing Song_, Jun 16 2024 and Jun 25 2024] The conjecture above is an analog of Hasse's result that the set {p prime : ord(2,p) is odd} has density 7/24 in the primes, where ord(a,m) is the multiplicative order of a modulo m; see A014663. - _Jianing Song_, Jun 26 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013559 | OEIS-A309581

Primes p with 2 zeros in a fundamental period of A000129 mod p.

Conjecturely, since (k+2)/2 = 4 is a square, this sequence has density 5/12 in the primes; see the end of Section 1 of my link. [Comment rewritten by _Jianing Song_, Jun 16 2024 and Jun 25 2024] The conjecture above is an analog of Hasse's result that the set {p prime : ord(2,p) is odd} has density 7/24 in the primes, where ord(a,m) is the multiplicative order of a modulo m; see A014663. - _Jianing Song_, Jun 26 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013560 | OEIS-A309586

Primes p with 1 zero in a fundamental period of A006190 mod p.

Conjecturely, this sequence has density 1/3 in the primes. [Comment rewritten by _Jianing Song_, Jun 16 2024 and Jun 25 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013561 | OEIS-A309616

a(n) is the number of ways to represent 2*n in the decibinary system.

Conjecture: a(n) = a(n-1) + a(ceiling(n/2)) - a(ceiling((n-5)/2)) if n > 5. The conjectured recurrence is true (see Fried link). - _Sela Fried_, Mar 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonas Hollm_, Aug 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013562 | OEIS-A309655

The smallest possible nonnegative difference between the sum of the first n primes (A007504) and the sum of any number of the directly following and consecutive primes.

Conjecture: a(0)=0, a(2)=0 and a(532)=0 are the only zeros in the sequence. a(n) has been computed for primes < 10^10. - _Bob Andriesse_, Oct 07 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bob Andriesse_, Aug 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013563 | OEIS-A309663

Primes that begin a run of consecutive primes whose first differences are nondecreasing.

Conjecture: Average length of each run = e, converging from above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Woodward_, Jun 06 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013564 | OEIS-A309666

a(n) is the least k such that the denominators of continued fraction convergents for sqrt(k) match the first n Fibonacci numbers.

Conjectures from _Colin Barker_, Aug 26 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Greg Dresden_, Aug 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013565 | OEIS-A309696

Numbers m such that the numerator of Sum_{k=1..m, gcd(k,m) = 1} 1/k^2 is divisible by m^2.

Conjecture: odd terms of this sequence are odd terms of A290815.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_ and _Thomas Ordowski_, Aug 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013566 | OEIS-A309746

Number of ways of placing 2*n-1 nonattacking queens on a hexagonal board with edge-length n in Glinski's hexagonal chess.

Conjecture: for n >= 12 is a(n) > 0. Proved for n <= 20. - _Vaclav Kotesovec_, Sep 06 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Aug 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013567 | OEIS-A309781

a(n) is the smallest even number m having n distinct odd prime divisors p_1, p_2, ..., p_n, each of which (p_i; i=1..n) has the property that there exists a k_i (0 < k_i < p_i-1) such that p_i - k_i | m - k_i.

Subsequence of A309769 and A309780. a(1) is the only nonsquarefree known term. Conjecture: For n > 1 the minimal condition requires m to be squarefree and for every odd prime divisor p of m to be such that m/p - 1 is composite with least prime divisor q < p (k=p-q). No term is divisible by

3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Aug 17 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013568 | OEIS-A309782

The sequence S is always extended with the smallest integer not yet present that begins with the sum of the two digits enclosing the last comma; a(1) = 1 is separated from a(2) = 2 by the first comma.

This sequence is conjectured to be a permutation of the integers > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Aug 17 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013569 | OEIS-A309791

Expansion of (1 + 8*x - 6*x^2 + 12*x^3 - 18*x^4)/(1 - x - 9*x^4 + 9*x^5).

Conjecture: Any finite subset of these maps can build chains of finite length only.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Georg Fischer_, Aug 17 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013570 | OEIS-A309815

a(n) is the smallest positive integer x such that sqrt(2) + sqrt(x) is closer to an integer than any other value already in the sequence.

If b(n) = round(sqrt(2) + sqrt(a(n))), then (b(n)^2 + 2 - a(n))/(2*b(n)) is an approximation for sqrt(2). Conjecture: all convergents of the continued fraction of sqrt(2) except 1 arise in this way. - _Robert Israel_, Aug 18 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ben Paul Thurston_, Aug 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013571 | OEIS-A309846

Number of degree n polynomials f with all nonzero coefficients equal to 1 such that f(k) is divisible by 4 for all integers k.

Conjecture: All terms are of the form 2^(n-5), 2^k*(2^(n-k-5) + 1), or 2^k*(2^(n-k-5) - 1) for some value of k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Nov 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013572 | OEIS-A309869

Let b(0)=n, b(1)=n+1, b(2)=n+2, and b(k)=(b(k-3)*b(k-2)*b(k-3)) mod (b(k-3)+b(k-2)+b(k-1)) for k>=3. Continue until either b(k)=0, in which case a(n)=k, or the ordered triple [b(k-2),b(k-1),b(k)] has appeared before, in which case a(n)=-k. If neither of these ever occur, then a(n)=0.

a(26-6*(-1)^k+132*k)=5. Conjecture: the only other n for which a(n)=5 are 14 and 32. - _Robert Israel_, Aug 21 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Aug 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013573 | OEIS-A309878

The real part of b(n) where b(n) = (n + b(n-1)) * (1 + i) with b(-1)=0; i = sqrt(-1).

Conjectures from _Colin Barker_, Aug 21 2019: (Start) E.g.f.: exp(x)*(-x + 2*sin(x)). - Conjectured by _Stefano Spezia_, Aug 21 2019 after _Colin Barker_ Conjecture: a(n) = 2*A009545(n)-n. - _R. J. Mathar_, Mar 06 2022 All conjectures stated above hold true. See links. - _Sela Fried_, Jul 27 2024.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jesse Fiedler_, Aug 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013574 | OEIS-A309879

Number of odd parts appearing among the fourth largest parts of the partitions of n into 5 parts.

Conjectures from _Colin Barker_, Aug 22 2019: (Start) (End) [Conjectures verified by _Wesley Ivan Hurt_, Aug 24 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Aug 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013575 | OEIS-A309906

a(n) is the smallest number of divisors of p^n - 1 that may possibly occur for arbitrarily large primes p.

The existence of infinitely many primes p such that p^n - 1 has exactly a(n) divisors is conjectured. E.g., although it is known that p-1 has fewer than 4 divisors for only finitely many primes p (see Example section), it is not known whether there exist infinitely many primes p such that p-1 has exactly 4 divisors. (Thanks to _Jianing Song_, who pointed out the need for this clarification.) - _Jon E. Schoenfield_, Mar 04 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Aug 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013576 | OEIS-A309948

Decimal expansion of the real part of the square root of 1 + i.

Conjecture: Equals Sum_{k>=0} A208850(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alonso del Arte_, Aug 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013577 | OEIS-A309985

Maximum determinant of an n X n Latin square.

a(10) = 36843728625, conjectured. See Stack Exchange link. - _Hugo Pfoertner_, Sep 29 2019 The conjectured value a(10) = 36843728625 above is false. The noncirculant Latin square in the Example section has absolute determinant 36860608125. Therefore a(10) >= 36860608125 > A328030(10) = 36843728625, and the maximum absolute determinant at order 10 cannot be attained by a circulant Latin square. - _Arthur Queiroz M._, Aug 06 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Aug 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013578 | OEIS-A310007

Coordination sequence Gal.4.31.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013579 | OEIS-A310018

Coordination sequence Gal.4.34.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013580 | OEIS-A310019

Coordination sequence Gal.6.110.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013581 | OEIS-A310025

Coordination sequence Gal.4.31.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013582 | OEIS-A310027

Coordination sequence Gal.5.41.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013583 | OEIS-A310031

Coordination sequence Gal.5.39.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013584 | OEIS-A310039

Coordination sequence Gal.3.7.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013585 | OEIS-A310040

Coordination sequence Gal.6.109.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013586 | OEIS-A310050

Coordination sequence Gal.4.36.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013587 | OEIS-A310068

Coordination sequence Gal.4.25.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013588 | OEIS-A310089

Coordination sequence Gal.6.42.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013589 | OEIS-A310096

Coordination sequence Gal.4.15.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013590 | OEIS-A310100

Coordination sequence Gal.5.19.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Colin Barker_, Apr 18 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013591 | OEIS-A310102

Coordination sequence Gal.4.16.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013592 | OEIS-A310117

Coordination sequence Gal.5.20.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013593 | OEIS-A310143

Coordination sequence Gal.3.6.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 04 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013594 | OEIS-A310146

Coordination sequence Gal.5.39.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013595 | OEIS-A310185

Coordination sequence Gal.4.15.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013596 | OEIS-A310190

Coordination sequence Gal.5.19.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013597 | OEIS-A310215

Coordination sequence Gal.3.2.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013598 | OEIS-A310218

Coordination sequence Gal.5.20.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 27 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013599 | OEIS-A310223

Coordination sequence Gal.6.12.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 18 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013600 | OEIS-A310244

Coordination sequence Gal.6.45.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 12 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013601 | OEIS-A310267

Coordination sequence Gal.4.12.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 12 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013602 | OEIS-A310279

Coordination sequence Gal.4.2.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 18 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013603 | OEIS-A310289

Coordination sequence Gal.4.19.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 30 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013604 | OEIS-A310319

Coordination sequence Gal.3.4.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 16 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013605 | OEIS-A310368

Coordination sequence Gal.6.230.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 04 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013606 | OEIS-A310369

Coordination sequence Gal.5.64.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 08 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013607 | OEIS-A310374

Coordination sequence Gal.6.245.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 19 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013608 | OEIS-A310378

Coordination sequence Gal.6.320.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013609 | OEIS-A310392

Coordination sequence Gal.6.366.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013610 | OEIS-A310393

Coordination sequence Gal.6.193.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 08 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013611 | OEIS-A310394

Coordination sequence Gal.6.568.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 13 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013612 | OEIS-A310400

Coordination sequence Gal.6.337.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 14 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013613 | OEIS-A310430

Coordination sequence Gal.3.41.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 16 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013614 | OEIS-A310440

Coordination sequence Gal.6.202.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 14 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013615 | OEIS-A310458

Coordination sequence Gal.4.78.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 03 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013616 | OEIS-A310462

Coordination sequence Gal.5.138.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 19 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013617 | OEIS-A310492

Coordination sequence Gal.6.249.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013618 | OEIS-A310499

Coordination sequence Gal.6.351.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013619 | OEIS-A310501

Coordination sequence Gal.6.328.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013620 | OEIS-A310502

Coordination sequence Gal.5.132.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013621 | OEIS-A310503

Coordination sequence Gal.4.73.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013622 | OEIS-A310504

Coordination sequence Gal.6.329.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013623 | OEIS-A310508

Coordination sequence Gal.4.123.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013624 | OEIS-A310509

Coordination sequence Gal.5.142.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013625 | OEIS-A310511

Coordination sequence Gal.6.354.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013626 | OEIS-A310512

Coordination sequence Gal.6.352.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013627 | OEIS-A310516

Coordination sequence Gal.6.599.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013628 | OEIS-A310518

Coordination sequence Gal.3.21.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013629 | OEIS-A310521

Coordination sequence Gal.5.133.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013630 | OEIS-A310525

Coordination sequence Gal.4.74.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013631 | OEIS-A310529

Coordination sequence Gal.4.75.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013632 | OEIS-A310530

Coordination sequence Gal.5.134.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013633 | OEIS-A310532

Coordination sequence Gal.6.333.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013634 | OEIS-A310556

Coordination sequence Gal.5.279.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 18 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013635 | OEIS-A310601

Coordination sequence Gal.6.32.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Nov 18 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013636 | OEIS-A310618

Coordination sequence Gal.4.121.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 02 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013637 | OEIS-A310640

Coordination sequence Gal.6.527.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures: for k >= 1, a(7*k-1) = a(7*k) = 28 + (k-1)*30 and a(7*k+1) = a(7*k+2) = 36 + (k-1)*30. - _Bernard Schott_, Dec 24 2019 Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013638 | OEIS-A310648

Coordination sequence Gal.5.253.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013639 | OEIS-A310650

Coordination sequence Gal.6.547.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013640 | OEIS-A310651

Coordination sequence Gal.6.548.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013641 | OEIS-A310668

Coordination sequence Gal.4.30.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013642 | OEIS-A310683

Coordination sequence Gal.4.15.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013643 | OEIS-A310704

Coordination sequence Gal.3.2.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013644 | OEIS-A310744

Coordination sequence Gal.4.11.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 15 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013645 | OEIS-A310784

Coordination sequence Gal.6.72.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 05 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013646 | OEIS-A310936

Coordination sequence Gal.3.7.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Mar 11 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013647 | OEIS-A310988

Coordination sequence Gal.4.16.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013648 | OEIS-A310999

Coordination sequence Gal.4.7.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013649 | OEIS-A311002

Coordination sequence Gal.3.3.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013650 | OEIS-A311007

Coordination sequence Gal.4.19.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013651 | OEIS-A311042

Coordination sequence Gal.4.11.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013652 | OEIS-A311057

Coordination sequence Gal.4.18.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 18 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013653 | OEIS-A311075

Coordination sequence Gal.6.72.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013654 | OEIS-A311088

Coordination sequence Gal.4.25.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013655 | OEIS-A311094

Coordination sequence Gal.6.32.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 12 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013656 | OEIS-A311101

Coordination sequence Gal.3.47.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013657 | OEIS-A311285

Coordination sequence Gal.4.70.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 28 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013658 | OEIS-A311332

Coordination sequence Gal.6.120.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 12 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013659 | OEIS-A311343

Coordination sequence Gal.4.60.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 03 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013660 | OEIS-A311344

Coordination sequence Gal.5.95.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 15 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013661 | OEIS-A311385

Coordination sequence Gal.6.221.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 13 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013662 | OEIS-A311416

Coordination sequence Gal.4.48.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Mar 05 2024: (Start) Conjecture: a(n) ~ 21*n/5. - _Stefano Spezia_, Mar 05 2024 after _Chai Wah Wu_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013663 | OEIS-A311491

Coordination sequence Gal.5.91.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 13 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013664 | OEIS-A311508

Coordination sequence Gal.3.11.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013665 | OEIS-A311533

Coordination sequence Gal.4.51.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013666 | OEIS-A311542

Coordination sequence Gal.5.155.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 11 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013667 | OEIS-A311553

Coordination sequence Gal.6.133.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013668 | OEIS-A311566

Coordination sequence Gal.4.93.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 15 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013669 | OEIS-A311593

Coordination sequence Gal.6.192.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013670 | OEIS-A311600

Coordination sequence Gal.4.65.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013671 | OEIS-A311601

Coordination sequence Gal.6.230.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013672 | OEIS-A311626

Coordination sequence Gal.6.232.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 24 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013673 | OEIS-A311627

Coordination sequence Gal.6.190.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Mar 06 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013674 | OEIS-A311647

Coordination sequence Gal.4.85.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Mar 09 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013675 | OEIS-A311665

Coordination sequence Gal.6.481.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Nov 18 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013676 | OEIS-A311666

Coordination sequence Gal.4.108.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013677 | OEIS-A311674

Coordination sequence Gal.6.58.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 15 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013678 | OEIS-A311686

Coordination sequence Gal.5.182.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Mar 13 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013679 | OEIS-A311714

Coordination sequence Gal.6.435.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 29 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013680 | OEIS-A311824

Coordination sequence Gal.5.158.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Nov 19 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013681 | OEIS-A311826

Coordination sequence Gal.5.91.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 28 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013682 | OEIS-A311861

Coordination sequence Gal.6.130.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 20 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013683 | OEIS-A311888

Coordination sequence Gal.4.45.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 12 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013684 | OEIS-A311936

Coordination sequence Gal.6.300.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013685 | OEIS-A311943

Coordination sequence Gal.4.80.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 18 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013686 | OEIS-A311993

Coordination sequence Gal.5.98.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 14 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013687 | OEIS-A312006

Coordination sequence Gal.4.23.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 25 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013688 | OEIS-A312105

Coordination sequence Gal.4.87.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 20 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013689 | OEIS-A312210

Coordination sequence Gal.6.159.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 01 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013690 | OEIS-A312226

Coordination sequence Gal.3.18.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 02 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013691 | OEIS-A312289

Coordination sequence Gal.6.527.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 19 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013692 | OEIS-A312402

Coordination sequence Gal.4.70.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013693 | OEIS-A312411

Coordination sequence Gal.3.39.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013694 | OEIS-A312484

Coordination sequence Gal.5.81.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 26 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013695 | OEIS-A312486

Coordination sequence Gal.3.30.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 09 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013696 | OEIS-A312508

Coordination sequence Gal.5.250.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 28 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013697 | OEIS-A312549

Coordination sequence Gal.4.61.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 24 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013698 | OEIS-A312556

Coordination sequence Gal.6.246.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 07 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013699 | OEIS-A312722

Coordination sequence Gal.5.146.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013700 | OEIS-A312773

Coordination sequence Gal.5.276.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

a(n) = - 2*a(n-1) - 2*a(n-2) - 2*a(n-3) - 2*a(n-4) + 3*a(n-6) + 4*a(n-7) + 4*a(n-8) + 4*a(n-9) + 3*a(n-10) - 2*a(n-12) - 2*a(n-13) - 2*a(n-14) - 2*a(n-15) - a(n-16) for n > 17 (conjectured). G.f.: (-2*x^17 - x^16 + 4*x^15 + 16*x^14 + 44*x^13 + 80*x^12 + 124*x^11 + 163*x^10 + 192*x^9 + 202*x^8 + 188*x^7 + 155*x^6 + 116*x^5 + 76*x^4 + 42*x^3 + 18*x^2 + 6*x + 1)/((x - 1)^2*(x + 1)^2*(x^2 - x + 1)*(x^2 + x + 1)*(x^4 + x^3 + x^2 + x + 1)^2) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013701 | OEIS-A312822

Coordination sequence Gal.6.577.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 21 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013702 | OEIS-A312860

Coordination sequence Gal.6.546.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 15 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013703 | OEIS-A312909

Coordination sequence Gal.6.150.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013704 | OEIS-A312926

Coordination sequence Gal.6.118.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013705 | OEIS-A312927

Coordination sequence Gal.6.149.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013706 | OEIS-A312928

Coordination sequence Gal.3.16.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013707 | OEIS-A312932

Coordination sequence Gal.5.81.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013708 | OEIS-A312933

Coordination sequence Gal.6.131.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013709 | OEIS-A312935

Coordination sequence Gal.4.105.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013710 | OEIS-A312938

Coordination sequence Gal.5.127.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013711 | OEIS-A312941

Coordination sequence Gal.5.14.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013712 | OEIS-A312943

Coordination sequence Gal.4.41.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013713 | OEIS-A312944

Coordination sequence Gal.6.245.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013714 | OEIS-A312948

Coordination sequence Gal.5.66.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013715 | OEIS-A312950

Coordination sequence Gal.6.151.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013716 | OEIS-A312956

Coordination sequence Gal.5.264.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013717 | OEIS-A312960

Coordination sequence Gal.4.106.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013718 | OEIS-A312961

Coordination sequence Gal.6.152.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013719 | OEIS-A312963

Coordination sequence Gal.6.153.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 23 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013720 | OEIS-A312969

Coordination sequence Gal.6.254.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013721 | OEIS-A312980

Coordination sequence Gal.6.248.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 01 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013722 | OEIS-A312981

Coordination sequence Gal.5.146.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 04 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013723 | OEIS-A312982

Coordination sequence Gal.5.232.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 16 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013724 | OEIS-A313025

Coordination sequence Gal.5.65.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 10 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013725 | OEIS-A313033

Coordination sequence Gal.5.247.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 08 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013726 | OEIS-A313059

Coordination sequence Gal.4.99.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 28 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013727 | OEIS-A313088

Coordination sequence Gal.6.120.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Mar 30 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013728 | OEIS-A313097

Coordination sequence Gal.5.195.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 31 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013729 | OEIS-A313121

Coordination sequence Gal.3.25.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 06 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013730 | OEIS-A313124

Coordination sequence Gal.6.250.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Nov 19 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013731 | OEIS-A313136

Coordination sequence Gal.6.256.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 24 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013732 | OEIS-A313144

Coordination sequence Gal.5.219.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 03 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013733 | OEIS-A313145

Coordination sequence Gal.6.264.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 02 2025: (Start)
Conjecture: a(n) ~ 29*n/6. - _Stefano Spezia_, Jun 05 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013734 | OEIS-A313150

Coordination sequence Gal.6.382.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 12 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013735 | OEIS-A313153

Coordination sequence Gal.5.113.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 08 2025: (Start)
Conjecture: a(n) ~ 46*n/9. - _Stefano Spezia_, May 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013736 | OEIS-A313154

Coordination sequence Gal.6.252.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 13 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013737 | OEIS-A313161

Coordination sequence Gal.5.237.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 12 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013738 | OEIS-A313172

Coordination sequence Gal.5.234.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Mar 03 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013739 | OEIS-A313216

Coordination sequence Gal.6.367.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 20 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013740 | OEIS-A313217

Coordination sequence Gal.6.382.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013741 | OEIS-A313241

Coordination sequence Gal.5.86.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 19 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013742 | OEIS-A313255

Coordination sequence Gal.5.87.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013743 | OEIS-A313259

Coordination sequence Gal.6.204.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 19 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013744 | OEIS-A313264

Coordination sequence Gal.5.216.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 28 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013745 | OEIS-A313265

Coordination sequence Gal.6.475.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Mar 13 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013746 | OEIS-A313295

Coordination sequence Gal.5.90.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 30 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013747 | OEIS-A313296

Coordination sequence Gal.6.209.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 16 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013748 | OEIS-A313297

Coordination sequence Gal.6.210.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Mar 22 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013749 | OEIS-A313321

Coordination sequence Gal.3.42.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 16 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013750 | OEIS-A313426

Coordination sequence Gal.5.220.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 25 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013751 | OEIS-A313442

Coordination sequence Gal.6.453.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 16 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013752 | OEIS-A313524

Coordination sequence Gal.5.219.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 17 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013753 | OEIS-A313556

Coordination sequence Gal.6.132.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 14 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013754 | OEIS-A313571

Coordination sequence Gal.5.169.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 16 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013755 | OEIS-A313577

Coordination sequence Gal.4.97.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 16 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013756 | OEIS-A313614

Coordination sequence Gal.5.129.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 30 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013757 | OEIS-A313615

Coordination sequence Gal.6.323.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 14 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013758 | OEIS-A313639

Coordination sequence Gal.6.204.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 17 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013759 | OEIS-A313646

Coordination sequence Gal.6.205.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 08 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013760 | OEIS-A313682

Coordination sequence Gal.5.55.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 15 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013761 | OEIS-A313685

Coordination sequence Gal.4.43.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 19 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013762 | OEIS-A313694

Coordination sequence Gal.4.63.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 13 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013763 | OEIS-A313697

Coordination sequence Gal.5.202.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 02 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013764 | OEIS-A313698

Coordination sequence Gal.6.157.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 08 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013765 | OEIS-A313707

Coordination sequence Gal.6.198.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 24 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013766 | OEIS-A313721

Coordination sequence Gal.5.129.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 08 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013767 | OEIS-A313730

Coordination sequence Gal.6.234.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 18 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013768 | OEIS-A313740

Coordination sequence Gal.5.309.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Mar 12 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013769 | OEIS-A313747

Coordination sequence Gal.6.200.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Oct 20 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013770 | OEIS-A313754

Coordination sequence Gal.4.142.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 29 2025: (Start)
Conjecture: a(n) ~ 16*n/3. - _Stefano Spezia_, May 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013771 | OEIS-A313764

Coordination sequence Gal.3.61.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 01 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013772 | OEIS-A313774

Coordination sequence Gal.6.622.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 10 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013773 | OEIS-A313782

Coordination sequence Gal.5.289.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 06 2025: (Start)
Conjecture: a(n) ~ 26*n/5. - _Stefano Spezia_, May 06 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013774 | OEIS-A313793

Coordination sequence Gal.4.134.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 06 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013775 | OEIS-A313796

Coordination sequence Gal.6.364.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013776 | OEIS-A313799

Coordination sequence Gal.3.53.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 12 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013777 | OEIS-A313824

Coordination sequence Gal.6.195.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 31 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013778 | OEIS-A313836

Coordination sequence Gal.3.23.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 16 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013779 | OEIS-A313837

Coordination sequence Gal.6.618.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 06 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013780 | OEIS-A313855

Coordination sequence Gal.5.87.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 13 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013781 | OEIS-A313859

Coordination sequence Gal.6.340.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 01 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013782 | OEIS-A313884

Coordination sequence Gal.6.347.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 09 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013783 | OEIS-A313896

Coordination sequence Gal.6.364.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 12 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013784 | OEIS-A313900

Coordination sequence Gal.6.621.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 11 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013785 | OEIS-A313912

Coordination sequence Gal.6.207.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 08 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013786 | OEIS-A313921

Coordination sequence Gal.5.313.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 09 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013787 | OEIS-A313922

Coordination sequence Gal.6.356.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Oct 16 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013788 | OEIS-A313929

Coordination sequence Gal.6.334.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 16 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013789 | OEIS-A313940

Coordination sequence Gal.4.138.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 19 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013790 | OEIS-A313955

Coordination sequence Gal.4.144.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 16 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013791 | OEIS-A313962

Coordination sequence Gal.3.53.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 02 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013792 | OEIS-A313965

Coordination sequence Gal.5.284.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 07 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013793 | OEIS-A313969

Coordination sequence Gal.6.489.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Mar 13 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013794 | OEIS-A313981

Coordination sequence Gal.5.319.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 15 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013795 | OEIS-A314015

Coordination sequence Gal.4.100.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 17 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013796 | OEIS-A314021

Coordination sequence Gal.5.84.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 13 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013797 | OEIS-A314024

Coordination sequence Gal.6.156.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 20 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013798 | OEIS-A314039

Coordination sequence Gal.5.299.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 16 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013799 | OEIS-A314048

Coordination sequence Gal.6.618.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 02 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013800 | OEIS-A314054

Coordination sequence Gal.4.142.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 25 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013801 | OEIS-A314055

Coordination sequence Gal.4.143.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 19 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013802 | OEIS-A314056

Coordination sequence Gal.4.135.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 03 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013803 | OEIS-A314065

Coordination sequence Gal.4.143.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 20 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013804 | OEIS-A314068

Coordination sequence Gal.3.55.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 03 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013805 | OEIS-A314070

Coordination sequence Gal.6.657.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 02 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013806 | OEIS-A314088

Coordination sequence Gal.5.111.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 22 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013807 | OEIS-A314093

Coordination sequence Gal.5.137.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 16 2025: (Start)
Conjecture: a(n) ~ 5*n. - _Stefano Spezia_, Apr 17 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013808 | OEIS-A314094

Coordination sequence Gal.6.345.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 16 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013809 | OEIS-A314098

Coordination sequence Gal.6.339.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 15 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013810 | OEIS-A314108

Coordination sequence Gal.6.198.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 29 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013811 | OEIS-A314121

Coordination sequence Gal.6.657.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 17 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013812 | OEIS-A314127

Coordination sequence Gal.6.326.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 16 2026: (Start)
Conjecture: a(n) ~ 58*n/11. - _Stefano Spezia_, Jun 17 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013813 | OEIS-A314140

Coordination sequence Gal.6.633.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 15 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013814 | OEIS-A314159

Coordination sequence Gal.6.624.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Colin Barker_, Aug 27 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013815 | OEIS-A314161

Coordination sequence Gal.4.140.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 20 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013816 | OEIS-A314167

Coordination sequence Gal.6.638.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 09 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013817 | OEIS-A314193

Coordination sequence Gal.4.141.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 15 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013818 | OEIS-A314203

Coordination sequence Gal.5.88.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 08 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013819 | OEIS-A314216

Coordination sequence Gal.6.624.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 16 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013820 | OEIS-A314219

Coordination sequence Gal.5.314.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 12 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013821 | OEIS-A314220

Coordination sequence Gal.5.306.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 14 2025: (Start)
Conjecture: a(n) ~ 28*n/5. - _Stefano Spezia_, Jul 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013822 | OEIS-A314228

Coordination sequence Gal.6.650.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 20 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013823 | OEIS-A314234

Coordination sequence Gal.6.332.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 29 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013824 | OEIS-A314236

Coordination sequence Gal.6.626.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 15 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013825 | OEIS-A314259

Coordination sequence Gal.3.60.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 13 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013826 | OEIS-A314287

Coordination sequence Gal.6.654.1 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 28 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013827 | OEIS-A314324

Coordination sequence Gal.5.252.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 13 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013828 | OEIS-A314377

Coordination sequence Gal.6.109.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 10 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013829 | OEIS-A314400

Coordination sequence Gal.6.392.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 12 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013830 | OEIS-A314460

Coordination sequence Gal.4.110.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 30 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013831 | OEIS-A314461

Coordination sequence Gal.6.568.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

a(n) = 2*a(n-7) - a(n-14) for n > 16 (conjectured). G.f.: (-x^16 - x^15 + 2*x^14 + 6*x^13 + 9*x^12 + 14*x^11 + 20*x^10 + 26*x^9 + 30*x^8 + 30*x^7 + 26*x^6 + 24*x^5 + 20*x^4 + 13*x^3 + 8*x^2 + 5*x + 1)/(x^14 - 2*x^7 + 1) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013832 | OEIS-A314466

Coordination sequence Gal.5.205.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Nov 19 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013833 | OEIS-A314477

Coordination sequence Gal.5.193.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 14 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013834 | OEIS-A314658

Coordination sequence Gal.5.14.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 30 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013835 | OEIS-A314697

Coordination sequence Gal.5.125.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Nov 19 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013836 | OEIS-A314712

Coordination sequence Gal.6.150.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 15 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013837 | OEIS-A314740

Coordination sequence Gal.5.116.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 13 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013838 | OEIS-A314742

Coordination sequence Gal.6.262.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 28 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013839 | OEIS-A314772

Coordination sequence Gal.4.126.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 15 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013840 | OEIS-A314807

Coordination sequence Gal.6.528.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 13 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013841 | OEIS-A314829

Coordination sequence Gal.4.41.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 08 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013842 | OEIS-A314838

Coordination sequence Gal.4.52.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Mar 08 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013843 | OEIS-A314839

Coordination sequence Gal.6.357.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Nov 19 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013844 | OEIS-A314840

Coordination sequence Gal.5.66.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 16 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013845 | OEIS-A314847

Coordination sequence Gal.6.193.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 18 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013846 | OEIS-A314849

Coordination sequence Gal.6.246.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Nov 19 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013847 | OEIS-A314929

Coordination sequence Gal.5.189.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 12 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013848 | OEIS-A314945

Coordination sequence Gal.5.84.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 19 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013849 | OEIS-A314955

Coordination sequence Gal.6.322.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 14 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013850 | OEIS-A314963

Coordination sequence Gal.6.199.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 02 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013851 | OEIS-A314994

Coordination sequence Gal.6.547.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 27 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013852 | OEIS-A315001

Coordination sequence Gal.6.190.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 18 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013853 | OEIS-A315045

Coordination sequence Gal.6.366.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Feb 03 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013854 | OEIS-A315047

Coordination sequence Gal.5.204.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 20 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013855 | OEIS-A315129

Coordination sequence Gal.6.401.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Nov 19 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013856 | OEIS-A315135

Coordination sequence Gal.4.122.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 18 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013857 | OEIS-A315187

Coordination sequence Gal.3.14.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 28 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013858 | OEIS-A315203

Coordination sequence Gal.6.321.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 19 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013859 | OEIS-A315213

Coordination sequence Gal.6.453.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 19 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013860 | OEIS-A315227

Coordination sequence Gal.5.169.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 27 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013861 | OEIS-A315266

Coordination sequence Gal.6.205.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Mar 29 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013862 | OEIS-A315271

Coordination sequence Gal.6.345.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 31 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013863 | OEIS-A315301

Coordination sequence Gal.6.584.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 30 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013864 | OEIS-A315303

Coordination sequence Gal.6.247.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 15 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013865 | OEIS-A315304

Coordination sequence Gal.5.84.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 14 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013866 | OEIS-A315306

Coordination sequence Gal.5.130.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 21 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013867 | OEIS-A315316

Coordination sequence Gal.6.324.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 12 2025: (Start)
Conjecture: a(n) ~ 56*n/11. - _Stefano Spezia_, Jun 12 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013868 | OEIS-A315317

Coordination sequence Gal.5.257.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Oct 19 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013869 | OEIS-A315318

Coordination sequence Gal.6.622.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Oct 19 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013870 | OEIS-A315319

Coordination sequence Gal.5.289.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Oct 19 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013871 | OEIS-A315320

Coordination sequence Gal.4.135.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Oct 19 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013872 | OEIS-A315322

Coordination sequence Gal.6.636.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Oct 19 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013873 | OEIS-A315327

Coordination sequence Gal.6.329.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Oct 19 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013874 | OEIS-A315339

Coordination sequence Gal.4.141.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 09 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013875 | OEIS-A315343

Coordination sequence Gal.5.304.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 04 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013876 | OEIS-A315368

Coordination sequence Gal.6.46.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 01 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013877 | OEIS-A315405

Coordination sequence Gal.3.15.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 02 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013878 | OEIS-A315418

Coordination sequence Gal.3.21.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 10 2025: (Start)
Conjecture: a(n) ~ 26*n/5. - _Stefano Spezia_, Jul 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013879 | OEIS-A315420

Coordination sequence Gal.6.355.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Nov 18 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013880 | OEIS-A315431

Coordination sequence Gal.6.669.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

a(n) = a(n-4) - a(n-5) + a(n-7) + a(n-9) - a(n-11) + a(n-12) - a(n-16) for n > 22 (conjectured). G.f.: (-x^22 - x^17 + x^16 + 8*x^15 + 11*x^14 + 16*x^13 + 21*x^12 + 23*x^11 + 30*x^10 + 34*x^9 + 31*x^8 + 34*x^7 + 28*x^6 + 23*x^5 + 21*x^4 + 15*x^3 + 11*x^2 + 6*x + 1)/(x^16 - x^12 + x^11 - x^9 - x^7 + x^5 - x^4 + 1) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013881 | OEIS-A315447

Coordination sequence Gal.5.112.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 01 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013882 | OEIS-A315460

Coordination sequence Gal.6.234.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 27 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013883 | OEIS-A315462

Coordination sequence Gal.6.339.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

a(n) = 2*a(n-1) - a(n-2) - a(n-4) + 2*a(n-5) - a(n-6) - a(n-8) + 2*a(n-9) - a(n-10) for n > 10 (conjectured). G.f.: (x^10 + 4*x^9 + x^6 + 3*x^5 + x^4 + 4*x + 1)/(x^8*(x - 1)^2 + x^4*(x - 1)^2 + (x - 1)^2) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013884 | OEIS-A315464

Coordination sequence Gal.6.346.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

a(n) = a(n-1) + a(n-11) - a(n-12) for n > 12 (conjectured). G.f.: (x^12 + 5*x^11 + 5*x^10 + 5*x^9 + 5*x^8 + 4*x^7 + 6*x^6 + 4*x^5 + 5*x^4 + 5*x^3 + 5*x^2 + 5*x + 1)/(x^12 - x^11 - x + 1) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013885 | OEIS-A315469

Coordination sequence Gal.4.134.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 20 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013886 | OEIS-A315471

Coordination sequence Gal.3.52.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 20 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013887 | OEIS-A315477

Coordination sequence Gal.5.293.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Jianing Song_, Jun 02 2026: (Start)
Conjecture: a(n) ~ 27*n/5. - _Stefano Spezia_, Jun 02 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013888 | OEIS-A315487

Coordination sequence Gal.5.214.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 22 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013889 | OEIS-A315492

Coordination sequence Gal.6.625.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 02 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013890 | OEIS-A315495

Coordination sequence Gal.5.135.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 14 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013891 | OEIS-A315496

Coordination sequence Gal.5.306.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 05 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013892 | OEIS-A315498

Coordination sequence Gal.6.334.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 10 2025: (Start)
Conjecture: a(n) ~ 62*n/11. - _Stefano Spezia_, Jun 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013893 | OEIS-A315501

Coordination sequence Gal.4.149.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 16 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013894 | OEIS-A315511

Coordination sequence Gal.4.139.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013895 | OEIS-A315517

Coordination sequence Gal.6.208.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013896 | OEIS-A315520

Coordination sequence Gal.4.140.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 11 2020: (Start)
Conjecture: a(n) = (22*n + 2*A188510(n))/4 for n > 0. - _Stefano Spezia_, Jul 30 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013897 | OEIS-A315522

Coordination sequence Gal.6.621.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013898 | OEIS-A315525

Coordination sequence Gal.3.49.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013899 | OEIS-A315532

Coordination sequence Gal.6.662.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 07 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013900 | OEIS-A315538

Coordination sequence Gal.3.50.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013901 | OEIS-A315540

Coordination sequence Gal.6.643.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 21 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013902 | OEIS-A315541

Coordination sequence Gal.4.136.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013903 | OEIS-A315543

Coordination sequence Gal.4.137.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013904 | OEIS-A315546

Coordination sequence Gal.5.294.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013905 | OEIS-A315548

Coordination sequence Gal.6.627.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 27 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013906 | OEIS-A315549

Coordination sequence Gal.6.628.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013907 | OEIS-A315557

Coordination sequence Gal.6.663.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 12 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013908 | OEIS-A315559

Coordination sequence Gal.4.149.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 09 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013909 | OEIS-A315563

Coordination sequence Gal.5.320.2 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013910 | OEIS-A315575

Coordination sequence Gal.4.84.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Sep 14 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013911 | OEIS-A315598

Coordination sequence Gal.3.57.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013912 | OEIS-A315605

Coordination sequence Gal.6.577.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 12 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013913 | OEIS-A315615

Coordination sequence Gal.6.609.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

a(n) = 2*a(n-1) - 3*a(n-2) + 4*a(n-3) - 5*a(n-4) + 6*a(n-5) - 7*a(n-6) + 8*a(n-7) - 7*a(n-8) + 6*a(n-9) - 5*a(n-10) + 4*a(n-11) - 3*a(n-12) + 2*a(n-13) - a(n-14) for n > 14 (conjectured). G.f.: (x^14 + 4*x^13 + 3*x^12 + 5*x^11 + 5*x^10 + 6*x^9 + 7*x^8 + 13*x^7 + 7*x^6 + 6*x^5 + 5*x^4 + 5*x^3 + 3*x^2 + 4*x + 1)/((x - 1)^2*(x^2 + 1)^2*(x^4 + 1)^2) (conjectured). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013914 | OEIS-A315617

Coordination sequence Gal.3.37.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 29 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013915 | OEIS-A315628

Coordination sequence Gal.4.104.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 09 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013916 | OEIS-A315633

Coordination sequence Gal.6.261.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 13 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013917 | OEIS-A315636

Coordination sequence Gal.6.346.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 13 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013918 | OEIS-A315640

Coordination sequence Gal.6.556.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Mar 12 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013919 | OEIS-A315644

Coordination sequence Gal.6.325.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Mar 08 2025: (Start)
Conjecture: a(n) ~ 26*n/5. - _Stefano Spezia_, Mar 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013920 | OEIS-A315645

Coordination sequence Gal.6.633.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 31 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013921 | OEIS-A315660

Coordination sequence Gal.5.320.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 21 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013922 | OEIS-A315669

Coordination sequence Gal.5.113.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 14 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013923 | OEIS-A315672

Coordination sequence Gal.5.139.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 16 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013924 | OEIS-A315679

Coordination sequence Gal.6.252.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 15 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013925 | OEIS-A315681

Coordination sequence Gal.6.159.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Nov 15 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013926 | OEIS-A315683

Coordination sequence Gal.6.634.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 21 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013927 | OEIS-A315685

Coordination sequence Gal.4.140.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 25 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013928 | OEIS-A315687

Coordination sequence Gal.6.210.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 25 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013929 | OEIS-A315688

Coordination sequence Gal.6.343.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 25 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013930 | OEIS-A315690

Coordination sequence Gal.5.135.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 25 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013931 | OEIS-A315691

Coordination sequence Gal.5.306.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 25 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013932 | OEIS-A315693

Coordination sequence Gal.3.50.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 17 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013933 | OEIS-A315696

Coordination sequence Gal.5.301.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 25 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013934 | OEIS-A315697

Coordination sequence Gal.6.621.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 25 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013935 | OEIS-A315701

Coordination sequence Gal.4.136.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 25 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013936 | OEIS-A315707

Coordination sequence Gal.5.295.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 25 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013937 | OEIS-A315732

Coordination sequence Gal.6.640.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 03 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013938 | OEIS-A315738

Coordination sequence Gal.6.343.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 08 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013939 | OEIS-A315741

Coordination sequence Gal.6.210.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 15 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013940 | OEIS-A315748

Coordination sequence Gal.5.294.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jul 27 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013941 | OEIS-A315749

Coordination sequence Gal.5.295.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 06 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013942 | OEIS-A315767

Coordination sequence Gal.6.627.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 18 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013943 | OEIS-A315801

Coordination sequence Gal.3.43.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, May 24 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013944 | OEIS-A315817

Coordination sequence Gal.4.111.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Dec 28 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013945 | OEIS-A315823

Coordination sequence Gal.6.507.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 01 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013946 | OEIS-A315832

Coordination sequence Gal.5.207.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 10 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013947 | OEIS-A315850

Coordination sequence Gal.6.104.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 16 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013948 | OEIS-A315860

Coordination sequence Gal.4.12.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 25 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013949 | OEIS-A315869

Coordination sequence Gal.5.247.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jun 20 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013950 | OEIS-A315902

Coordination sequence Gal.6.282.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Mar 11 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013951 | OEIS-A315907

Coordination sequence Gal.3.40.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _R. J. Mathar_, May 02 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013952 | OEIS-A315913

Coordination sequence Gal.6.599.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Apr 13 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013953 | OEIS-A315942

Coordination sequence Gal.6.507.5 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Jan 09 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013954 | OEIS-A315950

Coordination sequence Gal.3.47.3 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 03 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013955 | OEIS-A316022

Coordination sequence Gal.6.377.6 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Aug 01 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013956 | OEIS-A316068

Coordination sequence Gal.5.296.4 where Gal.u.t.v denotes the coordination sequence for a vertex of type v in tiling number t in the Galebach list of u-uniform tilings.

Conjectures from _Chai Wah Wu_, Nov 18 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Galebach_ and _N. J. A. Sloane_, Jun 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013957 | OEIS-A316111

a(n) is the smallest k > 1 such that gcd(k, n^k - n) = 1, for n > 1.

Conjecture: all the terms are in A121707. If k is a term, then k is an "anti-Carmichael number": p-1 does not divide k-1 for every prime p dividing k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Aug 13 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013958 | OEIS-A316141

Number of ordered pairs (k, m) with k > 1 and m > 0 such that n - F(k)*L(m) is an odd prime, where F(k) is the k-th Fibonacci number (A000045) and L(m) is the m-th Lucas number (A000204).

Conjecture: a(n) > 0 for all n > 3. In other words, any integer n > 3 can be written as p + F(k)*L(m), where p is an odd prime, and k and m are positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013959 | OEIS-A316142

Expansion of e.g.f. Product_{k>=1} (1 + (exp(x)-1)^k)^2.

Conjecture: Let k be a positive integer. The sequence obtained by reducing a(n) modulo k is eventually periodic with the period dividing phi(k) = A000010(k). For example, modulo 7 we obtain the sequence [1, 2, 1, 0, 0, 2, 2, 2, 1, 0, 0, 2, 2, 2, 1, 0, 0, 2, 2, ...], with a preperiod of length 1 and an apparent period thereafter of 6 = phi(7). - _Peter Bala_, Mar 03 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jun 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013960 | OEIS-A316144

Expansion of e.g.f. Product_{k>=1} ((1 + (exp(x)-1)^k) / (1 - (exp(x)-1)^k))^2.

Conjecture: Let k be a positive integer. The sequence obtained by reducing a(n) modulo k is eventually periodic with the period dividing phi(k) = A000010(k). For example, modulo 7 we obtain the sequence [1, 4, 0, 2, 5, 3, 2, 4, 0, 2, 5, 3, 2, 4, 0, 2, 5, 3, 2, ...], with a preperiod of length 1 and an apparent period thereafter of 6 = phi(7). - _Peter Bala_, Mar 03 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jun 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013961 | OEIS-A316189

Decimal expansion of Sum(1/p + 1/q) as (p, q) runs through the twin m^2 + 1 primes.

Conjecture: the series of all twin m^2 + 1 prime reciprocals converges to 0.357745147...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jun 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013962 | OEIS-A316316

Coordination sequence for tetravalent node in chamfered version of square grid.

From _N. J. A. Sloane_, Jun 30 2018: This conjecture is true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 29 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013963 | OEIS-A316317

Coordination sequence for trivalent node in chamfered version of square grid.

G.f. (assuming above conjecture): (1+x)^2*(1+3*x^2+x^4)/((1-x)^2*(1+x+x^2)*(1+x^2)). - _Robert Israel_, Jul 01 2018 a(n) = (30*n - 9*A056594(n-1) + 6*A102283(n))/9 for n > 0. - Conjectured by _Stefano Spezia_, Jun 12 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 29 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013964 | OEIS-A316357

Partial sums of A316316.

Conjectures from _Colin Barker_, Dec 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jun 30 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013965 | OEIS-A316460

Even integers not of the form prime(prime(x)) + prime(prime(y)) with x > y > 0.

Conjecture: Define an m-(prime-index-prime) as having "m" prime-on-prime iterations, For any m >= 0 and n >= 0, all sufficiently large even numbers are the sum of an m-(prime-index-prime) and an n-(prime-index-prime). See links.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrei-Lucian Dragoi_, Jul 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013966 | OEIS-A316472

Inverse permutation to A316385.

Conjecture: a(n) = (2*b(n) + 1)*2^(L(n) - L(b(n)) - 1) where b(n) = A053645(A153142(n)) and where L(n) = A000523(n) for n > 0 with L(0) = -1. - _Mikhail Kurkov_, Sep 09 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jul 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013967 | OEIS-A316584

Array read by antidiagonals: T(n,k) is the number of elements x in GL(2,Z_n) with x^k == I mod n where I is the identity matrix.

Conjecture: T(p,p) = p^2 for p prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Howroyd_, Jul 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013968 | OEIS-A316586

Array read by antidiagonals: T(n,k) is the number of elements x in SL(2,Z_n) with x^k == I mod n where I is the identity matrix.

Conjecture: T(p,p) = p^2 for p prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Howroyd_, Jul 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013969 | OEIS-A316676

Ordered set of products (P_s(k) + 1)*(P_s'(k') + 1), s,s' >= 3, k,k' in {3,4} with nontrivial polygonal numbers P_s(k).

Conjecture: All odd numbers d >= 17 excluding d in {P_s(k), s >= 3, k >= 5; P_s(k) - 1, s >= 3, k >= 4; a(n)} are accurate the primes p = d >= 17.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Steiner_, Jul 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013970 | OEIS-A316678

Smallest numbers leading in n steps to a term that repeats itself, according to the rule explained in A316650 (and hereunder in the Comment section).

It is conjectured, when iterating the idea explained in A316650 ("Result when n is divided by the sum of its digits and the resulting integer is concatenated to the remainder"), that all integers will end either on a fixed point (the first ones are listed in A052224) or grow forever [10 is an example of a simple pattern: 10,100,1000,10000,100000,. . .]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Jul 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013971 | OEIS-A316679

The integer 907 and its infinite growing pattern (when iterating the rule explained in A316650 and hereunder, in the Comment section).

It is conjectured, when iterating the idea explained in A316650 ("Result when n is divided by the sum of its digits and the resulting integer is concatenated to the remainder"), that all integers will end either on a fixed point (the first ones are listed in A052224) or grow forever (like 907 or 1358).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Jul 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013972 | OEIS-A316680

The integer 1358 and its infinite continuation (when iterating the rule explained in A316650 and in the Comment section here).

It is conjectured, when iterating the idea explained in A316650 ("Result when n is divided by the sum of its digits and the resulting integer is concatenated with the remainder"), that all integers will end either on a fixed point (the first ones are listed in A052224) or grow forever (like 907 or 1358).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Jul 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013973 | OEIS-A316833

Sums of four distinct odd squares.

Theorem (Conjectured by R. William Gosper, proved by M. D. Hirschhorn): Any sum of four distinct odd squares is the sum of four distinct even squares.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013974 | OEIS-A316908

a(n) is the smallest k with n prime factors such that 2^(k-1) == 1 (mod k) and p-1 does not divide k-1 for every prime p dividing k.

Conjecture: a(n) > A006931(n) for every n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Jul 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013975 | OEIS-A316918

For any k, the cumulative sum a(1) + a(2) + a(3) + ... + a(k) starts with a(k). Lexicographic first sequence of positive integers without duplicate terms having this property and extending itself forever.

The authors conjecture that it is impossible to extend the sequence forever if a(1) < 61.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Jul 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013976 | OEIS-A316942

a(n) = n - a(pi(n)) - a(n-pi(n)) with a(1) = a(2) = 1, where pi = A000720.

Conjecture : lim_{n->infinity} a(n)/n = 1/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Jul 17 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013977 | OEIS-A316971

Prime numbers of the form p1^3 + p2^2 + p3, with p1, p2 and p3 also prime and all different.

Conjecture: the only primes not in the sequence are

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierandrea Formusa_, Jul 17 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013978 | OEIS-A316973

a(n) is the number of occurrences of n in A316774.

These values may be only conjectural, in which case the b-file should be changed to an a-file and marked as conjectured. - _N. J. A. Sloane_, Dec 14 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 17 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013979 | OEIS-A316976

Numbers k such that some of the values (r0-r1+k) mod k for all pairs (r0,r1) of quadratic residues mod k are unique.

It is conjectured that this list is finite and limited to the terms given in the DATA section.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arnauld Chevallier_, Jul 17 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013980 | OEIS-A316984

a(n) is the index of the last occurrence of n in A316774.

[It may be that these values are only conjectural. If so, the b-file should be changed to an a-file and marked as conjectured. - _N. J. A. Sloane_, Dec 14 2019]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013981 | OEIS-A316993

Lexicographically earliest sequence of distinct positive terms such that a(1) = 2 and for any n > 0, the binary representation of Sum_{k=1..n} a(k) starts with the binary representation of a(n).

This sequence is conjectured to be infinite. This sequence is conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jul 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013982 | OEIS-A317023

Square array A(n,k), n >= 0, k >= 0, read by ascending antidiagonals, where the sequence of row n is the expansion of (1-x^(n+1))/((1-x)^(n+1)).

Conjecture: alternating row sums of the triangle give A106510 for n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Jul 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013983 | OEIS-A317175

Square array T(n, k) read by antidiagonals upwards, n > 0 and k > 0: T(n, k) is the least m > 0 such that m * n contains k as a substring in its decimal representation.

- the conjectured maximum values for the first 9 columns are: 5, 12, 17, 32, 25, 24, 35, 32, 72.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jul 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013984 | OEIS-A317190

Main diagonal of infinite Sudoku-type array A274528.

Conjectured to be a permutation of the nonnegative numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 30 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013985 | OEIS-A317193

First differences of A317192.

Conjecture: This is an encoded version of the ternary tribonacci word

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 31 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013986 | OEIS-A317297

a(n) = (n - 1)*(4*n^2 - 8*n + 5).

Conjecture: For n > 1, a(n) is the maximum eigenvalue of a 2*(n-1) X 2*(n-1) square matrix M defined as M[i,j,n] = j + n*(i-1) if i is odd and M[i,j,n] = n*i - j + 1 if i is even (see A317614). - _Stefano Spezia_, Dec 27 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Sep 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013987 | OEIS-A317345

E.g.f. A(x) satisfies: [x^n] exp(n^3*x) / A(x)^(n^2) = 0 for n >= 1.

Periodic modulo 10: a(5*n+k) = [1,1,5,5,5](k) (mod 10), for n>=0 and k = 0..4 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013988 | OEIS-A317518

Highly composite numbers k such that ceiling(sqrt(k))^2 - k is a square.

Conjecture: this sequence is finite, with a(120) = 76750440345035675071229376000 as its final term. - _Jon E. Schoenfield_, Aug 12 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaughn R Tiffany_, Jul 30 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013989 | OEIS-A317534

Numbers k such that the poset of factorizations of k, ordered by refinement, is not a lattice.

Conjecture: Let S be the set of all numbers whose prime signature is either {1,3}, {5}, or {1,1,2}. Then the sequence consists of all multiples of elements of S. - _David A. Corneth_, Jul 31 2018.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jul 30 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013990 | OEIS-A317551

Fertility numbers.

It is conjectured that there are infinitely many positive integers that are not fertility numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Defant_, Sep 14 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013991 | OEIS-A317587

a(n) is the smallest number m > n such that Sum_{k=1..n-1} k^(m-1) == n-1 (mod m).

Conjecture: If p is a prime, then odd a(p) is the next prime after p. - _Thomas Ordowski_, Aug 06 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Aug 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013992 | OEIS-A317625

a(n) = Sum_{k=1..n} phi(floor(n/k)) where phi is the Euler totient function.

Cautious conjecture: a(n) ~ n*log(n)/zeta(2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Randell G Heyman_, Aug 02 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013993 | OEIS-A317688

Absolute primes that are not repunits: primes where the number resulting from any permutation of the digits is also prime, excluding repunit primes.

Conjecture: The sequence is finite, with 991 being the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Aug 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013994 | OEIS-A317689

Largest nonrepunit base-n absolute prime (conjectured).

Largest nonrepunit base-n absolute prime (conjectured). These primes were found using lim=10^8 in the PARI program and match those found with lim=10^5, lim=10^6 and lim=10^7. Therefore I conjecture that they are the correct values for those n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Aug 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013995 | OEIS-A317800

G.f. A(x) satisfies: Sum_{n>=1} (A(x) - (-1)^n * A(-x))^n * (-1)^n / n = 0.

Odd terms occur at a(2^k - 1) and a(2^k - 2) for k > 1 and at a(1), while a(n) is even elsewhere (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013996 | OEIS-A317849

Number of states of the Finite State Automaton Gn accepting the language of maximal (or minimal) lexicographic representatives of elements in the positive braid monoid An.

Conjecture: g.f. -x*(1-x+x^2) / ((x^2-3*x+1)*(x-1)^3). a(n) = 2*A001519(n+1) -n*(n+1)/2 -2 = 2*A001519(n+1)-A152948(n+2). - _R. J. Mathar_, Aug 17 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Aug 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013997 | OEIS-A317850

The 10-adic integer x = ...603097394847873 satisfying x^7 + 1 = y and y^7 + 1 = x.

Data generated using calculator (first 15 terms) and MATLAB (next 85 terms). Conjecture: There exists a pair of 10-adic integers satisfying x^n + 1 = y and y^n + 1 = x iff n == 3, 7, or 15 (mod 20).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick A. Thomas_, Sep 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013998 | OEIS-A317851

Primes p such that A090086(p+1) < p.

Conjecture: the sequence contains all sufficiently large primes. Probably all primes except 2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 37, 41, 47, 53, 59, 71, 83, 89, 101, 109, 139, 149, 157, 179, 199, 307, 461, and 571.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Aug 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-013999 | OEIS-A317932

Denominators of certain "Dirichlet Square Root" sequences: a(n) = A046644(n)/(2^A007949(n)).

The original definition was: Denominators of the rational valued sequence whose Dirichlet convolution with itself yields A002487, Stern's Diatomic sequence. However, this definition depends on the conjecture given in A261179. a(n) = denominator of f(n), where f(1) = 1, f(n) = (1/2) * (b(n) - Sum_{d|n, d>1, d<n} f(d) * f(n/d)) for n > 1, where b can be A064989, A065769 or A234840 for example, conjecturally also A002487.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014000 | OEIS-A318008

G.f. A(x) satisfies: A(A(x - x^2)) = x + x^2.

a(2^k-1) = 1 (mod 2) and a(2^(k+1)-2) = 1 (mod 2) for k >= 1, and a(n) is even elsewhere (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 06 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014001 | OEIS-A318135

The 10-adic integer a = ...1588948901 satisfying a^2 + 1 = b, b^2 + 1 = c, c^2 + 1 = d, d^2 + 1 = e, e^2 + 1 = f, and f^2 + 1 = a.

Conjecture: Let r(k) = the smallest positive residue of A003095(6*k+1) mod 10^(6*k+1). Then the first 2*k + 2 digits of r(k), reading from right to left, give the first 2*k + 2 digits of this 10-adic number. For example with k = 5, r(k) = 2121286728960294(201588948901) gives the first 12 digits correctly. - _Peter Bala_, Nov 14 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick A. Thomas_, Aug 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014002 | OEIS-A318239

a(n) is the smallest primitive root of A133433(n).

If A023048(108) != 0 (which is implied assuming generalized Artin's conjecture) then the next term of this sequence is 108. Conjecturally 11 is the largest prime in this sequence, 42 is the largest squarefree term, and 45 is the largest odd term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014003 | OEIS-A318245

Scaled g.f. T(v) = Sum_{n>=0} a(n)*(3*v/64)^n satisfies 9*(5*v-4)*T + d/dv(16*v*(v-1)*(3*v-4)*T') = 0, and a(0)=1; sequence gives a(n).

For n > 0, a(n) mod 3 = 0 (conjecture, tested up to n=3*10^6).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bradley Klee_, Aug 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014004 | OEIS-A318246

a(n) = [x^n] Product_{k>=1} (1 + 3^n*x^k).

Conjecture: In general, if m > 1 and a(n) = [x^n] Product_{k>=1} (1 + m^n * x^k), then log(a(n)) ~ log(m)*(sqrt(2)*n^(3/2) - n/2). Conjecture: log(a(n)) ~ log(3)*sqrt(2)*n^(3/2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Aug 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014005 | OEIS-A318268

a(n) is the number of configurations of n indistinguishable pairs placed on the vertices of the ladder graph P_2 X P_n such that all but 3 such pairs are joined by an edge.

G.f.: x^2*(2*x + 20*x^2 + 46*x^3 + 40*x^4 + 30*x^5 + 4*x^6 + 4*x^7)/(1 - x)^3/(1 - x - x^2)^4 (conjectured). The above conjecture is true. The PARI program given in the links can be used to establish an upper limit on the order of the linear recurrence and sufficient number of terms to prove correctness. - _Andrew Howroyd_, Sep 03 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Donovan Young_, Aug 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014006 | OEIS-A318269

a(n) is the number of configurations of n indistinguishable pairs placed on the vertices of the ladder graph P_2 X P_n such that all but 4 such pairs are joined by an edge.

G.f.: x^2*(5*x^10 + 10*x^9 + 93*x^8 + 230*x^7 + 502*x^6 + 612*x^5 + 447*x^4 + 158*x^3 + 21*x^2)/(1 - x)^4/(1 - x - x^2)^5 (conjectured). The above conjecture is true. See A318268. - _Andrew Howroyd_, Sep 03 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Donovan Young_, Aug 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014007 | OEIS-A318270

a(n) is the number of configurations of n indistinguishable pairs placed on the vertices of the ladder graph P_2 X P_n such that all but 5 such pairs are joined by an edge.

G.f.: x^2*(6*x^13 + 20*x^12 + 228*x^11 + 888*x^10 + 3012*x^9 + 6612*x^8 + 10020*x^7 + 9636*x^6 + 5502*x^5 + 1620*x^4 + 186*x^3)/(1 - x)^5/(1 - x - x^2)^6 (conjectured). The above conjecture is true. See A318268. - _Andrew Howroyd_, Sep 03 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Donovan Young_, Aug 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014008 | OEIS-A318291

a(n) is the minimum k > 0 such that n*2^k - 3 is prime, or 0 if no such k exists.

Conjecture: For every odd prime p, there exist infinitely many numbers j that are non-multiples of p and have the property that j*2^k - p is composite for every k > 0. - _Martin Michael Musatov_, Sep 04 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Martin Michael Musatov_, Aug 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014009 | OEIS-A318314

Denominators of the sequence whose Dirichlet convolution with itself yields A068068, number of odd unitary divisors of n.

From _Antti Karttunen_, Sep 03-07 2018: (Start, conjectured formulas)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_ and _Andrew Howroyd_, Aug 29 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014010 | OEIS-A318351

a(n) is the smallest prime p such that the sum of the first 2*n + 1 odd primes starting with p is prime.

Conjecture: Sequence is bounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Aug 24 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014011 | OEIS-A318413

Expansion of Product_{i>=1, j>=1, k>=1} 1/(1 - x^(i*j*k))^(i*j*k).

Conjecture: log(a(n)) ~ (3*Zeta(3))^(1/3) * log(n)^(2/3) * n^(2/3) / 2. - _Vaclav Kotesovec_, Sep 02 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Aug 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014012 | OEIS-A318414

Expansion of Product_{i>=1, j>=1, k>=1} (1 + x^(i*j*k))^(i*j*k).

Conjecture: log(a(n)) ~ 3^(2/3) * Zeta(3)^(1/3) * log(n)^(2/3) * n^(2/3) / 2^(5/3). - _Vaclav Kotesovec_, Sep 02 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Aug 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014013 | OEIS-A318487

Even numbers that are not the sum of two lucky-indexed lucky numbers.

Conjecture (checked up to 10^7): every even number larger than 199254 is the sum of two lucky-indexed lucky numbers, so this sequence is finite with 1714 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Aug 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014014 | OEIS-A318495

Scaled g.f. T(u) = Sum_{n>=0} a(n)*(u/16)^n satisfies 5*(21*u-16)*T + d/du(4*u*(u-1)*(27*u-32)*T') = 0, and a(0)=1; sequence gives a(n).

For n > 0, a(n) mod 10 = 0 (conjecture, tested up to n=10^6).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bradley Klee_, Aug 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014015 | OEIS-A318496

Scaled g.f. T(v) = Sum_{n>=0} a(n)*(v/16)^n satisfies 15*(189*v-80)*T + d/dv(4*v*(27*v-5)*(27*v-32)*T') = 0, and a(0)=1; sequence gives a(n).

For n > 0, a(n) mod 30 = 0 (conjecture, tested up to n=10^6).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bradley Klee_, Aug 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014016 | OEIS-A318509

Completely multiplicative with a(p) = A002487(p).

Provided that the conjecture given in A261179 holds, then for all n >= 1, A007814(a(n)) = A007949(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 30 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014017 | OEIS-A318510

Completely multiplicative with a(prime(k)) = A002487(prime(k+1)).

Provided that the conjecture given in A261179 holds, then for all n >= 1, A007814(a(n)) = A007814(n), i.e., then the sequence preserves the 2-adic valuation of n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 30 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014018 | OEIS-A318698

a(n) is the smallest nonnegative integer of the same parity as n, not yet in the sequence, that shares a digit with a(n-1); a(0)=0.

Conjecture: This is a permutation of the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enrique Navarrete_, Aug 31 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014019 | OEIS-A318699

a(n) is the smallest nonnegative integer of opposite parity to n, not yet in the sequence, that shares a digit with a(n-1); a(0) = 1.

Conjecture: this is a permutation of the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enrique Navarrete_, Aug 31 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014020 | OEIS-A318712

Numbers k for which a riffle shuffle permutation with two or more packets permutes all except the first and the last of the k cards.

If the formula is true then Tiago Januario's conjecture on A217948 would be solved (see formula).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mauro Rigo_, Sep 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014021 | OEIS-A318764

Expansion of Product_{i>=1, j>=1, k>=1} ((1 + x^(i*j*k))/(1 - x^(i*j*k)))^(i*j*k).

Conjecture: log(a(n)) ~ (21*Zeta(3))^(1/3) * log(n)^(2/3) * n^(2/3) / 2^(5/3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Sep 03 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014022 | OEIS-A318778

Number of different positions that an elementary sphinx can occupy in a sphinx of order n.

Conjectures from _Colin Barker_, Nov 13 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Craig Knecht_, Sep 10 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014023 | OEIS-A318807

Numbers whose sum of squarefree divisors and sum of nonsquarefree divisors are both perfect squares.

The subsequence c(n) where the ratio r = (sigma(a(n)) - s)/s is an integer begins with 27, 88, 264, 280, 376, 594, 680, 840, 856, 1128, 1240, ... and the corresponding r are 3^2, 2^2, 2^2, 2^2, 2^2, 3^2, 2^2, 2^2, 2^2, 2^2, 2^2, 2^2, 2^2, 5^2, 3^2, 2^2, 7^2, 3^2, 2^2, 11^2, ... It is conjectured that r belongs to A001248.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Sep 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014024 | OEIS-A318858

Number of still-lifes synthesizable from n gliders in Conway's game of Life.

Most of the terms in this sequence are conjectural and will likely increase over time. - _Nathaniel Johnston_, Dec 03 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ed Pegg Jr_, Sep 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014025 | OEIS-A318921

In binary expansion of n, delete one symbol from each run. Set a(n)=0 if the result is the empty string.

Conjecture: For n in the range 2^k, ..., 2^(k+1)-1, the total value of a(n) appears to be 2*3^(k-1) - 2^(k-1) (see A027649), and so the average value of a(n) appears to be (3/2)^(k-1) - 1/2. - _N. J. A. Sloane_, Sep 25 2018 The above conjecture was proved by _Doron Zeilberger_ on Nov 16 2018 (see link) and independently by _Chai Wah Wu_ on Nov 18 2018 (see below). - _N. J. A. Sloane_, Nov 20 2018 Conjecture is correct for k > 0. Proof: looking at the least significant 2 bits of n, it is easy to see that a(4n) = 2a(2n), a(4n+1) = a(2n), a(4n+2) = a(2n+1) and a(4n+3) = 2a(2n+1)+1. Define f(k) = Sum_{i=2^k..2^(k+1)-1} a(i), i.e. the sum ranges over all numbers with a (k+1)-bit binary expansion. Thus f(0) = a(1) = 0 and f(1) = a(2)+a(3) = 1. By summing over the recurrence relations for a(n), we get f(k+2) = Sum_{i=2^k..2^(k+1)-1} (f(4i) + f(4i+1) + f(4i+2) + f(4i+3)) = Sum_{i=2^k..2^(k+1)-1} (3a(2i) + 3a(2i+1) + 1) = 3*f(k+1) + 2^k. Solving this first order recurrence relation with the initial condition f(1) = 1 shows that f(k) = 2*3^(k-1)-2^(k-1) for k > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014026 | OEIS-A318946

Column 1 of triangle A318945.

Conjectures from _Colin Barker_, Oct 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014027 | OEIS-A318947

Column 2 of triangle A318945.

Conjectures from _Colin Barker_, Oct 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014028 | OEIS-A318988

Totients whose inverses contain two or more consecutive tetrahedral pyramidal numbers.

Conjecture: There are at most three consecutive tetrahedral pyramidal numbers in the set of inverses of a totient number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Torlach Rush_, Sep 06 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014029 | OEIS-A319012

a(n) = Sum_{i=1..n} prime(n*(i - 1) + i).

Conjecture: a(2) and a(3) are the only terms that are perfect squares.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Sep 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014030 | OEIS-A319018

Number of ON cells after n generations of two-dimensional automaton based on knight moves (see Comments for definition).

The conjecture that A322050(2^k+1)=1 also suggests a fractal geometry. Let P_k be the associated set of eight points. It appears that P_k may be written as the intersection of four fixed lines, y = +-2*x and x = +-2*y, with a circle, x^2 + y^2 = 5*4^k (see linked image "Log-Periodic Coloring"). - _Bradley Klee_, Dec 16 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Sep 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014031 | OEIS-A319033

a(n) is the (conjectured) largest number k that is zeroless in every base b such that n <= b < k.

a(n) is the (conjectured) largest number k that is zeroless in every base b such that n <= b < k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Oct 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014032 | OEIS-A319036

a(n) is the smallest triangular number T(k) such that both it and its successor T(k+1) have exactly 2n divisors, or 0 if no such pair of consecutive triangular numbers exists.

Conjecture: The only primes p for which a(p) > 0 are 2 and 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Dec 05 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014033 | OEIS-A319053

a(n) is the exponent of the largest power of 2 that appears in the factorization of the entries in the matrix {{3,1},{1,-1}}^n.

The above conjecture is true. From the preceding formulas for the elements of M^n this claims that the Fibonacci numbers F(n-3), F(n) and F(n+3) are always odd for n == 1 or 2 (mod 3). This is true because F(n) is even iff n == 0 (mod 3) (see e.g. Vajda, p.73), and each of the three indices is == 1 or 2 (mod 3) for n == 1 or 2 (mod 3), respectively. - _Wolfdieter Lang_, Oct 09 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Greg Dresden_, Sep 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014034 | OEIS-A319116

Signs of the Maclaurin coefficients of 1/(exp(x) + Pi/2).

Conjecture: a(n) is the same as the sign of cos((n+1)*arctan(Pi/log(Pi/2))). We can also use arctan(Pi/log(Pi/2))/Pi ~ 1/2 ~ 5/11 ~ 3776/8307 ~ 3781/8318 and so on. If the numerator is odd, we have, for example, a(n) = -a(n+2) or a(n) = -a(n+11) with some counterexamples. For even numerators we have, for example, a(n) = a(n+8307), also with some counterexamples.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Sep 10 2018 [verification needed] | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014035 | OEIS-A319120

T(n, k) = binomial(n - k - 1, k)*binomial(2*n - 2*k, n)/(n + 1), for n >= 1 and 0 <= k <= floor((n - 1)/2), triangle read by rows.

Let L(u,v) be the set of integer partitions whose Young diagrams fit inside a u by v rectangle. Given lambda in L(u,v), let E(lambda) be the number of partitions whose Young diagrams fit inside the Young diagram of lambda. Also, for 1 <= i <= v, let x_i(lambda)-1 be the number of parts of lambda of length v+1-i. Let x_{v+1}(lambda) = u+v+1-Sum_{i=1..v} x_i(lambda) so that (x_1(lambda), ..., x_{v+1}(lambda)) is a composition of u+v+1 into v+1 parts. Let F(lambda) = Product_{i=1..v+1} Catalan(x_i(lambda)). Conjecturally, T(n,k) = Sum_{lambda in L(n-2k-1)} E(lambda) * F(lambda). Conjecturally, T(n,k) is the number of permutations pi of [n] such that s(pi) has k descents and avoids the patterns 231, 312, and 321, where s is West's stack-sorting map. Conjecturally, T(n,k) is the number of permutations pi of [n] that avoid the 4 patterns 4312, 4321, 4231, 3241 (more succinctly, that avoid 32x1 for all x) and contain k entries which are neither left-right maxima nor right-left minima (equivalently, contain k entries that serve as the "2" of a 321). - _David Callan_, Mar 05 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Defant_, Sep 17 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014036 | OEIS-A319171

Square array, read by antidiagonals, upwards: T(n,k) is the number of groups of order prime(k+1)^n.

In 1960, Higman conjectured that the function f(n,p) giving the number of groups of prime-power order p^n, for fixed n and varying p, is a "Polynomial in Residue Classes" (PORC), i.e., there exist an integer M and polynomials q_i(x) in Z[x] (i = 1, 2, ..., M) such that if p = i mod M, then f(n,p) = q_i(p). The conjecture is confirmed for n <= 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franck Maminirina Ramaharo_, Sep 12 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014037 | OEIS-A319197

All entries from a(3) to a(n) appear in addition to 2^n as factors in the conjectured factorization of Fibonacci(2^(n-2)*3*m) for n >= 3 and all m >= 0.

All entries from a(3) to a(n) appear in addition to 2^n as factors in the conjectured factorization of Fibonacci(2^(n-2)*3*m) for n >= 3 and all m >= 0. It appears that Fibonacci(2^(n-2)*3*m)/(2^n) is a nonnegative integer for n >= 3 and all m >= 0. The remaining factor for n >= 3 is given by A(n) := Product_{j=3..n} a(j). For n = 3 and 4 see A049660 and A253368, respectively. Thus the conjecture is that I(n; m) := Fibonacci(2^(n-2)*3*m) / ((2^n)*A(n)) is a nonnegative integer for all m >= 0. This is best possible for all m (because for m = 1 this becomes 1); special m may allow more factors. E.g., n = 3: 8 | Fibonacci(6*m), for all m >= 0, but for even m larger powers of 2 than 2^3 appear; for m = 0 any power of 2. I(n; m) := F(2^(n-2)*3*m) / ((2^n)* Product_{j=3..n} a(j)) is conjectured to be a nonnegative integer for n >= 3 and all m >= 0, where F = A000045. There are no more factors > 1 for all m >= 0 because I(n, 1) = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014038 | OEIS-A319204

Sequence used for the Boas-Buck type recurrence for Riordan triangle A319203.

a(n) = (1/(n+1)!)*[d^(n+1)/dx^(n+1) (1 - x^2 - x^3)^(n+1)] evaluated at x = 0, for n >= 0. (Cf. _Joerg Arndt_'s conjecture for A176806, which is proved there.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 29 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014039 | OEIS-A319248

Lesser of the pairs of twin primes in A001122.

Note that "there are infinitely many pairs of twin primes" and "there are infinitely many primes with primitive root 2" are two famous and unsolved problems, so a stronger conjecture implying both of them is that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Sep 15 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014040 | OEIS-A319316

Numbers k such that A090616(k) < A054861(k).

Note that A054861 and A090616 are both asymptotic to a(n) = n/2 + O(log(n)), nevertheless, it seems that the number of k such that A090616(k) is bigger predominates. Conjecture: the ratio of k <= N such that A090616(k) > A054861(k) tends to 1 as N tends to infinity, while the ratio of k <= N such that A090616(k) < A054861(k) and A090616(k) = A054861(k) both tend to 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Sep 17 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014041 | OEIS-A319357

Filter sequence combining A003415(d) from all proper divisors d of n, where A003415(d) = arithmetic derivative of d.

a(i) = a(j) => A101296(i) = A101296(j). [Conjectural, see notes above]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 02 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014042 | OEIS-A319371

Numbers k such that the characteristic polynomial of a wheel graph of k nodes has exactly one monomial with vanishing coefficient.

Conjecture: a(n) = A004772(n) for n> 1. [clarified by _Michel Marcus_, Apr 16 2019] Conjectures from _Colin Barker_, Nov 02 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierandrea Formusa_, Sep 17 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014043 | OEIS-A319421

Triangle read by rows: T(n,k) (1 <= k <= n) = one-half of the number of binary vectors of length n and cuts-resistance k.

Conjecture (Sloane): Sum_{i = 1..n} i * T(n,i) = A189391(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014044 | OEIS-A319424

Records in A175046 (but written in base 2).

Conjecture. a(n) is either an alternating string of 11's and 00's, or exactly one of the 00's is replaced by 000. Conjecture is true (see A319423). - _Chai Wah Wu_, Oct 04 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014045 | OEIS-A319430

First differences of the tribonacci representation numbers (A003726 or A278038).

Conjecture: All terms are of the form ceiling(2^k/7) for some k (cf. A046630), and all numbers of the form ceiling(2^k/7) occur. Conjecture (continued): Furthermore, new values of ceiling(2^k/7) (that is, new records) appear at n = 0, 6,12, 23, 43, 80, 148, 273, ..., which (apart from the start) are the tribonacci numbers minus 1, A000073 - 1, or A089068.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 30 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014046 | OEIS-A319476

a(n) is the minimum number of distinct distances between n non-attacking rooks on an n X n chessboard.

Conjecture: a(n^2) = A047800(n-1) - 1. - _Peter Kagey_, Nov 02 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Oct 12 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014047 | OEIS-A319480

Number of ordered pairs (i,j) with 0 < i < j < prime(n)/2 such that R(i^2,prime(n)) > R(j^2,prime(n)), where R(k,p) (with p an odd prime and k an integer) denotes the unique integer r among 0,1,...,(p-1)/2 for which k is congruent to r or -r modulo p.

Conjecture: Let p be any odd prime and let N(p) be the number of ordered pairs (i,j) with 0 < i < j < p/2 and R(i^2,p) > R(j^2,p). Then N(p) == floor((p+1)/8) (mod 2). See also A319311 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014048 | OEIS-A319506

Number of numbers of the form 2*p or 3*p between consecutive triangular numbers T(n - 1) < {2,3}*p <= T(n) with p prime.

1) It is conjectured that for k >= 1 each left-sided half-open interval (T(2*k - 1), T(2*k + 1)] and (T(2*k), T(2*(k + 1))] contains at least one composite c_2 = 2*p_i and c_3 = 3*p_j each, p_i, p_j prime, i != j. 2) It is conjectured that for k >= 3 each left-sided half-open interval (T(k - 1), T(k)] contains at least one composite c_2 = 2*p_i or c_3 = 3*p_j, p_i, p_j prime, i != j. 3) It is conjectured that for k >= 2 each left-sided half-open interval (T(2*k - 1), T(2*k)] contains at least one composite c_3 = 3*p_j, p_j prime. 4) It is conjectured that for k >= 1 each left-sided half-open interval (T(2*k), T(2*k + 1)] contains at least one composite c_2 = 2*p_i, p_i prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ralf Steiner_, Sep 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014049 | OEIS-A319515

Second term of the simple continued fraction of (Sum_{k=1..n} k^k)/(n^n).

Ignoring the first two terms, a(n+1) is very similar to floor(n*e) (the Beatty sequence for e, A022843). This breaks down at n=11, where a(12)=30 and floor(11*e)=29. The successive terms (or partial denominators) in the sequence differ by 3 except for semi-regular "glitches". For example, the pattern of differences is 3132333233323332332333233

233323332332333233233323332333233233233233332... A natural conjecture is that the mean of these differences is asymptotic to e. The glitches are positioned between terms n and n+1, for values of n which begin 2,4,8,12,16,19,23,26,30,34,37,41,44,48,.... If one counts the 3's immediately preceding the 2's in the pattern of differences, one gets the "derivative" sequence 233323233232323323....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Greg Huber_, Sep 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014050 | OEIS-A319524

a(n) is the smallest number that belongs simultaneously to the two arithmetic progressions prime(n) + m*prime(n+1) and prime(n+1) + m*prime(n+2), m >= 1, n >= 1.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexandra Hercilia Pereira Silva_, Sep 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014051 | OEIS-A319610

a(n) is the minimal number of successive OFF cells that appears in n-th generation of rule-30 1D cellular automaton started from a single ON cell.

G.f.: x (x + x/(1 - x) + x^3 + x^5 + x^7) (conjectured). It is conjectured that for all n>=10, a(n)=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philipp O. Tsvetkov_, Sep 24 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014052 | OEIS-A319658

a(n) is the minimal number of successive ON cells that appears in n-th generation of rule-30 1D cellular automaton started from a single ON cell.

G.f.: 1/(1 - x) + 2 x + x^3 + x^5 + x^7 + x^13 (conjectured). It is conjectured that for all n >= 15, a(n)=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philipp O. Tsvetkov_, Sep 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014053 | OEIS-A319674

a(n) = 1 + 2 + 3 - 4 - 5 - 6 + 7 + 8 + 9 - 10 - 11 - 12 + ... - (up to n).

Conjectures from _Bill McEachen_, Dec 19 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Sep 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014054 | OEIS-A319687

a(n) = A318509(n) - A002487(n).

All terms seem to be even. See the conjecture given in A261179.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 02 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014055 | OEIS-A319798

Smallest fixed points (>0) of the base-n Kaprekar map.

Conjecture: If n = 3*k - 1 (>2), a(n) = A000567(k). For example, a(29) = 10 * (3*10 - 2) = 280.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Sep 28 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014056 | OEIS-A319800

Numbers k such that Sum_{d|k} nphi(d) = k where the sum is over nonunitary divisors of k and nphi(k) is the nonunitary totient function (A254503).

Ligh and Wall found the first 6 terms and also the terms a(8) = 341863562880, a(9) = 357174165248, a(11) = 1018887932160, and 20993596382889043200. They showed that each term has a powerful part with at least 2 distinct prime factors, and conjectured that it is only even.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Sep 28 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014057 | OEIS-A319824

Numbers that are not the sum of distinct lucky-indexed lucky numbers.

Conjecture: this sequence is finite with the last term being a(98) = 373.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Sep 28 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014058 | OEIS-A319894

Number of ordered pairs (i,j) with 0 < i < j < prime(n)/2 such that R(i^4,prime(n)) > R(j^4,prime(n)), where R(k,p) (with p an odd prime) denotes the unique integer r among 0,...,(p-1)/2 with k congruent to r or -r modulo p.

Conjecture: Let p be an odd prime, and let r(p) be the number of ordered pairs (i,j) with 0 < i < j < p/2 and R(i^4,p) > R(j^4,p), where R(k,p) denotes the unique integer r among 0,...,(p-1)/2 with k congruent to r or -r modulo p. Then r(p) is even if p == 3 (mod 4). Also, r(p) == (p-5)/8 (mod 2) if p == 5 (mod 8). When p == 1 (mod 8), r(p) is even if and only if 2 is a quartic residue modulo p. See also A319311, A319480 and A319882 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 30 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014059 | OEIS-A319926

Isomer counts of compound perfect squared squares.

The isomer count of a compound perfect squared square (CPSS) is the number of ways its squared subrectangle and constituent squares can be arranged, up to symmetry of the CPSS. A squared square is perfect if none of its constituent squares are the same size. A squared square is compound if it contains a smaller squared subrectangle. Note that the squared subrectangle can be a squared square. Specific concrete examples of CPSSs with isomer counts under 100 of 4, 7, 8, 11, 12, 16, 19, 20, 23, 24, 28, 31, 32, 35, 36, 39, 40, 47, 48, 56, 60, 63, 64, 68, 72, 76, 80, 88 and 96 exist. Geometric constructions based on a suitable pair of perfect squared rectangles each with up to 4 isomers suggests additional isomer counts up to 100 of 14, 21, 22, 33, 42, 44, 66 and 99, but no actual examples are known. As the

number of squares in a squared square - the order - increases new arrangements appear. It is conjectured that expected CPSS subrectangle isomer arrangements will eventually appear if the order is high enough.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stuart E Anderson_, Oct 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014060 | OEIS-A319927

Numbers k such that the sum of the squares of the odd non-unitary divisors of k divides the sum of the squares of the non-unitary divisors of k.

Conjecture: For any nonnegative integer power p the sum of the p-th powers of the odd non-unitary divisors of a(n) divides the sum of the p-th powers of the non-unitary divisors of a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ivan N. Ianakiev_, Oct 02 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014061 | OEIS-A319940

O.g.f. A(x) satisfies [x^n] exp(-n*A(x)) / (1 - n*x)^n = 0, for n > 0.

a(n) is even for n > 2, with a(2^k + 1) = 2 (mod 4) for k >= 1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014062 | OEIS-A320031

Square array A(n,k), n >= 0, k >= 0, read by antidiagonals, where column k is the expansion of the e.g.f. exp(x)/(1 - k*x).

Conjecture: E.g.f. of row n is exp(x) * (Sum_{k=0..n} A371898(n, k) * x^k / k!). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Oct 03 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014063 | OEIS-A320043

Row sums of the triangle A322550.

Conjecture: a(n) is not a perfect square except for n = 1, 6 and 96.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Dec 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014064 | OEIS-A320044

Number of positive integers k < prime(n)/2 with {k^3/prime(n)} > 1/2, where {x} = x - floor(x) is the fractional part of a real number x.

Conjecture 1: For any prime p == 5 (mod 6), the difference card{0 < k < p/2: {k^3/p} > 1/2} - (p+1)/6 is nonnegative and even. Conjecture 2: For any prime p not congruent to 1 modulo 5, the number of positive integers k < p/2 with {k^5/p} > 1/2 is even. Conjecture 3: For any prime p == 5 (mod 12), the difference card{0 < k < p/2: {k^6/p} > 1/2} - (p-5)/12 is positive and odd.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014065 | OEIS-A320047

Consider coefficients U(m,l,k) defined by the identity Sum_{k=1..l} Sum_{j=0..m} A302971(m,j)/A304042(m,j) * k^j * (T-k)^j = Sum_{k=0..m} (-1)^(m-k) * U(m,l,k) * T^k that holds for all positive integers l,m,T. This sequence gives 2-column table read by rows, where n-th row lists coefficients U(1,n,k) for k = 0, 1 and n >= 1.

Conjectures from _Colin Barker_, Aug 03 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petro Kolosov_, Oct 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014066 | OEIS-A320051

Square array read by antidiagonals upwards: T(n,k) is the n-th positive integer with exactly k middle divisors, n >= 1, k >= 0.

Conjecture 1: T(n,k) is also the n-th positive integer j with the property that the difference between the number of partitions of j into an odd number of consecutive parts and the number of partitions of j into an even number of consecutive parts is equal to k. Conjecture 2: T(n,k) is also the n-th positive integer j with the property that the symmetric representation of sigma(j) has width k on the main diagonal.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Oct 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014067 | OEIS-A320097

Number of no-leaf subgraphs of the 4 X n grid.

Conjecture: a(n) = 36*a(n-1) - 7*a(n-2) - 201*a(n-3) + 49*a(n-4) + 20*a(n-5) - 5*a(n-6) for all n > 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Oct 05 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014068 | OEIS-A320099

Number of no-leaf subgraphs of the 5 X n grid.

Conjecture: a(n) = 103*a(n-1) + 1063*a(n-2) - 1873*a(n-3) - 20274*a(n-4) + 44071*a(n-5) - 10365*a(n-6) - 20208*a(n-7) + 5959*a(n-8) + 2300*a(n-9) - 500*a(n-10) for n > 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Oct 05 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014069 | OEIS-A320137

Numbers that have only one middle divisor.

Conjecture 1: sequence consists of numbers k with the property that the difference between the number of partitions of k into an odd number of consecutive parts and the number of partitions of k into an even number of consecutive parts is equal to 1. Conjecture 2: sequence consists of numbers k with the property that the symmetric representation of sigma(k) has width 1 on the main

diagonal. Conjecture 3: all powers of 2 are in the sequence. Every number in this sequence is a square or twice a square, i.e., this sequence is a subsequence of A028982, and conjectures 2 and 3 are true (see the link for proofs). Furthermore, all odd numbers in this sequence are squares and form subsequences of A016754 and of A319529.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Oct 06 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014070 | OEIS-A320142

Numbers that have exactly two middle divisors.

Conjecture 1: numbers k with the property that the difference between the number of partitions of k into an odd number of consecutive parts and the number of partitions of k into an even number of consecutive parts is equal to 2. Conjecture 2: numbers k with the property that symmetric representation of sigma(k) has width 2 on the main diagonal. By the theorem in A067742 conjecture 2 is true. - _Hartmut F. W. Hoft_, Aug 18 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Oct 06 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014071 | OEIS-A320192

Number of summation terms of the reciprocal integer squares series that gives the best approximation to n terms of the Euler product for zeta(2).

Conjecture: It appears that for integer n the number of summation terms of the reciprocal integer squares series that gives the best approximation to Pi^2/6 - 1/n is n. - _Andrew Howroyd_, Nov 24 2018 Conjecture: a(n) = floor(1/(Pi^2/6 - Product{k=1..n} 1/(1-1/prime(k)^2))). - _Andrew Howroyd_, Nov 24 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Horst H. Manninger_, Oct 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014072 | OEIS-A320223

First occurrence of the n-th prime in A075019.

Conjecture: All primes are present in A075019.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Oct 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014073 | OEIS-A320235

G.f.: Product_{k>=1, j>=1} (1 + x^(k*j))^2.

Conjecture: log(a(n)) ~ Pi*sqrt(n*log(n)/3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Oct 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014074 | OEIS-A320236

G.f.: Product_{k>=1, j>=1} 1/(1 - x^(k*j))^2.

Conjecture: log(a(n)) ~ Pi * sqrt(2*n*log(n)/3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Oct 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014075 | OEIS-A320237

G.f.: Product_{k>=1, j>=1} ((1 + x^(k*j)) / (1 - x^(k*j)))^2.

Conjecture: log(a(n)) ~ Pi * sqrt(n*log(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Oct 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014076 | OEIS-A320244

G.f.: Product_{k>=1, j>=1} (1 + x^(k*j))^2 / (1 - x^(k*j)).

Conjecture: log(a(n)) ~ Pi * sqrt(2*n*log(n)/3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Oct 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014077 | OEIS-A320384

Primes p such that 3/2 is a primitive root modulo p.

By Artin's conjecture, this sequence contains 37.395% of all primes, or 74.79% of all primes congruent to 7, 11, 13, 17 modulo 24.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Oct 12 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014078 | OEIS-A320416

G.f.: exp(Sum_{n>=1} sigma_2(n)*sigma_3(n)/sigma(n) * x^n / n), where sigma_{k}(n) equals the sum of the k-th powers of the divisors of n.

This sequence is conjectured to consist entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014079 | OEIS-A320422

The number of tiles inside a regular n-gon created by diagonals that run from each of the n vertices to the n-2 midpoints of opposite edges.

Conjecture: a(2n+1) = n*(2*n+1)*(4*n^2-3*n+5)/3. - Thomas Young (tyoung(AT)district16.org), Jan 05 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jan 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014080 | OEIS-A320465

a(n) = 2^n - (2^(n-1) mod n), where "mod" is the nonnegative remainder operator.

Conjecture: for n > 2, a(n) is prime if and only if n is in A000043. Note that a(n) = 2^n - 1 if and only if n is an odd prime or pseudoprime to base 2 (A001567), so a counterexample cannot be a Fermat pseudoprime to base 2. - _Thomas Ordowski_, Oct 14 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Oct 13 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014081 | OEIS-A320487

a(0) = 1; thereafter a(n) is obtained by applying the "delete multiple digits" map m -> A320485(m) to 2*a(n-1).

Conjecture: If we start with any nonnegative integer and repeatedly double and apply the "delete multiple digits" map m -> A320485(m), we eventually reach 0 or 1 (see A323835). - _N. J. A. Sloane_, Feb 03 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 24 2018, following a suggestion from _Eric Angelini_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014082 | OEIS-A320492

Coordination sequence of thinnest 5-neighbor packing of the plane with congruent triangles with respect to a hexavalent point with three-fold rotational symmetry.

Conjectures from _Colin Barker_, Oct 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014083 | OEIS-A320493

Coordination sequence of thinnest 5-neighbor packing of the plane with congruent triangles with respect to a hexavalent point with no symmetry.

Conjectures from _Colin Barker_, Oct 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014084 | OEIS-A320494

Coordination sequence of thinnest 5-neighbor packing of the plane with congruent triangles with respect to a tetravalent point.

Conjectures from _Colin Barker_, Oct 23 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014085 | OEIS-A320497

Coordination sequence of thinnest 5-neighbor packing of the plane with congruent hexagons with respect to a point of type C.

Conjectures from _Colin Barker_, Oct 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014086 | OEIS-A320498

Coordination sequence of thinnest 5-neighbor packing of the plane with congruent hexagons with respect to a point of type D.

Conjectures from _Colin Barker_, Oct 25 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 22 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014087 | OEIS-A320515

Squarefree k > 1 with sigma(sigma(k)) < 2*k + 1.

Conjecturally a subsequence of A085497. This conjecture is false, the first counterexample is a(113) = 2257 = 37 * 61 which is the least composite term in this sequence. - _Amiram Eldar_, Jun 17 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Oct 14 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014088 | OEIS-A320568

a(n) = Sum_{k=1..n} (-1)^(n-k)*binomial(n,k)*sigma(k).

Conjecture: a(n) ~ (-1)^n * Pi^2/48 * n * 2^n. - _Vaclav Kotesovec_, Oct 16 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Oct 15 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014089 | OEIS-A320572

The smallest integer m such that each nonzero digit appears in the decimal representation of the sequence n^1, n^2, ..., n^x, where 1 <= x <= m, or 0 if no such m exists.

Conjecture: a(n) <= 15, with a(n) = 15 only for numbers of the form 2 * 10^k (verified for n <= 10^7). - _Robert Israel_, Sep 03 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benjamin Knight_, Oct 15 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014090 | OEIS-A320586

Expansion of (1/(1 - x)) * Sum_{k>=1} k*x^k/(x^k + (1 - x)^k).

Conjecture: a(n) ~ c * 2^n * n, where c = Pi^2/24 = 0.411233516712... - _Vaclav Kotesovec_, Jun 26 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Oct 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014091 | OEIS-A320591

Expansion of Product_{k>=1} (1 + x^k/(1 + x)^k).

Conjectural o.g.f.: (1/2) * Sum_{n >= 0} x^(n*(n-1)/2)*(1 + x)^n/(Product_{k = 1..n} ((1 + x)^k - x^k)). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Oct 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014092 | OEIS-A320601

Exponents of powers of two having a digit zero in decimal.

Complement of A007377. It is a long-standing open problem to show that this sequence contains all numbers > 86.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Oct 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014093 | OEIS-A320632

Numbers k such that there exists a pair of factorizations of k into factors > 1 where no factor of one divides any factor of the other.

Mats Granvik has conjectured that these are all the positive integers k such that sigma_0(k) - 2 > (bigomega(k) - 1) * omega(k), where sigma_0 = A000005, omega = A001221, and bigomega = A001222. - _Gus Wiseman_, Nov 12 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Dec 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014094 | OEIS-A320666

a(n) is the maximum number of liberties a single group can have on an otherwise empty n X n Go board.

Exact for n <= 24, conjectured for n > 24 but it is at least a lower bound: Conjectures from _Colin Barker_, Jun 05 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ton Hospel_, Oct 28 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014095 | OEIS-A320722

Reduced numbers with multiplicative persistence 10 in base 15.

For any multiplicative persistence, except the multiplicative persistence 2, the set of class representative numbers with that multiplicative persistence is conjectured to be finite. Each class representative number represents an infinite set of numbers with the same multiplicative persistence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _A.H.M. Smeets_, Oct 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014096 | OEIS-A320907

Row sums of A320906.

Conjectures from _Colin Barker_, Oct 28 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Oct 28 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014097 | OEIS-A320926

Concatenation of successive segments generated by the morphism {0 -> {0, 0, 1}, 1 -> {0}}, starting with 0.

Conjecture: rho* = 0.085.... (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 24 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014098 | OEIS-A320949

G.f. A(x) satisfies: Sum_{n>=0} (1+x)^(n^2) / (1 + A(x))^(n+1) = 1.

a(n) = 0 (mod 3) for n > 0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 13 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014099 | OEIS-A321002

a(0)=3; thereafter a(n) = 20*6^(n-1)-2^(n-1).

Conjectured to be the sum of A175046(i) for 2^n <= i < 2^(n+1). Conjecture is true (see comments in A175046). - _Chai Wah Wu_, Nov 18 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014100 | OEIS-A321003

a(n) = 2^n*(4*3^n-1).

Conjectured to be the sum of A175046(i) for 1 <= i < 2^(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014101 | OEIS-A321005

Let p = prime(n); a(n) = number of pairs of primes (i,j) with i < j < p such that i*j == 1 (mod p).

See link for conjectured asymptotic behavior of this sequence, showing (red line) first asymptotic term and residual bounds (blue lines). It appears that a(n) ~ Li(p)^2/(2*p), based upon the assumption that the probabilities that the first and second numbers in a pair are primes are independent. The residuals (with minor exceptions for some small p) seem to be bounded by (2/Pi^2)*sqrt(p). These observations were contributed by Karl Love (via Mapleprimes). - _David James Sycamore_, Nov 05 2018 An alternative heuristic argument leads the conjecture that a(n) ~ (n-4)/(2*log(n)) ~ Pi(n)/2. (Essentially the same as Karl Love's conjecture in the Comments section, but simpler to calculate.) - _N. J. A. Sloane_, Nov 06 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 02 2018, following a suggestion from _David James Sycamore_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014102 | OEIS-A321013

a(n) = how many of {6,7,8} divide n.

Conjectures from _Colin Barker_, Nov 05 2018: (Start) The above conjectures are true. The sequence is periodic with period 168. Let f(x) be the g.f. above. Then f(x + 168) = f(x).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014103 | OEIS-A321032

Number of words of length 3n such that all letters of the binary alphabet occur at least once and are introduced in ascending order and which can be built by repeatedly inserting triples into the initially empty word.

Conjectures from _Mikhail Smirnov_, Oct 15 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Oct 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014104 | OEIS-A321084

Primes prime(n) such that 1 + Sum_{k=1..n} 2^(prime(k)-1) is prime.

Let S(n) = Sum_{k=1..n} 2^(prime(k)-1). Conjecture: q(n) = 1 + S(n) is prime if and only if 2^S(n) == 1 (mod q(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Oct 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014105 | OEIS-A321122

a(n) = n-th row common denominator of A321121.

a(n) = 52*a(n-6) - a(n-12) for n >= 15 (conjectured). G.f.: (4 + 2*x + 3*x^2 + 8*x^3 + 36*x^4 + 96*x^5 - 164*x^6 + 256*x^7 + 336*x^8 + 32*x^9 - 36*x^10 + 24*x^11 + 2*x^13 - 9*x^14)/(1 - 52*x^6 + x^12) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franck Maminirina Ramaharo_, Nov 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014106 | OEIS-A321137

Reduced numbers with multiplicative persistence 11 in base 14.

For any multiplicative persistence, except the multiplicative persistence 2, the set of class representative numbers with that multiplicative persistence is conjectured to be finite. Each class representative number represents an infinite set of numbers with the same multiplicative persistence.If there exists more reduced numbers with multiplicative persistence 11, it will be larger than 14^100.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _A.H.M. Smeets_, Oct 28 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014107 | OEIS-A321152

n such that all n - p are practical numbers where p is a practical number in range n/2 <= p < n.

This sequence is analogous to A320447. Instead of the sequence of primes it uses the sequence of practical numbers (A005153). It is conjectured that the sequence is finite and full.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Dec 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014108 | OEIS-A321170

Numbers k such that Fibonacci(k) in base 5 ends with k in base 5.

Conjecture: For m >= 4, a(39 + (m-4)*77) = 5^m and the 77 first differences to the next m are ((x,y,x)^6, x, 4, y-4, x, (x,y,x)^11, x, y-4, 4, x, (x,y,x)^6) with x = 5^(m-2)-5, y = 2*(5^(m-2)+5). Verified to m=30.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lars Blomberg_, Jan 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014109 | OEIS-A321225

Break the infinite word 1232112321... into chunks so that the sum of the digits in the n-th chunk is n.

Conjectures from _Chai Wah Wu_, Apr 18 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Oct 31 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014110 | OEIS-A321252

(Conjecturally) the lexicographically earliest infinite sequence over {0,1,2,3} avoiding abelian squares.

(Conjecturally) the lexicographically earliest infinite sequence over {0,1,2,3} avoiding abelian squares. An abelian square is a nonempty string where the second half is a rearrangement of the first half, like the English word "reappear". To "avoid" an abelian square means to have no contiguous block of that form. Although an easy compactness argument, combined with a result of Keränen (1992) shows that the lexicographically earliest infinite string avoiding abelian squares over the alphabet {0,1,2,3} must exist, the terms provided are only conjecturally part of the described sequence, because we have no proof currently that this particular string can be extended infinitely far to the right.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Nov 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014111 | OEIS-A321266

Smallest positive number for which the square cannot be written as sum of distinct squares of any subset of previous terms.

a(n) = 2 * a(n-3) for n > 9 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bert Dobbelaere_, Nov 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014112 | OEIS-A321291

Smallest positive number for which the 4th power cannot be written as sum of 4th powers of any subset of previous terms.

a(n) = 2 * a(n-12) for n > 25 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bert Dobbelaere_, Nov 02 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014113 | OEIS-A321303

a(n) = floor(d(n) * n^(11/2)) where d(n) is the number of divisors of n.

Ramanujan conjectured in 1916 that |tau(p)| <= 2 * p^(11/2) for all primes p and Pierre Deligne proved this conjecture in 1974. [Wikipedia] - _Bernard Schott_, Oct 24 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Nov 03 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014114 | OEIS-A321309

Coefficients of the power series expansion at p=1 of the growth rate C(p) of the length of the longest increasing path in an Erdös-Rényi graph with edge probability p.

The entries are known to be integers, they were conjectured to be nonnegative and increasing starting from index 2. The radius of convergence of the generating function is at least (sqrt(2)-1)/2 and at most 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sanjay Ramassamy_, Nov 03 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014115 | OEIS-A321321

Numbers n for which the "partition-and-add" operation applied to the binary representation of n results in only one power of 2.

Conjecture: With the exception of a(1) = 1 and a(17) = 31, all terms have a binary weight of 2 or 3. - _Peter Kagey_, Jun 14 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Nov 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014116 | OEIS-A321334

n such that all n - s are squarefree numbers where s is a squarefree number in range n/2 <= s < n.

This sequence is analogous to A320447. Instead of the sequence of primes it uses the sequence of squarefree numbers (A005117). It is conjectured that the sequence is finite and full.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Dec 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014117 | OEIS-A321341

An unbounded sequence which is 1 infinitely often, with the property that for any four consecutive terms the maximum term is the sum of the two minimum terms.

This sequence was constructed to show that there are many sequences, besides those merging with multiples of the Padovan sequence A000931, with the property that for any four consecutive terms the maximum term is the sum of the two minimum terms. This refutes a conjecture that was formerly in that entry.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Nacin_, Nov 05 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014118 | OEIS-A321480

Zeroless analog of triangular numbers (version 2): a(0) = 0, and for any n > 0, a(n) = noz(1 + noz(2 + ... + noz((n-1) + n))), where noz(n) = A004719(n) omits the zeros from n.

This sequence is conjectured to be bounded. This could be explained by the fact that the zeros appearing in the last steps of the calculation (when adding small values) erode the number of digits of the intermediate sums.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Nov 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014119 | OEIS-A321482

a(n) = log_2(A305677(n)).

It is conjectured that A305677(n) > 0. If A305677(n) = 0, then define a(n) = -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Nov 12 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014120 | OEIS-A321487

Numbers in A121707 (n^3 > 1 divides Sum_{k=1..n-1} k^n) which are not semiprimes.

Most terms of A121707 and its (conjectured) subsequence A267999 are semiprimes. This sequence lists the exceptions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Nov 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014121 | OEIS-A321518

Smallest k > 1 such that n^k + k^n is prime, i.e., a Leyland prime, or 0 if no such k exists.

Conjecture: a(6) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Nov 12 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014122 | OEIS-A321575

Nexus primary pretenders: a(n) is the smallest composite k such that n^k - (n-1)^k == 1 (mod k).

Conjecture: the only terms of the sequence that are not squarefree are 4, 9 and 49. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Nov 13 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014123 | OEIS-A321576

a(n) is the smallest b > 1 such that b^n - (b-1)^n has all divisors d == 1 (mod n).

If n is prime, then a(n) = 2. Conjecture: If n is composite, then a(n) > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Nov 13 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014124 | OEIS-A321612

Numbers k such that all k - t are triangular numbers where t is a triangular number in range k/2 <= t < k.

This sequence is analogous to A320447. Instead of the sequence of primes it uses the sequence of triangular numbers (A000217). It is conjectured that the sequence is finite and full.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Dec 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014125 | OEIS-A321614

Number of nonequivalent ways to place 2n nonattacking kings on a 4 X 2n chessboard under all symmetry operations of the rectangle.

Conjectures from _Colin Barker_, Dec 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anton Nikonov_, Dec 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014126 | OEIS-A321616

Primes p = k^2 + (k-1)^2 such that k^p - (k-1)^p is prime.

Conjecture: generally, these are primes p = a^2 + b^2 with a > b > 0 such that (a^p - b^p)/(a-b) is prime, so must be a-b = 1. It seems that there are no primes (a^q + b^q)/(a+b) for primes q = a^2 + b^2 > 5. Especially, there are probably no primes q = m^2 + 1 > 5 such that (m^q - 1)/(m-1) is prime or (m^q + 1)/(m+1) is prime. How to prove it?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Nov 15 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014127 | OEIS-A321691

Approximations up to 2^n for the 2-adic integer log(-3).

Conjecture: a(n) = 2*A309753(n-1). - _R. J. Mathar_, Aug 06 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Nov 17 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014128 | OEIS-A321728

Number of integer partitions of n whose Young diagram cannot be partitioned into vertical sections of the same sizes as the parts of the original partition.

Conjecture: a(n) is the number of non-half-loop-graphical partitions of n. An integer partition is half-loop-graphical if it comprises the multiset of vertex-degrees of some graph with half-loops, where a half-loop is an edge with one vertex, to be distinguished from a full loop, which has two equal vertices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Nov 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014129 | OEIS-A321729

Number of integer partitions of n whose Young diagram can be partitioned into vertical sections of the same sizes as the parts of the original partition.

Conjecture: a(n) is the number of half-loop-graphical partitions of n. An integer partition is half-loop-graphical if it comprises the multiset of vertex-degrees of some graph with half-loops, where a half-loop is an edge with one vertex, to be distinguished from a full loop, which has two equal vertices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Nov 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014130 | OEIS-A321773

Number of compositions of n into parts with distinct multiplicities and with exactly three parts.

Conjectures from _Colin Barker_, Dec 11 2018: (Start) Conjecture: a(n) = (3*n-k)/2 where k value has a cycle of 6 starting from n=3 of (7,6,3,10,3,6). - _Bill McEachen_, Aug 12 2025 The above conjectures are correct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Nov 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014131 | OEIS-A321774

Number of compositions of n into parts with distinct multiplicities and with exactly four parts.

Conjectures from _Colin Barker_, Dec 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Nov 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014132 | OEIS-A321775

Number of compositions of n into parts with distinct multiplicities and with exactly five parts.

Conjectures from _Colin Barker_, Dec 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Nov 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014133 | OEIS-A321776

Number of compositions of n into parts with distinct multiplicities and with exactly six parts.

Conjectures from _Colin Barker_, Dec 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Nov 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014134 | OEIS-A321777

Number of compositions of n into parts with distinct multiplicities and with exactly seven parts.

Conjectures from _Colin Barker_, Dec 11 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Nov 18 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014135 | OEIS-A321790

a(n) is the smallest base a > 2 such that a^(k-1) != 1 (mod k), where k = A001567(n), the n-th Fermat pseudoprime to base 2.

Conjecture: if k = A001567(n) is semiprime, then a(n) < lpf(k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Nov 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014136 | OEIS-A321859

Number of primes congruent to 3, 5, 6 modulo 7 and <= n minus number of primes congruent to 1, 2, 4 modulo 7 and <= n.

The first 10000 terms (except for a(2)) are nonnegative. a(p) = 0 for primes p = 3, 11, 211, 3371, 3389, ... The earliest negative term (besides a(2)) is a(48673) = -1. Conjecturally infinitely many terms should be negative.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Nov 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014137 | OEIS-A321862

a(n) = A321857(prime(n)).

The first 10000 terms are positive, but conjecturally infinitely many terms should be negative.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Nov 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014138 | OEIS-A321941

Scaled numerators in the asymptotic expansion of the Maclaurin coefficients in a Hadamard product involving the exponential integral.

a(n) is the numerator of the rational number called r_n in Brent et al. (2018). It is conjectured that r_n is an integer, so the denominators should all be 1 (this has been verified for n <= 1000). A stronger conjecture is given in Remark 12 of Brent et al. (2018). It is known that n!*r_n is an integer, see Theorem 18 of Brent et al. (2018).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard P. Brent_, Dec 08 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014139 | OEIS-A321942

A sequence related to the Euler-Gompertz constant.

Conjecture: Integral_{x = 0..oo} (x/(1 + x))^n*exp(-x) dx = 1/(n-1)!*(a(n) - A000262(n)*G), where G = Integral_{x = 0..oo} exp(-x)/(1 + x) dx is the Euler-Gompertz constant A073003. - _Peter Bala_, Mar 20 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard P. Brent_, Dec 12 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014140 | OEIS-A321966

Triangle read by rows, coefficients of a family of orthogonal polynomials, T(n, k) for 0 <= k <= n.

We conjecture that the polynomials have only negative and simple real roots.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Dec 20 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014141 | OEIS-A321982

Row n gives the chromatic symmetric function of the n-ladder, expanded in terms of elementary symmetric functions and ordered by Heinz number.

Conjecture: All terms are nonnegative (verified up to the 5-ladder).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Nov 23 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014142 | OEIS-A321992

a(n) is the least prime q different from p = prime(n) such that 2^(q-1) == 1 (mod p), or 0 if no such prime exists.

a(n) is the smallest prime q <> p such that q == 1 (mod ord_{p}(2)), where ord_{p}(2) = A002326((p-1)/2) = A014664(n). Strong conjecture: a(n) < A014664(n)^2. - _Thomas Ordowski_, Mar 15 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Mar 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014143 | OEIS-A321994

Number of different chromatic symmetric functions of hypertrees on n vertices.

Stanley conjectured that the number of distinct chromatic symmetric functions of trees with n vertices is equal to A000055, i.e., the chromatic symmetric function distinguishes between trees. It has been proven for trees with up to 25 vertices. If it is true in general, does the chromatic symmetric function also distinguish between hypertrees, meaning this sequence would be equal to A035053?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Nov 24 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014144 | OEIS-A322004

Index i of the smallest Fibonacci number > n such that Fib(i) - n is a prime, or 0 if no such index exists.

Motivated by a recent "conjecture" seen on internet, that any number > 1 is of the form prime(i) + Fib(j) + Fib(k). Actually the number of such representations increases that fast that this conjecture seems not interesting. The present sequence shows that any number is the difference of a Fibonacci number and a prime, or such that n + some prime = some Fibonacci number. (See A322005 for the corresponding prime.) Most of the terms correspond to the index of the smallest Fibonacci number > n or the subsequent one. Local maxima and/or a(n) > a(n+1) + 1 correspond to numbers for which one has to look further. a(25) = 102 is a noteworthy example.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Dec 12 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014145 | OEIS-A322008

1/(1 - Integral_{x=0..1} x^(x^n) dx), rounded to the nearest integer.

Conjectures from _Colin Barker_, Mar 07 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Mar 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014146 | OEIS-A322026

Lexicographically earliest infinite sequence such that a(i) = a(j) => A007814(i) = A007814(j) and A007949(i) = A007949(j), for all i, j, where A007814 and A007949 give the 2- and 3-adic valuations of n.

a(n) = A323884(25*n). [Conjectured by Sequence Machine]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 03 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014147 | OEIS-A322041

Triangle read by rows: let E denote the standard triangular 6-valent grid in the plane, regarded as a graph with the Eisenstein integers as vertices; row n gives the coordination sequence of the quotient graph E/nE.

When E is regarded as a lattice in R^2, E/nE has packing radius roughly n/2, but covering radius roughly n/sqrt(3) > n/2 (see Conway-Sloane, Chapter 4). This means that as n increases, the number of terms in the n-th row of the triangle will increase linearly with n. The largest k such that T(n,k) is nonzero is A322042(n), which is conjecturally n - ceiling(n/3). One can now formulate a specific conjecture for what row n is, for each of the six residue classes of n mod 6. There are similar conjectures for n = 6t+1, ..., 6t+5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 05 2018; corrected and extended Dec 06 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014148 | OEIS-A322042

Maximum edge-distance of a point in the quotient graph E/nE from the origin (see A322041 for further information).

Conjectures from _Colin Barker_, Dec 06 2018: (Start) Conjecture: a(n) = n - ceiling(n/3) = A004523(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 06 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014149 | OEIS-A322043

Numbers k such that the coefficient of x^k in the expansion of Product_{m >= 1} (1-x^m)^15 is zero.

Van der Blij, discussing the conjecture that the Ramanujan numbers tau(k) (see A000594) are never zero, mentions that a certain "Ferguson" had shown that 52 is a member of the current sequence. No details were given, and the 52 appears to be a typo for 53. Conjectures from _Colin Barker_, Dec 07 2018: (Start) E.g.f.: (1/2)*exp(x)*(106 + 858*x + 429*x^2). - conjectured by _Stefano Spezia_, Dec 07 2018 after the conjectures of _Colin Barker_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014150 | OEIS-A322048

Final elements in rows when A322050 is displayed as a triangle.

Conjectures: a(n) = 2*a(n-1) + 11 for n >= 5; G.f. = x*(6*x^4+2*x^2+2*x+1)/((1-x)*(1-2*x)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 15 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014151 | OEIS-A322049

When A322050 is displayed as a triangle the rows converge to this sequence.

Indices of records of a(n)/n are (1, 3, 7, 11, 23, 27, 43, 55, 87, 91, 119, 171, 183, 343, 347, 363, 367, 375, 439, 695, 731, 887, 1367, 1371, 1391, 1399, 1451, 1463, 2743, 2923, 2927, 2935, 3511, ...). The ratio a(n)/n increases roughly by 1 at each of these. We conjecture that this ratio is unbounded. We note that the record ratios occur in "clusters" at indices twice as large as the preceding cluster: 87, 91; 171, 183; 343..375; 695..731; 1367..1463; 2743..2935; ... This is compatible with the self-similar structure of the graph of this sequence, which starts over at a(2^k) = 8 for all k >= 4. (But note also the distinctive substructure repeating with period 2^10, cf. the "logarithmic plot" link.) - _M. F. Hasler_, Dec 18 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 15 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014152 | OEIS-A322051

a(n) is the number of initial terms in the row of length 2^n of A322050 that agree with the limiting sequence A322049.

Conjecture: For n >= 5, a(n) = 2*a(n-1)-1 if n is odd, 2*a(n-1) if n is even. Conjectures from _Colin Barker_, Dec 29 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 16 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014153 | OEIS-A322055

Number of ON cells after n generations of two-dimensional automaton based on knight moves (see Comments for definition; here a cell is turned ON if 1 or 2 neighbors are ON).

Conjectures from _Colin Barker_, Dec 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014154 | OEIS-A322056

First differences of A322055.

Conjectures from _Colin Barker_, Dec 22 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014155 | OEIS-A322069

Number of permutations f of {1,...,n} with f(1) < f(n) such that Sum_{k=1..n-1} 1/(f(k)*f(k+1)) = 1.

Conjecture 1: a(n) > 0 for all n > 5. In other words, for each n = 6,7,... we have Sum_{k=1..n-1} 1/(f(k)*f(k+1)) = 1 for some permutation f in the symmetric group S_n. Conjecture 2: For any integer n > 6, there is an undirected circular permutation g different from the circular permutation (1,2,...,n) such that 1/(g(1)*g(2)) + 1/(g(2)*g(3)) + ... + 1/(g(n-1)*g(n)) + 1/(g(n)*g(1)) = 1. We have verified both conjectures for n up to 11. For Conjecture 2 with n = 7, we may take (g(1),...,g(7)) = (3,2,1,6,5,4,7) since 1/(3*2) + 1/(2*1) + 1/(1*6) + 1/(6*5) + 1/(5*4) + 1/(4*7) + 1/(7*3) = 1. See also A322070 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014156 | OEIS-A322072

Row sums of the triangle A322071.

Conjecture: The difference a(n + 1) - a(n) between two consecutive terms is not a perfect square except for n = 1, 5 and 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Nov 25 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014157 | OEIS-A322099

Number of permutations f of {1,...,n} with f(1) < f(n) and Sum_{k=1..n-1} 1/(f(k)^2 - f(k+1)^2) = 0.

Conjecture: a(n) > 0 for all n > 7. In other words, for each n = 8,9,... we have Sum_{k=1..n-1} 1/(f(k)^2 - f(k+1)^2) = 0 for some permutation f in the symmetric group S_n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014158 | OEIS-A322108

Distance of n-th iteration in an alternating rectangular spiral.

Conjectures from _Colin Barker_, Feb 13 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Drew Edgette_, Nov 26 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014159 | OEIS-A322120

a(n) is the smallest composite k such that n^(k-1) == 1 (mod (n^2-1)*k).

Conjecture: if m is a composite number such that b^(m-1) == 1 (mod (b^2-1)m) for some b, then m is a strong pseudoprime to some base a in the range 2 <= a <= m-2. Thus, probably every term a(n) is in A181782.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Nov 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014160 | OEIS-A322121

Composite numbers m such that b^(m-1) == 1 (mod (b^2-1)*m) has a solution b.

Conjecture: If m is a composite number such that b^(m-1) == 1 (mod (b^2-1)m) for some base b, then m is a strong pseudoprime to some base a in the range 2 <= a <= m-2. Thus, these numbers m are probably a proper subset of A181782.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Nov 27 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014161 | OEIS-A322154

Numbers k satisfying gcd(k^2, sigma(k^2)) > sigma(k), where sigma is the sum-of-divisors function.

The assertion k = 1 is known as the Descartes-Frenicle-Sorli Conjecture on odd perfect numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jose Arnaldo Bebita Dris_, Nov 29 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014162 | OEIS-A322155

Consecutive terms that appear more than once in A014237.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enrique Navarrete_, Dec 11 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014163 | OEIS-A322165

Numbers k that give record values for s(k)*phi(k)/k^2, where s(k) is the sum of squares of the differences between consecutive totatives of k (A322144).

Erdős conjectured that this ratio is bounded and offered $500 for a proof. The conjecture was proved by Montgomery and Vaughan, who won the prize.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Nov 29 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014164 | OEIS-A322210

G.f.: P(x,y) = Product_{n>=1} 1/(1 - (x^n + y^n)), where P(x,y) = Sum_{n>=0} Sum_{k>=0} T(n,k) * x^n*y^k, as a square table of coefficients T(n,k) read by antidiagonals.

Conjecture 1: the triangular table T(n,k) is the number of ways to form the subsum k from the partitions of n, where n and k are integers such that 0 <= k <= n. For example, t(4,2)=10; the five partitions of 4 are (4), (3,1), (2,2), (2,1,1), (1,1,1,1) with subsum 2 occurring {0,0,2,2,6) times for a total of 10. - _George Beck_, Jan 03 2020 Conjecture 2: the square table T(n,k) is the coefficient of s_lambda in the sum over all partitions lambda |-n and nu |-k of (s_rho/mu) where s_lambda*s_mu = Sum(rho|-n+k; C(rho, lambda, mu) s_rho). Simply stated as: multiply lambda with mu, and, for each term in the result, take the skew Schur function with mu and count how often you get the original lambda back. Sum up over all lambda and mu of the size n and k. Conjecture 3: the triangular table T(n,k) is analogous to conjecture 2, but counting s_lambda in s_(lambda/mu) * s_mu with lambda |-n and mu |- k and 0<=k<=n. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 30 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014165 | OEIS-A322253

Number of circular permutations of the integers from 0 to n which generate a complete stepping-on sequence, when the stepping-on direction depends on the odd/even parity of the current value.

It is conjectured that the sequence continues indefinitely. For the complementary case where stepping-on is always in the same direction, no permutation of 0 to n with n even can generate a complete sequence. For odd n, the number of complete sequences corresponds to A141599((n + 1)/2) for n up to 11, as limited by available computing power - it is conjectured that this correspondence continues indefinitely. - _Ian Duff_, Dec 25 2018

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ian Duff_, Dec 01 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014166 | OEIS-A322286

Lexicographically earliest sequence of positive integers without 4 terms in a weakly increasing arithmetic progression.

Assuming Erdős's conjecture on arithmetic progressions, for a fixed positive integer r, the sum of the reciprocals of the numbers n for which the n-th term is r converges.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sébastien Palcoux_, Aug 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014167 | OEIS-A322341

The number of digits used in the sequence up to a(n) [a(n) included] is a multiple of a(n). The sequence starts with a(1) = 1 and is always extended with the smallest integer not yet present in the sequence that does not lead to a contradiction.

The sequence begins like A080676 but diverges from the 56th term on. The sequence is conjectured to be a permutation of A000027 (the positive integers): after computing 20000 terms, we see that the five smallest missing numbers are 2550, 2625, 3365, 3385 and 3399

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Dec 04 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014168 | OEIS-A322419

Number of n-step self-avoiding walks on L-lattice.

It is proved that a(n)^(1/n) has a limit mu called the "connective constant" of the L-lattice; approximate value of mu: 1.5657. It is only conjectured that a(n + 1) ~ mu * a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert FERREOL_, Dec 07 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014169 | OEIS-A322466

Lexicographically first sequence of distinct terms such that the a(n)th digit after a(n) shares a(n)'s parity.

This sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Dec 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014170 | OEIS-A322467

Lexicographically first sequence of distinct terms such that a(n) is duplicated a(n) digits to the right.

This sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-Marc Falcoz_ and _Eric Angelini_, Dec 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014171 | OEIS-A322469

Irregular table: row i = 1, 2, 3, ... starts with 4*i - 1; then, as long as the number is divisible by 3, the next two terms are the result of dividing it by 3, then multiplying it by 2.

Conjectures (verified for k = 0..11):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Georg Fischer_, Dec 09 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014172 | OEIS-A322523

a(n) is the least nonnegative integer k for which there does not exist i < j with i+j=n and a(i)=a(j)=k.

Proof is based on induction on n. Obviously a(n) != 0. For 1 <= e <= a(n/3), there exists i+j = n/3, i < j such that a(i) = a(j) = e-1, by induction hypothesis we have a(3*i) = a(3*j) = e, so a(n) != e. a(n) = a(n/3) + 1 is OK. Suppose otherwise that n = i+j, i < j and a(i) = a(j) = a(n/3) + 1 > 0, then i,j == 0 (mod 3), by induction hypothesis we have a(i/3) = a(j/3) = a(n/3), a contradiction. Obviously a(n) != 0. From (1) we know that a(3^e) = a(2*3^e) = t. For 1 <= e <= t, n - d_e*3^e = 3*(Sum_{0<=r<=t-1, r!=e-1} d_r*3^r + O(3^(t+1))) + 5 = 3*(Sum_{0<=r<=e-2} d_r*3^r + O(3^e)) + 5, so A215879((n-d_e*3^e-5)/3) = e-1. Since 5 <= n - d_e*3^e == 2 (mod 3), by induction hypothesis we have a(n-d_e*3^e) = e = a(d_e*3^e), so a(n) != e. a(n) = t+1 is OK. Suppose otherwise that n = i+j, i != j and a(i) = a(j) = t+1 > 0, then {i,j} == {0,2} (mod 3) and i,j > 2. Suppose that i == 2 (mod 3), by induction hypothesis A215879((i-5)/3) = t. Write i = 3*(Sum_{0<=r<=t-1} d'_r*3^r + O(3^(t+1))) + 5, 1 <= d'_r <= d_r. If d'_r = d_r for all r, then j = n-i is divisible by 3^(t+2), so a(j) >= t+2, a contradiction. If d'_r != d_r for some r, then j = 3^(r_0+1) + O(3^(r_0+2)) where r_0 is the smallest index such that d'_r != d_r, so a(j) = r_0+1 + a(1+O(3^1)) = r_0+1 <= t, also a contradiction.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aidan Clarke_, Aug 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014173 | OEIS-A322598

a(n) is the number of unlabeled rank-3 graded lattices with 3 coatoms and n atoms.

These imply the conjectured G.f. and recursion.(End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jukka Kohonen_, Dec 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014174 | OEIS-A322599

a(n) is the number of unlabeled rank-3 graded lattices with 4 coatoms and n atoms.

Conjectures from _Colin Barker_, Dec 19 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jukka Kohonen_, Dec 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014175 | OEIS-A322600

a(n) is the number of unlabeled rank-3 graded lattices with 5 coatoms and n atoms.

Conjectures from _Colin Barker_, Dec 20 2018: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jukka Kohonen_, Dec 19 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014176 | OEIS-A322631

a(n) = 2*binomial(7*n-1,2*n)/(7*n-1).

In 2012, Nakamigawa and Tokushige stated: Let A[x,y] = number of lattice paths starting at (0,0) that stay in y < 2*x/5 + 2/5 and B[x,y] = number of lattice paths starting at (0,0) that stay in y < 2*x/5 + 1/5, then a(t) = A[5*t-1,2*t-1] + B[5*t-1,2*t-1]. Their theorem was mentioned by D. Knuth in Problem 4 "Lattice Paths of Slope 2/5" in his lecture "Problems That Philippe (Flajolet) Would Have Loved". Knuth reported the empirical observation that A[5*t-1,2*t-1]/B[5*t-1,2*t-1] = a - b/t + O(t^-2), with constants a~=1.63026 and b~=0.159. Knuth's conjecture was proved by C. Banderier and M. Wallner, who also found the exact values of a and b. Numerical values of a and b are provided in A322632 and A322633.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 21 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014177 | OEIS-A322743

Least composite such that complementing one bit in its binary representation at a time produces exactly n primes.

Conjecture: a(n) mod 10 == 5 for n > 7. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_, Dec 24 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014178 | OEIS-A322844

a(n) = (1/12)*n^2*(3*(1 + n^2) - 2*(2 + n^2)*(n mod 2)).

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Dec 28 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014179 | OEIS-A322898

a(n) is the permanent of the matrix [((i + j)/(2*n + 1))]_{i,j=0..n}, where (k/m) denotes the Jacobi symbol.

Conjecture: a(n) is positive for any nonnegative integer n. This conjecture is false, a(26) < 0. - _Vaclav Kotesovec_, Aug 20 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 30 2018 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014180 | OEIS-A322944

Coefficients of a family of orthogonal polynomials. Triangle read by rows, T(n, k) for 0 <= k <= n.

We conjecture that the polynomials have only negative and simple real roots.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jan 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014181 | OEIS-A323013

Form of Zorach additive triangle T(n,k) (see A035312) where each number is sum of west and northwest numbers, with the additional condition that the first element T(n,1) is a Fibonacci number.

Conjecture: Let F(i) be the i-th Fibonacci number. Each number of T(n, k), k = 1, 2, 3 is the difference between two Fibonacci numbers F(i) - F(j) for some i, j, where F(i) is the smallest Fibonacci number greater than T(n, k). The case T(n, 1) is trivial. Examples: 10 = 13 - 3, 29 = 34 - 5, 20 = 21 - 1, 42 = 55 - 13, 84 = 89 - 5, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014182 | OEIS-A323016

a(n) is the number of ordered partitions of 24*n + 4 into four squares of primes (A001248).

a(n) > 0 for 4 <= n <= 2*10^4. Conjecture: a(n) > 0 for all n >= 4. A stronger conjecture: lim inf a(n) = +oo.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Jan 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014183 | OEIS-A323206

A(n, k) = hypergeometric([-k, k+1], [-k-1], n), square array read by ascending antidiagonals for n,k >= 0.

Conjecture: A(n, k) is odd if and only if n is even or (n is odd and k + 2 = 2^j for some j > 0).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Feb 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014184 | OEIS-A323210

a(n) = 9*J(n)^2 where J(n) are the Jacobsthal numbers A001045 with J(0) = 1.

Colin Barker conjectures that A208556 is a shifted version of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jan 09 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014185 | OEIS-A323250

Sequence lists numbers k > 1 such that k^3 == d(k) (mod sigma(k)), where d = A000005 and sigma = A000203.

Conjecture: All terms are even.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_, Jan 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014186 | OEIS-A323274

a(n) = ceiling(1/(e - 1/0! - 1/1! - 1/2! - ... - 1/n!)).

a(n) = least k such that 1/k > e - (n-th partial sum of the Maclaurin series for e). Let b(n) = a(n)/a(n+1). Conjectures: if n > 3, then n+1 < b(n) < n+2 and 0 < b(n+1)-b(n) < 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014187 | OEIS-A323359

a(n) = b(n+1)/b(n) - 1 where b(1)=2 and b(k) = b(k-1) + lcm(floor(sqrt(k^3)), b(k-1)).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pedja Terzic_, Jan 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014188 | OEIS-A323381

Number of solutions to n = x^2 + 2*y^2 + 5*z^2 + x*z in integers.

Rouse [2014] conjectures that the ternary quadratic form x^2 + 2y^2 + 5z^2 + xz represents all positive odd integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Jan 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014189 | OEIS-A323386

a(n) = b(n+1)/b(n) - 1 where b(1)=2 and b(k) = b(k-1) + lcm(floor(sqrt(2)*k),b(k-1)).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pedja Terzic_, Jan 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014190 | OEIS-A323388

a(n) = b(n+1)/b(n) - 1 where b(1)=3 and b(k) = b(k-1) + lcm(floor(sqrt(3)*k), b(k-1)).

Conjectures: The 2nd and 3rd conjectures are proved at the Mathematics Stack Exchange link. - _Sungjin Kim_, Jul 17 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pedja Terzic_, Jan 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014191 | OEIS-A323397

a(n) = (4^n + 15*n - 1)/9.

Conjecture: satisfies a linear recurrence having signature (6, -9, 4). (This is correct, see Formula section.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Harvey P. Dale_, Jan 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014192 | OEIS-A323417

When a prime-based mapping reaches 0.

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo van der Sanden_, Aug 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014193 | OEIS-A323576

Primes p such that 2 is a primitive root modulo p while 128 is not.

According to Artin's conjecture, the number of terms <= N is roughly ((6/41)*C)*PrimePi(N), where C is Artin's constant = A005596, PrimePi = A000720. Compare: the number of terms of A001122 that are no greater than N is roughly C*PrimePi(N).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014194 | OEIS-A323577

Primes p such that 2 is a primitive root modulo p while 2048 is not.

According to Artin's conjecture, the number of terms <= N is roughly ((10/109)*C)*PrimePi(N), where C is Artin's constant = A005596, PrimePi = A000720. Compare: the number of terms of A001122 that are no greater than N is roughly C*PrimePi(N).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014195 | OEIS-A323590

Primes p such that 2 is a primitive root modulo p while 8192 is not.

According to Artin's conjecture, the number of terms <= N is roughly ((12/155)*C)*PrimePi(N), where C is Artin's constant = A005596, PrimePi = A000720. Compare: the number of terms of A001122 that are no greater than N is roughly C*PrimePi(N).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014196 | OEIS-A323610

List of 5-powerful numbers (for the definition of k-powerful see A323395).

The sequence consists of the multiples of 8 that are greater than or equal to 64, together with 48. The result is due to D. Boyd, Berend, and Golan. It had been conjectured that the sequence began with 64, but Boyd discovered the set

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stan Wagon_, Jan 19 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014197 | OEIS-A323617

Primes p such that 3 is a primitive root modulo p while 243 is not.

According to Artin's conjecture, the number of terms <= N is roughly ((4/19)*C)*PrimePi(N), where C is Artin's constant = A005596, PrimePi = A000720. Compare: the number of terms of A001122 that are no greater than N is roughly C*PrimePi(N).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014198 | OEIS-A323653

Multiperfect numbers m such that sigma(m) is also multiperfect.

Conjecture 1 (a): This sequence consists of those m for which sigma(m)/m is an integer (thus a term of A007691), and coprime with m. Or expressed in a slightly weaker form (b): {1} followed by those m for which sigma(m)/m is an integer, but not a divisor of m. In a slightly stronger form (c): For m > 1, sigma(m)/m is always the least prime not dividing m. This would imply both (a) and (b) forms. Conjecture 2: This sequence is finite. Conjecture 3: This sequence is the intersection of A007691 and A351458. Conjecture 4: This is a subsequence of A349745, thus also of A351551 and of A351554.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jan 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014199 | OEIS-A323689

G.f.: Sum_{n>=0} (x^n + i)^n / (1 + i*x^n)^(n+1), in which the constant term is taken to be 1.

Odd terms occur only at n = k^2 for k >= 0 (conjecture);

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014200 | OEIS-A323693

G.f. A(x) satisfies: [x^n] A(x)^(n+1) = (n+1)^2 * [x^(n-1)] A(x)^(n+1) for n >= 1 with A'(0) = 1.

a(n) / 2^floor((n+1)/2) is odd for n >= 0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014201 | OEIS-A323695

G.f.: Sum_{n>=0} x^n * (1 - x^(n+1))^n / (1 + x^(n+1))^(n+1).

Odd terms occur only at n = k*(k+2) for k >= 0 (conjecture);

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014202 | OEIS-A323712

Number of the following described shuffles required to return a deck of n cards to its original state. Create two piles by alternating a card from the top of the deck left-right-left-right until the deck is exhausted. Then, placing the left pile on top of the right pile constitutes one shuffle.

Here are a few conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Lovler_, Jan 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014203 | OEIS-A323729

Non-prime-powers k at which the variance of the first differences of the logarithms of the divisors of k, scaled by log(k), reaches a new minimum.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Jan 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014204 | OEIS-A323739

a(n) is the number of residues modulo (4*primorial(n)) of the squares of primes greater than or equal to prime(n+1).

Conjecture: a(n) = 2^(1-n)*Product_{j=1..n} (prime(j)-1) for n >= 0, so a(n) = a(n-1)*(prime(n)-1)/2 for n >= 1. Conjecture is true. Since there exists a prime congruent to r modulo 4*primorial(n) for any r coprime to primorial(n), this set is precisely the set of coprime quadratic residues of 4*primorial(n). If n >= 1, each residue can be broken down into congruences modulo 8 and the first n-1 odd primes, each odd prime p has (p-1)/2 residue classes, and every combination eventually occurs, giving the formula. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Feb 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014205 | OEIS-A323742

a(n) is the smallest number that is a (2n+1)-digit palindrome in three consecutive integer bases, or 0 if no such number exists.

It is conjectured that a(n)=0 for all n>=4. - _Matej Veselovac_, Mar 25 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Mar 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014206 | OEIS-A323747

Smallest triangular number whose number of divisors is the n-th triangular number, or 0 if no such number exists.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, May 25 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014207 | OEIS-A323748

Square array read by ascending antidiagonals: the n-th row lists the Zsigmondy numbers for a = n, b = 1, that is, T(n,k) = Zs(k, n, 1) is the greatest divisor of n^k - 1 that is coprime to n^m - 1 for all positive integers m < k, with n >= 2, k >= 1.

There are many squares of primes in the third, fourth or sixth column (e.g., T(7,4) = 25 = 5^2, T(22,3) = T(23,6) = 169 = 13^2, T(41,4) = 841 = 29^2, etc.). Conjecturally all other prime powers with exponent >= 2 in the table excluding the first two columns are T(3,5) = 121 = 11^2, T(18,3) = T(19,6) = 343 = 7^3 and T(239,4) = 28561 = 13^4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Jan 25 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014208 | OEIS-A323749

Triangle read by rows: T(m,n) (1 <= n < m) is the number of moves of an (m,n)-leaper (a generalization of a chess knight) until it can no longer move, starting on a board with squares spirally numbered from 1. Each move is to the lowest-numbered unvisited square. T(m,n) = -1 if the path never terminates.

Are all the paths finite? This appears to be an open question.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_, Jan 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014209 | OEIS-A323756

a(1) = 2; for n >= 2, if a(n-1) has not yet been assigned, then a(n-1) = 1 and a(2*n-1) = 2, otherwise a(2*n) = 3.

The conjectured limit above is incorrect, this limit equals 7/4 = 1.75, instead of sqrt(3) = 1.732... This follows from the CLAIM below. The second conjecture above is correct. For a proof, we consider nu^2: We continue with induction, using a recursion formula for b from the Comments of A059448, the LEMMA, and the induction hypothesis: Looking at the positions of 3 in (a(n)), and the positions of 1 in (b(n)), one sees that the claim implies the second conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Jan 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014210 | OEIS-A323830

a(0) = 1; thereafter a(n) is obtained by doubling a(n-1) and repeatedly deleting any string of identical digits.

Conjecture: If we start with any nonnegative number, and repeatedly double it and apply the "repeatedly delete any run of identical digits" operation described here, we eventually reach one of 0, 1, or 5. In other words, the conjecture is that eventually we reach 0 or join the trajectory shown here or the trajectory shown in A323831.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 03 2019, following a suggestion from _Yukun Yao_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014211 | OEIS-A323831

a(0) = 5; thereafter a(n) is obtained by doubling a(n-1) and repeatedly deleting any string of identical digits.

Conjecture: If we start with any nonnegative number, and repeatedly double it and apply the "repeatedly delete any run of identical digits" operation described here, we eventually reach one of 0, 1, or 5. In other words, the conjecture is that eventually we reach 0 or join the trajectory shown here or the trajectory shown in A323830.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014212 | OEIS-A323832

Start with n and repeatedly double it and apply the "repeatedly delete any run of identical digits" operation described in A323830; a(n) is the number of steps needed to reach one of 0, 1, or 5, or -1 if none of these three numbers is ever reached.

Conjecture: every number will eventually reach one of 0, 1, or 5. Conjecture is true for n < 10^10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014213 | OEIS-A323844

Square array T(b,m), read by descending antidiagonals: Number of winning length m strings with a b-symbol alphabet in "same game" (b >= 2, m >= 0).

For binary strings (b = 2), the formula for the number of winning strings of length m (i.e., T(b=2, m) = 2^m - 2 * m * Fibonacci(m-2) - (-1)^m - 1 for m >= 2) was conjectured by Ralf Stephan (2004, p. 8) and proved by Burns and Purcell (2005, 2007). For b-ary strings with b >= 3, the same problem seems to be unsolved. It is not clear whether Kurz's formulas are statements of fact (with an easy proof) or just conjectures. From the results in the Crossrefs, we may also conjecture the following:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, Aug 31 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014214 | OEIS-A323910

Dirichlet inverse of the deficiency of n, A033879.

Following convolution formulas have been conjectured for this sequence by Sequence Machine, with each one giving the first 10000 terms correctly:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014215 | OEIS-A323912

Dirichlet inverse of A083254(n), where A083254(n) = 2*phi(n) - n.

Following convolution formulas were conjectured for this sequence by Sequence Machine, with each one giving the first 10000 terms correctly. The first one is certainly true, because A083254 is Möbius transform of A033879:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014216 | OEIS-A324017

Square array A(m,n) (m>=1, n>=1) read by antidiagonals: A(m,n) = (2*n - 1)^^m mod (2*n)^m (see Comments for definition of ^^).

Conjecture: For any three integers (greater than 1), m, n, and k, such that (2*n - 1)^^m == k (mod (2*n)^m), (2*n - 1)^k == k (mod (2*n)^m). For example, 5^^2 == 29 (mod 6^2) and 5^29 == 29 (mod 6^2). Conjecture: For n > 1 and m >= 2, floor(((2*n - 1)^^m)/(2*n)) == 2*(n - 1) (mod 2*n). For example, floor((13^^3)/14) == 12 (mod 14) and floor((15^^4)/16) == 14 (mod 16). Conjecture: For m > 1, where (2*n - 1)^^m == j (mod (2*n)^(m + 1)), A(m + 1,n) = j. For example, A(6,3) = 563 and A(6,4) = 16115; 11^^3 == 563 (mod 12^3) and 11^^3 == 16115 (mod 12^4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Davis Smith_, Mar 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014217 | OEIS-A324018

Consider the complete graph of an n-cube. a(n) is the number of 2-color colorings of the edges such that they do not contain a monochromatic planar K4 subgraph.

Conjecture: For n >= 3, a(n) < 31*(2^(binomial(2^n,2) - 5) - 2^(binomial(2^n,2) - 10)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Davis Smith_, Aug 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014218 | OEIS-A324152

a(0)=1; for n>0, a(n) = (3/((n+1)*(n+2)*(n+3))) * multinomial(4*n;n,n,n,n).

It is conjectured that a(n) is always an integer. [Proved in Sundaram (2026). - _Shiva Priyan Shanmuga Sundaram_, Sep 08 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_ and _N. J. A. Sloane_, Mar 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014219 | OEIS-A324221

Number of connected 2n-regular loopless multigraphs with five nodes.

Conjectures from _Colin Barker_, Feb 18 2019: (Start) Equivalent conjecture: 1152*a(n) = 6*n^5 + 30*n^4 + 220*n^3 + 540*n^2 + 1143*n - 353 + 72*A056594(n) + 128*A049347(n) + 153*A181983(n+1). - _R. J. Mathar_, Mar 09 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Natan Arie Consigli_, Feb 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014220 | OEIS-A324250

Sequence a(n) = 3*A002559(n) - 2 determining the principal reduced indefinite binary quadratic form [1, a(n), -a(n)] for Markoff triples.

The indefinite binary quadratic form F(n,x,y) = x^2 - 3*m(n)*x*y + y^2 = [1, -3*m(n), 1] representing -m(n)^2 with m(n) = A002559(n), determines Markoff triples MT(n) = (x(n) = A305313(n), y(n) = A305314(n), m(n)) with x(n) < y(n) < m(n), for n >= 3. For n = 1 and 2: x(n) = y(n) = 1. The Frobenius-Markoff conjecture is that this solution is unique. This form F(n,x,y) has discriminant D(n) = (3*m(n))^2 - 4 = a(n)*(a(n) + 4) = A305312(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Mar 04 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014221 | OEIS-A324252

Triangle T(n, k) read by rows from upwards antidiagonals of array A, where A(n, k) is the number of families (also called classes) of proper solutions of the Pell equation x^2 - D(n)*y^2 = k, for k >= 1.

These position lists are A057126 (conjecture) and A243655, for k = 1 and 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Apr 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014222 | OEIS-A324274

a(n) is the number of squares visited by a single pawn move for an even square and a double pawn move for an odd square on a diagonally numbered board and moving to the lowest available unvisited square of different parity at each step from subsequent starting squares n; or a(n) = 0 for an infinite length.

It is conjectured that all starting squares will either have a finite length or reach the top row of the board at square 2 first and then follow the sequence for a(2) to infinity. A324275 contains numbers n for which A324274(n) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jan Koornstra_, Feb 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014223 | OEIS-A324275

Numbers k for which A324274(k) is 0, i.e., starting squares in A324274 that yield a path of infinite length.

Note that the sequence up to a(n) (for its current known values) is actually the path of a(n) in reverse until it reaches square 2. It is therefore conjectured that all starting squares in A324274 either have a finite length or are part of this single sequence. Conjectures from _Colin Barker_, Mar 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jan Koornstra_, Feb 27 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014224 | OEIS-A324278

Exponential untouchable numbers: numbers that are not the sum of aliquot exponential divisors of any number.

The terms are conjectural and based on a search for solutions to esigma(x) - x = k for k in the range of the data section and x < 10^12 (esigma(x) - x = A051377(x) - x = A126164(x) is the sum of aliquot exponential divisors of x). - _Amiram Eldar_, Jan 22 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Feb 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014225 | OEIS-A324297

Positive integers k that are the product of two integers ending with 6.

Conjecture: lim_{n->oo} a(n)/a(n-1) = 1. The conjecture is true since it can be proved that a(n) = (sqrt(a(n-1)) + g(n-1))^2 where [g(n): n > 1] is a bounded sequence of positive real numbers. - _Stefano Spezia_, Aug 18 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Mar 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014226 | OEIS-A324298

Positive integers k such that 10*k+6 is equal to the product of two integers ending with 6 (A324297).

Conjecture: lim_{n->oo} a(n)/a(n-1) = 1. The conjecture is true since a(n) = (A324297(n) - 6)/10 and lim_{n->oo} A324297(n)/A324297(n-1) = 1. - _Stefano Spezia_, Aug 21 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Mar 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014227 | OEIS-A324300

G.f.: Sum_{n>=0} x^n * (x^(n+1) + i)^n / (1 + i*x^(n+1))^(n+1).

Odd terms occur only at n = k*(k+2) for k >= 0 (conjecture);

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014228 | OEIS-A324309

Smallest number m such that m is not a multiple of 10 and m^n has two or more identical adjacent decimal digits.

Conjecture: a(n) = 2 for n > 126, i.e., 2^n has two or more identical adjacent decimal digits for n > 126.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Feb 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014229 | OEIS-A324385

Distance from the n-th highly composite number, A002182(n), from the largest prime <= A002182(n).

Like in A141345 it appears (or is conjectured) that no composite numbers ever occur here. Taken together, this leads to McEachen's conjecture given in A117825. Here in range 2..10000 term 1 occurs for 313 times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014230 | OEIS-A324478

a(n) = (6/((n+1)*(n+2)*(n+3))) * multinomial(4*n;n,n,n,n).

(11!/12)*a(n) is divisible by ((n + 1)*(n + 2)*(n + 3))^2 for all n. More generally, we conjecture that there is a constant C(r) such that C(r)*(4*n)!/(n!*(n+r)!^3) is an integer for all n. Calculation suggests that we may take C(r) = (1/8)*(4*r)!/r! for r >= 1. - _Peter Bala_, Feb 28 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 10 2019, following a suggestion from Luis Fredes and Avelio Sepulveda | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014231 | OEIS-A324480

Consider the numbers on the x-axis in the spiral shown in A274641; a(n) is the distance from n to the origin, or -1 if n never appears on the x-axis.

It is conjectured that every nonnegative number appears on the x-axis exactly once.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014232 | OEIS-A324481

Index of first occurrence of n in the spiral shown in A274641.

Conjecture: a(n) = c(n)*n^2 with 0.32 <= c(n) <= 1 for all n, maybe lim c(n) ~ 0.4. - _M. F. Hasler_, Feb 01 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014233 | OEIS-A324601

Unique solution x of the congruence x^2 = -1 (mod m(n)), with m(n) = A002559(n) (Markoff numbers) in the interval [1, floor(m(n)/2)], assuming the Markoff uniqueness conjecture, for n >= 3.

Unique solution x of the congruence x^2 = -1 (mod m(n)), with m(n) = A002559(n) (Markoff numbers) in the interval [1, floor(m(n)/2)], assuming the Markoff uniqueness conjecture, for n >= 3. See the Aigner reference, Corollary 3.17., p. 58. If this congruence is solvable uniquely for integer x in the given interval then the Markoff uniqueness conjecture is true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jul 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014234 | OEIS-A324615

G.f. A(x) satisfies: 1 = Sum_{n>=0} x^n / A(x)^(n+1) * Product_{k=1..n} (1 + k*x).

a(n) is odd only when n is of the form 2^k or 3*2^k, or equal to 0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 19 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014235 | OEIS-A324643

Numbers k such that bitand(2k,sigma(k))/2 = k = bitand(k,sigma(k)-k), where bitand is bitwise-AND, A004198.

It is conjectured that there are no odd terms in this sequence, which is equivalent to the conjecture that there are no odd perfect numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014236 | OEIS-A324713

a(n) = 2*A156552(n) XOR A323243(n).

It is conjectured that a(n) may obtain value zero only when n is a power of prime, and especially for n > 1, it must be a prime power present in A324201.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014237 | OEIS-A324794

Conjectured value of N(2^n), defined by property that the Jensen polynomial J^{2^n,m}_p(X) is hyperbolic for m >= N(2^n).

Conjectured value of N(2^n), defined by property that the Jensen polynomial J^{2^n,m}_p(X) is hyperbolic for m >= N(2^n). Normally the OEIS does not list conjectured values in the Data section, but an exception has been made here in view of the importance of these numbers. The conjectured values of N(d) for d = 1,...,9 are 1, 25, 94, 206, 381, 610, 908, 1269, 1701.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014238 | OEIS-A324863

Binary length of A324866(n), where A324866(n) = A156552(n) OR (A323243(n) - A156552(n)).

Provided that the maximal value that A324861(d) attains among divisors d of n is attained an odd number of times, then a(n) gives that maximal value. It is conjectured that this always holds. Among n = 1..10000, there are only two such cases, where the maximal value occurs more than once among the divisors: 3675 and 7623, where it occurs three times in both (see the examples). a(n) = Max_{d|n} A324861(d) [conjectured].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014239 | OEIS-A324916

Triangle read by rows: T(n,k) is the number of 3-stack-sortable permutations of [n] with k descents (0 <= k <= n-1).

Bona has proven that the polynomial Sum_{k=0..n-1} T(n,k)*x^k is always symmetric and unimodal. He has conjectured that it has only real roots.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Defant_, Mar 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014240 | OEIS-A324918

Partial sums of A324917.

Conjecture: lim_{n--> infinity}(log(a(n))/log(n)) exists and is positive. Conjecture: a(n) >= A324965(n) for all n >= 13.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin Defant_, Mar 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014241 | OEIS-A324969

Number of unlabeled rooted identity trees with n vertices whose non-leaf terminal subtrees are all different.

The conjecture of Mathar is true. Proof: Given a rooted identity tree T, a terminal subtree T' with more than one vertex contains at least one edge that is also a terminal subtree of T'. Thus, if T has more than one branch with more than one vertex, then it fails the additional condition since it would have at least two non-leaf terminal subtrees (namely edges) that are the same. Also, T can't have under its root more than one branch with exactly one vertex because it is an identity tree. Now we know that under the root of T is exactly one branch of the same kind as T or else it has exactly one other branch with exactly one vertex. The leads immediately to the same recurrence as A000045 the Fibonacci sequence except for n=3. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Mar 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014242 | OEIS-A325023

Multiperfect numbers m from A007691 such that m*(m-tau(m))/sigma(m) is an integer h, where k-tau(k) is the number of nondivisors of k (A049820) and sigma(k) is the sum of the divisors of k (A000203).

Conjecture: Numbers m such that all values of sigma(m)/m, m*tau(m)/sigma(m) and m*(m-tau(m))/sigma(m) are any integers (f, g, and h respectively). Corresponding values of integers f: 1, 2, 2, 2, 3, 2, 4, 4, 4, 4, 2, 4, 4, 4, 2, ... Corresponding values of integers g: 0, 1, 11, 243, 216, 4057, 7536, 8166, ... Corresponding values of integers h: 1, 2, 3, 5, 8, 7, 24, 24, 54, 80, 13, 96, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Mar 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014243 | OEIS-A325173

Perfect squares of the form a + b^2 + c^3, where a,b,c are consecutive numbers.

E.g.f.: exp(x)*x*(9 + 63*x + 114*x^2 + 69*x^3 + 15*x^4 + x^5). - Conjectured by _Stefano Spezia_, Sep 05 2019 after _Colin Barker_ Above conjectures are true. Proof: k + (k+1)^2 + (k+2)^3 = (k+1)*(k+3)^2 and thus is a perfect square if and only if k+1 = n^2 is a perfect square. This implies that (k+1)*(k+3)^2 = n^2*(n^2+2)^2. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philip Mizzi_, Sep 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014244 | OEIS-A325260

Number of integer partitions of n whose omega-sequence covers an initial interval of positive integers.

Conjectures from _Chai Wah Wu_, Jan 13 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Apr 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014245 | OEIS-A325269

Number of integer partitions of n with 2 distinct parts or at least 3 parts.

conjecture: a(n) = A000041(n) - A000034(n-1), n>0. - _R. J. Mathar_, Dec 13 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Apr 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014246 | OEIS-A325286

G.f. satisfies: 1 = Sum_{n>=0} (1+x)^(n*(n-1)/2) / A(x)^n * 1/2^(n+1).

Conjecture: a(n) == 1 (mod 3) iff n is a number whose base-3 representation contains no 2 (cf. A005836), otherwise a(n) == 0 (mod 3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014247 | OEIS-A325350

Number of integer partitions of n whose augmented differences are weakly decreasing.

G.f.: Sum_{k>=0} x^k / Product_{j=1..k} (1 - x^(j*(j+1)/2)) (conjecture). - _Ilya Gutkovskiy_, Apr 25 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Apr 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014248 | OEIS-A325423

Numbers k such that sigma(2*k+1) >= sigma(2*k).

Kobayashi and Trudgian (2020) proved that the asymptotic density of this sequence exists, and that it is in the interval [0.0539171, 0.0549445]. They calculated the number of terms that do not exceed 10^k, for k = 3..9: 60, 551, 5490, 54603, 546879, 546537, and 54665173, and conjectured that the density equals 0.0546... . - _Amiram Eldar_, Aug 06 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giuseppe Melfi_, Sep 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014249 | OEIS-A325434

Row sums of A325433.

Conjecture: lim_{n->oo} a(n)/A000041(n) = 1/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Apr 27 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014250 | OEIS-A325444

Difference sequence of A325417.

See A325417 for a guide to related sequences. Conjecture: all the differences are in {1,2,3,4}; a count of differences d(n) = a(n)-a(n-1) for n=2..10000 follows: 4755 occurrences of d(n) = 1; 4332 of 2; 744 of 3; and 169 of 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014251 | OEIS-A325445

Difference sequence of A325418.

See A325417 for a guide to related sequences. Conjecture: all the differences are in {1,2,3,4,6}; 5 does not occur; a count of differences d(n) = a(n)-a(n-1) for n=2..10000 follows: 1082 occurrences of d(n) = 1; 2693 of 2; 744 of 3; 1609 of 4, 0 of 5, and 198 of 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014252 | OEIS-A325458

Triangle read by rows where T(n,k) is the number of integer partitions of n with largest hook of size k, i.e., with (largest part) + (number of parts) - 1 = k.

Conjectured to be equal to A049597. Franklin T. Adams-Watters has conjectured at A049597 that the k-th column gives the coefficients of the sum of Gaussian polynomials [k,m] for m = 0..k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, May 04 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014253 | OEIS-A325480

a(n) is the largest integer m such that the product of n consecutive integers starting at m is divisible by at most n primes.

Each term is only conjectured and has been verified up to 10^6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Onno M. Cain_, Sep 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014254 | OEIS-A325494

Difference sequence of A325419.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014255 | OEIS-A325495

Difference sequence of A325420.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3,4}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014256 | OEIS-A325496

Difference sequence of A077477.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3,4}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014257 | OEIS-A325497

Difference sequence of A325422.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3,4,6}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014258 | OEIS-A325498

Difference sequence of A036668.

Conjecture: every term is in {1,2,3,4}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014259 | OEIS-A325499

Difference sequence of A325424.

Conjecture: every term is in {1,2,3,4,5,6,7}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014260 | OEIS-A325518

Difference sequence of A325425.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3,4,5,6}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014261 | OEIS-A325519

Difference sequence of A325426.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3,4,5,6,7}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014262 | OEIS-A325520

Difference sequence of A325427.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3,4,5,6}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014263 | OEIS-A325522

Difference sequence of A325429.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014264 | OEIS-A325523

Difference sequence of A325430.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3,5}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014265 | OEIS-A325524

Difference sequence of A325431.

Conjecture: every term is in {1,2,3}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014266 | OEIS-A325525

Difference sequence of A325432.

Conjecture: every term is in {1,2,3,4,5,6,7,9}; note that 8 is missing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014267 | OEIS-A325526

Difference sequence of A325462.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3,4}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014268 | OEIS-A325527

Difference sequence of A325463.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3,4,6}; 5 is missing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014269 | OEIS-A325528

Difference sequence of A325464.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3,4}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014270 | OEIS-A325529

Difference sequence of A325465.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3,4,5}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014271 | OEIS-A325530

Difference sequence of A325440.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3,4}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014272 | OEIS-A325531

Difference sequence of A325441.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3,4,6}; note that 5 is missing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014273 | OEIS-A325532

Difference sequence of A325442.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014274 | OEIS-A325533

Difference sequence of A325443.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3,4}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014275 | OEIS-A325541

Difference sequence of A325539.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3,4}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014276 | OEIS-A325542

Difference sequence of A325540.

See A325417 for a guide to related sequences. Conjecture: every term is in {1,2,3,4,5,6,7}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014277 | OEIS-A325597

a(n) is the least number not 2a(m) + a(m-1) for any m < n.

Conjectures: Let d(n) = 3a(n) - 4n; then d(n) is bounded, and d(n) = 0 for infinitely many n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014278 | OEIS-A325618

Numbers m such that there exists an integer partition of m whose reciprocal factorial sum is 1.

Conjecture: 137 is the greatest integer not in this sequence. - _Charlie Neder_, May 14 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, May 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014279 | OEIS-A325629

Floor of number of n-dimensional degrees in an n-sphere.

The sequence has a maximum value at n = 20626 with a value of 1.3610489172...*10^4479, too large to be written here. I conjecture that the peak value of the function analytically is somewhere near 64800/Pi = 20626.48062...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eliora Ben-Gurion_, Sep 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014280 | OEIS-A325638

Numbers m such that sigma(m) can be obtained as the base-2 carryless product of 2m and some k.

Conjecture: all terms are even. If this is true, then there are no odd perfect numbers. See also conjectures in A325639 and in A325808.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, May 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014281 | OEIS-A325686

Number of strict length-3 compositions x + y + z = n satisfying x + y != z and x != y + z.

Conjectures from _Colin Barker_, May 14 2019: (Start) Above conjecture confirmed for n <= 5000. - _Fausto A. C. Cariboni_, Feb 17 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, May 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014282 | OEIS-A325688

Number of length-3 compositions of n such that every distinct consecutive subsequence has a different sum.

Conjectures from _Colin Barker_, May 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, May 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014283 | OEIS-A325689

Number of length-3 compositions of n such that no part is the sum of the other two.

Conjectures from _Colin Barker_, May 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, May 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014284 | OEIS-A325690

Number of length-3 integer partitions of n whose largest part is not the sum of the other two.

Conjectures from _Colin Barker_, May 15 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, May 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014285 | OEIS-A325691

Number of length-3 integer partitions of n whose largest part is not greater than the sum of the other two.

Conjectures from _Colin Barker_, May 16 2019: (Start) The above conjectured formulas are true. - _Stefano Spezia_, May 19 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, May 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014286 | OEIS-A325695

Number of length-3 strict integer partitions of n such that the largest part is not the sum of the other two.

Conjectures from _Colin Barker_, May 15 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, May 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014287 | OEIS-A325696

Number of length-3 strict compositions of n such that no part is the sum of the other two.

Conjectures from _Colin Barker_, May 16 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, May 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014288 | OEIS-A325804

Positions of nonzero terms of Product_{k=0..floor(log_2(n))} (1 + A004718(floor(n/(2^k)))).

Conjecture: a(n) - a(n-1) belongs to A094373. - _Mikhail Kurkov_, Feb 20 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, May 22 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014289 | OEIS-A325808

Numbers n such that sigma(n) can be obtained as the base-3 carryless product of 2n and some k.

Conjecture: If we select only those n of these for which sigma(n) >= 2n, then we get a subsequence which contains only even terms: 6, 28, 120, 496, 8128, 18990, 667296, 2097414, 33550336, etc. If this is true, then there are no odd perfect numbers. See also conjectures in A325638 and A325639.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, May 22 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014290 | OEIS-A325812

Numbers k such that gcd(A034448(k)-k, k-A048146(k)) is equal to abs(k-A048146(k)).

Conjecture: after the initial one all other terms are even. If this holds then there are no odd perfect numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, May 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014291 | OEIS-A325922

Number of Motzkin excursions of length n with an even number of humps and an even number of peaks.

conjecture: a(n)+A325924(n) = A307557(n). - _R. J. Mathar_, Jan 25 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrei Asinowski_, Jun 27 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014292 | OEIS-A325945

Let p(n) be the n-th composite squarefree number. a(n) is the smallest integer q that forms a pure idempotent product.

We conjecture that for all positive composite squarefree integers p, there exists a q such that pq is a pure idempotent product. Conjecture verified for all squarefree composite p < 2^15. The largest q correspond to the cases where p-1 is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Barry Fagin_, Sep 09 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014293 | OEIS-A325983

Row sums of the triangle A325982.

Conjecture: a(n) ~ 2^(n-2). - _Stefano Spezia_, Jun 16 2021 This conjecture is true. Its proof follows from _Vaclav Kotesovec_'s formula used in the 2nd Mathematica code. - _Stefano Spezia_, Jun 27 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, May 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014294 | OEIS-A326042

a(n) = A064989(sigma(A003961(n))), where A003961 shifts the prime factorization of n one step towards larger primes, and A064989 shifts it back towards smaller primes.

Conjecture: There are no other fixed points than a(1) = 1. If true, then there are no odd perfect numbers. This condition is equivalent to the condition that if A161942 has no fixed points larger than one, then there are no odd perfect numbers. This follows as whenever k is a fixed point, that is, a(k) = k, then we should also have A003961(a(k)) = A003961(A064989(sigma(A003961(k)))) = A161942(A003961(k)) = A003961(k). Note that A003961 is an injective and surjective mapping from natural numbers to odd numbers, A064989 is its (left) inverse, and composition A003961(A064989(n)) is equivalent to A000265(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jun 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014295 | OEIS-A326097

E.g.f. A(x) satisfies: 1 = Sum_{n>=0} ((1+x)^n - A(x))^n / n!.

a(4*n+2) = 1 (mod 2), otherwise the terms a(k) are even for k > 3 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014296 | OEIS-A326169

Iterations where ants mate in male/female Langton's ant variation.

Conjecture: a(n) = n + 128 for n >= 28. - _Jinyuan Wang_, Jul 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Jun 09 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014297 | OEIS-A326181

Numbers n for which sigma(sigma(n)) = 3*sigma(n).

If, as conjectured, A005820 has 6 terms, then this sequence is finite and has 756 terms. - _Giovanni Resta_, Jun 17 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jun 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014298 | OEIS-A326244

Number of labeled n-vertex simple graphs without crossing or nesting edges.

Conjectures from _Colin Barker_, Jun 28 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jun 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014299 | OEIS-A326247

Number of labeled n-vertex 2-edge multigraphs that are neither crossing nor nesting.

Conjectures from _Colin Barker_, Jun 21 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jun 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014300 | OEIS-A326251

Number of digraphs with vertices {1..n} whose increasing edges are not crossing.

Conjecture: Also the number of non-nesting digraphs with vertices {1..n} whose increasing edges are not crossing, where two edges (a,b), (c,d) are nesting if a < c < d < b or c < a < b < d.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jun 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014301 | OEIS-A326278

Number of n-vertex, 2-edge multigraphs that are not nesting. Number of n-vertex, 2-edge multigraphs that are not crossing.

Conjectures from _Colin Barker_, Jun 25 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jun 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014302 | OEIS-A326294

Number of connected simple graphs on a subset of {1..n} with no crossing or nesting edges.

Conjecture: a(n) = A052161(n - 2) + 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jun 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014303 | OEIS-A326326

T(n, k) = [x^k] Sum_{j=0..n} Pochhammer(x, j), for 0 <= k <= n, triangle read by rows.

Conjecture: T(n,k) = Sum_{i=0..n} A132393(i,k) for 0 <= k <= n. - _Werner Schulte_, Mar 30 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jul 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014304 | OEIS-A326367

Number of tilings of an equilateral triangle of side length n with unit triangles (of side length 1) and exactly two unit "lozenges" or "diamonds" (also of side length 1).

a(n) = (3/8)*(n-2)*(n-1)*(3*n^2 + 3*n - 4) (conjectured by _R. J. Mathar_, proved by _Greg Dresden_ and E. Sijaric).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Greg Dresden_, Jul 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014305 | OEIS-A326390

The number of ways of seating n people around a table for the second time so that k pairs are maintained. T(n,k) read by rows.

Weighted average of each row using k as weights converges to 2 for large n and is given with following formula: (Sum_{k} T(n,k)*k)/n! = 2/(n-1) + 2 (conjectured). T(n,n-2) = n^2*(n-3) for n > 3 (conjectured); T(n,n-3) = (3/4)*n^4 + 6*n^3 + (2/3)*n^2 - 14*n + 6 for n > 4 (conjectured); T(n,n-4) = (25/12)*n^5 + (73/6)*n^4 + (5/4)*n^3 - (253/6)*n^2 + (152/3)*n - 24 for n > 5 (conjectured); T(n,n-5) = (52/15)*n^6 + (77/3)*n^5 + 14*n^4 - (194/3)*n^3 + (4628/15)*n^2 - 273*n + 130 for n > 5 (conjectured); T(n,n-6) = (707/120)*n^7 + (2093/40)*n^6 + (2009/40)*n^5 - (245/8)*n^4 + (78269/60)*n^3 - (18477/10)*n^2 + (21294/10)*n - 684 for n > 6 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Witold Tatkiewicz_, Jul 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014306 | OEIS-A326397

Triangle T(n,k) read by rows: T(n,k) = number of ways of seating n people around a table for the second time so that k pairs are maintained. Reflected sequences are counted as one.

Weighted average of each row using k as weights converges to 2 for large n and is given by the following formula: (Sum_{k} T(n,k)*k)/(Sum_{k} T(n,k)) = 2/(n-1) + 2 (conjectured). T(n,n-3) = 1/2*n^3 + 3/4*n^2 - 2 (conjectured); T(n,n-3) = (2/3)*n^4 + 3*n^3 + (1/3)*n^2 - 7*n + 3 for n > 4 (conjectured); T(n,n-4) = (25/24)*n^5 + (73/12)*n^4 + (5/8)*n^3 - (253/12)*n^2 + (76/3)*n - 12 for n > 5 (conjectured); T(n,n-5) = (26/15)*n^6 + (77/6)*n^5 + 7*n^4 - (97/3)*n^3 + (2314/15)*n^2 - 273/2*n + 65 for n > 5 (conjectured); T(n,n-6) = (707/240)*n^7 + (2093/80)*n^6 + (2009/80)*n^5 - (245/16)*n^4 + (78269/120)*n^3 - (18477/20)*n^2 + (10647/0)*n - 342 for n > 6 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Witold Tatkiewicz_, Aug 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014307 | OEIS-A326404

Triangle T(n,k) read by rows: T(n,k) = number of ways of seating n people around a table for the second time so that k pairs are maintained. Rotated sequences are counted as one.

Conjecture: The weighted average of each row using k as weights converges to 2 for large n and is given by the following formula: (Sum_{k} T(n,k)*k)/(Sum_{k} T(n,k)) = 2/(n-1) + 2. Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Witold Tatkiewicz_, Aug 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014308 | OEIS-A326405

Minesweeper sequence of positive integers arranged on a 2D grid along ascending antidiagonals.

a(n) < 5 (conjectured). Set of n such that a(n) = 4 is unbounded (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Witold Tatkiewicz_, Sep 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014309 | OEIS-A326406

Minesweeper sequence of positive integers arranged on a 2D grid along a triangular maze.

a(n) < 5 (conjectured). Set of n such that a(n) = 4 is unbounded (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Witold Tatkiewicz_, Oct 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014310 | OEIS-A326407

Minesweeper sequence of positive integers arranged on a 2D grid along a square array that grows by alternately adding a row at its bottom edge and a column at its right edge.

a(n) < 6 (conjectured). a(n) = 5 for n={6} (conjectured). Set of n such that a(n) = 4 is unbounded (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Witold Tatkiewicz_, Oct 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014311 | OEIS-A326408

Minesweeper sequence of positive integers arranged on a 2D grid along a square maze.

Set of n such that a(n) = 4 is unbounded (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Witold Tatkiewicz_, Oct 04 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014312 | OEIS-A326409

Minesweeper sequence of positive integers arranged on a 2D grid along Hamiltonian path.

Set of n such that a(n) = 4 is unbounded (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Witold Tatkiewicz_, Oct 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014313 | OEIS-A326410

Minesweeper sequence of positive integers arranged on a square spiral on a 2D grid.

Set of n such that a(n) = 4 is unbounded (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Witold Tatkiewicz_, Oct 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014314 | OEIS-A326411

Triangle T(n,k) read by rows: T(n,k) = the number of ways of seating n people around a table for the second time so that k pairs are maintained. Reflected and rotated sequences are counted as one.

T(n,n-4) = (25/24)*n^4 + (23/12)*n^3 - (169/24)*n^2 + (85/12)*n - 3 for n > 5 (conjectured); [see Poulet] T(n,n-5) = (26/15)*n^5 + (25/6)*n^4 - (83/6)*n^3 + (221/6)*n^2 - (299/10)*n + 13 for n > 5 (conjectured); [see Poulet] T(n,n-6) = (707/240)*n^6 + (2037/240)*n^5 - (413/16)*n^4 + (2233/16)*n^3 - (2777/15)*n^2 + (3739/20)*n - 57 for n > 6 (conjectured). [See Poulet]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Witold Tatkiewicz_, Aug 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014315 | OEIS-A326478

a(n) = n*denominator(n*Bernoulli(n-1))/denominator(Bernoulli(n-1)).

Empirical: a(2*n) = [x^n] x*(2/(x - 1)^2 - 1) for n >= 1, implying the conjecture that a(2*n) = A103517(n+1) and/or A272651(n). Conjectural, the odd fixed points > 1 of this sequence are A121707; in other words, for n > 1, denominator(n*Bernoulli(n-1)) = denominator(Bernoulli(n-1)) <=> n | Sum_{k=1..n-1} k^(n-1). (See the conjectures of _Thomas Ordowski_ in A121707.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jul 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014316 | OEIS-A326579

a(n) = n*denominator(n*Bernoulli(n-1)) for n >= 1 and a(0) = 0.

Conjecture: For n>1: denominator(Bernoulli(n-1)) = n*denominator(n*Bernoulli(n-1)) <=> n is Korselt <=> n is prime or n is Carmichael.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jul 17 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014317 | OEIS-A326582

A signed variant of A309132.

See A309132 for background and conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jul 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014318 | OEIS-A326584

a(n) = gcd(n*N(n-1), D(n-1)), with N(n)/D(n) = B(n) the n-th Bernoulli number.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jul 19 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014319 | OEIS-A326602

G.f.: Sum_{n>=0} (x^(2*n-1) + 1)^n * x^n / (1 + x^(2*n+1))^(n+1), an even function.

a(n) is odd iff n is square (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014320 | OEIS-A326604

G.f. A(x) satisfies x / Series_Reversion(x*A(x)) = (2*A(x) + 1+x)/3.

a(k) = 6 (mod 9) when k = A191107(n) for n > 1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014321 | OEIS-A326663

Column 3 of the array at A309157; see Comments.

This is the sequence (c(n)) defined by the complementary equation c(n) = a(n) + b(2n), with initial value a(1) = 1. See A309157. Conjecture: 7n-c(n) is in {1,2} for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014322 | OEIS-A326664

Column 3 of the array at A326661 see Comments.

This is the sequence (c(n)) defined by the complementary equation c(n) = a(n) + b(3n), with initial value a(1) = 1. See A326661. Conjecture: 9n-c(n) is in {1,2} for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014323 | OEIS-A326691

a(n) = n/denominator(Sum_{prime p | n} 1/p - 1/n).

a(n) = gcd(n, 1+((n-1)*A003415(n))). [Conjectured, after an empirical formula found by LODA miner. This holds at least up to n=2^27] - _Antti Karttunen_, Mar 15 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Jul 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014324 | OEIS-A326714

a(n) = binomial(n,2) + (2-adic valuation of n).

2^a(n) is conjectured to be the order of the smallest n-symmetric graph.

猜想/问题提及段 | historical_or_current_mention_unreviewed | Sebastian Jeon and _Tanya Khovanova_, Dec 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014325 | OEIS-A326781

No position of a 1 in the reversed binary expansion of n is a power of 2.

Conjectures from _Colin Barker_, Jul 27 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jul 25 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014326 | OEIS-A326805

a(n) = floor(Sum_{k>=0} n^sqrt(k) / Gamma(sqrt(k) + 1)), where Gamma is Euler's gamma function.

Conjecture: a(n) ~ 2*n*exp(n). - _Vaclav Kotesovec_, Sep 16 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014327 | OEIS-A326823

a(n) is the number of iterations needed to reach 1 or 11 starting at n and using the map k -> (k/2 if k is even, otherwise k + (smallest square > k)). Set a(n) = -1 if the trajectory never reaches 1 or 11.

It is conjectured that this algorithm will always terminate at 1 or 11. _Matthijs Coster_ verified the conjecture for n <= 100000. _Jim Nastos_ verified the conjecture for n <= 43*10^6 (Oct 21 2019).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ali Sada_, Oct 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014328 | OEIS-A326825

a(n) is the number of iterations needed to reach 1 or 5 starting at n and using the map k -> (k/2 if k is even, otherwise k + (smallest triangular number > k)). Set a(n) = -1 if the trajectory never reaches 1 or 5.

It is conjectured that this algorithm will always terminate at 1 or 5. _Jim Nastos_ verified the conjecture for n <= 354999. The conjecture holds for all n <= 10^9. - _Jon E. Schoenfield_, Oct 20 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ali Sada_, Oct 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014329 | OEIS-A326896

Prime numbers p such that 0 < pi(p;10,(9,1)) = pi(p;10,(1,7)) where pi(x;q,(a,b)) is the number of primes p_n <= x such that p_n == a (mod q) and p_(n+1) == b (mod q).

Lemke Oliver and Soundararajan conjectured that pi(x;3,(1,1)) < pi(x,3,(1,2)) and pi(x;4,(1,1)) < pi(x;4,(1,3)) are robust biases, i.e., they always hold. Here we conjecture that pi(x;10,(9,1)) > pi(x;10,(1,7)) is a semi- robust bias, i.e., it holds from a certain value x_0 on. Here x_0 is conjectured to be 17839. This conjecture seems to be confirmed by the observation that pi(x;10,(9,1)) - pi(x;10,(1,7)) ~ (Li(x)/16) * (1/2) * log(2*Pi/5*log(x)) / log(x).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _A.H.M. Smeets_, Sep 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014330 | OEIS-A326897

Prime numbers p such that 0 < pi(p;10,(9,1)) = pi(p;10,(3,9)) where pi(x;q,(a,b)) is the number of primes p_n <= x such that p_n == a (mod q) and p_(n+1) == b (mod q).

Lemke Oliver and Soundararajan conjectured that pi(x;3,(1,1)) < pi(x,3,(1,2)) and pi(x;4,(1,1)) < pi(x;4,(1,3)) are robust biases, i.e., they always hold. Here we conjecture that pi(x;10,(9,1)) > pi(x;10,(3,9)) is a semi- robust bias, i.e., it holds from a certain value x_0 on. Here x_0 is conjectured to be 48539. This conjecture seems to be confirmed by the observation that pi(x;10,(9,1)) - pi(x;10,(3,9)) ~ (Li(x)/16) * (1/2) * log(2*Pi/5*log(x)) / log(x).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _A.H.M. Smeets_, Sep 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014331 | OEIS-A326923

a(n) is the number of iterations needed to reach 1 or 9 starting at n and using the map k -> (k/2 if k is even, otherwise k + (largest triangular number < k)). Set a(n) = -1 if the trajectory never reaches 1 or 9.

It is conjectured that this algorithm will always terminate at 1 or 9. _Jim Nastos_ verified the conjecture for n <= 64*10^5. _Jim Nastos_ verified the conjecture for n <= 45248000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ali Sada_, Oct 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014332 | OEIS-A326925

Irregular triangular array read by rows: row n shows the coefficients of this polynomial of degree n: (1/n!)*(numerator of n-th derivative of (1-x)/(1-x-x^2)).

Conjecture: The odd degree polynomials are irreducible; the even degree (= 2k) polynomials have exactly two irreducible factors, each of degree k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 22 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014333 | OEIS-A326929

Numbers whose divisors and arithmetic mean of divisors are palindromic.

(4*10^A259050(n)-7)/3 and (8*10^A259050(n)-14)/3 are terms. Conjecture: all terms > 10 are of these forms. - _Chai Wah Wu_, Nov 17 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ivan N. Ianakiev_, Oct 22 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014334 | OEIS-A327033

N(p-1)/p + D(p-1)/p^2 with p the n-th prime and B(k) = N(k)/D(k) the k-th Bernoulli number.

a(n) is an integer, as conjectured by Thomas Ordowski and proved by the author in A309132 and A326690. Ordowski also conjectured that the sequence is a subsequence of A174341.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan Sondow_, Aug 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014335 | OEIS-A327094

a(n) is the number of cells in the smallest polyomino that can contain all free n-ominoes.

Conjecture: a(9) = 26, a(10) = 31, a(11) = 37, and a(12) = 43. Conjecture: a(n) = floor((n+1)^2/4) + 1 for n >= 9. This matches the conjectured values a(11) = 37, a(12) = 43. The formula floor((n+1)^2/4) = A002620(n+1) counts cells in the "staircase" polyomino with row widths n, n-2, n-4, ..., which is optimal for n <= 6 and n = 8 but requires one extra cell for n >= 9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Sep 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014336 | OEIS-A327119

Sequence obtained by swapping each (k*(2n))-th element of the nonnegative integers with the (k*(2n+1))-th element, for all k>0 in ascending order, omitting the first term.

a(n) = A004442(A327420(n)) (conjectured). - _Jon Maiga_, May 31 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jennifer Buckley_, Sep 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014337 | OEIS-A327123

Expansion of Sum_{k>=1} phi(k) * x^k / (1 + x^(2*k)), where phi = A000010.

Conjecture: a(n) = Sum_{k=1..n} sin(GCD(k,n) * Pi/2). - _Velin Yanev_ and _Vaclav Kotesovec_, Jun 01 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Sep 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014338 | OEIS-A327136

Numbers k such that sin(2k) > sin(2k+2) < sin(2k+4).

Conjecture: 1.285 < n*Pi - a(n) < 1.286 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014339 | OEIS-A327138

Numbers k such that cos(2k) < cos(2k+2).

Conjecture: 2.07 < n*Pi - a(n) < 3.08 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014340 | OEIS-A327239

For n >= 1, a(n) = b(n+1) - b(n) where b is A013947.

Stepwise answer to the stepwise question "How far ahead is the next 1 in the Kolakoski sequence, starting from A000002(1) = 1?" Sequence contains only 1s, 2s and 3s. Conjecture: Densities of 1s, 2s, and 3s -> 1/3 as n->infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Sep 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014341 | OEIS-A327295

Numbers k such that e(k) > 1 and k == e(k) (mod lambda(k)), where e(k) = A051903(k) is the maximal exponent in prime factorization of k.

Conjecture: there are no composite numbers m > 4 such that m == e(m) (mod phi(m)). By Lehmer's totient conjecture, there are no such squarefree numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Dec 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014342 | OEIS-A327297

Terms in A003656 that are not prime powers (A246655).

Conjecture: if D is a term of this sequence, then D = uv, where u, v are 4, 8 or primes congruent to 3 modulo 4. For example, a(1) = 12 = 3*4, a(2) = 21 = 3*7, a(3) = 24 = 3*8, a(4) = 28 = 4*7, a(5) = 33 = 3*11, ... [This conjecture is correct: see Theorem 1 and Theorem 2 of Ezra Brown link; see also A003656. - _Jianing Song_, Dec 28 2021] Also, it is conjectured that the quadratic field with discriminant D has form class number 2, where D is a term of this sequence. This is equivalent to the conjecture above. [This can also be deduced from the first paragraph of Ezra Brown link: the norm of the fundamental unit of the field k is -1 if D = 8 or a prime congruent to 1 modulo 4, and 1 if D is in this sequence. Here k is the quadratic field with discriminant D. - _Jianing Song_, Dec 28 2021]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Sep 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014343 | OEIS-A327315

Irregular triangular array read by rows: row n shows the coefficients of this polynomial of degree n: (1/n!)*(numerator of n-th derivative of (x-2)/(x^2-x+1)).

Conjecture: The numbers n for which the n-th polynomial is irreducible are given by A069353.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014344 | OEIS-A327330

"Concave pentagon" toothpick sequence (see Comments for precise definition).

The behavior seems to suggest that this sequence can be calculated with a formula, in the same way as A139250, but that is only a conjecture. Conjecture: a(2^k) = A327332(2^k), k >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Sep 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014345 | OEIS-A327332

"Concave pentagon" toothpick sequence, starting with a V-toothpick (see Comments for precise definition).

The behavior seems to suggest that this sequence can be calculated with a formula, in the same way as A139250, but that is only a conjecture. Conjecture: a(2^k) = A327330(2^k), k >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Sep 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014346 | OEIS-A327342

a(n) gives the number of distinct odd prime divisors of m(n) = A002559(n) (Markoff numbers).

For more details see A327343(n) = 2^a(n). The Frobenius-Markoff uniqueness conjecture on ordered triples with largest member m(n) is certainly true for m(n) if a(n) = 0 (so-called singular cases) or 1. See the Aigner reference, p. 59, Corollary 3.20, for n >= 3 (the a(n) = 1 cases).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Sep 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014347 | OEIS-A327343

a(n) gives the number of representative parallel primitive forms for binary quadratic forms of discriminant Disc(n) = 9*m(n)^2 - 4 and representation of -m(n)^2, with m(n) = A002559(n) (Markoff numbers).

The Frobenius-Markoff uniqueness conjecture is certainly true for m(n) if a(n) = 1 or a(n) = 2. In the latter case the two rpapfs have to be equivalent to the principal form F_p(n), because the known solution implied by the ordered triple MT(n) = (x(n), y(n), m(n)) has an unordered partner solution which after ordering becomes (x(n), y'(n), m(n')) with y'(n) = m(n) and m(n') = 3*x(n)*m(n) - y(n) >= m(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Sep 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014348 | OEIS-A327345

Irregular triangle read by rows: Maximal numbers of the Markoff triples at level L of the Markoff tree, with members of the triples ordered increasingly.

The Frobenius-Markoff uniqueness conjecture is: each member of A002559 appears precisely once as maximal number of some triple in MT. Hence it is the conjectured that in MTreemax each member of A002559 appears only once.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Sep 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014349 | OEIS-A327421

In a Kolakoski n-chain, point at which term of penultimate sequence seq(n-1) differs from term of final sequence seq(n) in chain, when terms of seq(i) are run-lengths of seq(i+1) and the chain contains n sequences.

Conjectures: 1) In a Kolakoski n-chain of the form given, as n -> infinity, seq(n) converges on the Kolakoski sequence, A000002, whose terms always match its own run-lengths, while seq(1) converges on the anti-Kolakoski sequence, A049705, whose terms never match its own run-lengths. 2) As i -> infinity, a(i) / a(i+1) converges on 2/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anthony Sand_, Nov 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014350 | OEIS-A327444

a(n) is the maximum absolute value of the coefficients of the quotient polynomial R_(prime(n)#)/Product_{j=1..n} R_(prime(j)), where prime(n)# is the n-th primorial number A002110(n) and R_k = (x^k - 1)/(x - 1).

Conjecture: a(n) = exp((6n - 13 + (-1)^n)/8), approximately.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick A. Thomas_, Sep 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014351 | OEIS-A327463

a(1) = 1; thereafter, a(n) is the reversal of the next prime after a(n - 1) in base 6 if n is prime, and the reversal of the next composite after a(n - 1) in base 6 if n is composite.

Conjecture: a(n) <= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 27 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014352 | OEIS-A327464

a(1) = 1; thereafter, a(n) is the reversal of the next prime after a(n - 1) in base 5 if n is prime, and the reversal of the next composite after a(n - 1) in base 5 if n is composite.

Conjecture: the maximal value is 107, first attained at n = 883.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 27 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014353 | OEIS-A327468

Numbers m that divide 8^m + 7.

Conjecture: For k > 1, k^m == 1-k (mod m) has an infinite number of positive solutions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Oct 04 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014354 | OEIS-A327551

Number of cycles in the perfect shuffle permutation mapping (1,2,...,2n) to (1,n+1,2,n+2,...,n,2n).

Conjecture: a(n) = A006694(n-1) + 2 = A081844(n-1) + 1. - _N. J. A. Sloane_, Sep 16 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Sep 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014355 | OEIS-A327565

Number of transfers of marbles between two sets until the first repetition.

Conjecture: for m > 2, the first occurrence of a(n) = m is for n = 2^(m-3) + 1. For m > 1, first occurrence of a(n) = m is for n = A094373(m-1) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tristan Cam_, Sep 17 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014356 | OEIS-A327568

Exponent of the group GL(2, Z_n).

Conjecture: a(p^e) = (p^2-1)*p^e for primes p. If this is true, then 24 divides a(n) for n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Sep 17 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014357 | OEIS-A327569

Exponent of the group SL(2, Z_n).

Conjecture: a(p^e) = (p^2-1)*p^e/2 for primes p > 2 and 3*2^e for p = 2. If this is true, then 12 divides a(n) for n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Sep 17 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014358 | OEIS-A327593

Numbers m where an integer b that is a power of two > 2 with 1 < b < m exists such that m is a base-b repdigit.

Conjecture 1: Let c(n) = A001220(n)-1. This sequence is a supersequence of c. Conjecture 2: This is a supersequence of A240719.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Sep 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014359 | OEIS-A327597

a(n) = numerator((a(n-1) + a(n-2) + 1)/a(n-1)), with a(1)=1, a(2)=2.

As n goes to infinity, 32 < (a(n+1) + a(n))/(a(n+1) - a(n)) < 32.35 (conjectured). The average reduction 0.621 < a(n)/(a(n-1)+a(n-2)+1) < 0.622 (conjectured); the average reduction is the average of the individual reductions (The change from the numerator to the numerator in the simplest form):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vimal Vinod_, Sep 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014360 | OEIS-A327621

Sums of distinct powers of 3 and powers of 4 (greater than 1).

These gaps are closely related to the gaps in the set where 3^0 and 4^0 are (both) also allowed to be in the sum, in which case the first missing numbers are A367090 = (62, 63, 143, 144, 207, ...), see also Melfi's paper. It is obvious that the study of these gaps is crucial for the proof of Erdös conjecture. For A(x) the enumerating function, Erdős conjectured that A(x) > c*x.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giuseppe Melfi_, Sep 19 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014361 | OEIS-A327652

Intersection of A099011 and A327651.

They conjectured that this sequence is infinite, and their conjecture was proved by Grantham (2021).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Sep 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014362 | OEIS-A327692

Number of length-n phone numbers that can be dialed by a chess knight on a 0-9 keypad that starts on any number and takes n-1 steps.

Conjectures from _Colin Barker_, Oct 01 2019: (Start) Moreover, the matrix A is diagonalizable over the reals, with one zero eigenvalue, say L(0)=0. Combining this with the formula for the grand sum of a diagonalizable matrix in term of its eigenvalues, the above conjecture reduces to checking an algebraic condition on the nonzero eigenvalues L(1), ..., L(8) of A. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Derek Lim_, Sep 22 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014363 | OEIS-A327701

Maximum value for base-n variant of A326344.

The obvious conjecture is that a(n) = n-1 iff n-1 is 1 or an odd prime. - _N. J. A. Sloane_, Oct 02 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Dougherty-Bliss_, Sep 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014364 | OEIS-A327722

Number T(m,n) of permutations of [n] avoiding the consecutive pattern 12...(m+1)(m+3)(m+2), where m, n >= 0; array read by ascending antidiagonals.

Conjecture: T(m, m + 4) = A242569(m + 4) = (m + 4)! - 2*(m + 4) for m >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, Nov 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014365 | OEIS-A327777

Prime numbers whose binary indices have integer mean and integer geometric mean.

Conjecture: This sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Sep 27 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014366 | OEIS-A327836

Least k > 0 such that n^k == 1 (mod (n+1)^(n+1)).

Conjecture: a(n) <= 2*(n+1)^n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthias Baur_, Sep 27 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014367 | OEIS-A327859

a(n) = A276086(A003415(n)), where A003415 is the arithmetic derivative, and A276086 is the primorial base exp-function.

Conjecture: If n is in A013929, that is, if n is not squarefree, then a(n) does not divide n. See A370115. - _Antti Karttunen_, Aug 31 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014368 | OEIS-A327883

a(0)=a(1)=0; thereafter a(n+1) is the index of the earliest as yet unused occurrence of a(n) if a(n) has occurred before; otherwise a(n+1) = a(a(n)-1). Once an occurrence of a(n) has been used it cannot be used again.

There appears to be a proper copy subsequence given by a(4*m+2) = a(m-1); m >0 (noticed by Carl J Love). There may be other (independent) copies to be found (e.g. by selecting terms sequentially, one from each of the above mentioned intervals). A family of sequences similar to this one can be described using the same rule as above, with offset k >= 0 and initial terms a(k) = a(k+1) = k. Conjecture: Every such sequence contains a proper copy of itself as a(4*m+2+k) = a(m+k-1). Conjectured formulae:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Oct 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014369 | OEIS-A327931

Lexicographically earliest infinite sequence such that for all i, j, a(i) = a(j) => A327930(i) = A327930(j).

a(i) = a(j) => A101296(i) = A101296(j). [Conjectural, see notes above and in A319357]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014370 | OEIS-A327943

Numbers m that divide 6^m + 5.

Conjecture: For k > 1, k^m == 1 - k (mod m) has infinitely many positive solutions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Sep 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014371 | OEIS-A327966

Number of iterations of "tamed variant of arithmetic derivative", A327965 needed to reach 0 from n, or -1 if zero is never reached.

Conjecture: from all n, zero is eventually reached.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014372 | OEIS-A327967

Positions of records in A327966.

Starting offset is zero to align with the indexing used in A189760, as this is conjecturally also the least k such that A327966(k) = n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014373 | OEIS-A327969

The length of a shortest path from n to zero when using the transitions x -> A003415(x) and x -> A276086(x), or -1 if no zero can ever be reached from n.

When I created this sequence, I conjectured that by applying two simple arithmetic operations "arithmetic derivative" (A003415) and "primorial base exp-function" (A276086) in some combination, and starting from any positive integer, we could always reach zero (via a prime and 1). At the first sight it seems almost certain that the conjecture holds, as it is always possible at every step to choose from two options (which very rarely meet, see A351088), leading to an exponentially growing search tree, and also because A276086 always jumps out of any dead-end path with p^p-factors (dead-end from the arithmetic derivative's point of view). However, it should be realized that one can reach the terms of either A157037 or A327978 with a single step of A003415 only from squarefree numbers (or respectively, cubefree numbers that are not multiples of 4, see A328234), and in general, because A003415 decreases the maximal exponent of the prime factorization (A051903) at most by one, if the maximal exponent in the prime factorization of n is large, there is a correspondingly long path to traverse if we take only A003415-steps in the iteration, and any step could always lead with certain probability to a p^p-number. Note that the antiderivatives of primorials with a square factor seem quite rare, see A351029. For all of the above reasons, I now conjecture that there are natural numbers from which it is not possible to reach zero with any combination of steps. For example 128 or 5^5 = 3125.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014374 | OEIS-A327975

Breadth-first reading of the subtree rooted at 5 of the tree where each parent node is the arithmetic derivative (A003415) of all its children.

Question: Does this subtree contain infinitely long paths? How many? Cf. conjecture number 8 in Ufnarovski and Ahlander paper, and a similar tree starting from 7, A327977.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014375 | OEIS-A327977

Breadth-first reading of the subtree rooted at 7 of the tree where each parent node is the arithmetic derivative (A003415) of all its children.

Question: Does this subtree contain infinitely long paths? How many? Cf. conjecture number 8 in Ufnarovski and Ahlander paper. As an example of possible beginning of such a sequence they give: 1 ← 7 ← 10 ← 25 ← 46 ← 129 ← 170 ← 501 ← 414 ← 2045.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014376 | OEIS-A327978

Numbers whose arithmetic derivative (A003415) is a primorial number (A002110) > 1.

Conjecture: No multiples of 5 occur in this sequence, and no multiples of 3 after the initial 9. Of the first 10000 terms, all others are semiprimes (with 9 the only square one), except 1547371 = 7^2 * 23 * 1373 and 79332523 = 17^2 * 277 * 991, the latter being the only known term whose decimal expansion ends with 3. If all solutions were semiprimes p*q such that p+q = A002110(k) for some k > 1 (see A002375), it would be a sufficient reason for the above conjecture to hold. - _David A. Corneth_ and _Antti Karttunen_, Oct 11 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 09 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014377 | OEIS-A327995

Decimal expansion of Gamma(3/4)/Pi^(1/4).

Conjecture: Equals Sum_{k>=0} A004402(k) / exp(k*Pi). - _Simon Plouffe_, Sep 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Oct 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014378 | OEIS-A328002

a(n) = 2^n * Sum_{k=0..n} Product_{j=1..k} (2/j)^((-1)^j).

Conjecture: d(n) is periodic with period 12 and repetition pattern (1, 2, 3, 3, 3, 2, 1, 1, 3, 6, 3, 1). - _Peter Luschny_, Jan 10 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Oct 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014379 | OEIS-A328005

Number of distinct coefficients in functional composition of 1 + x + ... + x^(n-1) with itself.

Conjectured g.f.: (x^7 - x^6 - x^5 + 2*x^3 + 1)*x/((x + 1)*(1 - x)^3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Reshetnikov_, Oct 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014380 | OEIS-A328010

The 5x + 1 sequence beginning at 17.

Alex V. Kontorovich & Jeffrey C. Lagarias conjectured that there are very few periodic orbits. One of them is shown here.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antoine Beaulieu_, Oct 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014381 | OEIS-A328029

Lexicographically earliest permutation of [1,2,...,n] maximizing the determinant of an n X n circulant matrix that uses this permutation as first row, written as triangle T(n,k), k <= n.

For n <= 9 the corresponding circulant matrices are n X n Latin squares with maximum determinant A309985(n). It is conjectured that this also holds for n > 9. See Mathematics Stack Exchange link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Oct 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014382 | OEIS-A328030

Maximum determinant of a circulant n X n matrix whose rows are permutations of [1,2,..,n].

It is conjectured that A309985 is identical to this sequence. See Mathematics Stack Exchange link. The first rows of the corresponding circulant matrices are provided in A328029.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Oct 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014383 | OEIS-A328033

Numbers m that divide 7^m + 6.

Conjecture: For k > 1, k^m == 1 - k (mod m) has infinite number of positive solutions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Oct 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014384 | OEIS-A328034

a(n) = 3n minus the largest power of 2 not exceeding 3n.

Conjecture: iterations of a() stabilize at a power of 2, or at a 2-cycle (5*2^t,7*2^t).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Nov 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014385 | OEIS-A328040

a(n) is the number of integers b with 1 < b < p such that p = prime(n) is a base-b nonrepunit circular prime with at least two base-b digits.

Conjecture: a(n) > 0 for n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Oct 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014386 | OEIS-A328071

Difference triangle for A327460 read by upwards antidiagonals.

Conjecture: every positive number appears. (Probably false, see next comment. - _N. J. A. Sloane_, Oct 09 2019)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014387 | OEIS-A328078

Number of regions after n generations of Jim Conant's iterative dissection of a square.

Conjecture: The left half of the dissection at generation n is essentially the same as the whole of the dissection at generation n-1. To see this, take the dissection at generation n-1, rotate it counterclockwise by 90 degrees, then reflect it about a vertical line through its center, and compress it in the horizontal direction by a factor of 2. The result is essentially the left half of generation n. - _N. J. A. Sloane_, Oct 14 2019 Conjectures from _Colin Barker_, Oct 15 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 13 2019, based on emails from _Robert Fathauer_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014388 | OEIS-A328079

First differences of A328078.

Conjectured g.f.: (4*x^11-8*x^10+4*x^9+4*x^8-8*x^7+7*x^6-4*x^5-5*x^4+2*x^3-2*x^2+1) / ((x^2+1)*(2*x^2-1)*(4*x^7-4*x^6+4*x^4-2*x^3+2*x^2+x-1)). - _N. J. A. Sloane_, Oct 16 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014389 | OEIS-A328129

Finite cardinalities of equivalence classes of real intervals with respect to the symmetric transitive closure of R(x,y) = "x is an integer multiple of y".

The initially known sequence terms (n <= 20) have the following property: a(n) = 2 * a(n-1) or, for some i < n, a(n) = 2 * a(n-1) - a(i); the former case occurring if r_min(n) equals a prime number, the latter occurring if and only if r_min(n) = r_max(i). Having identified underlying reasons that look like the basis of a proof, the author conjectures this is true for all n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Munn_, Oct 04 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014390 | OEIS-A328133

Exponential (2,4)-perfect numbers: numbers m such that esigma(esigma(m)) = 4m, where esigma(m) is the sum of exponential divisors of m (A051377).

Conjecturally, a subsequence of A083207 (tested for the first 659 terms of this sequence). - _Ivan N. Ianakiev_, Oct 05 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Oct 04 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014391 | OEIS-A328138

Numbers m that divide 9^m + 8.

Conjecture: For n > 1, k^n == 1-k (mod n) has an infinite number of positive solutions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Oct 04 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014392 | OEIS-A328151

a(n) is the smallest nonnegative integer k where exactly n ordered pairs of positive integers (x, y) exist such that x^2 + y^2 = k.

Conjecture: a(2k) = A093195(k) for k >= 1, a(2k+1) = 2*A006339(k)^2 for k >= 0. - _Jon E. Schoenfield_, Jan 23 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Oct 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014393 | OEIS-A328190

Lexicographically earliest infinite sequence of distinct positive integers such that the sequence and its first differences have no values in common.

Conjecture from _N. J. A. Sloane_, Nov 04 2019: (Start) There is a similar conjecture for A328196. Call the three sets of conjectured terms S0, S1, and S3. The terms in S0 are == 0 or 3 mod 5; those are in S1 are == 1 or 2 mod 10; and those in S3 are == 7 mod 10. So the sets are disjoint, as required by the definition. This conjecture would imply that the points a(2t) lie on a line of slope 5/4 and the points a(2t+1) on a line of slope 5/2, as conjectured by _Peter Kagey_. (End) Let us DEFINE a sequence S by the conjectured formulas given here, and a sequence T by the conjectured formulas given in A328196. Then it is not difficult to prove that the first differences of S are given by T, and that the terms of S and T are disjoint.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Oct 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014394 | OEIS-A328196

First differences of A328190.

Conjecture from _N. J. A. Sloane_, Nov 05 2019: (Start) The conjecture would establish that the terms lie on two straight lines, of slopes +-5/4. There is a similar conjecture for A328190. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Oct 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014395 | OEIS-A328230

Numbers m that divide 3^(m + 1) + 1.

Conjecture: For k > 2, k^(m + 1) == -1 (mod m) has an infinite number of positive solutions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Oct 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014396 | OEIS-A328279

Smallest elements of the cycles of (iterations of) A329623: n -> |concat(sum of adjacent digits of n) - n|.

Conjectures from _Colin Barker_, Dec 05 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Dec 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014397 | OEIS-A328302

For n > 1, a(n) is the least number > 0 for which it takes n-2 steps to reach a squarefree number by applying arithmetic derivative (A003415) zero or multiple times. a(1) = 4 is the least number for which no squarefree number is ever reached.

The least number k such that A328248(k) = n. After the initial two terms, probably also the positions of records in A328248, that is, it is conjectured that the records in A328248 appear in order, with each new record one larger than previous.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014398 | OEIS-A328375

Numbers k such that the decimal expansion of 2^k contains the substring 777.

Conjecture: if n > 30536, then a(n) = n + 3623. - _Chai Wah Wu_, Oct 26 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eder Vanzei_, Oct 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014399 | OEIS-A328416

Smallest k such that (Z/mZ)* = C_2 X C_(2k) has exactly n solutions for m, or 0 if no such k exists, where (Z/mZ)* is the multiplicative group of integers modulo m.

Conjecture: a(n) > 0 for all n. That is to say, every number occurs in A328412.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Oct 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014400 | OEIS-A328417

Numbers k such that A328412(k) sets a new record; numbers k such that (Z/mZ)* = C_2 X C_(2k) has more solutions for m than all k' < k, where (Z/mZ)* is the multiplicative group of integers modulo m.

Conjecture: this sequence is infinite. That is to say, A328412 is unbounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Oct 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014401 | OEIS-A328418

Records in A328412.

Conjecture: this sequence is infinite. That is to say, A328412 is unbounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Oct 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014402 | OEIS-A328518

Number of terms k in A002182 that have A001222(k) = n.

The above hypothesis isn't true, e.g., t = A002182(815) is a counterexample. But it is surely true that a(n) > 0 for all n, see the graph of A112778. - _M. F. Hasler_, Jan 08 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David A. Corneth_, Jan 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014403 | OEIS-A328537

First members of a list of seven pairs connected with parity of sums of partition numbers in arithmetic progressions.

The seven pairs are [6,8], [8,12], [12,24], [15,40], [16,48], [20,120], [21,168]. The list is definite, but the conjecture is unproved. The conjecture asserts that

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014404 | OEIS-A328544

Numbers k such that k multiplied by any of its digits contains k as a substring.

Conjecture: consists of 0,5,6, and the numbers 11...1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 19 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014405 | OEIS-A328549

1, together with the numbers that are simultaneously superior highly composite (A002201), colossally abundant (A004490), deeply composite (A095848), and miserable average divisor numbers (A263572).

Conjecture: the solutions to this problem, if any, form an initial sequence of the present sequence. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014406 | OEIS-A328554

Coefficients in Göttsche's universal power series B_1(q) arising from enumeration of d-nodal curves in a linear system of dimension d on an algebraic surface.

The power series appears to be well defined, only the interpretation is conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014407 | OEIS-A328555

Coefficients in Göttsche's universal power series B_2(q) arising from enumeration of d-nodal curves in a linear system of dimension d on an algebraic surface.

The power series appears to be well defined, only the interpretation is conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014408 | OEIS-A328556

Expansion of Product_{p prime, k>=1} (1 - x^(p^k)).

Conjecture: the last zero (38th) occurs at n = 340.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Nov 01 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014409 | OEIS-A328610

Irregular triangular array read by rows: the rows show the coefficients of the first of two factors of even-degree polynomials described in Comments.

Let p(n) denote the polynomial (1/n!)*(numerator of n-th derivative of (1-x)/(1-x-x^2)). It is conjectured in A326925 that if n = 2k, then p(n) = f(k)*g(k), where f(k) and g(k) are polynomials of degree k. Row k of the present array shows the coefficients of f(k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014410 | OEIS-A328656

Numbers m such that d(m) < d(k) for all k < m, where d is the normalized delta defined as d(m) = (z(m+1) - z(m))*(log(z(m)/(2*Pi))/(2*Pi)) where z(k) is the imaginary part of the k-th Riemann zeta zero.

Conjectural next terms: 1217992279429, 4088664936219.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jan 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014411 | OEIS-A328685

Row sums of A309038.

Conjectures from _Colin Barker_, Oct 25 2019: (Start) a(n) ~ 5*n^4/8. - Conjectured by _Stefano Spezia_, Sep 08 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Oct 25 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014412 | OEIS-A328718

Square array T(n,k), n>=0, k>=0, read by antidiagonals, where T(n,k) is the constant term in the expansion of (1 + x_1 + x_2 + ... + x_n + 1/x_1 + 1/x_2 + ... + 1/x_n)^k.

Conjecture: Row r is asymptotic to (2*r+1)^(n + r/2) / (2^r * (Pi*n)^(r/2)). - _Vaclav Kotesovec_, Oct 27 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Oct 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014413 | OEIS-A328745

Dirichlet g.f.: Product_{p prime} 1 / (1 - p^(-s))^p.

Conjecture: Sum_{k=1..n} a(k) ~ c * n^2, where c = 0.40373... - _Vaclav Kotesovec_, Mar 28 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Oct 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014414 | OEIS-A328786

a(n) = (Sum_{k=0..n-1}(-1)^k*(39480*k+7321)*29700^(n-1-k)*T_k(14,1)*T_k(11,-11)^2)/n, where T_k(b,c) denotes the coefficient of x^k in the expansion of (x^2+b*x+c)^k.

Conjecture: (i) a(n) is a positive integer for each n > 0, and a(n) is odd if and only if n is a power of two. Moreover, we have the identity Sum_{k>=0}(39480*k+7321)(-29700)^(-k)*T_k(14,1)*T_k(11,-11)^2 = 6795*sqrt(5)/Pi. In 2011 the author formulated many conjectural series for 1/Pi involving T_k(b,c). The conjectural series for 1/Pi in A329073, A329105, A329107 and the present one are of a new type which might be called type VIII to distinguish from the 7 known types (types I-VII).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014415 | OEIS-A328825

Negative discriminants with form class group isomorphic to C_3 (negated).

Conjecture: this sequence is finite and this is the full list.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Dec 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014416 | OEIS-A328865

The first repeating term in the trajectory of n under iterations of A329623, or -1 if no such terms exists.

All -1 are so far conjectural, see A329624 and A329917 for more information.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014417 | OEIS-A328899

Numbers k such that the k-th repunit (10^k-1)/9 sets a new record for the number of distinct prime factors.

a(24) <= 840. Conjecture: a(24) = 840, a(25) = 1260, a(26) = 1680, a(27) = 1980, a(28) = 2520, a(29) = 3360, a(30) = 3780, a(31) = 3960, a(32) = 4620, a(33) = 5040, a(34) = 6300, a(35) = 7560, a(36) = 9240, a(37) = 10080. - _Chai Wah Wu_, Nov 01 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Resta_, Oct 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014418 | OEIS-A328919

Smallest index m such that from the m-th term on, the sequence {sigma_k(n) mod n: k >= 0} enters into a cycle.

Conjecture: a(n) = 0 only for n = 1, 12. If {sigma_k(n) mod n: k >= 0} is purely periodic, then sigma_0(n) == sigma_psi(n)(n) (mod n). Let p be any prime divisor of n, write n = p^e*s, then sigma_0(n) = d(n) = (e+1)*d(s), sigma_psi(n)(n) == Sum_{d|n, gcd(d,p)=1} 1 = d(s) (mod p^e), d = A000005. So sigma_0(n) == sigma_psi(n)(n) (mod n) means that p^e divides e*d(s) for every p dividing n, which seems highly impossible for n > 12. Of course, this shows that a(n) cannot be 0 if n is squarefree, so a(n) = 1 for squarefree n. Every number occurs in this sequence, because a(p^e) = e for primes p. Conjecture: a(2^e) is always the earliest occurrence of e.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Oct 31 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014419 | OEIS-A328934

Nonsquarefree N such that A328919(N) = 1; nonsquarefree N such that {sigma_t(N) mod N: t >= 1} is purely periodic, while {sigma_t(N) mod N: t >= 0} is not.

If N is squarefree then A328919(N) = 1, but the converse is not true. While it is conjectured that 12 is the only N such that 0 = A328919(N) < A051903(N), there are infinitely many N such that 1 = A328919(N) < A051903(N). This shows that A051903(N) - A328919(N) is unbounded, and can take every natural number as its value for infinitely many

times. It is conjectured that the smallest N such that A328919(N) = 1 and A051903(N) = k is N = 2^(k-1)*A002110(k-1), k >= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Oct 31 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014420 | OEIS-A328935

Imprimitive Carmichael numbers: Carmichael numbers m such that if m = p_1 * p_2 * ... *p_k is the prime factorization of m then g(m) = gcd(p_1 - 1, ..., p_k - 1) > sqrt(lambda(m)), where lambda is the Carmichael lambda function (A002322).

They conjectured that most Carmichael numbers are primitive and most 3-Carmichael numbers (A087788) are imprimitive.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Oct 31 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014421 | OEIS-A328936

The number of imprimitive Carmichael numbers (A328935) below 10^n.

Granville and Pomerance conjectured that most Carmichael numbers are primitive, i.e. lim_{n->oo} a(n)/A055553(n) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Oct 31 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014422 | OEIS-A328937

The number of imprimitive 3-Carmichael numbers (A087788 and A328935) below 10^n.

Granville and Pomerance conjectured that most Carmichael numbers are imprimitive, i.e. lim_{n->oo} a(n)/A132195(n) = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Oct 31 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014423 | OEIS-A328959

a(n) = sigma_0(n) - 2 - (omega(n) - 1) * nu(n), where sigma_0 = A000005, nu = A001221, omega = A001222.

Conjecture: All terms are nonnegative except for a(1) = -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Nov 02 2019. The idea for this sequence came from _Mats Granvik_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014424 | OEIS-A328960

Number of integer partitions of n whose number of nontrivial submultisets is greater than their number of distinct parts times their number of parts minus 1.

These partitions are conjectured to be precisely those that have a pair of multiset partitions such that no part of one is a submultiset of any part of the other (see A320632). For example, such a pair of partitions of {1,1,2,2} is ({{1,1},{2,2}}, {{1,2},{1,2}}).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Nov 02 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014425 | OEIS-A328982

Sorted list of the numbers of the form 5m+2 (m>=0) together with numbers of the form 5m-2+eps (m>=1), where eps = 1 if the binary expansion of m ends in an odd number of 0's and is otherwise 0.

Conjecture: these are the numbers k such that -k is a term of A328196.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014426 | OEIS-A328983

Complement of A328982.

Conjecture: These are the numbers k such that -k is not a term in A328196.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014427 | OEIS-A328987

The sequence C(n) defined in the comments (A and B smallest missing numbers, offset 0).

Conjectures from _Colin Barker_, Nov 08 2019: (Start) Conjecture: a(n) ~ 35*n/6. - _Stefano Spezia_, Nov 02 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014428 | OEIS-A329012

a(n) = p(1,n), where p(x,n) is the strong divisibility sequence of polynomials based on sqrt(5) as in A327322.

a(n) is a strong divisibility sequence; i.e., gcd(a(h),a(k)) = a(gcd(h,k)). Conjecture: there is no upper bound for the number of consecutive equal digits among numbers in this sequence, as suggested, for example, by 34 straight 1's in a(96) and 38 straight 6's in a(97).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014429 | OEIS-A329051

Positions of records in A329040.

Conjecturally also the position of the first occurrence of each n in A329040.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014430 | OEIS-A329073

a(n) = (1/n)*Sum_{k=0..n-1} (40k+13)*(-1)^k*50^(n-1-k)*T_k(4,1)*T_k(1,-1)^2, where T_k(b,c) denotes the coefficient of x^k in the expansion of (x^2+b*x+c)^k.

Conjecture 1: (i) a(n) is a positive integer for each n > 0; also, a(n) is odd if and only if n is a power of two. Moreover, we have the identity Sum_{k>=0} ((40k+13)/(-50)^k)*T_k(4,1)*T_k(1,-1)^2 = 55*sqrt(15)/(9*Pi). Conjecture 2: (i) For any n > 0, the number b(n):=(1/n)*Sum_{k=0..n-1} (40k+27)*(-6)^(n-1-k)*T_k(4,1)*T_k(1,-1)^2 is an integer. Moreover, b(n) is odd if and only if n is a power of two.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014431 | OEIS-A329105

a(n) = (Sum_{k=0..n-1} (1435*k+113)*3240^(n-1-k)*T_k(7,1)*T_k(10,10)^2)/(n*10^(n-1)), where T_k(b,c) denotes the coefficient of x^k in the expansion of (x^2+b*x+c)^k.

Conjecture 1: (i) a(n) is an integer for each n > 0. Moreover, we have Sum_{k>=0}((1435*k+113)/3240^k)*T_k(7,1)*T_k(10,10)^2 = 1452*sqrt(5)/Pi. Conjecture 2: (i) For each n > 0, the number (Sum_{k=0..n-1}(1435*k+1322)*50^(n-1-k)*T_k(7,1)*T_k(10,10)^2)*3/(2n*10^(n-1)) is an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 04 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014432 | OEIS-A329107

a(n) = (Sum_{k=0..n-1}(840*k+197)(-1)^k*2430^(n-1-k)*T_k(8,1)*T_k(5,-5)^2)/(n*5^(n-1)), where T_k(b,c) denotes the coefficient of x^k in the expansion of (x^2+b*x+c)^k.

Conjecture: (i) a(n) is a positive integer for each n > 0. Moreover, we have Sum_{k>=0} ((840*k+197)/(-2430)^k)*T_k(8,1)*T_k(5,-5)^2 = 189*sqrt(15)/(2*Pi).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 04 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014433 | OEIS-A329126

a(n) is the least positive number which yields a multiple of n when its decimal digits (which are necessarily 0's and 1's) are read in any base.

The number of 1's in a(n) is conjectured to be n; the number of 0's separating each one digit is usually A268336(n)-1 for small n. The number of trailing 0's is A051903(n). The first 21 terms have been found and verified to be minimal via an advanced search; the 13th term (see b-file) contains 146 digits, and in general for every prime n the corresponding a(n) is conjectured to contain 2+(n-1)^2 0's and 1's.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alon Ran_, Nov 05 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014434 | OEIS-A329168

Numbers m that divide 4^(m + 1) + 1.

Conjecture: For k > 1, k^(m + 1) == -1 (mod m) has an infinite number of positive solutions. Conjecture: For k > 1, if f(1) = p(1) equals one of the prime factors of k^2 + 1, p(i+1) equals one of the prime factors of k^(f(i)+1) + 1 greater than p(i), f(i+1) = f(i)*p(i+1), then k^(f(i) + 1) == -1 (mod f(i)) for all integers i. (Especially in this sequence, k = 4, so {f(i)} can be 17, 4097, 4298117633, ...) - _Jinyuan Wang_, Nov 16 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Nov 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014435 | OEIS-A329178

Sum of the products of pairs of Padovan numbers which are two apart, starting from A000931(5).

Conjectures from _Colin Barker_, Nov 09 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Nacin_, Nov 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014436 | OEIS-A329181

a(n) = n if n is 1 or prime; otherwise (1) let m = (concatenation of the two divisors in the middle of rows of A027750(n)), (2) if m is prime then a(n) = m, otherwise return to (1) with n=m.

Apparently it is only a conjecture that the process in the definition will alwats terminate. - _N. J. A. Sloane_, Feb 23 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Cobac_, Nov 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014437 | OEIS-A329203

Numbers k such that A329308(k) = 0.

No more terms up to 2*10^7. I conjecture that these are all the terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Nov 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014438 | OEIS-A329222

Numbers m that divide 5^(m + 1) + 1.

Conjecture: For k > 1, k^(m + 1) == -1 (mod m) has an infinite number of positive solutions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Nov 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014439 | OEIS-A329226

Numbers m that divide 6^(m + 1) + 1.

Conjecture: For k > 1, k^(m + 1) == -1 (mod m) has an infinite number of positive solutions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Nov 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014440 | OEIS-A329227

Products of consecutive terms of the Padovan sequence A000931.

G.f.: x^5 / ((1 - 2*x + x^2 - x^3)*(1 + x - x^3)) (conjectured). - _Colin Barker_, Nov 08 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Nacin_, Nov 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014441 | OEIS-A329230

The number of steps it takes to reach all n positions around a circle during the grasshopper procedure.

In the Mathematics Stack Exchange link, the author conjectures that a(n) = n if and only if n = 3, n = 7, or n = 2^k for some nonnegative k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Nov 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014442 | OEIS-A329231

The maximum number of times one reaches a single position during the grasshopper procedure.

Conjecture: a(n) = 1 if and only if n = 3, n = 7, or n = 2^k for some k. Conjecture: The largest values of n for which a(n) = 2, 3, 4, 5 respectively are n = 18, 68, 381, 1972. If the second conjecture is true, then 2, 3, 4, and 5 appear 2, 19, 87, and 313 times respectively. Conjecture: Every integer greater than 1 appears in this sequence a finite number of times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Nov 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014443 | OEIS-A329232

The number of counterclockwise steps during the grasshopper procedure.

Conjecture: a(n)=0 if and only if n = 2^k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Nov 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014444 | OEIS-A329233

The time of the first counterclockwise step during the grasshopper procedure, or 0 if no counterclockwise steps occur.

Conjecture: a(3^n) = A087503(n-1) + 1 for n > 0. (Checked up to 3^12.) - _Peter Kagey_, Nov 28 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Nov 10 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014445 | OEIS-A329254

Smallest palindrome k, not less than n, such that k is also a palindrome in base n.

a(10^k) = 10^k+1 for k > 0. Conjecture: a(n) in base n has 3 or fewer digits. - _Chai Wah Wu_, Nov 17 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Price_, Nov 09 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014446 | OEIS-A329263

Irregular triangle read by rows in which row n is the result of iterating the operation f(n) = n/8 if n == 0 (mod 8), otherwise f(n) = 8*(floor(n/8) + n + 1), terminating at the first occurrence of 1.

Conjecture: For any initial value of n >= 1 there is a number k such that f^{k}(n) = 1; in other words, every row of the triangle is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Davis Smith_, Nov 09 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014447 | OEIS-A329333

There is exactly one odd prime among the pairwise sums of any three consecutive terms: Lexicographically earliest sequence of distinct nonnegative integers with this property.

This is conjectured and designed to be a permutation of the nonnegative integers, therefore the offset is taken to be zero. Restricted to positive indices, this is a sequence of positive integers having the same property, then conjectured to be a permutation of the positive integers. (The word "odd" can be omitted in this case.) If the word "odd" is dropped from the original definition, the sequence starts (0, 1, 3, 6, 2, 7), and then continues from a(6) = 4 onward as the present sequence. This is again conjectured to be a permutation of the nonnegative integers, and a permutation of the positive integers when restricted to the domain [1..oo). The latter however no longer has the property of lexicographic minimality.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_, _Jean-Marc Falcoz_ and _M. F. Hasler_, Nov 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014448 | OEIS-A329343

Difference between the indices of the smallest and the largest primorial in the greedy sum of primorials adding to A108951(n).

Positions of the records (and conjecturally, the positions of the first occurrences of each n) begin as 1, 8, 27, 162, 289, 529, 841, 1369, 1681, 2209, 2809, 3481, 4489, 5041, 5329, 6889, ..., that after 162 all seem to be squares of certain primes. See also A329051.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014449 | OEIS-A329354

a(n) = Sum_{d|n} d*omega(d).

Conjecture: Sum_{k=1..n} a(k) ~ Pi^2*n^2*(log(log(n)) + A077761)/12. - _Vaclav Kotesovec_, Mar 03 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014450 | OEIS-A329387

Solution sequence (a(n)) of the complementary equation a(n) = 4 b(n+1) - b(n), with b(0) = 1.

The conditions that (a(n)) and (b(n)) be increasing and complementary force the equation a(n) = 4 b(n+1) - b(n), with initial value b(0) = 1, to have a unique solution; that is, a pair of complementary sequences (a(n)) = (7,10,13,16,19,26,28,...) and (b(n)) = (1,2,3,4,6,8,9,11,12,14,...). For n = 1..30000, a(n) - 4 n ranges between -12 and 15; conjecture: {a(n) - 4 n} is unbounded below and above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014451 | OEIS-A329388

Solution sequence (a(n)) of the complementary equation a(n) = 5 b(n+1) - b(n), with b(0) = 1.

The conditions that (a(n)) and (b(n)) be increasing and complementary force the equation a(n) = 5 b(n+1) - b(n), with initial value b(0) = 1, to have a unique solution; that is, a pair of complementary sequences (a(n)) = (9,13,17,21,25,29,...) and (b(n)) = (1,2,3,4,5,6,7,8,10, ...). Conjecture: {a(n) - 5 n} is unbounded below and above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014452 | OEIS-A329389

Solution sequence (a(n)) of the complementary equation a(n) = 6 b(n+1) - b(n), with b(0) = 1.

The conditions that (a(n)) and (b(n)) be increasing and complementary force the equation a(n) = 6 b(n+1) - b(n), with initial value b(0) = 1, to have a unique solution; that is, a pair of complementary sequences (a(n)) = (11,16,21,26,31,...) and (b(n)) = (1,2,3,4,5,6,7,8,9,10,12,...). Conjecture: {a(n) - 6 n} is unbounded below and above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014453 | OEIS-A329390

Solution sequence (a(n)) of the complementary equation a(n) = 6 b(n+1) - 2 b(n), with b(0) = 1.

The conditions that (a(n)) and (b(n)) be increasing and complementary force the equation a(n) = 6 b(n+1) - 2 b(n), with initial value b(0) = 1, to have a unique solution; that is, a pair of complementary sequences (a(n)) = (10,14,18,22,26,30,34,...) and (b(n)) = (1,2,3,4,5,6,7,8,9,11,12,...). Conjecture: {a(n) - 5 n} is unbounded below and above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 23 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014454 | OEIS-A329395

Numbers whose binary expansion without the most significant (first) digit has Lyndon and co-Lyndon factorizations of equal lengths.

Conjecture: also numbers k such that the k-th composition in standard order (A066099) is a palindrome, cf. A025065, A242414, A317085, A317086, A317087, A335373. - _Gus Wiseman_, Jun 06 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Nov 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014455 | OEIS-A329398

Number of compositions of n with uniform Lyndon factorization and uniform co-Lyndon factorization.

Conjecture: Also the number of compositions of n that are either weakly increasing or weakly decreasing. Hence a(n) = 2 * A000041(n) - A000005(n). - _Gus Wiseman_, Mar 05 2020 Theorem: a composition has a uniform Lyndon factorization and a uniform co-Lyndon factorization if and only if it is weakly increasing or weakly decreasing. This proves Wiseman's conjecture above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Nov 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014456 | OEIS-A329405

Among the pairwise sums of any three consecutive terms there is no prime: lexicographically earliest such sequence of distinct positive integers.

Conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Nov 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014457 | OEIS-A329412

Lexicographically earliest sequence of distinct positive numbers such that among the pairwise sums of any four consecutive terms there are exactly two prime sums.

Conjectured to be a permutation of the positive integers: a(10^5) = 10^5, a(10^6) = 999984 and all numbers below 99992 resp. 999963 have appeared by then. See A329452 for a more detailed discussion. - _M. F. Hasler_, Nov 15 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Nov 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014458 | OEIS-A329413

Lexicographically earliest sequence of distinct positive numbers such that among the pairwise sums of any five consecutive terms there are exactly two prime sums.

Conjectured to be a permutation of the positive integers: a(10^6) = 10^6 + 9 and all numbers below 10^6 - 7 are used at that point. - _M. F. Hasler_, Nov 15 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Nov 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014459 | OEIS-A329414

Lexicographically earliest sequence of distinct positive numbers such that among the pairwise sums of any six consecutive terms there are exactly two prime sums.

Condition a(1) = 1 follows from minimality. Conjectured to be a permutation of the positive integers: a(10^6) = 999994 and all numbers up to there have appeared at that point. - _M. F. Hasler_, Nov 15 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Nov 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014460 | OEIS-A329416

Among the pairwise sums of any ten consecutive terms there are exactly two prime sums: lexicographically earliest such sequence of distinct positive numbers.

Conjectured to be a permutation of the positive integers: a(10^6) = 10^6 + 2 and all numbers up to 10^6 - 7 have appeared at that point. - _M. F. Hasler_, Nov 15 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Nov 14 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014461 | OEIS-A329421

a(n) = gcd(A330050(n), A330051(n)).

Conjectures from _Colin Barker_, Dec 02 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Somos_, Nov 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014462 | OEIS-A329423

Each term is visited once according to the rule: go left (resp. right) a(n) places if a(n) is prime (resp. not prime), starting at a(1) = 1. Choose a(n) along this journey as the largest possible prime, or else the smallest possible composite, not occurring earlier and compatible with these rules.

The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Nov 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014463 | OEIS-A329425

For all n >= 0, six among (a(n+i) + a(n+j), 0 <= i < j < 5) are prime: lexicographically first such sequence of distinct nonnegative integers.

Conjectured to be a permutation, i.e., all n >= 0 appear. The restriction to [1, oo) is then the lexicographically first such permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, following an idea from _Eric Angelini_, Nov 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014464 | OEIS-A329440

Triangle read by rows: n-th row gives positions of ones in A329126(n) in decreasing order.

Conjecture: the last value in each row is A051903(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Nov 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014465 | OEIS-A329449

For any n >= 0, exactly four sums a(n+i) + a(n+j) are prime, for 0 <= i < j <= 3: lexicographically earliest such sequence of distinct nonnegative integers.

This map is defined with offset 0 as to have a permutation of the nonnegative integers in case each of these eventually appears, which is so far only conjectured, see below. The restriction to positive indices would then be a permutation of the positive integers, and as it happens, also the smallest one with the given property. (This is in contrast to most other cases where that one is not the restriction of the other one: see crossrefs). Concerning the existence of the sequence with infinite length: If the sequence is to be computed in a greedy manner, this means that for given P(n) := {a(n-1), a(n-2), a(n-3)} and thus 0 <= N(n) := #{ primes x + y with x, y in P(n), x < y} <= 4, we have to find a(n) such that we have exactly 4 - N(n) primes in a(n) + N(n). It is easy to prove that this is always possible when 4 - N(n) = 0 or 1. Otherwise, similar to A329452, ..., A329456, we see that P(n) is an "admissible constellation" in the sense that a(n-4) + P(n) already gave the number of primes required now. So a weaker variant of the k-tuple conjecture would ensure we can find this a(n). But the sequence need not be computable in greedy manner! That is, if ever for given P(n) no a(n) would exist such that a(n) + P(n) contains 4 - N(n) primes, this simply means that the considered value of a(n-1) (and possibly a(n-2)) was incorrect, and the next larger choice has to be made. Given this freedom, there is no doubt that this sequence is well defined up to infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, based on an idea from _Eric Angelini_, Nov 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014466 | OEIS-A329450

Lexicographically earliest sequence of distinct nonnegative integers such that neither a(n) + a(n+1) nor a(n) + a(n+2) is prime for any n.

Conjectured to be a permutation of the nonnegative integers. The restriction to positive indices is then a permutation of the positive integers with the same property, but not the lexicographically earliest given in A329405.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, based on an idea of _Eric Angelini_, Nov 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014467 | OEIS-A329452

There are exactly two primes in {a(n+i) + a(n+j), 0 <= i < j <= 3} for any n: lexicographically earliest such sequence of distinct nonnegative integers.

Conjectured to be a permutation of the nonnegative numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Nov 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014468 | OEIS-A329453

There are exactly two primes in {a(n+i) + a(n+j), 0 <= i < j <= 4} for any n: lexicographically earliest such sequence of distinct nonnegative integers.

Conjectured to be a permutation of the nonnegative numbers. (Therefore the offset is chosen to have a(0) = 0. The restriction to positive indices is then a permutation of the positive integers, but not the smallest one which is A329413, conjectured.) Concerning the existence of the sequence, if the sequence is to be computed in a greedy manner, this means the following: for given n, we assume given P(n) := {a(n-1), a(n-2), a(n-3), a(n-4)} and thus S(n) := #{ primes x + y with x, y in P(n), x < y} which may equal 0, 1 or 2. We have to find a(n) such that we have exactly 2 - S(n) primes in a(n) + P(n). It is easy to prove that this is possible when 2 - S(n) is 0 or 1. When S(n) = 0, we must find two primes which are at a distance of |x - y| for some x, y in P(n). This is much weaker than to require the existence of twin primes or cousin primes etc., generally believed to hold and in part proved under hypotheses weaker than RH: First, we have the choice of several distances. Second, we don't require that there be no other primes in between. But more than that, it is not at all needed that the sequence be computable in a greedy manner! I.e., in the extremely improbable event that for some n with S(n) = 0 there might be no a(n) such that a(n) + P(n) contains 2 primes, we have infinitely many other choices for a(n-1) or even a(n-2), etc.! Given this additional freedom, the existence of a(n) for any n is beyond any doubt (and maybe not even difficult to prove, by contradiction). Concerning the conjecture that all numbers will eventually occur: if a number m never appears, this means that m + P(n) never has the required number of 2 - S(n) primes. That is, for all n such that S(n) = 2, the set m + P(n) has at least one prime, and whenever S(n) = 1, the

set m + P(n) has never exactly 1 prime. Given that each of the sets P(n) contains 4 numbers which were chosen essentially independently of m, it appears extremely improbable that all these infinitely many constraints could hold simultaneously for any m. Computational results so far also give only strong evidence in favor of this conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, based on an idea from _Eric Angelini_, Nov 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014469 | OEIS-A329454

There are exactly three primes among a(n+i) + a(n+j), 0 <= i < j <= 3, for any n >= 0: lexicographically earliest such sequence of distinct nonnegative integers.

Concerning the existence of the sequence: If the sequence is to be computed in a greedy manner, this means that for given n, we assume given P(n) := {a(n-1), a(n-2), a(n-3)} and thus S(n) := #{ primes x + y with x, y in P(n), x < y} which may equal 0, 1, 2 or 3. We have to find a(n) such that we have exactly 3 - S(n) primes in a(n) + P(n). It is easy to prove that this is always possible when 3 - S(n) is 0 or 1, and similar to the case of A329452 and A329453 when S(n) = 1. When S(n) = 0, however, we must find three primes in the same configuration as the elements of P(n). It is almost surprising that this can always be found up to n = 10^6. However, the sequence does not need to be computable in a greedy manner. That is, if for given P(n) no a(n) would exist such that a(n) + P(n) contains 3 - S(n) primes, this simply means that the considered value of a(n-1) was incorrect, and the next larger choice has to be made. Given this freedom, well-definedness of this sequence up to infinity is far more probable than, for example, the k-tuple conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, based on an idea from _Eric Angelini_, Nov 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014470 | OEIS-A329455

There are exactly three primes in {a(n+i) + a(n+j), 0 <= i < j <= 4} for any n >= 0: lexicographically earliest such sequence of distinct nonnegative integers.

Concerning the existence of the sequence: If the sequence is to be computed in a greedy manner, this means that for given n, we assume given P(n) := {a(n-1), a(n-2), a(n-3), a(n-4)} and thus S(n) := #{ primes x + y with x, y in P(n), x < y} which may equal 0, 1, 2 or 3. We have to find a(n) such that we have exactly 3 - S(n) primes in a(n) + P(n). It is easy to prove that this is always possible when 3 - S(n) = 0 or 1, and for S(n) = 0 or 1, this is similar to the case of A329452 or A329453. However, the sequence does not need to be computable in a greedy manner. That is, if for given P(n) no a(n) would exist such that a(n) + P(n) contains 3 - S(n) primes, this simply means that the considered value of a(n-1) was incorrect, and the next larger choice has to be made. Given this freedom, well-definedness of this sequence up to infinity is far more probable than, for example, the k-tuple conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, based on an idea from _Eric Angelini_, Nov 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014471 | OEIS-A329456

For any n >= 0, exactly four sums a(n+i) + a(n+j) are prime, for 0 <= i < j <= 4: lexicographically earliest such sequence of distinct nonnegative integers.

Concerning the existence of the sequence with infinite length: If the sequence is to be computed in a greedy manner, this means that for given P(n) := {a(n-1), a(n-2), a(n-3), a(n-4)} and thus N(n) := #{ primes x + y with x, y in P(n), x < y} in {0, ..., 4}, we have to find a(n) such that we have exactly 4 - N(n) primes in a(n) + N(n). It is easy to prove that this is always possible when 4 - N(n) = 0 or 1. Otherwise, similar to A329452, ..., A329455, we see that P(n) is an "admissible constellation" in the sense that a(n-5) + P(n) already gave the number of primes required now. So a (weaker) variant of the k-tuple conjecture ensures we can find this a(n). But the sequence need not be computable in greedy manner! That is, if ever for given P(n) no convenient a(n) would exist, this just means that the considered value of a(n-1) (and possibly a(n-2)) was incorrect, and the next larger choice has to be made. Given this freedom, there is no doubt that this sequence is well defined up to infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, based on an idea from _Eric Angelini_, Nov 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014472 | OEIS-A329475

a(n) = Sum_{k=0..n} C(n,k)^2*T(k)*T(n-k), where T(k) = A002426(k) is the coefficient of x^k in the expansion of (x^2+x+1)^k.

The author introduced this sequence in arXiv:1911.05456 and made the following conjecture. Conjecture: Let p be an odd prime and let S = Sum_{k=0..p-1}a(k)/(-4)^k. If p == 1 (mod 12) and p = x^2 + 9*y^2 with x and y integers, then S == 4*x^2-2*p (mod p^2). If p == 5 (mod 12) and p = x^2 + y^2 with x == y (mod 3), then S == 4*x*y (mod p^2). If p == 3 (mod 4), then S == 0 (mod p^2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014473 | OEIS-A329478

a(n) = (Sum_{k=0..n-1}(-1)^k*(15*k+8)*beta(k)*t(k))/(2*n), where beta(k) = A005258(k), and t(k) is the coefficient of x^k in the expansion of (x^2+4*x-1)^k.

Conjecture 1: (i) a(n) is an integer for each n > 0. Moreover, a(n) is odd if and only if n is a positive power of two. Conjecture 2: Let p > 5 be a prime and let S(p) = Sum_{k=0..p-1}(-1)^k*beta(k)*t(k). If p == 1,4 (mod 5) and p = x^2 + 15*y^2 (with x and y integers), then S(p) == 4*x^2-2p (mod p^2). If p == 2,8 (mod 15) and p = 3*x^2 + 5*y^2, then S(p) == 12*x^2-2p (mod p^2). If Leg(-15/p) = -1, then S(p) == 0 (mod p^2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014474 | OEIS-A329479

Number of degree n polynomials f with all nonzero coefficients equal to 1 such that f(k) is divisible by 3 for all integers k.

Conjectured recurrence: a(n) = 2a(n-1) + 2a(n-2) - 5a(n-3) - 2a(n-4) + 10a(n-5) - 4a(n-6) - 4a(n-7) + 8a(n-8).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Nov 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014475 | OEIS-A329512

Array read by upward antidiagonals: row n = coordination sequence for cylinder formed by rolling up a strip of width 2*n hexagons cut from the hexagonal grid by cuts perpendicular to grid lines.

The g.f.s for the rows could be found using the "trunks and branches" method (see Goodman-Strauss and Sloane), as was done in A329508. This step has not yet been carried out, so the following g.f. is at present only conjectural. The g.f. G(n) for row n (n>=1) is (strongly) conjectured to be

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014476 | OEIS-A329515

Array read by upward antidiagonals: row n = coordination sequence for cylinder formed by rolling up a strip of width 2*n+1 hexagons cut from the hexagonal grid by cuts perpendicular to grid lines.

The g.f.s for the rows could be found using the "trunks and branches" method (see Goodman-Strauss and Sloane), as was done in A329508. This step has not yet been carried out, so the following g.f. is at present only conjectural. The g.f. G(n) for row n (n>=0) is (strongly) conjectured to be

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 25 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014477 | OEIS-A329545

After a(1) = 1, add the even terms and subtract the odd ones, the result must always be a palindrome. This is the lexicographically earliest sequence of distinct positive integers with this property.

a(63411) = 964. _Rémy Sigrist_'s comment in A329544 shows that terms in A104444 are not in the sequence. Conjecture: Sequence is a permutation of positive integers not in A104444. - _Chai Wah Wu_, Dec 11 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Nov 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014478 | OEIS-A329563

For all n >= 1, exactly five sums are prime among a(n+i) + a(n+j), 0 <= i < j < 5; lexicographically earliest such sequence of distinct positive numbers.

Conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014479 | OEIS-A329564

For all n >= 0, exactly five sums are prime among a(n+i) + a(n+j), 0 <= i < j < 5; lexicographically earliest such sequence of distinct nonnegative numbers.

Conjectured to be a permutation of the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014480 | OEIS-A329565

For all n >= 0, exactly five sums are prime among a(n+i) + a(n+j), 0 <= i < j < 6; lexicographically earliest such sequence of distinct nonnegative numbers.

Conjectured to be a permutation of the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014481 | OEIS-A329567

For all n >= 0, exactly seven sums are prime among a(n+i) + a(n+j), 0 <= i < j < 6: lexicographically earliest such sequence of distinct nonnegative numbers.

Conjectured to be a permutation of the nonnegative integers. The restriction to [1,oo) is then a permutation of the positive integers with the same property, but not the smallest one which is A329576 = (1, 2, 3, 4, 5, 8, 11, ...).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014482 | OEIS-A329568

For all n >= 1, exactly 9 sums are prime among a(n+i) + a(n+j), 0 <= i < j < 6: lexicographically earliest such sequence of distinct positive numbers.

Conjectured to be a permutation of the positive integers. See A329569 = (0, 1, 2, 5, 6, 11, 12, 17, ...) for the quite different variant for nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 10 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014483 | OEIS-A329569

For all n >= 0, exactly 9 sums are prime among a(n+i) + a(n+j), 0 <= i < j < 6: lexicographically earliest such sequence of distinct nonnegative numbers.

Conjectured to be a permutation of the nonnegative integers. The restriction to [1,oo) is then a permutation of the positive integers with similar properties, but different from the lexico-smallest one, A329568 = (1, 2, 3, 9, 4, 10, 27, ...).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 10 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014484 | OEIS-A329572

For all n >= 0, exactly 12 sums are prime among a(n+i) + a(n+j), 0 <= i < j < 7; lexicographically earliest such sequence of distinct nonnegative numbers.

Conjectured to be a permutation of the nonnegative integers. See A329573 for the "positive" variant: same definition but with offset 1 and positive terms, leading to a quite different sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014485 | OEIS-A329573

For all n >= 1, exactly 12 sums are prime among a(n+i) + a(n+j), 0 <= i < j < 7; lexicographically earliest such sequence of distinct positive numbers.

Conjectured to be a permutation of the positive integers. See A329572 for the nonnegative variant (same definition but with n >= 0 and terms >= 0), leading to a quite different sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014486 | OEIS-A329574

For every n >= 0, exactly 10 sums are prime among a(n+i) + a(n+j), 0 <= i < j < 7; lexicographically earliest such sequence of distinct nonnegative numbers.

Conjectured to be a permutation of the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 10 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014487 | OEIS-A329576

For all n >= 1, exactly seven sums are prime among a(n+i) + a(n+j), 0 <= i < j < 6; lexicographically earliest such sequence of distinct positive numbers.

Conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014488 | OEIS-A329616

Bitwise-OR of exponents of prime factors of A108951(n), where A108951 is fully multiplicative with a(prime(i)) = prime(i)# = Product_{i=1..i} A000040(i).

Positions of records are: 1, 2, 4, 6, 16, 24, 36, 54, 256, 384, 576, 864, 1296, 1944, 2916, 4374, 65536, ..., conjectured also to be the positions of the first occurrence of each n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 17 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014489 | OEIS-A329624

Number of iterations of A329623 for starting value n before a repeated value appears, or -1 if this never happens.

It seems the a(n) = -1 are conjectural, i.e., we have no proof that the terms for which the trajectory seems to "explode" do not eventually end up in a cycle. For example, the 8th iterate of 1373 is 5218725017016262626273. If the 2nd digit is changed from 2 to 0, then the further iterates appear to explode up to a length of 157 bits, but finally end up in a 2-cycle of 41-digit numbers (26...26273, 62...62637).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 19 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014490 | OEIS-A329649

Let D = A042948(n) be the n-th positive integer congruent to 0 or 1 mod 4, then a(n) = b(D) := Sum_{i=1..D} Kronecker(D,i)*i^2, where Kronecker(D,i) is the Kronecker symbol.

Conjecture (a): There are no 0's in this sequence. Conjecture (b): if D is not of the form k^2 or 5*k^2 or 2^t (t odd, t > 1), then b(D) is divisible by 4*D; if D = 5*k^2, then b(D) is divisible by 4*k^2 = 4*(D/5). Conjecture (c): let D be a fundamental discriminant > 1 (A003658), then Sum_{k>=1} Kronecker(D,k)/k^2 = Pi^2*b(D)/D^(5/2) = (4*Pi^2)/D^(3/2) * (b(D)/(4*D)), which is an integer times (4*Pi^2)/D^(3/2) if D > 8 by the conjecture (b). Here Sum_{k>=1} Kronecker(D,k)/k^2 is the value of the Dirichlet L-series of a non-principal character modulo D at s=2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Nov 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014491 | OEIS-A329654

a(n) = numerator(b(n)), where b(0) = b(1) = 1 and b(n) = n*b(n-1)/b(n-2) for n >= 1.

This sequence is derived from a particular case of a general recurrence relation expressed by B(0) = x, B(1) = y and B(n) = n*B(n-1)/B(n-2), for n > 1 and {x,y} any pair of nonzero real numbers. Scatter plots of sequences of this kind exhibit a particular pattern that suggests the following conjecture: This means that B(n)/n approaches a cycle of six different constant values which depend on the particular chosen seed {x,y}. In this particular case the seed is {1,1} and the corresponding conjectured constant limits {C_1, C_2, C_3, C_4, C_5, C_6} are approximately {0.431, 0.615, 1.426, 2.319, 1.626, 0.701}. The corresponding constant limits for a generic seed {x,y} are respectively {C_1*y, C_2*y/x, C_3/x, C_4/y, C_5*x/y, C_6*x}. If x and y are not both positive then four of these constants are negative and two are positive.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andres Cicuttin_, Nov 18 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014492 | OEIS-A329662

Position of first occurrence of n in A331410.

Conjecture: a(n) > a(n-1) and a(n) ~= sqrt(5)*a(n-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ali Sada_ and _Robert G. Wilson v_, Feb 28 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014493 | OEIS-A329725

a(1)=0, a(n) = n - (product of nonzero digits of n) - a(n-1).

a(10n+1)-a(10n-1)=1 for all positive integer n (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joshua Oliver_, Nov 19 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014494 | OEIS-A329733

Number of sequences n = b_1 < b_2 < ... < b_t = A329732(n) such that b_1*b_2*...*b_t is a perfect cube.

Conjecture: records are at 0 and the odd primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Nov 19 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014495 | OEIS-A329742

Indices n of Riemann zeta zeros for successive records of the normalized delta defined as d(n) = (z(n+1)-z(n))*(log(z(n)/(2Pi))/(2Pi)) where z(n) is the imaginary part of the n-th Riemann zero.

Conjectural next term: 77528045597.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Nov 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014496 | OEIS-A329799

Odd squarefree composite numbers k such that p-1 divides k-1 and p-1 does not divide (k-1)/2 for every prime p|k.

Conjecture: these are odd composite numbers k such that b^{(k-1)/2} == -1 (mod k) for some base b such that ord_{k}(b) = lambda(k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_ and _Thomas Ordowski_, Nov 21 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014497 | OEIS-A329917

Starting values for which iterations of A329623 diverge (conjectural).

Starting values for which iterations of A329623 diverge (conjectural). However, all these terms are so far only conjectural. We have no proof that the terms for which the trajectory seems to "explode" do not eventually end up in a cycle. For example, the 8th iterate of a(1) = 1373 is 5218725017016262626273. If the 2nd digit is changed from 2 to 0, then the further iterates grow to a length of 52 digits, but finally end up in a 2-cycle of 45-digit numbers (26...26273, 62...62637). (All members of cycles are listed in A328142.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014498 | OEIS-A329931

Reversal of base-n digits of largest prime < n^3.

Conjecture: a(n) < prevprime(n^3) for n >= 4. In other words, the most-significant base-n digit is larger than the least-significant base-n digit. This conjecture seems to hold for the analogous sequence with n^2, but fails for powers higher than 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Dougherty-Bliss_, Nov 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014499 | OEIS-A329934

a(1)=1, a(2)=1, a(n) = (number of times a(n-1) has appeared before) + (number of times a(n-2) has appeared before).

Conjecture: This sequence grows logarithmically.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joshua Oliver_, Nov 24 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014500 | OEIS-A330005

a(n) counts the square-words immediately before a(n), with a(1) = 0.

Conjecture: Every integer will appear in the sequence. (Although very slowly: after 10000 terms, the biggest term is 5.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Nov 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014501 | OEIS-A330024

a(n) = floor(n/z) where z is the number of zeros in the decimal expansion of 2^n, and a(n)=0 when z=0.

Conjecture: For every sequence which can be named as "digit k appears m times in the decimal expansion of 2^n", the sequences are finite for 0 <= k <= 9 and any given m >= 0. Every digit from 0 to 9 are inclined to appear an equal number of times in the decimal expansion of 2^n as n increases. Conjecture: a(n) = 33 (= floor(10/log_10(2))) for all sufficiently large n. - _Pontus von Brömssen_, Jul 23 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Metin Sariyar_, Nov 27 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014502 | OEIS-A330070

a(n) is the smallest integer k > 1 such that k cannot be expressed as a sum of distinct earlier terms nor is a multiple of any earlier term.

Conjecture: Composite terms k greater than 25 will have either two or six nontrivial divisors (that is, divisors d with 1 < d < k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Stauduhar_, Nov 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014503 | OEIS-A330073

Irregular triangle read as rows in which row n is the result of iterating the operation f(n) = n/5 if n == 0 (mod 5), otherwise f(n) = 5*(floor(n/5) + n + 1), terminating at the first occurrence of 1.

Conjecture: For any initial value n >= 1, there is a number k such that f^{k}(n) = 1, where f^{0}(n) = n and f^{k}(n) = f(f^{k - 1}(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Davis Smith_, Nov 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014504 | OEIS-A330076

Numbers k such that H(k) > d(k), where H(k) = Sum_{1 <= i < j <= d(k)} 1/(d_j - d_i) is sum over ordered pairs of divisors of k, and d(k) is the number of divisors of k.

Erdős and Nicolas conjectured that this sequence is finite with only 14 terms. Nicolas (2005) states that the conjecture has been checked up to 10^6. Conjecture checked up to 2*10^10. - _Giovanni Resta_, Dec 01 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Nov 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014505 | OEIS-A330077

a(n) = numerator of Sum_{1 <= i < j <= d(n)} 1/(d_j - d_i), sum over ordered pairs of divisors of n, where d(n) is the number of divisors of n.

Erdős and Nicolas conjectured that H(n) = a(n)/A330078(n) < d(n) for all n > 5040.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Nov 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014506 | OEIS-A330078

a(n) = denominator of Sum_{1 <= i < j <= d(n)} 1/(d_j - d_i), sum over ordered pairs of divisors of n, where d(n) is the number of divisors of n.

Erdős and Nicolas conjectured that H(n) = A330077(n)/a(n) < d(n) for all n > 5040.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Nov 30 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014507 | OEIS-A330154

Each term is visited according to the rule: move left (resp. right) a(n) places if a(n) is prime (resp. not prime), starting at n = 1: lexicographically earliest such sequence of distinct positive integers.

Conjectured to be a permutation of the positive integers: see Example for computational results.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Dec 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014508 | OEIS-A330155

Triangle read by rows. Given n enumerated cards in a stack, with 1 at the top and n at the bottom, repeat the following process k times: remove the card in the middle (at position (size of the stack)/2, rounding up), and move the card at the bottom of the stack to the top. T(n,k) is the number of the last card removed.

Conjecture: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wilmer Emiro Castrillon Calderon_, Dec 03 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014509 | OEIS-A330162

For imaginary fundamental discriminants -d, define b(-d) to be the smallest prime p such that Kronecker(-d,p) = 1. Sequence gives d such that b(-d)^3 > d/4 > b(-d)^2.

Note that 5460 is conjectured to be the largest term in A003644. Therefore, it seems that b(-d) < sqrt(d/4) for all d > 5460; it seems that b(-d) < (d/4)^(1/3) for all d > 1154008.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Dec 04 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014510 | OEIS-A330165

Odd terms in A003171: negated odd discriminants of orders of imaginary quadratic fields with 1 class per genus.

It is conjectured that this is the full list. Otherwise, there could be at most one more term d such that -d is a fundamental discriminant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Dec 04 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014511 | OEIS-A330191

For -d == 0, 1 (mod 4), let E(-d) to be the exponent of the class group of binary quadratic forms with discriminant -d, b(-d) to be the smallest prime p such that Kronecker(-d,p) = 1, then sequence gives d such that E(-d) > 2 and b(-d) > sqrt(d/4).

For -d == 0, 1 (mod 4), we want to determine the size of b(-d). Let R = Z[sqrt(-d)/2] if -d == 0 (mod 4), R = Z[(1+sqrt(-d))/2)] otherwise, note that R is not necessarily a Dedekind domain. It is conjectured that if Kronecker(-d,p) = 1, then p*R is the product of two distinct prime ideals of R (this is obviously true if -d is a fundamental discriminant). It also seems that if p*R = I*I', then I^(k*e) must be principal, e = E(-d) (again, this is true if -d is fundamental). If these statements are indeed true, let t*O_k = I^(k*e), then t/p is not in R, and the norm of t over R is p^e. Define f(x,y) = x^2 + (d/4)*y^2 if -d == 0 (mod 4), x^2 + x*y + ((d+1)/4)*y^2 otherwise, it is easy to see f(x,y) = p^(k*e) has integral solutions (x,y) such that gcd(x,y) = 1. Note that 7392 is conjectured to be the largest term in A003171. Therefore, it seems that b(-d) < sqrt(d/4) for all d > 7392.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Dec 04 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014512 | OEIS-A330221

Numbers d such that -d is a fundamental discriminant and all primes smaller than 2*sqrt(d)/Pi ramify or remain inert in the ring of integers of Q(sqrt(-d)).

But there are other d such that the ideal class group of O_K has exponent 2. In fact, the exponent is <= 2 if and only if: for all primes p < 2*sqrt(d)/Pi, either (a) Kronecker(-d,p) = 0 or -1, or (b) Kronecker(-d,p) = 1, and 4*p^2 - d is a square. Here 2*sqrt(d)/Pi can be replaced by sqrt(d); conjecturally, if 2*sqrt(d)/Pi is replaced by sqrt(d/4), we get exactly the sequence A003644.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Dec 06 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014513 | OEIS-A330266

Number of ways to shuffle a deck of 4n cards, with 4 cards in each of n ranks, so that adjacent cards have different ranks.

Conjecture: Limit_{n->oo} a(n)/(4n)! = 1/e^3. The conjecture is based on the observation of the asymptotic behavior of A007060 and A193624; it seems that it can be generalized in the following way. Let b(n) be the number of ways to shuffle a deck of k*n cards, with k cards in each of n ranks, so that adjacent cards have different ranks. Then, lim_{n->oo} b(n)/(kn)! = 1/e^(k-1); maybe we could prove it with the help of rook polynomials theory or in some other way. - _Sergey Kirgizov_, Sep 29 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Radcliffe_, Dec 07 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014514 | OEIS-A330276

NSW pseudoprimes: odd composite numbers k such that A002315((k-1)/2) == 1 (mod k).

They conjectured that this sequence is infinite, and their conjecture was proved by Grantham (2021). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Dec 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014515 | OEIS-A330277

a(n) = k such that A083016(k) = prime(n), or -1 if no such k exists.

Conjecture: every prime occurs in A083016, so k always exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Dec 08 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014516 | OEIS-A330288

Start with the list of numbers from 1 to n. As long as the list has odd and even elements, append to the list the sum of the smallest odd and smallest even elements, taken with multiplicity, and delete these terms from the list. a(n) is the largest element in the final list.

Conjecture: a(4)=10 is the only even term. - _Jon E. Schoenfield_, Dec 28 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ali Sada_, Dec 09 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014517 | OEIS-A330354

Expansion of e.g.f. Sum_{k>=1} log(1 + x)^k / (k * (1 - log(1 + x)^k)).

Conjecture: a(n) ~ n! * (-1)^n * Pi^2 * exp(n) / (24 * (exp(1) - 1)^(n+1)). - _Vaclav Kotesovec_, Dec 16 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Dec 11 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014518 | OEIS-A330358

a(n) = n mod 5 + n mod 25 + ... + n mod 5^k, where 5^k <= n < 5^(k+1).

Conjecture: For b >= 2, consider the function s(n,b) = Sum_{1 <= b^j <= n} (n mod b^j) from p. 8 in Dearden et al. (2011). Then s(b*n + r, b) = b*s(n,b) + r*N(n,b) for 0 <= r <= b-1, where N(n,b) = floor(log_b(n)) + 1 is the number of digits in the base-b representation of n. As initial conditions, we have s(n,b) = 0 for 1 <= n <= b. (This is a generalization of a result by _Robert Israel_ in A049802.) If A_b(x) = Sum_{n >= 1} s(n,b)*x^n is the g.f. of the sequence (s(n,b): n >= 1) and the above conjecture is correct, then it can be proved that A_b(x) = b * A_b(x^b) * (1-x^b)/(1-x) + x * ((b-1)*x^b - b*x^(b-1) + 1)/((1-x)^2 * (1-x^b)) * Sum_{k >= 1} x^(b^k). Conjecture: a(5*n+r) = 5*a(n) + r*A110592(n) = 5*a(n) + r*(floor(log_5(n)) + 1) for n >= 1 and r = 0, 1, 2, 3, 4. If the conjecture above is true, the g.f. A(x) satisfies A(x) = 5*(1 + x + x^2 + x^3 + x^4)*A(x^5) + x*(1 + 2*x + 3*x^2 + 4*x^3)/(1 - x^5) * Sum_{k >= 1} x^(5^k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, Dec 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014519 | OEIS-A330382

Composite numbers k such that k-1 divides 2^k-2.

Conjecture: k-1 | 2^k-2 for k = (2^n-1)^3 if and only if n(n-1) | 2^n-2 for n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_ and _Thomas Ordowski_, Dec 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014520 | OEIS-A330388

Expansion of e.g.f. Sum_{k>=1} (-1)^(k + 1) * log(1 + x)^k / (k * (1 - log(1 + x)^k)).

Conjecture: a(n) ~ n! * (-1)^(n+1) * Pi^2 * exp(n) / (24 * (exp(1) - 1)^(n+1)). - _Vaclav Kotesovec_, Dec 16 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Dec 12 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014521 | OEIS-A330408

Table of A(n,k) read by antidiagonals, where A(1,k)=k; A(n,k) is the least multiple of n >= A(n-1,k).

The main diagonal is conjectured to be A166447.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ali Sada_, Dec 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014522 | OEIS-A330411

a(n) is the index of the first 0 term in the rumor sequence with initial 0th term 1 and parameters b = 3 and n.

"A rumor sequence (running modulus recurrence sequence) is defined as follows: fix integer parameters b > 1 and n > 0. Set z[0] = any integer, and, for k > 0, define z[k] to be the least nonnegative residue of b*z[k-1] modulo (k+n). The rumor sequence conjecture states that all such rumor sequences are eventually 0." [Copied from the comments for A177356]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, Dec 13 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014523 | OEIS-A330445

Expansion of e.g.f.: Sum_{k>=1} log(1 + (exp(x) - 1)^k)/k.

Conjecture: a(n) ~ (n-1)! / (log(2))^(n-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Dec 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014524 | OEIS-A330450

Expansion of e.g.f. Sum_{k>=1} log(1 + x)^k / (k * (1 - log(1 + x)^k)^2).

Conjecture: a(n) ~ n! * (-1)^n * zeta(3) * n * exp(n) / (8 * (exp(1) - 1)^(n+2)). - _Vaclav Kotesovec_, Dec 16 2019

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Dec 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014525 | OEIS-A330466

Irregular triangle read by rows: T(n,k) is the number of parts in the partition of n into k consecutive parts that differ by 2, n >= 1, k >= 1, and the first element of column k is in row k^2.

Conjecture: row sums give A066839.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Apr 30 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014526 | OEIS-A330487

Sum of those x(p)*y(p) with p <= n, where p is a prime congruent to 1 modulo 4, and p = x(p)^2 + y(p)^2 with 1 <= x(p) <= y(p).

Conjecture: Let b(n) be the sum of all primes p <= n with p == 1 (mod 4). Then a(n)/b(n) = 1/Pi + O(1/sqrt(n)). A classical theorem of Euler (conjectured by Fermat) states that any prime p == 1 (mod 4) can be written uniquely as x^2 + y^2 with 1 <= x <= y.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 15 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014527 | OEIS-A330498

Expansion of e.g.f. Sum_{k>=1} log(1 + log(1/(1 - x))^k) / k.

a(n) ~ n! * c / (n * (1 - exp(-1))^n), where c = 0.6931..., conjecture: c = log(2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Dec 16 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014528 | OEIS-A330521

Lexicographically earliest sequence of distinct positive terms such that the digitsum of a(n) ends a(n+1).

This is conjectured to be a permutation of the integers > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carole Dubois_ and _Eric Angelini_, Mar 01 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014529 | OEIS-A330532

Triangular multiply-perfect numbers.

Corresponding values of numbers k and h: (1, 2, 3, 4, 5, 7, 10, 13, 17, 19, 31, 61, 89, 107, 127, ...), (1, 3, 7, 15, 31, 127, 1023, 8191, 131071, 524287, 2147483647, ...), where h = 2^k - 1. Conjecture: numbers h are numbers from A066175 (sigma(phi(sigma(h))) = h). Conjecture: union of even perfect numbers from A000396 and 3-perfect numbers 120 and 523776.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Dec 17 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014530 | OEIS-A330547

a(1)=2; thereafter a(n) = a(n-1) + (-1)^(n+1)*(prime(n)-prime(n-1)) (where prime(k) denotes the k-th prime).

Conjecture. Let k be an integer and X_k be the set of all n such that (-1)^n*a(n)=2k-1. If a, b are integers and a<>0, then X_k contains infinitely many terms of the arithmetic progression {a*n+b: n integer}. - _M. Farrokhi D. G._, Nov 12 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 19 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014531 | OEIS-A330607

Array read by rows: T(n,k) is the number of solutions to the equation Sum_{i=1..n} x_i^2 == k (mod 5) with x_i in 0..4, where n >= 0 and 0 <= k <= 4.

Conjecture: T(n,k=1) = A071304(n)/A071304(n-1) for n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, Dec 20 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014532 | OEIS-A330652

Decimal expansion of (phi^phi - 1/phi)/2, where phi = (1 + sqrt(5))/2 = A001622.

Conjecture: (phi^phi - 1/phi)/2 = integral_{x=0..oo} exp(-x*(2 + phi)) * (cosh(x) + sqrt(5) * sinh(x))^phi dx.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Dec 29 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014533 | OEIS-A330700

a(n) = (n - 1)*n*(2*n^2 + 4*n - 1)/6.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Dec 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014534 | OEIS-A330705

Numbers k such that A173419(k) < A173419(k-1).

Conjecture: there are no primes in this sequence. This conjecture is directly related to the conjecture in A173419.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Dec 26 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014535 | OEIS-A330715

a(1), a(2), a(3) = 1; a(n) = (a(n-1) mod a(n-3)) + a(n-2) + 1.

Conjectures from _Colin Barker_, Dec 28 2019: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthew Niemiro_, Dec 27 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014536 | OEIS-A330719

a(n) = denominator(Sum_{k=1..n} (2^(k-1) - 1)/k).

Conjecture: if p is an odd prime, then p | A330718(p+1) - a(p+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_ and _Thomas Ordowski_, Dec 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014537 | OEIS-A330724

Sum of prime factors (with multiplicity) of the n-th composite number coprime to 6.

It appears that odd numbers appear rarely in this sequence. I conjecture that all the odd numbers above 15 and all the even numbers above 10 will at least once show up in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nico Grimm_, Dec 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014538 | OEIS-A330731

Binary sequence created by greedily remaining as normal as possible: starting with the empty sequence, repeatedly find the longest tail (or suffix) which is followed by one digit more frequently than the other, and append the digit which follows said tail less often. 0 is appended if no such inequality is found.

a(n) is conjectured to be normal by virtue of its construction.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aresh Pourkavoos_, Dec 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014539 | OEIS-A330733

Triangle read by rows in which row n is the "complete rhythm" of n (see Comments for precise definition).

Right diagonal is A002033 (conjectured). Any prime column stripped of zeros also yields A002033 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Hood_, Dec 28 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014540 | OEIS-A330744

Primorial deflation of A133411(n), where A133411(n) is the smallest highly composite number of the form k*a(n-1) where k is an integer greater than 1.

Conjectured to be a subsequence of A330745.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 10 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014541 | OEIS-A330745

Primorial deflation of A019505(n), where A019505(n) is smallest number with same number of divisors as 2*A019505(n-1), starting from A019505(1) = 1.

Conjectured to be a subsequence of A329902.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 10 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014542 | OEIS-A330777

Numbers k such that k and Lucas(k) have the same number of divisors.

Conjecture: all terms are of the form 2^k*p for k >= 0 and p prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Dec 31 2019 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014543 | OEIS-A330807

a(1) = 2; a(n+1) = Sum_{k=1..n} {S(a(k)): S(a(k)) = S(a(n))}, where S is sopfr (A001414).

The primes do not appear in their natural order (e.g., 31 precedes 29). If the lesser of twin primes p is a(k) and the greater twin has not already occurred, then a(k+2) is the greater twin. Whereas if a composite number appears, it may appear more than once, a prime > 2, if it appears, can appear once only. Open question: Do all primes (except for 3) eventually appear?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jan 01 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014544 | OEIS-A330816

Numbers that reach 1 under the iterations of the map k -> k/d(k) if d(k) | k, and k -> k otherwise, where d(k) is the number of divisors of k (A000005).

Conjecture: the sequence is finite; all 45 terms are in Corneth's a-file. Heuristic evidence: I took the first few terms C and made the Cartesian product with C and the 101-smooth numbers <= 10^8, seeing which were terms and removing duplicates. This process was repeated a few times until no more new terms were found. The largest number of divisors of any of these terms is < 10^6. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Jan 01 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014545 | OEIS-A330849

First element of the first maximal isospectral chain of length n.

Isospectral Chain Conjecture: There exist isospectral chains of any positive length.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Walter Kehowski_, Feb 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014546 | OEIS-A330852

Numerator of the rational number A(n) that appears in the formula for the n-th cumulant k(n) = (-1)^n*2^n*(A(n) - (n - 1)!*zeta(n)) of the limiting distribution of the number of comparisons in quicksort, for n >= 2, with A(0) = 1 and A(1) = 0.

Hennequin conjectured his cumulant formula in his 1989 paper and proved it in his 1991 thesis.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, Apr 28 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014547 | OEIS-A330853

First occurrences of gaps between primes 6k+1: gap sizes.

Conjecture: the sequence is a permutation of all positive multiples of 6, i.e., all positive terms of A008588. Conjecture: a(n) = O(n). See arXiv:2002.02115 (sect.7) for discussion.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexei Kourbatov_, Apr 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014548 | OEIS-A330860

Denominator of the rational number A(n) that appears in the formula for the n-th cumulant k(n) = (-1)^n*2^n*(A(n) - (n - 1)!*zeta(n)) of the limiting distribution of the number of comparisons in quicksort, for n >= 2, with A(0) = 1 and A(1) = 0.

Hennequin conjectured his cumulant formula in his 1989 paper and proved it in his 1991 thesis.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, Apr 28 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014549 | OEIS-A330875

Numerator of the fraction fr(n) that appears in the n-th cumulant k(p) = fr(n) - (-2)^n*(n-1)!*zeta(n) of the limiting distribution of the number of comparisons in quicksort, for n >= 2, starting with fr(0) = 1 and fr(1) = 0.

Hennequin conjectured his cumulant formula in his 1989 paper and proved it in his 1991 thesis.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, Apr 29 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014550 | OEIS-A330876

Denominator of the fraction fr(n) that appears in the n-th cumulant k(n) = fr(n) - (-2)^n*(n-1)!*zeta(n) of the limiting distribution of the number of comparisons in quicksort, for n >= 2, starting with fr(0) = 1 and fr(1) = 0.

Hennequin conjectured his cumulant formula in his 1989 paper and proved it in his 1991 thesis.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, Apr 29 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014551 | OEIS-A330900

Number of terms of A066038 that do not exceed 10^n.

Luca & Moodley (2020) conjectured that a(n) ~ exp(gamma) * A006880(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, May 01 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014552 | OEIS-A330910

a(n-5) is the number of nonempty subsets of {1,2,...,n} such that the difference of successive elements is at least 5.

Conjectures from _Colin Barker_, May 17 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enrique Navarrete_, May 01 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014553 | OEIS-A331012

16 * squared area of triangles with integer sides i <= j <= k, such that more triples of sides produce the same area as for any smaller area.

Conjectured further terms are 226195200 (42 representations), 230630400 (52 representations), 372556800 (55 representations).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jan 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014554 | OEIS-A331020

Values k for successive maximal records of the function A defined as A(prime(k)) = log(prime(k)) - prime(k)/Pi(prime(k)) where Pi(prime(k)) is number of primes <= prime(k).

Chebyshev 1852, goes on to conclude that if we put Pi(x) = x/(log(x) - A(x)) has a limit as x -> +infinity, then a limit must be 1, not 1.08366 (A228211), as Legendre incorrectly conjectured in 1808.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jan 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014555 | OEIS-A331021

Numbers k such that gcd(k^2, 2^(k-1) - 1) > k.

Conjecture: each term is a multiple of a Wieferich prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_ and _Thomas Ordowski_, Jan 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014556 | OEIS-A331022

Numbers k such that the number of strict integer partitions of k is a power of 2.

Conjecture: This sequence is finite. Conjecture: The analogous sequence for non-strict partitions is: 0, 1, 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jan 10 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014557 | OEIS-A331041

Denominator of squared radius of inscribed circles of triangles with integer sides i <= j <= k, such that the number of triangles with this radius sets a new record. Numerators are A331040.

See A331040. Conjecturally, all terms a(n) are either 1 or 4 for n >= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jan 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014558 | OEIS-A331042

a(n) = 4 * squared radius of inscribed circles of triangles with integer sides i <= j <= k, such that the number of triangles with this radius sets a new record. If this radius is not a multiple of (1/4), a(n) = 0.

It is conjectured that all radii of incircles leading to records with the exception of the first two terms are multiples of 1/4, thus a(n) > 0 for all n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jan 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014559 | OEIS-A331164

Number of function evaluations when recursively calculating Fibonacci(n) with caching.

Conjecture: a(n) is O(log n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas König_, Jan 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014560 | OEIS-A331234

Triangular numbers having exactly 9 divisors.

Conjecture: a(5) is the final term of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Jan 12 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014561 | OEIS-A331272

Irregular triangle in which row n lists numbers m such that A330073(m,n) = 1.

The conjecture in A330073 claims that every positive integer appears in this triangle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Davis Smith_, Jan 13 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014562 | OEIS-A331294

Positions of records in A331293.

Conjectured: for n >= 9, a(n) = A000040(n-4)^2 = A001248(n-4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 17 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014563 | OEIS-A331330

a(n) is the number of sparse rulers of length n where the length of the first segment is unique.

Conjecture: a(n) ~ 2^n / (n * log(2)). - _Vaclav Kotesovec_, Nov 16 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jan 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014564 | OEIS-A331343

a(n) = lcm(1,2,...,n) * Sum_{k=1..n} (2^(k-1) - 1) / k.

Conjecture: for n > 3, if n^3 | a(n), then n is prime. If so, there are no such pseudoprimes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_ and _Thomas Ordowski_, Jan 14 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014565 | OEIS-A331404

a(0) = 1; a(n) = Sum_{k=1..n} (-1)^(n - k) * binomial(n,k) * a(k-1) * a(n-k).

Conjecture: the e.g.f. of this sequence is the inverse function of y = x - 1/x - log(x) in a neighborhood of x = 1 on the complex plane, and its convergence radius is sqrt(3) - Pi/3 = 0.6848532563... - _Jianing Song_, Sep 30 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Jan 16 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014566 | OEIS-A331432

Triangle T(n,k) (n >= k >= 0) read by rows: T(n,0) = (1+(-1)^n)/2; for k>=1, set T(0,k) = 0, S(n,k) = binomial(n,k)*binomial(n+k+1,k), and for n>=1, T(n,k) = S(n,k)-T(n-1,k).

For Table IV on page 93, it is simplest to ignore the minus signs. The present triangle then matches all the given terms in that triangle, so it seems best to define the triangle by the recurrences given here, and to conjecture (strongly) that this is the same as Ser's triangle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 17 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014567 | OEIS-A331460

Irregular triangle read by rows in which row n is the result of iterating the operation f(n) = n/7 if n == 0 (mod 7), otherwise f(n) = 7*(n + ceiling(n/7)), terminating at the first occurrence of 1.

Conjecture: For any number n >= 1, there exists a k such that f^{k}(n) = 1, where f^{0}(n) = n and f^{k + 1}(n) = f(f^{k}(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Davis Smith_, Jan 23 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014568 | OEIS-A331491

a(n) is the permanent of a 2n X 2n antisymmetric Toeplitz matrix M(2n) whose first row consists of a single zero followed by successive positive integers repeated (A004526).

Conjecture: for n > 0, det(M(2n)) = n^2 = A000290(n) with det(M(0)) = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jan 18 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014569 | OEIS-A331499

T(n,k) is the number of nonintersecting arch configurations with n arches on a horizontal axis that have a starting parent arch configuration of k arches.

T(n,n-2) = A000108(n-2) + A000108(n-3) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger Ford_, Feb 16 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014570 | OEIS-A331506

Least primitive root g < prime(n) of the n-th prime with g a product of two Fibonacci numbers, or 0 if such a number g does not exist.

Conjecture 1: a(n) > 0 for all n > 0. In other words, for each prime p there are two Fibonacci numbers F(k) and F(m) with F(k)*F(m) < p such that F(k)*F(m) is a primitive root modulo p. It seems that Conjecture 1 can be strengthened as follows: For any prime p, there is a primitive root g < p modulo p such that g/F(2) = g or g/F(3) = g/2 or g/F(4) = g/3 is a Fibonacci number. We have verified this strong version for all primes p < 5*10^9. We also have the following conjecture similar to Conjecture 1. Conjecture 2. For any prime p, there are two Lucas numbers L(k) and L(m) with k >= m >= 0 and L(k)*L(m) < p such that L(k)*L(m) is a primitive root modulo p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 18 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014571 | OEIS-A331544

Primes that begin a run of consecutive primes whose first differences are strictly increasing.

Conjecture: Average length of each run = e, converging from below.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Woodward_, Jan 19 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014572 | OEIS-A331546

a(n) = |{k^(k+1)+(k+1)^k (mod prime(n)): k = 0..prime(n)-1}|.

Conjecture: |{k^(k+1)+(k+1)^k (mod p): k = 0..p-1}| = (1-1/e)*p + O(p^(1/2)), where p denotes a prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 19 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014573 | OEIS-A331548

15-adic integer x = ...2AA66B44A40E43797853AD13 satisfying x^5 = x; also x^3 = -x; (x^2)^3 = x^2 = A331550; (x^4)^2 = x^4 = A331549.

Conjecture: If k is the number of prime factors congruent to 1 (mod 4) of an integer n, then there are exactly k n-adic integers x satisfying x^5 = x, while not satisfying x^h = x for h = 2, 3, or 4. This does not count -x, which also satisfies, in each case. - _Patrick A. Thomas_, Mar 31 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick A. Thomas_, Jan 20 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014574 | OEIS-A331562

Number A(n,k) of sequences with k copies each of 1,2,...,n avoiding absolute differences between adjacent elements larger than one; square array A(n,k), n>=0, k>=0, read by antidiagonals.

Row 2, the sequence of central binomial numbers A000984, satisfies the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and all positive integers n and r (see Meštrović, equation 39). This is also known to be true for row 3 (A103882) and row 4 (A177316). We conjecture that each row sequence of the table satisfies the same congruences. - _Peter Bala_, Oct 26 2024.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jan 20 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014575 | OEIS-A331564

Number of twin primes p < A014574(n), such that p*A014574(n) is in A014574.

Conjecture: a(n) > 0 for all n, except n = 10 where A014574(n) = 108. The conjecture is true for A014574(n) < 1000000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Jan 20 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014576 | OEIS-A331625

Numbers k such that both k and k+1 are exceptional (A072066).

Conjecture: for every p > 0, there exist infinitely many k such that k, k+1, ..., k+p-1 are all exceptional numbers. In specific, there exist infinitely many k such that both k and k+1 are exceptional.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Jan 22 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014577 | OEIS-A331761

a(n) = Sum_{i=1..n, j=1..n, gcd(i,j)=2} (n+1-i)*(n+1-j).

Conjecture: As n -> oo, a(n) ~ C*n^4/Pi^2, where C is about 0.3775. - _N. J. A. Sloane_, Jul 03 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014578 | OEIS-A331786

a(n) is the largest m such that there exists N such that none of S(N), S(N+1), ..., S(N+m-1) is divisible by n, where S(N) is the sum of digits of N.

Conjectures from _Colin Barker_, Jan 26 2020: (Start) (End) [This conjecture is correct.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Jan 25 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014579 | OEIS-A331788

a(n) is the smallest m such that for any N, at least one of S(N), S(N+1), ..., S(N+m-1) is divisible by n, where S(N) is the sum of digits of N.

Conjectures from _Colin Barker_, Jan 26 2020: (Start) (End) [This conjecture is correct.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Jan 25 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014580 | OEIS-A331807

a(n) is the smallest prime number p > n, not yet in the sequence, such that p is a palindrome when written in base n.

2) A conjecture: No lowest primes need more than three digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Colin M Ready_, Feb 22 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014581 | OEIS-A331810

Smallest integer x such that prime(n) divides 1 + x + x^2 + ... + x^(q-1) where q = A023503(n), or 0 if no such x exists.

Conjecture: every integer > 1 is in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014582 | OEIS-A331814

Fourier coefficients of the boundary of the Mandelbrot set (normalized by a power of two).

a(n) = 2^(2*n+1)*b(n) where b(n) is the unique sequence of complex numbers such that f(z) := z + Sum_{n>=0} (b(n)*z^-n) defines an analytic homeomorphism (biholomorphic bijection) between the complement of the unit disk and the complement of the Mandelbrot set, sometimes known as the "Jungreis function". (The b(n) are rationals, so we multiply them by the appropriate power of two to make them integers; this is equivalent to a simple rescaling of the complex plane.) It is conjectured that |b(n)| <= 1/n, so |a(n)| <= 2^(2*n+1)/n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David A. Madore_, Jan 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014583 | OEIS-A331859

The total number of elastic collisions between a block of mass n, a block of mass 1, and a wall.

Conjecture: The values of n for which a(n) != A121854(n) is a subset of A331903.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Jan 29 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014584 | OEIS-A331905

Number of spanning trees in the multigraph cube of an n-cycle.

I conjecture that a(n) = gcd(n,2) * n * (A005822(n))^2. [This is correct - see the Formula section. - _N. J. A. Sloane_, Feb 06 2020]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David J. Seal_, Jan 31 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014585 | OEIS-A331912

Lexicographically earliest sequence of positive integers that have at most one distinct prime index already in the sequence.

Conjecture: a(n)/A331784(n) -> 1 as n -> infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Feb 01 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014586 | OEIS-A331920

Total number of reduced pipe dreams (a.k.a. rc-graphs) for all permutations in S_n.

log_2(a(n))/n^2 -> c (conjectured by Stanley in arxiv:1704.00851).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sara Billey_, Feb 01 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014587 | OEIS-A331921

Number of iterations of the map G -> Aut(G) on the cyclic group of order n needed to reach stability; or -1 if the iteration never stabilizes.

Conjecture: a(n) != -1 for all n. The following two sequences are conjectured to be correct and to stabilize at the last term:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sébastien Palcoux_, Feb 01 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014588 | OEIS-A331944

a(n)/ceiling(6^(n-7)) is the expected number of rolls of a fair 6-sided die in a game where the player starts at 0, advances the position by the outcome of the die's roll until exactly position n is reached. Positions beyond n are avoided by staying at the last visited position, but counting the rolls.

Conjectures from _Colin Barker_, Feb 21 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Feb 19 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014589 | OEIS-A332009

Irregular triangle of numerators of the average value of the first letter over all derangements of {1, 2, ..., n} with k descents.

Conjecture: T(n, 1) = A083329(n - 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Feb 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014590 | OEIS-A332010

Irregular triangle of denominators of the average value of the first letter over all derangements of {1, 2, ..., n} with k descents.

Conjecture: T(n, 1) = 2^(n-2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Feb 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014591 | OEIS-A332018

a(n) = A038502(A000265(n)) if n is even or n == 0 (mod 3), a(n) = A038502(A000265(5*n + 1)) otherwise.

There is a conjecture which states that for any value of n > 0 there is a k such that a^{k}(n) = 1 or a^{k}(n) enters one of a finite number of periodic cycles, where a^{0}(n) = n and a^{k + 1}(n) = a(a^{k}(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Davis Smith_, Feb 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014592 | OEIS-A332020

Positive integers m which are quadratic residues modulo prime(m).

Conjecture: Let A(x) be the number of terms not exceeding x. Then A(x)/x has the limit 1/2 as x tends to the infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014593 | OEIS-A332021

Elements of the set {m > 0: m is a quadratic nonresidue modulo prime(m)}.

The conjecture in A332020 implies that the number of terms not exceeding x is about x/2 asymptotically.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014594 | OEIS-A332056

a(1) = 1, then a(n+1) = a(n) - (-1)^a(n) Sum_{k=1..n} a(k): if a(n) is odd, add the partial sum, else subtract.

Conjecture from Barker confirmed by recurrence formula. - _Ray Chandler_, Jul 31 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _M. F. Hasler_, Feb 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014595 | OEIS-A332067

a(n) is the square spiral number of the initial digit of the number placed at the n-th move of the Prime scrabble game: placing integers on a grid one digit per cell as to form primes, with minus sign in case of vertical placement.

Since the numbers cannot be split (cf. rule 1), we know that a run of only d consecutive empty squares in either direction will remain empty forever once we have used all numbers < 10^d (cf. EXAMPLE). This contrasts with a variant where the numbers can be split "around" already occupied squares. In that variant one can conjecture that each square will eventually be filled.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_ and _Eric Angelini_, May 05 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014596 | OEIS-A332084

Triangle read by rows: T(n,k) is the smallest m >= 0 such that floor(Pi*n^m) == k (mod n), -1 if one does not exist, k = 0..n-1.

Conjecture: Pi is absolutely normal, meaning that -1 will never appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Davis Smith_, Aug 22 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014597 | OEIS-A332088

Primes which yield again a prime when the digits are taken according to the lexicographically first superpermutation of corresponding order and of minimal length.

In March 2014, Ben Chaffin showed that minimal superpermutations of order n = 5 have length 153, and found all 8 distinct superpermutations of this kind (the lexicographically first being non-palindromic), so the 5-digit terms are known. For n = 6, Robin Houston has found a few months later several superpermutations of length 872 (one less than the previously conjectured minimal length), but we still don't know which is the shortest (and/or lexico-first) superpermutation for that case.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jul 28 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014598 | OEIS-A332089

Irregular table read by rows, where row n lists the lexicographically first superpermutation of minimal length over {1, ..., n}.

In March 2014, Ben Chaffin showed that minimal superpermutations of order n = 5 have length 153, and found all 8 distinct superpermutations of this kind; the (non-palindromic) lexicographically first one is row 5 of this table. For n = 6, Robin Houston has found a few months later several superpermutations of length 872 (one less than the previously conjectured minimal length), but we still don't know which is the shortest (and/or lexico-first) superpermutation for that case.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jul 31 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014599 | OEIS-A332101

Least m such that m^n <= Sum_{k<m} k^n.

a(n) = round(n / log(2)) + 2. (Conjectured; verified up to 10^4, in particular for 3525/log(2) = 5085.500019... and 7844/log(2) ~ 11316.49990...) The conjectured a(n) = round(n / log(2)) + 2 has been verified up to 5 * 10^4 and also for n=418079, 2639344, 10975455, 24590254, where n/log(2) ~ 0.5+integer. At n = 777451915729368, A078607(n) + 2 = ceiling(n / log(2)) + 2 = round(n / log(2)) + 3. So the two conjectures disagree. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 14 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014600 | OEIS-A332201

Sum of three cubes problem: a(n) = integer x with the least possible absolute value such that n = x^3 + y^3 + z^3 with |x| >= |y| >= |z|, or 0 if no such x exists.

It is known that there is no solution for n congruent to +-4 (mod 9), but it is now conjectured that there is a solution (and probably infinitely many such) for all other numbers. The numbers n = 0, 1 and 2 are the only cases for which infinite families of parametric solutions are known, for other n the solutions seem to be sporadic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014601 | OEIS-A332215

Mersenne-prime fixing variant of A243071: a(n) = A243071(A332213(n)).

For example, for n > 1, A007814(a(n)) = A007814(n) - A209229(n), that is, this map preserves the 2-adic valuation of n, except when n is a power of two, in which cases that value is decremented by one, and in particular, a(2^k * n) = 2^k * a(n) for all n > 1. Also, like A243071, this bijection maps primes to the terms of A000225 (binary repunits). However, the "tuning" (A332213) has a specific effect that each Mersenne prime (A000668) is mapped to itself. Therefore the terms of A335431 are fixed by this map. Furthermore, I conjecture that there are no other fixed points. For the starters, see the proof in A335879, which shows that at least none of the terms of A335882 are fixed.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014602 | OEIS-A332223

a(1) = 1, and for n > 1, a(n) = A005940(1+sigma(A156552(n))).

This rather meager empirical evidence motivates a conjecture that no quotient a(n)/n may be an even integer, and particularly, never a power of 2 larger than one, which (when translated back to the ordinary, unconjugated sigma) claims that it is not possible that sigma(n) = 2^k * n + 2^k - 1, for any n > 1, k > 0. See also A336700 and A336701, where this leads to a rather surprising empirical observation. a(A324201(n)) = A003961(A324201(n)). [It's an open problem whether A324201 gives all such solutions]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 12 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014603 | OEIS-A332242

Numbers k such that k! has exactly k nonzero decimal digits.

Conjecture: these 6 terms are the only terms of the sequence, i.e., there are no terms larger than 57.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Feb 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014604 | OEIS-A332300

The least prime factor of the numerator of Bernoulli(2*n), or 1 if the numerator is 1.

Conjecture: for prime p > 3, p is a prime factor of the numerator of Bernoulli(2*p), thus the conjecture implies that a(p) <= p for prime p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Feb 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014605 | OEIS-A332446

Numbers k for which A087808(sigma(k)) is equal to A087808(2*k).

Conjecture: includes all terms of A007520. - _Bill McEachen_, Dec 10 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 14 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014606 | OEIS-A332541

Denominators of coefficients in a series for Euler's constant gamma.

Conjecture: For n > 0, a(n) is a Zumkeller number (A083207). Verified form all n in [2,19]. - _Ivan N. Ianakiev_, Feb 17 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 16 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014607 | OEIS-A332553

a(n) = n + A082183(n) - A082184(n).

To study the lowest values taken by a(n), consider the record high values of n/a(n). The data suggests two conjectures. Conjecture 1: The record high values of n/a(n) are j/2 + 1 for j = 2,3,4,5,... and occur at n = j*(j+1)/2 - 1. Conjecture 2: Let j = 2,3,4,5,... For 1 <= n < T_j - 1, a(n) > 2*n/(j+2). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 21 2020, at the suggestion of _Bradley Klee_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014608 | OEIS-A332562

a(n) = number formed by concatenating the decimal digits of 44, 45, 46, ..., 44+n.

As of Feb 24 2020, it is an open question as to whether there is an N such that a(N) is divisible by 44+N+1. If such an N exists, N > 10^11, as shown by _Joseph Myers_ (see A332580).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Feb 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014609 | OEIS-A332602

Tridiagonal matrix M read by antidiagonals: main diagonal is 1,2,2,2,2,..., two adjacent diagonals are 1,1,1,1,1,...

For the proofs of the 2 X 2, 3 X 3 and 4 X 4 conjectures, see the comments in the respective A-numbers A001519, A080937 and A080938. - _Wolfdieter Lang_, Mar 30 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 06 2020, following a suggestion from _Gary W. Adamson_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014610 | OEIS-A332603

Working over the alphabet {1,2,3}, start with a(1) = 1; then a(n+1) is made by inserting a letter into a(n) at the rightmost possible position which makes a squarefree word (and the smallest letter if multiple letters are possible at that place).

Grytczuk et al. (2020) conjecture that this process never terminates. They also conjecture a(n) converges to a certain infinite word (the beginning of which is now given in A332604).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014611 | OEIS-A332604

Limiting word over the alphabet {1,2,3} defined by the process in A332603, or, in case that process terminates, the final term in A332603.

Grytczuk et al. (2020) conjecture that the process in A332603 never terminates, and report that they have computed the first 5000 terms of the limiting word.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014612 | OEIS-A332636

Two-parameter family of recursively defined triangles, T(m,t), whose rows right-padded with zeros appear in the limiting sequence of families of certain linear recursive sequences. The data presents the sequence of triangles T(2,t) by ascending antidiagonals.

The collection of k-th order recursions has a natural interpretation in terms of generalizations of the Fibonacci numbers. Define the Generalized Fibonacci numbers of order k by the recursion K(n) = Sum_{i=1..k} K(n-i), with initial conditions K(0)=0, K(i)=1, 1 <= i <= k-1. Further define G(n) = K(-n). (For k = 2, the [K(n)] are the Fibonacci numbers and for k = 3 they are the tribonacci numbers.) Then [G(n)] satisfies the recursion G(n) = G(n-k) - Sum_{i=1..k-1} G(n-k+i). (For purposes of reference later on we may suppose initial conditions G(i)=1, 1 <= i <= k-1, G(k)=0.) In other words, the recursions satisfied by the Generalized Fibonacci numbers over negative indices are

identical to the recursions we presented in this OEIS sequence above with t = 1 and m = 1 and the resulting triangle [T(i, j)] satisfies almost the same recursion as in sequence A118800 (the reason for the difference lies in the initial conditions). More precisely, the first row of the triangle [T(1, 1), T(1, 2) = [0, -(k-3)]. Boundary conditions for the rightmost and leftmost diagonals are given as follows: For i >= 1, T(i, 1) = -(2^(i-1)-1) and T(i, i+1) = (-1)^i (k-2+(-1)^i). Then for i >= 2, 2 <= j <= i, T(i, j) = 2*T(i-1, j) - T(i-1, j-1), the recursion used for the triangle in A118800 and used in this OEIS sequence for the case t = 1 and m = 1. Paul Young (see link) gives a p-adic proof of the lack of zeros among negative indices for certain Generalized Fibonacci sequences. The study presented in this OEIS sequence arose while attempting to find an inductive proof of this result which however failed. However, Hendel made the following conjecture in the problem session at West Coast Number Theory (see link). Consistent with the observations made in this OEIS sequence, when the recursion (after the initial row of length k-1) is arranged in blocks of k, then the first k-1 rows are the triangle entries right-padded with ones. It is easy to prove that these rows (for k >= 4) have no zeros. Consistent with the results in this OEIS sequence, one can show that G(k^2-1) = K(-(k^2-k-1)) = T(k-1, k). Hendel conjectured that the absolute value of G(k^2-1) or T(-(k^2-k-1)) is a minimum over the rest of the sequence, that is, |G(k^2-1)| < G(n), n >= k^2. Although it is stated on the WCNT conference that some headway had been made, these attempts failed (so the conjecture is still open). The reason for reformulating this OEIS sequence using initial conditions of mostly zeros instead of ones is that the triangle boundaries also satisfy the triangle recursion and the right-padding of zeros is more natural than the right-padding of ones.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Russell Jay Hendel_, Feb 17 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014613 | OEIS-A332664

a(n) = number of nonnegative integers that are not the sum of {2 squares, a nonnegative 5th power, and a nonnegative n-th power}.

a(2) = 0 by a theorem of Zhi-Wei Sun, see A273915. All terms beyond a(2) are conjectures and have only been checked to 4*10^9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _XU Pingya_, Feb 18 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014614 | OEIS-A332665

a(n) = largest number k that is not the sum of {2 squares, a nonnegative 5th power, and a nonnegative n-th power}.

All terms are conjectural and have only been checked up to 4*10^9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _XU Pingya_, Feb 19 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014615 | OEIS-A332710

Maximum value in n-th row of A332709.

Conjecture: the corresponding sequence for the minimum value in n-th row of A332709 is A258664.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Feb 20 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014616 | OEIS-A332803

Lexicographically earliest sequence of distinct terms > 0 such that the terms' cumulative sum and the sequence itself have the same digit succession.

This sequence is conjectured to be a permutation of the integers > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Feb 25 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014617 | OEIS-A332843

Lexicographically earliest sequence of positive terms such that a(4n+1) is the sum of the next three terms, those three terms having the property that each of them is a substring of a(4n+1).

The sequence is infinite as one can always multiply a(4n+1) by 10 and do the same with the next three terms. It is conjectured that at least two of those three terms must be equal.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Tom Duff_, Feb 26 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014618 | OEIS-A332847

a(n) is the smallest k such that exactly one of k*2^(2^n) - 2*k + 1 and k*2^(2^n) + 2*k - 1 is a prime.

Conjecture: a(n) > 1 for all n > 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Feb 26 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014619 | OEIS-A332872

Number of ordered set partitions of {1..n} where no element of any block is greater than any element of a non-adjacent consecutive block.

Conjectured to be the same as A007052, shifted right once.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Mar 06 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014620 | OEIS-A332887

a(n) is the number of partition numbers > 1 that are proper divisors of the n-th partition number.

Conjecture: every nonnegative integer occurs infinitely many times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 29 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014621 | OEIS-A332918

Average number of binary strings of length n with Levenshtein distance <= 3 from a uniform randomly sampled binary string of this length, rounded to nearest integer.

Conjectures from _Colin Barker_, Mar 05 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Mar 05 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014622 | OEIS-A332937

a(n) is the greatest common divisor of the first two terms of row n of the Wythoff array (A035513).

a(n) is also the gcd of every pair of consecutive terms of row n of the Wythoff array. Conjectures: the maximal number of consecutive 1's is 5, and the limiting proportion of 1's exists. See A332938.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014623 | OEIS-A332938

Indices of the primitive rows of the Wythoff array (A035513); see Comments.

In a row of the Wythoff array, either every two consecutive terms are relatively prime or else no two consecutive terms are relatively prime. In the first case, we call the row primitive; otherwise, the row is an integer multiple of a tail of a preceding row. Conjectures: the maximal number of consecutive primitive rows is 5, and the limiting proportion of primitive rows exists and is approximately 0.608.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014624 | OEIS-A332954

Triangle read by rows: T(n,k) is the number of permutations sigma of [n] such that sigma(j)/(j+k) > sigma(j+1)/(j+k+1) for 1 <= j <= n-1.

Conjecture: T(2*n+4,n) = A052955(n+2). This is true for n <= 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Mar 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014625 | OEIS-A332956

Symmetrical Zumkeller numbers: numbers s whose divisors can be partitioned into two disjoint sets with equal sum and whose non-divisors less than s can also be partitioned in such a way.

Conjecture 1: Except for 6, the perfect numbers (A000396) are in the sequence. Conjecture 2: At least one out of every twenty consecutive natural numbers is such a number. Conjecture 3: At least one out of every four consecutive Zumkeller numbers is such a number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ivan N. Ianakiev_, Mar 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014626 | OEIS-A332989

a(n) is the smallest number writable in n different ways as the sum of two distinct nonzero pentagonal numbers.

I conjecture this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_, Mar 05 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014627 | OEIS-A333042

G.f.: exp(Sum_{k>=1} (4*k)!/k!^4 * x^k/k).

Let A(x) denote the o.g.f. of the sequence. The sequence defined by b(n) := [x^n] A(x)^n for n >= 1 begins [24, 3672, 703968, 149835864, 33911355024, 7993981771488, 1940145241321920, ...]. We conjecture that b(n) satisfies the supercongruences b(n*p^r) == b(n*p^(r-1)) (mod p^(3*r)) for prime p >= 5 and all positive integers n and r. More generally, for a positive integer m, set A_m(x) = exp(Sum_{n >= 1} (m*n)!/(n!^m) * x^n/n) and define a sequence {b_m(n): n >= 1} by b_m(n) := [x^n] A_m(x)^n. Then we conjecture that b_m(n) is an integer sequence satisfying the same supercongruences. (End) Adamczewski includes proof abstract regarding the conjecture immediately above in linked article. - _Michael De Vlieger_, Aug 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Mar 06 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014628 | OEIS-A333086

Array read by antidiagonals: row n consists of the primes in row n of the array A333028.

The array shows, in order, the primes in the Wythoff array after deletion of all nonprimes. Every prime occurs exactly once; that is, every prime is uniquely expressible as F(k+1)*floor(n*phi) + (n-1)F(k), where phi is the golden ratio (A001622), F = A000045 (Fibonacci numbers), and n and k are positive integers. We assume as true the conjecture that each row is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 10 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014629 | OEIS-A333090

a(n) is equal to the n-th order Taylor polynomial (centered at 0) of S(x)^n evaluated at x = 1, where S(x) = (1 - x - sqrt(1 - 6*x + x^2))/(2*x) is the o.g.f. of the Schröder numbers A006318.

We conjecture that the sequence satisfies the stronger supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for all primes p >= 5 and positive integers n and k. Examples of these congruences are given below. More generally, for each integer m, we conjecture that the sequence {a_m(n) : n >= 0}, defined by setting a_m(n) = the n-th order Taylor polynomial of S(x)^(m*n) evaluated at x = 1, satisfies the same supercongruences. See A333091 for m = 2 and A333092 for m = 3. For similarly defined sequences see A333093 through A333097.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 22 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014630 | OEIS-A333091

a(n) is the n-th order Taylor polynomial (centered at 0) of S(x)^(2*n) evaluated at x = 1, where S(x) = (1 - x - sqrt(1 - 6*x + x^2))/(2*x) is the o.g.f. of the Schröder numbers A006318.

We conjecture that the sequence satisfies the stronger supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k. Examples of these congruences are given below. More generally, for each integer m, we conjecture that the sequence a_m(n),

defined as the n-th order Taylor polynomial of S(x)^(m*n) evaluated at x = 1 satisfies the same congruences. For cases see A333090 (m = 1) and A333092 (m = 3). For similarly defined sequences see A333093 through A333097.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 22 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014631 | OEIS-A333092

a(n) is the n-th order Taylor polynomial (centered at 0) of S(x)^(3*n) evaluated at x = 1, where S(x) = (1 - x - sqrt(1 - 6*x + x^2))/(2*x) is the o.g.f. of the Schröder numbers A006318.

We conjecture that the sequence satisfies the stronger supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k. Examples of these congruences are given below. More generally, for each integer m, we conjecture that the sequence a_m(n) defined as the n-th order Taylor polynomial of S(x)^(m*n) evaluated at x = 1 satisfies the same congruences. For cases, see A333090 (m = 1) and A333091 (m = 2). For similarly defined sequences see A333093 through A333097.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 22 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014632 | OEIS-A333093

a(n) is equal to the n-th order Taylor polynomial (centered at 0) of c(x)^n evaluated at x = 1, where c(x) = (1 - sqrt(1 - 4*x))/(2*x) is the o.g.f. of the Catalan numbers A000108.

We conjecture that the sequence satisfies the stronger supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k. Examples of these congruences are given below. More generally, for each integer m, we conjecture that the sequence

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014633 | OEIS-A333094

a(n) is the n-th order Taylor polynomial (centered at 0) of c(x)^(2*n) evaluated at x = 1, where c(x) = (1 - sqrt(1 - 4*x))/(2*x) is the o.g.f. of the Catalan numbers A000108.

We conjecture that the sequence satisfies the stronger supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k. Examples of these congruences are given below. More generally, for each integer m, we conjecture that the sequence a_m(n) defined as the n-th order Taylor polynomial of c(x)^(m*n) evaluated at x = 1 satisfies the same supercongruences. For cases A099837(m = -2), A100219 (m = -1), A000012 (m = 0), A333093 (m = 1), A333095 (m = 3), A333096 (m = 4), A333097 (m = 5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014634 | OEIS-A333095

a(n) = the n-th order Taylor polynomial (centered at 0) of c(x)^(3*n) evaluated at x = 1, where c(x) = (1 - sqrt(1 - 4*x))/(2*x) is the o.g.f. of the sequence of Catalan numbers A000108.

We conjecture that the sequence satisfies the stronger supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k. Examples of these congruences are given below. More generally, for each integer m, we conjecture that the sequence a_m(n) := the n-th order Taylor polynomial of c(x)^(m*n) evaluated at x = 1 satisfies the same supercongruences. For cases see A099837 (m = -2), A100219 (m = -1), A000012 (m = 0), A333093 (m = 1), A333094 (m = 2), A333096 (m = 4), A333097 (m = 5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014635 | OEIS-A333096

a(n) = the n-th order Taylor polynomial (centered at 0) of c(x)^(4*n) evaluated at x = 1, where c(x) = (1 - sqrt(1 - 4*x))/(2*x) is the o.g.f. of the sequence of Catalan numbers A000108.

We conjecture that the sequence satisfies the stronger supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k. Examples of these congruences are given below. More generally, for each integer m, we conjecture that the sequence a_m(n) := the n-th order Taylor polynomial of c(x)^(m*n) evaluated at x = 1 satisfies the same supercongruences. For cases see A099837 (m = -2), A100219 (m = -1), A000012 (m = 0), A333093 (m = 1), A333094 (m = 2), A333095 (m = 3), A333097 (m = 5). Adamczewski includes proof abstract regarding the above conjecture in linked article. - _Michael De Vlieger_, Aug 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014636 | OEIS-A333097

a(n) = the n-th order Taylor polynomial (centered at 0) of c(x)^(5*n) evaluated at x = 1, where c(x) = (1 - sqrt(1 - 4*x))/(2*x) is the o.g.f. of the sequence of Catalan numbers A000108.

We conjecture that the sequence satisfies the stronger supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k. Examples of these congruences are given below. More generally, for each integer m, we conjecture that the sequence a_m(n) := the n-th order Taylor polynomial of c(x)^(m*n) evaluated at x = 1 satisfies the same supercongruences. For cases see A099837 (m = -2), A100219 (m = -1), A000012 (m = 0), A333093 (m = 1), A333094 (m = 2), A333095 (m = 3), A333096 (m = 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014637 | OEIS-A333111

a(n) is the least positive integer not used earlier such that n+1 and the sum of terms up to a(n) are coprime.

Conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ali Sada_ and _M. F. Hasler_, Mar 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014638 | OEIS-A333122

Numbers m such that m = prime(k) + prime(k+5) = prime(k+1) + prime(k+4) for some k.

Conjecture: this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marcin Barylski_, Mar 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014639 | OEIS-A333197

Nonprime numbers k such that each nonprime divisor of k is 1 away from a prime number.

Conjecture: except for a(n) = 4, 8, 16 and 32, a(n) is of the form 2^i*p^j with p = 3, 5, 7, 11, 19, 23, 29, 31, ... ({A120628} minus {2}).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Mar 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014640 | OEIS-A333261

Numbers k such that A071324(k) = A071324(k+1).

Conjecture: There are infinitely many terms of this sequence. (Conjecture 15 in Jaiswal.) (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Mar 13 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014641 | OEIS-A333299

Number of canonical sequences of moves of length n for the Rubik's cube puzzle using the quarter-turn metric.

Conjectures from _Colin Barker_, Mar 23 2020: (Start) The above conjectured formulas are true. - _Ben Whitmore_, Dec 30 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 23 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014642 | OEIS-A333308

Numbers that are the sum of two distinct terms of A003622 (1st column of the Wythoff array, A035513).

Conjecture: the difference sequences of this sequence and its complement consist exclusively of Fibonacci numbers (A000045).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 01 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014643 | OEIS-A333331

Number of integer points in the convex hull in R^n of parking functions of length n.

It is observed by _Gus Wiseman_ in A368596 and A368730 that this sequence appears to be the complement of those sequences. If this is the case, then a(n) is the number of labeled graphs with loops allowed in which each connected component has an equal number of vertices and edges and the conjectured formula holds. Terms for n >= 9 are expected to be 167341283, 4191140394, 116425416531, ... - _Andrew Howroyd_, Jan 10 2024 An equivalent conjecture is that a(n) is the number of loop-graphs with n vertices and n edges such that it is possible to choose a different vertex from each edge. I call these graphs choosable. For example, the a(3) = 17 choosable loop-graphs are the following (loops shown as singletons): Conjectured e.g.f.: exp(-log(1-T(x))/2 + T(x)/2 - T(x)^2/4) where T(x) = -LambertW(-x) is the e.g.f. of A000169. - _Andrew Howroyd_, Jan 10 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard Stanley_, Mar 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014644 | OEIS-A333363

Horizontal visibility sequence at the onset of chaos in the 3-period cascade.

Conjectured: a(n) = 2*A007814(n/3) + 5 if 3|n and a(n) = 4 - (n mod 3) otherwise. - _Giovanni Resta_, Mar 16 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Francisco J. Muñoz_ and _Juan Carlos Nuño_, Mar 16 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014645 | OEIS-A333398

Partial sums of A333400.

Conjecture: This is a permutation of the integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alec Jones_, Mar 18 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014646 | OEIS-A333400

Lexicographically earliest infinite sequence of distinct integers whose partial sums are all distinct integers.

Conjecture: This is a permutation of the integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alec Jones_, Mar 18 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014647 | OEIS-A333406

Lexicographically earliest sequence of positive distinct integers such that if a(n) climbs over a(n) commas to the right, it is met by a term whose leftmost digit is nonprime.

This is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Carole Dubois_, Mar 19 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014648 | OEIS-A333456

a(n) is the smallest number with exactly n divisors that are Niven numbers.

Conjecture: For every n there is at least one number k with n divisors Niven numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius A. Burtea_, May 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014649 | OEIS-A333457

a(n) is the smallest number with exactly n divisors that are Moran numbers, or -1 if no such number exists.

Conjecture: For every n there is at least one number k with n divisors Moran numbers. Conjecture: The terms are divisible by 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius A. Burtea_, May 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014650 | OEIS-A333472

a(n) = [x^n] (c (x/(1 + x)))^n, where c(x) = (1 - sqrt(1 - 4*x))/(2*x) is the o.g.f. of the Catalan numbers A000108.

Here we take F(x) = c(x/(1 + x)) = 1 + x + x^2 + 2*x^3 + 4*x^4 + 9*x^5 + 21*x^6 + ... (cf. A001006 and A086246) and conjecture that the associated sequence a(n) = [x^n] (c(x/(1 + x)))^n satisfies the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(2*k)) for prime p >= 5 and positive integers n and k. Cf. A333473. More generally, we conjecture that for any positive integer r and any integer s the sequence a(r,s;n) := [x^(r*n)] (c(x/(1 + x)))^(s*n) also satisfies the above congruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 23 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014651 | OEIS-A333473

a(n) = [x^n] (S(x/(1 + x)))^n, where S(x) = (1 - x - sqrt(1 - 6*x + x^2))/(2*x) is the o.g.f. of the large Schröder numbers A006318.

Here we take F(x) = S(x/(1 + x)) = 1 + 2*x + 4*x^2 + 12*x^3 + 40*x^4 + 154*x^5 + 544*x^6 + ...(see A025227), where S(x) is the o.g.f. of the large Schröder numbers A006318. We conjecture that the associated sequence a(n) = [x^n] (S(x/(1 + x)))^n satisfies the congruences a(n*p^k) == a(n*p^(k-1)) (mod p^(2*k)) for prime p >= 5 and positive integers n and k. Cf. A333472. More generally, we conjecture that for a positive integer r and integer s, the sequence a(r,s;n) := [x^(r*n)] (S(x/(1 + x)))^(s*n) also satisfies the above congruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 23 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014652 | OEIS-A333481

a(n) = [x^n] S(x)^(2*n), where S(x) = (1 - x - sqrt(1 - 6*x + x^2))/(2*x) is the o.g.f. of the large Schröder numbers A006318.

We conjecture that the stronger supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) hold for prime p >= 5 and positive integers n and k. More generally, we conjecture that for any positive integer a and any integer b the sequence u(a,b;n) := [x^(a*n)] S(x)^(b*n) also satisfies the above congruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014653 | OEIS-A333510

Number of self-avoiding walks in the n X 2 grid graph which start at any of the n vertices on left side of the graph and terminate at any of the n vertices on the right side.

Conjecture: a(n) = (27*2^n - n^3 - 26*n - 24)/3. Conjecture: G.f.: x*(1+2*x-5*x^2+2*x^3+2*x^4)/((1-x)^4*(1-2*x)). Conjectures confirmed using the transfer matrix method. - _Andrew Howroyd_, Oct 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Mar 25 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014654 | OEIS-A333512

Numbers nontrivially palindromic in exactly three consecutive number bases.

Numbers which are strictly non-palindromic up to a set of k=3 consecutive number bases. It is conjectured that such a sequence for k>=4 is empty. For a special case of k=0, we have the sequence A016038.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matej Veselovac_, Mar 25 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014655 | OEIS-A333515

Number of self-avoiding closed paths on an n X 5 grid which pass through four corners ((0,0), (0,4), (n-1,4), (n-1,0)).

The order-11 linear and the g.f. by _Chai Wah Wu_ are correct, for all n. Proof sketch: sweeping the n X 5 grid one row at a time, the partial cycle meets the horizontal cut between consecutive rows in a non-crossing perfect matching of the 5 active stubs; tracking this boundary-connectivity state (with union-find enforcing degree <= 2, degree exactly 2 at the four corners, no premature loop closure, no dangling end, and a single component at the final CLOSE) gives an exact integer transfer matrix M on N = 20 reachable states with a(n) = cI^T M^(n-2) v1 for n >= 2. The minimal polynomial of v1 under M is x^11 - 13*x^10 + 45*x^9 - 66*x^8 + 17*x^7 + 209*x^6 - 151*x^5 - 140*x^4 + 112*x^3 + 48*x^2 - 50*x - 28, whose annihilation q(M)v1 = 0 yields the recurrence for every n in its domain (n >= 13; i.e., n > 12); its reciprocal is exactly the conjectured denominator, and the numerator forced by a(2..12) is exactly the conjectured numerator. - _Yiming Beckmann_, Jun 06 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Mar 25 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014656 | OEIS-A333560

Square array read by antidiagonals: T(n,k) = Sum_{j = 0..n*k} binomial(n+j-1,j)*2^j; n,k >= 0.

We conjecture that each column sequence satisfies the following supercongruences: Conjectural o.g.f. for column k: 2^(k+1)*x*f'(k, (2^k)*x)/(2*f(k, (2^k)*x) - 1) + 1/(1 + x), where f(k, x) = Sum_{n >= 0} 1/((k+1)*n+1)*binomial((k+1)*n+1, n) * x^n. The conjectural o.g.f. for column k is true for all k >= 1; equivalently, exp(Sum_{n>=1} T(n,k)*x^n/n) = (2*f(k, 2^k*x) - 1)/(1 + x). - _Samuel Lamphier_, Oct 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 26 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014657 | OEIS-A333561

a(n) = Sum_{j = 0..2*n} binomial(n+j-1,j)*2^j.

We conjecture that this sequence satisfies the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k. Some examples are given below. Conjectural o.g.f.: 1/(1 + x) + 8*x*f'(4*x)/(2*f(4*x) - 1), where f(x) = 1 + x + 3*x^2 + 12*x^3 + 55*x^4 + ... is the o.g.f. of A001764. Conjectural recurrence: n*(n - 1)*(2*n - 1)*(3098*n - 6455)*a(n) = (n - 1)*(172988*n^3 - 585840*n^2 + 550321*n - 169824)*a(n-1) - 12*(11825*n^4 - 168518*n^3 + 627675*n^2 - 853766*n + 350744)*a(n-2) - 36*(n - 3)*(3*n - 7)*(3*n - 8)*(991*n - 724)*a(n-3) with a(1) = 7, a(2) = 129, a(3) = 2815.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014658 | OEIS-A333562

a(n) = Sum_{j = 0..3*n} binomial(n+j-1,j)*2^j.

We conjecture that this sequence satisfies the congruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k. Some examples are given below. Conjectural o.g.f.: 1/(1 + x) + 16*x*f'(8*x)/(2*f(8*x) - 1), where f(x) = 1 + x + 4*x^2 + 22*x^3 + 140*x^4 + ... is the o.g.f. of A002293.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014659 | OEIS-A333563

a(n) = [x^n] G(x)^n, where G(x) is the o.g.f. of A079489.

The o.g.f. G(x) of A079489 is given by G(x) = c(sqrt(x))*c(-sqrt(x)), where c(x) = ((1 - sqrt(1 - 4*x))/(2*x)) is the o.g.f. of the Catalan numbers A000108. It is known that the sequence b(n) := [x^n] c(x)^n = 1/3*binomial(3*n,n) satisfies the supercongruences b(n*p^k) == b(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k - see Meštrović, equation 39. We conjecture that the present sequence satisfies the same congruences. Some examples are given below. [We note that the sequence u(n) := n^2 * Sum_{k = 0..2*n} 1/((n+2*k)*(5*n-2*k))*C(n+2*k,k)*C(5*n-2*k, 2*n-k) = (1/3)*C(6*n,2*n) is known to satisfy the congruences u(n*p^k) == u(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k - see Meštrović, equation 39. If, in the binomial sum formulas for a(n) and u(n) given above, we restrict the summation range to k = 0..n then we conjecture that the resulting pair of sequences satisfy the same congruences.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 12 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014660 | OEIS-A333564

a(n) = [x^n] (c(x)/c(-x))^n, where c(x) = (1 - sqrt(1 - 4*x))/(2*x) is the o.g.f. of the Catalan numbers A000108.

It can be shown that a(n) satisfies the Gauss congruences a(n*p^k) == a(n*p^(k-1)) (mod p^k) for all prime p. We conjecture that a(n) satisfies the stronger congruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 3 and positive integers n and k. Some examples are given below. More generally, for integer r and positive integer s, we conjecture that the sequence a(r,s;n) := 2^(r*n) * Sum_{k = 0..s*n-1} (-1)^(s*n-1+k)*(binomial(n+k,k))^r satisfies the same congruences. This is the case r = s = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014661 | OEIS-A333565

O.g.f.: (1 + 4*x)/((1 + x)*sqrt(1 - 8*x)).

We conjecture that this sequence satisfies the stronger congruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 3 and positive integers n and k. The particular case when n = k = 1 follows from the corresponding result for A333564. Some examples of these congruences are given below.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014662 | OEIS-A333574

Number of Hamiltonian paths in the n X 2 grid graph which start at any of the n vertices on left side of the graph and terminate at any of the n vertices on the right side.

Conjecture: Numbers k such that A339399(k) = A103128(k). - _Wesley Ivan Hurt_, Nov 19 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Mar 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014663 | OEIS-A333579

a(n) = [x^n] ((1 + x + x^2)/(1 - x + x^2))^n.

We conjecture that the present sequence satisfies the congruences a(p) == 2 (mod p^3) for prime p >= 5 (checked up to p = 499). Calculation suggests that a(p^k) == a(p^(k-1)) (mod p^(2*k)) for prime p >= 5 and k > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 29 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014664 | OEIS-A333592

a(n) = Sum_{k = 0..n} binomial(n+k-1, k)^2.

The sequence b(n) = C(2*n,n) of central binomial coefficients satisfies the supercongruences b(n*p^k) = b(n*p^(k-1)) (mod p^(3*k)) for all primes p >= 5 and positive integers n and k - see Meštrović. We conjecture that the present sequence also satisfies these congruences. Some examples of the congruences are given below.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014665 | OEIS-A333593

a(n) = Sum_{k = 0..n} (-1)^(n + k)*binomial(n + k - 1, k)^2.

The sequence b(n) = C(2*n,n) of central binomial coefficients satisfies the supercongruences b(n*p^k) = b(n*p^(k-1)) (mod p^(3*k)) for all prime p >= 5 and any positive integers n and k - see Mestrovic. We conjecture that the present sequence also satisfies these congruences. Some examples of the congruences are given below. For a

proof of the particular case of these congruences, a(p) == 0 (mod p^3) for prime p >= 5, see the Bala link. The sequence a(p)/(2*p^3) for prime p >= 5 begins [48, 304, 36519504, 4827806144, 109213719151680, 17975321574419440, ...]. Cf. A034602.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014666 | OEIS-A333603

Number of directed Hamiltonian walks from NW to SW corners of a 7 X (2*n+1) grid.

Conjecture: a(n)= 85*a(n-1) -1932*a(n-2) +20403*a(n-3) -116734*a(n-4) +386724*a(n-5) -815141*a(n-6) +1251439*a(n-7) -1690670*a(n-8) +2681994*a(n-9) -4008954*a(n-10) +3390877*a(n-11) -1036420*a(n-12) -178842*a(n-13) +92790*a(n-14) +17732*a(n-15) -5972*a(n-16) +1728*a(n-17) +144*a(n-18). - _R. J. Mathar_, Mar 13 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Mar 28 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014667 | OEIS-A333637

The number of cells which contain multiple squares of a Genealodron formed from 2^n - 1 equal-sized squares (when viewed from above).

Conjecture: for n>=6, a(n) = n^2 - n - 15. - _Vaclav Kotesovec_, Apr 07 2020 Conjectures from _Colin Barker_, Apr 07 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Smith_, Mar 30 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014668 | OEIS-A333650

Triangle read by rows: T(n,k) gives the number of domino towers of height k consisting of n bricks.

Conjecture: For n > 1, T(n,2) = A000071(n+2). Conjecture: T(n,n-1) = A339252(n-2). - _Peter Kagey_, Nov 21 2020 Conjecture: T(n,n-2) = A339254(n-3). - _Peter Luschny_, Nov 29 2020 Conjecture: T(n,n-3) = A339029(n-4). - _Peter Luschny_, Dec 01 2020 The above conjectures can be summarized as follows:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Mar 31 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014669 | OEIS-A333676

Least positive integer k such that k*prime(n) is an Ulam number (A002858).

Conjecture: For any prime number p there exists a positive integer k such that k*p is an Ulam number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Apr 01 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014670 | OEIS-A333686

Number of self-avoiding paths in (2*n+1) X 3 grid starting the upper left corner, passing through the center of grid and finishing the lower right corner.

Conjecture: G.f.: (1-3*x-x^2)*(1+3*x+x^2+x^3)/((1-x)*(1+x)*(1+x+x^2)*(1-11*x+x^2)). Conjecture: a(n) = 10*a(n-1) + 10*a(n-2) - 10*a(n-4) - 10*a(n-5) + a(n-6) for n>5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Apr 02 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014671 | OEIS-A333715

a(n) = [x^(3*n)] ((1 + x)/(1 - x))^n.

We conjecture that a(n) satisfies the stronger supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k. Some examples of these congruences are given below. For fixed m = 1,2,3,..., we conjecture that the sequence b(n) := a(m*n) satisfies a recurrence of the form P(6*m,n)*b(n+1) + P(6*m,-n)*b(n-1) = Q(3*m,n^2)*b(n), where the polynomials P(6*m,n) and Q(3*m,n^2) have degree 6*m. Conjecturally, the 6*m zeros of the polynomial Q(3*m,n^2) belong to the interval [-1, 1] and 6*m - 4 of these zeros appear to be approximated by the rational numbers +- k/(4*m), where 1 <= k <= 4*m - 3, k not a multiple of 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014672 | OEIS-A333794

a(1) = 1, for n > 1, a(n) = n + a(n-A052126(n)).

Conjecturally, also the largest path sum when iterating from n to 1 with nondeterministic map k -> k - k/p, where p is any prime factor of k. a(n) = n + Max a(n - n/p), for p prime and dividing n. [Conjectured, holds at least up to n=2^24]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Apr 05 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014673 | OEIS-A333820

a(n) is the number of pairs (q,r) such that q <= r and phi(q) + phi(r) = 2*n.

Paul Erdős and Leo Moser conjectured that, for any even numbers 2*n, there exist integers q and r such that phi(q) + phi(r) = 2*n. Therefore, they conjecture a(n) > 0 for all ns.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Apr 06 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014674 | OEIS-A333828

The 20-adic integer x = ...70D9AE7F1DI8 satisfying x^5 = x.

Conjecture: There exists a nontrivial n-adic integer x satisfying x^5 = x, and x^2, x^3, and x^4 are not x, if and only if n has a prime factor of the form 4k+1. Further, there is one nontrivial pair (x and -x) for each different prime factor of the form 4k+1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick A. Thomas_, Apr 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014675 | OEIS-A333852

Irregular triangle read by rows: representative simple difference sets of Singer type of order m, for m = A000961(n), for n >= 1.

A representative difference set of this type is one which uses 0 and 1 as elements. By adding each element of a given representative difference set by 1, 2, ..., (m+1)*m, modulo m^2 + m + 1, all m^2 + m + 1 members of a class of difference sets are obtained. This equivalence class is called Dev(D), the development of a difference set D. The

number of representative difference sets, hence the number of classes, has been conjectured by Singer, and is given in A335866(n). The difference sets will here be ordered increasingly. The set of all A335866(n) representative difference sets of this type will here be denoted by Sr(m) (also increasingly ordered). See the W. Lang link for these representatives of order m = m(n) = A000961(n), for n = 1, 2, ..., 11. The set of all difference sets of order m will be denoted by S(m).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jul 26 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014676 | OEIS-A333874

Numbers k such that A173557(k) = A173557(k+1).

Kim et al. (2019) conjectured that A173557(k) = A173557(k+1) is divisible by 12 for all the terms k > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Apr 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014677 | OEIS-A333897

Number of unlabeled loopless multigraphs with 4 nodes of degree n or less.

Conjectures from _Colin Barker_, Apr 10 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Howroyd_, Apr 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014678 | OEIS-A333905

Lexicographically earliest sequence of distinct positive integers such that a(n) divides the concatenation of a(n+1) to a(n+2).

Conjectures from _Colin Barker_, Apr 09 2020: (Start) Conjecture: a(n) = 2^((n-7)/2)*(5 + 2*sqrt(2) + (2*sqrt(2) - 5)*(-1)^n) for n > 6. - _Stefano Spezia_, Oct 23 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_, Apr 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014679 | OEIS-A333913

Numbers k such that lambda(k) is not the sum of 3 squares, where lambda is the Carmichael lambda function (A002322).

Pollack (2011) proved that this sequence has a lower and an upper asymptotic densities, and conjectured that they do not coincide.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Apr 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014680 | OEIS-A334001

The number of odd values in the n-th column of A279212.

Conjecture: a(n) <= 2*n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Apr 12 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014681 | OEIS-A334052

Variant of Van Eck's sequence: Set a(1) = 1, a(2) = 0, a(3) = 1. Thereafter, a(n) is the minimum positive offset i from a(n-1) such that some partial sum a(n-1-i)+...+a(n-1-i+k) = a(n-1) (0 <= k < i).

Conjecture: this sequence is defined. That is, each a(n-1) appears as a partial sum starting from some offset.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ethan Goldberg_, Sep 06 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014682 | OEIS-A334086

Positive numbers not of the form 2*x^4 + y*(y+1)/2 + z*(z+1)/2 with x,y,z nonnegative integers.

Conjecture: The sequence has totally 216 terms as listed in the b-file. As none of the 216 terms in the b-file is divisible by 3, the conjecture implies that for each nonnegative integer n we can write 3*n as 2*x^4 + y*(y+1)/2 + z*(z+1)/2 and hence 12*n+1 = 8*x^4 + (y+z+1)^2 + (y-z)^2, where x,y,z are integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 14 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014683 | OEIS-A334110

The squares of squarefree numbers (A062503), ordered lexicographically according to their prime factors. a(n) = Product_{k in I} prime(k+1)^2, where I are the indices of nonzero binary digits in n = Sum_{k in I} 2^k.

Conjecture: a(n) is the position of the first occurrence of n in A334109.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_ and _Peter Munn_, May 01 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014684 | OEIS-A334113

Positive numbers not of the form 4*x^4 + y*(y+1)/2 + z*(z+1)/2, where x,y,z are nonnegative integers.

Conjecture: The sequence only has 602 terms as listed in the b-file.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 14 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014685 | OEIS-A334125

Number of subsets of {1, 3, ..., 2*n-1} which sum to 0 modulo 2*n-1.

Conjecture: a(n) = 2*abs(A178738(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jinyuan Wang_, Apr 30 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014686 | OEIS-A334126

a(n) is the smallest positive integer not already in the sequence such that Sum_{i=k..n} a(i) is not a perfect power for 0 < k < n; start with a(1)=1.

I conjecture that every number eventually appears.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jinyuan Wang_, May 10 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014687 | OEIS-A334133

Numbers k > 2 such that gpf(A111076(k)^lambda(k) - 1) = gpf(lambda(k) + 1); where gpf is the greatest prime factor (A006530), and lambda = A002322 is the Carmichael function.

Conjecture: all odd terms {3, 5, 9, 13, 15, 21, 35, 39, 45, 63, 65, 91, 105, 117, 195, 273, 315, 455, 585, 819, 1365, 4095} are odd numbers k such that gpf(2^m-1) = gpf(m+1), where m = A002326((k-1)/2) is the multiplicative order of 2 mod 2k+1. - _Amiram Eldar_, Apr 15 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Apr 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014688 | OEIS-A334138

Number of ways to write n as x^4 + y*(2*y+1) + z*(3*z+1), where x is a nonnegative integer, and y and z are integers.

Conjecture 1: a(n) = 0 only for n = 64. In other words, any nonnegative integer n not equal to 64 can be written as x^4 + y*(2*y+1) + z*(3*z+1) with x,y,z integers. Conjecture 2: (i) The set {x^4+y^2+z(3z+1)/2: x,y,z are integers} contains all nonnegative integers except for 455. We have verified Conjecture 1 for n up to 10^8, parts (i) and (iii) of Conjecture 2 for n up to 5*10^7, and parts (ii), (iv) and (v) of Conjecture 2 for n up to 2*10^6. See also A334147 for the list of those numbers n with a(n) = 1. - _Zhi-Wei Sun_, Apr 16 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014689 | OEIS-A334147

Numbers which can be written uniquely as x^4 + y*(2y+1) + z*(3z+1), where x,y,z are integers with x>=0.

Conjecture: The sequence has no terms greater than a(37) = 1277405.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 16 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014690 | OEIS-A334164

a(n) is the number of ON-cells in the completed n-th level of a triangular wedge in the hexagonal grid of A151723 (i.e., after 2^k >= n generations of the automaton in A151723 have been computed).

Conjecture 1: Except for a(2^n + 1) = 2, n >= 1, for odd-numbered completed levels a lower bound of the ratio of ON-cells to the length of the level is (2^n + 2)/(3*2^(n+1) + 1) with limit 1/6, determined by the subsequence of levels starting with: 13, 25, 49, 97, 193, 385, 769, 1537, 3073, ..., and the associated ON-cell counts: 4, 6, 10, 18, 34, 66, 130, 258, 514, ..., as listed in the second column of each of the two triangles below. Conjecture 2: In an even-numbered completed level 2*n the fraction of ON-cells is bounded below by (23 * 2^n - 24)/(2^(n+5) - 36) with limit 23/32, determined by the subsequence of levels starting with: 28, 92, 220, 476, 988, 2012, 4060, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Apr 17 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014691 | OEIS-A334166

Numbers k having a divisor d, such that d*k is a Zumkeller number (A083207).

Conjecture: The difference between two consecutive terms is 6 at most.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ivan N. Ianakiev_, Apr 17 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014692 | OEIS-A334168

Numbers m whose leading digit and the other decimal digits appear respectively before and directly after the decimal point of its m-th root.

The last nonzero digits of the terms are conjectured to be the Beatty sequence for log(10), A059545. a(n+1) = floor((10^(n/10^n))*10^n) (conjectured). a(n) = 10^(n-1) + floor((n-1)*log(10)) (conjectured). - _David A. Corneth_, Apr 19 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alaa A. Nasr_, Apr 17 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014693 | OEIS-A334194

a(n) = n - d1*d2, where d1, d2 are the distances from n to the previous and the next prime number respectively.

So I conjecture that n > d1 * d2 for every N - {1, 2, 7, 23}. If it is true, it means that there is at least 1 prime number in the interval [n - sqrt(n), n + sqrt(n)]. I also conjecture that lim_{n -> oo} (n - d1*d2)/n = a(n)/n = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimitris Valianatos_, Apr 18 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014694 | OEIS-A334279

Irregular table read by rows: T(n, k) is the coefficient of x^k in the chromatic polynomial of the 1-skeleton of the n-dimensional cross polytope, 0 <= k <= 2n.

Conjectures from _Peter Kagey_, Apr 26 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Apr 21 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014695 | OEIS-A334339

Least positive integer m such that sigma(m * n) is a cube, where sigma(k) is the sum of the divisors of k.

Conjecture: a(n) exists for any n > 0. In other words, for any positive integer n, there is a positive integer m with sigma(m * n) equal to a cube. The author's conjecture in A259915 implies that for each n = 1, 2, 3, ... there is a positive integer m with sigma(m * n) equal to a square. See also A334337 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 23 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014696 | OEIS-A334350

Least positive integer m relatively prime to n such that phi(m*n) = phi(m)*phi(n) is a fourth power, where phi is Euler's totient function (A000010).

Conjecture: For any positive integers k and m, there is a positive integer n relatively prime to m such that phi(m*n) = phi(m)*phi(n) is a k-th power. This conjecture implies that a(n) exists for every n = 1,2,3,.... See also A334353 for a similar conjecture involving the sigma function (A000203).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014697 | OEIS-A334353

Least positive integer m relatively prime to n such that sigma(m*n) is a fourth power, where sigma(k) is the sum of the divisors of k.

Conjecture: For any positive integers k and m, there is a positive integer n relatively prime to m such that sigma(m*n) is a k-th power. See also A334350 for a similar conjecture involving Euler's totient function (A000010).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014698 | OEIS-A334386

a(n) is the number of ways to choose 3 points in a size n tetrahedral grid in such a way that the three points form an equilateral triangle that touches all four sides of the tetrahedron.

Conjecture: a(n) - 4*(n-1) is divisible by 12 for n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, May 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014699 | OEIS-A334413

First differences of A101803.

This is a non-homogenous Sturmian word, conjectured to be produced by the following recurrence relation: string S(0)=1; S(1)=10; S(n+2)=S(n+1)S(n)* for all natural n, where S(n)* denotes the reverse of S(n); a=S(infinity). Equivalently, a(n) is conjectured to be the fixed point of the morphism 0->101, 1->10101. See "A characterization of the quadratic irrationals."

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aresh Pourkavoos_, Apr 28 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014700 | OEIS-A334414

First differences of A334415.

See A334413 for conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aresh Pourkavoos_, May 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014701 | OEIS-A334501

Eventual period of a single cell in rule 190 cellular automaton in a cyclic universe of width n.

Conjectures from _Colin Barker_, May 09 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 05 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014702 | OEIS-A334502

Eventual period of a single cell in rule 62 cellular automaton in a cyclic universe of width n.

Conjectures from _Colin Barker_, May 13 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 05 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014703 | OEIS-A334503

Eventual period of a single cell in rule 131 cellular automaton in a cyclic universe of width n.

Conjectures from _Colin Barker_, May 13 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 05 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014704 | OEIS-A334509

Eventual period of a single cell in rule 41 cellular automaton in a cyclic universe of width n.

Conjectures from _Colin Barker_, May 09 2020: (Start)
Conjecture: a(n) = n*A176895(n) for n > 6. - _Stefano Spezia_, Oct 03 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 05 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014705 | OEIS-A334511

Eventual period of a single cell in rule 9 cellular automaton in a cyclic universe of width n.

Conjectures from _Colin Barker_, May 09 2020: (Start)
Conjecture: a(n) = (3 - (-1)^n)*n/2 for n > 10. - _Stefano Spezia_, Sep 11 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 05 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014706 | OEIS-A334513

Eventual period of a single cell in rule 25 cellular automaton in a cyclic universe of width n.

Conjectures from _Colin Barker_, May 09 2020: (Start)
Conjecture: for n>=10, a(n) = 2*n. - _Vaclav Kotesovec_, May 10 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 05 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014707 | OEIS-A334514

Eventual period of a single cell in rule 107 cellular automaton in a cyclic universe of width n.

Conjectures from _Colin Barker_, May 09 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 05 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014708 | OEIS-A334559

a(2n+1) = (2n)^2 - 1, a(4n+2) = (4n+2)^2 - 3, a(4n) = (4n)^2 - 5, except for a(1) = 1, a(4) = 2. (Conjectured outcome of an elimination game, see comments.)

a(2n+1) = (2n)^2 - 1, a(4n+2) = (4n+2)^2 - 3, a(4n) = (4n)^2 - 5, except for a(1) = 1, a(4) = 2. (Conjectured outcome of an elimination game, see comments.) Also (conjectured) the "winner" or largest remaining number in the following game:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ali Sada_ and _M. F. Hasler_, May 06 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014709 | OEIS-A334596

Number of values in A334556 with binary length n.

Conjectured formula:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, May 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014710 | OEIS-A334603

Period of the fraction 1/11^n for n >= 1.

Conjecture proposed by the authors in References page 205: if p is a prime with gcd(p,30) = 1 and if the period of 1/p is m then the period of 1/p^n is m*p^(n-1). a(n) = 2 * 11^(n-1) [conjectured, see comments].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, May 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014711 | OEIS-A334625

Maximal size of a subset T of S = {1,2,...,n} with a cyclic arrangement of T such that any three neighbors can be reordered in an arithmetic progression.

Conjectures from _Colin Barker_, Sep 10 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Fausto A. C. Cariboni_, Sep 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014712 | OEIS-A334701

Consider the figure made up of a row of n adjacent congruent rectangles, with diagonals of all possible rectangles drawn; a(n) = number of interior vertices where exactly two lines cross.

It would be nice to have a formula or recurrence. - _N. J. A. Sloane_, Jun 22 2020. Karlsson (see link) gives an explicit formula and establishes the conjectures made below. - _N. J. A. Sloane_, Jun 22 2026 Conjecture: As n -> oo, a(n) ~ C*n^4/Pi^2, where C is about 0.95 (compare A115004, A331761). - _N. J. A. Sloane_, Jul 03 2020 Both conjectures are correct - see Karlsson link. - _N. J. A. Sloane_, Jun 22 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, May 30 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014713 | OEIS-A334745

Starting with a(1) = a(2) = 1, proceed in a square spiral, computing each term as the sum of diagonally adjacent prior terms.

Conjecture: a(2n-1) = A247976(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alec Jones_ and _Peter Kagey_, May 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014714 | OEIS-A334763

Ceiling of circumradius of triangle whose sides are consecutive Ulam numbers (A002858).

Conjecture: Triangles whose sides are consecutive Ulam numbers are acute apart from (1,2,3), (2,3,4), (3,4,6), (4,6,8), (6,8,11) and (16,18,26).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, May 10 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014715 | OEIS-A334765

Numbers m such that the numbers of 1's in the binary expansion of m equals the negative sum of balanced ternary trits of m.

a(116) = 32770, a(117) = 32771 and a(118) = 32772 is the first time that three consecutive numbers appear in this sequence. Conjecture: There is no occurrence of four or more consecutive numbers. Tested by exhaustive search up to a(n) = 3^26. - _Thomas König_, Jul 19 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas König_, May 10 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014716 | OEIS-A334797

Primes q such that p-1 | q-1 or q-1 | p-1 for every prime p | 2^(q-1)-1.

Conjecture: the odd terms q for which 2^(q-1)-1 is squarefree are in A099596. - _Thomas Ordowski_, Jul 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Jun 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014717 | OEIS-A334811

Length of shortest overlap-free binary word having n periods.

Conjecture: a(n) = (17/6)*4^(n-3) + 2/3 for n >= 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, May 12 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014718 | OEIS-A334830

a(n) is the number of permutations of length n that have the same number of fixed points whether forwards or backwards.

Also relevant to the null hypothesis statistical test (NHST) called the matching method, used to evaluate the experimental hypothesis that two nominal or ordinal populations are mutually independent.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Cader Nelson_, Jun 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014719 | OEIS-A334838

Positive integers m with prime(m) in the form x^2 + m*y^2, where x and y are positive integers.

Conjecture: The current sequence has infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 13 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014720 | OEIS-A334856

E.g.f. A(x) satisfies: A(x) = sin(x * exp(A(x))).

Conjectural 2-adic values: v_2(a(8*n+4)) = 8*n + 4;

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, May 13 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014721 | OEIS-A334917

Indices in A334916 where records occur.

If the conjecture "A334916(n)>0 for n>3" is true, then the next term is a(12)=107.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matej Veselovac_, May 16 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014722 | OEIS-A334958

GCD of consecutive terms of the factorial times the alternating harmonic series.

Conjecture: a(n) = n! * Product_{p <= n} p^min(0, v_p(H'(n))), where the product ranges over primes p, H'(n) = Sum_{k=1..n} (-1)^(k+1)/k, and v_p(r) is the p-adic valuation of rational r (checked for n < 1100).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, May 17 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014723 | OEIS-A334981

Numbers k such that the numerator of the k-th alternating harmonic number H'(k) is divisible by the square of a prime less than k.

This sequence was inspired by unsolved conjectures related to the papers by Boyd (1994) and Krattenhaler and Rivoal (2007-2009, 2009) about the harmonic numbers H(k) = Sum_{i=1..k} 1/i. See also the comments for sequences A007757, A131657, A131658, and A268112. Here we are dealing with the alternating harmonic numbers H'(k) = Sum_{i=1..k} (-1)^(i+1)/i. If variations of the above conjecture are true, then some conditional results in Krattenhaler and Rivoal (2007-2009, 2009) would hold. Boyd (1994) found only 5 integers k such that there is a prime p < k with v_p(H(k)) >= 3. Since 1994 no other k's have been found that satisfy the latter inequality. We claim that a similar conjecture holds for the alternating harmonic numbers H'(k): there is no pair of an integer k and a prime p such that v_p(H'(k)) >= 4; i.e., there is no k for which the numerator of H'(k) is divisible by the fourth power of a prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, May 25 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014724 | OEIS-A334995

Twice the area of triangle with coordinates (Fn, Fn+k), (Fn+2k, Fn+3k) and (Fn+4k, Fn+5k), where Fn is the n-th Fibonacci number A000045.

Conjectures from _Colin Barker_, May 19 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, May 19 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014725 | OEIS-A334996

Irregular triangle read by rows: T(n, m) is the number of ways to distribute Omega(n) objects into precisely m distinct boxes, with no box empty (Omega(n) >= m).

Conjecture: Sum_{m=0..Omega(n)} T(n, m) = A002033(n-1) for n > 1. The above conjecture is true since T(n, m) is also the number of ordered factorizations of n into m factors (see Comments) and A002033(n-1) is the number of ordered factorizations of n. - _Stefano Spezia_, Aug 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, May 19 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014726 | OEIS-A335023

Ratios of consecutive terms of A334958.

Conjecture: a(n) = 1 if and only if n+1 is prime. Adamczewski gives a counterexample (1194648) to the above conjecture in linked paper. - _Michael De Vlieger_, Aug 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, May 19 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014727 | OEIS-A335024

Ratios of consecutive terms of A056612.

Conjecture 1: a(n) = 1 if and only if n + 1 = p^k for some prime p and some positive integer k < p. Conjecture 2 (due to _Alois P. Heinz_): a(n) = 1 <=> n+1 in A136327.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, May 19 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014728 | OEIS-A335038

a(n) is the smallest number m with exactly n divisors that are Zuckerman numbers, or -1 if there is no such m.

Conjecture: m always exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Jun 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014729 | OEIS-A335071

Numbers m such that the delta(m) = abs(sigma(m+1)/(m+1) - sigma(m)/(m)) is smaller than delta(k) for all k < m.

There is no disproof of existence, but this would require both of the consecutive numbers to be k-perfect with the same k >= 2, and it is conjectured that such numbers are multiples of k!. It is very unlikely that an odd k-perfect number will ever be found, and even much less probable that it will be just next to an even k-perfect number. - _M. F. Hasler_, Jun 06 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, May 22 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014730 | OEIS-A335080

First elements of maximal isospectral chains of length 1, or, equivalently, numbers with spectral basis of index 1.

Isospectral Chain Conjecture: There exist isospectral chains of any positive length.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Walter Kehowski_, May 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014731 | OEIS-A335081

First elements of maximal isospectral chains of length 2.

Isospectral Chain Conjecture: There exist isospectral chains of any positive length.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Walter Kehowski_, May 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014732 | OEIS-A335082

First elements of maximal isospectral chains of length 3.

Isospectral Chain Conjecture: There exist isospectral chains of any positive length.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Walter Kehowski_, May 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014733 | OEIS-A335083

First elements of maximal isospectral chains of length 4.

Isospectral Chain Conjecture: There exist isospectral chains of any positive length.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Walter Kehowski_, May 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014734 | OEIS-A335084

First elements of maximal isospectral chains of length 5.

Isospectral Chain Conjecture: There exist isospectral chains of any positive length.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Walter Kehowski_, May 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014735 | OEIS-A335085

First elements of maximal isospectral chains of length 6.

Isospectral Chain Conjecture: There exist isospectral chains of any positive length.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Walter Kehowski_, May 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014736 | OEIS-A335139

a(n) = (prime(n + 1) +- k) / 2 where k is the smallest possible odd number such that a(n) is prime and a(n + 1) >= a(n).

The sequence of k's begins {1, 1, -1, -1, 1, -3, 3, -1, -3, 3, 1, -3, 3, -1, ...}. I conjecture that the partial sums of the k's sequence change sign infinitely often and that their absolute value is less than the square root of n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimitris Valianatos_, May 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014737 | OEIS-A335159

a(n) = nim-product of 2^2 and 2^n.

Conjectures from _Colin Barker_, Jun 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014738 | OEIS-A335160

a(n) = nim-product of 2^3 and 2^n.

Conjectures from _Colin Barker_, Jun 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014739 | OEIS-A335161

a(n) = nim-product of 2^4 and 2^n.

Conjectures from _Colin Barker_, Jun 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014740 | OEIS-A335214

Divide the biggest term of the pair [a(n), a(n+1)] by the smallest one and keep the remainder; the successive remainders of the successive pairs rebuild the starting sequence, digit after digit. This is the lexicographically earliest sequence of distinct positive terms with this property.

This is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, May 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014741 | OEIS-A335226

Numbers m such that twice the number of unordered Goldbach partitions of 2m is less than the number of unordered Goldbach partitions of 4m.

It is conjectured that the last term in this sequence is a(114)=22564.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Craig J. Beisel_, May 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014742 | OEIS-A335250

Numbers m such that twice the number of unordered Goldbach partitions of 2m equals the number of unordered Goldbach partitions of 4m.

It is conjectured that the last term in this sequence is a(45)=11749.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Craig J. Beisel_, May 28 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014743 | OEIS-A335270

Numbers that are not powers of primes (A024619) whose harmonic mean of their proper unitary divisors is an integer.

Conjecture: all terms are of the form n*(usigma(n)-1) where usigma(n)-1 is prime. - _Chai Wah Wu_, Dec 17 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, May 29 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014744 | OEIS-A335315

Lexicographically earliest sequence of distinct positive integers such that the sum of digits of any consecutive pair of terms divides their consecutive concatenation.

Conjecture: This is a permutation of the natural numbers. The concatenation of any pair of adjacent terms is a composite number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, May 31 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014745 | OEIS-A335319

Decimal expansion of Sum_{n>=2} (-1)^n/(n*phi(n)), where phi(n) is the Euler totient function A000010.

The formula section of A000010 provides the following conjecture: Sum_{i>=2} (-1)^i/(i*phi(i)) exists and is approximately 0.558. - Orges Leka (oleka(AT)students.uni-mainz.de), Dec 23 2004 A more accurate value of the conjectured limit is provided.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, May 31 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014746 | OEIS-A335337

a(n) is the least k such that A335294(k) = n, or -1 if no such k exists.

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Jun 02 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014747 | OEIS-A335340

North-East paths from (0,0) to (n,n) with k cyclic descents.

I conjecture that the polynomial Sum_{k=1...n} T(n,k) t^k is real-rooted for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Per W. Alexandersson_, Jun 02 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014748 | OEIS-A335360

Numbers m such that the number of unordered Goldbach partitions of 2m is greater than the number of unordered Goldbach partitions of 4m.

It is conjectured that a(5)=353 is the last term in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Craig J. Beisel_, Jun 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014749 | OEIS-A335379

a(n) is the number of Mersenne prime (irreducible) polynomials M = x^k(x+1)^(n-k)+1 of degree n in GF(2)[x] (k goes from 1 to n-1) such that Phi_7(M) has an odd number of prime divisors (omega(Phi_7(M)) is odd).

It is surprising that a(n) be so small (conjecturally it is always 1 or 2). The sequence appeared when working the special case p=7 of a conjecture (see Links) about prime divisors in GF(2)[x] of the composed cyclotomic polynomial Phi_p(M), where p is a prime number and M is a Mersenne irreducible polynomial.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luis H. Gallardo_, Jun 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014750 | OEIS-A335402

Numbers m such that the only normal integer partition of m whose run-lengths are a palindrome is (1)^m.

Conjecture: The sequence consists of 0, 1, 4, and all primes except 3. The above conjecture is true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jun 06 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014751 | OEIS-A335411

a(n) is the number of vertices formed by n-secting the angles of an equilateral triangle.

Conjectures from _Colin Barker_, Jun 08 2020: (Start) Colin Barker's recurrence conjecture holds for 13 < n <= 500. _Lars Blomberg_, Jun 12 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lars Blomberg_, Jun 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014752 | OEIS-A335507

Index of the least Wendt determinant (A048954) divisible by prime(n).

It has been conjectured by Michael B Rees that there exists for every prime a Wendt determinant divisible by that prime. However the conjecture has been proved for all prime divisors equivalent to -1 (mod 6) - (see Lehmer link below).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_ and Michael B Rees, Jun 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014753 | OEIS-A335617

Let c(1) = c(2) = 0, c(3) = 1, and c(n + 3) = (c(n) - 2*c(n + 1) + c(n + 2))/n, then a(n) = ceiling (c(n)).

Conjectured quasiperiodicity with autocorrelation function R(x) = 1/2 if x = 0, 1/4 if x > 0. Some other proved or conjectured (or suspected) nonperiodic binary sequences where there are no more than two consecutive 0's or 1's include: A083035, A285305, A190843, A286059, A288213, A288551, A288473, A176405, A188321, A188398, A191162, A272170, A197879, A078588, A272532, A273129, A074937, A188297, A289128. Others?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andres Cicuttin_, Oct 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014754 | OEIS-A335624

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x + 3*y + 4*z a square, where x, y, z, w are nonnegative integers.

Conjecture: a(n) > 0 if n is not divisible by 8. Moreover, a(n) = 0 if and only if n has the form 2^(4k+3)*m (k >= 0 and m = 1, 3, 5, 43). We have verified this for n up to 3*10^6. The conjecture is similar to the author's 1-3-5 conjecture (cf. A271518). In his 2017 JNT paper, the author conjectured that any natural number not of the form 2^(4k+2)*7 (k = 0,1,...) can be written as w^2 + x^2 + y^2 + z^2 with w + 2*x + 3*y + 5*z a square, where w, x, y, z are nonnegative integers. See also A338019 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014755 | OEIS-A335636

Expansion of e.g.f. Product_{k>0} 1/(1 - tan(x)^k / k).

Conjecture: a(n) ~ A080130 * n * 2^(2*n+2) * n! / Pi^(n+2). - _Vaclav Kotesovec_, Oct 04 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Oct 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014756 | OEIS-A335638

Expansion of e.g.f. Product_{k>0} (1 + tan(x)^k / k).

Conjecture: a(n) ~ A080130 * 2^(2*n+1) * n! / Pi^(n+1). - _Vaclav Kotesovec_, Oct 04 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Oct 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014757 | OEIS-A335641

Number of ordered ways to write 2n+1 as p + x*(9x+7) with p prime and x an integer.

Conjecture 1: a(n) > 0 for all n > 0. Also, a(n) = 1 only for n = 1, 4, 5, 7, 8, 18, 24, 25, 42, 68, 70, 85, 117, 118, 196, 238, 287, 497, 628, 677, 732. Conjecture 2: Let f(x) be any of the polynomials x*(3x+1), x*(5x+1), 2x*(3x+1), 2x*(3x+2). Then, each odd integer greater than one can be written as p + f(x) with p prime and x an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014758 | OEIS-A335653

Smallest prime whose binary expansion has Hamming distance 1 from 2n+1, or 0 if no such prime exists.

2131099 is Sierpiński, and it is conjectured that the Sierpiński numbers are the same as the dual Sierpiński numbers. Furthermore 2131099 is the smallest Sierpiński number whose binary representation has the property stated above. If the dual Sierpiński conjecture holds, and if A076336 is complete up to 2131099, then a(1065549) = 0 and this is likely the first 0 in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ross Dempsey_, Jun 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014759 | OEIS-A335654

Numbers m such that the elements of all Pythagorean quadruples belonging to the set of divisors are exactly their first k divisors for some k.

Conjecture 1: a(n) == 0 (mod 72). Conjecture 2: if the numbers m such that the elements of all Pythagorean quadruples contained in the set of divisors of m are exactly the first k divisors of m, then k = tau(m) - 4 or tau(m) - 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jun 16 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014760 | OEIS-A335716

a(n) = pi(pi(n)) - pi(Sum_{k=1..n-1} a(k)) with a(1) = 0.

Conjecture: a(n) hits every nonnegative integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Jun 18 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014761 | OEIS-A335753

Maximal length of the shortest walk on any n X n maze that starts in one corner and visits the other three corners.

Daniel Mathias conjectures that a(n) cannot exceed 10*n*n/9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Jun 20 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014762 | OEIS-A335756

A cup filling problem starting with 2 empty cups of sizes 3 and n, where a(n) is the number of unreachable states (see details in comments).

Conjectures from _Colin Barker_, Jun 21 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Jun 20 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014763 | OEIS-A335793

Number of hare pop-stack sortable Cayley permutations.

Conjectures from _Colin Barker_, Jun 24 2020: (Start) The above conjectures are correct. - _Abigail Ollson_, Feb 12 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jun 23 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014764 | OEIS-A335817

a(k) is the index of the first occurrence of 2*k-1 in A208884, or 0 if it does not occur.

It is conjectured that A208884 contains all the odd numbers. If so, a(n) > 0 for every n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Resta_, Jun 25 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014765 | OEIS-A335866

Number of classes of simple difference sets of the Singer type (m^2 + m + 1, m + 1, 1) with m = m(n) = A000961(n), for n >= 1.

The formula given below was conjectured by Singer for n >= 2 on p. 383. See also the table on p. 384. This conjecture was later proved by Berman. If the conjectures in A058241 are true, we have a(n) = 2*A058241(A000961(n)+1) for n > 1. - _Pontus von Brömssen_, Aug 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jul 26 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014766 | OEIS-A335898

a(n) = a(floor((n-1)/a(n-1))) + a(floor((n-2)/a(n-2))) with a(1) = a(2) = 1.

Conjecture: This sequence hits every positive integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, following a suggestion from _Andrew R. Booker_, Jun 29 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014767 | OEIS-A335901

a(n) = 2*a(floor((n-1)/a(n-1))) with a(1) = 1.

Least k such that a(k) = 2^n are 1, 2, 5, 21, 169, 2705, ... (Conjecture: This sequence is A117261).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, following a suggestion from _Andrew R. Booker_, Jun 29 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014768 | OEIS-A335925

a(n) = a(floor((n-1)/a(n-1))) + 1 with a(1) = 1.

Least k such that a(k) = n are 1, 2, 5, 16, 65, 326, 1957, ... (Conjecture: This sequence is A000522).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Jun 30 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014769 | OEIS-A335970

Index of the least Wendt determinant (A048954(k)) that is divisible by the least prime of the form k*prime(n) + 1.

It has been conjectured by Michael B Rees that for every prime p there exists a Wendt determinant index j such that for all j < k primes of the form j*p + 1 will not divide Wendt(j). This sequence gives the least index k such that Wendt(k) is divisible by the least prime of the form k*p + 1 for each prime p = prime(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_ and Michael B Rees, Jul 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014770 | OEIS-A335976

Numbers k such that Fibonacci(6*k) is not a totient.

Conjecture: Sequence contains infinitely many primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Jul 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014771 | OEIS-A335990

The moment generating function of the limiting distribution of the number of comparisons in quicksort can be written in the form M(t) = m(-2*t)/(exp(2*gamma*t)*Gamma(1 + 2*t)) for |t| < 1/2, where m(z) = Sum_{n >= 0} B(n)*z^n/n! for |z| < 1. This sequence gives the numerators of the rational numbers B(n) for n >= 0.

The formula for M(t) appears as Theorem 6.1 in Tan and Hadjicostas (1993) and derives from the work of Hennequin (1991). Hennequin conjectured a cumulant formula for the limiting distribution of the number of comparisons in quicksort in his 1989 paper, and he proved it in his 1991 thesis.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, Jul 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014772 | OEIS-A336035

The leftmost digit d of a(n) jumps over d positions to the left and gets duplicated there. Lexicographically earliest sequence of distinct positive integers with this property and a(1) = 10.

This is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Carole Dubois_, Jul 05 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014773 | OEIS-A336088

k such that L(H(k,2)) = 2*L(H(k,1)) where L(x) is the number of terms in the continued fraction of x and H(k,r) = Sum_{u=1..k} 1/u^r.

Conjecture: this sequence is infinite. More generally for any fixed integers a,b,c,d >= 1, there are infinitely many k's such that c*d*L(H(k,a)^b) = a*b*L(H(k,c)^d) where L(x) is the number of terms in the continued fraction of x and H(k,r) = Sum_{u=1..k} 1/u^r. Here, (a,b,c,d) = (2,1,1,1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014774 | OEIS-A336089

k such that L(H(k,1)^2) = 2*L(H(k,1)) where L(x) is the number of terms in the continued fraction of x and H(k,r) = Sum_{u=1..k} 1/u^r.

Conjecture: this sequence is infinite. More generally, for any fixed integers a,b,c,d >=1, there are infinitely many k's such that c*d*L(H(k,a)^b) = a*b*L(H(k,c)^d) where L(x) is the number of terms in the continued fraction of x and H(k,r) = Sum_{u=1..k} 1/u^r. Here (a,b,c,d) =(1,2,1,1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014775 | OEIS-A336090

Number of k's, 1 <= k <= n, such that the continued fraction for n^2/k^2 has twice as many elements as the continued fraction for n/k.

Conjecture: a(n) >= 1 for n >= 15. Conjecture: a(n) << n/log(n) (see links for the graph of a(n)/n*log(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Oct 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014776 | OEIS-A336174

Number of non-symmetric binary n X n matrices M over the reals such that M^2 is the transpose of M.

Row sums of T(n,k) generate known terms of this sequence and the next term a(10) evaluates to 209920 (see conjectured formula below). a(n) = Sum_{k=0..n} A001471(k) * binomial(n, k). [Previously conjectured, for a proof see the link in A344912.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Torlach Rush_, Jul 10 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014777 | OEIS-A336186

Side length of a square block of integers, with 1 at the top-left corner, on a diagonally numbered 2D board such that the sum of the integers in the square is a perfect square.

Conjectures from _Colin Barker_, Jul 11 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _Eric Angelini_, Jul 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014778 | OEIS-A336205

Numbers k that can be expressed as x^3 + y^3 + z^3 with x^2 + y^2 + z^2 <= k where x, y, z are integers.

By definition, corresponding (x,y,z) variables are produced by equation x^3 + y^3 + z^3 = x^2 + y^2 + z^2 + t with t >= 0. That is, x^2*(x-1) + y^2*(y-1) + z^2*(z-1) >= 0. Conjecture: Every even integer can be represented as x^2*(x-1) + y^2*(y-1) + z^2*(z-1) where x, y, z are integers. - _Altug Alkan_, Jul 19 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Jul 12 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014779 | OEIS-A336244

Triangle read by rows: row n gives coefficients T(n,k), in descending powers of m, of a polynomial Q_n(m) (of degree n - 1) in an expression for the number of subdivisions A(m,n) of a grid with two rows.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_ and _Petros Hadjicostas_, Jul 14 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014780 | OEIS-A336280

Number of consecutive primes of the form k*prime(n) + 1, starting with the least such prime A035095(n), that divides the Wendt determinant A048954(prime(n)).

Michael B Rees has conjectured that:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_ and Michael B Rees, Jul 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014781 | OEIS-A336305

Alternating row sums of triangle A211343.

Conjecture: a(n) = (A000203(n) + A067742(n))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Oct 05 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014782 | OEIS-A336347

Least prime factor of 44745755^4*2^(4n+2) + 1.

There are k such that k*2^m + 1 is not prime for any m (then k is called a Sierpiński number). Erdős once conjectured that for such a k, the smallest prime factor of k*2^m + 1 would be bounded as m tends to infinitiy. The proven Sierpiński number k=44745755^4 is thought to be the first counterexample to this conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeppe Stig Nielsen_, Jul 19 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014783 | OEIS-A336351

Number of cyclic arrangements of S = {1,2,...,6n - 3} such that any three neighbors can be reordered in an arithmetic progression.

It seems that in S = {1,2,...,k} with k == 0 (mod 6) these cyclic arrangements are not possible, hence the definition of the sequence. Computationally, this conjecture is valid for k <= 114.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Fausto A. C. Cariboni_, Jul 19 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014784 | OEIS-A336397

Numbers k such that both k and A122111(k) [its conjugated prime factorization] are Ore's Harmonic numbers (A001599).

There also at least these additional terms present: 439151400324000, 889013810412000, 106713790278732000, that were found from the file of conjectured initial 10000 terms of A001599 provided by _David A. Corneth_ under that entry. However, there might be further terms in the range between.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jul 20 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014785 | OEIS-A336404

a(n) = floor(n*frac(prime(n)/pi(n))), where frac denotes the fractional part.

Conjecture: For any m >= 0 there is a k such that a(k) = m. Also for any reals x and epsilon such that 0 < x < 1 and epsilon > 0, there is a k such that abs(x - frac(prime(k)/pi(k))) < epsilon.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andres Cicuttin_, Jul 20 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014786 | OEIS-A336431

Number of ordered ways to write n as the sum of a practical number (A005153) and a generalized heptagonal number (A085787).

Conjecture 1: a(n) > 0 for all n > 0. Also, a(n) = 1 only for n = 1, 3, 4, 7, 10, 11, 14, 18, 23, 26, 27, 44, 51, 53, 65, 68, 77, 80, 92, 125, 143, 170, 179, 182, 185, 191, 206, 296, 362, 383, 425, 437, 674, 1340, 1622, 2273, 2558, 3167, 5591. Conjecture 2: Any positive integer can be written as the sum of a practical number and a generalized pentagonal number. Both conjectures are motivated by A208244.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014787 | OEIS-A336485

The least positive integer k for which there exist primes p, q, r such that phi(p*q*s^k) = phi(r*s^(2k+1)) and sigma(p*q*s^k) = sigma(r*s^(2k+1)), where s is the n-th prime.

Conjecture: For each prime s there are infinitely many positive integers k such that numbers p*q*s^k and r*s^(2k+1) are similar for some primes p, q, r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Letsko_, Jul 23 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014788 | OEIS-A336486

The least prime s for which there exist primes p, q, r such that phi(p*q*s^n) = phi(r*s^(2n+1)) and sigma(p*q*s^n) = sigma(r*s^(2n+1)).

Conjecture: For each positive integer n there are infinitely many primes s such that numbers p*q*s^n and r*s^(2n+1) are similar for some primes p, q, r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Letsko_, Jul 23 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014789 | OEIS-A336532

Square array read by antidiagonals upwards showing a stained glass windows with two colors and a hidden curve from the toothpick cellular automaton of A139250 (see Comments lines for definition).

Conjecture 1: after 2^k stages the number of cells of color 1 is equal to the number of cells of color 2. Conjecture 2: after 2^k stages, k >= 2, in the structure there are essentially one major region of color 1 and two major regions of color 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Oct 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014790 | OEIS-A336585

Integers m such that (m-d)*(m+d) < (m-1)^2, where d is the smallest number such that both m-d and m+d are primes.

It seems that the number of terms in this sequence is finite, with the last term being a(1225) = 1353559. Conjecture: there exist only 1225 semiprimes of the form (m-d)*(m+d), where d is the smallest number such that (m-d)*(m+d) < (m-1)^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Oct 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014791 | OEIS-A336597

a(n) is the least number k of the form (the number of nonnegative bases m < n such that m^k == m (mod n))/(the number of nonnegative bases m < n such that -m^k == m (mod n)) dividing n, or 0 if no such k exists.

Conjecture: For each natural number n, either there is a divisor k of the form (the number of nonnegative bases m < n such that m^k == m (mod n))/(the number of nonnegative bases m < n such that -m^k == m (mod n)) and it is unique, or it does not exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jul 26 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014792 | OEIS-A336614

Number of n X n (0,1)-matrices A over the reals such that A^2 is the transpose of A.

[The above formulas, first stated as conjectures, were proved by mjqxxxx at Mathematics Stack Exchange, see link.] (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Torlach Rush_, Jul 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014793 | OEIS-A336633

Triangle read by rows: T(n,k) is the number of generalized permutations related to the degenerate Eulerian numbers with exactly k ascents (0 <= k <= max(0,n-1)).

The following remarks are all conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Orli Herscovici_, Jul 28 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014794 | OEIS-A336660

a(n) is the maximal number of 1 X 1 squares in an arrangement of n squares, from 1 X 1 to n X n.

I myself conjecture that a(n) <= floor((n^2 - n)*(3/5)) for n >= 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rodolfo Kurchan_, Jul 28 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014795 | OEIS-A336687

Numbers m such that tau(sigma(m)) and sigma(tau(m)) both divide m, where tau(m) is the number of divisors function (A000005) and sigma(m) is the sum of divisors function (A000203).

Conjecture: The only m such that m = tau(sigma(m))*sigma(tau(m)) are in {1,468,3240}. Verified for m up to 1*10^9. - _Ivan N. Ianakiev_, Aug 06 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Jul 31 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014796 | OEIS-A336688

Primes p such that the Wendt determinant A048954(p) has prime factors less than p.

Michael B Rees has conjectured that for all primes p, each fully exponentiated prime factor less than p that divides the Wendt determinant W(p), if it exists, is of the form k*p + 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_ and Michael B Rees, Jul 31 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014797 | OEIS-A336700

Numbers k such that the odd part of (1+k) divides (1 + odd part of sigma(k)).

Conjecture: After 1, all terms are of the form 4u+3 (in A004767). If this could be proved, then it would refute at once the existence of both the odd perfect numbers and the quasiperfect numbers. Concentrating on the latter is probably easier, as it is known that all quasiperfect numbers must be odd squares, thus k is of the form 4u+1, in which case the condition given in A336701 that A000265(1+A000265(sigma(k))) must be equal to A000265(1+k) reduces to a simpler form, A000265(1+sigma(k)) = (1+k)/2, and as k = s^2, with s odd, so (s^2 + 1)/2 should divide 1+sigma(s^2). Does that condition allow any other solutions than s=1 ? See A337339.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 02 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014798 | OEIS-A336701

Numbers k for which A000265(1+A000265(sigma(k))) is equal to A000265(1+k).

Conjecture: There are no even terms. This is equivalent to claim that there are no k such that A336698(k) = 1+k: If we assume that k is even, then in above equations we set e=0, and the requirement will then become that A337194(k) = 2^A337195(k)*(1+k), thus 1+k = A336698(k) = A000265(1+A000265(sigma(k))).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 02 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014799 | OEIS-A336744

Integers b where the number of cycles under iteration of sum of squares of digits in base b is exactly three.

A 1978 conjecture of Hasse and Prichett describes the set L(x^2,2). New elements have been added to this set in the paper Integer Dynamics, by D. Lorenzini, M. Melistas, A. Suresh, M. Suwama, and H. Wang. It is natural to wonder whether the set L(x^2,3) is infinite. It is a folklore conjecture that L(x^2,1) = {2,4}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dino Lorenzini_, Aug 02 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014800 | OEIS-A336759

a(n) is the smallest positive number that is the sum of no fewer than n distinct repdigits.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Falcon Shapiro_, Aug 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014801 | OEIS-A336762

Integers b where the number of cycles under iteration of sum of squares of digits in base b is exactly two.

A 1978 conjecture of Hasse and Prichett describes the set L(x^2,2). New elements have been added to this set in the paper Integer Dynamics, by D. Lorenzini, M. Melistas, A. Suresh, M. Suwama, and H. Wang. The sequence contains all b <= 10^6 that are in L(x^2,2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Makoto Suwama_, Aug 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014802 | OEIS-A336764

Maximum number of order 3 subsquares in a Latin square of order n.

a(3^n) = 9*a(3^(n-1)) + 27^(n-1) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar Aceval Garcia_, _Cameron Byer_, _Eugene Fiorini_, _Nicholas Hanson_, _Brian G. Kronenthal_, _Lindsey Wise_, Aug 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014803 | OEIS-A336792

Values of odd prime numbers, D, for incrementally largest values of minimal positive y satisfying the equation x^2 - D*y^2 = -2.

For solutions of this Diophantine equation it is sufficient to consider the odd primes p(n) := A007520(n), for n >= 1, the primes 3 (mod 8). Also prime 2 has the fundamental solution (x, y) = (0, 1). If there is a solution for p(n) then there is only one infinite family of solutions because there is only one representative parallel primitive binary quadratic form for Discriminant Disc = 4*p(n) and representation k = -2. Only proper solutions can occur. The conjecture is that each p(n) leads to solutions. For the fundamental solutions (with prime 2) see A339881 and A339882. - _Wolfdieter Lang_, Dec 22 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christine Patterson_, Oct 14 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014804 | OEIS-A336893

Lexicographically earliest infinite sequence of distinct positive terms such that the sum of digits of the first n terms is coprime to their concatenation.

Conjecture: A permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_ and _Michael De Vlieger_, Aug 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014805 | OEIS-A336897

Infinite sum of the natural numbers, compacted (see comments for an explanation).

The sequence is a subset of A038550, numbers that can be expressed as the sum of k>1 consecutive positive integers in only one way. If the successive terms of the present sequence are expressed as the sum of k>1 consecutive integers and added, the result will be 1 + 2 + 3 + 4 + 5 + 6 + 7 + 8 + 9 + 10 + 11... (conjectured to extend ad infinitum).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Aug 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014806 | OEIS-A336943

Least prime factor of 44745755^4 + 2^(4n+2).

k = 44745755^4 has the property that k + 2^m is composite for all m. However, it is conjectured that this sequence is unbounded. This is the case if and only if A336347 is unbounded; because a full covering set for k*2^m + 1 would also be a full covering for k + 2^m, and vice versa.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeppe Stig Nielsen_, Aug 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014807 | OEIS-A336957

The Enots Wolley sequence: the lexicographically earliest infinite sequence {a(n)} of distinct positive numbers such that, for n>2, a(n) has a common factor with a(n-1) but not with a(n-2).

Conjecture 1: Every number with at least two distinct prime factors is in the sequence. In other words, apart from 1 and 2, this sequence is the complement of A000961. [It seems very likely that the arguments used to prove Theorem 1 of the Yellowstone Permutation paper can be modified to prove the conjecture.] [It appears that Conjecture 1 has been proved by Nathan Myles Nichols: see link. - _N. J. A. Sloane_, Oct 11 2026] We conjecture that even if p is introduced by some prime q>2, 2*p appears later. The differences between the two sequences are caused by the fact that although normally if p and q are odd primes with p < q, then 2p precedes 2q, this is not true for the following primes: (7,5), (31,29), and (109, 113, 107), which appear in the order shown. We conjecture that these are the only exceptions. Combining the above observations, we conjecture that for n >= 755 (at which point we have seen all the primes <= 367), every prime p is introduced by 2*p, and the terms 2*p appear in their natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Aug 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014808 | OEIS-A336972

Sum of the smallest two side lengths of all distinct integer-sided triangles with perimeter n.

Conjectures from _Colin Barker_, Aug 10 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Aug 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014809 | OEIS-A336973

Sum of the smallest and largest side lengths of all distinct integer-sided triangles with perimeter n.

Conjectures from _Colin Barker_, Aug 10 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Aug 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014810 | OEIS-A336974

Sum of the largest two side lengths of all distinct integer-sided triangles with perimeter n.

Conjectures from _Colin Barker_, Aug 10 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Aug 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014811 | OEIS-A336981

a(n) = (Sum_{k=0..n-1} (4290*k + 367)*3136^(n-1-k)*C(2*k, k)*T_k(14, 1)*T_k(17, 16)) / (n*C(2*n-1, n-1)), where T_k(b, c) denotes the coefficient of x^k in the expansion of (x^2 + b*x + c)^k.

Conjecture 1: a(n) is an integer for each n > 0. Moreover, a(n) is even for every n > 1. Conjecture 2: Denote (4290k+367)/3136^k*C(2k,k)*T_k(14,1)*T_k(17,16) by t(k). Conjecture 3. Let p > 7 be a prime and let S(p) denote the sum Sum_{k=0..p-1}C(2k,k)*T_k(14,1)*T_k(17,16). See also A336982 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014812 | OEIS-A336982

a(n) = (Sum_{k=0..n-1}(540*k + 137)*3136^(n-1-k)*C(2*k, k)*T_k(2, 81)*T_k(14, 81))/(2*n*C(2*n, n)), where T_k(b, c) denotes the coefficient of x^k in the expansion of (x^2 + b*x + c)^k.

Conjecture 1: a(n) is an integer for each n > 1. Moreover, a(n) is odd if and only if n = 2^k + 1 for some nonnegative integer k. Conjecture 2: Denote (540*k+137)/3136^k*C(2k,k)*T_k(2,81)*T_k(14,81) by t(k). Conjecture 3. Let p > 7 be a prime, and let S(p) denote the sum Sum_{k=0..p-1}C(2k,k)*T_k(2,81)*T_k(14,81). See also A336981 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014813 | OEIS-A337015

Number of distinct transitive subgroups of S_n, counting conjugates as distinct.

I have conjectured that log(A005432(n)/a(n)) > (n-1)/2 for n prime and log(A005432(n)/a(n)) < (n-1)/2 for n composite. L. Pyber shows c^{n^2*(1+o(1))} <= a(n) <= d^{n^2*(1+o(1))}, c=2^{1/16}, d=24^{1/6}; conjectures lower bound is accurate.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John Erickson_ and _Alexander Hulpke_, Nov 21 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014814 | OEIS-A337020

Maximum number of circles with unit area that can be packed into a square with an area of n.

Terms beyond a(n) = 30 (n = 38 & 39) except a(n) = 36 are conjectures supported by numerical results (see Packomania in the links) and terms for n from 40 through 70 are: 31, 31, 33, 33, 34, 35, 36, 36, 38, 39, 39, 40, 41, 42, 42, 43, 44, 44, 46, 46, 47, 48, 49, 50, 52, 52, 53, 53, 54, 55, 56.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Nov 06 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014815 | OEIS-A337036

Numbers m such that the elements of all quadruples (x, y, z, w) satisfying x^3 + y^3 + z^3 = w^3 contained in the set of divisors of m are exactly the first k divisors of m for some k.

Conjecture 1: a(n) == 0 (mod 144). Conjecture 2: if the numbers m such that the elements of all quadruples (x, y, z, w) satisfying x^3 + y^3 + z^3 = w^3 contained in the set of divisors of m are exactly the first k divisors of m, then k = tau(m) - 5 or tau(m) - 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Aug 12 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014816 | OEIS-A337046

Integers n such that n! = x^2 + y^3 + z^6 where x, y and z are nonnegative integers, is soluble.

Conjecture I: Natural density of this sequence is 1. Conjecture II: Any sufficiently large n is in the sequence. Conjecture III: There is a fixed value of t such that all integers >= t are terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Aug 12 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014817 | OEIS-A337065

Infinite sum of the prime numbers, compacted (see the Comments line for an explanation).

If the successive terms of the present sequence are expressed as the sum of k>1 consecutive primes in only one way and added, the end result will be 2 + 3 + 5 + 7 + 11 + 13 + 17 + 19 + 23 + ... (conjectured to extend ad infinitum).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Aug 14 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014818 | OEIS-A337066

a(n) is the smallest k such that A336957(k) = n, or -1 if n does not appear in A336957.

There is a strong conjecture that numbers n >= 3 appear in A336957 if and only if they are divisible by at least two different primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Aug 13 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014819 | OEIS-A337082

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with 2*x^2 + 4*y^2 - 7*x*y a power of two (including 2^0 = 1), where x, y, z, w are nonnegative integers with z <= w.

Conjecture: a(n) > 0 for all n > 0. Moreover, any positive integer n congruent to 1 or 2 modulo 4 can be written as x^2 + y^2 + z^2 + w^2 with x, y, z, w nonnegative integers such that 2*x^2 + 4*y^2 - 7*x*y = 4^k for some positive integer k. See also A338139 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 12 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014820 | OEIS-A337097

Infinite sum of the odd numbers, compacted (see the Comments line for an explanation).

When the successive terms of the present sequence are expressed as the sum of k>1 consecutive odd numbers and added, the end result will be 1 + 3 + 5 + 7 + 9 + 11 + 13 + 15 + 17..... (conjectured to extend ad infinitum).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Aug 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014821 | OEIS-A337121

a(n) is the number of ways the n-th prime number prime(n) can be represented as sum of two smaller odd prime numbers p1, p2 with prime(n) > p1 > (p2 minus the maximum odd prime factor of (p1-p2)).

Conjecture: a(n) > 0 when n >= 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Aug 17 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014822 | OEIS-A337136

a(1) = 1, a(2) = 2; for n > 1, a(n) = smallest positive number which has not yet appeared which has a common factor with a(n-2) + a(n-1).

Conjecture: This is a permutation of the natural numbers. Up to 1 million terms the only fixed points are 1,2,3,6,9,10, and it is likely that there are no others.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Aug 18 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014823 | OEIS-A337143

Numbers k for which there are only 3 bases b (2, k+1 and another one) in which the digits of k contain the digit b-1.

Conjecture: this sequence is finite and full. a(26) > 3.8*10^12 if it exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _François Marques_, Sep 14 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014824 | OEIS-A337184

Numbers divisible by their first digit and their last digit.

Conjecture: 3n < a(n) < 7n for n > 75. - _Charles R Greathouse IV_, Dec 02 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Jan 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014825 | OEIS-A337209

Triangle read by rows T(n,k), (n >= 1, k > = 1), in which row n has length A000070(n-1) and every column gives A000203, the sum of divisors function.

Conjecture: the sum of row n equals A066186(n), the sum of all parts of all partitions of n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Nov 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014826 | OEIS-A337212

Modulo 3 Pisano period of 'n-bonacci' series.

Conjecture: a(3^k-1)=a(3^k)-3^k-2=3^(2k)-1, a(3^k)=3^k(3^k+1)+1 for k>0. - _Chai Wah Wu_, Sep 15 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Adam Bascal_, Aug 19 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014827 | OEIS-A337226

Lexicographically earliest sequence of positive integers with the property that, for all k > 0, there is at most one j such that a(j) = a(j+k).

Conjecture: For all k > 0, there is exactly one j such that a(j) = a(j+k). For 0 < k < 11911, this conjecture holds.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Samuel B. Reid_, Aug 19 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014828 | OEIS-A337253

a(2n) = b(a(n)^2), a(2n+1) = b(a(n) + a(n - 1)) where b(k) = abs(k)*(-1)^k; a(0) = 0, a(1) = 1.

Conjecture: {abs(a(n))} contains every natural number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mario Cortés_, Aug 21 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014829 | OEIS-A337275

Index of appearance of 2*prime(n) in A336957, or -1 if 2*prime(n) never appears.

It is a strong conjecture that 2*prime(n) appears in A336957 for all n>1, and it is known that 4 does not appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014830 | OEIS-A337276

Index of appearance of 2*n in A336957, or -1 if 2*n never appears.

It is a strong conjecture that 2*n appears in A336957 for all odd n, and it is known that 2^i does not appear for i>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014831 | OEIS-A337300

Partial sums of the geometric Connell sequence A049039.

a(n) is Newey's "more complicated" conjectured length of the shortest sequence containing all permutations of 1..n (A062714). It agrees with A062714(n) for n <= 7. [But not for n=8. - _Pontus von Brömssen_, Aug 18 2025]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kevin Ryde_, Aug 22 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014832 | OEIS-A337332

a(n) = Sum_{k=0..n}C(n,k)*C(n+k,k)*C(2k,k)*C(2n-2k,n-k)*(-8)^(n-k).

Conjecture 1: Sum_{k>=0}(4k+1) a(k)/(-48)^k = sqrt(72+42*sqrt(3))/Pi. Conjecture 2: For each n > 0, the number (Sum_{k=0..n-1} (-1)^k*(4k+1)*48^(n-1-k)*a(k))/n is a positive integer. Conjecture 3: For any prime p > 3, the square of (Sum_{k=0..p-1} (4k+1)a(k)/(-48)^k)/p is congruent to 14*(3/p)-(p/3)-12 modulo p, where (a/p) is the Legendre symbol. Conjecture 4: Let p > 3 be a prime, and let S(p) = Sum_{k=0..p-1} a(k)/(-48)^k. If p == 1 (mod 4) and p = x^2 + 4y^2 with x and y integers, then S(p) == 4x^2-2p (mod p^2). If p == 3 (mod 4), then S(p) == 0 (mod p^2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 23 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014833 | OEIS-A337347

a(n) = a(n-2) + a(n-1) if that sum is prime. Otherwise, a(n) = a(n-2) + a(n-1) + prime(m-1) + prime(m-2) + ... + prime(s), where a(-1) = a(0) = prime(0) = 1, m = pi(a(n-2)), and s = max(0, 1, 2, ..., m-1) such that the sum is prime.

Conjecture: This sequence is infinite. - _Ya-Ping Lu_, Apr 02 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Aug 25 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014834 | OEIS-A337354

a(n) is the numerator of Product_{i=0..n-1} (n-i)^((-1)^ceiling(i/2)).

Conjecture: a(4*n)/A337355(4*n) ~ 0.5990701173677... (=A076390). - _Andrew Howroyd_, Aug 25 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Devansh Singh_, Aug 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014835 | OEIS-A337366

Number of representations of A036691(n) as a sum of 3 nonnegative cubes.

Conjecture I: a(n) = 0 only for n = 1. That is, any product of first n > 1 composite numbers is a sum of at most 3 positive cubes. For example, Conjecture II: For any term t >= 1, there are only finitely many values of n such that a(n) = t.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Aug 25 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014836 | OEIS-A337389

Square array T(n,k), n >= 0, k >= 0, read by antidiagonals, where column k is the expansion of sqrt((1+(k-4)*x+sqrt(1-2*(k+4)*x+((k-4)*x)^2)) / (2 * (1-2*(k+4)*x+((k-4)*x)^2))).

Conjecture: the k-th column entries, k >= 0, are given by [x^n] ((1 + (k-2)*x + x^2)*(1 + x)^2/(1 - x)^2)^n. This is true for k = 0 and k = 4. - _Peter Bala_, May 03 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Aug 25 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014837 | OEIS-A337390

Expansion of sqrt((1-2*x+sqrt(1-12*x+4*x^2)) / (2 * (1-12*x+4*x^2))).

Conjecture: a(n) = [x^n] ((1 + x^2)*(1 + x)^2/(1 - x)^2)^n. Equivalently, a(n) = Sum_{k = 0..n} Sum_{j = 0..n-2*k} binomial(n,k)*binomial(2*n,j)*binomial(3*n-2*k-j-1,n-2*k-j). If the conjecture is true then the Gauss congruences a(n*p^k) == a(n*p^(k-1)) (mod p^k) hold for all primes p and positive integers n and k. Calculation suggests that the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(2*k)) hold for primes p >= 3 and positive integers n and k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Aug 25 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014838 | OEIS-A337455

Numbers of the form m + bigomega(m) with m a positive integer.

Kucheriaviy proves that a(n) << n log log n and conjectures that a(n) ≍ n, that is, these numbers have positive lower density. - _Charles R Greathouse IV_, Dec 07 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nathan J. McDougall_, Aug 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014839 | OEIS-A337476

Position of the first occurrence of n in A337474.

For n >= 2, a(n) = A000040(A396829(n)). [Conjectured] - _Antti Karttunen_, Sep 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 28 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014840 | OEIS-A337479

Primitive elements of A337386: numbers k for which sigma(A003961(k)) >= 2*A003961(k), but none of the proper divisors of k satisfy the same condition.

Conjecture: every positive integer is either a (possibly trivial) multiple of a sequence term or divides infinitely many terms of this sequence. - _Peter Munn_, Sep 24 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 01 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014841 | OEIS-A337484

Number of ordered triples of positive integers summing to n that are neither strictly increasing nor strictly decreasing.

Conjectures from _Colin Barker_, Sep 13 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Sep 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014842 | OEIS-A337497

a(n) is the smallest integer k with exactly n bases b such that k in base b contains the digit b-1; or -1 if there is no such integer.

Conjecture: a(n) > log(n)^(sqrt(2)*log(n)) for n>1. This have been checked for n<3444, and for n<10275 unless if a(n)=-1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _François Marques_, Aug 29 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014843 | OEIS-A337499

a(n) is the number of ballot sequences of length n tied or won by at most 2 votes.

D-finite with recurrence +(n+2)*a(n) +n*a(n-1) +(-5*n-2)*a(n-2) +4*(-n+1)*a(n-3) +4*(n-3)*a(n-4)=0. - Conjectured by _R. J. Mathar_, Sep 27 2020, verified by _Robert Israel_, Oct 08 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nachum Dershowitz_, Aug 29 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014844 | OEIS-A337500

a(n) is the number of ballot sequences of length n tied or won by at most 3 votes.

Conjecture: D-finite with recurrence -(n+3)*(n-4)*a(n) +2*(n^2-2*n-11)*a(n-1) +4*(n-1)^2*a(n-2) -8*(n-1)*(n-2)*a(n-3)=0. - _R. J. Mathar_, Sep 27 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nachum Dershowitz_, Aug 30 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014845 | OEIS-A337536

Numbers k for which there are only 2 bases b (2 and k+1) where the digits of k contain the digit b-1.

This sequence has density 0. Conjecture: this sequence is finite and full. a(9) > 10^100 if it exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Aug 31 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014846 | OEIS-A337546

A binary sequence defined as follows: a(1)=0; thereafter choose a(n) so as to minimize the number of final consecutive repeated words, and if there is a choice, minimize the length of the repeated word.

Cousin of the so-called Linus sequence (A006345), which avoids the longest repeated suffix. Conjectures: 1. This is cubefree (the Linus sequence is not cubefree); 2. density of 0's exists and equals 1/2; 3. recurrent and mirror invariant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alessandro Della Corte_, Nov 22 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014847 | OEIS-A337606

Decimal expansion of the Gaussian twin prime constant: the Hardy-Littlewood constant for A096012.

Shanks (1960) conjectured that the number of these pairs with m <= x is asymptotic to c * li_2(x), where li_2(x) = Integral_{t=2..n} (1/log(t)^2) dt, and c is this constant. He defined c as in the formula section and evaluated it by 0.4876.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Sep 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014848 | OEIS-A337607

Decimal expansion of Shanks's constant: the Hardy-Littlewood constant for A000068.

Shanks (1961) conjectured that the number of primes of the form m^4 + 1 (A037896) with m <= x is asymptotic to c * li(x), where li(x) is the logarithmic integral function and c is this constant. He defined c as in the formula section and evaluated it by 0.66974.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Sep 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014849 | OEIS-A337608

Decimal expansion of Lal's constant: the Hardy-Littlewood constant for A217795.

Shanks (1967) conjectured that the number of primes of the form (m + 1)^4 + 1 such that (m - 1)^4 + 1 is also a prime (A217795 plus 1), with m <= x, is asymptotic to c * li_2(x), where li_2(x) = Integral_{t=2..n} (1/log(t)^2) dt, and c is this constant. He defined c as in the formula section, evaluated it by 0.79220 and named it after the mathematician Mohan Lal, who conjectured the asymptotic formula without evaluating this constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Sep 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014850 | OEIS-A337641

One-quarter of the number of regions in the central square of an equal-armed cross with arms of length n (as in A331456).

In the central square, the maximum number of sides in any region is (for n = 0, 1, 2, 3, ...) 3, 4, 6, 6, 6, 6, 6, 6, 6, 7, 7, 6, 6, 6, 6, 6, 7, 6, 7, 7, 6, 6, 6, 6, 6, 6, 6, 7, 7, 6, 6, 6, 6, 6, 6, 6, 7, 7, 6, 6, 6, ... We conjecture that 7 is the maximum. - _Lars Blomberg_, Sep 19 2020.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lars Blomberg_, _Scott R. Shannon_, and _N. J. A. Sloane_, Sep 17 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014851 | OEIS-A337645

a(1)=2; thereafter, a(n) = smallest number with at least two different prime factors that is missing from A336957 after A336957(n) has been found.

It is conjectured that every number with at least two different prime factors will eventually appear in A336957.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Sep 25 2020; corrected Oct 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014852 | OEIS-A337648

Odd primes p such that the first term in A336957 that is divisible by p is 2*p.

Conjecture 1: this sequence contains all primes > 367. Conjecture 2: The set of odd primes is partitioned into A337648, A337649, and {7}. (These conjectures have been checked for the first 161734 terms of A336957.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 26 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014853 | OEIS-A337649

Primes p such that the first term in {A336957(i): i >= 3} that is divisible by p is 3*p.

It is conjectured that there are no other terms besides the 34 terms shown here.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 26 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014854 | OEIS-A337705

Possible sums of orders of elements of finite groups.

Conjecture: If S is a simple group and G is a soluble group satisfying |S|=|G|, then psi(S) < psi(G).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. Farrokhi D. G._, Sep 16 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014855 | OEIS-A337742

Cycle length of applying XOR with the previous state and bit rotation on a string of n bits.

a(2*n) = 2*a(n) (conjectured). Conjecture: a(n) = 2^A007814(n) * (2^A007733(n) + q) * k, where q = -1 if n is in A038837, otherwise q = 1. General formula for k is unknown.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artem Yashin_, Sep 17 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014856 | OEIS-A337743

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x + 2*y a power of four (including 4^0 = 1), where x, y, z, w are nonnegative integers with z <= w.

Conjecture 1: a(n) > 0 if n is neither of the form 4^k*(4*m+3) (k>=0, m>=0) nor of the form 2^(4*k+3)*101 (k>=0). In particular, a(n^2) > 0 and a(2*n^2) > 0 for all n > 0. Conjecture 2: Any positive integer not of the form 16^k*m (k>=0, m = 1, 25, 46, 88) can be written as x^2 + y^2 + z^2 + w^2 (x,y,z,w >= 0) such that 2*x - y = 4^a for some nonnegative integer a. Conjecture 3: Any positive integer of the form 2^k*(2*m+1) (k>=0, m>=0) with k == floor(m/2) (mod 2) (such as positive squares) can be written as x^2 + y^2 + z^2 + w^2 (x,y,z,w >= 0) such that x + 3*y = 4^a for some nonnegative integer a.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 30 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014857 | OEIS-A337758

G.f. A(x) satisfies: [x^n] (1 + n*x + n*x^2 - A(x))^(n+1) = 0, for n > 0.

a(n) is odd iff n is a power of 2 (conjecture). a(n) = 2 (mod 4) iff n is twice the sum of two distinct powers of 2 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014858 | OEIS-A337818

Odd integers k>=3 such that k*(k-1)/2 divides 2^((k-1)/2)+1.

Conjecture: a(n) == 1 mod 162 for n >= 6. The next few larger terms of the form 162*k+1 are: 44165935747, 46696027123, 85683674179, 88567070707, 101297654083, 131264938963, 131315541283, 177876176419, 195689448883, 196838306227, 213339588643, 310256425603, 378897238243, 502106519683, 588454831747, 611537689459, 641551809187, 735075731107, 745956214867, 755236606483, 771153067603, 872146803043 (and they are all primes). - _Chai Wah Wu_, Oct 06 2020 The conjecture fails, as a(30) = 48269073907 (prime). - _Bill McEachen_, Jul 11 2025 For each term k, (k-1)/2 is a term of A006521. Conjecture: sequence consists of primes p == 3 (mod 8) such that (p-1)/2 is in A006521. - _Max Alekseyev_, Jul 26 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Sep 23 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014859 | OEIS-A337828

Odd integers k such that 3^((k-1)/2) + 1 == 0 (mod k*(k-1)/2).

Each term is a prime or Fermat pseudoprime base 3 (A005935). Furthermore, for each term k, (k-1)/2 is a term of A015949. Conjecture: sequence consists of primes p == 2 (mod 3) such that (p-1)/2 is in A015949. - _Max Alekseyev_, Jul 26 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Sep 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014860 | OEIS-A337830

Odd integers k such that 6^((k-1)/2) + 1 == 0 (mod k*(k-1)/2).

Conjecture: a(n) == 1 mod 406 for n > 5. - _Chai Wah Wu_, Oct 07 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Sep 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014861 | OEIS-A337832

Porous numbers: Numbers k which are not multiples of 10 such that every m with sum of digits = k and k a divisor of both m and rev(m) has a zero in its digits.

Up to k = 1000 no further porous numbers exist (see the linked file with the corresponding m where k | m and k | rev(m)). The steadily increasing number of possibilities to construct an m for a given k suggests that the 5 terms might be the only porous numbers, but a mathematical proof for this conjecture seems a big challenge.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ruediger Jehn_, Sep 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014862 | OEIS-A337856

Number of positive integers with n digits that are the product of two integers ending with 6.

Conjecture: lim_{n->oo} a(n)/a(n-1) = 10. The conjecture is true since the sequence A324297 has positive asymptotic density (see Corollary 1 in Spezia link). - _Stefano Spezia_, Mar 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Sep 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014863 | OEIS-A337868

Number of distinct residues of x^r (mod n), x=0..n-1, r=2, ..., n.

For n > 1, a(n) >= A000010(n) + 1 as all invertible elements of Z/nZ are powers, as is 0. (Conjecture: equality holds exactly for A000430, the primes and squares of primes.) - _Charles R Greathouse IV_, Dec 23 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Sep 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014864 | OEIS-A337912

G.f. A(x) satisfies: Sum_{n>=0} x^n * A(x)^(n*(n-1)/2) = 2 - Sum_{n>=0} (-x)^n * A(x)^(n*(n+1)/2).

a(2*n) = 0 (mod 16) for n>1 and a(2*n+1) = 8 (mod 16) for n>0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 29 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014865 | OEIS-A337924

Odd numbers, x, such that for all primes (p,q) such that x = p + 2q, x + 2p and x + 2q are composite.

The sequence is conjectured to be finite and full in A046927. The conjecture holds through 10^9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_, Oct 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014866 | OEIS-A337945

Numbers m with a solution (s,t,k) such that s^2 + t^2 = k*m, s + t = m, 1 <= s <= t and 1 <= k <= m - 1.

Conjecture: k is a term <=> k * Clausen(k, 1) <> 2 * Clausen(k, 0), (Clausen = A160014). In other words: k is in this sequence iff it is not an odd squarefree number. - _Peter Luschny_, Jun 08 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Oct 01 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014867 | OEIS-A337949

a(n) = 2^(n*(n-1)/2) + 2^(n*(n+1)/2) for n > 0, with a(0) = 1.

a(2*n+1) = 0 (mod 3), a(4*n+2) = 1 (mod 3), a(4*n+4) = 2 (mod 3), for n >= 0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014868 | OEIS-A337951

a(n) = 4^(n*(n-1)/2) + 4^(n*(n+1)/2) for n > 0, with a(0) = 1.

a(2*n+1) = 0 (mod 5), a(4*n+2) = 3 (mod 5), a(4*n+4) = 2 (mod 5), for n >= 0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014869 | OEIS-A337986

Prime numbers p such that v_p(A000166(k)) = v_p(k-1) for all k > 1, where v_p(k) is the p-adic valuation of k.

Miska (2016) proved that the complement of this sequence within the primes is infinite, and conjectured that this sequence is also infinite, and that its asymptotic density within the primes is 1/e (A068985). Numerically, he found that there are 28990 terms below 10^6, which are about 37% of all the primes less than 10^6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Jan 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014870 | OEIS-A337993

Numbers k such that L(k) < sigma(k) + k/Pi^2, where L(k) = floor(H(k) + exp(H(k)) * log(H(k))) and H(k) = Sum_{j=1..k} 1/j.

Conjecture: This sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Oct 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014871 | OEIS-A338019

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with 3*x + 10*y + 36*z a positive square, where x, y, z, w are nonnegative integers.

Conjecture: a(n) > 0 if n is not divisible by 8. Moreover, a(n) = 0 if and only if n has the form 2^(4k+3)*m (k >= 0 and m = 1, 3, 5, 61). We have verified this for n up to 5*10^6. See also A335624 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014872 | OEIS-A338020

a(n) is the number of circles of positive integer area with radii less than n and greater than n - 1.

Conjecture: k >= 2, each triple Tr(k) = {a(k), a(k+1), a(k+2)} gives the sides of an integer-sided triangle, and {(a(k+2) - a(k)), (a(k+2) - a(k+1)), (a(k+1) - a(k))} is a degenerate integer-sided triangle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Torlach Rush_, Oct 06 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014873 | OEIS-A338023

Primes q for which in the q-cyclotomic field the 2-dimension of the group of circular units over the group of totally positive circular units is not maximal.

Conjecture: a prime q is in this sequence if and only if A000927(A000720(q)) is even. - _Arkadiusz Wesolowski_, Apr 09 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kim Reece_, Oct 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014874 | OEIS-A338024

Minimal number of moves for the cyclic variant of Hanoi's tower for 4 pegs and n disks, with the final peg one step away.

Conjecture: a(n) = a(n-1) + a(n-2) + a(n-3) - 2*a(n-5) for n > 11. Conjectured g.f.: x*(1+5*x+8*x^2+6*x^3+8*x^4+8*x^6-4*x^8+4*x^9-4*x^10)/((1-x)*(1+x)*(1-x-2*x^3)). - _Stefano Spezia_, Oct 07 2020 after _Paul Zimmermann_

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Zimmermann_, Oct 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014875 | OEIS-A338040

E.g.f.: Sum_{j>=0} 4^j * (exp(j*x) - 1)^j.

Conjecture: Let p be prime. The sequence obtained by reducing a(n) modulo p for n >= 1 is purely periodic with period p - 1. For example, modulo 7 the sequence becomes [4, 6, 3, 0, 1, 0, 4, 6, 3, 0, 1, 0, 4, 6, 3, 0, 1, 0, ...], with an apparent period of 6. - _Peter Bala_, May 31 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Oct 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014876 | OEIS-A338041

Draw n rays from each of two distinct points in the plane; a(n) is the number of regions thus created. See Comments for details.

The conjectures are true (see Fried link). - _Sela Fried_, Sep 21 2025 a(n) = (n^2 + 8*n - 5)/4, n odd; (n^2 + 2*n)/4, n even (conjectured). Conjectured by _Stefano Spezia_, Oct 08 2020 after _Lars Blomberg_: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lars Blomberg_, Oct 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014877 | OEIS-A338042

Draw n rays from each of two distinct points in the plane; a(n) is the number of vertices thus created. See Comments for details.

The conjectures are true (see Fried link). - _Sela Fried_, Sep 21 2025 a(n) = (n^2 + 7)/4, n odd; (n^2 - 6*n + 16)/4, n even (conjectured). Conjectured by _Stefano Spezia_, Oct 08 2020 after _Lars Blomberg_: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lars Blomberg_, Oct 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014878 | OEIS-A338043

Draw n rays from each of two distinct points in the plane; a(n) is the number of edges thus created. See Comments for details.

The conjectures are true (see Fried link). - _Sela Fried_, Sep 21 2025 a(n) = (n^2 + 4*n - 1)/2, n odd; (n^2 - 2*n + 6)/2, n even (conjectured). Conjectured by _Stefano Spezia_, Oct 08 2020 after _Lars Blomberg_: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lars Blomberg_, Oct 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014879 | OEIS-A338059

The Enots Wolley sequence A336957 with the missing prime powers interpolated.

There is a strong conjecture that A336957 consists exactly of 1, 2, and all numbers with at least two different prime factors. [The only uncertainty is whether all numbers with at least two prime factors appear.] It is conjectured that this is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Oct 17 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014880 | OEIS-A338060

a(n) = m such that A338059(m) = n, or -1 if n does not appear in A338059.

It is conjectured that A338059 is a permutation of the positive numbers. This is (conjecturally) the inverse permutation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Oct 18 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014881 | OEIS-A338073

Rearrangement of positive integers except 3 defined by A083758.

This is conjectured to be a permutation of all positive integers except 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 20 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014882 | OEIS-A338075

Diagonal terms in the expansion of (1+x*y*z)/(1-x-y-z).

Suggested by Christol's Conjecture (see reference). Conjectures from _Robert Israel_, Oct 25 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 22 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014883 | OEIS-A338076

Diagonal terms in the expansion of 1/(1-x-2*y-3*z).

Conjectures from _Robert Israel_, Oct 25 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 22 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014884 | OEIS-A338089

Minimal number of moves for the cyclic variant of Hanoi's tower for 4 pegs and n disks, with the final peg three steps away.

Conjecture: a(n) = a(n-1) + a(n-2) + a(n-3) - 2*a(n-5) for n > 9 (the same recurrence as conjectured in A292764 and A338024). - _Pontus von Brömssen_, Oct 12 2020 a(n) ~ k*r^n, k = (725 + (310451786 - 3203949*sqrt(87))^(1/3) + (310451786 + 3203949*sqrt(87))^(1/3))/348, r=constant of A289265 (closed-form by Amiram Eldar via von Brömssen conjecture). - _Bill McEachen_, Aug 19 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Zimmermann_, Oct 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014885 | OEIS-A338094

Number of ways to write 2*n + 1 as x^2 + y^2 + z^2 + w^2 with x + y a positive power of two, where x, y, z, w are nonnegative integers with x <= y and z <= w.

Conjecture: a(n) > 0 for all n > 0. Moreover, any integer m > 1987 not congruent to 0 or 6 modulo 8 can be written as x^2 + y^2 + z^2 + w^2 with x, y, z, w nonnegative integers and x + y a positive power of 4. We have verified the latter version of the conjecture for m up to 3*10^7. See also A338121 for related information, and A338095 and A338096 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014886 | OEIS-A338095

Number of ways to write 2*n + 1 as x^2 + y^2 + z^2 + w^2 with x + y + 2*z a positive power of two, where x, y, z, w are nonnegative integers with x <= y.

Conjecture: a(n) > 0 for all n >= 0. Moreover, any integer m > 10840 not congruent to 0 or 2 modulo 8 can be written as x^2 + y^2 + z^2 + w^2 with x + y + 2*z = 4^k for some positive integer k, where x, y, z, w are nonnegative integers. We have verified the latter assertion in the conjecture for m up to 5*10^6. By Theorem 1.4(i) of the author's 2019 IJNT paper, any positive integer m can be written as x^2 + y^2 + z^2 + w^2 with x, y, z, w integers such that x + y + 2*z = 4^k for some nonnegative integer k. See also A338094 and A338096 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014887 | OEIS-A338103

Positive integers not congruent to 0 or 2 modulo 8 which cannot be written as x^2 + y^2 + z^2 + w^2 with x + 2*y + 3*z a positive power of 4, where x, y, z, w are nonnegative integers.

Conjecture: 4627 is the last term of this sequence. This is equivalent to Conjecture 2 in A338096.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 10 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014888 | OEIS-A338119

Positive integers not congruent to 0 or 2 modulo 8 which cannot be written as x^2 + y^2 + z^2 + w^2 with x + y + 2*z a positive power of 4, where x, y, z, w are nonnegative integers.

Conjecture: The sequence has exactly 124 terms as listed in the b-file with 10839 the last one.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 10 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014889 | OEIS-A338121

Positive integers not congruent to 0 mod 6 which cannot be written as x^2 + y^2 + z^2 + w^2 with x + y = 4^k for some positive integer k, where x, y, z, w are nonnegative integers.

Conjecture: The sequence only has 23 terms as listed.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014890 | OEIS-A338122

Place three points evenly spaced around a circle, draw n evenly spaced rays from each of the points, a(n) is the number of regions thus created. See Comments for details.

a(n) = 2160-a(n-4)+a(n-12)+a(n-16)+a(n-60)+a(n-64)-a(n-72)-a(n-76), n>78. (conjectured) Conjectured for 1 <= n <= 800.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lars Blomberg_, Oct 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014891 | OEIS-A338123

Place three points evenly spaced around a circle, draw n evenly spaced rays from each of the points, a(n) is the number of vertices thus created. See Comments for details.

a(n) = 2160-a(n-4)+a(n-12)+a(n-16)+a(n-60)+a(n-64)-a(n-72)-a(n-76), n>78. (conjectured) Conjectured for 3 <= n <= 800.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lars Blomberg_, Oct 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014892 | OEIS-A338124

Place three points evenly spaced around a circle, draw n evenly spaced rays from each of the points, a(n) is the number of edges thus created. See Comments for details.

a(n) = 4320-a(n-4)+a(n-12)+a(n-16)+a(n-60)+a(n-64)-a(n-72)-a(n-76), n>78. (conjectured) Conjectured for 3 <= n <= 800.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lars Blomberg_, Oct 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014893 | OEIS-A338136

a(n) is the least k >= 2 such that (n+1)^k mod n^j is a perfect power > 1 for some j >= 2 with n^j < (n+1)^k.

Is there a characterization of the numbers n where a(n) = n-2? Conjecture I: if n <> 7 is a safe prime (A005385), then a(n) = n-2. Conjecture II: if n <> 22, n <> 1822 and n+1 is a safe prime, then a(n) = n-2. - _Chai Wah Wu_, Oct 13 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Oct 12 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014894 | OEIS-A338139

Number of ways to write n as x^2 + y^2 + z^2 + w^2 with x^2 + 26*y^2 - 11*x*y a power of two (including 2^0 = 1), where x, y, z, w are nonnegative integers with z <= w.

Conjecture: a(n) > 0 for all n > 0. Moreover, any positive integer n congruent to 1 or 2 modulo 4 can be written as x^2 + y^2 + z^2 + w^2 with x, y, z, w nonnegative integers such that x^2 + 26*y^2 - 11*x*y = 4^k for some nonnegative integer k. See also A337082 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 12 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014895 | OEIS-A338153

a(n) is the number of acyclic orientations of the edges of the n-prism.

Conjectured linear recurrence and g.f. confirmed by Kagey's formula. - _Ray Chandler_, Mar 10 2024 Conjectures from _Colin Barker_, Oct 13 2020: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Oct 13 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014896 | OEIS-A338162

Number of ways to write 4*n + 1 as x^2 + y^2 + z^2 + w^2 with x^2 + 7*y^2 = 2^k for some k = 0,1,2,..., where x, y, z, w are nonnegative integers with z <= w.

Conjecture: a(n) > 0 for all n >= 0. Moreover, if m > 1 has the form 2^a*(2*b+1), and either a is positive and even, or b is even, then m can be written as x^2 + y^2 + z^2 + w^2 with x^2 + 7*y^2 = 2^k for some positive integer k, where x, y, z, w are nonnegative integers. We have verified the latter assertion in the conjecture for m up to 4*10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 14 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014897 | OEIS-A338169

A list of all finite sequences of distinct positive integers sorted by their maximum value, then lexicographically.

Conjecture: the sum of the sum of all sequences with a maximum value of n is given by A281912(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Oct 14 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014898 | OEIS-A338184

G.f. A(x) satisfies: 1 = Sum_{n>=0} (-1)^n * x^(n*(n+1)/2) * A(x)^(n+1) /(Product_{k=1..n+1} 1 + x^k*A(x)).

For n>0, a(n) = 2 (mod 4) iff n is a generalized pentagonal number (A001318), and a(n) = 0 (mod 4) elsewhere (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 16 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014899 | OEIS-A338186

Expansion of g.f. 2*(1-3*x-6*x^2)/((1-x)^2*(1-9*x)).

The locally small terms 4^k in A322469 occur at the positions a(k) (for k = 0..9, and probably in general; cf. conjectures in A322469).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Georg Fischer_, Oct 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014900 | OEIS-A338193

E.g.f. A(x) satisfies: A(x) = 1 + Integral (x/A(x))' / (x/A(x)^2)' dx.

Conjecture: a(n) = f(0, n-1) for n > 0 with a(0) = 1 where f(n, m) = f(n-1, m) + m*(f(n, m-1) + f(n+1, m-1)) for n > 0, m > 0 with f(0, m) = f(0, m-1) + m*f(1, m-1) for m > 0, f(n, 0) = 1 for n >= 0. - _Mikhail Kurkov_, Oct 26 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Oct 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014901 | OEIS-A338244

Triangle read by rows: T(n,k) is the coefficient of x^k in the ZZ polynomial of the hexagonal graphene flake O(3,4,n).

T(n,k) = Sum_{i=0..12} binomial(k+i,k)*(binomial(n,k+i)*binomial(12,k+i) + 18*binomial(n+1,k+i)*binomial(10,k+i-2) + 84*binomial(n+2,k+i)*binomial(8,k+i-4) + 126*binomial(n+3,k+i)*binomial(6,k+i-6) + 57*binomial(n+4,k+i)*binomial(4,k+i-8) + 4*binomial(n+5,k+i)*binomial(2,k+i-10) + Sum_{h=0..1} (4*binomial(n+1+h,k+i)*binomial(9,k+i-3) + 24*binomial(n+2+h,k+i)*binomial(7,k+i-5) + 36*binomial(n+3+h,k+i)*binomial(5,k+i-7) + 14*binomial(n+4+h,k+i)*binomial(3,k+i-9)) + Sum_{s=0..2} Sum_{h=0..2} binomial(2,s)*binomial(2,h)*binomial(n+2+s+h,k+i)*binomial(6-2*s,k+i-6-2*s)) (conjectured, explicitly confirmed for n=1..1000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henryk A. Witek_, Oct 18 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014902 | OEIS-A338259

Triangle read by rows: T(n,k) is the coefficient of (1+x)^k in the ZZ polynomial of the hexagonal graphene flake O(3,4,n).

T(n,k) = binomial(n,k)*binomial(12,k) + 18*binomial(n+1,k)*binomial(10,k-2) + 84*binomial(n+2,k)*binomial(8,k-4) + 126*binomial(n+3,k)*binomial(6,k-6) + 57*binomial(n+4,k)*binomial(4,k-8) + 4*binomial(n+5,k)*binomial(2,k-10) + Sum_{h=0..1} (4*binomial(n+1+h,k)*binomial(9,k-3) + 24*binomial(n+2+h,k)*binomial(7,k-5) + 36*binomial(n+3+h,k)*binomial(5,k-7) + 14*binomial(n+4+h,k)*binomial(3,k-9)) + Sum_{s=0..2} Sum_{h=0..2} binomial(2,s)*binomial(2,h)*binomial(n+2+s+h,k)*binomial(6-2*s,k-6-2*s) (conjectured, explicitly confirmed for n=1..1000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Henryk A. Witek_, Oct 19 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014903 | OEIS-A338263

Number of ways to write 8*n+7 as 2*w^2 + x^2 + y^2 + z^2 with w, x, y, z nonnegative integers such that w*(3*x+7*y+10*z) is a square and also one of w, x, y, z is a square.

Conjecture 1: Each nonnegative integer can be written as 2*w^2 + x^2 + y^2 + z^2 with w, x, y, z nonnegative integers such that w*(3*x+7*y+10*z) is a square and one of w, x, y, z is also a square. As all the nonnegative integers not of the form 4^k*(8*m+7) (k>=0, m>=0) can be written as 2*0^2 + x^2 + y^2 + z^2 with x, y, z integers, Conjecture 1 has the following equivalent version: a(n) > 0 for all n = 0,1,... Conjecture 2: If (a,b) is among the ordered pairs (1,2), (1,3), (2,4), (2,5), (2,8), (2,24), (6,8), (6,32), (9,12) and (18,24), then each n = 0,1,... can be written as 2*w^2 + x^2 + y^2 + z^2 with w, x, y, z nonnegative integers such that w*(a*x+b*y) is a square.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014904 | OEIS-A338272

Number of ways to write 4*n + 1 as x^2 + y^2 + z^2 + w^2 (0 <= x <= y and 0 <= z <= w) such that x*y + 32*z*w is a square.

Conjecture: Any positive integer congruent to 1 modulo 4 can be written as x^2 + y^2 + z^2 + w^2 with x, y, z, w nonnegative integers such that a*x*y + b*z*w is a square, provided that (a,b) is among the following ordered pairs: (1,-2), (1,2), (1,10), (1,32), (2,-1), (2,10), (2,14), (2,16), (2,36), (3,4), (4,-2), (4,18), (6,18), (8,9), (9,10), (16,-2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 19 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014905 | OEIS-A338276

a(n) is the number of odd terms in the n-th column of A334016.

Conjecture: a(2^n - 1) = 2^n for n > 0. - _Peter Kagey_, Oct 22 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Oct 20 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014906 | OEIS-A338278

G.f.: Sum_{n>=0} (Product_{k=1..n} 1 + (1+x)^k) / 3^(n+1).

a(n) = 3 (mod 9) for n >= 1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 20 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014907 | OEIS-A338309

a(1) = 4; thereafter a(n) is the least new semiprime such that a(n-1) + a(n) is a semiprime.

It is conjectured that every semiprime will appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, Oct 22 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014908 | OEIS-A338338

Lexicographically earliest infinite sequence of distinct positive numbers such that for any prime p, any run of consecutive multiples of p has length exactly 3.

Conjecture: This is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014909 | OEIS-A338347

a(n) = index k such that A338338(k) = n, or -1 if n does not appear in A338338.

There is a strong conjecture that A338338 is a permutation of the positive integers. If that is true, this is the inverse permutation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014910 | OEIS-A338378

a(1) = 4. a(n) is the smallest semiprime number, which is not an earlier term, for which a(n - 1) + a(n) is a brilliant semiprime number (A078972).

Conjecture: this sequence is a permutation of the semiprimes. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius A. Burtea_, Oct 26 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014911 | OEIS-A338421

Place four points evenly spaced on a circle, draw n evenly spaced rays from each of the points, a(n) is the number of regions thus created. See Comments for details.

Conjectured for 3 <= n <= 642.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lars Blomberg_, Oct 26 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014912 | OEIS-A338422

Place four points evenly spaced on a circle, draw n evenly spaced rays from each of the points, a(n) is the number of vertices thus created. See Comments for details.

Conjectured for 3 <= n <= 642.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lars Blomberg_, Oct 26 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014913 | OEIS-A338423

Place four points evenly spaced on a circle, draw n evenly spaced rays from each of the points, a(n) is the number of edges thus created. See Comments for details.

Conjectured for 3 <= n <= 642.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lars Blomberg_, Oct 26 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014914 | OEIS-A338441

Lexicographically earliest sequence of distinct squarefree numbers such that for any prime p, any run of consecutive multiples of p has length exactly 3.

Conjecture: this is a permutation of the squarefree numbers (A005117).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Oct 28 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014915 | OEIS-A338444

Lexicographically earliest sequence of distinct squarefree numbers such that for any prime p, any run of consecutive multiples of p has length exactly 2.

Conjecture: this sequence is a permutation of the squarefree numbers (A005117).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Oct 28 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014916 | OEIS-A338451

a(n) is the least positive number that does not appear among the first n terms of A338338.

Conjecture: this sequence is unbounded (and A338338 is a permutation of the positive numbers).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Oct 28 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014917 | OEIS-A338472

(1 + Sum_{k(even)=2..p-1} 2*k^(p-1))/p as p runs through the odd primes.

Conjecture: (1 + Sum_{k(even)=2..p-1} 2*k^(p-1))/p is an integer iff p is an odd prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Davide Rotondo_, Oct 29 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014918 | OEIS-A338486

Numbers n whose symmetric representation of sigma(n) consists of 3 regions with maximum width 2.

Conjecture: The combined extent of these sections in sigma(b(k)) is 2*3^(k-1) - 1 = A048473(k-1), k>=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Oct 30 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014919 | OEIS-A338488

Numbers n whose symmetric representation of sigma(n) has a maximum width of 2 that occurs exactly once (at the diagonal).

Conjecture: For every odd number 2k-1 there is an even number n in this sequence whose symmetric representation of sigma(n) has 2k-1 regions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Oct 30 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014920 | OEIS-A338489

Let t be the closest triangular number to n! (in case n=2, the only case where we have a tie, take the larger t); then a(n) = n! - t.

It is conjectured that 0! = 1, 1! = 1, 3! = 6 and 5! = 120 are the only numbers that are both factorial (A000142) and triangular (A000217) numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ruediger Jehn_, Nov 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014921 | OEIS-A338513

a(n) is the number of Chvátal-satisfying spurious graphical n-sequences.

Conjectures from Bauer et al.: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Nov 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014922 | OEIS-A338566

Primes p such that (p*q) mod r is prime, where q and r are the next primes after p.

Note that p*q == (r-p)*(r-q) (mod r). As soon as the prime gap grows slow enough, for all large enough p we have (p*q) mod r = (r-p)*(r-q), which is composite, implying finiteness of this sequence. In particular, finiteness would follow from Cramer's conjecture. - _Max Alekseyev_, Nov 09 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Nov 02 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014923 | OEIS-A338567

Primes p such that (q*r) mod p is prime, where q and r are the next primes after p.

Note that q*r == (q-p)*(r-p) (mod p). As soon as the prime gap grows slow enough, for all large enough p we have (q*r) mod p = (q-p)*(r-p), which is composite, implying finiteness of this sequence. In particular, finiteness would follow from Cramer's conjecture. - _Max Alekseyev_, Nov 09 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Nov 02 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014924 | OEIS-A338577

Primes p such that A013632(p)*A105161(p) > p.

Sequence is finite if Cramér's conjecture is true. - _Chai Wah Wu_, Nov 03 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Nov 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014925 | OEIS-A338578

Primes p such that (r-p)*(r-q) > r, where q and r are the next two primes.

As soon as the prime gap grows slow enough, for all large enough p we have (r-p)*(r-q) <= r, implying finiteness of this sequence. In particular, finiteness would follow from Cramer's conjecture. - _Max Alekseyev_, Nov 09 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Nov 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014926 | OEIS-A338613

Numbers given by a(n) = 1 + floor(c^(n^1.5)) where c=2.2679962677... is the constant defined at A338837.

Assuming Cramer's conjecture on largest prime gaps, it can be proved that there exists at least one constant 'c' such that all a(n) are primes for n as large as required. The constant giving the smallest growth rate is c=2.267996267706724247328553280725371774527042254 4...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Montaron_, Nov 03 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014927 | OEIS-A338618

Lexicographically earliest sequence of distinct positive integers such that three consecutive terms are never pairwise coprime.

Conjecture: this sequence is a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Nov 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014928 | OEIS-A338619

Lexicographically earliest sequence of distinct positive terms such that among three consecutive terms there is exactly one pair of terms that are not coprime.

Conjecture: this sequence is a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Nov 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014929 | OEIS-A338632

G.f. A(x) satisfies: 1 = A(x) - x/(A(x) - 3*x/(A(x) - 5*x/(A(x) - 7*x/(A(x) - 9*x/(A(x) - ...))))), a continued fraction relation.

a(n) = 2 (mod 4) for n > 1 (conjecture). For n > 0, a(n) = 1 (mod 3) iff n = A191107(k) for some k >= 1 (conjecture). For n > 0, a(n) = 2 (mod 3) iff n = A186776(k) for some k >= 2 where A186776 is the Stanley sequence S(0,2) (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014930 | OEIS-A338633

G.f. A(x) satisfies: 1 = A(x) - x/(A(x) - 2^3*x/(A(x) - 3^3*x/(A(x) - 4^3*x/(A(x) - 5^3*x/(A(x) - 6^3*x/(A(x) - ...))))), a continued fraction relation.

For n > 0, a(n) is odd iff n is a power of 2 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014931 | OEIS-A338634

G.f. A(x) satisfies: 1 = A(x) - x/(A(x) - 2^4*x/(A(x) - 3^4*x/(A(x) - 4^4*x/(A(x) - 5^4*x/(A(x) - 6^4*x/(A(x) - ...))))), a continued fraction relation.

For n > 0, a(n) is odd iff n is a power of 2 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014932 | OEIS-A338635

G.f. A(x) satisfies: 1 = A(x) - x/(A(x) - 3*x/(A(x) - 6*x/(A(x) - 10*x/(A(x) - 15*x/(A(x) - 21*x/(A(x) - 28*x/(A(x) - ... - (n*(n+1)/2)*x/(A(x) - ...)))))))), a continued fraction relation.

For n>0, a(n) is odd iff n = A118113(k) for some k >= 1, where A118113(k) = 2*Fibbinary(k) + 1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014933 | OEIS-A338636

G.f. A(x) satisfies: 1 = A(x) - x/(A(x) - 3^2*x/(A(x) - 5^2*x/(A(x) - 7^2*x/(A(x) - 9^2*x/(A(x) - ...))))), a continued fraction relation.

a(n) = 0 (mod 8) for n > 1 (conjecture). For n > 0, a(n) = 1 (mod 3) iff n = A191107(k) for some k >= 1 (conjecture). For n > 0, a(n) = 2 (mod 3) iff n = A186776(k) for some k >= 2 where A186776 is the Stanley sequence S(0,2) (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 04 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014934 | OEIS-A338686

Number of ways to write n as x^5 + y^2 + [z^2/7], where x,y,z are integers with x >= 0, y >= 1 and z >= 2, and [.] is the floor function.

Conjecture: a(n) > 0 for all n > 0. See also A338687 for a similar conjecture. Conjecture verified up to 2*10^9. - _Giovanni Resta_, Apr 28 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 23 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014935 | OEIS-A338687

Number of ways to write n as x^4 + y^2 + floor(z^2/7), where x,y,z are integers with x >= 0, y >= 1 and z >= 2.

Conjecture: a(n) > 0 for all n > 0. See also A338686 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 23 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014936 | OEIS-A338696

Number of ways to write n as x^3 + y^2 + z*(3*z+2), where x and y are nonnegative integers, and z is an integer.

Conjecture: a(n) > 0 except for n = 19. As z*(3*z+2) = floor((3*z+1)^2/3) and 19 = 0^3 + 4^2 + floor(3^2/3), the conjecture implies that each n = 0,1,... can be written as x^3 + y^2 + floor(z^2/3) with x,y,z nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014937 | OEIS-A338722

Row sums in triangle A338721.

Conjecture: Equals partial sums of A335616.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 30 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014938 | OEIS-A338751

Primes p such that (p*s) mod q and (p*s) mod r are a pair of twin primes, where q,r,s are the next primes after p.

Let g1 = q-p, g2 = r-q, g3 = s-r be the gaps between the consecutive primes p,q,r,s. Then p*s = (q-g1)*(q+g2+g3) == -g1*(g2+g3) (mod q) and similarly p*s == -(g1+g2)*g3 (mod r). According to Cramer's conjecture, g1*(g2+g3) < q and (g1+g2)*g3 < r for all but finitely many p: the only exception in this sequence appears to be p=3. When g1*(g2+g3) < q and (g1+g2)*g3 < r we have ((p*s) mod r) - ((p*s) mod q) = (r - (g1+g2)*g3) - (q - g1*(g2+g3)) = g2*(g1-g3+1). This is 2 (allowing the possibility of twin primes) if g2 = 2 and g3 = g1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Nov 06 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014939 | OEIS-A338757

Number of splitting-simple groups of order n; number of nontrivial groups of order n that are not semidirect products of proper subgroups.

Conjecture: a(n) = 0 for squarefree n which is not a prime. The conjecture that a(n) = 0 for nonprime squarefree n is true. Proof: It is known that every group G of squarefree order is supersolvable; hence G contains a normal series with prime cyclic factors. Since every Sylow subgroup of G is prime cyclic, these cyclic factors are isomorphic to the Sylow subgroups of G. Let P be one such factor; then for an appropriate M in G, P = G/M, where |G| = |P|*|M|. By the Schur-Zassenhaus theorem, G is a semidirect product of M and P, and a(n) = 0 when n is squarefree. - _Miles Englezou_, Oct 24 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Nov 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014940 | OEIS-A338790

a(n) = rad(n)^2 - sigma(n), where rad(n) is the squarefree kernel of n (A007947) and sigma(n) is the sum of divisors of n (A000203).

It is conjectured that only 1 and 1782 satisfy a(x) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Nov 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014941 | OEIS-A338857

With S(n,k) = Sum_{n<=j<=k} 1/(2*j+1), a(n)=k+1 such that S(n,k-1) < 1 <= S(n,k) for n>=0 and a(0)=1.

Conjecture: a(n) = ceiling(n*exp(2)+(exp(2)+exp(-2))/(24*n)), verified for n<=3000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerhard Kirchner_, Nov 12 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014942 | OEIS-A338861

a(n) is the largest area of a rectangle which can be dissected into n squares with integer sides s_i, i = 1 .. n, and gcd(s_1,...,s_n) = 1.

Due to the potential counterexamples to the minimal squaring conjecture (see MathOverflow link), terms after a(19) have to be considered only as lower bounds: a(20) >= 876696755, a(21) >= 2735106696. - _Hugo Pfoertner_, Nov 17 2020, Apr 02 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rainer Rosenthal_, Nov 12 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014943 | OEIS-A338869

Shortest most frequent distance among first n primes.

Conjecture: Shortest most frequent distance among first n primes is a primorial number (A002110) for n>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andres Cicuttin_, Nov 13 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014944 | OEIS-A338900

Difference between the two prime indices of the n-th squarefree semiprime.

In 2020 Gus Wiseman conjectured that this sequence is an anti-run, i.e., that there are no adjacent equal parts, and verified this up to n = 10^6. This conjecture was proved by Yongkuo Chen in 2026; see the Chen link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Nov 16 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014945 | OEIS-A338926

Decimal expansion of Stechkin's constant.

Let S_m(k, q) = Sum_{j=0..q-1} e(k*j^m/q), where q>=1, m>=2, and k are integers, and e(t) = exp(2*Pi*i*t) (i is the imaginary unit). This constant is A = sup_{m,q} max_{gcd(k,q)=1} |S_m(k, q)|/q^(1-1/m). Stechkin (1975) conjectured and Shparlinskii (1991) proved that A is finite, and Banks and Shparlinski (2016) determined its value.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Nov 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014946 | OEIS-A338942

Lexicographically earliest sequence of distinct positive terms such that a(n) is present in 3*a(n+1).

Conjecture: For n >= 222, a(n) = 100*a(n-16). - _Jason Yuen_, Dec 22 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Carole Dubois_, Nov 17 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014947 | OEIS-A338968

a(n) is the largest prime whose decimal expansion consists of the concatenation of a 1-digit prime, a 2-digit prime, a 3-digit prime, ..., and an n-digit prime.

It is a plausible conjecture that a(n) always exists and begins with 7. Conjecture: for n >= 3, a(n) is the concatenation of the largest k-digit primes with 1 <= k <= n-1: A003618(1)/A003618(2)/.../A003618(n-1) but the last concatenated prime with n digits is always < A003618(n). This conjecture has been checked by _Daniel Suteu_ until a(360), a prime with 64980 digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Dec 21 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014948 | OEIS-A338985

a(n) is the first prime p such that the sum of 2*n consecutive primes starting at p is an oblong number, or 0 if there is no such p.

Conjecture: all a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Dec 20 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014949 | OEIS-A338998

Composite numbers k such that phi(k) divides p*(k - 1) for some prime factor p of k - 1.

The "Lehmer numbers" above refers to composite 1-Lehmer numbers, that is, numbers n that would satisfy the equation y * phi(n) = n-1, for some integer y > 1. Lehmer conjectured that no such numbers exist. See the assorted Web-links. - _Antti Karttunen_, Dec 26 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tomohiro Yamada_, Nov 18 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014950 | OEIS-A339053

a(n) = least k such that the first n-block in A339051 occurs in A339052 beginning at the k-th term.

The sequence begins with 4 3's, 6 7's, 62 45's, 116 117's, 1104 799's, followed by 2091. Conjecture: the sequence includes infinitely many distinct numbers; if so, then every finite block in A339051 occurs infinitely many times in A339052.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014951 | OEIS-A339054

a(n) = least k such that the first n-block in A339052 occurs in A339051 beginning at the k-th term.

The sequence begins with 2 3's, 1 4, 7 8's, 6 12's, 28 29's, 47 46's, 117 118's, 116 190's, 494 495's, 493 800's, followed by 2092. Conjecture: the sequence includes infinitely many distinct numbers; if so, then every finite block in A339052 occurs infinitely many times in A339051.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014952 | OEIS-A339061

Least integer j such that H(k+j)>=n+1, where k is the least integer to satisfy H(k)>=n, and H(k) is the sum of the first k terms of the harmonic series.

Conjecture: a(n) = floor(1/2 + e^(n-gamma+1)) - floor(1/2 + e^(n-gamma)) for n > 1 where e is Euler's number and gamma is the Euler-Mascheroni constant. - _Adam Hugill_, Nov 06 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthew J. Bloomfield_, Dec 21 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014953 | OEIS-A339148

Insulated primes (see Comments for definition).

a(n) ~ c*n^(1+epsilon) (conjectured for some constant c,epsilon as n->oo).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Abhimanyu Kumar_, Nov 25 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014954 | OEIS-A339185

a(n) is the least prime p such that the sum of n consecutive primes starting with p has exactly n prime factors, counted with multiplicity, or 0 if no such p exists.

Conjecture: Such p exists for every n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Nov 26 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014955 | OEIS-A339210

a(n) = Sum_{k=0..(n+1)/2} A010815(n*k).

Conjecture: a(p) = 2 for prime p > 5. Tested for p up to 100000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gevorg Hmayakyan_, Nov 26 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014956 | OEIS-A339247

The primes that yield twice a prime when each bit of their binary expansion is inverted.

Conjecture: Each of the primes can be created by applying this transformation T(p) to at least one other prime. T(11)=2, T(549755813881)=3, T(53)=5, T(17)=7, T(41)=11.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bob Andriesse_, Nov 28 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014957 | OEIS-A339258

Triangle read by rows T(n,k), (n >= 1, k > = 1), in which row n has length A000070(n-1) and every column gives A000005, the number of divisors function.

Conjecture: the sum of row n equals A006128(n), the total number of parts in all partitions of n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Nov 29 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014958 | OEIS-A339261

Decimal expansion of the conjecturally maximum possible volume of a polyhedron with 9 vertices inscribed in the unit sphere.

Decimal expansion of the conjecturally maximum possible volume of a polyhedron with 9 vertices inscribed in the unit sphere.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 05 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014959 | OEIS-A339262

Decimal expansion of the conjecturally maximum possible volume of a polyhedron with 10 vertices inscribed in the unit sphere.

Decimal expansion of the conjecturally maximum possible volume of a polyhedron with 10 vertices inscribed in the unit sphere.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014960 | OEIS-A339263

Decimal expansion of the conjecturally maximum possible volume of a polyhedron with 11 vertices inscribed in the unit sphere.

Decimal expansion of the conjecturally maximum possible volume of a polyhedron with 11 vertices inscribed in the unit sphere.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 07 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014961 | OEIS-A339268

The smallest prime that becomes 2 * prime(n), when all the bits in its binary expansion are inverted, or -1 if no such prime exists.

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bob Andriesse_, Nov 29 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014962 | OEIS-A339269

a(n) is the least number that is the product of n primes (not necessarily distinct) and is the sum of n consecutive primes, or 0 if there are none.

Conjecture: a(n) > 0 for every n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Nov 29 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014963 | OEIS-A339387

a(n) = Sum_{k=1..n} (lcm(n,k)/gcd(n,k) mod k).

Conjecture: If a(n) = A339384(n), then n is squarefree. This appears to be true for at least the first 2000 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sebastian Karlsson_, Dec 02 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014964 | OEIS-A339451

Gray-code-like sequence in which, at each step, the least significant bit that has never been toggled from the previous value, is toggled.

Conjectured connections: the position of the bit that is toggled to derive a(n) from a(n-1) is A215020(n); the sequence of absolute differences of this sequence is A182105; there is some underlying connection to the "skew binary" counting system.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Allan C. Wechsler_, Dec 05 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014965 | OEIS-A339457

Decimal expansion of the smallest positive number d such that numbers of the sequence floor(2^(n^d)) are distinct primes for all n>=1.

Assuming Cramer's conjecture on prime gaps, it can be proved that there exists at least one constant d such that all floor(2^(n^d)) are primes for n>=1 as large as required. The constant giving the smallest growth rate is d=1.503928524069520633527689067897583199190738...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Montaron_, Dec 06 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014966 | OEIS-A339459

Prime numbers a(n) = floor(2^(n^d)) for all n>=1 where d=1.5039285240... is the constant defined at A339457.

Assuming Cramer's conjecture on prime gaps is true, it can be proved that there exists at least one constant d such that all terms of the sequence are primes. The constant giving the smallest growth rate is d=1.503928524069520633527689067897583199190738...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Montaron_, Dec 06 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014967 | OEIS-A339546

Number of at least 3-connected planar triangulations on n vertices such that the minimum valence of any vertex in the mesh is maximized and the number of vertices with this minimum valence is minimized.

It is conjectured that for a polyhedron with maximum possible volume inscribed in a sphere the stated condition is necessary. All known polyhedra with conjecturally maximum volume satisfy this condition. For n <= 15 the given condition determines a unique mesh topology for each n, which coincides with the proved optimal solutions for n <= 8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 08 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014968 | OEIS-A339570

Denote the van der Corput sequence of fractions 1/2, 1/4, 3/4, 1/8, 5/8, 3/8, 7/8, 1/16, ... (A030101/A062383) by v(n), n >= 1. Then a(n) = denominator of v(A014486(n)).

The initial values suggest the conjecture that this sequence consists exactly of Catalan(k) copies of 4^k for k >= 1. _Hugo Pfoertner_ tested this conjecture with the PARI program given below. The first column is A014137, the partial sums of the Catalan numbers, which is strong support for the conjecture. The conjecture has now been proved by _Raghavendra Tripathi_ - see link. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014969 | OEIS-A339579

a(n) = least nonnegative integer k such that n*2^k - 1 is composite.

Conjectured to grow without limit.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 24 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014970 | OEIS-A339599

Quadratically perfect numbers: numbers k such that (sigma(k) - 2k)^2 = sigma(k).

Lescot also confirms that there exists no odd integer between 5 and 10^6 in this sequence and conjectures that there exists no further odd term in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tomohiro Yamada_, Dec 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014971 | OEIS-A339602

a(n) = (a(n-2) XOR A030101(a(n-1))) + 1, a(0) = 0, a(1) = 1.

Conjecture: Let p be an odd number, then a(n) = p will be more frequently found in this sequence than a(n) = p+1 (tested for n = 0..10^7 with primes > 2, but seems to be true for all odd too).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Dec 09 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014972 | OEIS-A339607

a(1) = 1, a(n) is the least m not already in the sequence that contains the binary expansion of the binary weight of a(n-1) anywhere within its own binary expansion.

Conjecture: permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Dec 16 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014973 | OEIS-A339621

Sum of Fibonacci divisors of n^2 + 1.

Conjecture: If the sum of the Fibonacci divisors of m^2 + 1 is a Fibonacci number, then this number belongs to the sequence A001906(n) = F(2n) where F(n) is the Fibonacci sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Dec 10 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014974 | OEIS-A339624

Perfect powers p^k, k >= 2 of palindromic primes p when p^k is not a palindrome.

G. J. Simmons conjectured there are no palindromes of form n^k for k >= 5 (and n > 1) (see Simmons p. 98). According to this conjecture, these perfect powers are terms: {2^k, k>=4}, {3^k, k>=3}, {5^k, k>=2}, {7^k, k=2 and k>=4}, {11^k, k>=5}, {101^k, k>= 5}, {131^k, k>=2}, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Dec 10 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014975 | OEIS-A339636

Counterexamples to a conjecture about integers representable as the quotient of two Cantor numbers (A005823).

Counterexamples to a conjecture about integers representable as the quotient of two Cantor numbers (A005823). Conjecture: every number of the form 23*3^(4k+3) - 20 is not representable.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Dec 11 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014976 | OEIS-A339652

a(n) is the least number k such that 2*k+1 has exactly n expressions as p+2*q where p and q are primes, or 0 if there is no such k.

Conjecture: all a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Dec 13 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014977 | OEIS-A339659

Irregular triangle read by rows where T(n,k) is the number of graphical partitions of 2n into k parts, 0 <= k <= 2n.

Conjecture: The column sums 1, 0, 1, 2, 7, 20, 67, ... are given by A304787.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Dec 18 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014978 | OEIS-A339669

Number of Fibonacci divisors of Lucas(n)^2 + 1.

Particular attention must be paid to the regularity properties of the number of divisors of Lucas(n)^2 + 1 observed for n < 156, when a(n) = 1 or 2. From this observation, we propose two conjectures verified for n < 156. Conjecture 1: a(6*n+3) = 1. Conjecture 2: a(6*n+1) = a(6*n+5) = 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Dec 12 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014979 | OEIS-A339676

Nonpalindromic numbers that are products of repunits.

G. J. Simmons conjectured there are no palindromes of form n^k for k >= 5 (and n > 1) (see link, page 98). According to this conjecture, these perfect powers are terms: {11^k, k>=4}, {111^k, k>=4}, {1111^k, k>=3}, {11111^k, k>=3}, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Dec 12 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014980 | OEIS-A339709

a(n) is the least even number that has exactly n decompositions as the sum of an odd prime and a semiprime, or 0 if there is no such number.

Conjecture: a(12) = 0, but a(n) > 0 for all other n. Since this is only a conjecture, the Data stops at a(11)=92.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Dec 13 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014981 | OEIS-A339744

Numbers k such that rad(k)^2 < sigma(k), where rad(k) is the squarefree kernel of k (A007947) and sigma(k) is the sum of divisors of k (A000203).

The percentage of terms decreases as 10^k increases, and a plausible conjecture is that the asymptotic density of this sequence is 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Dec 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014982 | OEIS-A339752

Base-ten n containing a digit making its n-th appearance in the list of consecutive positive integers.

Conjectured to be "full": digit 0 (the final term) might have subsequents.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hans Havermann_, Dec 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014983 | OEIS-A339753

Base-ten n whose English number-word expression contains a letter making its n-th appearance in the list of consecutive positive integer number-words.

Conjecture: a(494) = 1001001001001998 (for the letter 'a').

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Claudio Meller_ and _Hans Havermann_, Dec 15 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014984 | OEIS-A339826

a(n) = least k such that the first n-block in A339825 occurs in A339824 beginning at the k-th term.

The sequence begins with 2 2's, 1 3, 7 7's, 6 11's, ... Conjecture: the sequence includes infinitely many distinct numbers, in which case, every finite block in A339825 occurs infinitely many times in A339824.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 19 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014985 | OEIS-A339869

Carmichael numbers k for which A053575(k) [the odd part of phi] divides k-1.

Lehmer conjectured that the equation k * phi(n) = n - 1 (with k integer) has no solutions for any composite n (i.e., when k > 1). If this sequence has no common terms with A339818, then the conjecture certainly holds.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 22 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014986 | OEIS-A339882

Fundamental positive solution y(n) of the Diophantine equation x^2 - A045339(n)*y^2 = -2, for n >= 1.

The conjecture is that all A045339 entries have solutions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Dec 22 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014987 | OEIS-A339883

Values of Euler's totient phi for A050498.

For the shown values the arithmetic progression with difference 6 has exactly 4 terms. Conjecture 1: this holds for all A050498 entries. Conjecture 2: All a(n) are divisible by 24, starting with [1, 3, 1060, 1410, 1620, 2040, 4140, 5130, 8680, 19110, 21280, 21640, 22200, 24310, 24850, 25920, 27660, 34950, 35290, 36400, 37240, 39270, 43770, 44400, 44950, 48460, ...].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jan 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014988 | OEIS-A339887

Number of factorizations of n into primes or squarefree semiprimes.

Conjecture: also the number of semistandard Young tableaux whose entries are the prime indices of n (A323437).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Dec 22 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014989 | OEIS-A339907

Odd squarefree numbers k > 1 for which the bigomega(phi(k)) <= bigomega(k-1), where bigomega gives the number of prime divisors, counted with multiplicity.

Natural numbers n > 2 that satisfy equation k * phi(n) = n - 1 (for some integer k) all occur in this sequence. Lehmer conjectured that there are no composite solutions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 21 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014990 | OEIS-A339909

Carmichael numbers k for which bigomega(phi(k)) < bigomega(k-1), where bigomega gives the number of prime divisors, counted with multiplicity.

Natural numbers n that satisfy equation k * phi(n) = n - 1, for some integer k > 1, should all occur in this sequence, if they exist at all. Lehmer conjectured that there are no such numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 22 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014991 | OEIS-A339929

a(n+1) = a(n-1-a(n)^2) + 1, starting with a(1) = a(2) = 0.

a(n) ~ (3*n)^(1/3) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rok Cestnik_, Dec 23 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014992 | OEIS-A339930

a(n+1) = a(n-2-a(n)^2) + 1, starting with a(1) = a(2) = a(3) = 0.

a(n) ~ (3*n)^(1/3) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rok Cestnik_, Dec 23 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014993 | OEIS-A339931

a(n+1) = a(n-3-a(n)^2) + 1, starting with a(1) = a(2) = a(3) = a(4) = 0.

a(n) ~ (3*n)^(1/3) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rok Cestnik_, Dec 23 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014994 | OEIS-A339932

a(n+1) = a(n-4-a(n)^2) + 1, starting with a(1) = a(2) = a(3) = a(4) = a(5) = 0.

a(n) ~ (3*n)^(1/3) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rok Cestnik_, Dec 23 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014995 | OEIS-A339946

a(n) = [x^n] 1/Legendre_P(n,(1 - x)/(1 + x)).

We conjecture that the following supercongruences hold for any prime p >= 5: a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for all positive integers n and k. For m a positive integer, define a_m(n) = [x^(m*n)] 1/Legendre_P(n,(1 - x)/(1 + x)). We conjecture that the supercongruence a_m(p) == a_m(1) (mod p^5) holds for all primes p >= 7. - _Peter Bala_, Mar 10 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Dec 23 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014996 | OEIS-A339950

Numbers k such that all k-sections of the infinite Fibonacci word A014675 have just two different run-lengths.

Conjecture: a(n) = A189378(n-1)+1 for n >= 2. - _Don Reble_, Apr 06 2021.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 26 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014997 | OEIS-A339987

The number of labeled graphs on 2n vertices that share the same degree sequence as any unrooted binary tree on 2n vertices.

Conjectured recurrence: 32*(1 + n)*(2 + n)*(1 + 2*n)*(3 + 2*n)*(5 + 2*n)*(7 + 2*n)*(9 + 2*n)*(11589 + 10844*n + 3300*n^2 + 328*n^3)*a(n) - 8*(2 + n)*(3 + 2*n)*(5 + 2*n)*(7 + 2*n)*(9 + 2*n)*(148119 + 232328*n + 129460*n^2 + 30664*n^3 + 2624*n^4)*a(n+1) - 16*(3 + n)*(5 + 2*n)*(7 + 2*n)*(9 + 2*n)*(341634 + 712135*n + 569267*n^2 + 219308*n^3 + 40852*n^4 + 2952*n^5)*a(n+2) + 8*(4 + n)*(7 + 2*n)*(9 + 2*n)*(527520 + 1057879*n + 818282*n^2 + 306380*n^3 + 55672*n^4 + 3936*n^5)*a(n+3) - 2*(5 + n)*(9 + 2*n)*(601452 + 1117119*n + 786236*n^2 + 264028*n^3 + 42472*n^4 + 2624*n^5)*a(n+4) + 3*(4 + n)*(6 + n)*(3717 + 5228*n + 2316*n^2 + 328*n^3)*a(n+5) = 0. - _Manuel Kauers_ and _Christoph Koutschan_, Mar 01 2023 Conjecture: a(n) ~ 2^(5*n - 1/2) * n^(2*n - 3/2) / (sqrt(Pi) * 3^(n-1) * exp(2*n + 21/16)), based on the recurrence by _Manuel Kauers_ and _Christoph Koutschan_. - _Vaclav Kotesovec_, Mar 07 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Atabey Kaygun_, Dec 25 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014998 | OEIS-A339991

The number of steps that n requires to reach 1 under the map: m -> m/2 if m is even, m-> m^2 - 1 if m is an odd prime, otherwise m -> m - 1. a(n) = -1 if 1 is never reached.

Conjecture: a(n) is never equal to -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Dec 25 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-014999 | OEIS-A340001

Number of ways prime(n) is a sum of five distinct primes.

Conjecture: all primes >= 43 are the sum of five distinct primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Dec 26 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015000 | OEIS-A340061

Irregular triangle read by rows T(n,k) in which row n lists n blocks, where the m-th block consists of A000041(n-m) copies of m, with n >= 1 and m >= 1.

Conjecture: all divisors of all terms of row n are also all parts of all partitions of n. The conjecture gives a correspondence between divisors and partitions (see example). It is conjectured that every section of the set of partitions of n has essentially the same correspondence. For more information see A336811.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Dec 28 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015001 | OEIS-A340063

The primes appear in their natural order and the absolute difference between two successive primes is the sum of the digits between them.

Lexicographically earliest sequence of distinct positive terms with this property. It is conjectured that the sequence is a permutation of the integers > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_, Dec 27 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015002 | OEIS-A340079

a(n) = n / gcd(n, 1+A018804(n)), where A018804(n) = Sum_{k=1..n} gcd(k, n).

It is conjectured that this is 1 iff n is 1 or a prime. See _Thomas Ordowski_'s Oct 22 2014 comment in A018804. Adamczewski provides a composite counterexample, n = 23492890653051, to the conjecture in linked paper. - _Michael De Vlieger_, Aug 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 30 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015003 | OEIS-A340088

a(n) = A091732(n) / gcd(n-1, A091732(n)), where A091732 is an infinitary analog of Euler's phi function.

Conjecture: a(n) = 1 iff n = 1 or in A050376. This is an infinitary analog of Lehmer's totient conjecture from 1932. For all i, j > 1: a(i) = a(j) => A302777(i) = A302777(j), if the above conjecture holds.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 31 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015004 | OEIS-A340115

Largest prime whose decimal expansion consists of the concatenation of a 1-digit cube, a 2-digit cube, a 3-digit cube, ..., and an n-digit cube, or 0 if there is no such prime.

We can conjecture that a(n) > 0 for all n > 1 and the terms converge to the concatenation of (c(1), c(2), c(3), ...) where c(k) is the largest k digit cube. The number of such primes between A215692(n) and a(n) is (0, 2, 2, 9, 177, 6909, 570166, ...). This is very close to what we expect given the number of concatenations of cubes of the respective length (product of 10^(k/3)-10^((k-1)/3), k=1..n) and the density of primes in that range according to the PNT. - _M. F. Hasler_, Dec 31 2020

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Dec 28 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015005 | OEIS-A340138

a(1)=0, a(n+1) = the least m not already in the sequence such that the decimal digital root r of a(n-1) is found somewhere among the decimal digits of m.

Conjecture: permutation of the nonnegative numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jan 21 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015006 | OEIS-A340162

a(1) = 7; thereafter a(n) is the smallest number k with exactly three 1-bits, not already in the sequence, for which k*a(n - 1) has exactly three 1-bits (A014311).

Conjectures from _Chai Wah Wu_, Jan 27 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius A. Burtea_, Dec 30 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015007 | OEIS-A340163

For n>=1, smallest integer k such that for all m>=k: m^(1/n)+(m+1)^(1/n) >= (2^n*m+2^(n-1)-1)^(1/n).

For k>1, a(n) <= ceiling(2^(k-3)). This sequence refers to a conjecture, which is a generalization of a Question 723. (iii) from "Collected Papers", Srinivasa Ramanujan.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrzej Kukla_, Dec 30 2020 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015008 | OEIS-A340224

a(n+1) = a(n-a(n)*(a(n)-1)/2) + 1, starting with a(1) = 0.

a(n) ~ (6*n)^(1/3) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rok Cestnik_, Jan 01 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015009 | OEIS-A340230

a(n) is the smallest number m such that numbers m, m + 1, m + 2, ..., m + n - 1 have k, 2*k, 4*k, 8*k, ..., (2^(n-1))*k divisors respectively.

Conjecture: a(7) = 213536830501.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Jan 01 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015010 | OEIS-A340234

Number of harmonious graphs with n edges and at most n vertices, allowing self-loops.

For n odd, a(n) = ceiling(n/2)^n; for n even, a(n) = ((n^2/4) + (n/2))^(n/2) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick D. Cone_, Jan 01 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015011 | OEIS-A340271

Positive integers with no digit equal to 0 or 1 such that when one iterates x -> A340270(x) starting with n, one eventually enters a cycle containing an integer greater than 9.

This was suggested by postings by Eric Angelini and Allan Wechsler to Math-Fun Mailing List, Dec 29 2020. It is an open question whether or not this sequence is complete. In fact, it is open whether or not the sequence of iterates of x -> A340270(x) starting with n always enters a cycle. There are no counterexamples up to 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _W. Edwin Clark_, Jan 02 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015012 | OEIS-A340274

Number of ways to write n as x + y + z with x, y, z positive integers such that 3*x^2*y^2 + 5*y^2*z^2 + 8*z^2*x^2 is a square.

Conjecture 1: a(n) > 0 for all n > 2. We have verified a(n) > 0 for all n = 3..10000. The conjecture holds if a(p) > 0 for every odd prime p. For any n > 0 we have a(3*n) > 0, since 3*n = n + n + n and 3 + 5 + 8 = 4^2. See also A343862 for similar conjectures. Conjecture 1 holds for all n < 2^15. Note a(1823) = 1. - _Martin Ehrenstein_, May 03 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015013 | OEIS-A340281

a(n) is the smallest prime p such that the number of distinct values of the ratio (number of nonnegative m < p such that m^k == m (mod p))/(number of nonnegative m < p such that -m^k == m (mod p)) is equal to n for some nonnegative k.

Conjecture: a(n) is the smallest prime p such that the number of distinct values of the ratio T(p, k) = (number of nonnegative m < p such that m^k == m (mod p))/(number of nonnegative m < p such that -m^k == m (mod p)) is equal to n for some 0 <= k <= floor((p + 2)/3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jan 02 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015014 | OEIS-A340347

An aware-like sequence; empty the first pit/integer and sow its content to the right, one seed/unit per pit/integer. When this is done the new sequence and the starting one share the same succession of digits.

This sequence is the lexicographically earliest one of distinct integers > 0 and is conjectured to be a permutation of the integers > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_, Jan 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015015 | OEIS-A340348

Add the two terms of the first pair of integers to produce b(1); add the two terms of the second pair of integers to produce b(2); add the two terms of the third pair of integers to produce b(3), etc. This sequence and the b(n) sequence share the same succession of digits.

This is the lexicographically earliest sequence of distinct integers > 0 with this property. The sequence is conjectured to be a permutation of the integers > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Carole Dubois_, Jan 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015016 | OEIS-A340352

Number of spanning trees of odd Aztec diamond OD_n.

R. P. Stanley conjectured that the even Aztec diamond has exactly four times as many spanning trees as the odd Aztec diamond. This conjecture was first proved by D. E. Knuth.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Jan 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015017 | OEIS-A340362

a(n) = A005940(n) - A324106(n), where A324106(n) is multiplicative with a(p^e) = A005940(p^e).

It is conjectured that A070776 gives the positions of all zeros after the initial a(1) = 0. If that holds, then for all i, j: a(i) = a(j) => A340363(i) = A340363(j).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015018 | OEIS-A340365

a(n) = A005940(n) / gcd(A005940(n), A324106(n)), where A324106(n) is multiplicative with a(p^e) = A005940(p^e).

It is conjectured that A070776 gives the positions of all ones after the initial one. If that holds, then for all i, j: a(i) = a(j) => A340363(i) = A340363(j).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015019 | OEIS-A340420

The number of steps that n requires to reach 1 under the map: m -> m/2 if m is even, m-> 3*m + 1 if m is an odd prime, otherwise m -> m - 1. a(n) = -1 if 1 is never reached.

Conjecture: a(n) is never equal to -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Jan 07 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015020 | OEIS-A340445

Number of partitions of n into 3 parts that are not all the same.

Conjecturally the same as A230059 (apart from the offset). - _R. J. Mathar_, Jan 14 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Jan 07 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015021 | OEIS-A340461

a(n) = 2*sigma(phi(n)) - n.

In 1964, A. Mąkowski and A. Schinzel conjectured that sigma(phi(n))/n >= 1/2 for all n (see links Mąkowski & Schinzel and Graeme L. Cohen), equivalent to a(n) >= 0. In 1992, K. Atanassov believed that he obtained a proof of this conjecture but his proof was valid only for certain special values of n (see link József Sándor).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Jan 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015022 | OEIS-A340462

Triangular numbers that are the hypotenuse of a primitive Pythagorean triple (PPT) such that another member of the triple is also triangular.

Conjecture: No PPT exists with all its members triangular numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Jan 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015023 | OEIS-A340487

a(n) is the least prime that is the concatenation of two n-digit primes, or 0 if there are none.

Conjecture: a(n) > 0 and the first n digits of a(n) = A003617(n). - _Chai Wah Wu_, Jan 13 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Jan 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015024 | OEIS-A340488

a(n+1) = XOR(a(n),y) for n>=1, where y is the number of m < n such that a(m) = a(n); a(1)=0.

It follows from the definition that when m appears for the first time, it is immediately followed by m, and then if m is even by m+1. A340494 gives a conjectured generating function for when a number appears for the first time. - _Rémy Sigrist_ and _N. J. A. Sloane_, Jan 10 2021
Conjectures from _Rémy Sigrist_, Jan 10 2021 (Start)
Conjecture: Let pi_i(n) denote the number of occurrences of i in the first n terms. Then pi_0(n) >= pi_i(n) for all i, and the first time pi_0(n) = m, pi_i(n) < m for i>0. - _N. J. A. Sloane_, Jan 13 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ian Hutchinson_, Jan 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015025 | OEIS-A340492

a(n) = A000041(n)*A000070(n-1), n >= 1.

Conjecture: a(n) is the size of a table of correspondence between the divisors of certain numbers and the partitions of n. The table contains A006128(n) positive integers and the rest are zeros (or empty cells). These positive integers can be interpreted in two ways as follows: For more information about the main conjecture see A336811.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jan 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015026 | OEIS-A340493

Sequence whose partial sums give A340492.

Conjecture: a(n) is the size of the n-th section of a table of correspondence between divisors and partitions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jan 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015027 | OEIS-A340497

Index where 2*n first appears in A340488.

A bisection of A340494. Note that there is a conjectured g.f. for A340494.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 11 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015028 | OEIS-A340498

Where 2^n appears in A340488 for the first time.

A subsequence of A340494 and A340497. Note that A340494 has a conjectured g.f.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 11 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015029 | OEIS-A340511

Numbers k such that there exists a group of order k which has no subgroup of order d, for some divisor d of k.

Conjecture: If k = p^a*q^b, where p, q are primes, q !== 1 (mod p), b >= ord(q,p), then k is a term of this sequence, unless k is an NSS-CLT order of the form described in MacHale-Manning, 2016, Theorem 8, Page 5. Here ord(q,p)

is the multiplicative order of q modulo p. Moreover, if k satisfies this condition, it seems that for each NCLT group of order k, the missing orders of subgroups are of the form p^a'*q^b' where either a' = a or b' = b, and a' = a if p == 1 (mod q) or a < ord(p,q). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Des MacHale_, Jan 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015030 | OEIS-A340518

Smallest order of a finite group with a commutator subgroup of order n.

The above conjecture is true. For even n, a(n) is even by Lagrange's theorem. For odd n, it follows from the fact that every dihedral group D_{2n} has a commutator subgroup of order n when n is odd; as no group of odd order is perfect, 2*n is the smallest possible order that such a commutator subgroup can be contained in. (For an extended proof see the Miles Englezou link.) - _Miles Englezou_, Mar 08 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Des MacHale_, Jan 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015031 | OEIS-A340530

Irregular triangle read by rows T(n,k) in which row n has length is A000070(n-1) and every column k is A006218, (n >= 1, k >= 1).

The sum of row n equals A284870(n), the total number of parts in all partitions of all positive integers <= n. It is conjectured that this property is due to the correspondence between divisors and partitions. For more information see A336811.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jan 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015032 | OEIS-A340539

a(n) is the least prime that is the concatenation of two n-digit primes, and such that the concatenation of the same primes in the other order is also prime or 0 if no such prime exists.

Conjecture: a(n) > 0 and for n > 1 the first n digits of a(n) = A003617(n). - _Chai Wah Wu_, Jan 13 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Jan 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015033 | OEIS-A340581

Irregular triangle read by rows in which row n has length A014153(n-1) and every column k lists the positive integers A000027, n >= 1, k >= 1.

Conjecture: all divisors of all terms in row n are also all parts of all partitions of all positive integers <= n. The conjecture is in accordance with the conjectures in A336811 and in A176206. A336811 contains the most elementary conjecture about the correspondence divisors/partitions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jan 14 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015034 | OEIS-A340630

Fill an n X n square with nonnegative integers so that all n^2 von Neumann neighborhoods have distinct sums; a(n) is smallest possible sum of the entries.

Conjecture: a(n) = ceiling((n^4 - n^2 + 14) / 10), matching the lower bound, for all n > n_0 for some constant n_0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo van der Sanden_, Jan 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015035 | OEIS-A340672

Number of repeated applications of the '4x+-1' operation needed to reach 1 from n, or -1 if 1 is never reached.

We can make a conjecture similar to that of Collatz to claim that the process will always reach 1. However, this is not proven yet.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Md. Towhidul Islam_, Jan 15 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015036 | OEIS-A340700

Lower of a pair of adjacent perfect powers, both with exponents > 2.

It is conjectured that the intersection of A340700 and A340701 is empty, i.e., that no 3 immediately consecutive perfect powers with all exponents > 2 (A076467) exist. No counterexample < 3.4*10^30 was found.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jan 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015037 | OEIS-A340706

Difference between upper and lower member of a pair of adjacent perfect powers A340700 and A340701, both with exponents > 2.

The differences are expected to be bounded below by the Lang-Waldschmidt conjecture (see Waldschmidt 2013, p. 6, Conjecture 6).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jan 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015038 | OEIS-A340708

Maximum denominator of resistances obtained by an electrical network with n unit resistors.

The author conjectures that this asymmetry will increase with n, and eventually the maxima will differ. Conjecture: a(19) = 22233, a(20) = 39918. It would be very desirable to know at which value of n > 18 the maximum values of numerators and denominators differ for the first time. - _Hugo Pfoertner_, Apr 19 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rainer Rosenthal_, Jan 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015039 | OEIS-A340726

Maximum power V_s*A_s consumed by an electrical network with n unit resistors and input voltage V_s and current A_s constrained to be exact integers which are coprime, and such that all currents between nodes are integers.

Take the set SetA337517(n) of resistances, counted by A337517. For each resistance R multiply numerator and denominator. Conjecture: a(n) is the maximum of all these products. The reason is that common factors of V_s and A_s are quite rare (see the beautiful exceptional example with 21 resistors).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rainer Rosenthal_, Jan 17 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015040 | OEIS-A340738

Denominator of a sequence of fractions converging to e.

The convergence is conjectured. Adamczewski includes a proof abstract regarding the conjecture immediately above in linked article. - _Michael De Vlieger_, Aug 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary Detlefs_, Jan 18 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015041 | OEIS-A340739

The stopping time sequence for the 3x+1 function, restricted to its range and adjusted. Each term is the number of iterations of the function until it decreases.

Observations, Conjectures: Conjecture: Ultimately every positive integer appears in the stopping time sequence. (verified up to 100, examples: a(6802394)=160, a(31229269)=161) And each positive integer is in the sequence an infinite number of times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sam E. Speed_, Jan 18 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015042 | OEIS-A340757

Counterexamples to a conjecture of Ramanujan about congruences related to the partition function.

Counterexamples to a conjecture of Ramanujan about congruences related to the partition function. For b in 5,7,11, and all integers n,e >= 1, Ramanujan conjectured that if (24*n-1) is divisible by b^e, the partition function p(n) = A000041(n) is also divisible by b^e. Chowla found the first counterexample a(1) = 243. Watson showed the conjecture holds for b=5, and Atkin showed it holds for b=11. Watson showed p(n) is divisible by 7^floor((d+2)/2) when 24n-1 is divisible by 7^d, so that exceptions here are restricted to 24n-1 == 0 (mod 7^3), which is n == 243 (mod 7^3). See A340957 for the converse, those n == 243 (mod 7^3) where the conjecture does hold.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Washington Bomfim_, Jan 19 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015043 | OEIS-A340797

Integers whose number of divisors that are Brazilian sets a new record.

Observation: the 58 consecutive highly composite numbers from A002182(12) = 240 to A002182(69) = 2095133040 (maybe more, according to conjectured terms) are also terms of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Jan 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015044 | OEIS-A340801

a(n) is the image of n under the map f defined as f(n) = n^2 - 2 if n is an odd prime, f(n) = n/2 if n is even, and f(n) = n - 1 otherwise.

Conjecture 1: Iterating map f on an integer n (n > 1) results in a different integer, or f^i(n) != f^j(n) if i != j, where f^i(n) and f^j(n) are the i-th and j-th iterations of map f on n respectively. Conjecture 2: An integer n eventually reaches 1 when map f is applied to n repeatedly.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Jan 21 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015045 | OEIS-A340862

Number of times the number n turns up in pseudo-Fibonacci sequences starting with [k, 1] (with k >= 1), excluding the starting terms.

Conjecture: for n > 2, a(n) appears to be equal to 1 + A067148(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robby Goetschalckx_, Jan 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015046 | OEIS-A340880

Irregular table: n-th row polynomial given by the formal power series expansion of Sum_{k >= 0} (1 + q)^(n*k + k*(k+1)/2)* Product_{j = 1..k} (1 - (1 + q)^j), n >= 1.

Conjecture: For fixed integers q and N, the sequence (R(n,q) mod N)n>=1 is eventually periodic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Feb 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015047 | OEIS-A340881

Row sums of A340880.

Conjectures: 1) For prime p, the sequence taken modulo p is purely periodic with minimum period dividing 2*(p - 1). For example, taken modulo 5 the sequence becomes [1, 3, 2, 3, 4, 2, 3, 2, 1, 3, 2, 3, 4, 2, 3, 2, ...], which appears to be a purely periodic sequence of period 8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Feb 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015048 | OEIS-A340882

Irregular table: the n-th row polynomial is given by the formal power series expansion of Sum_{k >= 0} (1 + q)^(n*k + n^2)*Product_{j = 1..k} (1 - (1 + q)^(2*j-1)), n >= 1.

Conjecture: For fixed integers q and N, the sequence mod(R(n,q),N)n>=1 is either purely periodic or periodic with a pre-period 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Feb 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015049 | OEIS-A340883

Row sums of A340882.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Feb 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015050 | OEIS-A340991

Triangle T(n,k) whose k-th column is the k-fold self-convolution of the primes; triangle T(n,k), n>=0, 0<=k<=n, read by rows.

Conjecture: row polynomials are x*R(n,1) for n > 0 where R(n,k) = R(n-1,k+1) + x*R(n-1,1)*R(1,k) for n > 1, k > 0 with R(1,k) = prime(k) for k > 0. The same recursion seems to work for self-convolution of any other sequence. - _Mikhail Kurkov_, Apr 05 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Feb 01 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015051 | OEIS-A341015

Numbers k such that A124446(k) = 1.

Conjecture: the only term not of one of those forms is 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Feb 02 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015052 | OEIS-A341037

The successive remainders of the division by (a(n) + a(n+1)) of concat(a(n),a(n+1)) rebuild the sequence itself.

The sequence is conjectured to be a permutation of the integers > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_, Feb 03 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015053 | OEIS-A341092

Rows of Pascal's triangle which contain a 3-term arithmetic progression of a certain form: a(n) = (2*n^2 + 22*n + 37 + (2*n + 3)*(-1)^n)/8.

Conjecture (67) in _Ralf Stephan_'s paper, "Prove or Disprove. 100 Conjectures from the OEIS" asks if it is true that: "The numbers n such that the n-th row of Pascal's triangle contains an arithmetic progresion are n = 19 or n = (1/8)*[2*k^2 + 22*k + 37 + (2*k + 3)*(-1)^k], k > 0." Conjecture 1: Row 19 is the only row that contains a 3-term AP that doesn't fit the pattern found here. Conjecture 2: No row contains an AP of more than three coefficients. A brute-force search of n<=1100 found no counterexample of either conjecture above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Stauduhar_, Feb 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015054 | OEIS-A341119

a(n) is the least positive number that has exactly n divisors d such that d-1 is prime.

The conjectured terms are exact if for 0 <= n <= 10000 we have a(n) / A046523(A000005(a(n))) <= 9. For the found terms, a(n) / A046523(A000005(a(n))) <= 7.3. - _David A. Corneth_, Jun 15 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Feb 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015055 | OEIS-A341209

a(n) = (n^3 + 6*n^2 + 17*n + 6)/6.

Conjecture: Mark each point on a 7^k or 11^k grid with the number of points that are visible from the point; for n > 0, a(n) is the number of distinct values in both grids.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Torlach Rush_, Feb 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015056 | OEIS-A341239

a(n) = floor(r*floor(s*n)), where r = 1 + sqrt(2) and s = sqrt(2).

Conjecture: 1 < r*s*n - a(n) < 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 07 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015057 | OEIS-A341249

a(n) = floor(r*floor(s*n)), where r = 2 + sqrt(2) and s = sqrt(2).

Conjecture: 1 < r*s*n - a(n) < 2 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015058 | OEIS-A341261

Numbers that occur twice in the Wythoff ceiling array (A341260).

No term occurs more than twice. Conjecture: the difference sequence, (13,3,5,13,13,3,5,...) has only 4 distinct terms: 3,5,10,13.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015059 | OEIS-A341262

Numbers not in the Wythoff ceiling array (A341260).

Conjecture: the difference sequence, (2,5,3,10,3,5,...) has only 5 distinct terms: 2,3,5,10,13; cf. 3,5,10,13 for the difference sequence of A341261.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015060 | OEIS-A341280

Numbers k such that A073837(k) is a multiple of k.

Conjecture: For n > 1, a(n) is prime if and only if a(n) is odd and not a multiple of 5. - _Chai Wah Wu_, Feb 17 2021 The conjecture is false because a(35) = 1704233306223 is divisible by 3 and a(36) = 3361207818001 is divisible by 11. - _Martin Ehrenstein_, Feb 21 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Feb 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015061 | OEIS-A341322

Primes p such that (p*r) == q (mod p+r) where q and r are the next primes after p.

If x = (p+r)/2 and d = (r-p)/2, p*r = x^2 - d^2 == -d^2 (mod x), while x-d < q < x+d. We can't have q = x-d^2, so it must be that q >= 2 x - d^2, and thus d^2 >= 2x-q > x-d. Cramér's conjecture implies d^2 = o(x), so that implies the sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Feb 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015062 | OEIS-A341334

Concatenation of all 01-words, in the order induced by A016777; see Comments.

positions of 0-free words in W: A134342 (conjectured);

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015063 | OEIS-A341362

a(n) begins the first sequence of n consecutive positive integers with the same h-value and the same d-value in the Collatz (or '3x + 1') problem.

It is conjectured that starting from any number, and repeatedly applying the function on its previous result, we will always reach 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lamine Ngom_, Feb 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015064 | OEIS-A341379

E.g.f. A(x) satisfies: A(x) = sqrt(Sum_{n>=0} (x^n/n!) * exp(x*A(x)^n)).

The sequence is surprisingly regular. Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 19 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015065 | OEIS-A341418

Triangle read by rows: T(n, m) gives the sum of the weights of weighted compositions of n with m parts from generalized pentagonal numbers {A001318(k)}_{k>=1}.

That the row sums give the number of partitions follows from the pentagonal number theorem. See also the Apr 04 2013 conjecture in A000041 by _Gary W. Adamson_, and the hint for the proof by _Joerg Arndt_. The INVERT map of A = {1, 1, 0, 0, -5, -7, ...}, with offset 1, gives the A000041(n) numbers, for n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Feb 15 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015066 | OEIS-A341522

a(n) = A156552(3*A005940(1+n)).

a(n) = 1 + n + A086799(1+n). - [Conjectured by LODA-miner, and easily seen to be correct]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 15 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015067 | OEIS-A341541

a(n) is the number of steps to reach square 1 for a walk starting from square n along the shortest path on the square spiral board without stepping on any prime number. a(n) = -1 if such a path does not exist.

Conjecture: There is no "island of two or more nonprimes" enclosed by primes on the square spiral board. If the conjecture is true, then numbers n such that a(n) = -1 are the terms in A341542.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Feb 14 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015068 | OEIS-A341577

Decimal expansion of log(658) * log(1398).

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michal Paulovic_, Feb 14 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015069 | OEIS-A341654

Table read by antidiagonals upward: |T(n,k)| is the smallest number j such that j and j+1 have n and k divisors, respectively, or 0 if no such number exists, and the sign of T(n,k) is positive iff there exists only one such number j.

The positive terms here are the terms of A161460 (although 50175 and 59049 there are conjectural). Conjecture: T(n,k) < 0 for all T(n,k) where both n and k are even, other than the case n=k=2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Feb 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015070 | OEIS-A341658

Primes p such that p^2 - 1 has 32 divisors.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Feb 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015071 | OEIS-A341659

Primes p such that p^3 - 1 has 8 divisors.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Feb 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015072 | OEIS-A341662

Primes p such that p^4 - 1 has 160 divisors.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Feb 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015073 | OEIS-A341665

Primes p such that p^5 - 1 has 8 divisors.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Feb 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015074 | OEIS-A341666

Primes p such that p^6 - 1 has 384 divisors.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Feb 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015075 | OEIS-A341669

Primes p such that p^7 - 1 has 8 divisors.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Feb 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015076 | OEIS-A341670

a(n) is conjecturally the largest prime p such that, for every prime q > p, q^n - 1 has more divisors than does p^n - 1, or -1 if no such prime p exists.

a(n) is conjecturally the largest prime p such that, for every prime q > p, q^n - 1 has more divisors than does p^n - 1, or -1 if no such prime p exists. Conjecture: a(n) is also the largest prime p for which there exists no prime q > p such that p^n - 1 and q^n - 1 have the same number of divisors.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Feb 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015077 | OEIS-A341684

Expansion of the 2-adic integer Sum_{k>=0} k!.

Conjecture: this constant is transcendental, which means that it is not the root of any polynomial with integer coefficients. Conjecture: this constant is normal, which means for every binary (base-2) string s with length k, if we denote N(s,n) as the number of occurrences of s in the first n digits, then lim_{n->inf} N(s,n)/n = 1/2^k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Feb 17 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015078 | OEIS-A341685

Expansion of the 3-adic integer Sum_{k>=0} k!.

Conjecture: this constant is transcendental, which means that it is not the root of any polynomial with integer coefficients. Conjecture: this constant is normal, which means for every ternary (base-3) string s with length k, if we denote N(s,n) as the number of occurrences of s in the first n digits, then lim_{n->inf} N(s,n)/n = 1/3^k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Feb 17 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015079 | OEIS-A341686

Expansion of the 5-adic integer Sum_{k>=0} k!.

Conjecture: this constant is transcendental, which means that it is not the root of any polynomial with integer coefficients. Conjecture: this constant is normal, which means for every quinary (base-5) string s with length k, if we denote N(s,n) as the number of occurrences of s in the first n digits, then lim_{n->inf} N(s,n)/n = 1/5^k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Feb 17 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015080 | OEIS-A341687

Expansion of the 7-adic integer Sum_{k>=0} k!.

Conjecture: this constant is transcendental, which means that it is not the root of any polynomial with integer coefficients. Conjecture: this constant is normal, which means for every septenary (base-7) string s with length k, if we denote N(s,n) as the number of occurrences of s in the first n digits, then lim_{n->inf} N(s,n)/n = 1/7^k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Feb 17 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015081 | OEIS-A341723

Triangle read by rows: coefficients of expansion of certain weighted sums P_1(n,k) of Fibonacci numbers as a sum of powers.

Conjectures from _Mélika Tebni_, Sep 09 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 04 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015082 | OEIS-A341724

Triangle read by rows: coefficients of expansion of certain sums P_2(n,k) of Fibonacci numbers as a sum of powers.

Conjectures from _Mélika Tebni_, Sep 04 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 04 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015083 | OEIS-A341742

Nodes read by depth in a binary tree defined as: Root = 1; an even node N has a left child N + 1 if N + 1 is not a prime, and an odd node N has a left child sqrt(N + 2) if sqrt(N + 2) is a prime; the right child of a node N is 2*N.

If Conjectures 1 and 2 made in A340801 hold, the sequence contains all positive integers and each integer appears once in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Feb 18 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015084 | OEIS-A341743

T(n,k) is the number of labeled Eulerian graphs with n vertices and k edges (according to Harary and Palmer) or the number of labeled connected Eulerian graphs with n vertices and k edges (according to Mallows and Sloane); irregular triangle T, read by rows (n >= 0 and 0 <= k <= n*(n-1)/2).

Conjecture: T(n,n+1) = n!*(n-4)/8 = 15*A062199(n-5) for n >= 4 (with A062199(-1) = 0).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, Feb 18 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015085 | OEIS-A341761

Triangle read by rows in which row n is the coefficients of the subword complexity polynomial S(n,x).

Conjecture: the coefficient of x^n in S(n,x) is n*(n+1)/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shiyao Guo_, Feb 19 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015086 | OEIS-A341763

Numbers whose trajectory under iteration of sum of cubes of digits (map) produce a narcissistic number greater than nine.

Conjecture: all multiples of 3 are terms of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José de Jesús Camacho Medina_, Feb 19 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015087 | OEIS-A341816

The a(n)-th digit of the sequence is present in a(n+1). Lexicographically earliest sequence of distinct terms > 0 with this property.

This sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Carole Dubois_, Feb 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015088 | OEIS-A341822

Length of the longest 2-increasing sequence of positive integer triples with entries <= n.

It is conjectured that a(n) <= n^{3/2} for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marcel K. Goh_, Feb 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015089 | OEIS-A341838

Number of permutations of degree n with highest Shannon entropy.

For 0 < n < 4, the Shannon entropy is 0 for every permutation, while for n >= 4 I found the maximum value of the entropy to be log(n) - log(10 - 6(-1)^n)/n. This holds for at least n=12, and I conjecture that it holds for n > 12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrea G. Amato_, Feb 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015090 | OEIS-A341843

Number of sexy consecutive prime pairs below 2^n.

For each n from 9 through 48, the most frequently occurring difference between consecutive primes is 6. On p. 108 of the article by Odlyzko et al., the authors estimate that around n=117, the jumping champion (i.e., the most frequently occurring difference between consecutive primes) becomes 30, and around n=1412 it becomes 210. Successive jumping champions are conjecturaly the primorial numbers A002110.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Feb 21 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015091 | OEIS-A341860

a(n) is the least k such that A341857(k) = psi(k^2)/k = n, or 0 if no such k exists.

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Feb 21 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015092 | OEIS-A341861

Number of primes among the (p-1)/2 numbers {2*p+1, 4*p+1, ..., (p-1)*p+1}, p = prime(n).

By Dirichlet's theorem on arithmetic progressions, we know there exists a prime q of the form 2*k*p+1. But the theorem does not give us any information about the size of the smallest q. It is conjectured that q < p^2. Moreover, it seems that a(n) goes to infinity as n increases.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Feb 21 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015093 | OEIS-A341883

Number of partitions of 2*n+1 into three parts from A339506.

Conjecture: for n > 0 this sequence is always positive (it has been verified up to 10^6).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lechoslaw Ratajczak_, Feb 22 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015094 | OEIS-A341920

Number of simple strong triangulations of a fixed pentagon with n interior nodes.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Howroyd_, Feb 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015095 | OEIS-A341941

T(n,k) is the number of unlabeled even graphs with n vertices and k edges; irregular triangular array T read by rows (n >= 0, 0 <= k <= n*(n-1)/2).

Conjecture (due to _Peter Luschny_): Sum_{k=0}^{n*(n-1)/2} (-1)^k*T(n,k) = A263626(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, Feb 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015096 | OEIS-A342000

a(1) = 0, and for n > 1, a(n) = 1 if G(A329886(n)) >= G(A329886(floor(n/2))), otherwise 0, where G(n) = sigma(n) / (n*log(log(n))), where sigma is the sum of the divisors.

It seems that the tree has infinitely long leftward branches that contain either only zeros or only ones after a while: The leftmost edge (that are primorials, A002110, in A329886) appears to consist of zeros only after its single 1 at the second term. This depends on the (so far conjectural) observation that (log(A002110(n)) + log(prime(1+n)))^(prime(1+n)) > log(A002110(n))^(1+prime(1+n)) for all n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_ and _Michael De Vlieger_, Mar 04 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015097 | OEIS-A342011

Lexicographically earliest infinite sequence such that a(i) = a(j) => f(i) = f(j), for all i, j >= 1, with f(1) = 2 and f(n) = A004490(n)/A004490(n-1) when n > 1, where A004490(n) is the n-th colossally abundant number.

This is also the restricted growth sequence transform of A073751, provided that quotient A004490(1+n)/A004490(n) is always prime, which is implied by a conjecture mentioned in Lagarias' paper. Note that the b-file of A073751 is computed based on the knowledge that the conjecture holds at least for the first 10^7 quotients.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015098 | OEIS-A342031

Starts of runs of 5 consecutive numbers that have mutually distinct exponents in their prime factorization (A130091).

Aktaş and Ram Murty (2017) gave an explicit upper bound to such a run of 23 numbers. They found the first 5 terms of this sequence (and stated that there are a few more known up to 7*10^8), and said that we may conjecture (based on numerical evidence) that there are no 6 consecutive numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Feb 25 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015099 | OEIS-A342042

When a digit d in the digit-stream of this sequence is even, the next digit is > d.

Conjecture 1. For k >= 1, the terms of length k in S form a consecutive block with the same starting and ending points as in P. In both P and S, the block begins with 10101... (1's and 0's alternate, length is k) and end with 99...9 (k 9's). Conjecture 2. We know every number appears in S. Suppose x = S(m) = 899...9 (with k-1 9's). Then x is the last term of length k in S that begins with a digit <= 8. The remaining terms of length k have leading digit 9 and appear in order, ending with 99...9 (k 9's). Let c1 = 7.422574840... and c2 = 1.3824387... be the constants defined in A377918. Then assuming Conjecture 1, the index of the last term of length k in the present sequence is close to (c2*c1^k, 10^k). [Thanks to _Sebastian Karlsson_ for pointing out that Conjecture 1 is required and is as yet unproved.] A proof can be found in the links. Since all k-digit numbers starting with 9 appears before any (k+1)-digit number, we get that terms of a certain length form a consecutive block. In particular, this proves Conjectures 1 and 2 above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_, Feb 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015100 | OEIS-A342064

Primes p such that p^8 - 1 has 384 divisors.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Feb 27 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015101 | OEIS-A342066

Primes p such that p^10 - 1 has 256 divisors.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Feb 27 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015102 | OEIS-A342067

Primes p such that p^11 - 1 has 8 divisors.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Feb 28 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015103 | OEIS-A342070

Numbers k such that there are more primes in the interval [2*k+1, 3*k] than there are in the interval [k+1, 2*k].

Conjecture: 1323 is the final term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Mar 23 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015104 | OEIS-A342071

Numbers k such that there are more primes in the interval [3*k+1, 4*k] than there are in the interval [2*k+1, 3*k].

Conjecture: a(194)=3977 is the final term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Mar 23 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015105 | OEIS-A342100

Abundant numbers k at which the ratio (number of abundant numbers in 1..k)/k reaches a new record high.

Conjecture: a(115) is the final term of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Feb 27 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015106 | OEIS-A342137

Least k such that k^8 - 16 has a root modulo prime(n) where n is the sequence index.

This sequence of least roots modulo prime(n) for k^8 - 16, represents the counterexample to a conjecture that stated if k^m - j has a root for modulo every prime, then j must be of the form j = y^m. The corrected theorem which excludes the counterexample is known as the Grunwald-Wang Theorem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Mar 01 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015107 | OEIS-A342158

Dividing a(n) by the first digit of the next term leaves the next digit as remainder.

However, we may conjecture that all other numbers (i.e., all positive integers whose second digit, if it exists, is strictly smaller than the first digit), will eventually appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _M. F. Hasler_, Mar 02 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015108 | OEIS-A342171

Nonnegative integers k such that k < sec(k)*csc(k).

Conjecture: 2*k/Pi is either a little more than an even integer or a little less than an odd integer. The conjecture is true. As k > 0 increases, satisfaction of the inequality k < sec(k)*csc(k) requires that sec(k)*csc(k) be a large positive number. Since sec(k)*csc(k) = 1/(sin(k)*cos(k)) = 2/sin(2*k), this requires that sin(2*k) be a small positive number, which occurs only when 2*k/Pi is a little more than an even integer or a little less than an odd integer. - _Jon E. Schoenfield_, Mar 06 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José Mauricio de Oliveira Neto_, Mar 04 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015109 | OEIS-A342172

a(n) is the smallest n-bit number having the most common prime signature among n-bit numbers. (In case more than one prime signature is tied for most common, choose the smallest n-bit number whose prime signature is one of those tied.)

Conjecture: for all n >= 1, there is only one prime signature in I_n that occurs with the maximum frequency.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Mar 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015110 | OEIS-A342175

a(n) is the difference between the n-th composite number and the smallest larger composite to which it is relatively prime.

Conjecture: The only nonprime terms are squares (based on checking the first 2 million terms). - _Ivan N. Ianakiev_, Mar 28 2021 The conjecture above is false (see A353203 for counterexamples). - _Ivan N. Ianakiev_, Jul 04 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _William C. Laursen_, Mar 04 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015111 | OEIS-A342202

T(n,k) = V(n,k)/k!, where V(n,k) = k^(n*k) - Sum_{t=1..k-1} binomial(k,t)*k^(n*(k-t))*V(n,t) for n, k >= 1; square array T read by upwards antidiagonals.

For the other compositions (s_1, ..., s_r) of k (of any size r >= 2), we group them according to the their last element s_r = s in {1, 2, ..., k - 1}, which gives rise to the factor k^(n*s)/s! = (Sum_{i=1..r} s_i)^(n*s_r)/s_r!. Using the inductive hypothesis, we substitute the expression for T(n,k-s) in the recurrence T(n,k) = k^(n*k)/k! - Sum_{s=1..k-1} k^(n*s)/s! * T(n,k-s). Each term in the expression for T(n,k-s) corresponds to a composition of k - s and is postmultiplied by k^(n*s)/s! = (Sum_{i=1..r} s_i)^(n*s_r)/s_r!. We thus get a term in the expression for T(n,k) that corresponds to a composition of the form (composition of k - s) + s, and the sign of this term is (-1)^((size of composition of k - s) + 1). The rest of the proof follows easily.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petros Hadjicostas_, Mar 04 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015112 | OEIS-A342235

Coordination sequence of David Eppstein's "Tetrastix" graph.

The graph in question is the subgraph of the adjacency graph of Z^3, induced by the set of vertices whose coordinates are not all of the same parity. The graph is regular of degree 4, and is vertex-transitive, so it only has one vertex coordination sequence. Conjecture: the second differences are 5, 8, 6, [4, 8], where the bracketed portion repeats.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Allan C. Wechsler_, Mar 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015113 | OEIS-A342309

When A123556(n) is the number of primes (not necessarily consecutive) in the longest arithmetic progression of primes with common difference n, a(n) is the smallest term of the first such arithmetic progression.

About the k-tuples conjecture in A123556: let p = A053669(n), if the arithmetic progression of p elements starting at p with difference n consists of primes, then A123556(n) = p, otherwise A123556(n) = p-1. The first 21 terms of this sequence are precisely the same as A053669, then a(22) = 7 (corresponding to the arithmetic progression (7,29)) while A053669(22) = 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Mar 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015114 | OEIS-A342326

a(n) is the smallest nonnegative integer that can be written as a sum of two distinct nonzero triangular numbers in exactly n ways or -1 if no such integer exists.

Conjecture: This sequence has a positive a(n) for every positive integer n, and each sequence in the infinite indexed family, of which this sequence offers the initial terms, is infinite, as well.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Root_, Mar 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015115 | OEIS-A342346

a(1) = 4, a(n) = smallest palindromic nontrivial multiple of a(n-1) containing all decimal digits of a(n-1).

For n = 5..8, we have a(n) = 7568 * A002275(2^(n-3)), and it follows that a(9) <= 7568 * A002275(64). Conjecture: for all n >= 5, a(n) = 7568 * A002275(2^(n-3)). Note that 7568 is a term of A370052 and A370053. - _Max Alekseyev_, Feb 08 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Mar 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015116 | OEIS-A342347

a(1) = 7, a(n) = smallest palindromic nontrivial multiple of a(n-1) containing all decimal digits of a(n-1).

For n=6..8, a(n) = 196930692 * A002275(2^(n-3)), and it follows that a(9) <= 196930692 * A002275(64). Conjecture: for all n >= 6, a(n) = 196930692 * A002275(2^(n-3)). Note that 196930692 is a term of A370052 and A370053. - _Max Alekseyev_, Feb 15 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Mar 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015117 | OEIS-A342348

Smallest number > 3 whose representation in all bases from 2 up to n consists only of '0's, '1's, '2's and '3's.

Conjecture: there are no more terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Mar 12 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015118 | OEIS-A342401

Numbers k such that the number of distinct sums of distinct unitary divisors of k sets a new record.

The analogous sequence of records with all the divisors (A119347) is conjecturally the highly abundant numbers (A002093), excluding 3 and 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Mar 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015119 | OEIS-A342559

Number of equal spheres setting a new density record in relation to the volume of the spherical layer that is occupied by the spheres when arranged touching the surface of a container sphere according to the criterion of maximizing their minimum mutual distance.

A conjectured continuation after the term 48 is 72, 78, 84, 92, 98, 120.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Apr 07 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015120 | OEIS-A342578

a(n) = n! * [x^n] (Sum_{j>=0} n^(j*(j+1)/2) * x^j/j!)^(1/n) for n > 0, a(0) = 1.

a(n) == 1 (mod n*(n-1)) for n >= 2 (see "general conjecture" in A178319 and link to proof by _Richard Stanley_ above).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Mar 15 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015121 | OEIS-A342591

Numbers k such that A340740(k) <= k.

Conjecture: these are all the terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Mar 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015122 | OEIS-A342594

Earliest occurrence of the next distinct width pattern (as listed in A342592) in the symmetric representation of sigma(n) not yet encountered as n increases.

Conjecture: If for some number n the symmetric representation of sigma(n) has the symmetric width pattern w in row n of A342592 then infinitely many numbers have that width pattern w.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Mar 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015123 | OEIS-A342833

Integers m such that the number of divisors whose last digit equals the last digit of m sets a new record.

For n >= 3, a(n) = 10 * A002182(n) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Mar 23 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015124 | OEIS-A342839

Numbers k such that there are more primes in the interval [4*k+1, 5*k] than there are in the interval [3*k+1, 4*k].

Conjecture: a(876) = 11895 is the final term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Mar 23 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015125 | OEIS-A342842

All positive integer solutions m of equation A342369^k(6*p - 2) = m*3 + 2, sorted by p and k in ascending order, p has higher priority than k. p and k are positive integers. "^k" means recursion here.

It is conjectured that this sequence is a permutation of the positive integers. If it does not contain all positive integers, then there exists a number of the form q = p*6 - 2, where no solution for j*3 - 1 = A006370^k(q) can be found for any j and any k. Such an example is not yet known.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Mar 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015126 | OEIS-A342843

a(n) is the number of edges of the regular polygon such that packing n nonoverlapping equal circles inside the regular polygon gives the highest packing density. a(n) = 0 if such a regular polygon is a circle.

Terms for n = 11, 12, 13 and 14 are conjectured values supported by numerical results (see Packomania in the links). Conjecture: As n tends to infinity, a(n) takes the value of 3 or 6 and the packing density approaches sqrt(3)*Pi/6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Apr 12 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015127 | OEIS-A342877

a(n) = 1 if the average distance between consecutive first n primes is greater than that of the first n-1 primes, otherwise a(n) = 0, for n > 2.

Conjecture: lim_{n->infinity} (1/n) Sum_{i=1..n} a(i) < 1/2. In support of the conjecture: If it is assumed, as an approximation, that position of primes follows a Poisson point process then the distance between consecutive primes is a stochastic variable with exponential probability distribution function. The probability that an exponentially distributed stochastic variable takes a value larger than the mean value is about 0.367879.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andres Cicuttin_, Mar 28 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015128 | OEIS-A342916

a(n) = (1+n) / gcd(1+n, A001615(n)), where A001615 is Dedekind psi, n * Product_{p|n, p prime} (1 + 1/p).

It is conjectured that a(n) = 1 only when n is a prime, A000040. See _Thomas Ordowski_'s May 21 2017 problem in A001615.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015129 | OEIS-A342965

Number of permutations tau of {1,...,n} with tau(n) = n such that tau(1)^tau(2) + ... + tau(n-1)^tau(n) + tau(n)^tau(1) is a square.

Conjecture: a(n) > 0 for all n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 31 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015130 | OEIS-A342992

Smallest k such that k*n contains only prime digits, or 0 if no such k exists.

Conjecture: a(n) > 0 for all n that are not divisible by 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Metin Sariyar_, Apr 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015131 | OEIS-A343009

a(n) = (n^(2n)-1)/(n^2-1) for n > 1, a(1) = 1.

Conjecture: for n > 2, a(n) is a Fermat pseudoprime to base n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Apr 02 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015132 | OEIS-A343017

a(1)=1. For n > 1, a(n) is the n-th positive integer after a(n-1) which cannot be written as a sum of distinct preceding terms in the sequence.

Theorem: for n >= 6, a(n) < (Sum_{i=1..n-1} a(i)) - a(n-3). Corollary: a(n) = O(x^n) where x is the positive real solution to x^4 - 2x^3 + x - 1 = 0, exact value (1 + sqrt(3 + 2*sqrt(5)))/2 = 1.86676.... I conjecture that this bound can be tightened further. - _Matthew R. Maas_, Apr 21 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthew R. Maas_, Apr 02 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015133 | OEIS-A343052

Table read by ascending antidiagonals: T(k, n) is the minimum vertex sum in a perimeter-magic k-gon of order n.

T(k, 3) = A265225(k-1) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Apr 03 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015134 | OEIS-A343073

a(n) is the number of integers 0 < b < n such that b^^x == 1 (mod n) has a solution; ^^ denotes the tetration operation (cf. A321312).

a(n) <= A000010(n), since gcd(b,n)=1 is a necessary condition. There is equality when n = 2 and n = 3. It is a conjecture that there are no more equality cases.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernat Pagès Vives_, Apr 04 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015135 | OEIS-A343093

Number of rooted toroidal maps with n edges and no isthmuses.

Conjectural g.f.: A(x) = (Sum_{n >= 1} (1/4)*binomial(4*n, n)*x^n)^2. - _Peter Bala_, Jul 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Howroyd_, Apr 04 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015136 | OEIS-A343099

Sums of 3 distinct odd squares.

Conjecture: contains all numbers == 3 (mod 8) except 3, 11, 19, 27, 43, 51, 67, 99, 123, 163, 187, 267, 627. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Apr 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015137 | OEIS-A343130

Number of permutations tau of {1,...,n} with tau(1) = 1 and tau(2) = 2 such that Product_{k=1..n} k^tau(k) = (p-1)^3 for some prime p.

Conjecture 1: a(n) > 0 for all n > 5. Conjecture 2: For any integer n > 5, there is a permutation tau of {1,...,n} with tau(1) = n - 1 and tau(n) = n such that tau(1)^tau(2)*...*tau(n-1)^tau(n)*tau(n)^tau(1) = q^2 for some integer q with q - 1 and q + 1 twin prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015138 | OEIS-A343144

a(n) is the smallest number m with n divisors such that m + 1 has n + 1 divisors; or 0 if no such number exists.

Conjecture: a(n) = 0 for more than 95% of the indices n in 1..1000. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Apr 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015139 | OEIS-A343249

a(n) is the least k0 <= n such that v_2(n), the 2-adic order of n, can be obtained by the formula: v_2(n) = log_2(n / L_2(k0, n)), where L_2(k0, n) is the lowest common denominator of the elements of the set S_2(k0, n) = {(1/n)*binomial(n, k), with 0 < k <= k0 such that k is not divisible by 2} or 0 if no such k0 exists.

Conjecture: a(n) is the greatest power of a prime different from 2 that divides n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dario T. de Castro_, Apr 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015140 | OEIS-A343250

a(n) is the least k0 <= n such that v_3(n), the 3-adic order of n, can be obtained by the formula: v_3(n) = log_3(n / L_3(k0, n)), where L_3(k0, n) is the lowest common denominator of the elements of the set S_3(k0, n) = {(1/n)*binomial(n, k), with 0 < k <= k0 such that k is not divisible by 3} or 0 if no such k0 exists.

Conjecture: a(n) is the greatest power of a prime different from 3 that divides n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dario T. de Castro_, Apr 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015141 | OEIS-A343251

a(n) is the least k0 <= n such that v_5(n), the 5-adic order of n, can be obtained by the formula: v_5(n) = log_5(n / L_5(k0, n)), where L_5(k0, n) is the lowest common denominator of the elements of the set S_5(k0, n) = {(1/n)*binomial(n, k), with 0 < k <= k0 such that k is not divisible by 5} or 0 if no such k0 exists.

Conjecture: a(n) is the greatest power of a prime different from 5 that divides n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dario T. de Castro_, Apr 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015142 | OEIS-A343252

a(n) is the least k0 <= n such that v_7(n), the 7-adic order of n, can be obtained by the formula: v_7(n) = log_7(n / L_7(k0, n)), where L_7(k0, n) is the lowest common denominator of the elements of the set S_7(k0, n) = {(1/n)*binomial(n, k), with 0 < k <= k0 such that k is not divisible by 7} or 0 if no such k0 exists.

Conjecture: a(n) is the greatest power of a prime different from 7 that divides n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dario T. de Castro_, May 31 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015143 | OEIS-A343253

a(n) is the least k0 <= n such that v_11(n), the 11-adic order of n, can be obtained by the formula: v_11(n) = log_11(n / L_11(k0, n)), where L_11(k0, n) is the lowest common denominator of the elements of the set S_11(k0, n) = {(1/n)*binomial(n, k), with 0 < k <= k0 such that k is not divisible by 11} or 0 if no such k0 exists.

Conjecture: a(n) is the greatest power of a prime different from 11 that divides n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dario T. de Castro_, May 31 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015144 | OEIS-A343260

a(n) = 2 * T(n,(n+1)/2) where T(n,x) is a Chebyshev polynomial of the first kind.

Conjecture: a(p^r) == 1 (mod p^(2*r)) for positive integer r and all primes p >= 5. - _Peter Bala_, Mar 11 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Apr 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015145 | OEIS-A343268

Numbers that are not the sum of exactly four terms from A020330 (not necessarily distinct).

Madhusudan et al. (2018) conjectured that a(112) = 1772 is the last term of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Apr 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015146 | OEIS-A343311

Numbers of the form x + y + z with distinct positive integers x,y,z such that (x+y+z) | x*y*z.

Conjecture: Consists of all composite numbers except 4, 8, 9, and 25. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Apr 11 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015147 | OEIS-A343326

Number of ways to write n as the integral part of (a^3+b^3)/2 + (c^3+d^3)/6, where a,b,c,d are nonnegative integers with a >= max{b,1} and c >= max{d,1}.

Conjecture: a(n) > 0 for any nonnegative integer n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 11 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015148 | OEIS-A343327

Start with a(1)=1; thereafter a(n+1) = Sum_{k=1..n such that d(a(k)) = d(a(n))} d(a(k)) where d = A000005.

Every nonzero number occurs in the sequence since all terms in A000005 are included. The multiplicity of any term m (> 2) is d(m)-1 (since m > 2 cannot be created by adding m 1s). Prime 2 occurs twice, consecutively (see example), and thereafter no two adjacent terms can be prime because the term prior to a prime q is the first occurrence of 2^(q-1) (composite for q > 2), and since d(q)=2, the following term is a multiple (> 1) of 2. Thus each prime > 2 is flanked by composite even numbers, and occurs once only. Conjecture: The primes, after the first appearance of 2, appear in their natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Apr 12 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015149 | OEIS-A343368

Number of ways to write n as floor((a^3+b^3)/3) + floor((c^3+d^3)/5), where a,b,c,d are nonnegative integers with a > b and c >= d.

Conjecture 1: a(n) > 0 for all n >= 0. Conjecture 2: Each n = 0,1,... can be written as floor((a^3+b^3)/4) + floor((c^3+d^3)/5) with a,b,c,d nonnegative integers. Both conjectures have been verified for all n = 0..10^5. We also conjecture that the pair (4,5) of denominators in Conjecture 2 can be replaced by some other pairs such as (4,6), (5,6), (3,7), (4,7), (5,7), (6,7).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 12 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015150 | OEIS-A343376

a(0) = 0, a(1) = 1. For n >= 2, a(n) = a(n-1)/(n-1) if n-1 divides a(n-1); otherwise, a(n) = a(n-1) + a(n-2).

2 divides a(10) through a(32), 3 divides a(134) through a(144), 5 and 7 divide a(116) through a(140), and 11 divides a(89) through a(297). Conjecture: For any prime p, there exist r and s = k*p from A343456 for which p divides a(r) through a(s).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Timothy L. Tiffin_, Apr 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015151 | OEIS-A343384

Number of ways to write n as [a^3/3] + [b^3/4] + [c^3/5] + [d^6/6] with a,b,c,d positive integers, where [.] is the floor function.

3-4-5-6 Conjecture: a(n) > 0 for all n >= 0. Conjecture verified up to 2*10^9. - _Giovanni Resta_, Apr 28 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015152 | OEIS-A343387

Number of ways to write n as x^2 + [y^2/2] + [z^4/8], where [.] is the floor function, x is a nonnegative integer, and y and z are positive integers.

Conjecture: a(n) > 0 for all n >= 0. We also conjecture that each n = 0,1,... can be written as x^2 + [y^2/3] + [z^4/7] with x,y,z nonnegative integers. See also A343391 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015153 | OEIS-A343391

Number of ways to write n as x^2 + [y^2/4] + [z^4/6] with x,y,z positive integers, where [.] is the floor function.

Conjecture: a(n) > 0 for all n > 0. See also A343387 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015154 | OEIS-A343397

Number of ways to write n as 2^x + [y^2/3] + [z^2/4] with x,y,z positive integers, where [.] is the floor function.

Conjecture: a(n) > 0 for all n > 1. Conjecture verified up to 10^10. - _Giovanni Resta_, Apr 14 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015155 | OEIS-A343400

Number of ways to write n as 3^x + [y^2/3] + [z^2/4], where [.] is the floor function, x is a nonnegative integer, and y and z are positive integers.

Conjecture: a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015156 | OEIS-A343411

Number of ways to write n as x^3 + [y^3/2] + [z^3/3] + 2^w, where [.] is the floor function, x,y,z are positive integers, and w is a nonnegative integer.

Conjecture: a(n) > 0 for all n > 1. Conjecture verified up to 10^10. - _Giovanni Resta_, Apr 14 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 14 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015157 | OEIS-A343419

Number of distinct sets { p(i) - p(j) : 1 <= i <= j <= n } where p ranges over all permutations of [n].

a(n) <= 2*a(n-1) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Baohua Tian_, Apr 15 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015158 | OEIS-A343460

Number of ways to write n as x^6 + y^3 + z*(3*z+1)/2 + 2^k, where x and y are nonnegative integers, z is an integer and k is a positive integer.

Conjecture: a(n) > 0 for all n > 1. Conjecture verified up to 10^10. - _Giovanni Resta_, Apr 18 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015159 | OEIS-A343464

The number of n-vertex graphs that are minimally non-Hamming-embeddable.

I conjecture (on admittedly weak evidence) that these graphs are always planar.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Don Knuth_, Apr 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015160 | OEIS-A343480

Decimal expansion of 9 * Product_{p prime >= 5} (1-3/p)/(1-1/p)^3.

This constant appears in the conjectured asymptotic formulas for A343478 and A343479.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Apr 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015161 | OEIS-A343491

Number of representations of n! as a sum of 3 tetrahedral numbers (A000292).

Conjecture I: There are infinitely many n such that a(n) >= 1. Conjecture II: Natural density of numbers n such that a(n) >= 1 is 1. Conjecture III: Numbers n such that a(n) = 0 is a finite sequence. Conjecture IV: a(n) >= 1 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Apr 17 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015162 | OEIS-A343502

Numbers k such that tau(tau(k)) and tau(k+1) are both prime, where tau is the number of divisors function.

Conjecture: all members except 2 and 3 are composite. Open question: is there a finite number of odd terms in this sequence?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Claude H. R. Dequatre_, Apr 17 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015163 | OEIS-A343528

Number of ways to write n as x^4 + T(y)^2 + T(z) + 2^w, where x,y,z are nonnegative integers, w is a positive integer, and T(m) denotes the triangular number m*(m+1)/2.

Conjecture: a(n) > 0 for all n > 1. Conjecture verified up to 10^10. - _Giovanni Resta_, Apr 21 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 18 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015164 | OEIS-A343564

a(n) is the sum of 2*n mod p for primes p such that 2*n-p is prime.

Conjecture: the only n for which a(n) <= n are 1, 2, 3, 5, 7, 11, 19, and 31.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Apr 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015165 | OEIS-A343635

10^n + a(n) is the least (n+1)-digit prime member of a prime 5-tuple, or a(n) = 0 if no such number exists.

Numerical evidence strongly suggests the conjecture that 0 < a(n) < 10^n for all n > 0, but not even the existence of infinitely many prime 5-tuples is proved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jul 17 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015166 | OEIS-A343636

10^n + a(n) is the smallest (n+1)-digit initial member of a prime sextuplet, or a(n) = 0 if no such number exists.

Numerical evidence strongly suggests the conjecture that 0 < a(n) < 10^n for all n > 5, but not even the existence of infinitely many prime sextuplets is proved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jul 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015167 | OEIS-A343637

10^n + a(n) is the least (n+1)-digit prime member of a prime septuplet, or a(n) = 0 if no such number exists.

Numerical evidence strongly suggests the conjecture that 0 < a(n) < 10^n for all n > 4, but not even the existence of infinitely many prime septuplets is proved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jul 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015168 | OEIS-A343705

Numbers that are the sum of five positive cubes in exactly three ways.

It is conjectured that the number of ways of writing N as a sum of 5 positive cubes grows like C(N)*N^(2/3), where C(N) depends on N but is bounded away from zero by an absolute constant (Vaughan, 1981; Vaughan and Wooley, 2002). So the number will exceed 3 as soon as N is large enough, and so it is very likely that this sequence is finite. However, at present this is an open question. - _N. J. A. Sloane_, May 15 2021 (based on correspondence with Robert Vaughan and Trevor Wooley).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Consiglio, Jr._, Apr 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015169 | OEIS-A343735

Odd palindromes having more divisors than all smaller odd palindromes.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Jun 22 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015170 | OEIS-A343755

Number of regions formed by infinite lines when connecting all vertices and all points that divide the sides of an equilateral triangle into n equal parts.

Conjectured formula: a(n) = A344279(n) + 6*(n^2 - n + 1). Conjectured formula: a(n) = A344279(n) + A121205(n-1), for n>=7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 28 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015171 | OEIS-A343784

Decimal expansion of Sum_{k>=1} A065043(k)/2^k.

Named "the Liouville number" and conjectured to be a transcendental number by Borwein and Coons (2008).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Apr 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015172 | OEIS-A343825

Table read by antidiagonals upward: T(n,k) is the least m such that there exists a sequence k = b_1 <= b_2 <= ... <= b_t = m such that no term appears n or more times and the product of the sequence is of the form c^n, where c is an integer; n >= 1 and k >= 0.

Conjecture: T(p,k) = A071537(k) for fixed k and sufficiently large prime p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Apr 30 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015173 | OEIS-A343829

Distinct values of gcd(B(2*j), B(2*j+2)) in the order of their appearance, where B(k) = A027641(k).

Conjecture: all terms greater than 1 are irregular primes (A000928) or a product of irregular primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Apr 30 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015174 | OEIS-A343862

Number of ways to write n as x + y + z with x, y, z positive integers such that x^2*y^2 + 5*y^2*z^2 + 10*z^2*x^2 is a square.

Conjecture 1: a(n) > 0 for all n > 2. We have verified a(n) > 0 for all n = 3..10000. Conjecture 1 holds if a(p) > 0 for each odd prime p. For any n > 0 we have a(3*n) > 0 since 3*n = n + n + n and 1 + 5 + 10 = 4^2. See also A340274 for a similar conjecture. Conjecture 2: There are infinitely many triples (a,b,c) of positive integers such that each n = 3,4,... can be written as x + y + z with x,y,z positive integers and a*x^2*y^2 + b*y^2*z^2 + c*z^2*x^2 a square. Conjecture 1 holds for all n < 2^15. - _Martin Ehrenstein_, May 02 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 02 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015175 | OEIS-A343881

Table read by antidiagonals upward: T(n,k) is the least integer m > k such that k^x * m^y = c^n for some positive integers c, x, and y where x < n and y < n; n >= 2, k >= 1.

Conjecture: T(p,k) = A064549(k) for fixed k > 1 and sufficiently large p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, May 02 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015176 | OEIS-A343883

Numbers k such that A023896(k) is a multiple of A340179(k).

Conjecture: there are only 8 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, May 02 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015177 | OEIS-A343897

Number of ways to write n as 2*x + y + z with x,y,z positive integers such that 16*x^2*y^2 + 19*y^2*z^2 + 29*z^2*x^2 is a square.

Conjecture: a(n) > 0 for all n > 3. We have verified a(n) > 0 for all n = 4..10000. The conjecture holds if a(p) > 0 for each odd prime p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 03 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015178 | OEIS-A343913

Positive integers m such that 2*m^2 - 1 = x^4 + y^4 for some nonnegative integers x and y with |x - y| > 1.

Conjecture: The sequence has infinitely many terms. See also A343917 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 03 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015179 | OEIS-A343917

Positive integers m with 2*m^2 - 2^4 = x^4 + y^4 for some nonnegative integers x and y with |x-y| > 2.

Conjecture: The sequence has infinitely many terms. See also A343913 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 04 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015180 | OEIS-A343950

Number of ways to write n as x + y + z with x^2 + 4*y^2 + 5*z^2 a square, where x,y,z are positive integers with y or z a positive power of two.

Conjecture 1: a(n) > 0 for all n > 7. Conjecture 2: For any integer n > 7, we can write n as x + y + z with x,y,z positive integers such that x^2 + 2*y^2 + 3*z^2 is a square. Conjecture 3: For any integer n > 4, we can write n as x + y + z with x,y,z positive integers such that 3*x^2 + 4*y^2 + 5*z^2 (or x^2 + 3*y^2 + 5*z^2) is a square.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015181 | OEIS-A343973

Primes p such that both p-1 and p+1 are in A093641.

Conjecture: these are all the terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, May 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015182 | OEIS-A343974

Even numbers k such that the two sets of primes in the Goldbach representation of k and k+2 as the sum of two odd primes do not intersect.

Conjecture: a(n) is congruent to 2 mod 6 with a(n)-3 not prime. a(n) = 6*A046953(n) + 2 (conjectured). - _Hugo Pfoertner_, Jun 09 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giorgos Kalogeropoulos_, Jun 07 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015183 | OEIS-A343981

a(n) is the least integer h such that there exists a Pythagorean triple whose hypotenuse is h and whose other legs z satisfy A176774(z) = n.

a(4)=0 is conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, May 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015184 | OEIS-A344018

Table read by rows: T(n,k) (n >= 1, 1 <= k <= 2^n) is the number of cycles of length k which can be produced by a general n-stage feedback shift register.

Conjectures by Bryant and Christensen (1983): Conjecture 1: T(k-4,k) = A001037(k) - 4*A346018(k,3) - 2*gcd(k,2) + 10 for k >= 8. Conjecture 2: T(k-5,k) = A001037(k) - 8*A346018(k,4) - gcd(k,3) + 19 for k >= 11. Conjecture 3: T(k-6,k) = A001037(k) - 16*A346018(k,5) - 4*gcd(k,2) - 2*gcd(k,3) + 48 for k >= 15. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 21 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015185 | OEIS-A344020

A sequence of prime numbers: a(1)=2, a(n+1) is the least prime dividing Product_{i in S} a(i)^2 + Product_{i not in S} a(i)^2, minimized over all subsets S of {1..n}.

For n > 1, a(n) >= p, where p is the smallest prime p such that p == 1 (mod 4) and a(2)*a(3)*...*a(n-1) is a nonzero square modulo p. Conjecture: a(n) = p. - _Jinyuan Wang_ and _Max Alekseyev_, Jul 04 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lucas Hoogendijk_, May 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015186 | OEIS-A344058

Number of ways to write n as x + y + z with x*y + 2*y*z + 3*z*x a square, where x,y,z are positive integers with x or y a power of two (including 2^0 = 1).

Conjecture: a(n) > 0 for all n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015187 | OEIS-A344083

a(n) = f(x)+f(y)+f(z), where (x,y,h) is the n-th Pythagorean triple listed in (A046083, A046084, A009000), and f(m)=A176775(m) is the index of m as k-gonal number for the smallest possible k.

Conjecture: there are no other Pythagorean triples that give this minimum. In other words, it is the only triple with 3 A090467 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, May 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015188 | OEIS-A344105

a(n) is the number of distinct numbers of transversals of order n diagonal Latin squares.

For all spectra of even orders all known values included in them are divisible by 2. For all spectra of orders n=6, n=10 and n=14 (and probably for all n=4k+2) all known values included in the corresponding spectra are divisible by 4. This leads to the following hypothesis: a(2k) <= (A287644(2k) - A287645(2k) + 2)/2 and a(4k+2) <= (A287644(4k+2) - A287645(4k+2) + 4)/4, where w(n) = A287644(n) - A287645(n) + 1 is a width of corresponding spectra and (w(n)+1)/2 is done to round the result of the division up. - _Eduard I. Vatutin_, Mar 21 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eduard I. Vatutin_, Jun 22 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015189 | OEIS-A344142

Lexicographically first irreducible polynomial over GF(2) of degree n with the lowest possible number of terms, evaluated at X = 2.

Note that it is conjectured that an irreducible polynomial of degree n with 5 terms exists for every n. It follows from the conjecture that A000120(a(n)) = 3 for n in A073571 and 5 for n in A057486.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, May 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015190 | OEIS-A344146

a(n) is the number of pentanomials x^n + x^a + x^b + x^c + 1 that are irreducible over GF(2) for n > a > b > c > 0.

It is conjectured that a(n) > 0 for all n >= 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, May 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015191 | OEIS-A344176

Alternating parity Yellowstone permutation: a(n) = n if n <= 3, otherwise the smallest number not occurring earlier having at least one common factor with a(n-2), but none with a(n-1), with the condition that the terms in the sequence alternate parity.

Even with the alternating parity condition, the primes seem to appear in their natural order as in the Yellowstone permutation (conjecture 8 in A098550), except for 11 and 13, which appear in reverse order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enrique Navarrete_, May 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015192 | OEIS-A344177

Numbers m > 3 such that x^m + x^3 + 1 is irreducible over GF(2) while x^m + x^3 + x^2 + x + 1 = x^m + (x + 1)^3 is not.

Conjecture: Given e >= 0, odd numbers r, k > 0, a > 2^e*r*k, consider the following two statements:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, May 11 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015193 | OEIS-A344185

a(n) = A344141(n) - 2^n.

It is conjectured that no term can be of the form P_m(2^k), where P_m(x) = Product_{i>=0} (1 + x^(2^(d_i)))^(c_i) if the binary representation of m is m = Sum_{i>=0} c_i * 2^(d_i), k is an odd number. See my conjecture in A344177.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, May 11 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015194 | OEIS-A344186

a(n) = A344142(n) - 2^n.

It is conjectured that an irreducible polynomial of degree n with 5 terms exists for every n. It follows from the conjecture that for n >= 2, a(n) is of the form 2^k + 1 or an odd number with Hamming weight 4. It is conjectured that no term can be of the form P_m(2^k), where P_m(x) = Product_{i>=0} (1 + x^(2^(d_i)))^(c_i) if the binary representation of m is m = Sum_{i>=0} c_i * 2^(d_i), k is an odd number. See my conjecture in A344177.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, May 11 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015195 | OEIS-A344191

a(n) = Catalan(n) * (n^2 + 2) / (n + 2).

Conjecture: These are the number of linear intervals in Pallo's comb posets. An interval is linear if it is isomorphic to a total order. The conjecture has been checked up to the term 36686 for n = 9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _F. Chapoton_, May 11 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015196 | OEIS-A344197

Numbers m > 5 such that x^m + x^5 + 1 is irreducible over GF(2) while x^m + x^5 + x^4 + x + 1 = x^m + (x + 1)^5 is not.

Conjecture: Given e >= 0, odd numbers r, k > 0, a > 2^e*r*k, consider the following two statements:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, May 11 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015197 | OEIS-A344198

Numbers m > 9 such that x^m + x^9 + 1 is irreducible over GF(2) while x^m + x^9 + x^6 + x^3 + 1 = x^m + (x^3 + 1)^3 is not.

Conjecture: Given e >= 0, odd numbers r, k > 0, a > 2^e*r*k, consider the following two statements: If the conjecture is true, this is also numbers m >= 10 such that x^m + x^9 + 1 is irreducible over GF(2) while x^m + x^9 + x^8 + x + 1 is not (the case r = 9, k = 1, e = 0).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, May 11 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015198 | OEIS-A344199

Numbers m > 15 such that x^m + x^15 + 1 is irreducible over GF(2) while x^m + x^15 + x^10 + x^5 + 1 = x^m + (x^5 + 1)^3 is not.

Conjecture: Given e >= 0, odd numbers r, k > 0, a > 2^e*r*k, consider the following two statements:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, May 11 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015199 | OEIS-A344200

Numbers m > 15 such that x^m + x^15 + 1 is irreducible over GF(2) while x^m + x^15 + x^12 + x^3 + 1 = x^m + (x^3 + 1)^5 is not.

Conjecture: Given e >= 0, odd numbers r, k > 0, a > 2^e*r*k, consider the following two statements:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, May 11 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015200 | OEIS-A344216

a(n) = n!*((n+1)/2 + 2*Sum_{k=2..n-1}(n-k)/(k+1)).

Conjecture: a(n) is the number of linear intervals in the weak order on the symmetric group S_n. An interval is linear if it is isomorphic to a total order. The above conjecture has been proved by Chenevière in his Ph. D. thesis (see Links). - _Ludovic Schwob_, Dec 31 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _F. Chapoton_, May 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015201 | OEIS-A344228

a(n) = binomial(2*n,n)*(2*n+1)/2+n*binomial(2*n-2,n)+(n-1)*binomial(2*n-2,n+1).

Conjecture: These are the number of linear intervals in the Cambrian lattices of type B_n. An interval is linear if it is isomorphic to a total order. The conjecture has been checked up to the term 34254 for n = 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _F. Chapoton_, May 12 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015202 | OEIS-A344275

Denominators of worst rational approximations to the golden ratio.

Conjecture: difference of consecutive terms is always a Fibonacci number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, May 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015203 | OEIS-A344321

a(n) = 2^(2*n - 5)*binomial(n-5/2, -1/2)*(36*n^4 - 78*n^3 + 54*n^2 - 48*n + 24)/((n + 1)*n*(n - 1)) for n >= 2 and otherwise 1.

Conjecture: These are the number of linear intervals in the Cambrian lattices of type D_n. An interval is linear if it is isomorphic to a total order. The conjecture has been checked up to the term a(8) = 101398.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _F. Chapoton_, May 15 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015204 | OEIS-A344338

Smallest number that is the sum of two or more consecutive positive n-th powers in more than one way.

Conjecture: no terms exist for n >= 4. - _Jon E. Schoenfield_, May 16 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, May 15 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015205 | OEIS-A344388

Decimal expansion of a constant related to the asymptotics of A048634.

Conjecture: It is equal to the limit of column "h^2" in the Table 1 in reference by Wright and Trefethen, p. 336.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Aug 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015206 | OEIS-A344409

Positive discriminants of orders of quadratic fields with class number 3.

Conjecture: if a term k is congruent to 4 modulo 16, then k/4 is either here or in A133315; if a term k is congruent to 0 modulo 16, then k/4 is in this sequence. Conjecture: a term k is in A006832 if and only if k/4 is not in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, May 17 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015207 | OEIS-A344422

Palindromes having more divisors than all smaller palindromes.

Conjecture: all terms after a(1)=1 are even. (End) Conjecture: all terms after a(5)=44 are divisible by 3. - _Bhupendra Kumar Singh_, Feb 05 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bhupendra Kumar Singh_, May 17 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015208 | OEIS-A344427

Decimal expansion of -zeta'(alpha), where alpha = A069995 is the fixed point of Riemann zeta function in (1, +oo).

Moreover, let s_0 be any real number > alpha, s_n = zeta(s_{n-1}) for n >= 1, then it seems that ... > s_{2n} > s_{2n-2} > ... > s_2 > s_0 > alpha > s_1 > s_3 > ... > s_{2n+1} > ..., and {s_{2n}} diverges to +oo, {s_{2n+1}} converges to 1. Moreover, the divergence of {s_{2n}} and convergence of {s_{2n+1}} should be really fast, see my conjecture in A344428.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, May 19 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015209 | OEIS-A344428

Decimal expansion of exp(-2/5).

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, May 19 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015210 | OEIS-A344448

Square array read by antidiagonals upwards: T(n,k) for integer k >= 0 is the n-th prime p such that p^(2*3^k) + p^(3^k) + 1 is prime.

T(n,k)^(3^k), for all n >= 1, k >= 0, arranged by increasing values, is A342690. It is conjectured that all columns are infinite. If 3^k was replaced by k in the definition, all additional columns would be empty, as x^(2*k) + x^k + 1 is reducible if k has prime factors other than 3. For checking the property, Pocklington-Lehmer type primality tests seem particularly effective, as n-1 always has a large smooth factor p^(3^k), cf. the paper of Brillhart, Lehmer and Selfridge (1975), Theorem 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Martin Becker_, May 19 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015211 | OEIS-A344502

a(n) = Sum_{k=0..n} binomial(n, k)^2 * hypergeom([(k-n)/2, (k-n+1)/2], [k+2], 4).

Conjecture D-finite with recurrence -4*(3035*n-11997) *(2*n+1) *(n+1) *a(n) +2*(153546*n^3-490325*n^2-62942*n+71982) *a(n-1) +2*(-452090*n^3+1745622*n^2-1405285*n-226200) *a(n-2) -2 *(n-2)*(83741*n^2-200458*n-19650) *a(n-3) -3*(n-2) *(n-3) *(46423*n+6938) *a(n-4)=0. - _R. J. Mathar_, Nov 02 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, May 23 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015212 | OEIS-A344503

a(n) = Sum_{k=0..n} (-1)^(n-k)*binomial(n, k)^2*hypergeom([(k-n)/2, (k-n+1)/2], [k+2], 4).

Conjecture D-finite with recurrence -18*(2*n+1) *(2*n-1) *(n+1) *a(n) +2*(-36*n^3+554*n^2-1128*n+27) *a(n-1) +6*(-12*n^3-188*n^2+1235*n-1618) *a(n-2) +9*(54*n^3-27*n^2-183*n+320) *a(n-3) +54*(n-3) *(9*n^2-125*n+75) *a(n-4) +81 *(n-3) *(n-4) *(6*n+127) *a(n-5)=0. - _R. J. Mathar_, Nov 02 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, May 23 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015213 | OEIS-A344598

a(n) = Sum_{k=1..n} phi(k) * (floor(n/k)^2 - floor((n-1)/k)^2).

Conjecture: a(n) = Sum_{k = 1..2*n} (-1)^k * gcd(k, 4*n). Cf. A344372. - _Peter Bala_, Jan 01 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, May 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015214 | OEIS-A344611

Number of integer partitions of 2*n with reverse-alternating sum >= 0.

Conjecture: a(n) <= A160786(n). The difference is 0, 0, 0, 0, 1, 2, 4, 9, 16, 28, 48, 79, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, May 30 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015215 | OEIS-A344620

Primes p such that there is no positive integer a with 2*a, a^2-1 and a^2+1 not only smaller than p but also quadratic residues modulo p.

Conjecture: No term is greater than 827. In other words, for any prime p > 828, there is a Pythagorean triple (2*a, a^2-1, a^2+1) with 2*a, a^2-1 and a^2+1 in the set {0 < r < p: r is a quadratic residue modulo p}. See also A344621 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015216 | OEIS-A344621

Primes p such that there is no positive integer a with 2*a, a^2-1 and a^2+1 not only smaller than p but also quadratic nonresidues modulo p.

Conjecture: No term is greater than 3119. In other words, for any prime p > 3120, there is a Pythagorean triple (2*a,a^2-1,a^2+1) with 2*a, a^2-1 and a^2+1 in the set {0 < r < p: r is a quadratic nonresidue modulo p}. See also A344620 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015217 | OEIS-A344640

Antidiagonal sums of A344639.

Conjecture: a(n) ~ n!. - _Vaclav Kotesovec_, May 26 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, May 25 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015218 | OEIS-A344695

a(n) = gcd(sigma(n), psi(n)), where sigma is the sum of divisors function, A000203, and psi is the Dedekind psi function, A001615.

Conjecture: the set of primes that appear in the sequence is A065091 (the odd primes). 5 does not appear as a term until a(366025) = 5, where 366025 = 5^2 * 11^4. At this point, the missing numbers less than 22 are 2, 10 and 17. 17 appears at the latest by a(17^2 * 103^16) = 17.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_ and _Peter Munn_, May 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015219 | OEIS-A344710

a(n)/2 is the smallest possible area of a non-obtuse triangle with coordinates in Z^2 and no side shorter than sqrt(n).

For [b(n)], it is conjectured that 1/x(n) is always an integer, so the ceiling function can be omitted. Conjectured next terms (from the conjectured inequality a(n) > sqrt(3/4)*n) are a(50,...,64) ?= 45, 48, 48, 52, 55, 55, 55, 55, 55, 56, 56, 56, 56, 56, 56. a(n) > sqrt(3/4)*n (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan F. Waldmann_, May 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015220 | OEIS-A344717

a(n) = (3n - 9/2 - 1/n + 6/(n+1))*binomial(2n-2,n-1).

Conjecture: These are the number of linear intervals in the tilting posets of type B_n. An interval is linear if it is isomorphic to a total order. The conjecture has been checked up to the term 295867 for n = 9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _F. Chapoton_, May 27 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015221 | OEIS-A344719

Minimum number of cells in an n X n grid that need to be painted so that each unpainted cell is orthogonally adjacent to exactly two painted cells.

Conjecture: for n >= 6 there is an optimal solution in which every row except the top 2 and bottom 2 rows, ignoring the 2 leftmost and 2 rightmost columns, paints exactly the cells in columns congruent to some s (mod 3). Verified for 6 <= n <= 23. Conjecture: a(n) = ceiling((n+2)^2/3) - 6 for n >= 7. Conjectured g.f.: x*(1 + 2*x^2 - x^3 + x^5 - 2*x^6 + 3*x^7 - 3*x^8 + 2*x^9 - x^10)/((1-x)^3*(1+x+x^2)). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, May 28 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015222 | OEIS-A344728

a(n) = (9*n/4 - 51/8 - 5/(16*n-24) + 1/n + 6/(n+1))*binomial(2*n-2,n-1).

Conjecture: a(n) is the number of linear intervals in the tilting posets of type D_n. An interval is linear if it is isomorphic to a total order. The conjecture has been checked up to the term 187187 for n = 9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _F. Chapoton_, May 27 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015223 | OEIS-A344734

a(n) is the smallest divisibility checking number for the n-th number coprime to 10.

Conjecture: the subsequence of divisibility checking numbers for numbers that end in 1 form an arithmetic sequence. This is also the case for numbers ending in 3, 7, and 9. Conjectures from _Chai Wah Wu_, Dec 29 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zach Pollard_, May 27 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015224 | OEIS-A344743

Number of integer partitions of 2n with reverse-alternating sum < 0.

Conjecture: a(n) >= A236914.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jun 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015225 | OEIS-A344810

Numbers that are the sum of six squares in six or more ways.

Conjectures from _Chai Wah Wu_, Apr 25 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Consiglio, Jr._, Jun 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015226 | OEIS-A344812

Numbers that are the sum of six squares in eight or more ways.

Conjectures from _Chai Wah Wu_, Jan 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Consiglio, Jr._, Jun 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015227 | OEIS-A344845

a(n)/2 is the smallest possible area of a non-obtuse triangle with integer coordinates in the plane and with shortest side of length sqrt(A001481(n+1)).

For [b(n)], it is conjectured that 1/x(n) is always an integer, making the ceiling function omittable for the definition. a(n) > sqrt(3/4)*m (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jonathan F. Waldmann_, May 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015228 | OEIS-A344857

Number of polygons formed when every pair of vertices of a regular n-gon are joined by an infinite line.

For odd n, a(n) = (n^4 - 7*n^3 + 19*n^2 - 21*n + 8)/8 = (n-1)^2*(n^2-5*n+8)/8. This was conjectured by _Scott R. Shannon_ and proved by Alexander Sidorenko on Sep 10 2021 (see link). - _N. J. A. Sloane_, Sep 12 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 30 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015229 | OEIS-A344899

Number of polygon edges formed when every pair of vertices of a regular n-gon are joined by an infinite line.

Conjectured formula odd n: a(n) = (n^4 - 7*n^3 + 17*n^2 - 11*n)/4 = (n-1)*n*(n^2-6*n+11)/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 02 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015230 | OEIS-A344902

Number of open tours by a biased rook on a specific f(n) X 1 board, where f(n) = A070941(n) and cells are colored white or black according to the binary representation of 2n.

Conjecture: a(n) = A284005(A073138(n)) for n >= 0 (noticed by Sequence Machine).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Jun 01 2021 [verification needed] | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015231 | OEIS-A344960

Positions in A341258 of the binary complement of the n-th word in A341258.

Conjecture: a(n) = A345253(A348366(n+1))-1 (noticed by Sequence Machine). - _Mikhail Kurkov_, Nov 28 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015232 | OEIS-A344962

a(n) is the position in A341258 of the reversal of the n-th word in A341258.

A permutation of the positive integers. Conjecture: a(n) = A344682(n) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015233 | OEIS-A345003

Numbers k for which A344998(k) = A344999(k).

Conjecture: Sequence is a disjoint union of A000396 and A301939.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jun 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015234 | OEIS-A345005

Odd numbers whose arithmetic derivative (A003415) is equal to Dedekind psi (A001615) applied to the same number.

Conjectured to be also the odd numbers k for which A344998(k) = A344999(k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jun 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015235 | OEIS-A345013

Triangle read by rows, related to clusters of type D.

As conjectured above by Chapoton we have

猜想/问题提及段 | historical_or_current_mention_unreviewed | _F. Chapoton_, Sep 30 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015236 | OEIS-A345051

Numbers k such that A345048(k) is equal to A345049(k).

Conjecture: Sequence is a disjoint union of A000396 and A166374, i.e., there are no terms of any other kind.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jun 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015237 | OEIS-A345053

Positions of zeros in A345055, which is the Dirichlet inverse of A011772.

A345055(d)=0 for the other proper divisors d of 2^i by the inductive hypothesis as d is a power of 2 and this implies A345033(2^i)=0 for i>=3. Conjecture: all terms are of the form 2^i, 2^i*p, 2^i*p*q or 7^2*p for some primes p and q. - _Chai Wah Wu_, Jul 05 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jul 01 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015238 | OEIS-A345079

Consider the coefficients in the expansion of the n-th cyclotomic polynomial. a(n) is the difference between the extremes.

Conjecture: every number occurs in this sequence. This is based on the fact that every integer is a coefficient in the expansion of a cyclotomic polynomial. See Chun-Gang Ji and Wei-Ping Li link below.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_ and _Robert G. Wilson v_, Jun 04 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015239 | OEIS-A345096

a(n) = Sum_{k=1..n} Sum_{i=1..k} k^(1 - ceiling(n/k^i) + floor(n/k^i)).

Conjecture: a(n) ~ n^2/2. - _Vaclav Kotesovec_, Jun 08 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Jun 07 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015240 | OEIS-A345114

Numbers whose trajectories under the map x -> A345111(x) do not reach a palindrome (conjectured).

Numbers whose trajectories under the map x -> A345111(x) do not reach a palindrome (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Jun 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015241 | OEIS-A345128

Number of squarefree products s*t from all positive integer pairs (s,t), such that s + t = n, s <= t.

Conjecture: a(n) > 0 for all n > 1. (End) A simple case of Zhou's conjecture: a(p) > 0 where p is prime. With some work this can be extended to a(n) > 0 for n with sum_{p | n} 1/p < 1/10 or so (the limit of the method is 6/Pi^2 - 1/2, so it can't prove the full conjecture). See my comment in A071068. - _Charles R Greathouse IV_, Dec 20 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Jun 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015242 | OEIS-A345135

Number of ordered rooted binary trees with n leaves and with minimal Sackin tree balance index.

Conjecture: Sum_{n>=0} a(n)*x^n = 1 + x + Sum_{n>=0} x^(2^n) * ((1 + x)^(2^n) - 1). - _Paul D. Hanna_, Jul 18 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mareike Fischer_, Jun 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015243 | OEIS-A345211

Numbers with the same number of odd / even, refactorable divisors.

All the terms are even. Conjecture: The sequence a(n)/2 is the complement of A157932 (verified for a(n) < 4*10^7). - _Amiram Eldar_, Oct 11 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Jun 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015244 | OEIS-A345255

Carter and Reyes sequence for the game of countably infinitely many hats.

The Carter and Reyes sequence represents a strategy for the game of countably infinitely many hats. This strategy is conjectured to be one of several optimal strategies. Actually it is possible to achieve a probability of success of 7/20 (35%) using the Carter and Reyes strategy described here, the Tiefenbruck strategy described in A345257, or variations thereof. Also it is an open problem whether strategies can achieve results better than 7/20.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frederic Ruget_, Jun 12 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015245 | OEIS-A345257

Tiefenbruck sequence for the game of countably infinitely many hats.

The Tiefenbruck sequence represents a strategy for the game of countably infinitely many hats. This strategy is conjectured to be one of several optimal strategies. Actually it is possible to achieve a probability of success of 7/20 (35%) using the Tiefenbruck strategy described here, the Carter and Reyes strategy described in A345255, or variations thereof. Also it is an open problem whether strategies can achieve results better than 7/20.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frederic Ruget_, Jun 12 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015246 | OEIS-A345262

a(n) is the order of the image of the J-homomorphism in the stable homotopy groups of spheres.

Im(J) is a finite cyclic subgroup of Pi_n^S and has known order a(n) calculated by Adams using the Adams conjecture, subsequently proven by Quillen. When n is 3 or 7 mod 8 the value a(n) is related to the Bernoulli numbers; the other values of a(n) are 8-periodic (after an exceptional n=0).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tom Harris_, Jun 12 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015247 | OEIS-A345370

a(n) is the number of distinct numbers of diagonal transversals that a diagonal Latin square of order n can have.

Conjecture: a(12) = A287648(12) - A287647(12) + 1. - _Natalia Makarova_, Oct 26 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eduard I. Vatutin_, Jun 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015248 | OEIS-A345397

a(n) is the least number k such that {k, k^2, ..., k^n} are all digitally balanced numbers in base 2 (A031443), or 0 if no such k exists.

Conjecture: all a(n) > 0. - _Alex Ratushnyak_, Apr 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Jun 17 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015249 | OEIS-A345453

Triangle of optimist numbers T(n,k) (n >= 1, 0 <= k <= n-1) read by rows: permutations needing k steps to be sorted by the "optimist" algorithm.

Conjecture: each row is unimodal.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Olivier Gérard_, Jun 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015250 | OEIS-A345476

Numbers that are the sum of six squares in nine or more ways.

Conjectures from _Chai Wah Wu_, Jan 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Consiglio, Jr._, Jun 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015251 | OEIS-A345477

Numbers that are the sum of six squares in ten or more ways.

Conjectures from _Chai Wah Wu_, Jan 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Consiglio, Jr._, Jun 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015252 | OEIS-A345482

Numbers that are the sum of seven squares in five or more ways.

Conjectures from _Chai Wah Wu_, Apr 25 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Consiglio, Jr._, Jun 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015253 | OEIS-A345483

Numbers that are the sum of seven squares in six or more ways.

Conjectures from _Chai Wah Wu_, Apr 25 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Consiglio, Jr._, Jun 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015254 | OEIS-A345484

Numbers that are the sum of seven squares in seven or more ways.

Conjectures from _Chai Wah Wu_, Apr 25 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Consiglio, Jr._, Jun 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015255 | OEIS-A345486

Numbers that are the sum of seven squares in nine or more ways.

Conjectures from _Chai Wah Wu_, Jan 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Consiglio, Jr._, Jun 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015256 | OEIS-A345487

Numbers that are the sum of seven squares in ten or more ways.

Conjectures from _Chai Wah Wu_, Jan 05 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Consiglio, Jr._, Jun 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015257 | OEIS-A345493

Numbers that are the sum of eight squares in six or more ways.

Conjectures from _Chai Wah Wu_, Apr 25 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Consiglio, Jr._, Jun 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015258 | OEIS-A345503

Numbers that are the sum of nine squares in six or more ways.

Conjectures from _Chai Wah Wu_, Apr 25 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Consiglio, Jr._, Jun 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015259 | OEIS-A345531

Smallest prime power greater than the n-th prime.

Take the family of correlated prime-indexed conjectures appearing in A343249 - A343253, in which an alternative formula for the p-adic order of positive integers is proposed. There, the general p-indexed conjecture says that v_p(n), the p-adic order of n, is given by the formula: v_p(n) = log_p(n / L_p(k0, n)), where L_p(k0, n) is the lowest common denominator of the elements of the set S_p(k0, n) = {(1/n)*binomial(n, k), with 0 < k <= k0 such that k is not divisible by p}. Evidence suggests that the primality of p is a necessary condition in this general conjecture. So, if a composite number q is used instead of a prime p in the proposed formula for the p-adic (now, q-adic) order of n, the first counterexample (failure) is expected to occur for n = q * a(i), where i is the index of the smallest prime that divides q.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dario T. de Castro_, Jun 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015260 | OEIS-A345647

Square array read by downward antidiagonals: A(n, k) = number of primes in the interval [n+1, n+k], n >= 1, k > 1.

Conjecture: If A(n, k) = 0, then A001221(Product_{x=1..k}(n+x)) >= k (cf. Wikipedia).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Fröhlich_, Jun 21 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015261 | OEIS-A345651

Fourth column of A008296.

Conjecture: D-finite with recurrence a(n) +2*(2*n-13)*a(n-1) +(6*n^2-84*n+295)*a(n-2) +(2*n-15)*(2*n^2-30*n+113)*a(n-3) +(n-8)^4*a(n-4)=0. - _R. J. Mathar_, Sep 15 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luca Onnis_, Aug 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015262 | OEIS-A345652

Expansion of the e.g.f. exp(-1 + (1 + x)*exp(-x)).

Conjecture: a(n) = 0 for only n = 1 and n = 4. Conjecture: For all p prime, a(p)^2 == 1 (mod p). Stronger conjecture: For n > 1, a(n) == -1 (mod n) iff n is a prime or 6. - _M. F. Hasler_, Jun 23 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mélika Tebni_, Jun 21 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015263 | OEIS-A345680

Nonnegative integers whose trajectory under iteration of taking the absolute value of the alternating sum of the squares of the digits (A257588) includes zero.

Conjecture: there are strings of consecutive terms of arbitrary length.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stephen Cross_, Jun 23 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015264 | OEIS-A345716

a(n) is the minimum number of nonattacking rooks required to attack every square on a size n right tetrahedral chessboard.

This sequence has been verified by brute force up to n = 27. It matches the covering numbers C(n+1, 3,2), A011975, for the first 14 terms, and it is conjectured that it matches for all even n. Link to the brute force program below. Computational runtime and increased complexity for minimum arrangements for bigger n account for the low number of terms. Other contributers: Fletcher Collins, Jennifer Gensler, Nick Jamesson, Chase Schwartzman, Evan

Young, Dr. Nat Theim. For n == 0,2 (mod 6): a(n) = n*(n+1)/6 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Connor Mason Weisburg_, Jun 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015265 | OEIS-A345754

Number of 2 X 2 matrices over Z_n whose permanent equals their determinant.

a(n) = A344372(n) * n^2 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Jul 19 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015266 | OEIS-A345877

a(1) = 1, a(n) = a(n-1)/2 if a(n-1) is even, otherwise a(n) = n - a(n-1).

Conjecture: a_i(n) hits every positive integers infinitely many times for all i >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Jun 28 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015267 | OEIS-A345928

Decimal expansion of Integral_{x>=0} (zeta(x)-1) dx (negated).

Robinson (1980) conjectured and Newman and Widder (1981) proved that this integral is equal to the limit given in the Formula section.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Jun 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015268 | OEIS-A345938

a(n) = uphi(n) / gcd(n-1, uphi(n)), where uphi is unitary totient (or unitary phi) function, A047994.

For all squarefree n (A005117), a(n) = A160595(n), thus if there are any composite solutions to the Lehmer's totient conjecture, then they give also a such a subset of positions of 1's in this sequence that are not powers of primes. See comments in A160595.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jun 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015269 | OEIS-A345997

Let m = A344005(n) be the smallest number such that n|m*(m+1); let X = A345992(n) = gcd(n,m); Y = A345993(n) = gcd(n,m+1). Sequence lists n such that neither X nor Y is equal to n/p^k, where p = largest prime divisor of n and k is its exponent in n.

Numbers n such that neither A345992(n) nor A345993(n) is equal to A051119(n). This disproves an obvious conjecture about A344005.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Dougherty-Bliss_ and _N. J. A. Sloane_, Jul 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015270 | OEIS-A346040

a(n) is 1w' converted to decimal, where the binary word w' is the result of applying Post's tag system {00,1101} to the binary word w, where 1w is n converted to binary (the leftmost 1 acts as a delimiter).

It is an important open question to decide whether there is any word whose orbit grows without limit.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carlos Gómez-Ambrosi_, Jul 02 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015271 | OEIS-A346061

A(n,k) = n! * [x^n] (Sum_{j=0..n} k^(j*(j+1)/2) * x^j/j!)^(1/k) if k>0, A(n,0) = 0^n; square array A(n,k), n>=0, k>=0, read by antidiagonals.

A(n,k) == 1 (mod k*(k-1)) for k >= 2 (see "general conjecture" in A178319 and link to proof by _Richard Stanley_ above).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 03 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015272 | OEIS-A346063

a(n) = primepi(A039634(prime(n)^2-1)).

This sequence looks at the effect on p^2 - 1 of A039634 with the primes represented by their indices. It seems that primes obtained by iterating the map A039634 on p^2 - 1 never fall into a cycle before reaching 2. Conjecture: Iterating the map k -> a(k) eventually reaches 1. For example, 1 -> 2 -> 1; 5 -> 4 -> 2 -> 1; and 27 -> 67 -> 16 -> 14 -> 4 -> 2 -> 1. If the conjecture holds, then A339991(n) != -1 and A340419 is a finite sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Jul 03 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015273 | OEIS-A346064

Number of primes that may be generated by changing any two digits of n simultaneously.

Indices of low records are given by A345289. By heuristic considerations it is conjectured that a(n) > 0 for all n >= 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Franz Vrabec_, Jul 03 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015274 | OEIS-A346074

a(n) = 1 + Sum_{k=0..n-5} a(k) * a(n-k-5).

Conjecture D-finite with recurrence (n+5)*a(n) +2*(-n-4)*a(n-1) +(n+3)*a(n-2) +2*(-2*n+5)*a(n-5) +4*(n-3)*a(n-6)=0. - _R. J. Mathar_, Feb 17 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Jul 04 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015275 | OEIS-A346115

Least number k such that k^2 can be expressed in exactly n ways as x^2 + y^4 with {x, y} >= 1.

These two conjectured values arise from the "green group". Up to term a(5) the least solutions are in the "blue group". Follow the links below to get more information about the different colored groups.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Karl-Heinz Hofmann_, Jul 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015276 | OEIS-A346119

Expansion of the e.g.f. sqrt(2*x*exp(x) - 2*exp(x) + 3).

Conjecture: a(n) = 0 for only n = 1 and n = 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mélika Tebni_, Jul 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015277 | OEIS-A346126

Numbers m such that no self-avoiding walk of length m + 1 on the hexagonal lattice fits into the smallest circle that can enclose a walk of length m.

Open and closed walks are allowed. It is conjectured that all optimal paths are closed except for the trivial path of length 1. See the related conjecture in A122226.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_ and _Markus Sigg_, Jul 31 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015278 | OEIS-A346149

a(n) is the least integer k > 1 such that n^k + n + 1 is prime, or 0 if there is no such k.

Conjecture: a(n) > 0 otherwise.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Jul 07 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015279 | OEIS-A346154

a(n) is the least prime of the form n^k+n+1 with k>1, or 0 if there is no such prime.

Conjecture: a(n) > 0 otherwise.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Jul 07 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015280 | OEIS-A346162

a(n) = (-1)^n*permanent[tan(Pi*(j+k)/(2n+1))]_{1<=j,k<=2n}.

Conjecture 1: a(n)/(2n+1) is always a positive integer congruent to 1 modulo 4. Conjecture 2: We have (n+1)*a(n) == 0 (mod (2n+1)!!) for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 22 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015281 | OEIS-A346211

Numbers m such that abs(K(m+1) - K(m)) = 1, where K(m) = A002034(m) is the Kempner function.

Florentin Smarandache (Muller, 1990) and Tutescu (1996) conjectured that there is no number m such that K(m) = K(m+1), and Weisstein (2004) verified it up to m = 10^9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Jul 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015282 | OEIS-A346212

Numbers m such that abs(K(m+1) - K(m)) = 2, and both m and m+1 are squarefree (A005117), where K(m) = A002034(m) is the Kempner function.

Since there are also no solutions to K(m) = K(m+1) below 10^9, it can be conjectured that the minimal difference abs(K(m+1) - K(m)) between consecutive numbers m and m+1 that are both squarefree is 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Jul 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015283 | OEIS-A346214

Triangular array read by rows. T(n,k) is the number of nilpotent n X n matrices over GF(2) with index k, 1 <= k <= n, n >= 1.

Define the co-index of an n X n matrix A to be n - index(A). Let X_n be the random variable that assigns to each nilpotent n X n matrix over GF(2) the value j in {0,1,...,n-1} of its co-index. Conjecture: lim_{n->inf} P(X_n = j) = Product_{i>=1}1-1/2^i * 2^((j-1)^2)/A002884(j). Moreover, for j < 2n, T(n,n-j) = A002884(n)/(A002884(j)*2^(n - (j-1)^2)). - _Geoffrey Critzer_, Jun 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Critzer_, Jul 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015284 | OEIS-A346275

Number of n-digit primes with digital product = 3.

Conjecture: for n >= 4, there is at least one composite number among the n-digit numbers with digital product = 3 <==> a(n) < n for n >= 4 (guess proposed by Derek Holton in reference).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Jul 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015285 | OEIS-A346298

a(n) is the smallest nonnegative number not in a(0..n-1) such that the sequence a(0..n) forms the starting row of an XOR-triangle with only distinct values in each row.

This sequence is conjectured to be a permutation of the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Jul 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015286 | OEIS-A346310

Positions of words in A076478 such that #0's - #1's is even.

a(n) = A053738(n+1) - 1, conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 28 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015287 | OEIS-A346370

Upper bound for the number of solutions of the TRINTUM cube puzzle n X 1 X 1 (cuboid formed by 4n + 2 parts) different by the set of parts, which are distinguished by the amount of surface area they contribute to the assembled cuboid.

Conjecture: upper bound can be reduced to n+2 (based on attempts to construct cuboid using nonnegative integer solutions with m > 0, where it looks like impossible to place at least all roosters and towers somewhere). Conjecture: a(n) = A008733(n-1) + A093907(n) for n > 0 (noticed by Sequence Machine). - _Mikhail Kurkov_, Oct 14 2021 [verification needed]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Jul 14 2021 [verification needed] | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015288 | OEIS-A346374

a(n) = f(n,n) where f(0,n) = f(n,0) = n! and f(m,n) = f(m-1,n) + f(m,n-1) + f(m-1,n-1).

a(n+1)/a(n) ~ n (conjectured). Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Jul 14 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015289 | OEIS-A346379

The least k such that the number of real roots of Sum_{i=0..j} (i+1)^k*x^i is j for j = 0,1,...,n or -1 if no solution for this n exists.

Conjecture: The sequence is defined for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _W. Edwin Clark_, Jul 14 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015290 | OEIS-A346385

a(n) = f(n,n) where f(0,n) = f(n,0) = n^n and f(m,n) = f(m-1,n) + f(m,n-1) + f(m-1,n-1).

a(n+1)/a(n) ~ e*n (conjectured). Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Jul 14 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015291 | OEIS-A346387

Number of permutations f of {1,...,prime(n)-1} with f(prime(n)-1) = prime(n)-1 and f(prime(n)-2) = prime(n)-2 such that 1/(f(1)*f(2)) + 1/(f(2)*f(3)) + ... + 1/(f(prime(n)-2)*f(prime(n)-1)) + 1/(f(prime(n)-1)*f(1)) == 0 (mod prime(n)^2).

Conjecture: a(n) > 0 for all n > 2. In other words, for any prime p > 3, there is a permutation f of {1,...,p-1} with f(p-1) = p-1 and f(p-2) = p-2 such that 1/(f(1)*f(2)) + 1/(f(2)*f(3)) + ... + 1/(f(p-2)*f(p-1)) + 1/(f(p-1)*f(1)) == 0 (mod p^2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 15 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015292 | OEIS-A346391

Number of permutations f of {1,...,n} with f(n) = n and f(n-1) > f(1) such that f(1)*f(2) + ... + f(n-1)*f(n) + f(n)*f(1) == 0 (mod n^2).

Conjecture: a(n) > 0 for all n > 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 15 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015293 | OEIS-A346401

a(n) is the minimum number of pebbles such that any assignment of those pebbles on a complete graph with n vertices is a next-player winning game in the two-player impartial (3, 2) pebbling game.

a(n) = 2n+3 when n >= 7 is odd (conjectured). a(n) = 2n+9 when n >= 6 is even (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kayla Barker_, _Mia DeStefano_, _Eugene Fiorini_, _Michael Gohn_, _Joe Miller_, _Jacob Roeder_, _Wing Hong Tony Wong_, Jul 15 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015294 | OEIS-A346501

With p = prime(n), a(n) is the least composite k such that A001414(k) = p and k+p is prime, or 0 if there is no such k.

Conjecture: a(n) > 0 for n >= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Jul 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015295 | OEIS-A346507

Positive integers k that are the product of two integers greater than 1 and ending with 1.

Conjecture: lim_{n->oo} a(n)/a(n-1) = 1. The conjecture is true since it can be proved that a(n) = (sqrt(a(n-1)) + g(n-1))^2 where [g(n): n > 1] is a bounded sequence of positive real numbers. - _Stefano Spezia_, Aug 21 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jul 21 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015296 | OEIS-A346508

Positive integers k such that 10*k+1 is equal to the product of two integers greater than 1 and ending with 1 (A346507).

Conjecture: lim_{n->oo} a(n)/a(n-1) = 1. The conjecture is true since a(n) = (A346507(n) - 1)/10 and lim_{n->oo} A346507(n)/A346507(n-1) = 1. - _Stefano Spezia_, Aug 21 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jul 21 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015297 | OEIS-A346509

Number of positive integers with n digits that are the product of two integers greater than 1 and ending with 1.

Conjecture: lim_{n->oo} a(n)/a(n-1) = 10. The conjecture is true since the sequence A346507 has positive asymptotic density (see Corollary 1 in Spezia link). - _Stefano Spezia_, Mar 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jul 21 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015298 | OEIS-A346575

a(n) is the number of 6-tuples (a_1,a_2,a_3,a_4,a_5,a_6) having all terms in {1,...,n} such that there exists a tetrahedron ABCD with those edge-lengths.

Conjecture 1: a(n)/n^6 tends to a limit which is 0.338170 +- 0.000017 (confidence level 95%). This number has been evaluated with a Monte-Carlo test on 3 billion sextuples with random values in (0,1) which simulate n -> oo. Conjecture 2: there is no polynomial formula for a(n), as finite difference method fails. Conjecture: Limit_{n->oo} a(n)/n^6 exists and is approximately 0.33817.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Corbelli_, Jul 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015299 | OEIS-A346608

Indices k such that A047994(k) != A344005(k).

Conjectures: (i) For all k in this sequence, A047994(k) >= A344005(k). The second conjecture asserts that this is equal to A265128 with {0, 1, 18} deleted. I believe I have a proof of both conjectures, although I have not yet written out all the details. Numbers k that are in A265128, but do not appear here are: 1, 18, 50, 54, 98, 162, 242, 250, 338, 486, 578, 686, ... probably given by {1} UNION A354929. Hence conjecture: the sequence consists of numbers that are neither a power of prime, or 2 * power of prime. - _Antti Karttunen_, Jun 14 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015300 | OEIS-A346643

Number of ways to write n as w^2 + 2*x^2 + 3*y^4 + 4*z^4, where w,x,y,z are nonnegative integers.

1-2-3-4 Conjecture: a(n) > 0 except for n = 158. See also A347865 and A350857 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015301 | OEIS-A346644

Least k >= 1 such that sigma(k)/tau(k) has denominator n or zero if no k exists.

Conjecture: k always exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Jul 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015302 | OEIS-A346645

Mersenne prime exponents p which are twin primes, so p-2 and/or p+2 is prime.

As of July 26, 2021, the sequence of Mersenne prime exponents with a twin prime above the exponent (A087858) contains 5 terms, whereas the sequence of Mersenne prime exponents with a twin prime below the exponent (A297674) contains 12 terms, which is 2.4x as many as the former sequence. This is an empirical observation. It is uncertain whether either sequence will increase in length, but given that it is conjectured that there are infinitely many twin primes, and also conjectured there are an infinite number of Mersenne primes, both sequences may increase in length, in which case this sequence might too.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David R. Kirkby_, Jul 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015303 | OEIS-A346669

Numbers r such that the number of nonnegative m < r such that m^k == m (mod r) is equal to k*(the number of nonnegative m < r such that -m^k == m (mod r)), where k = 2^A007814(r-1) + 1.

Conjecture: this sequence contains odd prime numbers A065091, but does not contain Carmichael numbers A002997. Conjecture: Let v(n) = A007814(n) be the 2-adic valuation of n. Define The conjecture is correct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jul 28 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015304 | OEIS-A346675

First differences of A088176.

Conjecture: each term > 4 appears at least twice.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luca Santarsiero_, Jul 28 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015305 | OEIS-A346692

a(n) = phi(n) - phi(n-phi(n)), a(1) = 1.

P. Erdős conjectured that a(n) > 0 on a set of asymptotic density 1, then Luca and Pomerance proved this conjecture (see link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Jul 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015306 | OEIS-A346695

Numbers with more divisors than digits in their binary representation.

It is natural to conjecture that this sequence has asymptotic density 0. However, after the first three terms where a(n)/n = 6 -- a function which would increase to infinity if the asymptotic density were zero -- it drops, and it seems to take a long time to get that high again. The first time it gets above 5.0 is at a(30243)=151216. Even as high as a(2188516)=10000000, the density is only ~1/4.57.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Meiburg_, Jul 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015307 | OEIS-A346696

a(n) is the least positive k such that A000041(n) divides A000041(n+k), or 0 if no such k exists.

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Altug Alkan_, Jul 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015308 | OEIS-A346778

Least k >= 1 such that {b(1), b(2), ..., b(k)} = {n, n-1, ..., n-k+1} and b(k+1) = n-k where b(1..n) is row n of A088643, or k = 0 if no such k >= 1 exists.

Since any row n of A088643 is a permutation of [1..n] having 1 as last term (conjectured), one always has a(n) <= n - 1. - _M. F. Hasler_, Aug 04 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sean A. Irvine_, Aug 03 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015309 | OEIS-A346784

Numerators of minimal squared radii of circular disks covering a record number of lattice points of the hexagonal lattice, when the centers of the circles are chosen to maximize the number of covered lattice points.

It is conjectured that the number of covered grid points is given by A346126(n-1) for n>2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Aug 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015310 | OEIS-A346802

Number of ways to start with set {1,2,...,n} and then repeat (n+1) times: partition each set into subsets.

Conjecture: a(n) ~ c * n^(2*n - 5/6) / (exp(n) * 2^n), where c = 42.345... - _Vaclav Kotesovec_, Aug 11 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Aug 04 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015311 | OEIS-A346807

Numbers that are the sum of ten squares in eight or more ways.

a(n) = n + 64 for n >= 8 (conjectured). - _Chai Wah Wu_, Dec 05 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Consiglio, Jr._, Aug 04 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015312 | OEIS-A346918

a(n) is 1 plus the number of iterations of the map prime(n)# + k -> prime(n)# + lpf(prime(n)# + k) required to reach a prime, starting at k=1, where prime(n)# is the n-th primorial and lpf() is the least prime factor, or 0 if no prime is ever reached.

Open question: Are there infinitely many n such that a(n)=0?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kim Hollesen_, Aug 07 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015313 | OEIS-A346934

Permanent of the 2n X 2n matrix with the (i,j)-entry i-j (i,j=1..2n).

Conjecture 1: (-1)^n*a(n) > 0 for all n > 0. Also, a(n) == 0 (mod 2n+1) if 2n+1 is composite. Conjecture 2: For any odd prime p, the sum of those 1/Product_{j=1..p-1}(j-f(j)) with f in the set D(p-1) is congruent to (-1)^((p-1)/2) modulo p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015314 | OEIS-A346952

Number of positive integers with n digits that are the product of two integers ending with 3.

Conjecture: lim_{n->oo} a(n)/a(n-1) = 10. The conjecture is true since the sequence A346950 has positive asymptotic density (see Corollary 1 in Spezia link). - _Stefano Spezia_, Mar 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Aug 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015315 | OEIS-A346973

a(n) = a(n-1) mod a(n-8) + a(n-2) mod a(n-7) + a(n-3) mod a(n-6) + a(n-4) mod a(n-5) where first 8 terms are 1, 2, 3, 4, 5, 6, 7, 8.

Conjecture: a(n)^(1/n) tends to c, where 1.11675 < c < 1.11690. - _Nicolas Bělohoubek_ and _Vaclav Kotesovec_, Aug 18 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nicolas Bělohoubek_, Aug 18 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015316 | OEIS-A346982

Expansion of e.g.f. 1 / (4 - 3 * exp(x))^(1/3).

O.g.f. (conjectural): 1/(1 - x/(1 - 4*x/(1 - 4*x/(1 - 8*x/(1 - 7*x/(1 - 12*x/(1 - ... - (3*n-2)*x/(1 - 4*n*x/(1 - ...)))))))) - a continued fraction of Stieltjes-type (S-fraction).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Aug 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015317 | OEIS-A346983

Expansion of e.g.f. 1 / (5 - 4 * exp(x))^(1/4).

O.g.f. (conjectural): 1/(1 - x/(1 - 5*x/(1 - 5*x/(1 - 10*x/(1 - 9*x/(1 - 15*x/(1 - ... - (4*n-3)*x/(1 - 5*n*x/(1 - ...)))))))) - a continued fraction of Stieltjes-type. - _Peter Bala_, Aug 22 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Aug 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015318 | OEIS-A346984

Expansion of e.g.f. 1 / (6 - 5 * exp(x))^(1/5).

O.g.f. (conjectural): 1/(1 - x/(1 - 6*x/(1 - 6*x/(1 - 12*x/(1 - 11*x/(1 - 18*x/(1 - ... - (5*n-4)*x/(1 - 6*n*x/(1 - ...)))))))) - a continued fraction of Stieltjes-type. - _Peter Bala_, Aug 22 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Aug 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015319 | OEIS-A346985

Expansion of e.g.f. 1 / (7 - 6 * exp(x))^(1/6).

O.g.f. (conjectural): 1/(1 - x/(1 - 7*x/(1 - 7*x/(1 - 14*x/(1 - 13*x/(1 - 21*x/(1 - ... - (6*n-5)*x/(1 - 7*n*x/(1 - ...)))))))) - a continued fraction of Stieltjes-type (S-fraction). - _Peter Bala_, Aug 25 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Aug 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015320 | OEIS-A347009

a(n) is the largest integer m such that e^-m > e - Sum_{k=0..n} 1/k!.

Conjecture: a(n) = floor((n + 3/2)*log(n) - n - zeta'(0) + (1/12)/n + 1/n^2) for n >= 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Fred Patrick Doty_, Aug 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015321 | OEIS-A347018

The number of unlabeled trees T on n vertices for which maximum multiplicity attained by any matrix whose graph is T implies the simplicity of its other eigenvalues.

Conjecture: For n>15, a(n) = 2*a(n-1) + a(n-2) - 3*a(n-3) + 2*a(n-4) - a(n-5) - 2*a(n-6) + a(n-7) + 3*a(n-8) - 4*a(n-9) - a(n-10) + 2*a(n-11) - 2*a(n-12) + 2*a(n-13) + a(n-14) - a(n-15), with initial terms a(1) through a(15) as given. If the above conjecture holds, then a(n) has o.g.f. (x - x^2 - 2*x^3 + 2*x^4 - x^5 - x^6 + 2*x^7 - 3*x^9 + 2*x^10 + x^11 - x^12 + 2*x^13 - x^15)/(1 - 2*x - x^2 + 3*x^3 - 2*x^4 + x^5 + 2*x^6 - x^7 - 3*x^8 + 4*x^9 + x^10 - 2*x^11 + 2*x^12 - 2*x^13 - x^14 + x^15). If the above conjecture

holds, then a(n) has an explicit Binet-like formula that can be obtained in terms of the roots of a fifth-degree polynomial.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Greyson C. Wesley_, Aug 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015322 | OEIS-A347063

Double Zumkeller numbers: numbers whose set of divisors can be partitioned into two disjoint subsets with equal sums and equal cardinalities.

If x is a Zumkeller number, then so is 2x. Conjecturally, if y is a term of this sequence, then so is 2y.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ivan N. Ianakiev_, Aug 15 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015323 | OEIS-A347171

Triangle read by rows where T(n,k) is the sum of Golay-Rudin-Shapiro terms GRS(j) (A020985) for j in the range 0 <= j < 2^n and having binary weight wt(j) = A000120(j) = k.

Doche and Mendès France form polynomials P_n(y) = Sum_{j=0..2^n-1} GRS(j) * y^wt(j) and here row n is the coefficients of P_n starting from the constant term, so P_n(y) = Sum_{k=0..n} T(n,k)*y^k. They conjecture that the number of real roots of P_n is A285869(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kevin Ryde_, Aug 21 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015324 | OEIS-A347186

Ziggurat sequence (see Comments lines for definition).

It appears that the geometric algorithm described here is equivalent to the numerical algorithm conjectured in A280850 in the sense that both allow the value of the subparts of the SRS(n) to be computed.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 21 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015325 | OEIS-A347205

a(2n+1) = a(n) for n >= 0, a(2n) = a(n) + a(n - 2^A007814(n)) for n > 0 with a(0) = 1.

Conjectured formulas: (Start) Conjecture: a(n) = T(n, wt(n)+1), a(2n) = Sum_{k=1..wt(n)+1} T(n, k) where T(2n+1, k) = T(n, k) for 1 <= k <= wt(n)+1, T(2n+1, wt(n)+2) = T(n, wt(n)+1), T(2n, k) = Sum_{i=1..k} T(n, i) for 1 <= k <= wt(n)+1 with T(0, 1) = 1. - _Mikhail Kurkov_, Dec 13 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Aug 23 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015326 | OEIS-A347225

Lesser of twin primes (A001359) being both half-period primes (A097443).

The twin primes conjecture being true should imply that these two sequences are infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lamine Ngom_, Aug 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015327 | OEIS-A347242

Numbers k such that when iterating the map x -> A000593(x), at some point before 1 is reached (after starting from x=k), a term is encountered whose largest prime factor is at least as large as A006530(k).

Conjecture: There are no primes in this sequence. Checked up to the 2^20-th prime, 16290047.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 28 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015328 | OEIS-A347255

Number of positive integers with n digits that are the product of two integers ending with 4.

Conjecture: lim_{n->oo} a(n)/a(n-1) = 10. The conjecture is true since the sequence A347253 has positive asymptotic density (see Corollary 1 in Spezia link). - _Stefano Spezia_, Mar 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Aug 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015329 | OEIS-A347259

a(n) is the least prime that can be written in exactly n ways as s+2*p where s is a semiprime and p is a prime, or 0 if no such prime exists.

Conjecture: First zeros of the sequence are a(41), a(666), a(1277), a(2701), ... - _Giorgos Kalogeropoulos_, Sep 23 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Aug 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015330 | OEIS-A347261

Numbers k for which sigma(k)/k = 32/11.

Conjecture: a(n) = 33*A341623(n) for n >= 1. Motivation: If no term of A341623 is divisible by 11 (which appears to be the case), then sigma(33*A341623(n))/(33*A341623(n)) = sigma(11)*sigma(3*A341623(n))/(33*A341623(n)) = 12*(8*A341623(n))/(33*A341623(n)) = 32/11. Does this sequence, though, contain any additional terms that are not generated by A341623?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Timothy L. Tiffin_, Aug 24 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015331 | OEIS-A347272

Main diagonal of the square array A347270.

Conjectures from _Alois P. Heinz_, Aug 31 2021: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 25 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015332 | OEIS-A347285

Irregular triangle T(n,k) starting with n followed by e_k = floor(log_p_k(p_(k-1)^e_(k-1))) such that e_k > 0.

Conjecture: let S be the sum of the absolute values of the first differences of terms in row n. For all n > 0, n - S = 1. - _Michael De Vlieger_, Aug 27 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Aug 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015333 | OEIS-A347298

Numbers that contain an even digit d immediately followed by a digit <= d.

Conjecture: This is precisely the list of numbers missing from A342042. The conjecture is correct, see A342042 for details. - _Sebastian Karlsson_, Oct 02 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015334 | OEIS-A347301

Let S be a set of n distinct integers in the range -n-3 to n+3, and consider the sums s+t of pairs of distinct elements of S; a(n) is the maximum number of such sums that are powers of 2, over all choices for S.

Conjecture: Construction (C1), with proper values of M, will compute the majority of maximal values for a(n). The exceptions, like 27, 28 above, seem to cluster near (7/8)*2^k, with a run from n = 54 to n = 59, and another from n = 111 to n = 124 (comparing values from (C1) to those provided by T. Scheuerle in A352178).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 28 2021, based on information supplied by _Rob Pratt_ and _Stan Wagon_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015335 | OEIS-A347306

When k appears in A347113, or -1 if k never appears.

It is conjectured that every positive number appears in A347113. - _N. J. A. Sloane_, Nov 08 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 01 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015336 | OEIS-A347313

Index of prime(n) in A347113, or -1 if that prime never appears.

Conjecture: every prime appears in A347113 (every number, in fact).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015337 | OEIS-A347344

Positive integers k such that k with the first (most significant) digit repeated is prime.

Conjecture: the sequence contains infinitely many twin prime pairs (k, k+2) such that (D(k), D(k+2)) is a twin prime pair, where D(x) = x with the most significant digit repeated. The first such k is 659: both (659, 661) and (6659, 6661) are twin prime pairs. All these k begin with either 3, 6, or 9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Aug 27 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015338 | OEIS-A347391

Numbers k such that sigma(k) is either their sibling in Doudna tree (A005940), or one of the sibling's descendants.

Conjecture: sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 30 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015339 | OEIS-A347409

Longest run of halving steps in the trajectory from n to 1 in the Collatz map (or 3x+1 problem), or -1 if no such trajectory exists.

Conjecture: every integer >= 4 appears in the sequence infinitely many times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Aug 30 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015340 | OEIS-A347469

For irrational r > 1, let B(r) denote the Beatty sequence for r. Let (s(n)) be the increasing sequence of numbers in both B(sqrt(2)) and B(sqrt(3)). Then this sequence gives the numbers k such that s(k) = floor(k*sqrt(6)).

It is conjectured that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 31 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015341 | OEIS-A347562

Number of ways to write n as 16^w + x^2 + (y^2 + 23*z^4)/324, where w,x,y,z are nonnegative integers.

Conjecture: a(n) > 0 for all n > 0. See also A347827 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015342 | OEIS-A347563

Binomial complement triangle, T(n,k) = LCM(1,...,n)/binomial(n,k) for 0 <= k <= n, a(0) = T(0,0) = 0, read by rows.

Conjecture: For all n, T(n,0) mod A213999(n-1,n-1) = 0, and T(n,k+1) mod A213999(n,k) = 0 for 0 <= k <= n-1 (computed and verified for rows = 0..2000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary Waters_, Sep 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015343 | OEIS-A347599

Irregular table read by rows, T(n, k) is the rank of the k-th Genocchi permutation of {1,...,n}, permutations sorted in lexicographical order. If no Genocchi permutation of {1,...,n} exists, then T(n, 1) = 0 by convention.

Let P(n) denote the number of Genocchi permutations of order n. _Zhi-Wei Sun_ conjectured, using permanents, that P(n - 1) = G(n), where G(n) are the Genocchi numbers A036968. From the well-known relation between Genocchi and Bernoulli numbers this implies, assuming the conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Sep 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015344 | OEIS-A347600

Irregular table read by rows, T(n, k) is the rank of the k-th Seidel permutation of {1,...,n}, permutations sorted in lexicographical order.

Let P(n) denote the number of Seidel permutations of order n. We conjecture that P(n) = A005439(n). This conjecture was inspired by the conjecture of _Zhi-Wei Sun_ in A036968. The name 'Seidel permutations' follows a comment of _Don Knuth_: "The earliest known reference for these numbers (A005439) is Seidel ...."

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Sep 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015345 | OEIS-A347603

Numbers k such that tau(k) = 2*tau(k-1) and sigma(k) = sigma(k-1), where tau(k) and sigma(k) are respectively the number and sum functions of the divisors of k.

Conjecture: the asymptotic density of terms is equal to 0 and this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Claude H. R. Dequatre_, Sep 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015346 | OEIS-A347637

Table read by ascending antidiagonals. T(n, k) is the minimum number of pebbles such that any assignment of those pebbles on a complete graph with n vertices is a next-player winning game in the two-player impartial (k+1, k) pebbling game. T(n, k) for n >= 5 and k >= 1.

It is conjectured that T(4, k) for odd k>=3 is infinite, so we start with n = 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kayla Barker_, _Mia DeStefano_, _Eugene Fiorini_, _Michael Gohn_, _Joe Miller_, _Jacob Roeder_, _Wing Hong Tony Wong_, Sep 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015347 | OEIS-A347668

Indices of records in A347409.

Conjecture 1: A347409(a(n)) is even for n >= 11. Conjecture 2: all even numbers > 2 appear as A347409(a(n)) for some n. - _Chai Wah Wu_, Sep 29 2021 If conjectures 1 and 2 are true, then A347409(a(n)) = 2n - 6 for n >= 11, and hence a(n) <= (4^(n-3)-1)/3 for n >= 11 since A347409((4^(n-3)-1)/3) = 2n - 6. - _Charles R Greathouse IV_, Oct 25 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Sep 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015348 | OEIS-A347684

Array read by antidiagonals: T(n,k) (n>=1, k>=1) = f(n,k), where f(x,y) = x*red_inv(x,y) + y*red_inv(y,x) if gcd(x,y)=1, or 0 if gcd(x,y)>1, and red_inv is defined in the comments.

Conjecture: The entries of this array can be expressed in terms of A347683 and A347687 as follows. Write k = j*n+i, where j >= 0 and 1 <= i <= n. Then T(n,k) = A347683(n,i) + 2*n*j*A347687(n,i). For example, suppose n=3, k=11, so k = 3*3+2, with j=3, i=2. Then A347683(3,2) + 2*3*3*A347687(3,2) = 5 + 2*3*3*1 = 23, which is indeed T(3,11).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 18 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015349 | OEIS-A347687

Triangle read by rows: T(n,k) (1<=k<=n) = (r(n+k)-r(k))/(2*n), where {r(i), i>=1} is the n-th row of array A347684.

Conjecture: Row n is not just the rescaled differences between the first two blocks of n terms of row n of A347684, but is in fact the rescaled differences between any two successive blocks of n terms that start at positions == 1 (mod n). See also the comments in A347684.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 19 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015350 | OEIS-A347699

Triangle read by rows: For n >= 1, 0 <= k <= n-1, T(n,k) = 0 if k=0, otherwise the number of inequivalent k X (n-k) 0,1 matrices having at least one 1 in each column.

It appears that this is essentially the same as the triangle in A055080, except that the 1's at the ends of the rows in A055080 have been moved to the start of the rows. (This is at present only a conjecture.) Even if the conjecture turns out to be true, this triangle deserves its own entry because the arrays look so different. The conjecture would imply the row sums are given by A048194.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015351 | OEIS-A347719

a(n) is the smallest prime(n)-rough number k that has more divisors than k-1 and more divisors than k+1.

Conjecture: no term is the product of more than 4 prime factors, counted with multiplicity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Sep 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015352 | OEIS-A347748

Number of positive integers with n digits that are equal both to the product of two integers ending with 4 and to that of two integers ending with 6.

Conjecture: lim_{n->oo} a(n)/a(n-1) = 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Sep 12 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015353 | OEIS-A347749

Number of positive integers with n digits and final digit 6 that are equal to the product of two integers ending with the same digit.

Conjecture: lim_{n->oo} a(n)/a(n-1) = 10. The conjecture is true since the sequence A347747 has positive asymptotic density (see Corollary 1 in Spezia link). - _Stefano Spezia_, Mar 17 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Sep 12 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015354 | OEIS-A347753

Number of polyhedra formed when a row of n adjacent cubes are internally cut by all the planes defined by any three of their vertices.

Conjectured formula for the number of internal cutting planes for n adjacent cubes is A007588(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 12 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015355 | OEIS-A347770

Conjectured list of numbers which are perfect, amicable, or sociable.

Conjectured list of numbers which are perfect, amicable, or sociable. Conjecture: There are no aliquot cycles containing even numbers and odd numbers simultaneously, i.e., every aliquot cycle either has only even numbers or has only odd numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, Sep 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015356 | OEIS-A347793

Intersection of Beatty sequences for 2^(1/3) and 2^(2/3).

Let d(n) = a(n) - 2n. Conjecture: (d(n)) is unbounded below and above, and d(n) = 0 for infinitely many n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 01 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015357 | OEIS-A347819

Minimal elements for the base-10 representations of the primes greater than 10.

b=7: {14, 16, 23, 25, 32, 41, 43, 52, 56, 61, 65, 113, 115, 131, 133, 155, 212, 221, 304, 313, 335, 344, 346, 364, 445, 515, 533, 535, 544, 551, 553, 1022, 1051, 1112, 1202, 1211, 1222, 2111, 3031, 3055, 3334, 3503, 3505, 3545, 4504, 4555, 5011, 5455, 5545, 5554, 6034, 6634, 11111, 11201, 30011, 30101, 31001, 31111, 33001, 33311, 35555, 40054, 100121, 150001, 300053, 351101, 531101, 1100021, 33333301, 5100000001, 33333333333333331} (conjectured, not proven)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Chen_, Sep 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015358 | OEIS-A347824

Number of ways to write n as x^4 + y^4 + (z^2 + 23*w^2)/16, where x,y,z,w are nonnegative integers with x <= y.

Conjecture: a(n) > 0 for all n = 0,1,2,.... For m = 32, 48, we also conjecture that every n = 0,1,2,... can be written as x^4 + y^4 + (z^2 + 23*w^2)/m, where x,y,z,w are nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015359 | OEIS-A347827

Number of ways to write n as w^4 + x^4 + (y^2 + 23*z^2)/16, where w is zero or a power of two (including 2^0 = 1), and x,y,z are nonnegative integers.

23-Conjecture: a(n) > 0 for all n = 0,1,2,.... This is stronger than the conjecture in A347824, and it has been verified for n up to 3*10^6. See also a similar conjecture in A347562.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015360 | OEIS-A347840

A surjective map of the positive numbers congruent to 5 modulo 8 (A004770) to the positive numbers congruent to 1, 3, or 7 modulo 8 (A047529).

This map f is of interest in the context of the Collatz 3*n+1 conjecture. The (modified) rooted tree with only odd labeled nodes has for each row k of the array A(k, m) (A347834) the same precursor (or (modified) Collatz map given in A075677(n+1), for 2*n+1). Therefore, all nodes with labels b(n) == 5 (mod 8) can be represented by a(n). This leads to a further restricted Collatz tree with only node labels congruent to {1, 3, 7} (mod 8) (A047529).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 30 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015361 | OEIS-A347854

a(n) = (6*n)!/((3*n)!*(2*n)!) * (n/2)!/(3*n/2)!.

Here we are essentially considering the sequence (u(n/2))n>=0. The sequence is conjectured to be integral. O.g.f.: A(x) = hypergeom([1/12, 5/12, 7/12, 11/12], [1/3, 1/2, 2/3], 27648*x^2) + 40*x*hypergeom([11/12, 13/12, 7/12, 17/12], [3/2, 5/6, 7/6], 27648*x^2) is conjectured to be algebraic over Q(x). Conjectural: a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015362 | OEIS-A347855

a(n) = (4*n)!/((2*n)!*(n)!) * (n/3)!/(4*n/3)!.

Fractional factorials are defined using the Gamma function; for example, (n/3)! := Gamma(1 + n/3). The sequence defined by u(n) = (12*n)!*n! / ((6*n)!*(4*n)!*(3*n)!) is one of the 52 sporadic integral factorial ratio sequences of height 1 found by V. I. Vasyunin (see Bober, Table 2, Entry 1). See A295431. Here we are essentially considering the sequence (u(n/3))n>=0. The sequence is conjectured to be integral. O.g.f.: A(x) = hypergeom([11/12, 7/12, 5/12, 1/12], [2/3, 1/2, 1/3], 27648*x^3) + 9*x*hypergeom([11/12, 5/4, 5/12, 3/4], [5/6, 4/3, 2/3], 27648*x^3) + 189*x^2*hypergeom([19/12, 13/12, 5/4, 3/4], [7/6, 5/3, 4/3], 27648*x^3) is conjectured to be algebraic over Q(x). Conjectural congruences: a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 17 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015363 | OEIS-A347856

a(n) = (6*n)!/((4*n)!*n!) * (n/2)!/(3*n/2)!.

The sequence defined by u(n) = (12*n)!/((8*n)!*(2*n)!) * n!/(3*n)! is one of the 52 sporadic integral factorial ratio sequences of height 1 found by V. I. Vasyunin (see Bober,

Table 2, Entry 3). See A295433. Here we are essentially considering the sequence (u(n/2))n>=0. The sequence is conjectured to be integral. O.g.f. A(x) = hypergeom([1/12, 1/6, 5/12, 7/12, 5/6, 11/12], [1/8, 3/8, 1/2, 5/8, 7/8], (19683/4)*x^2) + 20*x*hypergeom([17/12, 13/12, 11/12, 7/12, 4/3, 2/3], [11/8, 9/8, 7/8, 5/8, 3/2], (19683/4)*x^2) is conjectured to be algebraic over Q(x). Conjectural congruences: a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 17 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015364 | OEIS-A347857

a(n) = (6*n)!/((3*n)!*(2*n)!) * (3*n/2)!/(5*n/2)!.

Fractional factorials are defined using the Gamma function; for example, (3*n/2)! := Gamma(1 + 3*n/2). The sequence defined by u(n) = (12*n)!*(3*n)!/((6*n)!*(4*n)!*(5*n)!) is one of the 52 sporadic integral factorial ratio sequences of height 1 found by V. I. Vasyunin (see Bober, Table 2, Entry 5). See A295435. Here we are essentially considering the sequence (u(n/2))n>=0. The sequence is conjectured to be integral. O.g.f. A(x) = hypergeom([1/12, 1/3, 5/12, 7/12, 2/3, 11/12], [1/5, 2/5, 1/2, 3/5, 4/5], (2^10*3^9/5^5)*x^2) + 24*x*hypergeom([7/12, 5/6, 11/12, 13/12, 7/6, 17/12], [7/10, 9/10, 11/10, 13/10, 3/2], (2^10*3^9/5^5)*x^2) is conjectured to be algebraic over Q(x). Conjectural congruences: a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 18 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015365 | OEIS-A347858

a(n) = (9*n)!/((3*n)!*(2*n)!) * (n/2)!/(9*n/2)!.

The sequence defined by u(n) = (18*n)!*(n)!/((4*n)!*(6*n)!*(9*n)!) is one of the 52 sporadic integral factorial ratio sequences of height 1 found by V. I. Vasyunin (see Bober, Table 2, Entry 7). See A295437. Here we are essentially considering the sequence (u(n/2))n>=0. The sequence is conjectured to be integral. O.g.f.: A(x) = hypergeom([1/18, 5/18, 7/18, 11/18, 13/18, 17/18], [1/4, 1/3, 1/2, 2/3, 3/4], (2^4)*(3^12)*x^2) + 512*x*hypergeom([5/9, 7/9, 8/9, 10/9, 11/9, 13/9], [3/4, 5/6, 7/6, 5/4, 3/2], (2^4)*(3^12)*x^2) is conjectured to be algebraic over Q(x). Conjectural congruences: a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for prime p >= 5 and positive integers n and k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 19 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015366 | OEIS-A347865

Number of ways to write n as w^2 + 2*x^2 + y^4 + 3*z^4, where w,x,y,z are nonnegative integers.

Conjecture 1: a(n) > 0 except for n = 744. Conjecture 2: For any positive odd integer a, all sufficiently large integers can be written as a*w^4 + 2*x^4 + (2*y)^2 + z^2 with w,x,y,z integers. If M(a) denotes the largest integer not of the form a*w^4 + 2*x^4 + (2*y)^2 + z^2 (with w,x,y,z integers), then M(1) = 255, M(3) = 303, M(5) = 497, M(7) = 3182, M(9) = 4748, M(11) = 5662, M(13) = 5982, M(15) = 10526, M(17) = 4028 and M(19) = 11934. Conjecture 3: Let E(a,b,c) be the set of nonnegative integers not of the form w^2 + a*x^2 + b*y^4 + c*z^4 with w,x,y,z integers. Then E(1,2,4) = {135, 190, 510}, E(1,2,5) = {35, 254, 334}, E(2,1,4) = {190, 270, 590} and E(2,3,7) = {94, 490, 983} and E(3,1,2) = {56, 168, 378}. See also A346643 and A350857 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015367 | OEIS-A347878

Numbers k for which A003415(sigma(k)) is even.

Conjecture: A342923 is a subsequence of this sequence. See comments in A347870.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 19 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015368 | OEIS-A347941

For sets of n random points in the real plane, a(n) is an upper bound for the minimal number of nearest neighbors.

We conjecture that a(n) is optimal. This claim is true if the following assumptions hold:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Manfred Boergens_, Sep 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015369 | OEIS-A348032

Multiply-perfect numbers k that do not have a divisor d such that sigma(d)*d = k.

Question: Is A323653 a subsequence of this sequence? See also conjecture in A348033.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015370 | OEIS-A348055

Number of positive integers with n digits that are the product of two integers ending with 7.

Conjecture: lim_{n->oo} a(n)/a(n-1) = 10. The conjecture is true since the sequence A348054 has positive asymptotic density (see Corollary 1 in Spezia link). - _Stefano Spezia_, Mar 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Sep 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015371 | OEIS-A348058

a(n) = Min {k > n : A235137(k) == n (mod k)}, or -1 if no such minimum exists.

Conjecture: For all n, a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Sep 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015372 | OEIS-A348120

Starts of runs of 3 consecutive numbers whose powerful part is larger than their powerfree part (A328014).

It is conjectured that there are no runs of 3 consecutive numbers that are powerful (A001694), but if they do exist, their starts are contained in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Oct 01 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015373 | OEIS-A348126

a(n) is the least k for which x(k) > Sum_{j=1..k} 1/j, where x(k) is the Newton iterate (starting from x(1)=1) for exp(-P(n,x)) and P(n,x) is the n-th Maclaurin polynomial for exp(x).

Conjecture: a(n) is finite for every positive integer n, and a(n) approaches infinity with n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jennifer Sirkin_, Oct 01 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015374 | OEIS-A348168

Segment the list of prime numbers into sublists L_1, L_2, ... with L_1 = {2} and L_n = {p_1, p_2, ..., p_a(n)}, where a(n) is the largest m such that for 0 < i < m, p_1 - prevprime(p_1) > p_2 - p_1 >= p_{i+1} - p_i.

Conjecture 1: lim_{n->oo} N_i/n = k_i, where N_i is the number of the first n sublists consisting of i primes and k_i is a constant, with k_2 > k_1 > k_3 > k_4 > ... . Conjecture 2: lim_{n->oo} (Sum_{i=1..n} a(i))/n = Sum_{i=1..oo} i*k_i = e, meaning that, as n tends to infinity, the average length of sublists approaches 2.71828... (see the partial average - n plot in the links). The distribution of sublists with 1, 2, 3, 4 and 5 primes and the number of primes in the first n sublists are given in the table below. k_i's as defined in Conjecture 1 are: k1 = 0.281, k2 = 0.431, k3 = 0.127, k4 = 0.058, and k5 = 0.031, approximately. Sublists with length <= 5 account for about 93% of the terms and 70% of the primes, as n approaches infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Oct 03 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015375 | OEIS-A348176

Numbers m such that the number of symmetrically distributed consecutive primes centered at m sets a new record.

Conjecture: This sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Oct 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015376 | OEIS-A348197

Composition of the g.f. of A086246 with itself.

Conjecture: a(n) > A068875(n-1) for n > 7 (equivalently, a(n) > 2*A000108(n-1) for n > 7).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Burstein_, Oct 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015377 | OEIS-A348263

Orders of Parker fields.

It is conjectured that these are the only such fields. It appears that there is a mistake in the paragraph after Conjecture 7.2 of the Cain article. It claims that there are only 17 finite Parker fields, although Lemma 5.2 clearly shows that all fields of order 2^k are Parker. I think the corrected conjecture should state that there are only 16 finite Parker fields of odd order. - _Yevhenii Diomidov_, Jan 19 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Oct 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015378 | OEIS-A348264

a(n) is the number of iterations that n requires to reach a fixed point under the map x -> A348173(x).

Conjecture: essentially partial sums of A219977 (verified for n <= 5000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Oct 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015379 | OEIS-A348295

a(n) = Sum_{k=1..n} (-1)^(floor(k*(sqrt(2)-1))).

Conjecture: (1) Sequence is unbounded from above. Moreover, it seems that the earliest occurrence of m is A000129(m) for even m and A001333(m) for odd m (this has been confirmed for m <= 32 by _Chai Wah Wu_, Oct 21 2021). See A084068 for the conjectured indices of records. Adamczewski includes a proof abstract regarding Conjecture 1 in linked article. - _Michael De Vlieger_, Aug 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Oct 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015380 | OEIS-A348299

Indices of 0 in A348295: numbers m such that Sum_{k=1..m} (-1)^(floor(k*(sqrt(2)-1))) = Sum_{k=1..m} (-1)^A097508(k) = 0.

Conjecture: sequence is infinite. Conjecture: all terms are divisible by 4. There are no counterexamples <= 10^9. Conjecture: for n >= 2, a(2n-1) = ceiling(a(n) * (3+2*sqrt(2))), a(2n) = a(2n-1) + 4. This is correct for the first 2048 terms (all terms <= 10^9). Conjectured explicit formula: if the binary expansion of n-1 is n-1 = 2^(e_0) + 2^(e_1) + ... + 2^(e_k), then a(n) = 4*(A001109(1+(e_0)) + A001109(1+(e_1)) + ... + A001109(1+(e_k))). For example, since 28-1 = 27 = 2^0 + 2^1 + 2^3 + 2^4, a(28) = 4*(A001109(1) + A001109(2) + A001109(4) + A001109(5)) = 5600.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Oct 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015381 | OEIS-A348319

Perfect powers m^k, k >= 2 that are palindromes while m is not a palindrome.

G. J. Simmons conjectured there are no palindromes of form n^k for k >= 5 (and n > 1) (see Simmons link p. 98), according to this conjecture, we have 2 <= k <= 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Oct 12 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015382 | OEIS-A348320

Perfect powers m^k, k >= 2 of palindromes m when m^k is not a palindrome.

G. J. Simmons conjectured that there are no palindromes of form n^k for k >= 5 (and n > 1) (see Simmons link p. 98); according to this conjecture, every palindrome^k, k >= 5 is a term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Oct 12 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015383 | OEIS-A348339

a(n) is the number of distinct numbers of steps required for the last n digits of integers to repeat themselves by iterating the map m -> m^2.

Conjecture: For n >= 5, a(n) = a(n-1) + 2*n - 3 - ceiling(log_5 ((n-1)/16)), or a(n) = (n-1)^2 + 3 - Sum_{5..n} ceiling(log_5 ((i-1)/16)). The largest number of steps required is 4*5^(n-2) + (n-2) for n >= 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Oct 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015384 | OEIS-A348358

Primes which are not the concatenation of smaller primes (in base 10 and allowing leading 0's).

Conjecture. If p > 7 is a prime, then there exists a base b such that p in base b is the concatenation of smaller primes in base b.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. Farrokhi D. G._, Oct 14 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015385 | OEIS-A348410

Number of nonnegative integer solutions to n = Sum_{i=1..n} (a_i + b_i), with b_i even.

Conjecture: D-finite with recurrence +7168*n*(2*n-1)*(n-1)*a(n) -64*(n-1)*(1759*n^2-5294*n+5112)*a(n-1) +12*(7561*n^3-75690*n^2+165271*n-101070)*a(n-2) +5*(110593*n^3-743946*n^2+1659971*n-1232778)*a(n-3) +2680*(4*n-15)*(2*n-7)*(4*n-13)*a(n-4)=0. - _R. J. Mathar_, Oct 19 2021 Conjecture: the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) hold for all primes p >= 5 and positive integers n and k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _César Eliud Lozada_, Oct 17 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015386 | OEIS-A348424

a(n) is the difference between the index of the prime p = A035345(n)-A002110(n) and n, or 0 if the difference is composite.

Conjecture 1: The mechanism always returns a prime in a finite number of iterations (term always exists). So the sequence would be infinite. Conjecture 2: Terms will always be less than the associated offset. (So for n=33, the base primorial is 72047817630210000485677936198920432067383702541010310, with largest pf=137. The entry of 84 is less than 137.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bill McEachen_, Nov 04 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015387 | OEIS-A348429

Perfect powers m^k, m >= 1, k >= 2 such that m and m^k both are palindromes.

For k >= 5, G. J. Simmons conjectured there are no palindromes of the form m^k for k >= 5 and m > 1 (see Simmons link p. 98); according to this conjecture, all the terms are of the form (palindrome)^k, with 2 <= k <= 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Oct 18 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015388 | OEIS-A348474

Number of compositions of n into exactly 2n nonnegative parts such that each positive i-th part is odd if i is odd.

Conjecture: a(n) = [x^n] ((1 + x - x^2)/((1 + x)*(1 - x)^2))^n. The above conjectures of Bala are proved in Raisa (2026). - _Rubaiyat Zaman Raisa_, Aug 02 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Oct 19 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015389 | OEIS-A348482

Triangle read by rows: T(n,k) = (Sum_{i=k..n} i!)/(k!) for 0 <= k <= n.

T(n,1) = T(n,0)-1 = Sum_{k=0..n-1} T(n,k)/(k+2) for n > 0 (conjectured). (c) p(n,x) = (x+1) * p(n-1,x) + 1 + Sum_{i=1..n-1} (d/dx)^i p(n-1,x) for n > 0 (conjectured). Alternating row sums p(n,-1) equal A173184(n) for n >= 0 (conjectured). The three conjectures stated above are true. See links. - _Sela Fried_, Jul 11 2024.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Oct 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015390 | OEIS-A348484

Maximum number of squares on an n X n chessboard such that no two are two steps apart horizontally or vertically.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yang Hong_, Oct 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015391 | OEIS-A348506

Numbers k such that sigma(k) is a multiple of usigma(k), where sigma is the sum of divisors function, A000203, and usigma is the unitary sigma, A034448.

Conjectured to be the union of A005117 and A063880.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015392 | OEIS-A348511

Numbers k for which 2k+1 can be obtained with successive prime shifts towards larger primes (by iterating A003961, starting from k).

Conjecture: Apart from 4 all other terms in this sequence are squarefree. No counterexamples <= 2^22.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 30 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015393 | OEIS-A348514

Numbers k for which A003961(k) = 2k+1, where A003961 shifts the prime factorization of n one step towards larger primes.

It seems that after 4, all other terms are squarefree. See conjecture in A348511.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015394 | OEIS-A348546

Number of positive integers with n digits that are equal both to the product of two integers ending with 3 and to that of two integers ending with 7.

Conjecture: lim_{n->infinity} a(n)/a(n-1) = 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Oct 22 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015395 | OEIS-A348547

Number of positive integers with n digits and final digit 9 that are equal to the product of two integers ending with the same digit.

Conjecture: lim_{n->oo} a(n)/a(n-1) = 10. The conjecture is true since the sequence A348545 has positive asymptotic density (see Corollary 1 in Spezia link). - _Stefano Spezia_, Mar 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Oct 22 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015396 | OEIS-A348549

Number of positive integers with n digits that are the product of two integers ending with 8.

Conjecture: lim_{n->oo} a(n)/a(n-1) = 10. The conjecture is true since the sequence A348548 has positive asymptotic density (see Corollary 1 in Spezia link). - _Stefano Spezia_, Mar 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Oct 22 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015397 | OEIS-A348598

Least prime p of the form k^2+1 such that p == A002496(n) (mod A002496(n+1)) with p>A002496(n), or 0 if no such p exists.

Conjecture: Consider the smallest prime p of the form k^2+1 such that p is congruent to A002496(n) modulo q, q prime of the form m^2+1 > A002496(n). Then q = A002496(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Mar 20 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015398 | OEIS-A348618

a(n) = (1+(-1)^n)/2*4^n*(C((3*n)/2-1,n))+(1-(-1)^n)/2*((C((3*n-1)/2,n))*(C(3*n-1,(3*n-1)/2)))/(C(n-1,(n-1)/2)).

Conjecture: D-finite with recurrence n*(n-1)*a(n) -12*(3*n-2)*(3*n-4)*a(n-2)=0. - _R. J. Mathar_, Mar 06 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Oct 25 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015399 | OEIS-A348667

a(n) is the index of the first 0 term in the following sequence: X(1)=1, X(i+1) = (X(i)+i) mod (n+i). a(n)=-1 if there are no zeros in the X sequence.

Conjecture: a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Oct 28 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015400 | OEIS-A348713

Numbers whose divisors can be partitioned into two disjoint sets with equal arithmetic mean.

Conjecture: If a(n) and k are coprime, then a(n)*k is a term. No counterexamples found in the b-file of 872 terms. - _Ivan N. Ianakiev_, Jul 23 2026 This conjecture is true; see a proof in the Link section.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Oct 31 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015401 | OEIS-A348840

Triangle T(n,h) read by rows: The number of Motzkin Paths of n>=2 steps that start with an Up step and touch the horizontal axis h>=1 times afterwards.

Conjecture: T(n,n-2) = n-2. Conjecture: T(n,n-3) = A000124(n-3). Conjecture: T(n,n-4) = -11 + 19*n/3 - 3*n^2/2 + n^3/6. Conjecture: Riordan array (g(x)^2, x*g(x)), where g(x) = 1/(1 + x)*c(x/(1 + x)) and c(x) = (1 - sqrt(1 - 4*x))/(2*x) is the g.f. of the Catalan numbers A000108. - _Peter Bala_, Feb 04 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Nov 01 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015402 | OEIS-A348846

a(1) = 1. For n >=2 the number k in n-th position becomes a(n) only if all terms a(1)..a(n-1) have already been defined, and if the smallest number m, greater than k, not already defined and sharing greatest prime factor (gpf) p with k is reduced to m/p.

Let [p] = {m: m a fixed point with gpf = p}, then [2] = {2}, [3] = {3,9,18,27}, [5] = {5,15,25,40,50,90}, etc. Every odd multiple of odd prime p, up to and including p^2, is necessarily a fixed point. The number of terms in [p] is limited by reduction of q-smooth numbers (q>p) to those having gpf p. Conjecture: For odd prime p, [p] is a finite set with greatest term > p^2, and <= p^3. A variant based on least prime divisors is also possible.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Nov 07 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015403 | OEIS-A348889

Numbers m such that there exists at least one integer k < m where m^2 + 2 and k^2 + 2 have the same prime factors.

If we consider the pairs (m, k), k is not unique, in contrast to the same problem with m^2 + 1 (see A282092) where conjecturally k seems unique.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 28 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015404 | OEIS-A348890

Number of ways to write n as 5*w^4 + x^4 + y^2 + z^2, where w,x,y,z are nonnegative integers with y <= z.

Conjecture: a(n) = 0 only for n == 12 (mod 16).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 28 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015405 | OEIS-A348970

a(n) = A003959(n) - A129283(n), where A003959 is multiplicative with a(p^e) = (p+1)^e and A129283(n) is sum of n and its arithmetic derivative.

There are no negative terms. We prove this by induction over the prime factorization of n, showing that A348507(n) >= A003415(n) for all values of n >= 1. At n=1, both sequences have value 0, and at the primes both sequences obtain the value 1, so the base cases hold. We know that A348507(n)-(n/p) = (p+1)*A348507(n/p) for all prime factors p of n (see comment in A348507). With the arithmetic derivative we obtain respectively that A003415(n) = A003415(p*(n/p)) = A003415(p)*(n/p) + p*A003415(n/p) = (n/p) + p*A003415(n/p), for any prime factor p of n. Now A348507(p*(n/p)) >= A003415(p*(n/p)) iff A348507(p*(n/p)) - (n/p) >= A003415(p*(n/p)) - (n/p), that is, iff (p+1)*A348507(n/p) >= p*A003415(n/p), which indeed follows by the induction hypothesis, which assumes that A348507(x) >= A003415(x) for all proper divisors x of n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015406 | OEIS-A349044

Non-Brauer numbers.

These differences between l^*(n) and A003313(n) are highlighted by the following formulation: consider a machine which starts with a 1 in "cache" and can then at each step execute one of two operations: (1) Add any number that has ever been in cache to the current contents of cache, or (2) Restore any number that has previously been in cache to the cache, replacing its prior contents. Then n is in this sequence if and only if there is a shortest program that results in n in cache that includes a "Restore" step. Note further that if there is an entry in this sequence such that l^*(n) > A003313(n)+1, then all shortest programs producing n in cache would contain a "Restore" operation. The definition of A293771 is based on a similar machine with a separate "Store" operation that puts the cache value into "memory," and one could formulate an analogous conjecture here that the "Restore" operation is never necessary for a shortest program. The existence or not of an n in this sequence such that l^*(n) > A003313(n)+1 would settle this question and provide mild evidence one way or the other on the conjecture in A293771.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Glen Whitney_, Nov 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015407 | OEIS-A349093

a(n) is the smallest nonprime number m (m = A018252(t)) such that n divides the product P(t) of all nonprime numbers up to and including m (P(t) = A036691(t-1)).

Conjecture: the equation a(n) = a(n+1) has no solutions. This holds up to at least n = 10^7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lechoslaw Ratajczak_, Mar 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015408 | OEIS-A349106

Irregular triangle read by rows: T(n,k) is the number of permutations of {1,2,...,n} with cycle descent number equal to k.

T(n,k) = [z^k] Sum_{i=0..n} Stirling2(n,i)*(1 - z)^(n - i) Product_{j=0..i-1} (j*z + 1). - Conjectured by _Mikhail Kurkov_, Jun 13 2023; proved by _Max Alekseyev_, Mar 12 2025 (see MO link)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Dec 30 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015409 | OEIS-A349169

Numbers k such that k * gcd(sigma(k), A003961(k)) is equal to the odd part of {sigma(k) * gcd(k, A003961(k))}, where A003961 shifts the prime factorization one step towards larger primes, and sigma is the sum of divisors function.

Conjectures: Note: If any of the last four conjectures could be proved, it would refute the existence of odd perfect numbers at once. Note that it seems that gcd(sigma(k), A003961(k)) < k, for all k except these four: 1, 2, 20, 160.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 10 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015410 | OEIS-A349179

Numbers with a record number of nonempty subsets of divisors whose harmonic mean is an integer (A339665).

Conjecture: a(n) = A002182(n-2) for n >= 9. - _Chai Wah Wu_, Nov 11 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Nov 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015411 | OEIS-A349229

a(n) = Sum_{k=1..n} (-1)^A001222(k)*(-1)^A001222(k+1).

It is conjectured that a(n)=o(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Benoit Cloitre_, Nov 11 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015412 | OEIS-A349246

Number of ways to write n as w^8 + x^4 + 2*y^4 + 4*z^4 + t*(t+1), where w, x, y, z, and t are nonnegative integers.

Conjecture: a(n) > 0 for all n = 0,1,2,....

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015413 | OEIS-A349272

a(n) = Product_{k = 1..2*n+1} Fibonacci(2*k) / Sum_{k = 1..2*n+1} Fibonacci(2*k).

Let m be an even positive integer. We conjecture that the sequence defined by b_m(n) = Product_{k = 1..2*n+1} Fibonacci(m*k) / Sum_{k = 1..2*n+1} Fibonacci(m*k) is integral. The formula given below proves the conjecture in the present case m = 2. The cases m = 4 and m = 6 of the conjecture can be proved in a similar manner. More generally, if F(n,x) denotes the n-th Fibonacci polynomial we conjecture that, for each n, the rational function Product_{k = 1..2*n+1} F(m*k,x) / Sum_{k = 1..2*n+1} F(m*k,x) is an integral polynomial.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Nov 12 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015414 | OEIS-A349295

a(n) is the number of ordered 6-tuples (a_1,a_2,a_3,a_4,a_5,a_6) having all terms in {1,...,n} such that there exists a tetrahedron ABCD with those edge-lengths, taken in a particular order (see comments).

Conjecture: The ratio a(n)/n^6 decreases with n and tends to a limit which is 0.10292439+-0,00000024 (1.96 sigmas, 95% confidence level) evaluated for n=2^32 on 6.4*10^12 random 6-tuples.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Corbelli_, Nov 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015415 | OEIS-A349296

First differences of A349295.

Conjecture: for n tending to infinity the ratio a(n) / A097125(n) tends to 24 as the probability that all a_i's are different tends to 1 and there are 24 6-tuples corresponding to the same tetrahedron if all a_i's are different. For n=254 the ratio is 23.9936919.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giovanni Corbelli_, Nov 13 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015416 | OEIS-A349325

Number of times the Collatz plot started at n crosses the y = n line, or -1 if the number of crossings is infinite.

Conjecture: every positive integer appears in the sequence infinitely many times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Nov 15 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015417 | OEIS-A349425

n-th stable digit (in decimal system) of n^(n^(...^n)).

In 2011 Ripà conjectured that, for any n >= 1, if b >= n + 2, then the n rightmost digits of n^^b are stable. The above-mentioned paper, published in 2022, proved that this conjecture is true and also stated the stronger sufficient condition that the height of the hyperexponent is greater than or equal to tilde(v(a)) + 2, where tilde(v(a)) := v_5(a - 1) iff a == 1 (mod 5), v_5(a^2 + 1) iff a == {2, 3} (mod 5), v_5(a + 1) iff a == 4 (mod 5), v_2(a^2 - 1) - 1 iff a == 5 (mod 10), where v_2(x) = A007814(x) and v_5(x) = A112765(x) are the 2-adic and 5-adic valuations of x, respectively. - _Marco Ripà_, Jul 24 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Nov 17 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015418 | OEIS-A349459

Least positive integer m such that the n numbers k^2*(k^2-1) (k=1..n) are pairwise distinct modulo m.

Conjecture: For any integer n > 1, the term a(n) is the least prime p > 2*n with p dividing a^2 + b^2 - 1 for no 1 <= a < b <= n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 18 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015419 | OEIS-A349462

The a_i coefficients of the standard form decomposition of U(1,X), the Ulam sequence starting with 1, X in the ordered abelian group of linear integer polynomials in X, where the ordering is lexicographical.

It is conjectured that for all k and n > 3, U(1,n) intersect [1,c_k n + d_k + 1] = eval_n(U(1,X) intersect [1,c_k X + d_k + 1]). At the time of writing, this had been checked up to k = 217529.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arseniy (Senia) Sheydvasser_, Nov 19 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015420 | OEIS-A349485

Moran numbers whose arithmetic derivative is also a Moran number (A001101).

Conjecture: The sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius A. Burtea_, Nov 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015421 | OEIS-A349500

a(n) is the least number k such that A349474(k) = A349474(k+1) = n, or -1 if no such k exists.

The sequence begins at n = 2. a(1) != -1 if and only if two consecutive harmonic numbers exist. There are no odd harmonic numbers between 2 and 10^24 (Cohen and Sorli, 2010) and it is conjectured that they do not exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Nov 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015422 | OEIS-A349509

a(n) is the denominator of binomial(n^3 + 6*n^2 - 6*n + 2, n^3 - 1)/n^3.

Conjecture: 1 and 3 are the only terms that appear in this sequence. This conjecture is correct, see formula. - _Kevin Ryde_, Jul 01 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Nov 20 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015423 | OEIS-A349530

Least positive integer m such that the n numbers k*(k^4+1) (k=1..n) are pairwise distinct modulo m^2.

Conjecture 1: Suppose that 5^a < sqrt(n) <= 5^(a+1). Then a(n) = 3*5^a if sqrt(n) <= sqrt(3)*5^a, and a(n) = 5^(a+1) otherwise. Conjecture 2: Let f(n) be the least positive integer m such that the n numbers 18k*(k^4+1) (k=1..n) are pairwise distinct modulo m^2. Then f(n) is the least power of 5 not smaller than sqrt(n), except for 25 < n <= 45 (and in this case f(n) = 19). Conjecture 3: Let n be a positive integer not among 26, 27, 28, 626, 627, 628, 629, 630, and define D(n) as the least positive integer m such that the n numbers k*(k^4+1) (k=1..n) are pairwise distinct modulo m. If 5^a < n <= 3*5^a, then D(n) = 3*5^a. If 3*5^a < n <= 5^(a+1), then D(n) = 5^(a+1). We have verified the above conjectures for n up to 10^5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 21 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015424 | OEIS-A349536

Consider a circle on the Z X Z lattice with radius equal to the Pythagorean hypotenuse h(n) (A009003); a(n) = number of Pythagorean triples inside a Pi/4 sector of the circle.

Conjecture: the increment is a(n+1) - a(n) = 2^(m-1), where m is the sum of all powers of the Pythagorean primes (A002144) in the factorization of hypotenuse h(n+1) (see Eckert for PPT). However, starting from 58 the increment is 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Kritov_, Nov 21 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015425 | OEIS-A349537

Least positive integer m such that the n numbers 33*k^2*(k^3+1) (k = 1..n) are pairwise distinct modulo m.

Conjecture: a(n) is prime for each n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 21 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015426 | OEIS-A349576

Recurrence a(1) = 1, a(2) = 5; a(n) = (a(n-1) + a(n-2))/GCD(a(n-1),a(n-2)) + 1.

Conjecture: this sequence is not periodic. (Note that the analogous sequences with a(2) = 1, 2, 3, or 4 are periodic.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Dec 30 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015427 | OEIS-A349637

a(1) = 2 and a(n) is the smallest nonsquare positive integer not occurring earlier such that the intersection of the periodic parts of continued fractions for square roots of a(n) and a(n-1) is the empty set.

Conjecture: This is a permutation of the nonsquares (A000037).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giorgos Kalogeropoulos_, Nov 23 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015428 | OEIS-A349661

Number of ways to write n as x^4 + y^2 + (z^2 + 4^w)/2 with x,y,z,w nonnegative integers.

Conjecture: a(n) > 0 for all n > 0. As (x^2 + y^2)/2 = ((x+y)/2)^2 + ((x-y)/2)^2, the conjecture gives a new refinement of Lagrange's four-square theorem. See also A350012 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015429 | OEIS-A349753

Odd numbers k for which A003961(k)-2k divides A003961(k)-sigma(k), where A003961 shifts the prime factorization one step towards larger primes, and sigma is the sum of divisors function.

Conjecture (1): This is a subsequence of A319630, in other words, for all terms k, gcd(k, A003961(k)) = 1. Conjecture (2): Apart from 1, there are no common terms with A349169, which would imply that no odd perfect numbers exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 01 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015430 | OEIS-A349756

Numbers k such that the odd part of sigma(k) is equal to gcd(sigma(k), A003961(k)), where A003961 is fully multiplicative with a(p) = nextprime(p), and sigma is the sum of divisors function.

For any square s in this sequence, A349162(s) = 1, i.e. sigma(s) divides A003961(s), and also A286385(s). Question: Is 1 the only square in this sequence? (see the conjecture in A350072).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 03 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015431 | OEIS-A349778

Number of ways to write n as x^2 + y^k + 2*z^m, where x,y,z are nonnegative integers with x >= y, and k and m belong to the set {2,3}.

Conjecture 1: a(n) > 0 for all n >= 0, and a(n) = 1 only for n = 30, 120, 142. Conjecture 2: Let S = {x^k: k = 2,3 and x = 0,1,2,...}, and let a be 3 or 4 or 5. Then any nonnegative integer can be written as x + 2*y + a*z, where x,y,z are elements of the set S. Conjecture 3: Let T = {x^k: k = 2,3,4,... and x = 0,1,2,...}. If (b,c) is among the ordered pairs (1,2), (2,4), (2,5) and (3,2), then each n = 0,1,... can be written as x + b*y + c*z, where x and y are elements of T, and z is a square.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015432 | OEIS-A349787

Number of ways to write n as x^2 + y^k + 2^a + 2^b, where x,y,a,b are nonnegative integers with x >= y and a >= b, and k is either 2 or 3.

Conjecture 1: a(n) > 0 for all n >= 2. Below is our weaker version of Conjecture 1. Conjecture 2: Each n = 2,3,... can be written as a sum of two perfect powers (including 0 and 1) and two powers of 2 (including 2^0 = 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 30 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015433 | OEIS-A349791

a(n) is the median of the primes between n^2 and (n+1)^2.

Conjecture: lim_{n->oo} (a(n)/n - n) = 1 (see plot in Links). -_Ya-Ping Lu_, Sep 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015434 | OEIS-A349823

First differences of A230624.

This sequence could certainly be divided by 2, just as we divided A230624 itself by 2 to get A349821. But there is a reason for not dividing this by 2: it appears that, for any power of 2, from a certain point on, the sequence is divisible by that power of 2. At present this only a conjecture. But it may provide a clue to the structure of this sequence and therefore of A230624.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 31 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015435 | OEIS-A349824

a(0) = 0; for n >= 1, a(n) = (number of prime factors, counted with repetition) * (sum of prime factors, counted with repetition).

Conjecture (Mike Klein): Iterating n -> a(n) eventually leads to one of the fixed points {primes union 0, 27, 30} or the loop (28, 33).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015436 | OEIS-A349825

Number of steps when x -> A349824(x) is iterated starting at n needed to reach fixed point or 28, or -1 if trajectory increases for ever or ends in a nontrivial loop other than (28,33).

It is conjectured that every trajectory eventually reaches one of the fixed points {primes union 0, 27, 30} or the loop (28, 33).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015437 | OEIS-A349837

a(n) is the number of integers m > 1 such that sigma(m) = tau(m)^n, with a(1) = 1.

Conjecture: a(n) > 0 except for n = 76 and n = 81.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Dec 01 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015438 | OEIS-A349897

Start with b(0)=4. a(n) is the smallest integer such that b(n)=a(n)*b(n-1) has twice as many divisors as b(n-1) has.

Conjecture: a composite number is in this sequence if and only if it is the product of 2 consecutive terms of A050376.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Dec 04 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015439 | OEIS-A349931

Numbers that have an equal number of factorizations of even and odd length in both unordered and ordered manners.

Wilf's conjecture is equivalent to the statement that this sequence is the set difference of A319240 and A006881.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tian Vlasic_, Dec 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015440 | OEIS-A349937

Odd numbers k > 1 such that A309906(k-1) < A309906(k) > A309906(k+1).

Conjecturally, odd numbers k > 1 such that liminf_{n->oo} d(p(n)^(k-1)-1) < liminf_{n->oo} d(p(n)^k-1) > liminf_{n->oo} d(p(n)^(k+1)-1), where p(n) = prime(n), d = A000005.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Dec 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015441 | OEIS-A349938

Odd numbers k > 1 such that A309906(k-1) < A309906(k) > A309906(k+1) < A309906(k+2) > A309906(k+3).

Conjecturally, odd numbers k > 1 such that liminf_{n->oo} d(p(n)^(k-1)-1) < liminf_{n->oo} d(p(n)^k-1) > liminf_{n->oo} d(p(n)^(k+1)-1) < liminf_{n->oo} d(p(n)^(k+2)-1) > liminf_{n->oo} d(p(n)^(k+3)-1), where p(n) = prime(n), d = A000005.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Dec 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015442 | OEIS-A349941

Terms of A349937 that are not divisible by 3: numbers k > 1 not divisible by 2 or 3 such that A309906(k-1) < A309906(k) > A309906(k+1).

Conjecturally, numbers k > 1 not divisible by 2 or 3 such that liminf_{n->oo} d(p(n)^(k-1)-1) < liminf_{n->oo} d(p(n)^k-1) > liminf_{n->oo} d(p(n)^(k+1)-1), where p(n) = prime(n), d = A000005.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Dec 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015443 | OEIS-A349942

Number of ways to write n as a^4 + b^2 + (c^4 + d^2)/25 with a,b,c,d nonnegative integers.

Conjecture: a(n) > 0 for all n >= 0, and a(n) = 1 only for n = 0, 16^k*m (k = 0,1,2,... and m = 7, 8, 23, 44, 47). As m/n = (m*n^3)/n^4 for any nonnegative integers m and n > 0, the conjecture implies that each nonnegative rational number can be written as x^4 + 25*y^4 + z^2 + w^2 with x,y,z,w rational numbers. See also A349943 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015444 | OEIS-A349943

Number of ways to write n as a^4 + (b^4 + c^2 + d^2)/9, where a,b,c,d are nonnegative integers with c <= d.

Conjecture 1: a(n) > 0 for all n >= 0, and a(n) = 1 only for n = 0, 16^k*m (k = 0,1,2,... and m = 7, 8, 31, 39, 47, 79, 519). Conjecture 2: Each n = 0,1,2,... can be written as a^4 + (4*b^4 + c^2 + d^2)/81 with a,b,c,d nonnegative integers. We have verified Conjectures 1 and 2 for n <= 10^5. See also A349942 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 05 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015445 | OEIS-A349945

Number of ways to write n as a^4 + b^2 + (c^4 + d^2)/5 with a,b,c,d nonnegative integers.

Conjecture 1: a(n) > 0 for all n >= 0, and a(n) = 1 only for n = 0, 2^(4*k+3) (k = 0,1,2,...). Conjecture 2: Each n = 0,1,2,... can be written as a*x^4 + b*y^2 + (c*z^4 + w^2)/5 with x,y,z,w nonnegative integers, provided that (a,b,c) is among the four triples (1,2,4), (2,1,1), (6,1,1), (6,1,6). See also A349942 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015446 | OEIS-A349948

Tetrahedral-sided isosceles Heron triangle pairs.

Conjecture: for k a positive integer, the sequence {a(k^n): n >= 1} is a strong divisibility sequence; that is, for n, m >= 1, gcd(a(k^n), a(k^m)) = a(k^gcd(n,m)). - _Peter Bala_, Dec 03 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Randall L Rathbun_, Mar 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015447 | OEIS-A349956

Number of ways to write n as x^2 + 2*y^2 + (z^4 + 4*w^4)/5 with x,y,z,w nonnegative integers.

Conjecture: If m is 5 or 65 or 85, then each n = 0,1,2,... can be written as x^2 + 2*y^2 + (z^4 + 4*w^4)/m with x,y,z,w nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015448 | OEIS-A349957

Number of ways to write n as x^4 + y^2 + (z^2 + 11*16^w)/60, where x,y,z are nonnegative integers, and w is 0 or 1.

Conjecture 1: a(n) > 0 for all n > 0. Conjecture 2: (i) If a is 1 or 3, then each n = 0,1,2,... can be written as a*x^8 + y^2 + (7*z^4 + w^2)/64 with x,y,z,w nonnegative integers. Conjecture 3: If (a,b, m) is among the triples (1,7,8), (1,11,12), (2,7,8), (3,11,12), (5,7,8), then each n = 0,1,2,... can be written as a*x^4 + y^2 + (b*z^6 + w^2)/m with x,y,z,w nonnegative integers. Conjecture 4: (i) If F(x,y,z,w) is x^6 + y^2 + (5*z^4 + 3*w^2)/16 or 3x^6 + 2*y^2 + (11*z^4 + w^2)/60, then each n = 0,1,2,... can be written as F(x,y,z,w) with x,y,z,w nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015449 | OEIS-A349958

a(n) is the index of the first row in Pascal's triangle that contains a multiple of n.

If n is divisible by p^k where p is prime, then by Kummer's theorem a(n) >= p^k. In particular a(p^k) = p^k. Conjecture: a(n) = n if and only if n is a prime power (A246655). - _Robert Israel_, Aug 09 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nathan M Epstein_, Dec 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015450 | OEIS-A349960

Values taken by the function A067095 in the order of their appearance.

If we consider the sequence u(n) = a(n)/10^(k-1) where k = length(a(n)); we have u(n) is increasing with an upper bound 1.9; so, this sequence u(n) is convergent and, conjecture, this limit = 1.81757197277277957... found by _Giorgos Kalogeropoulos_; now, from this limit, it is possible to get the successive terms of this sequence here. a(n) = floor(k((n + 6)/2)*10^(n - 1 - ceiling(log_10(k((n + 6)/2))))) for k(n) = A019520(n)/A019519(n) and n >= 2 (conjectured). - _Giorgos Kalogeropoulos_, Dec 10 2021

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Dec 07 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015451 | OEIS-A349985

Primes of the form (product of 4 consecutive primes) + (sum of the same 4 consecutive primes).

It is conjectured that this sequence is infinite, and that similar lists of primes can be generated by using any even number of consecutive primes. Specifying 2 consecutive primes results in A096342. However, it should be noted that the percentage of resulting primes (as compared to all numbers generated in this manner) decreases as the number of consecutive primes to consider increases.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lyle Blosser_, Jan 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015452 | OEIS-A349992

Number of ways to write n as x^4 + y^2 + (z^2 + 2*4^w)/3, where x, y, z are nonnegative integers, and w is 0 or 1.

Conjecture 1: a(n) > 0 for all n > 0, and a(n) = 1 only for n = 1, 30, 64, 80, 302, 350, 472, 480, 847, 3497, 13582, 25630, 38064. Conjecture 2: If (a,b,c,m) is one of the ordered tuples (1,1,11,12), (1,1,11,60), (1,1,14,15), (1,1,23,24), (1,1,23,32), (1,1,23,48), (1,2,23,96), (2,1,11,60), (2,1,23,24), (2,1,23,48), (4,1,23,48), then each n = 1 2,3,... can be written as a*x^4 + b*y^2 + (z^2 + c*4^w)/m, where x,y,z are nonnegative integers, and w is 0 or 1. We have verified Conjecture 2 for n up to 2*10^5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

B08 A350000 - A401236 | 2,453 条

V3-015453 | OEIS-A350000

Table read by antidiagonals: T(n,k) (n >= 3, k >= 0) is the number of cells in a regular n-gon after k generations of mitosis.

Conjecture 1: For a fixed value of n, there are integers r and s, which are small compared to n, such that T(n,k) is a polynomial in k of degree r for all k >= s. Conjecture 2: Apart from the central n-gon when n is odd, any cell will eventually split into a mixture of triangles and pentagons. If we think of triangles and pentagons are harmless cells, and all other cells as dangerous, the conjecture states that (with the exception of the central odd cells), all cells eventually become harmless.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Dec 06 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015454 | OEIS-A350012

Number of ways to write n as 4*x^4 + y^2 + (z^2 + 4^w)/2 with x,y,z,w nonnegative integers.

Conjecture: a(n) > 0 for all n > 0. This is a new refinement of Lagrange's four-square theorem since (x^2 + y^2)/2 = ((x+y)/2)^2 + ((x-y)/2)^2. We have verified the conjecture for n up to 10^6. See also A349661 for a similar conjecture. We also have some other conjectures of such a type.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015455 | OEIS-A350021

Number of ways to write n as w^4 + x^2 + y^2 + z^2 with x - y a power of two (including 2^0 = 1).

Conjecture: a(n) > 0 for all n > 0. If x - y = 2^k, then x^2 + y^2 = ((x+y)^2 + (2^k)^2)/2 and x + y >= 2^k. So the above conjecture implies the conjecture in A349661.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 08 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015456 | OEIS-A350034

a(n) = n/gcd(n,6) if gcd(n,6)>1, 5n + 1 otherwise.

The 5x+1 problem is as follows: start with any number n. If gcd(n,6)>1, divide it by gcd(n,6), otherwise multiply it by 5 and add 1. Do we always reach 1? This is an unsolved problem. It is conjectured that the answer is yes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Dec 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015457 | OEIS-A350036

Numbers k such that (81^k + 1)/82 is prime.

These are the Repunits in base -81. Since 81=3^4, factors will be of the form p=8nk+1. (Negative) bases that are powers of small numbers appear to have a higher frequency of primes than Repunits in other bases. The best linear fit for this base is currently 0.29918 which is much lower (better) than the conjectured 0.56145948 (see link to conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Bourdelais_, Dec 09 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015458 | OEIS-A350050

a(n) = (2*n^4 - 6*(-1)^n*n^2 - 2*n^2 + 3*(-1)^n - 3)/96.

Conjectures: (Start) The same conjectures hold for an n X n square matrix A(n) defined as A[i,j,n] = (n mod 2) + abs((n + 1)/2 - j) + abs((n + 1)/2 - i) (see A349108).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Dec 11 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015459 | OEIS-A350053

a(n) = (2^(3*n + 3 + (-1)^n) - (6 + (-1)^n))/9, for n >= 1.

Sorted set of all A385109(A198584(i)), i>0 (conjectured but easy to see). - _Ralf Stephan_, Jun 18 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Jan 20 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015460 | OEIS-A350072

a(n) = sigma(n^2) / gcd(sigma(n^2), A003961(n^2)), where A003961 shifts the prime factorization of n one step towards larger primes, and sigma is the sum of divisors function.

Conjecture: There are no 1's after the initial term. Remark: If there were some k = x^2 > 1, for which a(x) = 1, then sigma(k) would be a divisor of A003961(k). In other words, d = A350073(k) = A064989(sigma(k)) would be a divisor of k. Then, if that divisor were also a unitary divisor [with gcd(d,k/d) = 1], it would need to satisfy the equation sigma(k) = sigma(d) * sigma(k/d) = sigma(A064989(sigma(k))) * sigma(k/A064989(sigma(k))), because sigma is a multiplicative function. (Minor correction by _Antti Karttunen_, Jul 11 2023)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 12 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015461 | OEIS-A350088

Number of ways to write n as t^2 + w^4 + 2*x^4 + 4*y^4 + 8*z^4, where t is a positive integer, and w,x,y,z are nonnegative integers.

Conjecture: If (a,b,c) is among the triples (2,3,4), (2,3,8), (2,4,6), (2,4,7), (2,4,8) and (2,4,11), then any positive integer n can be written as t^2 + w^4 + a*x^4 + b*y^4 + c*z^4, where t is a positive integer, and w,x,y,z are nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015462 | OEIS-A350120

a(n) is the least integer k such that A350090(k) = 2*n-1 or -1 if no such k exists.

Most terms of A350090 are odd. It is conjectured by _Klaus Purath_ that all odd numbers appear in A350090.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Dec 15 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015463 | OEIS-A350126

a(n) = (a(a(n-1)) mod 2) + a(n-2) with a(0) = 0 and a(1) = 1.

A similar formula is conjectured for A156253: (a(a(n-1)) mod 2) + a(n-2) + 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Maiga_, Dec 15 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015464 | OEIS-A350127

a(n) = a(a(n-1)) mod 3 + a(n-2) with a(0) = 0 and a(1) = 1.

Conjecture: The density of even terms is 1/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon Maiga_, Dec 15 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015465 | OEIS-A350132

a(n) is the smallest number m such that tau(m-1) = tau(m+1) = tau(m)^n, where tau(m) is the number of divisors of m (A000005).

Conjecture: a(n) is prime for all n >= 2, i.e., the sequence {a(n)} without the first term is the sequence of smallest primes p such that tau(p-1) = tau(p+1) = 2^n for n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Dec 15 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015466 | OEIS-A350142

Numbers m of the form 2^k + 1 such that tau(m-2) = tau(m-1) - 1.

Conjecture 1: Also numbers m of the form 2^k + 1 such that tau(m - 2) = k. Conjecture 2: If 6th Fermat prime F_p6 exists, then tau(F_p6 - 2) is a power of 2 and tau(F_p6 - 1) = tau(F_p6 - 2) + 1. Conjecture 3: Sequence is finite with 7 terms; supersequence of A262534.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Dec 16 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015467 | OEIS-A350189

Triangle T(n,k) read by rows: the number of symmetric binary n X n matrices with k ones and no all-1 2 X 2 submatrix.

T(n,5) = n*(n-1)*(n-2)*(13*n^2-n-24)/60. T(n,6) = n*(n-1)*(n-2)*(19*n^3-18*n^2-97*n+60)/180. T(n,7) = n*(n-1)*(n-2)*(n-3)*(58*n^3+75*n^2-223*n+180)/1260. - Conjectured by _R. J. Mathar_, Mar 11 2022; proved by _Max Alekseyev_, Apr 02 2022 Conjecture: the largest k such that T(n,k) is nonzero is k = A072567(n) = A001197(n) - 1. - _Max Alekseyev_, Apr 03 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Mar 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015468 | OEIS-A350191

Number of ways of placing maximum number of nonattacking rooks on a triangular board with edge-length n (not counting rotations and reflections as distinct).

Conjecture: a(3*n-2) = A309260(n). There is a nice analogy with A289893(3*n+1) = A002047(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Dec 19 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015469 | OEIS-A350249

a(n) is the constant term in expansion of Product_{k=1..n} (x^(k^2) + 1 + 1/x^(k^2)).

Conjecture: a(n) ~ sqrt(5) * 3^(n + 1/2) / (2*sqrt(Pi)*n^(5/2)). - _Vaclav Kotesovec_, Feb 04 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Jan 28 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015470 | OEIS-A350340

a(n) is the smallest k such that k^2 is an abelian order with precisely 2^n groups.

Except for a(1) = 2, no term can be divisible by 2 or 3. Conjecture: lpf(a(n+1)) >= lpf(a(n)) for all n, where lpf = least prime factor. - _David A. Corneth_ and _Jianing Song_, Jan 03 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Dec 25 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015471 | OEIS-A350341

a(n) is the smallest abelian order with precisely 2^n groups.

Except for a(1) = 4, no term can be divisible by 2 or 3. Conjecture: lpf(a(n+1)) >= lpf(a(n)) for all n, where lpf = least prime factor.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Dec 25 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015472 | OEIS-A350359

Lexicographically earliest infinite sequence of distinct positive integers such that for any four consecutive terms a,b,c,d, d is prime to a and c, but not to b.

Conjecture: The sequence is a permutation of the positive integers in which the primes appear in their natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Dec 26 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015473 | OEIS-A350399

a(n) is the number of prime pairs (p,q) with p <= q, p+q = 2*n, and p*q mod (2*n) prime.

Conjecture: a(n) > 0 for n >= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Dec 28 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015474 | OEIS-A350400

a(n) is the least k with A350399(k) = n, or 0 if there is no such k.

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Dec 28 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015475 | OEIS-A350402

Numbers k such that binomial(k, 2) divides binomial(2^k-2, 2).

Conjecture: aside from the first term, this is a subsequence of A094179 (numbers congruent to 3 mod 4 which are divisible only by primes congruent to 3 mod 4). The conjecture is false: a(2295) = 508606771 = 19531 * 26041 is not in A094179, nor even A004614. - _Charles R Greathouse IV_, Jan 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Dec 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015476 | OEIS-A350410

G.f. A(x) = Sum_{n>=0} a(n)*x^n satisfies: 0 = [x^n] Sum_{m=0..n} A(x)^(m^2) / m!, for n > 1.

This sequence is conjectured to consist entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015477 | OEIS-A350411

G.f. A(x) = Sum_{n>=0} a(n)*x^n satisfies: 0 = [x^n] Sum_{m=0..n+1} A(x)^(m^2) / m!, for n > 1.

This sequence is conjectured to consist entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015478 | OEIS-A350412

G.f. A(x) = Sum_{n>=1} a(n)*x^n satisfies: 0 = [x^n] Sum_{m=0..2*n} (A(x) - 1)^(m^2) / m!, for n > 1, with A(0) = 0.

This sequence is conjectured to consist entirely of integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015479 | OEIS-A350415

Number of acyclic digraphs on n unlabeled nodes with a global source (or sink).

Conjecture: a(n) ~ 2^(n*(n-1)/2) / A245654^n. - _Vaclav Kotesovec_, Sep 11 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Howroyd_, Dec 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015480 | OEIS-A350417

a(n) is the maximum number of rows of width n that can be filled by stacking squares of sizes 1 through n without creating voids or overhangs.

No square of any size can be used more than once. The initial terms listed are conjectured from what I can draw. Trying to stack squares in this way feels like playing Tetris and comes from the squared square idea (see A295130).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antoine Beaulieu_, Dec 29 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015481 | OEIS-A350430

a(n) is the smallest n-th power which can be represented as the sum of n distinct positive n-th powers in exactly n ways, or -1 if none exists.

Conjecture: a(6) = -1. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Dec 30 2021 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015482 | OEIS-A350495

a(n) is the constant term in expansion of Product_{k=1..n} (x^(k^2) + 1/x^(k^2))^2.

Conjecture: a(n) ~ sqrt(5) * 4^n / (sqrt(Pi) * n^(5/2)). - _Vaclav Kotesovec_, Feb 05 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Jan 28 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015483 | OEIS-A350499

Unsigned coefficients of free moment partition polynomials determining the free cumulants from the free moments of free probability theory. Irregular triangle with row lengths given by A000041, n >= 1.

Conjecture: free cumulants in terms of the free moments are R(n,1) for n > 0 where R(n,k) = R(n-1,k+1) - Sum_{j=1..n-1} R(j,k)*R(n-j,1) for n > 1, k > 0 with R(1,k) = m_k for k > 0. - _Mikhail Kurkov_, Mar 30 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tom Copeland_, Jan 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015484 | OEIS-A350520

The number of degree-n^2 polynomials over Z/2Z that can be written as f(f(x)) where f is a polynomial.

Conjecture: Conjecture from _Hugo Pfoertner_, Jan 09 2022: Terms starting at 3 coincide with {A156232}/8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Jan 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015485 | OEIS-A350541

Twin primes x, represented by their average, such that x is the first and x+18 the last of three successive twins.

The distribution of 35314 terms < 10^11 is in accordance with the k-tuple conjecture, see links "k-tuple conjecture" and "Test of the k-tuple conjecture".

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerhard Kirchner_, Jan 06 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015486 | OEIS-A350542

Twin primes x, represented by their average, such that x is the first and x+30 the last of four successive twins.

Subsequence of A014574. For x>6, d=30 is the least possible difference between the least and the greatest of four twins. With d=24, six primes would have the form 6*k+-1, 6*k+6+-1,6*k+12+-1 which is impossible because one of the six numbers would be divisible by 5. Therefore, d>24, except for x=6. The distribution of 1134 terms < 10^11 is in accordance with the k-tuple conjecture, see links

"k-tuple conjecture" and A350541, "Test of the k-tuple conjecture".

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerhard Kirchner_, Jan 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015487 | OEIS-A350543

Twin primes x, represented by their average, such that x is the first and x+48 the last of five successive twins.

Subsequence of A014574. The terms represent quintuples of twin primes. As there are only 31 terms < 10^11, the accordance with the k-tuple conjecture is not very good, see links "k-tuple conjecture" and A350541, "Test of the k-tuple conjecture". Moreover, the formalism of the conjecture allows the evaluation of the expected frequencies of eight types of quintuples relative to the frequency of all quintuples. The differences are considerable:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerhard Kirchner_, Jan 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015488 | OEIS-A350553

a(n) is the smallest number k for which A000120(k) = A000120(k') = n, or -1 if no such k exists, where k' is the arithmetic derivative of k (A003415).

Conjecture. a(n) >= 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marius A. Burtea_, Jan 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015489 | OEIS-A350577

Prime numbers in A036991.

Conjecture: The sequence is infinite. For example, in the first million primes (see A000040) 304208 numbers are terms of A036991.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gennady Eremin_, Jan 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015490 | OEIS-A350583

Coefficients of the expansion of Sum_{n,m>=1} (-1)^m*q^(2*n*m+m)/((1+q^n)*(1-q^(2*m-1))) (odd powers only).

It is conjectured that only odd powers appear in this expansion.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Jan 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015491 | OEIS-A350606

a(n) is the number of elements in row n of A350605.

Conjecture: a(n) is approximately = k*2^(n-5) for n>4. - _Bill McEachen_, Jul 30 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015492 | OEIS-A350636

a(n) is the number of partitions of n in which no part is divisible by 3 minus the number of basis partitions of n.

Andrews conjectures that a(n)>0 for n>3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Jan 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015493 | OEIS-A350689

a(n) = n*(1 - (-1)^n - 2*(3 + (-1)^n)*n^2 + 2*n^4)/384.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jan 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015494 | OEIS-A350693

Number of b > 0 which permit n^3 to be written as a sum of powers of b in n parts. Each exponent c is an integer >= 0, n^3 = b^c_1 + b^c_2 + ... + b^c_n.

a(n) <= A000005(n^3-n). Conjectured to become a(n) = A000005(n^3-n), if the definition would permit negative values for b and only the absolute value of the sum needs to be equal to n^3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Jan 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015495 | OEIS-A350694

Numbers that can be written as p*q+r where p,q,r are primes (not necessarily distinct) such that p+q+r is prime.

Conjecture: the sequence contains all even positive integers except 2, 4, 6, 10, 12, 30, 48, 60.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Jan 28 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015496 | OEIS-A350714

Least positive integer m such that m^12*n = x^4 + y^3 + z^2 for some nonnegative integers x,y,z.

4-3-2 Conjecture: a(n) exists for any nonnegative integer n. Equivalently, each nonnegative rational number can be written x^4 + y^3 + z^2 with x,y,z nonnegative rational numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015497 | OEIS-A350742

Sisyphus with squares: a(1) = 1; thereafter, if a(n-1) is even, a(n) = a(n-1)/2; otherwise a(n) = a(n-1) + k^2, where k^2 is the smallest square (starting from 1) that has not yet been added.

This variation on A350877 was proposed by _Keith F. Lynch_ on 2022 January 28. Because the former sequence uses prime increments, and because squares grow much faster than primes, Lynch felt that my conjecture about the former sequence (that all integers eventually appear) would not be true of this sequence. I responded that because squares nevertheless grow much more slowly than exponentials, we should still expect the same basic argument, that descents to the vicinity of 1 should happen on the order of once per "era", to apply, and I still thought that all values would appear, albeit perhaps very slowly.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Allan C. Wechsler_, Jan 28 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015498 | OEIS-A350747

Number of iterations required to terminate trajectory mapping described in A349824.

Conjecture from A349824: the iteration f: n -> A349824(n) becomes periodic for each n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary Detlefs_, Jan 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015499 | OEIS-A350755

Numbers k such that 11...111 (with k 1's) = (10^k - 1)/9 is a Kaprekar prime.

The terms that are repunit primes (including probable primes) are the only known Kaprekar primes, i.e., (10^19-1)/9 (prime), (10^109297-1)/9 (probable prime), (10^270343-1)/9 (probable prime), and (10^5794777-1)/9 (probable prime). Based on my investigations I conjecture that:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shyam Sunder Gupta_, Jan 14 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015500 | OEIS-A350759

a(n) = Sum_{k=0..n} (-1)^k*A345652(k)*Stirling1(n, k).

Conjectures: For all p prime, (a(p) + a(p+1) - 2) == 0 (mod p),

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mélika Tebni_, Jan 14 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015501 | OEIS-A350767

a(1)=1. Thereafter, a(n+1) is the least unused number k such that either d(j(n)) properly divides d(k) or d(k) properly divides d(j(n)), where j(n) = a(n)+1 and d is the divisor counting function A000005.

Conjecture: This is a permutation of the positive integers. Numbers with the same number (tau) of divisors appear in their natural orders (e.g., primes, semiprimes, squares).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jan 14 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015502 | OEIS-A350768

a(1)=1. Thereafter, if a(n) is a term that has appeared exactly k times prior to and including itself then a(n+1) is the number of terms prior to and including a(n) which have appeared exactly k times.

Conjecture: Every positive integer appears finitely many times (1 appears three times, 2 eight times, 3 seven times, and so on). It seems that after a number j has appeared for the final time in the sequence, it remains possible for subsequent terms to appear j times, but not in isolation, meaning that there are g>1 other numbers which have also appeared j times, so that the next term becomes j*m, rather than j. Henceforth j cannot appear again, but multiples of j can, until they too cease to appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jan 14 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015503 | OEIS-A350778

Numerators of fractions n/prime(n) sorted in decreasing order.

A permutation of the positive integers. Computation of the first n terms of the sequence requires the generation and sorting of more than n fractions; exactly how many more is an open question.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Petr Platais_, Jan 15 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015504 | OEIS-A350831

a(n) is the least positive integer not listed earlier such that its first digit together with a(n-1)'s last digit can form a prime; a(1) = 1.

Conjectured to contain all positive integers not multiple of 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_ and _Eric Angelini_, May 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015505 | OEIS-A350832

a(n) is the least integer > 9 not listed earlier whose first two digits together with a(n-1)'s last two digits form two 2-digit primes when "read vertically in the same direction"; a(1) = 11.

We are tempted to conjecture that this sequence contains all integers > 10 which don't have '0' as second or one of the last two digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_ and _Eric Angelini_, May 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015506 | OEIS-A350856

Initial members of prime triples (p, p+2, p+14).

According to the k-tuple conjecture this sequence is theoretically infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matt C. Anderson_, Jan 19 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015507 | OEIS-A350857

Number of ways to write n as 2*w^4 + x^4 + y^2 + z^2, where w,x,y,z are nonnegative integers with w or x a square.

Conjecture 1: a(n) > 0 except for n = 95, 255. Conjecture 2: For any positive integer a == 2 (mod 4), all sufficiently large integers can be written as a*w^4 + x^4 + (2*y)^2 + z^2 with w,x,y,z integers. If N(a) denotes the largest integer not of the form a*w^4 + x^4 + (2*y)^2 + z^2 (w,x,y,z = 0,1,2,...), then we have N(2) = 255, N(6) = 2716, N(10) = 598, N(14) = 8427, N(18) = 2463, N(22) = 3884, N(26) = 14988, N(30) = 10843. See also A346643 and A347865 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 19 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015508 | OEIS-A350860

Number of ways to write n as w^4 + (x^4 + y^2 + z^2)/81, where w,x,y,z are nonnegative integers with y <= z.

Conjecture 1: a(n) > 0 for any nonnegative integer n. Also, every n = 0,1,2,... can be written as 4*w^4 + (x^4 + y^2 + z^2)/81 with w,x,y,z integers. Conjecture 2: For any positive integer c, there is a positive integer m such that every n = 0,1,2,... can be written as w^4 + (c^2*x^4 + y^2 + z^2)/m^4 with w,x,y,z integers. Conjecture 3: For any positive odd number d, there is a positive integer m such that every n = 0,1,2,... can be written as 4*w^4 + (d^2*x^4 + y^2 + z^2)/m^4 with w,x,y,z integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 19 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015509 | OEIS-A350877

The Sisyphus sequence: start the sequence S with a(1) = 1 and extend S with a(n)/2 when a(n) is even, otherwise with a(n) + the smallest prime not yet added.

Conjecture: On naive probabilistic grounds, all integers should eventually appear. An up-step is always immediately followed by a down-step, and then, on average, by one more down-step. So we expect that every third step will be an up-step, by the next prime number, which will be around p(n/3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Carole Dubois_, Jan 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015510 | OEIS-A350878

Integers m that divide the sum of values d*p < m, where d is a divisor of m, p is a prime, and d*p does not divide m.

Conjecture: The sum of values d*p < m in the definition of the sequence is equal to m for m = 5 only. True for m <= 15000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Devansh Singh_, Jan 20 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015511 | OEIS-A350905

Numbers k such that binomial(k, 3) divides binomial(3^k-3, 2).

Conjecture: all terms of the intersection with A350176 are prime numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jan 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015512 | OEIS-A350963

Coordination sequence for the XXOXX tiling with respect to a tile of type R.

Conjectured g.f.: -(2*t^7-6*t^6-24*t^5-33*t^4-33*t^3-28*t^2-9*t-1)/((1-t^2)*(1-t^4)). Given the decomposition of this structure into eight sectors (see Sigrist's illustration of the first 100 generations), it should be possible to establish this g.f. and those of the other two coordination sequences by using the coloring book method. - _N. J. A. Sloane_, Feb 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015513 | OEIS-A351008

Alternately strict partitions. Number of even-length integer partitions y of n such that y_i > y_{i+1} for all odd i.

Conjecture: a(n+1) = A122129(n+1) - A122130(n). - _Gus Wiseman_, Feb 21 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jan 31 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015514 | OEIS-A351040

Lexicographically earliest infinite sequence such that a(i) = a(j) => A336158(i) = A336158(j), A206787(i) = A206787(j) and A336651(i) = A336651(j) for all i, j >= 1.

Conjectured to be equal to the lexicographically earliest infinite sequence such that b(i) = b(j) => A000593(i) = A000593(j), A336158(i) = A336158(j) and A336467(i) = A336467(j), for all i, j >= 1 (this was the original definition). In any case it holds that a(i) = a(j) => b(i) = b(j) for all i, j >= 1. See comment in A351461.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 31 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015515 | OEIS-A351064

Minimal number of positive perfect powers, with different exponents, whose sum is n (considering only minimal possible exponents for bases equal to 1).

Conjecture: the only numbers for which 5 addends are needed are 15, 23, 55, 62, 71. The numbers mentioned in the conjecture are also the first five terms of A111151. - _Omar E. Pol_, Mar 01 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alberto Zanoni_, Feb 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015516 | OEIS-A351087

Numbers k such that A327860(k) is a multiple of k, where A327860 is the arithmetic derivative of the primorial base exp-function.

Conjecture: For all terms k, A328581(k) = 1 <=> k = A327860(k). This is equal to the conjecture given in A400171. But compare also to the refuted conjecture given in A380527.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015517 | OEIS-A351124

a(n) is the least k > 0 such that the set { prime(n), ..., prime(n+k-1) } can be partitioned into two disjoint sets with equal sum, or -1 if no such k exists (prime(n) denotes the n-th prime number).

Conjecture: all terms are positive.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Feb 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015518 | OEIS-A351141

Pairs of primes (p,q) = (A002496(m), A002496(m+1)) such that q-p is a power r of the product of its prime factors for some m.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Feb 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015519 | OEIS-A351154

a(n) is the permanent of the n X n matrix M(n) that is defined as M[i,j,n] = A351153(n, min(i, j)) + abs(i - j).

Conjectures: (Start) The conjectures are true (see Fried link). - _Sela Fried_, Feb 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Feb 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015520 | OEIS-A351206

Least positive integer m such that n = x^4 + (y^4 + z^4 + 7*w^2)/m^4 for some nonnegative integers x,y,z,w with y <= z.

Conjecture: a(n) exists for any nonnegative integer n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015521 | OEIS-A351221

Least positive integer m such that m^6*n = x^6 + y^3 + z^2 for some nonnegative integers x,y,z.

6-3-2 Conjecture: a(n) exists for any nonnegative integer n. Equivalently, each nonnegative rational number can be written as x^6 + y^3 + z^2 with x,y,z nonnegative rational numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015522 | OEIS-A351228

Numbers k for which A003415(k) >= A276086(k), where A003415 is the arithmetic derivative and A276086 is the primorial base exp-function.

Conjecture: Apart from the initial 6, the rest of terms are the numbers k for which A003415(k) > A276086(k), thus giving the positions of zeros in A351232. In other words, it seems that only k=6 satisfies A003415(k) = A276086(k). See also comments in A351088. Sequence A276086(A391938(.)) sorted into ascending order, thus, if the above conjecture is correct, then apart from the initial 6 this is the complement of (A000040 U A391939).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015523 | OEIS-A351243

Counterexamples to a conjecture of Selfridge and Lacampagne.

Counterexamples to a conjecture of Selfridge and Lacampagne. The conjecture was that every natural number k not divisible by 3 can be written as the quotient of two terms chosen from A147991. It is not known if there are infinitely many counterexamples to the conjecture, but perhaps 3^m+4, for m >= 5 and odd, are.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Feb 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015524 | OEIS-A351259

First noncomposite number reached when iterating the map x -> x', when starting from x = A351255(n). Here x' is the arithmetic derivative of x, A003415.

For the initial 105367 19-smooth terms of A351255, the last 7 occurs here at a(54796), with A351255(54796) = 289993286583 = 3^2 * 7 * 11 * 13^2 * 19^5, and the last 5 occurs here at a(65777), with A351255(65777) = 391899820830375516750 = 2 * 3^2 * 5^3 * 7^3 * 13^3 * 17^3 * 19^6, already a moderately high starting value, in whose vicinity most ending primes for successful iterations are much larger. This observation motivates a conjecture: Even from large numbers with high exponents in their prime factorization it is sometimes possible to reach a small prime. Compare to the conjecture 8 in Ufnarovski & Åhlander paper.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015525 | OEIS-A351261

a(n) is the largest term encountered on the path to 0 when iterating the map x -> x', and starting from x = A351255(n). Here x' means the arithmetic derivative of x, A003415.

and A351255(104776) = 144537549602172859330715888995919357193998109417395984504745753750, the ratio a(n)/A351255(n) obtains another record (~ 42352.7947), which motivates a conjecture that it is not bound from above. See also A351079.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015526 | OEIS-A351306

Least positive integer m such that m^4*n = u^4 + v^4 - (x^4 + y^4) for some nonnegative integers u,v,x,y with x^4 + y^4 <= m^4*n^2.

Conjecture: Each n >= 0 can be written as u^4 + v^4 - (x^4 + y^4), where u,v,x,y are rational numbers with x^4 + y^4 <= n^2. In other words, a(n) exists for any nonnegative integer n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 06 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015527 | OEIS-A351321

Least positive integer m such that m^6*n = u^6 + v^3 - (x^6 + y^3) for some nonnegative integers u,v,x,y with x^6 + y^3 <= m^6*n^2.

6-6-3-3 Conjecture: Each rational number can be written as u^6 - v^6 + x^3 - y^3 with u,v,x,y nonnegative rational numbers. Moreover, a(n) exists for any nonnegative integer n. As a/b = (a*b^5)/b^6 for any integer a and nonzero integer b, the second assertion in the conjecture implies the first one.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015528 | OEIS-A351327

Numbers whose trajectory under iteration of the product of squares of nonzero digits map includes 1.

If k is a term, the number of iterations required to converge to 1 is less than or equal to 3 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luca Onnis_, Feb 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015529 | OEIS-A351336

Odd pseudoprimes to base 3; composite terms of A271116.

a(n) ~ (a(n-1)+a(n-2))/2 (conjectured). _Bill McEachen_, Nov 24 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bill McEachen_, Feb 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015530 | OEIS-A351338

Least nonnegative integer m such that n = x^3 + y^3 - (z^3 + m^3) for some nonnegative integers x,y,z with z <= m.

Conjecture: a(n) exists for any n >= 0. Equivalently, each integer can be written as x^3 + y^3 - (z^3 + w^3) with x,y,z,w nonnegative integers. This is stronger than Sierpinski's conjecture which states that any integer is a sum of four integer cubes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015531 | OEIS-A351341

Least nonnegative integer m such that n = x^4 + y^4 - (z^3 + m^3) for some nonnegative integers x,y,z with z <= m.

Conjecture 1: Let k be 4 or 5. Then each integer can be written as x^k + y^k - (z^3 + w^3) with x,y,z,w nonnegative integers. Conjecture 2: Let k be among 4, 5, 6 and 7. Then any integer can be written as x^k + y^k - (z^2 + w^2) with x,y,z,w nonnegative integers. Conjecture 3: For any integer k > 3, there are no nonnegative integers x,y,z,w such that x^k + y^k - (z^k + w^k) = 3. See also another similar conjecture in A351338.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015532 | OEIS-A351373

a(n) is the least prime that begins a sequence of 2*n consecutive primes whose sum is 6 times a prime.

Conjecture: each odd prime occurs infinitely many times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Feb 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015533 | OEIS-A351375

Least nonnegative integer m such that n = x^5 + y^5 - (z^3 + m^3) for some nonnegative integers x,y,z with z <= m.

Conjecture: a(n) exists for any nonnegative integer n. See also Conjecture 1 in A351341.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015534 | OEIS-A351376

Least nonnegative integer m such that n = x^3 + y^3 - (z^5 + m^5) for some nonnegative integers x,y,z with z <= m.

Conjecture: a(n) exists for any nonnegative integer n. See also Conjecture 1 in A351341.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015535 | OEIS-A351386

Sequence of balanced, multiplicative binary words, starting with a(1)=1, a(2)=10; for j > 2, if j is a prime, then a(j) is obtained by appending 0 at the end of a(j-1); otherwise, a(j) is obtained by appending a single digit at the end of a(j-1) such that the new word is multiplicative, but if the obtained a(j) is not balanced, then we change the digit at the rightmost possible prime position (and, eventually, some digits at following nonprime positions to maintain multiplicativity) so that a(j) becomes balanced.

Rejmer conjectures that this process never terminates (see Grytczuk (2021)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bartlomiej Pawlik_, Feb 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015536 | OEIS-A351410

Numbers m such that the decimal representation of 8^m ends in m.

Conjecture: if k(n) is "expomorphic" relative to "base" b, then the next one in the sequence, k(n+1), consists of the last n+1 digits of b^k(n). This conjecture is true. See A133618. - _David A. Corneth_, Feb 10 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Feb 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015537 | OEIS-A351413

a(n) is the maximum number of stable matchings in the Latin Stable Marriage Problem of order n.

In the Latin Stable Marriage Problem of order n, the sum of a man and woman's rankings of each other is n+1. This implies that the men's and women's ranking tables are Latin squares. As a subproblem of the Stable Marriage Problem, Latin instances provide lower bounds for the maximum number of stable matchings in the general problem, such as A005154 and A065982. For sizes 1 to 4, Latin instances provide exact bounds; they are conjectured to provide exact bounds for sizes a power of 2; they provide the best lower bounds known for sizes 6, 10, 12, and 24, of 48, 1000, 6472, and 126112960, respectively. The next term, a(8), is conjectured to be 268, consistent with A005154. The

minimum number of stable matchings for Latin instances of order n is n, and is realized for the cyclic group of order n. The average number of stable matchings is 7 for n=4 (cf. A351430 showing an average of about 1.5 for the general problem), and benefits from avoidance of mutual first choices and more generally the lack of overlap between the men's and women's preferred matchings. The Latin squares of A005154 and A065982 can be interpreted as multiplication tables of groups, n-th powers of the cyclic group C2 and n-th dihedral groups, respectively.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dan Eilers_, Feb 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015538 | OEIS-A351415

Intersection of Beatty sequences for (1+sqrt(5))/2 and sqrt(5).

Conjecture: every term of the difference sequence is in {2,3,4,5,6}, and each occurs infinitely many times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015539 | OEIS-A351458

Numbers k for which k * gcd(sigma(k), A276086(k)) is equal to sigma(k) * gcd(k, A276086(k)), where A276086 is the primorial base exp-function, and sigma gives the sum of divisors of its argument.

A324644 gives odd values for even numbers and for the odd squares. A324198 is odd on all arguments, therefore on odd squares the above equation reduces to odd * odd = odd * odd, and on odd nonsquares as odd * even = even * odd. It is an open question whether there are any odd terms after the initial a(1)=1. It is conjectured that the intersection of this sequence with the multiperfect numbers (A007691) gives A323653, see comments in the latter.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015540 | OEIS-A351461

Lexicographically earliest infinite sequence such that a(i) = a(j) => A206787(i) = A206787(j) and A336651(i) = A336651(j) for all i, j >= 1.

Conjectured to be equal to the lexicographically earliest infinite sequence such that b(i) = b(j) => A000593(i) = A000593(j) and A336467(i) = A336467(j) for all i, j >= 1. In any case, a(i) = a(j) => b(i) = b(j) for all i, j >= 1 [because both A000593(n) and A336467(n) can be computed from the values of A206787(n) and A336651(n)], but whether the implication holds to the opposite direction is still open. Empirically this has been checked up to n = 2^22. See also comment in A351040. - _Antti Karttunen_, Nov 23 2023 The conjecture in the Nov 23 2023 comment is false. A counterexample is given by i = 21570885 = 3^2*5*7*31*47^2 and j = 39651821 = 11^3*31^3. We have A000593(i) = A000593(j) = 45067776 and A336467(i) = A336467(j) = 27, but A206787(i) = 294912 != 384 = A206787(j) and A336651(i) = 141 != 116281 = A336651(j). The first divergence occurs at n = 39651821; an exhaustive search over odd n <= 2^26 found no earlier one. Since all four functions are invariant under multiplication by powers of 2, it is enough to check odd indices. - _Clément Garrot_, Aug 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015541 | OEIS-A351468

Irregular triangle read by rows where row n is Newey's sequence containing all permutations of 1..n.

For n = 3 to 7, Newey shows these sequences are as short as possible (A062714) for any sequence containing all permutations, and noted the "obvious" conjecture that maybe they would be shortest always. But this is not so, since _Jon E. Schoenfield_ gives a shorter n=16 in A062714, and Radomirovic constructs shorter for all n >= 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kevin Ryde_, Feb 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015542 | OEIS-A351494

a(n) is the least prime p such that n*(p+1)+1 is the square of a prime, or 0 if there is no such p.

Conjecture: all cases where a(n) = 0 arise in this way.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, May 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015543 | OEIS-A351495

a(1) = 1, for n > 1, a(n) is the smallest positive number that has not yet appeared that is a multiple of the smallest prime that does not divide a(n-1).

The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015544 | OEIS-A351496

Lexicographically earliest infinite sequence of distinct positive numbers such that, for n>2, a(n) has a common factor with the largest previous term but not with the second largest previous term.

The primes do not occur in their natural order, and the terms before and after prime values can be a large multiple of the prime, e.g. a(2147) = 3097, a(2148) = 19, a(2149) = 361. The sequence is conjectured to be a permutation of the positive integers although it may take many terms for some primes to appear, e.g., 29 has not occurred after 200000 terms. In the same range the fixed points beyond 2 are 92 and 40100, although it is possible more exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015545 | OEIS-A351497

Lexicographically earliest infinite sequence of distinct positive numbers such that, for n>2, a(n) has a common factor with the second largest previous term but not with the largest previous term.

The primes occur in their natural order, and in general are not divisors of the previous or following term. The sequence is conjectured to be a permutation of the positive integers although it takes many terms for some values to appear, e.g., a(176478) = 42. In the first 200000 terms the fixed points beyond 4 are 7968 and 18552, although it is possible more exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015546 | OEIS-A351509

G.f. A(x) = lim_{n->infinity} (P(n) - 1)/(2*x)^n, where P(1,x) = 1/sqrt(1-4*x), and P(n+1,x) = 1/sqrt(1 - 4*x + 4*x/P(n,x)) for n >= 1.

Conjecture: a(n) ~ 4^n/n. - _Vaclav Kotesovec_, Mar 05 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 14 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015547 | OEIS-A351510

G.f. A(x) = lim_{n->infinity} (1 - P(n,x))/(-2*x)^n, where P(1,x) = 1/sqrt(1 - 4*x), and P(n+1,x) = 1/sqrt(1 - 4*x + 4*x*P(n,x)) for n >= 1.

Conjecture: a(n) ~ 4^n/n. - _Vaclav Kotesovec_, Mar 05 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 14 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015548 | OEIS-A351511

G.f. A(x) = Product_{n>=1} P(n,x), where P(1,x) = 1/sqrt(1-4*x), and P(n+1,x) = 1/sqrt(1 - 4*x + 4*x/P(n,x)) for n >= 1.

Conjecture: a(n) ~ 4^n. - _Vaclav Kotesovec_, Feb 19 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015549 | OEIS-A351538

Numbers k such that both k and sigma(k) are congruent to 2 modulo 4 and the 3-adic valuation of sigma(k) is exactly 1.

All the terms of the form 4u+2 in A349745 (if they exist) are found in this sequence. As A351537 is the intersection of A191218 and A329963, and the latter has asymptotic density zero, so has this sequence also. It is conjectured that A351555(a(n)) is nonzero for all n, which would imply that the intersection with A349745 is empty. - _Antti Karttunen_, Feb 19 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 14 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015550 | OEIS-A351617

Number of ways to write n as 11^w + x^2 + 2*y^2 + 3*z^2 + x*y*z, where w,x,y,z are nonnegative integers.

Conjecture: (i) a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015551 | OEIS-A351625

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number that has not appeared that shares a factor with the largest previous term.

In the first 100000 terms the maximum run of even and odd terms is three and five respectively; it is unknown if these runs have a maximum number of terms or are unbounded. The fixed points beyond 2 are 15, 25, 35, and it is likely no more exist. The primes appear in their natural order, and it is conjectured that this is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015552 | OEIS-A351652

a(1) = 1; for n > 1, a(n) is the smallest positive integer not occurring earlier such that the intersection of the periodic parts of the continued fractions for square roots of a(n) and a(n-1) is the empty set.

Conjecture: This is a permutation of the positive integers. The conjecture is true: we can always extend the sequence with a square, so eventuality every square will appear; also, after a square, we can always extend the sequence with the least number not yet in the sequence. - _Rémy Sigrist_, Mar 12 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giorgos Kalogeropoulos_, Feb 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015553 | OEIS-A351692

a(n) is the least number k such that 1 <= k < n and prime(n) + 2*prime(n-k) and prime(n) + 2*prime(n+k) are both prime, or 0 if there is no such k.

a(n) = 0 for n = 1, 5, 7, 14, 16, 17, 18, 19, 33, 34, 43, 54, 77, 78, 80, 101, 127. Conjecture: these are all the n for which a(n) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, May 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015554 | OEIS-A351716

Starts of runs of 3 consecutive Lucas-Niven numbers (A351714).

Conjecture: 1 is the only start of a run of 4 consecutive Lucas-Niven numbers (checked up to 10^9).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Feb 17 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015555 | OEIS-A351721

Starts of runs of 3 consecutive lazy-Lucas-Niven numbers (A351719).

Conjecture: There are no runs of 4 consecutive lazy-Lucas-Niven numbers (checked up to 3*10^9).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Feb 17 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015556 | OEIS-A351723

Numbers of the form x^2 + y^2 + z^2 + x*y*z with x,y,z nonnegative integers.

Conjecture 1: a(n) < 2*n for all n > 0, and a(n)/n has a limit as n tends to the infinity. Also, a(n) <= a(n-1) + a(n-2) for all n > 4. Conjecture 2: Let S = {x^2 + y^2 + z^2 + x*y*z: x,y,z = 0,1,2,...}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 17 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015557 | OEIS-A351727

Decimal expansion of e^^e (tetration of the base e = 2.718281828... of height e), using H. Kneser's proposal for a tetration function.

A conjectured more accurate value of e^^e is 2075.968335058065833574392795134533608218230677029314\ 70763508334882303759310657791947791267737335609313482 1... obtained with S. Levenstein's PARI program fatou.gp, but its accuracy needs an independent assessment. - _Hugo Pfoertner_, Feb 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Feb 17 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015558 | OEIS-A351743

a(1)=1. Thereafter, a(n+1) is the greatest divisor of s(n) which is prime to a(n), where s(n) is the n-th partial sum.

Conjecture: a(n)=1 occurs infinitely many times, and therefore so does a(n)=2. Conjectures from _Rémy Sigrist_, Jun 20 2022: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jun 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015559 | OEIS-A351824

Irregular triangle read by rows: T(n,k) is the number of partitions of n into 2*k-1 consecutive parts, n >= 1, k >= 1. Column k lists 1's interleaved with 2*k-2 zeros, and the first element of column k is in row A000384(k).

Conjecture 1: T(n,k) is the number of subparts, in an octant of the symmetric representation of sigma(n), that arises from the (2*k-1)-th double-staircase of the double-staircases diagram of n described in A335616. Conjecture 2: Indices of 1's coincide with indices of nonzero terms in A347263, A347529, A351819. For the above conjectures see also the "ziggurat" diagram described in A347186.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Feb 20 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015560 | OEIS-A351832

Least nonnegative integer m such that n = x^6 + y^6 - (z^2 + m^2) for some nonnegative integers x,y,z with z <= m.

Conjecture: a(n) exists for each nonnegative integer n. See also Conjecture 2 in A351341.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015561 | OEIS-A351856

Number of nonnegative integer solutions to 2*n = x_1 + x_2 + ... + x_n + 2*y_1 + 2*y_2 + ... + 2*y_n.

Conjecture: the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) hold for all primes p >= 5 and positive integers n and k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Feb 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015562 | OEIS-A351871

a(1) = 1, a(2) = 2; a(n) = gcd(a(n-1), a(n-2)) + (a(n-1) + a(n-2))/gcd(a(n-1), a(n-2)).

Conjecture: In the first group the sequence always goes to infinity (as in A355898), and in the second group it always goes to a cycle (as in the present sequence).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Augusto Santi_, Feb 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015563 | OEIS-A351876

Numbers whose trajectory under iteration of the product of cubes of nonzero digits map includes 1 (conjectured).

Numbers whose trajectory under iteration of the product of cubes of nonzero digits map includes 1 (conjectured). If k is a term, the number of iterations required to converge to 1 is less than or equal to 10 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luca Onnis_, Feb 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015564 | OEIS-A351888

Number of spanning trees of the graph acquired by placing a vertex in each of the n^2 cells of a triangle of size n in a triangular grid and placing edges between two vertices whenever two cells share a side.

Conjecture: For each k there exists a sufficiently large N, such that for all n > N, if p_1, p_2, ..., p_k are the k largest primes that divide a(n) then (p_1*p_2*...*p_k)^2 also divides a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Feb 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015565 | OEIS-A351889

Table T(n,k) read by downward antidiagonals: period of n-step Fibonacci numbers mod k, n >= 1, k >= 1.

Conjecture 1: T(n,2^m) = (n+1)*2^(m-1) for n > 1. Conjecture 2: For p prime, if T(n,p) != T(n,p^2) then T(n,p^k) = p^(k-1)T(n,p). Conjecture 2 is true for n = 2, n = 3 and n = 4 (see [Wall, 1960], [Waddill, 1978] and [Waddill, 1992] resp.). It is easy to show that T(n,4) != n+1 for all n,

and thus Conjecture 2 implies Conjecture 1. Conjecture 3: T(p^m,p^k) = (p^(pm)-1)*p^(k-1)/(p^m-1) for p prime and k, m > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Feb 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015566 | OEIS-A351899

Integers k for which there exist two distinct prime nondivisors p, q < k such that, for all i, j >= 0, p^i*q^j mod k is either 1 or is divisible by p or q.

Conjecture: The prime nondivisors p and q are elements of the reduced residue system consisting of the totatives of k. Assume a triple (k,p,q) which satisfies the definition. If P and Q are the two subgroups generated by p and q respectively and p < q then P >= Q.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Craig J. Beisel_, Feb 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015567 | OEIS-A351902

Number of ways to write n as w^2 + x^2 + y^2 + z^2 + 3*x*y*z, where w is a positive integer, and x,y,z are nonnegative integers with x <= y <= z.

Conjecture: We have a(n) > 0 except for n = 23. In other words, any positive integer n not equal to 23 can be written as w^2 + x^2 + y^2 + z^2 + 3*x*y*z, where w is a positive integer and x,y,z are nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015568 | OEIS-A351913

Least k such that A352483(k) = n, or -1 if no such k exists.

Conjecture: a(102) and all "Unknown" values in the a-file equal -1. - _Paolo Xausa_, Aug 16 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Mar 18 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015569 | OEIS-A351927

Smallest positive integer k such that 2^k has no '0' in the last n digits of its ternary expansion.

Sloane (1973) conjectured that every power 2^n with n > 15 has a '0' somewhere in its ternary expansion (see A102483 and A346497).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Saye_, Feb 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015570 | OEIS-A351928

Smallest positive integer k such that 2^k has no '2' in the last n digits of its ternary expansion.

Erdős (~1978) conjectured that 1, 4, and 256 are the only powers of two whose ternary expansion consists solely of 0's and 1's.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Saye_, Feb 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015571 | OEIS-A351966

Coordination sequence for the XXOXX tiling with respect to a tile of type G.

Conjectured g.f.: -(-t^9+2*t^8+4*t^7-11*t^6-25*t^5-32*t^4-38*t^3-22*t^2-8*t-1)/((1-t^2)*(1-t^4)). - _N. J. A. Sloane_, Feb 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_ and _N. J. A. Sloane_, Feb 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015572 | OEIS-A351967

Coordination sequence for the XXOXX tiling with respect to a tile of type B.

Conjectured g.f.: -(2*t^9+6*t^8-5*t^7-13*t^6-27*t^5-37*t^4-30*t^3-19*t^2-8*t-1)/((1-t^2)*(1-t^4)). - _N. J. A. Sloane_, Feb 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Feb 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015573 | OEIS-A352040

a(n) is the least number k such that 2^k contains each of the 10 digits at least n times.

Conjecture: a(n) ~ c*n, where c = 10*log_2(10) = 33.21928... .

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Apr 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015574 | OEIS-A352057

Triangular numbers whose nonzero digits are all the same.

Conjecture: the largest term in this sequence whose nonzero digits are not 5 is 600060.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Lu_, Mar 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015575 | OEIS-A352092

Starts of runs of 4 consecutive tribonacci-Niven numbers (A352089).

Conjecture: There are no runs of 5 consecutive tribonacci-Niven numbers (checked up to 10^10).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Mar 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015576 | OEIS-A352097

a(1) = 4, a(2) = 9; let i = a(n-2) and j = a(n-1); a(n) = least k not already in the sequence such that gcd(j, k) = 1 and 1 < gcd(i, k) != i != k.

Conjecture: permutation of the composite numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jun 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015577 | OEIS-A352098

a(1) = 6, a(2) = 15; let i = a(n-2) and j = a(n-1); a(n) = least k not already in the sequence such that gcd(j, k) = 1 and gcd(i, k) = m > 1 and both omega(i) and omega(k) exceed omega(m), where omega = A001221.

Conjecture: permutation of A024619.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jun 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015578 | OEIS-A352110

Starts of runs of 4 consecutive lazy-tribonacci-Niven numbers (A352107).

Conjecture: There are no runs of 5 consecutive lazy-tribonacci-Niven numbers (checked up to 6*10^9).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Mar 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015579 | OEIS-A352117

Expansion of e.g.f. 1/sqrt(2 - exp(2*x)).

Conjectural o.g.f. as a continued fraction of Stieltjes type: 1/(1 - x/(1 - 4*x/(1 - 3*x/(1 - 8*x/(1 - ... - (2*n-1)*x/(1 - 4*n*x/(1 - ...))))))). Cf. A346982. - _Peter Bala_, Aug 22 2023 Conjecture: a(n) = Sum_{k>=1} 2^(n - 3*k - 1/2) * k^n * binomial(2*k, k) for n > 0. - _Velin Yanev_ and _Vaclav Kotesovec_, Jul 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Mar 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015580 | OEIS-A352148

Integers m such that nonzero digits of A000217(m) are all the same.

Conjecture: a(n) = 10^(n-13) for n >= 17.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Lu_, Mar 06 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015581 | OEIS-A352177

a(n) is the number of symmetric Toeplitz anti-Hadamard matrices of order n whose sum of the inverse squares of their singular values is maximal.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Mar 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015582 | OEIS-A352178

Let S = {t_1, t_2, ..., t_n} be a set of n distinct integers and consider the sums t_i + t_j (i<j); a(n) is the maximum number of such sums that are powers of 2, over all choices for S.

Conjecture A: The count of negative numbers in an optimal solution is either equal to the count of positive numbers or one less. This leads to Conjecture B: a(n) <= floor((n+1)*3/2). (End) _Max Alekseyev_ points out that both conjectures are wrong.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015583 | OEIS-A352185

a(n) = smallest integer m for which there is an integer k, with 0 < k < m and gcd(k,m)=1, such that the arithmetic progression m+k, 2*m+k, ..., n*m+k contains only composite numbers.

a(135) = 8207 is the first nonprime term and there are many other counterexamples to the conjecture: a(150..173) = 12311, a(193..195) = 40247, a(196..202) = 40951, ... - _Michael S. Branicky_, Mar 13 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015584 | OEIS-A352187

a(1)=1, a(2)=2; thereafter, a(n) is the smallest number m not yet in the sequence such that gcd(m,a(n-1)) > 1 and gcd(m,a(n-2))=1, except that the second condition is ignored if it would imply that no choice for m were possible.

Conjecture 1: Every positive number appears. Conjecture 2: The primes are the slowest numbers to appear. That is, when a prime p appears here, all numbers less than p have already appeared. To put this another way, the record high points in A352188 ("when does n appear") are exactly the primes (and 1). Conjecture 3: When a prime p first divides some term, that term is 2*p, which is followed by p then 3p. [It seems possible that we could see 2*a, 6*b, 3*p, p, 2*p. This does not happen in the first 10000 terms, but I can't prove it never happens. I can prove that every prime divides some term, and that the primes appear in increasing order. Conjecture 3 might be refuted by a more extensive calculation.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015585 | OEIS-A352189

Index of n-th prime in A352187.

Conjecture: This sequence, prefixed by 1, lists the numbers that take a record length of time to appear in A352188. That is, it appears that this sequence (prefixed by 1) and A352190 together give the RECORDS transform of A352188.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015586 | OEIS-A352190

Indices of records in A352188.

If conjecture (2) in A352187 is true, this is a duplicate of A008578 (1 together with primes). - _R. J. Mathar_, Mar 20 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015587 | OEIS-A352207

a(1) = 2, a(2) = 48; for n >= 1, a(n+2) = c(n)*a(n+1)^2/a(n) where c(n) = 256*(2*n+3)^2*(4*n+1)*(4*n+3)^2*(4*n+5) / (27*(3*n+1)*(3*n+2)^2*(3*n+4)^2*(3*n+5)).

See Problem 8 of the Propp article. Propp conjectured that a(n) is 2-adically continuous as a function of n. The formula is due to _David desJardins_. It is only a conjecture that the formula is the solution to the counting problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 31 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015588 | OEIS-A352233

Numbers that can be expressed as the sum of two primes in exactly 10 ways.

All terms are even. Conjecture: 632 is the last term. Hardy and Littlewood conjectured a growth rate of the number of decompositions for large even numbers (see Conjecture A in page 32 of Hardy and Littlewood reference), implying this sequence is finite. - _Chai Wah Wu_, Mar 10 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Mar 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015589 | OEIS-A352241

Maximal number of nonattacking black-square queens on an n X n chessboard.

Conjecture: a(5k)=4k-1, a(5k+1)=4k, a(5k+2)=4k+1, a(5k+3)=4k+1, a(5k+4)=4k+2. [This does not hold for n = 52 and n = 57. - _Andy Huchala_, Apr 02 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _George Baloglou_, Mar 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015590 | OEIS-A352259

Number of ways to write n as w^6 + x^2 + 2*y^2 + 3*z^2 + x*y*z, where w,x,y,z are nonnegative integers.

Conjecture 1: (i) a(n) > 0 for every n = 0,1,2,.... Moreover, 106, 744, 5469 and 331269 are the only nonnegative integers not in the set {w + x^2 + 2*y^2 + 3*z^2 + x*y*z: w = 0,1; x,y,z = 0,1,2,...}. Conjecture 2: Every n = 0,1,2,... can be written as 2*w^4 + 3*x^2 + y^2 + z^2 + x*y*z, where w,x,y,z are nonnegative integers. We have verified Conjectures 1 and 2 for all n <= 10^5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015591 | OEIS-A352269

Irregular triangle read by rows: T(n,k) is the number of parts in the partition of n into 2*k-1 consecutive parts, 1 <= k <= A351846(n). T(n,k) = 0 if no such partition exists.

Conjecture 1: row sums give A341309. Conjecture 2: A347529, A351819, A347263 have the same indices of zero and nonzero terms as A351824 and this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Mar 18 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015592 | OEIS-A352275

a(0) = 1 and a(n) = Sum_{k = 0..2*n} n/(n + 2*k)*binomial(n + 2*k,k) for n >= 1.

Conjecture: the supercongruences a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) hold for primes p >= 5 and positive integers n and r. Then we conjecture that each sequence u_m satisfies the above supercongruences. This is the case m = 2. See A333093 (case m = 1) and A352276 case (m = 3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015593 | OEIS-A352276

a(0) = 1 and a(n) = Sum_{k = 0..3*n} n/(n + 2*k)*binomial(n + 2*k,k) for n >= 1.

Conjecture: the supercongruences a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) hold for primes p >= 5 and positive integers n and r. Then we conjecture that each sequence u_m satisfies the above supercongruences. This is the case m = 3. See A333093 (case m = 1) and A352275 (m = 2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015594 | OEIS-A352286

Number of ways to write n as w + x^2 + 2*y^2 + 3*z^2 + x*y*z, where w is 0 or 1, and x,y,z are nonnegative integers.

Conjecture 1: a(n) = 0 only for n = 106, 744, 5469, 331269. Thus, for any nonnegative integer n not among 106, 744, 5469 and 331269, either n or n - 1 can be written as x^2 + 2*y^2 + 3*z^2 + x*y*z with x,y,z nonnegative integers. Conjecture 2: a(n) = 1 only for n = 0, 24, 346, 360, 664, 667, 1725, 2589, 3111, 4906, 5035, 8043, 8709, 16810, 18699, 34539, 39256, 51621, 59019, 62799, 108645, 136167, 562696. We have verified Conjectures 1 and 2 for n = 0..10^6. Conjecture 2 is false. In addition to the listed values, a(8710) = a(1269915) = a(1428184) = a(6504010) = a(6901288) = a(38355963) = 1. These were found by an exhaustive scan through 50000000 and verified by direct enumeration of n = w + x^2 + 2*y^2 + 3*z^2 + x*y*z, where w is 0 or 1 and x,y,z are nonnegative integers. - _Scott Moore_, May 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015595 | OEIS-A352296

Smallest number that can be expressed as the sum of two primes in exactly n ways or -1 if no such number exists.

Conjecture: a(n) != -1 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Mar 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015596 | OEIS-A352298

Conjectured largest number that can be expressed as the sum of two primes in exactly n ways or -1 if no such number exists.

Conjectured largest number that can be expressed as the sum of two primes in exactly n ways or -1 if no such number exists. Conjecture A in page 32 of the Hardy and Littlewood reference implies that a(n) != -1 for all n > 1. While the sequence is not monotonic, the plot of n versus a(n)/log(a(n))^2 has a linear trend which matches with the formula for the number of representations in Conjecture A.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Mar 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015597 | OEIS-A352305

a(n) is the (conjectured) largest even number that can be expressed as the sum of two distinct primes in exactly n ways.

a(n) is the (conjectured) largest even number that can be expressed as the sum of two distinct primes in exactly n ways.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Mar 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015598 | OEIS-A352322

Starts of runs of 3 consecutive Pell-Niven numbers (A352320).

Conjecture: There are no runs of 4 consecutive Pell-Niven numbers (checked up to 2*10^8).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Mar 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015599 | OEIS-A352329

Squares in A030299.

Conjecture: there are infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luca Onnis_ and _Marco Ripà_, Mar 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015600 | OEIS-A352337

Numbers m such that A_m(j) = 1 for some value of j, where A_m(k+1) = 2*A_m(k) - sigma(A_m(k)), sigma(m) = sum of the divisors of m, and A_m(1) = m.

Since it is conjectured that 2m - sigma(m) = 1 only for m which are powers of two (see comments at A237588) all numbers in this sequence must have one k for which A_m(k) is a power of two.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Wundling_, Mar 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015601 | OEIS-A352345

Starts of runs of 4 consecutive lazy-Pell-Niven numbers (A352342).

Conjecture: There are no runs of 5 consecutive lazy-Pell-Niven numbers (checked up to 10^9).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Mar 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015602 | OEIS-A352346

Common terms between A061078 and A061077.

Smarandache's conjecture: there are infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luca Onnis_, Mar 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015603 | OEIS-A352356

Number of ways to write 12*n + 5 as 2*x^2 + 5*y^2 + 9*z^2 + x*y*z, where x, y and z are nonnegative integers.

Conjecture: For each n = 0,1,2,... we can write 12*n + 5 as 2*x^2 + 5*y^2 + 9*z^2 + x*y*z with x,y,z nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015604 | OEIS-A352373

a(n) = [x^n] (1/((1 - x)^2*(1 - x^2)))^n for n >= 1.

Conjecture: the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) hold for all primes p >= 5 and positive integers n and k. Conjecture: for each r and s the above supercongruences hold for the sequence a(r,s;n) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 14 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015605 | OEIS-A352394

a(n) = n for n <= 3; let i = a(n-2) and j = a(n-1); a(n+1) = least k not already in the sequence such that (j, k) = 1 and (i, k) = m > 1 and only one of either omega(i) or omega(k) exceed omega(m), where omega = A001221, and either i | k or k | i.

Conjecture: sequence is not a permutation of natural numbers. Proof sketch: Since either i | k or k | i, and defining m as the smaller of the 2 terms, as n increases, it becomes harder to reach all numbers through multiplication or division by m. Therefore it would seem that there is strong tendency for the sequence to fall into multiplicative recurrence as does A113552.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jun 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015606 | OEIS-A352408

a(n) = [x^(n+1)] (1+x - x^2*C(x^3))^(n*(2*n+1)) / (n*(2*n+1)), for n >= 1, where C(x) = 1 + x*C(x)^2 is the Catalan function (A000108).

Conjecture: all terms are divisible by 3. Showing that a(n) = 0 (mod 3) for n >= 1 would consequently prove conjectures involving the g.f. G(x) of A352409, namely: G(x) = 1 + x^3*G(x)^2 (mod 3) and G(x) = C(x^3) (mod 3), where C(x) = 1 + x*C(x)^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015607 | OEIS-A352409

G.f. A(x) satisfies: [x^(n+1)] (1+x - x^2*A(x))^(n*(2*n+1)) = 0, for n >= 0.

Conjectures: Conjectures: g.f. A(x) = Sum_{n>=0} a(n)*x^n satisfies:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 15 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015608 | OEIS-A352425

Irregular triangle read by rows in which row n lists the partitions of n into an odd number of consecutive parts.

Conjecture: the total number of parts in all partitions of n into an odd number of consecutive parts equals the sum of odd divisors of n that are <= A003056(n). In other words: row n has A341309(n) terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Mar 15 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015609 | OEIS-A352446

Total number of parts in all partitions of n into an even number of consecutive parts.

a(n) = A204217(n) - A341309(n), conjectured. Conjecture: G.f.: Sum_{k>=1} 2*k*x^(k*(2*k+1))/(1-x^(2*k)). - _Vaclav Kotesovec_, Oct 23 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Mar 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015610 | OEIS-A352472

Triangle T(n,k) read by rows: the number of traceless symmetric binary n X n matrices with 2k one's and no all-1 2 X 2 submatrix.

T(n,2) = 3*A000332(n+1). T(n,3) = A093566(n+1). - Conjectured by _R. J. Mathar_, Mar 25 2022; proved by _Max Alekseyev_, Apr 02 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Mar 17 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015611 | OEIS-A352478

a(n) is the number of Toeplitz anti-Hadamard matrices of order n whose sum of the inverse squares of their singular values is maximal.

Conjecture: a(1) = 1 and a(3) = 3 are the only odd terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Mar 17 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015612 | OEIS-A352503

Number of ways to write n as w^3 + 2*x^3 + 4*y^3 + 5*z^3 + t^6, where w is a positive integer, and x,y,z,t are nonnegative integers.

Conjecture: a(n) > 0 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 28 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015613 | OEIS-A352511

Starts of runs of 4 consecutive Catalan-Niven numbers (A352508).

Conjecture: There are no runs of 5 consecutive Catalan-Niven numbers (checked up to 10^9).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Mar 19 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015614 | OEIS-A352540

Values for which the iteration of A352544 (half if even, add largest anagram if odd) does not end in a loop.

All the terms are only conjectured to have this property; we don't have a completely rigorous proof. But for all the listed initial terms, the trajectory quickly reaches numbers with many (>> 10) digits and grows larger with every iteration: When the number is odd and has a digit 0, then its successor is again odd and at least twice as large, most often more than 9 times larger. Roughly 1/10th of the digits are zeros, and similarly for 9s, so as the terms get larger, it becomes increasingly less probable that they could end up having no digit 0 at all, which is only a necessary condition that they might become even and not grow upon for one iteration, but still most likely resume growth immediately after. See sequence A352542, the trajectory of a(1) = 89, for an example studied in detail.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Mar 20 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015615 | OEIS-A352542

Trajectory of initial value 89 under iterations of the map A352544: half if even, add largest anagram if odd.

We conjecture that there is a strictly increasing sequence (b(k), k >= 0) = (32, 37, 46, 52, 88, 91, 118, 122, 141, ...) such that all terms a(n) with n >= b(k) have more than k digits 0. a(n+1) > 9*a(n) for all n > 50. (Conjectured.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Mar 20 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015616 | OEIS-A352568

Take n equally spaced points on circle, connect them by a path with n-1 line segments; sequence gives number of distinct multisets of segment lengths.

Say the type of a segment is its position in the increasing list of floor(n/2) possible segment lengths. The Buratti-Horak-Rosa conjecture is that the following condition is necessary and sufficient for a multiset to be realizable as a path: for each divisor d of n, the number of segments of type divisible by d is at most n-d. The sequence confirms the conjecture up to n=37. - _Brendan McKay_, May 14 2022 If the Buratti-Horak-Rosa conjecture is true, the values of a(38)-a(50) are 144079707346575, 212327989773900, 947309492837400, 1397281501935165, 6236646703759395, 9206478467454345, 41107996877935680, 60727722660586800, 271250494550621040, 400978991944396320, 1791608261879217600, 2650087220696342700, 11844267374132633700. - _Brendan McKay_, Jun 14 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015617 | OEIS-A352582

Two-column array read by rows, where the n-th row is the least pair of integers (p, q) such that f(p) = f(n) + q*f(n+1) where f(n) = A002496(n) is the n-th prime of the form k^2+1.

Conjecture: Given an integer n, there always exists a pair (p, q) such that f(p) = f(n) + q*f(n+1). Consequence: if the conjecture is true, then the set of prime numbers of the form k^2+1 is infinite because, by induction, there exists a pair

(p', q') such that f(p') = f(p-1) + q'*f(p), f(p') > f(p).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Mar 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015618 | OEIS-A352594

Expansion of square root of the golden ratio phi in base phi.

What is lim_{n->oo} (1/n)*Sum_{k=1..n} a(k)? (The value is near 0.2765 at n=10^6.) - _Vaclav Kotesovec_, Mar 23 2022 [Conjecture: This value is 1/(sqrt(5)*phi) (A244847). - _Amiram Eldar_, Mar 25 2022]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chittaranjan Pardeshi_, Mar 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015619 | OEIS-A352596

Conjecturally the number of positive even integers that can be expressed as the sum of two distinct primes in exactly n ways.

Conjecturally the number of positive even integers that can be expressed as the sum of two distinct primes in exactly n ways.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Mar 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015620 | OEIS-A352627

Number of ways to write n as a^2 + 2*b^2 + c^4 + 4*d^4 + c^2*d^2, where a,b,c,d are nonnegative integers.

Conjecture: a(n) > 0 for all n = 0,1,2,.... In other words, each nonnegative integer can be written as a^2 + 2*b^2 + c^4 + 4*d^4 + c^2*d^2 with a,b,c,d integers. See also A352628, A352629 and A352632 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015621 | OEIS-A352628

Number of ways to write n as a^2 + 2*b^2 + c^4 + 2*d^4 + 3*c^2*d^2, where a,b,c,d are nonnegative integers.

Conjecture: a(n) > 0 for all n = 0,1,2,.... In other words, each nonnegative integer can be written as a^2 + 2*b^2 + (c^2+d^2)*(c^2+2*d^2) with a,b,c,d integers. See also A352627, A352629 and A352632 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015622 | OEIS-A352629

Number of ways to write n as a^2 + 2*b^2 + c^4 + 9*c^4 + 3*c^2*d^2, where a,b,c,d are nonnegative integers.

Conjecture: a(n) > 0 for all n = 0,1,2,.... In other words, any nonnegative integer can be written as a^2 + 2*b^2 + c^4 + 9*d^4 + 3*c^2*d^2 with a,b,c,d integers. See also A352627, A352628 and A352632 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015623 | OEIS-A352632

Number of ways to write n as (a+b)*(a+2*b) + c^4 + 2*d^4, where a is a positive integer, and b,c,d are nonnegative integers.

Conjecture: a(n) > 0 for all n > 0. In other words, any positive integer n can be written as u*v + x^4 + 2*y^4, where u,v,x,y are nonnegative integers with 0 < u <= v < 2*u. See also A352627, A352628 and A352629 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015624 | OEIS-A352636

Number of ways to write n as a^2 + 3*b^2 + c^4 + 2*d^4 + a*b*c*d, where a,b,c,d are nonnegative integers.

Conjecture: a(n) > 0 except for n = 46.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Mar 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015625 | OEIS-A352651

a(n) = (binomial(5*n,2*n)*binomial(5*n/2,2*n)*binomial(2*n,n)^2) / binomial(5*n/2,n)^2.

Conjecture: the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) hold for all primes p >= 5 and positive integers n and k. The case n = k = 1 is easily proved. More generally, for an integer N not equal to 0 or 1, the height 2 factorial ratio sequence whose n-th term is given by (binomial(N*n,2*n)* binomial(N*n/2,2*n)* binomial(2*n,n)^2)/binomial(N*n/2,n)^2 is conjectured to be integral and satisfy the same supercongruences. This is the case N = 5. See A352652 (N = 7)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015626 | OEIS-A352652

a(n) = (binomial(7*n,2*n)*binomial(7*n/2,2*n)*binomial(2*n,n)^2) / binomial(7*n/2,n)^2.

Conjecture: the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) hold for all primes p >= 5 and positive integers n and k. The case n = k = 1 is easily proved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015627 | OEIS-A352653

a(n) = Sum_{k = 0..n-1} binomial(n,k)*binomial(n-1,k)*binomial(n+k,k)*binomial(n-1+k,k).

More generally, for nonnegative integers A, B and C with A >= 2 let S(n;A,B,C) = Sum_{k = 1..n} binomial(n,k)^A * binomial(n+k-1,k-1)^B * binomial(2*k,n)^C. Then we

conjecture the sequence (S(n;A,B,C))n>=1 satisfies the same supercongruences. This is the case A = B = 2, C = 0. Compare with Osburn et al., Theorem 1.2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 06 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015628 | OEIS-A352654

a(0) = 0, a(n) = [x^n] P(n-1,(1 + x)/(1 - x)) for n >= 1, where P(n,x) denotes the n-th Legendre polynomial.

Conjecture: a(n) = [(x*y)^(n-1)*z^n] 1/(1 - x - x*y - y*z - x*z - x*y*z) = [(x*z)^(n-1)*y^n] 1/(1 - x - x*y - y*z - x*z - x*y*z) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 14 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015629 | OEIS-A352655

a(n) = (1/2)*(A005258(n) + A005258(n-1)).

The Apéry numbers A005258 satisfy the supercongruences A005258(p) == 3 (mod p^3) and A005258(p-1) == 1 (mod p^3) for primes p >= 5. It easily follows that a(p) == 2 (mod p^3) for primes p >= 3. We conjecture that the stronger supercongruences a(p) == 2 (mod p^5) hold for primes p >= 5. See A212334 for the corresponding conjecture for the Apéry numbers A005259. Conjecture: for r >= 2, and all primes p >= 5, a(p^r) == a(p^(r-1)) (mod p^(3*r+3)). - _Peter Bala_, Oct 13 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 17 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015630 | OEIS-A352656

The number of lozenge tilings of a semiregular hexagon of side lengths n, n, 2*n, n, n and 2*n; equivalently, the number of plane partitions whose solid Young diagram fits inside an n X n X 2*n box.

Conjecture 1: the supercongruences F(a*p^r,b*p^r,c*p^r) == F(a*p^(r-1),b*p^(r-1),c*p^(r-1))^p (mod p^(4*k)) hold for all primes p, where r is a positive integer and a, b and c are nonnegative integers. Conjecture 2: the Gauss congruences a(n*p^r) == a(n*p^(r-1)) (mod p^r) hold for all primes p and positive integers n and r. If true, then the expansion of exp(Sum_{n >= 1} a(n)*x^n/n) has integer coefficients. Conjecture 3: the supercongruences a(n*p^r) == a(n*p^(r-1))^p (mod p^(4*r)) hold for all primes p and positive integers n and r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015631 | OEIS-A352657

The number of lozenge tilings of a semiregular hexagon of side lengths n, n, 3*n, n, n and 3*n; equivalently, the number of plane partitions whose solid Young diagram fits inside an n X n X 3*n box.

Conjecture 1): the Gauss congruences a(n*p^r) == a(n*p^(r-1)) (mod p^r) hold for all primes p and positive integers n and r. If true, then the expansion of exp(Sum_{n >= 1} a(n)*x^n/n) has integer coefficients. Conjecture 2): the supercongruences a(n*p^r) == a(n*p^(r-1))^p (mod p^(4*r)) hold for all primes p and positive integers n and r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015632 | OEIS-A352667

Maximum number of induced copies of the paw graph in an n-node graph.

|(conjectural)|

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pontus von Brömssen_, Mar 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015633 | OEIS-A352676

Intersection of Beatty sequences for sqrt(3) and 1+sqrt(3).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015634 | OEIS-A352700

G.f.: Sum_{n>=0} binomial((n+1)*(2*n+1),n)/(2*n+1) * x^n / C(x)^(n*(2*n+1)+1), where C(x) = 1 + x*C(x)^2 is the Catalan function (A000108).

Conjectures: given g.f. A(x), let C(x) = (1 - sqrt(1-4*x))/(2*x) be the Catalan power series (A000108), then

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 29 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015635 | OEIS-A352742

a(n) is the smallest number > 1 that is not divisible by 10 but is divisible by the n-th power of the sum of its digits.

Conjecture: No term is a multiple of 5. a(28) = 265^28, disproving the above conjecture. - _Charles R Greathouse IV_, Apr 02 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nicolas Bělohoubek_, Mar 31 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015636 | OEIS-A352743

a(n) = Product_{k=1..n} (p(k+1) + p(k))/(p(k+1) - p(k)), where p(k) = prime(k).

Conjecture: a(n) is an integer for every natural n. - _Thomas Ordowski_ a(0) = 1, a(n) is the largest k such that b(n+1) = b(n)*(k + a(n-1))/(k - a(n-1)) is prime, where b(1) = 2. By my conjecture, b(n) = prime(n). - _Thomas Ordowski_, Jul 30 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Apr 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015637 | OEIS-A352763

a(1) = 1; a(2) = 2; for n > 2, a(n) is the smallest positive number that has not appeared that shares a factor with a(n-1) and whose binary expansion has no 1-bit in common with the binary expansion of a(n-1).

In the first 50000 terms the fixed points are 1, 2, 105, 135, 225, 2157, 3972, 7009, 8531, although it is likely more exist. In the same range the lowest unseen number is 383; the sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 15 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015638 | OEIS-A352796

Numbers m such that {d + m/d : d | m } does not contain consecutive integers.

Conjecture: Complement of A072389.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _José María Grau Ribas_, Jun 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015639 | OEIS-A352797

Let S(k) be the subsequence of multiples of k from k*positive integers where each element of S(k) sets a new record of divisors in that sequence. Then f(k) is the first element from S(k)/k that is not a power of 2. Sequence lists records for f(k).

Conjecture: Subsequence of A147516.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Apr 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015640 | OEIS-A352801

Number of symmetric n X n 01-matrices with maximum number of 1s and no 2 X 2 all-1 submatrix.

Conjecture: the maximum number of 1s equals A072567(n), and so a(n) = A350189(n, A072567(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Apr 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015641 | OEIS-A352833

Irregular triangle read by rows where T(n,k) is the number of integer partitions of n with k fixed points, k = 0, 1.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Apr 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015642 | OEIS-A352834

Least k > 0 such that denominator(d(k*n)/(k*n)) = n, or 0 if no such k exists, where d = A000005 is the number-of-divisors function.

Conjecture: a(n) = O(n^(1/3)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015643 | OEIS-A352844

Smallest k > 1 such that sopfr(k) - tau(k) = n, or -1 if no such k exists.

Conjecture: There is no -1 in this sequence. Conjecture above is true. Assume k = 2^(n + 1). Then sopfr(k) = 2n + 2, tau(k) = n + 2. Therefore, sopfr(k) - tau(k) = n. - _Nicolas Bělohoubek_, Jul 09 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nicolas Bělohoubek_, Apr 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015644 | OEIS-A352849

a(n) is the least k not already in the sequence such that k is pairwise coprime to a(n-1) and a(n-2), starting with a(1) = 1, a(2) = 3, and a(3) = 5.

Conjecture: the sequence is a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Apr 14 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015645 | OEIS-A352905

Expansion of e.g.f. sin(x) * exp(exp(x) - 1).

Conjecture: a(n) = (i/(2*e))*Sum_{k=0..oo} ((k - i)^n - (k + i)^n)/(k!), where i = sqrt(-1) and e = exp(1). - _Velin Yanev_, Jul 06 2024 Yanev's conjecture is true (see Fried link). - _Sela Fried_, Dec 07 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Apr 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015646 | OEIS-A352920

Values of A109812(k) where k/A109812(k) reaches a new high point.

This suggests the following conjecture: Conjecture: r(k) > k for all k > 4. Combining this with the conjecture that A109812(k)/k is bounded (see A352919 and A352920), we have: Conjecture: k/A109812(k) is unbounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Broadhurst_, Aug 17 2022 (entry created by _N. J. A. Sloane_, Apr 23 2022) | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015647 | OEIS-A352931

a(n) = A093714(n) - n.

Sequence exhibits intervals where terms have same parity. Parity changes when a run of odd terms in A093714 with even run length occurs. These runs begin with A093714(A351498(k)). Conjecture: the last parity change begins with a(1036044) = -9. (End) Rival conjecture: The parity changes infinitely often. This would imply that A352932 contains infinitely many terms. - _N. J. A. Sloane_, May 04 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015648 | OEIS-A352943

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number that has not appeared that shares a factor with the sum of the largest and second largest value of all previous terms.

The sequence is conjectured to be a permutation of the positive integers, although it apparently takes many terms for some primes to appear, e.g., after 200000 terms 73 has not occurred. The primes do not occur in their natural order. Beyond the first three terms there are nine fixed points in the first 200000 terms, although it is likely more exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 06 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015649 | OEIS-A352968

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number that has not appeared that shares a factor with min(a(n-2),a(n-1)).

In the first 500000 terms there are thirty-eight fixed points - 1, 2, 3, 4, 14, 32, 85, ..., 3277, 8651, 9223. It is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Apr 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015650 | OEIS-A352976

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number that has not appeared that shares a factor with max(a(n-2),a(n-1)).

In the first 500000 terms the fixed points are 1, 2, 15, 25, 35. It is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Apr 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015651 | OEIS-A352991

Concatenation of all the distinct permutations of the first 1, 2, 3, ... (strictly) positive integers, arranged in ascending numerical order.

A001292 is a subsequence of the present sequence. An open problem, published by Kenichiro Kashihara in 1996 (see References, p. 25, #30, Problem 2), is to find how many terms of A001292 (which is a subsequence of A030299) are powers of integers; Kashihara conjectured that there are none (even if, clearly, A001292(1) = 1 should be disregarded in order to keep the conjecture alive). Currently, only the terms up to the prime a(409120) = 10123457689 have been directly checked by the author of this sequence and no nontrivial perfect power has been found. On the other hand, many (maybe infinitely many) terms of the present sequence are nontrivial powers of integers (e.g., A352329(2) to A352329(36) are squares of integers and belong to this sequence).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Apr 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015652 | OEIS-A353006

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number that has not appeared that shares a factor with (a(n-1) + a(n-2))*|a(n-1) - a(n-2)|.

The sequences is conjectured to be a permutation of the positive integers. In the first 500000 terms there are 637 fixed points: 1, 2, 3, 6, 9, ..., 180611, 189383, 298097. As the terms slowly move away from the line a(n) = n as n increases it is likely no more exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Apr 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015653 | OEIS-A353008

a(n) is the smallest positive k such that k^2 + 1 has 2*n divisors, or -1 if no such k exists.

Conjecture: a(n) is never -1. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, May 15 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015654 | OEIS-A353010

a(n) = maximal d such that Product_{k=0..m} binomial(m,k) is divisible by m^(m+d), where m = A276710(n).

It is conjectured that a(n) >= 0, computationally verified up to n = 10^7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hagen von Eitzen_, Apr 15 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015655 | OEIS-A353025

Terms of A352991 which are perfect powers.

a(n) = m^2, where the integer m := m(n) is not a perfect power itself (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Apr 17 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015656 | OEIS-A353026

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number that has not appeared that is a multiple of the smallest prime factor with minimal exponent of a(n-1) (cf. A067695).

The sequences is conjectured to be a permutation of the positive integers. In the first 500000 terms any time a prime p appears, where p>=5, the following term is 5p. It is unknown if this is true for all primes. In the same range the fixed points are 1, 2, 9, 39, 49, 1079, 4897, although it is possible more exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Apr 18 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015657 | OEIS-A353047

Number of length n words on alphabet {0,1,2} that contain each of the subwords 01, 02, 10, 12, 20, and 21 as (not necessarily contiguous) subwords.

Appears to equal 12 times A222993, except that sequence only has a conjectured formula. - _N. J. A. Sloane_, Jun 17 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Critzer_, Apr 19 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015658 | OEIS-A353050

G.f. A(x) satisfies: 0 = Sum_{n>=1} (Lucas(n) - A(x))^n * x^n/n, where Lucas(n) = A000204(n).

It is conjectured that this is an integer sequence. This sequence is a special case of a more general conjecture: 0 = Sum_{n>=1} (L(n) - F(x))^n * x^n/n is satisfied by an integer series F(x) if exp(Sum_{n>=1} L(n) * x^n/n) yields an integer series.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015659 | OEIS-A353071

Maximum number of clicks needed to solve any solvable Lights Out problem on an n X n grid.

If n = 6k-1 for some integer k, then a(n) <= 26k^2 - 12k + 1. This upper bound is equal to a(n) when A159257(n) = 2. Further, it is conjectured that if A159257(n) = 2, then n = 6k-1 for some integer k. It is conjectured that if A159257(n) = 4, then n = 5k-1 for some integer k, and a(n) = 17k^2 - 10k. It is conjectured that if A159257(n) = 6, then n = 12k-1 for some integer k, and a(n) = 88k^2 - 24k + 1 It is conjectured that if A159257(n) = 8, then either n = 10k-1 or n = 17k-1 for some integer k. If n = 10k-1, then a(n) = 60k^2 - 20k - 3. If n = 17k-1, then a(n) = 161k^2 - 34k - 3. It is conjectured that if A159257(n) = 10, then n = 30k-1 for some integer k, and a(n) = 506k^2 - 60k - 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _William Boyles_, Apr 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015660 | OEIS-A353077

Triangle read by rows, where the n-th row consists of the lexicographically earliest solution for n integers in 0..p-1 whose n*(n-1) differences are congruent to 1..p-1 (mod p), where p=n*(n-1)+1. If no solution exists, the n-th row consists of n -1's.

In the linked a-file, values a(n) > 3 from row 17 onwards depend on the conjecture that all perfect difference sets are Singer type, and were obtained through computer enumeration of Singer type sets.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Apr 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015661 | OEIS-A353138

Sum of (the number of digits in n to the power (each digit in n)).

In base 10, 1 and 4624 are the only numbers where a(n)=n (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _William Halpin_, Apr 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015662 | OEIS-A353154

Sum of the even-indexed terms in the n-th row of the triangle A196020.

Conjecture: indices of zeros give A082662.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Apr 27 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015663 | OEIS-A353195

Coefficients of the open mirror map of P2.

Conjecture: a(n) = (3*n - 1)*A364973(n). - - _Kyler Siegel_, Jul 06 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tim Graefnitz_, Apr 29 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015664 | OEIS-A353239

Lexicographically earliest infinite sequence of distinct positive numbers such that, for n > 2, a(n) has either a common factor with a(n-1) but not with a(n-2), or with a(n-2) but not with a(n-1).

The sequence is conjectured to be a permutation of the positive integers. Because of the selection rule at most two consecutive terms can be even, although the number of consecutive odd terms is likely arbitrarily large.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Apr 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015665 | OEIS-A353313

If n is of the form 3k, then a(n) = k, and if n is of the form 3k+r, with r = 1 or 2, then a(n) = 5*k + 3 + r.

It is conjectured that all iterations of this sequence starting from any n >= 0 will eventually reach a finite cycle, which by necessity then contains at least one multiple of three. See Drozd links and A349876.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Apr 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015666 | OEIS-A353408

a(n) is the least new prime number such that a(n) + a(n-1) is a semiprime.

For n >= 4, a(n) == 11 (mod 12). Conjecture: every prime == 11 (mod 12) occurs in the sequence. - _Robert Israel_, Mar 04 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_, May 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015667 | OEIS-A353448

Number w is in this sequence if every frame w X h, w >= h >= 3, contains more distinct distance quadrilaterals with corners interior to the 4 sides with concurrent diagonals, i.e., both ascending or both descending, than non-concurrent diagonals, or equivalently A353450(w,h) >= A353449(w,h).

It is conjectured that this sequence is a subsequence A160007 except for the small terms <= 46 (verified for all w < 661). The example section depicts the way this sequence matches the irregular pattern of A160007. Numbers w > 100 were computed by _Hugo Pfoertner_.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rainer Rosenthal_, May 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015668 | OEIS-A353638

a(n) = 1 if n is odd and A064989(sigma(n)) < A064989(n), otherwise 0. Here A064989 shifts the prime factorization one step towards lower primes, and sigma is the sum of divisors function.

For n > 1, a(n) = A000035(n) - A353639(n). [Conjectured, see comments in A336702]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, May 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015669 | OEIS-A353654

Numbers whose binary expansion has the same number of trailing 0 bits as other 0 bits.

Conjecture: 2*A247648(n-1) + 1 with rewrite 1 -> 1, 01 -> 0 applied to binary expansion is the same as a(n) without trailing 0 bits in binary. Conjectured formulas: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Jul 15 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015670 | OEIS-A353657

a(n) = A353655(n) - A353656(n).

Conjectures: a(n) = 0 for infinitely many n, and (a(n)) is unbounded below and above.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, May 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015671 | OEIS-A353709

a(0)=0, a(1)=1; thereafter a(n) = smallest nonnegative integer not among the earlier terms of the sequence such that a(n) and a(n-2) have no common 1-bits in their binary representations and also a(n) and a(n-1) have no common 1-bits in their binary representations.

Conjecture: This is a permutation of the nonnegative numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 06 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015672 | OEIS-A353717

a(n) = index of n in A353709, or -1 if n does not appear there.

If, as conjectured, A353709 is a permutation of the nonnegative integers, then this is the inverse permutation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015673 | OEIS-A353735

Number of n-digit terms in A333369.

Conjecture 1: lim_{n->oo} a(2n+1)/a(2n-1) = 100.
Conjecture 2: lim_{n->oo} a(2n+2)/a(2n) = 81. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael S. Branicky_, May 06 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015674 | OEIS-A353736

Number of n-digit terms in A353007.

Conjecture 1: lim_{n->oo} a(2n+1)/a(2n-1) = 100.
Conjecture 2: lim_{n->oo} a(2n+2)/a(2n) = 81. These conjectures are the same as for A353735. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael S. Branicky_, May 06 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015675 | OEIS-A353757

a(n) = A353750(n) - A353749(n).

It is conjectured that there are no other zeros after a(1) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, May 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015676 | OEIS-A353780

a(1) = 1; for n > 1, a(n) is the smallest positive number that has not yet appeared that is coprime to a(n-1), does not equal a(n-1)+1, and has no digit in common with a(n-1).

The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015677 | OEIS-A353794

a(n) = A353791(sigma(A003961(n))), where A353791(n) = A003958(n) * A064989(n).

It is conjectured that a(n) is not a multiple of A353793(n) on any other n except on n=1. See also A353795.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, May 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015678 | OEIS-A353890

a(n) is the period of the binary sequence {b(m)} defined by b(m) = 1 if (m+1)^n - m^n and (m+2)^n - 2*(m+1)^n + m^n are coprime, 0 otherwise.

For any n, consecutive n-th powers will never share a divisor > 1, so now consider the second differences. Specifically, each m > 0, define the binary sequence {b(m)} as follows: b(m) = 1 if the first difference (m+1)^n - m^n and the second difference (m+2)^n - 2*(m+1)^n + m^n are coprime, 0 otherwise. I conjecture that {b(m)} is periodic with period a(n). I conjecture that the greatest prime factor for any prime n >= 5 is phi(2^n+1)/2 + 1 = Jacobsthal(n). n*A069925 + 1 = A001045(n). I conjecture that all prime factors "f" are f=n*k+1, unless n is composite, in which case additionally all prime factors for any divisor of n will also be included in the prime factorization for a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Samuel Harkness_, May 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015679 | OEIS-A353904

a(1) = 1; for n > 1, a(n) is the smallest positive number that has not yet appeared that is coprime to a(n-1), does not equal a(n-1)+1, and differs from a(n-1) at every digit.

The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015680 | OEIS-A353905

a(1) = 1; for n > 1, a(n) = smallest positive number that has not appeared that has a common factor with a(n-1) + the smallest unseen positive number.

The sequence is conjectured to be a permutation of the positive integers. In the first 100000 terms the number 18869 holds the record for the greatest number of terms for which it is the lowest unseen number, 4769 terms in all. In the same range there are fifteen fixed points, the last being a(1204), and it is likely no more exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015681 | OEIS-A353916

a(1) = 1, a(2) = 2. Let j = a(n-1) and let m = omega(gcd(j, k)) with gcd(j, k) > 1. For n > 2, a(n) = least k such that min(omega(j), omega(k)) = m and m < max(omega(j), omega(k)).

Open question: do the primes appear in order? (They do for n <= 2^16).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, May 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015682 | OEIS-A353917

a(1) = 4. Let j = a(n-1) and let m = omega(gcd(j, k)) with gcd(j, k) > 1. For n > 1, a(n) = least k such that min(omega(j), omega(k)) = m and m < max(omega(j), omega(k)), but neither j | k nor k | j.

Conjecture: permutation of composite numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, May 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015683 | OEIS-A353944

Product_{n>=1} 1 / (1 - a(n)*x^n) = Sum_{n>=0} Bell(n)*x^n, where Bell = A000110.

Conjecture: a(n) ~ Bell(n). - _Vaclav Kotesovec_, May 12 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, May 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015684 | OEIS-A353989

a(1) = 1; a(2) = 3; for n > 2, a(n) is the smallest positive number that has not appeared that shares a factor with a(n-1) and whose binary expansion has a 1-bit in common with the binary expansion of a(n-1).

In the first 100000 terms the fixed points are 1, 16, 32, 209, 527, and it is likely no more exist. In the same range the lowest unseen number is 34849; the sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015685 | OEIS-A353990

a(1) = 1; for n > 1, a(n) is the smallest positive number that has not yet appeared that is coprime to a(n-1), does not equal a(n-1)+1, and whose binary expansion has no 1-bit in common with the binary expansion of a(n-1).

In the first 100000 terms the fixed points are 1, 3, 5, 12, 21, 26, 44, 49, 227, 3488, 5890, 9067, 9310, 37625, 74702, although it is likely more exist. In the same range the lowest unseen number is 30686; the sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015686 | OEIS-A354062

a(n) = Li(-2^n, 1/3), where Li(n, z) is the polylogarithm function.

Conjecture: for k >= 0, a(n) is divisible by 2^2^k+1 = A000215(k) for all n >= 2^max{k,1}. - _Jianing Song_, May 17 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, May 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015687 | OEIS-A354072

Perfect numbers that are the sum of the divisors of some number.

Conjecture: 8128 is the only perfect number that is not in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, May 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015688 | OEIS-A354073

Multiply-perfect numbers that are the sum of the divisors of some number.

Conjecture: 8128 is only multiply-perfect number that is not in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, May 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015689 | OEIS-A354074

Factorials that are the sum of the divisors of some m.

Conjecture: number 2 is the only factorial that is not in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, May 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015690 | OEIS-A354075

Lexicographically earliest infinite sequence of distinct positive integers such that A(a(n+1)) is prime to A(a(n)) but not to A(a(n-1)), where A is A001414.

Conjectured to be a permutation of N/{0,1}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jun 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015691 | OEIS-A354087

a(1) = 1; for n > 1, a(n) is the smallest positive number that has not yet appeared that shares a factor with a(n-1) and whose binary expansion has a single 1-bit in common with the binary expansion of a(n-1).

In the first 100000 terms the fixed points are 1, 3, 30, 38, 350, 1603, 1936, 10176, 11976, 46123, 58471, 89870, although it is likely more exist. In the same range the lowest unseen number is 1019; the sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 17 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015692 | OEIS-A354124

G.f. A(x) satisfies: 1 = Sum_{n>=0} x^n * (1 + x^n*A(x))^n / (1 + x^(n+1)*A(x))^(n+1).

sign(a(7*n+k)) = (-1)^n for k=1..7, n > 1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 17 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015693 | OEIS-A354139

a(n) is the least positive integer m such that (k+1)^n + (k+2)^n + ... + (k+m)^n == 0 (mod n) for every positive integer k.

Conjecture: a(n) = A007947(n) * A193267(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimitrios T. Tambakos_, May 18 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015694 | OEIS-A354150

Index of 2^n in A090252, or -1 if 2^n does not appear.

It is conjectured (see A090252) that the indices of even terms in A090252 are {3*2^i-1, i >= 0} (A083329), so the positive terms of the present sequence should be a subsequence of A083329.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015695 | OEIS-A354152

a(n) is the number of permutations pi in S_n such that pi(i) - i != 1 (mod n) and pi(i) - i != -1 (mod n) for all i.

Conjecture: a(2k) = A000179(2k) + 2 for k > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, May 27 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015696 | OEIS-A354159

Terms 2*p (p prime) in A090252, divided by 2, in order of appearance.

Conjecture: the indices of the terms 2*p in A090252 are terms in A083329.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 30 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015697 | OEIS-A354242

Expansion of e.g.f. 1/sqrt(5 - 4 * exp(x)).

Conjecture: Let k be a positive integer. The sequence obtained by reducing a(n) modulo k is eventually periodic with the period dividing phi(k) = A000010(k). For example, modulo 16 we obtain the sequence [1, 2, 14, 14, 6, 14, 6, 14, 6, ...], with an apparent period of 2 beginning at a(3). Cf. A354253. More generally, we conjecture that the same property holds for integer sequences having an e.g.f. of the form G(exp(x) - 1), where G(x) is an integral power series. (End) Conjectural o.g.f. as a continued fraction of Stieltjes type: 1/(1 - 2*x/(1 - 5*x/(1 - 6*x/(1 - 10*x/(1 - 10*x/(1 - 15*x/(1 - ... - (4*n-2)*x/(1 - 5*n*x/(1 - ...)))))))). - _Peter Bala_, Jul 07 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, May 20 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015698 | OEIS-A354246

Indices of coefficients of x^(2*k-1) in Integral exp(-x*tan(x))/cos(x) dx at which the signs of the coefficients change: list of k such that sign(A354245(k)) != sign(A354245(k-1)), starting with 1.

Conjecture: lim_{n->oo} a(n)/n^2 = Pi^2/8 = A111003 = 1.2337... - _Vaclav Kotesovec_, May 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 20 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015699 | OEIS-A354253

Expansion of e.g.f. 1/sqrt(9 - 8 * exp(x)).

Conjecture: Let k be a positive integer. The sequence obtained by reducing a(n) modulo k is eventually periodic with the period dividing phi(k) = A000010(k). For example, modulo 7 we obtain the sequence [1, 4, 3, 2, 3, 0, 0, 4, 3, 2, 3, 0, 0, 4, 3, 2, 3, 0, 0, ...], with a preperiod of length 1 and an apparent period thereafter of 6 = phi(7). Cf. A354242. - _Peter Bala_, Jul 08 2022 Conjectural o.g.f. as a continued fraction of Stieltjes type: 1/(1 - 4*x/(1 - 9*x/(1 - 12*x/(1 - 18*x/(1 - 20*x/(1 - 27*x/(1 - ... - (8*n-4)*x/(1 - 9*n*x/(1 - ...)))))))). - _Peter Bala_, Jul 08 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, May 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015700 | OEIS-A354255

Even numbers in A090252 in order of appearance.

Conjecture: The indices of even numbers in A090252 are precisely the numbers {A083329(n), n >= 1}. See A090252 for discussion. - _N. J. A. Sloane_, May 22 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael S. Branicky_, May 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015701 | OEIS-A354385

a(n) is the smallest odd number that has n middle divisors.

Conjecture: For every k, there is an n such that all >= a(n) are divisible by the first k odd primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, May 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015702 | OEIS-A354387

a(n) is the number of arch configuration solutions with n arches derived from 2 concentric arches using the exterior arch splitting algorithm.

Conjecture from _Roger Ford_, Aug 26 2022: (Start) Conjecture: For n >= 2, a(n) = Sum_{k = 2..floor((n+2)/2)} A339179 T(n,k)*(k-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger Ford_, May 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015703 | OEIS-A354399

List of k such that sign(A009273(k)) = sign(A009273(k+1)).

Conjecture: lim_{n->oo} a(n)/n^2 = Pi^2/8 = A111003 = 1.2337...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, May 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015704 | OEIS-A354408

Triangle read by rows of generalized ménage numbers: T(n,k) is the number of permutations pi in S_n such that pi(i) != i and pi(i) != i+k (mod n) for all i; n, 1 <= k < n.

Conjectures: (Start) Conjecture: T(n,j) < T(n,k) if gcd(n,j) < gcd(n,k) and (n,k) != (6,3). - _Pontus von Brömssen_, May 31 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, May 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015705 | OEIS-A354409

Maximum value in the n-th row of A354408.

Conjecture: a(n) = A000179(n) if and only if n is prime. Conjecture: a(n) = A354408(n, A032742(n)) for n != 6. - _Pontus von Brömssen_, May 31 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, May 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015706 | OEIS-A354414

a(n) is the smallest positive integer which does not occur in any Lucas sequence in which the first term is at most n and the second term is at most the first term.

Conjecture: a(n) = (n+1)*F(k+2) - F(k)*F(k+1) where F(k) are the Fibonacci numbers such that F(k) <= n < F(k+1). - _Muthu Veerappan Ramalingam_, Jun 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, May 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015707 | OEIS-A354420

Lexicographically earliest infinite sequence of distinct positive numbers such that, for n>3, a(n) has a common factor with a(n-2), shares a 1-bit in its binary expansion with a(n-2), has no common factor with a(n-1), and does not share a 1-bit in its binary expansion with a(n-1).

The sequence is similar to A098550 but with the addition restrictions that each new term a(n) must share a 1-bit in its binary expansion with a(n-2), while sharing no 1-bits with the binary expansion of a(n-1). Unlike A351691 no additional restrictions on the factors or 1-bits of a(n) are required for the sequence to be infinite. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015708 | OEIS-A354434

a(1) = 1; for n > 1, a(n) is the smallest unused square spiral number such that a(n) shares a factor with all existing numbers in its Moore neighborhood.

The sequence is conjectured to be a permutation of the positive integers, although it takes many terms for most primes to appear, e.g. a(1807) = 13, a(35156) = 179. The primes do not occur in their natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 28 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015709 | OEIS-A354478

a(n) is the numerator of Sum_{k=1..n} 1 / Stirling1(n,k).

Conjecture: a(n)/A354479(n) tends to 1 as n tends to infinity. For comparison: A112290(n)/A112291(n) tends to 2 as n tends to infinity. - _Vaclav Kotesovec_, Jun 02 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Jun 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015710 | OEIS-A354499

Number of consecutive primes generated by adding 2n to the odd squares (A016754).

Conjecture: a(n) <= 18 = a(326). I conjecture the opposite: a(n) is unbounded, and indeed for any k < 1 and any m there are >> x^k terms up to x with a(n) > m. At a very rough guess, there should be some n with 20-50 digits having a(n) > 18. - _Charles R Greathouse IV_, Oct 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven M. Altschuld_, Aug 15 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015711 | OEIS-A354516

Smallest k such that m - gpf(m) = k has exactly n solutions m >= 2, gpf = A006530; or -1 if no such k exists.

Conjectures (if no term equals -1): (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015712 | OEIS-A354528

Square array T(m,n) read by antidiagonals - see Comments for definition.

Conjecture: T(m, n) = M(P_m X P_n) where P_m X P_n is the grid graph of size m X n. The conjecture is proved if m = 1 or n = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sela Fried_, Aug 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015713 | OEIS-A354529

a(1) = 3, a(2) = 12 and a(n) = (3n^2+8n-2)/2 if n is even or = (3n^2+8n-5)/2, if n is odd, for n >= 3.

Conjecture: a(n) = M(P_3 X P_n) where P_3 X P_n is the grid graph of size 3 X n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sela Fried_, Aug 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015714 | OEIS-A354532

Numbers k that are not Mersenne exponents (A000043) such that 2*(2^k-1) is in A354525.

Conjecture: all terms beyond a(2) = 9 are primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015715 | OEIS-A354538

a(n) is the least k such that A322523(k) = n.

For n >= 0, the contents of the n-th box are (conjecture): [For the correctness and the formula and the conjecture see A322523. - _Jianing Song_, Oct 17 2022]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugh Williamson_, Aug 17 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015716 | OEIS-A354568

Irregular triangle read by rows: T(n,k) is the number of Hamiltonian cycles in the Kneser graph K(n,k), 1 <= k < n/2.

It has been conjectured that T(n,k) > 0 (i.e., that K(n,k) is Hamiltonian) for all 1 <= k < n/2, with the exception (n,k) = (5,2) (the Petersen graph). See Lovász 1970 for a more general conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pontus von Brömssen_, Aug 18 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015717 | OEIS-A354569

Ordered even leg lengths k (listed with multiplicity) of primitive Pythagorean triangles such that all odd prime factors of k are congruent to 1 (mod 4) and at least one prime factor is congruent to 1 (mod 4).

Conjecture: lim_{n->oo} a(n)/n -> 2*Pi.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lothar Selle_, Jun 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015718 | OEIS-A354570

Ordered odd leg lengths k (listed with multiplicity) of primitive Pythagorean triangles such that all prime factors of k are congruent to 3 (mod 4).

Conjecture: lim_{n->oo} a(n)/n = Pi. Also, lim_{n->oo} A354571(n)/n = Pi.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lothar Selle_, Jun 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015719 | OEIS-A354571

Ordered even leg lengths k (listed with multiplicity) of primitive Pythagorean triangles such that all odd prime factors of k are congruent to 3 (mod 4) and at least one prime factor is odd.

Conjecture: lim_{n->oo} a(n)/n = Pi. Also, lim_{n->oo} A354570(n)/n = Pi.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lothar Selle_, Jun 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015720 | OEIS-A354575

a(1) = 1; for n > 1, a(n) is the smallest positive number that has not yet appeared that is coprime to a(n-1) and the difference a(n) - a(n-1) is distinct from all previous differences.

In the first 200000 terms the fixed points are 1,2,8,35, and it is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015721 | OEIS-A354651

G.f. A(x) satisfies: 1/(1 - x) = Sum_{n>=1} (-1)^(n-1) * A(x)^(n^2).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 18 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015722 | OEIS-A354672

Numbers x with property that x is not the smallest possible value in the Pellian equation x^2 - D*y^2 = 1 with D = squarefree part of (x^2 - 1).

Conjecture: this sequence is equivalent to the sorted distinct values of cos(m*arccos(k)), where m and k are integers greater than 1. - _Jennifer Buckley_, Apr 23 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Herbert Kociemba_, Jun 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015723 | OEIS-A354687

a(1) = 1; for n > 1, a(n) is the smallest positive number that has not yet appeared that shares a factor with a(n-1) and the difference | a(n) - a(n-1) | is distinct from all previous differences.

In the first 200000 terms the fixed points are 1,2,6,10,68. It is plausible no more exist although this is unknown. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015724 | OEIS-A354717

Lexicographically earliest infinite sequence of distinct positive integers such that in any run of four consecutive terms there is one term which is prime to the other three, none of which are pairwise coprime.

Open question: Is it true that in any run of four consecutive terms there is always a prime or prime power (this being the term prime to the other three)? Conjecture: Sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jun 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015725 | OEIS-A354732

Lexicographically earliest infinite sequence of distinct positive integers such that in any run of four consecutive terms there is only one pair of terms which share a prime divisor, the rest are all pairwise coprime.

Conjecture: Sequence is a permutation of the positive integers in which the primes appear in their natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jun 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015726 | OEIS-A354749

a(1) = 1; for n > 1, a(n) is the smallest positive number greater than 1 that is coprime to a(n-1) and the product a(n) * a(n-1) is distinct from all previous products a(i) * a(i-1), i=2..n-1.

This is a variation of A354803 where all terms beyond the first must be greater than 1. In the first 1000000 terms the fixed points are 1,2,3,4,5,7, it is likely no more exist, while the smallest number not to have appeared is 2090. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 06 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015727 | OEIS-A354753

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number that shares a factor with a(n-1) and the product a(n) * a(n-1) is distinct from all previous products a(i) * a(i-1), i=2..n-1.

This sequence uses a similar rule to A088177 but here all neighboring terms also share a factor. In the first 500000 terms the fixed points are 1,2,4,6, it is likely no more exist, while the smallest number not to have appeared is 1153. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 06 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015728 | OEIS-A354766

1/4 of the total number of integral quadruples with sum = n and sum of squares = n^2.

Conjectures from _Colin Mallows_, Jun 12 2022: (Start) Conjecture 1: (Based on the data for n <= 5000) pq/4 and tq/4 are multiplicative sequences. Conjecture 2: When n = p^k, p prime and k >= 1: Conjecture: the numbers n for which a(n) = n have a positive asymptotic density.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 19 2022, based on an email from _Colin Mallows_, Jun 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015729 | OEIS-A354767

Indices of terms in A354169 that have Hamming weight 1.

Conjecture from _N. J. A. Sloane_, Jun 30 2022, modified Jul 30 2022: (Start) With b(n) defined in this way, we conjecture that a(n) = b(n) + n. This has been checked for the first 3296 terms. The conjecture is now known to be true. - _N. J. A. Sloane_, Aug 29 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015730 | OEIS-A354773

For terms of A354169 that are the sum of two distinct powers of 2, the exponent of the smaller power of 2.

Conjecture from _N. J. A. Sloane_, Jun 29 2022: (Start) The following is a conjectured recurrence for a(n). Basically a(n) = a(n/2-1) if n is even, and a(n) = (n+1)/2 if n is odd, except that there are four types of n which have a different formula, and there are 19 exceptional values for small n. Note that a(n) does not depend on earlier values when n is odd. The conjecture is now known to be true. See De Vlieger et al. (2022). - _N. J. A. Sloane_, Aug 29 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015731 | OEIS-A354775

Indices where A354169 is the sum of two consecutive powers of 2.

The first differences begin 4, 16, 12, 12, 18, 6, 18, 6, 48, 48, 96, 96, 192, 192, 384, 384, 768, 768, 1536, ..., which suggests that from the 9th term on, the differences have g.f. 48*(1+x)/(1-2*x^2), with an analogous conjecture for the sequence itself.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015732 | OEIS-A354778

Number of integer quadruples (u,v,w,x) such that u^2+v^2+w^2+x^2 = n^2 and u+v+w+x = n.

See A278085 and A354766 for some conjectural formulas.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 27 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015733 | OEIS-A354794

Triangle read by rows. The Bell transform of the sequence {m^m | m >= 0}.

Conjecture: T(n, k) = Sum_{i=0..n-k} A048994(n-k, i) * A048993(n+i-1, n-1) for 0 < k <= n and T(n, 0) = 0^n for n >= 0; proved by Mike Earnest, see link at A354797. (End) Conjectures from _Mikhail Kurkov_, Apr 20 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jun 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015734 | OEIS-A354797

Triangle read by rows. T(n, k) = |Stirling1(n, k)| * Stirling2(n + k, n) = A132393(n, k) * A048993(n + k, n).

Sum_{k=0..n} (-1)^(n - k)*T(n, k) = n^n. - _Werner Schulte_, Jun 03 2022 in A000312. [Formerly a conjecture, now proved by Mike Earnest, see link.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jun 06 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015735 | OEIS-A354798

Indices of terms in A354169 that are not powers of 2.

Conjecture from _N. J. A. Sloane_, Jul 15 2022: (Start) The following is a conjectured explicit formula for a(n). Then we conjecture that a(n) = V(i) + f(j) where f(0) .. f(3) are 0,2,4,8, and for j >= 4, f(j) = 3*j if j is even, f(j) = 3*j-1 if j is odd. The conjecture has been checked for the first 5000 terms. The conjecture is now known to be true. See De Vlieger et al. (2022). - _N. J. A. Sloane_, Aug 29 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_ and _N. J. A. Sloane_, Jun 06 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015736 | OEIS-A354803

a(1) = 1; for n > 1, a(n) is the smallest positive number that is coprime to a(n-1) and the product a(n) * a(n-1) is distinct from all previous products a(i) * a(i-1), i=2..n-1.

This sequences uses similar a similar rule to A088177 but here all neighboring terms are coprime. In the first 1000000 terms the only fixed point is the first term while the smallest number not to have appeared is 2054. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015737 | OEIS-A354846

a(n) is the first composite k such that exactly n primes are the sum of all but one of the numbers from 1 to k-1 that are coprime to k, or -1 if there is no such k.

Conjecture: for every composite k there is at least one such prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Jun 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015738 | OEIS-A354852

a(n) is the number of distinct ways to represent n using any mixed radix numeral system.

Conjecture: a(p) = a(p-1)+1 whenever p is a prime number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max J. Folkers_, Jun 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015739 | OEIS-A354896

A fixed point of the two-block substitution 00->001, 01->010, 10->101, 11->110.

Some more conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Dekking_, Jun 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015740 | OEIS-A354903

Lexicographically earliest infinite sequence of distinct positive integers such that the number of divisors of a(n+1) is prime to a(n).

Conjectured to be a permutation of the positive integers in which primes appear in natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jun 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015741 | OEIS-A354929

Numbers of the form 2*p^e, with p an odd prime and e >= 2.

Conjecturally numbers k > 1 such that A047994(k) = A344005(k) (see A354928), and k is in A265128. See comments in A346608.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jun 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015742 | OEIS-A354950

The number of squarefree numbers whose largest prime divisor is prime(n) and that are averages of twin prime pairs.

Conjecture: Limit_{n->oo} log(a(n))/(n*log(n)) = c ~ 0.13... .

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Jun 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015743 | OEIS-A354960

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number not occurring earlier that is a multiple of the number of proper divisors of a(n-1).

The sequence is conjectured to be a permutation of the positive integers although it may take an extremely large number of terms for the primes to appear; e.g., 263 has not occurred after 500000 terms. Also although the vast majority of primes will appear in their natural order, some may not; e.g., a(455) = 840, which has 31 proper divisors, so a(456) = 31, and then a(457) = 29.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jul 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015744 | OEIS-A354971

a(1)=1, a(2)=0; for n > 2, a(n) is the number of times a(n - 1 - a(n-1)) has appeared in the sequence.

Conjecture: every positive integer eventually appears in the sequence. Conjecture: The longest run of same, consecutive a(n) is 4 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Neal Gersh Tolunsky_, Jun 14 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015745 | OEIS-A355007

Triangle read by rows. T(n, k) = n^k * |Stirling1(n, k)|.

Conjecture: T(n,k) = A056856(n,k)*n. - _R. J. Mathar_, Mar 31 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jun 17 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015746 | OEIS-A355008

Triangle T(n, k), n > 0, k = 1..n, read by rows; each row contains the smallest distinct positive integers (in ascending order) such that a term of row n times a term of row n+1 does not appear in row n+2.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jun 15 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015747 | OEIS-A355039

Carmichael numbers whose number of prime factors is prime.

Wright shows that this sequence is infinite on the assumption of Heath-Brown's conjecture on the first prime in an arithmetic progression. - _Charles R Greathouse IV_, Aug 05 2022, corrected by _Amiram Eldar_, Mar 25 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Jun 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015748 | OEIS-A355041

Numbers k such that A152763(2^k) < A152763(2^k-1).

Conjecture: there is no prime in this sequence. Among the primes p <= 257, the prime p for which A152763(2^p)/A152763(2^p-1) is the smallest is p = 193, where A152763(2^p)/A152763(2^p-1) = 143/140.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Jun 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015749 | OEIS-A355072

a(0) = 0, a(1) = 1; for n > 1, a(n) is the smallest positive number whose sum a(n) + a(n-1) is distinct from all previous sums, a(i) + a(i-1), i=1..n-1, whose product a(n) * a(n-1) is distinct from all previous products, a(i) * a(i-1), i=1..n-1, and whose difference |a(n) - a(n-1)| is distinct from all previous differences, |a(i) - a(i-1)|, i=1..n-1.

For n up to ~35000 the vast majority of terms are concentrated along three lines, the lowest being near the x-axes; see the first linked image. In this same range there are many terms equal to 1; see A355135. Beyond this range the terms no longer fall along the upper-most line and the number of terms equal to 1 greatly diminishes. The reason for this change in behavior is unknown. The remaining upper-most line has a gradient close to 1 and contains multiple fixed points; see A355136 and the second linked image. The sequence it conjectured to contain all the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 18 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015750 | OEIS-A355076

a(n) is the denominator of Sum_{k = 0..n} fusc(k)/fusc(k+1) (where fusc is Stern's diatomic series A002487).

Conjecture: a(n) = 1 for n of the form 2*4^k - 1 or 2*4^k - 2 for some k >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jun 18 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015751 | OEIS-A355118

The number of nonequivalent root ancestral configurations in a recursively defined family of gene trees and species trees with at least n = 9 leaves, in which for n >= 12 leaves, 3-leaf trees are successively added at the root of the tree with n-3 leaves.

For 9 <= n <= 20, a(n) tabulates the number of nonequivalent root ancestral configurations for the pair (G, S) with the largest number of nonequivalent root ancestral configurations (among pairs with n leaves and G = S). Disanto and Rosenberg (2019) conjecture that a(n) provides this maximum for all n >= 9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Noah A Rosenberg_, Jun 19 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015752 | OEIS-A355144

Number T(n,k) of partitions of [n] having exactly k blocks of size at least three; triangle T(n,k), n>=0, 0<=k<=floor(n/3), read by rows.

Conjecture: Each row are coefficients of a polynomial with only real zeros. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jun 20 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015753 | OEIS-A355148

Numbers that are the concatenation of two palindromes and that have exactly two palindromic factors, all with the same number of decimal digits.

Conjecture: these are an infinite number of such terms. A subsequence of this sequence is {s(k)} where s(k) = (202/10989)*t(k)*u(k), t(k) = 10^(6*k + 3) - 1 and u(k) = 2099*10^(6*k + 1) + 988. s(k) can be decomposed in 2 different ways: the first is (202/333)*t(k) and (1/33)*u(k); the second is (101/111)*t(k) and (2/99)*u(k). And since {s(k)} is an infinite sequence, its existence proves _Chai Wah Wu_'s conjecture to be true. - _Nicolas Bělohoubek_, May 20 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nicolas Bělohoubek_, Jun 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015754 | OEIS-A355149

Partial sums of A351743.

Conjecture: a(n + 42) = 3645*a(n) + b(n) for n >= 138 (where b is 42-periodic).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, _Rémy Sigrist_ and _David James Sycamore_, Jun 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015755 | OEIS-A355150

The Hamming weight of A354169, a(n) = A000120(A354169(n)).

All the following conjectures are now known to be true. See De Vlieger et al. (2022). - _N. J. A. Sloane_, Aug 29 2022 Conjecture: It appears that this sequence may be computed by a fast algorithm: This conjecture was verified for n = 0..2^16 against the b-file provided by _Michael De Vlieger_. - _Thomas Scheuerle_, Jul 14 2022 Conjecture: It appears that this sequence is almost a periodic sequence, with period 6. Let x be the sequence defined below.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Jun 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015756 | OEIS-A355151

G.f. A(x) satisfies: (1 - x)/(1 + x) = Sum_{n=-oo..+oo} (-1)^n * A(x)^(n^2).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015757 | OEIS-A355175

Determinant of the n X n matrix [(i-j)^2 + d(i,j)]_{1<=i,j<=n}, where d(i,j) is 1 or 0 according as i = j or not.

Conjecture: a(n) = (n^2-4)*(n^2+2*n+3)*(n^5-2*n^4-n^3-28*n^2+60*n-90)/1080. The conjecture is true (see Fried link). - _Sela Fried_, Feb 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 28 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015758 | OEIS-A355178

Decimal expansion of 2^(-2/3)/L, where L is the conjectured Landau's constant A081760.

Decimal expansion of 2^(-2/3)/L, where L is the conjectured Landau's constant A081760.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Sep 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015759 | OEIS-A355187

Number of Collatz trajectories (A070165) for all positive integers <= 10^n that contain 2^4 as the greatest power of 2 within its trajectory.

It is conjectured that lim_{n->infinity} a(n)/10^n = 15/16. Empirically, 93.75% of all trajectories have 2^4 as the greatest power of 2 within its trajectory. Sequence A135282(n) is the maximum power of 2 reached in the Collatz trajectory for integer n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Jun 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015760 | OEIS-A355239

Starting values k > 4 of a Collatz iteration reaching either k-1 or k+1.

No further terms up to 2*10^9. It is conjectured that this is the full list of starting values of Collatz trajectories reaching k-1 or k+1, and that the number of steps until this happens is one of the 8 terms of A355240.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jul 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015761 | OEIS-A355240

Numbers of steps until the Collatz iteration started at k > 4 returns to either k-1 or k+1.

It is conjectured that no further terms exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jul 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015762 | OEIS-A355259

Triangle read by rows. Row k are the coefficients of the polynomials (sorted by ascending powers) which interpolate the points (n, A355257(n, k+1)) for n = 0..k.

Conjecture from _Mélika Tebni_: These polynomials generate column k + 1 of

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jul 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015763 | OEIS-A355269

Lexicographically earliest infinite sequence of distinct positive integers such that a(n+1) is prime to the number of divisors of a(n).

Conjectures: When a(n) is square a(n+1) is even; no two squares of equal parity can appear as adjacent terms; sequence is a permutation of the positive integers with primes in natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_ and _Michael De Vlieger_, Jun 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015764 | OEIS-A355323

Numbers k such that A007063(k) = A356026(k).

It is conjectured that this is an infinite sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015765 | OEIS-A355326

Determinant of the n X n matrix [(i-j)^3+d(i,j)]_{1<=i,j<=n}, where d(i,j) is 1 or 0 according as i = j or not.

Conjecture 1: a(n) = 1 + P(n^2)*n^2*(n^2-1)/672000, where P(n) = n^6 - 19*n^5 + 123*n^4 - 337*n^3 + 12376*n^2 - 44144*n + 40000. Conjecture 2: For any positive integers m and n, the determinant of the matrix [(i-j)^m+d(i,j)]_{1<=i,j<=n} has the form 1 + n^2*(n^2-1)*P(n), where P(n) is a polynomial in n with rational number coefficients whose degree is (m+1)^2-4. The conjectures are true (see Fried link). - _Sela Fried_, Feb 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jun 28 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015766 | OEIS-A355330

Numbers k such that A020696(2^k-1) < A020696(2^k+1).

Sándor (2021) showed that all the Mersenne exponents (A000043) are in this sequence and conjectured that both this sequence and its complement are infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Jun 29 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015767 | OEIS-A355339

a(n) is the number of central even terms in the n-th row of triangle A355320; a(0) = 0.

Conjecture: a(n) = 1 iff n > 0 and n belongs to A000918.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jun 29 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015768 | OEIS-A355350

G.f. A(x,y) satisfies: x*y = Sum_{n=-oo..+oo} (-1)^n * x^(n*(n+1)/2) * A(x,y)^n, with coefficients T(n,k) of x^n*y^k in A(x,y) given as a triangle read by rows.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 29 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015769 | OEIS-A355360

G.f. A(x,y) satisfies: x*y*A(x,y) = Sum_{n=-oo..+oo} (-1)^n * x^(n*(n+1)/2) * A(x,y)^n, with coefficients T(n,k) of x^n*y^k in A(x,y) given as a triangle read by rows.

Conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 19 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015770 | OEIS-A355372

Expansion of the e.g.f. log((1 - x) / (1 - 2*x)) / (1 - x)^3.

Conjecture: For p prime, a(p) == -1 (mod p). The conjecture is true (see Fried link). - _Sela Fried_, Dec 17 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mélika Tebni_, Jun 30 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015771 | OEIS-A355374

a(1) = 1; for n > 1, a(n) is the smallest positive number that has not yet appeared such that the number of proper divisors of a(n) equals the number of 1-bits in the binary expansion of a(n-1).

In the first 700 terms the fixed points are 1, 2, 3, 4, 5, 16, 21, 22, 35, 48, 168, 412, 428. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 30 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015772 | OEIS-A355406

Positive integers that are not powers of 2 and whose Collatz trajectory has maximum power of 2 different from 2^4.

It is conjectured that 15/16 (93.75%) of the positive integers that are not powers of 2 have 2^4 as the maximum power of 2 in their Collatz trajectory (see A232503 and A355187). {a(n)} lists the remaining positive integers. Consequently, it is conjectured that this sequence will have lim_{n->oo} a(n)/n = 1/16.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Jul 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015773 | OEIS-A355407

Expansion of the e.g.f. log((1 - x) / (1 - 2*x)) / (1 - x)^4.

Conjecture: For p prime, a(p) == -1 (mod p).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mélika Tebni_, Jul 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015774 | OEIS-A355409

Expansion of e.g.f. 1/(1 + exp(2*x) - exp(3*x)).

Conjecture: Let k be a positive integer. The sequence obtained by reducing a(n) modulo k is eventually periodic with period dividing phi(k) = A000010(k). For example, modulo 9 we obtain the sequence [1, 1, 7, 1, 4, 1, 1, 1, 7, 1, 4, 1, 1, 1, 7, 1, 4, 1, 1, ...] with an apparent period of 6 = phi(9) beginning at a(1). Cf. A354242. - _Peter Bala_, Apr 16 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Jul 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015775 | OEIS-A355414

Expansion of the e.g.f. log((1 - x) / (1 - 2*x)) / (1 - x)^5.

Conjecture: For p prime, a(p) == -1 (mod p).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mélika Tebni_, Jul 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015776 | OEIS-A355424

Positive integers m such that the real quadratic fields of the form Q(sqrt(m^2+4)) have class number 1.

Former Yokoi's conjecture, proved by Biró in 2003 (see References). There are only six real quadratic fields of the form Q(sqrt(a(n)^2+4)), where Q indicates the set of rational numbers, with class number one.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Jul 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015777 | OEIS-A355448

a(n) = 1 if the number of divisors of n^2 is coprime to 6, otherwise 0.

a(n) = abs(A307424(n)). [conjectured]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jul 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015778 | OEIS-A355461

Squarefree numbers d of the form r^2*m^2 + 4*r, where r and m are odd positive integers, such that Q(sqrt(d)) has class number 1.

In 1801, Gauss conjectured that there exist infinitely many real quadratic fields with class number one and the conjecture is still unproved, but there are only 12 real quadratic fields of class number one which are of the form Q(sqrt(r^2*m^2 + 4*r)), where the parameters r and m are odd integers. Those 12 values of d := r^2*m^2 + 4*r belong to the present sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Jul 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015779 | OEIS-A355476

a(1)=1. For a(n) a novel term, a(n+1) = A000005(a(n)). For a(n) seen already k > 1 times, a(n+1) = k*a(n).

Conjectures: (i) Every integer > 1 appears infinitely many times. (ii) The first occurrences of primes appear in natural order, starting with 2 as described above, and continuing with odd primes a(n) = p, following a(n-1) = 2^(p-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jul 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015780 | OEIS-A355518

Primes that cannot be represented as 2*p - q where p, q and 2*p^2 - q^2 are prime.

Conjecture: there are no other terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Jul 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015781 | OEIS-A355521

Primes that cannot be represented as 2*p+q where p, q and (2*p^2+q^2)/3 are prime.

Conjecture: there are no other terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Jul 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015782 | OEIS-A355585

T(j,k) are the numerators s in the representation R = s/t + (2*sqrt(3)/Pi)*u/v of the resistance between two nodes separated by the distance (j,k) in an infinite triangular lattice of one-ohm resistors, where T(j,k), j >= 0, 0 <= k <= floor(j/2) is an irregular triangle read by rows.

T(n,0)/A355586(n,0) = T(n-1,0)/A355586(n-1,0) + A084768(n-1)/3 for n>=1 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jul 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015783 | OEIS-A355611

a(0) = 0; for n > 0, a(n) is the smallest positive number not occurring earlier such that the binary string of |a(n) - a(n-1)| does not appear in the binary string concatenation of a(0)..a(n-1).

The sequence is conjectured to be a permutation of the positive integers. In the first 200000 terms the only fixed points are 1199 and 14767. It is unknown if more exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015784 | OEIS-A355702

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number that has not yet appeared that has the same number of prime divisors as the sum a(n-2) + a(n-1).

In the first 500000 terms on seventeen occasions the sum of the previous two terms equals the next term, these terms being 3, 5, 8, 11, 100,... ,131072, 262144. It in unknown if there are infinitely many such terms. In the same range there are seventy-three fixed points; see A356017. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jul 14 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015785 | OEIS-A355704

Indices k of partition function p where p(k) and p(k) + 2 are twin primes.

Because asymptotically size of partitions number function p(n) ~ O(exp(sqrt(n))), and probability of primality of p(n) ~ O(1/sqrt(n)) and combined probability of primality of p(n) and p(n)+-2 is ~ O(1/n), the sum of the prime probabilities is diverging and there are no obvious restrictions on primality; therefore, this sequence may be conjectured to be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Serge Batalov_, Jul 14 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015786 | OEIS-A355706

Indices k of partition function p where p(k) is a twin prime.

Because asymptotically the size of the partition number function p(n) is ~ O(exp(sqrt(n))), and the probability of primality of p(n) is ~ O(1/sqrt(n)) and the combined probability of primality of p(n) and p(n)+-2 is ~ O(1/n), the sum of the prime probabilities is diverging and there are no obvious restrictions on primality; therefore, this sequence may be conjectured to be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Serge Batalov_, Jul 15 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015787 | OEIS-A355728

Indices k of partition function where consecutive p(k) and p(k+1) are prime.

Because asymptotically the size of the partition number function p(n) is ~ O(exp(sqrt(n))), and the probability of primality of p(n) is ~ O(1/sqrt(n)) and the combined probability of primality of p(n) and p(n+1) is ~ O(1/n), the sum of the prime probabilities is diverging and there are no obvious restrictions on primality; therefore, this sequence may be conjectured to be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Serge Batalov_, Jul 15 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015788 | OEIS-A355760

a(n) is the number of grid points in a square lattice covered by the area enclosed by n loops of an Archimedean spiral with starting point (0, 0) and endpoint (n, 0).

Conjecture: Only lattice points on the positive x-axis are on the outer boundary of the spiral. It seems that the spiral passes all other grid points without hitting any.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Karl-Heinz Hofmann_, Jul 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015789 | OEIS-A355814

Smallest value t such that 1/s^2 + 1/t^2 = 1/p^2 + 1/q^2 has exactly n solutions (p,q) where p,q < t; or -1 if no such t exists.

Conjecture: a(n) is divisible by 35 for odd n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Jul 18 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015790 | OEIS-A355872

G.f. A(x) satisfies: x = Sum_{n=-oo..+oo} (-x)^(n^2) * A(x)^((n-1)^2).

Conjecture: a(n) == 2 (mod 4) for n >= 1. Conjecture: a(2*n-1) == 2 (mod 8) for n >= 1. Conjecture: a(2*n) == 6 (mod 8) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015791 | OEIS-A355878

Smallest p == 1 (mod 4) such that Q(sqrt(p)) has class number 2n+1.

Conjecture: a(n) > A002148(n) for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Jul 20 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015792 | OEIS-A355889

Concatenate the exponents of the powers of 2 in A354169(k) in increasing order, for k = 1, 2, 3, ...

It is conjectured that the Hamming weight of A354169(k) is always 0, 1, or 2. This is known to be true for at least the first 2^25 terms. (The present sequence is well-defined even if the conjecture is false.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_ and _N. J. A. Sloane_, Jul 20 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015793 | OEIS-A355895

Even terms in A354790, divided by 2, in order of appearance.

The even terms in A354790 appear at indices 2, 7, 15, 31, 63, 127, 255, 511, 1023, 2047, 4095, 8191, 16383, 32767, 65535, ... and conjecturally at indices 2^k-1 for all k >= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 27 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015794 | OEIS-A355896

Multiples of 3 in A354790, divided by 3, in order of appearance.

The multiples of 3 in A354790 appear at indices 3, 7, 17, 35, 71, 143, 287, 575, 1151, ... and conjecturally at indices 9*2^k-1 for all k >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 27 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015795 | OEIS-A355898

a(1) = a(2) = 1; a(n) = gcd(a(n-1), a(n-2)) + (a(n-1) + a(n-2))/gcd(a(n-1), a(n-2)).

Conjecture: For n >= 3775 a(n) can also be expressed in the following three ways: Adamczewski disproves the above conjecture in linked article. - _Michael De Vlieger_, Aug 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015796 | OEIS-A355920

Largest prime number p such that x^n + y^n mod p does not take all values on Z/pZ.

Conjecture: a(n) == 1 (mod n). - _Jason Yuen_, May 18 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Azuma_, Jul 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015797 | OEIS-A355956

Index k of partition function p such that p(k) is a member of a cousin prime pair.

Because asymptotically the size of the partition number function p(n) ~ O(exp(sqrt(n))), and the probability of primality of p(n) ~ O(1/sqrt(n)) and the combined probability of primality of p(n) and p(n)+-4 is ~ O(1/n), the sum of the prime probabilities is diverging and there are no obvious restrictions on primality; therefore this sequence may be conjectured to be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Serge Batalov_, Jul 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015798 | OEIS-A355987

a(n) = n! * Sum_{k=1..n} 1/floor(n/k)!.

Conjecture: c = Sum_{k>=1} 1/((k+1)!*k) = 2 - exp(1) - A001620 + A091725. - _Vaclav Kotesovec_, Sep 24 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Jul 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015799 | OEIS-A355997

Triangle read by rows: T(n,d) is the number of fixed orthoplex n-ominoes with cell centers determining d-space.

Conjecture: T(n,n-4) = 2^(n-7) * n^(n-9) * (n-4) * (n-5) * (n-6) * (n^6-14*n^5+65*n^4-189*n^3+594*n^2-1305*n+6832) / 6 ~ A259015(n) / 8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert A. Russell_, Jul 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015800 | OEIS-A356015

a(n) = n! * Sum_{k=1..n} 1/(k * floor(n/k)!).

Conjecture: a(n) ~ c * n!, where c = 0.95488757... - _Vaclav Kotesovec_, Aug 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Jul 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015801 | OEIS-A356026

Main diagonal of right-and-left variant of Kimberling expulsion array, A007063.

Conjectures involving a = A007063 and b = A356026:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015802 | OEIS-A356041

Numerator of the permanent of the matrix [m(j,k)]_{j,k=0..2*n}, where m(j,k) is 1 or (j+k)/(j-k) according as j = k or not.

Conjecture 1. If n is a positive odd integer, x(0) = 0, and x(1),...,x(n) are variables, then the permanent of the matrix [x(j,k)]_{j,k=0..n} vanishes, where x(j,k) is 1 or (x(j)+x(k))/(x(j)-x(k)) according as j = k or not. Conjecture 2: (-1)^n*a(n) > 0 for all n = 0,1,2,... Conjecture 3: For any odd prime p, the permanent of the matrix [m(j,k)]_{j,k=0..p-1} is congruent to 1 - (-1)^((p-1)/2) modulo p, where m(j,k) is defined as in the title.. Conjecture 4: For any odd prime p, the permanent of the matrix M = [m(j,k)]_{j,k=1..p-1} is congruent to ((p-2)!!)^2 modulo p^2, and det(M) is congruent to ((p-2)!!)^2*(-1)^((p+1)/2)/(p-2) modulo p^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015803 | OEIS-A356078

Highly composite numbers that are multiples of their number of divisors.

Conjecture: this sequence is finite. The prime factorization of the number of divisors of a large highly composite number usually has a high power of 2; this is not true of the number of divisors of a highly composite number itself.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Jul 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015804 | OEIS-A356081

Numbers k such that A356052(k) = A356056(k).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015805 | OEIS-A356109

Numbers k such that k^2 + {1,3,7,13} are prime.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jul 27 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015806 | OEIS-A356110

Numbers k such that k^2 + {1,3,7,13,31} are prime.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jul 27 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015807 | OEIS-A356111

The number of 1+1+1-free ordered posets of [n].

Conjectured g.f.: 2 - 2*x/(B(x)-1+x), where B(x) is the o.g.f. for A001181. - _Michael D. Weiner_, Oct 04 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Bevan_, Jul 27 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015808 | OEIS-A356145

Coefficients of the inverse refined Eulerian partition polynomials [E]^{-1}, partitional inverse to A145271. Irregular triangle read by row with lengths A000041.

Conjecture: row polynomials are R(n,1) for n > 0 where R(n,k) = R(n-1,k+1) - Sum_{j=1..n-1} binomial(n-1,j-1)*R(j,k)*R(n-j,1) for n > 1, k > 0 with R(1,k) = a_k for k > 0. - _Mikhail Kurkov_, Mar 22 2025 The above conjecture is true, see MathOverflow link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tom Copeland_, Jul 27 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015809 | OEIS-A356187

Number of permutations f of {1,...,n} with f(1) = 1 such that those k*f(k) + 1 (k = 1..n) are n distinct primes.

Conjecture: a(n) > 0 except for n = 3,4,5. Also, for any integer n > 5, there is a permutation f of {1,...,n} with f(1) = 3 such that those k*f(k) - 1 (k = 1..n) are n distinct primes. This is stronger than part (i) of the conjecture in A321597.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jul 28 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015810 | OEIS-A356211

Odd numbers that cannot be written as a product of an arbitrary number of rational factors of the form 2 + 1/t_k with integers t_k > 1.

It is conjectured that there are no further terms. This was checked up to 2^21.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_ and _Markus Sigg_, Aug 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015811 | OEIS-A356247

Denominator of the continued fraction 1/(2 - 3/(3 - 4/(4 - 5/(...(n-1) - n/(-1))))).

Conjecture 1: Every term of this sequence is either a prime number or 1. Conjecture 2: The sequence contains all prime numbers which ends with a 1 or 9. Conjecture 3: Except for 5, the primes all appear exactly twice. For a proof of Conjectures 1-3, see Cloitre (2025). - _Sean A. Irvine_, Aug 25 2025 If conjecture 3 is true, then we have:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mohammed Bouras_, Jul 30 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015812 | OEIS-A356252

The smallest number of straight lines that can be used to draw n non-overlapping pentagonal stars.

Conjecture: a(n+1) <= a(n)+2 for n > 1. - _David Consiglio, Jr._, Sep 15 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nicolay Avilov_, Jul 31 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015813 | OEIS-A356253

a(n) is the largest coefficient of P(x) := Product_{k} (x + p_k), where (p_k) are the primes dividing n listed with multiplicity.

Conjectured: a(A002110(n)) = A024451(n), for n > 2. Conjecture equality breaks down after n = 175, as a(A002110(176)) > A024451(176). - _Antti Karttunen_, Feb 08 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Jul 31 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015814 | OEIS-A356275

a(n) is the number of tuples (t_1,t_2,m) of integers 2 <= t_1 <= t_2 and 0 < m < n such that (3 + 1/t_1)^m * (3 + 1/t_2)^(n-m) is an integer.

The pairs (t_1,t_2) that arise for integer products suggest this conjecture: For integers t_1,t_2 >= 2 and m,k > 0, the product (3 + 1/t_1)^m * (3 + 1/t_2)^k can be an integer only when (t_1,t_2) is one of (2,7), (2,7^2), (2^2,13), (2^2,13^3), (5,2^3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Markus Sigg_, Aug 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015815 | OEIS-A356295

Numbers that are not the sum of a nonnegative cube and a prime.

It is conjectured that the subsequence of noncube terms, A045911, is finite (has 6195 terms). But there are infinitely many cubes in this sequence: k^3 if a term if and only if k^3 - (k-1)^3 = 3*k^2 - 3*k + 1 is a nonprime (k-1 is in A257772). For example, for k == 2, 6 (mod 7), 3*k^2 - 3*k

+ 1 is divisible by 7, so k^3 is a term for k == 2, 6 (mod 7) and k > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015816 | OEIS-A356360

Denominator of the continued fraction 1/(2-3/(3-4/(4-5/(...(n-1)-n/(n+1))))).

Conjecture: The sequence contains only 1's and the primes. a(n) = 1 positions appear to correspond to A104275(m), m > 2. Conjecture: all odd primes are seen in order after 11. -_Bill McEachen_, Aug 05 2024 From the conjecture: Except for n = 5, a(n)= 2n - 1 if 2n-1 is prime, 1 otherwise.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mohammed Bouras_, Oct 15 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015817 | OEIS-A356376

Main diagonal of the LORO variant of the array A035486; this is one of eight such sequences discussed in A007063.

Conjecture: every positive integer except 2 occurs exactly once.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015818 | OEIS-A356377

Main diagonal of the ROLI variant of the array A035486; this is one of eight such sequences discussed in A007063.

Conjecture: every positive integer occurs exactly once.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015819 | OEIS-A356378

Main diagonal of the RILO variant of the array A035486; this is one of eight such sequences discussed in A007063.

Conjecture: every positive integer occurs exactly once.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015820 | OEIS-A356379

Main diagonal of the LORI variant of the array A035486; this is one of eight such sequences discussed in A007063.

Conjecture: every positive integer except 2 occurs exactly once.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015821 | OEIS-A356380

Main diagonal of the LIRO variant of the array A035486; this is one of eight such sequences discussed in A007063.

Conjecture: every positive integer occurs exactly once.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015822 | OEIS-A356428

a(0) = a(1) = 0; for n > 1, a(n) is the number of distinct gpf(x)'s in the iterations x -> x - gpf(x) starting at n and ending at 0, where gpf = A006530.

Conjecture: sequence is unbounded. Since a(n) - a(n-gpf(n)) = 0 or 1 (see the formula below), this would imply that every number occurs in this sequence. But it seems that the bigger terms appear rather late: 6 does not appear until a(6664), and 7 does not appear until a(135450) (see A356429).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015823 | OEIS-A356429

Smallest m such that A356428(m) = n, or -1 if there is no such m.

Conjecture: a(n) != -1 for all n. This would be true if A356428 is unbounded; otherwise, this sequence consists of entirely -1's after some point.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015824 | OEIS-A356430

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number not occurring earlier that shares a factor with the number of divisors of a(n-1).

The sequence is conjectured to be a permutation of the positive integers although it may take a large number of terms for the primes to appear, e.g., 19 has not occurred after 100000 terms. In the same range the only fixed points are 1, 2, and 9, and it is likely no more exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Aug 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015825 | OEIS-A356431

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number not occurring earlier that shares a factor with both a(n-1) and the number of divisors of a(n-1).

The sequence is conjectured to be a permutation of the positive integers although it may take a large number of terms for the primes to appear, e.g., 17 has not occurred after 100000 terms. In the same range the only fixed points are the first two terms; it is likely no more exist although this is unknown.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Aug 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015826 | OEIS-A356432

a(1) = 1; for n > 1, a(n) is the smallest positive number not occurring earlier that shares a factor with a(n-1) plus the number of divisors of a(n-1).

The sequence is conjectured to be a permutation of the positive integers. In the first 250000 terms there are twenty-three fixed points: 1, 2, 12, 16, 27 ..., 2279, 5401, 7339. It is possibly no more exist although this is unknown.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Aug 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015827 | OEIS-A356446

Number of permutations f of {1,...,n} with f(1) = 2 and f(2) = 1 such that the numbers f(k)*f(k+1) (0 < k < n) are distinct and Sum_{k=1..n-1} 1/(f(k)*f(k+1)) = 1.

Conjecture: a(n) > 0 for all n > 5. This is stronger than Conjecture 1 in A322069.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015828 | OEIS-A356457

a(n) is the least number that can be written in exactly n ways as p*q + q*r + p*r where (p,q,r) is an unordered triple of distinct primes.

Conjecture: a(n) = A003415(A007304(k)) and a(n) is the least number where at least n solutions for k exist. - _Thomas Scheuerle_, Aug 08 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Aug 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015829 | OEIS-A356459

a(n) = n! * Sum_{k=1..n} Sum_{d|k} d/(k/d)!.

Conjecture: a(n) ~ c * n! * n^2, where c = 0.5732... - _Vaclav Kotesovec_, Aug 12 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Aug 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015830 | OEIS-A356492

a(n) is the determinant of a symmetric Toeplitz matrix M(n) whose first row consists of prime(n), prime(n-1), ..., prime(1).

Conjecture: a(n) is prime only for n = 1 and 2. Conjecture is true because a(n) is even for n >= 4. This is because all but two rows of the matrix consist of odd numbers. - _Robert Israel_, Oct 13 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Aug 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015831 | OEIS-A356493

a(n) is the permanent of a symmetric Toeplitz matrix M(n) whose first row consists of prime(n), prime(n-1), ..., prime(1).

Conjecture: a(n) is prime only for n = 1, 2, and 3. Conjecture is true because a(n) is even for n >= 4. This is because a(n) == A356492(n) (mod 2), and all but two rows of the matrix consist of odd numbers. - _Robert Israel_, Oct 13 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Aug 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015832 | OEIS-A356508

G.f. A(x) satisfies: 2 = Product_{n>=1} (1 + x^n*A(x)) * (1 + x^(n-1)/A(x)).

Conjecture: a(n) == 2 (mod 4) at n = 2*k for all k > 0 that have an odd number of partitions (A052002), otherwise a(n) == 0 (mod 4) when n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015833 | OEIS-A356567

Numbers that generate increasing numbers of consecutive primes when doubled and added to the sequence of odd squares. (Positions of records in A354499.)

Conjecture: this list is complete, since primes get farther apart as numbers increase. (2x-1)^2 + 2*29 generates many primes, with 38 of the first 43 and 105 of the first 156 values of x generating primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven M. Altschuld_, Aug 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015834 | OEIS-A356591

Numbers k such that A225205(k) is in A354513.

Conjecture: the odd numbers (numbers k such that A225204(k)/A225205(k) > sqrt(phi)) have relative density phi^(-1), and the even numbers (number k such that A225204(k)/A225205(k) < sqrt(phi)) have relative density phi^(-2). It is conjectured so because we have lim_{k->+oo} (m/k - sqrt((m^2+1)/(k^2+1)))/(sqrt((m^2+2)/(k^2+1)) - m/k) = phi if m/k -> sqrt(phi).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015835 | OEIS-A356638

Odd composite numbers k such that 2^((k-1)/2) == -1 (mod k).

Counterexamples (pseudoprimes) to the hypothesis that this congruence is sufficient to prove that an odd number is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeppe Stig Nielsen_, Aug 19 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015836 | OEIS-A356639

Number of integer sequences b with b(1) = 1, b(m) > 0 and b(m+1) - b(m) > 0, of length n which transform under the map S into a nonnegative integer sequence. The transform c = S(b) is defined by c(m) = Product_{k=1..m} b(k) / Product_{k=2..m} (b(k) - b(k-1)).

These transformation pairs are conjectured:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Aug 19 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015837 | OEIS-A356664

Numbers k such that A225205(k) is in A354549.

Conjecture: this and A356591 have the same natural density.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015838 | OEIS-A356677

The lexicographically earliest infinite squarefree sequence of nonnegative integers that starts with 1.

Conjecture: For any integer k>2, prepending a k at the beginning of this sequence gives the corresponding least word beginning k,1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joey Lakerdas-Gayle_, Aug 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015839 | OEIS-A356678

The lexicographically earliest infinite squarefree sequence of nonnegative integers that starts with 2.

Conjecture: For any integer k>2, prepending a k at the beginning of this sequence gives the corresponding least word beginning k,2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joey Lakerdas-Gayle_, Aug 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015840 | OEIS-A356714

Cardinality of the set{a_1+a_2+a_3+a_4: -floor((n-1)/2) <= a_1,a_2,a_3,a_4 <= floor(n/2), and a_1^2,a_2^2,a_3^2,a_4^2 are pairwise distinct}.

Conjecture 1: Let A_1,...,A_n (n > 3) be finite subsets of a field F with |A_i| > 2*n for all i = 1..n. Then the set {a_1+...+a_n: a_1,...,a_n belong to A_1,...,A_n respectively, and a_1^2,...,a_n^2 are pairwise distinct} contains at least min{p(F), |A_1|+...+|A_n|-n^2+1} elements, where p(F) is the positive infinity if F is of characteristic zero, and p(F) is p if the characteristic of F is a prime p. We also have the following conjecture extending Conjecture 1 to any group. Conjecture 2: Let G be any group with |G| > 1, and let p(G) be the minimum of orders of all nonzero elements of G (which is the positive infinity if each nonzero element of G has infinite order). If A_1,...,A_n (n > 3) are finite subsets of G with |A_i| > 2*n for all i = 1..n. Then the set {a_1+...+a_n: a_1,...,a_n belong to A_1,...,A_n respectively, and there are no 1<= i < j <= n with a_i = a_j or a_i = -a_j} contains at least min{p(F), |A_1|+...+|A_n|-n^2+1} elements.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015841 | OEIS-A356716

a(n) is the integer w such that (c(n)^2, -d(n)^2, -w) is a primitive solution to the Diophantine equation 2*x^3 + 2*y^3 + z^3 = 11^3, where c(n) = F(n+2) + (-1)^n * F(n-3), d(n) = F(n+1) + (-1)^n * F(n-4) and F(n) is the n-th Fibonacci number (A000045).

Conjecture: Several positive examples of conjecture: Then such c(n) and d(n) satisfy the conjecture. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _XU Pingya_, Aug 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015842 | OEIS-A356730

Conductor of the elliptic curve y^2 = x^3 + n.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015843 | OEIS-A356741

a(n) is the least prime(m) such that prime(n) + prime(m)# is prime, where prime(m)# denotes the product of the first m primes, or -1 if no such prime(m) exists.

Conjecture: Such a prime(m) exists for every n, i.e., a(n) is never -1 for n>1. Conjecture: Limit_{N->oo} (Sum_{n=2..N} a(n)) / (Sum_{n=2..N} log(prime(n))) = C with C constant between 0.5 and 1 inclusive.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Sep 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015844 | OEIS-A356751

Positive integers m such that x^2 - x + m contains more than m/2 prime numbers for x = 1, 2, ..., m.

Furthermore, it is conjectured that the present sequence consists of only 16 terms (it has been checked by brute force that there are only 16 terms which are smaller than 20000) and that they are all prime or semiprime (e.g., a(12) = 161, a(13) = 221, and a(16) = 377 are semiprime). Lastly, we trivially point out that all terms must be odd, since if m is even, then x^2 - x + m is also even (and x^2 - x + 2 has only one prime for x <= 2). Stronger conjecture: for every real number e > 0 and every integer m > 0, there are finitely many integer polynomials P(x) = Ax^2 + Bx + C with at least e*m primes in P(1), ..., P(m) and max(|A|, |B|, |C|) <= m. - _Charles R Greathouse IV_, Sep 11 2022 Altering the bounds for x in the definition to 0 <= x <= m-1 (and counting the same prime twice for x=0 and x=1 if m is prime) would result in an additional term 2. Conjecturally, there would be no more additional terms. - _Pontus von Brömssen_, Jun 20 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Aug 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015845 | OEIS-A356756

Positive integers m such that x^2 + x + m contains at least m/2 prime numbers for x = 1, 2, ..., m.

Furthermore, it is conjectured that the present sequence consists of only 15 terms (it has been checked by brute force that there are only 15 terms which are smaller than 20000). Lastly, we trivially point out that all terms must be odd, since if m is even, then x^2 + x + m is also even.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Aug 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015846 | OEIS-A356799

Table read by antidiagonals: T(n,k) (n >= 2, k >= 1) is the number of regions formed in a regular 2n-gon by straight line segments when connecting the k+1 points that divide each side into k equal parts to the equivalent point on the side diagonally opposite.

Conjectured formula for the rows for odd values of k for n>=3: Conjectured formula for the rows for even values of k for n>=3: Conjectured formula for the columns for odd values of k for n>=3: Conjectured formula for T(n,2):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Aug 28 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015847 | OEIS-A356810

Decimal expansion of the unique root of the equation x^(x^(((log(x))^(x-1) - 1)/(log(x) - 1))) = x+1 for x in the interval [1,2].

There are no other real solutions (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Flavio Niccolò Baglioni_ and _Marco Ripà_, Aug 29 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015848 | OEIS-A356837

Coordination sequence of the {3,5,3} hyperbolic honeycomb.

Conjectures from _Chai Wah Wu_, Nov 18 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eryk Kopczynski_, Aug 31 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015849 | OEIS-A356850

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number not occurring earlier such that a(n) is coprime to the previous Omega(a(n)) terms.

The two lowest unseen numbers after 1000000 terms are 32 and 48, for former indicating that the longest run of consecutive odd values is only four after 1000000 terms. Although the sequence is conjectured to be a permutation of the positive integers, if these missing terms, especially those of the form 2^k, only appear after the recombination of the lower lines with the upper line then it may take an extraordinarily large number of terms for some values, like 2^k for large k, to eventually appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Aug 31 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015850 | OEIS-A356851

a(1) = 1, a(2) = 2, a(3) = 4; for n > 3, a(n) is the smallest positive number not occurring earlier such that a(n) shares a factor with the previous Omega(a(n)) terms.

For n > 250 the terms are concentrated along seven lines, see the linked images. Unlike the other six lines, numbers along the second lowest line are somewhat spread out, and these terms contain all numbers with Omega(a(n)) > 1. The lowest line contains all the primes, while the upper five lines contain terms with Omega(a(n)) = 2, 3, and 4. The primes up to n=100000 occur in their natural order except for 11 and 13 which are switched. The only fixed point beyond the first two terms is 10, and it is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Aug 31 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015851 | OEIS-A356852

Minimum over all order two bases for the interval [1, n] of the maximum number of ways some number in the interval [1, n] can be written as a sum of at most two elements of the basis.

It is believed that a(n) tends to infinity. Erdős and Turán conjectured in 1941 that any basis of order two for all the natural numbers cannot avoid large multiplicities. The two conjectures can easily be proved to be equivalent.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Javier Múgica_, Aug 31 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015852 | OEIS-A356856

Primes p such that the least positive primitive root of p (A001918) divides p-1.

If Artin's conjecture is true then this sequence is infinite because it contains all primes with primitive root 2. Conjecture: This sequence has density ~0.548 in the prime numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giorgos Kalogeropoulos_, Aug 31 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015853 | OEIS-A356867

For n >= 1, write n = 3^m + k, where m >= 0 is the greatest power of 3 <= n, and k is in the range 0 <= k < 3^(m+1) - 3^m, then for n such that k=0, a(n)=n, and for n such that k > 0, a(n) is the smallest prime multiple p*a(k), p != 3, that is not already a term.

Conjectured to be a permutation of the positive integers in which the primes appear in order. The conjecture is true: Sequence is a permutation of natural numbers. By definition it is injective, and the surjectivity is guaranteed by the fact that there are infinitely many such n > k encountered by the greedy algorithm that a(n) will be a multiple of a(k), and "the smallest prime multiple" condition guarantees that all multiples of a(k) will eventually appear. That the primes and A100484 appear in order follows from the formulas a(3^m + 1) = prime(m+2), and a(3^m + 2) = 2*prime(m+2). Conjectures from _Jianing Song_, Nov 23 2022: (Start) (1) a(3^m+2) = 2*prime(m+2) for m >= 2. - [The conjecture is true because a(2) = 2 and 3^m + 2 < 3^(1+m) + (3^m) + 1 for all m - _Antti Karttunen_, Sep 16 2023] These are conjectures so far:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Sep 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015854 | OEIS-A356886

Write n as 2^m - k, where 2^m is the least power of 2 such that 2^m >= n, and k is a number in the range 0 <= k < 2^(m-1) - 1. Then for n such that k=0, a(n)=n, and for n such that k > 0, a(n) is the smallest odd prime multiple of a(k) that is not already a term.

Conjectured to be a permutation of the positive integers in which the primes appear in order. The even bisection, when divided by 2 reproduces the sequence. Has similar properties to the Doudna sequence, A005940.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Sep 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015855 | OEIS-A356903

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number not occurring earlier such that a(n) is coprime to the previous tau(a(n)) terms.

Numbers with a large number of divisors relative to the numbers close to it appear much later in the sequence. For example a(96) = 6, a(1873) = 12, a(2328) = 18, a(192) = 16. The sequence is conjectured to be a permutation of the positive integers although it may take a very large number of terms for some values to appear, e.g., after 500000 terms numbers such as 24, 30, 36 have not occurred. In the same range the longest run of consecutive odd values is seven, while the only fixed points are the first three terms, although it is possible others exist for very large values of n if the smaller terms continue to increase relative to the uppermost line.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015856 | OEIS-A356984

Number of regions in an equilateral triangle when n internal equilateral triangles are drawn between the 3n points that divide each side into n+1 equal parts.

Conjecture: a(n) = 3*n^2 + 1 for equilateral triangles that only contain simple vertices when cut by n internal equilateral triangles. This is never the case if (n + 1) mod 3 = 0 for n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015857 | OEIS-A356988

a(n) = n - a^[2](n - a^[3](n-1)) with a(1) = 1, where a^[2](n) = a(a(n)) and a^[3](n) = a(a(a(n))).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015858 | OEIS-A356989

a(n) = n - a^[3](n - a^[4](n-1)) with a(1) = 1, where a^[3](n) = a(a(a(n))) and a^[4](n) = a(a(a(a(n)))).

The sequence of plateau heights begins 4, 6, 9, 13, 19, 28, 41, 60, ..., conjecturally A000930. The plateaus start at absiccsa values n = 5, 8, 12, 17, 25, 37, 54, 79, ..., conjecturally A179070, and terminate at abscissa values n = 6, 9, 13, 19, 28, 41, 60, ..., conjecturally A000930.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015859 | OEIS-A356990

a(n) = n - a^[4](n - a^[5](n-1)) with a(1) = 1, where a^[4](n) = a(a(a(a(n)))) and a^[5](n) = a(a(a(a(a(n))))).

The plateaus start at absiccsa values n = 6, 9, 13, 18, 24, 33, 46, 64, ..., which appears to be A014101, and terminate at abscissa values 7, 10, 14, 19, 26, 36, 50, ..., conjecturally A003269.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015860 | OEIS-A356992

Then a(n) = n - b(n - b(n - b(n - b(n - b(n - b(n)))))) for n >= 2, where b(n) = A356988(n).

The sequence of plateau heights begins 4, 7, 11, 18, 29, 47, ..., conjecturally the Lucas sequence {A000032(k): k >= 3}. The plateaus end at abscissa values n = 7, 12, 19, 31, 50, 81, ..., conjecturally the sequence {A013655(k): k >= 3}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015861 | OEIS-A356995

a(n) = b(n) - b(b(n)) - b(n - b(n)) for n >= 3, where b(n) = A356988(n).

Starting at n = 7, the sequence consists of successive blocks of integers of the form 1, 2, 3, ..., F(k) - 1, F(k), F(k) - 1, ..., 3, 2, 1, where F(k), k >= 1, denotes the k-th Fibonacci number, followed by a string of zeros conjecturally of length 1 + 2*F(k+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015862 | OEIS-A356996

a(n) = b(n) - b(b(n)) - b(n - b(n)) for n >= 3, where b(n) = A356989(n).

The sequence of local peak values of the line graph of the sequence {a(n)} begins 1, 1, 1, 2, 3, 4, 6, 9, 13, 19, ..., conjecturally A000930; the local peaks occur at abscissa values n = 8, 12, 17, 25, 37, 54, 79, 116, 170, 249, ..., conjecturally {A179070(k): k >= 7}. Cf. A356995 and A356997.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015863 | OEIS-A357007

Number of vertices in an equilateral triangle when n internal equilateral triangles are drawn between the 3n points that divide each side into n+1 equal parts.

Conjecture: a(n) = 3*n^2 + 3 for equilateral triangles that only contain simple vertices when cut by n internal equilateral triangles. This is never the case if (n + 1) mod 3 = 0 for n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015864 | OEIS-A357008

Number of edges in an equilateral triangle when n internal equilateral triangles are drawn between the 3n points that divide each side into n+1 equal parts.

Conjecture: a(n) = 6*n^2 + 3 for equilateral triangles that only contain simple vertices when cut by n internal equilateral triangles. This is never the case if (n + 1) mod 3 = 0 for n > 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015865 | OEIS-A357039

Number of integer solutions to x' = 2n, where x' is the arithmetic derivative of x.

Conjecture: All terms are positive with the exception of a(1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Craig J. Beisel_, Sep 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015866 | OEIS-A357045

Lexicographically earliest sequence of distinct non-palindromic numbers (A029742) such that a(n)+a(n+1) is always a palindrome (A002113).

Conjecture: The sequence contains all non-palindromic numbers (A029742).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _M. F. Hasler_, Sep 14 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015867 | OEIS-A357057

a(n) = A356886(2^n+1)/A356886(2^n-1).

Conjecture: This sequence has the pattern: a prime repeated, a run of m times primes standing alone, the next prime repeated, a run of m+1 times primes standing alone, ... All primes will be repeated once or follow in sequence. We know that if A356886(2^n-1) = p1 then A356886(2^n+1) will be p1*p2. p2 will be the smallest possible prime such that p1*p2 is not yet in the sequence A356886, thus p2 = a(n). Let p1*pn be already in A356886 and pn < p2, then we know that p1*pn will be on a position A356886(2^n-(2^(k+1)-2)) with some k > 0. This should explain this pattern. - _Thomas Scheuerle_, Sep 14 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Curtz_, Sep 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015868 | OEIS-A357058

Number of regions in a square when n internal squares are drawn between the 4n points that divide each side into n+1 equal parts.

Conjecture: a(n) = 4*n^2 + 1 for squares that only contain simple intersections when cut by n internal squares. This is never the case for odd n >= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015869 | OEIS-A357060

Number of vertices in a square when n internal squares are drawn between the 4n points that divide each side into n+1 equal parts.

Conjecture: a(n) = 4*n^2 + 4 for squares that only contain simple intersections when cut by n internal squares. This is never the case for odd n >= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015870 | OEIS-A357061

Number of edges in a square when n internal squares are drawn between the 4n points that divide each side into n+1 equal parts.

Conjecture: a(n) = 8*n^2 + 4 for squares that only contain simple intersections when cut by n internal squares. This is never the case for odd n >= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015871 | OEIS-A357062

Number of ordered solutions to n = x*y*z + x + y + z in positive integers.

Conrey & Shah prove that a(n) << n^(1.3) * log n * (log log n)^4, and conjecture that a(n) << n^e for any e > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Oct 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015872 | OEIS-A357081

Leader at step n of the THROWBACK procedure (see definition in comments).

It is conjectured that every number (n >= 3) appears an infinite number of times in this sequence. a(n) = A087165(n+1) + 2 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anthony M. Kozar Jr._, Sep 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015873 | OEIS-A357082

a(0) = 0; for n > 0, a(n) is the smallest positive number not occurring earlier such that the binary string of a(n-1) + a(n) does not appear in the binary string concatenation of a(0)..a(n-1).

The main concentration of terms lie along the line a(n) = n, so numerous fixed points exists - there are 301 fixed points in the first 300000 terms. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015874 | OEIS-A357097

A multiplication table for the rows of the extended Wythoff array. See comments for definition.

A(A019586(x), A019586(y)) = A019586(A101330(x,y)). (conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Munn_, Sep 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015875 | OEIS-A357127

a(n) = A081257(n) if A081257(n) > n, otherwise a(n) = 1.

Conjecture 1: If a(n) != 1, then a(n) = a(a(n) - n - 1). Conjecture 2: If n != m and a(n) = a(m), then

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mohammed Bouras_, Sep 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015876 | OEIS-A357130

a(n) = 2*n - (-1)^n*(1+(n mod 2)).

Conjecture 1: Let G be an additive group with |G| > 1, and let p(G) be the minimum of those orders of nonzero elements of G (which is the positive infinity if each nonzero element of G has infinite order). Let n > 1 be an integer, and let A be a finite nonempty subset of G. Does the following extension of Conjecture 1 also hold? Conjecture 2: Let G be an additive group with |G| > 1, and let p(G) be as in Conjecture 1. Let A_1, ..., A_n (n > 1) be finite subsets of G with |A_i| > 1 for all i = 1..n. Conjectures 1 and 2 were partially confirmed by Han Wang and Zhi-Wei Sun in their preprint arXiv:2210.12044. - _Zhi-Wei Sun_, Oct 23 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Sep 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015877 | OEIS-A357196

Number of regions in a hexagon when n internal hexagons are drawn between the 6n points that divide each side into n+1 equal parts.

Conjecture: a(n) = 6*n^2 + 1 for hexagons that only contain simple intersections when cut by n internal hexagons.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 17 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015878 | OEIS-A357197

Number of vertices in a hexagon when n internal hexagons are drawn between the 6n points that divide each side into n+1 equal parts.

Conjecture: a(n) = 6*n^2 + 6 for hexagons that only contain simple intersections when cut by n internal hexagons.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 17 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015879 | OEIS-A357198

Number of edges in a hexagon when n internal hexagons are drawn between the 6n points that divide each side into n+1 equal parts.

Conjecture: a(n) = 12*n^2 + 6 for hexagons that only contain simple intersections when cut by n internal hexagons.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 17 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015880 | OEIS-A357216

Table read by antidiagonals: T(n,k) (n >= 3, k >= 0) is the number of regions in an n-gon when k internal n-gons are drawn between the n*k points that divide each side into k+1 equal parts.

Conjecture: the only n-gons that contain non-simple intersections are the 3-gon (triangle), 4-gon (square), and 6-gon (hexagon). Conjectured formula for all columns for n >= 7: T(n,k) = n*k^2 + 1. Conjectured formula for all rows except for n = 3, 4, 6: T(n,k) = n*k^2 + 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 18 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015881 | OEIS-A357235

Table read by antidiagonals: T(n,k) (n >= 3, k >= 0) is the number of vertices in an n-gon when k internal n-gons are drawn between the n*k points that divide each side into k+1 equal parts.

Conjecture: the only n-gons that contain non-simple intersections are the 3-gon (triangle), 4-gon (square), and 6-gon (hexagon). Conjectured formula for all columns for n >= 7: T(n,k) = n*k^2 + n. Conjectured formula for all rows except for n = 3, 4, 6: T(n,k) = n*k^2 + n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 19 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015882 | OEIS-A357254

Table read by antidiagonals: T(n,k) (n >= 3, k >= 0) is the number of edges in an n-gon when k internal n-gons are drawn between the n*k points that divide each side into k+1 equal parts.

Conjecture: the only n-gons that contain non-simple intersections are the 3-gon (triangle), 4-gon (square), and 6-gon (hexagon). Conjectured formula for all columns for n >= 7: T(n,k) = 2n*k^2 + n. Conjectured formula for all rows except for n = 3, 4, 6: T(n,k) = 2n*k^2 + n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 20 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015883 | OEIS-A357271

Lower bounds for the maximum number of stable matchings in the stable marriage problem based on composing smaller instances.

a(n) is from Appendix C of Thurber's 2002 paper, using the maximum from each row. At the time of publication, the bounds were known to be exact up to n=4. A357269 shows that they are also exact for n=5. This sequence is not to be confused with A069156, also from Thurber's Appendix C, which uses only the first column, making for looser bounds for n > 11. a(6), a(8), a(10), a(12), and a(16) are also conjectured to be exact.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dan Eilers_, Sep 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015884 | OEIS-A357291

a(n) = number of subsets S of {1,2,...,n} having more than 2 elements such that (sum of least two elements of S) < difference between greatest two elements of S.

a(n) = 2^n/21 - n^2/12 + n/6 + O(1). Conjecture: a(n) = round(2^n/21 - n^2/12 + n/6). - _Charles R Greathouse IV_, Oct 11 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015885 | OEIS-A357314

a(1) = 1; a(n) is the second smallest number k such that k > a(n-1) and concatenation of a(1), ..., a(n-1), k is a palindrome.

Conjecture: Length A055642(a(n)) = A000073(n+2), and A305393 is a sequence of digits in the concatenation of all terms in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gleb Ivanov_, Sep 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015886 | OEIS-A357315

Numbers m such that for all k < m, at least one of m*k - 1 and m*k + 1 is squarefree.

Conjecture: this sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Oct 17 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015887 | OEIS-A357318

Decimal expansion of 1/(2*L), where L is the conjectured Landau's constant A081760.

Decimal expansion of 1/(2*L), where L is the conjectured Landau's constant A081760.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Sep 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015888 | OEIS-A357377

a(0) = 0; for n > 0, a(n) is the smallest positive number not occurring earlier such that |a(n) - a(n-1)| does not appear in the string concatenation of a(0)..a(n-1).

The sequence is conjectured to be a permutation of the positive integers. In the first 200000 terms the only fixed points are 20, 24, 43 and 115. It is likely no more exist although this is unknown.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015889 | OEIS-A357385

a(n) = A071626(n+1) - A071626(n).

Conjectures: These conjectures are empirically supported by the following numbers of occurrences of the terms -1, 0 and 1 up to n = 10^k:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Sep 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015890 | OEIS-A357388

Numbers k such that A071626(k) < A071626(k+1).

Conjecture: This sequence has an asymptotic density 0.028..., which is the same as the asymptotic density of A357390 (see A357385).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Sep 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015891 | OEIS-A357409

a(n) is the maximum number of positive numbers in a set of n consecutive positive or negative odd numbers such that the number of pairs that add to a power of 2 is maximal.

Conjecture: a(n) = A274089(n) + 2^t - 1, where t is the number of solutions k > 0 to the inequality n >= 2^(2*(k+1) + 1) - 2^(k+1) - 1. For n < 27, (2^t - 1) is 0, it is 1 for 27 <= n < 119, and it is 3 for 119 <= n < 495.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Sep 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015892 | OEIS-A357432

a(1) = 1; a(2) = 2; for n > 2, a(n) is the smallest positive number not occurring earlier such that a(n) plus the sum of all previous terms appears in the string concatenation of a(1)..a(n-1).

The sequence is conjectured to be a permutation of the positive integers. In the first 20000 terms the fixed points are 393, 514, 1546, and 7854, although more likely exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 28 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015893 | OEIS-A357433

a(1) = 1; a(2) = 2; for n > 2, a(n) is the smallest positive number not occurring earlier such that the binary string of a(n) plus the sum of all previous terms appears in the binary string concatenation of a(1)..a(n-1).

The sequence is conjectured to be a permutation of the positive integers. In the first 20000 terms there are twenty-five fixed points starting 3, 37, 84, 99, 103, 166.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 28 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015894 | OEIS-A357448

Fixed point starting with 0 of the two-block substitution 00->010, 01->010, 10->101, 11->101.

Conjectures: (II) The frequencies mu[w] of the words w occurring in (a(n)) exist. Some conjectured values: mu[00] = 1/10, mu[01] = 4/10. Moreover, it appears that mu is invariant for taking binary complements.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Dekking_, Sep 29 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015895 | OEIS-A357449

a(0) = 0; for n > 0, a(n) is the smallest positive number not occurring earlier such that the binary string of a(n) plus the largest previous term does not appear in the binary string concatenation of a(0)..a(n-1).

The main concentration of terms lies near the line a(n) = n; there are 26 fixed points in the first 100000 terms. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 29 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015896 | OEIS-A357506

a(n) = A005258(n)^3 * A005258(n-1).

The Apéry numbers B(n) = A005258(n) satisfy the supercongruences B(p) == 3 (mod p^3) and B(p-1) == 1 (mod p^3) for all primes p >= 5 (see, for example, Straub, Example 3.4). It follows that a(p) == 27 (mod p^3) for all

primes p >= 5. We conjecture that, in fact, the stronger congruence a(p) == 27 (mod p^5) holds for all primes p >= 3 (checked up to p = 251). Compare with the congruence B(p) + B(p-1) == 4 (mod p^5) conjectured to hold for all primes p >= 5. See A352655. Conjecture: for r >= 2, a(p^r) == a(p^(r-1)) (mod p^(3*r+3)) for all primes p >= 5. - _Peter Bala_, Oct 13 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015897 | OEIS-A357507

a(n) = A005259(n)^5 * (A005259(n-1))^7.

The Apéry numbers A(n) = A005259(n) satisfy the supercongruences A(p) == 5 (mod p^3) and A(p-1) == 1 (mod p^3) for all primes p >= 5 (see, for example, Straub, Introduction). It follows that a(p) == 3125 (mod p^3) for all primes p >= 5. We conjecture that, in fact, the stronger congruence a(p) == 3125 (mod p^5) holds for all primes p >= 3 (checked up to p = 251). Compare with the congruence A(p) + 7*A(p-1) == 12 (mod p^5) conjectured to hold for all primes p >= 5. See A212334. Conjecture: a(p^r) == a(p^(r-1)) (mod p^(3*r+3)) for r >= 2 and all primes p >= 5. - _Peter Bala_, Oct 26 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015898 | OEIS-A357508

a(n) = binomial(4*n,2*n) - 2*binomial(4*n,n).

Let m be an integer and k a positive integer. Define u(n) = binomial((m+2)*n,(k+1)*n) - binomial(m,k)*binomial((m+2)*n,n). We conjecture that u(n) == u(1) (mod p^5) for all primes p >= 7. [added 22 Oct 2022: the conjecture is true: apply Helou and Terjanian, Section 3, Proposition 2.] Conjecture: for r >= 2, u(p^r) == u(p^(r-1)) (mod p^(3*r+3)) for all primes p >= 5. - _Peter Bala_, Oct 13 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015899 | OEIS-A357509

a(n) = 2*binomial(3*n,n) - 9*binomial(2*n,n).

For integers j and k, not necessarily positive, define u(n) = k^2*(k - 1)*binomial(j*n,n) - j^2*(j - 1)*binomial(k*n,n). We conjecture that u(p) == u(1) (mod p^5) for all primes p >= 7. This is essentially the case (j, k) = (3, 2). [Follows from Helou and Terjanian (2008), Section 3, Proposition 2.] Conjecture: for r >= 2, u(p^r) == u(p^(r-1)) (mod p^(3*r+3)) for all primes p >= 5. - _Peter Bala_, Oct 13 2022 Conjecture: the stronger supercongruence a(p) == a(1) (mod p^5) holds for all primes p >= 7. The conjecture is true: apply Helou and Terjanian, Section 3, Proposition 2. - _Peter Bala_, Oct 22 2022 The conjecture was proved by Aidagulov and Alekseyev; see the remarks following Corollary 2. - _Peter Bala_, Oct 29 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015900 | OEIS-A357510

a(n) = Sum_{k = 0..n} k * binomial(n,k)^2 * binomial(n+k,k)^2.

Conjecture: a(p-1) == 0 (mod p^4) for all primes p >= 5 (checked up to p = 499).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015901 | OEIS-A357511

a(n) = numerator of Sum_{k = 1..n} (1/k) * binomial(n,k)^2 * binomial(n+k,k)^2 for n >= 1 with a(0) = 0.

Conjecture: a(p-1) == 0 (mod p^4) for all primes p >= 7 (checked up to p = 499).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015902 | OEIS-A357512

a(n) = Sum_{k = 0..n} k^5 * binomial(n,k)^2 * binomial(n+k,k)^2.

Let m >= 2 and set u(n) = Sum_{k = 0..n} k^(2*m+1) * binomial(n,k)^2 * binomial(n+k,k)^2. We conjecture that there is a finite set of primes P(m) such that u(n-1) == 0 (mod n^4) for all odd numbers n not divisible by an element of P(m). For example, calculation suggests that P(2) = {3} (the present sequence), while conjecturally P(3) = {3, 5}, P(4) = {3, 7}, P(5) = {3, 5}, P(6) = {3, 11}, P(7) = {3, 5, 7, 13}, P(8) = {3}, P(9) = {3} and P(10) = {3, 7, 19}. Conjecture: a(n-1) == 0 (mod n^4) for all n congruent to 1 or 5 mod 6 (checked up to n = 499).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015903 | OEIS-A357518

Unique fixed point of the two-block substitution 00->111, 01->110, 10->101, 11->100.

Conjectures: (III) The frequencies mu[w] of the words w occurring in (a(n)) exist. Some conjectured values: mu[1] = 3/5, and mu[0000] = 0, mu[0001] = 0, mu[0010] = 1/30, mu[0011] = 1/15, mu[0100] = 1/30, mu[0101] = 1/15, mu[0110] = 2/15, mu[0111] = 1/15. Moreover, it seems that mu is reversal invariant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Dekking_, Oct 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015904 | OEIS-A357520

Expansion of Product_{k>=0} (1 - x^Lucas(k)).

Conjecture: These numbers are all in {-2,-1,0,1,2}. - _Ira M. Gessel_, Mar 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Oct 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015905 | OEIS-A357532

a(n) = Sum_{k=0..floor(n/3)} (n-2*k)!/(n-3*k)!.

a(n) ~ c * n^(n/3 + 1/2) / (3^(n/3) * exp(n/3 - n^(2/3)/3^(2/3) - 2*n^(1/3) / 3^(7/3))) * (1 + 1235/(729 * 3^(2/3) * n^(1/3)) + 9452027/(15943230 * 3^(1/3) * n^(2/3)) + 16015315669/(41841412812*n)), where c = 0.50682110703119..., conjecture: c = exp(4/81) * sqrt(2*Pi) / 3^(3/2). - _Vaclav Kotesovec_, Nov 25 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Nov 19 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015906 | OEIS-A357539

a(n) = coefficient of x^n/n! in: Sum_{n>=0} (x*exp(x))^(n*(n+1)/2).

Conjecture: Limit_{n->infinity} (a(n)/n!)^(1/n) = 1/LambertW(1). - _Vaclav Kotesovec_, Dec 06 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015907 | OEIS-A357558

a(n) = Sum_{k = 0..n} (-1)^(n+k)*k*binomial(n,k)*binomial(n+k,k)^2.

Conjecture: a(p-1) == 0 (mod (p - 1)*p^4) for all primes p >= 5 (checked up to p = 499).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015908 | OEIS-A357559

a(n) = Sum_{k = 0..n} (-1)^(n+k)*k^3*binomial(n,k)*binomial(n+k,k)^2.

Conjectures: Conjecture: 3*a(2*n) == 0 (mod n^2*(2*n + 1)^3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015909 | OEIS-A357560

a(n) = the numerator of (Sum_{k = 1..n} (-1)^(n+k)*(1/k)*binomial(n,k)* binomial(n+k,k)^2).

Conjectures: Conjecture: a(p-1) == 0 (mod p^4) for all primes p >= 7 (checked up to p = 499).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015910 | OEIS-A357561

a(n) = the numerator of (Sum_{k = 1..n} (-1)^(n+k)*(1/k^3)*binomial(n,k)* binomial(n+k,k)^2).

Conjecture: for odd m >= 1, S_m(p-1) == 0 (mod p^4) for all primes p > m+4. Conjecture: a(p-1) == 0 (mod p^4) for p = 5 and all primes p >= 11 (checked up to p = 499).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015911 | OEIS-A357565

a(n) = 3*Sum_{k = 0..n} binomial(n+k-1,k)^2 + 2*Sum_{k = 0..n} binomial(n+k-1,k)^3.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015912 | OEIS-A357566

a(n) = (Sum_{k = 0..n} binomial(n+k-1,k)^2)^3 * (Sum_{k = 0..n} binomial(n+k-1,k)^3)^2.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015913 | OEIS-A357567

a(n) = 5*A005259(n) - 14*A005258(n).

Conjectures: Additional conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 19 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015914 | OEIS-A357568

a(n) = 9*binomial(2*n,n)^2 - 8*binomial(3*n,n).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015915 | OEIS-A357569

a(n) = binomial(3*n,n)^2 - 27*binomial(2*n,n).

Conjectures: Adamczewski includes proof abstract regarding Conjecture 1 in linked article. - _Michael De Vlieger_, Aug 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015916 | OEIS-A357574

a(n) is the maximum number of pairs that sum to a power of 2 in a set of n consecutive odd numbers.

Conjecture: a(n+1) - a(n) = k, if n is even then A129868(k-1) < n < A129868(k), if n is odd then A020515(k) <= n < A020515(k+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Oct 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015917 | OEIS-A357578

Lexicographically earliest infinite sequence of distinct positive numbers with the property that a(n) is the smallest number not yet in the sequence with a Hamming weight equal to the Hamming weight of the XOR of previous two terms.

Conjecture: a(n) is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nathan Nichols_, Oct 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015918 | OEIS-A357579

Lexicographically earliest sequence of distinct numbers such that no sum of consecutive terms is a square or higher power of an integer.

An unproved conjecture (for the present sequence) is that all integers which are not nontrivial powers will eventually appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carl Witthoft_, Oct 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015919 | OEIS-A357581

Square array read by antidiagonals of numbers whose symmetric representation of sigma consists only of parts that have width 1; column k indicates the number of parts and row n indicates the n-th number in increasing order in each of the columns.

Conjecture: T(n,k) is the n-th number with k odd divisors and k 2-dense sublists of divisors. - _Omar E. Pol_, Feb 17 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Oct 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015920 | OEIS-A357583

Triangle read by rows. Convolution triangle of the Bell numbers.

Conjecture: row polynomials are x*R(n,1) for n > 0 where R(n,k) = R(n-1,k+1) + x*R(n-1,1)*R(1,k) for n > 1, k > 0 with R(1,k) = Bell(k) for k > 0. The same recursion seems to work for self-convolution of any other sequence. - _Mikhail Kurkov_, Apr 05 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Oct 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015921 | OEIS-A357595

Lexicographically earliest infinite sequence of distinct positive integers such that a(n+1) is the least k != j, for which gcd(k, j) > 1; j = n + a(n).

If n + a(n) = prime p, a(n+1) is the smallest multiple (>1) of p, which has not occurred earlier. Conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Oct 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015922 | OEIS-A357600

Largest number k such that C(-k) is the cyclic group of order n, where C(D) is the class group of the quadratic field with discriminant D; or 0 if no such k exists.

Conjecture: all terms are odd. Conjecture: all terms are congruent 19 (mod 24). - _Ralf Stephan_, May 23 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Oct 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015923 | OEIS-A357601

For n a power of 2, a(n) = n; otherwise, if 2^m is the greatest power of 2 not exceeding n and if k = n-2^m, then a(n) is the smallest number having d(a(k))+1 divisors which has not occurred earlier (d is the divisor counting function A000005).

Based on a similar recursion to that which produces the Doudna sequence, A005940. Conjectured to be permutation of the positive integers in which the primes appear in natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Oct 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015924 | OEIS-A357604

Number of prime powers in the sequence of the floor of n/k for k <= n, A010766.

Conjecture: a(n+1) - a(n) has all its record values at n = A135972(k) and the record values of a(n-1) - a(n) are a subsequence of A025487. This was verified for n = 1..20000. - _Thomas Scheuerle_, Oct 06 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Randell Heyman_, Oct 06 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015925 | OEIS-A357614

Lexicographically earliest infinite sequence of distinct positive integers such that a(n+1) is the least k != j, for which gcd(k, j) > 1, where j = a(n) + prime(n).

If a(n) + prime(n) is a prime p, then a(n+1) is the smallest multiple (>1) of p that has not occurred earlier. Conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Oct 06 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015926 | OEIS-A357637

Triangle read by rows where T(n,k) is the number of integer partitions of n with half-alternating sum k, where k ranges from -n to n in steps of 2.

Conjecture: The column sums are A029862.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Oct 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015927 | OEIS-A357638

Triangle read by rows where T(n,k) is the number of integer partitions of n with skew-alternating sum k, where k ranges from -n to n in steps of 2.

Conjecture: The columns are palindromes with sums A298311.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Oct 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015928 | OEIS-A357671

a(n) = Sum_{k = 0..n} (binomial(n+k-1,k) + binomial(n+k-1,k)^2).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015929 | OEIS-A357672

a(n) = Sum_{k = 0..n} binomial(n+k-1,k) * Sum_{k = 0..n} binomial(n+k-1,k)^2.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015930 | OEIS-A357673

a(n) = 4*Sum_{k = 0..2*n} binomial(n+k-1,k) + 3*Sum_{k = 0..2*n} binomial(n+k-1,k)^2.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015931 | OEIS-A357674

a(n) = (Sum_{k = 0..2*n} binomial(n+k-1,k))^4 * (Sum_{k = 0..2*n} binomial(n+k-1,k)^2)^3.

Conjectures: Adamczewski includes a proof abstract regarding Conjecture 1 in linked article. - _Michael De Vlieger_, Aug 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015932 | OEIS-A357678

Numbers k equal to the integer log of the sum of k and its digit reversal.

10^n + 7 is a term if 10^n + 1 is prime, however that is conjectured to occur only for n = 0, 1 and 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Oct 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015933 | OEIS-A357733

Integer lengths of the sides of such regular hexagons that a polyline described in A356047 exists.

Conjectures from _Chai Wah Wu_, Mar 13 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander M. Domashenko_, Oct 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015934 | OEIS-A357735

a(1)=1, a(2)=2. Thereafter a(n+1) is least k != partial sum s(n) which has not occurred earlier, such that gcd(k, s(n)) > 1.

It follows from the definition that if s(n) is prime then a(n+1) = 2*s(n). This happens only once in the sequence, when a(3)=6, following s(2)=3. For all n > 2 s(n) is composite. Conjectured to be a permutation of the positive integers (primes not in natural order).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Oct 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015935 | OEIS-A357741

Semiprimes k such that k is divisible by its index in the sequence of semiprimes.

Conjecture: a(18) > 10^17. - _Martin Ehrenstein_, Oct 28 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lucas A. Brown_, Oct 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015936 | OEIS-A357777

a(1)=1, a(2)=2. Thereafter a(n+1) is the smallest k such that gcd(k, a(n)) > 1, and gcd(k, s(n)) = 1, where s(n) is the n-th partial sum.

Conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Oct 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015937 | OEIS-A357825

Total number of n-tuples of semi-Dyck paths from (0,0) to (n,n-2*j) for j = 0..floor(n/2).

Conjecture: the superconguence a(2*p-1) == 1 (mod p^3) holds for all primes p >= 5 (checked up to p = 101). - _Peter Bala_, Mar 20 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Oct 14 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015938 | OEIS-A357880

a(1) = a(2) = 1; for n > 2, a(n) is the smallest positive number such that a(n) plus the sum of all previous terms appears in the string concatenation of a(1)..a(n-1).

It is conjectured that all numbers eventually appear. In the first 100000 terms the only fixed point is 210; it is likely no more exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Oct 18 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015939 | OEIS-A357888

a(n) is the minimal squared length of the longest side of a strictly convex grid n-gon of smallest area.

It is conjectured that at least one polygon of smallest area exists with 4 sides of length 1 for n >= 8 and additionally 4 sides of squared length 2 for n >= 12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Nov 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015940 | OEIS-A357943

a(0) = 0; a(1) = 1, a(2) = 2; for n > 2, a(n) is the number of times the term a(n - 1 - a(n-1)) has appeared in the sequence.

The sequence is conjectured to contain all positive numbers. See A357944 for the index of where a given number first appears.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Oct 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015941 | OEIS-A357955

a(n) = 3*binomial(4*n,n) - 20*binomial(3*n,n) + 54*binomial(2*n,n).

Conjectures: Conjecture 1) was proved by Aidagulov and Alekseyev; see the remarks following Corollary 2. - _Peter Bala_, Oct 29 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015942 | OEIS-A357956

a(n) = 5*A005259(n) - 2*A005258(n).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015943 | OEIS-A357957

a(n) = A005259(n)^5 - A005258(n)^2.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015944 | OEIS-A357958

a(n) = 5*A005259(n) + 14*A005258(n-1).

Conjectures: There is also a product version of these conjectures: 3) the sequence {u(n): n>= 1} defined by u(n) = A005259(n)^25 * A005258(n-1)^14 conjecturally satisfies the congruences in 1) and 2) above. For the sequence u(n) = A005259(n)^25 * A005258(n-1)^14 of conjecture 3), the congruence 2) fails at p = 3: the 3-adic valuation of u(3^r) - u(3^(r-1)) equals 3*r + 2 for r = 2, 3, 4, one short of the required 3*r + 3. (The main sequence a(n) does satisfy 2) at p = 3, with valuation exactly 3*r + 3 for r = 2, 3, 4.) The data suggest that for u(n) the congruence in 2) holds modulo 3^(3*r+2) at p = 3, or equivalently that in conjecture 3) the second congruence should be restricted to primes p >= 5. - _Sunsu Jeong_, Aug 10 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015945 | OEIS-A357959

a(n) = 5*A005259(n-1) + 2*A005258(n).

Conjectures: There is also a product version of these conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015946 | OEIS-A357960

a(n) = A005259(n-1)^5 * A005258(n)^6.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015947 | OEIS-A357963

a(1)=1, a(2)=2. Thereafter, if there are prime divisors p of a(n-1) which do not divide a(n-2), a(n) is the least novel multiple of any such p. Otherwise a(n) is the least novel multiple of the squarefree kernel of a(n-1). See comments.

Departs from A064413 and A352187 at a(19), a(31) respectively, and apparently shares the way odd primes are proven to appear in the former and conjectured to appear in the latter; namely as 2*p, p, 3*p. Conjectures: Permutation of the positive integers with primes in natural order, appearing in same way as in EKG.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Oct 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015948 | OEIS-A357990

Square array T(n, k), n >= 0, k > 0, read by antidiagonals, where T(n, k) = R(n, k+1) - R(n, k) for n > 0, k > 0 with T(0, k) = 1 for k > 0 and where R(n, k) = T(A053645(n), k)*k^(A290255(n) + 1).

Conjecture: T(n, 1) = A329369(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Nov 20 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015949 | OEIS-A357992

a(1)=1,a(2)=2,a(3)=3. Thereafter, if there are prime divisors p of a(n-2) which do not divide a(n-1), a(n) is the least novel multiple of any such p. Otherwise a(n) is the least novel multiple of the squarefree kernel of a(n-2).

Conjecture 1: A permutation of the positive integers with primes in natural order. Conjecture 2: The primes are the slowest numbers to appear (see also A352187).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Oct 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015950 | OEIS-A357994

a(1)=1, a(2)=2. Thereafter, if there are prime divisors p of a(n-1) which do not divide a(n-2), a(n) is the greatest least multiple of any such p which has not already occurred. Otherwise a(n) is the least novel multiple of the squarefree kernel of a(n-1). (see comments).

Conjecture: Permutation of the positive integers; primes being in natural order, and the slowest numbers to appear (as in A352187).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Oct 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015951 | OEIS-A358026

Let G(n) = gcd(a(n-2),a(n-1)), a(1)=1, a(2)=2, a(3)=3. Thereafter if G(n) = 1, a(n) is the least novel m sharing a divisor with both a(n-2) and a(n-1). If G(n) > 1 and every prime divisor of a(n-1) also divides a(n-2), a(n) is the least m prime to both a(n-1) and a(n-2). Otherwise a(n) is the least novel multiple of any prime divisor of a(n-1) which does not divide a(n-2).

Conjectured to be a permutation of the positive integers with the primes in natural order, and primes are the slowest numbers to appear (as in A352187).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Oct 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015952 | OEIS-A358035

a(n) = (8*n^3 + 12*n^2 + 4*n - 9)/3.

Conjecture: a(n) is the disorder number of the Aztec diamond of size n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sela Fried_, Oct 26 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015953 | OEIS-A358048

Lexicographically earliest sequence of distinct nonnegative integers on a square spiral such that every number shares a digit with each of its eight surrounding neighbors.

The sequence is conjectured to be a permutation of the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Scott R. Shannon_, Oct 27 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015954 | OEIS-A358082

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number not occurring earlier that shares a factor with Sum_{k=1..n-1} sigma(a(k)).

The sequence shows large jumps in value due to the sum occasionally forming a large prime, e.g., a(279) = 2650277753. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015955 | OEIS-A358093

a(n) = n for 1 <= n <= 2; thereafter a(n) is the least unused m such that rad(m) = rad(rad(a(n-1)) + rad(a(n-2))), where rad(m) = A007947(m).

Conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Nov 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015956 | OEIS-A358094

a(n) is the number of ways n can be reached in the following method: we start with 1, then add or multiply alternately, and each operand must be 2 or 3.

Conjecture: All numbers greater than 145 can be reached in at least 3 different ways. Numbers greater than 145 which can be reached in only three ways are all of the form 27*2^k - 5 (A304387). This is conjectured to arise from the relation: 27*2^(k+2) - 5 = 3*(27*2^(k+1) - 5) - 2*(27*2^k - 5) which is in some sense here the worst case in between *2 and *3.

(End) Proof of the conjecture: We use the identities in A358095. For n > 145, if a(n) <= 2, A358095(n) <= 2. Therefore, n must be a*3*2^k - 2, where a is in the set {3, 7, 26, 30, 32, 34, 40, 49} and k is a positive integer. However, A358096(a*3*2^k - 2) = A358095(a*3*2^(k-1) - 1) >= 3, a contradiction. Hence all numbers greater than 145 can be reached in at least 3 different ways. If a(n) = 3, the only possibilities are that A358095(n) = 0 and A358096(n) = 3 or A358095(n) = 3 and A358096(n) = 0. For the first one, n must be 9*2^k - 2, so A358096(n) = A358095(9*2^(k-1) - 1) >= 4. For the second one, if n is in the form a*3*2^k - 2, A358096(n) = A358095(a*3*2^(k-1) - 1) >= 4; if n = 108*2^k - 3, A358096(n) = A358095(36*2^k - 1) >= 4; if n is in the form 108*2^k - 5, A358096(n) = 0. Hence a(n) = 3 iff n = 27*2^k - 5 for n > 145. - _Yifan Xie_, Jan 07 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yifan Xie_ and _Thomas Scheuerle_, Oct 29 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015957 | OEIS-A358132

Numbers k such that there exists a pair of primes (p,q) with p+q = k such that p*q + k, p*q - k, p*q + A001414(k) and p*q - A001414(k) are all prime.

Conjecture: the sequence includes all sufficiently large k such that both k and A001414(k) are divisible by 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. M. Bergot_ and _Robert Israel_, Oct 31 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015958 | OEIS-A358151

Earliest infinite sequence of distinct integers on a square spiral such that every number equals the sum of its eight adjacent neighbors. See the Comments.

The sequence is conjectured to be a permutation of the integers. The values grow rapidly in size, e.g., in the first 1000 terms the largest magnitude number is a(899) = 3585008350, while the largest magnitude number in the first 100000 terms is a(99854) = -128...904, a number containing 103 digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015959 | OEIS-A358153

Lexicographically earliest infinite sequence of distinct positive integers on a square spiral such that each number shares a factor with its four orthogonally nearest neighbors but shares no factor with its four diagonal next-nearest neighbors.

Like A336957 is it conjectured that all nonprime powers eventually appear. In the first 10000 terms the largest value is a(6975) = 3005315.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 01 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015960 | OEIS-A358175

a(1) = 1, a(2) = 2; a(3) = 3; for n > 3, a(n) is the smallest positive number not occurring earlier that shares a factor with Sum_{k=1..n-1} A001065(a(k)), where A001065(m) is the sum of the proper divisors of m.

The sequence is conjectured to be a permutation of the positive integers. In the first 10000 terms, other than the first three terms, there are thirty-five fixed points, the last being 2051. It is plausible no more exist although this is unknown.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015961 | OEIS-A358176

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number not occurring earlier that shares a factor with sigma(a(n-1)).

The sequence is conjectured to be a permutation of the positive integers. In the first 500000 terms the fixed points are 1, 2, 3, 4, 6, 9, 12. It is unlikely more exist although this is unknown.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015962 | OEIS-A358200

Frequency ranking position of the ratio r(n) between consecutive prime gaps, among all previous ratios {r(i) : 2 < i < n, r(i) = (prime(i) - prime(i-1))/(prime(i-1) - prime(i-2))}. If the ratio r(n) is not among previous ratios, then a(n)=n.

Conjectures: 2) The sequence is unbounded. (Equivalent to conjecture in A001223 on Sep 29 2018.) 3) The relative frequencies (probabilities) of the consecutive prime-gap ratios approach constants as the length of the list of first primes approaches infinity. (Equivalent to conjecture in A001223 on Sep 01 2019.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andres Cicuttin_, Feb 22 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015963 | OEIS-A358201

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number not occurring earlier that shares a factor with sigma(max_{k=1..n-1}a(k)).

The sequence is conjectured to be a permutation of the positive integers. In the first 150000 terms the fixed points are 1, 2, 3, 4, 6, 9, 12, 93, 6003, 6881, 16269, 100707, 114839, 116999. It is likely more exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 03 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015964 | OEIS-A358208

a(1) = 1; a(2) = 2; a(3) = 3; for n > 3, a(n) is the smallest positive number not occurring earlier that shares a factor with Sum_{k=1..n-1} A001065(k), where A001065(k) is the sum of the proper divisors of k.

Conjecture: the sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015965 | OEIS-A358209

a(1) = 1; a(2) = 2; for n > 2, a(n) is the smallest positive number not occurring earlier that shares a factor with A024916(n-1) = Sum_{k=1..n-1} sigma(k).

Conjecture: the sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 04 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015966 | OEIS-A358213

The index of the first occurrence of A002110(n) in A356309.

Also, from a(2)=3 onward conjectured to be the positions of records in A356302 (after its initial zero), while A358214 gives the conjectured record values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015967 | OEIS-A358214

a(n) = A002110(n) - A358213(n).

From a(1) onward conjectured to be the record values in A356302. See also A358213.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 05 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015968 | OEIS-A358267

a(1) = 1, a(2) = 2. Thereafter:(i). If no prime divisor of a(n-1) divides a(n-2), a(n) is the least novel multiple of the squarefree kernel of a(n-1). (ii). If some (but not all) prime divisors of a(n-1) do not divide a(n-2), a(n) is the least of the least novel multiples of all such primes. (iii). If every prime divisor of a(n-1) also divides a(n-2), a(n) = u, the least unused number.

If a(n) = u = p from condition (iii), a(n+1) = 2*p. If p|a(n-1)-> a(n) = p we see m*p->p->u (and u may of course be prime, as in ...,13,17,...). 13 is the first prime to appear consequent to condition (ii), see Example. Consecutive primes appear often: (13,17); (53,59); (61,67); ... Sequence is conjectured to be a permutation of the positive integers with primes appearing slowest, and in natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Nov 06 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015969 | OEIS-A358268

a(n) is the least number k > 0 such that the binary weight of k^n is n times the binary weight of k.

Conjecture: lim_{n -> infinity} a(n) = infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Nov 06 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015970 | OEIS-A358272

Multiplicative sequence with a(p^e) = (-1)^e * p^(2*floor(e/2)) for prime p and e >= 0.

Conjecture: a(n) = Sum_{k=1..n} gcd(k, n) * lambda(gcd(k, n)) for n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Nov 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015971 | OEIS-A358277

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number not occurring earlier such that a(n) is coprime to the previous Omega(a(n-1)) terms.

Unlike A356850 all the terms are concentrated along three straight lines. In the first 100000 terms there are ten fixed points, 1, 2, 3, ..., 27, 57, and it is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015972 | OEIS-A358315

Primes p == 1 (mod 3) such that there exists 1 <= x <= p-2 such that (x+1)^p - x^p == 1 (mod p^2) and that p does not divide x^2 + x + 1.

Conjecture: this density of this sequence among the primes congruent to 1 modulo 3 is the same as that of A068209 among the primes congruent to 2 modulo 3. - _Jianing Song_,

Nov 08 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Nov 08 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015973 | OEIS-A358318

For n >= 5, a(n) is the number of zeros that need to be inserted to the left of the ones digit of the n-th prime so that the result is composite.

Conjecture: the sequence is bounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rida Hamadani_, Nov 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015974 | OEIS-A358319

Multiplicative sequence a(n) with a(p^e) = ((p-2) - (p-1) * e) * p^(e-1) for prime p and e > 0.

Conjecture: a(n) = Sum_{k=1..n} gcd(k, n) * A076479(gcd(k, n)) for n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Nov 09 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015975 | OEIS-A358337

Earliest infinite sequence of distinct integers on a square spiral such that every number equals the sum of its four adjacent neighbors. See the Comments.

The sequence is conjectured to be a permutation of the integers. The values grow rapidly in size, e.g., in the first 1000 terms the largest magnitude number is a(976) = -2492394, while the largest magnitude number in the first 500000 terms is a(499495) = -749...021, a number containing 147 digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015976 | OEIS-A358339

Array read by antidiagonals upwards: A(n,k) is the number of nonequivalent positions in the KRvK endgame on an n X n chessboard with DTM (distance to mate) k, n >= 3, k >= 0.

Conjecture 1: If A(n+1,k) is nonzero, it is always greater than A(n,k). Conjecture 2: As the increments to A225552 become periodic for n>24, so too do the differences between successive polynomials over n for each k>108, and for all j, Sum_{k=2*A225552(n)-j..2*A225552(n)}(A(n,k)) is o(Sum_{k>=0}(A(n,k))) (the same equation from the opposite side). Conjecture 3: (Start) For all n, assuming conjecture 2, because the polynomials provide upper bounds and all are of degree <=5, it is necessary for the sum of all nonzero A(n,k)'s polynomials (for k<=2*A225552(n)) to have n^2 coefficients summing to 3*n^4/2. Conjecture 4: The function o (defined in the FORMULA section), for each parity of k, contains only polynomials over n (without any terms with coefficients of n%2 or (-1)**n). For all m>=3, there exists a pair of polynomials over n (for even and odd k), such that when o adds them to the output of the k-specific fitting polynomial for n=k-m, the values returned are correct for all k>=5*(m-1). (Works for m=2,3,4.) Equivalently, for all k>10, while A(n,k) can be described exactly by a k-specific polynomial over n for all n>=A052928(k), it can be described exactly by the sum of that and a (2*k-n)-specific polynomial over n (or equivalently, over k) for all n>=k-1-floor(k/5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Natalia L. Skirrow_, Nov 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015977 | OEIS-A358354

a(n) = n for n <= 3. Thereafter a(n) is the least m such that rad(m) = rad(rad(a(n-3)) + rad(a(n-1))) where rad is A007947.

Conjecture: for n > 60, a(n) is squarefree. - _Michael De Vlieger_, Jul 01 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Nov 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015978 | OEIS-A358359

a(n) = number of occurrences of n in A128440; i.e., as a number [k*r^m], where r = golden ratio = (1+sqrt(5))/2, k and m are positive integers, and [ ] = floor.

Conjecture: every positive integers occurs infinitely many times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015979 | OEIS-A358360

The 3-adic valuation of the central Delannoy numbers (sequence A001850).

Conjectures from _Ridouane Oudra_, May 20 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Nov 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015980 | OEIS-A358386

Distinct values of A030717 in order of appearance.

Conjecture: 41 is not a term. This conjecture is related to A030720. Nowhere among the first 3*10^5 terms in A030717 does 41 appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Nov 13 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015981 | OEIS-A358444

a(1) = 1, a(2) = 2; for n > 2, a(n) = smallest positive number which has not appeared that has a common factor with a(n-2)^2 + a(n-1)^2.

Conjecture: the sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015982 | OEIS-A358452

The inverse Euler transform of p(n) = n if n is prime, otherwise 1.

Conjecture: signum(a(n)) + (-1)^n = 0 for n >= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Nov 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015983 | OEIS-A358535

a(1) = 1; let S(n) = Sum_{j=1..n-1} a(j) and let T(n) = Sum_{j=1..n-1} d(a(j)) where d(n) = A000005(n). a(n) = least novel k such that (T(n)+d(k)) | (S(n)+k).

Conjecture: 3 does not appear in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_ and _David James Sycamore_, Dec 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015984 | OEIS-A358560

a(n) = Sum_{k=0..floor(n/3)} (n-k)!/(k! * (n-3*k)!).

a(n) ~ c * 2^(2*n/3) * n^(n/3) / (3^(n/3) * exp(n/3 - 2^(1/3) * n^(2/3) / 3^(2/3) + n^(1/3) / (2^(4/3) * 3^(7/3)))) * (1 + 7795/(5832*6^(2/3)*n^(1/3)) + 135724109/(2040733440*6^(1/3)*n^(2/3)) - 5962064767253/(42845606719488*n)), where c = 0.46562048925..., conjecture: c = sqrt(2) * exp(-1/81) / 3. - _Vaclav Kotesovec_, Nov 25 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Nov 22 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015985 | OEIS-A358587

Number of n-node rooted trees of height equal to the number of internal (non-leaf) nodes.

Conjectures from _Chai Wah Wu_, Apr 15 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Nov 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015986 | OEIS-A358588

Number of n-node ordered rooted trees of height equal to the number of internal (non-leaf) nodes.

Conjectures from _Chai Wah Wu_, Apr 14 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Nov 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015987 | OEIS-A358612

Irregular triangle T(n, k), n >= 0, k > 0, read by rows of extended (due to binary expansion of n) Stirling numbers of the second kind.

Conjectured formulas: (Start) Conjecture: T(n, k) = (k-1)^g(n)*T(h(n), k-1) + k^(g(n)+1)*T(h(n), k) for n > 0, k > 1 with T(n, 1) = T(0, 2) = 1 where g(n) = A007814(n) and where h(n) = A025480(n-1). - _Mikhail Kurkov_, Jun 21 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Nov 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015988 | OEIS-A358616

a(n) is the position of the first occurrence of the least term in row n of the Gilbreath array shown in A036262.

Conjecture: Every positive integer occurs in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015989 | OEIS-A358617

a(n) is the number of zeros among the first n terms of row n of the Gilbreath array shown in A036262.

Conjecture: If = (a(n)/n), then (lim inf S) > 2/5 and (lim sup S) < 3/5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Nov 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015990 | OEIS-A358618

First differences of A258036.

Conjecture: All terms belong to {1, 2, 3}. See third comment in A258036.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ and _Robert G. Wilson v_, Oct 31 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015991 | OEIS-A358619

First forward difference of A258037.

Conjecture: All terms belong to {1, 2, 3}. See third comment in A258037.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_ and _Robert G. Wilson v_, Oct 31 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015992 | OEIS-A358628

Square array A(i,j), i >= 0, j >= 0, read by antidiagonals: A(i,j) = Sum_{|X|=0..i} Sum_{|Y|=0..i} Product_{k=1..j} (1+X(k)+Y(k)), where X and Y are multi-indices of length j.

It is conjectured that A(i,j) = binomial(i+j,j)^2 * p_j(i) for some polynomial function p_j(i) of degree j. G.f. for column j (conjectured): Sum_{k=0..2*j} binomial(2*j,k)^2 * x^k / (1-x)^(3*j+1)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas J. Radley_, Nov 27 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015993 | OEIS-A358631

Irregular triangle T(n, k), n >= 0, k > 0, read by rows of extended (due to binary expansion of n) Stirling numbers of the first kind.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Nov 24 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015994 | OEIS-A358696

Number of self-avoiding closed paths in the 5 X n grid graph which pass through all vertices on four (left, right, upper, lower) sides of the graph.

Conjecture: a(n) = 6 *a(n-1) + a(n-2) - 20 * a(n-3) + 37 * a(n-4) + 16 * a(n-5) - 12 * a(n-6) - 7 * a(n-7) + 5 * a(n-8) - a(n-9) for n > 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Nov 27 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015995 | OEIS-A358704

Numbers m such that the sum of the prime divisors and the sum of the nonprime divisors of m^2+1 are both prime.

Conjecture: when b(n) tends to infinity, q/p tends to 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Nov 27 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015996 | OEIS-A358746

The number of vertices formed when every pair of n points, placed at the vertices of a regular n-gon, are connected by a circle and where the points lie at the ends of the circle's diameter.

Conjecture: for odd values of n all vertices are simple, other than those defining the diameters of the circles. No formula for n, or only the odd values of n, is currently known.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 30 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015997 | OEIS-A358820

a(n) is the least novel k such that d(k)|n, where d is the divisor counting function A000005.

Conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Dec 02 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015998 | OEIS-A358971

a(1) = 1. Thereafter a(n) is least novel k != n such that rad(k) = rad(n), where rad is A007947.

Conjectured to be a permutation of the positive integers with primes appearing in natural order. Primes are minima, 1 and primes squared are records.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Dec 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-015999 | OEIS-A358973

Numbers of the form m + omega(m) with m a positive integer.

Kucheriaviy proves that a(n) << n log log n and conjectures that a(n) ≍ n, that is, these numbers have positive lower density.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Dec 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016000 | OEIS-A358976

Numbers that are coprime to the sum of their factorial base digits (A034968).

The numbers of terms not exceeding 10^k, for k = 1, 2, ..., are 7, 59, 601, 6064, 60729, 607567, 6083420, 60827602, 607643918, 6079478119, ... Conjecture: The asymptotic density of this sequence exists and equals 6/Pi^2 = 0.607927... (A059956), the same as the density of A094387.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Dec 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016001 | OEIS-A358977

Numbers that are coprime to the sum of their primorial base digits (A276150).

The numbers of terms not exceeding 10^k, for k = 1, 2, ..., are 7, 59, 603, 6047, 60861, 608163, 6079048, 60789541, 607847981, 6080015681... . Conjecture: The asymptotic density of this sequence exists and equals 6/Pi^2 = 0.607927... (A059956), the same as the density of A094387.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Dec 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016002 | OEIS-A358978

Numbers that are coprime to the number of terms in their Zeckendorf representation (A007895).

The numbers of terms not exceeding 10^k, for k = 1, 2, ..., are 7, 61, 614, 6028, 61226, 606367, 6041106, 61235023, 612542436, 6034626175, 60093287082, 609082612171, ... Conjecture: The asymptotic density of this sequence exists and equals 6/Pi^2 = 0.607927... (A059956), the same as the density of A094387.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Dec 07 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016003 | OEIS-A359005

Jane Street's infinite sidewalk's greedy walk.

It is conjectured that the sequence actually visits all nonnegative integers. This is related to the conjecture that given any positive integer n, there exists a minimal Jane Street's path (i.e., with minimal number of jumps) going from slab 0 to slab n, and ending either in "forwards" or in "forwards-forwards-backwards". This would make the sequence a one-to-one mapping of the nonnegative numbers onto themselves. See also A359008.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frederic Ruget_, Dec 10 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016004 | OEIS-A359008

Jane Street's infinite sidewalk's greedy walk inverse mapping.

It is conjectured that a(n) exists for all nonnegative n. This has been verified true with a computer for all n < 7884654. This would make {a(n)} a one-to-one mapping from the nonnegative integers onto themselves.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frederic Ruget_, Dec 11 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016005 | OEIS-A359009

Irregular table read by rows: T(n,k) is the number of k-gons formed, k>=2, when every pair of n points, placed at the vertices of a regular n-gon, are connected by a circle and where the points lie at the ends of the circle's diameter.

Conjectures: for odd values of n all vertices are simple, other than those defining the diameters of the circles. For n > 2 and (n-2) mod 4 = 0, T(n,2) = n. For n mod 4 = 0, T(n,2) = k*n, k>=2. For odd n, T(n,2) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Dec 12 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016006 | OEIS-A359061

Irregular table read by rows: T(n,k) is the number of k-gons formed, k>=2, among all circles that can be constructed on vertices of an n-sided regular polygon, using only a compass.

Conjecture: the only value for n which leads to the creation of 2-gons is n = 2. Despite values for n mod 6 = 0 forming intersecting arcs at the center of the n-gon, these are cut by other circles and thus create 3-gons or 4-gons. This is in contrast to values of n mod 4 = 0 in A359009 which do lead to the creation of 2-gons at the center of the figure from similar arcs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Dec 14 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016007 | OEIS-A359064

a(n) is the number of trees of order n such that the number of eigenvalues of the Laplacian matrix in the interval [0, 1) is equal to ceiling((d + 1)/3) = A008620(d), where d is the diameter of the tree.

Conjecture from Guo et al.: lim_{n->oo} a(n)/A000055(n) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Dec 15 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016008 | OEIS-A359087

a(n) is equal to the last point of a reverse pyramid summation with base 1, 2, 3, ..., n-2, n-1, n, n-1, n-2, ..., 3, 2, 1.

a(n) = n*3^(n-1) - 2*A132894(n-1) (conjectured). - _Bernard Schott_, Dec 20 2022 Israel's and Schott's conjectures are true (see Fried link). - _Sela Fried_, Nov 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Moosa Nasir_, Dec 15 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016009 | OEIS-A359114

a(1) = 1; for n > 1, a(n) is the smallest positive number which has not appeared that shares a factor with the sum of the first n bits of the binary Champernowne string starting from 1.

For the binary Champernowne string starting from 1 see A030302. In the first 100000 terms the fixed points are 1, 2, 12, 24, 25; it is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Dec 16 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016010 | OEIS-A359143

The sum-and-erase sequence starting at 11: a(0) = 11; for n>=1, let m = a(n-1), and if m < 0, change m to an improper decimal "number" by replacing the minus sign by a single leading zero; then a(n) = A359142(m).

It is conjectured that every starting number will eventually enter a cycle or reach 0 (see A359142 for small examples).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 31 2023, based on suggestions from _Eric Angelini_ and _Hans Havermann_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016011 | OEIS-A359185

Numbers k such that for any positive integers x,y, if x*y=k then (x+y)^2+1 is a prime number.

Conjecture: if a term k is a perfect square > 1, then sqrt(k) is in the sequence A236068 (Primes p such that f(f(p)) is prime, where f(z) = z^2 + 1). The conjecture is false. A counterexample is 296147^2 = 87703045609 where 296147 = 47 * 6301. - _Robert Israel_, Mar 05 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Dec 19 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016012 | OEIS-A359195

Positive integers k with a smaller fraction of powers (mod k) than any smaller positive integers.

Conjecture: terms are products of primorials (A025487). A proof would greatly speed the search for more terms. On this conjecture, the next terms are 352800, 529200, 1058400, 2116800, 4233600, 6350400, 10584000, 19051200, 21168000, 31752000, 63504000, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Dec 19 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016013 | OEIS-A359199

Least prime p such that 2n can be written as a signed sum of p and the next 3 primes, or -1 if no such prime exists.

Conjecture: a satisfactory prime p exists for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Karl-Heinz Hofmann_ and _Peter Munn_, Jun 04 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016014 | OEIS-A359217

Y-coordinates of a point moving along a counterclockwise undulating spiral on a square grid.

Conjecture: a(n) = T10 + T15 + T20 + T21 where

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hans G. Oberlack_, Dec 21 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016015 | OEIS-A359256

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number which has not appeared such that all the distinct prime factors of a(n-1) + a(n) are factors of a(n).

The primes do not occur in their natural order, and for all terms studied if a(n) is a prime p, then a(n-1) = p(p-1) and a(n+1) = p(p^2-1). In the first 10000 terms the fixed points are 22, 165, 710, 1005, 9003, although it is likely more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _Eric Angelini_, Jan 05 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016016 | OEIS-A359262

a(n) is the largest number m such that prime(n)^m is in A359260.

Apparently, all the terms are of the form prime(k)-2 (A040976). Conjecture: The asymptotic density of the occurrences of prime(k)-2 is (1/s(k-1)-1/s(k)), where s(k) = A005867(k) = phi(prime(k)#), and prime(k)# is the k-th primorial number (A002110). The sums of the first 10^k terms, for k = 1, 2, ..., are 15, 221, 2291, 23287, 233641, 2337007, 23379901, 233814475, 2338211029, 23382168187, ... If the mentioned above conjecture is correct, then the asymptotic mean of this sequence is Sum_{k>=1} (prime(k)-2)*(1/s(k-1)-1/s(k)) = 2.33821872365981424748... . Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Dec 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016017 | OEIS-A359263

Number of compositions of n into three parts, using only natural numbers not in A007283.

Strictly increasing, and hence the complement of A007283 gives a counterexample to a conjecture of Dombi (see Links) saying that a set with infinite complement cannot have its sequence of 3-compositions strictly increasing in cardinality.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Dec 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016018 | OEIS-A359264

First differences of A359263.

Values of this sequence are positive, thus refuting a 2002 conjecture of Dombi discussed more fully in A359263.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Dec 23 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016019 | OEIS-A359319

Maximal coefficient of (1 + x) * (1 + x^8) * (1 + x^27) * ... * (1 + x^(n^3)).

Conjecture: Maximal coefficient of Product_{k=1..n} (1 + x^(n^m)) ~ sqrt(4*m + 2) * 2^n / (sqrt(Pi) * n^(m + 1/2)), for m>=0. - _Vaclav Kotesovec_, Dec 30 2022 Conjecture: a(n) ~ sqrt(14) * 2^n / (sqrt(Pi) * n^(7/2)). - _Vaclav Kotesovec_, Dec 30 2022

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Dec 25 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016020 | OEIS-A359349

The initial bits, written from left to right, in the 2-adic limit of the mod 2^e value of the odd factor of (2^e)!.

Here we refer to the table in the Example, in which the bits of our number, called stable bits, appear to the left of the space. Here is a conjectured relationship between the stable bits starting at any point and the unstable bits on the previous line of the table. There is a 2-adic integer K such that, for any d and n > d, uns(n,d) + K = stab(n+1,d) mod 2^d. Here uns(n,d) is the number whose backwards binary expansion (BBE) is the first d bits after the space on line n, and stab(n+1,d) is the number whose BBE is bits (n+1) to (n+d) of our number, where the numbering of the bits starts at 0. The BBE of K begins 1011011. For example, the BBE of uns(17,7) is 0101001 and that of stab(18,7) is 1110110. Perform the binary addition uns(17,7) + K from left to right on these. But computing h(m) mod 2^B can become time-consuming as m gets larger, because there are 2^(m-2) consecutive odd numbers to be multiplied. However, if the following conjecture holds, then there is a much faster way to compute those products mod 2^B, depending on the values of e and B. Conjecture: for all e,B such that 3 <= e < B <= 3*e - 5, (When d < e-1, the product on the right-hand side of the conjectured congruence requires only 1/2^(e-1-d) as many multiplications to compute, after which that product merely needs to be squared one or more times, with the residue mod 2^B taken after each squaring, so this can be much faster than taking the product on the left-hand side of the conjectured congruence.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Donald M Davis_, Jul 05 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016021 | OEIS-A359356

a(n-1) + a(n) has only digits also in a(n); lexicographically earliest such sequence of distinct nonnegative integers.

Conjectured to be a permutation of the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_ and _Eric Angelini_, Dec 27 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016022 | OEIS-A359387

Primes p such that the smallest prime factor of (2^(p-1)-1)/(3*p) is greater than p.

It is conjectured that there are infinitely many terms in this sequence, and their estimated asymptotic density n/a(n) ~ C/(log(a(n)))^2 where C is a constant between 0.7 and 0.9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Dec 29 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016023 | OEIS-A359402

Numbers whose binary expansion and reversed binary expansion have the same sum of positions of 1's, where positions in a sequence are read starting with 1 from the left.

Conjecture: Also numbers whose binary expansion has as least squares fit a line of zero slope, counted by A222955.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jan 05 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016024 | OEIS-A359403

Primes that are the concatenation of the first m consecutive k-th powers.

Let cp(k, n) denote the concatenation of the first n k-th powers. Note that there are no primes in cp(1, n) = A007908(n) up to n = 64000 or cp(3, n) = A019522(n) up to n = 31152. But there is a prime in cp(2, n) = A019521(n): cp(2, 3) = A019521(3) = 149. Conjecture: if k is odd, no "smaller primes" (< 10000 digits, or non-gigantic primes) exist in cp(k, n); if k is even, then such primes may occur. So far, the expanded search supports this conjecture. For k

from 5 to 500, when k is an odd number, we have found no "smaller primes" in cp(k, n); for even k, the following primes have been found: However, the next search contradicted the above conjecture. For k from 501 to 1100, we get two counterexamples:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Dec 30 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016025 | OEIS-A359416

Write n as 2^m - k, where 2^m is the least power of 2 >= n (0 <= k <= 2^(m-1)-1). For n a power of 2 (k = 0), a(n) = n. For numbers with k > 0, a(n) is the least p*a(k) which has not occurred previously, the count of k being taken from right to left (backwards) from k = 1 at 2^m - 1.

Conjectured to be a permutation of the positive integers in which primes appear in natural order. The even bisection, when divided by 2, reproduces the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Dec 30 2022 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016026 | OEIS-A359480

Number of Q-isomorphism classes of elliptic curves E/Q with good reduction away from 2 and prime(n).

R. von Känel and B. Matschke conjecture that a(n) <= a(2) = 752 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robin Visser_, Mar 31 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016027 | OEIS-A359482

Lexicographically earliest sequence of distinct terms > 0 such that the sum a(n) + a(n+1) is a substring of the concatenation (a(n), a(n+1)).

I conjecture that it isn't, and more specifically, that a(1) = 1 is the only single-digit term, and a(2) = 10 the only multiple of 10 below 100. See Examples for other terms of the form x*10^k, 1 <= x <= 9. - _M. F. Hasler_, Jul 03 2023 I have been informed that Wenlin Zhang and Haobo Ma have produced an AI assisted formal proof of the first part of conjecture, see the link. I have now also written my own proof of the full conjecture, cf. link to the OEIS wiki page. - _M. F. Hasler_, Sep 29 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Hans Havermann_, Jul 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016028 | OEIS-A359493

Numbers k such that the bottom entry in the ratio d(i)/d(i+1) triangle of the elements in the divisors of n, where d(1) < d(2) < ... < d(q) denote the divisors of k, is equal to 1.

Observation according to a comment by Bernard Schott: 216, 1000, 1728, 2744, 5832, ... are the first terms of A124581 (abundant cubes), hence the following conjecture: terms in A124581 do not belong to this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016029 | OEIS-A359494

Zeroless numbers k which can be written as a product of the powers whose base is a digit of k and whose exponent is a nonnegative integer.

As this sequence is a subsequence of A238985 which is conjectured to be finite, this sequence is conjectured to be finite. - _David A. Corneth_, Jan 28 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Jan 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016030 | OEIS-A359504

a(n) is calculated by considering in ascending order all products P of zero or more terms from {a(1..n-1)} until finding one where P+1 has a prime factor not in {a(1..n-1)}, in which case a(n) is the smallest such prime factor.

Conjecture: P+1 has at most one prime factor not already in the sequence, so the requirement of taking "the smallest such" is unnecessary.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joel Brennan_, Jan 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016031 | OEIS-A359505

a(1)=2, a(2)=3, and for n >= 3, a(n) is calculated by considering in ascending order all products P of (distinct) terms from {a(1..n-1)} until finding one where P-1 has a prime factor not in {a(1..n-1)}, in which case a(n) is the smallest such prime factor.

Conjecture: P-1 has at most one prime factor not already in the sequence, so the requirement of taking "the smallest such" is unnecessary (verified up to n=10000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joel Brennan_, Jan 24 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016032 | OEIS-A359506

a(n) is the least integer m such that there exists a strictly increasing integer sequence n = b_1 < b_2 < ... < b_t = m with the property that b_1 XOR b_2 XOR ... XOR b_t = 0.

For n > 1, a(n) >= n + 3. a(4n) = 4n + 3 for n > 0. Conjecture: a(n) <= 5(n+1)/3. - _Charles R Greathouse IV_, Jan 12 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Jan 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016033 | OEIS-A359507

a(n) is the least integer k such that there exists a strictly increasing integer sequence n = b_1 < b_2 < ... < b_t = n + k with the property that b_1 XOR b_2 XOR ... XOR b_t = 0.

Conjecture: a(n) is of the form 2^k + 1 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Jan 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016034 | OEIS-A359508

a(n) = log_2(A359507(n) - 1).

Conjecture: A359507(n) is always of the form 2^m + 1. Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Jan 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016035 | OEIS-A359509

a(n) is the number of subsets {b_1, b_2, ..., b_t} of {n, n+1, ..., A359506(n)} containing n with the property that b_1 XOR b_2 XOR ... XOR b_t = 0.

Conjectured formula: a(n) = 2^A000295(A359508(n)) for n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Jan 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016036 | OEIS-A359517

Inverse permutation (conjectured) of A359356.

Inverse permutation (conjectured) of A359356.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016037 | OEIS-A359558

a(n) is the first number that is the sum of 5 distinct positive Fibonacci numbers in exactly n ways (with a single type of 1).

Conjecture: there are no other terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Jan 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016038 | OEIS-A359559

a(n) is the determinant of an n X n Hermitian Toeplitz matrix whose first row consists of 1, 2*i, ..., n*i, where i denotes the imaginary unit.

Conjectured formulas: (Start) a(n) = ((n^2+(4+i)*n+4+4*i)/8)*(1+i)^n + ((n^2+(4-i)*n+4-4*i)/8)*(1-i)^n, from which the conjectures above follow (see Fried link). - _Sela Fried_, Jan 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jan 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016039 | OEIS-A359568

Maximum number of distinct folds after folding a square sheet of paper n times.

Conjecture: a(n) = 2*a(n-1) + 4 - n, for n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _S. Brunner_, Jan 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016040 | OEIS-A359628

Triangle read by rows: T(n,k) is the maximum number of connected endofunctions that are spanning subgraphs of a semi-regular loopless digraph on n vertices each with out-degree k.

Conjecture: T(n,n-2) = A000435(n-1)*(n-2) for n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yali Harrary_, Jan 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016041 | OEIS-A359643

a(n) = Sum_{k=0..n} binomial(n,k) * binomial(4*k,k).

Conjecture D-finite with recurrence +81*n*(3*n-1)*(3*n-2)*a(n) +3*(243*n^3-8433*n^2+14984*n-7064)*a(n-1) +2*(-58607*n^3+297306*n^2-491401*n+269124)*a(n-2) +6*(n-2)*(56663*n^2-237722*n+252221)*a(n-3) -3*(n-2)*(n-3)*(111625*n-286402)*a(n-4) +110653*(n-2)*(n-3)*(n-4)*a(n-5)=0. - _R. J. Mathar_, Jan 09 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jan 09 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016042 | OEIS-A359663

a(1) = 1; for n > 1, a(n) is the smallest positive number which has not appeared that shares a factor with the sum of the first n terms of the Champernowne string starting from 1.

For the Champernowne string starting from 1 see A033307. In the first 100000 terms there are 45 fixed points: 1, 4, 5, 6, 7, ..., 252, 264, 319. It is plausible no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jan 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016043 | OEIS-A359664

Prime Maze Room 11, opposite parity of A059459 starting from prime room 11.

Conjecture: Room 2 and Room 11 are unlinked, i.e., two separate mazes or branches/trees, as they are of opposite parities.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gregory Allen_, Jan 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016044 | OEIS-A359734

Lexicographically earliest sequence of distinct nonnegative integers such that the sequence A051699(a(n)) (distance from the nearest prime) has the same sequence of digits.

Conjectured to be a permutation of the nonnegative integers. The inverse permutation would start (3, 0, 2, 4, 9, 39, 10, 48, 20, 6, 1, 58, 21, 75, 25, 14, ...).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_ and _Eric Angelini_, Jan 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016045 | OEIS-A359736

Lexicographically earliest sequence of distinct nonnegative integers such that the sequence d(n) = dist(a(n), SQUARES) has the same sequence of digits.

Conjectured to be a permutation of the nonnegative integers. The inverse permutation would start (0, 2, 3, 11, 9, 12, 4, 7, 19, 21, 1, 8, 16, 20, 18, 24, ...)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_ and _Eric Angelini_, Jan 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016046 | OEIS-A359737

Lexicographically earliest sequence of distinct nonnegative integers such that the sequence d(n) = A296239(a(n)) has the same sequence of digits, where A296239 gives the distance from the nearest Fibonacci number, cf. A000045.

Conjectured to be a permutation of the nonnegative integers. The inverse permutation would start (0, 4, 25, 37, 3, 54, 6, 7, 104, 12, 2, 13, 1, 106, 20, ...).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_ and _Eric Angelini_, Jan 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016047 | OEIS-A359740

Maximal number of moves needed by a knight to reach every cell from a fixed position on an n X n X n chessboard, or -1 if it is not possible to reach every square.

The author conjectures that, for any n greater than 3, a(n) = n, while all the 10 independent cases has been manually checked for n = 5 and n = 6 (if n = 4, there are only 4 cases to be checked in order to show that a(4) = 4). a(n) = n for n >= 4 (conjectured by author; proved above).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Jan 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016048 | OEIS-A359795

Dirichlet inverse of function f(n) = 1 + A048675(n), where A048675(n) is fully additive with a(p) = 2^(1-PrimePi(p)).

Conjecture: the only odd term that occurs more than once is 1 = a(1) = a(4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 26 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016049 | OEIS-A359799

a(1) = 1, a(2) = 3; for n > 2, a(n) is the smallest positive number which has not appeared that shares a factor with |a(n-1) - a(n-2)| while the difference |a(n) - a(n-1)| is distinct from all previous differences |a(i) - a(i-1)|, i=2..n-1.

In the first 100000 terms the only fixed point is a(1) = 1; it is unknown if more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Mar 07 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016050 | OEIS-A359804

a(1) = 1, a(2) = 2; thereafter let p be the smallest prime that does not divide a(n-2)*a(n-1), then a(n) is the smallest multiple of p that is not yet in the sequence.

Conjecture: The primes appear in their natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Mar 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016051 | OEIS-A359807

a(1) = 0; thereafter a(n) is the largest a(i) + i which is < n among i = 1..n-1.

Conjectured run lengths are A272729, and the terms which occur here are partial sums of A272729.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Neal Gersh Tolunsky_, Jan 13 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016052 | OEIS-A359836

Parity of A353418, where A353418 is Dirichlet inverse of the characteristic function for numbers k where A156552(k) is a multiple of 3.

a(n) <= A353269(n). [Conjectured. Note that as A329609 is not a multiplicative semigroup, the proof cannot be similar to that given in A353348 for A359826. See also A359835]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 17 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016053 | OEIS-A359842

a(n) = Sum_{k=0..n} binomial(n*k,n+k).

Conjectures: a(2^k) == 0 (mod 2^(k-1)) and a(3^k) == 0 (mod 3^(k+2)) for k >= 2; a(p^k) == 0 (mod p^(k+1)) for all primes p >= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jan 15 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016054 | OEIS-A359858

a(0) = 0; for n > 0, a(n) is the smallest positive number not occurring earlier such that the ones' complement of the binary string of a(n-1) + a(n) does not appear in the binary string concatenation of a(0)..a(n-1).

In the first 100000 terms the fixed points are 55, 123, 1779, 2009, although it is likely more exist. The sequence is conjectured to be a permutation of the positive integers. Note that 1 does not appear until the 160th term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jan 16 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016055 | OEIS-A359859

Number of vertices among all distinct circles that can be constructed from a 2 x n square grid of points using only a compass.

A circle is constructed for every pair of the 2 x n points, the first point defines the circle's center while the second the radius distance. The number of distinct circles constructed from the 2 x n points is conjectured to be 4*A001859(n-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jan 16 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016056 | OEIS-A359868

a(n) is the smallest prime q such that A305411(n) + q is a square.

A305411 is conjectured to have only 103 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Can Güllü_, Jan 16 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016057 | OEIS-A359914

a(n) = coefficient of x^n in A(x) such that 2 = Sum_{n=-oo..+oo} (-1)^n * x^(n*(3*n-1)/2) * A(x)^(3*n) * (1 + x^n*A(x)).

Conjecture: a(n)/2 == A000041(n) (mod 2) for n >= 1; that is, a(n) is even, and a(n)/2 has the same parity as the number of partitions of n, for n >= 1. Conjecture: a(n) == A361552(n+1) (mod 4) for n >= 0. - _Paul D. Hanna_, Mar 19 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 23 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016058 | OEIS-A359945

Largest k < n such that n! / k! = m! = A000142(m) for some m.

Surányi conjectured (cf. Erdős and Habsieger references and A003135) that a(10) = 7 corresponding to 10! = 7! * 6! is the only nontrivial solution, i.e., other than a(n) = n-1 for n = m! and a(n) = 1 otherwise.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jan 19 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016059 | OEIS-A359984

Number of non-crossing antichain covers of {1,...,n} without singletons.

Conjecture D-finite with recurrence 8*n*(n-1)*a(n) -4*(n-1)*(56*n-145)*a(n-1) +4*(101*n^2-682*n+996)*a(n-2) +2*(6200*n^2-47903*n+88131)*a(n-3) +2*(26985*n^2-234056*n+491978)*a(n-4) +2*(62749*n^2-628865*n+1584314)*a(n-5) +(n-5)*(121577*n-667756)*a(n-6) +38285*(n-5)*(n-6)*a(n-7)=0. - _R. J. Mathar_, Mar 10 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Howroyd_, Jan 20 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016060 | OEIS-A360042

Number of vertices in a Farey fan of order n.

The number of vertices along each edge is A005728(n), while the number of regions is conjectured to equal A005598(n) = 1 + Sum_{i=1..n} (n-i+1)*phi(i). The regions count the number of distinct approximate representations of straight lines y = mx + b that can be drawn on an x-y integer raster, where x, y, and b are restricted to [0,n) and 0 <= m <=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, _N. J. A. Sloane_ and _M. Douglas McIlroy_, Jan 23 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016061 | OEIS-A360053

Primes p such that each prime < p in the prime factorization of 2^(p-1) - 1 has exponent 1.

Conjecture: 2^(a(n)-1) - 1 is always squarefree.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Jan 23 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016062 | OEIS-A360068

Number of integer partitions of n such that the parts have the same mean as the multiplicities.

Conjecture: If n is squarefree, then a(n) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jan 27 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016063 | OEIS-A360070

Numbers for which there exists an integer partition such that the parts have the same mean as the multiplicities.

Conjecture: No term > 1 is squarefree.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jan 27 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016064 | OEIS-A360105

Numbers k such that sigma_2(k^2 + 1) == 0 (mod k).

Conjecture: the sequence contains infinitely many Fibonacci numbers (see A360107).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 26 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016065 | OEIS-A360116

a(n) = 1 if there are no prime factors p of n for which the 2-adic valuation of p-1 is less than that of n-1, otherwise 0.

a(n) = [A333570(n) <= 2] = [A360112(n) <= 2]. (Conjectured by Sequence Machine) - _Antti Karttunen_, May 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016066 | OEIS-A360145

Triangle read by rows where row n is the largest (or middle or n-th) column of the reverse pyramid summation of order n described in A359087.

T(n,k) = 3^(k-1)*n - 2*A132894(k-1) for 1 <= k <= n (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bernard Schott_, Jan 27 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016067 | OEIS-A360204

Primitive prime powers. p is a primitive prime power iff it is an odd prime power that exceeds the preceding odd prime power by more than any smaller odd prime power does. ('Prime power' defined in the sense of A246655.)

Conjecture: Primitive prime powers are primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Feb 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016068 | OEIS-A360209

Lexicographically earliest infinite sequence of distinct positive numbers such that, for n > 2, a(n) shares a factor with a(n-2) + a(n-1) but shares no factor with a(n-2).

In the first 100000 terms the fixed points for n > 2 are 3, 6, 441, 1677, 3629, 9701, 17131, although it is likely more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jan 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016069 | OEIS-A360278

Determinant of the matrix [L(j+k) + d(j,k)]_{1<=j, k<=n}, where L(n) denotes the Lucas number A000032(n), and d(j,k) is 1 or 0 according as j = k or not.

Conjecture 1: Let v(0) = 2, v(1) = A, and v(n+1) = A*v(n) + v(n-1) for n > 0. Then A^2*det[v(j+k) + d(j,k)]_{1<=j, k<=n} = v(n+1)^2 - (A^2 + 4)*(n mod 2) for any positive integer n. In particular, a(n) = L(n+1)^2 - 5*(n mod 2) for all n > 0. Conjecture 2: Let v(0) = 2, v(1) = A, and v(n+1) = A*v(n) - v(n-1) for n > 0. Then det[v(j+k) + d(j,k)]_{1<=j, k<=n} = u(n+1)^2 - n^2 for any positive integer n, where u(0) = 0, u(1) = 1, and u(n+1) = A*u(n) - u(n-1) for all n > 0. Conjecture 3: Let F(n) denote the Fibonacci number A000045(n). Then, for any positive integer n, we have det[F(j+k) + d(j,k)]_{1<=j, k<=n} = F(n+1)^2 + (n mod 2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Feb 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016070 | OEIS-A360374

Indices of the nonprimitive rows of the Wythoff array (A035513); see Comments.

In a row of the Wythoff array, either every two consecutive terms are relatively prime or else no two consecutive terms are relatively prime. In the first case, we call the row primitive, as in A332938. Otherwise, the row is an integer multiple of a tail of a preceding row, and we call the row nonprimitive. Conjecture: the limiting proportion of nonprimitive rows exists and is approximately 0.392.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 04 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016071 | OEIS-A360380

a(n) = number of the diagonal of the Wythoff array, A035513, that includes prime(n). See Comments.

The indexing of diagonals is given in A191360. Conjecture: every integer occurs infinitely many times in this sequence; i.e., every diagonal includes infinitely many primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 07 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016072 | OEIS-A360418

Numbers k such that, in a listing of all congruence classes of positive integers, the k-th congruence class contains k. Here the class r mod m (with r in {1,...,m}) precedes the class a' mod b' (with r' in {1,...,m'}) iff m < m' or r > r'.

Above conjecture confirmed with more terms and linear recurrence. See supporting formula below. - _Ray Chandler_, Feb 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James Propp_, Feb 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016073 | OEIS-A360425

Indices of records in A018804.

Conjecture: a(n) ~ exp(4*(n-1)/21). - _Matthew Russell Downey_, Jul 25 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthew Russell Downey_, Feb 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016074 | OEIS-A360519

Let C consist of 1 together with all numbers with at least two distinct prime factors; this is the lexicographically earliest infinite sequence {a(n)} of distinct elements of C such that, for n>2, a(n) has a common factor with a(n-1) but not with a(n-2).

Conjecture: The sequence is a permutation of C.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Mar 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016075 | OEIS-A360521

a(0) = 0; for n > 0, a(n) is the smallest positive number not occurring earlier such that neither the binary string a(n-1) + a(n) nor the same string reversed appear in the binary string concatenation of a(0)..a(n-1).

As in A357082 the main concentration of terms is along the line a(n) = n, so there are numerous fixed points - there are 24 fixed points in the first 200000 terms. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Feb 09 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016076 | OEIS-A360533

a(n) = index of the diagonal of the natural number array, A000027, that includes prime(n). See Comments.

The indexing of diagonals is given in A191360. Conjecture: Every odd-indexed diagonal contains infinitely many primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Feb 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016077 | OEIS-A360549

a(n) is the least prime p not already in the sequence such that a(n-1) + p is a triprime; a(1) = 2.

Conjecture: every prime eventually appears in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_ and _Robert Israel_, Mar 04 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016078 | OEIS-A360578

Expansion of g.f. A(x) satisfying A(x) = Series_Reversion(x - x*A'(x)*A(x)).

a(n) ~ c * n! * n^alfa / LambertW(1)^n, where alfa = 1.5447806483693... and c = 0.028888886141962894 96..., conjecture: alfa = 2*(2*LambertW(1) - 1 + 1/(1 + LambertW(1))). - _Vaclav Kotesovec_, Feb 22 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016079 | OEIS-A360581

Expansion of A(x) satisfying [x^n] A(x)^n / (1 + x*A(x)^n)^n = 0 for n > 0.

Conjecture: alpha = 3*LambertW(1) - 2 + 1/(1 + LambertW(1)) = 0.33953361459446... (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016080 | OEIS-A360587

a(n) is the least positive integer k such that k*(k+1)*...*(k+n-1) does not contain the digit 2, or -1 if there is no such k.

Conjecture: a(n) = -1 for n = 31 and all n >= 33.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Feb 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016081 | OEIS-A360589

Numbers k that set records in A355432.

Conjecture: for n > 2, subset of A364702. - _Michael De Vlieger_, Oct 04 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Feb 22 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016082 | OEIS-A360623

Largest k such that the decimal representation of 2^k is missing any n-digit string.

All terms are conjectural, checked up to 10^6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hans Havermann_, Feb 14 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016083 | OEIS-A360633

Square array A(n, k), n, k > 0, read by antidiagonals upwards; A(n, k) = A360613(2*n-1) * A360613(2*k).

Conjecture: all integers appear in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Feb 14 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016084 | OEIS-A360635

a(n) is the smallest nonnegative integer that satisfies p(a(n)+1) - p(a(n)) >= n, where p denotes the number partition function.

Conjecture: except for the values 2, 4 and 6, this sequence includes all the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Feb 14 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016085 | OEIS-A360753

Matrix inverse of A360657.

Conjectured formulas: All three conjectures are true (see Fried link in A360657). - _Sela Fried_, Aug 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Feb 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016086 | OEIS-A360807

Decimal expansion of Sum_{m>=1} 1/(1/4 + z(m)^2) where z(m) is the imaginary part of the m-th nontrivial zero of the Dirichlet beta function whose real part is 1/2.

Conjecture: Nontrivial zeros whose real part is not 1/2 do not exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Feb 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016087 | OEIS-A360821

Number of primes of the form k^2+1 between n^2 and 2*n^2 exclusive.

Conjecture: Except for n = 7, between the square of an integer n > 1 and the double of its square there is always at least one prime number of the form k^2+1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Feb 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016088 | OEIS-A360822

Numbers whose squares have at most 2 digits less than 8.

Conjecture: Sequence has 72 terms, with largest term 940206833.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Feb 22 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016089 | OEIS-A360917

Array read by antidiagonals: T(m,n) is the number of vertices in the longest induced path in the grid graph P_m X P_n.

T(4,n) = A047408(n+1) - [n = 4] (conjectured), with Iverson bracket [.] = 1 if true, 0 else. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Howroyd_, Feb 26 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016090 | OEIS-A360921

Maximum number of vertices in an induced tree in the n X n grid graph.

Conjecture: a(n) = n^2 - A354673(n). - _Andrew Howroyd_, Jul 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Howroyd_, Feb 26 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016091 | OEIS-A360923

Table T(i,j), i >= 0, j >= 0, read by antidiagonals giving the smallest number of moves needed to win Integer Lunar Lander, starting from position (i,j). Game rules in comments.

Conjecture 1: For all i, j, T(i,j) differs from its eight neighbors by at most 3. The array of differences {T(i+1,j) - T(i,j)} might be worth studying, as well as the array {T(i,j) mod 2}. - _N. J. A. Sloane_, Feb 25 2023 Conjecture 2 (Start) Conjecture 1 seems to hold for all (i,j) except (1,0) and (2,1). - _M. F. Hasler_, Feb 27 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Allan C. Wechsler_, Feb 25 2023, using information about the original game from _Eric Angelini_, _Hans Havermann_, and _M. F. Hasler_, and data from _Tom Karzes_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016092 | OEIS-A360925

Smallest number of moves needed to win Integer Lunar Lander from starting position (n,0).

Conjecture: For n >= 2, a(n) = n+1+floor(sqrt(2*n^2-2*n-3)). - _N. J. A. Sloane_, Feb 26 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Allan C. Wechsler_, Feb 25 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016093 | OEIS-A360926

Smallest number of moves needed to win Integer Lunar Lander with a starting position of (n,n).

The conjectured formula for A360923 implies a formula for a(n). - _N. J. A. Sloane_, Feb 26 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Allan C. Wechsler_, Feb 25 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016094 | OEIS-A360931

a(1) = 2, a(2) = 3; for n > 2, a(n) is the smallest number greater than 1 that has not appeared such that |a(n) - a(n-1)| has a common factor with a(n-2).

In the first 100000 terms the fixed points are 10, 16, 42, 52, 66; it is likely no more exist. The sequence is conjectured to be a permutation of the positive integers > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Feb 25 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016095 | OEIS-A361034

a(n) = 2520*(4*n)!/(n!*(n+2)!^3).

Conjecture: a(n) is odd iff n = 2^k - 2 for some k >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016096 | OEIS-A361035

a(n) = 9979200 * (4*n)!/(n!*(n+3)!^3).

Conjecture: a(n) is odd iff n = 2^k - 3 for some k >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016097 | OEIS-A361036

a(n) = n! * [x^n] (1 + x)^n * exp(x*(1 + x)^n).

We conjecture that a(n+k) == a(n) (mod k) for all n and k. If true, then for each k, the sequence a(n) taken modulo k is a periodic sequence and the period divides k. For example, modulo 7 the sequence becomes [1, 2, 4, 5, 6, 0, 4, 1, 2, 4, 5, 6, 0, 4, 1, 2, 4, 5, 6, 0, 4, ...], apparently a periodic sequence of period 7. More generally, let F(x) and G(x) denote power series with integer coefficients with F(0) = G(0) = 1. Define b(n) = n! * [x^n] exp(x*G(x)^n)*F(x)^n. Then we conjecture that b(n+k) == b(n) (mod k) for all n and k. The conjectures above are true, see Turazashvili links. - _Nikoloz Turazashvili_, Aug 22 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 13 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016098 | OEIS-A361047

Expansion of g.f. A(x) satisfying A(x) = Series_Reversion(x - x^3*A'(x)^2).

Conjecture: a(n) == 1 (mod 3) iff n = (3^k - 1)/2 for k >= 0, otherwise a(n) == 0 (mod 3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016099 | OEIS-A361077

a(n) = largest sqrt(2*n)-smooth divisor of binomial(2*n, n).

Binomial(2*n, n)/a(n) and A263931(n)/a(n) are coprime with a(n) and squarefree. By the Erdős squarefree conjecture, proved in 1996, no a(n) with n >= 5 is squarefree.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthieu Pluntz_, Mar 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016100 | OEIS-A361087

Maximum squared inverse distance from the origin to the hyperplane defined by hypercube points.

Because it is not known if the squared inverse distances are always integers, this is a sequence of numerators (and the known corresponding denominators are all 1). The next two terms are conjectured to be 59 and 203 (distinct from sequences already in the OEIS).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Veit Elser_, Jun 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016101 | OEIS-A361103

a(n) = k such that A360519(k) = A361102(n), or -1 if A361102(n) never appears in A360519.

Conjectured to be a permutation of the natural numbers (and if so, -1 will never appear). See A361104 for the putative inverse permutation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Mar 02 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016102 | OEIS-A361127

Let p = n-th odd prime; a(n) = index where 2*p appears in A360519, or -1 if 2*p never appears.

It is conjectured that every 2*prime(n) (n>1) appears in A360519. A proof of this would be a big step towards proving that every term of C appears in A360519.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Mar 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016103 | OEIS-A361132

Multiplicative with a(p^e) = e^4, p prime and e > 0.

Conjecture: log(log(c(k)))/k converges to a constant (around 0.315).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Mar 02 2023, following a suggestion from _Amiram Eldar_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016104 | OEIS-A361133

a(n) = n for n <= 3. Let h, i, j represent a(n-3), a(n-2), a(n-1) respectively. For n > 3, if there is a symmetric difference in the sets of distinct primes dividing h and j, with greatest member p then a(n) is the least novel multiple of p. Otherwise, a(n) is the least novel k such that (k,i) > 1.

Conjectured to be a permutation of the natural numbers, with primes in natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Mar 02 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016105 | OEIS-A361190

Number of 4n-step lattice paths starting and ending at (0,0) that do not go above the diagonal x=y or below the x-axis using steps in {(1,1), (1,-1), (-1,0)}.

Conjecture: alpha = 1 + Pi/arctan(sqrt(2)) = 1 + Pi/A195696. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 31 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016106 | OEIS-A361197

a(n) is the number of equations in the set {x^2 + 2y^2 = n, 2x^2 + 3y^2 = n, ..., k*x^2 + (k+1)*y^2 = n, ..., n*x^2 + (n+1)*y^2 = n} which admit at least one nonnegative integer solution.

Conjecture: a(n) = 2 for infinitely many n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luca Onnis_, Mar 04 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016107 | OEIS-A361281

a(n) = n! * Sum_{k=0..n} binomial(n*k,n-k)/k!.

It appears that a(n) == 1 (mod 4) and a(5*n+2) == 0 (mod 5) for all n. More generally we conjecture that a(n+k) == a(n) (mod k) for all n and k. If true, then for each k, the sequence a(n) taken modulo k is a periodic sequence and the period divides k. Let F(x) and G(x) be power series with integer coefficients with G(0) = 1. Define b(n) = n! * [x^n] F(x)*exp(x*G(x)^n). Then we conjecture that b(n+k) == b(n) (mod k) for all n and k. The present sequence is the case F(x) = 1, G(x) = 1 + x. Cf. A278070. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Mar 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016108 | OEIS-A361301

For the odd number 2n + 1, the least primitive practical number r such that 2n + 1 = r + p where p is prime.

Conjecture: every odd number, beginning with 3, is the sum of a prime number and a primitive practical number. This is a tighter conjecture than that posed by _Hal M. Switkay_ (see comments of A005153).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Mar 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016109 | OEIS-A361314

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number which has not appeared that shares a factor with a(n-2) + a(n-1) while the sum a(n) + a(n-1) is distinct from all previous sums a(i) + a(i-1), i=2..n-1.

In the first 100000 terms the fixed points are 1, 2, 3, 6, 9, 10, 39, 91, 112; it is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Mar 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016110 | OEIS-A361321

Lexicographically earliest infinite sequence of distinct elements of A000469 such that, for n > 2, a(n) has a common factor with a(n-1) but not with a(n-2).

Conjecture: The sequence is a permutation of A000469.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, _Rémy Sigrist_ and _N. J. A. Sloane_, Mar 09 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016111 | OEIS-A361323

a(n) = k such that A000469(k) = A361321(n).

Conjectured to be a permutation of the natural numbers (if not then there exists a k such that A000469(k) does not appear in A361321).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, _Rémy Sigrist_, and _N. J. A. Sloane_, Mar 11 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016112 | OEIS-A361324

a(n) = k such that A361321(k) = A000469(n), or -1 if A000469(n) never appears in A361321.

Conjectured to be a permutation of the natural numbers (and if so, -1 will never appear).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, _Rémy Sigrist_, and _N. J. A. Sloane_, Mar 11 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016113 | OEIS-A361337

Numbers that reach 0 after a suitable series of split-and-multiply operations (see Comments for precise definition).

a(2894 + k) = 3112 + k for all k >= 0 (conjectured). - _M. F. Hasler_, Apr 05 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 01 2023, based on a posting to the Sequence Fans mailing list by _Eric Angelini_, Mar 20 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016114 | OEIS-A361430

Multiplicative with a(p^e) = e - 1.

Conjecture: a(n) = abs(A298826(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Mar 11 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016115 | OEIS-A361460

a(n) = 1 if A135504(n+1) = 2 * A135504(n), otherwise 0.

a(n) = 1 if A135506(n) is not prime, otherwise 0. [Conjectured]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 13 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016116 | OEIS-A361461

Numbers k such that x(k+1) = 2 * x(k), when x(1)=1 and x(n) = x(n-1) + lcm(x(n-1),n), i.e., x(n) = A135504(n).

It is conjectured that this sequence gives also all numbers k for which A135506(k) is not a prime. See Ruiz-Cabello paper.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Mar 13 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016117 | OEIS-A361504

Index of n in A359804, or -1 if n never appears there.

Conjectured to be a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 18 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016118 | OEIS-A361511

a(1) = 1. Thereafter if a(n-1) is a novel term, a(n) = d(a(n-1)); otherwise, if a(n-1) is the t-th non-novel term, a(n) = a(n-1) + d(a(t)), where d is the divisor function A000005.

It is an obvious conjecture that every number eventually appears, but is there a proof?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016119 | OEIS-A361513

Novel terms in A361511, in order of appearance.

By definition the terms are distinct. It is an obvious conjecture that every number eventually appears, but is there a proof?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016120 | OEIS-A361515

a(n) = index of first appearance of n in A361511, or -1 if n never appears.

Conjecture: This is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016121 | OEIS-A361534

Let h,i,j be the latest 3 terms in the sequence, starting with a(1)=1, a(2)=2, a(3)=3. Let R = rad(h*i*j), where rad is A007947, and let S be the smallest number of terms in U = {h,i,j} which are divisible by any prime p dividing R. Then, a(n) is the least novel multiple of the greatest such prime p.

The sequence is conjectured to be a permutation of the positive integers with the primes appearing in their natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Mar 15 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016122 | OEIS-A361552

Expansion of g.f. A(x) satisfying 2*x = Sum_{n=-oo..+oo} x^(n*(3*n+1)/2) * (A(x)^(3*n) - 1/A(x)^(3*n+1)).

Conjecture: a(n) == A359914(n-1) (mod 4) for n > 1. Conjecture: a(n)/2 == A000041(n-1) (mod 2) for n >= 1; that is, a(n) is even, and a(n)/2 has the same parity as the number of partitions of n-1, for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 19 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016123 | OEIS-A361627

Positive integers such that GCD(A007504(n),n) != 1.

Conjecture: a(n) ~ C*n as n -> infinity, where 5.25 < C < 5.35.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luca Onnis_, Mar 18 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016124 | OEIS-A361629

For n <= 2, a(n) = n. Thereafter let p be the greatest prime which divides the least number of terms in U = {a(n-2), a(n-1)}, then a(n) is the smallest multiple of p that is not yet in the sequence.

Conjectured to be a permutation of the positive integers with primes in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Mar 18 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016125 | OEIS-A361683

a(n) is the least k such that tau(k) divides sigma_n(k) but not sigma(k), or -1 if no such k exists.

Conjecture: a(n) is of the form 2^b*p1^c*p2^d*...*pk^j with b > 0 and A020639(n) divides b*(c+1)*(d+1)*...*(j+1). (p1, p2, ..., pk are distinct odd prime numbers). (End) Conjecture: For primes q > p, a(q) > a(p). If true, we could replace "<=" with "=" in the above formula. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mohammed Yaseen_, Mar 20 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016126 | OEIS-A361710

a(n) = Sum_{k = 0..n-1} (-1)^k*binomial(n,k)*binomial(n-1,k)^2.

Conjecture: the supercongruence a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) holds for all primes p >= 5 and positive integers n and k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016127 | OEIS-A361711

a(1) = 1 and a(n) = Sum_{k = 0..n-2} (-1)^k * binomial(n,k)^2 * binomial(n-2,k) for n >= 2.

Conjecture: the supercongruence a(p^k) == a(p^(k-1)) (mod p^(3*k)) holds for all primes p >= 5 and positive integer k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016128 | OEIS-A361712

a(n) = Sum_{k = 0..n-1} binomial(n,k)^2*binomial(n+k,k)*binomial(n+k-1,k).

Conjecture 1: the supercongruence a(p) == a(1) (mod p^5) holds for all primes p >= 7 (checked up to p = 199). Conjecture 2: for r >= 2, the supercongruence a(p^r) == a(p^(r-1)) (mod p^(3*r+3)) holds for all primes p >= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016129 | OEIS-A361713

a(n) = Sum_{k = 0..n-1} binomial(n,k)^2 * binomial(n+k-1,k)^2.

Conjecture 1: the supercongruence a(p) == a(1) (mod p^5) holds for all primes p >= 7 (checked up to p = 199). Conjecture 2: for r >= 2, the supercongruence a(p^r) == a(p^(r-1)) (mod p^(4*r+1)) holds for all primes p >= 7. [added Aug 12 2026: The first conjecture is correct. The second conjecture is not correct as it stands - Thomas Adamczewski notes that a(7^6) - a(7^5) is divisible by 7^24 but not by 7^25.] Adamczewski disproves Conjecture 2 in linked article. - _Michael De Vlieger_, Aug 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016130 | OEIS-A361714

a(n) = Sum_{k = 0..n-1} (-1)^(n+k+1)*binomial(n,k)*binomial(n+k-1,k)^2.

Conjecture 1: the supercongruence a(p) == a(1) (mod p^5) holds for all primes p >= 7 (checked up to p = 199). Conjecture 2: for r >= 2, the supercongruence a(p^r) == a(p^(r-1)) (mod p^(3*r+3)) holds for all primes p >= 7. Conjectures 1 and 2 are true: a(p) == a(1) (mod p^5) and a(p^r) == a(p^(r-1)) (mod p^(3*r+3)) for r >= 2 and all primes p >= 7 (see Theorem 1 in the Blanco Gómez link). - _Roberto Blanco Gómez_, Oct 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016131 | OEIS-A361715

a(n) = Sum_{k = 0..n-1} binomial(n,k)^2*binomial(n+k-1,k).

Conjecture 1: the supercongruence a(p) == a(1) (mod p^5) holds for all primes p >= 7 (checked up to p = 199) [added Aug 15 2026: this is correct - see the Bala link.]. Conjecture 2: for r >= 2, the supercongruence a(p^r) == a(p^(r-1)) (mod p^(3*r+3)) holds for all primes p >= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 23 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016132 | OEIS-A361717

a(n) = Sum_{k = 0..n-1} binomial(n-1,k)^2*binomial(n+k,k).

Conjecture 1: the supercongruence a(p) == 0 (mod p^4) holds for all primes p >= 5 (checked up to p = 199). Conjecture 2: the supercongruence a(p-1) == 1 - 2*p - p^2 (mod p^3) holds for all primes except p = 3 (checked up to p = 199).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 26 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016133 | OEIS-A361747

Lexicographically earliest sequence of distinct positive integers such that a(n) and a(n-1) share at least one identical trit at the same position in their balanced ternary representations.

Conjecture: Suppose n is the x-th fixed point such that n-1 is not a fixed point. Then for all k such that n < k < (3^(x + 2) + 1)/2, k is also a fixed point.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jodi Spitz_, Mar 22 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016134 | OEIS-A361825

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number that has not yet appeared that is a multiple of the smallest prime that does not divide a(n-2) + a(n-1).

The sequence is conjectured to be a permutation of the positive integers, although it takes many terms for the primes to appear, e.g., a(191443) = 19.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Mar 25 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016135 | OEIS-A361826

a(n) is equal to the number of roots of the equation n*cos(x) = sqrt(x).

Conjecture: a(n) = 2*floor(n^2/(2*Pi)) + 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nicolay Avilov_, Mar 27 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016136 | OEIS-A361883

a(n) = (1/n) * Sum_{k = 0..n} (n+2*k) * binomial(n+k-1,k)^3.

The central binomial coefficients u(n) := binomial(2*n,n) = A000984(n) satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for positive integers n and r and all primes p >= 5. We conjecture that the present sequence satisfies the same congruences. Adamczewski includes a proof abstract of the above conjecture in linked article. - _Michael De Vlieger_, Aug 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016137 | OEIS-A361884

a(n) = (1/n) * Sum_{k = 0..n} (-1)^(n+k) * (n + 2*k) * binomial(n+k-1,k)^3.

The central binomial coefficients u(n) := binomial(2*n,n) = A000984(n) satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for positive integers n and r and all primes p >= 5. We conjecture that the present sequence satisfies the same congruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016138 | OEIS-A361885

a(n) = (1/n) * Sum_{k = 0..2*n} (n+2*k) * binomial(n+k-1,k)^3.

The binomial coefficients u(n) := binomial(3*n,n) = A005809(n) satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for positive integers n and r and all primes p >= 5. We conjecture that the present sequence satisfies the same congruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016139 | OEIS-A361887

a(n) = S(5,n), where S(r,n) = Sum_{k = 0..floor(n/2)} (binomial(n,k) - binomial(n,k-1))^r.

For r a positive integer define S(r,n) = Sum_{k = 0..floor(n/2)} (binomial(n,k) - binomial(n,k-1))^r. The present sequence is {S(5,n)}. Gould (1974) conjectured that S(3,n) was always divisible by S(1,n). See A183069 for {S(3,n)/S(1,n)}. In fact, calculation suggests that if r is odd then S(r,n) is always divisible by S(1,n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016140 | OEIS-A361888

a(n) = S(5,n)/S(1,n), where S(r,n) = Sum_{k = 0..floor(n/2)} (binomial(n,k) - binomial(n,k-1))^r.

For r a positive integer define S(r,n) = Sum_{k = 0..floor(n/2)} (binomial(n,k) - binomial(n,k-1))^r. Gould (1974) conjectured that S(3,n) was always divisible by S(1,n). See A183069 for {S(3,n)/S(1,n)}. In fact, calculation suggests that if r is odd then S(r,n) is always divisible by S(1,n). The present sequence is {S(5,n)/S(1,n)}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016141 | OEIS-A361889

a(n) = S(5,2*n-1)/S(1,2*n-1), where S(r,n) = Sum_{k = 0..floor(n/2)} (binomial(n,k) - binomial(n,k-1))^r.

Conjecture: the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) holds for positive integers n and r and all primes p >= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016142 | OEIS-A361890

a(n) = S(7,n), where S(r,n) = Sum_{k = 0..floor(n/2)} (binomial(n,k) - binomial(n,k-1))^r.

For r a positive integer define S(r,n) = Sum_{k = 0..floor(n/2)} (binomial(n,k) - binomial(n,k-1))^r. The present sequence is {S(7,n)}. Gould (1974) conjectured that S(3,n) was always divisible by S(1,n). See A183069 for {S(3,n)/S(1,n)}. In fact, calculation suggests that if r is odd then S(r,n) is always divisible by S(1,n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 30 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016143 | OEIS-A361891

a(n) = S(7,n)/S(1,n), where S(r,n) = Sum_{k = 0..floor(n/2)} (binomial(n,k) - binomial(n,k-1))^r.

For r a positive integer define S(r,n) = Sum_{k = 0..floor(n/2)} (binomial(n,k) - binomial(n,k-1))^r. Gould (1974) conjectured that S(3,n) was always divisible by S(1,n). See A183069 for {S(3,n)/S(1,n)}. In fact, calculation suggests that if r is odd then S(r,n) is always divisible by S(1,n). The present sequence is {S(7,n)/S(1,n)}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 30 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016144 | OEIS-A361904

Odd numbers k such that for all even divisors d of k^2+1, d^2+1 is a prime number.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Mar 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016145 | OEIS-A361919

The number of primes > A000040(n) and <= (A000040(n)^c + 1)^(1/c), where c = 0.567148130202... is defined in A038458.

Let c = 0.567148130202... (see A038458), the solution to 127^x - 113^x = 1. c is conjectured by Smarandache to be the smallest real number x such that A000040(n+1)^x - A000040(n)^x = 1 has a solution. This conjecture is equivalent to saying that the terms of the present sequence are always positive, but that if c were replaced by a larger real number, there would be zeros in the sequence. However, note that a(30) is not the last occurrence of 1: a(46) = a(61) = 1 as well.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hal M. Switkay_, Mar 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016146 | OEIS-A361929

a(1) = 2; for n > 1, a(n) is the smallest positive integer > 1 not to share a factor with terms a(n-c .. n-1) where c = gcd(n-1,a(n-1)).

Conjecture: All primes will occur. See A361931 for the first occurrences of primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Samuel Harkness_, Mar 30 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016147 | OEIS-A361930

a(n) is the greatest prime p such that p + q^2 + r^3 = prime(n)^4 for some primes q and r.

Conjecture: a(n) exists for all n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_ and _Robert Israel_, Mar 30 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016148 | OEIS-A361931

a(n) is the smallest number k such that A361929(k) = prime(n), or -1 if no such prime exists.

It is conjectured that all primes appear in A361929.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Samuel Harkness_, Mar 30 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016149 | OEIS-A361961

Total semiperimeter of 3-Fuss-Catalan polyominoes of length 3n.

Conjecture: D-finite with recurrence 3*n*(396221*n -410120) *(3*n-1) *(3*n+1) *a(n) +4*(-86981513*n^4 +457143117*n^3 -996839467*n^2 +906061905*n -279161658) *a(n-1) +32*(2*n-5) *(4*n-9) *(4*n-7) *(2282347*n -1795413)*a(n-2)=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Mar 31 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016150 | OEIS-A362001

Numbers k such that the digits of k^2 are a subsequence of the digits of 2^k.

Conjecture is true up to 10^6. - _Michael S. Branicky_, Apr 03 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Apr 02 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016151 | OEIS-A362006

a(n) is the minimum integer m such that floor(e^n) = floor(Sum_{k=0..m} (n^k)/(k!)).

Conjecture: a(n) ~ e*n as n->infinity. Conjecture: a(n) <= 3n for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luca Onnis_, Apr 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016152 | OEIS-A362017

a(n) is the leading prime in the n-th prime sublist defined in A348168.

If Conjecture 2 in A348168 is true, lim_{n->infinity} a(n)/prime(round((n-1)*e)+1) = 1, where e is Euler's number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Apr 04 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016153 | OEIS-A362018

Numbers k such that the digits of k^2 do not form a subsequence of the digits of 2^k.

Complement of A362001. It was conjectured that a(140)=327 is the last term (see A362001).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Apr 04 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016154 | OEIS-A362032

Lexicographically least counterexample to Dombi's conjecture.

Lexicographically least counterexample to Dombi's conjecture. Dombi's conjecture, now refuted, stated that there is no infinite set A such that the power series (sum(X^a, a>=0 not in A))^3 has strictly increasing coefficients. This sequence is conjecturally the lexicographically least such example---"conjecturally" because it is obtained by backtracking and there is currently no proof it can be extended to infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Apr 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016155 | OEIS-A362065

Write the English name of a(n); sum the ranks of its letters in the alphabet; divide the sum by a(n); the result is an integer.

Indices of zeros in A226911; and conjectured finite, full based on an observation there. - _Michael S. Branicky_, Apr 07 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and Mensanator, Apr 07 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016156 | OEIS-A362069

Numbers k such that k+digitsum(k^2) is a square.

Conjecture: there are infinitely many pairs of consecutive terms. Example: (296,297); (413,414); (863,864).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexandru Petrescu_, May 17 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016157 | OEIS-A362086

Denominator of the continued fraction 1/(2-3/(3-4/(4-5/(...(n-1)-n/(-3))))).

Conjecture: Except for 9, every term of this sequence is either a prime or 1. Conjecture: Record values correspond to A248697 (n>3). - _Bill McEachen_, Mar 06 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mohammed Bouras_, May 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016158 | OEIS-A362105

Numbers that take a record number of steps to appear in the Sisyphus sequence A350877.

It is conjectured that every number will eventually appear in A350877. The terms in this sequence are the numbers that are the slowest to appear. For example, 115 does not appear until term 45274461582754.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 12 2023, based on a suggestion from Siobhan Roberts | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016159 | OEIS-A362149

Decimal expansion of K, a constant arising in the analysis of the binary Euclidean algorithm.

Vallée (1998) conjectured that this constant times A362150 equals 4*log(2)/Pi^2; Brent (1999) supported the conjecture with numerical computations and Morris (2016) proved the conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Apr 09 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016160 | OEIS-A362167

a(n) = the hypergraph Catalan number C_2(n).

Gunnells gives several combinatorial interpretations of the hypergraph Catalan numbers, a method to compute their generating functions to arbitrary precision and some conjectural asymptotics. a(n) ~ e^(3/2) * 2^(n+1) * n!/sqrt(Pi*n) (conjectural).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016161 | OEIS-A362168

a(n) = the hypergraph Catalan number C_3(n).

Gunnells gives several combinatorial interpretations of the hypergraph Catalan numbers, a method to compute their generating functions to arbitrary precision and some conjectural asymptotics. a(n) ~ sqrt(3) * (9/2)^n * n!^2/(Pi*n) (conjectural).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016162 | OEIS-A362169

a(n) = the hypergraph Catalan number C_4(n).

Gunnells gives several combinatorial interpretations of the hypergraph Catalan numbers, a method to compute their generating functions to arbitrary precision and some conjectural asymptotics. a(n) ~ sqrt(2) * (32/3)^n * n!^3/(Pi*n)^(3/2) (conjectural).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016163 | OEIS-A362170

a(n) = the hypergraph Catalan number C_5(n).

Gunnells gives several combinatorial interpretations of the hypergraph Catalan numbers, a method to compute their generating functions to arbitrary precision and some conjectural asymptotics. a(n) ~ sqrt(5)/2 * (5^4/24)^n * n!^4/(Pi*n)^2 (conjectural).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016164 | OEIS-A362171

a(n) = the hypergraph Catalan number C_6(n).

Gunnells gives several combinatorial interpretations of the hypergraph Catalan numbers, a method to compute their generating functions to arbitrary precision and some conjectural asymptotics. a(n) ~ sqrt(3)/2 * (6^5/5!)^n * n!^5/(Pi*n)^(5/2) (conjectural)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016165 | OEIS-A362172

a(n) = the hypergraph Catalan number C_7(n).

Gunnells gives several combinatorial interpretations of the hypergraph Catalan numbers, a method to compute their generating functions to arbitrary precision and some conjectural asymptotics. a(n) ~ sqrt(7)/4 * (7^6/6!)^n * n!^6/(Pi*n)^3 (conjectural).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016166 | OEIS-A362191

Lexicographically earliest sequence of distinct nonnegative integers having the same concatenation of digits as the sequence 2^a(n).

We conjecture that this is a permutation of the nonnegative integers, but a proof seems without reach. Can it be disproved?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016167 | OEIS-A362202

Lexicographic earliest sequence of distinct positive integers having the same concatenation of digits as the sequence 2^a(n).

We conjecture that this is a permutation of the positive integers, but a proof seems without reach. Can it be disproved?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016168 | OEIS-A362248

a(n) is the number of locations 1..n-1 which can reach i=n-1, where jumps from location i to i +- a(i) are permitted (within 1..n-1); a(1)=1. See example.

Conjecture: a(n) is also the largest number such that starting point i=n can reach every previous location (with a(1)=1 and the same rule for jumps as in the current name).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Neal Gersh Tolunsky_, May 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016169 | OEIS-A362264

Numbers > 9 with increasingly large digit average of their square, in base 10.

It is conjectured but not known that there are only finitely many numbers whose square has a digit average above 8.3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 13 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016170 | OEIS-A362334

a(n) = A000010(n) + A000010(n+2), where A000010 is the Euler phi-function.

Conjecture: a(2*n) <= a(2*n-1) and a(2*n) < a(2*n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexandre Herrera_, Apr 16 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016171 | OEIS-A362372

Inventory of powers. Initialize the sequence with '1'. Then record the number of powers of 1 thus far, then do the same for powers of 2 (2, 4, 8, ...), powers of 3, etc. When the count is zero, do not record a zero; rather start the inventory again with the powers of 1.

Open question: will all positive integers appear in the sequence?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Damon Lay_, Apr 17 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016172 | OEIS-A362406

Denominators of the expansion of the series reversion of sin(x) + x, odd powers only.

Conjecture: a(n) = 2*A306254(n)/(4^(n+1)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Billy Bolton_, Apr 18 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016173 | OEIS-A362407

Numerators of the expansion of the series reversion of sin(x) + x, odd powers only.

Conjecture: a(n) = abs(A302977(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Billy Bolton_, Apr 18 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016174 | OEIS-A362408

a(n) = [x^n] (F(x)/F(-x))^n where F(x) = (1 + x)*(1 + x^3).

Let k and m be positive integers and let f(x) be a finite product of cyclotomic polynomials. Define b(n) = [x^(k*n)] (f(x)/f(-x))^(m*n). Then we conjecture that the supercongruences a(p) == a(1) (mod p^3) and, for n >= 2, a(n*p) == a(n) (mod p^2) hold for all primes p, with a finite number of exceptions depending on f(x). Conjectures: 1) the supercongruence a(p) == 2 (mod p^3) holds for all primes p >= 5 (checked up to p = 47).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 18 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016175 | OEIS-A362434

Numbers that can be written as A000045(i) + j^2 for i,j>=0 in 4 ways.

Conjecture: a(n) = 1 + A000045(6*n)^2/4. Conjecture: Sum_{n>=1} 1/a(n) = sqrt(5)/4 - 1/2 (true if the above formula is valid). - _Amiram Eldar_, Dec 25 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Apr 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016176 | OEIS-A362460

a(n) = A054978(n)/2 if that number is 0 or 1, otherwise -1.

Conjectured to be always 0 or 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 07 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016177 | OEIS-A362528

Numbers that can be written in at least 3 ways as the sum of a Lucas number (A000032) and a square.

Conjecture: there are never more than 3 solutions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Apr 23 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016178 | OEIS-A362534

Numerators of the ratio of the symmetry-constrained bound to the adiabatic bound on polarization transfer in AXn spin-1/2 systems.

Conjecture: the numerator of g(n) is the denominator of f(n)/g(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mohamed Sabba_, Apr 24 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016179 | OEIS-A362552

a(n) = n for n <= 2. For n > 2, a(n) is the least novel k (with rad(k) != rad(a(n-1))) such that k shares a nontrivial divisor with one of a(n-1), a(n-2), but not with the other.

Conjectured to be a permutation of the positive integers in which the primes appear in natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Apr 24 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016180 | OEIS-A362602

Integers in the interval [Pi*k - 1/k, Pi*k + 1/k] for some k > 0 that are numerators of convergents to 2*Pi.

Conjecture: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander R. Povolotsky_ and _Stefano Spezia_, Apr 27 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016181 | OEIS-A362623

Lexicographically earliest sequence of distinct positive terms such that for any n > 0, the initial digit "d" of a(n) divides a(n+d).

Conjecture: For n > 68, a(n) < 3*n. - _Charles R Greathouse IV_, Mar 13 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_, Apr 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016182 | OEIS-A362631

Lexicographically earliest infinite sequence of distinct positive integers with a(n) = n for n <= 3, and for n > 3 a(n) is the least novel multiple of the greatest prime divisor of a(n-2) which does not divide a(n-1).

Conjectured to be a permutation of the positive integers with the primes in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, May 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016183 | OEIS-A362676

a(n) = Sum_{k = 0..n} 4^(n-k)*binomial(n,k)*binomial(n-1,k)*binomial(2*k,k).

The Franel numbers satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and positive integers n and r. We conjecture that the present sequence satisfies the same supercongruences. We conjecture that the present sequence satisfies the same supercongruences. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016184 | OEIS-A362679

a(n) is the permanent of the n X n symmetric matrix M(n) defined by M[i, j, n] = min(i, j)*(n + 1) - i*j.

Conjecture: det(M(n)) = A000272(n+1). The conjecture is true (see proof in Links). - _Stefano Spezia_, May 24 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Apr 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016185 | OEIS-A362722

a(n) = [x^n] (E(x)/E(-x))^n where E(x) = exp(Sum_{k >= 1} A005258(k)*x^k/k).

We conjecture below that {a(n)} satisfies supercongruences similar to (but weaker than) the above supercongruences satisfied by the Apéry numbers. Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, May 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016186 | OEIS-A362724

a(n) = [x^n] E(x)^n, where E(x) = exp(Sum_{k >= 1} A005258(k)*x^k/k).

We conjecture below that {a(n)} satisfies supercongruences similar to (but weaker than) the above supercongruences satisfied by the Apéry numbers. We conjecture that the sequences {a(i,n) : n >= 0}, i >= 1, satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(2*r)) for all primes p >= 3, and positive integers n and r. Conjecture: the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(2*r)) holds for all primes p >= 3 and positive integers n and r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, May 02 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016187 | OEIS-A362725

a(n) = [x^n] E(x)^n, where E(x) = exp(Sum_{k >= 1} A005259(k)*x^k/k).

We conjecture below that the present sequence satisfies supercongruences similar to (but weaker than) the above supercongruences satisfied by the Apéry numbers. We conjecture that the sequences {a(i,n) : n >= 0}, i >= 1, satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(2*r)) for all primes p >= 3, and positive integers n and r. Conjecture: the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(2*r)) holds for all primes p >= 3 and positive integers n and r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, May 02 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016188 | OEIS-A362726

a(n) = [x^n] E(x)^n where E(x) = exp(Sum_{k >= 1} A208675(k)*x^k/k).

We conjecture that the sequences {a(i,n) : n >= 0}, i >= 1, satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 7, and positive integers n and r. Conjecture: the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) holds for all primes p >= 7 and positive integers n and r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, May 02 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016189 | OEIS-A362727

a(n) = [x^n] (E(x)/E(-x))^n where E(x) = exp(Sum_{k >= 1} A208675(k)*x^k/k).

Conjecture: the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) holds for all primes p >= 7 and positive integers n and r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, May 02 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016190 | OEIS-A362730

a(n) = [x^n] E(x)^n where E(x) = exp(Sum_{k >= 1} binomial(2*k,k)^2*x^k/k).

We conjecture that the sequences {a(i,n) : n >= 0}, i >= 1, satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5, and positive integers n and r. Conjecture: the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) holds for all primes p >= 5 and positive integers n and r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, May 05 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016191 | OEIS-A362731

a(n) = [x^n] E(x)^n where E(x) = exp(Sum_{k >= 1} A000172(k)*x^k/k).

Conjecture: the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(2*r)) holds for

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, May 05 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016192 | OEIS-A362732

a(n) = [x^n] E(x)^n, where E(x) = exp(Sum_{k >= 1} A006480(k)*x^k/k).

We conjecture that the sequences {a(i,n) : n >= 0}, i >= 1, satisfy the same supercongruences as above. Conjecture: the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) holds for all primes p >= 3 and positive integers n and r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, May 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016193 | OEIS-A362733

a(n) = [x^n] F(x)^n, where F(x) = exp(Sum_{k >= 1} A362732(k)*x^k/k).

Conjecture: the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) holds for all primes p >= 3 and positive integers n and r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, May 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016194 | OEIS-A362743

Positive integers which cannot be written as a sum of distinct numbers of the form 4^a + 5^b (a,b >= 0).

Conjecture 1: The only terms of the current sequence are 1, 3, 4, 10, 12, 18. Moreover, any positive integer not among 1, 3, 4, 8, 10, 12, 13, 18, 25, 39, 42 can be written as a sum of numbers of the form 4^a + 5^b (a,b>=0) with no one summand dividing another. Conjecture 2: Let k and m be positive integers greater than one with k*m even. Then, any sufficiently large integer n can be written as a sum of distinct numbers of the form k^a + m^b with a and b nonnegative integers. Conjecture 3: Let k and m be positive integers greater than one with k*m even. Then, any sufficiently large integer n can be written as a sum of numbers of the form k^a + m^b (a,b >= 0) with no summand dividing another. Clearly, Conjecture 3 is stronger than Conjecture 2. See also A362861 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016195 | OEIS-A362809

Numbers k for which the area of the first part of the symmetric representation of sigma(k) equals sigma(k)/3 and its width is 1.

Conjecture: The area of the first part of SRS(n) being equal to sigma(n)/3 implies that the first part has width 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, May 04 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016196 | OEIS-A362810

Define G(n, k) to be the n-th derivative of Gamma(x) at k. a(n)=floor(min(G(2n, x))), where min(f) is the local minimum of f in [0,oo).

Conjecture: limit_{n->oo} log(a(n)) / log(n!) < 1. - _Vaclav Kotesovec_, Nov 17 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jodi Spitz_, May 04 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016197 | OEIS-A362855

a(n) = n for n <= 3; for n > 3, a(n) is the least novel multiple of k, the product of all distinct prime factors of a(n-2) that do not divide a(n-1).

Conjecture: all primes appear eventually, but not in order. (End) Conjecture: permutation of natural numbers. Conjecture: the smallest missing number is always either prime or a powerful number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_ and _David James Sycamore_, May 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016198 | OEIS-A362861

Positive integers n such that 2*n cannot be written as a sum of distinct elements of the set {5^a + 5^b: a,b = 0,1,2,...}.

Conjecture 1: (i) The current sequence only has the listed 14 terms. Also, each positive even number can be written as a sum of distinct elements of the set {3^a + 3^b: a,b = 0,1,2,...}. Conjecture 2: Let k and m be positive odd numbers greater than one. Then, any sufficiently large even numbers can be written as a sum of distinct elements of the set {k^a + m^b: a,b = 0,1,2,...}. Conjecture 3: Let k and m be positive odd numbers greater than one. Then, any sufficiently large even numbers can be written as a sum of some numbers of the form k^a + m^b (a,b >= 0) with no summand dividing another. Clearly, Conjecture 3 is stronger than Conjecture 2. See also A362743 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, May 05 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016199 | OEIS-A362866

Numbers k with the property that the parts of the symmetric representation of sigma(k) are two octagons.

Conjecture 1: The octagons are S-shaped and they have width 1. Conjecture 2: This sequence is also the primes doubled (or even semiprimes) >= 10 (Cf. A100484). - _Omar E. Pol_, Aug 15 2023 For the symmetric representation of sigma(n) to consist of 2 octagons the first 3 entries in row n of the triangle of A249223 must be nonzero, hence must be 1's, indicating width 1, with the remaining entries zero. Therefore, row n of A237048 is 100100..., implying n = 2*p with p>3 prime. Both conjectures are true. - _Hartmut F. W. Hoft_, Aug 22 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, May 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016200 | OEIS-A362884

a(n) = (a(n-1)*a(n-2)*a(n-3)+64)/(4*a(n-4)) with a(0) = a(2) = a(3) = 2 and a(1) = 16.

Conjecture 1: all terms are integer. Conjecture 2: a(n) is never divisible by 64.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, May 07 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016201 | OEIS-A362889

a(n) = n for n <= 2. Let i = a(n-2), j = a(n-1), q = gpf(i*j) = prime(s), and k = product of all distinct primes < q which do not divide i*j. For n > 2 a(n) is the least novel multiple of either k (if k is not the empty product), or of prime(s+1) if it is.

Conjectured to be a permutation of the positive integers (but primes are not in order, e.g. 41 precedes 37).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, May 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016202 | OEIS-A362938

a(n) = minimum number of pieces needed to dissect a regular n-sided polygon into a monotile, i.e. a polygonal tile which tiles the plane (conjectured).

a(n) = minimum number of pieces needed to dissect a regular n-sided polygon into a monotile, i.e. a polygonal tile which tiles the plane (conjectured). The three related sequences A110312, A362938, A362939 are exceptions to the usual OEIS policy of requiring that all terms in sequences must be known exactly. These sequences are included because of their importance and in the hope that someone will establish the truth of some of the conjectured values. On Aug 31 2023 Gavin Theobald sent me two different solutions for a(9) = 3 and one solution for a(11) = 4 (see links). He

reports that he found these dissections in the 1990's. In his email and in a later email (Sep 04 2023) he also gives the values a(13) = 5, a(14) = 3, a(15) = 5 (with one piece turned over), a(16) = a(18) = 4, a(20) = 5. He conjectures that a(2t) = floor(t/2) for all t >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016203 | OEIS-A362939

a(n) = minimum number of pieces needed to dissect a regular n-sided polygon into a rectangle (conjectured).

a(n) = minimum number of pieces needed to dissect a regular n-sided polygon into a rectangle (conjectured). The three related sequences A110312, A362938, A362939 are exceptions to the usual OEIS policy of requiring that all terms in sequences must be known exactly. These sequences are included because of their importance and in the hope that someone will establish the truth of some of the conjectured values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 31 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016204 | OEIS-A362946

Positive integers that cannot be expressed as 1^e_1 + 2^e_2 + 3^e_3 ... + k^e_k with each exponent positive.

I conjecture that this list is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert C. Lyons_, Jul 05 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016205 | OEIS-A363011

Indices of record highs in A362816.

It is conjectured that only terms in A362815 appear in A362816, and novel terms in A362816 seen to be in ascending order. Does every term in A362815 appear in A362816? If the conjecture is true and novel terms are in ascending order, then these are the smallest k such that A362816(k) = A362815(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Samuel Harkness_, May 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016206 | OEIS-A363082

Numbers k neither squarefree nor prime power such that q*r > k, where q = A053669(k) is the smallest prime that does not divide k and r = A007947(k) is the squarefree kernel.

All terms are even. Conjectured to be a subsequence of A389216. - _Antti Karttunen_, Sep 30 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jul 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016207 | OEIS-A363104

Expansion of g.f. A(x) satisfying 4 = Sum_{n=-oo..+oo} (-x)^n * (4*A(x) + x^(n-1))^(n+1).

Conjecture: g.f. A(x) == theta_3(x) (mod 4); a(n) == 2 (mod 4) iff n is a nonzero square and a(n) == 0 (mod 4) iff n is nonsquare.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016208 | OEIS-A363110

G.f.: Sum_{n>=0} x^n * Product_{k=1..n} (k + (n-k+1)*x) / (1 + k*x + (n-k+1)*x^2).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 02 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016209 | OEIS-A363148

a(n) gives the number of equivalence classes of quaternary quadratic forms of discriminant A363147(n) not representing 2.

Conjecture: a(n) ~ c * A363147(n) ^ d where d is a constant which is roughly 1.51 and c is one of four constants, depending on the value of A363147(n) mod 24. See plots in files.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andy Huchala_, May 17 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016210 | OEIS-A363151

a(n) = denominator(Sum_{j=0..n} Bernoulli(j, 1) * Bernoulli(n - j, 1)).

Conjecture: a(n) is cubefree. (An integer is cubefree if it is not divisible by the cube of a prime number.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, May 18 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016211 | OEIS-A363162

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number that has not yet appeared that has the same number of distinct prime divisors as a(n-2) + a(n-1).

The terms with different numbers of distinct prime divisors are concentrated along different lines in the graph; see the attached colored image. There are numerous fixed points in the first one million terms, although the last nonprime fixed point is a(n) = 3495. Beyond that there are thirty-one more fixed points all with prime values; it is likely more exist although this is unknown. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jul 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016212 | OEIS-A363166

Bouton numbers: a(n) is the number of P positions in games of Nim with three nonzero heaps each containing at most n sticks.

Conjecture: a(n) = A000217(n) - A063915(n-1). - _Alan Michael Gómez Calderón_, Sep 18 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Rowlett_, Jul 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016213 | OEIS-A363184

Expansion of g.f. A(x) satisfying 4 = Sum_{n=-oo..+oo} (-1)^n * x^n * (4*A(x) + x^(2*n-1))^(n+1).

Conjecture: g.f. A(x) == theta_3(x^2) (mod 4); a(n) == 2 (mod 4) if n = 2*k^2 for integer k > 0, and a(n) == 0 (mod 4) if floor(n/2) is nonsquare.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 20 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016214 | OEIS-A363196

a(n) is the least k such that the rightmost 7 in the decimal expansion of 3^k is in the (10^n)'s position, or -1 if there is no such k.

Conjecture: a(n) = -1 only for n = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, May 20 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016215 | OEIS-A363198

a(n) = n for n <= 3; for n >= 4, a(n) is the smallest positive integer that has not appeared previously in this sequence and shares a factor with a(n-1) + a(n-2) + a(n-3).

Conjecture: This sequence is a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yifan Xie_, May 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016216 | OEIS-A363228

Exponent of 4 in 9^n - 1.

Conjecture: a(n) = A235127(A000045(6*n)), all other 4-adic 6-sections A235127(A000045(.)) = 0. - _R. J. Mathar_, Jun 28 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ruud H.G. van Tol_, May 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016217 | OEIS-A363265

Number of integer factorizations of n with a unique mode.

Conjecture: 9 is missing from this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jun 27 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016218 | OEIS-A363285

a(n) is the smallest multiple k of 2^n such that |sigma(k) - 2*k| = 2^n, where sigma = A000203.

I conjecture that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jesiah Darnell_, May 25 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016219 | OEIS-A363308

Expansion of g.f. C(x*C(x)^3), where C(x) = 1 + x*C(x)^2 is the g.f. of the Catalan numbers (A000108).

Conjecture: a(n) is odd iff n is a power of 2 or n = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016220 | OEIS-A363309

Expansion of g.f. A(x) = F(x*F(x)^5), where F(x) = 1 + x*F(x)^3 is the g.f. of A001764.

Conjecture: given A(x) = F(x*F(x)^(2*n-1)) where F(x) = 1 + x*F(x)^n, let B(x) = A(x*B(x)^(n-1)), then ((B(x) - 1)/x)^(1/(2*n-1)) is an integer series for n >= 1. Incidentally, the function A(x) = F(x*F(x)^(2*n-1)) is interesting because F(-x*F(x)^(2*n-1)) = 1/F(x) when F(x) = 1 + x*F(x)^n. This sequence illustrates the case for n = 3; for n = 2, see A363308.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016221 | OEIS-A363321

Number of fractions of the Farey sequence of order n, F_n, that coincide with those of the sequence of the #{F_n} equally distributed fractions between 0 and 1.

The conjectured formula below suggests that as the value of n increases, the proportion of terms in the Farey sequence F_n that align with the #F_n rationals, evenly distributed between 0 and 1, tends to decrease. Conjecture: lim_{n->infinity} a(n)/A005728(n) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andres Cicuttin_, May 27 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016222 | OEIS-A363364

Least nonnegative integer k such that all non-bipartite graphs with n nodes and at least k edges are weakly pancyclic.

Brandt (1997) conjectures that a(n) = floor((n-1)*(n-3)/4) + 5 for n >= 6. The conjecture is false for n = 8, since there exists a (unique) non-bipartite, not weakly pancyclic graph (shown below) with 8 nodes and 13 edges, showing that a(8) >= 14. This graph contains cycles of lengths 3, 4, 5, 6, and 8, but none of length 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pontus von Brömssen_, May 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016223 | OEIS-A363376

Determinant of the n X n matrix formed by placing 1..n^2 in L-shaped gnomons in alternating directions.

Conjecture: a(n) has trailing zeros for n > 3. - _Stefano Spezia_, May 31 2023 The conjecture is true and its proof follows easily from _Detlef Meya_'s formula. - _Stefano Spezia_, Apr 20 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nicolay Avilov_, May 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016224 | OEIS-A363381

a(n) is the number of distinct n-cell patterns (or quasi-polyominoes) that tile an n X n square.

Conjecture by Andrew Young: For an n X n square, where n is an odd prime, there is only one n-cell pattern. Conjecture by Andrew Young and Thomas Young: An odd integer n>=3 is prime iff there exists only one n-cell pattern for an n X n square. The second conjecture is a Corollary of the first: If n = p*q is not prime, then there is always a second tiling using rectangles, as explained above. Since the second conjecture implies the first, the two conjectures are actually equivalent. - _M. F. Hasler_, Jun 15 2025 For n > 1, a(n) = 1 iff n is an odd prime (conjectured: cf comments).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Young_, May 30 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016225 | OEIS-A363409

a(n) = the real part of Product_{k = 1..n} (1 + k*sqrt(-2)).

We conjecture that the set of type 1 primes begins {2, 5, 13, 23, 29, 31, 47, 53, 61, 71, 101, ...}. We conjecture that the set of type 2 primes consists of primes p == 1 or 3 (mod 8), i.e., rational primes that split in the field extension Q(sqrt(-2)) of Q. See A033200. Moll's conjecture 5.5 extends to this sequence and takes the form: for prime p == 1 or 3 (mod 8), the p-adic valuation v_p(a(n)) ~ n/(p - 1) as n -> oo. We conjecture that the set of type 3 primes begins {7, 37, 79, 103, ...}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jun 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016226 | OEIS-A363410

a(n)= 1/sqrt(2) * the imaginary part of Product_{k = 1..n} (1 + k*sqrt(-2)).

We conjecture that in this case, unlike in A105750, the set of type 1 primes is empty; that is, every prime p divides some term of this sequence. We conjecture that the set of type 2 primes consists of primes p == 1 or 3 (mod 8), equivalently, rational primes that split in the field extension Q(sqrt(-2)) of Q. See A033200. Moll's conjecture 5.5 extends to this sequence: for primes p of type 2, the p-adic valuation v_p(a(n)) ~ n/(p - 1) as n -> oo. We conjecture that the set of type 3 primes consists of primes p == 5 or 7 (mod 8), equivalently, primes that remain inert in the field extension Q(sqrt(-2)) of Q, together with the prime p = 2, which ramifies in Q(sqrt(-2)). See A033203.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jun 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016227 | OEIS-A363411

a(n) = the real part of Product_{k = 0..n} 1 + k*sqrt(-3).

We conjecture that the set of type 1 primes begins {3, 11, 17, 23, 29, 41, 47, 53, 59, 83, 89, 101, ...}. We conjecture that the set of type 2 primes consists of primes p == 1 (mod 6), i.e., rational primes that split in the field extension Q(sqrt(-3)) of Q, together with the prime p = 2. See A002476. Moll's conjecture 5.5 extends to this sequence and takes the form: We conjecture that the set of type 3 primes begins {5, 71, 179, 191, 227, 257, 263, ...}. Conjecture: the 5-adic valuation v_5(a(n+2)) = A079998(n) (checked up to n =

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jun 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016228 | OEIS-A363412

a(n) = 1/sqrt(3) * the imaginary part of Product_{k = 0..n} 1 + k*sqrt(-3).

We conjecture that in this case, unlike in A105750, the set of type 1 primes is empty; that is, every prime p divides some term of this sequence. We conjecture that the set of type 2 primes consists of primes of the form p == 1 (mod 6), i.e., rational primes that split in the field extension Q(sqrt(-3)) of Q, together with the prime p = 2. See A002476. Moll's conjecture 5.5 extends to this sequence and takes the form: We conjecture that the set of type 3 primes consists of primes of the form p == 5 (mod 6), i.e., the rational primes that remain inert in the field extension Q(sqrt(-3)) of Q, together with the prime p = 3, which ramifies in Q(sqrt(-3)). See A007528. Conjectures: 3 does not divide a(3*n+1) for all n; the 3-adic valuation v_3(a(3*n)) = v_3(a(3*n-1)) for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jun 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016229 | OEIS-A363413

a(n) = the real part of Product_{k = 0..n} 1 + k*sqrt(-4).

We conjecture that the set of type 1 primes begins {2, 3, 11, 19, 23, 31, 47, 59, 67, 71, 79, ...}. We conjecture that the set of type 2 primes consists of primes p == 1 (mod 4), equivalently, rational primes that split in the field extension Q(sqrt(-1)) of Q. See A002144. Moll's conjecture 5.5 extends to this sequence and takes the form: for primes of type 2, the p-adic valuation v_p(a(n)) ~ n/(p - 1) as n -> oo. We conjecture that the set of type 3 primes begins {7, 43, 83, 131, 163, 167, ...}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jun 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016230 | OEIS-A363414

a(n) = (1/2) * the imaginary part of Product_{k = 0..n} 1 + k*sqrt(-4).

In this case, unlike in A105750, the set of type 1 primes is conjecturally empty; it appears that every prime p divides some term of this sequence. We conjecture that the set of type 2 primes consists of primes p == 1 (mod 4), equivalently, rational primes that split in the field extension Q(sqrt(-1)) of Q. See A002144. Moll's conjecture 5.5 extends to this sequence: for the primes of type 2, the p-adic valuation v_p(a(n)) ~ n/(p - 1) as n -> oo. We conjecture that the set of type 3 primes consists of primes p == 3 (mod 4), equivalently, rational primes that remain inert in the field extension Q(sqrt(-1)) of Q, together with the prime p = 2, which ramifies in Q(sqrt(-1)). See A002145.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jun 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016231 | OEIS-A363415

a(n) = the real part of Product_{k = 0..n} 1 + k*sqrt(-5).

We conjecture that the set of type 1 primes begins {5, 11, 13, 31, 53, 79, 97, 113, 137, 157, 179, 193, 197, ...}. We conjecture that the set of type 2 primes begins {2, 3, 7, 23, 29, 41, 43, 47, 61, 67, 83, 89, 101, ...} and consists of primes p == 1, 3, 7 or 9 (mod 20), equivalently, rational primes that split in the field extension Q(sqrt(-5)) of Q, together with the prime p = 2. See A139513. Moll's conjecture 5.5 extends to this sequence: for type two primes p > 2, v_p(a(n)) ~ n/(p - 1) as n -> oo. We conjecture that the set of type 3 primes begins {17, 19, 37, 59, 71, 73, 131, 151, 173, 191, 199, ...}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jun 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016232 | OEIS-A363416

a(n) = 1/sqrt(5) * the imaginary part of Product_{k = 0..n} 1 + k*sqrt(-5).

We conjecture that in this case, unlike in A105750, the set of type 1 primes is empty; that is, every prime p divides some term of this sequence. We conjecture that the set of type 2 primes consists of primes p == 1, 3, 7 or 9 (mod 20),

equivalently, rational primes that split in the field extension Q(sqrt(-5)) of Q, together with the prime p = 2. See A139513. Moll's conjecture 5.5 extends to this sequence and takes the form: We conjecture that the set of type 3 primes consists of primes p == 11, 13, 17 or 19 (mod 20), equivalently, primes that remain inert in the field extension Q(sqrt(-5)) of Q, together with the prime p = 5, which ramifies in Q(sqrt(-5)). See A003626.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jun 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016233 | OEIS-A363418

Square array read by ascending antidiagonals: T(n,k) = [x^(n*k)] ((1 + x)/(1 - x))^k for n, k >= 1.

We conjecture that each row sequence satisfies the stronger supercongruences T(n, m*p^r) == T(n, m*p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and positive integers m and r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jun 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016234 | OEIS-A363435

Number of partitions of [2n] having exactly n blocks with all elements of the same parity.

Conjecture: Limit_{n->oo} (a(n)/n!)^(1/n) = A238258 = -2 / (LambertW(-2*exp(-2)) * (2 + LambertW(-2*exp(-2)))) = 3.0882773... - _Vaclav Kotesovec_, Oct 21 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jun 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016235 | OEIS-A363482

Denominator of the continued fraction 1/(2-3/(3-4/(4-5/(...(n-1)-n/(-5))))).

Conjecture: Except for 49, every term of this sequence is either a prime or 1. The conjecture holds through n=10000. The set of ordered primes appear to match A038901. - _Bill McEachen_, Jul 08 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mohammed Bouras_, Jun 04 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016236 | OEIS-A363504

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number that has not yet appeared that is a multiple of prime(Omega(a(n-1))).

The sequence is conjectured to be a permutation of the positive integers, although it takes many terms for the primes to appear, e.g., a(371221) = 61. After the first two terms the fixed points in the first 500000 terms are 10, 26076, 26151, 26223, 26307, 26379, 26406, although it is possible more exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016237 | OEIS-A363558

Expansion of g.f. A(x) = Sum_{n=-oo..+oo} x^n * (2 + x^n)^(2*n).

a(2^n) = 4^(2^n) for n > 0 (conjecture). a(p) = p*2^(p-1) + 4^p for primes p > 3 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016238 | OEIS-A363559

Expansion of g.f. A(x) = Sum_{n=-oo..+oo} x^n * (3 + x^n)^(2*n).

a(2^n) = 9^(2^n) for n > 0 (conjecture). a(p) = p*3^(p-1) + 9^p for primes p > 3 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016239 | OEIS-A363560

Expansion of g.f. A(x) satisfying A(x)^3 = 1 + x*(A(x) + A(x)^2 + A(x)^9).

Conjecture: a(n) == 0 (mod 3) for n > 0 except when n == 1 (mod 7).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016240 | OEIS-A363561

G.f.: Sum_{n=-oo..+oo} x^n * (sqrt(2) + x^n)^(2*n).

a(2^n) = 2^(2^n) for n > 0 (conjecture). a(p) = p*2^((p-1)/2) + 2^p for primes p > 3 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016241 | OEIS-A363568

Expansion of l.g.f. A(x) satisfying theta_4(x) = Sum_{n=-oo..+oo} x^n * (2*exp(A(x)) - x^n)^(n-1) where theta_4(x) = Sum_{n=-oo..+oo} (-1)^n * x^(n^2) is a Jacobi theta function.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 11 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016242 | OEIS-A363569

Expansion of g.f. A(x) = Sum_{n=-oo..+oo} x^n * (i + x^n)^(2*n), where i^2 = -1.

a(2^n) = 1 for n > 0 (conjecture). a(p) = p*(-1)^((p-1)/2) - 1 for primes p > 3 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 31 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016243 | OEIS-A363574

Expansion of g.f. A(x) satisfying theta_4(x) = Sum_{n=-oo..+oo} x^n * (2*A(x) - x^n)^(n-1) where theta_4(x) = Sum_{n=-oo..+oo} (-1)^n * x^(n^2) is a Jacobi theta function.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016244 | OEIS-A363593

Numbers k such that both A359804(k) and A359804(k+1) are odd.

Conjecture: for prime q > 11, even squarefree semiprimes 2q follow D(n) for some n. Consider that primes in A359804 appear late for q > 11, yet pairs of successive odd numbers in that sequence occur rather often. Conjectured to be an infinite sequence, meaning that consecutive odd terms appear infinitely many times in A359804. - _David James Sycamore_, Jun 21 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_ and _David James Sycamore_, Jun 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016245 | OEIS-A363609

Minimum sum of the visible pips on a polycube made from n dice.

Conjecture: a(k^3) = 6*(k+2)*k for k > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matt Donahoe_, Jun 11 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016246 | OEIS-A363657

Numbers m where A217854(m) is a record minimum.

It is conjectured that if m is a term, then abs((-m)^tau(m)) <= abs((-k)^tau(k)) for some k < m. See the link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Jensen_, Jun 13 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016247 | OEIS-A363683

Square array A(n, k), n, k > 0, read by antidiagonals; A(n, k) is the least e > 0 such that n^e and k^e have the same initial digit, or -1 if no such e exists.

Conjecture: all terms are positive.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jun 15 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016248 | OEIS-A363686

Irregular triangle T(n, k) read by rows which represents a predecessor number system for Collatz sequences (see Comments for precise definition).

Conjecture: With the associated residue classes, this sequence forms a complete decomposition of the odd integers. In other words: If we form the complete residue classes to each number from this sequence, then all odd numbers > 1 occur once. Conjecture: Every odd number > 1 is congruent to a(k) mod 2^(A056792(floor(k/2))) for some k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Conradi_, Jun 15 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016249 | OEIS-A363739

a(n) is the length of the n-th run of identical terms of A349509.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jun 19 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016250 | OEIS-A363793

Number of Q-isomorphism classes of elliptic curves E/Q with good reduction away from prime(n).

R. von Känel and B. Matschke conjecture that a(n) <= 24 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robin Visser_, Jun 22 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016251 | OEIS-A363798

Numbers k such that there is no prime p for which p^k + 2*k is prime.

Includes k = (q^2-1)/2 + j*(q^2-q) for odd primes q and nonnegative integers j such that q^k + 2*k is not prime, since for primes p <> q we have q | p^k + 2*k. Conjecture: all terms are of this form.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Jun 23 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016252 | OEIS-A363822

a(n) is the conjectured number of stable distinct centroidal Voronoi tessellations (CVTs) of a unit disk with n generators (seeds).

a(n) is the conjectured number of stable distinct centroidal Voronoi tessellations (CVTs) of a unit disk with n generators (seeds).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Denis Ivanov_, Oct 18 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016253 | OEIS-A363938

Lexicographically earliest sequence of numbers in a hexagonal spiral such that their neighbors have no common prime factor.

It is conjectured the sequence contains all positive numbers. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carole Dubois_, Jun 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016254 | OEIS-A363983

a(n) = Sum_{k = floor((n+1)/2)..n} (-1)^(n+k)*binomial(n,k)*binomial(n+k-1,k)*binomial(2*k,n).

The Franel numbers satisfy the supercongruences A000172(n*p^r) == A000172(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and positive integers n and r. We conjecture that the present sequence satisfies the same supercongruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016255 | OEIS-A363984

a(n) = Sum_{k = 0..n} (-1)^(n+k)*binomial(n,k)*binomial(n+k,k)*A363983(k).

We conjecture that the present sequence satisfies the same pair of congruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016256 | OEIS-A363985

a(n) = Sum_{k = 0..n} (-4)^(n-k)*binomial(n,k)*binomial(2*n+k,k)*binomial(2*k,k).

The Franel numbers satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and positive integers n and r. We conjecture that the present sequence satisfies the same supercongruences. More generally, define two families of sequences {u_m(n): n >= 0} and {v_m(n): n >= 0}, depending on an integer parameter m, by u_m(n) = Sum_{k = 0..n} (-4)^(n-k)*binomial(n,k)*binomial(m*n + k,k)*binomial(2*k,k) and v_m(n) = Sum_{k = 0..n} (-4)^(n-k)*binomial(n,k)*binomial(m*n + 2*k,2*k)*binomial(2*k,k). In this notation, the Franel numbers A000172 = v_1. Note that u_0(n) = v_0(n) = (-1)^n*binomial(2*n,n). We conjecture that, for all m in Z, the sequences u_m and v_m satisfy the above supercongruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 02 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016257 | OEIS-A363986

a(n) = Sum_{k = 0..n} (-4)^(n-k)*binomial(n,k)*binomial(2*n+2*k,2*k)*binomial(2*k,k).

The Franel numbers satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and positive integers n and r. We conjecture that the present sequence satisfies the same supercongruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 02 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016258 | OEIS-A364003

Integers K such that PSL_2(K) is a K_4-simple group, i.e., |PSL_2(K)| has 4 distinct prime divisors.

This sequence is conjectured to be infinite, see Bugeaud, Cao, & Mignotte.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lixin Zheng_, Jul 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016259 | OEIS-A364004

Orders of simple groups PSL(2,K) with exactly 4 prime divisors.

Sequence is conjectured to be infinite, see Bugeaud et al.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lixin Zheng_, Jul 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016260 | OEIS-A364086

Fixed points of A002326, i.e., numbers k such that A002326(k) = k.

The related case of A002326(k) = 2k is true if (and conjecturally only if) 2k+1 is a prime with primitive root 2, see A001122.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Haase_, Jul 09 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016261 | OEIS-A364087

Integers k for which A000594(k)^2 > 4 k^11, where A000594 is Ramanujan's tau function.

Ramanujan conjectured that for prime values of k, A000594(k)^2 <= 4 k^11, where A000594 is Ramanujan's tau function. This followed from Deligne's proof of the Weil Conjectures. This sequence demonstrates the sparsity of small integers k such that the absolute value of tau(k) is greater than this bound.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jack Gilchrist_, Jul 04 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016262 | OEIS-A364111

a(n) = Sum_{k = 0..n} binomial(n+k-1,k)^2 * binomial(2*n-2*k,n-k) * binomial(2*k,k).

We conjecture that the present sequence satisfies the same supercongruences. More generally, let A >= 2, B and C be positive integers. Then we conjecture that the sequence whose terms are given by Sum_{k = 0..n} binomial(n+k-1,k)^A * binomial(2*n-2*k,n-k)^B * binomial(2*k,k)^C also satisfies the same supercongruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 07 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016263 | OEIS-A364113

Square array read by ascending antidiagonals: T(n,k) = [x^k] 1/(1 - x) * Legendre_P(k, (1 + x)/(1 - x))^n for n, k >= 0.

We conjecture that each row sequence of the present table satisfies the same pair of supercongruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 07 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016264 | OEIS-A364114

a(n) = [x^n] 1/(1 - x) * Legendre_P(n, (1 + x)/(1 - x))^3 for n >= 0.

We conjecture that the present sequence also satisfies the supercongruences 1) and 2). Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 07 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016265 | OEIS-A364115

a(n) = [x^n] 1/(1 - x) * Legendre_P(n, (1 + x)/(1 - x))^4 for n >= 0.

We conjecture that the present sequence also satisfies the supercongruences 1) and 2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016266 | OEIS-A364116

a(n) = [x^n] 1/(1 - x) * Legendre_P(n, (1 + x)/(1 - x))^n for n >= 0.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016267 | OEIS-A364117

a(n) = [x^n] 1/(1 - x) * Legendre_P(n, (1 + x)/(1 - x))^(n+1) for n >= 0.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016268 | OEIS-A364118

a(n) = 3*A364114(n) - 11*A364114(n-1).

It is conjectured that the sequence {A364114(n)} and the shifted sequence {A364114(n-1)} both satisfy the supercongruences A364114(p^r) == A364114(p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and positive integers r. Stronger supercongruences may hold for the present sequence, a linear combination of A364114(n) and A364114(n-1). Conjectures: 1) the supercongruences a(p) == a(1) (mod p^5) hold for all primes p >= 7 (checked up to p = 101). There is also a multiplicative version of this sequence. Define a sequence of rational numbers {b(n) : n >= 1} by b(n) = A364114(n)^21 / A364114(n-1)^11. Then we conjecture that the above pair of supercongruences also hold for the sequence {b(n)}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016269 | OEIS-A364119

a(n) = 7*A364115(n) - 17*A364115(n-1).

It is conjectured that the sequence {A364115(n)} and the shifted sequence {A364115(n-1)} both satisfy the supercongruences A364115(p^r) == A364115(p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and positive integers r. Stronger supercongruences may hold for the present sequence, a linear combination of A364115(n) and A364115(n-1). Conjectures: 1) the supercongruences a(p) == a(1) (mod p^5) hold for all primes p >= 7 (checked up to p = 101). There is also a multiplicative version of this sequence. Define a sequence of rational numbers {b(n) : n >= 1} by b(n) = A364115(n)^63 / A364115(n-1)^17. Then we conjecture that the above pair of supercongruences also hold for the sequence {b(n)}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016270 | OEIS-A364131

Numbers k for which A348717(k) is a multiple of A348717(sigma(k)).

Conjecture: All terms apart from a(2) = 2 are squares.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jul 11 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016271 | OEIS-A364172

a(n) = (6*n)!*(n/3)!/((3*n)!*(2*n)!*(4*n/3)!).

A295437, defined by A295437(n) = (18*n)!*n! / ((9*n)!*(6*n)!*(4*n)!) is one of the 52 sporadic integral factorial ratio sequences of height 1 found by V. I. Vasyunin (see Bober, Table 2, Entry 7). Here we are essentially considering the sequence {A295437(n/3) : n >= 0}. This sequence is only conjecturally an integer sequence. Conjecture: the supercongruences a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) hold for all primes p >= 5 and all positive integers n and r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016272 | OEIS-A364173

a(n) = (9*n)!*(2*n)!*(3*n/2)!/((9*n/2)!*(4*n)!*(3*n)!*n!).

This sequence is only conjecturally an integer sequence. Conjecture: the supercongruences a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) hold for all primes p >= 5 and all positive integers n and r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 13 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016273 | OEIS-A364175

a(n) = (6*n)!*(2*n/3)!/((3*n)!*(2*n)!*(5*n/3)!).

This sequence is only conjecturally an integer sequence. Conjecture: the supercongruences a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) hold for all primes p >= 5 and all positive integers n and r. The integrality conjecture is true. See Links.

- _David Radcliffe_, Jul 19 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 13 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016274 | OEIS-A364197

a(n+1) = a(|n-a(n)^2|) + 1, a(0) = 0.

a(n) ~ (3n)^(1/3) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rok Cestnik_, Jul 13 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016275 | OEIS-A364198

Index of first occurrence of n in A364197.

a(n) ~ n^3/3 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rok Cestnik_, Jul 13 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016276 | OEIS-A364201

Lexicographically earliest sequence of distinct positive integers such that the sum of all terms a(1)..a(n) in binary is a substring of the concatenation of all terms a(1)..a(n) in binary.

In the first 10000 terms the smallest number that has not yet appeared is 7026; it is conjectured all numbers eventually appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jul 13 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016277 | OEIS-A364243

a(n) = A108625(2*n-1, n-1) for n >= 1.

The sequence of Apéry numbers A005258 forms the main diagonal of A108625, i.e., A005258(n) = A108625(n, n). The Apéry numbers satisfy the supercongruences A005258(n*p^r) == A005258(n^p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and positive integers n and r. We conjecture that the present sequence satisfies the same supercongruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 16 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016278 | OEIS-A364244

a(n) = A143007(2*n-1, n-1) for n >= 1.

The sequence of Apéry numbers A005259 forms the main diagonal of A143007, i.e., A005259(n) = A143007(n, n). The Apéry numbers satisfy the supercongruences A005259(n*p^r) == A005259(n^p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and positive integers n and r. We conjecture that the present sequence satisfies the same supercongruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 16 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016279 | OEIS-A364246

a(1) = 1. Thereafter, a(n) is the least novel multiple of either prime(k+1) if rad(a(n-1)) = A002110(k), or Product_{prime q; q < gpf(a(n-1)); and q!|a(n-1)} q otherwise.

In other words, if the squarefree kernel of a(n-1) is a primorial term then a(n) is the least novel multiple of the smallest prime which does not divide a(n-1). Otherwise a(n) is the least novel multiple of the product of all primes < gpf(a(n-1)) which do not divide a(n-1). Primes >= 5 arrive late (as least unused term), and a(k) is prime(m) iff a(k-1) is A002110(m-1). The pattern around a prime is P(k), prime(k+1), 2*P(k), m*prime(k+1) for some multiplier m, where P(k) = A002110(k). The sequence is conjectured to be a permutation of the positive integers, with primes in natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jul 15 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016280 | OEIS-A364264

The Parker Square, read by rows.

Named after Matt Parker, who attempted (and failed) to create a 3 X 3 magic square of squares (still an open problem). The sum of entries in the rows, columns and one diagonal is 3051, but in the other diagonal the sum is 4107. Moreover, three entries are repeated (1^2, 29^2 and 41^2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Jul 17 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016281 | OEIS-A364281

Number of permutations of [n] with distinct cycle lengths such that each cycle contains exactly one cycle length as an element.

Conjecture: a(n) ~ exp(1) * (n-1)!. - _Vaclav Kotesovec_, May 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Jul 17 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016282 | OEIS-A364284

Numbers k such that A261627(k) = 1.

Conjectured to be the only numbers k such that A261627(k) = 1. No other terms < 1.2 * 10^12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_, Aug 26 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016283 | OEIS-A364294

Difference k - A163511(k) computed for those odd numbers k for which the difference is nonnegative.

Conjecture: a(1) is the only zero in this sequence, which is equal to the claim that A007283 gives all fixed points of the map n -> A163511(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jul 25 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016284 | OEIS-A364303

Square array read by ascending antidiagonals: T(n,k) = [x^k] (1 - x)^(2*k) * Legendre_P(n*k, (1 + x)/(1 - x)) for n, k >= 0.

The first row of the table is a signed version of the central binomial coefficients A000984. The central binomial coefficients satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p ^(3*r)) for all primes p >= 5 and all positive integers n and r (see Meštrović, equation 39). We conjecture that each row sequence of the table satisfies the same supercongruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 19 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016285 | OEIS-A364305

a(n) = (8*n)!*(5*n)!*(3*n)! / ((6*n)!*(4*n)!^2*n!^2).

Conjecture: the supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) hold for all primes p >= 5 and all positive integers n and k [added 16 Oct 2024: the conjecture follows from Meštrović, equation 39, since a(n) = binomial(8*n, 2*n)*binomial(5*n, n)* binomial(2*n, n)/binomial(4*n, n)]. For integer r and positive integer s, define sequences u(n) = { [x^(s*n)] G(x)^(r*n) : n >= 0 } and v(n) = { [x^(s*n)] F(x)^(r*n) : n >= 0 }, with the power series F(x) and G(x) as defined above. We conjecture that both sequences {u(n)} and {v(n)} satisfy the above supercongruences mod p^(3*k). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016286 | OEIS-A364332

a(n) = f(prime(n)), where f(2) = 0 and for an odd prime p, f(p) = max{f(q)+1: q ranges over all prime factors of p-1}.

Conjecture: the smallest prime p such that f(p) = m (namely a(primepi(p)) = m) is p = A082449(m). Verified for m <= 13. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Lu_, Jul 18 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016287 | OEIS-A364364

For n <= 3 a(n) = n. Label a(n-3), a(n-2), a(n-1) i,j,k respectively. If i,j,k are pairwise coprime a(n) is the least unused number which shares a divisor with k. Otherwise a(n) is the least unused number coprime to j.

Conjectured to be a permutation of the positive integers, with primes occurring in natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jul 20 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016288 | OEIS-A364366

An irregular triangle read by rows, the 3rd row-symmetric Fibonaccian triangle: T(n,k) is the Whitney number of level k of the (3,n)-th symmetric Fibonaccian lattice (0 <= n, 0 <= k <= 2*n).

In [Donnelly, Dunkum, Lišková, and Nance, 2023], the (m,n)-th symmetric Fibonaccian lattice L is also described using semistandard tableaux of a specific ribbon shape; the irreducible components of the associated sl(m,C)-representation are in one-to-one correspondence with what are called the 'ballot-admissible' (aka Littlewood-Richardson) tableaux. In terms of Fibonaccian strings, an element S = (S(1),...,S(n)) in L is ballot-admissible iff for any integer q between 1 and n (inclusive) and any integer r between 1 and m-1 (inclusive), the following integer quantity is nonnegative: Sum_{k=n+1-q..n}([n+1-k is odd]*([r+(k-1)*m = S(k)] - [r+(k-1)*m+1 = S(k)]) + [n+1-k is even]*([k*m-r = S(k)]-[k*m+1-r = S(k)])), where '[]' denotes the Iverson bracket. Enumerating the ballot-admissible tableaux or Fibonaccian strings in L seems to be an interesting problem when m >= 3; when m=3, the sizes of the sets of ballot-admissible tableaux conjecturally agree with A004148.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Donnelly_ and _Molly W. Dunkum_, Jul 20 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016289 | OEIS-A364414

Numbers k with the property that the second part of the symmetric representation of sigma(k) is an octagon of width 1 and one of its vertices is also the central vertex of the first valley of the largest Dyck path of the diagram.

Conjecture 1: These are the numbers > 9 that are congruent to {3, 9, 21, 27} mod 30. Conjecture 2: These are the terms > 9 of A016945 except the terms ending in 5. Conjecture 3: The polygon mentioned in the definition is an "S"-shaped concave octagon. Conjecture 4: Every term of this sequence has as nearest neighbor a term of A091999. Conjecture 5: The terms of A091999 greater than 2 are the numbers k with the property that the first part of the symmetric representation of sigma(k) is an octagon. Conjecture 6: The octagon mentioned in the definition shares at least an edge with the octagon mentioned in conjecture 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jul 23 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016290 | OEIS-A364421

For n >= 3, r >= 0, y an integer, a(n) is the number of integral solutions to the elliptic equation y^2 = n^3 + n^2 + 2*r*n + r^2.

Factoring the equation y^2 = n^3 + n^2 + 2*r*n + r^2 yields (y+n+r)*(y-n-r) = n^3, which implies y+n+r = d and y-n-r = n^3/d for some divisor d of n^3. Thus a(n) is the number of divisors d of n^3 such that (d-n^3/d)/2 - n is a nonnegative integer. This resolves some of Thomas Scheuerle's conjectures. - _Robin Visser_, Sep 30 2023 Conjecture: a(A190300(n)) = 3. Conjecture: a(A196226(n)) = 4. Conjecture: a(p^3) = 5 if p is an odd prime. Conjecture: a(2*p^2) = 7 if p is an odd prime. But there exist other cases too, for example a(3*23) = 7. Conjecture: a(prime(n)^prime(n)) = A245685(n - 1) - 1. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Sep 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016291 | OEIS-A364456

Lexicographically earliest sequence of distinct primes such that a(1)*a(2)*...*a(n-1) - a(n) is prime for n>=1 with a(1)=5.

Conjecture: This sequence contains all primes. Conjecture: If f(n) = a(1)*a(2)*...*a(n-1) then f(n)-a(n) is the previous prime of f(n) - 1 for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giorgos Kalogeropoulos_, Jul 25 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016292 | OEIS-A364470

a(n) is the least k > a(n-1) such that Omega(k * (a(n-1)+k)) = n, where Omega = A001222; a(1) = 1.

Conjecture: a(n+4) = 4*a(n) for n >= 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Jul 25 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016293 | OEIS-A364481

The number of stones-and-bones tilings of the (3n+1,3n+1)-benzel when one orientation of bone is forbidden.

See Problem 9 of the Propp article. It is conjectured that a(n) is 2-adically continuous as a function of n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James Propp_, Jul 26 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016294 | OEIS-A364482

The number of stones-and-bones tilings of the (3n+1,3n+2)-benzel when one orientation of bone is forbidden.

See Problem 10 of the Propp article. It is conjectured that a(n) is 2-adically continuous as a function of n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James Propp_, Jul 26 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016295 | OEIS-A364506

Square array read by ascending antidiagonals: T(n,k) = (2*k)!/k! * ((2*n*k)! * ((2*n+1)*k)!)/((n*k)!^2 * ((n+1)*k)!^2).

It is known that both row 0, the central binomial numbers, and row 1, the de Bruijn numbers, satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and all positive integers n and r. We conjecture that all the row sequences of the table satisfy the same supercongruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 27 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016296 | OEIS-A364509

Square array read by ascending antidiagonals: T(n,k) = (2*k)!/k!^2 * ((2*n*k)! * ((n + 2)*k)!)/((n*k)! * ((n + 1)*k)!^2) for n, k > = 0.

It is known that both row 0, the squares of the central binomial numbers, and row 1, the de Bruijn numbers, satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and all positive integers n and

r. We conjecture that all the row sequences of the table satisfy the same supercongruences [added Oct 11 2024: follows from Meštrović, Section 6, equation 39, since T(n, k) = binomial(2*k, k) * binomial(2*n*k, n*k) * binomial((n+2)*k, k)/binomial((n+1)*k, k)].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016297 | OEIS-A364511

a(n) = (6*n)!*(5*n)!*(2*n)! / ((4*n)!^2 * (3*n)! * n!^2).

Conjecture: the supercongruences a(n*p^r) == a(n*p^(r-1)) (mod p^(3^r)) hold for all primes p >= 5 and all positive integers n and r [added Oct 11 2024: follows from Meštrović, Section 6, equation 39, since a(n) = binomial(6*n, 3*n)*binomial(5*n, n)* binomial(2*n, n)/binomial(4*n, n)].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016298 | OEIS-A364513

Square array read by ascending antidiagonals: T(n,k) = [x^k] (1 - x)^(2*k) * Legendre_P(n*k-1, (1 + x)/(1 - x)) for n, k >= 0.

The central binomial numbers A000984, row 1 unsigned, satisfy the supercongruences A000984(n*p^r) == A000984(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and all positive integers n and r. We conjecture that each row sequence of the table satisfies the same supercongruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 31 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016299 | OEIS-A364514

a(0) = 1 and a(n) = [x^n] (1 - x)^(2*n) * Legendre_P(n-1, (1 + x)/(1 - x)) for n >= 1.

Conjecture: the supercongruences a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) hold for all primes p >= 5 and all positive integers n and r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Aug 02 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016300 | OEIS-A364518

Square array read by ascending antidiagonals: T(n,k) = [x^(2*k)] ((1 + x)^(n+2)/(1 - x)^(n-2))^k for n, k >= 0.

It is known that the unsigned version of row 0 (the central binomial numbers A000984) and row 2 satisfy the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and all positive integers n and r. We conjecture that all the row sequences of the table satisfy the same supercongruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Aug 07 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016301 | OEIS-A364519

Square array read by ascending antidiagonals: T(n,k) = [x^(3*k)] ((1 + x)^(n+3)/(1 - x)^(n-3))^k for n, k >= 0.

It is known that A005810, the unsigned version of row 1, satisfies the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and all positive integers n and r. We conjecture that each row sequence of the table satisfies the same supercongruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Aug 07 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016302 | OEIS-A364554

a(n) = number of primes of the form T(k)/n, for some k, where T(k)=A000217(k) is a triangular number.

Conjecture: a(n) = number of primes in the union of sets {(2*r -+ 1)/d; (r -+ 1)/(2*d)}, with d divisor of n and r = n/d.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lamine Ngom_, Jul 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016303 | OEIS-A364555

Numbers m such that no triangular number is m times a prime.

Conjecture: Numbers m such that there is no prime number in the union of all sets {(2*r -+ 1)/d; (r -+ 1)/(2*d)}, where d is some divisor of m and r = m/d.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lamine Ngom_, Jul 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016304 | OEIS-A364581

Numbers k such that the number of iterations of psi(phi(x)) starting at x = k and terminating when psi(phi(x)) = x (k is counted), -1 otherwise is the same for phi(psi(k)).

Conjecture: For each a(n), n > 1, a(n)*7 is a term. Conjecture: For each even a(n), a(n)*2 is a term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Torlach Rush_, Jul 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016305 | OEIS-A364608

Smallest k such that there are as many 0's as 1's in the binary representation of (2*n+1)^k, or -1 if no such k exists.

Conjecture: a(n) != -1 for all n. For even m, the proportion of 1's in the binary representation of m^k conjecturally converges to less than 1/2, so unless the numbers of 0's and 1's are equal for some small k, it is likely that no such k exists. For m = 2, 6, 10, 12, 14, 38, 42, 44, 46, ... there do exist such small k = 1, 3, 1, 1, 3, 1, 1, 1, 5, ..., respectively.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pontus von Brömssen_, Jul 30 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016306 | OEIS-A364615

Numbers k such that the average of the decimal digits of 2^k is closer to 9/2 (the expected average for random digits) than for any smaller power of 2.

Conjecture: for each term k > 1, digitsum(2^k) - (9/2)*number_of_digits(2^k) = 1/2 if k is odd, -1/2 if k is even. - _Jon E. Schoenfield_, Jul 30 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pontus von Brömssen_ and _Jon E. Schoenfield_, Jul 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016307 | OEIS-A364639

Irregular triangle read by rows: T(n,k) = A237591(n,k) - A237591(n-1,k).

Conjecture: column k gives 1 together with period A002378(k) repeat S_k, where S_k is the k-th row of the triangle A364746. The conjecture of _Omar E. Pol_ is true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jul 30 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016308 | OEIS-A364648

Starting position of the first occurrence of the longest monochromatic arithmetic progression of difference n in the Fibonacci infinite word (A003849).

Conjecture: for n>0,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gandhar Joshi_, Jul 31 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016309 | OEIS-A364654

Numbers which are the sum or difference of two seventh powers.

Don Zagier's conjecture that the polynomial x^7 + 3y^7 is injective on rational numbers is equivalent to the non-existence of any term in this sequence that is exactly 3 times another term in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Caveney_, Jul 31 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016310 | OEIS-A364663

a(n+1) = a(|n-a(n)*a(n-1)|)+1; a(0) = 0.

a(n) ~ (3*n)^(1/3) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rok Cestnik_, Aug 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016311 | OEIS-A364675

Number of integer partitions of n whose nonzero first differences are a submultiset of the parts.

Conjecture: For subsets of {1..n} instead of partitions of n we have A101925. Conjecture: The strict version is A154402.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Aug 04 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016312 | OEIS-A364746

Irregular triangle read by rows T(n,k), n >= 1, 1 <= k <= A002378(n), which is mentioned in the conjecture of A364639 (see Comments lines for definition).

Irregular triangle read by rows T(n,k), n >= 1, 1 <= k <= A002378(n), which is mentioned in the conjecture of A364639 (see Comments lines for definition). It is conjectured that 1 together with infinitely many copies of the n-th row gives the n-th column of the triangle A364639.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 05 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016313 | OEIS-A364789

Initial digit of (n^n)^n (A002489(n)).

The author conjectures that this sequence obeys the well-known Benford's law.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Aug 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016314 | OEIS-A364822

Expansion of e.g.f. cosh(x) / (1 - 2*sinh(x)).

Conjectures: For p prime (p > 2), a(p) == 2 (mod p).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mélika Tebni_, Nov 07 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016315 | OEIS-A364824

Index of prime(n) in A067836, or -1 if prime(n) does not occur in it.

If the conjecture holds that A067836 contains only primes, then a(A062894(n)) = n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bert Dobbelaere_, Aug 09 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016316 | OEIS-A364847

Number of conjugacy classes in the group SL(2, Z_n), up to conjugacy in GL(2, Z_n).

Conjecture: a(n) = Sum_{d|n} d*A060594(n/d). - _Ridouane Oudra_, Feb 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robin Visser_, Aug 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016317 | OEIS-A364855

Initial digit of 3^(3^n) (A055777(n)).

The author conjectures that the distribution of the initial digits of the present sequence obey Benford's law or Zipf's law (see links).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Aug 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016318 | OEIS-A364883

Consider the Fermat quotient for base n: Fq(n,k) = (n^(p - 1) - 1)/p, where p = prime(k), for k >= 1. a(n) is the least k >= 1 such that Fq(n,j) is divisible by n^2 - 1 for all j >= k.

Conjecture: numbers appear in the sequence only a finite number of times. Terms appear in runs of length 1, 2, or 3, never more. The first time a term k appears is when the index is even. The terms appear for the first time in their natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Aug 17 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016319 | OEIS-A364892

Row sums of A364891.

Conjecture: lim_{n->oo} a(n)/A000041(n) = 1/4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Aug 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016320 | OEIS-A364902

Let x, y be the greatest exponents of 2, 3 respectively such that 2^x, 3^y do not exceed n and let k_2, k_3 be n - 2^x, and n - 3^y respectively. Then for n such that k_2 = 0 or k_3 = 0, a(n) = n, else a(n) is the least novel number Min{p*a(k_2), q*a(k_3)}, where p, q are primes not equal to either 2 or 3.

Conjectures: (i). This is a permutation of A047253 with primes in order; (ii). All terms between consecutive prime terms, prime(k), prime(k+1) are prime(k)-smooth.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_ and _Michael De Vlieger_, Sep 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016321 | OEIS-A364904

a(n) = |Aut^n(C_32)|: order of the group obtained by applying G -> Aut(G) n times to the cyclic group of order 32.

The sequence {Aut^n(C_m):n>=0} is well-known for m <= 31. It is conjectured that |Aut^n(C_32)| tends to infinity as n goes to infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016322 | OEIS-A364917

For each n, if the sequence defined by G_0 = C_n, G_k = Aut(G_{k-1}) for k >= 1 stabilizes, then a(n) is the order of G_k for sufficiently large k; otherwise a(n) = 0. Here C_n is the cyclic group of order n.

The following two sequences are conjectured to be correct and to stabilize at the last term:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016323 | OEIS-A364933

a(n) = Sum_{k=1..n} A191898(n,k)*[A191904(n,k) = A191898(n,k)].

Conjecture: a(n) = A057859(n) - A008472(n) - 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Aug 13 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016324 | OEIS-A364958

Fixed points of A356867, where A356867 is Sycamore's Doudna variant D(3).

Conjecture: All terms are of the form k*3^n, where k = 1,2,8,91, and n >= 0. - _David James Sycamore_, Aug 16 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_ and _Antti Karttunen_, Sep 15 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016325 | OEIS-A364969

a(n) = a(a(n-1)) if n is even, a(n) is the number of times a(n-1) occurs in the sequence so far if n is odd, with a(1) = 1.

Conjecture: All positive integers appear in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tanmaya Mohanty_, Oct 23 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016326 | OEIS-A365000

a(n) = n for n <= 2; for n > 2, rad(a(m)) != rad(a(n)), m < n, implies a(n+1) is the least novel k such that rad(k) | a(n), rad(a(m)) = rad(a(n)), m < n, implies a(n+1) = q*a(n), where q is the least novel prime such that gcd(q, a(n)) = 1 and rad(a(m)) != rad(q*a(n)), m < n.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_ and _David James Sycamore_, Sep 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016327 | OEIS-A365009

Semiprimes that are the concatenation of two or more semiprimes.

Conjecture: The fraction of semiprimes <= N that are in this sequence goes to 1 as N -> infinity. What is the first N for which that fraction >= 1/2?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_ and _Robert Israel_, Aug 15 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016328 | OEIS-A365020

Solutions k to the Diophantine equation k^5 = Sum_{i=1..7} y_i^5 with positive y_i.

Conjecture: sequence contains all integers >= 78 and a(n) = n+42 for n >= 36. - _Chai Wah Wu_, Aug 17 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Aug 16 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016329 | OEIS-A365021

a(n) is the largest prime of the form P+1 where P divides prime(n)# and p# denotes the product of all primes <= p.

Conjecture: a(n) > (1/2) * prime(n-1)#.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Aug 16 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016330 | OEIS-A365025

Square array read by antidiagonals: T(n, k) := (k/2)!/k! * ((2*n+1)*k)! * ((2*n+1/2)*k)! / ((n*k)!^2 * ((n+1/2)*k)!^2) for n, k >= 0.

Conjecture: each row sequence of the table satisfies the supercongruences u(n*p^r) == u(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and all positive integers n and r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Aug 17 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016331 | OEIS-A365028

a(n) = Sum_{k = 0..n} (-1)^(n+k) * binomial(n,k)*binomial(n+k-1,n)* binomial(3*n+k-1,n).

The central binomial coefficients A000984(n) satisfy the supercongruences A000984(n*p^r) == A000984(n*p^(r-1)) (mod p^(3*r)) for all primes p >= 5 and all positive integers n and r (see Meštrović, equation 39). We conjecture that the present sequence satisfies the same supercongruences. More generally, for M a positive integer and integers A, B and C, we conjecture that the sequence whose n-th term is given by Sum_{k = 0..M*n} (-1)^(n+k) * binomial(A*n,k)*binomial(B*n+k-1,n)*binomial(C*n+k-1,n) also satisfies the same supercongruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016332 | OEIS-A365048

a(n) is the number of steps required for the n-th odd prime number to reach 3 when iterating the following hailstone map: If P+1 == 0 (mod 6), then the next number = smallest prime >= P + (P-1)/2; otherwise the next number = largest prime <= (P+1)/2.

Conjecture: This hailstone operation on odd prime numbers will always reach 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Najeem Ziauddin_, Oct 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016333 | OEIS-A365059

a(1) = 2; for n > 2, a(n) is the smallest positive number that has not yet appeared that is a multiple of A008472(a(n-1)), the sum of the distinct primes dividing a(n-1).

In the first 500000 terms the only fixed points are 38 and 209, although it is likely more exist. In the same range the smallest missing numbers are 311, 313, 337. The sequence is conjectured to be a permutation of the integers >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Aug 19 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016334 | OEIS-A365060

a(1) = 2; for n > 2, a(n) is the smallest positive number that has not yet appeared that has a common factor with A008472(a(n-1)), the sum of the distinct primes dividing a(n-1).

In the first 100000 terms the only fixed point is 9, and it is likely no more exist. In the same range the smallest missing numbers are 503, 839, 877. The sequence is conjectured to be a permutation of the integers >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Aug 19 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016335 | OEIS-A365078

a(n) is the least divisor (d) of prime(n)# such that prime(n)# / d + 1 is prime where p# denotes the product of all primes <= p.

Conjecture: a(n) < 2*prime(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Aug 20 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016336 | OEIS-A365092

Write out the canonical factorization of n and factorize the exponents in the factorization, the exponents in the factorizations of the exponents, ... until there are only prime numbers left. Replace each p in the factorization by (p-1)+1 and factorize each p-1 by the same process if p-1 > 1. Continue this process until there are only 1s left. a(n) is the number of 1s used.

Conjecture: for k >= 2, the maximum n such that a(n) = k is n = A001144(k). n = A365093(k) is the minimum such n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016337 | OEIS-A365093

Smallest k such that A365092(k) = n.

For m,n >= 2, it is easy to see that if gcd(a(m),a(n)) = 1, then a(m+n) <= a(m)*a(n). This is conjectured to be true for all m,n. In particular, it is conjectured that a(n+2) <= 2*a(n). If a(m+n) <= a(m)*a(n) is true, then by Fekete's subadditive lemma, we have lim_{n->oo} a(n)^(1/n) = inf_{n>=1} a(n)^(1/n) <= a(29)^(1/29) = 2879^(1/29) = 1.3160857758...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Aug 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016338 | OEIS-A365095

Expansion of g.f. A(x) satisfying [x^(n-1)] (1 + (n-1)*x*A(x)^2)^n / A(x)^n = 0 for n > 1.

(2) [x^(n-1)] (1 + (k*n-1)*x*A(x)^2)^n / A(x)^n is divisible by n^2 for n > 0 and all integer k (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016339 | OEIS-A365214

Least k such that the binary representation of 3^k has exactly n 1's, or -1 if no such k exists.

The first 22 terms are from Dimitrov and Howe (2021). After a(22), the sequence continues (with the -1's conjectural but very likely correct) -1, 26, 28, 31, 36, 40, 34, 27, -1, 35, 49, 33, 53, 38, -1, 42, 41, 57, 48, 52, -1, -1, 46, 51, 67, 59, 62, -1, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pontus von Brömssen_, Aug 26 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016340 | OEIS-A365215

Largest k such that the binary representation of 3^k has exactly n 1's, or -1 if no such k exists.

The first 22 terms are from Dimitrov and Howe (2021). After a(22), the sequence conjecturally but very likely continues -1, 26, 30, 32, 36, 40, 34, 27, -1, 39, 49, 45, 53, 38, -1, 47, 56, 57, 50, 58, -1, -1, 66, 51, 67, 59, 62, -1, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pontus von Brömssen_, Aug 26 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016341 | OEIS-A365259

Lexicographically earliest sequence of distinct positive numbers such that, for n > 2, a(n) shares a factor with a(n-1) and a(n+a(n)).

In the first 10000 terms the fixed points are 1, 2, 10, 65, 1261, 6527, although more likely exist. The sequences is conjectured to be a permutation of the positive integers. Unlike A064413 the primes do not occur in their natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016342 | OEIS-A365265

Numbers k for which sqrt(k/2) divides k and the symmetric representation of sigma(k) consists of a single part and its width at the diagonal equals 1.

Conjecture: Let a(n) = 2^(2i+1) * s^2, i>=0 and s odd, be a number in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Aug 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016343 | OEIS-A365367

Number of steps for iteration of map x -> (5/3)*round(x) to reach an integer > n when started at n, or -1 if no such integer is ever reached.

Conjecture: an integer will always be reached, i.e. a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Sep 02 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016344 | OEIS-A365368

First integer > n reached under iteration of map x -> (5/3)*round(x) when started at n, or -1 if no such integer is ever reached.

Conjecture: an integer will always be reached, i.e. a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Sep 02 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016345 | OEIS-A365383

Triangle read by rows where T(n,k) is the number of integer partitions of n that can be linearly combined with nonnegative coefficients to obtain k.

Conjecture: The rows eventually become periodic with period n if extended further. For example, row n = 8 begins:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Sep 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016346 | OEIS-A365397

a(n) = 64 + A000720(n) - A365398(n).

It is conjectured that A365339(n) - PrimePi(n) = 64 for all n >= 31957 (Pollack et al.). Does a similar relation apply if one replaces Euler's totient by the sum of divisors function in A365339? In particular, note remark (4.4) by Terence Tao in the linked paper.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Sep 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016347 | OEIS-A365400

a(n) = 64 + A000720(n) - A365339(n).

It is conjectured that A365339(n) = PrimePi(n) + 64 for all n >= 31957 (Pollack et al.). Assuming this conjecture a(n) = 0 for n > 31956.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Sep 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016348 | OEIS-A365429

a(n) = 1 if the divisors of n increase by a factor of at most 2, otherwise 0.

a(n) <= A322860(n). [Conjectured, see comment in A174973]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 22 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016349 | OEIS-A365436

a(2^k) = 2^k for all k >= 0. let 2^r be the smallest power of 2 which exceeds n, then a(n) = the least novel m*a(k), where k = 2^r-n, and m is not a prior term.

Conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Nov 09 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016350 | OEIS-A365440

Square array read by upward antidiagonals: T(n,k) is the n-th number j with the property that the parts of the symmetric representation of sigma(j) are two s-gon of width 1, where s = 2^(k+1), n >= 1, k >= 1.

Conjecture 1: column k gives the row numbers of the triangle A364639 where the rows are [1, A036563(k+1) zeros, -1, 1] or where the rows start with [1, A036563(k+1) zeros, -1, 1] and the remaining terms are zeros. Conjecture 2: every column gives a subsequence of A246955. Conjecture 3: the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Sep 25 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016351 | OEIS-A365453

Lexicographically earliest sequence of distinct positive numbers such that, for n > 2, a(n) shares a factor with a(n-1), a(n + a(n)) and, if a(n) < n, with a(n - a(n)).

Other than the first two terms there are no other fixed points in the first 500000 terms, and it is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 04 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016352 | OEIS-A365468

Number of n X n nonnegative integer arrays with upper left entry 0, lower right entry n - 5, every value within 4 of its king-move distance from the upper left, and every value increasing by 0 or 1 with every step right, diagonally southeast, or down.

Conjecture: a(n) ~ (2^2 / (3^16 * Pi^2)) * 256^n / n^6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Dougherty-Bliss_, Sep 04 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016353 | OEIS-A365469

Number of n X n nonnegative integer arrays with upper left entry 0, lower right entry n - 6, every value within 5 of its king-move distance from the upper left, and every value increasing by 0 or 1 with every step right, diagonally southeast, or down.

Conjecture: a(n) ~ (2^4 / (3^23 * Pi^2)) * 1024^n / n^10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Dougherty-Bliss_, Sep 04 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016354 | OEIS-A365485

a(n) = A365399(10^n).

a(n)/10^n appears to be decreasing. Conjecture: a(n)/10^n converges to a nonzero value.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Sep 05 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016355 | OEIS-A365498

Dirichlet g.f.: zeta(s) * Product_{p prime} (1 + 1/p^s - 1/p^(3*s)).

Following formulas have been conjectured for this sequence by Sequence Machine, with each one giving the first 1000000 terms correctly:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Sep 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016356 | OEIS-A365536

a(n) = n for n <= 2. Thereafter a(n) is the least novel multiple of the greatest prior term which is coprime to a(n-1).

Conjecture: All terms are 7-smooth numbers. Is this a permutation of A002473?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Sep 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016357 | OEIS-A365555

Expansion of e.g.f. 1 / (7 - 6 * exp(x))^(1/3).

O.g.f. (conjectural): 1/(1 - 2*x/(1 - 7*x/(1 - 8*x/(1 - 14*x/(1 - 14*x/(1 - 21*x/(1 - ... - (6*n - 4)*x/(1 - 7*n*x/(1 - ...))))))))) - a continued fraction of Stieltjes-type (S-fraction).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Sep 09 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016358 | OEIS-A365556

Expansion of e.g.f. 1 / (7 - 6 * exp(x))^(2/3).

O.g.f. (conjectural): 1/(1 - 4*x/(1 - 7*x/(1 - 10*x/(1 - 14*x/(1 - 16*x/(1 - 21*x/(1 - ... - (6*n - 2)*x/(1 - 7*n*x/(1 - ...))))))))) - a continued fraction of Stieltjes-type (S-fraction). - _Peter Bala_, Sep 24 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Sep 09 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016359 | OEIS-A365557

Expansion of e.g.f. 1 / (7 - 6 * exp(x))^(5/6).

O.g.f. (conjectural): 1/(1 - 5*x/(1 - 7*x/(1 - 11*x/(1 - 14*x/(1 - 17*x/(1 - 21*x/(1 - ... - (6*n - 1)*x/(1 - 7*n*x/(1 - ...))))))))) - a continued fraction of Stieltjes-type (S-fraction). - _Peter Bala_, Sep 24 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Sep 09 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016360 | OEIS-A365573

Number of primes between prime(n) and prime(n)+log(prime(n)), exclusive.

Conjecture: Limit_{N->oo} (Ratio_{n=1..N} a(n)=0) = 1/e (A068985).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Sep 09 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016361 | OEIS-A365574

Expansion of g.f. A(x) satisfying [x^(n-1)] (1 + (n+1)*x*A(x))^n / A(x)^n = n*(n+2)^(n-2) for n > 1.

Conjecture 1: a(k) is odd iff k = 2^n - 2 for n >= 1.
Conjecture 2: a(2^n - 2) == 3 (mod 16) for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 11 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016362 | OEIS-A365593

Number of n X n Boolean relation matrices such that every block of its Frobenius normal form is either a 0 block or a 1 block.

Conjecture: lim_n->inf a(n)/(A003024(n)*2^n) = 1. In other words, almost all of the relations counted by this sequence have n strongly connected components. - _Geoffrey Critzer_, Sep 30 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Critzer_, Sep 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016363 | OEIS-A365619

a(n) is the least integer k such that A366110(k) = n, or 0 if there is no such k.

The nonzero values are sure, but the zeros are conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Nov 02 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016364 | OEIS-A365625

a(0) = 0, a(1) = 1. let i = a(n-2) and j = a(n-1), then if i,j have a digit in common a(n) is the least novel number having no digit in common with either i or j. If i,j have no common digit, a(n) is the least novel number having a digit in common with at least one of i or j. All digits are decimal.

The sequence is conjectured to be finite. It will end if for some m, a(m) is a pandigital number, or if the set {i,j} of distinct digits of i and j contains all nonzero digits, whichever happens first. In the latter case the definition would require the next term to consist entirely of 0's, but since 0 is already a term this is impossible. A sequence like this can be made using any base.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Sep 13 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016365 | OEIS-A365643

Number of permutations whose reverse-complement shares the same recording tableau in the Robinson-Schensted correspondence.

This is an open problem from Martin's "Algebraic Combinatorics" lecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dang-Son Nguyen_, Sep 14 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016366 | OEIS-A365658

Triangle read by rows where T(n,k) is the number of integer partitions of n with k distinct possible sums of nonempty submultisets.

Conjecture: Positions of strictly positive rows are given by A048166.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Sep 16 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016367 | OEIS-A365688

Primitive solutions k to k^2 = u^4 + v^4 + w^4, with u, v, w > 0.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jud McCranie_, Sep 16 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016368 | OEIS-A365739

a(n)=A365339(10^n)-A365740(10^n).

Pollack et al. conjectured that a(n) > 2 for n > 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Sep 17 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016369 | OEIS-A365740

Length of the longest subsequence of {m: 1<=m<=n, m not prime} on which the Euler totient function phi A000010 is nondecreasing.

Pollack et al. conjectured that a(n) < A365339(n)-2 for all n >= 31957.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Sep 17 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016370 | OEIS-A365777

Expansion of e.g.f. (exp(2*x) / (2 - exp(2*x)))^(3/4).

Conjecture: the sequence a(n) taken modulo a positive integer k is eventually periodic with the period dividing phi(k). For example, the sequence taken modulo 21 begins [1, 3, 15, 12, 18, 6, 9, 3, 15, 12, 18, 6, 9, 3, 15, 12, 18, 6, 9, ...] with an apparent period of 6 (= phi(21)/2) starting at n = 1. - _Peter Bala_, Oct 23 2025 Conjectural o.g.f. as a continued fraction of Stieltjes type: 1/(1 - 3*x/(1 - 2*x/(1 - 7*x/(1 - 4*x/(1 - 11*x/(1 - 6*x/(1 - 15*x/(1 - 8*x/(1 - 19*x/(1 - ...))))))))). - _Peter Bala_, Oct 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Nov 16 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016371 | OEIS-A365782

Expansion of e.g.f. 1 / (3 - 2 * exp(2*x))^(1/4).

Conjecture: the sequence a(n) taken modulo a positive integer k is eventually periodic with the period dividing phi(k). For example, the sequence taken modulo 21 begins [1, 1, 7, 16, 13, 7, 10, 1, 7, 16, 13, 7, 10, 1, 7, 16, 13, 7, 10, ...] with an apparent period of 6 (= phi(21)/2) starting at n = 1. - _Peter Bala_, Oct 23 2025 Conjectural o.g.f. as a continued fraction of Stieltjes type: 1/(1 - x/(1 - 6*x/(1 - 5*x/(1 - 12*x/(1 - 9*x/(1 - 18*x/(1 - 13*x/(1 - 24*x/(1 - ...))))))))). - _Peter Bala_, Oct 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Nov 16 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016372 | OEIS-A365794

Expansion of e.g.f. 1 / (3 - 2 * exp(2*x))^(3/4).

Conjecture: the sequence a(n) taken modulo a positive integer k is eventually periodic with the period dividing phi(k). For example, the sequence taken modulo 7 begins [1, 3, 6, 5, 3, 6, 5, 3, 6, 5, ...] with an apparent period of 3 (= phi(7)/2) starting at n = 1. - _Peter Bala_, Oct 23 2025 Conjectural o.g.f. as a continued fraction of Stieltjes type: 1/(1 - 3*x/(1 - 6*x/(1 - 7*x/(1 - 12*x/(1 - 11*x/(1 - 18*x/(1 - 15*x/(1 - 24*x/(1 - ...))))))))). - _Peter Bala_, Oct 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Nov 16 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016373 | OEIS-A365847

Expansion of (1/x) * Series_Reversion(x*(1-x)^4/(1+x)^4).

Conjecture: g.f.: B^4, where B is the g.f. of A260332.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Sep 20 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016374 | OEIS-A365861

a(1) = 4; for n > 1, a(n) is the first semiprime k not yet listed such that |k - a(n-1)| and k + a(n-1) are also semiprimes.

Conjecture: this is a permutation of A001358.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zak Seidov_ and _Robert Israel_, Sep 20 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016375 | OEIS-A365892

a(n) is the index of the n-th term of A365876 that includes at least one equation with at least one integer solution.

Conjectures in formula section hold for 2<=n<=300. - _Chai Wah Wu_, Oct 05 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Sep 22 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016376 | OEIS-A365959

Decimal expansion of Sum_{k>=2} zeta(k)/k^2.

Conjecture: Equals 2 - A001620 - Pi^2/12 + Integral_{x=0..oo} (2*x*arccot(x) - log(1/x^2 + 1))*log(1 - exp(-2*Pi*x))/(2*Pi*(x^2 + 1)) dx. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Sep 23 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016377 | OEIS-A365965

Numbers k such that A139315(k) = 0 but k is not in A138511.

Conjecture: includes all numbers that are 4 times a prime number greater than 13. If true, 188 would qualify for this sequence, but we don't know if any of 183-187 are in this sequence. - _J. Lowell_, Sep 27 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Lowell_, Sep 23 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016378 | OEIS-A365994

The n-th term of the sequence is the last term of n's trajectory under the "multiply with zero" rules explained in A365993.

It is conjectured that all trajectories will stop quite rapidly, mostly because the probability for 0 to appear in a divisor of a(n) increases with the size of a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Giorgos Kalogeropoulos_, Sep 25 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016379 | OEIS-A366021

a(1) = 1; a(2) = 2; for n > 2, a(n) is the smallest positive number that has not yet appeared that shares a factor with a(n-1) and, if a(n) < n, with a(n - a(n)).

The majority of terms lie near the line a(n) = 1.1*n. The only exceptions are the prime valued terms whose appearance in the sequence is delayed relative to their magnitude, e.g. a(1835) = 61. Other than the first two terms there are no other fixed points in the first 100000 terms, and it is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 26 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016380 | OEIS-A366029

Numbers k > 1 such that (k-1)^(2^k-2) == 1 (mod (2^k-1)*k^2).

Conjecture: for k > 1, (2^k-1)*k^2 divides (k-1)^(2^k-2)-1 if and only if 2^k-1 is (a Mersenne) prime or k = 2^(2^m)+1 (is a Fermat number). If my conjecture is true, primes in this sequence that are not Mersenne exponents are Fermat prime numbers k = 2^(2^m)+1 for m > 2, namely k = 257, 65537 and there are probably no more.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Oct 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016381 | OEIS-A366048

For n >= 1, a(n) is the least k >= 1 such that 1/d(k) + ... + 1/d(k + n - 1) is an integer, d(i) = A000005(i).

Conjecture: The sum 1/d(k) + ... + 1/d(k + n - 1) = C, C an integer, exists for all k >= 1, n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Sep 27 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016382 | OEIS-A366058

Number of n-step self-avoiding walks on a 3D cubic lattice where no step is to a lattice point closer to the origin than the current point.

Conjectured: a(n) = 6*(4*3^(n-1) - 4*2^(n-1) + 1), for n > 0. a(n) = Sum_{i=1..d} (-1)^(d-i) * binomial(d,i) * 2^i * i^n, where d=3, n>=1, which simplifies to 8*3^n - 12*2^n + 6, equivalent to conjectured formula (and row 3 of A371064). - _Shel Kaphan_, Mar 09 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Dec 15 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016383 | OEIS-A366063

Irregular triangle read by rows: T(n,k) is the number of partitions of n that have depth k.

Suppose that P is a partition of n. Let x(1), x(2),...,x(k) be the distinct parts of P, and let m(i) be the multiplicity of x(i) in P. Let f(P) be the partition [m(1)*x(1), m(2)*x(2),...,x(k)*m(k)] of n. Define c(0,P) = P, c(1,P) = f(P), ..., c(n,P) = f(c(n-1,P)), and define d(P) = least n such that c(n,P) has no repeated parts; d(P) is as the depth of P, as defined in A237685. Clearly d(P) = 0 if and only if P is a strict partition, as in A000009. Conjecture: if d >= 0, then 2^d is the least n that has a partition of depth d.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016384 | OEIS-A366105

a(n) is the number of parts in the n-th partition of n when the partitions are listed in graded reverse lexicographic order (cf. A080577, as in Mathematica).

Conjecture 1. Every integer m > 1 occurs infinitely many times. (For example, 2 occurs for n = 2,8,13,20,31,46,68,....) Conjecture 2. Let f(n) be the greatest (i.e., the first) part in the n-th partition of n. Then for every integer m, there exists an index i such that f(i+1), f(i+2), ..., f(i+m) are consecutive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Oct 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016385 | OEIS-A366111

a(1) = 1; a(2) = 2; for n > 2, a(n) is the smallest positive number that has not yet appeared such that |a(n) - a(n-1)| is a divisor of a(n)*a(n-1), and where |a(n) - a(n-1)| > 1.

In the first 10000 terms the fixed points are 16, 21, 48, 98, 105, 322, 3088, 7659, although more likely exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016386 | OEIS-A366133

Triangle read by rows: coefficients in expansion of another Asveld's polynomials Pi_j(x).

Conjectures: For 0 < k < p and p prime, T(p,k) == 0 (mod p).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mélika Tebni_, Sep 30 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016387 | OEIS-A366134

Number of primes between prime(n) and prime(n)+2*log(prime(n)), exclusive.

Conjecture: Limit_{N->oo} (Ratio_{n=1..N} a(n)=0) = 1/e^2 (A092553).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Sep 30 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016388 | OEIS-A366186

Positive integers k such that the third derivative of the k-th Bernoulli polynomial B(k, x) contains only integer coefficients.

As a published result on Oct 02 2023 (cf. A366169), all such sequences regarding higher derivatives of the Bernoulli polynomials having only integer coefficients are finite. We have an infinite chain of subsets: A094960 subset of A366169 subset of A366186 subset of A366187 subset of A366188 subset of ... . See Kellner 2023 (Theorem 13, Conjecture 14, and S_3 (this sequence)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Oct 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016389 | OEIS-A366189

a(n) is the positive integer k such that the k-th derivative of the n-th Bernoulli polynomial B(n, x) contains only integer coefficients but no lower derivative of B(n, x) has this property.

Conjecture: Every integer appears in this sequence only a finite number of times. (This generalizes the conjectures made in A366186-A366188.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Oct 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016390 | OEIS-A366190

Minimal lengths of prime knots formed by orthogonal unit line segments of the cubic lattice.

Conjecture: All even numbers >= 40 will appear in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tamas Sandor Nagy_ and _Thomas Scheuerle_, Oct 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016391 | OEIS-A366253

Table read by antidiagonals: Place k points in general position on each side of a regular n-gon and join every pair of the n*(k+1) boundary points by a chord; T(n,k) (n >= 3, k >= 0) gives number of regions in the resulting planar graph.

Conjectured:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Nov 09 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016392 | OEIS-A366278

Perfect powers k such that A052409(k) is equal to A052409(A366275(k)).

Conjecture: all terms are squares, and no higher powers occur.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016393 | OEIS-A366351

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number that has not yet appeared whose binary string value contains at least one binary prime factor of a(n-1).

This is the base-2 equivalent of A365703. Unlike that sequence is it conjectured all numbers appear. The only fixed point in the first 10000 terms is 6, and it is likely no more exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Oct 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016394 | OEIS-A366362

Triangle read by rows: T(n,k) = Sum_{y=1..n} Sum_{x=1..n} [GCD(f(x,y), n) = k], where f(x,y) = x^3 - x^2 - y^2 - y.

Conjecture: T(n,n) = A060457(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Oct 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016395 | OEIS-A366410

Number of linearly independent solutions to the neighbor sum problem on a cubical (3n-1) X (3n-1) X (3n-1) chessboard.

It is conjectured that if a(n) is nonzero, then n is divisible by 2 or 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sayan Dutta_, Oct 09 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016396 | OEIS-A366412

Number of nontrivial solutions to the P^*_k problem in base n.

It has been conjectured that for a given composite base B, if there are no new nontrivial solutions (except for extensions) in (2k + 1) digits, then there would be no new solutions in (2k + 3) digits (see Saha et al. link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sayan Dutta_, Oct 09 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016397 | OEIS-A366415

a(n) is the number of exterior top arches (no covering arch) for semi-meanders in generation n+1 that are generated by semi-meanders with n top arches and floor(n/2) exterior top arches using the exterior arch splitting algorithm.

b(n) = ((n-4)*2^floor((n-1)/2)+2)*floor(n/2) is the number of exterior top arches for all semi-meander solutions with n top arches and floor(n/2) exterior top arches. Conjecture: for n>=5, lim_{n->oo} a(n)/b(n) = 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger Ford_, Oct 10 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016398 | OEIS-A366704

Number of sphinx tilings of T(n+12) with a central T(n) defect where T(k) is an equilateral triangle with side length k.

Conjecture: a(3*k + 2) = 144. Conjecture: a(3*k + 1) = 180*k + 216.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Craig Knecht_, Oct 17 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016399 | OEIS-A366724

First index k where A366574(k) = n.

Conjectured: for n > 1, a(n) = (n-1)*(n^3 + 3*n^2 - 10*n + 72)/24 .

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Oct 17 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016400 | OEIS-A366833

Number of times n appears in A362965 (number of primes <= the n-th prime power).

Conjecture: a(n) can be only 1, 2, or 3 (with the first occurrences of 3 appearing at n = 4, 9, 30, 327 and 3512).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Oct 25 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016401 | OEIS-A366860

a(0) = 0; a(n) = n - a(floor(a(n-1)/3)).

Conjecture: a(n) ~ c*n, where c = 1/(1 + 1/(3 + 1/(1 + 1/(3 + ...)))) = (sqrt(21) - 3)/2 = 0.79128784747792...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Oct 26 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016402 | OEIS-A366870

a(0) = 0; a(n) = n - a(floor(a(n-1)/4)).

Conjecture: a(n) ~ c*n, where c = 1/(1 + 1/(4 + 1/(1 + 1/(4 + ...)))) = 2*(sqrt(2) - 1) = 0.82842712474619...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Oct 26 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016403 | OEIS-A366871

a(0) = 0; a(n) = n - a(floor(a(n-1)/5)).

Conjecture: a(n) ~ c*n, where c = 1/(1 + 1/(5 + 1/(1 + 1/(5 + ...)))) = (3*sqrt(5) - 5)/2 = 0.8541019662496845...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Oct 26 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016404 | OEIS-A366884

Number of branching factorizations of the least integer of each prime signature (A025487).

Sequence appears to be injective, but can it be proved? This would prove also the conjectures given in A277120 and A366377.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 02 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016405 | OEIS-A366891

Lexicographically earliest infinite sequence such that a(i) = a(j) => A365425(i) = A365425(j), A206787(A163511(i)) = A206787(A163511(j)) and A336651(A163511(n)) = A336651(A163511(j)) for all i, j >= 0.

Restricted growth sequence transform of the triplet [A365425(n), A206787(A163511(n)), A336651(A163511(n))], and also by conjecture, of sequence b(n) = A351040(A163511(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 04 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016406 | OEIS-A366908

Lexicographically earliest infinite sequence of distinct positive integers such that, for n > 1, a(n) shares a factor with n but does not equal n, while not sharing a factor with a(n-1).

For the terms studied the primes appear in their natural order except for 11 and 13 which are reversed. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Oct 27 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016407 | OEIS-A366932

Total number of edges formed after n points have been placed in general position on each edge of a triangle (as in A365929).

Conjecture: a(n) = 3*(3*n^4/2 - 2*n^3 + 3*n^2/2 + n + 1). [This is now a theorem - _N. J. A. Sloane_, Dec 31 2025]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 02 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016408 | OEIS-A366952

a(1) = 1, a(2) = 4; for n > 2, a(n) is the smallest positive number that has not yet appeared that shares a factor with n but does not equal n, and shares a factor with a(n-1).

The sequence is conjectured to be a permutation of the positive integers, although the primes typically take many terms to appear, e.g., a(95890) = 223. When a prime does appear it is often followed by a term that is significantly larger than the average-sized term. See the examples below. The primes do not occur in their natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Oct 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016409 | OEIS-A367015

Number of regions formed after n points have been placed in general position on each edge of a triangle (as in A365929).

Conjecture: a(n) = (9*n^4 - 12*n^3 + 15*n^2 + 4)/4. [This is now a theorem - _N. J. A. Sloane_, Dec 31 2025]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 01 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016410 | OEIS-A367052

Number of n X n matrices with elements {0, 1} whose characteristic polynomial has non-leading coefficients in {-1,0}.

Conjecture: The number of matrices with characteristic polynomial x^n - x^(n-1) is exactly n*A003024(n). (If so, (n+1)*A003024(n) is a lower bound for this sequence.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Nov 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016411 | OEIS-A367063

Least number k such that there exists a subset u of n distinct elements of {1..k} for which Sum_{i = 1..n} u(i) * prime(u(i)) mod Sum_{i = 1..n} u(i) = 0.

Conjecture: a(n) <= n + 3 for all n. - _Robert Israel_, Jun 07 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-Marc Rebert_, Nov 04 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016412 | OEIS-A367066

a(n) = ((Sum_{i=1..n} A367065(i))-2)/(n+2).

Conjecture: a(n) = floor(n/phi + 1/phi^3) - [n+2 = Fibonacci(2*j+1) for some j], where phi = (1+sqrt(5))/2 and [] is the Iverson bracket. - _Jon E. Schoenfield_, Nov 03 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zenan Sabanac_, Nov 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016413 | OEIS-A367082

a(1), a(2) = 2. Thereafter a(n) is the least novel multiple of the greatest prime which divides precisely one of a(n-1), a(n-2) but not the other. If no such prime exists a(n) is the least novel multiple of the smallest prime dividing neither a(n-1) nor a(n-2).

Conjecture: This is a permutation of the positive integers, with primes in order. Each prime p generates a trajectory T(p), dominated by multiples of p, see graph. Typically T(prime(k)) is initiated following the earliest term divisible by prime(k). Some trajectories are more prominent in the graph than others e.g. T(7),T(11),T(13) extend from p = prime(k) to prime(k+1)*prime(k). Others, e.g. T(3) terminate earlier but include prime(k)^2, whilst some do not reach as far as prime(k)^2 (T(5) has just two terms). Thus three categories of prime trajectory can be distinguished in the graph (respectively "full", "medium" and "small"). The graph resembles an inclined comb whose teeth correspond to full and medium trajectories; see Example.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_ and _Michael De Vlieger_, Nov 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016414 | OEIS-A367083

List of powers of 3 and powers of 4, in increasing order, starting with a(0) = 3^0 = 4^0 = 1.

We think these patterns are important for the study of Erdős's conjecture of a positive density of Sum(Pow({3,4})) mentioned in A327621.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Nov 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016415 | OEIS-A367118

Place n points in general position on each side of an equilateral triangle, and join every pair of the 3*n+3 boundary points by a chord; sequence gives number of regions in the resulting planar graph.

Conjecture: a(n) = (1/4)*(9*n^4 + 12*n^3 + 15*n^2 + 12*n + 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Nov 05 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016416 | OEIS-A367119

Place n points in general position on each side of an equilateral triangle, and join every pair of the 3*n+3 boundary points by a chord; sequence gives number of edges in the resulting planar graph.

Conjecture: a(n) = (3/2)*(3*n^4 + 4*n^3 + 3*n^2 + 4*n + 2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Nov 05 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016417 | OEIS-A367121

Place n points in general position on each side of a square, and join every pair of the 4*n+4 boundary points by a chord; sequence gives number of regions in the resulting planar graph.

Conjecture: a(n) = (17/2)*n^4 + 19*n^3 + (43/2)*n^2 + 14*n + 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Nov 05 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016418 | OEIS-A367122

Place n points in general position on each side of a square, and join every pair of the 4*n+4 boundary points by a chord; sequence gives number of edges in the resulting planar graph.

Conjecture: a(n) = 17*n^4 + 38*n^3 + 37*n^2 + 24*n + 8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Nov 05 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016419 | OEIS-A367147

Index of matching grid point in the bijection between two infinite triangular grids with one grid rotated by Pi/6 around the common point (0,0), using an enumeration of the grid points by A307014 and A307016.

While the method described only finds a minimum of the maximum distance of approximately 0.9659 by assigning the bijection partners using tiles, applying the Hopcroft-Karp algorithm to the bipartite graph corresponding to a sufficiently large section of the two infinite grids achieves significantly lower maximum distances. We conjecture that an upper bound for the maximum distance is sqrt(2)/2~=0.7071. See the corresponding link. A method that reduces the maximal occurring bijection distance to its conjectured minimum, and only requires local rearrangements, as described for the square grids in A307731, is currently not known in the present case of the triangular grids.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Klaus Nagel_ and _Hugo Pfoertner_, Nov 06 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016420 | OEIS-A367175

a(n) = Sum_{d|n} (-1)^[d is prime] * d, where [] denotes the Iverson bracket.

Conjecture: {k: a(k) divides sigma(k)} = {1, 2, 3, 6, 14, 15, 35}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Nov 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016421 | OEIS-A367182

Primes of the form (k-th primorial) - (k+1)st prime.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel D Gibson_, Nov 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016422 | OEIS-A367183

Table read by antidiagonals: Place k points in general position on each side of a regular n-gon and join every pair of the n*(k+1) boundary points by a chord; T(n,k) (n >= 3, k >= 0) gives number of vertices in the resulting planar graph.

Conjectured:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Nov 08 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016423 | OEIS-A367190

Table read by antidiagonals: Place k points in general position on each side of a regular n-gon and join every pair of the n*(k+1) boundary points by a chord; T(n,k) (n >= 3, k >= 0) gives number of edges in the resulting planar graph.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Nov 09 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016424 | OEIS-A367191

a(n) is the largest positive number k such that k^2 + k*n + n^2 is a perfect square, or 0 if no such k exists.

Conjectured formulas according to n mod 4 in first column.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-François Alcover_, Nov 09 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016425 | OEIS-A367306

Move bits in blocks in binary expansion of n where blocks are defined in Comments.

Conjecture: a(n) = A200714(p(n) + 1) where p(n) is an inverse permutation to A358654.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Nov 13 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016426 | OEIS-A367338

Comma-successor to n: second term of commas sequence if initial term is n, or -1 if there is no second term.

The following is a proof of a slight modification of a conjecture made by Ivan N. Ianakiev in A367341.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 15 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016427 | OEIS-A367341

Numbers without comma-successors: these are the numbers k such that if the commas sequence A121805 is started at k instead of 1, there is no second term.

(See link for proof.) This was stated without proof by _David W. Wilson_ in 2007 (see the Angelini link), and was conjectured (in a slightly less precise form) by _Ivan N. Ianakiev_, Nov 16 2023. This implies that the conjecture below is true, as well as the conjecture in A367342.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 15 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016428 | OEIS-A367342

a(n) = n+1 for n = 1 to 8; a(n) = 100 + a(n-8) for n = 9 to 16; thereafter a(8*i+j) = 10^(i+1) + a(8*(i-1)+j) for i >= 2, 1 <= j <= 8.

Conjectured to equal A367341(n)/9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael S. Branicky_ and _N. J. A. Sloane_, Nov 15 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016429 | OEIS-A367346

Numbers k such that there is more than one possible solution for A367338(k).

The sequence begins with the four exceptional terms 14, 33, 52, 71. It also includes all numbers with decimal expansions of the form d 9^i d (9-d), where juxtaposition is concatenation, ^ denotes repeated concatenation of digits, 1 <= d <= 8, and i >= 0, with associated next terms in the commas sequence being either d 9^(i+2) or (d+1) 0^(i+2). It is conjectured that there are no other terms. - _Michael S. Branicky_, Nov 16 2023 The conjecture is true; see link. - _Michael S. Branicky_, Nov 21 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 15 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016430 | OEIS-A367370

a(k) is the number of different widths patterns in the symmetric representation of sigma for numbers having k odd divisors.

Conjecture: a(9) = 28.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Dec 05 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016431 | OEIS-A367377

Square array T(n, k), n >= 1, k >= 1, read by antidiagonals, of the least numbers whose symmetric representation of sigma instantiate the unimodal width pattern 1, 2, ..., n, ..., 2, 1 repeated k times separated by instances of width 0.

Conjecture 1: None of the rows and columns are increasing. Conjecture 2: T(n, p) = T(n, 1) * A151800(2*T(n, 1))^(p-1) for n >= 1 and primes p. Conjecture 3: T(p, q), p and q primes, is a record for its upper left hand rectangle in the table. Only one prime number index generally is not sufficient as the inequality 4157280 = T(6, 2) < 5 * 10^6 < T(5, 2) shows.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Nov 15 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016432 | OEIS-A367396

Number of subsets of {1..n} whose cardinality is the sum of two distinct elements.

Conjectures from _Chai Wah Wu_, Nov 21 2023: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Nov 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016433 | OEIS-A367400

Number of subsets of {1..n} whose cardinality is not the sum of two distinct elements.

Conjectures from _Chai Wah Wu_, Nov 21 2023: (Start) The two conjectures of Chai Wah Wu above are true. Proof: write g(f,p,k) = [x^k] (1+2*x)^p*(1+x)^f and T_s(n) = Sum_{p=0..floor(n/2)} g(n-2*p, p, 2*p+s), so that a(n) = 1 + T_1(n) + T_2(n) by the formula above. Pascal's rule in f gives T_{s+1}(n+1) = T_{s+1}(n) + T_s(n), and in p it gives T_s(n+2) = binomial(n+2,s) + T_{s+2}(n) + 2*T_{s+1}(n). The first identity for s = 1, 2 and the second for s = 1 form a linear system from which the recurrence follows, and the initial terms 1, 2, 4, 7 then give the g.f. Both the formula and the recurrence are formalized in Lean 4 (see Links).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Nov 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016434 | OEIS-A367408

a(n) = n for n a power of 2; otherwise let 2^r be the greatest power of 2 which does not exceed n, then a(n) = the least novel m*a(k) where k = n - 2^r, and m is not a prior term.

Sequence is conjectured to be a permutation of the positive integers, A000027.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Nov 17 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016435 | OEIS-A367465

Lexicographically earliest infinite sequence of distinct positive numbers such that, when all terms are written as a product of their prime factors with specific primes as the first and last factor, the product of the two primes adjacent to the commas between the terms equals the magnitude of the difference between the terms.

See A367504 for the conjectured sequence when an additional requirement is added that the primes in the factorization of each term must be in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 18 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016436 | OEIS-A367467

Lexicographically earliest infinite sequence of positive integers such that a(n + a(n)) is distinct for all n.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Neal Gersh Tolunsky_, Nov 18 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016437 | OEIS-A367479

a(n) is the number of steps required for prime(n) to reach 2 when iterating the following hailstone map: If P == 5 (mod 6), then P -> next_prime(P + ceiling(sqrt(P))), otherwise P -> previous_prime(ceiling(sqrt(P))); or a(n) = -1 if prime(n) never reaches 2.

Conjecture: This hailstone operation on prime numbers will always reach 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Najeem Ziauddin_, Nov 19 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016438 | OEIS-A367484

Number of integers of the form (x^4 + y^4) mod 3^n; a(n) = A289559(3^n).

a(n) = 2*ceiling(3^(n+3)/80) - 1; a(n) = 2*3^(n-1) + a(n-4). - Conjectured by _Albert Mukovskiy_, Nov 19 2023; proved by _Max Alekseyev_, Mar 14 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Albert Mukovskiy_, Nov 19 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016439 | OEIS-A367504

a(1) = 2; for n > 1, a(n) = a(n-1) + 2*gpf(a(n-1)), where gpf(k) = A006530(k) = greatest prime dividing k.

Conjecture: with the requirement that the prime factorization of each term is written so that the primes are ordered from smallest to largest, the sequence is the lexicographically earliest infinite sequence of distinct positive numbers such that gpf(a(n-1)) * lpf(a(n)) = |a(n) - a(n-1)|, where gpf(k) = A006530(k) = greatest prime factor of k and lpf(k) = A020639(k) = least prime factor of k. In this way the sequence is the ordered prime factorization version of the 'Commas sequence', A121805. One can show that, for such a sequence to be infinite, no odd number can appear. Although for many terms a lower even number can be chosen for the following term, which can lead to even lower numbers for further terms, it is conjectured all such choices will ultimately halt the sequence as a number is eventually reached for which no unused next number exists which follows the required rule for the difference between the terms. Therefore all terms must be larger than the previous, and the earliest such infinite sequence is the given sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016440 | OEIS-A367558

Number of knight paths from opposite corners of an n X n matrix, using only turns that lower Manhattan distance.

Conjecture: a(n+1)/a(n) tends to d = 5.566315770083119..., where d is the real root of the equation d^3 - 4*d^2 - 8*d - 4 = 0. - _Vaclav Kotesovec_, Nov 28 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander Kuleshov_, Nov 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016441 | OEIS-A367621

The lexicographically earliest infinite sequence of positive numbers in which each term is a comma-child of the previous term in base 3.

We conjecture that choosing down-up-down-up-... is an infinite path, visiting the base-3 terms 1 2^{1+4*j} then 2 0^{2+4*j} for j in 0..oo, where ^ denotes repeated concatenation. This has been tested empirically up to j = 4300.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael S. Branicky_ and _N. J. A. Sloane_, Dec 20 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016442 | OEIS-A367646

a(0) = 0, a(1) = 1; for n > 1, a(n) = a(n-1) - GCT(a(n-2),a(n-1)) if nonnegative and not already in the sequence, else a(n) = a(n-1) + CT(a(n-2),a(n-1)), where CT(a,b) is the Comma transform (cf. A367360) of a and b, while GCT(a,b) is the largest possible generalized Command transform (cf. A367635) where at least one digit of both a and b can be chosen.

It is conjectured that all nonnegative numbers appear in the sequence. After the first 10 million terms the only fixed points are 0, 1, 2, 29, 65, 84, 222, 377, 491, 499, and it is likely no more exist. The first number to appear twice is a(35) = a(44) = 166.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 25 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016443 | OEIS-A367695

Numbers k such that k and k+1 are both exponentially odd numbers (A268335).

The conjecture above was proved by Srichan et al. (2026, p. 3, Corollary 1.3). They proved that the number of terms not exceeding x is d * x + O(x^(2/3+eps)), where d is the density and eps > 0. - _Amiram Eldar_, Jul 22 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Nov 27 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016444 | OEIS-A367708

Numbers k that are neither squarefree nor prime powers such that max(A119288(k), A053669(k)) <= A003557(k) < A007947(k).

Conjecture: contains most highly composite numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Feb 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016445 | OEIS-A367711

a(1) = 2; for n > 2, a(n) is the smallest positive number that has not yet appeared that has a common factor k > 1 with A001414(a(n-1)), the sum of the primes dividing a(n-1), with repetition.

In the first 100000 terms the only fixed point is 9, and it is likely no more exist. In the same range the smallest missing numbers are 4073, 5039, 5261. The sequence is conjectured to be a permutation of the integers >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016446 | OEIS-A367742

Lexicographically earliest infinite sequence of distinct positive numbers such that, for n>3, a(n) has a common factor with a(n-2) and n but not with a(n-1).

Unlike A098550 the primes do not appear in their natural order, and in general are delayed in their appearance relative to similarly sized numbers. In the first 100000 terms the fixed points are 1, 2, 3, 4, 9, 14, 16, 74, 76, 86, 88, 207, 320, 322, 901; it is unknown if more exist. The sequence is

conjectured to be a permutation of the positive numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016447 | OEIS-A367746

a(1) = 0, a(2) = 1. For n > 2, if a(n-1) is a novel term, a(n) = |a(n-1) - a(n-2) + 1|, else if a(n-1) is a repeat term, last seen at a(m), a(n) = n-m-1.

Conjecture: all numbers eventually appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Nov 28 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016448 | OEIS-A367750

a(1,2) = 1,2. let i,j represent a(n-2), a(n-1) respectively and let M(i) = {i,j}. For n > 2, if one of i,j is prime (p), and the other (m) is not, then if p!|m, a(n) is the least novel multiple of p. If p|m, a(n) = p*m. If both terms in M are nonprime a(n) is the least novel multiple of the smallest prime which does not divide i*j.

Conjectured to be a permutation of the positive integers with the primes occurring in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Nov 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016449 | OEIS-A367832

Array T(n, k) read by ascending antidiagonals is a dispersion based on A367467. Column 1 lists the numbers which cannot be represented by A367467(m) + m. For k >= 1, T(n, k+1) = A367467(T(n, k)) + T(n, k).

Conjectured: T(n, 3) = A328987(n-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Dec 02 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016450 | OEIS-A367857

a(n) is the smallest base b such that (b+1)^n in base b is a palindrome.

Conjecture: a(n) = binomial(n, floor(n/2))+1 for n>3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James Carruthers_, Dec 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016451 | OEIS-A367865

a(n) = Sum_{d|n} d * phi(d) * mu(d)^2.

Conjecture: a(n) = (rad(n)/n) * Sum_{k=1..n} rad(n*k)/rad(k), where rad is A007947. - _Velin Yanev_, Sep 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wesley Ivan Hurt_, Dec 03 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016452 | OEIS-A367949

Lexicographically earliest sequence of distinct positive integers such that the sum of the distinct prime factors (sopf) of a(n) + a(n + 1) is a perfect square.

Conjecture: this is a permutation of the natural numbers. - _Robert Israel_, Sep 15 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giorgos Kalogeropoulos_ and _Eric Angelini_, Dec 05 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016453 | OEIS-A367950

Lexicographically earliest sequence of distinct positive integers such that the sum of the distinct prime factors (sopf) of a(n) + a(n + 1) is a perfect cube.

Conjecture: this is a permutation of the positive integers. - _Robert Israel_, Sep 15 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giorgos Kalogeropoulos_ and _Eric Angelini_, Dec 05 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016454 | OEIS-A368108

a(1,2,3) = 1,2,3. For n > 3, a(n) is the smallest of the least novel multiples of all primes which divide an earlier term but do not divide a(n-1). If the prime divisors of all prior terms also divide a(n-1), a(n) is the least novel multiple of the smallest prime which does not divide a(n-1).

Conjectured to be a permutation of the positive integers with primes in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Dec 12 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016455 | OEIS-A368133

a(1,2,3) = 1,2,3; let j = a(n-1), M(n) = Product_{i = 1..n-2} { p a distinct prime: p | a(i), gcd(p, j) = 1 }. For n > 3, a(n) is the least novel multiple of M(n) if M(n) > 1; otherwise a(n) is the least novel multiple of A053669(j), the smallest prime which does not divide j.

Conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Dec 13 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016456 | OEIS-A368161

a(n) = least perfect power m that is the sum of n consecutive perfect powers, or -1 if m does not exist.

It is conjectured that m always exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. Stauduhar_ and _Tim Peters_, Dec 14 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016457 | OEIS-A368207

Bacher numbers: number of nonnegative representations of n = w*x+y*z with max(w,x) < min(y,z).

Conjectures: (1) a(n) >= sigma(n)/2, with equality if and only if n has no middle divisors, i.e., if and only if n is in A071561. (2) a(n)/sigma(n) converges to 1/2. - _Pontus von Brömssen_, Dec 18 2023 Conjecture: a(p^m) = sigma(p^m)/2 for odd m and all prime p > 2. a(p^m) = (sigma(p^m)-1)/2 + p^(m/2) for even m and all prime p.

a(2^m) = sigma(2^m)/2 + 1/2 for m odd. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Don Knuth_, Dec 16 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016458 | OEIS-A368235

Triangle read by rows: n-th row polynomial equals the numerator of the rational function (-1)^n*f(x) * (d/dx)^n (1/f(x)), where f(x) = sqrt(x + x^2).

Conjecture: the zeros of the polynomial R(n, -x) lie on the vertical line Re(x) = 1/2 in the complex plane.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Dec 18 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016459 | OEIS-A368246

Number of permutations of [n] whose cycle minima sum to n.

a(n) ~ c * (n-1)!, where c = 0.561459..., conjecture: c = exp(-gamma) = A080130, where gamma is the Euler-Mascheroni constant A001620. - _Vaclav Kotesovec_, Dec 29 2023

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alois P. Heinz_, Dec 18 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016460 | OEIS-A368281

Table T(n,k) n>=1, k>=1, read by downwards antidiagonals where the n-th row is the fixed point of the morphism 0->R(n), 1->R(n)0, where R(n) is the n-th repunit (A002275).

Conjecture: if a(k) is the fixed point of the morphism 0->R(n), 1->R(n)0, then the partial sum of a(k) is the Hofstadter-like sequence b(k): b(0)=0, b(k) = k - b(floor(b(k-1)/n)), i.e., the partial sum of the n-th row of A368281 is the n-th row of A368282. The cases n=1 and n=2 are known to be true (see A005206, A286389).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Dec 19 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016461 | OEIS-A368341

Fixed points of A368207.

Conjecture: 2^(2k+1) for k>=0 (A004171) are terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Dec 21 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016462 | OEIS-A368364

a(n) = number of s with n^k-n^2 <= s <= n^k-1, k >= 3, such that a comma sequence in base n with initial term s will not reach n^k.

Conjectured to have g.f. (Sum_{n>=1} x^((n^2+3*n)/2)/(1-x^n) - x^2)/(1-x). [Corrected by _N. J. A. Sloane_, May 14 2024] This is conjectured to equal A368363(n) - 1. Normally that would be enough to rule out this sequence. However, it is included because it is at present the only one of the nearly 100 OEIS entries based on comma sequences which has a connection with a sequence not connected with comma sequences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jan 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016463 | OEIS-A368460

a(n) = card(k: prime(n)^2 <= k < prime(n + 1)^2 and k term of A368458).

Conjecture: For all n >= 5, there is at least one j such that b(j) = 2 * (Bacher(j) - sigma(j)) + j + 1 is > 0 and prime(n)^2 < b(j) < prime(n + 1)^2. In other words, a(n) > 1 for n >= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Dec 26 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016464 | OEIS-A368539

Maximal sum of elements of A^2 where A is a square matrix of size n whose elements are a permutation of {1, 2, ..., n^2}.

In addition to the conditions (a)-(d) described in para 2.2 of Fried and Mansour (2023), conjecturally optimal matrices found using simulated annealing have the following additional property: If, using simultaneous row and column rearrangement, the matrix is brought into a form in which the terms of the main diagonal are sorted in ascending order, then every single row and every single column is monotonically increasing. See the linked file for examples from n=2 to n=14. - _Hugo Pfoertner_, Jan 25 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sela Fried_, Dec 29 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016465 | OEIS-A368546

Alternative version of the Markov tree A327345.

An open problem is to prove the uniqueness conjecture, which asserts that the largest element of a triple determines the other two.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _William P. Orrick_, Jan 04 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016466 | OEIS-A368558

Number of fractions i/n that are in the Cantor set.

Conjecture: The set of primes p such that a(p) > 2 has relative density 0 in the primes. This set begins 3, 13, 757, 1093, 3851, 4561, 34511, 363889, 368089, 797161, ... - _Charles R Greathouse IV_, Mar 12 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Yuen_, Dec 30 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016467 | OEIS-A368559

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest positive number not occurring earlier such that a(n) contains all the distinct digits of a(n-1) - a(n-2).

The sequence is conjectured to be a permutation of the positive integers. The fixed points begin 1, 2, 95, 122, 156, 318, 1644, 1964, 2189, 2740, 8264, 16904, ... although it is likely there are infinitely more.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Dec 30 2023 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016468 | OEIS-A368593

Expansion of g.f. A(x) satisfying A(x) = 1 + x*(A(x)^2 - A(-x)^2)/2 + x*sqrt((A(x)^4 + A(-x)^4)/2).

Conjecture: a(n) is odd when n = 2^k - 1 for k >= 0, and even elsewhere.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016469 | OEIS-A368630

Expansion of e.g.f. A(x) satisfying A(x/A(x)^2) = exp(x*A(x)).

Conjecture: a(n) == 1 (mod 3) for n >= 0. Conjecture: a(2*n) == 1 (mod 2) for n >= 0. Conjecture: a(2*n+1) == 0 (mod 2) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 02 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016470 | OEIS-A368631

Expansion of e.g.f. A(x) satisfying A(x/A(x)) = exp(x*A(x)^2).

Conjecture: a(n) == 1 (mod 3) for n >= 0. Conjecture: a(2*n) == 1 (mod 2) for n >= 0. Conjecture: a(2*n+1) == 0 (mod 2) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 02 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016471 | OEIS-A368748

a(n) is the number of numbers between prime(n) and prime(n+1) that are not prime powers.

Here "between" refers to numbers in the range [prime(n) + 1, prime(n+1) - 1], all of which are composite, and the sequence counts the numbers in each such range which are not prime powers. Whereas the corresponding number of prime powers seems bounded (see A080101), the number of numbers which are not prime powers is unbounded (see A014963). Conjecture: Every nonnegative integer appears in this sequence (at least once).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jan 04 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016472 | OEIS-A368749

a(n) is the smallest prime p such that there are n numbers between p and nextprime(p) which are not prime powers.

Conjecture: the sequence is infinite (number of nonprime powers between consecutive primes is arbitrarily large; see A014963).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_ and _Michael De Vlieger_, Jan 04 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016473 | OEIS-A368773

Antidiagonal sums of A059450.

Conjecture: D-finite with recurrence +(n+1)*a(n) +3*(-1)*a(n-1) +(-10*n+11)*a(n-2) +3*a(n-3) +9*(n-4)*a(n-4)=0. - _R. J. Mathar_, Mar 25 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joerg Arndt_, Jan 05 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016474 | OEIS-A368825

a(1) = 1; for n > 1, a(n) = A055231(a(n-1) + n), where A055231(k) is the powerfree part of k.

The sequence is conjectured to contain all the squarefree numbers; see A005117.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jan 07 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016475 | OEIS-A368846

Triangle read by rows: T(n, k) = (-1)^(n + k)*2*binomial(2*k - 1, n)* binomial(2*n + 1, 2*k) for k > 0, and k^n for k = 0.

Conjecture: |Sum_{j=0..k} T(k + j, k)| = A229580(k + 1) for k >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jan 07 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016476 | OEIS-A368916

a(n) = 1 if there is no prime p such that p^p divides the arithmetic derivative of A276086(n), and 0 otherwise, where A276086 is the primorial base exp-function.

Answer: If the conjecture that "For any given prime p, the distribution of the p-adic valuations of n' among those n that are not multiples of p, is the same as the distribution of the p-adic valuations among all the natural numbers", then the asymptotic mean of this sequence is equal to Product_{p prime} (1 - 1/p^(1+p)) = 0.864142072322... See also A368919, A360111 and A369653. - _Antti Karttunen_, Jan 29 and Feb 10 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016477 | OEIS-A368930

a(n) is the least k which is divisible by none of its digits in exactly n bases, or 0 if there is no such k.

Entries of 0 in the a-file are conjectural: if they are not 0, they are > 35000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Jan 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016478 | OEIS-A368959

Numbers m such that Sum_{k=0..m-1} exp(2*Pi*i*k^s/m) <> 0 for all s>=2.

Conjecture 1: This is a subset of A001694. Conjecture 2: This sequence contains all the terms from A338325, plus 243, 972, 1944, 2401, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jan 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016479 | OEIS-A369022

a(n) is the least start of a run of exactly n consecutive integers with the same maximal exponent in their prime factorization, or -1 if no such run exists.

a(8) <= 177001925537328703872748486819210922882 4 which is the conjectured value of A219452(8)+1. - _Giorgos Kalogeropoulos_, Jan 15 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Jan 12 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016480 | OEIS-A369040

Lexicographically earliest infinite sequence such that a(i) = a(j) => A003415(i) = A003415(j) and A342014(i) = A342014(j), for all i, j >= 1.

A369050(i) = A369050(j) => a(i) = a(j), [Conjectured]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 18 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016481 | OEIS-A369046

Lexicographically earliest infinite sequence such that a(i) = a(j) => A003415(i) = A003415(j) and A069359(i) = A069359(j), for all i, j >= 1.

A369050(i) = A369050(j) => a(i) = a(j), [Conjectured]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 18 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016482 | OEIS-A369054

Number of representations of n as a sum (p*q + p*r + q*r) with three odd primes p <= q <= r.

See the conjecture in A369055.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016483 | OEIS-A369083

Expansion of g.f. A(x) satisfying A(x) = 1 + x*(5*A(x)^2 - A(-x)^2)/4.

Conjecture: a(n) == binomial(4*n+3,n) (mod 2) for n >= 0 (cf. A263133).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 12 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016484 | OEIS-A369093

Numbers k >= 1 such that sigma(k) divides the sum of the triangular numbers T(k) and T(k+1), where sigma(k) = A000203(k) is the sum of the divisors of k.

Conjecture: The asymptotic density of primes in this sequence is 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Claude H. R. Dequatre_, Jan 13 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016485 | OEIS-A369096

Numbers k >= 2 such that omega(k) divides the sum of the triangular numbers T(k) and T(k+1), where omega(k) is the number of distinct primes dividing k (A001221).

Conjecture: the asymptotic density of primes in this sequence is 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Claude H. R. Dequatre_, Jan 13 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016486 | OEIS-A369110

a(n) is the number of distinct elements appearing in the sequence formed by recursively applying A063655 when starting from n.

A063655(n) gives the smallest semiperimeter of an integral rectangle with area n, which is the same thing as the minimum sum of two positive integers whose product is n. In this sequence, A063655 is applied recursively until a cycle is found. Then the number of distinct elements appearing in this process is given as a(n). Note that it's conjectured that a cycle will be found at some point. Conjecture: The cycle part of each sequence generated by the recursion is one of (4), (5, 6), or (6, 5). Confirmed through 1 millionth term. The conjecture is true. Proof: The conjecture holds for n <= 6. Suppose n >= 7 and the conjecture holds for lower values of n. If n is composite, then A063655(n) <= n/2+2. If n is prime, then A063655(n) = n+1 is even and A063655(A063655(n)) <= (n+1)/2+2. In both cases, n reaches a lower number and the conjecture holds for n. - _Jason Yuen_, Mar 30 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Adnan Baysal_, Jan 13 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016487 | OEIS-A369112

a(n) is the smallest number that has the n-th prime signature and is 1 greater than a prime, or -1 if no such number exists.

Conjecture: the converse is also true; i.e., if A025487(n) is of neither of the two forms above, then there exists at least one number that has the n-th prime signature and is 1 greater than a prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Jan 13 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016488 | OEIS-A369184

a(n) is the first positive number that has exactly n anagrams which have 3 prime divisors, counted by multiplicity, or 0 if there is no such number.

Conjecture: all a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Jan 15 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016489 | OEIS-A369255

Parity of A140773, where A140773 is the inverse Möbius transform of A038548.

Following convolution formulas have been conjectured for this sequence by Sequence Machine, with both computing the first 2^20 terms correctly:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 24 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016490 | OEIS-A369271

The smallest positive number that has not yet appeared that is a multiple of the smallest prime that does not divide n.

The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jan 18 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016491 | OEIS-A369272

a(1) = 1; for n > 1, a(n) is the smallest positive number that has not yet appeared that is a multiple of the smallest prime that does not divide n*a(n-1).

The sequence is conjectured to be a permutation of the positive integers, although it takes many terms for the primes to appear, e.g., a(63784) = 23.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jan 18 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016492 | OEIS-A369275

Nonnegative numbers k satisfying sin(k) < sin(k+1) < sin(k+2) < sin(k+3) < sin(k+4).

Conjecture: there is no followup sequence with sin(k) < sin(k+1) < ... < sin(k+5), i.e., there are no two consecutive integers in this sequence. Mathar's conjecture is true: these are just the numbers which are between Pi*3/2 - 1/2 and Pi*5/2 - 7/2 mod 2*Pi, and reducing the upper bound by 1 leaves it empty. - _Charles R Greathouse IV_, Dec 10 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _R. J. Mathar_, Jan 18 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016493 | OEIS-A369303

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest unused number whose string value contains the comma transform (cf. A367360) of the previous two terms.

The sequence is conjectured to be a permutation of the positive integers. The fixed points begin 1, 2, 10, 11, 2863, 3164, 3545, 3947, 6835, 6947, 7052, ... although it is likely there are infinitely more.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jan 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016494 | OEIS-A369366

Least number of inequivalent cells in a polyiamond with n cells.

Conjecture: a(6k+5) = 3(k+1) for all k >= 0. Observed for k = 0, 1, 2 (n = 5, 11, 17). The next test case is a(23), predicted to be 12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pontus von Brömssen_, Jan 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016495 | OEIS-A369375

Numbers m such that the Mersenne number 2^m - 1 is a de Polignac number (A006285).

Cf. A138290 (see _Chai Wah Wu_'s conjecture in the third comment). By Crocker's (1971) theorem: if m > 2 and a <> b, then 2^(2^m)-1 - 2^a - 2^b is not prime. Conjecture: if n > 5, then |2^(2^n-1)-1 - 2^m| is not prime for every m > 0. If so, then by the dual Riesel conjecture, 2^(2^n-1)-1 is a (dual) Riesel number, i.e., if n > 5, then (2^(2^n-1)-1)2^m-1 is composite for every integer m > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Jan 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016496 | OEIS-A369378

a(n) is the smallest k such that 2^(2^n+1)-1 + 2^k is prime, or -1 if no such k exists.

Conjecture: if n > 0, then 2^(2^n+1)-1 + 2^k is not prime for every k < 2^n. This conjecture seems provable, because it is not too strong for large n > 6. If, for n > 6, a(n) does not exist, then 2^(2^n+1)-1 + 2^k is composite for every natural k. Thus, by the dual Sierpinski conjecture, for n > 6, 2^(2^n+1)-1 is a Sierpinski number, i.e., if n > 6, then (2^(2^n+1)-1)2^k+1 is composite for every natural k. For example, the Mersenne number 2^(2^8+1)-1 may be a dual Sierpinski number. Similarly, if for n > 5, |2^(2^n-1)-1 - 2^m| is not prime for every m > 0, then by the dual Riesel conjecture, 2^(2^n-1)-1 is a Riesel number, i.e., if n > 5, then (2^(2^n-1)-1)2^m-1 is composite for every integer m > 0. For example, the double Mersenne prime 2^(2^7-1)-1 may be a dual Riesel number. By Crocker's theorem; if n > 2, then positive 2^(2^n-1)-1 - 2^m are composite. Let b(n) be the smallest k such that 2^k - (2^(2^n-1)-1) is prime, for n >= 0: {1, 2, 39, 47, 447, 191, ?}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Jan 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016497 | OEIS-A369435

Square array A(n, k) = n! * [t^n] (exp(t)/(1+k-k*exp(t))) for n >= 0 and k >= 0, read by antidiagonals upwards.

The following formulae are conjectures: Conjecture: A(n, n) = (n + 1) * A321189(n) for n >= 0. [This is true. - _Peter Luschny_, Apr 26 2024]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Jan 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016498 | OEIS-A369463

Numbers of the form 12*m-1 for which there is no representation as a sum (p*q + p*r + q*r) with three odd primes p <= q <= r.

Question: Is this a finite sequence, with the last term a(285) = 50688947 = (12*4224079)-1? See conjecture in A369055.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016499 | OEIS-A369580

a(n) := f(n, n), where f(0,0) = 1/3, f(0,k) = 0 and f(k,0) = 3^(k-1) if k > 0, and f(n, m) = f(n, m-1) + f(n-1, m) + 3*f(n-1, m-1) otherwise.

9*a(n) - a(n+1) = 2*A162326(n) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tadayoshi Kamegai_, Jan 26 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016500 | OEIS-A369595

Array read by downward antidiagonals: A(n,k) = A(n-1,k+2) + Sum_{j=0..k} binomial(k,j)*A(n-1,j) with A(0,k) = 1, n >= 0, k >= 0.

Conjecture: A(n,0) = A135920(n+1). - _Mikhail Kurkov_, Oct 27 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Jan 27 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016501 | OEIS-A369615

Powers of primes (A000961) whose neighbors have a prime number as their greatest odd divisor.

Conjecture: a(12) = 1594323 is the final term of the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jan 27 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016502 | OEIS-A369653

a(n) = 1 if A327860(n) is a multiple of 3, otherwise 0, where A327860 is the arithmetic derivative of the primorial base exp-function.

Conjecture: asymptotic mean is 4/9: One third of the primorial base representations (A049345 from which the intermediate value x = A276086(n) is built from) have "2" as their second rightmost digit (out of 0, 1, 2). That 2 is converted by A276086 to 3^2, so as x is a multiple of nine, it is guaranteed that x' [= A003415(x)] gives a multiple of three. If the second rightmost digit is "1", then x = A276086(n) is a multiple of 3 (but not of 9), and its arithmetic derivative, x' = 3*(x/3)' + (x/3), certainly is not a multiple of 3. This leaves the case where the second rightmost digit is "0", and conjecturing now in this case that the modulo 3 remainders are eventually evenly distributed for x' (see A359430 and A373256), at least among the number of the form A048103, it gives in this case 1/3 * 1/3 = 1/9 probability that x' is a multiple of 3, so the total probability would be 1/3 + 0 + 1/9 = 4/9. Note that Sum_{i=1..10^n} a(i), for n = 1..8 gives: 3, 43, 443, 4443, 44443, 444443, 4444444, 44444444, so empirically the value 4/9 seems very sharp. - revised Jun 18 2024.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016503 | OEIS-A369658

a(n) = 1 if n is not multiple of 3, but its arithmetic derivative is, otherwise 0.

Conjecture: the asymptotic mean of this sequence is (2/3)*(1/3) = 2/9. Compare to the comment at A369653, but consider also the four lowermost rows of the table given at A369252 (and further generalizations to various number of primes), and also A007352, A096629, and how they affect such probabilities.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016504 | OEIS-A369673

a(n) = Product_{k=0..n} (2^k + 2^(n-k)).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 06 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016505 | OEIS-A369686

LCM-transform of A359804 (see Comment and links).

The property S (as defined in A368900) refers to what is observed in the positive integers (A000027), and also in the Doudna sequence (A005940), whereby each prime power appears prior to any of its multiples. The present sequence does not have this property since, for example, 26 = a(31) precedes 13 = a(42). Thus A369804 represents a significant disturbance of A000027 in that whereas it is conjectured to be a permutation of the positive integers, it does not preserve one of the basic properties of that sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jan 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016506 | OEIS-A369692

Connected domination number of the n X n grid graph.

a(3*n) <= n*(3*n+1); a(3*n-1) <= 3*n^2 - 1; a(3*n-2) <= (n-1)*(3*n+1). Conjecturally these inequalities hold with equality for n > 1. - _Andrew Howroyd_, Mar 06 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alexander D. Healy_, Feb 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016507 | OEIS-A369713

a(n) is the sum over all multiplicative partitions k of n of the absolute value of the Möbius function evaluated at k,n in the poset of multiplicative partitions of n under refinement.

Conjecture: There exists c such that a(n) <= n^c for all natural numbers n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tian Vlasic_, Jan 29 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016508 | OEIS-A369786

Number of different coefficient values in expansion of Product_{k=1..n} (1+x^(k^2)).

Conjecture: a(n) ~ n^3/6. - _Vaclav Kotesovec_, Feb 02 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Feb 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016509 | OEIS-A369793

a(n) is the number of occurrences of n in A063655.

Construct a directed graph whose vertex set is the set of all positive integers, and a directed edge from k to n belongs to this graph iff A063655(k) = n. a(n) is the in-degree of the vertex n in this graph. As conjectured in A369110, it is also conjectured here that the only cycles in this graph are from 4 to itself and between 5 and 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Adnan Baysal_, Feb 07 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016510 | OEIS-A369797

Denominator of the continued fraction 1/(2-3/(3-4/(4-5/(...(n-1)-n/(n+2))))).

Conjecture: The sequence contains only 1's and the primes. Conjecture: The sequence of record values is A002476. - _Bill McEachen_, Mar 24 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mohammed Bouras_, Feb 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016511 | OEIS-A369802

Inversion count of the Eytzinger array layout of n elements.

Conjecture: a(n) = (A261692(n)-n)/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Darío Clavijo_, Feb 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016512 | OEIS-A369818

The sixth term of the greedy B_n set of natural numbers.

Conjectured that a(6n+i) is a quartic polynomial sequence with lead term (1/3)n^4 for each i in {1,2,3,5,6,10} in arxiv:2312.10910.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kevin O'Bryant_, Feb 03 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016513 | OEIS-A369825

a(1,2) = 1,2; thereafter let i = a(n-2) and r(n) = a(1)*a(2)*...*a(n-1): a(n) is the smallest novel multiple m of w = rad(r(n))/rad(i), where rad is A007947.

(i). Directly: rad(i) = rad(r(n)) implies a(n) is the least unused term, conjectured to be the smallest prime p not already in the sequence. In this case no prior term is divisible by p. This happens for 3, 5, 7. It is not known if this ever happens again (thought to be unlikely).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_ and _Michael De Vlieger_, Feb 02 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016514 | OEIS-A369831

a(n) is the number of distinct values of the permanent of an n X n symmetric Toeplitz matrix using the integers 1 to n.

Conjectured e.g.f.: 1/(1 - x) - x^2/2 - x^4/24.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Feb 03 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016515 | OEIS-A369833

a(n) is the number of distinct values of the permanent of an n X n symmetric Toeplitz matrix using the first n prime numbers.

Conjectured e.g.f.: 1/(1 - x) - x^2/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Feb 03 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016516 | OEIS-A369899

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest unused positive number that is a substring of the sum of all previous terms.

The fixed points begin 1, 2, 3, 94, 1420, 1423, 1425, 1426, 1427, 8592, although it is likely there are infinitely more. The sequence is conjectured to be a permutation of the positive numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Feb 05 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016517 | OEIS-A369967

a(n) = 1 if the arithmetic derivative of n is a multiple of 5, otherwise 0.

Question: Is the asymptotic mean of this sequence 1/5? See also conjecture in A369968.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016518 | OEIS-A369968

a(n) = 1 if n is not multiple of 5, but its arithmetic derivative is, otherwise 0.

Conjecture: the asymptotic mean of this sequence is (4/5)*(1/5) = 4/25 = 0.16. Compare to A369967 and the conjecture at A369658.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016519 | OEIS-A369988

Decimal expansion of Mallows's constant or stribolic constant kappa (of order 1).

In 1997, Colin Mallows observed and conjectured that the rows in Levine's triangle A012257 take on stribolic shape and that A011784(n+1)/(A011784(n)*A011784(n-1)) converges as n->oo. Presuming his conjecture, the limit would equal kappa, while Mallows estimated it to be "approximately ... 0.277", see A011784. Later, in 2006, Martin Fuller suggested 0.27887706... for the limit, based on a numerical iteration, see A012257. Set kappa_n := A369990(n) / A369991(n) and theta_n := (kappa_n-kappa_{n+1}) / (kappa_{n-1}-kappa_n). Under the hypothesis that theta_{2m} < theta_{2m+2} < theta_{2*m+3} < theta_{2*m+1} for m=1,2,... (verified for all values known so far), we would obtain 0.27887706136895087 < kappa_{21}' < kappa < kappa_{22}' < 0.27887706136898083, which is sharper than formula (3) below. Here, the transformed sequence (kappa_n') = G(kappa_n) is defined via kappa_n' := (kappa_{n-1}*kappa_{n+1} - kappa_n^2) / (kappa_{n-1} - 2*kappa_n + kappa_{n+1}). (See first arXiv article for a proof of this conjecture-dependent statement.) Feeling even more adventurous, we could apply the transformation G four times and would obtain 0.278877061368975064775 < kappa_{19}'''' < kappa < kappa_{18}'''' < 0.278877061368975064815. It is an open question whether kappa is rational or irrational, algebraic or transcendental.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roland Miyamoto_, Feb 07 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016520 | OEIS-A370046

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest unused positive number whose binary value is a substring of the binary value of the sum of all previous terms.

The fixed points begin 1, 2, 3, 16, 39, 42, 50, 79, 120, 361, although it is likely there are infinitely more. The sequence is conjectured to be a permutation of the positive numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Feb 08 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016521 | OEIS-A370047

a(1) = 1; a(2) = 2; for n > 2, a(n) is smallest unused number such that a(n) shares a factor with a(n-1) and sopfr(a(n)) shares a factor with sopfr(a(n-1)), where sopfr(k) is the sum of the primes dividing k, with repetition.

In the first 100000 terms the primes appear in their natural order, although they are delayed relative to similarly sized numbers. In the same range the fixed points are 1, 2, 9, 27, 165, 88812. The sequence is conjectured to be a permutation of the positive numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Feb 08 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016522 | OEIS-A370092

a(0) = 1, a(n) = (-1)^n + (1/2) * Sum_{j=1..n} (1-(-1)^j-(-2)^j) * binomial(n,j) * a(n-j) for n > 0.

Conjecture: Let k > 2 be a positive integer. The sequence obtained by reducing a(n) modulo k is eventually periodic with the period dividing phi(k) = A000010(k). For example, modulo 10 we obtain the sequence [1, 1, 3, 6, 5, 6, 3, 6, 5, 6, 3, 6, 5, 6, 3, 6, 5, 6, ...] with an apparent period of 4 beginning at a(2). See A000670 for a more general conjecture. - _Peter Bala_, Feb 16 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Prabha Sivaramannair_, Feb 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016523 | OEIS-A370097

a(n) = Sum_{k=0..n} binomial(3*n,k) * binomial(3*n-k-1,n-k).

Conjecture: the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) holds for all primes p >= 3 and positive integers n and r. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Feb 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016524 | OEIS-A370100

a(n) = Sum_{k=0..n} binomial(4*n,k) * binomial(2*n-k-1,n-k).

Conjecture: the supercongruences a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) hold for all primes p >= 5 and positive integers n and r. See A352373 for a more general conjecture. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Feb 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016525 | OEIS-A370101

a(n) = Sum_{k=0..n} binomial(4*n,k) * binomial(4*n-k-1,n-k).

We conjecture that the stronger supercongruences a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) hold for all primes p >= 5 and all positive integers n and r. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Feb 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016526 | OEIS-A370115

Numbers k for which k is a multiple of A276086(A003415(k)), where A003415 is the arithmetic derivative, and A276086 is the primorial base exp-function.

Conjecture: All nonzero terms of this sequence are squarefree, in A005117. This would follow from Aug 31 2026 conjecture given in A327859.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016527 | OEIS-A370164

The number of residues mod n that occur among the Markov numbers.

It is conjectured that, modulo n, all residues not forbidden by the constraints in the three comments above are actually realized by some Markov number. Specifically, mod n the forbidden residues include the following: (1) if p is a prime congruent to 3 (mod 4) that divides n, then any residue congruent to 0 (mod p) or congruent to either of the forbidden residues listed in the second comment (mod p), (2) if 4 divides n, then any odd residue congruent to 3 (mod 4), (3) if 2^r is the highest power of 2 dividing n and 2 <= r <= 5, then any even residue not congruent to 2 (mod 2^r), (4) if 2^r is the highest power of 2 dividing n and r > 5, then any even residue not congruent to 2 (mod 32). It is conjectured that all other residues occur. This has been verified for all n <= 38000. If the conjecture in the previous comment is correct, then it follows from the Chinese remainder theorem that a(n) may be found by writing n = 2^s * 3^t * u * p_1^r_1 * p_2^r_2 * ... * p_k^r_k, where p_1, ..., p_k are distinct primes greater than 3 and congruent to 3 (mod 4) and r_1, ..., r_k are positive and where the prime divisors of u are all congruent to 1 (mod 4). Then a(n) = u * C_s * D_t * Product_{j=1..k} (p_j - 3) * p_j^(r_j - 1), where C_s = 2^s if s < 2, 1 + 2^(s-2) if 2 <= s <= 5, and 2^(s - 5) + 2^(s - 2) if s > 5, and where D_t = 1 if t = 0 and 2 * 3^(t-1) if t > 0. If the conjecture holds then a(n) is a multiplicative function. Conjectured: write n = 2^s * 3^t * u * p_1^r_1 * p_2^r_2 * ... * p_k^r_k, where p_1, ..., p_k are distinct primes greater than 3 and congruent to 3 (mod 4) and r_1, ..., r_k are positive and where the prime divisors of u are all congruent to 1 (mod 4). Then a(n) = u * C_s * D_t * Product_{j=1..k} (p_j - 3) * p_j^(r_j - 1), where C_s = 2^s if s < 2, 1 + 2^(s-2) if 2 <= s <= 5, and 2^(s - 5) + 2^(s - 2) if s > 5, and where D_t = 1 if t = 0 and 2 * 3^(t-1) if t > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_ and _William P. Orrick_, Feb 26 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016528 | OEIS-A370206

Numbers j whose symmetric representation of sigma(j) consists of two copies of unimodal width pattern 121 separated by 0.

Conjecture: The subsequence of numbers m whose two parts of SRS(m) meet at the diagonal is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Feb 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016529 | OEIS-A370209

a(n) is the smallest number of the form 2^k * p * (2^(k+1) * p + 1) where 2 < p < 2^(k+1) is the n-th prime and 2^(k+1) * p + 1 is prime, or -1 if no such number exists.

Conjecture: The sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Feb 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016530 | OEIS-A370236

Triangle read by rows: T(n, k) is the number of partitions of genus 1 and k parts of the n-set (n >= 4, 2 <= k <= n-2).

The formula given below was conjectured by Martha Yip and proved by Robert Cori and Gábor Hetyei.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Coquereaux_, Feb 12 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016531 | OEIS-A370260

a(n) = sqrt(A370259(2*n+1)).

The sequence is conjectured to be integral [added 03 Mar 2024: now confirmed - see the Formula section].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016532 | OEIS-A370261

a(n) = sqrt(A370259(2*n)/(n+1)) for n >= 1.

The sequence is conjectured to be integral.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Mar 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016533 | OEIS-A370305

Numbers k such that the distance from exp(k) to the closest average of two consecutive primes is less than 1.

For k>1, the formula (prevprime(e^k)+nextprime(e^k))/2 either gives floor(e^k), for k = 61, 74, 2170, ..., or gives ceiling(e^k), for k = 3, 16, 91, 113, 1441, 1566, 2499, ...

This partitions {a(n)}\{1} into two subsequences each of which can be conjectured to have relative density 1/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeppe Stig Nielsen_, Feb 14 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016534 | OEIS-A370380

Array read by downward antidiagonals: A(n,k) = (k+2)*A(n-1,k+1) + Sum_{j=0..k} A(n-1,j) with A(0,k) = 1, n >= 0, k >= 0.

Conjecture: A(n,0) = A003319(n+2). - _Mikhail Kurkov_, Oct 27 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Feb 17 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016535 | OEIS-A370381

Array read by downward antidiagonals: A(n,k) = Sum_{j=0..k+1} binomial(k+2, j+1)*A(n-1,j) with A(0,k) = 1, n >= 0, k >= 0.

Conjecture: A(n,0) = A003319(n+2). - _Mikhail Kurkov_, Oct 27 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Feb 17 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016536 | OEIS-A370382

Array read by downward antidiagonals: A(n,k) = A(n-1,k+1) + (k+1)*Sum_{j=0..k} binomial(k, j)*A(n-1,j) with A(0,k) = 1, n >= 0, k >= 0.

Conjecture: A(n,0) = A249833(n+1). - _Mikhail Kurkov_, Apr 25 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Feb 17 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016537 | OEIS-A370388

First appearance of prime_i in A369797 or 0 if no such appearance occurs, with p_0 being 1. (A008578 but with a different offset.)

Conjectures from _Ridouane Oudra_, Apr 26 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Feb 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016538 | OEIS-A370394

Number of steps to reach 0 by applying the map f(x) = 5x + 1 if x is odd and floor(x/3) if x is even, or -1 if 0 is never reached.

Conjecture: for all n, a(n) != -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Feb 17 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016539 | OEIS-A370410

Number of length-n binary strings that are the concatenation of two nonempty palindromes.

a(n) = A007055(n) - A056458(n) (conjectured). - _Michael S. Branicky_, Feb 18 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Feb 18 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016540 | OEIS-A370488

a(n) is the smallest integer k such that the average deviation of previous terms and k is an integer, where a(n) > a(n - 1) and a(1) = 1.

Conjecture: a(n)/n^3 tends to c, where 0.08 < c < 0.1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nicolas Bělohoubek_, Feb 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016541 | OEIS-A370496

a(1) = 1; for n > 1, a(n) is smallest unused number such that a(n) is coprime to a(n-1) and sopfr(a(n)) is coprime to sopfr(a(n-1)), where sopfr(k) is the sum of the primes dividing k, with repetition.

In the first 100000 terms the primes appear in their natural order. In the same range the fixed points begin 1, 2, 3, 4, 5, 20, 134, 136, 403, 598, 608, 649, 667. The sequence is conjectured to be a permutation of the positive numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Feb 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016542 | OEIS-A370497

a(1) = 1; for n > 1, a(n) is smallest unused number such that a(n) is coprime to a(n-1), sopfr(a(n)) is coprime to sopfr(a(n-1)), and Omega(a(n)) does not equal Omega(a(n-1)), where sopfr(k) is the sum of the primes dividing k, with repetition.

In the first 100000 terms the primes 11, 7 and 5 appear in reverse order, but all others appear in their natural order. In the same range the fixed points begin 1, 2, 4, 13, 294, 295, 296, 299, 304, 309, 640, 649. The sequence is conjectured to be a permutation of the positive numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Feb 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016543 | OEIS-A370500

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest unused positive number such that a(n) does not share a factor with a(n-1) but sopfr(a(n)) does share a factor with sopfr(a(n-1)), where sopfr(k) is the sum of the primes dividing k, with repetition.

The fixed points begin 1, 2, 4, 56, 72, 138, 200, 438, 500, 540, 590, 3998. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Feb 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016544 | OEIS-A370501

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest unused positive number such that a(n) shares a factor with a(n-1) but sopfr(a(n)) does not share a factor with sopfr(a(n-1)), where sopfr(k) is the sum of the primes dividing k, with repetition.

The fixed points begin 1, 2, 64, 114, 116, 132, and it is plausible no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Feb 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016545 | OEIS-A370502

a(1) = 1; for n > 1, a(n) is the smallest unused positive number such that sopfr(|a(n) - a(n-1)|) = sopfr(a(n) + a(n-1)), where sopfr(k) is the sum of the primes dividing k, with repetition.

The fixed points begin 1, 8, 78, 314, 1373, 8822, 10247, 21935; it is likely there are infinitely more. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Feb 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016546 | OEIS-A370503

a(1) = 1; for n > 1, a(n) is the smallest unused positive number such that sopfr(|a(n) - a(n-1)|) = sopfr(a(n) + a(n-1)) and Omega(|a(n) - a(n-1)|) = Omega(a(n) + a(n-1)), where sopfr(k) is the sum of the primes dividing k, with repetition.

The fixed points begin 1, 306, 13442, 24566, 30607, 53557, 65136; it is likely there are infinitely more. The sequence is conjectured to be a permutation of the positive integers, although it takes many terms for some values to appear, e.g., a(177950) = 9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Feb 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016547 | OEIS-A370526

Square array read by descending antidiagonals: Define function b(i,n,k) where b(0,n,k) = n, b(1,n,k) = k, b(i,n,k) = A038502(b(i-1,n,k) + b(i-2,n,k)). T(n,k) is the number of steps until reaching the cyclic part of {b(i,n,k)}, or -1 if no cycle exists.

Conjecture: T(n,k) >= 0. This conjecture can be supported by a heuristic argument: Using dynamic programming, we can compute that for p's in A000057, the probability that b(i-1,n,k) + b(i-2,n,k) is not divisible by p is (p-2)/p, and the probability that valuation(b(i-1,n,k) + b(i-2,n,k), p) = x (x >= 1) is 2*(p-1)/p^(x+1). Therefore, the mathematical expectation of a(i) is (a(i-1,n,k) + a(i-2,n,k))*(p-1)/(p+1), which is exactly the average of the earlier two terms when p = 3, and larger when p >= 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yifan Xie_, Feb 21 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016548 | OEIS-A370574

a(n) is the least k > 0 such that n^2 XOR k^2 is a positive square number (where XOR denotes the bitwise XOR operator), or -1 if no such k exists.

Conjecture: a(n) > 0 for any n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Feb 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016549 | OEIS-A370578

Numbers k such that k + 1 divides 3^k + 1.

Apart from 0, the only possible residues mod 72 are 1, 3, 13, 25, 27, 37, 49, 51, and 61. It is conjectured that all appear eventually. (See John Omielan's answer to the author's question on Mathematics Stack Exchange.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Akiva Weinberger_, Feb 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016550 | OEIS-A370630

Lexicographically earliest sequence of distinct positive integers such that the Zeckendorf expansions of two consecutive terms have exactly one common term.

Conjecture: this sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, May 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016551 | OEIS-A370662

Numbers m such that 3 does not divide the m-th Catalan number A000108(m); m such that A067397(m) = 0.

Conjecture: the only terms of the form 2^r-1 are 0, 1, 3, 31 and 255. Since 2^r-1 !== 2 (mod 3), this is equivalent to saying that the only numbers of the form 2^r-1 that have no digits 2 in ternary are 0, 1, 3, 31, 255. The conjecture would imply that the n-th Catalan number is divisible by 2 or 3 other than n taking these values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Feb 24 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016552 | OEIS-A370691

Square array read by upward antidiagonals: T(n, k) = denominator(2*k!*(-2)^k*Sum_{m=1..n}(1/(2*m-1)^(k+1))).

Conjectured: T(n, 2) = A128507(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Apr 21 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016553 | OEIS-A370724

Determinant of n X n Fibonacci word matrix.

Conjectures: a(F(n)) = +-F(n-2) where F(n) is the n-th Fibonacci number, and a(L(n)) = +-L(n-2) where L(n) is the n-th Lucas number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Feb 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016554 | OEIS-A370726

Denominator of the continued fraction 1/(2-3/(3-4/(4-5/(...(n-1)-n/(n+3))))).

Conjecture: The sequence contains only 1's and the primes. Conjecture: Record values correspond to A002144 (n>3). - _Bill McEachen_, May 21 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mohammed Bouras_, Feb 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016555 | OEIS-A370740

a(1) = 1. Thereafter a(n) is the least novel k such that A007947(k*a(n-1)) is the smallest number in A002110 which is not already a term.

Conjectured to be a permutation of the positive integers (A000027), with primorials, primes and prime powers in natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Feb 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016556 | OEIS-A370789

Ore numbers whose nontrivial divisors are not Ore numbers.

Assuming odd perfect numbers do not exist, the above conjecture is true since: a) an even perfect number is not divisible by a smaller even perfect number, b) an even perfect number is not divisible by a smaller nonperfect Ore number k with omega(k)>=3, and c) all Ore numbers m with omega(m)=2 are even perfect numbers (see Carl Pomerance link at A001599). - _Ivan N. Ianakiev_, Apr 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ivan N. Ianakiev_, Mar 02 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016557 | OEIS-A370905

Partial sums of the alternating sum of divisors function (A206369).

Conjecture: also number of pairs (x, y) with 1 <= x <= y <= n such that gcd(x, y) is a perfect square. - _Michel Marcus_, Jun 12 2026 Conjecture is true as a consequence of Geoffrey Critzer's comment in A206369. - _Chai Wah Wu_, Jun 22 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Mar 05 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016558 | OEIS-A370911

a(n) is the smallest positive integer k with k != 10^m (m: nonnegative integer) for which 1/n can be obtained by incorrectly reducing k/(n*k) - by deleting the same digit in the numerator and denominator as often as possible, leaving one digit "1" in the numerator.

Conjectures from _Felix Huber_, Apr 28 2024: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Mar 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016559 | OEIS-A370955

Coefficients of the partition function per spin, x(k) (divided by 2), in the low temperature region, expressed as a power series in the parameter k^2, for the spin-1/2 Ising model on square lattice.

Conjecture: c = exp(2*G/Pi)/(8*Pi) = 0.07128640667426940835812 3..., where G is Catalan's constant A006752. (End) [proved by _Dominik Beck_, Jul 16 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016560 | OEIS-A371113

The sequence is a succession of triples of nonnegative integers. In each triple, the digits form a palindromic pattern. The sequence starts with a(0)=0 and is always extended with the smallest available integer not yet present in the sequence.

Conjecture 1: The only fixed points of this permutation are n = 0 and 1. Conjecture 2: For all k >= 0, a(3k+2) <= R(concatenate(a(3k), a(3k+1), 1)), where R = A004086 = reverse. Conjecture 3: For all k >= 0, a(3k+1) - a(3k) = 1 or 2. (Verified up to k = 3333. a(3k+1) - a(3k) = 2 for k = 15, 36, 62, 81, 110, 129, 138, 163, 182, ...) a(3k) = 2k + o(log k)

; a(3k) + 1 <= a(3k+1) <= a(3k) + 2 (conjectured). - _M. F. Hasler_, Apr 20 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Giorgos Kalogeropoulos_, Apr 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016561 | OEIS-A371176

Numbers k such that A000120(k) <= A001511(k).

Conjecture: lim sup_{n -> oo} log(a(n))/log(n) = log(2) / log((1 + sqrt(5))/2) = 1.440420090412556479... = A104287. - _Vaclav Kotesovec_, Aug 12 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Mar 14 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016562 | OEIS-A371195

List of values q^5-q^4 as q runs through A000961 (that is, 1, primes, and prime powers).

Parekh et al. conjecture that q^5-q^4 is the number of nonsingular generalized Weierstrass curves over the field with q elements, for q>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 14 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016563 | OEIS-A371279

Irregular triangular array of numerators of the set T of fractions generated by these rules: g(1) = (1), and if x and y are in T, then x/(y+1) is in T; see Comments.

Conjecture: every rational number in the interval (0,1] occurs exactly once in T.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 18 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016564 | OEIS-A371280

Irregular triangular array of denominators of the set T of fractions generated by these rules: g(1) = (1), and if x and y are in T, then x/(y+1) is in T; see Comments.

Conjecture: every rational number in the interval (0,1] occurs exactly once in T.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 18 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016565 | OEIS-A371303

Numbers k > 4 such that both k - 2^(2^m) and k + 2^(2^m) are prime for every natural m > 0 with 2^(2^m) < k.

Conjecture: numbers k > 2 such that both k - 2^(2^m) and k + 2^(2^m) are prime for every integer m >= 0 with 2^(2^m) < k are only 9, 15, and 195 (_Amiram Eldar_ checked that there are no more terms k < 10^8). 275105655 is the smallest counterexample to the finite m >= 0 conjecture. - _Sai Kambampati_, Sep 10 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Mar 18 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016566 | OEIS-A371440

Count of iteration steps for the iterates in A267371 that end in 1.

Conjecture: If iterate m terminates with prefix s ending in 1 then the subsequent iterate of A267371 is terminated by prefix s0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Mar 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016567 | OEIS-A371473

a(1) = 1; for n>1, if a(n-1) is squarefree, a(n) = a(n-1) + n, otherwise a(n) = squarefree kernel of a(n-1).

Some nonsquarefree numbers will not appear in this sequence. However, I conjecture that all squarefree numbers will appear. First occurrence of 2 is at a(766) = 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joseph C. Y. Wong_, Mar 24 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016568 | OEIS-A371511

a(n) is the smallest prime such that its representation in base n contains each of the digits 0,1,...,n-2 at least once and does not contain the digit n-1.

Conjecture: for n>3, a(n) has digit sum 2+(n-2)(n-1)/2 if n is of the form 4k+3 and has digit sum 1+(n-2)(n-1)/2 otherwise.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Apr 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016569 | OEIS-A371561

Numbers with multiplicative digital root of 5 that are free of 1s and have their digits in ascending order.

Conjectured to be complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sergio Pimentel_, Mar 27 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016570 | OEIS-A371567

Array read by downward antidiagonals: A(n,k) = A(n-1,k+1) + (k+1)*Sum_{j=0..k} A(n-1,j) with A(0,k) = k+1, n >= 0, k >= 0.

Conjecture: A(n,0) = A258173(n+1). - _Mikhail Kurkov_, Oct 27 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Mar 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016571 | OEIS-A371673

Expansion of g.f. A(x) satisfying [x^(n-1)] A(x)^(n^2) = A000108(n-1) * n^n for n >= 1, where A000108 is the Catalan numbers.

Conjecture: a(n) is odd for n > 0 iff n = 2*A003714(k) + 1 for some k, where A003714 is the Fibbinary numbers (integers whose binary representation contains no consecutive ones). See A263075, A263190, and A171791.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 02 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016572 | OEIS-A371710

Expansion of e.g.f. A(x) satisfying Sum_{n>=0} (A(x)^n - x)^n / n! = 1.

Conjecture: for n > 0, a(6*n + k) == [2,0,2,1,2,2] (mod 3) at k = [0,1,2,3,4,5], respectively.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016573 | OEIS-A371737

Number of quanimous strict integer partitions of n, meaning there is more than one set partition with all equal block-sums.

Conjecture: (1) Positions of 0's are A327782. (2) Positions of terms > 0 are A368459.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Apr 14 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016574 | OEIS-A371761

Array read by antidiagonals: The number of parades with n girls and k boys that begin with a girl and end with a boy.

Conjecture: Row n + 1 is row 2^n in table A136301, where a probabilistic interpretation is given (see the link to Parsonnet's paper below).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Apr 05 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016575 | OEIS-A371798

a(n) = Sum_{k=0..floor(n/2)} (-1)^k * binomial(2*n-2*k-1,n-2*k).

Conjectured g.f.: 1 + x*(4 - 10*x + 8*x^2)/(2 - 11*x + 14*x^2 - 8*x^3 + (2 - 3*x)*sqrt(1 - 4*x)) (see Elizalde et al. at p. 13). - _Stefano Spezia_, Dec 27 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Apr 06 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016576 | OEIS-A371823

Triangle T(n, k) read by rows: Maximum number of patterns of length k in a permutation from row n in A371822.

Conjecture: T(n, n-2) = ceiling(n*(n-1)/2), for n > 6. This is expected because this triangle does asymptotically approximate the factorial numbers from the left to the right and Pascal's triangle from right to the left.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Jun 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016577 | OEIS-A371835

Triangle read by rows: T(n,k) is the number of points (x,y,z) satisfying |x|+|y|+|z|<=n and max(|x|,|y|,|z|)<=k; 0<=k<=n.

Conjectures: The two conjectures are true. See links. - _Sela Fried_, Jul 05 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Apr 07 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016578 | OEIS-A371898

Triangle read by rows: T(n, k) = n * k * (T(n-1, k-1) + T(n-1, k)) for k > 0 with initial values T(n, 0) = 1 and T(i, j) = 0 for j > i.

Conjecture: E.g.f. of column k is exp(t) * t^k * k! / (Prod_{i=0..k} (1 - i*t)). Conjecture: Sum_{k=0..n} (-1)^(n-k) * T(n, k) = A000166(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Apr 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016579 | OEIS-A371904

a(1) = 1; for n > 1, a(n) is the smallest unused positive number such that 2^a(n) contains a(n-1) as a substring.

The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Apr 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016580 | OEIS-A371916

Zeroless analog of tetranacci numbers.

It is not known whether this sequence cycles, but it is conjectured to cycle just like A243063 and A371911 (have periods of 912 and 300056874, respectively) because the expected growth factor in the number of digits of successive terms is 0.9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bryle Morga_, Apr 12 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016581 | OEIS-A371918

The smallest unused positive number such that 2^a(n) contains n as a substring.

The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Apr 12 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016582 | OEIS-A372015

Product of Fibonacci and self-convolution of Fibonacci numbers: a(n) = A000045(n+1)*A001629(n+1).

Conjecture: a(n) is the total number of pairs of adjacent parts that are the same color in all n-color compositions of n+1. - _John Tyler Rascoe_, Jul 30 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vladimir Kruchinin_, Apr 15 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016583 | OEIS-A372018

G.f. A(x) satisfies A(x) = (1 + 4*x*A(x)/(1 - x*A(x)))^(1/2).

Conjecture: a(2n+1) = 2*A371364(). - _R. J. Mathar_, Apr 22 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Apr 15 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016584 | OEIS-A372031

Lexicographically earliest sequence of distinct positive integers such that for any n > 0, a(2*n+2) = a(2*n+1) * a(n+1).

Conjecture: this sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Apr 16 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016585 | OEIS-A372032

Lexicographically earliest sequence of distinct positive integers such that for any n > 0, a(2*n+1) = a(2*n) XOR a(n) (where XOR denotes the bitwise XOR operator).

Conjecture: this sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Apr 16 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016586 | OEIS-A372060

a(n) = smallest k such that A252867(k) = n, or -1 if no such k exists.

Both A252867 and this sequence are conjectured to be permutations of the nonnegative integers, in which case they are inverses of each other.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016587 | OEIS-A372063

a(n) = A280864(n) - A280866(n).

A280864 and A280866 are closely related. There is a proof that A280866 is a permutation of the positive integers, but for A280864 this is only a conjecture. The present sequence compares them term-by-term; A372064 compares their inverses; and A372065 compares where the primes appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016588 | OEIS-A372078

Primes p that are introduced in A373390 by 3*p.

Conjecture: there are no other terms. (See the comments and conjectures in A373790.) 157 is the 37th prime. The conjecture is that for the 38th and all later primes p, in A373390 there are three successive terms 2*p, x, p. The values of x are given in A373790.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 18 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016589 | OEIS-A372118

Square array A(n, k) = ((k+2)^(n+2) - 2 * (k+1)^(n+2) + k^(n+2))/2 for k, n >= 0 read by ascending antidiagonals.

Further properties are conjectures: The three conjectures are true, see links. - _Sela Fried_, Jul 07 2024 Conjectures: The three conjectures are true. See comments. - _Sela Fried_, Jul 09 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Apr 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016590 | OEIS-A372127

Lexicographically earliest sequence of distinct positive integers such that for any n > 0, a(2*n+2) = LCM(a(2*n+1), a(n+1)).

Conjecture: this sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Apr 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016591 | OEIS-A372128

Lexicographically earliest sequence of distinct positive integers such that for any n > 0, a(2*n+1) = a(2*n) OR a(n) (where OR denotes the bitwise OR operator).

Conjecture: this sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Apr 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016592 | OEIS-A372143

Lexicographically earliest sequence of distinct nonnegative integers such that for any n > 0, the binary expansions of a(n), a(2*n) and a(2*n+1) have a common 1 bit.

Conjecture: this sequence is a permutation of the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Apr 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016593 | OEIS-A372211

a(n) = [x^n] f(x)^n, where f(x) = (1 - x^5)^5/((1 - x^2)^2 * (1 - x^3)^3).

We conjecture that in this case the stronger supercongruences a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) hold for all primes p >= 5 and positive integers n and r. Some examples are given below. Cf. A351858.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016594 | OEIS-A372212

a(n) = [x^n] f(x)^n, where f(x) = (1 - x^7)^7/((1 - x^2)^2 * (1 - x^5)^5).

We conjecture that in this case the stronger supercongruences a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) hold for primes p >= 11 and positive integers n and r. Some examples are given below. Cf. A351858.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016595 | OEIS-A372214

a(n) is equal to the n-th order Taylor polynomial (centered at 0) of G(x)^n evaluated at x = 1, where G(x) = (1 - 2*x - sqrt(1 - 8*x + 4*x^2))/(2*x).

We conjecture that the sequence satisfies the stronger supercongruences a(n*p^k) == a(n*p^(k-1)) (mod p^(3*k)) for all primes p >= 5 and positive integers n and k. Examples of these supercongruences are given below. More generally, for each integer m, we conjecture that the

sequence {a_m(n) : n >= 0}, defined by setting a_m(n) = the n-th order Taylor polynomial of G(x)^(m*n) evaluated at x = 1, satisfies the same supercongruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Apr 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016596 | OEIS-A372224

The size of the smallest critical set of hints needed to uniquely solve a generalized n X n Sudoku board.

When n is prime, a(n) is the size of smallest critical set of an n X n Latin square, which is conjectured to equal A002620(n). When n is prime, a(n) is conjectured to equal A002620(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Agustin Gomez de la Vega_ and _Mithra Karamchedu_, Apr 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016597 | OEIS-A372239

Expansion of (1 + 2*x) / ((1 - 2*x)*sqrt(1 - 4*x)).

Conjecture: For p Pythagorean prime (A002144), a(p) - 6 == 0 (mod p). Conjecture: For p prime of the form 4*k + 3 (A002145), a(p) + 2 == 0 (mod p).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mélika Tebni_, Apr 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016598 | OEIS-A372313

Table read by antidiagonals: A(n,1) = 2n-1, and for k > 1, A(n,k) = A372289(A(n,k-1)+A(n,1)).

Conjecture: if A(n,1) is congruent to 33 (mod 100), then all terms on the n-th row are congruent to 33 (mod 100).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ali Sada_, Apr 26 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016599 | OEIS-A372314

Determinant of the matrix [Jacobi(i^2 + 3*i*j + 2*j^2, 2*n + 1)]_{1 < i, j < 2*n}, where Jacobi(a, m) denotes the Jacobi symbol (a / m).

Conjecture 1: Let n be any positive integer. The following conjecture is similar to Conjecture 1. Conjecture 2: (1) {2, 2}_p = 0 for any prime p == 13,19 (mod 24), and {2, 2}_p == 0 (mod p) for any prime p == 17,23 (mod 24). We have verified Conjecture 2 for p or n smaller than 2000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 27 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016600 | OEIS-A372368

a(1) = 1. Thereafter, if A007947(a(n-1)) is a term in A002110, a(n) is the smallest prime not already in the sequence. Otherwise a(n) is the smallest novel multiple of p, the greatest prime < g = gpf(a(n-1)).

A term whose radical is in A002110 occurs only if the multiplier m (of prime q) is in A002110, either with q|m, or with q the smallest prime > gpf(m). Thus all multiples of every prime will appear and it is therefore conjectured that this sequence is a permutation of the positive integers, A000027.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_ and _Michael De Vlieger_, Apr 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016601 | OEIS-A372385

Determinant of the matrix [Jacobi(i - j, 2n + 1)]_{0 < i, j < n}, where Jacobi(a, m) denotes the Jacobi symbol (a/m).

In a 2003 preprint, R. Chapman conjectured that if 2*n + 1 (with n > 1) is a prime congruent to 3 modulo 4, then a(n) = 1. Conjecture 1: If p = 2*n + 1 is a prime congruent to 1 modulo 4, then det[x + Jacobi(i - j, p)]_{0 < i, j < n} = r_p - 2*s_p + (-1)^{n/2}*x*(2*r_p - p*s_p), and det[x + Jacobi(i - j, p)]_{0 <= i, j <= n} = -r_p + (-1)^{n/2}*p*s_p*x, where r_p and s_p are rational numbers such that e_p^{(2 - Jacobi(2, p))*h(p)} = r_p + s_p*sqrt(p), e_p and h(p) are the fundamental unit and the class number of the real quadratic field Q(sqrt(p)), respectively. Conjecture 2: Suppose that p = 2*n + 1 is a prime greater than 3, and write e_p^{h(p)} = a_p + b_p*sqrt(p) with 2*a_p and 2*b_p integral. We have checked both conjectures for n up to 1000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 29 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016602 | OEIS-A372409

Determinant of the matrix obtained from [Jacobi(i-j, 2*n+1)]_{0<i,j<n} by replacing all the entries in the first row by 1, where Jacobi(a,m) denotes the Jacobi symbol (a/m).

Conjecture 1: If 2*n + 1 (with n > 1) is a prime p, then a(n)/floor((p-2)/3) coincides with (-1)^((p+3)/4) if p == 1 (mod 4), and (-1)^((h(-p)+1)/2) if p == 3 (mod 4), where h(-p) is the class number of the imaginary quadratic field Q(sqrt(-p)). Conjecture 2: Let p > 3 be a prime, and let S(p) and T(p) denote the matrices obtained from [Jacobi(i+j,p)]_{1<=i,j<=(p-3)/2} and [Jacobi(i+j,p)]_{0<=i,j<=(p-3)/2} (respectively) by replacing all the entries in the first row by 1. Then det S(p) = -det T(p) = 2^((p-5)/2)*s(p), where s(p) is (-1)^((p+3)/4) if p == 1 (mod 4), and (-1)^((h(-p)-1)/2) if p == 3 (mod 4). Both conjectures are motivated by Conjecture 4.6 in the author's 2019 FFA paper as well as the conjectures in A372385. They have been verified for primes p < 2000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 29 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016603 | OEIS-A372420

Expansion of (1 + x) / ((1 - 2*x)*sqrt(1 - 4*x)).

Conjecture: For p Pythagorean prime (A002144), a(p) - 5 == 0 (mod p). Conjecture: For p prime of the form 4*k + 3 (A002145), a(p) + 1 == 0 (mod p).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mélika Tebni_, Apr 30 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016604 | OEIS-A372430

Positive integers k such that the distinct prime indices of k are a subset of the binary indices of k.

Conjecture: The only number whose binary indices are a subset of its prime indices is 4100, with binary indices {3,13} and prime indices {1,1,3,3,13}. Verified up to 10,000,000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, May 02 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016605 | OEIS-A372457

a(n) is the least k such that k^2 + k + 1 is divisible by the n-th power of a prime.

For n > 1 the prime is in A002476. Conjecture: it is always 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, May 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016606 | OEIS-A372484

Positions of local maxima of the probability that the procedure described in A372422 successfully ends with the selection of a single person.

It is conjectured that lim_{n->oo} a(n+1)/a(n) = 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, May 07 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016607 | OEIS-A372485

Positions of local minima of the probability that the procedure described in A372422 successfully ends with the selection of a single person.

It is conjectured that lim_{n->oo} a(n+1)/a(n) = 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, May 07 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016608 | OEIS-A372514

Index k such that A280864(k) = A019565(n) or 0 if A019565(n) does not appear in A280864.

Conjecture: there are no zeros in this sequence, which is equivalent to the conjecture that A280864 is a permutation of natural numbers. Minor corollary: a(127) > 2^18.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jul 29 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016609 | OEIS-A372524

The smallest composite number k that shares exactly n distinct prime factors with sopfr(k), the sum of the primes dividing k, with repetition.

Conjecture: A001221(a(n)) = n+1, for n >= 2. - _Daniel Suteu_, May 08 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 05 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016610 | OEIS-A372527

Expansion of g.f. A(x) satisfying A(A(x)^2) = x*A(x)/(1 - 2*A(x)).

Conjecture: a(2^k + 1) == 2 (mod 4) for k >= 0, and a(n) == 0 (mod 4) for n > 0 when n is not of the form 2^k + 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 13 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016611 | OEIS-A372529

Expansion of g.f. A(x) satisfying A(x) = A(x^3 + x^4) / A(x^2).

Conjecture: lim_{n_>oo} abs(a(n))^(-1/n) = r where r = 0.75487766624669276... is the real root of x^3 + x^2 - 1 = 0; that is, r = (((25 + 3*sqrt(69))/2)^(1/3) + ((25 - 3*sqrt(69))/2)^(1/3) - 1) / 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016612 | OEIS-A372536

Expansion of g.f. A(x) satisfies A(A(x)^2/(1 + 3*A(x))^2) = x*A(x).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016613 | OEIS-A372547

Palindromic Zumkeller numbers in base 10.

Based on the conjecture that there are infinitely many repunit primes, one may conjecture there are infinitely many repdigit terms, since 2*3*y (where y is a repunit prime) is a repdigit palindromic Zumkeller number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ivan N. Ianakiev_, May 05 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016614 | OEIS-A372555

Least number of Jacobsthal numbers that add up to n.

See conjecture in A372556, and also in A372561.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, May 07 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016615 | OEIS-A372561

Array read by upward antidiagonals: A(n, k) = A265745(A372560(n, k)) for n > 1, k >= 1.

911, 58325, 933205, 238900565, 15656587449685, 67244531063362552157525, etc. I conjecture that for the former sequence of numbers x, from 933 onward, A372555(x) = 7, and for the latter sequence of numbers y, from 58325 onward, A372555(y) = 9, and that the array A372555(A372560(n, k)) is symmetric apart from its borders, i.e, that for n, k > 1, A372555(A372560(n, k)) = A372555(A372560(k, n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, May 08 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016616 | OEIS-A372577

Expansion of g.f. A(x) satisfying A(x)^2 = A(A(x*A(x) + x*A(x)^2)).

Conjecture: a(6*n - 3) == 3 (mod 4) and a(6*n - k) == 1 (mod 4) when k = {0,1,2,4,5} for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 30 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016617 | OEIS-A372607

Let a(1) = 2, f(n) = a(1)*a(2)*...*a(n-1) for n >= 1 and a(n) = nextludicnumber(f(n)+1) - f(n) for n >= 2, where nextludicnumber(x) is the smallest ludic number > x.

Conjecture: every element is a ludic number. This is the analog of Buss' conjecture (cf. A067836) for ludic numbers instead of primes, and similar to the idea of ludic Fortunate numbers (A376237) in analogy to the usual Fortunate numbers A005235. - _M. F. Hasler_, Nov 04 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Davide Rotondo_, May 07 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016618 | OEIS-A372611

Expansion of (1 + 3*x) / ((1 - 2*x)*sqrt(1 - 4*x)).

Conjecture: For p Pythagorean prime (A002144), a(p) - 7 == 0 (mod p). Conjecture: For p prime of the form 4*k + 3 (A002145), a(p) + 3 == 0 (mod p).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mélika Tebni_, May 07 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016619 | OEIS-A372697

Index k such that A280866(k) = A019565(n) or 0 if A019565(n) does not appear in A280866.

Conjecture: there are no zeros in this sequence, which is equivalent to the conjecture that A280866 is a permutation of natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jul 29 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016620 | OEIS-A372699

a(1) = 1, j = a(n-1). For n > 1, a(n) is the least novel m such that rad(m*j) is the smallest primorial number divisible by A007947(j).

Primes are in order and if p is prime and p|a(n) there is an i <= n such that a(i) = p (no multiple of p appears prior to p). Sequence is conjectured to have "Property S" of A368900. Also, for integers x, y with x < y and rad(x) = rad(y), x appears in the sequence before y. Conjecture: Sequence is a permutation of the positive integers which preserves the above mentioned properties of A000027.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_ and _Michael De Vlieger_, May 14 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016621 | OEIS-A372700

a(n) = Sum_{j=1..n} Sum_{k=1..n} mu(j*k), where mu is the Moebius function A008683.

Conjecture: a(n) changes sign infinitely often.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, May 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016622 | OEIS-A372706

a(n) = Sum_{k=1..n} mu(n*k), where mu is the Moebius function A008683.

Conjecture: a(n) = A008683(n) * A112399(n). - _Jamie Morken_, Sep 15 2026 The conjecture is correct - see my link. - _Peter Bala_, Sep 22 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, May 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016623 | OEIS-A372707

Smallest natural number requiring n applications of the map x -> 2^x mod x = A015910(x) to reach 0.

A015910 is conjectured to contain every natural number except for 1, which would imply that this sequence has infinitely many terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bryle Morga_, May 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016624 | OEIS-A372709

a(n) = Sum_{k=1..n} tau(n*k), where tau(n) = number of divisors of n, cf. A000005.

Conjecture: a(n) ~ A372674(n) / 2. - _Vaclav Kotesovec_, May 13 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, May 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016625 | OEIS-A372710

a(n) = Sum_{k=1..n} sigma(n*k).

Conjecture: a(n) ~ A372675(n) / 2 ~ Pi^4 * n^4 / (288*zeta(3)). - _Vaclav Kotesovec_, May 13 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, May 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016626 | OEIS-A372763

Denominator of the continued fraction 1/(2-3/(3-4/(4-5/(...(n-1)-n/(n+5))))).

Conjecture 1: The sequence contains only 1's and the primes. Conjecture 2: Except for 2 and 3, all primes appear in the sequence once. Conjecture: Record values correspond to A045375(m), m > 2. - _Bill McEachen_, Aug 03 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mohammed Bouras_, May 12 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016627 | OEIS-A372846

a(n) is the number of states in the smallest deterministic finite automaton that accepts the Zeckendorf representation of i and n*i, in parallel, for all integers i>=0.

Conjecture: a(n) <= 2*n^2 + g^2*n + 1, where g = (1+sqrt(5))/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Jul 04 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016628 | OEIS-A372864

Numbers k such that A372720(k) = 0.

Conjecture: proper subset of A361098, hence of A360765 and A360768. This is to say that k = a(n) is such that A003557(k) >= A119288(k), i.e., k/rad(k) >= second smallest prime factor of k, and A003557(k) > A053669(k), where A053669(k) is the smallest prime q that does not divide k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jun 02 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016629 | OEIS-A372865

a(n) = (Sum_{k = 1..n-1} k*a(k)) (mod n) with a(1) = 1.

More generally, define a sequence {a(n, s) : n >= 1} with starting parameter s by a(n, s) = (Sum_{k = 1..n-1} k*a(k, s)) (mod n) with a(1, s) = s. The sequence {a(n, s)} is conjectured to be one of 3 types as illustrated by the following examples for s in [1..100].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, May 15 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016630 | OEIS-A372867

Distinct terms in A242017, listed in the order of their appearance.

Conjecture: every term except 3 belongs to A366609. - _Bill McEachen_, Jun 12 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-Marc Rebert_, May 15 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016631 | OEIS-A372891

Anti-elite primes (A128852) that are not prime factors of Fermat primes (A023394).

Conjecture: All terms >= 97 are congruent to 1 modulo 8. (Note that every factor of Fermat numbers >= 17 is congruent to 1 modulo 8.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, May 15 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016632 | OEIS-A372916

Indices of record high values in A372915.

Conjectures (checked up to A372915(666)): (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Jun 17 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016633 | OEIS-A372974

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest unused positive number such that a(n) is coprime to a(n-1) and omega(a(n)) does not equal omega(a(n-1)).

The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 26 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016634 | OEIS-A372975

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest unused positive number such that a(n) shares a factor with a(n-1) while omega(a(n)) does not equal omega(a(n-1)).

The fixed points begin 1, 2, 22, 26, 36, 38, 1991, 2023, 2159, 2189, 2627; it is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 26 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016635 | OEIS-A372991

a(n) = (2n)!/(a(n-1)*a(n-2)), where a(0)=1, a(1) = 1.

Conjecture D-finite with recurrence a(n) -2*n*(2*n-1)*a(n-3)=0. - _R. J. Mathar_, Jul 16 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 15 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016636 | OEIS-A373049

Integers k such that the product of the nonzero digits of the k-th Fibonacci number (A000045) is a perfect power.

Conjecture: this sequence has infinitely many terms. Since the product of the nonzero digits of Fibonacci(k) is of the form 2^a * 3^b * 5^c * 7^d, a sufficient condition for Fibonacci(k) to belong to the sequence is that a, b, c and d are all even.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gonzalo Martínez_, May 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016637 | OEIS-A373097

a(n) = a[n/r] + a[n/r^2] + a[n/r^3] + ... , where a(0) = 0, a(1) = 1, r = 3/2, and [ ] = floor.

Conjecture: Let d(n) = a(n+1) - a(n) for n >= 0. Then d(n) = 1 for infinitely many n, and (d(n)) is unbounded.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016638 | OEIS-A373110

Number of distinct circles that can be constructed from the 4 vertices and the equally spaced 4*n points placed on the sides of a square when every pair of the 4 + 4*n points are connected by a circle and where the points lie at the ends of the circle's diameter.

Conjectured:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016639 | OEIS-A373116

Fibonacci numbers whose digits product is a positive perfect power (A001597).

Since the product of the digits of Fibonacci(k) is required to be positive, Fibonacci(k) does not have zero as a digit. For this reason this list is probably finite, since it is conjectured that there are only finitely many Fibonacci numbers that do not contain the digit zero (see A076564). If the conjecture is true, the largest number possessing the property would be Fibonacci(85) = 259695496911122585 whose digit product is 194400^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gonzalo Martínez_, May 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016640 | OEIS-A373162

a(n) = denominator of Sum_{k>=0} cos(2k*Pi/3)/n^k.

Conjecture: the sequence of fractions is completely monotonic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016641 | OEIS-A373163

a(n) = numerator of Sum_{k>=0} cos(2k*Pi/3)/n^k.

Conjecture: the sequence of fractions is completely monotonic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016642 | OEIS-A373226

Number of points in a diagonal Vicsek fractal subset of an n X n square.

Conjecture: a(n) = O(n^log_3(5)). This is motivated by the log_3(5) Hausdorff dimension of the diagonal Vicsek fractal. Conjecture: a(3n) = 5*a(n). This is motivated by the recursive construction of the diagonal Vicsek fractal. Conjecture: a(2n+1) = A151907(n)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sidharth Ghoshal_, May 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016643 | OEIS-A373308

Expansion of Product_{n>=0} (1 - x^(2^n))^3.

Conjecture: a(3*n) == A001285(n) (mod 3) for n >= 0, where A001285 is the Thue-Morse sequence: [1, 2, 2, 1, 2, 1, 1, 2, ...]. Conjecture: a(3*n+1) and a(3*n+2) are divisible by 3 for n >= 0. Conjecture: a(n) = 0 iff n = 3*A000695(k) - 1 for k >= 1, where A000695 lists sums of distinct powers of 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016644 | OEIS-A373312

Expansion of g.f. A(x) satisfying A(x)^2 = A(x*A(x)/(1 - A(x))^2).

Conjecture: a(n) == 1 (mod 2) iff n = 2^k - 1 for k >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016645 | OEIS-A373317

Numbers which are their own imbalance when written in English (see the definition of the imbalance in the Comments section).

This sequence is conjectured to be finite and full.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Jun 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016646 | OEIS-A373321

Numbers which are their own imbalance when written in French (see the definition of the imbalance in the Comments section).

This sequence is conjectured to be finite and full.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Angelini_ and _Jean-Marc Falcoz_, Jun 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016647 | OEIS-A373323

a(1,2) = 1,2; i = a(n-2), j = a(n-1). For n > 2 a(n) is least novel k (with A001221(k) != w if A001221(i) = A001221(j) = w), such that A007947(i*j*k) is a term in A002110.

In other words, a(n) is the smallest novel k satisfying the above constraint on omega(k) such that the squarefree kernel of i*j*k is a primorial number. Conjectured to be a permutation of the positive integers (A000027), with primes in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jun 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016648 | OEIS-A373348

Squarefree numbers k such that there exists a k-digit multiple of k that has k divisors.

Conjecture: a(3) = 206 = 2*103 and a(4) = 500015 = 5*100003 are the only semiprime terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Jun 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016649 | OEIS-A373350

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest unused positive number such that omega(a(n)) does not equal omega(a(n-1)) or omega(a(n-2)).

The fixed points begin 1, 2, 15, 18, 125, 137, 2737, 120051; it is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016650 | OEIS-A373390

a(n) = n for n <= 3; for n > 3, a(n) is the smallest unused positive number that is coprime to a(n-1) but has a common factor with at least one of a(1)...a(n-2).

A373790 has several as-yet unproved conjectures related to this sequence. - _N. J. A. Sloane_, Jun 23 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 03 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016651 | OEIS-A373409

Length of the n-th maximal antirun of nonsquarefree numbers differing by more than one.

Conjecture: The maximum is 9, and there is no antirun of more than 9 nonsquarefree numbers. Confirmed up to 100,000,000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jun 06 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016652 | OEIS-A373421

Positive integers that cannot be written as a sum of a practical number and a pentagonal number.

Somu and Tran (2024) conjectured that there are finitely many such integers. It was also conjectured that 2671 is the largest such integer. This conjecture was checked up to 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Duc Van Khanh Tran_, Jun 04 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016653 | OEIS-A373443

Positive integers that cannot be written as a sum of a practical number and a hexagonal number.

Somu and Tran (2024) conjectured that there are finitely many such integers. It was also conjectured that 1332329 is the largest such integer. This conjecture was checked up to 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Duc Van Khanh Tran_, Jun 05 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016654 | OEIS-A373444

Positive integers that cannot be written as a sum of a practical number and a heptagonal number.

Somu and Tran (2024) conjectured that there are finitely many such integers. It was also conjectured that 79445 is the largest such integer. This conjecture was checked up to 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Duc Van Khanh Tran_, Jun 05 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016655 | OEIS-A373487

Numbers k such that A276085(k) is a multiple of A003415(k), where A003415 is the arithmetic derivative and A276085 is fully additive with a(p) = p#/p.

Sequence A276086(A380527(.)) when sorted into ascending order gives the {intersection of this sequence with A048103}. As A380527 is a subsequence of A381035, this intersection is by the given correspondence a subsequence of (primes U A080364). Note that this applies also to the third known nonsquarefree term of this sequence, 855523 (= 67 * 113^2), which indeed disproves the conjecture I had made in A380527, while the first two nonsquarefree terms 28 and 33212 are both multiples of 4, and thus outside of the intersection.- _Antti Karttunen_, Jan 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jun 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016656 | OEIS-A373490

Numbers k for which A003415(k) is a multiple of A276085(k), where A003415 is the arithmetic derivative and A276085 is the primorial base log-function.

Conjecture: if term present here is squarefree, then it must be a term of A369650: 1, 2, 10, 15, 5005, and possibly no others. Or put in other words, if A003415(k) = e*A276085(k), then if e > 1, it is conjectured that k cannot be squarefree (in A005117). A stricter form of the above conjecture is given in A400171. Question: Does the intersection with A399379 contain any other terms after 1, 2, 10, 15, 20, 5005, ? - _Antti Karttunen_, Sep 23 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jun 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016657 | OEIS-A373516

Number of sets that can be represented as a length-n combination of commas and braces, with elements possibly repeated.

It is conjectured that there are only finitely many n such that a(n) > a(n+1), with n=11 being the last example.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bryle Morga_, Jun 08 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016658 | OEIS-A373545

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest unused positive number that shares a factor with a(n-1) if a(n-1) is odd otherwise is coprime to a(n-1) if a(n-1) is even.

The terms studied are all concentrated along lines of different gradient; the odd terms all being on the line a(n) = n or very close to it, while the even terms are distributed among the other lines dependent on the number of prime factors they contain. It is conjectured that this behavior is true for all n. The topmost line contains even semiprimes while the lower lines, those with gradient less than 1, contain powers of 2 multiplied by a larger single prime. However the topmost of these lower lines, which is more diffuse, also contains all other even numbers. Also noticeable is some of the lower lines terminating after which these values appear to move into the line contains all other even numbers. Conjecture: Odd prime p precedes 2*p. Conjecture: Odd prime p appears for odd n, but additionally, tends to occupy a(n) such that n mod A002110(k) is a reduced residue of the largest A002110(k) < n. Thus, a(n) = p for n <= p. Example, a(65) = 67 is displaced. However, generally primes represent fixed points. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016659 | OEIS-A373560

a(n) is the smallest multiple of prime(n)^2 that starts a run of 5 consecutive integers with 6 divisors, or -1 if no such multiple exists.

Conjecture: the converse is also true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Jun 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016660 | OEIS-A373561

a(n) = (1/3)*n^5 + (1/2)*n^4 + (1/6)*n^3.

Conjecture: a(n) = Sum_{k=1..n} Sum_{z=1..n} Sum_{y=1..n} Sum_{x=1..n} [GCD(f(x,y,z), n) = k] * f(x,y,z), where f(x,y,z) = x^2 + y^2 - z^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Jun 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016661 | OEIS-A373610

Positive integers that cannot be written as a sum of a practical number and an octagonal number.

Somu and Tran (2024) conjectured that there are finitely many such integers. It was also conjectured that 4005819 is the largest such integer. This conjecture was checked up to 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Duc Van Khanh Tran_, Jun 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016662 | OEIS-A373611

Positive integers that cannot be written as a sum of a practical number and a nonagonal number.

Somu and Tran (2024) conjectured that there are finitely many such integers. It was also conjectured that 325808 is the largest such integer. This conjecture was checked up to 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Duc Van Khanh Tran_, Jun 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016663 | OEIS-A373613

Positive integers that cannot be written as a sum of a practical number and a decagonal number.

Somu and Tran (2024) conjectured that there are finitely many such integers. It was also conjectured that 13613213 is the largest such integer. This conjecture was checked up to 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Duc Van Khanh Tran_, Jun 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016664 | OEIS-A373615

Positive integers that cannot be written as a sum of a practical number and an 11-gonal number.

Somu and Tran (2024) conjectured that there are finitely many such integers. It was also conjectured that 105712 is the largest such integer. This conjecture was checked up to 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Duc Van Khanh Tran_, Jun 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016665 | OEIS-A373616

Positive integers that cannot be written as a sum of a practical number and a 13-gonal number.

Somu and Tran (2024) conjectured that there are finitely many such integers. It was also conjectured that 1612172 is the largest such integer. This conjecture was checked up to 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Duc Van Khanh Tran_, Jun 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016666 | OEIS-A373617

Positive integers that cannot be written as a sum of a practical number and a 14-gonal number.

Somu and Tran (2024) conjectured that there are finitely many such integers. It was also conjectured that 106878 is the largest such integer. This conjecture was checked up to 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Duc Van Khanh Tran_, Jun 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016667 | OEIS-A373623

a(n) = n for n <= 3; for n > 3, a(n) is the smallest unused positive number that shares a factor with the most recently appearing even number if a(n-1) is odd, otherwise it shares a factor with the most recently appearing odd number if a(n-1) is even.

The terms studied are all concentrated along lines of different gradient; the prime numbers are all distributed on the lines with gradient < 1 while all other numbers are on the lines with gradient > 1, the highest concentration of terms appearing on a line with gradient ~ 1.04. The uppermost of these lines appear to consist entirely of numbers with two or three distinct prime factors. See the attached images. It is conjectured that this behavior is true for all n. In the first 200000 terms the longest run of consecutive even terms is eight, beginning at a(7681) = 8078, while the longest run of consecutive odd terms is seven, beginning at a(101234) = 105471. It in unknown if these runs can be of arbitrary length. The sequence is conjectured to be a permutation of the positive numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016668 | OEIS-A373633

Positive integers that cannot be written as a sum of a practical number and a 15-gonal number.

Somu and Tran (2024) conjectured that there are finitely many such integers. It was also conjectured that 1486748 is the largest such integer. This conjecture was checked up to 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Duc Van Khanh Tran_, Jun 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016669 | OEIS-A373634

Positive integers that cannot be written as a sum of a practical number and a 17-gonal number.

Somu and Tran (2024) conjectured that there are finitely many such integers. It was also conjectured that 9314 is the largest such integer. This conjecture was checked up to 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Duc Van Khanh Tran_, Jun 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016670 | OEIS-A373646

Positive integers that cannot be written as a sum of a practical number and an 18-gonal number.

Somu and Tran (2024) conjectured that there are finitely many such integers. It was also conjectured that 8541224 is the largest such integer. This conjecture was checked up to 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Duc Van Khanh Tran_, Jun 12 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016671 | OEIS-A373650

Integers k such that there are i groups of order k+i up to isomorphism, for i=1,2,3,4.

Conjectures (verified to 783 terms): If n = p-2 mod p, then n+2 = 2pm for some m, since n is even. Then if m > 1, one can show that this will always have 4 or more groups. Thus m=1 and n=2p-2. One must show that n can never be of the form 2p-2 if this conjecture is true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robin Jones_, Jun 12 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016672 | OEIS-A373686

a(n) is the number of ways n can be written as a sum of a practical number and two squares.

Somu and Tran (2024) proved that a(n) > 0 for sufficiently large n and conjectured that a(n) > 0 for all n > 0. The conjecture was checked up to 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Duc Van Khanh Tran_, Jun 13 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016673 | OEIS-A373698

a(n) = n for n <= 3; for n > 3, a(n) is the smallest unused positive number that shares a factor with the largest odd divisor of a(n-1).

The terms studied are all concentrated along lines of different gradient, the prime numbers, which appear in their natural order, forming the lowermost line. The central line has the highest concentration of terms and is composed of numbers with various numbers of prime factors, while many of the other lines consist of terms with the same number of prime factors. The odd terms all fall into the highest, lowest, and middle line, all other lines containing only even terms. It is conjectured the above behaviors are true for all n. See the attached images.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 13 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016674 | OEIS-A373725

Numbers k such that A011776(k) = A011776(k+1).

Conjecture: if p != 3 is a prime such that 2*p-1 is also a prime (p is in A005382 \ {3}), then 4*p^2 - 1 is a term of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Jun 15 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016675 | OEIS-A373737

a(n) is the smallest number k in the sorted sequence S(q) = {k : rad(k) = q}, q = A120944(n), such that tau(k) - A008479(k) is not positive, where rad = A007947 and tau = A000005.

Conjecture: a(n) <= s*gpf(s)^floor(log_gpf(s) s^2), where gpf = A006530.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jun 24 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016676 | OEIS-A373738

a(1) = 1, a(n) = floor((1/omega(n)!) * Product_{p | n} 1 + (log n)/(log p)), where omega = A001221.

Conjecture: A010846(k) - a(k) approaches 0 as k increases toward infinity, for k with omega(k) > 1 that have the same squarefree kernel r. Therefore, the difference is most significant for composite squarefree k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jul 16 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016677 | OEIS-A373774

a(n) is the number of terms of A000057 in [10^n].

Conjectured by Kumar under GRH: Limit_{n->oo} a(n)/A006880(n) = (10/19)*A005596 = 0.196818849273264362134067397024429692164015394...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jun 18 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016678 | OEIS-A373783

Number of ways that people can sit in n linearly arranged seats such that there are two or three empty seats between any two persons, zero, one or two empty seats at the start and end, and at least one person gets seated.

Conjecture: a(n) = (-7*a(n-1) - 7*a(n-2) + (n-2)*a(n-3) + 3*(n+1)*a(n-4) + 4*(n-1)*a(n-5) + 3*(n-3)*a(n-6) + (n-4)*a(n-7))/3 for n >= 9. - _Pontus von Brömssen_, Jun 23 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enrique Navarrete_, Jun 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016679 | OEIS-A373789

Indices of record high points in A372072.

Conjecture: Consists of 1, 4, 6, 10, and the primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016680 | OEIS-A373790

The term that immediately precedes prime(n) in A373390.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 21 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016681 | OEIS-A373791

a(n) = 0 if p = prime(n) is introduced in A373390 by 2*p, or 1 if p is introduced by 3*p.

It is conjectured (see A373790) that every prime p in A373390 is introduced by either 2*p or 3*p, and that exactly 11 primes (listed in A372078) are introduced by 3*p. This would imply that the present sequence is 0 after the 35th term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 21 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016682 | OEIS-A373848

Numbers k not divisible by p^p for any prime p, and for which 1 < A373842(k) <= k, where A373842 is the arithmetic derivative of the primorial base log-function.

It is conjectured that 5 is the only fixed point of A373842, which would imply that x=6 is the only number for which A003415(x) = A276086(x). See A351228.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jun 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016683 | OEIS-A373853

Row sums of A373557.

Conjecture: a(n) = A373346(n) + A007895(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Jun 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016684 | OEIS-A373860

a(n) = n for n <= 4; for n > 4, a(n) is the smallest unused positive number that shares a factor with the most recently appearing even number prior to a(n-1) if a(n-1) is even, otherwise it shares a factor with the most recently appearing odd number prior to a(n-1) if a(n-1) is odd.

The fixed points are 1, 2, 3, 4, 16, 32, 124, and it is likely that no more exist. The sequence is conjectured to be a permutation of the positive numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016685 | OEIS-A373867

Perfect powers of the form x^y + y^x, where x > 1 and y > 1.

Conjecture: 32 and 100 are the only terms for which x != y: 2^4 + 4^2 = 2^5 = 32 and 2^6 + 6^2 = 10^2 = 100.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gonzalo Martínez_, Jun 21 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016686 | OEIS-A373887

a(n) is the length of the longest arithmetic progression of semiprimes ending in the n-th semiprime.

Conjectures: a(n) >= 3 for n >= 16. If A001358(n) is divisible by A000040(m), then a(n) >= A373888(m). In particular, the conjectures above are implied by the corresponding conjectures for A373888. - _Robert Israel_, Aug 19 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Aug 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016687 | OEIS-A373888

a(n) is the length of the longest arithmetic progression of primes ending with prime(n).

Conjectures: a(n) >= 3 for n >= 13.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Aug 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016688 | OEIS-A373998

a(1) = 1, a(2) = 2, a(3) = 3, a(4) = 5; for n > 4, a(n) is the smallest unused positive number that is coprime to a(n-1) and a(n-2) but has a common factor with at least one of a(1)...a(n-3).

The fixed points are 1, 2, 3, 8, 45, 51, and it is likely no more exist. For the terms studied the primes appear in their natural order. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 24 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016689 | OEIS-A373999

a(1) = 1, a(2) = 2, a(3) = 3, a(4) = 5, a(5) = 7; for n > 5, a(n) is the smallest unused positive number that is coprime to a(n-1), a(n-2) and a(n-3) but has a common factor with at least one of a(1)...a(n-4).

Other than the first three terms the fixed points in the first 100000 terms are 35, 63, 219, 231, 1407, 2967, 3003, 6555, 14007, 14031, 32103, 77343, although it is likely more exist. For the terms studied the primes appear in their natural order. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 24 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016690 | OEIS-A374057

Integers k such that all k - p are primitive practical numbers where p is a primitive practical number in range k/2 <= p < k.

This sequence is analogous to A320447. Instead of the sequence of primes it uses the sequence of primitive practical numbers (A267124). It is conjectured that the sequence is finite and full.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frank M Jackson_, Jun 26 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016691 | OEIS-A374067

a(n) is the permanent of the symmetric Toeplitz matrix of order n whose element (i,j) equals the |i-j|-th prime or 1 if i = j.

Conjecture: a(n) is the minimal permanent of an n X n symmetric Toeplitz matrix having 1 on the main diagonal and all the first n-1 primes off-diagonal. - _Stefano Spezia_, Jul 08 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jun 27 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016692 | OEIS-A374068

a(n) is the permanent of the symmetric Toeplitz matrix of order n whose element (i,j) equals the |i-j|-th prime or 0 if i = j.

Conjecture: a(n) is the minimal permanent of an n X n symmetric Toeplitz matrix having 0 on the main diagonal and all the first n-1 primes off-diagonal. - _Stefano Spezia_, Jul 06 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jun 27 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016693 | OEIS-A374090

a(n) is the smallest nonnegative integer k where exactly n ordered pairs of positive integers (x, y) exist such that x^2 + x*y + y^2 = k.

Conjecture: Let q_i be the i-th prime of the form 3*k + 1 and let m = Prod_{j=1, t} b_j, a factorization of m into factors > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Jun 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016694 | OEIS-A374105

Numbers k such that (9^k + 3^k + 1)/13 is prime.

Conjecturally all terms are prime. There are no composite terms < 24000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aurelien Gibier_, Jun 29 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016695 | OEIS-A374140

a(n) is the permanent of the symmetric Toeplitz matrix of order n whose element (i,j) equals abs(i-j) or 1 if i = j.

Conjecture: a(n) is the minimal permanent of an n X n symmetric Toeplitz matrix having 1 on the main diagonal and all the integers 1, 2, ..., n-1 off-diagonal. - _Stefano Spezia_, Jul 05 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jun 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016696 | OEIS-A374153

Number of domino tilings of the order n Aztec diamond which are diagonally symmetric.

Finding a closed-form formula for a(n) is an open problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ludovic Schwob_, Jun 29 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016697 | OEIS-A374166

The set of primes p such that, for all integers m, the prime p divides at least one term of the recursive sequence (b_0, b_1, b_2, ...) given by b_0 = m and b_{k+1} = b_k^2 - 1 for k >= 0.

It is an open problem (proposed by Michael Stoll) whether this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robin Visser_, Jun 29 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016698 | OEIS-A374167

a(n) is the largest k such that the binary representation of prime(n), when interpreted as a base-b number, yields a prime for every b in 2..k.

Conjecture: every integer > 1 eventually appears.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Jun 29 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016699 | OEIS-A374175

a(n) is the conjectured number of occurrences of n in A373330.

a(n) is the conjectured number of occurrences of n in A373330. The sequence must be considered as conjectural, since so far no proof is known for the non-occurrence of arbitrarily small terms > 0 for very large n in A373330, despite the growing distance reserve observed in A373331 and A373332.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jun 30 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016700 | OEIS-A374192

Number of iterations required to reach 1 or 10 in a modified Collatz trajectory where x -> x/2 if x is even, x -> x+1 if x is odd and not divisible by 3 and x -> 3x+1 if x is odd and divisible by 3; or a(n) = -1 if 1 or 10 is never reached.

It is conjectured that this iteration always reaches one of two cycles [1,2] or [10,5,6,3].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary Detlefs_, Jun 30 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016701 | OEIS-A374242

a(n) is the minimal absolute value of the determinant of a nonsingular n X n symmetric Toeplitz matrix having 1 on the main diagonal and all the integers 1, 2, ..., n-1 off-diagonal.

Conjecture: all the terms are odd.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jul 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016702 | OEIS-A374258

Square array: T(n,k) = ((3^(n+1) + 1)^(k-1) + 2)/3, read by descending antidiagonals.

Conjecture: For each natural number n, the digits of the product of any (n+1) not necessarily distinct terms of the n-th row in the base (3^(n+1)+1) numeral system appear in nondecreasing order. Proof of the conjecture. Let b := 3^(n+1)+1. The product of any (n+1) terms of the n-th row has the form p/(b-1), where p is the product of (n+1) numbers of the form b^k+2. Let p = (dm, ..., d1, d0)_b, and we have d0+d1+...+dm = 3^(n+1) = b-1. Then p/(b-1) = (dm, ..., d2+...+dm, 1+d1+...+dm)_b, which do form a nondecreasing sequence. - _Max Alekseyev_, Jul 03 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ahmad J. Masad_, Jul 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016703 | OEIS-A374264

a(n) is the smallest m such that A374167(m) = n, or -1 if no such m exists.

Conjecture: a(n) is never -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jon E. Schoenfield_, Jul 02 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016704 | OEIS-A374284

a(1) = 1, a(2) = 2; a(n) = least k != a(m), m < n, such that omega(a(n-1)) >= omega(k) requires rad(k) | a(n-1), where omega = A001221 and rad = A007947.

Conjecture: this sequence is a permutation of natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jul 31 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016705 | OEIS-A374334

a(n) is the numerator of x(n) = (16*x(n-1) + (120*n^2 - 89*n + 16)/(512*n^4 - 1024*n^3 + 712*n^2 - 206*n + 21)) mod 1, with x(0) = 0.

The present sequence gives the numerators of the associated sequence (with b = 2) for alpha = Pi. See Bailey and Borwein (2005), p. 505 (second example of Theorem 3). In the same paper, on p. 513, they conjecture that, for n >= 1, y(n) = floor(16*x(n)) = A062964(n+1). See also Bailey and Crandall (2001), p. 176.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Jul 06 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016706 | OEIS-A374337

Start with two vertices and draw a circle around each whose radius is the distance between the vertices. The sequence gives the number of regions constructed after n iterations of drawing circles with this same radius around every new vertex created from all circles' intersections.

Conjectured:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jul 05 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016707 | OEIS-A374338

Start with two vertices and draw a circle around each whose radius is the distance between the vertices. The sequence gives the number of vertices constructed after n iterations of drawing circles with this same radius around every new vertex created from all circles' intersections. See the Comments.

Conjectured:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jul 05 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016708 | OEIS-A374351

Lexicographically earliest infinite sequence of distinct positive integers such that for any triple i,j,k of consecutive terms, gcd(i,k) = 1 and A007947(i*j*k) is a term in A002110.

Conjectures: A permutation of the positive integers, a(n) = prime p (> 2) iff p is least unused odd term, 2|i, and rad(i*j) is a primorial number divisible by all primes < p but not by p; primes appear in natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jul 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016709 | OEIS-A374365

Numbers m such that abs(Sum_{k=1..m} [k|m]*A008683(k)*(-1)^(2*k/5)) = 0.

Conjecture: Numbers having a prime factor congruent to 1 mod 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Jul 06 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016710 | OEIS-A374366

a(n) = Im(Sum_{d|n} mu(d)*(i^d)).

Conjecture 1: Numbers n such that a(n) = 0 is A009003. Conjecture 2: Numbers n such that a(n) = 1 is A000079. This means that Conjectures 1 and 2 above are true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Jul 06 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016711 | OEIS-A374379

a(1) = 1, a(2) = 2. Let i = a(n-2), j = a(n-1), then for n > 2 a(n) is the least novel k such that rad(k) !| rad(i*j) whereas rad(i*j*k) is a primorial number (A002110), where rad is A007947.

Conjectured to be a permutation of the natural numbers with primes in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jul 06 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016712 | OEIS-A374385

Irregular triangle read by rows: coefficients of polynomial P(n,x) (exponents in increasing order).

Conjecture 1: Sum_{j=1..(-1)^A001221(n)*A023900(n)} root(P(n,x) = 0, j)^k = (-1)^A001221(n)*A023900(GCD(n, k)), which means there is a loop: Conjecture 2: P(n,1) = A020500(n), for n >= 1. Conjecture 3: P(A005117(j),x) = Phi(A005117(j),x), j >= 2, where Phi(n,x) are the cyclotomic polynomials in A013595. The sum from the Formula section is equal to F_n(x) = -(-1)^omega(n)*f'(x)/f(x) where f(x) is the rad(n)-th cyclotomic polynomial (which implies f(x^A003557(n)) = Phi_n(x)), for n > 1. Conjectures 2 and 3 follow immediately. For p prime, F_{n*p}(x) = F_n(x) - [p does not divide n] * p * F_n(x^p). - _Andrei Zabolotskii_, Jul 15 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Jul 07 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016713 | OEIS-A374404

a(1) = 1, a(2) = 2. Let i = a(n-2), j = a(n-1). For n > 2 if rad(i*j) is primorial number A002110(t), a(n) is least novel k such that rad(i*j*k) = A002110(t+1). Otherwise, if rad(i*j) is not primorial, a(n) is least novel k such that rad(i*j*k) = A002110(s), where prime(s) = A006530(i*j); rad = A007947.

Conjectured to be a permutation of the positive integers A000027, with primes in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_ and _Michael De Vlieger_, Jul 07 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016714 | OEIS-A374412

Irregular triangle read by rows: Numerators of exponents of j-th root of the polynomial P(n,x) in A374385, and 1 if n is a power of 2, (numerators of exponents of roots in increasing order).

Conjecture 1: The j-th root of the n-th polynomial is: Root(P(n,x) = 0, j) = -(-1)^(j + n)*(-1)^(j/A204455(n))*[GCD(A204455(n),j) = 1], where 1 <= j <= A204455(n) and where terms equal to 0 are deleted. Conjecture 1 has been verified up to n = 200. Conjecture 1: a(n,j) = j*[GCD(A204455(n), j) = 1], 1 <= j <= A204455(n), where zeros are deleted. Verified up to n = 200.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Jul 08 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016715 | OEIS-A374416

a(n) is the number of ways n can be written as a sum of a practical number and two nonagonal numbers.

Conjecture (checked up to 10^8): a(n) = 0 if and only if n = 23.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Duc Van Khanh Tran_, Jul 08 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016716 | OEIS-A374445

Lexicographically earliest sequence of distinct positive integers such that any pair of consecutive terms are coprime whereas the squarefree kernel of their product is primorial.

{a(n); n >= 2} is conjectured to be a permutation of A047228.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jul 08 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016717 | OEIS-A374446

Positions of zeros in the expansion of Product_{k>=0} (1 - x^(2^k))^3; A373308(a(n)) = 0 for n >= 1.

Conjecture: a(n) = A001196(n) - 1 for n >= 1, where A001196 lists numbers with only even length runs in their binary expansion.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 08 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016718 | OEIS-A374464

Numbers k such that the odd parts of k and sigma(A003961(k)) are equal, where A003961 is fully multiplicative function with a(prime(i)) = prime(i+1), and sigma is the sum of divisors function.

Conjecture: the sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jul 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016719 | OEIS-A374486

Numbers k such that Taxicab(2,j,k) exists for large j.

Conjecture: Infinitely many positive integers are in this sequence, and infinitely many positive integers are not in this sequence. Conjecture: This sequence grows exponentially. Computationally it appears to have asymptotic a(n) = 1.03691*exp(0.594473*n^(1/2)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Oliver Lippard_ and _Brennan G. Benfield_, Aug 04 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016720 | OEIS-A374505

Maximum number of unit squares aligned with unit-spaced horizontal lines that can be enclosed by a circle of diameter n.

It is conjectured that this construction gives the maximal number of axis-parallel unit squares that can be packed into a circle of diameter n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Dewan_, Jul 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016721 | OEIS-A374525

T(n,k) is the number of distinct n X n {0,1}-matrices that reach a fixed point after k alternately applied sorts by rows and columns, where T(n,k), k>=0 is an irregular triangle read by rows.

It is conjectured that for n>=3 the last term > 0 in row n is T(n,2*n-3). This is consistent with the result of random draws, where T(7,11) is the last term in row 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_ at the suggestion of _Markus Sigg_, Jul 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016722 | OEIS-A374527

Odd numbers whose Collatz trajectory is a Sidon sequence.

Conjecture: because A375006 contains all powers of two, this sequence contains all numbers of form (4^k - 1)/3 (A002450) except 5. - _Ralf Stephan_, Mar 05 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Markus Sigg_ and _Hugo Pfoertner_, Jul 27 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016723 | OEIS-A374528

Conjectured number of unit squares setting a new density record when packed into the smallest possible enclosing circle.

Conjectured number of unit squares setting a new density record when packed into the smallest possible enclosing circle. All data after 7 have to be considered as conjectural. The sequence is based on the configurations shown on Erich Friedman's "Squares in Circles" webpage, see link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Jul 16 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016724 | OEIS-A374568

Expansion of g.f. A(x) satisfying x = A(x - x^2) - A(x + x^2)^2.

Conjecture: a(n) is odd iff n = 4^k for k >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 14 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016725 | OEIS-A374570

Expansion of g.f. A(x) satisfying A(x)^2 = A(A(x)*C(x)), where C(x) = x + C(x)^2 is the Catalan function (A000108).

Conjecture: for n > 1, a(n) is odd iff n = 2^k + 1 for k >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016726 | OEIS-A374571

Expansion of g.f. A(x) satisfying A(x) = A(x^2) - x*A(x^2)^2.

Conjecture: for n > 0, a(n) is odd iff n = A003714(k) for some k > 0, where A003714 lists Fibbinary numbers whose binary representation contains no two adjacent 1's. Conjectures. For n > 0, we have the following occurrences:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016727 | OEIS-A374575

Lexicographically earliest infinite sequence of distinct positive integers having the property that for any triple i,j,k of consecutive terms gcd(j,k) = 1 whereas A007947(i*j*k) is a term in A002110.

Conjectured to be a permutation of the positive integers with primes in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jul 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016728 | OEIS-A374594

Areas of trapezoids with integer sides, height and offsets whose area equals their perimeter.

Conjecture: in this sequence are only four terms which belong to trapezoids with exactly one pair of parallel sides: a(2) = 18, a(4) = 20, a(6) = 24, a(7) = 30.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Jul 13 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016729 | OEIS-A374602

Array of successive integer solutions to sqrt((d-c)*b^2 + c*(b+1)^2) for nonsquare integers d >= 2 (d=A000037(n) for n >= 1), where b and c are positive integers and c < d, read by antidiagonals.

Each row n divides into equal length, geometrically periodic subsequences, each with its own subsequence period length (A377290) and geometric growth factor (A377291); it is conjectured that this is the case for all n, and that all solutions conform as such and that there are no solutions that do not, but these are not proven.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles L. Hohn_, Jul 13 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016730 | OEIS-A374603

Numerators of sorted rationals r(n) of the form k/d, where d=(i+1)^m, 1 <= m < bigomega(k), bigomega(k) == 0 (mod i), bigomega(d) == 0 (mod i) and gcd(k, prime(j)) = 1 for all j <= i.

Conjecture: It appears that Pi*r(n) + sqrt(n) ~~ A002410(n), where '~~' means 'close to'. The relative error is less than +-0.001 for n ranging from 100000 to 400000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Friedjof Tellkamp_, Jul 13 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016731 | OEIS-A374605

a(n) = Sum_{k = 0..n} binomial(n, k)^2*binomial(n+k, k)*binomial(3*n+2*k, n).

It is easy to see that for odd prime p, binomial(2*n, n)^3 is divisible by p^3 for integer n in the interval [(p + 1)/2, p - 1]. A similar property appears to hold for the present sequence. We conjecture that for prime p >= 5, a(n) is divisible by p^3 for integer n in the interval [ceiling((2*p + 1)/3), p - 1] (checked up to p = 101).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jul 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016732 | OEIS-A374612

a(n) = n for n <= 3; for n > 3, a(n) is the smallest unused positive number that is coprime to a(n-1) but has a common factor with any other previous term that is also coprime to a(n-1).

The terms appear to follow a pattern similar to the A064413 and A373390, i.e., the terms are concentrated along just three lines of different gradient, and the lower line consists only of primes. In the first 10000 terms the primes appear in their natural order, and the fixed points are 1, 2, 3, 4, 28, 98; it is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jul 14 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016733 | OEIS-A374613

Prime numbers p with the property that neither of 2*(p+next prime to p)+/-1 is prime.

Conjecture: this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marcin Barylski_, Jul 14 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016734 | OEIS-A374621

Expansion of e.g.f. 1 - x^4/24 - log(1 - x).

Conjectures (confirmed up to n = 10): (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jul 14 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016735 | OEIS-A374641

Decimal expansion of log(9/10), negated.

Bailey and Crandall (2001), p. 185, present this constant as an example of an irrational number that, provided their "Hypothesis A" (p. 176) is true, is normal to base 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Jul 15 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016736 | OEIS-A374645

a(1) = 1, a(2) = 6, a(3) = 2; for n > 3, a(n) is the smallest unused positive number that is coprime to a(n-1) but has a common factor with any other previous term that has a common factor with a(n-1).

The terms appear to follow a pattern similar to the A064413 and A373390, i.e., the terms are concentrated along just three lines of different gradient, and the lower line consists only of primes. In the first 10000 terms the primes appear in their natural order except for a(37) = 19 and a(39) = 17 which are reversed. Many fixed points exist, 2228 in the first 10000 terms, these beginning 1, 12, 16, 18, 24, 32, 40, 48, ... The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jul 15 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016737 | OEIS-A374670

Generalized Collatz sequences: coefficients resulting in a cycle containing 1.

All terms are as yet only conjectures. Link includes github of code implementing CPU and GPU tests of these coefficients. Tests include reductions of integers up to 10^7 and random integers of size 2^2048.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John Laky_, Jul 14 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016738 | OEIS-A374717

For n a power of 2, a(n) = n. Otherwise let k = n - 2^j (> 0) where 2^j is the greatest power of 2 not exceeding n, then a(n) = least novel m*a(k); m a term in A033844.

Definition developed from the Doudna recursion A005940, see Example. Conjectured to be a permutation of A318400, numbers whose prime factorization consists only of primes with indices 2^k (terms in A033844).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jul 17 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016739 | OEIS-A374851

Obverse convolution A000040**A000040; see Comments.

See A374848 for the definition of obverse convolution and a guide to related sequences. a(2k+1) is a square for k>=0. Conjecture: every positive integer divides a(n) for infinitely many n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 05 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016740 | OEIS-A374853

Obverse convolution (n)**(Prime(n)); see Comments.

See A374848 for the definition of obverse convolution and a guide to related sequences. Conjecture: every positive integer divides a(n) for infinitely many n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 05 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016741 | OEIS-A374883

Decimal expansion of phi*(2*phi + 1) (i.e., (7 + 3*sqrt(5))/2), where phi is the golden ratio.

The author conjectures that this is the minimum volume of an axis-aligned bounding box which includes the shortest minimum-link circuit joining all the vertices of the cube {0,1}^3 (i.e., the closed polygonal chains consisting of exactly 6 edges visiting all the points of the set {(0,0,0),(0,0,1),(0,1,0),(0,1,1),(1,0,0),(1,0,1),(1,1,0),(1,1,1)}).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Jul 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016742 | OEIS-A374891

Obverse convolution (1)**A000984; see Comments.

a(n+1) = a(n)*A244174(n+1) for n>=0 (conjectured) = a(n)*A323230(n+2) for n>=0 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jul 31 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016743 | OEIS-A374908

Each term is the sum of the preceding term and its seven-segment total A006942.

Conjecture: Taking the least significant digit of each term is not an eventually periodic sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David J. Ellis_, Jul 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016744 | OEIS-A374911

a(n) = a(2^n mod n) + a(3^n mod n), with a(0) = 1.

Conjectured to contain all positive integers. Here are the indexes where each of the first few positive integers appear:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bryle Morga_, Jul 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016745 | OEIS-A374913

Numbers k such that k^(k + 1) == k + 1 (mod 2*k + 1).

Conjecture (Superseeker): a(n) = A263458(n)/2. - _R. J. Mathar_, Aug 02 2024 The conjectured formula is false. There exist numbers k such that 2*k + 1 is composite and k^(k + 1) == k + 1 (mod 2*k + 1). For example, when k = 1023: 1023^1024 == 1024 (mod 2047) and 2047 = 23*89 is composite. - _Jedrzej Miarecki_, Jan 16 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jul 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016746 | OEIS-A374916

Lexicographically earliest sequence of distinct positive integers in which any three consecutive terms are pairwise coprime whereas the squarefree kernel of their product is a primorial number (A002110).

6 cannot be a term in this sequence since if so a(i) = 6 would require Lpf(j) >= 5; contradiction. By similar arguments 10,15 and 30 are also not terms. Consequently no multiples of 6, 10, 15 or 30 are terms. 2,3,5 are the only prime terms and 1,2 are the only primorial terms. All terms are numbers divisible by one and only one of {2,3,5}, and the sequence is conjectured to be a permutation of such numbers, namely those congruent to (+ -){2,3,4,5,8,9,14}

(mod 30). Conjecture: As n approaches infinity all primorial numbers will eventually appear as rad(i*j*k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_ and _Michael De Vlieger_, Jul 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016747 | OEIS-A375006

Numbers whose Collatz trajectory is a Sidon sequence.

Hugo Pfoertner conjectured that the double of each odd term is a term, too. If this is true, then A375006 is the union of A374527, twice A374527, and A000079. Level-wise traversal of the corresponding subtree of the Collatz tree allows one to quickly generate many (unsorted) terms of the sequence. Up to at least level 1000 there is no counterexample to the conjectured occurrence of twice of all terms of A374527 in the present sequence. The results also suggest that A374527 is not finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Markus Sigg_, Jul 27 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016748 | OEIS-A375007

Numbers t which satisfy the equation: t mod k = floor((t - k)/k) mod k (1 <= k <= t) only for k = 1 and t.

Conjecture: a(n) + 1 is prime for n > 6. Verified for all terms < 10^8. Conjecture: nextprime(u(n)) - u(n), where u(n) = Product_{m=1..n} (a(m+1) - a(m)), is a noncomposite number. Verified for all terms < 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lechoslaw Ratajczak_, Jul 27 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016749 | OEIS-A375098

Diagonals of a Euclidean solid such that there exists a Pythagorean quadruple d^2=a^2+b^2+c^2 that is more cube-like than any prior value of d.

We conjecture this method provides complete solutions. This was confirmed directly by brute-force testing up to 1e7 (optimized using the sum-of-two-squares theorem for a given d^2-a^2 = b^2 + c^2; see link below).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian N. K. Anderson_, Mark K. Transtrum, and David D. Allred, Jul 29 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016750 | OEIS-A375147

a(1) = 0; for n >= 2, a(n) is the number of iterations needed for the map: x -> x / A000005(x) if x is divisible by A000005(x), x -> x + 1 otherwise, to reach 1.

The trajectory length is a repeated sum of steps up to the next refactorable number (A360778) and its refactoring "depth" (A374540). The sequence always reach 1 as soon as an iterate reaches the value x from A330816. Assuming A330816 to be finite (conjectured by David A. Corneth) and A360806 to be infinite, may there be a set of numbers n > 10^42, which is not reaching 1 ?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Aug 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016751 | OEIS-A375178

a(n) = Sum_{k = 0..n-1} binomial(n+k-1, k)^3 (same as A112028 with an extra 0 at the start).

For prime p, binomial(p+k-1, k) == 0 (mod p) for 1 <= k <= p-1. It follows that a(p) == 1 (mod p^3) for all primes p. We conjecture that, in fact, the stronger congruence a(p) == 1 (mod p^5) holds for all primes p >= 7. Further, we conjecture that for r >= 2 and prime p >= 5, a(p^r) == a(p^(r-1)) (mod p^(3*r+3)). More generally, for a positive integer m, define a sequence {b_m(n) : n >= 0} by setting b_m(n) = Sum_{k = 0..n-1} binomial(n+k-1, k)^(2*m+1). Then the congruence b_m(p) == 1 (mod p^(2*m+1)) clearly holds for all primes p. We conjecture that the stronger supercongruence b_m(p) == 1 (mod p^(2*m+3)) holds for all primes p >= 2*m + 5, and for r >= 2, the supercongruence b_m(p^r) == b_m(p^(r-1)) (mod p^(3*r+2*m+1)) also holds for all primes p >= 2*m + 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Aug 03 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016752 | OEIS-A375179

a(n) = Sum_{k = 0..n-1} (-1)^(n+k+1)*binomial(2*n, k)^3.

We conjecture that for the present sequence the stronger supercongruence a(p) == 1 (mod p^5) holds for all primes p >= 7. Further, we conjecture that for r >= 2 and prime p >= 5, a(p^r) == a(p^(r-1)) (mod p^(3*r+3)). More generally, for a positive integer m, define a sequence {b_m(n) : n >= 0} by setting b_m(n) = Sum_{k = 0..n-1} (-1)^(n+k+1)*binomial(2*n, k)^(2*m+1). We conjecture that the supercongruence b_m(p) == 1 (mod p^(2*m+3)) holds for all primes p >= 2*m + 5, and for r >= 2, the supercongruence b_m(p^r) == b_m(p^(r-1)) (mod p^(3*r+2*m+1)) also holds for all primes p >= 2*m + 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Aug 04 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016753 | OEIS-A375180

a(n) = Sum_{k = 0..n-1} (-1)^(n+k+1)*binomial(3*n, k)^3.

We conjecture that for the present sequence the stronger supercongruence a(p) == 1 (mod p^5) holds for all primes p >= 7. Further, we conjecture that for r >= 2 and prime p >= 5, a(p^r) == a(p^(r-1)) (mod p^(3*r+3)). More generally, for a positive integer m, define a sequence {b_m(n) : n >= 0} by setting b_m(n) = Sum_{k = 0..n-1} (-1)^(n+k+1)*binomial(3*n, k)^(2*m+1). We conjecture that the supercongruence b_m(p) == 1 (mod p^(2*m+3)) holds for all primes p >= 2*m + 5, and for r >= 2, the supercongruence b_m(p^r) == b_m(p^(r-1)) (mod p^(3*r+2*m+1)) also holds for all primes p >= 2*m + 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Aug 05 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016754 | OEIS-A375204

Record values in A375202.

Conjectures: All powers of 2 (A000079), 3*(powers of 2) (A007283) and 5*(powers of 2) (A020714) are terms. All terms are 5-smooth (A051037). - _Chai Wah Wu_, Oct 16 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Oct 15 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016755 | OEIS-A375215

Primes of the form p^16 + 2^16, where p is prime (see also A157764).

It is conjectured that solutions for p1^n + p2^n = p3 (where p1, p2, and p3 are all primes and n is a natural number) exist only when n is itself a power of two (when n is a number in A000079); and would have infinitely many solutions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mykhailo Papenko_, Oct 17 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016756 | OEIS-A375231

Interleaving A006040 and A228229.

Conjecture: a(n) is the permanent of the n X n matrix whose generic element is given by M_{i,j} = 1 if i = j or i + j = 1 (mod 2), with i,j in [n].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Aug 06 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016757 | OEIS-A375262

Least positive integer m such that sigma(m)/phi(m) = n + 1/2, where sigma(.) and phi(.) are given by A000203 and A000010, respectively.

Conjecture: Any rational number r >= 1 can be written as sigma(m)/phi(m) with m a positive integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 08 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016758 | OEIS-A375281

Lexicographically earliest infinite sequence of distinct positive integers such that for all n > 1 a(n-1), a(n) do not have the same number of distinct prime divisors, whereas the squarefree kernel of their product is a primorial number.

In other words omega(a(n-1)) != omega(a(n)) whilst rad(a(n-1)*a(n)) is a term in A002110, where omega is A001221 and rad is A007947. In general, primes appear consequent to primorial terms (a(11,12) = 18,5 is an exception). Conjectured to be a permutation of the positive integers, with primes in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Aug 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016759 | OEIS-A375286

a(n) = f(1) + f(2) + ... + f(n), where f(n) = (-2)^Omega(n) = A165872(n).

Johnston, Leong, & Tudzi prove that |a(n)| < 2260n. Sun conjectures that |a(n)| < n for n >= 3078. Mossinghoff & Trudgian verify this to 2.5 * 10^14.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Aug 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016760 | OEIS-A375345

a(n) = b(n, n-1) where b(n, m) = b(n, m-1) + gcd(b(n, m-1), n-m) for m > 0 with b(n, 0) = n.

It is conjectured that all terms are prime. Conjecture was checked up to 10^6. - _Vaclav Kotesovec_, Sep 10 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Aug 12 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016761 | OEIS-A375348

a(n) is the mode of the digits of n! not counting trailing zeros (using -1 if multimodal).

Conjecture: excluding -1, as n -> oo, all digits occur equally often.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_ and _Robert G. Wilson v_, Aug 12 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016762 | OEIS-A375351

Least squarefree integer m > 1 such that the product of all those (p + 4)/(p + 2) (with p a prime divisor of m) equals (2*n + 1)/(2*n - 1).

Conjecture: a(n) exists for any positive integer n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 14 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016763 | OEIS-A375417

a(1)=1; thereafter if n is odd we require that a(n) || a(n+1) be abundant and otherwise deficient. At each step, choose the smallest possible integer not yet in the sequence.

Conjecture: this is a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo P. Lava_, Aug 14 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016764 | OEIS-A375448

Expansion of the g.f. A(x) with the property that the sum of the first n coefficients in A(x/n)^n equals n^2 for n >= 1.

a(n) ~ c * n^(n+1), where c = 0.5698891168602..., conjecture: c = (exp(1)-1)/exp(3*exp(-1)). - _Vaclav Kotesovec_, Sep 13 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 12 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016765 | OEIS-A375449

Expansion of g.f. A(x) satisfying [x^n] x*A'(x) / (1 - n*A(x)) = n^(n+1) for n >= 1.

Conjecture: for n > 1, a(n) is odd iff n is of the form 2^k-2 or 2^k-1 for some k > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 12 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016766 | OEIS-A375452

Expansion of g.f. A(x) satisfying [x^n] x*A'(x) / (1 - n*A(x)) = n^n for n >= 1.

Conjecture: for n > 1, a(n) is odd iff n is of the form 2^k-2 or 2^k-1 for some k > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016767 | OEIS-A375457

Expansion of the g.f. A(x) with the property that the sum of the first n coefficients in A(x/n)^n equals n for n >= 1.

a(n) ~ c * n^n, where c = 1.189395759976..., conjecture: c = (exp(1)-1)/exp(exp(-1)). - _Vaclav Kotesovec_, Sep 13 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 08 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016768 | OEIS-A375472

Least k such that the ternary representation of 2^k has exactly 2*n 1's, or -1 if no such k exists.

Conjecture: a(n) ~ 6*log_2(3)*n = 6*A020857*n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pontus von Brömssen_, Aug 17 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016769 | OEIS-A375512

a(n) is the number of distinct compositions of four positive integers x, u, v, y (x < u <= v < y) for which x + u + v + y = n and u*v = x*y.

Conjecture: a(p) = 0 for primes p. The conjecture is true, in fact for any x,y,u,v as in the definition, n has proper divisor gcd(x,u) + gcd(v,y).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Aug 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016770 | OEIS-A375516

a(n) is the denominator of Sum_{k = 1..n} 1 / (k*A374663(k)).

One might be led to conjecture that the last 4 digits of the numbers from a(5) onwards are always 1200, but _Rémy Sigrist_ has observed that this does not hold for a(10) = 235541975237750435699516318092729420457559440943208103525303043995293765200.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016771 | OEIS-A375522

a(n) is the denominator of Sum_{k = 1..n} 1 / (k*A375781(k)).

suggesting the conjecture that they are always 1 except for n = 4 and 6 (compare the Theorem in A374983).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_ and _N. J. A. Sloane_, Aug 30 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016772 | OEIS-A375552

Yet unseen terms in the enumeration of A375553, prepended by [2, 5].

Conjecture: This is a permutation of the prime numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Sep 17 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016773 | OEIS-A375563

a(1) = 1; for n > 1, a(n) is the smallest unused positive number that shares a factor with a(n-1) if a(n-1) is prime, otherwise a(n) is coprime to a(n-1).

The terms are all concentrated along three lines with the central line, consisting entirely of all the odd terms, along the line a(n) = n. Surprisingly however there are no fixed points in the first 100000 terms. In the same range the primes appear in their natural order. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Aug 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016774 | OEIS-A375564

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest unused positive number that is coprime to a(n-1) if a(n-1) is prime, otherwise a(n) shares a factor with a(n-1).

The fixed points are 1, 2, 3, 4, 15, 51, 63, 363, 437, ... There are thirty-two fixed points in the first 100000 terms and it is likely there are infinitely many in total. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Aug 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016775 | OEIS-A375567

Length of the "exponential comma sequence" with n as the initial term, or -1 if that sequence is infinite.

If the decimal expansion of n is 1...0, its exponential comma sequence is trivially constant and therefore infinite so that a(n) = -1. Conjecture: these are the only infinite exponential comma sequences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nicholas M. R. Frieler_, Aug 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016776 | OEIS-A375575

a(n) is the least frequent digit of n! not counting trailing zeros, or -1 if there is more than one least frequent digit.

Conjecture: excluding -1, as n -> oo, the digits distribution is uniform as in A375348.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_ and _Robert G. Wilson v_, Aug 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016777 | OEIS-A375580

a(n) is the number of partitions n = x + y + z of positive integers such that x*y*z is a perfect cube.

Conjecture: for n > 176, a(n) > 0. - _Charles R Greathouse IV_, Aug 20 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Aug 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016778 | OEIS-A375583

a(n) is the number of ways n can be written as a sum of a practical number and two 11-gonal numbers.

Conjecture (checked up to 10^8): a(n) = 0 if and only if n = 11.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Duc Van Khanh Tran_, Aug 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016779 | OEIS-A375729

Irregular triangular array read by rows. T(n,k) is the number of monic irreducible polynomials of degree n in F_2[x] that are k-normal, n>=1, k>=0 .

Conjecture: Let alpha be in F_q^n. Write alpha as a q-ary word w with respect to the standard polynomial basis (1,x,x^2,x^3,...,x^(n-1)). Let beta in F_q^n be the q-ary word w interpreted with respect to any normal basis. Then beta is a root of a k-normal polynomial iff the period of w = n and deg(gcd(alpha,x^n-1))=k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Critzer_, Aug 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016780 | OEIS-A375792

Numbers k such that 2^k == 2 (mod k-th triangular number) and not 2^k == 2 (mod k-th oblong number).

Conjecture: all terms of the sequence are prime numbers A000040. The conjecture is false: 45812984491 = 1777 * 25781083 is in the sequence. - _Charles R Greathouse IV_, Aug 29 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Aug 29 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016781 | OEIS-A375824

Triangular numbers whose sum of digits is 9.

Conjecture: the last term not of one of those subsequences is a(53) = 210010000005.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Aug 30 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016782 | OEIS-A375828

a(1) = 1; for n > 1, a(n) is the smallest unused positive number such that (a(n-1) AND a(n)) = a(n-1) if a(n-1) is prime, otherwise (a(n-1) AND a(n)) = 0, where AND is the binary AND operation.

The terms form a pattern similar to that in A109812. In the first 250000 terms the fixed points are 1, 2, 3, 8, 3650, 50624, 203074. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Aug 30 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016783 | OEIS-A375853

Triangle read by rows: T(n, k) = k*(n - k)*binomial(2*n+2, 2*k+1)/(4*n + 2) for 1 <= k <= n-1.

Sum_{k>=0} T(n, k) = A002699(n-1) (conjectured by Bourn and Erickson).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mingjian Ding_, Aug 31 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016784 | OEIS-A375889

a(1) = 1, a(2) = 3; for n > 2, a(n) is the smallest unused positive number such that a(n) shares a factor with a(n-1) and (a(n) AND a(n-1)) = min(a(n), a(n-1)), where AND is the binary AND operation.

The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016785 | OEIS-A375890

a(1) = 1; for n > 1, a(n) is the smallest unused positive number such that a(n) is coprime to a(n-1) and (a(n) AND a(n-1)) = min(a(n), a(n-1)), where AND is the binary AND operation.

The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016786 | OEIS-A375925

Squares visited by a king moving on a walled, spirally numbered board, where a wall must be jumped on each move, always to the lowest available unvisited square.

However, it would be more natural to start the path at the origin, at a square labeled n = 0 (to which the king never moves). Then the sequence would conjecturally be a permutation of the nonnegative integers. This also leads to a more natural numbering for the squares in terms of the x,y coordinates - compare the Python function "square_number()". See A383185. (End) [Comment edited by _N. J. A. Sloane_, May 14 2025 following discussion with _Kevin Ryde_.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sameer Khan_, Sep 03 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016787 | OEIS-A375948

Expansion of e.g.f. 1 / (3 - 2 * exp(x))^(3/2).

Conjecture: a(n) = b(n-1) + 2 * Sum_{j=0..n-1} binomial(n, j) * a(j) for n > 0 where b(n) = b(n-1) + Sum_{j=0..n-1} binomial(n-1,j) * a(j+1) for n > 0 with b(0) = 1. Also it seems that b(n) = A305404(n+1). - _Mikhail Kurkov_, Jun 14 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Sep 03 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016788 | OEIS-A375986

Maximum number of edges in a simple polygon that is the regular union of n triangles.

A union of triangles is said to be regular if their vertices along the common border are arranged cyclically. If the union is non-regular, the maximal number of edges i(n) satisfies i(n) > c*A002004(n). So, it grows superlinearly. It is an open question to find n for which i(n) > a(n). - _Giedrius Alkauskas_, Jan 09 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giedrius Alkauskas_, Sep 07 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016789 | OEIS-A376055

Difference between numerator and denominator of the sum S(n) defined in A376052.

[Only the first 20 terms - shown here - have been calculated. It is only a plausible conjecture that all subsequent terms are also 1.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 14 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016790 | OEIS-A376084

Number of cryptarithmically unique primes with n decimal digits.

It is conjectured that a(n) is asymptotic to A006879(n) as n->oo based on the combinatorial probability estimate under the assumption that asymptotically for large n, the fraction of primes among integers that share a given digit pattern would be the same as among all integers with n digits, given by p(n)=1/(n*ln10) according to the prime number theorem. Since the number of integers sharing the same digit pattern cannot exceed 9*9!, the probability for a randomly chosen prime of length n to be cryptarithmically unique >= (1-p(n))^(9*9!-1), which is asymptotic to 1 as n->oo. The following terms are conjectured based on the assumption that at these lengths A374238 does not contain terms with 4 or more distinct digits, which follows from the vanishing probability of such terms estimated with combinatorial arguments: Further conjectured terms: a(20)=125975, a(21)=180471, a(22)=852579. - _Michael S. Branicky_, Oct 16 2024 lim_{n->oo} a(n)/A006879(n)=1 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmytro Inosov_, Sep 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016791 | OEIS-A376114

Refactorable numbers that are twice a square.

Conjecture: a(n) = A181795(n)/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Waldemar Puszkarz_, Sep 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016792 | OEIS-A376118

Cryptarithmically unique palindromic primes.

Number of terms < 100^k: 1, 1, 1, 1, 1, 4, 16, 19, (92), (249), (416), (1093)... . The numbers in brackets are conjectured based on the calculated terms with 1, 2, or 3 distinct digits and the vanishing combinatorial probability of terms with 4 or more distinct digits at these lengths.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmytro Inosov_, Sep 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016793 | OEIS-A376182

Triangle T read by rows: T(n, k) = (2*n^2 + 4*n + 1 - (-1)^n) / 4 - (1 + (-1)^k) * (n - k) - k.

Conjecture: This triangle seen as a sequence yields a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Sep 14 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016794 | OEIS-A376201

Fixed points in A376198.

Conjectures: the terms consist of runs of an even number (>1) of successive numbers that increase by 1 at each step; A376758(n) is one-half of the length of the n-th such run; and the run ends at A376751(n) - 1. - _N. J. A. Sloane_, Oct 07 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 03 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016795 | OEIS-A376225

G.f. A(x) satisfies [x^(2*n)] A(x)^(3*n) = -3 and [x^(2*n+1)] A(x)^(3*n+1) = [x^(2*n+1)] A(x)^(3*n+2) for n >= 1.

The g.f. A(x) of this sequence satisfies similar conditions, with the surprisingly simple formula [x^(2*n+1)] A(x)^(3*n+1) = (3*n+1) * Product_{k=1..n} (3*k + 2) for n >= 1, a conjecture which has been verified for the initial 1001 terms. Compare with A377095, the dual of this sequence. (3) [x^(2*n+1)] A(x)^(3*n+1) = (3*n+1) * Product_{k=1..n} (3*k + 2) for n >= 1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 26 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016796 | OEIS-A376226

G.f. satisfies A(x) = A(x^3 + 6*x*A(x)^3) / A(x^2 + 4*x*A(x)^2).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 04 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016797 | OEIS-A376238

Sequence obtained from A376239 by deleting the substrings of k-1 copies of k starting at index k(k-1)/2 + 1 for each k > 1.

In A376239 we conjecture that the sequences S[r+1] = D S[r], starting with S[1] = A376239, will have the property P(r) for all r >= 1. Here D means to delete the subsequences of k repeated k-1 times, starting at index (k-1)r + T(k-2) + 1, for all k > 1. The property P(r) means that for all k > 1, the sequence has k-1 copies of k, starting at index (k-1)r + T(k-2) + 1, and each of these runs is preceded by r smaller terms. (Thus, r is the number of 1s preceding the first 2 in the sequence.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Oct 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016798 | OEIS-A376239

a(n) = index k of the column in which n is found, if the columns k = 1, 2, 3, ... are initially filled with numbers T(k-1)+1, ..., T(k) = k(k+1)/2, and when n is found in column k, then the top element of each subsequent column (k+1, k+2, ...) is moved to the top of the previous column (k, k+1, ...).

Conjecture: Starting with S[1] = A376239, if S[r+1] = D(r) S[r] for all r >= 1, then each sequence S[r] has the property P(r). Here we denote by D(r) the operation of deleting the substring of k-1 copies of k starting at index (k-1)r + T(k-2) + 1, for all k > 1, and keeping only the intermediate chunks of length r. By property P(r) we mean that for each k > 1, the sequence contains k-1 consecutive terms equal to k, starting at index (k-1)r + T(k-2) + 1, each such run being preceded by r terms < k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ali Sada_ and _M. F. Hasler_, Oct 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016799 | OEIS-A376256

Numbers which are the minimum of a cycle in the map x -> phi(sigma(x)).

The status of 254731536 is unknown, but conjectured not a term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard R. Forberg_, Sep 16 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016800 | OEIS-A376287

Index of first occurrence of n in A129363, or 0 if no such number exists.

Conjecture: a(n) > 0. Checked to 1010. Conjectured last occurrence: 4208, 24536, 28916, 21278, 51806, 68078, 73538, 89216, 83978, ..., . Conjecture number of terms for A129363(k) = n: 35, 115, 285, 327, 557, 537, 723, 652, 882, ..., .

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert G. Wilson v_, Sep 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016801 | OEIS-A376317

a(n) = P(n+1, n+1) where P(n, m) = P(n, m-1) + P(n-1, m + f(m-n)) for n < m with P(n, m) = P(n-1, m) for 0 <= m <= n and P(0, m) = 1 for m >= 0 and where f(n) = [(n mod 4) > 0].

Conjecture: cases f(n) = n mod 2 and f(n) = [(n mod 3) > 0] both gives A006318. Conjecture: a(n+2) = a(n) + A215973(n+2) - A215973(n+1) (noticed by advanced OEIS search).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Sep 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016802 | OEIS-A376336

Least positive integer m such that tau(m) is divisible by n, where the Ramanujan tau function is given by A000594.

Conjecture: a(n) exists for any positive integer n. Moreover, a(n) <= n^2 for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016803 | OEIS-A376354

Numbers that end in 0 when written in base of triangular numbers (cf. A000462).

Conjecture: {a(n+1) - a(n) : n >= 1} = {1, 2, 3}. (See related conjectures at A376355 and A376356.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016804 | OEIS-A376355

Numbers that end in 1 when written in base of triangular numbers (cf. A000462).

Conjecture: {a(n+1) - a(n) : n >= 1} = {2,3,4,5,6}. (See related conjectures at A376354 and A376356.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016805 | OEIS-A376356

Positions of numbers in A000462 that end in 2.

Conjecture: {a(n+1) - a(n) : n >= 1} = {3,4,5,6,7,8,9}. (See related conjectures at A376354 and A376355.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016806 | OEIS-A376357

Positions of numbers in A007961 that end in 0.

Conjecture: {a(n+1) - a(n) : n >= 1} = {1,2,3,4}. (See related conjectures at A376358-A376360.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016807 | OEIS-A376358

Positions of numbers in A007961 that end in 1.

Conjecture: {a(n+1) - a(n) : n >= 1} = {2,3,4,5,6,7}. (See related conjectures at A376357, A376359, and A376360.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016808 | OEIS-A376359

Positions of numbers in A007961 that end in 2.

Conjecture: {a(n+1) - a(n) : n >= 1} = {3,4,5,6,7,8,9,10}. (See related conjectures at A376357, A376358, and A376360.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016809 | OEIS-A376360

Positions of numbers in A007961 that end in 3.

Conjecture: {a(n+1) - a(n) : n >= 1} = {4,5,6,7,8,9,10,11,13}. It has been checked that a(n+1) - a(n) is not 12 for 1<=n<=300000. (See related conjectures at A376357, A376358, and A376359.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016810 | OEIS-A376388

a(n) = P(n+1, n+1) where P(n, m) = P(n, m-1) + P(n-1, m + f(m-n)) for n < m with P(n, m) = P(n-1, m) for 0 <= m <= n and P(0, m) = 1 for m >= 0 where f(n) = [(n mod 5) > 0].

Conjecture: cases f(n) = n mod 2 and f(n) = [(n mod 3) > 0] both gives A006318.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Sep 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016811 | OEIS-A376395

a(n) = P(n+1, n+1) where P(n, m) = P(n, m-1) + P(n-1, m + f(m-n)) for n < m with P(n, m) = P(n-1, m) for 0 <= m <= n and P(0, m) = 1 for m >= 0 and where f(n) = [(n mod 6) > 0].

Conjecture: cases f(n) = n mod 2 and f(n) = [(n mod 3) > 0] both gives A006318.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Sep 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016812 | OEIS-A376399

a(0) = 1, and for n > 0, a(n) is the least k such that A276075(k) = a(n-1) + A276075(a(n-1)), where A276075 is the factorial base log-function.

Conjecture: there are infinitely many variants b of this sequence, such that A276075(b(n)) = partial sums of b (shifted once right). One way to construct them: set i for some value >= 4, construct b first as here, but at point i, set b(i+1) = b(i) * A276076(b(i)), and after that, proceed as before, always finding a minimal k satisfying the condition. Unless b(i+1) = a(i+1), then b differs from this sequence but satisfies the same general condition, except that it is not the lexicographically earliest one. See also A376400.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 02 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016813 | OEIS-A376446

Modular phase shift of the tetration base n written by juxtaposing its representative congruence classes modulo 10 taken by starting from height 2 + v(n), and a(n) = -1 if n is a multiple of 10.

The author conjectures that if n is not a multiple of 10, then a(n) is necessarily an element of the subset {2, 4, 5, 6, 8, 9, 19, 28, 46, 64, 82, 91, 1397, 1793, 2486, 2684, 3179, 3971, 4268, 4862, 6248, 6842, 7139, 7931, 8426, 8624, 9317, 9713}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Sep 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016814 | OEIS-A376452

Positions of numbers in A376450 that end in 0.

Conjecture: if (b(n)) denotes the complement of this sequence, then {b(n+1)-b(n) : n>=1} = {2,3,4,5}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016815 | OEIS-A376453

Positions of numbers in A376450 that end in 1.

{a(n+1)-a(n) : n>=1} = {2,3,4,5} (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016816 | OEIS-A376458

a(n) = Sum_{k = 0..n} (-1)^(n+k)*binomial(n, k)*binomial(n+k, k)*A108625(n-1, n-k).

We conjecture that the present sequence satisfies same supercongruences and in addition satisfies the stronger congruences a(p) == a(1) (mod p^5) for all primes p >= 7 (checked up to p = 199) and a(p^r) == a(p^(r-1)) (mod p^(3*r+3)) for all primes p >= 5 and integers r >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016817 | OEIS-A376459

a(n) = Sum_{k = 0..n} (-1)^(n+k)*binomial(n, k)*binomial(n+k, k)*A108625(n, n-k).

We conjecture that the present sequence satisfies the same pair of supercongruences. Some examples are given below.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 24 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016818 | OEIS-A376461

a(n) = Sum_{k = 0..n} binomial(n, k)^2*binomial(n+k, k)*A108625(n, k).

We conjecture that the present sequence satisfies the same pair of supercongruences. Some examples are given below. Conjecture: Sequence of constant terms in powers of the Laurent polynomial (v2 + 1) * (v2 + v3 + 1) * (v0 + 1) * (v0 + v1 + 1) * (v0 + v1 + v2 + v3 + 1) / (v0 * v1 * v2 * v3). - _F. Chapoton_, Sep 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Sep 24 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016819 | OEIS-A376478

a(1) = 1, a(2) = 2, and a(n) = 3^(n-2) for n > 2.

Graham's conjecture: also numbers k such sigma(k) - k = floor(k/2). See Guy.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Sep 24 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016820 | OEIS-A376484

Array read by ascending antidiagonals: A(n,k)=4^k*Sum_{j=1..n} sin(2*j*Pi/(2*n+1))^(2*k).

It is only a conjecture that A(n,k) is always an integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cheng-Jun Li_, Sep 24 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016821 | OEIS-A376498

Array read by ascending antidiagonals: A(n, k) = 2^k*Sum_{j=1..n} cos((2*j - 1)*Pi/(2*n + 1))^k.

It is only a conjecture that the A(n, k) are always integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cheng-Jun Li_, Sep 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016822 | OEIS-A376499

Array read by ascending antidiagonals: A(n,k) = A376484/(2*n+1).

It is only a conjecture that the A(n,k) are always integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Cheng-Jun Li_, Sep 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016823 | OEIS-A376510

a(n) is the number of pairs of primes p+q=2*(n+4) with 5 <= p <= n such that either p+6 or q+6 is also prime.

If the above hypothesis is true, the Goldbach Hypothesis is true, since for every even number 2n, if there is a Goldbach decomposition p+q=2n meets the condition of this sequence, p+q+6=2n+6 forms at least one Goldbach decomposition of 2n+6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Sep 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016824 | OEIS-A376511

a(1) = 2. For n > 1, if a(n-1) is a novel term, a(n)=A001414(a(n-1)), else if there are k (>1) occurrences of a(j) = a(n-1) (1<=j<=n-1), a(n) is the smallest novel m such that A001414(m) = k*a(n-1).

Row n of table T(n,k) in A064364 lists numbers j such that A001414(j) = n, with T(n,1) = A056240(n), and every term in this sequence is taken from the appropriate row of A064364. When a(n-1) is a novel term a(n) = A001414(a(n-1)), which is defined. Otherwise a(n) = smallest m such that A001414(m) = k*a(n-1), a number which is also defined since it is the smallest unused term in T(k*a(n-1),k) of A064364. Therefore the sequence is well defined and infinite. Conjecture: For any n > 1 every term in T(n,k) of A064364 appears eventually.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Sep 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016825 | OEIS-A376527

a(n) = Sum_{k=0..n*(n-1)/2} A227543(n,k)^2 for n >= 0.

Conjecture: a(n) is odd iff n = 2^k - 1 for some k >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016826 | OEIS-A376548

a(1)=1,a(2)=2. Let p be the smallest prime factor of j=a(n-1), then for n>2 a(n) is the smallest novel multiple of Sopfr(j) if j is in A046363, or of p if it is not.

Conjectured to be a permutation of the positive integers A000027, in which the primes do not appear in order (prime order starts:2,5,7,3,13,11,19,17,31,23,43..).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Sep 27 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016827 | OEIS-A376562

Second differences of consecutive non-perfect-powers (A007916). First differences of A375706.

Mihailescu's theorem implies that the only case where two consecutive non-perfect powers have difference > 2 is A375706(5) = 3. It is conjectured that the only case where A375706 has two consecutive 2's is A375706(19) = A375706(20) = 2. If this is the case, the nonzero terms of this sequence for n > 20 all fit the pattern of a 1 followed immediately by a -1. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Oct 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016828 | OEIS-A376586

Irregular triangle T(n, k), n >= 0, k >= 0, read by rows with row polynomials P(n-1, n-1) for n > 0 with T(0, 0) = 1 where P(n, k) = (-1)^(n-k)*z^f(n-k+1)*P(n, k-1) + P(k-1, k-1) for 0 < k <= n with P(n, k) = 0 for k > n, P(n, 0) = 1 for n >= 0 and where f(2n+1) = n, f(2n) = 2n for n >= 0.

Conjecture: antidiagonal sums equal A000041. Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Sep 29 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016829 | OEIS-A376628

G.f.: Sum_{k>=0} x^(k*(k+1)) / Product_{j=1..k} (1 - x^(2*j-1)).

Conjectural g.f.: (1 + q * nu(-q))/(1 + q) = 1 + Sum_{k >= 0} q^(k+2)*Product_{j = 1..k} 1 + q^(2*j+1), where nu(q) is the g.f. of A053254. - _Peter Bala_, Jan 03 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Sep 30 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016830 | OEIS-A376687

Numbers that set records in A376281.

Conjecture: Numbers k that set records in A376281 do not have type C divisor pairs, i.e., those that are coreful but neither divides the other. This, since type C requires k to be powerful and divisible by cubes of 2 distinct primes (i.e., in A376936). Therefore the record is achieved only through large numbers of type A and B.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jan 08 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016831 | OEIS-A376758

The terms of A376201 consist of runs of successive numbers that increase by 1 at each step: a(n) is one-half of length of the n-th such run.

[At present it is only a conjecture that the runs have even length, but the proof should not be difficult.]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Oct 07 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016832 | OEIS-A376798

Least number of inequivalent cells in a polycube with n cells.

Conjecture: The sequence (a(n)-n/48) is eventually periodic with period 48.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John Mason_, Oct 04 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016833 | OEIS-A376839

a(1) = 1. For n > 1 if A007947(a(n-1)) is in A002110, a(n) is the smallest prime not already a term. Otherwise a(n) is the least novel multiple of the smallest non divisor prime of a(n-1).

A non divisor prime of a(n-1) is any prime p < Gpf(a(n-1)) which does not divide a(n-1). A007947(a(n-1)) is in A002110 iff a(n-1) is a term in A055932. Sequence is conjectured to be a permutation of the natural numbers (A000027) with primes in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Oct 06 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016834 | OEIS-A376842

Asymptotic phase shift of the tetration base n written by juxtaposing its representative congruence classes modulo 10 (i.e., the 1, 2, or 4, distinct sfasamenti taken by starting from the minimum height such that the congruence speed of n is constant), and a(n) = -1 if n is a multiple of 10.

The author conjectures that if n is not a multiple of 10, then a(n) is necessarily an element of the subset {2, 4, 5, 6, 8, 9, 19, 28, 46, 64, 82, 2486, 2684, 3971, 4268, 4862, 6248, 6842, 7931, 8426, 8624}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Oct 06 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016835 | OEIS-A376868

a(n) = n! * 4^binomial(n, 2).

Conjecturally, the sum of the permanents of all nXn 0-1 matrices.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, Oct 07 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016836 | OEIS-A376907

a(n) is the least n-digit cuban prime.

Conjecture: a(n+1)/a(n) ~ 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Oct 08 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016837 | OEIS-A376917

Starting from Goldbach decomposition of 10 = p + q = 5 + 5, 12 = 7 + 5, and 14 = 7 + 7, a(n) is the first number in A001057 such that if 2n - 6 = p + q, 2n = p' + q', where p' = p + 6 * a(n) and q' = 2n - q' are both primes.

Hypothesis: a(n) is defined for all n >= 8 and for all n >=8, the corresponding Goldbach decomposition 2n = p + q has positive primes p and q.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Oct 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016838 | OEIS-A376927

Totients whose inverses can be separated into ordered pairs (x, 2*x) with no remainder.

Conjecture: Numbers having an even number of totient inverses, the odd inverses being confined to the lower half of the list.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Torlach Rush_, Oct 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016839 | OEIS-A376942

Irregular table read by rows: row(n) is the lexicographically earliest sequence of positive integers a(n,1), a(n,2), ... a(n,k) such that Sum_{m = n..(n+k-1)} 1/(m*a(n,m-n+1)) <= 1.

Conjecture: all rows have finite length.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Oct 12 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016840 | OEIS-A376971

Number of polycubes of size n and symmetry class G (full symmetry).

Conjecture: For n >= 62, a(n) > a(n-1) if and only if n is a multiple of 6. - _Pontus von Brömssen_, Oct 20 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John Mason_, Oct 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016841 | OEIS-A376992

a(n) is the least n-digit prime of the form j^2 + (j+1)^2.

Conjecture: a(n+1)/a(n) ~ 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Oct 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016842 | OEIS-A377014

a(n) is the number of primes p such that p - 6, p + 6 and 2*n - p are also primes.

Conjecture: a(n) = 0 only when n = 1, 2, 3, 4, 5, 6, 19, 31, 331, 499.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Oct 12 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016843 | OEIS-A377027

a(1) = 1. For n > 1, if a(n-1) is a novel term, a(n) = a(a(n-1)), and if a(n-1) seen k (>1) times (including a(n-1)), with k*a(n-1) < n then a(n) = k*a(n-1); else a(n) = 1.

The clause requiring a(n) = 1 if k*a(n-1) >= n is necessary to ensure continuation of the sequence, since without it there would be occasions of a(m) = a(a(m)) where a(m) is not yet defined, see Example. This is conjectured to occur infinitely many times as the sequence extends, which then implies that every positive integer will eventually enter the sequence (by incrementation of 1 if not before). Consequently every integer is conjectured to occur infinitely many times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Oct 13 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016844 | OEIS-A377078

Lexicographically earliest infinite sequence of distinct positive integers such that, for n > 2, a(n) shares a factor with a(n-2) mod a(n-1).

To ensure the sequence is infinite a(n) must be chosen so that a(n-1) mod a(n) is not 0 or 1. In the first 100000 terms the fixed points are 119, 205, 287, and it is likely no more exist. It is conjectured all numbers > 1 appear in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Oct 15 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016845 | OEIS-A377090

a(0) = 0; thereafter, for n > 0, a(n) is the least integer (in absolute value) not yet in the sequence such that the absolute difference between a(n-1) and a(n) is a prime number; in case of a tie, preference is given to the positive value.

Conjecture (after studying A383445 and A383446): The sequence contains every integer. - _N. J. A. Sloane_, Apr 30 2025 ||a(n)| - n/2| = O(log(n)), probably ||a(n)| - n/2| < 2*log(n+2) for all n. (Conjectured; verified up to n = 10^5.) - _M. F. Hasler_, Feb 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Oct 16 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016846 | OEIS-A377091

a(0) = 0; thereafter a(n) is the least integer (in absolute value) not yet in the sequence such that the absolute difference between a(n-1) and a(n) is a square; in case of a tie, preference is given to the positive value.

Conjecture 1: Every integer (positive or negative) appears in this sequence. Conjecture 2: For n > 16, |a(n)| is within sqrt(n/2) of floor(n/2). See A379071. - _N. J. A. Sloane_, Dec 29 2024 [Corrected by _Paolo Xausa_, Jan 21 2025] Conjecture 3: lim sup ||a(n)| - floor(n/2)|/sqrt(n) = 1/2. (See link.) - _N. J. A. Sloane_ and _Paolo Xausa_, Feb 03 2025 Conjecture 4: After a(n) has been found, the sequence contains all numbers in the range [0,f(n)], where lim sup f(n) = (n-sqrt(n))/2. There is a corresponding conjecture for the negative terms. See A379067. - _N. J. A. Sloane_ and _Paolo Xausa_, Feb 13 2025 These conjectures are still open, although I conjecture that all four are true. This sequence could be called the Heraclitus transform of the squares. The OEIS now contains many other examples of Heraclitus transforms. _N. J. A. Sloane_, Feb 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Oct 16 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016847 | OEIS-A377092

a(0) = 0, and for any n > 0, a(n) is the least integer (in absolute value) not yet in the sequence such that the absolute difference of a(n-1) and a(n) is a Fibonacci number (A000045); in case of a tie, preference is given to the positive value.

That question is still open, although I conjecture that the answer is Yes. This sequence could be called the Heraclitus transform of the Fibonacci numbers. For other examples of Heraclitus transforms, see A377091, etc. - _N. J. A. Sloane_, Feb 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Oct 16 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016848 | OEIS-A377096

G.f. A(x) satisfies [x^(2*n)] A(x)^(5*n) = -5 and [x^(2*n+1)] A(x)^(5*n+2) = [x^(2*n+1)] A(x)^(5*n+3) for n >= 1.

The g.f. A(x) of this sequence satisfies similar conditions, with the surprisingly simple formula [x^(2*n+1)] A(x)^(5*n+2) = (5*n+2) * Product_{k=1..n} (5*k + 3) for n >= 1, a conjecture which has been verified for the initial 1001 terms. Compare with A377097, the dual of this sequence. (3) [x^(2*n+1)] A(x)^(5*n+2) = (5*n+2) * Product_{k=1..n} (5*k + 3) for n >= 1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 04 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016849 | OEIS-A377098

G.f. A(x) = Sum_{n>=0} a(n)*x^n where a(n) = Sum_{k=0..n-1} [x^k] A(x)^(k*(n-k)) for n >= 1 with a(0) = 1.

a(n) ~ c * n! * n^alpha / LambertW(1)^n, where alpha = 0.68670155428... and c = 0.09981115968806..., conjecture: alpha = 2*LambertW(1) - 3 + 4/(1 + LambertW(1)) = 0.6867015542800108601...- _Vaclav Kotesovec_, Nov 23 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 10 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016850 | OEIS-A377100

G.f. A(x) satisfies A(x) = A(x^3)/A(x^2) + A(x)^2.

Conjecture: a(n) == 1 (mod 3) for n >= 1, which has been verified for the initial 2305 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 24 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016851 | OEIS-A377106

G.f. A(x) satisfies A(x)^3 = A(x^3 + 9*x*A(x)^3), with A(0)=0, A'(0)=1.

Conjecture: a(n) is odd iff n = 2^k for some k >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 08 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016852 | OEIS-A377157

a(n) = n for n <= 3; for n > 3, a(n) is the smallest unused positive number that is coprime to a(n-1) if a(n-1) is prime and coprime to a(n-2) if a(n-2) is prime, else a(n) shares a factor with a(n-1) and a(n-2) if both are composite.

The fixed points begin 1, 2, 3, 6, 32, 57, 119, 343, 3123, 3859, 120481. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Oct 18 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016853 | OEIS-A377182

Lexicographically earliest infinite sequence of distinct positive integers such that, for n > 2, a(n) shares a factor with a(n-2) mod a(n-1) while a(n-1) mod a(n) has not previously occurred as the mod value for any consecutive pair of terms.

Surprisingly the first prime to occur is a(94122) = 47857. The next is a(103105) = 26591, and no other primes appear in the first 500000 terms. It is unknown if more occur or why it takes so many terms for a prime to appear. Many small primes, like 5, can never occur as all mod values less than the prime have already appeared. It is conjectured all missing numbers are prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Oct 18 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016854 | OEIS-A377193

Lexicographically earliest infinite sequence of distinct positive integers such that any term j = a(n-1) with primorial kernel is followed by a prime, whereas any other term is followed by a number with prime factors p < q = Gpf(j) which do not divide j.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_ and _Michael De Vlieger_, Oct 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016855 | OEIS-A377217

Expansion of the o.g.f. A(x) defined by [x^n] A(x)^(6*n) = (3*n)!/n!^3 for n >= 0.

Conjecture: for positive integers r and s, the sequence {v(r,s; n) : n >= 0} defined by v(r,s; n) = [x^(s*n)] A(x)^(r*n) also satisfies the same supercongruences.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Oct 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016856 | OEIS-A377224

Number of ways to write n as x*(5*x+1) + y*(5*y+1)/2 + z*(5*z+1)/2, where x,y,z are integers with y*(5*y+1) <= z*(5*z+1).

Conjecture 1: a(n) = 0 only for n = 1. Also, a(n) = 1 only for n = 0, 2, 3, 5, 7, 14, 16, 19, 37, 43, 58, 61, 79. Conjecture 2: Let N be the set of all nonnegative integers. Then Conjecture 3: We have

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 13 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016857 | OEIS-A377236

The mod value of the consecutive pairs of terms in A377182.

It is conjectured the sequence contains all numbers > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Oct 21 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016858 | OEIS-A377251

G.f. A(x) satisfies A(x) = 1/A(-x*A(x)) such that [x^(2*n+1)] A(x)^n = 0 for n >= 0, with A(0) = A'(0) = 1.

(2.d) [x^(2*n+2)] A(x)^n = -[x^(2*n+3)] A(x)^n for n >= 0 (conjecture), (2.e) [x^(2*n+2*k+1)] (1+x)^k * A(x)^n = 0 for n >= 0, k >= 0 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 24 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016859 | OEIS-A377278

Denominators in a harmonic triangle; q-analog of A126615, here q = 2.

Conjecture: Row sums of the harmonic triangle are A204243(n) / A005329(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Oct 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016860 | OEIS-A377285

Position of first 0 in the n-th differences of the strict partition numbers A000009, or 0 if 0 does not appear.

Open problem: Do the 9th differences of the strict integer partition numbers contain a zero? If so, we must have a(9) > 10^5. a(12) = 47. Conjecture: a(n) = 0 for n > 12. - _Chai Wah Wu_, Dec 15 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Dec 12 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016861 | OEIS-A377288

Numbers k such that there are exactly two prime-powers between prime(k)+1 and prime(k+1)-1.

Is this sequence finite? For this conjecture see A053706, A080101, A366833.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Oct 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016862 | OEIS-A377307

Minimum number of consecutive pieces that must be added to the pattern given by the binary representation of n to produce a winning position in Gordon Hamilton's Jumping Frogs game, or -1 if there is no such position.

Gordon Hamilton conjectures in his reference below that a(n) is positive for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Glen Whitney_, Oct 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016863 | OEIS-A377334

Number of n-celled polycubes with full symmetry and the rotation point of the symmetries in the center of a cell (that may or may not be part of the polycube).

Conjecture: For n >= 62, a(n) > a(n-1) if and only if n is a multiple of 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pontus von Brömssen_, Oct 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016864 | OEIS-A377335

Number of polycubes with 8*n cells, full symmetry, and the rotation point of the symmetries at the common corner of 8 cells (that may or may not be part of the polycube).

Conjecture: a(n) < a(n-1) if and only if n mod 3 = 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pontus von Brömssen_, Oct 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016865 | OEIS-A377466

Numbers k such that there is more than one perfect power x in the range prime(k) < x < prime(k+1).

The Redmond-Sun conjecture (see A308658) implies that this sequence is finite. - _Pontus von Brömssen_, Nov 05 2024 a(n) = A000720(A116086(n)) = A000720(A116455(n)) for n <= 10. This would hold for all n if there do not exist more than two perfect powers between any two consecutive primes, which is implied by the Redmond-Sun conjecture. - _Pontus von Brömssen_, Nov 05 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Nov 02 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016866 | OEIS-A377654

Numbers m^2 for which the center part (containing the diagonal) of its symmetric representation of sigma, SRS(m^2), has width 1 and area m.

Conjectures (true for all a(n) <= 10^8):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Nov 03 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016867 | OEIS-A377714

Cogrowth sequence for the 8-element group C4 X C2 = <S,T | S^4, T^2, [S,T]>.

Conjectures from _Chai Wah Wu_, May 16 2026: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sean A. Irvine_, Nov 04 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016868 | OEIS-A377727

Number of digit patterns of length n that satisfy no divisibility rules but do not generate primes.

It is conjectured that the next terms are a(14)=362, a(15)=363, a(16)=1448. This is based on the calculated number of primonumerophobic digit patterns with only 2 or 3 distinct digits and the vanishingly small combinatorial probability for the existence of additional primonumerophobic digit patterns of this length with 4 or more distinct digits.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmytro Inosov_, Nov 05 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016869 | OEIS-A377729

a(n) is the smallest number which can be represented as the sum of n distinct nonzero n-gonal numbers in exactly 2 ways.

Conjecture: a(n) = (n^4 - 2*n^3 + 38*n^2 - 85*n + 72)/6 for n >= 5. (End) Conjectured g.f.: x^3*(19 - 5*x - 98*x^2 + 199*x^3 - 171*x^4 + 72*x^5 - 12*x^6) / (1 - x)^5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ilya Gutkovskiy_, Nov 05 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016870 | OEIS-A377757

Number of possibilities to place "hat" monotiles onto the first n hexagons in a counterclockwise order such that more rings of tiles can be placed around it.

An open question is: What is the asymptotic behavior of a(n) when n tends to infinity?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ruediger Jehn_ and _Kester Habermann_, Nov 06 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016871 | OEIS-A377879

Deficiency of squares: a(n) = 2n^2 - sigma(n^2).

It is conjectured that 1's occur only when n is two's power (A000079), and that there are no -1's in this sequence. See comments in A033879 and in A337339.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016872 | OEIS-A377903

Indices of records in A302656.

Conjecture: a(n) = A377904(n-9) for n >= 11.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Nov 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016873 | OEIS-A377979

List of exponents in the expansion of (1 - q)*Sum_{n >= 0} q^(2*n*(n+1))*Product_{k >= 2*n+1} 1 - q^k.

Conjectures: The following are conjectural:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Dec 16 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016874 | OEIS-A377992

Irregular triangle giving on row n all antiderivatives of A024451(n), for n >= 2.

Because A024451 is a subsequence of A048103, this is also. And if all terms of A024451 are squarefree as is conjectured, then all terms of this sequence are cubefree (A004709).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016875 | OEIS-A378143

a(n) is the smallest prime of the form (2*p)^(2^n) + 1 for some prime p.

Conjecture: the last digit of each value of a(n), where n >= 1, is 7. The conjecture is equivalent to the claim that a(n) is not 10^(2^n) + 1 for any n, which in turn is equivalent to the claim that, if 10^(2^n) + 1 is prime, then either 4^(2^n) + 1 or 6^(2^n) + 1 is prime. - _Charles R Greathouse IV_, Nov 17 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Nov 17 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016876 | OEIS-A378145

Riordan triangle (1 + x * C(x), x * C(x)), where C(x) is g.f. of A000108.

Conjecture: Sum_{i=0..n-k} binomial(i+m-1, i) * T(n, i+k) = T(n+m, m+k) for m > 0. Conjecture: Sum_{k=0..n} (1 + floor(k/2)) * T(n, k) = A000108(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Nov 17 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016877 | OEIS-A378174

Maximum number of clues in an n X n crossword puzzle where each word is at least 4 letters long and the puzzle has 180-degree rotational symmetry and no interlock.

Except for n = 9, 10, 12, and 14, conjectured recursive formula is a(n) = a(n-5) + 4(n-4). In particular, conjectured explicit formula is a(n) = 2n floor((n+1)/5) if mod(n,5) = 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Abigail Schnitzer_, Nov 18 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016878 | OEIS-A378233

a(1) = 1, a(2) = 2, let i = a(n-2), j = a(n-1). For n > 2, If i*j is a term in A055932, a(n) is the least novel multiple of the smallest prime which does not divide i*j. Otherwise a(n) is the least novel multiple of the greatest prime < Gpf(i*j) which does not divide i*j.

Conjectures: Sequence is a permutation of the natural numbers A000027, with primes in order. Prime powers and numbers that share the same squarefree kernel also appear in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_ and _Michael De Vlieger_, Nov 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016879 | OEIS-A378258

G.f. satisfies A(x) = B(x) + 5*A(x)^2 where B(A(x) - A(x)^2) = x.

Conjecture: a(n) is odd iff n is a power of 2. Conjecture: a(n) is not congruent to 1 nor 5 (mod 6) for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 03 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016880 | OEIS-A378259

G.f. satisfies A(x) = Sum_{n>=1} A(x^(2*n))/A(x^n), with A(0) = 0 and A'(0) = 1.

Conjecture: a(n) == 2 (mod 4) iff n = 2^k + 1 for some k >= 0; elsewhere, a(n) is divisible by 4 for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 04 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016881 | OEIS-A378277

Denominators in a harmonic triangle, based on products of Fibonacci numbers.

Conjecture: Alt. row sums of the harmonic triangle are Fibonacci(n-2) / Fibonacci(n+1), where Fibonacci(-1) = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Nov 21 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016882 | OEIS-A378288

Array read by antidiagonals: row k consists of the positive integers j for which the concatenation of 2^k - 1 and 2^j - 1 is prime.

Conjecture: all rows have infinitely many terms, and all positive integers not divisible by 4 appear in infinitely many rows.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Nov 26 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016883 | OEIS-A378311

Number of cyclic edge cuts in the n-prism graph.

a(n) = A378312(n) + 1 (conjectured). The above conjecture is true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Nov 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016884 | OEIS-A378312

Number of cyclic edge cuts in the n-Moebius ladder graph.

a(n) = A378311(n) - 1 (conjectured). The above conjecture is true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Nov 22 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016885 | OEIS-A378419

Positive integers in A376842, sorted according to their appearance in that sequence.

In Appendix of "Graham's number stable digits: an exact solution" (see Links) the author conjectures that this list is complete, and thus the present sequences has exactly 21 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Nov 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016886 | OEIS-A378421

Positive integers in A376446 sorted according to their appearance in that sequence.

If we merge A376446(n) and A377124(n*10), taking A376446(n) if and only if n is not a multiple of 10 and A376446(n*10) otherwise, we should get the sequence: 8, 64, 2486, 5, 4268, 8426, 2, 1, 4, 4862, 46, 82, 6248, 6, 6842, 8624, 2684, 28, 9, 7139, 3179, 19, 1397, 1793, 91, 3971, 7931, 9713, 9317 (which the author conjectures to be complete, as the present one). a(1) = 8, a(2) = 64, ..., a(28) = 9317 (and a(28) is the last term of the present sequence - conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Nov 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016887 | OEIS-A378424

Product_{n>=1} (1+x^n)^a(n) = Sum_{k>=0} C(k)*x^k, where C(k) = A000108(k).

Conjecture: A327937(n) divides a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Nov 26 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016888 | OEIS-A378444

a(n) is the number of divisors d of n such that A083345(d) is even, where A083345(n) is the numerator of Sum(e/p: n=Product(p^e)).

a(n) = ceiling(A174273(n)/2). [Conjectured] - _Antti Karttunen_, May 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Nov 27 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016889 | OEIS-A378550

a(n) = 1 if sigma(n) >= 2*n-1, otherwise 0.

a(n) = A209229(n) + A294936(n). [Conjectured]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016890 | OEIS-A378568

Lowest weight of rational fraction with denominator n.

It is conjectured by Kravitz and Sah that a(n) = O(log n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Dec 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016891 | OEIS-A378575

G.f. A(x) satisfies A(x) = x + x*A(A(A(A(A(x))))), so that this sequence shifts left under the 5th self-COMPOSE.

Conjecture: a(n) == 1 (mod 4) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016892 | OEIS-A378576

G.f. A(x) satisfies A(x) = x + x*A(A(A(A(A(A(x)))))), so that this sequence shifts left under the 6th self-COMPOSE.

Conjecture: a(n) == 1 (mod 5) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 01 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016893 | OEIS-A378578

G.f. A(x) equals the series obtained by removing all factors of 3 from the coefficients in 1 + x*A(x)^3.

Conjecture: a(n) == binomial(3*n,n)/(2*n+1) (mod 2) for n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016894 | OEIS-A378580

G.f. A(x) satisfies: A(x/A(x)) = theta_3(x) = 1 + 2*Sum_{n>=1} x^(n^2).

Conjecture: for n > 0, a(n) == 2 (mod 4) iff n is square, else a(n) is divisible by 4 if n is nonsquare.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 08 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016895 | OEIS-A378625

The first subsequence starting at the first digit after the decimal point in the decimal expansion of Pi that is divisible by n.

For every positive integer n there exists a contiguous subsequence of the decimal expansion of Pi that is divisible by n (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon R Blow_, Dec 02 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016896 | OEIS-A378722

a(1) = 1, a(2) = 2. For n > 2, a(n) is the smallest number not already a term which is divisible by A007947(a(n-1)/A020639(a(n-1))).

Conjectured to be a permutation of the positive integers with primes in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Dec 05 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016897 | OEIS-A378738

Primitively abundant numbers k for which A378665(k) > A378664(k).

Conjecture: A001222(a(n)) = 3 <=> 3|a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 07 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016898 | OEIS-A378771

a(n) is the least k such that the last k digits of m = A020666(n)^n contain all 10 possible digits (0 through 9).

A conjecture over at A020666 says that A020666(n) = 2 for n >= 189. Perhaps looking at last digits and some modular arithmetic leads to a proof.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David A. Corneth_, Dec 06 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016899 | OEIS-A378838

a(n) is the number of n-digit nonnegative integers with the mean of the digits equal to the digital root.

Conjecture: a(A007494(n)) = 1. The above conjecture is true, i.e. if n == 0 or 2 mod 3, then a(n) = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Dec 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016900 | OEIS-A378840

G.f. A(x) satisfies A(x) = (1 + x * A(x)^(4/3) * (1 + A(x)^(1/3)))^3.

Conjecture: g.f.: B(x)^3, where B(x) is the g.f. of A260332.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Dec 09 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016901 | OEIS-A378929

Number of steps it takes for the chain of f(n) = n * d(n) / gcd(n, d(n))^2 to reach a cycle from a starting number n.

It is conjectured that every number n eventually ends up in a cycle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Viliam Furík_, Dec 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016902 | OEIS-A378944

Triangle read by rows: T(n,k) = number of stamp foldings with stamp #1 first, n stamps and stamp #2 covered by exactly one fold. k = the stamp number before the fold covering stamp #2 divided by 2. See examples.

The conjectured formula for the numbers in T(n,k) involves two unsolved sequences, semi-meanders and meandric numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Roger Ford_, Dec 11 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016903 | OEIS-A378980

Numbers k such that (A003961(k)-2*k) divides (A003961(k)-sigma(k)), where A003961 is fully multiplicative with a(prime(i)) = prime(i+1), and sigma is the sum of divisors function.

Conjecture: Apart from a(5)=6, this is a subsequence of A319630, i.e., for all terms k<>6, gcd(k, A003961(k)) = 1. See also A372562, A372566.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 12 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016904 | OEIS-A378983

Numbers k such that (A003961(k)-2*k) divides (A003961(k)-(1+sigma(k))), where A003961 is fully multiplicative with a(prime(i)) = prime(i+1), and sigma is the sum of divisors function.

Conjecture: A202274 gives all terms of A028982 that occur in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 13 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016905 | OEIS-A379044

a(1) = 1, a(2) = 2. For n > 2 a(n) is the smallest positive integer k, not already a term in the sequence, such that A053669(k) = A006530(a(n-1)).

The terms of the sequence can be extracted from the table in A125703, where the k-th row gives numbers whose least non divisor prime p is the k-th prime. From a(5) onwards the first appearance of q = Gpf(a(n-1)) = prime(k) gives a(n) = A002110(k-1), the first term in the k-th row of A125703. Subsequent appearances of same q selects a(n) from the same row. The primes q = Gpf(a(n-1)) decrement from term to term if rad(a(n-1) is in A002110, and occasionally we see a full cycle of q from some higher prime terminating at q = 2 (see Example). However this is rare, because of interruptions when rad(a(n-1)) is not primorial, causing q to increment before decrementing again. This retards the arrival of odd numbers in the sequence. Whereas we may suppose that all numbers in A125703 will eventually appear, the blocking of full decremetation cycles of q produces a "firewall" effect opposing this possibility. Conjecture: Full cycles of decrementing q conclude with a(n) a power of 2 (producing the next odd term).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Dec 14 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016906 | OEIS-A379049

a(n) = prime(i)*dp(n,i) + prime(i)*dn(n,i) where dp(n,i) = 1 when the i-th trit of n is 1, dn(n,i) = 1 when the i-th trit of n is T, and dp(n,i) = dn(n,i) = 0 when the i-th trit of n is 0.

Conjecture: All positive integers greater than 1 appear in this sequence at least once.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Dec 14 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016907 | OEIS-A379059

Index of n in A377091, or -1 if n does not appear there.

It is conjectured that every integer appears in A377091. Conjecture 1: For n >= 10, |a(n) - 2*n| < 2*sqrt(n); for n >= 1000, |a(n) - 2*n| < 1.59*sqrt(n) (compare A379786). - _N. J. A. Sloane_, Jan 19 2025 Conjecture 2: lim sup |a(n) - 2*n|/sqrt(n) = sqrt(2) as n -> oo. - _N. J. A. Sloane_ and _Paolo Xausa_, Feb 02 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 26 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016908 | OEIS-A379061

First differences of A377091.

lim sup a(n)/n = - lim inf a(n)/n = 1. (Conjectured.) - _M. F. Hasler_, Feb 08 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 27 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016909 | OEIS-A379067

Smallest missing positive number after A377091(n) has been found.

Conjecture: lim sup |a(n) - n/2|/sqrt(n) = 1/2 (see links). This would imply that after A377091(n) has been found, A377091 contains all numbers in the range [0,f(n)], where lim sup f(n) = (n-sqrt(n))/2. There is a corresponding conjecture for A379068. - _N. J. A. Sloane_ and _Paolo Xausa_, Feb 03 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Dec 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016910 | OEIS-A379104

a(n) = third elementary symmetric function of the first n distinct Fibonacci numbers.

Conjecture: for k >=1, the k-th elementary symmetric function (esf) of first n distinct Fibonacci numbers (fndFn) is a homogeneous linear recurrence sequence of order (n+2)(n+1)/2. The conjecture and signatures also apply to the first n distinct Lucas numbers (A000032).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Dec 16 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016911 | OEIS-A379140

Numbers k such that the greatest prime < 10^k and the least prime > 10^k share no decimal digits.

Charles R Greathouse IV conjectures that A107801(n) = prime(n) for n sufficiently large (and similarly for other related sequences). If that is the case, this sequence must be finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Dec 16 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016912 | OEIS-A379151

The binary weights of the Catalan numbers (A000108).

Conjecture: Sum_{k=1..n} a(k) ~ n^2 / 2 (see the plot in the Links section).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Dec 16 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016913 | OEIS-A379153

The binary weights of the Apéry numbers (A005259).

Conjecture: Limit_{m->oo} (1/m^2) * Sum_{k=1..m} a(k) = log(sqrt(2) + 1)/log(2) = 1.2715533... (Knopfmacher and Luca, 2012).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Dec 17 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016914 | OEIS-A379181

a(n) is the number of n-digit nonnegative integers with mode and mean of the digits equal.

Conjecture: a(n+1)/a(n) ~ 10. - _Stefano Spezia_, Jul 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Dec 17 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016915 | OEIS-A379202

G.f. A(x) satisfies 1/x = Sum_{n=-oo..+oo} A(x)^n * (A(x)^n + 2)^(n+1).

Conjecture: a(n) is even for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016916 | OEIS-A379204

G.f. A(x) satisfies 1/x = Sum_{n=-oo..+oo} A(x)^n * (A(x)^n + 4)^(n+1).

Conjecture: a(n) is even for n > 1. Conjecture: a(n) == 2 (mod 4) iff n = (k-1)^2 + 1 for some k > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 20 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016917 | OEIS-A379240

Lexicographically earliest infinite sequence such that a(i) = a(j) => f(i) = f(j), for all i, j, where f(n) = [A003415(n), A085731(n)] if A359550(n) = 1, otherwise f(n) = n.

It is conjectured that this is also the lexicographically earliest infinite sequence such that a(i) = a(j) => A003415(i) = A003415(j), A085731(i) = A085731(j) and A376418(i) = A376418(j), for all i, j >= 1, i.e., the restricted growth sequence transform of the triple [A003415(n), A085731(n), A376418(n)]. This is true if for every pair of i and j for which i <> j, and A376418(i) = A376418(j) > 0, the ordered pairs [A003415(i), A085731(i)] and [A003415(j), A085731(j)] differ from each other. In other words, conjecture claims that for all x, y in A100716, when x != y, either x' != y' OR gcd(x, x') != gcd(y, y') OR A376418(x) != A376418(y). If the conjecture 2 given in A399369 holds, then it implies the above claim as well. Adamczewski's AI-assisted disproof of the above conjecture in the linked arXiv-article (e.g., in versions 1-2, August 2026) is incorrect because it was based on a misinterpretation of A-number A376418.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016918 | OEIS-A379248

a(1) = 1, a(2) = 2, for a(n) > 2, a(n) is the smallest unused positive number that shares a factor with a(n-1) while no exponent of each distinct prime factor of a(n) is the same as the exponent of the same prime factor of a(n-1).

The sequence is conjectured to be a permutation of the positive integers. See A379293 for the index where n first appears.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Dec 18 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016919 | OEIS-A379260

Index of first appearance of n in sequence A379049.

Conjecture: a(n) is defined for all integer n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Dec 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016920 | OEIS-A379263

a(n) = n for 1 <= n <= 3. For n >3 a(n) is the smallest positive integer not already in the sequence which is prime to a(n-3) but not to a(n-2)*a(n-1).

Conjecture: Sequence is a permutation of the positive integers, with primes in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Dec 19 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016921 | OEIS-A379379

Sum of odd divisors of n except the "e" odd divisors described in A005279.

Conjectures from _Omar E. Pol_, Aug 27 2025: (Start)
Conjectures from _Omar E. Pol_, Oct 30 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Dec 21 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016922 | OEIS-A379399

Numbers that can be written in exactly four different ways as a sum of at most nine positive third powers.

Conjecture: this sequence is finite and a(66) = 455 is the last term. Verified up to 10^8. - _Charles R Greathouse IV_, Dec 28 2024

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Patrick De Geest_, Dec 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016923 | OEIS-A379402

Rectangular array, read by descending antidiagonals: the Type 2 runlength index array of A039701 (primes mod 3); see Comments.

Conjecture: every column is eventually increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016924 | OEIS-A379404

Rectangular array, by descending antidiagonals: the Type 2 runlength index array of A039702 (primes mod 4); see Comments.

Conjecture: every column is eventually increasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016925 | OEIS-A379440

a(1) = 1, a(2) = 2, for a(n) > 2, a(n) is the smallest unused positive number that shares a factor with a(n-1) such that the exponents of each distinct prime factor of a(n) differ by one from those of the same prime factors of a(n-1).

Like A379442, for the terms studied, prime terms p are preceded by p^2 and followed by 2*p^2, can be divisors of terms before they appear as a term themselves, and are distributed in groups of primes, with many primes within the groups differing by six terms. Unlike A379442 not all primes appear in their natural order, although the occurrence of such primes is rare - only four primes are out of order in the first 250000 terms, namely a(6787) = 179, a(18355) = 353, a(43516) = 593, a(201498) = 1579. In the same range the fixed points are 1, 2, 30, 34, 46, 130, 352, 456, although more may exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Dec 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016926 | OEIS-A379441

a(1) = 1, a(2) = 2, for a(n) > 2, a(n) is the smallest unused positive number that shares a factor with a(n-1) such that the exponents of each distinct prime factor of a(n-1) differ by one from those of the same prime factors of a(n).

Like A379442, for the terms studied, prime terms p are preceded by p^2 and followed by 2*p^2, can be divisors of terms before they appear as a term themselves, and are distributed in groups of primes, with many primes within the groups differing by six terms. Unlike A379442 not all primes appear in their natural order, although the occurrence of such primes is rare - only three primes are out of order in the first 250000 terms, namely a(13350) = 149, a(18410) = 179, a(21382) = 191. The sequence contains numerous fixed points, these being 1, 2, 34, 46, 218, 370, 410, 462, 474, 1954, 5592, 19186,... . The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Dec 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016927 | OEIS-A379442

a(1) = 1, a(2) = 2, for a(n) > 2, a(n) is the smallest unused positive number that shares a factor with a(n-1) such that the exponents of each distinct prime factor of a(n-1) differ by one from those of the same prime factors of a(n), while the exponents of each distinct prime factor of a(n) differ by one from those of the same prime factors of a(n-1).

The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Dec 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016928 | OEIS-A379446

a(n) is the number of ones in the binary expansion of 10^(10^n).

Conjectured: Limit_{n->oo} a(n)/10^n = log(5)/log(4). (A153201)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Dec 26 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016929 | OEIS-A379458

Array read by downward antidiagonals: A(n,k) = A(n-1,k+1) + 2*(k+1)!*Sum_{j=0..k} A(n-1,j)/j! with A(0,k) = 1, n >= 0, k >= 0.

Conjecture: A(n,0) = A217061(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Dec 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016930 | OEIS-A379459

Array read by downward antidiagonals: A(n,k) = A(n-1,k+1) + 2*(k+1)!*Sum_{j=0..k} A(n-1,j)/j! with A(0,k) = 2*(k+1)!, n >= 0, k >= 0.

Conjecture: A(n,0) = A006351(n+2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Dec 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016931 | OEIS-A379460

Array read by downward antidiagonals: A(n,k) = A(n,k-1) + (k+1)*(A(n-1,k) + A(n-1,k+1)) with A(n,0) = A(n-1,0) + A(n-1,1), A(0,k) = 1, n >= 0, k >= 0.

Conjecture: A(n,0) = A006351(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Dec 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016932 | OEIS-A379463

a(n) is the total number of paths starting at (0, 0), ending at (n, 0), consisting of steps (1, 1), (1, 0), (1, -3), and staying on or above y = -1.

Conjecture D-finite with recurrence 3*n*(3*n+4)*(n-3)*(3*n+8)*a(n) +3*(-45*n^4+54*n^3+192*n^2-27*n-20)*a(n-1)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emely Hanna Li Lobnig_, Dec 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016933 | OEIS-A379464

a(n) is the total number of paths starting at (0, 0), ending at (n, 0), consisting of steps (1, 1), (1, 0), (1, -3), and staying on or above y = -2.

Conjecture D-finite with recurrence +3*(n+4)*(3*n+4)*(3*n+8)*a(n) +3*(-63*n^3-297*n^2-349*n-60)*a(n-1) +3*(189*n^3+270*n^2-229*n-140)*a(n-2) +15*(-63*n^3+117*n^2+44*n-64)*a(n-3) +(689*n^3-5372*n^2+6946*n-1288)*a(n-4) +(n-4)*(201*n^2+2767*n-3011)*a(n-5) -(n-5)*(579*n+257)*(n-4)*a(n-6) +229*(n-5)*(n-6)*(n-4)*a(n-7)=0. - _R. J. Mathar_, Jan 29

2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Emely Hanna Li Lobnig_, Dec 23 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016934 | OEIS-A379505

a(n) is the number of ways of partitioning the divisors of n into two disjoint sets with equal sum, when an extra 1-divisor is added to the divisor set, and the two 1-divisors are considered indistinguishable.

Conjecture 1: For all n >= 1, a(A156942(n)) > 0. Also, if a(A156942(n)) > 1 is true for all n, it would imply that there are no quasiperfect numbers, numbers x with sigma(x) = 2x+1, as such numbers must all reside in A156942 and have a(x) = 1. (See references in A336700). Conjecture 2: a(n) = 1 if and only if n = 2^k, with k >= 0. This claim is equal to the statement that there are neither quasiperfect numbers nor almost perfect (least deficient) numbers, numbers x with sigma(x) = 2x-1, others than those given by A000079.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016935 | OEIS-A379525

a(1) = 1. For n > 1, if a(n-1) is a novel term, a(n) = A003132(a(n-1)). If a(n-1) has been seen k (>1) times already, a(n) = k*a(n-1).

Conjecture: Every positive integer appears in the sequence at least once.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Dec 24 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016936 | OEIS-A379548

a(1) = 1. For n > 1, a(n) is the earliest novel squarefree number divisible by the smallest prime which does not divide a(n-1).

Squarefree version of A351495, conjectured to be a permutation of the squarefree numbers, A005117.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Dec 25 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016937 | OEIS-A379600

a(n) is the semiperimeter of the primitive Pythagorean triangle (x(n), y(n), z(n)) with x(n) < y(n) < z(n) and x(n) > x(n-1), y(n) > y(n-1), z(n) > z(n-1), which has the smallest perimeter (if there are several triangles with smallest perimeter: the one of these with the smallest area), starting from a(1) = (3 + 4 + 5)/2 = 6.

Conjecture: There is no primitive Pythagorean triangle that has a smaller semiperimeter than a(n) that can be drawn around the primitive Pythagorean triangle (x(n-1), y(n-1), z(n-1)) without touching it.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Feb 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016938 | OEIS-A379603

a(n) is the largest n-digit number whose square contains only digits greater than 5.

Conjecture: It appears that for all n >= 5,

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhining Yang_, Dec 27 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016939 | OEIS-A379630

Irregular triangle read by rows in which row n lists the smallest parts of the partitions of n into consecutive parts followed by the conjugate corresponding odd divisors of n in accordance with the theorem of correspondence described in the Comments lines.

Conjecture: the first A001227(n) terms in the n-th row are also the absolute values of the n-th row of A341971.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Dec 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016940 | OEIS-A379633

Irregular triangle read by rows in which row n lists the largest parts of the partitions of n into consecutive parts (with the partitions ordered by increasing number of parts).

Conjecture: the right border gives A212652.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Dec 31 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016941 | OEIS-A379643

List of x coordinates of prime numbers in a Cartesian grid, where the first prime 2 is placed at the origin (0,0) and the second prime 3 at (1,0). For the n-th prime prime(n), n >= 3, take a unit step in the direction (prime(n)-3)*45 degrees counterclockwise from the positive x-axis.

Conjecture: no prime appears on the negative y-axis.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Dec 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016942 | OEIS-A379652

a(1) = 2. For n > 1, a(n) = smallest prime factor of c=2*a(n-1)+1 that is not in {a(1), ..., a(n-1)}; if all prime factors of c are in {a(1), ..., a(n-1)}, then we try the next value of c, which is 2*c+1; and so on.

Conjecture: this sequence is a permutation of the primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert C. Lyons_, Dec 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016943 | OEIS-A379654

Least positive integer k <= n such that |tau(k)|*n + 1 is prime, or 0 if such k does not exist, where the tau function is given by A000594.

Conjecture 1: a(n) > 0 for all n > 0. In other words, for each positive integer n, there is a number k among 1,...,n such that |tau(k)|*n + 1 is prime. Conjecture 2: For each integer n > 1 not equal to 22, there is a number k among 1,...,n such that |tau(k)|*n - 1 is prime. We have verified both conjectures for n up to 10^8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Dec 28 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016944 | OEIS-A379665

Minimum number of intercalates in a Brown's diagonal Latin square of order 2n.

Hypothesis: minimum number of intercalates in Brown's diagonal Latin squares of order N=2n is equal to (N/2)^2 for N>4 (proved for N=6 and N=8 using Brute Force and for 10<=N<=24 using heuristic methods).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eduard I. Vatutin_, Dec 29 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016945 | OEIS-A379703

Antidiagonal sums of A343052.

Conjecture: a(n) = A273790((n-1)/2) for n odd and > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Dec 30 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016946 | OEIS-A379724

a(n) = A000668(n) - 1.

Conjecture; a(n) = A072868(n) - 2. - _Omar E. Pol_, Jan 16 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_ and _Antti Karttunen_, Dec 31 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016947 | OEIS-A379726

Minimum number of kings that must be placed on an n X n chessboard such that each square is attacked or occupied by at least two kings.

Every third term is conjectured to be A014105. Other terms are A001105. A093353 is conjectured to be this sequence with repeated terms removed. The above conjectures are true (see Beveridge link). - _Colin Beveridge_, Jan 13 2025 If n is a multiple of 3, it is conjectured that a(n)=2*(n/3)^2+n/3. The above conjectures are true (see Beveridge link). - _Colin Beveridge_, Jan 13 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matthew Scroggs_, Dec 31 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016948 | OEIS-A379732

Decimal expansion of 207/208.

Conjectured densest packing of truncated tetrahedra.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Dec 31 2024 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016949 | OEIS-A379748

a(n) is the number of ways to arrange any number of unit square cells into an i X j rectangle which contains exactly n square subarrays of all sizes.

It appears all natural numbers appear in the sequence. This is merely conjectured, but is provably true if there are an infinite amount of Sophie-Germain primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Adams_, Jan 02 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016950 | OEIS-A379753

Numbers that set records in A379752.

Conjecture: subset of A332785 = A126706 \ A286708.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jan 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016951 | OEIS-A379757

a(n) = a(n-1) + 1 with two exceptions: if a(n-1) is prime, a(n) = a(n-2) + a(n-1), or if a(n-1) is a power, a(n) = a(n-1) / (root factor), with initial three terms are 0, 1, 2.

Conjecture: log(a(n)) ~ k*sqrt(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bill McEachen_, Jan 02 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016952 | OEIS-A379764

G.f. A(x) satisfies Sum_{n=-oo..+oo} (4^n*A(x)^n - 4*A(x))^(n+1) = 2*theta_4(x) = 2*Sum_{n=-oo..+oo} (-x)^(n^2).

Conjecture: for n > 0, a(n) is odd iff n is a square. Conjecture: for n > 0, a(n) == (-1)^(n-1) (mod 4) iff n is a square, otherwise a(n) is divisible by 4 if n is nonsquare.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016953 | OEIS-A379773

Numbers that set records in A379772.

Conjecture: a(n) is powerful (i.e., in A286708) for n >= 68. Additionally, for some n much larger than 68, a(n) may be cubefull (i.e., in A372695).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jan 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016954 | OEIS-A379810

Composite numbers in A381019 which are immediately followed by another composite number, in order of their appearance.

Conjecture: For any k >= 1, A381019 contains k consecutive composite terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016955 | OEIS-A379817

Irregular triangle T(n, k), n >= 0, k >= 0, read by rows such that T(n,k) = f(n,k)/f(2^k-1,k) where f(n,k) is defined in Comments.

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Jan 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016956 | OEIS-A379854

a(0) = 0, and for any n > 0, a(n) is the least integer (in absolute value) not yet in the sequence such that the absolute difference of a(n-1) and a(n) is a cube; in case of a tie, preference is given to the positive value.

Conjecture: Every integer (positive or negative) appears in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jan 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016957 | OEIS-A379857

Number of values of k for which n can be written as a sum of k distinct positive squares.

a(n) < (3n)^(1/3) for n > 0. I conjecture that a(n) ~ (3n)^(1/3). - _Charles R Greathouse IV_, Feb 05 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luke E. Holland_, Jan 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016958 | OEIS-A379873

Nonnegative values in A377091, in order of appearance and with offset 0.

Conjecture: this sequence is a permutation of the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jan 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016959 | OEIS-A379874

Nonpositive values in A377091, negated, in order of appearance and with offset 0.

Conjecture: this sequence is a permutation of the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Jan 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016960 | OEIS-A379899

a(1) = 2. For n > 1, a(n) = smallest prime factor of c=a(n-1)+4 that is not in {a(1), ..., a(n-1)}; if all prime factors of c are in {a(1), ..., a(n-1)}, then we try the next value of c, which is c+4; and so on.

Conjecture: this sequence is a permutation of the primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert C. Lyons_, Jan 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016961 | OEIS-A380033

Numbers that set records in A380032.

Conjecture 1: Almost all numbers in this sequence are powerful squares. Only 12, 720, and 2880 are not powerful. Thereby this sequence is a proper subset of A368682 (intersection of A025487 and A131605, the latter a subset of A001597 and A286708), in turn a subset of A364710. Conjecture 2: 36, 900, and 44100 are the only squares of primorials (in A061742) in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jan 11 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016962 | OEIS-A380038

Triangle read by rows: T(n, k) = T(n-k, k) - T(n-k+1, k-1), T(n, 0) = A010815(n-1), and T(0, 0) = 1.

The initializing column T(n>0, 0) can be chosen at random. It is conjectured that T(n, 0) = A010815(n-1) minimizes the row-wise sum of absolute values for growing n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Friedjof Tellkamp_, Jan 12 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016963 | OEIS-A380049

Partial sums of A072203.

Conjecture: The average value of a(n) is 2*n^(3/2)/(-3*zeta(1/2)). - _Vaclav Kotesovec_, Jan 15 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tsuyoshi Hanatate_, Jan 10 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016964 | OEIS-A380056

E.g.f. (exp(x) - 1)/cos(2*x).

Conjecture: a(n) == 1 (mod 4) for n >= 1. Conjecture: a(4*n) is divisible by 5 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 28 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016965 | OEIS-A380066

G.f. A(x) satisfies 1/2 = Sum_{n=-oo..+oo} x^n*A(x)^n * (A(x)^n + x)^(n-1) * (A(x) + x^n)^(n-1).

Conjecture: for n > 0, a(n) == 2 (mod 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016966 | OEIS-A380070

Square pyramidal numbers sp(n) with a zeroless decimal representation such that (product of decimal digits of sp(n)) / n is an integer.

I conjecture the sequence has only finitely many members, i.e., for all n, sp(n) < 10^r for a large enough r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Jan 11 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016967 | OEIS-A380128

Triangle read by rows: Riordan array (1/(C(x)*sqrt(1-4*x)), x/C(x)) where C(x) is g.f. of A000108.

Conjecture: T(n, k) = Sum_{i=0..n-k} (-1)^i * binomial(2*n - 2*k, n - k - i) * binomial(k + i, i). The conjecture is true (see Fried link). - _Sela Fried_, Dec 05 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Jan 12 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016968 | OEIS-A380137

The conjectured maximum multiplicative persistence for bases b >= 2.

The conjectured maximum multiplicative persistence for bases b >= 2. The conjectured values for 32,33,35,36 and 39 are 12,17,16,13 and 19 respectively.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Jan 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016969 | OEIS-A380191

Triangle read by rows: Riordan array (2 - D(x), x * D(x)) where D(x) is g.f. of A001764.

Conjecture: Sum_{i=0..n-k} binomial(2*i, i) * T(n, i+k) = A110616(n, k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Jan 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016970 | OEIS-A380274

Sum of cubes of coefficients of q in the q-factorials.

Conjecture: a(n) ~ 6*sqrt(3) * n!^3 / (Pi * n^3). a(n) = 6*sqrt(3)/Pi * n!^3/n^3 * (1 - 93/(50*n) + 657703/(122500*n^2) - 18214629/(1225000*n^3) + o(n^(-3))), which proves above conjecture. See Wang link. - _Xinjun Wang_, May 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jan 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016971 | OEIS-A380275

Sum of the fourth powers of the coefficients of q in the q-factorials.

Conjecture: In general, sum of the k-th powers of the coefficients of q in the q-factorials is asymptotic to 2^((k-1)/2) * 3^(k-1) * n!^k / (sqrt(k) * Pi^((k-1)/2) * n^(3*(k-1)/2)). Conjecture: a(n) ~ 27*sqrt(2) * n!^4 / (Pi^(3/2) * n^(9/2)). This conjectured asymptotic formula is proved by Wang, who also gives the following third-order refinement.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jan 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016972 | OEIS-A380281

Triangle T(n, k) read by rows: T(n, k) = 2^n*binomial(2*n + 1, 2*k + 1) * Pochhammer(1/2, n - k) * Pochhammer(1/2, k).

Conjecture: Row sums are A076729.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Jan 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016973 | OEIS-A380290

a(n) = [x^n] G(x)^n, where G(x) = Product_{k >= 1} 1/(1 - x^k)^(k^2) is the g.f. of A023871.

We conjecture that a(p) == 1 (mod p^3) for all primes p >= 7 (checked up to p = 61). More generally, we conjecture that the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) holds for all primes p >= 7 and positive integers n and r. Some examples are given below.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jan 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016974 | OEIS-A380291

a(n) = [x^n] G(x)^n, where G(x) = Product_{k >= 1} (1 + x^k)^(k^2) is the g.f. of A027998.

We conjecture that a(p) == 1 (mod p^3) for all primes p >= 7 (checked up to p = 61). More generally, we conjecture that the supercongruences a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) hold for all primes p >= 7 and positive integers n and r. Some examples are given below.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jan 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016975 | OEIS-A380312

Primes not reaching 3 under iterations of p -> gpf(2*p-1).

Conjecture: a(n) ~ k*n log n for some constant k. Perhaps k ≈ 4.51. - _Charles R Greathouse IV_, Jan 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Johannes M.V.A. Koelman_, Jan 20 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016976 | OEIS-A380317

The lexicographically earliest sequence of positive numbers which is identical to the run lengths of its first differences.

Conjecture: Every positive integer eventually appears.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dominic McCarty_, Feb 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016977 | OEIS-A380333

Where prime race 8m+3 vs. 8m+7 is tied.

Conjecture: There is no prime on the negative y-axis of the Cartesian grid defined in A379643, meaning that prime p does not exist such that pi_{8,3}(p) - pi_{8,7}(p) = 0 and pi_{8,5}(p) - pi_{8,1}(p) < 0, where pi_{m,b}(x) is the number of primes <= x which are congruent to b (mod m).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Jan 21 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016978 | OEIS-A380346

Number of corona for a hexagon of edge n with diamonds of side 1.

a(n) = n^6 + 6*n^5 + 21*n^4 + 44*n^3 + 60*n^2 + 48*n + 18 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Craig Knecht_, Jan 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016979 | OEIS-A380354

a(n) = phi(2 + phi(3 + phi(5 + ... + phi(prime(n))))), where phi is Euler totient function (A000010).

Conjecture 1: a(n) can be only 1, 2, 4, 6, 8, 12, 16, 20, 18, 40, 72, 96, 64, 48 or 42. Conjecture 2: for n >= 187, a(n) can be only 20 or 64.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Jan 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016980 | OEIS-A380386

a(n) is the least number k that contains, in base n, every digit once, and such that k/(n-1) (if n is even) or 2*k/(n-1) (if n is odd) is prime.

Conjecture: a(n) exists for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Jan 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016981 | OEIS-A380392

Irregular triangle read by rows: T(n,k) is the number of n X n binary matrices containing k South-East paths of 1's connecting the top left and bottom right corners.

Conjecture: The average number of South-East paths of 1's in all n X n binary matrices is A001790(n-1)/A101926(n-1). - _John Tyler Rascoe_, Feb 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John Tyler Rascoe_, Jan 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016982 | OEIS-A380450

Number of integers k such that prime(n) - primorial(k) is prime.

Conjecture A: Each value occurs an infinite number of times in the sequence. Conjecture B: All natural numbers occur in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel D Gibson_, Jun 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016983 | OEIS-A380455

Maximum number of prime polyomino factors of an n-polyomino.

Conjectured terms: a(14)..a(17) = 10?, 3, 12?, 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gordon Hamilton_, Jun 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016984 | OEIS-A380464

Integers k such that A005245(m*k) < A005245(k) for some m.

Is this integer sequence infinite? This is an open problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John M. Campbell_, Jun 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016985 | OEIS-A380475

a(n) is the least term in A380468 that has exactly n prime factors.

Conjecture: Sequence is finite and a(4) is the last term. This is equivalent to the claim that no arithmetic derivative (A003415) of a product of five or more distinct primes, (i.e., the value of the (n-1)-st elementary symmetric polynomial formed from those n distinct primes) can be formed as a carry-free sum of those n summands in primorial base (A049345). See also A380476 and A380528, A380530.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016986 | OEIS-A380476

Numbers k with at least 4 prime factors such that A380459(k) is in A048103, i.e., has no divisors of the form p^p.

It is conjectured that there are no terms with more than four prime factors. See A380475 and A380528, A380530, also A380526.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016987 | OEIS-A380483

a(n) = least k such that A010846(6*prime(k)) = n, or -1 if no solution exists.

Conjecture: n = 30 is the largest number for which a(n) = -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jul 08 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016988 | OEIS-A380487

Numbers such that the sum of prime factors without repetition divides the product of prime factors without repetition and each division yields a greater quotient.

Conjecture: There exists m, the smallest positive integer such that a(k)|a(k+m), for all k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Torlach Rush_, Jan 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016989 | OEIS-A380494

a(1) = 1. For n > 1 a(n) is the smallest positive integer not yet in the sequence which is divisible by A007953(a(n-1)) + 1.

For n > 1 a(n) is the smallest novel multiple of digsum(a(n-1)) + 1. The "+1" is intended to avoid an infinite number of consecutive multiples of 9, which would otherwise occur after a(13). The only way a(n) = prime p can occur is if A007953(a(n-1)) is the first occasion of p-1. Sequence is conjectured to be a permutation of the natural numbers, with primes in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_ and _Ali Sada_, Jan 25 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016990 | OEIS-A380527

Numbers k such that k is a multiple of A327860(k), where A327860 is the arithmetic derivative of the primorial base exp-function.

It was conjectured that only terms of A276156 occur here. (This is false, see below.) Sequence obtained by applying A276085 to the {intersection of A373487 and A048103} and sorting the terms into ascending order. As that intersection contains at least one nonsquarefree term, 855523 = 67 * 113^2, the term x = A276085(855523) = 55946999363570987235666961212549502632097773 0 (with primorial base expansion A049345(x) = 200000000001000000000000000000) is a term of this sequence that is NOT in A276156, thus the conjecture made above is not true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 11 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016991 | OEIS-A380530

Positions of records in A380528.

Questions: Are all terms squarefree after 4, and do all terms end (in base ten) with digit 6 from the fifth term onward? Is the sequence infinite? Are there any terms with more than four prime factors? See also the conjecture given in A380475, and A380476.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 09 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016992 | OEIS-A380558

G.f. A(x) satisfies A(x - A(x)) = x^2/(1 - x^2).

Conjecture: a(n) is odd iff n = 2*A004760(k) for some k > 1, where A004760 lists numbers whose binary expansion does not begin 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016993 | OEIS-A380573

Number of distinct free polyominoes that are terraces in the first n levels of the stepped pyramid described in A245092.

Conjecture: the polyomino "I" of width 1 and length k appears for the first time in the level 2*k - 1 starting from the top of the stepped pyramid, k >= 1. In other words: that polyomino appears for the first time in the symmetric representation of sigma(2*k-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Mar 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016994 | OEIS-A380581

a(n) = [x^n] G(x)^n, where G(x) = Product_{k >= 1} 1/(1 - x^k)^(k^4) is the g.f. of A023873.

We conjecture that a(p) == 1 (mod p^3) for all primes p >= 5 (checked up to p = 61). More generally, we conjecture that the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) holds for all primes p >= 5 and positive integers n and r. Some examples are given below. Let A >= 1 and B be integers. Define a sequence {u(n) : n >= 0} by u(n) = [x^(A*n)] G(x)^(B*n), so the present sequence is the case A = B = 1. We conjecture that the above supercongruences also hold for the sequence {u(n)}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jan 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016995 | OEIS-A380582

a(n) = [x^n] G(x)^n, where G(x) = Product_{k >= 1} ((1 + x^k)/(1 - x^k))^(k^2) is the g.f. of A206622.

We conjecture that a(p) == 1 (mod p^3) for all primes p >= 5 (checked up to p = 61). More generally, we conjecture that the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) holds for all primes p >= 5 and positive integers n and r. Some examples are given below. Let A, B be integers and let C be a positive integer. Define u(n) = [x^(C*n)] Product_{k >= 1} ((1 + x^k)^A * (1 - x^k)^B)^(k^2). The present sequence is the case A = 1, B = -1 and C = 1. We conjecture that the above supercongruences also hold for the sequence {u(n)} for all primes p >= 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jan 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016996 | OEIS-A380583

a(n) = [x^n] G(x)^n, where G(x) = Product_{k >= 1} ((1 + x^(2*k))/(1 - x^k))^(k^2).

We conjecture that a(p) == 1 (mod p^3) for all primes p >= 7. More generally, we conjecture that the supercongruence a(n*p^r) == a(n*p^(r-1)) (mod p^(3*r)) holds for all primes p >= 7 and positive integers n and r. Some examples are given below. Let A and B be integers. Let C be a positive integer. Define u(n) = [x^(C*n)] Product_{k >= 1} ((1 + - x^(2*k))^A * (1 + - x^k)^B)^(k^2). The present sequence is the case A = 1, B = -1 and C = 1, with the appropriate choice of signs. We conjecture that the above supercongruences also hold for the sequence {u(n)} for all primes p >= 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Bala_, Jan 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016997 | OEIS-A380609

Primes a single step away from a cycle under the mapping p-> gpf(2*p+1).

Conjecture: under repeated mapping all primes ultimately enter the same cycle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Johannes M.V.A. Koelman_, Jan 28 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016998 | OEIS-A380645

The expansion of the Stieltjes continued fraction 1/(1 - x/(1 - a(A053645(0))*x/(1 - a(A053645(1))*x/(1 - a(A053645(2))*x/...)))) gives the sequence itself.

The Hankel transforms of this sequence starting with offset 0 and starting with offset 1 are both strictly positive, this means this sequence can be considered as a sequence of moments in the Stieltjes moment problem: a(n) = Integral_{x>=0} x^n*V(n, x) dx. It is conjectured that the measure V(n, x) is related to some kind of interesting distribution which shows self similarity over intervals of powers of two.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Feb 06 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-016999 | OEIS-A380671

a(n) is the smallest number not yet in the sequence which is coprime to n and shares at least one decimal digit with n.

Like A065190 but with the extra condition that n and a(n) must have at least one decimal digit in common. Definition implies that a(1) = 1 is the only fixed point. Let a(n) be even then 2|a(n) -> 2!|n-> 2|(n+1)->2!|a(n+1), therefore there are no consecutive even terms. Let [n] = |n - a(n)|, then it follows from the coprime conditions of the definition that n, a(n), [n] are pairwise coprime. Sequence is conjectured to be a permutation of the natural numbers with [n] < 100 (= base^2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_ and _Michael De Vlieger_, Jan 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017000 | OEIS-A380678

G.f. A(x) satisfies A(x - A(x)^2/(1 - A(x)^2)) = x.

Conjecture: for n>= 1, a(n) is odd iff n is a term of A027383, where A027383(2*m) = 3*2^m - 2 and A027383(2*m+1) = 4*2^m - 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 14 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017001 | OEIS-A380681

G.f. A(x) satisfies 2*x = Sum_{n=-oo..+oo} (-1)^n * x^(2*n) * (A(x) + x^n)^(2*n) with A(0) = 1.

Conjecture: if F(x) satisfies p*x = Sum_{n=-oo..+oo} (-1)^n * x^(p*n) * (F(x) + x^n)^(p*n) with F(0) = 1 for some fixed integer p, then F(x) is an integer series in x iff p is prime. The g.f. A(x) of this sequence is the case when p = 2; other cases include A380682 (p=3), A380683 (p=5), A380684 (p=7), A380685 (p=11), and A380686 (p=13).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017002 | OEIS-A380682

G.f. A(x) satisfies 3*x = Sum_{n=-oo..+oo} (-1)^n * x^(3*n) * (A(x) + x^n)^(3*n) with A(0) = 1.

Conjecture: if F(x) satisfies p*x = Sum_{n=-oo..+oo} (-1)^n * x^(p*n) * (F(x) + x^n)^(p*n) with F(0) = 1 for some fixed integer p, then F(x) is an integer series in x iff p is prime. The g.f. A(x) of this sequence is the case when p = 3; other cases include A380681 (p=2), A380683 (p=5), A380684 (p=7), A380685 (p=11), and A380686 (p=13).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017003 | OEIS-A380683

G.f. A(x) satisfies 5*x = Sum_{n=-oo..+oo} (-1)^n * x^(5*n) * (A(x) + x^n)^(5*n) with A(0) = 1.

Conjecture: if F(x) satisfies p*x = Sum_{n=-oo..+oo} (-1)^n * x^(p*n) * (F(x) + x^n)^(p*n) with F(0) = 1 for some fixed integer p, then F(x) is an integer series in x iff p is prime. The g.f. A(x) of this sequence is the case when p = 5; other cases include A380681 (p=2), A380682 (p=3), A380684 (p=7), A380685 (p=11), and A380686 (p=13).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017004 | OEIS-A380684

G.f. A(x) satisfies 7*x = Sum_{n=-oo..+oo} (-1)^n * x^(7*n) * (A(x) + x^n)^(7*n) with A(0) = 1.

Conjecture: if F(x) satisfies p*x = Sum_{n=-oo..+oo} (-1)^n * x^(p*n) * (F(x) + x^n)^(p*n) with F(0) = 1 for some fixed integer p, then F(x) is an integer series in x iff p is prime. The g.f. A(x) of this sequence is the case when p = 7; other cases include A380681 (p=2), A380682 (p=3), A380683 (p=5), A380685 (p=11), and A380686 (p=13).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017005 | OEIS-A380685

G.f. A(x) satisfies 11*x = Sum_{n=-oo..+oo} (-1)^n * x^(11*n) * (A(x) + x^n)^(11*n) with A(0) = 1.

Conjecture: if F(x) satisfies p*x = Sum_{n=-oo..+oo} (-1)^n * x^(p*n) * (F(x) + x^n)^(p*n) with F(0) = 1 for some fixed integer p, then F(x) is an integer series in x iff p is prime. The g.f. A(x) of this sequence is the case when p = 11; other cases include A380681 (p=2), A380682 (p=3), A380683 (p=5), A380684 (p=7), and A380686 (p=13).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017006 | OEIS-A380686

G.f. A(x) satisfies 13*x = Sum_{n=-oo..+oo} (-1)^n * x^(13*n) * (A(x) + x^n)^(13*n) with A(0) = 1.

Conjecture: if F(x) satisfies p*x = Sum_{n=-oo..+oo} (-1)^n * x^(p*n) * (F(x) + x^n)^(p*n) with F(0) = 1 for some fixed integer p, then F(x) is an integer series in x iff p is prime. The g.f. A(x) of this sequence is the case when p = 13; other cases include A380681 (p=2), A380682 (p=3), A380683 (p=5), A380684 (p=7), and A380685 (p=11).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017007 | OEIS-A380708

G.f. A(x) satisfies A(x) = 1 + x*abs(1/A(x))^2.

Conjecture: for n > 0, a(n) is odd iff n = 2*A003714(k) + 1 for k >= 0, where A003714 are the fibbinary numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 08 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017008 | OEIS-A380709

G.f. A(x) satisfies A(x) = 1 + x*abs(1/A(x))^3.

Conjecture: a(n) == binomial(4*n-1, n) (mod 2) for n >= 0 (cf. A263132).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 09 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017009 | OEIS-A380710

G.f. A(x) satisfies A(x) = 1 + x*A(x)*abs(1/A(x)^2).

Conjecture: for n > 0, a(n) is odd iff n is a power of 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017010 | OEIS-A380711

G.f. A(x) satisfies A(x) = 1 + x*A(x)*abs(1/A(x)^3).

Conjecture: a(n) == binomial(3*n-1,n)/(3*n-1) (mod 2) for n >= 0. Conjecture: for n > 0, a(n) is odd iff n = 2*A003714(k) + 1 for k >= 0, where A003714 are the fibbinary numbers. (This is equivalent to the prior conjecture.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017011 | OEIS-A380728

For n a power of 2, a(n) = n. Otherwise a(n) is the smallest number not yet in the sequence which is coprime to n and has the same binary weight as n.

Conjectured to be a permutation of the natural numbers (primes not in order).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jan 31 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017012 | OEIS-A380739

For n = 2^k (k>=0), a(n) = n. Otherwise a(n) != n is the smallest number not yet in the sequence having the same binary weight as n and such that gcd(n,a(n)) > 1.

Similar definition to A380728, conjectured to be a permutation of the natural numbers, with primes in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Jan 31 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017013 | OEIS-A380756

a(n) is the smallest number not yet in the sequence which is coprime to n and has the same number of 0's in its binary expansion as n.

a(n) is the smallest novel number coprime to n with A023416(a(n)) = A023416(n). Sequence is self inverse and conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Feb 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017014 | OEIS-A380760

Integers k with at least one proper factorization for which the sum of the same fixed integer power >= 2 of the factors equals k.

Superset of A381538 for values >= 16, and it is conjectured that the terms that match multiple nonequivalent factorizations here, such as a(5) = 256 (see Example), are exactly the terms of A381538 that can be produced as m^(m^e) by multiple m.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles L. Hohn_, Feb 02 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017015 | OEIS-A380787

Odd positive integers k whose continued fraction for sqrt(k) has a central term equal to either floor(sqrt(k)) or floor(sqrt(k)) - 1.

Conjecture: All terms are congruent to 3 mod 4 and all primes of this form (A002145) are terms of the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giorgos Kalogeropoulos_, Feb 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017016 | OEIS-A380835

Second column of Kimberling's ESC array.

Kimberling conjectured that every term of this sequence is even. This was proved by Behrend (2012).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Feb 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017017 | OEIS-A380876

a(1) = 1; a(2) = 4; for n > 2, a(n) = least unused positive y such that gcd(y,n-1) > 1 and |y-n| > 1.

For prime numbers p, a(2p+1) = p, and for p > 3, a(p+1) = 2p. (Conjectured.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ali Sada_, Feb 06 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017018 | OEIS-A380893

Triangle read by rows: T(n,m) = number of solid partitions of n with shape of a plane partition of m.

Conjecture: If A000219(k) == 0 (mod 3) then all terms of the k-th column of this triangle are also congruent to 0 (mod 3). - _John Tyler Rascoe_, Feb 08 2026 The above conjecture is false, it fails at column 14 see the Mathematics Stack Exchange link. - _John Tyler Rascoe_, Mar 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_, Feb 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017019 | OEIS-A380902

Integers k with at least 1 proper factorization for which the sum of the squares of the factors equals k.

It is conjectured that this sequence is infinite, that it does not contain any squarefree terms (A005117), and that with the exception of 16 (2^4) and 27 (3^3) it does not contain any squareful terms (A001694) or examples where the factors are all primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles L. Hohn_, Feb 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017020 | OEIS-A380944

a(n) = b(n,A000120(n)) for n >= 0 where b(n,k) is defined in Comments.

Conjecture: a(2^m*(2^n-1)) = (n+1)^m - n*n!*c(m,n+1) for n >= 0, m >= 0 where c(n,k) = Sum_{i=0..n-k} Stirling2(k+i,k) for n >= 0, k >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Feb 09 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017021 | OEIS-A380971

Irregular triangle T(n, k), n >= 0, k > 0, read by rows with row polynomials R(n, x) such that R(2n+1, x) = x*R(n, x) for n >= 0, R(2n,x) = wt(n)*x*((x+1)^wt(n) - x^wt(n)) + Sum_{k=1..wt(n)} k*x^k*T(n,k) for n > 0 with R(0,x) = 0 where wt(n) = A000120(n).

Conjectures: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Feb 10 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017022 | OEIS-A380974

Numbers k such that k*(k-1) is composed of exactly two different decimal digits.

Conjecture: all terms >= 334 are of the form 10...0, 10...01, 16...67, 3...34, or 6...67. Conjectured: for k >= 0, Conjectured G.f.: (4*x + 5*x^2 + 6*x^3 + 7*x^4 + 8*x^5 - 35*x^6 - 45*x^7 - 55*x^8 - 60*x^9 - 64*x^10 - 34*x^11 - 28*x^12 - 27*x^13 - 50*x^14 - 109*x^15 - 107*x^16 - 152*x^17 - 163*x^18 - 425*x^19 - 418*x^20 - 274*x^21 -

113*x^22 + 229*x^23 + 109*x^24 + 580*x^25 + 440*x^26 + 330*x^27 + 10*x^28 + 410*x^29)/(1 - 11 * x^5 + 10 * x^10).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Feb 11 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017023 | OEIS-A380984

Primes p such that p*(p-1) consists of exactly two different decimal digits.

Contains (10^k+2)/6 for k in A076850 and (2*10^k + 1)/3 for k in A096507. It is conjectured that these sequences are infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Feb 11 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017024 | OEIS-A381003

Lexicographically earliest sequence with a(0) = 0 and a(n) = a(n + a(n)) - a(n - a(n)) > 0.

We assume a(<0) = 0 in this definition but conjecture that a(n) < n for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Feb 11 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017025 | OEIS-A381005

Ordered short legs of the Pythagorean triangles defined by a = 2^(4n) + 2^(2n+1), b = 2^(4n) - 2^(4n-2) - 2^(2n) - 1, c = 2^(4n) + 2^(4n-2) + 2^(2n) + 1.

Conjecture: These Pythagorean triangles are primitive. Verified up to n=100000. The preceding conjecture is true, since, for n>=1, the values of a,b,c are given by Euclid's formula for generating Pythagorean triples: a=2xy, b=x^2-y^2, c=x^2+y^2 with x=2^(2n) and y=2^(2n-1)+1 and x and y are coprime and x is even and y is odd. - _Chai Wah Wu_, Feb 13 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert C. Lyons_, Feb 12 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017026 | OEIS-A381044

Primes prime(k) followed by a gap, prime(k+1)-prime(k), smaller than the local geometric average gap between consecutive primes: log(prime(k))/e^(gamma).

It has been conjectured that primes are distributed around their average spacing in a Poisson distribution (cf. D. A. Goldston and A. H. Ledoan). This is the basis of the conjecture that, for k tending to infinity, the asymptotic limit of the average of log(prime(k+1)-prime(k)) is log(log(prime(k))) - gamma (where gamma is Euler's constant). Also, the geometric mean of the gap between consecutive primes [p(k+1)-p(k)] is equivalent to log(prime(k)) / e^gamma.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Apr 14 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017027 | OEIS-A381054

a(n) is the least k such that floor(sqrt(n*k/d(n*k))) - floor(sqrt(d(n*k))) = 1, where d(k) is the largest divisor of k which is <= sqrt(k).

If we denote for each m the corresponding sequence a_m(n), it is conjectured that (a_m(n)=(A000196(n)-m)^2 iff n is a prime > m^2+1) the cases m=0:A382286,1 are answered affirmatively Conjecture: a(n) < (A000196(n)-1)^2 iff n>=9 is a composite number. Conjecture: a(n) = (A000196(n)-1)^2 iff n>=9 is a prime number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hassan Baloui_, Apr 14 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017028 | OEIS-A381055

a(n) = -n/2 if n is even, 3n + 1 if n is odd.

Conjecture: Iterating the map x -> a(x) on any negative or positive integer eventually reaches 1. Some iteration paths between -35 and 35 are illustrated in the plot of a(n) in LINKS.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Apr 14 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017029 | OEIS-A381169

List of twin prime averages (A014574) is partitioned by including as many elements as possible in the n-th partition, L_n, such that any gap in L_n is smaller than the gap between L_n and L_(n-1) but not bigger than the first gap in L_n. a(n) is the number of elements in L_n.

Conjecture 1: lim_{n->oo} N_i/n = k_i, where N_i is the number of partitions with i elements and k_i is a constant, with k_2 > k_1 > k_3 > k_4 > .... The values of k_i are the same as those in A348168. Conjecture 2: lim_{n->oo} Sum_{1..n} a(n)/n = lim_{i->oo} Sum_{1..i} i*k_i = e, or the average partition length approaches 2.71828... as n tends to infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Feb 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017030 | OEIS-A381190

Number of connected minimal dominating sets in the n-trapezohedral graph.

Conjectured g.f.: -2 * x^3 * (4*x^5 + 8*x^4 + 4*x^3 - 9*x^2 - 8*x - 3) / (x^3 + x^2 - 1)^2. - _Joerg Arndt_, Jan 07 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Feb 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017031 | OEIS-A381228

Smallest k such that n appears in row k of the triangle in A381227, or -1 if n never appears in A381227.

Conjecture: Every positive integer appears in A381227.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 25 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017032 | OEIS-A381238

a(n) = floor(b(n)), where b(n) = sec^2(b(n-1)), b(0)=1.

Conjecture: Every nonnegative number appears in this sequence. (See A382148.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 17 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017033 | OEIS-A381240

Indices of records in A381096.

Conjecture 1: For i > 1, A002110(i) is in this sequence. Conjecture 2: Intersection with A001694 (i.e., in A286708) is {900, 1800}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Feb 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017034 | OEIS-A381242

Lexicographically earliest sequence of distinct terms > 1 such that no term is a substring of the product of any two terms.

Conjecture: The digit 9 never appears.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dominic McCarty_, Mar 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017035 | OEIS-A381265

a(n) is the number of solid (3D) partitions of n with 2 layers and second layer a plane partition of 3.

Conjecture: equal to 3*(2*A000219 -A000990 -2*A000041 +1) tested up to n=20. G.f.: 3*(2*Product_{k>0} 1/(1-x^k)^k -(1-x)*Product_{k>0} 1/(1-x^k)^2 - 2*Product_{k>0} 1/(1-x^k) + 1/(1 - x)) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_, Feb 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017036 | OEIS-A381266

a(n) = least positive integer m such that when m*(m+1) is written in base n, it contains every single digit exactly once, or 0 if no such number exists.

Conjecture: a(n) = 0 if and only if n==3 (mod 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Mondot_ and _Ali Sada_, Feb 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017037 | OEIS-A381361

E.g.f. satisfies A(x) = exp(Integral abs(1/A(x)^2) dx), where abs(F(x)) equals the series expansion formed by the unsigned coefficients in F(x).

Conjecture: for n > 4, a(n) == 0 (mod 5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 26 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017038 | OEIS-A381362

G.f. A(x) satisfies 1/2 = Sum_{n=-oo..+oo} x^n*A(x)^n * (A(x)^n + x)^(2*n-1) * (x^n + A(x))^(2*n-1).

Conjecture: for n > 0, a(n) == 2 (mod 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 21 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017039 | OEIS-A381363

G.f. A(x) satisfies 1/2 = Sum_{n=-oo..+oo} x^n*A(x)^n * (A(x)^n + x)^(3*n-1) * (x^n + A(x))^(3*n-1).

Conjecture: for n > 0, a(n) == 2 (mod 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 21 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017040 | OEIS-A381364

G.f. A(x) satisfies 1/3 = Sum_{n=-oo..+oo} x^n*A(x)^n * (A(x)^n + 2*x)^(n-1) * (x^n + 2*A(x))^(n-1).

Conjecture: for n > 0, a(n) == 6 (mod 9).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 21 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017041 | OEIS-A381365

G.f. A(x) satisfies 1/3 = Sum_{n=-oo..+oo} x^n*A(x)^n * (A(x)^n + 2*x)^(2*n-1) * (x^n + 2*A(x))^(2*n-1).

Conjecture: for n > 0, a(n) == 6 (mod 9).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 21 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017042 | OEIS-A381420

a(1) = 1, a(2) = 3; for n > 2, a(n) is the smallest unused positive number that shares a factor with a(n-1) and has a different binary weight than a(n-1).

Unlike the EKG sequence A064413 the primes do not appear in their natural order. The only fixed points in the terms studied are 1 and 20, and it is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Feb 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017043 | OEIS-A381451

Triangle read by rows: T(n,k) is the clique covering number of the Johnson graph J(n, k), n >= 2, 0 < k < n.

T(2*k, k) = C(k) = A000108(k), the k-th Catalan number, for k = 1, 2, 4, 6, 8, 16; whether this holds for other values of k is an open question.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Søren Fuglede Jørgensen_, Feb 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017044 | OEIS-A381457

Integers encoding the recursive structure of a bitonic sorter network of n elements in their binary expansion.

A bitonic sorting network for n = 2^k elements requires (n/2)*(log_2(n))*(log_2(n) + 1) comparators (Batcher, 1968). This sequence extends the construction to arbitrary n by padding to the next power of 2 and pruning redundant comparators. The validity of this generalization as a sorting network remains conjectural; proofs are sought.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Darío Clavijo_, Mar 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017045 | OEIS-A381458

Primes p such that p/prevprime(p) < 1 + (1/PrimePi(p)).

Conjecture: Limit_{n->oo} n / PrimePi(a(n)) = 3/5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Feb 28 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017046 | OEIS-A381460

Smallest n-th perfect power greater than 1 satisfying A373387(a(n)) = n.

It is conjectured that if n > 2 is given, then a(n) is generated by {5^2^k}_oo or -{5^2^k}_oo (this probabilistic argument is based on the study of a(n) up to n = 50 since a(50) is a 762 digit number generated by {5^2^k}_oo = ...6259918212890625).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Feb 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017047 | OEIS-A381464

Lexicographically earliest positive integer sequence satisfying a(n) = a(a(n))/n.

While extending the sequence at a(k) we will check if k equals a previous term in the sequence. If such a term a(m) = k is found a(k) is determined as a(k) = a(m)*m. If no previous term matches k we may choose a(k) = k+c with the least c such that c > 0 and k+c does not equal any previous term in the sequence. It is conjectured that this check is sufficient. Reasoning behind this conjecture: The greatest common divisor of two consecutive Fibonacci numbers is 1, thus we know that (k-1)^F(m)*k^F(m+1) and (t-1)^F(n)*t^F(n+1) are all different for some m,n > 1 if k and t are chosen such that for m or n < 2 no solution for (k-1)^F(m)*k^F(m+1) = (t-1)^F(n)*t^F(n+1) exist, because this cannot be equal if (k-1)*k and (t-1)*t have different prime numbers as divisors and if the only difference is the exponent of the prime factors, then the distribution of these between (t-1) and t and thus their progression F(n) or F(n+1) is individually distinct. In this sequence we need also to consider the more general case (k-c)^F(m)*k^F(m+1) = (t-1)^F(n)*t^F(n+1) because sometimes we need to set a(k) = k+c. It is conjectured that in this case c is bounded to be < 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Feb 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017048 | OEIS-A381494

Smallest number with reciprocal of period length n in base 7.

Conjecture: a(n) = A218358(n) for n>=2. - _R. J. Mathar_, Mar 03 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Erich Friedman_, Feb 25 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017049 | OEIS-A381616

a(n) is the smallest prime that starts the first occurrence of exactly n consecutive primes in A381019.

Conjecture: a(n) always exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gonzalo Martínez_, Mar 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017050 | OEIS-A381636

Numbers whose prime indices cannot be partitioned into constant blocks with distinct sums.

Conjecture: These are the zeros of A382876.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Mar 10 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017051 | OEIS-A381658

Lexicographically earliest sequence of positive integers such that for each distinct positive integer t there is only one value of k such that t = a(n) = a(n+k) = a(n+2*k).

It is conjectured that the values of n for which a(n) = 1 is given by A092482.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Mar 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017052 | OEIS-A381670

The function A(x) = x+(1/2)*x^2-(1/16)*x^4... = Sum_{k >= 0} x^k*A381669(k)/a(k) satisfies the functional equation: x*(A(x)+1) = A(A(x)).

Conjecture: All terms are powers of two.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Mar 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017053 | OEIS-A381706

Three-dimensional array of the number b(n, k, i) of permutations of k chosen numbers in {1,2,...,n} with i-1 descents, n >= 1, 1 <= k <= n, 1 <= i <= n.

The recurrence b(n,k,i) = b(n,k-1,i) + (n choose k) Sum_{i=d-n+k}^{d-1} b(k-1,k-2,j) was conjectured by Ron Fertig and proved by Alex R. Miller.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Timothy Y. Chow_, Mar 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017054 | OEIS-A381707

Smallest initial value for unimodal Collatz (3x+1)/2 glide sequence that begins with exactly n increases.

a(n) mod 2^(n+2) = A283507(n+1) (conjectured). - _V. Barbera_, Apr 20 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Dewan_, Mar 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017055 | OEIS-A381717

Number of integer partitions of n that cannot be partitioned into constant multisets with distinct block-sums.

Conjecture: Also the number of integer partitions of n having no permutation with all distinct run-sums, ranked by zeros of A382876. In other words, a partition has a permutation with all distinct run-sums iff it has a multiset partition into constant blocks with all distinct block-sums, where the run-sums of a sequence are obtained by splitting it into maximal runs and taking their sums.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Mar 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017056 | OEIS-A381737

Orders k of Hermite polynomials whose maximal coefficient in absolute value appears twice.

Conjecture 1: a(n) = 2*n*(n + 2) + (n + 1)*(-1)^(n+1). Conjecture 2: a(n) = a(n-1) + 2*a(n-2) - 2*a(n-3) - a(n-4) + a(n-5). Conjecture 3: G.f.: (8*x + 5*x^2 + 5*x^3 - x^4 - x^5) / ((1 - x)^3 * (1 + x)^2). Terms < 20000 consistent with conjectures. - _Jinyuan Wang_, Mar 09 2025.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mike Sheppard_, Mar 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017057 | OEIS-A381762

Numbers k such that S(k) sets a new record, where S(k) denotes the sum of the reciprocals of odd elements in the Collatz sequence which starts at k.

This sequence is conjectured to be finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Barak Manos_, Mar 06 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017058 | OEIS-A381850

Primes p preceded and followed by primes whose difference is less than 2*log(p).

Conjecture: Limit_{n->oo} n / PrimePi(a(n)) = 1-(3/e^2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, May 06 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017059 | OEIS-A381888

Triangle read by rows: T(n, k) = (n + 1) * Sum_{j=k..n} binomial(n, j) * Eulerian1(j, j - k).

Consider A381706, the number of permutations of k chosen numbers in [n] with i-1 descents, as a sequence of squares of size 1x1, 2x2, 3x3, ..., as displayed in the example section of A381706. Conjecture: T(n, k) is the sum of column k+1 of the (n+1)th square; in other words: T(n, k) = Sum_{j=0..n} b(n+1, j+1, k+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Mar 11 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017060 | OEIS-A381927

Least positive integer k which has at least n divisors d for which tau(k) = sigma(d).

Conjecture: This sequence is infinite. Conjecture: a(n) has exactly n divisors d for which tau(a(n)) = sigma(d) or a(n+1) > a(n) respectively. If this conjecture is true, this sequence is a subsequence of A381925.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Mar 12 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017061 | OEIS-A381932

Triangular array T(n, k) read by rows: denominators of the coefficients for the iterated exponential F^{r}(x) = x + Sum_{n>=1} x^(n+1)*Sum_{k=1..n} r^(n+1-k)*T(n, k)/A381931(n, k) with F^{1}(x) = exp(x)-1 and F^{2}(x) = exp(exp(x)-1)-1.

Conjecture: abs(T(n, 2)) = A064169(n - 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Mar 12 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017062 | OEIS-A381949

a(n) is the smallest integer k greater than 1 and not a perfect power satisfying A373387(k^n) = n.

It is conjectured that a(n) is a multiple of 5 for any n > 2 (probabilistic argument).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Mar 10 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017063 | OEIS-A382050

a(n) = least positive integer m such that when m*(m+1) is written in base n, it is zeroless and contains every single nonzero digit exactly once, or 0 if no such number exists.

Conjecture: a(n) > 0 if n > 5 and n <> 3 (mod 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Mar 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017064 | OEIS-A382054

a(n) = least positive integer m such that when m*(m+1) is written in base n, it does not contain the digit n-1 and contains every single digit from 0 to n-2 exactly once, or 0 if no such number exists.

Conjecture: a(n) > 0 if n > 4 and n <> 3 (mod 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Mar 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017065 | OEIS-A382055

a(n) = least positive integer m such that when m*(m+1) is written in base n, it does not contain the digits 0 or n-1 and contains every single digit from 1 to n-2 exactly once, or 0 if no such number exists.

Conjecture: a(n) = 0 if and only if n == 3 (mod 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Mar 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017066 | OEIS-A382074

a(n) is the number of solutions to phi(x) + phi(n-x) = phi(n) where 1 <= x <= floor(n/2).

a(2*i) = 0 is not true for every positive odd i. For example, a(2*3) = 1. It is conjectured that a(2*A065381(n)) = 0 for n > 1. However, there are positive odd numbers i not in A065381 and for which a(2*i) = 0. For example, a(2*529) = a(2*1155) = 0. Conjecture: a(2*A065381(n)) = 0 for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Mar 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017067 | OEIS-A382082

F(k) such that F(k) + (F(k) reversed) is a palindrome, where F(k) is a Fibonacci number.

Conjecture: The sequence appears to be finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vincenzo Librandi_, Mar 21 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017068 | OEIS-A382091

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest unused positive number such that a(n) shares a factor with a(n-1) while the total number of prime terms of the form 4*k + 1 is never less than those of the form 4*k + 3.

In the first 100000 terms the fixed points are 1, 2, 77, 121; it is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Mar 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017069 | OEIS-A382121

Minimal polynomials of nimbers *(2^(2^n)-1), evaluated at 2.

Each of these polynomials from n=1 up to n=12 is primitive: if you make a finite field of order 2^(2^n) as GF(2)[x]/<P(x)> then x generates the field's multiplicative group. A natural conjecture is that this is true for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simon Tatham_, Mar 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017070 | OEIS-A382122

G.f. satisfies Sum_{n>=0} x^n * abs(1/A(x)^n) = C(x), where C(x) = 1 + x*C(x)^2 and abs(F(x)) equals the series expansion formed by the unsigned coefficients in F(x).

Conjecture: for n > 0, a(n) is odd iff n = 2^k for k >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017071 | OEIS-A382125

G.f. A(x) = exp(Sum_{n>=1} sigma(n)*sigma(2*n) * x^n/n), where sigma(n) = A000203(n) is the sum of the divisors of n.

Conjectures: a(3*n) == A382124(n) (mod 3) for n >= 0; a(3*n+1) == 0 (mod 3) and a(3*n+2) == 0 (mod 3) for n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 06 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017072 | OEIS-A382131

Complement of A381767.

a(n) ~ e^(n-1) (conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mike Sheppard_, Mar 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017073 | OEIS-A382237

Numbers that are not divisible by the sum of any subset of their digits.

Conjecture: after a sufficiently large n this sequence grows faster than the prime numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sergio Pimentel_, Mar 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017074 | OEIS-A382261

a(n) = floor(x^(phi^n)), where phi = (1+sqrt(5))/2 and x is the constant A382260.

Conjecture: All terms are prime numbers. For details see A382260.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Mar 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017075 | OEIS-A382285

Initial members of prime octuplets (p, p+4, p+12, p+24, p+28, p+40, p+48, p+52), where all primes are consecutive primes.

It is conjectured that there is an infinite number of primes for every admissible k-tuple.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Federico Salas_, Mar 20 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017076 | OEIS-A382296

Number of states in smallest DFAO computing t(i+n) on input n in base 2, msd-first, where t(n) = A010060(n), the Thue-Morse sequence.

This conjecture is proved. See Biswas link. - _Ranadeep Biswas_, Aug 02 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Mar 21 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017077 | OEIS-A382307

Position of start of first run of alternating bit values in the base-2 representation of Pi, or -1 if no such run exists.

Conjecture: no term is -1 (would follow from a proof that Pi is normal in base 2). - _Sean A. Irvine_, Mar 30 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James S. DeArmon_, Mar 21 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017078 | OEIS-A382316

G.f. satisfies A(x) = A(x^2) + A(x^2)^2*A(x^3)/A(x^6), with A(0) = 0 and A'(0) = 1.

Conjecture: there are no negative terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 08 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017079 | OEIS-A382318

G.f. A(x) satisfies A(x) = x + (Sum_{n>=1} A(x^n))^3.

Conjectures: a(3*n) == A382319(n) (mod 3) for n >= 0; a(3*n+1) == 0 (mod 3) and a(3*n+2) == 0 (mod 3) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 10 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017080 | OEIS-A382319

G.f. A(x) satisfies A(x) = x/(1-x) + Sum_{n>=1} A(x^n)^3.

Conjecture: a(n) == A382318(3*n) (mod 3) for n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 10 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017081 | OEIS-A382357

Lexicographically earliest sequence of distinct positive integers such that the 2-adic valuations of adjacent terms differ exactly by one.

Conjecture: this sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Mar 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017082 | OEIS-A382374

Lexicographically earliest sequence of distinct positive integers such that the number of prime factors counted with multiplicity of adjacent terms differ exactly by one.

Conjecture: this sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Mar 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017083 | OEIS-A382376

Lexicographically earliest sequence of distinct positive integers such that the number of distinct prime factors of adjacent terms differ exactly by one.

Conjecture: this sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Mar 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017084 | OEIS-A382407

a(n) is the number of partitions n = x + y + z of positive integers such that x*y + y*z + x*z is a perfect square.

Conjecture: a(k) = 0 iff k is an element of {2, 4, 8, 13} union A000244 union A005030.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Apr 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017085 | OEIS-A382427

Number of integer partitions of n that can be partitioned into constant blocks with distinct sums.

Conjecture: Also the number of integer partitions of n having a permutation with all distinct run-sums.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Mar 26 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017086 | OEIS-A382437

a(n) = a(n-1)^2 + 4 * a(n-1), with a(0) = 2.

Conjecture: a(n) = Sum_{k=1..2^n} (2^n * 2^k * binomial(2^n + k - 1, 2*k - 1) / k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _V. Barbera_, Mar 25 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017087 | OEIS-A382438

Numbers k in A024619 such that all residues r (mod k) in row k of A381801 are such that rad(r) divides k, where rad = A007947.

Conjecture: 12 and 24 are the only nonsquarefree numbers in this sequence, i.e., in A126706.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Mar 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017088 | OEIS-A382464

Positive integers that contain an even digit d immediately preceded by a digit >= d.

Conjecture: these are the numbers missing from A382462.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Mar 28 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017089 | OEIS-A382465

Positive integers such that every even digit except the first is immediately preceded by a smaller digit.

Conjecture: these are the terms of A382462, sorted.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Mar 28 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017090 | OEIS-A382518

Let N = A001481(n), the n-th number that is the sum of two nonnegative squares. a(n) is the index of the first lattice-edge sequence that will accept N so that no sequence contains the edges of a triangle, otherwise if no such sequence exists, a(n) = 0.

Conjecture 1: bin #1 contains the orthogonal and 45-degree diagonal lattice edges. Conjecture 2: After chessboard coloring the lattice, bin #3 contains only lattice edges that connect black and white points.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gordon Hamilton_, Mar 29 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017091 | OEIS-A382617

Numbers k such that k = m*(m^2 + 1) where m^2 + 1 is prime.

Numbers k that are the nontrivial solutions to A070304 for primes p = m^2 + 1. Conjecture: for such primes p, there are exactly two solutions, k = 0 and the nontrivial k = m * p. These k correspond to the terms of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Lee Benjamin_, Mar 31 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017092 | OEIS-A382624

Positive integers such that every even digit except the leftmost is immediately preceded by a larger digit.

Conjecture: these are the terms of A382621, sorted.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Apr 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017093 | OEIS-A382632

Numbers k such that one can make an equilateral triangle from a chain of linked rods of length 1, 2, 3, ..., k, with perimeter equal to the total length.

Conjecture: Out of some linked rods of length 1, 2...k, we can fold them in half (digon), or make equilateral triangles, but no other regular polygons (squares, regular pentagons, etc...) can be made.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Mondot_, Apr 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017094 | OEIS-A382655

Lexicographically earliest sequence of distinct positive integers such that if m = a(n-1), a(n) is the smallest novel number k such that: g = gcd(m, k) > 1, m/g > 1, k/g >1 and gcd(m/g, k) = gcd(k/g, m) = 1.

The relation described in Name between two distinct numbers m, k refers to what may be described as "symmetric non-coprimality" (expressed as m # k), which in this sequence requires that if a(n-1) = m is a term, then a(n) = k is the smallest novel number such that m # k. From the definition it follows that 1, primes and prime powers cannot be terms in this sequence, however powerful numbers which are not prime powers can be terms (see Example). Sequence is conjectured to be a permutation of A024619. For two numbers m, k symmetrically related in this way we may write (m # r) = 1, and if the relation is not symmetric we may write (m # r) > 1. a(n) --> n as n-->00.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Apr 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017095 | OEIS-A382681

Conjecturally, the numbers k (not multiples of 5) such that for all x >= 0, k*2^x has a '0' in its decimal expansion.

Conjecturally, the numbers k (not multiples of 5) such that for all x >= 0, k*2^x has a '0' in its decimal expansion.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Brian Kehrig_, Jun 02 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017096 | OEIS-A382715

The term in A377091 that immediately precedes n, or 0 if n does not appear in A377091.

It is a strong conjecture that every integer appears in A377091, so it is unlikely there will be a second 0 term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 09 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017097 | OEIS-A382814

Number of nachos that the first player gets when playing the "Fibonachos" game starting with n nachos.

Conjecture: Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Apr 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017098 | OEIS-A382815

Positive numbers k such that abs((sin(k))^k) sets a new record.

Conjecture: every term is either the numerator of an odd-denominator convergent (A096456) or an odd-denominator semiconvergent of Pi/2. Of the first 15 terms, 12 are convergents and 3 are semiconvergents: 8, 51464 and 860103093554. - _Jwalin Bhatt_, Aug 08 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Apr 28 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017099 | OEIS-A382837

Numbers k such that k - A071324(k) > A000010(k).

This sequence lists the counterexamples to Atanassov's conjecture that n - A071324(n) <= A000010(n) for all n. (See Atanassov link, page 2). Conjecture: For each even natural number n, infinitely many pairs (k,k+n) exist, such that both k and k+n are terms in the sequence. (Example - n = 876 has pairs (84, 960), (180, 1056), (264, 1140), (300, 1176)...)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shreyansh Jaiswal_, Apr 06 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017100 | OEIS-A382868

a(1) = 1, a(2) = 2. For n > 2 a(n) is the smallest novel number divisible by the smallest prime p which divides a(n-1) but does not divide a(n-2). If no such prime exists a(n) is the least novel k such that gcd(k, a(n-1)) > 1.

Similar to A064413, from which it departs at a(19) = 25. Any odd prime p is preceded by a multiple of itself. Conjectured to be a permutation of the natural numbers, with primes occurring in natural order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, Apr 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017101 | OEIS-A382943

Numbers k such that A382883(k) = 0.

Conjecture: A059404 is a subsequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Apr 12 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017102 | OEIS-A382946

a(n) is the least positive integer k having a proper divisor d such that the base n expansions of k and d, without leading zeros, have, up to order, the same digits, or a(n) = -1 if no such k exists.

Conjecture: a(n) > 0 for any n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Apr 09 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017103 | OEIS-A383016

Numbers k such that A382883(k) = 1.

Conjecture: A053810 is a subsequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Apr 12 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017104 | OEIS-A383061

Positive integers that contain an odd digit d immediately preceded by a digit >= d.

Conjecture: these are the numbers missing from A383059.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Apr 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017105 | OEIS-A383062

Nonnegative integers such that every odd digit except the leftmost is immediately preceded by a smaller digit.

Conjecture: these are the terms of A383059, sorted.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Apr 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017106 | OEIS-A383093

Number of integer partitions of n that can be partitioned into constant blocks with a common sum.

Conjecture: We have Sum_{d|n} a(d) = A323774(n), so this is the Moebius transform of A323774.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Apr 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017107 | OEIS-A383111

Number of integer partitions of n having more than one permutation with all distinct run-lengths.

A partition is Look-and-Say exactly when its distinct-part multiplicities m_i admit pairwise-disjoint strict partitions P_i of m_i (A239455). A valid word gives such a family from its run lengths. Conversely, choose such a family minimizing K = Sum_i |P_i|. If P_i = {r_1 < ... < r_k} and k > 1 + (K-k), then K <= 2k-2. The k values r_1,...,r_k and the k-1 sums r_j+r_k (1 <= j < k) are 2k-1 distinct integers because each sum exceeds r_k and the sums differ with j. Exactly K values are used globally and all r_1,...,r_k are used, so some sum r_j+r_k is unused. Replacing r_j,r_k by this sum preserves m_i, strictness and disjointness but lowers K, a contradiction. Thus |P_i| <= 1 + Sum_{h != i}|P_h| for every i. By the standard gap criterion, the block labels can be ordered with unequal neighbors; assigning these lengths yields a valid word. This proves the conjectured converse in A239455.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Apr 20 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017108 | OEIS-A383115

a(n) is the number of values m that satisfy floor(sqrt(m)) = n and A381054(m) = 1.

Conjecture: Sum_{n=1..N} a(n) ~ 4*N^(3/2)/3 for N large enough. Conjecture: Sum_{n=1..N} a(n) ~ 2*Sum_{n=1..N} A382306(n) for N large enough.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hassan Baloui_, Apr 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017109 | OEIS-A383131

a(n) is the number of iterations that n requires to reach 1 under the map x -> -x/2 if x is even, 3x + 1 if x is odd; a(n) = -1 if 1 is never reached.

Conjecture: a(n) != -1 for any positive or negative integer n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Apr 17 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017110 | OEIS-A383135

a(n) = number of iterations that n requires to reach 1 under the x -> A380891(x) map, or -1 if it never does.

The map x -> A380891(x) is conjectured to eventually reach 1 for all starting x > 0. Tested for x <= 7892481.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James C. McMahon_ and _Vikram Prasad_, Apr 17 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017111 | OEIS-A383165

Expansion of e.g.f. log(1 + (exp(2*x) - 1)/2)^2 / 2.

Conjecture: the sequence {a(n)} reduced modulo a positive integer k is eventually periodic. For example, modulo 5 the sequence becomes [0, 0, 1, 3, 3, 0, 0, 2, 3, 2, 0, 1, 3, 4, 0, 0, 3, 1, 0, 4, 3, 3, 0, 3, 3, 0, 0, 2, 3, 2, 0, 1, 3, 4, 0, 0, 3, 1, 0, 4, 3, 3, 0, ...], with an apparent period of 20 starting at n = 3. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Apr 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017112 | OEIS-A383166

Expansion of e.g.f. log(1 + (exp(2*x) - 1)/2)^3 / 6.

Conjecture: the sequence {a(n)} reduced modulo a positive integer k is eventually periodic. For example, modulo 5 the sequence becomes [0, 0, 0, 1, 1, 0, 0, 0, 3, 2, 0, 0, 1, 4, 3, 0, 0, 1, 4, 0, 0, 3, 3, 0, 1, 0, 0, 0, 3, 2, 0, 0, 1, 4, 3, 0, 0, 1, 4, 0, 0, 3, 3, 0, ...], with an apparent period of 20 starting at n = 4. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Apr 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017113 | OEIS-A383229

Indices of record low-water marks of the sequence abs((sin(k))^k).

Conjecture: Every forward difference of this sequence is a positive integer multiple of the numerator of either a convergent (A002485/A002486) or a semiconvergent to Pi.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Apr 28 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017114 | OEIS-A383245

Nonnegative integers that contain the digit 0, or an even digit d immediately followed by a digit >= d.

Conjecture: these are the numbers missing from A342043.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Apr 20 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017115 | OEIS-A383246

Positive integers without the digit 0 such that every even digit except the rightmost is immediately followed by a smaller digit.

Conjecture: these are the terms of A342043, sorted.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Apr 20 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017116 | OEIS-A383247

Positive integers that contain the digit 9, or an odd digit d immediately followed by a digit <= d.

Conjecture: these are the numbers missing from A342044.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Apr 25 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017117 | OEIS-A383248

Nonnegative integers without the digit 9 such that every odd digit except the rightmost is immediately followed by a larger digit.

Conjecture: these are the terms of A342044, sorted.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Apr 25 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017118 | OEIS-A383249

Positive integers ending with the digit 1, or containing an odd digit d immediately followed by a digit >= d.

Conjecture: these are the numbers missing from A342045.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Apr 26 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017119 | OEIS-A383250

Nonnegative integers not ending with the digit 1 and such that every odd digit except the rightmost is immediately followed by a smaller digit.

Conjecture: these are the terms of A342045, sorted.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Apr 26 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017120 | OEIS-A383257

Let p = prime(n), then a(n) is the non-p-smooth part of (p-1)!+1.

Let p = prime(n) and k = (p-1)!+1. If m<p, then k == 1 mod m, so a(n) = k/(p-adic valuation of k). By Wilson's theorem, a(n)<=k/p. Conjecture: a(n) = k/p^2 if n = 3, 6, or 103 and a(n) = k/p otherwise. - _Chai Wah Wu_, May 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mike Jones_, Apr 29 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017121 | OEIS-A383274

a(n) = Sum_{i,j = 0..n} C(n, i)^2*C(n, j)^2*C(i+j, i)*2^(i+j).

Conjecture 1: Let S(p) = Sum_{0<k<p} a(k) with p a prime. Conjecture 2: For any prime p == 1,3,7,9 (mod 20), we have Sum_{0<k<p}(8k+5)*a(k) == 14/5*p (mod p^2). Both conjectures have been verified for odd primes smaller than 1000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Apr 26 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017122 | OEIS-A383284

Lexicographically earliest infinite sequence such that a(i) = a(j) => A265576(i) = A265576(j), for all i, j >= 1, where A265576 is the LCM-transform of EKG-sequence.

a(n) = 1+A000720(A265576(n)). [Conjectured. True if there are no composites in A265576]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Apr 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017123 | OEIS-A383307

a(n) is number of n-digit nonnegative integers whose difference between the largest and smallest digits is equal to the mode of its digits.

Conjecture: lim_{n->oo} a(n+1)/a(n) = 10. - _Stefano Spezia_, Apr 26 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Apr 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017124 | OEIS-A383368

Number of intercalates in pine Latin squares of order 2n.

Hypothesis: For all known pine Latin squares of orders N=4k+2 number of intercalates a(n) = a(N/2)= a(2k+1) = (N/2)^3 = (2k+1)^3 = A016755((n-1)/2) (verified for N<29). Hypothesis: For all known pine Latin squares of orders N=4k number of intercalates a(n) = a(N/2) = a(2k) = (N/2)^2 + (N/2)^3 = 4*k^2 + 8*k^3 = (2k)^2 * (2k+1) = 2*A089207(n/2) = 4*A099721(n/2) (verified for N<29).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eduard I. Vatutin_, Apr 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017125 | OEIS-A383377

G.f. satisfies A(x) = Sum_{n>=0} x^n * abs(1/A(x)^n), where abs(F(x)) equals the series expansion formed by the unsigned coefficients in F(x).

Conjecture: a(n) is even for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017126 | OEIS-A383398

a(n) is the smallest number whose sum with any previous term is abundant.

I conjecture that all the terms are odd integers ending in 1 or 9. The odd nature of the terms seems particularly likely, as the sum of a(n) that's even with any previous term would need to be an odd abundant number (see A005231).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jakub Buczak_, Apr 25 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017127 | OEIS-A383410

Array read by downward antidiagonals: A(n,k) = Sum_{i=0..n-1} Sum_{j=0..k+1} binomial(n-1,i)*binomial(k+1,j)*A(i,j) with A(0,k) = 1, n >= 0, k >= 0.

Conjecture: A(n,0) = A005649(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Apr 26 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017128 | OEIS-A383423

Indices k such that A003849(k) = 0 and A383422(k) = 0.

Conjecture: {a(n) - a(n-1), n>=2} = {1, 3, 4, 5, 7}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017129 | OEIS-A383424

Indices k such that A003849(k) = 1 and A383422(k) = 1.

Conjecture: {a(n) - a(n-1), n>=2} = {2, 3, 5, 7, 8}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017130 | OEIS-A383425

Indices k such that A003849(k) = 0 and A383422(k) = 1.

Conjecture: {a(n) - a(n-1), n>=2} = {1, 2, 3, 4, 5}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017131 | OEIS-A383426

Indices k such that A003849(k) = 1 and A383422(k) = 0.

Conjecture: {a(n) - a(n-1), n>=2} = {3, 7, 8, 10}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017132 | OEIS-A383427

Indices k such that A003849(k) = A383422(k).

Conjecture: {a(n) - a(n-1), n>=2} = {1, 2, 3, 5, 6}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Apr 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017133 | OEIS-A383440

a(n) = (5*n + 3)!/((8*n^2 + 10*n + 3)*(n!)^2*(3*n + 2)!).

a(n) = A383453(2*n, n), conjectured by Wildberger-Rubine to be the main diagonal of the Geode Bi-Tri array G[m_2, m_3].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, May 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017134 | OEIS-A383442

a(0) = 0; thereafter a(n) is the least integer (in absolute value) not yet in the sequence such that the absolute difference between a(n-1) and a(n) is a triangular number; in case of a tie, preference is given to the positive value.

The present sequence is the Heraclitus transform of the triangular numbers A000217. For the squares, see A377091. Conjecture: both H(A000217) and H(A000290) contain every (positive or negative) integer. In fact it appears that this property holds whenever S is a monotonically strictly increasing sequence starting with 1. It does not hold for H(A000012), which is A001477.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017135 | OEIS-A383445

Index of n in A377090, or -1 if n does not appear there.

It is conjectured (see A377090) that every positive integer appears exactly once either here or in A383446.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Apr 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017136 | OEIS-A383456

a(n) = 2*binomial(2*n+4,n)*(n+3)/(n+2).

Conjectured to be column 1 of the array in A383453.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 02 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017137 | OEIS-A383466

a(0) = 1; thereafter a(n) = 10*n^2 - 5*n + 2.

Conjecture 2: a(n) is the maximum number of regions that can be formed in the plane by drawing n regular pentagrams with any radii and any centers. Adamczewski suggests a machine proof of Conjecture 2 in linked article. - _Michael De Vlieger_, Aug 16 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Jul 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017138 | OEIS-A383500

Positive integers that contain the digit 9, or an odd digit d immediately preceded by a digit <= d.

Conjecture: these are the numbers missing from A382935.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Apr 29 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017139 | OEIS-A383501

Nonnegative integers without the digit 9 such that every odd digit except the leftmost is immediately preceded by a larger digit.

Conjecture: these are the terms of A382935, sorted.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Apr 29 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017140 | OEIS-A383505

Least integer k >= 0 such that binomial(k*n,k+1) = -1 mod n, or -1 if no such integer exists.

Conjecture: a(A000430(n)) = A000430(n)-1. It is not hard to see that conjecture is true. - _Max Alekseyev_, Jul 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Bard_, May 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017141 | OEIS-A383540

Positive numbers k such that (sin k)^k sets a new record.

Conjecture: let p/q be a convergent (A096456/A096463) or a semiconvergent to Pi/2 with q odd. Then every term is either: p, when p is even or q == 1 (mod 4); or 3p, when p is odd and q == 3 (mod 4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Apr 29 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017142 | OEIS-A383541

Positive numbers k such that (cos k)^k sets a new record.

Conjecture: let p/q be a convergent (A002485/A002486) or a semiconvergent to Pi. Then every term is either: p, when p and q have opposite parity; or 2p, when p and q are both odd. Conjecture: a(n) = A002485(n+7) for n >= 9. - _Jakub Buczak_, May 05 2025 [This conjecture is false since a(14) != A002485(21). - _Jwalin Bhatt_, Aug 09 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Apr 29 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017143 | OEIS-A383570

Number of transversals in pine Latin squares of order 4n.

Hypothesis: number of transversals in pine Latin squares of all orders N=4k+2 is zero (verified for orders N<=18).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eduard I. Vatutin_, Apr 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017144 | OEIS-A383578

Let p = prime(n), then a(n) is the p-smooth part of (p-1)! + 1.

Let p = prime(n). If m<p, then (p-1)!+1 == 1 mod m, so a(n) = p-adic valuation of (p-1)!+1. By Wilson's theorem, a(n)>=p. Conjecture: a(n) = p^2 if n = 3, 6 or 103 and a(n) = p otherwise. - _Chai Wah Wu_, May 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mike Jones_, Apr 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017145 | OEIS-A383579

a(n) is the largest number t such that there exist numbers i,j,k such that, for all m <= t, there exist integers x,y,z with x*i + y*j + z*k = m and |x|+|y|+|z| <= n.

For n > 5, a(n) = max_{m=0..(n-1)} 4n^2*m - 12n*m^2 + 8m^3 + 3n^2 - 11n*m + 10m^2 - n + 3m (conjectured). - for n > 5, i(n) = 2n-2m-1 (conjectured); - for n > 5, j(n) = 4n*m-4m^2+3n-5m-2 (conjectured); - for n > 5, k(n) = 4n*m-4m^2+3n-5m-1 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zachary Barbanell_, Apr 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017146 | OEIS-A383580

a(n) is the largest number t such that there exist numbers i,j,k,l such that, for all m <= t, there exist integers x,y,z,w with x*i + y*j + z*k + w*l = m and |x|+|y|+|z|+|w| <= n.

For n > 3, a(n) = max_{m=0..(n-1)} 2n^2*m^2 - 4n*m^3 + 2m^4 + 4n^2*m - 6n*m^2 + 2m^3 + 2n^2 - 2n*m + m^2 + n (conjectured). - for n > 3, i(n) = 1 (conjectured); - for n > 3, j(n) = 2n-2m+1 (conjectured); - for n > 3, k(n) = 2n^2*m-4n*m^2+2m^3+2n^2-2n*m+m (conjectured); - for n > 3, l(n) = 2n^2*m-4n*m^2+2m^3+4n^2-6n*m+2m^2+2n-m+1 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zachary Barbanell_, May 14 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017147 | OEIS-A383586

a(n) is the minimum sum of a nonnegative integer 4-tuple that takes n moves to reach a 0 component, where a move picks two components, subtracts the smaller from the larger, and doubles the smaller.

Conjecture: the last sum S for which the diameter is d, noted (d,S), are (0,3), (1,9), (2,19), (3,39), (4,86), (5,204), (6,472), (7,1180), (8,2876), (9,6880) and (10,17496). Example: (1,9) means that for S>9, d>1. - _Karl Desfontaines_, Dec 14 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gerold Jager_, May 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017148 | OEIS-A383595

a(n) is the smallest prime k such that (prime(n), k, u, v) are the vertices of a square in Ulam's spiral, where k < u < v are all primes; or -1 if there is no such k.

Conjecture: if a prime is vertex of a square of prime vertices, then it is vertex of infinitely many squares whose vertices are prime. For example, in the case of 11, some of them are: (11, 59, 127, 131), (11, 137, 233, 239), (11, 769, 977, 991).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gonzalo Martínez_, May 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017149 | OEIS-A383619

Conjectured list of least elements of nontrivial arithmetic derivative orbits.

Conjectured list of least elements of nontrivial arithmetic derivative orbits. The values of a(n) are conjectural, contingent on the absence of further merges or the existence of nontrivial cycles beyond the computational horizon; this is analogous to Collatz dynamics.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimitris Cardaris_, May 02 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017150 | OEIS-A383650

Averages k of a twin prime pair such that 3*k*2^d is also the average of a twin prime pair for some divisor d of 3*k.

a(n) ~ b*n^c for some constants b and c as n tends to infinity (conjectured). - _Jakub Buczak_, May 04 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, May 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017151 | OEIS-A383652

Primes p preceded and followed by gaps whose product is less than (log(p))^2.

Conjecture: Limit_{n->oo} n / PrimePi(a(n)) = 0.720268...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, May 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017152 | OEIS-A383700

Coefficient of x^2 in expansion of (x+1) * (x+5) * ... * (x+4*n-3).

Conjecture D-finite with recurrence a(n) +3*(-4*n+7)*a(n-1) +(48*n^2-216*n+247)*a(n-2) -(4*n-11)^3*a(n-3)=0. - _R. J. Mathar_, May 07 2025 Conjecture verified using the differential equation y + (124*x - 31)*y' + (240*x^2 - 120*x + 15)*y'' + (64*x^3 - 48*x^2 + 12*x - 1)*y''' = 0 satisfied by the e.g.f. - _Robert Israel_, Mar 12 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, May 06 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017153 | OEIS-A383701

Coefficient of x^3 in expansion of (x+1) * (x+5) * ... * (x+4*n-3).

Conjecture D-finite with recurrence a(n) +4*(-4*n+9)*a(n-1) +2*(48*n^2-264*n+371)*a(n-2) -4*(4*n-13)*(16*n^2-104*n+173)*a(n-3) +(4*n-15)^4*a(n-4)=0. - _R. J. Mathar_, May 07 2025 Conjecture confirmed using differential equation y + (624*x - 156)*y' + (2656*x^2 - 1328*x + 166)*y'' + (1792*x^3 - 1344*x^2 + 336*x - 28)*y''' + (256*x^4 - 256*x^3 + 96*x^2 - 16*x + 1)*y'''' satisfied by the e.g.f. - _Robert

Israel_, Mar 12 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, May 06 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017154 | OEIS-A383702

Coefficient of x^2 in expansion of (x+3) * (x+7) * ... * (x+4*n-1).

Conjecture D-finite with recurrence a(n) +3*(-4*n+5)*a(n-1) +(48*n^2-168*n+151)*a(n-2) -(4*n-9)^3*a(n-3)=0. - _R. J. Mathar_, May 07 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, May 06 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017155 | OEIS-A383721

a(n) is the number of distinct rectangles with integer area that can be inscribed in a cube with edge length 4*n, as shown in the linked figure "Cube with inscribed rectangle".

Conjecture: There is no rectangle with an integer area if x is not a positive integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, May 08 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017156 | OEIS-A383730

a(0) = 0, a(n) = a(n-1) + A002260(n) * (-1)^(n-1) if not already in the sequence, otherwise a(n) = a(n-1) - A002260(n) * (-1)^(n-1).

Conjecture: the sequence is unbounded both above and below. Conjecture: each integer appears finitely often.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Markel Zubia_, May 06 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017157 | OEIS-A383777

a(n) is the number of steps that n requires to reach 0 under the map: x -> 2*x + 1 if x is even; 0 if x = 1; x - lpf(x) otherwise where lpf(x) is the least prime factor of x. a(n) = -1 if 0 is never reached.

Conjecture: a(n) != -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, May 17 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017158 | OEIS-A383780

a(n) is the number of n-digit terms in A383779.

Conjecture: A383779 is finite, since after the first two terms here, the ratios between successive odd terms and the ratios between successive even terms are quickly decreasing. - _Michael S. Branicky_, May 11 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, May 09 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017159 | OEIS-A383844

a(n) is the number of occurrences of n in A024934.

- It is likely that every n appears at least once in A383327, however this is currently conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Miles Englezou_, May 12 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017160 | OEIS-A383856

Dimension in which a random vector of length n has the highest probability to fall into a single hypercube with side length of 10.

Conjecture: a(11)..a(16) are 5, 6, 8, 9, 11, 12 with at least 99.7% confidence (based on numerical simulation discussed below).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ruediger Jehn_, May 12 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017161 | OEIS-A383877

a(n) is the smallest integer k such that the Diophantine equation x^3 + y^3 + z^3 + w^3 = k^3, where 0 < x < y < z < w has exactly n integer solutions, or 0 if there is no such k.

Conjecture: a(n) != 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhining Yang_, May 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017162 | OEIS-A383907

Echo primes: primes p such that the greatest prime factor of p-1 is a suffix of p.

Conjecture: There are infinitely many echo primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giorgos Kalogeropoulos_, May 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017163 | OEIS-A383922

a(n) = A002104(n) + A002104(n+1) - 1.

Junyan Xu gives a conjecture on the recursive formula of m(x), where x + m(x) is the least fusible greater than x. (A188545(n) is -log_2 m(n)). We have m(x) = m_0(x) for x < 33/16 (i.e., F_0 and F coincide on the interval (-oo,33/16]), but they differ for x >= 33/16, even if the intersection of F and (-oo,n) still has order type omega^^n if the conjecture is true. In fact, we have -log_2 m_0(3) = 1541023937, while -log_2 m(3) > 2^^^^^^^^^16 in Knuth's up-arrow notation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, May 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017164 | OEIS-A383941

a(0) = 0; for n >= 1, a(n) is the smallest positive integer not already in the sequence that is coprime to both a(n-1) and the sum of all preceding terms.

Conjecture: This is a permutation of the nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Sep 09 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017165 | OEIS-A383977

Sequence of successive merge positions when Fibonacci-sorting an infinite list.

Conjecture: a(n) is obtained by reading the binary number on the right column of page 10 in the Arndt reference, interpreted as Fibonacci base, reading from bottom to top; see example. - _Joerg Arndt_, Jan 31 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lucilla Blessing_, May 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017166 | OEIS-A384132

Integers k such that the Diophantine equation x^3 + y^3 + z^3 + w^3 = k^3, where 0 < x < y < z < w has no integer solutions.

Conjecture: a(27)=121 is the largest integer whose cube cannot be described as the sum of four distinct positive cubes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhining Yang_, May 20 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017167 | OEIS-A384146

Smallest squarefree order m > 0 for which there are n nonisomorphic finite groups of order m, or 0 if no such order exists.

It is conjectured that 0 never appears in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robin Jones_, May 21 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017168 | OEIS-A384149

Irregular triangle T(n, k) in which row n gives the 2-densely-aggregated composition of sigma(n).

From empirical evidence of the first 1000 rows, reinforced by some known related properties, it looks readily credible that if we take the average of row n and its reversal we get A237270 row n, a palindromic composition of sigma(n) that was defined using Dyck paths. Row lengths are therefore conjectured to be in the database as A237271. T(n,k) = A384226(n,k) * A038712(n) = A384226(n,k) * A000203(n) / A000593(n). (conjectured). - _Omar E. Pol_, Apr 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Munn_, May 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017169 | OEIS-A384162

Number of length n words over an n-ary alphabet such that a single letter in every run of letters is marked.

Conjecture: a(n) = n * A342168(n-1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _John Tyler Rascoe_, May 21 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017170 | OEIS-A384184

Order of the permutation of {0,...,n-1} formed by successively swapping elements at i and 2*i mod n, for i = 0,...,n-1.

Conjecture: a(2^n + 2^x) = 2^n * (x-n) if x > n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mia Boudreau_, May 29 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017171 | OEIS-A384193

Slice of elementary triangular automaton rule 210, starting from a lone 1 cell.

At some point in the sequence, the binary digits of a(n) repeat almost exactly every 4096 digits, starting from the 16622nd digit. See illustration of the binary digits of a(65534). I conjecture that exact periodicity is to be found later in the sequence, with a period that is a multiple of 12288.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul Cousin_, May 21 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017172 | OEIS-A384228

Numbers k such that 128 * 3^k - 1 is prime.

Conjecture: This sequence intersects with A388408 at k = 2 to form twin primes with center N = 2^7 * 3^2 = 1152 = A027856(9). Any such intersection has to be at an odd k because if k is even, N-1 will be the difference of two squares, so not prime for N > 4. A covering system can be constructed that eliminates all other intersections (see linked program), and the search up to k = 250000 makes the probability of another intersection vanishingly small.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ken Clements_, Sep 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017173 | OEIS-A384233

Square array read by upward antidiagonals: T(n,k) is the n-th number whose largest odd noncomposite divisor is its k-th divisor, n >= 1, k >= 1.

Conjecture: T(n,2) = A061345(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, May 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017174 | OEIS-A384267

a(n) = abs([x^n] 1 + x/A(x)^2), where A(x) is the g.f.

Conjectures: (2) A(x) = 1 + x * (2/A(w*x)^2 + 2/A(w^2*x)^2 - 1/A(x)^2)/3, where w = exp(i*2*Pi/3) = -1/2 + sqrt(3)/2*i (conjecture); this is implied by conjecture (C.2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017175 | OEIS-A384270

G.f. satisfies A(x) = A(x^4 + 4*x*A(x)^4) / A(x^3 + 3*x*A(x)^3).

Conjecture: a(n) is odd iff n = 2^k for k >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 10 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017176 | OEIS-A384275

a(1) = 1, a(2) = 2, a(3) = 4; for n > 3, a(n) is the smallest unused positive number that shares a factor with a(n-1) and at least one other previous term.

In the first 100000 terms the only fixed points are 1, 2, 9; it is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017177 | OEIS-A384309

a(1) = 1. Thereafter a(n) is the cardinality of the set of terms whose leading decimal digit is the same as that of a(n-1).

Conjecture: All positive integers appear precisely 9 times in this sequence, except for 1, which appears 10 times. For k >= 1, the last k digit term in the sequence is a(23[k-1]93) = [k]9 where "[m]9" means a run of m nines; see Example.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, May 25 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017178 | OEIS-A384318

Number of strict integer partitions of n that are not maximally refined.

Conjecture: Also the number of strict integer partitions of n such that it is possible in more than one way to choose a disjoint family of strict integer partitions, one of each part.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, May 28 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017179 | OEIS-A384336

a(1) = 1, a(2) = 2. For n > 2, a(n) = number of a(k), k = 1..n-2 such that a(k) divides a(n-1).

1 appears 3 times and 2 appears 2 times, all other numbers are conjectured to appear infinitely many times. The first occurrence of an odd prime is followed by 3, see Example, and the order of first appearances of primes starts: 2,5,3,7,11,19,23,13.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, May 28 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017180 | OEIS-A384350

Number of subsets of {1..n} containing at least one element that is a sum of distinct non-elements.

Conjecture: Also the number of subsets of {1..n} such that it is possible in more than one way to choose a disjoint family of strict integer partitions, one of each element.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jun 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017181 | OEIS-A384490

Prime numbers p such that both roots of x^2 - x - 1 modulo p are primitive roots modulo p.

Conjecture: these numbers are all in A106963. - _Michael Somos_, Oct 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jay Anderson_, May 31 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017182 | OEIS-A384503

Lexicographically earliest infinite sequence of distinct positive integers having the property that for any pair a(n-2) = i, a(n-1) = j of consecutive terms > 1, a(n) is the smallest novel k such that gcd(i,k) > 1 if gcd(i,j) = 1 or gcd(j,k) = 1 if gcd(i,j) > 1.

Conjectured to be a permutation of the positive integers, with primes occurring in order.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David James Sycamore_, May 31 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017183 | OEIS-A384531

Multiplicative sequence a(n) with a(p^e) = ((2*e+1) * p - 2*e) * p^(e-1) for prime p and e >= 0.

Conjecture: a(n) = Sum_{i=1..n} gcd(i, n) * gcd(i+1, n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Jun 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017184 | OEIS-A384537

Composite numbers that are equal to the concatenation of the primes and exponents in their prime factorizations in some bases.

Conjecture: a composite number can be a Davis number to at most one base.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Jun 02 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017185 | OEIS-A384539

Zeroless positive integers k for which for every pair of nonempty substrings that concatenate to give k one substring divides the other.

It is conjectured that this sequence is finite and contains 132 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Jun 09 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017186 | OEIS-A384603

Primes preceded and followed by gaps whose quotient (value greater or equal to 1) is less than 2.

Conjecture: Limit_{n->oo} n / PrimePi(a(n)) = 1/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Jun 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017187 | OEIS-A384618

Primes preceded and followed by gaps whose quotient (value greater or equal to 1) is greater than 2.

Conjecture: Limit_{n->oo} n / PrimePi(a(n)) = 2/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Jun 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017188 | OEIS-A384669

Positive integers setting a new record for the sum of the square roots of the prime exponents.

We provide facts, conjectures, and questions about the class of sequences A_x. Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hal M. Switkay_, Jun 06 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017189 | OEIS-A384698

The first prime number reached by iterating the map, x -> 2*x + 1 if x is even; x - lpf(x) otherwise where lpf(x) is the least prime factor of x, on n >= 2; or -1 if a prime is never reached.

Conjecture: a(n) != -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Jun 09 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017190 | OEIS-A384700

On a 2 X n grid of vertices, draw a circle through every unordered triple of non-collinear vertices: a(n) is the number of distinct circles created.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Jun 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017191 | OEIS-A384704

Triangle T(i, j), 1 <= j <= i, read by rows. T(i, j) is the smallest number k that has i odd divisors and whose symmetric representation of sigma, SRS(k), has j parts; when no such k exists then T(i, j) = -1.

The single value T(10, 4) = -1 has been verified; see the conjecture below. Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Jun 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017192 | OEIS-A384713

The number of steps that n requires to reach 1 under the map: x-> x^2 - 1 if x is an odd prime, x/2 if x is even, x - lpf(x) otherwise where lpf(x) is the least prime factor of x. a(n) = -1 if 1 is never reached.

Conjecture: a(n) != -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Jun 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017193 | OEIS-A384728

The number of different shuffle square roots of the prefix of length 2n of the infinite word 00110011001100...

Conjecture: There are infinitely many natural numbers n such that, among all shuffle squares of length 2n, the prefix of the word 001100110011... has the greatest number of distinct shuffle square roots.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bartlomiej Pawlik_, Jun 08 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017194 | OEIS-A384819

Nonnegative numbers a(n) < n for n >= 1 such that exp(Sum_{n>=1} (n^2 - a(n))*x^n/n) is a power series with integral coefficients.

Conjecture: a(p^n) = p - 1 when p is prime for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017195 | OEIS-A384830

G.f. satisfies A(x) = A(x^3 - 3*x*A(x)^3) / A(x^2 - 2*x*A(x)^2).

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 11 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017196 | OEIS-A384874

a(n) is the first prime encountered when iterating the map x -> x/2 if x is even, x*lpf(x) + 1 otherwise, where lpf(x) is the least prime factor of x, on n >= 2; or -1 if a prime is never reached.

Conjecture: a(n) != -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Jun 11 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017197 | OEIS-A384928

Number of 2-dense sublists of divisors of the n-th triangular number.

Conjecture: all odd indexed terms are odd. For odd n = 2k + 1 the n-th triangular number t(n) = (2k+1)*(k+1) satisfies A003056(t(n)) = 2k + 1; i.e., n is an odd divisor of t(n). Therefore SRS(t(n)) at the diagonal is positive and SRS(t(n)) has an odd number of parts. Since the number of parts equals the number of 2-dense sublists (see link in A384149), the conjecture is true. - _Hartmut F. W. Hoft_, Mar 25 2026 a(n) = A237271(A000217(n)) for n >= 1 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 08 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017198 | OEIS-A384930

Irregular triangle read by rows: T(n,k) is the sum of the terms of the (n-k+1)-th 2-dense sublist of divisors of n, with n >= 1, k >= 1.

Note that if the conjectures related to the 2-dense sublists of divisors of n are true so we have that essentially all sequences where the words "part" or "parts" are mentioned having cf. A237593 are also related to the 2-dense sublists of divisors of n, for example the square array A240062. T(n,k) = 2*A237270(n,k) - A384149(n,k), n >= 1, k >= 1, (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jul 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017199 | OEIS-A384931

Number of 2-dense sublists of divisors of the number of partitions of n.

a(n) = A237271(A000041(n)). Conjectured.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Jul 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017200 | OEIS-A385077

Primes x that cannot be written as x = p + 2*q where p < q are primes.

Conjectured to be finite (see also conjectures in A046927, A337924). Note that Lemoine representations (no restriction on order of p, q) hold for all odd numbers up to 10^13 (see Juhász et al.).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ian Shannon-Garvey_, Jul 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017201 | OEIS-A385093

Primes preceded and followed by gaps whose quotient (value greater or equal to 1) is greater than 2 and less than 3.

Conjecture: Limit_{n->oo} n / PrimePi(a(n)) = 1/6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Jun 17 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017202 | OEIS-A385156

The number of undirected, simple, unlabeled graphs G on n vertices which are prime, not split, and do not contain a vertex of degree 1 in G or in the complement of G, and has no induced P5 in G or in the complement of G.

Here, "prime" means with respect to modular decomposition (see link). A P5 is a path on 5 vertices. A split graph is a graph whose vertices can be partitioned into a clique and an independent set. Related to conjectures in the referenced paper.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jim Nastos_ and _Clara Elliott_, Jul 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017203 | OEIS-A385175

Cubes using at most three distinct digits, not ending in 0.

Conjecture: a(26) = 3163316636166336 is the largest term with nonzero digits (See comments of A030294 and the data of A155146, where a(26) = A155146(47)^3).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gonzalo Martínez_, Jun 20 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017204 | OEIS-A385192

Primes p such that multiplicative order of 5 modulo p is odd.

Conjecture: this sequence has density 1/3 among the primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Jun 20 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017205 | OEIS-A385194

Minimum number of steps to reach n by repeated doubling and/or digit reversal starting from 1, or -1 if n cannot be reached.

Conjecture: a(n) > -1 for all n not divisible by 3 or 11.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Radcliffe_, Jun 20 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017206 | OEIS-A385220

Primes p such that multiplicative order of 3 modulo p is odd.

Conjecture: this sequence has density 1/3 among the primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Jun 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017207 | OEIS-A385221

Primes p such that multiplicative order of 4 modulo p is odd.

Conjecture: this sequence has density 7/12 among the primes (see A014663).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Jun 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017208 | OEIS-A385252

Number of ternary strings of length 2*n that have at least one 0 but less 0's than the combined number of 1's and 2's.

Conjecture D-finite with recurrence n*a(n) +(-37*n+36)*a(n-1) +4*(131*n-245)*a(n-2) +16*(-221*n+605)*a(n-3) +192*(59*n-213)*a(n-4) +6912*(-2*n+9)*a(n-5)=0. - _R. J. Mathar_, Jul 31 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Enrique Navarrete_, Jul 28 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017209 | OEIS-A385330

The point numbers encountered by a rotating marker following the process described in the Comments.

Conjecture: The only time a label is replaced is laying down the second point of the 2-star which overwrites the first (and only point) from the 1-star.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tamas Sandor Nagy_, Jun 25 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017210 | OEIS-A385409

a(n) is the smallest positive integer k such that the Diophantine equation x^3 + y^3 + z^3 + w^3 = k^2, where 0 < x < y < z < w has exactly n integer solutions.

Conjecture: a(n) exists for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhining Yang_, Jun 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017211 | OEIS-A385431

Leading digit of the decimal expansion of the prime zeta function at n.

It is conjectured that a(n) = A111395(n) for all n >= 10 (see "Is the leading digit of the decimal expansion of the prime zeta function at n equal to the first digit of 5^n, for all integers n >= 10?" in Links). For all n > 9, a(n) = most significant (nonzero) digit of 5^n (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Jun 28 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017212 | OEIS-A385478

a(n) is the smallest divisor of sigma(n) that is not element of the union of the sets of divisors of sigma(k) for 1 <= k < n; or 0 if no such divisor exists.

Conjecture: The divisors counted in A385467(n) are multiples of a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Jul 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017213 | OEIS-A385525

Consider the graph whose vertices are the points of the n-dimensional cubic lattice with points connected by all integer-length diagonals that traverse all n dimensions and do not intersect intermediate points. a(n) is the total length of the shortest possible closed walk in this graph via noncongruent diagonals of the same length.

Other than a(2) whose walk is comprised of 8 diagonal segments, all known terms are produced by 3- or 4-segment walks, including some with examples of both. It is conjectured that this holds true for all n >= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles L. Hohn_, Jul 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017214 | OEIS-A385580

a(n) is the number of ways to partition an n X n X n cube into four noncongruent cuboids of different volumes.

Conjecture: a(p) = A384311(p) for any prime p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Janaka Rodrigo_, Aug 12 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017215 | OEIS-A385590

Triangle read by rows, based on Fibonacci numbers: Let i > 1 be such that F(i) <= n < F(i+1); i.e., i = A130233(n). Then T(n, k) = F(i-1)^2 + 1 - (i-1) mod 2 + (n - F(i)) * F(i-2) + (k-1) * F(i-1) where F(k) = A000045(k).

Conjecture: This triangle yields a permutation of the natural numbers. Conjecture: Sum_{k=1..n} (-1)^k * binomial(n-1, k-1) * T(n, k) = 0 for n > 2 and (-1)^n for n < 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Werner Schulte_, Jul 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017216 | OEIS-A385613

Number of steps that n requires to reach 0 under the map: x-> x^2 - 1 if x is an odd prime, floor(x/3) if x is even, otherwise x - 1. a(n) = -1 if 0 is never reached.

Conjecture: a(n) = -1 occurs only when starting number n runs into a loop.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ya-Ping Lu_, Jul 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017217 | OEIS-A385658

Least prime p < 2n*(n+1) such that the polynomial Sum_{k=1..n} tau(k)*x^(n-k) is irreducible modulo p, or 1 if such p does not exist, where tau is Ramanujan's tau function given by A000594.

Conjecture: a(n) > 1 for all n > 1. In other words, for each n = 2,3,... the polynomial x^(n-1) + tau(2)*x^(n-2) + ... + tau(n) is irreducible modulo some prime p < 2n*(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017218 | OEIS-A385676

Least prime p <= 2*n^2 - n + 1 such that the polynomial Sum_{k=1..n} sigma(k) * x^(n-k) is irreducible modulo p, or 1 if such p does not exist, where sigma is given by A000203.

Conjecture: a(n) > 1 except for n = 1, 5. See A385678 for a similar conjecture involving Euler's totient function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017219 | OEIS-A385678

Least prime p <= n^2 - 2*n + 4 such that the polynomial Sum_{k=1..n} phi(k)*x^(n-k) is irreducible modulo p, or 1 if such p does not exist, where phi is Euler's totient function given by A000010.

Conjecture: a(n) > 1 for all n > 5. See also A385658 and A385676 for similar conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017220 | OEIS-A385681

a(n) is the least k > 1 such that n^2 == k (mod sopfr(k)) and k^2 == n (mod sopfr(n)), or -1 if there is no such k, where sopfr = A001414.

a(n) = -1 if n is in A385679. Conjecture: a(n) > 0 if n is not in A385679.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Will Gosnell_ and _Robert Israel_, Aug 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017221 | OEIS-A385714

a(n) is the smallest positive integer k such that the Diophantine equation x^3 + y^3 + z^3 = k^n, where 0 < x < y < z has an integer solution, or -1 if no such integer exists.

Conjecture: if n >= 24 and n == 0 (mod 3), then a(n) = 2. Verified for n <= 69. Conjecture: if n >= 23 and n == 2 (mod 3), then a(n) = 3. Verified for n <= 44. Conjecture: if n >= 7 and n == 1 (mod 3), then a(n) = 6. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-Marc Rebert_, Jul 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017222 | OEIS-A385736

a(n) is the number of distinct nondegenerate triangles with perimeter n whose side lengths are triangular numbers.

0, 1, 6, 10, 28, 55 are the only triangular numbers <= 10^6 that are not perimeters of triangles whose side lengths are triangular numbers. Conjecture: There are no other triangular numbers that have this property.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Jul 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017223 | OEIS-A385812

Numbers k such that A063655(k) > A063655(k+1).

Lai and Reinfeld conjecture that:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard S. Chang_, Jul 09 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017224 | OEIS-A385881

Algebraic rank of elliptic curve y^2 = x^3 - n*x - n.

Terms from n = 43 onward are the analytic ranks (see PARI code) of the corresponding elliptic curves. By the BSD conjecture, these are expected to equal the algebraic ranks. Thus, the validity of these terms is conditional on BSD.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Shreyansh Jaiswal_, Aug 20 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017225 | OEIS-A385912

A diagonal (unsigned) of triangle A385910; a(n) = -A385910(n+2,n) for n >= 1.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 14 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017226 | OEIS-A385970

a(n) is the minimum number of powers needed to obtain the n-th term of sequence A385969.

The sequence is unbounded (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alberto Zanoni_, Jul 14 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017227 | OEIS-A386235

Number of partitions (p, q, r) of n into positive integers such that p + 11*q + 13*r is a perfect square.

Conjecture: a(n) ~ K * n^(3/2) where K = 0.0914... from a(10000) = 91413 and a(20000) = 258667.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hoang Xuan Thanh_, Jul 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017228 | OEIS-A386251

Consider the graph whose vertices are the points of the n-dimensional cubic lattice with points connected by all integer-length diagonals that traverse all n dimensions and do not intersect intermediate points. a(n) is the total length of the shortest possible closed walk in this graph via noncongruent diagonals.

It is provable that all such walks must be even in total length. It is also provable that 3-segment closed walks are impossible for n < 6, and conjectured that a(n) for all n >= 6 are produced by 3-segment walks.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles L. Hohn_, Jul 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017229 | OEIS-A386311

a(1) = 2, a(n+1) is the largest prime p such that b(n+1) = b(n)*(p + a(n))/(p - a(n)) is a positive integer, where b(1) = 1.

Conjecture: a'(n) = prime(n) for "the smallest prime p" and b'(n) = A352743(n-1) for "the largest k".

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Ordowski_, Jul 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017230 | OEIS-A386368

a(n) = Sum_{k=0..n-1} binomial(6*k,k) * binomial(6*n-6*k-2,n-k-1).

Conjecture D-finite with recurrence
48828125*(n-1)*(5*n-4)*(5*n-3) *(432862082629612805*n -769306661967834399) *(5*n-2)*(5*n-1)*a(n)
+1125000*(-40524540611581621957500 0*n^6
+26131807994689105103925 00*n^5
-76671644069685147952125 0*n^4
+13834502135358262506660375*n^3
-16251583347734702117341345*n^2
+11251247074043948959380314*n
-33956990693512417654957 20)*a(n-1)
+33592320*(14228169091832644053750 0*n^6
-12664243385216092720125 00*n^5
+52360412635832716879537 50*n^4
-12786608152035075786775875*n^3
+18838556229131595646260055*n^2
-15323925851720394901667853*n
+52406814069524168121612 36)*a(n-2)
-53444359913472*(6*n-17) *(395547729523405*n
-538181211711288)*(6*n-13) *(3*n-7)*(2*n-5)
*(3*n-8)*a(n-3)=0. - _R. J. Mathar_, Jul 30 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Jul 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017231 | OEIS-A386381

Main diagonal of A386363.

Conjecture 1: if d is a divisor of a(k), then d is also a divisor of a(m*d + k) where m is any natural number. In particular, a(k) is a divisor of a(m*a(k) + k) for any k >= 0 where m is any natural number. Conjecture 2: if d > 2 is a divisor of E(k), then d is also a divisor of E(m*phi(d)+k) where m is any natural number. In particular, E(k) is a divisor of E(m*phi(E(k)) + k) for any k >= 0 (with single exception at k = 3) where m is any natural number. Conjecture: 2*a(n) is divisible by A060818(n). (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Jul 20 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017232 | OEIS-A386424

Numbers k such that sigma(k) has the same powerful part as k has, where sigma is the sum of divisors function.

Conjecture 1: the initial 1 is the only square in this sequence, and a(2) = 2 is the only term that is twice a square. Conjecture 2: A323653 is a subsequence (which would follow from conjecture 1 (c) given there). Conjecture 3: The only common term with A387727 is 1. - _Antti Karttunen_, Sep 12 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 17 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017233 | OEIS-A386425

Odd composites k such that sigma(k) has the same powerful part as k, where sigma is the sum of divisors function.

By definition, the sequence contains all odd perfect numbers, and also includes any hypothetical odd triperfect number that is not a multiple of 3 (see A005820 and A347391), and similarly, any odd term of A046060 that is not a multiple of 5, etc. If there are no squares in this sequence (see conjecture in A386424), then the latter categories of numbers certainly do not exist, and this is then a subsequence of A228058.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 17 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017234 | OEIS-A386430

Odd numbers k such that there are no prime factors p of sigma(k) such that p does not divide A003961(k) and the valuation(k, p) is different from valuation(sigma(k), p), where A003961 is fully multiplicative with a(p) = nextprime(p), and sigma is the sum of divisors function.

Conjecture: After the initial 1, and apart from any hypothetical odd perfect numbers, all other terms are in A248150, i.e., sigma(k) == 0 (mod 4). This would imply (with the same caveat), that this sequence has no common terms with A228058 and no squares larger than one. This is true at least for the first 709203 terms (terms in range [1..2^34]).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017235 | OEIS-A386437

Alternating sum of the numbers in the trajectory of n for the 3x+1 problem.

Conjecture: Let P(n) be the set of distinct positive prime numbers that appear in the first n terms of the sequence. Let Q(n) be the subset of those primes that appear with frequency >= 2. Then, as n -> oo, the ratio |Q(n)|/|P(n)| tends to 11/100. Verified for n <= 5 * 10^7. Conjecture: Let S(n) be the set of positive prime numbers that appear in the first n terms of the sequence. Then, as n -> oo, the ratio |S(n)|/n tends to 0. Verified for n <= 10^6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Luca Santarsiero_, Jul 21 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017236 | OEIS-A386477

a(0) = 1; thereafter a(n) = 2*(6*n^2 - 3*n + 1).

Conjecture 2: a(n) is the maximum number of regions that can be formed in the plane by drawing n regular hexagrams with any radii and any centers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_ and _N. J. A. Sloane_, Jul 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017237 | OEIS-A386482

a(1)=1, a(2)=2; thereafter a(n) is either the greatest number k < a(n-1) not already used such that gcd(k, a(n-1)) > 1, or if no such k exists then a(n) is the smallest number k > a(n-1) not already used such that gcd(k, a(n-1)) > 1.

It is conjectured that every positive number appears.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Aug 15 2025, based on email messages from _Geoffrey Caveney_ | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017238 | OEIS-A386525

a(n) is the least k such that at least n terms of A063655 starting from index k are strictly decreasing.

Lai and Reinfeld conjecture that the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Richard S. Chang_, Jul 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017239 | OEIS-A386538

a(n) is the maximum possible area of a polygon within a circle of radius n, where both the center and the vertices lie on points of a unit square grid.

Conjecture: The maximum possible area of a polygon within the circle would be the same if only the vertices but not the center were fixed on grid points.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Aug 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017240 | OEIS-A386539

Triangle read by rows: T(n,k) is the maximum possible area of a polygon within a ellipse with integer axis n and k, where n >= k >= 1 and both the center and the vertices lie on points of a unit square grid.

Conjecture: The maximum possible area of a polygon within the ellipse would be the same if only the vertices but not the center were fixed on grid points.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Aug 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017241 | OEIS-A386566

a(n) = Sum_{k=0..n-1} binomial(5*k-1,k) * binomial(5*n-5*k,n-k-1).

Conjecture D-finite with recurrence 196608*n*(4*n-3)*(2*n-1)*(18270873280*n -32560150837) *(4*n-1)*a(n) +1280*(-1399185802400000*n^5 +1022280893000000*n^4 +17669158913120000*n^3 -48968110172924750*n^2 +49502057719349955*n -17877514345852392)*a(n-1) +125000*(-61298198200000*n^5 +1447969779032500*n^4 -7721498995066250*n^3 +17474948768595875*n^2 -18352567310653770*n +7399184154389181)*a(n-2) +48828125*(5*n-11) *(5*n-14)*(4958243695*n -6717884799) *(5*n-13)*(5*n-12)*a(n-3)=0. - _R. J. Mathar_, Jul 30 2025 Conjecture verified using differential equation (9558804241406250000*x^2 + 3478346131892625000*x - 65680378505199360)*y(x) + (1926966200765625000000*x^3 + 437226430372419375000*x^2 - 225114481533989760 0*x

- 8428158845779968)*y'(x) + (8668688191845703125000*x^4 + 126872614659839062500 0*x^3 - 55453256429062080000*x^2 + 90623583896076288*x)*y''(x) + (780185938037109375000 0*x^5 + 610975392048593750000*x^4 - 80710539920486400000*x^3 + 996406318397915136*x^2)*y'''(x) + (206469026568603515625 0*x^6 + 27750726879062500000*x^5 - 25555847863040000000*x^4 + 7722276038985646 08*x^3)*y''''(x) + (151313589324951171875*x^7 - 7662274775000000000*x^6 - 17909578270720000 00*x^5 + 114950395322695680*x^4)*y'''''(x) + 15353292436800000*x + 8428158845779968 = 0 satisfied by the G.f. - _Robert Israel_, Nov 09 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Jul 26 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017242 | OEIS-A386580

Number of normal multisets of size n having a permutation with all distinct run lengths.

Conjecture: Also the number of normal multisets of size n having a disjoint family of strict integer partitions, one of each multiplicity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Aug 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017243 | OEIS-A386608

a(n) is the smallest number k for which the smallest part of the symmetric representation of sigma of k, SRS(k), has size n. If no such number exists, a(n) = -1.

Conjecture: Every a(n) is positive. A351903(n) <= a(n), for all n >= 1 if the conjecture is true. Conjecture: For each n >= 1, max(k : n = min(SRS(k))) <= n^2, with equality holding for odd primes and their powers (See the irregular table below; true for n <= 100).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Oct 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017244 | OEIS-A386642

Roll n fair, distinguishable, n-sided dice and sort the dice in increasing order with the lowest on the left, highest on the right. Let m be the maximum value rolled and select the m-th die from the right. Then a(n) is the number of such rolls (over all n^n possible outcomes) where the selected die shows 1.

Let p be the probability that the selected die shows 1; i.e., p = a(n)/n^n. Jameson conjectured (2025) that the limiting value of p is between 0.46 and 0.47. Tim Large conjectured (2025) that the limiting value of p is 1 - exp((1/e) - 1) = 0.4685.... Conjecture: a(n) = Sum_{k=1..n} ((n-k+1)^n - Sum_{j=0..k-1} binomial(n, j)*(n-k)^(n-j) - (n-k)^n + Sum_{j=0..k-1} binomial(n, j)*(n-k-1)^(n-j)) + 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philip Jameson_, Jul 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017245 | OEIS-A386649

Product of first n central trinomial coefficients (A002426) for n > 0 with a(0) = 1.

Conjecture: a(n) = A214589(n) - 2 for n >= 1, where A214589(n) is the number of n X n X n triangular 0..2 arrays with every horizontal row having the same average value.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 08 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017246 | OEIS-A386650

Product of first n quadrinomial coefficients (A005725) for n > 0 with a(0) = 1.

Conjecture: 2*a(n) = A214590(n) - 2 for n >= 1, where A214590(n) is the number of nXnXn triangular 0..3 arrays with every horizontal row having the same average value.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Aug 08 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017247 | OEIS-A386666

G.f. A(x) satisfies: A(x)^2 = A(x^2 + 4*A(x)^3).

Conjecture: a(3*n) (mod 3) == 2*a(3*n+1) (mod 3) == a(3*n+2) (mod 3) == 2*A039966(n) for n > 0, where A039966(n) equals the number of partitions of n into distinct powers of 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017248 | OEIS-A386667

G.f. A(x) satisfies: A(x)^2 = A((x + 4*A(x)^2)^2).

Conjecture: [x^(2^n-2^k)] (x/A(x))^(2^n) = 0 for k = 0..n-2 and n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Sep 08 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017249 | OEIS-A386671

All integers k that can produce a closed walk in a square lattice via noncongruent primitive k-length hypotenuses, in ascending order.

Other series with their own such patterns occur with a series based on (29 * 61 * 89) * p beginning at a(30) = 787205, and another based on (13 * 109 * 229) * p beginning at a(60) = 1622465. It is conjectured that there are infinitely many of such series. It is conjectured that all terms are products of 4 or more Pythagorean primes, including at least 4 different ones, though not all such products produce closed walks. It is also conjectured that all walks have an even count of hypotenuse segments and a minimum of 6 segments, and are bilaterally symmetrical when arranged in order of increasing angle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles L. Hohn_, Jul 28 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017250 | OEIS-A386683

Number of unique n-element sum-distinct sets with minimal largest element.

Conjecture: there will be at least 2 solutions when n = m*(m+1)/2 + 2 (3,5,8,12,17,...) - exactly when A005230(n) = 2*A005230(n-1). In general, if (d(1),...,d(n)) is the difference vector of a set S (elements in decreasing order) that has distinct subset sums, and d(n) = 2*d(n-1), then (d(1),...,d(n-2) + d(n-1),d(n-1),d(n-1)) will also produce a valid solution (confirmed for ~10^7 solutions for n = 4 to 8, and for every entry in A005230).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ven Popov_, Nov 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017251 | OEIS-A386760

Numbers k such that the number of decimal digits of the Lucas number L(k) is greater than the number of decimal digits of the Fibonacci number F(k).

The difference in the number of decimal digits, A055642(L(k))-A055642(F(k)) = A060384(k)-A386758(k) is either zero or one. In fact, this difference is ceiling(beta-{k*alpha}), with alpha and beta as defined in the Formula section. This implies that, asymptotically, a fraction of beta=0.349485... of the Lucas numbers has one more decimal digit than the corresponding Fibonacci number. This gives the asymptotic behavior of the sequence as a(n)~n/beta. Conjecture: abs(a(n)-n/beta)<c, for some constant c.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hans J. H. Tuenter_, Aug 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017252 | OEIS-A386827

Least prime n < p < 2*n*(n-1) such that the polynomial Sum_{k=1..n} x^(n-k)/k is irreducible modulo p, or 1 if such a prime p does not exist.

Conjecture: a(n) > 1 for all n > 1. In other words, for each n = 2,3,... there is a prime p with n < p < 2*n*(n-1) such that the polynomial Sum_{k=1..n} x^(n-k)/k is irreducible modulo p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017253 | OEIS-A386828

Least prime n < p <= 2*n^2 + 1 such that the polynomial Sum_{k=1..n} x^(n-k)/k^2 is irreducible modulo p, or 1 if such a prime p does not exist.

Conjecture: a(n) > 1 for all n > 1. We also have similar conjectures for Sum_{k=1..n} x^(n-k)/k^s with other values of s.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017254 | OEIS-A386830

a(n) = Sum_{k=0..n} 3^k * 2^(n-k) * binomial(3*n+1,k) * binomial(3*n-k,n-k).

Conjecture D-finite with recurrence +8*n*(2*n-3)*a(n) +6*(-108*n^2+207*n-80)*a(n-1) +405*(3*n-2)*(3*n-4)*a(n-2)=0. - _R. J. Mathar_, Aug 19 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Aug 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017255 | OEIS-A386832

Triangle read by rows: T(n,k) is the number of the sets of partitions of n with length k grouped by largest part.

Row sums are equal to (n^2 + 3*n)/2 - Sum_{k=1..n} ceiling(n/k) which is conjectured to be equal to A309097: "Number of partitions of n avoiding the partition (4,2,1)." (cf. Geoffrey Caveney link). Conjecture is proven true by Kevin Ryde (see A309097).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wouter Meeussen_, Sep 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017256 | OEIS-A386834

a(n) = Sum_{k=0..n} binomial(4*n+1,k) * binomial(4*n-k-1,n-k).

Conjecture D-finite with recurrence +3*n*(16543753*n -26995933)*(3*n-1)*(3*n-2)*a(n) +(-1669899251*n^4 -26931977989*n^3 +131963667975*n^2 -188283072995*n +85757456660)*a(n-1) +2*(-61301926003*n^4 +515926265010*n^3 -1655392333929*n^2 +2311146075302*n -1165379619540)*a(n-2) -96*(39221117*n -50949760)*(4*n-9)*(2*n-5)*(4*n-7)*a(n-3)=0. - _R. J. Mathar_, Aug 19 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Aug 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017257 | OEIS-A386840

Number of fundamental one-dimensional discrete statistical models with rational maximum likelihood estimator supported on n states and of degree 2n-6.

Unlike A143107 and A143108 (and conjecturally A143109), there are infinitely many polynomials in H(2,d) of degree 2n-6. Nevertheless, this sequence consists of finite numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carlos Améndola_, Aug 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017258 | OEIS-A386850

Least prime n < p <= (n-1)*(2n-1) such that Sum_{k=1..n} x^(n-k)/k! is irreducible modulo p, or 1 if such a prime p does not exist.

Conjecture: a(n) > 1 for all n > 1. In other words, for any integer n > 1, there is a prime p with n < p <= (n-1)*(2n-1) such that the polynomial Sum_{k=1..n}x^(n-k)/k! is irreducible modulo p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Aug 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017259 | OEIS-A386885

a(n) is the number of iterations that -n requires to reach 1 under the map x -> -x/2 if x is even, 3x + 1 if x is odd.

Conjecture: Every nonzero integer reaches 1. Verified numerically up to +-5*10^7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ryosuke Miyazawa_, Aug 06 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017260 | OEIS-A386894

Markoff numbers that are powers of one odd prime or twice powers of one odd prime.

These Markoff numbers (see A002559) have been proved to obey the Frobenius-Markoff uniqueness conjecture. See Aigner, Corollary 3.20, p. 59, and there the references [4] A. Baragar, [18] J. O. Button, and [119] Ying Zhang.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Aug 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017261 | OEIS-A386921

Numerators of the partial sums of 1/d(prime(k)+1), where d is the number of divisors function.

a(n)/A386922(n) <= 4 * K * n/log(n)^(3/2) + O(n*log(log(n))/log(n)^(5/2)), where K = (1/sqrt(Pi)) * Product_{p prime} sqrt(p/(p-1)) * (p * log(p/(p-1)) - 1/(p-1)) = 0.25320111501639846923... (Iudelevich, 2022). Gabdullin et al. (2023) conjectured that a(n)/A386922(n) ~ K * n/log(n)^(3/2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Aug 08 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017262 | OEIS-A386963

Gaps between positions of odd terms in A065090.

Conjecture: a(n) tends to 2 in frequency (i.e., {n : a(n) = 2} has natural density 1). Conjecture is true because the primes have natural density 0. - _Robert Israel_, Aug 29 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aied Sulaiman_, Aug 11 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017263 | OEIS-A386970

Period of Langton's ant on an n X 3 torus.

Conjecture: for n>=2, a(3*n+1) = 96*n+16.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Carlos Moya Garcia_, Aug 11 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017264 | OEIS-A386984

Number of 2-dense sublists of divisors of the n-th hexagonal number.

Conjecture: all terms are odd. Since the hexagonal numbers are the odd-numbered triangular numbers, the proof in A384928 proves this conjecture. - _Hartmut F. W. Hoft_, Mar 25 2026 a(n) = A237271(A000384(n)) for n >= 1 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 11 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017265 | OEIS-A386989

Irregular triangle read by rows: T(n,k) is the product of terms in the k-th 2-dense sublist of divisors of n, with n >= 1, k >= 1.

It is conjectured that row lengths are given by A237271.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 12 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017266 | OEIS-A386992

Irregular triangle read by rows: T(n,k) is the number of nonprimes in the k-th 2-dense sublist of divisors of n, with n >= 1, k >= 1.

It is conjectured that row lengths are given by A237271.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017267 | OEIS-A386993

Number of 2-dense sublists of divisors of the n-th squarefree number.

a(n) = A237271(A005117(n)). (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017268 | OEIS-A386994

Number of 2-dense sublists of divisors of the n-th Fibonacci number.

a(n) = A237271(A000045(n)), n >= 1. (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017269 | OEIS-A387030

Irregular triangle read by rows: T(n,k) is the number of primes in the k-th 2-dense sublist of divisors of n, with n >= 1, k >= 1.

It is conjectured that row lengths are given by A237271.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Aug 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017270 | OEIS-A387031

All integers k that can produce a closed walk in an equilateral triangular lattice via noncongruent primitive k-length diagonals, in ascending order.

All observed terms are products of 4 different primes that are 1 mod 6 (A002476), though not all such products produce closed walks. It is conjectured that all terms are products of 4 or more such primes, including at least 4 different ones.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles L. Hohn_, Aug 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017271 | OEIS-A387043

Number of ways to write n as (1 + (n mod 2))*p + prime(k) + prime(k+1) + prime(k+2), where p is a prime and k is a positive integer.

Conjecture: (i) a(2*n) > 0 for all n > 12. In other words, any even number greater than 25 is the sum of a prime and three consecutive primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017272 | OEIS-A387045

Nonnegative numbers k for which there is no largest circle enclosing exactly k points (strictly inside) on a square grid, only a least upper bound on that circle size.

Conjecture: This sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianqiang Zhao_, Aug 14 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017273 | OEIS-A387048

Numbers k such that k^2 + sopfr(k)^2 is prime, where sopfr = A001414.

Includes 2*p where p is a prime such that 5 * p^2 + 4 * p + 4 is prime. The Generalized Bunyakowsky Conjecture implies there are infinitely many of these.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Aug 14 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017274 | OEIS-A387049

Numbers k such that both k + sopfr(k) and k^2 + sopfr(k)^2 are prime, where sopfr = A001414.

Includes 2*p if p is a prime such that 3*p + 2 and 5 * p^2 + 4 * p + 4 are prime. The Generalized Bunyakowsky Conjecture implies there are infinitely many of these.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Aug 14 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017275 | OEIS-A387052

Decimal expansion of -x, where x is the abscissa of the second local extremum of the Riemann zeta function on the negative real axis.

It is an open question whether the fractional part of x_n tends to 1 or some unknown constant c < 1 as n tends to infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Aug 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017276 | OEIS-A387060

Numbers k such that 16 * 3^k + 1 is prime.

Conjecture: The only intersection with A385115 is at k = 3 where 2^4 * 3^3 = 432 = A027856(8).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ken Clements_, Aug 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017277 | OEIS-A387064

Total number of entries in rows 0 to n of Pascal's triangle multiple of n.

a(p) = p-1, a(p^2) = p*(p-1) for p prime. Conjecture: a(p^k) = (p-1)*p^(k-1) for p prime. - _Chai Wah Wu_, Aug 21 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-Marc Rebert_, Aug 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017278 | OEIS-A387084

Expansion of B(x)/sqrt(1 + 4*(B(x)-1)/5), where B(x) is the g.f. of A001449.

Conjecture D-finite with recurrence 3902464*n*(8*n-5) *(8*n-3)*(8*n-1) *(8*n+1)*a(n) +80*(-12565760000*n^5 +68448000000*n^4 -163457516000*n^3 +200475354000*n^2 -122843089511*n +29804717943)*a(n-1) +125000*(134055000*n^5 -1109795000*n^4 +3726971625*n^3 -6307124125*n^2 +5325821766*n -1769460798)*a(n-2) +48828125*(-1556875*n^5 +15845625*n^4 -60659875*n^3 +103818375*n^2 -67764178*n +1391424)*a(n-3) -152587890625 *(5*n-16)*(n-3) *(5*n-19)*(5*n-18) *(5*n-17)*a(n-4)=0. - _R. J. Mathar_, Aug 19 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Aug 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017279 | OEIS-A387098

Triangle read by rows: T(n,k) is the number of ways to place k non-attacking kings in each row and column of an n X n board, 0 <= k <= floor(n/4) + [n=1].

Conjecture: T(n,k) = 0 for n < 4k, except for T(1,1)=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hamidreza Maleki Tirabadi_, Oct 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017280 | OEIS-A387112

Numbers with (strictly) choosable initial intervals of prime indices.

Conjecture: Also numbers whose weakly increasing prime indices y satisfy y(i) >= i for every index i. The conjecture above is true. Reordering the intervals does not affect the existence of a system of distinct representatives, so let y(1) <= ... <= y(z) be the weakly increasing rearrangement of the x_i. Then the intervals {1,...,y(i)} admit distinct representatives if and only if y(i) >= i for every i. Indeed, if y(i) >= i, choose i from the i-th interval. Conversely, if y(k) <= k-1 for some k, then the first k intervals are all contained in {1,...,y(k)}, which has at most k-1 elements, so distinct representatives are impossible by the pigeonhole principle. The same argument also proves the corresponding conjectures in A238873, A387118, and A388711. - _Clément Garrot_, Aug 31 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Aug 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017281 | OEIS-A387118

Number of integer partitions of n without choosable initial intervals.

Conjecture: Also the number of non-superdiagonal reversed integer partitions of n. - _Gus Wiseman_, Oct 04 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Sep 02 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017282 | OEIS-A387160

Terms of the form prime * m^2 in A351554.

Conjecture: This sequence has no common terms with A228058. See comments in A386430.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017283 | OEIS-A387171

Number of 4 element sets of distinct integer sided rectangles that fill an n X n square.

Conjectures from _Vaclav Kotesovec_, Aug 22 2025: (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Janaka Rodrigo_, Aug 20 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017284 | OEIS-A387186

Numbers k such that m!*k is a perfect square for some m < k-1.

Is a subsequence of A388851. Conjectured by Erdős and Graham to be the dominant component of that sequence (see also Erdős problem #374).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Terence Tao_, Sep 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017285 | OEIS-A387197

Numbers k such that 32 * 3^k - 1 is prime.

Conjecture: This sequence intersects with A387201 at k = 4 to form twin primes with center N = 2^5 * 3^4 = 2592 = A027856(10). Any such intersection has to be at an even k because if k is odd, either N-1 or N+1 has to be divisible by 5. A covering system can be constructed that eliminates all other intersections except where k = 4(mod 60), and for k > 4 with k = 4(mod 60), the search up to 10^5 makes the probability of another intersection in this residue class vanishingly small.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ken Clements_, Aug 21 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017286 | OEIS-A387205

a(n) = (n - 1)!*(2 + Harmonic(n - 1)) if n >= 1, and a(0) = 1.

Conjecture: Maple returns the exponential series expansion at x = 0:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Aug 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017287 | OEIS-A387231

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest unused positive number that shares a factor with a(n-1) if a(n-1) > a(n-2), else it is coprime to a(n-1) if a(n-1) < a(n-2).

In the first 100000 terms the fixed points are 1, 2, 8, 15, 32, 44, 50, 60, 90, 182, 198, 220; it is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Aug 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017288 | OEIS-A387232

Lexicographically earliest infinite sequence of distinct positive numbers such that a(2*n) > a(2*n-1) and a(2*n) > a(2*n+1) and all neighboring terms are coprime.

The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Aug 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017289 | OEIS-A387243

a(n) is the smallest positive k such that (k+n)^k < (k+1)^(k+1).

Conjecture: a(n) = A382131(n) - n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dwight Boddorf_, Aug 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017290 | OEIS-A387319

Numbers k for which modulo 3 and modulo another multiple of 3 are coincident among the fractions (2*k mod m)/m for 1 <= m <= k.

Conjecture: Every composite number, except 4, 8, 10, and 25, has at least one coincident pair linking m = 3 with another multiple of 3 among the fractions (2*k mod m)/m for 1 <= m <= k, while no prime has such a pair. Empirically verified to 10^5. Conjecture (based on comparing the first 10,000 terms): a(n) = A343311(n) for n>=3. - _R. J. Mathar_, Nov 10 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kenneth J Scheller_, Oct 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017291 | OEIS-A387331

Least k such that n*k + 1 is a prime > 2 with 2 as a primitive root; a(n) = 0 if no such k exists.

Computations show that for 1 <= n <= 2000 with 8 !| n, a(n) > 0. This agrees with Artin's conjecture: for each n with 8 !| n there should be infinitely many primes p == 1 (mod n) with 2 as a primitive root.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pablo Cadena-Urzúa_, Aug 26 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017292 | OEIS-A387349

Positions of 0's in A387348.

This sequence together with A387350 and A387351 partition the positive integers. Conjecture: the difference sequence (3, 2, 1, 2, 2, 1, 2, 3, 2, 1, 2, 3, 2, 3, 2, 1, ...) has exactly 3 distinct terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017293 | OEIS-A387350

Positions of 1's in A387348.

Conjecture: the difference sequence, (3, 5, 3, 2, 3, 5, 3, 2, 3, 3, 2, 3, 5, 3, 2, 3, 5, 3, 5, ...) has exactly 3 distinct terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017294 | OEIS-A387351

Positions of -1's in A387348.

Conjecture: the difference sequence, (5, 8, 5, 3, 5, 8, 5, 3, 5, 5, 3, 5, 8, 5, 3, 5, 8, 5, 8, ...) has exactly 3 distinct terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Aug 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017295 | OEIS-A387362

Cyclic numbers k such that k+2 is also a cyclic number.

Cohen (2025) conjectured and Pomerance (2025) proved that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Aug 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017296 | OEIS-A387405

a(n) is the least k that is a multiple of d=A240991(n), with abundancy ratio sigma(k)/k equal to (sigma(d)+1)/d, or -1 if no such k exists.

a(n) = A253560(A240991(n)) = A006530(A240991(n)) * A240991(n). - Conjectured, most likely true, especially if A240991 is a subsequence of A387406.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Aug 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017297 | OEIS-A387421

Numbers k such that sigma(k) >= 2*A057521(k), where A057521(n) gives the powerful part of n, and sigma is the sum of divisors function.

Conjectured to be equal to numbers k for which sigma(k) > 2*A057521(k), which holds if there are no such k that sigma(k) = 2*A057521(k). See comments in A386428, A387719.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017298 | OEIS-A387423

The length of binary expansion of n minus the length of the maximal common prefix of the binary expansions of n and sigma(n), where sigma is the sum of divisors function.

Positions of 0's in this sequence is given by such numbers n that sigma(n) = 2^k * n + r, for some n >= 1, k >= 0, 0 <= r < 2^k. These would include also quasi-perfect numbers and their generalizations, numbers n such that sigma(n) = 2^k * n + 2^k - 1, for some n > 1, k > 0 (see comments in A332223), if such numbers exist. However, it is conjectured that there are no other zeros than those given by A336702.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017299 | OEIS-A387426

Numbers k such that all nonzero fractions (2*k mod m)/m (for 1 <= m <= k) are distinct.

Conjecture: Every prime is in the sequence, and no composite greater than 24 is in the sequence. Empirically verified for k <= 10^6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kenneth J Scheller_, Oct 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017300 | OEIS-A387471

a(n) = number of interior triple intersections for the equal-angle model with parameter m = 2*n-1.

The closed form in FORMULA is conjectural but matches computations for n <= 200. Conjecture: a(n) = 6*n - 5 if n mod 5 != 0, and a(n) = 6*n+7 if n mod 5 = 0. Conjectured g.f.: 6*x/(1-x)^2 - 5*x/(1-x) + 12*x^5/(1-x^5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Constantinos Kourouzides_, Aug 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017301 | OEIS-A387499

a(n) is the conjecturally latest index where n occurs in A322345, or -1 if n doesn't occur.

a(n) is the conjecturally latest index where n occurs in A322345, or -1 if n doesn't occur.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Sep 08 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017302 | OEIS-A387523

Primes in the order in which they first divide a term of A386482.

Conjecture: if c is the first term in A386482 that is divisible by a prime p > 2, then c = 2*p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Sep 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017303 | OEIS-A387631

Sombor energy of the n-triangular honeycomb queen graph divided by sqrt(2).

Conjecture: a(n) = (3*n^3 - 9*n^2 + 24*n) / 2, for n > 4. - _Vincenzo Librandi_, Sep 07 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Sep 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017304 | OEIS-A387640

A387523(n) is the a(n)-th prime number.

If the conjecture in A387523 is true, then the present sequence is self-inverse.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Sep 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017305 | OEIS-A387681

Numbers that can be written in exactly seven ways as s_1^x_1 + ... + s_t^x_t, with 1 < s_1 < ... < s_t and {s_1,..., s_t} = {x_1,..., x_t} for some t > 0.

The sequence is infinite (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alberto Zanoni_, Dec 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017306 | OEIS-A387682

a(1) = 1, a(2) = 2, a(3) = 4; for n > 3, a(n) is the smallest unused positive number that shares a factor with a(n-1) and a(n-2) if a(n-1) > a(n-2), else it is coprime to a(n-1) and a(n-2) if a(n-1) < a(n-2).

For the terms studied the primes, other than 3 and 5, appear in their natural order, and the only two fixed points are the first two terms. The sequence is conjectured to be a permutation of the positive numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017307 | OEIS-A387724

Numbers k such that sigma(k)^2 >= k * sigma(A003961(k)), where A003961 is fully multiplicative with a(p) = nextprime(p).

Conjecture: 6 is the only nondeficient term, and all other terms are in A337382 (which is a subsequence of A005100).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017308 | OEIS-A387731

a(n) = n if n <= 3; thereafter a(n) is either the greatest number k < a(n-1) not already used such that gcd(k, a(n-1)) = 1 and gcd(k, a(n-2)) > 1, or if no such k exists then a(n) is the smallest number k > a(n-1) not already used such that gcd(k, a(n-1)) = 1 and gcd(k, a(n-2)) > 1.

Conjecture: this sequence is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rémy Sigrist_, Sep 06 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017309 | OEIS-A387737

Number of partitions of n such that the largest part is >= n/2 and if the partition has k parts then the smallest part is >= k.

Conjecture: a(n) = A387736(n) + 1, n>=4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sean A. Irvine_, Sep 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017310 | OEIS-A387757

a(1) = 1, a(2) = 3; for n > 2, a(n) is the smallest unused positive number that differs from a(n-1) by more than 1 and shares a factor with a(n-1) if a(n-1) < a(n-2), else it is coprime to a(n-1) if a(n-1) > a(n-2).

There are 73 fixed points in the first 500000 terms, these are 1, 16, 22, 28, 34, ..., 1901, 3174, 33376. It is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017311 | OEIS-A387758

Lexicographically earliest infinite sequence of distinct positive numbers such that, for n > 2, a(n) shares a factor with a(n-1) and is coprime to a(n-2) if a(n-1) > a(n-2), else it is coprime to a(n-1) and shares a factor with a(n-2) if a(n-1) < a(n-2).

In the first 250000 terms the fixed points are 1, 2, 8, 15, 25, 42, 49, 81, 90, 564; it is not known if more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017312 | OEIS-A387759

Lexicographically earliest infinite sequence of distinct positive numbers such that, for n > 2, a(n) shares a factor with a(n-1) and is coprime to a(n-2) if a(n-1) < a(n-2), else it is coprime to a(n-1) and shares a factor with a(n-2) if a(n-1) > a(n-2).. Start with a(1) = 1, a(2) = 4, a(3) = 9.

Of the terms studied there are no fixed points other than the first, and it is likely none exist. The sequence is conjectured to contain all positive integers other than the primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017313 | OEIS-A387760

Numbers k such that 64 * 3^k - 1 is prime.

Conjecture: This sequence intersects with A387761 at k = 1 and k = 7 to form twin primes with centers N = 2^6 * 3^1 = 192 = A027856(7) and N = 2^6 * 3^7 = 139968 = A027856(11). A covering system can be constructed that eliminates all other intersections (see linked program), and the search up to 250000 makes the probability of another intersection vanishingly small.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ken Clements_, Sep 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017314 | OEIS-A387790

Number of ways to write n as S(i) + S(j) + 2*S(k), where i,j,k are positive integers with i >= j, and S(k) is the sum of the first k primes.

Conjecture: a(n) > 0 for all n >= 5340. In other words, each integer n >= 5340 is a sum of four elements of the set {Sum_{k=1..m} prime(k): m = 1,2,3,...} two of which are equal.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 09 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017315 | OEIS-A387852

The final prime number in the base n sequence when starting from 2 and creating the smallest unused prime number, when read in base n, by either removing or adding a single digit, from 0 to n-1, anywhere in the base n value of the previous number. If the base n sequence does not terminate, a(n) = -1.

It is conjectured that the sequence of primes in base n terminates for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Oct 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017316 | OEIS-A387865

Number of ways to write 2*n as p + p' + q, where p is a prime, p' is the first prime after p, and q is a practical number.

Conjecture: a(n) > 0 for all n > 4. In other words, any even number not smaller than 10 is the sum of two consecutive primes and a practical number.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017317 | OEIS-A387907

a(n) = position of first odd digit of 2^n, starting from left, or 0 if no odd digit exists.

It is conjectured that a(n) is positive for n >= 12. This is sometimes known as the "2048 problem".

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Harry Richman_, Sep 12 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017318 | OEIS-A387957

Number of integer partitions of n with parts colored by {0, 1, 2, 3} such that the sum of colors is congruent to 1 (mod 4).

Conjectured g.f.: (1/4) * (Product_{j>=1} 1/(1-x^j)^4 - Product_{j>=1} 1/(1-x^(2*j))^2). Conjectured g.f.: (1/4) * (G.f. of A023003(x) - G.f. of A000712(x^2)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Hutton_, Nov 09 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017319 | OEIS-A387959

Least positive integer m such that the n numbers (prime(1)-1)^2,...,(prime(n)-1)^2 are pairwise incongruent modulo m.

Conjecture: a(n) = prime(n+1) except for n = 1, 2, 4, 9. This has been verified for n <= 55000. The conjectural recurrence for primes is equivalent to the conjecture in A387947, but different from the conjecture in A181901. The conjecture has been verified for n <= 10^6 by Nie Chen at the First Middle School of Nanjing via C++. - _Zhi-Wei Sun_, Oct 20 2025 Jiao-Min Lin, a former student of Nanjing University, has verified the conjecture for n <= 10^7. - _Zhi-Wei Sun_, Oct 23 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 12 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017320 | OEIS-A387991

Number of ways to write n as i*prime(i) + j*prime(j) + k*prime(k) + m*prime(m) with 1 <= i <= j <= k <= m.

Conjecture: a(n) > 0 for all n >= 3070. In other words, any integer n >= 3070 is a sum of four terms of A033286. This has been verified for n <= 2*10^5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017321 | OEIS-A388012

Numbers k such that sigma(k)/k >= 5/3.

It is not known whether sigma(k)/k = 5/3 for any k, so conjecturally, these are all numbers k such that sigma(k)/k > 5/3. However, if such a k exists, then it must by necessity be an odd square that is a multiple of 3, but not a multiple of 5 (and thus in A388016). Furthermore, 5*k is then an odd perfect number. See theorem 3 in Weiner paper.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017322 | OEIS-A388268

Numbers k for which A000005(k) >= A017665(k), where A000005 is the number of divisors function, and A017665 is the numerator of the abundancy ratio, sigma(k)/k.

After the initial 1, conjectured to be a subsequence of A023196. Note that for deficient numbers, k in A005100, we always have gcd(sigma(k),k) <= A032742(k), but not necessarily A000203(k) > A000005(k)*A032742(k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017323 | OEIS-A388269

Odd numbers k for which A000005(k) >= A017665(k), where A000005 is the number of divisors function, and A017665 is the numerator of the abundancy ratio, sigma(k)/k.

All odd terms of A001599 and of A007691 are also included in this sequence. It has been conjectured that both contain only one odd term, their initial 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017324 | OEIS-A388275

Numbers k such that k and sigma(k) have the same largest prime factor and also its exponent is equal in their prime factorizations, with a(1) = 1 included by convention.

Question: Is 1 the only common term of this sequence and A001694? If true, then it would imply the conjecture described in A386428, and also refute the existence of any odd terms of A007691 larger than 1 with an odd abundancy index (e.g., odd terms in A005820 and in A046060).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 20 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017325 | OEIS-A388425

First index k where A368558(k) = 2*n, or 0 if no such index exists.

Conjecture: This sequence does not contain 0. In other words, A368558 contains every positive even integer.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Yuen_, Sep 17 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017326 | OEIS-A388651

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest unused positive number that has at least one prime exponent in its prime factorization that equals a prime exponent in the prime factorization of a(n-1).

The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017327 | OEIS-A388652

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest unused positive number that shares a factor with a(n-1) and has at least one prime exponent in its prime factorization that equals a prime exponent in the prime factorization of a(n-1).

In the first 250000 terms the fixed points are 1, 2, 28, 35, 38, 65, 121, 215, 323; it is likely no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017328 | OEIS-A388711

Triangle read by rows where T(n,k) is the number of integer partitions of n into k parts with choosable initial intervals.

Conjecture: Also the number of superdiagonal reversed partitions of n into k parts, where a sequence (y_1, ..., y_k) is superdiagonal iff y_i >= i for all i = 1..k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Sep 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017329 | OEIS-A388712

Triangle read by rows where T(n,k) is the number of integer partitions of n into k parts without choosable initial intervals.

Conjecture: Also the number of non-superdiagonal reversed partitions of n into k parts, where a sequence (y_1, ..., y_k) is superdiagonal iff y_i >= i for all i = 1..k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Sep 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017330 | OEIS-A388713

Triangle read by rows where T(n,k) is the number of superdiagonal multisets summing to n (reversed partitions of n) with maximum k.

Conjecture: Also the number of integer partitions of n with maximum k and choosable initial intervals. For example, row n = 12 counts the following partitions:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Sep 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017331 | OEIS-A388724

Triangle read by rows where T(n,k) is the number of strictly superdiagonal reversed integer partitions of n with maximum k.

Conjecture: Column sums are A000245.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Sep 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017332 | OEIS-A388967

Smallest number in a group of 4 or more consecutive numbers such that, beyond the first number, no number's prime factor exponents equal any prime factor exponent of the previous number.

For other known terms, see the Examples. These are derived using the expression f(x) given below; it is not known if other terms exist which are not values of f(x). It is conjectured that only numbers that are values of f(x) for some x are terms of this sequence. The number 99612037019888, which lies between f(4) and f(5), is a term of this sequence and is a counterexample to the conjecture above that only numbers that are values of f(n) for some n are terms of this sequence. - _Peter Bala_, Sep 30 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Sep 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017333 | OEIS-A388986

Numbers k such that {sum of unitary divisors of k} plus {sum of squarefree divisors of k} is less than 2*k.

Conjecture: Numbers k such that at least 1 number less than k is a multiple of rad(k), but its prime factorization is not a subset of k's. As an example, rad(36)=6 and 30 is a multiple of 6 less than 36, but the prime factorization of 30 is not a subset of the prime factorization of 36, so 36 is in the sequence. - _Anthony Browne_, May 15 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 29 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017334 | OEIS-A389000

Smallest k such that the sums of digits of k and k^2 in base 10 are both divisible by n or a(n) = -1 if no such k exists.

Conjecture: a(9*m+5) = 2*10^(2*m+1)-1 for m >= 0. - _Chai Wah Wu_, Oct 01 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arjun Maneesh Agarwal_, Sep 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017335 | OEIS-A389079

Odd nonsquarefree numbers k such that {sum of unitary divisors of k} plus {sum of squarefree divisors of k} >= 2*k.

It is conjectured that there are no common terms with A228058 (see also A388986), and for all terms, A001221(a(n)) >= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017336 | OEIS-A389105

a(n) is the smallest value of k such that gcd(A381466(k-1), k) = n.

We conjecture that a(n) > 0 for all n; i.e., every positive integer is a divisor in A381466 at least once; indeed, we conjecture that every positive integer is a divisor infinitely many times. For positive values found so far, a(n) > 1.2^(n-1), and we also conjecture this is always true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sam Chapman_, Mar 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017337 | OEIS-A389106

Least positive integer k such that A386395(k*n) has the least possible number of digits.

It appears that one can indeed achieve a minimal final length of 2 digits if n is a multiple of 10, and of 1 digit in all other cases. When n has an even number 2m of trailing zeros, multiplication with n/100^m yields a starting point equivalent to n/10^m, so it is sufficient to consider just one or zero trailing '0's. More generally, multiplying by the squarefree kernel f of n yields a starting point equivalent to sqrt(n*f) <= n with equality iff n is squarefree. Thus, no nonsquarefree n can be the first counterexample to the question/conjecture. - _M. F. Hasler_, Sep 24 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ali Sada_, Sep 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017338 | OEIS-A389110

Least m such that the two-sided exact 95% Clopper-Pearson binomial confidence interval has worst-case width <= 1/n.

a(n) <= A182868(n) = 4*n^2 + n - 1 for 1 <= n <= 48 with equality at n = 2..5. Conjecture: this strict inequality holds for all n >= 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pablo Cadena-Urzúa_, Sep 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017339 | OEIS-A389142

Lower (1/3)-midsequence of square numbers and triangular numbers; see Comments.

Suppose that s = (s(n)) and t = (t(n)) are sequences of numbers and r > 0. The lower (r)-midsequence of s and t is given by u = floor(r*(s + t)); the upper r-midsequence of s and t is given by v = ceiling(r*(s + t)). If s and t are linearly recurrent and r is rational, then u and v are linearly recurrent. Conjecture: a(n+2) = A111097(n+2) for n>=0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Sep 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017340 | OEIS-A389219

Terms k of A228058 for which A048146(k)+A162296(k) >= 2*k, where A048146 is the sum of non-unitary divisors, and A162296 is the sum of divisors that have a square factor.

If A228058 is a subsequence of A388986 as is conjectured (see A388986), then A386427 (nondeficient terms of A228058) is a subsequence of this sequence. - _Antti Karttunen_, Oct 05 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 25 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017341 | OEIS-A389221

Numbers k with exactly four coincident values among the fractions (2*k mod m)/m for 1 <= m <= k.

Conjecture: All numbers of the form 4*p with p prime and p > 13 belong to the sequence, and no other numbers greater than 80 do. Empirically verified through 100000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kenneth J Scheller_, Oct 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017342 | OEIS-A389238

Odd numbers k whose symmetric representation of sigma, SRS(k), consists of an odd number, >1, of parts and whose center part is the minimum of all parts.

Conjecture: The width pattern for any k in this sequence for which SRS(k) consists of three parts is 10101 or 1012101; in addition, these numbers k form sequence A087718(n), n >= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Oct 28 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017343 | OEIS-A389280

a(n) is the least number k > 0 such that wt(k^n) >= wt(k) * T(n), or -1 if such k does not exist, where wt(x) = A000120(x) is the binary weight of x, and T(n) = n*(n+1)/2 = A000217(n) is the n-th triangular number.

Conjecture: a(n) > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Sep 28 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017344 | OEIS-A389302

Position of records in A389235.

Conjecture: a(n) = 1008*10^(n-2) for n>7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Karl-Heinz Hofmann_ and _Hugo Pfoertner_, Sep 29 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017345 | OEIS-A389343

Number of integer partitions of n with parts colored by {0, 1, 2} such that the sum of colors is congruent to 0 (mod 3).

Conjectured g.f.: (1/3) * (Product_{j>=1} 1/(1-x^j)^3 + 2*Product_{j>=1} 1/(1-x^(3*j))). Conjectured g.f.: (1/3) * (G.f. of A000716(x) + 2*G.f. of A000041(x^3)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Hutton_, Oct 29 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017346 | OEIS-A389354

a(n) = (A388289(n)^2 - 1)/(3*2^5), for n >= 1.

The reduced principal indefinite binary quadratic Markoff forms are F_p(n; X, Y) = X^2 + b(n)*X*Y - b(n)*Y^2, for n >= 1, representing -m(n)^2. The Frobenius-Markoff conjecture, that there is only one Markoff triple with largest number m(n), is trivially true for n = 1 and n = 2 (the so-called singular cases), and for n >= 3 it claims that there is only one proper solution F_p(n; X, Y) = -m(n)^2, with X < 0, Y >= 1, and Y - X < m(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 02 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017347 | OEIS-A389355

a(n) = (A388291(n)^2 - 1)/24.

See A389354 for the definitions of Disc(n)= A305312(n), D(n) = A308687(n), the reduced principal indefinte binary quadratic form F_p(n; X, Y), for n >= 1, and the Frobenius - Markoff conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Wolfdieter Lang_, Oct 02 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017348 | OEIS-A389364

Harshad numbers h (terms of A005349) such that A389235(h) + 2 = A055642(h) and A385105(h) is a 2 digit number with digit sum 10.

Conjecture: 9723402 and 74401740 are the last terms with really new digits not seen before in the sequence. Onward trivial terms are only previous terms multiplied by 10^x (See links for not trivial terms).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Karl-Heinz Hofmann_, Oct 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017349 | OEIS-A389368

a(n) = R(A007310(n)), where R(k) = (5*k+1)/(2^r*3^s), with r and s as large as possible, that is R(k) = A065330(A016861(k)).

Conjecture: By successive iterations x->a(x) we always end up arriving at 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Feb 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017350 | OEIS-A389392

a(n) is the number of 4 element sets of distinct integer sided trapezoids whose base angles are 60 degrees that fill an equilateral triangular grid of side n units.

Conjectured g.f.: x^5 * (1+9*x+19*x^2+29*x^3+27*x^4+14*x^5+3*x^6-6*x^7) / ((x^2+1) * (x+1)^2 * (x^2+x+1)^2 * (x-1)^4). - _Sean A. Irvine_, Oct 15 2025

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Janaka Rodrigo_, Oct 02 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017351 | OEIS-A389425

First differences of A199716.

Conjecture: a(n) tends to 1 in frequency (i.e., {n: a(n) = 1} has natural density 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aied Sulaiman_, Oct 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017352 | OEIS-A389472

G.f. satisfies A(x) = A(x^2 + x^3)/x^2 - 1.

Conjecture: a(2*n-1) = 0 (mod 2) for n > 1. Conjecture: a(3*n-1) = 0 (mod 3) for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017353 | OEIS-A389474

G.f. A(x) satisfies A(x^2 + 2*x^3 + 2*A(x)^2) = 3*A(x)^2.

(3) [x^n] A(x)^2 == 0 (mod 9) for n > 3 (conjecture). (4) [x^n] A(x)^(3*k) - A(x^3)^k == 0 (mod 3) for n >= 1 and k >= 1 (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017354 | OEIS-A389475

G.f. A(x) satisfies A(3*x^2) = 2*x^2 + A(x)^2 - 2*A(x)^3.

(4.a) A(x)^3 == A(x^3) (mod 3) (conjecture). (4.b) A(x)^2 == x^2 + 2*A(x^3) (mod 3) (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 26 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017355 | OEIS-A389524

Triangle read by rows. T(1, 1) = 1, T(n, k) = [n >= k](1 + Sum_{i=1..k-1} T(n - i, k - 1) - Sum_{i=1..n-1} T(n - i, k)).

Conjecture 1: A000005(n) = Sum_{k >= 1} T(n, k), verified up to n = 1000. Conjecture 2: A274628(n) = Sum_{k >= 1} T(n, n - k + 1)*(n - k + 1), verified up to n = 500. Conjecture 3: A049820(n) = T(2*n, n), verified up to n = 250.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mats Granvik_, Oct 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017356 | OEIS-A389525

Number of ways to write 2*n + 1 as prime(i^2) + prime(j^2) + prime(k^2) + 2*prime(m^2) with i >= j >= k >= m >= 1.

Conjecture: a(n) > 0 for n > 3200. See also A389517 for a similar conjecture. Conjecture: for all k, there exists n_k such that a(m)>k for all m >= n_k. k conjectured largest value of n for which a(n) = k

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 07 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017357 | OEIS-A389535

G.f. A(x) satisfies A(x*A(x)^4 + x*A(x)^5) = A(x)^5.

(7) A(x)^5 == A(x^5) (mod 5) (conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017358 | OEIS-A389536

G.f. A(x) satisfies: A(x) = A(x^2 + 2*x^3) / x.

Conjecture: for n > 1, a(n) == 2 (mod 4) when n = 2^k + 1 (for some k >= 0) otherwise a(n) is divisible by 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Oct 28 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017359 | OEIS-A389537

G.f. A(x) satisfies A(A(x) - x) = A(x)^2.

Conjecture: for n > 2, a(n) == 2 (mod 4) when n = 3*2^k for k >= 0 otherwise a(n) is divisible by 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017360 | OEIS-A389539

Maximum number of edges blocked from infinity by n vertices of a hexagonal grid.

Conjecture: for n >= 0, a(n) = max(0, 2*n - 3 + 3*floor(((n - 3)^2 + 3)/12)). - _Conner Silverstein_, Apr 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Conner Silverstein_, Jan 02 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017361 | OEIS-A389540

G.f. A(x) satisfies A(x)^2 = A(2*x - 2*A(x)) / 2.

Conjecture: a(n) is odd iff n = 2^k for k >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017362 | OEIS-A389542

G.f. A(x) satisfies A(A(x)^2 - x^2) = 4*A(x)^3.

Conjecture: for n > 1, a(n) == 2 (mod 4) if n = 2^k + 1 (for some k >= 0) otherwise a(n) is divisible by 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Nov 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017363 | OEIS-A389567

Number of integer exponents k such that in the expansion in powers of q of (q^(-1/24) * eta(q))^k the term q^n has a zero coefficient and eta is the Dedekind eta function.

Yet there are only 11 different types of zeros observed and it is conjectured that no others exist. A two digit value is first time reached at a(419) = 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Scheuerle_, Nov 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017364 | OEIS-A389591

Sum of the divisors r of n such that r is not greater than twice the adjacent previous divisor of n.

a(n) = A000203(n) - A379379(n). (Conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Oct 08 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017365 | OEIS-A389592

Irregular triangle read by rows: T(n,k) is the sum of the numbers that are not the smallest number in the k-th 2-dense sublist of divisors of n, with n >= 1, k >= 1.

Conjecture: T(n,k) = A384149(n,k) - A379288(n,k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Oct 08 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017366 | OEIS-A389614

a(n) = 2*n^2 + 1 - (m*n - 3*m*(m+1)/2) where m = floor(n/3).

Let Tbar denote a planar graph consisting of a distinguished point C (the center), an infinite line L through C, and a semi-infinite line through C perpendicular to L. Let M(n) denote the maximum number of regions that the plane can be divided into by drawing n Tbars. Cutler and Sloane (2025) conjectured that M(n) = a(n) for all n. This conjecture has now been established by _Jonas Karlsson_, see the reference. - _N. J. A. Sloane_, Jan 29 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | David O. H. Cutler and _N. J. A. Sloane_, Oct 17 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017367 | OEIS-A389657

a(n) is zero if n is a square, otherwise a(n) is the least triangular number m > 0 such that m*n is a triangular number.

Conjecture: Each positive rational number r with sqrt(r) irrational can be written as m*(m+1)/(n*(n+1)) = T(m)/T(n), where m and n are positive integers, and T(k) denotes the triangular number k*(k+1)/2. This conjecture implies that a(n) exists for any positive integer n. After the author posed the conjecture to MathOverflow, Alexander Kalmynin provided a proof of the conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 09 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017368 | OEIS-A389664

a(n) = 1 if n is a square, otherwise a(n) is the least integer m > 1 such that (m^2 - 1)*n + 1 is a square.

Conjecture: For any positive rational number r with sqrt(r) irrational, we have r = (m^2-1)/(n^2-1) for some integers m and n greater than one. This conjecture implies that a(n) exists for all n > 0. Clearly, 4*(m^2-1) = n^2-1 (i.e. (2m)^2 - n^2 = 3) fails for any m,n = 2,3,.... After the author posed the conjecture to MathOverflow, Alexander Kalmynin provided a proof of the conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 10 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017369 | OEIS-A389689

Length of the longest possible eventual period of the win/lose state of a subtraction game where the largest element of the subtraction set is n.

By Althöfer and Bültermann's paper, a(16n+1)>=56n^3+52n^2+9n+1 for 1<=n<=26. It is conjectured that a(38n+1) is superlinear using the subtraction set {n, 8n, 30n+1, 37n+1, 38n+1}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rui Yao_, Oct 10 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017370 | OEIS-A389793

Least positive integer m such that n*(prime(m) + prime(m+1)) = prime(k) + prime(k+1) for some positive integer k.

Conjecture: Each positive rational number can be written as (prime(k) + prime(k+1))/(prime(m) + prime(m+1)), where k and m are positive integers. The above conjecture is weaker than the strong conjecture in A259540, for, if prime(m) and prime(m+1) = prime(m) + 2 are twin prime then prime(m) + prime(m+1) = 2*k, where k = prime(m) + 1 with k - 1 and

k + 1 both prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017371 | OEIS-A389835

Numbers k such that wt(Fibonacci(k)) * e > k, where wt(x) is the binary weight of x (A000120), and e is the natural logarithm base.

Conjecture: the sequence is finite, and 4366 is the last term. Finiteness is likely since binary weight grows only as a factor log2(phi)/2 which is smaller than 1/e, where phi is the golden ratio.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Oct 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017372 | OEIS-A389852

Least positive integer d such that n + d^3 is the sum of 3 positive cubes.

Conjecture: a(n) exists for all positive integers n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhining Yang_, Jan 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017373 | OEIS-A389863

Numbers k with exactly two distinct repeated values among the fractions (2*k mod m)/m for 1 <= m <= k.

Conjecture: All numbers of the form 2*p with p prime and p > 5 belong to the sequence, and no other numbers greater than 36 do. Empirically verified for k <= 100000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kenneth J Scheller_, Oct 17 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017374 | OEIS-A389879

A005117(k) - 1 where k is the least k such that A389412(k) = n.

Probabilistic arguments (see the Englezou link in A389412) suggest the following conjecture: the set {a(r) : A056170(a(r)) = n} is finite for every n, and the non-unitary prime factors of a(r) are the first n primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Miles Englezou_, Oct 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017375 | OEIS-A389880

Number of permutations f of {1,...,n} with f(1) = 1 such that k*f(k) (k = 1..n) are distinct practical numbers.

Conjecture: a(n) > 0 except for n = 3, 5, 7. This is stronger than Conjecture 2 in A321651.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 15 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017376 | OEIS-A389914

Numbers k such that A048675(k) is in A186891\{1}.

If Ulas & Ulas conjecture (see A186891) is correct, then A277319 is a subsequence of this sequence, A277316 is a subsequence of A277318, A277317 is a subsequence of A389912, and A389917 contains only composite terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017377 | OEIS-A389917

Odd numbers m such that the Stern polynomial B(m,x) is reducible.

Ulas & Ulas conjecture implies that there should be only composite terms in this sequence. See A186891.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017378 | OEIS-A389926

The total length of the base n sequence when starting from 2 and creating the smallest unused prime number, when read in base n, by either removing or adding a single digit, from 0 to n-1, anywhere in the base n value of the previous number. If the base n sequence does not terminate, a(n) = -1.

It is conjectured that the sequence of primes in base n terminates for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Oct 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017379 | OEIS-A389927

The total length of the sequence when starting from n and creating the smallest unused prime number by either removing or adding a single digit anywhere in the value of the previous number. If the sequence does not terminate, a(n) = -1.

It is conjectured that the sequence of primes terminates for all starting values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Oct 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017380 | OEIS-A389940

Numbers k for which a sphere of radius k is a member of at least 1 set of 4 spheres of coprime positive integer radii which are tangent to a plane when arranged as mutually tangent to one another.

It is unknown if specific terms can otherwise be found algorithmically, though it is conjectured that the general increase in gap size between successive terms as n increases continues to infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles L. Hohn_, Oct 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017381 | OEIS-A390007

a(n) is the least possible difference between the areas of the largest and smallest rectangles in a 3 element set of distinct integer sided rectangles filling an n X n square.

Conjecture: except for n = 4, there is a unique set of three rectangles corresponding to the minimum defect.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Janaka Rodrigo_, Oct 21 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017382 | OEIS-A390088

a(n) is the least integer base b > 1 in which the representation of n uses only prime digits; or 0 if no such base exists.

Conjecture: Exactly the values k = 0, 1, 4, 6, 9 satisfy a(k) = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Oct 29 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017383 | OEIS-A390100

Numbers k with exactly three coincident values among the fractions (2*k mod m)/m for 1 <= m <= k.

Conjecture: All numbers of the form 3*p with p prime and p > 7 belong to the sequence, and no other numbers greater than 52 do. Empirically verified through 100000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kenneth J Scheller_, Oct 24 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017384 | OEIS-A390107

Least positive integer m such that m*(n-1) + 1 is relatively prime to C(m*n^2, m*n), where C(k, j) denotes the binomial coefficient k!/(j!*(k-j)!).

Conjecture: a(n) <= sqrt(3*n + 4) for all n > 0. In the linked 2012 paper, the author conjectured that for any integer k > 2 there is a positive integer n such that C(k*n, n) does not divide C(k*n, n - 1)*C(k^2*n, k*n) (i.e., (k - 1)*n + 1 does not divide C(k^2*n, k*n)). See also A390117 for a related conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 25 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017385 | OEIS-A390112

Smallest exponent b >= 2 such that the absolute value of the alternating 3-digit block sum of n^b equals n; 0 if none.

Conjectured to exist for all n that are not powers of 1000, for which S_3(1000^b) = 1 for all b.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pieter Post_, Oct 25 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017386 | OEIS-A390117

Least integer m > 1 such that m*n + 1 is a prime.

Conjecture: (i) a(n) <= 2*sqrt(n+6) for all n > 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 25 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017387 | OEIS-A390118

Least positive integer m such that m*n + 1 = p^2 for some prime number p.

Conjecture: a(n) <= (n+5)*(n-1) for all n > 1 (verified for n <= 12000).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Oct 25 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017388 | OEIS-A390148

Rectangular array read by rows of ordered quadruples of coprime positive integers {r1..r4} for which spheres of radius r1..r4 are tangent to a plane when arranged as mutually tangent to one another, with rows in lexicographic order.

Some pairs of quadruples contain coincident triples; the smallest are {52, 84, 156, 273} and {84, 156, 273, 364}. Quadruple {364, 741, 1092, 1596} has two of these: {228, 364, 741, 1092} and {741, 1092, 1596, 6916}. It is conjectured that arbitrarily large such chains of quadruples exist, but an infinite chain cannot, because the only possible such chain contains all successive terms with relative ratios of A289917, which is irrational. Observed for rows n -> 1000 and conjectured for n -> oo:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles L. Hohn_, Oct 26 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017389 | OEIS-A390208

Number of integer partitions of n with parts colored by {0, 1, 2} such that the sum of colors is congruent to 1 (mod 3).

Conjectured g.f.: (1/3) * (Product_{j>=1} 1/(1-x^j)^3 - Product_{j>=1} 1/(1-x^(3*j))). Conjectured g.f.: (1/3) * (G.f. of A000716(x) - G.f. of A000041(x^3)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Hutton_, Oct 29 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017390 | OEIS-A390217

Number of partitions of n with parts colored by {0, 1, 2, 3} such that the sum of colors is congruent to 0 (mod 4).

Conjectured g.f.: (1/4) * (Product_{j>=1} 1/(1-x^j)^4 + Product_{j>=1} 1/(1-x^(2*j))^2 + 2*Product_{j>=1} 1/(1-x^(4*j))). Conjectured g.f.: (1/4) * (G.f. of A023003(x) + G.f. of A000712(x^2) + 2*G.f. of A000041(x^4)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Hutton_, Oct 29 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017391 | OEIS-A390228

Least positive integer m such that the n numbers c(0),...,c(n-1) are incongruent modulo m, where c(k) = A000521(k).

Conjecture: a(n) is prime for each n > 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017392 | OEIS-A390319

Numbers k with exactly five coincident values among the fractions (2*k mod m)/m for 1 <= m <= k.

The sequence was conjectured to be finite and to consist exactly of the 11 listed terms, based on an empirical search through 10^5. An accompanying question asked whether the analogous sequences for other numbers of distinct repeated values are finite or empty.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kenneth J Scheller_, Nov 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017393 | OEIS-A390352

Least positive integer m such that (2*n-1)*(prime(m)+prime(m+1)+prime(m+2)) = prime(k)+prime(k+1)+prime(k+2) for some k > 0.

The existence of a(n) follows from the following general conjecture. Conjecture: Let m > 1 and let P(m) be the set of all sums of m consecutive primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 02 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017394 | OEIS-A390366

The total length of the base n sequence when starting from 2 and creating the smallest unused prime number, when read in base n, by either removing or changing or adding a single digit, from 0 to n-1, anywhere in the base n value of the previous number. If the base n sequence does not terminate, a(n) = -1.

It is conjectured that the sequence of primes in base n terminates for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017395 | OEIS-A390367

The final prime number in the base n sequence when starting from 2 and creating the smallest unused prime number, when read in base n, by either removing or changing or adding a single digit, from 0 to n-1, anywhere in the base n value of the previous number. If the base n sequence does not terminate, a(n) = -1.

It is conjectured that the sequence of primes in base n terminates for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Nov 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017396 | OEIS-A390424

Expansion of 1/(2 - g^2)^2, where g = 1+x*g^3 is the g.f. of A001764.

Conjecture D-finite with recurrence 10*n*(2*n-5)*a(n) +(-511*n^2+1968*n-1337)*a(n-1) +2*(2461*n^2-12759*n+14814)*a(n-2) +4*(-5447*n^2+35358*n-55483)*a(n-3) +16*(2651*n^2-20323*n+38832)*a(n-4) -2976*(3*n-11)*(3*n-13)*a(n-5)=0. - _R. J. Mathar_, Jan 26 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Dec 10 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017397 | OEIS-A390457

a(1) = 1 thereafter a(n) = floor(k(n)^n) where k(n) = a(1).a(2)a(3)...a(n-1) = 1.111... is the decimal digit concatenation of all preceding terms.

It is conjectured that a(n) = floor(k^n) where k = lim_{n->oo} k(n), which would be a constant k formed from its own powers. This conjecture fails if some floor(k^n) > floor(k(n)^n) due to k > k(n), but this seems unlikely since arbitrary decimals after k(n) would be a difference k^n - k(n)^n roughly equal to (n+1)*k(n)^n / 10^L(n) where L(n) is the length of k(n) after its decimal point, and that difference is small and rapidly decreasing.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jesus Pineiro Sanchez_, Nov 01 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017398 | OEIS-A390549

Smallest b>=2 for which Q_{n}(b) = prime(n) + Sum{k=1..n-1} b^(prime(n) - prime(k)) is prime, or -1 if no such b exists.

Conjecture: Q_{n}(b) takes infinitely many prime values for b >= 2 and n >= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Davide Oliveri_, Nov 10 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017399 | OEIS-A390550

Smallest b >= 2 for which P_{n}(b) = Sum_{k=1..n} b^(prime(k)-2) is prime or 0 if such a number does not exist.

Conjecture: the polynomials P_{n}(b) are irreducible in Z[b] for all but finitely many values of n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Davide Oliveri_, Dec 17 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017400 | OEIS-A390583

Rectangular array read by rows of ordered quadruples of coprime positive integers {r1..r4} for which circles of radius r1..r4 can be arranged as mutually externally tangent to one another, with rows in lexicographic order.

It is observed and generally conjectured that if the largest 2 terms of a quadruple are equal, then the absolute difference between the other 2 terms is 1 or 2 times a cube.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles L. Hohn_, Nov 11 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017401 | OEIS-A390626

Numbers k such that the symmetric representation of sigma, SRS(k), and SRS(2k), both considered as lists of the sizes of their parts, have the same number of parts.

Conjecture: SRS(2k) is a parts-multiple of SRS(k) for every k in this sequence (True for 2k <= 10^6).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Nov 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017402 | OEIS-A390671

Minimum number of hyperoperations to get n using only 1s.

Conjecture: Let b(k) be a sequence such that b(k) = max{n|a(n)=k}. b(0)=1; b(1)=2; b(2)=3; b(3)=4; b(k)=H_(b(k-4))(3,2). Conjecture: Let k = min{h|h is a natural number, h >= (min{#T[P]}-1)/4} b(k) >= min{x|P(x) is true}).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alessandro M. Pettinari_, Nov 14 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017403 | OEIS-A390730

Number of ways to write 6*n+2 as an unordered sum of four squares of elements in the set {m >= 0: m = 0 or 2*m is practical}.

Conjecture: Let S be the set of those natural numbers m such that 2*m is either zero or practical. Then any integer n >= 0 with n == 0,1 (mod 4) or n == 2,4 (mod 6) can be written as w^2 + x^2 + y^2 + z^2 with w,x,y,z in the set S. See also A389523 for related conjectures.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017404 | OEIS-A390762

Number of ways to write n as (p-1)^2 + (q-1)^2 + (r-1)^2/2 + (s-1)^2/4 with p <= q, where each of p,q,r,s is either 1 or an odd prime.

Conjecture: Let S = {(p-1)/2: p is 1 or an odd prime}. Then each nonnegative integer n can be written as w^2 + b*x^2 + c*y^2 + d*z^2 with w,x,y,z in the set S, provided that (b,c,d) is among the following 25 triples: (1,2,3),(1,2,4),(1,2,5),(1,2,6),(1,2,7),(1,2,12),(1,3,5),(1,3,6),(2,2,3),(2,2,5),(2,3,3),(2,3,4),(2,3,5),(2,3,7),(2,3,9),(2,3,10),(2,4,4),(2,4,5),(2,4,9),(2,4,10),(2,4,11),(2,4,12),(2,5,7),(2,5,8),(2,5,9). This conjecture holds for (b,c,d) = (2,4,4) if and only if a(n) > 0 for all n >= 0. We have verified a(n) > 0 for all n <= 10^6. It seems that a(n) = 1 only for n = 0, 1, 2, 3, 5, 7, 14, 15, 30, 120, 130, 754, 770, 1186, 2290.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017405 | OEIS-A390791

Number of Hamiltonian paths from vertex (1, 1) to (3, 2*n+2) on the product graph C_4 X P_{2*n+2}, using (row, column) numbering.

Conjecture: G.f.: 2 * (3 - 13*x - 2*x^2)/(1 - 21*x + 26*x^2 - 44*x^3 + 8*x^4).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Feb 12 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017406 | OEIS-A390793

Number of ways to write n as x^4 + y^2 + 2*z^2 + (p-1)^2/2, where x,y,z are in the set {m >= 0: 2*m is practical}, and p is either 1 or an odd prime.

Conjecture: a(n) > 0 for every natural number n. We have verified a(n) > 0 for n <= 4*10^5. As 2*z^2 + 2*w^2 = (z+w)^2 + (z-w)^2, the conjecture is stronger than the author's previous refinement of Lagrange's four-square theorem which states that each natural number can be written as x^4 + y^2 + z^2 + w^2 with x,y,z,w integers. The conjecture that a(n) > 0 for every natural number n is false. The following values of n have a(n) = 0: 3670016, 7864320, 12058624, 14680064, 15728640, 16252928, 19398656, 20447232, 22020096, 24641536, 28573696, 28835840, 30408704, 31457280, 32505856, 48234496, 49283072, 55574528, 58720256, 63963136, 65011712, 66060288, 74448896. These were verified by exhaustive enumeration of the equivalent representation n = x^4 + w^2 + 2*y^2 + 2*t^2, where x,w,y are nonnegative integers such that 2*x,2*w,2*y are practical numbers, and t = 0 or 2*t + 1 is prime. - _Scott Moore_, May 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017407 | OEIS-A390795

Number of ways to write n as ((p-1)/2)^4 + x^2 + 2*y^2 + 3*z^2, where p is 1 or an odd prime, and x,y,z are in the set {m >= 0: 2*m is practical}.

Conjecture: a(n) > 0 for every natural number n. This has been verified for n <= 2*10^5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017408 | OEIS-A390799

Number of ways to write n as (p-1)^4/16 + (q-1)^2/2 + u^2 + 3*v^2, where each of p and q is 1 or an odd prime, and u and v are in the set {m >= 0: 2*m is practical} with u > 0.

Conjecture: a(n) > 0 for all n > 0. This has been verified for n <= 2*10^5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017409 | OEIS-A390815

Denominators > 1 of ratio n!/sigma(n), where n! = factorial of n (A000142) and sigma(n) = sum of divisors of n (A000203).

Conjecture: all terms are prime (verified for n <= 10^5). Conjecture: for n >= 4 if sigma(n) does not divide n!, then n is a square. (verified for n <= 10^5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Claude H. R. Dequatre_, Feb 17 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017410 | OEIS-A390818

Number of ways to write n as 3*((p-1)/2)^4 + (q-1)^2/2 + (r-1)^2/4 + s^2, where each of p,q,r is either 1 or an odd prime, and s is half of a practical number.

Conjecture 1: a(n) > 0 for all n > 0. We also have the following similar conjecture. Conjecture 2: Each natural number n can be written as 3*u^4 + v^2 + (p-1)^2/2 + (q-1)^2/4, where each of u and v is either zero or half of an even practical number, and each of p and q is either 1 or an odd prime. Both conjectures have been verified for n <= 2*10^5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 20 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017411 | OEIS-A390839

a(n) = Product_{k=1..n} (prime(k) + 1)/A001223(k), where A001223 lists the prime gaps, rounded to the nearest integer if necessary.

The product is actually conjectured to be an integer for all n. Can anyone prove it?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Nov 21 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017412 | OEIS-A390871

Numbers k >= 1 such that k^2 - r^2 = 2^m - 1 for some 0 <= m <= k and 0 <= r < k.

For k != A000079(n), if there is a solution for k > 8, then m is from [A000523(k) + 3, 2*A000523(k) + 1] (I conjecture).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ctibor O. Zizka_, Nov 22 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017413 | OEIS-A390878

Convolution of A000041 and A237271.

Conjecture: a(n) is the total number of 2-dense sublists of divisors of the positive integers whose divisors are the parts of the partitions of n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Nov 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017414 | OEIS-A391017

Starting values k of Collatz orbits that achieve a new minimum of Product_{j == 4 mod 6 in "3x+1" orbit of k} (j-1)/j.

It is conjectured that A391016 is a lower bound of these products.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_ and _Markus Sigg_, Nov 26 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017415 | OEIS-A391037

Number of ways to write n as ((p-1)/2)^4 + (q-1)^2/2 + (r-1)^2/4 + (s+1)^2/4, where each of p,q,r,s is either 1 or an odd prime.

Conjecture: For any c = 1, 5, 7, every positive integer n can be written as c*((p-1)/2)^4 + (q-1)^2/2 + (r-1)^2/4 + (s+1)^2/4, where each of p,q,r,s is either 1 or an odd prime. See also A391038 for a smilar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 26 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017416 | OEIS-A391038

Number of ways to write n as (p-1)^4/8 + (q-1)^2/4 + (r-1)^2/4 + (s+1)^2/4, where each of p,q,r,s is either 1 or an odd prime.

Conjecture: a(n) > 0 for all n > 0. See also A391037 for a similar conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Nov 26 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017417 | OEIS-A391055

a(n) is the number of iterations of x-> gpf(3*x+1) starting at n until the value 2 is reached; a(n) = -1 if 2 is not reached.

Conjecture: a(n) is never equal to -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Nov 26 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017418 | OEIS-A391072

Numbers of twin prime candidates in primorial interval of prime(n)# after elimination by prime factor prime(n-1).

Conjecture: This number goes to infinity when n approaches infinity. Conjecture: The number of twin prime candidates remaining in the primorial(n)# interval after sieving with the prime(n) is given by A059861.This number goes to infinity when (n) approaches infinity.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Leif Nordqvist_, Nov 27 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017419 | OEIS-A391120

a(n) is the least k for which n^k contains at least one decimal digit 9, or -1 if there is no such k.

Conjecture: a(n) <= 12, with a(n) = -1 iff n is in A011557 and a(n) = 12 iff n/2 is in A011557.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Nov 29 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017420 | OEIS-A391205

a(n) is the least k for which n^k contains at least one decimal digit 7, or -1 if there is no such k.

Conjecture: a(n) <= 15, with a(n) = -1 iff n is in A011557 and a(n) = 15 iff n/2 is in A011557.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Dec 03 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017421 | OEIS-A391216

a(n) is the least k for which omega(k)*omega(k + 1) = n where omega = A001221.

Conjecture: For p prime, a(p) = A098026(p). - _Vaclav Kotesovec_, Jan 03 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jan 03 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017422 | OEIS-A391251

Numbers k such that the square of Stern polynomial B(k,x) is also a Stern polynomial.

The odd terms 1, 3, 7, 15, 27, 31, 63, 127, 255, 511, 1023, 2047, 4095, 8191, ..., are conjectured to be given by binary palindromes (A006995) that are equal to A000225(n)^(2*e-1), for some n, e >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017423 | OEIS-A391269

a(n) is the number of iterations of x-> gpf(2*x+1) starting at n until the value 3 is reached; a(n)=-1 if 3 is not reached.

Conjecture: a(n) is never equal to -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Dec 04 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017424 | OEIS-A391282

Primitive e-semiproper perfect numbers: powerful numbers k for which A323309(k) = 2*k.

Conjecture: a(n) = A064549(A002827(n)) (see the Comments).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Dec 05 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017425 | OEIS-A391317

a(n) is the smallest positive integer that has at least one subset of proper divisors that adds to the n-th weird number (A006037(n)).

This sequence is conjectured to be a subsequence of A005101.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Dec 16 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017426 | OEIS-A391346

Numbers k such that Stern polynomial B(5*k,x) is a multiple of B(5,x).

a(n) = A391566(n)/5. [Conjectured]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017427 | OEIS-A391473

a(n) is the smallest positive integer k such that k*R_n has digits with alternating parity, where R_n is the n-th repunit.

Conjecture:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gonzalo Martínez_, Dec 10 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017428 | OEIS-A391538

Guaranteed win if opponent chooses one of three baskets and the player chooses the coins with values from 1 to n (see Comments for details).

If n>4, a(n) = floor(n*(n-1)/6) + n (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ruediger Jehn_, Dec 12 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017429 | OEIS-A391563

a(n) is the number of ways to write n = p + q where p is prime and q is the reverse of a prime.

Conjecture: a(n) >= 1 for n >= 12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Feb 15 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017430 | OEIS-A391566

Numbers k such that Stern polynomial B(k,x) is a multiple of B(5,x).

a(n) = 5*A391346(n). [Conjectured]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 14 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017431 | OEIS-A391602

a(n) is the largest number m such that A391449(m) = n.

a(0) through a(2) computationally verified to m = 2*10^14 (A141399 verified a(0) to 2.29*10^25). We conjecture this is sufficient to justify the finality of a(0) and a(1), and likely the finality of a(2). See the distribution of delta values by running the Octave_Deltas.c program at the attached link. Conjecture: All polynomial generator functions of order 2 or greater, G(n), with integer roots, and that are not perfect powers, are bounded in prime-complete numbers (A055932). Consequences: Since primorials are prime-complete by definition and verification shows no primorial products m*(m+1) for 715 <= m <= 633555, this suggests (714, 715) is the last pair of consecutive integers with primorial product. For Brocard's equation n! + 1 = m^2, since factorials are prime-complete and n! = (m-1)*(m+1), note that for odd m, (m-1)*(m+1) = 4*k*(k+1) where k = (m-1)/2, and delta(4*k*(k+1)) = delta(k*(k+1)) (multiplying by 4 just adds more powers of 2). Thus prime-complete (m-1)*(m+1) requires delta(k*(k+1)) = 0. Since a(0) = 633555 is the last k with delta(k*(k+1)) = 0, the bound m <= 1267111 (the final term of A194099) closes the search for solutions to Brocard's equation.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ken Clements_, Dec 13 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017432 | OEIS-A391613

Number of ways to write n as x^2 + y + z with x >= 1 and y >= z >= 1 such that lcm(x^2,y,z) is a square.

Conjecture: a(n) > 0 for all n > 5

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 17 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017433 | OEIS-A391620

Number of integer partitions of n > 0 that are not the first sums of any composition with all parts > 1.

Conjecture: These are integer partitions of n whose least part is < 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Dec 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017434 | OEIS-A391629

Number of multisets summing to n (or reversed partitions of n) that are not the first sums of any integer partition.

Conjecture: a(n) = A000041(n) - A368684(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Dec 31 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017435 | OEIS-A391632

Number of mutual-visibility sets in the n-Andrásfai graph.

Conjectured g.f.: x*(4 - 19*x + 33*x^2 - 40*x^3 + 8*x^4)/(1 - 10*x + 29*x^2 - 32*x^3 + 8*x^4). - _Andrew Howroyd_, Jan 12 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Dec 14 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017436 | OEIS-A391714

Numerators of the convergents given by treating A072193 as continued fraction coefficients after the leading 0.

Conjecture: Limit_{n->oo} a(n)^(1/n) = Lévy's constant (A086702). - Corrected by _Jwalin Bhatt_, Jun 23 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Dec 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017437 | OEIS-A391715

Denominators of the convergents given by treating A072193 as continued fraction coefficients after the leading 0.

Conjecture: Limit_{n->oo} a(n)^(1/n) = Lévy's constant (A086702). - Corrected by _Jwalin Bhatt_, Jun 23 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Dec 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017438 | OEIS-A391716

Let f be an additive function with f(2^e) = (e + e mod 2)/2, f(p^e) = e*(p-1)/2, p an odd prime. a(n) is the number of iterations of f(n) required to reach 0.

Every n appears a finite number of times in this sequence. Let S_n be the set of numbers k such that a(k) = n. We can prove that S_n is finite by an induction on n = 1, since S_1 = {1} as k = 1 is the only k such that f(k) = 0, and thus a(k) = 1. Let the induction hypothesis be that S_i has a maximum element M_i for all 1 <= i <= n. If r is in S_{n+1}, then f(r) is in S_n. We will prove that if f(r) = M_n, then r <= 2^(2*M_n). We note that for an arbitrary prime power p^e, f(p^e) = e*(p-1)/2 >= log_2(p^e)/2. Let r be composite. Then f(r) = M_n = Sum f(p^e) >= (Sum log_2(p^e))/2 = log_2(r)/2. Hence log_2(r) <= 2*M_n, and r <= 2^(2*M_n). Since f(2^(2*M_n)) = M_n, therefore 2^(2*M_n) is in S_{n+1}. If t is in S_{n+1} and t > 2^(2*M_n) and f(t) < M_n, then log_2(t)/2 <= f(t) < M_n. But then log_2(t) < 2*M_n and t < 2^(2*M_n), which contradicts the assumption. Consequently if t is in S_{n+1} and f(t) < M_n, then t < 2^(2*M_n). Therefore 2^(2*M_n) is the maximum element of S_{n+1} and S_{n+1} is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Miles Englezou_, Jan 25 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017439 | OEIS-A391736

Square array read by descending antidiagonals: A(n, k) is the value of quotient A048720(n,i)/i for the k-th odd number i which divides A048720(n,i), where A048720 is carryless base-2 multiplication.

The number of distinct values that may occur on row n of this array is at least 1+A115861(n), and at most n, because A(n, k) <= n. We conjecture that 1+A115861(n) gives the exact value.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Dec 18 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017440 | OEIS-A391769

The number of gaps in the set of positive integers which need at most n steps of the Collatz iteration to reach 1.

It appears that this sequence has the same asymptotic growth rate (3 + sqrt(21))/6 (A176014) as is conjectured for A005186.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Markus Sigg_, Dec 19 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017441 | OEIS-A391775

Discriminants of totally real quartic fields in which every additively indecomposable element is a unit.

It is an open problem whether this sequence has infinitely many terms. All known terms correspond to totally real quartic fields of the form Q(sqrt(A + B*sqrt(D))) with D = 3 or 5. A special case of a conjecture of Siu Hang Man [Man 2024, Conjecture 1.7] states that there are only finitely many real biquadratic fields in which every additively indecomposable element is a unit.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robin Visser_, Mar 13 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017442 | OEIS-A391791

a(n)/2^(n-1) is the expected win if one of two baskets is chosen randomly and the player optimally chooses the coins with values from 1 to n (see Comments for details).

Conjecture: For n>9 the optimal strategy is to put coin floor((n+3)/2) in the first basket.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ruediger Jehn_, Dec 20 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017443 | OEIS-A391794

Coefficients in the expansion of A(q) = lim_{M->inf} Sum_{n=0..2M} (-1)^n / ((q^2;q)_n * (q^2;q^2)_n).

Conjectured from n <= 10000: log(a(n)) = Pi*sqrt(n) - (7/4)*log(n) + C + O(n^(-1/2)), C ~ -2.34. Conjectured: a(n) ~ (Pi/4)*b(n)/sqrt(n), where b(n) = A002513(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alejandro Radisic_, Feb 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017444 | OEIS-A391805

a(n) = greatest prime less than prime(n)*prime(n+1).

Conjecture: If k is an even integer greater than 4, then A391804(n) - a(n) = k for infinitely many n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 04 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017445 | OEIS-A391810

Irregular triangle, read by rows, where row n lists the coefficients of nonpositive powers of x in the doubly infinite series Sum_{k=-oo..oo} x^k * (1 - x^k)^(n+k) for n >= 1.

Conjectural formulas for the terms of this triangle: (2) Sum_{k=0..floor((n+1)/2)^2} T(n,k) = 0 for n >= 1 (conjecture). (3) Sum_{k=0..floor((n+1)/2)^2} k * T(n,k) = 1 for n >= 1 (conjecture). (4) Sum_{k=0..floor((n+1)/2)^2} k^2 * T(n,k) = -(2*n-5) for n >= 2 (conjecture). (5) Sum_{k=0..floor((n+1)/2)^2} -(-1)^k * T(n,k) = 2*round(2^n/3) = 2*A001045(n) for n >= 1 (conjecture), where A001045 is the Jacobsthal numbers (g.f. x/((1+x)*(1-2*x))). (6) Sum_{k=0..floor((n+1)/2)^2} -(-1)^k

* k * T(n,k), n >= 1, forms a sequence with g.f. x*(1-x+8*x^2)/((1+x)^2*(1-2*x)^2) (conjecture), and begins [1, 1, 13, 25, 81, 181, 453, 1025, ...].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Dec 21 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017446 | OEIS-A391906

Numerators of the convergents given by treating A390946 as continued fraction coefficients after the leading 0.

Conjecture: Limit_{n->oo} a(n)^(1/n) = Lévy's constant (A086702). - Corrected by _Jwalin Bhatt_, Jun 18 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Dec 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017447 | OEIS-A391907

Denominators of the convergents given by treating A390946 as continued fraction coefficients after the leading 0.

Conjecture: Limit_{n->oo} a(n)^(1/n) = Lévy's constant (A086702). - Corrected by _Jwalin Bhatt_, Jun 23 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Dec 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017448 | OEIS-A391938

Numbers k not divisible by p^p for any prime p, and for which A373842(k) >= k, where A373842 is the arithmetic derivative of the primorial base log-function.

After the initial 5, conjectured to be the intersection of A048103 with the complement of A373847, or equally, complement of (A100716 U A373848).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 15 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017449 | OEIS-A391939

Composite numbers k such that A003415(k) <= A276086(k), where A003415 is the arithmetic derivative and A276086 is the primorial base exp-function.

If the conjecture mentioned in A351228 is correct, then, apart from {6}, this is the intersection of {the complement of A351228} and A002808 (composite numbers).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 15 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017450 | OEIS-A391958

First column of A391957.

Conjectured: this sequence can be obtained by the following two rules:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _A.H.M. Smeets_, Dec 23 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017451 | OEIS-A391959

Second column of A391957.

Conjectured: this sequence can be obtained by the following two rules:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _A.H.M. Smeets_, Dec 28 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017452 | OEIS-A392002

Number of partitions of n with parts colored by {0, 1} such that the sum of colors is congruent to 1 (mod 2).

Conjectured g.f.: (1/2) * (Product_{j>=1} 1/(1-x^j)^2 - Product_{j>=1} 1/(1-x^(2*j))). Conjectured g.f.: (1/2) * (G.f. of A000712(x) - G.f. of A000041(x^2)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas Hutton_, Dec 26 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017453 | OEIS-A392003

a(n) = (7*2^n - 9 + 5*(-1)^n) / 6.

Conjecture: a(2*n) = A266753(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Karl-Heinz Hofmann_, Dec 26 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017454 | OEIS-A392058

Array read by downward antidiagonals: A(n,k) = A(n-1,k+1) + Sum_{j=0..k} binomial(k+1,j)*A(n-1,j)*A(k-j,0) with A(0,k) = 1.

Conjecture: A(n,0) = A233335(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Dec 29 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017455 | OEIS-A392095

Array read by downward antidiagonals: A(n,k) = A(n-1,k+1) + Sum_{j=0..k} A(n-1,j)*A(k-j,0) with A(0,k) = 1.

Conjecture: A(n,0) = A088713(n+1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, Dec 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017456 | OEIS-A392096

Numbers k for which the width pattern of the symmetric representation of sigma for k, SRS(k), has the form 1 2 1 0 1 2 1 0 . . . 0 1 2 1 0 1 2 1 containing at least one 0.

Conjecture: All numbers k in this sequence are abundant with 2k < sigma(k) < 8k/3; true for k <= 10^9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Dec 30 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017457 | OEIS-A392107

a(n) = position of first occurrence of the n-th binary word (using letters 1 and 2 as in A392106) in the sequence of residues modulo 3 of the primes.

Conjecture: every positive integer occurs infinitely many times. In particular, the words that start in position 1 (i.e., the positions of 4 in the sequence), are indexed as 1, 4, 9, 20, 41, 84, 170, 341, 683, 1368, 2737, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017458 | OEIS-A392116

a(n) = f(n+1) - f(n), where f(n) = floor(prime(n)/5).

Conjecture: every nonnegative integer occurs.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jan 18 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017459 | OEIS-A392131

a(n) is the determinant of the n X n circulant matrix whose rows are formed by successively rotating the vector (A000129(0), A000129(1), ..., A000129(n-1)) right.

Conjecture: lim_{n->oo} A392132(n)/a(n) = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Dec 31 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017460 | OEIS-A392132

a(n) is the permanent of the n X n circulant matrix whose rows are formed by successively rotating the vector (A000129(0), A000129(1), ..., A000129(n-1)) right.

Conjecture: lim_{n->oo} a(n)/A392131(n) = 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Dec 31 2025 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017461 | OEIS-A392135

Third column of A391957.

Conjectured: a(j*5^m + 5^(m-1)*k) = a(j*5^m) + k*(a(5^(m-1)) - 2*5^(m-1)*Sum_{i = 1..j} A055457(i)), 0 < k < 5, where a(j*5^m) = j*a(5^(m-1)) for j < 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _A.H.M. Smeets_, Jan 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017462 | OEIS-A392136

Fourth column of A391957.

Conjecture: a(j*7^m + 7^(m-1)*k) = a(j*7^m) + k*(a(7^(m-1)) - 2*7^(m-1)*Sum_{i = 1..j} A373217(i)), 0 < k < 7, where a(j*7^m) = j*a(7^(m-1)) for j < 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _A.H.M. Smeets_, Jan 05 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017463 | OEIS-A392163

Terms of A055932 that are products of two or more consecutive composite integers.

This sequence is particularly significant because it limits the possible solutions to the Suranyi-Hickerson Conjecture (see Hajdu et al.) to the exact known:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ken Clements_, Jan 30 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017464 | OEIS-A392184

a(n) is the least k for which omega(k)*omega(k + 1)*omega(k + 2) = n where omega = A001221, or -1 if no such k exists.

Conjecture: a(p) = A075590(p) - 1. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Juri-Stepan Gerasimov_, Jan 02 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017465 | OEIS-A392191

a(n) is the minimum determinant of an n X n circulant matrix whose rows are permutations of [0, 1, 2, ..., n-1].

Conjecture: a(n) = A084367((n-1)/2) for n odd.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jan 03 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017466 | OEIS-A392200

a(n) = gcd(A381466(n-1), n).

We conjecture that a(n) < log(n+1)/log(1.2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sam Chapman_, Mar 16 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017467 | OEIS-A392208

E.g.f. A(x) satisfies A'(x) = exp(A(x)*A'(x)^3), with A(0)=0.

Conjecture: a(n) is even iff n is divisible by 3 for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 24 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017468 | OEIS-A392211

G.f. satisfies A(x) = (1 - sqrt(1 + 4*A(-x) - 4*A(-x)^2))/2.

Conjecture D-finite with recurrence 3840*n*(7977321*n -32086520)*(n-1)*(n-2)*a(n) -64*(n-1)*(n-2)*(953294144*n^2 -8954569952*n +20118837885)*a(n-1) -15*(n-2)*(75138386499*n^3 -602776477876*n^2 +1515217257713*n -1239100034728)*a(n-2) +8*(280596173198*n^4 -4319298738497*n^3 +24426073694308*n^2 -60489429217342*n +55587559491135)*a(n-3) +24000*(2*n-9)*(15954642*n^3 -135968929*n^2 +388189337*n -380277463)*a(n-4) -12800*(29790442*n-101087659) *(n-5)*(2*n-9)*(2*n-11)*a(n-5)=0. - _R. J. Mathar_, Feb 24 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 10 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017469 | OEIS-A392241

a(n) is the 10^n-th term of A389544.

Illustrative of the density conjecture of the 421st Erdős Problem. a(7)..a(9) conjectured to be 10003376, 100010472, 1000032642, respectively, based on using the program in A389544, but only storing products <= 10^100 for a(7)..a(8) and <= 10^20 for a(9).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael S. Branicky_, Jan 04 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017470 | OEIS-A392283

Numbers k such that phi(k) divides k + 3.

Conjecture: terms in this sequence after 9 are identical to A039777.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tyler Yu_, Jan 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017471 | OEIS-A392319

Palindromes of the form b^k + 1 or b^k - 1, with b, k > 1.

This sequence explores the predecessors and successors of perfect powers. Sequence A075786 gives the perfect powers that are palindromes, where it is conjectured that there are no palindromic perfect powers with prime exponent > 3. Without considering the case when b = 10, this sequence records prime exponents greater than 3, for example, 2^5 + 1 = 33, 3^5 - 1 = 242, and 6^5 + 1 = 7777.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gonzalo Martínez_, Jan 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017472 | OEIS-A392329

a(n) is the number of iterations x->gpf(6*x+1) starting at n until the value 13 or 47 is reached; a(n) = -1 if neither of these two values is reached.

Conjecture: a(n) is never equal to -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Jan 07 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017473 | OEIS-A392338

a(n) is the minimum absolute value of determinant of a nonsingular n X n circulant matrix whose rows are permutations of [0, 1, 2, ..., n-1].

Conjecture: a(n) = A084367((n-1)/2) for n odd.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jan 07 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017474 | OEIS-A392344

a(n) is the greatest integer m such that m*(m+n) has a complete contiguous set of prime divisors from 2 to its greatest prime divisor.

All listed terms except a(1) are conjectural. For the initial terms given, it has been computationally verified that if a greater a(n) exists, it is at least 10000 times larger than the listed value.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ken Clements_, Jan 07 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017475 | OEIS-A392363

Smallest connected polyiamond containing all free n-iamonds under rigid motions of the triangular lattice.

Conjecture: lim_{n -> infinity} a(n)/n^2 = 1/5. The ratio a(n)/n^2 decreases from 0.240 at n=10 to 0.214 at n=14 (monotonic in this range). For all n <= 14, a(n) <= ceiling((n^2+2n)/5), with equality for n <= 8 and n=10, and a(n) is 1-3 less for n=9,11,12,13,14.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Exley_, Apr 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017476 | OEIS-A392387

Nullspace dimension of the periodic XX Heisenberg chain with n > 1 sites.

Conjecture: a(2*p) = 2*(6^((p-1)/2)+1) for odd prime p.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thore Posske_, Jan 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017477 | OEIS-A392406

a(n) is the maximum possible number of extensions of a sequence of any n integers, where an extension is the runs-resistance of the sequence including all previous extensions.

Conjecture: No other stopping criteria are necessary to ensure that any sequence will terminate, which also implies that a cycle with period > 1 will never appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Frederik G. Faye_, Jan 10 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017478 | OEIS-A392479

Pairs of prime numbers (p,q), p minimal, such that prime(n) = p^3 - 2*q, or (0,0) if no such pair exists.

Conjecture: Every prime number is of the form p^3 - 2*q for some primes p and q.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Jan 13 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017479 | OEIS-A392498

Numbers k > 1 with the smallest prime divisor p, where the sum of divisors d in the range 1 < d < k/p is less than k/p.

Conjecture: All perfect numbers are sparse. This conjecture is consistent with all known data (A000396).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jan 16 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017480 | OEIS-A392505

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest unused number such that a(n) shares a factor with a(n-1) while rad(a(n)*a(n-1)) has not previously occurred as the rad value of the product of adjacent terms, where rad = A007947.

In the first 500K terms there are no fixed points other than 1 and 2. The sequence is conjectured to be a permutation of the positive numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Mar 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017481 | OEIS-A392508

a(n) is the number of iterations prime(x) -> gpf(2*prime(x)+prime(x+1)) starting at prime(n) until the value prime(3) = 5 is reached where gpf(x) is the greatest prime factor of x; a(n) = -1 if the value 5 is not reached.

Conjecture: a(n) is never equal to -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Jan 14 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017482 | OEIS-A392525

G.f. satisfies: A(x) = A(x^3 + 15*x*A(x)^3)^(1/3), with A(0)=0, A'(0)=1.

Conjecture: a(n) is odd iff n = 2^k for k >= 0. Conjecture: a(n) == 5 (mod 10) iff n = 2^k for k >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jan 15 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017483 | OEIS-A392548

a(n) is the least positive integer k such that Mordell's equation y^2 = x^3 - k^2 has exactly n integer solutions with y >= 0.

Conjecture: a(n) is even for all n>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhining Yang_, Jan 15 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017484 | OEIS-A392592

Intersection of A391845 and A391866.

A369650 contains numbers k such that A003415(k) = A276085(k). It is an open question whether that sequence contains any other terms after 5005, and especially, if it contains any terms of A371083. On any such hypothetical terms x of A369650, the two functions A003415 and A276085 should satisfy all the conditions of the other function that function satisfies on the terms of A371083, namely, that A020639(x) is mapped to A053669(x), and that the divisor A003557(x) is retained in the result.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 17 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017485 | OEIS-A392603

Numbers k such that A380459(A276086(k)) has no divisors of the form p^p, for any prime p.

It is conjectured that A267263(a(n)), and equally, A276150(a(n)), is always <= 4. See A380475.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017486 | OEIS-A392604

Numbers k such that A380459(A019565(k)) has no divisors of the form p^p, for any prime p.

It is conjectured that A000120(a(n)) is always <= 4. See A380475.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017487 | OEIS-A392606

Intersection of A392605 and A392611.

It is conjectured that this has no common terms with A380527, even though both are subsequences of A381035. And the above conjecture was soon refuted, as x = A276085(855523) = 55946999363570987235666961212549502632097773O can be found in both sequences. See my Jan 28 2026 comment in A380527. - _Antti Karttunen_, Oct 10 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Jan 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017488 | OEIS-A392664

Triangle read by rows: T(n,k) (1 <= k <= n) given by T(n,1) = T(n,n) = 1, otherwise T(n,k) = (m*n-m*k+1)*T(n-1,k-1) + (m*k-m+1)*T(n-1,k), where m = -3.

Conjecture: T(n,k) >= 0 for all even n and all 1 <= k <= n. Verified for even n <= 400. Since all entries are nonneg for even n, the roots of R_n(y) lie in the annulus determined by consecutive coefficient ratios (Enestrom-Kakeya theorem).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Xin Chen_ and Yudian Chen, Jan 18 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017489 | OEIS-A392667

Least positive integer m such that tau(k)*(tau(k)-1) (k = 1..n) are incongruent modulo m, where tau(.) is Ramanujan's tau function.

Conjecture: a(n) is prime except for n = 1, 37.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 18 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017490 | OEIS-A392672

Smallest number m such that m*(m+1)*(m+2) has exactly n distinct prime factors.

Conjecture: this is an increasing sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhining Yang_, Apr 30 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017491 | OEIS-A392682

Primes p(k) > 3 such that, for every integer 1 < m < k (except m = k - 2 if p(k) > 5), there exists a set of m distinct primes < p(k) the sum of whose reciprocals forms a fraction that has a numerator divisible by p(k).

Conjecture: this is the intersection of A006512 and A293457.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Jan 19 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017492 | OEIS-A392684

Number of ways to write n as x + y + z with x >= y >= z >= 1 such that lcm(x,y,z) is a product of three consecutive integers.

Conjecture: (i) For any integer k > 2, each integer n >= 2^k - 2 can be written as x_1 + ... + x_k with x_1,...,x_k positive integers such that lcm(x_1,...,x_k) is a product of k consecutive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 19 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017493 | OEIS-A392694

Numbers whose weakly increasing prime indices are the 0-prepended first sums of some reversed integer partition.

Conjecture: Reversed integer partitions of this type (with weakly increasing 0-based partial alternating sums) are counted by A000009. For example, the A000009(1) = 1 through A000009(9) = 8 reversed partitions are:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jan 25 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017494 | OEIS-A392698

Numbers whose reversed prime indices are the 0-prepended first sums of some finite sequence of distinct integers.

Conjecture: Integer partitions of this type (distinct trimmed 0-based partial alternating sums) appear to be counted by A000009. For example, the A000009(1) = 1 through A000009(9) = 8 partitions are:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Jan 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017495 | OEIS-A392707

Numbers whose prime indices have strictly increasing 0-based partial alternating sums.

Conjecture: Reversed integer partitions of this type (with strictly increasing 0-based partial alternating sums) are counted by A053251.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gus Wiseman_, Feb 05 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017496 | OEIS-A392714

a(n) is the positive integer such that Sum_{sigma in S_{2n}} sign(sigma) * w_{sigma(1)}*D^n o w_{sigma(2)}*D^n o ... o w_{sigma(2n)}*D^n = a(n) * Wronskian(w_1,...,w_{2n}) * D^n, where D = d/dx and w_1,...,w_{2n} are smooth functions on the real line.

For prime p, p^(p-1) divides a(p) (proved). Conjecture: the p-adic valuation of a(p) equals exactly p-1 for every prime p. Among the contributing permutations, the counts of even and odd permutations differ by exactly 1 for n <= 13; conjectured to hold for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kian Shah_, Jul 05 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017497 | OEIS-A392732

Least positive integer m such that 2^(k*(k-1)/2) (k = 1..n) are pairwise incongruent modulo m.

Conjecture: Let c be any integer with |c| > 1, and let D(c,n) be the least positive integer m such that the n numbers c^(k*(k-1)/2) (k = 1..n) are pairwise incongruent modulo m.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017498 | OEIS-A392775

Least positive integer m such that 2*3^(k*(k-1)/2) (k = 1..n) are pairwise incongruent modulo m.

Conjecture: Let c be any integer congruent to 3 or -3 modulo 8. For any positive integer n, let m(c,n) be the least positive integer m such that the n numbers 2*c^(k(k-1)/2) (k = 1..n) are pairwise incongruent modulo m. This is a supplement to the conjecture in A392732.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhi-Wei Sun_, Jan 22 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017499 | OEIS-A392819

Numbers k such that p + e divides k for every prime power p^e in the canonical prime factorization of k.

Conjecture: This sequence has natural density 0: the divisibility conditions are stronger than for refactorable numbers (which already have density 0); in particular, for k = Product_{i} p_i^{e_i}, each (p_i + e_i) must be p_n-smooth and divide k. Natural density 0 0 (conjectured)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Feb 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017500 | OEIS-A392873

Nonsquarefree numbers k such that A276086(A003415(k)) <= k, where A003415 is the arithmetic derivative and A276086 is the primorial base exp-function.

It is conjectured that there are no common terms with A370115.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Feb 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017501 | OEIS-A392976

Expansion of (1/x) * Series_Reversion(x * ((1-x^2)^2 - x)).

Conjecture D-finite with recurrence 283*n*(n-1)*(n-2)*(83022305213*n -178385058984)*(n+1)*a(n) -6*n*(n-1)*(n-2)*(12196786904129*n^2 -45110300936020*n +52641391340361)*a(n-1) -6*(n-1)*(n-2)*(67790996421377*n^3 -248299346995054*n^2 +202497447227007*n +3061418504418)*a(n-2) -2*(n-2)*(192167357944360*n^4 -559010679688491*n^3 -2120668219560541*n^2 +912336136130610*n -8342981628089220)*a(n-3) +5*(15640122296875*n^5 -565964885220500*n^4 +4989857715427325*n^3 -18667678380008380*n^2 +32085126156301848*n -20925325629575136)*a(n-4) +10*(5*n-17)*(5*n-18) *(5*n-21)*(5*n-19)*(22509687685*n-60308240636)*a(n-5)=0. - _R. J. Mathar_, Jan 29 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Seiichi Manyama_, Jan 29 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017502 | OEIS-A392982

Run lengths of consecutive increases of prime(n)/n where prime(n) is the n-th prime.

Conjecture: For every positive integer k there exists m with a(m) = k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philip Jameson_, Mar 07 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017503 | OEIS-A393024

Numbers that can be written as s_1^x_1 + ... + s_t^x_t, with 0 < s_1 < ... < s_t and x_1 > ... > x_t > 1 for some t > 0, with the restriction that if s_1 = 1 then x_1 is the least possible.

All natural numbers greater than 1190 appear in this sequence (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alberto Zanoni_, Jan 31 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017504 | OEIS-A393028

Numbers that can be written in more than one way as s_1^x_1 + ... + s_t^x_t, with 0 < s_1 < ... < s_t and x_1 > ... > x_t > 1 for some t > 0, with the restriction that if s_1 = 1 then x_1 is the least possible.

Every integer greater than 1543 appears in the sequence (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alberto Zanoni_, Jan 31 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017505 | OEIS-A393029

Numbers that can be written in exactly two ways as s_1^x_1 + ... + s_t^x_t, with 0 < s_1 < ... < s_t and x_1 > ... > x_t > 1 for some t > 0, with the restriction that if s_1 = 1 then x_1 is the least possible.

Conjecture: The sequence is finite and its last number is 1446.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alberto Zanoni_, Jan 31 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017506 | OEIS-A393030

Numbers that can be written in exactly three ways as s_1^x_1 + ... + s_t^x_t, with 0 < s_1 < ... < s_t and x_1 > ... > x_t > 1 for some t > 0, with the restriction that if s_1 = 1 then x_1 is the least possible.

Conjecture: The sequence is finite and its last number is 1799.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alberto Zanoni_, Jan 31 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017507 | OEIS-A393031

Numbers that can be written in exactly four ways as s_1^x_1 + ... + s_t^x_t, with 0 < s_1 < ... < s_t and x_1 > ... > x_t > 1 for some t > 0, with the restriction that if s_1 = 1 then x_1 is the least possible.

Conjecture: The sequence is finite and its last number is 1919.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alberto Zanoni_, Jan 31 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017508 | OEIS-A393032

Numbers that can be written in exactly five ways as s_1^x_1 + ... + s_t^x_t, with 0 < s_1 < ... < s_t and x_1 > ... > x_t > 1 for some t > 0, with the restriction that if s_1 = 1 then x_1 is the least possible.

Conjecture: The sequence is finite and its last number is 2879.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alberto Zanoni_, Jan 31 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017509 | OEIS-A393039

Maximum Mahler measure of all quartic polynomials with coefficients in {-1, 0, 1}.

Conjecture 1: For every d M(d+1) > M(d). Conjecture 2: M(d) < sqrt(d+1). - _Charles R Greathouse IV_, Feb 05 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Jan 31 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017510 | OEIS-A393104

a(n) is the minimum total surface area of five-element sets of distinct integer-sided cuboids that fill an n X n X n cube.

Conjecture: a(n) = 2*(A054567(n+1) + 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Janaka Rodrigo_, Feb 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017511 | OEIS-A393110

a(n) = G(pi(10^n)) where pi is A000720 and G(n) is the "depth" of n in the sense of the Gilbreath-Proth conjecture.

a(n) = G(pi(10^n)) where pi is A000720 and G(n) is the "depth" of n in the sense of the Gilbreath-Proth conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Marcus_, Feb 02 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017512 | OEIS-A393111

Distance from 2^n to the nearest base-10 palindrome.

Gustavus Simmons has conjectured that there are no palindromes of the form b^k with k >= 5 and b >= 2 (see Wikipedia link). If the conjecture is true, a(n) > 0 for all n >= 4. Conjecture: a(n) = 2^(n/2+O(log n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pontus von Brömssen_, Feb 02 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017513 | OEIS-A393125

Maximum of the orbit of n under the map x -> 5x+1 if odd else x/2; -1 if the orbit is unbounded.

It is conjectured but still unproven that a(7) = -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Feb 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017514 | OEIS-A393128

Positive integers not of the form floor(k + log(k) + log(Pi)).

It was conjectured that for every prime p there is exactly one integer k such that p = k + floor(log(Pi*k)), but this is not true for the primes in this sequence, the first of which are 19069, 140837, 56812699, ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Mar 17 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017515 | OEIS-A393131

a(n) = EulerPhi(2*n+1)/A329726(n); index of witnesses for Solovay-Strassen primality test of 2*n+1 as a subgroup of (Z/(2*n+1)Z)*.

Conjecture: Every even number occurs infinitely many times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Mar 12 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017516 | OEIS-A393170

G.f. A(x) satisfies [x^n] 1/(A(x) - n*x) = 0 for n >= 1.

Conjecture: for n > 0, a(n) is odd iff n is a power of 2. Conjecture: a(n) == 2 (mod 4) iff n/2 is a sum of two distinct powers of 2 (A018900).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 05 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017517 | OEIS-A393171

G.f. satisfies A(x) = x + A(C(x))^2, where C(x) = x + C(x)^2 is a g.f. of the Catalan numbers (A000108).

Conjecture: for n > 2, a(n) == 4 (mod 8) iff n = 2^k or n = 2^k + 1 for k >= 1, otherwise a(n) is divisible by 8.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 04 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017518 | OEIS-A393172

G.f. A(x) satisfies A(x-x^2) = x + A(x)^2.

Conjecture: for n > 1, a(n) == 2 (mod 4) iff n is a power of 2, otherwise a(n) is divisible by 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Feb 04 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017519 | OEIS-A393210

Halting time for Wolfram's 2-state 2-symbol Turing machine 3626 when started with n on the tape.

Conjecture: a(n) = Sum_{d | n} m(d)^2 * phi(d), where m(n) = Sum_{d | n} (-1)^(d+1) * moebius(d) = A069517(n) and phi(n) denotes Euler's totient function A000010(n). Cf. A296306. - _Peter Bala_, Sep 11 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sean A. Irvine_, Feb 05 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017520 | OEIS-A393217

Perfect powers m^k, with m, k > 1, such that no digit of m^k is present in k^m.

Conjecture: For n > 20, a(n) is of the form 10^k or m^10. The sequence A062518 gives the conjectural largest exponent k such that n^k does not contain all digits from 0 to 9. Then, given m > 1, with the exception of m = 10, for large exponents it is very likely that both m^k and k^m are pandigital numbers, and therefore share digits. From this observation, and considering that 10^k contains only the digits 0 and 1, if one wants m^k and k^m to have no digits in common, this is more likely to occur when m = 10. For example, the numbers 10^5, 10^7, 10^14, 10^118, 10^976 and 10^71637 are terms of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gonzalo Martínez_, Feb 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017521 | OEIS-A393289

Genus of the n-path complement graph.

It is conjectured that a(n) = ceiling((n^2 - 9*n + 14)/12) for all n >= 8 except a(10) = 3. - _Eric W. Weisstein_, Jul 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Mar 10 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017522 | OEIS-A393306

a(n) is the greatest number k that contains, in base n, every digit exactly once, and such that k/(n-1) (if n is even) or 2*k/(n-1) (if n is odd) is prime.

Conjecture: a(n) exists for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Feb 10 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017523 | OEIS-A393317

Maximum halting time for any of the 4096 Wolfram 2-state 2-symbol Turing machines across all n-bit inputs.

We conjecture that the maximum runtime for machine #378 and #1351 is 2^(n+2)-3 = A036563(n+2) for the n-bit input 2^n-1, while for machine #1447 it is 2*(n+2)^2-1 = A056220(n+2), for the same input. _M. F. Hasler_, Feb 12 2026 a(n) = 2*(n+2)^2 - 1 = A056220(n+2) for n < 5, a(n) = 2^(n+2) - 3 = A036563(n+2) for n > 4 (conjectured). - _M. F. Hasler_, Feb 12 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sean A. Irvine_, Feb 10 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017524 | OEIS-A393331

Nonnegative integers k such that the Euclidean distance between the vector of proportions of the digits 0-9 in 2^k and the uniform distribution (1/10, ..., 1/10) sets a new minimum.

Conjecture: The digit proportions in 2^k can get arbitrarily close to, but not equal to, the uniform distribution. This implies that there are infinitely many terms in this sequence and in the related sequences A393332 and A393333.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pontus von Brömssen_, Feb 12 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017525 | OEIS-A393377

Minimum Mahler measure of all irreducible and non-cyclotomic quintic polynomials with coefficients in {-1, 0, 1}.

Conjecture: Mahler measure M(d) of non-cyclotomic polynomial for every d >= 1.17628081825991750654407...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Artur Jasinski_, Feb 13 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017526 | OEIS-A393434

a(0) = 0; thereafter a(n) is the least integer (in absolute value) not yet in the sequence such that the absolute difference between a(n-1) and a(n) is a factorial number k!, k >= 1; in case of a tie, preference is given to the positive value.

It is conjectured that every integer appears in this sequence. (It appears that the proof in A379719 can be modified to apply here. The start is somewhat irregular, but beginning at n = 25, the positive terms appear in blocks of consecutive numbers, where block B(k), k >= 4, has length k*k!/2, starts at n = k!/2 + 1 and ends at n = (k+1)!/2. The structure of

the negative blocks is less clear.)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Feb 17 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017527 | OEIS-A393438

Let x = smallest even number missing from A390314 after A390314(n) has been found; a(n) = x/2.

It is conjectured that every even number will appear in A390314.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Mar 10 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017528 | OEIS-A393460

a(n) is the least prime p such that n - p is the reverse of a prime, where leading 0's are not allowed, or -1 if there is no such p.

Conjecture: a(n) = -1 only for n = 11.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, Feb 15 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017529 | OEIS-A393487

Triangle read by rows: T(n,k) (0 <= k <= n) giving binomial basis coefficients of excess-tail polynomials p_n(s) = Sum_{k=0..n} T(n,k)*binomial(s-2,k) for connected defect clusters in the row-word inversion statistic on standard Young tableaux of shape (m,m,m).

Conjecture: the diagonal entries satisfy the factorization T(n,n-j) = 2^(n-2j) * Sum_{r=0..j} A393486(j,r)*binomial(n,r), where A393486(j,j) = 3^j. This has been verified numerically for j = 0..5. T(n,n) = 2^n (conjectured; verified for n = 0..10). Conjecture: T(n,n-j) = 2^(n-2j) * Sum_{r=0..j} A393486(j,r)*binomial(n,r) (verified for j = 0..5). Conjecture: the leading coefficient of p_n(s) is 2^n / n!.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Morten Brydensholt_, Mar 16 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017530 | OEIS-A393518

Number of unlabeled rooted trees with n nodes whose level sizes are equal, excluding the root level.

Open question: Does the sequence contain any value greater than 2 more than once?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert P. P. McKone_, Feb 18 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017531 | OEIS-A393585

The maximum number of distinct products from 1 to n^2 produced by the product of 2 sets, each with n positive integer elements.

Conjectured values:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rhys Feltman_, Mar 28 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017532 | OEIS-A393615

a(n) is the largest integer m such that every integer from 1 to m can be represented as a sum of at most n squares from the set {1^2, 2^2, 3^2, ..., n^2}.

n^3 - a(n) ~ C*n^(3/2) with C approximately 5.3 (conjectured, based on empirical data up to n = 100).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimas Saputra_, Mar 29 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017533 | OEIS-A393669

Size of the free implicative semilattice on n generators.

It is conjectured that log(a(n+1)) ~ a(n) as n tends to infinity (see the introduction from de Bruijn).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lorenzo Sauras Altuzarra_, Feb 24 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017534 | OEIS-A393685

Records in A393684.

Conjecture: these records occur at indices (3^n - 1)/2 = A003462(n); that is, they correspond to all 1's ternary inputs on the tape.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sean A. Irvine_, Feb 24 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017535 | OEIS-A393700

Numbers multiplied by 2 conjecturally not occurring in A393562.

Numbers multiplied by 2 conjecturally not occurring in A393562.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Feb 25 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017536 | OEIS-A393703

a(n) is the smallest index k such that the subsequence b(k), b(k+1),..., b(k+n-1) is a palindrome of length n, where b(k) = A206400(k).

The aim of this sequence is to study the local symmetries of the differences between prime numbers of the form m^2+1. To do this, we identify palindromes in subsequences of A206400. This research falls within the framework of the fourth problem posed by the mathematician Edmund Landau at the 1912 International Congress of Mathematicians (Landau, 1912). This problem, still unsolved to this day, conjectures the existence of infinitely many prime numbers of the form k^2+1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Feb 25 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017537 | OEIS-A393710

a(n) = number of triples (x, y, z) such that x^2 + y*z = n, where x, y, z are positive integers satisfying x < y < z.

Conjecture: each nonnegative positive integer occurs only finitely many times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Mar 31 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017538 | OEIS-A393794

Fixed points of the 1089-trick map T(n) = |n - reverse(n)| + reverse(|n - reverse(n)|).

a(n) is the unique fixed point of the 1089-trick map T among all (n+4)-digit numbers (verified exhaustively for n = 1..4; conjectured for all n >= 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Remco Havenaar_, Feb 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017539 | OEIS-A393798

a(n) is the least prime p such that a(n)-a(n-1) is a 3-smooth number, starting with a(1) = 2.

Conjecture: This sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Feb 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017540 | OEIS-A393831

Number of derangements p of n points on a circle such that if x is next to y, then p(x) is not next to p(y).

Conjecture: series for log(a(n)/n!) begins with -3 - 5/n - 11/n^2 - ...

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Liam Courtright_, Mar 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017541 | OEIS-A393833

Least integer k having exactly 2n+2 divisors such that the sequence of its 2n+1 proper divisors modulo n is a palindrome.

Conjecture: for every integer n>=1, there always exists at least one integer k such that its proper divisors are palindromic modulo n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Mar 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017542 | OEIS-A393840

Number of prime factors of the form k*2^A228845(n) + 1 (with k odd) that divide the Fermat number 2^(2^n) + 1.

Conjecture 1: the sequence is not monotonic. Conjecture 2: if A046052(n) = 2*m for some integer m > 1, then a(n) + 1 < A046052(n). According to this, if a Fermat number has an even number of prime factors and is not semiprime, at least two of them are in A351865.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadiusz Wesolowski_, Apr 02 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017543 | OEIS-A393850

G.f. A(x) satisfies: [x^n] A(x)^(n+1) = n*(n+1) * [x^(n-1)] A(x)^(n+1) for n >= 1 with A'(0) = 1.

Conjecture: a(n) is odd iff n is a power of 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017544 | OEIS-A393856

G.f. A(x) satisfies: A(x - x*A(4*x)/4) = x.

Conjecture: a(n) is odd iff n is a power of 2. Conjecture: a(n) == 1 (mod 5) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017545 | OEIS-A393857

G.f. A(x) satisfies A(x - x*A(5*x)/5) = x.

Conjecture: a(n) == 1 (mod 6) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017546 | OEIS-A393858

G.f. A(x) satisfies: A(x - x*A(6*x)/6) = x.

Conjecture: a(n) is odd iff n is a power of 2. Conjecture: a(n) == 1 (mod 7) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017547 | OEIS-A393859

G.f. A(x) satisfies A(x - x*A(7*x)/7) = x.

Conjecture: a(n) == 1 (mod 8) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017548 | OEIS-A393867

Logarithmic derivative of A393866.

Conjecture: a(n) is divisible by prime(n) for n > 1. Conjecture: all terms are odd.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Mar 29 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017549 | OEIS-A393870

G.f. A(x) satisfies [x^(n+2)] A^(2^n)(x) = n*2^n * [x^(n+1)] A^(2^n)(x) for n >= 1, where A^k(x) denotes the k-th iteration of A(x).

Note: it is conjectured that this sequence will continue to consist of integers (true for at least the initial 151 terms); if there are fractional coefficients eventually, then this sequence would list the numerators of those fractions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 08 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017550 | OEIS-A393879

G.f. satisfies Sum_{n>=0} (-1)^n * A(x)^(n^2) = 1 - x - x^2 - x^3.

Conjecture: a(n)^(1/n) tends to (2 + (53 - 3*sqrt(201))^(1/3) + (53 + 3*sqrt(201))^(1/3))/3 = 2.91963956539418145114139... - _Vaclav Kotesovec_, Apr 17 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017551 | OEIS-A393880

G.f. satisfies Sum_{n>=0} (-1)^n * A(x)^(n^2) = 1 - x - x^2.

Conjecture: a(n)^(1/n) tends to 1+sqrt(3). - _Vaclav Kotesovec_, Apr 17 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Apr 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017552 | OEIS-A393931

a(n) is the least positive integer k such that the elliptic curve y^2 = x^3 - k*x + 1 has exactly n integer solutions with y >= 0.

Conjecture: For all n > 3, a(n) is not prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhining Yang_, Mar 03 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017553 | OEIS-A393940

Numbers m such that tau(m) is a perfect number (A000396).

Conjecture: number m = 18 is only number such that tau(m) and phi(m) are both perfect numbers. Conjecture is false: with m = 2^61 * 17 * 65537 * ((2^2281-1)*2^2200+1), tau(m) and phi(m) are both perfect numbers. - _Michel Marcus_, Mar 12 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Mar 03 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017554 | OEIS-A393942

Numbers m such that phi(m) is a perfect number (A000396).

Conjecture: number m = 18 is the only number such that tau(m) and phi(m) are both perfect numbers. Conjecture is false: with m = 2^61 * 17 * 65537 * ((2^2281-1)*2^2200+1), tau(m) and phi(m) are both perfect numbers. - _Michel Marcus_, Mar 12 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaroslav Krizek_, Mar 03 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017555 | OEIS-A393997

Number of steps to reach the end of the cycle starting with n/(2n+1), or -1 if the end of the cycle is never reached, in the following process: If the numerator is even, divide the fraction by 2, otherwise multiply the fraction by 3 and add 1.

Conjecture: a finite cycle is always reached (checked for n <= 10^5).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, May 18 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017556 | OEIS-A394054

Irregular triangle read by rows: T(n,k) is the number of positive integers <= n with exactly k 2-dense sublists of divisors.

Conjecture: T(n,k) is the number of positive integers m <= n whose symmetric representation of sigma(m) has k parts.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Mar 12 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017557 | OEIS-A394057

Conjectured least positive integer k such that n^k has the largest average of digits and k>1.

Conjectured least positive integer k such that n^k has the largest average of digits and k>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Mar 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017558 | OEIS-A394058

Conjectured least integer k>1 such that n^k has the smallest average of digits, or -1 if no such k exists.

Conjectured least integer k>1 such that n^k has the smallest average of digits, or -1 if no such k exists.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmitry Kamenetsky_, Mar 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017559 | OEIS-A394059

a(n) is the smallest index k such that the subsequence b(k), b(k+1),..., b(k+n-1) is a palindrome of length n, where b(k) = prime(k) mod 6 = A039704(k).

Conjecture: There exist infinitely many palindromes of every length in the sequence of residues of primes modulo 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michel Lagneau_, Mar 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017560 | OEIS-A394082

Diagonal of the 4-dimensional geode array: a(n) = G(n,n,n,n).

Empirical: Numerical fitting of 120 terms suggests a(n) ~ K * alpha^n * n^beta where beta = -3.000 to at least 3 decimal places. The exponential base alpha has not fully converged at n=119 (successive ratios a(n+1)/a(n) ~ 1083000 and increasing); it is conjectured to be 14^14/10^10 = 1111200.6826... based on saddle-point heuristics.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Steven Kotlarz_, Mar 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017561 | OEIS-A394083

Genus of the n-cycle complement graph.

It is conjectured that a(n) = ceiling((n^2 - 9*n + 12)/12) for all n >= 8 except a(10) = 3. - _Eric W. Weisstein_, Jul 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Mar 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017562 | OEIS-A394090

Decimal expansion of the parameter c that minimizes the integral of squared distance between sin((Pi/2)x)^(1/c) and the quarter-circle sqrt(2x-x^2) on [0,1].

It is conjectured that this constant is transcendental.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Minyan Liu_, Mar 10 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017563 | OEIS-A394227

Positive integers whose decimal digits are 2, 3, or 6, and that are divisible by the product of their digits.

Conjecture: The sequence is infinite. Idea: If one assumes the heuristic that a "decimal with digits 2, 3, 6 is divisible by P, the product of its digits, with probability 1/P", then the expected number of terms with M digits is exactly 1 using 1/2 + 1/3 + 1/6 = 1 and the trinomial expansion formula (see Alkauskas link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giedrius Alkauskas_, Mar 13 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017564 | OEIS-A394258

Largest prime less than the n-th prime-complete number.

A double (two prime-complete numbers in one prime gap) occurs only for the shared previous-primes 31, 89, 479, and 4603; no further double exists below 10^12, and a heuristic based on the density of prime-complete numbers suggests this list of doubles is complete and that no triple occurs. That is, the counting function for prime-complete numbers, A377028(n), grows sub-polynomially (empirically exp(O(sqrt(log n)))), so prime-complete numbers thin out faster than any polynomial while prime gaps grow only logarithmically. The expected number of further doubles beyond x is therefore a convergent sum, supporting the conjecture that the complete list is 31, 89, 479, and 4603.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ken Clements_, May 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017565 | OEIS-A394263

Positive integers k for which the Euclidean distance between the vector of proportions of the first k decimal digits of Pi and the uniform distribution (1/10, ..., 1/10) sets a new minimum.

Conjecture: The digit proportions of an initial segment of the decimal expansion of Pi can get arbitrarily close to, but not equal to, the uniform distribution. This implies that there are infinitely many terms in this sequence and in the related sequences A394264 and A394265.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pontus von Brömssen_, Mar 14 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017566 | OEIS-A394382

Primes p such that the reversal of p^2 is the square of a prime not equal to p nor the reversal of p.

Conjecture 1: The first and last decimal digit of a(n) is 1. Conjecture 2: The nonzero decimal digits of a(n) are either 1, 2 or 9. Using Conjecture 2, larger terms such as 1001002109111, 1001011091011, 1001012009111, 1001100901111, 1001100911101, 1001199911111, 1002011091121, 1010109020111, 1010990110201, 1011010091111, ... are found.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Apr 07 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017567 | OEIS-A394399

Numbers k in A014574 (averages of twin prime pairs) such that gcd(k, sigma(k)) = 3.

Conjecture: all terms are of the form 2*m^2 (twice a perfect square; A001105).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aied Sulaiman_, Apr 24 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017568 | OEIS-A394407

Positive integers k such that there is no m < k with phi(m)/tau(m) = phi(k)/tau(k).

Conjecture: All primes p != 3 are in the sequence; equivalently, for every prime p != 3 there is no m < p with phi(m)/tau(m) = phi(p)/tau(p) = (p - 1)/2; equivalently, the sequence is a supersequence of A045344. Verified for all primes p < 10^6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Mar 28 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017569 | OEIS-A394409

Pascal-like triangle read by rows: top entry is 2; thereafter fill in entries from top to bottom, alternating between left to right and right to left, with the smallest number that is not yet in the sequence and has a common factor with any earlier entry that is adjacent to it in the triangle.

It is conjectured that every number greater than 1 appears.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017570 | OEIS-A394410

Quadratic recurrence sequence with a(1)=1, a(2)=3, and a(n) = (a(n-1)^2 + a(n-2)^2)/2 for n > 2.

Thomas Ordowski conjectured in A056826 that if a(n) is prime, then (a(n)^a(n) + 1)/(a(n) + 1) is also prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rayhan Ahmed_, Mar 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017571 | OEIS-A394413

a(1) = 1, a(2) = 2; for n > 2, a(n) is an unused positive number that shares a factor with a(n-1) while being as close as possible to a(n-2). If two such numbers exist then the smaller is chosen.

Overall the primes are rare, only 171 terms being prime in the first 10^8 terms, see A394414, and they do not occur in their natural order. In this range the smallest missing prime is 17, although given the pattern of the large drops in value it is conjectured eventually all numbers will appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Mar 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017572 | OEIS-A394414

The prime terms in A394413, in order of appearance.

It is conjectured that the sequence eventually contains all primes.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Apr 12 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017573 | OEIS-A394416

a(1) = 1, a(2) = 2; for n > 2, a(n) is the greatest unused number < a(n-1) such that it shares a factor with a(n-1) while rad(a(n)*a(n-1)) has not previously occurred as the rad value of the product of adjacent terms, where rad(k) = A007947(k). If no such number exists then a(n) is the smallest number > a(n-1) satisfying the same criteria.

The fixed points in the first 10^6 terms are 1, 2, 240, 723, 2123, 3263, 11885, 27305, 90145, 122419, 918023; more likely exist. The sequence is conjectured to be a permutation of the positive numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Mar 28 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017574 | OEIS-A394418

a(1) = 1, a(2) = 2; for n > 2, a(n) is an unused positive number that shares a factor with a(n-1) while being as close as possible to a(n-2). If two such numbers exist then the larger is chosen.

Overall the primes are rare, only 73 terms being prime in the first 10^8 terms, see A394419, and they do not occur in their natural order. In this range the smallest missing prime is 19, although, given the pattern of the lower line approaching the x-axis, it is conjectured eventually all numbers will appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Mar 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017575 | OEIS-A394431

The smallest base b < n where the sum of the digits for the number n in the base b is the largest, with 1 < b < n and a(1) = a(2) = 1.

Conjecture: a(n) = ceiling((n+1)/2) for n>8. - _Sean A. Irvine_, Mar 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jean-Marc Rebert_, Mar 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017576 | OEIS-A394432

a(n) is the least r such that ord(n,r) >= (log_2(n))^2, where ord(n,r) is the multiplicative order of n modulo r.

Conjecture: a(n) is always an odd prime power. It seems that most terms are of the form p^k for odd prime p and k <= 3. Other than a(110) = a(119) = 81 and a(n) = 625 for n = 3961388, 4132533, 4466762, 4610283, 4629398, 4961877, 4966163, 5006623, 5054222, 5063527, 5276203, 5314622, 5338663, is there any other term that is a prime power with exponent higher than 3? Conjecture: for n > 2 being a power of 2, there is no difference between ">= (log_2(n))^2" and "> (log_2(n))^2"; in other words, we actually have ord(2^e,a(2^e)) = ord(2,a(2^e))/gcd(ord(2,a(2^e)), e) > e^2 for e > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Mar 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017577 | OEIS-A394435

Primes p such that A001414(p-1)^2 + A001414(p+1)^2 is a perfect square, where A001414(n) is the sum of prime factors of n with multiplicity (sopfr).

Conjecture: this sequence is infinite (empirical observation; no proof is known).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rudra Jadhav_, May 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017578 | OEIS-A394450

Primes that survive the lucky number sieve when applied to the sequence of primes.

The survival fraction among primes up to N decreases slowly: 73/168 = 43.5% for N=1000, 452/1229 = 36.8% for N=10000, 3129/9592 = 32.6% for N=100000. The asymptotic density among primes is an open question.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nova Spivack_, Mar 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017579 | OEIS-A394456

Odd integers k such that k+2 is prime and the sequence of primes congruent to 2 modulo k is strictly stable under iterated absolute differences (all leading entries equal k).

All terms are currently conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Li Dong_, May 04 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017580 | OEIS-A394500

Ways to represent n = k + p where p is prime and k > 0 is not squarefree.

Lee & O'Clarey prove that a(n) > 0 for large enough n and conjecture that a(n) > 0 for n > 24.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, May 24 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017581 | OEIS-A394508

Write the primes in a column. Add new columns to the right, with each entry being the prime whose index is the entry to its left. The sequence a(n) is given by the antidiagonal sums.

Conjecture: a(n) is composite for n>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philip Jameson_, Mar 22 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017582 | OEIS-A394511

Minimum number of cells in a connected enclosing shell of an n-cell connected hole on the truncated-square (4.8.8) Archimedean tiling.

Conjectured (UNVERIFIED beyond n = 23): the minimum shell equals the corona of the optimal hole, |S| = |C(H)|; and a(n) = 2*ceiling((n + 17)/5) for n >= 6, equivalently a(n) = a(n - 5) + 2 for n >= 11. - _Peter Exley_, Jun 05 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Exley_, May 05 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017583 | OEIS-A394514

a(1) = 1, a(2) = 2; for n > 2, a(n) is the greatest unused number less than a(n-2) that shares a factor with a(n-1), else, if no such number exists, a(n) is the smallest unused number greater than a(n-2) that shares a factor with a(n-1).

The fixed points in the first 10^6 terms are 1, 2, 8, 25, 60, 74, 99, 714, 7424, and 149157, although more likely exist. The sequence is conjectured to be a permutation of the positive numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Mar 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017584 | OEIS-A394532

Numbers k such that every abelian group of order k appears as a group of units of the ring Z/mZ for at least one m.

For general totient numbers t, the map f_t is injective for t = 2^s, s >= 33, assuming five Fermat primes. See the Englezou link for a proof. This raises the interesting question whether f_r is injective for r <> 2^n. A negative claim would constitute a generalization of Carmichael's totient conjecture. The above generalization of Carmichael's conjecture is false. f_k is injective for many primes p such that k = p * 2^r: for example, it is injective for k = 11 * 2^203, k = 17 * 2^111, and k = 19 * 2^89. For a method of determining that f_k is noninjective for k = p * 2^r over all r >= 1, see the comment in A398282. - _Miles Englezou_, Jul 25 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Miles Englezou_, Mar 24 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017585 | OEIS-A394543

Numbers k that reach a power of 2 in zero or more iterations of k -> 2k - (sum of proper factors of 2k).

Conjecture: superset of A362684.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ruud H.G. van Tol_, Mar 24 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017586 | OEIS-A394555

Powerful k such that rad(Sum({m <= k : rad(m) = rad(k)})) = rad(k), where rad = A007947.

Conjecture: this sequence intersects A383394.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Mar 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017587 | OEIS-A394576

a(1) = 1; a(n+1) is the smallest positive integer greater than a(n) that avoids violating the condition a(k) + a(k+1) + a(k+2) is not a term.

Conjecture 1: a(n) <= floor((4*n-1)/3). Conjecture 2: The sequence alternates between "stable blocks" where a(n) strictly equals floor((4*n-1)/3) and "gaps" where the pattern temporarily breaks to avoid forbidden sums. Conjecture 3: Exact binary structure of the gaps. Let d(n) = floor((4*n-1)/3) - a(n). From Conjecture 1, d(n) = 0 within the stable blocks and fluctuates strictly in {0, 1} within the gaps.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hoang Xuan Thanh_, May 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017588 | OEIS-A394578

a(n) = 2^(2*n-1) * p(n, n+1/2), where p(n,x) is the Lagrange interpolating polynomial through (1,0) and (k,1) for k=2..n+1.

Conjecture: a(n+1)/a(n) -> 1/4. Conjecture D-finite with recurrence n*a(n) +2*(-2*n+1)*a(n-1) +16*(-n+3)*a(n-2) +32*(2*n-7)*a(n-3)=0. - _R. J. Mathar_, Jun 30 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017589 | OEIS-A394594

The largest positive integer for which there exist n arithmetic progressions whose union covers all smaller positive integers, but miss a(n) and a(n)+1.

By an explicit construction, a(2k-1)>=2^(k+1)-1, a(2k)>=3*2^k-1. From the result of Crittenden and Vanden Eynden it follows that for k>=2, a(k)<2^k. Numerical values support conjectural bounds c*2^(k/2)<a(k)<C*2^(k/2).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giedrius Alkauskas_, Mar 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017590 | OEIS-A394611

a(1) = 0. For n > 1, a(n) is the number of ways to write a(n-1) as the sum of two earlier terms with distinct positions in the sequence.

Conjecture: Many numbers, beginning with 5 and 9, never occur in this sequence (see A397471).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joshua B. Weinstein_, Jun 18 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017591 | OEIS-A394621

Numbers k of the form k = p1 + p2 = x^2 + y^2 + p1^2, where p1 and p2 are primes, x, y are integers.

Conjecture: odd a(n)-2 belong to A079739. Conjecture: odd a(n)-4 belong to A001481. Conjecture: even a(n) belong to A145979.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bill McEachen_, Mar 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017592 | OEIS-A394634

a(n) is the least k >= 0 such that the maximum length of a divisibility chain among the distinct entries in row k of Pascal's triangle is n, or -1 if no such k exists.

Conjecture: a(n) >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Oct 12 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017593 | OEIS-A394659

Number of independent vertex sets of size 7 in the n-triangular grid graph.

For n > 3, a(n) = (n - 4)*(n^13 + 25*n^12 + 9*n^11 - 3555*n^10 - 7745*n^9 + 260563*n^8 + 291075*n^7 - 11898033*n^6 + 10339924*n^5 + 303368632*n^4 - 909281184*n^3 - 2609966352*n^2 + 16737321600*n - 23213352960)/645120 (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric W. Weisstein_, Mar 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017594 | OEIS-A394660

a(n) is the largest prime number such that prime(n)-a(n) is a 3-smooth number where prime(n) denotes the n-th prime number, or zero if no such number exists.

Conjecture: No term is 0 (verified to n=10^7).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Lei Zhou_, Mar 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017595 | OEIS-A394666

a(n) = n! mod (2*n - 1).

Conjecturally, a(n) = 0 iff n > 5 and 2*n - 1 is not a prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sabuhi A. Amirov_, Apr 29 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017596 | OEIS-A394676

a(1) = 1, a(n+1) as the smallest positive integer greater than a(n) that avoids violating the condition a(k) + a(k+1) + a(k+2) + a(k+3) is not term.

Conjecture 1: a(n) <= floor((5*n-1)/4). Conjecture 2: The sequence L(k)=16^(k-1) + 3 describes the number of consecutive terms that match the formula a(n) = floor((5*n-1)/4): (Start)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hoang Xuan Thanh_, May 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017597 | OEIS-A394678

a(n) = (n! - 1) mod n^2.

Conjecture: Excluding the case a(4) = 7, when a(n) is a prime number, n is a prime such that n^2 - n - 1 is a prime (A091567). Verified for n <= 5*10^4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _C. Owen Prestwood_, Apr 29 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017598 | OEIS-A394682

Number of order-convex subsets of the cubic grid poset [n]^3 with componentwise partial order.

log a(n) = Theta(n^2) (dimension law for d=3). The growth constant c_3 = lim log(a(n))/n^2 is conjectured to equal 16*log(2)/7 = 1.5842... based on c_2/c_3 = 7/4 to 4 significant figures (where c_2 = log(16) is the growth constant for d=2, A393665).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas DiFiore_, Mar 28 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017599 | OEIS-A394689

Lexicographically earliest infinite sequence of distinct positive numbers such that, for n > 2, a(n) shares a factor with a(n-1) while the set of distinct prime factors shared between a(n) and a(n-1) differs from that shared between a(n-1) and a(n-2).

In the first 10^5 terms there are no fixed points other than the first two terms, and it is likely no more exist. The sequence is conjectured to contain all numbers other than the prime powers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Mar 29 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017600 | OEIS-A394694

Number of occurrences of n in A081493.

Conjecturally: a(n) = A000005(n-1)/2 + e(n), where e(n) = 1/2 if n-1 is a perfect square, 1 if n-1 is pronic, and 0 otherwise. Also conjecturally: a(n) + A033831(n-1) = A000005(n-1) for n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gokul Padmanabhan_, Mar 29 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017601 | OEIS-A394697

a(n) = max_k [x^k] I(G, x), where G is the n X n toroidal king graph and I(G,x) is its independence polynomial.

Conjecture: a(n)^(1/n^2) tends to A247413. - _Vaclav Kotesovec_, May 06 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Todd Hildebrant_, Apr 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017602 | OEIS-A394752

Indices of zeros (P-positions) in the Sprague-Grundy sequence for the octal game .404.

Conjecture: a(7) = 286 is the largest P-position; equivalently, every n >= 287 is an N-position (first player wins).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sijie Wu_, May 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017603 | OEIS-A394757

Numbers k in A014574 (averages of twin prime pairs) such that gcd(k, sigma(k)) is prime.

Conjecture: all terms are divisible by 18. Conjecture: gcd(k, sigma(k)) is in {2,3}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aied Sulaiman_, Mar 31 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017604 | OEIS-A394761

a(n) = the length of the n-th run of 1's in A392200.

We conjecture that no more than eight 1's, four 2's, and three 3's can appear in a row.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sam Chapman_, Mar 31 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017605 | OEIS-A394791

Smallest number k such that the symmetric representation of sigma(k) has 2*n parts and two parts meet at the center of the Dyck paths.

Conjecture: a(n) is the smallest number k with 2*n 2-dense sublists of divisors of k and 2*A033676(k) + 1 = A033677(k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Apr 04 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017606 | OEIS-A394815

Numbers k in A014574 (averages of twin prime pairs) such that gcd(k, sigma(k)) = 2.

Conjecture: every term is divisible by 36. Conjecturally, A394757 is the disjoint union of this sequence and A394399.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aied Sulaiman_, May 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017607 | OEIS-A394819

Number of distinct permutations generated by Faro and concatenation operations on n equal blocks (saturated case).

a(n) = max{|P(n*k, n)|: k >= 1}. If the saturation conjecture is true: a(1) = 1, a(2) = 4, for n>=3: a(n) = |P(n*(n-1), n)|.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Balbino S. Sobreviela Jiménez_, Apr 02 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017608 | OEIS-A394885

Terms of A099308 for which A051903 obtains value n for the first time.

I conjecture that the sequence is strictly growing, and is well defined for all n. At least terms a(9) .. a(14) exist, see example in A394887.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Apr 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017609 | OEIS-A394906

Representation of n in a positional system with the k-th prime as the k-th place value, choosing the representation that maximizes the decimal value.

Conjecture: For every d > 0 there exists n such that a(n) has exactly d nonzero digits. Conjecture: a(n) is defined for all n >= 2. Conjecture: The two least significant (rightmost) digits are in {0,1,2}; all other digits are in {0,1}. Proof of the first conjecture: If a(n) has m nonzero digits and p > n is a prime such that nextprime(p) > p + n, then a(p + n) has m+1 nonzero digits: a leading digit 1, followed by primepi(p) - length(a(n)) - 1 zeros, followed by the digits of a(n). Since prime gaps are known to become arbitrarily large, such a prime p always exists. - _M. F. Hasler_, Apr 08 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Dewan_, Apr 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017610 | OEIS-A394957

Numbers k for which the pairwise sumset of the distinct binomial coefficients in row k of Pascal's triangle is not maximal.

Conjecture: The sequence is {6, 7, 19, 52} union {d^2 - 4, d^2 - 3, d^2 - 2 : d >= 4}, sorted increasingly.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Jul 08 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017611 | OEIS-A394981

Given a set of n rectangular tiles, {1 X 1, 1 X 2, 1 X 3, 1 X 4, ..., 1 X n}, then a(n) is the number of tilings up to symmetry of the rectangle having smallest possible perimeter with area n*(n+1)/2 using all n tiles.

Conjecture: If n is odd, then a(n+1) = 4*a(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary Yane_, Apr 08 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017612 | OEIS-A395008

Numbers k representable as a sum of three squares at which the minimal three-square spread s3(k) reaches a new record value, where s3(k) is the least value of x-z over all representations k = x^2 + y^2 + z^2 with x >= y >= z >= 0.

Conjecture: a(n) = 4*a(n-5) for n >= 13, with initial terms a(1)..a(12) = 1, 4, 10, 16, 37, 58, 64, 130, 148, 232, 256, 445.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nenad Micic_, Jun 10 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017613 | OEIS-A395056

a(n) is the minimum, over all k >= 1, of the number of terms omitted from the first k terms of A008578 (1 followed by the primes) so that the remaining terms sum to n.

Conjectured: a(n) <= 3 for all n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ron Gordon_, Jul 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017614 | OEIS-A395088

Fixed point of the morphism 0->011, 1->0.

Conjecture: Let c(n) be the list with the length of 0-runs, then a(n) = (3-c(n+2))/2, i.e., c(1) = 1, c(n) = 3 - 2*a(n-2) for n > 1. If this conjecture is true, this is a self-generating sequence, starting with 011 whereafter the n-th 0-run has length 3 - 2*a(n-2) and the 1-runs all have length 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _A.H.M. Smeets_, Apr 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017615 | OEIS-A395095

Indices of records in A295741(n) = min {d > 0 | primorial(n)/d + d is a prime}.

It is still only conjectured that A295741(n) > 0 for all n, therefore it seems interesting to study those n which yield exceptionally high values.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Apr 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017616 | OEIS-A395102

a(1) = 3; a(n) is the smallest odd prime not occurring earlier so that a(n) mod a(n-1) is prime.

There are three ways that the next term may be generated. If there is at least an odd prime < a(n) not on the list, a(n+1) is the smallest among them. Otherwise, if a(n)+2 is prime and not on the list, a(n+1) = a(n)+2. Otherwise, a(n+1) takes the form 2*k*a(n)+p with prime p. It is conjectured that k is always 1, which has been confirmed until n = 10000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hoang Nguyen_, May 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017617 | OEIS-A395103

Numbers with exactly k prime factors, counting repeats, with k>1, that are formed by concatenating k primes.

Conjecture: The sequence has infinitely many terms. The conjecture is true if A117360 is infinite. - _Yehune Jeong_, May 24 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philip Jameson_, May 10 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017618 | OEIS-A395110

Number of functions on the grid [0,n+1]^2 which are zero on the boundary and satisfy f(i,j) = ceiling((f(i-1,j)+f(i,j-1)+f(i+1,j)+f(i,j+1))/4) for 1<=i,j<=n.

If Laplacian is bounded by 2, and f is 0 on the boundary, one can construct 2^(floor(n/2)) different examples of such functions. All of them have a dihedral symmetry. Conjecturally, this is the exact number of such functions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Giedrius Alkauskas_, May 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017619 | OEIS-A395126

a(n) = fourth coordinate d of the n-th P-position (a,b,c,d) of 4 X n Chomp in lexicographic order, where a>=b>=c>=d>=0 gives row lengths.

A position (a,b,c,d) with a>=b>=c>=d>=0 is a P-position of 4 X n Chomp if the player to move loses under optimal play. This sequence lists the d-values of all such positions sorted lexicographically by (a,b,c). The first 10 positions in order are: (1,0,0,0), (2,1,0,0), (2,2,1,0), (2,2,2,1), (3,1,1,0), (3,2,0,0), (3,3,1,1), (4,1,1,1), (4,2,2,0), (4,3,0,0). Computed for all n<=500, yielding 4316097 P-positions with zero violations of the Unique Extension conjecture: for any triple (a,b,c), at most one value of d completes a P-position. This is an empirical observation, not a proof.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arnav Garg_, Apr 12 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017620 | OEIS-A395143

Numbers with no representation as the quotient of two fibbinary numbers (A003714).

Meleshko has a conjectured characterization of the members of this sequence, but as of 2026 it has not been proved.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, May 13 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017621 | OEIS-A395149

Triangle of coefficients of characteristic polynomial of negative Pascal matrix with (i+1,j+1)-th entry -C(i+j+2,j).

Conjecture: The alternating row sum of index 2n is A266091(n)^4 / 3^(2*n) / (2*n+1). For example, 1 - 76 + 275 - 76 + 1 = 15^4 / 3^4 / 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _F. Chapoton_, Apr 14 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017622 | OEIS-A395156

Square array read by antidiagonals upwards: T(n,k) is the n-th number j with the property that the symmetric representation of sigma(j) has 2*k parts and two parts meet at the center of the Dyck paths, n >= 1, k >= 1, j >= 1.

Conjecture: T(n,k) is the n-th number j with exactly 2*k 2-dense sublists of divisors and 2*A033676(j) = A033677(j) - 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, May 16 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017623 | OEIS-A395160

Irregular triangle read by rows: T(n,k) is the average of the odd numbers of the k-th 2-dense sublist of divisors of n and of its conjugate, with n >= 1, k >= 1.

Conjecture 1: row lengths give A237271. Conjecture 2: T(n,k) = (A384226(n,k) + A384226(n,m+1-k)) / 2, where m = A237271(n) is the row length. Conjecture 3: T(n,k) = A237270(n,k) / A038712(n) = A237270(n,k) * A000593(n) / A000203(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar E. Pol_, Apr 15 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017624 | OEIS-A395235

Powerful numbers of the form k^2 - 2.

Note that k^2 - 4k + 4, k^2 - 2k + 1, and k^2 - 2 are powerful numbers in arithmetic progression. van Doorn proves that this sequence is infinite (by constructing an explicit infinite subsequence) and conjectures that, infinitely often, the three terms are consecutive as powerful numbers, which would disprove a conjecture of Erdős.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles R Greathouse IV_, May 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017625 | OEIS-A395261

Self-convolution of A000293.

Conjecturally, log(a(n)) ~ 2^(1/4) * C_3 * n^(3/4) as n -> infinity, where C_3 = 1.822... is the asymptotic constant for solid partitions (i.e., log(A000293(n)) ~ C_3 * n^(3/4)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Thomas DiFiore_, Apr 17 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017626 | OEIS-A395279

Jacobsthal's function H(n); the maximum, over all t having exactly n distinct prime divisors, of A048669(t).

It was conjectured (sometimes wrongly asserted) that H(n) = h(n) for all n >= 1. However, in 2012 Hajdu and Saradha showed that H(24) = 236 != 234 = h(24) is the first counterexample. The particular example achieving 236 given by Hajdu and Saradha is t = A002110(21)*73*89*101.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jeffrey Shallit_, Apr 18 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017627 | OEIS-A395284

Smallest semiprime that produces a chain of exactly n semiprimes when repeatedly replacing m=p*q (with primes p<=q) by the decimal concatenation of p followed by q, stopping when the result is not semiprime.

Conjecture: a(n) exists for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Philip Jameson_, Apr 18 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017628 | OEIS-A395320

Decimal expansion of the conjecturally maximum possible volume of a polyhedron with 14 vertices inscribed in the unit sphere.

Decimal expansion of the conjecturally maximum possible volume of a polyhedron with 14 vertices inscribed in the unit sphere.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Apr 19 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017629 | OEIS-A395321

Decimal expansion of the conjecturally maximum possible volume of a polyhedron with 16 vertices inscribed in the unit sphere.

Decimal expansion of the conjecturally maximum possible volume of a polyhedron with 16 vertices inscribed in the unit sphere.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Apr 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017630 | OEIS-A395322

Decimal expansion of the length of the longest edge of the polyhedron with 14 vertices with conjecturally maximum possible volume inscribed in the unit sphere.

Decimal expansion of the length of the longest edge of the polyhedron with 14 vertices with conjecturally maximum possible volume inscribed in the unit sphere.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Apr 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017631 | OEIS-A395323

Decimal expansion of the conjecturally maximum possible volume of a polyhedron with 32 vertices inscribed in the unit sphere.

Decimal expansion of the conjecturally maximum possible volume of a polyhedron with 32 vertices inscribed in the unit sphere.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Apr 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017632 | OEIS-A395324

Decimal expansion of the length of the shorter edges of a polyhedron with 32 vertices with conjecturally maximum possible volume inscribed in the unit sphere.

Decimal expansion of the length of the shorter edges of a polyhedron with 32 vertices with conjecturally maximum possible volume inscribed in the unit sphere.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Apr 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017633 | OEIS-A395338

Primes whose square is a concatenation of exactly 2 distinct nonzero squares.

Conjecture: sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pang Ern_, Apr 19 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017634 | OEIS-A395422

Minimum number of triangular cells in a connected polyiamond container that contains every fixed n-iamond as a translated subset.

a(n) is the minimum number of cells in a connected polyiamond containing every fixed n-iamond as a translated subset. The first differences are non-monotone, no low-degree polynomial fits, and a(n) exceeds the free-piece analog A392363(n). The reported a(n) is the minimum container size over the n X n triangular search window (1 X 2 for n=1); it is the conjectured global minimum, proved minimal within that window. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Exley_, Apr 22 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017635 | OEIS-A395424

Busy Beaver problem for FRACTRAN: a(n) is the maximal number of steps that a FRACTRAN program of size n can make on the integer 2 before eventually halting.

Conjecture: These 3 programs of size 22 are non-halting: [1/15, 27/77, 49/3, 10/49, 33/2], [1/15, 49/3, 27/77, 10/49, 33/2], and [27/35, 1/33, 25/3, 22/25, 21/2]. A proof would require solving a Collatz-like problem. For more details, see "Fenrir" in the bbchallenge wiki.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jason Yuen_, Apr 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017636 | OEIS-A395429

Least k such that {1^n, 2^n, ..., k^n} contains numbers with every leading digit from 1 through 9.

Conjecture 1: There exists some integer n > 1 satisfying a(n) = 9. Equivalently, for such n, the leading digits of 1^n, 2^n, ..., 9^n are exactly the nine distinct values 1 through 9. Conjecture 2: The sequence is uniformly bounded; there exists a fixed constant C such that a(n) <= C for all positive integers n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Maju Wu_, Jun 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017637 | OEIS-A395454

Triangle read by rows: T(n, k) = Sum_{j=0..n} C(-j-1, -n-1)*E2(j, n-k), E2 the Eulerian numbers A201637.

Conjecture: Let P(n) be the generating polynomial (in t) of row n. Then P(n) = (n - 2*t + n*t - 2)*P(n-1) + (t - t^2)*P'(n-1) + t*(2*n-4)*P(n-2). - _Per W. Alexandersson_, Jun 27 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Apr 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017638 | OEIS-A395473

Number of Q(chi) for chi running through the Dirichlet characters modulo n, where Q(chi) is the field generated by the values of chi.

Conjecturally each composite number k occurs infinitely many times. In particular, this would be true if there are infinitely many primes of the form 2^a*3^(b-1)*p1^(e1-1)*...*pr^(er-1), with a*b*e1*...*er = k, b > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, May 28 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017639 | OEIS-A395477

a(n) is the smallest semiprime that can be partitioned in exactly n unique ways into two coprime semiprimes.

a(542) is conjectured to not exist; if it does, its lower bound is 5600000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hoang Nguyen_, May 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017640 | OEIS-A395495

Minimum number of cells in a connected polyomino on the square grid that contains every fixed n-omino as a translated subregion.

Conjecture: a(n) = (n^2 - n + 6)/2 for n >= 4, with a(n) = 2n - 1 for n in {1, 2, 3}. Equivalently a(n) = A000217(n-1) + 3 for n >= 4. Predicts a(10) = 48.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Exley_, Apr 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017641 | OEIS-A395513

a(n) is the smallest number where the largest density of numbers with the same signature as A025487(n) among the first a(n) positive integers occurs.

Let Q(s, n) be the number of numbers <= n with prime signature s. Then a(n) is the (conjectured) smallest value k that maximizes the ratio Q(s, k)/k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David A. Corneth_, Apr 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017642 | OEIS-A395520

First differences of A036349.

Empirical observation: the maximum delay of 47205 steps is set near n = 48000 and never exceeded. This suggests the existence of an absolute constant C bounding all such delays (the Bounded Prime Debt Conjecture). See MathOverflow question 510477.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James W Worley_, Apr 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017643 | OEIS-A395522

Square table read by antidiagonals downwards: T(n,k) is the period of {F^(n)(x) mod k} for n >= 1, k >= 1, where F^(n) means the n-th iteration of the Fibonacci sequence (A000045).

Conjecture: each column is eventually constant. See the Math Stack Exchange link below for a more detailed introduction of the background of this conjecture. Note that, since T(n,k) can divide properly T(n-1,A001175(k)), this eventual period of {F^(n)(x) mod k} for sufficiently large n is not necessaily equal to A235249(k), the fixed point when A001175 is iterated with starting value k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Apr 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017644 | OEIS-A395531

Lay down bricks of length 1, 2, 3, ... on the top-right quadrant of the x-y plane. Place a brick as close to x=0 as possible, and on the highest row with bricks directly underneath where the new brick does not overhang. a(n) is the length of the leftmost brick on each row.

Conjecture: when n = a(m) for some m, we have a(n) = n(n+3)/2 - m. Verified experimentally up to m=1172, n=688491. - _Aleksei Udovenko_, May 07 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christian Perfect_, Apr 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017645 | OEIS-A395546

Number of satisfiable 3-SAT formulas with n variables and 5 clauses in the multiset clause model.

Using the conjectured formula for A396491, a(n) = binomial(binomial(2*n+2,3)+4,5) - (129332*n^6 - 2237100*n^5 + 16427761*n^4 - 63136975*n^3 + 130884126*n^2 - 135887564*n + 53820480)/3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ramin Mohammadi Masoudi_, Jun 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017646 | OEIS-A395565

Primes p preceded and followed by primes whose difference is greater than 2*log(p).

Conjecture: lim_{n->oo} n / pi(a(n)) = 3/e^2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Apr 28 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017647 | OEIS-A395605

a(n) = ((Lucas(prime(n)) - 1) / prime(n)) mod prime(n).

Conjecturally, for prime(n) != 2, 5, a(n) is the leading base-prime(n) digit of a p-adic invariant lambda(prime(n)) in Z_{prime(n)}, defined as the limit of the sequence u_k(p) = B(p^k)/p^k as k -> infinity. This p-adic interpretation is conjectural; the definition of a(n) is not.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Simone Cammarasana_, Apr 30 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017648 | OEIS-A395721

Largest k < m^2 such that both rad(k) and rad(k+1) divide m, where rad = A007947 and m = A005117(n), or -1 if such k does not exist.

Conjecture: for odd A005117(n), a(n) = -1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, May 04 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017649 | OEIS-A395792

Numbers k such that k-3 and k+3 are prime and gcd(k, sigma(k)) is prime.

Most computed terms satisfy gcd(k, sigma(k)) = 2; conjecture: this holds with natural density 1 within the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aied Sulaiman_, May 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017650 | OEIS-A395802

Minimum number of cells in a connected enclosing shell of an n-cell connected hole on the trihexagonal (3.6.3.6) Archimedean tiling.

Conjecture (UNVERIFIED, matches a(1..22)): a(n) grows as sqrt(18*n); equivalently the largest hole an s-cell shell can enclose is floor((s^2 + 6)/18), and the achievable shell sizes are 5, 11, and the odd integers >= 15 (so 7, 9, and 13 never occur). Each value is established as minimal over holes lying within a bounded region of the 3.6.3.6 tiling; that the region is large enough to exclude a smaller shell arising from a larger hole is supported by a wider-region check but not proved, so global minimality is conjectured. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Exley_, May 07 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017651 | OEIS-A395833

G.f. A(x) satisfies [x^n] A(x/A(x)^(2*n-1)) = 0 for n > 1.

Conjecture: (2*n-1) divides a(n) for n >= 1. Conjecture: a(n) == 0 (mod 3) for n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 07 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017652 | OEIS-A395835

Define F(n,x) as the power series in x satisfying F(n,x) = 1 + x*F(n,x)^2 - x^n*F(n,x)^(2*n), then for n >= 2, a(n) = M where x^M is the first power of x with a negative coefficient in F(n,x). Should no such M exist for some n, then there are no further terms and this sequence would be finite.

Conjecture: this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 14 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017653 | OEIS-A395836

Define F(n,x) as the power series in x satisfying F(n,x) = 1 + x*F(n,x)^2 - x^n*F(n,x)^(2*n), then for n >= 2, a(n) = abs([x^M] F(n,x)) where x^M is the first power of x with a negative coefficient in F(n,x). Should no such M exist for some n, then there are no further terms and this sequence would be finite.

Conjecture: this sequence is infinite. (4) lim_{n->oo} a(n)^(1/A395835(n)) = 4 (conjectural, if sequence is infinite).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 14 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017654 | OEIS-A395837

Define F(n,x) as the power series in x satisfying F(n,x) = 1 + x*F(n,x)^2 - x^n*F(n,x)^(2*n), then for n >= 2, a(n) = [x^(M-1)] F(n,x) where x^M is the first power of x with a negative coefficient in F(n,x). Should no such M exist for some n, then there are no further terms and this sequence would be finite.

Conjecture: this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 07 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017655 | OEIS-A395838

First differences of A395835.

Conjecture: this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 14 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017656 | OEIS-A395839

G.f. 10*C(x)^2 - C(x), where C(x) = 1 + x*C(x)^2 is the g.f. of the Catalan numbers (A000108).

Conjecture: a(n) is odd iff n is of the form 2^k - 1 (k >= 0). Conjecture: a(3*n+1) == A000108(n) (mod 3) for n >= 0 (see A111019 and A111020), with a(3*n) == a(3*n+2) == 0 (mod 3) for n >= 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 17 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017657 | OEIS-A395909

Triangle read by rows: row n gives marks on lexicographically earliest n-mark optimal ruler.

A ruler is optimal if it is perfect and no perfect ruler with the same number of segments has a greater length. Optimal rulers are rare. There are 155050 perfect rulers with length in [0, 123], but only 74 optimal rulers in this range. No blueprint for these rulers is known. It has been conjectured by the author that an optimal ruler with more than 13 segments is either a Wichmann ruler or the mirror image of a Wichmann ruler. For more definitions and references see A103294; more optimal rulers can be found under the Luschny link.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, May 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017658 | OEIS-A395970

Lexicographically earliest sequence of nonprime terms such that a(n) shares a factor with exactly n other terms.

There appears to be no easy method of generating each a(n) other than choosing the earliest semiprime that satisfies the condition that it can share with n other terms while not violating the same criteria for all other terms it may share a

factor with. To achieve this when a prime appears as a factor for the first time one must track the number of subsequent terms that must contain that as a factor so that a(n) can be satisfied. Each time that prime appears as a factor in subsequent terms that number is decremented; when it reaches zero no further terms can contain that prime as a factor. When a candidate semiprime is considered one must calculate the number of required terms still to come that must contain its primes as factors, and add to that the number of terms that have already appeared that contain these primes as factors. If that sum equals n then that implies the candidate semiprime can be satisfied so it can be chosen as the next term. Typically the sum will be less than n, implying that the primes for the candidate will not be able to be chosen as factors of subsequent terms before the candidate term has shared with n other terms. In this case the next lowest semiprime candidate must then be trialled. For the terms studied it appears this method never allows the square of primes to be terms, and it conjectured this is true for all n>1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 12 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017659 | OEIS-A395976

Numerators of rational coefficients related to asymptotic of A336293.

Conjecture: A395976(n)/A395977(n) ~ n^(n - 1/2) / (sqrt(Pi) * 2^(2*n - 3/2) * exp(n-2)). - _Vaclav Kotesovec_, May 21 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, May 12 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017660 | OEIS-A396013

a(1) = 2; for n > 1, a(n) is the smallest unused square spiral number such that a(n) shares a factor with all existing numbers in its Moore neighborhood.

A variation of A354434; here we start at 2 instead of 1. Like that sequence the primes, which do not occur in their natural order, are delayed relative to similarly sized terms, e.g. a(10507) = 173. It is conjectured that all numbers >= 2 eventually appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 18 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017661 | OEIS-A396018

Smallest number m such that m*(m+1)*(m+2)*(m+3)*(m+4) has exactly n distinct prime factors.

Conjecture: a(n) is an increasing sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhining Yang_, May 14 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017662 | OEIS-A396066

a(1) = 1; a(n) is the smallest positive integer not already in the sequence so that a(n)^4+a(n-1)^4 is prime.

It is conjectured that this list is a permutation of the natural numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hoang Nguyen_, May 15 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017663 | OEIS-A396081

Primes p such that A381466(2p) < 2p.

We conjecture that this is an infinite sequence. If so, then A394761 contains infinitely many 1's. It would also prove that there are infinitely many steps of A381466 at which we divide the index by 2. This would confirm one case of our conjecture that every positive integer is a divisor in A381466 infinitely often.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sam Chapman_, May 16 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017664 | OEIS-A396093

G.f. A(x) = B(B(B(x))), the third iteration of B(x) = x/(1-x)^2.

Conjecture: a(2*n) is even for n >= 1. Conjecture: a(2*n-1) is even iff n is of the form 5*k-2 (k >= 1) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 19 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017665 | OEIS-A396097

G.f. A(x) satisfies Sum_{k=0..n} [x^k] A(x)^n = Sum_{k=0..2*n} [x^k] C(x)^n for n >= 0, where C(x) = 1 + x*C(x)^2 is the g.f. of the Catalan function.

Conjecture: d is the largest real root of the equation -132400 + 2959788*d - 23123461*d^2 + 55640000*d^3 - 43613528*d^4 + 6629032*d^5 - 490000*d^6 + 12500*d^7 == 0. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, May 28 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017666 | OEIS-A396099

G.f. satisfies A(x) = x + A^2(x)*A^3(x).

Conjecture: all terms are odd. Conjecture: a(n) = [1,3,3,1] repeating (mod 4) for n > 2. Conjecture: [x^n] A(A(x)) == 0 (mod 4) for n > 2. Conjecture: [x^n] (A(x) - x*A(A(A(x)))) == 2 (mod 4) for n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 02 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017667 | OEIS-A396102

G.f. A(x) satisfies A(A(A(x))) = (1+x) * A(A(x)).

Conjecture: a(n) == 1 (mod 3) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017668 | OEIS-A396111

a(n) = n^4 mod (2*n+1).

Conjecture: a(n) contains every nonnegative integer k at least once (tested up to k=600000). Conjecture: a(n) contains every integer k only a finite number of times.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stanislav Sykora_, May 17 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017669 | OEIS-A396178

Lexicographically earliest sequence of the form 1, 2, followed by squarefree semiprimes (A006881), with the property that (a) for all k>=0, a(k) shares a factor with exactly k other terms, and (b) for all k >= 1, a(k) is divisible by a prime which does not divide any earlier term.

This is the same as A395795 except that the latter has a(1) = 4. Although the latter sequence has historical precedence, its definition was initially conjectural, and the present sequence is both simpler and belongs to a family of sequences (which is why the database now contains both versions).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, May 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017670 | OEIS-A396188

Index of 2^n in A394166, or -1 if 2^n does not appear.

It is a strong conjecture that every power of 2 appears in A394166.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017671 | OEIS-A396203

Primes p such that the gap to the next prime is floor(log_2(p)).

This sequence is conjectured to be infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sajid Hussain_, May 18 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017672 | OEIS-A396234

Smaller term of pairs of consecutive primes of the form p, p+6 where both primes eventually reach a value of 1 under iteration of the sums of the squares of their digits and the first iteration of both terms are also prime.

Conjecture: this sequence has an infinite number of terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Harry E. Neel_, May 19 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017673 | OEIS-A396265

Numbers k such that there exists a number r < k such that both gcd(r,k) > 1 and rad(r) does not divide k, yet rad(r+m*k) | k for some m, where rad = A007947.

Conjecture: includes A126706 \ {12, 24}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, May 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017674 | OEIS-A396324

Numbers k for which there exists a residue class r (mod k) with s = r+m*k, m >= 0, such that rad(s) | gcd(s,k), where rad = A007947.

Conjecture: A359280 is a proper subset (and thus, all powerful numbers greater than 4, i.e., A001694 \ {1, 4} = A359280 U A072777 / {4}).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jun 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017675 | OEIS-A396354

Number of unsatisfiable 3-SAT formulas with n variables and 4 clauses in the multiset clause model.

For fixed m, the corresponding column is expected to be a polynomial in n, since an m-clause formula involves only finitely many literal positions. The polynomial in the Formula section was obtained by interpolation from the listed terms and should be treated as conjectural unless independently verified. Conjecture: a(n) = n*(16*n^6 + 48*n^5 + 340*n^4 + 180*n^3 + 2818*n^2 - 10011*n + 6789)/18. The conjectured formula is true.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ramin Mohammadi Masoudi_, May 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017676 | OEIS-A396357

a(1) = 2; for n > 1, a(n) is the smallest unused number placed on a hexagon spiral that has a common factor with any earlier term that is adjacent to it on the spiral.

It is conjectured that every number greater than 1 appears.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017677 | OEIS-A396456

Smallest prime greater than the n-th prime-complete number.

A double (two prime-complete numbers in one prime gap) occurs only for the shared next-primes 37, 97, 487, and 4621; no further double exists below 10^12, and a heuristic based on the density of prime-complete numbers suggests this list of doubles is complete and that no triple occurs. That is, the counting function for prime-complete numbers, A377028(n) - 1, grows sub-polynomially (empirically exp(O(sqrt(log n)))), so prime-complete numbers thin out faster than any polynomial while prime gaps grow only logarithmically. The expected number of further doubles beyond x is therefore a convergent sum, supporting the conjecture that the complete list of doubles is 37, 97, 487, 4621.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ken Clements_, May 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017678 | OEIS-A396491

Number of unsatisfiable 3-SAT formulas with n variables and 5 clauses in the multiset clause model.

For fixed m, the corresponding column is expected to be a polynomial in n, since an m-clause formula involves only finitely many literal positions. The polynomial in the Formula section was obtained by interpolation from the listed terms and should be treated as conjectural unless independently verified.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ramin Mohammadi Masoudi_, May 28 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017679 | OEIS-A396493

Number of satisfiable 3-SAT formulas with n variables and 4 clauses in the multiset clause model.

Conjecture: a(n) = binomial(binomial(2*n+2,3)+3,4) - n*(16*n^6 + 48*n^5 + 340*n^4 + 180*n^3 + 2818*n^2 - 10011*n + 6789)/18, using the conjectured formula for A396354.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ramin Mohammadi Masoudi_, May 28 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017680 | OEIS-A396527

a(n) is the least number that is the sum of two distinct upper twin primes in exactly n ways, or -1 if there is no such number.

Conjecture: a(n) > 0 for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, May 28 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017681 | OEIS-A396563

Array read by ascending antidiagonals: A(n, k) is the determinant of the k X k matrix, where the (i, j)-th entry is |Stirling1(n + i + 2, n + j)|.

Conjecture: B(n, k, m) = (Sum_{j_1=0..n} Sum_{j_2=0..j_1} ... Sum_{j_m=0..j_{m-1}} (-1)^(j_1 + j_2 + ... + j_m) * (Product_{1 <= q < p <= m} (j_q - j_p + p - q)^2) * Product_{t=1..m} (n+t-1)_{j_t} * (n-j_t+t)^(n+k) / (j_t+m-t)!) / (Product_{i=0..m-1} (n+i)!) where B(n, k, m) is the determinant of the k X k matrix with (i, j)-th entry |Stirling1(n+m+i, n+j)|, where (x)_{n} is the falling factorial defined as (x)_{n} = Product_{k=0..n-1} (x-k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mikhail Kurkov_, May 29 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017682 | OEIS-A396570

Numbers == 2 (mod 6) that are not the sum of two distinct upper twin primes.

Conjecture: the sequence is finite and these are all the terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Robert Israel_, May 29 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017683 | OEIS-A396596

Natural numbers k such that the sum of the x-th powers of the divisors of k achieves a record, for some x <= 0.

This sequence includes the highly composite numbers (A002182) for which x = 0; the superabundant numbers (A004394) for which x = -1; and the deeply composite numbers (A095848) for which x << 0. We note that superabundant numbers less than 10^11 are either highly composite or deeply composite, or both. We conjecture that the present sequence can be constructed simply as a union of highly composite numbers and deeply composite numbers. All terms must be the least of their prime signature (A025487). The intersection of the above sequences is A095849. The conjecture above is false. For example, A004394(75) = 13974537376800 is superabundant, and like all terms of A004394, qualifies as a term of this sequence. However, it is neither highly composite (A002182) nor deeply composite (A095848); and there are other such terms. This is the least counterexample with x = -1. It would be interesting to know the least counterexample for any x < 0. This makes the production of additional terms of this sequence more difficult, because records could be set for the sum of the x-th powers of the divisors of k where k is neither highly composite nor deeply composite. - _Hal M. Switkay_, Oct 05 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hal M. Switkay_, Aug 31 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017684 | OEIS-A396603

Lexicographically earliest sequence of distinct nonnegative integers whose binary values, when written on a square spiral, one bit per square, high order bits first, contain no pair of 1 bits that are a knight's move apart.

It is conjectured that the sequence contains all nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 30 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017685 | OEIS-A396604

Lexicographically earliest sequence of distinct positive integers whose binary values, when written on a square spiral, one bit per square, high order bits first, contain no pair of 1 bits that are a knight's move apart.

It is conjectured that the sequence contains all positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 30 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017686 | OEIS-A396605

The smallest difference between n and the product of a nontrivial divisor of n with the totient of that divisor.

Conjecture: the set of n such that a(n) = 0 has natural density 0. Conjecture is true since Janous showed that the number of zero terms <= n ~ 1.36513.. * sqrt(n) (see A082473). - _Chai Wah Wu_, Jun 29 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pierre Colmant_, May 30 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017687 | OEIS-A396607

Lexicographically earliest sequence of distinct nonnegative integers whose binary values, when written on a square spiral, one bit per square, high order bits first, contain no pair of 1 bits that are diagonally adjacent.

It is conjectured that the sequence contains all nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 31 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017688 | OEIS-A396608

Lexicographically earliest sequence of distinct positive integers whose binary values, when written on a square spiral, one bit per square, high order bits first, contain no pair of 1 bits that are diagonally adjacent.

It is conjectured that the sequence contains all positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, May 31 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017689 | OEIS-A396659

Irregular triangle read by rows: T(n,k) is the number of edge-collision walks of length k from a fixed vertex on the complete directed graph with self-loops on n labeled vertices that visit all n vertices; n >= 1, n+1 <= k <= n^2+1.

Conjecture: T(n, n^2) / T(n, n^2+1) = (2n-1)/n for n >= 2. Verified for n = 2, 3, 4, 5, 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arkadeep Dutta_, Jun 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017690 | OEIS-A396695

Decimal expansion of the spectral radius of the matrix L[p_i, p_j] = 1/lcm(p_i, p_j), indexed over all primes p_i, p_j.

Conjecture (numerical evidence): lambda_max(W_N')/N -> lambda_inf, where W_N'[i,j] = floor(N/lcm(p_i,p_j)) for primes p_i, p_j <= N.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alessandro Munari_, Jun 02 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017691 | OEIS-A396696

a(n) is the smallest prime greater than n^2*a(n-1), with a(1) = 2.

Motivated by the following conjecture (see Guedes and Torres link): Is there a real number C so that [C*(n!)^2] is always a prime number for any positive integer?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Anderson Torres_, Aug 19 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017692 | OEIS-A396722

a(n) = (4^n/2) * p(n, 5/2), where p(n,x) is the Lagrange interpolating polynomial through (1,0) and (k,1) for k=2..n+1.

The recurrence conjectured by _R. J. Mathar_ on Jun 30 2026 follows directly from the interpolation conditions. Let p(n,x) be the polynomial in the definition. For n >= 1, we have n*p(n,x) = (n+1-x)*p(n-1,x) + x-1, since both sides are polynomials of degree at most n that agree at x = 1, 2, ..., n+1. Evaluating this at x = 5/2 and using a(n) = 4^n*p(n,5/2)/2 yields n*a(n) - (4*n-6)*a(n-1) = 3*4^(n-1). For n >= 2, subtracting four times this exact identity (with n replaced by n-1) gives n*a(n) - (8*n-10)*a(n-1) + (16*n-40)*a(n-2) = 0, proving the conjecture. - _Eiver Rodríguez_, Sep 28 2026 Conjecture D-finite with recurrence n*a(n) +2*(-4*n+5)*a(n-1) +8*(2*n-5)*a(n-2)=0. - _R. J. Mathar_, Jun 30 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017693 | OEIS-A396723

a(n) = (-4^n/2) p(n, -1/2), where p(n,x) is the Lagrange interpolating polynomial through (1,0) and (k,1) for k=2..n+1.

Conjecture D-finite with recurrence n*a(n) +2*(-4*n-1)*a(n-1) +8*(2*n+1)*a(n-2)=0. - _R. J. Mathar_, Jun 30 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017694 | OEIS-A396724

a(n) = (-4^n/2) p(n, -3/2), where p(n,x) is the Lagrange interpolating polynomial through (1,0) and (k,1) for k=2..n+1.

Conjecture D-finite with recurrence n*a(n) +2*(-4*n-3)*a(n-1) +8*(2*n+3)*a(n-2)=0. - _R. J. Mathar_, Jun 30 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Clark Kimberling_, Jun 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017695 | OEIS-A396758

Larger term of pairs of consecutive primes of the form p, p+6 where both primes eventually reach a value of 1 under iteration of the sums of the squares of their digits and the first iteration of both terms are also prime.

Conjecture: This sequence has an infinite number of terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Harry E. Neel_, Jun 04 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017696 | OEIS-A396760

Positive integers k at which A122921(k) - ceiling(sqrt(k/4)) reaches a new record value.

Conjecture: the sequence consists exactly of 1, 11, 53 and the three chains 96*4^m, 224*4^m, 2816*4^m (m >= 0), merged in increasing order. This formula gives all 37 listed terms below 10^9; beyond that bound it is conjectural.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nenad Micic_, Jun 04 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017697 | OEIS-A396785

Primes that do not appear at division steps of A381466.

We conjecture that there are infinitely many terms in this sequence. If this is true, then there exist infinitely many odd semiprimes that are addition steps, which, along with all odd primes and their squares, would be another infinite collection of odd addition steps that is "reasonably describable".

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sam Chapman_, Jun 05 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017698 | OEIS-A396793

G.f. A(x) satisfies A(x) * A(A(x)) = x^2 + 9*x^3.

Conjecture: a(n) == 0 (mod 3) for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 07 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017699 | OEIS-A396794

G.f. A(x) satisfies A(x) * A(A(A(x))) = x^2 + 16*x^3.

Conjecture: a(n) == 0 (mod 4) for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 07 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017700 | OEIS-A396795

G.f. A(x) satisfies A^2(x) * A^3(x) = x^2 + 25*x^3, where A^n(x) is the n-th iteration of A(x).

Conjecture: a(n) == 0 (mod 5) for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017701 | OEIS-A396797

G.f. satisfies A(x) = x + A^3(x)*A^4(x).

Conjecture: a(n) = 1 (mod 6) for n >= 1. Conjecture: [x^n] A^k(x) == k^(n-1) (mod 6) for n >= 1 and all integer k. The congruence conjectures are proved in Sundaram (2026). - _Shiva Priyan Shanmuga Sundaram_, Sep 13 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 15 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017702 | OEIS-A396798

G.f. satisfies A(x) = x + A^4(x)*A^5(x).

Conjecture: a(n) = [1,1,5,5] repeating (mod 8) for n >= 1. Conjecture: [x^n] A^2(x) == 0 (mod 8) for n > 3. Conjecture: [x^n] A^3(x) == [3,1,7,5] repeating (mod 8) for n > 1. Conjecture: [x^n] A^4(x) == 0 (mod 8) for n > 2. Conjecture: [x^n] A^5(x) == [5,1,1,5] repeating (mod 8) for n > 1. Conjecture: [x^n] A^6(x) == 0 (mod 8) for n > 3. Conjecture: [x^n] A^7(x) == [7,1,3,5] repeating (mod 8) for n > 1. Conjecture: [x^n] A^8(x) == 0 (mod 8) for n > 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 16 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017703 | OEIS-A396803

E.g.f. satisfies A(x) = x*exp(A^3(x)).

Conjecture: a(n) is odd iff n is odd for n >= 1. Conjecture: a(n) == [1,2,0] repeating (mod 3) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 08 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017704 | OEIS-A396804

E.g.f. satisfies A(x) = x*exp(A^4(x)).

Conjecture: a(n) is odd iff n is odd for n >= 1. Conjecture: a(n) == [1,2,0] repeating (mod 3) for n >= 1. [This fails at n=5, since a(5) = 41005 == 1 (mod 3), not 2. - _Danial Asaria_, Jul 26 2026] Conjecture: a(n) == [1,2,3,0] repeating (mod 4) for n >= 1. Conjecture: a(n) (mod 5) is never congruent to 3. Conjecture: a(n) == [1,2,0,1,1,0] repeating (mod 3) for n >= 1. This pattern holds through n=201. - _Danial Asaria_, Jul 26 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017705 | OEIS-A396805

E.g.f. satisfies A(x) = x*exp(A^5(x)).

Conjecture: a(n) is odd iff n is odd for n >= 1. Conjecture: a(n) == [0,1,0] repeating (mod 3) for n >= 3. Conjecture: a(n) == [1,2,3,4,0] repeating (mod 5) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017706 | OEIS-A396806

E.g.f. satisfies A(x) = x*exp(A^6(x)).

Conjecture: a(n) is odd iff n is odd for n >= 1. Conjecture: a(n) == [1,2,0] repeating (mod 3) for n >= 1. Conjecture: a(n) == [1,2,3,4,5,0] repeating (mod 6) for n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017707 | OEIS-A396807

G.f. satisfies A(x) = x + A^5(x)*A^6(x).

Conjecture: a(n) = 1 (mod 10) for n >= 1. Conjecture: [x^n] A^k(x) == k^(n-1) (mod 10) for n >= 1 and all integer k.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 16 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017708 | OEIS-A396808

G.f. A(x) satisfies (n+1) * [x^n] A(x)^(n+1) = n * [x^n] A(x)^(n+2) for n > 1.

Conjecture: for n > 0, a(n) is odd iff n is a power of 2. Conjecture: for n > 1, a(n) == 2 (mod 3) iff n is a power of 3. Conjecture: for n > 1, a(n) == 1 (mod 3) iff n = (3/2)*A038464(k) for some k, where A038464 lists the sums of 2 distinct powers of 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 18 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017709 | OEIS-A396836

Lexicographically earliest sequence of distinct positive integers whose binary values, when written on a square spiral, one bit per square, high order bits first, contain no pair of 0 bits that are a knight's move apart.

Although the terms equal to powers of 2 are delayed relative to similar size terms, e.g. a(4645) = 256, it is conjectured that the sequence contains all positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 07 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017710 | OEIS-A396837

Lexicographically earliest sequence of distinct nonnegative integers whose binary values, when written on a square spiral, one bit per square, high order bits first, contain no pair of 0 bits that are a knight's move apart.

Although the terms equal to powers of 2 are delayed relative to similar size terms, e.g. a(2214) = 128, it is conjectured that the sequence contains all nonnegative integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 07 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017711 | OEIS-A396843

G.f. A(x) satisfies A(x*A(x) - 3*x*A(x)^2) = x^2.

Conjecture: all terms are odd. Conjecture: for n > 3, a(n) == 1 (mod 9) iff n is of the form 4^k+1, otherwise a(n) is divisible by 9. [This conjecture is false: n=5=4^1+1, but

a(5)=975 == 3 (mod 9), not 1. - _Danial Asaria_, Jul 26 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017712 | OEIS-A396844

G.f. A(x) satisfies A(x*A(x) - 4*x*A(x)^2) = x^2.

Conjecture: for n > 8, a(n) == 12 (mod 16) iff n is of the form 2*4^k+1 otherwise a(n) is divisible by 16.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 03 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017713 | OEIS-A396857

Conjectured numbers of the form abs(x^^y - w^^z), where x, y, w, z are integers greater than 1 and ^^ denotes tetration.

Conjectured numbers of the form abs(x^^y - w^^z), where x, y, w, z are integers greater than 1 and ^^ denotes tetration. For each n >= 4, the conjectured a(n) has been obtained by an exhaustive computational search over all integer tetrations up to 10^50000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Marco Ripà_, Aug 19 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017714 | OEIS-A396859

Minimum number of cells in a connected enclosing shell of an n-cell connected hole on the snub-square (3.3.4.3.4) Archimedean tiling.

Conjecture: a(n) - a(n-1) is in {-1, 0, 1} for n >= 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Exley_, Jun 08 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017715 | OEIS-A396869

a(n) = 4*q - floor(2*sqrt(q))^2, where q = A246655(n).

Conjecturally, for every d == 0, 3, 4 (mod 8), there are only finitely many solutions (x,p,n) to x^2 + d = 4*p^n, with p >= 3 prime, n >= 3 odd. See the Muriefah et al link for the cases of small values d == 0 (mod 4), and the Luca et al link for the cases of small values d == 3 (mod 4). (Here only the case p not dividing d is treated in the Luca et al link, but the case p dividing d can be solved using elliptic curves, similar to the previous paragraph). In particular, the indices n of d <= 20 are as follows, where q = A246655(n):

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Jun 08 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017716 | OEIS-A396884

Number of distinct smallest connected polyiamonds that contain every free n-iamond, up to rotations, reflections and translations of the triangular lattice.

No closed form or low-order recurrence is known for the nine listed terms (conjectured to have none, UNVERIFIED). The listed values are the number of minimum-size containers found within a finite search window whose width equals n (forced by the straight n-iamond) and whose height grows as max(4, floor((n+2)/3)); global minimality over all bounding boxes is conjectured but not separately certified.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Exley_, Jun 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017717 | OEIS-A396889

Number of polynomials f(x) of degree <= n with f(0) = 1 and coefficients in {0, 1, ..., n} such that f(k) is prime for all k = 1, 2, ..., n.

Based on heuristic probabilistic models involving the Prime Number Theorem and Mertens' theorem, it is conjectured that a(n) grows exponentially. Conjecture: a(n) ~ c * exp(gamma * n) for some constant c > 0, where gamma is the Euler-Mascheroni constant A001620.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yuanhao Ding_, Jun 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017718 | OEIS-A396908

Number of decimal places which are known after using n terms of A390946 and treating them as the coefficients of a continued fraction.

Conjecture: Limit_{n->oo} n/a(n) = A086819 (Lochs's constant).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Jul 16 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017719 | OEIS-A396946

Primes p such that both Mordell equations y^2 = x^3 + p^2 and y^2 = x^3 - p^2 have nonzero integer solutions.

Conjecture: All terms can be expressed in the form of 4*m^2 + 1 or 12*m^2 - 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Zhining Yang_, Jun 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017720 | OEIS-A396960

Conjectured list of numbers whose trajectory under the 15x+1 map eventually reaches 1, where the map sends odd x to 15x+1 and even x to x/2.

Conjectured list of numbers whose trajectory under the 15x+1 map eventually reaches 1, where the map sends odd x to 15x+1 and even x to x/2. "Conjectured" in the same sense as A267703: every term shown provably reaches 1, but it is conjectural that the numbers not shown never do (a trajectory might in principle descend after exceeding the search bound). Heuristically each odd step multiplies by about 15/4 on average, so almost all positive integers should have divergent trajectories and this sequence should have density 0; neither statement has been proved (Kontorovich and Lagarias state the corresponding claim for the 5x+1 map as a conjecture). - _Karan Sharma_, Oct 03 2026 Conjecturally the number of terms <= X grows like X^(c+o(1)) with c = 0.13100579683..., the root in (0,1) of 15^(c-1) = 2^c - 1. Equivalently, c is the maximum over 0 < b < 1/L of (H(b) - b*L)/(1 - b*L), where L = log_2(15) and H(b) = -b*log_2(b) - (1-b)*log_2(1-b); the maximum occurs at b = 1/(1 + 15^(1-c)). The root equation comes from a branching-process model of the tree of preimages of 1; cf.

Kontorovich and Lagarias for stochastic models of this kind for the 5x+1 map. There are 57, 87, 123, 182 terms <= 10^4, 10^5, 10^6, 10^7, respectively. - _Karan Sharma_, Oct 03 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Karan Sharma_, Jun 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017721 | OEIS-A396966

a(n) is the number of obstruction residues modulo 2^n for the Collatz-type 3n-1 map T(x) = (3x-1)/2^A007814(3x-1).

a(n)/2^n is strictly increasing; lim_{n->oo} a(n)/2^n = c_W exists. The lower bound c_W >= 172651809822/2^40 = 0.157025906... is proved (from a(40) = 172651809822). The exact value of c_W is an open problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jakob Stemberger_, Jun 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017722 | OEIS-A396969

Intersection of A332785 and A396324.

Conjecture: A396324 is the union of this sequence and A001694 \ {1,4}.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jun 13 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017723 | OEIS-A397045

Number of distinct values in the interval [n,n+1) that can be represented as a sum of square roots of positive integers.

Conjecture: log a(n) ~ k*n^(2/3). Possibly k = (81*zeta(3)/Pi^2)^(1/3) = 2.1447.... - _Charles R Greathouse IV_, Jun 15 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ali Sada_, Jun 14 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017724 | OEIS-A397047

a(n) is the number of n-digit terms in A397046.

Conjecture: lim_{n->oo} a(n+1)/a(n) = 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jun 14 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017725 | OEIS-A397051

a(n) is the number of n-digit terms in A397050.

Conjecture: lim_{n->oo} a(n+1)/a(n) = 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jun 14 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017726 | OEIS-A397059

Number of free polyplets (king-move connected sets of n cells on the square grid, adjacent if they share an edge or a corner) whose king-adjacency graph is a tree.

Conjecture: a(n) grows at the same exponential rate as A030222, the tree restriction affecting only the sub-exponential factor (UNVERIFIED).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Exley_, Jun 15 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017727 | OEIS-A397127

Lexicographically earliest sequence of distinct numbers k that are neither squarefree nor prime powers (k in A126706), such that for any prime p, any run of consecutive multiples of p has length exactly 2.

Conjecture: There are no more duplex blocks {j,k} such that rad(j) = rad(k) but neither j nor k divides the other, apart from j = 12 and k = 18. Conjecture: permutation of A126706. Yumihiro Fujikawa provided a counterexample to the first conjecture above: a(25992) = 80675 = 5^2 * 7 * 461, a(25993) = 112945 = 5 * 7^2 * 461, using machine assisted tools and 2 independent program approaches. De Vlieger confirmed the find using Mathematica. - _Michael De Vlieger_, Oct 22 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jun 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017728 | OEIS-A397128

Lexicographically earliest sequence of distinct numbers k that are neither squarefree nor powerful (k in A332785), such that for any prime p, any run of consecutive multiples of p has length exactly 2.

Conjecture: permutation of A332785.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jun 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017729 | OEIS-A397150

Number of decimal places which are known after using n terms of A072193 and treating them as the coefficients of a continued fraction.

Conjecture: Limit_{n->oo} n/a(n) = A086819 (Lochs's constant).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Jul 15 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017730 | OEIS-A397153

Maximum length of a snake polyomino in a 12 X n rectangle.

Conjectured g.f.: x*(12 + 6*x + 8*x^2 + 7*x^3 + 7*x^4 + 7*x^5 - 4*x^6 + x^7 + 2*x^10 - x^11 - x^16 + x^17 - x^18)/((1 - x)^2*(1 + x)*(1 - x + x^2)*(1 + x + x^2)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Goupil_, Jul 17 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017731 | OEIS-A397155

Least prime p such that n^3 - p is semiprime.

This sequence is motivated by the computational conjecture that for every positive integer cube greater than 1, there exist primes p, q and r, such that n^3 = p*q + r.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Najeem Ziauddin_, Jul 24 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017732 | OEIS-A397174

The rows of A397173 converge to this sequence.

This sequence has the same status as A397471, A397175, and A397176: it is conjectural, although almost certainly correct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sean A. Irvine_ and _N. J. A. Sloane_, Jun 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017733 | OEIS-A397175

Numbers that appear in A394611.

The terms of A394611, with duplicates removed, sorted into increasing order. This is the complement of A397471. At present, since there is proof yet that 5, 9, ... (A397471) are missing from A394611, this sequence and A397176 have the same status as A397471: conjectural, but almost certainly correct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 27 2026. | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017734 | OEIS-A397176

Index in A394611 of the first appearance of A397175(n).

The terms are conjectural, but almost certainly correct (see A397471 and A397175).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jun 27 2026. | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017735 | OEIS-A397179

The rows of triangle A397178 converge to this sequence.

This formula (*) is unconditionally correct. But terms of the present sequence shown in the DATA section (since we do not know if 5, 9, ... will ever appear in A394611) have the same conjectural status as A397471, A397175, and A397176.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017736 | OEIS-A397185

a(n) is the smallest positive integer m such that every even integer from 2 to 2n can be expressed as a sum of two (not necessarily distinct) 1-bit positions in the binary expansion of m.

Conjecture: a(2k) = 2*a(2k-1) + 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jose Manuel Briceno Mendoza_, Jun 17 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017737 | OEIS-A397208

a(n) is the number of primes q < p = prime(n) such that (p + q)/4 is prime.

Conjecture: The only zeros occur for p = 2, 3, 11, 19, 43, 67. Verified numerically for n <= 10000 (p <= 104729). Conjecture: Every nonnegative integer occurs in this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Jun 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017738 | OEIS-A397209

a(n) is the smallest index k such that A397208(k) = n, or -1 if no such k exists.

It is conjectured that every nonnegative integer occurs in A397208.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Jun 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017739 | OEIS-A397218

Essentially the same as A397219, except insert an initial 1 and modify the definition accordingly.

It is conjectured that this is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 19 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017740 | OEIS-A397219

Pascal-like triangle read by rows: top entry is 2; thereafter fill in entries left to right, top to bottom, with smallest number m that is not yet in the sequence and has a common factor with any earlier entry that is adjacent to it in the triangle.

It is conjectured that every number greater than 1 appears (cf. A397218).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 19 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017741 | OEIS-A397241

G.f. A(x) satisfies n * [x^n] A(x)^n = (n-1) * [x^n] A(x)^(n+1) for n > 1.

Conjecture: all terms are odd. Conjecture: a(n) == 2 (mod 3) for n > 2. Conjecture: a(n) == 3 (mod 4) for n > 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 19 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017742 | OEIS-A397242

O.g.f. satisfies A(x) = exp(x + Sum_{n>=2} (n^2-1) * a(n)*x^n / n^2).

Conjecture: a(n) is odd iff n+1 is a power of 2. Conjecture: for n > 0, a(n) == 1 (mod 3) iff n+2 is of the form 3^k or 2*3^k. Conjecture: for n > 0, a(n) == 2 (mod 3) iff n+2 is a sum of 2 distinct powers of 3 (A038464).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 19 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017743 | OEIS-A397244

G.f. A(x) satisfies (2*n) * [x^n] A(x)^(2*n) = (2*n-1) * [x^n] A(x)^(2*n+1) for n > 1.

Conjecture: for n > 0, a(n) is odd iff n = 2*A003714(k) + 1 for some k >= 0, where A003714 lists the Fibbinary numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 24 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017744 | OEIS-A397245

O.g.f. satisfies A(x) = exp(x + Sum_{n>=2} (4*n^2-1) * a(n)*x^n / (4*n^2)).

Conjecture: a(n) == 1 (mod 3) iff n+2 is of the form 3^m or 2*3^m. Conjecture: a(n) == 2 (mod 3) iff n+2 is a sum of 2 distinct powers of 3 (A038464).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 24 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017745 | OEIS-A397257

Fixed points of A093811, sum of the digital products of the divisors.

We conjecture that there are no further terms, but we don't have a formal proof yet. We conjecture that for indices n ~ 10^k, record values of A093811 oscillate around 9^k. More precisely, A093811(n) < n for n > 10^3 (so this sequence is finite and complete), and A093811(10^k-2) = 9^k*4/3 sets a new record value for all k > 4, hence lim sup A093811(n)/n = 0.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jul 07 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017746 | OEIS-A397258

Indices of late birds in Buss's sequence B = A067836: numbers m such that all smaller primes have occurred before B(m).

To tackle Buss's conjecture (cf. A067836), it is useful to know what are the smallest primes that have not yet appeared up to a given point. They are given in A397259, and here we list the indices where they appear. It might seem more natural to consider "terms" rather than "primes" in the name. Conjecturally (i.e., if B lists all and only primes), the two are equivalent, i.e., this is the list of m such that B(k) >= B(m) for all k > m (and we know that equality is excluded). But if the conjecture isn't true, we can never know whether a smaller composite number might appear somewhere later in B. With the present definition, we know at once and for sure when a term can be listed.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017747 | OEIS-A397259

Late birds in Buss's sequence B = A067836: primes B(m) such that all smaller primes have appeared earlier.

To tackle Buss's conjecture (i.e., all and only primes appear in B = A067836), it is useful to know what are the smallest primes that have not yet appeared up to a given point. These are listed here, while the indices where they appear are given in A397258. Conjecturally this is equivalent to say that we list terms a(n) = B(m) such that B(k) > B(m) for all k > m. But we prefer the chosen definition because it makes the sequence unambiguously defined (and finite, in case one prime would never appear in A067836).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jun 22 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017748 | OEIS-A397260

Indices of records in A093811.

We conjecture that the following numbers are always in this sequence: A more complete list of simple patterns of numbers that we conjecture to be always in the sequence is as follows: {179...958, 179...976, 179...988, 19...96, 269...988, 29...9898, 29...988, 39...988,, 39...96, 479...988, 49...98, 579...98, 59...9898, 59...976, 59...988, 779...98, 79...988, 79...96, 79...98, 979...98, 9...96, 9...98}. Here, "9...9" means any positive number of digits 9.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. F. Hasler_, Jul 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017749 | OEIS-A397297

Index of n in A397219, or -1 if n never appears.

It is conjectured that every number >= 2 appears in A397219.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael S. Branicky_, Jun 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017750 | OEIS-A397315

Table read by antidiagonals: T(n,k) = maximum number of points that can be blacked out of an n X k grid of points so that every rectangle with all four vertices on grid points (tilted rectangles included) is uniquely identifiable from its set of visible vertices; n, k >= 2.

From Problem 001 of Peter Kagey's Open Problem Collection (https://peterkagey.com/problems/001/): black out as many grid points as possible while every rectangle with lattice corners remains identifiable from its visible corners; equivalently, injectivity of the map R -> C(R) \ S over all rectangles. Conjecture: T(3,m) = m+2 for all m >= 3 (verified for 3 <= m <= 10). Conjecture: T(n,m) = n+m whenever min(n,m) >= 4, except T(4,4) = 7 and T(4,5) = 8 (verified on seven cells through 6 X 6).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Arjun Pemmasani_, Sep 02 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017751 | OEIS-A397323

a(1) = 1; for n > 1, a(n) is the smallest unused number placed on a hexagon spiral that has a common factor with any earlier term that is adjacent to it on the spiral.

It is conjectured that all positive numbers appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017752 | OEIS-A397340

Lexicographically earliest sequence of distinct squarefree composite numbers, such that for any prime p, any run of consecutive multiples of p has length exactly 2.

Conjecture: a(n+1) is never coprime to a(n). This holds for n = 1..2^20. This would imply that this variant does not feature blocks in the sense they are defined for A280864.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael De Vlieger_, Jun 25 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017753 | OEIS-A397345

O.g.f. satisfies A(x) = exp(x + Sum_{n>=2} (5*n^2 - 1) * a(n)*x^n / (5*n^2)).

Conjecture: a(n) is odd iff n+1 is a power of 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 28 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017754 | OEIS-A397346

O.g.f. satisfies A(x) = exp(x + Sum_{n>=2} (2*n^2-1) * a(n)*x^n / (2*n^2)).

Conjecture: a(n) == [4,2,0,2] repeating (mod 8) for n >= 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017755 | OEIS-A397348

O.g.f. satisfies A(x) = exp(x + Sum_{n>=2} (3*n^2 - 1) * a(n)*x^n / (3*n^2)).

Conjecture: a(n) is odd iff n+1 is a power of 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jun 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017756 | OEIS-A397356

G.f. satisfies: [x^n] 1/A(x)^(n^2) = [x^n] 1/A(x)^(n^2-1) for n > 1 with A(0) = A'(0) = 1.

Conjecture: for n >= 0, a(n) is odd iff n+1 is a power of 2. Conjecture: for n >= 0, a(n) is not divisible by 3 iff 2*(n+1) is a sum of two powers of 3 (A055235).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 03 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017757 | OEIS-A397358

a(n) is the unique k >= 1 such that A400792(k) = n, or -1 if no such k exists.

A400792 is conjectured to be a permutation of the positive integers, so there should not be any -1 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Oct 13 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017758 | OEIS-A397369

a(n) = 64^n * [z^n] 2F1(1/4, 1/4; 1; z)^3.

Conjecturally, for every prime p == 1 (mod 4) and all m,r >= 1, a(m*p^r) == a(m*p^(r-1)) (mod p^(4*r)). The exponent is sharp: a(5) - a(1) = 5^4*43524, with 5 not dividing 43524. (The split restriction p == 1 (mod 4) mirrors the mixed-CM result of arXiv:2605.19773; the full depth-(4r) tower mirrors arXiv:2606.15462.) Conjecturally the exponent 4 is governed by the weight-5 control form C = (eta(tau)^12/eta(2*tau)^6)*(2*E_2(2*tau) - E_2(tau)), and the restriction p == 1 (mod 4) by the eta multiplier of self-Hecke modulus 4 and its 3/4-shifted Fourier lattice at the second cusp.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Shvets_, Jun 22 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017759 | OEIS-A397434

Number of monomials in the algebraic normal form (Zhegalkin polynomial over GF(2)) of the Boolean function that is 1 if and only if at least ceiling(n/2) of its n variables are 1.

Conjecture: a(n) = a(n+1) if and only if n == 2 (mod 4). Verified for n <= 998.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tanguy Gautier Loïc Le Mer_, Jul 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017760 | OEIS-A397443

Least integer such that floor(R(a(n))) = n, where R(x) is the Riemann prime counting function (slightly truncated).

Conjecture: n * (log(n)+log(log(n))-1+((log(log(n))-A)/log(n))) is asymptotic to a(n) if and only if A=2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Rocchelli_, Aug 08 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017761 | OEIS-A397445

a(n) is the permanent of the n X n Krawtchouk matrix.

Conjecture: a(n) = 0 iff n is congruent to {2, 3} mod 4 (cf. A042964).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Jun 25 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017762 | OEIS-A397456

Number of consecutive n-digit prime pairs with gap 6.

For 2 <= n <= 12, a gap of 6 is the most frequent gap ("jumping champion") among all occurring gaps between consecutive n-digit primes. This empirical dominance is supported by the Hardy-Littlewood prime-pair conjecture, as 6 is the first even number divisible by the two smallest primes 2 and 3, which maximizes its constellation constant over this range.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dieter Renz_, Jun 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017763 | OEIS-A397471

Numbers not appearing in A394611.

At present, all these terms are conjectural, but almost certainly correct.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sean A. Irvine_, Jun 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017764 | OEIS-A397519

Lexicographically earliest sequence of distinct nonnegative integers whose binary plot has no two 0 bits a knight's move apart.

It is conjectured that all nonnegative numbers appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 29 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017765 | OEIS-A397520

Lexicographically earliest infinite sequence of distinct nonnegative integers such that each 1-bit in the binary plot of the sequence is a knight's move apart from one, and only one, other 1-bit.

It is conjectured that all nonnegative numbers appear.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jun 29 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017766 | OEIS-A397551

a(n) is the number of repeated subset sums among binomial(n, 0), ..., binomial(n, floor((n - 1)/2)).

For odd n, a(n) appears in the conjectural formula A397501(n) = 2^((n - 1)/2) + 1 + a(n), where A397501 counts set partitions of {0, ..., n} whose block weights in row n of Pascal's triangle are powers of 2. Conjecture: a(n) = Sum_R 2^f(R), where the sum runs over all primitive relations R counted by A397559. The conjecture has been verified computationally for all n <= 55. In particular, if A397559(n) = 1 and the unique primitive relation R has support size s, then conjecturally a(n) = 2^f(R) = 2^(floor((n - 1)/2) + 1 - s).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Jul 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017767 | OEIS-A397578

a(n) is the least k such that sigma(k) > sigma(k-1)*n and sigma(k) > sigma(k+1)*n, where sigma(k) = A000203(k) is sum of divisors of k.

Conjecture: a(n) exists for all n.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Ratushnyak_, Jul 30 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017768 | OEIS-A397591

G.f. A(x) satisfies [x^(n-1)] (1 - A(x))^n / (1 - n*x) = 0 for n > 1.

Conjecture: for n > 3, a(n) is odd iff n is of the form 2^k-1 or 2^k+1 (k > 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017769 | OEIS-A397592

G.f. A(x) such that the sum of the first n coefficients in A(x/n)^n equals 2^(n-1) for n >= 1.

Conjecture: for n > 3, a(n) is odd iff n is of the form 2^k-1 or 2^k+1 (k > 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 10 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017770 | OEIS-A397594

G.f. A(x) such that the sum of the first n coefficients in A(x/n)^n equals 4^(n-1) for n >= 1.

Conjecture: for n > 3, a(n) is odd iff n is of the form 2^k-1 or 2^k+1 (k > 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 10 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017771 | OEIS-A397619

Number of permutations p of [n] such that p(i)*p(i+1) + 1 is prime for all 1 <= i <= n-1.

Since reversals are counted separately, a(n) is even for n>=2. No two odd numbers can be adjacent, as odd * odd + 1 is even and greater than 2. Therefore for odd n the permutation alternates parity. Conjecture: a(n)>0 for all n. Verified up to n = 27. For n>=2, a(n) > 0 implies a prime == 1 mod n, not exceeding n^2 - n + 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ethan Yu_, Jul 02 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017772 | OEIS-A397621

a(n) is the linear complexity over GF(2) of the binary expansion of n (as computed by the Berlekamp-Massey algorithm).

Conjecture: a(A001317(n)) = A080079(n).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Darío Clavijo_, Aug 02 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017773 | OEIS-A397656

a(n) is the smallest k such that A397224(k) = n or -1 if no such k exists.

I conjecture that k will always exist.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Žiga Pirc_, Jul 03 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017774 | OEIS-A397683

a(n) = f(2*n-1) where f(1)=1 and otherwise f(m) is the number of x, 1 <= x < m, such that gcd(x,m) = gcd(x+1,m) = 1 and ord_m(x) = ord_m(x+1).

Conjecture: for odd m, f(m) = 0 iff m is a power of 3 or a power of 7. This has been verified computationally for m < 20000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Donovan Sneider González May_, Sep 02 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017775 | OEIS-A397699

Associated primes of A397698.

Conjecture: All primes appear in this sequence. The conjecture is likely true (based on an argument by Allan Wechsler; see Links).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ali Sada_, Jul 05 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017776 | OEIS-A397711

Number of acyclic digraphs (or DAGs) on n labeled nodes in which every node has in-degree at most 2.

Conjecture (verified for n <= 100): a(n) == 1 (mod n) for all odd n >= 3, and a(n) == n/2 + 1 (mod n) for all even n >= 4. The conjecture is shown to be true for all n: the recurrence over sinks gives n | 2*(a(n) - 1), which settles the odd case; the even case follows from Burnside's lemma, using the fixed-point-free involutions. In the general case of in-degree <= r, n divides lcm(1,...,r)*(a_r(n) - 1). - _Roberto Blanco Gómez_, Oct 07 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tyler Satchel Orden_, Jul 05 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017777 | OEIS-A397730

a(n) is the number of distinct triples of triangles chosen from n^2 unit triangles of a triangular grid up to rotation such that no two triangles share common side.

Conjectured g.f. x^2 * (1+7*x+52*x^2+42*x^3+28*x^4-10*x^5-x^6+x^7) / ((1+x+x^2) * (1-x)^7). - _Sean A. Irvine_, Jul 14 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Janaka Rodrigo_, Jul 08 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017778 | OEIS-A397731

a(n) is the number of distinct triples of triangles chosen from n^2 unit triangles of a triangular grid up to rotation and reflection such that no two triangles share a common side.

Conjectured g.f.: x^2 * (1+6*x+32*x^2+75*x^3+78*x^4+49*x^5+4*x^6-3*x^7-3*x^8+x^9) / ((x^2+x+1) * (x+1)^2 * (1-x)^7). - _Sean A. Irvine_, Jul 14 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Janaka Rodrigo_, Jul 08 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017779 | OEIS-A397732

a(n) is the number of distinct triples of triangles chosen from n^2 unit triangles of a triangular grid up to rotation such that no two triangles share a common point.

Conjectured g.f.: 2*x^3 * (1+9*x+39*x^2+28*x^3+3*x^4-21*x^5-5*x^6+6*x^7) / ((1+x+x^2) * (1-x)^7). - _Sean A. Irvine_, Jul 14 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Janaka Rodrigo_, Jul 08 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017780 | OEIS-A397740

Partial sums of A399633.

Conjecture: limsup_{n->oo} a(n)/n <= 3.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Sep 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017781 | OEIS-A397755

Number of steps to reach 0 for the sequence b_n(m) defined in the A397774, or 0 if b_n(m) does not terminate.

Conjecture: all terms are positive.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bill McEachen_, Jul 08 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017782 | OEIS-A397757

Maximum length of a snake polyomino in a 6 X n rectangle.

Conjecture: a(n) = 2 + 15*n/4 + (n mod 4)/4 for n > 6.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alain Goupil_, Jul 08 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017783 | OEIS-A397774

Irregular triangle where the n-th row is the sequence b_n(m) defined in comments.

It is conjectured that b_n(m) reaches 0 for every n, at which point the sequence b_n(m) terminates because 0+p can never be composite. The index at which 0 occurs must necessarily be even. If this conjecture is false, then eventually we will encounter some row that does not terminate and the total number of rows in this sequence would then be finite, but the number of terms infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bill McEachen_, Jul 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017784 | OEIS-A397780

Lexicographically earliest numbers k such that the decimal digits of k and k^2 taken together form a set where each distinct digit d appears at least d times, and this set of digits has not appeared in any previous term.

Conjecture: The sequence has exactly 665 terms, distributed by the number of distinct digits in the target set (from 1 to 10) as: 2, 1, 2, 31, 243, 210, 120, 45, 10, 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dmytro Voievoda_, Sep 29 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017785 | OEIS-A397789

a(n) = nearest prime to prime(n)^(3/2).

Conjecture: A397790(n) = a(n)^2. [In other words, it is conjectured that there are no consecutive primes p1 < p2 and a prime q such that (p1 + p2)^2/4 < q^3 < (p1^2 + p2^2)/2. - _Jianing Song_, Jul 14 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James C. McMahon_ and _Vincenzo Manto_, Jul 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017786 | OEIS-A397790

a(n) = prime^2 closest to prime(n)^3.

Empirically, the closest prime square to prime(n)^3 is always unique. It's conjectured that no tie occurs. Conjecture: a(n) = A397789(n)^2. [In other words, it is conjectured that there are no consecutive primes p1 < p2 and a prime q such that (p1 + p2)^2/4 < q^3 < (p1^2 + p2^2)/2. - _Jianing Song_, Jul 14 2026]

猜想/问题提及段 | historical_or_current_mention_unreviewed | _James C. McMahon_ and _Vincenzo Manto_, Jul 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017787 | OEIS-A397806

Maximum number of lists of n distinct real numbers in [0, 1) that produce the same list when Gaston's algorithm is applied to them.

The following conjectures can be made: Assuming the third conjecture is true, the next two terms would be 7935 and 47181, that are respectively achieved for the lists [0.1, 0.23, 0.3, 0.45, 0.55, 0.65, 0.77, 0.9] and [0.1, 0.18, 0.26, 0.38, 0.48, 0.58, 0.7, 0.78, 0.9].

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Romain Malécot_, Jul 10 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017788 | OEIS-A397823

Decimal expansion of Product_{p prime} (1-1/p)^4 * (1 + 4/p + 1/p^2).

This constant, c, appears in the conjectured asymptotic formula for the sixth power moment of the Riemann zeta function on the critical line: Integral_{t=0..T} |zeta(1/2 + i*t)|^6 dt ~ (42/9!) * c * T * log(T)^9 (Conrey and Gosh, 1989; Keating and Snaith, 1998; Conrey and Gonek, 2001).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Jul 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017789 | OEIS-A397824

Decimal expansion of Product_{p prime} (1-1/p)^9 * (1 + 9/p + 9/p^2 + 1/p^3).

This constant, c, appears in the conjectured asymptotic formula for the eighth power moment of the Riemann zeta function on the critical line: Integral_{t=0..T} |zeta(1/2 + i*t)|^8 dt ~ (24024/16!) * c * T * log(T)^16 (Keating and Snaith, 1998; Conrey and Gonek, 2001).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Amiram Eldar_, Jul 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017790 | OEIS-A397826

a(1) = 2; for n > 1, a(n) is the smallest unused number greater than 2 placed on a hexagon spiral that is coprime with any earlier term that is adjacent to it on the spiral.

It is conjectured the sequence contains all positive numbers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jul 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017791 | OEIS-A397902

G.f. A(x) satisfies [x^(n-1)] (1 - A(x))^(n^2) / (1 - n^2*x) = 0 for n > 1.

Conjecture: for n > 2, a(n) is odd iff n is of the form 2^k, 2^k-1, 2^k-2, or 2^k-3 (k > 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paul D. Hanna_, Jul 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017792 | OEIS-A397903

a(n) is the minimum number of terms in a representation of prime(n) as a sum of distinct primes, each of which is itself recursively representable as a sum of distinct primes. The recursion starts with 2 and 3. a(n)=0 if no such representation exists.

So, based on the conjecture (see A124867) - "Every number n > 17 is the sum of 3 distinct prime numbers" - we can propose the following hypothesis: Conjecture: for all n > 9, a(n) = 2 or 3. This conjecture has been verified up to 10000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yves Debeuret_, Oct 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017793 | OEIS-A398024

Number of fixed 3-convex polyominoes with perimeter 2*n.

The conjectured result has since been proved by Thomas Prellberg, though the proof is still being examined as it's quite complex. [Anthony J. Guttmann] G.f.: x^2*(8*x^6 - 34*x^5 + 97*x^4 - 110*x^3 + 54*x^2 - 12*x + 1) / ((1-4*x)^2 * (1-2*x) * (2*x^2 - 4*x + 1)) - (1-3*x)^2 * x^4 / ((1-4*x)^(3/2) * (1-2*x) * (2*x^2 - 4*x + 1)). (Conjectured by Conway and Guttmann (2026).)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jul 17 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017794 | OEIS-A398025

Number of fixed 4-convex polyominoes with perimeter 2*n.

The conjectured result has since been proved by Thomas Prellberg, though the proof is still being examined as it's quite complex. [Anthony J. Guttmann] G.f.: x^2*(-12*x^7 + 49*x^6 - 178*x^5 + 282*x^4 - 208*x^3 + 77*x^2 - 14*x + 1) / ((1-4*x)^2 * (x^2 - 3*x + 1) * (5*x^2 - 5*x + 1)) - x^4*(2*x^2 - 4*x + 1)^2 / ((1-4*x)^(3/2) * (x^2 - 3*x + 1) * (5*x^2 - 5*x + 1)). (Conjectured by Conway and Guttmann (2026).)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jul 17 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017795 | OEIS-A398026

Number of fixed 5-convex polyominoes with perimeter 2*n.

The conjectured result has since been proved by Thomas Prellberg, though the proof is still being examined as it's quite complex. [Anthony J. Guttmann] G.f.: -x^2 * (8*x^7 - 38*x^6 + 136*x^5 - 248*x^4 + 198*x^3 - 76*x^2 + 14*x - 1) / ((1-x) * (1-3*x) * (1-4*x)^2 * (x^2 - 4*x + 1)) - (5*x^2-5*x+1)^2 * x^4 / ((1-4*x)^(3/2) * (1-x) * (1-2*x) * (1-3*x) * (x^2 - 4*x + 1)). (Conjectured by Conway and Guttmann (2026).)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jul 17 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017796 | OEIS-A398027

Number of fixed 6-convex polyominoes with perimeter 2*n.

The conjectured result has since been proved by Thomas Prellberg, though the proof is still being examined as it's quite complex. [Anthony J. Guttmann] G.f.: -x^2*(20*x^9 - 127*x^8 + 524*x^7 - 1292*x^6 + 1712*x^5 - 1267*x^4 + 544*x^3 - 135*x^2 + 18*x - 1) / ((1-4*x)^2 * (x^3 - 6*x^2 + 5*x - 1) * (7*x^3 - 14*x^2 + 7*x - 1)) - (1-2*x)^2 * (x^2 - 4*x + 1)^2 * x^4 / ((1-4*x)^(3/2) * (x^3 - 6*x^2 + 5*x - 1) * (7*x^3 - 14*x^2 + 7*x - 1)). (Conjectured by Conway and Guttmann (2026).)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jul 17 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017797 | OEIS-A398028

Number of fixed 7-convex polyominoes with perimeter 2*n.

The conjectured result has since been proved by Thomas Prellberg, though the proof is still being examined as it's quite complex. [Anthony J. Guttmann] G.f.: x^2*(24*x^10 - 198*x^9 + 867*x^8 - 2476*x^7 + 4072*x^6 - 3896*x^5 + 2251*x^4 - 798*x^3 + 170*x^2 - 20*x + 1) / ((1-2*x) * (1-4*x)^2 * (2*x^2 - 4*x + 1) * (2*x^4 - 16*x^3 + 20*x^2 - 8*x + 1)) - x^4*(7*x^3 - 14*x^2 + 7*x - 1)^2 / ((1-4*x)^(3/2) * (1-2*x) * (2*x^2 - 4*x + 1) * (2*x^4 - 16*x^3 + 20*x^2 - 8*x + 1)). (Conjectured by Conway and Guttmann (2026).)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vaclav Kotesovec_, Jul 17 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017798 | OEIS-A398064

Decimal expansion of Hadamard's Gamma function at x = 1/4.

Conjecture: H(x+1) = lim_{n->oo} n^{x+1} * Sum_{k=0..n} (-1)^k/(n+k) * binomial(x, k) such that H(1/4) = 0.79564036... . Changing the upper bound of the sum to oo yields Euler's product representation of the Gamma function.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Ulgenes_, Jul 18 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017799 | OEIS-A398139

Numbers m such that the distinct prime factors of the product m*(m+1)*(m+2) are consecutive primes beginning with 2 (that is, the product is a term of A055932).

Conjecture: There are no more terms in this sequence past 440. This conjecture is based on analysis of the growth rate of the minimum prime index values of the GPF of M_3(m) (call it iota_min(M_3(m))) in primorial intervals, such that P_r <= M_3(m) < P_(r+1). For intervals r=2 through r=22 the iota_min progression is: 2, 3, 3, 4, 4, 5, 6, 10, 8, 10, 13, 13, 16, 16, 20, 22, 26, 24, 30, 31, 39. The significance of iota_min in these intervals is that the maximum value of omega for M_3(m) in these intervals is r, so if iota_min exceeds r, it is impossible for there to be a prime-complete product in the interval. In the sequence above, iota_min exceeds r for r=14 and above. The links to the program and output data for this analysis are below.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ken Clements_, Jul 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017800 | OEIS-A398140

Numbers m such that the distinct prime factors of the product m*(m+1)*(m+2)*(m+3) are all consecutive primes beginning with 2 (that is, the product is a term of A055932).

Conjecture: There are no more terms in this sequence past 63. This conjecture is based on analysis of the growth rate of the minimum prime index values of the GPF of M_4(m) (call it iota_min(M_4(m))) in primorial intervals, such that P_r <= M_4(m) < P_(r+1). For intervals r=3 through r=27 the iota_min progression is: 3, 3, 4, 6, 7, 6, 8, 9, 11, 13, 14, 14, 21, 23, 27, 29, 33, 38, 31, 38, 62, 63, 72, 81, 73. The significance of iota_min in these intervals is that the maximum value of omega for M_4(m) in these intervals is r, so if iota_min exceeds r, it is impossible for there to be a prime-complete product in the interval. In the sequence above, iota_min exceeds r for r=15 and above. The links to the program and output data for this analysis are below.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ken Clements_, Jul 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017801 | OEIS-A398141

Numbers k such that (sin(k))^k sets a new record low.

Conjecture: every term after 0 is either the numerator of a convergent (A096456) or semiconvergent of Pi/2, or else p + 2p', where p/q and p'/q' are consecutive convergents of Pi/2. Of the 13 terms after 0, 7 are convergents, 4 are semiconvergents, and 2 are of the last form: 364863 = 260515 + 2*52174 and 5992555 = 4846147 + 2*573204.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Aug 25 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017802 | OEIS-A398149

a(1) = 1; for n > 1, a(n) is the smallest positive number on a square spiral such the resulting sums of the numbers in each row and column is distinct.

In the first ten million terms the lowest number not to appear is 153; it is conjectured all positive numbers eventually appear. In this range the largest value is a(6454127) = 248.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Jul 22 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017803 | OEIS-A398157

a(n) = n/3 if n == 0 (mod 3), 4n+1 if n == 1 (mod 3), 2n-1 if n == 2 (mod 3).

It is conjectured (not proved) that iterating this map from any positive integer eventually reaches one of exactly two cycles: (1, 3, 5, 9) and (7, 11, 17, 19, 21, 29, 33, 51, 57, 77, 153). Verified empirically for n up to 10^6 with no exceptions found.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nicholas Green_, Sep 13 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017804 | OEIS-A398183

a(n) = second-smallest prime p <= n such that 2n - p is also prime, or 0 if fewer than two such primes exist.

a(n) = 0 only for n = 3, 4, 6 (i.e., 2n = 6, 8, 12) among 2n <= 10^7; conjecturally these are the only even numbers with fewer than two Goldbach partitions.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Soroosh Hemmati_, Jul 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017805 | OEIS-A398186

a(n) = (1/n) * Sum_{k=0..n-1} A000897(k) * A000897(n-1-k), where A000897(k) = (4*k)!/((2*k)!*k!^2).

Sum_{n>=1} a(n)*x^n is the antiderivative vanishing at x = 0 of the hypergeometric square 2F1(1/4, 3/4; 1; 64*x)^2. Under the change of variable x = t/(1 + 64*t), where t = (eta(2*z)/eta(z))^24, it becomes D^(-1) C_4, where D = q*d/dq and C_4 is the weight-4 meromorphic modular form on Gamma_0(2) associated by Bönisch, Duhr and Maggio with a hypergeometric K3 family. Since x = q - 40*q^2 + ... has integer coefficients and linear coefficient 1, it has a compositional inverse over the integers. Hence the integrality of a(n) is equivalent to the denominator-one divisibility n | c_4(n). This strengthens the depth-one magneticity conjecture for C_4 and is proved in the Shvets reference below.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Shvets_, Jul 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017806 | OEIS-A398189

Table read by rows: T(n, k) = v2(A398187(n, k)) where v2(m) is the 2-adic valuation of m.

We conjecture that:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Luschny_, Jul 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017807 | OEIS-A398223

a(n) is the number of iterations for the map f(k) = k/3 if k == 0 (mod 3), 4k+1 if k == 1 (mod 3), 2k-1 if k == 2 (mod 3), to first reach a previously-visited value, starting from k=n.

This is a three-branch analog of the Collatz (3x+1) map, branching on residue mod 3 instead of mod 2. Empirically (not proved), every trajectory eventually reaches one of exactly two cycles: (1, 3, 5, 9) and (7, 11, 17, 19, 21, 29, 33, 51, 57, 77, 153). It is conjectured, but not proved, that no starting value diverges to infinity. Verified by direct computation for all n from 1 to 10^6 with no divergences and no cycles other than the two listed. Some trajectories transiently reach very large intermediate values (on the order of 10^15 or more) before collapsing into a cycle.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Nicholas Green_, Sep 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017808 | OEIS-A398259

a(1)=0; for n>=2, let s = digit sum of a(n-1); if s=a(m) for some m < n-1, let m be the greatest such index and set a(n)=n-1-m; otherwise set a(n) = 0.

Among the first 10^6 terms there are 26 zeros, see A398307. No further zero was found through n=10^6. It is conjectured that these are all the zeros. Empirically, a(n)/n appears to approach 1. In particular, a(10^4)=9365, a(10^5)=99353 and a(10^6)=999847. Conjecture: lim_{n->oo} a(n)/n=1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bence Bernáth_, Jul 24 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017809 | OEIS-A398270

Array read by antidiagonals: T(n,k) is the feedback vertex set number of the grid graph P_n X P_k.

Conjecture: T(n,k) = n*k - A360920(n,k).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Andrew Howroyd_, Jul 25 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017810 | OEIS-A398282

Let Ab_k be the set of abelian groups of order k, M_k the set of rings Z/mZ such that phi(m) = k, and let f_k : M_k -> Ab_k be the map such that f_k(Z/mZ) = (Z/mZ)*, the group of units of Z/mZ. a(n) is the least r such that f_k is injective for k = p*2^r, p the n-th prime. a(n) = -1 if no such number exists.

The map f_k can be considered as a refinement of Euler's phi function. For example, we may define another map g_k: Z -> Z where g_k(m) = phi(m) = k, i.e. the map mapping all m to their totient value. The image of g_k is obviously the singleton {k}; g_k is trivially surjective, and Carmichael's conjecture is that it is never injective. Since the image of f_k contains those unit groups (Z/mZ)* such that |(Z/mZ)*| = k, f_k exhibits richer mapping properties.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Miles Englezou_, Jul 24 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017811 | OEIS-A398300

a(n) is the minimum possible sum of the numbers on all non-mine cells when n mines are placed on an infinite Minesweeper board (each non-mine cell's number is the count of mines among its up to 8 neighbors).

This equals 8n - 2*M(n), where M(n) is the maximum number of edges in an induced subgraph of order n of the grid graph with diagonals added (vertex set Z^2, edges between points at Euclidean distance 1 or sqrt(2)). M(n) was conjectured by Taliceo and Fleron (2021) to equal 4n - ceil(sqrt(28n-12)), and this was proved by Vince (2024); see LINKS.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Dimas Saputra_, Jul 25 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017812 | OEIS-A398307

Indices of zero terms in the Van Eck-like sequence A398259 whose lookup key is the decimal digit sum of the previous term.

There are no further terms between 541 and 10^6. It is conjectured this list is complete.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bence Bernáth_, Jul 25 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017813 | OEIS-A398329

Number of partitions of the 2-dimensional vector (n, n) into strictly distinct parts with nonnegative components, excluding elementary basis vectors (1,0) and (0,1).

Conjecture: a(n) ~ A219554(n) / 4. - _Vaclav Kotesovec_, Sep 04 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jishnu Babu Ranitha_, Sep 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017814 | OEIS-A398343

a(n) = (Sum_{d|k} rad(d)) / rad(k), where k = A398342(n) and rad = A007947.

Conjecture: For each term m, m appears only a finite amount of times in the sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Žiga Pirc_, Sep 02 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017815 | OEIS-A398367

Values written by the walker in the trampled carousel process.

Empirically, the ratio L(k+1)/L(k) appears to converge to 1+x = 1.6573427020515763..., where x is the unique positive solution of x = Sum_{m>=2} (1/m) * Sum_{t=binomial(m,2)..binomial(m+1,2)-1} (1+x)^(-t); the last ratio in the computed range is 69020473/41645264 = 1.65734266..., which agrees with 1+x to 8 significant digits. The mean value written during greater-than-1 phase k appears to approach 1/x^2 = 2.3142821659504142717501025363582... (Conjectured.) The conjectured equation for x has an equivalent formulation in terms of Ramanujan's theta function. Let psi(q) = Sum_{n>=0} q^(n*(n+1)/2) = Product_{n>=1} (1-q^(2*n))/(1-q^(2*n-1)) and set q = 1/(1+x). Then the equation for x is equivalent to (1/2)*Integral_{t=0..q} (psi(t)-1)/t dt = 2-1/q. Indeed, if S(q) = Sum_{m>=2} q^binomial(m,2)/(m*(m-1)), then 2*q*S'(q) = psi(q)-1, and the conjectured equation for x is equivalent to S(q) = 2-1/q. The convergence to this solution is still conjectural. Both conjectures are true (see Epstein link). - _Andrew R. Epstein_, Sep 24 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Javier Alejandro Cabrera Otten_, Jul 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017816 | OEIS-A398383

a(n) = c_6a(n)/n^2, where c_6a(n) are the Fourier coefficients of the level-2 K3 meromorphic modular form C_6a defined below.

Bönisch, Duhr and Maggio introduced C_6a and conjectured that it is magnetic of depth 2. Shvets proved the stronger denominator-one statement a(n) in Z for every n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Shvets_, Jul 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017817 | OEIS-A398446

Number of permutations of [n] avoiding the pattern 1324 with the maximum element n in position n-2.

Conjecture: for d >= 1, A395725(n,n-d) = p_d(M)*binomial(2*M,M) + q_d(M)*4^M with M = n-1-d, where deg p_d = d-1 with leading coefficient 1/d!, q_1 = 0, and for d >= 2, deg q_d = d-2 with leading coefficient binomial(d,2)/d!, p_d(0) - q_d(0) = 1, and p_d(0) + q_d(0) = A061552(d). Verified for d = 1..9. This sequence is d = 2, where those conditions leave nothing free: p_2 = (M+3)/2 and q_2 = 1/2. The next case is d = 3, with p_3 = (M^2+11*M+21)/6 and q_3 = (M+5)/2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles Cornell Norton_, Jul 29 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017818 | OEIS-A398498

Let S(n) denote the sequence (b(i), i=0..n) sorted into increasing order, where b is A377091. Since 0 is in S(n), S(n) contains a maximal string of consecutive numbers that starts with a nonpositive term and ends with a nonnegative term. The present sequence is an irregular triangle, read by rows, in which row n is this maximal string.

It is conjectured that both A398499 and A398500 are infinite sequences. A proof that either of them is infinite would imply the long-standing conjecture that A377091 is a permutation of the integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _N. J. A. Sloane_, Jul 31 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017819 | OEIS-A398540

Decimal expansion of the asymptotic ratio r_infty = lim_{n->oo} W(n+1)/W(n), where W(n) is the winning probability in the continuous n-slot proportional placement game (derived from the social media game 'The Number Challenge').

Equals lim_{n->oo} W(n+1)/W(n) where W(n) = (1/n)*Sum_{k=1..n} W(k-1)*W(n-k)*(I(k/n;k,n-k+1) - I((k-1)/n;k,n-k+1)), W(0)=W(1)=1, I(x;a,b) = regularized incomplete Beta function (conjectured limit, supported by Richardson extrapolation).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alessandro Serra_, Aug 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017820 | OEIS-A398542

Number of 1324-avoiding L-griddings with n points.

Conjecture: for every m >= 1 and every b in S(m), d(b,k) = p(k)*binomial(2*k,k) + q(k)*4^k for polynomials p and q depending on b, with deg p <= m-1 and deg q <= m-2. Verified for m <= 6 and k <= 12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Charles Cornell Norton_, Aug 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017821 | OEIS-A398551

Even numbers that are not sums of two isolated primes.

628 is the conjectured to be the last term.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Liu Junhui_, Aug 02 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017822 | OEIS-A398563

Semiprimes of the form p^3 - q^3 where p and q are primes.

A disproof of the twin conjecture would imply that this sequence is finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Michael Shmoish_, Sep 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017823 | OEIS-A398564

Triangle read by rows: T(n,k) is the greatest common divisor of the numbers of ways to form j simultaneous cross-team handshakes between teams of n and k distinct people, with each person participating in at most one handshake, for 1 <= j <= k <= n.

Open question: Can M(n,k) be determined explicitly, and for which pairs (n,k) does M(n,k) > T(n,k)?

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ahaan Kallat_, Oct 15 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017824 | OEIS-A398581

Numbers k for which the lexicographically first integer solution (x,y,z) of 5/k = 1/x + 1/y + 1/z with 0 < x < y < z differs from a solution having maximal z.

Conjecture: every term is congruent to 1 modulo 5. Verified for all 86 terms with k <= 1500. The converse is false: many k == 1 (mod 5) are not terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Batista_, Aug 03 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017825 | OEIS-A398583

Earliest sequence of numbers k such that k * sin(k) is positive and decreases monotonically toward 0.

This sequence is infinite iff the sequence k * sin(k) accumulates on the positive side of 0, which is an open question. If k * sin(k) is dense on the real line, or if the irrationality measure of Pi exceeds 2, then the sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Matt Frank_, Aug 03 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017826 | OEIS-A398589

Irregular triangle read by rows, where row n is the lexicographically earliest sequence starting with T(n,1) = n, allowed integers are >= n, and each term T(n,k) is banned for the next T(n,k) terms in that row. If the sequence for row n is eventually periodic, the row terminates at the end of the first period; however, if it is not eventually periodic, the row is infinite.

Conjecture: There are no values of n > 2 that have rows without a pre-periodic block. Conjecture: All of the rows in this sequence are eventually periodic. It has been shown by Martin Fuller that for n up to 100 this conjecture holds true (see Weinstein link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Joshua B. Weinstein_, Sep 08 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017827 | OEIS-A398603

a(n) is the least number which is the end of an arithmetic progression of n numbers that are the sums of two squares (A001481).

Conjecture: a(n) = A093365(n) for n >= 13.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Aug 04 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017828 | OEIS-A398604

Gaps associated with the arithmetic progressions in A398603.

Conjecture: a(n) = A093366(n) for n >= 12.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hugo Pfoertner_, Aug 04 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017829 | OEIS-A398690

Triangle of polynomials related to Verlinde numbers.

Conjecture: The polynomials are palindromic.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _F. Chapoton_, Aug 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017830 | OEIS-A398693

Cardinality of the unique largest subdirectly irreducible factor of the lattice of partitions of n under the dominance order.

The set of partitions of n can be given a complete lattice structure as first shown by Brylawski. This lattice has an irredundant decomposition as a subdirect product of subdirectly irreducible factor lattices; for n >= 13, there is a unique largest factor lattice in this decomposition, the size of which is a(n). For n < 13 the subdirectly irreducible factors of highest cardinality are not always unique and have a different structure not fitting the size formula conjectured to hold below. Conjecture: a(n) = A000041(n) - 4*n + 7 - (-1)^n (verified for n <= 60).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Mike Behrisch_, Edith Vargas-García, and Andreas Wachtel, Aug 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017831 | OEIS-A398720

Triangle read by rows: T(n,k) in which the n-th row lists the weight distribution of square Single Parity Check Product (SPCP) [(n+1)^2, n^2, 4] code, 0 <= k <= n*floor(n/2).

Conjecture: T(n, n*(n-1)/2) = n! for n odd. The conjecture holds: the complement of such a matrix has n ones and odd row and column sums, so it is a permutation matrix. - _Andrew R. Epstein_, Oct 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _G. Gordon Thompson_, Aug 07 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017832 | OEIS-A398786

a(n) is the number of times n appears in A091508.

Conjecture: a(n) >= Product_{k=2..n} (A000005(k) - 1).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sam Chapman_, Sep 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017833 | OEIS-A398812

a(n) = n^2 - A138808(n): number of lattice points in the square [1,n] X [1,n] not covered by any box [1,p] X [1,q] with p*q = n.

Conjecturally a(n)/n^2 -> 1; i.e., A138808(n) = o(n^2). If n = p^k for p prime, then a(n) = n*(n+k(1/p-1)-1). The above conjecture is true, since tau(n) <= 2*sqrt(n) and A138808(n) <= n*tau(n). - _Chai Wah Wu_, Aug 20 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Soslan Daurov_, Aug 10 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017834 | OEIS-A398901

a(n) = c_6b(n)/n^2, where c_6b(n) are the Fourier coefficients of the level-2 K3 meromorphic modular form C_6b defined below.

Bönisch, Duhr and Maggio introduced C_6b and conjectured that it is magnetic of depth 2. Shvets proved the stronger denominator-one statement a(n) in Z for every n >= 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Alex Shvets_, Aug 13 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017835 | OEIS-A398916

a(n) is the Grundy value of "Bit Deletion" game at state n (see comments).

Conjecture: There are no terms exceeding 3. The first occurrences for the integers not exceeding 3 are: a(0) = 0; a(1) = 1; a(2) = 2; a(5) = 3. All n not exceeding 5*10^6 have been checked without finding a term for which a(n) > 3. The conjecture above is true; see the Theorem in the Formula section. - _Maxim Mikhisor_, Sep 07 2026 a(1) = 1. Inductive hypothesis: suppose that we have a(2^k) = (k mod 2) + 1. The binary expansion of 2^(k+1) is 1 followed by k+1 zeros. Thus, the only states reachable from a(2^(k+1)) are a(0) (by removing the leading 1) and a(2^k) (by removing any of the zeros). By the Sprague-Grundy theorem, we have a(2^(k+1)) = mex{a(0), a(2^k)} = mex{0, (k mod 2) + 1} = ((k+1) mod 2) + 1. Conjecture: a(n) = a(4n) for all n. If this conjecture is true, then the sequence a(n*2^k) for a fixed n and increasing k has a period dividing 2: a(n), a(2n), a(n), a(2n), ... All n not exceeding 10^6 have been checked without any

counterexamples to the conjecture found. Consequently a(n) <= 3 for all n, and a(n) is unchanged when any zero-run is altered by an even amount; in particular a(4n) = a(n), which proves both conjectures above. Also a(n) is odd iff n has an odd number of binary digits, and a(n) >= 2 iff m is odd. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Do Thanh Nhan_, Aug 14 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017836 | OEIS-A398920

Number of solutions of the Diophantine inequality a*b+b*c+c*d <= n, with a,b,c >= 1.

Conjecture: a(n) = n^2*log(n)^2/4 + O(n^2*log(n)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Bevan_, Aug 14 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017837 | OEIS-A399084

a(0) = 0; a(1) = 1; thereafter a(n) = a(n-1) - 1 if that is not already in the sequence, otherwise a(n) = a(n-1) + floor(sqrt(a(n-1))).

Conjecture: block lengths are 1,1,1,1,1 followed by m,1,m,1 for m = 2,3,4,..., i.e., 1,1,1,1,1, 2,1, 2,1, 3,1, 3,1, 4,1, 4,1, ... This holds for the first 25292 blocks, that is, for all terms below 4*10^7. The block tops fall into two interleaved orbits of the map x -> x + 1 + floor(sqrt(x+1)), which is where the alternation of long and unit blocks comes from. A proof along these lines should be routine.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Vasilios Mavroudis_, Aug 18 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017838 | OEIS-A399145

Number of distinct proper side-crossing points among regular polygons inscribed in a common circle and sharing a common vertex.

Conjecture: No three or more sides belonging to distinct polygons P_3, P_4, ..., P_n meet at the same proper crossing point. Hence the number of distinct proper crossing points equals the number of pairwise proper side-crossing events. Verified numerically for n up to 1000. Conjecture: a(n) = 2*Sum_{3 <= m < k <= n} (m - GCD(m,k)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eddie Lin Rui_, Aug 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017839 | OEIS-A399147

Number of distinct proper side-crossing points among regular polygons inscribed in a common circle and aligned by their side midpoints.

Conjecture: No three or more sides belonging to distinct polygons P_3, P_4, ..., P_n meet at the same proper crossing point. Hence the number of distinct proper crossing points equals the number of pairwise proper side-crossing events. Verified numerically for n up to 1000. Conjecture: a(n) = 2*Sum_{3 <= m < k <= n} (m - [val_2(m) = val_2(k)]*GCD(m,k)).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eddie Lin Rui_, Aug 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017840 | OEIS-A399155

a(n) = f(n) - g(n), where f(n) is the number of steps to reach 0 by repeatedly subtracting the smallest prime factor of the current value, and g(n) is the number of steps to reach 0 by repeatedly subtracting the largest prime factor.

Conjecture: a(n) >= 0 for all n >= 2; equivalently, subtracting the largest prime factor never takes more steps than subtracting the smallest.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jaylen Dilkhush_, Aug 20 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017841 | OEIS-A399208

Minimum over all symmetric n X n {-1,1} matrices M with zero diagonal of the maximum of |Sum_{1 <= i < j <= n} M[i,j]*x[i]*x[j]| over all vectors x in {-1,1}^n.

n*sqrt(n-1)/Pi <= a(n), and limsup a(n)/n^(3/2) <= 1/2. Whether a(n)/n^(3/2) converges is an open question of Paata Ivanisvili (see the MathOverflow link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rob Sneiderman_, Aug 22 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017842 | OEIS-A399306

Integers whose minimal Lucas representation (A130310) does not end in a 0 which if removed results in a valid minimal Lucas representation of another integer.

The similarity of this sequence with A094350 is striking, but the terms 41 and 117 (=76+41) occur in A094350 but not here. It is possible that the terms of this sequence comprise a subset of the terms of A094350, but this is merely a conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Caveney_, Aug 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017843 | OEIS-A399307

The periodic cycle of 23 in the "Lucas-Collatz" function: a(0)=23; for n>0, if a(n-1) is Lucas-odd (A399306), then a(n)=2*a(n-1); if a(n-1) is Lucas-even (A399305), then a(n) is the integer whose minimal Lucas representation (A130310) results from removing the final 0 of A130310(a(n-1)).

Conjecture: The orbit of every positive integer that is not a multiple of 5 reaches either this cycle of 23 or the cycle 2, 4, 3, 1, 2. - _Geoffrey Caveney_, Sep 03 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Caveney_, Aug 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017844 | OEIS-A399308

The periodic cycle of 70 in the "Lucas-Collatz" function: a(0)=70; for n>0, if a(n-1) is Lucas-odd (A399306), then a(n)=2*a(n-1); if a(n-1) is Lucas-even (A399305), then a(n) is the integer whose minimal Lucas representation (A130310) results from removing the final 0 of A130310(a(n-1)).

Conjecture: The orbit of every positive multiple of 5 reaches either this cycle of 70 or the cycle 5, 10, 5.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Geoffrey Caveney_, Sep 03 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017845 | OEIS-A399309

Deterministic version of the random Fibonacci sequence, with signs from A191232 and the seed as (0,1).

Conjecture: lim_{n->oo} |a(n)|^(1/n) = 1.1319882487943... (A078416), Viswanath's constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Oct 13 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017846 | OEIS-A399375

Odd numbers k for which A348717(A163511(k)) <= A348717(k).

It is conjectured that the equivalence is attained only with Mersenne primes (A000668). This is equivalent to the conjecture given in A364297.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017847 | OEIS-A399381

a(n) is the minimum positive integer k such that k*n-1 is squarefree.

I conjecture that the averages of the first n terms of this sequence and the first n terms of A399382 tend to the same constant as n grows arbitrarily large. (End)

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aidan Markey_, Aug 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017848 | OEIS-A399385

Number of 2n+1 X 2n+1 off-diagonally symmetric alternating sign matrices.

a(n) = 2^(n-1) * (3*n+2)! / (2*n+1)! * Product_{i=1..n} (6*i-2)! / (2*n+2*i+1)!. This formula was conjectured by Behrend, Fischer, and Koutschan and proved by Kumari.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Christoph Koutschan_, Aug 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017849 | OEIS-A399456

Decimal expansion of 2*(Pi - sqrt(3))/(sqrt(15) + sqrt(7) - 2*sqrt(3)).

Resnikoff (2015) proves that this is the density of a certain packing of Reuleaux triangles, and conjectures this to be the maximum packing density for any curve of constant width.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Aug 28 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017850 | OEIS-A399572

Triangular numbers in the Padovan sequence A000931.

Conjecture: the sequence is complete. No further terms: none of A000931(k) for 32 <= k <= 10^12 is triangular (8*A000931(k)+1 is a quadratic non-residue modulo at least one of the 40 smallest primes > 2^30, or, for k = 392532961617 and 972000569010, modulo 1000003 and 1000033 respectively). Any further term must have index k in one of 2365 residue classes modulo 55194480 (congruence obstructions from the primes 5, 7, 11, 19, 23, 89, 109, 271, 607, 691, 809, 829, 991, 1009, 3037, 3313, 3637). The corresponding indices k with A000931(k) triangular are 0, 1, 2, 3, 4, 5, 6, 7, 10, 17, 18, 27, 28, 31.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Ali Ashter_, Oct 01 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017851 | OEIS-A399627

a(n) is the number of residues modulo 2^n counted by A396966 that are not lifts of a residue counted by A396966 modulo 2^(n-1).

The new residues generate the whole family by lifting, and the asymptotic density of A396966 is exactly the generating series c_W = Sum_{n>=1} a(n)/2^n. The exact value of c_W is an open problem, governed by the growth of this sequence. c_W = Sum_{n>=1} a(n)/2^n is the asymptotic density of A396966. The ratios A396966(n)/2^n are non-decreasing lower bounds for c_W; at n = 40, c_W >= 172651809822/2^40 = 0.157025906... The exact value of c_W is an open problem.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jakob Stemberger_, Sep 05 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017852 | OEIS-A399633

The city-building sequence: a(n) = the height h of the shortest building which can see exactly h earlier buildings.

Conjecture: limsup_{n->oo} (Sum_{k=1..n} a(k))/n <= 3. See A397740. - _Paolo Xausa_, Sep 28 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric Fox_, Sep 05 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017853 | OEIS-A399659

Number of involutions of length n avoiding the patterns 4321 and 4231.

Equivalently, a(n) is the number of Motzkin paths of length n such that, for every pair of consecutive level steps, the number of down steps between them is at least the height of the first of the two (Barnabei, Castronuovo, Silimbani). The generating function satisfies a functional equation with one catalytic variable and can be expressed as an explicit continued fraction (see Formula). It is conjectured not to be D-finite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Niccolò Castronuovo_, Oct 05 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017854 | OEIS-A399681

a(1) = 1, a(2) = 2; for n > 2, a(n) is the smallest unused positive number that shares a factor with a(n-1) if a(n-1) <= n-1, else is coprime to a(n-1) if a(n-1) > n-1.

For the terms studied the fixed points are 1, 2, 33, 40, 42, 112, 663, 671, 962. It is plausible no more exist. The sequence is conjectured to be a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Scott R. Shannon_, Oct 10 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017855 | OEIS-A399714

a(n) = -1 - A010815(n-1) - Sum_{k=1..n-2} A010815(k)*2^(n-2-k) for n >= 3.

Conjecture: For n >= 3, a(n) is the number of linearly independent relations of weight n arising from the Ihara-Kaneko-Zagier derivation relation. See the Szűcs (2026) for details.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tamás Szűcs_, Sep 09 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017856 | OEIS-A399743

a(n) is the minimum number of entries from row n of Pascal's triangle needed to represent Fibonacci(n) as a signed sum, with each position used at most once.

Conjecture: a(n) <= ceiling(n/3) + 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Felix Huber_, Sep 16 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017857 | OEIS-A399755

Numbers k such that the generalized Pell equation x^2-(k^2+1)*(y^2+1) = 0 has more than one fundamental solution in (x, y).

Conjecture: A399491(k) < 4*k^3+3*k if and only if k is a term of this sequence.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Chai Wah Wu_, Sep 10 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017858 | OEIS-A399793

a(n) is the largest size of a subset S of {1, ..., n} such that there does not exist a function f: S -> {-1,1} satisfying |Sum_{x in S} x*f(x)| <= 1.

Conjecture: The equality is achieved when n >= 13.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Yifan Xie_, Sep 12 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017859 | OEIS-A399861

Deterministic version of the random Fibonacci sequence, with signs from A191232 and the seed as (1,0).

Conjecture: lim_{n->oo} |a(n)|^(1/n) = 1.1319882487943... (A078416), Viswanath's constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Oct 15 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017860 | OEIS-A399880

On Artin's conjecture, (a(n)/A399881(n))*C is the relative density of primes with primitive root n among the primes, where C = A005596 is Artin's constant.

On Artin's conjecture, (a(n)/A399881(n))*C is the relative density of primes with primitive root n among the primes, where C = A005596 is Artin's constant. For |n|>1, write n = (b^2*c)^k where b^2*c is not a prime power and c is squarefree. Artin's conjecture gives:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Sep 14 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017861 | OEIS-A399881

On Artin's conjecture, (A399880(n)/a(n))*C is the relative density of primes with primitive root n among the primes, where C = A005596 is Artin's constant.

On Artin's conjecture, (A399880(n)/a(n))*C is the relative density of primes with primitive root n among the primes, where C = A005596 is Artin's constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Sep 14 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017862 | OEIS-A399882

On Artin's conjecture, (a(n)/A399883(n))*C is the relative density of primes with primitive root -n among the primes, where C = A005596 is Artin's constant.

On Artin's conjecture, (a(n)/A399883(n))*C is the relative density of primes with primitive root -n among the primes, where C = A005596 is Artin's constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Sep 14 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017863 | OEIS-A399883

On Artin's conjecture, (A399882(n)/a(n))*C is the relative density of primes with primitive root -n among the primes, where C = A005596 is Artin's constant.

On Artin's conjecture, (A399882(n)/a(n))*C is the relative density of primes with primitive root -n among the primes, where C = A005596 is Artin's constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Sep 14 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017864 | OEIS-A399898

Partial sums missed by the grasshopper infinite barrycade (recursively defined family of permutations of the positive integers having pairwise disjoint sets of partial sums).

It is conjectured that this sequence is infinite (Dębski et al.).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bartlomiej Pawlik_, Sep 16 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017865 | OEIS-A399903

Partial sums missed by the greedy grasshopper infinite barrycade (recursively defined family of permutations of the positive integers having pairwise disjoint sets of partial sums).

It is conjectured in Dębski et al. that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bartlomiej Pawlik_, Sep 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017866 | OEIS-A399907

Smallest positive integers omitted from the rows of the greedy infinite quasi-barrycade (recursively defined family of quasi-permutations having pairwise disjoint sets of partial sums).

It is conjectured in Dębski et al. that every row except the first omits exactly one positive integer. Conjecturally, a(n) = A399908(n)-1 for n >= 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bartlomiej Pawlik_, Sep 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017867 | OEIS-A399908

First elements of the rows of the greedy infinite quasi-barrycade (recursively defined family of quasi-permutations having pairwise disjoint sets of partial sums).

It is conjectured in Dębski et al. that a(n)=A399907(n)+1 for n >= 4. Conjecturally, a(n)=A399907(n)+1 for n >= 4.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Bartlomiej Pawlik_, Sep 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017868 | OEIS-A399918

Decimal expansion of sqrt(6)/(1 + sqrt(2)).

It is conjectured that this constant is the Nieuwland number of the Tetrahedron (see Steininger and Yurkevich).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Sep 15 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017869 | OEIS-A399919

Decimal expansion of the least positive root of the equation 2025*x^8 - 11970*x^6 + 17009*x^4 - 9000x^2 + 2000.

It is conjectured that this constant is the Nieuwland number of both the dodecahedron and the icosahedron (see Steininger and Yurkevich).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Stefano Spezia_, Sep 15 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017870 | OEIS-A399926

Irregular triangle read by rows: T(n,k) = Product_{primes p divides n} (p - [m == 1 (mod p)]), where m = A038566(n,k) is the k-th number coprime to n, and [] is an Iverson bracket.

Let R(n,m) = Product_{primes p divides n} (p - [m == 1 (mod p)]) (then T(n,k) = R(n,A038566(n,k))). As described in A306198, we have the following conjecture: Let c != 1 be a squarefree number, b be any positive integer such that b^2*c != -1 is not a perfect power, and n be a positive integer divisible by c if c == 1 (mod 4) or by 4*c otherwise. Then for gcd(m,n) = 1, the relative density among the primes of primes congruent to m modulo n that have b^2*c as a primitive root is (2*R(n,m)/A306198(n))*C if kronecker(c/m) = -1 and 0 if kronecker(c/m) = +1, where C is Artin's constant = A005596.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jianing Song_, Sep 16 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017871 | OEIS-A399934

Six times the maximum area of the convex hull of a polyhex with n cells, where each hexagonal cell has area 1.

S. Kurz conjectured in 2008 that the maximum area is at most floor(n^2 + 14*n/3 + 1)/6. The formula below proves this conjecture and shows that the bound is attained exactly when n is not divisible by 3 (see the Gaur link).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Pragyaan Gaur_, Sep 16 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017872 | OEIS-A399952

Deterministic version of the random Fibonacci sequence, with signs from A030190 and the seed as (0,1).

Conjecture: lim_{n->oo} |a(n)|^(1/n) = 1.1319882487943... (A078416), Viswanath's constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Sep 16 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017873 | OEIS-A399970

Largest elements of integral Diophantine quadruples, among which one plus the product of any two distinct terms is a square.

Arkin, Hoggatt, and Straus found that when (a, b, c) is an integral Diophantine triple, then, putting r^2 = ab+1, s^2 = bc+1, t^2 = ac+1, and d+ = a + b + c + 2abc + 2rst, (a, b, c, d+) is an integral Diophantine quadruple and d+ > max(a, b, c); i.e., d+ belongs to this sequence. Such a quadruple is called regular. It is conjectured that any Diophantine quadruple must be regular.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Tomohiro Yamada_, Sep 17 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017874 | OEIS-A400127

Least m such that every binary word of length n occurs as a contiguous subword of the binary representation of 3^k for some 0 <= k <= m.

Conjecture: a(n) is finite for every n. This conjecture follows from the density of the fractional parts of k*log_2(3). Choose a binary de Bruijn sequence of order n, linearized and rotated to begin with 1. Since log_2(3) is irrational, infinitely many powers 3^k have this finite binary string as their initial digits. Such a power alone therefore contains every n-bit binary word, proving a(n) < infinity for every n. - _David Rabahy_, Oct 04 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _David Rabahy_, Sep 19 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017875 | OEIS-A400171

Numbers k not divisible by any p^p and for which A003415(k) is a multiple of A276085(k), where A003415 is the arithmetic derivative and A276085 is the primorial base log-function.

Conjecture: For all terms k > 1, A008966(k) = A063524(A003415(k)/A276085(k)), i.e., the quotient is 1 if and only if k is squarefree.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Sep 23 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017876 | OEIS-A400216

Scaled x-coordinates of the vertices of a Boerdijk-Coxeter helix starting at a tetrahedron with vertices (-1,-1,-1), (1, -1, 1), (-1, 1, 1), and (1, 1, -1).

G.f.: (54*x^6 - 84*x^5 - 66*x^4 + 23*x^3 + 9*x^2 + x - 1)/((3*x - 1)^2 * (9*x^2 + 4*x + 1)) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Sep 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017877 | OEIS-A400217

Scaled y-coordinates of the vertices of a Boerdijk-Coxeter helix starting at a tetrahedron with vertices (-1,-1,-1), (1, -1, 1), (-1, 1, 1), and (1, 1, -1).

G.f.: (-54*x^6 + 84*x^5 - 90*x^4 + 15*x^3 + 3*x^2 + 3*x - 1)/((3*x - 1)^2*(9*x^2 + 4*x + 1)) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Sep 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017878 | OEIS-A400218

Scaled z-coordinates of the vertices of a Boerdijk-Coxeter helix starting at a tetrahedron with vertices (-1,-1,-1), (1, -1, 1), (-1, 1, 1), and (1, 1, -1).

G.f.: (18*x^5 + 26*x^4 - 24*x^3 - 5*x^2 + 1)/(27*x^3 + 3*x^2 - x - 1) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Peter Kagey_, Sep 21 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017879 | OEIS-A400277

a(n) = phi(n)^2 mod (n+1), where phi is Euler's totient function (A000010).

Conjecture: no algebraic family exists for the composite n with a(n) = 4 (checked up to 10^8: 4, 6, 75, 132, 195, 272, 1275, 1332, 4246, 5187, 10947, 14755, 23493, 40795, 73776, 88101, 327675, 4144209, 5883744, 7478470, 33412795, 92644745).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sajid Khan Hussain_, Sep 22 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017880 | OEIS-A400359

Decimal expansion of 9 - 5*sqrt(3) - 7*sqrt(6)/2 + 6*sqrt(2).

This value was conjectured by Melissen (1993) and later proved by Joós (2021). See also Tedeschi (2021).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_, Sep 30 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017881 | OEIS-A400370

Deterministic version of the random Fibonacci sequence, with signs from A030190 and the seed as (1,0).

Conjecture: lim_{n->oo} |a(n)|^(1/n) = 1.1319882487943... (A078416), Viswanath's constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Sep 24 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017882 | OEIS-A400405

Deterministic version of the random Fibonacci sequence, with signs from A076478 and the seed as (0,1).

Conjecture: lim_{n->oo} |a(n)|^(1/n) = 1.1319882487943... (A078416), Viswanath's constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Sep 25 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017883 | OEIS-A400412

Triangle read by rows: T(n, k) = v is the smallest number which has n odd divisors and whose symmetric representation of sigma, SRS(v), consists of k parts, or 0 if no such number exists, 1 <= k <= n.

Conjectures:

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Hartmut F. W. Hoft_, Sep 25 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017884 | OEIS-A400454

Numbers k such that (2^k-1)*2^(k+3)+1 is a prime.

Conjecture: If 2^((N-1)/4) == -1 (mod N), then N is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary Wright_, Sep 26 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017885 | OEIS-A400469

Deterministic version of the random Fibonacci sequence, with signs from A076478 and the seed as (1,0).

Conjecture: lim_{n->oo} |a(n)|^(1/n) = 1.1319882487943... (A078416), Viswanath's constant.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jwalin Bhatt_, Sep 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017886 | OEIS-A400486

Number of positive integers m < 2^n - 1 such that the Collatz trajectories of m and m+1 meet within n steps, after equal numbers of halving steps and before either reaches 1.

Gao tabulated a(n)/2^n to three decimals for 4 <= n <= 24 (Table 3, from 0.063 to 0.389), proved that a(n)/2^n is nondecreasing, that is, a(n+1) >= 2*a(n) (his Corollary 2, from his Lemma 2), and conjectured that a(n)/2^n -> 1. Lagarias's annotated bibliography records the conjecture in another form: the set of m such that f^k(m) = f^k(m+1) for some k <= log_2(m) has natural density 1. A meeting within log_2(m) steps is always a coalescence in Gao's sense, so this form implies a(n)/2^n -> 1.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar Said_, Sep 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017887 | OEIS-A400487

Number of positive integers k < 2^n that take the same number of Collatz steps to reach 1 as k-1 or k+1.

Wu (1992) computed a(n) for n <= 24, observed that a(n)/(2^n-1) increases with n, and conjectured that the proportion of integers lying in such runs tends to 1. If a(n)/2^n -> 1, then the positive integers that reach 1 have natural density 1, because every k counted reaches 1. Whether they do is an open problem; the best known lower bound for the number of k <= x that reach 1 is x^0.84 (Krasikov and Lagarias).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar Said_, Sep 27 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017888 | OEIS-A400582

Convolution of the number of dissections of a convex (n+2)-gon into triangles and quadrilaterals by nonintersecting diagonals (A001002).

Conjecturally, a(n) is the number of rc-simple permutations of length 2n which are invariant under the reverse-complement map and which avoid 2413 (or equivalently, 3142). Here we say a permutation of length 2n which is invariant under the reverse-complement map is rc-simple whenever it contains no nontrivial intervals among its first n entries.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Eric S. Egge_, Sep 30 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017889 | OEIS-A400584

Irregular triangle read by rows: T(n,k) is the number of labelings of the internal vertices of the complete plane binary tree of depth n by + or min, with every leaf equal to 1, that evaluate to k at the root (n >= 1, 1 <= k <= 2^n).

A labeling has root value 1 exactly when it contains a root-to-leaf path whose internal vertices are all min. Choosing the leftmost such path gives a product decomposition of this class. This decomposition proves the product identity conjectured by Peter Bala in A076628 (Feb 11 2015); see Mendonça (2026), technical note linked below. Auffinger and Cable report unpublished work of Albert Chen proving that the root-value probabilities decrease with k, independently discovered by Tam Cheetham-West. Here monotonicity must be understood in the weak sense, since the first two rows contain ties. Conjecture: T(n,k) > T(n,k+1) for n >= 3 and 1 <= k < 2^n. This strict-monotonicity conjecture has been verified by exact computation for 3 <= n <= 12, but is not proved in general.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _J. R. G. Mendonça_, Sep 30 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017890 | OEIS-A400585

a(n) = Sum_{d|n, n/d odd} binomial(d, 2).

Conjecture: Sum_{k=1..n} a(k) ~ 7*zeta(3)*n^3/48. - _Vaclav Kotesovec_, Oct 01 2026

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Jishnu Babu Ranitha_, Sep 30 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017891 | OEIS-A400649

Indices q at which A327008 decreases: A327008(q+1) = A327008(q) - 1.

It is conjectured that this sequence is infinite.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Max Alekseyev_, Oct 02 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017892 | OEIS-A400701

Start with a set of five adjacent collinear points in a hexagonal lattice. Every iteration, add to the set the points in the lattice which are adjacent to at least two points already in the set. This sequence gives the number of points in the set after n iterations.

Conjectured: For n > 1: a(n) = A374338(n) + 4*n + 7.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Aidan Markey_, Oct 03 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017893 | OEIS-A400792

a(1) = 1. For n > 1, a(n) is the smallest unused positive integer k such that (number of earlier terms smaller than k that are coprime to k) - (number of earlier terms larger than k that share a factor with k) = 1.

Conjecture: this is a permutation of the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Paolo Xausa_ and _Michael De Vlieger_, Oct 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017894 | OEIS-A400795

Strongly nontriangular numbers, defined as the complement of the generalized triangular numbers in the reference below. Generalized triangular numbers include 1 and those in the range of the function P(i,j) = (i*j+i+j-gcd(i,j))/2 for integers i, j > 1.

It is conjectured that this sequence is infinite and has natural density zero. The sequence arises in an attempt to study "generalized triangular numbers" of the form P(i,j) = (i*j+i+j-gcd(i,j))/2. This formula counts the number of lattice points in the triangle with vertices (0,0), (i,0), (0,j), excluding points on the hypotenuse.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Kyle Petersen_, Oct 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017895 | OEIS-A400802

Number of orbits of exponents d, 1 <= d <= 2^n-2, under multiplication by 2 modulo 2^n-1, such that the power map x -> x^d on GF(2^n) is almost perfect nonlinear (APN).

Conjecture: for even n, a(n) = phi(n) - 1 + [5|n]; for odd n >= 11, a(n) = 2*phi(n) + 3 + 2*[5|n], where [5|n] = 1 if 5 divides n and 0 otherwise. These formulas agree with all terms for n >= 10.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Sajid Khan Hussain_, Oct 06 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017896 | OEIS-A400825

Numbers k such that (2^k-1)*2^(k+4)+1 is prime.

Let N = N(k) = (2^k-1)*2^(k+4)+1. Conjecture: If 2^((N-1)/4) == -1 (mod N), then N is prime.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Gary Wright_, Oct 07 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017897 | OEIS-A400868

Number of deals of n cards numbered 1 to n between two players for which the card game War never ends, when the winner of each turn puts the losing card and then the winning card at the bottom of their pile.

Conjecture: a(n) = 0 exactly when n is a power of 2.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Daniel Okwor_, Oct 07 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017898 | OEIS-A400887

a(n+1) = a(n-a(a(n))^2) + 1, a(0) = 0.

a(n) ~ 2^(1/4)*n^(1/2) (conjectured).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Rok Cestnik_, Oct 08 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017899 | OEIS-A400888

Contents of the quarter-turn corrected Stern vectors associated with the column words of the Gosper-Lafitte triangle.

Every positive integer divides some term of this unbounded sequence. Every positive even integer occurs infinitely often as a term. It is conjectured that the sequence is surjective onto the positive integers.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _M. Jeremie Lafitte (Levitas)_, Oct 08 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017900 | OEIS-A400983

Number of steps after which the Collatz trajectories of A078417(n) and A078417(n)+1 first coincide.

In the form recorded by Lagarias, Gao's conjecture (see A400486) states that the numbers m with f^k(m) = f^k(m+1) for some k <= log_2(m) have natural density 1. Among the m that reach 1, these are exactly the terms m = A078417(n) with a(n) <= log_2(m).

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar Said_, Oct 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017901 | OEIS-A400984

Numbers k > 1 whose trajectory under the Collatz map A014682 contains k+1.

Rozier and Terracol show that every solution with k > 3 is a paradoxical sequence in their sense: 3^q < 2^j, where q is the number of odd terms among k, T(k), ..., T^(j-1)(k), so k is in A393502. By their computations there is no further term below 2.8*10^19, and their Conjecture 6.1 (no paradoxical sequence starts above 4614) would imply that the sequence is finite with these 21 terms.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar Said_, Oct 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017902 | OEIS-A400985

Length of the longest run of consecutive positive integers < 2^n that take the same number of Collatz steps to reach 1.

Wu (1992) studied the runs of consecutive integers with the same number of steps in [1, 2^N), counting the integers that lie in runs of length at least 2 (A400487). He found that the longest run below 2^30 has 176 integers, starting at 722067240, and conjectured that the length of the longest run in [1, M) tends to infinity with M. Since a(n) is the largest k with A078441(k) + k <= 2^n, Wu's conjecture is the statement that A078441(k) exists for every k, that is, Garner's conjecture that there are arbitrarily long runs of consecutive integers with the same height.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Omar Said_, Oct 11 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017903 | OEIS-A401001

Numbers k such that the sum of digits of k in base b is not prime for any base b in the range 2 <= b <= floor(sqrt(k)).

Conjectured to be finite, with 1650 being the last term; no further terms exist up to k = 50000.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Guido Avagliano_, Oct 12 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017904 | OEIS-A401145

Number of partitions of n into distinct parts such that n plus the product of the parts is prime.

Conjecture: for n >= 7, a(n) > max(a(n-1), a(n+1)). Conjecture: a(n) > 0 for all n except for n = 2. This holds for n <= 170.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Akash Suresh_, Oct 16 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

V3-017905 | OEIS-A401236

a(n) = A276086(A035345(n)), where A276086 is the primorial base exp-function, and A035345 is the smallest prime > 1 + (prime(1)*prime(2)*...*prime(n)).

These terms are by definition all in A373846, while those of A401235 are there just by conjecture.

猜想/问题提及段 | historical_or_current_mention_unreviewed | _Antti Karttunen_, Oct 18 2026 | 2026-10-24

关键词命中原段；序列定义见题名及原页面 | 原始来源 | CC BY-SA 4.0

C 论坛与文献定位：2,455 条

论坛页是人类讨论的证据，不是开放状态认证。定位记录缺少正文，应打开原文核对；这里不把模型抽取或模型概述当成作者原句。

C01 MathOverflow | 603 条

V3-017906 | MO-10223

Linear equation with primes

Is there an integer n with an infinite number of representations of the form $n=2q-p$, where p and q are both primes? Given a positive integer $k>1$, I would like to know for which (if any) integers n the linear equation $q-kp=n$ admits an infinite number of solutions, where p and q are primes. (I'm not including $k=1$ because it reduces to well know open problems, $k=1$ and $n=2$ would be the twin primes conjecture) The density of the prime numbers implies that at least there are integers n with an arbitrarily large number of representations.

论坛问题帖 | has_answers | Manuel Silva | 2009-12-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-017907 | MO-102331

Sparse ramsey theory

It is known that for any graph H and all $k∈N$, there exists a graph G such that any k-coloring of the edges of G yields a monochromatic copy of H and ω(G)=ω(H) (the two graphs have the same clique numbers). My question is: Given any graph H with finite girth, is there a G with the same girth as H such that any 2-coloring of the edges of G yields a monochromatic copy of H? I think this is an open problem but if someone can confirm that and give some references concerning this I would be most obliged.

论坛问题帖 | has_answers | user14875 | 2012-07-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017908 | MO-102644

Actions on $\mathbb{H}^n$ generated by torsion elements

Let n be a large integer. I am looking for a cocompact properly discontinuous isometric action on n-dimensional Lobachevky space which is generated by elements of finite order. Or equivalently, I need a cocompact properly discontinuous isometric action $\Gamma\curvearrowright\mathbb{H}^n$ such that $\mathbb{H}^n/\Gamma$ is simply connected, here $\mathbb{H}^n/\Gamma$ stays for quotient space (NOT for orbifold). Comments. I am aware that for large n these is no cocompact action generated by reflections (Vinberg, 1984). Consider the group of matrices with integer coefficients from $\mathbb{Q}[\sqrt{5}]$ which preserve the quadratic form $$\tfrac{1+\sqrt{5}}{2}\cdot x_0^2-x_1^2-\cdots-x_n^2.$$ This gives cocompact properly discontinuous isometric action, say $\Gamma\curvearrowright\mathbb{H}^n$. (For $n=2$, the group Γ contains the Coxeter group of regular right-angled pentagon.) I would be very happy if the subgroup generated by the elements of finite order in Γ would have finite index. Or, equivalently, if $$|\pi_1(\mathbb{H}^n/\Gamma)|<\infty.$$ P.S. It seems that examples of such actions are not known. (In addition to Agol's comment, I've got a letter from Vinberg stating this.)

论坛问题帖 | no_answers_at_retrieval | Anton Petrunin | 2012-07-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017909 | MO-10356

Topological "Interpolation" ?

Let E be a normed space, and let T:E * $\rightarrow$ E * be a nonlinear operator. Suppose that : 1) T is continuous from (E *, ||.||) to itself (i.e., it is norm-continuous). and 2) T is continuous from (E *, w *) to itself (i.e., it is weakly-star continuous). Then does it follow that 3) T is continuous from (E *, w) to itself (i.e., it is weakly continuous) ? I guess the case E = $\ell\infty$ would be particularly interesting.

论坛问题帖 | has_answers | Ady | 2010-01-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-017910 | MO-10358

Solving NP problems in (usually) Polynomial time?

Just because a problem is NP-complete doesn't mean it can't be usually solved quickly. The best example of this is probably the traveling salesman problem, for which extraordinarily large instances have been optimally solved using advanced heuristics, for instance sophisticated variations of branch-and-bound. The size of problems that can be solved exactly by these heuristics is mind-blowing, in comparison to the size one would naively predict from the fact that the problem is NP. For instance, a tour of all 25000 cities in Sweden has been solved, as has a VLSI of 85900 points (see here for info on both). Now I have a few questions: 1) There special cases of reasonably small size where these heuristics either cannot find the optimal tour at all, or where they are extremely slow to do so? 2) In the average case (of uniformly distributed points, let's say), is it known whether the time to find the optimal tour using these heuristics is asymptotically exponential n, despite success in solving surprisingly large cases? Or is it asymptotically polynomial, or is such an analysis too difficult to perform? 3) Is it correct to say that the existence of an average-case polynomial, worst-case exponential time algorithm to solve NP problems has no importance for P=NP? 4) What can be said about the structure of problems that allow suprisingly large cases to be solved exactly through heuristic methods versus ones that don't?

论坛问题帖 | has_answers | DoubleJay | 2010-01-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-017911 | MO-10449

Artin-Schreier Theorem for Rings

This has been in my mind for quite some time. Looking at the Artin-Schreier Theorem for fields: If L is a field and K its algebraic closure and if $1< [K:L] < \infty$ then $K=L[i]$ and L is a real closed field. (Thus L has characteristic 0. Here i is just the square root of -1). I was wondering if a "generalized" Artin-Schreier exists or if someone could refer to me to some paper that attempts this. There is a concept of real closedness and "algebraic closedness" of reduced commutative rings, but I doubt that the statement would hold. So one

has the following conjecture: If L is a reduced commutative ring and K is its total integral closure (this is an equivalent notion of algebraic closure if K and L were fields) and if $1<[K:L]< \infty$ (here I mean that K is a finite L-module that is not the same as L) then L is real (thus its characteristic is 0, and one can add that L is real closed in the sense of reduced commutative rings). Can one easily show this, even at least prove that L has characteristic 0?

论坛问题帖 | no_answers_at_retrieval | Jose Capco | 2010-01-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-017912 | MO-104573

Open problems in sub-Riemannian geometry

What are some open problems in sub-Riemannian geometry? I am interested especially in problems concerning connections and curvature, but any contribution is welcomed.

论坛问题帖 | has_answers | Cristi Stoica | 2012-08-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017913 | MO-104598

A question concerns prime numbers

I have arrived to this conjecture in my work, I am not sure that is true or false. So I would appreciate if someone give a counterexample or prove it. My question: Let n be a non-prime such that n-1 be a prime. Is it true that there exist a prime p such that n-p is a prime as well?

论坛问题帖 | has_answers | Tina | 2012-08-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017914 | MO-104812

Sums of Strongly z-ideals

In the rings of continuous functions,i.e.$(C(X))$ an ideal I is called strongly z-ideal if it is an intersection of some maximal ideals of $C(X)$. i.e. $$I=\cap_{\alpha \in A} \mathcal{M_{\alpha}}$$ Where each $\mathcal{M_{\alpha}}$ is a maximal ideal of $C(X)$. Question: If I and J are two strongly z-ideals of $C(X)$, Is the ideal $I+J$ strongly z-ideal or all of $C(X)$? PS: If X is a P- space, then each ideal of $C(X)$ is strongly z- ideal, And conversly if each ideal of $C(X)$ is strongly z- ideal than X is a P-space.

论坛问题帖 | no_answers_at_retrieval | Ali Reza | 2012-08-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017915 | MO-105065

Minimizing the excursion of a sum of unit vectors

I have n unit-length vectors v_i in $\mathbb{R}^3$, whose sum is zero: $$ v_1 + v_2 + \cdots + v_n = 0 \; .$$ Now I form the closed polygon P in space by placing them head to tail. So the vertices of P are $$ 0, v_1, (v_1+v_2), \ldots, (v_1+\cdots +v_{n-1}), 0 \; .$$ My question is: What is the minimum excursion from the origin achievable by shuffling the vectors by a permutation of $(1,2,\ldots,n)$? For example, the 12 red vectors below wander $\sqrt{10}$ from the (purple) origin, but the light blue vectors-the same in a different order-stay within $\sqrt{3}$. (These particular vectors derive from the vertices of a cuboctahedron, so some are negations of others.) Is there some constant $r_{\min}$ independent of n such that the sum can always be arranged to be at most $r_{\min}$, i.e., lie within an origin-centered ball of that radius? Or must $r_{\min}$ depend on n? Is there some natural algorithm for minimizing the excursion, or must I (in the worst case) try all $n!$ permutations? Of course the same question can be asked in any dimension $\mathbb{R}^d$, but my focus is $\mathbb{R}^3$. Thanks for ideas and/or pointers! Update1. The suggestion (in the comments) that $r_{\min} = \sqrt{d}$ in $\mathbb{R}^d$, based on an answer to the previous MO question, "Bounding a signed sum of complex numbers," is intriguing, and may be true. But I do not see that it is proved by that answer. Update2. With key phrases suggested by Nik Weaver, I found a 1981 paper by Imre Bárány, "A Vector-Sum Theorem and its Application to Improving Flow Shop Guarantees" (Math. Oper. Res. link), which shows that $r_{\min} < \frac{3}{2} d$.

论坛问题帖 | has_answers | Joseph O'Rourke | 2012-08-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-017916 | MO-1051

Splitting Pythagorean triples

Can one partition the set of positive integers into finitely many Pythagorean-triple-free subsets? If so, what is the smallest number of such subsets? Taking a wild guess, I would be least surprised if the answer were 3. Notice that the 2 subsets of integers such that highest power of 5 that divides them is a) even b) odd manage to split most primitive triples, plus all the multiples of those. Notice also that Schur proved the positive integers cannot be split into any finite number of sum-free subsets (i.e. no finite partition can split all power-of-1 Fermat triples), while Fermat's theorem proves that all power-of-n (n>2) triples can be split by the trivial partition into 1 set. Edit: Since this turns out to be a known open problem, we're adding the tag [open-problem] and converting this question to community wiki. The idea is to have a separate answer for each possible approach to solving this problem. If you have some additional insight or a reference to contribute to an answer, you only need 100 rep to do so. We're still figuring out exactly how to handle open problems on MO. The discussion is happening on this tea.mathoverflow.net thread.

论坛问题帖 | has_answers | Yaakov Baruch | 2009-10-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-017917 | MO-105722

S-matrix conjecture: status?

Is the S-matrix conjecture still open? I mean the one listed as Problem 7 in this survey.

论坛问题帖 | has_answers | Felix Goldberg | 2012-08-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017918 | MO-106552

The Class Number One Problem for Real Quadratic Fields

An approach to the Gauß class number one problem for imaginary quadratic fields is to determine the integral points on the modular curve $Y_{nonsplit}(n)$ for a suitable n. Here follows a quick summary, but see Appendix A in Serre's book "Lectures on the Mordell-Weil Theorem" for more details: Given an imaginary quadratic K of class number 1, one considers an elliptic curve E with CM by O_K, which is unique up to $\mathbb{C}$-isomorphism. Its j-invariant lies in $\mathbb{Z}$. Given any integer n, all of whose prime divisors are inert in K, our E yields a unique integral point on $Y_{nonsplit}(n)$. That means, fixing an n and letting p be its largest prime divisor, any imaginary quadratic K of class number 1 whose discriminant is larger than $4p$ in absolute value will furnish a unique integral point on $Y_{nonsplit}(n)$. Hence, determining the integral points on $Y_{nonsplit}(n)$ for one n for which there are finitely many integral points solves the problem. Heegner used $n=24$, as did Stark. My question is whether a similar approach has been, or can be, used in the real quadratic case; the goal being to prove that there are infinitely many such fields of class number one. For what it's worth, here's what would happen in my pipe dream: To a real quadratic K of class number 1 one attaches uniquely an abelian surface (or some other kind of object) with real multiplication by O_K; One shows that this object gives rise to a unique rational point on some moduli space, and conversely, all such rational points would come from a K; ($Y_{nonsplit}$ plays this role in the imaginary case...) One shows that the moduli space has infinitely many rational points. (Since this is a well-known open problem, I'll follow the advice of the FAQ and make it community-wiki. I am a little embarrassed asking such a speculative question, but I also feel that speculation drives a lot of mathematical research.)

论坛问题帖 | has_answers | Giuseppe | 2012-09-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017919 | MO-106660

Kronecker's Jugendtraum for real quadratic fields?

Kronecker's Jugendtraum (or Hilbert's 12'th problem) is to find abelian extensions of arbitrary number fields by adjoining `special' values of transcendental functions. The Kronecker-Weber theorem was the first realisation of this: i.e. $\mathbb{Q}^{ab}=\mathbb{Q}^{cycl}=\mathbb{Q}(e^{2\pi i \mathbb{Q}})$. If K is an imaginary quadratic field, then the theory of complex multiplication realises Kronecker's dream: roughly $K^{ab}=K(j(\tau), \wp(\tau,z))$, where τ is a `special' value of the upper half plane corresponding to an elliptic curve $E\cong\mathbb{C} / \Lambda$ with complex multiplication by the ring of integers of K, and $z \in \Lambda \otimes \mathbb{Q}$ (this is like adjoining $j(E)$ and the x-coordinates of the torsion points of E). Is there some class of special analytic functions, and some special kind of objects which conjecturally play the role of the j-function, the $\wp$-function and CM elliptic curves for real quadratic fields?

论坛问题帖 | has_answers | Adam Harris | 2012-09-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017920 | MO-106694

Almost Complex Structures: 'Tame' versus 'Compatible'

Consider a symplectic manifold (M,ω) and the space of almost complex structures. These are $J:TM\to TM$ with $J^2=-\text{id}$. A given J is ω-tame when $\omega(v,Jv)>0$, and J is ω-compatible when it is ω-tame and $\omega(J\cdot,J\cdot)=\omega(\cdot,\cdot)$. The set of either such J forms a contractible subspace. Note that in either scenario we can form a Riemannian metric g_J by twisting ω in some way with J. In practice, I sometimes see J being tame, and more often I see J being compatible, but in either case I am unsure of the restrictions that each imposes. Is there any context in which it is necessary/helpful to use one condition over the other? This need not be related to pseudo-holomorphic curves in Floer theory. Do certain results fail when I relax "compatible" to "tame"? A fundamental difference I see (quoted from McDuff-Salamon's big textbook) is that compatible J minimize the energy of a J-holomorphic curve in its homology class, but not necessarily for tame J.

论坛问题帖 | has_answers | Chris Gerig | 2012-09-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017921 | MO-106815

Measures of full Hausdorff dimension for self-affine sets

Consider the iterated function system $T_{1}(x)=(\beta x,\tau y)$, $T_{2}(x,y)=(\beta x+(1-\beta),\tau y+ (1-\tau))$ for $\beta\in(1/2,1)$ and $\tau\in (0,1/2)$ with self affine set $\Lambda_{\beta,\tau}$. It is known that for almost all β and all τ $\dim_{H}\Lambda_{\beta,\tau}=\dim b=1-\log(2\beta)/\log(\tau)$ where $b=(1/2,1/2)$ is the Bernoulli measure on Λ. Essential this is due to the fact that the projection of b to the x-axis is an infinite Bernoulli convolution which is know to be generic absolutely continues (Solomyak theorem). On the other hand we have shown that if β^{-1} is a Pisot number we have $\dim b<\dim_{H}\Lambda_{\beta,\tau}<1-\log(2\beta)/\log(\tau)$ for all Bernoulli measures b. The projection of all b in this case is singular with a dimension drop (Erdös theorem). Now here comes the question: Is there a (ergodic) measure of full dimension and what is $\dim_{H}\Lambda_{\beta,\tau}$ in the later case?

论坛问题帖 | has_answers | Jörg Neunhäuserer | 2012-09-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017922 | MO-107534

Is $Q_n(x)=\sigma_{n+1}(x)/\sigma_n(x)$ logarithmically convex on $\mathbf{R}$?

In 1975 J. van de Lune considered the monotony properties of the canonical Riemann Upper and Lower sums for $\int_0^1 t^xdt$, with $x>0$. Writing $\sigma_n(x) := 1^x+2^x+\cdots+n^x$ these sums are $$U_n:=U_n(x)=\sigma_n(x)/n^{x+1} \quad \text{and}\quad L_n:= L_n(x)=\sigma_{n-1}(x)/n^{x+1}.$$ He proved (by mathematical induction) that $U_n>U_{n+1}$ and $L_n<L_{n+1}$. A later proof of $U_n>U_{n+1}$ was obtained by showing that the function $$h(x):= h_n(x)=(\sigma_{n+1}/\sigma_n(x))(n/(n+1))^x$$ is strictly decreasing on all of $\mathbf{R}$. Soon afterwards he came to realize that the monotonicity of $h(x)$ would be

a consequence of the logarithmic convexity of $Q(x):=Q_n(x)=\sigma_{n+1}(x)/\sigma_n(x)$ on all of $\mathbf{R}$. Various numerical tests on $Q(x)$ were performed, but no proof was found. Another application: From the logarithmic convexity of $Q(x)$ one may also obtain a simple proof of a conjecture made by H. Alzer and A. A. Jagers: $$f(x):=f_n(x)=\Bigl(\frac{\frac{1}{n+1}\sigma_{n+1}(x)}{\frac1n \sigma_n(x)}\Bigr)^{1/x}$$ is strictly increasing for $x>0$. Question: Is for every (fixed) integer $n\ge2$, the function $$x\in\mathbf{R}\mapsto Q_n(x):=(1^x+2^x+\cdots+n^x+(n+1)^x)/(1^x+2^x+\cdots+n^x)$$ logarithmically convex on all of $\mathbf{R}$?

论坛问题帖 | has_answers | juan | 2012-09-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017923 | MO-107573

The oriented homeomorphism problem for Haken 3-manifolds

Haken famously described an algorithm to solve the homeomorphism problem for the 3-manifolds that bear his name (fleshed out by many others, including Hemion and Matveev who fixed some gaps). But it's often natural to consider 3-manifolds equipped with an orientation, and so my question is: Question 1: Is there an algorithm in the literature that determines whether or not a pair of oriented Haken 3-manifolds M and N are orientedly homeomorphic? By Haken's Algorithm, we can reduce to the case in which M and N are (unorientedly) homeomorphic, and so the question can be reduced to: Question 2: Is there an algorithm in the literature that determines whether or not an orientable Haken 3-manifold M admits an orientation-reversing self-homeomorphism? Notes: Note the words 'in the literature'. I don't really doubt that this algorithm exists, but the details seem quite complicated, particularly in the non-geometric case, so I hope that someone will be able to supply a reference. A very short argument would also be nice. Answers of the form 'I don't know, but I'm sure it can be done if you work out each geometric case individually and then think hard enough about the JSJ decomposition', while appreciated, would be less useful. The hyperbolic case can be deduced from Sela's solution to the homeomorphism problem, which also computes the automorphism group. I've no doubt the other non-Haken cases can be handled similarly. Hence my interest in the Haken case. It may be that a simple modification of Haken's Algorithm handles the oriented case. This isn't at all clear to me, but I would be very interested to hear if it's true. (added later) I would also be happy to confirm that this question is `open' (in the sense that an answer is not in the literature, rather than that the answer is in doubt)---I very much suspect that this is the case. An authoritative pronouncement by someone who knows the literature very well would therefore be a valid answer.

论坛问题帖 | has_answers | HJRW | 2012-09-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017924 | MO-108000

Are Galois groups of Q with restricted ramification supposed to be finitely generated?

Fix a finite set S of places of $\mathbb Q$. Let $G_{\mathbb Q,S}$ be the Galois group of the maximal extension of $\mathbb Q$ unramified outside S$. I believe that it is an open question whether this group is topologically finitely generated, that is, contains a dense, finitely generated subgroup. Is there a standard conjecture about whether $G_{\mathbb Q,S}$ should be finitely generated or not? Has anyone published an opinion? JSE mentions a stronger conjecture: $|S|$ generators suffice; further, the existence of a special set of generators labeled by S and inert at the labeling place. Is this refinement standard? (Note that a finite extension would suffice to disprove this hypothesis.) Weakening the hypothesis, we could restrict the ramification at S to be tame.

论坛问题帖 | has_answers | Ben Wieland | 2012-09-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017925 | MO-108406

Why would dim primitive irrep divide size of some conjugacy class ?

From Isaacs et.al. 2005 Conjecture C. Let χ be a primitive irreducible character of an arbitrary finite group G. Then χ(1) divides | clG(g)| for some element g ∈ G. Here, of course, we have written clG(g) to denote the class of g in G. We have checked that Conjecture C holds for all irreducible characters (primitive or not) of all groups in the Atlas 1. Question 1 What is motivation for this ? Is it possible to describe what conjugacy class(es) should correspond to irreducible representation in this way ? (at least for some standard groups S_n, A_n, GL_n(F_q),...) What are representative examples? Question 2 Is it still open ? The authors write: We now digress to explain our original motivation for considering these questions. There are numerous parallels and analogies between theorems concerning the of set irreducible character degrees of a finite group and theorems concerning the set of conjugacy class sizes of such groups. This suggests that perhaps there are some subtle arithmetic connections between these two sets of integers associated with a given group. One such connection that is easy to see is that each prime number that divides an irreducible character degree of G must also divide some class size of G. If G is solvable, then S. Dolfi showed that more is true. He proved [2] that given any two distinct primes p and q such that pq divides some irreducible character degree of a solvable group G, then pq also divides some class size of G. One might conjecture that the analogous assertion for three or more distinct primes is also true, but as far as we know, this remains open. Partial result: In the following, we use the notation np to denote the p-part of a positive integer n, where p is a prime number. Corollary D. Let χ be a primitive irreducible character of a solvable group G, and let p be a prime divisor of |G|. Then χ(1)p divides (| clG(g)|p) 3 for some element g ∈ G. Not related results, for compltеness: Denote CV(g) fixed point subspace for g in V. Our main result is the following. Theorem A. Let V be a nonzero finite dimensional completely reducible F G-module, where F is any field and G is any finite group. Assume that CV (G) = 0 and let p be the smallest prime divisor of |G|. Then there exists some element g ∈ G such that $ dim CV (g) ≤ (1/p) ~ dim V $. The fraction 1/p cannot, in general, be replaced by any smaller quantity. In particular, this shows that Neumann’s conjecture is valid for odd-order groups, at least... Corollary B. Let V be a nonzero finite dimensional completely reducible F G-module, where F

is an arbitrary field and G is any finite group, and assume that CV (G) = 0. Then $1/ |G| \sum_{g∈G} dim CV (g) ≤ ((p + 1)/ 2p)~~ dim V$, where p is the smallest prime divisor of |G|.

论坛问题帖 | has_answers | Alexander Chervov | 2012-09-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017926 | MO-108530

A maximal element, where Schur gives a minimal element

Let me recall a result due to I. Schur, which I learnt from F. Goldberg's answer to my MO question Hadamard-like inequalites for positive definite symmetric matrices. If H is a subgroup of $\frak S_n$ and χ is an irreducible complex character over H, define $$d_\chi(S)=\frac1{\chi(e)}\sum_{g\in H}\chi(g)\prod_{i=1}^ns_{ig(i)}.$$ Then for every $S\in SPD_n$, we have $$\det(S)\le d_\chi(S).$$ Notice that if $H=\frak S_n$ and χ is the signature, then d_χ is the determinant. Thus $\det$ is the smallest element among the d_χ's. If instead $\chi={\bf1}$, then d_χ is the permanent. If $H=(e)$, Schur's inequality is just the Hadamard inequality $$\det S\le\prod_is_{ii}.$$ Given n, there are many distinct d_χ's, even though several choices of the pair (H,χ) yield the same function. For instance, there are only 11 distinct functions if $n=3$, among 13 pairs. My question is whether the permanent is the largest element among the d_χ's. In other words, is it true that for every $S\in SPD_n$, we have $$d_\chi(S)\le{\rm per}(S)\quad ?$$ I checked the truth of this assertion if $n=2$, $n=3$, and also in quite a complicated case of $n=4$, where $H={\frak A}_4$ and $\chi\ne{\bf1}$ is a linear character.

论坛问题帖 | has_answers | Denis Serre | 2012-10-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017927 | MO-108661

approximate uncertainty principle for finite abelian groups

Edit: we cannot find such an example. It would imply a negative solution to the KS${}_2$ conjecture which has now been proven by Marcus, Spielman, and Srivastava in this paper. In fact, their solution implies that there always exists S such that any f supported on S (or S^c) satisfies $\|\frac{1}{2}\hat{f} - \hat{f}|_T\|_2 \leq O(\sqrt{\epsilon})\cdot\|\hat{f}\|_2$. Can we find, for every $\epsilon > 0$, an example of the following? $\bullet$ a finite abelian group G $\bullet$ a small subset T of the dual group $\hat{G}$ (meaning $|T| \leq \epsilon |\hat{G}|$) such that for any subset S of G, there is a nonzero complex valued function f supported either on S or S^c whose Fourier transform is approximately supported on T (meaning $\|\hat{f} - \hat{f}|_T\|_2 \leq \epsilon \|\hat{f}\|_2$). It seems unlikely, but I don't know any version of the uncertainty principle that would forbid this. Two comments: (1) This arose out of discussions I had with Chuck Akemann about the Kadison-Singer problem. It wouldn't, as it stands, imply a negative solution to Kadison-Singer, but it would be close and (I believe) could probably be easily converted into a full negative solution. (2) Since either S or S^c has cardinality at least $|G|/2$, one could consider assuming $|S| \geq |G|/2$ and demanding that f be supported on S. But I know that this makes the problem too hard. (Given T, I can always find an S with $|S^c|/|G| = O(\sqrt{\epsilon})$ such that no nonzero function supported on S has Fourier transform approximately supported on T.)

论坛问题帖 | has_answers | Nik Weaver | 2012-10-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017928 | MO-108673

Open problems about CMC hypersurfaces with symmetries?

Recently, Andrews and Li announced a complete classification of CMC ($H=const.$) tori in S^3, confirming a conjecture of Pinkall and Sterling. Their main result is that any such torus is rotationally symmetric, and this reduces the problem to a previous classification due to Perdomo, also studied by Hynd, Park and McCuan. [As a side note, some of the techniques used are similar to Brendle's solution of the Lawson Conjecture, that states that the only minimal ($H=0$) torus in S^3, up to congruences, is the Clifford torus $S^1(\tfrac{1}{\sqrt2})\times S^1(\tfrac{1}{\sqrt2})$.] It is also known that there are CMC tori in S^3 that are not congruent to the family $S^1(r)\times S^1(\sqrt{1-r^2})$ of Clifford tori, so, in a certain sense, their only (continuous) symmetries are encoded in an isometric circle action, as confirmed by Andrews' and Li's result. Given this context, my question is: Are there open questions, conjectures, etc., regarding existence or non-existence of CMC hypersurfaces with symmetries in highly symmetric compact manifolds, analogously to the above case of CMC tori in S^3? Actually, I'd be particularly interested in questions regarding non-minimal CMC embeddings ($H=const\neq 0$). To be (slightly) more precise, I am interested in knowing if there are questions of the form "Let (M,g) be a compact manifold with many symmetries (i.e., a "big" Lie group acts isometrically), and consider embeddings of some manifold N into M. Then if $N\subset M$ is CMC, it must have certain symmetries (and perhaps cannot have other symmetries)." In other words, are there any analogues of (or questions similar to) the conjecture of Pinkall and Sterling that CMC tori in S^3 are rotationally symmetric, but for other (perhaps higher dimensional) compact manifolds? Perhaps some of these questions are disguised as questions about Mean Curvature Flow? Notice also that if $N\subset M$ is an orbit of an isometric group action, then it is automatically CMC. However, there could be other CMC embeddings of N into M, not congruent to any orbits (e.g., with less symmetries, as the tori above). Should one expect any general behavior for this other CMC embeddings? Are there any known results, questions or conjectures in this direction? I apologize for the somewhat vague question, but none of the above references seems to risk any further conjectures or indicates natural extensions of their results. Also, a quick google search and inverse citations on MathSciNet don't give much in terms of open questions. I was wondering if there is a reason behind it... Any references to recent survey-type papers related would also be very appreciated!

论坛问题帖 | has_answers | Renato G. Bettiol | 2012-10-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017929 | MO-109045

If all real conjugacy classes are strongly real, then all real irreps are "strongly real"(symmetric), true?

Question Is true that if all real conjugacy classes of a finite group are strongly real, then all its real irreducible representations (irreps) are "strongly real" (symmetric)? And vice versa? Definitions: An irrep is called real if its Frobenius-Schur indicator is either $+1$ or -1, but not 0. An irrep is called "symmetric" ("orthogonal" ("strongly real")) if it has FS-indicator $+1$, which is equivalent to it being possible to realize it over the real numbers. A conjugacy class is called "real" if it is preserved by $g\mapsto g^{-1}$. It is called "strongly real" if it is a product of two involutions. Context It is known that the number of real conjugacy classes is the same as the number of real irreps. A. Knutson asked: Are there “real” vs. “quaternionic” conjugacy classes in finite groups?, i.e., is it possible to split "real" conjugacy classes into two subclasses corresponding to the splitting of irreps. Tim Dokchitser answered that no such splitting may exist in general which will be automorphism covariant .... The question above is weakening A. Knutson's question. It has been asked explicitly in the paper Kulshrestha and Singh - Real elements and Schur indices of a group (MSN, arXiv), where the authors present some positive evidence for the question above. However, their examples do not cover GAP's SmallGroups List. SubQuestion How difficult is it to check this in GAP? (I just installed it, but never played.)

论坛问题帖 | has_answers | Alexander Chervov | 2012-10-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-017930 | MO-11091

A generalization of Gromov's theorem on polynomial growth

I was sure it is known, but it appears to be an open problem (see the answer of Terry Tao). Assume for a group G there is a polynomial P such that given $n\in\mathbb N$ there is set of generators $S=S^{-1}$ such that $$|S^n|\leqslant P(n)\cdot|S|\ \ (\text{or stronger condition}\ |S^k|\leqslant P(k)\cdot|S|\ \text{for all}\ k\le n) .$$ Then G is virtually nilpotent (or equivalently it has polynomial growth). Comments: Note that typically, $|S|\to \infty$ as $n \to\infty$ (otherwise it follows easily from the original Gromov's theorem). If it is known, then it would give a group-theoretical proof that manifolds with almost non-negative Ricci curvature have virtually nilpotent fundamental group (see Kapovitch--Wilking, "Structure of fundamental groups..."). This proof would use only one result in diff-geometry: Bishop--Gromov inequality. Edit: The actual answer is "Done now: arxiv.org/abs/1110.5008" --- it is a comment to the accepted answer. (Published reference: E. Breuillard, B. Green, T. Tao. The structure of approximate groups. Publ. Math. Inst. Hautes Études Sci. 116 (2012), 115-221. Link under Springer paywall)

论坛问题帖 | has_answers | Anton Petrunin | 2010-01-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-017931 | MO-112417

Why Donaldson's Four-Six Conjecture?

Simon Donaldson apparently made the following conjecture: Two closed symplectic 4-manifolds (X_1,ω_1) and (X_2,ω_2) are diffeomorphic if and only if $(X_1\times S^2,\omega_1\oplus\omega)$ is deformation-equivalent to $(X_2\times S^2,\omega_2\oplus\omega)$. Here ω is a symplectic structure on S^2, and a deformation-equivalence is a diffeomorphism $\phi:X_1\times S^2\to X_2\times S^2$ such that $\omega_1\oplus\omega$ and $\phi^*(\omega_2\oplus\omega)$ can be joined by a path of symplectic forms. However, where I read this did not contain any background or the original source. Where did Donaldson make this claim? And why did he make this claim? What is the motivation / are there good examples where this holds? Ivan Smith showed (through examples) that this conjecture fails when we replace S^2 by $\mathbb{T}^2$, so the statement itself seems pretty rigid. [Edit] Motivation and examples come from the 1994 paper "Symplectic Topology on Algebraic 3-Folds" of Ruan and the 1997 followup "Higher Genus Symplectic Invariants..." of Ruan-Tian, which states and proves the conjecture for simply-connected elliptic surfaces! And now we're starting to get counterexamples!

论坛问题帖 | has_answers | Chris Gerig | 2012-11-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-017932 | MO-11316

Smooth proper schemes over Z with points everywhere locally

This is a variation on Poonen's question, taking Buzzard's fabulous example into account. It was earlier a part of this other question. Question. Is there a smooth proper scheme $X\to\operatorname{Spec}(\mathbb{Z})$ such that $X(\mathbb{Q}_v)\neq\emptyset$ for every place v of $\ \mathbb{Q}$ (including $v=\infty$), and yet $X(\mathbb{Q})=\emptyset$? I believe the answer is No. The evidence is flimsy : X cannot be a curve or a torsor under an abelian variety (Abrashkin-Fontaine) or a twisted form of $\mathbb{P}_n$. I haven't gone through a list of smooth projective $\mathbb{Q}$-varieties which contradict the Hasse principle to check if any of them has good reduction everywhere, but the chances of such a thing are slim. Colliot-Thélène and Xu give a systematic treatement of many known examples of quasi-projective $\mathbb{Z}$-schemes Y such that $Y(\mathbb{Z}_p)\neq\emptyset$ for every prime p and $Y(\mathbb{R})\neq\emptyset$, but $Y(\mathbb{Z})=\emptyset$. Some of these schemes might even be smooth over $\mathbb{Z}$, but none of them is proper. Fontaine's letter to Messing (MR1274493) might be of some relevance here.

论坛问题帖 | no_answers_at_retrieval | Chandan Singh Dalawat | 2010-01-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017933 | MO-11349

Are most cubic plane curves over the rationals elliptic?

%This is a new version of the original question modified in the light of the answers and comments. The word 'most' in the title is ambiguous. The following is one way of making it precise. Question1: (This seems to be open. See Poonen's answer below) A cubic projective

curve over $\mathbb{Q}$ is given by ten relatively prime integers (the coefficients of its equation after clearing the denominators). Suppose we take a ten dimensional box $[-N,N]^{10}$ and choose points with integer coordinates with respect to the uniform measure and form the equation of the associated cubic curve. Suppose the number of points which give rise to a curve with a rational point is $E(N)$. Then what can we say about $E(N)/(2N+1)^{10}$ as $N\rightarrow \infty$? Should the limit exist and if it does, should it be one, zero, or some other number? Another question of interest is: Question 2: (There is a satisfactory answer to this. See Voloch's response below.) Are either of the sets {cubics with no rational point} and {cubics with at least one rational point} Zariski dense?

论坛问题帖 | has_answers | Idoneal | 2010-01-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-017934 | MO-113959

A function whose fixed points are the primes

If $a(n) = (\text{largest proper divisor of } n)$, let $f:\mathbb{N} \setminus \{ 0,1\} \to \mathbb{N}$ be defined by $f(n) = n+a(n)-1$. For instance, $f(100)=100+50-1=149$. Clearly the fixed points of f are the primes. Is every number preperiodic? In other words, is $f(f(\ldots(f(n)\ldots))$ eventually prime?

论坛问题帖 | has_answers | Rodrigo A. Pérez | 2012-11-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017935 | MO-114646

Intersecting family of triangulations

Let $\cal T_n$ be the family of all triangulations on an n-gon using $(n-3)$ non-intersecting diagonals. The number of triangulations in $\cal T_n$ is C_{n-2} the $(n-2)$th Catalan number. Let $\cal S \subset \cal T_n$ be a subfamily of triangulations with the property that every two triangulations of $\cal S$ have a common diagonal. Problem: Show that $|\cal S| \le |\cal T_{n-1}|$. Remark: This problem was raised independently around the same time by Karen Meagher and by me. Updates (and counter-updates) A few weeks after this problem was posted Gjergji Zaimi (private communication) proposed a more general conjecture: Conjecture: Let P be a polytope with no triangular face. Then the maximum number of vertices such that every two vertices belongs to a common facet is attained by all vertices of a single facet. The original question is the case of the associahedron. The case of the permutahedron is (update: not) known and very interesting. (It is similar in spirit but yet different from a result by Frankl and Deza- and it is related to extremal combinatorics on permutations.) For the cube the result is immediate but can serve as a good starting point for extremal combinatorics (Problem 1 here). Update: Bruno le Floch showed that the more general conjecture is false: He described a quadrangulation of S^2 with 15 vertices and 13 quadrangles having 5 vertices each two on a face. Update Paco Santos proposed the following intermediete conjecture: Conjecture: For every flag simplicial polytope, the maximum size of a set of pairwise intersecting facets is achieved by the facets containing some common vertex. A flag simplicial polytope is a simplicial polytops so that every set of vertices that any two form an edge is the set of vertices of a face. Santos's conjecture thus asserts that the conjecture about polytopes with no triangular faces still applies for dual-to-flag polytopes. Partially motivated by the MO question Jorge Olarte, Francisco Santos, Jonathan Spreer, Christian Stump wrote a paper The EKR property for flag pure simplicial complexes without boundary where they extend the conjecture to flag pure simplicial complexes without boundary and give some evidence for it.

论坛问题帖 | has_answers | Gil Kalai | 2012-11-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-017936 | MO-114758

Transcendentality of all irrationals in the Cantor set

Hi, I am a student researcher trying to prove that all irrationals within the Cantor set are transcendental. This is grounded, intuitively, in Cantor set members' being non-normal; since algebraic numbers are widely believed to be normal, this implies the transcendentality of the irrationals in the Cantor set. Now, I am at a dead end in trying to prove this, and I would appreciate any pointers (I will give you full credit in whatever end product comes out of this). I tried recasting the problem in the following way: if I can prove that any irrational algebraic number must have a 1 in its ternary expansion, then the result I'm after follows. With this, I realized that if an irrational algebraic number has at least one 1 in its ternary expansion, then it must have infinitely many of them (because irrational algebraics are a field, and otherwise I could add an appropriate irrational algebraic number and get another irrational algebraic without any 1's in their ternary expansions -- a contradiction). But I can't really get anywhere past this (tried playing with the field properties, too). Mostly, I just lack/do not know of the tools used in this kind of problem. Any help very much appreciated!

论坛问题帖 | has_answers | CantorSet | 2012-11-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017937 | MO-115001

Open problems in Birational Geometry, after BCHM

Rencently a breakthrough was made in the context of the Minimal Model Program by the work of Birkar-Cascini-Hacon-McKernan. They proved that the canonical ring of a smooth or mildly singular projective algebraic variety is finitely generated. Since I'm a master student and so I have no a wide view of the subject (I'm not an expert), I would like to know what are the main open problems in this direction (I mean, in the framework of the Mori Program). More generally, right now what are the driving forces, the big open questions in birational geometry? Feel free to close this question, if too generic for the purposes of the site. Thanks in advance.

论坛问题帖 | has_answers | Emanuele Ventura | 2012-11-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017938 | MO-11675

Borsuk pairs of Banach spaces

Given X, Y two real Banach spaces, let's say that $(X,\ Y)$ is a Borsuk pair if for any continuous mapping T : {x $\in$ X ; $||x||\leq1$} $\rightarrow$ Y s.t. T is odd on {x $\in$ X ; $||x||=1$}, it follows that the set T^{-1}({ 0 }) is nonempty. My conjecture is : $(X,\ Y)$ Borsuk pair $\Longleftrightarrow$ dim Y < ∞ and dim Y $\leq$ dim X. [Here "dim" stands for the algebraic dimension; it may be a natural number or ∞.] In other words, there is no Borsuk pair with X, Y both infinite-dimensional. Any thoughts ? [There exist several infinite-dimensional versions of the Classical Borsuk Theorem, e.g., when X is reflexive, Y is its dual, and T is a demicontinuous mapping of monotone type.] [I think the "big" problem (if it truly exists, of course ;-)) would be when both X, Y are infinite dimensional, and the density character dens(X) $\gg$ dens(Y), i.e., when X is "huge".]

论坛问题帖 | has_answers | Ady | 2010-01-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-017939 | MO-116947

Center of mass from the abstract point of view, or could the ancient Greeks invent modern analysis?

This is a very open-ended question, which may or may not have a perfect answer, and for which I have a few ideas but nothing like a clear picture. However, I guess it won't hurt to ask to see if people think about such things at all and if they do, what their ideas are. I don't know whether to make it CW or not: on one hand, it is pure mathematics, so we are within the usual set of standards to judge what's right and what's wrong, on the other hand, it is certainly not "a question of the type MO was designed for". So, I'm hesitant to check the community wiki option myself but have absolutely nothing against someone else doing so. I assume that the ancient Greeks had an idea of a complete normed space ($\mathbb R$ and $\mathbb R^2$ would be enough for our purposes for quite a while), a set, a linear transformation, and the center of mass. On the top of it, I assume they had as much common sense (probably more), as we have nowadays. The task is the usual one for Archimedes: given a reasonable non-empty set E in a complete linear space V, assign a point $C(E)$ to it that you can confidently call "the center of mass". For the purposes of this thread, let's consider bounded at most countable subsets in $V=\mathbb R$ first. If we can figure out what to do with this case to everyone's satisfaction, we can move to the next stage. It may be not a really illuminating model, but it has a few quite funny features already. The axioms of the center of mass are just the common sense ones: 1) The center of mass is never outside the closed convex hull of the set. 2) If A, B are disjoint, then $C(A\cup B)\in[C(A),C(B)]$. 3) If T is an affine transformation, then $C(TE)=TC(E)$. 4) (this is a bit tough, so feel free to drop or to modify it if it helps) If A,B are such that the sets $a+B$, $a\in A$ are disjoint, then $C(A+B)=C(A)+C(B)$ I don't know if we really need anything else (in particular, I'm not sure if the addition of the "obvious" definition of the center of mass of a finite set is needed, helpful, or hurtful), but feel free to play with this list in any reasonably way you want. The questions are the usual ones: A) Existence B) Uniqueness C) Way to find $C(E)$ given E. Any ideas, constructions, counterexamples, references, etc. (not necessarily restricted to the model I described) are welcome :). Edit: Thanks to everyone who responded! Let me clarify one thing. Yes, if we can create a meaningful notion of mass (say, if there exists a translation invariant Borel measure that is finite and positive on E), we can define the center of mass in the usual way and the only question will be if the definition is unambiguous. However, if I give you a symmetric set, you'll not hesitate to say that the center of symmetry is also the center of mass. Also, if I give you an infinite set and add one point to it, you'll probably (but not necessarily) agree that the center of mass won't feel this addition, the reason being that the set was infinitely times more massive than the point we added not as mush because we can measure the actual mass in some way but merely because infinitely many disjoint shifts of the point fit inside the set. In other words, quite often we can determine the relative size without being able to assign any meaning to the absolute size. This is one of the loopholes in the integration theory I'd like to exploit and see how far one can get with it. In a sense, that is a straight extension of the original Eudoxus line of thinking, which is why "ancient Greeks" entered the title of the question.

论坛问题帖 | has_answers | fedja | 2012-12-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017940 | MO-118667

Open problems in the theory of compact quantum groups

What are the important open problems in the theory of compact quantum groups? Or conjectures? Here is an example from An De Rijdt's Ph.D. thesis: Is every compact quantum group with the fusion rules of $SO(3)$ of the the form $A_{aut}(B,\varphi)$ for a finite dimensional C^*-algebra B and a δ-form φ on B? See page 83 at the end of Chapter 3 in An De Rijdt's Ph.D. thesis, available online from https://perswww.kuleuven.be/~u0018768/students/derijdt-phd-thesis.pdf

论坛问题帖 | has_answers | Uwe Franz | 2013-01-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017941 | MO-119352

Randomly switching street lights, in a square city

This is a combinatorics-probability question, best stated however in "recreational" terms. Imagine a $N\times N$ city, meaning that we have N horizontal streets, and N vertical streets. At each crossroads there is a street light. When evening comes, some of the lights are switched on, namely those corresponding to a certain given subset $E\subset(1,\ldots,N)\times(1,\ldots,N)$. Now assume that $2N$ kids come at night and start randomly playing with the switches: there is one such on/off switch at the end of each of the $2N$ streets. Problem: for each of the 4^N overall choices for the various switches, we count the number K of street lights that are switched on. What is the law of this random variable K, as a probability measure on $(1,2,\ldots,N^2)$, depending on the initial set E? [Edit, Jan 20. As signaled by Joseph O'Rourke in his answer below, computing the upper edge of the support of the measure μ_E in my problem is known as the Gale-Berlekamp game, a difficult question (details can be found via Google search). So I realise that my problem is probably extremely difficult, adding the "open-problem" tag. I'd be interested however in the case where E is an Hadamard matrix, cf. discussion with Gerhard Paseman in the comments below. Is there anything known about this measure μ_E? (I mean, not only about its support.) Btw here is the only non-trivial complete computation that I have so far: concerns the case $N=4$, where there are exactly $|E|=4$ street lights, positioned on

the main diagonal of the city. Here $\mu_E=\frac{1}{32}(\delta_4+12\delta_6+6\delta_8+12\delta_{10}+\delta_{12})$. Plus an experimental remark, that I'm not able to prove abstractly: the support of μ_E seems always to be an arithmetic progression.]

论坛问题帖 | has_answers | Fred L | 2013-01-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017942 | MO-120175

How many values a polynomial map misses?

Let F be a field. For a uni-variate polynomial $f(x)$ over F,let $M_f(F)$ denote the number of values that f misses, that is, the cardinality of the subset $F - f(F)$ in F. Assume that f is not surjective, that is, $M_f(F) > 0$. Is the number $M_f(F)$ big? If $F = {\mathbb F}_q$ is a finite field of q elements and $f\in \mathbb{F}_q[x]$ is a polynomial of degree $d>0$, then a simple result shows that if $M_f(F_q) > 0$, then $$M_f(F_q) \geq \frac{q-1}{d}. $$ As q goes to infinity, this lower bound goes to infinity. This suggests the following question for which I do not know the answer. Question. Assume that F is an infinite field. If $M_f(F)>0$. Is it true that $M_f(F) = \infty$? In other words, for an infinite field F, if $f \in F[x]$ misses one value, is it true that f misses infinitely many values? The answer is positive if F is big in the sense that F is algebraically closed (${\mathbb C}$ etc), or topologically closed ($\mathbb{R}$ or $\mathbb{Q}_p$ etc) or if F is a small field such as a Hilbertian field (number fields etc). Anyone has more examples or counter-examples? Gjergji-Zaimi points out that the above question has been asked before, see Can a non-surjective polynomial map from an infinite field to itself miss only finitely many points? To make my question somewhat newer, I will ask a multi-variable version of the question. It is unknown to me even in the case that the field is the complex numbers. Question: Assume that $f=(f_1,..., f_n): F^n -> F^n$ is a FINITE polynomial map in n variables over an infinite field F. If f misses one value, is it true that f misses infinitely many values? I do not know the answer even in the case when $F={\mathbb C}$ is the complex numbers if $n \geq 2$. Note that the finiteness assumption of the map f cannot be dropped, as Kharlamov (2012) has given the example $$f=(u(uv-1), u^2-(uv-1)^2)$$ which misses precisely one point, the origin (0,0) of ${\mathbb C}^2$.

论坛问题帖 | no_answers_at_retrieval | dwan | 2013-01-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017943 | MO-120819

open problems in Seiberg-Witten Theory on 4-Manifolds

What are some of the open problems in Seiberg-Witten Theory on 4-Manifolds.I tried googling but couldn't any. I tried googling it, but couldn't find any resources.The places where I can a survey or review of them would be welcome.

论坛问题帖 | has_answers | Koushik | 2013-02-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017944 | MO-120922

An open problem on general topology

There is an open problem in this paper: Classes defined by stars and neighbourhood assignments by van Mill and others. Problem 4.8. Is a regular (Tychonoff) star compact space metrizable if it has a G_δ-diagonal? A topological space X is said to be star compact if whenever $\mathscr{U}$ is an open cover of X, there is a compact subspace K of X such that $X = \operatorname{St}(K,\mathscr{U})$. More on star compactness see here. My question is this: Is always the cardinality of such regular (Tychonoff) star compact space less than 2^{ω_0}? See the related link here.

论坛问题帖 | has_answers | Paul | 2013-02-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017945 | MO-121010

Is there any o-minimal expansion of the real field with functions of growth higher than exponential?

Let $\bar{\mathbb{R}}$ be the structure of the real field, that is $(\mathbb{R},0,1,+,-,*,<)$. We say that a function f is of growth higher than exponential if for all $N\in \mathbb{N}$ there $f(x)$ is ultimately greater than $\exp^N(x)$. That is there is we can find $r\in \mathbb{R}$ such that for all $x>k$ we have $\exp^N(x) < f(x)$ where $\exp^N(x)$ stands for $\exp(\exp(\cdots \exp(x)\cdots))$ N times. Is it possible to find an o-minimal expansion (in the model theoretic sense) of $\bar{\mathbb{R}}$ where an f as above is definable?

论坛问题帖 | has_answers | Santiago | 2013-02-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017946 | MO-122115

problems from the scottish book

Which of the problems from the Scottish Book (pdf of English version) by Stefan Banach are still open? I know that one of the problems was solved by Per Enflo for which he got a live goose from Stanislaw Mazur.

论坛问题帖 | has_answers | Koushik | 2013-02-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017947 | MO-122199

Is it true that every f.g. infinite simple group has exponential growth?

Is it true that every finitely generated infinite simple group has exponential (word-)growth? Remark: As Mark Sapir has pointed out, the question whether every finitely generated group of subexponential growth is even residually finite has been answered in the negative in Anna Erschler. Not residually finite groups of intermediate growth, commensurability and non-geometricity, J. Algebra 272 (2004), no. 1, 154-172, http://www.sciencedirect.com/science/article/pii/S0021869303006410.

论坛问题帖 | has_answers | Stefan Kohl | 2013-02-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017948 | MO-12284

Topologically distinct Calabi-Yau threefolds

In dimensions 1 and 2 there is only one, respectively 2, compact Kaehler manifolds with zero first Chern class, up to diffeomorphism. However, it is an open problem whether or not the number of topological types of such manifolds of dimension 3 (Calabi-Yau threefolds) is bounded. I would like to ask what is known in this direction. In particular, is it known that the Euler characteristic or the total Betti number of Calabi-Yau threefolds can't be arbitrarily large? Are there any mathematical (or physical?) reasons to expect either answer? As a side question: I remember having heard several times something like "Calabi-Yau 3-folds parametrize (some kind of) vacua in string theory" but was never able to make precise sense of this. So any comments on this point or references accessible to mathematicians would be very welcome.

论坛问题帖 | has_answers | algori | 2010-01-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-017949 | MO-123317

Status of Grothendieck's conjecture on homomorphisms of abelian schemes

In [1] Grothendieck posits the following: Conjecture. Let S be a reduced connected scheme, locally of finite type over Spec($\mathbf{Z}$) or a field k, A and B two abelian schemes over S, l a prime number, $u_l: T_l(A) \rightarrow T_l(B)$ a homomorphism, and suppose that there exists a point $s\in S$ such that u_{ls} comes from a homormophism $u_s: A_s \rightarrow B_s$ of abelian schemes over $k(s)$. Then there exists an integer $n>0$ and a homomorphism $v: A \rightarrow B$ such that $T_l(v)=n u_l$. Next Grothendieck remarks that the conjecture follows from Tate's conjectures on algebraic cycles. He then proceeds to a give a proof, using the Serre-Tate theorem, for the case of a characteristic zero base field for which the statement holds with $n=1$. Question: What is the current status of this conjecture? If not resolved, are there any partial results for the positive characteristic case? Here is what I've spotted in the literature so far, all characteristic zero: Deligne in the second of three papers on Hodge theory, proves results about homomorphisms of abelian schemes over schemes S of finite type over $\mathbf{C}$. For instance: Proposition (Deligne 4.4.12 [2]) : Let $f: X \rightarrow S$ be an abelian scheme over a smooth scheme S. The following conditions are equivalent: (i) For every abelian scheme $g: Y \rightarrow S$, we have $$ \mathrm{Hom}_S(X,Y) \cong \mathrm{Hom}_S(R_1 f_* \mathbf{Z}, R_1 g_* \mathbf{Z}) $$ (ii) The condition (i) is verified for $X=Y$, and the centre Z of $\mathrm{End}_S(X)\otimes \mathbf{Q}$ does not admit a complex place $\rho: Z \rightarrow \mathbf{C}$ such that the direct factor $R_1 f_*\mathbf{Q} \otimes_{Z,\rho} \mathbf{C}$ of $R_1 f_* \mathbf{C}$ is of pure Hodge type (-1,0). Along similar lines, S.G. Tankeev in a 1976 paper [3] proves if S is a connected smooth curve over $\mathbf{C}$, and $\pi_i: X_i \rightarrow S, i=1,2$ are two abelian schemes, then under some natural conditions, the canonical homomorphism $$ \mathrm{Hom}_S(X_1,X_2) \rightarrow \mathrm{Hom}(R_1 \pi_{1*} \mathbf{Z}, R_1 \pi_{2*} \mathbf{Z})$$ is an isomorphism. In a paper the following year [4] Tankeev proves a Tate module variant of the above, of the form $$ \mathrm{Hom}_k(I_1,I_2)\otimes_\mathbf{Z} \mathbf{Z}_l \cong \mathrm{Hom}_G(T_l(I_1), T_l(I_2)),$$ where $G=\mathrm{Gal}(\overline{k}/k)$, and I_1, I_2 are abelian varieties over $k(t)$ whose Néron models admit compactifications with certain properties. Are there other well-known results along the same lines? I would especially like to know about the characteristic p case, as well any results that actually spread a homomorphism of abelian varities over a point of S to maps of abelian schemes over S, as in the conjecture. [1] A. Grothendieck Un théorème sur les homomorphismes de schémas abéliens Invent. Math. 2 59-78 (1966) [2] P. Deligne Théorie de Hodge: II Publications mathématiques de l'I.H.E.S, tome 40 (1971) p.5-57 [3] S.G. Tankeev On homomorphisms of abelian schemes Math. USSR Izvestija Vol. 10 (1976), No. 4 [4] S.G. Tankeev On homomorphisms of abelian schemes II Math. USSR Izvestija Vol. 11 (1977), No. 6

论坛问题帖 | has_answers | Zavosh | 2013-03-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017950 | MO-125049

Open problems in PDEs, dynamical systems, mathematical physics

(This question might not be appropriate for this site. If so, I apologize in advance. I would have posted to mathstack, but I'm looking for advice from active researchers.) I am an undergrad in math and physics getting ready to apply to grad school. So far, I know that I enjoy PDEs (and most kinds of analysis, generally), mathematical physics, and especially dynamical systems. But this is way too broad for choosing a Ph.D. program. My background in these subjects is somewhere between the undergraduate and graduate level, but certainly not 'up to date' or research-level. I want to get a clearer picture of these fields as they exist today. If I'm going to try to contribute to these topics in the next 5-10 years, I'd like to know what I'll be getting into. If you work in these fields, or you have colleagues who do, how would you describe the current state of affairs? Specifically: What are people trying to accomplish? How do those things fit into a larger picture? What are the obstructions? What are some of the major recent advances? What would you like to see happen over the next few decades? I'm looking for technical descriptions, preferably with references to actual papers. Pretend you were describing your work to a fellow mathematician from an entirely different specialty. Don't worry if it's over my head (I'm sure it will be); the point is to get a taste, to help narrow my interests, and to have a guide to come back to over the years. Thanks!

论坛问题帖 | has_answers | AJGibson | 2013-03-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-017951 | MO-127530

Surfaces filled densely by a geodesic

Which smooth, closed surfaces $S \subset \mathbb{R}^3$ have no single geodesic γ that fills S densely? Say a geodesic γ "fills S densely" if the closure of the set of points through which γ passes equals S. Some examples: A sphere: every geodesic is a great circle. Zoll surfaces, as discussed here: "Surfaces all of whose geodesics are both closed and simple." An ellipsoid. (Image from GeographicLib.) A torus generally has many geodesics that fill the surface. (Image by John Oprea) My assumption is that almost all surfaces have geodesics that fill them. Is this known, under any interpretation

of "almost all"? I would also be interested in extending the list of exceptional surfaces beyond {sphere, Zoll, ellipsoid}. Thanks for pointers! Answers Summary (18Apr2013): (Robert Bryant, Mikhail Katz) Any surface of revolution with poles has no dense geodesic. This holds for convex or nonconvex surfaces of revolution. (Robert Bryant) There are generalizations of Liouville surfaces (due to Goryachev-Chaplygin and to Dullin-Matveev) that have no dense geodesic. (Misha Kapovich) Every surface may be perturbed by gluing on "focusing caps" so that it has dense geodesics. (Keith Burns) Guess: There is always a dense geodesic on a closed Riemannian surface of genus ≥ 2.

论坛问题帖 | has_answers | Joseph O'Rourke | 2013-04-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017952 | MO-12758

Deligne-Simpson problem in the symmetric group

Question. Let $C_1,\dots,C_k$ be conjugacy classes in the symmetric group S_n. (More explicitly, each C_i is given by a partition of n; C_i consists of permutations whose cycles have the length prescribed by the partition.) Give a necessary and sufficient condition on C_i that would ensure that there are permutations $\sigma_i\in C_i$ with $$\prod\sigma_i=1.$$ Variant. Same question, but now σ_i's are required to be irreducible in the sense that they have no common invariant proper subsets $S\subset\lbrace 1,\dots,n\rbrace$. I am not certain how hard this question is, and I would appreciate any comments or observations. (I was unable to find references, but perhaps I wasn't looking for the right things.) This question is inspired by Jonah Sinick's question via the simple Geometric interpretation. Consider the Riemann sphere with k-punctures $X=\mathbb{CP}^1-\lbrace x_1,\dots,x_k\rbrace$. Its fundamental group $\pi_1(X)$ is generated by loops γ_i ($i=1,\dots,k$) subject to the relation $$\prod\gamma_i=1.$$ Thus, homomorphisms $\pi_1(X)\to S_n$ describe degree n covers of X, and the problem can be stated as follows: Determine whether there exists a cover of X with prescribed ramification over each x_i. The variant requires in addition the cover to be irreducible. Background. The Deligne-Simpson Problem refers to the following question: Fix conjugacy classes $C_1,\dots,C_k\in\mathrm{GL}(n,\mathbb{C})$ (given explicitly by k Jordan forms). What is the necessary and sufficient condition for existence of matrices $A_i\in C_i$ with $$\prod A_i=1$$ (variant: require that A_i's have no common proper invariant subspaces)? There are quite a few papers on the subject; my favorite is Simpson's paper, which has references to other relevant papers. The problem has a very non-trivial solution (even stating the answer is not easy): first there is a certain descent procedure (Katz's middle convolution algorithm) and then the answer is constructed directly (as far as I understand, there are two answers: Crawley-Boevey's argument with parabolic bundles, and Simpson's construction using non-abelian Hodge theory). The same geometric interpretation shows that the usual Deligne-Simpson problem is about finding local systems (variant: irreducible local systems) on X with prescribed local monodromy. So: any remarks on what happens if we go from $\mathrm{GL}(n)$ to S_n?

论坛问题帖 | has_answers | t3suji | 2010-01-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017953 | MO-129605

A conjecture on closed discrete subset

I am struggling with this old problem, which is also posted here: Let X satisfy countable chain condition(abbreviated as CCC) and X has a regular G_δ-diagonal. Then the cardinality of X is at most $\mathfrak c$. X has a regular G_δ-diagonal iff there is a collection of open sets of X^2, say $\lbrace U_n: n\in N\rbrace$, such that $\Delta=\bigcap\lbrace \overline{U_n}: n \in N\rbrace$, where $\Delta=\lbrace(x,x): x \in X\rbrace$. Note that the question is answered; however I hope to get new proof. By certain effort, If $|X|>\mathfrak c$, we can get an uncountable closed discrete subset S of X, and for any point $x \in X$, there exists an open set U_x such that $\overline{U_x} \cap S$ has at most one point. I would like to know whether the following conjecture is right, wrong, or neither: Let X be a Hausdorff space. If $S \subset X$ is an uncountable closed discrete subset of X, and for any point $x \in X$, there exists an open set U_x such that $\overline{U_x} \cap S$ has at most one point. Then could we obtain an uncountale collection of disjoint open sets in X? Thanks for your any help.

论坛问题帖 | has_answers | Paul | 2013-05-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017954 | MO-130255

Optimal inspection path on a sphere

Suppose you would like to "inspect" every point of a unit-radius sphere $S \subset \mathbb{R}^3$ by walking along a path γ on S, but you can only see a distance d from where you stand. Q1. For a given $d \in (0,\pi]$, what are the shortest such paths $\gamma(d)$? Two other ways to define $\gamma(d)$: The shortest path γ such that every point on S is no more than a distance d from some point of γ. The shortest path γ such that the union of disks of radius d centered on every point of γ cover S. For $d \in [\pi/2,\pi]$, $\gamma(d)$ is an arc of a great circle of length $2\pi - 2d$. So, for $d=\pi$, γ is a single point; for $d=3\pi/4$, γ is an arc of length $\pi/2$; for $d=\pi/2$, γ is a semicircle. As d approaches zero, it might be that γ is a type of spiral, e.g.: (Image from grabcad.com.) What is quite unclear to me is when d is less than but close to $\pi/2$. For example, suppose $d = \frac{5}{12}\pi = 75^\circ$. The union of disks of this radius centered on the equatorial semicircle that works for $\pi/2$ is shown in (a) below: a strip connecting the north and south poles is uncovered (blue circles are boundaries of disks of radius d centered on γ): One can construct a candidate γ as in (b) above: extend the semicircle at each end so that all points on the equator are reached, and add short vertical sections, red to reach the north pole, and green to reach the south pole. It does not seem likely that such sharp turns yield the shortest γ. Perhaps this question can be answered even without precise understanding of $\gamma(d)$ for all d: Q2. Does $\gamma(d)$ vary continuously with respect to d? I originally wanted to explore this for any (smooth, closed, compact) surface in $\mathbb{R}^3$, but it already seems not so straightforward for the sphere. Q3. Has this question been studied before? It feels classical. The question could also be posed for $(d{-}1)$-dimensional spheres in $\mathbb{R}^d$, again seeking an optimal inspection path for each d. (My interest in $\mathbb{R}^3$ stems from two related

concepts: Voronoi diagrams on a surface, and the cut locus with respect to a path, although neither seems to help here.) Thanks for any pointers or ideas! 9Aug2022. This issue was touched again by Fejes Tóth's Exploring a planet problem by Zhao, Yufei. "Exploring a planet, revisited." American Mathematical Monthly (2022): 1-3. But the Tóth version uses n great-circle satellites. Related to the Shortest closed curve to inspect a sphere, but quite distinct. As far as I know, my original questions remain unsolved. (Related earlier MO questions: Optimal paintbrush geodesics; Shortest closed curve to inspect a sphere.)

论坛问题帖 | has_answers | Joseph O'Rourke | 2013-05-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-017955 | MO-131528

How closed-form conjectures are made?

Recently I posted a conjecture at Math.SE: $$\int_0^\infty\ln\frac{J_\mu(x)^2+Y_\mu(x)^2}{J_\nu(x)^2+Y_\nu(x)^2}\mathrm dx\stackrel{?}{=}\frac{\pi}{2}(\mu^2-\nu^2),$$ where $J_\mu(x)$ and $Y_\mu(x)$ are the Bessel function of the first and second kind. It is supported by numerical calculation with hundreds of digits of precision for many different values of μ, ν. The question is open for several days with +500 bounty on it and is not resolved yet. But my question here is not about if this conjecture true or false. Obviously, several possible closed forms matched my numeric calculations, for example: $$\left(\frac{\pi}{2}+7^{-7^{7^{7^{7^{7^{\sqrt{5}+\sin \mu\nu}}}}}}\right)(\mu^2-\nu^2).$$ But for some reason that I cannot clearly explain (or even understand) I selected the simpler one, and I am strongly inclined to search for its proof rather than a disproof. I believe most people would feel and behave exactly the same way. (1) Are there any mathematical or philosophical reasons supporting this position? Why when we calculate some sum or integral (which do not contain explicit tiny quantities like $10^{-10^{10^{.^{.^{10}}}}}$) with thousands of digits of precision and it matches some simple closed-form expression, we inclined to believe this is the exact equality rather than an accidental very close value? (2) Are there known cases when such intuition turned out to be wrong? And one more question: (3) Do you believe there can be exact closed forms for some infinite sums or integrals, that cannot be proved in ZFC or any its reasonable extension (like adding some large cardinal axioms) - so to speak, equalities that hold without any reason.

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2013-05-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017956 | MO-131892

A conjecture on intersection of some intervals.

It was proved here that if $a\in \mathbb{N}_{\geq3}$ then $$\bigcap_{i = 1}^{a} \bigcup_{j = 0}^{i-1} \left[\frac{1+aj}{i},\frac{a(j+1)-1}{i}\right] = \varnothing \tag{1}$$ It may be conjectured that forcing $i\ne b$, where $1\leq b< a$, renders (1) untrue, that is, the result is not an empty interval. We look at the diagram here from the link above for $a=5$ to get a better picture: Red is the interval, Yellow are the gaps between the intervals that cause the intersection to be a null set, white gaps do not effect the intersection. The conjecture here says that if we were to remove any one of the top 4 strips, then there will form a region of intersection. I tried analyzing the gaps but everything seems to meet up at a dead end. What tools may one employ to handle such problems?

论坛问题帖 | has_answers | Maaz-ul-Haq | 2013-05-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017957 | MO-131973

Distinctive property of the primes 17 and 19?

Consider the question whether it is true that a prime number p divides $1^1+2^2+3^3+....+(p-1)^{p-1}$ if and only if $p \in \{17,19\}$. For the obvious heuristic reasons, for large n one would expect there to be roughly $\ln(\ln(n))$ such primes $p < n$, however it seems that presently no examples other than 17 and 19 are known. Is there a more efficient way of looking for examples than the brute force method of testing the primes one-by-one?

论坛问题帖 | has_answers | Mihir Sheth | 2013-05-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017958 | MO-131998

Needle probing for a convex body

Suppose there is an unknown closed convex body K of volume vol$(K) = V$ inside the unit cube $[-\frac{1}{2}, \frac{1}{2}]^d$ in $\mathbb{R}^d$. You are permitted to probe with a (one-dimensional) ray r, which detects whether r includes some point of K, or instead if $r \cap K = \varnothing$. My question is: Q1. What is the fewest number of such needle probe rays that guarantee hitting K, as a function of its volume V? I am especially interested in $\mathbb{R}^3$. For example, for $V \ge \frac{1}{2}$, one ray through the origin parallel to an axis suffices. Here such a ray touches the boundary of a body K with vol$(K)=\frac{1}{2}$: To be specific: Q2. How many needle probes suffice for $V \ge \frac{1}{4}$ in $\mathbb{R}^3$? Four parallel rays in a grid pattern do not suffice, but nine do suffice (I believe): Q3. In $\mathbb{R}^3$, is it sometimes more efficient to use nonparallel rays? This feels like a classic problem, but I am not finding literature. Thanks for pointers or ideas! Update 1 (28May13). Here is Yoav Kallus' example showing that dropping the middle of the 9 points of the grid permits a triangle with area larger than $\frac{1}{4}$ to avoid detection: Update 2 (30May13). Q2 and Q3 are now answered by Benjamin Dickman and Douglas Zare: for $V=\frac{1}{4}$, three needles suffice, and more are needed for parallel needles. Q1, in its full generality, is difficult, and so I've added the "open problem" tag.

论坛问题帖 | has_answers | Joseph O'Rourke | 2013-05-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017959 | MO-132269

An elementary expression for $_3F_2(1,1,9/4;2,2;-1)$

Consider the following series: $$S=\sum_{n=1}^\infty\frac{(-1)^n\ \Gamma\left(\frac{5}{4}+n\right)}{n^2\ \Gamma(n)}.$$ It can be expressed in terms of a hypergeometric function: $$S=-\frac{5}{16}\Gamma\left(\frac{1}{4}\right)\ { _3F_2}\left(1,1,\frac{9}{4};2,2;-1\right).$$ I tried to find an expression of S using elementary functions and ended up with this conjecture: $$S\stackrel{?}{=}\frac{\Gamma\left(\frac{1}{4}\right)}{16}\Bigg(8\sqrt[4]{8}-16-\ln\frac{24\sqrt{69708+49291\sqrt{2}}+3168\sqrt{2}+4481}{4096}\\\\+\arctan\frac{24\sqrt{49291\sqrt{2}-13260}}{6913}\Bigg).$$My derivation of this formula is quite long and uses a non-rigorous, heuristic approach involving heavy use of techniques like guessing sequence formulas using Mathematica command FindSequenceFunction and OEIS superseeker, and recognizing approximate numeric quantities using RootApproximant, TranscendentalRecognize, Inverse Symbolic Calculator and Mathematica command WolframAlpha[ToString[value], IncludePods -> "PossibleClosedForm", TimeConstraint -> ∞] The formula holds with at least 1800 decimal digits of precision. Now I am looking for a rigorous way to prove this formula. Can you suggest any ideas how to do that?

论坛问题帖 | has_answers | X.C. | 2013-05-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017960 | MO-133771

Does this modified Hasse principle hold for curves?

Let C be a curve over $\mathbb Q$ with a point P on Pic^1. For each $\mathbb Q$-rational point Q, $Q-P$ is a point on the Jacobian J. We can use the map $H^0(\mathbb Q, J) \to H^1(\mathbb Q,J[n])$ to obtain a class in $H^1(\mathbb Q,J[n])$ associated to each rational point. Rather than trying to find a local obstruction to existence of rational points on C, we could try to find a local obstruction to existence of rational points in each $H^1(\mathbb Q, J[n])$ class. A local obstruction at $\mathbb Q_p$ would occur when the class, projected down to $H^1(\mathbb Q_p, J[n])$ does not arise from any $\mathbb Q_p$-rational point of C. If C does not have rational points, must there be some n such that each class in $H^1(\mathbb Q, J[n])$ has a local obstruction at some p or at ∞?

论坛问题帖 | has_answers | Will Sawin | 2013-06-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-017961 | MO-13394

Finite Rank Commutators

My former student Detelin Dosev and I are interested in classifying the commutators in $L(X)$, the bounded linear operators on the Banach space X (see our joint paper on my home page or the ArXiv and the references therein, and recall that T is a commutator means that there are A and B in $L(X)$ so that $T=AB-BA$). If T is a finite rank operator in $L(X)$ with zero trace then it is classical that T is a commutator. The converse is obviously true when X is finite dimensional. Conjecture. There is an infinite dimensional Banach space X such that all finite rank commutators in $L(X)$ have zero trace. If true, this conjecture is very hard to prove, but maybe it is known to be false? We are also interested in conditions on X that guarantee that all finite rank operators in $L(X)$ are commutators. If X is isomorphic (meaning linearly homeomorphic) to its closed subspaces of codimension one, then rank one projections in $L(X)$ are commutators and hence every finite rank operator in $L(X)$ is the sum of two commutators, but we do not know whether every finite rank operator in $L(X)$ is a commutator. If also X contains a complemented subspace isomorphic to $X \oplus X$, then every diagonalizable finite rank operator in $L(X)$ is a commutator, but we still don't know whether all finite rank operators in $L(X)$ are commutators. Are we missing something easy? A Banach space X is said to have a Pelczynski decomposition provided X is isomorphic to $(X \oplus X \oplus X \oplus ...)_p$ for some $1 \le p \le \infty$ or $p = 0$. A general result is that if X admits a Pelczynski decomposition, then every compact operator on X is a commutator, so standard spaces do not give examples of finite rank non commutators.

论坛问题帖 | no_answers_at_retrieval | Bill Johnson | 2010-01-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-017962 | MO-13437

Hilbert subspaces of indefinite inner product spaces

Let E be a real linear space, endowed with a non-degenerate symmetric bilinear form $(.,.)$. Suppose that the [indefinite] inner product space $(E,(.,.))$ satisfies the following [sequential] properties: (WSC) If the sequence { ${(x_{n},\ y)}$ } is Cauchy for each y in E, then there exists some x in E such that $\left(x_{n}-x,y\right)$ $\rightarrow$ 0 whenever y in E. (That is to say, $(E,(.,.))$ is weakly sequentially complete.) and (DPG) If $\left(x_{n},y\right)$ $\rightarrow$ 0 for every y in E, then $\left(x_{n},x_{n}\right)$ $\rightarrow$ 0. (That would be sort of "Dunford-Pettis & Grothendieck'' property for indefinite inner product spaces.) Suppose also that $|E|$ is "big enough'' (at least $|E|$ $>|\mathbb{R}|$). Conjecture. $(E,(.,.))$ contains an infinite-dimensional Hilbert subspace. (That is, there exists a linear isometry from $(\ell^{2},<.,.>)$ into $(E,(.,.))$.)

论坛问题帖 | no_answers_at_retrieval | Ady | 2010-01-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-017963 | MO-13444

Prime divisors of numbers 2^n + 3

I'm interested in the following problem: do there exist infinitely many prime numbers p such that $p^2|2^{n}+3$ for some natural number n? Some motivation: If we replace the function $2^n + 3$ with the $f(n)$ where $f \in \mathbb{Z}[x]$ is non-constant that this is true (follows Hensel lemma). So, it's rather natural to try proving this for some other non-polynomial functions. $2^n + 3$ is an easy example of such function. There is also another good reason: sequence $a_n = 2^{n} + 3$ satisfies the reccurence relation: $a_{n+2} = 3a_{n+1} - 2a_{n}$. And for example this problem is true for Fibonacci sequence. So, for Fibonacci it's easier even if the closed form of Fibonacci numbers is more complicated. But I think that the reason of this is that the Fibonacci numbers satisfy some "good" identities which other sequences don't have to share. Now some remarks: It's easy

exercise to prove that there are infinitely many primes p such that $p|2^{n}+3$. Also, if we try "correcting" n to work also for p^2 and we try $m=n+k(p-1)$ we see that it is possible unless p is Wieferich prime, i.e. satisfies $p^2|2^{p-1}-1$. And this gives us nothing as we don't know much about Wieferich primes... This method can of course be generalized in such way: if $p|2^{n}+3$ and order of 2 mod p^2 is greater than order of 2 mod p then we can find m such that $p^2|2^{m}+3$. But I don't really think that it helps. I'm interested in some information about this problem (especially if it's open or not) and also related problems. We can ask a general question: for which functions f we know that this is true? Edit: Sorry for the confusion with k, deleted.

论坛问题帖 | has_answers | user3645 | 2010-01-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-017964 | MO-135608

A conjecture by Euler about $8n+3$

Euler's conjecture: For any positive integer n, $8n+3$ can be represented as a sum $$8n+3=(2k-1)^2+2p,$$ where k is a positive integer, and p is a prime. I want to know whether there has been progress on the problem. Could you recommend some references? Thanks a lot.

论坛问题帖 | has_answers | Ziang Chen | 2013-07-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017965 | MO-135765

Open problems in algebraic topology and homotopy theory

Some time ago (I see it was initially written before 1999?) Mark Hovey assembled a list of open problems in algebraic topology. The list can be found here. Some of the problems I know about have been worked on quite a bit since the time of writing. The list is very good as is, but there must also be a few good additions since 1999. Can someone point me to a more recent list of open problems in algebraic topology? My googling isn't turning up much else. Thank you!

论坛问题帖 | has_answers | Glen M Wilson | 2013-07-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-017966 | MO-135945

A question about the Axiom of Choice

Let AC denote the Axiom of Choice. Let PP denote the so-called "Partition Principle" which states that "If S is a non-empty set and T is a non-empty set of pairwise disjoint subsets of S, then S can be mapped onto T". It is well known that "AC implies PP" is provable in ZF, but the question of whether "PP implies AC" is provable in ZF has long been an open problem. What is the present status of this problem? Has any progress been made on it? Or-on the other hand-have any models of ZF been constructed in which pp is true and some consequence of AC -such as the existence of non-measurable sets of real numbers-is false? If "pp implies AC" could be proved in ZF, this would seem to provide a powerful philosophical argument for accepting AC. In my opinion, any set theory in which pp can be disproved yields a really counter-intuitive picture of the "Set-theoretical Universe".

论坛问题帖 | has_answers | Garabed Gulbenkian | 2013-07-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017967 | MO-136157

Set-theoretic reformulation of the invariant subspace problem

The invariant subspace problem (ISP) for Hilbert spaces asks whether every bounded linear operator A on l^2 (with complex scalars) must have a closed invariant subspace other than $\{0\}$ and l^2. A subspace E is invariant for A if $A(E) \subseteq E$. Some time ago I noticed a reformulation that has a set-theoretic flavor. Let P be the set of all linear operators A from a finite-dimensional subspace of l^2 into l^2 such that (1) $\|A\| < 1$ and (2) if E is a nonzero subspace of the domain of A then $A(E) \not\subseteq E$. Order P by reverse inclusion and for any unit vectors $v,w \in l^2$ define $$D_{v,w} = \{A \in P: \langle A^nv,w\rangle \neq 0\mbox{ for some n such that A^nv is defined}\}.$$ It is easy to see that each $D_{v,w}$ is dense in P. Having a counterexample to the ISP is the same as having a filter of P that meets each $D_{v,w}$. My question for the set theory experts on MO is simply whether this version of the problem suggests any possible approaches?

论坛问题帖 | no_answers_at_retrieval | Nik Weaver | 2013-07-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017968 | MO-136351

Small quadrilaterals containing a given convex region

It is easy to prove that (*) Every convex planar set of area 1 is contained in a quadrilateral of area 2. It is also easy to see that statement (*) remains true if the constant 2 is replaced with a somewhat smaller one. Contest: Find such a constant, the smaller the better. Update: Reaching $\sqrt 2$ and even a strictly smaller value was proved by Chakerian (1973) and Kuperberg (1983) and the research challenge offered is to improve it even further, and perhaps even to verify the conjecture that the minimum is attained by a regular pentagon. But any nice arguments for bounds below 2 are welcome.

论坛问题帖 | has_answers | Wlodek Kuperberg | 2013-07-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-017969 | MO-136925

Integer-distance sets

Let S be a set of points in $\mathbb{R}^d$; I am especially interested in $d=2$. Say that S is an integer-distance set if every pair of points in S is separated by an integer Euclidean distance. What are examples of maximal integer distance sets? (Maximal: no point can be added while retaining the integer-distance property between all pairs.) Of course the lattice points along any one line parallel to a coordinate axis in $\mathbb{R}^d$ constitute a countably infinite integer-distance set. What is an example of an infinite integer-distance set of noncollinear points? I know that Euler established that every circle contains a dense rational-distance set. Scaling any one circle by a large common denominator provides a rich,

but finite integer-distance set. For example, these five points on a circle are all separated by integer distances: $$ (1221025, 0), (781456, 586092), (439569, 586092), (270400, 507000), (180625, 433500) $$ I'm sure this is all known... Thanks for enlightening me! (This is tangentially related to my earlier question, "Rational points on a sphere in $\mathbb{R}^d$.") Update1. It turns out that determining the integer-distance sets is fundamentally open. What is known is nicely summarized by Robert Israel and "Daniel m3." In particular, via the Kreisel & Kurz paper, it is unknown (or was unknown in 2008) whether or not there exists an 8-point integer-distance set in $\mathbb{R}^2$, with no three of the points collinear and no four cocircular. Update2. Also open is a related problem identified by Nathan Dean: How many non-cocircular integer-distance points can be found on a parabola, a scaling of $y = x^2$? Nathan proved there are infinitely many sets of three such points; Garikai Cambell proved there are infinitely many sets of four such points. But the existence of five such points seems open. I just learned the parabola problem from this MSE question. Update3 (21 Jul 2013). I ran across this just-published paper, which explores the in-some-sense obverse of the question I asked: What are the largest point sets in $\mathbb{R}^d$ that avoid points an integral distance apart. Kurz, Sascha, and Valery Mishkin. "Open Sets Avoiding Integral Distances." Discrete & Computational Geometry (2013): 1-25. (Springer link) Update4 (29 Nov 2014). There is a nice article at Dick Lipton's blog on Ulam's 70-year-old un-resolved conjecture: If S is an rational-distance set, then it is not dense in the plane. And that article cites the Kurz-Mishkin paper above. Update5 (11 Mar 2024). Greenfeld, Rachel, Marina Iliopoulou, and Sarah Peluse. "On integer distance sets." arXiv:2401.10821 abstract (2024). "Our main result is that any integer distance set in the Euclidean plane has all but a very small number of points lying on a single line or circle. From this, we deduce a near-optimal lower bound on the diameter of any non-collinear integer distance set..." "It turns out that all so-far-known integer distance sets ... are of a similar special form: they have all but up to four of their points lying on a single line or circle. In this paper, we develop a new approach to the study of integer distance sets that enables us to prove a structure theorem partially explaining this phenomenon."

论坛问题帖 | has_answers | Joseph O'Rourke | 2013-07-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-017970 | MO-137287

Tiling the square with rectangles of small diagonals

For a given integer $k\ge3$, tile the unit square with k rectangles so that the longest of the rectangles' diagonals be as short as possible. Call such a tiling optimal. The solutions are obvious in the easy cases when k is the square of an integer and for a few small values of k only (unpublished). In each of the solved cases, the sides of all rectangles turn out to be rational and their diagonals are equal. Question. In an optimal tiling, must the sides of all rectangles be rational and their diagonals be equal? The analogous question for tiling the n-dimensional cube with rectangular boxes can be asked for every $n\ge3$ as well.

论坛问题帖 | has_answers | Wlodek Kuperberg | 2013-07-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017971 | MO-138368

Tarski Monster group with prime 5

Does the Tarski Monster group with prime 5 exist? I know that for 2 and 3, the group does not exist, but what about 5?

论坛问题帖 | no_answers_at_retrieval | D. N. | 2013-08-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017972 | MO-141744

Can six square numbers be simultaneously represented in a single sum of consecutive odd numbers?

I had some free time from my work to do a little exploration regarding the existence (or non existence) of perfect cuboids. A solution is represented by the set of Diophantine equations: $a^2 + b^2 = x^2$ $a^2 + c^2 = y^2$ $b^2 + c^2 = z^2$ $a^2 + b^2 + c^2 = R^2$ If there exists a perfect cuboid, it implies that there is a consecutive sum of odd numbers from 1 to the Rth odd number representing the space diagonal of a perfect cuboid, and that Six other perfect squares representing the face diagonals and edges also exist with the sum. It also means that each edge and face diagonal can be represented as a sum starting from 1 within R^2 I represent this consecutive arithmetic sequence like this: $1+3+5+...+ R_o = R^2$ Where R_o is the Rth odd number in a consecutive sequence. The set of edge, face diagonals and the body diagonals $[a,b,c,x,y,z,R]$ can each be represented similarly. By using the fourth equation in the set, we see that: $(1+3+5+...+a_o) + (1+3+5+...+b_o) + (1+3+5+...+c_o) = 1+3+5+...+R_o$ Then $(1+3+5+...+x_o) + (1+3+5+...+c_o) = 1+3+5+...+R_o$ Since by first equation and our definition, $(1+3+5+...+a_o) + (1+3+5+...+b_o) = (1+3+5+...+x_o)$ Which Implies $1+3+5+...+x_o = (c+1)_o + ... + R_o$ by subtracting $(1+3+5+...+c_o)$ Which implies: $x^2 = (c+1)_o + ... + R_o$ By using the same process, we can derive this for the others: $a^2 = (z+1)_o + ... + R_o$ $b^2 = (y+1)_o + ... + R_o$ $c^2 = (x+1)_o + ... + R_o$ $x^2 = (c+1)_o + ... + R_o$ $y^2 = (b+1)_o + ... + R_o$ $z^2 = (a+1)_o + ... + R_o$ Ultimately, the Perfect Cuboid Problem implies that there are two unique ways to represent each side and face diagonal in two different ways. Another way to see this is that it is possible to represent each side and face diagonal between $1+ ... + R_o$ both forwards and backwards. This also demonstrates that $a<b<c<x<y<z<R$ That is about as far as I got. My efforts now are attempts to demonstrate that it is not possible for one of these to be true, and by extension disproving others. I also believe that by extending the set of Diophantine equations, Perfect "Hypercuboids" can be disproved as well. Suggestions are appreciated.

论坛问题帖 | no_answers_at_retrieval | Richard Skinner | 2013-09-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-017973 | MO-143695

What are the divisors of $2n^2 - \sigma_{1}(n^2)$ for composite n?

What are the divisors of $2n^2 - \sigma_{1}(n^2)$ for composite n? Here, σ_{1} is the classical sum-of-divisors function. For example, $\sigma_{1}(3^2) = 1 + 3 + {3^2} = 13$. (The function $D(x) = 2x - \sigma_{1}(x)$ is called the deficiency of x [see OEIS Sequence A033879].) Now, for the motivation behind the question: Let $N = {q^k}{n^2}$ be an odd perfect number given in Eulerian form (i.e., q is prime with $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q, n) = 1$). Since N is perfect, we have the equation $$\sigma_{1}(N) = 2N \Longleftrightarrow \sigma_{1}({q^k}{n^2}) = \sigma_{1}(q^k)\sigma_{1}(n^2) = 2{q^k}{n^2}.$$ We can express $\sigma_{1}(q^k)$ as $$\sigma_{1}(q^k) = 1 + q + \ldots + q^{k-1} + q^k = \left(1 + q + \ldots + q^{k-1}\right) + q^k = \sigma_{1}(q^{k-1}) + q^k$$ whereupon we have $$\sigma_{1}(q^k)\sigma_{1}(n^2) = \left(\sigma_{1}(q^{k-1}) + {q^k} \right)\sigma_{1}(n^2) = 2{q^k}{n^2}.$$ Dividing through by q^k in $$\left(\sigma_{1}(q^{k-1}) + {q^k} \right)\sigma_{1}(n^2) = 2{q^k}{n^2},$$ we get $$2n^2 - \sigma_{1}(n^2) = I(q^{k-1})\cdot{\frac{\sigma(n^2)}{q}},$$ where $$I(x) = \frac{\sigma_{1}(x)}{x}$$ is the abundancy index of x. From the last equation, since $q \geq 5$ implies $1 \leq I(q^{k-1}) < \frac{5}{4} = 1.25$, we have the biconditional $$k = 1 \Longleftrightarrow 2n^2 - \sigma_{1}(n^2) = \frac{\sigma(n^2)}{q}.$$ In other words, we have the biconditional $$k = 1 \Longleftrightarrow 2n^2 - \sigma_{1}(n^2) = \frac{\sigma(n^2)}{q^k}.$$ [In the paper The Abundancy Index of Divisors of Odd Perfect Numbers, the assertion $k = 1$ for odd perfect numbers $N = {q^k}{n^2}$ was called as Sorli's conjecture -- although, in a more recent (proceedings) paper by B. Beasley of Presbyterian College, titled Euler and the Ongoing Search for Odd Perfect Numbers, he points out that it was in fact Descartes who initially made this conjecture (in a letter to Marin Mersenne in 1638), "with Frenicle's subsequent observation occurring in 1657".] Now, following Broughan, et. al., it is known that the index $\frac{\sigma_{1}(n^2)}{q^k}$ of the Euler factor q^k satisfies $$\frac{\sigma_{1}(n^2)}{q^k} \geq 315,$$ and $$\frac{\sigma_{1}(n^2)}{q^k}$$ cannot take any of the 11 forms $$\{p, p^2, p^3, p^4, p^5, p^6, {p_1}{p_2}, {p_1}^2{p_2}, {p_1}^3{p_2}, {p_1}^2{p_2}^2, {p_1}{p_2}{p_3}\}$$ where p is any odd prime and p_1, p_2, p_3 are any distinct odd primes. Now back to the original question: We want to know the divisors of $2n^2 - \sigma_{1}(n^2)$ for composite n. Note that, if n is prime, say r, then we have $$2n^2 - \sigma_{1}(n^2) = 2r^2 - \sigma_{1}(r^2) = 2r^2 - (r^2 + r + 1) = r^2 - r - 1 = r(r - 1) - 1,$$ which is, in general, odd (so that $2r^2 - \sigma_{1}(r^2) \equiv 1 \pmod 2$ for r prime.) [In fact, since $\sigma_{1}(n^2) \equiv 1 \pmod 2$ for all positive integers n, then we have $2n^2 - \sigma_{1}(n^2) \equiv 1 \pmod 2$ for all positive integers n.] Of course, the situation is more complicated when $\omega(n) \geq 2$ (where $\omega(x)$ is the number of distinct prime factors of x).

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2013-10-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017974 | MO-143743

Product of exponents of prime factorization

Let $p(n)$ be the product of the exponents of the prime factorization of n. For example, $$p(5184) = p(2^6 3^4) = 24 \;,$$ $$p(65536) = p(2^{16}) = 16 \;.$$ Define $P(n)$ as the number of iterations of $p(\;)$ to reduce n to 1. For example, $P(5184) = 3$ because $$p(5184)=24, \;p(24) = p(2^3 3^1) = 3, \;p(3)=1 \;;$$ and $P(65536)=4$ because $$p(65536) = 16, \;p(16)=p(2^4)=4, \;p(4)=p(2^2)=2, \; p(2)=1 \;.$$ Finally, define $m(k)$ to be the minimum value of n such that $P(n) = k$. Q1. What is $m(k)$? (I ask this question out of curiosity, not because it is part of a research program. It was previously posed on MSE.) It is easy to see that $m(1)=2$, $m(2)=4$, and $m(3)=16$, the latter because $16=2^{2^2}$. But, thanks to Calvin Lin's insight, $m(4)$ is not a power of 2, but instead is $m(4)=1296= 2^4 3^4$. I do not know the value of $m(5)$. Q2. More specifically: What is $m(5)$? I do know that $m(5) > 2 \times 10^8$. Update. Will Jagy showed that almost certainly $m(5) = 2^9 3^6 5^4 7^3 11^2 =9681819840000 \approx 10^{13}$. As it seems that an explicit expression for $m(k)$ is not in the offing, I will accept his resolution of Q2 and leave Q1 open.

论坛问题帖 | has_answers | Joseph O'Rourke | 2013-10-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017975 | MO-144080

Is $\varliminf_{n \rightarrow +\infty} |n \sin n| = 0$ correct, where n is an integer?

Is it true that $\varliminf_{n \rightarrow +\infty} |n \sin n| = 0$, where n runs over the integers? The existence of the limes inferior follows from Dirichlet's approximation theorem, but the problem is to prove that it is 0.

论坛问题帖 | has_answers | Y.X | 2013-10-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017976 | MO-145084

Euclidean real quadratic fields

It is known that, under GRH, a real quadratic field is Euclidean iff it is a UFD. So, assuming the conjecture of Gauss and GRH, we expect that there are infinitely many Euclidean real quadratic fields. I want to know if there are any approaches in showing that there are infinitely many Euclidean real quadratic fields by explicitly constructing the Euclidean functions. Can we hope for a single (or finitely many) Euclidean function(s) that will do the job or we need infinitely many different Euclidean functions? Can these Euclidean functions be obtained by modifying the norm function? Since very few non-norm Euclidean quadratic fields are known, can we at all hope for showing the infinitude of real UFD quadratic fields by finding infinitely many Euclidean quadratic fields?

论坛问题帖 | has_answers | Pritam Majumder | 2013-10-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017977 | MO-145299

Decidability of equality of elementary expressions

In the following definition the term expression is to be understood as a finite tree built from formal symbols without any predefined meaning assigned to them. Define the set $\mathcal{E}$ of elementary expressions as the minimal set such that: the expression i is in $\mathcal{E}$ if $x\in\mathcal{E}$, then the expression $\exp(x)\in\mathcal{E}$, if $x\in\mathcal{E}$, then the expression $\ln(x)\in\mathcal{E}$, if $x,y\in\mathcal{E}$, then the expression $(x\cdot y)\in\mathcal{E}.$ Now we can assign meaning (a numeric value) to expressions in $\mathcal{E}$ as follows: i is the imaginary unit, $\exp(x)$ is the exponent of x, $\ln(x)$ is the principal branch of the natural logarithm of x (unless the value of x is zero), $(x\cdot y)$ is the product of x and y. These rules do not assign a value to an expression if it contains a logarithm of an expression whose value is zero, e.g. $$\ln(\ln((i\cdot(i\cdot(i\cdot i)))))$$ in which case we say that the expression is invalid. Otherwise (if a value is successfully assigned), we say that the expression is valid. Note that expressions in $\mathcal{E}$ can represent all values that can be constructed from integers and elementary functions, e.g. $\sqrt[3]{3+\sqrt2},\ \sin\frac\pi{17},\ \arctan\frac27,\ \pi^e$, etc. Question: Is validity of expressions in $\mathcal{E}$ a decidable problem? In other words, is there an algorithm that, given an expression e in $\mathcal{E}$ as an input, always terminates and gives a correct yes/no answer indicating if the expression e is valid? Equivalent problems could be to check if a given expression is zero, or to check if two given expressions are equal to each other.

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2013-10-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-017978 | MO-145467

A challenging non homogenous fractional inequality

I have posted this question on Stackexchange but it has received no answer so far. It is a challenging generalization of several difficult inequalities, where none of the usual methods used in inequalities seems to work. I would like to know if someone has a deeper insight into what is happening here. Let $n\in\mathbb N$, $\lambda> 1$ and put $k:=\lambda n$ and $c:=\frac{\lambda-\frac 1n}{\lambda-1}=\frac{k-1}{k-n}$. For reals $a_1,...,a_n\ge 0,$ define $ r:=\prod_{i=1}^n a_i^c$. Then prove or disprove $$ {\sum\limits_{i=1}^n\displaystyle\dfrac {1-a_i }{1 - a_i^{\lambda n}}\geqslant \frac {1-r}{1 -r^{\lambda}}}.$$ For $\lambda\in\mathbb N$, we can write this inequality in a "purely reciprocal" form as$$ {\sum\limits_{i=1}^n\displaystyle\dfrac {1}{1 + a_i + \cdots + a_i^{k-1}}\geqslant \frac {1}{1 + r+ \cdots +r^{\lambda-1}}.}$$ The limit case $\lambda\to 1$ yields $\sum\limits_{i=1}^n\displaystyle\dfrac {1}{1 + a_i + \cdots + a_i^{n-1}}\geqslant 1 \qquad\qquad\ \ \text{ if}\ \ \prod a_i\leqslant 1$ $\sum\limits_{i=1}^n\displaystyle{\dfrac {1}{1 + a_i + \cdots + a_i^{n-1}}} \geqslant \prod a_i^\frac{1-n}n \qquad\text{ if}\ \ \prod a_i\geqslant 1$ The most interesting case of this is of course $\prod a_i=1$. I have proven it for this case for all $n\in\mathbb N$ by calculus, generalizing the (corrected) idea given here. Other special cases that have given rise to my general conjecture above: The case $n=2,k=3$ is equivalent to $ {\dfrac{1}{1+a+a^{2}}+\dfrac{1}{1+b+b^{2}}+\dfrac{1}{1+c+c^{2}}\ge 1 }$, i.e. the case $n=3,\lambda\to1$ with condition $abc=1$. The case $n=2,k=4$, i.e. $ { \dfrac{1}{1+a^2+a^4+a^6}+\dfrac{1}{1+b^2+b^4+b^6}\ge\dfrac{1}{1+a^3b^3} }$, is Vasc's "gm_33 lemma" and is already difficult to prove. There are several proofs of this around, all ad-hoc and rather technical. See http://www.artofproblemsolving.com/Forum/viewtopic.php?t=18857 http://www.artofproblemsolving.com/Forum/viewtopic.php?p=1701829#p1701829 The case $n=2,k=6$ has been suggested in one of those threads as a conjecture. The case $n=3,k=6$ could be called "gm_555 Lemma" and has been suggested by myself. This conjecture says: for $a,b,c\geqslant 0$, $ {\sum\dfrac{1}{1+a^3+a^6+a^9+a^{12}+a^{15}}\ge\dfrac{1}{1+a^5b^5c^5}.}$ One of the striking facts of these inequalities is a certain 'asymmetry' w.r.t. the change $a_i\mapsto\dfrac1{a_i}$. I think that it should be possible, at least for $k\in\mathbb N$, to find proofs similar to the case $n=2,k=4$, but certainly not without the help of a computer. Any ideas on how to tackle this?

论坛问题帖 | no_answers_at_retrieval | Wolfgang | 2013-10-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017979 | MO-145898

On Generalizations of Fermat's Conjecture

We know the following facts: (1) For all $1\leq n\leq 2$ the equation $x_{1}^{n}+x_{2}^{n}=x_{3}^{n}$ has a solution in $\mathbb{N}$. (2) For all $3\leq n$ the equation $x_{1}^{n}+x_{2}^{n}=x_{3}^{n}$ has no solution in $\mathbb{N}$. Question: Is the following generalization true? For all $2\leq m$ both of the following statements are true: (1) For all $1\leq n\leq m$ the equation $x_{1}^{n}+...+x_{m}^{n}=x_{m+1}^{n}$ has a solution in $\mathbb{N}$. (2) For all $m+1\leq n$ the equation $x_{1}^{n}+...+x_{m}^{n}=x_{m+1}^{n}$ has no solution in $\mathbb{N}$.

论坛问题帖 | has_answers | user36136 | 2013-10-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017980 | MO-145974

A Hot Betting On HOD

Remark: This question is based on an open question at the end of a paper by Hamkins, Kirmayer, and Perlmutter: "Generalizations of the Kunen Inconistency". HOD as an inner model of ZFC lies between L and V. But its nature shows a wide degree of flexibility between L -like and V-like behaviors up to different problems. One of the problems about any inner model of ZFC is consistency of existence of a non-trivial self elementary embedding. In this direction the following facts are well known: (a) There is no non-trivial elementary embedding from V to V. (b) Existence of a non-trivial elementary embedding from L to L is an acceptable large cardinal axiom. Here the open question is: Question (1): Is the existence of a non-trivial elementary embedding from HOD to HOD an acceptable large cardinal axiom? I usually discuss with my colleagues about it. I believe that HOD in this problem has a L - like behavior and existence of a non-trivial elementary

embedding from HOD to HOD is an acceptable (not too) large cardinal axiom but one of my friends insists that the facts shows there is no such embedding. Finally we bet on it. Here I want to know about any known partial result around the open problem. So: Question (2): Please introduce any known fact related to the possible answer of question (1) and explain that which one of the scenarios (a) or (b) for HOD seems more possible by this fact.

论坛问题帖 | has_answers | user36136 | 2013-10-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017981 | MO-146800

Forcing with c.c.c forcing notions, Cohen reals and Random reals

I think the following question is due to Prikry: Question. Is it consistent that any non-trivial c.c.c forcing notion adds a Cohen real or a Random real? Is the question still open? What partial results are known about this question (with references, please). Remark. I have the following simple observation which might be well-know. Theorem. Souslin hypothesis (SH) hols iff any non-trivial c.c.c forcing notion adds a new real. Proof. One direction is trivial, since a Souslin tree, considered as a forcing notion, is c.c.c and adds no new reals. For the other direction suppose there is a non-trivial c.c.c forcing notion P which adds no new reals. Let $B=R.O(P)$ be the completion of P. B is a c.c.c. complete Boolean algebra which is (ω, ω)-distributive, hence it is in fact $(\omega, \infty)-$distributive, thus it is a Souslin algebra, which implies the existence of a Souslin tree, and hence SH fails. Update. Can anyone say the first place where the above question has appeared?

论坛问题帖 | no_answers_at_retrieval | Mohammad Golshani | 2013-11-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017982 | MO-146876

A Special Pair of Models for ZFC (New Version)

Are there two models M and N for ZFC such that: (1) $M\subseteq N$ (2) $\aleph_{1}^{N}=\aleph_{1}^{M}$ (3) $\aleph_{2}^{N}=\aleph_{\omega +1}^{M}$ Update: According to Peter's useful answer it seems this problem is still open. So my question is about "partial" or "similar" results with different (but not too different) $\aleph$s. Please give me some references too.

论坛问题帖 | has_answers | user42090 | 2013-11-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017983 | MO-148471

Any results towards the irrationality of the sum of reciprocals of perfect numbers?

This question is a follow up to my comment to Sum of the reciprocal of perfect numbers. I would like to know which results have been published about the possible irrationality of the sum of reciprocals of perfect numbers. For example, do we know a lower bound for the denominator of this number provided it is rational? Thanks in advance.

论坛问题帖 | has_answers | Sylvain JULIEN | 2013-11-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-017984 | MO-149326

Second duals of Grothendieck spaces

The classical example of a Grothendieck space is ℓ_∞. It is also known that its even duals ℓ_∞^{**}, $\ell_\infty^{(4)}$, $\dots$ are Grothendieck spaces. (See, e.g., this note for the proof of the Grothendieck property of these even duals, which uses the fact that these spaces are $C(K)$ spaces for large Stonian K-s.) Is it known, in general, whether the second dual of a Grothendieck space is a Grothendieck space too? Is there a counterexample? Edit (T. Kania). Let me record here for the future reference that this very question was asked by J. Diestel on p. 105 in J. Diestel, "Grothendieck spaces and vector measures", Vector and Operator Valued Measures and Applications, Acad. Press (1973) pp. 97-108.

论坛问题帖 | no_answers_at_retrieval | Aleksei Lissitsin | 2013-11-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017985 | MO-149815

Automorphisms of the L-function associated to an elliptic $\mathbb{Q}$-curve

$\DeclareMathOperator\Aut{Aut}\newcommand{\alg}{\mathrm{alg}}\newcommand{\an}{\mathrm{an}}$Edited after Noam Elkies' comment: From what I understand (very little actually), there exist elliptic curves defined over some number fields $\mathbb{K}$ Galois over $\mathbb{Q}$ which are isogenous to all their Galois conjugates and such elliptic curves are called $\mathbb{Q}$-curves. Since two elliptic curves are isogenous if and only if they share the same L-function, can one define the automorphism group of the L-function corresponding to such an elliptic $\mathbb{Q}$-curve in such a way that this group is actually isomorphic to $\operatorname{Gal}(\mathbb{K}/\mathbb{Q})$? EDIT November 25th 2013: it seems rather clear that the automorphism group of the isogeny class of the $\mathbb{Q}$-curve E contains $\operatorname{Gal}(\mathbb{K}/\mathbb{Q})$ as a subgroup. As the L-function F of E determines its isogeny class entirely, the automorphism groups of F and of the isogeny class of E should be isomorphic. Therefore it remains to show that such a group doesn't contain any element not arising from $\operatorname{Gal}(\mathbb{K}/\mathbb{Q})$. EDIT December 6th 2013: There may be a way to provide a definite answer to my question. Suppose $g_1, g_2, \cdots g_n$ are elements of the considered automorphism group not arising from $\operatorname{Gal}(\mathbb{K}/\mathbb{Q})$ and let G be the group generated by $\operatorname{Gal}(\mathbb{K}/\mathbb{Q})\cup \{g_{1},\cdots g_{n}\}$. Then, provided G is realizable over $\mathbb{Q}$, there should exist a number field $\mathbb{L}$ containing $\mathbb{K}$ as a proper subfield such that the considered $\mathbb{Q}$-curve is actually defined over $\mathbb{L}$. Edit April 1rst 2015: I copy and paste here the content of a cross-post MSE question which, after getting one up vote, has received one down vote et two votes to close without any explanation even though I did my best to improve it. So here it is: "If I'm not mistaken, the isogeny class of E/K contains an abelian group $Z^{r_{\alg}}$ which is the torsion free part of the Mordell-Weil group of E/K. According to the BSD conjecture, the quantities $r_{\alg}$ and $r_{\an}:=\min\{k, L^{(k)}_{E/K}(1)\neq 0\}$ should be equal. What I would like to know is whether the set of sub Z-modules (taken in the increasing order of rank) of $Z^{r_{\alg}}$ can be viewed as

a Galois covering of Z, and equivalently if the set $\{L_{E/K}^{(k)}, L_{E/K}^{(k)}(1)=0\}$ (taken in the increasing order of derivation order) can be viewed as a Galois covering of $L_{E/K}$. If so, are these two Galois coverings isomorphic with Galois group $\operatorname{Gal}(K/Q)$? Of course, the objects considered in the present question are to be defined as sets endowed with a coherent topology to be viewed as genuine topological spaces. Perhaps the Zariski topology would do the job considering $\operatorname{spec} Z$ instead of Z, and the set of local factors of $L_{E/K}$ (i.e those that appear in the Euler product thereof) instead of $L_{E/K}$ itself. The key idea is to somehow identify: the group $\Aut_{B}(X)$ of automorphisms of the covering X of the base B preserving B (where B stands for the topological space corresponding to $L_{E/K}$ and X to the $r_{\an}$ topological spaces corresponding to $L_{E/K}, L'_{E/K},...,L^{(r_{\an}-1)}$ in this order) with $\Aut(L_{E/K})$ which, as showed in the link above, is isomorphic to $\operatorname{Gal}(K/Q)$, the group $\Aut_{B'}(Y)$ of automorphisms of the covering Y of the base B' preserving B' (where B' stands for the topological space corresponding to Z and Y to the $r_{\alg}$ topological spaces corresponding to $Z, Z^2,...,Z^{r_{\alg}}$ in this order) with the automorphism group of the isogeny class of E/K which, as showed in the link above, is isomorphic to $\operatorname{Gal}(K/Q)$." If some people here ever vote to close, please give a good reason to do so. Thanks in advance. Edit May 13th 2016: following http://www.math.wisc.edu/~ellenber/MCAV.pdf, it seems that the automorphism group of a $\mathbb{Q}$-curve is actually isomorphic to $\operatorname{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$. I find that rather interesting since I've been suspecting for long that if one calls 'L-function' any element of the set M generated by the Selberg class and automorphic L-functions closed under both the classical product and the Rankin-Selberg convolution, then the set of automorphisms thereof as the group of bijections from M to itself preserving this double structure should be isomorphic to the absolute Galois group of the rational. Still, I never managed to prove this. Can anyone please help me try to clear this out? Edit September 25th 2022: in Grothendieck's conception of Galois theory, the absolute Galois group of a number field is the automorphism group of a fiber functor, which may help reconcile the two points of view (automorphism group of the $\mathbb{Q}$-curve and of the L-function).

论坛问题帖 | no_answers_at_retrieval | Sylvain JULIEN | 2013-11-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-017986 | MO-150215

Who is attributed with the conjecture that every multiply-perfect number greater than 1 is even?

I know that Descartes is considered to be the first to ask whether or not odd perfect numbers exist (n such that $\sigma(n)=2n$, where $\sigma(n)$ is the sum of divisors of n), and he also discovered several multiply-perfect numbers (n such that $\sigma(n)=kn$ for some integer $k\geq 2$). No odd multiply-perfect numbers greater than one are yet known, and I've heard it called a conjecture that they don't exist several times, but I haven't found an attribution to anyone in particular. Is there an agreed upon origin for the conjecture? The context of my question (please, feel free to skip this): I'm looking for someone to attribute in a paper I'm writing that makes a refinement of this form of conjecture for a generalization of the multiply-perfect numbers. Specifically, let $$\dfrac{\sigma(n)}n=\dfrac{k}m$$ where $\gcd(k,m)=1$ and m is a practical number i.e. $m=1$, or $p_i^{a_i+1}\leq 1+\sigma(p_1^{a_1}p_2^{a_2}...p_i^{a_i})$ for every $i\in[1,\omega(m)]$, where $p_1^{a_1}p_2^{a_2}...p_{\omega(m)}^{a_{\omega(m)}}$ is the canonical prime factorization of m. I call these pad (practical abundancy denominator) numbers, and conjecture that every such number is practical (i.e. a practical number). $\dfrac{\sigma(n)}n$ is called the abundancy of n. It is easy to see that every even perfect number is practical, and every practical number greater than 1 is trivially even, so the conjecture that every perfect number is practical is equivalent to the conjecture that they're all even. From this perspective it's natural to ask whether every multiply-perfect number is practical. I've tested essentially every known multiply-perfect number using the factorizations from Achim Flammenkamp's database and all of them so far are practical, so already this seems like a promising lead. Clearly the multiply-perfect numbers are the special case of the displayed equation where $m=1$, which is trivially practical, so every multiply-perfect number is a pad number. For an idea of their relative size, there are 6484 pad numbers less than 10^6, all of them practical, compared with only 10 multiply-perfect numbers in the same range. An additional fact that makes this conjecture seem quite intuitive is that all small multiples of a practical number are also practical. Specifically, if a is practical and $b\leq \sigma(a)+1$ then ab is practical. Since, k/m is just the abundancy of n in lowest terms, there exists some positive integer r such that $n/m=\sigma(n)/k=r$. When $r=n$, the multiply perfect numbers, we have no idea why every term would be practical, and yet this is the special case I've tested the furthest, and no counterexamples have yet been found. On the other hand, if $r\leq 1+\sigma(m)$ then n is obviously practical. While in retrospective it seems rather intuitive to me, this was a purely numerical observation. Please let me know if you have ideas about this problem, since I believe it has the potential to remain open for a long time, but someone to attribute is what I'm asking for, since right now I feel I have very few people to attribute in my paper. This is for my first paper, which I think I will try to have published in Experimental Mathematics, since computational experiments were the method by which I reached this generalization. If you have strong opinions about other papers or people who should be attributed then please mention them at least in the comments. I'm not looking to downplay anyone's contributions, and I realize the importance of showing how your work ties in with that of others. Edit: Note that the same conjecture appears to hold with $\sigma(m)$ replaced by $\sigma^*(m)=\prod_{i=1}^{\omega(m)}(p_i^{a_i}+1)$, the sum of the unitary divisors of n, which might be simpler to work with.

论坛问题帖 | has_answers | Jaycob Coleman | 2013-11-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017987 | MO-151623

Representing the integers with powers of 2 and 3

Suppose that I have a number of the form $$ x = \frac{1}{3^m}(2^{h} - \sum\limits_{k=1}^{m}3^{m-k}2^{v_k}) $$ where m is a positive integer, and $v_1 = 0 $ $v_{k+1} = v_k +1$ for $1 \leq k < m-1 $ $v_{m} > v_{m-1} $ For a fixed m, can we find a relationship between h and v_m, such that x is an integer ?

论坛问题帖 | has_answers | A. Napster | 2013-12-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017988 | MO-153047

Stability of analytic Zariski structures

Noetherian Zariski structures are introduced by Hrushovski and Zilber(1996). An analytic Zariski structure is a generalization of Noetherian Zariski structures, introduced by Zilber and Peatfield. For more details you can see the following interesting papers: 1) Zariski Geometries, Zilber and Hrushovski. 2) Analytic Zariski structures and the Hrushovski construction, Peatfield and Zilber. Question : Under what conditions an analytic Zariski structure is stable ?

论坛问题帖 | no_answers_at_retrieval | Mostafa Mirabi | 2013-12-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017989 | MO-153426

Floors of rationals to powers: Infinite number of primes?

Let $r=a/b$ be a rational number in lowest terms, larger than 1, and not an integer (so $b > 1$). Q. Does the sequence $$ \lfloor r \rfloor, \lfloor r^2 \rfloor, \lfloor r^3 \rfloor, \ldots, \lfloor r^n \rfloor, \ldots$$ always contain an infinite number of primes? For example, for $r=13/4$, $$(r,r^2,r^3,r^4,r^5, \ldots) = (3.25, 10.56, 34.33, 111.57, 362.59, \ldots)$$ and so the floors are $$(3, 10, 34, 111, 362, \ldots)$$ The next prime after 3 occurs at $$\lfloor r^{34} \rfloor = 2535328703512709 71$$ and I only find four primes up to $\lfloor r^{1000} \rfloor$.

论坛问题帖 | has_answers | Joseph O'Rourke | 2014-01-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017990 | MO-15346

Distribution of the sum of the m smallest values in a sample of size n

Let $\mathbf X = [X_1, X_2, \ldots, X_n]^\mbox{T}$ be a vector random variable drawn from a known distribution with CDF $F(x)$. The CDF for the minimum value in $\mathbf X$ is clearly $P[\min_{i=1\ldots n} X_i \leq x]$ $=$ $1 - \prod_{i=1\ldots n} P[X_i>x]$ $=$ $1 - (1 - F(x))^n$. The expected value for the minimum value in $\mathbf X$ is therefore $\int_0^\infty 1 - (1 - (1 - F(x))^n)dx$. I am interested in the sum of the m smallest values (i.e., order statistics) in $\mathbf X$. It seems like the expected value for the sum should look something like $\sum_{i=0}^{m-1} \int_0^\infty 1 - (1 - (1 - F(x))^{n-i})dx$, however, my intuition tells me that that conflicts with the independence of the events. I therefore have two questions: What is the expected value of the sum of the m smallest values in $\mathbf X$? What is the PDF and/or CDF of the sum? An answer to #2 alone would be sufficient.

论坛问题帖 | has_answers | ESultanik | 2010-02-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-017991 | MO-153824

An open problem in convex geometry

Is it possible to find four norms $\| \cdot\|_k$ $(1 \leq k \leq 4)$ on the plane such that a three-dimensional normed space containing four subspaces isometric to these normed planes does not exist? An equivalent statement: Is it possible to find four 0-symmetric convex bodies in the plane such that a three-dimensional 0-symmetric convex body containing four central sections linearly equivalent to these four bodies does not exist? This seems to be an old problem, but I'm not sure of its origin. The earliest reference I've found is Rolewicz, S., Plane sections of centrally symmetric convex bodies. Israel J. Math. 4 1966 135-138. Does anyone know the status of this problem? Some earlier references? Just for completeness, let me add that Bessaga proved (C. Bessaga,A note on universal Banach spaces of a finite dimension, Bull. Acad. Polon. Sci.6 (1958), 97-101.) that Given $n > 2$, it is possible to find $N = N(n)$ norms $\| \cdot\|_k$ $(1 \leq k \leq N)$ on the plane such that an n-dimensional normed space containing N subspaces isometric to these normed planes does not exist. Rolewicz mentions that it is not clear whether we can take $N(n) = n+1$. Added on 08/01/2014. Trying to guess what possible four normed planes cannot be linearly and isometrically embedded in any one three-dimensional normed space: Does there exist a three-dimensional centrally symmetric convex body that contains four central sections that are linearly equivalent to a circle, a square, a regular hexagon, and a regular octagon?

论坛问题帖 | no_answers_at_retrieval | alvarezpaiva | 2014-01-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017992 | MO-1554

Point singularity of a Riemannian manifold with bounded curvature

Suppose you have an incomplete Riemannian manifold with bounded sectional curvature such that its completion as a metric space is the manifold plus one additional point. Does the Riemannian manifold structure extend across the point singularity? (Penny Smith and I wrote a paper on this many years ago, but we had to assume that no arbitrarily short closed geodesics existed in a neighborhood of the singularity. I was never able to figure out how to get rid of this assumption and still would like someone better at Riemannian geometry than me to explain how. Or show me a counterexample.) EDIT: For simplicity, assume that the dimension of the manifold is greater than 2 and that in any neighborhood of the singularity, there exists a smaller punctured neighborhood of the singularity that is simply connected. In dimension 2, you have to replace this assumption by an appropriate holonomy condition. EDIT 2: Let's make the assumption above simpler and clearer. Assume dimension greater than 2 and that for any r > 0, there exists 0 < r' < r, such that the punctured geodesic ball B(p,r'){p} is simply connected, where p is the singular point. The precludes the possibility of an orbifold singularity. ADDITIONAL COMMENT: My approach to this was to construct a differentiable family of geodesic rays emanating from the singularity. Once I have this, then it is straightforward using Jacobi fields to show that this family must be naturally isomorphic to the standard unit sphere. Then using what Jost and Karcher call "almost linear coordinates", it is easy to construct a C^1 co-ordinate chart on a neighborhood of the singularity. (Read the paper. Nothing in it is hard.) But I was unable to build this family of geodesics without the "no small geodesic loop" assumption. To me this is an overly strong assumption that is

essentially equivalent to assuming in advance that that differentiable family of geodesics exists. So I find our result to be totally unsatisfying. I don't see why this assumption should be necessary, and I still believe there should be an easy way to show this. Or there should be a counterexample. I have to say, however, that I am pretty sure that I did consult one or two pretty distinguished Riemannian geometers and they were not able to provide any useful insight into this.

论坛问题帖 | has_answers | Deane Yang | 2009-10-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-017993 | MO-156076

An infinite, amenable, finitely presentable group with no non-trivial finite quotients

My question is a simple one: is there a group with the properties in the title? In the absence of the 'finitely presentable' hypothesis, an example is provided by Juschenko--Monod's construction of a finitely generated, infinite, simple, amenable group. On the other hand, all the standard examples that I know of finitely presentable groups without finite quotients seem to be non-amenable. There are a number of participants on MO who have constructed interesting examples of amenable groups. My guess is that the answer to this question is unknown, but if so I'd also be interested in informed conjectures. Is it at all probable that such a group doesn't exist?

论坛问题帖 | has_answers | HJRW | 2014-01-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017994 | MO-156371

Conjectures in classical harmonic function theory

Because I'm doing research in the area of harmonic function theory I would like to know are there any conjectures in the theory of harmonic functions in $\mathbb{R}^{n}$ still open. I know that there are many conjectures related to manifolds, but I don't want them. I want conjectures that are just related to harmonic functions on open sets in $\mathbb{R}^{n}$. Are there some really famous ones and long open?

论坛问题帖 | has_answers | Alem | 2014-02-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017995 | MO-157184

Convergence in L^2 of iterated expectations

Take a probability space $(\Omega,\mathcal{F},\mathbf{P})$ and random variable $X \in L^2(\Omega,\mathcal{F},\mathbf{P})$. Define the iterated expectations of X as follows: $X_0 = X$, and, inductively, $X_k = \mathbf{E}[X_{k-1} \, | \, \mathcal{G}_k]$, where $\mathcal{G}_k \subseteq \mathcal{F}$ is some σ-algebra. Assume that all the $\mathcal{G}_k$ come from some finite set of distinct σ-algebras. In words, we are repeatedly taking conditional expectations with respect to various information. Can we conclude that the sequence (X_k) converges to some limit in L^2? The paper [1] says the question seems to be open (as of its publication) and reviews some of the literature, going back to Rota's Alternierende Verfahren. (The contribution of the paper is to show that if we drop the restriction of finitely many distinct σ-algebras, the statement is not true.) Has there been any progress since? [1] M. Akcoglu and J. King. An example of pointwise non-convergence of iterated conditional expectation operators. Israel J. Math. 94 (1996). This paper was pointed out to me by Anthony Quas at a related question -- When do iterated conditional expectations converge?

论坛问题帖 | no_answers_at_retrieval | Ben Golub | 2014-02-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017996 | MO-157800

Ramanujan's problem 754 still open?

In addition to the MO question The Ramanujan Problems. , I would like to ask the following. Problem 754 from the list of the Ramanujan's problems (http://www.imsc.res.in/~rao/ramanujan/collectedpapers/question/q754.htm) asks to show that $$e^xx^{-x}\pi^{-1/2}\Gamma(1+x)=(8x^3+4x^2+x+E)^{1/6},$$ where E lies between $\frac{1}{100}$ and $\frac{1}{30}$ for all positive values of x. According to http://citeseerx.ist.psu.edu/viewdoc/summary?doi=10.1.1.39.9228 (The Problems Submitted by Ramanujan to the Journal of the Indian Mathematical Society, by Bruce C. Berndt, Youn--seo Choi and Soon-Yi Kang) this problem was still open in 1997. What is the status of this problem at present?

论坛问题帖 | has_answers | Zurab Silagadze | 2014-02-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017997 | MO-157886

Real analytic ergodic diffeomorphisms of the two sphere

Does there exists a real analytic area preserving ergodic diffeomorphism on S^2? (Possibly by perturbing a rotation in the real-analytic topology?)

论坛问题帖 | has_answers | user42388 | 2014-02-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-017998 | MO-158022

An integral related to the Euler gamma function

The question is from the paper http://arxiv.org/abs/1312.7115 (A curious formula related to the Euler Gamma function, by Bakir Farhi): is it possible to express the integral $$\eta=2\int\limits_0^1 \ln{(\Gamma(x))}\cdot \sin{(2\pi x)}\,dx= 0.7687478924\ldots$$ in terms of the known mathematical constants as $\pi,\,e,\,\gamma,\;\ln{\pi},\, \ln{2},\,\Gamma{(1/4)}\ldots$? It is shown in the paper that $$\frac{\ln{1}}{1}-\frac{\ln{3}}{3}+\frac{\ln{5}}{5}-\ldots=\pi\ln{(\Gamma{(1/4)})}-\frac{\pi^2}{4}\eta-\frac{\pi}{2}\ln{\pi}-\frac{\pi}{4}\ln{2}.$$ So the question actually asks whether one can give a closed-form expression for the series in the l.h.s. in terms of the known mathematical constants.

论坛问题帖 | has_answers | Zurab Silagadze | 2014-02-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-017999 | MO-158039

Objections to and arguments for the simplicity of all Riemann zeros

It seems to be that the simplicity of all the zeros is quite widely accepted as a working hypotheses, and it is known that a positive proportion are as such. Titchmarsh explains in the last chapter on RH that the weak Mertens hypothesis, $M(x)=O(x^{1/2})$, implies that all zeros are simple. Though I think this is conjectured to be false by many authors. Is there a weaker bound whose optimality is known to imply that not all zeros are simple? Given the classical estimate $N(\sigma, T)=O(T^{4\sigma(1-\sigma)+\epsilon})$ combined with the fact that the number of solutions to $\zeta(s)=z$ in any rectangle of height T contained in $1/2\leq \sigma\leq 1$ is proportional to T, for any $z\neq 0$, simplicity of the zeros suggests that the derivative must vanish vastly more frequently in each such rectangle else one could presumably tie the number of such solutions to the number of zeros up to height T, for all sufficiently small $|z|$, by Rouche's theorem. One might therefore conjecture that $ \inf|\rho_{\zeta}-\rho_{\zeta'}|=0$ (is this known perhaps?). An old theorem of MacDonald (Whittaker and Watson, p131) says that the number of zeros of an analytic function on the interior of a simple closed curve on which the modulus of f is constant exceeds that of it's derivative by one. It can also be seen that the derivative necessarily vanishes on the largest such curve by the open and inverse mapping theorems. In the case of $\zeta(s)$ I'd expect that the largest of such curves around each zero would become small for large T, so I wonder if something in this direction is known? Anyway, as Titchmarsh points out, Ω-theorems for $M(x)$ seem to be more difficult to obtain than those for the prime counting functions and, since a significant distinction between the problems is that of the order of the zeros, I would like to enquire specifically about anything that is known in either direction, on any other hypotheses.

论坛问题帖 | has_answers | Kevin Smith | 2014-02-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018000 | MO-159494

Determine or estimate the number of maximal triangle-free graphs on n vertices

Among the collections of the open problems of Paul Erdős on the website of Professor Fan Chung, there is one called "number of triangle-free graphs". http://www.math.ucsd.edu/~erdosproblems/erdos/newproblems/NumberOfTriangleFreeGraphs.html Open Problem: Determine or estimate the number of maximal triangle-free graphs on n vertices. Is any one working on this problem? Any known related results? Now I am considering about using a "connection game" method to solve this problem: Given a set of n players, each one chooses to connect to other nodes, if any two neighbors of it connect a new edge(which means there would form a triangle), it has to choose delete either edges with those two neighbors. Then the question is how many different topology connections does it have? Any comments on this method?

论坛问题帖 | has_answers | user39815 | 2014-03-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018001 | MO-161731

practical algorithms for np complete problems

Inspired by: Conjecture on NP-completeness of tesselation of Wang Tile up to finite size And the practicality of this topic (solving tessellation on a lattice): coloring in lattice Computational approach deciding whether a set of Wang Tile could tile the space up to some size Reference for Wang Tile I want to ask for practitioner what kind of software/programmes/algorithms that are usually used to tackle NP complete problems?

论坛问题帖 | has_answers | user40780 | 2014-03-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018002 | MO-162020

Which region in the plane with a given area has the most domino tilings?

I just finished teaching a class in combinatorics in which I included a fairly easy upper bound on the number of domino tilings of a region in the plane as a function of its area. So this led to this question, which I suspect is an open problem: Does a $2n \times 2n$ square have the most domino tilings among all regions with area $4n^2$? I aired this question in a mailing list, which is one reason that I think that it's an open problem. The discussion there did lead to a great answer from Rick Kenyon, who says that using his methods, the conjecture is asymptotically true for dilated rectilinear regions. (In other words, if you take any fixed polyomino and dilate it by a sufficiently large factor, then it will have fewer tilings than a square with the same area or very slightly less area.) Still, I just came up with the question in my class, so I don't really know if it's open or previously stated. (This is my excuse for posting it MO.) I Googled some, without success. It is also easy to pose generalizations: (1) The inequality might also hold for regions that map to the plane with multiple sheets (immersions) or branch points, at least when such regions are topological disks and the branch points are at vertices of the tiling by unit squares. (Kenyon's methods might also apply to these cases.) (2) Does the d-dimensional cube $[0,2n]^d$ have the most tilings by domino bricks, among all regions in $\mathbb{R}^d$ with volume $(2n)^d$? (3) Given an integer $k > 2$, does a $kn \times kn$ square have the most tilings by $k \times 1$ rectangles, among regions with area k^2n^2? (4) Mutual generalizations of (1)-(3), and tilings by a rectangular brick with some other shape. Kenyon's work is a (deep) extension of the permanent-determinant method of Kasteleyn and Percus to count matchings of a planar graph, so it would not apply to cases (2) and (3).

论坛问题帖 | has_answers | Greg Kuperberg | 2014-04-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018003 | MO-16214

Is there an associative metric on the non-negative reals?

Recall that a function $f\colon X\times X \to \mathbb{R}_{\ge 0}$ is a metric if it satisfies: definiteness: $f(x,y) = 0$ iff $x=y$, symmetry: $f(x,y)=f(y,x)$, and the triangle inequality: $f(x,y) \le f(x,z) + f(z,y)$. A function $f\colon X\times X \to X$ is associative if it satisfies: associativity: $f(x,f(y,z)) = f(f(x,y),z)$. If $X=\mathbb{R}_{\ge 0}$, then it might be possible for the same function to be a metric and associative. Is there an associative metric on the

non-negative reals? Note that these demands actually make X into a group. The element 0 is the identity because $f(f(0,x),x) = f(0,f(x,x)) = f(0,0) = 0$ by associativity and definiteness, so again by definiteness $f(0,x) = x$. Every element is its own inverse because $f(x,x)=0$. In fact, the following question is equivalent. Is there an abelian group on the non-negative reals such that the group operation satisfies the triangle inequality? Note also that the answer is yes if $X=\mathbb{N}$, the non-negative numbers! Click here for a spoiler. "Also known as nim-sum." The question is originally due to John H. Conway. To my knowledge, the question is unsolved even for $X = \mathbb{Q}_{\ge 0}$, but he does not seem to care about that case. The spoiler above does extend to the non-negative dyadic rationals $\mathbb{N}[\frac 12]$, but apparently not to $\mathbb{N}[\frac 13]$.

论坛问题帖 | has_answers | user1079 | 2010-02-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018004 | MO-162245

Conjecture of a subset of Wang tile which might be decidable

From the two papers proving the undecidability of Wang tile in 1966 by Berger and in 1971 by RM Robinson, the tiles used in proving undecidability has a general common feature: The left color and right color do not uniquely determine the top color and bottom color. The tiles involved in practice in material science and physics for one big problem is actually more restricted and the tiling problem might actually turn out to be decidable. So the subset of Wang Tile is lattice tiles, shown in my previous post: Reference for Wang Tile The general feature of the tile is that the pair (left color, right color) uniquely determine the pair (top color, bottom color). I conjecture that a subset of Wang tiles which satisfies the restriction above might turn out to be decidable... Any thought on this problem? Thank you.

论坛问题帖 | has_answers | user40780 | 2014-04-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018005 | MO-163301

What is Chern-Simons theory expected to assign to a point?

Let G be a compact, connected, (simply connected?) Lie group and let $k \in H^4(BG, \mathbb{Z})$ be a cohomology class. Witten showed, at a physical level of rigor, that this data determines a 3-dimensional topological quantum field theory (going down to surfaces), Chern-Simons theory. Later other authors (Reshetikhin-Turaev, ???) described how to extend this theory down to 1-manifolds. It's known that such a theory is determined by what it assigns to a circle S^1, which must be a modular tensor category; this category can be described either as a certain category of representations of the loop group LG at level k or as a certain category of representations of the quantum group $U_q(\mathfrak{g})$, where q is a suitable function of k. The relationship between these two descriptions is unclear to me. My impression is that it's expected that Chern-Simons theory extends all the way down to 0-manifolds; that is, that it is a fully extended TQFT. By Lurie's classification, such a theory is completely determined by what it assigns to a point, which is a fully dualizable object in a suitable 3-category. What are some conjectural descriptions of this object? The nLab is somewhat vague on this subject. Here's what I know: The corresponding object for 3-dimensional Dijkgraaf-Witten theory is known, although I'm not sure exactly who this is due to. Here G is replaced by a finite group and k is thought of as a class in $H^3(BG, \text{U}(1))$. k is used to twist the associator on the monoidal category of G-graded vector spaces, giving a monoidal category (in fact a fusion category) of "twisted G-graded vector spaces," to be thought of as a fully dualizable object in the 3-category of monoidal categories and bimodule bicategories over these (Douglas-Schommer-Pries-Snyder?), and I think this is what fully extended 3-dimensional Dijkgraaf-Witten assigns to a point. Freed-Hopkins-Lurie-Teleman generalized this construction to the case that G is a torus; here G-graded vector spaces are replaced by skyscraper sheaves on G. I don't know if this is expected to generalize.

论坛问题帖 | has_answers | Qiaochu Yuan | 2014-04-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018006 | MO-16335

A generalization of Cauchy's mean value theorem.

The following simple theorem is known as Cauchy's mean value theorem. Let γ be an immersion of the segment $[0,1]$ into the plane such that $\gamma(0) \ne \gamma(1)$. Then there exists a point such that the tangent line at that point is parallel to the line passing through $\gamma(0)$ and $\gamma(1)$. So the boundary values of an immersion determine a direction such that for any immersion of a segment with given boundary values there exists a tangent line parallel to the direction. I would like to propose the following conjecture, generalizing this statement. Instead of immersions of a segment we consider immersions of a compact manifold M^n with non-empty boundary ∂M into ${\mathbf R}^{n+1}$. For a map $f\colon \partial M \to {\mathbf R}^{n+1}$ we consider the space $L(f)$ of all immersions $g\colon M \to {\mathbf R}^{n+1}$ such that $g|_{\partial M}=f$. The conjectural claim is the following: If f is sufficiently generic, then for every connected component L_0 of the space $L(f)$ there exists a hyperplane direction l such that for any immersion g from L_0 l is parallel to the tangent plane to $g(M)$ at some point. If the conjecture is true then it is very interesting how l depends on g.

论坛问题帖 | has_answers | Petya | 2010-02-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018007 | MO-16341

On Polynomials dividing Exponentials

EDIT, May 2015: in the second edition of the relevant book, the question was corrected, a single number had been mis-typed. The corrected question (thanks to Max) is to find all positive integer pairs such that $$ m^2 - 1 \; | \; 3^m + \left(n! - 2 \right)^m. $$ $$ $$ The version that caused me all that misery was $ m^2 - 1 \; | \; 3^m + \left(n! - 1 \right)^m. $ EDIT, 2010: it turns out that no answer to this is known, as the authors of the book it is in have now confirmed they do not know how to do it. Will Jagy. ORIGINAL: I have been wondering if there exist some general techniques to attack problems in which a polynomial divides and exponential equation. The motivation of this question came when trying to solve the following problem: Find all positive integers "m such that m^2-1 divides 3^m+5^m I haven't been able to come with a proof, and I am really considering the possibility that it cannot be solved using "elementary" methods. I would really apprecciate

some references (if there are any) to the general question as well as the solution to this very particular case.

论坛问题帖 | has_answers | Daniel Kohen | 2010-02-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018008 | MO-163845

Integral straight-line embeddings of planar graphs

Wikipedia says (in the article on Fáry's theorem), "Heiko Harborth raised the question of whether every planar graph has a straight line representation in which all edge lengths are integers. The answer remains unknown as of 2009." The reference is this: Kemnitz, Arnfried, and Heiko Harborth. "Plane integral drawings of planar graphs." Discrete Mathematics 236.1 (2001): 191-195. (Elsevier link) It is known that every n-vertex planar graph has a planar straight-line grid drawing on an $O(n) \times O(n)$ grid. But of course integer vertex coordinates do not imply integer edge lengths! The Kemnitz-Harborth article concludes with this sentence: "It seems to be a very hard problem to prove that every pentagon with sides of integral length contains an inner point with rational distances to its vertices." Here are my questions: Q. Have there been advances on this problem in the last five years? Perhaps for special cases beyond cubic graphs? Any progress on the specific pentagon problem identified above? Addendum. User fidbc cited two recent papers exhibiting progress: Biedl, Therese C. "Drawing some planar graphs with integer edge-lengths." CCCG. 2011. She shows that planar bipartite, series-parallel graphs, graphs with aboricity 2, outerplanar graphs, and-most interestingly-that every planar graph of degree at most four, but which is not 4-regular, has an integral embedding. Sun, Timothy. "Drawing some 4-regular planar graphs with integer edge lengths." CCCG. 2013. Timothy extends Therese's work and captures various families of (but not all) 4-regular planar graphs. Finally, TMA's mentions the four-square-corners problem, which is described in this MathWorld article, which says, "J.H. Conway and R. Guy found an infinite numbers of solutions to the problem of three such distances being integers."

论坛问题帖 | has_answers | Joseph O'Rourke | 2014-04-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018009 | MO-16410

$\operatorname{SL}_2(\mathbb R)$ Casson invariant?

$\DeclareMathOperator\SL{SL}$Casson's invariant is an invariant of a homology 3-sphere, obtained by “counting” representations of the fundamental group into $\operatorname{SU}(2)$. I was wondering if there is an analogous invariant counting representations into $\SL(2,\mathbb R)$? Curtis has an invariant counting representations into $\SL(2,\mathbb C)$. These invariants are obtained by taking a Heegaard splitting of the manifold, and considering the intersection of the representation varieties of the two handlebodies in the representation variety of the Heegaard surface. Casson has to perturb the resulting varieties to make them transverse, then counts the intersections. Curtis counts only the finite points of intersection using algebraic geometry to resolve any singularities, and ignoring any higher dimensional components of the intersection. Then they both have to show that this count is invariant under stabilization of Heegaard splittings, and therefore an invariant of the manifold. I was wondering whether one could combine the two approaches to get an analogous invariant in the case of $\SL(2,\mathbb R)$ representations? One would throw away higher dimensional components of intersection of the $\SL(2,\mathbb R)$ varieties of the two handlebodies, and perturb near the isolated intersections to get a count of intersection points. If this works, what about making an analogous Floer theory, by counting holomorphic disks between finite intersection points? I have't done a literature search, but I suspect this is an open question.

论坛问题帖 | has_answers | Ian Agol | 2010-02-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018010 | MO-164400

relationship between corner tile and edge tile of wang tile

It is clear that any corner type of Wang Tile could be converted to edge type of Wang Tile by defining the edge color according to the corner color. However, could we convert edge type of Wang Tile to corner tile so that each tiling of the corner tiles correspond to a (unique) tiling of the edge type of Wang tile?

论坛问题帖 | has_answers | user40780 | 2014-04-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018011 | MO-164665

Is there a precise notion of "almost all" such that almost all finite groups are Galois groups of extensions of the rationals?

I vainly tried to define a notion of "almost solvable group" such that every "almost solvable group" is the Galois group of a finite extension of the rationals, but I can't figure out the right way to do so, so I want to ask a probably less ambitious question. Is there a precise notion of "almost all" such that almost all finite groups are Galois groups of finite extensions of the rationals? If so, what is the considered notion exactly? Thanks in advance.

论坛问题帖 | has_answers | Sylvain JULIEN | 2014-04-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018012 | MO-164756

simple cycle analog in 2D (with application in tiling)

We know that any closed cycle of a graph could be decomposed into sum of simple cycles. To translate this theorem into tiling of 1D (Wang tile). We know that any 1D periodic tiling could be represented as a sum of basis tiling (correspondingly simple cycles). However, in 2D, the tiling problem could be represented as a graph with 2 different kinds of edges: Graph theoretical representation of Wang Tile Question: For any given graph with 2 kinds directed edges, is there some sort of simple 2D cycles? Such that any 2D cycles could be decomposed into sums of basis cycles?

论坛问题帖 | has_answers | user40780 | 2014-04-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018013 | MO-164983

Generating random finite groups

I would like a method to efficiently generate a random finite group of a given order n. If there are $g(n)$ non-isomorphic groups of order n, ideally each group would occur with probability $1/g(n)$. So if $n=64$, each of the 267 groups would be generated with equal probability. ($g(n)$ is A000001 in OEIS.) Groups of order $n=2^k$ would be of special interest. This is far from my expertise, and my searches must be using the wrong terminology, because I have not found such methods. I'd appreciate pointers-Thanks! Addendum. The comments indicate that this appears to be an open problem, with little chance of resolution in the near future. Now so tagged.

论坛问题帖 | has_answers | Joseph O'Rourke | 2014-05-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018014 | MO-168154

Rost Correspondence and minimal Pfister-Neighbors

In http://www.math.uiuc.edu/K-theory/0357/ Karpenko utters the following: Conjecture 1.6. If an anisotropic quadric $X = Q$ possesses a Rost correspondence, then the quadratic form (defining X) is a minimal Pfister neighbor. Since the Paper has been released 15 years ago, I wonder if there has been made any progress on the conjecture. Everything regarding Rost-Projectors is defined within the paper. I am quite new to the subject and I am "working" myself through most of the papers in order of their release, so it might be that my question is obsolete today. On the other hand, there is also a remark in "Rost Projectors and Steenrod Operations" by Karpenko and Merkurjev from 2002 that the problem is still not solved, except for the cases of $dim(X) = 3,7$.

论坛问题帖 | has_answers | nxir | 2014-05-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018015 | MO-16818

Are there Ricci-flat riemannian manifolds with generic holonomy?

This may well be an open problem, I'm not sure. In Berger's classification (refined by Simons, Alekseevsky, Bryant,...) of the holonomy representations of irreducible non-symmetric complete simply-connected riemannian manifolds, there are some cases which imply Ricci-flatness: namely, $\mathrm{SU}(n)$ (Calabi-Yau) in dimension $2n$, $\mathrm{Sp}(n)$ (hyperkähler) in dimension $4n$, G_2 in dimension 7 and $\mathrm{Spin}(7)$ in dimension 8. A natural question is the converse: whether Ricci-flatness implies a reduction of the holonomy. The other holonomy representations are known not to be Ricci-flat: $\mathrm{Sp}(n)\cdot \mathrm{Sp}(1)$ (quaternionic kähler) is known to be Einstein with nonzero scalar curvature, and in the case of $\mathrm{U}(n)$ in dimension $2n$ (Kähler) it is known that if a Kähler manifold is Ricci-flat then the holonomy is contained in $\mathrm{SU}(n)$, so that it is Calabi-Yau. So the remaining question is whether there exists any Ricci-flat riemannian manifolds with generic holonomy $\mathrm{SO}(n)$ in dimension n. I would like to know the present status of this question and if it's still open what the experts think: do people expect examples of Ricci-flat riemannian manifolds with generic holonomy? Bonus question: How about if the manifold is pseudoriemannian? Added Thanks to Igor's answer below, here are some further remarks. The question needs to be refined. The riemannian analogue of the Schwarzschild metric on $\mathbb{R}^2 \times S^2$ is an example of a complete, simply-connected noncompact Ricci-flat metric with generic holonomy. So the question is about compact examples. In fact, in Berger's 2003 book A panoramic view of Riemannian geometry (page 645) one reads at the bottom of the page: It remains a great mystery that no Ricci flat compact manifolds are known which do not have one of these special holonomy groups.

论坛问题帖 | has_answers | José Figueroa-O'Farrill | 2010-03-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018016 | MO-17058

Factorials in Pascal's triangle

I asked this question of Keith Conrad, and he suggested that I try posting here. One of my students observed that the only instances of factorials in the interior of Pascal's triangle are $\binom{4}{2}=3!$ and $\binom{10}{3}=\binom{16}{2}=5!$. I checked the first 500 rows, and he's right up to that point. This is a special case of the apparently unsolved problem of finding non-trivial solutions to $n!=a!b!c!\dots$ The special feature here is that I need $(a+b)!=a!b!c!$, and that seems like a special enough case to have been treated by someone. Unfortunately, literature searches have been fruitless, because every paper about Pascal's triangle contains the word "factorial" somewhere. My best idea (which I can't make work) is to show that the powers of 7 in the equation $(a+b)!=a!b!c!$ can't be made to match up unless neither side is a multiple of 7. Then exhaustive searching can show that the above are the only non-trivial solutions. Thanks greatly for any ideas that anyone might have.

论坛问题帖 | has_answers | Rob Gross | 2010-03-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018017 | MO-171803

Packing space by cones: Translates best?

Let C be a right circular cone, the convex hull of a unit-radius disk and a point directly above the disk center at height h. Is the most dense packing of $\mathbb{R}^3$ achieved by translates of C and its negative,* $-C$, i.e., upside-down C? Or is it known that rotations of the cones permit achieving a denser packing? * Barany, Imre, and Jiri Matousek. "Packing cones and their negatives in space." Discrete & Computational Geometry 38.2 (2007): 177-187. (Springer link) Here is TMA's idea: (27Jun14). Conjecture by Wlodek Kuperberg (see the comments), the expert on the topic: "For a generic cone, the densest packing by translates of the cone and its point-reflection is perhaps the best one." Now marked as an open problem.

论坛问题帖 | no_answers_at_retrieval | Joseph O'Rourke | 2014-06-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018018 | MO-172626

Is there a non-zero ghost map between finite suspension spectra?

A morphism $f\colon X\to Y$ of spectra such that for every integer n the induced map $\pi_n(f)\colon\pi_n(X)\to\pi_n(Y)$ on stable homotopy groups is zero is called a ghost map. Not every ghost map f is the zero object of the abelian group $\operatorname{Hom}_{SH}(X,Y)$ (more precisely, its image $\bar f\in \operatorname{Hom}_{SH}(X,Y)$ in the stable homotopy category SH). Suppose $X=\Sigma^\infty A$ and $Y=\Sigma^\infty B$ are the suspension spectra of finite CW-complexes, let $f=\Sigma^\infty f'$ for a map $f'\colon A'\to B'$ and suppose that f is a ghost map. Is f necessarily zero in this case?

论坛问题帖 | has_answers | user8463524 | 2014-06-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018019 | MO-17313

Is it possible to dissect a disk into congruent pieces, so that a neighborhood of the origin is contained within a single piece?

Problem: is it possible to dissect the interior of a circle into a finite number of congruent pieces (mirror images are fine) such that some neighbourhood of the origin is contained in just one of the pieces? It may be conceivable that there is some dissection into immeasurable sets that does this. So a possible additional constraint would be that the pieces are connected, or at least the union of connected spaces. A weaker statement, also unresolved : is it possible to dissect a circle into congruent pieces such that a union of some of the pieces is a connected neighbourhood of the origin that contains no points of the boundary of the circle? This is doing the rounds amongst the grads in my department. So far no one has had anything particularly enlightening to say - a proof/counterexample of any of these statements, or any other partial result in the right direction would be much obliged! Edit: Kevin Buzzard points out in the comments that this is listed as an open problem in Croft, Falconer, and Guy's Unsolved Problems in Geometry (see the bottom of page 87).

论坛问题帖 | has_answers | sobe86 | 2010-03-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018020 | MO-173446

Subspaces of l_{1} are not Lipschitz complemented in l_{1}

I have thought about the following question for several years. This question may be stupid or not interesting. My question is: Is there a subspace U of $l{_1}$ such that the quotient l_{1}/U is isomorphic to $L_{1}(\mu)$ for some measure μ, but U is not Lipschitz complemented in l_{1}.

论坛问题帖 | has_answers | Dongyang Chen | 2014-07-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018021 | MO-173510

Arithmetic progression and average of two prime numbers

Let $A=(a_n : n \in \mathbb{N})$ be the sequence given by: $$ \ a_n = a_1 + (n - 1)d,\quad a_1,\ d,\ n \in \mathbb N,\quad d\gt a_1,\quad \gcd(a_1,\ d)=1. $$ For all terms of A greater than $\ \varphi(d) \cdot d^2 \ $ either is a prime number or is the average of two primes which are terms of A. Are there similar conjectures to this conjecture? Source: http://www.mitbbs.com/article_t/Mathematics/31205723.html

论坛问题帖 | no_answers_at_retrieval | Mike | 2014-07-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018022 | MO-17532

Does linearization of categories reflect isomorphism?

Given a category C and a commutative ring R, denote by RC the R-linearization: this is the category enriched over R-modules which has the same objects as C, but the morphism module between two objects x and y is the free R-module on $\operatorname{Hom}_C(x,y)$. Thus in RC we allow arbitrary R-linear combinations of morphisms from the original category C. Question: if two objects in x, $y \in C$ are isomorphic in RC, are they already isomorphic in C? I do not know the answer to this question for any nontrivial ring R, but I'm particularly interested in $R=\mathbb{Z}$ and $R=\mathbb{Z}/2\mathbb{Z}$. What's obviously not true is that every isomorphism in RC comes from an isomorphism in C (take $-id_x$). (Thus the word "isomorphism" in the title refers to a relation on objects rather than to a property of morphisms.) Of course, it is enough to consider categories C with two objects x, y, but we cannot assume that C is finite. It's fairly elementary to see that if x and y are isomorphic in RC then in C, x is a retract of y and vice versa, but the latter does in general not imply that $x \cong y$. A more catchy way of phrasing this problem is: can we always classify objects in a category up to isomorphism by means of functors taking values in R-linear categories? (The inclusion $C \to RC$ is the universal such functor.) Edit: A lot of people have posted an "answer" that wasn't, and deleted it, so here's something that will not work, to save others going down the same road. I said that we cannot assume that the category is finite; in fact, it must be infinite. Here is an elementary argument: Since x and y are mutual retracts, there are maps $f,\;f'\colon x \to y$ and $g,\;g'\colon y \to x$ with $fg=\operatorname{id}$ and $g'f'=id$. Consider the powers of $fg' \in \operatorname{End}(y)$. If $\operatorname{End}(y)$ is finite then $(fg')^n = (fg')^m$ for some $m \neq n$; since fg' has a right inverse (viz, $f'g$), we must have that $(fg')^n=\operatorname{id}$ for some $n>0$. So we see that g' has not only a right inverse (f') but also a left inverse: $(fg')^{n-1}f$. So they are the same and g' is already an isomorphism.

论坛问题帖 | has_answers | Tilman | 2010-03-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018023 | MO-17560

If $2^x $and 3^x are integers, must x be as well?

I'm fascinated by this open problem (if it is indeed still that) and every few years I try to check up on its status. Some background: Let x be a positive real number. If n^x is an integer for every $n \in \mathbb{N}$ then x must be an integer. This is a fun little puzzle. If 2^x, 3^x and 5^x are integers then x must be an integer. This requires fairly sophisticated tools, and can be derived from the results in e.g. Lang, Algebraic values of meromorphic functions. II., Topology 5, 1966. Finally, if all you know is that 2^x and 3^x are integers, then as far as I know it is not known if x is forced to be an integer (unbelievable, isn't it?). Although of course one can never be certain, I am quite sure this was still the case as recently as 2003. So the question is, is that still an open problem, and is there any sort of relevant progress that may provide some hope?

论坛问题帖 | has_answers | Alon Amit | 2010-03-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018024 | MO-176219

If there are two primes at an even distance k is there another disjoint pair of primes also at distance k?

Suppose p,q are two primes at even distance k. Must there necessarily exist a different pair p',q' composed of entirely different numbers such that p' and q' are also at distance k? Edit: None of the primes need to be consecutive.

论坛问题帖 | has_answers | Gorka | 2014-07-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018025 | MO-179138

A possible minimal aperiodic set of corner Wang Tile

From one of my previous question Aperiodic set of corner Wang Tile (although it is put on hold), I realize there is a systematic way to construct aperiodic corner type of Wang tile from edge type aperiodic Wang Tiles. But what I am more interested in is this: For edge type of Wang tile, there is common knowledge of a very small set of Wang tile with 13 elements: http://www.sciencedirect.com/science/article/pii/S0012365X96001185 (doi: 10.1016/S0012-365X(96)00118-5 or with 14 elements: http://www.sciencedirect.com/science/article/pii/0012365X9500120L (doi: 10.1016/0012-365X(95)00120-L It seems there is a competition on this arena of achieving smaller set of aperiodic edge type of Wang tiles. On the other hand, I interested on a documented small set of corner Wang tiles. Do someone know some reference on this? Thank you:)

论坛问题帖 | has_answers | user40780 | 2014-08-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018026 | MO-179206

reduction to np hard ordering problem

I am trying to show a reduction from a problem of ordering problem to an np-hard problem that has approximation poly-time algorithm. My problem is: I have M auctions and in each auction I have N bidders. A bidder can bid in all the auctions until he wins one, but if he won one of the auctions he can't win other items. assuming that the auctioneer that plans the auctions has information about all the values that the bidders are willing to bid on all the items before the auctions, he needs to find an ordering that maximizes his profit - meaning the sum he get from all the auctions. I tried finding an NP-hard problem that I can reduct this problem to, but with no succes. I thought about max-weighted-IS problem by setting the vertices as : Xij - the value that bidder i is willing to bid on item j, and then add edges between all the vertices that have the same j-th value and between all the vertices that have the same i-th value, thus promising that the vertices in the independent set will have preserve the constraing of the auctio, but the problem in this reduction is that given an max weighted indepndent set the max ordering is not neccesserily optimal. Any help would be appriciated. Thanks

论坛问题帖 | has_answers | jhon | 2014-08-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018027 | MO-179790

Is the fixed point property for posets preserved by products?

Recall that a partially ordered set (poset) P has the fixed point property (FPP) if any order preserving function $f:P\longrightarrow P$ has a fixed point. Theorem. Suppose P and Q are posets with the FPP and at least one of them is finite. Then $(P\times Q)$ has FPP. Note: $(a,b)\le(c,d)$ if and only if $a\le c$ and $b\le d$. Question. Suppose P and Q are two infinite posets with the FPP. Does $(P\times Q)$ have the FPP ?

论坛问题帖 | has_answers | Mostafa Mirabi | 2014-08-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018028 | MO-179898

Does S^4 have a "symplecto-homeomorphic" structure?

The 4-sphere cannot be a symplectic manifold. In particular, it does not admit an atlas whose transition maps are symplectomorphisms $(\mathbb{R}^4,\omega_\text{std})\to(\mathbb{R}^4,\omega_\text{std})$. Eliashberg-Gromov proved that if the C^0-limit of symplectomorphisms is a diffeomorphism, then it is a symplectomorphism. We can now ask about the next best thing: Does S^4 have a symplectohomeomorphic structure, i.e. are there charts (equipped with the standard symplectic form on $\mathbb{R}^4$) whose transition maps are homeomorphisms that are C^0-limits of symplectomorphisms? (This question may have originated from Alan Weinstein, who told me it after conversations concerning symplectic rigidity.)

论坛问题帖 | no_answers_at_retrieval | Chris Gerig | 2014-09-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018029 | MO-181853

Non-negative, monotone polynomial sequences without combinatorial interpretation

I am wondering what sequences of integers there are, that are known to grow polynomially, are non-negative, monotone but lacks a combinatorial interpretation. By combinatorial interpretation, they are given by counting some discrete class of objects, (say lattice points in a polytope, Young tableaux, lattice paths, etc). For example, the function $k \mapsto c^{k\lambda}_{k\mu,k\nu}$ where c is the Littlewood-Richardson coefficient, λ,μ,ν are partitions, and $k\geq0$ is an integer, is a sequence that satisfies all above properties, but it has a combinatorial description. Replace c with the more general Kronecker coefficients, and we obtain a sequence of integers that satisfy all listed properties, and it is one of the big unsolved problems in combinatorial representation theory to give a combinatorial description of the Kronecker coefficients. Another non-example is the super-Catalan numbers, $$C(m,n) = \frac{(2m)!(2n)!}{(m+n)!m!n!}$$ which lacks a combinatorial description in general, and is non-negative, integer and monotone for each fixed m. However, the growth is not polynomial (the case $m=1$ corresponds to 2 times the usual Catalan numbers). Of course, there are many sequences of numbers above, so to clarify, I mean sequences that belong to some natural family, parametrized by some auxiliary parameter(s), for example partitions, as above.

论坛问题帖 | no_answers_at_retrieval | Per Alexandersson | 2014-09-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018030 | MO-182

Is there a finite-index finite-depth II$_1$ subfactor which is more than 7-super-transitive?

Background: See Noah and Emily's posts on subfactors and planar algebras on the Secret Blogging Seminar. There are plenty of examples of 3-super-transitive (3-ST) subfactors; Haagerup, $S_4 < S_5$, and others. There's exactly one known example of a 5-ST subfactor, the Haagerup-Asaeda subfactor, and one 7-ST subfactor, the extended Haagerup subfactor. Below index 4 there are the A_n and D_n families, which are arbitrarily super-transitive. Ignore those; I'm just interested above index 4. Is there anything that's even more super-transitive?

论坛问题帖 | no_answers_at_retrieval | Kim Morrison | 2009-10-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018031 | MO-18264

ℓ^2, ℓ^3, and ℓ^4: a possible fixed point result?

The question below is related to the classical Browder-Goehde-Kirk fixed point theorem. Let K be the closed unit ball of ℓ^{2}, and let $T:K\rightarrow K$ be a mapping such that $$\Vert Tx-Ty\Vert _{\ell^{4}}\leq\Vert x-y\Vert _{\ell^{3}}$$ for all $x,y\in K$. Is it true that T has fixed points?

论坛问题帖 | has_answers | Ady | 2010-03-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018032 | MO-184611

Gauss Codes that produce classical knots as opposed to virtual knots

I have been doing some research in Gauss codes and have been reading Kauffman's paper Virtual Knot Theory. In section 3.3, Theorem 2, he states that If K is a virtual knot whose underlying Gauss code is planar and whose sign sequence is standard, then K is equivalent to a classical knot. He then goes on to say that "The fundamental problem in Gauss codes is to give an algorithm for determining whether a given code can be realized by a planar shadow." One problem I have is that I can't figure out what exactly the word standard means in the theorem - he says he will define it later and never does. And does he actually solve this "fundamental problem" in the rest of the section 3.3. If anyone knows a technical definition of standard, I would be very appreciative. The point is to figure out if the the follow question is open: Given a Gauss Code, is there an (explicit) algorithm for detecting if it can be realized as a classical knot? So if you know that it is or is not open, I would love to know too. In the case that it is known, a reference would be wonderful too. Thanks.

论坛问题帖 | has_answers | N. Owad | 2014-10-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018033 | MO-186534

Randomly placing nonoverlapping unit cuboids

Suppose one places unit cuboids of dimension d with min-corners uniformly distributed to lie in $[0,n]^d$, but with cuboid (strict) overlap forbidden. At some point, the region is "saturated," without room for more cuboids to fit. For example, for $d=2$ squares in $[0,10]^2$, it appears that about 63 squares on average will fit (whereas $n^d=100$ could fit): So here the packing density is about 63%. Q. Is the expected number of cuboids known, as a function of n and of d? Is the growth rate in n known for fixed d? Update. Benjamin Dickman identified the $d=1$ version as the Rényi parking problem, where it is known that, for large n, the packing density approaches $m \approx 0.7476$, i.e., the expected number of cars(intervals) is $m n$. It is conjectured that in $d=2$, the expected packing density approaches $m^2 \approx 0.559$. Yoav Kallus points to a related paper that uses the apt phrase sequential random packing for the process.

论坛问题帖 | no_answers_at_retrieval | Joseph O'Rourke | 2014-11-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018034 | MO-187421

Inverted pair of complex analytic families

I read the following "problem" in an old set of notes of Morrow and Kodaira which focused on deformations of complex manifolds: Find a pair of complex analytic families $\lbrace M_t\rbrace$ and $\lbrace N_t\rbrace$ with $|t|<1$ such that $M_t=N_0$ for $t\ne 0$ and $N_t=M_0$ for $t\ne0$, with $M_0\ne N_0$. (Every manifold assumed compact.) As of 1971 there were no known examples. Up to date, is this possibly true? Naively it seems impossible. In the spirit of Lefschetz fibrations, we have M_0 as a critical level set amongst the M_t (thinking of cylinders which get pinched via a vanishing cycle). It is then not generic, and any perturbation of this fiber (locally about the nodal point) would alter the topology. So to me

the inverted family $\lbrace N_t\rbrace$ couldn't exist, because it would force a generic M_{t_0} to take over the role as the critical level set amongst the N_t, but there is nothing "non-regular" about it. If the manifolds are allowed to be noncompact, perhaps it's easy, again via a rough sketch: take $\lbrace M_t\rbrace$ to be the family of open cylinders which pinch the vanishing cycles at $t=0$, and $\lbrace N_t\rbrace$ to be the family of "pinched" cylinders where the pinching point pushes off to infinity as $|t|\to 0$. Edit: The problem is more subtle than I thought, and my example with Lefschetz fibrations doesn't apply. From the definition of a "complex analytic family", all M_t are diffeomorphic to each other (in particular, they're all nonsingular spaces), and so we can only distinguish them by their holomorphic structures. In other words, $M_0\ne N_0$ means diffeomorphic but not complex-analytically homeomorphic.

论坛问题帖 | has_answers | Chris Gerig | 2014-11-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018035 | MO-187680

Reduction argument from a general vertex set V(G) to a prime power in Prof. Keevash's proof on the Existence of Designs

The proof flow of the paper "On the Existence of Designs" by Prof. Keevash as I understand it is the following: -- Reduction from the general case to $V = \mathbb{F}_{p^a}$ (Lemma 6.3) -- Covering all the linearly dependent r-sets of $V = \mathbb{F}_{p^a}$ using Φ'. This also covers "bounded" number of r-sets which have linearly independent entries(dimension r over $\mathbb{F}_{p}$)but does not cover any set from the template G_r^{*}. -- Cover the r-sets of the template G_r^* using $M(G^*)$. The template is algebraically defined and it basically is a refined version of all q-tuples of the form My where the $q\times r$ matrix M has all the sub matrices of size r to be full rank and $y \in \mathbb{F}_{p^a}^r$. -- Now cover the r-sets that are full rank, not in the template G_r^* and not covered by Φ' using the nibble. -- Now to cover the r-sets that are not covered by the nibble, edges are picked from G_q (such that all r-sets except one belongs to G_r^*) in a greedy way randomly. This uses the processes defined in section 4 and generalised in section 5. This could cover few r-sets from the template G_r^* again. -- Express the sets of G_r^* that are covered twice as the difference $\partial\Psi^+ - \partial\Psi^-$. This uses Lemma 5.28 which is a generalisation of Graver and Zurkat's result. -- Apply local modifications on the covering of the template $M(G^*)$ to include $\partial\Psi^+$. This does not change G_r^*. This uses the cascade machinery built in the last subsection of section 5. -- Replace $\partial\Psi^+$ with $\partial\Psi^-$ in the template cover and this completes the construction. I was trying to understand the reduction as mentioned in the first step (Lemma 6.3). I do understand the way the complex is decomposed according to the intersection pattern with the set V' but I find myself stuck in the technicalities of the part where he proves that the greedy algorithm can always be completed(Page 46 last 2 paragraphs). The exact places at which I am not very clear are: 1) The motivation for keeping track of the full sets in the reduction argument to p^a. (Lemma 6.3). I am not very sure as to where exactly this is used but I think this is used in getting the degree regularity requirements to apply the nibble on A^x when $x^0 < r$. However I do understand why full sets were needed to be kept track of in the initial elimination of the dependent sets after the reduction to p^a in the proof of Theorem 6.1(Page 48). 2) Also I am not very clear on the analysis of G^2 in the proof of Lemma 6.3(page 46). Where he analyses G^2 by repeatedly restricting to a random subset (p-binomial) and a bounded subset (ie using Lemma 3.19 repeatedly). Could you please provide an intuitive explanation as to why the greedy algorithm here can be completed.

论坛问题帖 | no_answers_at_retrieval | Sankeerth | 2014-11-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018036 | MO-189431

Long paths in the supercritical percolation.

I have a question on the length of the longest path, denoted by ℓ_n, in the supercritical percolation on $[0,n]^d$, denoted by C_n. We know that C_n has a giant component whose size is of order n^d. The question is that what is the order of ℓ_n. Is it n^d? For $d=2$, Grimmett proved that ℓ_n is of order n^2. The idea is interesting. He showed that there are many (of order n) edge-disjoint paths crossing the square from the left side to the right side. Then near the left and right sides, we can find crossings from the bottom to the top. By geometry property of the dimension 2, the two crossings cut all above edge-disjoint paths. Hence we get a long path whose size is of order n^2. For $d\geq 3$, it is shown that there are many (of order n^{d-1}) edge-disjoint paths crossing the square from the left side to the right side. However, I can not glue these paths. Do you have any idea to deal with my problem? Thank you in advance!

论坛问题帖 | no_answers_at_retrieval | HHH | 2014-12-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018037 | MO-190938

Is every projective space curve a set-theoretic intersection of two surfaces? What is known about this question?

I am sorry if this question has been already asked, couldn't find anything similar myself. I have recently recalled this long standing open problem of whether every irreducible curve in $\mathbb{P}^3(\mathbb{C})$ is a set-theoretic intersection of two surfaces in $\mathbb{P}^3(\mathbb{C})$. I know that there are results in characteristic $p > 0$. My question is what is the best result regarding this problem known today? Some references would be appreciated.

论坛问题帖 | has_answers | shamovic | 2014-12-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018038 | MO-191191

Open problems in compressed sensing

What are the main open problems in compressed sensing? I am interested in theoretical as well as in numerical point of view.

论坛问题帖 | has_answers | Felix Goldberg | 2014-12-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018039 | MO-191386

Ruth-Aaron triples, etc

A Ruth-Aaron pair is two numbers $(n,n+1)$ such that their sum of prime factors is equal, counting repeated prime factors. (The name refers to Hank Aaron's 715 homeruns surpassing Babe Ruth's 714!) So $$15 = 3 \cdot 5 \;;\; 3+5 = 8$$ $$16 = 2^4 \;;\; 2+2+2+2 = 2 \cdot 4 = 8$$ The notion can be generalized: $$417,162 = 2 \cdot 3 \cdot 251 \cdot 277 \;;\; 2+3+251+277 = 533$$ $$417,163 = 17 \cdot 53 \cdot 463 \;;\; 17+53+463 = 533$$ $$417,164 = 2^2 \cdot 11 \cdot 19 \cdot 499 \;;\; 2+2+11+19+499 = 533$$ My question is: Q. Are there k-tuples, for all $k \ge 2$, for which there exists at least one n such that $(n, n+1, n+2, \ldots, n+k)$ has the same sum of prime factors (counting repeated prime factors as above)? Addendum. I was out of date, relying on a "claimed proof by Erdős" that there were an infinite number of R-A pairs, which turned out not to be sound, as Benjamin Dickman explained. So the problem is open even for pairs, let alone k-tuples. Apparently no Ruth-Aaron quadruple is known. David Stork wrote Mathematica search code and found there are no such quadruples up to 10^{10}.

论坛问题帖 | has_answers | Joseph O'Rourke | 2014-12-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018040 | MO-194832

Open problems in Berkovich geometry

I would like to know if there is a state of the art recent reference on non-archimedean analytic spaces mentioning/listing open problems, conjectures, unresolved questions in the theory (*). I have looked for such a reference a bit, but didn't find anything. In case there is really nothing, I would be glad if experts in the field present here could enlighten me. (*) By "theory" I mean "mainstream" theory over a non-archimedean complete valued field (trivial valuations are accepted), including cohomological questions concerning the spaces themselves, but also theory over Banach rings, as mentioned in Berkovich's AMS monogograph. I must precise that even if dynamics and potential theory do perfectly fit for an answer, they are not my first target of interest, I am indeed looking for open problems, conjectures, unresolved or partially resolved questions that are more linked to Berkovich's or Temkin's works, that is, to works developing the theory or its links with schemes and formal schemes, cohomology, vanishing cycles.

论坛问题帖 | has_answers | Olórin | 2015-01-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018041 | MO-196033

Can any finite lattice be realized as an intermediate subgroups lattice?

Let G be a finite group and H a subgroup. Let $\mathcal{L}(H \subset G)$ be the lattice of all the intermediate subgroups between H and G. Question: Can any finite lattice be realized as an intermediate subgroups lattice? Remark: It's true for all the finite distributive lattices (see theorem 2.1 here).

论坛问题帖 | has_answers | Sebastien Palcoux | 2015-02-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018042 | MO-1975

Is a smooth closed surface in Euclidean 3-space rigid?

Classical theorem of Cohn-Vossen: A closed convex surface in Euclidean 3-space cannot be deformed isometrically. Robert Connelly found an example of a polyhedral surface that can be deformed isometrically. A metal hinged model of it can be found at IHES. But what about an arbitrary not-necessarily-convex smooth closed surface? Is it necessarily rigid? Or maybe it might be possible to make a smooth version of Connelly's example? It's easy to make smooth "hinges". The real challenge is finding a smooth model of the vertices, which is where two or more hinges meet.

论坛问题帖 | has_answers | Deane Yang | 2009-10-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018043 | MO-198082

Are there any interesting open questions having to do with submodularity, specially in the intersection of theoretical machine learning?

I was interested in knowing about open research topics related with sub modularity, specially within its intersection with theoretical machine learning (and related topics). It seems to me that much of what I've seen, specially related with greedy algorithms and theoretical guarantees has been solved. Is there anything open questions related with theoretical guarantees that is open in sub modularity?

论坛问题帖 | no_answers_at_retrieval | Charlie Parker | 2015-02-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018044 | MO-199111

On Descartes / spoof odd perfect numbers

Descartes, Frenicle, and subsequently Sorli, conjectured that $k = 1$, if $N = {q^k}{n^2}$ is an odd perfect number given in Eulerian form (i.e., q is prime with $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$). The inequality $q^k < n^2$ follows from a result of Pomerance. In particular, this implies that $n < q \Longrightarrow k = 1$ is true. (Note that, if $n < q$, then the Euler prime q becomes the largest prime factor of the odd perfect number N. In comparison to the case of even perfect numbers, the Mersenne prime $2^p - 1$ is the largest prime factor with exponent 1.) Taking off from a recent MO question, we have the following related papers: Descartes Numbers by Banks, et.al. and SPOOF ODD PERFECT NUMBERS by Dittmer My question is this: Does anybody here know of more recent papers that tackle whether the quasi-Euler prime of a spoof odd perfect / Descartes number is the largest quasi-prime factor and also has exponent 1? Quick searches via Google for the keywords "Descartes number" or "spoof odd perfect number" do not seem to yield much results. I have likewise e-mailed Professors Banks and Dittmer, hoping to get some information. [Edit (March 5, 2015): Just for the sake of comparison - In OEIS sequence A228059, T. D. Noe lists odd numbers of the form ${r^s}{t^2}$, where r is prime with $r \equiv s \equiv 1 \pmod 4$ and $\gcd(r,t)=1$, that are closer to being perfect than previous terms. He notes that, coincidentally the first 9 numbers in this sequence have the

exponent $s=1$. Additionally, except for the first number in the sequence (45), all of the succeeding numbers listed in A228059 satisfy $r < t$.]

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2015-03-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018045 | MO-201419

What is the status on questions related to Bhargava's factorial function?

In Manjul Bhargava's The Factorial Function and Generalizations he motivates a new type of factorial $n!_S$ using by generalizing a few theorems like: For $k, l \in \mathbb{Z}$, we have $k! \times l!$ divides $(k+l)!$. For any primitive polynomial $f(x) \in \mathbb{Z}[x]$ with $\deg f = k$ then $\mathrm{gcd}\{ f(a): a \in \mathbb{Z}\}$ divides $k!$ In the process of solving generalizing these two results, he invents a factorial for any set of integers $S \in \mathbb{Z}$. For any prime p, order the element of S by: choose $a_0 \in S$ find a_1 giving the smallest power of $(a_1 - a_0)$ find a_2 giving the smallest power of $(a_2 - a_0)(a_2 - a_1)$ ... find a_k giving the smallest power of $\prod_{i< k} (a_k - a_i)$ One could look for analogues of the gamma function, stirling's approximation, binomial theorem and taylor series expansion of e and indeed, Bhargava mentions these questions towards the end of the paper. Have any of these questions been answered (partially or otherwise)?

论坛问题帖 | no_answers_at_retrieval | Asvin | 2015-03-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018046 | MO-201922

The open problem in General Relativity?

Q. Is there a single, clear mathematical question that has emerged as the open problem in General Relativity? I ask this on the ~100th anniversary of Einstein's (4-page!) 1915 paper, "Die Feldgleichungen der Gravitation," Preussische Akademie der Wissenschaften, Sitzungsberichte, 1915 (part 2), 844-847. (Wikisource:The Field Equations of Gravitation.) Einstein's notebook, Riemann curvature tensor. It is difficult to surpass Willie Wong's thorough 2011 answer to the MSE question, "Open problems in General Relativity." But perhaps a core mathematical question has risen to the fore since? Naked singularities? Beyond the Cauchy horizon? "Electrodynamics of moving bodies?" Florentin Smarandache's 2013 Unsolved Problems in Special and General Relativity (PDF download.)? Or perhaps Willie's inventory cannot be condensed or sorted further at this point in time?

论坛问题帖 | has_answers | Joseph O'Rourke | 2015-04-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018047 | MO-201958

Open problems in continued fractions theory

I propose to collect here open problems from the theory of continued fractions. Any types of continued fractions are welcome.

论坛问题帖 | has_answers | Alexey Ustinov | 2015-04-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018048 | MO-20200

distribution of degree of minimum polynomial for eigenvalues of random matrix with elements in finite field

This is an attempt to extend the current full fledged random matrix theory to fields of positive characteristics. So here is a possible setup for the problem: Let $A_{n,p}$ be an $n \times n$ matrix with entries iid taking values uniformly in F_p. Then one should be able to find its eigenvalues together with multiplicities, which might lie in some finite extension of the field F_p. To ensure diagonalizability, one might even take $A_{n,p}$ to be symmetric or antisymmetric (I am not so sure if that guarantees diagonalizability in F_p but I have no counterexamples either). Now the question is if we associate to each eigenvalue λ the degree of its minimal polynomial $d(\lambda)$, then does the distribution of $d(\lambda)$ as n goes to infinite converge to some law upon normalization (say maybe Gaussian)? I am very curious whether others have studied this problem before. Maybe it's completely trivial.

论坛问题帖 | has_answers | John Jiang | 2010-04-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018049 | MO-203764

What are the strongest conjectured uniform versions of Serre's Open Image Theorem?

This question concerns the uniform conjectured effective versions and generalizations of these two results of Serre on ℓ-adic Galois representations $\rho_{E,\ell}$ associated to a non-CM elliptic curve E defined over a number field K: For each prime ℓ, the index $I(E,\ell)$ of the image of $\rho_{E,\ell}$ in $\operatorname{GL}_2(\mathbb{Z}_\ell)$ is finite (and bounded independent of ℓ), and The representation $\rho_{E,\ell}$ is surjective for all but finitely many primes ℓ. Remark: As I understand it, these two statements together say that the representation $\rho_E:G_K \to \operatorname{GL}_2(\hat{\mathbb{Z}})$ made up of all the ℓ-adic ones has open image, and hence "Serre's Open Image Theorem" refers to both of these statements, although many papers written on this topic seem to only discuss one statement or the other. Let $p_0(E)$ denote the smallest prime such that $\rho_{E,\ell}$ is surjective for all $\ell > p_0$. My question: What are the strongest conjectured ("conjectured" means either officially conjectured or in a folklore way, or even someone writing down that they hope it's true) generalizations of Serre's Theorem which are uniform in E? Here are examples of the kind of statements I am referring to: Conjecture 1: $p_0(E)$ is bounded independently of E, where E ranges over all elliptic curves defined over $\mathbb{Q}$ (and the bound is 37). Conjecture 2: $I(E,\ell)$ is bounded independently of E and ℓ, where E ranges over all elliptic curves defined over $\mathbb{Q}$ (and the bound is...?). Conjecture 1': Like Conjecture 1, but for elliptic curves defined over an arbitrary number field K, and the bound only depends on K. Conjecture 2': Like Conjecture 2, but for elliptic curves defined over an arbitrary number field K, and the bound only depends on K. Conjecture 1'': Like Conjecture 1', but the bound only depends on $[K:\mathbb{Q}]$. Conjecture 2'': Like Conjecture 2', but the bound only depends on $[K:\mathbb{Q}]$. I am aware that a lot of

progress has been made on Conjecture 1, and we might be close to knowing that one.

论坛问题帖 | has_answers | Bobby Grizzard | 2015-04-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018050 | MO-204361

Open problems in Banach spaces, universality

I have gathered a list of universality problems in Banach spaces which have been solved: 1.The non existence of a separable reflexive space universal for the class of separable reflexive spaces. 2.If a space is universal for the class of separable reflexive spaces, then it is universal for the class of separable Banach spaces. 3.There is a separable reflexive space which is universal for all separable uniformly convex spaces. My question is, are there any open problems in a similar vein to these?

论坛问题帖 | has_answers | Tom | 2015-04-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018051 | MO-204662

Open problems books

As the title might indicate , I would like to look for recommendations for mathematical book that present open problems in depth with commentary. The only book of this type that I've come across is Arnold's Problems by V.I Arnold, are there any similar ones? The subjects that these problems belong to should preferably be the following: Algebraic Geometry. Algebraic Topology. Intersections of pure mathematics with theoretical computer science or ''theoretical'' biology.

论坛问题帖 | has_answers | Theristo | 2015-05-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018052 | MO-20497

Is it known that the ring of periods is not a field?

I have just learned here that we know numbers that are not periods; is it known meanwhile that the ring of periods is not a field? I know that it is conjectured that $1/\pi$ is not a period, but the existence of a period whose inverse is not a period seems to be still open. Is this correct? More generally: is it believed that the unit group of the ring of periods is bigger than the nonzero algebraic numbers?

论坛问题帖 | has_answers | Franz Lemmermeyer | 2010-04-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018053 | MO-209190

Open problems in hyperplane/subspace arrangements?

What are some open problems in hyperplane/subspace arrangements, preferably of the combinatorial algebraic topology kind, and where can one read about them? That is, where are they discussed, and where can one learn the necessary background knowledge to understand them properly and in context? I have been working one one such problem for a while, but I'm ignorant about the big questions, and more generally, interesting problems in this field, and also current trends. (I'm aware of a few unresolved questions in Richard P. Stanley's An Introduction to Hyperplane Arrangements (available online, the unresolved problems are signaled by [5]))

论坛问题帖 | has_answers | Olivier Bégassat | 2015-06-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018054 | MO-21003

Polynomial bijection from $\mathbb Q\times\mathbb Q$ to $\mathbb Q$?

Is there any polynomial $f(x,y)\in{\mathbb Q}[x,y]{}$ such that $f\colon\mathbb{Q}\times\mathbb{Q} \rightarrow\mathbb{Q}$ is a bijection?

论坛问题帖 | has_answers | Z.H. | 2010-04-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018055 | MO-210167

Arctangents of odd powers of the golden ratio

While trying to answer this MSE question, I found that arctangents of many odd powers of the golden ratio $\varphi=\frac{1+\sqrt5}2$ are expressible as rational linear combinations of arctangents of positive integers: $$\begin{align}
\arctan\varphi&=2\arctan1-\frac12\,\arctan2\\
\arctan\varphi^3&=\arctan1+\frac12\,\arctan2\\
\arctan\varphi^5&=4\arctan1-\frac32\,\arctan2\\
\arctan\varphi^7&=3\arctan1+\frac12\,\arctan2-\arctan5\\
\arctan\varphi^9&=\arctan1-\frac12\,\arctan2+\arctan4\\
\arctan\varphi^{11}&=5\arctan1+\frac12\,\arctan2-\arctan5-\arctan34\\
\arctan\varphi^{13}&=3\arctan1-\frac12\,\arctan2+\arctan4-\arctan89\\
\arctan\varphi^{15}&=-2\arctan1+\frac32\,\arctan2+\arctan11 \end{align}$$ I was not able to find such a representation for $\arctan\varphi^{17}$ though. Question 1. Can we prove that it does not exist? I also could not find such a representation for any positive even power. Question 2. Can we prove that it does not exist for any positive even power? Question 3. Is there a simple way to determine if such a representation exists for a given power?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2015-06-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018056 | MO-212659

Connes Embedding Conjecture and Fusion Categories

I was recently introduced to Connes' Embedding Conjecture (CEC) which states: Every separable type II_{1} factor is embeddable into R^{ω}. Where ω is a generic free ultrafilter on $\mathbb{N}$, R is a hyperfinite type II_{1} factor, and R^{ω} is the ultrapower of R with respect to ω. I'm not well versed in the theory of factors and this conjecture is mysterious and difficult for me to understand. However, there do seem to be a lot of equivalent formulations of CEC. For some examples of

this see: N. Ozawa, "About the Connes Embedding Conjecture---Algebraic Approaches---, arXiv:1212.1700 [math.QA] My background is in fusion categories, and given the relationship between subfactors and fusion categories I was led to wonder: Question: Are there statements in the theory of fusion categories that are equivalent to CEC, or does CEC have implications for fusion categories? My understanding of the connections between fusion categories and subfactors, I'm sorry to say, is quite weak. So it very well could be that there is simply no relation between CEC and fusion categories, which would be a completely acceptable answer.

论坛问题帖 | has_answers | Paul | 2015-07-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018057 | MO-216174

Conjectured new primality test for Mersenne numbers

How to prove that this conjecture about a new primality test for Mersenne numbers is true ? Definition: Let $M_{q}=2^{q}-1 , S_{0} = 3^{2} + 1/3^{2} , \ and: \ S_{i+1} = S_{i}^{2}-2 \pmod{M_{q}}$ Conjecture: $$M_{q}\text{ is a prime iff: } \ S_{q-1} \equiv S_{0} \pmod{M_{q}}$$ $$\text{ and iff: } \prod_{0}^{q-2} S_i \equiv 1 \pmod{M_{q}}$$ $$\text{ and iff: } S_i \not\equiv S_0 \pmod{M_{q}} , 0 < i < q-1$$ The easy part part (if M_{q} is prime then $S_{q-1}\equiv S_{0} \pmod{M_{q}}$), has already been proven. See: http://tony.reix.free.fr/Mersenne/ConjectureLLTCyclesMersenne.pdf . Now, a proof of the converse is needed. The "classic" LLT by Lucas and Lehmer for Mersenne numbers is based on the binary tree of the digraph (under $x^2-2 \pmod{M_q}$). This new conjectured test makes use of a cycle of the same digraph. The goal of this question is to validate (or not) the method (use a cycle of the digraph) for Wagstaff numbers ($\dfrac{2^q+1}{3}$, q prime) for which we also lack a proof for the converse. See: https://trex58.files.wordpress.com/2009/01/wagstaffandfermat.pdf for a proof of the easy part, by Robert Gerbicz . I and a team (Vincent and Paul) had found a very large Wagstaff PRP with this algorithm. See: http://www.primenumbers.net/prptop/prptop.php (2^4031399+1)/3 , rank 9 for now.

论坛问题帖 | no_answers_at_retrieval | Tony Reix | 2015-09-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018058 | MO-217461

A conjecture like Cayley-Bacharach theorem

Let six points A, A', B, B', C, C' lie on a conic and a cubic. Let a conic through B, B', C, C' and meets the cubic again at A_1, A_2. Let a conic through C, C', A, A' and meets the cubic again at B_1, B_2. Let a conic through A, A', B, B' and meets the cubic again at C_1, C_2. Then six points $A_1,A_2, B_1, B_2, C_1, C_2$ lie on a conic. In this case if we let A_1, B_1, C_1 are collinear then A_2, B_2, C_2 are collinear. Check the conjecture by ggb Special case 1, The cubis are three lines. Let six points A, A', B, B', C, C' lie on a conic. Let a conic through B, B', C, C' and meets line AA' at A_1, A_2 Let a conic through C, C', A, A' and meets line BB' at B_1, B_2 Let a conic through A, A', B, B' and meets line CC' at C_1, C_2. Then six points $A_1,A_2, B_1, B_2, C_1, C_2$ lie on a conic. In this case if A_1, B_1, C_1 are collinear then A_2, B_2, C_2 are collinear. Check the special case 1 by ggb Special case 2, A conic and a cubic: Let a conic meets a cubic at six points A, B, C, A', B', C'. Let $AB', A'B$ cut the cubic at C_1, C_2 respectively; Let $BC', B'C$ cut the cubic again at A_1, A_2; Let $AC', A'C$ cut the cubic at B_2, B_1 then A_1, B_1, C_1 are collinear and A_2, B_2, C_2 are collinear. Check the special case 2 by ggb Special case 3, The conics are lines: Let two lines meet a cubic at A, B, C; and A', B', C' respectively. Let $AB', A'B$ cut the cubic at C_1, C_2 respectively; Let $BC', B'C$ cut the cubic again at A_1, A_2; Let $AC', A'C$ cut the cubic at B_2, B_1 then A_1, B_1, C_1 are collinear and A_2, B_2, C_2 are collinear. Check the special case 3 by ggb Special case 4, The cubic are a line and a conic: Let the cubic is a conic (γ) through four points A, A', B, B' and line CC' on the plane). Let points A, B, C, A', B', C' lie on another conic. Let $AB', A'B$ meet CC' at C_1, C_2; let $AC', A'C, BC', B'C$ meet conic γ at B_2, B_1, A_1, A_2. Then six points $A_1, A_2, B_1, B_2, C_1, C_2$ lie on a conic. Check the special case 4 by ggb

论坛问题帖 | has_answers | Oai Thanh Đào | 2015-09-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018059 | MO-217567

Functions on [0,1] with positive fractional series coefficients and symmetric under x->1-x

Suppose a function $g(x)$ has a convergent power series expansion with real (not necessarily integer or rational) exponents on $[0,1]$ of the form $$ g(x)=\frac{1}{x^\delta}(1+\sum_{i=1}^\infty c_i x^{\Delta_i}) $$ where $\delta>0$, $0<\Delta_1<\Delta_2<\ldots$ and, crucially, $c_i>0$. Suppose further that $$ g(x)=g(1-x) $$ What is known about such functions? Do they occur anywhere in mathematics? (They do occur in theoretical physics, namely in the study of conformal field theories. In fact this condition is a toy model for operator product expansion associativity.) I can use the Hardy-Littlewood tauberian theorem to show that c_i's should have a certain asymptotic behavior. But this does not use the full strength of the condition $g(x)=g(1-x)$. Can I learn more about the asymptotics, e.g. about the subleading terms, via a smart use of this condition? I'm also interested in questions which are not asymptotic. E.g. I am interested how large Δ_1 can be for a fixed δ. Expanding around $x=1/2$ to third order I can easily show that $\Delta_1\le 2\delta+3$. I know that this bound is not best possible, since expanding to higher order and doing some numerics I can somewhat improve it. I'm interested in the exact bound on Δ_1 as a function of δ and which functions $g(x)$ saturate it. I can also study the question for δ integer under an extra assumption that all Δ_i's are integers. E.g. for $\delta=1$ I can find a function for which $\Delta_1= 3$: $$ g(x)=\frac{P(x)+P(1-x)}{2x(1-x)}=x^{-1}(1+2x^3+x^4+x^5+\ldots) $$ where $P(x)=3x^2-x^4$. I can also show that $\Delta_1= 4$ is impossible. But I'm most interested in the case when δ and Δ_i's are real.

论坛问题帖 | no_answers_at_retrieval | Slava Rychkov | 2015-09-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018060 | MO-218508

Sets not containing the vertices of unit triangles (Question posed by Erdős)

Following this post, I have been thinking about the problem posed by Erdős, Does there exist a constant $c > 0$ such that every subset A of the plane of area more than c contains the vertices of a triangle of unit area? I think the spirit of the problem justifies assuming A is measurable (counterexamples exist for non-measurable sets, see post), and so we can use the following to simplify: If A is measurable, with finite measure a, then for every $\epsilon >0$ there exists an $n \in \mathbb{N},$ $r > 0,$ such that A can be filled with n non-overlapping disks of radius r with combined area $a' = n \pi r^2 > a(\eta_h -\epsilon)$, where $\eta_h \approx 0.9$ is the optimal packing density of disks in the plane. So we can reduce to just considering sets of nonoverlapping disks of equal radius r. The problem becomes, Does there exist a constant $c > 0$ such that every set A of n disks of radius r, with $n \pi r^2 > c$, contains the vertices of a triangle of unit area? Now, we can scale such a set A by $1/r$ (that is, probably expanding it, because probably $r < 1$.) and ask of the new set of n unit disks, if $n\pi r^2>c$, for some c, must A contain the vertices of a triangle of area $1/r^2$? Defining $\ell = c/{(\pi r^2)}$, if $\ell<n$, must A contain the vertices of a triangle of area $\pi \ell/c$? That is, must A contain vertices of triangles of all areas $< \pi n/c$? So, setting $c' = c/\pi$ (to get rid of the constant), we get the equivalent question, Does there exist a constant $c' > 0$ such that every set of n unit disks in the plane must contain the vertices of triangles of all areas $< n/c'$? Say we are given some such c', and we must determine the conditions under which a triangle of area $c < c'$ must have vertices in A. Now, since two unit disks with centers a distance $> n/c$ apart always contain the vertices of a triangle of area n/c, we can assume that these disks are all within a distance of n/c of each other. Thus, they all lie in a big disk D of radius $\frac{2n}{\sqrt{3}c}+2$, and area d_c. We can then look at pairs of points within this disk that must not both lie in A. That is, pairs of points in D which, together with some point in A, form the vertices of a triangle of area n/c. We denote this set $\mathcal{F}_2$. Using the usual product measure, we denote the measure of $\mathcal{F}_2 \in D \times D$, m_2. The measure of $D \times D$ itself is d_c^2. We can show that, m_2 = $\Omega(cd_c^2/\sum_{1 \leq i \leq j \leq n} {s_{ij}})$ as a function of c and n, where s_{ij} is the distance between the centers of the i^th and j^th disks. Letting $S = \sum_{1 \leq i \leq j \leq n} {s_{ij}}$, we win if $S = o(c)$ (S a function of both c and n). Now, we can also look at the set of individual points in D that are "forbidden," i.e., together with two points from A, form a triangle of area n/c. We denote this set $\mathcal{F}_1$, and take it as a subset of D, with measure denoted m_1. It can be shown that the set of points f_{ij} that are forbidden due to forming a triangle of area n/c with points in disks i and j, which we say has measure $m_{1,ij}$, satisfies, $m_{1,ij} = \Omega(d_c/s_{ij})$ $\mathcal{F}_1$ is the union of the f_{ij}. Naïvely summing the $m_{1,ij}$, we get the sum m_1' of their measures is, $m_1' = \Omega(d_cS)$. If this were indeed $\mathcal{F}_2$, we would win if $S \neq o(c)$, since $d_c \sim \frac{2n}{\sqrt{3}c}$ (winning meaning $m_1 > d_c$, a contradiction). Since, from before, if $S = o(c)$ we also win, the problem would be finished. You could simply pick c large enough, and either $\mathcal{F}_1$ or $\mathcal{F}_2$ would exceed the measure of its superset, a contradiction. But of course the area of a union is not the sum of the areas. And I can't find a way to deal with the intersections of the $m_{1,ij}$. I also can't tell if I'm really right next to a solution, or if I'm just hitting the brick wall surrounding the problem in a different spot. I hope you found this interesting, at least. If the way I put growth rates is too confusing, I'll post some updates to clarify. If you have any ideas, please let me know.

论坛问题帖 | has_answers | Jason Eliot | 2015-09-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018061 | MO-218674

A conjecture associated with n-gon cut curve of degree m

This conjecture is a generalization of A theorem for cubic-A generalization of Carnot theorem, in MSE question. I'm an electrical engineer, I am not a mathematician. I don't know how to prove this result. Let $A_1A_2....A_n$ be a n-gons, Let m points B_{ij} lie on A_iA_{i+1} for j=1, 2,...., m. Then $m.n$ points B_{ij} for $i=1, 2, 3,...., n$ and $j=1, 2,...., m$ lie on a curve of degree m if only if: $$\prod_{i=1}^n \prod_{j=1}^m \frac{\overline{B_{ij}A_i}}{\overline{B_{ij}A_{i+1}}}=1$$ Note that: $A_{m+1}=A_1$

论坛问题帖 | has_answers | Oai Thanh Đào | 2015-09-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018062 | MO-218884

Bang's open question strengthening Tarski's planks problem

Tarski's Planks problem, solved by Thøger Bang in 1951, says (in a simplified $\mathbb{R}^2$ version) that it requires "planks" (parallel strips) of total width $\ge d$ in order to completely cover a disk of diameter d. The last paragraph of Bang's short paper poses an open problem: (Snapshot from Bang's paper.) Does anyone know if this problem has been solved? I am especially interested in $\mathbb{R}^2$. Bang, Thøger (1951), "A solution of the 'plank problem'", Proc. Amer. Math. Soc. 2 (6): 990-993, doi:10.2307/2031721, JSTOR 2031721, MR 0046672.

论坛问题帖 | no_answers_at_retrieval | Joseph O'Rourke | 2015-09-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018063 | MO-219344

Gerstenhaber conjecture for free loop space

I- Is the following statement still a conjecture see this article ? Conjecture (?) Let M be a simply connected compact oriented d-manifold (smooth), then $HH^{\ast}(C^{\ast}(M))$ the Hochschild cohomology of cochain complexes associated to M is isomorphic as a Gerstenhaber algebra to $H_{\ast+d}(LM)$ the d-shifted homology of the free loop space on M. II- What are the recent advances in solving the conjecture ? III- Is there a short proof of the following fact: $HH^{\ast}(C^{\ast}(M))$ is isomorphic to $H_{\ast+d}(LM)$ as a graded vector space ?

论坛问题帖 | has_answers | lilia | 2015-09-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018064 | MO-219594

Progress in Guy's "Unsolved problems in Number Theory"?

I often peruse through Guy's book whenever I'm not being boggled down by my research. It crossed my mind today if any of these "unsolved problems" have become indeed solved. I thought about doing a literature review to find out, but I realised that posting this question here is a much more a efficient way of finding the current status of these problems. I'm also open to finding out about any recent breakthroughs concerning said problems.

论坛问题帖 | no_answers_at_retrieval | user119264 | 2015-09-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018065 | MO-22

Can N^2 have only digits 0 and 1, other than $N=10^k$?

Pablo Solis asked this at a recent 20 questions seminar at Berkeley. Is there a positive integer N, not of the form 10^k, such that the digits of N^2 are all 0's and 1's? It seems very unlikely, but I don't have a proof. It's easy to see that such a number must end in 1 or 9, and then easy to see that it must end in 01, 49, 51 or 99, and you can continue recursively for as long as you like, determining possible "suffixes". Using this, I had a computer check for me that there are no such N up to about 10^{24}. If you pretend that the digits of N^2 are randomly distributed, and N has n-digits, there's a $(2/10)^{2n}$ chance of satisfying this condition. There are only 10^n n-digit numbers, so you might expect a $(4/10)^n$ chance of having a some n-digit number. This suggests we shouldn't expect to find anything. (If you try the same problem in other bases, where the probabilities are better, you do find a few: in base 5, 222112144, 22222111221444 and 100024441003001 work.)

论坛问题帖 | has_answers | Kim Morrison | 2009-09-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018066 | MO-222478

A conjecture about $\lfloor n!\cdot q/e\rfloor-\,!n\cdot q$

I was thinking about this question asked at Math.SE, when I came up with the following conjecture. For every $q\in\mathbb Q$ consider a sequence $s_n^{(q)}$ (terms within the sequence are indexed by $n\in\mathbb N$): $$s_n^{(q)}=\left\lfloor\frac{n!\cdot q}{e}\right\rfloor-\,!n\cdot q$$ ($!n$ denotes the subfactorial). Conjecture: For every q the sequence $s_n^{(q)}$ is periodic. Furthemore, if q is a positive integer is a reciprocal of a positive integer m, then the period is: $$p_{1/m}=\begin{cases}m,\quad m\text{ is even}\\2m,\quad m\text{ is odd}\end{cases}$$ Could you suggest any ideas how to prove this conjecture?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2015-11-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018067 | MO-224125

Looking for (information about) long diamonds

I was given an open problem as a birthday present recently. While I can probably handle spoilers at this point, what I really want are literature and other references. Also acceptable would be suggestions for approaches. Ideally, search terms for the web are most welcome. (It is my lack of imagination for search terms for this problem that is most keeping me from doing preliminary research on it.) Question: Where and how can I find out more about the Problem ($C_p \lt 2$?) below? Problem: Let n be a an integer parameter sufficiently large. Let x and y be vectors in n-dimensional real space. (Do complex if it makes it easier.) We will be using a family of L_p norms $|z|$ for p real and greater than 1 (pth root of the sum of the pth powers of the n coordinates of the vector z), although other norms are also of interest. Fixing p and its associated norm, there is a value C_p (which may be independent of n and all other parameters) such that for all x and y with $|x|=|y|=|x-y|=1$, one has $|x+y|\leq C_p$. The triangle inequality gives $C_p \leq 2$, and for $p=2$ we have $C_2 = \sqrt{3}$. Is it true that $C_p \lt 2$? (Can we also hope that lim sup over all $p \gt 1$ of C_p is also less than 2?) C_p to me represents the length of the long diagonal of a rhombus (diamond) with short diagonal and sides length 1. Please retag this post as appropriate. Gerhard "Still Enjoys His Birthday Presents" Paseman, 2015.11.20

论坛问题帖 | has_answers | Gerhard Paseman | 2015-11-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018068 | MO-224711

All compact surfaces $S\subseteq \mathbb{R}^3$ are rigid?

Recently I've come across a lecture in differential geometry by Fernando Codá (in portuguese!) in which he stated that the following problem is (at least, up to 2014) open: Given $S\subseteq\mathbb{R}^3$ a smooth (i.e. C^∞) compact boundaryless connected surface (therefore orientable), let $f:S\times[0,1]\to \mathbb{R}^3$ be a differentiable map with $S_t:=f(S\times\{t\})$ being a surface and $f_t:S\to S_t$ an isometry (where $f_t(p):=f(p,t)$), where $f_0\equiv \mathrm{Id}_S$. Then there are rigid motions $A_t:\mathbb{R}^3\to\mathbb{R}^3$ with $f_t(S) = A_t(S)$. More generally, in the cases of the sphere S^2 (Liebmann's Theorem) and "ovaloids", i.e. surfaces with Gaussian curvature K strictly positive (Cohn-Vossen's rigidity theorem), it is known that isometric embeddings of it in $\mathbb{R}^3$ are exactly the restrictions of the rigid motions from $\mathbb{R}^3$. Not all compact connected surfaces $S\subseteq\mathbb{R}^3$ satisfies this, as pointed out in the book Curves and Surfaces, by S. Montiel and A. Ros: Anyway, it is possible to show that this isometry in particular cannot be obtained by a continuous family of isometries from the identity map to the latter, so it's not a counterexample to the conjecture. Although this seems to me to be a very studied problem, and I imagine that there must be several approachs to it, I just couldn't find enough information about it on the internet (the Montiel & Ros book was recommended by Codá later in the same lecture linked above), for "rigidity" seems to be a very comprehensive term. So my actual question is: Actual question: Where can I find more information about this problem and the current approaches going on?

论坛问题帖 | has_answers | Alufat | 2015-11-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018069 | MO-225039

Open questions on (finite) tensor categories

I would like to know about problems on (finite) tensor categories. I have read Etingof´s notes from his course at MIT. I have a question: There exists any reference where I can find an open problem about this topic? Thank you very much for your attention.

论坛问题帖 | has_answers | F.R | 2015-12-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018070 | MO-225459

A conjecture based on Wilson's theorem

Definitions: Lagrange's theorem implies that for each prime p, the factors of $(p - 1)!$ can be arranged in unequal pairs, with the exception of $±1$, where the product of each pair $≡ 1 \pmod p$. See Wiki article on Wilson's theorem. From the example in the link above, for $p=11$ we have $$(11-1)!=[(1\cdot10)]\cdot[(2\cdot6)(3\cdot4)(5\cdot9)(7\cdot8)] \equiv [-1]\cdot[1\cdot1\cdot1\cdot1] \equiv -1 \pmod{11}$$ Let the products of the pairs that $≡ 1 \pmod p$ be the multiset A_p, and A_{p_n} the multiset for the nth prime. For the above example then, $A_{p_5}=\{(2\cdot6),(3\cdot4),(5\cdot9),(7\cdot8)\}=\{12,12,45,56\}$. Conjecture: $$\lim\limits_{n\rightarrow\infty}\dfrac{\sum\limits_{k \in A_{p_n}}(k-1)}{(p_n)^3}\approx\frac18$$ where p_n is the nth prime. Examples: For $p=11$ we have $$\dfrac{11+11+44+55}{11^3}=\dfrac{1}{11}$$ For $p=997$ we have $$\dfrac{123218233}{997^3}=\dfrac{123218233}{991026973}$$ Comments: As @YCor noted below, the -1 in the $k-1$ can be removed, since its contribution tends to 0. The conjecture can therefore be simplified to $$\lim\limits_{n\rightarrow\infty}\dfrac{\sum\limits_{k \in A_{p_n}}k}{(p_n)^3}\approx\frac18$$ I have no idea whether the above statement is correct, or how to go about trying to find a proof. Any comments on the any of the above are most welcome.

论坛问题帖 | has_answers | martin | 2015-12-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018071 | MO-225572

On cubic reciprocity for $x^3+y^3+z^3 = 996$?

I. The Diophantine equation, $$x^3+y^3+z^3 = 3w^3\tag1$$ with $x\geq y \geq z$ and $w=1$ has only two known solutions, namely $1,1,1$ and $4,4,-5$. Are there larger ones? As Noam Elkies points out in this post, Cassels in a 1985 paper showed that $w=1$ must have, $$x\equiv y\equiv z \bmod 9\tag2$$ For general w, Heath-Brown in a 1992 paper showed that, $$\text{either}\;x\equiv y\equiv z \bmod 9,\; \text{or one of}\; x,y,z\;\text{is}\;9m\tag3$$ Checking the Elsenhans-Jahnel list for $x^3+y^3+z^3=N$ with $N<1000$, one finds large solutions for $w=2,3,4$. (The list only covers 10^{14}, so if it can be raised higher, maybe a large one can be found for $w=1$ as well.) II. Trying to find something similar to (1), and if I did my search right, there are at least two other N that obeyed (2), namely $N=996$, $$x^3+y^3+z^3 = 996\tag4$$ with, $$x,y,z = 11,2,\,-7$$ $$x,y,z = 2169364505441,\, -631266388780,\, -2151398424325$$ and $N = 2^4\times3^3+3=435$, $$x^3+y^3+z^3 = 435\tag5$$ with seven known (and rather large) solutions, all of which satisfied (2). Questions: Is it true that all x,y,z of (4) and (5) also obey $x\equiv y\equiv z \bmod 9$? In general, for what N does this condition hold?

论坛问题帖 | has_answers | Tito Piezas III | 2015-12-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018072 | MO-225600

Diagonal argument for even perfect numbers

Following this, let's define the notion of perfect sequence as follows: $(u_{i})_{i}$ is a perfect sequence if and only if it is the sequence of divisors of an even perfect number in increasing order or the infinite sequence $(u_{i})_{i}, \ \ u_{i}=2^{-i}$. Let's now list the set of finite perfect sequences by increasing number of terms, so that $(v_{1,i})=(1, 2, 3, 6)$, $(v_{2,i})=(1, 2, 4, 7, 14, 28)$, And so on. My question is: is the diagonal sequence $(v_{i,i})_{i}$ such that $\sum_{i}{v_{i,i}}^{-1}=2$? Many thanks in advance.

论坛问题帖 | has_answers | Sylvain JULIEN | 2015-12-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018073 | MO-225870

Open Problems for Undergraduates

Suppose: I am a 'problem-solver' rather than a 'theory-builder' I am an undergraduate student I have a passion for solving mathematical problems The homework I get is not satisfying (in the sense that the problems are computing-problems rather than problems that require creative thinking), and I get far too little homework Where can I find interesting problems (that require creative thinking) if I want to have fun solving mathematical problems and to practice problem-solving? Are there lists/books of such problems? Furthermore, suppose I want to know how it is to do research. Are there lists of the kind "open problems which can be understood by undergraduates". I guess these open problems should be in the fields of "discrete mathematics/combinatorics" and "graph theory". I only found: http://dimacs.rutgers.edu/~hochberg/undopen/

论坛问题帖 | has_answers | sdfaewrFAAEA | 2015-12-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018074 | MO-226580

Does every model of $I\Delta_0$ has an end extension to a model of $I\Delta_0+\Omega_1$?

Does every model of $I\Delta_0$ has an end extension to a model of $I\Delta_0+\Omega_1$? In End extensions of models of linearly bounded arithmetic paper the author said this problem is open. I want know is there any progress on this problem? Thanks.

论坛问题帖 | no_answers_at_retrieval | Erfan Khaniki | 2015-12-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018075 | MO-229999

The complexity of Max-K interval selection

I came up with the following problem, but do not know how to analyze it. Let S be an ordered set of integers with size n (i.e., $S=\{1,2,...,n\}$). An interval $INV(a,b)$ covers the elements in S from a to b $(1 \leq a < b \leq n)$. For example, $INV(3,5)$ covers the element of $\{3,4,5\}$. Now I am trying to analyze the following problem: Given a set of intervals I, select k intervals R from I (i.e., $R\subseteq I$ and $|R|=k$), such that R covers the most of S. I am also interesting in a more generalized version where each element has a weight. But I wish to analyze the simpler version first. Specifically, I wish to know whether there exists polynomial time (wrt. n) for this problem. If not, is this problem NP-hard, and how to prove it ? Any hints or discussions would be very helpful. Thank you.

论坛问题帖 | no_answers_at_retrieval | fanqi | 2016-02-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018076 | MO-230174

If $N = qn^2$ is an odd perfect number with $\gcd(q,n)=1$, is it possible to have $q + 1 = \sigma(n)$?

The title says it all. Question If $N = qn^2$ is an odd perfect number with Euler prime q and $\gcd(q,n)=1$, is it possible to have $q + 1 = \sigma(n)$? Heuristic From the Descartes spoof, with quasi-Euler prime q_1: $$n_1 = 3003 < \sigma(n_1) = 5376 < q_1 = 22021$$ So it appears that it might be possible to prove that $q + 1 \neq \sigma(n)$. Some Essential Estimates Acquaah and Konyagin recently obtained the estimate $q < n\sqrt{3}$. We will use this estimate to obtain an upper bound for $\sigma(q)/n$. Ochem and Rao recently obtained the lower bound $N > {10}^{1500}$ for the magnitude of an odd perfect number. Using this bound, together with the inequality $n < q$, gives $$I(q) < 1 + {10}^{-500}.$$ Motivation We wish to prove the following proposition: If $N = {q^k}{n^2}$ is an odd perfect number with Euler prime q, then $3 \nmid N$ implies that $q < n$. If $q + 1 \neq \sigma(n)$, then it follows that $$I(q) + I(n) \neq \frac{\sigma(q)}{n} + \frac{\sigma(n)}{q}$$ from which we obtain $$I(q) + I(n) < \frac{\sigma(q)}{n} + \frac{\sigma(n)}{q},$$ since the reverse inequality $$\frac{\sigma(q)}{n} + \frac{\sigma(n)}{q} < I(q) + I(n)$$ will violate the inequality $I(q) < \sqrt[3]{2} < I(n)$ (see this paper). (Edit February 8 2016: Assuming $$\frac{\sigma(q)}{n} + \frac{\sigma(n)}{q} < I(q) + I(n)$$ is true, then $$\left(q < n\right) \land \left(\sigma(n) < \sigma(q)\right)$$ is false. However, I am currently unable to rule out $$\left(n < q\right) \land \left(\sigma(q) < \sigma(n)\right).$$ This particular case remains open.) But the inequality $$I(q) + I(n) < \frac{\sigma(q)}{n} + \frac{\sigma(n)}{q}$$ implies that the biconditional $$q < n \Longleftrightarrow \sigma(q) < \sigma(n)$$ holds. This biconditional is then a key ingredient in the proof of the proposition mentioned earlier. My method is able to rule out $\sigma(q) = q + 1 = \sigma(n)$ if $3 \nmid n$, since we obtain $$2.799 \approx 1 + 2^{\frac{\log(6/5)}{\log(31/25)}} \leftarrow \frac{q + 1}{q} + \left(\frac{2q}{q + 1}\right)^{\frac{\log(I(5))}{\log(I(5^2))}} \leq I(q) + \left(I(n^2)\right)^{\frac{\log(I(u))}{\log(I(u^2))}}$$ $$< \frac{\sigma(q)}{q} + \frac{\sigma(n)}{n} = \frac{\sigma(q)}{n} + \frac{\sigma(q)}{q} < \sqrt{3}\left(1 + {10}^{-500}\right) + \left(1 + {10}^{-500}\right) \approx 2.732,$$ (where the smallest prime factor u of N satisfies $u \geq 5$), whence we arrive at a contradiction. Further Considerations If $\sigma(q) = q + 1 = \sigma(n)$ and $3 \mid n$, then the same method does not force a contradiction, because we then have $$2.7199 \approx 1 + 2^{\frac{\log(4/3)}{\log(13/9)}} \leftarrow \frac{q + 1}{q} + \left(\frac{2q}{q + 1}\right)^{\frac{\log(I(3))}{\log(I(3^2))}} \leq I(q) + \left(I(n^2)\right)^{\frac{\log(I(u))}{\log(I(u^2))}}$$ $$< \frac{\sigma(q)}{q} + \frac{\sigma(n)}{n} = \frac{\sigma(q)}{n} + \frac{\sigma(q)}{q} < \sqrt{3}\left(1 + {10}^{-500}\right) + \left(1 + {10}^{-500}\right) \approx 2.732,$$ where u is the smallest prime factor of N. Added February 7 2016 If $q + 1 = \sigma(n)$, then $\sigma(n) \equiv 2 \pmod 4$, so that n takes the form $$n = {p^r}{m^2}$$ where p is a prime with $p \equiv r \equiv 1 \pmod 4$ and $\gcd(p, m) = 1$. If $3 \mid n$, then $p \neq 3$, so that $3 \mid m$. So I have $$\sigma(n) = \sigma(p^r)\sigma(m^2)$$ where $\sigma(p^r) \equiv r + 1 \equiv 2 \pmod 4$. Since $\gcd(q, q+1) = 1$, then $\gcd(q, \sigma(n)) = 1$, so that $$\gcd(q, \sigma(p^r)\sigma(m^2)) = 1.$$ Thus, $q \nmid \sigma(p^r)$ and $q \nmid \sigma(m^2)$. However, note that $$q \mid \sigma(n^2) = \sigma(p^{2r})\sigma(m^4).$$ Alas here is where I get stuck.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2016-02-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018077 | MO-231160

Open problems in Hopf algebras

I couldn't find a list of open problems in Hopf algebras. So my question is the following: In the theory of Hopf algebras, what are the (big) open problems? Any kind of problem/question will be welcome. I am interested in any kind of problem involving Hopf algebras, say going from combinatorics or topology to abstract algebra.

论坛问题帖 | has_answers | david | 2016-02-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018078 | MO-231769

A generalization of Frankl's conjecture?

Would it be reasonable to conjecture what follows : there is a real constant $c > 1/2$ such that, for every natural number n, if $X_{1}, \ldots , X_{n}$ is a union-stable family of distinct finite sets with at least two elements in their union U, then there is at least one subset of U with two elements that intersects at least $cn - 1$ sets X_{i} ? Thanks in advance.

论坛问题帖 | has_answers | Panurge | 2016-02-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018079 | MO-231922

Torsion-freeness of two groups with 2 generators and 3 relators and Kaplansky Zero Divisor Conjecture

Let G_1 and G_2 be the groups with the following presentations: $$G_1=\langle a,b \;|\; (ab)^2=a^{-1}ba^{-1}, (a^{-1}ba^{-1})^2=b^{-2}a, (ba^{-1})^2=a^{-2}b^2 \rangle,$$ $$G_2=\langle a,b \;|\; ab=(a^{-1}ba^{-1})^2, (b^{-1}ab^{-1})^2=a^{-2}b, (ba^{-1})^2=a^{-2}b^2 \rangle,$$ Are these groups torsion-free? Motivation: In both of these groups $1+a+b$ as an element of the group algebra $\mathbb{F}_2[G_i]$ over the field with two elements is a zero divisor. Thus one has a counterexample for the Kaplansky zero divisor conjecture if one of G_is is torsion-free! $$(1+a+b)(b^{-1}a^{-2}ba^{-1}+a^{-1}ba^{-2}ba^{-1}+a^{-1}ba^{-1}+b^{-1}a^{-1}+a^{-1}b^2a^{-1}ba^{-1}+aba^{-1}ba^{-1}+1+a^{-2}ba^{-1}+a^{-1}b^{-1}a^{-1}+b+baba^{-1}ba^{-1}+ba^{-1}ba^{-1}+a^{-1}+b^{-1}a)=0$$ $$(1+a+b)(aba^{-1}ba^{-1}+a^{-1}b^2a^{-1}ba^{-1}+a^{-1}ba^{-1}+b^{-1}a^{-1}+b^{-1}a^{-2}ba^{-1}+ab+1+ba^{-1}ba^{-1}+baba^{-1}ba^{-1}+b+bab+a^{-2 }ba^{-1}+a^{-1}+b^{-1}a)=0$$

论坛问题帖 | has_answers | Alireza Abdollahi | 2016-02-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018080 | MO-2323

Hamiltonian S^1 actions with isolated fixed points

I have in mind the following question for some time. Is there an example of a compact symplectic manifold with a Hamiltonian S^1-action with isolated fixed points, that does not admit a compatible S^1-invariant Kahler strucutre? One would say, of course there should be such an example. But I have not seen any... Added, 2009. Apparently this is an open problem. PS, 2019. Not anymore!

论坛问题帖 | has_answers | Dmitri Panov | 2009-10-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018081 | MO-232342

A cubic and six conics problem

I am an electrical engineer system. I live in Viet Nam. I am not a Mathematician. I construct and found a problem as follows: Let a cubic, and five conics (C_1), (C_2), (C_3), (C_4), (C_5). The Cubic and a conic (C_i) have six common points. Conic (C_i) and conic $(C_{i+1}$ and the cubic have three common points for $i=1,2,3,4$. Let P_1, P_2, P_3 lie on (C_1) and the cubic but don't lie on (C_2) and P_4, P_5, P_6 lie on (C_5) and cubic but don't lie on (C_4). Then $P_1, P_2, P_3, P_4, P_5, P_6$ lie on a conic. I need a proof? Could you help me give your proof?

论坛问题帖 | has_answers | Oai Thanh Đào | 2016-02-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018082 | MO-232430

A problem of four curves

This is a generalization of my previous question, a problem of a cubic and six conics. Let a curve (K) of degree m and three curves (C_i) of degree n, for $i=1,2,3$. Let (C_1) meets (K) at mn points. Let (C_2) meets (K) at mn points. Let (C_3) meets (K) at mn points. Let $(C_1), (C_2)$ and K have d common points. Let points $P_1, P_2, ...,P_{mn-d}$ lie on (C_1) and (K) but don’t lie on (C_2). Let $(C_2), (C_3)$ and K have $mn-d$ common points. Let points $Q_1, Q_2, Q_3,....,Q_d$ lie on C_3 and (K) but don’t lie on C_2. The problem: show that the mn points $P_1, P_2, ...,P_{mn-d}, Q_1, Q_2, Q_3,....,Q_d $ lie on a curve of degree n, where $mn-d < \frac{n^2+3n}{2}$ and $d < \frac{n^2+3n}{2}$. The first figure I, make with $m=3$, $n=2$ and $d=2, mn-d=4$ The second figure I, make with $m=4$, $n=2$ and $d=4, mn-d=4 $. In this case 8 points lie on a new conic. The third figure, I make with $m=4$, $n=3$ and $d=6, mn-d=12-6=6 $. In this case, I make curve C_1, C_2, C_3 = three cubics, one cubic = three line, the curve (K) = 4 line, the new curve is a conic through 8 points and a line share a same line of cubic 1.

论坛问题帖 | has_answers | Oai Thanh Đào | 2016-02-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018083 | MO-2328

When does the converse to Schur's Lemma hold?

Let R be a commutative ring, let A be an R-algebra, and let M be an A-module. If M is simple, then End$_{A-mod}(M)$ is a division ring. A common use is when R is the complex numbers $\mathbb{C}$, and M is such that End$_{A-mod}(M)$ is finite dimensional. Then End$_{A-mod}(M) = \mathbb{C}$. Under what circumstances (regarding R and/or A) is the converse true, that the endomorphism ring being a division ring, or being just R itself, implies that M is simple?

论坛问题帖 | has_answers | cdouglas | 2009-10-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018084 | MO-234762

A conjecture on six planes

When I read Cox's Theorem, and Clifford's Circle Theorem and Miquel six circles theorem, I found the conjecture as folowing. And I checked the conjecture by the Geogebra sofware, the conjecture is right. I am looking for a proof of the conjecture. Conjecture: Let four points P_1, P_2, P_3, P_4 on a plane σ. Let six planes σ_{12}, σ_{13}, σ_{14}, σ_{23}, σ_{24} such that the plane σ_{ij} through two points P_i and P_j. Let P_{ijk} be a common points of three planes $\sigma_{ij}, \sigma_{jk}, \sigma_{ki}$. Then show that four points $P_{123}, P_{124}, P_{134}, P_{234}$ lie on a plane. See also Cox's Theorem

论坛问题帖 | no_answers_at_retrieval | Oai Thanh Đào | 2016-03-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018085 | MO-235576

On even almost perfect numbers other than powers of two

(Note: This question is an improved version of and has been cross-posted from this MSE post.) Let $\sigma(x)$ denote the sum of the divisors of x. If $\sigma(x) = 2x - 1$, then we call x an almost perfect number. If b is an odd composite and $\dfrac{b^2 - 1}{\sigma(b^2) - b^2} = \sigma(2^r)$ with $r \geq 1$, then $M = {2^r}{b^2}$ is an even almost perfect number. Now, if $\sigma(2^r)$ is prime, then ${2^r}{\sigma(2^r)}$ is an even perfect number. In an answer to the linked MSE question, Giovanni Resta asserts that: If $b = 3^k$ for $k > 1$ then $\dfrac{b^2 - 1}{\sigma(b^2) - b^2} = q = 2$. Up to ${10}^8$ there are no other values of b that make q prime. However, by work of Antalan, it is known that $3 \nmid b$, if $\dfrac{b^2 - 1}{\sigma(b^2) - b^2} = \sigma(2^r)$ with $r \geq 1$. Hence, I cannot help but ask: (1) Is it possible to prove that if b is an odd composite and $\dfrac{b^2 - 1}{\sigma(b^2) - b^2} = q$ is prime, then necessarily $b = 3^k$ with $k > 1$? If it is not possible by current mathematical methods, what is the main obstruction? (2) If b is an odd composite and $r \geq 1$, is it possible to prove that $\dfrac{b^2 - 1}{\sigma(b^2) - b^2} \neq \sigma(2^r)$? If it is not possible by current mathematical methods, what is the main obstruction?

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2016-04-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018086 | MO-235800

A long-lasting conjecture of Pólya & Szegő

There is a conjecture by Pólya & Szegő (~1950, stated in p. 159 of their book Isoperimetric Inequalities in Mathematical Physics) which is as follows: "Of all n-gons of a fixed area, the regular n-gon minimizes the first Dirichlet eigenvalue." Surprisingly, this is still open (to my knowledge) for the general case. The only settled cases are the triangles and the quadrilaterals (see Henrot's survey). Is there any progress on the general case?

论坛问题帖 | has_answers | BigM | 2016-04-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018087 | MO-236420

A conjecture of Blakley and Dixon about odd powers of positive matrices

In a 1966 paper Blakley and Dixon conjecture the following. Let S be a symmetric matrix with nonnegative entries and let u be a unit vector with nonnegative entries. For integers $k\ge j$ both odd, $$\langle u, S^ku\rangle^j\ge \langle u, S^ju\rangle^k.$$ They verify this conjecture under the condition that RHS is sufficiently large. I am wondering if there has been any progress towards this conjecture. A year earlier, Blakley and Roy verify the $j=1$ case. This also implies that the conjecture is true whenever j divides k. There are no papers known to Google scholar citing this paper that solves the conjecture.

论坛问题帖 | no_answers_at_retrieval | MERTON | 2016-04-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018088 | MO-2369

In a Banach algebra, do ab and ba have almost the same exponential spectrum?

Let A be a complex Banach algebra with identity 1. Define the exponential spectrum $e(x)$ of an element $x\in A$ by $$e(x)= \{\lambda\in\mathbb{C}: x-\lambda1 \notin G_1(A)\},$$ where $G_1(A)$ is the connected component of the group of invertibles $G(A)$ that contains the identity. Is it true that $e(ab)\cup\{0\} = e(ba)\cup\{0\}$ for all $a,b \in A$? Equivalently, is it true that $1-ab$ is in $G_1(A)$ if and only if $1-ba$ is in $G_1(A)$, for all $a,b \in A$? Note: The usual spectrum has this property. Just an additional note: We have $e(ab)\cup\{0\} = e(ba)\cup\{0\}$ for all $a,b \in A$ if 1) The group of invertibles of A is connected, because then the exponential spectrum of any element is just the usual spectrum of that element. 2) The set $Z(A)G(A) = \{ab: a \in Z(A), b\in G(A)\}$ is dense in A, where $Z(A)$ is the center of A. (One can prove this). In particular, we have $e(ab)\cup\{0\} = e(ba)\cup\{0\}$ for all $a,b \in A$ if the invertibles are dense in A. 3) A is commutative, clearly. But what about other Banach algebras? Can someone provide a counterexample?

论坛问题帖 | has_answers | Malik Younsi | 2009-10-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018089 | MO-237796

What explains the asymptotic and the pattern in this sequence related to Riemann zeta zeros?

As the starting point for my experiment I assumed that the imaginary parts of the Riemann zeta zeros are of the form: $$\Im \{ \rho_n \} = \frac{2\pi}{\log x_n}$$ where x_n is unknown. Therefore I solved for x_n and got this integer sequence: $$\Large a_1(n) = (n+1) \left(\left[\frac{1}{e^{\frac{2 \pi }{\Im(\rho _{n+1})}}-1}\right]-\left[\frac{1}{e^{\frac{2 \pi }{\Im(\rho _n)}}-1}\right]\right)$$ where $\left[\text{number}\right]$ is the round or nearest integer function. This is a sequence starting: $a_1 =\;$2, 3, 0, 5, 0, 7, 0, 9, 0, 11, 0, 13, 0, 15, 0, 17, 0, 19, 0, 0, 22, 0, 0, 25, 0, ... $ \; a_1(n)$ I then deleted the zeros in that sequence to get: $a_2 =\;$2, 3, 5, 7, 9, 11, 13, 15, 17, 19, 22, 25, 27, 30, 33, 36, 39, 42, 45, 48, 51, 54, 57, 61, 64,... Now a search in The On-Line Encyclopedia of Integer Sequences (OEIS) suggests that the first few terms are sequence A092919. Therefore the sequence: $$b(k)=\sum _{n=1}^{k+2} \text{Round}[\log (n)]$$ where k is a natural number: $$k=1,2,3,4,5,...$$ agrees with the beginning of $a_2 \;$. $b = \;$ 2, 3, 5, 7, 9, 11, 13, 15, 17, 19, 22, 25, 28, 31,... With Odlyzko's list of the 100000 first Riemann zeta zeros I plotted the difference $a_2 - b$: . . . The smoothness is only apparent at a such zoomed out plot. Actually when zooming in at the beginning it looks like this: Just for clarity I plotted the sequence a_2 and its conjectured asymptotic b separately also: Mathematica 8 code for the plots: (*start*) (*Mathematica 8*) nn = 1000; Monitor[a1 = Table[(n + 1)*(Round[(Exp[1/(Im[ZetaZero[n + 1]]/(2*Pi))] - 1)^-1] - Round[(Exp[1/(Im[ZetaZero[n]]/(2*Pi))] - 1)^-1]), {n, 1, nn}], n] a2 = DeleteCases[a1, 0] b = Table[Sum[Round[Log[n]], {n, 1, k}], {k, 3, Length[a2] + 2}] cc = a2 - b; cc[[1 ;; 20]] Label g1 = ListLinePlot[a2, PlotStyle -> Red, PlotLabel -> "sequence a2"] g2 = ListLinePlot[b, PlotLabel -> "asymptotic b"] g3 = Show[g1, g2, PlotLabel -> "sequence a2 and asymptotic b superimposed"] g4 = ListLinePlot[cc, PlotLabel -> "difference between sequence a2 and asymptotic b"] GraphicsGrid[{{g1, g2}, {g3, g4}}] (*end*) Is the sequence

b asymptotic to sequence a_2 and what explains the large scale repeating pattern in the difference $a_2 - b$?

论坛问题帖 | has_answers | Mats Granvik | 2016-05-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018090 | MO-23859

Covers of Riemann surfaces which become arbitrary close in Teichmuller space

Suppose S and S' are two compact Riemann surfaces of genus g. Does there exist a sequence of genera $g_i \to \infty$ and covers S_i, S_{i}' of S,S', both of genus g_i, such that $d(S_i,S_{i}')\to 0$? Here d a "natural" distance function on Teichmuller space, of which I suppose there are many, but for definiteness let's take it to be induced by the Teichmuller metric. This question was asked to me by Rick Kenyon last year, and some brief thought on it got me nowhere.

论坛问题帖 | has_answers | David Hansen | 2010-05-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018091 | MO-238628

Prove this conjecture inequality

This following problem is from my Conjecture many years ago, Question : Let $a,b>0,n\in N^{+},n\ge 3$,such $$a^n+b^n+(2n+2)(ab)^n\le 2n$$ Conjecture: then $a+b\le 2$ or $a+b>2.a>0.b>0,n\ge 3$,then we prove $$a^n+b^n+(2n+2)a^nb^n-2n>0$$ or it suffuce to prove $$a^n+b^n+(2n+2)a^nb^n-2n>0\rm{when}~ a+b=2$$ This inequality is from this conjecture $$\sqrt[n]{\dfrac{x}{1+y}}+\sqrt[n]{\dfrac{y}{1+x}}\le 2,\rm{where}~ x+y\le 2n,n\ge 3,n\in N^{+}$$ it is not hard to prove this case $n=3,4$,but for $n\ge 5$, use this $$\sqrt[n]{\dfrac{x}{1+y}}=a,\sqrt[n]{\dfrac{y}{1+x}}=b$$ then $$x=\dfrac{a^n(b^n+1)}{1-(ab)^n},y=\dfrac{b^n(a^n+1)}{1-(ab)^n}$$take $x+y\le 2n\Longrightarrow a^n+b^n+(2n+2)(ab)^n\le 2n$

论坛问题帖 | has_answers | math110 | 2016-05-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018092 | MO-239355

Pro-V topology on a free group

Let $W=G_p*G_q$, where G_p and G_q are pseudovarieties of all finite p-groups and all finite q-groups respectively, with p and q fixed prime numbers. The pro-W topology on a group G is the unique group topology such that the set of normal subgroups N with G/N in W is a fundamental system of neighborhoods of the identity. Let $FG(A)$ be a finitely generated free group. Steinberg and Auinger proved that the pro-W closure of a finitely generated subgroup H of $FG(A)$ is also finitely generated. Question. Let H be a finitely generated subgroup of a finitely generated free group $FG(A)$. Is there an algorithm to compute a basis for the pro-W closure H given as input a generating set for H? We know that every minimal normal subgroup of a finite p-group G is isomorphic to $\mathbb{Z}_p$, hence G has a finite normal series $$1=N_1\leq N_2\leq \ldots N_m=G$$ such that $N_{i+1}/N_{i}\simeq \mathbb{Z}_p$. Hence, we can write $G_q=\bigcup_{i=1}^{\infty} V^i$ where V is the locally finite pseudovariety generated by $\mathbb{Z}_q$ and $V^i={V*V*\ldots*V}$. By Proposition 4.6 in "HALL VARIETIES OF FINITE SUPERSOLVABLE GROUPS", the G_p*V-closure of a finitely generated subgroup H of $FG(A)$ is computable. By induction on i, we can prove that the G_p*V^i- closure of H is computable. So, we have a chain $$Cl_{Gp*V}(H)\geq Cl_{Gp*V^2}(H)\geq\ldots\geq Cl_{Gp*V^n}(H)\geq \ldots $$ of subgroups of $FG(A)$ containing $Cl_{G_p*G_q}(H)$. By Theorem 7.5 and Theorem 5.14 in "the geometry of profinite graphs with applications to free groups and finite monoids", every pseudovariety G_p*V^i is Hall pseudovariety that means the automaton of $Cl_{G_p*V^i}(H)$ is a quotient of the automaton of H. Since H is a finitely generated subgroup, the automaton of H is finite and so, the chain must stop somewhere and therefore there is i such that $Cl_{G_p*G_q}(H)=Cl_{Gp*V^i}(H)$. But the question is "Is there any bounded on i?

论坛问题帖 | no_answers_at_retrieval | user182085 | 2016-05-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018093 | MO-239671

Can we solve the FGF problem by finding an appropriate action?

If we can find an action of the free group $\mathbb{F}_2$ on a measure space X such that the crossed product $M=L^∞(X)⋊\mathbb{F}_2$ is a ${\rm III}_1$ factor with core isomorphic to $L(\mathbb{F}_2)⊗B(H)$, then the fundamental group of $L(\mathbb{F}_2)$ is $\mathbb{R}_+$, and the free group factors isomorphism problem is solved. Remark: There is an attempt here. Question: Can we rule out to solve this problem like that? Remark: One obstruction could be that M admits a Cartan subalgebra whereas $L(\mathbb{F}_2)$ not. To validate this obstruction, it is necessary to prove that the process of taking the core and to truncate it by a projection of trace 1, keeps the existence of a Cartan subalgebra. If we can rule out this method: Can we improve it by replacing $L^∞(X)$ by a more appropriate von Neumann algebra? More generally: Could we expect to be able to build by hand a ${\rm III}_1$ factor with core $L(\mathbb{F}_2)⊗B(H)$?

论坛问题帖 | has_answers | Sebastien Palcoux | 2016-05-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018094 | MO-24144

Is there an uncountable, non-discrete, Hausdorff Toronto space?

We call a topological space X a Toronto space if for any subspace $Y \subseteq X$ such that Y and X have the same cardinality it follows that Y is homeomorphic to X. Does anybody know what is known about the following question?: Is there an uncountable, non-discrete, Hausdorff Toronto space? It is not hard to show that if X is countable, Hausdorff and Toronto then X has the discrete topology. I have been thinking about the uncountable case for a while and it turns out it is a much harder question.

论坛问题帖 | has_answers | Manuel Rivera | 2010-05-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018095 | MO-24281

Current status of Bloch constant and Landau constant bounds

The Bloch constant B (based on a theorem introduced by André Bloch in 1925 on the maximum radius of a one-to-one disk in the image of a normalized analytic function of the unit disk, see for instance Remmert "Funktionentheorie II" or Steven Finch marvelous "Mathematical constants") was conjectured by Ahlfors to be $$ \frac{1}{\sqrt{1+\sqrt{3}}} \frac{\Gamma(\frac{1}{3})\Gamma(\frac{11}{12})}{\Gamma(\frac{1}{4})}$$ (This value, if I remember well Ahlfors' article corresponds to a particular function that he constructed for this purpose). The Bloch Constant B is currently known to be at least slightly greater than $\frac{\sqrt{3}}{4}$ (several articles improving upon each other by Mario Bonk, Chen and Gauthier, Xiong). Has there been some progress since 1998 on the lower bound ? Same question for the closely related (univalent) Landau constant (quite often called Bloch-Landau constant, sometimes seen as B_∞) ? The conjectured upper bound is $$\frac{\Gamma(\frac{1}{3})\Gamma(\frac{5}{6})}{\Gamma(\frac{1}{6})}$$ What can be said of the various adaptations or specializations of this constant to various class of functions, and extensions of these constants to several complex variables or other functional spaces ? I give as background the original article from Bloch, Ahlfors and Grunsky. (1) A. Bloch, Les théorèmes de M. Valiron sur les fonctions entières et la théorie de l'uniformisation, Ann. Fac. Sci. Univ. Toulouse, vol. 17, (1925), pp1-22. (2) L. V. Ahlfors and H. Grunsky, Über die Blochsche Konstante, Math. Zeitschrift 42 (1937), pp671-673. (3) L. V. Ahlfors, An extension of Schwarz's lemma, Trans. Amer. Math. Soc. 43 (1938), pp359-364. (these two are reprinted in Ahlfors Works vol 1) Ahlfors life and works are evocated in an [AMS Notices][1] of 1998.

论坛问题帖 | has_answers | ogerard | 2010-05-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018096 | MO-243330

If $N = q^k n^2$ is an odd perfect number, and $n < q^{k+1}$, does it follow that $k > 1$?

Let $N = q^k n^2$ be an odd perfect number with Euler prime q. According to Dickson (as pointed out recently by Beasley), Descartes conjectured $k=1$ in a letter to Mersenne in 1638, with Frenicle's subsequent observation occurring in 1657. In recent times, Sorli (page 89) conjectured that $k=1$ in his PhD thesis titled Algorithms in the study of multiperfect and odd perfect numbers. Following Brown's recent announcement of a proof for $q < n$, and a partial proof that $q^k < n$ holds under many cases, I recently realized that the biconditional $$q < n \iff \sigma(q) < \sigma(n) \iff \dfrac{\sigma(q)}{n} < \dfrac{\sigma(n)}{q}$$ would follow suit, if Brown's proof for $q < n$ is correct. (Similarly, $$q^k < n \iff \sigma(q^k) < \sigma(n) \iff \dfrac{\sigma(q^k)}{n} < \dfrac{\sigma(n)}{q^k}$$ would follow, if $q^k < n$ is true.) In this MSE post, I asked the following question (back in March 2014): If $N = q^k n^2$ is an odd perfect number, if $n < q^{k+1}$, does it follow that $k > 1$? Brown actually proves $q < n$ by proving $k = 1 \implies q < n$, so that we obtain that the implication $$k = 1 \implies \sigma(q) = \sigma(q^k) < n$$ holds, since $q < n < q+1 = \sigma(q)$ is impossible. By the contrapositive, $n < \sigma(q^k) \implies k > 1$. My question here is: To what extent can we then prove that $n < q^{k+1} \implies k > 1$? I note that $n < \sigma(q^k) \implies n < q^{k+1}$, but the converse may not necessarily be true. Notice that the implication $$n < q^{k+1} \implies k > 1$$ will prove that $$q^2 < n,$$ since it is known that $$n < q^2 \implies k = 1.$$

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2016-06-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018097 | MO-24352

Cutting convex sets

Any bounded convex set of the Euclidean plane can be cut into two convex pieces of equal area and circumference. Can one cut every bounded convex set of the Euclidean plane into an arbitrary number n of convex pieces having equal area and circumference? The solution of this problem for $n=2$ is generically unique. Are there other values of n (assuming that the problem is possible) where this happens? More generally, given a $d-$dimensional bounded convex set C in the Euclidean space $\mathbf E^d$ of dimension d. for which values of n can one cut C into n convex pieces of equal area with boundaries of equal $(d-1)-$dimensional area? ($n=2$ is again easy, but the solution is no longer generically unique if $d>2$). Are there values for (n,d) for which the solution always exists in a generically unique way (or for which the number of solutions is generically finite)?

论坛问题帖 | has_answers | Roland Bacher | 2010-05-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018098 | MO-243771

Primes and the factorial

Conjectures: For all prime numbers $p>5$ there exist a prime number $q<p$ such that $q\equiv m!\!\pmod p$, for some m with $2<m<p$. and For all primes p there is a prime $q<p^2$ such that $q\equiv m!\pmod {p^2}$, for some m with $2<m<p^2$. (Even $2<m<p$ as pointed out in comment). Both tested for all $p<100,000,000$. There are reasons to believe that the conjectures are true. If p is a large prime there are a lot of nonzero solutions to $n\equiv m!\!\pmod p$ and to $n\equiv m!\!\pmod {p^2}$. The probability for one of those n to be a prime increase with p. https://math.stackexchange.com/questions/1838931/conjecture-about-primes-and-the-factorial

论坛问题帖 | has_answers | Lehs | 2016-07-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018099 | MO-244984

An inequality in cyclic polygon and tangential polygon

I proposed my conjecture, it is strengthened version of the Erdős-Mordell inequality as following: Let $A_1A_2.....A_n$ be a cyclic polygon and $B_1B_2....B_n$ be the its tangential polygon. Let P be an arbitrary point inside of $A_1A_2.....A_n$. Let d_i be the distances from P to A_iA_{i+1}, for $i=1,...,n$ and $A_{i+1}=A_1$, and D_i be the distances from P to B_iB_{i+1}, for $i=1,...,n$ and $B_{i+1}=B_1$. Then show that: $$\sum_{1}^{n}{ D_i} \ge sec{\frac{\pi}{n}}\sum_{1}^{n}{d_i} $$ Equality holds when

$A_1A_2....A_n$ be the regular polygon. A proof of the case polygon is a triangle in the paper A strengthened version of the Erdős-Mordell inequality", Forum Geometricorum, 16: 317-321, MR 3556993. I am looking for a proof of general case.

论坛问题帖 | no_answers_at_retrieval | Oai Thanh Đào | 2016-07-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018100 | MO-245016

An inequality of real continuous function with f'>0 and f''>0

I proposed my conjecture as follows: Let $f(x)$ is a real continuous function on $[m, M]$ and $f'>0, f''>0$ on $[m, M]$, let $m \le x_i \le M$, for $i=1, 2,..., n$. Then $$\frac{f(x_1)+f(x_2)+.....+f(x_n)}{n}-f(\frac{x_1+x_2+....x_n}{n}) \le \frac{f(M)+f(m)}{2}-f(\frac{M+m}{2}) $$ Equality holds if only if $m=x_1=x_2=....=x_n=M$

论坛问题帖 | has_answers | Oai Thanh Đào | 2016-07-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018101 | MO-245092

A conjecture generalization of Karamata inequality

Fist I observe function $f(x)=x^2$ in the figure as following I found that when $x_1 \ge y_1$ and $x_2 \le y_2$ $\Rightarrow$ $AB \ge CD$ $\Rightarrow$ $$\frac{f(x_1)+f(x_2)}{2}-f(\frac{x_1+x_2}{2}) \ge \frac{f(y_1)+f(y_2)}{2}-f(\frac{y_1+y_2}{2})$$ (1). Note that: When $x_1+x_2=y_1+y_2$ the inequality (1) is Karamata inequality (with case n=2) $$f(x_1)+f(x_2) \ge f(y_1)+f(y_2)$$ From above observation, I am looking for a proof of a conjecture generalization of Karamata inequality as following: Let I be an interval of the real line and let f denote a real-valued, convex function defined on I. If $x_1, . . . , x_n$ and $y_1, . . . , y_n$ are numbers in I such that: $x_1 \ge x_2 \ge x_3...\ge x_n,$ and $y_1 \ge y_2 \ge y_3...\ge y_n$ 2.x_1+x_2+...+x_i \ge y_1+y_2+...+y_i for i=1,...,n-1 and 3. x_n \le y_n 2'. $x_1+...+x_i \ge y_1+...+y_i$ and $x_{i+1}+...+x_n \le y_{i+1}+...+y_n$ for $i=1,...,n-1$ then $$\frac{f(x_1)+f(x_2)+...+f(x_n)}{n}-f(\frac{x_1+x_2+...+x_n}{n}) \ge \frac{f(y_1)+f(y_2)+...+f(y_n)}{n}-f(\frac{y_1+y_2+...+y_n}{n}) $$ The inequality holds with equality if and only if $x_i=y_i$ for all $i \in {1, 2,...,n}$

论坛问题帖 | has_answers | Oai Thanh Đào | 2016-07-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018102 | MO-245516

Does every separable Banach space have a Markushevich-Auerbach basis?

Let X be a separable Banach space and X^* be its dual, let $\{x_i\}$ be a sequence in X with dense linear span and such that there exists a sequence $\{x_i^*\}$ in X^* satisfying $x_i^*(x_j)=\delta_{i,j}$ (Kronecker delta). It is clear that $\{x_i^*\}$ is uniquely determined. Let us call $\{x_i\}$ a Markushevich-Auerbach basis if the linear span of $\{x_i^*\}$ is weak* dense in X^* and $1=\|x_i\|_X=\|x_i^*\|_{X^*}$ for all i. Question: Is the problem about the existence of a Markushevich-Auerbach basis in an arbitrary infinite-dimensional separable Banach space still open? I am asking this question since I noticed that some recent papers (see Theorem 3.3 in http://arxiv.org/pdf/1604.03547.pdf) claim a solution of this problem, but I do not understand the argument.

论坛问题帖 | no_answers_at_retrieval | August Cleaner | 2016-07-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018103 | MO-2460

Important (interesting) unsolved problems

I think it would be interesting to have a list of important unsolved problems in mathematics. What are the important (interesting) problems in your field of work? It would be especially nice, to have a list of "non-mainstream-problems" in the miscellaneous areas of mathematics.

论坛问题帖 | has_answers | Spinorbundle | 2009-10-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018104 | MO-249567

Looking for an appropriate reference(s) for two conjectures on odd perfect numbers

(I apologize in advance if this question is unsuitable for MO. If so, please let me know and I will migrate it to MSE.) Let $\sigma(M)$ be the sum of the divisors of the positive integer M. For example, $\sigma(6)=1+2+3+6=12$. A number N is called perfect if $\sigma(N)=2N$. Euler proved that an odd perfect number N must have the form $N=q^k n^2$ where q is prime with $q \equiv k \equiv 1 \pmod 4$. We call q the Euler prime of N. Here are my questions: (1) Who is credited with making the conjecture that q is the largest prime factor of N? (2) Additionally, to whom should I attribute the conjecture that $k=1$? Should it be Descartes, Frenicle, or Sorli? (Added September 28 2016 For (2): In an edit to this MO post, it is stated that (per Beasley), "Dris $\ldots$ refers to Descartes' and Frenicle's claim (that $k=1$) as Sorli's conjecture; Dickson has documented Descartes's conjecture as occurring in a letter to Marin Mersenne in 1638, with Frenicle's subsequent observation occurring in 1657". As commented by Gerry, one would need to double-check Dickson's History of Number Theory to verify Beasley's statements.)

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2016-09-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018105 | MO-249569

Remaining models conjectured to converge to SLE(6) or CLE(6)

I am wondering which models are conjectured (eg. numerically) to converge to SLE(6) (Schramm-Loewner evolution with $\kappa=6$) or CLE(6) (conformal loop ensemble). I am searching for a research topic in this very interesting area. The following are from Smirnov's 2007 review and elsewhere: 1) Critical percolation for lattices other than triangular eg. the square lattice. Any detailed review on this? 2) Fortuin-Kasteleyn random cluster model for $n=1$ and $x>x_{c}$, where x_{c} the critical edge weight. 3) For $O(1)$ but for $x\neq 1$ and $p\neq 1/2$, which correspond to the critical percolation case. 4) In the article "conformal invariance of hydrodynamic turbulence", they showed numerically that

zero-vorticity isolines correspond to SLE(6). 5) Random-field Ising model by Stevenson (2011).

论坛问题帖 | no_answers_at_retrieval | user133100 | 2016-09-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018106 | MO-251145

What arguments do exist against defining completeness in NP using injective Karp reductions?

It is crucial to use the right notion of reduction to define completeness inside NP. Different notions of completeness inside NP may have significant impact on the properties of complete languages. Initially, P vs NP problem was formulated by Cook using polynomial-time Turing reductions. Later Karp refined the notion of completeness using polynomial-time many-one reductions. It is not known whether complete problems are equivalent under different notions of completeness (Cook vs Karp reductions). This post on CS Theory shows that defining completeness using injective Karp reductions would prove $P \ne NP$. The reason is that SAT is dense language and can not be reduced to a sparse language using injective Karp reductions. Mahaney's Theorem states that $P=NP$ if and only if there is a Karp reduction from SAT to a sparse language. Also, one of the answers highlights the fact that all known Karp reductions to NP-complete problems are injective. There is no known natural NP-complete problem via Karp reduction that can not be made one-to-one. If Cook had defined NP-completeness using injective Karp reductions then the P vs NP problem could not existed. What arguments do exist against defining completeness in NP using injective Karp reductions? Proving the existence (or non-existence) of polynomial-time algorithms for SAT would be more satisfactory way to resolve P vs NP but it is important to link it to the existence (or non-existence) of incomplete languages inside NP. Ladner's theorem states the $P\ne NP$ if and only if there exists an incomplete language in NP \ P (under Karp reductions).

论坛问题帖 | has_answers | Mohammad Al-Turkistany | 2016-09-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018107 | MO-251557

Localizing Bondy's metaconjecture on hamiltonicity

Definitions: Let G be a graph on n vertices. G is Hamiltonian provided G has a cycle of length n. G is pancyclic provided G has a cycle of length ℓ for every $3 \leq \ell \leq n$. Background: Dirac [1952] showed that if G has $n \geq 3$ vertices and every vertex has degree at least $\frac{n}{2}$, then G is Hamiltonian. Bondy [1971] showed this condition implies that either G is the complete bipartite graph $K_{\frac{n}{2},\frac{n}{2}}$ or G is pancyclic. Bondy [1973] then proposed the following metaconjecture (which I quote from Kevash and Sudakov 2010): Metaconjecture. Almost any non-trivial condition on a graph which implies that the graph is Hamiltonian also implies that the graph is pancyclic. (There may be a simple family of exceptional graphs.) Another instance of this metaconjecture is the following corollary of a theorem of Clark [1981]. Let G be a connected, locally connected, claw-free graph on at least 3 vertices. Then G is pancyclic. Given property $\mathcal{P}$ (such as "connected"), a graph is locally $\mathcal{P}$ provided the neighborhood of every vertex is $\mathcal{P}$. A graph is claw-free provided it has no induced copies of $K_{1,3}$. Being claw-free is equivalent to being locally $\overline{K_3}$-free. It seems to me that such local properties might eliminate bothersome exceptional cases. Might we be able to prove a localized version of Bondy's metaconjecture such as the following? Tentative Subconjecture. Let $\mathcal{P}$ be a graph property such that every connected graph on at least 3 vertices that is locally $\mathcal{P}$ is Hamiltonian. Then every connected graph on at least 3 vertices that is locally $\mathcal{P}$ is pancyclic. A monotonicity assumption on the property--for example, if A satisfies $\mathcal{P}$ and A is a subgraph of B and $|A|=|B|$ then B satisfies $\mathcal{P}$--might also be useful. (This question is inspired by Ortrud Oellermann's presentation at the 2016 Prairie Discrete Math Workshop.)

论坛问题帖 | has_answers | D. Ror. | 2016-10-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018108 | MO-251996

Is the Gromov conjecture still open?

Today I read about Gromov's definition of minimal volume for smooth manifolds. $$\min {\rm Vol}(M):=\inf_{|K_g|\leq1}\{{\rm Vol}(M,g)\}.$$ Gromov's conjecture states that for every closed simply connected odd-dimensional manifold $\min {\rm Vol}(M)=0$. Is the Gromov conjecture still open? Can anybody give an example for this conjecture? Update: My teacher told to me that this conjecture can be solved by the Ricci flow method but I don't know how to use it. Can anybody give me an explanation to this? and how it works? Thanks.

论坛问题帖 | has_answers | C.F.G | 2016-10-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018109 | MO-252424

Current state of Straus's illumination problem

In George W. Tokarsky's Polygonal Rooms Not Illuminable from Every Point (1995) it is stated that the problem Is a polygonal region illuminable from at least one point in the region? was still open at the time. What is its current state? Has it been settled or is it still open?

论坛问题帖 | has_answers | AlterTim | 2016-10-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018110 | MO-253088

Expectation of changing the gift choice

Suppose we are given two boxes, with one of gift valued n dollars and the other one valued twice as much. We can pick a box, and after open it we have the choice of switching to another box. Shall we switch? I was asked about this during some graduate student seminar. My reasoning is since two choices are equally likely, it is equally likely I will get a box half of the original value and a box twice the original value. So I should switch as the net expectation is positive. However, this implies that my original choice is somehow inferior to the choice after switch, which violates symmetry principle. Can someone help?

论坛问题帖 | has_answers | Zhexiu Tu | 2016-10-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018111 | MO-25316

Square powers of hemicontinuous operators

Let H be an infinite dimensional real Hilbert space. A [not necessarily linear] mapping of H into itself is said to be hemicontinuous if it is continuous from each line segment of H to the weak topology of H (F.E. Browder / G.J. Minty). [Obviously, any linear operator is hemicontinuous.] Intuitively speaking, this is an extremely weak continuity requirement, still very useful, e.g., in the study of nonlinear elliptic boundary value problems. Now, here is my problem. Let U be an arbitrary (i.e., possibly discontinuous) selfmap of H. Is it true that there exists some hemicontinuous selfmap of H, say V, such that U 2 = V 2 ?

论坛问题帖 | no_answers_at_retrieval | Ady | 2010-05-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018112 | MO-254323

Any counter example for this: ${\phi(2^n-1)} \bmod \tau(2^n-1)=0$ for every integer $n \geq 1$?

I asked this question here In S.E but i don't received any resposnes for it, I would like to know if it is appropriate for M.O. I'm always interesting for properties of the following series : $ \sum_{n=1}^{\infty }\frac{\tau(2^n-1)}{\phi(2^n-1)}$ such that: $\tau(N)$ is the number of divisors of N and ϕ is Euler totient function, In this question i w'd be interest to know a little bit about the reciprocal of each term of the titled series at $N=2^n-1$ for every $n\geq 1$. Note:01 According to my calculations that i run in wolfram alpha from $n=1$ to 100 I have got this result as shown here such that it works which letting me to ask the following question. My question here is: Is ${\tau(2^n-1)}$ always divides $\phi(2^n-1)$ for every integer $n\geq 1$ and any counterexample for it? Thank you for any help !!!

论坛问题帖 | no_answers_at_retrieval | zeraoulia rafik | 2016-11-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018113 | MO-255384

A conjecture about algebraic values of $(-q;\,-q)_\infty/(q;\,q)_\infty$

Recall that $(a;\,q)_\infty$ is the q-Pochhammer symbol: $$(a;\,q)_\infty=\prod_{n=0}^\infty(1-a \, q^n).\tag1$$ Its important special case $(q;\,q)_\infty=\prod_{n=1}^\infty(1-q^n)$ is sometimes called the Euler function. It appears in Euler's pentagonal number theorem, and its reciprocal $(q;\,q)_\infty^{-1}$ is the generating function for the partition numbers. It is also related to Jacobi theta functions and Ramanujan theta functions. Let $$f(x) = \frac{(-q;\,-q)_\infty}{(q;\,q)_\infty},\quad\text{where}\,\,q=e^{-\pi\sqrt x}.\tag2$$ In the OEIS entry A080054 there is an empirical observation by Simon Plouffe that apparently $f(1)=\sqrt[8]2$. Empirical (Simon Plouffe, Feb. 20, 2011): $$\sum_{n=0}^{\infty}e^{-\pi n}a(n) =\sqrt[8]2.$$ I did some numerical experiments related to this observation, and the outcomes suggest a fascinating stronger conjecture: Conjecture: For every $p\in\mathbb Q,\,p>0$, the value $f(p)$ is an algebraic number. For example, it appears that $$f(3/5) = \sqrt[8]{2} \cdot \sqrt[4]{9 \sqrt{5}+5 \sqrt{15}-11 \sqrt{3}-19},$$ and $f(13/7)$ is an algebraic number of degree 96 whose minimal polynomial is $$x^{96}-647442063456 \, x^{88}+16702438371168 \, x^{80}-529345497357824 \, x^{72}+4159684203040512 \, x^{64}-12099397290541056 \, x^{56}+16408771708010496 \, x^{48}-10607690933600256 \, x^{40}+2651923007078400 \, x^{32}-367001600 \, x^{24}+257949696 \, x^{16}-100663296 \, x^8+16777216$$ and an isolating rational interval is $(37/36,\,6/5)$. Is this conjecture new? Is it known to be true? If not, can you suggest any ideas how to (dis-)prove it?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2016-11-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018114 | MO-257015

Decidable open problems

Are there any significant open problems in mathematics which are clearly decidable (in that it is easy to write a clearly corresponding program which will eventually output either Yes or No (or whatever sort of answer, out of finitely many possibilities, is appropriate), though it may take an implausibly long time to do so) but which remain open? Dropping the qualifier "significant", examples of this sort of thing would be determining whether chess between perfect players results in a white win, black win, or stalemate; determining the $10^{10^{100}}$th decimal digit of π; etc. But none of these are of particular significance in mathematics, such that anyone would ordinarily list them as an open problem of note. So, though it is inherently a subjective judgment: Are there any good examples of significant open problems of this sort? (Edit: I nominate this question for reopening as NOT a duplicate, in that the question it has been marked a duplicate of specifically excludes problems that "naturally resolve after a finite computation", being interested only in problems which nontrivially reduce to finite computation. My interest is in either, but for me, the ideal examples are those that are manifestly finite computations from the start!)

论坛问题帖 | has_answers | Sridhar Ramesh | 2016-12-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018115 | MO-257494

The derivative of an integral function with indicator and max function as integrand

I encounter the following type of problem: \begin{equation} F(x) = \int_a^b \mathbf{1}_{\{v+x-h(v)\geq 0\}}\max\{h(v)-y-x,0\}dv \end{equation} where $\mathbf{1}_{\{z\geq 0\}}=1$ if event $z\geq 0$ occurs, and zero otherwise. $y>0$ is constant. I am not sure how to apply fundamental theorem of calculus (or may be something else?) to get $\frac{dF(x)}{dx}$. I think in $\frac{dF(x)}{dx}$, it should contain something like $\delta (x)$?

论坛问题帖 | no_answers_at_retrieval | kim kevin | 2016-12-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018116 | MO-257628

Theorems demoted back to conjectures

Many mathematicians know the Four Color Theorem and its history: there were two alleged proofs in 1879 and 1880 both of which stood unchallenged for 11 years before flaws were discovered. I am interested in more such examples, especially, in former (famous or interesting) "theorems" where no remedy could be found to the present day.

论坛问题帖 | has_answers | Moritz | 2016-12-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018117 | MO-258097

there exist infinite many $n\in\mathbb{N}$ such that $S_n-[S_n]<\frac{1}{n^2}$

Let $S_n:=1+\frac12+\frac13+\ldots+\frac1n$. Is it true that the set of $n\in\mathbb N$ such that $$S_n-[S_n]<\dfrac{1}{n^2}$$ is infinite? Here, $[x]$ represents the largest integer not exceeding x. This question has been asked previously on math.SE without receiving any answers.

论坛问题帖 | no_answers_at_retrieval | math110 | 2016-12-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018118 | MO-258353

Proving $k = 1 \implies q = 5$, if $q^k n^2$ is an odd perfect number with Euler prime q

Let $N = q^k n^2$ be an odd perfect number with Euler prime q. We want to show that the biconditional $k = 1 \iff q = 5$ holds. It suffices to prove one direction, as the implication $q = 5 \implies k = 1$ was proved by Iannucci (Lemma 12, page 873). Per a comment from Pace: "Jose, you are misinterpreting the result from Iannucci. His number N is restricted to be an odd perfect number which additionally has all but at most two of its prime divisors less than 100. He goes on to prove that there are no such numbers, but along the way, in the lemma you cite, he shows that such numbers have restricted form." To this end, we show first that $$k = 1 \implies I(n^2) \leq 2 - \frac{5}{3q}.$$ The proof is easy. We refer the reader to this preprint. Next, we observe that $$I(n^2) = 2 - \frac{5}{3q}$$ implies that $$k = 1 \land q = 5,$$ (The assertion that follows only holds when $k=1$.) while $$I(n^2) < 2 - \frac{5}{3q}$$ implies $q > 5$. In particular, we have $$I(n^2) \leq 2 - \frac{5}{3q} \implies \bigg(\left(k=1\right) \lor \left(q>5\right)\bigg).$$ But we know that $$\bigg(\left(k=1\right) \lor \left(q>5\right)\bigg) \iff \bigg(q = 5 \implies k = 1\bigg).$$ Consequently, we have the chain of implications $$q = 5 \implies k = 1 \implies q = 5 \implies k = 1,$$ and we are done. Now here is my question: Is this "proof" logically sound? Update (December 30 2016) As in the given answer, we could only have the implication $$k = 1 \implies \bigg(q = 5 \implies k = 1\bigg),$$ which I agree is not the same as $$k = 1 \implies q = 5 \implies k = 1.$$

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2016-12-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018119 | MO-259278

An explicit series representation for the analytic tetration with complex height

Tetration is the next hyperoperation after more familiar addition, multiplication and exponentiation. It can be seen as a repeated exponentiation, similar to how exponentiation can be seen as a repeated multiplication, and multiplication - as a repeated addition. Tetration is denoted $^n a$, where a is called the base and n is called the height, and is defined for $n\in\mathbb N\cup\{-1,\,0\}$ by the recurrence $$ {^{-1} a} = 0, \quad {^{n+1} a} = a^{\left({^n a}\right)},\tag1$$ so that $${^0 a}=1, \quad {^1 a} = a, \quad {^2 a} = a^a, \quad {^3 a} = a^{a^a}, \, \dots \quad {^n a} = \underbrace{a^{a^{{.^{.^{.^a}}}}}}_{n\,\text{levels}},\tag2$$ where power towers are evaluated from top to bottom. To simplify matters, we restrict our attention to real bases in the interval $1<a<e^{1/e}$. Under this restriction, the sequence $\{^n a\}$ is strictly increasing and converges to a limit: $$^\infty a=\lim_{n\to\infty} {^n a} = \frac{W(-\ln a)}{-\ln a},\tag3$$ where $W(z)$ denotes the principal branch of the Lambert W-function. To simplify our following exposition it is convenient to introduce a new variable q linked to a: $$q = \ln({^\infty a}) = {^\infty a} \cdot \ln a = -W(-\ln a).\tag4$$ The bounds on a imply $0<q<1$. The value q is closely related to the asymptotic growth of the tetration. In particular, it appears in remarkable limits: $$ \lim_{n\to\infty} \, \frac{{^{n+k} a}-{^n a}}{{^{n-k} a}-{^n a}} = -q^k, \quad \lim_{n\to\infty} \, \frac{{^{n+k} a}-{^\infty a}}{{^n a}-{^\infty a}} = q^k .\tag5$$ Definitions of addition, multiplication and exponentiation for fractional, negative and complex arguments are well-established. A natural extension of the tetration to fractional and complex heights is a long-standing question. It appears to be much less straightforward problem, mainly because exponentiation, unlike addition and multiplication, lacks both commutativity and associativity, and so tetration does not obey simple laws similar to $a^m \cdot a^n = a^{m+n}$ or $(a^m)^n = a^{m\,n}$. Despite these obstacles, several proposals aiming to extend the definition of the tetration to fractional and complex heights have been put forth. Their common goal is to define a function that agrees with the discrete tetration for all positive integer heights, and satisfies the functional equation $${^{z+1} a = a^{\left({^z a}\right)}\tag6$$ for all z in its domain. These conditions alone are not sufficient to determine a unique function, so additional natural restrictions (such as continuity, smoothness or analyticity) have been proposed, aiming to narrow down the set of candidate functions to a single and, in some sense, "the most natural" extension. The Wikipedia article on tetration is very cautious and indicates that, apparently, so far there is no consensus about which extension is "the one". Some of the proposals seem to lead to the same function, although apparently no rigorous proof of their equivalence has been published yet. The Citizendum article is written in a much bolder style, and refers to tetration as a unique well-defined function. But, overall, the study of tetration seems to be a quite isolated area of mathematics, with few connections to other topics. A few months ago I made some interesting conjectures related to tetration that I hoped I would prove soon. It appears that there are some connections between tetration and the theory of q-analogs. Unfortunately, I have not made much progress in proving them since then (although I have learnt a lot about q-series), so I finally decided to post them here and ask for your help. First, let's recall some definitions. The q-Pochhammer symbol: $$(p;\,q)_\infty = \prod_{k=0}^\infty(1-p\,q^k)\tag7$$ $$(p;\,q)_z = \frac{(p;\,q)_\infty}{(p\,q^z;\,q)_\infty},

\quad z\in\mathbb C\tag8$$ It easily follows that $$(p;\,q)_n = \prod_{k=0}^{n-1}(1-p\,q^k), \quad n\in\mathbb N\tag9$$ The q-binomial coefficients (also known as the Gaussian binomial coefficients): $${r \brack s}_q = \frac{(q;\,q)_r}{(q;\,q)_s \, (q;\,q)_{r-s}}.\tag{10}$$ From now on we assume that a and, consequently, q are fixed. Let us define an analytic function $t(z)$ by the following series: $$ t(z) = \sum_{n=0}^\infty \sum_{k=0}^n (-1)^{n-k} \, q^{\binom {n-k} 2} {z \brack n}_q {n \brack k}_q ({^k a}).\tag{11} $$ This formula can be seen as a combination of the direct and reverse q-binomial transforms. It can be proved by induction that for $n\in\mathbb N, \, t(n)={^n a}$, so $t(n)$ satisfies the functional equation for tetration (6) at least for positive integer arguments. Indeed, partial sums of (11) can be seen as Lagrange interpolating polynomials in powers of q^z that exactly reproduce tetration values for progressively larger initial segments of $\mathbb N$. Numeric evidence suggests that adding new points to the interpolating polynomial of this form does not result in erratic oscillations of its arcs between existing points (i.e. Runge's phenomenon does not occur), but rather makes them converge to a monotone function satisfying the functional equation for the tetration (6). So we have: Conjecture 1. The series $t(z)$ converges at least in the half-plane $\Re(z)>-2$ to an analytic function that satisfies the functional equation for tetration $t(z+1)=a^{t(z)}$ for all arguments in its domain of analyticity. Conjecture 2. The function $t(z)$ is periodic with a purely imaginary period $\tau = 2\pi n i/\ln q$, and admits an analytic continuation to the half-plane $\Re(z) \le -2$ by repeated backward application of the functional equation (6). There it has a countable set of isolated singularities at $z = m + \tau\,n, \,m,n\in\mathbb Z, \, m \le -2$, and a countable set of branch cuts that can be made along horizontal rays at $z = x + \tau\,n, \, n\in\mathbb Z, \, x\in\mathbb R, \, x < -2$. Conjecture 3. For real $x>-2$, the derivative $t'(x)$ is a completely monotone function. Conjecture 4. There is only one analytic function that vanishes at $z=-1$ and satisfies both the functional equation for the tetration (6), and the complete monotonicity condition from the Conjecture 3. Related questions: [1][2][3].

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2017-01-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018120 | MO-259467

Reconstructing analytic tetration with a complex height from a thinner set of points

This is a follow-up to my previous question An explicit series representation for the analytic tetration with complex height. Recall the definition (11) from there: $$t(z) = \sum_{n=0}^\infty \sum_{k=0}^n (-1)^{n-k} \, q^{\binom {n-k} 2} {z \brack n}_q {n \brack k}_q ({^k a}).\tag1$$ Let us introduce additional parameters $p,s\in\mathbb Z^+$ into this formula (changes are shown in red): $$t(\color{red}p,\color{red}s,z) = \sum_{n=0}^\infty \sum_{k=0}^n (-1)^{n-k} \, q^{\,\color{red}p \, \binom {n-k} 2} {(z-\color{red}s)/\color{red}p \brack n}_{q^{\color{red}p}} {n \brack k}_{q^{\color{red}p}} ({^{\color{red}p\,k+\color{red}s} a}).\tag2$$ Basically, it means that we build our function from a thinner and shifted sample of tetration values. Conjecture 1. Values of integer parameters p,s do not matter, i.e. $$\forall p,s\in\mathbb Z^+, \, t(p,s,z) = t(z).$$ Let us generalize the definition of $t(p,s,z)$ to real-valued parameters $p,s\in\mathbb R^+$ (note that we switch to continuous tetration $t(z)$ in the last factor): $$t_{\mathbb R}(p,s,z) = \sum_{n=0}^\infty \sum_{k=0}^n (-1)^{n-k} \, q^{\,p \, \binom {n-k} 2} {(z-s)/p \brack n}_{q^p} {n \brack k}_{q^p} \color{red}t(p\,k+s).\tag3$$ Conjecture 2. Values of real parameters p,s do not matter, i.e. $$\forall p,s\in\mathbb R^+, \, t_{\mathbb R}(p,s,z) = t(z).$$ Can we prove these conjectures? If yes, it means that we do not need all values of the discrete tetration (represented by the factor $(^k a)$ in (1)) to build a formula for the analytic tetration, but a very thin (although still infinite) subset of them already contains enough information to exactly reconstruct its values at all points in between (including skipped integer points). Can we make a step further and find a way to construct an explicit series for the analytic tetration that relies only on a finite set of values of the discrete tetration?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2017-01-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018121 | MO-259637

A seemingly simple inequality

Let $a_i,b_i\in\mathbb{R}$ and $n>1$, does the inequality $$ \left(\sum_{i=1}^n a_i^2\right)\left(\sum_{i=1}^n b_i^2\right)+\left(\sum_{i=1}^na_i b_i\right)^2\ge \sqrt{\left(\sum_{i=1}^n a_i^4\right)\left(\sum_{i=1}^n b_i^4\right)}+\sum_{i=1}^na_i^2b_i^2 $$ hold true? Observe that a_i and b_i are not required to be non-negative. I ran an extensive number of numerical simulations and no counterexample showed up yet. Note 1. The inequality holds true for $n=2$, as showed here. Note 2. This conjecture was formulated by Fedor Petrov in an attempt to provide a solution to a particular case of this question. Note 3. I have posted this question on math.SE some time ago but it has received no answer, so I cross-posted it here. Note 4. As Fedor Petrov rightly observed in his answer below, the inequality also follows by an argument used in one of his answers in the above-cited question. However, I decided to accept Markus Sprecher's answer because of its conciseness and clarity.

论坛问题帖 | has_answers | Ludwig | 2017-01-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018122 | MO-260379

A curious series related to the asymptotic behavior of the tetration

The tetration is denoted $^n a$, where a is called the base and n is called the height, and is defined for $n\in\mathbb N\cup\{-1,\,0\}$ by the recurrence $$ {^{-1} a} = 0, \quad {^{n+1} a} = a^{\left({^n a}\right)},\tag1$$ so that $${^0 a}=1, \quad {^1 a} = a, \quad {^2 a} = a^a, \quad {^3 a} = a^{a^a}, \, \dots \quad {^n a} = \underbrace{a^{a^{{.^{.^{.a}}}}}}_{n\,\text{levels}},\tag2$$ where power towers are evaluated from top to bottom. Let a be a real number in the interval $1<a<e^{1/e}$. It is convenient to introduce a notation $$\lambda = -W(-\ln a),\tag3$$ where $W(z)$ denotes the principal branch of the Lambert W-function, satisfying $W(z) \, e^{W(z)}=z$. Note that the restrictions on a imply $0<\lambda<1$. We may also observe that $\ln\ln a = - \lambda + \ln \lambda$. The sequence $\{^n a\}$ converges to a limit $$\lim_{n\to\infty} {^n a} = e^\lambda = \frac\lambda{\ln a}.\tag4$$ Its asymptotic behavior can be represented as $${^n a} = e^\lambda - c_{\lambda} \cdot \lambda^n + O(\lambda^{2n}),\tag5$$ where c_{λ} is some positive coefficient depending on λ (or, equivalently, on

a). It is known that the coefficients in $O(\lambda^{2n})$ and all higher-order terms can be expressed in a closed form via c_{λ}. But, apparently, the dependency between λ and c_{λ} has no known simple representation. Here is how its graph looks: It appears to have a maximum near (or exactly at?) $\lambda = \ln 2$ that corresponds to $a=\sqrt2$. Numerical evidence suggests that c_λ has a Taylor-Maclaurin series expansion with rational coefficients $$c_\lambda = \frac{\lambda }{1!}+\frac{\lambda ^2}{2!}-\frac{2 \lambda ^3}{3!}-\frac{11 \lambda ^4}{4!}-\frac{44 \lambda ^5}{5!}-\frac{89 \lambda ^6}{6!}-\frac{636 \lambda ^7}{7!}-\frac{615 \lambda ^8}{8!}-...\tag6$$ (more numerators can be found here) I could not find a formula for the coefficients (apparently, they are not yet in the OEIS), and I do not even have a proof that the coefficients given above are exact, so I am asking for your help with it. Is it possible to sum this series in terms of known special functions? Is it actually unimodal as is suggested by its graph? What is the exact location of its maximum? Related questions: [1][2][3][4][5][6].

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2017-01-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018123 | MO-260511

Can there be a polymath project for mathematical physics?

My hunch is that it might be possible to create something like https://polymathprojects.org/ for mathematical physics and I'd like to know whether MathOverflow users can recommend some appropriate problems. As Abdelmalek Abdessalam pointed out in the comments below such a platform can potentially be used to develop research programs in mathematical physics that are suitable for polymath type collaborative work. The gist of this project would be to use collaborate problem-solving to tackle open problems in mathematical physics. Timothy Gowers wrote a seminal blog post on the subject here: Is massively collaborative mathematics possible? There's a good point made by a commenter below as to why not simply use the current polymath platform. Well, I think that it would be easier for such a platform to succeed and gain support among mathematical physicists if it was maintained by mathematical physicists and focused mainly on problems in the area of mathematical physics. This is not the current setup of the Polymath project.

论坛问题帖 | has_answers | Aidan Rocke | 2017-01-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018124 | MO-260783

What is the status on this conjecture on arithmetic progressions of primes?

The Green-Tao theorem states that for every n, there is an arithmetic sequence of length n consisting of primes. For primes, p, let $P(p)$ be the maximum length of an arithmetic progression of primes whose least element is p. Is it known whether $P(p)=p$ for every prime? (This clearly generalizes the Green-Tao theorem, asserting that long progressions show up "as soon as possible." Note that $P(p) \leq p$ by viewing the progression mod p.)

论坛问题帖 | has_answers | Gorka | 2017-01-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018125 | MO-261408

Several conjectured identities for polylogarithms

I asked a question at M.SE a couple of years ago about polylogarithms $\!^{[1]}$$\!^{[2]}$$\!^{[3]}$$\!^{[4]}$ where I conjectured $$720\,\operatorname{Li}_4\!\left(\tfrac12\right)-2160\,\operatorname{Li}_4\!\left(\tfrac13\right)+2160\,\operatorname{Li}_4\!\left(\tfrac23\right)+270\,\operatorname{Li}_4\!\left(\tfrac14\right)+540\,\operatorname{Li}_4\!\left(\tfrac34\right)+135\,\operatorname{Li}_4\!\left(\tfrac19\right)\\ =19\pi^4+30\left(\pi^2-\beta^2\right)\left(10\alpha^2-12\alpha\beta+3\beta^2\right)-30\,\alpha^2\left(19\alpha^2-24\alpha\beta+8\beta^2\right)\!,\\ \text{where $\alpha=\ln2,\,\beta=\ln3$.}$$ I put a bounty on it recently, but it still remains unanswered. So, I decided to re-post it here. Also, I recently discovered several similar conjectures for $\operatorname{Li}_5(z)$ involving powers of the golden ratio $\phi=\frac{1+\sqrt5}2$: $$4455\,\operatorname{Li}_5\left(\phi^{-2}\right)-1215\,\operatorname{Li}_5\left(\phi^{-4}\right)-360\,\operatorname{Li}_5\left(\phi^{-6}\right)+15\,\operatorname{Li}_5\left(\phi^{-12}\right)\\ =3015\,\zeta(5)-702\ln^5\phi+180\,\pi^2\ln^3\phi-38\,\pi^4\ln\phi$$ $$45000\,\operatorname{Li}_5\left(\phi^{-2}\right)-16875\,\operatorname{Li}_5\left(\phi^{-4}\right)-144\,\operatorname{Li}_5\left(\phi^{-10}\right)+9\,\operatorname{Li}_5\left(\phi^{-20}\right)\\ =28944\,\zeta(5)-6000\ln^5\phi+1600\,\pi^2\ln^3\phi-356\,\pi^4\ln\phi$$ $$15660\,\operatorname{Li}_5\left(\phi^{-2}\right)-19440\,\operatorname{Li}_5\left(\phi^{-4}\right)+7680\,\operatorname{Li}_5\left(\phi^{-6}\right)+2430\,\operatorname{Li}_5\left(\phi^{-8}\right)-15\,\operatorname{Li}_5\left(\phi^{-24}\right)=5025\,\zeta(5)+2088\ln^5\phi-240\,\pi^2\ln^3\phi-28\,\pi^4\ln\phi$$ How can we prove these conjectures? Are there other similar identities of this kind?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2017-02-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018126 | MO-261649

A conjecture about certain values of the Fabius function

The Fabius function is a smooth monotone function $F:[0,1]\to[0,1]$, satisfying functional equations $$F(0)=0, \quad F(1-x)=1-F(x)\tag1$$ and $$F'(x) = 2 \,F(2 x) \quad \text{for} \,\, 0<x<1/2.\tag2$$ The function F assumes rational values at dyadic rational arguments. In particular, it is known $\!^{[1]}$$\!^{[2]}$ that $$F\left(2^{-n}\right) = \frac1{n! \, 2^{\binom{n+1}2}} \, \sum_{m\ge0}\binom n {2m} \, c_m,\tag3$$ where c_m are defined by the recurrence $$c_0 = 1, \quad c_n = \frac1{(4^n - 1)(2n+1)} \, \sum_{m\ge1} \binom{2n+1}{2m+1} \, c_{n-m}.\tag4$$ Note that only finite number of terms in each sum are non-zero. The values $F\left(2^{-n}\right) $ appear as A272755 / A272757 in the OEIS. Let $$a_n = F\left(2^{-n}\right) \, 2^{\binom {n-1}2} \, (2n)! \, \prod_{m=1}^{\lfloor n/2\rfloor}\left(4^m - 1\right).\tag5$$ This sequence begins $$1, \, 5, \, 15, \, 1001, \, 5985, \, 2853675, \, 26261235, \, 72808620885, \, 915304354965 \, ...\tag6$$ (see more terms here) I conjecture that all terms of this sequence are integers. How can we prove (disprove) this conjecture?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2017-02-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018127 | MO-262456

Primes dividing functions defined by linear recurrence relations with constant coefficients

For Fibonacci numbers F_n it holds that $p|F_{p-(\frac{5}{p})}$, if p is an odd prime (Legendre symbol). I guessed that the number 5 came from the roots of the characteristic polynomial and tried the following approach: Given $a,b\in\mathbb Z^+$ (for simplicity), and let $F_{a,b}:\mathbb N\to\mathbb N$ be a function such that $F_{a,b}(0)=0$, $F_{a,b}(1)=1$ and $F_{a,b}(n+1)=a\cdot F_{a,b}(n)+b\cdot F_{a,b}(n-1)$. Conjecture: If a,b are co-prime and $p>2$ is a prime that doesn't divide b, then $F_{a,b}\left(p-\left(\frac{a^2+4b}{p}\right)\right)\equiv 0\pmod p$, Is this conjecture correct and known? I have tested the conjecture with 50000 randomly selected a,b,p. That is, three numbers $1\le a',b'\le 100$ and $3\le p'\le 10000$ are selected with uniform distribution, then $a=\frac{a'}{gcd(a',b')},b=\frac{b'}{gcd(a',b')}$ and p is set to the least prime $\ge p'$.

论坛问题帖 | no_answers_at_retrieval | Lehs | 2017-02-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018128 | MO-26330

Dissecting trapezoids into triangles of equal area

[Lightly edited for copy and proper formatting of mathematics. -- Pete L. Clark] The Background: Let T be a trapezoid. Sherman Stein, using valuation theory, showed that if T is dissectible into n triangles of equal area, then $\frac{n}{(r+1)}$, where r is the ratio of the parallel sides, is an algebraic integer. See Monsky and Jepsen, Constructing equidissections for certain classes of trapezoids, Discrete Mathematics, 308 23 (2008) 5672-5681. In particular r is algebraic. If r is in $\mathbb{Q}$, a diagonal of T divides T into two triangles of commensurable area, so T has an "equidissection". Stein showed that if r is a root of a degree 2 element of $\mathbb{Q}[z]$ [i.e., the univariate polynomial ring over $\mathbb{Q}$ -- PLC] with both roots positive, then T is equidissectible. Using ideas of Stein and Jepsen I showed that if r is a root of a degree 3 or 4 element of $\mathbb{Q}[z]$ with all roots in the open right half-plane, then once more T is equidissectible. (It's not clear that my proof can be adapted to degree greater than 4). In contrast, whenever the algebraic r has a conjugate over $\mathbb{Q}$ that's not in the open right half-plane, nothing is known about the equidissectibilty of T. Stein conjectured that when r has a $\mathbb{Q}$-conjugate not in this half plane, then T is not equidissectible. I ask: The Question: Is it true that if r has a $\mathbb{Q}$-conjugate not in the open right half-plane then T cannot be dissected into triangles of equal area? (For no such r has an equidissection of T or a proof of non-equidissectibility been found).

论坛问题帖 | no_answers_at_retrieval | paul Monsky | 2010-05-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018129 | MO-26358

Can we color Z^+ with n colors such that a, 2a, ..., na all have different colors for all a?

For example for n=2 coloring odd numbers red, numbers of the form 4k+2 blue and so on works. This problem was posed in the KoMaL for n+1 prime, by Peter Pach Pal. I verified it for all n<30, I think with a computerprogram one can easily verify it for much bigger numbers by trying certain periodic colorings. Ideas. As there is a lot of discussion going on, I thought I share here my attempts as Ewan and Gowers took similar paths. First of all, if we denote the primes by p_i and the largest prime not bigger than n by p_d, then it is sufficient to color the numbers of the form $\Pi p_i^{\alpha_i}$. This is equivalent to coloring $\mathbb Z^d$ with n colors such that the translates of a special poliomino are all rainbow colored, meaning they contain all n colors. This is also equivalent to tiling the space with translates of this poliomino. The easiest way to give a coloring is if we have some nice periodicity, eg. if n+1 is prime, then $\Pi p_i^{\alpha_i} \mod (n+1)$ is such, whenever we go in a direction, it corresponds to an authomorphism of Z_{n+1}^*. Another possibility is to give a "linear" coloring using the addition $\mod n$, for example for $n=5$, one can take $x+3y+4z \mod 5$. So far I could always find such a linear coloring but I cannot prove that it always exist, eg. we have too many constraints to use the combinatorial nullstellensatz.

论坛问题帖 | has_answers | domotorp | 2010-05-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018130 | MO-263982

Maximum matching in a graph with no "shortcuts"

For a directed acyclic graph (DAG) G, denote by $G^\star$ the undirected graph obtained from G by ignoring direction of its arcs. Let $e(G)=e(G^\star)$ be the number of arcs in G (or edges in $G^\star$). An arc (u,v) in G is called a shortcut if there exists a directed path from u to v in G different from (u,v). In our research, we came up with the following conjecture: Conjecture. Let G be a DAG without shortcuts such that the indegree and outdegree of every vertex is each ≤ 2. Then there exists a matching in $G^\star$ of size at least $\frac{1}{4}e(G)$. Small examples seem to support this conjecture, although a general proof appears quite elusive. The lower bound $\frac{1}{5}e(G)$ here is almost trivial (and holds even with shortcuts), and with some effort we were able to prove the bound $\frac{2}{9}e(G)$. Any help in proving or disproving the conjecture will be appreciated. P.S. Apparently it is crucial that $G^\star$ results from a DAG. E.g., a similar statement for a generic undirected graph of degree at most 4 and no cycles of length 3 (which follows from the absence of shortcuts G) does not hold.

论坛问题帖 | has_answers | Max Alekseyev | 2017-03-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018131 | MO-26525

Coiling Rope in a Box

What is the longest rope length L of radius r that can fit into a box? The rope is a smooth curve with a tubular neighborhood of radius r, such that the rope does not self-penetrate. For an open curve, each endpoint is surrounded by a ball of radius r. For a box of dimensions

$1{\times}1{\times}\frac{1}{2}$ and rope with $r=\frac{1}{4}$, perhaps $L=\frac{1}{2}+\frac{\pi}{4} \approx 1.3$, achieved by a 'U': I know packing circles in a square is a notoriously difficult problem, but perhaps it is easier to pack a rope in a cube, because the continuity of the curve constrains the options? (I struggle with this every fall, packing a gardening hose in a rectangular tub.) I am more interested in general strategies for how to best coil the rope, rather than specific values of L. It seems that if r is large w.r.t. the box dimensions (as in the above example), no "penny-packing" cross-sectional structure is possible, where one layer nestles in the crevices of the preceding layer. This is a natural question and surely has been explored, but I didn't find much. Edit 1. It seems a curvature constraint is needed to retain naturalness: The curve should not turn so sharply that the disks of radius r orthogonal to the curve that determine the tubular neighborhood interpenetrate. Edit 2 (26Jun10). See also the MO question concerning decidability. Edit 3 (12Aug10). Here is an observation on the 2D version, where a $1 {\times} 1 {\times} 2r$ box may only accommodate one layer of rope. If $k=\frac{1}{2r}$ is an even integer, then I can see two natural strategies for coiling the rope within the box: $\qquad \qquad \qquad \qquad \qquad$ Red is rope core curve, blue marks the rope boundary. Interestingly, if I have calculated correctly, the length of the red rope curve is identical for the two strategies: $$L = 2 (k-1)[r \pi/2] + 2(k-1)^2 r \;. $$ For $r=\frac{1}{16}$, $k=8$ as illustrated, $L=\frac{7\pi}{16} + \frac{49}{8} \approx 7.5$. (As a check, for $r=\frac{1}{4}$, $k=2$, and L evaluates to $\frac{\pi}{4}+\frac{1}{2}$ as in the first example above.)

论坛问题帖 | has_answers | Joseph O'Rourke | 2010-05-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018132 | MO-265401

Surface bundles over surfaces with(out) flat structure

I vaguely remember that I once attended a seminar or conference talk in which it was mentioned that the following question is open. Is there a (smooth) surface bundle over a surface $\Sigma_h \to E \to \Sigma_g$ that does not admit a flat structure? (Equivalently, one can ask whether there is a homomorphism $\pi_1(\Sigma_g) \to \pi_0 \text{Diff}(\Sigma_h)$ that cannot be lifted to the group of diffeomorphisms.) What is the status of this question, and could somebody point out a reference that mentions this as an open question? Finally, is something known for small values of g,h (but still $g,h \geq 2$)? (In case you do not understand what I am asking, I can very much recommend reading the last chapter of Morita's book Geometry of Characteristic Classes.)

论坛问题帖 | has_answers | Jens Reinhold | 2017-03-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018133 | MO-266242

Hausdorff's question on ω_1-gap

I read here that the following problem of Hausdorff is apparently still open. Is there a maximal branch C in the poset ω^ω with the eventual domination order, such that C has no ω_1-gap. My questions concerning the above problem are: What is its current status? What are the consequences of its solution? negative, independence or positive.

论坛问题帖 | has_answers | Rahman. M | 2017-04-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018134 | MO-267017

Generalization of Legendre`s conjecture

Legendre`s conjecture states that there is always a prime between n^2 and $(n+1)^2$ for every natural n. It is natural to create following generalization: Is it true that for every $\varepsilon \in (0,1]$ there exists natural number $n_0(\varepsilon)$ such that for every $n \geq n_0(\varepsilon)$ there is always a prime between $n^{1+\varepsilon}$ and $(n+1)^{1+\varepsilon}$? I do not know if some results from number theory do not allow this generalization to be true but if it is not known is it true or not what is known, if anything, about this generalization of Legendre`s conjecture?

论坛问题帖 | has_answers | Paladin | 2017-04-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018135 | MO-267936

Is the fundamental group of any compact hyperbolic 3-manifold embeddable into a p-adic group?

Is it true that for every compact hyperbolic 3-manifold M there exists a prime p, a finite field extension $K/\mathbb{Q}_p$, and an injective group homomorphism $$\tau \colon \pi_1(M) \to \mathrm{PSL}_2(\mathcal{O}_K) ?$$ Here $\mathcal{O}_K$ is the ring of integers of K. This question is obviously related to the existence of faithful low dimensional representation of hyperbolic groups (see the following survey by Aschenbrenner, Friedl, and Wilton). An affirmative solution will give a (partial) confirmation of Conjecture 1 from this paper by Luo. A paper by Calegari and Emerton also discusses such group homomorphisms τ. Note that the answer is obviously 'yes' if M is arithmetic.

论坛问题帖 | has_answers | Pablo | 2017-04-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018136 | MO-26821

Is Thompson's Group F amenable?

Last year a paper on the arXiv (Akhmedov) claimed that Thompson's group F is not amenable, while another paper, published in the journal "Infinite dimensional analysis, quantum probability, and related topics" (vol. 12, p173-191) by Shavgulidze claimed the exact opposite, that F is amenable. Although the question of which, if either, was a valid proof seemed to be being asked by people, I cannot seem to find a conclusion anywhere and the discussion of late seems to have died down considerably. From what I can gather, Shavgulidze's paper seems unfixable, while the validity of Akhmedov's paper is undecided (although it may of course be decided now). So, does anyone know if either of these papers is valid?

论坛问题帖 | has_answers | ADL | 2010-06-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018137 | MO-268929

Is there a composite number for which this game with composite numbers never ends?

I created some "game" a couple of days ago and modified it during the last hour or so and now I want to present it to you. Choose some natural number >1. If the chosen natural number is of the form p^n, where $p \in \mathbb P$ and $n \in \mathbb N$ then the game is over. But if the chosen number has $k \geq2$ different prime factors (abbreviated as dpf in the rest of the text) then we add to that number the next number which also has k dpf. If after adding we arrive at the number of the form p^n then the game is over. But if after adding we arrive at a number that has $l \geq 2$ dpf then we add to it the next number which has l dpf. We continue in this way until we arrive at the number of the form p^n and then the game is over. To clarify if something is not clear, let us take as an example the number 12. It has two dpf and the next number that has two dpf is 14 so we arrive at $12+14=26$ which has two dpf, the next number that has two dpf is 28 so we arrive at $26+28=54$ which has two dpf and the next one with two dpf is 55 so we arrive at $54+55=109$ which is prime and the game is over. Does there exist at least one composite number for which this game is never over? This seems (at least to me) as a very hard question and I would like to call the composite numbers for which the game never ends an awesome numbers, although maybe the number of those numbers equals zero, but that is yet to be proved.

论坛问题帖 | no_answers_at_retrieval | Living Hell | 2017-05-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018138 | MO-269354

Which motivic cohomology groups of complex numbers are non-torsion?

I would like to know which motivic cohomology groups of complex numbers are non-zero and ("better") non-torsion, i.e., for which (i,j) the ith cohomology of the complex ${\mathbb{Q}}(j)$ over $\mathbb{C}$ is not zero. I would like to know both which of these groups are known to be non-zero (and also known to be "large") unconditionally and also which of them shouldn't vanish according to some conjectures. Certainly, my question can be easily translated into the language of K-theory.

论坛问题帖 | has_answers | Mikhail Bondarko | 2017-05-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018139 | MO-270616

Open bilinear maps that are not uniformly open

A map $f\colon X\to Y$ between metric spaces is uniformly open whenever for each $\varepsilon >0$ there is $\delta >0$ such that for any $x\in X$ one has $$B_Y\big(f(x),\delta\big)\subseteq f\big(B_X(x,\varepsilon)\big), $$ where B_X, B_Y denote open balls in the respective spaces. Uniformly open maps have the property that they are surjective as long as the codomain is connected. The exponential function is an example of an open map that is not uniformly open. A continuous, surjective bilnear map $A\colon X\times X\to Y$ between Banach spaces need not be open, contrary to the linear case. (Already matrix multiplication is an example; another example is multiplication in $C[0,1]$ in the case of real scalars; this is due to Fremlin). Is there an example of a continuous, bilinear and surjective map between Banach spaces that is open but not uniformly open? I have a suspicion that it could be already convolution in $\ell_1(\mathbb Z)$ but I am not sure how to approach this (in the latter case I know that it is not uniformly open as $\ell_1(\mathbb{Z}/n\mathbb{Z})$ do not have equi-uniformly open convolutions). Edit (27.06.2017). This question was also asked in a paper by Balcerzak, Behrends, and Strobin (Banach J. Math. Anal. 10 (2016), no. 3, 482-494). I would also welcome examples of n-linear maps with this property, where n is arbitrary.

论坛问题帖 | has_answers | Tomasz Kania | 2017-05-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018140 | MO-270728

Solve this diophantine equation: $m^4+n^4=10m^2n^2+1$

t's probably common knowledge that there are Diophantine equations which do not admit any solutions in the integers, but which admit solutions modulo nn for every nn. This fact is stated, for example, Conjecture Let $m,n $be integer,find all $$m^4+n^4=10m^2n^2+1$$ I this equality have only foursolution$(m,n)=(0,1),(0,-1),(1,0),(-1,0)$.But I can't prove it

论坛问题帖 | has_answers | math110 | 2017-05-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018141 | MO-272386

Random N-body problem

Suppose there are N unit-mass particles whose initial positions are uniformly distributed in a unit-radius disk. Each particle is assigned a randomly oriented initial velocity vector v_i of length 1 (green below). Added: Robert Israel's incisive comment suggests that I should also stipulate that $\sum_i v_i = 0$. Then the particles act upon one another via inverse-square gravity. Dots show initial positions inside unit disk. Green vectors: initial velocity; red vectors: final velocity. Q. What is the probability that $k \ge 1$ of the N particles remain within a disk of some radius $R \ge 1$ forever? In the illustration above, $N=8$ and $R=3$. (But I do not trust my crude simulations.) I am wondering if the answer is: zero, independent of k and R and the gravitational constant? Perhaps the answer differs for points in $\mathbb{R}^2$ vs. points in $\mathbb{R}^3$ (or $\mathbb{R}^d$, $d > 3$)?

论坛问题帖 | has_answers | Joseph O'Rourke | 2017-06-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018142 | MO-275011

Open problems in mathematical physics

What are good, still unsolved problems in mathematical physics that are in vogue? I always get the same answers: reference to Millennium Problems by the Clay Institute, or "there's still a lot to do in this or that..." or the three body problem. Could you be more specific?

论坛问题帖 | has_answers | CristinaSardon | 2017-07-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018143 | MO-27716

Does there exist a comprehensive compilation of Erdos's open problems?

Fan Chung and Ron Graham's book Erdos on Graphs: His Legacy of Unsolved Problems (A. K. Peters, 1998) collects together all of Erdos's open problems in graph theory that they could find into a single volume, complete with bounties where applicable. Of course Erdos posed many other open problems in combinatorics and number theory that do not appear in this book. I once heard a rumor that some people were working on a project to publish a similar but more comprehensive book or series of books, covering all of Erdos's open problems, but I don't know if the rumor is true. Does such a compilation exist? If not, is there anything else like this besides Chung and Graham's book?

论坛问题帖 | has_answers | Timothy Chow | 2010-06-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018144 | MO-277495

Limit cycles as closed geodesics(2)

Hilbert 16th problem asks for a uniform upper bound $H(n)$ for the number of limit cycles of a polynomial vector field of degree n on the plane. Here is an updated proof of the finitness part of the Hilbert 16th problem: http://www.mathnet.ru/php/archive.phtml?wshow=paper&jrnid=im&paperid=8352&option_lang=eng This problem is open even for $n=2$. In this question, by a quadratic vector field we mean a polynomial vector field $$P(x,y)\partial_x+Q(x,y) \partial_y\;\;\;(V)$$ where $P,Q\in \mathbb{R}[x,y]$ are polynomials of degree 2 with $P(0,0)=Q(0,0)=0$. So the question is that (Hilbert 16th problem for n=2) Is $H(2)$ a finite number? As a possible approach to this question we would like to look at limit cycles of a vector field as closed geodesics. Namely we would like to put a Riemannian metric on the (regular part of the) phase space such that the trajectories of the vector field would be unparametrized geodesics. By regular part we mean the complement of singularities of the vector field. Thus each limit cycle would be a closed geodesic. Then using Gauss Bonnete theorem we try to count the number of closed geodesics. So the sign of the Gaussian curvature play a very important role. But the first problem is that, regardless of the curvature sign, is there a Riemannian metric on the regular part of the phase space such that the trajectories of the vector field would be geodesic. This example shows that, for a polynomial vector field of arbitrary degree such metric does not necessarily existed. But for quadratic vector field the situation is different as follows. In the following C is the algebraic curve $$yP(x,y)-xQ(x,y)=0\;\;\;\;\;(C)$$ Observation: There is a Riemannian metric on $\mathbb{R}^2 \setminus C$ such that all solutions of Quadratic vector field (V) are geodesics. Furthermore every limit cycle of $(V) $ which surround the origin can not intersect the curve C The proof of this observation is based on Proposition 6.7 and 6.8 as follows: Geometry of foliation So we are sure that, on the complement of the above algebraic curve C, we have a Riemanian metric such that all solutions of (V) are geodesics. The methods of the proof of the Propositions 6.7 and 6.8 in the book Geometry of foliation suggest that we choose a Riemannian metric whose orthonormal base is the following: $$\left\{\frac{x^2+y^2}{yP(x,y)-xQ(x,y)}V,\ \frac{1}{x^2+y^2}W\right\}.$$ where $V=P\partial_x +Q\partial_y$ is the quadratic vector field as in (V) and W is the radial vector field $W=x\partial_x+y\partial_y$. In fact W, as it is required in the proof of the above two propositions, lies in the kernel of $1_$ form $\psi=\frac{1}{x^2+y^2}(ydx-xdy)$. This $1_$ form is very essential to apply propositions 6.7 and 6.8 in the above reference. In fact this is a closed form which is identically equal to 1 on the first vector of the above orthonormal frame.That is $\psi(\frac{x^2+y^2}{yP-xQ}(V))=1$. I am really indebted to Ben McKay who suggested the $1_$form $d\theta$ as a required $1_$ form for possible satisfactions of proposition 6.8. On the other hand the method of the proof of the Proposition 6.7 shows that, on the complement of C, all trajectories of V are geodesics for the metric arising from the above orthonormal frame. Moreover we are free to rescale the second vector of the frame, arbitrarily. Now the main problem is that we control the sign of the curvature of such metric to make facilities to count the number of closed geodesics. In fact we use the Gauss Bonnete theorem. Call the curvature of this metric κ. Question: When a quadratic vector field V does not have a center on the plane, is the curve $$\{(x,y)\mid \kappa(x,y)=0\}$$ transverse to V? If not, what appropriate rescaling of the the second vector $ \frac{1}{x^2+y^2} W$ of the orthonormal frame would give a positive answer? Notes: 1)A center is a singularity which is surrounded by a band of closed orbits. For quadratic vector fields they are classified at this paper. 2)Of course existence of a positive answer to this question implies that $H(2)$ is finite. Because the number of limit cycles of V which suround the origin can be uniformly bounded since $\kappa=0$ is an algebraic curve so in every connected component of this algebraic curve, we have at most on limit cycle surrounding origin. On the other hand there are at most 2 nest of limit cycles. So this would imply that $H(2)<\infty$ The motivation for this post is mentioned in this answer and this post. However the initial motivation is mentioned in page 3, item 5 of this arxiv note.

论坛问题帖 | no_answers_at_retrieval | Ali Taghavi | 2017-07-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018145 | MO-277570

A cubic system with two nested limit cycles with opposite orientations(2)

The second part of Hilbert's 16th problem not only concerns "The number of limit cycles of a polynomial vector field", but also the position and configuration of of those limit cycles with respect to each other. However the primary and the usual interpretation of the concept "Relative Position" is that "Whether two given limit cycles are nested closed curves or not" but the following post is a motivation to consider other types of questions about configuration of limit cycles, for example "Can two nested limit cycles have opposite orientation?" A cubic system with two nested limit cycles with opposite orientations In this regard we focus on cubic polynomial system and ask the following question: What is an example of Cubic polynomial vector field $$\begin{cases} x'=P(x,y)\\ y'=Q(x,y) \end{cases}$$ such that two nested closed orbits C_1, C_2 of the system surrounds an annular region R such that R does not contain any singular point and the flow orientation of C_1 is opposite to the flow-orientation of C_2? There is a very interesting example of degree 5 here but we search for degree 3

论坛问题帖 | no_answers_at_retrieval | Ali Taghavi | 2017-07-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018146 | MO-27851

Polynomials having a common root with their derivatives

Here is a question someone asked me a couple of years ago. I remember having spent a day or two thinking about it but did not manage to solve it. This may be an open problem, in which case I'd be interested to know the status of it. Let f be a one variable complex polynomial. Supposing f has a common root with every $f^{(i)},i=1,\ldots,\deg f-1$, does it follow that f is a power of a degree 1 polynomial? upd: as pointed out by Pedro, this is indeed a conjecture (which makes me feel less badly about not being able to do it). But still the question about its status remains.

论坛问题帖 | has_answers | algori | 2010-06-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018147 | MO-27861

Periodic Automorphism Towers

In Scott's classic textbook on Group Theory, he asks: Suppose that G is a finite group. Is the sequence of isomorphism types of the groups $Aut^{(n)}(G)$ for $n \in \mathbb{N}$ eventually periodic? Here $Aut^{(2)}(G) = Aut(Aut(G))$ etc. Equivalently, is the sequence $|Aut^{(n)}(G)|$ always bounded above? It apparently remains opens whether the sequence of automorphism types of $Aut^{(n)}(G)$ is in fact always eventually constant. (A wonderful theorem of Wielandt says that if G is a finite centerless group, then the sequence is eventually constant.) So I would like to ask: Does there exists a finite group such that $Aut(G) \not\cong G$ but $Aut^{(n)}(G) \cong G$ for some $n \geq 2$? Edit: Joel has pointed out that my question is perhaps even open for infinite groups. This sounds like an interesting question which doesn't seem amenable to the standard tricks.

论坛问题帖 | has_answers | Simon Thomas | 2010-06-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018148 | MO-278686

Splitting of ordinals of oscillation ranks of a Baire 1 function

Denny and Tang proved that Theorem 2.3 Let (f_n) be a sequence in $\mathfrak{B}_1(K)$ converging pointwise to a function $f.$ Suppose $\sup\{ \beta(f_n):n\in\mathbb{N} \} \leq \beta_0$ and $\gamma((f_n)) \leq \gamma_0.$ Then f is Baire-1 and $\beta(f) \leq \beta_0 \cdot \gamma_0.$ They asked problem 4.7 (still open) in the same paper. Open Problem: Is it true that if $f\in\mathfrak{B}_1(K)$ with $\beta(f)\leq\beta_0 \cdot \gamma_0$ for some countable ordinals β_0 and $\gamma_0.$ then there exists a sequence (f_n) converging pointwise to f so that $\sup_n \beta(f_n) \leq \beta_0$ and $\gamma((f_n)) \leq \gamma_0?$ They showed that the problem has positive answer if $\beta_0 = 1$ (Theorem 5.5) or γ_0 is a limit ordinal (Theorem 4.5). Question: What are the known techniques to construct real-valued functions that satisfy $\sup_n\beta(f_n) \leq \beta_0$ and $\gamma((f_n)) \leq \gamma_0?$ Background and Definitions: Let K be a compact metric space. We say that $f:K\rightarrow \mathbb{R}$ is of Baire 1 if there exists a sequence of continuous functions (f_n) converging to f pointwise. The set $\mathfrak{B}_1(K)$ contains all real-valued Baire 1 functions f on K, both bounded and unbounded. For a Baire 1 function $f:K\rightarrow \mathbb{R}$, its oscillation rank β is defined as follows: For any $\varepsilon>0,$ let $K^0(f,\varepsilon) = K.$ If $K^\alpha(f,\varepsilon)$ is defined for some countable ordinal $\alpha,$ let $K^{\alpha+1}(f,\varepsilon)$ be the set of all those $x\in K^\alpha(f,\varepsilon)$ such that for every open set U containing $x,$ there are two points x_1 and x_2 in $U \cap K^\alpha(f,\varepsilon)$ with $|f(x_1) - f(x_2)| \geq \varepsilon.$ For a countable limit ordinal $\alpha,$ we let $$K^\alpha(f,\varepsilon) = \bigcap_{\alpha^{\prime}<\alpha}K^{\alpha^{\prime}}(f,\varepsilon).$$ The index $\beta_K(f,\varepsilon)$ is taken to be the least α with $K^\alpha(f,\varepsilon) = \emptyset$ if such an α exists, and ω_1 otherwise. The oscillation rank of f is $$\beta_K(f) = \sup\{ \beta_K(f,\varepsilon):\varepsilon>0 \}.$$ If (f_n) is a sequence of real-valued functions on $K,$ let $K^0((f_n),\varepsilon) = K$ for any $\varepsilon>0.$ If $K^\alpha((f_n),\varepsilon)$ has been defined for some countable ordinal $\alpha,$ let $K^{\alpha+1}((f_n),\varepsilon)$ be the set of all those $x\in K^\alpha((f_n),\varepsilon)$ such that for every open set U containing x and any $m\in\mathbb{N},$ there are two integers n_1,n_2 with $n_1>n_2 > m$ and $x' \in U \cap K^\alpha((f_n),\varepsilon)$ such that $|f_{n_1}(x') - f_{n_2}(x')| \geq \varepsilon.$ Define $$K^\alpha((f_n),\varepsilon) = \bigcap _{\alpha' < \alpha}K^{\alpha'}((f_n),\varepsilon)$$ if α is a countable limit ordinal. Let $\gamma_K((f_n),\varepsilon)$ be the least α with $K^\alpha((f_n),\varepsilon) = \emptyset$ if such α exists, and ω_1 otherwise. The convergence index of f is the ordinal $$\gamma_K((f_n)) = \sup\{ \gamma_K((f_n),\varepsilon): \varepsilon>0 \}.$$

论坛问题帖 | no_answers_at_retrieval | Idonknow | 2017-08-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018149 | MO-27944

Do there exist chess positions that require exponentially many moves to reach?

By "chess" here I mean chess played on an $n\times n$ board with an unbounded number of (non-king) pieces. Some care is needed if you want to generalize some of the subtler rules of chess to an $n\times n$ board, but I will not dwell on this point because the answer to the question I'm interested in should be the same under any reasonable generalization. Namely, does there exist an infinite sequence (A_n, B_n) of pairs of chess positions on an $n\times n$ board such that the minimum number of legal moves required to get from A_n to B_n is exponential in n? Here I allow any legal moves and not only strategically intelligent moves. Technically this question might be classified as an "open problem" (which is illegal on MO) because it was implicitly asked by A. Fraenkel and D. Lichtenstein in "Computing a perfect strategy for $n\times n$ chess requires time exponential in n," J. Comb. Th. A 31 (1981), 199-214. However, I think it is fair game for MO because I'm pretty confident that this has not been looked at much. Fraenkel and Liechtenstein showed that determining whether a given chess position is won for White (with best play) is EXPTIME-complete and asked for the computational complexity of the chess reachability problem ("is B_n reachable from A_n?"). Clearly chess reachability is in NPSPACE = PSPACE, and Hans Bodlaender has shown that it is NP-hard. If the answer to the question I've posed above is "No, it can always

be done with polynomially many moves" then it would solve this problem by showing that chess reachability is NP-complete, because exhibiting the sequence of moves yields a short certificate. If you have some experience with retrograde chess problems and if you've read Hearn and Demaine's lovely book Games, Puzzles, and Computation then you may get the intuition that shuffling chess pieces around is reminiscent of other rearrangement puzzles that have been shown to be PSPACE-complete. However, I've asked both Demaine and Hearn and neither of them saw immediately how to show that chess reachability is PSPACE-complete. [Edit: Searching more carefully through Hearn and Demaine's book, I see that they list this problem in their list of open problems at the end of the book under the name "Retrograde Chess." I didn't notice it before because for some reason that page is not listed in the index under "chess." I can perhaps be blamed for using the name "Retrograde Chess" for this problem because that's what I called it when I first posted this question to USENET way back when. I think that "reachability" is a better name for it.]

论坛问题帖 | has_answers | Timothy Chow | 2010-06-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018150 | MO-280123

The set of numbers $a+b$ such that ma^2+nb^2 is prime

Conjecture: If m,n are coprime it exist a minimal natural number N_{mn} such that: $\{a+b>N_{mn}\mid a,b\in\mathbb N^+\wedge ma^2+nb^2\in\mathbb P_{>2}\} = \{ k > N_{mn} \mid \gcd(k,m+n)=1\}$. Test for $k\leq 10^8$ suggest that $N_{11}=1$. The table below shows the largest minimal N_{mn} for $k\leq 10^5$. In the table 0 stands for $\gcd(m,n)>1$. 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 1 1 1 1 1 1 4 1 1 1 7 1 3 1 1 1 4 1 5 1 1 13 1 1 17 1 2 1 0 1 0 1 0 4 0 1 0 1 0 1 0 31 0 18 0 20 0 10 0 1 0 1 3 1 1 0 1 1 0 1 19 0 10 1 0 3 11 0 6 7 0 1 1 0 9 3 0 1 4 1 0 1 0 8 0 1 0 4 0 1 0 5 0 1 0 5 0 14 0 6 0 1 0 6 5 1 1 1 8 0 16 1 21 1 0 1 1 1 1 0 1 1 22 1 0 1 8 1 18 0 6 4 0 0 0 16 0 20 0 0 0 38 0 1 0 0 0 1 0 6 0 0 0 5 0 16 7 1 4 1 1 1 20 0 1 1 1 1 14 1 0 1 1 1 12 1 19 0 15 1 12 1 8 1 0 19 0 21 0 1 0 10 0 1 0 4 0 19 0 9 0 5 0 13 0 18 0 8 9 1 1 0 4 1 0 1 10 0 17 1 0 1 14 0 1 1 0 3 5 0 6 1 0 1 10 7 0 10 0 0 0 0 1 0 17 0 1 0 1 0 0 0 11 0 1 0 10 0 28 0 0 11 1 1 1 1 1 38 1 1 1 1 0 11 1 16 1 4 1 12 1 6 3 0 1 29 1 12 3 0 0 0 1 0 14 0 0 0 11 0 9 0 0 0 18 0 1 0 0 0 24 0 16 13 1 1 3 5 1 1 1 4 1 1 1 9 0 10 1 20 1 23 1 1 1 1 5 1 1 14 1 0 11 0 1 0 0 0 14 0 16 0 10 0 1 0 33 0 37 0 0 0 1 0 1 15 1 31 0 1 0 0 1 19 0 0 1 0 1 1 0 6 1 0 1 0 0 1 1 0 0 16 4 0 6 0 1 0 1 0 1 0 4 0 20 0 6 0 5 0 9 0 16 0 1 0 1 17 1 18 7 5 1 1 1 9 1 11 1 18 1 33 1 5 0 16 1 7 1 20 1 1 1 18 5 0 0 0 22 0 12 0 0 0 12 0 23 0 0 0 16 0 22 0 0 0 23 0 11 19 1 20 1 14 1 6 1 5 3 1 1 1 1 37 1 9 1 22 0 8 1 5 1 18 1 20 1 0 1 0 0 0 19 0 5 0 6 0 1 0 0 0 7 0 8 0 17 0 8 0 0 21 13 10 0 6 1 0 0 13 0 10 3 0 1 0 0 16 1 0 1 17 0 1 1 0 1 22 1 0 9 0 8 0 15 0 6 0 0 0 1 0 1 0 20 0 5 0 1 0 7 0 1 23 1 1 3 1 1 5 1 18 1 28 1 24 5 1 1 1 1 23 1 8 1 7 0 16 1 24 17 0 0 0 18 0 12 0 0 0 29 0 1 0 0 0 1 0 18 0 0 0 16 0 10 25 1 1 1 6 0 16 1 8 1 0 1 16 1 1 0 1 1 11 1 0 1 1 1 10 0 My question is: are there heuristic reasons for or against the conjecture?

论坛问题帖 | no_answers_at_retrieval | Lehs | 2017-09-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018151 | MO-280192

Is there always one integer between these two rational numbers?

It appears that for each integer $k\geq2$, there is always one integer c that satisfies the inequalities below. Can this be proved? $$\frac{3^k-2^k}{2^k-1}<c\leq \frac{3^k-1}{2^k}.$$ Note that for $k\geq2$ the lower bound is always a proper fraction and will never match an integer. Edit 28/1/2018 I have a short proof here on Overleaf. <--- It's done! Edit #28 is perfect! edit It looks like Waring's problem already has a solution. The sequence: https://oeis.org/A060692 "Corresponds to the only solution of the Diophantine equation 3^n = x*2^n + y*1^n with constraint 0 <= y < 2^n." This is equivalent to our $a+c$ for the Waring if statement. Oops, there is no proof that the diophantine equation is always $\leq 2^n.$ Here is a proof sketch of Waring's Here is a brute force function: aplusc[k_] := Module[{c}, c = 1; While[0 < 3^k - 2^k (++c)]; 3^k - 2^k (--c) + c] where we increment c until the calculation becomes negative, then we decrement by one to get c. We recalculate using that c and add them together. A060692(k) equals this value. Closed form: here. The brute force function illustrates the sawtooth pattern. We can also create a(k), b(k), and c(k) using the same module. And we can use b(k) for the proof. Trivially, we can show b$(k) + 1 \leq 2^k.$ edit. Some more information and references: https://oeis.org/A002379 PM

论坛问题帖 | has_answers | Fred Daniel Kline | 2017-09-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018152 | MO-281356

Does the image of the exponential map generate the group?

Let G be a connected Fréchet-Lie group and let $\mathfrak g$ be its Lie algebra. Does the image $\exp(\mathfrak g) \subset G$ of the exponential map generate G?

论坛问题帖 | has_answers | Jarek Kędra | 2017-09-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018153 | MO-283109

Converse of the Archimedean property of the sphere

In his remarkable book On the Sphere and Cylinder, where he came close to discovering calculus, Archimedes showed that the area of the portion of the sphere contained between a pair of parallel planes cutting the surface depends only on the distance between the planes (see p. 625 of this paper of King). This fact, which has been dubbed Archimedes hatbox theorem, is now a standard exercise in many calculus texts, and is even commemorated on the back of the Fields medal (if you look closely there, you can see the figure below in the background). Conversely, Blaschke showed that the only convex surface with this slab area property is the sphere. Indeed it is a simple exercise in classical differential geometry to check that any smooth convex surface with the slab area property must have constant curvature (Let $A(h)$ be the area trapped between a tangent plane and a parallel plane at distance h, and compute the limit of $A(h)/h$ as $h\to 0$). But can one still characterize the sphere if we fix the distance between the planes: Question: Let $S\subset R^3$ be a convex surface with diameter d. Suppose that for some constant $0<h<d$, the area of the

portion of S trapped between every pair of parallel planes separated by the distance h is constant, whenever both planes intersect S. Does it follow then that S is a sphere? A convex surface is the boundary of a compact convex set with interior points, and its diameter is the distance between its farthest points in the ambient space. Although this is known to be an open problem, I am not aware if it has been explicitly mentioned anywhere. Addendum: See this question for a closely related problem on plank invariant measures for convex planar's domains, and more references in this area.

论坛问题帖 | has_answers | Mohammad Ghomi | 2017-10-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018154 | MO-283370

Plank invariant measures on convex bodies

Let $K\subset R^2$ be a convex body, i.e., a compact convex set with interior points. A plank P is the region between a pair of parallel lines in R^2. Let us say that P intersects K properly provided that both boundary lines of P intersect K. Question: Let $K\subset R^2$ be a convex body with diameter d. Suppose there exists a measure μ on K such that $\mu(P\cap K)$ is constant for all planks of some fixed width $0<h<d$ which properly intersect K. Must K then be a disk? A disk $D\subset R^2$ does indeed admit such a measure. Let $S\subset R^3$ be a sphere with D as an equatorial disk, and μ be the measure on D induced by the projection of S (see the picture below which is from this paper of Kupavskii and Pach). Then μ will be plank invariant due to the Archimedes hatbox theorem. The motivation for the above question is that a positive answer here would yield a positive answer to this earlier question, on the Converse of the Archimedean property of the sphere, as follows. Suppose that there exits a convex surface $S\subset R^3$ with the Archimedean slab invariant area property mentioned in the earlier question. Then the projection of S into any plane would yield a convex body with the plank invariant property with respect to the induced measure. Hence the projection would be a disk, if the answer to the above question were yes, which would in turn imply that S is a sphere (it is well-known that the sphere is the only convex surface with round projections). For some results in the direction of the above problem see this paper of Gardner. Another nice paper discussing some relevant background and questions is the expository paper of King in the Monthly. More references can be found in the recent paper of Kupavskii and Pach mentioned above. Also see the answer to this related question asked earlier, and another similar question which has remained unanswered. Questions of this type have connections to Tarski's plank problem and an open question of Bang.

论坛问题帖 | no_answers_at_retrieval | Mohammad Ghomi | 2017-10-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018155 | MO-283476

Converse to Euclid's fifth postulate

There is a fascinating open problem in Riemannian Geometry which I would like to advertise here because I do not think that it is as well-known as it deserves to be. Euclid's famous fifth postulate, or more precisely Playfair's version of it, states that, in the Euclidean plane, through every point outside a line ℓ there passes one and only one line which does not intersect ℓ. Question: Is the Euclidean plane the only complete Riemannian manifold homeomorphic to R^2 which satisfies the fifth postulate, i.e., through any point outside a complete geodesic γ there passes one and only one complete geodesic which does not intersect γ? In other words, does the fifth postulate force the curvature to be zero? The reason this is interesting, or historically significant, is that it was the attempts to reduce the fifth postulate to the other axioms of Euclid, throughout the middle ages, which eventually led to the discovery of non Euclidean geometries and the notion of curvature by Gauss and Riemann. This problem appears to be originally due to Burns and Knieper in 1991: see the survey paper by Burns and Matveev, which also includes other nice problems on geometry of geodesics. The problem is also mentioned in papers of Croke, and Bangert and Emmerich, and has been studied most recently by Ge, Guijarro, and Solorzano.

论坛问题帖 | has_answers | Mohammad Ghomi | 2017-10-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018156 | MO-283568

Intrinsic vs Extrinsic geometry of convex surfaces

By Alexandrov's isometric embedding theorem, any locally convex metric prescribed on the sphere admits a realization as a convex surface in Euclidean 3-space, which, by Pogorelov's rigidity result, is unique up to an isometry of the ambient space. Thus, the intrinsic geometry of a convex surface completely determines its extrinsic geometry, and in principle it should be possible to describe all geometric quantities in terms of the intrinsic metric. On the other hand, Alexandrov's proof is not constructive, and does not provide any hints as to how one might be able to do this. Hence my first question is: Question 1: Is there any way to compute, recognize, or characterize intrinsically any of the geometric quantities or features of a convex surface which are not invariant under local isometric deformations, e.g., mean curvature, principal directions, principal curvatures, or umbilic points? I recognize this might be too much to ask, because the procedures which the above question asks for would somehow need to take into account the whole metric, not just a neighborhood of a point. So let me try to make things a bit more concrete and maybe a little more accessible. It is well known that any (smooth) convex surface must have at least one umbilic, i.e., a point where the principal curvatures are equal (if not then the direction of the larger principal curvature yields a nonvanishing line field on the surface which violates the Poincare-Hopf index theorem). Can one prove this without reference to the ambient space, or quantities that we only know how to compute extrinsically: Question 2: Is it possible to prove the existence of an umbilic point of a smooth convex surface in a purely intrinsic way? If so, can one also find or approximate the location of the umbilics? One motivation behind this questions is the famous conjecture of Caratheodory, which states that each convex surface must have at least two umbilics. The extrinsic or local approaches to this problem have always been problematic. Adendum: As Robert Bryant correctly points out below, the topological argument for the existence of at least one umbilic described above is really intrinsic and independent of the notion of principal directions. So the main point of Question 2 is its second part, i.e., to somehow get a handle on the location of the umbilic. See also this related question.

论坛问题帖 | has_answers | Mohammad Ghomi | 2017-10-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018157 | MO-283626

What is an umbilic point of a convex polyhedron?

An umbilic point of a smooth (C^2) convex surface in Euclidean 3-space is a point where the principal curvatures are equal. Is there some good way to generalize this notion to convex polyhedra? See this earlier question, specially the comments by Igor Rivin and Ivan Izmestiev for the motivation behind the current question. First, what do I mean by "good"? Here are some requirements that come to mind (which may be negotiable): It is not necessary for the notion of umbilic to be defined for all convex polyhedra but it would be enough if it is well-defined for generic polyhedra, or a class of polyhderda which is dense in the space of all convex polyhedra, and possibly open as well. Each convex polyhedron (or each polyhedron in the generic subclass) should have at least one umbilic point. It would be better if each polyhedron had at least two umbilics, or otherwise there was a notion of index associated to each umbilic so that the sum of the indices was constant for all polyhedra (in parallel with the index formula for the singularities of a line field on a sphere). It would be reassuring if there were some convex polyhedra with exactly two umbilics. (Bonus) Ultimately it would be nice if this notion were stable as a family of convex polyhedra converges to a smooth convex surface, but this can be set aside for now. Question: Does there exist a notion of umbilic point for convex polyhedra which would satisfy the first four requirements listed above? As was suggested by Ivan Izmestiev, maybe one should consider the Gauss image around vertices of the polyhedron. More precisely, consider the outward unit normals at the faces around a vertex. These normals form the vertices of a convex spherical polygon whose edges are geodesic arcs which join a pair of normals if and only if those normals correspond to adjacent faces of the polyhedron. So a vertex would be considered umbilic if its spherical image is somehow "circular" or relatively round (as is the case in the smooth case). Perhaps a useful guide here could be the generalized four vertex theorem for convex (spherical) polygons; see for instance Section 21 of Igor Pak's Lecture Notes and the corresponding papers of Serge Tabachnikov cited there. I first heard about the question in the title when Brian White asked me that in 2010. My idea back then was to consider the class of convex polyhedra with quadrilateral faces, and define an umbilic point as a vertex with degree different from 4. It is a simple exercise to show, via Euler's formula, that all convex polyhdera with quadrilateral faces must have at least 8 umbilics in this sense. The intuition behind this definition was the notion of "circular planar quadrilateral meshes", or "circular nets" which are discrete analogues of principal curvature lines. See for instance the book of Bobenko and Suris. This then was a purely topological candidate, and I did not get much chance to think about it more since then. Perhaps it might be more fruitful to consider geometric alternatives mentioned above, or some kind of a hybrid.

论坛问题帖 | no_answers_at_retrieval | Mohammad Ghomi | 2017-10-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018158 | MO-283758

Conjecture: The number of points modulo p of certain elliptic curve is p or $p+2$ for p of form $p=27a^2+27a+7$

Numerical evidence suggests a conjecture that the number of points of certain elliptic curve over $\mathbb{F}_p$ is either p or $p+2$ for p of certain form. Let p be prime of the form $p=27a^2+27a+7$ and (a/b) denote the Kronecker symbol. For integer k nonzero modulo p define $E_k / \mathbb{F}_p : y^2=x^3+2k^3$. E_k is the quadratic twist of $y^2=x^3+2$. Conjecture 1: $\#E(\mathbb{F}_p)=p+1 + (2k^3 / p)$. In other words for p of the given form $\#E(\mathbb{F}_p)$ is either p or $p+2$. Is Conjecture 1 true? Example sage session with 200 bit p sage: a=2^100+8;p=27*a^2+27*a+7;k=3;Kp=GF(p);E=EllipticCurve([Kp(0),Kp(2*k^3)]) sage: o=E.order();o2=p+1+kronecker(2*k^3,p);o==o2 True

论坛问题帖 | has_answers | joro | 2017-10-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018159 | MO-283921

Equilaterally triangulated surfaces with prescribed boundary

There is a problem in Richard Kenyon's list (Wayback Machine) which I would like to post here, because although I have thought about it from time to time, I have not been able to make the slightest progress on it: Question: "Given a closed polygonal path p in R^3 composed of unit segments, is there an immersed polygonal surface whose faces are equilateral triangles of edge length 1, spanning p?" The condition that the configuration is an "immersed" surface means that the whole configuration may be parametrized by a continuous locally one-to-one mapping from a compact connected 2-dimensional manifold with boundary into R^3. In other words at most two triangles may share an edge, and triangles sharing a vertex form an embedded topological disk, but two triangles which do not share an edge or a vertex may intersect. I cannot answer the question even for the case where p is a quadrilateral. Has there been any partial results for this problem?

论坛问题帖 | has_answers | Mohammad Ghomi | 2017-10-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018160 | MO-284612

Can one find a Jordan curve which has exactly one inscribed rectangle?

In On the number of inscribed squares of a simple closed curve in the plane it is shown that Theorem: For every positive integer n there is a simple closed curve in the plane (which can be taken infinitely differentiable and convex) which has exactly n inscribed squares. this is my question: Can one find a Jordan curve which has exactly one inscribed rectangle? More generally, what is the result if we use "rectangle" instead of "square" in the foregoing theorem?

论坛问题帖 | has_answers | C.F.G | 2017-10-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018161 | MO-284876

On statistical bases in Banach spaces

Let K be a subset of the positive integers $\mathbb{N}$. For each $n\in \mathbb{N}$, K_{n} denotes the set $\{k\in K: k\leq n\}$ and $|K_{n}|$ denotes the number of the elements in K_{n}. The natural density of K is defined by $$\delta(K)=\lim_{n\rightarrow \infty}\frac{|K_{n}|}{n}.$$ A sequence $(x_{k})_{k}$ in a Banach space X is said to be norm statistically convergent to $x\in X$ if $\delta(\{k\in \mathbb{N}:\|x_{k}-x\|\geq \epsilon\})=0$ for every $\epsilon>0$. It is natural to define a series $\sum_{k=1}^{\infty}x_{k}$ to be norm statistically convergent to $x\in X$ by $(s_{n})_{n}=(\sum_{k=1}^{n}x_{k})_{n}$ to be norm statistically convergent to x. We say that a sequence $(x_{k})_{k}$ in a Banach space X is a statistical basis for X if, for each $x\in X$, there exists a unique $(a_{k})_{k}$ such that the series $\sum_{k=1}^{\infty}a_{k}x_{k}$ norm statistically convergent to x. Clearly, statistical basis is a generalization of the notion of Schauder basis. Question: For each n, we can define a linear functional f_{n} in X by $f_{n}(x)=a_{n}$. If $(x_{k})_{k}$ is a Schauder basis, we know that f_{n} is continuous. However, if $(x_{k})_{k}$ is a statistical basis for X, is f_{n} continuous?

论坛问题帖 | has_answers | Dongyang Chen | 2017-10-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018162 | MO-284932

Portability of Thompson theorem about solvability to Moufang loops

Say we have a finite Moufang Loop Q, $|Q|<\infty$. There is a theorem proved by Thompson that states: Group G, $|G|<\infty$ is solvable $\iff$ $\forall a, b \in G$, $\langle a, b\rangle$ is solvable. My question is: can we translate the theorem to the finite Moufang loops (maybe with some extra constraints)? The statement is reasonable because it is well-known that every pair of Moufang loop's elements forms a group structure (Moufang's theorem). What is interesting, Feit-Thompson theorem about solvability of finite groups with odd order translates to the Moufang loop case. The proof can be found here. There is an idea to translate the task to the case of $U(R)$, where R is alternative algebra, $1 \in R$.

论坛问题帖 | has_answers | Evgeny | 2017-11-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018163 | MO-285396

Two questions on "Table problem on $\Bbb S^2$"

The following conjecture is known as "Table problem on $\Bbb S^2$" Conjecture (Table problem on $\Bbb S^2$): Suppose $x_1, x_2,x_3,x_4 \in\Bbb S^2 \subseteq \Bbb R^3$ are the vertices of a square that is inscribed in the standard 2-sphere, and let $h : \Bbb S^2\to \Bbb R$ be a smooth function. Then there exists a rotation $\rho\in \rm{SO}(3)$ such that $h(\rho(x_1)) =\cdots= h(\rho(x_4))$. Question 1: What is the meaning of "standard 2-sphere"? topological 2-sphere or geometrical? Question 2: Is this conjecture open still?

论坛问题帖 | has_answers | C.F.G | 2017-11-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018164 | MO-287011

A conjecture regarding prime numbers

For $n,m \geq 3$, define $ P_n = \{ p : p$ is a prime such that $ p\leq n$ and $ p \nmid n \}$. For example : $P_3= \{ 2 \}$ $P_4= \{ 3 \}$ $P_5= \{ 2, 3 \}$, $P_6= \{ 5 \}$ and so on. Claim: $P_n \neq P_m$ for $m\neq n$. While working on prime numbers I formulated this problem and it has eluded me for a while so I decided to post it here. I am not sure if this is an open problem or solved one. I couldn't find anything that looks like it. My attempts haven't come to fruition though I have been trying to prove it for a while. If m and n are different primes then it's clear. If $m \geq 2n$, I think we can find a prime in between so that case is also taken care of. My opinion is that it eventually boils down to proving this statement for integers that share the same prime factors. My coding is kind of rusty so would appreciate anybody checking if there is a counterexample to this claim. Any ideas if this might be true or false? Thanks. PS: I asked this question on mathstackexchage and somebody recommended I post it here as well. Here is the link to the original post https://math.stackexchange.com/questions/2536176/a-conjecture-regarding-prime-numbers

论坛问题帖 | has_answers | BRP | 2017-11-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018165 | MO-289148

Asymptotic form of pdf of Escape Time of arithmetic fBm

I am trying to apply the Girsanov formula and Doobs optional sampling theorem to obtain an asymptotic form of first passage density of an fbm process with drift, but the answer i am getting seems strange, it does not match the brownian motion case result as well as i assumed it would. I am seeking help in debugging these computations. I am also very much interested in learning about analytical ways of verifying this result. Thank You Some known results The first passage density of Brownian motion case is given by the following theorem. Theorem: Let the arithmetic Brownian motion process $X \left(t\right)$ be defined by the following Brownian motion driven SDE \begin{equation} \mbox{d}X \left(t\right) = a \mbox{d}t + b \mbox{d}{W}\left(t\right). \end{equation} with initial value X_0. Let $\tau =\inf \left(u |X(u) \le B\right)$ denote the first passage time for the barrier $X_0 < B$. Then the first passage time τ is distributed as Inverse Gaussian Distribution \begin{equation} \tau \sim IG\left(\frac{B - X_0}{a}, \frac{\left(B - X_0\right)^2}{b^2}\right),\label{abmFirstPassageDist} \end{equation} and for $t > 0$ the pdf of τ is \begin{equation} f(t) = \sqrt{\frac{(B - X_0)^2}{2 \pi b^2 t^3}} \exp\left[-\frac{ \left(at - B + X_0\right)^2}{2 b^2 t}\right]\label{abmFirstPassageDensity}. \end{equation} The Girsanov formula for fBm Let $B^H \left(t\right)$ denote fractional brownian motion with mean 0 and variance t^{2H}, for all $H \in \left(0, 1\right)$ defined on $\left(\Omega, \mathcal{F}, \mathbb{P}\right)$. Let a be a scalar. Define a new probability measure $\mathbb{Q}$ on $\left(\Omega, \mathcal{F}\right)$ via the Radon Nikodym derivative with respect to $\mathbb{P}$ \begin{equation} \frac{\mbox{d}{\mathbb{Q}}}{\mbox{d}{\mathbb{P}}} = \exp\left\{a M_T - \frac{1}{2}{a}^2 \langle M, M\rangle_T \right\}\label{girsanovRadonNikodymDerivativeWithInnerProduct} \end{equation} where \begin{equation} M_T = \frac{1}{2 H \Gamma \left(\frac{3}{2} - H\right) \Gamma \left(H +

\frac{1}{2}\right)} \int_0^T \left(s\left(T - s\right)\right)^{\frac{1}{2} - H} \mbox{d}{{B_s}^H}. \end{equation} The process M_T is a martingale with independent increments, zero mean and variance function $c^2 T^{2 - 2H}$ where \begin{equation}\label{c} c = \sqrt{\frac{\Gamma\left(\frac{3}{2} - H\right)}{2 H \left(2 - 2H\right)\Gamma\left(H + \frac{1}{2}\right)\Gamma\left(2 - 2H\right)}}. \end{equation} Then the process defined, for all $t \in \left[0, T\right]$, by $B^H \left(t\right) + a t$ is the standard $\mathbb{Q}$-fractional Brownian motion on $\left[0, T\right]$. In other words, under probability measure $\mathbb{Q}$, $B^H \left(t\right)$ restricted to $t \in \left[0, T\right]$ is distributed as an arithmetic fractional Brownian motion with drift a. A proof of the Girsanov formula for fractional Brownian motion can be found in Norros's paper, where the term the fundamental martingale is also coined for the process M_t. It's noteworthy that using the variance of M_t, Radon Nikodym Derivative can also be re-written as \begin{equation} \frac{\mbox{d}{\mathbb{Q}}}{\mbox{d}{\mathbb{P}}} = \exp\left\{a M_T - \frac{1}{2}{a}^2 c^2 T^{2 - 2H} \right\}.\label{girsanovRadonNikodymDerivative} \end{equation} corollary Let $X\left(t\right) = b B^H \left(t\right)$ be an arithmetic fractional brownian motion with volatility b, for all $H \in \left(0, 1\right)$ defined on $\left(\Omega, \mathcal{F}, \mathbb{P}\right)$. Let a be a scalar. Define a new probability measure $\mathbb{Q}$ on $\left(\Omega, \mathcal{F}\right)$ via the Radon Nikodym derivative with respect to $\mathbb{P}$ \begin{equation} \frac{\mbox{d}{\mathbb{Q}}}{\mbox{d}{\mathbb{P}}} = \exp\left\{\frac{a}{b} M_T - \frac{1}{2}\frac{a^2}{b^2} c^2 T^{2 - 2H}\right\}.\label{scaledGirsanovRadonNikodymDerivative} \end{equation} Then the process defined, for all $t \in \left[0, T\right]$, by $Z \left(t\right) = X \left(t\right) + a t$ is an arithmetic $\mathbb{Q}$-fractional Brownian motion process on $\left[0, T\right]$ with volatility b. In other words, under probability measure $\mathbb{Q}$, $X\left(t\right)$ restricted to $t \in \left[0, T\right]$ is distributed as an arithmetic fractional Brownian motion with drift a and volatility b. Proposition Let $B^H \left(t\right)$ denote scaled fractional brownian motion with mean 0 and variance $b^2 t^{2H}$, for all $H \in \left(0, 1\right)$ with respect to measure $\mathbb{P}$. Define $\tau_k = \inf \left\{t \ge 0 : B^H \left(t\right) = k\right\}$ for $k > 0$. Then the conditional mean and variance of M_t given B_t are \begin{equation} E\left(M_t | B^H \left(t\right) = k \right) = {t^{1-2H} k \over b} \end{equation} and \begin{equation} \hbox{Var}\left(M_t | B^H \left(t\right) = k\right) = t^{2-2H}\left(c^2-1\right). \end{equation} Proof Both $B^H \left(t\right)$ and M_t have mean zero, the variance of $B^H \left(t\right)$ is $b^2 t^{2H}$, the variance of M_t is $c^2 T^{2 - 2H}$, and their covariance bt, can be derived similarly to, as in Proposition 3.2, in Norros's paper. Hence the correlation coefficient ρ between M_t, $B^H \left(t\right)$ is $1/c$. Therefore, using elementary results for the bivariate normal distribution, we find \begin{eqnarray} E\left(M_t | B^H \left(t\right) = k \right) &=& {\rho \sigma_{M_t}\over \sigma_{B^H \left(t\right)}}k\\\nonumber &=& {t^{1-2H} k \over b}\nonumber \end{eqnarray} and \begin{eqnarray} \hbox{Var}\left(M_t | B^H \left(t\right) = k\right) &=& {\sigma^2}_{M_t}\left(1 - \rho^2\right) \\\nonumber &=& t^{2-2H}\left(c^2-1\right).\nonumber \end{eqnarray} The Derivation Attempt By Doob's optional sampling theorem (justified by the uniform integrability of the martingale \begin{equation} \exp\left\{\frac{a}{b} M_T - \frac{1}{2}\frac{a^2}{b^2} \langle M, M\rangle_T\right\}\nonumber \end{equation} on $\left[0, T\right]$ and the fact that $\left\{\tau_k \le T \right\} \in \mathcal{F}_{\tau_k} \cap \mathcal{F}_{T} = \mathcal{F}_{\tau_k \bigwedge T} \subseteq \mathcal{F}_T$), \begin{equation} \mathrm{E}\left[\left.\exp\left\{\frac{a}{b} M_T - \frac{1}{2}\frac{a^2}{b^2} \langle M, M\rangle_T\right\} \right| \mathcal{F}_{\tau_k \bigwedge T}\right] = \exp\left\{\frac{a}{b} M_{\tau_k \bigwedge T} - \frac{1}{2}\frac{a^2}{b^2} \langle M, M\rangle_{\tau_k \bigwedge T}\right\}. \end{equation} Therefore, \begin{eqnarray} \mathbb{P}^{a,T}\left[\tau_k \in (t,t+dt)\right] &=& \mathbb{E}^{a,T} \left[1_{\left\{\tau_k \in (t,t+dt)\right\}} \right]\\ \nonumber &=& \mathbb{E}\left[\exp\left\{\frac{a}{b} M_T - \frac{1}{2}\frac{a^2}{b^2} \langle M, M\rangle_T\right\}1_{\left\{\tau_k \in (t,t+dt)\right\}}\right] \end{eqnarray} by Corollary above \begin{eqnarray} \nonumber &=& \mathbb{E}\left[\left.\mathbb{E}\left[\exp\left\{\frac{a}{b} M_{\tau_k \bigwedge T} - \frac{1}{2}\frac{a^2}{b^2} \langle M, M\rangle_{\tau_k \bigwedge T}\right\} \right|{\cal F}^{B^H}_{\tau_k\wedge T}\right]1_{\left\{\tau_k \le T\right\}}\right]\\ \nonumber &=& \mathbb{E}\left[\exp\left\{\frac{a}{b} M_{\tau_k} - \frac{1}{2}\frac{a^2}{b^2} \langle M, M\rangle_{\tau_k}\right\}1_{\left\{\tau_k \le T\right\}}\right]\\ \end{eqnarray} by Doob's sampling theorem \begin{eqnarray} \nonumber &=& \mathbb{E}\left[\exp\left\{\frac{a k}{b^2}{\tau_k}^{1-2H} - \frac{1}{2}\frac{a^2}{b^2} {\tau_k}^{2-2H}\left(c^2 - 1\right)\right\}1_{\left\{\tau_k \le T\right\}}\right]\\ \end{eqnarray} by propostion above \begin{eqnarray} \nonumber &=& \int_0^T \exp\left[\frac{a k t^{1-2H}}{b^2} - \frac{1}{2}\frac{a^2 t^{2 - 2H}}{b^2} \left(c^2 - 1\right)\right] \mathrm{P}\left[\tau_k \in \mbox{d}{t}\right]. \end{eqnarray} On the other hand, $\left\{B^H \left(t\right) + a t\right\}_{t \in \left[0, T\right]}$ is a scaled fractional Brownian motion under $\mathbb{P}^{a,T}$, so \begin{equation} \mathbb{P}^{a,T} \left[\tau_k \le T\right] = \mathbb{P}\left[\hat{\tau_k} \le T\right], \end{equation} where $\hat{\tau_k}$ is the first hitting time of the level k of the scaled fractional Brownian motion with drift a. Using the formula for asymptote of the first passage density of fBM without drift due to Molchan, it follows immediately that the long-time form of the first passage time density for fBm with drift is given by \begin{eqnarray} f\left(t\right) &=& \exp\left[\frac{a k t^{1-2H}}{b^2} - \frac{1}{2}\frac{a^2 t^{2 - 2H}}{b^2} \left(c^2 - 1\right)\right] t^{H-2}.\label{firstPassageDensityArithmeticFbm} \end{eqnarray} It is noteworthy that C^2 has a minimum at $H =\frac{1}{2}$, where $c^2 = 1$. Verification Against Brownian motion case Upon substituting $H = \frac{1}{2}$ in c^2 evaluates to 1, and the function reduces to \begin{eqnarray} f_{H = \frac{1}{2}}\left(t\right) &=& \exp\left[\frac{a k}{b^2}\right] t^{-\frac{3}{2}}. \end{eqnarray} This seems a bit off as according to "Brownian motion and stochastic calculus" books, it should be \begin{eqnarray} f'_{H = \frac{1}{2}}\left(t\right) &=& \exp\left[\frac{a k}{b^2} - \frac{a^2}{2b^2}t \}\right] t^{-\frac{3}{2}}. \end{eqnarray} The presence of $c^2 - 1$ seems to prevents a perfect match with the Brownian motion case.

论坛问题帖 | has_answers | Comic Book Guy | 2017-12-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018166 | MO-289293

The difference between two coprime semiprimes

Conjecture: Any positive integer can be written as the difference between two coprime semiprimes. Tested up to 1,000,000. See also: https://math.stackexchange.com/questions/2579578/the-difference-of-two-coprime-composites I'm not surprised that there are heuristics backing up the conjecture, but think it would be strange if the proof was so far away when the question above had an easy answer.

论坛问题帖 | has_answers | Lehs | 2017-12-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018167 | MO-289554

If $a^3+b^3+c^3=N$, then $x^3+y^3+z^3+t^3 = N$ in infinitely many ways?

It is well-known that, $$a^3+b^3+c^3 = N\tag1$$ for $N=1,\,2$ is solvable in the integers in infinitely many ways . However, it is an open question (but is conjectured) that if for general N it has a solution, then it must have infinitely many. I'd like to propose a variant, Conjecture: If $a^3+b^3+c^3=N$, then $x^3+y^3+z^3+t^3 = N$ in infinitely many ways. Part I: Use the identity, $$x^3+y^3+z^3+t^3 = (a^3+b^3+c^3)\big(p^2+(a+b)cq^2\big)^3$$ where, $$\begin{aligned} t &= a p^2 - c^2 p q + b (a + b) c q^2 \\ x &= b p^2 + c^2 p q + a (a + b) c q^2\\ y &= c p^2 + (a^2 - b^2) p q\\ z &= -(a^2 - b^2) p q + (a + b) c^2 q^2\end{aligned}$$ Part II: Solve the Pell equation, $$p^2+(a+b)cq^2 = 1\tag2$$ This is the tricky part since we require $(a+b)c<0$. Perusing through S. Huisman 15254 solutions, I find that all his solvable N have solutions where both a,b are negative, or c is negative. Of course, (2) can then be easily solved as a Pell equation. Q: If (1) has a solution in positive integers, then how do we prove that implies a subsequent solution where one or two of the variables are negative? P.S. The second solution may be huge, $$4^3+ 4^3+ 7^3 =-17863863023^3+ 13242033973^3+ 15005424841^3 = 471$$ but it guarantees infinitely many solutions to (2).

论坛问题帖 | no_answers_at_retrieval | Tito Piezas III | 2017-12-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018168 | MO-2897

A group action of the Heisenberg group with special symmetries

Suppose we look at the Heisenberg group H_{d} as a matrix group of upper triangular matrices over the ring $\mathbb{Z}/d\mathbb{Z}$. You can even choose d to be prime if you want. A natural irrep of H_{d} acting on $\mathbb{C}^{d}$ maps the group elements into the "shift" and "phase" operators, plus roots of unity. More specifically, the two natural generators map the orthonormal basis vectors from $j \to j+1\mod d$, and the Fourier transform of that operation, plus overall phases by roots of unity. The question is this: Can you find a unit vector v such that $|(v,U_g v)| = c$ for all g not in the center of $H_{d}\ ?$ One can solve for the constant: $c=\frac{1}{\sqrt{d+1}}$. Numerics suggests that these vectors exist in all the dimensions < 67, hence they may exist in every dimension, but the form of the vectors contains no (obvious) hint as to how to prove this. This problem seems extremely truculent and any help is greatly appreciated!

论坛问题帖 | has_answers | Steve Flammia | 2009-10-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018169 | MO-29000

Volumes of sets of constant width in high dimensions

Background The n-dimensional Euclidean ball of radius $1/2$ has width 1 in every direction. Namely, when you consider a pair of parallel tangent hyperplanes in any direction the distance between them is 1. There are other sets of constant width 1. A famous one is the Reuleaux triangle in the plane. The isoperimetric inequality implies that among all sets of constant width 1 the ball has largest volume. Let's denote the volume of the n-ball of radius $1/2$ by V_n. The question Is there some $\varepsilon >0$ so that for every $n>1$ there exist a set K_n of constant width 1 in dimension n whose volume satisfies $\mathrm{vol}(K_n) \le (1-\varepsilon)^n V_n$. This question was asked by Oded Schramm who also asked it for spherical sets of constant width r. A proposed construction Here is a proposed construction (also by Schramm). It will be interesting to examine what is the asymptotic behavior of the volume. (And also what is the volume in small dimensions 3,4,...) Start with $K_1=[-1/2,1/2]$. Given K_n consider it embedded in the hyperplane of all points in R^{n+1} whose $(n+1)$-th coordinate is zero. Let K^+_{n+1} be the set of all points x, with nonnegative $(n+1)$-th coordinate, such that the ball of radius 1 with center at x contains K_n. Let K^-_{n+1} be the set of all points x, with nonpositive $(n+1)$-th coordinate, such that x belongs to the intersection of all balls of radius 1 containing K_n. Let K_{n+1} be the union of these two sets K^-_{n+1} and K^+_{n+1}. Motivation Sets of constant width (other than the ball) are not lucky enough to serve as norms of Banach spaces and to attract the powerful Banach space theorist to study their asymptotic properties for large dimensions. But they are very exciting and this looks like a very basic question. References and additional motivation In the paper: "On the volume of sets having constant width" Isr. J Math 63(1988) 178-182, Oded Schramm gives a lower bound on volumes of sets of constant width. Schramm wrote that a good way to present the volume of a set $K \subset R^n$ is to specify the radius of the ball having the same volume as K, called it the effective radius of the set K and denote it by $\operatorname{er} K$. Next he defined r_n as the minimal effective radius of all sets having constant width two in R^n. Schramm proved that $r_n \ge \sqrt {3+2/(n+1)}-1$. He asked if the limit of r_n exists and if r_n is a monotone decreasing sequence. Our question is essentially whether r_n tends to 1 as n tends to infinity. In the paper: O. Schramm, Illuminating sets of constant width. Mathematika 35 (1988), 180--189, Schramm proved a similar lower bound for the spherical case and deduced the best known upper bound for Borsuk's problem on covering sets with sets of smaller diameter.

论坛问题帖 | has_answers | Gil Kalai | 2010-06-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018170 | MO-291018

Mathematical conjectures on which applications depend

What are some examples of mathematical conjectures that applied mathematicians assume to be true in applications, despite it being unknown whether or not they are true?

论坛问题帖 | has_answers | Christopher King | 2018-01-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018171 | MO-291754

What is the smallest sphere whose surface includes 100 integer points?

Let $S(r)$ be the surface of the origin-centered sphere in $\mathbb{R}^3$. A point is an integer point if all its coordinates are integers. What is the smallest radius r_n such that $S(r_n)$ includes $\ge n$ integer points? What is the growth rate of r_n with respect to n? Is there an algorithm that could compute r_n for a specific given n? It is known that rational points (all coordinates rational) are dense on $S(1)$: see, e.g., the MO question Rational points on a sphere in $\mathbb{R}^d$. One possible approach is via rational points of bounded height. The height of a rational a/b in lowest terms is max$(|a|,|b|)$, and the height of a rational point is the max of the heights of its coordinates. Rationals of height ≤ 2048 on sphere. Image due to Stefan Kohl in this answer. Choose an appropriate $h_\max$, perhaps using an estimate of the number of rational points of height at most $h_\max$ on $S(1)$. Then scale all coordinates by the LCM of the points' denominators. For example, for $h_\max=10$, scaling by $2^3 \cdot 3^2 \cdot 5 \cdot 7 = 2520$ would suffice to clear all denominators, so that $S(2520)$ includes all those points at integer coordinates. But this would not necessarily result in the minimum r_n for a given n. It would likely be better to use rational points that result in a small LCM. Exact calculations on a sphere (for example, computing Voronoi diagrams on a sphere) often need integer points of bounded size. Update. The exact answer to the question in the title, due to Dap and Gerhard Paseman, is that a sphere of radius $\sqrt{74} \approx 8.6$ includes 120 integer points on its surface, and smaller spheres include fewer than 100 points.

论坛问题帖 | has_answers | Joseph O'Rourke | 2018-01-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018172 | MO-291868

Open problems in the theory of manifolds relating to construction

A while ago I stumbled across a paper of Thurston: Some Simple Examples of Symplectic Manifolds, where Thurston constructs closed symplectic manifolds with no Kaehler structure. My question is: What are the most rewarding and beautiful (potential) manifolds that one might seek to construct, that have not been constructed thus far? (Does there exist [insert]?). Obviously potential constructions are not restricted to manifolds , however I am mostly interested in constructions of manifolds or spaces in the most current and/or new areas of research in geometry and topology. I would also be interested in conjectures (interconnected or otherwise) relating to manifolds that can be resolved via constructions, what constructions?

论坛问题帖 | no_answers_at_retrieval | nicolas bourbaki | 2018-01-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018173 | MO-293005

What is known about the relationship between Fermat's last theorem and Peano Arithmetic?

As far as I know, whether Fermat's Last Theorem is provable in Peano Arithmetic is an open problem. What is known about this problem? In particular, what is known about the arithmetic systems $PA + \text{Fermat's Last Theorem}$ and $PA + \lnot \text{Fermat's Last Theorem}$? Note in particular that if $PA \vdash \text{Fermat's Last Theorem}$, the first system will just be PA and the second system with be inconsistent. If FLT is independent of PA, then both will be consistient systems distinct from PA. $PA \vdash \lnot \text{Fermat's Last Theorem}$ is not possible, since PA and Fermat's last theorem are both true in $\mathbb N$.

论坛问题帖 | has_answers | Christopher King | 2018-02-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018174 | MO-293247

Open problems concerning Araujo's biseparating maps

Araujo stated the following four open questions at the end of his paper, page 518 and $519.$ Question $1:$ Assume that there exists a biseparating map $T:A^n(\Omega:E)\to A^m(\Omega',F)$ which is not linear. Can we deduce that the support map $h:\Omega'\to\Omega$ is a diffeomorphism of class $n?$ Problem $2:$ Suppose that $C^\infty(\Omega,E)$ is the space of E-valued functions which are of class C^∞ in $\Omega,$ and that $C^\infty(\Omega',F)$ is defined in a similar way. Describe the linear biseparating maps from $C^\infty(\Omega,E)$ onto $C^\infty(\Omega',F).$ Must such map be continuous? Problem $3:$ Let Ω and Ω' be unbounded open subsets of $\mathbb{R}^p$ and $\mathbb{R}^q$ respectively. Describe the linear biseparating maps from $C^n_*(\Omega,E)$ onto $C^m_*(\Omega',F).$ Problem $4:$ Determine all subspaces $A(\Omega,E)\subseteq A^n(\Omega,E)$ and $B(\Omega',F)\subseteq A^m(\Omega',F)$ such that the existence of a linear biseparating map from $A(\Omega,E)$ onto $B(\Omega',F)$ implies that E and F are isomorphic as Banach spaces. What is the status of these questions? Any progress on each of them? If yes, any reference is appreciated. Definitions: Let $\Omega\subseteq \mathbb{R}^p$ be an open subset and E be a Banach space. Denote $C^n(\Omega,E)$ the space of E-valued functions f on Ω that are of class $C^n.$ $T:C^n(\Omega,E)\to C^m(\Omega',F)$ is called biseparating map if T is bijective and $$\|Tf\| \|Tg\|=0 \text{ if and only if }\|f\|\|g\|=0$$

论坛问题帖 | no_answers_at_retrieval | Idonknow | 2018-02-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018175 | MO-296328

Discriminant of numerator of inverse logarithmic derivative operator iteration

Let $T:\mathbb Q(x)\to \mathbb Q(x)$ be the operator of inverse logarithmic derivative, i.e. $$Tf=\frac{f}{f'}.$$ Define $$p_n(x)=T^n\left(x-\frac{x^2}{2}\right).$$ Let $f_n(x) \in \mathbb Z[x]$ be the numerator of $p_n(x)$, taken with positive leading term. The sequence D_n of discriminants of our polynomials starts with $$\begin{aligned} D_0&=4,\\ D_1&=4,\\ D_2&=4,\\ D_3&=-256=2^8,\\ D_4&=2^{32}5^2,\\ D_5&=2^{128}5^6331^2, \\D_6&=2^{512}5^{20}331^646599695357^2.\\ \end{aligned} $$ Is it true that we always have $D_n=\pm a_n^2$ for some integer a_n? What else can be said about the arithmetic properties of a_n? For example, is it true that for all $n \geq 2$ we have $\nu_2(D_n)=2^{2n-3}$?

论坛问题帖 | has_answers | Alexander Kalmynin | 2018-03-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018176 | MO-296661

A "conjectured" concentration inequality for operators, probably related with random matrix theory

I am working on some open problem. And I have reduced the original problem to the "conjecture" (actually I am not familiar with random matrix theory or other fields that may have such a result) as follows. Consider a random variable X supported on $\mathbb{R}^{d}$ with distribution P_X. Assume we have constructed a cover of $\mathbb{R}^{d}$ denoted as $\{\Lambda(\omega):\omega\in\Omega\}$, where Ω is a finite set. We then make Ω into some probability space and introduce an extra r.v. σ that takes value in Ω. Now since we have (or not, why?) $$ \mathbb{E}_{\sigma}\int_{\Lambda(\sigma)}XX^{T}dP_{X} = \mathbb{E}_{X}[XX^{T}] $$, there should exist some constant $\lambda >0$, such that the following operator inequality is true with probability (in the order of positive semi-definiteness) $$ P_{X}(\Lambda(\sigma))\mathbb{E}_{X}[XX^{T}] - \lambda{I} \preceq \int_{\Lambda(\sigma)}XX^{T}dP_{X}\preceq P_{X}(\Lambda(\sigma))\mathbb{E}_{X}[XX^{T}] + \lambda{I} $$ From my perspective, the existence may be easy as we could somehow add some additional assumptions on P_X to assert the "boundedness" of XX^{T}. Thus we could take λ larger than the upper bound. Could anyone provide some reference or hints relevant to this problem? That will help a lot.

论坛问题帖 | no_answers_at_retrieval | Morino_Hikari | 2018-03-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018177 | MO-297054

The Krzyż Conjecture

What is the state of the Krzyż Conjecture? It states for that for all $f:\mathbb{D}\to \bar{\mathbb{D}}$ holomorphic and non-vanishing, the coefficients a_n in the power series of f are at most $2/e$ in absolute value seen from the maximal function $$f(z)=e^{\frac{z^n+1}{z^n-1}}$$ I have read The Krzyż Conjecture Revisited where the authors (in 2014) say the problem is open for coefficients $n>5$. A paper by S. Krushkal claims to have proved it, but in the former paper Krushkal's work is mentioned: Krushkal's unpublished preprint, in spite of some gaps, contains a wealth of geometric and analytic ideas which may be useful for a further study of problems of this type I could not spot any obvious holes but it has yet to be published in a reputable journal as far as I know, so I wonder if someone with a more penetrating and experienced eye could verify or point the "gaps".

论坛问题帖 | no_answers_at_retrieval | mysatellite | 2018-04-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018178 | MO-298012

An upper bound for a vector with given norm 1 and norm 2

Suppose $X = (x_1, \ldots , x_n)$ is given and we know that x_i's are nonnegative, $\sum_{i=1}^n x_i = n$ and $\sum_{i=1}^n x_i^2 = m $. Just by this information, is it possible to find a vector that majorizes X? the meaning of majorization can be found here: https://en.wikipedia.org/wiki/Majorization. Certainly, $(n , 0 , \ldots , 0)$ is an answer, but I want a nontrivial vector. In particular,I'm looking for some nontrivial known results . Thanks.

论坛问题帖 | has_answers | user115608 | 2018-04-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018179 | MO-298194

Lattices that top is the top of its join-irreducibles, such that a random element is almost surely greater then any given join-irreducible

Let $(L,\leq,0,1)$ be a lattice, and let's denote by $JI(L)$ the set of its join-irreducibles (i.e. elements that are not the lowest grater bound of two other elements). We suppose that $\sup JI(L)=1$. and that $\mathbb P$ is a probability on L such that ideals and filters are measurable and singleton have probability 0. Let's call such a lattice a "reach-irreducible lattice" (RIB). If the cardinality of L is that of the continuum" we can say "continuum reach-irreducible lattice" (CRIL) Does there exist in any L that is a continuum-reach-irreducible lattice, $g\in JI(L)$ such that $\mathbb P(\left\{x\in L,\, g\leq x\right\})<1$. Related discussions on the MSE are https://math.stackexchange.com/questions/2737618/lattice-generated-by-join-irreducible-elements-such-that-any-principal-filter-c and https://math.stackexchange.com/questions/2742357/probability-martin-axiom-and-the-weak-frankl-conjecture The first link is an imperfect attempt to fit with the motivation of this question, that is explained in the second link. If the answer to this question is yes, then a weak version of the Frankl Conjecture ("WFC")- that is also an open problem - is true (see MSE links). Note that in both links, the lattices that I built assuming that WFC is wrong are not exactly CRIL, but up to small details, one can easily get CRIL from them. Note also that the cardinality condition is just given to fit with WFC, but it would be nice indeed if the answer of the question is yes for any RIL. And if it is not, it might also give some useful informations, so I would also be happy to read answers and comments about general RIL.

论坛问题帖 | has_answers | jcdornano | 2018-04-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018180 | MO-298600

PhD dissertations that solve an established open problem

I search for a big list of open problems which have been solved in a PhD thesis by the Author of the thesis (or with collaboration of her/his supervisor). In my question I search for every possible open problem but I prefer (but not limited) to receive answers about those open problems which had been unsolved for at least (about) 25 years and before the appearance of the ultimate solution, there had been significant attentions and efforts for solving it. I mean that the problem was not a forgotten problem. If the Gauss proof of the fundamental theorem of algebra did not had a gap, then his proof could be an important example of such dissertations. I ask the moderators to consider this question as a wiki question.

论坛问题帖 | has_answers | Ali Taghavi | 2018-04-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018181 | MO-298635

Status of two Banach space theory open problems posted by Pełczyński

In the book 'Open Problems in the Geometry and Analysis of Banach Spaces', I am interested in the following two problems. Problem 1: Let X be a separable infinite-dimensional Banach space that is not isomorphic to a Hilbert space. (i) (A. Pełczyński) Does there exist an infinite dimensional subspace of X with Schauder basis that is not complemented in $X?$ (ii) (A. Pełczyński) Do there exist two infinite-dimensional subspaces of X with Schauder basis that are not isomorphic? I would like to know status of the two problems above. Are they solved or still open?

论坛问题帖 | no_answers_at_retrieval | Idonknow | 2018-04-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018182 | MO-298636

Status of the three-body problem

I find many numerical results on the three-body problem, but what is rigorously proved? Especially I would be interested in the parameter domains for which we have rigorous lower bounds on the topological entropy or Lyapunov exponent of the system.

论坛问题帖 | has_answers | Jörg Neunhäuserer | 2018-04-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018183 | MO-299056

Strengthened version of Isoperimetric inequality with n-polygon

Let $ABCD$ be a convex quadrilateral with the lengths a, b, c, d and the area S. The main result in our paper equivalent to: \begin{equation} a^2+b^2+c^2+d^2 \ge 4S + \frac{\sqrt{3}-1}{\sqrt{3}}\sum{(a-b)^2}\end{equation} where $\sum{(a-b)^2}=(a-b)^2+(a-c)^2+(a-d)^2+(b-c)^2+(b-d)^2+(c-d)^2$ and note that $\frac{\sqrt{3}-1}{\sqrt{3}}$ is the best constant for this inequality. Similarly with the same form above apply to n-polygon, I propose a conjecture that: Let a convex polygon $A_1A_2...A_n$ with the lengths are $a_1, a_2, ...a_n$ and area S we have: \begin{equation} \sum_i^n{a_i^2} \ge 4\tan{\frac{\pi}{n}}S + k\sum_{i < j}{(a_i-a_j)^2}\end{equation} I guess that $k=tan{\frac{\pi}{n}}-tan{\frac{\pi}{n+2}}$ I am looking for a proof of the inequality above. Note that using Isoperimetric inequality we can prove that: \begin{equation} \sum_i^n{a_i^2} \ge 4\tan{\frac{\pi}{n}}S\end{equation} Case n=3: Hadwiger-Finsler inequality, $k=1 \approx 1.00550827956352=tan{\frac{\pi}{3}}-tan{\frac{\pi}{5}}$: \begin{equation}a^{2} + b^{2} + c^{2} \geq 4 \sqrt{3}S+ (a - b)^{2} + (b - c)^{2} + (c - a)^{2}\end{equation} Case n=4: our paper, $k=\frac{\sqrt{3}-1}{\sqrt{3}}$, $k=tan{\frac{\pi}{4}}-tan{\frac{\pi}{6}}$ \begin{equation} a^2+b^2+c^2+d^2 \ge 4S + \frac{\sqrt{3}-1}{\sqrt{3}}\sum{(a-b)^2}\end{equation} See also: Weitzenböck's inequality Hadwiger-Finsler inequality Isoperimetric inequality.

论坛问题帖 | has_answers | Đào Thanh Oai | 2018-04-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018184 | MO-299323

Relative to Isoperimetric inequality with n-polygon

Since Brahmagupta's formula and Bretschneider's formula we have the inequality: Any two quardrilaterals $A_1A_2A_3A_4$ and $B_1B_2B_3B_4$ with the same sidelengths and $A_1A_2A_3A_4$ is a cyclic but $B_1B_2B_3B_4$ is not a cyclic then area of $A_1A_2A_3A_4$ $\ge$ area of $B_1B_2B_3B_4$ Isoperimetric inequality state that: For the length L of a closed curve and the area A of the planar region that it encloses, that $L^2 \ge 4\pi .Area$ and that equality holds if and only if the curve is a circle. I conjecture that: Let $A_1A_2...A_n$ and $B_1B_2...B_n$ be two n-polygons with the lengths $a_1, a_2,...,a_n$ and $b_1, b_2,...,b_n$ such that with $i \in n$ then exist $j \in n$ such that $a_i=b_j$. If $A_1A_2...A_n$ is a cyclic and $B_1B_2...B_n$ is not a cyclic then area of $A_1A_2...A_n$ $\ge$ area of $B_1B_2...B_n$. I am looking for the proof of the inequality above.

论坛问题帖 | has_answers | Đào Thanh Oai | 2018-05-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018185 | MO-300505

Status of an open problem in isometric aspect of Banach space theory

The following open problem is taken from the book Open Problems in the Geometry and Analysis of Banach Spaces, page $40.$ Problem $84:$ Assume that X is an infinite-dimensional separable Banach space such that for any pair of points x and y in the unit sphere of X there exists a linear isometry from X onto X such that $Tx=y.$ Is X linearly isometric to a Hilbert space? After stating the problem, authors mentioned that the problem holds true for finite-dimensional spaces but not for non separable spaces. Then they point to the reference,page 255 which contains the problem above. My question: Does anyone know whether the problem is still open?If yes, what is a good reference? The authors of the book mentioned that problem 84 is an old classical open problem. So I suspect that there should be some papers available on solving the problem partially or fully.

论坛问题帖 | no_answers_at_retrieval | Idonknow | 2018-05-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018186 | MO-301110

Density of the Klarner-Rado Sequence

Consider the Klarner-Rado sequence OEIS A005658 defined by the rule: the sequence starts with 1, and if it contains n it also contains $2n$, $3n+2$ and $6n+3$. According to R. Guy's popular article, Klarner conjectures that this set has positive density. Lagarias in The Ultimate Challenge book says the conjecture is open. Question: should we believe or disbelieve the conjecture? Is there a heuristic argument explaining what's going on here?

论坛问题帖 | has_answers | Igor Pak | 2018-05-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018187 | MO-301135

Upper bound of the expectation of sum of the absolute value pairs

We have two arrays A,B of length n. All values are i.i.d drawn from same distribution on $[0,1]$. If we sort A,B in non-decreasing order and let $A_{(i)},B_{(i)}$ denote the i-th value in the sorted array. Then we have $S=\sum_{i=1}^n|A_{(i)}-B_{(i)}|$. What is the upper bound of $\mathbb{E}[S]$? I believe it must be the uniform distribution in order to get the upper bound, but I can't prove that. I've searched order statistics and I think that may give me some upper bound, but the expression is too complex and I can't complete the derivation.

论坛问题帖 | has_answers | user124861 | 2018-05-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018188 | MO-301691

Algebraic version of unilateral shift

It was confirmed that Wold-type decomposition can be extended from von Neumann algebras to Baer*-rings (see this paper). By algebraic tools the notion of unilateral shifts is successfully transmitted to Baer*-rings (see Def. 2.3 of the paper). I have a question in this context. To state it, some notations are given from the paper. $\bullet$ For a given y in A, the left projection of y is the smallest projection $[y]$ satisfying in $[y]y=y$. When A is a von algebra contained in $B(H)$ for some Hilbert space H then, $[y]$ is just the range projection of the operator y. $\bullet$ Let x be an isometry element in A that is $x^*x=1$. A projection p is called wandering for x if $[x^np][x^mq]=0$ for every different pair of natural numbers m and n. $\bullet$ An isometry x is called a unilateral shift in A if there exists a wandering projection p with $\sup_{n\geq0} \sum_0^n[x^np]=1$. Wold-decomposition in Baer*-rings. Let x be an isometry in A. There exists a commuting projection q with x such that x is decomposed to a direct sum $x=xq+(1-q)x$ of a unitary xq and unilateral shift $(1-q)x$ in the corner sub-rings Aq and $A(1-q)$ respectively. Question. Let A be a Baer*-ring. Let x be a unilateral shift and q be a finite projection in A. Can we conclude that $\inf_{n\geq1}[x^{*n}q]=0$? The answer is clearly yes in von Neumann algebras and seems to be YES in general case.

论坛问题帖 | no_answers_at_retrieval | ABB | 2018-05-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018189 | MO-302598

A question related to Conways 99 graph problem

I have observed that the number of triangles $\frac{vk}{6}$ of a strongly regular graph with parameters $(v,k,1,2)$ is given by the coefficient $2(k-1)$ in the molien series of the "4-D extraspecial group $2^{1+2\cdot 2}$": For example the beginning of the molien series is: $1 + t^2 + 5t^4 + 6t^6 + 15t^8 + 19t^{10} + 35t^{12} + 44t^{14} + 69t^{16} + 85t^{18}\\ + 121t^{20} + 146t^{22} + 195t^{24} + 231t^{26} + 295t^{28} + 344t^{30} + 425t^{32} + O(t^{34})$ We know that for $srg(9,4,1,2)$ we have $\frac{9\cdot4}{6}=6$ triangles and the coeffient $6 \cdot t^6$ tells us that at $6 = 2\cdot(4-1)$ we have 6 triangles. We know that for $srg(99,14,1,2)$ we have $\frac{99\cdot14}{6}=231$ triangles and the coeffient $231 \cdot t^{26}$ tells us that at $26 = 2\cdot(14-1)$ we have 231 triangles. And so on ... I have checked this for all known parameters for such hypothetical graphs: The parameters of all such graphs are: $(9,4,1,2),(99,14,1,2),(243,22,1,2),(6273,112,1,2),(494019,994,1,2)$ of which only the $n=9$ and $n=243$ are known. The Conway 99-graph problem asks if there exists such graph with 99 vertices. The number of triangles in such a (hypthetical) graph is: $6,231,891,117096,81842481$ (1) My (naive) question is, if there is a (deeper) connection, and maybe one can find unified way to construct all such graphs using the group above and the invariant homogenous polynomials (if this is not too much to ask)? (2) The 6 homogenous invariants of degree 6 are: $\frac14x_0^2x_1^2x_2^2 + \frac14x_0^2x_1^2x_3^2 + \frac14x_0^2x_2^2x_3^2 + \frac14x_1^2x_2^2x_3^2,$ $ \frac14x_0^4x_2^2 + \frac14x_0^2x_2^4 + \frac14x_1^4x_3^2 + \frac14x_1^2x_3^4,$ $ \frac14x_0^6 + \frac14x_1^6 + \frac14x_2^6 + \frac14x_3^6,$ $ \frac14x_0^3x_1x_2x_3 + \frac14x_0x_1^3x_2x_3 + \frac14x_0x_1x_2^3x_3 + \frac14x_0x_1x_2x_3^3,$ $\frac14x_1^4x_2^2 + \frac14x_1^2x_2^4 + \frac14x_0^4x_3^2 + \frac14x_0^2x_3^4,$ $ \frac14x_0^4x_1^2 + \frac14x_0^2x_1^4 + \frac14x_2^4x_3^2 + \frac14x_2^2x_3^4$ My question is, if one can find a way to associate to each such a homogenous invariant a unique triangle. I am struggling to find out what the set of vertices V with $|V|=9$ is. Edit: I found some way for the case $\deg=6$ with a brute force attack in SAGEMATH. But this attack doesn't work for $\deg=26$ since there I have to compute the common zeros of the 231 homogenous polynomials over some finite field. The problem I am facing is that the ideal generated by the homogenous invariants has dimension 2. But for a finite field there should be at most finitely many zeros. Any idea how to solve this in SAGEMATH (Computing the common zeros over some finite field?) J is the ideal of interest: sage: [I.gens() for I in J.minimal_associated_primes()] [[x1 + x2, x0 + x3], [x2 + x3, x0 + x1], [x1 + x3, x0 + x2]] sage: J.dimension() 2 https://oeis.org/A030533 https://pure.tue.nl/ws/portalfiles/portal/2157517 https://people.maths.bris.ac.uk/~matyd/GroupNames/1/ES+(2,2).html

论坛问题帖 | no_answers_at_retrieval | user6671 | 2018-06-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018190 | MO-303141

Is the conjecture A+B=C following correct?

Is the conjecture on A+B=C following correct ? Conjecture: Let A, B, C be three positive integer numbers such that $A+B=C$ with $\gcd(A, B, C) = 1$. By Fundamental theorem of arithmetic we write: $A=a_1^{x_1}a_2^{x_2}...a_n^{x_n}$, $B=b_1^{y_1}b_2^{y_2}...b_m^{y_m}$, $C=c_1^{z_1}c_2^{z_2}...c_k^{z_k}$ Let $d = \min\{x_i, y_j, z_h \}$ where $1 \le i \le n, 1\ \le j \le m, 1\le h \le k$ then $$d \le 5$$ PS: I read above one hunded papers, I observed that in any case $\min\{x_i, y_j, z_h \} \le 3$ Example 1: Ten solutions of Catalan-Fermat equation Example 2: $2^4.3^5.7^6+5^9.11^8=19^1.23^1.47^1.6679^1.3051977^1$ See also: Minimal exponent in prime factorization of n A051904 Nivens Constant

论坛问题帖 | no_answers_at_retrieval | Đào Thanh Oai | 2018-06-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018191 | MO-304819

$\sum_{i<j} x_{ij}^\alpha \ge \sum_{i<j} y_{ij}^\alpha $ for arbitrary cyclic polygons

I am looking for a proof of the inequality as follows: Conjecture: Let $A_1A_2....A_n$ be the regular polygon inscribed in a circle (O). Let $B_1B_2....B_n$ be another polygon inscribed in the circle (O). Denote $x_{ij}=A_iA_j$ and $y_{ij}=B_{i}B_{j}$ then: $$\sum_{i<j} x_{ij}^\alpha \ge \sum_{i<j} y_{ij}^\alpha $$ Where $ 1 \leq \alpha \leq 2$. The case $ \alpha = 1 $ was proved in our paper in here The case $ \alpha = 2 $ was proved in here Example: $n=3$, let ABC be a triangle with sidelength a, b, c then we have the inequality as follows: $$a^\alpha + b^\alpha+ c^\alpha \leq 3\times 3^{\frac{\alpha}{2}}R^\alpha$$ Where $ 1 \leq \alpha \leq 2$ and R is circumradius. See also:

论坛问题帖 | has_answers | Đào Thanh Oai | 2018-07-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018192 | MO-30590

Is there a progress on a solution of the inequality $\pi (m+n) \leq \pi (m) + \pi (n)$

in 1923 Hardy and Littlewood proposed the conjecture $\pi (m+n) \leq \pi (m) + \pi (n)$. Is there any progress towards solving this conjecture?

论坛问题帖 | has_answers | Hashem sazegar | 2010-07-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018193 | MO-306204

A symmetric polynomial inequality

I improve my previous question. Because this conjecture is exactly natural development of A Muirhead Like Inequality and Muirhead's Inequality so I think the conjecture is true. But I can not prove it. So I am looking for a proof of a conjecture as follows. Inequality 1: Let $n>2$ and $1 \le m \le n$ be integers. Let $x_1, \dots, x_n$ and $y_1,\dots, y_n$ be nonnegative real numbers such that $(x_1,\dots, x_n)$ majorizes $(y_1,\dots, y_n)$. Then for all reals $0 \leq a_1, a_2,\cdots,a_n \leq 1$, $$\sum\limits_{sym}\left(\sum\limits_{sym} x_{i_1}^{a_{p_1}} \cdots x_{i_m}^{a_{p_m}} \right) \leq \sum\limits_{sym}\left(\sum\limits_{sym} y_{i_1}^{a_{p_1}} \cdots y_{i_m}^{a_{p_m}} \right) $$ The summations are of course meant to be respectively over all m-tuples $(i_1,\dots,i_m),(p_1,\dots,p_m)$ with pairwise distinct entries. When $m=n$ this inequality is A Muirhead Like Inequality Inequality 2: Let $n>2$ and $1 \le m \le n$ be integers. Let $x_1, \dots, x_n$ and $y_1,\dots, y_n$ be nonnegative real numbers such that $(x_1,\dots, x_n)$ majorizes $(y_1,\dots, y_n)$. Then for all reals $ a_1, a_2,\dots,a_n \geq 0$, $$\sum\limits_{sym}\left(\sum\limits_{sym} a_{i_1}^{x_{p_1}} \cdots a_{i_m}^{x_{p_m}} \right) \geq \sum\limits_{sym}\left(\sum\limits_{sym} a_{i_1}^{y_{p_1}} \cdots a_{i_m}^{y_{p_m}} \right)$$ When $m=n$, this inequality is just Muirhead.

论坛问题帖 | has_answers | Đào Thanh Oai | 2018-07-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018194 | MO-306439

A rearrangement inequality for exponentiation function

Update: A year ago, but the first answer is not clear with me. I bounty this question again. My question: I am looking for a proof or counterexample to the following inequality: If $n \in \mathbb{N}$, and $a_1 \ge a_2 \ge \cdots \ge a_n \ge 0$ and $\alpha_1 \ge \alpha_2 \ge \cdots \ge \alpha_n \ge 0$ then $${\left(\sum_{i=1}^{n}{a_i^{\alpha_i}} \right)}^n \ge \prod_{i=1}^{n}{\left(\sum_{j=1}^{n}{a_i^{\alpha_j}} \right)}$$ If $n \in \mathbb{N}$ and $a_1 \ge a_2 \ge \cdots \ge a_n \ge 0$ and $0 \le \alpha_1 \le \alpha_2 \le \cdots \le \alpha_n$ then $${\left(\sum_{i=1}^{n}{a_i^{\alpha_i}} \right)}^n \le \prod_{i=1}^{n}{\left(\sum_{j=1}^{n}{a_i^{\alpha_j}} \right)}$$ Moreover equality holds in either case if and only if $a_1=a_2=a_3=...=a_n$. [The original question claimed a proof when $n=2$ or $n=3$.]

论坛问题帖 | has_answers | Đào Thanh Oai | 2018-07-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018195 | MO-309496

A non-rational variety with a full exceptional collection?

Does there exist a smooth non-rational projective variety whose bounded derived category of coherent sheaves admits a full exceptional collection? I could not find any examples in the literature (for instance projective spaces and intersections of quadrics, which admit full exceptional collections, are rational).

论坛问题帖 | has_answers | user74900 | 2018-08-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018196 | MO-309985

Existence of a Lipschitz map from a positive measure set to a ball

Question. If $A\subset \mathbb{R}^n$ is any set of positive Lebesgue n-measure, does there exists a Lipschitz map $f:A\to\mathbb{R}^n$ such that $f(A)$ is a ball with the same measure? In dimension $n=1$, such a map is easily found by defining $f(x) = \mathcal{L}^1(A\cap [0,x])$. In higher dimensions the problem becomes much more difficult. There is a short discussion on the topic in Ch. 7.10 of Mattila's 1995 book Geometry of Sets and Measures in Euclidean Spaces, where it is said that "Recently Preiss [6] proved that if A is an $\mathcal{L}^2$ measurable subset of $\mathbb{R}^2$ with $\mathcal{L}^2(A) > 0$, there is a Lipschitz map $f:A\to \mathbb{R}^2$ such that fA is a disc. In $\mathbb{R}^n$ for $n\geq 3$ the problem is unsolved." The reference given is [6] D. Preiss. Lipschitzian images of planar sets of positive measure, preprint However I was unable to find the mentioned paper of Preiss anywhere. At least no paper with such a title seems to exist on MathSciNet, arXiv or Preiss's list of publications. Given that Mattila referred to the result as "recent" in 1995, it seems unlikely that the paper would still be a preprint, so a few possibilities come to mind: The title changed before publication An error in the proof was found and the paper was never published The result was included in a different paper Any information about what has happened to the paper of Preiss, or any other references or information about the current state of the problem would be much appreciated.

论坛问题帖 | has_answers | Eero Hakavuori | 2018-09-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018197 | MO-31262

Smallest dilation of a quadrilateral?

What is the smallest dilation of a quadrilateral in $\mathbb{R}^d$? This may be an open problem; my question is: Is this indeed open? It will take me some time to explain the terms. The notion of dilation derives from Gromov, as far as I know (He defines a version in Metric Structures for Riemannian and Non-Riemannian Spaces, p.11, although he called it distortion). I came upon it myself via t-spanners. The version in which I am interested is this. Let P be a polygon (its boundary, not its interior), and x,y two points on P. You can think of P in $\mathbb{R}^2$, but also $\mathbb{R}^3$ and $\mathbb{R}^d$ for $d>3$ are interesting. Define $\delta(x,y)$ as the maximum (supremum) of $d_P(x,y) / | x y |$, where $d_P(x,y)$ is the distance between x and y following P (the shortest path staying on the closed path that consitutes P), and $|xy|$ is the Euclidean distance in $\mathbb{R}^d$. Thus $\delta(x,y)$ measures how much P dilates w.r.t. Euclidean distance. I am interested in the minimum value $\delta(P)$ of $\delta(x,y)$ over all $x,y \in P$, for all n-gons P, for fixed n. Example 1. If P is a unit square, then $\delta(x,y)$ for x,y opposite corners is $\sqrt{2}$, but $\delta(P)=2$ because with x,y midpoints of opposite sides, $\delta(x,y)= 2/1$. Example 2. If P is an equilateral triangle, $\delta(P)=2$, as shown in the figure. In fact, the dilation of any triangle is ≥ 2 [Lemma 7 in the 2nd paper below]. Example 3. It is known the the dilation of any closed curve C satisfies $\delta(C) \ge \pi/2$, with equality achieved only by the circle. [Corollary 23 in the first paper below.] This is (apparently) due to Gromov. So I finally come to my question. By reading these two papers, "Geometric Dilation of Closed Planar Curves: New Lower Bounds," and "On Geometric Dilation and Halving Chords," it appears to me that the minimum dilation of a quadrilateral in $\mathbb{R}^2$ (and $\mathbb{R}^d$) is not known. I had heard this was the case three years ago in a seminar in Brussels, but (a) I didn't quite believe it, (b) it was hearsay, and (c) it is now out of date. I am trying to clarify with the authors of these papers, but in parallel I would appreciate any information on the status of this question. The latter paper cited above proves a lower bound of $4 \tan(\pi/8) \approx 1.66$ (if I have interpreted it correctly). Addendum. I don't want to close-out this question, but I have heard from one of the authors of the above cited papers, and indeed it appears that the dilation of a planar quadrilateral is unknown [as of July 2010, the original posting date]. So I have tentatively tagged this as an open-problem, and I will update if new information surfaces. Thanks for everyone's interest and input!

论坛问题帖 | has_answers | Joseph O'Rourke | 2010-07-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018198 | MO-31315

Is Lebesgue's "universal covering" problem still open?

The following problem has been attributed to Lebesgue. Let "set" denote any subset of the Euclidean plane. What is the greatest lower bound of the diameter of any set which contains a subset congruent to every set of diameter 1? There are a number of interesting geometric problems of this type. Is it possible that some of them may be difficult to solve because the solution is a real irrational number which (when expressed in decimal form) is not even recursive-and so cannot be approximated in the usual way?

论坛问题帖 | has_answers | Garabed Gulbenkian | 2010-07-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018199 | MO-313313

Integral equality of 1st intrinsic volume of spheroid

Computations suggest that $$\int_{0}^{\infty}\int_{0}^{\infty} \sqrt{x+y^2} \cdot e^{-\frac{1}{2}(\frac{x}{s}+s^2y^2)}dxdy=\frac{2}{s}+\frac{2s^2\arctan(\sqrt{s^3-1})}{\sqrt{s^3-1}}.$$ The question is how to prove this equality. Background: this is the mean width of an ellipsoid with semiaxes $s,s,1/s^2$ and this almost completes the proof of the uniqueness hypothesis of an ellipsoid with given intrinsic volumes.

论坛问题帖 | has_answers | Alexander Tarasov | 2018-10-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018200 | MO-313639

Zero divisors in complex group algebras of residually finite groups

Conjecture. There exists a function $f:\mathbb{N} \rightarrow \mathbb{N}$ such that if α and β are non-zero elements of the complex group algebra $\mathbb{C}[G]$ of a finite group G such that $1\in \text{supp}(\alpha) \cap \text{supp}(\beta)$, $|\text{supp}(\alpha)|\leq |\text{supp}(\beta)|$ and $\alpha \cdot \beta =0$, then $\text{exp}\big (\langle \text{supp}(\alpha) \rangle \big) \leq f(|\text{supp}(\beta)|)$, where $\text{exp}(H)$ denotes the exponent of a finite group H and $\text{supp}(\gamma)$ for

an element $\gamma=\sum_{g\in G} \gamma_g \; g \in \mathbb{C}[G]$ denotes the set $\{ g\in G \;|\; \gamma_g \not=0\}$. Motivation. If the conjecture is true, then the support of any zero divisor of the complex group algebra of any residually finite group generates a finite subgroup. Proof. Let α be a non-zero element of $\mathbb{C}[G]$ for some residually finite group G such that $\alpha \cdot \beta =0$ for some non-zero $\beta \in \mathbb{C}[G]$. Let $H=\langle \text{supp}(\alpha) \rangle$. By Zelmanov's celebrated result on restricted Burnside problem, it is enough to show that the exponent of H is finite. Since H is finitely generated residually finite, there exists a descending series $H=N_1\geq N_2 \geq \cdots$ of normal subgroups H of finite index such that $\cap_{i\in\mathbb{N}} N_i=1$. Note that there exists $k\in\mathbb{N}$ such that $\bar{\alpha} \cdot \bar{\beta}=0$ in $\mathbb{C}[H/N_i]$ for all $i\geq k$ and $|\text{supp}(\bar{\beta})|=|\text{supp}(\beta)|=:t$, where $\bar{}$ is the natural ring epimorphism from $\mathbb{C}[H]$ onto $\mathbb{C}[H/N_i]$. Now if the above conjecture is true then $\text{exp}(H/N_i)\leq f(t)$ and so $\text{exp}(H)$ is finite. This completes the proof. If the above proof and conjecture are true then the complex group algebras of torsion-free residually finite groups have no zero divisor.

论坛问题帖 | has_answers | Alireza Abdollahi | 2018-10-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018201 | MO-313670

Zero divisors with support size 3 in complex group algebras of residually finite groups

Conjecture. There exists a function $f:\mathbb{N} \rightarrow \mathbb{N}$ such that if β is a non-zero element of the complex group algebra $\mathbb{C}[G]$ of a finite group G such that $1\in \text{supp}(\beta)$, and $(1+x+y) \cdot \beta =0$ or $(1+x-y) \cdot \beta =0$ for some distinct and nontrivial $x,y\in G$, then $\text{exp}\big (\langle x,y \rangle \big) \leq f(|\text{supp}(\beta)|)$, where $\text{exp}(H)$ denotes the exponent of a finite group H and $\text{supp}(\gamma)$ for an element $\gamma=\sum_{g\in G} \gamma_g \; g \in \mathbb{C}[G]$ denotes the set $\{ g\in G \;|\; \gamma_g \not=0\}$. Motivation. If the conjecture is true, then the support of any zero divisor of the form $1+x+y$ or $1+x-y$ of the complex group algebra of any residually finite group generates a finite subgroup. Proof. Let $\alpha=1+x+y$ or $1+x-y$ be a non-zero element of $\mathbb{C}[G]$ for some residually finite group G such that $\alpha \cdot \beta =0$ for some non-zero $\beta \in \mathbb{C}[G]$. Let $H=\langle \text{supp}(\alpha) \rangle$. By Zelmanov's celebrated result on restricted Burnside problem, it is enough to show that the exponent of H is finite. Since H is finitely generated residually finite, there exists a descending series $H=N_1\geq N_2 \geq \cdots$ of normal subgroups H of finite index such that $\cap_{i\in\mathbb{N}} N_i=1$. Note that there exists $k\in\mathbb{N}$ such that $\bar{\alpha} \cdot \bar{\beta}=0$ in $\mathbb{C}[H/N_i]$ for all $i\geq k$ and $|\text{supp}(\bar{\beta})|=|\text{supp}(\beta)|=:t$, where $\bar{}$ is the natural ring epimorphism from $\mathbb{C}[H]$ onto $\mathbb{C}[H/N_i]$. Now if the above conjecture is true then $\text{exp}(H/N_i)\leq f(t)$ and so $\text{exp}(H)$ is finite. This completes the proof. If the above proof and conjecture are true then the integral group algebras of torsion-free residually finite groups have no zero divisor with support size 3.

论坛问题帖 | no_answers_at_retrieval | Alireza Abdollahi | 2018-10-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018202 | MO-31406

Does listing the prime factors always stop?

Take a natural number's prime factors and list them increasingly and repeating them according to multiplicity. Concatenate their decimal (or in any base) representation to get a new number and repeat the process. Does this always end in a prime number for any input?

论坛问题帖 | has_answers | O.R. | 2010-07-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018203 | MO-314561

Asymptotics of Littlewood polynomials

Littlewood in [L] states several conjectures regarding asymptotics of polynomials with ±1 coefficients. He considers the class $\mathscr F$ of polynomials of form $\sum^n\pm z^m$ and asks whether there exists a sequence f_n in $\mathscr F$ such that $$A\leq\frac{|f_n(\theta)|}{\sqrt{n+1}}\leq B$$ for large n's. 1. I'd like to know if this is still open and would be grateful if someone could provide references for recent survey papers. 2. If the conjecture is not settled yet, I was thinking if one can apply integer programming methods to tackle the problem? Update 1 As pointed out by Fedor, J-P. Kahane's work gives an elegant proof for polynomials with unimodular coefficients. His result is as follows(I didn't translate it to English as seems rather easy to understand): Theoreme. Il existe une suite de polynomes $$P_n(z)=\sum_{m=1}^{n}a_{m,n}z^m,\,(|a_{m,n}| = 1; \, n=1,2,\ldots,m=1,\ldots,n)$$ et une suite ϵ_n positive tendant vers 0 telles que pour tout z de module 1 on ait$$(1-\epsilon_n)\sqrt{n}\leq|P_n(z)|\leq(1+\epsilon_n)\sqrt{n}.$$ Numerical experiments suggest there might be hope to construct a sequence of $P_n\in\mathbb {Z}_2[x]$ which satisfies Kahane's asymptotics. [L]Littlewood, J. E., On polynomials $\sum^n\pm z^m, \sum^n e^{\alpha_mi}z^m, z = e^{\theta i}$, J. Lond. Math. Soc. 41, 367-376 (1966). ZBL0142.32603.

论坛问题帖 | no_answers_at_retrieval | BigM | 2018-11-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018204 | MO-31482

The "sensitivity" of 2-colorings of the d-dimensional integer lattice

Consider the d-dimensional integer lattice, Z^d. Call two points in Z^d "neighbors" if their Euclidean distance is 1 (i.e., if they differ by 1 on exactly one coordinate). Let C be a two-coloring of Z^d, which makes each point either red or blue. We'll assume C has the following "nontriviality" property: the origin is colored red, but on each of the d axes through the origin, there's a point on that axis that's colored blue. Let the "sensitivity" of a point x with respect to C, or $s_x(C)$, be the number of x's neighbors that are colored differently from x. Then let $s(C) = \max_{x \in Z^d} s_x(C)$. QUESTION: Can you give me any decent lower bound on $s(C)$ in terms of d? For example, that $s(C) \ge k \sqrt{d}$ for some constant $k > 0$? REMARK 1: If you prove a lower bound of the form $k d^l$ (for

constants $k,l > 0$), then you'll have solved an old open problem in the study of Boolean functions, namely the "sensitivity versus block sensitivity" problem (posed by Noam Nisan in 1991). But please don't let that discourage you! My variant feels more approachable, and maybe something is even already known about it. (I'll be happy to supply full details of the reduction on request. But here's the basic idea: let $f : \lbrace 0,1 \rbrace ^n \rightarrow \lbrace 0,1 \rbrace$ be a Boolean function such that the block sensitivity $bs(f)$ is much much larger than the sensitivity $s(f)$. Then there must be an input x of f that has $bs(f)$ disjoint sensitive blocks. Let $d=bs(f)$. Then we can construct a two-coloring of Z^d with the properties listed above, and such that $s(C) \le 2 s(f)$ where $s(f)$ is the sensitivity of f. The input x gets mapped to the origin of Z^d, while each of the d sensitive blocks of x gets mapped to one of the axes of Z^d. To map a Boolean assignment to an integer, in a way that preserves the sensitivity, we use the Gray Code. Finally, we color each point $y \in Z^d$ either red or blue, depending on whether $f(x)$ is 0 or 1 for the corresponding Boolean point x.) REMARK 2: I can give an example of a coloring with $s(C) = O(\sqrt{d})$, meaning that $s(C) \ge k \sqrt{d}$ really is the best lower bound one can hope for. This coloring can be obtained by starting from "Rubinstein's function" -- a Boolean function $f : \lbrace 0,1 \rbrace ^n \rightarrow \lbrace 0,1 \rbrace$ with $bs(f) = n/2$ and $s(f) = 2 \sqrt{n}$ -- and then applying the reduction sketched in Remark 1. (For those who are interested, let me now go ahead and describe a coloring with $s(C) = O(\sqrt{d})$ explicitly. Assume for simplicity that d is a perfect square. Partition the d coordinates of x into $\sqrt{d}$ "blocks" of $\sqrt{d}$ coordinates each. Then we'll color x blue, if and only if at least one of the blocks has a single coordinate equal to 2 and all other coordinates equal to 0. I'll leave it as an exercise for you to verify that $s(C) = 2 \sqrt{d}$.) Note: I edited the above paragraph a little, to simplify the construction and insert a missing factor of 2. REMARK 3: At the moment, I don't even have a proof that $s(C)$ has to grow with d (!!). But I suspect at least $s(C) \ge k \log d$ ought to be doable. EDIT: Sorry to switch notations in the middle of the game, but I have a better one if you want to talk about low dimensions (per domotorp's question below)! Let's let $r_x(C)$ be the number of axes (up/down, left/right, etc.) along which x has a neighbor that's colored differently than x is. Then let $r(C) = \max_x r_x(C)$. Clearly $r(C) \le s(C) \le 2r(C)$ for all C. In fact, something even stronger than that is true: given any coloring C, one can create a new coloring C' that satisfies $s(C')=r(C)$, by simply "blowing up" each point x into a cube of 2^d points, which are all colored the same way x was colored in C. The nontriviality and sensitivity properties will clearly be preserved; all this transformation does is to eliminate the problem of a point having two differently-colored neighbors along the same axis. So without loss of generality, we can shift attention to $r(C)$. Now let $r_d = \min_C r(C)$ over all nontrivial colorings C of Z^d. Then here's what I know: $r_1 = 1$ $r_2 = 2$ $r_3 = 2$ $r_4 = 2$ $r_5 \in \lbrace 2,3 \rbrace$ UPDATE: I created an image that shows an explicit coloring of Z^3 that satisfies both the nontriviality condition and $r(C)=2$. (That is, from every point, you can change color by moving along at most 2 different axes.) As explained above, this can easily be converted into a coloring with $s(C)=2$ as well. (source) domotorp is right that proving $r_5=3$ could be a great start...

论坛问题帖 | has_answers | Scott Aaronson | 2010-07-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018205 | MO-315051

Is it known how the Sigma Algebra generated by Jordan measurable sets compares to universally measurable sets and analytic sets?

Unlike the collection L of Lebesgue measurable sets, the collection J of Jordan measurable sets do not form a Sigma algebra. (A set is Jordan measurable if and only if its characteristic function is Riemann integrable, and the characteristic function of a singleton is Riemann integrable, but the characteristic function of the rational numbers is not.) But we can still talk about $\sigma(J)$, the Sigma algebra generated by J. Now in this 1993 journal paper, K.G. Johnson shows that the Borel Sigma algebra is a proper subset of $\sigma(J)$ which is a proper subset of the Lebesgue Sigma algebra, and that a set is in $\sigma(J)$ if and only if it can be written as a union of a Borel set and a subset of a measure 0 F_σ set. But Johnsons says there are two questions he doesn’t know the answer to: Where does $\sigma(J)$ stand relative to the analytic sets ... and relative to the universally measurable sets[?] My question is, have either of these questions been answered in the two decades since this paper was published? Is it known whether there are analytic sets which are not in $\sigma(J)$ or vice versa? Is it known whether there are universally measurable sets which are not in $\sigma(J)$ or vice versa? Or are these still open problems?

论坛问题帖 | has_answers | Keshav Srinivasan | 2018-11-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018206 | MO-315664

For what sets does the Lebesgue Differentiation Theorem hold in one dimension?

Lebesgue's differentiation theorem states that if x is a point in $\mathbb{R}^n$ and $f:\mathbb{R}^n\rightarrow\mathbb{R}$ is a Lebesgue integrable function, then the limit of $\frac{\int_B f d\lambda}{\lambda(B)}$ over all balls B centered at x as the diameter of B goes to 0 is equal almost everywhere to $f(x)$. But if you replace balls with other kinds of set with diameter going to 0, this need not be true. For instance it need not be true if you replace balls with rectangles. But my question is, what is known about what collections of sets the Lebesgue Differentiation Theorem holds for in one dimension? We know it holds for bounded intervals. Does it hold for finite unions of bounded open intervals? What about bounded Borel sets? What about bounded Lebesgue measurable sets in general? Or are all these open problems?

论坛问题帖 | has_answers | Keshav Srinivasan | 2018-11-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018207 | MO-316075

Is a maximal set of rectangles known for which Lebesgue’s Differentiation Theorem holds true?

Lebesgue's differentiation theorem states that if x is a point in $\mathbb{R}^n$ and $f:\mathbb{R}^n\rightarrow\mathbb{R}$ is a Lebesgue integrable function, then the limit of $\frac{\int_B f d\lambda}{\lambda(B)}$ over all balls B centered at x as the diameter of B goes to 0 is equal almost everywhere to $f(x)$. But if you replace balls with other kinds of set with diameter going to 0, this need not be true. For instance it need not be

true if you replace balls with rectangles, even rectangles with sides are parallel to the coordinate axes. But if we restrict things to L^p functions where $1<p<\infty$, then Lebesgue's differentiation theorem does hold true for rectangles with sides parallel to the coordinate axes, but it does not hold true for arbitrary rectangles. So my question is, is a maximal set of directions known for which Lebesgue's differentiation theorem holds for L^p functions for the collection of all rectangles oriented in those directions? Now this journal paper says that a maximal set of directions is not known for tubes in $\mathbb{R}^n$ for which Lebesgue's differentiation theorem holds (though the paper makes a conjecture about it). And a rectangle is nothing but a tube in $\mathbb{R}^2$. But it's possible that the answer is known for this special case.

论坛问题帖 | no_answers_at_retrieval | Keshav Srinivasan | 2018-11-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018208 | MO-316543

An open problem in Sobolev spaces

Let $\Omega\subset\mathbb{R}^n$ be a bounded domain. Suppose that there there is a bounded extension operator $$ E:W^{1,p}(\Omega)\to W^{1,p}(\mathbb{R}^n) \quad \text{and} \quad E:W^{1,q}(\Omega)\to W^{1,q}(\mathbb{R}^n), $$ where $1\leq p<q\leq\infty$ (or $1<p<q<\infty$ if you find it easier). Open problem. Does it follow that $$ E:W^{1,r}(\Omega)\to W^{1,r}(\mathbb{R}^n) $$ for all $p<r<q$? Some partial results are known, but the general case seems to be difficult. It looks like a simple interpolation exercise, but it is not. Later, when I have time I will add some references to known results.

论坛问题帖 | no_answers_at_retrieval | Piotr Hajlasz | 2018-11-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018209 | MO-318917

A tantalizing Gamma quotient to challenge the Rohrlich-Lang Conjecture

The Rohrlich-Lang Conjecture for polynomial relations in Gamma values predicts that all polynomial relations between Gamma values over $\mathbb Q$ come from the functional equations satisfied by the Gamma function. An other statement, somewhat narrower: if a quotient of products of gamma values over $\mathbb Q$ is an algebraic number, then it can be evaluated by using only the well-known reflection and multiplication formulae. Lang, S. - Relations de distributions et exemples classiques. Séminaire Delange-Pisot-Poitou, 19e année: 1977/78, Théorie des nombres, Fasc. 2, Exp. N° 40, 6 p. Collected Papers, vol. III, Springer (2000), 59-65. I am wondering what that means for the following. Define $$A:=\prod_{k=0}^5{\Gamma(\frac{49^k \pmod {78}}{78})}= {\Gamma(\frac{1}{78})\Gamma(\frac{25}{78})\Gamma(\frac{43}{78})\Gamma(\frac{49}{78})\Gamma(\frac{55}{78})\Gamma(\frac{61}{78})} $$ and $$B:=\prod_{k=0}^5 {\Gamma(\frac{7\cdot49^k\pmod {78}}{78})}= \Gamma(\frac{7}{78})\Gamma(\frac{19}{78})\Gamma(\frac{31}{78})\Gamma(\frac{37}{78})\Gamma(\frac{67}{78})\Gamma(\frac{73}{78}).$$ Then numerically, it appears that $$\frac AB=4.$$ Equivalently, as $AB=(2\pi)^6\;\sqrt[3\ \ ]{13}$ by the multiplication formula, we'd have to show $$A=16\pi^3\;\sqrt[6\ \ ]{13}\ \ \text{ and / or }\ \ B=4\pi^3\;\sqrt[6\ \ ]{13}.$$ So the product of the 12 Gamma terms $\Gamma(\frac{6j+1}{78})$ with $j=0,1,3,...,12$ (excluding $j=2$ where $\frac{13}{78}=\frac16$) splits into two halves, one 4 times as big as the other. How to "get rid of" the numerators, which are all $\equiv1\pmod6$? Applying the multiplication formula with step $\frac16$ to each term as suggested in this comment has the same effect (in terms of producing algebraic factors that can then be "ignored") as applying the reflection formula to each term: applying it for instance to A will leave us with $\Gamma(\frac{77}{78})\Gamma(\frac{53}{78})\Gamma(\frac{35}{78})\Gamma(\frac{29}{78})\Gamma(\frac{23}{78})\Gamma(\frac{17}{78})$ in the denominator, no gain. I am aware that there can be surprises like the fact that $\frac{ \Gamma\left(\frac{11}{42}\right)\Gamma\left(\frac{12}{42}\right)}{\Gamma\left(\frac{21}{42}\right)\Gamma\left(\frac2{42}\right)} $ and $\frac{ \Gamma\left(\frac{18}{42}\right)\Gamma\left(\frac{4}{42}\right)}{\Gamma\left(\frac{21}{42}\right)\Gamma\left(\frac1{42}\right)} $ are algebraic and I would not dare to claim that the identity $\frac AB=4$ provides a counterexample to the Rohrlich-Lang conjecture. It is just that the simplicity of the result, being an integer, is particularly intriguing. Is there any other way to evaluate this, using only the functional equations of the Gamma function? Has anybody ever written a program for that (which is rather a combinatorial issue that should not be awfully hard)?

论坛问题帖 | has_answers | Wolfgang | 2018-12-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018210 | MO-318972

Percentage of Ramanujan's conjectures that were proven correct

Today I read the following brief but insightful account of Ramanujan's approach to mathematics: https://www.imsc.res.in/~rao/ramanujan/images/KSRchap3.pdf and while reading this I wondered whether we have a lower-bound on the percentage of Ramanujan's conjectures which are correct. I'm planning to get a copy of Ramanujan's notebooks. Meanwhile, the above question intrigues me.

论坛问题帖 | has_answers | Aidan Rocke | 2018-12-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018211 | MO-31920

Curves of constant curvature on S^2

Most probably this is a well known question. Consider S^2 with a Riemannian metric. I would like to ask what is known about the structure of the set of simple (without self-intersections) closed curves on it of constant geodesic curvature. Here is a series of questions. 1) Is this true that through each point of S^2 passes a simple closed curve of constant curvature? If not, can one estimate from below the proportion of the area of S^2 covered by such curves? 2) Is it true that for each value of curvature there are at least 2 simple closed curves on S^2 of this curvature? Or maybe even more than 2? 3) What can be said about the global structure of these curves on a generic S^2? Taking the union of all such closed curves we could try to cook up from them a surface (that maps naturally to S^2). Is something known about the topology of this surface? Comments 1) The theorem of Birkhoff states that each Riemannian S^2 contains at least three simple closed geodesics, as Joseph remarks below. 2) For a generic metric on S^2 the set of such curves this set should be one dimensional. Indeed for each fixed value of curvature you can consider

an analogue of the geodesic flow on the space of unite tangent vectors to S^2 and you expect that closed orbits will be isolated. ADDED. Is seems indeed that these are open (and I guess hard) questions. Macbeth gave a very nice reference, that tells in particular that similar questions were raised previously by Arnold, I copy the Macbeth's reference here, so it is visible to everyone: http://count.ucsc.edu/~ginzburg/ARNOLD/mag-post.pdf Update. The following reference : http://arxiv.org/abs/0903.1128 gives a positive answer to question 2) for spheres of non-negative Gaussian curvature provided we consider not only simple curves on S^2 but also curves that bound immersed disks. One more update. There is a new nice article http://arxiv.org/abs/1105.1609 that provides some further results concerning question 2) for S^2 of positive curvature. This article also gives all necessary references from which one can conclude that question 2) was considered by Poincare in 1905, as it is written in the article of S.P. Novikov
http://iopscience.iop.org/0036-0279/37/5/R01/pdf/0036-0279_37_5_R01.pdf

论坛问题帖 | has_answers | Dmitri Panov | 2010-07-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018212 | MO-320323

Integer valued polynomials over several variables

For simplicity this is about polynomials in just two variables. Any $f\in\mathbb Q[X,Y]$ can be written as a linear combination of monomials X^iY^j and therefore as a sum of polynomials $p_{ij}\in\mathbb Q[X^iY^j]$ over one variable: $$\displaystyle f(X,Y)=p_{00}+\sum_{\gcd(i,j)=1}p_{ij}(X^iY^j).$$ My question: is the subring of all integer valued polynomials f over two variables identical with the subring of all sums of integer valued polynomials p_{ij} over one variable as above? An integer valued polynomial in one variable is a polynomial p with rational coefficients such that $p(\mathbb Z)\subseteq \mathbb Z$. And corresponding for polynomials over several variables. It might be some abuse of language to call p_{ij} polynomials over one variable, they may rather be polynomials over one variable applied to monomials X^iY^j.

论坛问题帖 | has_answers | Lehs | 2019-01-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018213 | MO-321267

Open Problems in Random Graphs

I am a PhD student in mathematics. I'm interested in probabilistic methods in combinatorics and especially random graphs. I am looking for an open problem in this area for my PhD proposal. I know that there are many open problems and conjectures, but I would like to find a "good" problem such that I can get some results during my PhD. It would be very helpful for me if you could suggest a bunch of open problems. Thank you!

论坛问题帖 | has_answers | Henry Zagreb | 2019-01-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018214 | MO-322240

Why is $\sup f_- (n) \inf f_+ (m) = \frac{5}{4} $?

This question is an old question from mathstackexchange. Let $f_- (n) = \Pi_{i=0}^n (\sin(i) - \frac{5}{4}) $ And let $ f_+(m) = \Pi_{i=0}^m (\sin(i) + \frac{5}{4}) $ It appears that we have $$\sup f_- (n) \inf f_+ (m) = \frac{5}{4} $$ Why is that so ? Notice $$\int_0^{2 \pi} \ln(\sin(x) + \frac{5}{4}) dx = Re \int_0^{2 \pi} \ln (\sin(x) - \frac{5}{4}) dx = \int_0^{2 \pi} \ln (\cos(x) + \frac{5}{4}) dx = Re \int_0^{2 \pi} \ln(\cos(x) - \frac{5}{4}) dx = 0 $$ $$ \int_0^{2 \pi} \ln (\sin(x) - \frac{5}{4}) dx = \int_0^{2 \pi} \ln (\cos(x) - \frac{5}{4}) dx = 2 \pi^2 i $$ That explains the finite values of $\sup $ and $ \inf $.. well almost. It can be proven that both are finite. But that does not explain the value of their product. Update This is probably not helpful at all , but it can be shown (not easy) that there exist a unique pair of functions $g_-(x) , g_+(x) $, both entire and with period $2 \pi $ such that $$ g_-(n) = f_-(n) , g_+(m) = f_+(m) $$ However i have no closed form for any of those ... As for the numerical test i got about $ln(u) (2 \pi)^{-1}$ correct digits , where $u = m + n$ and the ratio m/n is close to 1. Assuming no round-off errors i ended Up with $1.2499999999(?) $. That was enough to convince me. I often get accused of " no context " or " no effort " but i have NOO idea how to even start here. I considered telescoping but failed and assumed it is not related. Since I also have no closed form for the product I AM STUCK. I get upset when people assume this is homework. It clearly is not imho ! What kind of teacher or book contains this ? --- Example : Taking $m = n = 8000 $ we get $$ max(f_-(1),f_-(2),...,f_-(8000)) = 1,308587092.. $$ $$ min(f_+(1),f_+(2),...,f_+(8000)) = 0,955226916.. $$ $$ 1.308587092.. X 0.955226916.. = 1.249997612208568.. $$ Supporting the claim. I'm not sure if $sup f_+ = 7,93.. $ or the average of $f_+ $ ($ 3,57..$) relate to the above $1,308.. $ and $0,955..$ or the truth of the claimed value $5/4$. In principle we could write the values $1,308..$ and $0,955..$ as complicated integrals. By using the continuum product functions $f_-(v),f_+(w)$ where v,w are positive reals. This is by noticing $ \sum^t \sum_i a_i \exp(t \space i) = \sum_i a_i (\exp((t+1)i) - 1)(\exp(i) - 1)^{-1} $ and noticing the functions f_+,f_- are periodic with 2π. Next with contour integration you can find min and max over that period 2π for the continuum product functions. Then the product of those 2 integrals should give you $\frac{5}{4}$. -- Maybe all of this is unnecessarily complicated and some simple theorems from trigonometry or calculus could easily explain the conjectured value $\frac{5}{4}$ .. but I do not see it. -- -- Update This conjecture is part of a more general phenomenon. For example the second conjecture : Let $g(n) = \prod_{i=0}^n (\sin^2(i) + \frac{9}{16}) $ $$ \sup g(n) \space \inf g(n) = \frac{9}{16} $$ It feels like this second conjecture could somehow follow from the first conjecture since $$-(\cos(n) + \frac{5}{4})(\cos(n) - \frac{5}{4}) = - \cos^2(n) + \frac{25}{16} = \sin^2(n) + \frac{9}{16} $$ And perhaps the first conjecture could also follow from this second one ? Since these are additional questions and I can only accept one answer , I started a new thread with these additional questions :
https://math.stackexchange.com/questions/3000441/why-is-inf-g-sup-g-frac916 ---EDIT--- I want to explain better how to get a closed form for these numbers. I already mentioned that the periods of these functions are 2π and how to use that. But those few lines deserve more attention. Basically this is what we do : We use the fourier series $$ f(x) = \ln(\sin(x) + 5/4) = \sum_{k=1}^{\infty} \frac{-2 \cos(k(x + \pi/2))}{2^k k} $$ Now we use the inverse of the backward difference operator (similar but distinct from the so-called "indefinite sum"

which is defined as inverse of the forward difference operator) In other words we solve for $F(x)$ such that $$F(x) - F(x-1) = f(x)$$ We call this the " continuum sum " (CS) and write/define : $$ CS(f(x),x) := F(x) $$ $$ CS(f(x),y) := F(y) $$ For clarity : $$ \sum_0^y f(x) = CS(f(x),y) - CS(f(x),-1) $$ $$ \sum_0^0 f(x) = CS(f(x),0) - CS(f(x),-1) = f(0) $$ This operator is linear so we make use of that : $$CS(2 \cos(k(x+\pi/2),x) = \csc(k/2) \sin(k(x+1/2 + \pi/2)) -1.$$ This implies that : $$ CS(f(x),x) = F(x) = - \sum_{k=1}^{\infty} \frac{\csc(k/2)\sin(k(x + 1/2 + \pi/2)) - 1}{2^k k} $$ and $$ G(x) = \frac{d F(x)}{dx} = - \sum_{k=1}^{\infty} \frac{\csc(k/2) \cos(k(x+1/2+\pi/2))}{2^k} $$ Now $\sup f_+ = 7.93.. $ is the supremum of $ \exp(F(x) - F(-1) $ and $\inf f_+ = 0.95..$ is the infimum of $ \exp(F(x) - F(-1) $. And both these values are achieved at x such that $G(x) = 0$. The analogue for f_- works. So the numbers from the OP can (more or less) be given by these infinite series and hence the whole conjectures can be stated by these infinite series. We also know for instance $$ \sum_{k=1}^{\infty} \frac{1}{k 2^k} = \ln(2) , \sum_{k=1}^{\infty} \frac{(-1)^k}{k 2^k} = - \ln(3/2) $$ So that is hopefull. Also the max and min of functions can be given by contour integrals but that might not make things easier ? Many trig identities and symmetry are probably related. But I see no clear proof. So that is how we compute the values and it might just help. Also notice : $$ t(x) = \ln(\sin(x) - 5/4) = \ln(-1) + \ln(\sin(-x) + 5/4) = \ln(-1) + \sum_{k=1}^{\infty} \frac{-2 \cos(k(x + \pi/2))}{2^k k} $$ So the other case is no mystery. And ofcourse the case $ \ln(\cos^2(x) - 9/16) $ is also simply related. (trig addition identities can be used) EDIT : I wanted to make as little conjectures as possible when I posted this. Just one question/conjecture per post is the usual rule. But many related conjectures exist. Many might turn out to be equivalent or have already been shown to be equivalent. (such as the analogue cosine cases with $\frac{3}{4}$) I thought it would be best not to flood with related conjectures and post the most important one. Which I did. But one comment of Richard can not be ignored. Richard Stanley wrote $$ \sup f_-(n)^2 - \inf f_+(n)^2 = \frac{4}{5} $$ He might not be the first to notice this, even if I exclude my mentor who came up with the main conjecture and related ones. (I think it was also mentioned on chat and comments on MSE) Anyways it is also a nice conjecture. Is it equivalent ? No. Lets use shorthand notations : $x = \sup f_-(n), y = \inf f_+$. Then combining both conjectures (the original and the "Richard" variant) we get $$x y = \frac{5}{4} , x^2 - y^2 = (x-y)(x+y) = \frac{4}{5}$$ Now this implies we can compute the value of x (or y). We get an interesting situation here. If 2 conjectures are true than so is the third : $$x y = \frac{5}{4}$$ $$ x^2 - y^2 = (x-y)(x+y) = \frac{4}{5} $$ x is the positive real solution to $5 * 4^2 x^4 - 4^3 x^2 - 5^3 $ or $80 x^4 - 64 x^2 - 125$. NOTICE the 5's and 4's all over again. (or closed form for $x = \frac{1}{2} \sqrt {\frac{8 + \sqrt {689}}{5}}$) ($689 = 13*53$ if that matters to anyone) This was all known in the $90's$ by my mentor. I call it the Raes-Stanley conjecture. Update ! I had a talk with my mentor why he did not mention the $$ \sup f_-(n)^2 - \inf f_+(n)^2 = \frac{4}{5} $$ part of the Raes-Stanley conjecture. Although he noticed the apparant identity, he does not actually believe that last part. He said that the value converges fast for $n,m > 210000$ to $$ \sup f_-(n)^2 - \inf f_+(n)^2 = \frac{3.999789007..}{5} $$ In fact increasing n,m from 210000 to 314314 both gives $$ \sup f_-(n)^2 - \inf f_+(n)^2 = \frac{3.999789007..}{5} $$ so barely any noticeable change, while $$\sup f_- (n) \inf f_+ (m) = \frac{5}{4} $$ seems already heuristically confirmed. Testing for n,m smaller than 100000 might give the wrong impression and might be the cause of the mistake. Numerical coincidence might then lead to wrong conclusions. Roundoff errors might create an effect in the computations but he does not believe the "Stanley part " $$\sup f_- (n) \inf f_+ (m) = \frac{5}{4} $$ can be saved by those errors. If anyone can prove or argue or compute a higher value than $\frac{3.999789007..}{5}$, please inform me. Finally the Full Raes conjecture is the slight generalization : Let $f_- (n) = \Pi_{i=0}^n (\sin(i) - \frac{5}{4}) $ And let $ f_+(m) = \Pi_{i=0}^m (\sin(i) + \frac{5}{4}) $ It appears that $$\sup f_- (n) \inf f_+ (m) = \frac{5}{4} $$ and $$\inf |f_- (n)| \sup f_+ (m) = \frac{5}{4} $$ Why is that so ? $|*|$ means the absolute value here. Indeed $$\inf |f_- (n)| = 0.157559...$$ $$\sup f_- (n) = 1.308592...$$ $$\inf f_+ (n) = 0.955225...$$ $$\sup f_+ (n) = 7.933553...$$ $$0.955225... * 1.308592... = 1.25 = \frac{5}{4}$$ $$0.157559... * 7.933553... = 1.25 = \frac{5}{4}$$ Let $g_- (n) = \Pi_{i=0}^n (\cos(i) - \frac{5}{4}) $ And let $ g_+(m) = \Pi_{i=0}^m (\cos(i) + \frac{5}{4}) $ It appears that $$\sup g_- (n) \inf g_+ (m) = \frac{3}{4} $$ and $$\inf |g_- (n)| \sup g_+ (m) = \frac{3}{4} $$

论坛问题帖 | has_answers | mick | 2019-02-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018215 | MO-32226

Lower bounds (or less) for the period of sqrt(D) and related sequences

This is a continuation of Lower bounds for period length of continued fraction of square root which is a continuation of Upper bound of period length of continued fraction representation of very composite number square root. Yesterday, I came across a nice article with many problems on continued fractions of Michel Mendes France (from 1993): http://retro.seals.ch/digbib/fr/view?rid=ensmat-001:1993:39::498&id=&id2=&id3= (broken link) In problem 6 he asks: Let x be quadratic irrational and $\pi(x)$ the length of its period. Is $sup_n(\pi(x^n))=\infty$? Is this problem still open? I saw many articles about upper bounds for $\pi(\sqrt{D})$ but none on lower bounds (expect the Golubeva one that is mentioned in the first mentioned question, but due to lack of background, I fail to understand the results there). As Will Jagy asked: Do we even know whether $[\pi(\sqrt{n})):n\in \mathbb N]$ is bounded or not?

论坛问题帖 | has_answers | Menny | 2010-07-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018216 | MO-322598

Open questions about posets

Partially ordered sets (posets) are important objects in combinatorics (with basic connections to extremal combinatorics and to algebraic combinatorics) and also in other areas of mathematics. They are also related to sorting and to other questions in the theory of computing. I am asking for a list of open questions and conjectures about posets.

论坛问题帖 | has_answers | Gil Kalai | 2019-02-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018217 | MO-324214

Open problems in Sobolev spaces

What are the open problems in the theory of Sobolev spaces? I would like to see problems that are yes or no only. Also I would like to see problems with the statements that are short and easy to understand for someone who has a basic knowledge in the theory, say at the level of the book by Evans and Gariepy. The problems do not have to be a well know ones. Just the problems you think are interesting. Please, list each problem as a separate answer. That will allow people to leave comments related exclusively to this particular problem. I have been working with Sobolev spaces for most of my adult live and I have some of my favorite problems that I will list below. But I will do it later, because first I would like to see your problems.

论坛问题帖 | has_answers | Piotr Hajlasz | 2019-02-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018218 | MO-32477

Smallest area shape that covers all unit length curve

On a euclidean plane, what is the minimal area shape S, such that for every unit length curve, a translation and a rotation of S can cover the curve. What are the bounds of the shape's area if this is a open problem? When I asked this problem few years ago, someone told me it's open. I don't know if this is still open and I can't find any reference on it. I don't even know what branch of mathematics it falls under. so I can't even tag this question.

论坛问题帖 | has_answers | Chao Xu | 2010-07-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018219 | MO-325899

Symmetric function transition matrix and a non-conjecture by Clifford and Stanley

Consider the transition matrix $R = \left(R_{\lambda,\mu}\right)$, defined by $$ p_\lambda = \sum_{\mu} R_{\lambda\mu}m_\mu , $$ between the power-sum and the monomial basis of the ring of symmetric functions. There are plenty of combinatorial descriptions of $R_{\lambda\mu}$, it is for example the number of ordered brick-tableaux of shape λ and brick-sizes determined by μ, see e.g. this excellent paper with transition matrices by Eğecioğlu and Remmel. In the paper on Bottom Schur functions by P. Clifford and R. Stanley (2004), the authors consider the dimension of the null-space of $R-D$, where D is the diagonal of R. As the size n of the partitions increase, they computed (by computer I presume) the sequence of dimensions $d(n)$ is $$ 1 , 1 , 1 , 2 , 2 , 3 , 4 , 5 , 7 , 9 , 11 , 15 , 19 , 24 $$ but they do not make any conjecture of what this sequence of integers might be. However, a quick search in OEIS gives one result, A129528, entered in 2007, after Clifford and Stanley's paper was published. The sequence A129528, $a(n)$ is defined as follows: Consider the set of all Dyck paths, consisting of steps $(1,1)$ and $(1,-1)$. The peak-abscissa-sum of a path is the sum of the x-coordinates of all peaks. Then $a(n)$ is the number of Dyck paths of any length with peak-abscissa-sum n. Question: Is there some neat way to show that A129528 is at least a lower bound of $d(n)$, by explicitly constructing $a(n)$ linearly independent vectors in the null-space? A good start is to examine the Bottom Schur functions defined in the paper, as they can be identified with elements in the null space of $R-D$.

论坛问题帖 | no_answers_at_retrieval | Per Alexandersson | 2019-03-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018220 | MO-326199

Open problems concerning all the finite groups

What are the open problems concerning all the finite groups? The references will be appreciated. Here are two examples: Aschbacher-Guralnick conjecture (AG1984 p.447): the number of conjugacy classes of maximal subgroups of a finite group is at most its class number (i.e. the number of conjugacy classes of elements, or the number of irreducible complex representations up to equiv.). K.S. Brown's problem (B2000 Q.4; SW2016 p.760): Let G be a finite group, μ be the Möbius function of its subgroup lattice $L(G)$. Then the sum $\sum_{H \in L(G)}\mu(H,G)|G:H|$ is nonzero. There are two types of problems, those involving an upper/lower bound (like Aschbacher-Guralnick conjecture) and those "exact", involving no bound (like K.S. Brown's problem). I guess the first type is much more abundant than the second, so for the first type, please restrict to the main problems.

论坛问题帖 | has_answers | Sebastien Palcoux | 2019-03-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018221 | MO-327040

Is the exact solution of the wave equation for the scattering of waves by a disk/cylinder an open problem?

The solution exact solution of the Helmholtz equation for the scattering of waves by a sphere is relatively straightforward and has been known since the time of Lord Rayleigh. The exact solution of the wave equation for scattering of waves by a sphere is much trickier and has been dealt with in two relatively recent papers - here and here. I have been trying to determine whether there exists an exact solution for the wave equation in the two-dimensional case, i.e. for a disk/cylinder. It is not mentioned in the recent papers I listed above for the case of the sphere, and I haven't been able to find anything despite an extensive search of the literature. Does anyone know if an exact solution of the wave equation for the scattering of waves by a disk/cylinder exists, or is this still an open problem?

论坛问题帖 | no_answers_at_retrieval | electroscience | 2019-04-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018222 | MO-327172

Seeking proof of the Cuckoo Cycle Conjecture

Allow me to introduce the Cuckoo Cycle Proof of Work, and conclude with a closely related Conjecture about random graphs, which I hope someone can find a proof for. Cuckoo Cycle is named after the Cuckoo Hashtable, in which each data item can be stored at two possible locations, one in each of two arrays. A hash function maps the pair of item and choice of array to its location in that array. When a newly stored item finds both its locations already occupied, one is kicked out and replaced by the new item. This forces the kicked-out item to

move to its alternate location, possibly kicking out yet another item. Problems arise if the corresponding Cuckoo graph, a bipartite graph with array locations as nodes and item location pairs as edges, has cycles. While n items, whose edges form a cycle, could barely be stored in the table, any $n+1$st item mapping within the same set of locations (a chord in the cycle) would not fit, as the pigeonhole principle tells us. In the Proof-of-Work problem, we typically set $n \ge 29$, and use edge indices (items) $0 \dots N-1$, where $N = 2^n$. The endpoints of edge i are (siphash(i|0) % N, siphash(i|1) % N), with siphash being a popular keyed hash function. A solution is a cycle of length L in this bipartite Cuckoo graph, where typically $L = 42$. Cuckoo Cycle solvers spend nearly all cycles on edge trimming; identifying and removing edges that end in a leaf node (of degree 1), as such edges cannot be part of a cycle. Trimming rounds alternate between the two node partitions. The fraction f_i of remaining edges after i trimming rounds (in the limit as N goes to infinity) appears to obey The Cuckoo Cycle Conjecture: $f_i = a_{i-1} * a_i$, where $a_{-1} = a_0 = 1$, and $a_{i+1} = 1 - e^{-a_i}$ f_i could equivalently be defined as the fraction of edges whose first endpoint is the middle of a (not necessarily simple) path of length $2i$. So far I have only been able to prove the conjecture for $i \le 3$. For instance, for i = 1, the probability that an edge endpoint is not the endpoint of any other edge is (1-1/N)N-1 ~ 1/e. Here's hoping someone finds an elegant proof...

论坛问题帖 | has_answers | John Tromp | 2019-04-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018223 | MO-32785

Cyl(E) = Borel(E) for E non-reflexive Grothendieck Banach space

This is sort of a follow-up to Borel(X) = \sigma(X') for X non-separable PROBLEM: Given a Banach space E over $\mathbb{K} \in \{\mathbb{C}, \mathbb{R}\}$ that has the Grothendieck property. Does $\hat{C}(E) = \mathcal{B}(E)$ imply E is reflexive? (This would in turn imply that E is separable). Some definitions: A Banach space is a Grothendieck space if a sequence in E' which is $\sigma(E', E)$-convergent is automatically $\sigma(E', E'')$-convergent. Or equivalently: every $\sigma(E', E)$ zero sequence has subsequence which is $\sigma(E', E'')$-convergent, or equivalently: every linear, bounded operator from E to c_0 (or any separable Banach space) is automatically weakly compact. The σ-algebra $\hat{C}(E)$ is the σ-algebra generated by sets of the form $\mathcal{C}_{u_1, \cdots, u_n; C} := \{x \in E : (u_1(x), \cdots, u_n(x)) \in C\}$ where $u_1, \cdots, u_n \in E'$, $C \in \mathcal{B}(\mathbb{K}^n)$ and $n \in \mathbb{N}$. Notes: The σ-algebra $\hat{C}(E)$ equals the σ-algebra of weak Baire sets $\mathcal{B}_0(E, \sigma(E, E'))$ for every locally convex space E (see [2], Theorem 2.3). The inclusion $\hat{C}(E) \subset \mathcal{B}(E)$ is trivially true. If E is separable then $\hat{C}(E) = \mathcal{B}(E)$. [To see this use the Hahn-Banach theorem to show that $\mathcal{B}_E \in \hat{C}(E)$. As translations and scalar multplications are measurable with regard to the cylindrical σ-algebra the other inclusion follows.] A reflexive space is automatically Grothendieck. For a separable Grothendieck space E we have that the identity is weakly compact so E becomes reflexive A reflexive space E with $\hat{C}(E) = \mathcal{B}(E)$ is automatically separable ([1], Prop. 2.6, p.19). Without reflexivity the equality $\hat{C}(E) = \mathcal{B}(E)$ does not imply E is separable in general. The example $E = \ell^2(\mathbb{R})$ shows that there is a reflexive and non-separable space with $\mathcal{B}(E) \not= \hat{C}(E)$ Edgar's example below or $E = \ell^{\infty} = C(\beta \mathbb{N})$ gives a non-reflexive Grothendieck space with $\mathcal{B}(E) \not= \hat{C}(E)$. The question therefore: does every non-reflexive Grothendieck space have that property? There are non-reflexive Grothendieck spaces which do not contain ℓ^{∞} (cf. [3]). So we can't simply reduce to this case. I don't know much more about Grothendieck spaces though or characterizations of them that might be helpful. REFERENCES: [1] N. N. VAKHANIA, V. I. TARIELADZE, S. A. CHOBANYAN, Probability Distributions on Banach Spaces, Mathematics and its applications (D. Reidel Publishing Company), 1987 [2] http://www.iumj.indiana.edu/IUMJ/FULLTEXT/1977/26/26053 [3] R. HAYDON, A non-reflexive Grothendieck Space that does not contain ℓ^{∞}, Israel Journal of Mathematics, Vol 40, No. 1, 1981 EDIT: I rephrased the question and added some information.

论坛问题帖 | has_answers | santker heboln | 2010-07-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018224 | MO-330350

Coloring a graph formed by cliques sharing at most one point

Consider a graph formed by k k order cliques sharing at most one point. Consider thedegenerate case of all cliques disjoint, which is trivially k colorable. Now, to colour any other such graph, we first color any one clique. Now, if any vertex of the colored clique is shared by m cliques, then, we could put at maximum $k-(m-1)$ vertices to expand the color class containing that vertex, which are chosen one from each clique not sharing that vertex. Such a vertex exists, by construction(as otherwise, other cliques would be sharing that vertex) . This corresponds to the number of reduced vertices from k^2 vertices, which is the degenerate case of all cliques being disjoint. Proceeding so on for all vertices of the colored clique, I think we can cover all the vertices of the graph by expanding the color classes to the required deficiency. Thus, the graph as a whole would be $k-$ colorable. Is the above assertion right? Or are there any counterexamples? Any light on this. Thanks beforehand.

论坛问题帖 | no_answers_at_retrieval | vidyarthi | 2019-04-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018225 | MO-3332

Two commuting mappings in the disk

Suppose that f and g are two commuting continuous mappings from the closed unit disk (or, if you prefer, the closed unit ball in R^n) to itself. Does there always exist a point x such that $f(x)=g(x)$? If one of the mappings is invertible, then it is just a restatement of Brouwer's fixed point theorem but I do not know the answer in the general case and would not even dare to guess what it must be. Also, the answer is well-known to be "Yes" in dimension 1.

论坛问题帖 | has_answers | fedja | 2009-10-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018226 | MO-333425

Hilbert 16th problem and dynamical Lefschetz trace formula

I would like to apply the known version of the conjectural formula (11) page 10 of the paper Number theory and dynamical Lefschetz trace formula. Disclaimer: I do not have a complete understanding of this formula but I can get an sketch of it. I just know that both sides of the formula are not number but distribution. I also know the meaning of ingredient of the formula. I wish to apply it to find an appropriate application in limit cycle theory. To start this way, I have three precise questions: Question 1: In the right hand side of the formula (11), is not necessary to assume that there are only a finite number of non degenerate periodic orbits γ(As they are appeared in the sum $\sum$ of the right side of the formula)? Is it implicitly included in the assumptions of that formula? I think that even the non degeneracy assumption of periodic orbits does not easily imply the finite ness of such periodic orbits. Is not possible that a sequence of non degenerate periodic orbits accumulate to a non periodic orbit which is a kind of complicated and strange attractor? Question 2:(This question is completely different from the previous one but there are some motivations from this posts And also this post Lifting a Quadratic System to a non Vanishing vector field on S^3) A polynomial vector field on the plane gives us an analytic vector field X on S^2. Put $\tilde{X}$ for the obvious lifting of X on $S^2\times S^1$ with $\tilde{X}=X+\partial/\partial{\theta}$ Is there a quadratic polynomial vector field on $\mathbb{R}^2$ with Poincare compactification X such that $\tilde{X}$ on $S^2\times S^1$ does not admit a 2 dimensional transversal foliation which is invariant under the flow of $\tilde{X}$. The reason we consider Quadratic system: Note that every quadratic system is a geodesible vector field but it is not the case for higher degree polynomial vector field. On the other hand, in dimension 2, if a vector field X is geodesible then there is a transversal field Y with $[X,Y] \parallel Y$ this implies that orbits of Y are invariant under flow of X. So this make us to be hopeful a little that the product vector field $\tilde{X}=X+\partial/\partial \theta$ admit a transversal 2 dimensional foliation which is invariant under the flow of $\tilde{X}$. Existence of such transversal foliation is the key condition in the paper we linked in the first lines of this post. The reason we consider $S^2\times S^1$ rather than S^3: The lifting of the simplest vector field $X=0$ to the Hopf vector field on S^3 does not admit a transversal foliation. Question 3: When we lift a vector field X to a non vanishing vector field $\tilde{X}$ on $S^3, S^2\times S^1\quad\text{or}\quad T^1 S^2$, it is possible that the preimage of a closed orbit, which is an invariant torus, would not contain any closed orbit, so we loose our closed orbits.please see the comment by Sebastian Goette in this post. With terminologies in the linked paper, let we have a 3 manifold foliated by 2 dimensional leaves whch is compatible with a flow X. So is there an analogy of the formula (11) in page 10 of the linked paper in the first lines of this post whose right side depends on invariant torus of X as well as closed orbits of X? Remark: The concept "Blue Sky catastrophe" and difficulties occured in "Finitness theorem for Limit Cycles" are motivations for the first question of this post.

论坛问题帖 | no_answers_at_retrieval | Ali Taghavi | 2019-06-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018227 | MO-33396

Razborov's response to Almost Natural Proofs

This post is about Natural Proofs barrier in computational complexity. There are two recent papers related to this. They are: Amplifying lower bounds by means of self-reducibility by Eric Allender and Michal Koucký, Almost-natural proofs by Timothy Chow. Alexander Razborov mentioned them at the end of his talk in the first Barriers Workshop at IAS in 2009. The video of his talk is available here. He mentioned that a possible way to respond to these would be to make the largeness condition relative. So here is my question: Has there been any progress on this?

论坛问题帖 | no_answers_at_retrieval | Kaveh | 2010-07-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018228 | MO-334188

An triangle inequality $\sum_{i=1}^n b_i^\alpha \ge \sum_{i=1}^na_i^\alpha $ if $\alpha >1$

Using my computer I discovered that: if a,b,c are sidelengths of a triangle, then $(a+b-c)^\alpha+(b+c-a)^\alpha+(c+a-b)^\alpha \ge a^\alpha+b^\alpha+c^\alpha $ if $\alpha >1$ $(a+b-c)^\alpha+(b+c-a)^\alpha+(c+a-b)^\alpha \le a^\alpha+b^\alpha+c^\alpha $ if $0< \alpha <1$ My question: The generalization inequality as follow is corect? Let and $\alpha>0$, $a_i>0$ for $i=\overline{1,n}$ and $S=a_1+a_2+....+a_n$. Let $b_i=S-(n-1)a_i \ge 0$ for $i=\overline{1,n}$, then $\sum_{i=1}^n b_i^\alpha \ge \sum_{i=1}^na_i^\alpha $ if $\alpha >1$ $\sum_{i=1}^n b_i^\alpha \le \sum_{i=1}^na_i^\alpha $ if $0< \alpha <1$

论坛问题帖 | has_answers | Đào Thanh Oai | 2019-06-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018229 | MO-334951

Hyperbolic groups and spaces of negative curvature

Mikhail Gromov states that he "tried for about 10 years to prove that every hyperbolic group is realizable by a space of negative curvature" in his interview with Martin Raussen and Christian Skau (page 6, link below). Is there any information available about the current status of this conjecture?
https://www.ihes.fr/~gromov/wp-content/uploads/2018/08/rtx100300391p.pdf

论坛问题帖 | has_answers | spiramirabilis | 2019-06-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018230 | MO-335820

A cohomology associated to a (not necessarily integrable) distribution (Hilbert 16th problem and dynamical Lefschetz trace formula 2)

Let (M,D) be a pair consisting of a manifold M and a distribution D on M. The de Rham complex $\Omega^*(M)$ has the following subcomplex $$\Omega^*(M,D)=\{ \alpha \in \Omega^*(M)\mid \alpha_{| D}=d\alpha_{|D}=0\}.$$ When D is an integrable

distribution, the last condition $d\alpha_{|D}=0$ is redundant. The quotient complex $\Omega^*(M)/\Omega^*(M,D)$ produces a cohomology. In this complex we can replace the space of coboundaries(exact forms along distribution) with its smooth topology closure. The reason for closure is explained in this paper which is linked in the MO post in the next paragraph. So we get a cohomology $H^*(M,D)$, the cohomology of M along distribution D. Now assume that the flow of a vector field $\tilde{X}$ on M preserves D. Is the dynamical Lefschetz trace formula discussed in "Hilbert 16th problem and dynamical Lefschetz trace formula" still valid? Motivation: Theorem 9.3 page 29 of the following paper says that if a vector field on S^3 is transversal to a codimension one foliation, then the vector field must have a parameter family of periodic orbits. On the other hand one of the main concerns of the above MO post is finiteness of closed orbits for a vector field on S^3 which is a result of lifting of a (generic) vector field on 2-sphere. So we should almost give up finding a foliation transversal to our vector field. Instead of a foliation, we search a (non integrable) transversal distribution which is preserved by the flow of our vector field. So we need an appropriate analogy of leafwise de Rham cohomology when there is not any leaf, that is a non-integrable distribution. The above cohomology we introduce is a candidate for such appropriate cohomology. So we hope that the lifting $\tilde{X}$ (to S^3 or $S^2\times S^1$) of a Poincare compactification of a quadratic plannar vector field would be a geodesible vector field. Then the orthogonal direction D(with respect to an appropriate metric adapted to $\tilde{X}$) is obviously preserved by flow of $\tilde{X}$. (For more explaination on the later paragraph please see [this post](Finding a 1-form adapted to a smooth flow) So we try to investigate the dynamics of $\tilde{X}$ via a possible distribution analogy of Dynamical trace formula). Precise and detailed motivations are mentioned in the above linked MO post entitled "Hilbert 16th problem and dynamical Lefschetz trace formula".

论坛问题帖 | no_answers_at_retrieval | Ali Taghavi | 2019-07-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018231 | MO-335960

Have new conjectures generated by the Ramanujan machine been proven?

Recently the following preprint was published with new automatically generated conjectures on generalizes continuous fractions, e.g., for the Euler constant: Raayoni, Pisha, Manor, Mendlovic, Haviv, Hadad, and Kaminer - The Ramanujan machine: Automatically generated conjectures on fundamental constants. Have these conjectures been proven in the meantime? Are there any partial results?

论坛问题帖 | has_answers | Andreas Rüdinger | 2019-07-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018232 | MO-337954

How many solutions of x^3 +y^3 = z^3+3 are known?

I've been told that there is reason to think that the equation $x^3 + y^3 = z^3 + 3$ has solutions in positive integers other than $$4^3 + 4^3 = 5^3 +3.$$ Can someone tell me the current status of this problem? Is there another solution known? Here's a reference to the problem in greater generality. MATHEMATICS OF COMPUTATION Volume 66, Number 218, April 1997, Pages 841-851 S 0025-5718(97)00830-2 ON SEARCHING FOR SOLUTIONS OF THE DIOPHANTINE EQUATION $x^3 + y^3 + z^3 = n$

论坛问题帖 | no_answers_at_retrieval | David S. Newman | 2019-08-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018233 | MO-33796

Existence of a pseudo-polynomial time algorithm for a counting problem.

Let T={1,...,n} be a set of tasks. Each task i has associated a non negative processing time p_i and a deadline d_i. A feasible schedule of the tasks consists of a permutation of n elements pi, such that \sum_i=1^k p_(pi(i)) <= d_(pi(i)) for all k=1,...n. Does there exists a pseudo-polynomial time algorithm for computing the total number of feasible schedules? A pseudo-polynomial time algorithm is an algorithm whose running time is bounded by a polynomial on the size of the input, given that the input is written in unary notation (2=II, 3 =III). (e.g., the size of a number n in unary notation is O(n), and not O(log(n)). This is an open question from an article published in 2009 at Operations Research Letters.

论坛问题帖 | has_answers | Gerardo | 2010-07-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018234 | MO-33817

Work on independence of π and e

It is an open problem to prove that π and e are algebraically independent over $\mathbb{Q}$. What are some of the important results leading toward proving this? What are the most promising theories and approaches for this problem?

论坛问题帖 | has_answers | muad | 2010-07-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018235 | MO-338925

The Total Graph is similar to a line graph

Consider the total graph of a regular graph. From the structure, it seems that it has a similar structure to the line graph (two different sub-cliques joining at a single point) except that, in addition, there is an edge between two different sub-cliques. Note that the sub-cliques in the total graph are the just the sub-cliques which are present in the line graph of the graph with a vertex added to each sub-clique make it adjacent to all the vertices of a particular sub-clique, thereby increasing the order of the sub-cliques by one. This means that the clique number of the total graph of a graph is one more than the clique number of the line graph of the same graph. This leads me to infer that the total graph of a graph has similar chromatic properties as a line graph of a graph. Specifically, I expect the total graph of a graph to satisfy a similar bound to Vizing's theorem for line graphs. Will this be true? Thanks beforehand.

论坛问题帖 | has_answers | vidyarthi | 2019-08-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018236 | MO-339030

Is the conjecture $min(A,B) \le rad(ABC)$ new and correct?

$\DeclareMathOperator\rad{rad}$Conjecture: If A, B, C are positive integers with $\gcd(A, B)=1$, $\gcd(B, C)=1$, and $\gcd(C, A)=1$, and if $A+B=C$, then $\min(A,B) \le \rad(ABC)$. If the conjecture is valid, then we can use the conjecture to prove the Fermat last theorem as follows: Proof of the Fermat last theorem: We consider the Fermat equation: $x^n+y^n=z^n$ with $\gcd(x, y)=1$, $\gcd(y, z)=1$, and $\gcd(z, x)=1$. There is no loss of generality in assuming that $x \le y <z$. By the conjecture, we get $x^n \le \rad(x^n y^n z^n)=\rad(x y z)\le x y z $. So $x^n+y^n \le xyz+y^n<z^3+(z-1)^n$. But we can easily prove that $z^3+(z-1)^n < z^n$ whenever $n > 3$ and $z>1$. So now we only need to prove the Fermat last theorem with $n=3$. My question: Is the conjecture above new and correct?

论坛问题帖 | has_answers | Đào Thanh Oai | 2019-08-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018237 | MO-341313

Which knot invariants have no known diagram-independent descriptions?

Many knot invariants in knot theory are discovered by finding a property of knot diagrams which is invariant under the three Reidemeister moves. Now in principle, any knot invariant can be described in a diagram-independent way, that is, as a property of the three-dimensional knot itself without reference to diagrams of the knot. But in practice, it can take years between the development of a knot invariant and the discovery of a diagram-independent description of it. So my question is, for what knot invariants is a diagram-independent description not yet known?

论坛问题帖 | has_answers | Keshav Srinivasan | 2019-09-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018238 | MO-341362

What is the state of research on finding all prime knots with 17 crossings?

In this 1998 journal paper, all the prime knots with 16 or fewer crossings are found (some of which were found earlier by others). There are over 1.7 million such knots. But the prime knots with 17 crossings have not yet been tabulated. Here is what this book says: This is probably hard and requires new ideas. But this book was written in 2004, so things may have changed since then. There have certainly been a lot of developments in knot theory over the past 15 years. So my question is, what is the state of research on finding all prime knots with 17 crossings? Are we relatively close to doing so, and have partial results been discovered?

论坛问题帖 | has_answers | Keshav Srinivasan | 2019-09-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018239 | MO-34145

Can we cover the unit square by these rectangles?

The following question was a research exercise (i.e. an open problem) in R. Graham, D.E. Knuth, and O. Patashnik, "Concrete Mathematics", 1988, chapter 1. It is easy to show that $$\sum_{1 \leq k } \left(\frac{1}{k} \times \frac{1}{k+1}\right) = \sum_{1 \leq k } \left(\frac{1}{k} - \frac{1}{k+1}\right) = 1.$$ The product $\frac{1}{k} \times \frac{1}{k+1}$ is equal to the area of a $\frac{1}{k}$by$\frac{1}{k+1}$ rectangle. The sum of the areas of these rectangles is equal to 1, which is the area of a unit square. Can we use these rectangles to cover a unit square? Is this problem still open? What are the best results we know about this problem (or its relaxations)?

论坛问题帖 | has_answers | Kaveh | 2010-08-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018240 | MO-345388

Decision problems for which it is unknown whether they are decidable

In computability theory, what are examples of decision problems of which it is not known whether they are decidable?

论坛问题帖 | has_answers | Dominic van der Zypen | 2019-11-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018241 | MO-345680

What's an example of an ∞-topos not equivalent to sheaves on a Grothendieck site?

My question is as in the title: Does anyone have an example (supposing one exists) of an ∞-topos which is known not to be equivalent to sheaves on a Grothendieck site? An ∞-topos is as in Higher Topos Theory (HTT) 6.1.0.4: an ∞-category which is an accessible left-exact localization of presheaves on a small ∞-category. A Grothendieck site is a small ∞-category $\mathcal{C}$ equipped with the ∞-categorical variant of the classical notion of a Grothendieck topology $\mathcal{T}$, as in HTT 6.2.2: a collection of sieves (subobjects $U\to j(C)$ of representable presheaves on $\mathcal{C}$) satisfying some axioms. Sheaves on $(\mathcal{C},\mathcal{T})$ are presheaves of ∞-groupoids on $\mathcal{C}$ which are local for the sieves in $\mathcal{T}$. Such form a full subcategory $\mathrm{Shv}(\mathcal{C},\mathcal{T})$ of the ∞-category of presheaves. Note: the question Examples of $(\infty,1)$-topoi that are not given as sheaves on a Grothendieck topology appears superficially to be equivalent to this one. In practice it is not exactly the same. As answers to that question show, many interesting ∞-topoi exist which can be described without reference to any Grothendieck site. But it is still conceivable that a suitable site exists. Also note: any ∞-topos $\mathcal{X}$ can be obtained as an accessible left-exact localization of some $\mathrm{Shv}(\mathcal{C},\mathcal{T})$ with respect to a suitable class of ∞-connected morphisms (HTT 6.2.2, 6.5.3.14), e.g., the class of hypercovers. However, this does not immediately preclude $\mathcal{X}$ being equivalent to $\mathrm{Shv}(\mathcal{C}',\mathcal{T}')$ for some other Grothendieck site

$(\mathcal{C}',\mathcal{T}')$. Added remark. I asked this question because I had, for a long time, tacitly assumed that such examples were plentiful, until I thought about it and realized I had no basis for thinking that. As no answers have yet been given, and I'm not aware of any tools which would likely lead to a resolution one way or the other, it looks to me that this should be regarded as an open question.

论坛问题帖 | has_answers | Charles Rezk | 2019-11-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018242 | MO-345913

Is it an open problem whether fast-growing hierarchies can be defined without fundamental sequences?

Googology Wiki says this, concerning the relation between fast-growing hierarchies defined for all countable ordinals, and the existence of a system of assigning a canonical fundamental sequence to each countable limit ordinal: It is an open problem whether an ordinal hierarchy can be defined without using fundamental sequences. More explicitly, the problem concerns whether there is a model of ZF such that there exists an $F : \omega_1 \rightarrow (\mathbb{N} \rightarrow \mathbb{N})$ where for all $\alpha > \beta, F(\alpha)$ eventually outgrows $F(\beta)$, but there does not exist an $S: \omega_1 \cap \text{Lim} \rightarrow (\mathbb{N} \rightarrow \omega_1)$ such that for all $\alpha, \sup(R) = \alpha$ where R is the range of $S(\alpha)$. My question is, is it true that the question of whether there exists such a model of ZF is an open problem?

论坛问题帖 | no_answers_at_retrieval | Keshav Srinivasan | 2019-11-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018243 | MO-34651

Permute Wada Lakes keeping the coastline intact? (still open in dim >2)

Wada Lakes are three disjoint open subsets of $\mathbb R^2$ with common boundary. Originally they were constructed by hand, but they also arise naturally in the real life, that is, theory of dynamical systems. (See, for example, this http://www.math.cornell.edu/~hubbard/pendulum.pdf paper of John Hubbard.) From dynamical construction it is clear that one can have a homeomorphism that permutes the lakes. For example, this expository paper http://www.ams.org/notices/200601/fea-coudene.pdf gives such a homeomorphism (in fact, a diffeomorphism) for lakes on the sphere S^2. The dynamics on the boundary of the lakes is chaotic. == Question 1: Let U_1, U_2 and U_3 be open subset of $\mathbb R^d$ with common boundary C. Assume that each U_i is an image of an injective continuous map $\mathbb R^d\to\mathbb R^d$. Is there a homeomorphism of $\mathbb R^d$ that permutes open sets U_i and is identity on C? Question 2: Let $U_i\subset \mathbb R^d$, $i\in \mathbb Z$, be a disjoint collection of open sets each of which is homeomorphic to $\mathbb R^d$. Is there a homeomorphism of $\mathbb R^d$ that satisfies the following h maps U_i onto U_{i+1} every point in ∂U_i is periodic under h == Edit: Below André Henriques has produced two examples when such a homeomorphism does not exist. Both examples make use of some local structure that cannot be permuted. It is still not clear what happens for "abstact Wada Lakes": are these examples exceptions or arbitrary Wada Lakes cannot be permuted as well? Edit2 (Jan. 9th 2012) In is clear now that there is no such homeomorphism if the dimension $d=2$. (See the comments below). The answer to Q1 is also negative in higher dimension as explained in the reference given by Andres Koropecki. For $d\ge 3$ there are examples by André Henriques of open sets for which one cannot find such a homeomorphism. In general, Question 2 is still open.

论坛问题帖 | has_answers | Andrey Gogolev | 2010-08-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018244 | MO-3468

Do torsion-free groups give projectionless group ($C^\ast$) algebras?

One of the reasons I study von Neumann algebras is that they always have plenty of projections. There are many projectionless $C^\ast$-algebras (0 and possibly 1 are the only projections), but the von Neumann algebras they generate must have nontrivial projections (unless it's just the complex numbers, of course). A good example of this is the reduced group $C^\ast$-algebra of any free group F_n. If $n=1$, then $C_r^\ast(Z)\cong C(S^1)$ via the Gelfand transform, which is clearly projectionless. If $n\geq 2$, the proof is fairly complicated. See Davidson's book for a proof when $n=2$. If G is a torsion-free group, is the reduced group $C^\ast$-algebra of G projectionless? This $C^\ast$-algebra always contains the group algebra $C[G]$, so a simpler question is whether $C[G]$ is projectionless if G is torsion-free. Note that torsion-free is a necessary condition as one gets a projection from summing up the elements in the cyclic group generated by a torsion element and dividing by the order of the element. EDIT: changed typestting. still some bugs... help please?

论坛问题帖 | has_answers | Dave Penneys | 2009-10-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018245 | MO-346842

Is every finite-order unimodular matrix conjugate to a $0,1,-1$ matrix?

Problem. Given a matrix $A\in\mathrm{GL}(n,\mathbb{Z})$ such that $A^k=1$ for some $k\geq 1$, is there a matrix $g\in\mathrm{GL}(n,\mathbb{Z})$ such that gAg^{-1} has only 0, 1, and -1 as possible entries? Edit: after the remark by Mark Sapir that it is a famous open problem (which for me was already sufficient as an answer), I changed my question into the following ones, so now maybe it is more suitable for staying on MO without being closed. What is known about this problem? Which other parts of mathematics is it connected to?

论坛问题帖 | has_answers | Qfwfq | 2019-11-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018246 | MO-348055

When do two knots have isomorphic fundamental bikeis?

A kei, also known as an involutive (or involutory) quandle, is a quandle $(Q,*)$ satisfying the involution condition that $(x*y)*y=x$ for all x and y. Just like we can define a fundamental quandle of a knot using arcs of an oriented knot diagram, we can define the fundamental kei of a knot using arcs of an unoriented knot diagram. A bikei, also known as an involutive biquandle, is a biquandle satisfying similar involution conditions. Just as we can define the fundamental biquandle of a knot using semiarcs of an oriented knot diagram, we can define the fundamental bikei of a knot using the semiarcs of an unoriented knot diagram. (These are all defined precisely in Elhamdadi and Nelson's book Quandles: An Introduction to the Algebra of Knots.) My question is, under what circumstances are the fundamental bikeis of two knots isomorphic? Now this thesis Martina Vaváčková, Algebraic Structures for Knot Coloring, Masters thesis, Charles University (2018) (abstract page, pdf) shows that the fundamental keis of the granny knot and square knot are isomorphic, whereas their fundamental quandles are not isomorphic. The recent journal paper Katsumi Ishikawa, Knot Quandles vs. Knot Biquandles, International Journal of Mathematics (accepted to appear) doi:10.1142/S0129167X20500159, on the other hand, shows that the fundamental biquandles of two knots are isomorphic if and only if their fundamental quandles are isomorphic. So this raises two natural questions: Can two knots have isomorphic fundamental bikeis without having isomorphic fundamental biquandles? And can two knots have isomorphic fundamental keis without having isomorphic fundamental bikeis? Note that the answers to both can't be no since isomorphic fundamental keis don't imply isomorphic fundamental quandles. (The converses are clear: isomorphic biquandles imply isomorphic bikeis, and isomorphic bikeis imply isomorphic keis.) Do the two results I mentioned shed any light on these questions? Are either of these questions open problems?

论坛问题帖 | no_answers_at_retrieval | Keshav Srinivasan | 2019-12-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018247 | MO-34909

Conjugacy classes and reduced group C^*-algebra of an amenable group

The reduced C^*-algebra of a non-abelian free group G has a unique trace. Hence, there is no chance to separate conjugacy classes of group elements using traces on $C^\star_{red} G$. On the other side, for the group ${\mathbb Z}$, separation is clearly possible. Question: Let G be an amenable group. Does the reduced group $C^\star$-algebra of G support sufficiently many traces to distinguish between conjugacy classes of group elements? EDIT: The question seems already interesting for $S_{\infty} = \cup_n S_n$. Let's get explicit and pick $g \in S_n$ (for some n) and consider the canonical trace $\tau_{g,n}$ which sends every conjugate of g to 1 and all other elements to zero. (This can be done for any $n' \geq n$ since $S_n \subset S_{n'}$.) The function $\tau_{g,n} \colon {\mathbb C}S_n \to {\mathbb C}$ is a conjugation invariant function and hence, it must be a linear combination of the normalized traces of irreducible representations of S_n. Question: What is the sum of the absolute values of the coefficients that come up in this linear combination of traces? This is (as one can check) the norm of $\tau_{g,n}$, call it $c(g,n)$. So, we see that the compatible family of maps $\tau_{g,n}$ extends from ${\mathbb C}S_{\infty}$ to $C^* S_{\infty}$ if and only if $c(g,n)$ remains bounded.

论坛问题帖 | has_answers | Andreas Thom | 2010-08-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018248 | MO-34980

How can an approach to P vs NP based on descriptive complexity avoid being a natural proof in the sense of Raborov-Rudich?

EDIT: This question has been modified to make it a stand-alone question. Feel free to retract your votes for the previous version. Here are Vinay Deolalikar's paper, and Richard Lipton's first post about it, and the wiki page on polymath site summarizing the discussions about it. His approach is based on descriptive complexity. One of famous barriers for separating NP from P is Razborov-Rudich Natural Proofs barrier. Richard Lipton remarked about his paper and the natural proofs barrier that apparently "it exploits a uniform characterization of P that may not extend to give lower bounds against circuits". A question which is mentioned in one of the comments on Lipton's post is: How essential is the uniformity of P to his proof? i.e is the uniformity of P used in such an essential way that the barrier will not apply to it? (By essential I mean that the proof does not work for the non-uniform version.) So here is my questions: Are there any previous computational complexity results based on descriptive complexity that avoid the Razborov-Rudich natural proofs barrier (because of being based on descriptive complexity)? How can an approach to P vs NP based on descriptive complexity avoid being a natural proof in the sense of Raborov-Rudich? A related question is: What are the complexity results using uniformity in an essential way other than proofs by diagonalization? Related closed MO posts: When would you read a paper claiming to have settled a long open problem like P vs. NP? https://mathoverflow.net/questions/34953/whats-wrong-with-this-proof-closed Discussion on meta: http://mathoverflow.tqft.net/discussion/590/whats-wrong-with-this-proof/

论坛问题帖 | no_answers_at_retrieval | Kaveh | 2010-08-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018249 | MO-35151

What impact would P!=NP have on the characterization of BQP?

Many complexity theorists assume that $P\ne NP.$ If this is proved, how would it impact quantum computing and quantum algorithms? Would the proof immediately disallow quantum algorithms from ever solving NP-Complete problems in Quantum Polynomial time? According to Wikipedia, quantum complexity classes BQP and QMA are the bounded-error quantum analogues of P and NP. Is it likely that a proof that $P\ne NP$ can be adapted to the quantum setting to show that $BQP \ne QMA?$

论坛问题帖 | has_answers | user8347 | 2010-08-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018250 | MO-351688

Conjectured primality test for specific class of $N=4kp^n+1$

Can you provide a proof or counterexample for the following claim? Let $P_m(x)=2^{-m}\cdot((x-\sqrt{x^2-4})^m+(x+\sqrt{x^2-4})^m)$. Let $N= 4kp^{n}+1 $ where k is a positive natural number , $ 4k<2^n$, p is a prime number and $n\ge3$.

Let a be a natural number greater than two such that $\left(\frac{a-2}{N}\right)=-1$ and $\left(\frac{a+2}{N}\right)=-1$ where $\left(\frac{}{}\right)$ denotes Jacobi symbol. Let $S_i=P_p(S_{i-1})$ with S_0 equal to the modular $P_{kp^2}(a) \text{mod} N$. Then N is prime if and only if $S_{n-2} \equiv 0 \pmod{N}$. You can run this test here. I have verified this claim for $k \in [1,500]$ with $p \leq 97$ and $n \in [3,50]$. Further generalization of the claim A Let $P_m(x)=2^{-m}\cdot((x-\sqrt{x^2-4})^m+(x+\sqrt{x^2-4})^m)$. Let $N= 2kp^{n} + 1 $ where k is a positive natural number , $ 2k<2^n$, p is a prime number and $n\ge3$. Let a be a natural number greater than two such that $\left(\frac{a-2}{N}\right)=-1$ and $\left(\frac{a+2}{N}\right)=-1$ where $\left(\frac{}{}\right)$ denotes Jacobi symbol. Let $S_i=P_p(S_{i-1})$ with S_0 equal to the modular $P_{kp^2}(a) \text{mod} N$. Then N is prime if and only if $S_{n-2} \equiv -2 \pmod{N}$. You can run this test here. B Let $P_m(x)=2^{-m}\cdot((x-\sqrt{x^2-4})^m+(x+\sqrt{x^2-4})^m)$. Let $N= 2kp^{n} - 1 $ where k is a positive natural number , $ 2k<2^n$, p is a prime number and $n\ge3$. Let a be a natural number greater than two such that $\left(\frac{a-2}{N}\right)=1$ and $\left(\frac{a+2}{N}\right)=-1$ where $\left(\frac{}{}\right)$ denotes Jacobi symbol. Let $S_i=P_p(S_{i-1})$ with S_0 equal to the modular $P_{kp^2}(a) \text{mod} N$. Then N is prime if and only if $S_{n-2} \equiv -2 \pmod{N}$. You can run this test here.

论坛问题帖 | has_answers | Pedja | 2020-02-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018251 | MO-35207

Finite-dimensional subalgebras of $C^\star$-algebras

Let A be a unital $C^\star$-algebra and let $a_1,\dots,a_n$ be a finite list of normal elements in A which (together with their adjoints) generate a norm-dense $\star$-subalgebra $B \subset A$. Clearly, if A is finite-dimensional, then every element in A (and hence B) has finite spectrum. I am asking for the converse. Question: Assume that each element of B has finite spectrum. Is it true that A has to be finite-dimensional? The existence of finitely-generated infinite torsion groups shows that this might be a highly non-trivial problem. In this case one would consider the reduced group $C^\star$-algebra and note that all monomials in the generators of the group and their inverses (which are equal to the adjoints of the generators) would have finite spectrum. However, the generated algebra would still be infinite-dimensional. I do not know of any simpler way to come up with such an example. In this case it is conceivable that the random-walk operator associated with the generating set (which is an element in the real group ring) has infinite spectrum, even though I did not prove this. Maybe there is also need to consider the spectra of elements in matrices over B (which of course follow to be finite if A is finite-dimensional.) In view of this, I am not only asking for an answer to the question but also for the right question (or a better one) if the answer to the original question is negative. A stronger assumption would be to assume that A itself consists only of elements with finite spectrum. This case seems much easier to approach and the answer seems to be positive. In fact every infinite-dimensional $C^\star$-algebra should contain an element with infinite spectrum. Just to get started, a more concrete instance of the question above is: Question: Let p_1, p_2 and p_3 be three projections in a C^*-algebra with the property that every (non-commutative) polynomial in p_1, p_2 and p_3 has finite spectrum. Is it true that the projections generate a finite-dimensional algebra?

论坛问题帖 | no_answers_at_retrieval | Andreas Thom | 2010-08-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018252 | MO-353150

Schur positive expression involving border-strip tableaux

Recall the power-sum expansion of Schur functions, $$ s_\lambda = \sum_\mu \chi^{\lambda}(\mu) \frac{p_\mu}{z_\mu}, $$ in terms of Sn-character. These can be calculated by the Murnaghan-Nakayama rule, stating that characters are a signed sum over border-strip tableaux: $$ \chi^{\lambda}(\mu) = \sum_{T \in BST(\lambda,\mu)} (-1)^{ht(T)} . $$ Now, what if we instead consider the 'cancellation-free' expression (yes, the omission of z_μ is intended): $$ CF_\lambda := \sum_\mu |BST(\lambda,\mu)| p_\mu. $$ Is the symmetric function CF_λ also Schur-positive? I have verified that this is the case for all $|\lambda| \leq 8$. This is the data for all λ with at most size 4. 1->ss[{1}] 2->2 ss[{2}] 11->2 ss[{2}] 3->3 ss[{3}]+ss[{2,1}]+ss[{1,1,1}] 21->5 ss[{3}]+3 ss[{2,1}]+ss[{1,1,1}] 111->3 ss[{3}]+ss[{2,1}]+ss[{1,1,1}] 4->5 ss[{4}]+3 ss[{2,2}]+2 ss[{3,1}]+2 ss[{2,1,1}]+ss[{1,1,1,1}] 31->10 ss[{4}]+6 ss[{2,2}]+10 ss[{3,1}]+6 ss[{2,1,1}]+2 ss[{1,1,1,1}] 22->7 ss[{4}]+7 ss[{2,2}]+6 ss[{3,1}]+2 ss[{2,1,1}]+3 ss[{1,1,1,1}] 211->10 ss[{4}]+6 ss[{2,2}]+10 ss[{3,1}]+6 ss[{2,1,1}]+2 ss[{1,1,1,1}] 1111->5 ss[{4}]+3 ss[{2,2}]+2 ss[{3,1}]+2 ss[{2,1,1}]+ss[{1,1,1,1}]

论坛问题帖 | no_answers_at_retrieval | Per Alexandersson | 2020-02-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018253 | MO-355009

About semiprimal representations of 1

Conjecture A_1 : For every $m \in \mathbb N \setminus \{1\}$ there exist mutually different primes $p_{r_1},...,p_{r_m}$ and mutually different primes $b_{w_1},...,b_{w_m}$ and numbers $i_1,...,i_m \in \{1,2\}$ such that $$\displaystyle \sum_{k=1}^m (-1)^{^{i_k}}b_{w_k}p_{r_k}=1$$ Is this known to be true? If not known to be true, I conjecture it is true. According to your knowledge of the subject, is the number of representations infinite for all $m\geq 2$? This I would like to formulate as: Conjecture B : The number of representations is infinite for all $m \geq 2$ And, with the help and computational efforts of Peter it seems that the conjecture A_1 can be strengthened to: Conjecture A_2: For every $m \in \mathbb N \setminus \{1\}$ there exist mutually different primes $p_{r_1},...,p_{r_m}$ and mutually different primes $b_{w_1},...,b_{w_m}$ such that $$\displaystyle p_{r_1}b_{w_1-} \sum_{k=2}^m b_{w_k}p_{r_k}=1$$ Some observations and computations even suggest that in conjecture A_1 and A_2 there are examples where all the mutually different primes $p_{r_1},...,p_{r_m}$ also can be mutually different from all the mutually different primes $b_{w_1},...,b_{w_m}$. What is already known about these and similar problems?

论坛问题帖 | has_answers | user153451 | 2020-03-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018254 | MO-355567

Isometries on the unit sphere

Suppose that X and Y are two Banach spaces, S_{X} and S_{Y} their unit spheres, and f an onto isometry between S_X and S_Y. Does it follow that X and Y are isometric?

论坛问题帖 | no_answers_at_retrieval | Markus | 2020-03-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018255 | MO-358940

Effective radius of section of a convex set compared to that of the convex itself

The effective radius $er(A)$ of a n-solid A, is defined by Schramm (see the question by Gil Kalai Volumes of Sets of Constant Width in High Dimensions) to be the radius of the n-ball that has the same volume. Question(s) : Let A be a n-convex (convex set in the n dimentionnal euclidian affine space $\mathbb A_n$) and H a hyperplane of $\mathbb A^n$ such that $B=A\cap H$ has maximal volume (resp. such that the boundary of $B'=A\cap H$ is maximal in volume) a) Is $er(B)\geq er(A) $? b) Is $er(B')\geq er(A)$? I think the answer (especially the b) question) if positive implies the resolution of the question in Volumes of Sets of Constant Width in High Dimensions podted by Gil Kalai. Let me try an heuristic for the first inequality, in 3 dimensions (I write vol(A) as well as for a volume or an area, as soon as there is no ambiguities) : Take $t\mapsto H_t$ a path of plans parallele to H that go throught C and that intersect C in B_t. Now we associate to B_t a disk D_t that radius is $er(B_t)$ such that its center is $t\in T$ and that T is a line. Consider $E=\bigcup_{t\in T} D_t$. It is clear that $Vol (E) = Vol (C)$. It is also clear that E is the π- rotation solid generated by , say E_0 : union of coplanar "effectives diameters" The question a) for dim 3 is then equivalent to : Is $vol(E_0)$ smaller then $vol(B)$? This would be automatically the case if E_0 can be isometrically embeded in C whom a greater section (in volume, well in this case "area") is B - it appears that the question that translate the possibility of this transversal embedding is : can we put the line T inside C and take in each "efficient disk" an "efficient diametre" inside C in such a way that they are co-planar.. we could imagine that there is some torsion around T that make this operation impossible, despite convexity condiderations... never the less, this intermediary request is stronger then the one of the (here part a) but also wrt part b)) question... The problem of this heuristic is not only the torsions phenomenons that I think they might not occure in 3 dim (and that we can probably get rid of, considering that the case that we are dealing with are well behaving, especially throught heredity) but the fact that I'm not sure it can be adapted to greater dimensions, because E_0 is 2-dimensional in any case... that does not go against the relevance of the questions but that does not give at least to me a clear hope or a clue, well maybe this discussion can be adapted to the general case... or not.

论坛问题帖 | no_answers_at_retrieval | jcdornano | 2020-04-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018256 | MO-359244

Which curves and surfaces are realizable by linkages? references?

Ok, so I try to formulate rigorously the question in the title, for which I am asking for references. My definitions may be flawed, so feel free to adjust/correct them! I care about dimensions 2 and 3 below, but feel free to mention general d as well. Let $d\ge 1$ be an integer, $G=(V,E)$ be a graph, $w:E\to [0, \infty)$ be a weight function. A realization in $\mathbb R^d$ with graph G and weight w is a map $P:V\to\mathbb R^d$ with the further property that $|P(v)-P(v')|=w(v,v')$ whenever $\{v,v'\}\in E$. I will identify such P with its image, I hope it's not a problem. Edit (May 11, 2020): As pointed out by Misha, this below definition is not correct. The action of isometries makes the set of all realizations of a linkage always cover all $\mathbb R^d$. He indicates a paper in which a more inclusive definition is formulated in $d=2$. (Previous "wrong" definition: I say that a set $A\subset \mathbb R^d$ is realizable by linkages if there exists G,w as above and an cover of A by open sets of $\mathbb R^d$ such that for every $U\subset\mathbb R^d$ in the cover there exist G,w such that that the union of all (images of) realizations of G,w in $\mathbb R^d$ intersected with U coincides with $A\cap U$.) To "fix the problem", following the we will allow a subset of vertices of G to be kept fixed in $\mathbb R^d$. In dimension 2 this apparently generalizes the definition in the above paper, but I think that the result of the paper still allows to reply positively to the $d=2$ case of the question, with little extra work. Revised definition: We say that $A\subset \mathbb R^d$ is realizable by linkages if there exists a cover of A by open sets of $\mathbb R^d$ such that for every $U\subset\mathbb R^d$ in the cover there exist $G=(V,E)$ and w as above, a subset $F\subset V$, and a map $\phi: F\to\mathbb R^d$, such that the union of all (images of) those realizations of G,w which restricted to F equal ϕ, intersected with U, coincides with $A\cap U$. Question: Say $d=2$ or $d=3$. Is it true that all algebraic sets $A\subset\mathbb R^d$ are realizable by linkages? What are references for this? (Note: as of May 11 2020, it appears to me that the case $d=2$ is nicely treated in the answers given, while the case $d=3$ is not yet treated, possibly due to the previously bad definition.) I found some mention of this, without references on Branko Grünbaum's "Lectures on lost mathematics", dated around 1975, and he says there that $d=2$ case is known, but does not give references, and $d=3$ case is a question by Hilbert which is open (but again no references there).

论坛问题帖 | has_answers | Mircea | 2020-05-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018257 | MO-361474

Basic research problems references

I have been looking for research problems in pure mathematics that I can try to solve for publishing papers. I am quite aware that it takes a lot of time and effort to get to a level where I can do some "serious" research work. Meanwhile, I was thinking that, may be, I can try to solve problems which are at the level of 1st year graduate level (or even Master's level). By "level", I mean that a 1st year grad student should know all the terms used in the paper. I understand that maintaining a list of unsolved/open problems is an extremely tough task and probably the best method is to look for journals/papers where authors include open/unsolved/extension of the problems. I would appreciate if you can include here: (i) List

of problems (I have also collected a list of unsolved/open problems from the internet which I would be happy to share here in one of the answers). (ii) Journals which concerns articles at the "level" mentioned above (For e.g. I hope that this list will exclude annals of math, Inventiones Mathematicae or any top journals which mostly include high level research papers. The American Mathematical Monthly is an excellent source.) Remarks: 1) I admit that it seems like a very vague question but I am also confident of the fact that many other fellow students are looking for such a source. If the community feels that it should be closed, I will also second it. 2) One of the great sources for finding research problems are conferences (or proceedings of ICM) where the leading researchers include open problems, but those problems, in my view, seem to be untouchable for most of the 1st year graduate students. 3) From one point view, there is a related question about letters in mathematics. This is definitely one of the answer I would have expected for this question. I will be glad to provide any details if needed. Kindly let me know if I should post this question somewhere else.

论坛问题帖 | has_answers | MUH | 2020-05-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018258 | MO-361595

Generalized figures of constant width

Is it known which plane figures Q can rotate touching three given circles A, B, and C? This question was asked by Lazar Lyusternik in 1946, there is only one reference to this paper that solves the problem in one limit case. Do you know of any other research on this question? (I learned this question from Sergei Tabachnikov.)

论坛问题帖 | has_answers | Anton Petrunin | 2020-05-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018259 | MO-362679

A constant bizarrely related to the Fibonacci Numbers

For roughly the past month, I have been studying denesting radicals. For example: the expression $\sqrt[3]{\sqrt[3]2-1}$ is a radical expression that contains another radical expression, so this radical is nested. Is there a way to express this with radicals that are not (or not as) nested? Writing it in this way is referred to as denesting, and indeed, there is one such way. Ramanujan mysteriously found that $$\sqrt[3]{\sqrt[3]2-1}=\sqrt[3]{\frac 19}-\sqrt[3]{\frac 29}+\sqrt[3]{\frac 49}.$$ I found that this is also equal to $\sqrt{\sqrt[3]{\frac 43}-\sqrt[3]{\frac 13}}$, which does not denest it, but it does write the expression under a radical of a coprime degree, which fascinates me. I have found an abundance of results, one of which fascinates me the most, and is on the constant, $$1-\sqrt[3]{\frac 12}+\sqrt[3]{\frac 14}.$$ This constant is equal to all of the expressions below. Note that the degree of the radicals are, lo and behold, the Fibonacci numbers!
$$\sqrt{\frac 32\bigg(\sqrt[3]2-\frac 1{\sqrt[3]2}\bigg)}$$
$$\sqrt[3]{\frac{3^2}{2^2}\big(\sqrt[3]2-1\big)}$$ $$\sqrt[5]{\frac{3^3}{2^3}\bigg(\frac 3{\sqrt[3]2}-\sqrt[3]2-1\bigg)}$$
$$\sqrt[8]{\frac{3^5}{2^5}\bigg(4-\frac{5}{\sqrt[3]2}\bigg)}$$
$$\sqrt[13]{\frac{3^8}{3^8}\bigg(1+\frac{17}{\sqrt[3]2}-\frac{23}{\sqrt[3]4}\bigg)}$$
and, slightly breaking the pattern,
$$\sqrt[21]{\frac{3^{14}}{2^{14}}\big(41-59\sqrt[3]2+21\sqrt[3]4\big)}$$ and presumably, this list goes on forever (but the numbers start becoming pretty big). So... what on earth is going on here? It appears that for some nth Fibonacci number F_n, this is equal to (for at least most of the time),
$$\sqrt[F_n]{\frac{3^{F_{n-1}}}{2^{F_{n-1}}}\big(a+b\sqrt[3]2+c\sqrt[3]4\big)}$$ for some $\{a, b, c\}\subset \mathbb R$, with a minimal polynomial of $4x^3-12x^2+18x-9$. Can anybody explain these wild affairs? Don't know of any other appropriate tags

论坛问题帖 | has_answers | Mr Pie | 2020-06-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018260 | MO-364611

Frechet-Urysohn quotient of second countable locally compact Hausdorff space

In this paper from 2010 https://cmuc.karlin.mff.cuni.cz/pdf/cmuc1001/arhangav.pdf Arhangelskii asks if there is a quotient of a second countable locally compact Hausdorff space which is Frechet-Urysohn and Tychonoff but not first countable at any point (Problems 5.14 and 5.15). Is this problem still open?

论坛问题帖 | no_answers_at_retrieval | Douglas Somerset | 2020-07-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018261 | MO-366687

Conjecture about minimal number of edge crossings in complete bipartite graphs

I am interested in the status of the conjecture about the minimum number of edge crossings $cr(K_{m,n})$ in a drawing of the complete bipartite graph $K_{m,n}$. The Wikipedia article https://en.wikipedia.org/wiki/Tur%C3%A1n%27s_brick_factory_problem led me to study the original papers of Zarankiewicz (On a problem of P. Turan concerning graphs) from 1954 and of Urbanik (Solution du problème posé par P. Turán) from 1955. I wondered whether someone could tell me whether an asymptotic approach has been successfully attempted (letting $n\to\infty$). If so, I would be very interested in any references for that.

论坛问题帖 | has_answers | Ruth-NO | 2020-07-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018262 | MO-36738

Detecting almost-primes quickly

There are many fast algorithms (deterministic and probabilistic) for detecting primality. Are there any fast algorithms (probabilistic ones allowed) known for detecting whether a number is the product of at most two primes? In this context, can anybody think of a property fulfilled by all or most products of two primes, but by few other numbers? (You can't explicitly use divisor sums or the Moebius function, obviously, as then the game becomes (a) trivial (b) of doubtful computational utility.)

论坛问题帖 | has_answers | H A Helfgott | 2010-08-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018263 | MO-369168

Two more counterexamples to a conjecture from 1975 about hamiltonicity of digraphs

Question from 2013 gives one counterexample to Nash-Williams's conjecture 1975 about hamiltonicity of dense digraphs. In the linked answer, @LouisD "reverse engineered" the counterexample and pointed out it is special case in another paper. We found 2 more digraph counterexamples with directed edges: g1=[(0, 3), (0, 4), (1, 3), (1, 5), (2, 4), (2, 5), (3, 0), (3, 1), (3, 4), (3, 5), (4, 1), (4, 2), (4, 3), (4, 5), (5, 0), (5, 2), (5, 3), (5, 4)] g2=[(0, 2), (0, 3), (0, 4), (0, 5), (1, 3), (1, 4), (2, 0), (2, 4), (3, 0), (3, 1), (3, 4), (3, 5), (4, 0), (4, 3), (5, 0), (5, 1), (5, 2), (5, 3), (5, 4)] g3=[(0, 2), (0, 3), (0, 4), (0, 5), (1, 2), (1, 5), (2, 0), (2, 1), (2, 4), (2, 5), (3, 1), (3, 5), (4, 1), (4, 2), (5, 0), (5, 1), (5, 2), (5, 3)] Q1 Are these counterexamples true? Q2 Can we get infinitely many counterexamples from the small ones? All three counterexamples are planar.

论坛问题帖 | no_answers_at_retrieval | joro | 2020-08-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018264 | MO-370047

Does the equation $\sigma(\sigma(x^2))=2x\sigma(x)$ have any odd solutions?

This question was posted in MSE in early August 2020. It did garner several upvotes, but did not receive any responses. I have therefore cross-posted it here, hoping that it gets answered. Let $\sigma(x)$ denote the classical sum of divisors of the positive integer x. Here is my question: Does the equation $\sigma(\sigma(x^2))=2x\sigma(x)$ have any odd solutions? MY ATTEMPT I tried searching for solutions to the equation in Sage Cell Server, in the range $1 < x \leq {10}^6$, here is the Pari-GP code: for(x=1, 1000000, if(sigma(sigma(x^2))==2*x*sigma(x),print(x,factor(x)))) Here are the results: 9516[2, 2; 3, 1; 13, 1; 61, 1] 380640[2, 5; 3, 1; 5, 1; 13, 1; 61, 1] Note that both results obtained $x_1 = 9516$ and $x_2 = 380640$ are even. The Pari-GP interpreter of Sage Cell Server crashes as soon as a search limit of ${10}^7$ is specified. CONJECTURE The equation $\sigma(\sigma(x^2)) = 2x\sigma(x)$ does not have any odd solutions. Alas, I have no proof. SOME PARTIAL RESULTS Per Will Jagy's answer (and a subsequent comment by Erick Wong) to the following MSE question, we have the conjectured (sharp?) upper bound $$\frac{\sigma(x^2)}{x\sigma(x)} \leq \prod_{\rho}{\frac{{\rho}^2 + \rho + 1}{{\rho}^2 + \rho}} = \frac{\zeta(2)}{\zeta(3)}.$$ Claim: $\sigma(x^2) \neq p^r$ if prime $p \geq 5$. Suppose to the contrary that the equation $\sigma(\sigma(x^2))=2x\sigma(x)$ has an odd solution, and that $\sigma(x^2) = p^r$ for some prime $p \geq 5$. Then we have $$\frac{\sigma(p^r)}{p^r}=\frac{\sigma(\sigma(x^2))}{\sigma(x^2)}=\frac{2x\sigma(x)}{\sigma(x^2)} \geq \frac{2\zeta(3)}{\zeta(2)} \approx 1.4615259388.$$ But we know that the abundancy index $\sigma(p^r)/p^r$ is bounded above by $$\frac{\sigma(p^r)}{p^r} < \frac{p}{p - 1} \leq \frac{5}{4}.$$ This is a contradiction. Claim: $\sigma(x^2) \neq p^r q^s$ if primes $p \geq 5$ and $q \geq 7$. Suppose to the contrary that the equation $\sigma(\sigma(x^2))=2x\sigma(x)$ has an odd solution, and that $\sigma(x^2) = p^r q^s$ for some primes $p \geq 5$ and $q \geq 7$. Then we have $$\frac{\sigma(p^r)}{p^r}\cdot\frac{\sigma(q^s)}{q^s}=\frac{\sigma(\sigma(x^2))}{\sigma(x^2)}=\frac{2x\sigma(x)}{\sigma(x^2)} \geq \frac{2\zeta(3)}{\zeta(2)} \approx 1.4615259388.$$ But we know that the product of abundancy indices $(\sigma(p^r)/p^r)(\sigma(q^s)/q^s)$ is bounded above by $$\frac{\sigma(p^r)}{p^r}\cdot\frac{\sigma(q^s)}{q^s} < \frac{p}{p - 1}\cdot\frac{q}{q - 1} \leq \frac{5}{4}\cdot\frac{7}{6} = \frac{35}{24} = 1.458\overline{333}.$$ This is a contradiction. Edit (August 27, 2020 - 4:21 PM Manila time) MOTIVATION FOR THE PROBLEM Slowak (1999) proved that an odd perfect number $N = u^t v^2$ (with special prime u satisfying $u \equiv t \equiv 1 \pmod 4$ and $\gcd(u,v)=1$) must be of the form $$\frac{u^t \sigma(u^t)}{2}\cdot{d},$$ where $d > 1$. Dris (2017) showed further that d must have the form $$\frac{D(v^2)}{\sigma(u^{t-1})}=\gcd(v^2,\sigma(v^2))=\frac{\sigma(v^2)}{u^t}=\frac{v^2}{\sigma(u^t)/2},$$ where $D(y)=2y-\sigma(y)$ is the deficiency of y. If the odd perfect number $N = u^t v^2$ is of the form $$\frac{u^t \sigma(u^t)}{2}\cdot{v},$$ then by the results of Slowak and Dris, we have the equations $$\sigma(v^2) = u^t v$$ and $$2v = \sigma(u^t),$$ which led me to my question in this post.

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2020-08-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018265 | MO-370207

The action of the unitary divisors group on the set of divisors and odd perfect numbers

Let n be a natural number. Let $U_n = \{d \in \mathbb{N}\mid d\mid n \text{ and } \gcd(d,n/d)=1 \}$ be the set of unitary divisors, D_n be the set of divisors and $S_n=\{d \in \mathbb{N}\mid d^2 \mid n\}$ be the set of square divisors of n. The set U_n is a group with $a\oplus b := \frac{ab}{\gcd(a,b)^2}$. It operates on D_n via: $$ u \oplus d := \frac{ud}{\gcd(u,d)^2}$$ The orbits of this operation "seem" to be $$ U_n \oplus d = d \cdot U_{\frac{n}{d^2}} \text{ for each } d \in S_n$$ From this conjecture it follows (also one can prove this directly since both sides are multiplicative and equal on prime powers): $$\sigma(n) = \sum_{d\in S_n} d\sigma^*(\frac{n}{d^2})$$ where σ^* denotes the sum of unitary divisors. Since $\sigma^*(k)$ is divisible by $2^{\omega(k)}$ if k is odd, where $\omega=$ counts the number of distinct prime divisors of k, for an odd perfect number n we get (Let now n be an odd perfect number): $$2n = \sigma(n) = \sum_{d \in S_n} d \sigma^*(\frac{n}{d^2}) = \sum_{d \in S_n} d 2^{\omega(n/d^2)} k_d $$ where $k_d = \frac{\sigma^*(n/d^2)}{2^{\omega(n/d^2)}}$ are natural numbers. Let $\hat{d}$ be the largest square divisor of n. Then: $\omega(n/d^2)\ge \omega(n/\hat{d}^2)$. Hence we get: $$2n = 2^{\omega(n/\hat{d}^2)} \sum_{d \in S_n} d l_d$$ for some natural numbers l_d. If the prime 2 divides not the prime power $2^{\omega(n/\hat{d}^2)}$, we must have $\omega(n/\hat{d}^2)=0$ hence $n=\hat{d}^2$ is a square number, which is in contradiction to Eulers theorem on odd perfect numbers. So the prime 2 must divide the prime power $2^{\omega(n/\hat{d}^2)}$ and we get: $$n = 2^{\omega(n/\hat{d}^2)-1} \sum_{d \in S_n} d l_d$$ with $l_d = \frac{\sigma^*(n/d^2)}{2^{\omega(n/d^2)}}$. Hence the odd perfect number, satisifies: $$n = \sum_{d^2\mid n} d \frac{\sigma^*(n/d^2)}{2^{\omega(n/d^2)}}=:a(n)$$ Hence an odd perfect number satisifies: $$n = a(n)$$ So my idea was to study the function $a(n)$, which is multiplicative on odd numbers, on the right hand side and what properties it has to maybe derive insights into odd perfect numbers. The question is if it ever can happen that an odd number n satisfies: $n=a(n)$? (checked for $n=2k+1$ and $1 \le k \le 10^7$) Edit: Conjecture: For all odd $n \ge 3$ we have $a(n)<n$. This would prove that there exists no odd perfect number. This conjecture could be proved as follows: Since $a(n)$ is

multiplicative, it is enough to show that for an odd prime power p^k we have $$a(p^k) < p^k$$ The values of a at prime powers are not difficult to compute and they are: $$a(p^{2k+1})= \frac{p^{2(k+1)}-1}{2(p-1)}$$ and $$a(p^{2k}) = \frac{p^{2k+1}+p^{k+1}-p^k-1}{2(p-1)}$$ However, I am not very good at proving inequalities, so: If someone has an idea how to prove the following inequalities for odd primes p that would be very nice: $$p^{2k+1} > \frac{p^{2(k+1)}-1}{2(p-1)}, \text{ for all } k \ge 0$$ and $$p^{2k} > \frac{p^{2k+1}+p^{k+1}-p^k-1}{2(p-1)}, \text{ for all } k \ge 1$$ Thanks for your help! The inequalities have been proved here: https://math.stackexchange.com/questions/3807399/two-inequalities-for-proving-that-there-are-no-odd-perfect-numbers

论坛问题帖 | has_answers | user6671 | 2020-08-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018266 | MO-370905

$π(x+y) - π(x) ≤ c·y/\ln(y)$ for some constant c?

(I posted this question on Math SE but it has had no answer for a year now so I would like to ask if anyone here can provide one.) Thinking about the prime number theorem, I wondered whether it is known that there is some constant c such that $π(x+y) ≤ π(x) + c·y/\ln(y)$ for every integers $x,y > 1$. I read that experts believe $π(x+y) ≤ π(x) + π(y)$ fails for some y, since it fails for $y = 3159$ if the k-tuple conjecture holds, but it is just barely false, so I am curious if it is known to be true if the inequality is relaxed by a constant factor. If so, is it also known that $π(x+y) ≤ π(x) + π(y) + c·\!\sqrt{y}·\ln(y)$ for some constant c? I simply do not know how to search for such conjectures, and neither Wikipedia nor Wolfram seem to state any results that would affirm or refute these two conjectures easily, so any references would be appreciated!

论坛问题帖 | has_answers | user21820 | 2020-09-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018267 | MO-371135

Inductively computing Mersenne primes / perfect numbers?

For two sets A,B set $A+B = \{a +b | a \in A,b \in B\}$. Let $(x_n)_{n \in \mathbb{N}}$ be independent variables. Let $\sigma(n)$ be the sum of divisors of n. Set $\hat{\phi}(1) = \{x_1\}$ and then inductively: $$\hat{\phi}(n) = \{x_n\}$$ if $\sigma(k) \neq n$ for all $k \in \mathbb{N}$ and $$\hat{\phi}(n) = \bigcup_{ \sigma(m) = n} \left\{ \sum_{d|m} \hat{\phi}(d) \right\}$$ otherwise. Examples: 1 [x1] 2 [x2] 3 [x1 + x2] 4 [2*x1 + x2] 5 [x5] 6 [x1 + x5] 7 [3*x1 + 2*x2] 8 [4*x1 + 2*x2] 9 [x9] 10 [x10] 11 [x11] 12 [x1 + x11, 3*x1 + 2*x2 + x5] 13 [2*x1 + x2 + x9] 14 [3*x1 + x2 + x9] 15 [7*x1 + 4*x2] 16 [x16] 17 [x17] 18 [x1 + x10 + x2 + x5, x1 + x17] 19 [x19] 20 [x1 + x19] 21 [x21] 22 [x22] 23 [x23] 24 [9*x1 + 5*x2 + x5, x1 + x23, 7*x1 + 4*x2 + x9] 25 [x25] 26 [x26] 27 [x27] 28 [8*x1 + 5*x2 + 2*x5, 6*x1 + x11 + 3*x2 + x5] Conjecture: Let $n=2^{p-1}(2^p-1)$ be an even perfect number. Then there exists a $x \in \hat{\phi}(2n)$, such that, if we plug in 1 for all free variables in x we get: $$ y = x(1,\cdots,1)$$ such that: $$ y+1 = 2^q-1$$ is a Mersenne prime. Furthermore we have: $$N=2^{q-1}(2^q-1)$$ is the next even perfect number after n. I have tested this for $n=6,28,496$. Unfortunately, my naive computation method hits the limit for the next perfect number. So I am asking if somebody can check if the next perfect number satisfies this conjecture or not? It would be nice if you share also your method to check the number. Thanks for your help! Here is some SAGE code I used. And here is some output: n y 6 2 6 6 28 30 496 2 496 126 Edit: Here are the trees corresponding to $2\cdot 6$: [( )] [((12, 6, 1), (12, 6, 2), ____(12, 6, 3) ____(12, 6, 6))] [(/ / / /) ( )] [((3, 2, 1) (3, 2, 2) , (6, 5, 1) (6, 5, 5)), ((12, 11, 1), (12, 11, 11))] And here are the corresponding trees to $2\cdot 28$: [([((56, 28, 1), (56, 28, 2), _____(56, 28, 4) _____(56, 28, 7)_____ _____(56, 28, 14)____ [(/ / / / / / / [((4, 3, 1) ____(4, 3, 3) (7, 4, 1) (7, 4, 2) _____(7, 4, 4) (14, 13, 1) _____(14, 13, 13)_____ [(/ / / / / / / [((3, 2, 1) (3, 2, 2) , (4, 3, 1) ____(4, 3, 3) (13, 9, 1) ____(13, 9, 3) (13, 9, 9) [(/ / / / [((3, 2, 1) (3, 2, 2) , (3, 2, 1) (3, 2, 2) ,) (_____(56, 28, 28)_____) ((56, 28, 1), (56, 28, 2), _____(56, 28, 4) / / / / / /) (/ / (28, 12, 1) (28, 12, 2) ____(28, 12, 3) _____(28, 12, 4) ____(28, 12, 6) _____(28, 12, 12)_____) ((4, 3, 1) ____(4, 3, 3) / / / / / / / / / /) (/ / (3, 2, 1) (3, 2, 2) (4, 3, 1) ____(4, 3, 3) (6, 5, 1) (6, 5, 5) (12, 6, 1) (12, 6, 2) ____(12, 6, 3) ____(12, 6, 6)) ((3, 2, 1) (3, 2, 2) , / / / / / /) ((3, 2, 1) (3, 2, 2) (3, 2, 1) (3, 2, 2) (6, 5, 1) (6, 5, 5)), ( ) _____(56, 28, 7)_____ _____(56, 28, 14)____ _____(56, 28, 28)_____) / / / / / / / / / /) ((7, 4, 1) (7, 4, 2) _____(7, 4, 4) (14, 13, 1) _____(14, 13, 13)_____ (28, 12, 1) (28, 12, 2) ____(28, 12, 3) _____(28, 12, 4) ____(28, 12, 6) _____(28, 12, 12)) ((56, 39, 1), / / / / / / / / / / / / /) ((4, 3, 1) ____(4, 3, 3) (13, 9, 1) ____(13, 9, 3) (13, 9, 9) (3, 2, 1) (3, 2, 2) (4, 3, 1) ____(4, 3, 3) (6, 5, 1) (6, 5, 5) (12, 11, 1) (12, 11, 11)) (/ / / / / /) ((3, 2, 1) (3, 2, 2) , (3, 2, 1) (3, 2, 2) , (3, 2, 1) (3, 2, 2)), ( ) (____(56, 39, 3) _____(56, 39, 13)_____ _____(56, 39, 39)_____) ((56, 39, 1), ____(56, 39, 3) / / / / / / / / / / /) (/ / (3, 2, 1) (3, 2, 2) , (13, 9, 1) ____(13, 9, 3) (13, 9, 9) (39, 18, 1) (39, 18, 2) ____(39, 18, 3) ____(39, 18, 6) (39, 18, 9) _____(39, 18, 18)_____) ((3, 2, 1) (3, 2, 2) , / / / / / / / / / /) ((3, 2, 1) (3, 2, 2) , (3, 2, 1) (3, 2, 2) (6, 5, 1) (6, 5, 5) (18, 10, 1) (18, 10, 2) (18, 10, 5) (18, 10, 10)), ( ]])] _____(56, 39, 13)_____ _____(56, 39, 39)_____)] / / / / / / / / /)] (13, 9, 1) ____(13, 9, 3) (13, 9, 9) (39, 18, 1) (39, 18, 2) ____(39, 18, 3) ____(39, 18, 6) (39, 18, 9) _____(39, 18, 18))] / / / / / / / /)] (3, 2, 1) (3, 2, 2) , (3, 2, 1) (3, 2, 2) (6, 5, 1) (6, 5, 5) (18, 17, 1) (18, 17, 17))] The trees were constructed with the following code: def inductiveGraph(n): from itertools import product ll = [i for i in range(1,n) if sigma(i)==n] #print(ll) if len(ll)==0: return [LabelledOrderedTree([],label=n)] else: phll = [] for m in ll: ml = list(product(*[[LabelledOrderedTree(ph,label=(n,m)) for ph in inductiveGraph(d)] for d in divisors(m)])) #print(n,ml) for x in ml: phll.append(x) return phll print(ascii_art(inductiveGraph(2*28))) Furthermore, plugging in 1 for all variables in $x(1,\cdots,1)$ counts the number of leafs in the tree which corresponds to $x \in \hat{\phi}(2n)$. Related question: Additive number theory, Hilbert spaces and polynomial rings?

论坛问题帖 | has_answers | user6671 | 2020-09-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018268 | MO-372254

What groups should I test my conjecture on?

I have a conjecture that a certain criterion is enough for two groups to be isomorphic. I tested it on all pairs of groups up to size 12, and it worked like a charm. I know, however, that groups are strange and that it is very likely that my conjecture will break for larger groups. I have made myself a python library that handles making groups and seeing they have certain properties/are isomorphic, but I do not know what pairs of groups to test my conjecture on. What groups form a "representative sample" of all groups? What paris of groups would it be absolutely crucial for me to test my hypothesis on? I don't see the need for any answer to be complete, but any ideas for pairs that would be interesting to test on would be appreciated. EDIT: The conjecture is that if two groups have the same count of elements of every order, then they must be isomorphic. I proved this for abelian groups, and am now wondering if it is true in general.

论坛问题帖 | has_answers | Milo Moses | 2020-09-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018269 | MO-373497

Prove that there are no composite integers $n=am+1$ such that $m \ | \ \phi(n)$

Let $n=am+1$ where $a $ and $m>1$ are positive integers and let p be the least prime divisor of m. Prove that if $a<p$ and $ m \ | \ \phi(n)$ then n is prime. This question is a generalisation of the question at https://math.stackexchange.com/questions/3843195/let-n-apq1-prove-that-if-pq-phin-then-n-is-prime. Here the special case when m is a product of two distinct odd primes has been proven. The case when m is a prime power has also been proven here https://arxiv.org/abs/2005.02327. How do we prove that the proposition holds for an arbitrary positive integer integer $m>1 $? (I have not found any counter - examples). Note that if $n=am+1$ is prime, we have $\phi(n)= n-1=am$. We see that $m \ | \ \phi(n) $. Its the converse of this statement that we want to prove i.e. If $m \ | \ \phi(n) $ then n is prime. If this conjecture is true, then we have the following theorem which is a generalisation (an extension) of Lucas's converse of Fermat's little theorem. $\textbf {Theorem} \ \ 1.$$ \ \ \ $ Let $n=am+1$, where a and $m>1$ are positive integers and let p be the least prime divisor of m with $a<p$. If for each prime q_i dividing m, there exists an integer b_i such that ${b_i}^{n-1}\equiv 1\ (\mathrm{mod}\ n)$ and ${b_i}^{(n-1)/q_i} \not \equiv 1(\mathrm{mod}\ n)$ then n is prime. Proof. $ \ \ \ $ We begin by noting that ${\mathrm{ord}}_nb_i\ |\ n-1$. Let $m={q_1}^{a_1}{q_2}^{a_2}\dots {q_k}^{a_k}$ be the prime power factorization of m. The combination of ${\mathrm{ord}}_nb_i\ |\ n-1$ and ${\mathrm{ord}}_nb_i\ \nmid (n-1)/q_i$ implies ${q_i}^{a_i}\ |\ {\mathrm{ord}}_nb_i$. $ \ \ $${\mathrm{ord}}_nb_i\ |\ \phi (n)$ therefore for each i, ${q_i}^{a_i}\ |\ \phi (n)$ hence $m\ |\ \phi (n)$. Assuming the above conjecture is true, we conclude that n is prime. Taking $a=1$, $m=n-1$ and $p=2$, we obtain Lucas's converse of Fermat's little theorem. Theorem 1 is thus a generalisation (an extension) of Lucas's converse of Fermat's little theorem. This question was originally asked in the Mathematics site, https://math.stackexchange.com/questions/3843281/prove-that-there-are-no-composite-integers-n-am1-such-that-m-phin. On recommendation by the users, it has been asked here.

论坛问题帖 | has_answers | ASP | 2020-10-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018270 | MO-375219

Updated bounds or references for an old Erdős problem -- coloring the plane with multiple forbidden distances?

Define a graph G_1 where the vertices of G_1 are the points of the plane $\mathbb{R}^2$, and a pair of vertices p, q is connected by an edge if and only if the Euclidean distance $d(p,q) =1$. The Hadwiger-Nelson unit-coloring problem asks for the "chromatic number of the plane": i.e. what is $\chi(G_1)$? The best known bounds are that $5 \le \chi(G_1) \le 7$, with the lower bound improved from 4 to 5 in a breakthrough result by Aubrey de Grey in 2018. Apparently Erdős asked about the following variant. Let $S =\{ d_1, d_2, \dots, d_k \}$ be a set of positive numbers. Define G_S with vertices corresponding to points in the plane, and edges corresponding to pairs of points at any distance d_1, d_2, ..., or d_k. Let $f(k)$ denote the maximum chromatic number $\max \chi(G_S)$ as S ranges over all sets of k positive numbers. Clearly $f(k) \le 7^k$, so $f(k)$ is well defined and is finite for every k. Roughly, how does $f(k)$ grow as $k \to \infty$? According to Soifer (in The Mathematical Coloring Book), Erdős wrote that "it is not hard to see" that $$ \lim_{k \to \infty} \frac{f(k)}{k} \to \infty.$$ As pointed out in the comments, this follows from the fact that a square grid with $\sqrt{n} \times \sqrt{n}$ points only realizes $O \left(n / \sqrt{\log n} \right)$ distinct distances. What are the best known upper and lower bounds on $f(k)$? It was recently shown by Exoo and Ismailescu that $f(2) \ge 6$, but I'm not aware of any nontrivial lower bounds for $f(k)$ even for $k \ge 3$. Any more information for small k would be great, but I'm especially interested in the rate of growth as $k \to \infty$. Erdős apparently asked whether $f(k)$ grows only polynomially fast, and I am wondering whether this is still an open problem.

论坛问题帖 | has_answers | Matthew Kahle | 2020-10-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018271 | MO-375607

Profound but not popular mathematical topics and notions

The algebraic Theory of Invariants used to be a hot topic until David Hilbert proved his two theorems about invariants. Then for tens of years, the popularity of the topic went down a long time before it picked up again. Question What are today's mathematical known topics and/or notions that are profound but not popular? Together with an example, could you add an explanation of such a situation? Example from Computer Science -- geometric SIMD (fine grain parallel processing) was a popular and hot topic till the middle of 1980'. Then you hardly hear about it while the idea is as fundamental as always. The explanation is two-folded but very simple. On the one hand, there is some learning and new understanding involved in geometric SIMD processing; one needs to acquire new habits, new reflexes. On the other hand, the technology progress was such that people were satisfied with the results obtained without bothering with the SIMD ways. (Underneath, the new traditional computer architecture is not that traditional -- these days, it incorporates quite a bit of parallelity). We see that the geometric SIMD is not popular for the wrong reasons, and a lot of potentials is

wasted.

论坛问题帖 | has_answers | Wlod AA | 2020-11-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018272 | MO-376217

A conjectured upper bound for the mean value of prime divisors inside prime gaps

In 1969 C.A. Grimm stated this interesting conjecture: the prime gap $\,G_n=\{x\in N:p_n\lt x\lt p_{n+1}\}\,$ contains at least $\,\#G_n=(p_{n+1}-p_n)-1=g_n-1\,$ distinct prime divisors, that is if $\,V(n)=(p_n+1)(p_n+2)...(p_{n+1}-1)$ $$\omega\,[V(n)]\ge g_n-1$$ Let's define the following additive arithmetic function: $$\sigma_p(n)=\sum_{k=1}^r p_k\;\;\;\;\;(\,n=\prod_{k=1}^r p_k^{a_k}\,)$$ Well, I have made some computations in order to conjecture a possible estimate for $\,\sigma_p(V(n))$. The following graph reports the values ($x10^6$) of $\,\sigma_p(V(n))-p_n\,$ till $\,n=10^5$: A possible estimate should be $$\sigma_p(V(n))=2n\,(\log n)^{3/2}+O(n\log n)$$ But under Cramér's conjecture $\,g_n=p_{n+1}-p_n=O((\log p_n)^2)$, therefore we can deduce the following upper bound for the mean value of prime divisors inside the gap $\,g_n$: $$\frac {\sigma_p(V(n))} {\omega\,[V(n)]}\lt\frac {2n\sqrt {\log n}} {\log n+2\log {\log n}}\lt\frac {2n} {\sqrt {\log n}}$$ Notice that the max prime divisor inside the gap is less than $$\frac {p_{n+1}} 2\sim\frac {n\log n+g_n} 2\sim\frac {n\log n+(\log n)^2} 2\sim\frac {\log n\,(n+\log n)} 2$$ I ask your opinions and suggestions about the above arguments. Many thanks.

论坛问题帖 | no_answers_at_retrieval | Augusto Santi | 2020-11-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018273 | MO-376480

How to put a monetary incentive on an open research problem?

Since the old days, many famous mathematicians have been attaching monetary rewards to problems. I would like to put monetary bounties on some research mathematics problems. Some of those problems might be solved or partially solved by a hard-working college graduate in pure math. The ideal platform to put these questions would be a website like MathOverflow: high visibility, acceptable ranking algorithm, and little distractions (Ads). However, I have yet to find a single forum for posting these kind of questions, let alone being "ideal". I am not sure if we could attach a monetary incentive in the body of the question here on this site. I could also hire a research assistant but I am not sure if I could attract a high quality one as I am not a tenure-track faculty at a renowned institute. Another alternative will be freelancing websites. However, no one put research math questions there: those sites are full of coding problems and homework problems. And of course, we could publish an advertisement just like what have done by those famous mathematicians. However, with little reputation, we are unlikely to attract enough attention. What resources do we have for posting the open research problems?

论坛问题帖 | no_answers_at_retrieval | dodo | 2020-11-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018274 | MO-377606

Open problems in matroid theory

I read Oxley's book on matroid theory and found the theory fascinating. At the end, Oxley stated some open problems and conjectures in matroid theory. Are there any modern lists about such problems? Are there more open conjectures in matroid theory?

论坛问题帖 | has_answers | LogicTheorist | 2020-11-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018275 | MO-378730

How small the radical of $xyz(x+y+z)$ can be infinitely?

This is an open problem. Let x,y,z be coprime integers (not necessarily pairwise coprime) and no proper subset sum of $\{x,y,z,-(x+y+z)\}$ is zero. For a quadruple $(x,y,z,-(x+y+z))$ define the quality
$$q(x,y,z,-(x+y+z))=\frac{\log{(\max(|x|,|y|,|z|,|x+y+z|))}}{\log(rad(xyz(x+y+z))}$$ where rad is the radical. We have quality $q= 3+o(1)$ given by $$x=1, y=3 (t - 1) t, z=(t - 1)^3, -(x+y+z)=-t^3$$ and at integers set $t=2^{large}$. Can we get quality $3+C$ infinitely often for positive C?

论坛问题帖 | no_answers_at_retrieval | joro | 2020-12-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018276 | MO-37902

Walking to infinity on the primes: The prime-spiral moat problem

It is an unsolved problem to decide if it is possible to "walk to infinity" from the origin with bounded-length steps, each touching a Gaussian prime as a stepping stone. The paper by Ellen Gethner, Stan Wagon, and Brian Wick, "A Stroll through the Gaussian Primes" (American Mathematical Monthly, 105: 327-337 (1998)) discusses this Gaussian moat problem and proves that steps of length $< \sqrt{26}$ are insufficient. Their result was improved to $\sqrt{36}$ in 2005. My question is: Is the analogous question easier for the prime spiral (a.k.a. Ulam spiral)-Can one walk to infinity using bounded-length steps touching only the spiral coordinates of primes? What little I know of prime gaps suggest that should be easier to walk to infinity. For example, the first gap of 500 does not occur until about 10^{12} (more precisely, 499 and 303,371,455,241). I ask this primarily out of curiosity, and have tagged it 'recreational.' Edit1. In light of Gjergji's remarks below, I have tagged this as an open problem. Edit2. Just for fun, I computed which primes are reachable on a small portion of the spiral, for step distances $d \le 3$ (left below) and $d \le 4$ (right below); red=reached, blue=not reached. The former does not reach 83, the 23rd prime blue dot barely discernable at spiral coordinates (5,-3); the latter does not reach 5087, the 680th prime blue dot at spiral coordinates (36,10).

论坛问题帖 | has_answers | Joseph O'Rourke | 2010-09-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018277 | MO-381214

Dense sets in $\Bbb{R}^2$ with rational distance

We call a subset $S\subset \Bbb{R}^2$ rationally distanced if all $s_1,s_2 \in S$ have rational Euclidean distance. The Erdos-Ulam conjecture asks if there is a dense subset of $\Bbb{R}^2$ which is rationally distanced. Inspired by this, I came up with the following toy problem. A subset $X \subset \Bbb{R}^2$ is good if there exists a rationally distanced $S \subset \Bbb{R}^2\setminus \overline{X}$ such that each $x\in X$ is a limit point of S. I have three questions: Is $[0,1]\times \{0\}$ good? If $[0,1] \times \{0\} $ is not good, does this imply a negative solution to the Erdos-Ulam conjecture? (i.e. can we show that no rationally distanced subset S of $\Bbb{R}^2$ can intersect all possible line segments in $\Bbb{R}^2$) Is there an infinite good set that is not a subset of a straight line? (to avoid trivial examples like $\{n+\sqrt{2}:n\in \Bbb{Z}\}\times \{0\}$ for which $\Bbb{Q}\times\{0\}$ is a solution) Edit: Question 2 is straight-forward. Obviously, for S to intersect each of the line segments $\{[0,1]\times \{\alpha\}:\alpha \in \Bbb{R}\}$, we must have S be uncountable. Originally I saw this much, but didn't know how to deal with the distance function not being injective. For S to be dense in $\Bbb{R}^2$, there must distinct $s_1,s_2,s_3 \in S$ which are not collinear. We then have that the map $\phi:x \mapsto (|x-s_1|,|x-s_2|,|x-s_3|)$ is a injection from $\Bbb{R}^2$ to $\Bbb{R}^3$. The uncountability of S then immediately gives that $\phi(S) \not\subseteq \Bbb{Q}^3$, hence there exists some $i \in \{1,2,3\}, s^* \in S$ such that $|s_i-s^*|$ is irrational, making S not rationally distanced. Edit 2: Wojowu pointed out that since there are rationally distanced subsets of the unit circle, we can get infinite good sets that are not subsets of a line. So Question 3 as stated here is resolved in the positive. I'm still interested to see other constructions, but I will ask about that in a future post, after I've collected my thoughts.

论坛问题帖 | no_answers_at_retrieval | Zach Hunter | 2021-01-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018278 | MO-38356

What heuristic evidence is there concerning the unboundedness or boundedness of Mordell-Weil ranks of elliptic curves over $\Bbb Q$?

Some experts have a hunch that for any nonnegative integer r there are infinitely many elliptic curves over $\Bbb Q$ with Mordell-Weil rank at least r. The best empirical evidence for this hunch can be found in Andrej Dujella's tables here and here, the strongest of this evidence being provided by Elkies using constructions involving K3 surfaces. The only heuristic evidence I know of supporting this hunch is the results of Tate and Shafarevich (strengthened by Ulmer) that the Mordell-Weil ranks of elliptic curves over the a fixed function field $\Bbb F_q(t)$ can be arbitrarily large. What other heuristic evidence (if any) is there that the ranks of elliptic curves over $\Bbb Q$ should be unbounded? Seeing as how some experts do no believe this conjecture, I'd also accept answer to the companion question: What heuristic evidence (if any) is there that the ranks of elliptic curves over $\Bbb Q$ should be uniformly bounded?

论坛问题帖 | has_answers | James Weigandt | 2010-09-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018279 | MO-383855

Is it a reasonable way to write a research article assuming truth of a conjecture?

I have found a conjecture in a research article (published in a good journal) on number theory, which is not well known but very reasonable. Let me be clear that, there is no counter-example that vote down the conjecture, rather, its trueness has been proved in some special situation so far. I need this conjecture to develop a tool in order to write a research article. Unfortunately, I am unable to prove the conjecture as of now. Maybe others can prove it after some time. My question: Is it a reasonable way to write a research article assuming truth of the conjecture?

论坛问题帖 | has_answers | MAS | 2021-02-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018280 | MO-384170

Status of a conjecture of Hirzebruch

I was reading a paper from 1994 which claimed that the following statement was a conjecture of Hirzerbruch: If a complex surface X is homeomorphic to either $S^2 \times S^2$ or $\mathbb{C}P^2 \# \overline{\mathbb{C}P^2}$ then it is biholomorphic to one of the Hirzerbruch surfaces. I would like to know if this conjecture has been resolved or what the current status about it. Also, I would like to know the state of the art about the variant of the conjecture in which "homeomorphic" is replaced by "diffeomorphic": If a complex surface X is diffeomorphic to either $S^2 \times S^2$ or $\mathbb{C}P^2 \# \overline{\mathbb{C}P^2}$ then is it biholomorphic to one of the Hirzerbruch surfaces? If this has been proved, what is a reference?

论坛问题帖 | has_answers | cr1t1cal | 2021-02-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018281 | MO-38632

Projective Plane of Order 12

I asked this question on the new Theoretical Computer Science "overflow" site, and commenters suggested I ask it here. That question is here, and it contains additional links, which I doubt I can embed here because I don't have enough reputation. Anyway, here goes: Objective: Settle the conjecture that there is no projective plane of order 12. In 1989, using computer search on a Cray, Lam proved that no projective plane of order 10 exists. Now that God's Number for Rubik's Cube has been determined after just a few weeks of massive brute force search (plus clever math of symmetry), it seems to me that this longstanding open problem might be within reach. I'm hoping this question can serve as a sanity check. The Cube was solved by reducing the total problem size to "only" 2,217,093,120 distinct tests, which could be run in parallel. Questions: There have been special cases of nonexistence shown (again, by computer search). Does anyone know, if we remove those and exhaustively (cleverly?) search the rest, if the problem size is on the order of the Cube search? (Maybe too much to hope for that someone knows this....) Any partial information in this vein?

论坛问题帖 | has_answers | Aaron Sterling | 2010-09-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018282 | MO-391898

Cramer-Castillon problem like

Special case of Golden ratio as a property of conic section (is it known?) as follows: Let ABC be arbitrary triangle and DEF is the its tangential triangle. Let CF meets AB at G and BE meets AC at H, let GH meets the circumcircle at I. A line through I and parallel to EF meets AB, AC at J, K then $$\frac{JK}{JI}=\frac{\sqrt{5}+1}{2}$$ Open problem: Cramer-Castillon problem being very difficult problem. Related to the Cramer-Castillon problem configuration, I posed a problem as follows: Let ABC be arbitrary triangle and let three collinear points E', F', D' such that $\frac{E'F'}{E'D'}=k$ where k is positive real number. How can construct points D in the circumcircle of $ \triangle ABC$, E in AB, F in AC such that D, E, F are collinear and $\frac{EF}{ED}=k$?

论坛问题帖 | has_answers | Đào Thanh Oai | 2021-05-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018283 | MO-3929

Infinite dimensional Newlander-Nirenberg theorem

The Newlander-Nirenberg theorem states that an almost complex structure is integrable if and only if the Nijenhuis tensor vanishes. I heard that this statement is not true in infinite dimensions, since for example the Loop space of a Riemannian 3-manifold is counterexample. (In fact, I think NN fails for Fréchet manifolds in general(?)) So my question is: Is the Newlander Nirenberg theorem valid for Banach- or Hilbertmanifolds? If not, is it possible to weaken the statement (or some conditions) such that it remains true for some class of infinite dimensional manifolds? EDIT: Added the tag "open-problem", since NN for Hilbertmanifolds seems to be an open problem.

论坛问题帖 | has_answers | Spinorbundle | 2009-11-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018284 | MO-392994

Factoring integers of the form $n=p q^2$ using elliptic curves

We got argument and strong experimental support that integers of the form $n=p q^2$ can be factored using elliptic curves easier than general integers Q1 Is this known? Added This is known since at least 2002: Factoring N = p q^2 with the Elliptic Curve Method Q3 Can the method be improved for general integers? Assume E is elliptic curve modulo n with known point P of the order of P over $\mathbb{F}_p$ is $\rho = S U$ where S is B_1 smooth. We can find the factor p working over the curve modulo n in time $O(B_1 + \sqrt{U})$ and constant memory. It is an open problem if this can be generalized to arbitrary n. Stage 1: This is the first stage of the elliptic curve factoring algorithm. Set $Q=P$ and for primes r, $r \le B_1$ let $Q=r^{\log_r{n}} Q$. If we reach the point at infinity modulo p we are done and the algorithm stops with factor p. Stage 2: This is based on the birthday paradox and Pollard's rho algorithm for discrete logarithms in groups. Let $\mathrm{kronecker}(a,n)$ denote the kronecker symbol and for a point a on an elliptic curve $X(a)$ denote the X coordinate of a. For points a,b on E define the function $f(a,g)$: If $\mathrm{kronecker}(X(a),n)=1,f(a,g)=2 a + 5 g$, else $f(a,g)=3 a + g$. For Q from stage 1, set $a=Q,b=Q,g=Q$. For n from 1 to B_2 set $a=f(a,g),b=f(b,g),b=f(b,g)$. If $\gcd(X(a)-X(b),n) \bmod n>1$ return the factor. Stage 2 finds multiple of $U Q$ in time B_2 while $U \sim B_2^2$. Since the kronecker symbol is multiplicative, we have $\mathrm{kronecker}(a,n)=\mathrm{kronecker}(a,p)$ for a coprime to n. Iterates of $f^k(a,g)$ are deterministic random walk modulo p, so $f^N(a,g)=f^{2N}(a,g)$ happens for $N = O(\sqrt{U}) \sim B_2$.

论坛问题帖 | no_answers_at_retrieval | joro | 2021-05-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018285 | MO-393993

0's in 81591528324789773434561126959611589427200000000

Is 40 the largest number for which all the 0 digits in the decimal form of $n!$ come at the end? Motivation: My son considered learning all digits of 40! for my birthday. I told him that the best way to remember them would be to come up with a mnemonic. He asked me what word he should come up with if the digit is 0. The rest is history.

论坛问题帖 | has_answers | domotorp | 2021-05-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018286 | MO-394427

Is the unit-stick number of a knot equal to its stick number?

Define the unit-stick number $\sigma_1(K)$ of a knot K to be the fewest unit-length sticks that can realize K. Clearly $\sigma_1(K)$ is at least the stick number $\sigma(K)$. It is known that the trefoil knot K_{3_1} has stick number 6; see the image below. I believe, although I don't have a formal proof, that $\sigma_1(K_{3_1})=6$: the trefoil also has unit-stick number 6. (Wikipedia image by Wiebew. The sticks are not all the same length.) Q. Is $\sigma_1(K) = \sigma(K)$ for every knot K? If not, what is a knot for which it can be proved that the unit-stick number is strictly greater than the stick number, $\sigma_1(K) > \sigma(K)$? Related: Which knots' stick numbers are twice their crossing numbers?. Update. User @aorq's comment shows that my "unit-stick number" has been studied under the name "equilateral stick number": Rawdon, Eric J., and Robert G. Scharein. "Upper bounds for equilateral stick numbers." Contemporary Mathematics 304 (2002): 55-76. These authors proved that indeed the trefoil has equilateral stick number 6. They identify the 8_{19} knot as possibly one whose equilateral stick number is greater than its stick number. So, about 20 years ago, it appears that the question I posed was an open problem.

论坛问题帖 | no_answers_at_retrieval | Joseph O'Rourke | 2021-06-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018287 | MO-394507

Must Mersenne numbers be divisible by arbitrary large primes with exponent one?

Let M_n denote the Mersenne numbers $M_n=2^n-1$. As n varies, must M_n be divisible by arbitrary large prime p with exponent one, i.e. $p \mid M_n, p^2 \nmid M_n$? In other words, must the exponential diophantine equation $2^n-1 = A x^2 y^3$ for constant A has only finitely many solutions (n,x,y) and $n >1$? Related question is this. Also related question that might show there are infinitely many non-Wieferich primes is here. Looking for unconditional results, abc easily implies it. Added The paper Remarks on Exponential Congruences and Powerful Numbers P. RIBENBOIM on p7. (M') There exist infinitely many Mersenne numbers which are not powerful. (M') implies B_2 implies infinitely many non-Wieferich primes. One easy way to construct infinitely many non-powerful Mersenne numbers is to observe that 3 divides M_{6n+2} with exponent one, so M_{6n+2} is not powerful. Doesn't this approach give infinitely many non-Wieferich primes?

论坛问题帖 | has_answers | joro | 2021-06-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018288 | MO-395973

Conjectures or Results?

There is a paper (not accepted for publication yet) that contains several conjectures. Some of these conjectures were proven recently. The referee of the original paper requires to substitute the proven "Conjectures" with the "Results". However, there are several papers that cite these conjectures, so I feel it would be wrong to rename them. What is the best (or standard) way to indicate in the original paper that the conjectures were proven in the subsequent papers? Are there any good examples of doing that?

论坛问题帖 | has_answers | Sasha | 2021-06-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018289 | MO-396004

Simple example of Hammerstein integral equation

I'm currently reading this paper (and working on a similar one). The main goal is to study the Hammerstein integral equation (in $\mathcal{C}(I,E)$: $$x(t) = \int_{0}^{t} K(t,s)f\big(s,x(s)\big)ds,\quad t\in I;$$ where $I=[0,1]$, $K $ is a scalar kernel, E a Banach space and $f:I \times E \rightarrow E$ is a given function. Let's say that in my hypothesis, I need: H_1. There exist $a:I \rightarrow (0,+\infty)$ with $a(.)\in L^{\infty}(I)$ and $b:[0,+\infty) \rightarrow(0,+\infty)$ a nondecreasing function such that $\|f(s, x)\| \leq a(s) b\big(\|x\|\big)$ for a.e. $s \in I$ and $x\in E$, H_2. there exists at least one solution $r \in \mathcal{C}\big(I,(0, \infty)\big)$ to the inequality $$ b\left(\|r\|_{\infty}\right) \int_{0}^{t} \left |K(t, s) \right |a(s) d s \leq r(t), \quad t \in I$$ where $\|.\|_{\infty}$ is the sup norm in $\mathcal{C}\big(I,(0, \infty)\big)$. But I'm wondering if those are reasonable assumptions. I'm looking for an example of the Hammerstein integral equation with the hypothesis (H_1) and (H_2), take $\color{blue}{E=\mathbb{R}}$ for simplification.

论坛问题帖 | has_answers | Motaka | 2021-06-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018290 | MO-3965

Minimal surface in a ball

Assume a minimal surface Σ has boundary on the unit sphere in the Euclidean space and r is the distance from Σ to the center of the ball. Is it true that $$\mathop{\rm area} \Sigma\ge \pi\cdot(1-r^2).$$ Comments: The problem is solved in all dimensions and codimension, see "Area bounds for minimal..." by Brendle and Hung in 2016. (Thanks Rbega for the ref.) If $r=0$, the statement follows directly from the monotonicity formula. If Σ is topological disc the answer is YES, see answer of Oleg Eroshkin below. The general question is formulated as a conjecture in 1975 --- see comment of Ian Agol. There is an analog in all dimension and codimension for area minimizing surfaces, see Alexander, H.; Hoffman, D.; Osserman, R. Area estimates for submanifolds of Euclidean space. 1974.

论坛问题帖 | has_answers | Anton Petrunin | 2009-11-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018291 | MO-397159

Model theory and dynamical system (open problems)

I am curious about the open problems which are between model theory and dynamical system. I mean the open problems that are interesting for both groups and there are some evidences showing there might be a nice interaction between model theory and dynamical systems. Thank you! Any references would be appreciated!

论坛问题帖 | no_answers_at_retrieval | Alex Israel | 2021-07-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018292 | MO-400809

Inscribed square and convexity

Let $b(X)$ be the boundary of any X subset of the plane. Does there exist A,B convex compact sets of the plane, such that $C:=A\setminus B$ is simply connected and not empty, and such that there is no square inscribed in $b(C)$ that has a vertex in $b(C)\setminus B$? If the answer is no, this would kind of mean that the Toeplitz conjecture is kind of linked to convexity (one of the first intersting resolved cases was the case of the boundary of a convex) If the answer is yes... then it suggests a "fractal" way to built some counterexample. Do not hesitate to say your "prognosis" in the comments, even if it is not argued so far, I have absolutally no intuition myself about the pronostic (I would say it must be "NO" but I would be influenced by the fact that according to me, if it is "YES", then there is a counterexample not far, and this would have been found - but I might as well underevaluate the difficulty of "YES=> counterexample" ...)

论坛问题帖 | no_answers_at_retrieval | jcdornano | 2021-08-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018293 | MO-401018

The 4th Lemoine circle

The first and second Lemoine circles are well-known to geometers. According to this article the third Lemoine circle has been first discovered by Jean-Pierre Ehrmann in 2002. It is worth noting that the centres of these circles belong to a line that is going through the Lemoine point and the circumcenter of the triangle. I would like to present the construction of another circle, which I believe might be called the 4th Lemoine circle. The idea behind its construction is slightly similar to that proposed by Ehrmann: Symmedians AA', BB', CC' intersect each other at the point M (the Lemoine point of the triangle ABC). The circumcircle of the triangle A'B'C' was drawn (also known as the 'symmedial circle'). A'', B'', C'' are the first intersection points of the symmedians with the symmedial circle. Finally, the circumcircles of the triangles A''B''M, B''C''M, A''C''M always intersect the sides of the original triangle ABC at six points, that are conclycic. Theorem illustration As far as I can tell the point O (the center of our six point circle) is not included into the C.Kimberling's Encyclopedia, so it must be unknown or uncatalogued. O belongs to the line that contains the centres of the first three Lemoine circles, so by my reckoning, the dotted red circle that is shown in the picture above perfectly qualifies for being called the 4th Lemoine circle. Could you please give a synthetic proof of this theorem ? Geogebra dynamic sketch.

论坛问题帖 | has_answers | A.Zakharov | 2021-08-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018294 | MO-401348

Planar subsets with many pairs of points on distance 1

Let X be a subset of $\mathbb R^2$ consisting of n distinct points. Let $d_1(X)$ be the number of pairs of points of X on distance 1 from each other. Define $$d_1(n)=\sup_{X\subset \mathbb R^2|, |X|=n}d_1(X).$$ In particular $d_1(1)=0$, $d_1(2)=1$, $d_1(3)=3$, $d_1(4)=5$, etc. Question. I wonder what is known about the asymptotic behaviour of $d_1(n)$ and about its upper bound? It should be at least $n\log(n)$ as the following example shows: Example. It is easy to see that $d_1(2^d)\ge d2^d$, indeed, one can take the set Y consisting of d unit vectors and then take 2^Y consisting of sums of vectors over all 2^d subsets of Y. Then for generic Y we have $|2^Y|=2^d$ and $d_1(2^Y)=d2^d$. By fiddling a bit with Y one can increase $d_1(2^Y)$.

论坛问题帖 | has_answers | aglearner | 2021-08-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018295 | MO-401358

Heuristics for the very little torsion in the cohomology of Shimura variety

Consider the following statement which is a part of Conjecture 1.3 in the paper titled "The asymptotic growth of torsion homology for arithmetic groups" authored by N. Bergeron and A. Venkatesh. Let G be a $\mathbb{Q}$-semisimple group of zero defect(includes the Shimura variety case) and let $K \subset G(\mathbb{R})$ be a maximal compact subgroup and $\Gamma \subset G(\mathbb{Q})$ be a congruence subgroup. Denote the locally symmetric space $\Gamma \backslash G(\mathbb{R})/K$ by X_{Γ}. Let Γ_N denote the descending family of congruence subgroups such that $\cap \Gamma_N = \lbrace 1 \rbrace$. Then $$ \lim_{N \to \infty} \frac{\log |\mathrm{H}^i(X_{\Gamma_N}, \mathbb{Z})_{tors}|}{[\Gamma : \Gamma_N]} = 0. $$ I was wondering what are heuristics for this conjecture? What are the evidence available for this conjecture? Is something known in the case of Shimura varieties? Any help of reference is appreciated. Thank you.

论坛问题帖 | no_answers_at_retrieval | random123 | 2021-08-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018296 | MO-401456

The 4th vertex of a triangle?

I was immensely surprised and amused by the idea of the fourth side of a triangle that was introduced by B.F.Sherman in 1993. 'Sherman's Fourth Side of a Triangle' by Paul Yiu is available here. Naturally a question arises whether it is possible to determine the '4th vertex' of a triangle in a somewhat similar way? Basically a vertex of the given triangle can be described as a point that is lying on its circumcircle. Then the middle point of the segment connecting this point to the Orthocenter must belong to the nine point circle of the triangle. (This holds true for any point that is lying on the circumcircle.) Finally it appears that we should just add another equation somehow linking our 'vertex' to the inscribed circle. In other words, there might exist some triangle center on the circumcircle that is also satisfying a certain condition related to the inscribed circle, so that eventually this point can be called the fourth vertex of the triangle. Luckily I found in my files a construction of a triangle center X that in a sense satisfies these conditions: It belongs to the circumcircle of ABC The midpoint of the segment XX(4) belongs to the nine point circle. The constriction of X primarily relies on the Incenter of the triangle (i.e. the inscribed circle) Last but not least, this point X is not included in Kimberling's encyclopaedia: A',B',C' is the circumcevian triangle with respect to the Incenter I. Lines AB and A'B' intersect at point C'', points B'', A'' are defined cyclically. Circumcircles for the triangles IRA, IRB, IRC were drawn. These 3 circles intersect the circumcircle at the points A''', B''', C'''. Finally A''A''',B''B''',C''C''' always cross each other at some point X that conveniently belongs to the circumcircle of the original triangle ABC. Geogebra dynamic sketch. It is highly speculative to call our point X the fourth vertex of a triangle, of course. Presumably this attribution will be outright dismissed or proven wrong. However I assume that perhaps a better fit for the 4th vertex of a triangle can be found? What would its construction be?

论坛问题帖 | has_answers | A.Zakharov | 2021-08-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018297 | MO-401841

Are there infinitely many n such that $n!-1$ and $n!+1$ are prime numbers?

Are there infinitely many n such that $n!-1$ and $n!+1$ are prime numbers?

论坛问题帖 | has_answers | Kai Gu | 2021-08-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018298 | MO-402310

Lemoine-Lozada circles

I made some rookie attempt to define the 4th Lemoine circle recently. The alternative name for this circle was suggested yesterday. Further investigation revealed a family of circles associated with the Brocard axis. Supposedly the name of Ehrmann-Lozada circles (or Lemoine-Lozada circles) can be safely attached to them. Specifically, it appears that for each Lozada circle it is possible to construct at least one circle of the Lemoine type, with its center on the line that is going through the Symmedian point and the circumcenter of the triangle. step 1: Let A'B'C' be the midheight triangle of ABC and let BA and CA be the orthogonal projections of B' and C' on BC, respectively. Build AB, CB, AC and BC similarly. These last six points lie on a circle with center X(9729). The circle is called the 1st Lozada circle. (César Lozada, March 17, 2016) step 2: Circumcircles of the triangles BcAcX(6), AbCbX(6), BaCaX(6) cut the sides of the original triangle at six points, that define the 1st Ehrmann-Lozada circle. Its center O_1 is not in the ETC. Ehrmann-Lozada circles can be constructed in the similar manner for each(?) of the ten remaining Lozada circles:
https://www.geogebra.org/geometry/pg9mrmzx 2 new circles of the Lemoine type
https://geogebra.org/geometry/uqnpqjeh 2 new circles of the Lemoine type
https://www.geogebra.org/geometry/t4jpn9dp only 1 new circle of the Lemoine type ... 11. https://www.geogebra.org/m/kppjdcny gives us only 1 new circle of the Lemoine type... More links will be added soon. None of these points O_i is included in Clark Kimberling's encyclopaedia. Finding a strict proof of existence of these new circles is the main question here. Bit I am also interested in what makes Lozada points/circles so special? (Taylor point belongs to the Brocards Axis as well, but for some reason it is impossible to construct an Ehrmann-Taylor circle in the same way!) Are these Ehrmann-Lozada circles also Tucker circles?

论坛问题帖 | has_answers | A.Zakharov | 2021-08-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018299 | MO-403955

A special case of Frankl's conjecture. A question about known results

Let's recall a Frankl's conjecture. Consider a finite family of finite sets $\mathcal{F}$, such for every pair of sets $A\in \mathcal{F}$ and $B\in\mathcal{F}$, we have $A\cup B\in\mathcal{F}$(the family $\mathcal{F}$ is said to be union closed). The conjecture says that there is an element $a\in \bigcup_{A\in\mathcal{F}}A$, so that $a\in A$ for at least $\frac{|\mathcal{F}|}{2}$ sets A. Now, let's take a set $U=\{1,2,...,n\}$ and an union closed family $\mathcal{F}'\subseteq\mathcal{P}(U)$, where $\{U,\emptyset\}\cup\mathcal{P}_{n-1}(U)\subseteq\mathcal{F}'$. Is it known whether this case is solved? I would like to see a proof of this specific case. I was trying to figure it out myself by considering an induction in respect to $|U|$, but i failed. Thank you for any help or hint.

论坛问题帖 | no_answers_at_retrieval | mkultra | 2021-09-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018300 | MO-404209

Is there an even number a such that $a^{2^{n}}+1$ is prime for infinitely many n?

Is there an even number a such that $\{n: a^{2^{n}}+1 \text{ is prime} \}$ is an infinite set? Let a be even. Is there infinitely many n such that $a^{2^{n}}+1$ is composite?

论坛问题帖 | has_answers | LMP | 2021-09-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018301 | MO-405177

Open problems in symbolic dynamics

I would like to know which are some noticeable open problems in symbolic dynamics, including substitution dynamics. I'm especially interested in connections with topological chaos of various forms. Thanks!

论坛问题帖 | has_answers | user393632 | 2021-09-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018302 | MO-406096

On improving the upper bound $I(m^2) \leq \frac{2p}{p+1}$, if $p^k m^2$ is an odd perfect number with special prime p

(Preamble: This question is an offshoot of this answer to an MSE question with the same title.) Denote the classical sum of the divisors of the positive integer x by $\sigma(x)=\sigma_1(x)$ and the abundancy index of x by $I(x)=\sigma(x)/x$. Let $p^k m^2$ be an odd perfect number with special prime p satisfying $p \equiv k \equiv 1 \pmod 4$ and $\gcd(p,m)=1$. Since we have $$I(m^2) \leq \dfrac{2p}{p+1}$$ and because $p^k m^2$ is perfect, then we have $$I(p^k)I(m^2) = I(p^k m^2) = 2$$ where we have used the fact that the abundancy index function is multiplicative. Hence, we obtain (via iteration) $$\text{First iteration: } I(m^2) \leq I(p^k)I(m^2)\cdot\dfrac{p}{p+1} \leq 2I(p^k)\cdot\dfrac{p^2}{(p+1)^2}$$ $$\text{Second iteration: } I(m^2) \leq \bigg(I(p^k)\bigg)^2 {I(m^2)} \cdot\dfrac{p^2}{(p+1)^2} \leq 2\bigg(I(p^k)\bigg)^2 \cdot\dfrac{p^3}{(p+1)^3}$$ $$\ldots$$ $$\ldots$$ $$\ldots$$ by recursively replacing 2 with $I(p^k)I(m^2)$ and then bounding $I(m^2)$ from above by $2p/(p+1)$. Note that, at the n^{th} iteration, we have the inequality $$I(m^2) \leq 2\bigg(I(p^k)\bigg)^n \cdot \dfrac{p^{n+1}}{(p+1)^{n+1}}.$$ Repeating the process ad infinitum, we get $$I(m^2) \leq \lim_{n \rightarrow \infty}{2\bigg(I(p^k)\bigg)^n \cdot \dfrac{p^{n+1}}{(p+1)^{n+1}}} = \lim_{n \rightarrow \infty}{\dfrac{2\bigg(I(p^k)\bigg)^n}{\dfrac{(p+1)^{n+1}}{p^{n+1}}}}$$ $$= \dfrac{2p}{p+1}\cdot\lim_{n \rightarrow \infty}{\dfrac{\bigg(I(p^k)\bigg)^n}{\bigg(I(p)\bigg)^n}},$$ where the limit is of the indeterminate form $$\dfrac{+\infty}{+\infty}.$$ Hence, we may apply L'Hôpital's rule, by considering $$h(p) = \dfrac{\bigg(I(p^k)\bigg)^n}{\bigg(I(p)\bigg)^n}$$ as a function of p. (Also, note that there is "currently no L'Hôpital's rule for multiple variable limits". The closest thing that I could find on arXiv is this preprint by Gary R. Lawlor.) I tried working on applying L'Hôpital's rule to $h(p)$, by asking for help from WolframAlpha, however the

computations are a mess. Here is my inquiry: QUESTION: Is it possible to simplify the computations, via say, some recursive rule that I do not know about?

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2021-10-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018303 | MO-406311

A counterexample to a conjecture of Lawson

Yau quotes Lawson as having formulated the following conjecture [1]: Let M be an embedded minimal surface in $\mathbf{S}^3$. Prove that the two domains in $\mathbf{S}^3$ divided by M have equal volume. This is easily seen to hold for the simplest two minimal surfaces in $\mathbf{S}^3$, namely the equatorial spheres and the Clifford tori; it was also known at the time that the analogous statement fails in higher-dimensional spheres. (Whether it holds for the Lawson surfaces in $\mathbf{S}^3$ I do not know.) Question. I remember reading a statement claiming that some surface had been found to contradict the proposed statement. Does somebody know which surface this is? Unfortunately I cannot seem to remember where I read this, and online searches are obscured by Lawson's other, more prominent conjecture. [1] Shing-Tung Yau. Problem section. Seminar on Differential Geometry. 102 (1982) pp.669-706.

论坛问题帖 | has_answers | Leo Moos | 2021-10-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018304 | MO-40725

Base locus of divisors on blowings up of the projective space

Let X be the blowing-up of $\mathbb{P}^n$ in r points in general position. Let $\{H,E_1,...,E_r\}$ be the standard basis of $Pic(X)$. Further let $D=dH-\sum m_j E_j$, be an effective divisor, with $m_j \in \mathbb{Z}$. Is it true that if $m_1\geq 0$ then $E_1 \not\subseteq Bs(|D|)$? If we suppose all the m_i's are non negative, this corresponds to say that $dim |dH-m_1E_i...-m_rE_r|>dim |dH-(m_1+1)E_1-...-m_rE_r|$. In other words this means that if we consider all the hypersurfaces of $\mathbb{P}^n$ of degree d passing through r general points $\{p_1,...,p_r\}$ with multiplicities $m_1,...,m_r$, then they do not all pass through p_1 with multiplicity m_1+1. My intution brings me to say this is obvious but I cannot prove it. In fact everything is trivial for nonspecial linear systems, but, a priori, special linear systems may be a problem. Note that the generality of the points is necessary: for example you can simply take $\mathbb{P}^2$ blown up in 3 collinear points and note that $E_1 \subseteq Bs(|H-E_2-E_3|)$.

论坛问题帖 | has_answers | Gianni Bello | 2010-10-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018305 | MO-408481

An open triangle problem in plane geometry

Some years ago, I asked some 'famous' people in an advanced Plane Geometry forum about the following: Let ABC be arbitrary triangle, how can one construct a point P in the plane such that P is the circumcenter of the cevian triangle of P with respect to $ABC?$ The answer was negative, even in the sense of calculations by a Computer (can't construct by rule and compass, even can not calculations by computer). My conjecture: Let ABC be arbitrary triangle, then there exist a point P such that P is the circumcenter of the cevian triangle of P with respect to ABC. Question: How can prove conjecture and how construct this point?

论坛问题帖 | has_answers | Đào Thanh Oai | 2021-11-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018306 | MO-413332

Distinct exponents in the factorization of the factorial, a problem of Erdős

In the 1982 paper below, Paul Erdős proved that if $h(n)$ is the number of distinct exponents in the prime factorization of $n!$ then $$c_1\Big(\frac{n}{\log n}\Big)^{1/2} < h(n) < c_2\Big(\frac{n}{\log n}\Big)^{1/2}$$ for all sufficiently large n, where c_1, c_2 are positive constants. Then he said that "there is no doubt" that $$h(n) = (c + o(1))\Big(\frac{n}{\log n}\Big)^{1/2}$$ as n goes to infinity, for some positive constant c, but that "we do not know enough about the difference of consecutive primes" to conclude so. I wonder if now, after 40 years, "we" know enough to prove Erdős conjecture (?) P. Erdős, Miscellaneous problems in number theory, Congressus Numerantium 34 (1982), 25-45. https://users.renyi.hu/~p_erdos/1982-08.pdf

论坛问题帖 | has_answers | sebaztian | 2022-01-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018307 | MO-41358

Special values of Artin L-functions

This question might be naive and might carry the heuristic that we are living in the best possible world a little too far. If so, I appreciate being told so. Background: Stark's conjecture interprets the leading coefficients of Artin L-functions around $s=0$ as certain regulators. Let F/K be a Galois extension of number fields and ρ an Artin representation of the absolute Galois group of K factoring through F. Let $L(F/K,\rho,s)$ be the Artin L-function attached to ρ and let $c(\rho)$ be the leading coefficient of its Taylor expansion around $s=0$. Then Tate's formulation of Stark's conjecture does the intuitive thing and "picks out the ρ-isotypical bit" of the units $\mathcal{O}_F^\times$ and measures its volume in a suitable sense. But there are various problems with this intuition and it doesn't quite work the way one would like to (see below). The result is that one only gets a complex number Reg(ρ) that is well defined up to a rational multiple. The conjecture is that this complex number accounts for the transcendental part of $c(\rho)$, more precisely that Reg$(\rho)/c(\rho)$ is algebraic and that Galois acts on this quotient through its action on the Artin characters. There is a different circle of conjectures, due to Beilinson, Bloch, Burns, Flach, Fontaine, Kato and Perrin-Riou, subsumed under the bracket of "Tamagawa number conjectures", which gives exact predictions for special values of much more general motivic L-functions. Some of these conjectures work up to units, but some of them give actual

numbers. The regulators involved are rather more complicated motivic things and, being a hands-on person, I find it difficult to discern the arithmetic in them (I know, this is embarassing to admit for someone who works on Galois module structures). Question: Does anyone know of an explication of the Tamagawa number conjectures in the setting of Artin L-functions, which would interpret the exact value of $c(\rho)$ (possibly up to units) in terms of the objects that one intuitively expects to enter into such a formulation? Special cases would already help. In an ideal world, I would like a formulation, which makes it visible how the "ρ-isotypical parts" of the units and of the class numbers are picked out and how one measures the volume of a suitable lattice in the ρ-isotypical part of the units. Two extra demands that I would have of such a formulation are It should be easily visible that for trivial ρ, we recover the analytic class number formula over K and that the product of these conjectures for all irreducible ρ that factor through F, raised to the powers dim(ρ), gives the analytic class number formula over F. If you don't know of a reference, could you give a hint, how one goes about picking the right sublattice of the units? Difficulties: I see two big difficulties in what I am asking for, so if you can't answer the full question, maybe you can address the difficulties one by one or give me examples of special cases where one or the other has been solved? Firstly, if ρ is not defined over $\mathbb{Q}$, then it's not at all clear, how to define the right volume on the sublattice we pick out. My naive picture is that volumes are determinants of some non-degenerate pairing. If I restrict a non-degenerate pairing to the ρ-component of a lattice (I'm happy to base change the lattice to the ring of definition of ρ, even at the risk of losing some information) and ρ is not self-dual, then the pairing will become degenerate, so the determinant that I would expect in the formulation of a precise Stark conjecture would be 0. Secondly, even if all Artin representations are defined over $\mathbb{Q}$, e.g. is the Galois group is C_2, then although it is easy to say what we mean by the 1-isotypical component and the -1-isotypical component of a lattice, the product of the volumes of these will almost never be the volume of the whole lattice, since their direct sum will have finite non-trivial index in the whole lattice. Many thanks in advance!

论坛问题帖 | no_answers_at_retrieval | Alex B. | 2010-10-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018308 | MO-415982

Status of finite-dimensional Ando's theorem

Szőkefalvi-Nagy's theorem says the following: if A is a contraction on a Hilbert space H, then there exists a unitary U on a Hilbert space $H'\supset H$ for which $A^n=P_HU^nP_H$ for all $0\le n$. (P_H denotes projection onto H.) If H is finite-dimensional and we only need this to be true for $0\le n\le N$ then this can be modified such that H' is also finite-dimensional (cf. Levy and Shalit - Dilation theory in finite dimensions: the possible, the impossible and the unknown). Ando's theorem (see e.g. Levy-Shalit) says the following: if A and B are commuting contractions on H, then there exist commuting unitaries U and V on $H'\supset H$ for which $A^nB^m=P_HU^nV^mP_H$ for all $0\le n,m$. The construction in the proof is naturally infinite-dimensional even if H is finite-dimensional so a natural question (Problem C in Levy-Shalit) is: if we only need this to be true for $0\le n+m\le N$ then does an analogous modification work to make H' finite-dimensional? Levy-Shalit make this out to be wide-open, as of 2011. My question here is if there are any recent developments in the years since then on this or related problems.

论坛问题帖 | has_answers | zjs | 2022-02-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018309 | MO-41676

Milnor's isotopy invariant using spectral sequence?

I'm reading stalling's article "the augmented ideal in group ring" in Ann. Math. Studies 84(R. H. Fox memorial volume) In his final remark, he says that Milnor's link invariant could be interpreted by using Spectral sequence.(see Milnor, Isotopy of links, Algebraic geometry and topology, Princeton press) Are there anybody who knows about further advances in this story??

论坛问题帖 | has_answers | Topologieee | 2010-10-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018310 | MO-420253

$\mathbf{P} = \mathbf{NP}$, what's the problem?

Let's take the problem of the backpack: $A_1,\ldots ,A_n$ the weights that are integers, and we want to know if we can achieve a total weight of V. We take $$I=\dfrac{1}{2\pi}\int_0^{2\pi} \exp(-iVt)\times (1+\exp(iA_1t))\times\cdots\times(1+\exp(iA_nt)) \, dt.$$ The question then becomes what $I=0$ or $I\geq 1$. Can we not approach the value of I with the method of Monte Carlo, or others? Why are these approaches not succeeding? Remark : it's not difficult to find a good approximation of $B\times t \bmod 2\pi$, where B is a big integer and $t\in [0,2\pi]$, because we know excellent approximation of π.

论坛问题帖 | has_answers | Dattier | 2022-04-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018311 | MO-421151

Resources where I can find open problems in number theory along with their level of difficulty

NOTE: I will not accept an answer because a lot of answers are really good and if anyone want to post under this question later then they are most welcome to post as comment or answer because it will certainly help many people. Thanks to all that contributed. I have completed my master's in mathematics a couple of years ago and due to very strong personal and professional reasons I couldn't get admitted to grad school despite having a good academic background. I have a really good background in number theory. I don't have guidance of any professor right now and I want to try working on an open problem. Can you please let me know of resources(websites/ blogs/books) where I can find open problems in number theory to work on along with their estimated level of difficulty (if possible)? Thanks!

论坛问题帖 | has_answers | Arnold | 2022-04-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018312 | MO-422876

Source for conjectures in commutative algebra

Do you know some books/survey papers/ websites on conjectures or open problems in commutative ring theory? I want to see if there are very famous open problems or conjectures in commutative ring theory. Thank you.

论坛问题帖 | no_answers_at_retrieval | user 1 | 2022-05-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018313 | MO-42352

Midpoint geodesic polygon / Birkhoff curve shortening

I would like to know under what conditions the process of creating a midpoint piecewise geodesic polygon converges on a surface $S \subset \mathbb{R}^3$. S may be assumed smooth, closed, and oriented; or it may be assumed a Riemannian manifold. Let P be a closed geodesic polygon each of whose finite number of edges is a shortest path connecting its endpoint vertices v_i. Moreover, the edges of P are "short" in a sense specified below. P need not be simple, i.e., it may intersect and cross itself. Define the midpoint polygon $M(P)$ for P to have vertices u_i, each the midpoint of the geodesic segment $v_i v_{i+1}$ of P, where u_i is connected to u_{i+1} by a shortest path. I would like the segment $u_i u_{i+1}$ to be the unique shortest path between those points. So assume that $|u_i u_{i+1}|$ is smaller than the injectivity radius of the manifold. (This might require adding new vertices to ensure this condition holds for the next iteration.) In the plane, iterating this midpoint polygon construction is well studied. On a surface (or Riemannian manifold) it plays a role in curve shortening, in particular Birkhoff curve shortening, which he used to prove the existence of a simple closed geodesic [B27]. Subsequently the technique has many uses. For example, Christopher Croke used Birkhoff shortening to find the length of a shortest closed geodesic on a sphere [C88]. The uses I have seen of Birkhoff shortening are on the sphere, not on more general surfaces. Which brings me to the question: Q. Does Birkoff shortening, repeatedly applied to a geodesic polygon on a closed surface S, always converge (perhaps to a point)? If so, always to a closed geodesic? If not, under what conditions might it not converge? References. [B27] George D. Birkhoff, Dynamical Systems, AMS, 1927. p.135ff. [C88] Christopher B. Croke, "Area and the length of the shortest closed geodesic." J. Differential Geom., Volume 27, Number 1 (1988), 1-21.

论坛问题帖 | has_answers | Joseph O'Rourke | 2010-10-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018314 | MO-424613

Mark Hovey's open problems in the theory of model categories

Mark Hovey maintains a list of open problems in model category theory. I think this list is quite old, and I don't know if Hovey is still updating it or not. My question is: i) which of the 13 problems in that list are still considered open and still important in homotopy theory? ii) or dually, which of the problems are considered settled or are perceived as no longer relevant given the current status of homotopy theory?

论坛问题帖 | has_answers | user139854 | 2022-06-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018315 | MO-425498

Convex polygon shadows: Shortest equivalent segments

Let P be a convex polygon. Q1. What is the shortest collection of line segments S inside P with the property that both P and S have the same sequence of orthogonal shadows as P and S rotate? An orthogonal shadow is cast by parallel light rays. For example, for P a unit-square, the X below has length $2 \sqrt{2}$, less than the perimeter 4 of P, but casts the same shadows: every lightray that intersects ∂P intersects S; every lightray that intersects S intersects ∂P: Q2. Is S always connected? Clearly it must span the vertices of P. Q3. Is S the medial axis of P? Medial axis. Fig.5.1(b) from Discrete and Computational Geometry. I feel I should know or be able to find the answers to these questions, but I am not hitting on the appropriate search terms. My real interest is in $\mathbb{R}^3$, but first let's explore $\mathbb{R}^2$. Answered by Gerry Myerson: Q1: Open problem Q2: Not necessarily connected Q3: No

论坛问题帖 | has_answers | Joseph O'Rourke | 2022-06-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018316 | MO-425925

The "Smallest" open Diophantine Equation, a potential approach

In this question, Bogdan Grechuk found the "Smallest" open Diophantine equation, where size is determined by taking absolute values of all coefficients, then substituting 2 into all variables. The equation is $y(x^3-y)=z^3+3$. There are a couple of different reformulations of this problem, for example "Is there any two numbers whose sum is a cube and whose product is three more than a cube?" However, one direction where I have begun to make progress is to reframe the equation as: $yx^3=z^3+y^2+3$. Taking this equation modulo 27, it can be shown that (proof on request): 1) y is not 0, 1, or 8 mod 9 (note in particular this forces y to not be a cube) 2) If y is divisible by 3, then y is 6 or 21 mod 27 Let v be the square-free part of y^2+3. Modulo v, the equation becomes $yx^3=z^3$, so either v divides x and z, or $y=(\frac zx)^3$. In the former case, this implies $v^3|y^2+3$, which seems extremely unlikely due to how v was defined (though I'm not sure how to prove that). In the latter case, we have a third condition on y: 3) y must be a cube mod v. I have tested all values of y up to 1000 so far, and no values of y satisfy all three conditions. So my question is: Can it be proven that there is no integer y which satisfies all three constraints? And a follow up question for the extra case: Can y^2+3 be a 3-Powerful number?

论坛问题帖 | no_answers_at_retrieval | Thomas Blok | 2022-07-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018317 | MO-426277

Stronger conjectured inequality for area of a polygon

Four years ago, I proposed an inequality related to area and sides of a polygon. After computer checking, I conjecture that the previous inequality can be strengthened as follows: Let $A_1A_2\cdots A_n$ be an arbitrary polygon, then:

$$4(\tan\frac{\pi}{n})Area(A_1A_2\cdots A_n) \le \sum_{i=1}^n(A_iA_{i+1})^2-\frac{1}{n-2}\sum_{i=1}^n(A_iA_{i+1}-A_{i+1}A_{i+2})^2$$ Question: Can you prove this conjectured inequality? PS: I also found that for an equiangular convex pentagon, $Area(ABCDE)=\frac{1}{4}(\cot\frac{\pi}{5})(a^2+b^2+c^2+d^2+e^2-\frac{(a-b)^2+(b-c)^2+(c-d)^2+(d-e)^2+(e-a)^2}{\sqrt{5}})$

论坛问题帖 | no_answers_at_retrieval | Đào Thanh Oai | 2022-07-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018318 | MO-426401

Resources where one can find open problems in Analytic Number Theory

Edit: I have asked this question separately because I am in need of some resources which focus on analytic number theory only.So, this question is not a duplicate of:Resources where I can find open problems in number theory along with their level of difficulty as I am in looking for websites/books/blogs in Analytic Number Theory only. I don't have much knowledge of other branches of number theory. I have completed my master's in mathematics a couple of years ago from a good university of my country but due to very strong personal and professional reasons I couldn't get admitted to grad school despite having a good academic background. I have a really good background in analytic number theory. I have studied 1 course in Sieve Theory, 2 courses which were based on 2 volumes of Apostol's Number Theory books. I will start a course in Automorphic Forms soon. I don't have guidance of any professor right now and I want to try working on an open problem in Analytic Number Theory. Can you please let me know of resources(websites/ blogs/books) where I can find open problems in analytic number theory to work on? Since, I don't have a guidence of a prof. due to various reasons, these resources will really help me. Thank you!

论坛问题帖 | has_answers | Arnold | 2022-07-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018319 | MO-427595

A conjecture on consistent monotone sequences of polynomials in Bernstein form

A Conjecture In the following, a polynomial $P(x)$ is written in Bernstein form of degree n if it is written as- $$P(x)=\sum_{k=0}^n a_k {n \choose k} x^k (1-x)^{n-k},$$ where $a_0, ..., a_n$ are the polynomial's Bernstein coefficients. The degree-n Bernstein polynomial of an arbitrary function $f(\lambda)$ has Bernstein coefficients $f(k/n)$. In general, this Bernstein polynomial differs from f even if f is a polynomial. Let $f(\lambda):[0,1]\to(0,1)$ have a continuous r-th derivative, where $r\ge 1$, and let M be the maximum of the absolute value of f and its derivatives up to the r-th derivative. Let $W_{2^0}(g; \lambda), W_{2^1}(g; \lambda), ..., W_{2^i}(g; \lambda),...$ be a sequence of operators that map a continuous function g on $[0, 1]$ to a bounded function on $[0, 1]$ and converge uniformly to f. Prove or disprove the following conjecture. For each integer $n\ge 1$ that's a power of 2, suppose that there is $D>0$ such that- $$\text{abs}(f(\lambda)-B_n(W_n(f; \lambda); \lambda)) \le DM/n^{r/2},$$ whenever $0\le \lambda\le 1$. Then there is $C_0\ge D$ such that the polynomials (g_n) in Bernstein form of degree 2, 4, 8, ..., 2^i, ..., defined as $g_n=B_n(W_n(f; \lambda); \lambda) - C_0 M/n^{r/2}$, converge from below to f and satisfy: $(g_{2n}-g_{n})$ is a polynomial with nonnegative Bernstein coefficients once it's rewritten to a polynomial in Bernstein form of degree exactly $2n$. Equivalently (see also Nacu and Peres (2005)), there is $C_1>0$ such that the inequality- $$\sum_{i=0}^n \left(W_{2n}\left(f; \frac{k}{2n}\right) - W_n\left(f; \frac{i}{n}\right)\right)\sigma_{n,k,i}\ge -C_1 M/n^{r/2},\tag{PB}$$ holds true for each integer $n\ge 1$ that's a power of 2 and whenever $0\le k\le 2n$, where $\sigma_{n,k,i} = {n\choose i}{n\choose {k-i}}/{2n \choose k}=\mathbb{P}(X_k=i)$ and X_k is a hypergeometric($2n$, k, n) random variable. C_0 or C_1 may depend on r and the sequence (W_n), but not on f, λ, or n. When C_0 or C_1 exists, find a good upper bound for it. It is also conjectured that the same value of C_0 (or C_1) suffices when f has a Lipschitz continuous $(r-1)$-th derivative and M is the maximum absolute value of f and the Lipschitz constants of f and its derivatives up to the $(r-1)$-th derivative. Background I asked this question in order to solve the so-called Bernoulli factory problem, described next. We're given a coin that shows heads with an unknown probability, λ. The goal is to use that coin (and possibly also a fair coin) to build a "new" coin that shows heads with a probability that depends on λ, call it $f(\lambda)$. This is the Bernoulli factory problem, and it can be solved only if f is continuous (Keane and O'Brien 1994). However, since I asked this question I have found a Bernoulli factory algorithm that I believe is general enough to cover all the cases that this conjecture would help solve. Since this conjecture may be of broader interest, though, I leave this question open. See also my other open questions about the Bernoulli factory problem. Special Cases The conjecture may be trivial to prove if the convergence rate of W_n is $O(1/n^{r/2})$. It may also be easy to prove if W_n or $B_n(W_n)$ preserves polynomials of degree $(r-1)$ or less. The conjecture is not so easy to prove in many other cases (e.g., when W_n converges arbitrarily slowly to f). Results Here are results that correspond to the conjecture. If $r=1$ and $W_n=f$, then $C_0 = (1+\sqrt{2})$ (Nacu and Peres 2005), even in the Lipschitz case. If $r=2$ and $W_n=f$, then $C_0 = 1/2$ (Nacu and Peres 2005). If $r=2$ and $W_n=f$, then $C_0 = 1/7$ for every $n\ge 4$ (see "Proofs for Polynomial Building Schemes"). An example for $r=3$ and $W_n=2 f - B_n(f)$*, suggests $C_1 = 0.3078$. In the linked page, it's stated that it's useful to find raw and central moments of hypergeometric random variables, and to find images of λ^i ($k=2n\lambda$) under the left-hand side of (PB), if that left-hand side is a linear operator. References Keane, M. S., and O'Brien, G. L., "A Bernoulli factory", ACM Transactions on Modeling and Computer Simulation 4(2), 1994. Nacu, Şerban, and Yuval Peres. "Fast simulation of new coins from old", The Annals of Applied Probability 15, no. 1A (2005): 93-115. C.S. Güntürk, W. Li, "Approximation of functions with one-bit neural networks", arXiv:2112.09181 [cs.LG], 2021. Micchelli, C. (1973). The saturation class and iterates of the Bernstein polynomials. Journal of Approximation Theory, 8(1), 1-18. Butzer, P.L., "Linear combinations of Bernstein polynomials", Canadian Journal of Mathematics 15 (1953). * Corresponds to the iterated Boolean sum of order 2 (Güntürk and Li 2021).

论坛问题帖 | no_answers_at_retrieval | Peter O. | 2022-07-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018320 | MO-42902

The determinant of the sum of normal matrices

Given two normal matrices $A,B\in M_n({\mathbb C})$ whose respective spectra are $(\alpha_{1},\ldots,\alpha_{n})$ and $(\beta_{1},\ldots,\beta_{n})$, is it true that $\det(A+B)$ belongs to the convex hull of the set of numbers $$\prod_{i=1}^n(\alpha_i+\beta_{\sigma(i)}),$$ as σ runs over the set ${\mathfrak S}_n$ of permutations of $\{1,\ldots,n\}$? Nota. It is known (see Exercise 101) that the trace of AB belongs to the convex hull of the set of numbers $$\sum_{j=1}^n\alpha_{j}\beta_{\sigma(j)},\qquad\sigma\in {\mathfrak S}_n.$$

论坛问题帖 | has_answers | Denis Serre | 2010-10-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018321 | MO-430424

What is the current status on bad tangent cones at isolated singularities?

Let $M^8 \subset B^9 \setminus \{ 0 \} \subset \mathbf{R}^9$ be a properly embedded, stable minimal hypersurface. Suppose that $0 \in \overline{M}$ is an isolated singularity of the surface. Question. (i) Are the tangent cones at the origin regular, that is $\operatorname{sing} \mathbf{C} = \{ 0 \}?$ (ii) If one cone in $\operatorname{VarTan}(\lvert M \rvert,0)$ is regular, is there a 'direct' way of showing that they must all be regular? (By 'direct' I would mean without going via uniqueness of tangent cones à la Lojasiewicz-Simon.) I believe (i) used to be an open question, but I am curious about its current status or partial progress. The Schoen-Simon regularity theory implies that the cones are either regular or cylindrical of the form $\mathbf{C}^7 \times \mathbf{R}$. Correction. Otis Chodosh pointed out in his answer that this is incorrect. The aforementioned regularity theory of Schoen-Simon shows that the singular set of every tangent cone $\mathbf{C} \in \operatorname{VarTan}(\lvert M \rvert,0)$ is one-dimensional, but it can consist of several rays, one for each singularity in the link of the cone. It's the tangent cones taken at these rays that are cylindrical, of the form described above.

论坛问题帖 | has_answers | Leo Moos | 2022-09-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018322 | MO-430672

Are there any open sourced or crowd source math research projects?

I’m looking for open source math research projects as I know computer science students can easily have access to material they can contribute to but this is not the case with math as far as I’m aware. But I am looking for a possible answer with open source or crowd sourced math research projects that someone in their last year of getting a bachelors or during a Masters could contribute to. Given they have the ability that is.

论坛问题帖 | has_answers | Jayden Maryott | 2022-09-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018323 | MO-430830

Conjecture about the equality : $f(y)=y\ln(y)+\sum_{n=2}^{\infty}\pm\frac{\left(y\ln y\right)^n}{2^{a_n}}$

I try here because I expect I cannot have any answer on MSE : Problem : Let : $$f\left(x\right)=\frac{\left(1+\ln27\right)x!\ln x!}{x+1},g\left(x\right)=x\ln x$$ Then it seems $\exists y\in(0,1)$ and a_n be an infinite set of positive integer such that : $$f(y)=y\ln(y)+\sum_{n=2}^{\infty}\pm\frac{\left(y\ln y\right)^n}{2^{a_n}},g(x)=x\ln(x)+\sum_{n=2}^{\infty}\pm\frac{\left(x\ln x\right)^n}{2^{a_n}},f'(y)-g'(y)=0$$ For consolidate my conjecture see : $$H\left(x\right)=x\ln x+\frac{x^{2}\ln^{2}\left(x\right)}{4}-\frac{x^{3}\left(\ln x\right)^{3}}{8}+\frac{x^{4}\left(\ln x\right)^{4}}{4\cdot8}-\frac{x^{5}\left(\ln x\right)^{5}}{4\cdot32}+\frac{x^{6}\left(\ln x\right)^{6}}{4\cdot16}-\frac{x^{7}\left(\ln x\right)^{7}}{4\cdot64}+\frac{x^{8}\left(\ln x\right)^{8}}{2048}+\frac{x^{9}\left(\ln x\right)^{9}}{2048\cdot4}-\frac{x^{10}\left(\ln x\right)^{10}}{2048\cdot2}-\frac{x^{11}\left(\ln x\right)^{11}}{2048\cdot2}-\frac{x^{12}\left(\ln x\right)^{12}}{2048\cdot8}-\frac{x^{12}\left(\ln x\right)^{12}}{2^{22}}-\frac{x^{13}\left(\ln x\right)^{13}}{2^{24}}+\frac{x^{14}\left(\ln x\right)^{14}}{2^{24}}-\frac{\left(1+\ln27\right)x!\ln\left(x!\right)}{x+1}$$ Then : $$|H\left(0.35925959\right)|<3\cdot10^{-14},|H'\left(0.35925959\right)|<6\cdot10^{-8}$$ For this conjecture I have two question (if true): How to show the convergence ?If yes : How to find the a_n ? Thanks everyone ! Ps: for $\pm$ it's not $(-1)^n$ as you can see for order $8,9$ I cannot go further with Desmos Software but look at the following partial sum : $$H\left(x\right)=x\ln x+\frac{x^{2}\ln^{2}\left(x\right)}{4}-\frac{x^{3}\left(\ln x\right)^{3}}{8}+\frac{x^{4}\left(\ln x\right)^{4}}{4\cdot8}-\frac{x^{5}\left(\ln x\right)^{5}}{4\cdot32}+\frac{x^{6}\left(\ln x\right)^{6}}{4\cdot16}-\frac{x^{7}\left(\ln x\right)^{7}}{4\cdot64}+\frac{x^{8}\left(\ln x\right)^{8}}{2048}+\frac{x^{9}\left(\ln x\right)^{9}}{2048\cdot4}-\frac{x^{10}\left(\ln x\right)^{10}}{2048\cdot2}-\frac{x^{11}\left(\ln x\right)^{11}}{2048\cdot2}-\frac{x^{12}\left(\ln x\right)^{12}}{2048\cdot8}-\frac{x^{12}\left(\ln x\right)^{12}}{2^{22}}-\frac{x^{13}\left(\ln x\right)^{13}}{2^{24}}+\frac{x^{14}\left(\ln x\right)^{14}}{2^{24}}+\frac{x^{15}\left(\ln x\right)^{15}}{2^{24}}-\frac{x^{16}\left(\ln x\right)^{16}}{2^{25}}+\frac{x^{17}\left(\ln x\right)^{17}}{2^{26}}+\frac{x^{18}\left(\ln x\right)^{18}}{2^{26}}+\frac{x^{19}\left(\ln x\right)^{19}}{2^{26}}-\frac{\left(1+\ln27\right)x!\ln\left(x!\right)}{x+1}$$ Where : $x=0.35925959\cdots$ Update 08/10/2022 : We have a similar problem with the Binet formula slighlty modified see : $$g\left(x\right)=\sqrt{2\pi x}\left(\frac{x}{e}\right)^{x}e^{\frac{47000000000}{116593560186976815022080000x^{13}}+\frac{4388413659}{9716130015581401251840000x^{12}}+\frac{310}{13494625021640835072000x^{11}}+\frac{390}{86504006548979712000x^{10}}+\frac{800}{514904800886784000x^{9}}+\fra

c{1}{86684309913600x^{8}}+\frac{2}{902961561600x^{7}}+\frac{7}{75246796800x^{6}}+\frac{1}{836075520x^{5}}+\frac{2}{24883200x^{4}}-\frac{1}{10368x^{3}}-\frac{1}{291x^{2}}+\frac{1}{12x}}-x!$$ And $g(0.285)$

论坛问题帖 | has_answers | DesmosTutu | 2022-09-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018324 | MO-431769

Open problems in type theory

I am only a beginner in the field of type theory, and I'm wondering if the community could point me out a few open problems in the field. I have a good background in logic, in particular, proof theory and I have a reasonable knowledge on category theory. While I am interested in homotopy type theory, I'd like to receive answers that point to flavors of type theory other than HoTT because open problems in HoTT are not difficult to find on the web (problems in HoTT are still welcome though). This is my very first question in this site so please feel free to edit, point mistakes and criticize my question. Thank you for your attention!

论坛问题帖 | has_answers | Guimarães | 2022-10-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018325 | MO-431810

Approximating spectra of (finite rank pertubations of) Laurent operators by spectra of (pertubations of) periodic finite operators

A tridiagonal matrix is a matrix which only has elements on three diagonals. So for $\alpha, \beta, \gamma \in \mathbb{C}$ consider the bi-infinite tridiagonal Laurent operator T with $\beta $ on the diagonal given by. \begin{pmatrix} \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \dots & \alpha & \beta & \gamma & 0 & 0 & 0 & \dots \\ \dots & 0 & \alpha & \beta & \gamma & 0 & 0 & \dots\\ \dots & 0 & 0 & \alpha & \beta & \gamma & 0 & \dots\\ \dots & 0 & 0 & 0 & \alpha & \beta & \gamma & \dots \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ \end{pmatrix} General theory tells us that T is invertible if and only the symbol curve (which is in this case is an ellipsis) given by $ \{ z \in \mathbb{T} \mid \frac{\alpha}{z} + \beta + z \gamma \} $ does not enclose 0. In finite dimensions, we can consider the periodic version T_n. For example T_5 looks as follows: \begin{pmatrix} \beta & \gamma & 0 & 0 & \alpha \\ \alpha & \beta & \gamma & 0 & 0 \\ 0 & \alpha & \beta & \gamma & 0 \\ 0 & 0 & \alpha & \beta & \gamma \\ \gamma & 0 & 0 & \alpha & \beta \\ \end{pmatrix} Now, when n tends to ∞ then the spectra of the periodic versions $\sigma(T_n)$ approximate the spectrum of the Laurent operator T. In the sense that in Haussdorff distance it holds that $$ \sigma(T_n) \to \sigma(T). $$ This can for example be seen by Fourier Transform. Now, suppose that I add finitely many diagonal terms. I.e. the matrix $K = \text{diag}(K_{1,1}, K_{2,2} , \dots, K_{m,m})$. Then it is a Theorem (see https://link.springer.com/article/10.1007/BF01275512, Corollary 1.3) that $$ lim_{n \to \infty} \left(\sigma(T_n + P_n K P_n) \cup \sigma(T) \right) \to \sigma(T+K). $$ Here P_n is the projection onto the n sites of T_n. But in general it is conjectured (https://link.springer.com/article/10.1007/BF01275512, Conjecture 7.3) that $$ lim_{n \to \infty} \left(\sigma(T_n + P_n K P_n) \right) \to \sigma(T+K). $$ Meaning that the finite system can also approximate the symbol curve as was the case without the pertubation. What is the status of this conjecture? And in the case $m=1$ where the pertubation is only a single entry, how could one go about proving it? The reason I am interested is given on page 18 in here https://arxiv.org/pdf/2206.09879.pdf, where one can see that these question can help answer to what extend the spectra of some infinite volume open quantum systems are approximated by their finite volume periodic counterparts.

论坛问题帖 | no_answers_at_retrieval | Frederik Ravn Klausen | 2022-10-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018326 | MO-432645

Finding an optimal covering trail for the set $\{0,1,2,3\}\times\{0,1,2,3\}\times\{0,1,2,3\}$

Here is a key question (i.e., Question 2 below) that, if correctly answered, would let me support a very general conjecture on a wide class of related problems, a conjecture that I have never shared before. In the Euclidean space $\ \mathbb{R}^3,\ $ let $$ \mathcal H(4,3)\ :=\ \{0,1,2,3\}\times\{0,1,2,3\}\times\{0,1,2,3\} \,\ \subseteq\,\ \mathbb R^3 $$ We consider the family of all the distinct straight lines that joins at least two elements of $\mathcal{H}(4,3)$ (i.e., we need to take into account all the lines that pass through two distinct points among the 64 given elements of the above mentioned cubic lattice). Now, let us call $\mathcal{I_2}(4,3):=\{S_1,S_2, \dots, S_{t-1}\, S_t\}$ the set of all the (distinct) intesections of the family of lines above. Question 1: Which is t, the cardinality of $\mathcal{I_2}(4,3)$? Question 2: If we check by brute force all the possible polygonal chains consisting of 21 or 22 line segments (how many?), whose vertices always belong to $\mathcal{I_2}$, will we find at least one covering trail for $\mathcal{H}(4,3)$ (i.e., is it possible to draw at most 22 consecutive links starting and ending on the vertices of $\mathcal{I_2}$ and joining in this way all the 64 elements of the set $\mathcal{H}(4,3)$?)? Current results: I have showed that we can constructively solve the problem by Question 2 with a polygonal chain of length 23 or less, 23 lines" /> whereas it is not possible to find any solution spending less than 21 lines (trivial lower bound)1. By improving my current upper bound, I would finally be able to strongly conjecture that the link length of any minimum-link covering trail for any set of $\{0,1,2,\dots,n-1\}\times\{0,1,2,\dots,n-1\} \times \dots \times\{0,1,2,\dots,n-1\}$ in $\mathbb{R}^k$ is equal to $\frac{n^k-1}{n-1}+n-3$ (here we are assuming that $n \in \mathbb{N}-\{0,1,2\}$ and $k \in \mathbb{N}-\{0,1\}$). Otherwise, if we will be able to show that it is not possible to improve my upper bound of 23 lines, I would infer that the general solution for the $\{0,1,2,\dots,n-1\}\times\{0,1,2,\dots,n-1\}\times \dots \times\{0,1,2,\dots,n-1\}\subset \mathbb{R}^k$ points problem is given by $\frac{n^k-1}{n-1}+(k-1)\cdot(n-3)$. I am persuaded that the 22 lines solution is likely to exist (or at least I strongly believe that it does not exist any covering trail consisting of only 21 links that covers $\mathcal{H}(4,3)$), but I would not have many arguments to support my general claim without confirming the above by brute force testing. Moreover, I have constructively proven that the link-length of the optimal covering trail for the set $\{0,1,2,3,4\}\times\{0,1,2,3,4\}\times\{0,1,2,3,4\} \subset \mathbb{R}^3$ belongs to $\{31, 32, 33, 34, 35, 36\}$, see the figure below 2. Related OEIS sequence: A318165. P.S. It is trivial to point out that all the S_i, for any $i=1,2,...,t$, necessarily belong to the

Axis-Aligned Bounding Box $[-6,9] \times [-6,9] \times [-6,9]$. Some useful References concerning the general problem: 1 - Proved untrivial bounds for the general problem) S. Bereg, P. Bose, A. Dumitrescu, F. Hurtado, and P. Valtr. Traversing a set of points with a minimum number of turns. Discrete & Computational Geometry, 41(4):513-532, 2009. 2 - Proof that my general conjecture holds for $k=2$) B. Keszegh. Covering paths and trees for planar grids. Geombinatorics Quarterly, 24(1):5-10, 2014. 3 - General problem under some simplifying constraints) E. Kranakis, D. Krizanc, and L. Meertens. Link length of rectilinear Hamiltonian tours in grids. Ars Combinatoria, 38:177-192, 1994. 4 - Conjectures in conclusion) M. Ripà, Minimum-Link Covering Trails for any Hypercubic Lattice, arXiv, 2022. 5 - Proof that my general conjecture holds for $n=3$) M. Ripà. Solving the 106 years old 3k points problem with the clockwise-algorithm. Journal of Fundamental Mathematics and Applications, 3(2):84-97, 2020. Announcement) We have also proven that it is possible to solve the general case ($n=2, k=k$) with exactly $3 \cdot 2^{k-2}$ lines and the full paper will be released in a couple of weeks.

论坛问题帖 | has_answers | Marco Ripà | 2022-10-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018327 | MO-433060

Conjecture on the unsolvability of the $\{3 \times 3 \times \cdots \times 3\} \subseteq \mathbb{R}^k$ dots problem starting from the central point

In 2020 (see Solving the 106 years old 3^k points problem with the clockwise-algorithm, JFMA, 3(2), p. 96), I conjectured that, in the Euclidean space $\mathbb{R}^k$, we can cover any given set of 3^k points $\mathcal{H}(3,k):=\{3 \times 3 \times \cdots \times 3\}\subseteq \mathbb{R}^k$ with an optimal polygonal chain $\mathcal{P}\left(\frac{3^k-1}{2}\right):=\rm{S}_1$-$\rm{S}_2$-$\rm{S}_3$-$\cdots$-$\rm{S}_{\left(\frac{3^k-1}{2}\right)}$-$\rm{S}_{\left(\frac{3^k+1}{2}\right)}$ having minimum link-lenght, whose two endpoints belong to $\mathcal{H}(3,k)$, if and only if $$\rm{S}_1 \not\equiv (1,1,\dots,1) \wedge \rm{S}_{\left(\frac{3^k+1}{2}\right)} \not\equiv (1,1,\dots,1).$$ I suspect that the above would not be so hard to be proven, since Theorem 1 by 1 and Theorem 2.1 by Minimum-Link Covering Trails for any Hypercubic Lattice imply that any optimal covering trail for $\mathcal{H}(3,k)$ has to be a sequence of consecutive (connected) line segments joining $\frac{3^k-3}{2}$ times 2 unvisited points of $\mathcal{H}(3,k)$ and only one time 3 new points (no exceptions). Conjecture: For any $k \in \mathbb{N}-\{0,1\}$, there does not exist any $\frac{3^k-1}{2}$ links (optimal) polygonal chain $\mathcal{P}\left(\frac{3^k-1}{2}\right):=\rm{S}_1$-$\rm{S}_2$-$\rm{S}_3$-$\cdots$-$\rm{S}_{\left(\frac{3^k-1}{2}\right)}$-$\rm{S}_{\left(\frac{3^k+1}{2}\right)}$ that visits all the points of the set $\mathcal{H}(3,k)$ and such that $\rm{S}_1 \not\equiv (1,1,\dots,1) \wedge \rm{S}_{\left(\frac{3^k+1}{2}\right)} \not\equiv (1,1,\dots,1)$. Now, since we need to start from the central point $C:=(1,1,\dots, 1) \in \mathbb{R}^k$ (otherwise, we could simply reverse the chosen sequence of the nodes of $\mathcal{P}$), we observe that the link $\rm{S}_1$-$\rm{S}_2$ visits no more that two points and the same thing must necessarily occur for $\rm{S}_2$-$\rm{S}_3$, but is that true up to $\rm{S}_{\left(\frac{3^k-1}{2}\right)}$-$\rm{S}_{\left(\frac{3^k+1}{2}\right)}$, for any $k \in \mathbb{N}-\{0,1\}$?

论坛问题帖 | no_answers_at_retrieval | Marco Ripà | 2022-10-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018328 | MO-433862

Do primes of the form $4k+1$ ever lead the greatest prime factor race?

Analogous to Chebyshev's race between primes, I examined the race between primes in the greatest prime factors, GPF, of natural numbers. Similar to the regular prime race, in the GPF race, the proportion of prime of both forms is roughly 50% each. However, unlike the regular prime race where the race is equal (A038691) or changes lead between primes of the form $4k+1$ and $4k+3$ (A096628) in the GPF race, $4k+3$ dominates. For $n \le 3 \times 10^{10}$, there are 14970975209 numbers the GPF is of the form $4k+1$ and 15029024755 numbers where GPF is of the form $4k+3$ and up till this point, there is not a single instance where primes of the form $4k+1$ equal or take the lead in the race. Let $f_1(n)$ and $f_3(n)$ be the number of natural numbers $\le n$ in which the GPF is of the form $4k+1$ and $4k+3$ respectively. The difference between the number of GPFs on these two forms appeared to increase in a very orderly manner as shown in the graph below. Question 1: Does the GPF race ever become equal or change the lead i.e. is there an n such that $f_1(n) = f_3(n)$? Clearly, the graph is sub-linear. I applied different curve fitting models to this data to have an idea of the growth rate. The best fit was given by $$ f_3(n) - f_1(n) \approx an^b $$ where $a \approx 0.1785$ and $b \approx 0.8122$. This experimental model has a very high $R^2 = 0.9999958$. If this is close to the true growth rate then, prime of the form $4k+1$ will always lag behind in the GPF race. Question 2: What is the true growth rate of $f_3(n) - f_1(n)$? Check out the interesting comments by Greg Martin (1 2) and Eric Naslund (1 2 3) in the linked question in MSE. Note: Was posted on MSE but was unanswered. Hence posting in MO.

论坛问题帖 | no_answers_at_retrieval | Nilotpal Kanti Sinha | 2022-11-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018329 | MO-434511

Is it still not known whether the construction of shortest nonzero vector of a lattice w.r.t. l^2-norm is NP-hard?

It was shown in P. van Emde Boas, Another NP-complete partition problem and the complexity of computing short vectors in a lattice that the construction of a shortest nonzero vector of a Euclidean lattice w.r.t. the L^{∞}-norm is NP-hard. But for the L^2 norm, is this question still open? Can anyone explain the difficulty and progress in this problem?

论坛问题帖 | has_answers | taylor | 2022-11-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018330 | MO-435725

Traveling salesperson problem algorithm

I was wondering something, let's say in a symmetric distance matrix of a sample of TSP, there was a sure algorithm that could remove between 30% to 80% of the values (distances) that wouldn't definitely be used in order to get the shortest routes in the end, would such algorithm be valuable and something new? It will definitely reduce many combinations of

routes but I don't think 80% would be always even close to enough to turn samples like a million cities into a P problem.

论坛问题帖 | has_answers | Ehsan Javanbakht | 2022-12-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018331 | MO-438258

Places where one can post open problems

(This must have been asked before and exist somewhere in Community Wiki, but I can't find it...) Where can you post open (math) problems? And what are the advantages and disadvantages? Example: This place (and Math StackExchange), duh. Example: The journal AMM had a corner "Unsolved Problems", but no longer. (Editor Stan Wagon told me that he doesn't know any math journal accepting unsolved problems.) Example: But then, probably any journal will accept a small "Open Questions" section after a research article. Example: USENET once had an useful group. The group still exists. (Yes, I'm that old...) Example: Other social media, e.g. Reddit has a math sub.

论坛问题帖 | has_answers | Hauke Reddmann | 2023-01-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018332 | MO-43923

Is there a 7-regular graph on 50 vertices with girth 5? What about 57-regular on 3250 vertices?

The following problem is homework of a sort -- but homework I can't do! The following problem is in Problem 1.F in Van Lint and Wilson: Let G be a graph where every vertex has degree d. Suppose that G has no loops, multiple edges, 3-cycles or 4-cycles. Then G has at least d^2+1 vertices. When can equality occur? I assigned the lower bound early on in my graph theory course. Solutions for $d=2$ and $d=3$ are easy to find. Then, last week, when I covered eigenvalue methods, I had people use them to show that there were no solutions for $d=4$, 5, 6, 8, 9 or 10. (Problem 2 here.) I can go beyond this and show that the only possible values are $d \in \{ 2,3,7,57 \}$, and I wrote this up in a handout for my students. Does anyone know if the last two exist? I'd like to tell my class the complete story.

论坛问题帖 | has_answers | David E Speyer | 2010-10-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018333 | MO-441001

Conjecture about primes and Fibonacci numbers

I posted this conjecture on math.stackexchange, but I received no answer proving or disproving it: if $ m > 4 $ is a positive integer not divisible by $ 2 $ or $ 3 $, it's ever possible to find a positive integer $ n $ such that the difference of the $ 2 $ Fibonacci numbers $ F_{m+n}-F_n $ is a prime number. Various example of $ [m, n] $, with the smallest n, are: $[5,1],[7,3], [11,4], [13,5], [17,3],..., [619,1353],...,[2011,6629]...$. The numerical evidence supporting that is quite good; I need to thank @Peter for his great findings searching for counterexamples. Here you'll find all the values of n for $m < 3000$ supporting the conjecture (the hardcases are at the bottom). Here the Oeis sequence about the difference of Fibonacci numbers being a prime I converted the primes found that way in the Zeckendorf representation and they are an alternating sequence of $ 1 $ and $ 0 $ ending with two adjacent $ 0 $ and $ 1 $ (and eventually just various $ 0 $). Looking for some known result, I found this article: "Primes as Sums of Fibonacci Numbers", I asked about this conjecture to one of the authors and this was the answer: "The conjecture can be stated as follows. First, for $ k≥1 $, we define K to be the string of length $2k-1$, starting with 1 and containing $1s$ and $0s$ alternatingly.Is there always a prime number p having the Zeckendorf expansion $K0010..0$? This would imply the "showcase result" of our paper, "for each large enough k, there is a prime number p that is the sum of exactly k different, non-adjacent Fibonacci numbers". I don't see a way how to prove this, as it amounts to finding a prime number in a very sparse subsequence (growing exponentially). In our paper the problem was easier, since there are "many" integers whose Zeckendorf expansion has length Ck (C some constant involving the golden ratio) and exactly k digits equal to 1." Do you have any idea?

论坛问题帖 | no_answers_at_retrieval | user967210 | 2023-02-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018334 | MO-441599

$\lim_{b \rightarrow \infty} {^{b}a} \in \mathbb{Q}_p$ for any $a \in \mathbb{Z}^+$?

$\newcommand\tetra[2]{{^{#1}{#2}}}$In a recent discussion on the Tetration Forum (see https://math.eretrandre.org/tetrationforum/showthread.php?tid=1703&page=2), it has been pointed out how my results on the constancy of the "congruence speed" of the integer tetration $\tetra b a$ (a peculiar property of hyper-4 described in The congruence speed formula), would imply that $\lim_{b \rightarrow \infty} \tetra b a$ is always an element of the set $\mathbb{Q}_p$. At the time, I did not fully realize the power of this breakthrough, since my original goal was only to find a function $V(a) : \mathbb{Z}^+-\{M\} \rightarrow \mathbb{Z}^+$, where $M = \{a : a \not\equiv 0 \pmod {10} \land a \neq 1\}$, but after a little search, I have seen that this is a general open problem, explicitly solved only for a few cases (since 1953, for $a:=2$). Furthermore, I have not managed to find any proof that even $\lim_{b \rightarrow \infty} \tetra b 6$ is a rational p-adic number (or rather an irrational one, disproving the aforementioned claim). Now, my first concern is if I am wrong and $\lim_{b \rightarrow \infty} \tetra b 6$ has already been proven to be an element of $\mathbb{Q}_p$ (or not), while my general request here is to know more about the implications of Equation 16 on page 15 of Ripà and Onnis - Number of stable digits of any integer tetration on the above mentioned general open problem (and related topics).

论坛问题帖 | has_answers | Marco Ripà | 2023-02-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018335 | MO-44234

Furstenberg's Conjecture on 2-3-invariant continuous probability measures on the circle

Hillel Furstenberg conjectured that the only 2-3-invariant probability measure on the circle without atoms is the Lebesgue measure. More precisely: Question: (Furstenberg) Let μ be a continuous probability measure on the circle such that $$\int_{S^1} f(z) d\mu = \int_{S^1} f(z^2) d\mu = \int_{S^1} f(z^3) d\mu, \quad \forall f \in C(S^1).$$ Is μ the Lebesgue measure? The assumption implies that the Fourier coefficients $\hat\mu(n)$ satisfy $$\hat\mu(n) = \hat \mu(2^k3^l n), \quad \forall k,l \in \mathbb N, n \in \mathbb Z.$$ Furstenberg's question is known to have an affirmative answer if one makes additional assumptions on the entropy of the measure. The basic strategy is usually to show that a non-vanishing (non-trivial) Fourier coefficient implies the existence of an atom. The standard tool to construct atoms in a measure on the circle is Wiener's Lemma, which says that as soon as there exists $\delta>0$ such that the set $\lbrace n \in \mathbb Z \mid |\hat\mu(n)| \geq \delta \rbrace$ has positive density in $\mathbb Z$, μ has an atom. More precisely, the following identity holds: $$\sum_{x \in S^1} \mu(\lbrace x \rbrace)^2 = \lim_{n \to \infty} \frac1{2n+1} \sum_{k=-n}^n |\hat \mu(k)|^2.$$ Clearly, $$|\lbrace 2^k3^l \mid k,l \in \mathbb N \rbrace \cap [-n,n] | \sim (\log n)^2 $$ so that Wiener's Lemma can not be applied directly. My question is basically, whether this problem can be overcome for any other subsemigroup of $\mathbb N$. Question: Is there any subsemigroup $S \subset \mathbb N$ of zero density known, such that every S-invariant continuous probability measure on S^1 is the Lebesgue measure? What about the subsemigroup generated by $2,3$ and 5?

论坛问题帖 | has_answers | Andreas Thom | 2010-10-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018336 | MO-445819

Normal curvature of Veronese embedding

Is it true that the Veronese embedding minimizes the maximal normal curvature among all smooth embeddings $\mathbb{R}\mathrm{P}^n\hookrightarrow \mathbb{S}^N$? Remarks The Veronese embedding $\mathbb{R}\mathrm{P}^n\hookrightarrow \mathbb{S}^{(n+3)\cdot n/2}$ describes a minimal surface. Its induced metric tensor $=2\times$(canonical metric on $\mathbb{R}\mathrm{P}^n$). It sends geodesics of $\mathbb{R}\mathrm{P}^n$ to circles with curvature $\kappa_n=\sqrt{\tfrac{n-1}{n+1}}$ in the sphere; in particular, its normal curvature κ_n at every point in every direction. The same question can be asked about $\mathbb{C}\mathrm{P}^n$ and $\mathbb{H}\mathrm{P}^n$. The value κ_n can be visualized the following way. Suppose d_n is the spherical distance between vertices of an inscribed equilateral n-simplex in $\mathbb{S}^{n-1}$. Then κ_n is the curvature of a spherical circle of radius $d_n/2$. Note that $d_n\to \tfrac\pi2$ and $\kappa_n\to 1$ as $n\to\infty$. I can solve the case $n=2$. Now it is written; any comments are welcome. In fact any submanifold with normal curvature $<\kappa_2=\tfrac1{\sqrt{3}}$ is diffeomorphic to the standard sphere. (In this case the curvature of the induced metric is $\tfrac14$-pinched; so it is sufficient to show that the submanifold is simply connected.) This question is from my paper (see 1.4), and it was motivated by Gromov's "Isometric immersions with controlled curvatures".

论坛问题帖 | no_answers_at_retrieval | Anton Petrunin | 2023-04-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018337 | MO-447472

Open problem: $\log n$ factor in Binomial empirical process

The following problem was considered in Cohen and Kontorovich, "Local Glivenko-Cantelli", https://arxiv.org/abs/2209.04054, to appear in COLT 2023 (henceforth, CK'23). Let Y_j, $j\in\mathbb{N}$ be a sequence of independent $\operatorname{Binom}(n,p_j)$ random variables, where $n\in\mathbb{N}$ and $1/2\ge p_j\downarrow 0$ as $j\to\infty$. Since $\mathbb{E} Y_j=np_j$, let us consider the centered, normalized process $\bar Y_j:= n{^{-1}} Y_j-p_j $. Finally, we define Δ_n to be the expected uniform absolute deviation: $$ \Delta_n := \mathbb{E}\sup_{j\in\mathbb{N}}|\bar Y_j|. $$ CK'23 gave an exact characterization of the $p\in[0,{\frac12}]^{\mathbb{N}}_{\downarrow0}$ (that is, $p\in[0,{\frac12}]^{\mathbb{N}}$ with $p_j\downarrow 0$) for which $\Delta_n\to0$ as $n\to\infty$. Namely, they defined the functional $$ T(p) := \sup_{j\in\mathbb{N}}\frac{\log (j+1)}{\log(1/p_j)} . $$ and showed that $\Delta_n\to0$ iff $T(p)<\infty$. They also gave the finite-sample bound $$ \Delta_n \le c\left(\sqrt{\frac{S(p)}{n}} +\frac{T(p)\log n}{n} \right), \qquad n\ge \mathrm{e}^3 \qquad(*) $$ where $c>0$ is an absolute constant and $$ S(p) := \sup_{j\in\mathbb{N}}p_j\log (j+1). $$ We conjecture that the $\log n$ factor in $(*)$ is superfluous. This conjecture is motivated by the asymptotic lower bounds $$ \liminf_{n\to\infty} \sqrt n \Delta_n \ge c\sqrt{S(p)}, $$ $$ \liminf_{n\to\infty} n \Delta_n \ge cT(p) , $$ where $c>0$ is a universal constant. This shows that beyond the conjecturally removable $\log n$ factor, $(*)$ is tight up to constants. Open problem. Is the $\log n$ factor in $(*)$ necessary, or can it be removed? UPDATE 21-Jul-2023. This was presented as an open problem at COLT'23: https://proceedings.mlr.press/v195/cohen23b.html Within days of the announcement and independently of each other, Moïse Blanchard, Václav Voráček, and Jaouad Mourtada obtained essentially the same substantial improvement. Formally, their results show that there is an absolute $c_1>0$ such that for any constant $c>0$ there is an n_0 such that for all $n>n_0$ there is a sequence $p\in[0,{\frac12}]^{\mathbb{N}}_{\downarrow0}$ for which $$ \Delta_n(p) > c\sqrt{\frac{S(p)}{n}} + c_1\frac{T(p)\log n}{n\log\log n}. \qquad(**) $$ This certainly dashes any hope of removing the $\log n$ factor in $(*)$ entirely; the best one could aim for is replacing it by $\log(n)/\log\log(n)$. This raises a number of fascinating questions. First: Is there a fixed sequence p_j (i.e., independent of n) for which $$ \liminf_{n\to\infty} \frac{n\log\log n}{\log n} \Delta_n \ge cT(p) , $$ where $c>0$ is a universal constant? Second: Is there a functional $\tilde T(p)$ and a monotonically increasing $\varphi:\mathbb{N}\to \mathbb{N}$ such that $$ c\tilde T(p) \le \liminf_{n\to\infty} \varphi(n) \Delta_n \le \limsup_{n\to\infty} \varphi(n) \Delta_n \le c'\tilde T(p) $$ for absolute constants $c,c'>0$? UPDATE 25-Jul-2023. We congratulate Moïse Blanchard on resolving the gap entirely, by exhibiting a p for which $$ \Delta_n(p) > c\sqrt{\frac{S(p)}{n}} + c_1\frac{T(p)\log n}{n} $$ (all constants and quantifiers as in $(**)$). Additionally, it has been pointed out to us by Moïse, Václav, Jaouad that the "lower bound" $ \liminf_{n\to\infty} n \Delta_n \ge cT(p) $ is trivial, since for nontrivial p, Δ_n will be at least $1/\sqrt n$ for n large enough, meaning that $\liminf_{n\to\infty} n \Delta_n =\infty$. We have

corrected this with the following non-trivial lower bound: Theorem (Cohen, K., 2023): There is an absolute constant $c>0$ such that $$ \Delta_n(p) \ge c\left(1\wedge\frac{T}{n}\right), $$ for all $n\in\mathbb{N}$ and $p\in[0,{\frac12}]^{\mathbb{N}}_{\downarrow0}$. Thus, currently, the two open questions are: Is there a fixed sequence p_j (i.e., independent of n) for which $$ \Delta_n(p) \ge c\left(1\wedge\frac{T\log n}{n}\right), $$ for some $c>0$? Is there a functional $\tilde T(p)$ and a monotonically increasing $\varphi:\mathbb{N}\to \mathbb{N}$ such that $$ \Delta_n \le c\left(\sqrt{\frac{S(p)}{n}} +\frac{\tilde T(p)}{\varphi(n)} \right) $$ and $$ \Delta_n(p) \ge c'\left(1\wedge\frac{\tilde T(p)}{\varphi(n)}\right), $$ for absolute constants $c,c'>0$ and all $n\in\mathbb{N}$, $p\in[0,{\frac12}]^{\mathbb{N}}_{\downarrow0}$?

论坛问题帖 | has_answers | Aryeh Kontorovich | 2023-05-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018338 | MO-450190

Books/blogs/websites that have open problems in Algebraic geometry

I got admitted in a PhD program in Europe last year. But due to serious mental health issues , I was deemed unfit by the mathematics department to continue the program. I am from a 3rd world nation and I got these problems there due to some issues I would not like to mention. So, I have decided to take a break and work on my health and to take a job to sustain myself before I apply again. The reason why I want to return to mathematics (in future years only not in current or upcoming year) because I love studying mathematics, solving exercises. I have a fine background in abstract algebra and algebraic geometry and I was wondering if you can let me know some books/ websites/ blogs which lists some open problems in Algebraic Geometry? Having to know some problems in algebraic geometry and current literature will help me to keep in touch with the field and a little motivated.

论坛问题帖 | has_answers | Arnold | 2023-07-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018339 | MO-450700

An open problem of Hardy and Littlewood on p-integral means

In Duren's book "Theory of H^p spaces" (MSN) in the comment section after Section 4, it is mentioned that Littlewood and Hardy proved in Some properties of conjugate functions that if u is a harmonic function in the unit disc belonging to the harmonic Hardy space h^p, $0<p\leq 1$. Then the harmonic conjugate of u, call it v satisfies $$ M_p(r,v):= \biggl(\frac{1}{2\pi} \int_{0}^{2\pi} \lvert v(re^{i\theta})\rvert^p d\theta \biggr)^\frac1p= O\bigl(\bigl(\log\frac{1}{1-r}\bigr)^\frac1p \bigr), \,\,\, r\to 1. $$ Furthermore they show that for $p=1,\frac12,\frac13,\dotsc$ the "big-O" cannot be improved to "little-o" by giving a concrete example. Duren then claims that this is an open question for other values of $p\in(0,1]$. Given that the book is written in the '70s I would like to know if this is still an open problem, or if there has been some progress.

论坛问题帖 | no_answers_at_retrieval | an_ordinary_mathematician | 2023-07-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018340 | MO-45177

What polynomials biject from $\mathbb{N}^{2}$ to $\mathbb{N}$?

Perhaps there are none with integral coefficients; so let us admit rational coefficients. The map $(x, y) \mapsto x + \frac{1}{2}(x + y)(x + y + 1)$ is well known, and swapping x and y in the formula yields another, so we have two for starters.

论坛问题帖 | has_answers | John Bentin | 2010-11-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018341 | MO-452403

Status of Problems in 102 problems in mathematical logic

Is there any location that records the current status of the problems in 102 problems in mathematical logic? Or, better yet, serves as a status board for open problems in mathematical logic? Mathoverflow is nice, but I'd leave it in a heartbeat for a variant that had an 'open problems' section which people could add problems to or mark them as solved as papers got published. I almost just embarrassingly claimed to solve Q62 while writing up my solution to this question even though I should have remembered Harrington solved it. Especially, as I agreed to write up beyond ω parts of his notes up at some point but haven't gotten to it yet.

论坛问题帖 | no_answers_at_retrieval | Peter Gerdes | 2023-08-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018342 | MO-452603

Different flavours of Vassiliev Conjecture

There is something that puzzles me about "Vassiliev's Conjecture". I am sure I am missing some detail which is obvious to the community, since there are several tightly related kind of results around Vassiliev invariants that I am not sure how to connect. Genuine Vassiliev's Conjecture [GVC] Let $\mathcal{K}$ be the space of knots in 3 dimensions. Let $\sim_n$ be the equivalence relation on $\pi_0(\mathcal{K})$ that identifies knots that cannot be distinguished by Vassiliev invariants of order $\le n$. We have a tower of surjections: $$ \pi_0(\mathcal{K}) \to \ldots \to \pi_0(\mathcal{K})/ \sim_n \to \pi_0(\mathcal{K})/\sim_{n-1} \to \ldots \to \pi_0(\mathcal{K})/ \sim_1 $$ The Genuine Vassiliev Conjecture states that two different knots can always be distinguished by a Vassiliev invariant. In other words, we require the injectivity of the maps $$ \pi_0(\mathcal{K}) \to \varprojlim_{n \to \infty} \pi_0(\mathcal{K})/\sim_n $$ Chord Vassiliev's Conjecture [CVC] A first line of results in this direction are given by things like Konsevitch Theorem. If $\mathcal{V}_n$ is the set of Vassiliev invariants of order $\le n$, Konsevitch proved that $$\mathcal{V}_n \otimes \mathbb{R} \cong \bigoplus\limits_{s \le n} A^*_s \otimes \mathbb{R} $$ where A_s is a suitable algebra of chord diagrams. There is a little trick we can do to reformulate Konsevitch theorem in terms of our tower. If we consider $\mathcal{V}_n \otimes \mathbb{Q}$ as a subspace of the dual of $\pi_0(\mathcal{K}) \otimes \mathbb{Q}$, we can write (see here): $$ (\pi_0(\mathcal{K})/ \sim_n) \otimes \mathbb{Q} \cong \frac{\pi_0(\mathcal{K}) \otimes \mathbb{Q}}{(\mathcal{V}_n \otimes \mathbb{Q})^{\perp}} \cong (\mathcal{V}_n \otimes \mathbb{Q})^* \cong \bigoplus_{s

\le n} A_s \otimes \mathbb{Q} $$ De Brito and Horel state that there exists a map $A_n \to \ker(\pi_0(\mathcal{K})/\sim_n \to \pi_0(\mathcal{K})/\sim_{n-1})$, where knots up to n-equivalence can be regarded as an abelian group under connected sum. When tensoring over $\mathbb{Q}$, it becomes an isomorphism; this fact, by an easy induction, yields the above formula for $\pi_0(\mathcal{K})/\sim_n$. Thus we will call the Chord Vassiliev's Conjecture the fact that $A_n \to \ker(\pi_0(\mathcal{K})/\sim_n \to \pi_0(\mathcal{K})/\sim_{n-1})$ is an isomorphism. Tower Vassiliev's Conjecture [TVC] At last, there are a few results linking the above tower with the taylor tower of knots. Specifically, Budney et al. proved that there exist maps $\pi_0(\mathcal{K})/\sim_n \to \pi_0(T_{n+1} \mathcal{K})$ that commutes with the tower maps. We call Tower Vassiliev's Conjecture (TVC) the fact that such a map is an isomorphism. Implications Let us denote by [CHL], [CCH], [CHT] the collapse at the second page of the homology, cohomology and homotopy spectral sequence associated with the taylor tower, at least over the antidiagonal. I want to prove some implications between the three vassiliev conjectures, eventually assuming some of the collapsing conditions. Here is what I have tried: [TVC] implies [GVC]. Since we have in isomorphism of towers, we can analogously show the injectivity of the map $$\pi_0(\mathcal{K}) \to \varprojlim_{n \to \infty} \pi_0(T_n \mathcal{K})$$, which is the composite of $\pi_0(\mathcal{K}) \to \pi_0(T_{\infty} \mathcal{K})$ with the projection $$\pi_0(T_{\infty} \mathcal{K}) \to \varprojlim_{n \to \infty} \pi_0(T_n \mathcal{K})$$ I suspect that these maps should be injective for a general reason, which I can't see. For example, it should be the case that $\pi_0(T_{\infty} \mathcal{K})$ is uncountable and $\pi_0(\mathcal{K})$ is countable, which makes injectivity plausible. Unluckily, it seems to me that the connectivity estimates of Goodwillie are not enough to conclude its injectivity. The injectivity of the other map should have to do with the relation between the total homotopy spectral sequence and the truncated ones (converging to $T_n \mathcal{K}$). It seems to me that both sides are given by $ \oplus_p E^{-p,p}_{\infty}$ where the latter is the homotopy spectral sequence; but π_0 of a homotopy limit is not always the limit of π_0, so I am definitely missing something. [TVC] implies [CVC]. The map reformulates as $ A_n \to \ker(\pi_0 T_{n+1} \mathcal{K} \to \pi_0(T_n\mathcal{K}))$. We know that in the homotopy spectral sequence $ A_n \cong E_{n+1,n+1}^2$ and also that $E^1_{n+1,n+1} = \pi_0(\textrm{fib}(T_{n+1}\mathcal{K} \to T_n \mathcal{K}))$. I suspect that using the long exact sequence for the fiber seq and analyzing a bit the spectral sequence should give the result, maybe up to assuming something else on the way. [TVC] implies [CHT]. This is stated explicitly by De Brito-Horel in theorem 6.1. [CCH] implies [CVC]. Indeed, Vassiliev invariants are $\oplus_{p} E^{-p,p}_{\infty}$, where the latter is the cohomology spectral sequence, while chord algebras pop up at the second page of Sinha Spectral sequence (first of Vassiliev) on the same line. Questions How many wrong things have I said? Can you fill in the gaps in my implications? Are there other implications that I am missing? Could you share your perspective on this matter? I feel like mine is somewhat goofy deformation of a solid one. How does the homology spectral sequence fit in this framework? Why has it been so intensively studied? Thank you. Of course, since it is a lot of material, partial answers are ok too.

论坛问题帖 | has_answers | Andrea Marino | 2023-08-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018343 | MO-454329

A question on Giles Gardam counter example to the Unit conjecture of Kaplansky

The unit version of the Kaplansky conjecture is about units in FG where F is a field and G is a torsion free group. In a recent counter example by Giles Gardam, it is given an example of a non trivial unit when $F=F_2$ is the field with 2 elements and G is a finitely presented torsion free group. See this MO post for some discussions on this counter example. The group G in Giles Gardam counterexample is a result of a group extension $Z^3\to G\to Z_2\times Z_2$. So it would be helpful to study some other extensions of Z^3 by this quotient group? Do they generate other examples?

论坛问题帖 | no_answers_at_retrieval | Ali Taghavi | 2023-09-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018344 | MO-455052

Initial conditions to falsify Rowland's conjecture

Based on the Rowland's paper (A natural prime-generating recurrence), is there any theorem to show that for which initial condition $a(1) = k$ the conjecture can be falsified? For example, for k equals 532 or 801, the conjecture of "only 1 or prime for the recurrent formula $g(n) = a(n)-a(n-1) = \gcd(n, a(n-1))$" does not hold, mentioned in the same paper. In sum, I am wondering if there is any theorem or conjecture to show some properties for (a subset of) such ks.

论坛问题帖 | no_answers_at_retrieval | OmG | 2023-09-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018345 | MO-45511

Density of values of polynomials in two variables

This question is a reposting of a comment I made on Polynomial representing all nonnegative integers. Let $f(x,y)\in \mathbb Q[x,y]$ such that $f(\mathbb Z\times \mathbb Z)$ is a subset of $\mathbb N$ (the nonnegative integers). Let $g(n)$ be the number of elements of $f(\mathbb Z\times \mathbb Z)\cap \lbrace 0,1,\dots, n\rbrace$. How fast can $g(n)$ grow? Is it always true that $g(n)=O(n/\sqrt{\log(n)})$? If true this is best possible since if $f(x,y)=x^2+y^2$ then $g(n)\sim cn/\sqrt{\log(n)}$.

论坛问题帖 | has_answers | Richard Stanley | 2010-11-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018346 | MO-455750

Has there been any progress on this open problem about co-well-poweredness of accessible categories?

On the relations between accessible categories and large cardinal axioms, one big example is the following: Assume the existence of a proper class of strongly compact cardinals. Then every accessible category is co-well-copowered. This can be found, for instance, in the standard reference Makkai-Paré (Acessible categories: The Foundations of Categorical Model Theory). We have Proposition 6.2.2: If pushouts are detectable in $\mathcal{K}$ (that is, the

category of diagrams which have pushouts is accessible), $\mathcal{K}$ is co-well-powered This also follows imitating the argument in Theorem 2.49 of Adamek-Rosicky (Locally presentable and accessible categories). The important result is: Theorem 6.3.8 (M-P): Assume there is a proper class of strongly compact cardinals. Then colimits are uniformly detectable in every accessible category, that is, the subcategory of $\mathrm{Diag}(A)$ (the total category) whose objects are diagrams admiting colimits is accessible. From there we can deduce detectability of pushouts and the result follows. Furthermore, every accessible category being co-wellpowered implies the existence of a proper class of measurable cardinals, as Example A.19 of Locally presentable and accessible categories shows with a counterexample using transitive models of ZC and a formal initial object, assuming there's a maximal measurable cardinal. There's probably an example in Makkai-Paré too but I don't recall any. Thus the large cardinal strength of the assertion "every accesible category is co-well-powered" lies somewhere between "there's a proper class of strongly compacts" and "there's a proper class of measurables", but as far as I know this is an open problem, and even though I by no means have made an exhaustive research on its status, I did some for my bachelor thesis (on the topic of accessible categories, AECs and large cardinals), and haven't been able to find anything at all (except for an article of Bagaria using supercompacts). Has there been any development on this open problem? What is the most likely direction: an argument assuming a proper class of measurables or deriving a proper class of strongly compacts?

论坛问题帖 | no_answers_at_retrieval | interregno | 2023-10-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018347 | MO-458564

An open problem about simple Noetherian rings

The following is a well-known open problem in ring theory (see, for instance, Goodearl, Warfield, 'An introduction to noncommutative Noetherian rings, Appendix, Problem 19) Question: Let R be a left Noetherian simple ring. Can every left ideal be generated by two elements? The history of this problem goes back to the seminal work of Stafford 'Module structure of the Weyl algebra'. In this paper, Stafford showed, among other things, that k being a field of zero characteristic, every left ideal for the Weyl algebras $A_n(k)$ can be generated by two elements. In the same paper, he stated as problem to find large classes of algebras for which his results on the Weyl algebras also hold; in particular, the two-generation property of left ideals. A lot of research has been done on this topic. There are some natural generalizations of Stafford's result for algebras that are similar to the Weyl algebra: the two-generation property of left ideals holds for $k(x_1,\ldots,x_n)\langle \partial_1, \ldots, \partial_n \rangle$, and the subalgebra of $End_k k((x_1,\ldots,x_n))$ generated by $k((x_1,\ldots,x_n))$ and the usual partial derivatives $\partial_1, \ldots, \partial_n$. Bavula showed that the two-generation property for left ideals holds for many generalized Weyl algebras - for example, it holds for the quantum torus $k_q[x^\pm,y^\pm]$, when q is not a root of unity. A natural question, raised in 1980 by Björk, is if we have the two-generation property for left ideals for rings of differential operators on smooth affine varieties. Coutinho and Holland worked on this problem, and showed that every left ideal for such rings can be generated by three elements - and Björk's question is still open. If we consider the Weyl algebra $A_n(D)$ over a division ring D of zero characteristic, we have a similar result, obtained by Byun: every left ideal can be generated by three elements. All of the above are particular cases of rings. The only general result I know, also due to Stafford, is the following: let R be a simple left Noetherian ring with Krull dimension d. Then every left ideal can be generated by $d+1$ elements. I am curious for what reasons there are to believe or not that every left ideal for a simple left Noetherian ring can be generated by two elements. Sure, the results I described above count as positive evidence to believe this fact. However, in my opinion, all those rings above are rather similar in nature: all of them are some natural generalization of the Weyl algebra. So, besides the above discussed positive results and lack of counter-examples, are there other reasons to believe in a positive solution to this open problem?

论坛问题帖 | no_answers_at_retrieval | jg1896 | 2023-11-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018348 | MO-459295

Conjecture: Given any five points, we can always draw a pair of non-intersecting circles whose diameter endpoints are four of those points

The following question resisted attacks at Math SE, so I thought I would try posting it here. Is the following conjecture true or false: Given any five coplanar points, we can always draw at least one pair of non-intersecting circles coplanar with the points, such that two of the given points are diameter endpoints of one circle, and another two of the given points are diameter endpoints of the other circle. Tangent circles are considered to be non-intersecting. Coincident circles are considered to be intersecting. Example: Another example: I cannot find a counter-example, nor can I prove the conjecture. Remarks: I made a generator of five (pseudo)random points. If, instead, we were given four points, the conjecture would not be true: for example, if the four points were the vertices of an equilateral triangle plus the centre, then we could not draw a pair of non-intersecting circles. The generalized conjecture with $2n+1$ points and n circles is not true. I have asked, and answered, a similar question, which might provide ideas for this question. Context: I was thinking about this question about random points in a disk. Staring at various sets of five points, I came up with this conjecture.

论坛问题帖 | has_answers | Dan | 2023-11-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018349 | MO-459516

Does the interior of Pascal's triangle contain three consecutive integers?

This question defeated Math SE, so I am posting it here. Consider the interior of Pascal's triangle: the triangle without numbers of the form $\binom{n}{0},\binom{n}{1},\binom{n}{n-1},\binom{n}{n}$. A006987 (which is described here) is a list of the smallest 10000 numbers in the interior of Pascal's triangle, omitting repetitions. It contains 15 instances of two consecutive integers. It contains only one instance of integers separated by 2. But it does not contain three consecutive integers. So I wonder: Does the interior of Pascal's triangle contain three consecutive integers? Conjecture 3.1 of Binomial Collisions and Near Collisions is the conjecture that a list contained in that paper is the complete list of binomials that differ by 1. If this conjecture

is true, this would imply that there are no three consecutive binomial coefficients. But even if this conjecture is unresolved, there might be a way to show that there are no three consecutive binomial coefficients. Maybe proof by contradiction, or a combinatorial argument?

论坛问题帖 | no_answers_at_retrieval | Dan | 2023-11-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018350 | MO-461473

What is the current status of research on the von Neumann's inequality for $n \ge 3$?

Problem Let $(T_1, \ldots, T_n)$ be a tuple of commuting contractions in Hilbert space H. Does a constant $C_n \ge 1$ exist, for which it would be true, that: $$\forall_{p \in \mathbb{C}[x_1, \ldots, x_n]} \quad \| p(T_1, \ldots, T_n) \| \le \ C_n \sup_{(z_1, \ldots, z_n) \in \overline{\mathbb{D}^n}} | p (z_1, \ldots, z_n) |$$ I've heard numerous anecdotes from professors and other scientific employees, that there has been some more recent results in this field. One of them being "using graph theory for $n \ge 3$", and the other one "using spectral theory". I couldn't find anything new, the only "newest" thing is Shalit's work from 2021 if I remember correctly. Could anyone help me out?

论坛问题帖 | no_answers_at_retrieval | S-F | 2024-01-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018351 | MO-462321

5n+1 sequence starting at 7

Consider the following variant of the Collatz function: $f:\mathbb N\rightarrow\mathbb N$ is defined by \begin{equation} f(n):=\begin{cases} n/2 & \text{if n is even}\\ 5n+1 & \text{if n is odd}\\ \end{cases} \end{equation} Contrary to the Collatz function with $3n+1$, many orbits of this function seem to diverge. The smallest one is probably 7. After 32282 iterations, $f^k(7)$ is already larger than 10^{1000}. Can one prove that this orbit diverges to infinity? Here is the Python code that I used: from decimal import * getcontext().prec=1001 def collatz5(n): if n%2 == 0: return Decimal(n)/Decimal(2) else: return Decimal(5)*Decimal(n)+Decimal(1) n=7 for i in range(100000): if n>10**1000: print("bound exceeded") print("number of steps: ",i) break n=collatz5(n)

论坛问题帖 | has_answers | Riemann | 2024-01-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018352 | MO-462557

Optimal covering trails for every k-dimensional cubic lattice $\mathbb{N}^k := \{(x_1, x_2, \dots, x_n) : x_i \in \mathbb{N} \wedge n \geq 3\}$

After a dozen years spent investigating this particular class of problems, I finally give up and I wish to ask you if any improvement is achievable from here on. The general problem is as follows: Let the k-dimensional grid $\mathcal{H}(n,k):=\{0,1,2,\ldots,n-1\} \times \cdots \times \{0,1,2,\ldots,n-1\}$ be given in the Euclidean space $\mathbb{R}^k$. How many line segments connected at their endpoints have the minimum-link covering trails that join all the points of $\mathcal{H}(n,k)$ (i.e., the optimal polygonal chain that visits all the given n^k points by spending the minimum number of line segments)? Then, let us call $h(n,k)$ the mentioned minimum number of links. Accordingly, let us denote with $h_l(n,k)$ the proven lower bound for $h(n,k)$ and with $h_u(n,k)$ the corresponding upper bound so that we can state the following set of relations. \begin{equation} h(n,1)=1, \forall n. \end{equation} \begin{equation} h(n,2)=2 \cdot (n-1), \forall n \geq 3. \end{equation} \begin{equation} h(1,k)=1, \forall k. \end{equation} \begin{equation} h(2,k)=3 \cdot 2^{k-2}, \forall k \geq 2. \end{equation} \begin{equation} h(3,k)=\frac{3^k-1}{2}, \forall k. \end{equation} \begin{equation} h(n,k) \leq \Bigg(\Bigl\lfloor{\frac{3}{2} \cdot n^2}\Bigl\rfloor - \Bigl\lfloor{\frac{n-1}{4}}\Bigl\rfloor + \Bigl\lfloor{\frac{n+1}{4}}\Bigl\rfloor - \Bigl\lfloor{\frac{n+2}{4}}\Bigl\rfloor + \Bigl\lfloor{\frac{n}{4}}\Bigr\rfloor + n - 1 \Bigg) \cdot n^{k-3} - 1, \forall n\geq 3 \wedge \forall k \geq 3 \end{equation} \begin{equation} h(n,k) \geq \frac{n^k-1}{n-1}, \forall n \geq 3 \wedge \forall k. \end{equation} \begin{equation} h(n,k) \leq (h_u(n,3)+1) \cdot n^{k-3}-1, \forall n\geq 3 \wedge \forall k \geq 3. \end{equation} Thus, what I have currently found is resumed in the table below, where the strictly proven results are highlighted in the green cells, while the proven bounds are shown in the orange cells. $\{0,1,2,\dots,n-1\}\times\{0,1,2,\dots,n-1\}\times \dots \times\{0,1,2,\dots,n-1\}$ points problem in $\mathbb{R}^k$." /> Conjecture: $h(n,k)=\frac{n^k-1}{n-1}+c \cdot (n-3)$, where $c = k-1$ iff $h(4,3)=23$, $c = 1$ iff $h(4,3)=22$, or even $c = 0$ iff $h(4,3)=21$ (see also this Previous discussion about finding the value of $h(4,3)$). Any idea to go further (can we see some kind of pattern in the table)? Thanks in advance!

论坛问题帖 | no_answers_at_retrieval | Marco Ripà | 2024-01-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018353 | MO-462733

A more complete set of open problems

Over time, there have been a number of posts on open problems remaining in different fields of math, both here and on the MathSE. So I had the idea of trying to construct a “list of lists” of problems from the many that exist - and it turns out someone else has already started such an effort! I’ve found something quite neat, so I’ll be posting an answer below.

论坛问题帖 | has_answers | Aidan W. Murphy | 2024-01-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018354 | MO-462981

Open problems which might benefit from computational experiments

Question: I wonder what are the open problems , where computational experiments might be helpful? (Setting some bounds, excluding some cases, shaping some expectations). Grant program: The context of the question is the following - in some sense there is a way to crowdsource some (quite specific, but still) computational experiments - the platform for data science challenges Kaggle (Google's subsidiary) - offers grants to researchers in various fields to organize the challenges. Properly organized it might attract thousands of participants, some of whom are extremely skilled in solving computational challenges (like NVIDIA grandmaster's team - 6 super-skillful guys who are paid by NVIDIA just to participate in challenges (and promote some NVIDIA tools) - kind of profi-sportsmen). There already

many research oriented challenges took place - mainly around in bioinformatics (e.g. CAFA5, Stanford-Ribonanza, Single Cell Perturbations, etc..). Example: The one quite related to mathematics - is ongoing challenge on finite permutation groups - the task is to find as short paths as possible for sets of given points on Cayley graphs. The groups include N-dimensional Rubik's cube and other puzzle motivated groups like "Globe". For example even the diameters of the N-dimensional Rubik's cube are still unknown, so computational experiments may shed light on these questions. The other questions - like better understanding of the random walks, growth patterns, patterns of mixing, fast scrambling algorithms - seems to be of interest from both sides - machine learning and mathematical. So fruitful idea exchanges and interconnections might be possible. It would be great if some mathoverflow colleagues experienced in such kind of permutation group questions may share experience - please post on forum e.g. here or/and contact e.g. me. PS I am a kind of a fan of many sides of Kaggle - it managed to create a community of people who are not only competing but sharing lots of experience. Gamified "reputation" scores - similar to mathoverflow - play a role, moreover Kaggle rankings are appreciated in the data science world, so can help to find a better job - another stimulus for many to participate, in addition to networking. Many people think that the best way to learn practical data science - participate in Kaggle challenges. It is not only for competition, but a free cloud service - like Google Colab, but more devoted to team work - offering free computing power, storage, sharing, version control, team work. Would be great to have something similar for mathematical community or to use Kaggle itself for mathematical community - one may think of a competition as a kind of mini-"Polymath" project. Where all the infrastructure - is ready - forum, community, free computing power - the only thing one needs - is to create a challenge. Kaggle offers everybody to create a challenge for free, but only "official" challenges attract most attention.

论坛问题帖 | has_answers | Alexander Chervov | 2024-01-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018355 | MO-464357

Squares whose differences are squares

EDIT. I've just noticed a thread from 2011 in the "Related" column on the right (click me), where a closely related question is being discussed (the main difference seems to be that, in Question 2 below, I'm only asking for the existence of an upper bound). A subproblem of Problem D18 in the 2004 softcover reprint of the 3rd edition of Guy's Unsolved Problems in Number Theory asks if there are four squares the absolute values of whose differences in pairs are squares (see p. 278 in the book, the subproblem is titled "Four squares whose differences are square"). Guy comments that "Although a solution is unlikely, there do not appear to be any congruence conditions which forbid it." I have two (main) questions: (Q1) What is the status of the problem? Has there been any substantial progress since 2004? (Q2) What about the existence of an integer $N \ge 4$ such that there is no N-tuple $(a_1, \ldots, a_N)$ of positive integers with the property that $|a_i^2 - a_j^2|$ is a square for all $i, j = 1, \ldots, N$? More precisely, what about $N = 5$? The question is motivated by a problem in graph theory posed by Sergey Goryainov. I'm mainly looking for references.

论坛问题帖 | no_answers_at_retrieval | Salvo Tringali | 2024-02-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018356 | MO-46444

Natural numbers n which satisfy gnu(n)=n?

Are there any natural numbers n (other than 1) for which $gnu(n)=n$? We define $gnu(n)$ to be the number of isomorphism classes of groups of order n. This question popped into my head today, and I couldn't come up with a proof one way or another. In the paper by Conway, et al. entitled "Counting Groups: Gnus, Moas, and other Exotica, Conjecture 10.1 implies that there should be no such natural number n which satisfies $gnu(n)=n$. Does anybody have an argument to show that there is no natural number n (other than 1) for which $gnu(n)=n$? If it is really straightforward, just say so and I'll work it out for myself when I find time. Thanks!

论坛问题帖 | no_answers_at_retrieval | Glen M Wilson | 2010-11-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018357 | MO-46445

Zeroes of a tricky function.

I am attempting to show that there does not exist an N past which every open unit interval (k, k+1) -where k is an integer- contains a zero of the following function: $h(x)=\sum_{n=2}^{[\sqrt(x)]} \frac{\cot(x\pi/n)}{n}+\frac{\cot((x+2)\pi/n)}{n}$ Where the $[\sqrt(x)]$ is the lowest integer of the square root of x. Any thoughts? I had figured that I could consider the interval $(i^2, (i+1)^2)$ in which h(x) is described by the function $h(x)=h_n(x)=\sum_{n=2}^{i} \frac{\cot(x\pi/n)}{n}+\frac{\cot((x+2)\pi/n)}{n}$ Then try to see on what unit intervals (k, k+1) in here contain a zero of h_n then try to show that $((i+1)^2, (i+2)^2)$ also has such an interval, but I am running out of ideas.

论坛问题帖 | no_answers_at_retrieval | Alex Botros | 2010-11-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018358 | MO-464644

On the number of complete Boolean algebras

In their 1972 paper On the number of complete Boolean algebras Monk and Solovay showed that if λ is an infinite cardinal, then there are 2^{2^λ} many isomorphism types of complete Boolean algebras of power $2^\lambda.$ On the other hand, by a result of Pierce, a complete Boolean algebra of infinite power κ exists iff $\kappa=\kappa^{\aleph_0}$. In the above mentioned paper, the following is asked: Question. Suppose $\kappa=\kappa^{\aleph_0}$, but it is not of the form 2^λ for any λ. Are there ${2^\kappa}$ many isomorphism types of complete Boolean algebras of power $\kappa?$ What is known about the above question?

论坛问题帖 | has_answers | Mohammad Golshani | 2024-02-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018359 | MO-4672

level sets of multivariate polynomials

Let $p:\mathbb R^n \rightarrow \mathbb R$ be a polynomial of degree at most d. Restrict p to the unit cube $Q=[0,1]^n\subset\mathbb R^n$. We assume that p has mean value zero on the unit cube Q: $$\int_Q p(x) dx = 0.$$ For $\alpha>0$ consider the sublevel sets of P, $$E_\alpha= \{x\in Q: |p(x)|\leq \alpha\}$$ There are several known estimates for the Lebesgue measure of this set which in some sense or another are uniform over some classes of polynomials. For example, we have that $$|E_\alpha| \lesssim \min(pd,n) \frac{ \alpha^{1/d} }{ \|p\|_{ L^p(Q) }^{1/d} } $$ This particular estimate is due to Carbery and Wright and can be found here. I'm interested in studying the (induced Lebesgue) measure of the boundary of this set $$|\partial E_\alpha|=|\{x\in Q: |P(x)|=\alpha\}| $$ Consider first the easy case of dimension $n=1$. Then the set E_α is a finite union of closed intervals and the question is trivial. It is obvious that in this case there are at most $O(d)$ intervals so the 0-dimensional measure of the boundary is $O(d)$. Now in many variables things will be much more complicated. For example can we say that the set E_α has $O(d)$ connected components? Is there an estimate for the measure of the boundary $\partial E_\alpha $ in terms of α, d and n, assuming (say) that $\|p\| _ {L^1(Q)}=1$? This question comes up naturally if one tries to study an oscillatory integral with phase p and apply integration by parts (i.e Gauss theorem) imitating the one dimensional method of proving the van der Corput lemma (for example).

论坛问题帖 | has_answers | ioannis.parissis | 2009-11-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018360 | MO-46769

Covering the primes by arithmetic progressions

Define the length of a set of arithmetic progressions of natural numbers $A=\lbrace A_1, A_2, \ldots \rbrace$ to be $\min_i | A_i |$: the length of the shortest sequence among all the progressions. Say that A exactly covers a set S if $\bigcup_i A_i = S$. Let P' be the primes excluding 2. What is the longest set of arithmetic progressions that exactly covers the primes P'? In other words, I want to maximize the length of a set of such arithmetic progressions. Call this maximum $L_{\max}$. $L_{\max} \ge 2$ because $$ P' \;=\; \lbrace 3,5 \rbrace \cup \lbrace 7,11 \rbrace \cup \cdots \cup \lbrace 521,523 \rbrace \cup \cdots $$ Perhaps it is possible that $$P' \;=\; \lbrace 3, 11, 19 \rbrace \cup \lbrace 5, 17, 29,41,53 \rbrace \cup \lbrace 7,19,31,43 \rbrace \cup \cdots \;,$$ but I cannot get far with sequences of length ≥ 3. (I know Green-Tao establishes that there are arbitrarily long arithmetic progressions in P, but I don't know if that helps with my question.) I am number-theoretically naïve, and apologize if this question is nonsensical or trivial. In any case, I appreciate the tutoring! Addendum. Although my question should be revised (as Idoneal suggests) in light of George Lowther's proof that 3 cannot be in a progression of length 4, George has shown that it is likely that $L_{\max}=3$ but certification requires resolving an open problem. So I've added the open-problem tag. Thanks for everyone's interest!

论坛问题帖 | has_answers | Joseph O'Rourke | 2010-11-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018361 | MO-468737

Could I possibly exploit distinct odd primes raised to 6 to solve Exact Three Cover, when reducing it in Subset Sum?

I'm solving Exact 3 Cover, given a list with no duplicates S of $3m$ whole numbers and a collection C of subsets of S, each containing exactly three elements. The goal is to decide if there are $\operatorname{len}(S)/3$ subsets that cover every element in S one time. N is the $len(S)$ I'm going to use Subset Sum to solve Exact 3 Cover. I reduce Exact Three Cover into Subset Sum by transforming S into N odd distinct primes raised to 6, and easily map out the collection of subsets in the same manner. I then get the sums of the transformed collection of subsets. And the sum of the transformed S will be our target sum. So now I have a transformed $S = {p_1^6, p_2^6, p_3^6...}$ into the first N distinct odd primes raised to 6. There are rules for combinations of size three to ensure distinctness. All combinations of size three must be sorted in ascending order. There will be no duplicate subsets. There will be no subsets that have elements not in S . There will be no subsets that have duplicate elements. This does not affect correctness, as it is either impossible or not necessary that a solution could use these subsets. We easily filter the input to get rid of these unnecessary subsets. The conjecture is that no distinct subset that was transformed into primes raised to 6 could have duplicate sums. Because duplicates could cause collision and the reduction would fail. This is a variant of the unsolved problem of $a^6 + b^6 + c^6 ≠ d^6 + e^6 + f^6$ where variables are distinct primes. In this case both sides of the equation is supposed to be distinct subsets of size three. Of course, we have to tweak it to encapsulate all combinations of size 3. For example, When a variable is shared between two distinct subsets. $A^6 + b^6 + c^6 ≠ A^6 + e^6 + f^6$ $b^6 + c^6 ≠ e^6 + f^6$ is another open problem for distinct primes, it also implies $A^6 + b^6 + c^6 ≠ A^6 + e^6 + f^6$ When two variables are shared between two distinct subsets. $A^6 + B^6 + c^6 ≠ A^6 + B^6 + f^6$. What about when two subsets share all variables? Per the rules of combinations no duplicate subsets. Because $a^6 + b^6 + c^6 = a^6 + b^6 + c^6$ For generality the position of shared variables could be at any location on either side of the equation. Also the unique factorization of each p^6 seems to imply unique sums for all combinations of size 3. My question is, could I possibly exploit properties of numbers to solve Exact Three Cover more efficiently? Everybody's looking to use conventional techniques to solve NP complete problems, but nobody seems to looking for how numbers work. And, I think that could yield interesting finds. Even if the reduction is dependent on conjecture, its still nonetheless interesting. I have written code for empirical analysis, and I've had no duplicate sums all the way up to 500. This was originally asked on Theoretical Computer Science

论坛问题帖 | has_answers | The T | 2024-04-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018362 | MO-468771

Congruence obstructions for three consecutive powerful numbers

A powerful number is an integer m such that if p is prime and $p \mid m$ then $p^2 \mid m$. Powerful numbers can be represented in the form $m=u^2 v^3$. Erdos conjectured that three consecutive powerful numbers don't exist. Assume $(a-1,a,a+1)$ is a

triple of consecutive powerful numbers. Choose integers n,A and set $a=A$ and assume the system $$x_1^2 x_2^3=A-1, y_1^2 y_2^3=A+1$$ doesn't have solution modulo n. Then over the integers powerful triple with $a \equiv A \pmod{n}$ doesn't exist. Some examples with n square: n | A congruence obstruction 4 [1, 3] 9 [2, 4, 5, 7] 16 [1, 3, 5, 7, 9, 11, 13, 15] 25 [4, 6, 9, 11, 14, 16, 19, 21] 36 [1, 2, 3, 4, 5, 7, 9, 11, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 25, 27, 29, 31, 32, 33, 34, 35] 49 [6, 8, 13, 15, 20, 22, 27, 29, 34, 36, 41, 43] Q1 Can we find n and congruence obstructions to show non-existence of more naturally defined powerful triples? Q2 Can we show that for some even B the triple with $a=B^k$ doesn't exist for all k? (For odd B this is visible with naked eye)

论坛问题帖 | has_answers | joro | 2024-04-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018363 | MO-468899

No starter "accessible" well known open problems

We all know very famous open problems. Usually the ones that become famous are well studied and lots of progress was achieved and conjectures, partial results, reductions and so on exist. This is one of the reasons they become famous. This kind of problems are usually hopeless to attack without lots of investment of time and dedication to learn the basics and beyond. I am interested in the other extreme, that is, I would like relatively easy to state problems which are well known (at least in the relevant area), but no one has a clue how to start solving them, preferably no one even has a good guess what the right answer is. Let me give you two examples from my research area: A pro-p group in which there exists a bound on the number of generators (topologically) for all closed subgroups is said to have finite rank. These are well known due to the works of Lazard, and Lubtozky and Mann, for instance, they are p-adic analytic. However, as far as I know known no-one has a clue whether a pro-p group in which every closed subgroup is finitely generated has finite rank? In other words, if a pro-p group is not of finite rank does it contain a closed subgroup which is not finitely generated? Zelmanov solved the Restricted Burnside Problem that is there are finitely many finite p-groups with d-generators and exponent at most e. This is equivalent to saying that every finitely generated pro-p of finite exponent is finite. He also generalized and proved the Burnside Problem for pro-p groups, that is, every finitely generated torsion pro-p is finite. However, the Generalized Burnside Problem for pro-p group is completely open, that is, is it true that a torsion pro-p group (not necessarily f.g.) has finite exponent? (I'll just comment that Burnside kind of problems are usually very hard.) Motivation: Well I recently retired and while I do not have the energy (or talent) to try to solve most famous open problem, it would be nice to have a list of famous problems that either by luck one might encounter a counter example or by luck one might find a very different approach.

论坛问题帖 | has_answers | Yiftach Barnea | 2024-04-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018364 | MO-470132

A probability involving areas in a random pentagram inscribed in a circle: Is it really just $\frac12$?

This question was posted at MSE but was not answered. The vertices of a pentagram are five uniformly random points on a circle. The areas of three consecutive triangular "petals" are a,b,c. The petals are randomly chosen, but they must be consecutive, either clockwise or anticlockwise. A simulation of 10^7 such random pentagrams yielded a proportion of 0.5000179 satisfying $a^2<bc$. Is the following conjecture true: $P(a^2<bc)=\frac{1}{2}$ Remarks Note that the three petals must be consecutive. Calling the areas of consecutive petals a,b,c,d,e, simulations suggest that: $P(a^2<bd)\approx0.468$ $P(a^2<be)\approx0.505$ $P(a^2<cd)\approx0.460$ Curiously, the probability that seems to equal $\frac{1}{2}$, i.e. $P(a^2<bc)$, does not involve a symmetrical arrangement of three petals. As for other random star polygons $\{\frac{n}{2}\}$ inscribed in a circle, simulations suggest that Star polygon $\left\{\frac{6}{2}\right\}$: $P(a^2<bc)\approx0.505$. Star polygon $\{\frac{7}{2}\}$: $P(a^2<bc)\approx0.504$. I used the shoelace formula to calculate the areas of the triangular petals. One might expect that a probability of $\frac12$ should have an intuitive explanation, but sometimes probabilities of $\frac12$ are hard to explain. Underlying reason? Simulations suggest that the random pentagram/pentagon shape is teeming with probabilities of powers of $\frac12$. In the following diagram, the letters represent areas of the regions. $P(g+b+h<a+f+c)\overset{?}{=}\color{red}{\frac{1}{2}}$ $P(g+h<f)\overset{?}{=}\color{red}{\frac{1}{4}}$ $P((g+a+k)(h+c+i)>(b+f+e+d+j)^2)\overset{?}{=}\color{red}{\frac{1}{8}}$ $P(\text{each region with area g,b,f,l contains the centre of the circle})=\color{red}{\frac{1}{16}}$ (proved) $P(\text{areas of petals increase going around once, clockwise or anticlockwise})\overset{?}{=}\color{red}{\frac{1}{32}}$ I am not asking to prove these other probability claims. I am presenting them to suggest that there may be an underlying reason why the pentagram has probabilities of powers of $\frac12$, which may be relevant to my conjecture. These simulations make me more inclined to believe that my conjecture is true, but I could be wrong.

论坛问题帖 | has_answers | Dan | 2024-04-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018365 | MO-470539

Probability of finding a prime number between $x-\ln(x)$ and $x+\ln(x)$

Using my computer, I found that in the interval $[1, N]$ the probability of finding a prime number between $x-\ln(x)$ and $x+\ln(x)$ is greater than constant c where $N=10^2, 10^3,...,10^{9}$, x is the positive integer number in $[1, N]$ and $c > 0.8$. Question: Does the result hold for $N \to +\infty $?

论坛问题帖 | has_answers | Đào Thanh Oai | 2024-05-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018366 | MO-470571

Conjecture: x^4+1 is never Wieferich prime

Related to this question and Alexander Kalmynin's answer. For natural n define $J(n)=(2^{n-1}-1) \bmod n^2$ and if n is power of two define $J(2^n)=1$ (this is artificial, just to avoid triviality and by $\bmod$ we mean take the smallest residue modulo n^2). Prime p is Wieferich prime iff $J(p)=0$. The answer in the linked question use the doubly exponential sequence of Fermat numbers and we are trying to generalize this. Conjecture 1: For even n we have $J(n^2+1) < J(n^4+1)$ and hence $J(n^4+1)>0$, so n^4+1 can't be Wieferich prime. This holds for n up to 10^6 and several very large n. Further we have: Conjecture 2 For even n large enough we have $\frac{\log(J(n^4+1))}{\log{J(n^2+1)}} \sim 2$. Q1 If Conjecture 1 is false, can we use this approach to show primes of certain form are not Wieferich?

论坛问题帖 | no_answers_at_retrieval | joro | 2024-05-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018367 | MO-472263

Rearrangement inequality for sum

Rearrangement inequality: Assume we have finite ordered sequences of nonnegative real numbers $0 \le a_1 \le a_2 \le\cdots\le a_n \quad\text{and}\quad 0\le b_1 \le b_2 \le\cdots\le b_n, \cdots\,, \quad\text{and}\quad 0\le z_1 \le z_2 \le\cdots\le z_n$ and a permutation $b_{\sigma(1)}, b_{\sigma(2)}, \ldots,b_{\sigma(n)}$ of $b_1, b_2\dots,b_n$,$\cdots$, and another permutation $z_{\tau(1)}, z_{\tau(2)}, \dots,z_{\tau(n)}$ of $z_1, z_2, \dots,z_n$, then $$ a_1 b_{\sigma(1)}\cdots z_{\tau(1)} + \cdots + a_n b_{\sigma(n)} \cdots z_{\tau(n)} \le a_1 b_1 \cdots z_1 + \cdots + a_n b_n \cdots z_n.$$ Does this inequality hold if we replace multiplication by sumation? detail as follows: Assume we have finite ordered sequences of real numbers $ a_1 \le a_2 \le\cdots\le a_n \quad\text{and}\quad b_1 \le b_2 \le\cdots\le b_n, \cdots\,, \quad\text{and}\quad z_1 \le z_2 \le\cdots\le z_n$ and a permutation $b_{\sigma(1)}, b_{\sigma(2)}, \ldots,b_{\sigma(n)}$ of $b_1, b_2\dots,b_n$,$\cdots$, and another permutation $z_{\tau(1)}, z_{\tau(2)}, \dots,z_{\tau(n)}$ of $z_1, z_2, \dots,z_n$, then $$ |a_1 + b_{\sigma(1)}+\cdots +z_{\tau(1)}| + \cdots + |a_n + b_{\sigma(n)}+ \cdots + z_{\tau(n)}| \le |a_1 + b_1 + \cdots + z_1| + \cdots + |a_n + b_n + \cdots + z_n|$$

论坛问题帖 | has_answers | Đào Thanh Oai | 2024-05-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018368 | MO-473926

Open problems in complete theories

It is well-known that every complete recursively enumerable first-order theory is decidable. Does that mean that such theories are "trivial", or are there still interesting open problems that we potentially know the answer of (by running a computer for an (insane) amount of time)? Examples of complete theories include: Presburger arithmetic Tarski's axioms for Euclidean geometry The theory of dense linear orders without endpoints The theory of algebraically closed fields of a given characteristic The theory of real closed fields Most (all?) examples given here about Euclidean geometry do not seem to be expressible in Tarski's language.

论坛问题帖 | has_answers | user507115 | 2024-06-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018369 | MO-473957

Does the sequence formed by Intersecting angle bisector in a pentagon converge?

I asked this question on MSE here. Given a non-regular pentagon $A_1B_1C_1D_1E_1$ with no two adjacent angle having a sum of 360 degrees, from the pentagon $A_nB_nC_nD_nE_n$ construct the pentagon $A_{n+1}B_{n+1}C_{n+1}D_{n+1}E_{n+1}$ as follows: A_{n+1} is the intersection between the angle bisectors of $\angle C_n $ and $\angle D_n$. B_{n+1} is the intersection between the angle bisectors of $\angle D_n$ and $\angle E_n$. C_{n+1} is the intersection between the angle bisectors of $\angle E_n$ and $\angle A_n$. D_{n+1} is the intersection between the angle bisectors of $\angle A_n$ and $\angle B_n$. E_{n+1} is the intersection between the angle bisectors of $\angle B_n$ and $\angle C_n$. (the intersection of the angle bisectors of the two opposite angles) I am allowing self-intersecting polygons in this constructions. The reason why I chose this construction is that the point A_{n+1} is the only point that doesn't depend on A_n. My question is if this process is repeated indefinitely would the sequences ${A_n}$, ${B_n}$, ${C_n}$, $D_n $ and E_n converge ? and if they converge what is the limit? if they din't always converge what are the necessary conditions that the pentagon $A_1B_1C_1D_1E_1$ should satisfy so the five sequences converge ? There are only four possible scenarios: The points will converge. The points will eventually trapped on a loop. The points will diverge completely. Two or more points coincide, or two adjacent angle have a sum of 360 degrees ending the sequence. I tried to draw the first few pentagons to see if the point will converge or not. Here is the first 50 pentagons in a zoom-in animation: I conjecture that the sequences converge to a single point for all non-regular pentagons. If convergence occurs, how can we determine the limit point based on the initial pentagon? If the limit point exist what interesting properties does it have for the pentagon? Here is a Geogebra file that have the first 50 pentagons. I wrote a python code than can calculate A_n, B_n , C_n ,D_n ,E_n given the initial pentagon: import math print("Number of pentagons") n=int(input()) m=max(n,50) from decimal import Decimal import decimal decimal.getcontext().prec = m Decimal(10)**(-m) # enter your coordinates here A=[Decimal(0),Decimal(0)] B=[Decimal(0),Decimal(0)] C=[Decimal(0),Decimal(0)] D=[Decimal(0),Decimal(0)] E=[Decimal(0),Decimal(0)] print("Enter the coordinates of A_1") A[0]=Decimal(float (input())) A[1]=Decimal(float (input())) print("Enter the coordinates of B_1") B[0]=Decimal(float (input())) B[1]=Decimal(float (input())) print("Enter the coordinates of C_1") C[0]=Decimal(float (input())) C[1]=Decimal(float (input())) print("Enter the coordinates of D_1") D[0]=Decimal(float (input())) D[1]=Decimal(float (input())) print("Enter the coordinates of E_1") E[0]=Decimal(float (input())) E[1]=Decimal(float (input())) X=[Decimal(0),Decimal(0)] Y=[Decimal(0),Decimal(0)] Z=[Decimal(0),Decimal(0)] W=[Decimal(0),Decimal(0)] T=[Decimal(0),Decimal(0)] #defining some useful functions---------- def v(a,b): result= [b[0]-a[0], b[1]-a[1]] return result def crs(a, b): result = [-Decimal(0.5)*((Decimal(a[0]*b[1])-Decimal(a[1]*b[0])))] return result def dis(a,b): result =[Decimal(math.sqrt((a[0]-b[0])**2+(a[1]-b[1])**2))] return result for i in range (2,n+1): AB= dis(A,B)[0] BC= dis(B,C)[0] CD= dis(C,D)[0] DE= dis(D,E)[0] EA=

dis(E,A)[0] #Here to calculate A_n---------- x= CD*crs(v(D,C),v(D,E))[0] y= -CD*crs(v(E,D),v(E,B))[0]+ 2*DE*crs(v(C,B),v(C,D))[0] z= -CD*crs(v(C,B),v(C,E))[0]+ 2*BC*crs(v(D,C),v(D,E))[0] t= CD*crs(v(C,B),v(C,D))[0] X[0]=(x *B[0]+y*C[0]+z*D[0]+t*E[0])/(x+y+z+t) X[1]=(x *B[1]+y*C[1]+z*D[1]+t*E[1])/(x+y+z+t) #Here to calculate B_n---------- x= DE*crs(v(E,D),v(E,A))[0] y= -DE*crs(v(A,E),v(A,C))[0]+ 2*EA*crs(v(D,C),v(D,E))[0] z= -DE*crs(v(D,C),v(D,A))[0]+ 2*CD*crs(v(E,D),v(E,A))[0] t= DE*crs(v(D,C),v(D,E))[0] Y[0]=(x *C[0]+y*D[0]+z*E[0]+t*A[0])/(x+y+z+t) Y[1]=(x *C[1]+y*D[1]+z*E[1]+t*A[1])/(x+y+z+t) #Here to calculate C_n---------- x= EA*crs(v(A,E),v(A,B))[0] y= -EA*crs(v(B,A),v(B,D))[0]+ 2*AB*crs(v(E,D),v(E,A))[0] z= -EA*crs(v(E,D),v(E,B))[0]+ 2*DE*crs(v(A,E),v(A,B))[0] t= EA*crs(v(E,D),v(E,A))[0] Z[0]=(x *D[0]+y*E[0]+z*A[0]+t*B[0])/(x+y+z+t) Z[1]=(x *D[1]+y*E[1]+z*A[1]+t*B[1])/(x+y+z+t) #Here to calculate D_n---------- x= AB*crs(v(B,A),v(B,C))[0] y= -AB*crs(v(C,B),v(C,E))[0]+ 2*BC*crs(v(A,E),v(A,B))[0] z= -AB*crs(v(A,E),v(A,C))[0]+ 2*EA*crs(v(B,A),v(B,C))[0] t= AB*crs(v(A,E),v(A,B))[0] W[0]=(x *E[0]+y*A[0]+z*B[0]+t*C[0])/(x+y+z+t) W[1]=(x *E[1]+y*A[1]+z*B[1]+t*C[1])/(x+y+z+t) #Here to calculate E_n---------- x= BC*crs(v(C,B),v(C,D))[0] y= -BC*crs(v(D,C),v(D,A))[0]+ 2*CD*crs(v(B,A),v(B,C))[0] z= -BC*crs(v(B,A),v(B,D))[0]+ 2*AB*crs(v(C,B),v(C,D))[0] t= BC*crs(v(B,A),v(B,C))[0] T[0]=(x *A[0]+y*B[0]+z*C[0]+t*D[0])/(x+y+z+t) T[1]=(x *A[1]+y*B[1]+z*C[1]+t*D[1])/(x+y+z+t) A[0]=X[0] A[1]=X[1] B[0]=Y[0] B[1]=Y[1] C[0]=Z[0] C[1]=Z[1] D[0]=W[0] D[1]=W[1] E[0]=T[0] E[1]=T[1] print(f"A_{i}= {A}") print(f"B_{i}= {B}") print(f"C_{i}= {C}") print(f"D_{i}= {D}") print(f"E_{i}= {E}") print("\n") Another thing that supports the convergence of the series that no matter how many times I used that random function in python with the sequence the result always converge (suggesting it always converge except some zero measure set)

论坛问题帖 | has_answers | pie | 2024-06-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018370 | MO-47419

Difficult examples for Frankl's union-closed conjecture

Frankl's well-known union-closed conjecture states that if F is a finite family of sets that is closed under taking unions (that is, if A and B belong to the family then so does $A\cup B$), then there must be an element that belongs to at least half the sets. I know that pretty well any naive approach one takes to this conjecture is known to fail. By "naive approach" I suppose I mean something like an observation that it would follow from such-and-such a stronger conjecture -- it seems that all sensible stronger conjectures one thinks of are false. A very simple example of a stronger conjecture would be that if you pick a random element then on average it will belong to at least half the sets. That is completely false: take the family that consists of the empty set, {1}, and {1,2,3,4,5,6,7,8,9,10}, for example. One can try to "correct" this strengthening by devices such as insisting that for any two elements there is a set that contains one and not the other (which WLOG is the case), but such corrections don't get one very far. What I am asking for is examples, either small ones or ones that are constructed theoretically, of union-closed families that defeat more sophisticated strengthenings of the original conjecture. I'm fairly sure they are out there but I am not an expert on this problem so I don't know them myself. Apologies in advance if this resembles an existing question (which it feels as thought it easily might). But I've looked and not found anything.

论坛问题帖 | has_answers | gowers | 2010-11-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018371 | MO-475220

A generalization of Barrow's inequality

More than seven years ago. I posted this problem in stackexchange: Let ABC be a triangle, P be arbitrary point inside of ABC. Let $A_1B_1C_1$ be the tangential traingle of ABC. Let A', B', C' be three be arbitrary points on B_1C_1, C_1A_1, A_1B_1 respectively. Let internal angle bisectors of three angles $\angle B'PC'$, $\angle C'PA'$, $\angle A'PC'$ meet BC, CA, AB at A'', B'', C'' respectively. Then $$PA'+PB'+PC' \ge 2(PA''+PB''+PC'')$$. Equality hold iff ABC be an equilateral triangle and $\angle B'PC'= \angle C'PA'=\angle A'PC' =120^0$ When $A' \equiv A, B' \equiv B, C' \equiv C$ we have Barrow's inequality I am looking for a proof the conjecture. See also: Erdős-Mordell inequality Barrow's inequality

论坛问题帖 | has_answers | Đào Thanh Oai | 2024-07-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018372 | MO-475326

Open problems in differential algebra and affine algebraic geometry

I am currently doing a PhD in differential algebra and affine algebraic geometry at the University of Buenos Aires. I've been struggling to find a list of interesting and big open problems in these areas. I would be interested in a list of problems or also a list of subtopics that are currently on trend together with a list of reasons about their importance/influence in the area.

论坛问题帖 | no_answers_at_retrieval | Leonardo Lanciano | 2024-07-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018373 | MO-475413

Papers showing two open problems are connected

I'm currently writing an article in which I connect two open problems from different fields of mathematics (group theory and combinatorics). While writing the introduction, I was trying to think of other examples of two problems from different fields of mathematics that are connected. I could only think of the modularity theorem (formerly known as the Taniyama-Shimura-Weil conjecture) and Fermat's Last Theorem. Both are now solved, but historically, it was first shown that they are related; more specifically, that a negative answer to Fermat's Last Theorem would yield a negative answer to the Taniyama-Shimura-Weil conjecture. My question is: What are other examples of such connections? Edit: based on coLaidernotte's suggestion, the type of connections I'm looking for can better be described as follows: examples of two (or more) open problems that are not considered related at first, but which are eventually proved in a uniform way or proved to be closely related.

论坛问题帖 | no_answers_at_retrieval | P. Senden | 2024-07-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018374 | MO-475733

Solving functional analysis problems by using Algebraic geometry

I am thinking about some open problems in nonlinear functional analysis and I just wanted to know if there are any problems that have been solved by using Algebraic geometry techniques in these fields.

论坛问题帖 | no_answers_at_retrieval | Abdullah M Al-jazy | 2024-07-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018375 | MO-477402

Poisson binomial conjecture

Let $X_i\in\{0,1\}$ be mutually independent and distributed according to $\mathrm{Bernoulli}(p_i)$ and similarly, $Y_i\sim\mathrm{Bernoulli}(q_i)$, for some parameters $p,q\in[0,1]^n$. Put $X:=\sum_{i=1}^n X_i$ and $Y:=\sum_{i=1}^n Y_i$; both take values in $\{0,\ldots,n\}$. Assume $p_i\ge q_i$ and even n, put $\gamma:=\frac1n\sum_{i=1}^n(p_i-q_i)$, and let $\bar X\sim\mathrm{Binomial}(n,1/2+\gamma/2)$ and $\bar Y\sim\mathrm{Binomial}(n,1/2-\gamma/2)$. Conjecture: $$\max_{k\ge 0} | \mathbb{P}(X\ge k) - \mathbb{P}(Y\ge k) | \ge \mathbb{P}(\bar X\ge n/2) - \mathbb{P}(\bar Y\ge n/2) .$$ This would imply, in particular, a positive resolution of the conjecture in Minimizing total variation under constraint

论坛问题帖 | has_answers | Aryeh Kontorovich | 2024-08-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018376 | MO-477603

Inequality $(a_1^x+a_2^x+\cdots+a_n^x)^y \ge (a_1^y+a_2^y+\cdots+a_n^y)^x$

Conjecture: Let $a_1, a_2, \cdots , a_n>0$ and $y \ge x $ then $$(a_1^x+a_2^x+\cdots+a_n^x)^y \ge (a_1^y+a_2^y+\cdots+a_n^y)^x$$ Equality iff $x=y$ Is the conjecture right? Have you ever seen this inequality before?

论坛问题帖 | has_answers | Đào Thanh Oai | 2024-08-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018377 | MO-47974

Artin's conjecture for $n=2$

I am interested in the following question: Is it known that 2 is a primitive root modulo p for infinitely many primes p? There is some information about Artin's conjecture in Wikipedia. I need to know if it is up-to-date and if one can say something about the case $n=2$.

论坛问题帖 | has_answers | Kate Juschenko | 2010-12-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018378 | MO-480443

Have the open Questions 1 and 2 from Section 7 of the paper "Integrals with values in Banach spaces" been answered?

Some context: I had previously asked the post below on MSE, but someone suggested I ask it here and delete the original post. In section 7 of the paper Integrals with values in Banach Spaces and locally convex spaces by Piotr Mikusiński (2014), two open questions are presented: Question 1: If $f: X \to E$ (where E is a complete locally convex space) and $\pi_\alpha f$ (where $\{|\cdot|\}_\alpha$ is a family of seminorms for the topology of E and π_α the projection from E to the quotient normed space E_α) is Bochner integrable for every $\alpha \in I$, is f integrable? Question 2: Is the space $L^1(X, \mu, E)$, which consists of E-valued functions that are integrable with respect to a measure μ on X, complete under the topology induced by the family of semi-norms $|\cdot|_\alpha$? Have these questions been answered since the paper was written? Any updates or references would be appreciated.

论坛问题帖 | no_answers_at_retrieval | Sam | 2024-10-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018379 | MO-480839

Are local complete intersections of small codimension necessarily (global) complete intersections?

Hartshorne's 1974 conjecture states that a smooth closed subvariety X of $\mathbb{CP}^r$ of dimension $>2/3r$ is necessarily (globally) a complete intersection. Is anything interesting known about this conjecture? Can it (maybe, in a weaker form) be true if one only assumes X to be a local (set-theoretic) complete intersection? If any counterexamples (that is, local set-theoretic complete intersections of small codimension that are not complete intersections) are known, I would like to know their lower cohomology.

论坛问题帖 | no_answers_at_retrieval | Mikhail Bondarko | 2024-10-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018380 | MO-481090

A new Conjecture at OEIS sequence A376842

Here is a new conjecture of mine from the appendix of an unpublished manuscript currently under review. Let $b \in \mathbb{Z}^+$ and assume that n is an integer greater than 1 and not a multiple of 10. Let $^b{n}$ be $n^{n^{{\cdots}^n}}$ (b-times $-$ see tetration). Then, let us call $\bar{b}$ the smallest hyperexponent of n such that the congruence speed of n is constant (i.e., $\bar{b} := \bar{b}(n)$) and matches the result of Equation (16) of This Paper. Finally, in radix-10, we define as asymptotic phase shift of n (APH(n)) the n-th term of the OEIS sequence A376842 (i.e., A376842(n)), where, if n is not a multiple of 10, the value of the most significant digit of each A376842(n) corresponds to the congruence class modulo 10 of the difference between the rightmost non-stable digit of $^\bar{b}{n}$, say the m-th by counting digits from right to left, and the m-th rightmost digit of $^{\bar{b}+1}{n}$ (which is always a stable digit since the constant congruence speed of n necessarily belongs to $\mathbb{Z}^+$). E.g., if $n =

57$ is given, then we calculate APS(57) by looking at the following sequence of iterations: $\ldots 00000000000000000000000000057$ (height $b = 1$), $\ldots 860230813777543677048556880 57$ (height $b = 2$), $\ldots 443464175207610790048796880 57$ (height $b = 3$), $\ldots 666446504611015011568796880 57$ (height $b = 4$), $\ldots 671511645122477971568796880 57$ (height $b = 5$), $\ldots 190953270434557971568796880 57$ (height $b = 6$), $\ldots 328651320274557971568796880 57$ (height $b = 7$), $\ldots 933653640274557971568796880 57$ (height $b = 8$), $\ldots$ Since the constant congruence speed of 57 is 3 and $\bar{b}(57)=1$, we can finally state that APS$(57) = 2684 = A376842(57)$. Question: Does the following conjecture hold for every $n \in \mathbb{N}-\{1\} : n \not\equiv 0 \pmod{10}$? Conjecture: For any given $n \in \mathbb{N}-\{1\} : n \not\equiv 0 \pmod{10}$, A376842$(n) \in$ $\{2, 4, 5, 6, 8, 9, 19, 28, 46, 64, 82, 1397, 1793, 2486, 2684, 3971, 4268, 4862, 6248, 6842, 7931, 8426, 8624\}.$ Note that each term of the set $\{2, 4, 5, 6, 8, 9, 19, 28, 46, 64, 82, 1397, 1793, 2486, 2684, 3971, 4268, 4862, 6248, 6842, 7931, 8426, 8624\}$ corresponds to some value of n (a direct proof is included in Section 3 of the mentioned manuscript). Moreover, if $2 \mid n$ or $5 \mid n$ (and $n \neq 5$) we have that $\bar{b}$ does not exceed 3 while APS$(n) \not\in \{9,19,1397,1793,3971,7931,\}$.

论坛问题帖 | no_answers_at_retrieval | Marco Ripà | 2024-10-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018381 | MO-48141

A Game of Knights and Queens

Let $m,n,u,v \in \mathbb{N}$ be parameters with $m,n \geq 3$. Suppose two players play a game on a $m \times n$ chess board and we denote the squares of the board by the set of points $ (i,j) $ such that $1 \leq i \leq m, 1 \leq j \leq n $. Player 1 has u queens, and player 2 has v knights. The initial configuration is some given subset of the board where no piece is placed at a position that can be immediately attacked by another piece of the opposing player, and the player with at least one piece when the game terminates is the winner. Both players must make a move when it is their turn. The game terminates when either player captures all of the pieces of the opposing player. If Player 1 goes first, under what circumstances does either player have a winning strategy? What can be said about the probability as a function of the parameters given with both players playing optimally, that the game will terminate in finitely many steps? I have obtained the result for $m = 3$ and arbitrary n with $u= v = 1$. Ed Dean proved that Player 1 always has a winning strategy for arbitrary m,n and $u = v = 1$ (see below), and he sketched a winning strategy for Player 1 in the $u = 1, v = 2$ case. He also gave an argument for the lack of a winning strategy for Player 1 for the $u = 1, v = 3$ case. All other cases are still unknown as of yet. Edit: Previous suggestion on a way to show in the $u = v = 1$ case that the knight can avoid capture for sufficiently large m,n is removed. Edit 2: Thanks to Ed Dean for resolving some cases, as indicated above. Edit 3: Included a new question regarding a related probability distribution. Edit 4: Moved edit 3 to a new question: A random variable in a game of knights and queens

论坛问题帖 | has_answers | Stanley Yao Xiao | 2010-12-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018382 | MO-481786

Does a matrix ring over a ring satisfy the Koethe conjecture if the coefficient ring itself satisfies the Koethe conjecture?

I just want to know whether the following statement is true or false. If R is a ring satisfying the Koethe conjecture, then the matrix ring over R also satisfies the Koethe conjecture. Or is it equivalent to the Koethe conjecture itself? I will be grateful if you can link to a paper or a book.

论坛问题帖 | no_answers_at_retrieval | Eunnaya First | 2024-11-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018383 | MO-48203

On Sorli's Conjecture Re: OPNs (Circa 2003)

In the PhD dissertation titled "Algorithms in the Study of Multiperfect and Odd Perfect Numbers" (hyperlinked here) and completed in 2003, Ronald Sorli conjectured that the exponent k on the Euler prime p for an odd perfect number $N = (p^k)(m^2)$ is one (i.e. we can drop k). Assuming Sorli's conjecture is true, does anyone know if there exist (any) "effective" results (pardon my use of the term, I just could not think of a better word) in the literature, particularly with respect to relations between the Euler prime p, the exponent k and the number $\sqrt{\frac{N}{p^k}}$? I have, so far, only been able to get hold of Paolo Starni's article titled "Odd Perfect Numbers: A Divisor Related to the Euler's Factor". In particular, note that Sorli's conjecture implies the following relations: $$I(p^k) = I(p) = \frac{p+1}{p}$$ $$I(m^2) = \frac{2}{I(p)} = \frac{2p}{p + 1}$$ which, in turn, gives the (trivial) algebraic identity: $$\frac{1}{p} = \frac{1}{p+1} + \frac{1}{p}\left(\frac{1}{p+1}\right)$$ where p is the Euler prime (i.e. p^k is the Euler's Factor) and $$I(x) = \frac{\sigma(x)}{x}$$ is the abundancy index of x. [Edit (September 18 2013) - Per Professor Beasley's paper titled "Euler and the Ongoing Search for Odd Perfect Numbers" from this hyperlink: Before proceeding with Euler's proof, we pause to note that his result was not quite what Descartes and Frenicle had conjectured, as they believed that $k = 1$, but it came very close. In fact, current research continues in an effort to prove $k = 1$. For example, Dris has made progress in this direction, although his paper refers to Descartes' and Frenicle's claim (that $k = 1$) as Sorli's conjecture; Dickson has documented Descartes's conjecture as occurring in a letter to Marin Mersenne in 1638, with Frenicle's subsequent observation occurring in 1657. End Edit.]

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2010-12-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018384 | MO-48209

Reference Request - Sharp Estimates for a Logarithmic Sum

Can anybody suggest a good (e.g. "non-technical") introduction to estimating bounds for logarithmic sums of the form $$\sum_{i=1}^{r}{{\alpha_i}{\log(q_i)}}$$ where the $$\alpha_i$$ are positive integers (not necessarily distinct) and the $$q_i$$ are odd primes?

The reason why I ask this question is because I am trying to estimate (both lower and upper) bounds for the logarithm of a number-theoretic function, specifically $$\sigma_{1}(N)$$ I was able to show (in 2008) that $$\sigma_{1}({q_i}^{\alpha_i}) \le \frac{2}{3}\frac{N}{{q_i}^{\alpha_i}}$$ for all $i = 1, 2, ..., r = \omega(N)$, where N is an OPN (i.e. Odd Perfect Number) and ${q_i}^{\alpha_i} || N$ for all i. You will be able to get an upper bound for the logarithmic sum given above, by first taking logarithms of both sides of the inequality, then summing over all i. Unfortunately, for the "numbers" N that I am considering, the current literature (on OPNs) do not point to an approach "strong enough" to prove nonexistence of such "numbers" N. (This is because the upper bound alluded to in the previous paragraph is still (of course) dependent on $r = \omega(N)$). (And that is the reason why --) I'd be particularly interested in an analytic-number-theoretic approach. =) (Thanks to Gerry Myerson for encouraging this "clarification".)

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2010-12-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018385 | MO-48215

Re: Mordell's equation $y^2 = x^3 + k$ and perfect numbers

I have already tried a somewhat exhaustive search of the literature, but couldn't find anything close to the problem that I am working on. My question is: When does Mordell's equation $$Y^2 = X^3 + K$$ have only finitely many solutions over the field of rational numbers, if we allow K itself to be a rational number? I've seen a "criterion" (i.e. a set of sufficient conditions) related to the class number of the (real/imaginary) quadratic field $\mathbb{Q}(\sqrt{K})$, but it is limited only to K being either 1 or 2 modulo 4. (The actual "criterion" (as stated in the Japanese[?] paper that I allude to) is: Mordell's equation $Y^2 = X^3 + K$ has finitely many solutions in $\mathbb{Q}$ if (1) $-K$ is not of the form $3t^2 + 1$ or $3t^2 - 1$; and (2) $K \equiv 1 \pmod 4$ or $K \equiv 2 \pmod 4$; and (3) 3 does not divide the class number of the (real/imaginary) quadratic field $Q(\sqrt{K})$.) Edit: Please refer to this hyperlink for more information as to the context of the previous "criterion". These have since been refuted by Kevin Buzzard (@Kevin - thank YOU!). Thanks to Kevin for pointing out some of the subtle errors in my original post, particularly in the third condition. I was considering the case $K > 0$ (i.e. for real quadratic fields). Now for the context: Let $$Y = W + Z$$ and $$X = WZ$$ where W and Z are defined as: $$W = I(p^k) = \frac{\sigma_{1}(p^k)}{p^k}$$ $$Z = I(m^2) = \frac{\sigma_{1}(m^2)}{m^2}$$ Let $$N = {p^k}{m^2}$$ be a perfect number. (At this point, we don't have to distinguish between even or odd N because the Euclid-Euler model for perfect numbers fits both cases. That is, the Eulerian form for an odd perfect number and the Euclidean form for an even perfect number have very similar multiplicative forms.) We "know" that the exponent k allows us to distinguish between even and odd N in the sense that: (1) If k = 1, then N is even. (2) If k > 1, then N is odd. hyperlink Perhaps the important feature that should be used to distinguish between even perfect numbers $M=2^{q-1}(2^q - 1)$ and odd perfect numbers $N=p^k m^2$ (where $2^q - 1$ and p are the Mersenne and special primes, respectively) is the index of a perfect number: (1) $\gcd(2^{q-1},\sigma(2^{q-1}))=1$ (2) $\gcd(m^2,\sigma(m^2))>1$ There is a known formula for computing $\gcd(m^2,\sigma(m^2))$. For example, if $k=1$, then it is equal to $2m^2 - \sigma(m^2)$. Thus, a (possibly) feasible and modern approach to the OPN problem (i.e. determining nonexistence or otherwise) will be to try establishing a finiteness result first (for particular values of K). In other words, checking for finiteness of OPNs amounts to checking for finiteness of solutions for Mordell's equation $$Y^2 = X^3 + K$$ for particular values of K. And you will only have to check for values of K in the range $[50, 399]$ (for a total of 350 elliptic curves), per the previous answer to this MathOverflow question. K falls in that range because the sum $$Y = W + Z$$ is known to lie in the open interval $(57/20, 3)$. Of course, the "juicy" implication is that: If you will be able to find a condition (e.g. equation, inequality, etc.) relating k to K and you are also able to further show that the number of solutions to the corresponding Mordell equation $Y^2 = X^3 + K$ is finite for all such K, then it would follow that there are only finitely many perfect numbers (odd and even). Disclaimer: This is a "naive" approach based on my current understanding of elliptic curve theory. I am well-aware, of course, that the rationals are dense over the real numbers. (Edit: In addition, the abundancy indices and the abundancy outlaws are both dense over the rationals.) Which is why I was kind of surprised that there is no need to assume ("strict") rationality (i.e. $K \in \mathbb{Q} \setminus \mathbb{Z}$) for K when checking for finiteness of solutions to Mordell's equation.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2010-12-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018386 | MO-483085

Conjecture about Euler quotients related to non-Wieferich numbers $W(n)=\frac{2^n+1}{3}$

For odd natural n define the Euler quotient: $$ a(n)=\frac{(2^{\phi(n)}-1) \bmod n^2}{n}=\frac{2^{\phi(n)}-1}{n} \bmod n$$ $a(n)=0$ is n being Wieferich number (not necessarily prime). For odd n, define $W(n)=\frac{2^n+1}{3}$. If $W(n)$ is prime, n is Wagstaff prime. Conjecture 1 $a(W(n)) \bmod 2=1$ for all odd $n>1$. Conjecture 2 $a(W(n))$ is nonzero all odd $n>1$. The Conjecture 1 will imply all $a(W(n))$ are non-zero and thus $W(n)$ non-Wieferich numbers. If n is prime all prime factors of $W(n)$ are congruent to 1 modulo n, and we will get infinite sequence of non-Wieferich numbers with unbounded smallest prime factor. Experimentally the conjecture holds for n up to 219 and factorization for computing the Euler totient is expensive. Computing the parity of other sequences gives random parity. Added per request for the code in comments. Here is sagemath code which you can run in a browser def W(n): return (2**n+1)//3 def V(n): return 2**n+1 def a(n,B=2): return lift((Integers(n**2)(B))**(euler_phi(n))-1)/n LAST=50 #limit for n for n in range(3,LAST,2): v1=n*(-V(n) + a(V(n))) + euler_phi(V(n)) v2=n*(-W(n) + a(W(n))) + 3*euler_phi(W(n)) print(n,v1==0,v2==0) #,a(V(n))/a(W(n))) """ pari/gp code ? a(N)=lift(Mod(2,N^2)^eulerphi(N) -1)/N %6 = (N)->lift(Mod(2,N^2)^eulerphi(N)-1)/N ? a(2^101+1) %7 = 25185665390673072597393245202 53 ? """

论坛问题帖 | has_answers | joro | 2024-11-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018387 | MO-483874

What is the status of this conjecture on symplectic forms "standard-at-infinity" on $\mathbb{R}^{2n}$?

In McDuff and Salamon's Introduction to Symplectic Topology, the following open problem is mentioned. Problem 50 (Standard-at-infinity) Let $n \ge 3$ and let ω be a symplectic form on $\mathbb{R}^{2n}$ that agrees with ω_0 on the complement of a compact set. Is $(\mathbb{R}^{2n}, \omega)$ symplectomorphic to $(\mathbb{R}^{2n}, \omega_0)$? In case it isn't clear, ω_0 is the standard symplectic form. The authors go on to mention some related results: The same question in dimension four has a positive answer by a celebrated theorem of Gromov [287]. A theorem of Floer-Eliashberg-McDuff [451] asserts that if (M, ω) is a symplectic manifold with $\pi_2(M) = 0$ that is symplectomorphic to $(\mathbb{R}^{2n}, \omega_0)$ outside of a compact set, then M is necessarily diffeomorphic to $\mathbb{R}^{2n}$. Mark McLean and others constructed many examples of symplectic structures on $\mathbb{R}^{2n}$ that are convex at infinity, but are not symplectomorphic to $(\mathbb{R}^{2n}, \omega_0)$ (see McLean [474, 475], Abouzaid-Seidel [6], and Seidel [576]). Since another open problem concerning the symplectic topology of Euclidean space was recently resolved, I became curious if any progress has been made on the problem stated above. I suppose this will likely be work performed since 2017, since that is when the most recent edition of McDuff-Salamon was published. I am particularly interested in the special case of $\mathbb{R}^6$. My questions are: What progress (if any) has been made beyond what is mentioned above? Is this problem known to be "very hard" by experts in the field? Does anyone have intuition on whether this is anything close to approachable by an average graduate student? References [6] Abouzaid, M., Seidel, P. (2010). Altering symplectic manifold by homologous recombination. http://arxiv.org/abs/1007.3281v3 [287] Gromov, M. (1985). Pseudo holomorphic curves in symplectic manifolds. Inventiones Mathematicae, 82, 307-347. [451] McDuff, D. (1991). Symplectic manifolds with contact type boundary. Inventiones Mathematicae, 103, 651-671. [474] McLean, M. (2009). Lefschetz fibrations and symplectic homology. Geometry & Topology 13, 1877-1944. http://arxiv.org/abs/0709.1639 [475] McLean, M. (2012). The growth rate of symplectic homology and affine varieties. Geometric and Functional Analysis, 22, no. 2, 369-442. http://arxiv.org/abs/1109.4466 [576] Seidel, P. (2011). Simple examples of distinct Liouville type symplectic structures. Journal of Topology and Analysis, 3 no. 1, 1-5. http://arxiv.org/abs/1011.0394v2

论坛问题帖 | no_answers_at_retrieval | Brendan Langfield | 2024-12-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018388 | MO-484182

Maximum number of connected components of surfaces in three dimensions, what is known?

Part of Hilbert's 16th problem is: It seems to me that a thorough investigation of the relative positions of the upper bound for separate branches is of great interest, and similarly the corresponding investigation of the number, shape and position of the sheets of an algebraic surface in space (...) The solution for a quartic surface is contained in Kharlamov: The topological type of nonsingular surfaces in RP3 of degree four. What is the status of this problem? In particular, is there a conjectural upper bound for the maximum number of connected components of a smooth degree d surface in $\mathbb{RP}^3$?

论坛问题帖 | has_answers | user548513 | 2024-12-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018389 | MO-484415

A simple group with 3 generators

All finite simple groups are generated by two elements, though apparently the proof is tens of thousands of pages long. Here’s an obvious follow-up: What’s an example of a finitely presentable simple group with 3 generators (and not 2)?

论坛问题帖 | has_answers | Geoffrey Irving | 2024-12-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018390 | MO-484586

Conjecture on a cyclic quadrilateral associated with central line of triangle

Using computer I found a conjecture as follows (click to check by geogebra): Conjecture: Let A, B, C, D lie on a circle such that A, B, C, D are not formed an Isosceles trapezoid. Let AB meets CD at F and AD meets BC at E. Define $\ell(n, m, \triangle T)$ is a line through triangle center $X(n)$, $X(m)$ of triangle $\triangle T$, then $\ell(n, m, \triangle ABE)$, $\ell(n, m, \triangle CDE)$, $\ell(n, m, \triangle ADF)$, $\ell(n, m, \triangle BCF)$ form a cyclic quadrilateral for every positive integers n, m ($n \neq m$). Where $X(n)$ refer to Kimberling triangle center. Question: I am looking for a proof the conjecture above.

论坛问题帖 | has_answers | Đào Thanh Oai | 2024-12-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018391 | MO-484639

Conjecture closed form of summation over an integer lattice

Conjecture: $$\forall \Lambda,\ \exists P(x)\in \mathbb{Z}[x],\ S(\Lambda):=\sum_{k\in\Lambda}\prod_{j=1}^{n}\frac{1}{1+k_{j}^2}=\frac{\pi^{n}}{\sinh^{n}(\pi)\operatorname{d}(\Lambda)}P\left(\cosh\left(\frac{\pi}{\operatorname{d}(\Lambda)}\right)\right)$$ where $\Lambda=\bigl\{\sum_{j=1}^{n}a_j v_j|\ a_j \in \mathbb{Z}\bigl\}$, $V=\{v_1,\ldots,v_n\}$, $\operatorname{d(\Lambda)}=|\det(V)|\ $, V is an integer matrix. The conjecture is known to be true for $n=1$ or $\operatorname{d}(\Lambda)=1$, I also checked some few cases when $n=2$: $$\operatorname{d(\Lambda)}=2\implies P(x)\in \{4x^4-4x^2+2,\ 4x^4-2x^2\} \tag{1}$$ This corresponds to 2 lattice structures with $\operatorname{d(\Lambda)}=2$ in dimension 2. And for $V=\{(4,1)^T,(1,1)^T\}\implies P(x)=16x^6-24x^4+11x^2$; $V=\{(3,1)^T,(2,2)^T\}\implies P(x)=64x^8-128x^6+88x^4-24x^2+4$ As I mentioned in this previous question, I don't have a good way to compute this sum, all calculations and simplifications was done by calculator and each result took me quite a lot of time. Unfortunately, I cannot compute the sum when $n\geq 3$ because it is too much for my calculator to handle. However, partial calculation ($n=3$) suggested that closed form is achievable. Update Slightly deviated from

the main goal, I found out that $S(\Lambda)$ has a nice integral representation: $$\sum_{k\in \Lambda}\int_{\mathbb{R}^n}d^{n}x\exp{(-\|x\|_{1}+i\langle k,x \rangle)}=2^{n}S(\Lambda)$$ After some investigation, I think that $\sum_{k\in \Lambda} e^{i\langle k,x \rangle}$ is a distribution similar to Dirac comb, I couldn't find its formula though. If the formula of distribution is known, the integral can be turned into something like a geometric sum. I also noticed if that is the case, $\operatorname{d}(\Lambda)$ should be the "angular frequency". () (Edited) My question is: How do we find $$\sum_{k\in \Lambda} e^{i\langle k,x \rangle}$$ Does this give an answer to the conjecture?

论坛问题帖 | has_answers | user500489 | 2024-12-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018392 | MO-485389

Is it possible to discover a "Universal formula" that generates and generalizes all odd Collatz numbers?

I am just an ordinary student, and I have never had the chance to ask this question to mathematicians. Perhaps this is the first opportunity I have to ask it here. When we refer to "Collatz numbers", we mean all natural numbers that eventually reach 1 under the Collatz process. By odd-step Collatz numbers, we consider only the odd numbers that appear at each step of the sequence, skipping all even numbers. For example, for $n=4$, one possible sequence of odd-step Collatz numbers is: $$11 \rightarrow 17 \rightarrow 13 \rightarrow 5 \rightarrow 1 $$ Similarly, for $n = 6$, another example sequence could be: $$ 19 \rightarrow 29 \rightarrow 11 \rightarrow 17 \rightarrow 13 \rightarrow 5 \rightarrow 1 $$ Now, my question is: Is it possible to find a "Universal formula" (or something "magic formula") that generates all sequences of odd numbers with exactly n odd-steps in the Collatz process? For instance, consider a function like $f(n) = 2^n$. While this generates a sequence, it does not satisfy the criterion of producing a complete sequence of odd-step Collatz numbers, as it results in only one odd step. I use $f(n) = 2^n$ merely as an example to illustrate the kind of formula I am seeking. To summarize, for a given n, is there a general formula - similar to $f(n) = 2^n$ - that universally produces all sequences of n-step odd Collatz numbers? Has such a formula been discovered or studied by mathematicians? Thank you so much . I see that I needed to phrase the question more clearly. Let me rewrite what I mean by "Universal Formula". For example, when $n = 1$ is a finite number, we have a formula like $\dfrac {4^n-1}{3}$. We can go backwards and construct such a formulas for $n = 2$. But, it would be a great experience for me to learn whether a "formula" that directly depends on n exists or - if it is already known - to understand it .

论坛问题帖 | has_answers | nonuser | 2025-01-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018393 | MO-487645

The behaviour of $\mathsf{HOD}^{L[x]}$ on a cone

In "Introduction to Q-Theory" by Kechris-Martin-Solovay in the proceedings of the Cabal Seminar 1979 - 1981, back when it was believed axioms up to I_3 could be compatible with Δ^1_2-nondeterminacy, the following ultraproduct is introduced: $$\prod_{d \in \mathcal{D}} \mathsf{HOD}^{L[d]} / \mu$$ where μ is the Martin (cone) measure on Turing degrees $\mathcal{D}$, which is an ultrafilter assuming AD. Since the Martin measure is ω_1-complete, this ultraproduct is well-founded and its behaviour can be analyzed by analyzing the behaviour of $\mathsf{HOD}^{L[x]}$ for a cone of reals x (as if $x \in d \in \mathcal{D}$ then $L[x] = L[d]$). They write "this ultraproduct is understood as taken within $L(\mathbb{R})$", although I'm not sure how you would prove that every $f \in \prod_{d \in \mathcal{D}} \mathsf{HOD}^{L[d]}$ agrees with such an $f \in L(\mathbb{R})$ on a cone (maybe they just mean that you should compute it in $L(\mathbb{R})$ when you don't want to assume AD in V). Regardless, back then, it was expected that this would be "$\mathcal{M}^\infty$", an ultimate generalization of Mitchell's $L[\mathcal{U}]$ models that contains all Δ^1_3 reals. Indeed, they were partially right! In FSIT, the theory of inner models with a few Woodin cardinals was introduced, and M_1, the smallest inner model with a Woodin cardinal, was isolated as being a "canonical" Σ^1_3-correct model (a higher analogue of the Σ^1_2-correctness for $M_0 = L$ proven by Shoenfield absoluteness). In that paper, they prove $\mathcal{M}^\infty \cap \omega^\omega = M_1 \cap \omega^\omega = Q_3$ (the set of reals Δ^1_3 in a countable ordinal) and, if $\pi: \prod_{d \in \mathcal{D}} \mathsf{HOD}^{L[d]} / \mu \to \mathcal{M}^\infty$ is the Mostowski collapsing map, that $\pi(\langle \omega_1^{L[d]}: d \in \mathcal{D} \rangle)$ is measurable in $\mathcal{M}^\infty$. However, this 2018 answer by Goldberg indicates that the behaviour of $\mathsf{HOD}^{L[x]}$ is still mostly unknown. I am asking: Can $M_1^{L(\mathbb{R})} = (\mathcal{M}^\infty)^{L(\mathbb{R})}$, assuming $\mathrm{AD}^{L(\mathbb{R})}$? Has anything changed in the last 7 years? Apparently, $\mathsf{HOD}^{L[x][G]}$, for G an $L[x]$-generic for $\mathrm{Coll}(\omega, < \kappa)$, where κ is $L[x]$'s least inaccessible cardinal, was analyzed by Woodin, and I recall this 2020 paper by Müller-Sargsyan generalizing this to $\mathsf{HOD}^{M_n(x)[G]}$, but I still cannot find anything regarding $\mathsf{HOD}^{L[x]}$. Are there any papers describing the guess, mentioned in the above MO answer, that $\omega_2^{L[x]}$ is Woodin in $\mathsf{HOD}^{L[x]}$ on a cone (and hence $\pi(\langle \omega_2^{L[d]}: d \in \mathcal{D} \rangle)$ is Woodin in $\mathcal{M}^\infty$) due to iterated ultrapowers, or the result of Woodin, mentioned in "Introduction to Q-Theory" that there is a single, Sacks-like real s over $L(\mathbb{R})$ with $(\mathcal{M}^\infty)^{L(\mathbb{R})} = \mathsf{HOD}^{L[s]}$? Are these all in handwritten notes and personal communications, or have some facts been published since 1981?

论坛问题帖 | no_answers_at_retrieval | Jayde SM | 2025-02-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018394 | MO-488756

Conjecture on the preservation of unfragmented sets under element addition

A set A_k contains k elements $a_i, i=1..k$. The a_i are incomparable (For example, functions, matrices, etc. they cannot be connected using > or <), but an operation $\otimes$ can be defined such that $a_i \otimes a_j$ holds for any a_i and a_j and satisfies the commutative law, and $a_i \otimes a_j$ are comparable. Therefore, from $a_i, i=1..k$, we can obtain $k(k-1)/2$ elements $b_{ij} \equiv b_{ji} \equiv a_i \otimes a_j \equiv a_j \otimes a_i$. These b_{ij} form a new set B_k. By sorting all elements in B_k from smallest to largest, we obtain the indices of the smallest $k-1$ elements, which total $2k-2$ numbers. Among these, the mode is s. If the count of s is $k-1$, then remove a_s from set A_k to obtain a new set A_{k-1} and repeat the above steps; if the count of s is less than $k-1$, then terminate. If the above operation removes all elements from A_k to obtain A_0, then A_k is called "fragmented"; otherwise, it is "unfragmented". Conjecture: For an unfragmented set A_k, after adding any element that satisfies the definition of A_k to A_k, A_k remains unfragmented. Please prove or disprove this conjecture.

论坛问题帖 | no_answers_at_retrieval | Yu-Fu Shen | 2025-03-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018395 | MO-48908

Is the Invariant Subspace Problem interesting?

There's an amusing comment in Peter Lax's Functional Analysis book. After a brief description of the Invariant Subspace Problem, he says (paraphrasing) "...this question is still open. It is also an open question whether or not this question is interesting." To avoid lengthy discussions involving subjective views about what makes math interesting, I'd simply like to know if there are examples of math papers out there that begin with something like, "Suppose the invariant subspace problem has a positive answer..." Of course, papers that are about the ISP itself don't count!

论坛问题帖 | has_answers | William DeMeo | 2010-12-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018396 | MO-489375

Why is Erdős' conjecture on arithmetic progressions not discussed much, and is there an active pathway to its resolution?

Erdős' conjecture on arithmetic progressions (also known as the Erdős-Turán conjecture) is a major open problem in arithmetic combinatorics. It asserts that if the sum of the reciprocals of the members of a set A of positive integers diverges, then A contains arbitrarily long arithmetic progressions. Despite its fundamental nature and other well known theorems being a special case of it, the conjecture does not seem to be widely discussed in literature as much as it should. While I know some weaker results have been proven,the full conjecture remains open. My questions are: Are researchers actively working towards its resolution, and is there a specific pathway being pursued? What are the key difficulties that prevent a proof using current methods? Is there any expectation that this problem might be resolved in the near future, or is the problem considered beyond the reach of existing techniques? Any insights on the obstacles, possible strategies, and the current research landscape regarding this conjecture would be greatly appreciated

论坛问题帖 | has_answers | HasIEluS | 2025-03-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018397 | MO-48953

Perfect Numbers - On Mersenne and Euler Primes

Hi, I apologize if there is already an (obvious) answer to my question, but please bear with me for the moment as I find it hard to see a good way to answer this question: In the same way that the Mersenne primes uniquely determine the even perfect numbers (i.e. in the "bijective" sense), do the Euler primes likewise uniquely determine the odd perfect numbers? That is, while an Euler prime obviously maps to a single odd perfect number, it is not obvious to me that an odd perfect number could only map to a single Euler prime. This is because: (1) In the most general case, it is not known whether squares are friendly or solitary. (2) It is not even known if the smallest possible Euler prime 5 does divide an odd perfect number. (3) It is conceivable that two distinct numbers P and Q which satisfy a particular number-theoretic equation do not necessarily have to satisfy $\gcd(P, Q) = 1$. I hope you guys can help me out on this one. Establishing this very last gap will provide a proof for the implication: Assuming Sorli's Conjecture for OPNs is true, then there are no OPNs.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2010-12-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018398 | MO-489799

Reference request: Open problems about finite free products of finite groups

I'm working with finite free products of finite groups, i.e. a group G given by $$G = F_1 \ast \ldots \ast F_n$$ where each F_i is finite. Do you know of any open problems as well as references about them about these kinds of groups? In particular, I'm interested in finitely generated subgroups of such groups and related conjectures.

论坛问题帖 | has_answers | Eduardo Magalhães | 2025-03-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018399 | MO-491675

Furstenberg's topological proof method (of prime infinitude) might be more powerful than initially thought. $\sigma_0(\frac{p_n - 1}{4})$ odd i.o.?

Let $\sigma_0(n)$ be from number theory, i.e. the total number of divisors of an integer n. Let p_n denote the nth prime number. Let $S =$ the set of prime numbers $p \in 4\Bbb{Z} + 1$ such that $\sigma_0(\frac{p - 1}{4})$ is odd. Now generate the multiplicative submonoid M of $(\Bbb{Z}, \times)$ with $M = \langle 1,S \rangle$. Take as open topological basis the "Furstenberg-like balls" of the form $M\cap (a\Bbb{Z} + r)$, where $p \in S, a \in M$, and $r = 0..a-1$. And more importantly, where $M \cap (a\Bbb{Z} + r)$ is infinite. By convention $0 \notin \Bbb{N}$. Clearly $(M \cap a\Bbb{Z} + r) \cap (M\cap b\Bbb{Z} + s) = M\cap \operatorname{lcm}(a,b)\Bbb{Z} + t$, or the intersection is empty; where t is any element of the intersection. Satisfying the definition of topological basis.

Clearly, we retain the property that any open set is either empty or infinite, which will become important later. And so no complement of a non-empty finite set can be closed. The open ball $a\Bbb{Z} + r \cap M$ is also closed, it's complement clearly being equal to the union of all open balls over the residues of $s \neq r$ modulo a. Just as in Furstenberg's prime infinitude proof. Now if S were finite, then: $$ M\setminus \{1\} = \bigcup_{p \in S} M\cap p\Bbb{Z} $$ Which would lead to a contradiction, because the set of the right is closed and the set on the left is a complement of a finite non-empty set. As long as the equation is true. Clearly the left hand contains the right hand side. Now let $z \in$ the left hand side. Then $z =\prod_{i = 1}^m q_i^{n_i}$ such that $q_i \in S$. Then clearly $z \in q_i\Bbb{Z}$ for each $i=1..m$, hence $z \in M\cap q_i\Bbb{Z}$ and so is clearly in their union which is a subset of the right hand side. And we're done. Therefore, what I think I've proven is that: $$ \sigma_0(\frac{p_n - 1}{4}) \text{ is odd i.o.} $$ Question. Since this proof is elementary like Furstenberg's topological proof of the infinitude of primes, and it follows it pretty analogously, and there exists three-sentence proofs of the infinitude of primes; can you think of a very simple proof of the fact that $\sigma_0(\frac{p_n - 1}{4})$ is odd infinitely often? Note. I think the above proof might fail only at $M\cap a\Bbb{Z} + r$ always being infinite. So please don't take it as a real proof (yet).

论坛问题帖 | has_answers | TotoposDiagramChaseApp | 2025-04-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018400 | MO-492159

The strong Mersenne conjecture

The strong twin conjecture: For every number $a≥0$, there exist two prime numbers p and $p+2$ such that $$a+4<p<2^{a+4}$$ and it was believed that this conjecture implies the twin conjecture. Then my question: Does there exist a similar stronger form of Mersenne conjecture formulated by using a double inequality. More generally, I am asking about the methods used to get a formulation like the one given in this question.

论坛问题帖 | has_answers | John2000 | 2025-05-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018401 | MO-495160

Can a uniformly-bounded Klein-Gordon wave sweep its nodal set across a rigid obstacle?

Consider Minkowski space $\mathbb R^{3+1}$ and the massive Klein-Gordon equation $$ \square\varphi + m^{2}\varphi = 0 . $$ Let $\Omega\subset\mathbb R^{3}$ be a fixed smooth bounded “obstacle’’ with boundary Σ. Does there exist a one-parameter family of smooth real solutions $\{\varphi_\varepsilon\}_{\varepsilon>0}$ such that: for every compact $K\subset\mathbb R^{3+1}$ the stress-energy tensor $T_{\mu\nu}[\varphi_\varepsilon]$ is bounded independently of ε. The canonical stress-energy tensor for a real massive Klein-Gordon field φ on Minkowski space $(\mathbb R^{3+1},\eta)$, with the mostly-plus signature $\eta = \operatorname{diag}(-1,+1,+1,+1)$, is given by: $$ T_{\mu\nu}[\varphi] = \partial_\mu\varphi\,\partial_\nu\varphi - \tfrac12\,\eta_{\mu\nu} \bigl(\partial^\alpha\varphi\,\partial_\alpha\varphi + m^{2}\varphi^{2}\bigr), \qquad \mu,\nu=0,1,2,3. $$ for each time t the nodal set $N_\varepsilon(t)=\{x:\varphi_\varepsilon(t,x)=0\}$ is a smooth surface; as $\varepsilon\to0$, the family $\{N_\varepsilon(t)\}$ converges (in the Hausdorff sense) to a prescribed smooth isotopy of embedded surfaces Σ_t that starts wholly outside Ω and ends wholly inside, so that the zero level set “passes through’’ the rigid boundary Σ. Informally, can one “vibrate’’ a Klein-Gordon field whose instantaneous nodal surface slides through an obstacle without the field’s energy blowing up? Standard geometric-optics constructions $$ \varphi_\varepsilon=\mathrm{Re}(a\,e^{iS/\varepsilon}) $$ appear to lose the required uniform bounds, while unique-continuation phenomena suggest obstructions when the evolving surface becomes characteristic. Note: The motivation for this problem came from the idea of “phasing” in fictional superhero stories. In particular, in the show The Flash, the main character has the ability to phase through solid objects by vibrating his molecules sufficiently fast.

论坛问题帖 | no_answers_at_retrieval | Alex Cooper | 2025-05-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018402 | MO-496298

Pillai's arithmetical function and primality

Pillai's arithmetical function is defined by $P(n)=\sum_{k=1}^n\gcd(k,n)$. Is it true that $1+P(n)\equiv 0\pmod{n}$ iff n is prime? It was stated without proof in 2014 on the OEIS A018804 and has since been reverted as a conjecture. A question posted last week on MSE didn't get a full proof: the case of squarefree composites with more than 3 factors is completely open. Do you have a reference or a proof for that case?

论坛问题帖 | has_answers | Jon23 | 2025-06-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018403 | MO-496925

Primality test using the Golden Ratio

Background and Motivation The golden ratio, $$ \phi = \frac{1 + \sqrt{5}}{2}, $$ is a well-known irrational constant that appears frequently in geometry, algebra, and in the Fibonacci and Lucas sequences. Though fundamentally irrational and geometric, it surprisingly shows up in discrete number theory as well. Primality testing, on the other hand, is a central topic in number theory with both theoretical and practical importance, especially in cryptography. Most known primality tests rely on modular arithmetic, integer sequences, or algebraic identities - but very few, if any, directly involve irrational constants like ϕ. While experimenting numerically with powers of ϕ, I noticed an unexpected and elegant pattern. The simplicity and strength of this condition - despite its seemingly irrational origin - suggest a potentially deep connection between the golden ratio and prime numbers, possibly via the Lucas or Fibonacci sequences. This motivates the following natural questions: Why does this work? Is there a theoretical explanation rooted in recurrence sequences or algebraic number theory? Could this inspire a new class of primality tests? I am sharing this in hopes of uncovering deeper insights or connections that could explain the phenomenon. The Golden Ratio Primality Test (GRPT) If any positive integer $n>5$ is not divisible by 2 or 5, and satisfies the condition: $$ \lfloor \phi^n \rfloor \mod n = 1, $$ then, $$ n \text{ is

prime}. $$ Here, $$ \phi=\frac{1+\sqrt{5}}{2}.$$ Empirical Evidence I have verified GRPT for the range $5<n\leq 1{,}000{,}000 $ using Wolfram GPT. No false positives or false negatives found. Let's see how this applies to the first few $n>5$ (excluding the multiples of 2 and 5): $$\begin{array}{|c|c|c|} \hline n & \lfloor \phi^n \rfloor & \lfloor \phi^n \rfloor \mod n & \\ \hline 7 & 29 & 1 & \text{Prime}\\ \hline 9 & 76 & 4\neq1 & \text{Not prime}\\ \hline 11 & 199 & 1 & \text{Prime}\\ \hline 13 & 521 & 1 & \text{Prime}\\ \hline 17 & 3571 & 1 & \text{Prime}\\ \hline 19 & 9349 & 1 & \text{Prime}\\ \hline 21 & 24476 & 11\neq1 & \text{Not prime}\\ \hline 23 & 64079 & 1 & \text{Prime}\\ \hline 27 & 439204 & 22\neq1 & \text{Not prime}\\ \hline 29 & 1149851 & 1 & \text{Prime}\\ \hline 31 & 3010349 & 1 & \text{Prime}\\ \hline 33 & 7881196 & 4\neq1 & \text{Not prime}\\ \hline 37 & 54018521 & 1 & \text{Prime}\\ \hline 39 & 141422324 & 17\neq1 & \text{Not prime}\\ \hline 41 & 370248451 & 1 & \text{Prime}\\ \hline 43 & 969323029 & 1 & \text{Prime}\\ \hline 47 & 6643838879 & 1 & \text{Prime}\\ \hline 49 & 17393796001 & 29\neq1 & \text{Not prime}\\ \hline 51 & 45537549124 & 4\neq1 & \text{Not prime}\\ \hline 53 & 119218851371 & 1 & \text{Prime}\\ \hline 57 & 817138163596 & 4\neq1 & \text{Not prime}\\ \hline 59 & 2139295485799 & 1 & \text{Prime}\\ \hline 61 & 5600748293801 & 1 & \text{Prime}\\ \hline 63 & 14662949395604 & 41\neq1 & \text{Not prime}\\ \hline 67 & 100501350283429 & 1 & \text{Prime}\\ \hline 69 & 263115950957276 & 50\neq1 & \text{Not prime}\\ \hline 71 & 688846502588399 & 1 & \text{Prime}\\ \hline 73 & 1803423556807921 & 1 & \text{Prime}\\ \hline 77 & 12360848946698171 & 73\neq1 & \text{Not prime}\\ \hline 79 & 32361122672259149 & 1 & \text{Prime}\\ \hline 81 & 84722519070079276 & 22\neq1 & \text{Not prime}\\ \hline 83 & 221806434537978679 & 1 & \text{Prime}\\ \hline 87 & 1520283919093591604 & 62\neq1 & \text{Not prime}\\ \hline 89 & 3980154972736918051 & 1 & \text{Prime}\\ \hline 91 & 10420180999117162549 & 3\neq1 & \text{Not prime}\\ \hline 93 & 27280388024614569596 & 35\neq1 & \text{Not prime}\\ \hline 97 & 186982561199565069121 & 1 & \text{Prime}\\ \hline 99 & 489526700523968661124 & 76\neq1 & \text{Not prime}\\ \hline 101 & 1281597540372340914251 & 1 & \text{Prime}\\ \hline \end{array}$$ Why does this work? Can this become a deterministic primality test? Any insights, references, counterexamples, or theoretical developments would be greatly appreciated.

论坛问题帖 | has_answers | Dev Sharma | 2025-06-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018404 | MO-49894

The Invariant Subspace Problem: examples

Question. Is there a concrete example of a bounded linear operator on a Hilbert space for which it is not known if it has a non-trivial closed invariant subspace? [Added 24.01.2011: According to Bernard Beauzamy (Introduction to Operator Theory and Invariant Subspaces, Elsevier (1988), p. 345), the operator which is "closest" to a counter-example is the one built by the present author: it has one hypercyclic point x_0, and for every polynomial p with complex coefficients, $p(T)x_0$ is also hypercyclic. Therefore, the operator has a vector space of hypercyclic points (thus solving a question raised by P. Halmos), but it may still have points which are not cyclic at all, thus having Invariant Subspaces. Beauzamy refers to his manuscript "The orbits of a linear operator". I have not been able to find an electronic version of this manuscript (or paper) online. Does anyone know where one may find a description of the example? Is it presently known whether the operator in Beauzamy's example has an invariant subspace?]

论坛问题帖 | has_answers | Andrey Rekalo | 2010-12-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018405 | MO-499155

Which pairs of mutually contradicting conjectures are there?

Years ago I had the pleasure of witnessing Simon Thomas giving a wonderful talk about Martin's conjecture, which I just now fondly remembered reading this question. Even though I am not well-versed in all (or even many) fields of mathematics I find it quite curious that it was the only time I ever heard about two contradictory "official" conjectures. In chit-chat I definitely heard colleagues expressing different sentiments about which way a certain problem might finally be resolved but my guess is that they usually would not have been eager to put it in writing, possibly not even be happy to "go on the record". Also, the problem in question may have been a quite specific one relevant only to a small number of people... So...my question is: What are other example pairs(triples?) of that kind you are aware of? They ideally should have appeared in writing but I guess that I would be satisfied with conjectures which are ascribed to people by a nontrivial part of the (respective) mathematical (sub)community.

论坛问题帖 | has_answers | TLo | 2025-08-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018406 | MO-500262

How can referees verify computationally intensive results when HPC resources are required?

This question is somewhat related to my previous question and is also inspired from this other question concerning the credibility of extensive computations (although from a different perspective). In the course of preparing a paper, I verified a conjecture computationally for a large range of cases. For small cases, I was able to run the computations on my laptop, since they required little RAM. However, at some point this was no longer feasible. At my university we have access to a high-performance computing (HPC) cluster, which allowed me to perform the larger calculations. This raises a concern: when the paper is submitted, the referee may wish to check the correctness of these computations. But what if they do not have access to an HPC cluster? In that case, they can only verify a portion of the claims. Is it sufficient to provide the raw output together with the code on GitHub/Zenodo, or is this not considered credible? After all, the raw output will consist of a .txt file that could, in principle, be tampered with (although doing so would clearly be dishonest, against all academic standards, and would eventually be detected by the community). My concern is precisely how referees can reasonably test such results in order to avoid this potential issue.

论坛问题帖 | has_answers | Chess | 2025-09-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018407 | MO-50071

Prepending strings to primes.

Hello, we all know that 31,331,3331,33331,333331,3333331,33333331 all are primes, and that 333333331 is not. Here we prepend the digit 3 to 31, to get a list of 7 primes.This gives me the following thought: Let $$D = \{\text{all possible nonnull finite digit strings}\},$$ $$D' = \{\text{all things in D that do not start with 0}\}.$$ Define a function $m: D' \times D \to N \cup {\infty}$ by: $$m(A,B)= \min \{ k\geq 1 \colon A^kB \text{ is composite} \} - 1,$$ i.e., the number of consecutive primes at the beginning of the list $AB, AAB, AAAB, \dots$. For example, $m(3,1)=7$. Which values does m take? Is it unbounded? Is it ever ∞? This question has been posted to math.stackexchange too, and I got one comment talking about that it might involve Tao-Ziegler extension to the Green-Tao theorem, and so I thought it might be more appropriate here. So please, excuse me if it's posted wrongly, or if one shouldn't post to both channels.

论坛问题帖 | has_answers | Dedalus | 2010-12-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018408 | MO-501474

An inequality that implies Frankl's union-closed conjecture

Let $\mathcal{F}$ be an union-closed family of subsets of $[n]=\{1,2,...,n\}$, assume $\varnothing\in\mathcal{F}$. Let $l_i=|\{S|S\in\mathcal{F},i\notin S\}|,u_i=|\{S|S\in\mathcal{F},i\in S\}|$, then $l_i+u_i=|\mathcal{F}|,\forall i\in[n]$. For a set $S\subset[n]$ (not necessary in $\mathcal{F}$), let $P_S=\prod_{i\notin S}l_i\prod_{i\in S}u_i$, M_S is the maximum set in $\mathcal{F}$ such that $M_S\subset S$ (which always exists because $\mathcal{F}$ is union-closed and $\varnothing\in\mathcal{F}$). Is the following inequality true? $$\sum_{S\subset[n]}P_{S/M_S}\leq|\mathcal{F}|^n$$ We could reduce Frankl's union-closed conjecture to the case $l_i,u_i>0,\forall i\in[n]$, then if the conjecture is false ($l_i>u_i,\forall i\in[n]$), then $P_S>P_R,\forall S\nsubseteq R\subset[n]$, so $P_S<P_{S/M_S}=P_{\varnothing}$ for non-empty $S\in\mathcal{F}$. We then have: $$|\mathcal{F}|^n=\sum_{S\subset[n]}P_{S}<\sum_{S\subset[n]}P_{S/M_S}$$ contradicting the above inequality.

论坛问题帖 | has_answers | Veronica Phan | 2025-10-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018409 | MO-501564

Polynomial identification of natural numbers

Let n be an integer parameter. Is there a polynomial $f(x,y,n)\in \mathbb{Z}[x,y,n]$ such that $$ n\in \mathbb{N} \Longleftrightarrow \exists x,y\in \mathbb{Z}, f(x,y,n)=0? $$ This is a generalized form of the infamous Poonen Christmas problem. Bjorn's question boils down to the special case when $f(x,y,n)$ is of the form $cn+g(x,y)$ for some nonzero integer c and some $g(x,y)\in \mathbb{Z}[x,y]$. A formula by Raphael Robinson in Arithmetical definitions in the ring of integers, modified slightly as by Zhi-Wei Sun in A new relation-combining theorem and its applications (and reiterated by Mihai Prunescu in this paper) gives the following near example: $$ n\in \mathbb{N} \Longleftrightarrow \exists x,y\in \mathbb{Z}, (x^2-(4n+2)y^2-1=0) \land (y\neq 0). $$ "Open Question A" in Prunescu's linked paper is exactly my question.

论坛问题帖 | no_answers_at_retrieval | Pace Nielsen | 2025-10-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018410 | MO-503219

No real roots of $\frac{1}{2^{1+it}} + \frac{1}{3^{1+it}} + \frac{1}{5^{1+it}}$

Prove or disprove that $$\frac{1}{2^{1+it}} + \frac{1}{3^{1+it}} + \frac{1}{5^{1+it}} \neq 0$$ for any real number t. I find it surprising that so simple looking equations involving complex numbers might turn out to be quite challenging to solve. For more context, see here, so I hope this gives a good motivation to discuss the problem here.

论坛问题帖 | has_answers | DesmondMiles | 2025-10-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018411 | MO-503437

Minimizing an expression with sum of digits

Conjecture. If $S_b(n)$ denotes the sum of digits of n with base b then $$10 \cdot S_5(n)+12 \cdot S_2(n)-5 \cdot S_3(n) \geq 0$$ and the equality holds if and only if $n = 19925000$. The expression we look to minimize in this case is $$P(n) = 10 \cdot S_5(n)+12 \cdot S_2(n)-5 \cdot S_3(n).$$ It seems the minimium value of this is 0 and the maximum tends to infinity. Intuitively, to minimize $P(n)$ we want to maximize $\{\log_3(n)\}$ while simultanously minimizing $\{\log_2(n)\}$ and $\{\log_5(n)\}$. Of course, we have coefficients so it is not just as simple as this. But even if you look at cases like $n = 3^m-1$ it still does not get below 0. In this case my question would be to minimize the expression $$Q(n) = 10 \cdot S_5(3^n-1)+12 \cdot S_2(3^n-1)-2n.$$ In general, is it known how to minimize the expression $$P(n) = \sum_{i = 1}^m a_i S_{b_i}(n)$$ for some $a_i \in \mathbb{Z}$ and $b_i \in \mathbb{N}$ such that $b_i > 1$? In other words, under which conditions does the minimum value exist?

论坛问题帖 | has_answers | John C | 2025-11-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018412 | MO-503646

A conjecture on prime distribution (PKD Conjecture)

I would like to propose the following conjecture The PKD Conjecture (PKD) Let p,d be positive integers with $\gcd(p,d)=1$. There exists a function $$ f:\mathbb{N}\to\mathbb{N}, \quad f(N)<N, $$ such that there always exists an integer k satisfying $$ 0 < k \le f(\max\{p,d\}), \quad q = pk + d \text{ is prime}. $$ Special cases Case d,p: This reduces to the existence of the smallest prime in the residue class $d \pmod p$. This is related to Linnik’s theorem on the least prime in an arithmetic progression. Case $p=1$: This is equivalent to Cramér’s prediction, i.e. that there is a prime in the interval $$ (d,\, d+f(d)). $$ Heuristic A reasonable growth for f is expected to be $$ f(N) \sim e (\log N)^2, $$ which is consistent with Cramér’s conjecture on prime gaps. Motivation This conjecture can be viewed as a “meta-conjecture” that generalizes both: the distribution of

primes in arithmetic progressions (Linnik-type problems), and the distribution of primes in short intervals (Cramér-type problems). It suggests a unified framework for understanding prime distribution both along progressions and within bounded gaps. Question Is this conjecture a consequence of Montgomery’s conjecture? When $$ f(N) \sim e (\log N)^2, $$ Any references or insights would be greatly appreciated.

论坛问题帖 | has_answers | Lazy fish | 2025-11-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018413 | MO-5041

Pencils with many completely decomposable fibers

Let $F= \frac{G}{H} : \mathbb P^n \to \mathbb P^1$ be a non-constant rational function (G and H homogenous polynomials of the same degree in $\mathbb C^{n+1})$. The fiber over $(\lambda:\mu) \in \mathbb P^1$ is completely decomposable if all the irreducible factors of the polynomial $\mu G - \lambda H$ are linear polynomials. In a joint work with Sergey Yuzvinsky, we proved that when $n=4$ either (a) F, after a linear change of coordinates, belongs to $\mathbb C(x_0,x_1,x_2,x_3)$; or (b) F has at most three completely decomposable fibers. We couldn't come up with an example not satisfying (a), but with three completely decomposable fibers. Is anyone aware of an example like that ? Almost one year later: Let me describe the known examples of rational functions on $\mathbb P^3$ with three completely decomposable fibers. For every natural number $d \ge 1$, consider the rational function $F_d : \mathbb P^3 \to \mathbb P^1$ defined as $$ F_d(x_0:x_1:x_2:x_3) = \frac{(x_0^d - x_1^d)(x_2^d - x_3^d)}{(x_0^d - x_2^d)(x_1^d - x_3^d)} . $$ Clearly the fibers over $(0:1)$ and $(1:0)$ are completely decomposable. The fiber over $(-1:1)$ is also completely decomposable since $$ F^{-1}(1:-1) = \lbrace (x_0^d - x_3^d)(x_1^d - x_2^d) = 0 \rbrace \, . $$ For $d=1$, F_d can be written in fewer variables, but for $d \ge 2$ this is not the case.

论坛问题帖 | no_answers_at_retrieval | Jorge Vitório Pereira | 2009-11-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018414 | MO-50570

Real approximation for homogeneous spaces of linear algebraic groups

Let X be a smooth geometrically integral variety over $\mathbf{Q}$, having a $\mathbf{Q}$-point. We say that X has the real approximation property if $X(\mathbf{Q})$ is dense in $X(\mathbf{R})$. Let now X be a homogeneous space of a connected linear algebraic group G defined over $\mathbf{Q}$. Since X has a $\mathbf{Q}$-point, we may write $X=G/H$, where H is a $\mathbf{Q}$-subgroup of G, not necessarily connected. Question 1. Does real approximation hold for any homogeneous space $X=G/H$ of any connected linear algebraic $\mathbf{Q}$-group G, where H is not necessarily connected? Serre proved real approximation for algebraic tori over $\mathbf{Q}$ (i.e. when G is a torus and $H=1$). Sansuc proved real approximation for connected linear algebraic groups over $\mathbf{Q}$ (i.e. when G is any connected linear algebraic group over $\mathbf{Q}$ and $H=1$). One can prove real approximation for homogeneous spaces with connected stabilizer (i.e. when G is a connected linear algebraic group over $\mathbf{Q}$, and $H\subset G$ is any connected $\mathbf{Q}$-subgroup). In the general case (i.e. when H is non-connected) I expect the negative answer, but I cannot construct a counter-example. The question can be stated in terms of Galois cohomology. Assume for simplicity that $G=\mathrm{SL}_n$, then $H^1(\mathbf{Q},G)=1$ and $H^1(\mathbf{R},G)=1$. Any orbit of $G(\mathbf{R})$ in $X(\mathbf{R})$ is open. If such an orbit contains a $\mathbf{Q}$-point, then $\mathbf{Q}$-points contained in this orbit are dense in this orbit, because our $G=\mathrm{SL}_n$ has real approximation. We denote by $G(\mathbf{R})\backslash X(\mathbf{R})$ the set of orbits of $G(\mathbf{R})$ in $X(\mathbf{R})$. Now our question is, whether the canonical map $G(\mathbf{Q})\backslash X(\mathbf{Q})\to G(\mathbf{R})\backslash X(\mathbf{R})$ is surjective. There is a canonical bijection $$ G(\mathbf{Q})\backslash X(\mathbf{Q})=\mathrm{ker}[H^1(\mathbf{Q},H)\to H^1(\mathbf{Q},G)]=H^1(\mathbf{Q},H) $$ and similarly $G(\mathbf{R})\backslash X(\mathbf{R})=H^1(\mathbf{R},H)$, so in this case Question 1 can be stated as follows: Question 2. Is it true that for any linear algebraic group H, defined over $\mathbf{Q}$ (not necessarily connected or abelian) the localization map $\mathrm{loc}_\infty\colon H^1(\mathbf{Q},H)\to H^1(\mathbf{R},H)$ is surjective? If the group H is connected or abelian, then the localization map loc_∞ is surjective (Sansuc's proof in the case of finite abelian H uses the Chebotarev density theorem and the fact that a decomposition group of ∞ is isomorphic to $\mathbf{Z}/2\mathbf{Z}$, hence cyclic). In the general case I expect the negative answer, i.e. that there exists a finite non-abelian algebraic group H over $\mathbf{Q}$ for which the localization map loc_∞ is not surjective, but I cannot construct such a counter-example.

论坛问题帖 | no_answers_at_retrieval | Mikhail Borovoi | 2010-12-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018415 | MO-506499

A 2-fully normal space which is not finitely-fully normal

An open cover $\mathcal{U}$ has a n-star open refinement $\mathcal{V}$ if its a refinement such that if $V_1\cap ...\cap V_n\neq\emptyset$ and $V_k\in\mathcal{V}$ then $\bigcup_k V_k\subseteq U$ for some $U\in\mathcal{U}$. It has a finite-star refinement if it has a refinement which is an n-star refinement for all $n\in\omega$ (at the same time). A space is called n-fully normal if every open cover has n-star open refinement. It's a theorem of Goldman that this is equivalent to 2-fully normal. (here of course $n\in \omega$) A space is called finitely-fully normal if every open cover has a finite-star refinement. M. J. Mansfield conjectures the following: Not every 2-fully normal space is finitely-fully normal. I did try to search for it, but what came up are conjectures relating to the almost 2-fully normal property (strongly collectionwise normal) instead. In Almost n-fully normal spaces by K. P. Hart, this is question 4.9a, so it wasn't known in 1982. I assume this has been answered by now. Where can I find out about this? What do I read?

论坛问题帖 | no_answers_at_retrieval | Jakobian | 2025-12-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018416 | MO-50661

unboundedness of number of integral points on elliptic curves?

If $E/\mathbf{Q}$ is an elliptic curve and we put it into minimal Weierstrass form, we can count how many integral points it has. A theorem of Siegel tells us that this number $n(E)$ is finite, and there are even effective versions of this result. If I'm not mistaken this number $n(E)$ is going to be a well-defined invariant of $E/\mathbf{Q}$ (because different minimal Weierstrass models will have the same number of integral points). Is it known, or conjectured, that $n(E)$ is unbounded as E ranges over all elliptic curves? Note: the question is trivial if one does not put E into some sort of minimal form first: e.g. take any elliptic curve of rank 1 and then keep rescaling X and Y to make more and more rational points integral.

论坛问题帖 | has_answers | Kevin Buzzard | 2010-12-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018417 | MO-506831

Are there infinitely many k such that $6k-1, \; 6k+1, \;$ and $12k-1$ are all prime?

I have been looking into the intersection of Twin Primes and Sophie Germain primes. Specifically, I am interested in whether a prime p can be both the "lower" member of a twin prime pair and a Sophie Germain prime. If p is the upper member of a twin prime pair, $p = 6k+1$ (for $k≥1$). Then $2p+1 = 2(6k+1)+1 = 12k+3 = 3(4k+1)$. This is always a multiple of 3, so $2p+1$ is never prime for k≥1. If p is the lower member, $p=6k-1$. Then $2p+1=2(6k-1)+1=12k-1$ My question: Is it conjectured (or known) that there are infinitely many k such that $6k-1$, $6k+1$, and $12k-1$ are all prime?

论坛问题帖 | has_answers | nifemibosun | 2026-01-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018418 | MO-508168

Status of linearity of mapping class group

This is a followup to this question on the status of the linearity of the mapping class group, which doesn't appear to me to have been satisfyingly answered. It seems to be agreed that this paper by Bigelow and Budney (2001) establishes the linearity of the mapping class group of a surface Σ_g for $g=2.$ On the other hand, this paper by Nikolaev (2003) claims a positive resolution of the question for any g. But Benson Farb in chapter 2 of this book (2006) and also Ozsvath, Lipshitz, and Thurston in this paper (2010) claim the problem is open. Has a mistake been found in the proof by Nikolaev? Why does there not seem to be much confidence in it?

论坛问题帖 | has_answers | subrosar | 2026-02-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018419 | MO-508398

How far is Scholze's vision for shtuka over $\operatorname{Spec} (\mathbb{Z})$ from realization?

How speculative is Scholze's vision for shtuka over $\operatorname{Spec} (\mathbb{Z})$? Intuitively, it appears very speculative. Primarily because shtuka are very much function field objects and in order to have a "shtuka" over a number field one needs very strict analogues of (1) the Frobenius and (2) a field of constants both of which are not available for number fields. Very recently Scholze gave a seminar on a new object he calls Gestalten. From what I understand, one of the major motivations for this was to create a "place" or "setting" that can unify or simultaneously handle both non-archimedean and archimedean algebraic geometry. I am wondering: does this new setting get one closer to this dream or does it make headway on either of the two major obstacles? Moreover, I am confused as to how such an object (Gestalten) should succeed in unifying the non-archimedean and archimedean worlds in a non-trivial way. Any answers both more generally and more narrowly relating Gestalten would be wonderful.

论坛问题帖 | has_answers | Luka | 2026-02-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018420 | MO-509408

A 2D lattice generalization related to prime gaps and Cramér's conjecture

Conjecture: Let x and y be integers such that $1 < x < y$ and $\gcd(x, y) = 1$. There always exists at least one prime p of the form: $$p = n \cdot x + m \cdot y$$ where $n, m \ge 1$ are integers satisfying $n \le \lceil \ln^2(y) \rceil$ and $m \le \lceil \ln^2(x) \rceil$. Restricted Conjecture (Weak Form): Let x and y be primes ($x < y$). There always exists at least one prime p of the form:$$p = n \cdot x + m \cdot y$$ Where $n, m \ge 1$ are integers satisfying $n \le \lceil \ln^2(y) \rceil$ and $m \le \lceil \ln^2(x) \rceil$. The logarithmic bounds are structurally inspired by Cramér's conjecture. However, by strictly enforcing $x > 1$ and incorporating the parameters (x, y) as independent weights, this formulation intentionally shifts the problem from a classical 1D prime gap search into a 2D discrete lattice equipped with strict coprimality sieves. I have computationally verified this existence for all primes $x \le 10^5$ and $y \le 10^8$ with absolutely no counter-examples found. Moreover, I stress-tested the conjecture against extreme "death cases" where x and y are highly composite primorials (e.g., setting x as the product of the first 200 primes, $x \approx 10^{512}$, and y as the product of the next 5 primes). Interestingly, the empirical data shows that despite the coprimality filters severely throttling the available grid points (the survival rate of (n, m) pairs drops to approximately 15%), a prime is almost always found immediately at $n=1$ and a very small m (utilizing less than 0.1% of the m-bound), suggesting that the conjecture holds with a comfortable safety margin. Questions: Q1. Is it possible to give heuristic support for this conjecture for all values of x and y? Q2. Are existing bounds on prime gaps or primes in arithmetic progressions capable of fully resolving this problem, or does it require new sieve methods?

论坛问题帖 | no_answers_at_retrieval | Đào Thanh Oai | 2026-03-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018421 | MO-510194

A mysterious statement by Martin and Solovay

D. A. Martin and R. M. Solovay, "Internal Cohen extensions", Annals of Mathematical Logic - Volume 2, No. 2 (1970) pp. 143--178, (MR270904, Zbl 0222.02075) say on page 166: Solovay will show elsewhere that II implies that $\aleph_1$ is a $0, 1$-measurable cardinal, so that II contradicts the axiom of choice. Here II is Lusin's hypothesis from N. Lusin, "Sur les ensembles analytiques nuls", Fund. Math. 25 (1935) 109-131 (JFM 61.0222.03, Zbl 0012.34401) saying that: (II) every union of a collection of constituents of a coanalytic set is again a coanalytic set. The supposed result by Solovay sounds very unusual as compared to typical solutions of problems of classical descriptive set theory. Is there anything known about any further development?

论坛问题帖 | has_answers | Vladimir Kanovei | 2026-04-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018422 | MO-512480

On $a^3+b^3+c^3 = N$ and $a^5+b^5+c^5+d^5+e^5 = N$

(Update, July 21, 2026): For $N<100$ and using a higher search radius, Simon Goater found, 55 = 38800^5 - 37045^5 + 22356^5 - 29644^5 - 15378^5 59 = 32064^5 - 32521^5 + 25162^5 - 23778^5 + 1078^5 while Jeff Braun found (June 27, 2026), 49 = -22403^5 + 21596^5 + 15669^5 + 4698^5 - 4001^5 Guruprasad's earlier search radius of $|x|<20000$ found 6 (and more) like, 38 = 9265^5 - 9258^5 - 3035^5 - 98^5 - 46^5 reducing unknown $N\leq100\,$ to $\,29-6-3 = 20$. I. Problem We are looking for integer near-misses of the Fermat quintic threefold, $$a^5+b^5+c^5+d^5+e^5 = 0\qquad$$ $\hskip1.5in$ where the sum, instead of vanishing, is some small integer $N<300$, preferably $N<100$. This threefold is a Calabi-Yau manifold. Note: Jeff Braun (mentioned above) found the fourth primitive integer solution with $N=0$. See paper. II. Third Powers The equation $a^3+b^3+c^3 = N$ has often been discussed in MO. For a small sample, see this, this, this. In a 1955 paper, Miller and Woollett used the early computer EDSAC to search $x_1^3+x_2^3+x_3^3 = N$ for positive $N<100$ with $N \not\equiv \pm\,4\, \pmod9.\,$ If they had used a very modest range of $|x|<30$, they wouldn't have found 12 N. Arranged by form, \begin{array}{|c|c|} \hline \text{Form} & \text{Missing N (3rd powers)} \\ \hline N\equiv 2 \pmod{9} & 74 \\ \hline N\equiv 3 \pmod{9} & 30, 39, \color{blue}{48}, 75, 84\\ \hline N\equiv 6 \pmod{9} & 33, 42,\color{blue}{51}, \color{blue}{78}, 87\\ \hline N\equiv 7 \pmod{9} & 52 \\ \hline \end{array} with none missing for $N\equiv 0 \pmod{9}$ and $N\equiv \pm1 \pmod{9}$. Miller and Woollett mentioned some $N\equiv \pm3 \pmod{9}$ were "difficult". But they actually used a search radius 100x higher at $|x|<3000$, and found just three additional numbers (in blue above) with the largest terms for $N=51$, $$51 = -796^3 + 659^3 + 602^3$$ so $12-3=9$ missing N. The next value, $N = 87$ was found ten years later in 1964 by Gardiner and Lazarus. Over the years, the search radius increased from $10^4, 10^6, 10^8,\dots,10^{18}$ (though not exhaustive) to find all 9 culminating in Booker and Sutherland's 2019, $$-80538738812075974^3+80435758145817515^3+12602123297335631^3 = 42$$ III. Fifth Powers A similar phenomenon seems to be at work for $a^5+b^5+c^5+d^5+e^5 = N$ which does not bode well for finding "smallish" solutions. Initial searches for small positive $N<100$ were done by DanielV and Guruprasad. If terms are $|x|< 10^3$ or less, then one can easily find 70 N but wouldn't find 29 N. Arranged by form, these are, \begin{array}{|c|c|} \hline \text{Form} & \text{Missing N (5th powers)} \\ \hline N\equiv 0 \pmod{11} & \color{red}{55} \\ \hline N\equiv 1 \pmod{11} & \color{blue}{89}\\ \hline N\equiv 2 \pmod{11} & 90 \\ \hline N\equiv 3 \pmod{11} & \color{blue}{58}, \color{blue}{80}, \color{blue}{91}\\ \hline N\equiv 4 \pmod{11} & 15, 26, 37, 48, \color{red}{59}, 81\,\;\; \quad \\ \hline N\equiv 5 \pmod{11} & 16, 27, \color{blue}{38},\color{red}{49}, 60, 82, 93\; \\ \hline N\equiv 6 \pmod{11} & 39, 72\; \\ \hline N\equiv 7 \pmod{11} & 7, 40, 51, 73, 84 \\ \hline N\equiv 8 \pmod{11} & 41 \\ \hline N\equiv 9 \pmod{11} & 9,\color{blue}{42} \\ \hline N\equiv 10 \pmod{11} & none\\ \hline \end{array} Note: Adam Bailey pointed out that the difficult N would be the four forms $N\equiv \pm4 \pmod{11}$ and $N\equiv \pm5 \pmod{11}$ as seen in the table above. Guruprasad extended the search radius to $|x|<\color{blue}{10^4}$ and found six additional numbers (in blue), though only N = 38 has terms $|x|>5000$, $$38 = 9265^5 - 9258^5 - 3035^5 - 98^5 - 46^5$$ Jeff Braun and Simon Goater went higher and found N = 49, 55, 59 (in red). So $29-6-3=20$ are missing, namely, N = 7, 9, 15, 16, 26, 27, 37, 39, 40, 41, 48, 51, 60, 72, 73, 81, 82, 84, 90, 93 IV. Higher N For the higher range $N<300$, Simon Goater searched terms $|x_i|<4000$ and Guruprasad extended to terms $|x_i|<20000$, \begin{array}{|c|c|} \hline \text{Range}\; R_n & \text{# of missing}\, N & \text{# of missing}\, N & \text{# of missing}\, N\\ & \text{if}\, |x|<10^3 & \text{if}\, |x|<10^4 & \text{if}\, |x|<10^5 \\ \hline \quad 1\leq N \leq100 & 29 & 23 & 20\,?\\ \hline 101\leq N\leq200 & 39 & 30 & 27\,?\\ \hline 201\leq N\leq300 & 36 & 28 & 25\,?\\ \hline \end{array} Notice that, just like for 3rd powers, with increasing search radius $10^3\to10^4$, the number of missing N reduces. So what happens when $10^4\to10^5$? It seems they go down. The counts in the third search radius reduced when N = 49 was later found by Jeff Braun, N = 126, 169, 184, 289 by Eugene Go, and N = 55, 59, 228, 233 by Simon Goater. Updated missing N of range R_n are: R_1 = 7, 9, 15, 16, 26, 27, 37, 39, 40, 41, 48, 51, 60, 72, 73, 81, 82, 84, 90, 93 R_2 = 103, 104, 105, 106, 112, 114, 116, 117, 132, 134, 135, 136, 137, 138, 139, 149, 153, 158, 159, 161, 170, 171, 172, 183, 191, 194, 196 R_3 = 203, 205, 215, 222, 224, 226, 229, 235, 236, 238, 248, 249, 258, 264, 268, 269, 270, 279, 280, 281, 290, 291, 292, 293, 297 V. Question What is a good method to reach higher bounds and find more missing $N<300$? Is it Guruprasad's sieve-based, meet-in-the-middle algorithm, or Jeff Braun's method to find $x_1^5+x_2^5+x_3^5+x_4^5+x_5^5 = N$ for $N=0$ described in his paper, or can the methods for $x^3+y^3+z^3=N$ be modified for 5th powers? Note: By the Easier Waring's Problem, then $5\leq v(5)\leq 10$. There is no mod obstruction, so we assume $v(5)=5$.

论坛问题帖 | has_answers | Tito Piezas III | 2026-06-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018423 | MO-513215

Can an integer reciprocal series remain rational after a unit shift in the denominator?

Let a_n be an infinite sequence of positive integers (not necessarily distinct), and for this sequence, $$ \sum_{n=1}^{\infty}\frac{1}{a_n} $$ converges. Define $$ S(k)=\sum_{n=1}^{\infty}\frac{1}{a_n+k}, \qquad k\in\mathbb{Z}_{\ge0}. $$ My question is whether there exists any such sequence a_n for which both $$

S(0)\in\mathbb{Q} \quad\text{and}\quad S(1)\in\mathbb{Q}. $$ Equivalently, can one prove that this is impossible? The sequence is allowed to have repeated values; the only assumptions are that every a_n is a positive integer and the series converges. I know that if the a_n are allowed to be arbitrary positive real numbers, then such examples exist by construction, so does if negative integers are allowed using telescoping series, so the positive integrality of the sequence is the essential restriction here. If this is known, I'd appreciate either a reference or an explicit example. Otherwise, a proof that no such sequence exists would also answer the question.

论坛问题帖 | has_answers | bihariforces | 2026-07-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018424 | MO-513304

Conjecture : $\forall n>3 , \text{gpf}(y_n)>\text{gpf}(y_1)$

Let $n \in \mathbb{Z}_{>1}$, we call $\text{gpf}(n)$ the greatest prime factor of n Take $d \in \mathbb{Z}_{>1}$ a squarefree integer, we call (x_n,y_n) the n-th pell solution of $x^2-dy^2=1$ Can you show that $\forall n>3 , \text{gpf}(y_n)>\text{gpf}(y_1)$? For instance, If $d=263$ we have $x_1=139128$ and $y_1=8579=23 \times \color{red}{373}$ $$y_2=2^4 \times 3 \times 11 \times 17 \times 23 \times 31 \times \color{red}{373}$$ $$y_3=5 \times 7 \times 19 \times 23 \times 29 \times 101 \times 127 \times 313 \times \color{red}{373}$$ $$y_4=2^5 \times 3 \times 11 \times 17 \times 23 \times 31 \times \color{red}{373} \times 184913 \times 209359 $$ However, the next values of y_n seem to be divisible by really huge primes ! e.g : $\text{gpf}(y_{17})=152978144089051842195133050091524167 41 \approx 1.53 \times 10^{37}$ and so can we find in general a lower bound like $\text{gpf}(y_n)>f(n,y_1)$ for $n\ge n_0$? I know only few results about the size of primes dividing y_n, like this one by D. H. Lehmer in Lemma 5 : if $n>3$, y_n is divisible by a prime $\ge 2n-1$ and so for $n>3$, we can say $\text{gpf}(y_n) \ge 2n-1$ However, this lower bound doesn't account for the size of primes dividing y_1 (in our case 23 and 373) so a lower bound like $\text{gpf}(y_{17})\ge 2\times 17-1=33$ is so useless :-(because $y_1 \mid y_n$, so $\forall n\ge 1$ we have $\text{gpf}(y_n)\ge \text{gpf}(y_1)=373$ already bigger than 33 So an elementary function f we can deduce for $n>3$ is $f(n,y_1)=\max(2n-1,\text{gpf}(y_1))$ and could one points me towards published results of the same nature ? Thanks for help :-)

论坛问题帖 | has_answers | Lhachimi | 2026-07-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018425 | MO-513417

Is Gardam's support-21 unit in $\mathbb{F}_2[P]$ (Promislow group) support-minimal?

Gardam [1] disproved the Kaplansky unit conjecture by exhibiting a nontrivial unit α of support size 21 in $\mathbb{F}_2[P]$, where $P = \langle a,b \mid b^{-1}a^2b = a^{-2},\ a^{-1}b^2a = b^{-2}\rangle$ is the Promislow group, and remarks that it is "plausible" that every nontrivial unit has support size at least 21. Craven-Pappas [2] had verified the unit conjecture in $K[P]$ for supports of dihedral length at most 3. Since then, computational searches for units and zero divisors of small support in a group-agnostic setting have been carried out by Garg-Mineyev [4], with a structural obstruction to that framework proved by Shin [5]; as far as I can tell, these do not directly address support minimality inside $\mathbb{F}_2[P]$ itself. Setup. Write elements of P as $x^i y^j z^k \sigma$ with $x=a^2$, $y=b^2$, $z=(ab)^2$, $\sigma \in \{1,a,b,ab\}$, and fix the asymmetric boxes B_1 (the 108 elements with $|i|,|j|,|k|\le 1$) and B_2 (the 500 elements with $|i|,|j|,|k|\le 2$); Gardam's unit and its inverse have supports inside B_1 and B_2. Over $\mathbb{F}_2$, the equation $\alpha\beta=1$ is a parity condition: every group element $w \ne 1$ must have an even number of factorizations $w = gh$ with $g \in \operatorname{supp}\alpha$, $h \in \operatorname{supp}\beta$, and $w=1$ an odd number. Existence of such supports inside the boxes is a finite decision problem, which we solve with SAT solvers, together with a bound on $|\operatorname{supp}\alpha|$ and, optionally, the support symmetry of [1, Lemma 1]. The setup is validated by reproducing $\alpha\alpha^{-1}=1$ for Gardam's unit in an independent implementation of P. Current computations (certificates and code: github.com/Divide-By-0/promislow-unit-minimality): A support-21 unit is re-found in $B_1 \times B_2$ and verified by exact multiplication in $\mathbb{F}_2[P]$. No unit pair exists with $\operatorname{supp}\alpha \subseteq B_1$, $\operatorname{supp}\beta \subseteq B_2$, $|\operatorname{supp}\alpha| \le 20$, and both supports symmetric in the sense of [1, Lemma 1] (solver-level, all $k \le 20$). For $|\operatorname{supp}\alpha| \le 16$ in the symmetric class (with no gauge normalization), the impossibility additionally carries machine-checkable proofs of unsatisfiability, validated with an independent proof checker. The computation aiming at the full range $|\operatorname{supp}\alpha| \le 20$ over $B_1 \times B_2$ with no symmetry assumption, with certificates throughout, is in progress. We regard the symmetric-class results as computational evidence and validation of the pipeline; even the completed unrestricted computation would be a finite-region statement, not the global theorem. Hence the actual questions: Question 1. Is a global lower bound on the support size of nontrivial units of $\mathbb{F}_2[P]$ (beyond the length-based results of [2]) known or announced? Is the global statement "every nontrivial unit has support size at least 21" believed to be within reach of current structural methods? Question 2. Are there structural arguments - via the dihedral-length analysis of [2], the structure theory of $(\mathbb{F}_2[P])^\times$ in [3], or otherwise - that would reduce every hypothetical unit of support size < 21 to finitely many configurations, e.g. by confining its support to an explicit finite region of P? That reduction is what would turn finite certified computations like the above into a proof of global minimality, and seems to be the conceptually valuable step. Question 3. Is anything known toward classifying the support-21 units up to multiplication by trivial units, inversion, and automorphisms of P? Independently found units (Gardam's, Murray's char-p variants, the char-0 units, and the SAT re-discoveries) land at support exactly 21, and a classification would explain the recurrence of 21 rather than merely confirm it numerically. [1] G. Gardam, A counterexample to the unit conjecture for group rings, Ann. of Math. 194 (2021), 967-979. [2] D. A. Craven, P. Pappas, On the unit conjecture for supersoluble group algebras, J. Algebra 394 (2013), 310-356. [3] L. Bartholdi, On Gardam's and Murray's units in group rings, Algebra Discrete Math. 35 (2023), 22-29. [4] M. Garg, I. Mineyev, arXiv:2501.07646 (2025). [5] Shin, A global girth obstruction for Garg-Mineyev taiko product structures, arXiv:2607.01716 (2026).

论坛问题帖 | no_answers_at_retrieval | August Sha | 2026-07-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018426 | MO-513661

Was a computational hardness argument ever used to solve a mathematical conjecture?

Let P be a property of positive integers. Suppose that $P(n)$ can be decided in finite time, but all known algorithms take a long time. Based on experiments done so far, it is conjectured that $P(n)$ holds for all n. Now someone proves, using computer science methods, that deciding $P(n)$ is NP-hard (or hard in some other complexity class). This immediately refutes the above conjecture, since if the conjecture were true, the decision problem would have been trivial. Has such a story ever happened?

论坛问题帖 | has_answers | Erel Segal-Halevi | 2026-07-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018427 | MO-513707

Checking one group for torsion

Let's consider such a group $$ G = \langle a, b, c, d \mid r_1, r_2, r_3, r_4, r_5, r_6, r_7 \rangle, $$ where $$ \begin{aligned} r_1 &= dcb c^{-1}d^{-1}bab^{-1}a^{-2},\\ r_2 &= cdc^{-1}d^{-1}a^{-1}b^{-1}acdb^{-2},\\ r_3 &= babdb^{2}d^{-1}c^{-1}a^{-2}c^{-1}d^{-1},\\ r_4 &= dcdb^{-1}d^{-1}a^{-1}c^{-1}db^{2}d^{-1}c^{-1}a^{-1},\\ r_5 &= dcdacdb^{-2}d^{-1}a^{-1}b^{-1}d^{-1}c^{-1},\\ r_6 &= dcabd^{-1}b^{-2}d^{-1}acdb^{-2},\\ r_7 &= badcb^{-1}d^{-1}a^{-1}c^{-1}acdb^{-2}d^{-1}. \end{aligned} $$ I found this group in my records. It is noteworthy that its group ring $Z[G]$ contains non-trivial zero-divisors. I can't provide them yet, because I'm not near the notebook where it was written out. In the future, I will update the post and provide them. First, a divisor of zero was built, and then a group G on relations that allow $\alpha\beta = 0$ in $Z[G]$. The motivation for considering this group is connected with Kaplansky's hypothesis about zero-divisors in group rings: Let G be a torsion-free group and K be a field. Then $K[G]$ does not contain zero-divisors. I examined this group for torsion, but I could not find an obvious torsion. I know that it is incredibly difficult to check a group with a random representation for torsion. Therefore, my question is: are there any techniques applicable specifically to this group that could prove\refute the presence of torsion? Is it possible to provide any relatively easily calculated signs that can give some information about the torsion in it? Is it possible to get any indirect information about torsion in this group using a computer? Can any GAP\Sage functions help with this? So far, all the functions I used have not been able to complete their work and give the final answer. Still, it's quite possible that I didn't see an obvious torsion. A couple of facts about this group: It's perfect. That is, $G = [G, G]$. Some relations contain quite large common bites. For example, $acdb^{-2}$. UPD: There is a zero divisor in $Z[G]$: $$AB = (d^2 c a + d^2 c d + d c b a - d b a b - d c d b a - a^2 b^2) (d b^2 d + d^2 c a b + a c d^2 - d c d - b a d c - c a d b)$$ Let's rename the terms. $$AB = (w_1 + w_2 + w_3 - w_4 - w_5 - w_6)(w_7 + w_8 + w_9 - w_{10} - w_{11} - w_{12})$$ In G, it is satisfied: \begin{align} &1. \quad w_1 w_7 = w_4 w_8 \\ &2. \quad w_1 w_8 = w_4 w_9 \\ &3. \quad w_1 w_9 = w_4 w_7 \\ &4. \quad w_2 w_7 = w_5 w_8 \\ &5. \quad w_2 w_8 = w_5 w_9 \\ &6. \quad w_2 w_9 = w_5 w_7 \\ &7. \quad w_3 w_7 = w_6 w_8 \\ &8. \quad w_3 w_8 = w_6 w_9 \\ &9. \quad w_3 w_9 = w_6 w_7 \\ &10. \quad w_4 w_{10} = w_1 w_{11} \\ &11. \quad w_4 w_{11} = w_1 w_{12} \\ &12. \quad w_4 w_{12} = w_1 w_{10} \\ &13. \quad w_5 w_{10} = w_2 w_{11} \\ &14. \quad w_5 w_{11} = w_2 w_{12} \\ &15. \quad w_5 w_{12} = w_2 w_{10} \\ &16. \quad w_6 w_{10} = w_3 w_{11} \\ &17. \quad w_6 w_{11} = w_3 w_{1} \\ &18. \quad w_6 w_{1} = w_3 w_{10} \end{align} You can set up a group with these relations in Sage, and using the G.simplified() function, get exactly the same representation as in the post.

论坛问题帖 | has_answers | Maximka123 | 2026-07-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018428 | MO-514251

Are there infinitely many primes of the form $\hspace{1mm} {}^b a \pm {}^a b$, with integers $a,b\geq 2$?

Let ${}^b a$ denote the tetration of a to height b (e.g., ${}^3 3=3^{3^3}=3^{27}=7625597484987$). I am interested in primes of the forms ${}^b a+{}^a b$ and ${}^b a-{}^a b$, where $a,b \geq 2$ are integers (I came across this problem while I was computing the first terms of the more general sequences of primes of the form ${}^b a \pm {}^d c$, with $a,b,c,d\geq 2$, for submission to the OEIS). Now, the only two terms I found of the more restrictive forms ${}^b a \pm {}^a b$ are: ${}^3 2+{}^2 3=16+27=43$ and ${}^2 3-{}^3 2=27-16=11$ (trivially, to be an odd prime, a and b must have opposite parity). Question. Are there any other primes of the form ${}^b a + {}^a b \hspace{1mm}$ or $\hspace{1mm} {}^b a - {}^a b$? If yes, are there infinitely many primes of either form? Remark. At a naive heuristic level, it is not clear to me what to expect: it seems that there are no immediate obstructions to primality beyond parity, while the extremely rapid growth of tetration makes prime values increasingly unlikely. Even partial results for either the sum or the difference would be of interest.

论坛问题帖 | has_answers | Marco Ripà | 2026-08-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018429 | MO-5144

Is Murasugi's conjecture still open?

Normalize the Alexander polynomial (in t) so that the positive and negative exponents are balanced. For example in the Conway normalization, make the substitution $z = t^{1/2} - t^{-1/2}$. The trefoil gives $t^{-1} - 1 + t$. Conjecture: Suppose that K is an alternating knot. Then the sequence of absolute values of the coefficients is unimodal. Specifically, if $\Delta_K(t) = \sum_i a_i t^i$, then $|a_0| \ge |a_1| \ge |a_2| \ge \cdots$. This is a conjecture due to Murasugi, I believe. Where is it written? Has this been proved or disproved?

论坛问题帖 | has_answers | Sammy Black | 2009-11-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018430 | MO-514729

A recursive overlap construction of finite Bernoulli-sum laws and an entropy-concavity conjecture

Let (a_{n}) be a strictly increasing sequence of integers with $a_{1}=2$. Define probability vectors $P_n\in\mathbb R^{a_n}$ by $ P_1=(p,1-p)$ and $$ P_{n+1} = p\,L(P_n)+(1-p)\,R(P_n), $$ where the “left’’ and “right’’ embedding operators $ L:\mathbb R^{a_n}\to\mathbb R^{a_{n+1}}, R:\mathbb R^{a_n}\to\mathbb R^{a_{n+1}} $ are given by $$ L(P_n)=(P_n,0,\ldots,0), \quad R(P_n)=(0,\ldots,0,P_n). $$ Is the entropy of P_{n} concave in p? This is easy to prove for the non overlapping case $a_{n}\ge 2^n$ and fairly easy to prove for $a_{n}=n+1$. I tried to prove it for $a_{n}=n^2+1$ and Fibonacci like sequences given by $a_{n+1}=a_{n}+a_{n-1}+1$, which are of special interest for me, but I did not succeed. I ask ChatGpt to give me a Mathematica Code to test the conjecture. For any sequence that I tested up to $n=30$ (and for some sequences up to n=100) it seems to be true. But honestly I do not understand the mechanism. I have now written an preprint https://arxiv.org/abs/2609.05546 on the problem.

论坛问题帖 | has_answers | Jörg Neunhäuserer | 2026-08-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018431 | MO-515053

On the Integer solutions to $w^5+x^5+y^5+z^5=1$

Let $w^5+x^5+y^5+z^5=1$ such that w,x,y,z are all Integers. Are there any non trivial Solutions ? EDIT: Nontrivial means $|wxyz| > 1$. Computational Search : According to Jeffrey M. Braun Paper ,The fourth known primitive solution to $a^5+b^5+c^5+d^5=e^5$, we can conclude that $max(|w|,|x|,|y|,|z|)<1.45×10^6$ has no non trivial solutions (search is exhaustive) and $1.45×10^6<|w|,|x|,|y|,|z|<2×10^6$ no non trivial solutions (partial search).

论坛问题帖 | no_answers_at_retrieval | Guruprasad | 2026-09-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018432 | MO-515297

Adaptive prime window conjecture, does a previous prime gap correlate with the following one?

I am looking for feedbacks on my conjecture which I worked on for almost a year now. I'll define the conjecture with the following definition: For every integer $x>2$, let p_x denote the largest prime strictly less than x, and let $d_x=x-p_x$ denote the distance from x to the preceding prime. Let q_x denote the smallest prime strictly greater than x, and define $g_x=q_x-x$ as the distance from x to the following prime. I conjecture: $${ g_x \leq \left\lceil 2e^{-\gamma} \left[\left(\ln\frac{x}{d_x}\right)^2+2 \right] \right\rceil }$$ where γ is the Euler--Mascheroni constant. Originally I was not using x but rather p, meaning I thought this was working only if the input was a prime number, however since the computation works with any integer I believe x represents even a stronger claim. Let's look at an example: $ x=23 $ $ p_x=19 $ $ q_x=29 $ $ d_x=23-19=4 $ $ g_x=29-23=6 $ $$\left\lceil 2e^{-\gamma} \left[\left(\ln\frac{23}{4}\right)^2+2 \right] \right\rceil =6$$ ${g_x\leq6}$$ Findings connected to this conjecture: A larger preceding gap makes the upper bound smaller It holds from 2 up to 10^8, and it has been tested with random samples (window sizes of 1.000.000) up to 10^{20} The Nicely gap (x = 1693182318746371) would have broken the conjecture without adapting it to the Cramér-Granville conjectural model. No other exception was found by removing the factor $2e^{-\gamma}$ It is on average tighter than Cramér's conjecture by 15%, scaling according to the size of d_x, reaching tightness discrepancies above 90% in some cases Pattern driven by negative lag-1 autocorrelation (r ≈ -0.03) On point 5, Gideon Afriyie (2026) independently computed lag-1 autocorrelation on 50 million primes and reported r ≈ -0.0276 in his published paper (https://www.sciencedirect.com/science/article/pii/S0167715226002622) While the formula holds for any X bigger than 2, the striking observation is that, in the vast majority of cases, the tightest value in the whole interval occurs at the prime at the left endpoint of the gap. On point 7: Defining "normalized tightness" a following: $$[R(x)= \frac{g_x}{2e^{-\gamma}\left[(\ln(x/d_x))^2+2\right]}]$$ From my current computation, the tightest value in the whole interval occuring at the prime at the left endpoint of the gap it is already clearly above 95% of the prime-gap intervals. I am in the process of rerunning the calculation carefully, because I suspect the actual figure is around 99%, but I don't want to claim the exact percentage until I have checked it again. There are interesting exceptions. For example, $$[31397<31411<31469]$$ where (31397) and (31469) are consecutive primes and (31411) is composite. At the prime (x=31397) $$[d_x=4,\qquad g_x=72]$$ giving $$[R(31397)\approx0.778]$$ But at the interior composite (x=31411), $$[d_x=14,\qquad g_x=58]$$ and $$[R(31411)\approx0.839]$$ So this is a genuine example where an interior composite point is tighter than the preceding prime, although this appears to be relatively uncommon. Any feedback would be appreciated. I don't have a formal math education and I am spending my spare time on this project. Therefore I will try my best to engage with whoever wants to spend his/her time in having a look at what I assembled.

论坛问题帖 | has_answers | Davide Ravasi | 2026-09-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018433 | MO-515446

Almost-primes produced by $n^2 + n + (d+1)/4$ for a positive proportion of fundamental discriminants?

I arrived at the question below through an extended exploration that began with the classical Euler polynomials $n^2 + n + p$ and the Rabinowitsch-Heegner connection, and gradually moved toward more approachable variants. I am not a number theorist; the formulation and the suggested attack emerged from a long dialogue with an AI system (Grok), which I used to survey the landscape, test successive weakenings, and identify plausible analytic obstacles. I am sharing both the problem and a brief account of the process in case either is useful. Motivation. It is classical that $n^2 + n + p$ takes prime values for all $n = 0, \dots, p-2$ if and only if $4p-1$ is a Heegner number. Only six such primes p exist. For a general prime p (or a general fundamental discriminant $d = 4p-1$) one still expects the polynomial to produce primes for many n, but even the existence of a single extra prime value for every large p appears open and is essentially a special case of Bunyakovsky-type problems. A natural weakening is to ask only for almost-primes, and only for a positive proportion of

discriminants. Open problem. Do there exist constants $\theta > 0$ and $\alpha > 0$ such that for a positive proportion of fundamental discriminants $d \le Y$ (as $Y\to\infty$), there is an integer n with $1\le n\le d^\theta$ for which $$n^2 + n + \frac{d+1}{4}$$ is either a $P_{r-\text{number}}$ for some fixed r, or at least has a prime factor larger than d^α ? Suggested approach and apparent obstacles. One can view the values as norms of elements $n + \frac{1+\sqrt{-d}}{2}$ in the quadratic order of discriminant -d Averaging over fundamental discriminants places the problem in a setting where tools from the theory of L-functions of quadratic characters and class-group characters are available. A plausible route is: establish a usable averaged level of distribution for the sequence (or for the principal binary quadratic form of discriminant -d) when summing over fundamental discriminants; apply a modern sieve (combinatorial or multidimensional) to detect almost-primes in the short range $n\le d^\theta$; control the fluctuation of the class number $h(-d)$ and possible exceptional zeros. The most serious gaps appear to be obtaining a sufficiently strong and uniform averaged level of distribution, and handling the variation of $h(-d)$ while retaining a positive-density set of good discriminants. Questions for the community. Is a result of this strength already known, or does it follow quickly from existing theorems on almost-primes represented by binary quadratic forms averaged over discriminants? Are there stronger partial results (for example, for a thinner set of discriminants, or with a larger number of prime factors) that are already in the literature? Does the suggested approach look viable with current technology, or are there fundamental obstructions I have missed? Are there natural further weakenings (or strengthenings) that would make the problem more attractive? I am aware that the problem is far from my own expertise. My hope is that the formulation is clear enough to be useful, and that the transparent account of how it was reached may illustrate one way AI tools can assist in exploring and refining questions in analytic number theory. Thank you for any comments or references.

论坛问题帖 | has_answers | Kouros Goodarzi | 2026-09-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018434 | MO-515517

What open problems seems to be difficult enough not to be resolved soon (despite AI advances)?

What open problems seems to be difficult enough not to be resolved soon (despite AI advances) ? Resolved by any means - AI, humans, AI+humans.

论坛问题帖 | has_answers | Alexander Chervov | 2026-09-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018435 | MO-515678

Is every generalized variety finitely accessible?

Is every generalized variety (definition below) finitely accessible? If not, what is a counterexample? A finitely accessible category is a category with filtered colimits and an essentially small collection of finitely presentable objects that generate the category under filtered colimits; here an object is finitely presentable when its hom-functor preserves filtered colimits. Replacing "filtered colimits" with "sifted colimits", we obtain the notion of a generalized variety, studied by Adamek-Rosicky in On sifted colimits and generalized varieties (TAC): this is a category with sifted colimits and an essentially small collection of strongly finitely presentable objects that generate the category under sifted colimits; here an object is strongly finitely presentable when its hom-functor preserves sifted colimits. In the end of the mentioned paper, the authors remark that every generalized variety is $\aleph_1$-accessible, and they ask if every generalized variety is finitely accessible. This was an open problem when the paper was written. But this was 25 years ago, so maybe we have more insights now. Among the 52 generalized varieties currently recorded in CatDat, all are finitely accessible. But there are many finitely accessible categories that are no generalized varieties. Every cocomplete generalized variety is a variety (Theorem 3.10 in op. cit.) and therefore finitely accessible.

论坛问题帖 | has_answers | Martin Brandenburg | 2026-10-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018436 | MO-516336

Has AI made any progress as a mathematical theory-builder, rather than just a problem-solver?

In his essay The Two Cultures of Mathematics, Timothy Gowers distinguishes between two broad mathematical cultures: problem-solvers and theory-builders. Recent advances in AI-assisted mathematics seem to demonstrate increasingly impressive problem-solving abilities, including progress on difficult open problems. However, I wonder whether comparable progress has been made on the other side of Gowers's distinction. By theory-building, I mean something more than proving new theorems or solving existing conjectures: for example, introducing genuinely useful mathematical concepts, definitions, structures, or frameworks that unify previously disconnected results, suggest new research directions, or fundamentally change how a mathematical subject is understood. Are there documented examples of AI systems making original contributions of this kind to mathematical research? I am particularly interested in cases where the AI contributed substantially to the conceptual framework itself, rather than merely helping prove results within a framework already developed by humans. I realize that the distinction is not sharp, and that the originality and mathematical significance of a new conceptual framework can be difficult to assess. I would welcome references to concrete examples, including unsuccessful but instructive attempts, and any published assessments of how close current systems have come to mathematical theory-building in this sense.

论坛问题帖 | has_answers | Nick Belane | 2026-10-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018437 | MO-516346

Are all odd abundant numbers nearly half-representable by distinct divisors?

Let n be an odd abundant number, that is, $ \sigma(n)>2n, $ where $\sigma(n)$ denotes the sum of the positive divisors of n. Question. Is it true that for every odd abundant number n, there exists a subset S of its positive divisors such that $$ \sum_{d\in S}d=\frac{n+1}{2}\,? $$ For example, the smallest odd abundant number is 945, and $

\frac{945+1}{2}=473=315+105+35+15+3, $ where all five summands are distinct divisors of 945. I was unable to find counterexemples. Indeed, I suspect that a much stronger property holds, and namely that there is a suitable (quite large?) interval centered in $n/2$ of positive integers all representable as a sum of distinct divisors of n. This question is related to pseudoperfect and weird numbers, but differs from the usual subset-sum representation of n by its proper divisors. However a positive answer could give an insight to the longstanding question of the existence (or non-existence) of odd weird numbers.

论坛问题帖 | has_answers | G. Melfi | 2026-10-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018438 | MO-51680

Amenability of groups in terms of a perturbation condition

Let G be a countable group and $\lambda \colon G \to U(\ell^2 G)$ its left-regular representation. Suppose that there exists a constant $C>0$ such that for all $T \in B(\ell^2 G)$ $$\inf \lbrace\|T-S\| \mid S \in \lambda(G)' \rbrace \leq C \cdot \sup\lbrace \|\lambda(g)T-T \lambda(g) \| \mid g \in G\rbrace.$$ (Here $\lambda(G)'$ denotes the commutant of G in $B(\ell^2 G)$.) Question: Is G amenable? It is fairly easy to see that amenability of G implies the existence of such a constant. Indeed, one may take S to be some fixed point for the conjugation action on $\overline{\rm conv}\lbrace \lambda(g)T\lambda(g)^* \mid g \in G\rbrace$. I am asking for the converse of this statement. EDIT: Since the derivation problem came up in Kate's comment, I want to clarify to what version of it my question is related. The inequality above holds for some C if and only if the first bounded cohomology of G with coefficients in $B(\ell^2 G)$ (with the conjugation action induced by λ) is reduced. This is a straightforward application of the open mapping theorem. Now, two things are unclear: Question: Can $H^1_b(G,B(\ell^2 G))$ be reduced without being zero? and Question: Can $H^1_b(G,B(\ell^2 G))$ be zero without G being amenable?

论坛问题帖 | has_answers | Andreas Thom | 2011-01-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018439 | MO-52552

Nontrivial finite group with trivial group homologies?

I stumbled across this question in a seminar-paper a long time ago: Does there exist a positive integer N such that if G is a finite group with $\bigoplus_{i=1}^NH_i(G)=0$ then $G=\lbrace 1\rbrace$? I believe this to still be an open problem. For $N=1$, any perfect group (ex: A_5) is a counterexample. For $N=2$, the binary icosahedral group $SL_2(F_5)$ suffices (perfect group with periodic Tate cohomology). And I found in one of Milgram's papers a result for $N=5$, the sporadic Mathieu group M_{23}. Note that this question is answered for infinite groups, because we can always construct a topological space (hence a BG for some discrete group G) with prescribed homologies. Is there another known group with a larger $N\ge 5$ before homology becomes nontrivial? Are there any classifications of obstructions in higher homology groups? [[Edit]]: Another view. A group is $\textit{acyclic}$ if it has trivial integral homology. There are no nontrivial finite acyclic groups. Indeed, a result of Richard Swan says that a group with p-torsion has nontrivial mod-p cohomology in infinitely many dimensions, hence nontrivial integral homology.

论坛问题帖 | has_answers | Chris Gerig | 2011-01-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018440 | MO-52913

Free subgroups vs law

Consider the following two conditions for a group G: (1) G does not satisfy a nontrivial law. (2) G contains a non-abelian free subgroup. Obviously (2) implies (1) and it is easy to construct torsion groups that do not satisfy any law (e.g., the direct product of all finite groups). Thus (1) does not imply (2) in general. However the following question seems open: Are (1) and (2) equivalent for profinite groups? Here is a similar question: Suppose that a residually finite group does not satisfy a law. Does its profinite completion contain a nonabelian free subgroup? These questions may be thought of as possible generalizations of the Tits alternative to residually finite (or profinite) groups. The answer to the second question is positive for finitely generated p-groups. This follows from [Wilson, Zelmanov, Identities for Lie algebras of pro-p groups, JPAA 81(192), 103-109], where the authors prove that if G is a finitely generated residually finite p-group, then either it is finite (and hence satisfies a law), or its profinite completion contains a nonabelian free group. In the general case they only prove that if the profinite completion of a residually finite group does not contain a nonabelian free subgroup, then some Lie algebra associated to the group satisfies a law.

论坛问题帖 | has_answers | Denis Osin | 2011-01-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018441 | MO-5340

Is every matching of the hypercube graph extensible to a Hamiltonian cycle

Given that Q_d is the hypercube graph of dimension d then it is a known fact (not so trivial to prove though) that given a perfect matching M of Q_d ($d\geq 2$) it is possible to find another perfect matching N of Q_d such that $M \cup N$ is a Hamiltonian cycle in Q_d. The question now is - given a (non necessarily perfect) matching M of Q_d ($d\geq 2$) is it possible to find a set of edges N such that $M \cup N$ is a Hamiltonian cycle in Q_d. The statement is proven to be true for $d \in\{2,3,4\}$.

论坛问题帖 | has_answers | Jernej | 2009-11-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018442 | MO-53534

Comparing the Kazhdan-Lusztig and Steinberg pre-orders

Both Kazhdan-Lusztig and Steinberg have defined pairs of preorders on S_n. Kazhdan and Lusztig's preorders come from their basis: We write $x\leq_L y$ if any left ideal spanned by K-L basis vectors which contains C_y also contains C_x. Similarly for $x\leq_R y$ and right ideals. Note that $x\leq_L y$ if and only if $x^{-1}\leq_R y^{-1}$, so all the information is in one of these preorders. Now, we can define a geometric preorder satisfying similar properties, following Steinberg. Let $\mathfrak{n}$ be the usual strictly upper-triangular matrices in $\mathfrak{sl}_n$ and let $\mathfrak{n}\cap\mathfrak{n}^w$ be its

intersection with itself conjugated by w. We let $x\leq_l y$ if $B\cdot (\mathfrak{n}\cap\mathfrak{n}^y)\subset B\cdot (\mathfrak{n}\cap\mathfrak{n}^x)$, and $x\leq_r y$ if $x^{-1}\leq_ly^{-1}$. Do the preorders $\leq_L$ and $\leq_l$ coincide? It's a theorem (look, for example in Appendix B of Borho and Brylinski, Differential... III) that the equivalence relations induced by them (the left cells) do; they're both distinguished by their Q-symbol under Robinson-Schensted. Do these preorders have a nice description in terms of Robinson-Schensted? EDIT: After speaking with various people, I believe this is an open question, and quite an interesting one in my opinion.

论坛问题帖 | no_answers_at_retrieval | Ben Webster | 2011-01-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018443 | MO-53797

Open problems in Euclidean geometry?

What are some (research level) open problems in Euclidean geometry ? (Edit: I ask just out of curiosity, to understand how -and if- nowadays this is not a "dead" field yet) I should clarify a bit what I mean by "Euclidean geometry". By this term I mean, loosely, the study of the geometry of certain subsets of Euclidean space $\mathbb{E}^n$ from a point of view which is either the "classical" one (i.e. axiomatic), or one that involves more modern tools, but the problem in question has not to be "clearly" a problem within some other branch of maths such as differential or algebraic geometry, algebraic or general topology, analysis, or measure theory. Some examples to clarify: the study of configurations of lines or affine subspaces is EG; but the algebro-topological study of hyperplane arrangements is not. plane conics as defined via their metric property are objects of EG; but "algebraic curves" are not, unless they're defined by some "elementary enough property" (intentionally vague) involving the Euclidean metric. root systems of Lie algebras are EG. polyhedral cones are EG. polytopes are EG. tessellations of space with polytopes or analogous objects are in EG. minimal surfaces in $\mathbb{E}^3$ are not EG. fractal geometry (Julia sets, self-affine fractals...) is not EG. not sure about convex bodies. If they're polyhedral I'd say their study fits in EG. packings of spheres are EG.

论坛问题帖 | has_answers | Qfwfq | 2011-01-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018444 | MO-54143

What's the Kirby Diagram of a universal $\mathbb{R}^4$?

What's the Kirby diagram of a universal $\mathbb{R}^4$? Background Define $\mathcal{R}$ as the set of smoothings of $\mathbb{R}^4$. For two oriented elements R_1, R_2 in $\mathcal{R}$ we can define their end sum $R_1 \natural R_2$ if we are given two proper embeddings $\gamma_i : [0, \infty) \rightarrow R_i$. We remove a tubular neighborhood of $\gamma_i((0, \infty))$ from each R_i and glue the resulting $\mathbb{R}^3$ boundaries together respecting orientations. The result is the end sum $R_1 \natural R_2$ of R_1 and R_2. As γ_i is unique up to ambient isotopy, $R_1 \natural R_2$ is well defined up to diffeomorphism. In "A universal smoothing of four-space" Freedman and Taylor proved the existence of an element $U \in \mathcal{R}$ such that for any $R \in \mathcal{R}$ the end sum $U \natural R$ is diffeomorphic to U. This U is the universal $\mathbb{R}^4$. Foreground In "An invariant of smooth 4-manifolds" Taylor defines an invariant $\gamma(R) \in \{0,1,2,\ldots,\infty \}$ for $R \in \mathcal{R}$. Taylor defines $\gamma(R)$ to be $sup_K \{ min_X\{ \frac{1}{2} b_2(X) \} \}$, where K ranges over compact 4-manifolds smoothly embedding in R and X ranges over closed, spin 4-manifolds with signature 0 in which K smoothly embeds. (Actually, Taylor defines γ for all smooth 4-manifolds, but we don't need this detail here.) Taylor goes on to prove that if $R \in \mathcal{R}$ and $\gamma(R) > 0$, then any handle decomposition of R has infinitely many three handles. In "4-Manifolds and Kirby Calculus" Stipsicz and Gompf prove, see page 376, that $\gamma(U) = \infty$. Thus, any Kirby Diagram of U must have infinitely many three handles. Question What's the Kirby diagram of a such a U?

论坛问题帖 | has_answers | Kelly Davis | 2011-02-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018445 | MO-5479

Milnor's cartography problem

Let Ω be a round disc of radius $\alpha<\frac{\pi}{2}$ on the unit sphere $\mathbb{S}^2$. It is easy to construct a $(1,\tfrac{\alpha}{\sin\alpha})$-bi-Lipschitz map from Ω to the plane. Is it true that any convex domain Ω' on $\mathbb{S}^2$ with the same area as Ω also admits a $(1,\tfrac{\alpha}{\sin\alpha})$-bi-Lipschitz map to the plane? Comments: This problem appears in Milnor's A problem in cartography. Amer. Math. Monthly 76 1969 1101--1112. I spent quite a bit of time to solve it, but without success. I only noticed that if one exchange "area" above to "perimeter" then the answer is YES. (To prove the statement appy the idea from "A New Area Maximization Proof for the Circle" by Lawlor.)

论坛问题帖 | has_answers | Anton Petrunin | 2009-11-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018446 | MO-55890

Card game / options pricing / Brownian bridge question

We play a game. I shuffle a deck of cards and start dealing them face up. After any card you can say "stop", at which point I pay you 1 dollar for every red card dealt and you pay me 1 for every black card dealt. What is your optimal strategy, and how much would you play to pay this game? Clearly the game is worth at least 0, since you can follow the strategy 'wait until all the cards are dealt' and get paid 0. We can conclude that at any stage of the game, we will ask for another card if the expected value of continuing to play is greater than the amount of money already won; otherwise we will say stop. Mathematically, if r is the number of red cards remaining and b is the number of black cards remaining, and $f_{r,b}$ is the expected value of the game at this point, then the expected value of taking another card is $e_{r,b} = \frac{r}{r+b} f_{r-1,b} + \frac{b}{r+b} f_{r,b-1}$ and hence the value of the game at this point, assuming we play the optimal strategy, is $f_{r,b} = \max (\frac{r}{r+b} f_{r-1,b} + \frac{b}{r+b} f_{r,b-1}, b-r)$ with $f_{r,0} = 0$ (if there are only red cards remaining, we have a guaranteed payout of 0) and $f_{0,b} = b$ (if there are only black cards remaining we won't take any more cards). Solving numerically gives an

expected value for this game of 2.624 with 52 cards. I'm interested in what the value is for a deck of $2n$ cards, and if the optimal strategy can be expressed analytically as a function of r and b (ie keep taking cards until you have made $g(r,b)$ dollars, then stop). I've tried changing coordinates to $u=b-r$, $v=b+r$, and I've tried approximating the difference equation as a PDE, but no luck with an analytical solution yet - or even any decent approximations.

论坛问题帖 | has_answers | Chris Taylor | 2011-02-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018447 | MO-57078

A question on a special type of function

Suppose I have a function f, and positive integers x and y such that x is a square, $x \ne y$ and $y \ne \sqrt{x}$. Now, assume that: $|\frac{f(x)}{y} - \frac{f(y)}{x}| > 2$ $|\frac{f(\sqrt{x})}{y} - \frac{f(y)}{\sqrt{x}}| > 1$ for all x, y in the domain of f. (Note that $|N|$ is the absolute value of N.) My question is: What properties would this function f necessarily possess? In particular, can f be injective?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2011-03-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018448 | MO-57158

Existence of Solutions to an Equation Involving the Sum-of-Divisors Function [Reference Request]

Let $\sigma(x) = \sigma_1(x)$ denote the sum of all the positive divisors of x. If $n \in \mathbb{N}$ is odd and $\gcd(n, \sigma(n)) = 1$, then do there exist any solutions to the following equation? $$2{n^2}\sigma(n) = \sigma({n^2})\sigma(\sigma(n))$$ In other words, does there exist such an odd $N = {n^2}\sigma(n)$ which is also a perfect number? Here, I also list some (open?) problems mentioned by other researchers: (1) Suryanarayana: Is it true that every odd perfect number is of the form $m\sigma(m)$ for some odd integer m; if so, is $\gcd(m, \sigma(m)) = 1$ necessarily? (2) M. V. Subbarao: Does every odd perfect number n (if such exist) have the representation $$n = \frac{1}{2}m\sigma(m) \hspace{0.5in} (*)$$ Another question: Whenever n given by $(*)$ is perfect, does it follow that n is odd and $\gcd(m, \sigma(m)) = 1$? I would appreciate it if anybody could point me to reference(s) to recent results on either Problem (1) or Problem (2).

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2011-03-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018449 | MO-57436

A simple question regarding the sum-of-divisors function

A good day to everyone. Consider the following "Conjecture": If $a, b \in M \subset \mathbb{N}$, then $1 < a < b$ and ... [plus some more conditions on a, b and M...] if and only if $\sigma(a) < \sigma(b)$. I have two questions at this point: (1) What properties should elements of the set M have to satisfy this "Conjecture"? (2) What happens to the sets in M if you restrict to the case $\gcd(a, b) = 1$? (Of course, the sets in (1) should have greater asymptotic density than the sets in (2), but can this notion be made more precise?) Now, for the motivation: An example of a set M satisfying this "Conjecture" is any pair (A, B) of relatively prime factors of an odd perfect number given in the Eulerian form $N = {q^k}{n^2}$ (modulo some exceptions), where $\gcd(q, n) = 1$ and $q \equiv k \equiv 1 \pmod4$. Let $$I(x) = \displaystyle\frac{\sigma(x)}{x}$$ be the abundancy index of $x \in \mathbb{N}$. Then by enumerating all possible permutations of the set $$q^k, n, \sigma(q^k), \sigma(n) \in \mathbb{N}$$ we know that exactly one of the following holds (because of $1 < I(q^k) < I(n)$): $$[1] \hspace{0.2in} q^k < n < \sigma(q^k) < \sigma(n)$$ $$[2] \hspace{0.2in} q^k < \sigma(q^k) < n < \sigma(n)$$ $$[3] \hspace{0.2in} n < q^k < \sigma(q^k) < \sigma(n)$$ $$[4] \hspace{0.2in} n < q^k < \sigma(n) < \sigma(q^k)$$ $$[5] \hspace{0.2in} n < \sigma(n) < q^k < \sigma(q^k)$$ Last question: Which of these five "configurations" could we eliminate?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2011-03-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018450 | MO-58113

Kronecker Approximation theorem and Fibonacci numbers

There is a famous old theorem by Kronecker that for every positive real α and $\epsilon>0$ there exists a positive integer n such that αn is within ϵ of an integer. Recently I found that the same result is true if we replace αn by αn^2 or any polinomial p such that $p(0)=0$. Could this result be generalised to other functions? Particularly I'm curious about sequences $\alpha 2^n$ and αF_n where by F_n I denote n-th Fibonacci number. Does anyone know anything about it?

论坛问题帖 | has_answers | Ostap Chervak | 2011-03-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018451 | MO-58794

Bounds on number of conjugacy classes in terms of number of elements of a group ?

What are bounds on number of conjugacy classes in terms of number of elements of a group ? (I allowed myself to edit the question in spirit of remarkable answers given to it by Gerry Myerson and Geoff Robinson. Below is original text of the question. (Alexander Chervov)). It's about the first step to find an upper bound to the order of a finite group with h conjugacy classes (right or left) that depends only on h. (h a natural non nul integer). I have some doubts about the rigor of my proof that I am sharing with you so that you can help me find a likely error or an omited step. I have attached the scan of my proof to this post. Many thanks

论坛问题帖 | has_answers | Averroes | 2011-03-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018452 | MO-59046

Can you flip the end of a large exotic $\mathbb{R}^4$

Can you flip the end of a large exotic $\mathbb{R}^4$ Background Definition (Exotic $\mathbb{R}^4$): An exotic $\mathbb{R}^4$ is a smooth manifold R homeomorphic but not diffeomorphic to $\mathbb{R}^4$, where $\mathbb{R}^4$ is equipped with its

standard smooth structure. Definition (Large exotic $\mathbb{R}^4$): A large exotic $\mathbb{R}^4$ is an exotic $\mathbb{R}^4$ containing a four-dimensional compact smooth submanifold K' that can not be smoothly embedded into $\mathbb{R}^4$. Definition (End of a large exotic $\mathbb{R}^4$): If R is a large exotic $\mathbb{R}^4$ and D^4 is a four-dimensional disk topologically embedded into R such that $K' \subset D^4$, then $R - D^4$ is an end of R. Remark: The previous definition varies slightly from the standard definition of "end", however it will be used for the remainder of this question. (See Gompf and Stipsicz Exercise 9.4.11 for the standard definition.) Remark: If $R - D^4$ is the end of a large exotic $\mathbb{R}^4$, then $R - D^4$ is a smooth manifold that inherits a smooth structure from R. Definition (Flip of the end of a large exotic $\mathbb{R}^4$): Given $R - D^4$, the end of a large exotic $\mathbb{R}^4$, a flip of $R - D^4$ is a diffeomorphism $f: R - D^4 \rightarrow R - D^4$ that maps the "inner region" of $R - D^4$, that "near" the removed D^4, to the "outter region", that "near infinity", and vica-versa. Remark: The previous definition is also non-standard. I am not aware of any standard definitions that carry, more-or-less, the same meaning. So, at this stage the meaning of the question is hopefully clear. Foreground In our attempt to flip the end of a large exotic $\mathbb{R}^4$ an inconvenient truth stands in our way: Theorem 1 (Uncountably many flips fail): There are uncountably many large exotic $\mathbb{R}^4$'s that one can not flip the end of. Quickly, let us see why this is true. Lemma 9.4.2 along with Addendum 9.4.4 of Gompf and Stipsicz state: Lemma 1: There exist pairs (X,Y) and (L,K) of smooth, oriented four-manifolds with X simply connected, Y and K compact, X and L open (i.e. noncompact and boundaryless), L homeomorphic to $\mathbb{R}^4$, and X with negative definite intersection form not isomorphic to $n\langle-1\rangle$, such that $X - int(Y)$ and $L - int(K)$ are orientation-preserving diffeomorphic. Theorem 9.4.3 of Gompf and Stipsicz states: Theorem: Any L as appears in Lemma 1 is a large exotic $\mathbb{R}^4$. The two statements above lead to: Lemma: One can not flip the end of any L as appears in Lemma 1. Proof: Assume one could flip the end of L. Thus, one could use this flip to glue L to the "end" of X and obtain a simply connected closed smooth four-manifold with negative definite intersection form not isomorphic to $n\langle-1\rangle$. However, according to Donaldson's Theorem (Gompf and Stipsicz Theorem 1.2.30) there exists no such manifold. Thus, there exists no such flip. QED Now we have shown that L can not be flipped. Before we show how uncountably many large exotic $\mathbb{R}^4$'s can not be flipped, we need the definition: Definition (Radial Family): Let R be an exotic $\mathbb{R}^4$. Thus, there exists a homeomorphism $h:\mathbb{R}^4 \rightarrow R$. Define R_t as the image under h of the open ball of radius t centered at 0 in $\mathbb{R}^4$. A radial family is a set of the form $\{R_t | 0 < t \le \infty \}$. Remark: If R_t is a member of a radial family, then R_t is a smooth manifold as it inherits a smooth structure from R Theorem 9.4.10 of Gompf and Stipsicz states: Theorem: If $\{L_t | 0 < t \le \infty \}$ is a radial family for an L as appears in Lemma 1 and r is such that $K \subset L_r$, then $\{L_t | r \le t \le \infty \}$ is an uncountable family of non-diffeomorphic large exotic $\mathbb{R}^4$'s. This leads directly to a proof of Theorem 1. Proof: Assume one could flip the end of L_t for $r \le t \le \infty$, where all notation is as in the previous theorem. Thus, one could use this flip to glue L_t to the "end" of X less the image of $L - L_t$ and obtain a simply connected closed smooth four-manifold with negative definite intersection form not isomorphic to $n\langle-1\rangle$. Again, according to Donaldson's Theorem (Gompf and Stipsicz Theorem 1.2.30) there exists no such manifold. Thus, there exists no such flip.QED Things are seeming rather hopeless at this point. In fact, things are worse than they seem! But, before we can revel in this despair, we must introduce two definitions: Definition (Simply Connected at Infinity): Let Z be a topological manifold. Z is simply connected at infinity if for any compact subset C of Z there exists a compact subset C' of Z that contains C and is such that the inclusion $Z - C' \rightarrow Z - C$ induces the trivial map $\pi_1(Z - C') \rightarrow \pi_1(Z - C)$. Definition (End Sum): Let Z_1 and Z_2 be non-compact oriented smooth four-manifolds that are simply connected at infinity. Choose two proper smooth embeddings $\gamma_i : [0, \infty) \rightarrow Z_i$. Remove a tubular neighborhood of $\gamma_i((0, \infty))$ from each Z_i and glue the resulting $\mathbb{R}^3$ boundaries together respecting orientations. The result is the end sum $Z_1 \natural Z_2$ of Z_1 and Z_2. Remark: The requirement that Z_i is simply connected at infinity guarantees that γ_i is unique up to ambient isotopy and thus $Z_1 \natural Z_2$ is unique up to diffeomorphism (Gompf and Stipsicz Definition 9.4.6). Remark: If R_1 and R_2 are exotic $\mathbb{R}^4$, then they are non-compact oriented smooth four-manifolds that are simply connected at infinity and $R_1 \natural R_2$ is a smooth manifold homeomorphic to $\mathbb{R}^4$. Remark: X of Lemma 1 is simply connected at infinity. Theorem 2: If $\{L_t | 0 < t \le \infty \}$ is a radial family for an L as appears in Lemma 1 and r is such that $K \subset L_r$, where K is as in Lemma 1, then for R an exotic $\mathbb{R}^4$ and t such that $r \le t \le \infty$ there exists no flip of $R \natural L_t$. Proof: The proof is basically a slight variation on the above theme. Assume one could flip the end of $R \natural L_t$ for $r \le t \le \infty$. Thus, one could use this flip to glue $R \natural L_t$ to the "end" of X less the image of $L - L_t$ end summed with R, in other words with the flip glue $R \natural L_t$ to $R \natural (X - (L - L_t))$, and obtain a simply connected closed smooth four-manifold with negative definite intersection form not isomorphic to $n\langle-1\rangle$. Again, according to Donaldson's Theorem (Gompf and Stipsicz Theorem 1.2.30) there exists no such manifold. Thus, there exists no such flip.QED Now we can revel in this despair! However, other ways of creating large exotic $\mathbb{R}^4$'s exist. For example, given a topologically slice knot that is not smoothly slice one can create a large exotic $\mathbb{R}^4$. (See, for example, Davis.) Such a large exotic $\mathbb{R}^4$, as far as I can see, might admit an end flip. But, I'm not sure. Thus, we end where we began. Question Can you flip the end of a large exotic $\mathbb{R}^4$?

论坛问题帖 | has_answers | Kelly Davis | 2011-03-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018453 | MO-59082

A question concerning products of finite cyclic groups

Let $m_1,\ldots, m_n$ be pairwise coprime natural numbers ≥ 1. We consider the product $$G(m_1,\ldots,m_n) := \prod_{i=1}^{n} \mathbb{Z} / m_i \mathbb{Z}.$$ We define $M(n)$ as the set n-tuple of natural numbers ≥ 1 with the property that the entrys are pairwise coprime. We define $l : M(n) \rightarrow \mathbb{N}_0$ by $$l(m_1,\ldots,m_n) = \max_{(x_1,\ldots,x_n) \in G(m_1,\ldots,m_n)} \min \{ k \in \mathbb{N}_0 | x_i+k \neq 0 \forall 1 \leq i \leq n \},$$ hence $l(m_1,\ldots,m_n)$ may be regarded as the maximal distance of the elements of G from the elements which have no identity in their entrys. It is clear, that $l(m_1,\ldots,m_n)$ is always a natural number. The question is now: Does there exist a real number r, such that $r \cdot n \cdot \ln (n) \geq

\sup_{(m_1,\ldots,m_n) \in M(n)} l(m_1,\ldots,m_n)$$ holds for all $n >> 0$ and is it possible to take $r=2$?

论坛问题帖 | has_answers | tobias | 2011-03-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018454 | MO-60011

Trees in groups of exponential growth

Question: Let G be a finitely generated group with exponential growth. Is there a finite generating set $S \subset G$, such that the associated Cayley graph $Cay(G,S)$ contains a binary tree? Some background: The existence of such a tree clearly implies exponential growth. Kevin Whyte showed in Amenability, Bilipschitz Equivalence, and the Von Neumann Conjecture, Duke Journal of Mathematics 1999, p. 93-112, that such trees exist if G is non-amenable. So the question is only open for amenable groups of exponential growth. One good reason for such a binary tree to exist is the existence of a free semigroup inside G. In fact, if G is solvable, then the existence of such a semigroup is known to be equivalent to exponential growth (and equivalent to being not virtually nilpotent). This is part of some version or extension of the Tits alternative. Grigorchuk constructed an amenable torsion group with exponential growth, which does not contain such a semigroup, but it contains a binary tree. EDIT: Al Tal pointed out in an answer below that Benjamini and Schramm covered the non-amenable case (this is 2. from above) already in Benjamini and Schramm "Every Graph With A Positive Cheeger Constant Contains A Tree With A Positive Cheeger Constant", GAFA, 1997.

论坛问题帖 | has_answers | Andreas Thom | 2011-03-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018455 | MO-60864

Symplectic structures on a homotopy complex projective space

For $n>2$, there are infinitely many differentiable structures on the homotopy type of $\mathbb{C}P^n$. I want to know which differentiable structures support a symplectic form. My question is as follows. Let M be a closed symplectic manifold which is homotopy equivalent to a complex projective space. Can we say that M is homeomorphic or diffeomorphic to the standard one?

论坛问题帖 | has_answers | Hwang | 2011-04-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018456 | MO-61309

Fixed point theorem for convex, closed multivalued mapping

There is well-known fixed point theorem theorem for multivalued l.s.c. maps, based on Michael selection theorem: Suppose, that X is compact, convex and metrizable in locally convex Hausdorff topological vector space. Then any l.s.c. map $F:X\rightarrow X$ which is closed and convex valued has a fixed point, i.e. $x\in F(x)$. The question is, what happens if we drop the assumption that X is metrizable. Xian Wu in his paper "A new fixed point theorem and its application" left it as an open problem, after giving the proof using a metrizability.

论坛问题帖 | no_answers_at_retrieval | Maciej Skorski | 2011-04-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018457 | MO-62548

Zero divisor conjecture for finite fields

I believe that the most attractive "zero-divisor" conjecture is the existence of non-trivial zero-divisors in a group ring $\mathbb{C}[G]$ for a torsion free group G. For the sake of knowledge let me ask if the zero conjecture is known for finite fields instead of $\mathbb{C}$. More precisely: Let G be a torsion-free group and K be a finite field. Is it true that the group ring $K[G]$ has no non-trivial zero divisors? In particular: Let K be the field with two elements, G be a torsion free group and let $rank(a)$ be the smallest number of elements in the expression of a in the sum $a=\sum g_i$, $g_i\in G$. Is there a constant $R>0$ such that we know for sure that $rank(a)>R$ for every zero-divisor $a\in K[G]$? Here is related question: Group ring and left zero divisor.

论坛问题帖 | has_answers | Kate Juschenko | 2011-04-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018458 | MO-64099

The maximum of a polynomial on the unit circle

Encouraged by the progress made in a recently posted MO problem, here is a "conceptually related" problem originating from a 2003 joint paper of Sergei Konyagin and myself. Suppose we are given n points $z_1,...,z_n$ on the unit circle $U=\{z\colon |z|=1\}$ and n weights $p_1,...,p_n\ge 0$ such that $p_1+....+p_n=n$, and we want to find yet another point $z\in U$ to maximize the product $$ \prod_{i=1}^n |z-z_i|^{p_i}. $$ How large can we make this product by the optimal choice of z? Conjecture. For any given $z_1,...,z_n\in U$ and $p_1,...,p_n\ge 0$ with $p_1+...+p_n=n$, there exists $z\in U$ with $$ \prod_{i=1}^n |z-z_i|^{p_i} \ge 2. $$ Here are some comments. If true, the estimate of the conjecture is best possible, as evidenced by the situation where the points are equally spaced on U and all weights are equal to 1. We were able to resolve a number of particular cases; say, that where the points z_i are equally spaced on U, and also that where all weights are equal to 1. The case $n=2$ is almost trivial, but already the case $n=3$ is wide open. In the general case we have shown that the maximum is larger than some absolute constant exceeding 1. Although this is not obvious at first glance, this conjecture is actually about the maxima of polynomials on the unit circle. I would be very interested to see any further progress!

论坛问题帖 | has_answers | Seva | 2011-05-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018459 | MO-65194

Majorization and Schur Polynomials

Let me first define the majorization order (or dominance order) on partitions as $\lambda \succeq \mu$ iff $$\sum _{i=1}^{k}\lambda_i \geq \sum_{i=1}^{k}\mu_i$$ for all $1\le k\le l-1$ and $$\lambda_1+\cdots+\lambda_l=\mu_1+\cdots+\mu_l.$$ While playing around with some variation on Muirhead's inequality which says that $$m_{\lambda}(x_1,x_2,\dots,x_n)\geq m_{\mu}(x_1,x_2,\dots,x_n)$$ for all $x_i\geq 0$ when $\lambda \succeq \mu$ (here the m_{λ}'s are the monomial symmetric polynomials) I ended up investigating if such an inequality holds for Schur polynomials, which are another famous basis of the ring of symmetric functions. I tried a few special cases, and the following seems to hold, but I don't have a proof or a counterexample. So the question is: Let $x_1,\dots,x_n \geq 0$ and $\lambda\succeq \mu$, is it always true that $$\frac{s _{\lambda}(x _1,x _2,\dots,x _n)}{s _{\lambda}(1,1,\dots,1)}\geq \frac{s _{\mu}(x _1,x _2,\dots,x _n)}{s _{\mu}(1,1,\dots,1)}?$$ If not, does majorization induce some similarly nice order on values of Schur polynomials?

论坛问题帖 | has_answers | Gjergji Zaimi | 2011-05-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018460 | MO-6536

Motivation for uniform surjectivity of mod l representations associated to elliptic curves

Background Let E be an elliptic curve over $\mathbb{Q}$ and let $G_{\mathbb{Q}}$ be the absolute Galois group $Aut(\overline{\mathbb{Q}})$. For any positive integer n the n-torsion subgroup $E[n](\overline{\mathbb{Q}})$ is stable under the $G_{\mathbb{Q}}$-action. Since $E[n](\overline{\mathbb{Q}})$ is isomorphic to $(\mathbb{Z}/n\mathbb{Z})^2$ one gets a continuous (with respect to the profinite topology on the left and the discrete on the right) homomorphism $$ \overline{\rho_{E,\, n}}\colon G_{\mathbb{Q}} \to GL_2(\mathbb{Z}/n\mathbb{Z}) $$ which one calls the mod n representation associated to E. As n varies these are compatible and taking limits gives representations $\rho_{E,\ell^{\infty}}$ and ρ_E with values in $GL_2(\mathbb{Z}_l)$ and $GL_2(\widehat{\mathbb{Z}})$ which one calls respectively the ℓ-adic and adelic representations associated to E. Alternatively, $\rho_{E,n}$ is isomorphic to the representation induced by the action of $G_{\mathbb{Q}}$ on the etale cohomology $H^1_{\text{et}}(E_{\overline{\mathbb{Q}}}; \mathbb{Z}/n\mathbb{Z})$; the description of $\rho_{E,n}$ via torsion generalizes to give representations $\rho_{A,n}$ for higher dimensional abelian varieties, but for a general variety one must instead use cohomlogy. Serre famously proved that for E an elliptic curve with $End(E) = \mathbb{Z}$, $\rho_E(G_{\mathbb{Q}})$ has finite index in $GL_2(\widehat{\mathbb{Z}})$. In particular for ℓ large $\rho_{l^{\infty}}$ is surjective; how large one must take ℓ depends on E. Conjecture This last fact does not depend on E. I.e. there exists a constant N such that for every $E/\mathbb{Q}$ with $End(E) = \mathbb{Z}$ and $\ell \geq N$, the mod ℓ representation $\overline{\rho_{E,\ell}}$ is surjective; equivalenty there is an upper bound (independent of E) on the index of $\rho_E(G_{\mathbb{Q}})$. By a recent paper of Bilu and Parent, one knows that the image is either surjective or contained in a non-split cartan subgroup. My Question Why do people expect this to be true? Does it follow from other believable conjectures? Is there some heuristic that predicts this? One fact is that one can phrase this question as the failure of various modular curves (e.g. $X_{ns}(l)$, which have increasingly large genus) to have non-trivial rational points. Is there a geometric reason that one expects these modular curves to have no non-trivial rational points?

论坛问题帖 | has_answers | David Zureick-Brown | 2009-11-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018461 | MO-65858

Series whose convergence is not known

For most of the mathematical concepts I learn, it has more or less always been possible to find (at least google and find) unsolved problems pertaining to that specific concept. Keeping a bag of unsolved problems on most topics I know has been to my benefit in that it reaffirms me that mathematics is a thriving subject. Coming to point, I am unable to find an elementary series of the kind we know on real analysis courses whose convergence is an unsolved problem. Please, share if you have any. Thanks.

论坛问题帖 | has_answers | Chulumba | 2011-05-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018462 | MO-6627

Convex hull in CAT(0)

Let X be complete $\mathop{CAT}(0)$-space and $K\subset X$ be a compact subset. Is it true that convex hull of K is compact? Comments: Convex hull of K = intersection of all closed convex sets containing K. The space is NOT assumed to be locally compact This problem was mentioned in 6.B$_1$(f) of Gromov's "Asymptotic invariants of infinite groups" (1993). I believe there is a counterexample, maybe even in case of negatively pinched curvature (in the sense of Alexandrov).

论坛问题帖 | has_answers | Anton Petrunin | 2009-11-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018463 | MO-66613

A toy model for the t-section problem

Let $S(x)$ be the area of the yellow curvilinear triangle. I'd like to find a graph for which $S(x)=H(x)$ where $H $ is some prescribed function (small, smooth, vanishing near the endpoints to any order you wish, etc.). Is it always possible or there are some non-obvious hidden restrictions? alt text http://www.artofproblemsolving.com/Forum/latexrender/pictures/b/4/6/b46dbb397d96f17cce1e421bfaf29b9a80d26c81.png The question comes from the infamous t-section problem (if you know the areas of all sections of a symmetric convex body by the hyperplanes at some fixed small distance t from the origin (so small that all sections are non-empty), can you recover the body?). The problem is open even on the plane. I do not say that this toy question is directly relevant here but an answer to it will certainly make a few things clearer for me.

论坛问题帖 | has_answers | fedja | 2011-06-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018464 | MO-6690

What is known about polyhedra nets that allow overlapping?

It is an open problem that the net of any convex polyhedron can be unfolded onto a flat plane with no overlapping. Is anything known if we allow x faces to overlap? For example, is it known if any convex polyhedron can be unfolded with a maximum of 2 faces overlapping? What about the more general case of a polyhedron that is topologically convex (that is, its graph is isomorphic to the graph of a convex polyhedron)? This paper provides an example of an open polyhedron without a net that is topologically convex, and the closed case appears to be taken care of by this paper (mentioned in an answer below). One solution to the topologically convex case would then to be to find a procedure to modify either polygon so that the number of overlapping sides increases without bound. I have been unable to do so without breaking the topologically convex property, but it seems a reasonable task.

论坛问题帖 | has_answers | Jason Dyer | 2009-11-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018465 | MO-68950

Axiom of choice and bases of vector spaces over a fixed field

Let k be a field. In 1984 Andreas Blass proved that the axiom "for every extension $K|k$, every vector space over K has a basis" implies the axiom of choice. He also raised the question Does the axiom "every vector space over k has a basis" imply the axiom of choice ? What's the current status of the question ? Has there been progress ?

论坛问题帖 | no_answers_at_retrieval | Ralph | 2011-06-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018466 | MO-69099

Shortest closed curve to inspect a sphere

Let S be a sphere in $\mathbb{R}^3$. Let C be a closed curve in $\mathbb{R}^3$ disjoint from and exterior to S which has the property that every point x on S is visible to some point y of C, in the sense that the segment xy intersects S at precisely the one point x. I am interested in the shortest C with this property. In computational geometry, such paths are called watchman tours, and there are many results concerning polygons in the plane finding such tours. This question arose at a conference I'm attending, and I was pointed to a paper by V. A. Zalgaller: "Shortest Inspection Curves for the Sphere" (Journal of Mathematical Sciences, Volume 131, Number 1, 5307-5320; Translated from Zapiski Nauchnykh Seminarov POMI, Vol. 299, 2003, pp. 87-108.) I cannot access the paper from the conference, but from the abstract it appears he focused on open rather than closed curves. Has anyone heard of this natural question? Can you point me to relevant literature? Thanks! Addendum. Here is the 4π saddle / baseball-stitches curve suggested by Gjergji Zaimi:

论坛问题帖 | has_answers | Joseph O'Rourke | 2011-06-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018467 | MO-7016

When shorter means smaller?

Assume a convex figure $F\subset \mathbb R^2$ satisfies the following property: if $f:F\to \mathbb R^2$ is a distance-non-increasing map then its image $f(F)$ is congruent to a subset of F. Is it true that F is a round disk? Comments: It is easy to see that the round disk has this property. One can reformulate the property: if for some set $G\subset\mathbb R^2$ there is a distance-non-contracting map $G\to F$ then there is a distance-preserving map $G\to F$. (The equivalence follows from Kirszbraun's theorem) Some figures, such as the Reuleaux triangle are bad (see the comments below). The construction with two folds along parallel lines kills many examples, in particular, the following figure: the intersection of two discs, say the unit disc with center at (0,0) and the disc of radius 1.99 and center at (0,1) --- see the comments by Martin M. W. below. This construction also gives the following: If F is good then for any point $x\in \partial F$ the restriction of $\textrm{dist}_x$ to ∂F has no local minima other than x. (This property holds for any convex figure C^2-close to a round disc.) This problem was meant to be an exercise for school students, but I was not able to solve it :). It appeared in print in 2008 (in Russian), see Problem 5 in Плоское оригами и длинный рубль. One answer is accepted, BUT it only provides a solution for unbounded figures.

论坛问题帖 | has_answers | Anton Petrunin | 2009-11-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018468 | MO-70391

Convex bodies with constant maximal section function in odd dimensions

In 1970 or so, Klee asked if a convex body in $\mathbb R^n$ ($n\ge 3$) whose maximal sections by hyperplanes in all directions have the same volume must be a ball. The counterexample in $\mathbb R^4$ is trivial and can be described as follows: Let $f:[-1,1]\to\mathbb R$ be continuous, strictly concave and satisfy $f(-1)=f(1)=0$. For every such function, let Q_f be the body of revolution given by $y^2+z^2+t^2\le f(x)^2$. Then Q_f and Q_g have the same maximal sections in every direction if (and, actually, only if) f and g are equimeasurable, i.e., $|\{f>t\}|=|\{g>t\}|$ for all $t>0$. (Of course, there are plenty of concave functions equimeasurable with $\sqrt{1-x^2}$). With some extra work, one can construct something like this in $\mathbb R^n$ when n is even though I do not know any similarly nice geometric description of such bodies for $n\ge 6$. What I (and my co-authors) are currently stuck with is the case of odd n (say, the usual space $n=3$). In view of such simple example in $\mathbb R^4$, I suspect that we are just having a mental block. Can anybody help us out?

论坛问题帖 | has_answers | fedja | 2011-07-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018469 | MO-71949

Totally rational polytopes

Define a convex polytope in $\mathbb{R}^d$ as totally rational (my terminology) if its vertex coordinates are rational, its edge lengths are rational, its two-dimensional face areas are rational, etc., and finally its (positive) volume is rational. So: rational coordinates, and the measure of every k-dimensional face, $1 \le k \le d{-}1$, is rational, and the d-dimensional volume is positive and rational. (Scaling could then convert all these rationals to integers.) For example, the hypercube with vertex coordinates $\{0,1\}^d$ is totally rational. Similarly an axis-aligned box with integral vertex coordinates is totally rational. Q1. Are there other classes of totally rational polytopes, classes defined for all d? In particular, Q2. Do there exist totally rational simplices in $\mathbb{R}^d$ for arbitrarily large d? Pythagorean triples yield totally rational triangles. I am not even certain that the Heronian tetrahedra described in this MathWorld article are totally rational, because it is unclear (to me) if they can be realized with rational vertex coordinates. All this is likely known, in which case key search phrases or other pointers would be welcomed. Thanks! Addendum. Gerry Myerson's useful summary of Problem D22 in Unsolved Problems In Number Theory answers Q2: The problem is open! Q1 remains (apparently) interesting; see the comments by Steve Huntsman and Gerhard Paseman.

论坛问题帖 | has_answers | Joseph O'Rourke | 2011-08-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018470 | MO-72189

Once differentiable, piecewise degree three polynomials on triangulated planar domains

Here is an easily described, but very difficult, problem that I (and a number of other people) really would like to see solved during our life times. The basic problem is to compute the dimension of a certain vector space. Suppose we are given a triangulation (a tessellation by triangles where any two triangles share at most a common edge or a common vertex) of a polygonal domain in the plane. Let S be the space of once differentiable functions on that triangulation that on each triangle can be represented as a bivariate polynomial of degree 3. S is clearly a vector space. It is known that the dimension of that space is greater than or equal to $3V_B + 2V_I + 1 + \sigma$ where V_B is the number of boundary vertices, V_I is the number of interior vertices, and σ is the number of singular vertices of the underlying triangulation. A singular vertex is an interior vertex that has exactly four edges attached where those edges form two parallel pairs. (In other words, a singular vertex is the intersection of the diagonals of a convex quadrilateral.) It is known that generically the dimension of S equals the given expression, and there is no case known where the dimension is larger than that expression. Many people conjecture that the lower bound equals the dimension for all triangulations. Prove, or disprove, that conjecture. Some background: triangulations are the natural generalization of a partition of an interval to two variables, and S is a spline space with potential for a wide range of practical problems, such as data fitting or solving partial differential equations. The problem has been known among approximation theorists since the early 1970s, and despite efforts by a number of people the problem is still unsolved. For more information on spaces like S see the recent book by Lai and Schumaker [Spline Functions on Triangulations, Cambridge University Press] in particular section 9.9. The basic issue with the kind of spline space considered here is that the dimension depends not just on the topology of the triangulation, but also on its geometry. An arbitrarily small change of the location of the vertices can change the dimension. If the polynomial degree is four (instead of three) it is known that the dimension can change only when a vertex switches between being singular and non-singular. On the other hand, if the polynomial degree is two (instead of three) many configurations other than singular vertices are known where the dimension changes with the geometry. So the polynomial degree three straddles the boundary, and we would like to know to which camp it belongs.

论坛问题帖 | no_answers_at_retrieval | Maritza Sirvent | 2011-08-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018471 | MO-72206

Open problems in "Algebraic geometry by robin hartshorne"

Hi, Is there a list of which of the open problems in algebraic geometry by robin hartshorne are still open ? (I searched the internet and didn't find one) Thanks from advanced, Matan Fattal

论坛问题帖 | no_answers_at_retrieval | Matan Fattal | 2011-08-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018472 | MO-73001

Is there a smooth 4-manifold homeomorphic but not diffemorphic to CP^2?

Is there a smooth 4-manifold homeomorphic but not diffemorphic to CP^2? Are there known non-smooth examples homeomorphic CP^2?

论坛问题帖 | has_answers | Yang | 2011-08-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018473 | MO-7304

Complex vector bundles that are not holomorphic

Is there an example of a complex bundle on $\mathbb CP^n$ or on a Fano variety (defined over complex numbers), that does not admit a holomorphic structure? We require that the Chern classes of the bundle are (k,k) Hodge classes (which is automatic for $\mathbb CP^n$ or Fanos of dimension<4). If by any chance such examples are known, what is the smallest dimension of the variety (or the bundle)? For $\mathbb CP^1$ it is elementary to see that all bundles are holomorphic. In the book of Okonek and Schneider it is stated, that all complex bundles on $\mathbb CP^2$ and $\mathbb CP^3$ are also holomorphic. But for $\mathbb CP^n$, $n\ge 4$ this is stated as an open problem (as for 1980).

论坛问题帖 | has_answers | Dmitri Panov | 2009-11-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018474 | MO-73428

When is a compact topological 4-manifold a CW complex?

Freedman's E_8-manifold is nontriangulable, as proved on page (xvi) of the Akbulut-McCarthy 1990 Princeton Mathematical Notes "Casson's invariant for oriented homology 3-spheres". Kirby showed that a compact 4-manifold has a handlebody structure if and only if it is smoothable: 1 and 2. When is a compact topological 4-manifold a CW complex?

论坛问题帖 | no_answers_at_retrieval | Andrew Ranicki | 2011-08-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018475 | MO-74191

What is the degree of a symmetric boolean function?

(previous title " Zero sum of binomials coefficients - a stronger version ") This is a stronger version of another question. Is there an $N\in \mathbb N$ and a sequence of non-constant functions $ \left\{ p_n:[n] \to \{ 1,-1 \} \right\}_{n=N} ^{\infty}$ such that for all $n>N$ we have: $$ \sum _{i=0} ^{n} (-1)^{i} p_n(i) \cdot \binom {n} {i} = 0$$ For instance, for all odd values of n, we may choose $p_{n}(i)=\begin{cases} (-1)^{i} & i\leq\frac{n-1}{2}\\\ (-1)^{i+1} & i\geq\frac{n+1}{2}\end{cases}$. This simply means we sum the first half of the binomial coefficients and subtract the second half. The fact that for odd values we can partition the set of binomial coefficients evenly allows us to do that, so I don't see how the same trick may be applied for even values. To my understanding, the methods that solved the previous question (for which I thank darij grinberg and Mikael de la Salle) are not applicable here. My guess, as before, is that there is no such sequence (in which all functions are non-constant), any ideas on how to prove it? (a counter example would surprise me, but is of interest as well)

论坛问题帖 | has_answers | Shir | 2011-08-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018476 | MO-74544

The quaternion moat problem

"One cannot walk to infinity on the real line if one uses steps of bounded length and steps on the prime numbers. This is simply a restatement of the classic result that there are arbitrarily large gaps in the primes." So begins the paper by Gethner, Wagon, and Wick, "A Stroll Through the Gaussian Primes" (American Mathematical Monthly 105(4): 327-337 (1998).) They explain that it is unknown if one can walk to infinity on the Gaussian primes with steps of bounded length. Paul Erdős was reported to have conjectured this is possible ("A conjecture of Paul Erdős concerning Gaussian primes." Math. Comp 24: 221-223 (1970); PDF). Later Erdős is reported to have conjectured the opposite: that no such walk-to-∞ is possible [GWW98, p.327]. This has become known as the Gaussian moat problem, apparently still unresolved. My question is: Is there an analogous quaternion moat problem? Is it solved? open? Is it easier or harder than the Gaussian moat problem? Define a nonzero quaternion $q = a + bi + cj + dk$ as prime iff (a) it is a Hurwitz quaternion (all components integer, or all components half-integer) and (b) its norm $a^2 + b^2 +c^2 + d^2$ is prime. (Part (b) is a consequence of the inability to factor q; see, e.g., Theorem 15 in *"A proof of Lagrange's four square theorem using quaternion algebras." Drew Stokesbary, 2007; PDF, DOI: 10.2139/ssrn.2396123). Can one "walk-to-∞" on the quaternion primes using steps of bounded length? Perhaps relevant here is Lagrange's four-square theorem, which states that any natural number can be represented as the sum of four squares. I ask this question in relative naïveté, and appreciate being enlightened.

论坛问题帖 | has_answers | Joseph O'Rourke | 2011-09-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018477 | MO-76280

Exact consistency-strength of "all projective sets are Ramsey"

I wonder if the exact consistency strength of "All projective sets have the Ramsey property" is still open. In Solovay's model, all sets have the Ramsey property, so the consistency strength of this is below an inaccessible. As far as I know, there are some implications under forcing axioms. And all sets up to a certain level of the projective hierarchy having the Ramsey property is still do-able from just ZFC. But I can't seem to find anything which shows Solovays inaccessible is really necessary in case of the Ramsey property (like Raisonnier/Shelahs result for Lebesgue measure), or oppositely, that you can get a model where "All projective sets have the Ramsey property" from just ZFC (like for the Baire property).

论坛问题帖 | has_answers | David Schrittesser | 2011-09-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018478 | MO-77434

Convergence speed of Jacobi eigenvalue algorithm for parallel ordering(Brent-Luk) ?

Is there estimate for convergence of the Jacobi eigenvalue algorithm for Hermitian matrices for "parallel ordring" (Brent-Luk ordering (see comment below)) ? For example for 4 4 matrices parallel ordering is the following 1a) 12 1b) 34 2a) 23 2b) 14 3a) 13 3b) 24 [EDIT] Moreover convergence itself is not known for such ordering even in 4x4 case. I have consulted with many experts in the field - it is not proved. Numerical simulations (checked more than 10^10 matrices of different form) shows that the convergence exists. There is certain subtlety in the definition of method. Which lead some authors to claim that there is NO convergence. But actually counter-example is not for "reasonable" implementation of details. The detail is the following consider 2x2 matrix such that diagonal elements are equal. Then the rotation can be either +45 either -45 - no unique choice. What the authors claim that if we have a freedom to choose +45 or -45 by our own wish, in each step where ambiguity occurs - then there will be counterexample ! However this counter-example does NOT work if we fix +45 (or -45) once and forever ! I.e. in the case of ambiguity we ALWAYS choose angle to be the same. Simulations shows - that than there is no problem. I spent about 2 weeks trying to prove this just in the 4x4 example - but I was unable to prove it. The difficulty is that we need to analyse about 3-4 sweeps. It can be shown that there always exists a matrix that can be arbitrary "BAD" after 1-2 sweeps... [END of EDIT on 21 Jan. 2012] As far as I can expect that there should be ultimate quadratic convergence [EDIT] actually as works of Walter F. Mascarenhas suggests their will be cubic ultimate convergence[EDIT] but I

am interested at the first iteration - they should be at most linear convergence, but it is not clear for is there uniform convergence speed or there can be some matrices where convergence can be arbitrary bad ? (From simulation we see that probably there is NO bad examples - convergence seems rather fast, but there are certain difficulties in proving this theoretically). Actually even the convergence for arbirary ordering is not clear for me. Paper by Walter Mascarenhas: SIAM. J. Matrix Anal. & Appl. 16, pp. 1197-1209 (13 pages) On the Convergence of the Jacobi Method for Arbitrary Orderings Walter F. Mascarenhas States only convergence of the diagonal elements. Non-diagonal elements may not converge, for some sophisticated orderings. He constructed examples in his PhD at MIT unpublished (private communication from him)

论坛问题帖 | has_answers | Alexander Chervov | 2011-10-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018479 | MO-77446

What is known about Ulam's problem of metric spaces with isometric squares?

Background In the book Problems in Modern Mathematics, S. Ulam asks the following question: Suppose A and B are metric spaces, such that A^2 and B^2 equipped with the 2-metric $d((x_1, y_1),(x_2, y_2)) = \sqrt{d(x_1, x_2)^2 + d(y_1, y_2)^2}$ are isometric. Does it follow that A and B must also be isometric themselves? The answer to this general question is negative. Take for example the space of the rational numbers $\mathbb{Q}$ and the space $\mathbb{Q} \sqrt{2}$ of rational multiples of $\sqrt{2}$. These two spaces are obviously not isometric, but their squares are, isometry being just the rotation by $\frac{\pi}{4}$. See http://www.ams.org/journals/proc/1971-029-03/S0002-9939-1971-0278262-5/S0002-9939-1971-0278262-5.pdf for the proof. Question What is the status of this problem for slightly less general metric spaces? I am especially interested in the following cases: What is known about this problem for complete metric spaces? What is known for finite metric spaces? A good survey if it exists would be more than welcome.

论坛问题帖 | has_answers | Dejan Govc | 2011-10-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018480 | MO-78026

Compact surfaces smoothly immerse in: $\mathbb{R}^4$ or $\mathbb{R}^5$?

I wonder if someone can clarify whether it is known that every closed, orientable surface (2-manifold) has a smooth isometric immersion in $\mathbb{R}^4$? This topic has been discussed rather thoroughly in the earlier MO question, "Nash embedding theorem for 2D manifolds." However, I am puzzled by two claims that seem at odds, but are surely not-rather I just don't understand the terminology. First, Gromov established that every compact surface has a smooth embedding in $\mathbb{R}^5$. This is from his 1986 book, Partial Differential Relations. See the above MO question for a quote. Second, as user jc said, "Poznyak proved that $\mathbb{R}^4$ works for any compact part of a complete surface." Here is a more precise statement of Poznyak's result, from the book by Jia-Xing Hong, Isometric embedding of Riemannian manifolds in Euclidean spaces (p.40): Theorem 2.3.1. Let g be a smooth complete metric defined in $\mathbb{R}^2$. Then for any compact set Ω in $\mathbb{R}^2$, (Ω,g) admits a smooth isometric immersion in $\mathbb{R}^4$. Since Poznyak's $\mathbb{R}^4$ result is from 1973, and Gromov's $\mathbb{R}^5$ from 1986, it must be that Poznyak's theorem does not improve Gromov's to $\mathbb{R}^4$. Is this because "a compact part of a complete surface" is not the same as a "compact surface"? Can anyone explain this to me? Thanks! E.G. Poznyak, "Isometric immersions of two-dimensional Riemannian metrics in Euclidean space." Russian Math. Surveys, 28:4 (1973), pp. 47-77. Uspekhi Mat. Nauk, 28:4 (1973), pp. 47-76. Answered. Robert Bryant resolved my confusion: "A compact surface without boundary can't be identified with a subset of the plane." It is my understanding now that it remains an open question of whether or not every compact surface (without boundary) has a smooth immersion in $\mathbb{R}^4$: neither Gromov's nor Poznyak's results settle this question, and as far as I know there is no known counterexample. I'm adding the open-problem tag, because although Robert completely answered my specific question, the question suggested by the title is open.

论坛问题帖 | no_answers_at_retrieval | Joseph O'Rourke | 2011-10-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018481 | MO-78838

What is the maximal density vertex subset of the 8-connected grid with induced vertex degrees ≤ 4?

Let G be the infinite graph defined by 8-connection of $\mathbb{Z}^2$. What is the maximal density vertex subset whose induced subgraph has maximum degree ≤ 4? More precisely, what is the maximal density $\delta(4)$ as defined in this paper which gives the bounds $\frac{3}{5} \leq \delta(4) \lt \frac{5}{8}$? This question could be known to be open, and it has been asked on math stackexchange. To make this question less dry I am adding an example of a periodic tiling which meets the $\frac{3}{5}$ lower bound, and additional tilings with this density are given in the paper linked above. I do not know if a pattern has been found with a higher density subject to the degree constraint, or whether the strictness of the $\frac{3}{5}$ bound has been proved. . . X X X . . X X X X X . . X X X . . X . X X X . . X X X . X . . X X X . . X X X X X . . X X X . . . . X X X . . X X X X X . . X X X . . X . X X X . . X X X . X . . X X X . . X X X X X . . X X X . .

论坛问题帖 | no_answers_at_retrieval | nom-de-mo | 2011-10-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018482 | MO-7969

Irreducible polynomials with constrained coefficients

Over at the Cafe, after reading about TWF 285, I asked more-or-less About how many polynomials with coefficients in $\{\pm 1\}$ and of degree d are irreducible? and that's what I want to ask here. The first no-go analysis: since if P is reducible, then it is reducible mod 3, we get that the number of reducible such is $O(\frac{d-1}{d}3^d)$; but that's clearly too large already to help much; reducing mod 2 we can't distinguish polynomials anymore!

论坛问题帖 | has_answers | some guy on the street | 2009-12-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018483 | MO-80101

3-piece dissection of square to equilateral triangle?

At a workshop it was suggested that it likely remains an open problem whether or not there is a 3- or 2 -piece dissection of a square to an equilateral triangle. Can anyone confirm that this is unresolved? Four-piece dissections are known, the most famous being Henry Dudeney's century-old gem: [Maple animation from this link.] Update (6Dec2024). Now proved impossible. "Dudeney's Dissection is Optimal." Erik D. Demaine, Tonan Kamata, Ryuhei Uehara. arXiv abs.

论坛问题帖 | has_answers | Joseph O'Rourke | 2011-11-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018484 | MO-83004

Algorithmic decidability of equality in the ring of periods

Suppose two elements of the ring of periods are given by their systems of polynomial inequalities with rational coefficients. Is there a known algorithm deciding their equality? Is it known if their equality is algorithmically decidable at all? It is quite obvious that if they are not equal, then it can be shown after finite number of steps just by evaluating them with big enough precision. The difficult part is when they are equal.

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2011-12-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018485 | MO-83262

Square and cube?

Gowers and Maurey proved in their remarkable paper(s), that there is a Banach space X such that X is isomorphic to its cube $X\oplus X\oplus X$ but not to isomorphic to its square $X\oplus X$. This space seems to be not a dual space, am I correct? Is there a solution to this problem which is a dual space? And the second quick question: what is the status of the following conjecture: There is a Banach space X such that X is not isomorphic to X^2 but X^2 is isomorphic to X^3? Thank you very much. S.

论坛问题帖 | no_answers_at_retrieval | Sellapan Nathan | 2011-12-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018486 | MO-83550

Has n^2*|sin(n)| limit?

Is there any sub-sequence n_k of natural numbers such as $\lim(n_k^2|\sin(n_k)|) = 0$? when k tends to infinity. In other words, does $\lim(n^2|\sin(n)|)$ exist (equal to infinity)? or there is no such limit?

论坛问题帖 | no_answers_at_retrieval | John | 2011-12-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018487 | MO-8534

Diameter of universal cover

Let M be Riemannian manifold and $\tilde M$ be its universal cover (with induced metric). What is the upper bound for $k=\mathop{diam}\tilde M/\mathop{diam} M$ in terms of $m=|\pi_1(M)|$ (or $\pi_1(M)$)? Comments: There is a similar answered question here, but there cover is NOT universal. So we get $k\leqslant m$, but $k\ll m$ is expected. The question is open even in case $\pi_1=\mathbb Z_m$ (even asymptotics is not known). Clearly, $\sup k$ for given finite group $\Gamma=\pi_1(M)$ is an invariant of Γ. Is it an interesting invariant? Examples: For $\pi_1=\mathbb Z_{3\cdot 2^n}$ one can make $k\sim n$ or $k=O(\log m)$ (see my answer below). For $\pi_1=S_n$, one can make k of order n^2 or (see Greg's answer and also this question). It is much more than $\log m$, but still $k=o(\log^2 m)$.

论坛问题帖 | has_answers | Anton Petrunin | 2009-12-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018488 | MO-85724

Given a lattice L with n elements, are there finite groups H < G such that L $\cong$ the lattice of subgroups between H and G?

If there is no restriction on n, this is a famous open problem. I'm wondering if any recent work has been done for small $n>6$. I believe the question is answered (positively) for $n=6$ by Watatani (1996) MR1409040 and Aschbacher (2008) MR2393428. I also believe we can answer it for $n=7$, with one possible exception. The exceptional case is shown below. So my two questions are these: Does anyone know of recent work on this special case of the problem (specifically for $n=7$ or $n=8$)? Has anyone found a finite group G with a subgroup H such that the interval $[H, G] = \{K : H \leq K \leq G \}$ is the lattice shown above? If not, I'd be happy to hear if anyone has ideas about how we might prove that such a group exists (without necessarily producing the group). Or, if you can put restrictions on such a group, that might be useful. For example, I believe I can show that there is no finite solvable group with this property. (By the way, I have searched many of the groups in GAP's small groups library, and this was helpful for finding some of the other difficult 7-element cases.) Correction: We don't really have an answer to this question for $n=7$. Rather, we can show that every lattice with at most 7 elements (except possibly the one above) is the congruence lattice of a finite algebra. Pálfy and Pudlák MR593011 showed the two questions are equivalent when n is not restricted, but their proof doesn't go through for fixed n. Nonetheless, we can prove that a (minimal) finite algebra with the lattice above as congruence lattice must be a transitive G-set (modulo constant operations), so the existence of such a finite algebra is equivalent to the existence of finite groups H < G with [H,G] isomorphic to the lattice above. (This is the reason I phrased my question in terms of subgroup lattices.) Update: (2012/02/22) If the interval $[H,G]$ is the lattice above, with H core-free, I believe I can prove that a minimal normal subgroup of G must be non-abelian, which isn't much, but it's a start! In particular, I think it rules out the affine case of the Aschbacher-O'Nan-Scott Theorem.

论坛问题帖 | no_answers_at_retrieval | William DeMeo | 2012-01-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018489 | MO-8729

Are the C(S^n, S^n)'s homeomorphic ?

Let m, n > 1. Is it true that C(S^m, S^m), and C(S^n, S^n) are homeomorphic ? [both endowed with the sup metric (or equivalently the compact-open topology)] Generally, C(S^n, S^n), with n >= 1, is a countably infinite (disjoint) union of path-connected (due to Hopf) components C_{n,k}, k in Z. I think each of these components may be viewed as an infinite dimensional Frechet manifold. Unfortunately, they are not contractible. However, the question is [somewhat vaguely] motivated by the Henderson's Theorem. Also, I have some related questions : - Fixing n, are the C_{n,k}'s homeomorphic to each other (at least for k <> 0) ? - Are there some m, n, k, l with m <> n s.t. C_{m,k} ~ C_{n,l} ? What I was able to do in this direction until now is to show the existence of a proper, one-to-one, degree-preserving map from C(S^m, S^m) into C(S^n, S^n). Even for m >> n, but far to be surjective.

论坛问题帖 | has_answers | Ady | 2009-12-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018490 | MO-87342

Are acyclic subcomplexes of finite contractible 2-complexes contractible?

Let Y be a contractible finite simplicial 2-complex. Let X be an acyclic subcomplex of Y (i.e. X connected, $H_1(X)=0$, $H_2(X)=0$). Is X contractible? (Equivalently, is $\pi_1(X)$ trivial?) I asked this question among others here before, but it remained unanswered. Apparently this question is equivalent to the following one. If $m < n$, $G = \langle a_1,\dotsc,a_m \mid r_1 = \dotsb = r_m =1 \rangle$ and $J=\langle a_1,\dotsc,a_n \mid r_1 = \dotsb = r_n =1 \rangle$ are two groups given by (balanced) presentations, the presentation of G is a sub-presentation of the presentation of J, and J is trivial, is G trivial? (It can be shown that the abelianization of G is trivial.)

论坛问题帖 | has_answers | Alexey Muranov | 2012-02-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018491 | MO-87437

Die-rolling Hamiltonian cycles

Let R be a rectangular region of the integer lattice $\mathbb{Z}^2$, each of whose unit squares is labeled with a number in $\lbrace 1, 2, 3, 4, 5, 6 \rbrace$. Say that such a labeled R is die-rolling Hamiltonian, or simply rollable, if there is a Hamiltonian cycle obtained by rolling a unit die cube over its edges so that, for each square $s \in R$, the cube lands on s precisely once, and when it does so, the top face of the cube matches the number in s. For example, the 4×4 "board" shown below is rollable. Q. Is it true that, if R is die-rolling Hamiltonian, then the Hamiltonian cycle is unique, i.e., there are never two distinct die-rolling Hamiltonian cycles on R? This "unique-rollability" question arose out of a problem I posed in 2005, and was largely solved two years later, in a paper entitled, "On rolling cube puzzles" (complete citation below; the 4×4 example above is from Fig. 17 of that paper). Although the original question involved computational complexity, the possible uniqueness of Hamiltonian cycles is independent of those computational issues, so I thought it might be useful to expose it to a different community, who might bring different tools to bear. It is known to hold for R with side lengths at most 8. If not every cell of R is labeled, and unlabeled cells are forbidden to the die, then there are examples with more than one Hamiltonian cycle. Edit1. Rolling a regular tetrahedron on the equilateral triangular (hexagonal) lattice is not as interesting. See the Trigg article cited below. Edit2. Serendipitously, gordon-royle posted a perhaps(?) relevantly related question: "Uniquely Hamiltonian graphs with minimum degree 4." The computational version is Open Problem 68 at The Open Problems Project. "On rolling cube puzzles." Buchin, Buchin, Demaine, Demaine, El-Khechen, Fekete, Knauer, Schulz, Taslakian. Proceedings of the 19th Canadian Conference on Computational Geometry, Pages 141-144, 2007. PDF download. Charles W. Trigg. "Tetrahedron rolled onto a plane." J. Recreational Mathematics, 3(2):82-87, 1970.

论坛问题帖 | has_answers | Joseph O'Rourke | 2012-02-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018492 | MO-87860

Example of a compact Kähler manifold with non-finitely generated canonical ring?

A celebrated recent theorem of Birkar-Cascini-Hacon-McKernan and Siu says that the canonical ring $R(X)=\oplus_{m\geq 0}H^0(X,mK_X)$ of any smooth algebraic variety X over $\mathbb{C}$ is a finitely generated $\mathbb{C}$-algebra. On the other hand P.M.H. Wilson (using a construction of Zariski) gave an example of a compact complex manifold X with $R(X)$ not finitely generated. However his manifold X is not Kähler. Does anyone know an example of a compact Kähler manifold X with $R(X)$ not finitely generated? Or is this an open problem?

论坛问题帖 | has_answers | YangMills | 2012-02-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018493 | MO-88087

Reference Request - Jakob Weisblat's "The Search for the Odd Perfect Number"

Hi All! I am currently trying to locate an online copy of Jakob Weisblat's paper titled "The Search for the Odd Perfect Number". I could only get hold of the abstract: "A perfect number is a number whose proper divisors add up to the number itself. All known perfect numbers are of the form $(2^{p}-1)(2^{p-1})$ where p is prime and $2^{p}-1$ is prime. This talk will consider the possibility of an odd perfect number. After considering criteria for the prime factorization of an odd perfect number, many categories of odds will be eliminated. The remaining categories will then be more thoroughly considered. The first part of the talk will be my personal research and investigations without having looked at anyone's previous work. Next, interesting previous studies on this matter will be discussed. Finally, my personal work will be compared to the previous analyses, and my research will be extended using this new information." If anybody out there has a copy or has a web-link to an online copy, would you please be willing to share it? Thanks! Edit: After taking to Ivars Peterson of MAA, it appears that the author was not required to make a paper submission for his talk. Likewise, the author does not have a publicly available e-mail address. Hence, reflecting quid's suggestion, is a write-up of the author's notes for this talk available somewhere? The author is currently unreachable through Facebook messaging.

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2012-02-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018494 | MO-8970

Number of valid topologies on a finite set of n elements

I've heard that the problem of counting topologies is hard, but I couldn't really find anything about it on the rest of the internet. Has this problem been solved? If not, is there some feature that makes it pretty much intractable?

论坛问题帖 | has_answers | Harry Gindi | 2009-12-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018495 | MO-89798

Is S^2 x S^4 a complex manifold?

As observed by Calabi a long time ago, the manifold $S^2\times S^4$ admits an almost-complex structure (obtained by embedding it in $\mathbb{R}^7$ and using the octonionic product), which however is not integrable. Is it known whether $S^2\times S^4$ admits an integrable complex structure? A few remarks. This is stated as an open problem in Calabi's paper, but perhaps it has been solved in the meantime? This is similar to the case of S^6, which is still open (see this question). One can also ask the same question for $\Sigma\times S^4$ for Σ any compact Riemann surface It seems that some people believe that every almost-complex manifold of real dimension 6 or more admits an integrable complex structure (see this other question). Even more generally (and this is obviously still open), one can ask about an arbitrary finite product of even dimensional spheres (excluding S^0). It is known that this is almost complex iff the only factors that appear are S^2, S^6 and $S^2\times S^4$. If one allows connected sums, then for example $(S^2\times S^4)\#2(S^3\times S^3)$ is a complex manifold, and in fact it has complex structures with trivial canonical bundles (see for example here and here).

论坛问题帖 | has_answers | YangMills | 2012-02-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018496 | MO-90277

When is the norm of all positive operators on an ordered Banach space determined by their values on the positive cone?

I'm trying to investigate the interplay between the norm and cone of positive elements in ordered Banach spaces. In particular, I would like a nice characterization of when the norm of a positive operator can be obtained as a supremum of norms over all norm 1 positive elements (see 1. below). It is easy to show that this always holds for Banach lattices, but gets tricky when trying to prove this more generally. To be a bit more detailed, let X be a Banach space with a cone (or wedge) of 'positive' elements, denoted by X_{+}; X is then called an ordered Banach space and $x\leq y$ means $y-x\in X_{+}$. I will call X_{+} generating if $X=X_{+}-X_{-}$; proper if $X_{+}\cap(-X_{+})=\{0\}$; and normal if $0\leq x\leq y$ implies $\|x\|\leq\|y\|$. A bounded linear operator $T:X\to X$ is called positive, if $x\geq 0$ implies $Tx\geq 0$. An ordered Banach space with the property that every positive operator satisfies $$\|T\|=\sup\{\|Tx\|:x\in X_{+},\|x\|=1\}$$ I will say has the positive operator property, i.e., the norms of positive operators are completely determined by their behavior on the cone. My question is: Let X be an ordered Banach space with a closed, proper, generating, normal cone. Is there a characterization of the positive operator property in terms of the cone-norm interaction? Thus far I have found a few sufficient conditions: For all $x\in X$ there exist $X\ni x_{1},x_{2}\geq0$ such that $x=x_{1}-x_{2}$ and $\|x_{j}\|\leq\|x\|$ for $j=1,2$. For all $x\in X$ and $X\ni a,b\geq0$, $-a\leq x\leq b$ implies $\|x\|\leq\max \{ \|a\|,\|b\| \}$. together imply the positive operator property. Another sufficient condition is having the property that for any $x\in X$ it holds that $\|x\| = \inf \{\|z_+ + z_-\|:z_\pm \geq 0;x=z_+ -z_-\}$. Being a Banach lattice is sufficient to have this property (by invoking the property $\|x\|=\||x|\|$), but seems to be a bit more general since $\mathbb{R}^3$ with the Euclidean norm and the `ice-cream cone' $\{(x_1,x_2,x_3):x_1\geq (x_2^2+x_3^2)^{1/2} \}$ (which is not a lattice) also has this property (so being a lattice is not necessary for the positive operator property). I've been unable to prove that any of these conditions are necessary. The last one had been my best bet so far, but I'm beginning to doubt the existence of a nice list of conditions on the norm and cone that together are necessary and sufficient. Any suggestions, counterexamples or pointers to the literature that anyone may have will be greatly appreciated. UPDATE: Cleaned it up a bit, and posted some new information I found since first posting. I found no mention in any literature to this question. I also asked around a bit with no hits from people who are a bit in the know, so it seems to be wide open. Gets the 'open-problem' tag.

论坛问题帖 | has_answers | Miek Messerschmidt | 2012-03-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018497 | MO-90977

Interesting result on the Euler-Maschroni constant - what is the background?

Today I entered the following expression in maple: $$a_i = H_{10^i} - ln(10^i) - \gamma$$ Here H_j equals $\sum_{k=1}^{j} 1/k$ and γ is the Euler-Mascheroni constant. When I computed a_n for $i = 0$ to 10 I obtained the following results: $i = 0$;
4.22784335098467139393487909917597568957840664060076401194232765115132 3 * 10^{-1} $i=1$;
4.91674960726754236294647092014873296107074293995573934148731181158 13 * 10^{-2} $i=2$;
4.99166674999603216262267620712231166460981351098210230411091976720 6 * 10^{-3} $i=3;$
4.99916666674999996031750198405122676215367882561138867875812170113 3 * 10^{-4} $i=4$;
4.99991666666674999999960317460734126976551226762154503821179264423 * 10^{-5} $i=5$;
4.99999166666666674999999999603174603216269841262265512267621545 23 * 10^{-6} $i=6$;
4.99999916666666666674999999999996031746031750198412698405122652 3 * 10^{-7} $i=7$;
4.99999991666666666666674999999999999603174603174607341269841 23 *

10^{-8} $i=8$; 4.99999999166666666666666674999999999999999960317460317460321623 * 10^{-9} $i=9$; 4.99999999916666666666666666674999999999999999960317460317423 * 10^{-10} $i=10$; 4.999999999916666666666666666666749999999999999999960317423 * 10^{-11} So we see that the periodic strips of ...99999..., of ...66666... and ...99999... an many other periods increase for even larger i. The question is now: Is there any rule behind it that the remainder term a_i behaves that way?

论坛问题帖 | has_answers | tobias | 2012-03-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018498 | MO-93891

What is the status of the subadditivity problem for analytic capacity?

Hi, Here is another question that concerns analytic capacity. For a compact set K in the plane, define the analytic capacity of K by $$\gamma(K):=\sup|f'(\infty)|,$$ where the supremum is taken over all functions f holomorphic and bounded by 1 in the complement of K. Here $$f'(∞)=\lim_{z \rightarrow \infty}z(f(z)-f(\infty)).$$ It was conjectured by Vitushkin, in view to applications in approximation theory, that analytic capacity is semi-additive, i.e. there exists a universal constant C such that $$\gamma(E \cup F) \leq C(\gamma(E) + \gamma(F)).$$ This was proved by Tolsa in 2003. From what I know, wether or not analytic capacity is subadditive, i.e. if we can take $C=1$, is still open. However, I can't find much about this problem in the literature, hence the question : What is the status on this problem? Has it been proved that analytic capacity is indeed subadditive in some particular cases? The only thing I found about this is an article by Suita, "On subadditivity of analytic capacity for two continua.", in which it is proved that analytic capacity is subadditive in the case where E,F are disjoint connected compact sets. The proof relies deeply on the fact that for compact connected sets, analytic capacity equals logarithmic capacity, which does not hold in the disconnected case. Any reference about this is welcome. Thank you, Malik

论坛问题帖 | no_answers_at_retrieval | Malik Younsi | 2012-04-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018499 | MO-93997

Avoidable words

Let $u(x_1,...,x_n)$ be a word, $k\in \mathbb{N}$. We say that u is k-avoidable if there exists an infinite word in k letters $\{a_1,...,a_k\}$ which does not contain values of u (i.e. words of the form $u(v_1,...,v_n)$, $v_i\in \{a_1,...,a_k\}^+$ as a subword. For example x^3 is 2-avoidable and x^2 is 3-avoidable but not 2-avoidable because two infinite words constructed by Thue and Morse avoid x^2 and x^3. All avoidable words have been described by Bean-Ehrenfeucht-McNulty and Zimin (see these slides of George McNulty's talk, for example. Is there an example of a 6-avoidable but not 5-avoidable word? The word x^3 is 2-avoidable but not 1-avoidable. The word x^2 is 3-avoidable (Thue) but not 2-avoidable (obvious). Walter's word $$abwbcxcayba$$ is 4-avoidable but not 3-avoidable. Ronald Clark in his PhD thesis proved that $$abubawacxbcycdazdcd$$ is 5-avoidable but not 4-avoidable (see Clark, Ronald J. The existence of a pattern which is 5-avoidable but 4-unavoidable. Internat. J. Algebra Comput. 16 (2006), no. 2, 351-367.) It is of course hard to believe that every avoidable word is 5-avoidable (or C-avoidable for any constant C). There were rumors that an example of a 6-avoidable word which is not 5-avoidable has also been constructed. What is the status of that question? Update Since according to James Currie, the above is almost as much as we know, as far as the lower bound is concerned, here is the best upper bounds I know. Let k ibe the number of letters in an avoidable word u, $n=n(k)$ is the minimal n such that all such u are n-avoidable. An exponential upper bound for $n(k)$ was found in the original papers by Bean-Ehrenfeucht-McNulty and Zimin, quadratic upper bound - in my paper, "Problems of Burnside type and the finite basis property in varieties of semigroups". Math. USSR-Izv. 30 (1988), no. 2, 295-314 (I considered a more complicated problem of avoidability of identities) and a linear upper bound $9k+20$ in Baker, McNulty, Taylor, Growth problems for avoidable words. Theoret. Comput. Sci. 69 (1989), no. 3, 319-345. The best upper bound I know is $n(k)\le 3\lfloor k/2\rfloor+3$. It was proved by Irina Mel'nichuk in "On the existence of infinite finitely generated free semigroups in some varieties of semigroups", Leningrad Algebraic Systems with One Operation and Relations (1985), pp. 74-83. George McNulty claims in the talk I linked to above that Mel'nichuk proved the upper bound $k+6$ in 1988, but I could not find a reference. The question is whether $k(n)$ is bounded by a constant.

论坛问题帖 | no_answers_at_retrieval | user6976 | 2012-04-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018500 | MO-9558

The shortest path in first passage percolation

Update (January 17): The problem has now been solved by Daniel Ahlberg and Christopher Hoffman. (Thanks to Matt Kahle for informing us.) Consider a square planar grid. (The vertices are pair of points in the plane with integer coordinates and two vertices are adjacent if they agree in one coordinate and differ by one in the other.) Give every edge a length one with probability a half and length two with probability a half. Consider a shortest path between the origin and the vertex $(n,0)$. Show that with probability that tends to one as n tends to infinity the shortest path will not contain the "middle edge" on the x-axis inbetween the orgin and $(n,0)$. (Namely, the edge between the vertices $(\lfloor\frac{n}{2}\rfloor,0)$ and $(\lfloor\frac{n}{2}\rfloor+1,0)$.) This question is in the category of "a missing lemma". It is not really a full fledged open problem but rather a statement which looks correct that was needed in some paper and resisted proof. Of course, some such "lemmas" turn out to be very difficult, but sometimes maybe a simple argument was simply missed. The relevant paper is Itai Benjamini, Gil Kalai, and Oded Schramm, First Passage Percolation Has Sublinear Distance Variance, Ann. Probab. 31(4) (2003) pp 1970-1978, doi:10.1214/aop/1068646373, arXiv:math/0203262. While MO have chosen to accept one answer, and there were some nice suggestions, the problem is still wide open.

论坛问题帖 | has_answers | Gil Kalai | 2009-12-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018501 | MO-96202

Open problems/questions in representation theory and around?

What are open problems in representation theory? What are the sources (books/papers/sites) discussing this? Any kinds of problems/questions are welcome - big/small, vague/concrete. Some estimation of difficulty and importance, as well as, small description, prerequisites and relevant references, ... are welcome. To the best of my knowledge, there are NO good lists of representation theory problems on the web. E.g. the sites below contain lots of unsolved problem in other areas, but not in representation theory: http://en.wikipedia.org/wiki/Unsolved_problems_in_mathematics http://garden.irmacs.sfu.ca/ http://maven.smith.edu/~orourke/TOPP/ MO questions also discuss other fields, but not representation theory: What are the big problems in probability theory? What are some open problems in algebraic geometry? What are some open problems in toric varieties? More open problems Open problems with monetary rewards Open problems in Euclidean geometry? Open Questions in Riemannian Geometry What are some of the big open problems in 3-manifold theory? Open problems in continued fractions theory

论坛问题帖 | has_answers | Alexander Chervov | 2012-05-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018502 | MO-96347

Entropy of Bernoulli walks on semi-groups.

Consider the Fibonacci semi-group $<L,R|LRR=RLL>$ with a Bernoulli walk $P(R)=p, P(L)=1-p$. Is the entropy $H(p)$ an unimodal function with maximum at p=0.5? Is this true for all finitely generated semi-groups (with some symmetries)? For the free semi-group $<L,R>$ (the Shift space) it is well known and easy to prove that $H(p)=-(plog(p)+(1-p)log(1-p))$.

论坛问题帖 | no_answers_at_retrieval | Jörg Neunhäuserer | 2012-05-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018503 | MO-96782

vector balancing problem

I believe the solution posted to the arXiv on June 17 by Marcus, Spielman, and Srivastava is correct. This problem may be hard, so I don't expect an off-the-cuff solution. But can anyone suggest possible proof strategies? I have vectors $v_1, \ldots, v_k$ in ${\bf R}^n$. Each of them has euclidean length at most $.01$, and for every unit vector $u \in {\bf R}^n$ they satisfy $$\sum_{i=1}^k |\langle u,v_i\rangle|^2 = 1.$$ Is it possible to find a set of indices $S \subset \{1, \ldots, k\}$ such that $$.0001 < \sum_{i \in S} |\langle u,v_i\rangle|^2 < .9999$$ for every unit vector u? This will imply the same bounds when summing over the complement of S. The $.01$ and $.0001$ aren't important; I just need the result for some positive δ and ϵ. But they have to be independent of k and n. (This may seem unlikely, until you try to construct a counterexample.) The motivation is that this is a (very slightly simplified) equivalent version of the famous Kadison-Singer problem. A solution would have important consequences in operator theory, harmonic analysis, and C*-algebra. Many people have worked on this problem, but perhaps not in the above form, which I feel exposes the combinatorial difficulty which is the real root of the problem.

论坛问题帖 | has_answers | Nik Weaver | 2012-05-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018504 | MO-97086

Does smooth and proper over $\mathbb Z$ imply rational?

Does smooth and proper over $\mathbb Z$ imply rational? I think someone told me that this is a standard conjecture. Is it a widely held? held at all? Did someone in particular make this conjecture? write it down somewhere? give evidence for it? Would you like to provide evidence? Why rational and not, say, unirational? Fontaine's non-digitized letter to Messing shows that if X is smooth and proper over $\mathbb Z$, then the low degree nondiagonal Dolbeault groups vanish $H^q(X_{\mathbb Q}; \Omega^p)=0$ for $p\ne q$, $p+q\leqslant 3$. An obvious conjecture is to remove the degree restriction. Is this expected? Does this conjecture imply the rationality one? (This hypothesis on a single space does not force rationality: there are fake projective planes that mimic the cohomology of the real projective plane, but they have finite covers that would again be smooth and proper, but which can't have such cohomology by the Atiyah-Bott fixed point formula. a little more here) What I'm looking for: A citation of a survey would be ideal. Short of that, I would accept a citation of someone making this conjecture (which may well be the paper I cite above). In any event, I would be happy if people provide evidence for or against the conjecture. Some definitions: propriety is a relative version of compactness; let's just say that the scheme is cut out of projective space by homogeneous equations with coefficients in $\mathbb Z$. Smoothness means that the fibers, that is, the schemes defined by the same equations considered over finite fields (or their algebraic closures) are smooth. This is stronger than regularity of the total space and more like a submersion, so by analogy with Ehresmann's theorem, all the fibers are supposed to be "the same." For example, $\operatorname{Spec}\mathbb Z[i]$ is regular, but the map to $\operatorname{Spec}\mathbb Z$ is not smooth, but ramifies at (2). (Geometrically) rational means that that the field of rational functions with complex coefficients are the same as on $\mathbb CP^n$, namely $\mathbb C(x_1,\ldots,x_n)$. I think this question shows that strict rationality, a fraction field of $\mathbb Q(x_1,\ldots,x_n)$, is too much to ask for. To complete the list of relevant MO questions, would (geometric) rationality imply a Hasse principle, answering this question?

论坛问题帖 | has_answers | Ben Wieland | 2012-05-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018505 | MO-9731

Polynomial representing all nonnegative integers

Lagrange proved that every nonnegative integer is a sum of 4 squares. Gauss proved that every nonnegative integer is a sum of 3 triangular numbers. Is there a 2-variable polynomial $f(x,y) \in \mathbf{Q}[x,y]$ such that $f(\mathbf{Z} \times \mathbf{Z})=\mathbf{N}$?

论坛问题帖 | has_answers | Bjorn Poonen | 2009-12-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018506 | MO-99144

Local structure of rational varieties

I've been asked this question by a colleague who's not an algebraic geometer; we both feel that the answer should be "no", but I don't have a clue how to prove it. Here's the question: let X be a smooth rational variety (over the complex numbers, say). Is it true that every point of X has a Zariski open neighbourhood that is isomorphic to an open subset of ${\mathbb P}^n$?

论坛问题帖 | has_answers | rita | 2012-06-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018507 | MO-99352

Important open questions in the field of Tropical geometry

What are some of the important unanswered questions in the field of tropical geometry?

论坛问题帖 | has_answers | john | 2012-06-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018508 | MO-9957

distance regular metric spaces

A metric space (V,d) will be called distance regular if for every distances a>0, b, c a nonnegative integer p(a,b,c) is defined, so that whenever d(B,C)=a, there are precisely p(a,b,c) points A such that d(A,B)=c, d(A,C)=b. The Euclidean plane is an example: p(a,b,c)=0,1, or 2 when the triangle inequality for a,b,c, correspondingly, fails, turns into equality, or is strict. If we also require that p(a,b,c)>0 whenever the triangle inequality does not fail, then I conjecture that this is the only possibility for the parameters p(a,b,c). That is, there may be many non-isomorphic examples, but the parameters will be the same for all of them. (Thanks to Heather for this clarification.) Has anybody formulated/proved/refuted this conjecture before? It looks very natural. UPD. I should have mentioned this: "for every distances a>0, b, c" means all nonnegative reals, and the same for "whenever the triangle inequality does not fail". In particular this means that all positive real distances actually occur. UPD2. After a week trial, the question seems to be new, open, and interesting. Anton suugested a line of attack, and I believe I can write down the proof of the first step: that p(a,b,c)=1 when the triangle inequality turns into equality. Fedja produced examples showing that this first step is indeed essential. I'm adding the "open problem" tag to the question.

论坛问题帖 | has_answers | Dima Fon-Der-Flaass | 2009-12-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

C02 Mathematics Stack Exchange | 1,034 条

V3-018509 | MSE-1017636

Inequality with Euler's totient function

In A conjecture concerning primes and algebra on MSE, I defined a multiplicative function $\omega:\mathbb Z_+\!\!\to\mathbb Z_+$ with $\omega(p_n)=n$, for the n-th prime p_n. It was conjectured that $\omega(N)\leq\pi(N)$ for $N>49$, where π is the prime-counting function (which seems to be solved by a nice lemma by Greg Martin). Now I have found another conjecture concerning ω. I have tested $\varphi(N)\geq\omega(N)$ for all $N<1.000.000$, where φ is the Euler totient function, without finding a counterexample. The conjecture, for which I would like a proof or a counter proof, is then: $\displaystyle n\prod_{p|n}\left(1-\frac{1}{p}\right)\geq\prod_{k\geq 1}k^{n_k}$, where $\displaystyle n=\prod_{k\geq 1}p_k^{n_k}$. Notes: φ is multiplicative, $\varphi(mn)=\varphi(m)\varphi(n)$, for $(m,n)=1$ ω is completely multiplicative, $\omega(mn)=\omega(m)\omega(n)$ for all $m,n\in\mathbb Z_+$. The inequality $\varphi(N)\geq\pi(N)$ should also be true, for sufficiently large N, but I haven't found anything about this on internet. Jaycob Colemans comment seems to prove the inequality: $\displaystyle \prod_{p_k,n_k\ne0}\left(1-\frac{1}{p_k}\right)\geq\prod_{p_k,n_k\ne0}\frac{k^{n_k}}{p_k^{n_k}}$, since $\displaystyle \frac{p_k-1}{p_k}\geq\left(\frac{k}{p_k}\right)^{n_k}$.

论坛问题帖 | no_answers_at_retrieval | Lehs | 2014-11-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018510 | MSE-1039259

Conjecture on the value of limit and related primality testing

Just I made a curious conjecture when I was playing with my calculator. We will use $\displaystyle\prod_{i=1}^n p_i$ where p_i is the i-th prime. Then I have noted that, $$\left\lvert\cos \left(\displaystyle\prod_{i=1}^3 p_i\right)\right\rvert=\left\lvert\cos \left(\displaystyle\prod_{i=1}^4 p_i\right)\right\rvert=\left\lvert\cos \left(\displaystyle\prod_{i=1}^5 p_i\right)\right\rvert=\dfrac{\sqrt{3}}{2}$$ The typical thing is that if in case of the product of consecutive primes we take any composite number then the equality doesn't hold. For example $\left\lvert\cos (2\cdot 3 \cdot 5 \cdot 6)\right\rvert \neq \dfrac{\sqrt{3}}{2}$, $\left\lvert\cos (2\cdot 3 \cdot 5 \cdot 7 \cdot 8)\right\rvert \neq \dfrac{\sqrt{3}}{2}$. Now since I am calculating on a 12-digit calculator, I don't have plenty of evidence. Basing upon this very little evidence I am putting forward a conjecture. Conjecture $$\displaystyle\lim_{n\to \infty}\left\lvert\cos \left(\displaystyle\prod_{i=1}^n p_i\right)\right\rvert=\dfrac{\sqrt{3}}{2}$$ I think that both the conjectures are probably false but I can't prove or disprove anyone of them. Though counterexamples are welcome but in that case I will be glad to know if there are infinitely many counterexamples or not.

论坛问题帖 | has_answers | user170039 | 2014-11-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018511 | MSE-1042333

Conjecture: three or more decompositions into powers with a base differing by 1 means its a perfect power

If $$(i_1)^{a_1}(i_1+1)^{b_1}=n $$ $$(i_2)^{a_2}(i_2+1)^{b_2}=n $$ $$(i_3)^{a_3}(i_3+1)^{b_3}=n $$ where all the terms are positive integers and the groups $(i_1,a_1,b_1),(i_2,a_2,b_2),(i_3,a_3,b_3)$ are different (i.e. n is being decomposed in 3 different ways) then n is a perfect power, i.e. it is expressible in the form $j^c = n$, where $c>1$. I have tested this up to 10^{14} and not found a counterexample. Furthermore, I have never found an integer which can be expressed like this in four ways, i.e. with the addition of $(i_4)^{a_4}(i_4+1)^{b_4}=n$. Is it generally true, and if so can it be proved? Examples $$576=2^{6}.3^{2}=3^{2}.4^{3}=8^{2}.9^{1}$$
$$5184=2^{6}.3^{4}=3^{4}.4^{3}=8^{2}.9^{2}$$
$$36864=2^{12}.3^{2}=3^{2}.4^{6}=8^{4}.9^{1}$$
$$46656=2^{6}.3^{6}=3^{6}.4^{3}=8^{2}.9^{3}$$
$$331776=2^{12}.3^{4}=3^{4}.4^{6}=8^{4}.9^{2}$$
$$419904=2^{6}.3^{8}=3^{8}.4^{3}=8^{2}.9^{4}$$
$$810000=5^{4}.6^{4}=9^{2}.10^{4}=15^{4}.16^{1}$$
$$2359296=2^{18}.3^{2}=3^{2}.4^{9}=8^{6}.9^{1}$$
$$2985984=2^{12}.3^{6}=3^{6}.4^{6}=8^{4}.9^{3}$$
$$3779136=2^{6}.3^{10}=3^{10}.4^{3}=8^{2}.9^{5}$$
$$16003008=6^{6}.7^{3}=27^{2}.28^{3}=63^{3}.64^{1}$$
$$21233664=2^{18}.3^{4}=3^{4}.4^{9}=8^{6}.9^{2}$$
$$26873856=2^{12}.3^{8}=3^{8}.4^{6}=8^{4}.9^{4}$$
$$34012224=2^{6}.3^{12}=3^{12}.4^{3}=8^{2}.9^{6}$$
$$150994944=2^{24}.3^{2}=3^{2}.4^{12}=8^{8}.9^{1}$$
$$191102976=2^{18}.3^{6}=3^{6}.4^{9}=8^{6}.9^{3}$$
$$241864704=2^{12}.3^{10}=3^{10}.4^{6}=8^{4}.9^{5}$$
$$306110016=2^{6}.3^{14}=3^{14}.4^{3}=8^{2}.9^{7}$$
$$1358954496=2^{24}.3^{4}=3^{4}.4^{12}=8^{8}.9^{2}$$
$$1719926784=2^{18}.3^{8}=3^{8}.4^{9}=8^{6}.9^{4}$$
$$2176782336=2^{12}.3^{12}=3^{12}.4^{6}=8^{4}.9^{6}$$
$$2754990144=2^{6}.3^{16}=3^{16}.4^{3}=8^{2}.9^{8}$$
$$9663676416=2^{30}.3^{2}=3^{2}.4^{15}=8^{10}.9^{1}$$
$$12230590464=2^{24}.3^{6}=3^{6}.4^{12}=8^{8}.9^{3}$$
$$15479341056=2^{18}.3^{10}=3^{10}.4^{9}=8^{6}.9^{5}$$
$$19591041024=2^{12}.3^{14}=3^{14}.4^{6}=8^{4}.9^{7}$$
$$24794911296=2^{6}.3^{18}=3^{18}.4^{3}=8^{2}.9^{9}$$
$$86973087744=2^{30}.3^{4}=3^{4}.4^{15}=8^{10}.9^{2}$$
$$110075314176=2^{24}.3^{8}=3^{8}.4^{12}=8^{8}.9^{4}$$
$$139314069504=2^{18}.3^{12}=3^{12}.4^{9}=8^{6}.9^{6}$$
$$176319369216=2^{12}.3^{16}=3^{16}.4^{6}=8^{4}.9^{8}$$
$$223154201664=2^{6}.3^{20}=3^{20}.4^{3}=8^{2}.9^{10}$$

$$618475290624=2^{36}.3^{2}=3^{2}.4^{18}=8^{12}.9^{1}$$
$$656100000000=5^{8}.6^{8}=9^{4}.10^{8}=15^{8}.16^{2}$$
$$782757789696=2^{30}.3^{6}=3^{6}.4^{15}=8^{10}.9^{3}$$
$$990677827584=2^{24}.3^{10}=3^{10}.4^{12}=8^{8}.9^{5}$$
$$1253826625536=2^{18}.3^{14}=3^{14}.4^{9}=8^{6}.9^{7}$$
$$1586874322944=2^{12}.3^{18}=3^{18}.4^{6}=8^{4}.9^{9}$$
$$2008387814976=2^{6}.3^{22}=3^{22}.4^{3}=8^{2}.9^{11}$$
$$5566277615616=2^{36}.3^{4}=3^{4}.4^{18}=8^{12}.9^{2}$$
$$7044820107264=2^{30}.3^{8}=3^{8}.4^{15}=8^{10}.9^{4}$$
$$8916100448256=2^{24}.3^{12}=3^{12}.4^{12}=8^{8}.9^{6}$$
$$11284439629824=2^{18}.3^{16}=3^{16}.4^{9}=8^{6}.9^{8}$$
$$14281868906496=2^{12}.3^{20}=3^{20}.4^{6}=8^{4}.9^{10}$$
$$18075490334784=2^{6}.3^{24}=3^{24}.4^{3}=8^{2}.9^{12}$$
$$39582418599936=2^{42}.3^{2}=3^{2}.4^{21}=8^{14}.9^{1}$$
$$50096498540544=2^{36}.3^{6}=3^{6}.4^{18}=8^{12}.9^{3}$$
$$63403380965376=2^{30}.3^{10}=3^{10}.4^{15}=8^{10}.9^{5}$$
$$80244904034304=2^{24}.3^{14}=3^{14}.4^{12}=8^{8}.9^{7}$$

论坛问题帖 | has_answers | Paul Richards | 2014-11-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018512 | MSE-1043020

Conjecture about the powers of 2 in the prime factorizations of the series $a_n = n(n+1)(n+2)$

I found the following law and would like to know what do you think about it and if anyone can explain why this is so. Also, is this already known and proven? Consider the following series: $$1\times2\times3, 2\times3\times4, 3\times4\times5,\ldots,n(n+1)(n+2)$$ or more compactly: $$a_n = n(n+1)(n+2)$$ Lets take the prime factorization for every element of a_n: $$2\times3, 2^3\times3, 2^2\times3\times5,2^3\times3\times5,\ldots$$ Lets take all the powers of 2 in every prime factorization in the above series and create a sequence out of them, and then group the elements of that sequence in tuples with size 4. The result of this we will call p_1. $$p_1 = (1,3,2,3),(1,4,3,4),(1,3,2,3),(1,5,4,5),\ldots$$ It seems that every 4-tuple with odd index in p_1 is $(1,3,2,3)$. Lets remove all 4-tuples with odd positions from p_1 and call the result p_2: $$p_2 = (1,4,3,4),(1,5,4,5),(1,4,3,4),(1,6,5,6),\ldots$$ Now p_2 will have $(1,4,3,4)$ at every odd position. If we take the p_3 sequence to have only those elements of p_2 that are with even indices, p_3 will have $(1,5,4,5)$ at every odd position. And so on... p_n will have $(1, n+2, n+1,n+2)$ at every odd position.

论坛问题帖 | has_answers | mzdravkov | 2014-11-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018513 | MSE-1065941

Two conjectures regarding $\varphi(n)$

There is a famous unsolved problem called Lehmer's Totient Problem which states that, $\varphi(n)\mid n-1 \implies n$ is a prime. Where $\varphi(n)$ is Euler's Totient Function. I was wondering about two conjectures of my own which I found unable to prove or disprove myself. The conjectures are, Conjecture 1 If n is an odd integer then $\varphi(n)+1\mid n \implies n$ is a prime. Conjecture 2 For odd n, $\varphi(n)+1\mid n\iff \varphi(n)\mid n-1$ Note that both the conjectures are independent of one another and the proof of any one willn't shed light on the proof of disproof the other unless a proof or disproof of Lehmer's original conjecture is done. So, can anyone help me in proving anyone of the conjectures? Note:- I don't know whether my conjectures are already well-known conjectures or not. If that's the case then any reference to the authentic source(s) will be enough. Added:- The conjectures are now edited taking in view the suggestions of fretty and A. Nicolas.

论坛问题帖 | has_answers | user170039 | 2014-12-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018514 | MSE-107911

Problems about consecutive semiprimes

I was playing around with semi-prime numbers and I made two conjectures, which are: Given any integer a, at least one of $a,(a+1),(a+2)$ or $(a+3)$ is not semi-prime. There are infinitely many integers a, such that $a,(a+1)$ and $(a+2)$ are semi-primes. I've written a computer program to verify the conjectures for values up to 700,000 (so, there's a high chance that they are both true). Can anyone give a proof or counter example for any of these problems, or a link to a paper on the subject? Thanks!

论坛问题帖 | has_answers | Obinna Okechukwu | 2012-02-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018515 | MSE-1086967

Prove or disprove this upper bound on chromatic number.

Let G be a simple connected finite graph and let $v \ge 4$ be the number of vertices, E the number of edges, $\chi(G)$ the chromatic number , $\omega(G)$ the clique number and Δ the maximum degree of G. Let $C(G)$ be a function of G which is equal to 3 when G contains cycles of odd length. Let $ {(d_1,d_2,...)}$ be the degree sequence of G. Let $i > 4$ be the smallest integer such that $$ \left\lceil \frac{(0.5\sum\limits_{j=1}^i d_j) -i}{i-3} \right\rceil < \frac{i}{2}$$ and $$ 2\frac{(0.5\sum\limits_{j=1}^i d_j) -i}{i-3} < \Delta $$ Take $i = v$ if a minimum value does not exist. Let $ B = 4 $ if $$ ^{**}1.5\leq\frac{(0.5\sum\limits_{j=1}^i d_j) -i}{i-3} < 2 $$ Otherwise $$ B = \left\lfloor 2\frac{(0.5\sum\limits_{j=1}^i d_j) -i}{i-3} \right\rfloor $$ Then $$\chi(G) \le \max\left\{C(G)\ , B,\ \omega(G)\right\}$$ ========== I've edited it completely because of bof's answer. ** I'm not too sure if this lower bound should be 1 or 1.5.

论坛问题帖 | has_answers | Napster | 2014-12-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018516 | MSE-1125563

Explanation for relationship between hypotenuse segments and leg lengths?

|` | ` x | ` | ` c a | z /` | / ` y |_/ ` |/|_____` b I'm new to this SE, but I have an SO account, so hello! Assume that z is the altitude that intersects the hypotenuse, and x and y are the resulting line segments. This is what my friend had to say about this. $$ z = \sqrt{xy} $$ You cannot find z with a and b. Here's what I said. Somehow, a and b must be related to x and y. At first I thought that $$\frac{a}{a+b} * c = x$$ That was wrong, as the altitude did not bisect the right angle. Through some thinking, but not proving, I said this, which was correct, AFAIK. $$\frac{a^2}{a^2+b^2} * c = x\\\\\frac{b^2}{a^2+b^2} * c = y $$ Some simplifying ensued. $$\frac{a^2}{c^2} * c = x\\\\\frac{b^2}{c^2} * c = y $$ More. $$\frac{a^2}{c} = x\\\\\frac{b^2}{c} = y $$ Now that a and b were used to find x and y, I got z with the geometric mean. $$ z = \sqrt{\frac{a^2}{c} * \frac{b^2}{c}} $$ That goes to: $$ z = \frac{ab}{c} $$ This was all and well, but then I was asked how did you come up with the first step?

论坛问题帖 | has_answers | user211599 | 2015-01-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018517 | MSE-1148543

Does any sum of twin primes, where the sum is greater than 12, also represents the sum of 2 other distinct primes?

I was in the midst of proving a conjecture when I came across an observation that led me to forming a potentially new conjecture. The conjecture goes as follows: Any given sum of twin primes (specifically the two primes in a twin prime pair, ie. 11 and 13) where the sum is greater than 12, also represents the sum of 2 other distinct primes. I've proven this for the first 1000 twin primes and my computer is calculating beyond that set. Anyways, does anybody have any ideas of how I could go about proving this conjecture? I apologize if this conjecture has already been posed.

论坛问题帖 | has_answers | user1939991 | 2015-02-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018518 | MSE-1156594

Generalizing conjecture of Bertrand

Joseph Bertrand's conjecture for primes from 1845, $\;p_{n+1} < 2p_n$, proved by Chebyshev 1852, can be generalized as follows: $$\forall a\in\Bbb Z_+\exists N\in\Bbb Z_+:(n\ge N\implies\exists p\in\Bbb P: ap_n<p<(a+1)p_n)$$ It seems to be proved for $a=2,3$ that $N=1$ works (Wikipedia). Computer tests for $p_n< 10,000,000$ $(n\le 664,579)$: $a=5,9,14$ holds for all primes $p_n\ge 2$, that is, it seems to hold for $N=1$. The case $a=4$ is tested for $p_n\ge 5\; (N=3)$. The case $a=25$ is tested for $p_n\ge 23\; (N=9)$. Computer tests for $p_n<1,000,000\;(n\le 78,498)$: Computer tests for $p_n<100,000\;(n\le 9,592)$: What is known about this? Is the conjecture possible to prove?

论坛问题帖 | has_answers | Lehs | 2015-02-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018519 | MSE-1157530

In the Collatz function, why $3^{2k}-1$ and $3^{2k-1}-1$ always share the same trailing trajectory?

Why are the trajectories always the same for numbers of the form $3^{2k}-1$ and $3^{2k-1}-1$ for the Collatz function? For example, let $k = 3$. So, $3^6-1 = 728$ and $3^5-1 = 242$. The trajectories are 242 728 121 364 364 same trajectory starts 182 182 91 91 274 ... 137 412 206 103 310 155 ...

论坛问题帖 | has_answers | user217938 | 2015-02-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018520 | MSE-1157556

In the Collatz function, why does 2^k-1 reach 3^k-1 after $2k$ steps, and could it be used to find divergent trajectories?

If you start calculating the Collatz function from an integer of the form 2^k-1, you will reach 3^k-1 after $2k$ steps. So, it is possible to pick a starting value that continuously zig-zags upwards for a period of time that can be easily defined. Has this property been used in trying to find divergent trajectories? One could say something like that when the starting value grows exponentially, the longest possible continuously growing sub-sequence expands linearly. Well.. you get the point

论坛问题帖 | has_answers | user217938 | 2015-02-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018521 | MSE-1178112

Prove conjecture using premises

I have three premises with me defined: $(B \land L) \implies A$ $(A \land D) \implies \lnot H$ $\lnot J \implies (D \land \lnot H)$ I need to prove the following conjecture with the help of the above premises: $(B \land L) \implies J$ and $(A \land H) \implies J$ Please help me out.

论坛问题帖 | has_answers | TheGaME | 2015-03-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018522 | MSE-1196327

Power towers of 2 and 3 - looking for a proof

Let $\uparrow$ denote the right-associative exponentiation operator: $a\uparrow b\uparrow c=a\uparrow(b\uparrow c)=a^{b^c}$ There is a sequence $A248907$ recently submitted to OEIS (see also $A256179$): $2, 3, 22, 23, 32, 222, 33, 322, 223, 232, 323, 332, 2222, 3222, 233, 333, 2322, ...$ It represents all power towers of numbers 2 and 3 in increasing order. In other words, it consists of all non-empty finite strings over the alphabet $\{2,3\}$ ordered in such a way, that once each string of digits is interspersed with the $\uparrow$ operators and evaluated, the sequence of results is monotonically increasing, i.e. $2=2$ $3=3$ $2\uparrow2=4$ $2\uparrow3=8$ $3\uparrow2=9$ $2\uparrow2\uparrow2=16$

$3\uparrow3=27$ $3\uparrow2\uparrow2=81$ $...$ I'm interested in a reasonably efficient algorithm for generating n^th element of $A248907$ (denote it a_n). It should avoid direct evaluation of power towers, otherwise it would easily run into huge numbers that would make the algorithm unfeasible in practice. I have a recursive algorithm that I suppose does the right thing, but I lack a rigorous proof of that. If $n\le12$, the corresponding elements $a_1\,..\,a_{12}$ are $2, 3, 22, 23, 32, 222, 33, 322, 223, 232, 323, 332;$ Otherwise, if n is odd, a_n is $a_{(n-1)/2}$ with the digit 2 prepended to it; Otherwise (n is even), a_n is $a_{(n-2)/2}$ with the digit 3 prepended to it. This algorithm builds the sequence from an "irregular" initial segment, and a "regular" tail consisting of pairs of elements that differ by the first digit (2 or 3, interleaved) that is prepended to previous elements taken sequentially from a certain offset, each element is used twice. Could you please help me prove (or disprove) its correctness?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2015-03-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018523 | MSE-1200325

Asymptotic behavior of the zeros of the digamma function

The gamma function has just one extremum on each interval $(k,k+1)$, where k is a negative integer. These extrema occur at the zeros of the derivative of the gamma function. Let z_n denote the n-th greatest zero, or the zero found on the interval $(-n, -n+1)$. As n grows, z_n nears $-n$. In other words, $z_n+n=\{z_n\}=z_n-\lfloor z_n\rfloor$ nears 0. I have numerically found $\{z_n\}^{-1}$ to be asymptotically equivalent to $\log(n)$: Here's the kicker, though. It seems that $\lim\limits_{n\to\infty}\left(\{z_n\}^{-1}-\log(n)\right)\approx 1-\gamma$. I appear to be reaching the limits of double precision. At $n=1754$, I start getting (presumably) wildly inaccurate values for z_n. If my approximations can be trusted before this point, however, then $\{z_n\}^{-1}-\log(n)+\gamma$ decreases and reaches about 1 at $n=1713$, but continues decreasing below 1. If this is truly the case, then the claim in the previous paragraph is not a strict equality. Does anyone know of a proof of $\{z_n\}^{-1}\sim\log(n)$ or if $\lim\limits_{n\to\infty}\left(\{z_n\}^{-1}-\log(n)\right)= 1-\gamma$ is true? I welcome any analytical or numerical evidence hinting at the confirmation or negation of either of these two statements. I myself know next to nothing about the gamma function or its relatives, but I noticed the zeros drifting left on their intervals and decided to investigate.

论坛问题帖 | has_answers | Regret | 2015-03-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018524 | MSE-1203370

Can I demonstrate a proposition without a rigorous proof?

Writing a paper, I want to declare a certain proposition, but is only supported by other research's empirical result. In this sort of case, what can I name to my argument in my paper? CONJECTURE is so much informal and seems to be so lazy. Otherwise, is it possible just to write down PROPOSITION in my paper without rigorous proof, but with some procedures attaining to that proposition including previous empirical results? Thank in advance!

论坛问题帖 | has_answers | kurtkim | 2015-03-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018525 | MSE-1232772

Conjectured diagonal Ramsey numbers

While doing some reading on the Wikipedia page for Ramsey numbers, I stumbled upon OEIS sequence A120414, which lists the allegedly conjectured values of the diagonal Ramsey numbers $R(n,n)$. I haven't found any indications in the standard web resources (WP, Mathworld) that suggest that any of the numbers in the known ranges for $R(n,n)$, $n\geq 5$ is more probable than others. What is this conjectured list based on? Considering the age of references on the OEIS page, the answer is probably elsewhere.

论坛问题帖 | has_answers | Daniel R | 2015-04-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018526 | MSE-1235390

Euler's Totient function $\forall n\ge3$, if $(\frac{\varphi(n)}{2}+1)\ \mid\ n\ $ then $\frac{\varphi(n)}{2}+1$ is prime

While I was studying Euler's Totient function, $\varphi(n)$, I stumbled upon the marvelous book "Index to Mathematical Problems, 1980-1984" By Stanley Rabinowitz. In this page of the book (link to Google books sample), the open problems and conjectures about the Totient function appear. All are really awesome, but one of them was specially interesting to me: CMB34 by M.V.Subbarao: if $(\varphi(n)+1)\ \mid\ n$ then, n or $n/2 \in \Bbb P$. Well it has two parts, the 'trivial' one is that if n is indeed a prime number, $\varphi(n)=n-1$, so $\varphi(n)+1 = n-1+1 = n$, and then $n \mid n$. But the other part is not trivial: when $(\varphi(n)+1)$ divides n (even), then $n/2$ is a prime number. Simply amazing and still open (if I am not wrong). I wanted to learn more, so I did my own try-outs of other combinations like CMB34, and I did find one that is very interesting, indeed I did not find any open conjecture or open problem about it, so here is my own proposed statement: (1) $\forall n\ge3$, if $(\frac{\varphi(n)}{2}+1)\ \mid\ n\ $ then $(\frac{\varphi(n)}{2}+1)\in \Bbb P$ And it really seems to work! I have made a Python test in the interval $[1,50000]$ and the statement is always true, no counterexamples. Here are some samples: $n = 52,\ \varphi(n)=24,\ \frac{\varphi(n)}{2}+1=13,\ 13\mid52$, and $13 \in \Bbb P$ $n = 57,\ \varphi(n)=36,\ \frac{\varphi(n)}{2}+1=19,\ 19\mid57$, and $19 \in \Bbb P$ $n = 66,\ \varphi(n)=20,\ \frac{\varphi(n)}{2}+1=11,\ 11\mid66$, and $11 \in \Bbb P$ $n = 68,\ \varphi(n)=32,\ \frac{\varphi(n)}{2}+1=17,\ 17\mid68$, and $17 \in \Bbb P$ $etc.$ Having those results in mind, I tried to generalize to $\frac{\varphi(n)}{2^i}+1, i\in\Bbb N$. And for instance for the case $i=2, (2^2)$, apart from some specific initial small n counterexamples $n = \{3,4,6,28,36,66\}$, for $n\gt66$ seems to be always true as well. I still did not try other values of i. I have uploaded the test written in Python here. I have continued trying to generalize, and the following statement seems also to be true always: (2) $\forall n\ge3, n=2k,\ k\in \Bbb N$, if $(\frac{\varphi(n)}{2}+1)\ \mid\ \frac{n}{2}\ $ then $(\frac{\varphi(n)}{2}+1)\in \Bbb P$ There are two differences with my main statement (1): in this case, n can only be an even number ≥3, so it can be divided by 2, and the relationship is true when a factor of $n/2$ is found (instead of a factor of n). I will update

the question if I find more successful combinations. The same can be applied to those n multiples of 4, and in this case, the expression is (trivially) true when $n = 4p\ \forall p \in \Bbb P, p\ge3$, so the list of odd primes appears in the original natural order of the prime numbers: (3) $\forall n\ge3, n=4k,\ k\in \Bbb N$, if $(\frac{\varphi(n)}{2}+1)\ \mid\ \frac{n}{4}\ $ then $(\frac{\varphi(n)}{2}+1)\in \Bbb P$ This is in other words: (3) $\forall n\ge3, n=4k,\ k\in \Bbb N$, if $\frac{n}{4} = (\frac{\varphi(n)}{2}+1)$ then $(\frac{\varphi(n)}{2}+1)\in \Bbb P$ In this case, the relationship is trivial for those $n=4p$ multiples of any p prime, because $\frac{\varphi(4p)}{2}+1 = \frac{\varphi(4)*\varphi(p)}{2}+1 = \frac{2*\varphi(p)}{2}+1 = \varphi(p)+1 = p-1+1 = p$ But the interesting point is that if n is a multiple of $4\ /\ n\not=4p, p \in \Bbb P$ then the relationship is false always, so no primes are found. Anyway, the expressions (1) and (2) seem more interesting than (3). I did not find any n able to make the relationship true for $\frac{n}{8}$ or $\frac{n}{16}$. The prime numbers $p = (\frac{\varphi(n)}{2}+1)$ generated by the expression (1): Except for the case $n=6$ in the rest of cases the prime number p seems to be always the greatest prime factor of n. Trivial but interesting, if p is factor of n, then (n/p) is an upper bound of the other prime factors of q, so $\forall q, q\in \Bbb P\ /\ q \not= p, \ q\mid n, \ q\le(n/p)$. And n is a subset of the numbers following the pattern: $n=(\frac{\varphi(n)}{2}+1)*k_1 /\ k_1 \in \{2,3,4,6\}$ It could be possible that I can not see the wood for the trees, and my statement is trivial, or already known, or nonsense, so I would appreciate any help to understand those results. I would like to share with you the following questions: Is my proposed statement about the Totient function already known or trivial? Is there a counterexample of it? Could it be generalized? Thank you! ** UPDATE 2015/04/22 ** My definition about the numbers n covering the properties explained at the statement (1) has been accepted at OEIS! (link here). ** UPDATE 2015/05/01 ** Related question about other possible interesting property (link here): (2)$\ \forall n\in [3,29]\cup[31,1000000]$, if $[(\frac{\varphi(n)}{2}+1)\ \mid\ n]\ \land\ [(\frac{\tau(n)}{2}+1)\ \mid\ n]\ $ then $[(\frac{\varphi(n)}{2}+1)\cdot(\frac{\tau(n)}{2}+1)] = n$

论坛问题帖 | has_answers | iadvd | 2015-04-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018527 | MSE-1236944

Euler's Totient function $\forall n\ge3$, if $(n-\phi(n)) = \sqrt{n}\ $,$\ $then $(n-\phi(n)) \in \Bbb P$

Recently I opened a question about what it might be a new property of Euler's Totient function. I am still studying the Totient function and I found another interesting relationship, it is very different from the ones I wrote in my former question, so I would like to ask about it apart because this might be another interesting topic. As far as I have checked, I did not find any reference to this property, but if I am wrong I will remove the question. This is my new proposed statement: $\forall n\ge3$, if $(n-\phi(n)) = \sqrt{n}\ $,$\ $then $(n-\phi(n)) \in \Bbb P$ Meaning that $n=p^2$ is the perfect square of a prime p if and only if the distance between n and $\phi(n)$ is exactly p. I have also tested this using Python, and no counterexamples are found in the interval $[1..7000]$. Usually when I find these kind of possible properties I test them, but I do not have enough theoretical knowledge to prove them, so any comment, hint or explanation about it is very appreciated. So as usual my questions are: Is my proposed statement about the Totient function already known or trivial? Is there a counterexample of it? Thank you!

论坛问题帖 | has_answers | iadvd | 2015-04-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018528 | MSE-1239863

Prove that $\det(A) \cdot v \, A^{-1} = \det(A+uv) \cdot v \, (A+uv)^{-1}$.

Let A be a $n \times n$ matrix, u a $n \times 1$ matrix and v a $1 \times n$ matrix. If A and $(A+uv)$ are invertible, prove that $$ \det(A) \cdot v \, A^{-1} = \det(A+uv) \cdot v \, (A+uv)^{-1}. $$ I have numerical evidence that this is true, but I cannot prove it. Note that the term on the left does not depend on u! Thank you.

论坛问题帖 | has_answers | teide4 | 2015-04-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018529 | MSE-1258928

Euler's totient and divisors count function relationship when $[(\frac{\varphi(n)}{2}+1)\cdot(\frac{\tau(n)}{2}+1)] = n$

I am studying the Euler's totient function $\varphi(n)$ and the divisors count function, $\tau(n)$, also named $d(n)$, and recently opened a question (link here) about the following condition: (1)$\ \forall n\ge3$, if $(\frac{\varphi(n)}{2}+1)\ \mid\ n\ $ then $(\frac{\varphi(n)}{2}+1)\in \Bbb P$ I wanted to do the same exercise, but adding an extra restriction regarding the divisors count function, $\tau(n)$, and I found the following to be true in the interval $[3,29]\cup[31,10^9]$ except for only one isolated counterexample at $n=30$: (2)$\ \forall n\in [3,29]\cup[31,10^9]$, if $[(\frac{\varphi(n)}{2}+1)\ \mid\ n]\ \land\ [(\frac{\tau(n)}{2}+1)\ \mid\ n]\ $ then $[(\frac{\varphi(n)}{2}+1)\cdot(\frac{\tau(n)}{2}+1)] = n$ So basically when the condition (1) happens and also the restriction $(\frac{\tau(n)}{2}+1)\mid n$ happens, then both $(\frac{\varphi(n)}{2}+1)$ and $(\frac{\tau(n)}{2}+1)$ are factors of n and if they are multiplied they are exactly n. E.g.: $n=21\ ,\ (\frac{\varphi(21)}{2}+1)=7 = a\ ,\ (\frac{\tau(21)}{2})+1=3 = b\ ,$ and $a\cdot b = 7\cdot3 = 21$ $n=999993\ ,\ (\frac{\varphi(999993)}{2}+1)=333331 = a\ ,\ (\frac{\tau(999993)}{2})+1=3 = b\ ,$ and $a\cdot b = 333331\cdot3 = 999993$ The only counterexample I found so far is: $n=30\ ,\ (\frac{\varphi(30)}{2}+1)=5 = a\ ,\ (\frac{\tau(30)}{2})+1=5 = b\ ,$ and $a\cdot b = 5\cdot 5 = 25 \not= 30$ I have been reading about the properties of both the totient and the divisor count functions, but I do not understand why always seems to happen (2), except for $n=30$, so I wanted to share with you these questions: Is it trivial? can be obtained by knowing the properties of $\varphi(n)$ and $\tau(n)$? Is there another counterexample for $n\gt10^9$? If somebody could check further would be very appreciated. Any hint or explanation about why (2) happens is very welcomed, thank you! UPDATE 2015/05/25 Tested up to 10^9, no counterxamples found apart from $n=30$. Here is the PARI/GP code: eulerdiv(limitval)={ for (n = 1, limitval, mytot = (eulerphi(n)/2)+1; mydiv = (numdiv(n)/2)+1; if ((n%mytot==0) && (n%mydiv==0), if((mytot*mydiv)==n,,print("Counterexample: "," ",n," ",mytot," ",mydiv)))) }

论坛问题帖 | has_answers | iadvd | 2015-04-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018530 | MSE-126583

Does $|n^2 \cos n|$ diverge to $+\infty$?

I was recently exposed to the problem of deciding whether $$ \lim_{n \to +\infty} |n \cos n| = +\infty$$ where the limit is taken over the integers. As $|\cos n|$ oscillates throughout the interval $[0,1]$, it seems plausible that every real number ought to be a limit point of this sequence. But to show the limit doesn't exist, it's enough to show the sequence is less than 2 infinitely often, and this can be done with continued fractions: we can find arbitrarily large p and q with q odd such that $$ \left| \frac{\pi}{2} - \frac{p}{q} \right| < \frac{1}{q^2} $$ which implies $$ \left| \frac{q-1}{2} \pi + \frac{\pi}{2} - p \right| < \frac{1}{q} $$ making p a good approximation of $\pi/2$ modulo π, and thus $$ \left| p \cos p \right| < \frac{p}{q} < 2 $$ This method fails if we consider instead the sequence $n^{1+\epsilon} \cos n$, since the final inequality is an increasing function of n. In fact, a heuristic statistical argument suggests that $$ \lim_{n \to +\infty} |n^{1 + \epsilon} \cos n| = +\infty $$ really does hold: imagine the n's are actually uniformly randomly selected from $[0, \pi]$, there is a roughly $2p/\pi$ chance that $|\cos n| < p$. For any positive constant B, the expected number of terms of $n^{1 + \epsilon} |\cos n|$ which are less than B is $$ \sum_{n=1}^{+\infty} \frac{2B}{\pi n^{1 + \epsilon}} $$ which, in particular, is finite. Of course, this is just heuristic. Does anyone know of a proof or disproof of the conjecture $$ \lim_{n \to +\infty} |n^{1 + \epsilon} \cos n| = +\infty $$ for positive real numbers ϵ, where the limit is taken only over integers? EDIT: Since there have been a few erroneous answers, I'll point out that observing $|\cos n|$ oscillates from 0 to 1 is not enough to show the limit does not exist. Consider the sequence $g(n)$ defined by $g(0) = 1$ $g(n) = \max\{ g(n-1), n / |\cos n| \} + 1$ Then $g(n)$ is an unbounded increasing sequence such that $|g(n) \cos n| > n$ for all n, and thus $$ \lim_{n \to +\infty} |g(n) \cos n| = +\infty $$ Roughly speaking, $g(n)$ diverges to $+\infty$ faster than $n \pmod \pi$ can approximate $\pi / 2$. The question of this post is whether n^2 (or $n^{1 + \epsilon}$) also diverges to $+\infty$ fast enough.

论坛问题帖 | has_answers | user14972 | 2012-03-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018531 | MSE-1273725

Why is it that if you square two prime numbers and add them, you get a number that is even and is not a perfect square?

If you do $x^2 + y^2 = n$ where x and y are both prime numbers and are both greater than 3, why is n always an even number that isn't a perfect square?

论坛问题帖 | has_answers | Parzivalz13 | 2015-05-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018532 | MSE-1283244

Conjecture about the product of the primitive roots modulo a prime number ($\prod Pr_p$)

While I was learning about the primitive roots modulo $p \in \Bbb P$ (I will call Pr_p to the complete list of the primitive roots module p) and having in mind the conjecture explained in this nice question about the Fortunate numbers and variations of that conjecture for factorials, I found (only empirically tested with Python) that the following expressions are true $\forall p \in [1,1000]$, $p \in \Bbb P$: Let $\mathcal{P}(n)$ be the previous prime smaller than n: $\prod Pr_p\:-\mathcal{P}(\prod Pr_p) = d_1$ is either 1 or a prime. Let $\mathcal{N}(n)$ be the next prime greater than n: $\mathcal{N}(\prod Pr_p)-\prod Pr_p\:= d_2$ is either 1 or a prime. So it seems that the product of the primitive roots modulo a prime p is at a distance d_1 of its previous closer prime and a distance d_2 of its closer next prime, being d_1 and d_2 = 1 or a prime number. E.g.: $Pr_{17}=\{3, 5, 6, 7, 10, 11, 12, 14\}$ $\prod Pr_{17}=11642400$ $\mathcal{P}(\prod Pr_{17})=11642387$, so $d_1=11642400-11642387=13$ $\mathcal{N}(\prod Pr_{17})=11642419$, so $d_2=11642419-11642400=19$ (*) Notice that in the sample by chance d_1 and d_2 incidentally are exactly the previous and next prime numbers of the original $p=17$. I just like to play with the numbers and find relationships between them that I did not know, and usually that is empirical. My theoretical knowledge is quite basic (usually I learn from the answers at MSE very much!). I wanted to share with you the following questions to know more about the properties of the primitive roots modulo p: Does it make sense that according to the definition or properties of the primitive roots modulo a prime p, the product of them was able to have those properties? Is there a counterexample of them? (my computer takes time for bigger values of p so if somebody with Mathematica or other software, a good computer and interested in the topic could give it a try, that would be really appreciated) Is it a trivial property? UPDATE 2015/05/18: added in my blog the Python code for the test in the interval [1,1000]. Thank you!

论坛问题帖 | has_answers | iadvd | 2015-05-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018533 | MSE-1283375

Conjecture: only one even Fibonacci term divided by two gives a prime: $F(9) = 34 = 2 \times 17$

Every Fibonacci term $F(3n)$ is divisible by two $F(3) = 2$ $F(6) = 8$ $F(9) = 34$ $...$ After seeking Fibonacci tables factorization until $F(10000)$, for every term $\frac{F(3n)}{2}$, it appears there is just one result which is prime $\frac{F(9)}{2} = \frac{34}{2} = 17$ all other $\frac{F(3n)}{2}$ = composite numbers I can't see some reason for this to happen, but since there is no more primes in so huge quantity of terms, I suppose there is not more any prime for $\frac{F(3n)}{2}$ http://mersennus.net/fibonacci Miguel Velilla

论坛问题帖 | has_answers | Miguel Velilla | 2015-05-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018534 | MSE-1292012

Prove that $\lim_{n\rightarrow \infty} \frac{\log_{10}\lfloor\text{Denominator of } H_{10^n}\rfloor+1 }{10^n}=\log_{10} e$

In short, my question is asking to prove that the $$\lim_{n\to\infty}\frac{\text{number of digits in the denominator of} \sum_{k=1}^{10^n} \frac 1k}{10^n}=\log_{10} e$$ I know that the number of digits in a number is $\lfloor \log_{10} n\rfloor +1$ and that the Harmonic numbers are given by $\gamma+\psi_0 (n+1)$, where $\psi_0 (x)=\frac{\Gamma'(x)}{\Gamma(x)}$ but I don't see how to find the number of digits in the denominator of $\psi_0(n+1)+\gamma$. Context:From Wolfram MathWorld, "The numbers of digits in the denominator of H_{10^n} for $n=0, 1, \ldots$ is given by $1, 4, 40, 433, 4345, 43450, 434110, 4342302, 43428678, \ldots$ (OEIS A114468). These digits converge to what appears to be the decimal digits of $\log_{10}e=0.43429448\ldots$ (OEIS A002285)."

论坛问题帖 | has_answers | Teoc | 2015-05-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018535 | MSE-1293679

A question and a conjecture on $USp(N)$ group

$USp(N)$ with N an even integer is defined as the group of unitary matrices M that satisfy $M^TJM=J$, where M^T is the transpose of M and J is the anti-symmetric N-by-N matrix \begin{equation} J= \left(\begin{array}{cc} &I\\ -I& \end{array} \right) \end{equation} where I is the $\frac{N}{2}$-by-$\frac{N}{2}$ identity matrix. $USp(N)$ group has $\frac{N(N+1)}{2}$ generators. For $USp(4)$, the generators can be represented as Γ_{ab} which are defined through another set of matrices Γ_a by \begin{equation} \Gamma_{ab}=\frac{1}{2i}(\Gamma_a\Gamma_b-\Gamma_b\Gamma_a) \end{equation} and Γ_a themselves satisfy $\Gamma_a\Gamma_b+\Gamma_b\Gamma_a=2\delta_{ab}$. In this case we can verify that $J\Gamma_{ab}J=\Gamma_{ab}^T$ and $J\Gamma_{a}J=-\Gamma_{a}^T$. Moreover, $\sum_{ab}\Gamma_{ab}\Gamma_{ab}$ and $\sum_{a}\Gamma_{a}\Gamma_{a}$ are invariant under the fundamental representation of $USp(4)$, namely, $\sum_{ab}\Gamma_{ab}\Gamma_{ab}=U^\dagger\sum_{ab}\Gamma_{ab}\Gamma_{ab}U$ and $\sum_{a}\Gamma_{a}\Gamma_{a}=U^\dagger\sum_{a}\Gamma_{a}\Gamma_{a}U$ for any $USp(4)$ matrix U. My question is: for any even $N>2$, can we always find some Hermitian matrices Γ_a so that $J\Gamma_aJ=-\Gamma_a^T$ and $\sum_a\Gamma_a\Gamma_a$ is invariant under the fundamental representation of $USp(N)$ in the above sense? My conjecture to this question is that there are indeed such Γ_a, and more precisely, there are $N+1$ of them and they can still be used to derive the $\frac{N(N+1)}{2}$ generators of $USp(N)$ by $\Gamma_{ab}\propto\Gamma_a\Gamma_b-\Gamma_b\Gamma_a$. I appreciate if any one will check this conjecture and answer this question.

论坛问题帖 | no_answers_at_retrieval | Mr. Gentleman | 2015-05-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018536 | MSE-1297960

How I can make a proof to this conjecture if it is possible?

Is there someone who can show me the way helping me to proof this conjecture. If it's not open, at a least show me links or papers which to be helpful. Conjecture: Assume $c > 0$ and that an equilibrium point $\bar{z}$ of the difference equation: $$z_{n+1}=\frac{\alpha+z_{n}\beta+z_{n-1}\lambda }{A+Bz_{n}+cz_{n-1}}, \ n=0,1,\dots$$ is locally asymptotically stable. My question is: Show that $\bar{x}$ is a global attractor of all positive solutions of precedent equation. Note: this conjecture deal with the following simpler system: $$\begin{cases} x_{n+1}=\dfrac{\alpha_{1}}{y_{n}} \\ y_{n+1}=\dfrac{\alpha_{2}}{z_{n}} \\ z_{n+1}=\dfrac{\alpha_{3+}+\beta_{3}x_{n}+\sigma_{3}y_{n}+\lambda_{3}z_{n}}{A_{3}+B_{3}x_{n}+c_{3}y_{n}+D_{3}z_{n}}{}\end{cases} \quad n=0,1,\dots,$$ with non-negative parameters and non-negative initial conditions such that the denominators are always positive. any help is very welcom , thank you for your replies or any comments

论坛问题帖 | no_answers_at_retrieval | salimmath15 | 2015-05-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018537 | MSE-130070

I noticed a pattern, does this have a name?

First of all I am a programmer, not a mathematician, so I may articulate what I am trying to say very poorly. I was working with powers of 2 when I noticed a relationship I had never noticed before. Basically, if you look at the powers of 2, every other power of two can be added together make a multiple of 10. Furthermore, if the two numbers are 2^{10} and 2^8, adding them together will result in $2^7\cdot 10$. Put another way, $2^n + 2^{n-2} = 2^{n-3}\cdot 10$. Does this pattern have a name? I am attaching an image just to clarify my poor explanation. Thanks all! $$ \begin{matrix} & & \vdash & 20 & \dashv & \vdash & 160 & \dashv & \vdash & 1280 & \dashv \\ 1 & 2 & 4 & 8 & 16 & 32 & 64 & 128 & 256 & 512 & 1024 \\ & \vdash & 10 & \dashv & \vdash & 80 & \dashv & \vdash & 640 & \dashv \\ \end{matrix} $$

论坛问题帖 | has_answers | Brendon Dugan | 2012-04-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018538 | MSE-1302206

Why are there so many conjectures in number theory and comparatively less in others?

My question is that : Why are there so many conjectures in elementary number theory and comparatively less in others? This is particularly weird because every topic in maths should have its equal share of conjectures. On the other hand there are so many conjectures in number theory. Also (most of) these conjectures are so easy to state but literally impossible to prove. I thought of this question because I myself have two open conjectures in number theory.

论坛问题帖 | has_answers | user210387 | 2015-05-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018539 | MSE-131281

Is it known or new?

Possible Duplicate: Starting digits of 2^n While I was randomly working with number patterns, I came along with some interesting pattern which seems to turn to a conjecture in fact. My conjecture can be stated as follows : For any number n there exists any number p such that the number np can be a power of two or 2^k (here the np doesn't represent product of n an p but the number p is placed to the right of n , for eg: if the number n is 11 then if p is 12 then np is 1112 not $11 \cdot 12$). For a practical demonstration let us consider a number 11 there exists any number you choose for example $805916207174113034 24$ (I have selected this number deliberately and I need to find an effective way of choosing numbers) such that if the number is placed to right of 11 it becomes $11805916207174113034 24$ which in turn makes the complete entity as 2^{70} . For a practical formulation, it took some time for me in the mathematical representation. My conjecture can be stated as " Consider a number N, then there exist a number P such that $$((N\cdot 10^{\lceil\,\log_{10} P\,\rceil})+P) \equiv 0\pmod{2^k} $$for some K. " Does this already exists or it is new? , if it exists please give me a reference. P.S : I have been trying for a proof, and I have a proof with me. Is it worth of publishing ? Thank you.

论坛问题帖 | has_answers | IDOK | 2012-04-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018540 | MSE-1313318

Conjecture about linear diophantine equations

I've been dabbling with linear Diophantine equations and came across a rather interesting pattern that I would like to conjecture as true but I have no idea how about to come up with a proof. Let $ax+by=c$ be the linear Diophantine equation such that $gcd(a,b)=1$. Obviously $1|c$ so there are solutions to the equation. If we consider only positive solution pairs (x,y) for the equation, then the number of ordered pairs that satisfy the equation is equal to (c/a)/b, rounded down to the nearest integer. Can anyone verify this by counterexample or by proof? Thanks

论坛问题帖 | has_answers | Jeffrey L. | 2015-06-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018541 | MSE-1318254

Is the conjecture $Per A $ is the largest eigenvalue of $\tilde{A}$ being solved?

Let A be a positive semidefinite matrix of order n. Is the conjecture $Per A $ is the largest eigenvalue of $\tilde{A}$ is being solved? Where $\tilde{A}$ is the matrix of order $n!\times n!$whose $(\sigma, \tau)^{th}$ entry is given by $\prod _{i=1}^{n} a_{{\sigma (i)}\tau(i)}$ $Per (A)=\sum _{\sigma\in S_n} \prod_{i=1}^{n} a_{i\sigma (i)}$ The conjecture has being proved for $n\leq 3$ in Bapat, R. B. ; Sunder, V. S. An extremal property of the permanent and the determinant. Linear Algebra Appl. 76 (1986), 153-163. Is there any further development in this problem?

论坛问题帖 | has_answers | Babai | 2015-06-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018542 | MSE-1323497

What do you think about this conjecture?

Beforehand, please know that I'm a little bit of an amateur mathemitician, so this could be very wrong. But, I have tested it over and over and over again and it seems to be plausible, but there's still other things that I haven't tried that might possibly prove it wrong. That's why it's a 'conjecture' and not a 'theorem' or definitely not a 'law'. This conjencture involves functions. Any function. Let's say we have the function $f(x)$. Obviously, this doesn't mean I just multiply f by x, it is a function of x. But, what if there was a way to separate the two? Well, I came up with this simple equation as the base of this entire conjecture: $$f(x)=f*ƨx$$ where f is the function modifier and $ƨx$ is the terminal, meaning what the function modifier is in terms of. The first problem with this is $ƨx$. This is not a number or anything like that. It is its own concept, but it can be utilized in an equation. The second problem with this is f. How do he know that f is used in a function, and that it's not variable? Well, we don't know, unless it is blatantly defined as a function modifier. Usually, it is assumed that it's a function modifier, since it is in front of $ƨx$ and it's 'f', probably the most commonly used function modifier. Simply put, $f(x)$ is the same as a function modifier f multiplied by a value that has $ƨ$ in front of it. There is no operation that can add $ƨ$ in front of a value, $ƨ$ can only be taken from an actually function. Since: $$f(x) = f*ƨx$$ Therefore: $$f = f*ƨx/ƨx$$ and Also: $$ƨx = f*ƨx/f$$ With this conjecture, I can find the solution to some problems: find $g(x)$ in terms of a and b, assuming f and g are both function modifiers $$f(a)+f(b)=g(a+b)$$ Solve: $$f(a)+f(b)=g*ƨ(a+b)$$ $$(f(a)+f(b))/ƨ(a+b)=g$$ $$(f(a)+f(b))/ƨ(a+b)*ƨx = g*ƨx$$ $$(f(a)+f(b))ƨx/ƨ(a+b)=g(x)$$ Assuming $f/f=1$ (which it should) $$f/f *(f(a)+f(b))ƨx/ƨ(a+b)=g(x)$$ $$((f(a)+f(b))f*ƨx)/f*ƨ(a+b)=g(x)$$ $$((f(a)+f(b))f(x))/f(a+b)=g(x)$$ So my solution is: $$((f(a)+f(b))f(x))/f(a+b)=g(x)$$ Now after some tests, this proves to be right. I'll try to prove it here by saying $f(x)=2x-1$. This next problem will be similar to the last problem, excpet that $f(x)$ will now be defined. find $g(x)$, assuming f and g are both function modifiers. $f(x)=2x-1$ $$f(a)+f(b)=g(a+b)$$ Solve: $$((f(a)+f(b))f(x))/f(a+b)=g(x)$$ $$((2a-1)+(2b-1))(2x-1)/(2(a+b)-1)=g(x)$$ $$(2a-1+2b-1)(2x-1)/(2a+2b-1)=g(x)$$ $$(2a+2b-2)(2x-1)/(2a+2b-1)=g(x)$$ $$(4ax-2a+4bx-2b-2x+2)/(2a+2b-1)=g(x)$$ Now let's see if $g(x)$ really is equal to that... $$f(a)+f(b) = 2a-1+2b-1 = 2a+2b-2$$ Let's plug in $a+b$ into $g(x)$ to see if it's equal to $f(a)+f(b)$ $$((2a-1)+(2b-1))(2(a+b)-1)/(2(a+b)-1)=g(a+b)$$ The $2(a+b)-1$ in the nominator cancels out with the $2(a+b)-1$ in the denominator: $$(2a-1)+(2b-1)=g(a+b)$$ $$2a+2b-1=g(a+b)$$ And, it works! If you're thinking this, I also thought that it was a little suspicious that the nominator and the denominator cancel out, but considering I used $ƨx$ as the basis, that suspicion decreased. I've also used this conjecture to find another way to define the Gamma Function($\Gamma(x)$) using this definition: $$\Gamma(1-x)\Gamma(x)=\pi/sin(\pi*x)$$ $$\Gamma*ƨ(1-x)\Gamma*ƨ(x)=\pi/sin(\pi*x)$$ $$\Gamma^2*ƨ(1-x)ƨx)= \pi/sin(\pi*x)$$ $$\Gamma^2*ƨx=\pi/sin(\pi*x)ƨ(x-1)$$ $$\Gamma^2*ƨx*ƨx=\pi/sin(\pi*x)ƨ(x-1) *ƨx$$ $$(\Gamma*ƨx)^2=\pi/sin(\pi*x)ƨ(x-1) *ƨx$$ $$(\Gamma(x))^2=\pi/sin(\pi*x)ƨ(x-1) *ƨx$$ $$\Gamma(x)=sqrt((\pi*ƨx)/(sin(\pi*x)ƨ(x-1)))$$

However, this solution can not be used to directly utilize the function, since we don't know what $2x$ and $2(x-1)$ are, so it stays like that. Since it doesn't involve any other function, I can't multiply the numerator and denominator by a function operator. I've also used this conjecture to attempt to a find a way to define antiderivation. Now I COULD be very wrong on this because I don't really take into account the limit, which I have a feeling I should be, and I'm only assuming that h is still approaching 0 after all of this. $$f'(x)= lim(h>0) (f(x+h)-f(x))/h$$ $$f'(x)= lim(h>0) (f*2(x+h)-f*2(x))/h$$ $$f'(x)= lim(h>0) (f*(2(x+h)-2(x)))/h$$ $$f'(x)*h= lim(h>0) f*(2(x+h)-2(x))$$ $$(f'(x)*h)/(2(x+h)-2x) = lim(h>0) f$$ $$(f'(x)*h)/(2(x+h)-2x)*2x = lim(h>0) f*2x$$ $$(hf'(x)*2x)/(2(x+h)-2x) = lim(h>0) f(x)$$ $$ f(x) = lim(h>0)(hf'(x)*2x)/(2(x+h)-2x)$$ This is proven to be wrong, since h is not defined outside of the limit, but still, it shows the use of my conjecture. And so, this has been my conjecture. I don't know if it's solid, and I hope it doesn't get disproved (most likely will), but even if I'm redefining something that I didn't know existed, I hope it shows another way of thinking it.

论坛问题帖 | has_answers | Sam | 2015-06-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018543 | MSE-1326999

Where $ax + b$ prime infinitely often, is $ax + b - 2$ semiprime at least once?

I'm trying to figure out a way to prove this: Given arithmetic progression $ax + b$ where a, b coprime and $ax + b$ is prime infinitely often, it is the case at least once that $ax + b - 2$ is semiprime when $ax + b$ is prime. I'm looking for a way either to prove it, or whether an existing theorem is already known from which this would follow. Stated another way, given arithmetic progression $ax + b$ where a, b coprime and $ax + b$ is prime infinitely often, it is at least once the case that $ax + b$ produces a cd + 2 prime, where c, d are prime. The list of cd + 2 primes is A063638 on OEIS. What we know: A Chen prime is a prime p such that $p + 2$ is either prime or semiprime. Chen proved in 1966 there are infinitely many of these. There are also infinitely many primes p such that $p - 2$ is specifically semiprime (not just prime-or-semiprime). Friedlander and Iwaniec proved this in their 2010 book. Note that I'm not asking whether $ax + b - 2$ is semiprime infinitely often. I just need it to happen once, that $ax + b$ is prime while $ax + b - 2$ is semiprime. A more specific version of the problem is this: Given arithmetic progression $P(P + 2)x + (P + 4)$ where P, $P + 2$ are twin primes, it is the case at least once that $P(P + 2)x + (P + 2)$, i.e., $(Px + 1)(P + 2)$ is semiprime when $P(P + 2)x + (P + 4)$ is prime. The deeper problem I'm trying to show is that it is the case at least once that $Px + 1$ is prime when $P(P + 2)x + (P + 4)$ [or, said another way, $(Px + 1)(P + 2) + 2$] is also prime. I'm trying to use Chen's theorem as a strategy to show this--- in other words, to show that $(Px + 1)(P + 2)$ must be semiprime at least once (and so, ergo, $Px + 1$ must be prime at least once) when $P(P + 2)x + (P + 4)$ is prime. All my efforts to prove one just leads to a tautology where I must assume the other first. Any thoughts, suggestions or help would be appreciated.

论坛问题帖 | has_answers | Robert Gross | 2015-06-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018544 | MSE-135133

Does this $\zeta(s)$ identity have a name?

I have generalized the product from this thread: Let $s=2n+1$ for $n\ge1$, $$\zeta (s)=\frac{\zeta (2 s)}{\zeta (2)} \prod _{n=1}^{\infty } \frac{\sum _{i=0}^{s-1}(-p_n){}^i{(p_{n}-1)p_{n}^{s-2}}$$ This is a $\zeta(s)$ identity for each odd s. Does it have a name? I can only find $\zeta(3)$ in the literature.

论坛问题帖 | has_answers | Fred Daniel Kline | 2012-04-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018545 | MSE-1360575

Inequality in 3 variables (conjecture)

Let a, b, c be nonnegative real numbers such that $a+b+c=3$. If $0<k\leq 3+2\sqrt{3}$, then $$\frac{a}{b^2+k}+\frac{b}{c^2+k}+\frac{c}{a^2+k}\geq \frac{3}{1+k}$$ If $k=3+2\sqrt{3}$, then equality occurs if $a=b=c=1$ or $a=0$, $b=1-\sqrt{3}$ and $c=\sqrt{3}$ or any cyclic permutation thereof. This inequality can be found in Vasile Cirtoaje's Discrete inequalities, Volume 4 - page 61. Any idea for this inequality? Thank you!

论坛问题帖 | has_answers | Marlo | 2015-07-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018546 | MSE-1365712

Prove that there exist infinitely many integers $(n^{2015}+1)\mid n!$

I conjecture that there exist infinitely many integers n such that $$(n^{2015}+1)\mid n!.$$ I have seen a simpler problem that there exist infinitely many integers n such that $(n^2+1)\mid n!$. Alternatively, I considered the Pell equation $n^2+1=5m^2$, $2m<n$, but for 2015 I can't figure it out.

论坛问题帖 | has_answers | user225250 | 2015-07-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018547 | MSE-1373123

A conjectured identity for tetralogarithms Li_4

I experimentally discovered (using PSLQ) the following conjectured tetralogarithm identity: $$720 \,\text{Li}_4\!\left(\tfrac{1}{2}\right)-2160 \,\text{Li}_4\!\left(\tfrac{1}{3}\right)+2160 \,\text{Li}_4\!\left(\tfrac{2}{3}\right)+270 \,\text{Li}_4\!\left(\tfrac{1}{4}\right)+540 \,\text{Li}_4\!\left(\tfrac{3}{4}\right)+135 \,\text{Li}_4\!\left(\tfrac{1}{9}\right)\\ \stackrel?=19 \pi ^4+30 \left(\pi ^2-\beta ^2\right) \left(10 \alpha ^2-12 \alpha \beta +3 \beta ^2\right)-30 \,\alpha ^2 \left(19 \alpha ^2-24 \alpha \beta +8 \beta ^2\right)\!,\\ \text{where $\alpha=\ln 2, \,\, \beta=\ln 3$.}$$ Numeric computations show that the absolute value of the difference between the left and right sides is smaller than 10^{-10^5}, so I believe the identity must hold exactly. Moreover, this is the simplest vanishing linear combination with integer coefficients of elementary terms and tetralogarithm values at rational points in the interval $(0,1)$ that I found. How can we prove it?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2015-07-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018548 | MSE-1379464

Sum involving the number of zeros of k

I'm currently interested in sums involving digit-functions. Especially, I'd like to calculate the following sum: $$ s=\sum_{k=1}^{\infty}{\frac{a_0(k)}{k\left(k+1\right)}} $$ Where $a_0(k)$ is the number of zeros in the base 10 representation of k. Normally, such sums can be tackled by considering products of the form $P(x,q)=\prod_{k=0}^{\infty}{\left(1+f_1(x)q^{10^k}+...+f_9(x)q^{9\cdot 10^k}\right)}$, but this approach seems not to work here, because the zero is a bit problematic. Numerically, it seems that $s=\frac{10+\gamma+\psi{(0.1)}}{9}$ where ψ is the digamma function.

论坛问题帖 | has_answers | Redundant Aunt | 2015-07-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018549 | MSE-1384588

Improved Betrand's postulate

I want to show that $2p_{n-2} \geq p_{n}-1$... Bertand's postulate shows us that $4p_{n-2}\geq p_{n}$ but can we improve on this? any ideas?

论坛问题帖 | has_answers | Brad Graham | 2015-08-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018550 | MSE-1385325

If (a_n) is increasing and $a_n^{1/c^n}\to\infty$, then $\sum\frac1{a_n}$ is irrational?

I $\DeclareMathOperator\lcm{lcm}$am trying to generalise the result from this question: If (a_n) is increasing and $\lim_{n \to \infty} a_n^{1/2^n} = \infty$, show that $\sum_{k=1}^{\infty} 1/{a_n}$ is irrational and I try to prove that Conjecture (strong). If (a_n) is an increasing natural sequence and $c>1$ such that $a_n^{1/c^n}\to\infty$, then $\sum\frac1{a_n}$ is irrational. (Note that for $c\geq2$ we are done already.) Though I will also be happy if I only manage to prove that Conjecture (weak). If (a_n) is an increasing natural sequence such that $a_n^{1/c^n}\to\infty$ for all $1<c<2$, then $\sum\frac1{a_n}$ is irrational. In particular I am wondering whether $\sum_{n=0}^\infty\frac1{2^{2^n}+1}$ is irrational. (just curiosity) I'll summarize the proof in the linked question: It suffices to prove that $$\liminf_{k\to\infty}\sum_{n=k+1}^\infty\frac{\lcm(a_1,\ldots,a_k)}{a_n}=0;$$ we'll prove that in fact $$\liminf_{k\to\infty}\sum_{n=k+1}^\infty\frac{a_1\cdots a_k}{a_n}=0.$$ Let $a_n=b_n^{2^n}$ (so $b_n\to\infty$) and let r be such that $b_n\leq b_r$ for all $n\leq r$. Then $$\frac{a_1\cdots a_{r-1}}{a_{r+n}}=\frac{b_1^2\cdots b_{r-1}^{2^{r-1}}}{b_{r+n}^{2^{r+n}}}\leq\frac{b_r^{2+\cdots+2^{r-1}}}{b_{r+n}^{2^{r+n}}}\leq b_{r+n}^{2^n(2^r-2)-2^{r+n}}=b_{r+n}^{-2^{n+1}}.$$ Now choose r such that $b_n\geq M>1$ for all $n\geq r$ and we get $$\sum_{n=r}^\infty\frac{a_1\cdots a_{r-1}}{a_n}\leq\sum_{n=0}^\infty M^{-2^{n+1}}\leq\sum_{n=0}^\infty M^{-(n+1)}=\frac1M.$$ Because $M>1$ was arbitrary, we get $$\liminf_{k\to\infty}\sum_{n=k+1}^\infty\frac{\lcm(a_1,\ldots,a_k)}{a_n}=0.$$ It's not hard to see that this strategy (proving $\liminf_{k\to\infty}\sum_{n=k+1}^\infty\frac{\lcm(a_1,\ldots,a_k)}{a_n}=0$) does not work for $c<2$. For example for $a_n=2^{2^n}+1$ we have $\lcm(a_0,\ldots,a_k)=a_1\cdots a_k=2^{2^{k+1}}-1\approx a_{k+1}^2$ so even controlling the size of the first term is hopeless. What else can we do? Note: using the same strategy as in the linked question I was able to control the terms of $\sum_{n=k+1}^\infty\frac{a_1\cdots a_k}{a_n}$ (for suitable k) except the first $l-1$, where l is such that $c^l\geq c^{l-1}+1$. I'm not mentioning the proof here because controlling the first $l-1$ terms seems hopeless anyway.

论坛问题帖 | has_answers | Bart Michels | 2015-08-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018551 | MSE-1386628

Can the sum of 3 unique primes be expressed as the sum of 2 primes?

Let's consider the example, $$ 3 + 11 + 19 = 33 \\ 2 + 31 = 33 $$ we can see that there are cases where the sum of 3 primes be expressed as the sum of 2 primes. However, I couldn't find a case where a sum of 3 primes couldn't be expressed as a sum of 2 primes. Is there such a conjecture already in the world of mathematics? Or is there a counter example?

论坛问题帖 | has_answers | Valera Rozuvan | 2015-08-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018552 | MSE-1395811

conjecture about primes and a certain q-series.

Using wolfram Mathematica ,I observed an interesting and surprising property concerning prime numbers and q-series which I could not prove.Yet there is strong evidence supporting it. I would be happy if anyone can find a counter-example.On the other hand,if it could turn out to be true,it would no doubt have amazing consequences in prime number theory. Let p be an odd prime and $$\frac{1}{(q;q^4)_{\infty}^p}=\prod_{n=0}^{\infty}\frac{1}{(1-q^{4n+1})^p}=1+\sum_{n=1}^{\infty} \phi(n)\,q^n$$ with the usual convention $$(a;q)_{\infty}=\prod_{k=0}^{\infty} (1-aq^{k})$$ then $$\phi(n)\equiv 0\pmod{p}$$ is true for all natural numbers $\{1,2,3,\dots\}$ except at multiples of p. For example, $$\frac{1}{(q;q^4)_{\infty}^3}=1 + 3q +6q^2 + \color{brown}{10}q^3 + 15q^4 + 24q^5+ \color{brown}{37}q^6+\dots$$ $$\frac{1}{(q;q^4)_{\infty}^5}=1 + 5q +15q^2 + 35q^3 + 70q^4 + \color{brown}{131}q^5+ 235q^6+405q^7+\dots$$ Note: $\phi(n)$ is just a notation I chose as a matter of personal taste. Can any one try to prove the conjecture or rather verify it by numerical methods,thanks in advance.

论坛问题帖 | has_answers | Nicco | 2015-08-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018553 | MSE-1397188

connection between odd primes and a certain q-series

I posed a conjecture about odd primes and a certain q-serieshere.I thought it would be more appropriate ,if I could ask the converse of the aforementioned problem . Is p an odd prime iff $$\prod_{n=0}^{\infty}\frac{1}{(1-q^{4n+1})^p}=1+\sum_{n=1}^{\infty}\phi(n)\,q^n$$ such that $$\phi(n)\equiv 0\pmod{p}$$ is true for all natural numbers $n\in\mathbb{N}$ except at multiples of p. I've experimentally verified the conjecture for many values of p,and it seems to hold.Independent verification would be greatly appreciated.This conjecture seems to point to a deep connection between primes and modular forms,which I was not able to prove.Outlining a proof or connection between the particular q-series above and odd primes will be appreciated.

论坛问题帖 | has_answers | Nicco | 2015-08-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018554 | MSE-1400244

conjectured new generating function of fibonacci numbers

I conjecture a new generating function for the fibonacci numbers F_{n}. Given,the following conjectured q-continued fraction $$\chi(q)=\cfrac{1}{1+q-\cfrac{(1+q^2)}{1+q^3+\cfrac{q^2(1-q)(1-q^3)}{1+q^5-\cfrac{q^3(1+q^2)(1+q^4)}{1+q^7+\cfrac{q^4(1-q^3)(1-q^5)}{1+q^9-\ddots}}}}}$$ how do we show that, $\chi\Big(\frac{1}{q}\Big)= \sum_{n=1}^\infty (-1)^{n-1} F_{n}q^n$ is true?

论坛问题帖 | has_answers | Nicco | 2015-08-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018555 | MSE-1400700

a conjectured new generating function of narayana's sequence

In the 14th century ,an Indian mathematician T.V Narayana came up with a sequence now named after him.The sequence satisfies the recurrence $$a_{n}=a_{n-1}+a_{n-3}$$ Starting with $a_{0}=a_{1}=1$, and $a_{2}=2$ hence, $a_n = 1, 1, 2, 3, 4, 6, 9, 13,\dots$ I would like to conjecture a new generating function for narayana's sequence a_{n}. Given $$\psi\Big(q\Big)=\cfrac{1}{1+q-\cfrac{(q^2)}{1+q^3+\cfrac{q^2(1-q)(1-q^3)}{1+q^5-\cfrac{q^3(1+q^2)(1+q^4)}{1+q^7+\cfrac{q^4(1-q^3)(1-q^5)}{1+q^9-\ddots}}}}}$$ How can we show that $$\psi\Big(q\Big)= \sum_{n=0}^\infty (-1)^{n} a_{n}q^n = 1 -1q +2q^2-3q^3+4q^4-6q^5+\dots$$ is true,such that the functional equation holds $$\psi\Big(q\Big)=\frac{1}{q}\psi\Big(\frac{1}{q}\Big)$$

论坛问题帖 | has_answers | Nicco | 2015-08-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018556 | MSE-1400920

Infinitely many correct solutions to equation?

I conjecture that there are infinitely many correct solutions to this equation: (Where we are assuming $a,b \in \Bbb{N}$) $$a!+1=b^2$$ I chose to list the first three solutions below: $4!+1=5^2$ $5!+1=11^2$ $7!+1=71^2$ and so on... Could I have a proof or disproof of my conjecture? (I don't even know where to start)

论坛问题帖 | has_answers | user253055 | 2015-08-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018557 | MSE-1401747

an interesting q-series and a certain continued-fraction

My aim is to find a rigorous proof of the following conjectured identity.Given $$1+q+q^2-q^4-q^5+q^7+q^8-q^{10}-q^{11}+\ddots=\cfrac{1}{1-q+\cfrac{(q^3)}{1-q^3+\cfrac{q^2(1+q)(1+q^3)}{1-q^5+\cfrac{q^3(1+q^2)(1+q^4)}{1-q^7+\cfrac{q^4(1+q^3)(1+q^5)}{1-q^9+\ddots}}}}}$$ The exponents on the left-hand side $${1,2,4,5,7,8,\ddots}$$ are positive integers not divisible by 3. How do we show that, the identity is true? Independent verification and attempts at proving/disproving the result are very welcome.

论坛问题帖 | has_answers | Nicco | 2015-08-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018558 | MSE-1402023

A relation related with odd perfect numbers

It is easy to prove, using the relation $\prod_{d\mid n}d=n^{\sigma_0(n)/2}$ holds for $n\geq 1$ where $\sigma_0(n)$ is the number of divisors, the following Proposition. The integer $n\geq 1$ is a perfect number if and only if $$2^{\sigma_0(n)}\cdot\left(\prod_{d\mid n}d\right)^2=\left(\sum_{d\mid n} d\right)^{\sigma_0(n)}$$ Thus it is easy to prove, noting the sum of divisors as $\sigma(n)$, that if $n\geq 1$ is an odd perfect number then $$\sigma\left(\sigma(n)^{\sigma_0(n)}\right)=\left(2^{\sigma_0(n)+1}-1\right)\cdot\sigma\left(\prod_{d\mid n}d^2\right)$$ My attempt was prove the converse statement, Question. Prove that if an integer $n\geq 1$ satisfies $$\sigma\left(\sigma(n)^{\sigma_0(n)}\right)=\left(2^{\sigma_0(n)+1}-1\right)\cdot\sigma\left(\prod_{d\mid n}d^2\right)$$ then n is odd and a perfect number (this is, an odd perfect number), or give a counterexample. Thanks in advance.

论坛问题帖 | has_answers | user243301 | 2015-08-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018559 | MSE-1405982

A false conjecture by Goldbach

In 1752 Goldbach send this conjecture to Euler: "Every odd integer can be written in the form $p+2a^2$ where p is a prime or 1 and a is a natural number (can be even 0)." This conjecture turned out to be false and my book asks me to prove that 5777 cannot be

written in such manner. What I did is simply noted that if exists such p it must be of the form $5777-2a^2$ and so a mustn't be greater than 53. Then I simply checked that for every values of a from 0 to 53 p is not prime. Btw this is a very tedious way to prove this and may in a test I wouldn't be able to do this so I was wondering if there was any shorter way? (or my book just wanted me to make a lot of calculations for some reason.)

论坛问题帖 | has_answers | AlienRem | 2015-08-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018560 | MSE-1406122

Conjectured value of a harmonic sum $\sum_{n=1}^\infty\left(H_n-\,2H_{2n}+H_{4n}\right)^2$

There is a known asymptotic expansion of harmonic numbers H_n for $n\to\infty$: $$\begin{align}H_n&=\gamma+\ln n+\sum_{k=1}^\infty\left(-\frac{B_k}{k\cdot n^k}\right)\\ &=\gamma+\ln n+\frac1{2n}-\frac1{12n^2}+\frac1{120n^4}-\frac1{252n^6}\,+\,\dots,\end{align}\tag1$$ where B_k are Bernoulli numbers. We can take a linear combination of harmonic numbers to cancel constant and logarithmic terms, compensate for $O(n^{-1})$ term, and get the following series that is possible to evaluate in a closed form (e.g. using generating function): $$\sum_{k=1}^\infty\left(H_n-\,2H_{2n}+H_{4n}-\frac1{8n}\right)=\frac18-\frac\pi{16}.\tag2$$ Rather than compensating for $O(n^{-1})$ term, we can take a series with alternating signs, that is also possible to evaluate in a closed form: $$\sum_{n=1}^\infty\,(-1)^n\,\Big(H_n-\,2H_{2n}+H_{4n}\Big)=\frac{3\pi}{16}-\frac\pi{4\sqrt2}-\frac{\ln2}8.\tag3$$ Generalizing, we can consider two families of series: $$\mathcal A_m=\sum_{n=1}^\infty\,(-1)^n\,\Big(H_n-\,2H_{2n}+H_{4n}\Big)^m,\tag4$$ $$\mathcal B_m=\sum_{n=1}^\infty\Big(H_n-\,2H_{2n}+H_{4n}\Big)^m,\tag5$$ and try to evaluate them in a closed form. So far I have the following conjectured result: $$\large\sum_{n=1}^\infty\Big(H_n-\,2H_{2n}+H_{4n}\Big)^2\stackrel{\normalsize\color{gray}?}=\frac\pi8-\frac\pi{16}\,\ln2-\frac{\pi^2}{96}+\frac3{16}\,\ln^22-\frac{G}{4},\tag{$\diamond$}$$ where G is the Catalan constant. Could you please help me to prove this result and, possibly, find other values of $\mathcal A_m,\mathcal B_m$?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2015-08-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018561 | MSE-1406400

A proof that $\frac{(2\phi)^n-(-1)^n}{\phi^{2n}-(-1)^n}\cdot\left(2^n-\phi^n\right)\cdot\sqrt5\in\mathbb Q$ for all $n\in\mathbb Z$

During computation of some series (with help of a CAS), at an intermediate step I encountered an expression, that after dropping non-essential parts looks like this:$$\mathcal E_n=\frac{(2\phi)^n-(-1)^n}{\phi^{2n}-(-1)^n}\cdot\left(2^n-\phi^n\right)\cdot\sqrt5$$ where $\phi=\frac{1+\sqrt5}2$ is the golden ratio (for $n=0$ take a limit and let $\mathcal E_0=0$). Although it is not immediately obvious from its form, it appears that $\mathcal E_n$ takes only rational values for all integer n. So far it is only an observation, and I am asking you to help me to find a proof. For example, it could be a recurrence relation with only rational coefficients.

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2015-08-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018562 | MSE-1407637

Conjecture on different type of triangle in a complete graph?

How many different triangles are there in K_5? The Answer is 35.(The Moscow Mathematics Puzzle) Then I asked what about K_6, K_7 and so on ...? With my intuition I arrived at this conjecture Conjecture: The number of different triangles in a complete graph of order n (K_n) is \begin{cases} \frac{n^2(n-2)^2(n-1)^2)}{6^2},&\mbox{ where } \binom{n}{3} \mbox{ is odd.}\\ \frac{(n^2-n)(n^2-4)(n^2-5n+12)}{12^2} ,&\mbox{where }\binom{n}{3} \mbox{is even.} \end{cases} What do you say about this?

论坛问题帖 | no_answers_at_retrieval | albo | 2015-08-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018563 | MSE-1412649

Can the determinant of an integer matrix with k given rows equal the gcd of the $k\times k$ minors of those rows?

I'm interested if the following is true: Let $n> k\geq1$ be integers, let $A\in\mathbb Z^{k\times n}$ and denote the $\binom nk$ $k\times k$ minors of A by $A_1,\ldots,A_N$. Then the equation $$\left|\begin{array}{c}A\\\hline X\end{array}\right|=\gcd(A_1,\ldots,A_N)$$ always has a solution $X\in\mathbb Z^{n-k\times n}$. This is a generalisation of Can the determinant of an integer matrix with a given row be any multiple of the gcd of that row? (which is the case $k=1$). For $\style{text-decoration:line-through}{k=n}$ and $k=n-1$ the above conjecture is also (obviously) true. Note that, as in the linked question, finding the minimal positive value of that determinant is equivalent to finding the minimal hypervolume of an n-dimensional simplex whose vertices are in an integer lattice and $k+1$ vertices are fixed. Idea: The case $\gcd(A_1,\ldots,A_N)=1$ may follow from Elementary proof that if A is a matrix map from $\mathbb{Z}^m$ to $\mathbb Z^n$, then the map is surjective iff the gcd of maximal minors is 1

论坛问题帖 | has_answers | Bart Michels | 2015-08-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018564 | MSE-141308

Primality test for specific class of $3\cdot 2^n+1$

I am interested in approaches which can be used to prove (disprove) conjecture given below . Let's define number N as : $N=3\cdot 2^n+1$ such that $n \equiv 2,6,8,10 \pmod{12}$ or $n \equiv 1 \pmod 4$. Let's define starting seed S as : $S = \begin{cases} 50542, & \text{if } n\equiv 8 \pmod {12} \\ 32672, & \text{if } n\equiv 1 \pmod 4 \\ 1692, & \text{if } n\equiv 2 \pmod {12} \\ 21868, & \text{if }n\equiv 6,10 \pmod {12} \end{cases}$ Let's define sequence S_i as : $S_i=S^2_{i-1}-2~$ with $~S_0=S$ Conjecture : $N ; (n > 2)~$ is a prime iff $S_{n-2} \equiv 0 \pmod N$ I have checked this claim for all exponents under 100000 from this list .

论坛问题帖 | no_answers_at_retrieval | Pedja | 2012-05-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018565 | MSE-1419912

A series summation

How should I calculate the following expression? \begin{equation} T\sum_{n=-\infty}^\infty\frac{1}{\sqrt{k^2+(2\pi nT)^2}}-\int_{-\infty}^\infty\frac{d\omega}{2\pi}\frac{1}{\sqrt{k^2+\omega^2}} \end{equation} Since neither of the two terms converges, so what this expression really means is \begin{equation} \lim_{N\rightarrow\infty}\left(T\sum_{n=-N}^N\frac{1}{\sqrt{k^2+(2\pi nT)^2}}-\int_{-2\pi NT}^{2\pi NT}\frac{d\omega}{2\pi}\frac{1}{\sqrt{k^2+\omega^2}} \right) \end{equation} The answer to the above expression must be a function of k/T, say it is $f\left(\frac{k}{T}\right)$. I think this function f has pretty good properties so that the following integral converges: \begin{equation} \int_0^\infty dk\cdot k\cdot f\left(\frac{k}{T}\right) \end{equation} I suspect the final answer to the above integral is $\pi T^2\ln\left(\frac{1+\sqrt{5}}{2}\right)$ up to a factor of 2.

论坛问题帖 | no_answers_at_retrieval | Mr. Gentleman | 2015-09-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018566 | MSE-1422179

Does this conjecture about prime numbers exist? If n is a prime, then there is exist at least one prime between n^2 and n^2+n.

I made an observation on prime numbers, want to check if any conjecture already exist or not? I am a computer programmer by profession and I am interested in number theory. As like many others I am intrigued by prime numbers. Based on my observation, I found following to be true If n is a prime, then there is exist at least one prime between n^2 and n^2+n I am not sure if this conjecture already exist? I tried searching in the internet but did not find any exact conjecture. I would like to know, first of all is my above statement is correct? if not, can any provide me with a counter example where it fails. If this statement is correct, does this conjecture already proposed by someone?

论坛问题帖 | has_answers | K.P. | 2015-09-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018567 | MSE-1423086

Understanding Erdös Discrepancy Conjecture $\left| \sum_{i=1}^k x_{i\cdot d} \right| > C$ for random series of -1 and 1

In today's newspaper NRC from The Netherlands a two-page article was dedicated to computer-aided mathematical proofs (sorry, paid link and in Dutch). Interesting in itself, but as an example it used the partial proof for divergence of the series consisting of a random distribution of -1 and 1. The Wikipedia lemma about Erdös conjecture shows this formula: $$\left| \sum_{i=1}^k x_{i\cdot d} \right| > C$$ But I have trouble understanding that in common English and applying it to the text of the article, which translates as: given an infinite sequence of -1 and 1, take a finite set of terms at equal distance, i.e. the first, third and fifth, or the fifth, tenth and fifteenth term. Add all those terms together. How far can they differ from 0? Erdös conjecture: infinitely far. My misunderstanding comes from combining infinite, random and finite. I think that if the sequence is random, then any given finite set can potentially exist of only 1 or only -1, meaning that the difference to zero is (potentially) equal to the amount of terms taken. The article further explains that Boris Konev and Alexei Lisitsa have proven that within the first 1161 terms, the discrepancy is always at least 3. But I can easily come up with a sequence that alternates on exactly the numbers taken, which would obviously result in a possible discrepancy of at most 1 for at least some series or slices. From the link, quote: The proof that the computer came up with proves, the two researchers claim, "that no sequence of length 1161 and discrepancy 2 exists." This seems contradictory, but no-one in the articles I found seem to wonder about this, and given that it requires so much research, I assume I simply misunderstand the problem. I assume I am misunderstanding the problem, but can anybody explain me where I am going wrong?

论坛问题帖 | has_answers | Abel | 2015-09-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018568 | MSE-1424600

Conjecture $\Re\,\operatorname{Li}_2\left(\frac12+\frac i6\right)=\frac{7\pi^2}{48}-\frac13\arctan^22-\frac16\arctan^23-\frac18\ln^2(\tfrac{18}5)$

I numerically discovered the following conjecture: $$\Re\,\operatorname{Li}_2\left(\frac12+\frac i6\right)\stackrel{\color{gray}?}=\frac{7\pi^2}{48}-\frac{\arctan^22}3-\frac{\arctan^23}6-\frac18\ln^2\!\left(\frac{18}5\right).$$ It holds numerically with a precision of more than 30000 decimal digits. Could you suggest any ideas how to prove it? Can we find a closed form for $\Im\,\operatorname{Li}_2\left(\frac12+\frac i6\right)$? Is there a general method to find closed forms of expressions of the form $\Re\,\operatorname{Li}_2(p+iq)$, $\Im\,\operatorname{Li}_2(p+iq)$ for $p,q\in\mathbb Q$?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2015-09-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018569 | MSE-1426061

A number-theory question on the deficiency function $2x - \sigma(x)$

Let $\sigma(x)$ be the sum of the divisors of a (positive) integer x. (For example, $\sigma(2) = 1 + 2 = 3$.) Define the deficiency function $D(x)$ to be the number $$D(x) = 2x - \sigma(x).$$ Let y be a (positive) integer. Now I compute the difference: $$D(xy) - D(x)D(y) = 2xy - \sigma(xy) - (2x - \sigma(x))(2y - \sigma(y))$$ $$=2xy - \sigma(xy) - 4xy + 2y\sigma(x) + 2x\sigma(y) - \sigma(x)\sigma(y)$$ $$=-2xy - 2\sigma(x)\sigma(y) + 2y\sigma(x) + 2x\sigma(y) + (\sigma(x)\sigma(y) - \sigma(xy))$$ $$=2(x - \sigma(x))(\sigma(y) - y) + (\sigma(x)\sigma(y) - \sigma(xy)).$$ This is because I want to compare $D(xy)$ and $D(x)D(y)$. Note that, in general we have $$2(x - \sigma(x))(\sigma(y) - y) \leq 0$$ and $$\sigma(x)\sigma(y) - \sigma(xy) \geq 0.$$ Lastly, observe that if we have $D(xy) = D(x)D(y)$, then it follows that $$2(\sigma(x) - x)(\sigma(y) - y) = \sigma(x)\sigma(y) - \sigma(xy).$$ My question is: Will it be possible to prove either $$D(xy) \leq D(x)D(y)$$ or

$$D(x)D(y) \leq D(xy),$$ subject to additional assumptions, if necessary? [Updated Sept. 8, 2015] If x and y are relatively prime (i.e., $\gcd(x, y) = 1$), then it follows that $D(xy) \leq D(x)D(y)$. By the contrapositive, we have the equivalent implication: "If $D(x)D(y) < D(xy)$, then $\gcd(x, y) > 1$." Finally, I conjecture that if the equation $D(xy) = D(x)D(y)$ is true, then either x or y is 1, or both x and y are powers of 2.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2015-09-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018570 | MSE-1428124

conjectured identity of the product of two theta functions

Looking into the discussion in this post, I was naturally led to consider the following general identity Given the two well known jacobi theta functions, namely $\theta_2(q)=\sum_{n=-\infty}^\infty q^{(n+1/2)^2}$ and $\theta_3(q)=\sum_{n=-\infty}^\infty q^{n^2}$ where $q=e^{2 \pi i\tau}$, $|q|\lt1$ It is then conjectured that the following identity is true $\frac{\cfrac{2\,q^{\frac{1}{2}}}{1-q^2+\cfrac{q^2(1-q^2)^2}{1-q^6+\cfrac{q^4(1-q^4)^2}{1-q^{10}+\cfrac{q^6(1-q^6)^2}{1-q^{14}+\ddots}}}}}{\cfrac{1}{1-q+\cfrac{q(1+q)^2}{1-q^3+\cfrac{q^2(1+q^2)^2}{1-q^5+\cfrac{q^3(1+q^3)^2}{1-q^7+\ddots}}}}}=\theta_2(q^2)\,\theta_3(q^2)$

论坛问题帖 | has_answers | Nicco | 2015-09-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018571 | MSE-1429201

a conjecture of certain q-continued fractions

Given the squared nome $q=e^{2i\pi\tau}$ with $|q|\lt1$, define, $$\begin{aligned}F(q)=\cfrac{1-q^2}{1-q^3+\cfrac{q^3(1-q)(1-q^5)}{1-q^9+\cfrac{q^6(1-q^4)(1-q^8)}{1-q^{15}+\cfrac{q^9(1-q^7)(1-q^{11})}{1-q^{21}+\ddots}}}}\overset{\color{red}{?}}=\prod_{n=1}^\infty\frac{(1-q^{12n-2})(1-q^{12n-10})}{(1-q^{12n-4})(1-q^{12n-8})}\\[1.5mm]&\end{aligned}$$ and $$\begin{aligned}G(q)=\cfrac{1-q^4}{1-q^3+\cfrac{q^3(1-\frac{1}{q})(1-q^7)}{1-q^9+\cfrac{q^6(1-q^2)(1-q^{10})}{1-q^{15}+\cfrac{q^9(1-q^5)(1-q^{13})}{1-q^{21}+\ddots}}}}\overset{\color{red}{?}}=\prod_{n=1}^\infty\frac{(1-q^{12n-4})(1-q^{12n-8})}{(1-q^{12n-2})(1-q^{12n-10})}\\[1.5mm]&\end{aligned}$$ Q: How do we prove rigorously that the two q-continued fractions are equal to the q-series? (if true, then the two q-continued fractions are reciprocals such that their product is unity, $F(q)\,G(q)=1$.)

论坛问题帖 | has_answers | Nicco | 2015-09-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018572 | MSE-1430169

Conjectured closed form for $\operatorname{Li}_2\!\left(\sqrt{2-\sqrt3}\cdot e^{i\pi/12}\right)$

There are few known closed form for values of the dilogarithm at specific points. Sometimes only the real part or only the imaginary part of the value is known, or a relation between several different values is known: [1][2][3][4][5][6]. Discovering a new identity of this sort is always of a great interest. I numerically discovered the following conjectured closed form and now am looking how to prove it: $$\operatorname{Li}_2\!\left(\sqrt{2-\sqrt3}\cdot e^{i\pi/12}\right)=\operatorname{Li}_2\!\left(\tfrac12+i\left(1-\tfrac{\sqrt3}2\right)\right)\stackrel{\color{gray}?}=\\\frac{23\pi^2}{288}-\frac18\ln^2\!\left(2+\sqrt3\right)+i\left[\frac{2\;\!G}3+\frac{\pi^2}{12\sqrt3}-\frac\pi{24}\ln\left(2+\sqrt3\right)-\frac1{8\sqrt3}\psi^{\small(1)}\!\left(\tfrac13\right)\right]$$ If you have got any ideas please share them. G is the Catalan constant, and $\psi^{\small(1)}(z)$ is the trigamma function. To respond questions in comments: my approach to find closed forms like this is to evaluate the expression with hundreds or sometimes thousands digits of precision and then use integer relation algorithms to find a matching linear combination with rational coefficients from a pool of candidate terms. The tricky part is that the pool cannot be too big, otherwise the algorithms begin to work extremely slowly or fail and produce false positives, so I need some intuition to select a restricted set of candidates that are likely to appear in a closed form I'm looking for. I usually select candidates from known closed forms of similar expressions, or from known closed forms of integrals that contain the expression in question along with other simpler terms (its "siblings"). Sometimes I also add some variations of existing terms, e.g. multiply them by a simple irrational coefficient like $\sqrt3$, or replace occurences of $\ln2$ with $\ln3$. When I find a matching linear combination, I verify it with much higher precision (usually tens or hundreds of thousand of decimal digits) to reduce chances that it is an accidental close but non-exact expression.

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2015-09-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018573 | MSE-1431200

the ratio of jacobi theta functions and a new conjectured q-continued fraction

Given the squared nome $q=e^{2i\pi\tau}$ with $|q|\lt1$, define $$\begin{aligned}H(q)=\cfrac{2(1+q^2)}{1-q+\cfrac{(1+q)(1+q^3)}{1-q^3+\cfrac{2q^2(1+q^4)}{1-q^5+\cfrac{q^3(1+q)(1+q^5)}{1-q^7+\cfrac{q^4(1+q^2)(1+q^6)}{1-q^9+\cfrac{q^5(1+q^3)(1+q^7)}{1-q^{11}+\ddots}}}}}\end{aligned}$$ Question: How do we show that $$H(q)\overset{\color{red}{?}}=\frac{2\,q^{1/2}\,\vartheta_3(0,q^2)}{\vartheta_2(0,q^2)}$$ where $\vartheta_n(0,q)$ are jacobi theta functions

论坛问题帖 | has_answers | Nicco | 2015-09-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018574 | MSE-1436130

I conjecture that there are infinitely many linear relations $p_n + p_{n + 3} = 2 p_{n + 2}$ in the sequence of primes!

Let $p_1 < p_2 < p_3 < \ldots < p_n < \ldots$ be the sequence of primes (with $p_1 := 2$ as usual). Conjecture: There are infinitely many positive integers n for which \begin{eqnarray} p_n + p_{n + 3} = 2 p_{n + 2}. \end{eqnarray} Or equivalently, There are infinitely many quadruplets of consecutive primes of the form $$(p, q, p + h, p + 2h)$$

论坛问题帖 | has_answers | dohmatob | 2015-09-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018575 | MSE-1446170

a conjectured continued fraction for $\tan\left(\frac{z\pi}{4z+2n}\right)$

Given a complex number $\begin{aligned}\frac{z}{n}=x+iy\end{aligned}$ and a gamma function $\Gamma(z)$ with $x\gt0$, it is conjectured that the following continued fraction for $\displaystyle\tan\left(\frac{z\pi}{4z+2n}\right)$ is true $$\begin{split}\displaystyle\tan\left(\frac{z\pi}{4z+2n}\right)&=\frac{\displaystyle\Gamma\left(\frac{z+n}{4z+2n}\right)\Gamma\left(\frac{3z+n}{4z+2n}\right)}{\displaystyle\Gamma\left(\frac{z}{4z+2n}\right)\Gamma\left(\frac{3z+2n}{4z+2n}\right)}\\&=\cfrac{2z}{2z+n+\cfrac{(n)(4z+n)}{3(2z+n)+\cfrac{(2z+2n)(6z+2n)}{5(2z+n)+\cfrac{(4z+3n)(8z+3n)}{7(2z+n)+\ddots}}}}\end{split}$$ Corollaries: By taking the limit(which follows after abel's theorem) $$ \begin{aligned}\lim_{z\to0}\frac{\displaystyle\tan\left(\frac{z\pi}{4z+2}\right)}{2z}=\frac{\pi}{4}\end{aligned}, $$ we recover the well known continued fraction for π $$\begin{aligned}\cfrac{4}{1+\cfrac{1^2}{3+\cfrac{2^2}{5+\cfrac{3^2}{7+\ddots}}}}=\pi\end{aligned}$$ If we let $z=1$ and $n=2$,then we have the square root of 2 $$\begin{aligned}{1+\cfrac{1}{2+\cfrac{1}{2+\cfrac{1}{2+\cfrac{1}{2+\ddots}}}}}=\sqrt{2}\end{aligned}$$ Q: How do we prove rigorously that the conjectured continued fraction is true and converges for all complex numbers z with $x\gt0$? Update:I initially defined the continued fraction $\displaystyle\tan\left(\frac{z\pi}{4z+2}\right)$ for only natural numbers,but as a matter of fact it holds for all complex numbers z with real part greater than zero.Moreover,this continued fraction is a special case of the general continued fraction found in this post.

论坛问题帖 | has_answers | Nicco | 2015-09-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018576 | MSE-1446449

about conjecture "every finite nonabelian p-group admits a noninner automorphism of order p"

by "The Kourovka Notebook. Unsolved Problems in Group Theory" there is a strong conjecture asserts that every finite nonabelian p-group admits a noninner automorphism of order p. What about finite abelian p-group? Is there a finite abelian p-group without noninner automorphism of order p? If yes can we for example by changing "order p" in conjecture make it for all finite p-group?

论坛问题帖 | has_answers | user148528 | 2015-09-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018577 | MSE-1446792

a conjectured continued-fraction for $\displaystyle\cot\left(\frac{z\pi}{4z+2n}\right)$ that leads to a new limit for π

Given a complex number $\begin{aligned}\frac{z}{n}=x+iy\end{aligned}$ and a gamma function $\Gamma(z)$ with $x\gt0$, it is conjectured that the following continued fraction for $\displaystyle\cot\left(\frac{z\pi}{4z+2n}\right)$ is true $$\begin{aligned}\displaystyle\cot\left(\frac{z\pi}{4z+2n}\right)=\frac{\displaystyle\Gamma\left(\frac{z}{4z+2n}\right)\Gamma\left(\frac{3z+2n}{4z+2n}\right)}{\displaystyle\Gamma\left(\frac{z+n}{4z+2n}\right)\Gamma\left(\frac{3z+n}{4z+2n}\right)}=\cfrac{2z+2n}{2z+n+\cfrac{(0z-n)(4z+3n)}{3(2z+n)+\cfrac{(2z+0n)(6z+4n)}{5(2z+n)+\cfrac{(4z+n)(8z+5n)}{7(2z+n)+\cfrac{(6z+2n)(10z+6n)}{9(2z+n)+\ddots}}}}}\end{aligned}$$ Or in gauss's notation $$\begin{aligned}\displaystyle\cot\left(\frac{z\pi}{4z+2n}\right)=-\frac{1}{2z+2n}\underset{m=0}{\overset{\infty}{\mathbf K}}\frac{((2m-2)z+(m-2)n)((2m+2)z+(m+2)n)}{((2m+1)(2z+n)}\end{aligned}$$ Corollaries: 1):let $z=1$ and $n=2$,then we obtain a beautiful continued fraction for square root 2 $$\begin{aligned}{-1+\cfrac{3}{2+\cfrac{\frac{(-1)(5)}{(1)(3)}}{2+\cfrac{\frac{(1)(7)}{(3)(5)}}{2+\cfrac{\frac{(3)(9)}{(5)(7)}}{2+\cfrac{\frac{(5)(11)}{(7)(9)}}{2+\ddots}}}}}=\sqrt{2}\end{aligned}$$ 2):However the most interesting case(for me at least),occurs when we take the limit to zero $$\begin{aligned}\lim_{z\to0}\cfrac{z(z+1)}{2z+1+\cfrac{(0z-1)(4z+3)}{3(2z+1)+\cfrac{(2z+0)(6z+4)}{5(2z+1)+\cfrac{(4z+1)(8z+5)}{7(2z+1)+\cfrac{(6z+2)(10z+6)}{9(2z+1)+\ddots}}}}}=\frac{1}{\pi}\end{aligned}$$ yielding a new limit for π from which one obtains the first few convergents $\begin{aligned}0,\frac{3}{8},\underline{\frac{5}{16}},\frac{15}{47},\frac{7}{22}},\frac{1365}{4288},\frac{3015}{9472},\frac{1575}{4948},\ddots\end{aligned}$. Where the underlined convergents appear in the stern-brocot tree for $\pi $ associated to its simple continued fraction. Q: Is the conjectured continued fraction true (for all complex numbers z with $x\gt0$)? Update:I initially defined the continued fraction $\displaystyle\cot\left(\frac{z\pi}{4z+2}\right)$ for only natural numbers,but as a matter of fact it holds for all complex numbers z with real part greater than zero.This continued fraction is one special case of the general continued fraction found here.

论坛问题帖 | has_answers | Nicco | 2015-09-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018578 | MSE-1454163

Why Navier-Stokes solutions are expected to be C^∞?

I am curious, why the Millennium problem about Navier-Stokes is about smooth (C^∞) not just C^1 solutions?

论坛问题帖 | no_answers_at_retrieval | porton | 2015-09-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018579 | MSE-1462239

Proving the existence of consecutive quadratic residues modulo $p>5$ by means of Pell's equation

When I read Existence of Consecutive Quadratic residues I thought we might be able to prove the existence of non-zero consecutive quadratic residues modulo a prime $p>5$ using the theory of Pell-type equations: if $d>1$ is a quadratic residue modulo p, then any solution (x,y) to $x^2-dy^2=1$ gives rise to consecutive quadratic residues (x^2,dy^2) modulo p. It seems a promising method to me; the only problem lies in making sure that x and y aren't divisible by p. Can we show that for any prime $p>5$, there exists a quadratic residue $d>1$ modulo p and a solution (x,y) to $x^2-dy^2=1$ s.t. $p\nmid x,y$?

论坛问题帖 | has_answers | Bart Michels | 2015-10-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018580 | MSE-1464339

Up to which value is Rassias' conjecture verified?

I came across this conjecture: Rassias' conjecture Up to which p has this conjecture be verified ? Are there intermediate results related to this conjecture ? The conjecture can be formulated in this way : For every prime $p>2$, there is a prime q, such that $(p-1)q-1$ is prime. The following PARI/GP-program shows, that for every p with $2<p<10^8$, there is a prime $q\le 1187$ doing the job. ? maxi=0;p=2;while(p<10^8,p=nextprime(p+1);q=1;gef=0;while(gef==0,q=nextprime(q+1);u=(p-1)*q-1;if(isprime(u)==1,gef=1;if(q>maxi,maxi=q;print(p," ",q," ",u))))) 3 2 3 19 3 53 73 5 359 109 11 1187 139 19 2621 179 23 4093 467 29 13513 1229 71 87187 2447 83 203017 2819 269 758041 8699 281 2444137 24419 311 7593997 57977 443 25683367 266117 503 133856347 374399 641 239989117 711089 653 464340463 4099493 719 2947534747 4313873 821 3541688911 8466209 881 7458729247 13187129 1031 13595928967 52172843 1187 61929163453 ? Update : The conjectrue remains true for $2<p<10^9$ and a prime $q\le 1709$ does the job in that range. ? while(p<10^9,p=nextprime(p+1);q=1;gef=0;while(gef==0,q=nextprime(q+1);u=(p-1)*q-1;if(isprime(u)==1,gef=1;if(q>maxi,maxi=q;print(p," ",q," ",u))))) 103784147 1193 123814486177 112564253 1361 153199946971 174454277 1367 238478995291 280721033 1511 424169479351 623020787 1613 1004932527817 855177437 1697 1451236108891 929408729 1709 1588359516151 ?

论坛问题帖 | no_answers_at_retrieval | Peter | 2015-10-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018581 | MSE-1475399

Does the following equation have any solutions in $\mathbb{N}$?

Let $\mathbb{N}$ be the set of positive integers. The function $\sigma(N)$ gives the sum of the divisors of N. My question is: Does the following equation have any solutions for $x \in \mathbb{N}$? $$\sigma(3x + 1) = 4x + 1$$ Notice that $3x + 1$ must be deficient, and that if we allow $x = 0$, then it is a (trivial) solution. Furthermore, suppose that $3x + 1 = p^{\alpha}$, where p is prime. Then we have $$\sigma(3x + 1) = \frac{p^{\alpha+1} - 1}{p - 1} = \frac{4(p^{\alpha} - 1)}{3} + 1 = 4x + 1,$$ from which we obtain $$3(p^{\alpha+1} - 1) = 4(p - 1)(p^{\alpha} - 1) + 3(p - 1).$$ Simplifying and collecting like terms, we get $$0 = p^{\alpha+1} - 4p^{\alpha} + p + 4.$$ Rewriting the last equation, we have $$(p^{\alpha} + 1)(4 - p) = 8.$$ As $p^{\alpha} + 1 \geq 3$, we have the possibilities $$(p^{\alpha} + 1, 4 - p) \in \{(4,2),(8,1)\}.$$ Both of them are impossible under the given constraints. Consequently, $\omega(3x + 1) \geq 2$, where $\omega(y)$ is the number of distinct prime factors of y. (That is, $3x + 1$ must be composite.) Lastly, I tried checking for equality (in the range $0 \leq x \leq 100$) using a spreadsheet, and found only the solution for $x=0$. I am therefore compelled to predict that: The equation $\sigma(3x + 1) = 4x + 1$ does not have any solutions for $x \in \mathbb{N}$.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2015-10-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018582 | MSE-1491378

Number of Magic squares and their applications

A magic square is a square array of numbers consisting of the distinct positive integers $1, 2, ..., n^2$ arranged such that the sum of the n numbers in any horizontal, vertical, or main diagonal line is always the same number I was wondering if the numbers of magic squares of order n is still a conjecture (For a general n) ? And I have some elementary questions regarding the magic squares. (1) What is the smallest n for a magic square ? I guess for $n=1$ we just have one 1×1 matrix $[1]$ (2)And for $n=2$, They don't exists !! (right) (3)For $n=3$ we have 8 different magic squares and so on. Now my question is, after $n=3,4$ The numbers of magic squares grows rapidly, and I want to know if there is a systematic way to get those numbers or does it remain to be a conjecture ? Is there some interesting applications where those magic squares are used ? I have a project to do on combinatorics and I was thinking of considering this (what do you guys think ?) It will be interesting to try some code and algorithms on it too. I would also love if someone knows a good resource or papers on those magic squares.

论坛问题帖 | no_answers_at_retrieval | alkabary | 2015-10-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018583 | MSE-1513841

4 Divides x Proofs of conjectures

Hi there I'm working on a set of problems and I'm having some difficulty proving and disproving these examples. I know that #1 is essentially (There exists K where [x=4k]) I'm lost after that. I'm not sure about 2 or 3 I believe that 2 is false and 3 is true but I don't know how to prove or disprove formally. Any advice would be wonderful thank you! Define what it means for a number to be divisible by 4, that is, the concept 4 | x. Answer: There exists k where [x=4k] Determine whether the following conjecture is true or false. If it is true, prove it, otherwise construct a counterexample to disprove it. Conjecture 1. Let x and y be arbitrary integers. If 4 | (x-y), then 4 | (3x+y). Determine whether the following conjecture is true or false. If it is true, prove it, otherwise construct a counterexample to disprove it. Conjecture 2. For integers x and y, if 4 | x and 8 | y, then 8 | (x+y).

论坛问题帖 | has_answers | mm19 | 2015-11-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018584 | MSE-1514255

Iterated $\sigma^k(n)$, excluding the possibility that primes $p|\sigma^t(n)$, $t< k$ divides an odd perfect number n

Let $m\geq 1$ an integer and $\sigma(m)=\sum_{d|m}$ the sum of positive divisors function. A positive integer is said to be perfect if and only if $\sigma(n)=2n$. We have that $\sigma(m)$ is a multiplicative function (see this site or [1]), thus we have the known Fact. If n is an odd perfect number then $\sigma(\sigma(n))=6n$. Proof. We don't know if there are odd perfect numbers, is an unsolved problem, but assuming the existence then $\sigma(2n)=\sigma(2)\cdot\sigma(n)=3\cdot 2n=6n$, since $gcd(2,n)=1$, and this is the end of the proof. Hypothesis to work. In this post we assume that $n=n_1$ is the first odd perfect number, if there is one. Now we define iterated of the sum of divisor function, as is ususal $$\sigma^k(m)=\sigma(\sigma^{k-1}(m))=\sum_{b|\sigma^{k-1}(m)}b$$ for integers $k>1$ and $\sigma^1(m)=\sigma(m)$. Hypothetical algorithm. And I compute terms of the sequence $$\sigma^k(n)$$ always excluding the possibility that an prime previously computed in the factorization of all previous terms of the sequence $\sigma^t(n)$, are $t<k$, divides n. (Please if you want explain my idea in a better sense or words, too the question bellow feel free to edit, thanks.) Thus my Question. For the first odd perfect number $n=n_1$ where we can find/estimate the barrier of using this algorithm? Thanks in advance. I say, $\sigma^3(n)=8\cdot 3\cdot n$ and I assume to add as hypothesis that $3\nmid n$. Now I compute using that the sum of positive divisor function is multiplicative another time... $\sigma^4(n)=\frac{2^4-1}{2-1}\cdot 4\cdot 2n=8\cdot 3\cdot 5\cdot n$. Well Tochard theorem say that perhaps is possible that there are odd perfect numbers of the form $12\lambda+1$, but perhaps I cannot assume yet that $5\nmid n$ (looking the barrier yes I assume and add the hypothesis that $5\nmid n$). Going to the behaviour of the sum of divisor function, and the fact that if n_1 exists, then can I apply the Fundamental Theorem of the Arithmetic and Euler's Theorem for odd perfect numbers to bound the barrier of use in previous algorithm? References: [1] Wikipedia pages for Divisor function and Odd Perfect Numbers.

论坛问题帖 | has_answers | user243301 | 2015-11-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018585 | MSE-1514471

a conjecture of two equivalent q-continued fractions related to the reciprocal of the Göllnitz-Gordon continued fraction A111374-OEIS

Given the square of the nome $q=e^{2i\pi\tau}$ and ramanujan theta function $f(a,b)=\sum_{k=-\infty}^{\infty}a^{k(k+1)/2}b^{k(k-1)/2}$ with $|q|\lt1$, define, $$\begin{aligned}M(q)=\cfrac{1-q^3}{1-q^2+\cfrac{q^2(1-q^{-1})(1-q^5)}{1-q^6+\cfrac{q^4(1-q)(1-q^7)}{1-q^{10}+\cfrac{q^6(1-q^3)(1-q^{9})}{1-q^{14}+\ddots}}}}\overset{\color{blue}{?}}=\prod_{n=1}^\infty\frac{(1-q^{8n-3})(1-q^{8n-5})}{(1-q^{8n-1})(1-q^{8n-7})}\\[1.5mm]&\end{aligned}$$ $$=\frac{f(-q^3,-q^5)}{f(-q,-q^7)}$$ and $$\begin{aligned}N(q)=\cfrac{1-q^3}{1+q^2-\cfrac{q^2(1+q^{-1})(1+q^5)}{1+q^6+\cfrac{q^4(1-q)(1-q^{7})}{1+q^{10}-\cfrac{q^6(1+q^3)(1+q^{9})}{1+q^{14}+\ddots}}}}\overset{\color{blue}{?}}=\prod_{n=1}^\infty\frac{(1-q^{8n-3})(1-q^{8n-5})}{(1-q^{8n-1})(1-q^{8n-7})}\\[1.5mm]&\end{aligned}$$ $$=\frac{f(-q^3,-q^5)}{f(-q,-q^7)}$$ Q: How do we prove that the two q-continued fractions are equal to the q-products such that the continued fractions are equivalent $M(q)=N(q)$?

论坛问题帖 | has_answers | Nicco | 2015-11-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018586 | MSE-1515684

Is it true/known/important that $(\log p_n)/n$ is nonincreasing, where p_n is the nth odd prime number?

First thing first, I would like to apologize in advance for my poor knowledge of Maths and English. I'm an Italian student and after asking to all the mathematicians and Maths teachers in my town, I hope that the internet might help me regarding the following conjecture. Conjecture: If p_{n} is the nth odd prime, then $$ \frac{\log_{10}p_{n}}{n} \geq \frac{\log_{10}p_{n+1}}{n+1}. $$ To date, I've tested it for the first 64k prime numbers. My questions are: -Does it deserve to be shown to other people/tried to prove, or is it just "one among thousands"? -If it's important, who could I show it to? -If it isn't important, what makes some conjectures noteworthy?

论坛问题帖 | has_answers | samuelemarro | 2015-11-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018587 | MSE-1522550

Is $p(p + 1)$ always a friendly number for p a prime number?

Let $\sigma(x)$ denote the sum of the divisors of x. We call the ratio $I(x) = \sigma(x)/x$ as the abundancy index of x. A positive integer N is friendly if there exists a positive integer $M \neq N$ such that $I(M) = I(N)$. My question is the following: Is $Z = p(p + 1) = p\sigma(p)$ always a friendly number for p a prime number? Note that $$\gcd(Z, \sigma(Z)) = \gcd(p(p + 1), (p + 1)\sigma(p + 1)) \geq p + 1 \geq 3,$$ so that Greening's Theorem fails to establish solitude. OEIS sequence A074902 lists the following as friendly: $$6, 12, 30, 56, 132$$ but does not list 182 and 306, for example.

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2015-11-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018588 | MSE-1522619

Why is proving that 10 is solitary considered very difficult?

The title says it all. We denote the sum of the divisors of x by $\sigma(x)$. The ratio $I(x)=\sigma(x)/x$ is called the abundancy index of x. If $I(m)=I(n)$, then $\{m,n\}$ is called a friendly pair. A number which is not a member of a friendly pair is called solitary. So: Why is proving that 10 is solitary considered very difficult? Is there no (known) partial converse to Greening's Theorem (i.e., $\gcd(N, \sigma(N)) = 1 \implies N$ is solitary)? What is the main obstruction? OEIS sequence A014567 contains the following statement from Dean Hickerson: It is easy to show that if N and $\sigma(N)$ are relatively prime then N is solitary. But the converse is not true; for example, 18, 45, 48 and 52 are solitary.

Probably also 10, 14, 15, 20, 22 and many others are solitary, but I do not think that will ever be proved. Update (November 11, 2015): I am aware of the paper Does Ten Have a Friend?, which gives some necessary conditions for a number to be a friend of 10. This question is essentially an offshoot of the following earlier MSE posts: Can distinct odd perfect numbers $N = {p^k}{m^2}$ share the same Euler factor p^k? If $\sigma(N) = aN + b$, where $\gcd(a, b) = 1$, does it follow that at least one of N's factors is solitary? Is $p(p + 1)$ always a friendly number for p a prime number?

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2015-11-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018589 | MSE-1522912

If $\gcd(Z,\sigma(Z))=1$ and $1<N=Z\sigma(Z)$, is N always friendly?

This question is a generalization / offshoot of this earlier MSE post: If $\gcd(Z,\sigma(Z))=1$ and $1<N=Z\sigma(Z)$, is N always friendly? Here, $\gcd(a,b)$ is the greatest common divisor of a and b, and $\sigma(x)$ is the sum of the divisors of x. Note that $$\gcd(N ,\sigma(N))=\gcd(Z\sigma(Z),\sigma(Z\sigma(Z)))=\gcd(Z\sigma(Z),\sigma(Z)\sigma(\sigma(Z))) \geq \sigma(Z) > 1$$ where the last inequality follows from $1<N=Z\sigma(Z)$, so that Greening's Theorem fails to establish that N is solitary. (However, this does not conclusively prove that N is friendly.) For an example: Consider $Z = 2^{p-1}$ where p and $2^p - 1$ are primes. There does not seem to be a sequence in OEIS for Numbers n such that $\gcd(n,\sigma(n))=1$ and $n\sigma(n)$ is friendly.

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2015-11-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018590 | MSE-1544221

Twin primes sums conjecture

I have found an interesting conjecture between twin primes sums. I don't know if it is already described by someone else. I have checked in internet, but I didn't find any mention of such conjecture. Could you please tell me if the following conjecture is well known for mathematicians or not? Let's take twin prime pair 11 and 13 and do the sum: 11+13=24 Now find out sums of two prime numbers which gives the result 24. 5+19=24 7+17=24 You can see that sums 5+19 and 7+17 contain twin primes 5, 7 and 17, 19. Sum of twin primes 11 + 13 can be written by sum of combinations of twin primes 5, 7 and 17, 19. Another example: Let's take sum of twin prime pair 17, 19: 17+19=36 Its sum can be written with using of other twin primes 5, 7 and 29, 31 as: 5+31=36 7+29=36 We can define twin prime sums conjecture as follow. Let’s p and q are twin prime numbers; q=p+GAP. For each p+q = a+b = c+d. Where a, b, c, d are prime numbers, where a, c and b, d are prime pairs where c=a+GAP ; d=b-GAP. For twin primes sums conjecture GAP is equal to 2. Twin prime sums conjecture doesn’t work for prime pairs p, q equals (3, 5), (5, 7), (197, 199), (347, 349). I have checked the first 100 000 twin primes. The largest checked twin prime pair was 18409199 + 18409201 = 1019 + 36817381 = 1021 + 36817379. It has been found that there are exceptions of p, q for which this conjecture doesn’t work. For sums 3+5; 5+7; 197+199; 347+ 349 it doesn’t exist a, b and c, d which fulfill conjecture. If twin prime sum conjecture is true then there are unlimited twin prime pairs, because it shows that each twin prime pair sum can be written by sum of combination of smaller and larger twin prime pairs. For twin primes gap between p,q is equal to 2. I have found that similar conjecture works also for larger even gaps. Example for GAP equals 6 23+29=52 We can see that GAP between primes 23 and 29 is equal to 6. 29 - 23 = 6. Now we will find sums of other primes which have equal sum to 52. 5+47=52 11+41=52 We have combination of primes 5, 11 and 41, 47. We can see that GAP between those primes is equal to 6, (11 - 5 = 47 - 41 = 6) and is same as the GAP between primes 29 and 23 . I have also checked this generalized conjecture for primes below 100 000 and even gaps below 100. All of the gaps have few exception pairs p, q for which conjecture doesn’t work, but in major amount of cases conjecture works and for sum of primes with gap we can find another two sums of prime pairs with same gap.

论坛问题帖 | no_answers_at_retrieval | Ivan Sas | 2015-11-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018591 | MSE-1545993

With $s(n)=\sum_{k=1}^n n \bmod k$, can be justified that $\forall\epsilon>0$ let us $\lim_{n\to\infty}\frac{s(n-1)}{\epsilon+s(n)}=1?$

Denoting as $$s(n)=\sum_{k=1}^n n \bmod k$$ the sum of remainders function (each remainder is defined as in the euclidean division of integers $n\geq 1$ and k). See [1] for example. For examples $s(1)=0$ and $$s(6)=0+6\bmod4+6\bmod5+0=2+1=3.$$ After some computational experiments I am interesting in Question. Can you prove or refute (too are welcome heuristics) that $\forall\epsilon>0$ $$\lim_{n\to\infty}\frac{s(n-1)}{\epsilon+s(n)}=1?$$ Similarly that $\forall\epsilon>0$ $$\lim_{n\to\infty}\frac{s(n)}{\epsilon+s(n-1)}=1?$$ Thanks in advance, it seems curious, and I believe that previous is a strong condition in the sense that its proof could be easy, or well in the sense that find a counterexample could be easy (I am refering to the presence of such ϵ). I don't know if previous exercises are in the literature. There are formulas involving this function (see [1] for example) with the sum of divisors function $\sigma(n)=\sum_{d\mid n}d$, that holds for each integer $n>1$ $$\sigma(n)+s(n)=s(n-1)+2n-1,$$ and a companion formula that can be deduced inductively from previous (see too in [1]). Well my experiements aren't the bests, neither I have enough resolution to see the limit, but there is a difference when I plot $\frac{s(n)}{s(n-1)}$ ($n\geq 5$) or $\frac{\sigma(n)}{\sigma(n-1)}$ or $\frac{\phi(n)}{\phi(n-1)}$, where this last is Euler's totient function. References: [1] Math Overflow, How to calculate the sum of remainders of N? https://mathoverflow.net/questions/195325/how-to-calculate-the-sum-of-remainders-of-n I've read the notes from two authors Spivey and Cross.

论坛问题帖 | has_answers | user243301 | 2015-11-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018592 | MSE-1560968

Among any $2n$ consecutive integers below n^2+2n at least one has no prime divisor less than n?

What is your idea about my conjecture? Consider a sequence of $2n$ consecutive natural numbers, all the terms less than $n^2 + 2 n$. Then there exists at least one number in the sequence which is not divisible by any prime less than or equal to n.

论坛问题帖 | has_answers | some one | 2015-12-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018593 | MSE-158293

Density of odd numbers in a sequence relating base 2 and base 3 expansion

Define the function $$f(4n)=6n+1\\ f(4n+1)=6n+2\\ f(4n+2)=6n+3\\ f(4n+3)=6n+5$$ and the sequence $u_0=2$, $u_{k+1}=f(u_k)$. Let $d_1\le d_2$ be the lower and upper asymptotic density of odd numbers in u_k. Since $f(n)$ is odd for even n, no two consecutive terms can be even so obviously $d_1\ge 1/2$. Experimentally, it seems reasonable to conjecture $d=d_1=d_2=2/3$. Heuristically, when n is odd, $f(n)$ has "probability 1/2" to be even, so we get this Markov chain: which is consistent with $d=2/3$. But can we prove rigorously that the heuristic works, that is that u_k is "random enough" for this to be true? Failing that, a sharper lower bound on d_1 could still be an interesting result. (One approach could be to let $a_k=1$ if u_k is even and 0 otherwise, and examine the probabilities of the p-bit subwords $(a_k,\dots,a_{k+p-1})$. Unfortunately every Fibonacci word, that is one not containing the pattern 11, seems to appear infinitely frequently in a_k. This is consistent with the Markov chain model.) PS: If you're wondering, the problem arose from this question. Response to comments: Someone suggested to work modulo some larger number, such as 12. The problem with this approach is that if the input is considered mod $2^a 3^b$, the output will only be known mod $2^{a-1} 3^{b+1}$ so you're not going to be able to say much about the behavior of f when iterating more than a times. And it seems that $010101\dots$ can always appear as a subword of a_k, so you won't be able to prove anything non-trivial about the density by proving something about the density after k iterations starting from an arbitrary state (that is, by examining f^k for bounded k). The first few terms are 2, 3, 5, 8, 13, 20, 31, 47, 71, 107, 161, 242. The sequence grows as $\Theta((3/2)^n)$. Interestingly, notice how closely the first few terms match the Fibonacci sequence (which can be explained by how close $3/2$ is to the golden ratio and by the fact that modulo 2, neither sequence contains the pattern 00).

论坛问题帖 | has_answers | Generic Human | 2012-06-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018594 | MSE-1583609

The four runner problem/conjecture

I've recently read here the following problem, called « four-runner problem » : Suppose four runners (represented by labeled spheres) run around a circular track. Their speeds are constant positive rationals $v_1<v_2<v_3<v_4$. At time 0, all the runners are together at the starting gate; shortly after, they are arranged in the order $\{1, 2, 3, 4\}$, counting from the starting gate. The order changes to $\{4, 1, 2, 3\}$ when runner 4 passes the starting gate. Then either runner 3 passes the starting gate, runner 4 laps runner 1, or the two events happen at the same time, giving possible orders $\{3, 4, 1, 2\}$, $\{1, 4, 2, 3\}$, or $\{3, 1, 2, 4\}$. [...] The answer to this question is unknown: which rates generate all 24 possible orders? For three runners, all six orders are achieved if and only if the two slower rates do not add up to the fast rate. For instance, I found that the rates $0.1 < 0.2 < 0.61 < 0.9$ generate the 24 possible combinations. I would like to know where to find more information on that problem. Are there some research articles about it? I didn't find anything, even for the case with 3 runners (which does not seem obvious to me). Thank you in advance!

论坛问题帖 | has_answers | Watson | 2015-12-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018595 | MSE-1591741

A conjecture about the prime counting function

Using this lemma it can be proved that $\Delta(m,n)=\pi(m\cdot n)-\pi(m)\cdot\pi(n)+1$ (where π is the prime counting function) is a function $\Delta:\mathbb N\times\mathbb N\to\mathbb N$. Reformulated conjecture: Given $m\in\mathbb N,m>2$, then $n\in \mathbb N$ is an odd prime if $\Delta(m,n-1)>\Delta(m,n)<\Delta(m,n+1)$ and if n is an odd prime then $\Delta(m,n-1)\ge\Delta(m,n)\le\Delta(m,n+1)$. The conjecture is tested for $m,n<3000$. Counterexample: 5879 is a prime, but $\Delta(5878,5879)=1525672>\Delta(5878,5878)=1523414$

论坛问题帖 | has_answers | Lehs | 2015-12-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018596 | MSE-1595838

Are all totient values of Fibonacci Numbers distinct?

This question was inspired while I was seeing how certain recurrence relations would behave when I applied Multiplative Functions. Let F_{n} be a sequence for which $F_{1}=1,F_{2}=1$, and $F_{n}=F_{n-1}+F_{n-2}$. For any $n,m>8$, is it always true that $\phi(F_n)$ is not equal to $\phi(F_m)$, when ϕ is the Euler Totient Function? A case when n divides m appeared be rather simple. $n|m$ would imply $F_{n}|F_{m}$. Then let F_{m} be F_{n} multiplied by k. $\phi(F_{m}) \ge \phi(F_{n})\phi(k)$. $F_{n+1}=F_{n}+F_{n-1}<2F_{n}$, and $F_{n+2}=2F_{n}+F_{n-1}>2F_{n}$. Thus k may never be equal to 2. Therefore, $\phi(k)$ will always be greater than 1 implying $\phi(F_{m})=\phi(k)\phi(F_{n})>\phi(F_{n})$. Therefore, for the case when n divides m, $\phi(F_m)$ is always not equal to $\phi(F_n)$. However, other cases proved to be rather difficult, such as proving $\phi(F_n) \neq \phi (F_{n+1})$. Can anybody prove or disprove this question? Any Help would be greatly appreciated.

论坛问题帖 | has_answers | Chad Shin | 2016-01-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018597 | MSE-1598092

Conjecture $\sum_{m=1}^\infty\frac{y_{n+1,m}y_{n,k}}{[y_{n+1,m}-y_{n,k}]^3}\overset{?}=\frac{n+1}{8}$, where $y_{n,k}=(\text{BesselJZero[n,k]})^2$

While solving a quantum mechanics problem using perturbation theory I encountered the following sum $$ S_{0,1}=\sum_{m=1}^\infty\frac{y_{1,m}y_{0,1}}{[y_{1,m}-y_{0,1}]^3}, $$ where $y_{n,k}=\left(\text{BesselJZero[n,k]}\right)^2$ is square of the k-th zero of Bessel function J_n of the first kind. Numerical calculation using Mathematica showed that $S_{0,1}\approx 0.1250000$. Although I couldn't verify this with higher precision I found some other cases where analogous sums are close to rational numbers. Specifically, after some experimentation I found that the sums $$ S_{n,k}=\sum_{m=1}^\infty\frac{y_{n+1,m}y_{n,k}}{[y_{n+1,m}-y_{n,k}]^3} $$ are independent of k and have rational values for integer n, and made the following conjecture $\bf{Conjecture:}\ $ for $k=1,2,3,...$ and arbitrary $n\geq 0$ $$\sum_{m=1}^\infty\frac{y_{n+1,m}y_{n,k}}{[y_{n+1,m}-y_{n,k}]^3}\overset{?}=\frac{n+1}{8},\\ \text{where}\ y_{n,k}=\left(\text{BesselJZero[n,k]}\right)^2. $$ How one can prove it? It seems this conjecture is correct also for negative values of n. For example for $n=-\frac{1}{2}$ one has $y_{\frac{1}{2},m}=\pi^2 m^2$, $y_{-\frac{1}{2},k}=\pi^2 \left(k-\frac{1}{2}\right)^2$ and the conjecture becomes (see Claude Leibovici's answer for more details) $$ \sum_{m=1}^\infty\frac{m^2\left(k-\frac{1}{2}\right)^2}{\left(m^2-\left(k-\frac{1}{2}\right)^2\right)^3}=\frac{\pi^2}{16}. $$

论坛问题帖 | has_answers | user294724 | 2016-01-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018598 | MSE-1602212

Conjecture for the maximum number of rooms.

The puzzle I have is essentially this, but for n rows.For this instance of $n=5$, quick tallying reveals the answer to be 21. For $n=4$, it is 13. For $n=3$, it is 7. For $n=2$ it is 3. And for $n=1$, it is obviously 1. I conjectured a formula to calculate the maximum number of rooms: $n^2 - n +1$. I have tested this formula upto $n=7$ and it works. (I don't want to test anymore, too lazy) Now it is my understanding that all conjectures must be proven, but I don't know how to prove mine. I'm thinking induction, but as I haven't formally learnt it yet, I'm not well versed in induction. I'd appreciate it if you could show me a proof. It need not be induction. Thanks :)

论坛问题帖 | has_answers | Airdish | 2016-01-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018599 | MSE-1602312

Sum and product of k real numbers > 0 is unique?

Can we prove that $\sum_{i=1}^k x_i$ and $\prod_{i=1}^k x_i$ is unique for $x_i \in R > 0$? I stated that conjecture to solve CS task, but I do not know how to prove it (or stop using it if it is false assumption). Is the pair of Sum and Product of array of k real numbers > 0 unique per array? I do not want to find underlying numbers, or reverse this pair into factorisation. I just assumed that pair is unique per given set to compare two arrays of equal length to determine whether numbers inside with possible multiplicities are the same. The order in array does not matter. 0 is discarded to give meaningful sum and product. At the input I get two arrays of equal length, one of which is sorted, and I want to determine if the elements with multiplicities are the same, but I do not want to sort the second array (this will take more than linear time, which is the objective). So I thought to make checksums, sum and product seemed unique to me, hence the question. For example: I have two arrays: v=[1,3,5,5,8,9.6] and the second g=[9.6,5,3,1,5,8]. I calculate sum = 31.6, and product = 5760. For both arrays, since one is sorted version of another. And now I calculate this for b=[1,4,4,5,8,9.6], sum is 31.6 but product is 6144. So I assumed that if both sum and product are not the same for two given arrays, than the element differs. Getting back to question, I am thinking that the pair {sum, product} is the same for all permutations of array (which is desired), but will change when elements are different (which is maybe wishfull thinking).

论坛问题帖 | has_answers | Evil | 2016-01-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018600 | MSE-1606340

Recreational conjecture on factoring groups

Consider the following: For a group G with identity e, define $s: G \to \mathbb{N} \cup \{ \infty \}$ by $s(g) = \min \{ k \in \mathbb{N} : g^{k} = e \}$, where $ \min \emptyset = \infty$. Moreover, let $\Theta(G) = \sup \{ s(g) : g \in G \}$. Refer to G as bounded if there exists $N \in \mathbb{N}$ such that $g^{N} = e$ for all $g \in G$. We see that G is bounded iff $\Theta(G) < \infty$. If $\Theta(G) $ is finite, then $g^{\Theta(G) !} = e$; if $\Theta(G) = \infty$, then for every N there exists $g \in G$ for which $s(g) > N$, so $g^{N} \neq e$. Moreover, any group of order n is bounded. To see this, the claim is made that $s(g) \leq n$ for all $g \in G$. To see this, consider the set $ \{ g, g^{2}, \ldots, g^{n + 1} \}$. By pigeonhole, there exist $1 \leq i < j \leq n + 1$ such that $g^{i} = g^{j}$, so $g^{j - i} = e$; then $s(g) \leq j - i \leq n$. Thus $\Theta(G) \leq n$. Initially, I conjectured that a set was bounded off it was finite, considering the multiplicative group $\{ e^{iq} : q \in \mathbb{Q} \}$, where $s(g)$ is always finite, but unbounded. Then I dismissed this considering $G = \mathbb{Z}_{2}^{I}$, where I is an infinite indexing set and $\Theta(G) = 2$. I revised this to the claim that a group G is bounded if and if it can be written as a product $G = \prod_{i \in I} H_{i}$ of finite groups for some index set I, where $\sup \{ \# H_{i} : i \in I \} < \infty$. I believe I know how to show any factoring of G into finite groups would satisfy this, but I can't show that every bounded group is factorable into finite groups, i.e. can't show there doesn't exist a bounded group which cannot be expressed as a product of finite groups. Is this correct? If so, how might I show it? If not, is there a counterexample? Thanks.

论坛问题帖 | has_answers | AJY | 2016-01-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018601 | MSE-1609840

Integral $\int_0^\infty\Big[\log\left(1+x^2\right)-\psi\left(1+x^2\right)\Big]dx$

I found this intriguing integral: $$\int_0^\infty\Big[\log\left(1+x^2\right)-\psi\left(1+x^2\right)\Big]dx\approx0.84767315533332877726581...$$ where $\psi(z)=\partial_z\log\Gamma(z)$ is the digamma. A lookup in ISC+ (in "advanced" mode) returned a possible closed form $\frac\pi2\Big[\zeta\!\left(\frac12\right)+2\Big]$ that seems to agree with the numeric value of the integral when calculated to a higher precision. Any ideas how to prove it?

论坛问题帖 | has_answers | july | 2016-01-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018602 | MSE-1611869

Fibonacci Numbers and the Harmonic Series

$$\sum_{k=1}^{n} \frac{1}{k}=H_n=\frac{p_n}{q_n}$$ Where p_n,q_n are coprime intergers. The first few values for p_n+q_n are $2,5,7,37,197,69,504,1041,9649$. When are p_n+q_n Fibonacci numbers? Due to the inequality, $\ln(n+1)<H_n<\ln(n)+1$, $\ln(n+1)q_n<p_n<(\ln(n)+1)q_n$. Therefore, this would imply that $(\ln(n+1)+1)q_n<p_n+q_n<(\ln(n)+2)q_n$. However, $\frac{\ln(n)+2}{\ln(n+1)+1}<\phi$ where ϕ is the golden ratio. Though this is sloppy logic, because of this I thought it was likely are few cases where p_n+q_n are Fibonacci numbers. The only solutions appeared to be $2,5$. Any help would be appreciated.

论坛问题帖 | no_answers_at_retrieval | Chad Shin | 2016-01-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018603 | MSE-1619365

Is there a typo in the formula or does my GAP-package sglppow fail?

This site deals with group formulas for prime powers p^k for $k\le 7$. The formula for $k=7$ seems to be wrong. I compared the results with GAP and the formula is off by 2453 for $p=13,17,19,23$ and 29. Moreover the lcm of the numbers $3,4,5,7,8,9$ is not 360, but 2520 and as I understood the column "modulus relative ...", the value 2520 should be right. The values for $p=7$ and $p=11$ coincide with the given formula. Does anyone know the correct formula ? Does GAP give the wrong values, or is the given formula wrong ? The formula given in the link is $$gnu(p^7)=3p^5+12p^4+44p^3+170p^2+707p+2455$$ $$+(4p^2+44p+291)gcd(p-1,3)+(p^2+19p+135)gcd(p-1,4)$$ $$+(3p+31)gcd(p-1,5)+4gcd(p-1,7)+5gcd(p-1,8)+gcd(p-1,9)$$ What I get with GAP : gap> LoadPackage("sglppow"); LieRing a package for working with Lie rings by Serena Cicalo' and Willem de Graaf ---------- Loading LiePRing 1.8 by Michael Vaughan-Lee and Bettina Eick ---------- ---------- Loading SglPPow 1.0 by Michael Vaughan-Lee and Bettina Eick ---------- true gap> gnup7:=p->3*p^5+12*p^4+44*p^3+170*p^2+707*p+2455+ > (4*p^2+44*p+291)*Gcd(p-1,3)+(p^2+19*p+135)*Gcd(p-1,4)+ > (3*p+31)*Gcd(p-1,5)+4*Gcd(p-1,7)+5*Gcd(p-1,8)+Gcd(p-1,9); function(p) ... end gap> p:=Filtered([7..100],IsPrimeInt); [7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43, 47, 53, 59, 61, 67, 71, 73, 79, 83, 89, 97] gap> r:=List(p, x -> [x, gnup7(x), NrSmallGroups(x^7)]); [[7, 113147, 113147], [11, 750735, 750735], [13, 1600573, 1598120], [17, 5546909, 5544456], [19, 9380741, 9378288], [23, 23316851, 23314398], [29, 71271069, 71268616], [31, 98488755, 98486302], [37, 233043067, 233040614], [41, 384847485, 384845032], [43, 485930975, 485928522], [47, 751588475, 751586022], [53, 1356370173, 1356367720], [59, 2299880351, 2299877898], [61, 2710679045, 2710676592], [67, 4306310927, 4306308474], [71, 5734323819, 5734321366], [73, 6578172579, 6578170126], [79, 9721485395, 9721482942], [83, 12413061671, 12413059218], [89, 17537591045, 17537588592], [97, 26866372821, 26866370368]] gap> List(r, x -> [x[2]-x[3]]); [[0], [0], [2453]] gap>

论坛问题帖 | has_answers | Peter | 2016-01-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018604 | MSE-1628000

Group conjecture

Conjecture: Given a finite group G and a subset $A\subset G$. Then $\{A,A^2,A^3,\dots\}$ is a group iff $\forall n\in \mathbb N: |A^n|=|A^{n+1}|$. Given that the composition between the subsets $A,B\subset G$ is $A\cdot B=\{g\in G|\exists a\in A\exists b\in B:g=a\cdot b\}$. Example: suppose that $N\subset G$ is a normal subgroup, then the cosets $\{Ng,Ng^2,...\}$ have the same cardinality and constitutes a group, while for random sets the cardinality seems to grow when multiplying: { 3412 2143 4321 1234 } { 2143 1234 } nswap pnormal . -1 ok { 3412 2143 4321 1234 } { 2143 1234 } pquotient set. {{3412,4321},{2143,1234}} ok { 4321 3412 } go ok gen. {4321,3412} ok gen. {2143,1234} ok gen. {4321,3412} ok ndrop ok { 2431 2341 } go ok gen. {2431,2341} ok gen. {4132,3142,4312,3412} ok gen. {1234,1243,3214,4213,1324,1423,3124,4123} ok gen. {2413,2314,3421,4321,1423,1324,2341,2431,2143,2134,3241,4231,1243,1234} ok gen. {4123,3124,4321,3421,2143,2134,4132,3142,4213,3214,4231,3241,3412,4312,1432,1342,2413,2314,2431,2341} ok gen. {4231,3241,1234,1243,3214,4213,1432,1342,1324,1423,2134,2143,2314,2413,4123,3124,4321,3421,4132,3142,4312,3412} ok gen. {3412,4312,2314,2413,2341,2431,2143,2134,4321,3421,3241,4231,1342,1432,3142,4132,1234,1243,3214,4213,1324,1423,3124,4123} ok gen. {4123,3124,3142,4132,3412,4312,1432,1342,3214,4213,2413,2314,3421,4321,1423,1324,2341,2431,2143,2134,3241,4231,1243,1234} ok ndrop ok

论坛问题帖 | has_answers | Lehs | 2016-01-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018605 | MSE-1628437

Does the antipode of a f.d. Hopf algebra have finite order

In a lecture I have heard that the antipode of a finite-dimensional Hopf algebra is conjectured to be finite, and that this has only been proven in characteristic 0 by Larson and Radford in 1988. I couldn't find such a result (only stronger statements that the antipode has order 2 or 4 in some nice situations). But I have found the older paper D. E. Radford, The antipode

of a finite-dimensional Hopf algebra over a field has finite order. Bull. Amer. Math. Soc. 81 (1975), no. 6, 1103--1105 which gives a proof sketch (which I cannot verify) that the conjecture is true in general. What is the actual status of the conjecture? Related (weaker) question: Is the antipode of a finite-dimensional Hopf algebra always invertible? If so, is there an easy proof for this statement, which one can understand without much background? Takeuchi has constructed an infinite-dimensional Hopf algebra whose antipode is not invertible.

论坛问题帖 | no_answers_at_retrieval | Martin Brandenburg | 2016-01-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018606 | MSE-1639468

Rational solutions for $\sin(n)$ in radians

This is completely for my own curiosity. Does $y = \sin(n)$ have rational solutions for n, an integer number of radians. I know that this is strange because usually integers are only used in degrees. I know that there will be convergent sub series of n's to any rational number in $[-1,1]$. Would we ever get an algebraic integer? It would have something like $$ e^\left(n i\right)=e^\left(\frac{n}{\pi}\pi i\right) $$ I suspect that it will never have a rational solution but I don't know how to prove it.

论坛问题帖 | has_answers | Matttonk | 2016-02-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018607 | MSE-1645595

Conjecture: $\lim_{x\to\infty} \frac{256}{163x}\sum_{n=1}^{\lfloor x\rceil} |\sin n|=1$

I conjecture: $$\lim_{x\to\infty} \frac{256}{163x}\sum_{n=1}^{\lfloor x\rceil} |\sin n|=1$$ Is this provable? If this is false, then can I have a function f such that: $$\lim_{x\to\infty} f(x)\sum_{n=1}^{\lfloor x\rceil} |\sin n|=1$$

论坛问题帖 | has_answers | user311151 | 2016-02-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018608 | MSE-1645997

Does this conjecture in 1-d real analysis seem reasonable

Hello I am currently trying to prove a result and I have basically whittled it down to showing the following is true. Let $I\subset\mathbb{R}$ be an interval and fix $\alpha>1$ real number. Fix $x\in I$ and suppose $f:I \rightarrow \mathbb{R}$ is a strictly increasing function. Let $\{x_k\}$ be a sequence contained in I such that $x_k \neq x$ for all k and $x_k \rightarrow x$. Now suppose that the limit $\lim_{k \to \infty} \frac{f(x_k)-f(x)}{f(x+\alpha(x_k-x))-f(x)} $ exists and is equal to l. Then $l \neq 0$. To me it makes sense that this is true. Since f is strictly increasing it can't do anything too mental near x, since f is locally bounded both must converge to 0 and surely the numerator can't dominate the denominator enough for the limit to be zero. My argument in trying to prove it so far goes: It is clear that $0\leq l \leq 1$ so suppose $l=0$. Then since f is locally bounded we deduce that $f(x_k)-f(x) \to 0$. Since f is strictly increasing we deduce that f must be continuous (here we either get full continuity or left continuity or right continuity depending on the sequence x_k but I don't think this really matters much) at x. Then I've been playing around with functions and this seems to hold for everything but I don't really know how to proceed.

论坛问题帖 | has_answers | th0masb | 2016-02-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018609 | MSE-1671836

If b is an odd composite number and $\dfrac{b^2 - 1}{\sigma(b^2) - b^2} = q$ is a prime number, what happens when $q = 2^{r + 1} - 1$?

(Note: An improved version of this question has been cross-posted to MO.) Let $\sigma(X)$ be the sum of the divisors of X. For example, $\sigma(2) = 1 + 2 = 3$, and $\sigma(4) = 1 + 2 + 4 = 7$. My question is: If b is an odd composite and $\dfrac{b^2 - 1}{\sigma(b^2) - b^2} = q$ is a prime number, what happens when $q = 2^{r + 1} - 1$ (with $r \geq 1$)? Without the restriction on $q = 2^{r + 1} - 1$ being prime, I only know that $M = {2^r}{b^2}$ must be an even almost perfect number. (That is, it must satisfy $\sigma(M) = 2M - 1$.) I guess my question can be rephrased as follows: (1) If b is an odd composite, how often is $\dfrac{b^2 - 1}{\sigma(b^2) - b^2} = q$ a prime number? (2) If b is an odd composite, and $\dfrac{b^2 - 1}{\sigma(b^2) - b^2} = q$ is a prime number, how often is q a Mersenne prime?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2016-02-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018610 | MSE-1672261

Does the Descartes number $D = 198585576189$ have a friend?

Let $\sigma(X)$ be the sum of the divisors of X, and denote the abundancy index $\sigma(X)/X$ by $I(X)$. If the equation $I(X) = r/s$ has no solution $X \in \mathbb{N}$, then r/s is said to be an abundancy outlaw. A number $Y \neq X$ is said to be a friend of X if $I(Y) = I(X)$. If a number Z does not have a friend, then it is said to be solitary. Greening's Theorem provides a sufficient but not necessary condition for a number to be solitary - namely, $\gcd\left(Z, \sigma(Z)\right) = 1 \implies Z$ is solitary. Here is my first question: (1-A) Does the Descartes number $D = 198585576189$ have a friend? Greening's Theorem fails to establish the number's solitude, because $$\gcd\left(D, \sigma(D)\right) = \gcd(198585576189, 426027470778) = \gcd\left({3^2}\cdot{7^2}\cdot{{11}^2}\cdot{{13}^2}\cdot{{19}^2}\cdot{61},{2}\cdot{3^3}\cdot{7}\cdot{13}\cdot{{19}^2}\cdot{31}\cdot{61}\cdot{127}\right) = {3^2}\cdot{7}\cdot{13}\cdot{{19}^2}\cdot{61} = 18035199.$$ Here is a way to rephrase my first question: (1-B) Is there a number $\overline{D} \neq 198585576189$ such that $$I(\overline{D}) = I(D) = \dfrac{{2}\cdot{3}\cdot{31}\cdot{127}}{{7}\cdot{{11}^2}\cdot{13}} = \dfrac{23622}{11011}?$$ Notice that 11011 is a palindrome (in base-10). Note also that 23622 is almost a palindrome (again, in base-10). This brings me to my second question: (2) Is the ratio $22622/11011$ an abundancy index or outlaw?

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2016-02-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018611 | MSE-1673006

Can we capture the definition to be a Mersenne prime in an identity only involving the arithmetic function $S(n)=\sum_{k=1}^n\text{nmod k}$?

Let $S(n)=\sum_{k=1}^n\text{nmod k}$ the sum of remainder function, denoting $\sigma(n)$ as the sum of divisors function, it is know that for each $n>1$ $$S(n)-S(n-1)=2n-1-\sigma(n),$$ is as curiority an statement due to Lucas. Too it is know the identities that we can show for perfect nubmers, almos perferct numbers and prime numbers. See here. On the other hand it is know some attempts and theorems that look capture the definition of a sequence only in terms of an arithmetic function. My attempt was use previous function to show that Fact 1. If $m=2^p-1$ is a Mersenne prime then $$m\left(S\left(m\right)-S\left(m-1\right)\right)+S\left(\frac{m-1}{2}\right)-S\left(\frac{m+1}{2}\right)=m(m-2).$$ $$$$ The proof is show that then $m(m-2)-\left(m-(2^p-1)\right)=m(m-2)$. Notice that the second summand $-\left(m-(2^p-1)\right)$ is zero. Fact 2. Conversely if $m>3$ is an odd integer and safisties the equation in previous fact, then this satisfies $$m^2+\sigma\left(\frac{m+1}{2}\right)=m\sigma(m),$$ and thus denoting $m=2k-1$ one has $$4k^2+1-4k+\sigma(k)=(2k-1)\sigma(2k-1).$$ $$$$ Computational facts. I don't know if previous identity involving the sum of divisor function for odd integers is in the literature. My computations were that only obtain Mersenne primes when I use a computer. On the other hand as I've said there are relationships between the formulas for primes and almost perfect numbers, for this reason I am confusing. Also I know equation, a relation problem, here in spanish, PROBLEMA 55, page 156-157, where * means unsolved, for which I don't know how clarify this $$$$ Question. Can you get a closed formula involving only the sum of remainders function, that isn't superfluous, and does capture the definition of Mersenne primes? Is this $$m\left(S\left(m\right)-S\left(m-1\right)\right)+S\left(\frac{m-1}{2}\right)-S\left(\frac{m+1}{2}\right)=m(m-2)?$$ If you can show an odd integer m, as counterexample for this last equation I will apprecite, since I don't made computations for this equation. Can you clarify my doubts about if my computations were empty by comparision with the cited literature? Thanks in advance.

论坛问题帖 | no_answers_at_retrieval | user243301 | 2016-02-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018612 | MSE-1675709

I think I've found all roots to $f_k(x)=\sum_{j=1}^k x^j-x^{-j}$ for any k - how to prove it?

Conjecture: The set of unique roots of $$f_k(x)=\sum_{j=1}^k x^j-x^{-j} \;,\;\; x \not=0$$ is given by $e^{i \pi \phi_k}$, where $$\frac{1}{2}\phi_k=\{0, \frac{1}{2}, \underbrace{\frac{\pm1}{k+1},\frac{\pm1}{k},\frac{\pm2}{k+1},\frac{\pm2}{k},\frac{\pm3}{k+1},...}_\text{$k-1$ elements} \}$$ Q: If true, how can this be proven? I unfortunately don't have much experience with proving stuff (I'm a physicist), so any insights would greatly appreciated! This may very well be some lemma, but I thought it might be useful since Mathematica wasn't able to solve this kind of equation for $k>12$. Note that the roots of f_k are equivalent to the roots of $f_{k,n}=\sum_{j=1}^k x^{n+j}-x^{n-j}$. Also, if $x \not = 1$ $f_k=\frac{x^{-k}(x^{k}-1)(x^{k+1}-1)}{x-1}$. The process through which I reached this conjecture (btw, sorry if "conjecture" sounds pretentious, but I didn't know what else to call it) is described below. Just to get an overview, here's one way of writing out $f_k=0$: \begin{array}{c|c} k & \\ \hline 1 & x^2-1=0\\ 2 & x^4+x^3-x-1=0 \\ 3 & x^6+x^5+x^4-x^2-x-1=0 \\ \vdots &\vdots \end{array} I noticed a high degree of regularity when plotting the solutions. The solutions to the first 12 k are shown below, plotted in the complex plane. $\hskip1.4in$ (I have no idea why Mathematica doesn't recognize -1 as a solution to f_1, but oh well) I now organized the solutions such that one might see a pattern: If one looks at the list of the roots for $k\in \{1,...,12\}$, they can all be written as a phase only, $e^{i \pi \phi}$, where ϕ is some rational number. For instance, the roots of f_2 are $\{1,e^{\pm \frac{2}{3} i \pi},-1\}$ so $\phi_2=\{0,\pm \frac{2}{3},1 \}$. I'm assuming that this pattern continues indefinitely. All $0<\phi<1$ appears with both a positive and a negative sign, and all sets of solutions include $\phi=\{0,1\}$. The non-negative and non-trivial (0 and 1) values for ϕ_{k} for $k\in \{1,...,12\}$ are listed here (written as reduced fractions, going from smallest to largest values): \begin{array}{c|c} k & \phi_k \\ \hline 2 & \frac{2}{3}\\ 3 & \frac{1}{2} \;\; \frac{2}{3}\\ 4 & \frac{2}{5} \;\; \frac{1}{2} \;\; \frac{4}{5}\\ 5 & \frac{1}{3} \;\; \frac{2}{5} \;\; \frac{2}{3} \;\; \frac{4}{5}\\ 6 & \frac{2}{7} \;\; \frac{1}{3} \;\; \frac{4}{7} \;\; \frac{2}{3} \;\; \frac{6}{7}\\ 7 & \frac{1}{4} \;\; \frac{2}{7} \;\; \frac{1}{2} \;\; \frac{4}{7} \;\; \frac{3}{4} \;\; \frac{6}{7}\\ 8 & \frac{2}{9} \;\; \frac{1}{4} \;\; \frac{4}{9} \;\; \frac{1}{2} \;\; \frac{2}{3} \;\; \frac{3}{4} \;\; \frac{8}{9}\\ 9 & \frac{1}{5} \;\; \frac{2}{9} \;\; \frac{2}{5} \;\; \frac{4}{9} \;\; \frac{3}{5} \;\; \frac{2}{3} \;\; \frac{4}{5} \;\; \frac{8}{9} \\ 10 &\frac{2}{11} \;\; \frac{1}{5} \;\; \frac{4}{11} \;\; \frac{2}{5} \;\; \frac{6}{11} \;\; \frac{3}{5} \;\; \frac{8}{11} \;\; \frac{4}{5} \;\; \frac{10}{11}\\ 11 & \frac{1}{6} \;\; \frac{2}{11} \;\; \frac{1}{3} \;\; \frac{4}{11} \;\; \frac{1}{2} \;\; \frac{6}{11} \;\; \frac{2}{3} \;\; \frac{8}{11} \;\; \frac{5}{6} \;\; \frac{10}{11}\\ 12 & \frac{2}{13} \;\; \frac{1}{6} \;\; \frac{4}{13} \;\; \frac{1}{3} \;\; \frac{6}{13} \;\; \frac{1}{2} \;\; \frac{8}{13} \;\; \frac{2}{3} \;\; \frac{10}{13} \;\; \frac{5}{6} \;\; \frac{12}{13} \end{array} I now rewrote this (by extending by 2 here and there) to get \begin{array}{c|c} k & \phi_k \\ \hline 2 & \frac{2}{3}\\ 3 & \frac{2}{4} \;\; \frac{2}{3}\\ 4 & \frac{2}{5} \;\; \frac{2}{4} \;\; \frac{4}{5}\\ 5 & \frac{2}{6} \;\; \frac{2}{5} \;\; \frac{4}{6} \;\; \frac{4}{5}\\ 6 & \frac{2}{7} \;\; \frac{2}{6} \;\; \frac{4}{7} \;\; \frac{4}{6} \;\; \frac{6}{7}\\ 7 & \frac{2}{8} \;\; \frac{2}{7} \;\; \frac{4}{8} \;\; \frac{4}{7} \;\; \frac{6}{8} \;\; \frac{6}{7}\\ 8 & \frac{2}{9} \;\; \frac{2}{8} \;\; \frac{4}{9} \;\; \frac{4}{8} \;\; \frac{6}{9} \;\; \frac{6}{8} \;\; \frac{8}{9}\\ \vdots & \vdots \end{array} which clearly reveals the pattern given above. Here are the first 100 sets of solutions: And 500:

论坛问题帖 | has_answers | Bobson Dugnutt | 2016-02-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018613 | MSE-168334

Is this conjecture on Fibonacci sums correct?

I have the following conjecture, need to know a proof just in case mine is wrong, or the conjecture itself is wrong. The sum of k distinct Fibonacci numbers can be written in at most k ways as the sum of another k distinct numbers from the sequence Proof: Take, say 4, Fibonacci numbers with distinct values: $$144+34+3+2$$ We can see that we must maintain the distinctness and number of the Fibonacci numbers. So when we split the numbers; $$[89+55]+[13+21]+3+2$$ It is immediately apparent that if some numbers are split into smaller Fibonacci numbers, others must be merged to keep only 4 numbers. Thus, 3+2 becomes 5, and either 144 or 34 kept as-is. Thus, $$144+34+3+2=89+55+34+5=144+21+13+5$$ I know it's a rudimentary proof, and k ways of writing the same sum is probably a wrong upper bound, but at least it works for now. Please do let me know what's wrong. Thanks.

论坛问题帖 | has_answers | Simian | 2012-07-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018614 | MSE-1696714

If $|G_1|=|G_2|<\infty$ and $|G_1'|<|G_2'|$, then $|Z(G_1)|\geq |Z(G_2)|$? where G' is the commutator subgroup of G.

We know that G' characterization how ``abelian'' of a group because we have a theorem: if $G'=\{e\}$, then G is abelian. I have a conjecture. If there are two finite groups G_1 and G_2, $|G_1|=|G_2|$ and $|G_1'|< |G_2'|$, where G' is the commutator subgroup of G, does it imply that $|Z(G_1)|\geq |Z(G_2)|$? Or find a counterexample, is there an example satisfing $|G_1|=|G_2|<\infty$, $|G_1'|< |G_2'|$ and $|Z(G_1)|< |Z(G_2)|$? This conjecture is true for $|G|\leq 30$ by verify manually myself and this website. Remarks There is a similar theorem: if $\text{inn }G=\{e\}$, then G is abelian, where $\text{inn }G$ is the group of inner automorphism on G. If there are two finite groups G_1 and G_2, $|G_1|=|G_2|$ and $|\text{inn }G_1|< |\text{inn }G_2|$, which imply that $|Z(G_1)|> |Z(G_2)|$ because $G/Z(G)\cong \text{inn }G$.

论坛问题帖 | no_answers_at_retrieval | bfhaha | 2016-03-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018615 | MSE-1697145

Are there infinitely many primes in any sequence determined by a k that is not a Sierpinski number?

Consider the sequence of numbers ranging over n of the form $k\cdot a^n + 1$ for a fixed, odd natural number k. The number k is considered a Sierpinski number if the sequence determined by that k contains no primes. One way to show that such a k is a Sierpinski number is to find a covering system of congruences for the sequence proving there are no primes in the sequence. The way to show that such a k is not a Sierpinski number is to find one prime among the numbers in the sequence. For those sequences where a prime has been found, that is, k is not a Sierpinski number, is it known whether there could be infinitely many primes in that sequence? If only a finite number of primes are expected to be in the sequence, then a covering system of congruences might work after a certain n to show that the rest of the sequence contains only composite numbers. Sierpinski showed that there are an infinite number of k whose sequences contain no primes. My conjecture is that there can be at most a finite number of primes for any k, but I wonder if this has not already been resolved by someone or if someone has already made the conjecture.

论坛问题帖 | no_answers_at_retrieval | Frank Hubeny | 2016-03-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018616 | MSE-1706416

$\sum\limits_i a_i^2\sum\limits_i b_i^2+\left(\sum\limits_ia_i b_i\right)^2\geq \sqrt{\sum\limits_i a_i^4\sum\limits_i b_i^4}+\sum\limits_ia_i^2b_i^2$

I have no idea about how to prove (or disprove) the following inequality: $$ \left(\sum_{i=1}^n a_i^2\right)\left(\sum_{i=1}^n b_i^2\right)+\left(\sum_{i=1}^na_i b_i\right)^2\geq \sqrt{\left(\sum_{i=1}^n a_i^4\right)\left(\sum_{i=1}^n b_i^4\right)}+\sum_{i=1}^na_i^2b_i^2,\quad a_i,b_i\in\mathbb{R}, \ n>1. $$ I ran some numerical simulations and no counterexample showed up yet. Note 1. The inequality holds true for $n=2$, as shown here. Note 2. This conjecture was formulated by Fedor Petrov in an attempt to provide a solution to a particular case of this question. EDIT. The inequality has been finally proved here.

论坛问题帖 | has_answers | Ludwig | 2016-03-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018617 | MSE-1710951

Is the following inequality involving the sum-of-divisors and Euler totient functions true?

First Question Is the following inequality involving the sum-of-divisors σ and Euler totient ϕ functions true? $$\frac{\sigma(N)}{N} \leq \frac{N}{\phi(N)}$$ Second Question When N satisfies $\sigma(N) = 2N - p$ for some $p > 1$, we say that N is p-deficient and we know that $$\frac{2N}{3N - \sigma(N)} < \frac{\sigma(N)}{N}.$$ Assuming the previous inequality is correct, this gives the upper bound $$\frac{\phi(N)}{N} < \frac{3}{2} - \frac{\sigma(N)}{N}.$$ (Consider $N = 2^r$ where $r \geq 1$. (Note that N is deficient.) Then $$\frac{\sigma(N)}{N} = \frac{\sigma(2^r)}{2^r} \geq \frac{\sigma(2)}{2} = \frac{3}{2}.$$ This agrees with the fact that $p = 2N - \sigma(N) = 2(2^r) - \sigma(2^r) = 1$.) These considerations bring me to my next question: Is there a nice lower bound for $\phi(N)/N$ that will allow us to rule out many, if not most, deficient values for N?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2016-03-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018618 | MSE-1719749

Uniqueness or non uniqueness of a pair of natural numbers

Let $1<m<n$ be two natural numbers. Let us call (m,n) a math.se pair if the prime factors of m are the same as those of n and the prime factors of $m+1$ are the same as those of $n+1$, e.g. $(2,8)$. For every $s\geq 2$, the pair $((2^s-2),2^s(2^s-2))$ is such a pair (for $s=2$ we have the previous pair). The question I asked myself was: are there other math.se pairs, i.e., math.se pairs not of the form $((2^s-2),2^s(2^s-2))$? So, I made a

fortran program that verifies if (i,j) is a math.se pair for $1<i<j$ and for $1<i\leq m$ and $m<j\leq n$ where m and n are input values, but the only different math.se pair that I (actually, it was the computer) found is $(75,1215)$ (we have that $75=3\cdot5^2$, $1215=3^5\cdot 5$, $76=2^2\cdot19$ and $1216=2^6\cdot19$). The problem is that the greater i or j is, the greater is the time the computer takes. Now I wonder if that $(75,1215)$ is the only math.se pair not of the form $((2^s-2),2^s(2^s-2))$. If there are others, are there infinitely many of them? If yes, can we list them all? If you look carefully, you'll see that $(a_s+1)^2=b_s+1$, where $a_s=2^s-2$ and $b_s=2^s(2^s-2)$. With this in mind, I tried to obtain a math.se pair (a,b) that satisfies $(a+1)^3=b+1$. In this case, we need to find a number b, such that $b=a^3+3a^2+3a$ and such that b and a have the same prime factors. But even in this case I don't know how to show if there is or if there isn't such a math.se pair.

论坛问题帖 | has_answers | Larara | 2016-03-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018619 | MSE-1725549

A conjecture about traces of projections

Let M_n denote the space of all $n\times n$ complex matrices. Define $\tau:M_n\rightarrow \mathbb{C}$ by $$\tau(X)=\frac{1}{n}\sum_{i=1}^n x_{ii},$$ where of course $X=[x_{ij}]\in M_n$. Recall that a matrix $P\in M_n$ is called an "orthogonal projection" if $P=P^*=P^2$. Let A, B, C be orthogonal projections in M_n and define two quantities - $$x=\frac{1}{3}\tau(A+B+C)$$ and $$y=\frac{1}{3}\tau(AB+BC+CA).$$ It is easy to see that for any orthogonal projections $A,B,C\in M_n$ the value of x lies in $[0,1]$. I want to investigate the case when $x\in [\frac{1}{3},\frac{1}{2}]$. My goal is to compute the infimum of y under this constrain on x. My observation is that the minimum value of y is $\frac{3x-1}{4}$ when $x\in [\frac{1}{3},\frac{1}{2}]$ and for all $n\in \mathbb{N}$. However I am unable to prove this and I wondered if people here may be able to see why this might hold, or provide a counterexample to this conjecture.

论坛问题帖 | has_answers | Nirakar Neo | 2016-04-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018620 | MSE-1726236

How close are we to knowing the rate of convergence to 0 of $\prod_{p\le x}(1-1/p)^{-1}-e^\gamma\log x $?

This is a question related to an earlier one of mine, which I may answer myself eventually, as I have learnt more about the topic. Despite what one can read on the MathWorld page about Mertens' third theorem, in 2009 Diamond and Pintz showed$$\underset{p\ \text{prime}}{\prod_{p\le x}}\left(1-\frac1p\right)^{-1}-e^\gamma\log x=\Omega_{\pm}\left(\frac{\log\log\log x}{\sqrt{x}}\right);$$on the other hand, by a theorem of Dusart, for all $x\ge2973$ $$\left\lvert\prod_{p\le x}\left(1-\frac1p\right)^{-1}-e^\gamma\log x\right\rvert <\frac{0.2}{\log x}.$$ In the end of 2014, using a probabilistic argument, Lamzouri conjectured $$\left\lvert\prod_{p\le x}\left(1-\frac1p\right)^{-1}-e^\gamma\log x\right\rvert \sim \frac{e^\gamma}{2\pi}\frac{(\log\log\log x)^2}{\sqrt{x}}.$$ Has it been established that$$-\infty<\liminf_{x\to\infty}\frac{\sqrt{x}}{(\log\log\log x)^2}\left(\prod_{p\le x}\left(1-\frac1p\right)^{-1}-e^\gamma\log x\right)<0 \ ?$$I googled quite a lot, but I could find nothing. Has there been no progress yet?

论坛问题帖 | no_answers_at_retrieval | Vincenzo Oliva | 2016-04-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018621 | MSE-1739431

conjectured general continued fraction for the quotient of gamma functions

Given complex numbers $a=x+iy$, $b=m+in$ and a gamma function $\Gamma(z)$ with $x\gt0$ and $m\gt0$, it is conjectured that the following general continued fraction which is symmetric on a and b is true $$\frac{\displaystyle\Gamma\left(\frac{a+3b}{4(a+b)}\right)\Gamma\left(\frac{3a+b}{4(a+b)}\right)}{\displaystyle\Gamma\left(\frac{3a+5b}{4(a+b)}\right)\Gamma\left(\frac{5a+3b}{4(a+b)}\right)}=\cfrac{4(a+b)}{a+b+\cfrac{(2a)(2b)}{3(a+b)+\cfrac{(3a+b)(a+3b)}{5(a+b)+\cfrac{(4a+2b)(2a+4b)}{7(a+b)+\ddots}}}}}\tag{1}$$ And can be further generalised to $$\frac{1}{\sqrt{c}}\tan\left(\frac{b-a}{b+a}\tan^{-1}\left(\frac{\sqrt{c}}{d}\right)\right)=\cfrac{(b-a)}{d(a+b)+\cfrac{c(2a)(2b)}{3d(a+b)+\cfrac{c(3a+b)(a+3b)}{5d(a+b)+\cfrac{c(4a+2b)(2a+4b)}{7d(a+b)+\ddots}}}}\tag{2}$$ Corollaries: (i) Specializing to $a=1/2$ and $b=2z/n+1/2$,we obtain the continued fraction for $\tan\left(\frac{z\pi}{4z+2n}\right)$ found in this post after applying the functional equation of the gamma function (ii) and specializing further to $a=-1/2$ and $b=2z/n+3/2$,we obtain immediately the continued fraction for $\cot\left(\frac{z\pi}{4z+2n}\right)$ found here after applying the functional equation of the gamma function.This continued fraction was proved by @GEdgar. (iii) letting $2a=m-n$ and $2b=m+n$,we find $$\displaystyle\tan\left(\frac{\pi n}{4m}\right)=\cfrac{n}{m+\cfrac{m^2-n^2}{3m+\cfrac{4m^2-n^2}{5m+\cfrac{9m^2-n^2}{7m+\cfrac{16m^2-n^2}{9m+\ddots}}}}}$$ From which we obtain its hyperbolic companion $$\displaystyle\tanh\left(\frac{\pi n}{4m}\right)=\cfrac{n}{m+\cfrac{m^2+n^2}{3m+\cfrac{4m^2+n^2}{5m+\cfrac{9m^2+n^2}{7m+\cfrac{16m^2+n^2}{9m+\ddots}}}}}$$ (iv)and if $2a=-n$ and $2b=2m+n$,then it follows that $$\displaystyle\tan\left(\frac{\pi(m+n)}{4m}\right)=\frac{1+\tan\Big(\frac{\pi n}{4m}\Big)}{1-\tan\Big(\frac{\pi n}{4m}\Big)}=\cfrac{m+n}{m+\cfrac{(-n)(2m+n)}{3m+\cfrac{(m-n)(3m+n)}{5m+\cfrac{(2m-n)(4m+n)}{7m+\cfrac{(3m-n)(5m+n)}{9m+\ddots}}}}}$$ Q: Is the conjectured general continued fraction true (for all complex numbers a,b with $x\gt0$ and $m\gt0$)?

论坛问题帖 | has_answers | Nicco | 2016-04-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018622 | MSE-1763562

f has to be open in order for continuous bijection to be a homeomorphism?

Claim : Let $f: X\to Y$ be a continuous bijection, then f is a homeomorphism if f is open Is the claim true? I thought f was a homemorphism if f^{-1} is a continuous function

论坛问题帖 | has_answers | Your neighbor Todorovich | 2016-04-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018623 | MSE-1773546

Endowing module with algebra

Given a module M over a ring R, is it possible to endow M with an operation $M^{2} \to M$ that turns M to an algebra that is not the trivial $m n \equiv \mathbf{0}$ identically? So far I have that if M is a free module, i.e. it has a basis, then we can define a pointwise multiplication that non-trivially makes M an algebra. My first thought after this was to consider M as a direct sum $R \oplus N$ for some submodule $N \subseteq M$, and define it by $(r, n)(r', n') = (r r', 0)$, but even this (if my module can be written as the direct sum as earlier supposed) feels kinda trivial. So the next question: can any module M be endowed with multiplication such that N is a nontrivial subalgebra for every nontrivial submodule $N \subseteq M$? I'm quite at a loss on how to show it. Thanks.

论坛问题帖 | no_answers_at_retrieval | AJY | 2016-05-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018624 | MSE-1782472

Factoring the factorials

Just for the fun of it, I've started factoring $n!$ into its prime divisors, and this is what I got for $2\leq n\leq20$: $$\begin{align} 2! &= 2^\color{red}{1} &S_e=1\\ 3! &= 2^\color{red}{1} \cdot 3^\color{red}{1} &S_e=2\\ 4! &= 2^3 \cdot 3^\color{red}{1} &S_e=4\\ 5! &= 2^3 \cdot 3^\color{red}{1} \cdot 5^\color{red}{1} &S_e=5\\ 6! &= 2^4 \cdot 3^2 \cdot 5^\color{red}{1} &S_e=7\\ 7! &= 2^4 \cdot 3^5 \cdot 5^\color{red}{1} \cdot 7^\color{red}{1} &S_e=8\\ 8! &= 2^7 \cdot 3^2 \cdot 5^\color{red}{1} \cdot 7^\color{red}{1} &S_e=11\\ 9! &= 2^7 \cdot 3^4 \cdot 5^\color{red}{1} \cdot 7^\color{red}{1} &S_e=13\\ 10! &= 2^8 \cdot 3^4 \cdot 5^2 \cdot 7^\color{red}{1} &S_e=15\\ 11! &= 2^8 \cdot 3^4 \cdot 5^2 \cdot 7^\color{red}{1} \cdot 11^\color{red}{1} &S_e=16\\ 12! &= 2^{10} \cdot 3^5 \cdot 5^2 \cdot 7^\color{red}{1} \cdot 11^\color{red}{1} &S_e=19\\ 13! &= 2^{10} \cdot 3^5 \cdot 5^2 \cdot 7^\color{red}{1} \cdot 11^\color{red}{1} \cdot 13^\color{red}{1} &S_e=20\\ 14! &= 2^{11} \cdot 3^6 \cdot 5^3 \cdot 7^2 \cdot 11^\color{red}{1} \cdot 13^\color{red}{1} &S_e=22\\ 15! &= 2^{11} \cdot 3^6 \cdot 5^3 \cdot 7^2 \cdot 11^\color{red}{1} \cdot 13^\color{red}{1} &S_e=24\\ 16! &= 2^{15} \cdot 3^6 \cdot 5^3 \cdot 7^2 \cdot 11^\color{red}{1} \cdot 13^\color{red}{1} &S_e=28\\ 17! &= 2^{15} \cdot 3^6 \cdot 5^3 \cdot 7^2 \cdot 11^\color{red}{1} \cdot 13^\color{red}{1} \cdot 17^\color{red}{1} &S_e=29\\ 18! &= 2^{16} \cdot 3^8 \cdot 5^3 \cdot 7^2 \cdot 11^\color{red}{1} \cdot 13^\color{red}{1} \cdot 17^\color{red}{1} &S_e=32\\ 19! &= 2^{16} \cdot 3^8 \cdot 5^3 \cdot 7^2 \cdot 11^\color{red}{1} \cdot 13^\color{red}{1} \cdot 17^\color{red}{1} \cdot 19^\color{red}{1} &S_e=33\\ 20! &= 2^{18} \cdot 3^8 \cdot 5^4 \cdot 7^2 \cdot 11^\color{red}{1} \cdot 13^\color{red}{1} \cdot 17^\color{red}{1} \cdot 19^\color{red}{1} &S_e=36\\ \end{align}$$ where S_e is the sum of the exponents. There are lots of interesting patterns to chase here, but I've focused on the prime factors with exponent 1 (highlighted). My conjecture is, In the sequence above, prime factor p^1 will appear p times starting from $p!$. Or, otherwise stated, $\forall k = 2p-1$, $k!$ is the last term of the sequence $\{a_n\}=n!$ to have p^1 as one of its prime factors. [I could not find any reasonable pattern concerning p^2.] This might be some well-known fact, but as someone who hasn't seen prime factorization since elementary school I was wondering how this could be proven (or whether it is even right, although it looks like it is).

论坛问题帖 | has_answers | giobrach | 2016-05-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018625 | MSE-1783949

Conjecture about odd primes

For each odd prime p there exist $n\in\mathbb{N}$ such that $p\equiv n^2 \text{ (mod }\varphi(n^2))$, where φ is Euler's totient function. I'm developing my Forth based computational system Zet to be used for testing conjectures and it seems that it generate conjectures as well... What I have tried is to test the modular condition for squares of all integers less than n. All odd primes less than n are represented then. It's not so that all odd number is represented - which seems to be the case if dropping the squaring: $p\equiv n \text{ (mod }\varphi(n))$ - instead the most of the numbers represented are primes.

论坛问题帖 | has_answers | Lehs | 2016-05-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018626 | MSE-1794203

Number of primes of a certain form

Let p_n be the nth prime. Are there an infinite number of primes of the form $2p_n+1$? Is something known about questions like this?

论坛问题帖 | no_answers_at_retrieval | Lehs | 2016-05-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018627 | MSE-1794448

A conjecture about primes: if a,b are coprime and not both odd, is $A(a,b,m)$ finite for some m?

Let p_n be the nth prime and define $p_n^{(m)}$ by $p_n^{(1)}=p_n$ and $p_n^{(m+1)}=p_{p_n^{(m)}}$: $p_n^{(2)}=p_{p_n}$, $\;p_n^{(3)}=p_{p_{p_n}}$ and so far... For some coprime numbers a,b, not both odd, do there exist a number $m\in \mathbb Z_+$ such that $A(a,b,m)=\{ap_n^{(m)}+b\in\mathbb P|n\in\mathbb Z_+\}$ is finite? $\mathbb P$ is the set of primes. I have done some tests and there is no sign of finiteness. Conjecture: If a and b are coprime and $a+b$ is odd, then $A(a,b,m)$ is infinite for all $m\in\mathbb Z_+$. One consequence of the conjecture, which I don't expect to be proved, is that any even gap just not only exists but exists infinitely, not only for primes, but also for the case when one of the primes is of the form $p_n^{(m)}$ any m. So perhaps there is a possibility of counterexamples? See also this paper.

论坛问题帖 | no_answers_at_retrieval | Lehs | 2016-05-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018628 | MSE-1797863

A conjecture about the prime function p_n: $p_m \cdot p_n >p_{m \cdot n}$

While testing my system Zet for computational mathematics I find possible relations now and then. The latest is: Conjecture: For all $(m,n)\in\mathbb Z_+^2$ except $(3,4),(4,3) \text{ and } (4,4)$ it holds that $p_m\cdot p_n > p_{m\cdot n}$

论坛问题帖 | has_answers | Lehs | 2016-05-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018629 | MSE-1803721

A very difficult Diophantine problem $n^2 \mid 3^n+2^n+1$

Prove that $n=3$ is the only positive integer greater than 1, for which$$n^2 \mid 3^n+2^n+1.$$ This is a conjecture.

论坛问题帖 | has_answers | Ghartal | 2016-05-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018630 | MSE-1804279

A stronger form of Rolle's Theorem in the direction of number of roots of $f'(x)$

Today I read an interesting generalization of the Rolle's Theorem for Polynomials in $E. 28$ of E. J. Barbeau's book on Polynomials. It says that if a, b are two consecutive zeroes of polynomial $P(x)$ then the number of roots $P'(x)$ (counting multiplicities) in (a, b) is odd in number. I saw the proof and all. Now, this made me conjecture a similar thing for any differentiable function, given consecutive roots exist (roots are countable). ALso, I had to assume that the number of roots of $f'(x)$ in (a, b) is finite in number. Also, I don't know if this makes sense, but I also defined multiplicity of roots for $f(x)$ as if $f(a)=f'(a)= \cdots = f^{(n)}(a)=0$ and $f^{(n+1)} \neq 0$, then I say that multiplicity of a is n and I count it as n roots. My conjecture is that the number of roots of $f'(x)$ (counting orders) stictly between any two consecutive roots of f is odd in number. Also, I think I have proved it given the assumptions. Because clearly the extremums of $f(x)$ between a and b are odd in number and any point critical point in (a, b) that is not an extremum will be a root with even multiplicity by nth derivative test which I can prove. So, the conjecture holds. Now, my main questions are, is there a better form of the above conjecture? Also, are any of my assumptions on the countability and infinitude of roots "unecessary"? Are there any analogous results when the roots of $f'(x)$ are infinitude? Please throw light. Now my question is not very specific or may its asking for too many things, so any related results or links will also be immensely appreciated. Also I am in high school and don't much analysis analysis (i know uptil spivak calculus). UPDATE: Since my conjecture (with the assumptions stated seems true) and we are trying to find and analogous extension of it for the case of infinitely many derivatives, I thought we must try to extend the notion of cardinality of a set with odd number of elements to sets with infinitely many elements. Here is my proposed extension: Call the cardinality of a set S with infinitely many elements, odd iff we cannot cannot partition S into A and B such that there exists a bijection from A to B. When the cardinality is finite, then the above definition is clearly true by definition. So, does the above extension make sense (even when the set is countable)? Now with the above definition can we say that the number of roots of $f'(x)$ in (a, b) odd? (at least when the roots are countably infinite?)

论坛问题帖 | has_answers | Subham Jaiswal | 2016-05-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018631 | MSE-1804411

Conjecture about Rabin-Miller pseudo prime test

I tested the Rabin-Miller pseudo prime algorithm using a single test value and found that the number of false calls depends on the size of the number to test, reducing to a (conjectured) negligible probability of error for very large numbers. A number is randomly chosen to be tested. If the number is $0,1,2,3$ or even the correct result is returned. If the number is odd and >3 the Rabin-Miller algorithm specified below is used. The result is compared with table values. This algorithm use a random number a for the test. It would never miss a prime but occasionally report a composite to be a prime. If the test is erroneous an error is returned. I used the method at oeis A014233 where a method for exact testing of primes using multiple Rabin-Miller tests is described, together with squeezed prime tables, to assess the truth of primality. For each b I ran the test $10,000,000$ times with pseudo random numbers less than 2^b and then got the following number of errors e: e b f 78144 8 78125 61425 9 55243 43642 10 39063 32688 11 27621 22201 12 19531 15190 13 13811 10789 14 9766 7176 15 6905 4814 16 4883 3227 17 3453 2122 18 2441 1422 19 1726 926 20 1221 574 21 863 400 22 610 247 23 432 194 24 305 112 25 216 65 26 153 46 27 108 24 28 76 21 29 54 11 30 38 9 31 27 2 32 19 4 33 13 4 34 10 0 35 7 2 36 5 0 37 3 0 38 2 0 39 2 0 40 1 0 41 1 The f-column is the function $f(b)=1250000\cdot 2^{-\frac{b}{2}}$. The algorithm used is: To test an odd number n, write $n=2^r\cdot s+1$, where s is odd. Given a random number a such that $1<a<n$, if 1. $a^s\equiv 1(\text{mod}\; n)$ or 2. it exists an integer j: $0\le j<r$ with $a^{2^j\cdot s}\equiv -1(\text{mod}\; n)$ then n is pseudo prime. The conjecture is: For b big enough $\text{e}<1250000\cdot 2^{-\frac{b}{2}}$. And specially, $\text{e}\sim 2^{-\frac{b}{2}}$. I intend to use the algorithm for testing big integers and it seems like just one test is enough for $b>256$ since $f(256)=\frac{1}{272225893536750770770699685945414}\,$, which is supposed to be the probability for an error within 10,000,000 random primary tests. For $b<<256$ one can use multiple tests or an exact algorithm. But how to prove the conjecture or at least make it resonable? I will also reward an indepentent computer test, that support or not support my own computer test. I made a test with double check of Rabin-Miller and made a diagram:

论坛问题帖 | has_answers | Lehs | 2016-05-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018632 | MSE-180606

Integer Partition by Counting Repetition : Conjecture ??

I would like to find informations regarding this way of doing Integer Partitions or this conjecture, Suppose you have all the ordered partitions of 5: 5 4 1 3 2 2 2 1 3 1 1 2 1 1 1 1 1 1 1 1 Then extend those partitions to 5 digits by adding 0: 5 0 0 0 0 4 1 0 0 0 3 2 0 0 0 2 2 1 0 0 3 1 1 0 0 2 1 1 1 0 1 1 1 1 1 And finaly for each extended partitions, count the digit repetitions, like this: 5 0 0 0 0 -> 1 4 4 1 0 0 0 -> 1 1 3 3 2 0 0 0 -> 1 1 3 2 2 1 0 0 -> 2 1 2 3

1 1 0 0 -> 1 2 2 2 1 1 1 0 -> 1 3 1 1 1 1 1 1 -> 5 And you can repeat ! 5 0 0 0 0 -> 1 4 -> 1 4 0 0 0 -> 1 1 3 -> 1 1 3 0 0 -> 2 1 2 -> 2 1 2 0 0 -> 1 1 1 2 -> .. 4 1 0 0 0 -> 1 1 3 -> 1 1 3 0 0 -> 2 1 2 -> 2 1 2 0 0 -> 1 1 1 2 -> 1 1 1 2 0 -> 3 1 1 3 2 0 0 0 -> 1 1 3 -> 1 1 3 0 0 -> 2 1 2 -> 2 1 2 0 0 -> 1 1 1 2 -> 1 1 1 2 0 -> 3 1 1 2 2 1 0 0 -> 2 1 2 -> 2 1 2 0 0 -> 1 1 1 2 -> 1 1 1 2 0 -> 3 1 1 -> 3 1 1 0 0 -> 1 2 2 3 1 1 0 0 -> 1 2 2 -> 1 2 2 0 0 -> 1 2 2 -> 1 2 2 0 0 2 1 1 1 0 -> 1 3 1 -> 1 3 1 0 0 -> 1 1 1 2 -> .... -> 1 2 2 1 1 1 1 1 -> 5 -> 5 0 0 0 0 -> 1 4 -> 1 4 0 0 0 -> 1 1 3 -> ... -> 1 2 2 All of them finish in 1 2 2! I want informations, documentations Thanks a lot!

论坛问题帖 | has_answers | Yvan | 2012-08-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018633 | MSE-1828091

Conflicting conjectures

I feel like when two conjectures are inconsistent with one another, it's a clear sign of our misunderstanding of deeper mathematics. I was wondering if anyone knew of a comprehensive list of conjectures that contradict other conjectures.

论坛问题帖 | no_answers_at_retrieval | JasonM | 2016-06-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018634 | MSE-1828323

Seeking proof to a Hyperbolic polygon conjecture

In the course of writing a(n Honours) thesis, I'm searching for a proof to a conjecture that appears very likely to be true. Many results will rely upon it. My own attempts to prove it have been unfruitful, and at this point I'd be happy to settle for a reference to something in the literature. Here's the conjecture: Let $O,V_1, \dots V_{n-2} ,f$ be an $n-$pointed convex polygon in H^2. Given all outer angles (those between adjacent edges of the polygon), and angles $\angle V_n O V_m$ and $\angle V_{n-2} O f$, the entire polygon is uniquely determined up to isometry. Pictorial statement of the conjecture: Given all outer angles of an $n-$pointed convex polygon, and all α_i, the full geometry of the shape is determined up to isometry.

论坛问题帖 | no_answers_at_retrieval | Myridium | 2016-06-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018635 | MSE-1830280

There are at least two solutions such that $2p_n=p_a+p_b$ (p being prime)

I've stumbled across this playing around and summing primes at random during a boring lecture. Is this a known conjecture? Can it be proven? My conjecture: There exists at least one non trivial solution such that $2p_n = p_a + p_b$ (the trivial being obviously $a=b=n$) for $n > 2$. Tested by starting at the trivial $2p_n = p_n + p_n$ then incremented the right as I decremented the left until both were prime again or I've ran below 2 with the left number. The second condition never occurred though, and I've tested for the first 1000 primes by writing a simple program. It fascinated me for the fact that this would mean that $p_b = 2p_n - p_a$ where $b > n > a$ so by knowing primes up to the nth you would have enough information to evaluate the $(n + 1)$-th?

论坛问题帖 | has_answers | Ilhan | 2016-06-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018636 | MSE-1830821

Closed form and limit of the integral of a rational function

While trying to answer this question, I wondered whether there could be a way to: (A) Find the closed form of the generalization of integrals I and J, that is $$I_n=\int_{-\infty}^{+\infty}\frac{dx}{x^{4n}+x^{2n}+1}$$ (B) Prove my own conjecture that $$\lim_{n \to \infty}I_n = 2$$ Is there a clean way to do these? EDIT: I have managed to find a closed form for the poles of the integrand that lie in the upper half-plane, which would be the ones lying within the usual semicircular contour: $$\omega^+_n(j)=\exp\left(i\frac{\pi}{4n}\big(2j-1-(-1)^n\big)\right) \qquad 1 \leq j \leq 2n, \quad j \in \mathbb{N}$$ My integral would become $$\def\res{\mathop{\text{Res}}} \begin{align} \oint_\Gamma \frac{dz}{z^{4n}+z^{2n}+1}&=\lim_{R\to\infty} \int_{-R}^{+R}\frac{dx}{x^{4n}+x^{2n}+1}+\lim_{R\to\infty}iR\int_{0}^{\pi}\frac{e^{i\theta}d\theta}{R^{4n}e^{i\theta4n}+R^{2n}e^{i\theta2n}+1} \\ &=2\pi i \sum_{j=1}^{2n} \res_{z=\omega^+_n(j)} \left(\frac{1}{z^{4n}+z^{2n}+1}\right) \end{align}$$ The limit of the integral in θ is 0, as one would expect, so my question reduces to: Is there a way to express $$\sum_{j=1}^{2n}\res_{z=\omega^+_n(j)} \left(\frac{1}{z^{4n}+z^{2n}+1}\right)$$ in terms of n and j? The answers to (A) and (B) above should follow from this in a relatively straightforward way.

论坛问题帖 | has_answers | giobrach | 2016-06-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018637 | MSE-1834262

Is the Fermat primality test secure enough for very big numbers?

The random variable X_m is the number of trials before $n\notin\mathbb P\wedge n\mid 2^{n-1}-1$ where n is an odd random integer $2^{m-1} < n < 2^m$. Computer simulations makes me believe that $\text E[\log X_m]=\frac{m}{6}$ and that $\operatorname{Var}[\log X_m]<1$. I'm looking for some kind of proof of this conjecture and would like to know how to compute or estimate $P(X_{1000}=1)$, given that the conjecture is true. The context is: how secure is the Fermat primality test with base 2 on numbers with 1000 binary bits? Compared with the probability of hardware errors? Well, perhaps the 10 in the logarithm doesn't flag for an exact $\frac{m}{6}$. The regression line is $\log N= 0.1666\cdot m+0.006$ which is interpreted as $N=10^{\frac{m}{6}}$ but might also be interpreted as $N=\pi^{\frac{m}{3}}$ within the marginals. $\overset{..}{\smile}$ 3251 simulations total so far. Some lower experiments $(m=10)$ has been removed, since lower intervalls gives more irregular results. In some intervalls there are no discrepancies. Also long time running results from $m=40$ is included, so the equation of the line of regression has changed a little. Diagram of the mean values of each m

论坛问题帖 | has_answers | Lehs | 2016-06-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018638 | MSE-1838931

Conjecture about primes and the factorial: for all primes $p>5$, must there exist a prime $q<p$ such that $q\equiv m!\pmod p$ for some $2<m<p$?

Below $0\notin\mathbb N$. Further corrected conjecture: For all prime numbers $p>5$ there exist a prime number $q<p$ such that $q\equiv m!\!\pmod p$, $2<m<p$. or Given a prime $p>5$ there exist a prime $q<p$ and $k,m\in\mathbb N$, $2<m<p$, such that $kp+q=m!$ I want help to prove the conjecture (which is tested for all $p<100,000,000$) or to find a counter-example. p=7 q=3 k=3 m=4 p=11 q=2 k=2 m=4 p=13 q=11 k=1 m=4 p=17 q=7 k=1 m=4 p=19 q=5 k=1 m=4 p=23 q=5 k=5 m=5 p=29 q=23 k=173 m=7 p=31 q=7 k=23 m=6 p=37 q=17 k=19 m=6 p=41 q=23 k=17 m=6 p=43 q=29 k=937 m=8 p=47 q=11 k=107 m=7 p=53 q=31 k=13 m=6 p=59 q=2 k=2 m=5 p=61 q=59 k=1 m=5 p=67 q=53 k=1 m=5 ... Up to the prime 1020361 For several reasons I have had major problems extracting my real observations. Also, the primes are not unique having this property, a lot of semiprimes, but not all, and occasionally some other numbers, also have it. It seems like less than one third of all natural numbers have it and there is a secondary conjecture: For all primes p there is a prime $q<p^2$ such that $q\equiv m!\pmod {p^2}$, $2<m<p^2$. tested for all $p<100,000,000$. There are strong reasons to believe that the conjecture is true. Suppose $0\equiv m!\!\pmod n$, then $0\equiv r!\!\pmod n$ for all $r>m$. And suppose $n=sp^t$, where p is the largest prime dividing n, $p\nmid s$ and $t>0$, then $0\equiv (pt)!\!\pmod n$. If p is a large prime there are a lot of nonzero solutions to $x\equiv m!\!\pmod p$ and the probability for one of those solutions to be a prime increase with p.

论坛问题帖 | has_answers | Lehs | 2016-06-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018639 | MSE-1839944

Conjecture concerning modular arithmetic

Below $0\notin\mathbb N$. I want a proof or a counter-example of the following (corrected) conjecture: Suppose p is the smallest prime dividing $n\in\mathbb N$ and suppose $kn+ap=m!$, where $ap<kn$, for some $a,k,m\in\mathbb N$, then there are $k^{\prime},a^{\prime}\in\mathbb N$ such that $k^{\prime}n+a^{\prime}p=(m+1)!$, where $a^{\prime}p<k^{\prime}n$. A more simple way to write the conjecture is (with some abuse of language): Suppose p is the smallest prime dividing $n\in\mathbb N$ and suppose $p|(m!\pmod n)$, then $p|((m+1)!\pmod n)$

论坛问题帖 | has_answers | Lehs | 2016-06-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018640 | MSE-1844809

What are the mathematical consequences if 10 is proved to be solitary?

Let $\sigma(x) = \sigma_{1}(x)$ denote the sum of the divisors of x, and let $$I(x) = \dfrac{\sigma(x)}{x}$$ be the abundancy index of x. For example, $$\sigma(10) = 1 + 2 + 5 + 10 = 18$$ so that $$I(10) = \dfrac{\sigma(10}{10} = \dfrac{18}{10} = \dfrac{9}{5}.$$ Consider the equation $$I(x) = I(a)$$ where a is fixed. If $x \neq a$, then x is said to be a friend of a. We will then call x and a as friendly numbers. On the other hand, if $x = a$ whenever $$I(x) = I(a),$$ then a is called a solitary number. In this paper by Ward, necessary conditions for the existence of a friend of 10 is derived. It is conjectured that 10 is in fact solitary, although a proof appears to be very difficult. My question in the present post is as follows: What would be the mathematical consequences, if any, of proving that 10 is solitary?

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2016-06-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018641 | MSE-1846289

Is $\zeta(2n+1)\notin (2\pi)^{2n+1}\mathbb{Q}$ already known?

Is it already shown or at least conjectured that $$\zeta(2n+1)\notin (2\pi)^{2n+1}\mathbb{Q}?$$ You have any names and years who proved or conjectured it?

论坛问题帖 | has_answers | user337073 | 2016-07-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018642 | MSE-1854859

A condition for being a prime: $\;\forall m,n\in\mathbb Z^+\!:\,p=m+n\implies \gcd(m,n)=1$

If $\;p=m+n$ where $p\in\mathbb P$, then m,n are coprime, of course. But what about the converse? Conjecture: p is prime if $\;\forall m,n\in\mathbb Z^+\!:\,p=m+n\implies \gcd(m,n)=1$ Tested (and verified) for all $p<100000$.

论坛问题帖 | has_answers | Lehs | 2016-07-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018643 | MSE-1855497

Graph Laplacian rank-one update

Can anyone help me prove/disprove this conjecture? Let G be an undirected nonnegative weighted connected graph with n nodes and Laplacian matrix L. Also, let $0 = \lambda_1 < \lambda_2 \leq \dots\leq \lambda_n$ be eigenvalues of the Laplacian matrix L and assume that $\max(d_i)<1$, where d_i is the degree of i^th node. If $\lambda_i>1$ for some $2\leq i \leq n$, there exists an edge e such that increasing weight of e while having maximum degree less than one, does not increase number of eigenvalues larger than one. Here $d_i = \sum_i a_{ij}$ where a_{ij} is the ij^th component of the adjacency matrix of the graph.

论坛问题帖 | has_answers | user353321 | 2016-07-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018644 | MSE-1857467

For every prime $p > 3$ that is 3 mod 4, does $q+1 \mid p-q$ for some other prime q?

Yet another random conjecture about primes: Given a prime $p>3$ of the form $4n+3$. Then there exist a prime $q<p$ such that $q+1\mid p-q$. Verified for all $p<100000$.

论坛问题帖 | has_answers | Lehs | 2016-07-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018645 | MSE-1859617

Primes in the binomial transform of $ [1, 1, 2, 1, -1, 1, -1, 1, -1, 1, -1, 1, -1, ...]$.

This question is related to this sequence A139482. A commentator gives the following formula for a_m $$a_m = {3m^2-9m+10 \above 1.5pt 2}$$ I have that you should consider the sequence $b_n =3n+2$ and arrange the terms as follows: $$B_3 = \begin{matrix} &&&&&65&\ldots \\ &&&&47&62&\ldots \\ &&&32&44&59&\ldots \\ &&20&29&41&56&\ldots \\ &11&17&26&38&53&\ldots \\ 5&8&14&23&35&50&\ldots \end{matrix}$$ Then a_m appears to be the the sequence on the uppermost slope of B_3 for $m>2$. In particular a_m can be written as b_n for some suitable n. Dirichlet tells us there are infinitely many primes in B_3. The obvious question is are there infinitely many primes on the uppermost slope? If that is too much to ask can we show that every column or row of B_3 has at least one prime number? Update: I believe I have the differences of consecutive terms on the uppermost slope: $a_{m+1}-a_m=3(m+1)$. So that $a_{m+1} = 3(m+1)+a_m$. I also have that $(3,a_m) =1$ for every m. Would not Dirichlet's theorem tell us there are infinitely many primes of the form $3(m+1)+a_m$? <--- The answer to this question is NO!

论坛问题帖 | has_answers | Anthony | 2016-07-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018646 | MSE-1860437

Is this a known conjecture? Given odd primes p,q with $p + q$ sufficiently large, must there exist a different pair p',q' with $p+q = p'+q'$?

Conjecture: There is a natural number $N\in\mathbb N$ such that given odd primes p,q with $p+q>N$ there are primes p',q' where $p' \notin \{p,q\}$ such that $p+q=p'+q'$. Is this known?

论坛问题帖 | has_answers | Lehs | 2016-07-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018647 | MSE-1860716

Combining a working hypothesis for odd perfect numbers with an inequality for logarithms

Euler's theorem for odd perfect numbers states that if there exists and odd perfect number, that is an odd positive integer n satisfying $\sigma(n)=2n$, where $\sigma(m)=\sum_{d\mid m}d$ denotes the sum of divisor function, then this integer has the form $$n=p^{4\lambda+1}\cdot M^2$$ where the prime $p\equiv 1\text{ mod }4}$ and $\gcd(p,M)=1$. We assume the factorization $$n=p^{4\lambda+1}\prod_{\substack{q\text{ prime} \\ q\mid M}}q^{2e_q}$$ then it known the following Fact. Using Euler's theorem for odd perfect numbers, the condition to be perfect is written as $$2\cdot p^{4\lambda+1}\prod_{\substack{q\text{ prime} \\ q\mid M}}q^{2e_q}=\frac{p^{4\lambda+2-1}}{p-1}\prod_{\substack{q\text{ prime} \\ q\mid M}}\frac{q^{2e_q+1}-1}{q-1},$$ since the sum of divisor function is multiplicative. I've used the Lemma 7.9 from page 73 from Krizek, Luca and Somer, 17 Lectures on Fermat Numbers From Number Theory to Geometry Springer CMS (2001) (you can see this page as a Google Book, type here in Google Books the words: 17 lectures on fermat numbers lemma 7.9) with the specialization $y=q^{2e_q+1}-1$ and $x=p-1$ on assumption of the following Hypothesis A. For all prime $q\mid M$ $$p<q^{2e_q+1}-1$$ holds. I don't know if the following simple calculations were interesting. My purpose was do calculations and ask you Question. It is possible discard from the literature previous Hypothesis A as true? What other different hypothesis could be interesting to work from my approach? I am asking with my first question, if my hypothesis to work was absurd, and with the second question what hypotheis can I work with following calculations. I hope heuristics, that you cite the literature or mathematical reasonings. Thanks in advance. My approach was take the sum over all prime $q\mid M$ from the result of the specialization of cited Lemma 7.9, to show $$\frac{\log(p-1)}{\log p}\sum_{q\mid M}\log q^{2e_q+1}<\sum_{q\mid M}\log (q^{2e_q+1}-1),$$ and secondly take logarithms in Fact to combine showing that Claim. On assumption of Hypothesis A one has $$\log(2p^{4\lambda+1})+\sum_{q\mid M}\log q^{2e_q}>\log\frac{p^{4\lambda+2}-1}{p-1}+\frac{\log(p-1)}{\log p}\sum_{q\mid M}\log q^{2e_q+1}-\sum_{q\mid M}\log(q-1).$$ I am not saying that previous claim be interesting.

论坛问题帖 | has_answers | user243301 | 2016-07-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018648 | MSE-1861380

Does every power of two arise as the difference of two primes?

Conjecture: For each $n\in\mathbb N$ there are primes $q<p$ with $p-q=2^n$. Verified for $n\leq 26$: n p q 0 3 2 1 5 3 2 7 3 3 11 3 4 19 3 5 37 5 6 67 3 7 131 3 8 263 7 9 523 11 10 1031 7 11 2053 5 12 4099 3 13 8209 17 14 16421 37 15 32771 3 16 65539 3 17 131101 29 18 262147 3 19 524341 53 20 1048583 7 21 2097169 17 22 4194371 67 23 8388619 11 24 16777259 43 25 33554473 41 26 67108961 97 Proofs or counterexamples may be far away, but is something known about this topic?

论坛问题帖 | has_answers | Lehs | 2016-07-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018649 | MSE-1862245

Does this category have a name? (Relations as objects and relation between relations as morphisms)

Given two relations $R\subseteq A\times B$ and $R'\subseteq A'\times B'$. Is it known/used that every relation $r\subseteq R\times R'$ can be characterized by two relations $\alpha\subseteq A\times A'$ and $\beta\subseteq B\times B'$ so that $((a,b),(a',b'))\in r \iff \Big((a,a')\in\alpha\wedge (b,b')\in\beta\wedge (a,b)\in R\implies (a',b')\in R'\Big)$ and if $R''\subseteq A''\times B''$, $\,r'\subseteq R'\times R''$, where r' is characterized by $\alpha'\subseteq A'\times A''$ and $\beta'\subseteq B'\times B''$, then the composition $r'\circ r$ is characterized by the relations $\alpha'\circ\alpha\subseteq A\times A''$ and $\beta'\circ\beta\subseteq B\times B''$? (Where $\circ$ denote the composition of relations). $\;$In diagram form: $\require{AMScd}$ \begin{CD} A@>\alpha>>A'@>\alpha'>>A''@. A@>\alpha'\circ\alpha>>A''\\ @VRVV r @VVR' V r'@VVR''V \quad\implies\quad @VRVV r'\circ r @VVR''V\\ B@>>\beta>B'@>>\beta'>B'' @. B@>>\beta'\circ\beta>B'' \end{CD} Does this category of relations as objects and relation between relations as morphisms have a name? The reason for my interest in this category: Suppose $A=B\times B$ and that $R\subseteq A\times B$ is the composition in a magma. Then the functions among the morphisms between two such objects defines magma morphisms $B\to B'$. Suppose $B=\mathcal P(A)$ and that $R\subseteq A\times B$ is the relation $(a,S)\in R\iff a\in\overline{S}$ for some topology on A. Then the functions among the morphisms between two such objects define continuous functions $A\to A'$. Edit: The category of relations as objects and pairs of relations defined by some α,β (as above) as morphisms, is a subcategory of the category of relations as objects and relations of relations as morphisms. Very few relations between relations can be defined with two relations α,β, but those relations which can seems to be an interesting category.

论坛问题帖 | has_answers | Lehs | 2016-07-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018650 | MSE-1863300

Does the sequence $q(n)=3n+1+\frac{1-(-1)^n}{2}$ generate all the prime numbers?

Define $$q(n)=3n+1+\frac{1-(-1)^n}{2} \quad, \quad n\in \mathbb N.$$ $$1,5,7,11,13,17,19,23,25,29,31,35,\dots$$ It seems like this formula gives all primes >3 (although not just primes of course), which is verified for all $n<1000$. Is it provable or are there counter examples?

论坛问题帖 | has_answers | Lehs | 2016-07-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018651 | MSE-1865207

Conjecture: Every prime number is the difference between a prime number and a power of 2

Conjecture: $\forall p\in\mathbb P\exists q\in\mathbb P\exists n\in \mathbb N: q-p=2^n$ Verified for the 100 first primes.

论坛问题帖 | has_answers | Lehs | 2016-07-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018652 | MSE-1867875

For each prime $p>3$ there are non twin primes q,r with $p^3=2q+r$

Define $\mathbb P'=\{n\in\mathbb P|n-2,n+2\notin \mathbb P\}$. Conjecture: Given a prime $p>3$, then $\exists q,r\in\mathbb P':p^3=2q+r.$ Tested for the first 10000 primes. The solutions are not unique in general: 5 set-of-solutions cardinality . 1 ok 7 set-of-solutions cardinality . 1 ok 11 set-of-solutions cardinality . 10 ok 13 set-of-solutions cardinality . 23 ok 17 set-of-solutions cardinality . 41 ok 19 set-of-solutions cardinality . 38 ok 23 set-of-solutions cardinality . 82 ok 29 set-of-solutions cardinality . 166 ok 31 set-of-solutions cardinality . 171 ok 37 set-of-solutions cardinality . 253 ok 41 set-of-solutions cardinality . 391 ok 43 set-of-solutions cardinality . 411 ok 47 set-of-solutions cardinality . 604 ok 103 set-of-solutions cardinality . 4130 ok 107 set-of-solutions cardinality . 5755 ok 503 set-of-solutions cardinality . 382264 ok Calculated using BigZ

论坛问题帖 | has_answers | Lehs | 2016-07-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018653 | MSE-1869835

Even natural numbers are sums of two primes with twins or of two primes without twins

I seems to be very few even numbers that can't be written as a sum of two primes with twins or as a sum of two primes without twins. That is, suppose that $\mathbb P'$ is the set of the primes not belonging to a twin and that $\mathbb P''$ is the set of the primes belonging to a twin, then all but a finite number of even natural numbers can be written as $2n=p'+q',\quad p',q'\in\mathbb P'$ or as $2n=p''+q'',\quad p'',q''\in\mathbb P''$, due to a conjecture. So far the only exceptions are $0,2,96,98$ of even numbers $<100,000,000$. I'm interested in counterexamples and references. Below the number of ways to write 1000 as sum of two twin primes is 3, and the number of ways to write 1000 as sum of two non twin primes is 10. 100 terms cardinality . cardinality . 1 2 ok 1000 terms cardinality . cardinality . 10 3 ok 10000 terms cardinality . cardinality . 63 10 ok 100000 terms cardinality . cardinality . 415 64 ok 1000000 terms cardinality . cardinality . 3455 167 ok 10000000 terms cardinality . cardinality . 27081 1259 ok 100000000 terms cardinality . cardinality . 210124 5854 ok I see now that it seems to be a conjecture for the case $2n=p'+q',\quad p',q'\in\mathbb P'$, but then with a larger number of small exceptions: {0,2,4,6,8,10,12,14,16,18,20,22,24,26,28,30,32,34,36,38,40,42,44,46,48,50,52,54,56,58,62,64,66,68,72,78,80,82,86,88,92,96,98,108,110,118,122,124,128,138,140,148,152,170,182,188,208,212,218,222,232,238,266,268,272,282,292,296,328,332,358,362,392,482,512,628}. The corresponding set of small exceptions in the case $2n=p''+q'',\quad p'',q''\in\mathbb P''$ is: {0,2,94,96,98,400,402,404,514,516,518,784,786,788,904,906,908,1114,1116,1118,1144,1146,1148,1264,1266,1268,1354,1356,1358,3244,3246,3248,4204,4206,4208}. And the intersection is {0,2,96,98}.

论坛问题帖 | no_answers_at_retrieval | Lehs | 2016-07-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018654 | MSE-1870482

If $|G|=p^3q^2$ then $\Phi(G)$ is cyclic for primes $p\neq q$.

I have conjectured this result for the Frattini subgroup by doing some calculations in GAP. I think this is even true if $|G|=p_1^{i_1}\cdots p_n^{i_n}$ for $i_j\leq 3$ holds, but I would like to stick with this example first. I know the following: If $p\mid |G|$, then $p\mid |G/\Phi(G)|$, thus the order of $\Phi(G)$ is at most p^2q. Since $\Phi(G)$ is nilpotent, it must be abelian, too. That's why I think that it suffices to show this result for only abelian G (which is fairly easy), as considering all abelian groups with that order would lead to all possible abelian $\Phi(G)$. But I still need a valid argument for that. I also tried the assumption $\Phi(G)=C_p\times C_p$ or $\Phi(G)=C_p\times C_p\times C_q$ but I have not yet found a contradiction. I would be glad to see some ideas or approaches for this conjecture.

论坛问题帖 | has_answers | Nogard | 2016-07-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018655 | MSE-1874180

Any even $n> 10$ can be written as $n = pp'+p''$ for some primes p,p',p''

Conjecture: Any even integer $n>10$ can be written as $n=p\cdot p'+p''$, for some $p,p',p''\in\mathbb P$. Verified for all $n<1,000,000$. I first intended to post the weaker conjecture: $\mathbb P\subset\Big(\mathbb P\cdot\mathbb P+\mathbb P+1\Big)\cup\{2,3,5,11\}$.

论坛问题帖 | has_answers | Lehs | 2016-07-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018656 | MSE-1875948

Conjectured Compositeness Test for $E_n(b)=\frac{b^{2^n}+1}{2}$

How to prove that following conjecture is true? Definition Let $P_m(x)=2^{-m}\cdot \left(\left(x-\sqrt{x^2-4}\right)^{m}+\left(x+\sqrt{x^2-4}\right)^{m}\right)$, where m and x are nonnegative integers. Conjecture Let $E_n(b)=\Large\frac{b^{2^n}+1}{2}$ such that $n>1$, b is odd number greater than one . Let $S_i=P_b(S_{i-1})$ with $S_0=P_{b}(6)$, thus If $E_n(b)$ is prime, then $S_{2^n-1} \equiv 6\pmod{E_n(b)}$. I have checked this conjecture for all odd $b \in [3,1999] $ with $n \in [2,10]$. Searching for counterexample (PARI/GP) : CEGF(g1,g2)= { forstep(b=g1,g2,[2], for(n=2,10,my(s=Mod(2*polchebyshev(b,1,3),(b^2^n+1)/2)); for(i=1,2^n-1, s=2*polchebyshev(b,1,s/2)); if(!s==6 && isprime((b^2^n+1)/2),print("b=",b,"n=",n)))) } PARI/GP implementation of test . GFOdd(b,n)= { my(s=Mod(2*polchebyshev(b,1,3),(b^2^n+1)/2)); for(i=1,2^n-1,s=2*polchebyshev(b,1,s/2)); s==6 } Any hint would be greatly appreciated .

论坛问题帖 | has_answers | Pedja | 2016-07-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018657 | MSE-1878644

The relationship between the order of an element in a group and the order of its image

I am wondering if the following is true: Let G be a finite group, $g \in G$, and let $\phi : G \rightarrow H$ be a group homomorphism to another finite group H. Then $|g| = |\phi(g)| \cdot |k|$ for some element k in $\ker \phi$. To be specific, the element $k \in \ker \phi$ is taken to be $g^{|\phi(g)|}$. This clearly satisfies $\phi(g^{|\phi(g)|}) = \phi(g)^{|\phi(g)|} = e$. I came up with a possible proof of this statement, but I wanted to see if anyone had counter-examples or insights on this issue. The proof I had is to show that $|g|$ divides $|\phi(g) \cdot |k|$ and conversely, that $|\phi(g)|$ divides $\frac{|g|}{|k|}$, using the fact $|x|$ divides n iff $x^n = e$.

论坛问题帖 | no_answers_at_retrieval | Tac-Tics | 2016-08-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018658 | MSE-1882613

Every prime number divide some sum of the first k primes.

Let $S_n=\Sigma^n_{k=1}p_k$, where p_k is the k-th prime number. Conjecture: $$\forall p\in\mathbb P\exists n\in\mathbb N: p|S_n$$ Verified for the 1000 first primes. Is there a proof for this result in general? In the diagram primes are projected on the x-axis and n (as in S_n) on the y-axis. As H.H.Rugh commented there is a stronger conjecture for all positive integers m. Below a table of n-records for different $m\in\mathbb Z^+$ (some of them primes): m n factorization of Sn 1 (1) 1 (2) 3 (3) 10 (3,43) 6 (2,3) 57 (2,3,5,229) 12 (2,2,3) 97 (2,2,3,1879) 18 (2,3,3) 113 (2,3,3,41,43) 35 (5,7) 180 (5,7,2531) 42 (2,3,7) 305 (2,2,2,3,7,7,239) 90 (2,3,3,5) 357 (2,2,2,3,3,3,5,367) 101 (101) 422 (5,101,1129) 137 (137) 861 (2,137,9739) 163 (163) 902 (5,7,11,47,163) 195 (3,5,13) 907 (2,3,3,5,13,2551) 202 (2,101) 1207 (2,2,19,101,719) 222 (2,3,37) 1359 (2,2,2,3,3,37,2671) 252 (2,2,3,3,7) 1683 (2,2,3,3,7,17,37,71) 326 (2,163) 1765 (2,2,3,23,163,277) 474 (2,3,79) 2077 (2,2,2,3,5,79,1861) 504 (2,2,2,3,3,7) 2133 (2,2,2,3,3,3,7,53,233) 522 (2,3,3,29) 2379 (2,3,3,3,3,3,7,29,239) 643 (643) 2529 (2,3,3,11,211,643) 647 (647) 3092 (11,647,5791) 658 (2,7,47) 3353 (2,3,7,43,47,577) 700 (2,2,5,5,7) 3593 (2,2,5,5,7,103,787) 817 (19,43) 4683 (2,3,11,19,43,1847) 995 (5,199) 5329 (2,3,5,17,199,1291) 1004 (2,2,251) 6415 (2,2,2,251,96643) 1204 (2,2,7,43) 6533 (2,2,2,2,7,7,31,43,193) 1459 (1459) 7241 (2,2,3,3,5,5,191,1459) 1488 (2,2,2,2,3,31) 7307 (2,2,2,2,2,3,31,85909) 1610 (2,5,7,23) 8079 (2,5,7,23,43,4567) 1677 (3,13,43) 10171 (2,2,2,3,3,3,13,43,4259) 1870 (2,5,11,17) 10331 (2,5,11,17,71,4003) 2035 (5,11,37) 11459 (2,2,3,3,5,11,37,9029) 2616 (2,2,2,3,109) 11753 (2,2,2,3,3,3,3,3,37,89,109) 2672 (2,2,2,2,167) 18137 (2,2,2,2,7,167,93047) 3420 (2,2,3,3,5,19) 21709 (2,2,3,3,3,5,13,19,137,139) 3830 (2,5,383) 27617 (2,5,53,383,20749) 4232 (2,2,2,23,23) 38861 (2,2,2,2,3,3,23,23,113189) 7394 (2,3697) 45381 (2,107,3697,15083) 7450 (2,5,5,149) 47323 (2,3,3,5,5,7,41,149,677)

论坛问题帖 | has_answers | Lehs | 2016-08-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018659 | MSE-188362

Integer solutions to $2^x+3^y=9^z+8^w+8^t$

currently in number theory is there a method to solve for integer solutions to equations like $2^x+3^y=9^z+8^w+8^t$? For example, $2^{13}+3^6=9^3+8^4+8^4$. (obtained using computer brute force) In general, is there a way to solve for ${A_1}^{x_1}+{A_2}^{x_2}+\ldots+{A_n}^{x_n}={B_1}^{y_1}+{B_2}^{y_2}+\ldots+{B_m}^{y_n}$ or is it out of scope of the current number theory methods? (A_i and B_i are the given numbers, want to solve for x_i and y_i) Sincere thanks for help!

论坛问题帖 | has_answers | yoyostein | 2012-08-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018660 | MSE-1884143

About the distribution of Gaussian prime pairs

The primes in the ring of Gaussian integers $\mathbb Z[i]$ has a simple connection to the ordinary primes: either one of the components is zero and the other is a prime of type $4n+3$ or the norm $\|a+ib\|=a^2+b^2$ is a prime $a\ne 0\ne b$, that is a prime of the type $4n+1$. All $n\in\mathbb Z^+$ with $n=a^2+b^2$, e.g. all ordinary primes of the form $4n+1$, are composites in $\mathbb Z[i]$: $n=(a+ib)(a-ib)$. There is essentially one pair of Gaussian primes such that $|z_1-z_2|=1$, and that is $(1+i,2+i)$. If defining prime pairs in $\mathbb Z[i]$ with $|z_1-z_2|=2$, then it seems like for each integer $n>0$ there is a prime $n+im\in\mathbb Z[i]$ with a prime as a pair. Can this be proved? Verified for all $a+ib\in\mathbb Z[i]$ with $0<a,b\leq1000$. I'll elaborate. If $|z_1-z_2|=2$ one can assume e.g. that $\Re (z_1-z_2)=0$ and $|\Im (z_1-z_2)|=2$. Hence, the conjecture can be formulated as: For each $a\in\mathbb N^+$ it exists $b\in\mathbb Z$ such that $a+ib$ and $a+i(b+2)$ are Gaussian primes. Equivalent conjecture: $\forall n\in\mathbb N^+\exists m\in\mathbb Z: n^2+m^2,n^2+(m+2)^2\in\mathbb P$.

论坛问题帖 | has_answers | Lehs | 2016-08-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018661 | MSE-1887562

Counting even and odd numbers in the columns of a triangular arrangement of the integers

I write the positive numbers starting at 1 in a triangle:$$\mathbb{N}_\triangle = \begin{matrix} &&&&&21&\ldots \\ &&&&15&20&\ldots \\ &&&10&14&19&\ldots \\ &&6&9&13&18&\ldots \\ &3&5&8&12&17&\ldots \\ 1&2&4&7&11&16&\ldots \end{matrix}$$ I would like some help proving the following claim I denote the count of even numbers in column n of $\mathbb{N}_\triangle$ by $e(n)$. Then for $n>1$ $$e(n) = \Bigg\lfloor{n+2 \above 1.5pt 4}\Bigg\rfloor+\Bigg\lfloor{n+1 \above 1.5pt 4}\Bigg\rfloor$$ Note the n^{th}-column$ of $\mathbb{N}_\triangle$ has n integers so I can set $f(n)$ to count the number of odd integers in $\mathbb{N}_\triangle$ and $$f(n) = n-e(n)$$ It appears that $e(n)\in$ A004524 This is being driven by hunches and computer checks. $e(n)$ can be simplified but I think it is easier to remember it the way it is written.

论坛问题帖 | has_answers | Anthony | 2016-08-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018662 | MSE-1889549

Counting multiple of 3 in a triangular arrangement of the integers.

I write :$$\mathbb{N}_\triangle = \begin{matrix} &&&&&21&\ldots \\ &&&&15&20&\ldots \\ &&&10&14&19&\ldots \\ &&6&9&13&18&\ldots \\ &3&5&8&12&17&\ldots \\ 1&2&4&7&11&16&\ldots \end{matrix}$$ Can we determine how many numbers in any given column of $\mathbb{N}_\triangle$ are divisible by 3. I need help proving the following claim: If $f_3(n)$ counts the numbers in the n^{th} column of $\mathbb{N}_\triangle$ that are divisible by 3 then $$f_3(n) =\Bigg\lfloor{n+2 \above 1.5pt 3}\Bigg\rfloor$$ For example $$f_3(6) = \Bigg\lfloor{6+2 \above 1.5pt 3}\Bigg\rfloor=\Bigg\lfloor{8 \above 1.5pt 3}\Bigg\rfloor=\Bigg\lfloor 2.66666\Bigg\rfloor =2$$ and surely there are 2 numbers in column six that are divisible by 3. I suspect there are no multiples of 3 on the base of the triangle since possibly $$\Bigg(3, 1+ {n(n-1) \above 1.5pt 2}\Bigg)=1$$ In fact if $(3,T_n)=1$ then I suspect that there are no numbers in the n^{th} row that are divisible by 3. Surely there are consecutive triangular numbers divisible by 3.

论坛问题帖 | has_answers | Anthony | 2016-08-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018663 | MSE-1890264

Conjectured Compositeness Test for Generalized Fermat Numbers

How to prove that following conjecture is true? Definition Let $P_m(x)=2^{-m}\cdot \left(\left(x-\sqrt{x^2-4}\right)^{m}+\left(x+\sqrt{x^2-4}\right)^{m}\right)$, where m and x are nonnegative integers. Conjecture Let $F_n(b)=b^{2^n}+1$ such that $n \ge 2$ and b is even number . Let $S_i=P_b(S_{i-1})$ with $S_0=P_{b}(6)$, thus If $F_n(b)$ is prime, then $S_{2^n-1} \equiv 2\pmod{F_n(b)}$. I have checked this conjecture for all even $b \in [2,2000] $ with $n \in [2,10]$. PARI/GP implementation of test : CTGF(b,n)= { my(s=Mod(2*polchebyshev(b,1,3),b^2^n+1)); for(i=1,2^n-1,s=2*polchebyshev(b,1,s/2)); s==2 } Any hint would be greatly appreciated .

论坛问题帖 | has_answers | Pedja | 2016-08-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018664 | MSE-1897785

A conjecture about integer polynomials and primes

Conjecture: For all sets of non constant polynomials $\{f_1,\dots,f_n\}\subset \mathbb Z[X]$, it exists $m\in\mathbb N$ such that $f_k(m)\notin\mathbb P$, for all $1\leq k\leq n$. Where $\mathbb P$ is the set of primes. Can this be proved? The original conjecture with $\mathbb R[X]$ seems too difficult and may depend on an open conjecture.

论坛问题帖 | has_answers | Lehs | 2016-08-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018665 | MSE-1898800

A conjecture about primes and holomorphic functions

Inspired by this question and this answer I raise the following conjecture: If $\{f_1,\dots,f_n\}$ is a set of entire holomorphic functions and if for every $m\in\mathbb N$, it exists a function with $f_i(m)\in\mathbb P$, then some function f_k is a constant. Can this be proved?

论坛问题帖 | has_answers | Lehs | 2016-08-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018666 | MSE-1899443

An accountancy of the natural numbers

I guess this formula is known since I believe it's true $$\sum_{k=2}^\infty\frac{(-1)^{1+\Omega(k)}}{k}=1$$ where $\Omega(k)$ is the number of prime factors of k (not necessary different primes). I can't find it and want a formal proof or a reference. My intuition about this sum is something like this: The probability of a number to have the prime factor p is $\frac{1}{p}$ and to have the prime factors p or q is $\frac{1}{p}+\frac{1}{q}-\frac{1}{pq}$ etc. The probability of a number >1 to have a prime factor is 1. Is this sum really not known? Haven't anything been published about it before? I finally realize that the series $$\sum_{k=1}^\infty\frac{(-1)^{\Omega(k)}}{k}=0$$ has been known for a long time, and that solves my confusion.

论坛问题帖 | has_answers | Lehs | 2016-08-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018667 | MSE-1900947

Rearrange convergent series

It was long time ago I studied analyze and I do have forgot a lot, but isn't it so that any convergent series $$\sum_{i=1}^\infty a_i$$ can be rearranged to an absolut convergent series $$\sum_{i=1}^\infty \left|\sum_{j=1}^{n_i} b_{ij}\right|=\sum_{k=1}^\infty a_k$$ where the b_{ij}-s is a rearrangement of the a_k-s?

论坛问题帖 | no_answers_at_retrieval | Lehs | 2016-08-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018668 | MSE-1902386

A series with the non square free numbers $q_n:\$ $\sum_{k=1}^\infty\frac{(-1)^{1+\Omega(q_k)}}{q_k}=0$

Let q_n be the n-th natural number that can be divided by a square >1 (https://oeis.org/A013929). Conjecture: $$\sum_{k=1}^\infty\frac{(-1)^{1+\Omega(q_k)}}{q_k}=0$$ where $\Omega(n)$ is the number of (not necessarily different) prime factors of n. I'm sure that the conjecture is true, but I can't prove it and that's what I need help with. The conjecture is related with An accountancy of the natural numbers. The corresponding series for squarefree numbers s_n $$\sum_{k=2}^\infty\frac{(-1)^{1+\Omega(s_k)}}{s_k}=1$$

论坛问题帖 | has_answers | Lehs | 2016-08-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018669 | MSE-190666

how to prove this extended prime number theorem?

A Generalized Prime Number Theorem? Conjecture Let n and k be positive integers with $n - 50 > k^2 > 0$ and n sufficiently large. Then for the odd primes we have, when p is the biggest odd prime $\le n$, $$ 3^k + 5^k + 7^k + 11^k + ... + p^k \sim \frac{n^{k+1}}{(k+1) (\log(n) - \log(k)) } $$ I wonder if you guys have seen it before ? How to prove it ? Any useful references for $k > 1$?

论坛问题帖 | has_answers | mick | 2012-09-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018670 | MSE-1910024

$P^2 + P + 1$ is prime if P is prime.

I just thought of it. I don't know if there is one such conjecture or a proven problem or if this is a new conjecture. If it already exists where can I find this problem. And most importantly how to prove it??

论坛问题帖 | has_answers | jnyan | 2016-08-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018671 | MSE-1922972

Who is credited with the conjecture that all odd squares are solitary?

Let $\sigma(x)$ be the sum of the divisors of the positive integer x. For example, $\sigma(6)=1+2+3+6=12$ and $\sigma(28)=1+2+4+7+14+28=56$. Denote the abundancy index I of x by $$I(x)=\frac{\sigma(x)}{x}.$$ If a positive integer y is one of at least two solutions of $$I(y)=\frac{a}{b}$$ for a given rational number a/b, then y is called a friendly number. Here is a formal definition: DEFINITION. Let x and y be distinct positive integers. If x and y satisfy the equation $I(x)=I(y)$ then (x,y) is called a friendly pair. Each member of the pair is called a friendly number. (In other words, x is a friend of y, and y is also a friend of x.) A number which is not friendly is called a solitary number. Checking the first few odd squares N gives us $\gcd(N,\sigma(N))=1$, which by Greening's Theorem means that such N are solitary. Since Greening's Theorem that $\gcd(N,\sigma(N))=1$ implies that N is solitary. is sufficient (but not necessary), "it's easy to find odd squares for which $\gcd(n, \sigma(n))$ is not equal to 1; e.g. if $n = {21}^2 = 441$ then $\gcd(n, \sigma(n)) = 3$. But 441 is solitary." (Dean Hickerson, personal communication via e-mail, Mar. 12, 2011) Here is my question: Who is credited with the conjecture that all odd squares are solitary?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2016-09-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018672 | MSE-1923695

If $N = q^k n^2$ is an odd perfect number with Euler prime q, does the following implication hold?

Let $\sigma(x)$ denote the sum of the divisors of x. For example, $\sigma(6)=1+2+3+6=12$. Numbers N satisfying the equation $\sigma(N)=2N$ are called perfect numbers. Euler showed that an odd perfect number must have the form $N = q^k n^2$ where q is prime with $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. Here is my question: If $N = q^k n^2$ is an odd perfect number with Euler prime q, does the following implication hold? $$\bigg(q^k < n \iff \sigma(q^k) < \sigma(n) \iff \dfrac{\sigma(q^k)}{n} < \dfrac{\sigma(n)}{q^k}\bigg) \implies \left(q^k < n\right)$$ From an answer to this earlier MSE question, I know that the converse $$\left(q^k < n\right) \implies \bigg(q^k < n \iff \sigma(q^k) < \sigma(n) \iff \dfrac{\sigma(q^k)}{n} < \dfrac{\sigma(n)}{q^k}\bigg)$$ holds.

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2016-09-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018673 | MSE-1923741

Irrationality of the Riemann zeta-function on odd arguments

It is proved that infinitly many of the numbers $\zeta(2n+1)$ are linearly independent over $\mathbb{Q}$. Is there a conjecture such as 'the numbers $\zeta(2n+1)$ are linearly independent over $\mathbb{Q}$'? Thank you!

论坛问题帖 | has_answers | user368031 | 2016-09-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018674 | MSE-1926517

A property of v

Let n be a natural number. Denote by $v(n)$ the sum of the numbers appearing in the canonical factorization of n (like in A curious property of integers). Then I believe that for $n>1$ the sequence $ n, v(n), v(v(n)), v(v(v(n))),\ldots $ is eventually the same prime number. Was this stated anywhere?

论坛问题帖 | has_answers | yhsai | 2016-09-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018675 | MSE-1931161

A general criterion of convergence of series?

Given a non degenerated function $f:\mathbb N^+\to\mathbb Z_2 $. (It doesn't stabilize and become constant for all n larger than some natural number). Define the function $\sigma(i,n,f)=|f^{-1}(i)\cap\mathbb Z_n|\,$ and $c_n= \left\{ \begin{array}{l} a_{\sigma(o,n,f)}\;\text{ if } f(n)=0\\ b_{\sigma(1,n,f)}\;\text{ if } f(n)=1 \end{array} \right. $ Then $(c_n)_n$ is a shuffle of $(a_n)_n$ and $(b_n)_n$ for each function f as above. All the terms a_i and b_i occur exactly once in the sequence $c_1,c_2,\dots$ in preserved index order. I would like to formulate this conjecture: If $\displaystyle a=\sum^\infty_{i=1} a_i\;$ and $\displaystyle \; b=\sum^\infty_{i=1} b_i\;$ are convergent and c_i arise from a shuffle of a_i and b_i, then $\displaystyle a+b=\sum^\infty_{i=1} c_i$. I'm asking for a proof of the conjecture or a counterexample and will award the three first correct answers with a bounty worth 200.

论坛问题帖 | has_answers | Lehs | 2016-09-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018676 | MSE-1932639

Conjecture: if $a_{n+1}-a_{n}=c $ $\forall n\in\mathbb{N}$ and $b_{n+1}-b_{n} \rightarrow \infty$ then $a_{n}/b_{n} \rightarrow 0$

Is the following proposition true or false? Given two positive and strictly increasing sequences a_n and b_n, if $a_{n+1}-a_{n}=c \in \mathbb{R}$ for every $n\in \mathbb{N}$ and $b_{n+1}-b_{n} \rightarrow \infty$, then $\frac{a_n}{b_n}$ converges to 0

论坛问题帖 | has_answers | Adrian | 2016-09-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018677 | MSE-1933452

A conjecture on tensor product vectors

Consider a real vector space $V^{(1)}\otimes V^{(2)}$ where $\otimes$ is the tensor product. Product vectors are of the form $v_1\otimes v_2$ where $v_1\in V^{(1)}$, $v_2\in V^{(2)}$, anything else is a non-product vector. I conjecture the following I have a set of product vectors $a_i\otimes b_i$, $i=1,\dots,n$ that sum to make a product vector $A\otimes B$, $A\otimes B=\sum_i \alpha_i a_i \otimes b_i$ where α_i are real, non-zero constants and there are no repetitions of a_i. All of these product vectors must have $b_i= \lambda \, B$ $\forall$ i, where λ is some non-zero real. I believe this is true, but am not sure how to proceed. For example, if $n=2$, then $A\otimes B=a_1\otimes b_1 + a_2 \otimes b_2$ is not a product vector unless $a_1=a_2$ or $b_1 =b_2$, and then it is clear that either $a_1=a_2=A$ or $b_1=b_2=B$

论坛问题帖 | has_answers | jdizzle | 2016-09-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018678 | MSE-1937193

Conjecture $\sum_{n=1}^\infty\frac{n^2}{(-1)^n \cosh(\pi n\sqrt{3})-1}=\frac{1}{12\pi^2}$

Wolfram Alpha numerical calculation shows that the quantity $$ \sum_{n=1}^\infty\frac{12\pi^2n^2}{(-1)^n \cosh(\pi n\sqrt{3})-1} $$ is 1 with high accuracy. Can anybody prove the resulting conjecture: $$ \sum_{n=1}^\infty\frac{n^2}{(-1)^n \cosh(\pi n\sqrt{3})-1}\overset{?}{=}\frac{1}{12\pi^2} $$

论坛问题帖 | has_answers | Tyrell | 2016-09-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018679 | MSE-1938947

Is any mathematican more famous for their conjecture(s) than their theorem(s)?

I'm wondering if some mathematicians gained more fame from their (perhaps visionary) conjectures, than from the positive results they proved? I would say this is not true of Fermat, despite his famous eponymous conjecture (now settled), because he established so many results independent of his conjecture. And this seems not true of Poincaré, whose famous conjecture (also now settled) bears his name. But he was incredibly accomplished independent of that conjecture. Atiyah has formulated wonderfully productive conjectures (one leading to a Witten advance), but he has also established major results, e.g., the Atiyah-Singer theorem. I am interested to explore whether some mathematicians have specific conjecture-talent that is not evidently reflected in theorem-proving talent.

论坛问题帖 | has_answers | Joseph O'Rourke | 2016-09-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018680 | MSE-1940934

$S_n=\sum^n_{k=1} (-1)^{\Omega(q_k)}\cdot{q_k}$ change sign infinitely many times?

Let q_k be the k-th positive natural number that is divisible by a square larger than one. How to prove that $S_n=\sum^n_{k=1} (-1)^{\Omega(q_k)}\cdot{q_k}$ change sign infinitely many times as n increases? $\Omega(n)$ is the number of (not necessarily different) prime factors of n. I have made some numerical experiments that indicates this conjecture. Adequate answers will be rewarded with bounties.

论坛问题帖 | no_answers_at_retrieval | Lehs | 2016-09-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018681 | MSE-1973585

Does a basis b exist such that only a finite number of palindromes in base b are primes?

Can we prove that a base b must necessarily exist so that only a finite number of palindromes in base b are primes? My friend came up with this, I hope it isn't one of those ubiquitous "innocent" number theory problems which are years beyond any hope for solution. So far our ideas have not yielded any fruits: The number of palindromes under n is roughly $\sqrt{n}$ The number of primes under n is roughly $\frac{n}{\log n}$ Any such prime must have odd length (work $\bmod b+1$)

论坛问题帖 | no_answers_at_retrieval | Asinomás | 2016-10-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018682 | MSE-1978786

Solving a high school conjecture

Right, so this was something I stumbled upon whilst looking at forms of the golden ratio (if you replace x with one in the diagram above then you get phi) and then I wondered about what if you messed with the value of x, did a massive amount of trial and error (lost in a notebook somewhere) then came up with the equation in the right, which worked exactly for every value I tested (about up to 100) Now I have found stack exchange, in wondering if there is a way to prove this, I plugged in some values into wolfram alpha and geogebra, which seems to show it works, but I want a proof, not just trial and error answers.

论坛问题帖 | has_answers | Cursed | 2016-10-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018683 | MSE-1987944

Complete monotonicity of a sequence related to tetration

Let Δ denote the forward difference operator on a sequence: $$\Delta s_n = s_{n+1} - s_n,$$ and Δ^m denote the forward difference of the order m: $$\Delta^0 s_n = s_n, \quad \Delta^{m+1} s_n = \Delta\left(\Delta^m s_n\right).$$ We say that a sequence s_n is completely monotone iff $$(-1)^m \Delta^{m} s_n > 0,$$ i.e. the sequence s_n itself is positive and decreasing, its first differences Δs_n are negative and increasing (that is, decreasing in absolute values), its second differences $\Delta^2 s_n$ are positive and decreasing (like the sequence s_n itself), and so on. So, the sequence of second differences of a completely monotone sequence is also completely monotone (and so is the sequence of its differences of any even order). Let a be a real number in the interval $1<a<e^{1/e}$. Let ${^n a}$ denote tetration: $${^0 a}=1, \quad {^{(n+1)} a} = a^{\left({^n a}\right)}.$$ The sequence ${^n a}$ is strictly increasing and converges to the limit $${^\infty a} = \lim_{n\to\infty}\left({^n a}\right) = \frac{W(-\ln a)}{-\ln a},$$ where $W(x)$ is the Lambert W-function. Let $s_n = {^\infty a} - {^n a}$. It appears that the sequence s_n is completely monotone. How can we prove it?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2016-10-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018684 | MSE-1996047

$\forall m\in\mathbb N\exists n>m+1\exists N\in\mathbb N:m^2+n^2+(mn)^2=N^2$

Prove the conjecture or give a counter-example: For each $m\in\mathbb N$ there exist a $n>m+1$ such that $m^2+n^2+(mn)^2$ is a perfect square. I have just tried it out numerically and it holds for $m<1000$. I can't see any pattern for the smallest n: m n 1 12 2 8 3 18 4 32 5 50 6 72 7 98 8 30 9 162 10 200 11 242 12 119 13 338 14 392 15 450 16 512 17 578 18 105 19 722 20 800 21 208 22 968 23 1058 24 1152 25 1250

论坛问题帖 | has_answers | Lehs | 2016-11-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018685 | MSE-1999562

Recycling a false conjecture of Goldbach

The conjecture of Goldbach (1752) "Every odd integer can be written in the form $p+2a^2$ where p is a prime or 1 and a is a natural number (can be even 0)." has the smallest exception 5777. While not allowing $p=1$ the formula give the following exceptions less than $10,000,000$: 1 5777 5993 4946453 5293163 5651423 6017447 6098663 6392393 6420563 6429467 7071803 7072865 7081223 7103447 7217417 7277753 7341677

7499447 7532687 7571537 7642637 7727657 7809383 7825643 7844183 7860815 7935173 7970267 8063393 8111033 8226863 8286527 8344247 8409437 8464367 8498393 8542397 8552147 8562857 8654093 8725973 8772227 8848883 8851403 8863703 8865593 8867663 8869907 8885675 8952797 8957747 8971775 8985503 9166637 9200573 9233333 9309857 9325103 9341147 9409787 9447533 9470147 9481763 9492203 9519269 9534653 9552155 9553367 9564347 9707303 9760307 9843275 9882023 9897017 9911393 9916337 9948617 9958307 It's trivially true that all odd numbers greater than 1 can be written in the form $p+2a$, p prime, and I therefore conjecture: There is a real number $\gamma>1$ such that every odd number greater than 1 can be written in the form $p+2\lfloor a^\gamma\rfloor$. Where $p\in\mathbb P$ and $a\in\mathbb N$ Is this possible to prove? For $\gamma=1.5\;$ I found no exceptions less than $100,000,000$.

论坛问题帖 | has_answers | Lehs | 2016-11-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018686 | MSE-2004746

Make a conjecture about the values of k for which multiplicative inverses exist for every element in $\mathbb{Z}_k$

I know that $gcd(k, \bar{k}) = 1$ for k to have a multiplicative inverse. Can someone give me pointers where to start?

论坛问题帖 | no_answers_at_retrieval | phoxd | 2016-11-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018687 | MSE-2006506

High School Calculus: Find cut out length x which gives Maximum volume of can.

This is a pretty long question so I will try to shorten it as much as possible. Part A: Given: a square piece of tin plate with side length l, cut out corners with shape of square of side length x. Fold the resulting shape into an open-ended cuboid. Determine x for maximum volume (deduce conjecture) Next, prove, using the principles of calculus (and the quadratic formula),the conjecture. Pictues of the problem that I drew: https://i.sstatic.net/YvXk5.png What I did: $V(x)=Length×Width×Height$ (Equation for cuboid) $V(x)=x(l-2X)(l-2x)$ (substitute known values) $=x[(l-2X)]^2$ $=x[(l-2X)]^2$ $=x(l^2-4lx+4x^2)$ $=l^2 x-4lx^2+4x^3$ $V'(x)=l^2-8lx+12x^2$ (differentiate) $0=l^2-8lx+12x^2$ ($V'(x)=0$ to find out maximum/minimum) $0=(2x-l)(6x-l)$ (factorise) $2x-l=0$ $2x=l$ $x=l/2$ (first answer is not correct, if $x=1/2$ then cuboid cant even be made) $6x-l=0$ $6x=l$ $x=l/6$ (correct answer) I am unable to do the next part, Im not sure if i am overthinking it or not. But I cant seem to figure out how do i prove this conjecture? Haven't i already proved it? Part B https://i.sstatic.net/jxYWf.png It ask the same thing as above x when volume is at maximum, however, as you can see from this picture. This time it is a rectangular plate with side length r and s. What I did: $V(x)=x(r-2x)(s-2x)$ (substitute values into equation) $=x(rs-2rx-2sx+4x^2)$ $=rsx-2rx^2-2sx^2+4x^3$ $V'(x)=rs-4rx-4sx+12x^2$ (differentiate) $=rs-4x(r+s)+12x^2$ $=12x^2-4x(r+s)+rs$ $0=12x^2-4x(r+s)+rs$ (set $V'(x)$ as 0 to find out maximum/minimum) $x=(--4(r+s)±√([(-4(r+s))]^2-4\times12rs))/(2\times12)$ (use quadratic formula as it is impossible to factorize) $x=(4(r+s)±√([(-4(r+s))]^2-48rs))/24$ $=(4(r+s)±√([16(r+s)]^2-48rs))/24$ $=(4(r+s)±√([16[(r+s)]^2-3rs]))/24$ $=(4(r+s)±4√([(r+s)^2-3rs]))/24$ $=((r+s)±√([(r+s)]^2-3rs))/6$ $=((r+s)±√(r^2+2rs+s^2-3rs))/6$ $=(r+s±√(r^2-rs+s^2))/6$ And from what I think (only when the $+√(r^2-rs+s^2$)gives the maximum volume) But again, I am suppose to prove the conjecture that I just deduced. Just How? Any advice are welcome.

论坛问题帖 | no_answers_at_retrieval | Nin. than | 2016-11-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018688 | MSE-2021539

Is there a maximal prime number not being a sum of consecutive primes?

Let $\mathcal S=\{p\in\mathbb P|\exists m,n\in\mathbb Z_+:p=p_n+\cdots+p_{n+m}\}$. For "small" primes, not much greater than one million, it seems that about half of the primes belong to $\mathcal S$. But the greater primes the more combinations of consecutive primes available. There are $\pi(p)$ primes less than or equal to p and the number of available prime sequences is $1\cdot\pi(p)+2\cdot(\pi(p)-1)+3(\pi(p)-2)+\cdots+(\pi(p)-1)\cdot 2=$ $\displaystyle\frac{\pi(p)^3+3\pi(p)^2+4\pi(p)}{6}$ (If I calculated it right). I suppose it's possible to use the Prime Number Theorem to estimate the probability of $p\in\mathcal S$, but I'm pretty sure that I would mess it up. My questions are: Does $P(p\in\mathcal S)\to 1$ as $p\to\infty$? Can it be proved that there is a maximal prime number not being a sum of consecutive primes? See also How often is a sum of k consecutive primes also prime? which is a similar but not an equivalent question. This is what I got so far. It doesn't support my intuition, but I will let it run up to $10,000,000$ or more.

论坛问题帖 | has_answers | Lehs | 2016-11-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018689 | MSE-2024107

Is it true that, for two infinite functions, $\lim_{x->x_0}\frac{f(x)}{g(x)}=1$ entails $\lim_{x->x_0}[f(x)-g(x)]=0$?

Let f,g be real functions defined in a neighborhood of $x_0\in \mathbb{R}$ such that $g(x)>0$,$\lim_{x\rightarrow x_0}f(x)= \lim_{x\rightarrow x_0}g(x)=+\infty$, and $\lim_{x\rightarrow x_0}\frac{f(x)}{g(x)}=1$. Is it true that $$\lim_{x\rightarrow x_0}[f(x)-g(x)]=0?$$

论坛问题帖 | has_answers | Adrian | 2016-11-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018690 | MSE-2030706

Sets of natural numbers closed under addition of successors

Conjecture: Given integers $0<a_1<a_2,\dots\,$ without a common divisor, such that all sums $\sum_{k=r}^q a_k\in A=\{a_1,a_2,\dots\}$. Then there is an integer N such that for all integers $n>N$ it holds that $n\in A$. I would like to see a proof or a counter-example. I have computationally investigated some minimal sets A as above which includes the primes or the squares or the exponents of two or the faculties, and it seems like $\,\mathbb

Z_+\!\!\setminus\! A\,$ always is a finite set. If A is such a set that is minimal including the primes, then $\{n\in A|n \leq 97\}=$ $\{2,3,5,7,8,10,11,12,13,15,17,18,19,20,21,23,24,25,26,28,29,30,31,32,33,35,36,$ $37,39,40,41,42,43,44,45,46,47,48,49,50,51,52,53,54,56,57,58,59,60,61,63,64,$ $65,66,67,68,69,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,86,87,88,89,90,$ $91,92,93,94,95,96,97\}$ And if A is the minimal set including the faculty numbers $1,2,6,24,120,\dots\,$ then $\{n\in A|n \leq 120\}=$ $\{1,2,3,5,6,8,9,10,11,12,14,16,17,19,20,21,22,23,24,25,26,27,28,29,30,31,32,$ $33,34,36,37,38,39,40,41,42,43,44,45,46,47,49,50,51,52,53,54,55,56,57,59,60,$ $61,62,63,64,65,66,67,68,69,70,71,72,73,74,75,76,77,78,79,80,81,82,83,84,85,$ $86,87,88,89,90,91,93,94,95,96,97,98,99,101,102,103,104,105,106,107,108,109,$ $110,111,112,113,114,115,116,117,118,119,120\}$

论坛问题帖 | no_answers_at_retrieval | Lehs | 2016-11-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018691 | MSE-2038555

Why does the same convergent (relative to the period) of the continued fraction of $\sqrt{n}$ give solutions to the Pell equation?

Let n be a natural number which isn't a square and $$\sqrt{n}=[a_0;\overline{a_1,a_2\dotsb a_r}]$$ then consider the convergents $$\frac{x_1}{y_1}=[a_0;a_1,a_2\dotsb a_{r-1}]$$ $$\frac{x_2}{y_2}=[a_0;a_1,a_2\dotsb a_{r-1},a_r,a_1\dotsb a_{r-1}]$$ etc I conjecture that $x_k^2-ny_k^2=\pm 1$. And that all solutions of the positive and negative Pell equations have this form. If $p^2-nq^2=1$ then $0<\frac{p}{q}-\sqrt{n}<\frac{1}{2\sqrt{n}q^2}$. And if $p^2-nq^2=-1$ then $0<\sqrt{n}-\frac{p}{q}<\frac{1}{2\sqrt{n}q^2}$. In both cases $q^2\left|\frac{p}{q}-\sqrt{n}\right|<\frac{1}{2\sqrt{n}}$, so $\frac{p}{q}$ is in some sense a very good approximation of $\sqrt{n}$, and the last number in the period, a_r seems to always be $2\lfloor\sqrt{n}\rfloor$ and larger than all of $a_1,a_2\dotsb a_{r-1}$, so the next best rational approximation of $\sqrt{n}$ would have a "much" bigger denominator than q, so it should make sense for the solutions to the Pell equation to appear before a_r. $$\sqrt{397}=[19;\overline{1,12,3,4,9,1,2,1,2,1,1,2,1,2,1,9,4,3,12,1,38}]$$ $$\frac{20478302982}{1027776565}=[19;1,12,3,4,9,1,2,1,2,1,1,2,1,2,1,9,4,3,12,1]$$ This is the fundamental solution of the negative Pell equation $$20478302982^2-397\times 1027776565^2=-1$$ Getting the second convergent of this type $$\frac{838721786045180184649}{42094239791738433660}=[19;1,12,3,4,9,1,2,1,2,1,1,2,1,2,1,9,4,3,12,1,38,1,12,3,4,9,1,2,1,2,1,1,2,1,2,1,9,4,3,12,1]$$ Gives the fundamental solution to the positive Pell equation $$838721786045180184649^2-397\times 42094239791738433660^2=1$$ It is hard not to prove by induction that the convergents of this form alternate between generating solutions to the positive and negative equations, although very laborious. I have also proven this for the first 20 naturals, 61, and 92. So that's a bit of computational evidence. Heuristically it seems far too unlikely that the conjecture isn't true. If the conjecture is true then the numbers for which the negative Pell equation has a solution are those with an even number of numbers in their period, in particular primes congruent to 1 mod 4.

论坛问题帖 | has_answers | Sophie | 2016-12-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018692 | MSE-2040398

A nice pattern for the regularized beta function $I(\alpha^2,\frac{1}{4},\frac{1}{2})=\frac{1}{2^n}\ $?

In this post, the problem was given integer/rational N, to solve for algebraic number z in the equation, $$\begin{aligned}\frac{1}{N} &=I\left(z^2;\ a,b\right)\\[1.5mm] &= \frac{B\left(z^2;\ a,b\right)}{B\left(a,b\right)} \end{aligned} $$ using the beta function $B(a,b)$, incomplete beta $B(z;a,b)$ and regularized beta $I(z;a,b)$. It seems for some $a,b,$ there can be a pattern for the solutions. Given the equation for $n>1$, $$I\left(x_1^2;\ \tfrac14,\tfrac12\right)=\frac{1}{2^n}\tag1$$ First define $\color{blue}{\gamma = u+\sqrt{-1+u^2}},$ with fundamental unit $u=1+\sqrt{2}.\ $ Then, For $n=2$: $$x_1=x_2-\sqrt{1+x_2^2},\quad \color{blue}{x_2 =\gamma}$$ For $n=3$: $$x_1=x_2-\sqrt{1+x_2^2},\quad \color{green}{x_2 = x_3+\sqrt{-1+x_3^2},\quad x_3=x_4+\sqrt{1+x_4^2},}\quad \color{blue}{x_4 = \gamma}$$ For $n=4$: $$x_1=x_2-\sqrt{1+x_2^2},\quad x_2 = x_3+\sqrt{-1+x_3^2},\quad x_3=x_4+\sqrt{1+x_4^2},\\ \quad \color{green}{x_4 = x_5+\sqrt{-1+x_5^2},\quad x_5=x_6+\sqrt{1+x_6^2},}\quad \color{blue}{x_6 = \gamma}$$ and so on, where we add the same two nested layers (in green) each time and so end with even index x_m. Questions: Does this pattern really hold for all 2^n and $n>1?$ Why the regularity? What is the integral associated with (1) similar to the ones in the post cited above?

论坛问题帖 | has_answers | Tito Piezas III | 2016-12-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018693 | MSE-2045783

Prove that if $n \mid 1 + \sum_{k=0}^{n-1} k^{n-1}$ then n is prime.

How can I show that if $$n \mid 1 + \sum_{k=0}^{n-1} k^{n-1}$$ then n is prime? I was able to prove that n must be squarefree as follows: since $n \mid 1 + \sum_\limits{k=0}^{n-1} k^{n-1}$ then $\sum_\limits{k=0}^{n-1} k^{n-1} \equiv -1 \pmod n.$ Suppose some prime $p \mid n$, and let $q = n/p$. Now $\sum_\limits{k=0}^{n-1} k^{n-1} \equiv -1 \pmod p$, so $$\sum_{k=0}^{n-1}k^{n-1} = \sum_{m=0}^{q-1}\sum_{k=0}^{p-1}(mp+k)^{n-1} \equiv \sum_{m=0}^{q-1}\sum_{k=0}^{p-1}k^{n-1} = q\sum_{k=0}^{p-1}k^{n-1} \equiv -1 \pmod p.$$ Since -1 is a unit in $\mathbb Z/p\mathbb Z$, then so must be q, so since p is prime we have $p \nmid q$, and thus $p \mid n$ but $p^2 \nmid n$. I have not been able to show that n must be prime. By reducing exponents using Fermat's Little Theorem, I can show that for every prime $p \mid n$ (again $q = n/p$) we have $$q \sum_{k=0}^{p-1} k^{q-1} \equiv -1 \pmod p$$ but I am not sure how to deduce (if it is even possible) that for some $p \mid n$ we must have $q = 1$ (which would immediately imply $n = p$).

论坛问题帖 | has_answers | feralin | 2016-12-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018694 | MSE-2061175

How to show an infinite number of algebraic numbers α and β for $_2F_1\left(\frac13,\frac13;\frac56;-\alpha\right)=\beta\,$?

(Note: This is the case $a=\frac13$ of ${_2F_1\left(a ,a ;a +\tfrac12;-u\right)}=2^{a}\frac{\Gamma\big(a+\tfrac12\big)}{\sqrt\pi\,\Gamma(a)}\int_0^\infty\frac{dx}{(1+2u+\cosh x)^a}.\,$ There is also $a=\frac14$ and $a=\frac16$.) In a post, Reshetnikov considered some integrals and the surprising evaluations, $$ \frac{1}{48^{1/4}\,K(k_3)}\,\int_0^1 \frac{dx}{\sqrt{1-x}\,\sqrt[3]{x^2+\color{blue}{4}x^3}}=\,_2F_1\big(\tfrac{1}{3},\tfrac{1}{3};\tfrac{5}{6};-\color{blue}{4}\big)= \frac3{5^{5/6}}\tag1$$ $$ \frac{1}{48^{1/4}\,K(k_3)}\,\int_0^1 \frac{dx}{\sqrt{1-x}\,\sqrt[3]{x^2+\color{blue}{27}x^3}}=\,_2F_1\big(\tfrac{1}{3},\tfrac{1}{3};\tfrac{5}{6};-\color{blue}{27}\big)=\frac{4}{7}\tag2$$ We postulate these are just the first of an infinite family of algebraic numbers α and β such that, $$_2F_1\left(\frac13,\frac13;\frac56;-\alpha\right)=\beta\tag3$$ $$\begin{array}{|c|c|c|c|c|} \hline p&\tau&\alpha&\beta&\text{Deg}\\ \hline 3&\frac{1+3\sqrt{-3}}2& \large \frac13& \large \frac{2}{3^{2/3}}&1\\ 5&\frac{1+5\sqrt{-3}}2&\color{blue}{4}& \large\frac3{5^{5/6}} &1\\ 7&\frac{1+7\sqrt{-3}}2&\color{blue}{27}&\large\frac47&1\\ 11&\frac{1+11\sqrt{-3}}2& \sqrt{11}\big(2\sqrt3 + \sqrt{11}\big)^3& \large\frac6{11^{11/12}} \frac1{U_{33}^{1/4}} &2 \\ 13&\frac{1+13\sqrt{-3}}2& 4\sqrt{13}\big(4 + \sqrt{13}\big)^3&\large\frac7{13}\frac1{U_{13}}&2\\ 17&\frac{1+17\sqrt{-3}}2& \frac4{729}\left[(1 + \sqrt[3]{17})^2 + 6\right]^6& \large\frac9{17^{5/6}}\left(\frac{18}{17^{1/3}}-7\right)^{1/3}&3\\ 19&\frac{1+19\sqrt{-3}}2& \frac1{27}\left[(1 + \sqrt[3]{19})^2 + 5\right]^6 &\large \frac{10}{19} \Big(1-\frac{(1-19^{1/3})^2}{3}\Big)&3\\ 29&\frac{1+29\sqrt{-3}}2& 4(u_1)^6&\frac{15}{29^{5/6}}(u_2)^{1/3} &5\\ 31&\frac{1+31\sqrt{-3}}2& \frac1{27}(v_1)^6 &\frac{16}{31} (v_2)&5\\ \hline \end{array}$$ and so on, where $U_{13} = \frac{3+\sqrt{13}}2$, $U_{33} = 23+4\sqrt{33}\,$ are fundamental units, while u_i and v_i are roots of quintics, etc. (While the quintics were solvable in radicals, unfortunately they don't have the simple form as the others. The original forms for $p=17,19$ were found by yours truly while alternative ones were suggested by Reshetnikov.) And Deg is degree of $\alpha(\tau)$ and $\beta^6(\tau)$. Conjecture 1: Let $\tau = \frac{1+p\sqrt{-3}}{2}$. Using the Dedekind eta function quotient $\lambda=\frac{\eta\big(\tfrac{\tau+1}{3}\big)}{\eta(\tau)}$, then α is just a quadratic, $$16\cdot27\,\alpha(1+\alpha)=\left(\lambda^6 -27\, \lambda^{-6} \right)^2$$ or more simply, $$\alpha = \frac1{4\sqrt{27}}\big(\lambda^3-\sqrt{27}\,\lambda^{-3}\big)^2\tag4$$ And if $p=6k\pm1$ is a prime, then α and β^6 of (3) are algebraic numbers of degree k. Alternatively, one can use the well-known j-function $j(\tau)$, $$j(\tau) = {1 \over q} + 744 + 196884 q + 21493760 q^2 + 864299970 q^3+\dots$$ which is easily calculated in Mathematica as 12^3KleinInvariantJ[tau]. Conjecture 2: Let $\tau = \frac{1+p\sqrt{-3}}{2}$. Then α is an appropriate root of, $$j(\tau) = \frac{432}{1+f}\left(\frac{5+4f}{1 - f}\right)^3,\quad \text{where}\quad f = \frac{2\alpha+1}{2\sqrt{\alpha(1+\alpha)}}$$ P.S. Conjecture 2 is indebted to the answer by Noam Elkies, though the nature of τ which should provide the correct $\alpha(\tau)$ seems to have been left out.

论坛问题帖 | has_answers | Tito Piezas III | 2016-12-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018695 | MSE-2063783

Are these sequences zero for the greatest of twin, cousin, and prime triples and so on?

Let: $$x=2$$ $$(x=2 \text{ gives greater of twin primes})$$ $$(x=4 \text{ gives greater of cousin primes and so on})$$ $$\text{ If }n=k \text{ then } t(n,k)= 1$$ $$\text{ else if } k=1 \text{ then } t(n,k)= 1-\frac{1}{2} \prod _{i=1}^{n-1} t(n,i+k)-\frac{1}{2} \prod _{i=1}^{n-1-x} t(n-x,i+k)$$ $$\text{ else if } n \bmod k=0 \text{ then } t(n,k) = t\left(\frac{n}{k},1\right) \text{ else } 1 \text{ else } 1$$ This is a table starting: As a Mathematica program this is: (*start*) Clear[t, n, k, i, nn, a]; nn = 1000; x = 2; t[n_, k_] := t[n, k] = If[n == k, 1, If[k == 1, (1 - Product[t[n, k + i], {i, 1, n - 1}]/2 - Product[t[n - x, k + i], {i, 1, n - 1 - x}]/2), If[Mod[n, k] == 0, t[n/k, 1], 1], 1]]; Monitor[a = Table[t[n, 1], {n, 1, nn}];, n] Flatten[Position[a, 0]] (*end*) which gives the first column from the table in the picture above: $$a=1,0,0,\frac{1}{2},0,1,0,1,\frac{1}{2},1,\frac{1}{2},1,0,1,\frac{1}{2},1,\frac{1}{2},1,0,1,...$$ The positions of zeros in that sequence a is: 2, 3, 5, 7, 13, 19, 31, 43, 61, 73, 103, 109, 139, 151, 181, 193, 199, 229, 241, 271, 283, 313, 349, 421, 433, 463, 523, 571, 601, 619, 643, 661, 811, 823, 829, 859, 883 which matches sequence https://oeis.org/A006512 in the oeis starting: 5, 7, 13, 19, 31, 43, 61, 73, 103, 109, 139, 151, 181, 193, 199, 229, 241, 271, 283, 313, 349, 421, 433, 463, 523, 571, 601, 619, 643, 661, 811, 823, 829, 859, 883 and known as the greater of twin primes. Looking up the positions that are fractions (not necessarily 1/2, that breaks down later) I get a match with the sequence https://oeis.org/A093514 starting: 4, 9, 11, 15, 17, 21, 23, 25, 29, 33, 37, 39, 41, 45,... The title says it has something to do with the Sierpinski gasket given by the Rule 90 cellular automaton. Unfortunately I don't understand the Pari GP program given by the author. Question: Is it so that sequence a (the first column in the table from the program) is always 0 for $n>3$ if and only if the position is the greater of the twin primes? An example of a prime triplet: $$\text{ If }n=k \text{ then } t(n,k)= 1$$ $$\text{ else if } k=1 \text{ then } t(n,k)= 1-\frac{1}{3} \prod _{i=1}^{n-1} t(n,i+k)-\frac{1}{3} \prod _{i=1}^{n-1-2} t(n-2,i+k)-\frac{1}{3} \prod _{i=1}^{n-1-6} t(n-6,i+k)$$ $$\text{ else if } n \bmod k=0 \text{ then } t(n,k) = t\left(\frac{n}{k},1\right) \text{ else } 1 \text{ else } 1$$ https://oeis.org/A098413 Greatest members p of prime triples (p-6,p-2,p). {2, 3, 5, 7, 13, 19, 43, 73, 103, 109, 199, 229, 283, 313, 463, 619, \ 829, 859, 883} Compared with: {13, 19, 43, 73, 103, 109, 199, 229, 283, 313, 463, 619, 829, 859, 883,...} (*start*)Clear[t, n, k, i, nn, a]; nn = 1000; t[n_, k_] := t[n, k] = If[n == k, 1, If[k == 1, (1 - Product[t[n, k + i], {i, 1, n - 1}]/3 - Product[t[n - 2, k + i], {i, 1, n - 1 - 2}]/3 - Product[t[n - 6, k + i], {i, 1, n - 1 - 6}]/3), If[Mod[n, k] == 0, t[n/k, 1], 1], 1]]; Monitor[a = Table[t[n, 1], {n, 1, nn}];, n] Flatten[Position[a, 0]] (*end*)

论坛问题帖 | no_answers_at_retrieval | Mats Granvik | 2016-12-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018696 | MSE-2063872

Is it cheating to use the sign function when sieving for twin primes?

Let: $$x=2$$ $$t(1,1)=1$$ $$t(2,1)=0$$ $$t(3,1)=0$$ $$\text{ If }n=k \text{ then } t(n,k)= 1$$ $$\text{ if } n>3 \text{ else if } k=1 \text{ then }$$ $$t(n,k)= \text{sgn}((2- \prod _{i=1}^{n-1} t(n,i+k)- \prod _{i=1}^{n-1-x} t(n-x,i+k)) \cdot (2- \prod _{i=1}^{n-1} t(n,i+k)- \prod _{i=1}^{n-1+x} t(n+x,i+k)))$$ $$\text{ else if } n \bmod k=0 \text{ then } t(n,k) = t\left(\frac{n}{k},1\right) \text{ else } 1 \text{ else } 1$$ where $$\text{sgn}(x)$$ is the sign function. This produces a characteristic sequence of non-twin primes for $n>2$ starting: 1, 0, 0, 1, 0, 1, 0, 1, 1, 1, 0, 1, 0, 1, 1, 1, 0, 1, 0, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 0, 1, 0, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 0, 1, 0, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1

\frac{j(\tau)-1728}{j(\tau)}$. Examples: $$_2F_1\left(\tfrac14,\tfrac14;\tfrac12;\,\tfrac{(3-6\sqrt2)^2}{(1+\sqrt2)^4}\right)=\tfrac{3}{4\sqrt2}(1+\sqrt2)$$ $$_2F_1\left(\tfrac16,\tfrac13;\tfrac12;\,\tfrac{25}{27}\right)=\tfrac{3\sqrt3}{4}$$ $$_2F_1\left(\tfrac18,\tfrac38;\tfrac12;\tfrac{2400}{2401}\right)=\tfrac{2\sqrt7}{3}$$ $$_2F_1\left(\tfrac1{12},\tfrac5{12};\tfrac12;\tfrac{1323}{1331}\right)=\tfrac{3\,\sqrt[4]{11}}{4}$$ using $\tau_1=2\sqrt{-4},\;\tau_2=2\sqrt{-3},\;\tau_3=3\sqrt{-2},\;\tau_4=2\sqrt{-1}$, respectively. And so on for any integer $N>1$. II. Type $a+b=c=\color{blue}{\tfrac23}$. $$\begin{aligned} &\,_2F_1\big(\tfrac14,\tfrac5{12};\tfrac23;(1-2\beta_1)^2\big),\quad \color{red}{\beta_1 =\,?} \\[2mm] &\,_2F_1\big(\tfrac16,\tfrac12;\tfrac23;(1-2\beta_2)^2\big),\quad \frac{1}{\beta_2}-1=\sqrt{\frac{-1}{1728}\big(2k+\sqrt{4k^2-1728^2}\big)}\\[2mm] &\,_2F_1\big(\tfrac18,\tfrac{13}{24};\tfrac23;(1-2\beta_3)^2\big),\quad \color{red}{\beta_3 =\,?} \\[2mm] &\,_2F_1\big(\tfrac1{12},\tfrac7{12};\tfrac23;(1-2\beta_4)^2\big),\quad \frac{1}{\beta_4}-1=\frac{-1}{1728}\big(2k+\sqrt{4k^2-1728^2}\big)\end{aligned}$$ and where $k=j(\tau)-864$. Examples: Use $\tau = \frac{1+N\sqrt{-3}}2$, like $\tau = \frac{1+3\sqrt{-3}}2$, $$_2F_1\big(\tfrac16,\tfrac12;\tfrac23;\tfrac{125}{128}\big) =\tfrac43\,2^{1/6}$$ $$_2F_1\big(\tfrac1{12},\tfrac7{12};\tfrac23;\tfrac{64000}{64009}\big) =\tfrac23\,253^{1/6}$$ III. Type $a+b=c=\color{blue}{\tfrac34}$. $$\begin{aligned} &\,_2F_1\big(\tfrac14,\tfrac12;\tfrac34;(1-2\gamma_1)^2\big), \quad\frac1{\gamma_1}-1=\sqrt{\frac1{64}\Big(\tfrac{\sqrt2\,\eta(2\tau)}{\zeta_{48}\,\eta(\tau)}\Big)^{24}}\\[2mm] &\,_2F_1\big(\tfrac16,\tfrac7{12};\tfrac34;(1-2\gamma_2)^2\big),\quad\color{red}{\gamma_2 =\,?} \\[2mm] &\,_2F_1\big(\tfrac18,\tfrac58;\tfrac34;(1-2\gamma_3)^2\big),\quad\frac1{\gamma_3}-1=\frac1{64}\Big(\tfrac{\sqrt2\,\eta(2\tau)}{\zeta_{48}\,\eta(\tau)}\Big)^{24}\\[2mm] &\,_2F_1\big(\tfrac1{12},\tfrac23;\tfrac34;(1-2\gamma_4)^2\big),\quad\color{red}{\gamma_4 =\,?} \end{aligned}$$ where $\zeta_{48} = e^{2\pi i/48}$. Examples: Use $\tau = \frac{1+N\sqrt{-1}}2$, like $\tau = \frac{1+5\sqrt{-1}}2$, $$_2F_1\big(\tfrac14,\tfrac12;\tfrac34;\tfrac{80}{81}\big)=\tfrac95$$ $$_2F_1\big(\tfrac18,\tfrac58;\tfrac34;\tfrac{25920}{25921}\big)=\tfrac35\,161^{1/4}$$ IV. Type $a+b=c=\color{blue}{\tfrac56}$. $$\begin{aligned} &\,_2F_1\big(\tfrac12,\tfrac13;\tfrac56;(1-2\delta_1)^2\big),\quad\;\delta_1 = \delta_2\\[2mm] &\,_2F_1\big(\tfrac13,\tfrac12;\tfrac56;(1-2\delta_2)^2\big),\quad\;\frac1{\delta_2}-1=\sqrt{\frac1{27}\left(\tfrac{\eta\big(\frac{\tau+1}{3}\big)}{\eta(\tau)}\right)^{12}}\\[2mm] &\,_2F_1\big(\tfrac14,\tfrac7{12};\tfrac56;(1-2\delta_3)^2\big),\quad\color{red}{\delta_3 =\,?} \\[2mm] &\,_2F_1\big(\tfrac16,\tfrac23;\tfrac56;(1-2\delta_4)^2\big),\quad\;\frac1{\delta_4}-1=\frac1{27}\left(\tfrac{\eta\big(\frac{\tau+1}{3}\big)}{\eta(\tau)}\right)^{12} \end{aligned}$$ Note: Of course, $_2F_1\big(\tfrac12,\tfrac13;\tfrac56;z\big) =\,_2F_1\big(\tfrac13,\tfrac12;\tfrac56;z\big)\,$ so the first form is superfluous. Examples: Use $\tau = \frac{1+N\sqrt{-3}}2$, like $\tau = \frac{1+5\sqrt{-3}}2$, $$\begin{aligned} &\,_2F_1\big(\tfrac13,\tfrac12;\tfrac56;\tfrac45\big)=\;\tfrac35\sqrt5 \\[2mm] &\,_2F_1\big(\tfrac16,\tfrac23;\tfrac56;\tfrac{80}{81}\big)=\tfrac35\,(9\sqrt5)^{1/3}\end{aligned}$$ with the last one discussed in this post. Q: How do we find the five missing formulas (?) for the variables in red?

论坛问题帖 | no_answers_at_retrieval | Tito Piezas III | 2016-12-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018699 | MSE-2069793

How to show an infinite number of algebraic numbers α and β for $_2F_1\left(\frac14,\frac14;\frac34;-\alpha\right)=\beta\,$?

(Note: This is the case $a=\frac14$ of ${_2F_1\left(a ,a ;a +\tfrac12;-u\right)}=2^{a}\frac{\Gamma\big(a+\tfrac12\big)}{\sqrt\pi\,\Gamma(a)}\int_0^\infty\frac{dx}{(1+2u+\cosh x)^a}.\,$ There is also $a=\frac13$ and $a=\frac16$.) In a post, Reshetnikov considered some integrals and their equivalent forms, $$\frac{1}{2\sqrt2\,K(k_1)}\,\int_0^1 \frac{dx}{\sqrt{1-x}\,\sqrt[4]{x^3+\color{blue}{3}x^4}}=\,_2F_1\big(\tfrac{1}{4},\tfrac{1}{4};\tfrac{3}{4};-\color{blue}3\big) = \frac{2}{3^{3/4}}\tag1$$ $$\frac{1}{2\sqrt2\,K(k_1)}\,\int_0^1 \frac{dx}{\sqrt{1-x}\,\sqrt[4]{x^3+\color{blue}{80}x^4}}=\,_2F_1\big(\tfrac{1}{4},\tfrac{1}{4};\tfrac{3}{4};-\color{blue}{80}\big) = \frac35\tag2$$ Just like for the case $a=\tfrac13$, we postulate these are just the first of an infinite family of algebraic numbers α and β such that, $$_2F_1\left(\frac14,\frac14;\frac34;-\alpha\right)=\beta\tag3$$ The table below summarizes results and show some quartic trends, $$\begin{array}{|c|c|c|c|c|} \hline p&\tau&\alpha(\tau)&\beta(\tau)&\text{Deg}\\ \hline 3&\frac{1+3\sqrt{-1}}2&4\cdot1^4-1=\color{blue}{3}& \large\frac2{3^{3/4}} &1\\ 5&\frac{1+5\sqrt{-1}}2&3^4-1=\color{blue}{80}& \large\frac35 &1\\ 7&\frac{1+7\sqrt{-1}}2&4(2+\sqrt7)^4-1& \large\frac4{7^{7/8}} \frac1{(8+3\sqrt7)^{1/4}} &2\\ 11&\frac{1+11\sqrt{-1}}2&4u^4 - 1 &\large \frac6{11^{11/12}}\Big(\frac{11^{2/3}-(14-6\sqrt3)^{1/3}-(14+6\sqrt3)^{1/3}}{3}\Big) &3 \\ 13&\frac{1+13\sqrt{-1}}2& v^4-1 &\large \frac7{13^{2/3}}\Big(\frac{13^{-1/3}+(-2+6\sqrt3)^{1/3}-(2+6\sqrt3)^{1/3}}{3}\Big) &3 \\ 17&\frac{1+17\sqrt{-1}}2& x_1^4 - 1 & \large \frac9{17}x_2 &4 \\ 19&\frac{1+19\sqrt{-1}}2& 4y_1^4 - 1 & \large \frac{10}{19}y_2^{1/4} &5 \\ \hline \end{array}$$ and fundamental unit $U_7=8+3\sqrt7$, with u,v as the real root of the cubics, $$u^3 - 23u^2 + 15u - 9=0\\v^3 - 67v^2 - 159v - 99=0$$ and x_i as roots of quartics, and y_i as roots of quintics, and so on. Deg is degree of $\alpha(\tau)$ and $\beta^4(\tau)$. Conjecture: Let $\tau = \frac{1+p\sqrt{-1}}{2}$. To find α, define the eta quotient $\displaystyle\lambda=\frac{\sqrt2\,\eta(2\tau)}{\zeta_{48}\,\eta(\tau)}$ with $\zeta_{48} = e^{2\pi i/48}$. Then, like the case $a=\tfrac13$, α also is a quadratic of similar form, $$16\cdot64\,\alpha(1+\alpha)=\left(\lambda^{12} -64\, \lambda^{-12} \right)^2$$ with analogous root, $$\alpha= \frac1{4\sqrt{64}}\big(\lambda^6-\sqrt{64}\,\lambda^{-6}\big)^2\tag4$$ And if $p=4k\pm1$ is a prime, then α and β^4 of (3) are algebraic numbers of degree k. Questions: How do we prove the conjecture? (And the analogous one in the other post?) Is there a third family of this group with infinitely many algebraic numbers α_3 and β_3? $\color{green}{Update}:$ It seems there is a third family. The obvious choices are $a=\frac12$ and $a=\frac16$. For the latter, and using a better search method, I found, $$_2F_1\Big(\frac{1}{6},\frac{1}{6};\frac{2}{3};-\frac{125}3\Big) = \frac{2}{3^{5/6}}$$ See case $a=\frac16$ here.

论坛问题帖 | has_answers | Tito Piezas III | 2016-12-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018700 | MSE-2072468

How to prove $_2F_1\big(\tfrac16,\tfrac16;\tfrac23;-2^7\phi^9\big)=\large \frac{3}{5^{5/6}}\,\phi^{-1}\,$ with golden ratio ϕ?

(Note: This is the case $a=\frac16$ of ${_2F_1\left(a ,a ;a +\tfrac12;-u\right)}=2^{a}\frac{\Gamma\big(a+\tfrac12\big)}{\sqrt\pi,\Gamma(a)}\int_0^\infty\frac{dx}{(1+2u+\cosh x)^a}.\,$ There is also $a=\frac13$ and $a=\frac14$.) After investigating $a=\frac13$ and $a=\frac14$, I wondered if there was for $a=\frac16$. And happily there was, $$\frac{1}{\color{red}{432}^{1/4}\,K(k_3)}\,\int_0^1 \frac{dx}{\sqrt{1-x}\,\sqrt[6]{x^5+\color{blue}{\tfrac{125}3}x^6}}=\,_2F_1\big(\tfrac16,\tfrac16;\tfrac23;-\color{blue}{\tfrac{125}{3}}\big)=\frac{2}{3^{5/6}}$$ $$\frac{1}{\color{red}{432}^{1/4}\,K(k_3)}\,\int_0^1 \frac{dx}{\sqrt{1-x}\,\sqrt[6]{x^5+\color{blue}{2^7\phi^9}\, x^6}}=\,_2F_1\big(\tfrac16,\tfrac16;\tfrac23;-\color{blue}{2^7\phi^9}\big)=\frac{3}{5^{5/6}}\phi^{-1}$$ The first was found by computer search and, from previous posts, the denominator with $K(k_3)$ was enough to give me a clue that $\tau=\frac{1+3\sqrt{-3}}2$ was involved. After fiddling around with some equations, a third conjecture can be made, that there is an infinite family of algebraic numbers α and β such that, $$_2F_1\left(\frac16,\frac16;\frac23;-\alpha\right)=\beta$$ Conjecture: "Let $\tau = \frac{1+p\sqrt{-3}}{2}$ with integer $p>1$. Then α is the root of an analogous quadratic, $$16\cdot\color{red}{432}\,\alpha(1+\alpha)=-j(\tau)$$ with j-function $j(\tau)$. And if odd $p=3k\pm1$ is a prime, then α and β^6 are algebraic numbers of degree k." $$\begin{array}{|c|c|c|c|c|} \hline p&\tau&\alpha(\tau)&\beta(\tau)&\text{Deg}\\ \hline 3&\frac{1+3\sqrt{-3}}2&\frac{125}3& \large\frac2{3^{5/6}} &1\\ 5&\frac{1+5\sqrt{-3}}2&2^7\phi^9& \large\frac3{5^{5/6}}\phi^{-1} &2\\ 7&\frac{1+7\sqrt{-3}}2&\Big(\frac{129 + 29\sqrt{21}}2\Big)^3& \large\frac47 \frac1{U_{21}^{1/2}} &2\\ 11&\frac{1+11\sqrt{-3}}2& x_1 & \large\frac6{11} x_2 &4 \\ 13&\frac{1+13\sqrt{-3}}2& y_1 & \large\frac7{13} y_2 &4 \\ \hline \end{array}$$ $U_{21}=\frac{5+\sqrt{21}}2$ is a fundamental unit, while x_i,y_i are roots of quartics which are rather tedious to write down. And so on. Q: How do we prove this conjecture? (And the other two?)

论坛问题帖 | has_answers | Tito Piezas III | 2016-12-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018701 | MSE-2081467

On $\int_0^1\arctan\,_6F_5\left(\frac17,\frac27,\frac37,\frac47,\frac57,\frac67;\,\frac26,\frac36,\frac46,\frac56,\frac76;\frac{n}{6^6}\,x\right)\,dx$

Reshetnikov gave the remarkable evaluation, \begin{align} I&= \int_0^1\arctan{_4F_3}\left(\frac15,\frac25,\frac35,\frac45;\frac24,\frac34,\frac54;\frac{1}{64}\,x\right)\,dx \\ &=\frac{3125}{48}\left(5+3\pi+6\ln2-3\alpha^4+4\alpha^3+6\alpha^2-12\alpha\\-12\left(\alpha^5-\alpha^4+1\right)\arctan\frac1\alpha-6\ln\left(1+\alpha^2\right)\right)\\ &=0.7857194\dots \end{align} where α is a quartic root. However, it seems this can be simplified a bit and generalized. I. $p=5$: Given, $$I(n)=\int_0^1\arctan\,_4F_3\left(\frac15,\frac25,\frac35,\frac45;\,\frac24,\frac34,\frac54;\,\frac{n}{4^4}\,x\right)\,\mathrm dx$$ Is it true that, in general, $$\frac{12n}{5^5}\, I(n) = -12\Big(1\color{blue}-\frac{n}{5^5}\Big)\arctan\frac1\alpha+6\ln\Big(\frac{2}{1+\alpha^2}\Big)+3\pi+P(\alpha)$$ where $P(\alpha)=(1-\alpha)^3(5+3\alpha)$ and α is the largest positive root of the quintic, $$\alpha^5-\alpha^4+\frac{n}{5^5}=0$$ provided real number $n<4^4\,$? (Note: By sheer coincidence, the choice of $n=4$ in the other post caused the quintic to factor.) II. $p=7$: Given, $$J(n)=\int_0^1\arctan\,_6F_5\left(\frac17,\frac27,\frac37,\frac47,\frac57,\frac67;\,\frac26,\frac36,\frac46,\frac56,\frac76;\,\frac{n}{6^6}\,x\right)\,\mathrm dx$$ is it true that, $$\frac{60n}{7^7}\, J(n) = 60\Big(1\color{blue}+\frac{n}{7^7}\Big)\arctan\frac1\alpha-30\ln\Big(\frac{2}{1+\alpha^2}\Big)-15\pi-P(\beta)$$ where $P(\beta) = (1-\beta)^2(27-6\beta-9\beta^2+8\beta^3+10\beta^4)$ and β is the largest positive root of, $$\beta^7-\beta^6+\frac{n}{7^7}=0$$ provided real $n<6^6\,$? Questions: How do we prove the two conjectured equalities? What's the formula for $p=3$? (Mathematica takes too long to evaluate the integral that I couldn't use an integer relations subroutine on it.)

论坛问题帖 | has_answers | Tito Piezas III | 2017-01-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018702 | MSE-2084716

Connecting $\sum\limits_{n=0}^\infty\frac{n!\,(2n)!}{(3n+2)!}$ and $\sum\limits_{n=1}^\infty\frac{1}{n \binom{3n}{n}}$

Define three generalized hypergeometric functions (which differs in the starting numerator in blue): $$A_k =\,_kF_{k-1}\left(\left.\begin{array}{c} 1,\frac{\color{blue}{k-2}}{k-1},\frac{k-1}{k-1},\dots\\ \frac{k+1}{k}, \frac{k+2}{k},\dots \end{array}\right| \frac{(k-1)^{k-1}}{k^k}\right)$$ $$B_k =\,_kF_{k-1}\left(\left.\begin{array}{c} 1,\frac{\color{blue}{k-1}}{k-1},\frac{k}{k-1},\dots\\ \frac{k+1}{k}, \frac{k+2}{k},\dots \end{array}\right| \frac{(k-1)^{k-1}}{k^k}\right)$$ $$C_k =\,_kF_{k-1}\left(\left.\begin{array}{c} 1,\frac{\color{blue}{k}}{k-1},\frac{k+1}{k-1},\dots\\ \frac{k+1}{k}, \frac{k+2}{k},\dots \end{array}\right| \frac{(k-1)^{k-1}}{k^k}\right)$$ The smallest common non-trivial case would then be $k=3$, $$A_3={_3F_2}\left(1,\tfrac12,\tfrac22;\ \tfrac43,\tfrac53;\ \tfrac4{27}\right)$$ $$B_3={_3F_2}\left(1,\tfrac22,\tfrac32;\ \tfrac43,\tfrac53;\ \tfrac4{27}\right)$$ $$C_3={_3F_2}\left(1,\tfrac32,\tfrac42;\ \tfrac43,\tfrac53;\ \tfrac4{27}\right)$$ The first two were addressed in this and this posts by different OPs from different perspectives, but a common approach may be possible. First, let x_n be the roots of $x^k-x+1=0$. Question 1: Is it true, $$\frac{A_k}{(k-1)(k-2)} = \int_1^\infty \frac{-k+(k-1)x}{x^k-x+1}dx =-\sum_{n=1}^k x_n\ln(1-x_n)\tag1$$ Note: This implies a simple solution to the case $k=3$ discussed in by Reshetnikov in this post as, $$\sum\limits_{n=0}^\infty\frac{n!\,(2n)!}{(3n+2)!}= -x_1\ln(1-x_1)-x_2\ln(1-x_2)-x_3\ln(1-x_3) = 0.5179778\dots$$ where the x_n are the three roots of $x^3-x+1=0\;^\color{red}*$ and which is the minpoly of the negated plastic constant. Question 2: Is it true, $$\frac{B_k}{k} = \int_1^\infty\frac1{x(x^k-x+1)}dx=-\sum_{n=1}^k \frac{\ln(1-x_n)}{-k+(k-1)x_n}\tag2$$ Note: This implies a similarly simple solution to the case $k=3$ in this post, $$\sum_{n=1}^\infty\frac1{n\binom{3n}n} = \frac{\ln(1-x_1)}{3-2x_1}+

\frac{\ln(1-x_2)}{3-2x_2}+ \frac{\ln(1-x_3)}{3-2x_3}=0.371216\dots$$ and the x_n again are the three roots of $^\color{red}*$. Question 3: See this post.

论坛问题帖 | no_answers_at_retrieval | Tito Piezas III | 2017-01-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018703 | MSE-2090850

When p is an odd prime, is $(p+2)/p$ an outlaw or an index?

Let $\sigma(x)$ denote the sum of the divisors of x, and denote the abundancy index of x as $$I(x) = \dfrac{\sigma(x)}{x},$$ and the deficiency of x as $$D(x) = 2x - \sigma(x).$$ If the equation $I(a)=b/c$ has no solution $a \in \mathbb{N}$, then b/c is said to be an abundancy outlaw. Statement of the Problem When p is an odd prime, is $(p+2)/p$ an outlaw or an index? Preliminary Results The following lemmas are easy to show: Lemma 1. If p is an odd prime, and $(p+2)/p$ is the abundancy index of some integer n, then n is deficient. Lemma 2. If p is odd, then $\gcd(p,p+2)=1$. Lemma 3. If p is an odd prime and $I(n) = (p+2)/p$, then $D(n) = 2n - \sigma(n) \neq 1$. Lemma 4. If p is an odd prime and $I(n) = (p+2)/p$, then $p < n$. Lemma 5. If p is an odd prime and $I(n) = (p+2)/p$, then $\gcd(n,\sigma(n)) \neq 1$. Remarks In fact, one can show that, if p is an odd prime and $I(n) = (p+2)/p$, then $$\sigma(n) = \bigg(\dfrac{n}{p}\bigg)\cdot(p+2)$$ and $$n = \bigg(\dfrac{\sigma(n)}{p+2}\bigg)\cdot(p).$$ (Note that n/p and $\sigma(n)/(p+2)$ are (equal) integers because of Lemma 2.) Consequently, we obtain $$\gcd(n,\sigma(n)) = \dfrac{n}{p} = \dfrac{\sigma(n)}{p+2}.$$ (Note further that both $\gcd(n,\sigma(n)) \leq n/3$ and $\gcd(n,\sigma(n)) \leq \sigma(n)/5$ hold.) Added September 16 2017 Given that $X = A/B = C/D$ ($B \neq 0$, $D \neq 0$, and $B \neq D$), we can make use of the algebraic identity $$\frac{A}{B}=\frac{C}{D}=\frac{C-A}{D-B}$$ to get another expression for $$\gcd(n,\sigma(n)) = \dfrac{n}{p} = \dfrac{\sigma(n)}{p+2}.$$ Indeed, $$\gcd(n,\sigma(n)) = \dfrac{n}{p} = \dfrac{\sigma(n)}{p+2} = \frac{\sigma(n) - n}{2}.$$ This last finding implies that $$\bigg(\frac{\sigma(n) - n}{2}\bigg) \mid n \iff (\sigma(n) - n) \mid (2n) \iff 2n = (\sigma(n) - n){d_1}$$ and $$\bigg(\frac{\sigma(n) - n}{2}\bigg) \mid \sigma(n) \iff (\sigma(n) - n) \mid (2\sigma(n)) \iff 2\sigma(n) = (\sigma(n) - n){d_2}.$$ Note that $2 \mid (\sigma(n) - n)$. Additionally, notice that $$2\gcd\left(n,\sigma(n)\right) = \gcd\left(2n, 2\sigma(n)\right) = \gcd\left((\sigma(n) - n){d_1},(\sigma(n) - n){d_2}\right)$$ $$= \left(\sigma(n) - n\right)\gcd({d_1},{d_2}) \iff \frac{2\gcd\left(n,\sigma(n)\right)}{\left(\sigma(n) - n\right)}=1=\gcd({d_1},{d_2}).$$ In fact, $${d_1} = \frac{2n}{\sigma(n) - n} = p$$ and $${d_2} = \frac{2\sigma(n)}{\sigma(n) - n} = p+2.$$ Double-checking if it is indeed the case that ${d_1}+2={d_2}$: $${d_1} = \frac{2n}{\sigma(n) - n} + 2 = \frac{2n + 2(\sigma(n) - n)}{\sigma(n) - n} = \frac{2\sigma(n)}{\sigma(n) - n} = d_2.$$ So far so good! More is actually true. One can also show that $$p(2n - \sigma(n)) = (p - 2)n$$ so that $$D(n) = (p - 2)\cdot\bigg(\dfrac{n}{p}\bigg) = (p - 2)\cdot\bigg(\dfrac{\sigma(n)}{p + 2}\bigg) = (p - 2)\cdot\gcd(n,\sigma(n)).$$ We therefore conclude that $$\dfrac{D(n)}{n} = \dfrac{p - 2}{p} = \bigg(\dfrac{p - 2}{p + 2}\bigg)\cdot{I(n)}.$$ Added October 8 2017 We deduce that $$\dfrac{p-2}{p}=\dfrac{D(n)}{n}<\dfrac{\phi(n)}{n}<\dfrac{n}{\sigma(n)}=\dfrac{p}{p+2},$$ whence there is still no contradiction. Motivation It is conjectured that $(p+2)/p$ is an outlaw, since if it were an index, then we would be able to produce an odd perfect number for $p=3$. Here is my question: To what extent can the following theorem be improved to hopefully produce some results towards proving the aforementioned conjecture? Theorem If n is a positive integer satisfying $D(n) = 2n - \sigma(n) > 1$, then we have the following bounds for the abundancy of n in terms of the deficiency of n: $$\dfrac{2n}{n + D(n)} < I(n) < \dfrac{2n + D(n)}{n + D(n)}.$$

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2017-01-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018704 | MSE-2102804

If $q^k n^2$ is an odd perfect number with Euler prime q, does $I(n^2) \geq 5/3$ imply $k=1$?

Let $\sigma(x)$ denote the sum of the divisors of $x \in \mathbb{N}$. Denote the abundancy index of $y \in \mathbb{N}$ by $I(y)=\sigma(y)/y$. If $\sigma(N)=2N$, then N is said to be perfect. Euler proved that every odd perfect number has the form $q^k n^2$ where q is prime (called the Euler prime) with $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. If $k=1$, then since $q \geq 5$, then we have $$I(q)I(n^2)=2 \Longleftrightarrow I(n^2)=\dfrac{2q}{q+1} \geq \dfrac{5}{3}.$$ If $k>1$, then we can use the bound $$I(q^k) < \dfrac{q}{q-1} \leq \dfrac{5}{4}$$ so that $$I(n^2)=\dfrac{2}{I(q^k)} > \dfrac{2(q-1)}{q} \geq \dfrac{8}{5}.$$ Here is my question: If $q^k n^2$ is an odd perfect number with Euler prime q, does $I(n^2) \geq 5/3$ imply $k=1$?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2017-01-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018705 | MSE-2106690

Modular relationship between a number and a copy with removed figures

I want a proof of this conjecture: Given a number n with more than two figures written in a positional base-b numeral system and let n' be a base-b number obtained by removing a consecutive pair of equal b-figures in n. Then $$\,n\!\!\mod (b+1)=\,n'\!\!\mod (b+1)$$ Example: $123345667 \mod 11 = 1245667 \mod 11$ for $b\ge 8$.

论坛问题帖 | has_answers | Lehs | 2017-01-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018706 | MSE-2112707

A limit related to asymptotic growth of tetration

The tetration is denoted $^n a$, where a is called the base and n is called the height, and is defined for $n\in\mathbb N\cup\{-1,\,0\}$ by the recurrence $$ {^{-1} a} = 0, \quad {^{n+1} a} = a^{\left({^n a}\right)},\tag1$$ so that $${^0 a}=1, \quad {^1 a} = a, \quad {^2 a} = a^a, \quad {^3 a} = a^{a^a}, \, \dots \quad {^n a} = \underbrace{a^{a^{{.^{.^{.^a}}}}}}_{n\,\text{levels}}.\tag2$$ Let a be a real number in the interval $e^{-e} < a < e^{1/e}$. It is known that the following limit exists $$L(a) = \lim_{n\to\infty} {^n a},\tag3$$ where $L(a)$ satisfies $a^{L(a)}=L(a)$. For example, $L\left(\!\sqrt2\right)=2$. It is also known that $$\lim_{n\to\infty} \, \frac{L(a) -

{^{n+1} a}}{L(a) - {^n a}} = \ln L(a).\tag4$$ Finally, it is known that the following limit exists $$C(a) = \lim_{n\to\infty} \, \frac{L(a) - {^n a}}{\left(\ln L(a)\right)^n}.\tag5$$ Apparently, no closed form for the function $C(a)$ is known. But the numerical evidence suggests the following conjecture (basically, this is the coefficient of the linear term in the Taylor series expansion of $C(a)$ near $a=1$): $$C'(1) = \lim_{a\to1} \, \frac{C(a)}{a-1} \stackrel?= 1.\tag{$\diamond$}$$ How can we prove it? Can we find values of some higher-order derivatives? Are they all integers? Is there a general formula, recurrence or an efficient algorithm to compute them? Related questions: [1][2][3].

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2017-01-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018707 | MSE-2113387

Primitive pythagorean triple conjecture with pi (π)

An Ordered Primitive Pythagorean Triple (a,b,c) is one in which $a \le b \le c$ are coprime and $a^2+b^2 = c^2$. $f(n) = |\{(a,b,c)~|~ a^2+b^2=c^2,a\le b\le c,~c \le n\}|$. Function $f(n)$ defines the number of all distinct Ordered Primitive Pythagorean Triple (a,b,c) with $c \le n$. For example, $f(4) = 0$ $f(5) = 1$ with triple $(3,4,5)$ Conjecture: for any $\epsilon > 0$, there exists N_0 such that $\forall n \ge N_0, \frac{n}{f(n)} \in (2 \pi - \epsilon, 2 \pi + \epsilon)$. Question:I'd like to ask is there anyone has proposed such conjecture or is there any proof (true or false) for this conjecture? Here are the first several triples n f(n) n/f(n) 5 1 5 13 2 6.5 17 3 5.66666666666667 25 4 6.25 [3, 4, 5] [5, 12, 13] [8, 15, 17] [7, 24, 25]

论坛问题帖 | has_answers | Kevin | 2017-01-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018708 | MSE-2115058

Prime gap around p_n

As ${\rm li}(x)\sim \pi(x) $ and Cramer's conjecture predicts that the maximal prime gap around $ p_n $ is $ O(\log^{2}p_n) $, does a strong heuristics suggest that this prime gap is approximately $ \int_{1}^{p_n}\left(\frac{dx}{dy}\right)\frac{dy}{y} $ where $ y=\pi(x) $? Indeed the derivative of $ \log^{2}x $ is $ 2\frac{\log x}{x} $ which is approximately $ \frac{2}{\pi(x)} $.

论坛问题帖 | has_answers | Sylvain Julien | 2017-01-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018709 | MSE-2117122

About square free semiprimes which are sums of two squares

This conjecture seems to be true: If the product of two different primes is a sum of two squares, the so are the primes. I've tested it for $pq<1000$, but would like to see a proof. Edit: If $p\neq q$ are primes and $pq=a^2+b^2$, then there are integers c,d such that $p=c^2+d^2$.

论坛问题帖 | has_answers | Lehs | 2017-01-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018710 | MSE-2129245

Two times the sum of two squares is a sum of two squares

For all $a,b\in\mathbb Z$ there are $c,d\in\mathbb Z$ such that $2(a^2+b^2)=c^2+d^2$. The conjecture is tested for $a^2+b^2<1,000,000$ but I have problems with proving it.

论坛问题帖 | has_answers | Lehs | 2017-02-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018711 | MSE-2130737

Can this heuristic on odd perfect numbers be made rigorous?

Can this heuristic on odd perfect numbers be made rigorous using the "Axiom of Choice"? Preliminaries Euler showed that an odd perfect number, if one exists, must take the form $N=q^k n^2$ where q is prime with $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. (It follows that $q \geq 5$.) Let $\sigma(x)$ be the sum of the divisors of $x \in \mathbb{N}$. Then by definition, $$\sigma(q^k)\sigma(n^2) = \sigma(q^k n^2) = \sigma(N) = 2N = 2 q^k n^2$$ from which it follows that $$I(q^k)I(n^2) = \dfrac{\sigma(q^k)}{q^k}\cdot\dfrac{\sigma(n^2)}{n^2} = 2$$ where $I(x)=\sigma(x)/x$ is the abundancy index of x. Numerics By a well-known formula for the sum of divisors of a prime power, we have $$I(q^k) = \dfrac{\sigma(q^k)}{q^k} = \dfrac{q^{k+1} - 1}{q^k(q - 1)}.$$ This is bounded by $$\dfrac{q+1}{q} = I(q) \leq I(q^k) < \dfrac{q^{k+1}}{q^k(q - 1)} = \dfrac{q}{q - 1}.$$ Now, note that, when $k=1$, we have the bounds $$1 < I(q^k) = \dfrac{q+1}{q} = 1 + \dfrac{1}{q} \leq 1 + \dfrac{1}{5} = \dfrac{6}{5},$$ from which it follows that $$\dfrac{5}{3} \leq I(n^2) = \dfrac{2}{I(q^k)} = \dfrac{2}{I(q)} < 2.$$ Additionally, when $k>1$, we have the bounds $$1 < I(q^k) < \dfrac{q}{q - 1}.$$ Since $$\dfrac{q}{q - 1} = \dfrac{1}{1 - (1/q)}$$ we obtain $$q \geq 5 \iff 1/q \leq 1/5 \iff 1 - (1/q) \geq 4/5 \iff \dfrac{1}{1 - (1/q)} \leq 5/4$$ so that $$1 < I(q^k) < \dfrac{q}{q - 1} \leq \dfrac{5}{4}.$$ This then implies that $$\dfrac{8}{5} < I(n^2) = \dfrac{2}{I(q^k)} < 2.$$ Implications Note that we have proved the following implications: (1) If $N=q^k n^2$ is an odd perfect number with $k=1$, then $I(n^2) \geq 5/3$. (2) If $N=q^k n^2$ is an odd perfect number with $k>1$, then $I(n^2) > 8/5$. Contrapositives (3) Contrapositive of (1) If $N=q^k n^2$ is an odd perfect number with $I(n^2) < 5/3$, then $k>1$. (4) Contrapositive of (2) If $N=q^k n^2$ is an odd perfect number with $I(n^2) \leq 8/5$, then $k=1$. Analysis By (3) and (2), we have $$I(n^2) < 5/3 \implies k>1 \implies I(n^2) > 8/5 \implies 8/5 < I(n^2) < 5/3$$ resulting in no contradiction, since ${8}\cdot{3} = 24 < 25 = {5}\cdot{5}$. By (4) and (1), we get $$I(n^2) \leq 8/5 \implies k=1 \implies I(n^2) \geq 5/3 \implies 5/3 \leq I(n^2) \leq 8/5$$ resulting in a contradiction, since ${5}\cdot{5}= 25 \nleq 24 = {8}\cdot{3}$. This implies that $I(n^2) > 8/5$ is unconditionally true. Now here is the heuristic: Heuristic If we use the "Axiom of Choice" and identify the lower bound $I(n^2) \geq 5/3$ with $k=1$ and similarly identify the lower bound $I(n^2) > 8/5$ with $k>1$, and since assuming the negation of the latter lower bound contradicts (4) and (1) above, does this mean that we can predict $k>1$ to be true? That is, can we "assume" or prove that the following biconditionals hold? $$k=1 \iff I(n^2) \geq 5/3$$ $$k>1 \iff I(n^2) > 8/5$$

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2017-02-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018712 | MSE-2135432

Help finding the flaw in this proof

Today I spent my afternoon trying to understand why the Hardy-Littlewood's Second Conjecture is considered as a problem of such a great difficulty. I got a fairly strange result, so I decided to post it here and wait for opinion. So sorry if I made any kind of terrible mistake or so. The conjecture itself states that: $$\pi(x+y) - \pi(x) \le \pi(y).$$ Or, in the same way, $$\frac{\pi(x+y)}{y(y^2-1)} - \frac{\pi(x)}{y(y^2-1)} \le \frac{\pi(y)}{y(y^2-1)}.$$ Then, integrating both parts from 2 to ∞ over the variable y, $$\int_2^\infty \frac{\pi(x+y)}{y(y^2-1)}dy - \pi(x) \frac{\log\big(\frac{4}{3}\big)}{2} < \frac{\log(\zeta(2))}{2}.$$ Integrating by parts in the first integral, $$\pi(x+2)\frac{\log\big(\frac{4}{3}\big)}{2} - \int_2^\infty \frac{\log\big(1-\frac{1}{y^2}\big)}{2}\pi '(x+y) dy - \pi(x) \frac{\log\big(\frac{4}{3}\big)}{2} < \frac{\log(\zeta(2))}{2}.$$ Then $$\log\bigg(\frac{4}{3}\bigg) \left(\pi(x+2) - \pi(x) \right) - \int_2^\infty \log\bigg(1-\frac{1}{y^2}\bigg)\pi '(x+y) dy < \log(\zeta(2)).$$ And, as the difference between both Prime Counting Functions will be at most 1, $$-\int_2^\infty \log\bigg(1-\frac{1}{y^2}\bigg)\pi '(x+y) dy < \log(\zeta(2))-\log\bigg(\frac{4}{3}\bigg) $$ $$-\int_2^\infty \log\bigg(1-\frac{1}{y^2}\bigg)\pi '(x+y) dy < \log\bigg(\frac{\pi^2}{8}\bigg) $$ Which seems to hold true, since the integral seems to monotonically decrease for large x. Where is the flaw? Thank you!

论坛问题帖 | no_answers_at_retrieval | user3141592 | 2017-02-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018713 | MSE-2137627

There are infinitely many prime numbers p for which $p + 2$ and $p + 4$ are also prime numbers

I need help with this question: There are infinitely many prime numbers p for which $p + 2$ and $p + 4$ are also primes. Where should I start? what proof techniques will be useful? Any input will be very useful to me.

论坛问题帖 | has_answers | Jamie John | 2017-02-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018714 | MSE-2145404

The $\gcd$ operator commutes with functions defined by linear recurrence relations

Given $a,b\in\mathbb Z^+$, and let $F_{a,b}:\mathbb N\to\mathbb N$ be a function such that $F_{a,b}(0)=0$ and $F_{a,b}(n+1)=a\cdot F_{a,b}(n)+b\cdot F_{a,b}(n-1)$. $F_{1,1}$ correspond to the Fibonacci function if $F_{1,1}(1)=1$. Conjecture which I would like to see proved: If a,b are co-prime and $F_{a,b}(1)\in \mathbb Z^+$, then $F_{a,b}(\gcd(m,n))=\gcd(F_{a,b}(n),F_{a,b}(m))$ I guess that some formulation of the conjecture is true for all homogeneous linear recurrence relations with constant coefficients of degree > 1, but I have only (randomly) tested it for degree 2. See also A generalization of a divisibility relation for Fibonacci numbers

论坛问题帖 | no_answers_at_retrieval | Lehs | 2017-02-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018715 | MSE-2145704

A generalization of a divisibility relation for Fibonacci numbers

Given $a,b\in\mathbb Z^+$, and let $F_{a,b}:\mathbb N\to\mathbb N$ be a function such that $F_{a,b}(0)=0$, $F_{a,b}(1)=1$ and $F_{a,b}(n+1)=a\cdot F_{a,b}(n)+b\cdot F_{a,b}(n-1)$. $F_{1,1}$ correspond to the Fibonacci function. Conjecture for which I would like to see a proof: If a,b are co-prime and $p>2$ is a prime that doesn't divide b, then $F_{a,b}\left(p-\left(\frac{a^2+4b}{p}\right)\right)\equiv 0\pmod p$, where $\left(\frac{a^2+4b}{p}\right)$ is the Legendre symbol. I have tested the conjecture with $10,000$ randomly selected a,b,p. Observation: for Fibonacci numbers $a=b=1$ and $a^2+4b=5$. Wikipedia See also The $\gcd$ operator commutes with functions defined by linear recurrence relations I've never been interested in the Fibonacci numbers and when I recently read about them on Wikipedia I realized that all this interesting theorems for Fibonacci numbers could be generalized to linear recurrence relations with constant coefficients. This funny relation $p|F_{p-(\frac{5}{p})}$ promted for an explanation. I guessed that the '5' came from the solution of the characteristic polynomial of the recurrence relation for the Fibonacci relation with the solutions $\frac{1\pm\sqrt{5}}{2}$. Subtracting the greater root from the smaller and square gives 5 and this seems to be correct. For the general case $x_{n+2}-ax_{n+1}-bx_n=0$ the characteristic polynomial is $t^2-at-b=0$ with the solutions $\frac{a\pm\sqrt{a^2+4b}}{2}$. Since I have an excellent system to check hypotheses (BigZ) the conjecture was easy to test computationally.

论坛问题帖 | has_answers | Lehs | 2017-02-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018716 | MSE-2147588

Do finitely generated ideals have a "dimension"?

In our algebra class, we recently talked briefly about finitely generated modules, and the instructor referred to how even if a R-module admits a basis, we can easily have that different bases of the module are not in bijection with each other. This brought to mind a different question: For a ring R and a finitely generated ideal I, need all minimal generators of I be in bijection? That is, if $S, T \subseteq I$ are two finite sets such that $I = (S) = (T)$, and moreover such that any proper subset of S or T will not generate I (i.e. if S and T are minimal), then does there exist a bijection between S and T? I raised the question with a friend in my class who's often interested in these kinds of questions, and he brought up that any (left) ideal I would carry a natural structure as a (left) R-module, so if there existed a finite ideal I that would induce a module with non-bijective bases, then the answer would be to the negative, but I have no intuition for whether that would be possible. If the answer is in general to the negative, then is there a natural assumption that could be made about R so that the conjecture does hold?

论坛问题帖 | has_answers | AJY | 2017-02-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018717 | MSE-2160267

Given polynomial $P(x)$, find polynomial $Q(x)$ that defines the same field with fixed degree terms

A conjecture about polynomials $P(x)$ and $Q(x)$ defining the same field is believed to be true, but unproven: Let $P(x)$ be an n degree polynomial. There exists a polynomial $Q(x)$ of degree n, with fixed of terms from degrees n to $(n/2)$ if n is even, and $(n+1)/2$ if n is odd, and varied terms degrees from $(n/2)-1$ to 0 if n is even, and degrees from $((n+1)/2)-1$ to 0 if n is odd such that $Q(x)$ defines the same field as $P(x)$ and also infinitely many polynomials of this form, but does not have the same discriminant as $P(x)$. Note that the fixed coefficients in $Q(x)$ do not have to necessarily match those in $P(x)$. Example of conjecture (assuming true): Let $P(x)$ $=$ x^3-16x^2+8x+1, since $P(x)$ is a 3 degree, or cubic polynomial, the first 2 coefficients can be fixed in $Q(x)$, and there exists infinitely solutions (a, b) such that $P(x)$ and $Q(x)$ define the same field where: $P(x)$ $=$ x^3-16x^2+8x+1 $Q(x)$ $=$ $4x^3+7x^2+ax+b$ Note the coefficients in any such $Q(x)$ may be chosen at random, and infinitely many polynomials of this form $Q(x)$ would define the same field $P(x)$. Second, can someone please explain an algorithm for solving the example problem assuming conjecture is true? Thanks for looking into this.

论坛问题帖 | has_answers | J. Linne | 2017-02-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018718 | MSE-2161989

Can we prove this conjecture concerning Pell equations?

For every positive integer n, not being a perfect square, denote the fundamental solution of the Pell equation $$x^2-ny^2=1$$ with (a,b). In other words, b is the smallest positive integer such that nb^2+1 is a perfect square. Define $f(n)=b$. Searching for n with large b , I came to the following conjecture : "If $s>46$ is a composite non-square number, there is a prime $p<s$ with $f(p)>f(s)."$ To formulate it in another way : If we write down every non-square s for which we get a new record for the largest number b (for all smaller numbers, we get a smaller b), the only composite numbers that are written down are 10 and 46. The other numbers written down are prime numbers. The conjecture is true for $s\le 10^6$, but can we prove it for all $s>46$?

论坛问题帖 | no_answers_at_retrieval | Peter | 2017-02-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018719 | MSE-2167296

Is there a Collatz-like sequence ending with $1,1,1,\ldots$?

I found a conjecture of some mathematics teacher. Is this known or solved problem? Is there a non-constant function f such that for any n, iterating $f(n)$ end to the sequence $1,1,1,1,1,\ldots$. By iteration I mean the similar way as in Collatz problem we end to the sequence $2,1,2,1,\ldots$. For example, does the function $f(n)=\begin{cases}n/3\text{ if }n\equiv 0\pmod 3 \\(4n-1)/3\text{ if }n\equiv 1\pmod 3\\(5n-7)/3\text{ if} n\equiv 2\pmod 3\end{cases}$ end to the sequence $\ldots, 1,1,1,\ldots$ This is from the Finnish mathematical journal from college students called Solmu, http://matematiikkalehtisolmu.fi/2017/1/otaksuma.pdf

论坛问题帖 | no_answers_at_retrieval | Jaakko Seppälä | 2017-03-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018720 | MSE-2170526

Consecutive prime powers $p^m<q^n$ such that $q^n-p^m=11$

Define two prime powers $p^m<q^n$, ($m,n\in\mathbb N$) to be consecutive if it doesn't exist a prime power r^k such that $p^m\!<r^k\!<q^n$. It seems that every even number is the difference between two consecutive prime powers, I haven't ran any large tests yet, but what about the odd numbers? Are there two consecutive prime powers $p^m<q^n$ and $i\in\mathbb N^+$ such that $q^n-p^m=11^i$? I have tested all consecutive prime powers less than $10,000,000$ without finding any solution to $q^n-p^m=11^i$. (A pattern seems to suggest that no odd numbers $N\ge 31$ have a solution to $q^n-p^m=N$, consecutive prime powers. But this must be wrong since prime powers has similar asymptotic distribution as the primes).

论坛问题帖 | has_answers | Lehs | 2017-03-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018721 | MSE-2172102

If p is a prime then p^2+26 is not a prime

I have tested for all primes less than $40,000$ and it seems that, if p is a prime then p^2+26 is not. I would like to see a proof or a counter-example.

论坛问题帖 | has_answers | Lehs | 2017-03-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018722 | MSE-2172126

For which $n\in\mathbb N$ exist a unique prime p such that $p^2+2n\in \mathbb P$

If $n\in\mathbb N$ is such that there is only one prime p such that $p^2+2n\in\mathbb P$ and if $m\in\mathbb N$ is such that there is only one prime q such that $q^2+2m\in\mathbb P$, then $3|n-m$. Tested computationally and I would like to see a proof or a counter-example. Alternative formulation of the test result below: If $n\in\mathbb N$ is such that there is only one prime p such that $p^2+2n\in\mathbb P$, then $n\equiv 1\pmod 3$. 0 value n : p^2+2n_isprime \ p -- flag locals| p | p prime p p * n 2* + prime and ; 1 to n { 1 10000 | p^2+2n_isprime } cardinality . 1 ok 2 to n { 1 10000 | p^2+2n_isprime } cardinality . 183 ok 3 to n { 1 10000 | p^2+2n_isprime } cardinality . 125 ok 4 to n { 1 10000 | p^2+2n_isprime } cardinality . 1 ok 5 to n { 1 10000 | p^2+2n_isprime } cardinality . 195 ok 6 to n { 1 10000 | p^2+2n_isprime } cardinality . 221 ok 7 to n { 1 10000 | p^2+2n_isprime } cardinality . 1 ok 8 to n { 1 10000 | p^2+2n_isprime } cardinality . 160 ok 9 to n { 1 10000 | p^2+2n_isprime } cardinality . 329 ok 10 to n { 1 10000 | p^2+2n_isprime

} cardinality . 1 ok 11 to n { 1 10000 | p^2+2n_isprime } cardinality . 286 ok 12 to n { 1 10000 | p^2+2n_isprime } cardinality . 129 ok 13 to n { 1 10000 | p^2+2n_isprime } cardinality . 0 ok 14 to n { 1 10000 | p^2+2n_isprime } cardinality . 340 ok 15 to n { 1 10000 | p^2+2n_isprime } cardinality . 232 ok 16 to n { 1 10000 | p^2+2n_isprime } cardinality . 1 ok 17 to n { 1 10000 | p^2+2n_isprime } cardinality . 123 ok 18 to n { 1 10000 | p^2+2n_isprime } cardinality . 182 ok 19 to n { 1 10000 | p^2+2n_isprime } cardinality . 1 ok 20 to n { 1 10000 | p^2+2n_isprime } cardinality . 174 ok 21 to n { 1 10000 | p^2+2n_isprime } cardinality . 295 ok 22 to n { 1 10000 | p^2+2n_isprime } cardinality . 1 ok 23 to n { 1 10000 | p^2+2n_isprime } cardinality . 161 ok 24 to n { 1 10000 | p^2+2n_isprime } cardinality . 249 ok 25 to n { 1 10000 | p^2+2n_isprime } cardinality . 1 ok 26 to n { 1 10000 | p^2+2n_isprime } cardinality . 198 ok 27 to n { 1 10000 | p^2+2n_isprime } cardinality . 129 ok 28 to n { 1 10000 | p^2+2n_isprime } cardinality . 0 ok 29 to n { 1 10000 | p^2+2n_isprime } cardinality . 453 ok 30 to n { 1 10000 | p^2+2n_isprime } cardinality . 362 ok 31 to n { 1 10000 | p^2+2n_isprime } cardinality . 1 ok I'm using BigZ on https://forthmath.blogspot.se Anyone is welcome to criticize the test or reformulate the question. Due to the comment of Ivan Neretin it seems like the following formulation is equivalent: If p is a unique prime such that $p^2+2n\in\mathbb P$, then $p=3$. And then I belive that the proof essentially is in this proof: Conditions on $n\in\mathbb N^+$ for $p\in\mathbb P\implies p^2+2n\notin\mathbb P$

论坛问题帖 | no_answers_at_retrieval | Lehs | 2017-03-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018723 | MSE-2172856

Conditions on $n\in\mathbb N^+$ for $p\in\mathbb P\implies p^2+2n\notin\mathbb P$

The condition on $n\in\mathbb N^+$ for $$\forall p\in\mathbb P:\, p^2+2n\notin\mathbb P$$ seems to be: $$n\equiv 1\pmod 3\wedge 9+2n\notin\mathbb P$$ I would like to see a proof or a counter-example.

论坛问题帖 | has_answers | Lehs | 2017-03-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018724 | MSE-2172922

There exists infinitely many N such that $\{\sum_{n=2}^N\log(n)\}<\varepsilon$

I am wondering whether or not the following result is true: For all $\varepsilon>0$, there exists infinitely many $N\in \mathbb N$ such that $$S_N:=\left\{\sum_{n=2}^N\log(n)\right\}<\varepsilon$$ where $\{\cdot\}$ denote the fractional part, i.e. $\{x\}=x-[x]$. It seems to be true, looking at these drawing of S_N for N up to 100 and $1\,000$ respectively. But I really don't think $1\,000$ is enough to forge an intuition regarding the veracity of the result. I tried a comparison series-integral, to evaluation $\sum_n \log(n)$: $$I_-=\int_2^{N-1}\log (x)\mathrm d x\leqslant \sum_{N=2}\log (n)\leqslant \int_2^{N} \log(x)\mathrm dx=I_+.$$ And we have: \begin{align*}\int_2^k \log (x) \mathrm d x&=[(-1+\log(x))x]_2^k \\ &=(-1+\log (k))k+(1-\log 2)2,\end{align*} so we can find an equivalent of both I_- and I_+: $$\begin{cases} I_-\underset{N\to\infty}\sim N\log N \\ I_+\underset{N\to\infty}\sim N\log N. \end{cases}$$ I then tried to find if $N\log(N)$ was equidistributed modulo 1, which would prove that there is infinitely many N such that $$N\log N-[N\log N]<\varepsilon.$$ It seems true looking a these images for $N\log N$ modulo 1 up to 520, $5\,000$ and $30\,000$ respectively: But I don't know how to prove it. We win if we show that $$\lim_{N\to\infty}\frac 1N\sum_{n=1}^N e^{2\pi i \ell n\log n}=0$$ for all $\ell\in\mathbb Z\setminus \{0\}$ by Weil's criterion, but I can't prove it. Plus, I don't think the equivalent would actually provide a proof of the result, would it? What other approach could be taken?

论坛问题帖 | no_answers_at_retrieval | E. Joseph | 2017-03-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018725 | MSE-2173314

What is wrong with this proof that there are no odd perfect numbers?

Main Question What is wrong with this proof that there are no odd perfect numbers? The "Proof" Euler proved that an odd perfect number N, if any exists, must take the form $N = q^k n^2$ where q is the Euler prime satisfying $\gcd(q,n)=1$ and $q \equiv k \equiv 1 \pmod 4$. Denote the sum of divisors of $x \in \mathbb{N}$ by $\sigma(x)$, and the deficiency by $D(x)=2x-\sigma(x)$. Finally, denote the abundancy index by $I(x)=\sigma(x)/x$. Consider the quantity $D(q^k)/\sigma(q^k)$: $$\dfrac{D(q^k)}{\sigma(q^k)}=\dfrac{D(q^k)}{q^k}\cdot\dfrac{1}{I(q^k)}=\dfrac{1}{I(q^k)}\cdot\bigg(2-I(q^k)\bigg)=I(n^2)-1.$$ Multiplying both sides by n^2, we get $$\dfrac{D(q^k)\cdot{n^2}}{\sigma(q^k)}=\sigma(n^2)-n^2 \in \mathbb{N}.$$ It is easy to show the following claims: Claim 1 $\sigma(q^k) \nmid D(q^k)$ Proof Suppose to the contrary that $\sigma(q^k) \mid D(q^k)=2q^k - \sigma(q^k)$. This implies that $\sigma(q^k) \leq 2q^k - \sigma(q^k)$, which contradicts $q^k + 1 \leq \sigma(q^k)$. QED Claim 2 $\sigma(q^k) \nmid n^2$ Proof Suppose to the contrary that $\sigma(q^k) \mid n^2$. Then $$\dfrac{n^2}{\sigma(q^k)}=\dfrac{\sigma(n^2)}{2q^k}$$ is an integer, contradicting the fact that $\sigma(n^2)$ is odd. QED Hence, $\sigma(q^k) \nmid \bigg(D(q^k)\cdot{n^2}\bigg)$, contradicting $$\dfrac{D(q^k)\cdot{n^2}}{\sigma(q^k)}=\sigma(n^2)-n^2 \in \mathbb{N}.$$

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2017-03-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018726 | MSE-217774

Has every finite group a minimal presentation?

For a finite group G let $d(G)$ be the minimal number of generators of G and let $r(G)$ be the minimal number such that G has a finite presentation with $r(G)$ relators. Call a presentation with $d(G)$ generators and $r(G)$ relators a minimal presentation. Question: Has every finite group a minimal presentation?

论坛问题帖 | no_answers_at_retrieval | Ralph | 2012-10-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018727 | MSE-2178323

Peculiar pattern concerning primes and semi-primes?

Define the function $f:\mathbb N^+\to\mathbb N^+$ as: $f(n)$ is the least semi-prime pq such that $n+p+q$ is a prime. (One must prove that this really is a function, but it should be and my tests doesn't contradicts that). There is a sequence of numbers $$21\to 14\to 15\to 10\to 9\to 6\to 4$$ such that if $f(n)\neq 4$ is a number in this sequence, then $f(n+1)$ equals to the successor of $f(n)$ in the sequence. This is tested for all values of $n<100,000$. I would like a proof of that f really is a function, that is, for all $n\in\mathbb N^+$ it exists a semi-prime pq with the property that $n+p+q\in\mathbb P$. Also, if anyone can find a counter-example from the peculiar pattern, I would like to see that. And off course, if anyone can explain the pattern I would be very glad.

论坛问题帖 | has_answers | Lehs | 2017-03-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018728 | MSE-2192217

Is $\frac{b^2 - 1}{\sigma(b^2) - b^2} + 1$ always an odd prime for odd composite b?

Let $\sigma(x)$ be the sum of the divisors of x. Here is my original question: If $\bigg(\sigma(b^2) - b^2\bigg) \mid \bigg(b^2 - 1\bigg)$, is $$\frac{b^2 - 1}{\sigma(b^2) - b^2} + 1$$ always an odd prime for b odd? Preliminary computations using Mathematica show that this must be true. Is it? Added March 18 2017 When $b = p^k$ where p is an odd prime and $k \geq 1$, we have $$\frac{b^2 - 1}{\sigma(b^2) - b^2} + 1 = \frac{\sigma(b^2) - 1}{\sigma(b^2) - b^2} = \frac{\sigma(p^{2k}) - 1}{\sigma(p^{2k}) - p^{2k}} = \frac{p\sigma(p^{2k-1})}{\sigma(p^{2k-1})} = p,$$ validating didgogns's claim in the comments. Hence, I modify my original question slightly as follows: MODIFIED QUESTION If $\bigg(\sigma(b^2) - b^2\bigg) \mid \bigg(b^2 - 1\bigg)$, is $$\frac{b^2 - 1}{\sigma(b^2) - b^2} + 1$$ always an odd prime for b an odd composite?

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2017-03-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018729 | MSE-219359

Does giving a counterexample to a conjecture prove it to be true or false?

If a problem asks me to give a counterexample to a conjecture, am I proving the conjecture true or false by giving a counterexample? I am leening towards proving it false, because if I were to prove it true, it would involve another technique (contradiction, direct, ...)

论坛问题帖 | has_answers | user29163 | 2012-10-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018730 | MSE-2211090

A new property of Rowland's sequence or there is a counterexample? Each element a_i is a witness of a greater a_j whose index j is always prime

Rowland's sequence, named after Professor Eric Rowland, is defined as: $$a_1=7$$ $$a_n=a_{n-1}+gcd(n,a_{n-1}), n \gt 1$$ Rowland already proved ("A Natural Prime-Generatin Recurrence", Journal of Integer Sequences, Vol.11, 2008, Article 08.2.8) that for any index n of the sequence, the difference $a_{n}-a_{n-1}$ is a prime number or 1. I would like to ask about an observation regarding Rowland's sequence, that might be a new property of the sequence. I found it just by trial and error, but as far I have tested it, it seems to be true up to index $i=498270791$ (after that point the computer calculations are very slow). The questions are at the end of the explanation. Basically the observation is (Expression 1): $$\forall i \ge 2, \ \exists j \gt i \ :\ \lfloor \frac{lcm(1..i)}{a_i} \rfloor \lt a_j \ \ ,\ \ j \in \Bbb P$$ $$ \land \not \exists k \lt j \ :\ \lfloor \frac{lcm(1..i)}{a_i} \rfloor \lt a_k$$ In other words, any a_i ($i \ge 2$) of the sequence seems to be a kind of "witness" or "marker" of a greater element of the sequence, a_j, whose index j is always a prime number. Here is the method of calculation: We take any index $i \ge 2$, and calculate the least common multiple of all the natural numbers up to the index: $lcm(i)$. Then we divide $lcm(i)$ by the current element of the sequence ${a_i}$. Our witness will be the floor of the number resulting of the division at step 2. Now we look for the first element a_j of Rowland's sequence greater than the witness of step 3 (the index j is also greater than i). The index of that number, j, seems to be always a prime number. For instance here is the list of the first 27 primes confirmed by this method (in other words, the index j that is found and is always prime). Some of them are repeated, for instance 2 and 47, because for those cases lcm([1..i])=lcm([1..i+1]) and the floor of the division in those cases is always smaller than the same a_j, and thus the index j is the same prime number: Another way to write the (Expression 1) is using the second Chebyshev function, which is exactly the natural logarithm of the least common multiple of the integers from 1 to n, so for that reason we can write as follows (Expression 2): $$lcm(1..n)=e^{\psi(n)}$$ $$\forall i \ge 2, \ \exists j \gt i \ :\ \psi(i)-log(a_i) \lt log(a_j)\ ,\ \ j \in \Bbb P$$ $$ \land \not \exists k \lt j \ :\ \psi(i)-log(a_i) \lt log(a_k)$$ or $$\forall i \ge 2, \ \exists j \gt i \ :\ \sum_{p^t \le i} p-log(a_i) \lt log(a_j)\ ,\ \ j \in \Bbb P$$ $$ \land \not \exists k \lt j \ :\ \sum_{p^t \le i} p-log(a_i) \lt log(a_k)$$ This is the PARI\GP code, please feel free to use it and modify it: \p200000; print();print("Calculating first 30 Rowland's witnesses:");print();testlimit = 30;current_rowland_seq_element=7;lcm_list=listcreate();listput(lcm_list,1);witnesses_list=listcreate();for(sequence_index=2,testlimit,current_rowland_seq_element=current_rowland_seq_element+gcd(sequence_index,current_rowland_seq_element);listput(lcm_list,sequence_index);listput(witnesses_list,truncate(lcm(lcm_list)/current_rowland_seq_element)););print(witnesses_list);print();print("Starting test:");print();sequence_index=2;current_rowland_seq_element=7;testlimit = 31247916400;was_found_prime=0;for(prospect=1,length(witnesses_list),if(witnesses_list[prospect]>testlimit,break);while(1,if(was_found_prime==0,current_rowland_seq_element=current_rowland_seq_element+gcd(sequence_index,current_rowland_seq_element),was_found_prime=0);if(current_rowland_seq_element>witnesses_list[prospect],print("(timer:",getabstime(),")\tThe index of the closest element of Rowland's sequence greater than:\t",witnesses_list[prospect],"\t is:\t",sequence_index,"\t . Primality test of the index (0=False, not prime / 1:True, is prime).

Result:\t",isprime(sequence_index));print();was_found_prime=1;break;);sequence_index=sequence_index+1;);); It seems quite interesting, but still do not know if there is a counterexample or not. My computer is quite slow, and the calculation of the next prime index will take it around 6 hours (probably my computer will run out of memory, but I am trying). I would like to ask the following questions: Is there a counterexample? If somebody reading this wants to use the PARI\GP code or create a quicker version in another language and try to look for a counterexample would be great. Maybe the observation does not hold for the next still not calculated indexes of the sample list of the picture and included in the PARI\GP code as well. If there is not a counterexample, why is a_i able to be the witness of a greater a_j whose index j is prime? It might be related with the relationship between the recurrence relation used by Rowland's sequence, which is based on the manipulation of the greatest common divisor, gcd, versus the value lcm($[1,i]$). The link of the expression with the second Chebyshev function is also interesting because it is very related with the primes in the interval $[1,i]$. Maybe the reason is very straightforward but I cannot see it. Any insights are very appreciated, thank you!

论坛问题帖 | has_answers | iadvd | 2017-03-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018731 | MSE-2216324

If $q^k n^2$ is an odd perfect number with Euler prime q, is it possible to have $n\sigma(n)=q\sigma(q^k)$?

QUESTION If $q^k n^2$ is an odd perfect number with Euler prime q, is it possible to have $n\sigma(n)=q\sigma(q^k)$? BACKGROUND If $\sigma(N)=2N$ (where $\sigma(N)$ is the sum of the divisors of N), then N is said to be perfect. (Denote the abundancy index of $x \in \mathbb{N}$ by $I(x)=\sigma(x)/x$.) Euler proved that every odd perfect number has the form $N=q^k n^2$, where q is prime (called the Euler prime) satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. MY ATTEMPT TO ANSWER THE QUESTION Let $N=q^k n^2$ be an odd perfect number with Euler prime q, and assume to the contrary that $n\sigma(n)=q\sigma(q^k)$. Since $\gcd(q,n)=1$, this implies that $$\frac{\sigma(n)}{q}=\frac{\sigma(q^k)}{n} \in \mathbb{N}.$$ Since $\sigma(q^k) \equiv k+1 \equiv 2 \pmod 4$ and n is odd, then $\sigma(q^k) \neq n$. This implies that $$2 \leq \frac{\sigma(n)}{q}=\frac{\sigma(q^k)}{n}.$$ But qn and $q^k n$ are both deficient. This means that $$\frac{\sigma(q)}{n}\cdot{2} \leq \frac{\sigma(q)}{n}\cdot\frac{\sigma(n)}{q} = I(qn) < 2 \implies \frac{\sigma(q)}{n} < 1 \implies q < n,$$ and $$\frac{\sigma(n)}{q^k}\cdot{2} \leq \frac{\sigma(n)}{q^k}\cdot\frac{\sigma(q^k)}{n} = I({q^k}n) < 2 \implies \frac{\sigma(n)}{q^k} < 1 \implies n < q^k.$$ Together, $q < n$ and $n < q^k$ imply that $k > 1$. Additionally, the equation $$n\sigma(n) = q\sigma(q^k)$$ is equivalent to $$\frac{\sigma(n)}{\sigma(q^k)}=\frac{q}{n},$$ which together with $q < n$, implies that $$\sigma(n) < \sigma(q^k).$$ Consequently, we have $$q < \sigma(q) < n < \sigma(n) < q^k < \sigma(q^k)$$ so that $$\frac{\sigma(q)}{n} < 1 < \frac{\sigma(n)}{q}$$ and $$\frac{\sigma(n)}{q^k} < 1 < \frac{\sigma(q^k)}{n},$$ whereupon I do not obtain any contradictions.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2017-04-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018732 | MSE-2228451

Can a Graph Edge participate in more than two Faces?

After learning some very useful Graph Theory in this question (What is the correct term for cycles more fundamental than Simple Cycles?), I realized that in my work I have been considering a conjecture, which I would like to know if it is a theorem. Has this been proven? For planar graphs, any edge of an undirected cyclic graph may participate in a maximum of two faces. Can anyone confirm or correct me?

论坛问题帖 | has_answers | philologon | 2017-04-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018733 | MSE-2230651

A conjectured continued fraction for ϕ^ϕ

As I previously explained, I am a "hobbyist" mathematician (see here or here); I enjoy discovering continued fractions by using various algorithms I have been creating for several years. This morning, I got some special cases for a rather heavy expression I was working on, finding the beautiful identities: $$ \phi^\phi \;=\; 2 + \displaystyle\frac{2\left(1-1/\phi\right)/\phi} {2\left(2-1/\phi\right) + \displaystyle\frac{3\left(1-2/\phi\right)/\phi} {3\left(2-1/\phi\right) + \displaystyle\frac{4\left(1-3/\phi\right)/\phi} {4\left(2-1/\phi\right) + \displaystyle\frac{5\left(1-4/\phi\right)/\phi} {5\left(2-1/\phi\right) + \;\ddots }}}} $$ Equivalently $$ \phi^\phi \;=\; 2 + \displaystyle\frac{2/\phi\left(\phi-1\right)} {2\sqrt5 + \displaystyle\frac{3\left(\phi-2\right)} {3\sqrt5 + \displaystyle\frac{4\left(\phi-3\right)} {4\sqrt5 + \displaystyle\frac{5\left(\phi-4\right)} {5\sqrt5 + \;\ddots }}}} $$ where ϕ is the golden ratio. I was very excited to have found this ϕ^ϕ formula and my question is about how original it is: did someone already see it or could I be confident in assuming it is new? Similarly $$ \phi^{2/\phi} \;=\; 2 + \displaystyle\frac{2\left(1-2/\phi\right)} {2\phi + \displaystyle\frac{3\left(1-3/\phi\right)} {3\phi + \displaystyle\frac{4\left(1-4/\phi\right)} {4\phi + \displaystyle\frac{5\left(1-5/\phi\right)} {5\phi + \;\ddots }}}} $$ By using a more compact notation (see at the end of this message for an explanation about how to read it), other identities for $\sqrt{3}$, $\sqrt[3]{4}$, $\sqrt[4]{5}$, etc. would be: $$ \sqrt[a]{a+1} \;=\;2+ \displaystyle\operatorname*{K}_{n=1}^{\infty} \frac {\left(n+1\right)\left(1-an\right)/\left(a+1\right)} {\left(n+1\right)\left(1+a/\left(a+1\right)\right)} $$ These identities are special case of the following conjectured identity (of course, when asking whether the previous formulae are known or not, it also means the formula below should be new also): $$ \left(\displaystyle\frac{x}{x-1}\right)^{x-1} \;=\;2+ \displaystyle\operatorname*{K}_{n=1}^{\infty} \frac {\left(n+1\right)\left(x-n-1\right)/x} {\left(n+1\right)\left(x+1\right)/x} \tag{1}\label{1} $$ that I found from a more general conjecture being related to the following continued fraction: $$ g\left(k,x\right) = \displaystyle\operatorname*{K}_{n=1}^{\infty} \frac {\left(n+1\right)\left(k-n-k/x\right)/x} {\left(n+1\right)\left(1+1/x\right)} $$ for which I empirically noticed the relation (with $\textrm{B}$ the beta function): $$ \begin{array}{l} 2\;+\; g\left(\alpha,\xi\right) \;+\;

g\left(\alpha,\displaystyle\frac{\xi}{\xi-1}\right) \\[8pt] \qquad\qquad =\; \alpha\left(\xi-1\right)^{\alpha/\xi-1}\left(\displaystyle\frac{\xi}{\xi-1}\right)^{\alpha-2} \textrm{B}\left(\alpha/\xi, \alpha-\alpha/\xi\right) \end{array} \tag{2}\label{2} $$ where it is easy to notice that the case $\alpha=\xi/\left(\xi-1\right)$ makes $g\left(\alpha,\xi\right)=0$ leading to an identity containing a single continued fraction (which can later be simplified as above). The previous notation is the one I use; I find it convenient and it can be found for instance in Continued Fractions with Applications by Lorentzen & Waadeland, but I know that some people don't like it; it has to be read the following way: $$ a_0 + \operatorname*{K}_{n=1}^{\infty} \frac{b_n}{a_n} = a_0 + \cfrac{b_1}{a_1 + \cfrac{b_2}{a_2 + \cfrac{b_3}{a_3 + \ddots}}} $$

论坛问题帖 | has_answers | Thomas Baruchel | 2017-04-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018734 | MSE-2251201

Correlation matrix inequality conjecture

Let A_{ij} be a correlation matrix, in other words it has the properties that $\forall i\, A_{ii}=1$ $|A_{ij}|\leq1$ $A_{ij}=A_{ji}$ A_{ij} is positive semi-definite, ie. $\forall y_i\,\sum_{ij}A_{ij}y_iy_j\geq 0$ I conjecture that if x_i is any vector of positive values, ie. $\forall i\,x_i\geq 0$, then $$ \forall i\quad\sum_jA_{ij}x_j\leq\sqrt{\sum_{jk}A_{jk}x_jx_k} $$ provided that the RHS is greater than 0 (not sure whether this last condition can be relaxed). Can someone please point me to a proof of this conjecture, or provide a counterexample?

论坛问题帖 | has_answers | Bob Mortimer | 2017-04-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018735 | MSE-2259725

Limits in higher dimension: is this double implication true?

Let $f:\mathbb{R}^n\rightarrow \mathbb{R}^m$ be a vector-valued function of n variables, and $x_0\in $ Dom$f $. Is the following proposition true? $$ \lim_{x\rightarrow x_0}\frac{f(x)}{||x-x_0||}=\underbrace{(0,..,0)}_{m\space times}\in \mathbb{R^m} \iff \lim_{x\rightarrow x_0}\frac{||f(x)||}{||x-x_0||}=0\in \mathbb{R}$$ If yes, how do you prove it? If not, can you show a counterexample?

论坛问题帖 | has_answers | Adrian | 2017-05-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018736 | MSE-2263932

Adding digits in this way to primes to obtain another primes

I just created the following "game": Choose some prime number. If it has k digits then add another k digits to the right to obtain another prime number. Now we have a prime number with $2k$ digits. Now add $2k$ digits to the right of the new number to obtain a new prime number. Repeat the procedure until it is no longer possible to produce new primes in this way. To clarify, let us take for example 5. We must add one digit to the right to obtain another prime number, it can be 53. Now we must add two digits to the right of 53 to obtain a new prime. It could be 5323. Now we must add four digits to the right of 5323 to obtain a new prime. It could be 53231117 and so on and so on... Is it true that whatever prime we choose at the beginning that this game never ends? That is, that it is always possible to build larger primes from the smaller ones in this way?

论坛问题帖 | has_answers | Living Hell | 2017-05-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018737 | MSE-2264657

Is this known about π?

Suppose that we travel along the digits of π and when we arrive at 2 or 3 or 4 or 5 or 6 or 7 or 8 or 9 we replace those digits with the number 1 and when we arrive at 1 or 0 we leave them unchanged. When we do this with all the digits will resulting number be irrational? Since this looks quite hard you can, if you want, assume that π is normal in order to prove this. But I do not know is normality enough? (The statement that we can suppose that π is normal was not in the original version of the question.)

论坛问题帖 | has_answers | Living Hell | 2017-05-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018738 | MSE-2266143

Is this limit relation true, and, if so, how to prove it?

Let $f:A\rightarrow \mathbb{R} $ be a function, where $A\subset \mathbb{R}$, and x_0 a limit point of A in which f is continuous, with $y_0=f(x_0)$. Suppose moreover that $g:\mathrm {Im}f\rightarrow \Bbb R$ is such that $g(y_0)=0$ and $$\lim_{y\rightarrow y_0} \frac{g(y)}{y-y_0}=0$$. Now, is it true that $$\lim_{x\rightarrow x_0}\frac{g(f(x))}{x-x_0}=0$$? If it is, how do you prove it rigorously?

论坛问题帖 | no_answers_at_retrieval | Adrian | 2017-05-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018739 | MSE-2269128

The conjecture: $\binom{2n-1}{n-1} \equiv 1 \pmod{n^3} \quad \Longrightarrow \quad n \in \mathbb{P}$

I recall seeing the following conjecture somewhere, but I cannot find the reference any more. Where can I find more information about this conjecture? Does it have a name? Conjecture: For any natural number n it holds that $$ \binom{2n-1}{n-1} \equiv 1 \pmod{n^3} \quad \Longrightarrow \quad n \in \mathbb{P}, $$ where $\mathbb{P}$ denotes the set of primes.

论坛问题帖 | has_answers | Mårten W | 2017-05-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018740 | MSE-2270159

Set conjectures concerning the asymptotic behaviour of erratic arithmetic functions, related to the Möbius function and the Liouville function

I get from an artificious way, but simple, two similar statements that I belive that are true (I had that identify some arithmetic functions), that is my Claim below. In next question I am asking about how to create good conjectures concerning two erratic arithmetic function, that will be introduced in this statement (see my motivation). Motivation. M1) For $n\geq 1$ was defined in the literature the sum of remainders function $$s(n)=\sum_{k=1}^n n\operatorname{mod}k.$$ Professor Spivey did a comparison in [1] between the properties that the sum of divisor function has versus this function. Is known that there are interesting identities with weights involving functions like Pillai function (I try to explore different ways to know more about the sum of remainders function). M2) From simple identities that appears in [2] multiplying by, corresponding the Möbius function $\mu(n)$ or the Liouville function $\lambda(n)$, and taking the sum, I can state the following (notice of the typography $\nmid$ in the first terms of LHS) Claim. A) For each $n\geq 1$ $$\varphi^{-1}(n)+\sum_{\substack{1\leq k\leq n \\ k\nmid n}}\mu(k)(n\operatorname{mod}k)=M(n)+\sum_{1\leq k\leq n}\mu(k)((n-1)\operatorname{mod}k)$$ where $\varphi^{-1}(n)$ is the Dirichlet inverse of Euler's totient function, and $M(n)$ is the Mertens function. B) For each $n\geq 1$ $$f(n)+\sum_{\substack{1\leq k\leq n \\ k\nmid n}}\lambda(k)(n\operatorname{mod}k)=L(n)+\sum_{1\leq k\leq n}\lambda(k)((n-1)\operatorname{mod}k)$$ where $f(n)$ is the Sloane's sequence A061020 (see [3] for the rest of references of our sequences), and $L(n)$ is the summary function $$L(n)=\sum_{k=1}^n\lambda(k).$$ M3) I have curiosity also about how to create good conjectures involving erratic functions. I am asking about how to combine mathematical reasonings, knowledges of the behaviour of the partial sums of Möbius and Liouville functions, and/or experiments with a computer to set conjectures that would be difficult to rule out.$\square$ Question. I was doing experiments and seems that the arithmetic functions defined for integers $n\geq 1$ $$R_1(n):=\sum_{1\leq k\leq n}\mu(k)((n-1)\operatorname{mod}k),$$ and $$R_2(n):=\sum_{1\leq k\leq n}\lambda(k)((n-1)\operatorname{mod}k)$$ are very erratic*. Imagine that I have a friend that ask me about a reasoning/method to get good conjectures about the asymptotic behaviour of these functions as $n\to\infty$. Are possible deductions to set such conjectures using mathematical ideas (reasonings, heuristics, numerical evidence)? What should be such conjectures? Many thanks. *You can see the behaviour of the sequence typying similar codes than these in Wolfram Alpha online calculator: sum mu(k)mod(10000-1,k), from k=1 to 10000 sum LiouvilleLambda(k)mod(50000-1,k), from k=1 to 50000 References: [1] Spivey, The Humble Sum of Remainders Function, Mathematics Magazine Vol. 78, No. 4 (2005). [2] James T. Cross, A Note on Almost Perfect Numbers, Mathematics Magazine, Vol. 47, No. 4 (1974). [3] The sequences A002321, A002819 and A023900 from The On-Line Encyclopedia of Integer Sequences.

论坛问题帖 | has_answers | user243301 | 2017-05-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018741 | MSE-2276799

Open Problems in Semigroup Theory.

NB: If this is a duplicate in any way or is too broad, I'm sorry. What are some open problems in Semigroup Theory? I'm reading Howie's "Fundamentals of Semigroup Theory" and I'm looking for problems accessible from that point of view.

论坛问题帖 | has_answers | Shaun | 2017-05-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018742 | MSE-2280915

proving of the given conjecture

let us suppose that there is given such kind of conjecture if their division is square of integer then we will have $\frac{a^2+b^2}{ab+1}=k^2$ for some integer k, then we would have $a^2+b^2=k^2*ab+k^2$ or we would have $a^2-k^2*a*b+b^2=k^2$ compare to the equation $a^2-2*a*b+b^2=(a-b)^2$ we would have $k^2=2$ but is that proof ? i could not understand if there is another option or alternative ways

论坛问题帖 | has_answers | dato datuashvili | 2017-05-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018743 | MSE-2297901

A Conjectured Mathematical Constant For Base-10 Normal Numbers.

Question 1: Let a be a real number with a base-10 decimal representation $a_1a_2\ldots a_n \ldots$ Denote the number of ways to write a_n as the sum of positive integers as $p(a_n)$ - also called the partition of a_n. I write $A(n)=\sum_{k=1}^n p(a_k)$ and $B(n)= \sum_{k=1}^n a_k$ and set $$\beta(a)=\lim_{n\to \infty}{A(n)\above 1.5 pt B(n)}$$ If a is a base-10 simply normal number is it true that $\beta(a)={97 \above 1.5 pt 45}$? Attempted Solution: Let a be a base-10 simply normal number with decimal representation $a_1a_2\ldots a_n \ldots$. Explicitly we can write \begin{equation}\lim_{n\to \infty}{A(n)\above 1.5 pt B(n)} =\lim_{n\to \infty}{p(a_1)+p(a_2)+\ldots +p(a_n)\above 1.5pt a_1+a_2\ldots +a_n }\end{equation} Surely $a_k \in [0,9]$ for every $1\leq k \leq n$. Now let $\nu_n(a_k)$ count occurrences of the digit a_k up to and including the n-th decimal place of a. Observe the following: the numerator in the above equation can be written explicitly as: $$ \underbrace{(p(0)+\ldots +p(0))}_{\nu_n(0)\text{ times}}+\ldots+\underbrace{(p(9)+\ldots + p(9))}_{\nu_n(9)\text{ times}}$$ and similarly the denominator may be written as $$\underbrace{(0+\ldots +0)}_{\nu_n(0)\text{ times}}+\ldots+\underbrace{(9+\ldots + 9)}_{\nu_n(9)\text{ times}}$$ Combining those two observation together we can rewrite the RHS of our first equation as $$ \lim_{n \to \infty}{\nu_n(0)\times p(0)+\nu_n(1)\times p(1)+\ldots + \nu_n(9)\times p(9)\above 1.5pt \nu_n(0)\times 0+ \nu_n(1)\times1+\ldots + \nu_n(9)\times9} $$ Recall that every digit of a occurs with equal frequency. In particular since a is simply normal we have that $\lim_{n \to \infty}\nu_n(a_j)=\lim_{n \to \infty}\nu_n(a_k)$ for every j and k. With out loss to generality we can substitute $\nu_n(j)$ with $\nu_n(1)$ in the numerators and denominators of the equation above and evaluate the limit, $$\lim_{n \to

\infty}{\nu_n(1)\times p(1)+\ldots + \nu_n(1)\times p(9) \above 1.5pt \nu_n(1)\times1+\ldots + \nu_n(1)\times9}$$ We can then pull out the term $ \nu_n(1)$, noticing those terms cancel out in the limit, and make a final simplification to show that $$\lim_{n \to \infty}{\nu_n(1) \above 1.5pt \nu_n(1)}{p(1) +\ldots +p(9)\above 1.5 pt 1+ \ldots +9}={1+1+1+2+3+5+7+11+15+22+30 \above 1.5pt 0+1+2+3+4+5+6+7+8+9}={ 97 \above 1.5pt 45}.$$ I believe the above solution to Question 1 is correct? Now my understanding is that we have the following implication: base-10 normal $\implies$ base-10 simply normal. So the follow up question seems natural. Question 2: If a is a base-10 normal is it true that $\beta(a)={97 \above 1.5 pt 45}$? The answer to Question 2 should be yes by implication alone. Moreover it should be true for both algebraic and transcendental base-10 normal numbers. Since I believe I have shown every base-10 simply normal number converges to ${97 \above 1.5 pt 45}$ under β and every base-10 normal number is simply normal then so to will every base-10 normal number converge to ${97 \above 1.5 pt 45}$ under β. It should be clear that the converse two Question 2 is false ! If we believe Questions 1 and 2 are true then we would expect several well known provably base-10 normal constants to converge to ${97 \above 1.5pt 45}$ under β. So I ran an experiment using data from Sloan's Database on the Erdos-Copeland constants which is proven to be normal in base-10. Using the first $2\text{ }000$ digits I plotted the values of $\beta(\text{Erdos-Copeland})$. The graph below illustrates the convergence. The blue line is the value of ${97 \above 1.5 pt 45}$ and the green graph are the values of the Erdos-Copeland constant under β. Up to the $2\text{ }000$-th digit $\beta(\text{Erdos-Copeland})=2.12999037999\ldots$ I believe given enough terms one can plot a similar graph for the Champernowne constant and one should expect the convergence to be significantly slower. Finally it is not known if π is normal in base-10. Out of curiosity I computed $\beta(\pi)$ up to the first $500\text{ }000$ digits of π. See the graph below. Again the blue line is the value of ${97 \above 1.5 pt 45}$ and the orange graph are the values of the π under β. Up to the $500\text{ }000$-th digit $\beta(\pi)=2.153781\ldots$. π appears to converge faster under β than any other mathematical constant I have looked into. I have numerous plots for zeta values and square roots and they all appear to approach ${97 \above 1.5 pt 45}$ which can be written numerically as $2.15555\ldots$. I have not attempted to plot numbers that are known to be abnormal in base-10.

论坛问题帖 | has_answers | Anthony | 2017-05-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018744 | MSE-2298823

Squares in $(\operatorname{rad}(1)^2+1)\cdot(\operatorname{rad}(2)^2+1)\ldots(\operatorname{rad}(n)^2+1)$

For integers $n\geq 1$, $\operatorname{rad}(n)$ denotes the radical of an integer, see in Wikipedia this definition $$\operatorname{rad}(n)=\prod_{p\mid n}p,$$ if $n>1$ with factorization $n=\prod_{p\mid n}p^{e_p}$, and defining $\operatorname{rad}(1)=1$. I'm inspired in a similar conjecture due to Amdeberhan, and Medina and Moll. Question. I believe that the following conjecture holds. Can you prove or refute it? There exists an integer N such that $\forall n>N$ $$\prod_{k=1}^n(1+(\operatorname{rad}(k))^2)$$ is not a square. Many thanks. References: Amdeberhan, and Medina and Moll, Arithmetical properties of a sequence arising from an arctangent sum, J. Number Theory (2007).

论坛问题帖 | has_answers | user243301 | 2017-05-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018745 | MSE-2304557

A Matrix With Eigenvalues Equal to The Golden Ratio and the Golden Conjugate

Question : Let $A(n)$ be a finite square $n \times n$ matrix with entries $a_{i,j}=1$ if $i+j$ is a perfect power; otherwise equals to 0. Can we determine the numbers n such that characteristic polynomial of $A(n)$ has at least two real roots: one equal to the golden ratio and the other equal to the golden conjugate ? For an examlple consider $$A(8)= \text{ }\begin{pmatrix} 0&0&1&0&0&0&1&1\\ 0&1&0&0&0&1&1&0\\ 1&0&0&0&1&1&0&0\\ 0&0&0&1&1&0&0&0\\ 0&0&1&1&0&0&0&0\\ 0&1&1&0&0&0&0&0\\ 1&1&0&0&0&0&0&0\\ 1&0&0&0&0&0&0&1\\ \end{pmatrix}$$ I denote the characteristic polynomial of $A(n)$ by $\chi_{A(n)}(X)$. So for example $$\chi_{A(8)}(X)=X^8-3X^7-5X^6+19X^5+2X^4-31X^3+X^2+10X-3$$ I use the standard notation φ to denote the golden ratio and it is equal to ${1+\sqrt{5}\above 1.5pt 2}$. The "Golden Conjugate" is written as $-{1\above 1.5 pt \varphi}$ and is equal to ${1-\sqrt{5}\above 1.5pt 2}$. I use λ to denote an eigenvalue of $\chi_{A(n)}(X)$. In the example above the characteristic polynomial $\chi_{A(8)}(X)$ has two eigenvalues $\lambda_1=-{1\above 1.5 pt \varphi}$ and $\lambda_2=\varphi$. In particular $\chi_{A(8)}(X)$ has exactly two roots: one equal to the golden ratio and the other equal to the golden conjugate. Below is a tabulation of data for small values of n. Calculations were performed in WOLFRAM ALPHA. The calculations should be correct but errors are inevitable. Only real eigenvalues are listed: \begin{array}{| l | l | l |l|} \hline n & \text{characteristic polynomial} &\text{eigenvalues}\\ \hline 2 & X^2-X & (0,1)\\ 3 & -X^3+X^2+X-1 & (-1,1,1)\\ 4 & X^4-2X^3+2X-1 & (-1,1,1,1)\\ 5 & -X^5+2X^4+2X^3-5X^2+X+1 &(-\lambda_1,\lambda_2,1,1,\lambda_3)\\ \color{blue}{6} & \color{blue}{X^6-2X^5-4X^4+8X^3+2X^2-4X-1} & \color{blue}{(1,-{1\above 1.5 pt \varphi},\varphi,\lambda_1,\lambda_2,\lambda_3)}\\ 7 & -X^7-2X^6+6X^5-11X^4-9X^3+15X^2+2X-4 & (1,\lambda_1,\lambda_2,\lambda_3,\lambda_4,\lambda_5,\lambda_6) \\ \color{blue}{8} & \color{blue}{X^8-3X^7-5X^6+19X^5+2X^4-31X^3+X^2+10X-3} & \color{blue}{(-{1\above 1.5 pt \varphi},\varphi,\lambda_1,\lambda_2,\lambda_3,\lambda_4,\lambda_5,\lambda_6)}\\ \color{blue}{9} & \color{blue}{-X^9+3X^8+6X^7-22X^6-5X^5+45X^4-12X^3-22X^2+5X+3} & \color{blue}{(\lambda_1,\lambda_2,\lambda_3,-\varphi,\varphi,1,-{1\above 1.5 pt \varphi},1-\varphi,\lambda_4)}\\ \hline \end{array} Surely $n=6,8,9$ are shown to answer the question. I am certain that if $A(n)$ has φ as a eigenvalue then it must also have $-{1\above 1.5 pt \varphi}$ as an eigenvalue. The motivation here is an understanding of the matrix $A(n)$. Edit 1: Correction to tabulated eigenvalues when n=6. There are actually 6 real eigenvalues and I listed only 3. Three of the eigenvalues can be written : 1, φ and $1-\varphi$. The other three however have exact forms that can be written with $i=\sqrt{-1}$. $$\lambda_1 = {\sqrt[3]{\frac{1}{2}\bigg(9 + i \sqrt{687}\bigg)}\above 1.5 pt 3^{2/3}} + {4 \above 1.5pt {\sqrt[3]{\frac{3}{2} \bigg (9 + i \sqrt{687}\bigg)}}}$$

$$\lambda_2 = {\bigg(1-i\sqrt{3}\bigg)\sqrt[3]{\frac{1}{2}\bigg(9 + i \sqrt{687}\bigg)}\above 1.5 pt 2\times 3^{2/3}} - {2(1+i\sqrt{3})\above 1.5pt {\sqrt[3]{\frac{3}{2} \bigg (9 + i \sqrt{687}\bigg)}}}$$ $$\lambda_3 = -{\bigg(1+i\sqrt{3}\bigg)\sqrt[3]{\frac{1}{2}\bigg(9 + i \sqrt{687}\bigg)}\above 1.5 pt 2\times 3^{2/3}} - {2(1-i\sqrt{3})\above 1.5pt {\sqrt[3]{\frac{3}{2} \bigg (9 + i \sqrt{687}\bigg)}}}$$ Numerically the eigenvalues λ_1,λ_2 and λ_3 are real - in particular the imaginary terms cancel out. Explicitly $\lambda_1\approx 2.11491\ldots$, $\lambda_2\approx -.618034\ldots$,$\lambda_3\approx -.254102\ldots$ Edit 2: $A(n)$ never has complex eigenvalues for any n. I included the missing eigenvalues in the table. Also I corrected errors in the table for $n=5$. None of the corrections change the result/conjecture of the problem. Namely that $6,8$ and 9 are the only integers such that $A(n)$ has a eigenvalues equal to φ or $-{1\above 1.5 pt \varphi}$. For $n=5$ the corrected values are numerically $\lambda_1 \approx 1.87939\ldots$, $\lambda_2\approx -1.53209\ldots$,$\lambda_3=1$,$\lambda_4=$, $\lambda_5 \approx -0.347296\ldots$ These values have exact forms similar to the radicals shown in Edit 1 above For $n=7$ the 6 missing values are numerically $\lambda_1 \approx 2.38839\ldots$, $\lambda_2\approx -1.9041\ldots$,$\lambda_3\approx 1.77217\ldots$,$\lambda_4\approx -1.33388\ldots$, $\lambda_5 \approx 0.0.64993\ldots$,$\lambda_6 \approx -0.572499 \ldots$. For $n=8$ the 6 missing values are numerically $\lambda_1 \approx 2.48767\ldots$, $\lambda_2\approx -1.98486\ldots$,$\lambda_3\approx 1.77268\ldots$,$\lambda_4\approx -1.39899\ldots$, $\lambda_5 \approx 0.827404\ldots$,$\lambda_6 \approx 0.296099 \ldots$. For $n=9$ the 4 missing values are numerically $\lambda_1 \approx 2.54926\ldots$, $\lambda_2\approx -2.03439\ldots$,$\lambda_3\approx 1.80552\ldots$,$\lambda_4\approx 0.320385\ldots$ These values have exact forms similar to the radicals shown in Edit 1 above. Conjecture: $6,8$ or 9 are the only numbers n such that characteristic polynomial of $A(n)$ has eigenvalues: one equal to the golden ratio and another equal to the golden conjugate.

论坛问题帖 | has_answers | Anthony | 2017-05-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018746 | MSE-2306090

A conjectured continued fraction related to a 2F1 hypergeometric function (and a formula for π)

In a previous question, some months ago, I described a continued fraction for which I was trying to find an identity. I finally found what I was looking for, and I will publish an answer to this question in the days or weeks to come by focusing to what was asked at that time. But now, I am opening a separate question for a new interest: Could the conjectured statements below be proved? Do these identities sound familiar to anyone? etc. My identity finally is: $$ \begin{array}{lcl} \mathcal{K}\left(a,x\right)&=&\displaystyle\operatorname*{K}_{n=1}^{\infty} \frac{ \left(n+1\right)\left(\left(a-1\right)x-n\right)a}{\left(n+1\right)\left(a+1\right)}\\[12pt] &=& \displaystyle\frac{\left(a-1\right)x-1}{x+1}\; \displaystyle{}_{2}F_{1}\left(1,ax\;; 2+x\;; a^{-1}\right) \end{array} \tag{1} \label{1} $$ Two functional identities I empirically noticed two functional identities, namely: $$ 2+\mathcal{K}\left(a,x\right)+\mathcal{K}\left(1-a,x\right) \;=\; 0 \tag{2} \label{2} $$ and (with $\textrm{B}$ the Beta function): $$ \begin{array}{l} 2+\displaystyle\frac{1}{a}\mathcal{K}\left(a,x\right) +\displaystyle\frac{a-1}{a}\mathcal{K}\left(a/\left(a-1\right), \left(a-1\right)x\right)\\[8pt] \qquad\qquad\;=\; ax\left(a-1\right)^{x-1}\left(\displaystyle\frac{a}{a-1}\right)^{ax-2}\textrm{B}\left(x,\left(a-1\right)x\right) \end{array} \tag{3} \label{3} $$ I already told about the latter in another question. A finite summation While studying the polynomials from the initial question, I finally found the following summation: $$ \mathcal{K}\left(a,x\right)\;=\; \displaystyle\frac{\displaystyle 2\sum_{k=2}^{\left(a-1\right)x-1} \left(k-1\right){{ak/\left(a-1\right)}\choose{k}} {a\left(\left(a-1\right)x-k\right)/\left(a-1\right) \choose{\left(a-1\right)x-k}} } {\displaystyle\left(\left(a-1\right)x-2\right){{ax}\choose{x}}} \tag{4} \label{4} $$ (when $(a-1)x$ is an integer greater than 2). A formula for π I tried several computer algebra systems and found the hyperexpand function in Sympy to be able of simplifying the ${}_2F_1$ hypergeometric function for many rational cases. My favorite case comes from $\mathcal{K}\left(-3, 1/2\right)$, after some cosmetic changes: $$ \pi \;=\; \operatorname*{K}_{n=1}^{\infty} \frac{ 81\left(n+1\right)\left(n+2\right)/16}{3\sqrt{3}\left(n+1\right)/2} \tag{5} \label{5} $$ The formula for ϕ^ϕ in the previous question was also a specific case for the same continued fraction.

论坛问题帖 | no_answers_at_retrieval | Thomas Baruchel | 2017-06-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018747 | MSE-2306801

If $N=q^k n^2$ is an odd perfect number, which is bigger, $\sigma(n^2)/n^2$ or $\sigma(n)/q^k$?

QUESTION If $N=q^k n^2$ is an odd perfect number, which is bigger, $\sigma(n^2)/n^2$ or $\sigma(n)/q^k$? BACKGROUND If $\sigma(N)=2N$ (where $\sigma(N)$ is the sum of the divisors of N), then N is said to be perfect. (Denote the abundancy index of $x \in \mathbb{N}$ by $I(x)=\sigma(x)/x$.) Euler proved that every odd perfect number has the form $N=q^k n^2$, where q is prime (called the Euler prime) satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. MY ATTEMPT TO ANSWER THE QUESTION Let $N=q^k n^2$ be an odd perfect number with Euler prime q. Suppose that $$\frac{\sigma(n^2)}{n^2}=\frac{\sigma(n)}{q^k}.$$ Then $$\frac{\sigma(q^k)}{n}\cdot t\frac{\sigma(n)}{q^k}<2=\frac{\sigma(q^k)}{q^k}\cdot\frac{\sigma(n^2)}{n^2}=\frac{ \sigma(q^k)}{q^k}\cdot\frac{\sigma(n)}{q^k}$$ so that $q^k < n$. Additionally, $$\frac{\sigma(n^2)}{n^2}=\frac{\sigma(n)}{q^k}$$ implies that $${q^k}\sigma(n^2) = {n^2}\sigma(n).$$ Therefore, $n^2 \mid {q^k}\sigma(n^2)$ and $q^k \mid {n^2}\sigma(n)$. Since $\gcd(q,n)=\gcd(q^k, n^2)=1$, then $n^2 \mid \sigma(n^2)$ and $q^k \mid \sigma(n)$. This implies that $$\frac{\sigma(n^2)}{n^2}=\frac{\sigma(n)}{q^k}$$ is an integer, contradicting $$1 < \frac{\sigma(n^2)}{n^2} < 2.$$ (In fact, we have the stronger bound $8/5 < \sigma(n^2)/n^2$.) Consequently, $$\frac{\sigma(n^2)}{n^2} \neq \frac{\sigma(n)}{q^k}.$$ Therefore, we either have $$\frac{\sigma(n)}{q^k} < \frac{\sigma(n^2)}{n^2} < 2$$ or $$\frac{8}{5} < \frac{\sigma(n^2)}{n^2} < \frac{\sigma(n)}{q^k}.$$ The second case implies $q^k < n$ since $$\frac{\sigma(n)}{n} < \frac{\sigma(n^2)}{n^2} < \frac{\sigma(n)}{q^k}.$$ The first case implies $$n <

{2^{\frac{\ln(12/13)}{\ln(9/13)}}}\cdot{(q^k)^{(\frac{\ln(13/12)}{\ln(13/9)}+\ln(13/9))}} \approx (1.162854)\cdot(q^k)^{0.5853949291}$$ if we use the estimate $$\frac{\sigma(n^2)}{n^2} < \bigg(\frac{\sigma(n)}{n}\bigg)^{\frac{\ln(13/9)}{\ln(4/3)}}.$$ Note that the inequality $$n < {2^{\frac{\ln(12/13)}{\ln(9/13)}}}\cdot{(q^k)^{(\frac{\ln(13/12)}{\ln(13/9)}+\ln(13/9))}} \approx (1.162854)\cdot(q^k)^{0.5853949291}$$ almost violates the inequality $q^k < n^2$. References The Abundancy Index of Divisors of Odd Perfect Numbers The Abundancy Index of Divisors of Odd Perfect Numbers - Part II

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2017-06-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018748 | MSE-2322336

How to prove that $p_{n+1}p_{n+2}p_{n+3}>p_{n+4}p_{n+5}$?

p_n is the n-th prime number: If $n > 3$ then: $$p_{n + 1} p_{n + 2} p_{n + 3} > p_{n + 4} p_{n + 5}$$ I checked the conjecture above true for first fifty million primes. If the conjecture above is true, it is stronger than Bonse's inequality. Could you give your remark, reference, or your proof of conjecture above?

论坛问题帖 | has_answers | Cố Gắng Lên | 2017-06-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018749 | MSE-2323482

Length of curve $X_2(t) = (t\cos t, t\sin t)$ where $(0 \le t \le 2\pi)$

Here I got $X_2(t) = (t\cos t, t\sin t)$ where $(0 \le t \le 2\pi)$ Let $X_2(t) = (x, y)$ then $x^2 + y^2 = t^2 \;\;\forall t\in[0, 2\pi]$ now will divide $[0,2\pi]$ into $n-identical$ segments. then we get $[0, \frac{2\pi}{n},2\frac{2\pi}{n},..k\frac{2\pi}{n}, (k+1)\frac{2\pi}{n},...n\frac{2\pi}{n}]$ then we take kth and $(k+1)th$ term, and measure the distance between two: $d(X_2(k\frac{2\pi}{n}),X_2((k+1)\frac{2\pi}{n})) = \sqrt{[k\frac{2\pi}{n}\cos k\frac{2\pi}{n}-(k+1)\frac{2\pi}{n}\cos((k+1)\frac{2\pi}{n})]^2+[k\frac{2\pi}{n}\sin k\frac{2\pi}{n}-(k+1)\frac{2\pi}{n}\sin((k+1)\frac{2\pi}{n})]^2}=\sqrt{k^2(\frac{2\pi}{n})^2+(k+1)^2(\frac{2\pi}{n})^2-2k(k+1)(\frac{2\pi}{n})^2[\cos k\frac{2\pi}{n}\cos (k+1)\frac{2\pi}{n}+\sin k\frac{2\pi}{n}\sin (k+1)\frac{2\pi}{n}]} $ then if we let $n\rightarrow \infty$, we can estimate that $[\cos k\frac{2\pi}{n}\cos (k+1)\frac{2\pi}{n}+\sin k\frac{2\pi}{n}\sin (k+1)\frac{2\pi}{n}] \rightarrow 1$ thus $d(X_2(k\frac{2\pi}{n}),X_2((k+1)\frac{2\pi}{n})) \rightarrow \frac{2\pi}{n}$ thus, total length would be $\frac{2\pi}{n}\cdot n = 2\pi$ I had developed my own conjecture as denoted above, and wants to be verified and get some correction for illogical reasoning. Any harsh ones are welcomed.

论坛问题帖 | has_answers | Beverlie | 2017-06-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018750 | MSE-2324643

Conjectured formula for the standard Clausen function

I often guess generalizations based on subtle patterns. I found the following formula in such a way and I wonder how it can be proven. Let n be a non-negative integer. Then \begin{align} \text{Cl}_{n}(x)&=(-1)^{\lfloor\frac{n-1}{2}\rfloor}x^{n-1}\left(\frac{H_{n-1}-\ln x}{\Gamma(n)}+\sum_{k=1}^{\lfloor\frac{n-1}{2}\rfloor}(-1)^k\frac{\zeta(2k+1)}{x^{2k}\Gamma(n-2k)}+2\sum_{k=1}^\infty\frac{\zeta(2k)\Gamma(2k)}{\Gamma(n+2k)}\left(\frac{x}{2\pi}\right)^{2k}\right) \\&=(-1)^{\lfloor\frac{n-1}{2}\rfloor}\left(\psi^{(-n)}(x)-\psi^{(-n)}(x+1)+\sum_{k=1}^{\lfloor\frac{n-1}{2}\rfloor}(-1)^k\frac{x^{n-2k-1}\zeta(2k+1)}{\Gamma(n-2k)}+2\sum_{k=1}^\infty\frac{x^{n+2k-1}\zeta(2k)\Gamma(2k)}{(2\pi)^{2k}\Gamma(n+2k)}\right) \end{align} where $\text{Cl}_{n}(x)$ is one of the standard Clausen functions defined by $$ \text{Cl}_{n}(x)= \begin{cases} \sum_{k=1}^\infty\frac{\sin(kx)}{k^n}& n\space\text{even}\\\\ \sum_{k=1}^\infty\frac{\cos(kx)}{k^n}& n\space\text{odd} \end{cases} $$ Edit: Upon further investigation, the proof can likely be derived from equations 2.7 and 2.8 of this paper.

论坛问题帖 | no_answers_at_retrieval | tyobrien | 2017-06-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018751 | MSE-2326215

A conjecture about primes

Conjecture: For each prime p there are an infinite number of primes q such that $p+q$ is a perfect square. I have done a lot of tests using Bigz and I believe that it's possible to prove. I change from possible to prove to not possible to disprove. Define for all primes q, $w_q(n)=|\{p\in\mathbb P|p<n\wedge \exists k\in\mathbb N:p+q=k^2\}|$. Then I've observed that it seems that for all primes q there exists a natural number N and a $\alpha\in\mathbb R_+$ such that $m>N$ implies that $w_q(10^m)>\alpha2^m$. This would imply the conjecture and is maybe possible to disprove.

论坛问题帖 | has_answers | Lehs | 2017-06-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018752 | MSE-232808

Primes + Inetvel + conjecture on primes

a) Can we establish a proof, there exists infinitely many primes of the form n^2 + 1. Why is the unit digit of such a prime p always 1 or 7? Is there any reasonable procedure or concept for the fact that the unit digit 7 occur essentially twice as often, when we identify the primes < 10000? b) can we prove or disprove that, there exists an interval of the form [n^2, $(n+1)^2$] containing at least 1000 prime numbers. c) We know that even integer > or = 4 can be written as sum of two primes and integers > 5 can be written as sum of three primes. Of course, those are conjectures. I am not asking the proof of those conjectures. I would like to know those statements are equivalent or not. If yes, how you will justify?

论坛问题帖 | has_answers | vmrfdu123456 | 2012-11-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018753 | MSE-2328423

A $3n-2$ Conjecture for the Redheffer Matrix.

The divisor matrix $D=(d_{i,j})_{i,j\in\mathbb{N}}$ is defined by \begin{equation*} d_{i,j}=\begin{cases}1,\quad \text{if i divides j,}\\ 0\quad \text{otherwise.}\end{cases} \end{equation*} A finite truncation of the divisor matrix obtained from the upper left $n\times n$ submatrix of D, setting each entry in the first column to 1 was first studied by Raymond Redheffer. For our purpose we denote such matrices by $D[n]$ and eponomously refer to them as Redheffer matrices. Question: Can we show that $\text{tr}\left(D[n]^{2}\right)=3n-2$ For example, if $n=3$ then $$ D[3]=\begin{pmatrix} 1 & 1 & 1\\ 1 & 1 & 0 \\ 1 & 0 & 1 \\ \end{pmatrix}.$$ If we denote by $p(X)$ the characteristic polynomial of $D[3]$, then its zeros are given by the cubical equation, $ -X^3+3X^2-X-1=0. $ This equation has roots: $1, 1+\sqrt{2}$ and $1-\sqrt{2}$. The eigenvalues of $D[3]$ are precisely the roots of $p(X).$ We can check that $$ 3\cdot 3-2=7=1+(1+\sqrt{2})^2+(1-\sqrt{2})^2. $$

论坛问题帖 | has_answers | Anthony | 2017-06-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018754 | MSE-2333801

If $N=q^k n^2$ is an odd perfect number and $q = k$, why does this bound not imply $q > 5$?

Let $\mathbb{N}$ denote the set of natural numbers (i.e., positive integers). A number $N \in \mathbb{N}$ is said to be perfect if $\sigma(N)=2N$, where $\sigma=\sigma_{1}$ is the classical sum of divisors. For example, $\sigma(6)=1+2+3+6=2\cdot{6}$, so that 6 is perfect. (Note that 6 is even.) Denote the abundancy index of $x \in \mathbb{N}$ as $I(x)=\sigma(x)/x$. Euler proved that an odd perfect number N, if any exists, must take the form $N=q^k n^2$, where q is prime satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. Suppose that $k=q$. Since q is prime and $q \equiv 1 \pmod 4$, this implies that $k \geq 5$. (In particular, $k \neq 1$, so that the Descartes-Frenicle-Sorli conjecture is false in this case.) Using WolframAlpha, we get the upper bound $$I(q^k)=I(q^q)=\frac{q^{q+1}-1}{{q^q}(q-1)} \leq \frac{3906}{3125} = 1.24992$$ which corresponds to the lower bound $$I(n^2)=\frac{2}{I(q^k)} \geq \frac{3125}{1953} \approx 1.6001\ldots.$$ Consider the product $$\bigg(I(q^q) - \frac{3906}{3125}\bigg)\bigg(I(n^2) - \frac{3906}{3125}\bigg).$$ This product is nonpositive. Therefore, $$I(q^q)I(n^2) + \bigg(\frac{3906}{3125}\bigg)^2 \leq \frac{3906}{3125}\cdot\bigg(I(q^q) + I(n^2)\bigg).$$ Since $N=q^k n^2$ is perfect with $q=k$, then $I(q^k)I(n^2)=I(q^q)I(n^2)=2$, so that $$I(q^q) + I(n^2) \geq \frac{3906}{3125} + \frac{3125}{1953} = \frac{17394043}{6103125} \approx 2.850022406554\ldots.$$ But in the paper [Dris, 2012 (pages 4 to 5)], it is proved that $$I(q^k) + I(n^2) \leq \frac{3q^2 + 2q + 1}{q(q+1)} = 3 - \frac{q-1}{q(q+1)}$$ with equality occurring if and only if $k=1$. In our case, since $k = q \geq 5$, we obtain $$\frac{17394043}{6103125} \leq I(q^q) + I(n^2) = I(q^k) + I(n^2) < 3 - \frac{q-1}{q(q+1)}$$ which implies that $$q > \frac{3125}{781} \approx 4.00128\ldots.$$ Here is my question: Why does the bound $$I(q^q) + I(n^2) \geq \frac{3906}{3125} + \frac{3125}{1953} = \frac{17394043}{6103125} \approx 2.850022406554\ldots$$ not imply that $q > 5$? I am thinking along the lines that: (1) $57/20 < I(q^k) + I(n^2) < 3$ is best-possible. (2) Improving the upper bound 3 would result in a finite upper bound for the Euler prime q. (3) Therefore, improving the lower bound $57/20$ would result in a lower bound for q better than the currently known $q \geq 5$. REFERENCES If $N=q^k n^2$ is an odd perfect number, does $q \leq 97$ imply that $I(q^k)+I(n^2) \leq 2.99$?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2017-06-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018755 | MSE-2340824

How often does $D(n^2) = m^2$ happen, where $D(x)$ is the deficiency of x?

Let $\sigma=\sigma_{1}$ denote the classical sum of divisors. For instance, $\sigma(12) = 1 + 2 + 3 + 4 + 6 + 12 = 28$. Let $x \in \mathbb{N}$ ($\mathbb{N}$ is the set of natural numbers/positive integers). Denote the number $2x - \sigma(x)$ by $D(x)$. We call $D(x)$ the deficiency of x. Now, let $m, n \in \mathbb{N}$. Here is my question: How often does $D(n^2) = m^2$ happen, where $D(x)$ is the deficiency of x? MY ATTEMPT I note (from the OEIS sequence listing) that $D(n^2) = m^2$ holds when $$(m, n) = \left\{(1, 1), (1, 2), (1, 4), (1, 8), \ldots \right\}$$ So, $n=2^r$ for integers $r \geq 0$ is an infinite family of numbers satisfying $D(n^2) = m^2$ (in particular, $m = 1$). Are $n=2^r$ for $r \geq 0$ the only numbers for which $D(n^2) = m^2$?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2017-06-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018756 | MSE-2341424

Do there exist deficient-perfect square numbers whose deficiency is also a square (and not one)?

This is a follow-up to this MSE question: How often does $D(n^2) = m^2$ happen, where $D(x)$ is the deficiency of x? Here is my question in this post: Do there exist examples for which $D(n^2) = m^2$ and $m \mid n$, where $m, n \in \mathbb{N}$ and $m > 1$? MY ATTEMPT In the linked MSE question, the following examples for $D(n^2) = m^2$ (without the divisibility constraint) were given: $$(m_1, n_1) = (19, 46), m_1 \nmid n_1$$ $$(m_2, n_2) = (53, 284), m_2 \nmid n_2$$ $$(m_3, n_3) = (3509, 149728), m_3 \nmid n_3$$ $$(m_4, n_4) = (6479, 242656), m_4 \nmid n_4$$ $$(m_5, n_5) = (1583, 1633), m_5 \nmid n_5$$ $$(m_6, n_6) = (223309, 260495), m_6 \nmid n_6$$ Rephrasing my question a bit, I have: Do there exist deficient-perfect square numbers whose deficiency is also a square (and not one)? Added on July 09 2017 I have the following bounds for $D(n^2)/n^2$ in terms of an auxiliary variable q: $$\frac{2}{q+1} \leq \frac{D(n^2)}{n^2} < \frac{2}{q}.$$ The conditions specified in my problem imply that $$\frac{2}{q+1} = \frac{m^2}{n^2}.$$ Added on July 12 2017 - MOTIVATION When $N=q^k n^2$ is an odd perfect number, then the following equations hold: $$\gcd\left(n^2,\sigma(n^2)\right)=\frac{D(n^2)}{\sigma(q^{k-1})}=\frac{\sigma(n^2)}{q^k}=\frac{2n^2}{\sigma(q^k)}.$$ (This is proved in this preprint, to appear in NNTDM.) In particular, when the Descartes-Frenicle-Sorli conjecture that $k=\nu_{q}(N)=1$ is true, then $D(n^2) \mid n^2$. In the other direction, Broughan, et. al (page 7, Lemma 8) proved that if $$\frac{\sigma(N/q^k)}{q^k}=\frac{\sigma(n^2)}{q^k}$$ is a square, then $k=1$.

Putting these all together, under the assumption that $\sigma(n^2)/q^k$ is a square, we have that $$1 < m^2 = D(n^2) \mid n^2.$$ Now, MSE user Matthew Conroy checked all $n < {10}^7$ and found only one ODD deficient-perfect square (other than 1), namely $n^2 = {3003}^2 = 9018009$, with deficiency 819. (Notice that the deficiency is not a square.) All other deficient-perfect squares are powers of 2, up to $n < {10}^7$. Indeed, if $n = 2^k$ for some integer $k \geq 0$, then n^2 is deficient-perfect with deficiency 1.

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2017-06-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018757 | MSE-2346852

Area of enclosed smooth curve is always irrational for rational dimensions

I have the intuition that smooth curves that are enclosed cannot be possible in real world , so it must be the case that either the dimensions(parameters) that define the curve (like the radius of circle) are irrational or for real dimensions the area of the curve is irrational like for example for circle () and ellipse (). Is there any such proof (for or against) for this conjecture for generic smooth enclosed curve ?.

论坛问题帖 | has_answers | Vikram Bhat | 2017-07-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018758 | MSE-2347178

Do all terms n of OEIS sequence A228059 have a p with exponent 1?

Do all terms n of OEIS sequence A228059 have a p with exponent 1? OEIS sequence A228059: Odd numbers of the form $p^{1+4k}{r^2}$, where p is prime of the form $1+4m$, $r > 1$, and $\gcd(p,r) = 1$ that are closer to being perfect than previous terms. Here are the first couple of terms: $$45 = 5\cdot{3^2}$$ $$405 = 5\cdot{3^4}$$ $$2205 = 5\cdot(3\cdot7)^2$$ $$26325 = 13\cdot({3^2}\cdot5)^2$$ $$236925 = 13\cdot({3^3}\cdot5)^2$$ $$1380825 = 17\cdot(3\cdot5\cdot19)^2$$ $$1660725 = 61\cdot(3\cdot5\cdot11)^2$$ $$35698725 = 61\cdot({3^2}\cdot5\cdot17)^2$$ $$3138290325 = 53\cdot({3^4}\cdot5\cdot19)^2$$ UPDATE - July 05 2017 (7:30 PM - Manila time): I am currently running the Mathematica code referenced in the OEIS sequence to compute more terms past 3138290325. UPDATE - July 06 2017 (1:00 AM - Manila time): The Mathematica code is still running and is currently at 35698725. It has not displayed 3138290325 yet.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2017-07-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018759 | MSE-2348700

A conjecture on diophantine equation

Let A, B, C be given three integer numbers, $A^2+B^2+C^2 \ne 0$; and $f(x,y,z)=0$ be given diophantine equation. Then exist $N_0 > 2$ such that $m, n, k \ge N_0$ the equation as follows: $f(x,y,z)+Ax^n+By^m+Cz^k=0$ with condition $gcd(x,y)=gcd(y,z)=gcd(z,x)=1$ has only finitely solutions Question: The conjecture above is true?

论坛问题帖 | has_answers | Cố Gắng Lên | 2017-07-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018760 | MSE-234877

Fibonacci conjecture: $(F_{n+5})^2 - (F_n)^2 = 3((F_{n+3})^2 - (F_{n+2})^2) + 8 F_{n+2} F_{n+3} $.

So this is the question I have The Fibonacci sequence is a recurrence system given by $$F_1 = 1, \ F_2 = 1, \ F_{n+2} = F_{n+1} + F_n \qquad (n = 1, 2, 3, \ldots).$$ This question concerns the following conjecture: $$(F_{n+5})^2 - (F_n)^2 = 3((F_{n+3})^2 - (F_{n+2})^2) + 8 F_{n+2} F_{n+3} \qquad (n = 1, 2, 3, \ldots).$$ (a) Confirm that the conjecture is true when $n = 6$. (b) Prove that the conjecture is true for all integers $n \geq 1$. But when I go through the solution I don't understand one line (separated and marked). (b) Let n be any integer such that $n \geq 1$. $$\begin{align} \left(F_{n+5} \right)^2 - \left(F_n \right)^2 &= \left(F_{n+5} - F_n \right) \left(F_{n+5} + F_n \right) \\ &= \left(F_{n+4} + F_{n+3} - F_{n+2} + F_{n+1} \right) \left(F_{n+4} + F_{n+3} + F_{n+2} - F_{n+1} \right) \\ \\ &= \left(\color{red}{2} F_{n+3} + F_{n+1} \right) \left(2 F_{n+3} + 2 F_{n+2} - F_{n+1} \right) \tag{*} \\ \\ &= \left(2 F_{n+3} - F_{n+3} - F_{n+2} \right) \left(2 F_{n+3} + 2 F_{n+2} - F_{n+3} + F_{n+2} \right) \\ &= \left(3 F_{n+3} - F_{n+2} \right) \left(F_{n+3} + 3 F_{n+2} \right) \\ &= 3 \left(\left(F_{n+3} \right)^2 - \left(F_{n+2} \right)^2 \right) + 8 F_{n+2} F_{n+3} \end{align}$$ Hence, the conjecture is true for all integers $n \geq 1$ In the marked line, where does the red $\color{red}{2}$ come from?

论坛问题帖 | has_answers | JackyBoi | 2012-11-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018761 | MSE-2353721

Parity-dependent pth-powers conjecture

For $p\in\mathbb{N}$ a monic polynomial q_p exists with $S_p(n):=\sum_{k=1}^nk^p=\frac{n(n+1)}{p+1}q_p(n)$. After $q_1=1$ we seem to have an alternating pattern, with $$q_2=n+\frac{1}{2},\\q_3=n(n+1),\\q_4=(n^2+n-\frac{1}{3})q_2,\\q_5=(n^2+n-\frac{1}{2})q_3,\\q_6=(n^4+2n^3-n+\frac{1}{3})q_2,\\q_7=(n^4+2n^3-\frac{1}{3}n^2=\frac{4}{3}n+\frac{2}{3})q_3,\\q_8=(n^6+3n^5+n^4-3n^3-\frac{1}{5}n^5+\frac{9}{5}n-\frac{3}{5})q_2,\\q_9=(n^2+n-1)(n^4+2n^3-\frac{1}{2}n^2-\frac{3}{2}n+\frac{3}{2})q_3.$$In particular we have the following conjectures: For $k\ge 1$ degree-($2k-2$) monic polynomials $A_k,\,B_k$ exist for which $q_{2k}=q_2A_k,\,q_{2k+1}=q_3B_k$ so that S_{2k} has repeated roots $-1,\,0$ and S_{2k+1} has roots including $-1,\,-\frac{1}{2},\,0$; For $k\ge 2$ the polynomials $A_k,\,B_k$ have the same next-to-leading coefficients. Can we prove, or at least make some sense of, any of these results? It feels like the $-\frac{1}{2}$ root and the repeated roots have some interpretation, and the fact that A_k-B_k is of degree $\le 2k-4$, should have some significance.

论坛问题帖 | no_answers_at_retrieval | J.G. | 2017-07-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018762 | MSE-2354517

The minimal combination $\big|\pm p_1\pm\dots\pm p_n\big|$ for primes

Conjecture: The minimal combination $\big|\pm p_1\pm\dots\pm p_n\big|$ is 0 for odd $n>1$ and 1 for even n, where p_n is the n-th prime. Tested for $n\leq 17$ by me and for $n\leq 28$ by user AugSB. $\min \big|\pm p_1\pm\dots\pm p_n\big|=\frac{1}{2}\big(1+(-1)^n\big)$, if $n>1$.

论坛问题帖 | has_answers | Lehs | 2017-07-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018763 | MSE-2355197

conjectured continued fraction related to certain polynomials

Let n be any complex number and q be a real number such that $q\in\Big(-\frac{1}{4};\frac{1}{4}\Big)$,$n\in\Big(-\infty;\infty\Big)$,it is then conjectured that the following is true $$G(q,n)=\cfrac{1}{1-q+\cfrac{q(1-q)^n}{1-q^3+\cfrac{q(1-q^3)^n}{1-q^5+\cfrac{q(1-q^5)^n}{1-q^7+\cfrac{q(1-q^7)^n}{1-q^9+\ddots}}}}}\\=1+(n+1)\frac{q^2}{1!}-(n^2+n+4)\frac{q^3}{2!}+(n^3+6n^2+23n+30)\frac{q^4}{3!}-(n^4+22n^3+71n^2+218n+384)\frac{q^5}{4!}+(n^5+65n^4+265n^3+1315n^2+3634n+6000)\frac{q^6}{5!}-(n^6+171n^5+1225n^4+6765n^3+24334n^2+70464n+118080)\frac{q^7}{6!}+(n^7+420n^6+5908n^5+33810n^4+159859n^3+595770n^2+1638552n+2766960)\frac{q^8}{7!}-(n^8+988n^7+27146n^6+179200n^5+1057049n^4+4580212n^3+16233244n^2+44611440n+75519360)\frac{q^9}{8!}+ \dots$$ Special cases ($n=-1$) $$\cfrac{1}{1-q+\cfrac{\frac{q}{\Big(1-q\Big)}}{1-q^3+\cfrac{\frac{q}{\Big(1-q^3\Big)}}{1-q^5+\cfrac{\frac{q}{\Big(1-q^5\Big)}}{1-q^7+\ddots}}}}=1-2q^3+2q^4-9q^5+29q^6-92q^7+316q^8-\dots$$ ($n=i$) $$\cfrac{1}{1-q+\cfrac{q(1-q)^i}{1-q^3+\cfrac{q(1-q^3)^i}{1-q^5+\cfrac{q(1-q^5)^i}{1-q^7+ \dots}}}}=1+\Big(1+i\Big)q^2-\Big(\frac{3}{2}+\frac{i}{2}\Big)q^3+\Big(4+\frac{11i}{3}\Big)q^4-\Big(\frac{157}{12}+\frac{49i}{6}\Big)q^5+\Big(\frac{475}{12}+\frac{337i}{12}\Big)q^6- \dots$$ where $i=\sqrt{-1}$ Note that by analysing the conjectured form it is evident that the coefficients of the series expansion are polynomial sequences. The form of the RHS suggests that the continued fraction is the exponential generating function $G(q)=\sum_{n=1}^{\infty}\frac{a_{n}}{n!}q^{n}$ for the given polynomials a_{n}(assuming we ignore matters of convergence). How do we prove that the conjecture is true? and if true,can it be related to some well established theory? PS:The radius of convergence had to be revised based on @mercio's argument supported by @Einar Rødland

论坛问题帖 | has_answers | Nicco | 2017-07-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018764 | MSE-2359790

k-tuple Admissibility?

Admissibility is defined for the integer k-tuples by the non-existence of a prime number p, for which each element of the k-tuple under union forms a complete residue system $\pmod{p}$. If we define $H \subset \mathbb{Z}$ as a k-tuple and $H' = (\mathbb{Z}/m\# \mathbb{Z})^\times$, the multiplicative group of residue classes with a primorial modulus, such that the modulus $m\#$ is greater than the diameter of H. Is it equivalent to say that if there exists a number $n \in \mathbb{Z}$, such that the translation on the number line $H \oplus n \subseteq H'$, that H is admissible?

论坛问题帖 | no_answers_at_retrieval | user56983 | 2017-07-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018765 | MSE-2360165

Euler Totient of Numbers Between Twin Primes.

Are there any known special properties of a number located between twin primes? The question came up in the discussion. (The expression below has been rephrased to a weaker form for clarity) In studying admissibility of k-tuples I observed that $$\varphi(\lambda) \le \lim_{\lambda < n \le \infty} \inf \varphi(n) $$ where we use the set theoretical representation of a twin prime pair as $\{\lambda-1, \lambda+1\} \subset \mathbb{P}$ and the definition of $\lim \inf x_n$ is given by $\lim_{N\rightarrow \infty} \inf x_k: k \in [N, \infty]$, and I'm not quite sure how to prove the statement above on the totient of a "lumenal" (?) number beyond conjecture. I think amWhy is curious as well. This post may be related, because this is a special case of the same problem. Now, of course, with the explicit formula for Euler totient, $$ \varphi(n) = n \prod_{p|n} \bigg(1 - \frac{1}{p} \bigg) $$ we do, in fact, end up with $\frac{ \varphi(n)}{n}$ having the product form in terms of the prime divisors of n giving the value $n-1$ when $n \in \mathbb{P}$. If we were not prime, then there would be another factor less than one in the formula for totient. Similar logic applies for the number $n+1$, because $n+1$ cannot be prime and as a result $\varphi(n+1) < n$. But this doesn't entirely explain the conjectured expression. For context, we arrive at a similar expression for the totient of a prime number $$\varphi(p) > \sup_{0 < n < p} \varphi(n) $$

论坛问题帖 | no_answers_at_retrieval | user56983 | 2017-07-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018766 | MSE-2363374

Number of monomials in $a_n=a_{n-1}+(a_{n-2})^2$ with $a_1=a$, $a_2=b$

I was playing around with the sequence $a_1=a$, $a_2=b$ and the recurrence $a_n=a_{n-1}+(a_{n-2})^2$ and just listed out a few general terms. For instance, $a_1=a$, $a_2=b$, $a_3=b+a^2$, $a_4=b+b^2+a^2$, $a_5=a^4+2a^2b+a^2+2b^2+b$, and $a_6=b^4+2b^3+3b^2+2a^2b^2+4a^2b+b+2a^4+a^2.$ When I did that, I realized that the number of terms in a_n appears to be F_n where F represents the Fibonacci numbers. I conjecture that this is true, but I am not sure how to even approach this.

论坛问题帖 | has_answers | Kai Williams | 2017-07-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018767 | MSE-2368370

Does the Rogers-Ramanujan continued fraction $R(q)$ satisfy this conjectured infinite series

Given the Rogers-Ramanujan continued fraction $R(q)= \cfrac{1}{1+\cfrac{q}{1+\cfrac{q^2}{1+\cfrac{q^3}{1+\ddots}}}}$ where $q=\exp(2\pi i \tau)$, $|q|\lt1$ for the sake of brevity, let us introduce the following notation $R_{0}=\frac{1}{R(q)}$ $R_{1}=\frac{1}{\Big(\frac{1}{q}\Big(\frac{1}{R(q)}-1\Big)\Big)}$ $R_{2}=\frac{1}{\Big(\frac{1}{q^2}\Big(\frac{1}{\Big(\frac{1}{q}\Big(\frac{1}{R(q)}-1\Big)\Big)}-1\Big)\Big)}$ $R_{3}=\frac{1}{\Big(\frac{1}{q^3}\Big(\frac{1}{\Big(\frac{1}{q^2}\Big(\frac{1}{\Big(\frac{1}{q}\Big(\frac{1}{R(q)}-1\Big)\Big)}-1\Big)\Big)}-1\Big)\Big)}$ up to R_{n}, $\frac{1}{R_{n}}=R(q^n,q)$ for natural number n, where $R(a,q)$ is the Generalized Rogers-Ramanujan continued fraction as pointed out by @ccorn It is then conjectured that the following infinite series holds $\displaystyle -\frac{R'(q)}{R(q)}=\frac{1}{R_{0}R_{1}}-\frac{2q^2}{R_{0}R^2_{1}R_{2}}+\frac{3q^5}{R_{0}R^2_{1}R^2_{2}R_{3}}-\frac{4q^9}{R_{0}R^2_{1}R^2_{2}R^2_{3}R_{4}}+\frac{5q^{14}}{R_{0}R^2_{1}R^2_{2}R^2_{3}R^2_{4}R_{5}}-\dots\tag1$ It has the ascending continued fraction equivalent $\displaystyle -\frac{R'(q)}{R(q)}= \frac{\frac{1}{R_1}+\large{\frac{\frac{-2q^2}{R_2}+\large{\frac{\frac{3q^5}{R_3}+...}{R^2_2}}}{R^2_1}}}{R_0} \tag2$ After considering the Rogers-Ramanujan continued fraction with the factor $q^{1/5}$ and applying the identity $\frac{R'(q)}{R(q)}=\frac{1}{5q}\frac{(q;q)^5_{\infty}}{(q^5;q^5)_{\infty}}$ due to Ramanujan,we are led to the following beautiful identity $\displaystyle \frac{(q)^5_{\infty}}{(q^5)_{\infty}}=1-\frac{5q}{(R_{0}R_{1})}+\frac{10q^3}{(R_{0}R_{1})(R_{1}R_{2})}-\frac{15q^6}{(R_{0}R_{1})(R_{1}R_{2})(R_{2}R_{3})}+\frac{20q^{10}}{(R_{0}R_{1})(R_{1}R_{2})(R_{2}R_{3})(R_{3}R_{4})}-\frac{25q^{15}}{(R_{0}R_{1})(R_{1}R_{2})(R_{2}R_{3})(R_{3}R_{4})(R_{4}R_{5})}+\dots\tag3$ $\displaystyle \frac{(q)^5_{\infty}}{(q^5)_{\infty}}=1+5\sum_{n=1}^{\infty}\frac{(-1)^n nq^{\frac{n(n+1)}{2}}}{\prod_{k=1}^{n-1}\Big(R_{k}+q^k\Big)}$ Question:How do we prove that the conjecture is true? Remark As entry 9 in chapter 19 of Ramanujan's second notebook we have the ff well known identity $\displaystyle \frac{(q)^5_{\infty}}{(q^5)_{\infty}}=1-5\sum_{n=1}^{\infty}\left(\frac{n}{5}\right) \frac{n q^n}{1-q^n}\tag{4}$ Combining (1) and Ramanujan's identity (4), we are immediately led to $\displaystyle \sum_{n=1}^{\infty}\left(\frac{n}{5}\right) \frac{n q^n}{1-q^n}=\sum_{n=1}^{\infty}\frac{(-1)^{n-1} nq^{\frac{n(n+1)}{2}}}{\prod_{k=1}^{n-1}\Big(R_{k}+q^k\Big)}\tag{5}$

论坛问题帖 | has_answers | Nicco | 2017-07-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018768 | MSE-2369296

Conjecture about distribution of certain primes

Conjecture: Let $f(x,y,z)=xy+xz+yz\:$ and $\: S_m=\{n\in\mathbb N|n<m\wedge f(p_n,p_{n+1},p_{n+2})\in\mathbb P\}$. Then $|S_m|\sim \frac{m}{\ln m}$. Computationally tested only, but I would like to see a proof. It seems like I made a mistake with the inequality in the comment, but here are some data: m m/ln m |Sm| pi(m) li m |Tm| 10 4.34 5 4 6.1655995 6 100 21.71 26 25 30.1261416 41 1000 144.76 171 168 177.6096580 295 10000 1085.74 1292 1229 1246.1372159 2234 100000 8685.89 10102 9592 9629.8090011 12024 1000000 72382.41 82114 78794 78627.5491595 5000000 324150.19 366716 348513 348638.1150413 For comparision: $T_m=\{n\in\mathbb N|n<m\wedge f(a_n,a_{n+1},a_{n+2})\in\mathbb P\}$, where $a_n=2n+1$

论坛问题帖 | has_answers | Lehs | 2017-07-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018769 | MSE-2370039

How can we prove that $F_p \in \mathcal {P} \lor Φ?$

I have created a conjecture which I do not know how to prove. The conjecture is written like: $F_p \in \mathcal {P} \lor Φ$ p = prime number in correspondence to the appearance of a number in the Fibonacci Sequence. $\mathcal {P}$ = The set of prime numbers in the Fibonnaci Sequence. $Φ$ = The set of Fibonacci numbers that have $1+$ divisors which cannot be divided into any earlier Fibonacci numbers. My conjecture, is that if you have a prime-th Fibonacci number, then that number is either going to be prime or have factors of which are not factors of any earlier Fibonacci numbers. Here is a list of the first 10 Fibonacci numbers. 1) 1 2) 1 3) 2 4) 3 5) 5 6) 8 7) 13 8) 21 9) 34 10) 55 As we can see, all prime-th Fibonacci numbers in the list $\in$ $\mathcal {P}$ except for the 2^{nd} Fibonacci number because 1 is not considered a prime, but we will consider it a prime to all numbers. Now as we list the following 20 Fibonacci numbers this time, we will start to notice that not all prime-th Fibonacci numbers are prime. The 19^{th} Fibonacci number is 4181 which is not a prime number. $4181 = 37 \times 113$ $\therefore$ $4181 \in Φ$ $\because$ it's factors $(37, 113)$ are not factors of any Fibonacci number before 19. All prime-th Fibonacci numbers after 4181 are prime, but then when we come across the 31^{st} Fibonacci number being 1346269, which is also not a prime number. $1346269 = 557 \times 2417$ $\therefore$ $134629 \in Φ$ $\because$ it's factors $(557, 2417)$ are not factors of any Fibonacci number before 1346269. The most common factor I assume to be divided into fibonnaci numbers is 2^3 where I assume that $2^3 | F_{6 + 12n} : n \in \mathbb{W}$. Now at first you would believe that $α(F_p \in \mathcal {P}) > α(F_p \in Φ)$: $α(χ) =$ amount of times $χ$ appears. But as $P_y \rightarrow \infty$ the gap between $(F_p \in \mathcal {P})_y \land (F_p \in \mathcal {P})_{y + 1} \rightarrow \infty$ at a much faster rate the greater the value of y : $y \in \mathbb{Z^+}$ $\therefore$ $α(F_p \in \mathcal {P}) < α(F_p \in Φ)$ the greater the value of y, and when $y > 137$ then $α(F_p \in \mathcal {P}) \ll α(F_p \in Φ)$. $φ$ can be expressed as a limit: $$\lim_{k \rightarrow \infty} \frac{F_k + 1}{F_k} = φ : k \in \mathbb{Z^+}$$ $\because$ $F_k = F_{k + 1} - F_{k - 1}$ (you can carry out the mathematics from there). However even if I substitute k as p, and then substitute F_p with their unique prime factorisation $\iff$ $F_p \in Φ$, it does not get me anywhere. Is there a way to prove that this conjecture works either via this approach or another? The largest Fibonnaci prime I know in the sequence so far is $F_{9311}, (9311 = prime)$ so if that is a key fact to finding a proof, please provide it in your answer if you have one. It would be much appreciated. If you do not know what any of the various symbols in this question mean, I will post an edit to this question tomorrow to explain their definition which will be when I see your answers $-$ I have to go to sleep now. G'night.

论坛问题帖 | has_answers | George N. Missailidis | 2017-07-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018770 | MSE-2373253

How is moving the last digit of a number to the front and multiplying related to multiplicative orders?

There seems to be a relationship between multiplicative orders modulo n and a puzzle Presh Talwalkar gave a few days ago at https://www.youtube.com/watch?v=1lHDCAIsyb8 I'm hoping someone can give a more rigorous explanation of the pattern I see. $\bullet\textbf{ The Puzzle }\bullet$ He states the puzzle as: "What Positive Number Doubles When Its Last Digit Becomes Its First Digit?" For example, the smallest solution - assuming base-10 representation - is 105263157894736842. Which means that $$(2)\cdot105263157894736842 \quad\ \ \ $$ $$||$$ $$2105263157894736842$$ Naturally, one can proceede to find solutions in other bases. The base-5 solution is $$(2)\cdot102342_5 \quad\ \ \ $$ $$||$$ $$210234_5$$ $\bullet\textbf{ Multiplicative Order Connection }\bullet$ The base-10 solutions is 18 digits and the base-5 solution is 6 digits. I wrote a program to generate the smallest solution in all bases and then counted the number of digits in each such solution. Here's the sequence, starting with base-2 $$2,4,3,6,10,12,4,8,18,6,11,20,...$$ $$\uparrow \quad\quad\quad\quad\ \uparrow \quad\quad$$ $$\text{Base-} 5 \quad\quad\quad \text{Base-} 10 \quad\quad$$ A quick search on http://oeis.org/ shows that these numbers are the multiplicative order of $$2\ (\text{mod } 2B-1)$$ where B is the representation base we are working in. See http://oeis.org/A002326 This puzzle can be further generalized by finding numbers that triple instead of double upon moving the last digit to the front. Let's introduce an arbitrary multiplying factor: k So the solutions mentioned above and Talwalkar's puzzle are a particular case when $k=2$ For a different example, let's look at $k=5$ in base-7. The smallest solution is $$(5) \cdot 1013042156536245_7 \quad\ \ \ $$ $$||$$ $$5101304215653624_7$$ This solution is 16 digits long. As we did before for base-10, we can write out a sequence of the number of digits in each base's solution starting with base-2 $$6,6,9,2,14,16,4,5,42,18,...$$ $$\uparrow\ \ $$ $$\text{Base-}7\ \ $$ Another search on http://oeis.org/ quickly shows that these numbers are the multiplicative order of $$5\ (\text{mod } 5\cdot7-1)\quad\text{or rather}\quad 5\ (\text{mod }34)$$ $\bullet\textbf{ A Conjecture }\bullet$ The pattern might be clear now. After checking other cases, it seems that the smallest positive number in any base B - which is k times the value gotten by moving it's last digit to the front - has a number of digits equal to the multiplicative order of $$k\ (\text{mod } Bk-1)$$ I can see how multiplicative order would be related to this problem. But I can't find an exact reason why this relationship should be so. Is there an intuitive reason?

论坛问题帖 | has_answers | user377597 | 2017-07-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018771 | MSE-2374156

Does anyone know the name of this conjecture?

Given p and q are two different prime numbers. Does there exist a positive integer n such that $p^n = 1 \pmod q$ Is this conjecture true? If so, any source of the prove. What is the name of this conjecture or theorem (if it is true)?

论坛问题帖 | has_answers | Wan Chap | 2017-07-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018772 | MSE-2375260

If N is a sum of two squares, then so is the odd part of N

Given two natural numbers m,n, where m is odd. Please prove that if $m\cdot2^n$ is a sum of two squares, then m is a sum of two squares. Tested for all $m\cdot2^n<100,000$. I love the natural numbers but feels number theory being inapproachable. So I'm experimenting and mostly (mis)use ME as a "proof calculator". While testing and using my BigZ I analyze sets and sequences of numbers like below. 1 1000 ~ :| n | n sqrsum ; intcond \ create all square sums in intervall function odd-part transform-set \ transform to odd parts zdup cardinality . 153 condition sqrsum filter-set \ only square sums filter zdup cardinality . 153 \ conclude the conjecture in the question condition noprime filter-set \ remove all primes from the set condition nosqr filter-set \ remove all squares set-of-factors \ create the set of all prime factor-tuples cr zet. \ and print it: {(5,197),(3,3,109),(5,193),(13,73),(5,5,37),(3,3,101),(5,181),(17,53),(3,3,97),(5,173),(5,13,13),(7,7,17),(3,3,89),(13,61),(5,157),(3,3,5,17),(5,149),(5,5,29),(17,41),(13,53),(5,137),(3,3,73),(7,7,13),(17,37),(5,11,11),(3,3,5,13),(5,113),(3,3,61),(5,109),(13,41),(5,101),(17,29),(5,97),(13,37),(3,3,53),(5,89),(5,5,17),(3,3,3,3,5),(13,29),(3,3,41),(5,73),(3,3,37),(5,5,13),(5,61),(5,53),(3,3,29),(5,7,7),(13,17),(5,41),(5,37),(3,3,17),(5,29),(5,5,5),(3,3,13),(5,17),(5,13),(3,3,5)} ok From the remaining set the Sum of Two Squares Theorem is almost visible. Also see my blog about using the most low level high level language Forth for mathematical calculations with sets and big integers.

论坛问题帖 | has_answers | Lehs | 2017-07-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018773 | MSE-2379242

A conjecture inspired by a famous contest problem

There is a famous difficult problem: For all natural numbers a,b it's true that: $\displaystyle(ab+1)|(a^2+b^2)\implies \frac{a^2+b^2}{ab+1}$ is a perfect square. I've noticed $(a=1,\dots 100)$ that for each a there's a b such that $(ab+1)|(a^2+b^2)$ and my question is if this can be proved generally? Induction seems to lead nowhere.

论坛问题帖 | has_answers | Lehs | 2017-08-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018774 | MSE-2380758

What must the value of y be so that a and b are integers?

I have made some sort of conjecture that in the equation: $$a^2 = b^2 + 2y^2.$$ y must equal $(2n + 1)\sqrt {8}$ such that $n \in \mathbb{W}$ for a and b to be integers, or at least Pythagorean Triples. All I believe that is most correct is that for every value of n there are at least four solutions to satisfy the equation. And that for some other integer x, if $a^2 = x + y$ and $b^2 = x - y$ then this will work out, but it doesn't seem to be that $x + y$ and $x - y$ can both be squared numbers $\in \mathbb{W}$. Could somebody please help me prove/disprove this? I am a little bit stuck. Thanks.

论坛问题帖 | has_answers | George N. Missailidis | 2017-08-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018775 | MSE-238607

Prime clasfication by some constructive function

How to prove or justify the following: $$ f(g)= \frac1{1-g^2} \prod_{k=1}^{\infty} \left(\frac{\sin(\pi \frac gk)}{\pi \frac gk} \cdot \frac 1{1-\frac{g^2}{k^2}}\right). $$ The above statement can illustrate the following facts: (1) if $f(g) = 0$, then g is composite number (2) If $f(g)$ is not equal to 0, the g is prime number (3) if $f(1+g) = 0$, then g is prime. Please justify.

论坛问题帖 | no_answers_at_retrieval | vmrfdu123456 | 2012-11-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018776 | MSE-2386860

Any odd number is of form $a+b$ where a^2+b^2 is prime

This conjecture is tested for all odd natural numbers less than 10^8: If $n>1$ is an odd natural number, then there are natural numbers a,b such that $n=a+b$ and $a^2+b^2\in\mathbb P$. $\mathbb P$ is the set of prime numbers. I wish help with counterexamples, heuristics or a proof. Addendum: As in Greg Martins answer $$k(n)\approx k'(n)=\frac{3}{4}\frac{n}{\ln n} \displaystyle \prod_{\substack{p\equiv1\pmod 4 \\ p\mid n}} \frac{p-1} {p-2}\prod_{\substack{p\equiv3\pmod 4 \\ p\mid n}} \frac {p-1}p$$ which gives the diagram where k is the blue curve and k' is the red curve.

论坛问题帖 | has_answers | Lehs | 2017-08-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018777 | MSE-2390542

A conjecture about an unlimited path

Conjecture: For all primes of the form a^2+b^2, there are natural numbers s,t,u,v such that $\quad s^2+t^2,u^2+v^2$ are primes $\quad a+b=s+t$ $\quad u+v=s+t+2$ $\quad |s-u|+|t-v|=2$ This conjecture implies Any odd number is of form $a+b$ where a^2+b^2 is prime and seems to be much stronger. I have tested it for $a+b<50.000$ without founding a counterexample, but I guess there may exist counterexamples. In the diagram the vertices correspond to points (x,y) where x^2+y^2 is prime, the red edges correspond to pairs of vertices with taxi cab distance equal to 2 and the gray diagonal lines is the (taxi cab) equidistance lines to origin. The conjecture says there are paths following red edges and gray diagonal lines up to any distance from origin. Addendum: Further tests show that there exists red vertical edges that together with the gray diagonal lines make such paths for all odd $n=a+b=3,\dots,49,999$ except for $n\in\{9,19,49,75,149\}$. In this case I want help with finding counterexamples.

论坛问题帖 | has_answers | Lehs | 2017-08-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018778 | MSE-2391825

A question related to the existence of a prime

Let $I_m=[m!+2,m!+m]\cap\mathbb N$ an "interval" of $\mathbb N$; it can obviously be as long as we want and it is easy to prove I_m does not contain any prime. Prove the following: $$\text { if }n^2+(n+1)^2\in I_m\text{ then }4n^2+1\notin I_M$$ Note that if in a large interval I_m could exist n denying what is proposed here, then we would have found a counterexample to the conjecture in here

论坛问题帖 | has_answers | Ataulfo | 2017-08-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018779 | MSE-2392150

Yet an other conjecture about odd numbers $n=a+b$ such that a^2+b^2 is prime

This question is related to A conjecture about an unlimited path and Any odd number is of form $a+b$ where a^2+b^2 is prime but I present it on its own if anyone would like to help finding counterexamples. Conjecture: For all odd numbers $n>159$ there are positive integers $a\le b$ such that: $\quad n=a+b$ $\quad a^2+b^2$ and $a^2+(b+2)^2$ are primes Let $S_n=\{a\in\mathbb Z^+|\exists b\in\mathbb Z^+:n=a+b\wedge a^2+b^2\in\mathbb P\}$. Below there is a list of n and $S_n\cap S_{n+2}$, the set of the possible $a\,$ that fulfills the conjecture for the odd number n, $1<n<1000$ (0 denotes the empty set): 3 {1} 5 {1,2} 7 {2} 9 0 11 {3,5} 13 {4} 15 {1,2,7} 17 {2,7} 19 0 21 {5} 23 {9} 25 {1,7} 27 {5,10} 29 {5,10} 31 {5} 33 {4,13} 35 {2,6,12} 37 {2,5,10} 39 {5,10} 41 {3,8,13} 43 {8,13} 45 {16,22} 47 {2,15,22} 49 0 51 {8,20,25} 53 {4,8,18,23} 55 {1,17} 57 {7,17} 59 {15} 61 {23} 63 {8,19,23,29} 65 {11,12,16,17,21} 67 {2,20} 69 {10,20} 71 {13,20} 73 {13,14,29,34} 75 0 77 {5,17,25,27,30} 79 {5,25} 81 {8,25} 83 {4,9,14,18,23,24,39} 85 {7,26,32,41} 87 {2,32} 89 {40} 91 {40,43} 93 {4,28,29,34} 95 {11,17,26,37,41} 97 {2,17} 99 {5} 101 {3,5,18,35,38,50} 103 {19,43,44} 105 {26} 107 {17,25,37,40,45,50} 109 {10,35,50} 111 {5,10,20,23,28,35,50} 113 {3,18,28,39,48,54} 115 {47} 117 {2,10,40,47,55} 119 {10,30,40} 121 {13,23,50,58} 123 {4,13,19,38,58} 125 {1,21,27,47} 127 {35,47,50,55} 129 {5,20} 131 {15,45,53,60,65} 133 {43,53} 135 {31,32,37,56} 137 {2,5,10,22,32,42,52,57} 139 {5} 141 {5,25,28,50,58} 143 {3,23,24,34,53,54,64} 145 {11,17,37,41,62} 147 {10,17,20,32,37,50,67} 149 {30} 151 {5,40,53,65} 153 {8,74} 155 {12,17,26,46,52,57,66,71,72} 157 {40,52,67,77} 159 0 161 {10,13,43,50,65,78} 163 {23,28,58,68,74} 165 {31,32,61,76} 167 {12,32,37,50} 169 {5} 171 {10,23,43,65,68,85} 173 {19,23,29,43,48,59,68,73} 175 {16,17,22,82} 177 {7,40,55}

179 {35,40,50} 181 {38,65,83} 183 {8,34,38,49,53,83} 185 {6,21,36,52,56,62,76,82,91}
187 {20,37,40,50,62} 189 {20,50,55,85} 191 {3,18,23,35,68,70} 193
{14,19,23,34,53,58,68,79,94} 195 {46,47,92} 197 {20,27,40,55,60,72,75,92,95} 199
{20,95} 201 {20,25,43,65,73,85} 203 {3,38,53,54,64,93,94} 205 {1,52,86,91} 207
{5,32,82} 209 {5,40,45,75,80} 211 {13,25,53,70,103} 213 {13,44,59,88} 215
{11,37,46,61,72,76,81,102,106} 217 {2,25,55,65,107} 219 {5,10,25,40} 221
{3,20,23,25,33,40,50,63,70,83,98} 223 {4,23,88,98,104} 225
{26,41,46,52,67,86,91,97,107} 227 {7,17,52,55,65,72,77,95,97} 229 {85} 231
{68,83,85,103} 233 {4,14,39,48,59,74} 235 {2,26,31,62,116} 237 {65,70,97} 239
{35,65,70,95} 241 {13,80,88,100,113} 243 {13,58,64,104} 245
{2,11,16,47,51,71,101,122} 247 {7,10,40,47,115,122} 249 {50,115} 251
{8,65,78,85,98,100,123} 253 {59,74,98,113} 255 {2,7,26,52,112} 257
{12,25,52,60,72,85,100,102,125} 259 {85} 261 {23,28,40,103,130} 263
{23,24,34,38,49,63,69,99,109,118} 265 {32,37,46,56,62,71,82,92} 267
{2,20,22,32,55,95,100,107,127} 269 {25,45,50,55,75,100} 271
{8,25,55,68,73,88,100,103,115,128} 273 {8,23,73,83,103,128} 275 {36,42,47,92,111,112}
277 {2,10,20,65,67,97,112,125,127} 279 {20,25,55} 281
{18,25,28,38,55,68,73,78,83,113,118,135} 283 {29,64,68,73,79,89,118} 285 {22,86,116}
287 {5,45,47,57,62,67,127,130} 289 {40,130} 291 {40,83,98,115,118,128} 293
{4,13,19,24,29,39,98,109,128} 295 {61,67,106,136,142} 297 {10,32,47,62,67,112,145}
299 {5,120,145} 301 {23,65,95,125,130,143,145} 303 {4,34,58,89,104,113,119} 305
{17,21,32,46,71,81,82,116,117,132,136} 307 {25,40,47,107,122} 309 {20,25,40,100,125}
311 {8,20,30,40,70,90,98,110,113,125,138} 313 {23,38,44,88,104,118} 315
{37,71,76,101,142} 317 {5,10,17,30,97,102,135,137,152} 319 {5,10,50,85} 321
{5,50,58,70,83,115,128,140,155} 323 {23,44,49,58,63,74,88,118,129,138,148,159} 325
{37,76,92,106,122,151} 327 {20,40,50,67,82,92,115,137} 329 {55,75,125,130,145} 331
{28,38,55,58,85,113,125,130,133,145} 333 {14,29,38,83,113,139} 335
{11,12,22,26,31,37,42,66,77,102,111,116,126,142,156,161} 337
{7,32,77,85,92,100,152,155,160} 339 {5,70,145} 341 {5,10,23,60,73,83} 343
{19,58,83,124,134,139,143} 345 {61,67,97,101,121,146,152,172} 347
{5,25,42,45,75,87,90,102,120,132,172} 349 {10,40,110} 351 {80,88,118,133,148,158}
353 {8,13,59,99,119,133,158,164} 355 {26,67,76,127,137,146,166,172} 357
{20,25,55,67,137,172} 359 {20,55,60,85,100,105,170} 361 {5,53,85} 363 {14,53} 365
{27,52,86,91,106,132,151,166} 367 {10,17,37,40,92,112,125,157,170,175} 369
{20,85,110} 371 {5,15,43,45,55,65,83,88,93,113,145,150,160,165} 373
{13,49,68,83,94,133,139,169,179} 375 {2,31,32,37,47,61,76,86,107,142,151,166,187} 377
{7,10,20,25,27,30,60,90,92,102,142,155,160,185} 379 {80,95,130,160} 381
{73,95,130,140,143,160,178} 383 {3,8,73,74,78,113,123,124,169} 385 {1,122,146,151}
387 {5,22,47,70,112,155} 389 {5,25,65,70} 391 {13,43,53,70,103,130,140,148} 393
{4,13,43,49,53,58,64,103,134,148,188} 395 {37,56,71,87,97,126,131,137,166,167} 397
{97,160,197} 399 {80} 401 {5,35,38,58,63,68,73,80,105,123,145,153,175,188,193} 403
{14,29,38,43,44,73,109,128,164,179,193} 405 {17,41,62,82,131,151,172} 407
{15,17,35,42,65,80,97,107,117,152,162,175} 409 {35,65,95,110} 411
{8,13,20,40,53,65,85,113,145,200} 413
{3,8,13,24,39,53,64,109,124,163,164,183,194,198,204} 415 {47,61,137,142,161} 417
{10,47,52,67,110,140,142,170,187} 419 {95,105,110,140,180} 421
{5,35,85,100,110,115,118,128,140,163} 423 {13,43,44,59,74,83,128,133,163,179} 425
{2,11,16,32,52,106,117,127,156,162,191,207} 427 {2,25,32,50,67,92,115,145,152,185}
429 {5,50,70,115,160} 431 {28,43,63,68,78,95,118,140,143,160,183,193,208} 433
{34,68,89,94,134,169,179,184,188,193,208} 435 {22,32,52,107,136,151,191} 437
{20,25,70,85,90,117,155,157,165,195} 439 {215} 441 {38,118,220} 443
{13,18,19,78,88,173,179,184} 445 {17,26,31,71,91,92,97,112,116,121,131,142,182,212}
447 {5,7,92,95,115,127,130,200,212,215,217} 449 {5,20,30,85,160,175,200,210,215} 451
{23,25,35,58,100,103,178,193,200} 453 {38,44,73,79,103,149,178} 455
{2,11,36,51,62,81,116,146,157,186,192} 457 {7,10,62,107,130,137,157,160,202,205,215}
459 {35,80,130,160,190} 461
{35,43,55,68,80,93,100,110,120,138,168,178,188,190,208,210,225,228} 463
{73,74,109,118,139,143,169,208} 465 {1,11,46,47,52,112,116,131,142,157,196,206,221}
467 {12,60,65,97,117,137,157,160,177} 469 {40,145,160} 471
{8,38,53,70,95,128,148,190,235} 473 {48,49,59,63,104,113,124,148,174,208} 475
{17,31,32,46,67,122,136,137,167,206,221} 477 {2,22,65,155,172,185,187,205,217,230}
479 {90,100,165,210,230,235} 481 {5,10,50,73,158,193,223,233} 483
{8,13,34,68,79,158,163,164,223} 485
{11,32,41,72,86,107,117,126,136,146,152,171,186,227,231,237} 487
{25,35,47,82,110,137,185} 489 {25,145,230} 491
{8,25,48,60,70,73,90,100,120,130,135,138,163,180,198,230} 493 {8,19,59,109,194,199}
495 {92,101,106,136,157,236} 497
{12,30,50,75,102,122,135,157,160,162,190,205,207,222} 499 {85,115,205,230} 501
{28,133,163,185,193,205,215} 503
{23,58,74,103,109,139,154,168,174,184,193,213,218,233} 505
{31,76,122,131,146,187,191,197,211} 507
{5,37,50,70,77,107,122,160,170,185,212,215,232,250} 509 {15,145,150,250} 511
{85,100,125,163,178,205,230} 513 {13,163,178,194,199,214,224,233} 515
{16,41,61,81,107,116,131,142,152,161,182,217,226,241} 517
{50,107,142,152,155,172,232} 519 {5,40,190,215} 521
{13,65,93,105,118,135,148,158,190,218,225,238,255,260} 523
{13,89,103,118,164,218,239} 525 {26,61,82,101,127,166,226,227,251} 527
{10,32,37,72,75,80,97,122,140,145,152,162,165,197,210,212,227,240,247} 529
{35,80,145,220} 531 {35,38,50,58,158,188,190,200,220,223,230,233,250} 533
{24,29,63,94,98,109,119,163,184,213,218,223,249,254,259,264} 535
{17,47,91,211,232,242,247} 537 {10,52,80,130,160,167,185,197,215,227,232,247} 539
{20,90,130,145,155,160,215} 541 {5,8,20,28,100,130,133,145,148,155,158,170,235} 543
{8,13,14,29,133,139} 545 {6,12,17,31,52,56,86,87,112,117,137,162,182,216,241,256,267}
547 {47,62,65,95,142,170,187,197,247,260,265} 549 {25,50,65,70,115,125,130,160,170}
551 {33,45,78,88,113,115,123,135,183,205,225,240} 553 {23,68,89,139,173,193,208,244}
555 {67,101,136,166,187,197,272} 557
{5,45,62,70,87,90,100,107,115,152,157,167,220,235,270,272} 559
{25,50,70,115,185,205} 561 {53,115,118,125,173,185,190,193,203,205,215} 563
{29,34,43,58,59,63,69,74,93,98,104,153,154,173,193,204,219,228,238,258,263,268} 565
{22,52,121,137,256,272,277,281} 567 {67,130,142,152,235} 569 {50,100,245,255,265}
571 {38,95,125,160,173,233,253,260,265} 573
{8,14,28,109,148,173,179,208,229,233,238,244} 575

{17,32,157,182,186,196,206,212,232,257,271} 577
{17,22,32,35,115,137,157,175,187,212,247,257,260,280} 579 {100,260} 581
{10,48,90,93,130,148,153,178,180,193,200,233,243,260} 583
{8,23,28,113,118,148,149,158,193,214,224,268,269} 585
{11,67,122,157,166,172,181,197,212,226,241,242} 587
{10,17,20,50,55,85,130,140,177,195,212,222,232,242,245,275} 589 {10,20,35,250,275}
591 {35,38,73,100,103,110,145,158,223,263} 593
{73,74,94,124,163,169,174,183,194,209,234,243,249,254,263} 595
{47,86,131,142,151,157,172,232,271} 597 {47,92,122,157,247,280,290} 599
{30,65,205,240,285} 601 {10,40,103,125,128,143,148,155,188,208,233,245,283} 603
{43,49,74,79,113,169,188,199,284} 605 {7,26,57,62,82,96,117,192,201,227,241,267,276}
607 {2,10,62,167,172,197,227,230,275,292,295} 609 {25,110,170,220} 611
{15,25,30,33,38,68,70,73,80,83,88,133,138,145,153,170,173,218,223,298} 613
{34,59,88,128,133,218,223,224,248,263} 615
{2,11,56,71,107,167,176,182,202,232,241,266,292,296} 617
{12,17,45,85,95,142,147,165,172,267,282,285,292,305} 619 {35,70,130,215,265} 621
{8,13,25,130,220,293,295} 623
{8,13,34,59,64,69,99,123,129,148,183,188,199,243,253,293,299} 625
{26,106,127,181,226,236,251,257,272,287} 627
{47,50,82,125,127,155,167,200,212,235,245,260} 629 {10,20,50,115,165,245,270,290}
631 {23,35,50,58,110,115,128,145,178,193,238,250,275,278} 633
{103,104,128,143,154,179,194,218,224,278} 635
{1,16,81,92,152,157,167,191,197,232,261,282,291} 637
{80,115,167,185,215,242,262,265,317} 639 {10,170,185,245,250,275} 641
{8,20,55,73,78,100,128,138,158,165,185,225,228,240,258,275,280,318} 643
{38,73,158,173,194,203,233,289} 645 {1,16,67,107,122,151,187,196,202,236,281,287}
647 {25,27,42,87,125,195,207,230,235,265,312} 649 {125,190,200,235} 651
{73,95,110,118,173,200,218,220,235,275} 653
{13,18,23,44,73,103,138,144,173,174,203,229,244,248,279,294} 655
{7,11,52,71,76,136,137,166,176,187,227,256,271} 657
{52,67,82,107,130,137,175,197,205,280,310,322} 659
{55,130,145,155,175,195,270,285,310,315} 661
{28,53,55,70,118,125,133,140,145,155,298,310} 663
{53,64,79,118,164,179,268,278,298,313} 665 {2,17,26,41,47,66,106,117,121,131,157,166,
177,212,216,222,241,257,262,267,271,297,326} 667
{2,17,52,82,172,205,217,257,262,277,292} 669 {125,160,215,230,320} 671
{48,60,118,120,135,185,215,233,235,248,313} 673 {29,34,128,133,143,193,328} 675
{17,22,46,56,77,82,142,151,157,182,332} 677
{17,62,67,75,117,127,130,155,157,247,260,262,272,300,305,317,325,335} 679 {65,260}
681 {20,65,98,110,163,170,230,253,308} 683
{3,9,19,24,39,43,54,83,139,148,163,213,214,219,224,264,303,308,329} 685
{31,37,41,61,71,76,107,127,136,142,161,172,187,197,226,232,257,266,271,272,337} 687
{112,127,187,200,242,272,307,322} 689 {5,15,20,55,75,90,105,200,225,255,290,340} 691
{20,85,208,218,268,290,293,310,323,340} 693
{4,13,59,128,149,158,188,218,229,283,299,323,334} 695
{6,42,47,61,81,117,121,132,162,171,191,251,262,276,302,331} 697
{20,40,80,100,130,145,182,302,307,325} 699 {40,310,325} 701
{40,108,125,135,160,170,173,183,188,203,240,255,273,293,325} 703
{4,14,28,53,109,169,184,203,223,239,283} 705
{26,97,151,166,167,181,187,251,316,317,346} 707
{10,32,52,65,100,160,180,205,222,270,300} 709 {10,55,85,100,160,185,295,335} 711
{50,58,88,110,113,185,233} 713
{19,48,53,73,103,158,168,224,233,239,243,284,304,309,314,353,354} 715
{1,112,161,212,277,281,302,307,311} 717
{7,40,202,212,232,245,250,262,277,302,320,332,355} 719 {115,180,250,295,320} 721
{8,25,85,160,188,220,295,305,358} 723
{13,34,79,124,158,163,169,208,259,268,328,334,338,343,358} 725
{16,22,107,122,131,137,156,161,207,237,246,252,267,282,286,287,302,341,351} 727
{5,22,82,212,220,230,235,265,280,325} 729 {125,140,175,220,235,290,325} 731 {18,40,5
3,63,125,140,148,173,193,200,203,208,220,240,245,248,270,273,298,318,338,350,355}
733 {19,83,88,89,134,169,173,193,254,274,319,338,349} 735
{17,52,71,101,122,221,251,296} 737 {12,35,65,80,115,135,140,142,197,295,305,315,350}
739 {35,40,145,230,305,320} 741
{20,73,83,145,158,160,163,215,218,248,265,283,290,293,313,358} 743
{4,28,39,49,79,99,114,119,138,163,193,219,263,264,284,289,313,323,329,364} 745
{2,11,32,37,107,121,152,167,182,236,247,257,286,331,347} 747
{20,52,110,152,242,275,277,305,340,370} 749
{25,80,125,170,195,230,265,340,345,360,365} 751
{13,25,70,73,115,125,130,133,143,163,170,233,278,328,338,365} 753
{13,73,74,143,149,173,214,233,254,274,278,343,359,364} 755
{2,52,57,106,132,147,181,187,242,307,316,322,326,327,356,361} 757
{10,52,62,112,200,265,355} 759 {50,100,140,265,355,365} 761 {13,45,65,85,100,120,145
,153,158,163,178,180,198,230,260,265,270,283,338,340,345,355} 763
{38,58,158,184,188,214,278,283,298,304,353} 765
{47,71,92,101,112,161,172,211,256,257,266,296,307} 767
{17,47,72,75,85,105,115,142,190,220,285,307,310,352,355,365,375,382} 769
{55,85,115,145,190,235,260,310} 771 {85,115,145,193,218,223,250,323,325,340} 773
{4,9,38,48,74,94,108,149,163,233,259,299,333} 775
{22,52,107,137,142,146,181,187,191,256,272,281,332,337,386} 777
{65,107,122,187,215,230,260,317,337} 779 {25,100,135,170,230,255,280,305,345} 781
{5,8,73,83,188,200,223,235,268,305,325,383} 783
{8,19,83,149,188,209,214,229,253,278,329,383} 785 {32,37,56,61,66,96,97,117,121,166,1
67,182,212,226,241,261,266,277,306,312,336,351,352,371} 787
{2,70,85,95,110,167,187,212,250,262,292,380} 789 {55,380} 791
{8,15,18,45,68,73,100,115,220,243,253,285,288,295,300,348,353,360,373} 793
{14,98,133,154,164,199,209,218,229,233,254,259,289,304,314,353,379,388} 795
{16,47,52,56,86,101,157,196,206,232,307,316,332,341,361,391} 797
{30,42,82,87,117,145,147,150,175,205,212,240,270,310,312,322,362,367} 799
{25,145,215,295,350} 801 {8,25,70,85,100,148,185,203,223,263,280,295,298,340} 803
{3,39,74,148,204,258,269,298,328,369,398} 805 {137,191,242,251,286,397} 807
{10,47,65,67,125,130,182,220,262,265,335,392,395} 809
{10,125,130,190,300,320,330,375,395} 811

{10,43,113,128,130,178,208,238,325,383,395,400} 813
{29,43,68,113,119,128,323,373,379} 815
{32,97,101,106,116,276,286,311,331,336,357,377,406} 817
{32,97,142,170,320,352,365,367} 819 {25,125,145,170,185,215,320,340} 821
{95,98,125,150,170,203,250,263,268,288,300,323,340,345,368,388,390,410} 823
{4,8,34,188,203,223,229,263,268,299,323,334,364,368,373} 825
{17,71,86,91,142,151,197,211,241,377,406} 827
{22,80,85,120,122,152,157,165,187,197,207,215,217,220,237,255,275,287,317,320,377} 829 {130,170,215,310} 831 {58,130,143,155,163,215,248,265,268,295,313,365} 833
{38,54,58,123,148,173,174,183,234,248,279,298,353,359,369,389,394,398,409} 835
{52,76,212,221,271,272,326,376,382,407} 837
{5,52,82,160,175,182,187,200,230,332,347,407} 839
{10,35,55,70,115,160,165,200,225,295,405} 841
{10,35,38,68,103,133,193,253,295,323,383,388} 843
{14,29,73,103,133,148,179,224,233,244,259,284} 845
{27,37,51,72,76,92,107,116,172,212,226,241,246,277,317,327,332,337,362,367,376} 847
{2,20,50,127,247,310,320,332,355,410} 849 {35,50,80,260,290,310,340,410} 851
{13,25,30,35,48,98,103,110,120,168,175,190,240,260,275,290,308,350,363,410,418,420} 853 {28,34,53,74,119,139,164,248,268,274,308,313,403} 855
{37,61,112,116,146,167,176,182,232,326,367,371,386,401,427} 857
{47,85,87,105,110,112,122,135,160,177,305,342,345,360,370,382,395} 859
{25,110,170,260,265,290,355,370,395,415} 861
{25,38,83,170,200,208,218,235,248,335,355,395,403,428} 863
{4,43,59,83,149,154,178,213,219,248,254,293,334,349,374,384,429} 865
{22,41,62,91,121,172,226,232,356,392,397,412,431} 867
{10,25,32,52,67,70,97,142,155,185,197,217,232,295,347,370,377,380,412} 869
{80,105,120,185,200,380,425} 871
{23,73,80,128,133,160,233,245,275,290,293,298,313,380,395,400,428} 873
{128,164,184,244,289,293,349,359,368,418} 875
{97,101,146,151,166,167,177,191,197,216,256,267,282,327,332,386,397,407} 877
{22,32,65,95,97,112,145,227,230,275,277,332,347,355,385} 879
{55,130,145,230,265,275,280,350,385,395,430} 881
{33,53,83,118,130,168,178,213,230,263,280,298,310,313,393,395,398,403,415,430,440} 883 {4,14,103,104,169,203,254,259,298,403,428,434} 885
{31,56,91,122,166,281,292,382,416} 887
{45,62,65,82,102,110,115,117,122,155,160,192,235,247,285,365,387,430} 889
{115,335,365} 891 {70,115,130,175,205,260,340,343,358,368} 893 {18,23,44,93,103,123,143,158,168,174,198,214,219,224,243,244,249,263,273,343,348,373,414,419} 895
{31,37,71,86,106,116,131,146,166,236,317,361,386,392,401,406,421} 897
{25,77,137,172,220,257,280,337,365,407,410} 899
{10,15,55,125,150,185,195,215,260,300,320,365} 901
{10,88,110,125,185,188,193,205,325,338,358,383,400,433} 903
{19,59,104,143,178,253,254,319,338,359,389} 905 {2,7,26,47,57,67,116,157,171,176,177,222,241,247,276,291,311,326,392,412,432,441,442,447,451} 907
{112,157,167,187,247,272,295,320,332,347,350,395} 909 {160,320,370} 911
{18,58,73,75,90,130,140,148,158,160,190,205,238,255,278,318,335,338,355,378} 913
{23,148,233,283,338,413,454} 915
{11,26,92,97,121,127,206,232,277,326,331,382,412,422} 917
{5,32,82,97,202,225,247,277,302,377,382,395,410,425,445} 919
{80,155,250,295,340,370,400,425,445} 921
{68,113,133,155,223,233,248,250,283,295,328,340,398,425,440} 923
{14,44,108,124,159,164,183,199,218,223,224,229,258,283,309,328,339,369,393,454} 925
{86,91,92,127,137,232,271,326,431,452} 927
{55,127,152,160,232,242,290,295,305,440,452} 929 {25,30,55,65,165,215,345,370,445} 931 {53,55,83,158,220,265,293,298,320} 933
{53,79,113,164,193,203,259,268,269,293,299,304,358,359,389,434,464} 935
{26,27,91,111,112,152,161,202,212,217,307,316,332,347,372,381,397,441,467} 937
{17,77,142,145,170,182,200,212,242,322,325,355,455} 939
{5,40,125,140,170,220,290,325,395,455,460} 941
{13,153,163,240,305,375,383,388,395,420,455,468} 943
{13,109,113,143,148,169,188,323,353,388,418} 945
{32,41,62,136,232,286,296,317,367,401,412} 947
{32,42,62,90,105,175,190,197,240,242,255,282,297,327,357,372,385,402,405,445} 949
{80,85,340,385,415,445} 951
{20,43,65,85,88,140,185,250,290,325,328,353,358,403,440,443,475} 953 {29,43,49,88,99,133,134,149,159,188,199,229,234,268,273,284,294,308,309,318,323,328,333,353,379,384,418,464} 955 {17,32,47,56,82,107,136,146,202,211,241,247,281,397,431,442,467} 957
{5,40,92,107,122,125,265,340,397,425,430,442} 959 {125,265,365,445} 961
{13,35,85,98,103,118,173,253,260,278,290,365,368,433,448} 963
{13,49,64,73,118,119,134,169,173,208,268,283,368,374,403,424,433,439} 965
{21,31,86,102,136,186,222,247,271,317,357,372,401,447,452,461} 967
{10,40,200,215,230,317,362,382,400,422,452} 969 {55,145,200,215,230,295,370,400} 971 {18,23,55,80,88,110,123,143,193,200,223,230,233,253,295,335,338,360,370,373,400,408,425,438,460,465} 973
{58,74,103,148,149,223,239,298,379,383,388,404,424,428,479} 975
{62,131,176,187,196,206,271,281,301,316,322,332,346,427,472} 977 {7,10,17,30,62,95,112,120,160,210,222,237,270,310,322,332,387,390,417,432,437,442,472} 979
{125,160,205,260,310,355,400} 981
{25,28,58,68,110,133,155,178,208,275,298,328,355,368,380,433} 983 {28,29,73,178,208,228,229,233,243,254,304,308,313,328,378,389,398,399,424,429,444,449} 985
{22,26,31,151,172,202,352,377,412,421} 987 {20,40,52,95,142,265,352,377,397} 989
{10,65,125,160,255,265,270,340,365,415,450,485} 991
{10,65,110,115,143,173,278,323,368,370,428,473,478} 993
{19,104,109,124,143,173,299,308,404,443,449,478} 995 {6,7,61,76,101,107,117,127,146,152,192,217,221,241,256,266,296,327,341,382,387,406,416,442,446,466,472,496} 997
{97,107,127,130,152,217,232,310,337,367,395,422,425,442,460} 999
{230,395,400,430,470} So far tested for $n<50,000$.

论坛问题帖 | no_answers_at_retrieval | Lehs | 2017-08-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018780 | MSE-2397926

Hard for-all-integer-problems proveable for very great integer limits

Computational experiments suggests the conjecture: All big enough odd numbers N is of the form $N=\sum_{k=1}^n m_k$, where $\sum_{k=1}^n m_k^2$ is prime and all $m_k\in\mathbb Z^+$. Since the possible combinations of partitions of N increase with increasing N and n, the case with $n=2$ is tested for $N<10^8$ and seems to be supported by heuristics, and the cases $n=3,4,5$ are tested for N up to 200, I'm interested to understand the following, which is my question: Are there reasons to believe that a proof for some "big" n with N "big enough" would be within reach while proofs for "small" n isn't, at the moment?

论坛问题帖 | no_answers_at_retrieval | Lehs | 2017-08-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018781 | MSE-2398642

Conjectures (or intuitions) that turned out wrong in an interesting or useful way

The question What seemingly innocuous results in mathematics require advanced proofs? prompts me to ask about conjectures or, less formally, beliefs or intuitions, that turned out wrong in interesting or useful ways. I have several in mind, but will provide just one here now, as an example. For centuries mathematicians tried to show that Euclid's parallel postulate followed from his others. When Lobachevsky, Bolyai and Gauss discovered that you could do interesting geometry just as consistent as Euclid when the parallel postulate failed a whole new world was open for exploration. One example per answer, please. If you want to post several, answer repeatedly. Related: Conjectures that have been disproved with extremely large counterexamples?

论坛问题帖 | has_answers | Ethan Bolker | 2017-08-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018782 | MSE-2398917

Any integer is of the form $n=a_1^2+a_2^2+a_3^2-a_4^2-a_5^2$

I've struggled with this conjecture, which probably can be proved: Any natural number n can be written as $n=a_1^2+a_2^2+a_3^2-a_4^2-a_5^2$ for some $a_1,a_2,a_3,a_4,a_5\in\mathbb Z^+$. I guess there are some polynomials $a_k=p_k(n)$ that solves the problem by substitution. But which? I'm almost sure that ME can prove the conjecture.

论坛问题帖 | has_answers | Lehs | 2017-08-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018783 | MSE-2399035

Most even numbers is a sum $a+b+c+d$ where $a^2+b^2+c^2=d^2$

I have no idea how hard this conjecture is to prove: Any even number $n\ge 36$ can be written as $n=a+b+c+d$ where $a^2+b^2+c^2=d^2$ and $a,b,c,d\in\mathbb Z^+$. Small exceptions are $n=2, 4, 6, 10, 12, 14, 20, 26, 34.\,$ Tested for $n\le 10,000$. A counter-example would be as interesting as a proof. n as above must be even $d-c,c+d|a^2+b^2$

论坛问题帖 | has_answers | Lehs | 2017-08-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018784 | MSE-2402029

$\forall n \in \mathbb{N}_{>5}\implies\exists (a,b)\in \mathbb Z^+:a^2+b^2\notin\mathbb P\;\wedge\;n=a+b$

Conjecture: $\forall n \in \mathbb{N}_{>5} \implies\exists (a,b)\in \mathbb Z^+:a^2+b^2\notin\mathbb P\;\wedge\;n=a+b$ Tested $\forall n\leq 100,000$. Small exceptions: {1,2,3,5}. I would like to see proofs or counterexamples. The conjecture seems to be related to: Most even numbers is a sum $a+b+c+d$ where $a^2+b^2+c^2=d^2$ Any odd number is of form $a+b$ where a^2+b^2 is prime

论坛问题帖 | has_answers | Lehs | 2017-08-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018785 | MSE-2406659

A characteristic property of soluble groups

In the paper A characteristic property of soluble groups, P. Hall proved that in a finite soluble group, if one has elements $a,b,c,\ldots$ of coprime orders, and you suppose that $abc\ldots=1$, then actually $$a=b=c=\ldots=1$$ He conjecture that this statement is actually equivalent to the property of being soluble. Is this result already been established? If so, where I can find it?

论坛问题帖 | no_answers_at_retrieval | W4cc0 | 2017-08-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018786 | MSE-2406937

If $q^k n^2$ is an odd perfect number with Euler prime q, does $\sigma(n^2)/q^k < q$ hold?

If $q^k n^2$ is an odd perfect number with Euler prime q, then it is known that $$\frac{\sigma(n^2)}{q^k}=\frac{2n^2}{\sigma(q^k)}=\frac{D(n^2)}{\sigma(q^{k-1})}=\gcd\left(n^2,\sigma(n^2)\right),$$ where $\sigma(x)$ is the sum of divisors of $x \in \mathbb{N}$ and $D(x)=2x-\sigma(x)$. (See this paper for a proof.) Here is my question: Does $\sigma(n^2)/q^k < q$ hold? MY ATTEMPT It is easy to prove that $\sigma(n^2)/q^k \neq q$. For suppose to the contrary that $$q=\frac{\sigma(n^2)}{q^k}=\frac{2n^2}{\sigma(q^k)}.$$ This means that $$2n^2 = q\sigma(q^k)$$ which implies that $q \mid n^2$, contradicting $\gcd(q,n)=1$. So now, assume that $\sigma(n^2)/q^k > q$. We obtain $$\frac{\sigma(n^2)}{q^k}=\frac{2n^2}{\sigma(q^k)}=\frac{D(n^2)}{\sigma(q^{k-1})}=\gcd\left(n^2,\sigma(n^2)\right)>q,$$ from which we get $$2n^2 > \sigma(n^2) > q^{k+1}$$ $$2n^2 > q\sigma(q^k)$$ $$2n^2 - \sigma(n^2) > q\sigma(q^{k-1}) \geq q$$ $$\gcd(n^2, \sigma(n^2)) > q,$$ whence I do not arrive at a contradiction.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2017-08-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018787 | MSE-2408529

conjectured relations of mock theta functions of order 3

Given the following mock theta functions of order 3,found in this wikipedia article $f(q)=\sum_{n=0}^{\infty} \frac{q^{n^2}}{(-q;q)^2_{n}}$,$\phi(q)=\sum_{n=0}^{\infty} \frac{q^{n^2}}{(-q^2;q^2)_{n}}$,$\omega(q)=\sum_{n=0}^{\infty} \frac{q^{2n(n+1)}}{(q;q^2)^2_{n+1}}$ and $\nu(q)=\sum_{n=0}^{\infty} \frac{q^{n(n+1)}}{(-q;q^2)_{n+1}}$ where $q=e^{2\pi i\tau}$,$|q|\lt1$ and $(a;q)_n=\prod_{j=0}^{n-1}(1-aq^j)$ I propose the conjecture that the following relations hold $\Big(\phi(q)-1\Big)^2+\Big(f(q^2)-1\Big)=2\sum_{k=1}^{\infty}\Big(\sum_{n=1}^{\infty} \frac{q^{(n-1)^2+2k(n-1)+2k^2}}{(-q^2;q^2)_{k} (-q^2;q^2)_{n+k-1}}\Big)\tag1$ and $\Big(\nu(q)-\frac{1}{(1+q)}\Big)^2+\Big(\omega(-q)-\frac{1}{(1+q)^2}\Big)=2\sum_{k=1}^{\infty}\Big(\sum_{n=1}^{\infty} \frac{q^{n^2+(2k-1)n+2k^2}}{(-q;q^2)_{k+1} (-q;q^2)_{n+k}}\Big)\tag2$ How do we prove the conjecture? Note:comparing coefficients on the RHS and LHS of (1) and (2) respectively,we observe that the coefficients are equal and the identities are both special cases of the more general identity $\Big(\sum_{n=1}^{\infty} \frac{x^n z^{n^2}}{(a;q)_{n}}\Big)^2+\sum_{n=1}^{\infty} \Big(\frac{x^n z^{n^2}}{(a;q)_{n}}\Big)^2=2\sum_{k=1}^{\infty}\Big(\sum_{n=1}^{\infty} \frac{x^{n+2k-1} z^{(n-1)^2+2k(n-1)+2k^2}}{(a;q)_{k} (a;q)_{n+k-1}}\Big)\tag3$ with complex number x and z,$z\lt1$whereby a transformation($(a;q)_n\rightarrow (a;q)_{n+c}$ or $(a;q)_n\rightarrow (a;q)_{cn}$) on the LHS results in an equivalent transformation($(a;q)_k\rightarrow (a;q)_{k+c}$ or $(a;q)_k\rightarrow (a;q)_{ck}$) on the RHS for natural number c

论坛问题帖 | has_answers | Nicco | 2017-08-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018788 | MSE-2410720

Even numbers has the form $a+b$ where $\frac{a^2+b^2}{2}$ is prime

Conjecture: All even numbers $n>2$ can be written $n=a+b$ where $a,b\in\mathbb N^+$ and $\frac{a^2+b^2}{2}\in\mathbb P$. Is tested for $n<10,000,000$. This conjecture is related to and maybe dependent on Any odd number is of form $a+b$ where a^2+b^2 is prime.

论坛问题帖 | no_answers_at_retrieval | Lehs | 2017-08-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018789 | MSE-2411538

$\{a+b|a,b\in\mathbb N^+\wedge ma^2+nb^2\in\mathbb P\}=\{k>1|\gcd(k,m+n)=1\}$

Conjecture: $\{a+b|a,b\in\mathbb N^+\wedge ma^2+nb^2\in\mathbb P^{>2}\}=\{k>2|\gcd(k,m+n)=1\}$ if $m,n\in \mathbb N^+$ and $\gcd(m,n)=1$. This is a generalization of Any odd number is of form $a+b$ where a^2+b^2 is prime. Perhaps the generalization will spread some light of what is going on? There is a perfect match of the formula for all tests I've done. https://mathoverflow.net/questions/280123/the-set-of-numbers-ab-such-that-ma2nb2-is-prime

论坛问题帖 | has_answers | Lehs | 2017-08-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018790 | MSE-2413425

When does the Ext-quiver of a finite-dimensional algebra have no oriented cycles, or what structure does it have?

Let H be a finite-dimensional algebra and $S(1), \dots, S(n)$ be the simple modules of H. Let Q be the Ext-quiver of H, i.e. Q has as vertices the simple modules $S(1), \dots, S(n)$ and an arrow $S(i) \to S(j)$ provided Ext$^1_H(S(i), S(j)) \neq 0$. Question: What is known or conjectured about the question when there cannot be an oriented cycle in the Ext-quiver? The no loop conjecture (which is, as far as I know, only proven for algebraically closed fields, see for example A proof of the strong no loop conjecture by Igusa, Liu and Paquette) says that in finite global dimension there cannot be a loop in the Ext-quiver. Is there an analogous statement or conjecture for oriented cycles? I'm also interested in references, if you know any. Added later: As pointed out by Stephen, there cannot be a general analogous statement. Nevertheless, I'm additionally interested in anything you know about the structure of the Ext-quiver whenever H has finite global dimension, whether it is related to cycles or not.

论坛问题帖 | has_answers | Leon Lang | 2017-09-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018791 | MSE-2414536

Two different polynomial expressions for certain primes

Here is an other random conjecture which I have no clue how to prove: $a,b\in\mathbb N^+\wedge a^2+b^2+ab\in\mathbb P\implies\exists$ $A,B\in\mathbb N^+:A^2+B^2-AB=a^2+b^2+ab$. Tested for $a,b<20,000$ on my 32 bit tabloid. I would like to see a proof or an counter-example.

论坛问题帖 | has_answers | Lehs | 2017-09-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018792 | MSE-2415330

The set of primes of the form a^2+8b^2, where $a,b\in\mathbb N_+$

Due to the Brahmagupta identity the product of numbers of the form a^2+nb^2, with a fixed n, is on the same form. For $a,b,n\in\mathbb N_+$, these semigroups B_n are generated bye sets of primes G_n of the same form. It's well known that $G_1=\mathbb P\setminus(4\mathbb N+3)$ and easy to see that $G_4=\mathbb P\cap(4\mathbb N+1)$. My computations suggest that $G_8=\mathbb P\cap(8\mathbb N+1)$. I would like a proof or a counterexample? Sorry about the errors in the first editions. It's OK to redraw the votes.

论坛问题帖 | has_answers | Lehs | 2017-09-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018793 | MSE-2417479

Two dissimilar continued fractions that are equivalent $F(q)=G(q)$

Given the following two continued fractions,with $|q|\lt1$ $F(q)= \cfrac{1}{1+\cfrac{q}{1+\cfrac{q^2}{1+\cfrac{q^5}{1+\cfrac{q^8}{1+\cfrac{q^{12}}{1+\cfrac{q^{16}}{1+\cfrac{q^{24}}{1+\dots}}}}}}}}\tag1$ where the exponents are given by the Generating function $f(z)=\frac{-2z^4-4z^3-3z^2-2z-1}{2z^2-1}$ and $G(q)= 1-\cfrac{q}{1+\cfrac{q}{1-\cfrac{q}{1+\cfrac{q}{1-\cfrac{q^4}{1+\cfrac{q^{4}}{1-\cfrac{q^{4}}{1+\cfrac{q^{4}}{1-\dots}}}}}}}}\tag2$ in which case the exponents are given by the generating function $g(z)=\frac{-2z^7-2z^6-2z^5-2z^4-z^3-z^2-z-1}{2z^4-1}$ and satisfy the recurrence relation $a(n+4)=2a(n)$ for $n\geq4$ How do we prove that $F(q)=G(q)$? Note:by comparing coefficients of each continued fraction,we observe that they are equal. Curiously though,by plotting each generating function on the complex plane yields beautiful images(f(z) and g(z)) of bugs

论坛问题帖 | has_answers | Nicco | 2017-09-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018794 | MSE-2422504

Is there a deficient N such that $\sqrt{D(N)} \mid A(N)$ holds, where $D(N)$ is the deficiency and $A(N)$ is the sum of the aliquot parts of N?

Do there exist deficient numbers N such that $$\sqrt{D(N)} \mid A(N)$$ holds, where $$D(N) = 2N - \sigma(N)$$ is the deficiency and $$A(N) = \sigma(N) - N$$ is the sum of aliquot parts of N?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2017-09-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018795 | MSE-2423712

If the equation $x^2 = (2^{2^y} + 1)^s + 2^t$ holds, can t be odd?

If $x, y, s, t \in \mathbb{N}$ and the equation $$x^2 = (2^{2^y} + 1)^s + 2^t$$ holds, can t be odd if s is odd? I know that t cannot be odd when $s=1$. How about if $s>1$? Added September 10 2017: I was mistaken.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2017-09-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018796 | MSE-2423769

When does $p^s = 2^t + 1$ have a solution, for t odd and p prime?

If $p, s, t \in \mathbb{N}$, when does $$p^s = 2^t + 1$$ have a solution, for t odd and p prime? Note that, when $s = 1$, we have $p \equiv 1 \pmod 4$, and therefore $t = 2^s$ for some $s \in \mathbb{N}$ (i.e., p is a Fermat prime). I do not know what happens when $s > 1$, or if it is even possible.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2017-09-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018797 | MSE-2424059

If $F_m = 2^{2^m} + 1$ is prime (with $m \geq 1$), does it follow that $3 \mid \frac{F_m + 1}{2}$?

If $$F_m = 2^{2^m} + 1$$ is prime with $m \geq 1$, does it follow that $$3 \mid \frac{F_m + 1}{2}?$$ I have verified this to be true for $1 \leq m \leq 4$.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2017-09-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018798 | MSE-2426255

How to guess a formula

"The 'proof by induction' works well provided that we think we know the formula in advance. In other words, it is a good technique to transform a Conjecture in a Theorem." Biggs, N. L., Discrete Mathematics, ch. 4.4. Sometime, creating the proper Conjecture is on our shoulder. But, what is the criteria? What elements should I keep considered in order to develop a formula? E.g. Guess a formula for the sum $\sum_{r = 1}^n(br + c),$ where b and c are given numbers, and prove it by using the principle of induction. First, I tried to make a table: $n:\qquad 1\qquad\qquad 2\qquad\qquad 3\qquad\qquad 4\qquad\qquad 5 $ $S_n:\quad 1b+c\quad 3b + 2c\quad 6b + 3c\quad10b + 4c\quad 15b + 5c $ but I failed to transform this pattern in an usable formula. The solution reports $\frac{1}{2}bn^2 + (\frac{1}{2}b + c)n$. Which steps should I follow to get from the table I created to the correct formula? Apart from this specific case I reported as an example, is there a general method? Are there criteria? Guides? Is it mainly based on intuition and personal creativity and knowledge?

论坛问题帖 | has_answers | Worice | 2017-09-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018799 | MSE-2427561

Has this conjecture been discovered before? (Concerns prime numbers and not $\#(n)$ like in a question similar to this)

This is my second question today (I want to keep my major questions seperate): Let's denote the n^{th} prime number as p_n. Given the equation: $$\frac{p_1p_2p_3 \cdots p_n}{1 + p_1 + p_2 + p_3 + \cdots + p_n} = \{x_m : x_m \in\mathbb{Z}^+\}$$ Now the LHS is

not always equal to a positive integer, but when it is, it is equal to x_m. The first positive integer it will be equal to (which is 1) is denoted as x_1 such that $m = 1$ and $n = 2$, and the second integer (which is 715) is denoted as x_2 such that $m = 2$ and $n = 6$ etc. Conjecture: $$δ(x_m) = \{2^{2m - 1} : m > 1 \ \land \ δ(x_m) = \text{number of divisors of x_m including x_m itself}\}$$ $$m > 1 \because δ(x_1) = δ(1) = 1 \neq 2^{2\times 1 - 1} = 2$$ Is there a way to prove/disprove this? This is my own conjecture and I put my own little symbols to it. Could somebody please explain to me why this works and any examples where it doesn't (if they exist)? If this is somebody else's conjecture then I will delete this post. I am in Year 9, so my skill level in mathematics is not too high (compared to most users on this site), so I am sure somebody else must have discovered this. Thanks to this site, the only non-Year 9 maths I am familiar with include: Sets Matrices Sum/product series Σ and Π Imaginary (lateral) numbers $\sqrt{\text{negative}}$ (not particularly complex numbers $a + bi$) Quadratic equations Logarithms $\log_bn = m$ Trigonometry ΔABC and Linear Graphs $y = mx + c$. I guess these are the only things that may make you wonder if I actually am in Year 9. Everything else that what a Year 9 is usually taught, I also know. If you are using other techniques and ways of proving/disproving my conjecture, please be clear and try not to skip too many steps. I have not decided to name my conjecture as of yet (and I don't want to be selfish by naming it based on my own name) but I want to see what you think of it. Thank you in advance.

论坛问题帖 | no_answers_at_retrieval | Mr Pie | 2017-09-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018800 | MSE-2428281

some property of the simple continued fraction for the riemann-zeta function

Define the well known Riemann-zeta function $$\zeta(n)=\sum_{k=1}^{\infty}\frac{1}{k^n}$$ for natural number $n\gt1$ and let $a_1(n)$ be a sequence $a_1(1),a_1(2),a_1(3),\dots$ of the first partial quotients of the simple continued fraction for $\zeta(2n+1)$, $$\zeta(2n+1)=\{a_0(n);a_1(n),a_2(n),a_3(n),a_4(n),a_5(n),\dots\}$$ The sequence $a_0(n)=1$, and the other sequence $a_1(n)$ increases with n,while the rest of the sequence of partial quotients $a_2(n),a_3(n),a_4(n),\dots$ seem to be random How do we prove that $\lim_{n\rightarrow\infty} \frac{a_1(n)}{a_1(n+1)}=\frac{1}{4}$ Example $\zeta(151)=\{1;2854495385411919762116571931558091627803291516,1,4,1,1,\dots\},$ see wolframalpha and $\zeta(153)=\{1;1141798154164767904846628774254547461112916991 5,7,1,15,1,\dots\},$ see wolframalpha then $$\frac{a_1(75)}{a_1(76)}=\frac{2854495385411919762116571931558091627803291516}{1141798154164767904846628774254547461112916991 5}=0.249999999999999999999999999996428\dots$$

论坛问题帖 | has_answers | Nicco | 2017-09-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018801 | MSE-2439707

Does a counterexample always disprove a conjecture?

I'm sincerely confused on this topic (the title). I was wondering if a counterexample always disproved a conjecture. I'm leaning towards true because it's not really a counterexample if it evaluates to false... right? Any insight or help would be dearly appreciated.

论坛问题帖 | has_answers | NerdOfCode | 2017-09-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018802 | MSE-2441516

Two equivalent simple continued fractions $A\overset{\color{red}?}=B$

Fix a sequence a_n of complex numbers for natural number n,such that $|a_n|\gt1$ and then define the following two simple continued fractions $A= \cfrac{1}{a_{0}-\cfrac{1}{a_{1}+\cfrac{1}{a_{2}-\cfrac{1}{a_{3}+\cfrac{1}{a_{4}-\cfrac{1}{a_{5}+\cfrac{1}{a_{6}-\cfrac{1}{a_{7}+\dots}}}}}}}}\tag1$ which converges by Śleszyński-Pringsheim Theorem and $B= \cfrac{1}{a_{0}-1+\cfrac{1}{1+\cfrac{1}{a_{1}-1+\cfrac{1}{a_{2}-1+\cfrac{1}{1+\cfrac{1}{a_{3}-1+\cfrac{1}{a_{4}-1+\cfrac{1}{1+\cfrac{1}{a_5-1+\cfrac{1}{a_{6}-1+\cfrac{1}{1+\dots}}}}}}}}}}\tag2$ How do we prove that $A=B$? Special case (i) Define $R(q)=q^{1/5}\frac{(q;q^5)_{\infty}(q^4;q^5)_{\infty}}{(q^2;q^5)_{\infty}(q^3;q^5)_{\infty}}$ the Rogers-Ramanujan continued fraction where $|q|\lt1$,then $R(q)=\cfrac{q^{1/5}}{1+\cfrac{q}{1+\cfrac{q^2}{1+\cfrac{q^3}{1+\cfrac{q^4}{1+\cfrac{q^{5}}{1+\dots}}}}}}$ and $R(q)=\cfrac{q^{1/5}}{1+\cfrac{q}{1-q+\cfrac{q}{1-\cfrac{q}{1+q+\cfrac{q^3}{1+q^2-\cfrac{q^2}{1+\cfrac{q^2}{1-q^2+\cfrac{q^5}{1-q^3+\cfrac{q^3}{1-\cfrac{q^3}{1+q^3+\cfrac{q^7}{1+q^4-\cfrac{q^4}{1+\dots}}}}}}}}}}}$ (ii) Let $a_n=2$,then $A= \cfrac{1}{2-\cfrac{1}{2+\cfrac{1}{2-\cfrac{1}{2+\cfrac{1}{2-\cfrac{1}{2+\dots}}}}}}=\frac{1-\sqrt{5}}{2}$ and $B=\cfrac{1}{1+\cfrac{1}{1+\cfrac{1}{1+\cfrac{1}{1+\cfrac{1}{1+\cfrac{1}{1+\dots}}}}}}=\frac{1-\sqrt{5}}{2}$ Remark: this general identity leads to another continued fraction for the Generalized Rogers-Ramanujan continued fraction Edited 17 Oct 2017:I've just noticed that this oeis article also discusses the same continued fraction for the special case of the Generalized Rogers-Ramanujan continued fraction $R(-1,q)$

论坛问题帖 | has_answers | Nicco | 2017-09-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018803 | MSE-2444412

A question about Brown Numbers $(m, n) : n! + 1 = m^2$

So I recently looked at a video on YouTube by Numberphile which is accessible down below: Brown Numbers $-$ Numberphile The video was about so-called numbers called Brown Numbers. These types of numbers are a pair of integers (m, n) such that $$n! + 1 = m^2$$ When you have a term looking a little like this $\longrightarrow n!$ it is not read as shouting the term before the exclamation mark (and the term before the exclamation mark isn't called the coefficient either). It is read as, in this case, n factorial and is equal to the product of every number from 1 to n including n itself. $$n! = 1\times 2\times 3\times \cdots \times n = \prod_{k = 1}^{n}k$$ Understanding what $n!$ means, we can now find a

pair of Brown Numbers to substitute for m and n in the equation, however in the video, it was revealed that only three pairs of Brown Numbers have been found, and it has been conjectured that there only exists three pairs of Brown Numbers. $$\begin{align} (5, 4) \qquad 4! + 1 &= 1\times 2\times 3\times 4 + 1 = 24 + 1 = 25 = 5^2 \\ (11, 5) \qquad 5! + 1 &= 1\times 2\times 3\times 4\times 5 + 1 = 120 + 1 = 121 = 11^2 \\ (71, 7) \qquad 7! + 1 &= 1\times 2\times 3\times 4\times 5\times 6\times 7 + 1 = 5040 + 1 = 5041 = 71^2 \end{align}$$ It was also revealed in the video that this conjecture has not been proven, so I accepted the challenge of proving it, but came across a problem. My Attempt: If $$n! + 1 = m^2$$ then $$n! = m^2 - 1 = m^2 - 1^2 = (m + 1)(m - 1)$$ Assume that $m = 2r$ for some integer r. $$n! = (2r + 1)(2r - 1) = (2r)^2 - 1^2 = 4r^2 - 1$$ $$\implies n! + 1 = 4r^2$$ We arrive at a contradiction. Since $n! = 1\times 2\times \cdots \times n$ then if $n > 1$, n is even. Any even number plus 1 is an odd number, and an even number cannot divide an odd number, thus 4 does not divide an odd number, so m cannot be even. Now let $m = 2r \pm 1$. $$n! = (2r \pm 1 + 1)(2r \pm 1 - 1) = (2r \pm 2)(2r) = 2(r \pm 1)(2r) = 4r(r \pm 1)$$ We now arrive at another contradiction. 4 now divides m, but 4 is even and m is odd! (This attempt can serve as a Lemma to proving that for any integer x, if x is even/odd, then x^y is even/odd.) Did I do something wrong? I must have. Besides, every Brown Number so far is odd and not even.

论坛问题帖 | has_answers | Mr Pie | 2017-09-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018804 | MSE-2453376

Feedback on a general conjecture for A = Bp - Cq where p and q are primes?

I previously asked for feedback on the following conjectures (similar in appearance to Goldbach ‘s and Lemoine’s see: Two new conjectures related to Lemoine's and Goldbach's) : $O = 2n + 1 = p - 2q$ always has a solution in distinct odd primes p and q for $O \ge 5$ $O = 2n + 1 = 2q - p$ always has a solution in p and q with $q < p$, p being an odd prime and q being 1 or an odd prime and $O \ge 1$ Those two conjectures, even though they do not seem easy to prove, are very weak: the number of possible solutions is possibly infinite. (Since increases in p and q have opposite effect.) Also, while it is possible by concrete calculations to increase the domain of proven validity, it is not possible to find a concrete counter example since a solution may lay with higher values of p and q. Since they are so easy to satisfy, I thought that they may become more interesting if they were generalized by changing the component $2q$ into $2nq$. It is easy to see that in $O = p-2nq$ we will not have solutions in primes p and q as long as n and O have common divisors. So the revised conjectures become: For O, odd natural number, and n ,natural number ,without common divisors: $O = p - 2nq$ always has a solution in distinct odd primes p and q for $o \ge 5$ $O = 2nq - p$ always has a solution in p and q with $q < p$, p being an odd prime and q being 1 or an odd prime and $o \ge 1$ I verified that it is the case for all $O < 2000000$ and $n < 1000$ But better, a further generalization is possible that envelops these conjectures: $A = Bp - Cq$ where the natural numbers A,B,C have no common divisor and one of them is even has solutions in p and q odd primes or 1 I verified this with values of A, B and C ranging from 1 to 1000.

论坛问题帖 | has_answers | Raymond Duchesne | 2017-10-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018805 | MSE-2459057

Is this conjecture correct ${(2\pi)^{m-1}\Gamma(m)\over m^m}=\prod_{k=1}^{\infty}{mk\over mk-(m-1)}\cdot{mk\over mk+(m-1)}?$

The Wallis's Product $${\pi\over 2}={2\over 1}\cdot{2\over 3}\cdot{4\over 3}\cdot{4\over 5}\cdot{6\over 5}\cdot{6\over 7}\cdots\tag1$$ Conjecture: The generalised of (1) $${(2\pi)^{m-1}\Gamma(m)\over m^m}=\prod_{k=1}^{\infty}{mk\over mk-(m-1)}\cdot{mk\over mk+(m-1)}\tag2$$ $m\ge2$ and $\Gamma(m)$ is the Gamma function How do we prove $(2)?$

论坛问题帖 | has_answers | gymbvghjkgkjkhgfkl | 2017-10-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018806 | MSE-2472205

Is every number a sum of 3 tetrahedral numbers?

It is known that every number can be represented by a sum of 3 triangular numbers. According to Gauss (see formula 35 in mathworld article) $$ \text{num}=\Delta+\Delta+\Delta $$ I did some numerical experiments that suggest the above formula is correct when triangular numbers are replaced by tetrahedral numbers $$ \Delta=\frac{n(n+1)(n+2)}6 $$ if n is allowed to be negative. Is this conjecture correct? I tried to google representation of integers by tetrahedral numbers but didn't find anything.

论坛问题帖 | has_answers | Tyrell | 2017-10-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018807 | MSE-2476870

About three conjectures concerning the Pythagorean Theorem.

I have two main conjectures relating to the Pythagorean Theorem that I desperately want to find out if they are true or not. Could somebody please help me? If $ \ a^2 + b^2 = c^2 \ $ for which c is a prime number, then for some $n \in \mathbb{N}, \ c = 4n + 1$. $\forall \{x, y, z\} \subset \mathbb{N}, \ \big(x^2 + (x + 1)^2\big)^2 = y^2 + z^2 \ $ for which $x \neq y \neq z$. Tested for $c, x \leqslant 10^6$. I did also have another conjecture that if $ \ a^2 + b^2 = c^2 \ $ for which a is odd (and greater than 1), b is even, and $a < b$ then there existed a solution such that $c = b + 1$, but I am 100% certain this has been discovered before (assuming so, since it was not very difficult to notice). However I am not sure if the other two conjectures have been discovered before, although if it had to be one of them, I would pick the second one. So could somebody also please clarify that for me? Thank you in advance.

论坛问题帖 | has_answers | Mr Pie | 2017-10-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018808 | MSE-2481335

If $a,b,c,a^2+b^2+c^2$ are primes, then a or b or c is equal to 3

Given primes a,b,c such that $a^2+b^2+c^2$ is prime, then $3\in\{a,b,c\}$. Tested for $a,b,c<500$.

论坛问题帖 | has_answers | Lehs | 2017-10-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018809 | MSE-2488065

Conjectured continued fraction for the Jacobi theta function $\vartheta_{4}(q)$

Given the Jacobi theta function $$\vartheta_{4}(q)=1+2\sum_{n=1}^{\infty}(-1)^nq^{n^2}$$ where $q=e^{2\pi i\tau}$, and $|q|\lt1$. It is conjectured that it has the following continued fraction $\vartheta_{4}(q)=1-\cfrac{2q}{1-q+\cfrac{q}{1-\cfrac{q^2}{1-q^3+\cfrac{q^2}{1-\cfrac{q^3}{1-q^5+\cfrac{q^3}{1-\cfrac{q^{4}}{1-q^7+\cfrac{q^{4}}{1-\dots}}}}}}}}\tag{1a}$ and its reciprocal, $\begin{aligned}\frac{1}{\vartheta_{4}(q)}=1+\cfrac{2q}{1-q-\cfrac{q}{1+\cfrac{q^2}{1-q^3-\cfrac{q^2}{1+\cfrac{q^3}{1-q^5-\cfrac{q^3}{1+\cfrac{q^{4}}{1-q^7-\cfrac{q^{4}}{1+\dots}}}}}}}}\end{aligned}\tag{1b}$ How do we prove the conjecture? Edited later: If we define the q-pochhammer symbol for any complex number a, $$(a;q)_{\infty}=\prod^{\infty}_{n=0}\left(1-aq^n\right) $$ the conjectured continued fractions can be generalized to the following form $\begin{aligned}\prod_{n=0}^\infty\frac{\big(1-aq^n\big)}{\big(1+aq^n\big)}=1-\cfrac{2a}{1-q+\cfrac{a}{1-\cfrac{aq}{1-q^3+\cfrac{aq}{1-\cfrac{aq^2}{1-q^5+\cfrac{aq^2}{1-\cfrac{aq^{3}}{1-q^7+\cfrac{aq^{3}}{1-\dots}}}}}}}}\end{aligned}\tag{1c}$ which obviously leads to the continued fraction for the reciprocal of $\vartheta_{4}(q)$ when $a\rightarrow -q$.

论坛问题帖 | has_answers | Nicco | 2017-10-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018810 | MSE-2488843

$\gcd(p_1+p_2,p_1+p_3,p_2+p_3)\leq 2$ if $p_1<p_2<p_3$ are primes

Conjecture about greatest common divisors and primes: $\gcd(p_1+p_2,p_1+p_3,p_2+p_3)\leq 2$ if p_1,p_2,p_3 are different primes. Tested for all $p_1,p_2,p_3<500$. I'm pretty sure that someone can find an easy proof of this, but I have no clue myself.

论坛问题帖 | has_answers | Lehs | 2017-10-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018811 | MSE-2490483

The set of numbers of the form $q_1+q_2^2+q_3^3+q_4^4+q_5^5$ where all q_k are primes.

Conjecture: All prime numbers greater than 109 is of the form $\displaystyle\sum_{k=1}^5q_k^k$, where all q_k are primes. The conjecture is extracted from the question and answer here: Annoying primes Tested for all primes $<70,000$. The exceptional primes seems to be $\{2,3,5,7,11,13,17,19,23,29,31,37,41,43,47,53,59,61,109\}$ Are there heuristic arguments for the conjecture? Also: All primes except $\{2,3,5,7,11,13,17,19,23,29,37,43,53,61,67\}$ seems to be of the form $\displaystyle\sum_{k=1}^4q_k^k$. All primes except $\{2,3,5,7,11,13,37,61,127\}$ seems to be of the form $\displaystyle\sum_{k=1}^3q_k^k$. Both cases tested for primes less than $10,000$.

论坛问题帖 | has_answers | Lehs | 2017-10-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018812 | MSE-2491672

Conjecture: $a^3 + b^3 + c^3 = p^3 \Rightarrow x^3 + y^3 + z^3 = \big(\frac{a + b + c}{2}\big)^3$

Conjecture: Let p be an even perfect number, and a, b, c be positive natural numbers. There exists values for a, b, c to satisfy the following equation $$a^3 + b^3 + c^3 = p^3$$ for which $\bigg(\dfrac{a + b + c}{2}\bigg)^3 = x^3 + y^3 + z^3$ for natural numbers x, y, z. Tested for all perfect numbers up to $2^{30}(2^{31} - 1) = 2305843008139952128$. I posted something similar relating to this a while ago on the Math Overflow and now I need better computer-power. Ideally, I want a better way to approach this problem in order to prove/disprove this, but I do not know how or where to begin. Could somebody please help me? Also, this is just a speculation, but every prime number can be written of the form $a^3 - b^3 - c^3$ for some natural a, b, c. This includes 2, but there is already a formula for 2: $$2 = (6t^3 + 1)^3 - (6t^3 - 1)^3 - (6t^2)^3$$ Also, if you have the following equation: $$\qquad \ \ \ a^3 + b^3 + c^3 = \big\{d^3 : \min\{a, b, c\} = 6\big\}$$ $$\implies 6 \mid d + (-1)^{n + 1}$$ for which $a^3 + b^3 + c^3 = d^3$ is the n^{th} equation such that $\min\{a, b, c\} = 6$. Thank you in advance. Edit: Here is the link to my post on the $MO \longrightarrow$ About prime numbers and the sum of three cubes And here are a few examples of a, b, c, x, y, z as part of the conjecture in the yellow box: Let $a = 3, b = 4, c = 5$ then $x = 3, y = 4, z = 5$. Let $a = 18, b = 19, c = 21$ then $x = 11, y = 15, z = 27$. Let $a = 57, b = 82, c = 495$ then $x = 15, y = 213, z = 281$ or $x = 162, y = 173, z = 282$. Let $a = 2979, b = 4005, c = 7642$, this was too high for me to calculate a while ago so I went here to find the value of $\big((a + b + c)/2\big)^3$ which I did, being equal to 7313, meaning that 7313^3 can be written as the sum of three positive cubes.

论坛问题帖 | no_answers_at_retrieval | Mr Pie | 2017-10-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018813 | MSE-2493889

The fractal dimension of the Kolakoski sequence is $2-1/e$

The Kolakoski sequence, which is defined as the infinite sequence of symbols {1,2} that is its own run-length encoding (Wikipedia), has been suggested to be self-similar^{1}. The fractral dimension of a self-similar time-series is directly related to its Hurst exponent^{2}. I have estimated the Hurst exponent of the Kolakoski sequence for different sequence lengths, and found that the answer converges near $1 / e$ when the sequence length increases. This suggests that the fractral dimension of the Kolakoski sequence is $D=2-H = 2-1/e.$ Here is a small subset of the data sequence length hurst exponent 1e2 .1167 1e3 .1796 1e4 .2236 1e5 .2579 1e6 .3108 1e7 .3464 1e8 .3657 2e8 .3720 3e8 .3766 My question is: Can you find a proof for this conjecture? If yes, I would be interested to write a brief note about this to a mathematical journal. References: ^{1} https://maths-people.anu.edu.au/~brent/pd/Kolakoski-ACCMCC.pdf ^{2} https://en.wikipedia.org/wiki/Hurst_exponent#Relation_to_Fractal_Dimension

论坛问题帖 | no_answers_at_retrieval | tui | 2017-10-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018814 | MSE-2504263

A conjecture concerning numbers of the form $k\sqrt{2}$ where $k \in \mathbb{N}$

I was experimenting with radicals on my calculator when I noticed that the units digit and the hundredths digit of $\sqrt{2}, \, 2\sqrt{2}, \, ... \, 11\sqrt{2} \,$ are the same. The next one does not work since in $12\sqrt{2} = 16.97...$ ($6 \neq 7$). On doing this up to $50\sqrt{2}$, I found that the numbers $\alpha\sqrt{2}$ which don't work are when $\alpha = 12, 19, 24, 31, 36, 38, 41, 43, 48, 50$. My conjecture: The units digit and the hundredths digit of a number of the form $k\sqrt{2} \,$ ($k \in \mathbb{N}$) are the same, except when k is of the form $12a+19b+41c$ where $a, b, c \in \mathbb{Z^+} \cup \left \{0 \right \}$. Does anyone know how to prove this (or give a counterexample)? Also, if this is true, are there similar conjectures for numbers of the form $k\sqrt{3}, \, k\sqrt{5}, \, ...$? Thanks!

论坛问题帖 | has_answers | TheSimpliFire | 2017-11-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018815 | MSE-2509517

Repdigit Conjecture

I noted a conjecture down a while back after doing a fair few calculations on repdigits and noticing patterns. Define some digit repetition syntax for a 'string of integers' x and an integer y, let $x;y$ be x repeated and concatenated y times. For example $$ 123;3 = 123123123\\ 9;10 = 9999999999 $$ then it seemed that for all strings I have tried the following was true: For a prime b, if $\frac{s;a}{b}$ is an integer then $\frac{s;k}{b}$ is also an integer for any k such that $k \bmod (b-1) = a$ This is best shown with some examples, for $s=165$, $a=3$, $b=5$, $$ \frac{s;a}{b} = \frac{165165165}{5} = 33033033 \in \mathbb{Z}\\ \{k:k\bmod(b-1)=a\} = \{k:k\bmod 4 =3\} = \{3,7,11,\cdots\} $$ so we can predict that $$ \frac{165;7}{5}=\frac{16516516516516516565}{5} \in \mathbb{Z} \;\;(true)\\ \frac{165;11}{5}=\frac{165165165165165165165165165165}{5} \in \mathbb{Z} \;\;(true)\\ $$ and so on. Moreover, it seems we can concatenate (denoted by the $|$ symbol) some other strings around this pattern take for example \begin{equation} \frac{11222221}{97}= \frac{11|2;5|1}{97} \in \mathbb{Z} \end{equation} then we have $a=5,b=97,k=5,101,197,\cdots$ \begin{equation} \frac{11|2;101|1}{97} \in \mathbb{Z} \end{equation} (which appears to be true) This leads to a modified conjecture For a prime b and strings A and B, if $\frac{A|s;a|B}{b}$ is an integer then $\frac{A|s;k|B}{b}$ is also an integer for any k such that $k \bmod (b-1) = a$ This seems to work if the number is a repdigit in any integer base for example in binary $427=110101011_2=(11|01;3|1)_2$ we have $$ \frac{(11|01;3|1)_2}{7} = \frac{427}{7} = 61 \in \mathbb{Z}\\ k = \{3,9,15,\cdots\}\\ \frac{(11|01;9|1)_2}{7} = \frac{1747627}{7} \in \mathbb{Z} \\ \frac{(11|01;15|1)_2}{7} = \frac{7158278827}{7} \in \mathbb{Z} $$ This makes me suspect it is something to do with the pattern rather than the numeric content per se. Questions: Can the conjecture be proved true or false? Can a counter example be found? Why does this work/not work?

论坛问题帖 | has_answers | Benedict W. J. Irwin | 2017-11-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018816 | MSE-2510783

Can this summation be proved to equal 1 for any upper bound 0 or greater?

$$\sum_{i=0}^n \frac{(n-i+1)^n (-1)^i}{i!(n-i)!} =1$$ Prove that the summation above always equals 1 for any integer n greater than or equal to zero. I believe that it probably does equal one always. I've tested low numbers for n and used computers to get slightly higher values confirmed. The problem comes when the factorials and number of terms get to high for accurate computation. I'm looking for a way to prove it is true for all numbers that meet the integer and greater than zero stipulations. If there is a counter example, then that also works to disprove the statement. I'm guessing that simplification of the summation would be useful for a proof, but I've found it very difficult to create a more simplified summation than the one above. If you try out a few numbers, you'll notice that the way that the summation works out to be one is pretty cool.

论坛问题帖 | has_answers | Luke Surak | 2017-11-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018817 | MSE-252950

On expressing a square as a sum of two cubes

Given $a,b,c \in \mathbb{N}$ which satisfy the following conditions: $a^3 + b^3 = c^2$ $ a \neq b$ =-=-=-=-=-=-=-=-=-=-=-=-=-= EDIT, Will Jagy: The conjecture is that, for a given $c,$ there are at most two distinct pairs (a,b) with $1 \leq a < b$ and $a^3 + b^3 = c^2.$ Note the example $77976^2=1026^3+1710^3=228^3+1824^3$ that Gerry found with exactly two pairs. =-=-=-=-=-=-=-=-=-=-=-=-=-= There are maximum 2 possible pairs of (a, b) which satisfy this conditions. I have verified it for $a, b \leq 200\,000$. Note: the situation when a and b just take each others value is discarded.

论坛问题帖 | has_answers | Alex Filipovici | 2012-12-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018818 | MSE-2533599

Confirmation of a conjecture: $3\mid (8n)^5 - k$

I have made a conjecture: $$\forall n\in \mathbb{N} \setminus \{6r : r \in \mathbb{N}\}, \ 3\mid (8n)^5 - k$$ Such that $k =$ a constant containing all the digits of $(8n)^5$ except for the last 3. The order of digits is unecessary. e.g. Let $n = 1$. $(8\times 1)^5 = 32768 \Rightarrow k_1 = 32\require{cancel}\cancel{768} \ \land \ k_2 = 23\cancel{768}$ $32768 - 32 = 32736 = 3\times 10912$ or $32768 - 23 = 32745 = 3\times 10915$ $\therefore 3\mid (8n)^5 - k_1, k_2$ I do not know how to prove/disprove it apart from using computer power and trying to find a counter-example, however this has been tested for all $n \leqslant 10^6$ and I have not found one. Could somebody please help me? Thank you in advance.

论坛问题帖 | has_answers | Mr Pie | 2017-11-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018819 | MSE-2562400

Infinitely many simultaneous palindromes in "exponentially close" number bases?

I've observed regular patterns that suggest that there are infinitely many numbers which are simultaneously palindromic in n number bases of form: $$ b_1,b_2,b_3,\dots,b_n={x^{y_1},x^{y_2},x^{y_3},\dots,x^{y_n}} $$ For every combination of natural numbers $(n,x,y_1,\dots,y_n)\in\mathbb N$ you can pick. Where of course, $x\ge2$, since we are observing integer number bases $b_n\ge2$. For one example, this conjecture implies that there are infinitely many numbers simultaneously palindromic in number bases $3,9,81$; Some examples for this example: 10 ,20 ,82 ,164 ,820 ,1640 ,6562 ,6643 ,6724 ,13124 ,13205 ,13286 ,65620 ,66430,... The 66430 is an example since it can be written as $10101010101_3=111111_9=AAA_{81}$ Are there similar, more/less powerful(general) claims made/proven somewhere? Is there anything that would suggest a way this can be proven? (or perhaps disproven/reduced to a less general statement?) It is worth noting that the patterns were observed for $n\le3$, and $n=4$ for smaller x,y variables, since bigger values are too hard to compute for more than couple examples using my python code. (And thus hard to see/confirm whether the pattern extends to larger n) I was observing the gaps between palindromes, and palindromes produced by this conjecture, to formalize the patterns for smaller and simpler cases of variable combinations. Simplest case (pattern) in context For simplest example, we can generate infinitely many such palindromes for $b_1=2,b_2=4$ case using $a(x)=$A007583 and $b(x)=$A001511 to predict the gaps with: $$ g(k,d) = \begin{cases} 2\cdot a(b(k)), & \text{$d=4t-1 \land b(k)>2$} \\ a(b(k)), & \text{otherwise} \end{cases} $$ (See g values in the output below) The k goes from 1 to $2^{\lfloor(d/4)\rfloor}$ where d is the number of digits of odd digit palindromic examples behind the generated gaps (in b_1 representation). There are finitely many examples (one for each k) per d digit palindromes for each d, but infinitely many as d tends to infinity - grows larger. For more clarity, each following row contains gaps produced by g function, between examples for this case of "$d :$" digits: (In other words, each "," represents a number palindromic in both bases 2 and 4, and the values between them are the number of "$d :$" digit palindromes that are only palindromic in base 2) 3 : 1 5 : 1 ,1 7 : 1 ,2 9 : 1 ,1 ,3 ,1 11 : 1 ,2 ,6 ,2 13 : 1 ,1 ,3 ,1 ,11 ,1 ,3 ,1 15 : 1 ,2 ,6 ,2 ,22 ,2 ,6 ,2 17 : 1 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 19 : 1 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 21 : 1 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 23 : 1 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,342 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 25 : 1 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,683 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 27 : 1 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,342 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,1366 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,342 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 29 : 1 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,683 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,2731 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,683 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 31 : 1 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,342 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,1366 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,342 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,5462 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,342 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,1366 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,342 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 33 : 1 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,683 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,2731 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,683 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,10923 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,683 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,2731 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,683 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 35 : 1 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,342 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,1366 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,342 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,5462 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,342 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,1366 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,342 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,21846 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,342 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,1366 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,342 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,5462 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,342 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,1366 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,342 ,2 ,6 ,2 ,22 ,2 ,6 ,2 ,86 ,2 ,6 ,2 ,22 ,2 ,6 ,2 37 : 1 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,683 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,2731 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,683 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,10923 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,683 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43691 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,683 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,2731 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,683 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,10923 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,683 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,2731 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,683 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,171 ,1 ,3 ,1 ,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1

,11 ,1 ,3 ,1 ,43 ,1 ,3 ,1 ,11 ,1 ,3 ,1 By picking large enough d we can get as many examples as we want. All this is for the simplest case of bases $(2,4)$. Similar patterns in the observed gaps (between palindromic representations) seem to exist for other combinations of variables, but they get rarer in examples and less predictable as n grows. For every odd d there seem to always be examples for any x,y's for small n, but for larger n we start getting more and more (always periodic?) gaps in these sets (rows above) of d digit odd palindromic examples gaps. By these gaps in gaps, I mean that some rows start getting blank (no examples), but others still seem to follow in increasing number of examples (so far non blank rows seem to appear periodically).

论坛问题帖 | has_answers | Vepir | 2017-12-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018820 | MSE-2562525

Confirmation of Proof: If p is a perfect number, then $1 + 8p$ is a squared number.

I have created a hypothesis: $$\text{If } p \text{ is a perfect number, then } 1 + 8p \text{ is a squared number.}$$ Could anyone prove/disprove the above statement? I have no idea where to begin, and I discovered this by pure accident when trying to solve a quadratic equation. Edit: I believe that all odd numbers m squared can be expressed as the form $8n + 1$ for some $n \in \mathbb{N}$. $$\begin{align} 3^2 &= 9 = 8\times 1 + 1 \\ 5^2 &= 25 = 8\times 3 + 1 \\ 7^2 &= 49 = 8\times 6 + 1 \end{align}$$ Yes, it is obvious that n is also a triangular number $(1 + 2 + 3 + \cdots)$. Let $m = 2k + 1$, then one has the following: $$(2k + 1)^2 = 8n + 1$$ And $(2k + 1)^2 = 4k(k + 1) + 1$, so it follows, then, that: $$4k(k + 1) \require{cancel}\cancel{+1} = 8n \cancel{+1} \Rightarrow k(k + 1) = k^2 + k = 2n$$ And since the parity of $r \in \mathbb{N}$ remains the same for r^k for all $k \in \mathbb{N}$ then our proof stands valid.

论坛问题帖 | has_answers | Mr Pie | 2017-12-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018821 | MSE-25660

Status of a conjecture about powers of 2

I recently saw a conjecture on a blog (http://blog.tanyakhovanova.com/?p=311) which the author refers to as the 86 conjecture. The conjecture claims that all powers of 2 greater than 2^{86} have a zero in their base 10 representation. At first glance this seems like a numerical curiosity, and I wasn't sure that this was an interesting or deep mathematical question. I wanted to ask the community if they knew of any existing work that deals with either this type of question or something similar. I tried searching for 86 conjecture on Google and didn't get anything useful.

论坛问题帖 | has_answers | svenkatr | 2011-03-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-018822 | MSE-2579148

Sequences $p,p+d,p+2d,\dots,p+N\cdot d$ of prime numbers

A conjecture that I believe is possible to prove: For any $N\in\mathbb N$ there is a $d\in\mathbb N$ such that $p,p+d,p+2d,\dots,p+N\cdot d$ all are primes (for some prime p). My context: it seems that greater $\omega(d)$ (number of different prime factors) gives greater N.

论坛问题帖 | has_answers | Lehs | 2017-12-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018823 | MSE-2579578

The difference of two coprime composites

I guess this is a simple task for them who know some number theory: Any natural number is the difference between two coprime composites. Tested up to 1000. I develop some computer tools to investigate sets and now and then find some patterns or connections that makes me formulate a conjecture, which in my mind just is a word for an unproved statement.

论坛问题帖 | has_answers | Lehs | 2017-12-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018824 | MSE-2590035

A conjecture for an inequality for every odd number.

Let $E(n) = \sum_{d|n} d \log(d) $. Then for coprime n,m we have $$E(mn) = \sigma(m) E(n) + \sigma(n) E(m).$$ Suppose there exists an odd perfect number n. Then$$E(2n) = E(2)\sigma(n)+E(n)\sigma(2) = 4 n \log(2) + 3 E(n).$$ But looking at a few numerical examples, one can see that the right hand side seems to be alway bigger than the left hand side. Consider the upper bound for logarithm:$$\log(x) \le x^s/s. \quad \forall s>0$$ Making use of this upper bound we can derive: $$E(2n) = 2 \sum_{d|n} d \log(2d) + E(n) \le 2 \sum_{d|n} \frac{(2d)^s}{s} + E(n). \quad \forall s > 0$$ Now the question is, how do we choose $s>0$ such that we have $$2 \sum_{d|n} \frac{(2d)^s}{s} + E(n) < 4 n \log(2) + 3 E(n).$$ My conjecture is, that it is always possible to choose one such $s>0$ such that the last inequality is valid.

论坛问题帖 | has_answers | user276611 | 2018-01-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018825 | MSE-2602144

Conjectured value of a difficult integral with Dedekind eta functions

In my research on quantum groups I have the following conjecture: \begin{equation} \int_0^\infty\frac{\eta (2 i x)^8}{\eta (i x)^2 \eta (4 i x)^2}\,dx\,{\stackrel?=}\,\frac{K(\tfrac{1}{\sqrt{2}})}{\pi}\tag{1} \end{equation} where \begin{equation} \eta(ix)=e^{-\frac{\pi x}{12}}\prod_{n=1}^\infty (1-e^{-2n\pi x}) \end{equation} is the Dedekind eta function and \begin{equation} K(\tfrac{1}{\sqrt{2}})=\frac{\Gamma^2(\tfrac14)}{4\sqrt{\pi}} \end{equation} is the

Elliptic integral singular value. This value has been guessed using Inverse symbolic calculator and then checked numerically to a high precision, but I do not have a proof. Question: Is (1) true?

论坛问题帖 | has_answers | user520487 | 2018-01-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018826 | MSE-2603021

Where is the flaw in my proof?

Possible proof of the conjecture that there always exists a prime between n^2 and $(n+1)^2$? Assume the statement is invalid. Therefore, for some $n \in \mathbb{N}$, there does not exist a prime that ranges from n^2 and $(n+1)^2$ exclusive. Let $\pi (x)$ denote the amount of prime numbers less than or equal to x, then we have that if $\pi ((n + 1)^2) - \pi (n^2) = k$, there exists k primes that range between n^2 and $(n + 1)^2$ exclusive. Therefore, by assuming the conjecture is false, we consider $k = 0$, which consequently gives us the following: $$\begin{align} \pi ((n + 1)^2) - \pi (n^2) &= 0 \\ \Leftrightarrow \pi ((n+1)^2) &= \pi (n^2). \end{align}$$ It is well known that $\pi (x) \approx \dfrac{x}{\ln x}$ so by order of substitution, $$\frac{n^2}{\ln n^2} \approx \frac{(n + 1)^2}{\ln (n + 1)^2}.$$ For some m, one can see that $\ln x^2 = m \Rightarrow e^m = x^2 \Rightarrow \sqrt {e^m} = e^{m/2} = x \Rightarrow \ln x = m/2$. Therefore, we conclude that $\ln x^2 = 2\ln x$. $$\begin{align} \therefore \frac{n^2}{2\ln n} &\approx \frac{(n + 1)^2}{2\ln (n + 1)} \\ \\ \Leftrightarrow \bigg(\frac n2\bigg)\bigg(\frac{n}{\ln n}\bigg) &\approx \bigg(\frac{n + 1}{2}\bigg)\bigg(\frac{n + 1}{\ln (n + 1)}\bigg) \\ \\ \Leftrightarrow \bigg(\frac n2\bigg)\pi (n) &\approx \bigg(\frac{n + 1}{2}\bigg)\pi (n + 1) \\ \\ \Leftrightarrow \frac{n}{n + 1} &\approx \frac{\pi (n + 1)}{\pi (n)}. \end{align}$$ Now, by taking the reciprocal of the equation, it follows then that, $$\frac{n + 1}{n } = 1 + \frac 1n \approx \frac{\pi (n)}{\pi (n + 1)}.$$ However, although true, $$\square \ \dfrac{n + 1}{n} > \dfrac{\pi (n)}{\pi (n + 1)}.$$ By definition of the function π, we have that $\pi (n) = \pi (n + 1)$ if and only if n and $n + 1$ are not prime and $\pi (n) < \pi (n + 1)$ if and only if $n + 1$ is prime, therefore $\pi (n) \leqslant \pi (n + 1)$. We thus arrive at the following conclusion: $$\begin{align} \frac{\pi (n)}{\pi (n + 1)} &\leqslant 1 \\ \\ \Leftrightarrow \frac{n + 1}{n} &> \frac{\pi (n)}{\pi (n + 1)} \\ \\ \Leftrightarrow \pi ((n + 1)^2) &> \pi (n^2) \\ \\ \Leftrightarrow \pi ((n + 1)^2) - \pi (n^2) &\geqslant 1 \\ \\ \therefore k &= 1. \end{align}$$ Thus, the conjecture is true.$\qquad\qquad\qquad

V3-018830 | MSE-2613252

A variation on Giuga's conjecture

Can you provide a counterexample to the following claim : Let n be an odd natural number greater than one , let k be a natural number such that $k \le n$, then n is prime if and only if : $\displaystyle\sum_{i=0}^{k-1}i^{n-1} + \displaystyle\sum_{j=0}^{n-k}j^{n-1} \equiv -1 \pmod{n}$ You can run this test here . I was searching for a counterexample using the following two PARI/GP codes : VariationGiuga1(k,ub)={ forcomposite(n=k,ub, if(Mod(n,2)==1, s=sum(i=0,k-1,lift(Mod(i,n)^(n-1)))+sum(j=0,n-k,lift(Mod(j,n)^(n-1))); if((Mod(s,n)==n-1),print("n="n)))) } VariationGiuga2(k,ub)={ forprime(n=k,ub, s=sum(i=0,k-1,lift(Mod(i,n)^(n-1)))+sum(j=0,n-k,lift(Mod(j,n)^(n-1))); if(!(Mod(s,n)==n-1),print("n="n))) }

论坛问题帖 | has_answers | Pedja | 2018-01-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018831 | MSE-2615963

A congruence involving Chebyshev polynomials

Can you provide a proof or a counterexample to the following claim : Let n be a natural number greater than one and let $T_{n}(x)$ be Chebyshev polynomial of the first kind , then n is prime if and only if :$\displaystyle\sum_{k=0}^{n-1}2T_{n-1}\left(\frac{k}{2}\right) \equiv -1 \pmod n$. You can run this test here . I have tested this claim up to 10000 . I was searching for counterexample using the following two Pari/GP codes : Chebyshev1(lb,ub)={ forprime(n=lb,ub, if(!(Mod(sum(k=0,n-1,2*polchebyshev(n-1,1,k/2)),n)==n-1),print(n))) } Chebyshev2(lb,ub)={ forcomposite(n=lb,ub, if((Mod(sum(k=0,n-1,2*polchebyshev(n-1,1,k/2)),n)==n-1),print(n))) }

论坛问题帖 | has_answers | Pedja | 2018-01-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018832 | MSE-2624073

The quotient of two palindromes

Besides the multiples of 10, are there infinitely many positive integers which are not the quotient of two palindromes? Is 12 such an exception?

论坛问题帖 | has_answers | Bernardo Recamán Santos | 2018-01-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018833 | MSE-2624685

A congruence involving Lucas polynomials

Can you provide a proof or a counterexample to the following claim : Let n be a natural number greater than two and let $L_{n}(x)$ be Lucas polynomial , then n is prime if and only if :$\displaystyle\sum_{k=0}^{n-1}L_{n-1}(k) \equiv -1 \pmod n$. You can run this test here . I was searching for counterexample using the following two Pari/GP codes : Lucas1(lb,ub)={ my(lucas(m,x)= p0=2; p1=x; l=2; while(l<=m, p=x*p1+p0; p0=p1; p1=p; l++); return(p)); forprime(n=lb,ub, if(!(Mod(sum(k=0,n-1,lucas(n-1,k)),n)==n-1),print(n))) } Lucas2(lb,ub)={ my(lucas(m,x)= p0=2; p1=x; l=2; while(l<=m, p=x*p1+p0; p0=p1; p1=p; l++); return(p)); forcomposite(n=lb,ub, if((Mod(sum(k=0,n-1,lucas(n-1,k)),n)==n-1),print(n))) }

论坛问题帖 | has_answers | Pedja | 2018-01-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018834 | MSE-2627914

Constant related to $f(n) = f(n-1) + \frac{1}{n f(n-1)}$

Consider the sequence f : $$f(1) = 1$$ $$ f(n) = f(n-1) + \frac{1}{n f(n-1)}$$ Now we have for large n : $$ f(n) = \sqrt {2 \ln(n)} + \frac{1}{19} + C + eps(n) $$ Where C is a constant and $eps(n)$ Goes to 0 for large n. Main question : Is $C = 0 $? If not What is Its value and do we have a closed form ?? Second question : How does eps behave ? Do we have a closed form ?

论坛问题帖 | has_answers | mick | 2018-01-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018835 | MSE-2634029

Primality test for repunits in bases ± 3

Can you provide a proof or a counterexample to the following claim : Let p be a prime greater than three , let $R_p(+3)=\frac{3^p-1}{2}$ and $R_p(-3)=\frac{3^p+1}{4}$. Let $S_i=S_{i-1}^3+3S_{i-1}$ with $S_0=36$, then $R_p(\pm 3)$ is prime iff $S_{p-1} \equiv 36 \pmod{R_p(\pm 3)}$. You can run this test here . I have tested this claim up to 10000 .

论坛问题帖 | has_answers | Pedja | 2018-02-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018836 | MSE-2636332

Bounding Numbers for $N>2$

Conjecture: "For any N number of points placed arbitrarily in the plane, the minimum number of lines need to bound each point in a separate polygon(which we will call B), is equal to: 2+ ceiling($n/2$),regardless of the placement of the N Points." Restrictions: N is an integer greater than 2 2.All of the points together may not be co-linear (Although subgroups of points may be co-linear) 3.You may only use straight lines 4.Lines drawn may not intersect your starting points The questions is, Can you prove or disprove the Conjecture above, and if it is false, What is the formula for finding B? Here's an Example using three points in an orderly arrangement: Example for $N=3$ In the example, $N=3$ and $B=4$, as we were able to bound each of the individual three points into heir own separate polygon using the minimum number of lines:4 Example of what was said in comment, n=10,b=8, formula holds Counter example to all co linear but one:

论坛问题帖 | no_answers_at_retrieval | Rdog60 | 2018-02-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018837 | MSE-2637546

What is the name of the conjecture related to what Matt Parker has called "The Square-Sum Problem"?

I read a book called Things to Make and Do in the Fourth Dimension, written by Australian mathe-matician and comedian, Matt Parker. In one part of the book, I remember him explaining about a conjecture such that for all $n\geqslant 89$, you can arrange the elements of the set $\{1, 2,\ldots,n\}$ in a certain way where each adjacent pair of elements sums to a squared number. For example, let $n = 17$. Al-though $17<89$, below is a good example to demonstrate what I mean: We have the set $\{1, 2, 3,\ldots, 17\}$ which can also be written as $\mathbb{N}_{\leqslant 17}$. How can we order each number in a certain way such that every adjacent pair of numbers in the ordered sequence add up to a squa-red number? Well, we order it like so: Let $S_{17} = [\ldots]$ be our sequence that orders the elements from $\{1, 2, 3,\ldots, 17\}$ in a certain way as mentioned in the foregoing then, $$S_{17} := \big[17, 8, 1, 15, 10, 6, 3, 13, 12, 4, 5, 11, 14, 2, 7, 9, 16\big].$$ Here, $17 + 8 = 5^2$, $8 + 1 = 3^2$, $1 + 15 = 4^2,\ldots$ The sequence in the sandbox above is a special case where it has 17 elements and begins with 17. Take the sequence S_{16} then this sequence ends with 16. In fact, it is exactly the same as S_{17} except it does not start from 17, but 8 instead. However, the sequence S_{18} does not exist, and there are many sequences with this property that do not exist. The conjecture is interesting because if true, it will prove that there are only finitely many sequences S_n that do not exist, also proving the contrapo-sitive. I did some research and it seems like this is true for $89\leqslant n \leqslant 300$ thus far, but I do not know the name of this conjecture. Does it even have a name? I also haven’t stumbled across any attempts of proving this conjecture. Can it be done? I guess that is two questions, then. Thank you in advance.

论坛问题帖 | has_answers | Mr Pie | 2018-02-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018838 | MSE-2641688

How large will the smallest counter-example probably be?

Here : Primes p of the form $10^{m}(2^{k}-1)+2^{k-1}-1$, where m is the number of decimal digits of $ 2^{k-1}$. it is conjectured that the number $$(2^k-1)\cdot 10^m+2^{k-1}-1$$ where m is the number of decimal digits of $2^{k-1}-1$ (See also the title), is never prime when it is of the form $7s+6$. Amazingly, primes with the other residues exists although the residue 6 occurs twice often than 1. Upto $k=131\ 000$, there is no counter-example, but probably a counter-example exists because the growth rate is roughly 4^k and 1 out of 9 $k's$ lead to a number of the form $7s+6$. How large will the smallest counterexample probably be ?

论坛问题帖 | no_answers_at_retrieval | Peter | 2018-02-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018839 | MSE-2641690

On integer sequences of the form $\sum_{n=1}^N (a(n))^2H_n^2,$ where H_n is the nth harmonic number: refute my conjecture and add yourself example

After I've read a problem posed by Édouard Lucas from the first paragraph of the biography from this Wikipedia's article I am trying to get different variations of this problem using number theoretic functions. After I did some experiments (I've calculated some terms using Wolfram Alpha online calculator) I posed a surprising conjecture. Conjecture. Here H_n denotes the nth harmonic number. For each integer $n\geq 1$, the integer $$\sum_{k=1}^n (k!)^2(H_k)^2\tag{1}$$ is a square-free integer (has no repeated primes in its factorization, see this Wikipedia; we get 1 as $n=1$). Question 1. Can you refute numerically my conjecture? Can you provide us some reasoning about if our sequence (1) has infinitely many terms being square-free integers? Many thanks. As you see I tried to get a variation of the problem posed by Lucas, since my proposal in the RHS of my problem is the sequence of square-free integers instead of the square numbers, and the terms of my finite series are the squares of the integers $k!\cdot H_k$. I tried more examples using number theoretic functions. With the purpose of making this post more interesting I ask to this commmunity about other possible variations, and in order to ask a concise question, I am going to limit the nature of the sequences that are terms in our summation, now the square of sequences of the form $$a(k)\cdot H_k,$$ where $a(k)$ is a sequence of rational numbers. The RHS of your statement/conjecture can be any sequence (I prefer some interesting sequence of figurate numbers, see this MathWorld). Question 2. Provide us yourself example of a conjecture or statement from a sequence $a(k)$ or rational numbers, such that for each $n>N$ (for a suitable positive integer in your proposal) N $$\sum_{k=1}^n (a(k)\cdot H_k)^2=\text{ a sequence of integers* }.\tag{2}$$ *I prefer figurate numbers, or famous sequences of integers like prime numbers... Many thanks.

论坛问题帖 | no_answers_at_retrieval | user243301 | 2018-02-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018840 | MSE-2643802

For prime numbers p_n it holds $m\equiv p_n^2\pmod {p_{n+1}}\implies$ m is a square

Conjecture about primes: For all $n\notin\{4,9,30\}$ it holds that if $m\equiv p_n^2\pmod {p_{n+1}}$ where $0 \leq m < p_{n+1}$, then m is a square. Tested for $n<1,000,000$. The only exceptions seems to be $p_4=7$, $p_9=23$ and $p_{30}=113$. Would like help to prove, explain or contradict this.

论坛问题帖 | has_answers | Lehs | 2018-02-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018841 | MSE-2644264

All even squares are congruent with a prime square $p_n^2\pmod {p_{n+1}}$

A conjecture about primes related to the question For prime numbers p_n it holds $m\equiv p_n^2\pmod {p_{n+1}}\implies$ m is a square For every even $m>0$ there is a $n\in \mathbb Z_+$ such that $m^2\equiv p_n^2\pmod {p_{n+1}}$, $0\leq

m^2<p_{n+1}$. Tested for $m\leq 152$. Are there some heuristics about this? Or counterexamples?

论坛问题帖 | has_answers | Lehs | 2018-02-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018842 | MSE-2645263

On the conjecture that, for every n, $\lfloor e^{\frac{p_{n^2}\#}{p_{n^2 + 1}}}\rfloor $ is a square number.

I was playing around with numbers and I discovered that $$\left\lfloor e^{\frac{p_4\#}{p_5}}\right\rfloor=\left\lfloor e^{\frac{210}{11}}\right\rfloor =13981^2,$$ with floor function $\lfloor x \rfloor := \max\{m\in\mathbb{Z} : m\leqslant x\}$; $ \ p_n$ denotes the n^{th} prime number; $ \ $and primorial $$p_n\#_c := \prod_{i=1}^n (p_i + c).\tag{$p_n\#_0 = p_n\#$}$$ Furthermore, $$\left\lfloor e^{\frac{p_1\#}{p_2}}\right\rfloor = 1^2.$$ I then made a conjecture with very few support, that $$\left\lfloor e^{\frac{p_{n^2}\#}{p_{n^2 + 1}}}\right\rfloor \tag1$$ is always a square number, say $k_n^{\ \ 2}$. Does somebody have a big enough computer to find the value of $(k_n)_{n\geqslant3}$? Or can my conjecture be proven/disproven with a pen and paper? And perhaps, to support the potential of this not being a coincidence, every divisor of 13981 takes the form $(13981 - 10x)$ for some $x\geqslant 0$. Maybe if this conjecture is true, $k_n > 1$ has this certain property? Edit 1: With some computational power, I found that $k_3^{\ \ 2}$ has $\approx 176,000$ digits. It turned out, however, that I was wrong, and really, $k_3^{ \ \ 2}$ has $\approx 3,340,970$ digits. Edit 2: Expressions equivalent to (1), avoiding e, for any base $b\in\mathbb{N}_{>1}$:$$\left\lfloor b^{\frac{p_{n^2}\#}{p_{n^2 + 1} \cdot Ln(b)}}\right\rfloor=k_n^{ \ \ 2}$$E.g. base $b=2$:$$\left\lfloor 2^{\frac{p_{n^2}\#}{p_{n^2 + 1} \cdot Ln(2)}}\right\rfloor=k_n^{ \ \ 2}$$E.g. base $b=n^2$:$$\left\lfloor (n^2)^{\frac{p_{n^2}\#}{p_{n^2 + 1} \cdot Ln(n^2)}}\right\rfloor=k_n^{ \ \ 2}$$

论坛问题帖 | has_answers | Mr Pie | 2018-02-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018843 | MSE-2645854

A conjecture concerning primes and perhaps prime gaps

I found out this conjecture which is tested for $m\leq 100$: For all natural numbers $n>\!30$, for all $\alpha\in\mathbb Z$ and all $\mu=0,\pm 1,\pm 2$: $m\equiv p_n^2+\mu p_n+\alpha\!\pmod{p_{n+1}}$ $\wedge\; m<p_{n+1}\iff$ $\exists k\in\mathbb Z_+:m=4k^2-2\mu k +\alpha$ Since this is a more general case than in the question For prime numbers p_n it holds $m\equiv p_n^2\pmod {p_{n+1}}\implies$ m is a square and that (due to an answer) was related to an unproved conjecture about prime gaps, I suspect that this conjecture is also related to prime gaps and is to deep to be proved. I would like to know: Can this conjecture relate to prime gaps? Are there counterexamples? Do someone have an intuition about the logic structure of the conjecture? Some context: I use bigz implemented in ANS-Forth with numbers and sets on stacks and all operations post-fix. First the case with $\alpha=\mu=0$: 1 100 :| n | n pnr@ ^2 n 1+ pnr@ umod dup sqr and ; build-set cr z. {0,1,4,16,36,64,100,144,196,324} ok DUP SQR AND mask the result from UMOD so all non square results are transformed to 0. SQR results in -1 or 0 depending on if the top of the number stack is a perfect square or not. Z. prints (and consumes) the set of the top of the set stack. 30 1000 :| n | n pnr@ ^2 n 1+ pnr@ umod dup sqr and ; build-set cr z. {0,4,16,36,64,100,144,196,256,324,400,484,576,676,784,900,1024,1156} ok Still exceptions indicated by 0 when $n\geq 30$. 31 1000 :| n | n pnr@ ^2 n 1+ pnr@ umod dup sqr and ; build-set cr z. {4,16,36,64,100,144,196,256,324,400,484,576,676,784,900,1024,1156} ok No exceptions when $n>31$. 31 10000 :| n | n pnr@ ^2 n 1+ pnr@ umod dup sqr and ; build-set cr z. {4,16,36,64,100,144,196,256,324,400,484,576,676,784,900,1024,1156,1296,1444,1600,1764,1936,2116,2304,2500,2704,2916,3136,3364,3600,3844,4096,5184} ok 31 50000 :| n | n pnr@ ^2 n 1+ pnr@ umod dup sqr and ; build-set cr z. {4,16,36,64,100,144,196,256,324,400,484,576,676,784,900,1024,1156,1296,1444,1600,1764,1936,2116,2304,2500,2704,2916,3136,3364,3600,3844,4096,4356,4624,4900,5184,5476,5776,6084,6400,6724,7056,7396,7744,8100,9216,9604,10000,12544,12996} ok The unbroken sequence of $4k^2$ values get longer for greater intervalls. Now an example with $\alpha=3$ and $\mu=1$: 31 10000 :| n | n pnr@ ^2 n pnr@ + 3 + n 1+ pnr@ umod ; build-set cr z. {5,15,33,59,93,135,185,243,309,383,465,555,653,759,873,995,1125,1263,1409,1563,1725,1895,2073,2259,2453,2655,2865,3083,3309,3543,3785,4035,5115} ok And an example with $\alpha=7$ and $\mu=-2$: 31 10000 :| n | n pnr@ ^2 n pnr@ 2* - 7 + n 1+ pnr@ umod ; build-set cr z. {15,31,55,87,127,175,231,295,367,447,535,631,735,847,967,1095,1231,1375,1527,1687,1855,2031,2215,2407,2607,2815,3031,3255,3487,3727,3975,4231,5335} ok https://forthmath.blogspot.se

论坛问题帖 | no_answers_at_retrieval | Lehs | 2018-02-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018844 | MSE-2647183

Is this function prime infinitely often?

Is it provable that there exist infinitely many primes of the form, $$f(n) = \left\{-1 + (p_{n+2} - 1)\prod_{i=2}^np_i : (p_n)_{n\in\mathbb{N}} = n^{\text{th}}\text{ prime number.}\right\}?$$ This is prime for $n\in\{1,2,3,4,6,7,8,12,13,15,21,48\}$. I have checked for $n\leqslant 54$. The reason I want to know if there is inf many, is because I conjecture that for every counter-example which we can call $c_1, c_2, c_3,\ldots$ then $\Omega(c_k) \leqslant 4$ such that we denote by $\Omega(c_k)$ the number of prime factors of c_k for which the function does not count the same prime factor twice. For instance, $\Omega(64) = 1$ because $64 = 2^6 = 2\times 2\times 2\times\cdots \times 2$, and since 2 is prime, we only count it once, so the function in this case is equal to 1 (and not 6). Is there a way to prove/disprove my conjecture? If there are infinitely many primes $f(n)$ then perhaps I could apply that to proving my conjecture. Thanks in advance.

论坛问题帖 | has_answers | Mr Pie | 2018-02-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018845 | MSE-2649633

On a conjecture and other numbers like 144.

Yesterday, I noticed that the following numbers were prime: $$67, 167, 467, 967, 1667.$$ I realised that the digits of these primes, apart from 67, were square numbers: $$0^2, 1^2, 2^2, 3^2, 4^2.$$ I then wanted to find values a_n such that when we concatenate $a_n^{\ \ 2}$ with 67, it is a prime number. This means, we join $a_n^{\ \ 2}$ with 67 to form a new number, and we will denote this as $a_n^{\ \ 2}\mid\mid67$. For example, 1 concatenated with 2 is denoted as $1\mid\mid 2$ which equals 12. Below is a sequence of all the values a_n that range from 0 to 160 inclusive: $0,1,2,3,4,7,9,10,15,16,18,23,24,28,30,31,33,42,43,45,48,53,55,61,65,68,69,72,76,77,87,88,93,94,96,98,105,106,109,110,112,113,116,117,119,122,125,130,132,133,138,140,144,145,147,151,155,159.$ Here, $a_1 = 0$, $a_2 = 1$, $a_3 = 2$, $\ldots$ Conjecture 1. I made a conjecture that for every number $b_n^{\ \ 2}\mid\mid 67$ that is not prime, it will either have 2 or 3 divisors such that b_n does not appear in this sequence. I did find a counter-example, namely 158. This was the first counter-example thus far such that, $$158^2\mid\mid 67 = 24964\mid\mid 67 = 2496467 = 17\times 19\times 59\times 131.\tag{4 divisors}$$ Conjecture 2. I also made another conjecture that for $n\geqslant 3$, if $a_n^{\ \ 2}\mid\mid 67$ is prime, and $(a_n + 1)^2\mid\mid 67$ is also prime, then not only will $(a_n + 2)^2\mid\mid 67$ not be prime, but it will be semi-prime, which is a prime composed of strictly two prime numbers. However, I also found another counter-example, namely 144. This was also the first counter-example to this conjecture, such that $144^2\mid\mid 67$ is prime, $(144^2 + 1)\mid\mid 67$ is also prime, but $$(144^2 + 2)\mid\mid 67 = 19\times 151\times 743.\tag{3 divisors}$$ This number is not semi-prime because it is composed of three prime numbers. I extended my first conjecture for which every number $b_n^{\ \ 2}\mid\mid 67$ that is not prime will have at most 4 prime divisors. But my question is, do there exist other counter-examples to the second conjecture? Particularly counter-examples that are square numbers, like 144? And is there a counter-example to my extended first conjecture? Thank you in advance. Edit: Conjecture 2 has been proven to be false, and lots of the counter-examples of conj$_2$ are square numbers. Here are the first few among the list: $$\begin{align} 1 = 1^2 &\quad 2916 = 54^2 &\quad 5776 = 76^2 &\quad \\ 837225 = 915^2 &\quad 3111696 = 1764^2 &\quad 10259209 = 3203^2\end{align}$$

论坛问题帖 | has_answers | Mr Pie | 2018-02-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018846 | MSE-2651638

On the conjecture that if $\,\forall\{x, y, z\}\subset \Bbb{Z^+}$ squarefree, and $\ x^3 + y^3 = z^2$, then $\gcd(3, x, y) = 1$.

I made a conjecture that if $\,\forall\{x, y, z\}\subset\mathbb{Z}^+$ with x, y, z squarefree (not raised to a power greater than 1; not a perfect power), and $$x^3 + y^3 = z^2,$$ then $\gcd(3, x, y) = 1$. I was able to prove my conjecture for all z prime. Proof: Let z be a prime number. Since $(x+y)^3 = x^3 + y^3 + 3xy(x+y),$ then $$\begin{align} x^3 + y^3 &= (x+y)^3 - 3xy(x+y) \\ &= (x+y)\big((x+y)^2-3xy\big) \\ &=(x+y)(x^2 + y^2 - xy).\tag1\end{align}$$ Therefore, we need integer solutions such that $(*)\,\, x + y = x^2 + y^2 - xy$ because $(1) = z^2$, follows that $$\begin{align} 0 &= x^2 + y^2 - xy - x - y \\ &= x(x - 1) + y(y - 1) - xy \\ \Leftrightarrow 3xy &= x(x-1) + y(y-1) + 2xy.\end{align}$$ Since $3xy$ is not a square number, this means that $x \neq 3y$, $y\neq 3x$ and $3\neq xy$. Therefore $\gcd(3, x, y) = 1$ for this case, but there is also another case where $(**)\,\, x + y = z$ and $x^2 + y^2 - xy = 1$. Of course $x + y\neq 1$ because $x, y\in\mathbb{Z}^+$, thus we need to consider statement $(**)$. $$\begin{align} x^3 + y^3 &= (x+y)^2 \\ \Leftrightarrow (x+y)(x^2 + y^2 - xy) &= (x+y)^2 \\ \Leftrightarrow x^2 + y^2 - xy &= x + y\tag*{$\unicode{x21af}$ (Contradiction)}\end{align}$$ Therefore, statement $(*)$ must always be true, i.e. $\gcd(3, x, y) = 1$.$\quad\quad\quad\quad\qquad\qquad\qquad\quad\Box$ Of course if z is not prime, then $x+y$ does not necessarily have to equal $x^2 + y^2 - xy$. Take $z = 6$ for example, then $z^2 = 6^2 =6\times 6$ but it also equals 9×4, etc. Could somebody prove/disprove my conjecture for all $z\in\mathbb{Z^+}$? Along with that, is my proof valid? Thank you in advance. Edit: The symbol $\unicode{x21af}$ (Harry Potter's Scar) is used to symbolise a contradiction in this case. Most people prefer $\Rightarrow\Leftarrow$, so if you prefer that instead, treat the Scar as if it were the arrows.

论坛问题帖 | has_answers | Mr Pie | 2018-02-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018847 | MSE-2654826

A trivial prime conjecture?

For all polynomials $q\in\mathbb Z[X]$ and all $n\in \mathbb N$ $$\exists k\in\mathbb N: q(p_n)\equiv q(-2k)\pmod{p_{n+1}}$$ Where p_n is the n-th prime. This conjecture is tested for random polynomials and seems to hold. Are there counterexamples? Is it a known conjecture? Is it trivial?

论坛问题帖 | has_answers | Lehs | 2018-02-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018848 | MSE-2655385

Compositeness tests for specific classes of generalized repunit numbers

Can you provide proofs or counterexamples for the following two claims: First claim Let $P_m(x)=2^{-m}\cdot \left(\left(x-\sqrt{x^2-4}\right)^{m}+\left(x+\sqrt{x^2-4}\right)^{m}\right)$. Let $N_p(b)=\frac{b^p+1}{b+1}$, where p is an odd prime and b is an odd natural number greater than one . CASE(1). $b \equiv 1,9 \pmod{12}$, or $b \equiv 3,7 \pmod{12}$ and $p \equiv 1 \pmod 4$, or $b\equiv 5 \pmod{12}$ and $p \equiv 1,7 \pmod{12}$, or $b \equiv 11 \pmod{12}$ and $p \equiv 1,11 \pmod{12}$. CASE(2). $b \equiv 3,7 \pmod{12}$ and $p \equiv 3 \pmod 4$, or $ b\equiv 5 \pmod{12}$ and $p \equiv 5,11 \pmod{12}$, or $b \equiv 11 \pmod{12}$ and $p \equiv 5,7 \pmod{12}$. Let $S_i=P_b(S_{i-1})$ with $S_0=P_b(4)$. Suppose $N_p(b)$ is prime , then : $\bullet$ $S_{p-1} \equiv P_{b}(4) \pmod {N_p(b)}$ if Case(1) holds ; $\bullet$ $S_{p-1} \equiv P_{b+2}(4) \pmod {N_p(b)}$ if Case(2) holds ; You can run this test here . Second claim Let $P_m(x)=2^{-m}\cdot \left(\left(x-\sqrt{x^2-4}\right)^{m}+\left(x+\sqrt{x^2-4}\right)^{m}\right)$. Let $M_p(a)=\frac{a^p-1}{a-1}$, where p is an odd prime and a is an odd natural number greater than one . CASE(1). $a \equiv 3,11 \pmod{12}$, or $a \equiv 5,9 \pmod{12}$ and

$p \equiv 1 \pmod 4$, or $a\equiv 7 \pmod{12}$ and $p \equiv 1,7 \pmod{12}$, or $a \equiv 1 \pmod{12}$ and $p \equiv 1,11 \pmod{12}$. CASE(2). $a \equiv 5,9 \pmod{12}$ and $p \equiv 3 \pmod 4$, or $ a\equiv 7 \pmod{12}$ and $p \equiv 5,11 \pmod{12}$, or $a \equiv 1 \pmod{12}$ and $p \equiv 5,7 \pmod{12}$. Let $S_i=P_a(S_{i-1})$ with $S_0=P_a(4)$. Suppose $M_p(a)$ is prime , then : $\bullet$ $S_{p-1} \equiv P_{a}(4) \pmod {M_p(a)}$ if Case(1) holds ; $\bullet$ $S_{p-1} \equiv P_{a-2}(4) \pmod {M_p(a)}$ if Case(2) holds ; You can run this test here . I have tested these claims for many random values from this list . EDIT A command line program that implements these two tests can be found here .

论坛问题帖 | has_answers | Pedja | 2018-02-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018849 | MSE-2660281

Show $\lim_{n\to \infty}{|A_{n}|+|A_{n-2}| \above 1.5pt |A_{n-1}|}=2$

Question: Let $A(n)$ be a finite square $n \times n$ matrix with entries $a_{ij}=1$ if $i+j$ is a prime; otherwise equals to 0. I write $|A(n)|$ to count the number of 1's in $A(n)$. Is it true, possibly trivially, and how can we show that $$\lim_{n\to \infty}{|A_{n}|+|A_{n-2}| \above 1.5pt |A_{n-1}|}=2$$ The sequence of $A(n)$'s can be found here in Sloan. The graph below illustrates the limit. I believe I have shown the question is true, conditionally using a parity argument along with Bertrand's Postulate. Solution: It is easy to verify that $A(n)$ has exactly $2n-1$ skew diagonal. Two small lemmas. $\underline{\text{lemma 1}}:$ Observe $a_{ii}=1$ if and only if $i=1$. Consequently a_{11} is the only entry on the main diagonal of $A(n)$ equal to 1. Note $A(n)$ is symmetric since addition is commutative. So the count of 1's below the main diagonal equals the counts of 1's above the main diagonal and the two counts combined form an even number. If we remember to add the sole 1 for the entry a_{11} then there are an odd number of entries in $A(n)$. In particular $|A(n)|$ is always odd. $\underline{\text{lemma 2}}:$ As mentioned earlier every matrix $A(n)$ has $2n-1$ skew diagonals. Let S_n be the n-th skew diagonal of $A(n)$. A skew analogue of Bertrand's Postulate shows that at least one of the skews between S_n and S_{2n-2} has all of its entries equal to 1. Consequently for $n>2$ we have that there is always a skew diagonal with entries equal to 1 below the main skew diagonal. In particular $\{A(n)\}_{n=1}^\infty$ is a monotonically increasing sequence of odd integers. Lemma's 1 and 2 control the growth of $A(n)$ as $n \to \infty.$ By parity check alone as $A(n)$ grows to $A(n+1)$ the count of new 1's must be an nonzero even number. Because of this we have that $|A_n|=|A_{n-1}|+2j$ for some $j\in\mathbb{N}$, $j\neq0$. So we have the following from direct substitution \begin{align} {|A_{n}|+|A_{n-2}| \above 1.5pt|A_{n-1}|}&={\left(|A_{n-1}|+2j\right)+\left(|A_{n-1}|-2k\right)\above 1.5pt |A_{n-1}|}\\ &=2\cdot{|A_{n-1}|+j-k\above 1.5pt |A_{n-1}|}\\ &=2\cdot\left(1+{j-k\above 1.5pt |A_{n-1}|}\right)\\ \end{align} Taking limits we have \begin{align} \lim_{n\to \infty}2\cdot\left(1+{j-k\above 1.5pt |A_{n-1}|}\right)&=2+\lim_{n\to \infty}\left({j-k\above 1.5pt |A_{n-1}|}\right)\\ &=2+0\\ &=2 \end{align} This completes the proof.

论坛问题帖 | no_answers_at_retrieval | Anthony | 2018-02-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018850 | MSE-2660289

What is the difference between intuition and conjecture in mathematics?

I don't know any difference between intuition and conjecture. In my opinion: A conjecture is a mathematical statement that has not yet been rigorously proved. Conjectures arise when one notices a pattern that holds true for many cases. Conjectures must be proved for the mathematical observation to be fully accepted. When a conjecture is rigorously proved, it becomes a theorem, and intuition is prediction of any thing and how it will be in the future. Question: What is the difference between intuition and conjecture in mathematics ?

论坛问题帖 | has_answers | zeraoulia rafik | 2018-02-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018851 | MSE-2663327

About a very interesting constant g.

I was playing around with numbers like I always do, and I came up with a conjecture: $$\ln(e^{\pi + n} - e^{\pi + n - 1} - \ldots - e^{\pi + 1} - e^{\pi}) - \frac{\pi}{\sqrt{2}} \approx n$$ such that $n\in\mathbb{N}$. I then realised that the larger the value of n, the more accurate the LHS was getting closer to the RHS... ...and then it stopped. Strangely enough, it appeared that $$\ln\left(e^{\pi + n} - \sum_{i = 0}^{n-1}e^{\pi + i}\right) - \frac{\pi}{\sqrt{2}} \sim n + k$$ such that, as far as I have tested, $k \approx 0.0479330616936374724102$ for all $n\leqslant 61$. Subtracting this constant k from both sides, I see that $$\frac{\pi}{\sqrt{2}} + k \approx 2.26937453077282059591 8141.$$ Now by letting $g = \dfrac{\pi}{\sqrt{2}} + k$, is there some algebraic formula (like an infinite series) to find k? If there is, and we find it, we can find the value of g, and we would be able to write down that, $$\ln\left(e^{\pi + n} - \sum_{i=0}^{n-1}e^{\pi + i}\right) - g\sim n$$ with whatever g is equal to (algebraically speaking). For now, g is just a special constant. Thanks in advance.

论坛问题帖 | has_answers | Mr Pie | 2018-02-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018852 | MSE-2668104

A sum of sum involving binomial coefficent.(my conjecture)

While using MATLAB I found out that $\displaystyle\sum_{k=0}^{n}\frac{(\sum_{r=\max (0,k+m-n)}^{\min (k,m)}(-1)^{r}\binom{n-m}{k-r}\binom{m}{r})^{2}}{\binom{n}{k}} =\frac{2^{n}}{\binom{n}{m}} $ hold for all $m\leq n\leq 100\ $ and I proved that the equality hold for all $m=0,1,n-1,n\ $ but I couldn't prove it in general (or disprove). Any help is appreciated! Thanks in advance.

论坛问题帖 | has_answers | LightKnight | 2018-02-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018853 | MSE-2675905

Equiangular lines in $\mathbb{C}^d$.

Zauners' conjecture regarding equiangular lines for any $\mathbb{C}^{d}$, holds for many dimesnions, like 17 or more(43, 31, and so on). This conjecture gives the number of equiangular lines in a given complex space $\mathbb{C}^d$. A similar result shows that for Euclidean spaces, given an angle, the number of equiangular lines can be found, result here this result is for the case of a given angle, finding the number of equiangular lines like in Zauners' conjecture (over $\mathbb{C}^{d}$), but for a given angle. Is there a number(it could be a lower/upper bound, or like Zauners' conjecture - exact), but with the constraint that it is for a given angle over $\mathbb{C}^{d}$?

论坛问题帖 | no_answers_at_retrieval | user3483902 | 2018-03-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018854 | MSE-2676103

If $\frac{σ(x)}{x}=\frac{p+2}p$ where p is an odd prime, does it follow that x is an odd square?

(This question is related to the latest comment underneath this MSE question.) Let $$\sigma(x)=\sum_{d \mid x}{d}$$ be the sum of the divisors of x. Denote the abundancy index of x by $$I(x)=\frac{\sigma(x)}{x}.$$ Here is my question: If $I(x)=\dfrac{p+2}{p}$, where p is an odd prime, does it follow that x is an odd square? From the equation $$p\sigma(x)=(p+2)x$$ I can only prove that x is an odd square when x is odd. But what about if x is even? That is, if $I(x) = \dfrac{p+2}{p}$, where p is an odd prime, must it necessarily be the case that x is odd?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2018-03-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018855 | MSE-2683513

An extension of prime number theorem

Suppose $r:\Bbb N\to (0,1)$ is a function given by $r(n)$ is obtained as put a point at the beginning of n instance $r(34880)=0.34880$ and $\Bbb P$ is the set prime numbers. Theorem: For each subinterval (a,b) of $[0.1,1),\,\exists m\in \Bbb N$ that $\forall k\in \Bbb N$ with $k\ge m$ then $\exists t\in (a,b)$ that $t\cdot 10^k\in \Bbb P$. ($\color{fuchsia}{\text{proof}}$ at my previous account and it is shown $\color{fuchsia}{\text{here}}$ too.) Corollary: $r(\Bbb P)$ is dense in the $[0.1,1]$. $$\\$$ and let $\forall n\in\Bbb N,$ and for each subinterval (a,b) of $[0.1,1),$ that $a\neq b,$ $\begin{cases} U_{(a,b)}:=\{n\in\Bbb N\mid a\le r(n)\le b\},\\ \\V_{(a,b)}:=\{p\in\Bbb P\mid a\le r(p)\le b\},\\ \\U_{(a,b),n}:=\{m\in U_{(a,b)}\mid m\le n\},\\ \\V_{(a,b),n}:=\{m\in V_{(a,b)}\mid m\le n\},\\ \\w_{(a,b),n}:=(\#U_{(a,b),n})^{-1}\cdot\#V_{(a,b),n}\cdot\log n,\\ \\w_{(a,b)}:=\lim _{n\to\infty} w_{(a,b),n}\end{cases}$ $$\\$$ Now regarding to Prime number theorem and above corollary I offer following conjecture so please guide me about possibility of this conjecture: Conjecture: $\forall (a,b)\subset [0.1,1),\,w_{(a,b)}=0.9^{-1}\cdot (b-a)$. Thanks in advance!

论坛问题帖 | has_answers | user538968 | 2018-03-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018856 | MSE-2692080

Conjecture over this series, and its generalisation

PART I The following series, according to W. Mathematica, does converge to $$\sum_{k = 1}^{+\infty} \frac{e^{-k}}{k^k \sqrt{k}} = 0.3929049383779132(...)$$ The previous result can be written in terms of elementary numbers plus the Euler constant, as follows: $$0.3929049383779132 \approx \frac{-92-95 e+86 e^2}{2 \left(130-4 e+33 e^2\right)}$$ Question Is that conjecture true? Probably it's a lack of mine but I cannot manage to make W. Mathematica to spit out more digits of the previous number. PART II The previous series was actually a special case for $x = 1$ of the more general series: $$\sum_{k = 1}^{+\infty} \frac{x^k e^{-k}}{k^k \sqrt{kx}}$$ Question: Is there a close form for this? I tried with many values of x and the series always gets a numerical result, but Mathematical cannot give me a close form. I am not assuming a priori it does exist, but many times people found out close forms whereas software could not.

论坛问题帖 | has_answers | user266764 | 2018-03-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018857 | MSE-2698479

If $q^k n^2$ is an odd perfect number with Euler prime q, does this equation imply that $k=1$?

Let $\sigma(x)$ be the sum of the divisors of x. Denote the deficiency of x by $D(x) := 2x - \sigma(x)$, and the sum of the aliquot divisors of x by $s(x) := \sigma(x) - x$. Here is my question: If $q^k n^2$ is an odd perfect number with Euler prime q, does this equation imply that $k=1$? $$D(q^k)D(n^2)=2s(q^k)s(n^2)$$ I only know that $k=1$ is true if and only if (one of) the following conditions hold: (1) $\sigma(n^2)/q \mid n^2$ (2) $D(n^2) \mid n^2$ (3) $\gcd(n^2, \sigma(n^2)) = D(n^2)$ Source of Equation From the fundamental equation $$\frac{\sigma(n^2)}{q^k} = \frac{2n^2}{\sigma(q^k)} = \gcd(n^2, \sigma(n^2))$$ we obtain $$\frac{2n^2 - \sigma(n^2)}{\sigma(q^k) - q^k} = \frac{D(n^2)}{s(q^k)} = \gcd(n^2, \sigma(n^2))$$ and $$\frac{\sigma(n^2) - n^2}{\frac{2q^k - \sigma(q^k)}{2}} = \frac{2s(n^2)}{D(q^k)} = \gcd(n^2, \sigma(n^2)),$$ using the identity $$\frac{A}{B}=\frac{C}{D}=\frac{C-A}{D-B}.$$ Reference Conditions Equivalent to the Descartes-Frenicle-Sorli Conjecture on Odd Perfect Numbers

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2018-03-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018858 | MSE-2717456

A Simple Calculus view on Fermat's Last Theorem

My question (more of a hypothesis really) is basically this: If a function $f(x)$ is defined such that $f'(x)$ is not constant and never the same for any 2 values of x. Then there do not exist positive integers a,b,c and $a\le b\le c$ such that, $$\int_0^{a} f(x)dx = \int_b^{c} f(x)dx\tag1$$ Taking $f(x)=x^n$ then for $n \gt 1$, (1) becomes Fermat's Last Theorem,

while for $n=1$, (1) becomes $a^2 + b^2 = c^2$ Maybe this has something to do with the "curviness" (I am 16 so please forgive me for non-technical language) of the graphs of x^n since for $n=1$ the graph is linear and for $n>1$ the graph is curvy. My school teachers are not talking to me about this because "it is out of syllabus", so I thought maybe this community could help.

论坛问题帖 | has_answers | Doctorwho2311 | 2018-04-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018859 | MSE-2720878

A conjectured vanishing sum involving the Thue-Morse sequence

Let's define t_n by the recurrence $$t_0 = 1, \quad t_n = (-1)^n \, t_{\lfloor n/2\rfloor}.\tag1$$ It is easy to see that $|t_n|=1$, and the signs follow the same pattern as the Thue-Morse sequence: $$1,\,-1,\,-1,\,1,\,-1,\,1,\,1,\,-1,\,-1,\,1,\,1,\,1,\,-1,\,1,\,-1,\,-1,\,1,\,1,\,...\tag2$$ (see this question for an example of a non-recursive formula for t_n). Now, define: $$\mathcal{S}(m, n)=\sum_{k=0}^{2^n-1}t_k\,k^m, \quad \color{gray}{m,\,n\in\mathbb{Z}^+.}\tag3$$ Computing some values of this sum immediately leads to the following conjecture: $$\text{For all }\, n>m,\,\,\mathcal{S}(m, n)=0.\tag{$\diamond$}$$ (e.g. for $m=2$ and $n=3$ we have $0-1-4+9-16+25+36-49=0$). How can we prove it? Is it a known result?

论坛问题帖 | no_answers_at_retrieval | Vladimir Reshetnikov | 2018-04-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018860 | MSE-2722327

A limit involving the Thue-Morse sequence

(it is a follow-up on my previous question) Let's define t_n by the recurrence $$t_0 = 1, \quad t_n = (-1)^n \, t_{\lfloor n/2\rfloor}.\tag1$$ It is easy to see that $|t_n|=1$, and the signs follow the same pattern as the Thue-Morse sequence: $$1,\,-1,\,-1,\,1,\,1,\,-1,\,1,\,1,\,1,\,-1,\,-1,\,1,\,1,\,1,\,-1,\,1,\,1,\,-1,\,-1,\,1,\,1,\,...\tag2$$ (see this question for an example of a non-recursive formula for t_n). Now, let: $$\mathcal{L}(z)=\lim_{n\to\infty}\,\sum_{k=1}^{2^n-1}t_k\,k^z.\tag3$$ I conjecture that the following propositions hold: $\color{gray}{\text{(a)}}$ The limit $\mathcal{L}(z)$ exists for all $z\in\mathbb C$. $\color{gray}{\text{(b)}}$ $\mathcal{L}(z)$ is an entire function of z. $\color{gray}{\text{(c)}}$ $\mathcal{L}(z)=0$ if and only if $z\in\mathbb Z^+$. Can we prove these conjectures? Can we find a different representation of $\mathcal{L}(z)$? Does this function have any interesting properties? Update: The "if" part of the conjecture $\color{gray}{\text{(c)}}$ is certainly true. I am now pretty sure the "only if" part fails at some points on the imaginary axis (e.g. near $z\approx i\,4.53236...$; is it exactly $i\,\pi/\ln2$?). I am still wondering if there are any zeros except positive integers and off the imaginary axis. Update: I found a very relevant and interesting paper (preprint) by Giedrius Alkauskas, Dirichlet series associated with Thue-Morse sequence$^{[1]}$$\!^{[2]}$ (be aware that the author claims$^{[3]}$ some results in it may be incorrect). Here is my attempt to plot the function for a range of real arguments:

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2018-04-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018861 | MSE-2727194

A few conjectured limits of products involving the Thue-Morse sequence

(related to my previous questions$^{[1]}$$\!^{[2]}$) Let's define the signed Thue-Morse sequence t_n by the recurrence $$t_0 = 1, \quad t_n = (-1)^n \, t_{\lfloor n/2\rfloor},\tag1$$ or by the generating function $$\sum_{n=0}^\infty t_n \, x^n=\prod_{n=0}^\infty\left(1-x^{2^n}\right).\tag{$1^\prime$}$$ It seems that the following conjectures hold: $$\lim_{n\to\infty}\prod_{k=0}^{2^n-1}\left(k+\tfrac12\right)^{t_k}\stackrel{\color{gray}?}=\frac12\tag2$$ $$\lim_{n\to\infty}\prod_{k=0}^{2^n-1}\left(k+1\right)^{t_k}\stackrel{\color{gray}?}=\frac1{\sqrt2}\tag3$$ $$\lim_{n\to\infty}\prod_{k=0}^{2^n-1}\left(k+1\right)^{(-1)^k\,t_k}\stackrel{\color{gray}?}=\frac1{2\sqrt2}\tag4$$ How can we prove these? Are there any other limits of products similar to these?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2018-04-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018862 | MSE-2730611

A congruence involving Fibonacci polynomials

Can you provide a proof or a counterexample to the following claim : Let n be a natural number greater than one and let $ F_{n}(x)$ be Fibonacci polynomial , then n is prime if and only if : $ \displaystyle\sum_{k=0}^{n-1}F_{n}(k) \equiv -1 \pmod n$. You can run this test here .

论坛问题帖 | has_answers | Pedja | 2018-04-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018863 | MSE-2733783

Does there exist an odd natural number $N = xy$ satisfying $D(x)D(y) = 2s(x)s(y)$?

Preamble: I apologize in advance if what I am asking for in this question, I could get an answer easily so myself, for example by coding a short Mathematica script. It is just that I have not yet studied the rudiments of Mathematica (or even Python) scripting, so I am not there yet. This question is a follow-up to this earlier MSE question. Let $\sigma(z)$ be the sum of the divisors of $z \in \mathbb{N}$. Denote the deficiency of z by $D(z) := 2z - \sigma(z)$ and the sum of the aliquot parts of z by $s(z) := \sigma(z) - z$. Here is my question: Does there exist an odd natural number $N = xy$ satisfying $D(x)D(y) = 2s(x)s(y)$?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2018-04-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018864 | MSE-2734161

Is this tiny conjecture true concerning α true?

As part of a problem that i am trying to solve i would like to know if the following conjecture is true. Conjecture. Given a non-zero vector α in a finite dimensional vector space W there exist a basis for W say $w_1,w_2,...,w_m$ such that $\alpha = a_1w_1+a_2w_2+\cdot\cdot\cdot+a_mw_m$ where $a_j\neq 0$ $\forall j\in\{1,2,...,m\}$ and $m = \dim W$. The Field of Scalars is either $\mathbf{R}$ or $\mathbf{C}$.

论坛问题帖 | has_answers | atifcppprogrammer | 2018-04-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018865 | MSE-2747063

An inequality involving three consecutive primes

Can you provide a proof or a counterexample to the following claim : Let p,q,r be three consecutive prime numbers such that $p\ge 11 $ and $p<q<r$, then $\frac{1}{p^2}< \frac{1}{q^2} + \frac{1}{r^2}$. I have tested this claim up to 10^{10} . For $p>5$ we get $\pi(2p)-\pi(p) \ge 2$, a result by Ramanujan . This means that $q<2p$ and $r<2p$, so $\frac{1}{2p}<\frac{1}{q}$ and $\frac{1}{2p}<\frac{1}{r}$ which implies $\frac{1}{p} < \frac{1}{q} + \frac{1}{r}$. If we square both sides of inequality we get $\frac{1}{p^2} < \frac{1}{q^2} + \frac{2}{qr} + \frac{1}{r^2}$. Now , I don't know how to rule out term $\frac{2}{qr}$.

论坛问题帖 | has_answers | Pedja | 2018-04-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018866 | MSE-2750429

"Schäffer's conjecture" on equation $1^k+2^k+\cdots+x^k=y^n$

In 1956 J. J. Schäffer proved that the equation $$1^k+2^k+\cdots+x^k=y^n$$ for fixed integers $k\ge1,n\ge2$ has only finitely many solutions in positive integers unless $(k,n)\in\{(1,2),(3,2),(3,4),(5,2)\}$ [see Theorem 2, p. 27 in the file's numbering of pages]. Furthermore, he conjectured the equation has only trivial solution $x=y=1$ except $(k,n,x,y)=(2,2,24,70)$ (and excluding those four pairs of (k,n)). I was able to find partial results, that the conjecture is true for $k\le11$ [see this] $n=2,k\le58$ [mentioned there as well] $k<170$ odd and even n [see this one] The newest paper (3.) was published in 2007 and I didn't find newer articles. So here is my question, is Schäffer's conjecture still an open problem?

论坛问题帖 | no_answers_at_retrieval | byk7 | 2018-04-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018867 | MSE-2751545

A conjecture about $8n+6$

Can you provide a counterexample to the claim given below ? Inspired by Euler's conjecture about $8n+3$ (see page 4 of this note) I have formulated the following claim : For any nonnegative integer n , $8n+6$ can be represented as a sum : $$8n+6 = a^2 +2p$$, where a is a nonnegative integer , and p is a prime . I have tested this claim for all n up to $2 \cdot 10^6 $.

论坛问题帖 | has_answers | Pedja | 2018-04-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018868 | MSE-2759813

A question on entropy of natural numbers

This question is more of a conjecture, but I am not posting it to mathoverflow, since I am not a professional mathematician and I do not know if it is research related: Let $n=p^{a} s$, p be the smallest prime dividing n, $v_p(n) = a$, $n\equiv 1(2)$,$\sigma(n) = 2n$. Then we must have $p>2$. Let $m=2^a s $. Then we have $$\sigma(m)= \sigma(2^a)\sigma(s)= \frac{\sigma(2^a)\sigma(n)}{\sigma(p^a)}$$. Since $2<p$ we have $H(2^a) < H(p^a)$ where H is the entropy of n defined here: Entropy of a natural number. It follows that: $$H(n) = H(p^a s) = H(p^a)+H(s) > H(2^a) + H(s) = H(m) = \log(\sigma(m))-\frac{1}{\sigma(m)}\sum_{d|m}{d \log(d)} = \log(\frac{\sigma(2^a)\sigma(n)}{\sigma(p^a)})-\frac{1}{\sigma(m)}E(m)$$, where $E(k) = \sum_{d|k} {d \log(d)}$. Since $\log(x) \ge 1 - 1/x$ we have $$ H(n) > H(m) = \log(\sigma(n))+\log(\frac{\sigma(2^a)}{\sigma(p^a)})-\frac{E(m)}{\sigma(m)}\ge\log(\sigma(n))+1-\frac{\sigma(p^a)}{\sigma(2^a)}-\frac{E(m)}{\sigma(m)}$$ which is equal to: $$\log(\sigma(n))+1-\frac{\sigma(n)}{\sigma(m)}-\frac{E(m)}{\sigma(m)}$$ If we can prove that, the last quantity is greater than or equal to $\log(\sigma(n))-\frac{E(n)}{\sigma(n)}= H(n)$, we would get the contradiction: $$H(n)>H(m) \ge H(n)$$ So we have to prove that: $$1 - \frac{\sigma(n)}{\sigma(m)}-\frac{E(m)}{\sigma(m)} \ge - \frac{E(n)}{\sigma(n)}$$ We have the following identities: $$E(m) = \sigma(2^a)E(s)+\sigma(s)E(2^a)$$ $$E(n) = \sigma(p^a)E(s)+\sigma(s)E(p^a)$$ Furthermore one can show that: $$E(p^a) = p \log(p) \frac{p^{a+1}-1-(a+1)p^a(p-1)}{(p-1)^2}$$ which can be applied to compute $E(2^a)$. So my question is if somebody can help me prove the inequality above or show an "example" where it fails, if this is not too much to ask.

论坛问题帖 | has_answers | user276611 | 2018-04-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018869 | MSE-2761488

Where is the mistake in the argument? (And two conjectures)

Let n be a natural number. $\Omega = \{ d : d | n \}$, For $A\subset \Omega$ define the probability for A as : $$P(A) = \frac{1}{\sigma(n)}\sum_{d\in A}{d}$$ Consider the set $B = \{d \in \Omega |d \equiv 0 (2) \}$. Let $a = v_2(n)$. Then (first conjecture) $$P(B) = \frac{2^{a+1}-2}{2^{a+1}-1}$$ We have $E(d) = \sum_{d|n} {d \cdot P(d)} = \sum_{d|n} { d \cdot \frac{d}{\sigma(n)}} = \frac{\sigma_2(n)}{\sigma(n)}$ Second conjecture: $|B| = a \cdot b$, where $n = 2^a \cdot p_1^{a_1}\cdots p_r^{a_r}$ is the factorization of n, with a possibly $=0$ and $b = (a_1+1)\dots(a_r+1)$. Consider $Z = \sum_{d \in B} {d}$ and consider the random variable: $Y_d = 1$ if $d \equiv 0 (2) $, otherwise $= 0$. Then $Z = \sum_{d|n} Y_d d$ is a random variable. Then on the one hand we have: $E(Z) = \sum_{d \in B}{E(d)} = \sum_{d \in B}{\frac{\sigma_2(n)}{\sigma(n)}} = a b \frac{\sigma_2(n)}{\sigma(n)}$ On the other hand we have $P(B) = \frac{Z}{\sigma(n)}$ hence solving for Z we get: $$Z = \sum_{d|n}{d} \cdot \frac{2^{a+1}-2}{2^{a+1}-1}$$ From this it follows that: $$E(Z) =

\frac{2^{a+1}-2}{2^{a+1}-1} \sum_{d|n}{E(d)} = \frac{2^{a+1}-2}{2^{a+1}-1} \sum_{d|n}{\frac{\sigma_2(n)}{\sigma(n)}} = \frac{2^{a+1}-2}{2^{a+1}-1} \tau(n) \cdot \frac{\sigma_2(n)}{\sigma(n)}$$ Hence we get: $$a b \frac{\sigma_2(n)}{\sigma(n)} = \frac{2^{a+1}-2}{2^{a+1}-1} \tau(n) \cdot \frac{\sigma_2(n)}{\sigma(n)}$$ from which it follows: $$ab = \tau(n) \cdot \frac{2^{a+1}-2}{2^{a+1}-1}$$ But for example for $n=6$ this is wrong, so where is the mistake in the argument? If you happen to have a proof for one of the conjectures, that would also be fine.

论坛问题帖 | has_answers | user276611 | 2018-05-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018870 | MSE-2763787

A conjecture on bounded complex partial sums

A friend of mine has made the following conjecture, but we don't know how to prove it. Let $(a_n)_{n\in \mathbb {N}}$ be a strictly increasing sequence of natural numbers. Suppose that for every complex $z\neq 1 $ with $|z|=1$, there exists a constant $C_z>0 $ verifying $$\forall N\geq 0, \qquad \left|\sum_{i=0}^N z^{a_i}\right| <C_z$$ Then the sequence $(a_n) $ contains all but finitely many natural numbers. For exemple, if $a_n=2n$, the property is not verified with $z=-1$. Can you find a proof or a counterexample?

论坛问题帖 | has_answers | Friedrich | 2018-05-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018871 | MSE-2764901

A conjecture concerning the number of divisors and the sum of divisors.

I stumbled upon the following conjecture: Let n be a positive integer. Let $\sigma\left(n\right)$ be the sum of all (positive) divisors of n, and let $\tau\left(n\right)$ be the number of these divisors. Then, $$\tau(n)+\sigma(n)\equiv 1 \mod 2 \iff n = 2m^2 \text{ for some integer } m .$$ Does anybody have an idea on how to prove this, maybe in parts? Or maybe someone has a counterexample?

论坛问题帖 | has_answers | user276611 | 2018-05-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018872 | MSE-2765028

Can the following argument be pushed to a full proof that $(p + 2)/p$ is an outlaw if p is an odd prime?

This is related to this earlier MSE question. Let $\sigma(x)$ be the sum of the divisors of x, and denote the abundancy index of x by $I(x):=\sigma(x)/x$. If the equation $I(a) = b/c$ has no solution $a \in \mathbb{N}$, then the rational number b/c is said to be an abundancy outlaw. In what follows, let p be an odd prime. In the earlier question, it is shown that: (a) If $I(n) = (p+2)/p$, then n is an odd square. (b) If $I(n) = (p+2)/p$, then $p \mid n$. So now suppose to the contrary that $I(n) = (p+2)/p$. From (a) and (b), since p is a prime number, then $p^2 \mid n$. It follows that $$\frac{p^2 + p + 1}{p^2} = I(p^2) \leq I(n) = \frac{p+2}{p},$$ whence there is still no contradiction. From the divisibility constraint $p^2 \mid n$, we have that $p^2 \leq n$. We claim that $p^2 \neq n$. Suppose to the contrary that $p^2 = n$. Then we have $$\frac{\sigma(n)}{n} = \frac{p+2}{p} = \frac{p(p+2)}{p^2}$$ so that $$p^2 + p + 1 = \sigma(p^2) = \sigma(n) = p^2 + 2p,$$ which contradicts the fact that p is an odd prime. This implies that $p < \sqrt{n}$, which further means that $$\gcd(n, \sigma(n)) = \frac{n}{p} > \sqrt{n} > {10}^8$$ where the lower bound $n > {10}^{16}$ is due to Richard F. Ryan, "Results concerning uniqueness for $\sigma(x)/x = \sigma(p^n q^m)/(p^n q^m)$ and related topics, International Math. J., 2002 , V2#5pp497-514". Here is my question: Can the preceding argument be pushed to a full proof that $(p + 2)/p$ is an outlaw if p is an odd prime?

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2018-05-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018873 | MSE-2768683

On a conjecture that if $2\left(p_n+p_{n-1}\right)\equiv -1\pmod {n^2}$ then $n=11$ only.

Conjecture: We denote by p_n the n^{th} prime number. If $2\left(p_n+p_{n-1}\right)\equiv -1\pmod {n^2}$, then $n=11$ only. How must one be able to prove / disprove this supposed claim? I had come up with this conjecture, but am a little stuck in trying to prove it. Attempt: I know that the congruence can be similarly written as, $2\left(p_n+p_{n-1}\right) = mn^2 - 1$ for some $m\in\mathbb{Z}$, the set of all integers. However, I will consider the case where $m=1$, or where $2\left(p_n+p_{n-1}\right)$ $ = (n+1)(n-1)$. Of course now, we find that n must be odd and greater than 1. Replace n with n_k, namely, the k^th odd number. Then, $$\frac{n_k+1}{2} = k \ \land \ \frac{n_k-1}{2} = k-1.\tag{$\land$ is read as and}$$ Therefore, $p_{n_k}+p_{n_k-1} = k(n_k-1) = (k-1)(n_k+1)$. This means that $\left\{k,k-1\right\}\mid p_{n_k} + p_{n_k-1}$. So, $p_{n_k}+p_{n_k-1}$ has to be squarefree, leaving out primes 3 and 5 since $5+3=8=2^3$ and also 17 and 19 since $19+17=36=6^2$. I do not know where to go from here. Am I taking the correct approach? Is there anything important I should know abut prime numbers (e.g. any lemmas and such) that will be useful in (dis)proving my conjecture? Hints are mostly appreciated, but you can provide a full and disclosed answer if you wish. Thank you in advance.

论坛问题帖 | has_answers | Mr Pie | 2018-05-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018874 | MSE-2770339

Is 387420547 another counter example to John H. Conway's Problem 5?

John H. Conway is a mathematician who created these five $\$1000$ problems. They are just problems that he discovered himself and that relate to mathematics, mainly on a recreational level. Anyone to have "solved" one of the problems (by "solved", I mean provide a proof or counterexample, etc) will win $\$1000$, and will win that for every problem solved. I looked at Problem 5 in particular, and someone had found a counterexample to the proposition and won the prize. You can go here to check out the problems, and Problem 5 is the very last one. It mentions that 20 seems to (also) be a counterexample, but it has not been proven to be. However, I believe I might have found myself another potential counterexample, namely, $$387420547.$$ This number cannot be generated from any other

numbers, as what can happen in Problem 5. This is because in the problem, you can join primes together to make a new number that might be prime. In the number above, there are n prime numbers such that when you concatenate them, they make this number. After going to the link and reading the problem, you will understand why this is very important. If you cannot access the link, please let me know, and I will write out the problem in this post for all to see. My question is, Is the number above an actual counterexample? It seems like it never climbs to a prime, no matter how many iterations I make. Can it be checked for certain? Thank you in advance.

论坛问题帖 | no_answers_at_retrieval | Mr Pie | 2018-05-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018875 | MSE-2773046

Confirmation of Proof: $\pi_n + p_a + p_b \geqslant \sum_{i=1}^4 x_i$.

I was messing around with numbers and I made the following conjecture: Conjecture: Let π_n be the n^{th} perfect number; p_a be the prime after π_n and p_b be the prime before π_n. Then, one may always see that$$\begin{align}\pi_n^{ \ \ 2}+p_a^{ \ \ 2}+p_b^{ \ \ 2}&=\sum_{i=1}^4x_i^{ \ 2}\tag{$\exists x_i\in\mathbb{Z}^+$} \\ \Leftrightarrow \pi_n + p_a + p_b&\geqslant\sum_{i=1}^4x_i\tag{with equality $\Leftrightarrow n = 1$}\end{align}$$ (Developed this conjecture from here.) Does anybody know how to prove /disprove this? I know that by Lagrange's Four-Square Theorem, the first equality is true. However, I am not particularly good at inequalities, strict and nonstrict. I know that every perfect number is of the form $2^{q-1}\left(2^q-1\right)$ for a prime q and for all primes 2^q-1, thus far. (It is not proven whether or not there exists an odd perfect number, which would mean that 2^{q-1} does not divide that, supposing it exists). Thus, $\pi_n^{ \ \ 2} = 4^{q-1}(4^q - 2^{q+1}+1)$ but this does not really help me. So, I made a new substitution: $$\pi_n = (2^q-1)^2 - \sum_{i=1}^{2^{q-1}-1}(4k-1).\tag{provable by induction}$$ This looks much more appropriate to work with. Squaring this and simplifying, $$\begin{align}\pi_n^{ \ \ 2} &= \left((2^q-1)^2-\frac 13(3\cdot 2^q-4^q-2)\right)^2 \\ &=\cdots \tag{lots of working out}\\ &= 4^{q-1}(2^q-1)^2.\end{align}$$ But this is exactly what we had before (also proving that it is a perfect number). What do I do in order to (dis)prove my conjecture? Thank you in advance.

论坛问题帖 | has_answers | Mr Pie | 2018-05-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018876 | MSE-2783900

Do odd numbers n satisfying $\gcd(n, \sigma(n)) > \sqrt{n}$ have a special form?

Let $\sigma(n)$ denote the sum of divisors of the positive integer n. Using Sage Cell Server, I was able to get the following odd numbers $n < 5000$ satisfying $\gcd(n, \sigma(n)) > \sqrt{n}$: $$117 = {{3}^2}\cdot{13}$$ $$135 = {{3}^3}\cdot{5}$$ $$585 = {{3}^2}\cdot{5}\cdot{13}$$ $$775 = {{5}^2}\cdot{31}$$ $$819 = {{3}^2}\cdot{7}\cdot{13}$$ $$891 = {{3}^4}\cdot{11}$$ $$1287 = {{3}^2}\cdot{11}\cdot{13}$$ $$1305 = {{3}^2}\cdot{5}\cdot{29}$$ $$1485 = {{3}^3}\cdot{5}\cdot{11}$$ $$1989 = {{3}^2}\cdot{13}\cdot{17}$$ $$2295 = {{3}^3}\cdot{5}\cdot{17}$$ $$2793 = {3}\cdot{{7}^2}\cdot{19}$$ $$3515 = {5}\cdot{19}\cdot{37}$$ $$4095 = {{3}^2}\cdot{5}\cdot{7}\cdot{13}$$ $$4455 = {{3}^4}\cdot{5}\cdot{11}$$ $$4655 = {5}\cdot{{7}^2}\cdot{19}$$ Is this surprising? Or is there an underlying explanation for this phenomenon? Updated Question (Added October 30, 2018) Under what suitable conditions does it follow that $3 \mid n$, if n is an odd number satisfying $\gcd(n,\sigma(n)) > \sqrt{n}$? Note that it follows from the inequality $\gcd(n,\sigma(n)) > \sqrt{n}$ that n must be composite.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2018-05-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018877 | MSE-2791535

Is this integral $\int_0^\infty\frac{\cos(a x+ 2b \arctan x)}{x^2+1}dx$ exactly zero when $b\in\mathbb{N}$?

I recently encountered this integral $$\int_0^\infty\frac{\cos(a x+ 2b \arctan x)}{x^2+1}dx$$ which suspiciously close to 0 for nonzero integer values of b, as indicated by numerical calculations. When b is not an integer it is not 0. Based on my experiments I conjecture that $$\int_0^\infty\frac{\cos(a x+ 2b \arctan x)}{x^2+1}dx=0,~~~~b\in \mathbb{Z}\setminus \{0\}.$$ Question. Is the above conjecture true?

论坛问题帖 | has_answers | Tyrell | 2018-05-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018878 | MSE-2793989

Is it true that $2^{2^{n}}$ always ends in the digit 6?

As the question says, is it true, that for $n >1$, $2^{2^{n}}$ ends in the digit 6? How would one prove this? It seemed true. I considered writing it as $$2^{2^{n}} = \prod_{k=0}^{n} 2^{ n\choose k }$$ but I'm not sure that's useful. Any help, ideas, or even hints?

论坛问题帖 | has_answers | Ryan Goulden | 2018-05-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018879 | MSE-2794208

Is my proof correct on how k must be a power of 2? Are there other proofs?

So I was looking at the Fermat Primes. These are primes of the form 2^k+1 for a natural number k, such that I define by $\mathbb{N}:=\big\{1,2,3,\ldots\big\}$ and $0\notin \mathbb{N}$. We denote by F_1 the first Fermat Prime; F_2 the second Fermat Prime; et cetera up to F_k. Thus far, however, only 5 Fermat Primes are known. They are the following, respectively: $$3,5,17,257,65537,\ldots$$ I write $(\ldots)$ because I assume that there exist more Fermat Primes, but it is a conjecture that the above Fermat primes are the only ones. I introduce this to you in the event that you do not know of them, because my question very much relates to these Fermat Primes. If 2^k+1 is prime, then must it be true that k is strictly a power of 2? I noticed that, $$\begin{align}3&=2^{2^0}+1\\ 5&=2^{2^1}+1 \\ 17&=2^{2^2}+1 \\ 257&=2^{2^3}+1 \\

65537&=2^{2^4}+1.\end{align}$$ Attempt at Proof: I proved that k is even, because supposing otherwise implies that $k=2j+1\,\exists j\in\mathbb{N}$ and $$2^k+1=2^{2j+1}+1=2^{2j+1}-(-1)^{2j+1}\stackrel{\centerdot}{:}2-(-1)=3.\tag*{$\bigg(\begin{align}&\text{$a\stackrel{\centerdot}{:}b$ is read as a} \\ &\text{is divisible by b.}\end{align}\bigg)$}$$ Ergo, 2^k+1 is not prime if k is odd; k must be even. $\;\bigcirc$ Now if k is even, then $k=2j$ and $$2^k+1=2^{2j}+1=4^j+1.$$ If j is odd, then likewise, as similarly demonstrated before, $$4^j+1=4^j-(-1)^j\stackrel{\centerdot}{:}4-(-1)=5.$$ Ergo, $l\nmid k\,\exists l$ odd; k must be a power of 2 as it will only then have no odd divisors. $\;\bigcirc$ Is my proof correct? I don't see anything wrong with it, but if it is correct, are there other (better) proofs? Also, is there anything new on Fermat Primes; particularly, are there any other general constraints on k? How far has the conjecture been tested for? I checked the first 250 Fermat Primes and it seems like the conjecture is true. (If you want, you can go here, type x=2;x=2x-1;c<=250;x and then wait a minute before the first 250 Fermat Primes appear. I still have to thank one of the users on the MSE who introduced me to the site accessible via the link.) Thank you in advance.

论坛问题帖 | has_answers | Mr Pie | 2018-05-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018880 | MSE-2795345

Does the prime sequence satisfy $p_n+p_m \le p_{n+m} < p_n p_m$?

I did some experiments in SAGE and it seems like the prime sequence p_n satisfies: $$p_n+p_m \le p_{n+m} < p_n p_m$$ for $(n,m) \neq (1,1)$. For $n=1$ the last inequality is Bertrand's postulate. Here is some SAGE code to test this for some primes: lp = list(primes(10000)) for n in range(len(lp)): for m in range(len(lp)): pn = lp[n] pm = lp[m] pnm = nth_prime(n+m+2) print pn*pm>=pnm,pnm>=pn+pm, n+1,m+1,n+m+2,pn,pm,pn*pm,pnm Is this something known (if it is true)? And if so, how does one prove or disprove it, are there heuristics?

论坛问题帖 | has_answers | user276611 | 2018-05-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018881 | MSE-2795392

Conjectured inequality: $\frac{p_{2n+1}-1}{2}\geqslant p_{n+1}+2(n+1)^{2/(n-1)}+\frac12\log_{4}^2(n-6)\qquad(n>6)$

I was inspired by this post to look at some prime numbers and see if I can find a bound. In the post, a conjecture was made with variables m and n such that $(m,n)\neq (1,1)$ respectively. I was able to prove that if it was true, then $m=n$ would imply a tighter upper bound for p_{n+1}. A current upper bound for p_{n+1} is $2p_n$, but the OP's conjecture would imply that $2p_{n+1}<p_{2n+1}$ which is tighter and better than the inequality, $p_{n+1}<2p_n$. Doing some tests, I thought that a better inequality would be that $2p_{n+1}\le p_{2n+1}-1$, but then I decided to do some more tests and I came up with a conjecture. Conjecture: $$\begin{align}\frac{p_{2n+1}-1}{2}&\geqslant p_{n+1}+2(n+1)^{2/(n-1)}\underbrace{+\,\frac 12\log_4^2(n-6)}_{\text{recent edit}}\tag{$n>6$} \\ \text{with equal}&\text{ity iff $n=7$.}\end{align}$$ Can this be proven/disproven? Thank you in advance. Edit: For a tighter bound, I added the term ${}^1/{}_2\log_{4}^{2}(n-6)^\dagger$ to the RHS (Right-Hand Side). †By $\log_b^2(x)$, I mean $\big(\log_b(x)\big)^2$.

论坛问题帖 | has_answers | Mr Pie | 2018-05-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018882 | MSE-2796633

Conways 99 graph problem

The Conway 99 graph problem is stated here as one of 5 problems. My question is this: Is there somewhere a list of examples with "Conway graphs" with fewer then 99 vertices? Or can you provide examples with fewer than 99 vertices? Definition of "Conway graph": It is a simple undirected graph $G=(V,E)$ with the following property: (1) $\forall (x,y) \in E: \exists ! z \in V: (x,z),(y,z) \in E$ (2) $\forall (x,y) \in E^c: \exists ! (z,w) \in V \times V: (x,z),(x,w),(y,z),(y,w) \in E$ (The definition is from here: https://arxiv.org/pdf/1707.08047.pdf) The Conway 99 graph problem asks if there exists a "Conway graph" with 99 vertices. I am searching for examples of "small" Conway graphs. Edit: Given the answer below, the next non trivial graph with 9 vertices is: $G(2,1)$ http://mathworld.wolfram.com/GeneralizedQuadrangle.html or the same graph: $T(3,3)$: http://mathworld.wolfram.com/TorusGridGraph.html This paper of Berlekamp might also be of interest where the graph with 243 vertices is constructed.

论坛问题帖 | has_answers | user276611 | 2018-05-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018883 | MSE-2796868

On a conjecture that $P_n^{\,2}+5^2+2^k=(P_n-1)^2+l^2$.

I was looking at perfect numbers and came across something that might serve a little interesting. Denote by P_n the n^th perfect number, then there appears to always exist $k\in\mathbb{W}$ and $l\in\mathbb{Z}$ such that $$P_n^{\,2}+5^2+2^k=(P_n-1)^2+l^2.\tag{$\verb|Conjecture|$}$$ for which $\mathbb{W}:= \mathbb{N}\cup\big\{0\big\}=\big\{0,1,2,3,\ldots\bigr\}$; id est, the whole numbers. Here is what I have discovered thus far, given that $(6,28,496)=(P_1,P_2,P_3)$ respectively: $$\begin{align}6^2+5^2+2^6&=5^2+10^2 \\ 28^2+5^2+2^0&=27^2+9^2 \tag*{$\begin{pmatrix}(k,l)\,\,=&(6,10)&\text{resp.}\\ (k,l)\,\,=&(0,9)&\text{resp.} \\ (k,l)\,\,=&(3,32)&\text{resp.}\end{pmatrix}$}\\ 496^2+5^2+2^3&=495^2+32^2.\end{align}$$ This is as far as I have tested. I know that for every perfect number P_n, there exists a prime number p such that $P_n=2^{p-1}(2^p-1)$ thus far (or at least, for every even perfect number), but this does not help me prove the conjecture. It is also conjectured that every perfect number is the sum of three cubes, but supposing this is valid does not help me prove this conjecture either. And, writing the equation like the following does not help as well. $$2^k+(P_n+l)(P_n-l)=(P_n+4)(P_n+6).$$ Can it be verified whether or not my conjecture holds water? Thank you in advance.

论坛问题帖 | has_answers | Mr Pie | 2018-05-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018884 | MSE-2797642

Approaches to Brocard's problem

The brocard problem is an unsolved problem that asks how many integers m and n exist such that $n! + 1 = m²$ More specifically it conjectures there are only three such numbers. What are some interesting ways to approach this conjecture? I've being playing around with it for over an hour and I tried the following approach. For any n, what is the smallest integer a, such that $n! + a = m²$? I created a function $v(n!)=a$ and now restated the conjecture in the following terms. $ v(n!) > 0 $ for all n except 4,5,7 I checked to see if v(n!) had any interesting pattern and as expected I found none. The next natural thing I wanted to do was to graph the function. To do this I need to somehow make n!, naturally we can do this with the gamma function. $v(\gamma(n))$. I am no expert on the gamma function, but I'd be really interested to see what patterns could emerge from such extensions. Do you know any papers/books discussing interesting approaches to resolving this conjecture?

论坛问题帖 | has_answers | Ryan Cole | 2018-05-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018885 | MSE-2801433

On a conjecture that $\sum\limits_{n=1}^k\frac{1}{\pi^{1/n}p_n}\stackrel{k\to\infty}{\longrightarrow} 2$.

I have made the following conjecture, and I do not know if this is true. Conjecture: \begin{equation*}\sum_{n=1}^k\frac{1}{\pi^{1/n}p_n}\stackrel{k\to\infty}{\longrightarrow}2\verb| such that we denote by | p_n\verb| the | n^\text{th} \verb| prime.|\end{equation*} Is my conjecture true? It seems like it, according to a plot made by Wolfram|Alpha, but if it does, then it converges.... very.... very, slowly. In fact, let $k=5000$, then the sum is approximately equal to 1.97, which just proves how slow it would be. Is there a way of showing whether or not this is indeed convergent? For any other higher values of k, it seems that it is just too much for Wolfram|Alpha to calculate, and it does not give me a result when I let $k=\infty$. Also, for users who might not understand the notation, we can similarly write that $$\sum_{n=1}^\infty\frac{1}{\pi^{1/n}p_n}=2\qquad\text{ or }\qquad\lim_{k\to\infty}\sum_{n=1}^k\frac{1}{\pi^{1/n}p_n}=2.$$ Also, without Wolfram|Alpha, I have no idea how to approach this problem in terms of proving it or disproving it. Does the sum even converge at all? If so, to what value? Any help would be much appreciated. Thank you in advance. Edit: I looked at this post to see if I could rewrite my conjecture as something else in order to help myself out. Consequently, I wrote that $$\sum_{n=1}^k\frac{1}{\pi^{1/n}p_n}\stackrel{k\to\infty}{\longleftrightarrow}4\sum_{n=1}^\infty\frac{1}{n^k+1}\tag{$\text{LHS}=2$}$$ since both sums look very similar. Could this be of use?

论坛问题帖 | has_answers | Mr Pie | 2018-05-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018886 | MSE-2802872

Trigamma identity $\psi_1\left(\frac{11}{12}\right)-\psi_1\left(\frac{5}{12}\right)=4\sqrt 3 \pi^2-80G$

Regarding this integral: Integral $\int_0^1 \frac{\sqrt x \ln x} {x^2 - x+1}dx$ The following conjecture comes: $$\psi_1\left(\frac{11}{12}\right)-\psi_1\left(\frac{5}{12}\right)=4\sqrt 3 \pi^2-80G$$ Where G is Catalan's constant. How can we show this? Wolfram-alpha agrees on this: https://www.wolframalpha.com/input/?i=trigamma(11%2F12)-trigamma(5%2F12)%3D4sqrt3pi%5E2-80Catalan

论坛问题帖 | has_answers | Zacky | 2018-05-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018887 | MSE-2804109

A conjecture regarding a strange infinite sum

Numerical calculations indicate that the following conjecture is true $$ \sum _{n=0}^{\infty} \frac{\prod _{k=0}^{n-1} \left(x+e^{-k}\right)}{\prod _{k=0}^n \left(1+te^{-k}\right)}t^n =\frac{1}{1-xt}.\tag{1} $$ Here the $n=0th$ term of the sum is $\frac{1}{1+t}$ to avoid confucion. Question. Is the conjecture (1) correct? This sum emerged in certain birth and death process calculations. I have no idea how to prove it.

论坛问题帖 | no_answers_at_retrieval | Tyrell | 2018-06-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018888 | MSE-2806590

A conjectured formula for the polylogarithm of a negative integer order

I discovered the following formula while working on the sequence A141697 from the OEIS. I have no idea whether it is something trivial or not. I would be very happy to know more about it. $$ \textrm{Li}_\nu\left(z\right)= \frac { 6 (1+z)^{-\nu-1} + \displaystyle\sum_{k=0}^{-\nu-1} \displaystyle\left(-6 \displaystyle{{-\nu-1}\displaystyle\choose k}+7\sum_{j=0}^{k+1}(-1)^j (k-j+1)^{-\nu} {{-\nu+1}\ \choose j} \right) z^k } { 7 (1 - z)^{-\nu+1} }z $$ when ν is a negative integer. In case I mistyped the formula, here are my codes in two different languages: For Pari-GP: mypolylog(n, x) = { (6*(x+1)^(-n-1) + sum(k=0,-n-1, (-6*binomial(-n-1,k) + 7*sum(j=0,k+1, (-1)^j * (k-j+1)^(-n) * binomial(-n+1,j)))*x^k)) * x / (7*(1-x)^(-n+1)) } For Mathematica: mypolylog[n_, x_] := (6*(x+1)^(-n-1) + Sum[(x^k*(-6*Binomial[-n-1, k] + 7*Sum[(-1)^j*(k-j+1)^(-n)*Binomial[-n+1, j], {j, 0, k+1}])), {k, 0, -n-1}]) / (7*(1 - x)^(-n+1)) * x

论坛问题帖 | has_answers | Thomas Baruchel | 2018-06-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018889 | MSE-2815447

Conjecture about the prime number function

Define $p_{n,m}$ by $p_{n,1}=p_n$ and $p_{n,m}=p_{p_{n,m-1}}$, where p_n is the prime number function. Conjecture: $n^n<p_{n,n}< (n+1)^{(n+1)}$ I've only tested it for $n=1,\dots,11.$ I need help testing the conjecture for $n>11$ and to prove it. Perhaps the conjecture is a consequence of PNT? n n^n p_{n,n} (n+1)^{n+1} 1 1 2 4 2 4 5 27 3 27 31 256 4 256 277 3125 5 3125 5381 46656 6 46656 87803 823543 7 823543 2269733 16777216 8 16777216 50728129 387420489 9 387420489 1559861749 10000000000 10 10000000000 64988430769 285311670611 11 285311670611 2428095424619 8916100448256 Calculate $p_{n,m}$ by m times iteration: 1. $p_{n,1}=p_n$ 2. $p_{n,2}=p_{p_{n,1}}$ .. m. $p_{n,m}=p_{p_{n,m-1}}$

论坛问题帖 | has_answers | Lehs | 2018-06-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018890 | MSE-2825945

If $a+b+c$ divides the product abc, then is (a,b,c) a Pythagorean Triple?

Firstly, I will define what Pythagorean Triples are for those who do not know. Definition: A Pythagorean Triple is a group of three integers a, b and c such that $a^2+b^2=c^2$, since the Pythagorean Theorem asserts that for any 90° (right-angle) triangle ABC with sides a, b and c, one will always have the equation, $a^2+b^2=c^2$. I was looking at Pythagorean Triples and noticed another property apart from how $a^2+b^2=c^2$. Here are the first 30 Pythagorean Triples (a,b,c) ordered from smallest to greatest value, i.e. $$(a,b,c)\qquad\text{ s.t. }\qquad a<b<c.\tag*{$\big(\text{s.t. = such that}\big)$}$$ I noticed that $a^2=(c+b)(c-b)$, but that is trivial since $$\begin{align}a^2&=(c+b)(c-b)\tag{given} \\ &=c^2-b^2 \\ \Leftrightarrow\,\,\,\, a^2+b^2&=c^2.\end{align}$$ However, I also noticed that by having "$u\mid v$" be read as "u divides v", it appears that $$a+b+c\mid abc.$$ For example, $(a,b,c)=(3,4,5)$ is a classic Pythagorean Triple; $3^2+4^2=5^2$. Also, $$\begin{align}3+4+5&=12 \\ \& \quad3\times 4\times 5 &= 60. \\ \\ 12 &\,\mid 60 \\ \Leftrightarrow \,\,\,\,\,3+4+5&\,\mid 3\times 4\times 5.\end{align}$$ This, I cannot prove to be true $-$ but I tested with all the 30 Pythagorean Triples above, and I have come across no counter-example. Is there a proof? I do not know where to begin myself. Conjecture: Given three positive integers a, b and c, if $a < b<c$ and $a^2+b^2=c^2$, then $$a+b+c\mid abc.$$ Thank you in advance. Edit: My conjecture was originally the other way round; i.e. if $a+b+c\mid abc$ then $a^2+b^2=c^2$. But 6 is a counter-example, namely because it is a Perfect Number.

论坛问题帖 | has_answers | Mr Pie | 2018-06-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018891 | MSE-2828167

Two sequences which have an average tend to zero

Let $(a_n)_{n\geq 1}, (b_n)_{n\geq 1}$ be two sequence of positive real numbers such that $$ \lim_{n\to +\infty}\frac{a_1+a_2+\cdots+a_n}{n}=\lim_{n\to +\infty}\frac{b_1+b_2+\cdots+b_n}{n}=0.$$ Conjecture. For all $\epsilon>0$, there are infinitely many values of indices k such that $a_k<\epsilon$ and $b_k<\epsilon.$ I think that this is true but I can not prove it now. In the special case where $a_n = b_n$, that is, there is only one sequence, then one can argue easily using a contradiction argument. In the general case, the hard part is to show that the same set of indices are shared by both sequences $(a_n)_{n\geq 1}, (b_n)_{n\geq 1}$.

论坛问题帖 | has_answers | Plimpton 322 | 2018-06-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018892 | MSE-2834013

A modular conjecture about odd primes

Conjecture: $n^{2p-1}\equiv n\pmod {2p}$ for all $n\in \mathbb N$ and all odd primes p. I started investigate the least x_n such that $p_n^{p_{n+1}}\equiv x_n\pmod{p_{n+2}}$ and ended up with this for me unknown conjecture. I've tested it for a lot of random values of n and p.

论坛问题帖 | has_answers | Lehs | 2018-06-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018893 | MSE-2834518

The equation $\varphi(n)=n-\log_2(1+\operatorname{gpf}(n))-\operatorname{gpf}(n)+1$ and Mersenne primes

Let $n\geq 1$ an integer, we denote the Euler's totient function as $\varphi(n)$ and the greatest prime dividing n as $\operatorname{gpf}(n)$ (that it the arithmetic function defined in the MathWorld's article Greatest Prime Factor). I know that is it possible to prove the following claim (I was inspired in [1] to ask a similar question). Claim. If n is an integer of the form $n=p\cdot(2^p-1)$, where 2^p-1 is a Mersenne prime, then satisfies $$\varphi(n)=n-\log_2(1+\operatorname{gpf}(n))-\operatorname{gpf}(n)+1.\tag{1}$$ In this post I ask about if those are the only solutions of the equation (1). Previous equation (1) can be written also as $$\left(1+\operatorname{gpf}(n)\right)\cdot 2^{\varphi(n)+\operatorname{gpf}(n)}=2^{n+1}.\tag{2}$$ Question. Prove or disprove* the following conjecture: If an integer $N\geq 1$ satisfies our equation $$\left(1+\operatorname{gpf}(N)\right)\cdot 2^{\varphi(N)+\operatorname{gpf}(N)}=2^{N+1}$$ then $N=p(2^p-1)$, where 2^p-1 is a Mersenne prime. Many thanks. *To disprove it show a counterexample. References: [1] Tripathi, A note on products of primes that differ by a fixed integer, The Fibonacci Quarterly, Vol. 48, No. 2 (MAY, 2010).

论坛问题帖 | has_answers | user243301 | 2018-06-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018894 | MSE-283960

Question regarding the status of Erdős' conjectures

Has Erdős' conjecture on arithmetic progressions or his conjecture on Sylvester's sequence been proven?

论坛问题帖 | has_answers | user1825464 | 2013-01-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018895 | MSE-2847779

On Sierpinski's Conjecture

The statement of Sierpinski's Conjecture is 'for every integer $n>1$, there exist three integers a,b,c such that $\frac{5}{n}=\frac{1}{a}+\frac{1}{b}+\frac{1}{c}$ Would this conjecture be proved if we could show this for every prime number n ? If this is true, how do we prove this?

论坛问题帖 | no_answers_at_retrieval | Clarkson | 2018-07-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018896 | MSE-2855047

Prove that $\sum\limits_{i=1}^{n} a_i\geq n^2$.

A hint can be helpful, but not a whole solution. The Problem (conjecture): Given a natural number $n \geq 1$ and a sequence of natural numbers $(a_i)_{1 \leq i \leq n}$ in which for every pair (i,j) with $i \neq j,$ we have $$\gcd(a_i,a_j)\nmid i-j$$ prove that $\sum\limits_{i=1}^{n} a_i\geq n^2$. What I have done: During my research, I ran into this problem and I am not quite sure if it is true. It is clear if we put $a_i=n$ then the problem will be solved and the summation will be equal to n^2. I tried to solve this problem. For example, I showed that $$a_i> max(i,n-i)$$ otherwise, I can put $j=i+a_i$ or $j=i-a_i$ and considering the fact $\gcd(a_i,a_j) \mid a_i,$ then we conclude that $\gcd(a_i,a_j)\mid i-j,$ hence, $a_i> max(i,n-i)$ which means that $a_i\geq \dfrac{n} {2}$. Moreover, if $a_i\leq n$ and p are prime divisors of a_i by putting $j=i-\dfrac {a_i}{p}$ for $i\geq \dfrac {n} {2}$ and $j=i+\dfrac {a_i}{p}$ for $i\leq \dfrac {n} {2}$ we could conclude that $a_i \mid a_j$. I could go further, but it is not enough to prove the conjecture. I also tried Induction and considered that the property holds for every $n\leq k$ and then try to prove the theorem for $n= k+1$ but again, there are some issues that I could not go further.

论坛问题帖 | has_answers | farshid evazabadian | 2018-07-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018897 | MSE-2857099

Is ${n^2}/D(n^2) \in \mathbb{N}$, if $q^k n^2$ is an odd perfect number?

Let $x \in \mathbb{N}$, the set of positive integers. The sum of the divisors of x is denoted by $\sigma(x)$. Denote the deficiency of x by $D(x):=2x-\sigma(x)$, and the sum of the aliquot parts of x by $s(x):=\sigma(x)-x$. Finally, denote the abundancy index of x by $I(x):=\sigma(x)/x$. If m is odd and $\sigma(m)=2m$, then m is called an odd perfect number. Euler proved that an odd perfect number (if one exists) must have the form $m=q^k n^2$ where q is the special / Euler prime satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. Here is my question: Is ${n^2}/D(n^2) \in \mathbb{N}$, if $q^k n^2$ is an odd perfect number? MY ATTEMPT From the fundamental equality $$\frac{\sigma(n^2)}{q^k}=\frac{2n^2}{\sigma(q^k)}$$ one can derive $$\frac{\sigma(n^2)}{q^k}=\frac{2n^2}{\sigma(q^k)}=\gcd(n^2,\sigma(n^2))$$ so that we ultimately have $$\frac{D(n^2)}{s(q^k)}=\frac{2s(n^2)}{D(q^k)}=\gcd(n^2,\sigma(n^2)).$$ Thus, we have $$y:=\frac{n^2}{D(n^2)}=\frac{{n^2}}{s(q^k)\gcd(n^2,\sigma(n^2))}=\frac{{n^2}D(q^k)}{2s(q^k)s(n^2)}.$$ In particular, we obtain $$\frac{n^2}{D(n^2)}=\frac{{n^2}}{s(q^k)\gcd(n^2,\sigma(n^2))}=\frac{\sigma(q^k)}{2s(q^k)}.$$ Using the reasoning in this blog post, we obtain $$\frac{q}{2} + \frac{1}{2{q^{k-1}}} - \frac{1}{2q^k} < y \leq \frac{q}{2} + \frac{1}{2{q^{k-1}}}.$$ Equality holds if and only if $k=1$. That is, it appears that $y \in \mathbb{N}$ if and only if $k=1$ (i.e., the Descartes-Frenicle-Sorli conjecture holds). So the question "Is ${n^2}/D(n^2) \in \mathbb{N}$?" is equivalent to asking whether $k=1$, for $m=q^k n^2$ an odd perfect number with special / Euler prime q.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2018-07-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018898 | MSE-2861604

Conjecture about special grid of numbers

Consider you have created grid of numbers like following image starting from any positive integer (in this case 8) To create such grid, follow this steps; Pick a number greater than one and write down $n, n-1, n-2, ..., 1$ as column headers Likewise, write down $n, n-1, n-2, ..., 1$ as row headers Fill each cell with $row_header * column_header$ If you group numbers diagonally starting from upper left, you get these groups of numbers {64} {56, 56} {48, 49, 48} {40, 42, 42, 40} etc. Let G_n denote nth group. I want to prove if following theory is correct. $$ i < k \Rightarrow\forall x \in G_i, \forall y \in G_k, x > y $$ It seems to hold for small enough starting numbers like 8. It is trival to prove that any cell will be bigger then the cells to the right on the same row and will be bigger then the cells below it on the same column. But, I can't be sure about diagonal groups.

论坛问题帖 | has_answers | yasar | 2018-07-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018899 | MSE-2866629

Show that $\int_0^1 4 \space\operatorname{li}(x)^3 \space (x-1) \space x^{-3} dx = \zeta(3) $

My mentor tommy1729 wrote $\int_0^1 4 \space \operatorname{li}(x)^3 \space (x-1) \space x^{-3} dx = \zeta(3) $ I wanted to prove it thus I looked at some methods for computing integrals and also representations of $\zeta(3)$ that might be useful. But nothing was very helpful to me. In particular the fact that the RHS is so short - Just Apery’s constant - was surprising. I expected it longer and more complicated. So I tend to believe that either the

integral computation requires many steps and Then Finally we Get a long expression but alot of cancellation until we are left with Apery’s constant only. Or There is a simple way to get Apery’s constant directly with a trick I missed. In either case it is amazing I would say. So How to show that $$\int_0^1 4 \space \operatorname{li}(x)^3 \space (x-1) \space x^{-3} dx = \zeta(3) $$ I would like to see different ways to show it. I assume real analysis methods are simpler than complex analysis methods (on the complex plane like contour integration). I Also wondered If not knowing the RHS in advance would change the difficulty of this question. Also I wonder about $$ \int_0^1 5 \space \operatorname{li}(x)^4 \space (x-1) \space x^{-4} dx = ?? $$

论坛问题帖 | has_answers | mick | 2018-07-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018900 | MSE-2870886

On the quantity ${n^2}/D(n^2)$ where n^2 is the non-Euler part of members of the OEIS sequence A228059

Let $\sigma(x)$ denote the sum of the divisors of the number $x \in \mathbb{N}$, the set of positive integers. Denote the deficiency of x as $D(x):=2x-\sigma(x)$. This afternoon I noticed some interesting fact about the quantity ${n^2}/D(n^2)$ where n^2 is the non-Euler (i.e., square) part of members of the OEIS sequence A228059. Specifically, we have the following values of n^2 from members of OEIS sequence A228059 and the corresponding (apparently almost increasing, and non-integral) values of ${n^2}/D(n^2)$: $${n_1}^2 = 3^2 = 9, {{n_1}^2}/D({n_1}^2) = 9/5 = 1.8$$ $${n_2}^2 = 3^4 = 81, {{n_2}^2}/D({n_2}^2) = {81}/{41} \approx 1.97561$$ $${n_3}^2 = {21}^2 = 441, {{n_3}^2}/D({n_3}^2) = {147}/{47} \approx 3.12766$$ $${n_4}^2 = {45}^2 = 2025, {{n_4}^2}/D({n_4}^2) = {2025}/{299} \approx 6.77258$$ $${n_5}^2 = {135}^2 = 18225, {{n_5}^2}/D({n_5}^2) = {18225}/{2567} \approx 7.09973$$ $${n_6}^2 = {285}^2 = 81225, {{n_6}^2}/D({n_6}^2) = {27075}/{2969} \approx 9.11923$$ $${n_7}^2 = {165}^2 = 27225, {{n_7}^2}/D({n_7}^2) = {27225}/{851} \approx 31.9918$$ $${n_8}^2 = {765}^2 = 585225, {{n_8}^2}/D({n_8}^2) = {585225}/{18893} \approx 30.9758$$ $${n_9}^2 = {7695}^2 = 59213025, {{n_9}^2}/D({n_9}^2) = {19737675}/{731333} \approx 26.9886.$$ Here is my question: Is it always the case that ${n^2}/D(n^2)$ is non-integral where n^2 is the non-Euler (i.e., square) part of members of the OEIS sequence A228059? Updated August 28 2018 The short answer to my original question is NO. (See the answer below.) I ask because it is known that the exponent of the special / Euler prime of an odd perfect number is 1 if and only if the non-Euler part is deficient-perfect. Coincidentally, in OEIS sequence A228059, all of the special / Euler primes for the first 9 terms have exponent 1.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2018-08-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018901 | MSE-2873204

Positive integers have the form $a^2+b^2+c^2+2^d$

For any positive integer n there seems to be non-negative integers a,b,c,d such that $$n=a^2+b^2+c^2+2^d.$$ Due to Legendre's three-square theorem a natural number can be represented as the sum of three squares of integers $$m=a^2+b^2+c^2$$ if and only if m is not of the form $m=4^a(8b+7)$, for integers a and b. If the conjecture is true, then for all $n\geq 1$ it must exist a k such that $n-2^k$ is not of the form $m=4^a(8b+7)$. I want help to prove the conjecture or to find a counter-example.

论坛问题帖 | has_answers | Lehs | 2018-08-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018902 | MSE-2880738

Is there any conjecture that has been proved to be solvable/provable but whose direct solution/proof is not yet known?

In mathematics, is there any conjecture about the existence of an object that was proven to exist but that has not been explicitly constructed to this day? Here object could be any mathematical object, such as a number, function, algorithm, or even proof.

论坛问题帖 | has_answers | lone student | 2018-08-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018903 | MSE-2886597

Question about a result on odd perfect numbers

In the paper titled Improving the Chen and Chen result for odd perfect numbers (Lemma 8, page 7), Broughan et al. show that if $$\frac{\sigma(n^2)}{q^k}$$ is a square, where $\sigma(x)$ is the sum of divisors of $x \in \mathbb{N}$ and $q^k n^2$ is an odd perfect number with special/Euler prime q, then $k=1$. Is it then possible to derive the implication $$D(n^2) = 2n^2 - \sigma(n^2) \text{ is a square } \implies k=1?$$ I could only derive the equation $$\frac{\sigma(n^2)}{q^k} = \frac{D(n^2)}{s(q^k)}$$ where $s(q^k) = \sigma(q^k) - q^k = \sigma(q^{k-1})$. Added August 25 2018 I also currently know that $$D(n^2) \mid n^2 \iff k=1.$$ That is, the Descartes-Frenicle-Sorli conjecture is true for the odd perfect number $q^k n^2$ if and only if the non-Euler factor n^2 is deficient-perfect. Added July 07 2019 If $$\frac{\sigma(n^2)}{q^k}=\frac{D(n^2)}{s(q^k)}$$ is a square, then by Broughan et al.'s result, we have $k=1$, which implies that $$s(q^k) = \sigma(q^k) - q^k = \sigma(q) - q = (q+1) - q = 1$$ is a square. Consequently, if $\sigma(n^2)/q^k$ is a square, then $D(n^2)$ is a square too.

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2018-08-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018904 | MSE-2887045

Conjecture that $ \frac{\gcd(a+b,ab)}{\gcd(a,b)} \mid \gcd(a,b)$

I have discovered some exercise type conjectures which I can't prove and this is one of them: Given positive integers a,b, then $$ \frac{\gcd(a+b,ab)}{\gcd(a,b)}\ \bigg|\ \gcd(a,b)$$ Can this be proved or disproved? From time to time, when testing my growing math packages BigZ and Forthmath, I recognize some patterns which I can't prove or disprove (or even have the ambition to). I post them here with the hope that it will not annoy too much. I hope you can bear with it.

论坛问题帖 | has_answers | Lehs | 2018-08-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018905 | MSE-2887078

Conjecture: $ \gcd(\operatorname{rad}(a+b) ,ab)= \operatorname{rad}(\gcd(a,b))$

I have discovered some exercise type conjectures which I can't prove and this is one of them: Given positive integers a,b, then $$ \gcd(\operatorname{rad}(a+b) ,ab)= \operatorname{rad}(\gcd(a,b))$$ Can this be proved or disproved? From time to time, when testing my growing math packages BigZ and Forthmath, I recognize some patterns which I can't prove or disprove (or even have the ambition to). I post them here with the hope that it will not annoy too much. I hope you can bear with it.

论坛问题帖 | has_answers | Lehs | 2018-08-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018906 | MSE-2889532

A conjecture of exercise type: $\gcd(a,b)^2=\gcd(a^2+b^2,ab)$

This must be known, but I haven't found it and want help to prove it: For all $a,b\in \mathbb Z$, $\gcd(a,b)^2=\gcd(a^2+b^2,ab)$ Tested for $10,000,000$ pseudo random number pairs. From time to time, when testing my growing math packages BigZ and Forthmath, I recognize some patterns which I can't prove or disprove (or even have the ambition to). I post them here with the hope that it will not annoy too much. I hope you can bear with it.

论坛问题帖 | has_answers | Lehs | 2018-08-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018907 | MSE-2895522

There are primes p,q such that $\gcd(a,b)=|ap-bq|$

Conjecture: Given $a,b\in\mathbb Z^+$ there are primes p,q such that $\gcd(a,b)=|ap-bq|$. I would like help with a proof or a counter-example. Tested for millions of pseudo random numbers.

论坛问题帖 | has_answers | Lehs | 2018-08-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018908 | MSE-2902568

Subsubgroups are subgroups of subgroups / Multiplicative Property of the Index

Algebra by Michael Artin Prop 2.8.14 Multiplicative Property of the Index Statement of Prop 2.8.14 Let $G \supseteq H \supseteq K$ be subgroups of a group G. Then $[G:K] = [G:H][H:K]$. Proof of Prop 2.8.14 Here's the original Counting Formula (Formula 2.8.8) Question: Could we perhaps prove Prop 2.8.14 by using Counting Formula 2.8.8? I think we could do so (Tower Law for Subgroups) if we show that K is a subgroup of H which I guess hasn't been proven yet, if it's even true.

论坛问题帖 | has_answers | BCLC | 2018-09-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018909 | MSE-2915675

Free action of finite cyclic group on S^3

I'm starting to read Allen Hatcher's Space of Knots and at the end of the first paragraph he says that his work would apply to all knots if the following conjecture is true: Every free action of a finite cyclic group on S^3 is equivalent to a standard linear action. The paper is from 1999, so I don't know if this is still a conjecture. Where can I read about it and the progress on it? I'm also interested in knowing how this conjecture applies to knot theory, in particular to the results Hatcher gives in the paper.

论坛问题帖 | has_answers | Javi | 2018-09-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018910 | MSE-2931716

Is there are similar conjecture like this??

Talking with my friend, my friend suggest impressive conjecture that For $i\in\mathbb{N}$, there are always exist natural number r that satisfies $$\sum_{n=1}^{i} \frac{1}{n^r}=\frac{p}{q} , \gcd(p,q)=1$$ and $p+q$ is a prime number. For example, $$\sum_{n=1}^{2} \frac{1}{n}=\frac{3}{2}$$ and $3+2=5$. And $i=3$, $$\sum_{n=1}^{3} \frac{1}{n^4}=\frac{1393}{1296}$$ and $1393+1296=2689$. $$\sum_{n=1}^{4} \frac{1}{n^2}=\frac{205}{144}$$ and $205+144=349$ which is 70th prime number. In $i=5$ $$\sum_{n=1}^{5} \frac{1}{n^3}=\frac{256103}{216000}$$ and $256103+216000=472103$. Also $$\sum_{n=1}^{6} \frac{1}{n^3}=\frac{28567}{24000}$$ and $24000+28567=52567$. There are similer like this? If not, What about your think, is it true? Or false.... I would glad you guys to think about this thing and share opinions for me and my friend's curiosity. Note: ID:metamorphy suggest that the sequence of the smallest values of r begins with $1, 1, 1, 1, 1, 2, 1, 3, 1, 1, 1, 2, 34, 1, 1, 5, \ldots$. (There are full comment in bellow with more comments.)

论坛问题帖 | has_answers | user366725 | 2018-09-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018911 | MSE-2931943

Manufacturing desired planks from an existing pile of planks

Suppose there is a pile of commensurable planks that only may differ in lengths $0<a_1\leq\cdots\leq a_m$, which are to be used to manufacture planks of length: $0<b_1\leq\cdots\leq b_n$. Algorithm: A is the list of existent planks B is the list of wanted planks If B is empty return 'success' If $a_m<b_1$ return 'failure' Remove the smallest plank $a_i\geq b_1$ from list A Manufacture the greatest plank $b_j\leq a_i$ by, if necessary, cut a_i into planks of length b_j and $a^\prime_i$ Return what's left ($a^\prime_i$) to A Remove b_j from the list B Goto 1. I conjecture that this algorithm is optimal in two ways: If the manufacturing is possible in any way, it is also possible by the algorithm. The square sum of the lengths of the remaining planks in A is maximal compared to other cuttings. Is this possible to prove or disprove?

论坛问题帖 | has_answers | Lehs | 2018-09-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018912 | MSE-2939248

What is the common (and simplified) value for $D(q^k)D(n^2) = 2s(q^k)s(n^2)$ when $q^k n^2$ is an odd perfect number?

In an answer to an earlier question, it is shown that $$D(2^p - 1)D(2^{p-1}) = 2s(2^p - 1)s(2^{p-1}) = 2^p - 2,$$ if $2^{p-1}(2^p - 1)$ is an even perfect number, $D(x) = 2x - \sigma(x)$ is the deficiency of x, $s(x) = \sigma(x) - x$ is the sum of the aliquot divisors of x, and $\sigma(x)$ is the sum of divisors of $x \in \mathbb{N}$, the set of positive integers. (Lastly, denote the abundancy index of x by $I(x) = \sigma(x)/x$.) Here is my question in this post: What is the common (and simplified) value for $D(q^k)D(n^2) = 2s(q^k)s(n^2)$ when $q^k n^2$ is an odd perfect number? MOTIVATION Since $\gcd(q^k, \sigma(q^k)) = 1$, we have $$\frac{\sigma(n^2)}{q^k} = \frac{2n^2}{\sigma(q^k)} = \gcd(n^2, \sigma(n^2))$$ $$\frac{D(n^2)}{s(q^k)} = \frac{2s(n^2)}{D(q^k)} = \gcd(n^2, \sigma(n^2))$$ $$D(q^k)D(n^2) = 2s(q^k)s(n^2) = D(q^k)s(q^k)\gcd(n^2, \sigma(n^2)).$$ Then, we have from this earlier question the following expression for $\gcd(n^2, \sigma(n^2)$ in terms of q, k, n^2, and $\sigma(n^2)$: $$\gcd(n^2,\sigma(n^2)) = \frac{I(q^k)}{s(q^k)}\sigma(n^2) - \frac{1}{I(q^k)s(q^k)}(2n^2).$$ Plugging in this value for $\gcd(n^2, \sigma(n^2))$ into our equation for $$D(q^k)D(n^2) = 2s(q^k)s(n^2) = D(q^k)s(q^k)\gcd(n^2, \sigma(n^2)),$$ we obtain $$D(q^k)D(n^2) = 2s(q^k)s(n^2) = D(q^k)s(q^k)\Bigg(\frac{I(q^k)}{s(q^k)}\sigma(n^2) - \frac{1}{I(q^k)s(q^k)}(2n^2)\Bigg) = D(q^k)\Bigg(I(q^k)\sigma(n^2) - \frac{1}{I(q^k)}(2n^2)\Bigg).$$ Note that "if" $I(q^k) = -1$ then the equation would be trivial and the common value would just be $D(q^k)D(n^2)$. But, of course, we know that $$1 < I(q^k) < \frac{5}{4}.$$ Alas, this is where I get stuck.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2018-10-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018913 | MSE-293966

Number of subsets of U whose arithmetic mean is integral

Question is :- n is a positive integer. Call a non-empty subset S of $\{1,2,\dots,n\}$ "good" if the arithmetic mean of elements of S is also an integer. Further Let t_n denote the number of good subsets of $\{1,2,3,\ldots,n\}$. Prove that t_n and n are both odd or both even. And my solution to this problem is :- We deal with problem in 2 cases : n is even and other when n is odd. If n is odd , we first calculate n={1,2} by brute force. We get numbers of solutions for n=1 as 1 & n=3 as 5. Thus we conjecture that t_n=$t_{n-2}+2(n-1)$ for n as even. So by this recurrence we calculate t_5 which comes to be 15. A brute force manual calculation confirms this answer . Thus we prove our conjecture by induction. Now t_1 is odd (=1) so t_2 has to be odd since t_2=$t_1+2(3-1)$,here t_1 is odd & other term is even thus overall parity is odd. So t_2 is odd. Again t_3 has to be odd as t_3=t_2+2(3-1), t_2 is odd & other term even , so t_3 is odd. So , inductively we conclude that t_n has to be odd . For the other case we conjecture same recurrence relation and prove in the same way.

论坛问题帖 | has_answers | Maggi Iggam | 2013-02-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018914 | MSE-2949092

Primes of the form $p_1p_2\dotsm p_k+1$

Let $p_1,p_2,...,p_k$ be the first k primes and $k>1$. Does there exist a prime p_{k+1} such that $p_{k+1} = p_1p_2...p_k + 1$?

论坛问题帖 | has_answers | Tom Carter | 2018-10-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018915 | MSE-2952820

Calculating the $\sum_{x=1}^\infty\left[\prod_{k=0}^s{s\choose k}x^k\right]^{-1}$

Question: If $s \in \mathbb{N}$ is it true: $\sum_{x=1}^\infty\left[\prod_{k=0}^s{s\choose k}x^k\right]^{-1}={\zeta\left[{s+1 \choose 2}\right] \above 1.5pt \prod_{k=1}^s{s \choose k}};$ where $\zeta\left[{s+1 \choose 2}\right]$ is the Riemann zeta function^1 evaluated at the triangular numbers^2 ? I believe I can show the answer to the question is yes by way of inspection. I cannot get an actual explicit calculation that answers the question affirmatively - that is what I am asking for ! Solution to question by inspection: For short handedness I write $$\zeta_{\times}(s)=\sum_{x=1}^\infty\Bigg[\prod_{k=0}^s{s\choose k}x^k\Bigg]^{-1}$$ I computed a small list of values of $\zeta_{\times}(s)$ for small s: \begin{array}{ l | l } s & 2 & 3 & 4 & 5 & 6 & 7\\ \hline \zeta_{\times}(s) & \frac{\zeta(3)}{2} & \frac{\zeta(6)}{9} & \frac{\zeta(10)}{96} & \frac{\zeta(15)}{2500} & \frac{\zeta(21)}{162000}& \frac{\zeta(28)}{26471025}\\ \end{array} Inspection suggests that $\zeta_{\times}(s)$ has numerator equal to the Riemann zeta function at ${s+1 \choose 2}.$ The denominators of $\zeta_{\times}(s)$ can be written explicitly as $2,9,96,2500,162000,\ldots$; which appear to be the integer sequence A001142^3. Listed in the reference section is the following paper by Lagarias: Products of Binomial Coefficients

and Unreduced Farey Fractions.^{4}. Lagarias defines the unreduced Farey fractions^5 to be the ordered sequence of all reduced and unreduced fractions between 0 and 1 with denominator of size at most $s.$ Following Lagarias' notation: I write G_s for the set of unreduced Farey fractions and let $|G_s|$ and G_s^* denote the cardinality and the product of all elements of G_s respectively. Lagarias shows the following results: $|G_s|={s+1 \choose 2}$ and $G_s^*=\prod_{k=1}^s{s \choose k}.$ I put everything together and get $$\zeta_{\times}(s)={\zeta\big(|G_s|\big) \above 1.5pt G_s^*}$$ and I am done. $\blacksquare$ Source and motivation of the problem: Curiosity is the main driving source of the problem. Here it goes: If x and s are positive numbers I write $\zeta(s)$ for the Riemann zeta function evaluated at the number $s.$ For ease of typing I write Pascal's Triangle^{6} like this: $$\text{ }\begin{matrix} 1&&&&\\ 1&1&&&\\ \color{red}{1}&\color{red}{2}&\color{red}{1}&&\\ \color{blue}{1}&\color{blue}{3}&\color{blue}{3}&\color{blue}{1}&\\ 1&4&6&4&1\\ \vdots \end{matrix}$$ and recall that the first row in Pascal's triangle is numbered at 0. Now look at the second and third rows of the triangle, highlighted in red and blue respectively and observe: $\sum_{x=0}^\infty{ 1 \above 1.5pt \color{red}{1}+\color{red}{2}x+\color{red}{1}x^2}=\zeta(2)$ and $\sum_{x=0}^\infty{ 1 \above 1.5pt \color{blue}{1}+\color{blue}{3}x+\color{blue}{3}x^2+\color{blue}{1}x^3}=\zeta(3).$ This observation shows that $\zeta(s)$ can be recovered from the rows of Pascal's triangle - in particular I can write $\sum_{x=0}^\infty\frac{1}{(1+x)^s}=\zeta(s).$ On the other hand using the binomial theorem^{7} allows me to compute $(1+n)^s$ explicitly with the formula: $\sum_{k=0}^s{s\choose k}x^k$ in which case after substitution: $$\zeta(s)=\sum_{x=0}^\infty\bigg[\sum_{k=0}^s{s\choose k}x^k\bigg]^{-1}$$ Now out of total curiosity I decided to swap the inner summation inside the big brackets and swap it out with a product, noting carefully that if I do that I need to shift the starting point in the outer summation to 1 otherwise I would be dividing by zero. For short handedness I wrote $$\zeta_{\times}(s)=\sum_{x=1}^\infty\Bigg[\prod_{k=0}^s{s\choose k}x^k\Bigg]^{-1}$$ Numerical inspection suggested $$\zeta_{\times}(s) ={\zeta\left[{s+1 \choose 2}\right] \above 1.5pt \prod_{k=1}^s{s \choose k}}$$ I double checked my numerical hunch against Sloan's Database and encountered the paper by Lagarias. In Lieu of the fact that Lagarias has an explicit formulae for $\log G_s^*$ (see Lagarias paper or reference above for notation) I primarily became interested in computing $\log \zeta_{\times}(s).$ If I could answer the question affirmatively then I would know that $$\log \zeta_{\times}(s)=\log \zeta(|G_s|)-\log(G_s^*)$$

论坛问题帖 | has_answers | Anthony | 2018-10-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018916 | MSE-2954161

Conjecture for $\ln(x)$ and $\frac{d^ny}{dx^n}$

I'm not really sure where to start, I found the first, second, third and fourth derivatives of $\ln(x)$ to be $\frac{1}{x}$,-$\frac{1}{x^2}$, $\frac{2}{x^3}$, and -$\frac{6}{x^4}$, respectively. Letting n be a natural number, I have to formulate a conjecture for a formula for $\frac{d^ny}{dx^n}$. Afterward, I have to use mathematical induction to prove the conjecture.

论坛问题帖 | has_answers | Claire | 2018-10-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018917 | MSE-2954401

Validation for a conjecture about Chinese Remainder Theorem for groups

I was wondering if the following statement is true: Let G be a group with normal subgroups $H_1,H_2,...H_n$. Suppose $H_iH_j=G$ for all $i\neq j$. Then $G/H_1\cap H_2...\cap H_n\cong G/H_1 \times...\times G/H_n.$ There is a similar question here, which is the case when $n=2$. There is also a similar question here, but the conditions involve the index of a subgroup. I want to get rid of such condition so that the conclusion is applicable to some other situations. But when I tried to proceed proof by induction to get the conclusion with arbitrary n, things became not so approachable. If you think this is false, please give a counter-example. If you think this is true, please share your ideas of the proof. Thank you!

论坛问题帖 | has_answers | J. Wang | 2018-10-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018918 | MSE-2959445

Schwarz inequality for unital positive maps on C*-algebras

I was recently studying this paper by Man-Duen Choi about inequalities for positive maps on C*-algebras. He demonstrates that Let $\phi : \mathcal{A} \to \mathcal{B}$ be a unital positive linear map between the two C*-algebras $\mathcal{A}, \mathcal{B}$. Then $$ \phi(A^* A) \ge \phi(A)^* \, \phi(A) $$ and $$ \phi(A^* A) \ge \phi(A) \, \phi(A)^* $$ for every subnormal $A \in \mathcal{A}$. He then conjectures that the same result might apply also for hyponormal opertators, i.e. operators such that $A^* A \ge A A^*$ (this conjecture is equivalent to the Woronowicz's conjecture). I was wondering whether this conjecture has been better investigated or not. Also, is there a characterization of hyponormal operators?

论坛问题帖 | has_answers | luigi | 2018-10-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018919 | MSE-2960158

Is a domain a UFD if all products of two irreducibles have unique factorization?

Let A_p be an Integral domain. Conjecture : If every a in A_p that equals $b \space c$ for irreducible elements b,c in A_p , has Unique factorization then the Integral domain A_p is a Unique factorization domain (UFD). Is this true ? How to show that ?

论坛问题帖 | has_answers | mick | 2018-10-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018920 | MSE-2960648

Does mathlove's answer imply that $D(n^2) \mid n^2$?

In what follows, set $\sigma(x)$ to be the sum of divisors of $x \in \mathbb{N}$, and let $$D(x) = 2x - \sigma(x)$$ be the deficiency of x, and let $$s(x) = \sigma(x) - x$$ be the sum of the aliquot divisors of x. An earlier question asks: What is the common (and

simplified) value for $D(q^k)D(n^2) = 2s(q^k)s(n^2)$ when $q^k n^2$ is an odd perfect number? In the accepted answer, MSE user mathlove writes: $$\begin{align}D(q^k)D(n^2) &= 2s(q^k)s(n^2) \\\&=\frac{2n^2(q^k-1)(q^{k+1}-2q^k+1)}{(q-1)(q^{k+1}-1)}\end{align}.$$ Now, I know that $$s(q^k) = \sigma(q^k) - q^k = \sigma(q^{k-1}) = \frac{q^k - 1}{q - 1}$$ and that $$\gcd(D(q^k), s(q^k)) = 1.$$ Here is my question in this post: Does mathlove's answer imply that $D(n^2) \mid n^2$? I ask because the divisibility condition $D(n^2) \mid n^2$ for the non-Euler part n^2 is known to be equivalent to the Descartes-Frenicle-Sorli Conjecture (that is, $k=1$) for odd perfect numbers.

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2018-10-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018921 | MSE-2967348

Can this bound for the abundancy index of n be improved, given that $q^k n^2$ is an odd perfect number with $k=1$?

In what follows, set $I(x)=\sigma(x)/x$ to be the abundancy index of $x \in \mathbb{N}$, where $\sigma(x)$ is the sum of divisors of x. If $I(y)=2$ and y is odd, then y is called an odd perfect number. Euler proved that an odd perfect number, if one exists, must have the so-called Eulerian form $N = q^k n^2$ where q is the special/Euler prime satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. The following result was communicated to me (via e-mail, by Pascal Ochem) on April 17, 2013. If $N = {q^k}{n^2}$ is an odd perfect number given in Eulerian form, then $$I(n) > {\left(\frac{8}{5}\right)}^{\frac{\ln(4/3)}{\ln(13/9)}} \approx 1.44440557.$$ The proof of this result uses the following lemma. Let $x(n) = \ln(I(n^2))/\ln(I(n))$. If $\gcd(a, b) = 1$, then $$\min(x(a), x(b)) < x(ab) < \max(x(a),x(b)).$$ Proof of Lemma: First, note that $x(a) \neq x(b)$ (since $\gcd(a , b) = 1$). Without loss of generality, we may assume that $x(a) < x(b)$. Thus, since $$x(n) = \frac{\ln(I(n^2))}{\ln(I(n))},$$ we have $$\frac{\ln(I(a^2))}{\ln(I(a))} < \frac{\ln(I(b^2))}{\ln(I(b))}.$$ This implies that $$\ln(I(a^2))\ln(I(b)) < \ln(I(b^2))\ln(I(a)).$$ Adding $\ln(I(a^2))\ln(I(a))$ to both sides of the last inequality, we get $$\ln(I(a^2))\bigg(\ln(I(b)) + \ln(I(a))\bigg) < \ln(I(a))\bigg(\ln(I(b^2)) + \ln(I(a^2))\bigg).$$ Using the identity $\ln(X) + \ln(Y) = \ln(XY)$, we can rewrite the last inequality as $$\ln(I(a^2))\ln(I(ab)) < \ln(I(a))\ln(I((ab)^2))$$ since $I(x)$ is a weakly multiplicative function of x and $\gcd(a, b) = 1$. It follows that $$\min(x(a), x(b)) = x(a) = \frac{\ln(I(a^2))}{\ln(I(a))} < \frac{\ln(I((ab)^2))}{\ln(I(ab))} = x(ab).$$ Under the same assumption $x(a) < x(b)$, we can show that $$x(ab) < x(b) = \max(x(a),x(b))$$ by adding $\ln(I(b^2))\ln(I(b))$ (instead of $\ln(I(a^2))\ln(I(a))$) to both sides of the inequality $$\ln(I(a^2))\ln(I(b)) < \ln(I(b^2))\ln(I(a)).$$ This finishes the proof. From this lemma, we note that $1 < x(n) < 2$ follows from $$I(n) < I(n^2) < {\left(I(n)\right)}^2$$ and $I(n^2) = {\left(I(n)\right)}^{x(n)}$. The trivial lower bound for $I(n)$ is $${\bigg(\frac{8}{5}\bigg)}^{1/2} < I(n).$$ Note that decreasing the denominator in the exponent gives an increase in the lower bound for $I(n)$. Proof of Result: We sketch a proof for the mentioned result here, as communicated to the author by Pascal Ochem. Suppose that $N = {q^k}{n^2}$ is an odd perfect number given in Eulerian form. We want to obtain a lower bound on $I(n)$. We know that $$I(n^2) = 2/I(q^k) > 2/(5/4) = 8/5.$$ We need to improve the trivial bound $I(n^2) < {\left(I(n)\right)}^2$. Let $x(n)$ be such that $$I(n^2) = {\bigg(I(n)\bigg)}^{x(n)}.$$ That is, $x(n) = \ln(I(n^2))/\ln(I(n))$. We want an upper bound on $x(n)$ for n odd. By the lemma, we consider the component r^s with r prime that maximizes $x(r^s)$. We have $$I(r^s) = \frac{r^{s + 1} - 1}{{r^s}(r - 1)} = 1 + \frac{1}{r - 1} - \frac{1}{{r^s}(r - 1)}.$$ Also, $$I(r^{2s}) = \frac{r^{2s + 1} - 1}{{r^{2s}}(r - 1)} = I(r^s)\bigg(1 + \left(\frac{1 - r^{-s}}{r^{s + 1} - 1}\right)\bigg).$$ So, $$x(r^s) = \frac{\ln(I(r^{2s}))}{\ln(I(r^s))} = \frac{\ln(I(r^s)) + \ln(1 + \left(\frac{1 - r^{-s}}{r^{s + 1} - 1}\right))}{\ln(I(r^s))},$$ from which it follows that $$x(r^s) = 1 + \frac{\ln(1 + \left(\frac{1 - r^{-s}}{r^{s + 1} - 1}\right))}{\ln(1 + \frac{1}{r - 1} - \frac{1}{{r^s}(r - 1)})}.$$ We can check that $$x(r^s) > x(r^t)$$ if $s < t$ and $r \geq 3$. Therefore, $x(r^s)$ is maximized for $s = 1$. Now, $$x(r) = 1 + \frac{\ln(1 + (1/(r(r + 1))))}{\ln(1 + (1/r))} = \frac{\ln(1 + (1/r) + (1/r)^2)}{\ln(1 + (1/r))} = \ln(I(r^2))/\ln(I(r)),$$ which is maximized for $r = 3$. So, $$x(3) = \ln(I(3^2))/\ln(I(3)) = \ln(13/9)/\ln(4/3) \approx 1.27823.$$ The claimed result then follows, and the proof is complete. Motivation for My Inquiry Note that, since q is prime with $q \equiv 1 \pmod 4$, then $q \geq 5$. In particular, when $k=1$, then we obtain $$I(q^k) = 1 + \frac{1}{q} \leq \frac{6}{5} \iff I(n^2) = \frac{2}{I(q^k)} \geq \frac{5}{3},$$ from which we get $$I(n) > \bigg(I(n^2)\bigg)^{1/{x(3)}} \geq \bigg(\frac{5}{3}\bigg)^{\frac{\ln(4/3)}{\ln(13/9)}} \approx 1.4912789897463723558.$$ Here is my question: Would it be possible to tweak the argument in this post in order to come up with a proof for the implication $$k=1 \implies I(n) > \frac{3}{2}?$$

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2018-10-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018922 | MSE-2973765

Divisibility rule

Example: 2^1=2 --> $2\mid2$ If a number has their last digit divisible by 2, than the number is divisible by 2 2^2=4--> $4\mid2$, $4\mid4$ If a number has their last two digit divisible by 4, than the number is divisible by 4 2^3=8--> $8\mid2$, $8\mid4$, $8\mid8$ If a number has their last three digit divisible by 8, than the number is divisible by 8 and so on... How would you word this into a conjecture?

论坛问题帖 | has_answers | user608997 | 2018-10-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018923 | MSE-2977138

If x and y are solitary numbers satisfying $\gcd(x,y)=1$, under what conditions does it follow that xy is also solitary?

Let $\sigma(z)$ denote the sum of divisors of $z \in \mathbb{N}$. Denote the abundancy index of z by $I(z) = \sigma(z)/z$. If the equation $I(z)=I(a)$ has the lone solution $z=a$, then a is said to be solitary. Greening's Theorem states that if $\gcd(b,\sigma(b))=1$, then b is solitary. (Note that the converse does not necessarily hold.) Here is my question: If x and y are solitary numbers satisfying $\gcd(x,y)=1$, under what conditions does it follow that xy is also solitary? The latest paper on solitary numbers that I could find on Google seems to be New families of solitary numbers by Paul A. Loomis. However, my present inquiry is not covered in this paper.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2018-10-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018924 | MSE-2983179

$\sum_i X_i^2$ has χ^2_{n} distribution and X_i i.i.d. imply X_i normal

Let $X_1,\ldots,X_n$ be i.i.d. random variables with distribution F. It is known that if F is the standard normal distribution then $$ S:=\sum_{i=1}^n X_i^2 $$ has a chi square distribution with n degrees of freedom. I remeber that the converse is an open problem: if S has a chi square distribution with n degrees of freedom then F has to be the standard normal. Do you have some references on this problem? (I remember that some instances has been solved, but I couldn't find them anymore)

论坛问题帖 | no_answers_at_retrieval | Paolo Leonetti | 2018-11-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018925 | MSE-2993265

A weaker conjecture than a known conjecture

I really apologize if my question is not appropriate here, though I hope it is. Let C be any known conjecture in mathematics, which is still open. Let D be another conjecture such that a positive answer to conjecture C implies a positive answer to conjecture D. Now assume that one proved conjecture D. Do you think that mathematicians working on conjecture C will find such a result interesting? Of course, if one has found a counterexample to conjecture D, then conjecture C is false, but this is not the situation I am asking about. Thank you very much!

论坛问题帖 | has_answers | user237522 | 2018-11-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018926 | MSE-3006379

The sum of all divisors of N is a 2 potency iff N is a product of different Mersenne primes

As far as I have controlled: $\sigma(a)=2^n$, for some $n\in\mathbb N \iff $ a is a product of different Mersenne primes. The $\Leftarrow$-part is an immediate consequence of that σ is multiplicative, but the other part seems more complicated, if it is true. Maybe this is interesting in the view of that it is unknown if there are only a finite number of Mersenne primes or not.

论坛问题帖 | has_answers | Lehs | 2018-11-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018927 | MSE-3017227

What proportion of the positive integers satisfies $\gcd(n^2,\sigma(n^2))>\sigma(n)$, where σ is the sum-of-divisors function?

The title says it all. What proportion of the positive integers satisfies $\gcd(n^2,\sigma(n^2))>\sigma(n)$, where σ is the sum-of-divisors function? I tried using Sage Cell Server to search for positive integers n satisfying the said inequality, in the range $1 \leq n \leq {10}^6$, but it was unable to find any. I dare to conjecture that the proportion in question is zero, but I have no proof. Updated November 28 2018 I got the same output as Julian, after running the following (presumably correct) GP code in Sage Cell Server: for (x=1, 1000000,if(gcd(x^2,sigma(x^2))>sigma(x),print(x))) Updated November 29 2018 The number $n=1$ satisfies $$\gcd(n^2,\sigma(n^2))=\sigma(n)$$ while almost all integers $n>1$ (except for the 111 numbers less than ${10}^6$ mentioned by Julian in his answer below) satisfy $$\gcd(n^2,\sigma(n^2))<\sigma(n).$$ Therefore, I think it might indeed be possible to prove that the set $$\bigg\{n \in \mathbb{N} \mid \gcd(n^2,\sigma(n^2)) > \sigma(n)\bigg\}$$ has asymptotic density 0. But then again, as I have already mentioned, I have no proof.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2018-11-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018928 | MSE-3027572

For all $n>1$ there are positive $a+b=n$ such that $a+ab+b\in\mathbb P$

For all integers $n>1$ there are positive integers a,b such that $a+b=n$ and such that $a+ab+b\in\mathbb P$. Tested for all $n\leq 1,000,000$. Hopefully, someone can explore and explain the heuristics about this conjecture. This question is related to: Any odd number is of form $a+b$ where a^2+b^2 is prime Does every power of two arise as the difference of two primes? Most even numbers is a sum $a+b+c+d$ where $a^2+b^2+c^2=d^2$ Natural numbers large enough can be written as $ab+ac+bc$ for some $a,b,c>0$ $\{a+b|a,b\in\mathbb N^+\wedge ma^2+nb^2\in\mathbb P\}=\{k>1|\gcd(k,m+n)=1\}$ Even numbers has the form $a+b$ where $\frac{a^2+b^2}{2}$ is prime Is every positive integer greater than 2 the sum of a prime and two squares? It's about a relation $R\subseteq \mathbb N^m$, a function $f:\mathbb N^m\to \mathbb N$, and an image of a restriction $\operatorname{Im}(f|R)$. In Goldbachs conjecture the relation is $p,q\in\mathbb P$, the function is $(p,q)\mapsto p+q$ and the image of the restriction is $2\mathbb N\setminus\{2\}$. Maybe some of the conjectures can be generalized?

论坛问题帖 | has_answers | Lehs | 2018-12-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018929 | MSE-3030627

Does the infinite product of the reciprocals of a decreasing (or increasing) function equal zero?

I've made a short document explaining what I've just claimed. I'd like to know if the criteria for the infinite product to be zero is enough to hold for all decreasing and increasing functions. Theorem.

论坛问题帖 | has_answers | muhammad | 2018-12-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018930 | MSE-3048705

Every 'decreasing' or 'increasing' infinite sequence whose sum converges contains at least one term of magnitude 0

Note on the title: This conjecture is not restricted to real numbers, 'decreasing' and 'increasing' were used because of the character limit. The actual conjecture is that every sequence whose terms strictly increase or decrease in magnitude contains at least one term whose magnitude is 0. Warning: the following contains a gratuitous excess of '∞' it is recommended that you be familiar with at least one of the following: extended real/complex numbers (projective real, hypereal, Riemann sphere, etc.), non-standard numbers, or "Indian" infinities (Jaina infinities, Bhaskara, etc.). Let $\mathbb{U}$ denote a universe (e.g. $\mathbb{R}^n,\mathbb{C},\mathbb{H}\setminus\mathbb{R},\ldots$ etc.), where $u\in\mathbb{U}\implies |u|\neq\infty$ Definition: finite - a number* $z\in\mathbb{U}$ is finite iff $|z|\in(0,\infty)$ Definition: infinitesimal - a number $z\in\mathbb{U}$ is infinitesimal iff $\forall w\in\mathbb{U}\setminus\{\mathbb{U}_0\}.|z|<|w|$, where $\mathbb{U_0}$ is the additive identity (zero) of $\mathbb{U}$ Definition: infinite - a number $z\in\mathbb{U}$ is infinite iff $\forall w\in\mathbb{U}.|z|>|w|$ Definition: convergence - the sum of an infinite sequence $A\subseteq \mathbb{U}$ converges iff $\sum A\in\mathbb{U}$ Let A be an infinite sequence in $\mathbb{U}$, indexed by a set $I\subseteq\mathbb{N}\cup\{\infty\}$ where: $$A=\bigcup_{i\in I}\{a_i\}:\qquad\forall i,j\in I.i<j\implies(|a_i|\leq|a_j|\oplus |a_i|\geq|a_j|)$$ and where $\lim_{i\to n}a_i=a_n$, i.e. A is equal to its own limit at every point and is strictly increasing xor decreasing in magnitude. Conjecture: $$\sum_{i\in I}a_i\in\mathbb{U}\implies\exists k\in I:|a_k|=0$$ Reasoning: The sum of infinitely many finite numbers is always infinite (proof needed). Thus, if A contains only finite terms, then the sum of A does not converge. If a_1 is finite and A is increasing, then either all $a\in A$ are finite, or $|a_\infty|=\infty$. In the case of the former, $\sum A$ is the sum of infinitely many finite terms, and thus infinite. In the case of the latter, A contains at least one infinite term, therefore $|\sum A|=|a_1|+\cdots+\infty=\infty$. This is intuitively reinforced (though not proven) by treating the sum of A as the integral of a continuous function.** If A is increasing and the sum of A converges, then the first term must be infinitesimal. Since the magnitude of a number is given by a standard, real number, it follows that $|a_1|=0$. If A is decreasing and the sum of A converges, then $|a_\infty|=0$. Otherwise, $|a_\infty|$ is finite, making all $a\in A$ finite; if this were the case, then sum of A would be the sum of A would not converge. While this is at least true for all integer sequences (unless there is a proof that the sum of an infinite sequence of increasing integers converges), I'm not absolutely certain that it holds for all sequences. If it does, the how should I prove this? Can I prove this? I'm particularly skeptical about what would happen if I applied this to sequences of matrices or to sequenences with multiple infinite terms (e.g. where $|a_1|=|a_\infty|=\infty$ but $|a_1+a_\infty|=0$) *the term "number" can be replaced with "element" as this can also apply to the elements of any $\mathbb{U}$ equipped with a unary operation $|\cdot|:\mathbb{U}\to\mathbb{R}\cup\{\infty\}$ for which $u\in\mathbb{U}\implies|u|\neq\infty$ **Imagine a function $f:\mathbb{R}\to\mathbb{U}$ such that $n\in\mathbb{N}\implies f(n)=a_n$ and $\forall x,y,z\in\mathbb{R}. x<y<z\implies |f(x)|\leq|f(y)|\leq|f(z)|$. For any such function $\int_0^\infty f(x)\ dx=\infty$, even if $f(\infty)$ is finite (proof needed). This is actually one way to show that $\int\frac{1}{z}\ dx=\ln{z}$.

论坛问题帖 | has_answers | R. Burton | 2018-12-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018931 | MSE-3051098

On an interesting assertion in the OeisWiki page on multiply-perfect numbers

The following (interesting) assertion appears in the OeisWiki page on multiply-perfect numbers: ---------- Forwarded message ---------- From: Georgi Guninski <guninski@guninski.com> To: Sequence Fanatics Discussion list <seqfan@list.seqfan.eu> Cc: Date: Mon, 16 Jul 2012 13:14:33 +0300 Subject: [seqfan] Re: Reference that "A027687 4-perfect numbers" is finite Thank you. Asked because an odd perfect number and infinitely mersenne primes implies 4-perfect numbers are infinite (and a lot of other 2k-perfect numbers) - take the product of the OPN and coprime to it EPN. On the other hand 4-perfect being finite and infinitely mersenne primes implies no OPN. What is the reason to believe all 4-perfect are discovered (even if they are finite)? (This post is taken from the following SeqFan thread.) Honestly, I cannot seem to wrap my head around the first assertion (and therefore, also the second). Why is it that the existence of an odd perfect number and infinitely many Mersenne primes implies that there are infinitely many 4-perfect numbers (and a lot of other $2k$-perfect numbers)? It says "take the product of the OPN and coprime to it EPN". However, an OPN and an EPN may not always be coprime, as a Mersenne prime (for example, 3) may divide an OPN. (See this MSE question.) Note: OPN = Odd Perfect Number, EPN = Even Perfect Number

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2018-12-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018932 | MSE-3055907

Probability for composite n to have prime factor $\geq \sqrt n$

Let $\operatorname{GPF}(n)$ denote the largest prime factor of $n\in\mathbb N_{>1}$. My computer tests for intervalls $[m,n]$, where $n<10,000,000$, suggests that the probability $\operatorname{P}(\operatorname{GPF}(n)\geq\sqrt n)$ is asymptotically equivalent with a constant $\approx\frac{8}{11}$. Can this be proved, heuristically understood or falsified?

论坛问题帖 | has_answers | Lehs | 2018-12-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018933 | MSE-3056044

I have found a way of computing Euler's number. Is there any possible intuition of how that might be the case?

So a few days ago I just kind of messed around with my calculator, when I had an idea about a new continued fraction. I inputted it, and I found that it converged really quickly, and, quite wondrously, converged to the base of the natural logarithm. Then, the day later, I tried computing a similar continued fraction, this time starting with the number 2 instead of 1, without the increment of the numerator by 1. My first computation My second computation Interestingly, the second computation converges to EXACTLY the same values, except it's faster in the sense that the first method is delayed by 1 value per time. Does anybody possibly have an intuition / explanation as for why this might be the case? I've looked at the general ways of computing e: the sum of integer factorials and (1+1/n)^n, which I suspect might be more relevant to this.

论坛问题帖 | has_answers | Avtarás Karîm Elyméşer | 2018-12-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018934 | MSE-3057512

What is the sufficient and necessary conditions that $-1 \in G$, where G is a multiplicative group of a ring.

I am trying to prove the following conjecture. Let $(R, +,\times)$ be a finite ring with an identity. Let G be a subgroup of $(R,\times)$ with order d. Then $-1\in G$, if and only if $2\mid d$ or $\text{Char}(R)=2$. My attempt: $\Leftarrow$: Case 1:If $\text{Char}(R)=2$, then $-1=1$. Since G is a multiplicative group, the indentity $1\in G$. Hence $-1\in G$. Case 2: If $2\mid d$, then ...(I do not know how to prove in this case.) $\Rightarrow$: Suppose that $-1 \in G$. Then $(-1)^2=1\in G$. The multiplicative order of -1 is either 2 or 1. If $\text{Ord}(-1)=1$, then $-1=1$. Consequently, $\text{Char}(R)=2$. If $\text{Ord}(-1)=2$, then it must have $\text{Ord}(-1) \mid \text{Ord}(G)$, i.e., $2\mid d$. ========== So does $2\mid d$ implies that $-1\in G$? If the conjecture does not hold, then can we have a sufficient and necessary condition of $-1 \in G$? Thanks to Arthur, we have the following result using the fact that -1 is the only element of order 2. Let $(R, +,\times)$ be an integral domain. Let G be a subgroup of $(R,\times)$ with order d. Then $-1\in G$, if and only if $2\mid d$ or $\text{Char}(R)=2$.

论坛问题帖 | has_answers | Zongxiang Yi | 2018-12-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018935 | MSE-3060811

Conjecture about polynomials $f_n\in\mathbb Q[X_1,\dots,X_n]$ defining bijections $\mathbb N^n\to\mathbb N$

This is inspired by an answer of a question of mine: Bijective polynomials $f\in\mathbb Q[X_1,\dots,X_n]$ There is a polynomial $f_1\in\mathbb Q[X_1]$ which define a bijection $f_1:\mathbb N\to\mathbb N$, $f_1(X_1)=X_1$. There is a polynomial $f_2\in\mathbb Q[X_1,X_2]$ which define a bijection $f_2:\mathbb N^2\to\mathbb N$, $\displaystyle f_2(X_1,X_2)=\frac{(X_1+X_2)(X_1+X_2+1)}{2}+f_1(X_1)$. And as far as I understand and have tested there is a polynomial $f_3\in\mathbb Q[X_1,X_2,X_3]$ which define a bijection $f_3:\mathbb N^3\to\mathbb N$ $\displaystyle f_3(X_1,X_2,X_3)=\frac{(X_1+X_2+X_3)(X_1+X_2+X_3+1)(X_1+X_2+X_3+2)}{6}+f_2(X_1,X_2)$. This seems possible to generalize as $\displaystyle f_{n+1}(X_1,\dots ,X_{n+1})=f_{n}(X_1,\dots ,X_{n})+\frac{1}{(n+1)!}\prod_{k=1}^{n+1}\Big(k-1+\sum_{i=1}^{n+1}X_i\Big)$. This is a generalisation of the diagonalization in case of $n=2$ with "triangularization", "tetraederization" or higher. Now the conjecture is $\displaystyle f_n(X_1,\dots,X_n)$ define a bijection $\mathbb N^n\to\mathbb N$ for all $n>0$. Induction seems natural but how to prove that f_n is a bijection implies that f_{n+1} is a bijection? What I want are proofs, parts of proofs or counter proofs.

论坛问题帖 | has_answers | Lehs | 2019-01-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018936 | MSE-3069196

Conjecture about fixed prime divisors of polynomials with integer coefficients

While experimenting with random polynomials I've found this conjecture: A polynomial $f\in\mathbb Z[X]$ of degree n with co-prime coefficients have no fixed prime divisor $p> n$. A fixed prime divisor is a prime p such that $p|f(m)$ for all $m\in\mathbb Z$. Is this known? Proved? Or are there counterexamples?

论坛问题帖 | has_answers | Lehs | 2019-01-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018937 | MSE-3069987

Proof of $ x^2 + y = y^2 + x$ when $ x+ y =1$ and x is larger than y

I know that whatever numbers you choose for x and y and their sum equals to 1 will satisfy the equation $x^2 + y = y^2 + x$ Algebraic proof: Given: $x + y = 1$ $$LS = x^2+ y = (1-y)^2 + y = 1 - 2y+y^2 + y = y^2 - y + 1$$ $$RS = y^2 + x = y^2 + (1-y) = y^2 - y + 1$$ Therefore,$$ LS = RS $$ How can this be proved geometrically? (Ex. in a diagram of rectangular areas) I tried to add a square piece with side lengths y with a rectangle with side lengths x and x+y but I can't seem to prove it geometrically. Can someone help?

论坛问题帖 | has_answers | Andrew Nguyen | 2019-01-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018938 | MSE-3070332

Can Erdős-Turán $\frac{5}{8}$ theorem be generalised that way?

Suppose for an arbitrary group word w ower the alphabet of n symbols $\mathfrak{U_w}$ is a variety of all groups G, that satisfy an identity $\forall a_1, ... , a_n \in G$ $w(a_1, ... , a_n) = e$. Is it true, that for any group word w there exists a positive real number $\epsilon (w) > 0$, such that any finite group G is in $\mathfrak{U_w}$ iff $$\frac{\lvert\{(a_1, ... , a_n) \in G^n : w(a_1, ... , a_n) = e\}\rvert}{{|G|}^n} > 1 - \epsilon(w)?$$ How did this question arise? There is a widely known theorem proved by P. Erdős and P. Turán that states: A finite group G is abelian iff $$\frac{|\{(a, b) \in G^2 : [a, b] = e\}|}{{|G|}^2} > \frac{5}{8}.$$ This theorem can be rephrased using aforementioned

terminology as $\epsilon([a, b]) = \frac{3}{8}$. There also is a generalisation of this theorem, stating that a finite group G is nilpotent of class n iff $$\frac{|\{(a_0, a_1, ... , a_n) \in G^{n + 1} : [... [[a_0, a_1], a_2]... a_n] = e\}|}{{|G|}^{n + 1}} > 1 - \frac{3}{2^{n + 2}},$$ thus making $\epsilon([... [[a_0, a_1], a_2]... a_n]) = \frac{3}{2^{n + 2}}$. However, I have never seen similar statements about other one-word varieties being proved or disproved, despite such question seeming quite natural . . . Actually, I doubt that the conjecture in the main part of question is true. However, I failed to find any counterexamples myself.

论坛问题帖 | no_answers_at_retrieval | Chain Markov | 2019-01-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018939 | MSE-3072341

Multiplicative order of $a\bmod c^{k+1}$

I have some questions about moving from $\mathbb Z_{c^k}$ into $\mathbb Z_{c^{k+1}}$-specifically, with regard to the order of elements. Suppose a (which is coprime to c) has multiplicative order b in $\mathbb Z_{c^k}$. Now if a has multiplicative order p in $\mathbb Z_{c^{k+1}}$, then, $c^{k+1}\mid (a^p-1)\implies c^k\mid(a^p-1)\implies b\mid p$ So $p=bn$ for some positive integer n. $$\require{cancel}a^b\equiv 1\mod c^k\implies a^{bn}\equiv 1\mod c^{k+1}$$ $$(a^b)^n\equiv 1\mod c^{k+1}\implies ((a^b-1)+1)^n\equiv 1 \mod c^{k+1}$$ $$\sum_{i=0}^n\Biggl(\binom{n}{i}(a^b-1)^i1^{n-i}\Biggr)\equiv 1 \mod c^{k+1}$$ Since $c^k|(a^b-1):$ $$\binom{n}{0}(a^b-1)^01^{n-0}+\binom{n}{1}(a^b-1)^11^{n-1}+\sum_{i=2}^n\cancelto{0}{\Biggl(\binom{n}{i}(a^b-1)^i1^{n-i}\Biggr)}\equiv 1\mod c^{k+1}$$ $$1+n(a^b-1)\equiv 1\mod c^{k+1}$$ So, finally: $$n(a^b-1)\equiv 0\mod\ c^{k+1}$$ This gives us that $n|c$. More specifically, if $g.c.d.\Bigl(\frac{a^b-1}{c^k},c\Bigr)=d$, then $n=\frac{c}{d}$. Now, looking at something like the base-a Wieferich primes, there are clearly examples when $n\neq c$ (technically, Wieferich primes don't care specifically about the multiplicative order of the bases, but it happens frequently enough that $p-1$ is the base's order). I have two main questions. For any non-perfect power, c, can we always find an a and a k such that, if $ \operatorname{ord}(a)=b$ in $\mathbb Z_c$ and $ \operatorname{ord}(a)=n$ in Z_{c^k}, $n\neq bc^{k-1}$? And, second, (with c still being a non-perfect power$\neq2$) is it possible that, if, for some given a, that $ \operatorname{ord}(a)=b$ in $\mathbb Z_c$ and $ \operatorname{ord}(a)=bc$ in $\mathbb Z_{c^2}$, but for some $k>2$, $ \operatorname{ord}(a)\neq bc^{k-1}$ in $\mathbb Z_{c^k}$? Edit: I think I have an answer to the first question-it looks to be a bit more general. If we look at the set of elements in $\mathbb Z_{c^{k+1}}$ that are invertible, we have a group under the operation of multiplication- denote it by $V_{c^{k+1}}$. Of course, these are exactly the elements that are coprime to c^{k+1} so the order of this group is $$\phi(c^{k+1})=c^{k+1}\prod_{p:p=prime,p\mid c^{k+1}}\Bigl(1-\frac{1}{p}\Bigr)=c^{k+1}\prod_{p:p=prime,p\mid c}\Bigl(1-\frac{1}{p}\Bigr)=\\\frac{c^{k+1}}{\prod_{p:p=prime,p\mid c}p}\prod_{p:p=prime,p\mid c}(p-1)=d\prod_{p:p=prime,p\mid c}(p-1)$$ where d is clearly an integer. Now, consider any prime factor, q, of $p_{min}-1$ (where p_{min} is the smallest prime that divides c). $$q\mid (p_{min}-1)\mid\biggl(\prod_{p:p=prime,p\mid c}(p-1)\biggr)\mid \phi(c^{k+1})=\Bigl|V_{c^{k+1}}\Bigr|$$ So $q\mid \Bigl|V_{c^{k+1}}\Bigr|$. But by Cauchy's theorem for groups, this means that $V_{c^{k+1}}$ has an element of order q- denote it by x. Clearly the multiplicative of order of x in $\mathbb Z_{c^{k+1}}$ is also q. Referring back to the way q was defined, it's clear that q and c are coprime (since q must be less than any prime factor of c). But, if x has order b in $\mathbb Z_{c^k}$, then the order of x in $\mathbb Z_{c^{k+1}}$ is (as shown before) of the form bn where n is some divisor of c. That means $bn=q$. Since q and c are coprime, $n=1$, and $b=q$, i.e., the order of x is the same in $\mathbb Z_{c^{k+1}}$ as it is in $\mathbb Z_{c^k}$. The reason this seems more general is that it works for any k. Of course, there's a small flaw when it comes to c being even, i.e. that $p_{min}-1=2-1=1$ always. But, the above argument could still easily be extended to any even numbers where the 2^{nd} smallest prime divisor isn't a Fermat prime (just let q be some prime other than 2 that divides the second smallest prime-1). Even if the second smallest prime divisor is Fermat, we'll still have $\operatorname{ord}(a)=2b\neq bc^k$ (I just can't guarantee an element with the same order in $\mathbb Z_{c^k}$ and $\mathbb Z_{c^{k+1}}$ exists). That seems to just leave $c=2$. Still have no clue how to deal with the second question. Edit 2: The above argument (in the Edit) can be tweaked a bit- once we have an x of order q in $\mathbb Z_{c^{k+1}}$, instead of returning to $\mathbb Z_{c^k}$ and examining x's order there, we can look directly at $\mathbb Z_c$ instead. If the order of x in $\mathbb Z_c$ is b, then by extending the reasoning presented above, the order of x in $\mathbb Z_{c^{k+1}}$ is bN, where N is a divisor of c^k. That means $bN=q$ and (since q and c are coprime) $b=q$ but now b is the order of x in $\mathbb Z_c$. This reasoning can be extended so that for every $1\leq h\leq k+1$, the order of x in $\mathbb Z_{c^h}$ is the same. This seems to be suggesting that the answer to the second question is generally no- but that would be to assume that elements that do not have order bc^k in $\mathbb Z_{c^{k+1}}$ are only constructed by the method in the first Edit (it also isn't true in the case when $c=2$ just by the fact that all odd numbers have order 1 in $\mathbb Z_2$).

论坛问题帖 | no_answers_at_retrieval | CardioidAss22 | 2019-01-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018940 | MSE-3079407

A conjecture about irreducible polynomials with integer coefficients

Let $f\in\mathbb Z[X]$, define $\operatorname{P}^+(f)$ as the number of primes >0 that f assumes at distinct integral arguments. Theorem: If $f\in\mathbb Z[X]$ is non constant and reducible of degree n, then $\operatorname{P}^+(f)\leq n$. And for all n there are non constant reducible polynomials of degree n such that $\operatorname{P}^+(f)=n$. [Acta Arith.,104.2 (2002) 117-127.] Most polynomials are irreducible but using the theorem for an irreducibility test would be inefficient since a lot of irreducible polynomials has a fixed divisor >1 and wouldn't pass the test. A conjecture related to the conjecture of Bunjakowsky states: Conjecture: If $ f\in\mathbb Z[X]$ is non constant and irreducible, then $f(a)/d$ assumes primes for an infinit number of distinct integral arguments a, where d is the largest fixed divisor of f. Irreducibility of Polynomials Whose Coefficients are Integers Page 32. This makes me wonder if the following hypothesis is true: $f\in\mathbb Z[X]$ of

degree $n>0$ with coprime coefficients is irreducible, iff $\;\operatorname{P}^+(d^{-1}\cdot f)> n$ or $\;\operatorname{P}^+(d^{-1}\cdot (-f))> n$, where d is the greatest fixed divisor of f. P^+ is extended above and defined even for integer-valued polynomials with rational coefficients. Proofs or counter-examples? With a test program using the hypothesis on Eisenstein polynomials f with random coefficients between -19 and 19 and random degree between 2 and 5, testing both f and $-f$, resulted in no miss in $1,000,000$ polynomials. The only drawback is the risk of overflow when evaluating the polynomials for higher degrees and greater coefficients.

论坛问题帖 | has_answers | Lehs | 2019-01-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018941 | MSE-310026

On Ramanujan's Question 359

In JIMS 4, p.78, Question 359 was asked by Ramanujan. (See The Problems Submitted by Ramanujan to the Journal of the Indian Mathematical Society, p. 9, by Bruce Berndt, et al.) If, $$\sin(x+y) = 2\sin\big(\tfrac{1}{2}(x-y)\big)\tag1$$ $$\sin(y+z) = 2\sin\big(\tfrac{1}{2}(y-z)\big)\tag2$$ prove that, $$\big(\tfrac{1}{2}\sin x\cos z\big)^{1/4}+\big(\tfrac{1}{2}\cos x\sin z\big)^{1/4} =\big(\sin 2y)^{1/12}\tag3$$ The example by Ramanujan was, $$\begin{aligned} x &= \frac{\pi-\arcsin\big((\sqrt{5}-2)^3(4+\sqrt{15})^2\big)}{2}=1.094\dots\\ y &=\frac{\arcsin\big(\sqrt{5}-2\big)}{2}=0.119\dots\\ z &=\frac{\arcsin\big((\sqrt{5}-2)^3(4-\sqrt{15})^2\big)}{2}=0.0001\dots \end{aligned}$$ Ten years later, a 3-page proof was given in JIMS 15, p.114-117. I got an email asking if there was a shorter proof. Considering the problem, I observed the following. Given the quartic, $$a^3w^4+(1-3a^2)w^3+3a(1-a^2)w^2+a^2(3-a^2)w-a=0\tag4$$ with $a=\tan(\color{blue}y/4)$. Define, $$x=4\tan^{-1} u\\z=4\tan^{-1} v$$ where u,v are two appropriate roots of the quartic, then we get the same bizarre relation as Ramanujan, $$\big(\tfrac{1}{2}\sin x\cos z\big)^{1/4}+\big(\tfrac{1}{2}\cos x\sin z\big)^{1/4} =\big(\sin 2\color{blue}y)^{1/12}\tag5$$ Equivalently, those two roots u,v obey, $$\left(\frac{1}{2}\,\frac{4(u-u^3)}{(u^2+1)^2}\frac{v^4-6v^2+1}{(v^2+1)^2} \right)^{1/4}+\left(\frac{1}{2}\,\frac{4(v-v^3)}{(v^2+1)^2}\frac{u^4-6u^2+1}{(u^2+1)^2} \right)^{1/4}=\big(\sin 2\color{blue}y\big)^{1/12}\tag6$$ For example, let $y=1$, so $a=\tan(1/4)$, then u,v are the two real roots of (4). Question: Anyone knows a short proof for (3) and (6)?

论坛问题帖 | has_answers | Tito Piezas III | 2013-02-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018942 | MSE-3102286

A consequence of assuming the Descartes-Frenicle-Sorli Conjecture on odd perfect numbers

Let x be a positive integer. Denote the sum of divisors of x by $$\sigma(x) = \sum_{d \mid x}{d},$$ and the deficiency of x by $$D(x) = 2x - \sigma(x).$$ A number N is said to be perfect if $\sigma(N)=2N$. Euler proved that an odd perfect number, if one exists, must have the so-called Eulerian form $N = q^k n^2$, where q is the special prime satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. The Descartes-Frenicle-Sorli Conjecture on odd perfect numbers predicts that $k=1$. In the paper titled Conditions equivalent to the Descartes-Frenicle-Sorli Conjecture on odd perfect numbers - Part II, the following results are proved: If $N = q^k n^2$ is an odd perfect number given in Eulerian form, then $k=1$ if and only if $$N = \bigg(\frac{q(q+1)}{2}\bigg)\cdot{D(n^2)}.$$ If $N = q^k n^2$ is an odd perfect number given in Eulerian form, then $k=1$ if and only if $$N = \frac{n^2 \sigma(n^2)}{D(n^2)}.$$ Putting these two results together, we have that If $N = q^k n^2$ is an odd perfect number given in Eulerian form, then $k=1$ if and only if $$\bigg(\frac{q(q+1)}{2}\bigg)\cdot{D(n^2)} = N = \frac{n^2 \sigma(n^2)}{D(n^2)}.$$ This last equation implies that $k=1$ if and only if $$\bigg(D(n^2)\bigg)^2 = \frac{n^2}{(q+1)/2}\cdot\frac{\sigma(n^2)}{q}.$$ Here is my question: Does the equation $$\bigg(D(n^2)\bigg)^2 = \frac{n^2}{(q+1)/2}\cdot\frac{\sigma(n^2)}{q}$$ imply that $(q+1)/2$ and $\sigma(n^2)/q$ are both squares?

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2019-02-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018943 | MSE-3125022

Sum of digits of a^b equals ab

The following conjecture is one I have made today with the aid of computer software. Conjecture: Let $s(\cdot)$ denote the sum of the digits of $\cdot$ in base 10. Then the only integer values $a,b>1$ that satisfy $$s(a^b)=ab$$ are $(2,2),(3,3),(3,6),(3,9)$ and $(3,27)$. Remarks: The number of digits of an integer n is $1+\lfloor\log_{10}n\rfloor$ where $\lfloor\cdot\rfloor$ denotes the floor function. This means that $s(a^b)<9+9\lfloor b\log_{10} a\rfloor$ as each digit takes a value of at most 9. From this plot it is evident that the equality will never hold for $a\ge 9$, after checking the first 21 values of b. Therefore it suffices to consider $2\le a\le 8$. I have excluded the case $b=1$ as it is trivial - it forces a to be a single-digit integer. In PARI/GP the code is given by mfun(b)={for(i=2,8,for(j=2,b,if(sumdigits(i^j,10)==i*j,print(i," ",j))));} and tests up to $b\le 10^6$ verify the above conjecture. Aside: The equation $s(ab)=a+b$ is much simpler to solve. From here, the upper bound $a\le 23$ is apparent and for these values of a, we can form the upper bound $b\le 22$ as there will be no solutions when the red line lies above the blue lines. This 23×22 grid can be computed through software and it is found that the only solutions are $$(a,b)=(2,2),(3,6),(6,3).$$ The similarity in the first two solutions with the conjectured solutions may only be coincidental. Advances on this will be appreciated.

论坛问题帖 | has_answers | TheSimpliFire | 2019-02-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018944 | MSE-3125087

Could prime numbers be defined like this?

While playing with prime numbers, I found the following definition. Let p be an integer. Then p is a prime number if and only if there is some integer $b \neq 1$ such that $$ \frac{b^p - 1}{b - 1} $$ is also a prime number. It is easy to show that the primality of p is a necessary condition for primality of $(b^p - 1)/(b - 1)$. I am however stuck to prove that, for given p, there is always at least one prime number of the form $(b^p - 1)/(b - 1)$. Is my definition correct, and if so, how to prove the second part? Any suggestion is welcome.

论坛问题帖 | has_answers | DaBler | 2019-02-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018945 | MSE-3128677

Is This a New Property I Have Found Pertaining to Mersenne Primes?

While playing with Mersenne numbers, I found the following property distinguishing Mersenne prime numbers from Mersenne composite numbers. A Mersenne number, $\text{M}p$, is a number of the form $2^p - 1$, where p is prime. Property For $p > 2$, Mersenne primes can be expressed as \begin{align*} \text{M}p = \frac{a^3 + b^3}{a + b}\text{,} \end{align*} where a and b are integers, $a \neq -b$, with exactly 12 different solutions. So far, also $\operatorname{gcd}(a,b)=1$ holds for Mersenne primes. Mersenne composites have either no integer solution or more than 12 solutions (24 so far). Also $\operatorname{gcd}(a,b)=1$ does not hold if the integer solutions exist so far. Examples \begin{align*} \text{M}5 &= \frac{6^3 + 5^3}{11} = 31 \\ \text{M}7 &= 7^3 - 6^3 = 127 \end{align*} The M11 has no integer solution for (a,b). The M37 has 24 solutions and also $\operatorname{gcd}(a,b)=1$ does not hold. Remarks Except the M2, twelve solutions exist for each Mersenne prime. If (a,b) is a solution, then also $(-a,-b)$, (b,a), and $(-b,-a)$ are. Since \begin{align*} \frac{a^3 + b^3}{a + b} = a^2 -ab + b^2 \text{,} \end{align*} each Mersenne prime has an ellipse intersecting integer grid associated with it. For example, $-a^2 + ab - b^2 + 127 = 0$ is the ellipse for M7. Results Solutions for the first few Mersenne numbers: $$\begin{matrix} p & \text{M}p & (a,b) \\ \hline 2 & 3 & (1,2) \\ 3 & 7 & (1,-2), (1,3), (2,3) \\ 5 & 31 & (1,-5), (1,6), (5,6) \\ 7 & 127 & (6,-7), (6,13), (7,13) \\ 11 & 2047 & \text{no solution} \\ 13 & 8191 & (1,-90), (1,91), (90,91) \\ 17 & 131071 & (6,-359), (6,365), (359,365) \\ 19 & 524287 & (83,-679), (83, 762), (679, 762) \\ 23 & 8388607 & \text{no solution} \\ 29 & 536870911 & \text{no solution} \\ 31 & 2147483647 & (4698, 43813), (4698,48511), (43813, 48511) \\ 37 & 137438953471 & \text{24 solutions} \\ 41 & 2199023255551 & \text{no solution} \\ 43 & 8796093022207 & \text{no solution} \\ ... & ... & ... \\ \end{matrix}$$ I have verified the conjecture using WolframAlpha for all p below 100. Question Can you confirm this result? Is this known? Any feedback is welcome.

论坛问题帖 | has_answers | DaBler | 2019-02-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018946 | MSE-3131506

Which other unsolved problems, have necessary restrictions on the prime gaps?

We all know of Unsolved problems, like Goldbach,Legendre, and Grimm's conjectures. Goldbach has the necessary condition of: There exists a prime between n and $2n-2$, which means prime gaps are limited to $n-3$ Legendre has a necessary condition of: Ratio's of primes never exceed $\left(\frac{n+2}{n} \right)^2$ Grimm's has a necessary condition (infinitely often) of: A prime gap never exceeds in length the number of primes less than the prime proceeding the gap. My Question is: What other unsolved problems, have necessary conditions on the size of prime gaps?

论坛问题帖 | has_answers | user645636 | 2019-03-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018947 | MSE-313282

Finding a counterexample to a Prime Factorization Conjecture

Let $\mathbb{Z}_{\geq 2}$ be the set of natural numbers starting at 2: $$\mathbb{Z}_{\geq 2}= \{2, 3, 4, 5,\ldots\}.$$ An natural number's prime factorization is odd if the total number of primes in its factorization is odd. It is even if the total number of primes in its factorization is even. Let $N(k) = \{j \mid j\in \mathbb{Z}_{\geq 2}, j\leq k\text{, the prime factorization for j is odd}\}$. Let $n(k) = |N(k)|$ Let $A(k) = \{j \mid j\in \mathbb{Z}_{\geq 2}, j\leq k\text{, the prime factorization for j is even}\}$. Let $a(k) = |A(k)|$ Conjecture: $n(k) \geq a(k)$ for all prime numbers k in $\mathbb{Z}_{\geq 2}$.

论坛问题帖 | has_answers | user63813 | 2013-02-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-018948 | MSE-3133571

Equivalence of Frankl's Conjecture

Frankl’s conjecture is one of the most famous problems in combinatorics. Frankl's conjecture claims: For every finite non-empty set A and for every Frankl's family F on A exists $a\in A$ such that $$|\{X\in F | a\in X \}|\geqslant \frac{|F|}{2}.$$ How can be proved that Frankl's conjecture is equivalent to next statement: For every finite lattice L exists $a \in\mathcal{M}(L),$ such that $|a\downarrow|\leqslant \frac{|L|}{2},$ where $\mathcal{M}(L)$ is the set of all strictly $\wedge$-irreducible elements of L and $a ↓ = \{b ∈ L | b ≤ a\}?$

论坛问题帖 | no_answers_at_retrieval | math_ | 2019-03-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018949 | MSE-3134700

Almost a prime number recurrence relation

For the number of partitions of n into prime parts $a(n)$ it holds $$a(n)=\frac{1}{n}\sum_{k=1}^n q(k)a(n-k)\tag 1$$ where $q(n)$ the sum of all different prime factors of n. Due to https://oeis.org/A000607. I have found that $$a(p_n)\approx a(p_{n-2})+a(p_{n-1})\tag 2$$ and conjecture the asymptotic relation. $$\log a(p_n)\sim

\log \big(a(p_{n-2})+a(p_{n-1})\big)\tag 3$$ On x-axis the prime numbers p_n are plotted. The blue lines correspond to $a(p_n)$ and the red lines correspond to $a(p_{n-2})+a(p_{n-1})$. On the oeis site above there is also a formula $$a(n)\sim e^{2\pi\sqrt{n/\log n}\,/\sqrt{3}}\tag 4$$ but I don't know if this really helps? Can this (reformulated) conjecture be proved?

论坛问题帖 | no_answers_at_retrieval | Lehs | 2019-03-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018950 | MSE-3148772

Group (mathematics) Conjecture

Given $(G,•)$ as a Group with finite set G, operator •. Define: subset $S \subset G$ is called the core of the Group if and only if $$ \{ x•y ~|~ x \in S, y \in S \} = G \setminus S$$ Conjecture 1: if G has at least two elements, core exists. Conjecture 2: if core exists, any two cores have the same cardinality. Conjecture 2.1: given prime $p \ge 3$, for $G = \{1,2,\ldots,p-1\}$ and operator • is 'multiplication modulo p', then any core $S \subset G$ has cardinality $(p-1)/2$. Example: $p = 7$, $S = \{3,5,6\}$ $1 \equiv 3*5 \equiv 6*6\pmod{7}$ $2 \equiv 3*3 \equiv 5*6\pmod{7}$ $4 \equiv 3*6 \equiv 5*5 \pmod{7}$ Question: anyone can find any (related) reference of the definition of core?

论坛问题帖 | has_answers | Kevin | 2019-03-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018951 | MSE-3159134

An explicit "formula" for the prime counting function?

It is known that $\log(p_1),\cdots,\log(p_n)$ are linearly independet over $\mathbb{Q}$, where p_i denotes the i-th prime. For a number $1 \le k \le n$ let $Log(k)$ denote the vector with respect to this basis for the numbers $1,\cdots,n$. For a subset A of $1,\cdots,n$ let $rank(A)$ denote the rank of the matrix build by numbers k in A of vectors $Log(k)$. Through experimentation I found the following explicit "formula" for prime-pi function: $$ \Pi(n)= (-1)^{n+1} \sum rank(A) \cdot (-1)^{|A|}$$ where A runs through the subsets of $1\cdots n$ of size $< n$. But I have no proof for this. The formula above has some similarity with the inclusion-exclusion principle: $$\left| \bigcup_{i=1}^n A_i\right| = \sum_{\emptyset\neq J\subseteq\{1,\ldots,n\}}(-1)^{|J|-1} \left |\bigcap_{j\in J} A_j\right|$$ But how does one define the sets A_i? The formula has also similarity with Euler characteristic. For a set of (possibly repeating) vectors, define: $$\chi(v_1,\cdots,v_n) = \sum_{A} rank(A) \cdot (-1)^{|A|}$$ where A runs through the subsets (with repetion) of $v_1,\cdots,v_n$. It is not difficult to show, that if those vectors are linearly independent then $$\chi=0$$ My conjecture is that if $\chi(v_1,\cdots,v_n)=0$ then $$\chi(v_1,\cdots,v_n,w)=0$$ for any vector w. This would prove the prime-counting formula. If you can think of a proof, that would be great. Related-but-not duplicate question: https://mathoverflow.net/questions/325880/is-this-line-of-thought-using-linear-algebra-to-get-number-theoretic-results-a Here is some sage code to play around with: MAXN=100 def Log(a,N=MAXN): return vector([valuation(a,p) for p in primes(N)]) def findsubsets(S,m): return Set(S).subsets(m) def getMatrixA(someSubset,N=MAXN): return matrix([Log(xx,N=N) for xx in someSubset]) def eulerCharPrimePi(n,N=MAXN): return (-1)^(n+1)*sum([getMatrixA(x).rank()*(-1)^k if k < n else 0 for k in range(0,n+1) for x in findsubsets(range(1,n+1),k)])

论坛问题帖 | has_answers | user276611 | 2019-03-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018952 | MSE-3159577

Conjecture about primes and the greatest common divisor

Conjecture: Given $m,n\in\mathbb N^+$, one odd and one even, there are two primes p,q such that $|mp-nq|=\gcd(m,n)$. I hope MSE can determine its validity. From time to time, when testing my growing math packages BigZ and Forthmath, I recognize some patterns which I can't prove or disprove (or even have the ambition to). I post them here with the hope that it will not annoy too much. I hope you can bear with it.

论坛问题帖 | has_answers | Lehs | 2019-03-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018953 | MSE-3164268

Sum on GCD and prime numbers

I was studying gcd then I encountered this sum $(1).$ A conjecture: If $(1)=1$ for any values of $N\ge3$, then N is a prime number. Let: $$f(N)=\frac{1}{N^{1-s}(N-1)}\sum_{j=1}^{N}(-1)^jj^s\frac{{N \choose j}}{gcd(j,N)}=1\tag1$$ where gcd is the greatest common divisor and $s\ge1$ If this sum is correct, can we uses it to test for prime numbers or look for a largest prime number?

论坛问题帖 | has_answers | user550260 | 2019-03-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018954 | MSE-3166140

Symmetry conditions for symmetric random vectors

While formulating the properties for a certain statistical model I'm dealing with, I came up with the following question (with credit going to MikeEarnest in comments for the proper formulation). A random variable X is symmetric (about zero) if X and $-X$ are identically distributed. This in particular applies to the case of a random vector. The question is then: Suppose $X=(X_1,X_2,\ldots, X_n)$ is a random vector such that $(X_{i(1)},X_{i(2)},\ldots,X_{i(m)})$ is symmetric for any $\{i(1),i(2),\ldots,i(m)\}\subseteq \{1,2,\ldots,n\}$. Does it follow that X is symmetric? This statement is probably too generic, given the lack of assumptions on X beyond symmetry. If I specialize to the case where each X_k can only take values ± 1, then my understanding is that the above is true when $m=1$ regardless of n. (This would imply that it is true for arbitrary m, since the subsets of $\{1,2,\cdots, n\}$ of size at most m includes the singletons.) However, if I move to X_k taking values in $\{1,0,-1\}$ then the case of $m=1$ fails (see the comments for a counter-example due to MikeEarnest) and I don't know what can be said about $m>1$. This gives me the following (admittedly imprecise) question: What assumptions on the maximum size m and the distribution of the random variables $\{X_k\}_{k=1}^n$ must be made in order for the above statement to be valid? I would also be interested in additional

counter-examples.

论坛问题帖 | has_answers | Semiclassical | 2019-03-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018955 | MSE-3166862

On a conjecture that two intersections are at $\big(\pm\frac{S}m, S\big)$ (about linear equations)

Suppose we have two linear equations, $y=mx$ and $y=-mx$, inserted on the same plane for $m>0$. These equations are going to have an intersection at $(0,0)$. Now, let $$\begin{align}S&=1+\sum_{i=2}^{m-1}\left(i-\frac12\right)\\ &\stackrel{\small\text{or}}{=}1+\left\{\bigg(2-\frac12\bigg)+\bigg(3-\frac12\bigg)+\cdots+\bigg(m-1-\frac12\bigg)\right\}.\end{align}$$ If we insert the equation, $y={m^2+1 \over 2}$, on the selfsame plane, then this horizontal line will pass through both the aforementioned linear equations, thus forming two more intersections. Conjecture: The two intersections are at $\big(\pm\frac{S}m, S \big)$, such that the positive x coordinate is on the RHS of the y axis, and the negative x coordinate is on the LHS of the y axis. This is my own conjecture actually... and I can't seem to prove it. The following is my attempt, in which I did make some progress... but it is not a proof (...yet). Attempt: $$\begin{align}S&=1+\left\{\bigg(2-\frac12\bigg)+\bigg(3-\frac12\bigg)+\cdots+\bigg(m-1-\frac12\bigg)\right\}\tag{given} \\ \\ 2\cdot S&=2+(4-1)+(6-1)+\cdots+(2m-2-1)\tag{$\times 2$} \\ &=\require{cancel}{\cancel2}+(4-1)+(6-1)+\cdots+(2m\cancel{-2}-1)\tag{$2-2=0$} \\ &=(4-1)+(6-1)+\cdots+(2m-1)\tag{simplifying} \\ &=\sum_{i=2}^m(2i-1).\tag{$\ast$}\end{align}$$ Here, we can use the following lemma to enable us to write Eq. $(\ast)$ in closed form: $$\sum_{j=1}^n(2j-1)=n^2.$$ The proof of this lemma can be found in Intermediate Mathematics: Proof by Induction in Page 8, Section 2, Example 3. By letting $n=m$ and $j=i$, we can rewrite this lemma and obtain the following: $$\begin{align}\sum_{i=1}^m(2i-1)&=(2\times 1-1)+\sum_{i=2}^m(2i-1) \\ \Leftrightarrow m^2&= 1+\sum_{i=2}^m(2i-1) \\ \Leftrightarrow \sum_{i=2}^m(2i-1)&= m^2-1 \\ \Leftrightarrow 2S&=m^2-1\end{align}$$ $$\boxed{\therefore S=\frac{m^2-1}{2}}\tag*{$\quad\quad\quad

infinity}$$ Therefore, evaluating an upper bound should be a simple as solving the equation $$L_n=L_{n+1}\stackrel{\text{in the example}}{\implies}L_n=\frac{L_n+1}{L_n+2}\implies L_n=\frac{-1+\sqrt5}{2}$$ Another example would be, for instance, the sequence defined recursively as follows $$M_{n+1}=\frac{M_n}{3}+\frac{1}{M_n}\qquad \text{where}\quad M_0=4$$ which converges to $\sqrt{\frac{3}{2}}$. This leads to Conjecture Suppose there's a recursive sequence $\{L_n\}^\infty_{n=0}$ that converges to k, where $L_{n+1}-L_n\to 0$ as n approaches infinity. Then, k is one of the solutions to the equation $$L_n=L_{n+1}$$

论坛问题帖 | has_answers | Dr. Mathva | 2019-04-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018958 | MSE-3181469

Conjecture on the growth of $q_1 = 1, q_{n+1}=q_n + f(q_n) $

This is a generalization of my answer to Calculate the limit of the following recurrent series in the form suggested by Will Jagy. $q_1 = 1, q_{n+1}=q_n + f(q_n) $ where $f(x) > 0$ and $f'(x) < 0$. I show that q_n is unbounded and derive a conjecture for the growth of q_n as a function of $f(x)$. To make my question explicit (since some people seem to want that): Is my conjecture on the growth of q_n true? My conjecture for $f(x) = \dfrac{a}{x^b} $ is $q_n \approx (a(b+1)n)^{1/(b+1)} $ and for $f(x) = \dfrac{a}{b^x} $ is $q_n \approx \dfrac{\ln(a\ln(b)n)}{\ln(b)}$. First, a proof that q_n is unbounded. $q_n > 1$ for $n > 1$ and q_n is increasing. If q_n is bounded, there is an L such that $q_n \le L$ for all n so that, since $f'(x) < 0$, $f(q_n) \ge f(L)$. Writing $q_{n+1}=q_n + f(q_n) $ as $q_{n+1}-q_n = f(q_n) $ and summing $\begin{array}\\ q_{n+1}-q_1 &=\sum_{k=1}^n (q_{k+1}-q_k)\\ &=\sum_{k=1}^n f(q_k)\\ &\ge\sum_{k=1}^n f(L)\\ &= nf(L)\\ \end{array} $ which is unbounded, a contradiction. Now to get an estimate for q_n in terms of $f(x)$. Suppose $q_n = g(n)$ where g is a reasonably smooth function (i.e., has at least a first derivative). Then $\begin{array}\\ q_{n+1}-q_n &=g(n+1)-g(n)\\ &=g'(n+c) \qquad\text{where } 0 \le c \le 1\\ &\approx g'(n) \qquad\text{this is the big assumption}\\ \end{array} $ Therefore $g'(n) \approx f(q_n) =f(g(n)) $. This is the functional equation that g has to satisfy. For example, if $f(x) = \frac{2}{x}$ as in the original problem, the equation is $g'(n) =\frac{2}{g(n)} $ so $2 = g(n)g'(n) =\frac12(g^2(n))' $. Integrating, $4n+c =g^2(n)$ so $g(n) = 2\sqrt{n+c_1} $. This agrees with my conjecture in my answer to the original question. More generally, if $f(x) = ax^{-b}$ where $a, b > 0$, the equation is $g'(n) =a(g(n))^{-b} $ or $a = g'g^b =\frac1{b+1}(g^{b+1})' $. Integrating, $a(b+1)n+c =g^{b+1}(n) $ so $g(n) =(a(b+1)n+c)^{1/(b+1)} $. As a check, simplifying with $c=0$, $g(n) = (a(b+1)n)^{1/(b+1)}$ so $\begin{array}\\ g(n+1)-g(n) &=(a(b+1)(n+1)^{1/(b+1)}-(a(b+1)n)^{1/(b+1)}\\ &=(a(b+1)n)^{1/(b+1)}((1+1/n)^{1/(b+1)}-1)\\ &=(a(b+1)n)^{1/(b+1)}(\frac1{(b+1)n}+O(\frac1{n^2}))\\ &=a^{1/(b+1)}((b+1)n)^{1/(b+1)-1}+O(\frac1{n^{2-1/(b+1)}})\\ &=a^{1/(b+1)}((b+1)n)^{-b/(b+1)}+O(\frac1{n^{2-1/(b+1)}})\\ &=\dfrac{a^{1/(b+1)}}{((b+1)n)^{b/(b+1)}}+O(\frac1{n^{1+b/(b+1)}})\\ \text{and}\\ f(g(n)) &=\dfrac{a}{g^b(n)}\\ &=\dfrac{a}{((a(b+1)n)^{1/(b+1)})^b}\\ &=\dfrac{a}{(a(b+1)n)^{b/(b+1)}}\\ &=\dfrac{a^{1/(b+1)}}{((b+1)n)^{b/(b+1)}}\\ \end{array} $ Another class of functions, suggested by Will Jagy, is $f(x) = \frac{a}{b^x}$ where $a > 0, b > 1$, the equation is $g'(n) =\frac{a}{b^{g(n)}} $ or $a = g'b^g = g'e^{g\ln(b)} =\frac1{\ln(b)}(e^{g\ln(b)})' =\frac1{\ln(b)}(b^g)' $. Integrating, $a\ln(b)n+c =b^g $ so $g(n)\ln(b) =\ln(a\ln(b)n+c) $ or $g(n) =\dfrac{\ln(a\ln(b)n+c)}{\ln(b)} $. If $a=1, b = e$ as in Will Jagy's example, $g(n) =\ln(n+c) $. To check, if $c = 0$, $g(n) = \dfrac{\ln(a\ln(b)n)}{\ln(b)}$ so $\begin{array}\\ g(n+1)-g(n) &=\dfrac{\ln(a\ln(b)(n+1))}{\ln(b)}-\dfrac{\ln(a\ln(b)n)}{\ln(b)}\\ &=\dfrac1{\ln(b)}(\ln(a\ln(b)(n+1))-\ln(a\ln(b)n))\\ &=\dfrac1{\ln(b)}\ln(1+1/n))\\ &=\dfrac1{n\ln(b)}+O(\frac1{n^2})\\ \text{and}\\ f(g(n)) &=\dfrac{a}{b^{g(n)}}\\ &=\dfrac{a}{b^{\frac{\ln(a\ln(b)n)}{\ln(b)}}}\\ &=\dfrac{a}{a\ln(b)n}\\ &=\dfrac1{\ln(b)n}\\ \end{array} $

论坛问题帖 | no_answers_at_retrieval | marty cohen | 2019-04-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018959 | MSE-3185268

How does one subtract from concatenation in order to prove that $4\times 5 + 67 = 45 + 6\times 7$?

I noticed that if we get the numbers 4, 5, 6 and 7, they have an interesting property! $$4 \times 5 + 67 = 45 + 6 \times 7\tag*{= 87.}$$ I then conjectured that these were the only four consecutive positive integers with this property. Hmm... Let the four integers be n, $n+1$, $n+2$ and $n+3$. Then we have $$\begin{align} &n(n+1) + (n+2)\| (n+3) \\ =\;&n\| (n+1) + (n+2)(n+3).\tag{$\|\;\small{ \rm stands\;for}\;concatenation$}\end{align}$$ Or $$\begin{align} (n+2)\|(n+3)-n\|(n+1)&=(n+2)(n+3)-n(n+1) \\ &=n^2 + 5n + 6 - n^2 - n \\ &= 4n+ 6 \\ &= 2(2n+3).\end{align}$$ Now this is when I noticed that if $n=4$ then $2(2n+3)=22$. I may be able to reach 22 like so: $$\begin{align}(n+2)\|(n+3)-n\|(n+1)&=(n+2-n)\|(n+3-n-1)\\ &=2\|2 \\ &= 22\end{align}$$ If that is true, then this happens: $$\begin{align}22 &= 2(2n+3)\\ \Leftrightarrow \; 11 &= 2n+3 \\ \Leftrightarrow \; 8 &= 2n \\ \\ \therefore \; n &= 4.\end{align}$$ This is my desired result, thus proving the conjecture. But is the equation in the second sandbox true? How does one subtract from concatenation? This, in particular, is new to me; I would mostly appreciate a full, explanatory answer. Thank you in advance.

论坛问题帖 | has_answers | Mr Pie | 2019-04-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018960 | MSE-3193902

For every b in the power a^{b}, does there exist an a such that the digit sum of this power is equal to a?

$1^0 = 1\to 1 =1$ $x^1=x\to x=x\;\forall x$. $9^2 = 81\to 8+1=9$ $8^3=512\to 5+1+2=8$. $7^4=2401\to 2+4+0+1=7$ $46^5 = 205962976\to 2+0+5+9+6+2+9+7+6=46$ $64^6 = 68719476736\to 6+8+7+1+9+4+7+6+7+3+6 = 64$ $68^7= 6722988818432\to 6+7+2+2+9+8+8+8+1+8+4+3+2 = 68$ $54^8 = 72301961339136\to 7+2+3+0+1+9+6+1+3+3+9+1+3+6=54$ $71^9 = 45848500718449031$ $\downarrow$ $4+5+8+4+8+5+0+0+7+1+8+4+4+9+0+3+1 =

71$ Conjecture: Given a positive integer b, there exists a positive integer a such that the digit sum of a^b is equal to a. Can this be proven? I don't know how to prove it; it was about 3:45am in the morning and I couldn't go to sleep, so I just went on my calculator and messed around because I was bored. That's when I noticed this cool property and decided to share it here. It is now 4:35am so... I gotta go to bed. I'll see you in some hours and hopefully gather the time to work on this. Sorry about that! Oh, incidentally, I also noticed that the digit sum of 29^5 is 23 and the digit sum of 23^5 is 29, so... there are cycles here. Same for 31 and 34. Also, the digit sum of 13^2 is 16 and the digit sum of 16^2 is 13. These cycles seem to only have two numbers involved, but I think regarding the seventh power, there is more than two involved (start with 72^7 I think), however there is also a cycle between two numbers of seventh powers (between 44 and 62). Does this help? I don't know. I have to go to bed. Good night! Thank you in advance.

论坛问题帖 | has_answers | Mr Pie | 2019-04-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018961 | MSE-3196077

Minimal order of a counterexample to Wall’s conjecture

There used to be once a rather well known and interesting conjecture, that was formulated by Gordon E. Wall: The number of maximal subgroups of a finite group G does not exceed $|G|$ That conjecture appeared to be false. However, the only information about it, that I was able to find, was this MO answer. It contains the statement, that the conjecture was disproven, and a link to the research in which the counterexamples were found. That link, however, seems to be already dead by the time I got there. So, my question is, what is the minimal possible order of a counterexample to Wall’s conjecture? Or, in case, if that is unknown yet, the minimal order of a known counterexample to Wall’s conjecture? I need this kind of information for my collection of disproven conjectures in combinatorics

论坛问题帖 | has_answers | Chain Markov | 2019-04-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018962 | MSE-3196785

Are there infinitely many solutions such that the digit sum of a prime power is a smaller power of the same prime?

While discussing prime powers and divisors, I came up with the following problem. Examples $\to$ prime $p=3$ digit sum (in base ten) of $p^3=27$ is $p^2=9$, a power of p,. $\to$ prime $p=7$ digit sum of $p^4=2401$ is $p^1=7$, a power of p. We can get even crazier: $\to$ prime $p=5$ digit sum of p^{208} is equal to $p^4=625$. and so on. The PARI/GP code used to generate the examples is below: sfun(p,k,n)={for(q=2,n,if(sumdigits(p^q,10)==p^k,print(q)))}. Questions Main question: Denote the digit sum (in base ten) of an integer m as $s(m)$. Are there infinitely many solutions such that $$s(p^n)=p^k$$ where p is a prime, and k,n are positive integers? When $p=3$, it has the nice property that $s(3^n)$ is always divisible by 3, so the first few solutions are not very hard to find. For example, when $k=3$, we obtain $n=9,10,11,13,16,17,21$. Some experimentation with the code reveals that the solutions get sparser as p increases, as expected. But the solutions themselves are very unexpected, such as $$s(5^{4938})=5^6,\quad s(89^{898})=89^2.$$ Thus I believe in the finitude of the solutions, but I think evaluating them will be out of the question. To disprove my claim, it suffices to show that for every p there is a solution, since we know that the number of primes is infinite.

论坛问题帖 | no_answers_at_retrieval | TheSimpliFire | 2019-04-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018963 | MSE-3197125

Are 2, 3 the only prime numbers that don't have the digit 1 and are palindromes whose squares are also palindromes?

While thinking about prime numbers, I noticed that: (1) Very few prime numbers have squares that are palindromes. Ex: 2, 3, 11, 101, 307 (2) Even rarer are prime numbers that are palindromes whose square are palindromes. Ex. 2, 3, 11 This inspired me to ask the following questions: (1) Are $2, 3$ the only prime numbers that don't have the digit 1 and are palindromes whose squares are also palindromes? (2) If not, then are there a finite number of these types of these prime numbers.

论坛问题帖 | no_answers_at_retrieval | Mathphile | 2019-04-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018964 | MSE-3198101

Pairs of palindromic primes without 1 and have a palindromic product

While discussing about prime numbers with other users, I noticed that: (1) There are very few pairs of palindromic prime numbers that do not contain the digit 1 and that have products which are palindromes. Ex : $[2, 3], [2, 30203], [2, 30403]$ (2) For the large range that was tested on PARI/GP, I noticed that all the palindromic primes that yielded palindromic products were composed only of the digits $0, 2, 3, 4$ (3) The palindromic primes of these pairs are equal to 2 or always exist in the range of 3×10^k to 4×10^k where $k \in \Bbb{+Z}, 0$ User Peter helped me get the following results for $a, b \lt 10^7$ on PARI/GP: [2, 3] [2, 30203] [2, 30403] [2, 32323] [2, 32423] [2, 3002003] [2, 3222223] [2, 3223223] [2, 3233323] [2, 3304033] [2, 3343433] [2, 3400043] [2, 3424243] [2, 3443443] [2, 3444443] [3, 30203] [3, 32323] [3, 3002003] [3, 3222223] [3, 3223223] [3, 3233323] [30203, 3002003] Questions: (1) Are there a finite number of pairs of a, b, where a, b are palindromic primes that do not contain the digit 1 and ab is a palindrome? (2) Are all the palindromic primes that yielded palindromic products composed only of the digits $0, 2, 3, 4$? (3) Are all the palindromic primes of these pairs equal to 2 or always exist in the range of 3×10^k to 4×10^k where $k \in \Bbb{+Z}, 0$?

论坛问题帖 | no_answers_at_retrieval | Mathphile | 2019-04-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018965 | MSE-3200433

New results on Identical Binomial Coefficient?

Are there any nontrivial identical binomial coefficients found other than: $$ {16 \choose 2}={10 \choose 3}=120 \\ {21 \choose 2}={10 \choose 4}=210 \\ {56 \choose 2}={22 \choose 3}=1540 \\ {120 \choose 2}={36 \choose 3}=7140 \\ {153 \choose 2}={19 \choose 5}=11628 \\ {221 \choose 2}={17 \choose 8}=24310 \\ {78 \choose 2}={15 \choose 5}={14 \choose 6}=3003$$ and the infinite family (where F_n is the nth fibonacci number): $$ {F_{2i+2}F_{2i+3} \choose F_{2i}F_{2i+3}}={F_{2i+2}F_{2i+3}-1 \choose F_{2i}F_{2i+3}+1}$$ By non-trivial I mean, of the shape ${n \choose k}$ where $2\leq k \leq \frac{1}{2}n $. This was conjectured to be the complete list in a paper from 1996: https://pdfs.semanticscholar.org/69fe/a6ef6f5ac5818538a86e98ddd9236a3310f7.pdf Are there any new results?

论坛问题帖 | no_answers_at_retrieval | Hyunbok Wi | 2019-04-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018966 | MSE-3203320

Proof that the error between $W(\lambda(\sqrt{n}))$ and the fraction of composite integers in $\mathbb{O}_n$ declines as n increases.

Please provide assistance for the proof that the error between the function $W(\lambda(\sqrt{n}))$ and the fraction of composite integers in $\mathbb{O}_n$ declines as n increases. Let $\mathbb{O}_n$ represent the set of odd integers less than or equal to n excluding 1. $\mathbb{O}_n$ = {3,5,7,9,11,13,15,17,19,...n} Let the function $l(x)$ equal the largest prime number less than x. For example: $l(10) = 7, l(20) = 19$ and $l(23) = 19$ Let the function $\lambda(x)$ equal the largest prime number less than or equal to x. For example: $\lambda(10) = 7, \lambda(20) = 19$ and $\lambda(23) = 23$ The function $W(\lambda(\sqrt{n}))$ was derived to represent the fraction of composite integers in the set $\mathbb{O}_n$ for large values of n. \begin{eqnarray*} W(\lambda(\sqrt{n})) = \sum_{\substack{p=3\\p\ \text{prime}}}^{\lambda(\sqrt{n})}\left((\frac{1}{p})\prod_{\substack{q=3\\q\ \text{prime}}}^{l(p)}\frac{(q-1)}{q}\right) \end{eqnarray*} or $W(\lambda(\sqrt{n}))= \frac{1}{3} + (\frac{1}{5})(\frac{2}{3}) + (\frac{1}{7})(\frac{2}{3})(\frac{4}{5}) + ... + (\frac{1}{ \lambda(\sqrt{n}))})(\frac{2}{3})(\frac{4}{5})(\frac{6}{7})(\frac{10}{11})...(\frac{(l(\lambda(\sqrt{n})) - 1)}{l(\lambda(\sqrt{n}))})$.\ I have proven that the maximum error between $W(\lambda(\sqrt{n}))$ and the actual fraction of composite integers in the set $\mathbb{O}_n$ declines as n increases in sections 6 and 7 in this paper http://vixra.org/pdf/1901.0436v5.pdf. Is this proof sufficient? If not, what needs to be done? Please only post helpful information.

论坛问题帖 | has_answers | temp watts | 2019-04-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018967 | MSE-3206134

Can the product of n positive integers, where $n \gt 5$ in A.P. be a palindrome?

Reading the question can the product of four positive integers in A.P. be a square?, also made me question whether the product of n positive integers, where $n \gt 5$ in arithmetic progression be a palindrome? Me and user Peter tried to find solutions for various n in PARI/GP, and found that there were no solutions for a large range of numbers we tested when $n \gt 5$. This led to the following questions: (1) Can the product of n positive integers, where $n \gt 5$ in A.P. be a palindrome? (2) If yes then for what value of n is there no solution?

论坛问题帖 | no_answers_at_retrieval | Mathphile | 2019-04-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018968 | MSE-3208273

Is the set of elementary functions which do not have elementary integrals bigger than set of elementary functions which have elementary integrals?

It increasingly seems to me that the functions that have elementary integrals are quite rare in comparison to the ones that don't have them. Even raising an elementary function to a different power may result in it not having an elementary integral . Ex. $\sqrt{\arctan (x)}$ Also many seemingly simple functions do not have elementary integrals. Ex. $\frac {\sin (x)}{x}$ or $ \sin \left(\frac{1}{x} \right) $ So my question is that can we write a formal proof to prove/disprove that the set of elementary functions which do not have elementary integrals is bigger than set of elementary functions which have elementary integrals?

论坛问题帖 | has_answers | Mathphile | 2019-04-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018969 | MSE-3210806

A conjectured identity for the fractional iterates of $\sqrt{x}+2$

Deeply impressed by the Fractional iterates of $x²-2$, a problem by Ramanujan post one month ago, I decided to investigate a little by myself on the topic and discovered that using Carleman matrices would be of great help for that purpose. After various numerical computations, and helped by several algorithms for recognizing patterns and guessing formulas, I finally came accross a differential equation detected in a Carleman matrix (by the guessHolo function in the FriCAS computer algebra system) which I could easily solve in order to write down the following identity: $$ F^{[n]}(x) = 2 + 2\cosh\left(2^{1-n}\,\textrm{arcsinh}\!\left(\frac {\sqrt{x-4}}{2}\right)\right) $$ with $F(x) = \sqrt{x} + 2$. Plotting this function for various values of n between 1 and 2 together with the functions F and $F^{[2]}$ shows how smoothly the interpolation works. Now, I don't know if this is something trivial or not and would be very happy to learn about it. Is such a result already known?

论坛问题帖 | has_answers | Thomas Baruchel | 2019-05-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018970 | MSE-3211298

Asymptotic frequency of $[0,\,1,\,1]$ in the Thue-Morse sequence

Let t_n be the Thue-Morse sequence: $$[\color{blue}{0,\,1,\,1,\,0,}\,\color{red}{1,\,0,\,0,\,1,}\,\color{blue}{1,\,0,\,0,\,1,}\,\color{red}{0,\,1,\,1,\,0,}\,...].\tag1$$ See this question for a definition and formula for t_n. Let's split the sequence into non-overlapping runs of length 3: $$[\underline{[\color{blue}{0,\,1,\,1}]},\,[\color{blue}{0,}\,\color{red}{1,\,0}],\,[\color{red}{0,\,1,}\,\color{blue}1],\,[\color{blue}{0,\,0,\,1}],\,\underline{[\color{red}{0,\,1,\,1}]},\,...].\tag2$$ There are 6 different sorts or runs in this sequence: all possible combinations of 0 and 1 except $[0,\,0,\,0]$ and $[1,\,1,\,1]$ - the Thue-Morse sequence is cube-free. What is the asymptotic frequency of the $\underline{[0,\,1,\,1]}$ run? Empirically, it seems to be around $\style{color:#bbbbbb;text-decoration:line-through}{1/5}\,1/6$, but the convergence is quite slow and erratic.

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2019-05-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018971 | MSE-3212925

Integrals of the Bessel function $J_0(x)$ over the intervals between its zeros

Let $J_0(x)$ be the Bessel function of the first kind. It has an infinite number of zeros on the positive real semi-axis. Let's denote them as $j_{0,n}$: $$j_{0,1}=2.40482...,\quad j_{0,2}=5.52007...,\quad j_{0,3}=8.65372...,\quad\small...\tag1$$ We are interested in absolute values of the integrals of $J_0(x)$ over the intervals between its consecutive zeros: $$\sigma_n=(-1)^n\int_{j_{0,n}}^{j_{0,n+1}}\!\!J_0(x)\,dx.\tag2$$ Their values are: $$\sigma_1=0.80145...,\quad\sigma_2=0.59932...,\quad\sigma_3=0.49904...,\quad\small...\tag3$$ We are interested in the asymptotic behavior of this sequence. Empirically, it seems that $$\sigma_n\,\stackrel{\color{#a0a0a0}?}\sim\,\frac{2\sqrt2}{\pi\sqrt n}\left(1-\frac1{8n}+O\!\left(\frac1{n^2}\right)\right)\!.\tag4$$ Can we prove this? Can we find next coefficients in this expansion?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2019-05-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018972 | MSE-3213216

A question on discrete Fourier transform of some function

Let $\sigma(n) = \sum_{d|n} d$ and $\tau(n) = $ number of divisors of n. For each $k, 0 \le k \le n-1$ we can look at the discrete Fourier transform of the numbers $\sigma(\gcd(n,k))$ given by: $$\hat{\sigma}(k) = \sum_{l=0}^{n-1} \sigma(\gcd(n,l))\exp\left(\frac{-2 \pi i k l }{n}\right)$$ I implemented this in SAGEMATH, and it occured me that : Conjecture: $$\hat{\sigma}(k) = n \tau(\gcd(n,k))$$ Is there any way to prove this? Here is some sage-code which implements this: def tau(n): return len(divisors(n)) def FTS(n,k): # Fourier Tranformierte von sigma(gcd(n,k)) return sum([sigma(gcd(n,l))*exp(-2*pi*I/n*k*l) for l in range(n)]) N = 8 for k in range(N): print k,abs(FTS(N,k).N()),N*tau(gcd(N,k)) which gives: 0 32.0000000000000 32 1 8.00000000000000 8 2 16.0000000000000 16 3 8.00000000000000 8 4 24.0000000000000 24 5 8.00000000000000 8 6 16.0000000000000 16 7 8.00000000000000 8 Edit: From the conjecture above it follows, that, by setting $k=0$: $$\tau(n) = \frac{1}{n}\sum_{l=0}^{n-1} \sigma(\gcd(n,l))$$ for which I have a proof. By taking the inverse Fourier transform, and setting again $k=0$ we get: $$\sigma(n) = \sum_{l=0}^{n-1} \tau(\gcd(n,l))$$

论坛问题帖 | has_answers | user276611 | 2019-05-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018973 | MSE-3214649

Is it still unknown whetever any $\mathscr{D}$-class D being a semigroup is bisimple?

Introduction: It is stated in my book from year 1961 that it is unknown whatever a subsemigroup D of semigroup S, which is also a $\mathscr{D}$-class of S is always bisimple, but it is known for $\mathscr{D}$-classes which are regular. Is it still unknown to this day? If it have been solved, can you provide sources? Definitions: Definition 1: We say that the elements x and y belonging to the semigroup S are in the same $\mathscr{L}$-class ($\mathscr{R}$-class) if and only if they generate the same left (right) principal ideal, that is, if $S^1x = S^1y$ ($xS^1 = yS^1$) holds. Definition 2: We say that elements x and y of S belong to the same $\mathscr{D}$-class if and only if $(x, y)\in\mathscr{L}\lor \mathscr{R}$ (the joint of relations $\mathscr{L}$ and $\mathscr{R}$). As an alternative definition, it can be shown that relations $\mathscr{L}$ and $\mathscr{R}$ commute, and hence $\mathscr{D} = \mathscr{L}\circ\mathscr{R} = \mathscr{R}\circ\mathscr{L} $, that is x and y belong to the same $\mathscr{D}$-class if and only if there exists z such that $x\mathscr{L}z\mathscr{R}y$. Definition 3: We say that the semigroup S is bisimple, if and only if there is a single $\mathscr{D}$-class in S.

论坛问题帖 | has_answers | Jakobian | 2019-05-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018974 | MSE-3216455

Circular Happy Palindromic Primes

(1) A circular prime is a prime number with the property that the number generated at each intermediate step when cyclically permuting its (base 10) digits will be prime. For example, 1193 is a circular prime, since 1931, 9311 and 3119 all are also prime. (2) A happy number is defined by the following process: Starting with any positive integer, replace the number by the sum of the squares of its digits in base-ten, and repeat the process until the number either equals 1 (where it will stay), or it loops endlessly in a cycle that does not include 1. Those numbers for which this process ends in 1 are happy numbers. For example, 19 is happy, as the associated sequence is $1^2 + 9^2 = 82,$ $8^2 + 2^2 = 68,$ $6^2 + 8^2 = 100,$ $1^2 + 0^2 + 0^2 = 1.$ (3) A palindromic prime is a prime that remains the same when its digits are reversed. For example, 11 A Circular happy palindromic prime is a prime number that has all three of the above properties. According to my testing in PARI/GP and my own results, I have found the following circular happy palindromic primes: $7, 1111111111111111111$ (19 digits)$, 1111111111111111111111$ (23 digits) (Haven't

tested complete range of 10^{24} in PARI/GP) Note that 19 and 23 are happy primes themselves which explains why $111111111111111111, 11111111111111111111111$ are also happy primes. Questions: (1) Are $7, 111111111111111111, 11111111111111111111111$ the only circular happy palindromic primes? (2) Are $1111111111111111111, 11111111111111111111111$ the only happy primes of the form 11111.... ? (3) If not then are there a finite number of circular happy palindromic primes? Can we prove/disprove this?

论坛问题帖 | no_answers_at_retrieval | Mathphile | 2019-05-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018975 | MSE-3216962

Create & Prove Conjecture - Discrete Math (Proofs)

I am stuck on the following problem: Imagine that a building has been overrun with snakes and rats. To help curb the problem, the building manager decides to offer employees brownie points for capturing and relocating them. They offer 13 points for each snake and 6 points for each rat captured and removed. Make a conjecture about what totals for points are possible. For example, an employee could not earn 7 points in this scenario. Prove that your conjecture is true. What "conjecture" am I supposed to be making? I assume I am trying to find a fact/pattern; something like "points can only be obtained in multiples of 6", but I am lost. I only managed to find that the two provided numbers, 6 and 13, are relatively prime. Of course, I also need to figure out how to prove such conjecture, I just can't do that until after I make it. Edit: Okay, I totally misread the example conjecture. So my conjecture will be "an employee could not earn 14 points in this scenario". I write the following proof: Proof: The two values of obtainable points are 6 and 13. The first 5 possible combinations of sums of 6 and 13 are, in order, as follows: { (0), (6), (6+6), (13), (13+6), (6+6+6) } = { 0, 6, 12, 13, 18 } There are no possible combinations of sums of 6 and 13 that equate to 14. Therefore, it is impossible for an employee to earn 14 points in this scenario.

论坛问题帖 | no_answers_at_retrieval | m450n | 2019-05-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018976 | MSE-3218966

Is 29 the only prime of the form p^p+2?

I searched for primes of the form p^p+2, where p is prime for a range of $p \le 10^5$ on PARI/GP and found that 29 is the only prime of this form in this range. Questions: (1) Is 29 the only prime of the form p^p+2, where p is prime? (2) If not, then are there a finite number of primes of the form p^p+2? Can you prove/disprove this? Edit: Since p^p grows really fast and primes get rarer and are spread farther out for large numbers, I conjecture that 29 is the only prime of the form p^p+2 where p is a prime.

论坛问题帖 | has_answers | Mathphile | 2019-05-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018977 | MSE-3219050

Are 3, 9 the only natural numbers n for which both 2^n-n and 2^n+n are prime?

I searched for natural numbers n, where 2^n-n and 2^n+n are both prime for the range of $n \le 10^5$ on PARI/GP and found that 3, 9 are the only solutions in this range. Note that since 2^n-n, 2^n, 2^n+n are in arithmetic progression and 2^n is not divisible by 3, n must be a multiple of 3. Questions: (1) Are 3, 9 the only natural numbers n for which both 2^n-n and 2^n+n are prime? (2) If not, then can you prove or disprove that there are finite natural numbers of this type?

论坛问题帖 | no_answers_at_retrieval | Mathphile | 2019-05-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018978 | MSE-3219130

Will p always be prime if $p^p+(p-1)!$ is prime?

While finding primes of the form $p^p+(p-1)!$ on PARI/GP, I noticed that p is always prime if $p^p+(p-1)! \gt 2$ is prime. The search range was $p \le 10^5$. Here are the solutions for $p\in\Bbb{+Z}$ for which $p^p+(p-1)!$ is prime that I got on PARI/GP: 1 2 3 11 43 Questions: (1) Will p always be prime if $p^p+(p-1)! \gt 2$ is prime? (2) Are there finite primes of the form $p^p+(p-1)!$, where $p\in\Bbb{+Z}$? How would you prove/disprove this? Edit: Just realized that the answer for the 1st question was obvious. But I think the second question will be much harder to answer.

论坛问题帖 | has_answers | Mathphile | 2019-05-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018979 | MSE-3221569

Conjecture: all complex roots of $\sum_{k=0}^\infty \frac{z^k}{\left(nk\right)!}$ are real

Conjecture: $$\left[n\in\mathbb{Z}^+,z\in\mathbb{C},0=\sum_{k=0}^\infty \frac{z^k}{\left(nk\right)!}\right]\Rightarrow z\in\mathbb{R}$$ This conjecture has been verified for $n\in\{1,2,4\}$. The motivation for this conjecture arose during the study of the exponential sum function which has applications to exponentiation in rings with abelian multiplication: $$\text{rues}_n\left(z\right)=\sum_{k=0}^\infty \frac{z^{nk}}{\left(nk\right)!}=\frac{1}{n}\sum _{k=1}^n \exp\left(ze^{2ki\pi/n}\right)$$

论坛问题帖 | has_answers | Nazgand | 2019-05-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018980 | MSE-3230707

Is there a maximum number of consecutive sexy prime pair sums that are all divisible by 10?

I was finding the sums of pairs of sexy primes (prime numbers that differ by 6) and noticed that there are a lot of pairs who's sum is divisible by 10. Ex. $(7, 13)$ as $7+13=20$ and 20 is divisible by 10. I then wondered if there are consecutive sexy prime pairs who's

sum is divisible by 10 and ran some PARI/GP code to find these kind of consecutive pairs. I found upto 17 consecutive pairs divisible by 10 on PARI and tested for a search limit of 10^{12}. Here is the output I got after running my code, the first column displays the smallest prime of the lowest pair of consecutive sexy primes and the second column displays the number of consecutive pairs divisible by 10. 7 1 167 2 2237 3 2267 4 108187 5 1004057 6 3281777 7 32895377 8 65947927 9 569959037 10 602817437 11 5476396897 12 16842019627 13 16842019637 14 17004549137 15 312318208577 16 382560132847 17 This leads to my question: If there are an infinite number of sexy primes, then is there a maximum number of consecutive sexy prime pair sums that are all divisible by 10?

论坛问题帖 | has_answers | Mathphile | 2019-05-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018981 | MSE-3231142

Infinitude of super happy primes

Similar to happy primes, I define super happy primes by the following process: (1) Find the sum of the digits raised to the power of themselves. Ex. 13 gives sum $ = 1^1 + 3^3 = 28$ (2) If the digit root of the sum is equal to 1 then it is a super happy number. Ex. digitroot(sum)$=$digitroot(28)$=1$. So 13 is a super happy prime. Here are some super happy primes I got by searching using PARI/GP with the help of user Peter. 13 19 31 61 83 89 139 157 163 193 199 313 331 383 389 571 587 613 619 631 661 68 3 691 751 839 857 863 919 983 991 1087 1117 1171 1187 1277 1399 1567 1579 1597 1 657 1663 1669 1693 1699 1747 1753 1759 1871 1933 1993 1999 2141 2281 2411 2447 Note: Since digitroot(sum) is nothing but $sum \mod 9$, the sum will be of the form $9k+1$, for some natural number k. Now coming to the question: Can we prove the infinitude of super happy primes?

论坛问题帖 | has_answers | Mathphile | 2019-05-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018982 | MSE-3232391

There cannot be more than three primes of the form $n^{n^2}-k$ for the same k?

I was searching for primes of the form $n^{n^2}-k$ on PARI/GP and noticed that primes of this form for same k are quite rare. The probability of finding a prime of this form is $\frac{1}{n^2 \log (n)}$, which makes me believe that the number of primes for same k of this form would be finite. I also noticed that the maximum number of primes of the form $n^{n^2}-k$ for the same k seems to be three. I searched for a range of $k \le 3 \times 10^5$ and $n \le 50$ on PARI/GP and didn't find more than three primes for same k. Here are the some values of k, for $k \le 10^4$ I found on PARI/GP, that have three primes of the form $n^{n^2}-k$. The first column displays the values of k and the second column shows the values of n for which $n^{n^2}-k$ is prime. 1409 [4, 6, 8] 3083 [4, 6, 8] 5496 [5, 7, 13] 7427 [4, 6, 8] 8902 [3, 9, 15] Questions: (1) Can there not be more than three primes of the form $n^{n^2}-k$ for the same k? (2) If there can be more than three, then is there a maximum number of primes of the form $n^{n^2}-k$ for the same k? Edit 1: Following the same process, I believe that the maximum number of primes of the form $n^{n^2}+k$, for same k is four. Edit 2: Peter showed provided an example with four primes for the same k which solves my question 1. So now I only look for answers of question 2.

论坛问题帖 | has_answers | Mathphile | 2019-05-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018983 | MSE-3237810

A really nice and elementary conjecture involving numbers

Yesterday, i discovered a nice thing while playing with numbers. It is trivial to note that $\forall n\in \mathbb{Z^+},\exists x,y\in \mathbb{Z}$ such that $3^n=5x^2+y$ has solutions. Also, undoubtedly, x,y are of different parity. But the good thing i noticed is really tough to prove i think. Conjecture If we choose to consider only the x,y such that $\mid x-y \mid$ is minimum, then if we make a table whose rows are the different values of this predefined x,y for consecutive values of n, like this where we make another row $y_{i-1}+y_i$(mainly for convenience) which will attain the sum of the current $y ~+$ the y of the previous entry. Now, interestingly, if we fix y_i, then i noticed upto $n=11$ using my calculator that $y_{i-1}+y_i\equiv y_{i+1}[5]$ (I purposely did not add the $n=11$ because i love even numbers :D But for $n=11$, $(x,y)=(188,427)$ and again $-518 \equiv 427 [5]$) I cannot understand the reason of its occurence. Can there be any elementary proof? Thanks for giving time! where the two ends of $/$ denotes congruency between the two $\mod 5$

论坛问题帖 | has_answers | user675504 | 2019-05-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018984 | MSE-3238379

Trigonometric integrals related to Gieseking's constant

This question at MathOverflow conjectures certain relation between fast converging hypergeometric series and Gieseking's constant. The Gieseking constant (G) is the volume of the hyperbolic 3-manifold, also called Gieseking's manifold. The hypergeometric series under consideration has trigonometric integral representations, which give two different forms of the conjecture: $$\int_0^{{\pi }/{3}} \frac{x \left({\sqrt{3}-{\sin x}}\right)\, dx}{\sin x \cdot \sqrt{3-2 \sqrt{3} \sin x}}\overset{?}{=}\frac{5}{2}\text{Im}\left(\text{Li}_2\left(e^{\frac{2 i \pi }{3}}\right)\right) = \frac{5}{3}G\tag{1}$$ $$ \int_0^{{\pi }/{3}} P(x)\, \frac{xdx}{\sin x \sqrt{\frac{3}{4}-\sin ^2x}}\overset{?}{=}5 \sqrt[4]{3}\, \rm{Im}\left(\rm{Li}_2\left(e^{\frac{2 i \pi }{3}}\right)\right) = \frac{10\sqrt[4]{3}}{3} G \tag{2} $$ where, $$P(x)=\left(\frac{\sqrt{3}}{2}-\sqrt{\frac{3}{4}-\sin ^2x}\right)^{3/2}+\left(\frac{\sqrt{3}}{2}+\sqrt{\frac{3}{4}-\sin ^2x}\right)^{3/2}$$ The 2nd representation is interesting because Lobachevsky has studied similar integrals in relation to volumes in hyperbolic geometry in his work "Application of imaginary geometry to certain integrals" (1836). Also see equation 3.842.7 in Gradshteyn and Ryzhik. For example he proves the following integration formula: $$\int_0^\beta \frac{x\sin x\, dx}{(1-\sin^2 \alpha \sin^2 x)\sqrt{\sin^2 \beta -\sin^2 x}} = \frac{\pi\ln\frac{\cos \alpha +\sqrt{1-\sin^2\alpha \sin^2\beta}}{2\cos \beta \cos^2 \frac{\alpha}{2}}}{2\cos \alpha \sqrt{1-\sin^2\alpha \sin^2 \beta}}$$ This formula in the limit $\alpha\to\pm i\infty $

becomes $$ \int_0^\beta\frac{xdx}{\sin{x}\sqrt{\sin^2{\beta}-\sin^2{x}}}=\frac{\pi}{4\sin{\beta}}\ln{\frac{1+\sin{\beta}}{1-\sin{\beta}}}.\tag{3} $$ One can see that when $\beta=\pi/3$, then both (2) and (3) have the same factor $\frac{x}{\sin{x}\sqrt{\sin^2{\beta}-\sin^2{x}}}$, and if it wasn't for the convoluted factor $P(x)$ these integrals would be the same. Lobachevsky also studied other integrals and in particular he shows that the integral with the variable upper limit $y<\beta$ $$ \int_0^y\frac{xdx}{\sin{x}\sqrt{\sin^2{\beta}-\sin^2{x}}} $$ can be expressed as a sum of elementary functions and Clausen functions. Maybe (2) is related to hyperbolic geometry and there is certain substitution that can transform this integral into a sum of Clausen functions or maybe it has some geometric interpretation? Q: How to prove conjectures (1) and (2)?

论坛问题帖 | has_answers | anon | 2019-05-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018985 | MSE-3241568

An asymptotic series for $\small\prod_{k=n}^\infty\operatorname{sinc}\left({2^{-k}}\pi\right),\,n\to\infty$

Using empirical methods, I conjectured that$^{[1]}$\!^{[2]}$ $$\small\prod_{k=n}^\infty \operatorname{sinc}\left({2^{-k}}\pi\right)=1-\frac{2\pi^2}9\,4^{-n}+\frac{38 \,\pi ^4}{2025}\,4^{-2n}-\frac{2332\,\pi ^6}{2679 075}\,4^{-3 n}+\frac{265618\,\pi^8 }{10247461875}\,4^{-4 n}+O\!\left(4^{-5 n}\right)\!.$$ Can we prove this? Can we find more coefficients in this expansion, or a general formula for those coefficients?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2019-05-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018986 | MSE-3252403

Does every simply-connected reductive group have trivial Galois cohomology?

Let G be a linear algebraic group over a field k with separable closure k^s and absolute Galois group $\Gamma\!_k$. Consider the Galois cohomology group $$ H^1(\Gamma\!_k,G(k^s)). $$ This group is known to be trivial in the following cases: k arbitrary, $G = SL_n$ or Sp_{2n} (Serre, Galois Cohomology, III.1). k nonarchimedean local, G semisimple and simply-connected (Bruhat-Tits). Based on these examples, we should expect the following conjecture to hold. Conjecture: Let G be a simply-connected reductive algebraic group. Then $$ H^1(\Gamma\!_k,G(k^s)) = 1. $$ Is this conjecture known to be true? Is it known to be true with additional assumptions on G or k?

论坛问题帖 | has_answers | user134824 | 2019-06-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018987 | MSE-3254331

Prove that $\int_0^\infty\left(\arctan \frac1x\right)^2 \mathrm d x = \pi\ln 2$

Prove $$\int_0^\infty\left(\arctan \frac1x\right)^2 \mathrm d x = \pi\ln 2$$ Out of boredom, I decided to play with some integrals and Inverse Symbolic Calculator and accidentally found this to my surprise $$\int_0^\infty\Big(\arctan \frac1x\Big)^2 \mathrm d x = \pi\ln 2 \quad (\text{conjectural}) \,\,\, {\tag{1}} $$ Here is Wolfram Alpha computation which shows (1) to be true to 50 digits. Is (1) true and how to prove it? I can calculate $$\int_0^\infty\arctan \frac{1}{x^2}\mathrm d x = \frac{\pi}{\sqrt2}$$ easily by expanding $\arctan$ into Maclaurin series. But how to proceed with $\arctan^2$?

论坛问题帖 | has_answers | Tyrell | 2019-06-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018988 | MSE-3257373

Is $\lfloor \zeta(-n) \rfloor$ only prime for $n=23$?

I searched for primes of the form of $\lfloor \zeta(-n) \rfloor$, where $n \in \Bbb{N}$, for a range of $n \le 10^4$ on PARI/GP and found $\lfloor \zeta(-n) \rfloor$ is only prime for $n=23$. My PARI code: for(n=1, 10^4, if(ispseudoprime(floor(-bernfrac(2*n)/(2*n)))==1, print([2*n-1, floor(-bernfrac(2*n)/(2*n))])); print(2*n-1)) Note that $\zeta(-n)$ for odd n (for even n, $\zeta(-n)=0$) can be also expressed as: $$\zeta(-(2n-1))=-\frac{B_{2n}}{2n}$$ where B_{2n} is the $2n$th Bernoulli number and written as brenfrac(2*n) in PARI/GP. Questions: (1) Is $\lfloor \zeta(-n) \rfloor$, where $n \in \Bbb{N}$, only prime for $n=23$? (2) Are there finite primes of the form of $\lfloor \zeta(-n) \rfloor$? I would appreciate any counterexamples(can be a probable prime)/proofs/papers. Extra: I also searched for primes of the form $\lceil\zeta(-n)\rceil$, $\lfloor\zeta(-n)\rceil$, $\lceil B_n \rceil$ and $\lfloor B_n \rceil$. For $n \le 10^4$: (1) $\lceil\zeta(-n)\rceil$ seems to be only prime for $n=691$. (2) There seems to be no prime of the form $\lfloor\zeta(-n)\rceil$. (3) $\lceil B_n \rceil$ seems to be only prime for $n=14$. (4) $\lfloor B_n \rceil$ seems to be only prime for $n=38$.

论坛问题帖 | no_answers_at_retrieval | Mathphile | 2019-06-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018989 | MSE-3271457

Conjectures on the primality of $\sum\limits_{i=1}^n p_i^{p_i}$ and $\sum\limits_{i=1}^n (-1)^ip_i^{p_i}$

Inspired by Is 29 the only prime of the form p^p+2?. Claims on prime powers and their alternating sums Consider the expressions $\mathcal P=\sum\limits_{i=1}^n p_i^{p_i}$ and $\mathcal Q=\sum\limits_{i=1}^n (-1)^ip_i^{p_i}$ over the prime numbers $(p_i)_{i\in\Bbb N}$. Then I claim that $n=2,4,24$ are the only occasions when $\mathcal P$ is prime. $n=2,4$ are the only occasions when $\mathcal Q$ is prime. PARI/GP code is for(n=1,+oo,if(isprime(sum(i=1,2*n,prime(i)^prime(i)))==1,print(n))). Note that we need only check even n as all prime powers are odd excluding $p_1=2$. Frankly I have no clue where to start. However, $\mathcal P$ and $\mathcal Q$ can be represented as $$\pm2^2+3^3+x\,\text{terms of the form}\,(6k-1)^{6k-1}+(n-x-2)\,\text{terms of the form}\,(6k+1)^{6k+1}.$$ Now working in modulo 6, $$\mathcal P\equiv 4+0+x\cdot(-1)^{-1}+(n-x-2)\cdot1^1\equiv2-2x+n\pmod6$$ so $\mathcal P$ is always composite when $n\equiv2(x-1)\pmod6$. Similarly, $$\mathcal Q\equiv

-4+0+x\cdot(-1)^{-1}+(n-x-2)\cdot1^1\equiv n-2x\pmod6$$ so $\mathcal Q$ is always composite when $n\equiv2x\pmod6$. Despite this, it is difficult to verify these congruences since x can be hard to determine when n is large. Nonetheless, I suspect a modular approach like the one above is too simple to solve the problem. I would appreciate any advances on this, though I would like further values of n through computation to be made in the comments and not in the answers.

论坛问题帖 | no_answers_at_retrieval | TheSimpliFire | 2019-06-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018990 | MSE-3274454

Uniform residue limits over $\{1,2,\dots,p-1\}$

Let p be a prime and $m \ge 1$. Define the (p residue of a residue) function $\tag 1 \gamma_{(p,m)}: [1, pm] \to \{0,\dots,p-1\}$ $\tag 2 \quad \quad \quad \quad \quad \quad \quad \quad \quad \quad \; n \mapsto [pm \pmod n] \pmod {p}$ Define $\quad u_{(p,m)} = pm - \text{#}\bigl(\gamma_{(p,m)}^{-1}(0)\bigr)$ For $1 \le r \le p -1$ define $\quad v_{(p,m,r)} = \frac{\text{#}\bigl(\gamma_{(p,m)}^{-1}(r)\bigr)}{u_{(p,m)}}$ Please prove that $\tag 3 \lim_{m \to \infty} v_{(p,m,r)} = \frac{1}{p-1}$ While the proof is missing (or (3) is refuted), allow me to make it a conjecture. My Work This came about from my investigations over here.

论坛问题帖 | no_answers_at_retrieval | CopyPasteIt | 2019-06-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018991 | MSE-3280127

Infinite $\frac{3}{2}$-generated groups?

Let’s call a group $\frac{3}{2}$-generated iff for every $g \in G \setminus \{1\}$ there is some $h \in G \setminus \langle g \rangle$ with $G = \langle g,h \rangle$. There is a conjecture by Breuer, Guralnick and Kantor, that a finite group is $\frac{3}{2}$-generated iff all its proper quotients are cyclic. Why is the word “finite” in the conjecture necessary? Is there an infinite counterexample? If yes, what it is? A $\frac{3}{2}$-generated group with a non-cyclic proper factor? Or a non-$\frac{3}{2}$-generated group, all proper cycles of which are cyclic?

论坛问题帖 | has_answers | Chain Markov | 2019-07-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018992 | MSE-3283519

Conjectured formula for the Fabius function

The Fabius function is the unique function ${\bf F}:\mathbb R\to[-1, 1]$ satisfying the following conditions: a functional-integral equation$\require{action} \require{enclose}{^{\texttip{\dagger}{a poet or philosopher could say "it knows and replays its own future"}}}$ $\displaystyle{\bf F}(x)={\small\int_0^{2{\tiny\text{ }}x}}{\bf F}\left(t\right)dt,\,$ and a normalization condition ${\bf F}(1)=1.$ It is noteworthy that despite being smooth (class C^∞)${^{\texttip{\dagger}{i.e. continuous and having continuous derivatives of any order}}}$ everywhere, ${\bf F}(x)$ is not real-analytic at any point $x\ge0$ - its Taylor series there is either divergent, or a polynomial with a finite number of terms.${^{\texttip{\dagger}{the latter happens at dyadic rational points}}}$ Unlike some other functions that appear specifically constructed for that purpose, the Fabius function naturally occurs in research of several seemingly unrelated problems - perhaps, this is why it has been discovered and re-discovered independently many times by several mathematicians. It also has some practical applications in computational mathematics. The Fabius function ${\bf F}(x)$ is constant 0 for all $x\le0,$ but some authors prefer to define it only on $[0,\infty)$ or only on $[0,1].$ Some authors study a very closely related Rvachev${^{\texttip{\dagger}{also variously spelled as Rvachëv or Rvachyov, a.k.a. "atomic function"}}}$ function $\operatorname{up}(x)$ defined as $\operatorname{up}(x) = {\bf F}(x+1)$ for $-1\le x\le1$ and $\operatorname{up}(x) = 0$ otherwise. It is known that the Fabius function assumes rational values at dyadic rational arguments - efficient algorithms to compute those values have been published or posted online, and have been discussed on this site.$^{[1]}$$\!^{[2]}$$\!^{[3]}$$\!^{[4]}$$\!^{[5]}$$\!^{[6]}$$\!^{[7]}$$\!^{[8]}$$\!^{[9]}$ I have been looking for a non-recursive, self-contained formula for the Fabius function for quite a long time. After lots of experimentation and looking for patterns in its values I came up with a conjectured empirical formula.
Let${^{\texttip{\dagger}{the superscript m is just an index here, not to be confused with a power}}}$ $$\mathscr F^m_n = \frac1{2^{n^2}\left(\frac12;{\tiny\text{ }}\frac12\right)_n}\,\sum _{k=0}^n\frac{\binom n k_{1/2}}{2^{{\tiny\text{ }}k{\tiny\text{ }}(k-1)}(n+k)!}\,\sum _{\ell=0}^{2^k{\tiny\text{ }}m-1}\,(-1)^{s_2\left(\ell\right)}\,\left(\ell-2^km+\tfrac12\right)^{n+k}{\small,}\tag{$\small\spadesuit$}$$ where $k,\,\ell,\,m,\,n$ are non-negative integers, $\displaystyle\small\left(a;{\tiny\text{ }}q\right)_n=\prod_{k=0}^{n-1} (1-a{\tiny\text{ }}q^k)$ is the q-Pochhammer symbol${^{\texttip{\dagger}{it assumes only rational values in this formula}}}$, ${\binom n k}_q=\displaystyle\small\frac{\left(q;{\tiny\text{ }}q\right)_n}{\left(q;{\tiny\text{ }}q\right)_k\left(q;{\tiny\text{ }}q\right)_{n-k}}\vphantom{\Huge|}$ is the q-binomial coefficient${^{\texttip{\dagger}{it assumes only rational values in this formula}}}$, and $s_2(n)\vphantom{\Huge|}$ is the sum of binary digits${^{\texttip{\dagger}{i.e. the number of 1's in the base-2 representation}}}$ of n (note that $(-1)^{s_2\left(n\right)} = t_n\vphantom{\Huge|}$ is just the signed version of the Thue-Morse sequence, satisfying recurrence $t_0 = 1,\,t_n = (-1)^n \, t_{\lfloor n/2\rfloor};\vphantom{\Huge|}$ see also${^{[10]}}$${\!^{[11]}}$${\!^{[12]}}$). I conjecture that for all non-negative integers $m,\,n,$ the following identity holds: $${\bf F}\!\left({\small\frac m{2^n}}\right)=\mathscr F^m_n.\tag{$\small\diamondsuit$}$$ Note that there is no requirement for $\frac m{2^n}$ to be a proper irreducible fraction - it might well be that m is even, or $m>2^n$. I have not been able to rigorously prove it yet, but it produces exact rational values that agree with those computed using known correct algorithms for all dyadic rational arguments I have tried. Of course, this formula did not just appear out of the blue - its structure was inspired by known algorithms, and it went a long way to this relatively concise form (at some point it had about 5 levels of nested summations/products). Can we prove that $\small(\diamondsuit)$ is indeed a correct representation of the Fabius function at dyadic rationals? If the conjecture is difficult to tackle at once, we can try to at least prove that the function defined by $\small(\spadesuit)$ shares some known properties with the Fabius function, e.g.: $0\le\mathscr F^m_n\le1,$ $\mathscr F^m_n = \mathscr F^{2{\tiny\text{ }}m}_{n+1}\vphantom{\Large|}$ - the result is the same across all

representations of a rational number $\frac m{2^n}$, $\mathscr F^m_n = 1-\mathscr F^{(2^n-m)}_n\vphantom{\Large|}$ for all $\small0\le m\le2^{n}$ - rotation symmetry, $\mathscr F^m_n = \mathscr F^{(2^{n+1}-m)}_n\vphantom{\Large|}$ for all $\small0\le m\le2^{n+1}$ - reflection symmetry, for a fixed $n,$ and $\small0\le m\le2^n$ the values of $\mathscr F^m_n\vphantom{\Large|}$ strictly increase, all points $\small\left(\frac m{2^n},\,\mathscr F^m_n\right)\vphantom{\Large|}$ lie on a continuous curve - for any convergent sequence of dyadic rationals $\frac m{2^n}\vphantom{\Large|}$, the sequence of corresponding values $\mathscr F^m_n\vphantom{\Large|}$ also converges. If the conjecture $\small(\diamondsuit)$ it true, then for all $0\le x\le1$ (not necessary rational) $${\bf F}\!\left(x\right)=\lim_{n\to\infty}\,\frac1{2^{n^2}\left(\frac12;{\tiny\text{ }}\frac12\right)_n}\,\sum _{k=0}^n\frac{\binom n k_{1/2}}{2^{{\tiny\text{ }}k{\tiny\text{ }}(k-1)}(n+k)!}\,\sum _{\ell=0}^{\left\lfloor2^{n+k}{\tiny\text{ }}x-1\right\rfloor}\,(-1)^{s_2\left(\ell\right)}\,\left(\ell-2^{n+k}{\tiny\text{ }}x+\tfrac12\right)^{n+k}.\tag{$\small\heartsuit$}$$ For dyadic rational x the limit is trivial, because the sequence under the limit becomes constant for large enough n. Also, there is a known series (by Rvachev) that expresses values of the Fabius function for all $0\le x\le1$ via its values at negative powers of 2: $${\bf F}\!\left(x\right)=\sum _{n=1}^\infty\frac{(-1)^{\left\lfloor 2^n{\tiny\text{ }}x\right\rfloor }-1}{2}\,(-1)^{s_2\left(\lfloor2^n{\tiny\text{ }}x\rfloor\right)}\;\sum_{k=0}^n\frac{2^{\frac{k{\tiny\text{ }}(k+1)}2}}{k!}\,\left(x-2^{-n}\left\lfloor 2^n{\tiny\text{ }}x\right\rfloor\right)^k\;{\bf F}\!\left(2^{{\tiny\text{ }}k-n}\right).\tag{$\small\clubsuit$}$$ Again, for dyadic rational x this series terminates after a finite number of terms, producing an exact rational value.

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2019-07-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018993 | MSE-3284709

If q is prime, can $\sigma(q^{k-1})$ and $\sigma(q^k)/2$ be both squares when $q \equiv 1 \pmod 4$ and $k \equiv 1 \pmod 4$?

This is related to this earlier MSE question. In particular, it appears that there is already a proof for the equivalence $$\sigma(q^{k-1}) \text{ is a square } \iff k = 1.$$ Let $\sigma(x)$ denote the sum of divisors of the positive integer x. Here is my question: If q is prime, can $\sigma(q^{k-1})$ and $\sigma(q^k)/2$ be both squares when $q \equiv 1 \pmod 4$ and $k \equiv 1 \pmod 4$? MY ATTEMPT Suppose that $$\sigma(q^{k-1}) = a^2$$ and $$\frac{\sigma(q^k)}{2} = b^2$$ for $q \equiv 1 \pmod 4$ and $k \equiv 1 \pmod 4$. Since $\sigma(q^k) = q^k + \sigma(q^{k-1})$, it follows that $$2b^2 = \sigma(q^k) = q^k + \sigma(q^{k-1}) = q^k + a^2.$$ Additionally, congruence-wise we obtain $$a^2 = \sigma(q^{k-1}) \equiv 1 + (k-1) \equiv k \equiv 1 \pmod 4,$$ from which it follows that a is odd, and $$2b^2 = \sigma(q^k) = q^k + \sigma(q^{k-1}) \equiv 1^1 + 1 \equiv 2 \pmod 4,$$ which implies that b is likewise odd. Now, using the definition of $\sigma(q^k)$ and $\sigma(q^{k-1})$ for q prime, we derive $$\frac{1}{2}\cdot\frac{q^{k+1} - 1}{q - 1} = b^2$$ and $$\frac{q^k - 1}{q - 1} = a^2.$$ Assume to the contrary that $$\frac{1}{2}\cdot\frac{q^{k+1} - 1}{q - 1} = b^2 \leq a^2 = \frac{q^k - 1}{q - 1}.$$ This assumption leads to $$q^{k+1} - 1 \leq 2(q^k - 1)$$ which implies that $$16 = {5^1}(5-2) + 1 \leq q^k(q - 2) + 1 = q^{k+1} - 2q^k + 1 \leq 0,$$ since q is a prime satisfying $q \equiv k \equiv 1 \pmod 4$. This results in the contradiction $16 \leq 0$. Consequently, we conclude that $a < b$. Furthermore, I know that $$(q+1) = \sigma(q) \mid \sigma(q^k) = 2b^2$$ so that $$\frac{q+1}{2} \leq b^2.$$ Finally, I also have $$\frac{q^{k+1} - 1}{2b^2} = q - 1 = \frac{q^k - 1}{a^2}.$$ Alas, here is where I get stuck. CONJECTURE (Open) If q is a prime satisfying $q \equiv k \equiv 1 \pmod 4$, then $\sigma(q^{k-1})$ and $\sigma(q^k)/2$ are both squares when $k = 1$. SUMMARY OF RESULTS SO FAR zongxiangyi appears to have proven the implication $$\sigma(q^k)/2 \text{ is a square} \implies k = 1.$$ The proof of the following implication is trivial $$k = 1 \implies \sigma(q^{k-1}) \text{ is a square}.$$ The truth value of the following implication is currently unknown: $$\sigma(q^{k-1}) \text{ is a square} \implies k = 1.$$ Together, the two results give $$\sigma(q^k)/2 \text{ is a square} \implies k = 1 \iff \sigma(q^{k-1}) \text{ is a square},$$ so that $\sigma(q^{k-1})$ is a square if $\sigma(q^k)/2$ is a square. Therefore, $\sigma(q^{k-1})$ and $\sigma(q^k)/2$ are both squares (given $q \equiv 1 \pmod 4$ and $k \equiv 1 \pmod 4$) when $\sigma(q^k)/2$ is a square.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2019-07-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018994 | MSE-3285468

Uniqueness of non-completable latin square of size n

It is known that partial latin squares of order n and size $n-1$ can always be completed to a latin square. I want to know if non-completable partial latin squares of order n, size n satisfy this property: There exists $n-1$ entries with the same row, column, or color. Is this a known result or maybe a false statement? If not, would it be decent for a research topic?

论坛问题帖 | no_answers_at_retrieval | Hyunbok Wi | 2019-07-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018995 | MSE-3285692

An arithmetic pattern

Define the odd part of a positive integer $\operatorname{od}(2^i(2j+1))=2j+1$ and define the function $f(n)=\operatorname{od}(n!-2^n)$. For $3\leq n \leq 10$ it holds that: $$f(2^n)-f(2^n-1)=2^n-2 \tag{1}$$ Does this equality holds for all $n\in\mathbb N_{>2}$? I don't like to try to prove random conjectures because I'm to bad at it, but I do like finding arithmetical patterns and test them with a computer. In the table below n and $f(n)$ are listed and the pattern is visible. Of course, there could be a rather easy explanation, only that I haven't found it. 4 1 5 11 6 41 7 307 8 313 9 2831 10 14171 11 155917 12 467771 13 6081067 14 42567517 15 638512859 16 638512873 17 10854718871 18 97692469871 19 1856156927617 20 9280784638121 21 194896477400617 22 2143861251406867 23 49308808782358109 24 147926426347074371 25 3698160658676859367 26 48076088562799171867 27 129805439119557764060 9 28 908638073836904348 4367 29 263505041412702261046859 30 395257562119053391570 3109 31 122529844256906551386796843 32 1225298442569065513867968 73 33 4043484860477916195764296871 34 687392426281245753279930 46871 35 240587349198436013647975 6640617 36 216528614278592412283178 09765621 37 801155872830791925447758961328117 38 152219615837850465835074202652 34367 39

593656501767616816756789390344140609 40
296828250883808408378394695172070312 1 41
121699582862361447435141825020548828117 42
255569124010959039613797832543152539061 7 43
109894723324712387033933067993555591796859 44
120884195657183625737326374792911150976561 7 45
543978880457326315817968686568100179394531 09 46
125115142505185052638132797910663041260742185 9 47
588041169774369747399224150180116293925488280 93 48
176412350932310924219767245054034888177646484 371 49
864420519568323528676859500764770952070467773 4367 50
216105129892080882169214875191192738017616943 359367 51
110213616244961249906299586347508296388984641 11328109 52
143277701118449624878189462251760785305680033 447265617 53
759371815927783011854404149934332162120104177 2705078109 54
205030390300501413200689120482269683772428127 863037109359 55
112766714665275777260379016265248326074835470 32467041015593 56
789367002656930440822653113856738282523848292 27269287109367 57
449939191514450351268912274898340821038593526 595434936523435 9 58
130482365539190601867984559720518838101192122 7126761315917968 59 59
769845956681224551021108902351061144797033524 0047891763916015593 60
115476893502183682653166335352659171719555028 6007183764587402343 59 61
704409050363320464184314645651220947489285674 4643820963983154296843 62
218366805612629343897137540151878493721678559 0839584498834777832030 93 63
137571087535956486655196650295683451044657492 2228938234265910034179681 1 64
137571087535956486655196650295683451044657492 2228938234265910034179687 3 65
894212068983717163258778226921942431790273699 4488098522728415222167968 71 66 295089982764626663875396814884241002490790320818107251250037702331542968 71 67 197710288452299864796515865972441471668829514948131858337525260562 1337890617 68 336107490368909770154076972153150501837010175411824159173792 94295562744140621 69 231914168354547741406313110785673846267537021034158669 829917130639382934570 3117

论坛问题帖 | has_answers | Lehs | 2019-07-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018996 | MSE-3286093

Conjecture: No positive integer can be written as a^b+b^a in more than one way

Today, I came up with the following problem when trying to solve this. Are there distinct integers $a,b,m,n>1$ such that the equation $$a^b+b^a=m^n+n^m$$ holds? That is, is there ever an integer that can be written as a^b+b^a in more than one way? I claim that the answer is No, but I think solving this is beyond my knowledge. For a very preliminary observation, the simplest case is to consider the powers of $1,5,6,0$, since they end in those same digits. For example, $$\begin{cases}a\equiv5\pmod{10}\\b\equiv6\pmod{10}\end{cases}\implies a^b+b^a\equiv1\pmod{10}.$$ However, this brings about an issue, since there is hardly any indication as to what values x and y can take other than them having opposite parity. PARI/GP code is intfun(a,b,m,n)={for(i=2,a,for(j=2,b,for(k=2,m,for(l=2,n,if(i<>k && i<>l && j<>k && j<>l && i^j+j^i-k^l-l^k==0,print(i," ",j," ",k," ",l))))));} No solutions have been found up to $a,b,m,n\le100$.

论坛问题帖 | no_answers_at_retrieval | TheSimpliFire | 2019-07-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018997 | MSE-3289757

Searching for a conjecture that is true until the 127 power of n.

I am searching for a mathematical conjecture that is true until something like n = 117, or n = 127, or a number close. It is an equality, a formula implying calcultations to the power of n, IIRC. What is the name of this conjecture ? Why do I search for this formula ?: I had read an article a few years ago that was discussing about experiments around the validation of a theory in Science, namely in Physics. The example of this conjecture was given so as to convince that you can choose to repeat an experiment with different parameters a certain number of time, ultimately in the end, it does not give the entire certainty a physical theory is true: it just happens we still haven't found any outlier measurement that contracdicts the physical theory. As a matter of fact, we can test this mathematical conjecture for the first 127 integers, which is a bunch of times, and believe it is always true, whereas it is false for all n. I would like the name of it so as to be able to name it / source it in discussion. Does it ring any bell for you ?

论坛问题帖 | has_answers | Stephane Rolland | 2019-07-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-018998 | MSE-3290798

A conjectured value for $\operatorname{Re} \operatorname{Li}_4 (1 + i)$

In evaluating the integral given here it would seem that: $$\operatorname{Re} \operatorname{Li}_4 (1 + i) \stackrel{?}{=} -\frac{5}{16} \operatorname{Li}_4 \left (\frac{1}{2} \right) + \frac{97}{9216} \pi^4 + \frac{\pi^2}{48} \ln^2 2 - \frac{5}{384} \ln^4 2$$ I arrived at a result involving the $\operatorname{Re} \operatorname{Li}_4 (1 + i)$ term for the value of the integral while the OP is convinced the integral in question has a simple, elementary answer. If both of us are right, then the conjecture holds. So my question is, is it possible to either (i) prove the conjecture true analytically or (ii) disprove the conjecture based on (very high precision) numerical evidence?

论坛问题帖 | has_answers | omegadot | 2019-07-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-018999 | MSE-3293832

Does there exist some prime k for which there will be exactly two primes of the form $n!+k$?

This is a question related to my recent question Conjecture: “For every prime k there will be at least one prime of the form $n!\pm k$” true? Using PARI/GP I searched for the number of primes of the form $n!+k$ possible for each prime $k\le 2300$. I observed that for some prime k, all numbers less than 24 were found except 2. Note that when $n \ge k$, $n! \pm k$ cannot be prime as k will be a factor of $n! \pm k$. This means that there are a finite number of primes of the form $n! \pm k$ for each k. User Peter and I, searched for some prime k for which $n!+k$ has two primes upto $k\le 10^5$ and found none in the range. Here is the code we used in PARI/GP if anyone wants to extend the search: forprime(k=2,10^7,s=0;for(n=1,k,if(s<3,if(ispseudoprime(n!+k)==1,s=s+1)));if(s==2,print(k))) Conjecture: For any prime k there cannot be exactly two primes of the form $n!+k$. Can prove/disprove this? Any ideas, heuristics, guesses are welcome. Update: Upto $k \le 10^6$, $k=714377$ is the only candidate remaining for which we have only found one prime of the form $n!+k$ till $n\lt 3000$. If a second prime of the form $n!+714377$ is found for the range $3000 \lt n \lt 714377$, then the conjecture is disproved!

论坛问题帖 | no_answers_at_retrieval | Mathphile | 2019-07-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019000 | MSE-3294459

On Composite Numbers of the Form $p_{1}p_{2} \ldots p_{k} - 1$

This question is related to D. H. Lehmer's 1932 conjecture on Euler's totient function: Are there any composite n for which $\phi(n)$ divides $n-1$? See, for example: On Lehmer's Totient Conjecture I would like to ask what is known regarding the factors of $p_{1}p_{2} \ldots p_{k} - 1$, where the p_{i} are distinct odd prime numbers? I was not able to find much in some number theory books I consulted. Thank you.

论坛问题帖 | has_answers | DDS | 2019-07-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019001 | MSE-3295143

What methods show that a number is transcendental?

I've been doing a lot of research on such theories lately and these are all I've found so far: Liouvilles criterion (here) Lindemann-Weierstrass theorem (here) Gelfond-Schneider theorem (here) Brownawell-Waldschmidt theorem (here or here) Schanuels conjecture (no proof yet) (here) Six and five exponentials theorem (here) Bakers theorem (here) Roths theorem (here) Or theorems such as: If you take two transcendental numbers, a and b, then at least one of ab and $a+b$ is transcendental (here) or x^{x^x} is transcencental if $x\in\mathbb{Q}, x\notin\mathbb{N},x>0$ (here) What other methods show that a number is transcendental besides these ones? If you know any other proven or unproven theorems it would be nice to share them. Edit 1 K.Dilcher and K.B. Stolarsky (here) Carl Ludwig Siegel (here) Edit 2 The Rogers-Ramanujan continued fraction (here) is transcendental for all algebraic $-1<q<1$ All the transcendental functions (here) Edit 3 All about trancendental Infinite Products (here) (here) and (here)

论坛问题帖 | has_answers | Kinheadpump | 2019-07-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019002 | MSE-3297995

Recurrence relation for the Thue-Morse sequence

I made a curious observation. Let a_n be the sequence of numbers determined by a recurrence relation $$\begin{cases} \vphantom{\large|}a_0=0\\ \vphantom{\large|}a_1=1\\ \vphantom{\Large|}n\,a_n=(5-2 n)\,a_{n-1}+3{\tiny\text{ }}(n-1)\,a_{n-2}+1 \end{cases}\tag{$\small\spadesuit}$$It also can be generated using an equivalent homogeneous recurrence relation $$\begin{cases} \vphantom{\large|}a_0=0\\ \vphantom{\large|}a_1=a_2=1\\ \vphantom{\Large|}n\,a_n = (4-n)\,a_{n-1}+(n-2)\left(5{\tiny\text{ }}a_{n-2} - 3{\tiny\text{ }}a_{n-3}\right) \end{cases}\tag{$\small\clubsuit}$$ 0, 1, 1, 2, 1, 4, -2, 13, -23, 68, -164, 439, -1146, 3067, -8231, 22306, -60791, 166684, ... It appears that this sequence modulo 2 gives the Thue-Morse sequence, meaning that if we denote $t_n=(-1)^{a_n}\,$ then it satisfies $$t_0 = 1,\quad t_n = (-1)^n \, t_{\lfloor n/2\rfloor}.\tag{$\small\diamondsuit}$$ How can we prove this?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2019-07-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019003 | MSE-3302459

Prime number function inequality conjecture

In my intuitive random search for conjectures I found $$n\neq 4\implies p_{n^2}\leq p_n^2<p_{4n^2}$$ It's tested for $n<1000$. I've looked at it the point of view of PNT but haven't the skills to prove or disprove it. Could there be counter examples out of reach of my conjecture seeking routines?

论坛问题帖 | has_answers | Lehs | 2019-07-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019004 | MSE-3303015

Any positive integer can be written as sum / difference of consecutive squares

How should one go about proving that $x \in \mathbb{N}$ can be written (with the right combination of signs) as $\pm 1^2 \pm 2^2 \pm \ldots \pm n^2$ for any $x : x, n \in \mathbb N$? I have tried for hours but can't figure out an appropriate approach. So what I ask of you specifically is, how should one be able to determine a reasonable approach for such a conjecture? Thanks.

论坛问题帖 | has_answers | Hassebae | 2019-07-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019005 | MSE-3304587

Conjecture: ${_2 F_1} \left(\frac{1}{2}- \frac{x}{2},1- \frac{x}{2};2;1 \right)=\prod_{n=2}^\infty \frac{(2n-2+x)(2n-1+x)}{2n (2n-3+2x)}$

Is the following conjecture for $0<x<1$ true and how do we prove it? $${_2 F_1} \left(\frac{1}{2}- \frac{x}{2},1- \frac{x}{2};2;1 \right)=\prod_{n=2}^\infty \frac{(2n-2+x)(2n-1+x)}{2n (2n-3+2x)}$$ I encountered this function when answering a question a while back, where I have shown that: $$S=\sum_{k=1}^\infty \frac{\Gamma \left(k+ \frac{1}{2} \right)}{\Gamma \left(k+1 \right)} G_{2k-1}=\frac{\sqrt{\pi}}{2} \int_0^1 x ~{_2 F_1} \left(\frac{1}{2}- \frac{x}{2},1- \frac{x}{2};2;1 \right) dx$$ Where G_k are so called Gregory coefficients. Now I returned to this problem and considered the ratio: $$\frac{{_2 F_1} \left(\frac{1}{2}- \frac{x}{2},1- \frac{x}{2};n;1 \right)}{{_2 F_1} \left(\frac{1}{2}- \frac{x}{2},1- \frac{x}{2};n+1;1 \right)}$$ For several cases for $n \geq 2$ Wolfram Alpha shown me the terms from the product above. This, together with an obvious identity: $$\lim_{n \to \infty} {_2 F_1} \left(\frac{1}{2}- \frac{x}{2},1- \frac{x}{2};n;1 \right)=1$$ Makes me believe that the infinite product converges. However, I don't know how to prove the ratio. Gauss's contiguous relations connect the two functions and the derivative, but I'm not sure how to use them in this case. The product is not telescoping (which is obvious, otherwise this hypergeometric function would be a rational function). Another question: are there more infinite product identities like this one for hypergeometric functions? We can rewrite the product in a more convenient way: $$\prod_{n=2}^\infty \frac{\left(1-\frac{2-x}{2n}\right)\left(1-\frac{1-x}{2n}\right)}{\left(1-\frac{3-2x}{2n}\right)}$$ Which looks a lot like a more complicated Gamma function. Claude Leibovici proposed a closed form for the product which means that the following holds: $$\sum_{k=1}^\infty \frac{\Gamma \left(k+ \frac{1}{2} \right)}{\Gamma \left(k+1 \right)} G_{2k-1}=\int_0^1 2^x \frac{x}{1+x} \frac{\Gamma \left(x+ \frac{1}{2} \right)}{\Gamma \left(x+1 \right)}~ dx$$ The series converges extremely slow, for example: $$\sum_{k=1}^{500} \frac{\Gamma \left(k+ \frac{1}{2} \right)}{\Gamma \left(k+1 \right)} G_{2k-1}=0.506676104408579236 96$$ While the integral gives: $$\int_0^1 2^x \frac{x}{1+x} \frac{\Gamma \left(x+ \frac{1}{2} \right)}{\Gamma \left(x+1 \right)}~ dx=0.507181289066845641 10$$ Which is the correct numerical value. Unfortunately, there's no in-built Gregory coefficient list in Mathematica, so it's hard to compute them for large k.

论坛问题帖 | has_answers | Yuriy S | 2019-07-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019006 | MSE-3305985

An iteration formula I found (please don't jump at me if it's already been discovered)

$\sin^2 \theta$"> Where the modulus-like symbol actually denotes iteration of a radical function. Sorry for the messy work everyone- I am new to this stuff, and I literally just found this iteration formula like 15 minutes ago. $X[i]$ starts as being an arbitrary number greater than $n!/(n-2)!$ which I should have mentioned earlier but didn't have space to (apologies). So the business is: you start with a number e.g. 1, then if you want to find say 50, you compute $sqrt(1 + 50!/48!)$ then replace 1 on your calculator with the result you just got a.k.a. Ans etc. and keep iterating till you get (at a surprisingly fast convergence rate!) the very number you started with, n. I appreciate this might not be the most interesting insight or $n!/(n-2)!$ is probably just me not realising that it could also be presented as $n(n-1) = n^2-n$. But still, please don't mass-downvote it all, remember it's a 15-year old person who figured this out...

论坛问题帖 | has_answers | Avtarás Karîm Elyméşer | 2019-07-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019007 | MSE-3319421

Conjecture $\sum_{n=0}^\infty a_n= \frac{1}{2}-\frac{7 \zeta(3)}{2 \pi^2}$

Working with some integrals I stumbled upon the following slowly converging series: $$ S = \sum_{n = 0}^{\infty}\left(-1\right)^{n} \left[n + \frac{3}{2} + \left(n + 1\right)\left(n + 2\right) \log\left(1 - \frac{1}{n + 2}\right)\right] $$ I have reasons to suspect that the series has a closed form: $$S=\frac{1}{2}-\frac{7 \zeta(3)}{2 \pi^2}=0.073721601182494209 \ldots$$ The actual proof eludes me so far. Can you prove or disprove this conjecture? Writing the logarithm as a series we have: \begin{align} &\left(n + 2\right) \log\left(1 - \frac{1}{n+2}\right) = -\sum_{k = 1}^{\infty} \frac{1}{k\left(n + 2\right)^{k - 1}} \\ = &\ -1-\sum_{k=1}^\infty \frac{1}{(k+1) (n+2)^k} \end{align} Which turns the series into: $$S=\sum_{n=0}^\infty (-1)^n \left(\frac{1}{2}-(n+1) \sum_{k=1}^\infty \frac{1}{(k+1) (n+2)^k} \right)$$ I can provide the way I came to this expression, but it's very long and complicated, as usual. I'd like some clear proof, if possible. To get the sense of how slowly the series converges, for 20000 terms the result agrees with the stated closed form in 4 first significant digits. The integral from which this series was obtained is (again, conjectured): $$\int_0^1 {_2 F_1} (1,-t;2-t;-1) dt = \frac{7 \zeta(3)}{\pi^2}+\frac{1}{2}$$ I don't think it's very useful, except for numerical confirmation.

论坛问题帖 | has_answers | Yuriy S | 2019-08-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019008 | MSE-3331664

Conjecture on Prime Numbers and Exponential Equipment

How to generate a rigorous proof for - The specific conjecture which states that we can always frame every Prime number in an exponentially framed equation with perfect cubes & squares. I'll present the respective segments of the conjecture and its Prototype. Conjecture's statement - The square of every Prime number can be expressed as a sum or difference of some perfect cube and any another Prime Mathematical Structure $a^3 \pm P_{k} = P^2_n$, $a,n, k\in\mathbb{N}$, Where P_m denotes the mth Prime respectively . Progress on Initial inputs and outputs- $\begin{array}{||c || c ||} \hline a^3 \pm p_k& p_n^2 \\ \hline 1+3& 2^2\\ \hline 512-503& 3^2\\ \hline 8+17& 5^2\\ \hline 8+41& 7^2\\ \hline 8+113& 11^2\\ \hline 216-47& 13^2\\ \hline 8+281& 17^2\\ \hline 8+353& 19^2\\ \hline 8+521& 23^2\\ \hline 1728-887& 29^2\\ \hline 8+953& 31^2\\ \hline 8+1361& 37^2\\ \hline 1728-47& 41^2\\ \hline 8000-6151& 43^2\\ \hline 216+1993& 47^2\\ \hline 8+2801& 53^2\\ \hline 512+2969& 59^2\\ \hline 512+3209& 61^2\\ \hline 8+4481& 67^2\\ \hline 1728+3313& 71^2\\ \hline 216+5113& 73^2\\ \hline 1728+4513& 79^2\\ \hline 216+6673& 83^2\\ \hline 8000-79& 89^2\\ \hline ...\pm...& 97^2\\ \hline ...\pm...& 101^2\\ \hline ...\pm...& 103^2\\ \hline \end{array}$ Conjecture's Prototype The sum or difference of

square of every Prime Number with some Perfect cube yields any another Prime number. Mathematical Structure $(m^3 \pm P^2_{b} = P_c)$, $m,b,c \in\mathbb{N}$ and P_n is the nth Prime Number respectively. The above parent equation can be framed in other 3 ways aslike - $P_c - m^3 = \pm P^2_b$ which clearly generates the expression for square of every Prime number as per the conjecture constraints. $|P_c - m^3| = |P^2_b|$ where taking modulus yields positive results, $P_c \mp P^2_b =m^3$ Initial inputs and outputs- $\begin{array}{||c || c ||} \hline P_c - m^3 & \pm P_b^2 \\ \hline\hline 31-27& 2^2\\ \hline 17-8& 3^2\\ \hline 89-64& 5^2\\ \hline 113-64& 7^2\\ \hline 337-216& 11^2\\ \hline 233-64& 13^2\\ \hline 353-64& 17^2\\ \hline 577-216& 19^2\\ \hline 593-64& 23^2\\ \hline 887-1728& -29^2\\ \hline 2689-1728& 31^2\\ \hline 1433-64& 37^2\\ \hline 1063-2744& -41^2\\ \hline 1913-64& 43^2\\ \hline 2273-64& 47^2\\ \hline 3023-5832& -53^2\\ \hline 3697-216& 59^2\\ \hline 5449-1728& 61^2\\ \hline 6217-1728& 67^2\\ \hline 8783-13824& -71^2\\ \hline 5393-64& 73^2\\ \hline \end{array}$ The parameter a in Primary conjecture and the parameter m in its prototype takes infinitely many but unique values . In order to prove it , one needs to overcome a new level of challenges it brings in Number theory , So far many efforts have been made to prove the Conjecture but we couldn't prove it for every value of a & m respectively. Aops community on internet has made efforts on it here Providing hypothesis... Proving a special case ... Hope we get to see a well defined constructive proof which even generalizes the Special case within itself.

论坛问题帖 | no_answers_at_retrieval | Alphatrion | 2019-08-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019009 | MSE-3334566

Determinant Optimization Conjecture

Let the maximum value of $n \times n$ determinant with $n \in\mathbb{N} >1$ be D_n . If each of the elements in determinant is either $+1$ or -1 , Then the successive values of D_n makes a Geometric Progression . D_2 $\begin{vmatrix} 1&-1 \\ 1&1 \\ \end{vmatrix}=2$ D_3 $\begin{vmatrix} 1&-1&\pm1 \\ 1&1&-1 \\ 1&1&1 \end{vmatrix}=4$ D_4 $\begin{vmatrix} 1&-1&1&-1 \\ 1&-1&-1&\pm1 \\ 1&1&1&-1\\ 1&1&1&1\\ \end{vmatrix}=8$ D_5 $\begin{vmatrix} 1&1&-1&-1&1 \\ -1&1&-1&1&1 \\ 1&1&1&1&1\\ -1&1&1&1&-1\\ 1&-1&1&1&-1\\ \end{vmatrix}=16$ D_6 $\begin{vmatrix} 1&-1&1&1&1&1 \\ 1&1&1&-1&-1&-1 \\ 1&1&1&-1&1&1\\ -1&-1&-1&1&1&-1\\ 1&1&1&1&-1&-1\\ -1&-1&1&1&1&1\\ \end{vmatrix}=32$ And so on... CRUX - Optimization- The determinant being optimized to maximum value always yields a number which follows up a geometric progression , Why isn't it even a bit higher or lower than it , to remove our this specific qualm , it's always saturated to a G.P. and strictly follows the same algorithm . Next thing is the Result of the Optimized Determinant - D_n $=$ Always a G.P. , how does it strictly follow up being a G.P. , suppose that we frame it algebraically , then will it show the same behavior ?! It's very necessary for algebraic frameworks . Configuration- Where to plot the elements so that in order to optimize it ?! The order increases ... So does the difficulty. Does it holds true for even more than 2 elements ?! And will it also yield G.P. as output ? Can we also have negative numbers and later non-integral Reals & which also follows the same G.P. tradition ?! Configuration also includes the order of configuration of the determinant itself , so can we have $n\times n$ type determinant with at most 2 distinct elements which follows the same above argument ?! Till now we saw that as the order of the determinant increases , the difficulty in optimization & it's respective element's configuration also increases ... Generalization- Generalizing it & framing it to a well finished conjecture ! Yes , but there always exists the Qualm about optimization & configuration issues as per mentioned above ... I did tried distinct elements - 1 & 2 this time , and surprisingly it came out to generate Geometric progressive numbers as output ! Holy Jesus ! Before generalizing one should keep all the above issues in mind & Is it a NP-HARD /COMPLETE problem which I end up with? ... Proof Positive- The ultimate explanation for this Enigmatic behavior of such determinant's optimization and it also includes the strict algorithm for the configuration of elements and generation of the final & the absolute proof dealing with all the above arguments. My dear colleague mrtaurho has made significant efforts towards the conjecture and I'll phrase his words here , and that'll count in the latest progress made in my conjecture over the sphere of internet. [Source - Aops community Forum named - Mathematics Research Station] mrtaurho's article - As brilliantly found by Alphatrion there are some matrices of size $n\times n$ of the given type with a determinant precisely 2^{n-1}. However, this value is far away from being maximal. I have not found a version of D_3 with determinant $\geqslant4$ but for all values $n>3$ (we may distinguish between odd and even n) it is worth considering matrices of the type $$D_{2k+1}^*=\begin{pmatrix}-1&\dots&1\\1&\ddots&\vdots\\1&\dots&-1\end{pmatrix}~~~D_{2k}^*=\begin{pmatrix}1&\dots&-1\\1&...&\vdots\\-1&\dots&1\end{pmatrix}$$ For those we gain $D^*_4=2^4=16,~D^*_5=s^4\cdot3=48,~D^*_6=2^7,~D^*_7=2^6\cdot5,~D^*_8=2^8\cdot3,~D^*_9=2^8\cdot7,\dots$ from where one can guess that in general we have $D^*_n=(n-2)2^{n-1}$ which is clearly greater than 2^{n-1} alone. This seems to work out but I have no idea how to prove this relation. [Small Remark: sticking to the type D^*_n given for odd n the formula changes to $D_n^*=(-1)^{n-1}(n-2)2^{n-1}$; so the alternate form for even n is just for aesthetics]. I have found the values using this online calculator Another observation worth being mentioned: $|D_n|\leqslant|D_{n+1}|$. This is quite simple by using block matrices as we have that $$|\det(D^\star_n)|=\left|\begin{pmatrix}\mathbf{1}&\mathbf{0}\\\mathbf{0}&\mathbf{D_{n-1}}\end{pmatrix}\right|=|1\cdot\det(D_n)|=|\det(D_n)|$$ And given D_n^* we know that there are configurations for D_n whose determinant exceeds D_{n-1}. To end this post I would like to mention Leibniz Formula for Determinants which states that $$\det(A)=\sum_{\tau\in S_n}\operatorname{sign}(\tau)\prod_{i=1}^na_{i,\tau(i)}$$ Here τ is a permutation of the symmetric group (on n letters) S_n and $\operatorname{sign}(\tau)$ returns the sign of the permutation τ. Note that there are precisely $|S_n|/2=n!/2=|A_n|$ even and thus $n!/2$ odd permutations. However, I do not see how one may apply this formula here but I thought it may be good to have it linked here. So the above article ends up here and the contribution of user Zatanna on Aops (which he gave values of D_5 & D_6 have been merged) . I choose ±1 to initially explain my observations because it's easier than picking up any other set of elements available to us, The results holds true for other specific cases too. Source - Mathematics Research Station (MARS)

论坛问题帖 | no_answers_at_retrieval | Alphatrion | 2019-08-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019010 | MSE-3336205

n^{th} Dimensional Prime Nexus Conjecture (Hamilton's Path & Cycle)

The system and proposition of Prime Nexus - We have to arrange $n>1$ Distinct natural numbers in a sequence such that the adjacent elements sums upto any Prime number. Convenience is when we set this problem to Graph Theoretic setting and specifically by using Hamilton's path and cycles. This system of clusters of natural numbers summing upto primes is called the Prime Nexus . Leave the question that at which extent it's helpful , likewise if the numbers are more then it starts getting tough . n^{th} Dimensional Prime Nexus Conjecture We can always generate any regular & definite n-th dimensional shape out of the nexus of Primes , specifically out of the Hamilton's path such that the nodes sums upto any Prime number. Suppose in order to arrange Distinct natural numbers in a sequence such that their adjacent elements sums upto any prime , we can have $1D$ Shape , $2D$, $3D$ and so on... generated out of the nexus , (Preferring the extension of squares & cubes...) I've worked out till $3D$ (can also replace 1 by 9 In $3D$) as the number of numbers increases in the cluster , the probability of getting more higher dimensional nexus gets increased. I strongly believe that the system never saturates at any k^{th} dimension. So If someone comes up with any counter example, actually he/she hasn't ! Because if we try generation of prime nexus for more higher number of numbers , we can always do that ! Please lemme know your opinions on it , how could anyone explain this enigmatic behavior of prime numbers making up higher dimensional shapes... And how could anyone even attack this conjecture! I'm very excited to learn new things regarding this conjecture, Everyone else here on MSE community holds more knowledge that I do , so please forgive me for any inconveniences followed up in presentation of my contents here. Regards

论坛问题帖 | has_answers | Alphatrion | 2019-08-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019011 | MSE-3344080

Conjecture: Any sufficiently big sum of three squares can be written as a square sum of three different natural numbers greater than zero

A natural number can be written as a sum of three squares if and only if it's not of the form $4^m(8n+7)$ for natural numbers m,n. I'm curious about such numbers that can be written as the square sum of three different natural numbers >0. My conjecture is: Any sufficiently big sum of three squares can be written as a square sum of three different natural numbers greater than zero. That is, there is a largest number $a^2+b^2+c^2$ which can not be written so that $0<a<b<c$. I would like to see a proof or a counter-proof or a related reference request. See also Primes of the form $a^2+b^2+c^2$, $0<a<b<c$

论坛问题帖 | has_answers | Lehs | 2019-09-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019012 | MSE-3346154

Natural numbers that are square sums of four different nonzero integers

It's known that any natural number can be expressed as a sum of four squares. Conjecture: A natural number can be written as a square sum of four different nonzero integers if and only if it not is a potency of two and not is a member of the set: {3,5,6,7,9,10,11,12,13,14,15,17,18,19,20,21,22,23,24,25,26,27,28,29,31,33,34,35,36,38,40,41,43,44,47,48,49,52,53,56,59,68,73,76,80,83,88,96,104,112,128,144,160,192,208,224,304,320,384,416,448,640,768,832,896,1280,1536,1664,1792,2560,3072,3328,3584,5120,6144,6656,7168,10240,12288,13312,14336,20480,24576,26624,28672}. So far tested up to 2^{16}.

论坛问题帖 | has_answers | Lehs | 2019-09-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019013 | MSE-3347920

Conjecture: $n\in\mathbb{Z}^+\Rightarrow 0=\int_{0}^\infty x^{4n-1}\sin(x)e^{-x}dx$

Conjecture: $n\in\mathbb{Z}^+\Rightarrow 0=\int_{0}^\infty x^{4n-1}\sin(x)e^{-x}dx$ This has been verified with WolframCloud for $1\leq n\leq 500$.

论坛问题帖 | has_answers | Nazgand | 2019-09-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019014 | MSE-3348582

Manipulating sets into periodic sets

Consider the set $[0,1,1,0,1,0,1,0,0,0,1,...]$. The nth element in this set is equal to 1 if n is prime and is else equal to 0. Here are my question: Is this Set non-periodic (If you write it as the fractional part of a binary number, the number is irrational)? To me it seems pretty obvious, because if it would be periodic we would have formula for the nth prime, but i cant come up with a explicite proof If we turn an infinite amount of 1's into 0's so that the set still contains of an infinite amount of 1's, can we create an periodic set? I think the answer is no, but i cannot come up with a proof of this either

论坛问题帖 | has_answers | Kinheadpump | 2019-09-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019015 | MSE-3353489

Is there a Collatz 'branch' stopping time equation?

Definitions: -Collatz: odd: $\frac{3x+1}{2}$ | even: $\frac{x}{2}$ -Branch: Starting at an odd number and increasing until reaching an even number. Then decreasing until reaching another odd number (ex: 7,11,17,26,13 -> 4 steps). Question: As far as I'm aware, there is not an equation where one can plug in a specific starting number and get the stopping time of Collatz for that number, but is there a known equation that gives the stopping time for the first 'branch' of Collatz for a given number (or in the case of 5, the full stopping time)?

论坛问题帖 | has_answers | JohnDNoone | 2019-09-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019016 | MSE-3355706

$\int_{0}^{\infty}\sin({x f({x})}) d x$ converges for positive monotone increasing unbounded continuous $f({x})$

Let a continuous speed function $f:\mathbb{R}^+\to\mathbb{R}^+$ exist such that $[{\{a,b\}\subseteq\mathbb{R}^+\land a<b}] \Rightarrow f(a)<f(b) $. Then $\int_{0}^{\infty}\sin({x f({x})}) d x$ converges. Looking at the positive and negative parts in pairs, if the integral starts with a positive chunk, the integral is positive, and if the integral starts with a negative chunk, then the integral is negative. Thus the first chunk is an upper bound if the chunk is positive, and the first chunk is a lower bound if the chunk is negative. The chunks approach zero because $f(x)\to\infty$ as $x\to\infty$, thus the upper and lower bounds approach each other. My problem with this argument is that even though this argument is obvious, the argument seems non-rigorous. I need the proof to convince Wolfram staff about a bug report: Integrate[Sin[x*Log[x+1]], {x, 0, Infinity}] is claimed to not converge because it converges too slowly.

论坛问题帖 | no_answers_at_retrieval | Nazgand | 2019-09-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019017 | MSE-3368938

A pattern on finding squares for which their sum is $\Big(\sum\limits_{i=0}^j x^i\Big)^2$

Apologies for the inactivity. I haven't been doing so well in life, lately, but I'm glad to be back at some maths! Here's a pattern I discovered. I'm not good at explaining with words, so hope you get it. It shouldn't be too difficult. Can it be proven? For all $x\in\mathbb{R}$, $$\begin{align}\bigg(\sum_{i=0}^1 x^i\bigg)^2&=x^2+2x+1\\ &=(x+1)^2\end{align}$$ $$\begin{align}\bigg(\sum_{i=0}^3 x^i\bigg)^2&=x^6+2x^5+3x^4+4x^3+3x^2+2x+1\\ &=(x+1)^2+2(x^2+x)^2+(x^3+x^2)^2\end{align}$$ $$\begin{align}\bigg(\sum_{i=0}^5 x^i\bigg)^2&= x^{10} +2x^9+3x^8+4x^7+5x^6+6x^5+5x^4+4x^3+3x^2+2x+1\\ &=(x+1)^2+2(x^2+x)^2+3(x^3+x^2)^2+2(x^4+x^3)^2+(x^5+x^4)^2\end{align}$$ Clearly there is a pattern with the exponents in the square of each partial sum, as well as the coefficients. Can one prove this pattern continues ad infinitum? For $\sum\limits^j$ I think the pattern also applies to even j and not just odd j as shown above, but I wrote this all on a napkin as I was eating at a restaurant, and now I've lost the napkin. (True story.) Any help would be much appreciated. Thank you :)

论坛问题帖 | has_answers | Mr Pie | 2019-09-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019018 | MSE-3369746

Rare gaps between integers having the same number of co-primes

Let $\varphi(x)$ be the Euler totient function. For $k = 1,2,3,\ldots$ I calculated the number of solutions of $\varphi(x) = \varphi(x+k)$. I observed that we have very few solutions when $k = 3,9,15,21,\ldots$ and a very large number of solutions when $k = 6,12,18,24,\ldots$ as shown below for $n \le 10^7$. Question 1: Why do we have a very few solutions when k is an odd multiple of 3 and a very large number of solutions when k is an even multiple of 3? Question 2: Is the following claim true? For any six consecutive positive integer values of k, and a sufficiently large x, the number of integers $n \le x$ such that $\varphi(n) = \varphi(n+k)$, is the lowest when $k \equiv 3 \text{ (mod 6)}$ and is the highest when $k \equiv 0 \text{ (mod 6)}$. Related question: Gap between integers having the same number of co-primes

论坛问题帖 | no_answers_at_retrieval | Nilotpal Kanti Sinha | 2019-09-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019019 | MSE-3370010

Can this heuristic argument be useful to prove: Sum of digits of a^b equals ab conjecture?

I've been trying to write a proof for the following conjecture (from this question): Let $s\left(a^{b}\right)$ denote the sum of the digits of a^{b} in base 10. Then the only integer values a,$b>1$ that satisfy $s\left(a^{b}\right)=ab$ are $(2,2),(3,3),(3,6),(3,9)$ and $(3,27)$. I found what I think is a heuristic argument, but i'm not sure if it can be useful in proving the conjecture. Let $d\left(n\right)$ denote the number of digits of integer n in base 10: $$d\left(n\right)=1+\left\lfloor \log_{10}n\right\rfloor $$ Let $s\left(n\right)$ denote the digit sum of integer n in base 10. Now from the conjecture, take for example the case $a=2$. I've been looking at the following sum: $$\sum_{n=1}^{b}\frac{s\left(2^{n}\right)}{\sum_{k=1}^{n}d\left(2^{k}\right)}$$ The plot of that sum for $1\leq b\leq20000$ looks like that: And now the same plot, with in orange, $9\log b$: The difference between the 2 curves quickly converges to a value c, and we see that: $$ \lim_{b\rightarrow\infty}\left(9\log b-\sum_{n=1}^{b}\frac{s\left(2^{n}\right)}{\sum_{k=1}^{n}d\left(2^{k}\right)}\right)=c\approx12.721\ldots $$ From this, we can also conclude that: $$ \frac{s\left(2^{b}\right)}{\sum_{k=1}^{b}d\left(2^{k}\right)}\sim9\log\left(\frac{b-1}{b}\right)\sim\frac{9}{b} $$ And since: $$ d\left(n\right)=1+\left\lfloor \log_{10}n\right\rfloor \approx1+\log_{10}n $$ Then: $$ \sum_{k=1}^{b}d\left(2^{k}\right)\approx\frac{b^{2}\log_{10}2}{2} $$ And: $$ s\left(2^{b}\right)\sim\left(\frac{9}{b}\right)\left(\frac{b^{2}\log_{10}2}{2}\right)\sim\left(\frac{9}{2}\right)b\log_{10}2s\left(2^{b}\right)\sim1.3546\times b $$ The same applies to other values of a, so more generally: $$ s\left(a^{b}\right)\sim\left(\frac{9}{2}\right)b\log_{10}a $$ Looking at plots of $s(a^b)$ for each values of a from 2 to 8, we can see this asymptotic relation seems to be very accurate. Now I have 2 questions: 1: Does the asymptotic relation above is correct, or is there some errors in my reasoning? 2: Since $a>\left(\frac{9}{2}\right)\log_{10}a$, does an asymptotic relation like that is enough to prove $s\left(a^{b}\right)<ab$, for sufficiently large b? Any help or advice would be appreciated.

论坛问题帖 | has_answers | François Huppé | 2019-09-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019020 | MSE-3383921

Yet a conjecture about the prime counting function.

This is a conjecture emanating from random tests with my set routines written in Forth. $a>b \;\wedge \pi(a+b)=\pi(a)+\pi(b)\implies b<11$ Furthermore, if $b\in\mathbb N$, then $b\in\{0,1,2,3,4,9,10 \}$ I would like (partly) proofs or counterexamples. Tested for $b<a<100,000$. Testing the numbers of hits for the non trivial values of b and for some upper limits: 1 2 3 4 9 10 100 74 48 8 14 2 2 1000 832 332 34 66 4 6 10000 8771 2454 204 406 11 20 100000 90408 19180 1223 2444 37 72 1000000 921502 156992 8168 16334 165 328

论坛问题帖 | has_answers | Lehs | 2019-10-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019021 | MSE-3383965

Convergence of Laplace transform

Suppose that $F: \mathbb{C} \to \mathbb{C}$ is analytic when $\operatorname{Re}(z)>0$. Assume that it is possible to show that F can be represented as a Laplace transform $$ F(z) = \int_0^{ + \infty } {\rm e}^{ - zt} f(t)\,{\rm d}t, $$ for $\operatorname{Re}(z)>a$ with some non-negative constant a and f being analytic in a neighbourhood of the positive real line. Does it follow from the analyticity of F for $\operatorname{Re}(z)>0$ that $a=0$?

论坛问题帖 | has_answers | Gary | 2019-10-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019022 | MSE-3394755

Some lesser known open problems/ conjectures in number theory

What are some lesser known problems/ conjectures in number theory (especially on prime numbers) which have evolved in these recent years and didn't got much of attention, and obviously wasn't featured in any global mathematical mainstream , It includes equivalences of pre existing conjectures/ results... Please let me know, thanks in advance

论坛问题帖 | has_answers | Alphatrion | 2019-10-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019023 | MSE-3396925

Prime twins $ (3^n - 2, 3^n - 4) $ conjecture

Let n be a positive integer. Conjecture There are infinitely many prime twins of the form $$ (3^n - 2, 3^n - 4) $$ Examples include $$(3^2 - 2,3^2 - 4) = (7,5) $$ $$ (3^{37} - 2 , 3^{37} - 4) $$ $$ (3^{41} - 2 , 3^{41} - 4) $$ Notice $2,37,41$ are primes themselves. Coincidence ? This resembles conjectures about prime twins , fermat primes and mersenne primes. So I do not expect a proof :) I already know the typical statistical arguments why this should be false (zeta(2) type sums as expected amount) , so I Will not ask for that either. My mentor believes this. How many values are known apart from these ? Are there equivalent conjectures known ? Is a disproof possible ? I was not able to disprove it. I tried mod arguments. Any ideas ?

论坛问题帖 | no_answers_at_retrieval | mick | 2019-10-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019024 | MSE-3402845

Does this procedure always generates at least one prime?

Suppose that p_n is the n-th prime and $n \neq 1,2$. To every p_n we can associate $(n-2)$-tuple $(2p_n+p_{n-1},...,2p_n+p_2)$ and from some calculations that I have done it seems that it could be that at least one of the numbers from that $(n-2)$-tuple for every p_n is a prime number. Is that true?

论坛问题帖 | has_answers | user716491 | 2019-10-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019025 | MSE-3409889

A conjecture about the number of divisors of a natural number

Conjecture: $\tau(n)\mid\tau(n^2)\iff$ n is a perfect square and $\sqrt n=p^2s$, where p is prime and s is a non prime squarefree number such that $\gcd(p,s)=1$. $\tau(n)$ is the number of factors of n. See this question on MSE: Number theory problem on divisors! It's tested for $n<10,000$.

论坛问题帖 | has_answers | Lehs | 2019-10-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019026 | MSE-3411896

Has this basic number theory conjecture been discovered or proved before?

So I was sitting in a car bored for two hours playing around with the calculator on my phone, and I discovered that by picking two numbers (at least one being odd and both being not divisible by each other) I could always add them together to make many different numbers. Furthermore, I could always add and subtract them together make 1; this is significant because I could then negate and indefinitely duplicate the series to create any positive and negative integers. Here are some examples: 8-7=1 16+16+16+16-9-9-9-9-9-9-9=1 17+17+17-5-5-5-5-5-5-5-5-5-5=1 and so on. For convenience, I preferred to write these equations in the form a*x-b*z=1 8*1-7*1 = 1 16*4-9*7 = 1 17*3-5*10 = 1 Here is my Conjecture: Any two non-zero integers indivisible of each other, if at least one of them is odd, can be added or subtracted together indefinitely to make any integer. Another way of writing it (though it is more limiting) is this: Given that a and b are two non-zero integers indivisible of each other, if at least one of them is odd, there always exists some value of x and z for any given integer n that makes the equation ax-bz=n true. My question is, has anyone ever discovered or proved this conjecture before, and how can it be proven? It almost feels like the solution should be as obvious as the equation a*x-b*z=0 (which just requires finding the lcm of the two numbers) and that it's just a simple quirk in numbers based upon the already laid down basic laws of mathematics.

论坛问题帖 | has_answers | Zach K | 2019-10-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019027 | MSE-3417044

If $q^k n^2$ is an odd perfect number with special prime q, then can $n^2 - q^k$ be a cube?

Let $q^k n^2$ be an odd perfect number with special prime q satisfying $q \equiv k \equiv 1 \pmod 4$. This question is an offshoot of the following earlier post: If $q^k n^2$ is an odd perfect number with special prime q, then $n^2 - q^k$ is not a square. My question in the present post is the following: If $q^k n^2$ is an odd perfect number with special prime q, then can $n^2 - q^k$ be a cube? Not much can be said without some additional hypotheses. In the paper titled Improving the Chen and Chen result for odd perfect numbers (Lemma 8, page 7), Broughan et al. show that if $$\frac{\sigma(n^2)}{q^k}$$ is a square, where $\sigma(x)$ is the sum of divisors of $x \in \mathbb{N}$ and $q^k n^2$ is an odd perfect number with special/Euler prime q, then $k=1$. It follows that, if $\sigma(n^2)/q^k$ is a square, then $\sigma(n^2) \equiv 1 \pmod 4$, which implies that $q \equiv k = 1 \pmod 8$. Since n is odd, $n^2 \equiv 1 \pmod 8$. This implies that $8 \mid (n^2 - q^k)$. Alas, this is where I get stuck. I do not even know how to rule out the simplest case $$n^2 - q^k = 8.$$ An Elementary Attempt to Rule Out $n^2 - q^k = 8$ Subtract 9 from both sides, and transfer q^k to the RHS: $$n^2 - 9 = q^k - 1.$$ Factoring both sides, we get $$(n+3)(n-3)=(q-1)Q$$ where $Q=\sigma(q^{k-1})$. We get two cases: Case 1 $n+3 \mid (q^k - 1)$ This implies that $n + 3 \leq q^k - 1$, from which we obtain $$n < n + 4 \leq q^k.$$ But we know from work of Brown (2016), Dris (2017), and Starni (2018) that $q < n$ holds unconditionally. Thus, we get $$q < n < q^k$$ which contradicts $k=1$. Case 2 $n - 3 \mid (q^k - 1)$ This implies that $$n - 3 \leq q^k - 1$$ which means that $$n - 2 \leq q^k.$$ Again, by the results of Brown, Dris, and Starni, we have $$q - 2 < n - 2 \leq q^k,$$ which does not contradict $k=1$. Can anything be said about the general case, preferably by relaxing the hypotheses? UPDATE (1 NOV 2019 - 08:32 AM Manila time) Blimey! I did not realize that Case 1 $n+3 \mid (q^k - 1)$ holds in general. Thus, under the assumption that $\sigma(n^2)/q^k$ is a square, $n^2 - q^k \neq 8$.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2019-10-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019028 | MSE-3418018

Why is that a conjecture is written and its proof can not be done for years?

Why is it that if a conjecture is written and its proof can not be done for years, means if its statement is understandable in language, why is the proof not possible if the conjecture is correct? E.g., the proof of Fermat's last theorem (which was very clear what is it saying) was done 350 years later than it was conjectured and it took 130 pages. Fermat couldn't think for 130 pages and then give the conjecture. So why is the proof so horrible for such simple equation?

论坛问题帖 | has_answers | ANUJ GUPTA | 2019-11-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019029 | MSE-342124

Conjecture to start a proof

In an inductive proof example, my book starts with the identity stating $$\sum\limits_{i=1}^n \sum\limits_{j=1}^i j = \frac{n(n+1)(n+2)}{6}\;.$$ As a sidenote, it says they reached this identity by conjecture, calculating the double sum for low values of n. How on earth did they see this? I attempted this expansion (don't know if it's right): $$\sum\limits_{i=1}^n \sum\limits_{j=1}^i j = \sum\limits_{i=1}^n (1 + 2 +\ldots+i) = 1 + 2 + \ldots + \sum\limits_{i=1}^n i = 1 + 2 + \ldots + (1 + 2 + \ldots + n)\;.$$ From this, I don't see how they arrived at the fractional cubic polynomial. Any help would be appreciated! Thanks.

论坛问题帖 | has_answers | user67241 | 2013-03-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019030 | MSE-3437560

For all $n\in\mathbb{N}$ with $n>1$ there exists an $m\in\mathbb{N}$ with $m<n$ such that $mn+1$ is prime.

I was using Python to explore some interesting lexicographically earliest sequences. I was looking into the sequence of numbers such that $a(1)=1$ and $a(n)$ is the smallest number such that $a(k)n+1$ is not prime for any $k<n$. After computing this sequence, it appeared to be exactly the odd integers. The absence of even integers interested me, as that suggested that for every even integer n there exists an odd $m<n$ such that $mn+1$ is prime. After checking with Mathematica, this appeared to be true. In fact, this suggested the more general conjecture in the title: For all $n\in\mathbb{N}$ with $n>1$ there exists an $m\in\mathbb{N}$ with $m<n$ such that $mn+1$ is prime. This doesn't sound too difficult, but then again there are plenty of nice-sounding prime conjectures that remain unsolved, so this could be one of those. How could this be proven? And if its unsolved / really difficult, what is the name of the conjecture?

论坛问题帖 | has_answers | volcanrb | 2019-11-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019031 | MSE-3452492

Conjecture about generalized means

Conjecture Every continuous function $f(v_1,\dots,v_n)$ of n arguments such that it lies in the closed interval $[\min(v_{1},\dots ,v_{n});\max(v_{1},\dots ,v_{n})]$, is symmetric for all permutations of arguments, and is homogeneous (with the degree 1) necessarily is a generalized mean. Help me to prove or disprove (with a counterexample) this conjecture. Is it a known result?

论坛问题帖 | no_answers_at_retrieval | porton | 2019-11-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019032 | MSE-3452518

Does "finite average" exist?

I will call an average any continuous function $f(v_1,\dots,v_n)$ of n arguments such that it lies in the closed interval $[\min(v_{1},\dots ,v_{n});\max(v_{1},\dots ,v_{n})]$, is symmetric for all permutations of arguments, and is homogeneous with the degree 1. Question: Can an average not to tend to infinity when one of the arguments tends to infinity and the rest arguments are fixed nonnegative reals? (We can assume $n\geq 2$.) My original question had a trivial solution $\min(v_{1},\dots ,v_{n})$, so let's add an additional requirement: our average not to tend to zero when one of the arguments tends to zero and the rest arguments are fixed positive reals? (We can assume $n\geq 2$.)

论坛问题帖 | has_answers | porton | 2019-11-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019033 | MSE-3452562

Does “smooth finite average” exist?

(That question was answered. I realized I asked a wrong question. So here is a modified question.) I will call an average any continuous function $f(v_1,\dots,v_n)$ of n arguments such that it lies in the closed interval $[\min(v_{1},\dots ,v_{n});\max(v_{1},\dots ,v_{n})]$, is symmetric for all permutations of arguments, and is homogeneous with the degree 1. Question: Can an average provided the below additional requirements (I ask for the case if all of them are required but also for milder cases of just a part of these requirements) not to tend to infinity when one of the arguments tends to infinity and the rest arguments are fixed nonnegative reals? (We can assume $n\geq 2$.) Our average does not tend to infinity when all arguments except one tend to infinity. Our average is C^1-smooth in every argument (maybe except the argument(s) value equal to zero). Our average is C^∞-smooth in every argument (maybe except the argument(s) value equal to zero).

论坛问题帖 | has_answers | porton | 2019-11-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019034 | MSE-3453668

Lower bound for the n-th record gap between primes in an arithmetic progression

(This is a natural counterpart to question 3132001 which deals with upper bounds.) Let q and r be coprime integers, $1\le r < q$, and consider the arithmetic progression $$ r, \ r+q, \ r+2q, \ r+3q, \ldots \tag{P} $$ Dirichlet proved that there are infinitely many primes in progression (P) when $\gcd(q,r)=1$. This is Dirichlet's theorem on arithmetic progressions. Let $R(n,q,r)$ be the nth record gap between primes in progression (P). For example, with $q=6$ and $r=5$, we have $R(n,6,5)=\mbox{A268928}(n)$; see http://oeis.org/A268928. Conjecture: Almost all record gaps $R(n,q,r)$ satisfy $$ R(n,q,r) > {1\over6} \varphi(q) n^2. \tag{L} $$ Here $\varphi(q)$ is Euler's totient function. Question: Find counterexamples to inequality (L). (Please find as many as you can. You will likely need to write a program and run it long enough.)

论坛问题帖 | no_answers_at_retrieval | Alex | 2019-11-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019035 | MSE-3455970

Conjecture: smallest missing mod value always yields previous prime

I've come up with a conjecture that seems similar in strength to Legendre's or Oppermann's, but maybe subtly different. Let a_n be the smallest nonnegative value such that there is no m in $1<m<n/2$ where $n \equiv a_n \pmod m$. Then for all $n>2$, we have $n-a_n=p_{\pi(n)-1}$, the closest previous prime to n. Take $n=16$ as an example: $$\begin{eqnarray} 16 &\equiv 0 \pmod 2 \\ &\equiv 1 \pmod 3 \\ &\equiv 0 \pmod 4 \\ &\equiv 1 \pmod 5 \\ &\equiv 4 \pmod 6 \\ &\equiv 2 \pmod 7 \end{eqnarray}$$ The smallest value not seen is $a_n=3$, and $16-3=13$ is the previous prime. In cases where n itself is prime, e.g. 17 yielding the values $\{1,2,1,2,5,3,1\}$, you can either interpret 0 as the missing value and 17 as the prime, or 4 giving $17-4=13$. (I'm not sure which is the more consistent interpretation.) I've verified this empirically through 10^5, but cannot come up with a proof. In fact, I suspect a proof would be very difficult since what this seems to come down to is whether there is always a prime in the interval $(n,n+d)$ for a composite n, where d is the largest proper divisor of n. This has its worst case for forms of p^2, which seem to require a prime in $\left(p^2, p(p+1)\right)$. Note that when $a_n < \lfloor \sqrt{n} \rfloor$, it is easily provably true; the problem is you can't guarantee it will be in that range, despite the fact that it almost certainly is for all $n \geq 127$. I'm curious whether this conjecture already exists somewhere or is actually equivalent to one of the better-known prime gap conjectures. Better yet would be a proof, but that's obviously wishful thinking.

论坛问题帖 | has_answers | Trevor | 2019-11-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019036 | MSE-3455999

Justification for the conjectures on self-avoiding walks

There are two conjectures on self-avoiding walks that I haven't seen any justification of. One is the asymptotic growth of the number of length n self-avoiding walks, the form I've seen most often is $$ c_n \sim A\mu^n n^{\gamma-1}, $$ where A, μ, γ are constants. Of all the literature I've seen, I haven't read about any justification for this formula, rigorous or not. The other conjecture is stated in the Wikipedia page for self-avoiding walks and says that finding the number of such walks in a given lattice is NP-hard. There was no citation given for this, so I must ask where does this assertion appear in the literature. Edit: Any papers referencing the NP-hardness conjecture should be about counting SAWs in the lattice $\mathbb{Z}^2$ or $\mathbb{Z}^n$

论坛问题帖 | no_answers_at_retrieval | zetapenguin | 2019-11-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019037 | MSE-3456415

p, q and $\sqrt[n]{p} + \sqrt[n]{q}$ are rational, with the latter being non-zero. Are $\sqrt[n]{p}$ and $\sqrt[n]{q}$ rational?

Let $p, q \in \mathbb Q$, $n \in \mathbb Z^+$ and label $a = \sqrt[n]p, b=\sqrt[n]q$. Conjecture: If $a + b$ is a non-zero rational, then both a and b are rational. (Preliminary question: is this a known result I'm not aware of?) I believe I have found a partial proof of the above. Namely, I first proved that ab is rational for $n = 1, 2, 3$. For $n = 2$, $(a + b)^2 = a^2 + 2ab + b^2$, so $$ab = \frac{(a+b)^2-a^2-b^2}{2} \in \mathbb Q$$ For $n = 3$, $(a + b)^3 = a^3 + 3ab(a + b) + b^3$, so $$ab = \frac{(a+b)^3-a^3-b^3}{3(a+b)} \in \mathbb Q$$ As it turns out, an additional assumption that $ab \in \mathbb Q$ allows proving the conjecture. The question: how can I prove $ab \in \mathbb Q$ for $n > 3$? (or what numbers are a counterexample?) The details of the proof assuming $ab \in \mathbb Q$ follow. Consider the polynomial $(x - a)(x - b) = x^2 - (a + b)x + ab$. Note that its coefficients are rational. This means that $\Pi \in \mathbb Q[x]$, the minimal polynomial of a, is of degree at most 2. Hence, $\deg \Pi \in \{1, 2\}$. If $\deg \Pi = 1$, then a is rational, which was to be proven, so let's assume that $\deg \Pi = 2$ and hope for a contradiction. As the minimal polynomial is unique, $\Pi = (x - a)(x - b)$. Moreover, Π divides $x^n - p$, since the latter has a root at a. Hence, the roots of Π are a subset of the roots of $x^n - p$. For odd n, we have $\{a, b\} \subseteq \{a\}$, so $a = b$. For even n, we have $\{a, b\} \subseteq \{a, -a\}$. b can't be equal to $-a$, since roots of even degree are nonnegative (or imaginary, but in that case $a + b$ wouldn't be rational). Hence, $a = b$. In both cases, from the assumption $a + b \in \mathbb{Q} \setminus \{0\}$, we get $2a \in \mathbb{Q} \implies a \in \mathbb Q$ and therefore $\deg \Pi = 1$, which is a contradiction.

论坛问题帖 | has_answers | Maya | 2019-11-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019038 | MSE-3465494

The crosshatch conjecture, on primes in (p,p^2)

If the first p^2 integers are laid out in a $p\times p$ square, every row and column will have at least one prime. Easily visualized as so: I recognize this should maybe be packaged as two conjectures, but hey, I like the aesthetics. Individually, we have: Given some prime p, there will always be at least one prime in $[kp+1,kp+p-1]$ for all $1 \leq k \leq p-1$. Given some prime p, there will always be at least one prime q in (p,p^2) such that $q \equiv k \pmod{p}$ for all $1 \leq k \leq p-1$. I'm asking if a) this is somehow wrong or someone finds a counterexample, and b) if either of these conjectures already exist.

论坛问题帖 | has_answers | Trevor | 2019-12-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019039 | MSE-3467455

A pre-successor hyperoperation: subplus

I wrote up these notes about a month ago and just found them. I could really use a second opinion or two. I notice there's a whole question about how my whole thesis is wrong, but that doesn't automatically mean it's without merit. That said, forgive my layman's take on everything. Addition is a special case of multiplication. That follows from the whole set theory interpretation where units are indistinct in addition, but have types in multiplication. By that logic, you could also argue that multiplication is a special case of exponentiation. After all, $(a^b)^c=a^{bc}$, and $a^b \cdot a^c = a^{b+c}$. So, it's interesting; if you ignore the base, if you're taking a power, you're multiplying, as in the first case; and if you're multiplying, you're adding, as in the second case. What about if you're adding? It'd be nice if it turned out to be a^{b+1}, but I don't see how that's easily compatible. Adding 1 seems to require $a^b \cdot a=a^b \cdot a^{a^0}=a^{b+1}$. And on the other side, what operation gives exponentiation? Tetration? Hmm. Maybe adding requires another level of implicit base below everything. Even if it doesn't "exist" in a useful sense, other than to paint a more complete logical framework, or to serve as an intermediary tool. Putting this into action, addition is still just exponentiation, just with one deeper level of implicit base. Asking what's below addition is meaningless unless we allow that addition is just a label we've applied to that level of abstraction. We'll appropriate the symbol $\oplus$ for this hypothetical operation, and let γ serve as an exponential base. From powers on down, the four operations we've got now, reframed as exponentiation, look like: $$ \begin{align} x\textrm{ ^ }y&\Rightarrow&& x^y \\ x\times y&\Rightarrow&& (\alpha^x)^y = \alpha^{xy} \\ x+y&\Rightarrow&& (\beta^{\alpha^x})^{\alpha^y} =\beta^{\alpha^x \cdot \alpha^y} =\beta^{\alpha^{x+y}} \\ x\oplus y&\Rightarrow&& (\gamma^{\beta^{\alpha^x}})^{\beta^{\alpha^y}} =\gamma^{\beta^{\alpha^x} \cdot \beta^{\alpha^y}} =\gamma^{\beta^{(\alpha^x + \alpha^y)}} =\gamma^{\beta^{\alpha^{x \oplus y}}} \end{align} $$ One actual effect of this is we can infer that $x \oplus y$ is arbitrarily close to $\max(x,y)$. This is using $x\oplus y := \log_\alpha (\alpha^x+\alpha^y)$, but there are other possible interpretations of the extrapolation above. Mine was just an illustrative guess. But if we go ahead with the implication, I'm tempted to assign a real base value for more detail. More specifically, if the base is e, it's very close to $\max(x,y)$; if the base is $-e$, it's very close to $-\min(x,y)$. Upon some reflection though, I think the correct choice of base is $\alpha=2$, because then if $x=y$, you get $x \oplus y = x+1=y+1$, the successor function, which is what one expects for this level of hyperoperation. I acknowledge it's unusual to have a binary successor operation, it being famously unary and all. However, rather than break it, I think we've arguably improved it. By the same token, it might be reasonable to replace all the undefined base variables above with 2s. Note we have $x \oplus 0=\log_2 (2^x+1)$. As with every binary operation, all else being equal, the closer the two operands are to each other, the more the value is maximized. As an aside, it is awfully easy to represent everything computationally as exponentiation (even if it is just wrapping operations up in logs and powers). Observe: xTimes[a_,b_]:=Log2[(2^a)^b] xTimes[3,5] -> 15 xPlus[a_,b_]:=Nest[Log2,(2^2^a)^2^b,2] xPlus[3,5] -> 8 xSucc[a_,b_]:=Nest[Log2,(2^2^2^a)^2^2^b,3] N@xSucc[-100,-50] -> -50. N@xSucc[-5,19] -> 19. xSucc[-5,-5] -> -4 xSucc[4,4] -> 5 FYI - the Nest function

$\mathsf{nest}(f,n,k):=f(f(\ldots f(f(n))\ldots \textrm{k times} \ldots))$ So generally, I'm looking for responses like: a) "interesting ideas, worth considering further", or b) "crackpot idea, go read a book". I did find a post alluding to a well-regarded professor that seems to have had a similar thought, although I'm still trying to get details on that.

论坛问题帖 | no_answers_at_retrieval | Trevor | 2019-12-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019040 | MSE-3477463

2'nd Law of thermodynamics in a relativistic gas?

Question I'm trying to understand in the context of a gas if a particular boundary condition can be used be to be the source of the 2'nd law of thermodynamics: "Given an isolated system the entropy will increase or remain the same." Is the attempt correct? Background Entropy is a statement of the likelihood of a configuration. If that is reflected in the action as well I think I have a chance? I try to make use of the fact it is only in thermal equilibrium (maximum entropy) that collisions can be neglected. Usually, the Hamiltonian of a gas at thermal equilibrium does not include a collision term which would imply at a collision: $$ \underbrace{\cdot}_{A} \rightarrow \leftarrow \underbrace{\cdot}_{B}$$ $$ \underbrace{\cdot}_{B} \leftarrow \rightarrow \underbrace{\cdot}_{A}$$ They actually go through each other. My attempt Consider a relativistic gas (point) particles with a 2 particles A and B in a box and the only collide once. The line element of the $A$and B before a collision is given by $ds_i^2 $ where $i=A$ or $i=B$. Similarly, the action is given by: $$ S_i = - m_ic \int_P d s_i$$ Where P is the world line before the collision $s_A^\mu = s_B^\mu$ where s_i^μ is the four position vector. After the collision, we know the momentum p^μ is conserved: $$ p_A^\mu + p_B^\mu = {p'}_A^\mu + {p'}_B^\mu$$ where ${p'}_i^\mu$ denotes the momentum after the collision. Differentiating with respect to $\frac{d}{ds_i}$ and using $ \frac{d p^\mu_i}{ds_i} =0$ then: $$ \frac{d p_j^\mu}{ds_i} = \frac{d {p'}_A^\mu}{ds_i} + \frac{d {p'}_B^\mu}{ds_i}$$ with $j \neq i$ After the collision the action is given by: $$ S'_i = - m_i c \int_{P'} d s'_i$$ where P' is the world line after the collision and $d s'_i$ is defined by $\frac{dp'_i}{ds'_i} = 0$. Let us write ds_A in terms of ds'_A. We proceed with: $$ \frac{d {p}_A^\mu}{ds_i} + \frac{d {p}_B^\mu}{ds_i} = \frac{d {p'}_j^\mu}{ds_i}$$ Using the chain rule: $$ \frac{d {p}_A^\mu}{ds_i} + \frac{d {p}_B^\mu}{ds_i} = \frac{d {p'}_j^\mu}{ds'_j} \frac{d s'_j}{ds_i} = \frac{\frac{d {p'}_j^\mu}{ds'_j}}{(\frac{d s'_j}{ds_i})^{-1}}$$ And as the L.H.S above is finite: $$\frac{d {p'}_j^\mu}{ds'_j} \to 0 \implies (\frac{d s'_j}{ds_i})^{-1} \to 0$$ Using L' Hopital Rule: $$ \frac{d {p'}_j^\mu}{ds_i} =\frac{d {p}_A^\mu}{ds_i} + \frac{d {p}_B^\mu}{ds_i} = \frac{\frac{d^2 {p'}_j^\mu}{{ds'_j}^2}}{\frac{d}{ds'_j}(\frac{d s'_j}{ds_i})^{-1}}$$ Or: $$\frac{d}{ds'_j}(\frac{d s'_j}{ds_i})^{-1} \frac{d {p'}_j^\mu}{ds_i} = \frac{d^2 {p'}_j^\mu}{{ds'_j}^2}$$ The above should be solvable as we have 2 boundary conditions (conservation of momentum and $s_i = s_j$). Hence: $$ {ds_i} = {d s'_j} \int \frac{d^2 {p'}_j^\mu}{{ds'_j}^2} (\frac{d {p'}_j^\mu}{ds_i})^{-1} {ds'_j}$$ Hence, the action of particle i is: $$ S_i = - m_ic \int_P {d s'_j} \Big (\int \frac{d^2 {p'}_j^\mu}{{ds'_j}^2} (\frac{d {p'}_j^\mu}{ds_i})^{-1} {ds'_j} \Big) - m_i c \int_{P'} d s'_i$$ One can take sum over i to find the net action. Heuristically? Since, the action depends on where the gas collides. It is possible to use this as a source for the second law in the limit of an infinite collisions?

论坛问题帖 | no_answers_at_retrieval | More Anonymous | 2019-12-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019041 | MSE-3479074

Conjecture: All x^2+1 primes for $x>90$ can be represented as the sum of five x^2+1 primes.

For all $x>90$, I assert that any x^2+1 prime may be written as the sum of five smaller x^2+1 primes. In fact, above that bound, I think a stronger conjecture holds where one of said primes can always be 5. e.g. $$\begin{array}\\ 26^2+1&=677\\ &=401+197+37+37+5 \\ &=(20^2+1)+(14^2+1)+2(6^2+1)+(2^2+1) \end{array}$$ The only exceptions appear to be $x\in\{1,2,16,20,90\}$. The primes need not be distinct, although it seems that there is always at least one such representation for $x>170$. All statements above have been confirmed up to at least 10^6 with an apparent strong trend toward increasing numbers of representations. Below is a plot of the number of different representations for each x^2+1 prime, found by a brute force check using any five x^2+1 primes up through $1060^2+1=1123601$ (the 117th Landau prime). As expected, it only begins tapering off around that bound, and if I hadn't stopped there, I'd expect both minima and maxima to continue to rise. As an informal observation, it also seemed that the largest prime representable was gradually approaching $5(x^2+1)$. I'm interested in insight on the conjecture and/or confirmation/counterexamples. I am also curious whether this is related to (or even follows from) Lagrange's four-square theorem.

论坛问题帖 | has_answers | Trevor | 2019-12-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019042 | MSE-3487963

Do all the zeroes of this arithmetic function occur at even integers?

Let $a(n)$ be the number of natural numbers $\le n$ which have an odd number of distinct prime factors. Let $b(n)$ be the number of natural numbers $\le n$ which have an even number of distinct prime factors. Conjecture: If $a(n) = b(n)$ then n is even. I have verified this conjecture for each of 8883 zeroes of $s(n) = a(n)-b(n)$ for $n \le 1.25 \times 10^8$. Is this true in general? Sage code n = 1 s = f = 0 target = set = 10^6 while True: s = s + (-1)^(len(prime_divisors(n))) if s == 0: f = f + 1 if n%2 == 1: print "Odd zero", n break if n > target: print n,f target = target + set n = n + 1

论坛问题帖 | has_answers | Nilotpal Kanti Sinha | 2019-12-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019043 | MSE-3497098

Can gcd separate primes from composite numbers?

Let $x_i=i+1, 1\le i \le n$, $y_i = +1,$ if x_i is prime, -1 otherwise. Conjecture: Then (?) there exists $b,c_i \in \mathbb{R}, 1 \le i \le n$ such that for all $j=1,\ldots,n$: $$y_j = \text{ sgn }\left(\left[\sum_{i=1}^n c_iy_i\gcd(x_i, x_j)\right] - b\right)$$ Motivation for asking this question: https://mathoverflow.net/questions/349589/are-primes-linearly-separable Thanks

for your help! this question has been answered at mo.

论坛问题帖 | no_answers_at_retrieval | user276611 | 2020-01-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019044 | MSE-3499800

A conjecture about a smothness concerning consecutive primes

This is a conjecture inspired by the relative but piecewise smoothness of the sequence https://oeis.org/A068901/graph?png=1, about the least number a_n, such that $n\mid p_n+a_n$, where p_n is the n-th prime. Conjecture: $n\mid p_n+a \wedge 0≤a<n \Rightarrow \exists b\in\mathbb Z: n+1\mid p_{n+1}+b$, where $|a-b|<2\cdot\sqrt n$. Are there counter-examples, or can the conjecture be proved? Tested for all primes $p_n<100,000,000$. The conjecture badly describe the real smothness. Making some more experiments lead me to comment my question with an other conjecture: As above, given $\varepsilon>0$ there exist N such that $n>N \Rightarrow |a-b|<\varepsilon\sqrt n$.

论坛问题帖 | no_answers_at_retrieval | Lehs | 2020-01-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019045 | MSE-3507935

Greatest common divisor of consecutive square free numbers

I guess that every prime number occurs as the greatest common divisor of two consecutive square free numbers, which I don't expect a proof of. But I've done some experiments indicating that: If m, n are consecutive square free numbers then $\gcd(m, n)$ is not composite. Is that true and can it be proved?

论坛问题帖 | has_answers | Lehs | 2020-01-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019046 | MSE-3509799

Prove or Disprove the Conjecture on Cousin Primes

Here is the problem in a nutshell - Ethers or Ether of n, $\mathbf{\eth(n)}$: Ethers of any number n is the set of all the values of every possible permutations of its digits. Illustration Ethers of 123, $\eth(123)$= $123, 132, 213, 231, 312$ and 321. Similarly it follows that $\eth(123)=\eth(132)=\eth(213)=\eth(231)=\eth(312)=\eth(321)$ Magnitude of Ether n, denoted by $|\eth(n)|$ is the total count of Prime Numbers we obtain by considering all of its Ethers individually. Illustration $|\eth(38)|=\mid\eth(83)|=1$, $|\eth(67)|=1$, $|\eth(251)|=2$, $|\eth(157)|=3$, $|\eth(973)|=4$, $|\eth(2183)|=6$, $|\eth(1379)|=7$, $|\eth(7309)|=9$, $|\eth(5917)|=10$, $|\eth(1237)|=11$ and so on... Zeroes of Ether Any natural number n such that $| \eth(n)|= 0$ is termed as zeroes of Ether, and if such n can be explained, trivially, then it's termed as Trivial Zeroes of Ether. Trivial Zeroes of Ether are classified as follows 1.) All the multiples of 3, $| \eth(3m)| = 0$, $m \in{W}$ 2.) The number having any of these digits $(0, 2, 4, 5, 6, 8)$ in any combinations. 3.) Any number (>11) entirely made up of only a single digit (>1). 4.) All the numbers of form $k \cdot 10^\alpha$, $\mathit{| \eth(k\cdot 10^\alpha)| =0, \, 0 \leq k \leq 9, \, \alpha\in{W}}$ CONJECTURE 1 If $P_{n+1} - P_n =2$ then there always exists k such that $| \eth(k)| = 0$ for $P_n < k < P_{n+1}$. CONJECTURE 2 If $P_{n+1} - P_n = 4$ then there exists minimum one k such that $| \eth(k)| \geq 1$ for $P_n < k < P_{n+1}$ where all of its Exceptions are Trivial Zeroes of Ether. It's just a segment from my original post On The Distribution of Prime Numbers I tried my best to handle this Conjecture 2 but couldn't get anywhere. I thought the representation of Primes in base 10 would incinerate this problem but no progress was made. Any progress will be highly appreciated. Remark \eth symbol is being utilitized for the new notions of digit permutations. Edit: Rephrased the Conjecture for better clarification.

论坛问题帖 | has_answers | Crystal Wolftron | 2020-01-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019047 | MSE-3510517

Do all elements of $[n+1,2n]$ have strictly higher gpf than elements of $[1,n]$ when sorted by gpf?

For any $n\in\mathbb N$, let $A=\{x\in\mathbb N \mid 1 \leq x \leq n\}$ and $B=\{x\in\mathbb N \mid n+1 \leq x \leq 2n\}$. Order the sets by greatest prime factor of each element, ascending. Let A_1 be the element in A with the smallest gpf and A_n be the largest, and likewise for B. Conjecture: $\mathrm{gpf}(A_k)<\mathrm{gpf}(B_k)$ for all $1 \leq k \leq n$. e.g. Let $n=5$, and we have: $$ \begin{array}{|||||||} \hline A_k&\mathrm{gpf}(A_k)&\mathrm{gpf}(B_k)&B_k\\ \hline 1&1&2&8=2^3\\ \hline 2&2&3&6=2\cdot 3\\ \hline 4&2&3&9=3^2\\ \hline 3&3&5&10=2\cdot 5\\ \hline 5&5&7&7 \\ \hline \end{array} $$ where $\mathrm{gpf}(B_k)$ is strictly greater than $\mathrm{gpf}(A_k)$. I confirmed this empirically for $n<3000$. Informally, it looked like the trends strongly suggest it won't fail at any point. If this is a known result or conjecture, please share. Otherwise, I'm looking for counterexamples or any advice on how a proof could be approached$-$or a reason why a proof is likely to be difficult or impossible, as seems more typical with this stuff. Quicky data tables for $n<70$ are here.

论坛问题帖 | has_answers | Trevor | 2020-01-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019048 | MSE-3512721

A conjecture in number theory with twin primes

It's a conjecture found with the help of Wolfram Alpha : Let p_i be the first primes and $n>5$ with n an odd natural number numbers we are interested by the quantity : $$A=(1+p_1\times p_2(1+p_3\times p_4(1+p_5\times p_6(\cdots(1+p_n\times p_{n+1})^{\frac{1}{2}})\cdots)$$ Or $$A=(1+2\times 3(1+5\times 7(1+11\times 13(\cdots(1+p_n\times p_{n+1})^{\frac{1}{2}})\cdots)$$ Where p_n and p_{n+1} are twin primes numbers Example $$(1+2\times3(1+5\times7(1+11\times13(1+17\times19(1+23\times29(1+31\times37(1+41\times43)^{\frac{1}{2}})))))=311677481085187=7×43×433×2391393439$$ Conjecture 1 The last digit of A is seven . Conjecture 2 If A is not a prime number then A is divisible by 7. I try to work with divisibility rule for small numbers but it becomes insane with big numbers . I'm a very beginners in number theory so if you could use elementary tools it will be cool . Thanks a lot for your time and patience .

论坛问题帖 | has_answers | Barackouda | 2020-01-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019049 | MSE-3513415

A conjecture concerning Pollard rho

I'm investigating the Pollard rho factorization algorithm, searching for a proof that any odd composite can be factorized for some start number. I'm only studying the standard function $f(x)=x^2+1\mod n$, meaning the rest of x^2+1 when divided with n and have found this conjecture: If n is odd then $\;2*|\operatorname {Im} f|=n+1\iff n$ is prime. I would like to see a proof or a counter-example.

论坛问题帖 | has_answers | Lehs | 2020-01-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019050 | MSE-3517807

The Rook Conjecture: arrangement of p primes being distinct $\pmod{p}$ through p^2

For any prime p, divide $[1,p^2]$ into p equal intervals of length p, so that the first interval is $[1,p]$, the next $[p+1,2p]$, and so on. It is definitely unproven but seems likely that there will always be a prime in every one of these intervals. It is even probably true when you substitute any natural n in place of the prime p above. I've noticed a stricter condition that seems to hold. If you divide p^2 as described above, not only will there be a prime in every interval of length p, but there will always be at least one way to select those primes such that no value is repeated $\bmod{p}$. A good name for this might be the Rook Conjecture, as this is equivalent to saying that if you number the squares of a $p\times p$ sized chessboard, there is always at least one way to place p rooks on prime-numbered squares yet have no rooks share a rank or file. Examples $p=3,\qquad S=\{3,5,7\}\equiv\{0,2,1\} \pmod{3}$ $p=5,\qquad S=\{5,7,11,19,23\}\equiv\{0,2,1,4,3\} \pmod{5}$ etc. I've verified this for $p\leq 1000$. As usual, I am curious whether this is a known result and/or where to look for related work, and of course any counterexamples should they exist. I'm also curious whether this result seems to others to be surprising, or expected; I'm having a hard time deciding the answer for myself.

论坛问题帖 | has_answers | Trevor | 2020-01-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019051 | MSE-3520446

proof of prime in every interval (p^2,p^2+p)

Overview We'll introduce a sort of little hack called the missing modulo conjecture which can identify an integer's previous prime. We then show that this may not be perfectly reliable on account of how it interacts with all prime squares. In fact, we show that we would need a prime in every $(p^2,p(p+1))$ to be safe: we'd be forced to use them. Flip that around, though, and it means that proving the MMC incidentally guarantees us that prime existence (which we do still need). So we prove it, and since we know it works, it follows that those primes must exist after all. The missing modulo conjecture (MMC) The gist here is that you can take any integer, look at the results you get from taking that integer mod every smaller integer, and determine the nearest prime by noting which small factors could not be generated. For a given $n\in\mathbb N$, let a be the smallest nonnegative integer such that $n \not\equiv a \pmod {m}$, where m can take any value in the interval $[2,n/2]$. I assert that $n-a$ will always be the nearest prime $\leq n$. (See this question for more details.) Prime squares While the MMC mechanism most likely works perfectly, there's one big obstacle preventing its use in a proof. Let p be any prime. Then take $p(p+1) \not\equiv p \pmod{m}$ for $m\in \left[2,\frac{p(p+1)}{2}\right]$. This is the same operation as before aside from fixing p; it erroneously fails to return a factor of p, which is problematic because the typical range of p in these cases could conceivably overlap or interfere with a, and here, p does not point to a previous prime, meaning our a values could also be suspect. When $n=p(p+1)$, either there exists an $a<p$ and life is good, or we must default to $a=p$, since by definition we cannot have $a>p$ for the same n. If $a=p$, then since $n-a=p(p+1)-p=p^2 \not\in\mathbb P$, MMC would fail. For MMC to hold, we must always have our $a<p$. However, this requires $n-a=p(p+1)-a$ be prime. And since $a<p$, this prime would need to be found within $\left(p^2,p(p+1)\right)$. Given this conflict arises for every prime p, if we want the MMC to bear out, we would need a prime to be located in every similar prime square interval. Conversely, if MMC were independently found to always hold, the existence of these primes would necessarily follow. Solution: the MMC proof Let there be some prime gap $q'<\ldots<q$, with $c=q-q'-1$ being the number of composite values in (q',q). We're going to show that MMC works without issue in this instance, and by extension, everywhere else. The lowest missing mod value a on a prime n is always 0, and increments by 1 with each higher n until resetting to 0 at the next higher prime. At $q-1$, we'll be up to $a=c$; this requires we establish that no values $0 \leq v < a$ are possible, where v would be a hypothetical smaller missing mod result that we can't necessarily trust, like the prime square above. $v<a$ iff $q-1 \not\equiv v \pmod {m\in[2,\frac{q-1}{2}]}$. In other words, we're making sure that any v that exists is congruent to every last one of the c possible lower values than a, or more precisely, making sure it would have had to be congruent$-$really, we're ruling out possibilities: if none of those values are missing from the results of our mods, then no v using them could exist. To check that, we establish that $q-k\equiv [0,a-k] \pmod {m}$, meaning that for each $k\leq c$, we want to map it to one of several 0s we're about to find, at least one on each composite in the gap. The lower bound of our range means that any starting-at-0 mod value in the range $[k+1,(q-k)/2]$ won't cycle around to 0 until at least reaching q, so k will be removed from the pool of could-have-been v values. The best part is that because of the huge permissible range, you can simply rely on the greatest proper divisor of $q-k$ to qualify with no actual check or calculation required. Repeat this until all choices for v have been confirmed as observed, and therefore excluded. We'll have demonstrated that there cannot be a $v<a$, not even a bogus one like the prime square. With that addressed, there is no reason the same process cannot be implicitly applied to all prime gaps, which would prove MMC. And that in turn would prove there's always at least one prime in $\left(p^2,p(p+1)\right)$. (example) Consider the prime gap $(13,17)$. Identify three appropriate k_i such that $2\leq \gcd(k_1,16) \leq 8\qquad\qquad 3\leq\gcd(k_2,15) \leq 7\qquad \qquad 4\leq \gcd(k_3,14) \leq 7.$ We can go ahead and use $k_{1,2,3}=\{8,5,7\}$, the greatest proper divisors. These divisors correspond to mod values m which would yield 0, as they divide in cleanly: now, we know where some zeros are. Again, we're not changing anything; we're simply confirming things. Also, now we know that these are the

mod values which are possible to receive within our parameters: $ 14 \mapsto \{0,2,4\} \\ 15 \mapsto \{0,1,3\} \\ 16 \mapsto \{0,1,2,4\} $ Note the conspicuous absences of $\{1,2,3\}$, respectively. This is because those are the values a takes when given each n. And note we have accomplished our goal of confirming the impossibility of 0 for 14, $\{0,1\}$ for 15, and $\{0,1,2\}$ for 16 as potential values for v. Collectively they comprise all values lower than our own $a=\{1,2,3\}$, meaning MMC definitely works for this gap interval. Remember, this is only to illustrate the process, and because finding the divisors is guaranteed, the proof can be automatically applied to all gaps without any further work. I'm looking forward to having someone confirm or point me at the mistake. Much thanks for either. Further updates/improvements to come.

论坛问题帖 | no_answers_at_retrieval | Trevor | 2020-01-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019052 | MSE-3520913

Erdős-Strauss Conjecture: Is there a simpler way to solve?

I'm a huge fan of conjectures, and I'm fascinated by this new one. The Erdős-Strauss conjecture is that $\frac{4}{n} = \frac{1}{a} + \frac{1}{b} + \frac{1}{c}$, where n is greater than or equal to 2 and is a natural number. Or, $\frac{4}{n}$, where n is greater or equal to two and is a natural number, can be written as the sum of three unit fractions. I know that I can eliminate all even numbers, because $\frac{4}{n}$ where n is even can be written as $\frac{1}{0.5n} + \frac{1}{n} + \frac{1}{n} = \frac{2}{n} + \frac{1}{n} + \frac{1}{n} = \frac{4}{n}$. The way I test the odd numbers is I subtract the nearest unit fraction less than $\frac{4}{n}$ and see if the difference can be written as the sum of two unit fractions. But as n keeps getting higher, my strategy doesn't work for long. So, my question is, what's a better strategy?

论坛问题帖 | no_answers_at_retrieval | Mathemagician314 | 2020-01-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019053 | MSE-3521679

A Weak Schönflies Theorem in $\mathbb{R^n}$?

I was reading about the classical topological result of Schönflies that Jordan curve in a plane can be extended homeomorphically onto the whole plane. The well-known counterexample of Alexander's horned sphere tells us that the Schönflies' theorem does not extend into higher dimensions, but can we have a weaker version? Any homeomorphism of the sphere $\mathbb{S^{n-1}}$ onto a simple $(n-1)$-dimensional polygonal surface of $\mathbb{R^n}$ can be extended to a homeomorphism from $\mathbb{R^n}$ onto $\mathbb{R^n}$. EDIT: J. W. Alexander showed in 1920 that any simple $(n-1)$-dimensional polyhedral (polygonal surface) in $\mathbb{R^3}$ can be extended to a homeomorphism from $\mathbb{R^3}$ onto $\mathbb{R^3}$. The general result was proved in 1958 by B. Mazur in his PhD thesis.

论坛问题帖 | has_answers | Hulkster | 2020-01-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019054 | MSE-3527292

The Generalized Bergström’s inequality

Prove the Generalized Bergström’s inequality For all $x_1,x_2,\ldots,x_n>0$ and $r \geq \frac{1}{n}$, it holds that $$\sum_{i=1}^n\frac{x_i}{\sqrt[nr]{x_i^{nr}+(n^{nr}-1)\prod \limits_{j=1}^nx^r_j}} \ge 1.$$ This is a slightly modified version of my earlier question here (which corresponds to the case $r=1$). The case $n=3$ and $r=\frac{2}{3}$ can be reduced to the inequality problem of IMO 2001 (Problem 2).

论坛问题帖 | has_answers | Hulkster | 2020-01-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019055 | MSE-3536226

How do I solve the Continued Fraction $[1,x,x^2,x^3,\ldots]$

My function $M(x)$ is defined as $$M(x)=1+\cfrac{x^0}{1+\cfrac{x^1}{1+\cfrac{x^2}{1+\cfrac{x^3}{1+\cfrac{x^4}{1+\cfrac{x^5}{1+\cfrac{x^{6}}{1+\cfrac{x^{7}}{1+\cfrac{x^{8}}{1+\cfrac{x^9}{1+\cfrac{x^{10}}{1+\ddots}}}}}}}}}}$$ I understand the Rogers-Ramanujan Continued Fraction $R(a,q)$ is very similar but it is slightly different. I would like to know what $M(2)$ is, but I learned that any value $|q|>1$ starts to split and become two different values. What are the decimal approximations of $M(2)$ for both n=even and n=odd and what are their exact values? This is my first question on this site so if it's janky or similar to an answer already I'm sorry.

论坛问题帖 | no_answers_at_retrieval | Jondoolio | 2020-02-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019056 | MSE-3536534

Conjecture about a continued fraction

Conjecture: $$\large 2^{n-1}+\frac{1}{2+\cfrac{1}{2^{n}-1+\cfrac{1}{2+\cfrac{1}{2^{n}-1+\cfrac{1}{2+\ddots}}}}}=\frac{1+\sqrt{3a_n}}{2}\tag*{[1]}$$ such that $a_n=4a_{n-1}+1$ and $a_0=0$.$\quad(n\geqslant 1)$ Ex. If $n=1$, then $a_n=4a_0+1=4\times 0 + 1 = 1$. $$\therefore 1+\frac{1}{2+\cfrac{1}{1+\cfrac{1}{2+\cfrac{1}{1+\cfrac{1}{2+\ddots}}}}}=\frac{1+\sqrt{3}}{2}\tag*{[2]}$$ This can be proven using the formula $x=a+\dfrac{1}{b+\dfrac{1}{x}}=a+\dfrac{1}{b+\dfrac{1}{a+\dfrac{1}{b+\dfrac{1}{a+\ddots}}}}$ Solving for x results in a quadratic equation for which $x=\dfrac12\left\{a+\sqrt{a\left(a+\dfrac4b\right)}\right\}$. Substituting $a=1$ and $b=2$ yields $[2]$ as required. Problem is, I am unsure on how to (dis)prove this conjecture given the recursive sequence involved. How do I appropriately go about this? Any suggestions or counter-examples? Thanks :)

论坛问题帖 | has_answers | Mr Pie | 2020-02-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019057 | MSE-3545934

Conjectured summation inequality

I was playing around with numbers and noticed that the following series is quite close to $\sqrt 2$... but not quite. So I have conjectured that $\sqrt 2$ is arguably the closest highest bound; $$1+\frac 12\left(1-\frac{1}{2^2}\right)+\frac{1\cdot 3}{2\cdot 4^2}\left(\frac{1}{2}-\frac{1}{3^2}\right)+\frac{1\cdot 3\cdot 5}{2\cdot 4\cdot 6^3}\left(\frac{1}{3}-\frac{1}{4^2}\right)+\cdots < \sqrt{2}$$ Can this be proven? Is there a closed form of the infinite series? I... have no idea how to tackle this problem. I've never seen anything like this and have just made it up because I was bored in physics class, but after some iterations, methinks this series is convergent. Are there any series that look like this, particularly for $\sqrt 2$, that may serve a good use of application to measuring the truth of this conjecture? Thanks. :)

论坛问题帖 | has_answers | Mr Pie | 2020-02-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019058 | MSE-3546252

Is this elementary-number-theoretic conjecture already mentioned somewhere or it is entirely new?

While I was doing some research in elementary number theory I discovered some regularities that seem to be very promising. First, function d can be defined as $d(n)=d(\prod_{i=1}^{k(n)} p_i^{r_i})=\sum_{i=1}^{k(n)}r_i$ where $n=\prod_{i=1}^{k(n)} p_i^{r_i}$ is unique prime factorization of n and it can be said that natural number $n>1$ is of degree w if and only if $d(n)=w$. So, for example, prime numbers are natural numbers of first degree. The conjecture I would like to propose is the following: Every prime number ≥7 is the sum of a prime number and number of second degree. As noted in the comments, the conjecture can be rephrased very simply as: Every prime number $p\geq 7$ can be written as $p=qr+s$ where q,r,s are primes.

论坛问题帖 | no_answers_at_retrieval | user750262 | 2020-02-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019059 | MSE-3550096

Minimal counterexample to $\frac{p - 1}{p^2}$-conjecture

There existed once a “folklore” conjecture that stated: Suppose p is a prime. Then any finite group G with $> (1 - \frac{p-1}{p^2})|G|$ elements of order p has exponent p This conjecture was disproved by G.E.Wall in 1965, who constructed a group of exponent 25, where all elements of order 25 lie in the subgroup of index 25. My question is: What is the minimal possible size of a counterexample to this conjecture? I am interested in it because I collect disproven conjectures with large minimal counterexamples Note, that p in this counterexample has to be 5 of greater, as the conjecture was proven to be true for $p = 2$ and $p = 3$ by Thomas Laffey in 1976.

论坛问题帖 | no_answers_at_retrieval | Chain Markov | 2020-02-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019060 | MSE-3555736

Formula for Feigenbaum’s constant?

I have conjectured a formula to calculate the Feigenbaum constant $\delta \approx 4.66920$. $\delta\stackrel{?}{=}$ $$4+\cfrac{1\times 2 -1}{1+\cfrac{2\times 3 -1}{2^2+\cfrac{3\times 4 -1}{1+\cfrac{4\times 5 -1}{3^2+\cfrac{5\times 6 -1}{1+\ddots}}}}}$$ which after several iterations is ≈ 4.66919 (I believe). This discovery, whether exact or approximate, was completely accidental. Can this be numerically verified? Thanks.

论坛问题帖 | has_answers | Mr Pie | 2020-02-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019061 | MSE-3556952

Prove that $\sum_{k=2}^{n}\{\sqrt{a_{k-1}}+\sqrt{a_{k}}\}^{-1}=\frac{n-1}{\sqrt{a_{1}}+\sqrt{a_{n}}}$

If $a_{1}, a_{2}, a_{3}, \dots, a_{n}$ are consecutive terms of an arithmetic progression (None of which are 0) , prove that $\sum_{k=2}^{n}\{\sqrt{a_{k-1}}+\sqrt{a_{k}}\}^{-1}=\frac{n-1}{\sqrt{a_{1}}+\sqrt{a_{n}}}$ I started out doing this problem by approaching it using mathematical induction. If my calculations are right, it must be proved through induction that $\frac{k-1}{\sqrt a_{1}+\sqrt{a_{n}}}+\frac{1}{\sqrt{a_{k}}+\sqrt{a_{k-1}}}=\frac{k}{\sqrt{a_{1}}+\sqrt{a_{k-1}}}$ Trying to simplify this gives me a very long expression. I assume that we have to use the fact that the terms are in AP. Any help is appreciated.

论坛问题帖 | has_answers | NoLand'sMan | 2020-02-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019062 | MSE-3558182

On conjectured continued fractions and e

Playing around with numbers, I conjectured three incredibly interesting things: $$9+\cfrac{1}{18+0\times 12\cfrac{1}{18+1\times 12+\cfrac{1}{18+2\times 12+\cfrac{1}{18+3\times 12+\ddots}}}}=\frac{4e^{1/3}-2}{e^{1/3}-1}$$ $$6+\cfrac{1}{9+0\times 6+\cfrac{1}{9+1\times 6+\cfrac{1}{9+2\times 6+\cfrac{1}{9+3\times 6+\ddots}}}}=\frac{4e^{2/3}-2}{e^{2/3}-1}$$ $$5+\cfrac{1}{6+0\times 4+\cfrac{1}{6+1\times 4+\cfrac{1}{6+2\times 4+\cfrac{1}{6+3\times 4+\ddots}}}}=\frac{4e-2}{e-1}$$ So, what in God's name is going on behind the scenes? Why does this seem to be true, why does it involve e, so many questions! All I did was play around on a calculator with some continued fractions, went to Wolfram Alpha and asked for the result to be written in terms of e and then I noticed some patterns. But what is really going on? Well, apart from a bit of luck, I have no idea. Any ideas? Thanks. Edit: This question may be of help, as it reveals general continued fractions regarding the hyperbolic tangent, which to those who don't know, is a function with respect to some value α defined as $\tanh(\alpha):=\frac{e^{2\alpha}-1}{e^{2\alpha}+1}.$

论坛问题帖 | has_answers | Mr Pie | 2020-02-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019063 | MSE-3560689

On expressing $\frac{\pi^n}{4\cdot 3^{n-1}}$ as a continued fraction.

It is a celebrated equation that $$\frac{\pi}{4}=\cfrac{1}{1+\cfrac{1^2}{3+\cfrac{2^2}{5+\cfrac{3^2}{7+\ddots}}}}$$ However, there are two other conjectured equations that I found which, if true (they seem to be), might reveal a pattern. $$\frac{\pi^2}{12}=\cfrac{1}{1+\cfrac{1^4}{3+\cfrac{2^4}{5+\cfrac{3^4}{7+\ddots}}}}$$ $$\frac{\pi^3}{36}=\cfrac{1}{1+\cfrac{1^6}{3+\cfrac{2^6}{5+\cfrac{3^6}{7+\ddots}}}}$$ Conjectured General Formula: For natural $n\geqslant 1$, $$\frac{\pi^n}{4\cdot 3^{n-1}}=\cfrac{1}{1+\cfrac{1^{2n}}{3+\cfrac{2^{2n}}{5+\cfrac{3^{2n}}{7+\ddots}}}}$$ Can these be numerically verified? I have not the skill to by-hand prove/disprove these, and have only been using Wolfram Alpha to arrive at these conjectures. It would also be much appreciated if one could suggest a program I could install in order to evaluate these continued fractions independently, as well as the code required. Will PARI/GP suffice? Thanks.

论坛问题帖 | has_answers | Mr Pie | 2020-02-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019064 | MSE-3583435

Interesting patterns related to the sums of the remainders of integers

Let us define, $$r(b)=\sum_{k=1}^{\lfloor \frac{b-1}{2} \rfloor} (b \bmod{k})$$ After reading the posts Surprising fact about a certain number-theoretic function and Do primes have special sums... I decided to play around with the $r(b)$ function and after performing some computations using PARI/GP, I found some very interesting patterns. $r(a^n)-r(a^n-1)=0$ when $a\in {2, 3}$ for any $n\in \Bbb{N}$. (Proved by user Haran in the first post) For fixed a and any $n\in \Bbb{N}$, $r(a^n)-r(a^n-1)$ is either always positive or always negative when a is not of the form $2p$ or $3q$ where p,q are prime and $p\gt 3, q\gt 7$. For fixed a, $r(a^n)-r(a^n-1)$ is initially postive and then becomes negative as $n\in \Bbb{N}$ increases when a is the form of $2p$ or $3q$ where p,q are prime and $p\gt 3, q\gt 7$. If $a \notin \{2p, 3p\}$ is prime or semiprime then $r(a^n)-r(a^n-1)$ is positive for all $n\in \Bbb{N}$. (edit) What is the reason for these weird characteristics or $r(a^n)-r(a^n-1)$? I would really appreciate if someone could prove/disprove any of the last 3 observations. Here is some of the initial data from where I made my observations. The first column gives the a values, the second gives the n values, and the third gives the corresponding values of $r(a^n)-r(a^n-1)$. 1 1 0 1 2 0 1 3 0 1 4 0 1 5 0 2 1 0 2 2 0 2 3 0 2 4 0 2 5 0 3 1 0 3 2 0 3 3 0 3 4 0 3 5 0 4 1 0 4 2 0 4 3 0 4 4 0 4 5 0 5 1 1 5 2 6 5 3 31 5 4 156 5 5 781 6 1 -1 6 2 -20 6 3 -169 6 4 -1160 6 5 -7381 7 1 2 7 2 16 7 3 114 7 4 800 7 5 5602 8 1 0 8 2 0 8 3 0 8 4 0 8 5 0 9 1 0 9 2 0 9 3 0 9 4 0 9 5 0 10 1 1 10 2 -18 10 3 -341 10 4 -4212 10 5 -46079 11 1 4 11 2 48 11 3 532 11 4 5856 11 5 64420 12 1 -5 12 2 -116 12 3 -1625 12 4 -20360 12 5 -247445 13 1 5 13 2 70 13 3 915 13 4 11900 13 5 154705 14 1 3 14 2 -8 14 3 -513 14 4 -10000 14 5 -159657 15 1 -2 15 2 -66 15 3 -1178 15 4 -18564 15 5 -282722 16 1 0 16 2 0 16 3 0 16 4 0 16 5 0 17 1 7 17 2 126 17 3 2149 17 4 36540 17 5 621187 18 1 -4 18 2 -200 18 3 -4732 18 4 -95120 18 5 -1800964 19 1 8 19 2 160 19 3 3048 19 4 57920 19 5 1100488 20 1 -3 20 2 -162 20 3 -3813 20 4 -79092 20 5 -1595583 21 1 -1 21 2 -80 21 3 -2109 21 4 -47200 21 5 -1011161 22 1 7 22 2 36 22 3 -665 22 4 -30744 22 5 -853565 23 1 10 23 2 240 23 3 5530 23 4 127200 23 5 2925610 24 1 -13 24 2 -500 24 3 -13273 24 4 -327560 24 5 -7929493 25 1 6 25 2 156 25 3 3906 25 4 97656 25 5 2441406 26 1 9 26 2 70 26 3 -549 26 4 -45220 26 5 -1577991 27 1 0 27 2 0 27 3 0 27 4 0 27 5 0 28 1 -1 28 2 -200 28 3 -6897 28 4 -202000 28 5 -5716841 29 1 13 29 2 390 29 3 11323 29 4 328380 29 5 9523033 30 1 -13 30 2 -1022 30 3 -39601 30 4 -1309532 30 5 -40972393 31 1 14 31 2 448 31 3 13902 31 4 430976 31 5 13360270 32 1 0 32 2 0 32 3 0 32 4 0 32 5 0 33 1 1 33 2 -96 33 3 -4655 33 4 -169824 33 5 -5781695

论坛问题帖 | has_answers | Mathphile | 2020-03-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019065 | MSE-3587066

For fixed hypotenuse, can the number of primitive Pythagorean triples exceed the number of non-primitive ones?

For the equation, $$a^2+b^2=c^2$$ if c is fixed and the number of natural solutions for a, b is greater than 1, then can the number of primitive solutions (solutions in which a, b, c are coprime) exceed the number of non-primitive ones? After testing a large number of cases I believe that the number of non-primitive solutions will always exceed the number of primitive ones although I have no proof for this. If false, what is the smallest counter-example for c?

论坛问题帖 | has_answers | Mathphile | 2020-03-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019066 | MSE-3587292

Interesting property related to the sums of the remainders of integers

Cross-posted on Math Overflow too Let us define, $$r(b)=\sum_{k=1}^{\lfloor \frac{b-1}{2} \rfloor} (b \bmod{k})$$ After playing around with the $r(b)$ function for sometime I noticed that $r(b)$ appreared to be more even than odd. So to see the difference between the number of even and odd terms of $r(b)$, I defined a function, $$z(x)=\sum_{n=1}^x(-1)^{r(n)}$$ When user Peter ran a program for computing values of $z(x)$ in PARI, I observed that for $x\le 10^{10}$, $z(x)\gt 0$. This suggests that there are always more even terms of $r(n)$ than odd terms for any x. This leads to my two questions: Is $z(x)$ always positive? If so, then how do we prove this? Is $|z(x)|$ bounded by some maximum value? If so, then what is this maximum value? Till now the maximum value of $|z(x)|$ found was 49 for $x = 5424027859$. I find it odd that $|z(x)|$ goes to these large values and then returns back to small values as small as 1. Edit: With the help of user Vepir I was able to plot $z(x)$ and noticed that the function has a sinusoidal-fractal-like appearance and it seems to grow without bounds, although very very slowly. Is there any reason for this sinusoidal and fractal nature?

论坛问题帖 | has_answers | Mathphile | 2020-03-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019067 | MSE-3587363

For any $k \gt 3$, if $n!+k$ is a perfect power then does there exist any $n\gt k$?

A while ago, I asked a similar question For any $k \gt 1$, if $n!+k$ is a square then will $n \le k$ always be true? where users mathworker21 and WE Tutorial School proved that for non-square k, $n\le k$ is always true when $n!+k$ is sqaure. Recently I got the idea to check if the property is true for all perfect powers and using PARI, I observed that if, $$n!+k=a^b$$ where $n, k, a, b\in \Bbb{N}$, $k\gt 3$ and $b\ge 2$, then $n\le k$ is always true. In search for a counter-example, I covered a range of $k\le 2500$ and $n\le 10^4$ for each k and found none. So my question is, If true, can we prove/partially prove that $n\le k$ always holds? If false, then what is the smallest counter-example where $n\gt k$?

论坛问题帖 | no_answers_at_retrieval | Mathphile | 2020-03-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019068 | MSE-3589121

Pattern in Number of Conjugacy Classes of p-groups

I was playing around with the number of conjugacy classes of p-groups in GAP and made the following conjecture: If there is a group of order p^{2n} with k conjugacy classes then there is a group of order p^{2n+1} with $k+p-1$ conjugacy classes. The only exception is $p=2$, $n=2$, $k=7$. For example, there are groups of order 64 with $\{13,16,19,22,25,28,34,40,64\}$ many conjugacy classes so there are groups of order 128 with $\{14,17,20,23,26,29,35,41,65\}$ many conjugacy classes. Can someone prove this or find a counterexample? For reference, here is the data that I was using. [1] [2] [4] [5.8] [7.10,16] [11,14,17,20,32] [13,16,19,22,25,28,34,40,64] [14,17,20,23,26,29,32,35,38,41,44,50,56,65,68,80,128] [19,22,25,28,31,34,37,40,43,46,49,52,55,58,61,64,67,70,73,76,82,85,88,100,112,130,136,160,256] [1] [3] [9] [11,27] [17,33,81] [19,35,51,83,99,243] [41,57,73,89,105,121,153,249,297,729] [43,59,75,91,107,123,139,155,171,187,219,251,267,299,315,363,459,731,747,891,2187] [1] [5] [25] [29,125] [49,145,625] [53,149,245,629,725,3125] [73,169,265,361,649,745,841,1225,3145,3625,15625] [1] [7] [49] [55,343] [97,385,2401] [103,391,679,2407,2695,16807] n:=16; l:=[]; for i in [1..NrSmallGroups(n)] do G:=SmallGroup(n,i); if not \in(NrConjugacyClasses(G),l) then Add(l,NrConjugacyClasses(G)); Sort(l); fi; od; Print(l); One well-known pattern in the above data is a result of Hall that if $n=2k+e$ (with $e=0,1$) then $k=p^e+(p^2-1)(n+(p-1)t)$ for some $t\geq0$. This explains why the above lists jump by $(2-1)(2^2-1)=3$ when $p=2$, by $(3-1)(3^2-1)=16$ when $p=3$, by $(5-1)(5^2-1)=96$ when $p=5$, and by $(7-1)(7^2-1)=288$ when $p=7$.

论坛问题帖 | no_answers_at_retrieval | Thomas Browning | 2020-03-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019069 | MSE-3591424

Conjecture $\frac{a}{a^r+b^r}+\frac{b}{b^r+c^r}+\frac{c}{c^r+a^r}\geq \frac{a}{a^r+c^r}+\frac{c}{c^r+b^r}+\frac{b}{b^r+a^r}$

following this kind of inequality One of my old inequality (very sharp) I propose this because I don't see it on the forum : Let $a,b,c>0$ and $a+b+c=1$ with $r\in(\frac{1}{2},1)$ and $a\geq b \geq c$ then we have : $$\frac{a}{a^r+b^r}+\frac{b}{b^r+c^r}+\frac{c}{c^r+a^r}\geq \frac{a}{a^r+c^r}+\frac{c}{c^r+b^r}+\frac{b}{b^r+a^r}$$ First of all it's a conjecture where I don't find counter-examples . Secondly when $r\in(0,\frac{1}{2})$ the inequality is reversed .I use Pari-gp for that .Furthermore (if it's true) I think it's really not new so I add the tag reference request.We have an equality case when $r=0.5$ whenever $a,b,c>0$. So if you have idea to prove it or disprove it... Thanks a lot .

论坛问题帖 | has_answers | Barackouda | 2020-03-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019070 | MSE-3598714

Expectation of arbitrary product of Brownian motions.

Let (B_t) be a standard Brownian motion, $n\in\mathbb{N}$ even and $t_1,\ldots,t_n\in\mathbb{N}$. I was wondering whether there is a general formula for $$\mathbb{E} \left[B(t_1)\cdots B(t_n)\right].$$ By successively applying the property of independent increments, it is straightforward to calculate this expression for some fixed $n\in\mathbb{N}$. Now my conjecture is the following. Let $\mathcal{P}_n$ be the set of partitions of $\{1,\ldots,n\}$ into two-element subsets. Then $$\mathbb{E} \left[B(t_1)\cdots B(t_n)\right] \stackrel{?}{=} \sum_{P\in\mathcal{P_n}} \prod_{\{i_1,i_2\}\in P} t_{i_1} \wedge t_{i_2}.$$ I confirmed this by hand for n up to 6 and tried to prove it by induction , but failed. I would be grateful for any advice on how to prove this or a known formula for $\mathbb{E} \left[B(t_1)\cdots B(t_n)\right].$ For illustration, this is the formula for $n=4$: There are three partitions of $\{1,2,3,4\}$, namely $\{\{1,2\},\{3,4\}\}$, $\{\{1,3\},\{2,4\}\}$ and $\{\{1,4\},\{2,3\}\}$. Their contributions to the sum are $t_2 t_4$, $t_3 t_4$ and $t_4 t_3$ respectively (assuming $t_1>t_2>t_3>t_4$), yielding the correct result $2t_3 t_4 + t_2 t_4$.

论坛问题帖 | no_answers_at_retrieval | Peter Koepernik | 2020-03-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019071 | MSE-3619244

Is the last digit of $2^{2^{n-1}(2^n-1)}-1$ always 5 for $n >3$?

Is the last digit of $2^{2^{n-1}(2^n-1)}-1$ always 5 for $n >3$? I did modification to the Mersenne numbers (Even perfect numbers) foruma I put that formual to be the power of 2 have got : $2^{2^{n-1}(2^n-1)}-1$, Now for some values of $n >3$ that i have tried i have got the produced integer with last digit equal 5 like this example, Now to do more n or so large to conjecture that it is beyond of current tecknology , Really am interesting to prove that if it is true ?

论坛问题帖 | has_answers | zeraoulia rafik | 2020-04-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019072 | MSE-3620230

Conjectured continued fraction formula for Catalan's constant

Yesterday I posted this conjecture, but then deleted it thinking it was false. Turns out Python doesn't define a^b as a^b, but rather as a**b. Conjecture: Denote by G Catalan's constant, then $$G=\cfrac{1}{1+\cfrac{1^4}{8+\cfrac{3^4}{16+\cfrac{5^4}{24+\cfrac{7^4}{32+\cfrac{9^4}{40+\ddots}}}}}}$$ Given the connection G has with the number 8 shown here, as well as this continued fraction reaching nearly the first five decimal places of G after around 200 iterations (vinculums), I am confident this is true. However, I do not know how to code a continued fraction on Python or Pari/GP (a friend of mine gave it a go, but also to no avail) up to an iteration n without having to write it out manually, which is really tedious. Here is some python code from a friend, coding this fraction up to 12 iterations to be ≈ 0.9151, reaching the first three decimal places of G. The only 'local' behaviour that I can say about continued fractions is that most of them are convergent, and that they all converge via oscillation at each iteration. But, more importantly, I'd like to know that if this be true, can it be shown from here that G is irrational (or even transcendental, if you are willing)? I am aware this is an unsolved problem, which was what inspired me to write G in another closed form. Any thoughts? Thank you in advance.

论坛问题帖 | has_answers | Mr Pie | 2020-04-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019073 | MSE-3625652

Attempt to get a characterization of even perfect numbers from an equation involving the Dedekind psi function

In this post we denote the Dedekind psi function as $\psi(n)$ for integers $n\geq 1$. This is an important arithmetic fuction in several subjects of mathematics. As reference I add the Wikipedia Dedekind psi function. An even perfect number is an even integer $n\geq 1$ satisfying that $\sigma(n)=2n$, where $\sigma(n)=\sum_{1\leq d\mid n}d$ is the sum of divisors function. Claim. Invoking Euclid-Euler theorem it is easy to prove that each even perfect number satisfies $$4\psi(n)^2-(12n+3)\psi(n)=-9n^2,\tag{1}$$ alternatively $$\psi(n)=\frac{12n+3+3\sqrt{1+8n}}{8}.\tag{2}$$ Question. I would like to know if it is possible to get a characterization for even perfect numbers from (1): what I evoke is if we can to prove that if an integer $n\geq 1$ satisfies (1) then n is an even perfect number. Many thanks. The case $n=1$ is easy. By contradiction also is easy to rule out/discard that n is a power of two and that n is odd. Thus the only discussion is what happens with numbers of the form $$n=2^{\alpha}m\tag{3}$$ where $\alpha\geq 1$ is integer and $(2,m)=1$ for integers $m>1$. Our case (3) leads to $$2^{\alpha+1}m\prod_{\substack{\text{primes }p\\p\mid m}}(p+1)^2-(2^{\alpha+2}m+1)\prod_{\substack{\text{primes }p\\p\mid m}}p(p+1)=2^{\alpha+1}m \prod_{\substack{\text{primes }p\\p\mid m}}p^2,\tag{4}$$ and if I was right from (4) on assumption that $m=p$ is a prime (that is we assume that the number of distinct primes dividing m is $\omega(m)=1$) I get that $p+1=\lambda 2^{\alpha+1}$ with $\lambda=1$ using the congruence $\prod_{\substack{\text{primes }p\\p\mid m}}(p+1)\equiv 0\text{ mod }2^{\alpha+1}$. Thus if my argument works I need to rule out/discard the case of integers m with more than 1 distinct primes factors in their prime factorization (I want to remove the case $\omega(m)>1$).

论坛问题帖 | has_answers | user759001 | 2020-04-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019074 | MSE-3626667

If $q^k n^2$ is an odd perfect number with special prime q, then $k \neq 1$ (MSE version)

Note that $\gcd(\sigma(q^k),\sigma(n^2))=i(q)=\gcd(n^2,\sigma(n^2))$ if and only if $\gcd(n,\sigma(n^2))=\gcd(n^2,\sigma(n^2))$. I have therefore undeleted this question, in view of this closely related MSE question: When does $\gcd(m,\sigma(m^2))$ equal $\gcd(m^2,\sigma(m^2))$? What are the exceptions?. (Cross-posting this question from MO, as I think it is about to be closed there, due to some reasons yet unknown to myself.) This question is directly related to this earlier one ("On odd perfect numbers and a GCD - Part II") in MSE. That question has all the details. Now, let $N = q^k n^2$ be an odd perfect number with special prime q satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. Descartes, Frenicle, and subsequently Sorli conjectured that $k=1$ always holds. In the hyperlinked MSE question, we have the equation $$\gcd(\sigma(q^k),\sigma(n^2))=i(q)=\gcd(n^2,\sigma(n^2))$$ where $$i(q)=\frac{\sigma(n^2)}{q^k}=\frac{2n^2}{\sigma(q^k)}.$$ Assume to the contrary that $k=1$. By Brown (2016), it follows that $q < n$, whence it follows that $\sigma(q^k) = \sigma(q) = q + 1 < n$. But $i(q) = \gcd(\sigma(q^k),\sigma(n^2))$; in particular, this means that $$i(q) \mid \sigma(q^k) \implies \frac{2n^2}{\sigma(q^k)} = \frac{\sigma(n^2)}{q^k} \leq \sigma(q^k).$$ We know that $\sigma(q^k) < n$, so we have $$2n < \frac{2n^2}{\sigma(q^k)} = \frac{\sigma(n^2)}{q^k} \leq \sigma(q^k) < n,$$ resulting in a contradiction. We therefore conclude that $k \neq 1$.

论坛问题帖 | no_answers_at_retrieval | Jose Arnaldo Bebita Dris | 2020-04-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019075 | MSE-3627784

Does the fraction of distinct substrings in prefixes of the Thue-Morse sequence of length 2^n tend to $73/96$?

Recall that the Thue-Morse sequence$^{[1]}$$\!^{[2]}$$\!^{[3]}$ is an infinite binary sequence that begins with $\,t_0 = 0,$ and whose each prefix p_n of length 2^n is immediately followed by its bitwise complement (i.e. obtained by flipping $0\to1$ and $1\to0$): $$ \begin{array}{c|cc}&t_0&t_1&t_2&t_3&t_4&t_5&t_6&t_7&\!\!\!\dots\\\hline p_0&0\\ p_1&0&\color{red}1\\ p_2&0&1&\color{red}1&\color{red}0\\ p_3&0&1&1&0&\color{red}1&\color{red}0&\color{red}0&\color{red}1\\ \cdots&\cdots\!\! \end{array} $$ We are interested in contiguous substrings of these prefixes. For a string $\mathcal{S}$ of length ℓ, the total number of its substrings, including the empty substring $\langle\unicode{x202f}\rangle$ and the string $\mathcal{S}$ itself, is

$(\ell^2+\ell+2)/2.$ Hence, the total number of substrings in p_n is $(4^n+2^n+2)/2.$ Clearly, not all of those substrings are distinct for $n>1$. For example, $p_2 = \langle0\,1\,1\,0\rangle$ has 11 substrings in total, but only 9 distinct substrings: $$ \begin{array}{|cc}&\langle\!\!\!&0&\color{#808080}1&\color{#b8b8b8}1&\color{#c8c8c8}0&\!\!\!\rangle\\\hline 1&\langle\!\!\!&&&&\!\!\!\rangle\\\hdashline 2&\langle\!\!\!&0&&&\!\!\!\rangle\\ &\langle\!\!\!&&&&\color{#c8c8c8}0&\!\!\!\rangle\\\hdashline 3&\langle\!\!\!&&\color{#808080}1&&&\!\!\!\rangle\\ &\langle\!\!\!&&&\color{#b8b8b8}1&&\!\!\!\rangle\\\hdashline 4&\langle\!\!\!&0&\color{#808080}1&&&\!\!\!\rangle\\ 5&\langle\!\!\!&&\color{#808080}1&\color{#b8b8b8}1&&\!\!\!\rangle\\ 6&\langle\!\!\!&&&\color{#b8b8b8}1&\color{#c8c8c8}0&\!\!\!\rangle\\ 7&\langle\!\!\!&0&\color{#808080}1&\color{#b8b8b8}1&&\!\!\!\rangle\\ 8&\langle\!\!\!&&\color{#808080}1&\color{#b8b8b8}1&\color{#c8c8c8}0&\!\!\!\rangle\\ 9&\langle\!\!\!&0&\color{#808080}1&\color{#b8b8b8}1&\color{#c8c8c8}0&\!\!\!\rangle \end{array} $$ Among these, $\langle0\rangle$ and $\langle1\rangle$ appear in p_2 twice, so the fraction of distinct substrings in p_2 is $\,\stackrel9{}\!\!\unicode{x2215}_{\!\unicode{x202f}11}\!.$ Can we find a simple general formula for $\mathscr D_n$, the number of distinct substrings in p_n? Let's try to compute a few terms: $$2,\,4,\,9,\,28,\,101,\,393,\,1561,\,6233,\,24921,\,99673,\,398681,\,1594713,\,6378841,\,\dots$$ These few terms can be computed by a brute-force approach, but using Coolwater's program from here we can compute hundreds of thousands more. It is not too difficult to discover that for $n>2$ all known terms match a simple formula: $$\mathscr D_n\stackrel{\color{#d0d0d0}?}=\frac{73\cdot4^n+704}{192}\color{#d0d0d0}{\,\,\text{for}\,\,n>2}\tag{$\diamond$}$$ Somewhat oddly, the three initial terms $\mathscr D_0=2,\,\mathscr D_1=4,$ and $\mathscr D_2=9$ do not match the general formula $(\diamond)$, which results in non-integer rational values for these indexes. I conjecture that the general formula $(\diamond)$ is valid for all $n>2$. $$\bbox[LemonChiffon]{\begin{array}{c} \\ \hspace{1in}\text{Could you suggest a way to prove this conjecture?}\hspace{1in}\\ \vphantom. \end{array}}$$ If the conjecture turns out to be true, then we have a remarkable corollary that for $n\to\infty$ the fraction of distinct substrings in the prefixes p_n tends to a quite surprising limit: $$\mathscr L=\lim_{n\to\infty}\frac{73\cdot4^n+704}{192}{\large/}\frac{4^n+2^n+2}2=\frac{73}{96}.\tag{$\small\spadesuit$}$$

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2020-04-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019076 | MSE-3627916

Prove that $1=\sqrt{2\cdot \color{red}{-1} + \sqrt{2\cdot \color{red}2 + \sqrt{2\cdot \color{red}{9}+\sqrt{2\cdot\color{red}{20}+\cdot \cdots}}}}$

I was playing around with square roots and I noticed that the number 1 can be seemingly expressed as an infinite nested radical with an easy pattern. I then noticed that if this is true, this would mean that each nested radical inside is equal to the next odd number, respectively. I have tried to denote this using LaTeX but it did not display as I thought it might, though I hope you can understand what I am trying to concisely say. Conjecture: $$\begin{align}\color{green}1 &= \sqrt{2\cdot \color{red}{-1} + \color{green}{3}} \\ &= \sqrt{2\cdot \color{red}{-1} + \sqrt{2\cdot \color{red}2 + \color{green}{5}}} \\ &= \sqrt{2\cdot \color{red}{-1} + \sqrt{2\cdot \color{red}2 + \sqrt{2\cdot \color{red}{9}+\color{green}7}}}\\ &= \sqrt{2\cdot \color{red}{-1} + \sqrt{2\cdot \color{red}2 + \sqrt{2\cdot \color{red}{9}+\sqrt{2\cdot\color{red}{20}+\color{green}9}}}}\\ &= \sqrt{2\cdot \color{red}{-1} + \sqrt{2\cdot \color{red}2 + \sqrt{2\cdot \color{red}{9}+\sqrt{2\cdot\color{red}{20}+\sqrt{2\cdot \color{red}{35}+\cdots}}}}}\end{align}$$ $\color{red}{-1}+(4\cdot 0+3)=\color{red}{2}$ (radical $=\color{green}1$) $\color{red}2+(4\cdot 1+3)=\color{red}9$ (radical $=\color{green}3$) $\color{red}9+(4\cdot 2+3)=\color{red}{20}$ (radical $=\color{green}5$) $\color{red}{20}+(4\cdot 3+3)=\color{red}{35}$ (radical $=\color{green}7$) $\cdots = \cdots$ Can this be proven? If it is a result of a generalised identity for odd numbers $2n+1$ (for some n, this case being $n=0$), please let me know. Thanks. Edit: @TheSimpliFire kindly gave the recurrence relation: $1=2a_0+b_n$ where $b_n=\sqrt{2a_n+b_{n+1}}$ and $a_n+4n+3=a_{n+1}$ with $a_0=-1$ and $n>0$.

论坛问题帖 | has_answers | Mr Pie | 2020-04-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019077 | MSE-3631561

Conjecture on a functional equality and nested radical of Ramanujan :$3=\sqrt{1+f(2)\sqrt{1+f(3)\sqrt{1+\cdots}}}$

Hi i was wondering something about the great Ramanujan : I think moreover I'm not the only one who propose this kind of problem (so if you have a link related to this subject). We have : $$3=\sqrt{1+2\sqrt{1+3\sqrt{1+\cdots}}}$$ Now the problem : Let $f(x)$ be a positive,continuous and differentiable on function$(0,\infty)$ and non-constant then the functional equation : $$3=\sqrt{1+f(2)\sqrt{1+f(3)\sqrt{1+\cdots}}}$$ Have as unique solution $f(x)=x$ I have tried to build a counter-example without success .First it seems obvious for strictly increasing\decreasing functions such that $f(x)>x$ or $f(x)<x$. I would like to create an counter-example of the form : $$f(x)=x+g(x)$$ Where $g(x)$ is a periodic function . I have tried more general representation without success . If it's true we can see how Ramanujan was great . Any helps are highly appreciated . Thanks a lot .

论坛问题帖 | has_answers | Barackouda | 2020-04-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019078 | MSE-3649763

Conjectured series representation of elliptic integral powers $K(k)^n$

This is a very simple question which has probably already been answered; I apologize if that is the case. We know that the following equation holds: $$K(k)=\frac{\pi}{2}\sum\limits_{n=0}^\infty a_n^{\;2}k^{2n}\tag{1}$$ where $a_n=\frac{(2n-1)!!}{(2n)!!}=\frac{(2n)!}{4^n(n!)^2}$. Additionally, we have that: $$K(k)^2=\frac{\pi^2}{4}\sum_{n=0}^\infty a_n^{\;3}(2kk')^{2n}\tag{2}$$ ($k'=\sqrt{1-k^2}$). Generalization to $K(k)^n$ seems a natural possibility. To slightly

support this, I have in the past formally derived (part of the derivation here although identity not explicitly stated) the following identity: $$\sum_{n=0}^{\infty}\frac{(2n)!^m}{(n!)^{2m}}x^n\\=\left(\frac{2}{\pi}\right)^{m-1}\int_0^\frac{\pi}{2}...\int_0^\frac{\pi}{2}\frac{1}{\sqrt{1-x\left(2^m\prod\limits_{k=1}^{m-1}\cos\theta_k\right)^2}}\;d\theta_1...d\theta_{m-1}\tag{3}$$ (please correct me if this formula is wrong); this is an $m-2$-tuple integral of an elliptic integral. A special case of this is: $$\sum_{n=0}^{\infty}\frac{(2n)!^3}{(n!)^{6}}x^n=\frac{4}{\pi^2}\int_0^\frac{\pi}{2}\int_0^\frac{\pi}{2}\frac{1}{\sqrt{1-64x\cos^2\theta\cos^2\varphi}}\;d\theta\;d\varphi\tag{4}$$ By comparing (2) and (4), we have: $$\int_0^\frac{\pi}{2}\int_0^\frac{\pi}{2}\frac{1}{\sqrt{1-64(kk')^2\cos^2\theta\cos^2\varphi}}\;d\theta\;d\varphi=K(k)^2\tag{5}$$ The symmetry of (5) makes me guess that the integrand can be split into a product of functions depending only on θ and φ and turned into a product of two elliptic integrals (although for some reason I cannot see how to do it and Wolfram Alpha won't check (5) for me). If this is indeed the case it makes me wonder whether the RHS of (3) may also be written as the product of $m-1$ complete elliptic functions $K(k)$ for some $k(x)$. In (2), $(kk')^2=k^2-k^4$, so I wonder in particular whether the following conjecture is true: $\forall m\in\mathbb{N}$ there exists a polynomial $\Pi_m(k)$ of order m^2 such that:
$$K(k)^m=\left(\frac{\pi}{2}\right)^m\sum_{n=0}^\infty a_n^{\;m+1}\Pi_m(k)^n\tag{*}$$ Question: Is $(*)$ true? If so, what is $\Pi_m(k)$? If not where does the pattern break down (if I am right in thinking that there is a pattern)?

论坛问题帖 | no_answers_at_retrieval | Anon | 2020-04-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019079 | MSE-3654530

Conjecture: every set of n equally spaced integers $\leq n^2$ contains a prime.

My claim: Any set of exactly n integers in the range $[1,n^2]$ which are an arithmetic sequence and share no common factor will contain at least one prime. Let $n\geq 2$ be a natural number, and let $a,b\in\mathbb{N}$ where $1\leq a \leq n$ and $1 \leq b \leq n^2$, such that $an+b\leq n^2$. Then the set $\{ax+b \mid 1 \leq x \leq n\}$ will contain between 1 and n primes, inclusive. There are three exceptions: $(n=12,a=1,b=113),(n=12,a=1,b=114),(n=13,a=1,b=113)$, but I strongly suspect those are the only ones. (You could eliminate them by requiring $a > 1$, but I don't think it's worth the cost in generality.) In a nutshell, this conjecture claims that given an $n \times n$ square, any line you draw (that intersects the integers appropriately) must contain a prime, including for starters all rows and columns-something I called the Rook Conjecture, but have since seen referred to as the Calendar Conjecture. I've empirically tested this out to $n=150$ or so, and the trend is strong enough that I'm almost sure it's correct. Of course, finding a proof is another story. I'm looking for feedback. Good answers would be: counterexamples indicating that this is previously known and/or conjectured providing reasons why this does or does not seem likely to be true any other insights

论坛问题帖 | no_answers_at_retrieval | Trevor | 2020-05-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019080 | MSE-3656615

Is the following an existing conjecture or a conjecture at all?

the following floated to my mind today, can you verify if it stands to be true, or is a pre-existing conjecture. If not, can you correct me? And if it is one, can you prove it? Prime factorisation of any perfect square P of the form $$P=a_1^{x_1}\times a_2^{x_2}\times a_3^{x_3}\times.....\times a_n^{x_n} $$ where $$x_1,x_2≠0$$ which implies that atleast two non unity factors exist,then $$a_1^{x_1}+ a_2^{x_2}+ a_3^{x_3}+.....+ a_n^{x_n}=Z$$ where Z is perfect square, is only applicable for $P=144$. Example: for $P=144$, prime factorisation is as follows: $$2^4\times 3^2$$ and $$2^4+ 3^2=25=5^2$$

论坛问题帖 | has_answers | Anmoldeep | 2020-05-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019081 | MSE-3659875

A pattern with third coefficients of sums of powers.

So I have heard countless times of Bernoulli Numbers and its relation to the sums of powers. Inspired by this (as well as Chebyshev polynomials), I decided to look at the sums of powers myself and find patterns in the coefficients. Let $$S_{n,k}=1^k+2^k+\cdots + n^k$$ then, thanks to wikipedia, these are the first five formulae. However, I will highlight the following. $$\begin{align}S_{n,1}&=\frac 12n^2 + \frac 12n\\S_{n,2}&=\frac 13n^3+\frac 12n^2+\color{red}{\frac 16}n\\ S_{n,3}&=\frac 14n^4+\frac 12n^3+\color{red}{\frac 14}n^2+0n\\ S_{n,4}&= \frac 15n^5+\frac 12n^4+\color{red}{\frac 13}n^3+0n^2-\frac 1{30}n\\S_{n,5}&=\frac 16n^6+\frac 12n^5+\color{red}{\frac 5{12}}n^4+0n^3-\frac{1}{12}n^2+0n\\&{\;}\vdots\end{align}$$ Let $R_k=$ the red coefficient of $S_{n,k}$. Then, it appears that, for all $m\in\{0,1,2,3,4\}$, we have $$R_{6+m}+R_{6-m}=1.$$ e.g. $R_6=\frac 12$, thus $m=0\Rightarrow \frac 12 + \frac 12 = 1$; and $R_7=\frac 7{12}$, thus $m=1\Rightarrow \frac 7{12}+\frac 5{12}=1$. Is there some kind of "reason" why this is happening? I mean, upon speculation, I think $R_k=\frac k{12}$ so if that is true, then my conjecture is true. How can this be (dis)proven? Any thoughts? Thanks. Incidentally, I did not include bernoulli-numbers because, although they may be related, I do not specifically focus on them in this question.

论坛问题帖 | has_answers | Mr Pie | 2020-05-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019082 | MSE-3669368

How do we prove this continued fraction for the quotient of gamma functions

Given complex numbers $a=x+iy$, $b=m+in$ and a gamma function $\Gamma(z)$ with $x\gt0$ and $m\gt0$, it is conjectured that the following continued fraction holds $$\frac{\displaystyle4\Gamma\left(\frac{2a+3}{4}\right)\Gamma\left(\frac{2b+3}{4}\right)}{\displaystyle\Gamma\left(\frac{2a+1}{4}\right)\Gamma\left(\frac{2b+1}{4}\right)}=\cfrac{(2a+1)(2b+1)}{a+b+2+\cfrac{(a-b+1)(b-a+1)}{a+b+4+\cfrac{(2a+3)(2b+3)}{a+b+6+\cfrac{(a-b+3)(b-a+3)}{a+b+8+\ddots}}}}\tag{1a}$$ Corollary $$\frac{4}{\pi}=\cfrac{(2)^2}{3+\cfrac{(1)^2}{5+\cfrac{(4)^2}{7+\cfrac{(3)^2}{9+\ddots}}}}\tag{1b}$$ Q: How do we

prove the continued fraction $(1a)$ rigorously?

论坛问题帖 | has_answers | Nicco | 2020-05-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019083 | MSE-3678064

Conjecture about Jensen's inequality and polynomials

Hi it's related to the following conjecture An inequality for polynomials with positives coefficients : We have the first conjecture : Let $x,y>0$ then we have : $$(x+y)f\Big(\frac{x^2+y^2}{x+y}\Big)(f(x)+f(y))\geq 2(xf(x)+yf(y))f\Big(\frac{x+y}{2}\Big)$$ Where :$$f(x)=\sum_{i=0}^{n}a_ix^i$$ And $a_0 \ge a_1 \ge ... \ge a_n>0$ (Thanks to p4sch) Second conjecture : Let $x_i>0$ be n real numbers then we have : $$(\sum_{i=1}^{n}x_i)f\Big(\frac{\sum_{i=1}^{n}x_i^2}{\sum_{i=1}^{n}x_i}\Big)(\sum_{i=1}^{n}f(x_i))\geq n(\sum_{i=1}^{n}x_if(x_i))f\Big(\frac{\sum_{i=1}^{n}x_i}{n}\Big)$$ Where :$$f(x)=\sum_{i=0}^{n}a_ix^i$$ And $a_0 \ge a_1 \ge ... \ge a_n>0$ It seems to be true in the case where $f(x)=\tan(x)$ see A sharp inequality for tangent (refinement of Jensen's inequality) I have tried the approach of River Li in the case $n=3$ without success . I think it's really hard and interesting since at the beginning I have put together two applications of Jensen's inequality . Putting $f(x)=e^x$ gives a nice refinement of Am-Gm (as applications). If it's true I would like just one approach to prove it . Any helps is greatly appreciated . Thanks a lot for all your contributions . Ps:Is it a good idea to put it on maths overflow ? Reference : https://en.wikipedia.org/wiki/Stable_polynomial

论坛问题帖 | has_answers | Barackouda | 2020-05-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019084 | MSE-3691308

Prove a conjecture, balls in boxes, n steps

My uncle gave me the following problem to work on (just for fun), he doesn't know whether the problem has a solution. I haven't been able to solve it and I give up, I don't think my current knowledge is enough to solve it. Problem statement: Initially there are three boxes, labelled A, B and C. A contains $3n$ balls, with $n\geq 5$ a natural number. At step number k, one withdraws k balls from any box and place it into any other box. Show that it is always possible to end up with n balls in each boxes after exactly n steps. The first thing that came to my mind was induction. I.e. build a solution for $n+p$ given a solution for n. It turns out that this will not work because if we have a solution for a given n, then the step $k=n+1$ would be to withdraw $n+1$ balls from any box and place them into another box. However this would make the number of balls negative in a box, so this is not possible. Hence I think it is impossible to construct a solution based on another solution, i.e. just continue the game from the case where there was an equal amount of balls at step n. I have written a Python program to output the solutions for various n's, and I stared hard at the solutions to see whether there are "trivial" ones. I couldn't find such a thing. Sometimes there is a rather "simple looking" solution for a given n, but it is usually not that easy to guess either, and other times I cannot find such nice answers. By the way, the number of answers for a given n grows with n in a strange way (to me), that would make a nice problem on its own (i.e. determine the relationship between the number of solutions and n, but I might ask it in another question.). So I am thinking that maybe the problem can be tackled with number theory, i.e. by finding a methodical/algorithmic way to construct a solution for a given n that is not based on any solution for a lesser n. But I have no knowledge in number theory, so I am not even sure. I am curious to see if people here have ideas and possibly solution(s).

论坛问题帖 | has_answers | untreated_paramediensis_karnik | 2020-05-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019085 | MSE-3707157

To prove the existence of solution(s) of $\frac{a}{b+c}+\frac{b}{a+c}+\frac{c}{a+b}=4$

To prove if a certain equation has a solution, we do not need necessarily solve that equation. Example: Is there any point of the curve $y=\sin^2(x)/x$ between $x=0$ and $x=\pi$ so that the slope of the tangent line at that point is zero? The answer is yes, because y is continuous, differentiable, and $y(0)=y(\pi)=0$ (limits take place). This is (Rolle's Theorem). And we are finish. We did not solve it and find where is that point. My question is about a false conjecture that the equation $$\frac{a}{b+c}+\frac{b}{a+c}+\frac{c}{a+b}=4$$ has no positive integer solutions for a,b,c. Without finding that/those solution(s), which is/are really large numbers, can we prove that the equation has a solution? In other words, can we prove the existence of solution(s) but not by giving an examples like: $a=154476802108746166441951315019919837485664325669565431700026634898253202035277999$ $b=368751317941299998271978115652254748254929799689719709962831374716372246340555795$ $c=437361267792869725786125260237139015281653755816161361862143799337842346777203 6$ The proof can be by contradiction, or direct proof, or any way. Thanks in advance

论坛问题帖 | no_answers_at_retrieval | Hussain-Alqatari | 2020-06-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019086 | MSE-3708545

A conjectured upper bound for $\left(\frac{x^n+1}{x^{n-1}+1}\right)^n+\left(\frac{x+1}{2}\right)^n$ and $x\geq 1$

Hi I have (related https://mathoverflow.net/questions/337457/prove-that-left-fracxn1xn-11-rightn-left-fracx12-rightn): Let $x\geq 1$ a real number and $n\geq 2$ a natural number then we have : $$\Big(\frac{x^{n+1}+1}{x+1}\Big)\Bigg(\frac{\Big(\frac{x+1}{2}\Big)^n}{\Big(\frac{x^2+1}{x+1}\Big)^n}+\frac{(x^n+1)^2}{(x^{n-1}+1)(x^{n+1}+1)}\Bigg)\geq \left(\frac{x^n+1}{x^{n-1}+1}\right)^n+\left(\frac{x+1}{2}\right)^n$$ It's a conjecture so I have tested for $n\leq 20$. Maybe it works also for n a positive real number . If it's true I think we can make something similar to the Peter Mueller's answer. For the case $n=3$ here we have the derivative . Clearly the numerator (of the derivative) is positive for $x\geq 1$.But in some way it's not so easy with the general case . So how to solve it ? Any helps is greatly appreciated. Thanks in advance for all your contribution .

论坛问题帖 | has_answers | Barackouda | 2020-06-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019087 | MSE-3709461

Is there an open mathematical conjecture which has been shown not to be provable in Peano Arithmetic before (possibly) being proved true?

There are several mathematical statements φ about the natural numbers which are known to be true and known not to be provable in Peano Arithmetic (c.f. Gödel's incompleteness theorem, Paris-Harrington theorem, Kirby-Paris theorem...). In all such instances I know of, φ was first shown to be true in "standard mathematical practice" and only later logicians showed that φ is not provable in PA. Is there any example of statement φ about the natural numbers which is not known to be true but already known not to be provable in PA? Or, at least, is there any such example which had such a status in the past (i.e., until actually being proven true)?

论坛问题帖 | no_answers_at_retrieval | Lorenzo | 2020-06-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019088 | MSE-3709724

Conjectured upper bound $\frac{\Gamma(x+n)}{\Gamma(x)}$ and limit at infinity

Let $x\geq 1$ a real number and $n\geq 3$ a natural number then we have : $$\frac{\Gamma(x+n)}{\Gamma(x)}\leq \left(\frac{nx+\left(\frac{n(n-1)}{2}\right)}{\frac{n}{2}}-\left(x^x(x+1)^{x+1}(x+2)^{x+2}\cdots(x+n-1)^{x+n-1}\right)^{\frac{1}{nx+\left(\frac{n(n-1)}{2}\right)}}\right)^{n}$$ It's a conjecture but first for me :$\frac{\Gamma(x+n)}{\Gamma(x)}=x(x+1)\cdots(x+n-1)$.So if there are problems of notations feel free to correct me . I have tested for $n\leq 25$. If it's true I find this inequality beautiful . I conjecture also that the difference tends to 0 when x tends to infinity . I have tried Gautschi's inequality and more without success . Update 08/12/2026: GM and Gw are both of the form $a\bar{} \exp(f(u))$, where $u=\frac{1}{\bar a}$, with $f(u)=\sum_{j\geq 1}c_j u^j$, where $f(u)$ comes from the expansion of $\ln(1+d_k u)$. Faà di Bruno's formula then gives the coefficients of $\exp(f(u))$ in terms of the complete Bell polynomials B_n, evaluated at the derivatives of f. By the exact symmetry of $d_k=k-\frac{n-1}{2}$, namely the invariance under $k\mapsto n-1-k$, all odd moments $m_{2j+1}=\frac{1}{n}\sum_k d_k^{,2j+1}$ vanish exactly. Therefore, f contains only even powers of u, and consequently $G_M+G_w-2\bar a$ does as well. If you have an idea to solve it... Any helps is greatly appreciated . Thanks in advance for all your contributions .

论坛问题帖 | has_answers | Barackouda | 2020-06-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019089 | MSE-3714660

A conjectured criterion for a generalized ramanujan tau function

Given Ramanujan tau function $\tau(n)$, which is the nth Fourier coefficient of the modular discriminant $\displaystyle \Delta(q)=q\prod_{m=1}^\infty (1-q^m)^{24} = \sum_{n=1}^\infty \tau(n)\,q^n\tag{1a}$ Nothing forbids us from generalizing the generating function to $\displaystyle \Delta_{k}(q)= \sum_{n=1}^\infty \tau((2k-1)n)\,q^n\tag{1b}$ for integers $k\gt0$ After some experimentation with wolfram mathematica, one discovers the following intriguing patterns $\Delta_{1}(q)=q^{\color{blue}{1}}-24q^2+252q^3-1472q^4+4830q^5-6048q^6-16744q^7+84480q^8-113643q^{\color{blue}{9}}-115920q^{10}+543612q^{11}-\cdots$ $\Delta_2(q)=252q-6048q^2-113643q^{\color{brown}{3}}-370944q^4+1217160q^5+2727432q^6-4219488q^7+21288960q^8-73279080q^9-29211840q^{10}+134722224q^{11}+167282496q^{12}-145589976q^{13}+101267712q^{14}-548895690q^{15}+248758272q^{16}-1740295368q^{17}+1758697920q^{18}+2686677840q^{19}-1791659520q^{20}+1902838392q^{21}-3233333376q^{22}+4698104544q^{23}-9600560640q^{24}-6425804700q^{25}+3494159424q^26+1665188361q^{\color{brown}{27}}+\cdots$ $\vdots$ From which we deduce the following conjecture The nth coefficient $\tau((2k-1)n)$ is odd if and only if the powers of q in the generating function $(1b)$ are $n\stackrel{\mathrm{def}}{=} \begin{cases}{\color{brown}{(2k-1),9(2k-1),25(2k-1),49(2k-1),...}} & \mbox{ where $2k-1=1,3,5,7,11,13,...$ is any odd square-free integer}\\ \end{cases}$ Or stated differently $\displaystyle\sum_{n=1}^\infty \tau((2k-1)n) q^{n}\equiv \sum_{m=1}^\infty q^{(2k-1)(2m-1)^2}\bmod 2$ for positive integers k That is, $\tau((2k-1)n)$ is odd iff the powers of q in $(1b)$ are $n=(2k-1)(2m-1)^2$ given that $(2k-1)$ is any odd square-free integer valid for $k\gt0$ Question: How do we prove the conjectured criterion? Remark: The Möbius function $\mu(n)$ is either 1 or -1 if n is square-free depending on whether it has a number of even prime factors or odd respectively and $\mu(n)=0$ if it is non square-free. It happens to be a very important arithmetic function in number theory. For example, the Dirichlet series that generates the Möbius function is the reciprocal of the Riemann zeta function $\displaystyle \sum_{n=1}^{\infty} \frac{\mu(n)}{n^s}=\frac{1}{\zeta(s)}$. The fact that this conjectured congruence has something to do with the möbius function is exactly what makes it interesting.

论坛问题帖 | has_answers | Nicco | 2020-06-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019090 | MSE-3718571

Little conjecture in elementary number theory

I was calcultaing quadratic residues. I noticed nice pattern: quadratic residues modulo p^n where p is prime are well-behaved; they always repeat after $\frac{p^n - 1}{2}$-th place (starting with $1^2 \equiv 1 \pmod {p^n}$. I know this is proven for $n = 1.$ Also, I noticed that many composite numbers have random-looking quadratic residues. I found only few exception: 10 (1, 4, 9, 6, 5, 6, 9, 4, 1), 6 (1, 4, 3, 4, 1). Three examples for $n = 2$: (1) $p = {2^2} = 4$: 1, 0, 1, 0, etc. (2) $p = {5^2} = 25$: 1, 4, 9, ..., 19, 19, ... 9, 4, 1. (3) $p = 3^2 = 9$: 1, 4, 0, 7, 7, 0, 4, 1. Two examples for $n = 3$: (1') $p = 2^3 = 8$: 0, 1, 4, 1, 0, 1, 4, 1. (2') $p = 3^3 = 27$: 1, 4, 9, 16, 25, 9, ..., 7, 7 ... 9, 25, 16, 9, 4, 1. So, I see that residues can repeat (see example (2')) but they always have this property that sequance repeat after half terms. Is there proof of this?

论坛问题帖 | no_answers_at_retrieval | 1b3b | 2020-06-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019091 | MSE-3720368

Conjecture on $\pi(n)$ and other arithmetic functions

Notations Let $\pi(n)$ be the prime countiong function. Let $\alpha(n)$ denote the number of prime factors of n and $\beta(n)$ the sum of the prime factors of n. In other words, if $$n=p_1^{x_1}p_2^{x_2}...p_m^{x_m}$$ then $\alpha(n)=m$ and $\beta(n)=p_1+p_2+...+p_m$ (I changed the notations; It was pointed out in the comments that ψ and ω are other functions and they were misleading) Statement Let $\pi(n)$ be the prime counting function, $\psi(n)$ the number of prime factors and $\omega(n)$ the sum of the prime factors of n. Prove or disprove that there are infinitely many n such that: $$\pi(n)=\alpha(n)\beta(n)$$ $\text{Some other observations}$ Here are the first solutions: 4, 120, 437, 546, 620, 8144, 11509, 170049, 170907, 340655, 478476, 789575 Also, if we use the Prime Number Theorem ($\pi(n)$ is aproximatively $\frac{n}{\ln(n)}$) we can derive solutions in an exhaustive manner. For example, find at what point $\frac{n}{\pi(n)}$ is aproximatively $\frac{47}{2}$ and then subtract 47 from n until you get the solution. Using this method we get 45764089927 which in fact works. $\text{Why is it an interesting problem?}$ I find it rather interesting due to the unusual relationship between these 3 functions. I played a while with several arithmetic functions and couldn't find strong conection between them, but the equation $\pi(n)=\alpha(n)\cdot\beta(n)$ seems promising.

论坛问题帖 | has_answers | user799688 | 2020-06-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019092 | MSE-3720572

Conjecture regarding the sum of prime factors

Notations Let $\pi(n)$ be the prime countiong function. Let denote $\alpha(n)$ the sum of the prime factors of n. In other words, if $$n=p_1^{x_1}p_2^{x_2}...p_m^{x_m}$$ then $\alpha(n)=p_1+p_2+...+p_m$ (I changed the notation; It was pointed out in the comments that ω is another function and it was misleading) Statement Prove or disprove that there exist infinitely many composite positive integers n such that $\alpha(n)+1|n+1$. Important I made a new thread in which the question is posted with some new conditions. I am now interested in the squarefree solutions of the above equation. New problem link: Conjecture on the sum of prime factors

论坛问题帖 | has_answers | user799688 | 2020-06-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019093 | MSE-3720724

Conjecture on the sum of prime factors

Notations Let $\pi(n)$ be the prime counting function. Let denote $\alpha(n)$ the sum of the prime factors of n. (i.e. In other words, if $$n=p_1^{x_1}p_2^{x_2}...p_m^{x_m}$$ then $\alpha(n)=p_1+p_2+...+p_m$.) (I changed the notation; It was pointed out in the comments that ω is another function and it was misleading) Statement Find whether the equation $\alpha(n)+1|n+1$ has infinitely many squarefree solutions. Moreover, show that one can find an infinitude of composite, non-squarefree solutions. Progress The original question was posted here and Peter found the following general solution for non-squarefree integers: Claim : $n=2\cdot 3\cdot 5^{5m+4}$ is a solution for all $m\ge 1$ Proof : The sum of the prime factors is obviously 10. Because of $5^5\equiv 1\mod 11$, we have $n\equiv 2\cdot 3\cdot 5^4\equiv -1\mod > 11$, hence $11\mid n+1$ Hence there are infinite many solutions. So it has been proven that there are in fact infinitely many non-squarefree solutions (and the prime numbers are trivial solutions). The question about composite squarefree solutions still holds.

论坛问题帖 | no_answers_at_retrieval | user799688 | 2020-06-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019094 | MSE-3722533

Problem from Discrete Mathematics and its application for Rosen section 4.4

This exercise outlines a proof of Fermat’s little theorem. a) Suppose that a is not divisible by the prime p. Show that no two of the integers 1 · a, 2 · a, . . . , (p - 1)a are congruent modulo p. b) Conclude from part (a) that the product of 1, 2, . . . , p - 1 is congruent modulo p to the product of a, 2a, . . . , (p - 1)a. Use this to show that (p - 1)! ≡ a p-1 (p - 1)! (mod p). I already solved the first problem (a) but I couldn't solve Problem (b) So can any one please help me to solve it, mean problem (b) ? Note : I can't use Fermat’s little theorem to prove it as this problem is to prove the Fermat’s little theorem

论坛问题帖 | has_answers | Blue Fire | 2020-06-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019095 | MSE-3723793

Conjecture on prime generating sequence

Statement Denote p_n the nth prime. Consider the set $$\mathcal{P}=\{\sum_{i=1}^np_i:n\in\mathbb{N}\}$$ Prove that $|\mathcal{P}\cap\mathbb{P}|=\infty$ $\text{Why am I studying this}$ First of all, I find it an itriguing and rather unusual result. I also want to see how we can generate primes by adding numbers, so this was a natural idea. To conclude, I also highly highly suspect these primes have some extra properties from normal ones. $\text{Computational arguments}$ First of all, here are the first 200 values for $\sum_{i=1}^{n}p_i$: 5 10 17 28 41 58 77 100 129 160 197 238 281 328 381 440 501 568 639 712 791 874 963 1060 1161 1264 1371 1480 1593 1720 1851 1988 2127 2276 2427 2584 2747 2914 3087 3266 3447 3638 3831 4028 4227 4438 4661 4888 5117 5350 5589 5830 6081 6338 6601 6870 7141 7418 7699 7982 8275 8582 8893 9206 9523 9854 10191 10538 10887 11240 11599 11966 12339 12718 13101 13490 13887 14288 14697 15116 15537 15968 16401 16840 17283 17732 18189 18650 19113 19580 20059 20546 21037 21536 22039 22548 23069 23592 24133 24680 25237 25800 26369 26940 27517 28104 28697 29296 29897 30504 31117 31734 32353 32984 33625 34268 34915 35568 36227 36888 37561 38238 38921 39612 40313 41022 41741 42468 43201 43940 44683 45434 46191 46952 47721 48494 49281 50078 50887 51698 52519 53342 54169 54998 55837 56690 57547 58406 59269 60146 61027

61910 62797 63704 64615 65534 66463 67400 68341 69288 70241 71208 72179 73156 74139 75130 76127 77136 78149 79168 80189 81220 82253 83292 84341 85392 86453 87516 88585 89672 90763 91856 92953 94056 95165 96282 97405 98534 99685 100838 102001 103172 104353 105540 106733 107934 109147 110364 These are the primes that were generated: 2, 5, 17, 41, 197, 281, 7699, 8893, 22039, 24133, 25237, 28697, 32353, 37561, 38921, 43201, 44683, 55837, 61027, 66463, 70241, 86453, 102001, (and some others: 109147, 116533, 119069, 121631, 129419, 132059, 263171, 287137, 325019, 329401, 333821, 338279, 342761) Generalizations Of course, there are many ways to generalize this. I will state 3, which might be of interest. However, for the sake of simplicity, i will only leave the first statement as a conjecture. These new results are quite similar so I think a hypothetical proof would not differ too much. $\text{Generalization } 1:$ The same question as the original one, but this time $$\mathcal{P_k}=\{\sum_{i=k}^np_i:n\in\mathbb{N}\}$$ Prove that $\forall k$, $|\mathcal{P_k}\cap\mathbb{P}|=\infty$ $\text{Generalization } 2:$ The same question as the original one, but this time $$\mathcal{P^l}=\{\sum_{i=1}^np_i^l:n\in\mathbb{N}\}$$ Prove that $\forall k$, $|\mathcal{P^l}\cap\mathbb{P}|=\infty$ $\text{Generalization } 3:$ A combination of generalization 1 and 2 $$\mathcal{P_k^l}=\{\sum_{i=k}^np_i^l:n\in\mathbb{N}\}$$ Prove that $\forall k$, $|\mathcal{P_k^l}\cap\mathbb{P}|=\infty$ I am not providing computational arguments for these ones. More Resources https://arxiv.org/abs/1804.04198

论坛问题帖 | no_answers_at_retrieval | user799688 | 2020-06-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019096 | MSE-3738458

If $N = q^k n^2$ is an odd perfect number with special prime q, then can N be of the form $q^k \cdot (\sigma(q^k)/2) \cdot {n}$?

Let $\sigma(x)$ denote the sum of divisors of the positive integer x. A number M is said to be perfect if $\sigma(M)=2M$. For example, 6 and 28 are perfect since $$\sigma(6) = 1 + 2 + 3 + 6 = 2\cdot{6}$$ and $$\sigma(28) = 1 + 2 + 4 + 7 + 14 + 28 = 2\cdot{28}.$$ It is currently unknown if there are infinitely many even perfect numbers. It is also an open problem whether any odd perfect numbers exist. It is widely believed that there are no odd perfect numbers. Euler proved that an odd perfect number N, if one exists, must necessarily have the so-called Eulerian form $$N = q^k n^2$$ where q is the special/Euler prime satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. QUESTION Here is my question: Can the odd perfect number N be of the form $$\frac{q^k \sigma(q^k)}{2}\cdot{n}?$$ I would certainly appreciate it if anybody could point me to papers/articles/publications in the literature where this particular inquiry is covered. CONTEXT Slowak (1999) proved that the odd perfect number N must be of the form $$\frac{q^k \sigma(q^k)}{2}\cdot{d},$$ where $d > 1$. Dris (2017) showed further that d must have the form $$\frac{D(n^2)}{\sigma(q^{k-1})}=\gcd(n^2,\sigma(n^2))=\frac{\sigma(n^2)}{q^k}=\frac{n^2}{\sigma(q^k)/2},$$ where $D(x)=2x-\sigma(x)$ is the deficiency of x.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2020-06-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019097 | MSE-3746635

On Erdös-Szekeres convex polygons lower bound

I have problems with the construction of 2^{n-2} points that contain no n-gon, particulary, the proof of the book "Open Problems in Mathematics". The proof sais that: For $i = 0, ..., n-2$ let T_i be a set of ${{n-2}\choose i}$ points in general position in the plane, containing no $i+2$-cap and no $n-i$-cup and having the property that no two points in the set are connected by a line having slope of absolute value greater than 1. (The existence of T_i follows from a previous theorem, and the condition on slopes can be satisfied by “compressing” T_i vertically.) Place a small copy of T_i in a neighborhood of the point on the unit circle making an angle of $\frac{\pi}{4} - \frac{i\pi}{(n-2)2}$ with the positive x-axis. Let X be the union of $T_1, T_2, ..., T_{n-2}$. Clearly, $|X| = 2^{n-2}$. Suppose that Y is a convexly independent subset of X. Let k and l be the smallest and the largest values of i so that $Y\cap T_i\neq \emptyset$. If $k=l$, then Y contains no $k+2$-cap and no $n-k$-cup. The construction guarantees that: $Y \cap T_k$ is a cap of at most $k + 1$ points, $Y \cap T_l$ is a cup of at most $n + l- 1$ points, $|Y \cap T_i|\leq 1$ for all $i = k + 1, ..., l-1$ Summing up, $|Y| = (k + 1) + (l + k - 1) + (n -l - 1) = n-1$. I understood all the reasoning except the part that say: $|Y\cap T_i|\leq 1$ for all $k+1\leq i\leq l-1$. Why in the middle sets the intersection with Y have at least one element?

论坛问题帖 | no_answers_at_retrieval | Eduardo Quintana | 2020-07-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019098 | MSE-3754664

Reference request for conjecture about bridge and crossing number of knots

Murasugi in his book (Knot theory and its applications, page 60) writes: Conjecture. If K is a knot, then $c(K) \ge 3(br(K) - 1)$, where equality only holds when K is the trivial knot, the trefoil knot, or the (connected) sum of trefoil knots. I would like to know who came up first with this conjecture, if there was any progress on it in recent years and if it has some special name. Murasugi's book is quite tacit here.

论坛问题帖 | has_answers | user471311 | 2020-07-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019099 | MSE-3757936

Pretty conjecture $x^{\left(\frac{y}{x}\right)^n}+y^{\left(\frac{x}{y}\right)^n}\leq 1$

inspired (again) by an inequality of Vasile Cirtoaje I propose my own conjecture : Let $x,y>0$ such that $x+y=1$ and $n\geq 1$ a natural number then we have : $$x^{\left(\frac{y}{x}\right)^n}+y^{\left(\frac{x}{y}\right)^n}\leq 1$$ First I find it very nice because all the coefficient are 1 . I have tested with Geogebra until $n=50$ without any counter-examples. Furthermore we have an equality case as $x=y=0.5$ or $x=1$ and $y=0$ and vice versa . To solve it I have tried all the ideas here My main idea was to make a link with this inequality (my inspiration) see here So if you can help me to solve it or give me

an approach... ...Thanks for all your contributions ! Little update I think there is also an invariance as in question here Conjecture $a^{(\frac{a}{b})^p}+b^{(\frac{b}{a})^p}+c\geq 1$ Theoretical method Well,Well this method is very simple but the result is a little bit crazy (for me (and you ?)) Well ,I know that if we put $n=2$ we can find (using parabola) an upper bound like $$x^{\left(\frac{1-x}{x}\right)^2}\leq ax^2+bx+c=p(x)$$ And $$(1-x)^{\left(\frac{x}{1-x}\right)^2}\leq ux^2+vx+w=q(x)$$ on $[\alpha,\frac{1}{2}]$ with $\alpha>0$ and such that $p(x)+q(x)<1$ In the neightborhood of 0 we can use a cubic . Well,now we have (summing) : $$x^{\left(\frac{1-x}{x}\right)^2}+(1-x)^{\left(\frac{x}{1-x}\right)^2}\leq p(x)+q(x)$$ We add a variable ε such that $(p(x)+\varepsilon)+q(x)=1$ Now we want an inequality of the kind ($k\geq 2$): $$x^{\left(\frac{1-x}{x}\right)^{2k}}+(1-x)^{\left(\frac{x}{1-x}\right)^{2k}}\leq (p(x)+\varepsilon)^{\left(\frac{1-x}{x}\right)^{2k-2}}+q(x)^{\left(\frac{x}{1-x}\right)^{2k-2}}$$ Now and it's a crucial idea we want something like : $$\left(\frac{x}{1-x}\right)^{2k-2}\geq \left(\frac{1-(p(x)+\varepsilon)}{q(x)}\right)^y$$ AND : $$\left(\frac{1-x}{x}\right)^{2k-2}\geq \left(\frac{1-q(x)}{p(x)+\varepsilon}\right)^y$$ Now it's not hard to find a such y using logarithm . We get someting like : $$x^{\left(\frac{1-x}{x}\right)^{2k}}+(1-x)^{\left(\frac{x}{1-x}\right)^{2k}}\leq q(x)^{\left(\frac{1-q(x)}{q(x)}\right)^{y}}+(1-q(x))^{\left(\frac{q(x)}{1-q(x)}\right)^{y}}$$ Furthermore the successive iterations of this method conducts to 1 because the values of the differents polynomials (wich are an approximation of the initial curve) tend to zero or one (as abscissa). The extra-thing (and a little bit crazy) we can make an order on all the values. My second question Is it unusable as theoretical\practical method ?

论坛问题帖 | has_answers | Barackouda | 2020-07-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019100 | MSE-3769796

If a is deficient-perfect, can ab be deficient-perfect if $b > 1$, and ab is odd?

Let $\sigma(x)$ denote the sum of divisors of the positive integer x. The deficiency of x is given by the formula $$D(x)=2x-\sigma(x).$$ A number y is said to be deficient-perfect if $D(y) \mid y$. Note that a deficient-perfect number z is automatically deficient (i.e., z satisfies $D(z) \geq 1$). Here is my question: If a is deficient-perfect, can ab be deficient-perfect if $b > 1$, and ab is odd? The reason for the condition $ab \equiv 1 \pmod 2$ is to exclude the case $$(a_1,b_1)=\bigg(2^k,2^{k+1}+2^t-1\bigg),$$ where $2^{k+1}+2^t-1$ is an odd prime and $t \leq k$, as well as the case $$(a_2,b_2)=(2^r,2^s).$$ MY ATTEMPT Let ab be odd. This implies that a is odd and b is odd. Let a be deficient-perfect, and suppose to the contrary that ab is also deficient-perfect, where $b>1$. Then we have the divisibility constraints $$D(a) \mid a \iff a = (2a - \sigma(a))c$$ $$D(ab) \mid ab \iff ab = (2ab - \sigma(ab))d$$ Substituting the first equation into the variable a in the LHS of the second equation, we obtain $$(2a - \sigma(a))bc = (2ab - \sigma(ab))d \iff 2abc - bc\sigma(a) = 2abd - d\sigma(ab)$$ $$\iff 2ab(c-d) = bc\sigma(a) - d\sigma(ab).$$ But $$c - d = \frac{a}{D(a)} - \frac{ab}{D(ab)} = a\cdot\bigg(\frac{D(ab)-bD(a)}{D(a)D(ab)}\bigg) = a\cdot\bigg(\frac{2ab-\sigma(ab)-2ab+b\sigma(a)}{D(a)D(ab)}\bigg)$$ $$=a\cdot\bigg(\frac{-\sigma(ab)+b\sigma(a)}{D(a)D(ab)}\bigg)$$ which implies that $c - d \leq 0$ since $$b\sigma(a) \leq \sigma(ab).$$ We therefore conclude that $$2ab(c-d) = bc\sigma(a) - d\sigma(ab) \leq 0.$$ Alas, this is where I get stuck. Postscript: Note that $$\mathscr{D} = 9018009 = {3^2}\cdot{7^2}\cdot{{11}^2}\cdot{{13}^2}$$ is the lone odd deficient-perfect number that we know of, to this day. So essentially, the question asks whether we could produce another odd deficient-perfect number from $\mathscr{D}$ (via multiplying $\mathscr{D}$ by $\mathscr{D'} > 1$), or if it could be proven that this is impossible.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2020-07-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019101 | MSE-3771678

$\pi(n)$ is always more than the sum of the prime indices of the factors of composite $n \geq 12$

Let $n=p_1^{a_1} p_2^{a_2} \cdots p_k^{a_k} \geq 12$ be any composite integer. Then it seems that this is true: $$\pi(n) > \sum_{i=1}^{k}{\pi(p_i)a_i}\ .$$ You get equality instead iff n is prime. I also assume that if it is true, it's a known result. Can anyone point me towards a resource discussing it if so? Alternatively, if I've made a mistake and/or this is a trivial result, please point out how. Also, if true, I think Bertrand's Postulate immediately follows: For prime p, it gives $\pi(2p)>\pi(p)+1$, implying at least one prime between p and $2p$. By the same token, $\pi(3p)>\pi(p)+2$, and $\pi(p^2)>2 \pi(p)$. (See my answer below for additional thoughts.)

论坛问题帖 | has_answers | Trevor | 2020-07-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019102 | MSE-3776712

Is this a valid proof for $I(n^2) \geq \frac{5}{3}$, if $q^k n^2$ is an odd perfect number with special prime q?

Let $\sigma(x)$ denote the sum of divisors of the positive integer x, and let $I(x)=\sigma(x)/x$ be the abundancy index of x. Note that both σ and I are multiplicative functions. A number m is said to be perfect if $\sigma(m)=2m$. Equivalently, $I(m)=2$. Euler proved that an odd perfect number, if one exists, must have the form $$m = q^k n^2$$ where q is the special prime satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. Since q is prime, we have $$\frac{q+1}{q} = I(q) \leq I(q^k) = \frac{\sigma(q^k)}{q^k} = \frac{q^{k+1} - 1}{q^k (q - 1)} < \frac{q^{k+1}}{q^k (q - 1)} = \frac{q}{q - 1}$$ from which it follows that $$\frac{2(q-1)}{q} < I(n^2) = \frac{2}{I(q^k)} \leq \frac{2q}{q+1}.$$ Note that we then have the lower bound $$I(n^2) > \frac{2(q-1)}{q} \geq \frac{8}{5}$$ since q is a prime satisfying $q \equiv 1 \pmod 4$. Here is my initial question: Can we improve the lower bound for $I(n^2)$ to $$I(n^2) \geq \frac{5}{3}$$ using the following argument? $$\bigg(\frac{2q}{q+1} \geq I(n^2) > \frac{5}{3}\bigg) \implies q > 5 \implies q \geq 13 \implies \bigg(I(n^2) > \frac{2(q-1)}{q} \geq \frac{24}{13} > \frac{5}{3}\bigg)$$ Thus, we

have the biconditional $$I(n^2) > \frac{5}{3} \iff q > 5.$$ Next, we have the implication $$I(n^2) = \frac{5}{3} \implies q = 5.$$ It then suffices to prove the implication $$q = 5 \implies I(n^2) = \frac{5}{3}$$ to finally show that $$I(n^2) \geq \frac{5}{3},$$ since $q \geq 5$ holds. But note that, if $q=5$, then $$\frac{5}{3} = I(n^2) = \frac{2}{I(5^k)} = \frac{2\cdot{5^k}(5-1)}{5^{k+1}-1}$$ which implies that the Descartes-Frenicle-Sorli Conjecture that $k=1$ holds. Still, notice that we have $$k=1 \implies I(q^k) = I(q) = \frac{q+1}{q} = 1 + \frac{1}{q} \leq \frac{6}{5} \implies I(n^2) = \frac{2}{I(q^k)} = \frac{2}{I(q)} \geq \frac{2\cdot{5}}{6} = \frac{5}{3},$$ which is what we set out to prove. Here is my final question: Would it be possible to remove the reliance of the proof on the Descartes-Frenicle-Sorli Conjecture?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2020-08-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019103 | MSE-3779332

Properties of subsets for which $\sum 1/k$ diverges

The well-known Erdos-Turan conjecture is the following. Let $V \subset \mathbb{N}$ be such that $\sum_V k^{-1}$ diverges. Then V contains arithmetic progressions of every possible length. A recent result showed such a set V must contain infinitely many length-3 progressions. I am wondering: if $V \subset \mathbb{N}$ is such that $\sum_V k^{-1}=\infty$, what are some properties of V? This Wikipedia page gives a small number of properties, but I am curious whether there are others (being non-trivial).

论坛问题帖 | has_answers | Descartes Before the Horse | 2020-08-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019104 | MSE-3780183

Proving $\pi(ab) \geq \pi(a)\pi(b)$?

It occurs to me that a stronger version of my related post the other day would be: $$\pi(ab) \geq \pi(a) \cdot \pi(b)\ ,$$ which I believe holds for all naturals a,b except for the two pairs $(5,7)$ and $(7,7)$. My first thought on a proof was to point out that $$\pi(ab)\sim\frac{ab}{\log(ab)}>\frac{ab}{\log(a)\log(b)}\sim \pi(a)\pi(b)\ ,$$ given $a>e$ and $b>a^{1-\log{a}}$ to rule out the small log problems, though I'm not sure that's necessary. My question is whether that's enough to establish that the conjecture is always true for sufficiently large ab, or if it merely implies that it will tend towards being true in the limit. If this isn't enough and someone has a different way to go at it, I'd be interested to hear it. And if this approach is valid, is it safe to change the conjecture to the stricter $\pi(ab)>\pi(a)\pi(b)$ (for large enough ab)?

论坛问题帖 | has_answers | Trevor | 2020-08-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019105 | MSE-3785911

Conjecture about graph with points and arrows with weights

Postulates Each point has arrow(s), and each arrow is pointing to another point. An arrow of a point cannot point to itself. Arrows can only point to any direction on the right. That means $\uparrow$, $\downarrow$ are not allowed, and of course $\leftarrow$ is not allowed. Each arrow has a number x on it with $0<x<1$ and $x\in\mathbb{R}$. Of a point, say A, arrows pointing out to the right must have their numbers summing up to 1, only if the arrows exist. (That means if a point does not have arrows pointing to the right, the condition does not have to be satisfied) Same for arrows pointing to A. Conjecture Suppose there are a few points and arrows. Prove that if there are some points each with some arrows pointing to the right, there must be the same case but being pointed from the left, in order to satisfy the conditions listed above. (At least this is the pattern I can see.) For example, if there are 2 points each with 3 arrows each pointing to another point, there must be two points each being pointed with 3 arrows. I think it is related to Graph Theory, but I have no idea of proving it, or better say I am not familiar with Graph Theory. Evidences Example 1: There exists $A\xrightarrow{1} B$. Since there is 1 point with 1 arrow pointing to another point, there must be and there is 1 point being pointed with 1 arrow. Example 2: point A and B both have two arrows pointing to C and D. Hence, both point C and D are being pointed by two arrows from A and B. Example 3: point A and C both have two arrows pointing out, so two points D and F are being pointed by two arrows. point B has three arrows pointing out, so a point E is being pointed by three arrows. This graph also satisfies the conditions above: For point A: $0.8+0.2=1$ For point B: $0.2+0.3+0.5=1$ For point C: $0.3+0.7=1$ For point D: $0.8+0.2=1$ For point E: $0.2+0.3+0.5=1$ For point F: $0.3+0.7=1$ These are just simple cases. The conjecture will not be obvious or maybe even false in more complex graphs.

论坛问题帖 | has_answers | Bernard | 2020-08-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019106 | MSE-3790631

Rare interesting primes such as Wolstenholme and Wieferich primes

Take a look at the following two examples of "rare interesting primes"; Wolstenholme prime is a special prime number related to a stronger version of Wolstenholme's theorem. They have connections to Fermat's last theorem. The only known such primes are 16843 and 2124679, but it is believed (conjectured) that there are infinitely many. The corresponding OEIS sequence is A088164. Definition. A Wolstenholme prime is a prime number $p > 7$ that satisfies the congruence $$\left(\begin{array}{l}2 p-1 \\p-1\end{array}\right) \equiv 1 \quad\left(\bmod p^{4}\right).$$ Wieferich prime is a prime number related to a stronger version of Fermat's little theorem. They appear in works pertaining to Fermat's last theorem. The only known such primes are 1093 and 3511, but it is believed that there are infinitely many. The corresponding OEIS sequence is A001220. Definition. Wieferich prime is a prime number p such that p^2 divides $2^{p - 1} - 1$. I was wondering, Question. Are there any other types of "rare interesting primes"? I.e. primes related to known results in number theory and have been studied (there exist compelling references, i.e. such primes are "interesting"), but do not have many examples (are "rare"). For non-example, Twin primes are "interesting", but are not "rare" since one can easily list say 10^4 examples.

For example, Fermat primes are "interesting" and are "rare". Only known are $3, 5, 17, 257, 65537$. It is conjectured that these are the only terms (unlike the Wolstenholme and Wieferich primes). Another example may be of Mersenne Primes. We can list around 50 examples at the moment, where the largest example holds the current record for the largest prime. Are there any other examples?

论坛问题帖 | has_answers | Vepir | 2020-08-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019107 | MSE-3792709

A conjecture about Catalan sequence

For $n=2$, consider the free abelian group generated by polynomials corresponding to $\frac{(2 n)!}{2^n n!}=3$ partitions of $2n=4$ vertices into pairs (chords): $$ (x_1-x_3)(x_2-x_4),(x_1-x_2)(x_3-x_4),(x_1-x_4)(x_2-x_3) $$ Euler's Identity: $ (x_1-x_3)(x_2-x_4)=(x_1-x_2)(x_3-x_4)+(x_1-x_4)(x_2-x_3)$. So$\{(x_1-x_2)(x_3-x_4),(x_1-x_4)(x_2-x_3)\}\tag1\label2$$ is a basis. So the rank of the free abelian group is 2. For $n=3$, consider the free abelian group generated by polynomials corresponding to $\frac{(2 n)!}{2^n n!}=15$ partitions of $2n=6$ vertices into pairs (chords): \begin{array}r \{(x_1-x_2) (x_3-x_4) (x_5-x_6),\\(x_1-x_2) (x_3-x_5) (x_4-x_6),\\(x_1-x_2) (x_4-x_5) (x_3-x_6),\\(x_1-x_3) (x_2-x_4) (x_5-x_6),\\(x_1-x_3) (x_2-x_5) (x_4-x_6),\\(x_1-x_3) (x_4-x_5) (x_2-x_6),\\(x_2-x_3) (x_1-x_4) (x_5-x_6),\\(x_1-x_4) (x_2-x_5) (x_3-x_6),\\(x_1-x_4) (x_3-x_5) (x_2-x_6),\\(x_2-x_3) (x_1-x_5) (x_4-x_6),\\(x_2-x_4) (x_1-x_5) (x_3-x_6),\\(x_3-x_4) (x_1-x_5) (x_2-x_6),\\(x_2-x_3) (x_4-x_5) (x_1-x_6),\\(x_2-x_4) (x_3-x_5) (x_1-x_6),\\(x_3-x_4) (x_2-x_5) (x_1-x_6)\} \end{array} Using Euler's Identity we can decompose the product of intersecting chords into the sum of products of the non-intersecting chords. For example $(x_1-x_4)(x_2-x_5)(x_3-x_6)$ as in Fuhrmann's Theorem, There are 5 partitions of 6 vertices into non-intersecting chords $$\tag2\label1\begin{array}l\{(x_1-x_6)(x_2-x_5)(x_3-x_4),\\ (x_1-x_6)(x_2-x_3)(x_4-x_5),\\ (x_1-x_4)(x_2-x_3)(x_5-x_6),\\(x_1-x_2)(x_3-x_6)(x_4-x_5),\\(x_1-x_2)(x_3-x_4)(x_5-x_6)\}\end{array}$$ Under lexicographic order, the leading terms of \eqref{1} are distinct: $\{x_1x_2x_3,x_1x_2x_4,x_1x_2x_5,x_1x_3x_4,x_1x_3x_5\}$. Suppose $c_1(x_1-x_6)(x_2-x_5)(x_3-x_4)+\dots+c_5(x_1-x_2)(x_3-x_4)(x_5-x_6)=0$ for some $c_1,\dots,c_5\in\mathbb{Z}$. Comparing coefficients of term $x_1x_2x_3$ to get $c_1=0$, comparing coefficients of term $x_1x_2x_4$ to get $c_2=0$, comparing coefficients of term $x_1x_2x_5$ to get $c_3=0$, comparing coefficients of term $x_1x_3x_4$ to get $c_4=0$, comparing coefficients of term $x_1x_3x_5$ to get $c_5=0$, so $c_1=\dots=c_5=0$, \eqref{1} is linearly independent. So the rank of the free abelian group is 5. Here is a proof that the number of non-intersecting chords is Catalan number $\frac1{n+1}\binom{2n}n$. Each polynomial in \eqref{1} can be written as a $2×n$ "non-intersecting tableaux" $$ \Biggl\{\begin{array}{|c|c|}\hline1&2&3\\\hline6&5&4\\\hline\end{array}, \begin{array}{|c|c|}\hline1&2&4\\\hline6&3&5\\\hline\end{array}, \begin{array}{|c|c|}\hline1&2&5\\\hline4&3&6\\\hline\end{array}, \begin{array}{|c|c|}\hline1&3&4\\\hline2&6&5\\\hline\end{array}, \begin{array}{|c|c|}\hline1&3&5\\\hline2&4&6\\\hline\end{array}\Biggr\} $$ To show that a "non-intersecting tableux" is determined by its first row, we give a procedure to fill in the second row. For example $\begin{array}{|c|c|}\hline1&2&5&7\\\hline?&?&?&?\\\hline\end{array}$ to fill $3,4,6,8$ in the second row such that there are no intersecting chords, we must put 3 under the largest number less than $3:\begin{array}{|c|c|}\hline1&2&5&7\\\hline?&3&?&?\\\hline\end{array}$ then we must put 4 under the largest remaining number less than $4:\begin{array}{|c|c|}\hline1&2&5&7\\\hline4&3&?&?\\\hline\end{array}$, then we must put 6 under the largest remaining number less than $6:\begin{array}{|c|c|}\hline1&2&5&7\\\hline4&3&6&?\\\hline\end{array}$, then we must put 8 under the largest remaining number less than $8:\begin{array}{|c|c|}\hline1&2&5&7\\\hline4&3&6&8\\\hline\end{array}$ For all n we have shown the first rows of $\frac1{n+1}\binom{2n}n$ number of "non-intersecting tableaux" are distinct. Under lexicographic order, the leading terms of \eqref{1} correspond to the first row of tableaux, so the leading terms are distinct, so the set of polynomials corresponding to "non-intersecting tableux" are linearly independent for all n. So we have proved: For all n, the rank of the above free abelian group in $2n$ variables is $\frac1{n+1}\binom{2n}n$. For $n=4$, similarly decompose the product of intersecting chords into the sum of products of the non-intersecting chords. For example $(x_1-x_5)(x_2-x_6)(x_3-x_7)(x_4-x_8)$ As you can see, there will be two identical pairs in the end, so after collecting terms there will be two terms with coefficient 2 \begin{align} (x_1 - x_5)(x_2 - x_6)(x_3 - x_7)(x_4 - x_8)\\ = (x_1 - x_4)(x_2 - x_3)(x_5 - x_8)(x_6 - x_7)\\ + \color{red}{2}(x_1 - x_8)(x_2 - x_3)(x_4 - x_5)(x_6 - x_7)\\ + (x_1 - x_2)(x_3 - x_8)(x_4 - x_7)(x_5 - x_6)\\ + (x_1 - x_2)(x_3 - x_8)(x_4 - x_5)(x_6 - x_7)\\ + \color{red}{2}(x_1 - x_2)(x_3 - x_4)(x_5 - x_6)(x_7 - x_8)\\ + (x_1 - x_2)(x_3 - x_4)(x_5 - x_8)(x_6 - x_7)\\ + (x_1 - x_4)(x_2 - x_3)(x_5 - x_6)(x_7 - x_8)\\ + (x_1 - x_8)(x_2 - x_3)(x_4 - x_7)(x_5 - x_6)\\ + (x_1 - x_2)(x_3 - x_6)(x_4 - x_5)(x_7 - x_8)\\ + (x_1 - x_8)(x_2 - x_7)(x_3 - x_4)(x_5 - x_6)\\ + (x_1 - x_8)(x_2 - x_7)(x_3 - x_6)(x_4 - x_5)\\ + (x_1 - x_8)(x_2 - x_5)(x_3 - x_4)(x_6 - x_7)\\ + (x_1 - x_6)(x_2 - x_3)(x_4 - x_5)(x_7 - x_8)\\ + (x_1 - x_6)(x_2 - x_5)(x_3 - x_4)(x_7 - x_8)\end{align} Conjecture. Express $(x_1-x_{n+1})(x_2-x_{n+2})\dots(x_n-x_{2n})$ as a linear combination of polynomials corresponding to the non-intersecting chords, then the term $(x_1-x_2)(x_3-x_4)\cdots(x_{2n-1}-x_{2n})$ and the term $(x_{2n}-x_1)(x_2-x_3)\cdots(x_{2n-2}-x_{2n-1})$ have largest coefficient in absolute value. Mathematica code to compute the largest coefficients in absolute value Needs["Combinatorica`"]; coeff[n_]:=Max[Abs[PolynomialReduce[Product[Indexed[x,i]-Indexed[x,n+i],{i,n}],Table[Module[{t1=t[[1]],t2=t[[2]],poly=1,i=1},Do[i=1;While[Length[t1]>i&&t1[[i+1]]<t2[[1]],i++];poly*=Indexed[x,t1[[i]]]-Indexed[x,t2[[1]]];t1=Delete[t1,i];t2=Rest[t2],n];poly],{t,Tableaux[Table[n,2]]}],Table[Indexed[x,i],{i,2n}]][[1]]] the terms $n=2,\dots,9$ in the sequence agree with A099960: \begin{array}{c|c} n&2&3&4&5&6&7&8&9\\\hline \text{largest abs coefficient}&1&1&2&3&8&17&56&155 \end{array}

论坛问题帖 | has_answers | hbghlyj | 2020-08-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019108 | MSE-3793239

Can composition of integer polynomial and rational polynomial with a non-integer coefficient result in integer polynomial?

Can we find two polynomials $p(x)$ and $q(x)$, where $p(x)$ is a non-constant monic polynomial over integers and $q(x)$ is a monic polynomial over rationals with at least one non-integer coefficient, such that their composition $p(q(x))$ is a polynomial over integers? If not, how to prove it? For example let $q(x)=x^2+\frac{1}{2}x+1$ and $p(x)=x^3+a_2x^2+a_1x+a_0$, then $p(q(x))=x^6+\frac{3}{2}x^5+\dots$, so no matter what integers a_i we choose, the resulting polynomial will have a non-integer coefficient. The monic condition is important, since otherwise we could multiply $p(x)$ with such integer that would guarantee all coefficients to be integers. I've tried to look at the coefficient in composition for general polynomials, which I believe should follow this formula: \begin{align} [x^r]p(q(x))=\sum_{k_1+2k_2+\dots+mk_m=r}\sum_{k_0=0}^{n-(k_1+\dots+k_m)}\binom{k_0+k_1+\dots+k_m}{k_0,k_1,\dots,k_m}a_{k_0+k_1+\dots+k_m}\left(\prod_{j=0}^{m}b_j^{k_j}\right) \end{align} (here a_i and b_i are the coefficients of $p(x)$ and $q(x)$ with degrees n and m, respectively). However it is not at all clear on which coefficient to focus to prove it will give the non-integer number. This arose when trying to solve the Infinitely many solutions leads to existence of a polynomial, but it seems interesting enough by itself.

论坛问题帖 | has_answers | Sil | 2020-08-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019109 | MSE-3797253

Is Gromov's conjecture about optimal Betti number estimate true for compact manifolds?

Gromov's conjecture state that: Conjecture: Any complete manifold (M, g) with $\sec\geq 0$ and for any field $\Bbb F$ of coefficients satisfies $$\sum_{i=0}^n b_i(M,\Bbb F)\leq 2^n.$$ Is Gromov's conjecture true for compact non-negatively curved manifolds?

论坛问题帖 | has_answers | C.F.G | 2020-08-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019110 | MSE-3798002

Is it true that $\sum_{i=1}^n (nGx_i^{G} + G^{x_i}) \ge n^2G + G^2n$, for all $x_i>0$, where $G=\prod_{j=1}^nx_j$?

Source: https://math.stackexchange.com/questions/3688041/sum-i-1n-ngx-ig-gx-i-ge-n2g-g2n-for-all-x-i0-where-g Prove or disprove that, for all $x_1,x_2,\ldots,x_n>0$, it holds that $$\sum_{i=1}^n (nGx_i^{G} + G^{x_i}) \ge n^2G + G^2n, \space \space \space \text{where} \space \space \space G=\prod_{j=1}^nx_j.$$ The case $n=2$ is equlvalent to $$2\sqrt{xy}\left(x^{\sqrt{xy}}+y^{\sqrt{xy}}\right)+\left(\sqrt{xy}^{x}+\sqrt{xy}^y\right)\geq 4\sqrt{xy}+2xy\,.$$ We will show that the inequality above is true at least when $xy\geq 1$. By AM-GM, $$x^{\sqrt{xy}}+y^{\sqrt{xy}}\geq 2\,\sqrt{x^{\sqrt{xy}}y^{\sqrt{xy}}}=2\,\sqrt{xy}^{\sqrt{xy}}\geq 2\,\sqrt{xy}\,.$$ Also by AM-GM, $$\sqrt{xy}^x+\sqrt{xy}^y\geq 2\,\sqrt{\sqrt{xy}^x\sqrt{xy}^y}=2\,\sqrt{xy}^{\frac{x+y}{2}}\geq 2\,\sqrt{xy}^{\sqrt{xy}}\geq 2\,\sqrt{xy}\,.$$ Thus, $$\begin{align}2\sqrt{xy}\left(x^{\sqrt{xy}}+y^{\sqrt{xy}}\right)+\left(\sqrt{xy}^{x}+\sqrt{xy}^y\right)&\geq 2\sqrt{xy}(2\sqrt{xy})+(2\sqrt{xy})\\&=4xy+2\sqrt{xy}\geq 4\sqrt{xy}+2xy\,.\end{align}$$

论坛问题帖 | has_answers | Batominovski | 2020-08-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019111 | MSE-3839696

Van der Pol's identity for the sum of divisors and a quartic polynomial equation for odd perfect numbers

In Touchard (1953) it is mentioned that the sum of divisors $\sigma(n)$ satisfies the following recurrence relation ($n>1$): $$n^2(n-1) = \frac{6}{\sigma(n)} \sum_{k=1}^{n-1}(3n^2-10k^2)\sigma(k)\sigma(n-k)$$ Substituting in this equation an odd perfect number $n = \sigma(n)/2$ we find that it is a root of the quartic polynomial: $$ \begin{align} f(x) &= x^4-x^3+12a_0x^2-60a_1x+60a_2 \\ &= (x^3+(n-1)x^2+(12a_0+n^2-n)x+12a_0n+n^3-n^2-60a_1)(x-n) \end{align} $$ where $$a_i = \sum_{k=1}^{\frac{n-1}{2}}k^i \sigma(k)\sigma(n-k),\text{ } i=0,1,2$$ We have: $$n^3(n-1) = 3 \sum_{k=1}^{n-1} (3n^2-10k^2)\sigma(k) \sigma(n-k)$$ and by symmetry of $\sigma(k)\sigma(n-k)$ in $k,n-k$ we get: $$= 3 \sum_{k=1}^{(n-1)/2} (6n^2-10k^2-10(n-k)^2)\sigma(k) \sigma(n-k)$$ which might be simplified to: $$=-12n^2a_0 -60a_2+60na_1$$ which proves that the odd perfect number n is a root of the polynomial $f(x)$ above. However, computations with Sagemath (it takes a few seconds to do the computation) seem to suggest, that for an odd number m, we have $$f(m) > 0, \text{for } m>1 \text{ odd}$$ This observation would maybe prove, that there are no odd perfect numbers. I am aware that $\sigma(k)\sigma(n-k)$ is kind of like a black box to handle, but it would be nice, if maybe for some special cases where m is odd the last inequality could be proven. It might be noticed that it seems like for $n>1$, odd the polynomial $f(x)$ is irreducible over the rational numbers. Here are the first polynomials: n, f(n,t), f(n,n)>0? 3 t^4 - t^3 + 36*t^2 - 180*t + 180 f(n,n)>0 ? True 5 t^4 - t^3 + 228*t^2 - 1860*t + 3300 f(n,n)>0 ? True 7 t^4 - t^3 + 696*t^2 - 7920*t + 20160 f(n,n)>0 ? True 9 t^4 - t^3 + 1548*t^2 - 22500*t + 72900 f(n,n)>0 ? True 11 t^4 - t^3 + 2940*t^2 - 51600*t + 204600 f(n,n)>0 ? True 13 t^4 - t^3 + 4956*t^2 - 102360*t + 478920 f(n,n)>0 ? True 15 t^4 - t^3 + 7752*t^2 - 185760*t + 1004400 f(n,n)>0 ? True 17 t^4 - t^3 + 11376*t^2 - 307620*t + 1900260 f(n,n)>0 ? True 19 t^4 - t^3 + 16020*t^2 - 481980*t + 3309420 f(n,n)>0 ? True 21 t^4 - t^3 + 22080*t^2 - 735120*t + 5559120 f(n,n)>0 ? True Is there a irreducibility criterion which might be applied to all ($n=3,\ldots,21$) of these example polynomials? Thanks for your help! (Originally asked on MO: https://mathoverflow.net/questions/372476/van-der-pols-identity-for-the-sum-of-divisors-and-a-quartic-polynomial-equation) Edit: The odd perfect number is also a root of the polynomial: $$g(x) = x^4-x^3-9A_0x^2+30A_2$$ where $$A_i = \sum_{k=1}^{n-1}k^i \sigma(k)\sigma(n-k),i=0,2$$ But numerical computations suggest, that: $$\gcd(f(x),g(x))=1$$ which would lead to a contradiction. Can this last claim about the $\gcd$ be proven? An answer to the question: It would be nice if someone can spot the error in this answer, otherwise it would follow that there are no odd perfect numbers...: Let $$A_i =

\sum_{k=1}^{n-1} k^i \sigma(k)\sigma(n-k),\text{ } i=0,1,2,3$$. Using the symmetry of $\sigma(k)\sigma(n-k)$ in n,k for A_i we get: $$A_0 = 2a_0$$ $$A_1 = na_0$$ $$A_2 = a_0n^2-2a_1n+2a_2$$ $$A_3 = n^3a_0-3n^2a_1+3na_2$$ On the other hand we can evaluate A_i with Eisenstein series (https://mathoverflow.net/questions/372766/is-there-a-similar-formula-like-ramanunjans-eisenstein-series-identity-for-su): $$A_0 = \tfrac5{12}\sigma_3(n)-\tfrac12n\sigma(n)+\tfrac{\sigma(n)}{12} = \tfrac5{12}\sigma_3(n)-\tfrac12n \cdot 2n +\tfrac{2n}{12}$$ $$A_1 = \frac{5n}{24}\sigma_3(n) - \frac{6n^2-n}{24}\sigma(n) = \frac{5n}{24}\sigma_3(n) - \frac{6n^2-n}{24}2n$$ $$A_2 = \frac{n^2}{8}\sigma_3(n) - \frac{4n^3-n^2}{24}\sigma(n) = \frac{n^2}{8}\sigma_3(n) - \frac{4n^3-n^2}{24}2n$$ $$A_3 = \frac{n^3}{12}\sigma_3(n) - \frac{3n^4-n^3}{24}\sigma(n) = \frac{n^3}{12}\sigma_3(n) - \frac{3n^4-n^3}{24}2n$$ We can solve the set of equations in a_i,A_i for a_i and get: $$a_0 = 1/2 A_0$$ $$a_2=\frac{A_{2}-nA_1}{2} +na_1$$ We can further substitute for A_i the last equations in $\sigma_3(n)$ and then substitute these a_i in $f(x)$. We get (here are the calculations done with Sagemath, I hope there is no bug in this): $$0 = f(n) = n^4 + 6A_0n^2 - n^3 + 30A_1n + 30A_2 $$ $$ = -5/2(12n^2 - 5\sigma_3(n) - 2n)n^2$$ Hence: $$0 = 12 \, n^{2} - 2 \, n - 5 \, \sigma_{3}(n)$$ But $\sigma_3(n)>n^3+1$ and we get the contradiction for $n>1$: $$5(n^3+1) < 5 \sigma_3(n)= 12n^2-2n < 5(n^3+1) $$

论坛问题帖 | no_answers_at_retrieval | Orges Leka | 2020-09-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019112 | MSE-3857649

Nice lemma regarding factorials and quadratic residues

IMPORTANT: I posted this as I thought I found a generalized result, which turned out to be false (by a trivial flaw), but a nice lemma still remains (however, this can be closed now) Lemma: If $p\equiv 1\pmod{8}$, then $$\bigg(\frac{(\frac{p-1}{4})!}{p}\bigg)=-1$$ This problem below appeared in a USA contest (I think) some time ago: Let p be a prime number and $k\in\mathbb{N}$ such that $p=8k+1$. Suppose that $$\binom{4k}{k}\equiv r\pmod{p}$$ Prove that $\sqrt{r}$ is not an integer. Now this is actually true (By the way, try the problem yourself, it is very beautiful). So under this assumptioin, lets prove the lemma Proof: When $v_2(p-1)=3$ This gives us $p\equiv 9\pmod{16}\Rightarrow p\equiv 1\pmod{8}$. Lets write $p=8k+1$. So by the above problem i just stated, we get: $$\bigg(\frac{\binom{4k}{k}}{p}\bigg)=-1$$ Lets start interpreting this: $$\bigg(\frac{\binom{4k}{k}}{p}\bigg)=-1\Leftrightarrow \bigg(\frac{(4k)!}{p}\bigg)=-\bigg(\frac{\frac{1}{k!\cdot(3k)!}}{p}\bigg)$$ Notice that $$1 =\bigg(\frac{1}{p}\bigg)=\bigg(\frac{\frac{1}{k!\cdot(3k)!}}{p}\bigg)\cdot\bigg(\frac{k!\cdot(3k)!}{p}\bigg)$$ so $$\bigg(\frac{k!\cdot(3k)!}{p}\bigg)=\frac{1}{\bigg(\frac{\frac{1}{k!\cdot(3k)!}}{p}\bigg)}=\bigg(\frac{\frac{1}{k!\cdot(3k)!}}{p}\bigg)$$ Thus, substituting in $$\bigg(\frac{(4k)!}{p}\bigg)=-\bigg(\frac{\frac{1}{k!\cdot(3k)!}}{p}\bigg)$$ we have $$\bigg(\frac{(4k)!}{p}\bigg)=-\bigg(\frac{k!\cdot(3k)!}{p}\bigg)\Leftrightarrow\bigg(\frac{(3k)!}{p}\bigg)\cdot\bigg(\frac{(3k+1)(3k+2)...4k}{p}\bigg)=-\bigg(\frac{(3k)!}{p}\bigg)\cdot\bigg(\frac{k!}{p}\bigg)$$ so we get $$\bigg(\frac{(3k+1)(3k+2)...4k}{p}\bigg)=-\bigg(\frac{k!}{p}\bigg)$$ which is equivalent to $$\big((3k+1)(3k+2)...4k\big)^{\frac{p-1}{2}}\equiv-\big(k!\big)^{\frac{p-1}{2}}\pmod{p}$$ Multiplying both sides by $2^{\frac{p-1}{2}}$, we have: $$\big((6k+2)(6k+4)...8k\big)^{\frac{p-1}{2}}\equiv-\big(2\cdot4\cdot6...\cdot2k\big)^{\frac{p-1}{2}}\pmod{p}$$ But $$\big((6k+2)(6k+4)...8k\big)^{\frac{p-1}{2}}\equiv\big(-(2k-1)\cdot-(2k-3)...\cdot-1\big)^{\frac{p-1}{2}}\equiv \big((-1)^k\cdot1\cdot3...\cdot(2k-1)\big)^{\frac{p-1}{2}}\pmod{p}$$ $\frac{p-1}{2}=4k$, so $$\big((-1)^k\cdot1\cdot3...\cdot(2k-1)\big)^{\frac{p-1}{2}}\equiv\big(1\cdot3...\cdot(2k-1)\big)^{\frac{p-1}{2}}\pmod{p}$$ Substituting in $$\big((6k+2)(6k+4)...8k\big)^{\frac{p-1}{2}}\equiv-\big(2\cdot4\cdot6...\cdot2k\big)^{\frac{p-1}{2}}\pmod{p}$$ we get the following: $$\big(1\cdot3...\cdot(2k-1)\big)^{\frac{p-1}{2}}\equiv-\big(2\cdot4\cdot6...\cdot2k\big)^{\frac{p-1}{2}}\pmod{p}$$ Finally, by multiplying both sides by $\big(2\cdot4\cdot6...\cdot2k\big)^{\frac{p-1}{2}}$, we get: $$(2k)!\equiv-1\big(2\cdot4\cdot6...\cdot2k\big)^{p-1}\equiv-1\pmod{p}$$ So we have $$\bigg(\frac{(\frac{p-1}{4})!}{p}\bigg)=\bigg(\frac{(2k)!}{p}\bigg)=-1$$ Which is exactly what we wanted to prove. Thank you!

论坛问题帖 | has_answers | user799688 | 2020-10-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019113 | MSE-3867967

Given $A\subset \mathbb{N}$, probability of a $B\subset A$, $s(B)\in\mathbb{P}$

To clarify the notation, for a finite set K, let $s(K)$ be the sum of the elements of K. My question Quite straight forward, I want to ask this: Given $A\subset\mathbb{N}$ such that A is finite, what the the probability that there exists $B\subset A$ such that $s(B)$ is a prime number. Motivation This is mainly inspired by the fact that the probability of a number, N, being a prime is $\sim\frac{1}{\log{N}}$ and the fact that in a recent post, I talked about the probability of the sum of n randomly chosen integers being divisible by p, a prime (which was later on generalized). To be honest, I have almost no knowledge in probability or advanced number theory whatsoever, so the inly background I can provide you with is this idea I got: When I first thought of the probability of a number being a prime, I had this train of thought (more or less, this is a summary): So i have $\pi(n)$ favorable cases and n total cases, as n goes to infinity, so the probability would be: $$\lim_{n\to\infty}\frac{\pi(n)}{n}=\lim_{n\to\infty}\frac{1}{\log{n}}$$ But that gives us 0, which is a contradiction. So then I thought of something very similar but simpler: instead of having $\pi(m)$ favorable cases and m total cases as m gets arbitrarely big, we can just have n favorable cases and p_n total cases, because $n\leq p_n$. So the probability now is $$\sim\frac{n}{p_n}\sim\frac{n}{n\log{n}}=\frac{1}{\log{n}}$$ But after doing this, I thought I could use an almost identical approach to see what is the probability for a number n such that $n+c$ is a prime (for a given constant c), but I am not sure this actually works. However, I am not sure how to generalize this to sums or sets. To sum up, I would appreciate any help or ideas. Thank you for reading.

论坛问题帖 | no_answers_at_retrieval | user799688 | 2020-10-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019114 | MSE-3870551

Carmichael number divisible by a given positive integer

Prove or disprove this conjecture : If $k>2$ is an integer with $\gcd(k,\varphi(k))=1$, then there is a Carmichael number N with $k\mid N$ The condition $\gcd(k,\varphi(k))=1$ is necessary. Otherwise, k would not be squarefree and therefore also N , which is impossible. Or there would be prime factors p,q of k and therefore N, with $p\mid q-1$. This is also impossible since this would imply $p\mid N-1$ because of $q-1\mid N-1$ contradicting $p\mid N$ But is it also sufficient ? And if yes, is there an efficient method to construct a Carmichael number divisible by k ? A particular hard case seems to be $k=885=3\cdot 5\cdot 59$. Is there a Carmichael number divisble by 885 ?

论坛问题帖 | has_answers | Peter | 2020-10-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019115 | MSE-387215

Conjecture regarding trapping rational numbers in some special intervals

Conjecture: Let $b\in\mathbb{N}_{\geq3}$ and $\{x_i\}$ be a collection of $b-2$ rational numbers greater than 1. Does there always exist a natural number a such that for all i there exists some natural number $1\leq c\leq a$ such that: $$\displaystyle \frac{ab-1}{(c-1)b+1}\geq x_i \geq \frac{ab-1}{cb-1}\tag{1}$$ One may think of this as having a UNION of a intervals $\left[\frac{ab-1}{cb-1},\frac{ab-1}{(c-1)b+1}\right]$ on the real line and proving that, for some a, these can trap any finite discrete set of rational numbers having cardinality $b-2$. For example, Let's assume $b=10$, $x=\{\frac{12}{5},\frac{17}{5},\frac{59}{10},\frac{187}{20},15,21,\frac{67}{2},39 \}$, then (1) is true for $a=4$ and $a=5$ and therefore the intervals for $a=4,5$ contain or "trap" all the members of set x. In the diagram, the red/green regions are intervals $\left[\frac{ab-1}{cb-1},\frac{ab-1}{(c-1)b+1}\right]$ for all $c<a$. What mathematical tools can one employ to tackle this? WHAT I DID: I solved the inequality (1) for a and got $$\large x_i\left(1-\frac{2}{b}\right)+\frac{x_i+1}{b}+cx_i\geq a\geq \frac{x_i+1}{b}+cx_i\tag{2}$$ Note that $\frac{x_i+1}{b}+cx_i$ is common on both sides of the inequality. Now I'll just have to plug in the values of x_i to get $b-2$ inequalities and check for a common value of a for any c. For example, Let $x_i \in \{ \frac{101}{4},\frac{3001}{7} \}$ and $b=4$ then we have inequalities $\frac{307}{16}+\left(\frac{101}{4}\right)c_1\geq a\geq \frac{105}{16}+\left(\frac{101}{4}\right)c_1$ $\frac{911}{32}+\left(\frac{301}{8}\right)c_2\geq a\geq \frac{309}{32}+\left(\frac{301}{8}\right)c_2$ Put $c=2$ in the first inequality and $c=1$ in the second inequality to get a common $a=58$. But how should one go about to prove that there will always be some value of a which satisfies the above set of inequalities? I calculated numerically for some random b and x_i and always managed to get some a, i.e., there existed some a which trapped random x_i given some random b. A counterexample may prove the negative but I was unable to find any.

论坛问题帖 | has_answers | Maazul | 2013-05-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019116 | MSE-3873500

Is this proof regarding odd perfect numbers valid?

(Edited in response to a comment.) Here are my: QUESTIONS (1) Is this proof regarding odd perfect numbers valid, particularly the part where it says $$\dfrac{2n^2}{D(n^2)} \neq (q + 1)?$$ (2) If the proof is incorrect, how can the argument be mended to produce a valid proof? Let $N = q^k n^2$ be an odd perfect number with special/Euler prime q. (That is, $\gcd(q,n)=1$ and $q \equiv k \equiv 1 \pmod 4$.) Let $\sigma(x)$ denote the sum of the divisors of $x \in \mathbb{N}$. Define $$D(n^2) := 2n^2 - \sigma(n^2)$$ to be the deficiency of the non-Euler part n^2. Define $$I(n^2) := \dfrac{\sigma(n^2)}{n^2}$$ to be the abundancy index of n^2. Since N is perfect, then we have $$\sigma(q^k)\sigma(n^2)=\sigma(q^k n^2)=\sigma(N)=2N=2q^k n^2$$ from which it follows that $$I(N)=2 \iff I(q^k)I(n^2)=2.$$ But since q is prime, $I(q^k)$ can be rewritten as $$I(q^k) = \dfrac{q^{k+1} - 1}{q^k (q - 1)}$$ which can be bounded as follows (since $k \equiv 1 \pmod 4$ implies that $k \geq 1$) $$\frac{q+1}{q} = I(q) \leq I(q^k) < \dfrac{q^{k+1}}{q^k (q - 1)} = \frac{q}{q - 1}.$$ This implies that $$\dfrac{2(q - 1)}{q} < I(n^2) = \dfrac{2}{I(q^k)} \leq \dfrac{2q}{q + 1}$$ from which it follows that $$\dfrac{2}{q+1} \leq 2 - I(n^2) = \dfrac{D(n^2)}{n^2} < \dfrac{2}{q},$$ which means that $$q < \dfrac{2n^2}{D(n^2)} \leq (q + 1).$$ Since $2n^2$ is even and $D(n^2) = 2n^2 - \sigma(n^2)$ is always odd (since n^2 is a square), then $$\dfrac{2n^2}{D(n^2)}$$ cannot be an integer. This implies that $$\dfrac{2n^2}{D(n^2)} \neq (q + 1)$$ which implies that $$\dfrac{n^2}{D(n^2)} \neq \dfrac{q+1}{2}$$ and $$2 - I(n^2) = \dfrac{D(n^2)}{n^2} \neq \dfrac{2}{q+1}.$$ Finally, we obtain $$I(n^2) \neq \bigg(2 - \dfrac{2}{q+1}\bigg) = \dfrac{2q}{q+1}$$ which implies that $$I(q^k) = \dfrac{2}{I(n^2)} \neq \dfrac{q+1}{q},$$ thus resulting in $$k \neq 1.$$

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2020-10-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019117 | MSE-3879709

Sketch of proof concerning a conjecture of Vasile Cirtoaje : $(1-x)^{(2x)^k}+x^{(2(1-x))^k}\leq 1$ with $k\geq1$ and $0<x\leq \frac{1}{2}$

Claim Prove that [1]: Let $0<x\leq 0.5$ and $k\geq 1$ a real number then we have : $$h(k)=(1-x)^{(2x)^k}+x^{(2(1-x))^k}\leq 1$$ Sketch of proof : The case $k\geq 1+\alpha$: We define α as : $$\lim_{x\to 0.5^-}=\frac{\ln\Big(2(2^{2x}x^2(1-x)2)\Big)}{\ln(2x)}=\frac{\ln\Big(2(1-(2^{2x}x^2(1-x)2))\Big)}{\ln(2(1-x))}=\alpha=1.69315\cdots$$ We have : $$x^{2(1-x)}\leq2^{2x}(1-x)2x^2$$ And : $$(1-x)^{2x}\leq 1-2^{2x}(1-x)2x^2$$ For a sketch of proof see Refinements of the inequality $f(x)=x^{2(1-x)}+(1-x)^{2x}\leq 1$ for $0<x<0.5$ So we have : $$x^{(2(1-x))^k}\leq f(x)^{(2(1-x))^{k-1}}=(2^{2x}(1-x)2x^2)^{(2(1-x))^{k-1}}$$ And : $$(1-x)^{(2x)^k}\leq (1-2^{2x}(1-x)2x^2)^{(2x)^{k-1}}$$ Now we want $y\geq 1$ a real number : $$(2x)^{k-1}\geq (2f(x))^{y}$$ And $$(2(1-x))^{k-1}\geq (2(1-f(x)))^{y}$$ Solving this we have $$\frac{\ln\Big(2(1-(2^{2x}x^2(1-x)2))\Big)}{\ln(2(1-x))}\leq \frac{k-1}{y}\leq \frac{\ln\Big(2(2^{2x}x^2(1-x)2)\Big)}{\ln(2x)}\quad (1)$$ The constraint (1) is honoured if :$$\alpha=\frac{k-1}{y}$$ So we have : $$x^{(2(1-x))^k}\leq

(f(x))^{(2(1-f(x))^{\frac{k-1}{\alpha}} }\quad (2)$$ And : $$(1-x)^{(2x)^k}\leq (1-f(x))^{(2f(x))^{\frac{k-1}{\alpha}}}\quad (3)$$ Adding (2) and (3) we have : $$h(k)\leq (1-f(x))^{(2f(x))^{\frac{k-1}{\alpha}}}+(f(x))^{(2(1-f(x))^{\frac{k-1}{\alpha}}}$$ We can repeat the reasoning p times ($p\geq 1$ a natural number) and the last exponent becomes : $$1\leq \frac{k-1}{\alpha^p}\leq \alpha$$ It's true because we can always choose k such that : $$\alpha^p\leq k \leq \alpha^{p+1}$$ So we reduce this case to the case $1\leq k\leq 1+\alpha $ Case $k\leq 1+\alpha $ Using Bernoulli's inequality and derivatives we have $1\leq k\leq 2$ and $0<x\leq \frac{1}{4}$ $$(1-x)^{(2x)^k}\leq 1-x(2x)^k$$ And $$x^{(2(1-x))^k}\leq x(2x)^k$$ In the same idea we have $2\leq k\leq 1+\alpha$ and $0<x\leq \frac{1}{3}$ $$(1-x)^{(2x)^k}\leq 1-x(2x)^k$$ And $$x^{(2(1-x))^k}\leq x(2x)^k$$ Now to finalize the sketch of proof we have : $h(k)$ is decreasing for $1\leq k\leq 2$ and $ \frac{1}{4}\leq x\leq 0.5\quad (4)$ $h(k)$ is decreasing for $2\leq k\leq 1+\alpha$ and $ \frac{1}{3}\leq x\leq 0.5\quad (5)$ Questions : It is good and coherent ? How to show (4) and (5)? Thanks in advance ! Regards Max [1] Vasile Cirtoaje, "Proofs of three open inequalities with power-exponential functions", The Journal of Nonlinear Sciences and its Applications (2011), Volume: 4, Issue: 2, page 130-137. https://eudml.org/doc/223938 https://link.springer.com/article/10.1186/1029-242X-2013-468

论坛问题帖 | has_answers | Barackouda | 2020-10-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019118 | MSE-3891195

Two homotopy equivalent closed n-manifolds are homeomorphic

I heard somewhere that there is a wrong conjecture that states: Two homotopy equivalent closed n-manifolds ($n\geq 3$) are homeomorphic. and the lens spaces are the first counterexample to this conjecture. I want to know the author of this wrong conjecture, and a reference for it. I am also curious about other counterexamples? (Here is a related question)

论坛问题帖 | has_answers | C.F.G | 2020-11-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019119 | MSE-3897185

Is $X = (n-1)^n + n$ always composite for $n \geq 4 \in \Bbb Z$?

$$\textit{Proposition}: \text{Given} \ X = (n-1)^n + n,\ \forall \ n \in \Bbb Z, \ n \geq 4, X \ \text{is composite.}$$ This came up sort of at random on a mailing list and it's jammed its way into being stuck in my brain. It's easy to test that it's true for the first few hundred thousand integers, suggesting it's likely true. There are a few things that are easy to prove: X is always odd: if $(n-1)^n$ is odd, n is even and vice versa. $(n - 1) \equiv -1 \pmod n $. Since $n \equiv 0 \pmod n$, we can see that $X \equiv (-1)^n \pmod n $. Therefore, $X \equiv -1 \pmod n$ for odd n, and $X \equiv 1 \pmod n $ for even n. If $n \equiv 2 \pmod 3$, then $n-1 \equiv 1 \pmod 3$, so $(n-1)^n \equiv 1 \pmod 3$. If we add $n \equiv 2 \pmod 3$, we have $0 \pmod 3$, so $3 \mid X$. So a third of possible values for n are easily shown to be composite, namely $n = 3m + 2$. If $n \equiv \{4,13\} \pmod {20}$, then $5 \mid X$, as follows: If $n \equiv 4 \pmod {20}$, then $4 \mid n$, $n \equiv 4 \pmod{10}$, and $(n - 1) \equiv 3 \pmod{10}$. Then we have $4^3+4 \equiv 5 \pmod{10}$ If $n \equiv 13 \pmod {20}$, then $4 \mid (n-1)$, $(n-1) \equiv 12 \pmod{20}$, and $(n-1) \equiv 2 \pmod {10}$. Additionally, $n \equiv 1 \pmod 4$ and $n \equiv 3 \pmod{10}$. Therefore, in decimal, $(10k + 2)^{4m+1} \equiv 2 \pmod{10}$. Adding $n \equiv 3 \pmod{10}$ gives $X \equiv 5 \pmod{10}$. That takes care of 40% of the integers. But (1) I feel like we ought not to have to go through the integers modulus by modulus, and (2) well, we have a lot of integers left. Some other interesting tidbits that may or may not help: For n even and $n \not\equiv 2 \pmod 3$, X is often divisible by $(n+1)$, or by p if $(n+1) = p^2$, p prime. Up to $n = 72$, this fails for $n = 34, 54, 64$. For $n \not\equiv 2 \pmod 3$, X is sometimes divisible by $(2n-1)$. Again up to $n = 72$, this is true for even $n = 6, 10, 22, 30, 34, 42, 54, 66, 70$. (Interesting, for all of those, $n \equiv 2 \pmod 4$. Just noticing that.) It's true for odd $n = 9, 21, 37, 45, 49, 57$... which are all $n \equiv 1 \pmod 4$. Of course, being sometimes divisible by those numbers doesn't tell us why they're divisible by them. I tried using partial binomial expansions (learning about Stirling numbers along the way) but couldn't get anything useful. Has anyone seen or worked with this proposition before? Is it proven and I just can't find it anywhere?

论坛问题帖 | has_answers | Eric Snyder | 2020-11-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019120 | MSE-3901694

A new method for finding a solution (when they exist) to $x^2 = a \pmod p$?

For prime $p \ge 2$ define a set $S \subset \Bbb N$ by $\quad S = \{t^2 \mid t \ge 1 \land t^2 \lt p\}$ Let a be an integer satisfying $\quad 1 \le a \lt p$ Conjecture: The equation $x^2 \equiv a \pmod p$ has a solution if and only if there exists an integer s belonging to S such that, when applying Euclidean division by setting the dividend to sa and the divisor to p, the remainder also belongs to S. When a solution exists, it can be calculated by solving the corresponding $tx \equiv 1 \pmod{p}$ equation. If this is known please point to some references or supply a proof. Otherwise please help in finding a counter-example; see also Jaap Scherphuis' comments. It is certainly possible that S has to be expanded, up, to say, $\; S = \{t^2 \mid 1 \le t \lt p\}$ when forming the divisor sa, but the 'tighter the better' when formulating algorithms for this idea. But we really want the aesthetically pleasing duality statement, $\quad S^\text{*} (\text{residue side of } sa) = S (\text{multiply side of } sa$). Example 1: $x^2 = 2 \pmod{17}$. Here $S = \{1,4,9,16\}$. Dividing $4 \cdot 2:\quad$ $8 = 0\cdot 17 + 8$ and $8 \notin S$ Dividing $9 \cdot 2:\quad$ $18 = 1\cdot 17 + 1$ and $1 \in S$ We have $[3]^{-1} = [6]$ in the $\text{modulo-}17$ system. So $\; x^2 = 2 \pmod{17} \; \text{ iff } \; 3^2 x^2 \equiv 1 \pmod{17} \; \text{ iff } \; \large x^2 \equiv 3^{-2} \pmod{17} \; \text{ iff } \; x^2 \equiv 6^{2} \pmod{17} $ and we've found a solution, $x = 6$. Example 2: $x^2 \equiv 3\pmod {10007}$. Here is a Python program: S =[1, 4, 9, 16, 25, 36, 49, 64, 81, 100, 121, 144, 169, 196, 225, 256, 289, 324, 361, 400, 441, 484, 529, 576, 625, 676, 729, 784, 841, 900, 961, 1024, 1089, 1156, 1225, 1296, 1369, 1444, 1521, 1600, 1681, 1764, 1849, 1936, 2025, 2116, 2209, 2304, 2401, 2500, 2601, 2704, 2809, 2916, 3025, 3136, 3249, 3364, 3481, 3600, 3721, 3844, 3969, 4096, 4225, 4356, 4489, 4624, 4761, 4900, 5041, 5184, 5329, 5476, 5625, 5776, 5929, 6084, 6241, 6400, 6561, 6724, 6889, 7056, 7225, 7396, 7569, 7744, 7921, 8100, 8281, 8464, 8649, 8836, 9025, 9216, 9409, 9604, 9801, 10000] for m in S: r = m*3 % 10007 if r in S: print(m, m**.5, r) raise SystemExit with output $3721\; 61.0\; 1156$ We have $1156 = 34^2$. We

have $[61]^{-1} = [3281]$ in the $\text{modulo-}10007$ system. So $\; x^2 = 3 \pmod{10007} \; \text{ iff } \; 61^2 x^2 \equiv 34^2 \pmod{10007} \; \text{ iff } \; \large x^2 \equiv 61^{-2} 34^2 \pmod{10007} \; \text{ iff } \; x^2 \equiv (3281 \cdot 34)^2 \pmod{10007} \; \text{ iff } \; x^2 \equiv 1477^2 \pmod{10007}$ and we've found a solution, $x = 1477$. Example 3: $x^2 \equiv 3 \pmod{7}$. Here $S = \{1,4\}$. Since $4 \times 3 \equiv 5 \pmod{7}$ is not in S, there are no solutions. Example 4: $x^2 = 19 \pmod{127}$ $\; x^2 \equiv 19 \pmod{127} \; \text{ iff }$ $\;6^2 x^2 \equiv 6^2 \cdot 19 \pmod{127}\; \text{ iff }$ $\;6^2 x^2 \equiv 7^2 \pmod{127}$ We have $[6]^{-1} = [106]$ in the $\text{modulo-}127$ system. So, $\;x^2 \equiv 6^{-2} 7^2 \pmod{127}\; \text{ iff }$ $\;x^2 \equiv \bigr(6^{-1} 7^1 \bigr)^2 \pmod{127}\; \text{ iff }$ $\;x^2 \equiv \bigr(106 \cdot 7 \bigr)^2 \pmod{127}\; \text{ iff }$ $\;x^2 \equiv 107^2 \pmod{127}$ and we see that $x \equiv 107 \pmod{127}$ is a solution. My Work After seeing lonza leggiera's counter-example answer to my question here, I went back to the drawing board and this kind of just 'popped out'. In this question I wanted to focus on the extensively studied modulo prime systems, but if it works here further research would be warranted.

论坛问题帖 | no_answers_at_retrieval | CopyPasteIt | 2020-11-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019121 | MSE-3902896

On the nearest-square function - Part 2 and the quantity $m^2 - p^k$ where $p^k m^2$ is an odd perfect number

(Note: This question has been cross-posted to MO.) This question is an offshoot of this earlier one and this other question. Let $n = p^k m^2$ be an odd perfect number with special prime p satisfying $p \equiv k \equiv 1 \pmod 4$ and $\gcd(p,m)=1$. It was conjectured in Dris (2008) and Dris (2012) that the inequality $p^k < m$ holds. Brown (2016) showed that the Dris Conjecture (that $p^k < m$) holds in many cases. It is trivial to show that $m^2 - p^k \equiv 0 \pmod 4$. This means that $m^2 - p^k = 4z$, where it is known that $4z \geq {10}^{375}$. (See this MSE question and answer, where the case $m < p^k$ is considered.) Note that if $p^k < m$, then $$m^2 - p^k > m^2 - m = m(m - 1),$$ and that $${10}^{1500} < n = p^k m^2 < m^3$$ where the lower bound for the magnitude of the odd perfect number n is due to Ochem and Rao (2012). This results in a larger lower bound for $m^2 - p^k$. Therefore, unconditionally, we have $$m^2 - p^k \geq {10}^{375}.$$ We now endeavor to disprove the Dris Conjecture. Consider the following sample proof argument: Theorem If $n = p^k m^2$ is an odd perfect number satisfying $m^2 - p^k = 8$, then $m < p^k$. Proof Let $p^k m^2$ be an odd perfect number satisfying $m^2 - p^k = 8$. Then $$(m + 3)(m - 3) = m^2 - 9 = p^k - 1.$$ This implies that $(m + 3) \mid (p^k - 1)$, from which it follows that $$m < m + 3 \leq p^k - 1 < p^k.$$ We therefore conclude that $m < p^k$. QED So now consider the equation $m^2 - p^k = 4z$. Following our proof strategy, and the formula in the accepted answer to the first hyperlinked question, we have: $$m^2 - \bigg(\lfloor{\sqrt{m^2 - p^k} + \frac{1}{2}}\rfloor\bigg)^2 = p^k + \Bigg(4z - \bigg(\lfloor{\sqrt{m^2 - p^k} + \frac{1}{2}}\rfloor\bigg)^2\Bigg).$$ So the only remaining question now is whether it could be proved that $$\Bigg(4z - \bigg(\lfloor{\sqrt{m^2 - p^k} + \frac{1}{2}}\rfloor\bigg)^2\Bigg) = -y < 0$$ for some positive integer y? In other words, is it possible to prove that it is always the case that $$\Bigg((m^2 - p^k) - \bigg(\lfloor{\sqrt{m^2 - p^k} + \frac{1}{2}}\rfloor\bigg)^2\Bigg) < 0,$$ if $n = p^k m^2$ is an odd perfect number with special prime p? (Additionally, note that it is known that $m^2 - p^k$ is not a square, if $p^k m^2$ is an OPN with special prime p. See this MSE question and the answer contained therein.) If so, it would follow that $$\Bigg(m + \lfloor{\sqrt{m^2 - p^k} + \frac{1}{2}}\rfloor\Bigg)\Bigg(m - \lfloor{\sqrt{m^2 - p^k} + \frac{1}{2}}\rfloor\Bigg) = p^k - y$$ which would imply that $$\Bigg(m + \lfloor{\sqrt{m^2 - p^k} + \frac{1}{2}}\rfloor\Bigg) \mid (p^k - y)$$ from which it follows that $$m < \Bigg(m + \lfloor{\sqrt{m^2 - p^k} + \frac{1}{2}}\rfloor\Bigg) \leq p^k - y < p^k.$$ Update (November 11, 2020 - 10:21 PM Manila time) Please check out the recently posted answer for a minor adjustment to the logic that should make the general proof argument work.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2020-11-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019122 | MSE-3907167

Conjecture about :$\frac{a}{\sqrt[4]{8(b^4+c^4)}}+\frac{b}{a+c}+\frac{c}{a+b}\geq\sqrt{\frac94+\frac32 \cdot \frac{(a-b)^2}{ab+bc+ca}}$

It's a conjecture . I combine two inequalities : Stronger than Nesbitt inequality and Stronger than Nesbitt's inequality $\frac{a}{\sqrt[4]{8(b^4+c^4)}}+\frac{b}{a+c}+\frac{c}{a+b}\geq\frac{3}{2}$ So now the conjecture : Let $a,b,c>0$ such that $a\geq b \geq c$ then we have : $$\frac{a}{\sqrt[4]{8(b^4+c^4)}}+\frac{b}{a+c}+\frac{c}{a+b}\geq\sqrt{\frac94+\frac32 \cdot \frac{(a-b)^2}{ab+bc+ca}}$$ I cannot find any counter-example until now . To prove it I have tried the same method as here Prove this refinement of Nesbitt's inequality based on another without success . My question : Have you an element of proof or an counter-example ? Thanks in advance !! Max .

论坛问题帖 | has_answers | Barackouda | 2020-11-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019123 | MSE-3910671

Can the Infinite Product $\prod_{n=1}^\infty \frac{a_n}{a_n + 1}$ Converge or Diverge to Zero?

Can the infinite product $\prod_{n=1}^\infty \frac{a_n}{a_n + 1}$ converge or diverge to zero when the sequence a_n is monotonically increasing? I conjecture that this is impossible given that $\lim_{n\to\infty}\frac{a_n}{a_n+1}$ by the monotonic increase of the a_n. However, I don't know of a rigorous proof or disproof of this, and analyzing convergence of infinite series and infinite products is not something I am very familiar with. If someone could provide a proof or disproof of this conjecture (preferably with a reference to a known theorem), or explain what additional restrictions need to be imposed on a_n for the conjecture to hold, I would be very appreciative!

论坛问题帖 | has_answers | Math Rules | 2020-11-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019124 | MSE-391360

Integrate $\int_0^1 \Bigl(\mathrm{Li}_2\bigl(\frac{x-1}{x}\bigr)\Bigr)^2 \mathrm{d}x$

This question has some relationship to this integral: Let Li_2 be the dilogarithm. Then, numerically, $$ \int_0^1 \Bigl(\mathrm{Li}_2\bigl(\frac{x-1}{x}\bigr)\Bigr)^2 \mathrm{d}x = \frac{17}{180}\pi^4, $$ as can be checked by Mathematica code like NIntegrate[PolyLog[2, (x-1)/x]^2, {x,0,1}, WorkingPrecision -> 100] and N[17/180 * Pi^4, 100] both of which output the same digits 9.19974748654467462788603142059992717358538 3 ... Is this a known identity? Some related results appear in Freitas and Laurenzi, but without this specific argument in the dilogarithm.

论坛问题帖 | has_answers | heiner | 2013-05-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019125 | MSE-3914281

Conjecture: $a^{(2(1-a))^{\frac{1}{n}}}+b^{(2(1-b))^{\frac{1}{n}}}+c^{(2(1-c))^{\frac{1}{n}}}+\frac{c}{n}\leq 1$

Hi it's a conjecture inspired by a work of Vasile Cirtoaje : Claim : Let $0.5\geq a \geq b \geq c\geq 0$ such that $a+b+c=1$ and $n\geq 1$ a natural number then we have : $$a^{(2(1-a))^{\frac{1}{n}}}+b^{(2(1-b))^{\frac{1}{n}}}+c^{(2(1-c))^{\frac{1}{n}}}+\frac{c}{n}\leq 1$$ To prove it I have tried Bernoulli's inequality as we have : $$a^{(2(1-a))^{\frac{1}{n}}}\leq 1-(1-a)(2(1-a))^{\frac{1}{n}}$$ But it's not enough and wrong. I have tested for $n=1,2,3,4,5$.Moreover the equality case is $a=b=0.5$. Case $n=1$: As in my previous posts we have the inequality $x\in[0,0.5]$: $$x^{2(1-x)}\leq 2^{2x+1}x^2(1-x)$$ applying this for each variable we want to show : $$2^{2a+1}a^2(1-a)+2^{2b+1}b^2(1-b)+2^{2(1-(a+b))+1}(1-(a+b))^2(a+b)+1-(a+b)\leq 1$$ Now by Bernoulli's inequality we have: $$2^{2x+1}\leq 2(1+2x)$$ Remains to show : $$2(1+2a)a^2(1-a)+2(1+2b)b^2(1-b)+2(1+2(1-(a+b)))(1-(a+b))^2(a+b)+1-(a+b)\leq 1$$ Wich is wrong because we have an equality case at $b=c=0.25$ and $a=0.5$! My question : Have you a counter-example or later (if it's true) a proof ? Thanks in advance !

论坛问题帖 | has_answers | Barackouda | 2020-11-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019126 | MSE-3948617

Bézout giving some spare coins to Frobenius? (the Coin Problem)

Let a and b be relatively prime positive integers and write the two minimal Bézout's identity pairs, $\tag 1 (s)\,a + (t)\,b = 1$ $\tag 2 (u)\,a + \;(v)\,b = 1$ where the extended Euclidean algorithm produces one of them (see this discussion on the structure of solutions). Set $\quad \; \; x = s - 1$ if $s \gt 0$, else set $x = u-1$ Set $\quad \; \; y = t - 1$ if $t \gt 0$, else set $y = v-1$ Let $n = ab - a - b$ be the solution to the Frobenius coin problem for a and b. Conjecture: The numbers x and y derived from the extended Euclidean algorithm satisfy the equation $n + 1 = (a-1)(b-1) = (x)\,a + (y)\,b$ My Work I noticed this relationship occured while working on this and that (not spelled out in this answer). Here is another example. Let $a = 227$ and $b =2011$ and write $\tag{Z1} (-691)\,227 + (78)\,2011 = 1$ $\tag{Z2} (1320)\,227 + \;(-149)\,2011 = 1$ and $x = 1319$ and $y = 77$. The statement $ (1319)\,227 + \;(77)\,2011 = 227\cdot2011 - 227 - 2011 + 1$ is true. If the conjecture is true one could use the connection to calculate Bézout's coefficients (in the coprime setting) with new algorithms. Example: Use an ad-hoc method (algorithm?) to solve for x and y in the equation $\quad (x)\,227 + \;(y)\,2011 = 227\cdot2011 - 227 - 2011 + 1$ Examining $\quad 227\cdot2011 - 227 - 2011 + 1 = (2011)\, 227 - 2237$ the problem is reduced to solving $\quad 227 k \equiv 2237 \pmod{2011}$ That gives $k \equiv 692 \pmod{2011}$ and $\quad (2011)\, 227 - 2237 =$ $\quad (2011 -692)\, 227 + 692\cdot227 - 2237 =$ $\quad (1319)\, 227 + 154847$ Now use our 'best' algorithm for dividing (exactly) 2011 into 154847, $\quad \large \frac{154847}{2011} = 77$ so $\quad (1319)\, 227 + 154847 =$ $\quad (1319)\, 227 + (77) \, 2011 $ and we can now easily calculate Bézout's coefficients by solving the linear equations $\quad (z)\,227 + (78)\,2011 = 1$ and $\quad (1320)\,227 + (z)\,2011 = 1$

论坛问题帖 | has_answers | CopyPasteIt | 2020-12-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019127 | MSE-3951341

Conjecture : A lower bound for the prime counting function .

Well it take my a little bit of time to find it but now I think that my conjecture is ready . Conjecture : Let $n\geq 100$ and then define the sum : $$S(n)=\sum_{k=1}^{n}\frac{1}{\operatorname{argtanh}\left(\frac{k}{k+1}\right)}$$ Then we have : $$\lceil{S(n)}\rceil\leq 2\pi(n)$$ Where we see the ceiling function and the prime counting function .Nicely the equality case is $n=100$. I have checked my conjecture with wolfram alpha up to $n=1000$ and extra values of n up to $n=1000000$ Obviously I'm inspired by the prime number theorem and it's not a chance if I choose $\operatorname{argtanh}$ in my conjecture . I have several questions: Well first I would like to know if it's true for larger n . Is my conjecture equivalent to other conjecture (stronger or weaker conjecture) . Can we improve the conjecture by adding (by example) a fixed exponent ? What my conjecture involves ? If it's not true for small n can we take a larger n to start with ? Thanks!

论坛问题帖 | has_answers | Barackouda | 2020-12-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019128 | MSE-3953462

Can you prove this number theory conjecture?

I think I can prove the following (using a conjecture): $$ 2n (\sum_{i=2}^n p_i) \geq p_n(p_n+1)/2 - 6$$ where p_i is the ith prime. Can one prove this without any conjecture?

论坛问题帖 | no_answers_at_retrieval | More Anonymous | 2020-12-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019129 | MSE-3966201

Conjecture regarding the sum of prime factors of $x!$ and $(x-1)!$?

I think using dodgy means I can show the following: Let $\lambda(x)$ be the sum of primes in x. For example: $$ \lambda(2) = 2$$ $$ \lambda(4) = 2 + 2 = 4$$ $$ \lambda(6) = 3 + 2 = 5$$ Then for large n: $$ \frac{\lambda (n!)}{\lambda ((n-1)!)} \sim \frac{\ln (n!)}{\ln (n-1)!}$$ Or more precisely substracting 1: $$ \frac{\lambda (n)}{\lambda ((n-1)!)} \sim \frac{\ln (n)}{\ln (n-1)!}$$ Or even better: $$ \frac{\lambda (n)}{\lambda ((n-1)!)} - \frac{\ln (n)}{\ln (n-1)!} = o(1)$$ Can someone prove or disprove this conjecture? For example: $$ \frac{\lambda(8!)}{\lambda(7!)} = \frac{31}{26} \approx 1.23$$ And $$ \frac{\ln(8!)}{\ln(7!)} \approx 1.244 $$ Substracting 1: $$ .23 \approx .24$$

论坛问题帖 | has_answers | More Anonymous | 2020-12-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019130 | MSE-3970629

Sums of integer powers similar to Prouhet-Tarry-Escott problem

Recall Prouhet-Tarry-Escott problem. Its solution shows that certain sums of powers of integers can be made to vanish simultaneously if their signs are chosen to follow the Thue-Morse sequence, e.g. $$\sum_{k=0}^{2^4-1}(-1)^{\sigma_2(k)}\,k^n=0\quad\text{for}\;n<4,\quad\text{i.e.}\\ \small \begin{array}{l} 0^1+3^1+5^1+6^1+9^1+10^1+12^1+15^1=1^1+2^1+4^1+7^1+8^1+11^1+13^1+14^1 \\ 0^2+3^2+5^2+6^2+9^2+10^2+12^2+15^2=1^2+2^2+4^2+7^2+8^2+11^2+13^2+14^2 \\ 0^3+3^3+5^3+6^3+9^3+10^3+12^3+15^3=1^3+2^3+4^3+7^3+8^3+11^3+13^3+14^3, \\ \end{array}$$ where $\sigma_2(k)$ is the sum of digits of k in base-2. I am interested in sums of powers of a similar form, but with larger exponents (such that the sum does not vanish anymore). I was able to make a few conjectures: $$\begin{align} &\sum _{k=0}^{2^n-1} (-1)^{\sigma _2(k)}\, k^{n+2}\stackrel{\color{gray}?}{=}\frac{2^{\binom{n}{2}}}{36}\, (-1)^n\, \left(2^n-1\right)\, \left(5\times2^n-4\right)\, (n+2)!\\ \\ &\sum _{k=0}^{2^n-1} (-1)^{\sigma _2(k)}\,k^{n+3}\stackrel{\color{gray}?}{=}\frac{2^{\binom{n}{2}}}{72}\,(-1)^n\,\left(2^n-1\right)^2\,\left(2\times2^n-1\right)\,(n+3)!\\ \\ &\small\sum _{k=0}^{2^n-1} (-1)^{\sigma_2(k)}\,k^{n+4}\stackrel{\color{gray}?}{=}\frac{2^{\binom{n}{2}}}{32400}\,(-1)^n\,\left(2^n-1\right)\,\left(143\!\times\! 8^n-307\!\times\!4^n+193\!\times\! 2^n-32\right)\,(n+4)!\end{align}$$ This pattern seems to continue within increasingly complicated polynomials of 2^n on the right-hand side. Can you suggest an approach to prove these, and to find a more general formula for an arbitrary exponent k^{n+p}?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2021-01-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019131 | MSE-3971819

A curious infinite product of factorials

I found the following infinite product of factorials without proof: $$\small\prod_{n=1}^\infty\frac{{(2 n)!}^{20}\,{(8 n)!}^{32}\,{(32 n)!}^2}{{n!}^4\,{(4 n)!}^{37}\,{(16 n)!}^{13}}=\\\frac{\sin ^{14}\!\frac{\pi }{8}\cdot\sin \frac{\pi }{32} \cdot\sin \frac{3 \pi }{32} \cdot\sin \frac{5 \pi }{32} \cdot\sin \frac{7 \pi }{32}}{\sin ^6\!\frac{\pi }{16} \cdot\sin ^6\!\frac{3\pi }{16}}\cdot \frac{2^{1283/64}\,\pi^{14}\,\Gamma^{10} \!\left(\frac{1}{8}\right) \Gamma^2\! \left(\frac{5}{32}\right) \Gamma^2 \!\left(\frac{7}{32}\right)}{\Gamma^{18} \!\left(\frac{5}{8}\right) \Gamma^{10} \!\left(\frac{1}{16}\right) \Gamma^{10} \!\left(\frac{3}{16}\right) \Gamma^2 \!\left(\frac{17}{32}\right) \Gamma^2 \!\left(\frac{19}{32}\right)}.$$ We can verify that $$\small\frac{{(2 n)!}^{20}\,{(8 n)!}^{32}\,{(32 n)!}^2}{{n!}^4\,{(4 n)!}^{37}\,{(16 n)!}^{13}}=1+\mathcal O\left(n^{-3}\right),$$ so the product indeed converges. Can you suggest how to prove its closed form on the right? Is it possible to further simplify it? Is it possible to find a simpler convergent infinite product of this form (involving only integer powers of factorials of integer multiples of n)?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2021-01-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019132 | MSE-3982312

Integer solutions to $r\cdot a^2+t=s\cdot b^2$ forms sequences with $a_{n+1}\cdot b_n-a_n\cdot b_{n+1}$ is invariant

Conjecture: Given natural numbers r,s,t. Suppose there are an infinite number of solutions $a_n,b_n \in \mathbb N$ to $ra_n^2+t=sb_n^2$, where a_n, b_n correspond to increasing sequences of solutions, then $a_n\cdot b_{n+1} - a_{n+1}\cdot b_n$ is invariant equal for each n greater than some $N>0$. I've done a lot of tests without exceptions. The number N is typically ≤ 1. I would like to see ideas how to proceed. Some (r,s,t) values plus sets of (a,b) values: (1,2,1) {(1,1),(7,5),(41,29),(239,169),(1393,985),(8119,5741)} (2,1,1) {(0,1),(2,3),(12,17),(70,99),(408,577),(2378,3363)} (1,2,2) {(0,1),(4,3),(24,17),(140,99),(816,577),(4756,3363)} (3,2,2) {(0,1),(4,5),(40,49),(396,485),(3920,4801)} (2,3,1) {(1,1),(11,9),(109,89),(1079,881),(10681,8721)} There seems to be a matrix $ \begin{pmatrix} \alpha & ms \\ mr & \alpha \\ \end{pmatrix} $ that transform (a_{n},b_{n}) to (a_{n+1},b_{n+1}), but I haven't proved that and I'm insecure about α and m. It could be that $\alpha=\sqrt{m^2rs+1}$ and that the determinant is unital. The conjecture is false with counter-example (1,2,7) {(1,2),(5,4),(11,8),(31,22),(65,46),(181,128),(379,268)} What's left is Given $r,s,t\in\mathbb R_+$ and $S=\{(a,b)\in\mathbb R_+^2|rs^2+t=sb^2\}$, then it exist a sequence $(a_n,b_n)\in S$ such that: $$a_{n+1}b_n-a_nb_{n+1}=2t \tag 1$$ $$ \begin{pmatrix} a_{n+1} \\ b_{n+1} \\ \end{pmatrix} = \begin{pmatrix} \sqrt{4rs+1}& 2s \\ 2r & \sqrt{4rs+1} \\ \end{pmatrix} \begin{pmatrix} a_n\\ b_n \\ \end{pmatrix} \tag 2$$ which I consider as virtually proved by the accepted answer.

论坛问题帖 | has_answers | Lehs | 2021-01-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019133 | MSE-3987996

Can you spot the mistake here?

assume that : $$x^{3} + y^{3} = z^{3}$$ so : $$z^{3} - y^{3} = x^{3}$$ $$z^{3} ≡ y^{3}\bmod x$$ $$z ≡ y\bmod x $$ $$z-y = x $$ and now let's plug this result to the original equation: $$(z-y)^{3} + y^{3} = z^{3}$$ $$z^{3}-3z^{2}y+3zy^{2}-y^{3}+y^{3}=z^3$$ by simplfying, we get this: $$3zy^{2} = 3z^{2}y$$ dividing by $3zy$: $$y=z$$ so this equation has solutions if and only if $y = z$, and now let's plug our results to the original equation: $$x^{3}+y^{3} = y^{3}$$ $$x^{3} = 0$$ $$x = 0$$ Therefore :There is no solution when: $$x,y,z\in\mathbb{N}$$ and by mathematical induction we can prove it (instead of the power 3) for all $ n\in\mathbb{N}$

论坛问题帖 | has_answers | PNT | 2021-01-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019134 | MSE-3998982

Does the following lower bound improve on $I(q^k) + I(n^2) > 3 - \frac{q-2}{q(q-1)}$, where $q^k n^2$ is an odd perfect number?

Let $N = q^k n^2$ be an odd perfect number with special prime q satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. Define the abundancy index $$I(x)=\frac{\sigma(x)}{x}$$ where $\sigma(x)$ is the classical sum of divisors of x. Since q is prime, we have the bounds $$\frac{q+1}{q} \leq I(q^k) < \frac{q}{q-1},$$ which implies, since N is perfect, that $$\frac{2(q-1)}{q} < I(n^2) = \frac{2}{I(q^k)} \leq \frac{2q}{q+1}.$$ By considering the negative product $$\bigg(I(q^k) - \frac{2(q-1)}{q}\bigg)\bigg(I(n^2) - \frac{2(q-1)}{q}\bigg) < 0,$$ since we obviously have $$\frac{q}{q-1} < \frac{2(q-1)}{q},$$ then after some routine algebraic manipulations, we arrive at the lower bound $$I(q^k) + I(n^2) > 3 - \frac{q-2}{q(q-1)} = \frac{3q^2 - 4q + 2}{q(q - 1)}.$$ Now, a recent MO post improves on the lower bound for $I(n^2)$, as follows: $$I(n^2) > \bigg(\frac{2(q-1)}{q}\bigg)\bigg(\frac{q^{k+1} + 1}{q^{k+1}}\bigg)$$ Added February 3, 2021 - 8:10 PM (Manila time) Here is a quick way to show the improved lower bound for $I(n^2)$. We have $$I(n^2)=\frac{2}{I(q^k)}=\frac{2q^k (q - 1)}{q^{k+1} - 1}=\frac{2q^{k+1} (q - 1)}{q(q^{k+1} - 1)}=\bigg(\frac{2(q-1)}{q}\bigg)\bigg(\frac{q^{k+1}}{q^{k+1} - 1}\bigg)$$ $$=\bigg(\frac{2(q-1)}{q}\bigg)\bigg(\frac{(q^{k+1} - 1) + 1}{q^{k+1} - 1}\bigg)=\bigg(\frac{2(q-1)}{q}\bigg)\bigg(1 + \frac{1}{q^{k+1} - 1}\bigg)$$ $$>\bigg(\frac{2(q-1)}{q}\bigg)\bigg(1 + \frac{1}{q^{k+1}}\bigg)$$ since $q^{k+1} > q^{k+1} - 1$. But, of course, we obtain $$I(n^2)>\bigg(\frac{2(q-1)}{q}\bigg)\bigg(1 + \frac{1}{q^{k+1}}\bigg)=\bigg(\frac{2(q-1)}{q}\bigg)\bigg(\frac{q^{k+1} + 1}{q^{k+1}}\bigg).$$ QED. Repeating the same procedure as above, we have the negative product $$\Bigg(I(q^k) - \left(\frac{2(q-1)}{q}\bigg)\bigg(\frac{q^{k+1} + 1}{q^{k+1}}\right)\Bigg)\Bigg(I(n^2) - \left(\frac{2(q-1)}{q}\bigg)\bigg(\frac{q^{k+1} + 1}{q^{k+1}}\right)\Bigg) < 0.$$ This implies, after some algebraic manipulations, that $$I(q^k) + I(n^2) > \frac{q^{k+2}}{(q - 1)(q^{k+1} + 1)} + \frac{2(q-1)(q^{k+1} + 1)}{q^{k+2}}.$$ But WolframAlpha says that the partial fraction decomposition of the new lower bound is given by $$\frac{q^{k+2}}{(q - 1)(q^{k+1} + 1)} + \frac{2(q-1)(q^{k+1} + 1)}{q^{k+2}} = \frac{3q^2 - 4q + 2}{q(q - 1)} + \frac{2(q - 1)}{q^{k+2}} - \frac{q}{(q - 1)(q^{k+1} + 1)}.$$ So essentially, my question boils down to: QUESTION: Is it possible to produce an unconditional proof (that is, for all $k \geq 1$ and for all special primes $q \geq 5$) for the following inequality? $$\frac{2(q - 1)}{q^{k+2}} > \frac{q}{(q - 1)(q^{k+1} + 1)}$$ MY ATTEMPT I tried to ask WolframAlpha for a plot of the above inequality, it gave me the following GIF image: So it does appear that the inequality is unconditionally true, which would mean that the new lower bound for $I(q^k) + I(n^2)$ improves on the old. Is it possible to prove this analytically? And lastly: Based on this answer to a closely related question, since we appear to have obtained an improved lower bound for $I(q^k) + I(n^2)$, can we then say that there is indeed an integer a such that $k \leq a$?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2021-01-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019135 | MSE-4000653

Let $n=am+r$ with $m \ge a>5$ and $r \ge 0$. Prove that $\phi(n) \ge m$

I understand that for all $n>6$, we have $\phi(n) \ \ge \ \sqrt n $. This fact however is not getting me anywhere towards the proof of the claim in question. (a, m, r are all integers)

论坛问题帖 | has_answers | ASP | 2021-01-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019136 | MSE-4000705

Conjecture: primes non-decreasing from $(n,2n)$ to $(2n,4n)$.

More generally, I assert: $$\pi(na^k)-\pi(na^{k-1})\leq \pi(na^{k+1})-\pi(na^k)\quad\text{for }n,a,k \in \mathbb N.$$ The most useful case of this is probably to say that if there are m primes in $[n,2n]$, then there will be at least m primes in $[2n,4n]$. I thought it worth noting this can also be written as $$\pi(2n) \leq \frac{\pi(4n)+\pi(n)}{2}.$$ I doubt there are counterexamples, but if someone finds one, please share. Barring that, since this seems like an intermediate-strength statement, I'm curious how easy or difficult this might be to prove, and whether there are known similar results. Update Looks like a much tighter bound can also be specified below as $ \pi(3n)+\pi(n) \leq 2\pi(2n)$ for $n>1323$. Also, note this can be used in reverse. E.g. suppose we've worked out that the only primes in $[8,16]$ are 11 and 13. Then, even knowing nothing whatsoever about primes other than this rule (and that they start at 2), it follows that there could be at most 2 more primes in $[4,8]$, and another 2 in $[2,4]$, so we'd know that $\pi(13)\leq 6$.

论坛问题帖 | has_answers | Trevor | 2021-01-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019137 | MSE-4004640

Conjecture: if collatz(n) reaches 1, it must do so in at most 2^{n+1} steps

Let $f(n)$ give the number of Collatz iterations it takes to yield a number less than n, using the "shortcut" version of Collatz which applies $n\rightarrow (3n+1)/2$ for odds. Suppose it can be shown that $$f(n)=k \iff f(n+a2^k)=k\qquad \forall a\in\mathbb N.$$ So if a starting value of n decreases, it must happen in at most 2^n steps, and then we can say the same thing about that lower number, and so on, summing to a worst case upper bound of 2^{n+1} steps. My question: Is my reasoning correct, and if so, is this a known result? Per request, the reason I think I can show the above is based on the empirical results for $n<5\times 10^6$, which gives: The leftmost column is k, the number of steps required to reach a smaller number. The fourth column, obviously going as 2^k, is the period after which appearances of k repeat. There are multiple appearances of a given k within that period, e.g. for $k=5$ which shows up with alternating intervals of 12 and 20, but that's not required for my conjecture. This pattern strikes me as regular enough that I suspect a good reason for it would not be hard to work out. Of course, I could be wrong about that, but my question is whether the rest of my logic makes sense assuming a proper proof for this part exists. In fact, Wikipedia gives "... it can be shown that the parity sequences for two numbers m and n will agree in the first k terms if and only if m and n are equivalent modulo 2^k." Parity in this context refers to whether a step increases or decreases the number, so if the parity sequences are the same for two numbers, they will indeed both decrease below their original value on the same step number, which seems like it confirms my suspicion that the above is solid, but as Collag3n points out, is not quite proof. Couple of thoughts on $f(n)=k \iff f(n+a2^k)=k\qquad \forall a\in\mathbb N$: First, we still assert that $f(n)=k \implies f(n+a2^k)=k$. Then, let $f(n)=j$ and $f(n+2^k)=k$, where $j \neq k$. Note that $f(n)=j \implies f(n+a2^j)=j$. If $j<k$, then let $a=2^{k-j}$ and we have $f(n+a2^k)=f(n+(2^{k-j})2^j)=j=f(n+2^k)=k$ by the asserted implication, which is a contradiction. Experimentally, it looks like $f(n+2^m)>m$ until hitting $f(n+2^k)=k$. Note this would give $f(n+2^k)=k\implies f(n+2^{k-1})>f(n+2^k)$.

论坛问题帖 | no_answers_at_retrieval | Trevor | 2021-01-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019138 | MSE-4007078

There are $2^{\omega(a)}$ pairs (b,c) such that $a=bc$ and $\gcd(b,c)=1$

It looks as there are $2^{\omega(a)}$ ordered pairs (b,c) such that $a=bc$ and $\gcd(b,c)=1$, where $\omega(a)$ is the number of different prime factors of a. Proof? It's also true that there are $n^{\omega(a)}$ n-tipples $(a_1,\dots ,a_n)$ such that $a=a_1\cdots a_n$.

论坛问题帖 | has_answers | Lehs | 2021-01-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019139 | MSE-400708

How to prove $4\times{_2F_1}(-1/4,3/4;7/4;(2-\sqrt3)/4)-{_2F_1}(3/4,3/4;7/4;(2-\sqrt3)/4)\stackrel?=\frac{3\sqrt[4]{2+\sqrt3}}{\sqrt2}$

I have the following conjecture, which is supported by numerical calculations up to at least 10^5 decimal digits: $$4\times{_2F_1}\left(-\frac{1}{4},\frac{3}{4};\frac{7}{4};\frac{2-\sqrt{3}}{4}\right)-{_2F_1}\left(\frac{3}{4},\frac{3}{4};\frac{7}{4};\frac{2-\sqrt{3}}{4}\right)\,\stackrel?=\,\frac{3\sqrt[4]{2+\sqrt{3}}}{\sqrt{2}},$$ where $_2F_1$ denotes the hypergeometric function. Can you suggest any ideas how to prove it? The conjectural closed form was obtained using WolframAlpha query ToRadicals[RootApproximant[2.9484457662642558059990881423857006769923 3]]

论坛问题帖 | has_answers | Laila Podlesny | 2013-05-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019140 | MSE-4008273

Conjecture: For almost all n, the sum of $n \bmod k$ for $k<n$ is unique.

It looks to me like $$f(n):= \sum_{k=1}^{n}{(n\bmod k)}$$ is almost unique to n, with the only systematic exceptions being $(2^m-1, 2^m)$ pairs, which share a value for any $m \in \mathbb N$. In other words, the set $\{f(n): n\in\mathbb N\land n\neq 2^m \ \forall m\in\mathbb N\}$ shouldn't have many duplicate values. As discussed in the answers, a handful of counterexamples have been found, but they seem more sparse than one would expect, and nobody has yet suggested why that might be, or what's special about the counterexamples found, apart from a moderate preponderance of small factors in the mean of the pair. Although there's not much data to go on, it also looks like counterexamples may come in the form $(n,n+2i)$, or possibly even $(n,n+2^i)$, and I would be interested if anyone can find a case outside of those. The latter form would also include the $(2^m-1,2^m)$ exceptions, which seems elegant. Also, if someone could set me straight on this: My understanding is that the term 'almost all' in this context means approaching 0 density in the limit; so if one could show that the exceptions aren't necessarily finite, but they are less common than powers of two, would that be sufficient?

论坛问题帖 | has_answers | Trevor | 2021-02-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019141 | MSE-4015940

Conjecture of Differential Geometry Line Element identity?

Question Lets say I have a line element $ds^2 = \sum_{\mu , \nu = 0}^n g^{\mu \nu} d \tilde x_\mu d \tilde x_\nu $ and $1.$ It is of the form $ds^2 = \sum_{\mu=0}^n g^{\mu \mu} d \tilde x_\mu d \tilde x_\mu= \sum_{\mu=0}^n (d (\int \sqrt{g^{\mu \mu}} d\tilde x_\mu))^2 = \sum_{\mu=0}^n d x_\mu^2$. This form of the line element $\sum_{\mu=0}^n d x_\mu^2$ is achieved via coordinate transformation. If the square root is negative one can use $*$ $2.$ There exists a path whose path-length extends to infinity We define $s = \int_{\lambda_0}^{\lambda_1} \sqrt{\sum_{\mu=0}^n \frac{ds_\mu}{d \lambda} \frac{d s^\mu}{d\lambda}} d \lambda $ with $s(0)= 0 = \lambda_0$ 3. There is a unique one to one mapping between s and its coordinates x_i . For example,

$s(x_{0c},x_{1c},\dots) = c$ then $s = c \implies (x_0,x_1,x_2,\dots) = (x_{0c},x_{1c},x_{2c},\dots)$ Further the path taken has if s is expressed coordinates x_i $4.$ Then also impose $d x_i \neq 0 $. For example if $ds^2 = dx^2 + dy^2$ then $s_\mu = (x,0)$ is not a valid path because $ds_1= 0 $ (counting μ from 0). However, $s_\mu= (x,-y)$ is a valid path. Then we have the following identity: $$(\int_{s=0}^\infty e^{-\alpha s^2} ds)^2 = \sum_{i=0}^n (\int_{x_{i0}}^\infty e^{-\alpha s(x_i)^2} dx_i)^2 $$ $*$Note if $g^{j j}$ is negative then we use the same procedure with a negative sign. For example, $s_\mu = (t,x) = (\lambda, \lambda)$ $$ds^2 = - dt^2 + dx^2\implies (\int_{s=0}^\infty e^{-\alpha s^2} ds)^2 = -(\int_{t=0}^\infty e^{-\alpha s^2} dt)^2 + (\int_{x=0}^\infty e^{-\alpha s^2} dx)^2 =0 $$ Note: ds^2 is also 0 Are there any counterexamples? Can I use this to find the mapping between s and x_i? Does the formula still work if there isn't a unique mapping between s and its coordinates? Heuristic Rough Derivation and Example The following proof is difficult to follow but is meant to give only a rough sketch. We define $s = \int_{\lambda_1}^{\lambda_2} \sqrt{\sum_{\mu=0}^n \frac{ds_\mu}{d \lambda} \frac{d s^\mu}{d\lambda}} d \lambda $ For example, let's say I have the line element: $$ ds^2 = dx^2 + dy^2 + dz^2$$ Now we multiply a factor of $\lambda_s e^{-\alpha s(x,y,z)}$ both sides. Note: λ_s should not be thought as a function $$ \lambda_s e^{-\alpha s(x,y,z)}ds^2 = \lambda_s e^{-\alpha s(x,y,z)} dx^2 +\lambda_s e^{-\alpha s(x,y,z)} dy^2 + \lambda_s e^{-\alpha s(x,y,z)}dz^2 $$ Now we assume the path takes is continuous and has a mapping such that $s(x,y,z) = c $ implies there exists a unique set of coordinates (x_c,y_c,z_c) and $s(x_c ,y_c ,z_c) = c$. Further, the path chosen cannot be such that $dx = 0$ or $dy= 0$ or $dz= 0$. As an example let the path be $s(x,y,z)$. We then do a transformation based on this and cancel the common coefficient: $$(\int_{s=0}^\infty e^{-\alpha s^2} ds)^2 = (\int_{x_{0}}^\infty e^{-\alpha s(x,y(x),z(x))^2} dx)^2 + (\int_{y_{0}}^\infty e^{-\alpha s(x(y),y,z(y))^2} dy)^2 + (\int_{z_{0}}^\infty e^{-\alpha s(x(z),y(z),z))^2} dz)^2 $$ $s = \sqrt{3} \lambda$ with $x=y=z=\lambda$ with $x_0 = y_0 =z_0 = \lambda_0 = 0$ when $s=0$. Thus, $$ (\int_{s=0}^\infty e^{-\alpha s^2} ds)^2 = 3(\int_0^{\infty} e^{ - 3 \alpha x^2} dx)^2 = (\int_0^{\infty} e^{ - 3 \alpha x^2} d(\sqrt{ 3 }x))^2 $$ Hence the identity holds.

论坛问题帖 | no_answers_at_retrieval | More Anonymous | 2021-02-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019142 | MSE-402432

How can I prove this closed form for $\sum_{n=1}^\infty\frac{(4n)!}{\Gamma\left(\frac23+n\right)\,\Gamma\left(\frac43+n\right)\,n!^2\,(-256)^n}$

How can I prove the following conjectured identity? $$\mathcal{S}=\sum_{n=1}^\infty\frac{(4\,n)!}{\Gamma\left(\frac23+n\right)\,\Gamma\left(\frac43+n\right)\,n!^2\,(-256)^n}\stackrel?=\frac{\sqrt3}{2\,\pi}\left(2\sqrt{\frac8{\sqrt\alpha}-\alpha}-2\sqrt\alpha-3\right),$$ where $$\alpha=2\sqrt[3]{1+\sqrt2}-\frac2{\sqrt[3]{1+\sqrt2}}.$$ The conjecture is equivalent to saying that $\pi\,\mathcal{S}$ is the root of the polynomial $$256 x^8-6912 x^6-814752 x^4-13364784 x^2+531441,$$ belonging to the interval $-1<x<0$. The summand came as a solution to the recurrence relation $$\begin{cases}a(1)=-\frac{81\sqrt3}{512\,\pi}\\\\a(n+1)=-\frac{9\,(2n+1)(4n+1)(4n+3)}{32\,(n+1)(3n+2)(3n+4)}a(n)\end{cases}.$$ The conjectured closed form was found using computer based on results of numerical summation. The approximate numeric result is $\mathcal{S}=-0.06339748327393640606333225108136874...$ (click to see 1000 digits).

论坛问题帖 | has_answers | Hanna K. | 2013-05-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019143 | MSE-4024390

Yet another Collatz generalization and conjecture

Observe that Collatz (the $\frac{3n+1}{2}$ version) can be written as: $$n_{i+1}:= \begin{cases}n_i+\left\lceil\frac{n_i}{2}\right\rceil\quad\text{ if odd }n_i \\ n_i-\left\lceil\frac{n_i}{2}\right\rceil\quad\text{ if even }n_i \end{cases}$$ Looking at it this way gives the following pleasingly simple table for the difference between between consecutive terms, where $\Delta n := n_{i+1}-n_i$: $$ \begin{array}{|l|l|} \hline n_i&\Delta n\\ \hline 1&+1 \\ \hline 2&-1 \\ \hline 3&+2 \\ \hline 4&-2 \\ \hline 5&+3 \\ \hline \ldots&\ldots\\ \hline \end{array} $$ I apologize for not having worked out a nice short-form version of this, but consider a generalization of the above to give a construction like: $$ \begin{array}{|l|l|} \hline n_i&\Delta n\\ \hline 1&+1 \\ \hline 2&+2 \\ \hline 3&-1 \\ \hline 4&-2 \\ \hline 5&+3 \\ \hline 6&+4 \\ \hline 7&-3 \\ \hline 8&-4 \\ \hline 9&+5 \\ \hline \ldots&\ldots\\ \hline \end{array} $$ In fact, let k be the number determining that cyclic count, so that $k=1$ for the standard Collatz in the first table, and $k=2$ in the second table, and $k=3$ would give $\{1,2,3,-1,-2,-3,4,5,6,-4\ldots\}$. And by that definition, let $f(n,k)$ return the corresponding Δn using k as described. Then the next term for regular Collatz could be found as $C(n):=n+f(n,1)$. More generally, let's define $$C_k(n):=n+f(n,k),$$ so e.g. by examining that second table, we see that $C_2(8)=8-4=4$ and $C_2(9)=14$. This construction seems like it might have some interesting properties. In particular, I conjecture that $\forall k,n\in\mathbb N : C_k(n)$ will eventually reach k, assuming Collatz itself holds. Furthermore, if you take any such sequence and number it starting with $S_0=n$, it seems we have $$\gcd(S_i,k) \leq \gcd(S_{i+1},k).$$ This means that once you have $k|S_j$ for some j, all future terms can be divided through by k to give a standard Collatz sequence, but this is definitely not the case before reaching S_j. I'm interested to see if anyone can find a counterexample or simple reason why this should or should not hold. Any reasonably constructive feedback would be acceptable as an answer, and of course if this has been investigated before, by all means let me know. Sample Mathematica code if anyone wants it: Clear[f, c]; SetAttributes[{f, c}, Listable]; f[n_, k_] := With[{m = Mod[n, 2 k, 1]}, If[m <= k, 1, -1] (Floor[(n - 1)/(2 k)] (k) + Mod[n, k, 1])] c[n_, k_] := NestWhileList[# + f[#, k] &, n, # != k &, 1, 1000] c[27, Range[6]] // Column will give you sequences for $C_k(27)$ for $k \leq 6$.

论坛问题帖 | has_answers | Trevor | 2021-02-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019144 | MSE-4024470

Differential Geometry Inequality Conjecture?

Question and Conjecture In a Lorentzian metric (Only one time dimension) of signature: $(+,-,-,-,\dots)$ Lets say I have a line element $ds^2 = \sum_{\mu , \nu = 0}^n g^{\mu \nu} d \tilde x_\mu d \tilde x_\nu $ and $1.$ It is of the form $ds^2 = \sum_{\mu=0}^n g^{\mu \mu} d \tilde x_\mu d \tilde x_\mu$ $2.$ There exists a continuous path

parametrized by λ whose path-length extends to infinity We define $s = \int_{\lambda_0}^{\lambda_1} \sqrt{\sum_{\mu=0}^n \frac{d \tilde x_\mu}{d \lambda} \frac{d \tilde x^\mu}{d\lambda}} d \lambda $ with $s(0)= 0 = \lambda_0$ $3.$ It must be of the Lorentzian only g_{00} must be positive and the path must be timelike. Then: We define: $$ g^{00} ds'^2 = ds^2 $$ and define coordinates: $d x_i = \sqrt{|g^{ii}|} d \tilde x_i$. Then: $$(\int_{s(\lambda_0)=0}^\infty e^{-\alpha s'^2} ds')^2 \geq (\int_{\lambda_0}^\infty e^{-\alpha s(x (\lambda),y(\lambda),z(\lambda))^2} \frac{dx_0}{d \lambda} d \lambda)^2 - \sum_{\mu=1}^n (\int_{\lambda_0}^\infty e^{-\alpha s(x(y),y,z(y))^2} \frac{d x_\mu}{d \lambda} d \lambda)^2 $$ Are there any counter examples? Heuristic Rough Derivation and Example The following proof is difficult to follow but is meant to give only a rough sketch. We define s as done already previously now let's say I have the line element: $$ ds^2 = f(r,t)^2 dt^2 - r^2 d\theta^2 - dr^2 $$ We compare this to: $$ f(r,t) ds' = ds $$ Hence, $$ ds'^2 = dt^2 - \frac{dr^2}{f(r,t)^2} - \frac{r^2}{f(r,t)^2} d \theta^2 $$ Now we multiply a factor of $\beta_s e^{-\alpha s'(x,y,z)}$ both sides. Note: β_s should not be thought as a function $$ \beta_s e^{\alpha s'} ds'^2 = e^{\alpha s'} \beta_s dt^2 - \beta_s e^{\alpha s'} \frac{dr^2}{f(r,t)^2} - \beta_s e^{\alpha s'} \frac{r^2}{f(r,t)^2} d \theta^2 $$ Let: $$ \beta_s e^{\alpha s'} ds'^2 = e^{\alpha s'} \beta_s dt^2 - \beta_s e^{\alpha s'} \frac{dr^2}{f(r,t)^2} - \beta_s e^{\alpha s'} \frac{r^2}{f(r,t)^2} d \theta^2 $$ We can also say: $$\beta_s e^{\alpha s'} ds'^2 \geq e^{\alpha s'} \beta_s dt^2- \beta_s e^{\alpha s'} (d \int \frac{ dr}{f(r,t)})^2 - \beta_s e^{\alpha s'} (d \int \frac{r}{f(r,t)} d \theta)^2 $$ We define $y = \int \frac{ dr}{f(r,t)} $ and $z = \int \frac{r}{f(r,t)} d \theta$ Now we assume the path takes is continuous. We then do a transformation based on this and cancel the common coefficient: $$(\int_{s(\lambda_0)=0}^\infty e^{-\alpha s'^2} ds')^2 \geq (\int_{\lambda_0}^\infty e^{-\alpha s(x (\lambda),y(\lambda),z(\lambda))^2} \frac{dt}{d \lambda} d \lambda)^2 $$ $$- (\int_{y(\lambda_0)}^\infty e^{-\alpha s(x(y),y,z(y))^2} \frac{dy}{d \lambda} d \lambda)^2 - (\int_{\lambda_0}^\infty e^{-\alpha s(x(z),y(z),z))^2} \frac{dz}{d \lambda} d \lambda)^2 $$

论坛问题帖 | no_answers_at_retrieval | More Anonymous | 2021-02-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019145 | MSE-4027277

Is this Pythagorean triple formula original?

I am hesitant because I have asked this question before in a different form here. In 2009, I knew nothing about Pythagorean triples, not even Euclid's formula. In my ignorance, I did a brute force search, and the patterns I saw led to a formula for generating them. Years later, when I learned about Euclid's formula, it was confusing and I questioned it here. I'm an amateur but what I have found has kept me plodding along for over a decade. I developed a formula that should have been discovered in the 2300 years since Euclid. It generates the subset of triples where $GCD(A,B,C)$ is an odd square and this includes all primitives--without the trivials, the doubles and the even-square multiples of Euclid's formula. The formula is like Euclid's $ \quad A=m^2-k^2,\quad B=2mk,\quad C=m^2+k^2\quad$ but, as I discovered later, with $\space m\space$ replaced by $\space (2m-1+k)\space$ and this is why I show them both with a $\space k.\space$ This formula generates non-trivial triples for every pair of natural numbers (n,k) and very few non-primitives. Here is \begin{align*} A=(2n-1)^2+ &\quad 2(2n-1)k \\ B= \qquad &\quad 2(2n-1)k+ \quad 2k^2\\ C=(2n-1)^2+ &\quad 2(2n-1)k+ \quad 2k^2\\ \end{align*} It produces "sets" of triples where non-primitives like $\space(27,36,45)\space$ are in predictable/avoidable locations. \begin{array}{c|c|c|c|c|c|} n & k=1 & k=2 & k=3 & k=4 & k=5 \\ \hline Set_1 & 3,4,5 & 5,12,13& 7,24,25& 9,40,41& 11,60,61 \\ \hline Set_2 & 15,8,17 & 21,20,29 &27,36,45 &33,56,65 & 39,80,89 \\ \hline Set_3 & 35,12,37 & 45,28,53 &55,48,73 &65,72,97 & 75,100,125 \\ \hline Set_{4} &63,16,65 &77,36,85 &91,60,109 &105,88,137 &119,120,169 \\ \hline Set_{5} &99,20,101 &117,44,125 &135,72,153 &153,104,185 &171,140,221 \\ \hline Set_{6} &43,24,145 &165,52,173 &187,84,205 &209,120,241 &231,160,281 \\ \hline \end{array} $\text{If }\space(2n-1)=1\space $ or $\space k=1,\space $ triples are primitive: $\quad C-B=1\space \lor\space C-A=2.$ $\text{If }\space(2n-1)\space$ is prime, the following formula will generate only primitives. It repeatedly generates $\quad\big((2n-1)-1\big)\quad $ primitives and then skips a predicted non-primitive. \begin{align*} &A=(2n-1)^2+&2(2n-1)\bigg(k+\bigg\lfloor\frac{(k-1)}{(2n-2)}\bigg\rfloor\bigg)&\qquad\\ &B=&2(2n-1)\bigg(k+\bigg\lfloor\frac{(k-1)}{(2n-2)}\bigg\rfloor\bigg)&\qquad+2\bigg(k+\bigg\lfloor\frac{(k-1)}{(2n-2)}\bigg\rfloor\bigg)^2\\ &C=(2n-1)^2+&2(2n-1)\bigg(k+\bigg\lfloor\frac{(k-1)}{(2n-2)}\bigg\rfloor\bigg)&\qquad+2\bigg(k+\bigg\lfloor\frac{(k-1)}{(2n-2)}\bigg\rfloor\bigg)^2 \end{align*} 3) Else, there is a non-primitive whenever $\space k\space$ is a multiple of any factor of $\space(2n-1).$ I have not been able to find reference anything like this formula or the sets it generates in the many books I've bought on the subject or on web sites or in forums like this. Is this formula original with me or is there someone, somewhere or something I should cite in my references?

论坛问题帖 | no_answers_at_retrieval | poetasis | 2021-02-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019146 | MSE-404222

How to prove $\int_0^\infty J_\nu(x)^3dx\stackrel?=\frac{\Gamma(1/6)\ \Gamma(1/6+\nu/2)}{2^{5/3}\ 3^{1/2}\ \pi^{3/2}\ \Gamma(5/6+\nu/2)}$?

I am interested in finding a general formula for the following integral: $$\int_0^\infty J_\nu(x)^3dx,\tag1$$ where $J_\nu(x)$ is the Bessel function of the first kind: $$J_\nu(x)=\sum _{n=0}^\infty\frac{(-1)^n}{\Gamma(n+1)\Gamma(n+\nu+1)}\left(\frac x2\right)^{2n+\nu}.\tag2$$ Mathematica gives the following result: $$\int_0^\infty J_\nu(x)^3dx=\frac1{\pi\,\nu}{}_3F_2\left(\frac12,\frac12-\nu,\frac12+\nu;1-\frac\nu2,1+\frac\nu2;\frac14\right)+\frac{\Gamma\left(-\frac\nu2\right)\Gamma\left(\frac{1+3\nu}2\right)\cos\frac{\pi\,\nu}2}{2^{\nu+1}\ \pi^{3/2}\ \Gamma(\nu+1)}{}_3F_2\left(\frac{1-\nu}2,\frac{1+\nu}2,\frac{1+3\nu}2;1+\frac\nu2,1+\nu;\frac14\right),\tag3$$ which can be significantly simplified for odd and half-integer values of ν. The results at those points allow to conjecture another, simpler general formula: $$\int_0^\infty J_\nu(x)^3dx\stackrel?=\frac{\Gamma\left(\frac16\right)\ \Gamma\left(\frac16+\frac\nu2\right)}{2^{5/3}\ 3^{1/2}\ \pi^{3/2}\ \Gamma\left(\frac56+\frac\nu2\right)},\tag4$$ which agrees with (3) to a very high precision for many different values of ν. It also has an advantage that it is defined for all $\nu>-\frac13$ whereas (3) is undefined at every even ν and requires calculating a limit at those points. Is it possible to prove the formula (4)? Mathematica is able to

evaluate (1) for even values of ν in terms of the Meijer G-function. Plugging those expressions into (4) we get another form of the conjecture: $$G_{3,3}^{2,1}\left(\frac14\left|\begin{array}{c}2a,1,2-2a\\\frac12,1-a,a\\\end{array}\right.\right)\stackrel?=\frac{\Gamma\left(\frac16\right)\ \Gamma\left(\frac23-a\right)}{2^{5/3}\ 3^{1/2}\ \pi\ \Gamma\left(\frac43-a\right)}.\tag5$$ Incidentally, the case $a=\frac12$ would positively resolve my another question.

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2013-05-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019147 | MSE-404391

Is $K\left(\frac{\sqrt{2-\sqrt3}}2\right)$ equal to $\frac{\Gamma\left(\frac16\right)\Gamma\left(\frac13\right)}{4\ \sqrt[4]3\ \sqrt\pi}$?

Working on this conjecture, I found its corollary, which is also supported by numeric calculations up to at least 10^5 decimal digits: $$K\left(\frac{\sqrt{2-\sqrt3}}2\right)\stackrel?=\frac{\Gamma\left(\frac16\right)\Gamma\left(\frac13\right)}{4\ \sqrt[4]3\ \sqrt\pi},$$ where $K(x)$ is the complete elliptic integral of the 1st kind. I did not find this specific value at MathWorld, Wolfram Functions Site, Wikipedia or DLMF. Is it a known value?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2013-05-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019148 | MSE-4048386

Product of Consecutive integers problem

Product of Consecutive integers problem I've begun pondering the following problem, and have found myself unable to advance on it; it goes like this: Prove that the product of three consecutive integers is never the product of two consecutive integers (with a finite number of exceptions). Now, I basically have very little idea of what to work with. Obviously, I have two exceptions: {1, 2, 3} ($1 \times 2 \times 3 = 2 \times 3$) and {5, 6, 7} ($5 \times 6 \times 7 = 14 \times 15$). I formed the following equations: $x(x-1)(x+1) = y(y+1)$ ==> $x^3 - x = y^2 + y$ But I have no idea where to go from here. Graphing this on desmos yields the following elliptical curve, which I don't think helps me: How do I prove/ this conjecture?

论坛问题帖 | no_answers_at_retrieval | user895383 | 2021-03-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019149 | MSE-4050308

A Question on Integers and Fractions

Given that a,b,c,d,e,f are all distinct positive integers, prove that there are no solutions for the following knowing their fractions are non-integers and completely simplified: $$\left(\frac{a}{c}\right)^2 + \left(\frac{b}{d}\right)^2 = e^2 \tag{1}$$ I coulden't find a proof for this one but I managed to prove this for the case of $$\left(\frac{a}{c}\right)^2 + \left(\frac{b}{d}\right) ^2 = \left(\frac{e}{f}\right)^2 \tag{2}$$ by simplifying: $$\frac{(ac)^2 + (bd)^2}{(cd)^2} = (\frac{e}{f})^2 $$ Since $(ac)^2 + (bd)^2$ must satisfy a Pythagorean triple, we know that each side length(exculding the hypotenuse) must be the product of two distinct integers, which can be proven through Euclid's formula : $$ac = (2m)(n)$$ $$bd = (m -n)(m + n)$$ Q.E.D. I also noted that (1) can be written as: $$(ac)^2 + (bd)^2 = (ecd)^2$$ This lead me to conjecture that there are no Pythagorean triples where the hypotenuse has 2 distinct integer factors where one of each is shared with its side lengths. Furthermore, I also conjecture the hypotenuse's remainder must be a prime such that those conditions can never be satisfied. Can anyone prove or disprove this?

论坛问题帖 | has_answers | SpoonedBread | 2021-03-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019150 | MSE-4067953

Proof of conjecture about orthogonalized Specht polynomials.

Let $\lambda=(\lambda_1,\lambda_2,\ldots,\lambda_r)$ be a r-part partition of integer N ($\lambda\vdash N$), i.e., $$\sum_{i=1}^r{\lambda_i}=N,$$ such that $\lambda_i\leq\lambda_j$ for $r\geq j>i\geq1$. Thus, λ defines a Young diagram. In addition, let $\mathbf{y}^{(k)}_{\lambda}$ be the kth standard Young tableau, SYT, of shape λ, i.e., the array of numbers $1\leq y_{i,j}\leq N$ that fulfills $y_{i,j}>y_{i-1,j}$ $y_{i,j}>y_{i,j-1}$ for indices $1\leq i \leq r$ and $1 \leq j \leq\lambda_i$. The number of possible SYT of shape λ is $\dim S^\lambda$, the dimension of the Specht module. Moreover, let us define the permutation operator $\hat{\pi}=(\hat{\pi}_1,\hat{\pi}_2,\ldots,\hat{\pi}_r)$, such that $$\mathbf{w}=\hat{\pi}\mathbf{y},$$ where $\textrm{row}_i[\mathbf{w}]=\hat{\pi}_i\textrm{row}_i[\mathbf{y}]$, with $\textrm{row}_i[\mathbf{w}]$ the ith row of array $\mathbf{w}$, and $\hat{\pi}_i\in S_{\lambda_i}$. Finally, let $$\textrm{Spe}\left(\mathbf{w}\right)=\prod_{i=1}^{\lambda_1}V\left(\textrm{col}_i[\mathbf{w}]\right)$$ be the Specht polynomial for the (not-necessarily standard) Young tableau $\mathbf w$, where $\textrm{col}_i[\mathbf{w}]$ is the ith column of array $\mathbf{w}$, and $$V\left(a_1,a_2,\ldots,a_n\right)=\prod_{1\leq j<k\leq n}(x_{a_j}-x_{a_k})$$ is the Vandermonde determinant of set $\{a_1,a_2,\ldots,a_n\}$. After playing around with Gram-Schmidt orthogonalization, I have formulated the following conjecture: The set of polynomials $$\left\{p_\lambda^{(k)}\left(x\right):1\leq k\leq \dim S^\lambda\right\},$$ where $x=\{x_i:1\leq i\leq r\}$, $$p_\lambda^{(k)}\left(x\right)=\sum_{c}\textrm{Spe}\left(\mathbf{w}_\lambda^{(c,k)}\right),$$ and $\mathbf{w}_\lambda^{(c,k)}=\hat{\pi}^{(c)}\mathbf{y}_\lambda^{(k)}$, with the index c running over all possible permutation operators that produce Young tableaux with the constraint $$w_{i,j}>w_{i,j-1},$$ forms an orthogonal basis in the domain $-1/2\leq x_{1\leq i\leq r} \leq 1/2$, i.e., $$\int_{-1/2}^{1/2} p_\lambda^{(k)}(x)p_\lambda^{(k')}(x)\textrm{d}^{r}x\propto\delta_{k,k'}.$$ My question is, how can I go about a general proof that does not involve checking case by case and performing all inner products one by one?

论坛问题帖 | no_answers_at_retrieval | GeoArt | 2021-03-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019151 | MSE-4075551

Progress on a conjecture of Burnside...

Given a group $G $, the set of automorphisms of $G $ also forms a group, $\rm {Aut}(G) $,with composition as the operation (recall that an automorphism of a group is a bijective endomorphism) . An inner automorphism is one determined by conjugation by some element $g\in G $. That's we have the automorphism $i_g $ given by $i_g (h)=ghg^{-1}\,\forall h $. I learned from a comment by @LeeMosher on this site that Burnside once conjectured that any class-preserving automorphism is inner. Does anyone know about any progress on this? Of course the converse is trivial. (Btw, the reference here is to conjugacy classes. Of course the class equation gives the sizes of these classes. For example, in the case of an abelian group, the class equation consists in all ones.) For reference, this would have been early in the 20-th century.

论坛问题帖 | has_answers | user403337 | 2021-03-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019152 | MSE-4081683

Is there any algorithm generating "all" chrodal graphs of order n?

I defined a procedure Γ to generate graphs following some rules. The obtained graphs form a class of graphs $\mathcal{G}$. I found out that these graphs are chrodal. In the other hand, I noticed that every chordal graph of order 3 and 4 can be constrcuted by the procedure Γ. I implemented an algorithm that generates all $\mathcal{G}$-graphs with n vertices. But I need some algorithm that constructs all the chordal graphs with n vertices in order to make the following conjecture more plausible. Let G be a graph. Then G belongs to $\mathcal{G}$ if and only if G is chordal. I want to show that every graph obtained by such an algorithm can also be obtained by the procedure Γ. So, my question is: Does any algorithm generating all chordal graphs exist?

论坛问题帖 | no_answers_at_retrieval | N. Chikh | 2021-03-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019153 | MSE-4081975

Asymptotic series for the inverse Riemann ζ-function

Consider the positive branch of the Riemann ζ-function: $\zeta:(1,\infty)\to(1,\infty)$ and its inverse $\zeta^{\small(-1)}:(1,\infty)\to(1,\infty)$ satisfying $\zeta\left(\zeta^{\small(-1)}(x)\right) = x$ for $x>1$. Here are a few initial terms of an asymptotic expansion of $\zeta^{\small(-1)}$ in powers of x (except for the initial term that is logarithmic): $$\begin{align}\zeta^{\scriptscriptstyle(-1)}(1+x)=-\log_2x+{\small\frac1{\ln2}}\Big(&x^{\log_2\left(\frac32\right)}+x+{\small\log_2\left(\tfrac{\sqrt2}3\right)}\cdot x^{2\log _2\left(\frac32\right)}\\+\,&x^{\log _2\left(\frac52\right)}-{\small\log_2 3}\cdot x^{\log_23}\Big)+\mathcal O\Big(x^{3\log_2\left(\frac32\right)}\Big)\color{gray}{,\,x\to0^+,}\end{align} $$ or, equivalently, $$\begin{align}\zeta^{\scriptscriptstyle(-1)}\left(1+2^{-z}\right)=z+{\small\frac1{\ln2}}\Big(&\left(\tfrac32\right)^{-z}+{\small2^{-z}}+{\small\log_2\left(\tfrac{\sqrt2}3\right)}\cdot \left(\tfrac32\right)^{-2z}\\+\,&\left(\tfrac52\right)^{-z}-{\small\log_2 3}\cdot {\small3^{-z}}\Big)+\mathcal O\Big(\left(\tfrac32\right)^{-3z}\Big)\color{gray}{,\,z\to\infty.}\end{align} $$ Can we compute more terms of this series or find a general formula for its terms? Does this series converge?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2021-03-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019154 | MSE-4087586

Addition to previously asked question on Generalized Carmichael Numbers

I had previously asked a question on mathstack exchange (Conjecture on The Generalized Carmichael Numbers) concerning with a conjecture I had discovered. I worked on the problem for a long time and discovered one more conjecture. With the same terms and notations explained in the previous question (Conjecture on The Generalized Carmichael Numbers) The new conjecture goes as follows: Let $k \geqslant 1$, we have $$\lim_{n \to \infty}\dfrac{\mid C_{-k} \text{ }\cap \text{ } (0, X]\mid}{\mid C_{k} \text{ }\cap \text{ } (0, X]\mid - \left(\pi\left(\dfrac{X}{k}\right)\dfrac{\gcd(\lambda(k), k)}{\lambda(k)}\right)} = 1$$ where $\pi(X)$ is the prime-counting function. Also Note that For $k \geqslant 1$, we have: $C_k = $ Generalized Carmichael Numbers + $k\left(\text{primes} \equiv 1 \mod\left(\dfrac{\gcd(\lambda(k), k)}{\lambda(k)}\right)\right)$ For $k \leqslant 0$, we have C_k = Generalized Carmichael Numbers. I have worked on this conjecture for a long time but I'm not sure why the phenomenon is happening. I have used lot of brute force (computer program) for checking and verifying the conjecture for some large values and seems like it works. Any help or advice on how to proceed further would be appreciated. I really appreciate your time and efforts. Thanks.

论坛问题帖 | no_answers_at_retrieval | BookWick | 2021-04-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019155 | MSE-4093971

Is there an odd x such that $2x^2 \equiv 0 \pmod {\sigma(x)}$ and $\sigma(x^2) \equiv 0 \pmod {\sigma(x) - 1}$?

CONTEXT This question is a result of considerations stemming from this closely related MO question. INITIAL QUESTION My question is as is in the title: Is there an odd x such that $$2x^2 \equiv 0 \pmod {\sigma(x)}$$ and $$\sigma(x^2) \equiv 0 \pmod {\sigma(x) - 1}?$$ Here, $\sigma(x)=\sigma_1(x)$ is the classical sum of divisors of x. MY ATTEMPT I tried searching for solutions to the congruences using GP in Sage Cell Server, I obtain: For both congruences for(x=2, 10000000, if((Mod(2*x^2,sigma(x)) == 0) && (Mod(sigma(x^2),sigma(x)-1) == 0),print(x,factor(x)))) No output returned. for(x=2, 1000000, if((Mod(2*x^2,sigma(x)) == 0) && (Mod(sigma(x^2),sigma(x)-1) == 0),print(x,factor(x)))) No output returned. For the congruence $2x^2 \equiv 0 \pmod {\sigma(x)}$ for(x=2, 10000000, if((Mod(2*x^2,sigma(x)) == 0),print(x,factor(x)))) Several lines returned. Will confirm if everything so far are even: 6[2, 1; 3, 1] 28[2, 2; 7, 1] 30[2, 1; 3, 1; 5, 1] 84[2, 2; 3, 1; 7, 1] 120[2, 3; 3, 1; 5, 1] 364[2, 2; 7, 1; 13, 1] 496[2, 4; 31, 1] 672[2, 5; 3, 1; 7, 1] 840[2, 3; 3, 1; 5, 1; 7, 1] 1080[2, 3; 3, 3; 5, 1] 1488[2, 4; 3, 1; 31, 1] 1782[2, 1; 3, 4; 11, 1] 2280[2, 3; 3, 1; 5, 1; 19, 1] 3276[2, 2; 3, 2; 7, 1; 13, 1] 3472[2, 4; 7, 1; 31, 1]

7440[2, 4; 3, 1; 5, 1; 31, 1] 7560[2, 3; 3, 3; 5, 1; 7, 1] 8128[2, 6; 127, 1] 8736[2, 5; 3, 1; 7, 1; 13, 1] 8910[2, 1; 3, 4; 5, 1; 11, 1] 9480[2, 3; 3, 1; 5, 1; 79, 1] 10920[2, 3; 3, 1; 5, 1; 7, 1; 13, 1] 11880[2, 3; 3, 3; 5, 1; 11, 1] 12400[2, 4; 5, 2; 31, 1] 16368[2, 4; 3, 1; 11, 1; 31, 1] 16380[2, 2; 3, 2; 5, 1; 7, 1; 13, 1] 18360[2, 3; 3, 3; 5, 1; 17, 1] 18600[2, 3; 3, 1; 5, 2; 31, 1] 20832[2, 5; 3, 1; 7, 1; 31, 1] 24384[2, 6; 3, 1; 127, 1] 24840[2, 3; 3, 3; 5, 1; 23, 1] 24948[2, 2; 3, 4; 7, 1; 11, 1] 25296[2, 4; 3, 1; 17, 1; 31, 1] 26208[2, 5; 3, 2; 7, 1; 13, 1] 30240[2, 5; 3, 3; 5, 1; 7, 1] 30256[2, 4; 31, 1; 61, 1] 30294[2, 1; 3, 4; 11, 1; 17, 1] 32760[2, 3; 3, 2; 5, 1; 7, 1; 13, 1] 35640[2, 3; 3, 4; 5, 1; 11, 1] 37200[2, 4; 3, 1; 5, 2; 31, 1] 45136[2, 4; 7, 1; 13, 1; 31, 1] 55692[2, 2; 3, 2; 7, 1; 13, 1; 17, 1] 55860[2, 2; 3, 1; 5, 1; 7, 2; 19, 1] 56896[2, 6; 7, 1; 127, 1] 57240[2, 3; 3, 3; 5, 1; 53, 1] 66960[2, 4; 3, 3; 5, 1; 31, 1] 76680[2, 3; 3, 3; 5, 1; 71, 1] 81480[2, 3; 3, 1; 5, 1; 7, 1; 97, 1] 84360[2, 3; 3, 1; 5, 1; 19, 1; 37, 1] 86800[2, 4; 5, 2; 7, 1; 31, 1] 90768[2, 4; 3, 1; 31, 1; 61, 1] 94446[2, 1; 3, 4; 11, 1; 53, 1] 115560[2, 3; 3, 3; 5, 1; 107, 1] 121920[2, 6; 3, 1; 5, 1; 127, 1] 122668[2, 2; 7, 1; 13, 1; 337, 1] 131040[2, 5; 3, 2; 5, 1; 7, 1; 13, 1] 149856[2, 5; 3, 1; 7, 1; 223, 1] 170688[2, 6; 3, 1; 7, 1; 127, 1] 173628[2, 2; 3, 2; 7, 1; 13, 1; 53, 1] 199584[2, 5; 3, 4; 7, 1; 11, 1] 201960[2, 3; 3, 3; 5, 1; 11, 1; 17, 1] 211792[2, 4; 7, 1; 31, 1; 61, 1] 215760[2, 4; 3, 1; 5, 1; 29, 1; 31, 1] 235600[2, 4; 5, 2; 19, 1; 31, 1] 249480[2, 3; 3, 4; 5, 1; 7, 1; 11, 1] 251968[2, 6; 31, 1; 127, 1] 268224[2, 6; 3, 1; 11, 1; 127, 1] 270816[2, 5; 3, 1; 7, 1; 13, 1; 31, 1] 288288[2, 5; 3, 2; 7, 1; 11, 1; 13, 1] 293760[2, 7; 3, 3; 5, 1; 17, 1] 332640[2, 5; 3, 3; 5, 1; 7, 1; 11, 1] 334800[2, 4; 3, 3; 5, 2; 31, 1] 336784[2, 4; 7, 1; 31, 1; 97, 1] 360360[2, 3; 3, 2; 5, 1; 7, 1; 11, 1; 13, 1] 393120[2, 5; 3, 3; 5, 1; 7, 1; 13, 1] 406224[2, 4; 3, 2; 7, 1; 13, 1; 31, 1] 409200[2, 4; 3, 1; 5, 2; 11, 1; 31, 1] 414528[2, 6; 3, 1; 17, 1; 127, 1] 441936[2, 4; 3, 4; 11, 1; 31, 1] 445536[2, 5; 3, 2; 7, 1; 13, 1; 17, 1] 453840[2, 4; 3, 1; 5, 1; 31, 1; 61, 1] 514080[2, 5; 3, 3; 5, 1; 7, 1; 17, 1] 520800[2, 5; 3, 1; 5, 2; 7, 1; 31, 1] 523776[2, 9; 3, 1; 11, 1; 31, 1] 556920[2, 3; 3, 2; 5, 1; 7, 1; 13, 1; 17, 1] 560832[2, 6; 3, 1; 23, 1; 127, 1] 574560[2, 5; 3, 3; 5, 1; 7, 1; 19, 1] 602784[2, 5; 3, 2; 7, 1; 13, 1; 23, 1] 605880[2, 3; 3, 4; 5, 1; 11, 1; 17, 1] 622440[2, 3; 3, 2; 5, 1; 7, 1; 13, 1; 19, 1] 629640[2, 3; 3, 3; 5, 1; 11, 1; 53, 1] 632400[2, 4; 3, 1; 5, 2; 17, 1; 31, 1] 677160[2, 3; 3, 4; 5, 1; 11, 1; 19, 1] 695520[2, 5; 3, 3; 5, 1; 7, 1; 23, 1] 698760[2, 3; 3, 3; 5, 1; 647, 1] 706800[2, 4; 3, 1; 5, 2; 19, 1; 31, 1] 739648[2, 6; 7, 1; 13, 1; 127, 1] 763308[2, 2; 3, 2; 7, 1; 13, 1; 233, 1] 812448[2, 5; 3, 2; 7, 1; 13, 1; 31, 1] 819000[2, 3; 3, 2; 5, 3; 7, 1; 13, 1] 819720[2, 3; 3, 4; 5, 1; 11, 1; 23, 1] 828816[2, 4; 3, 1; 31, 1; 557, 1] 853440[2, 6; 3, 1; 5, 1; 7, 1; 127, 1] 876960[2, 5; 3, 3; 5, 1; 7, 1; 29, 1] 884520[2, 3; 3, 5; 5, 1; 7, 1; 13, 1] 899808[2, 5; 3, 1; 7, 1; 13, 1; 103, 1] 950040[2, 3; 3, 2; 5, 1; 7, 1; 13, 1; 29, 1] 979600[2, 4; 5, 2; 31, 1; 79, 1] 997920[2, 5; 3, 4; 5, 1; 7, 1; 11, 1] 1033560[2, 3; 3, 4; 5, 1; 11, 1; 29, 1] 1078800[2, 4; 3, 1; 5, 2; 29, 1; 31, 1] 1097280[2, 6; 3, 3; 5, 1; 127, 1] 1108560[2, 4; 3, 1; 5, 1; 31, 1; 149, 1] 1128400[2, 4; 5, 2; 7, 1; 13, 1; 31, 1] 1146048[2, 6; 3, 1; 47, 1; 127, 1] 1230120[2, 3; 3, 3; 5, 1; 17, 1; 67, 1] 1231776[2, 5; 3, 2; 7, 1; 13, 1; 47, 1] 1239840[2, 5; 3, 3; 5, 1; 7, 1; 41, 1] 1270752[2, 5; 3, 1; 7, 1; 31, 1; 61, 1] 1341120[2, 6; 3, 1; 5, 1; 11, 1; 127, 1] 1343160[2, 3; 3, 2; 5, 1; 7, 1; 13, 1; 41, 1] 1421280[2, 5; 3, 3; 5, 1; 7, 1; 47, 1] 1441440[2, 5; 3, 2; 5, 1; 7, 1; 11, 1; 13, 1] 1503376[2, 4; 7, 1; 31, 1; 433, 1] 1543056[2, 4; 3, 1; 17, 1; 31, 1; 61, 1] 1602720[2, 5; 3, 3; 5, 1; 7, 1; 53, 1] 1675080[2, 3; 3, 4; 5, 1; 11, 1; 47, 1] 1731264[2, 6; 3, 1; 71, 1; 127, 1] 1784160[2, 5; 3, 3; 5, 1; 7, 1; 59, 1] 1854360[2, 3; 3, 3; 5, 1; 17, 1; 101, 1] 1860768[2, 5; 3, 2; 7, 1; 13, 1; 71, 1] 1888920[2, 3; 3, 4; 5, 1; 11, 1; 53, 1] 1932840[2, 3; 3, 2; 5, 1; 7, 1; 13, 1; 59, 1] 2056320[2, 7; 3, 3; 5, 1; 7, 1; 17, 1] 2102760[2, 3; 3, 4; 5, 1; 11, 1; 59, 1] 2147040[2, 5; 3, 3; 5, 1; 7, 1; 71, 1] 2178540[2, 2; 3, 2; 5, 1; 7, 2; 13, 1; 19, 1] 2194800[2, 4; 3, 1; 5, 2; 31, 1; 59, 1] 2209680[2, 4; 3, 4; 5, 1; 11, 1; 31, 1] 2218944[2, 6; 3, 1; 7, 1; 13, 1; 127, 1] 2296476[2, 2; 3, 2; 7, 1; 13, 1; 701, 1] 2316480[2, 6; 3, 1; 5, 1; 19, 1; 127, 1] 2340360[2, 3; 3, 3; 5, 1; 11, 1; 197, 1] 2388960[2, 5; 3, 3; 5, 1; 7, 1; 79, 1] 2467600[2, 4; 5, 2; 31, 1; 199, 1] 2489760[2, 5; 3, 2; 5, 1; 7, 1; 13, 1; 19, 1] 2509920[2, 5; 3, 3; 5, 1; 7, 1; 83, 1] 2530440[2, 3; 3, 4; 5, 1; 11, 1; 71, 1] 2594592[2, 5; 3, 4; 7, 1; 11, 1; 13, 1] 2691360[2, 5; 3, 3; 5, 1; 7, 1; 89, 1] 2699424[2, 5; 3, 2; 7, 1; 13, 1; 103, 1] 2719080[2, 3; 3, 2; 5, 1; 7, 1; 13, 1; 83, 1] 2804160[2, 6; 3, 1; 5, 1; 23, 1; 127, 1] 2815560[2, 3; 3, 4; 5, 1; 11, 1; 79, 1] 2863080[2, 3; 3, 3; 5, 1; 11, 1; 241, 1] 2901696[2, 6; 3, 1; 7, 1; 17, 1; 127, 1] 2915640[2, 3; 3, 2; 5, 1; 7, 1; 13, 1; 89, 1] 2944032[2, 5; 3, 1; 7, 1; 13, 1; 337, 1] 3013920[2, 5; 3, 2; 5, 1; 7, 1; 13, 1; 23, 1] 3171960[2, 3; 3, 4; 5, 1; 11, 1; 89, 1] 3231360[2, 7; 3, 3; 5, 1; 11, 1; 17, 1] 3235680[2, 5; 3, 3; 5, 1; 7, 1; 107, 1] 3243240[2, 3; 3, 4; 5, 1; 7, 1; 11, 1; 13, 1] 3276000[2, 5; 3, 2; 5, 3; 7, 1; 13, 1] 3310800[2, 4; 3, 1; 5, 2; 31, 1; 89, 1] 3328416[2, 5; 3, 2; 7, 1; 13, 1; 127, 1] 3392928[2, 5; 3, 4; 7, 1; 11, 1; 17, 1] 3535680[2, 6; 3, 1; 5, 1; 29, 1; 127, 1] 3666432[2, 9; 3, 1; 7, 1; 11, 1; 31, 1] 3800160[2, 5; 3, 2; 5, 1; 7, 1; 13, 1; 29, 1] 3813480[2, 3; 3, 4; 5, 1; 11, 1; 107, 1] 4021920[2, 5; 3, 3; 5, 1; 7, 2; 19, 1] 4062240[2, 5; 3, 2; 5, 1; 7, 1; 13, 1; 31, 1] 4203360[2, 5; 3, 3; 5, 1; 7, 1; 139, 1] 4241160[2, 3; 3, 4; 5, 1; 7, 1; 11, 1; 17, 1] 4253200[2, 4; 5, 2; 7, 3; 31, 1] 4324320[2, 5; 3, 3; 5, 1; 7, 1; 11, 1; 13, 1] 4357080[2, 3; 3, 2; 5, 1; 7, 2; 13, 1; 19, 1] 4553640[2, 3; 3, 2; 5, 1; 7, 1; 13, 1; 139, 1] 4590432[2, 5; 3, 4; 7, 1; 11, 1; 23, 1] 4687200[2, 5; 3, 3; 5, 2; 7, 1; 31, 1] 5005728[2, 5; 3, 2; 7, 1; 13, 1; 191, 1] 5050080[2, 5; 3, 3; 5, 1; 7, 1; 167, 1] 5412960[2, 5; 3, 3; 5, 1; 7, 1; 179, 1] 5518912[2, 6; 7, 1; 97, 1; 127, 1] 5524200[2, 3; 3, 4; 5, 2; 11, 1; 31, 1] 5542800[2, 4; 3, 1; 5, 2; 31, 1; 149, 1] 5654880[2, 5; 3, 3; 5, 1; 7, 1; 11, 1; 17, 1] 5691600[2, 4; 3, 3; 5, 2; 17, 1; 31, 1] 5864040[2, 3; 3, 2; 5, 1; 7, 1; 13, 1; 179, 1] 5929560[2, 3; 3, 2; 5, 1; 7, 1; 13, 1; 181, 1] 6106464[2, 5; 3, 2; 7, 1; 13, 1; 233, 1] 6158280[2, 3; 3, 1; 5, 1; 19, 1; 37, 1; 73, 1] 6158880[2, 5; 3, 2; 5, 1; 7, 1; 13, 1; 47, 1] 6187104[2, 5; 3, 4; 7, 1; 11, 1; 31, 1] 6187600[2, 4; 5, 2; 31, 1; 499, 1] 6299200[2, 6; 5, 2; 31, 1; 127, 1] 6320160[2, 5; 3, 3; 5, 1; 7, 1; 11, 1; 19, 1] 6379560[2, 3; 3, 4; 5, 1; 11, 1; 179, 1] 6420960[2, 5; 3, 2; 5, 1; 7, 3; 13, 1] 6630624[2, 5; 3, 2; 7, 1; 11, 1; 13, 1; 23, 1] 6656832[2, 6; 3, 2; 7, 1; 13, 1; 127, 1] 6658800[2, 4; 3, 1; 5, 2; 31, 1; 179, 1] 6683040[2, 5; 3, 3; 5, 1; 7, 1; 13, 1; 17, 1] 6756480[2, 7; 3, 3; 5, 1; 17, 1; 23, 1] 6759060[2, 2; 3, 1; 5, 1; 7, 2; 11, 2; 19, 1] 6770400[2, 5; 3, 1; 5, 2; 7, 1; 13, 1; 31, 1] 6911760[2, 4; 3, 1; 5, 1; 31, 1; 929, 1] 6998208[2, 6; 3, 1; 7, 1; 41, 1; 127, 1] 7044840[2, 3; 3, 3; 5, 1; 11, 1; 593, 1] 7193280[2, 6; 3, 1; 5, 1; 59, 1; 127, 1] 7227360[2, 5; 3, 3; 5, 1; 7, 1; 239, 1] 7234920[2, 3; 3, 4; 5, 1; 7, 1; 11, 1; 29, 1] 7242048[2, 6; 3, 4; 11, 1; 127, 1] 7469280[2, 5; 3, 3; 5, 1; 7, 1; 13, 1; 19, 1] 7512912[2, 4; 3, 4; 11, 1; 17, 1; 31, 1] 7590240[2, 5; 3, 3; 5, 1; 7, 1; 251, 1] 7633080[2, 3; 3, 2; 5, 1; 7, 1; 13, 1; 233, 1] 7731360[2, 5; 3, 2; 5, 1; 7, 1; 13, 1; 59, 1] 8134560[2, 5; 3, 3; 5, 1; 7, 1; 269, 1] 8150688[2, 5; 3, 2; 7, 1; 13, 1; 311, 1] 8182944[2, 5; 3, 4; 7, 1; 11, 1; 41, 1] 8222760[2, 3; 3, 2; 5, 1; 7, 1; 13, 1; 251, 1] 8386560[2, 11; 3, 2; 5, 1; 7, 1; 13, 1] 8419600[2, 4; 5, 2; 7, 1; 31, 1; 97, 1] 8517960[2, 3; 3, 4; 5, 1; 11, 1; 239, 1] 8717200[2, 4; 5, 2; 19, 1; 31, 1; 37, 1] 8802624[2, 6; 3, 1; 19, 2; 127, 1] 8910720[2, 7; 3, 2; 5, 1; 7, 1; 13, 1; 17, 1] 8936928[2, 5; 3, 2; 7, 1; 11, 1; 13, 1; 31, 1] 9020256[2, 5; 3, 1; 7, 1; 31, 1; 433, 1] 9041760[2, 5; 3, 3; 5, 1; 7, 1; 13, 1; 23, 1] 9106560[2, 7; 3, 3; 5, 1; 17, 1; 31, 1] 9380448[2, 5; 3, 4; 7, 1; 11, 1; 47, 1] 9587160[2, 3; 3, 4; 5, 1; 11, 1; 269, 1] 9646560[2, 5; 3, 3; 5, 1; 7, 1; 11, 1; 29, 1] 9694080[2, 7; 3, 4; 5, 1; 11, 1; 17, 1] 9709200[2, 4; 3, 3; 5, 2; 29, 1; 31, 1] 9767520[2, 5; 3, 3; 5, 1; 7, 1; 17, 1; 19, 1] 9828000[2, 5; 3, 3; 5, 3; 7, 1; 13, 1] for(x=2, 10000000, if((Mod(2*x^2,sigma(x)) == 0) && (Mod(x, 2) == 1),print(x,factor(x)))) No output returned. This confirms that the output for the preceding GP code are all even. For the congruence $\sigma(x^2) \equiv 0 \pmod {\sigma(x) - 1}$ for(x=2, 1000000,

if((Mod(sigma(x^2),sigma(x)-1) == 0),print(x,factor(x)))) 52686[2, 1; 3, 2; 2927, 1] 54549[3, 2; 11, 1; 19, 1; 29, 1] 318528[2, 6; 3, 2; 7, 1; 79, 1] 520768[2, 6; 79, 1; 103, 1] 696045[3, 1; 5, 1; 7, 2; 947, 1] 925677[3, 2; 163, 1; 631, 1] CONJECTURE If one could show the truth of the following conjecture, then the problem in the hyperlinked MO question would be solved: Conjecture: There is no odd $x > 1$ such that $2x^2 \equiv 0 \pmod {\sigma(x)}$. FINAL QUESTION Would you be able to prove the Conjecture?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2021-04-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019156 | MSE-4099636

Term for example that is counterexample for "Conjecture A AND Conjecture B"

I am looking at something where there are two conjectures, Conjecture A and Conjecture B. I have Example X. It isn't a counterexample for Conjecture A or for Conjecture B but it is a counterexample for "Conjecture A and Conjecture B". That is, if Conjecture A is true, then Example X is a counterexample to Conjecture B; and if Conjecture B is true, then Example X is a counterexample to Conjecture A. It is (alas) not true that Example X implies that Conjecture A exclusive-or Conjecture B is true. It probably isn't natural to bundle Conjectures A and B into Conjecture C, and call Example X a counterexample to Conjecture C. Is there a name for such a counterexample? I have a short note and it would be nice to have a nice title and an answer to the question might help.

论坛问题帖 | no_answers_at_retrieval | JP McCarthy | 2021-04-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019157 | MSE-4100554

Could the discovery of a counterexample to the Unit Conjecture change mathematicians' understanding of spinors in general?

Recently it has been reported that Giles Gardham has found a counterexample to the Unit Conjecture for group rings, as given in https://arxiv.org/abs/2102.11818 (and also in a popular article https://www.quantamagazine.org/mathematician-disproves-group-algebra-unit-conjecture-20210412/) Being from a chemistry background with an interest in spinors (not myself a mathematician), I have been advised that a spinor can be defined, from a mathematician's viewpoint, as "a unit length element of a division algebra". (I do know what division algebras are, but am not a competent algebraist.) Assuming division algebras can contain group rings, then could the discovery of a counterexample to the Unit Conjecture lead to significant changes in the way mathematicians view spinors in general?

论坛问题帖 | has_answers | iSeeker | 2021-04-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019158 | MSE-4110221

Continued fraction for Apéry's constant conjectured by The Ramanujan Machine

Recently the following identity was conjectured by The Ramanujan Machine: $$ \frac{8}{7\zeta(3)}=1-\frac{u_1}{v_1-\frac{u_2}{v_2-\frac{u_3}{v_3-\ddots}}}, $$ where $u_n=n^6$ and $v_n=(2n+1)(3n^2+3n+1)$. It has not yet been proven (authors are not aware of a proof), however this is somewhat similar to known cfrac in this answer, the main difference there being $v_n=(2n+1)(17n^2+17n+5)$. I couldn't prove it (to be expected), but I've found a following observation. The polynomial $(2n+1)(17n^2+17n+5)$ above is in various related articles used in certain recurrence which is satisfied by an integer sequence (see for example section 3 in Apéry's theorem. Thirty years after): $$ (n+1)^3r_{n+1}-(2n+1)(17n^2+17n+5)r_n+n^3r_n=0. $$ But in the paper of A Proof that Euler Missed (footnote page 202) we can find the conjectured polynomial $(2n+1)(3n^2+3n+1)$ to be in similar recurrence as well (after shift on n): $$ (n+1)^3r_{n+1}-2(2n+1)(3n^2+3n+1)r_n-(4n+1)4n(4n-1)r_{n-1}=0. $$ This recurrence is mentioned to be satisfied by integer sequence $r_n=\sum_{k=0}^n\binom{n}{k}^4$ (apparently with $r_0=1$ and $r_1=2$). Can this observation be any helpful in proving the conjecture, or is this just coincidence? Maybe do you see another way to prove it? (the problem has not been open for that long, so who knows...) EDIT: One more instance of the same polynomial, we have $$ \sum_{k=0}^{\infty} \frac{1}{(3k+1)^3}-\sum_{k=0}^{\infty} \frac{1}{(3k+2)^3}=\frac{4\pi^3\sqrt{3}}{243}=\frac{1}{P(1)+\frac{3\cdot 1^6}{P(2)+\frac{3\cdot 2^6}{P(3)+\dots}}} $$ where $P(n)=6n^3-9n^2+5n-1=(2n-1)(3n^2-3n+1)$ (the first equality is well known, the second equality is found in https://lfant.math.u-bordeaux.fr/seminar/slides/2019-11-05T10:00--Henri_Cohen.pdf). Also notice $$ \frac{1}{9}((3n+1)^3+(3n+2)^3)=(2n+1)(3n^2+3n+1). $$

论坛问题帖 | has_answers | Sil | 2021-04-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019159 | MSE-411706

"Goldbach's other conjecture" and Project Euler - writing 1 as a sum of a prime and twice a square

From Problem 46 of Project Euler : It was proposed by Christian Goldbach that every odd composite number can be written as the sum of a prime and twice a square. $$9 = 7 + 2 \cdot 1^2$$ $$15 = 7 + 2 \cdot 2^2$$ $$21 = 3 + 2 \cdot 3^2$$ $$25 = 7 + 2 \cdot 3^2$$ $$27 = 19 + 2 \cdot 2^2$$ $$33 = 31 + 2 \cdot 1^2$$ It turns out that the conjecture was false. What is the smallest odd composite that cannot be written as the sum of a prime and twice a square? As soon as I read this problem, I've immediately fought that the smallest odd number 1 was not on the list of examples. Then I've thought that 1 can be the answer. The only way that I could write 1 close to the problem $1=1+2 \cdot 0^2$, Which is of course false. I've also tried to give 1 as answer, but the site didn't accepted. So how can we write 1 as a sum of a prime and twice a square?

论坛问题帖 | has_answers | moray95 | 2013-06-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019160 | MSE-4118761

A conjecture about binary palindromes and arithmetic derivatives

Corrected question. From the sequence of binary palindromes A006995 (eg. 1001001001001) the sequence of possible gaps between consecutive palindromes contain the elements: S={2,4,6,8,12,16,24,32,48,64,96,128,192,256,384,512,768,1024,1536,2048,3072,4096,6144,8192,12288,16384,24576,49152,...} Compare with A131117 that correspond to the set of positions of the records in the sequence of arithmetic derivatives. T={2,4,6,8,12,16,24,32,48,64,96,128,192,256,384,512,640,768,960,1024,1280,1536,1920,2048,2560,3072,3584,3840,4096,5120,6144,7168,7680,8192,10240,12288,14336,15360,16384,20480,24576,28672,30720,32768,40960,...} This prompts for the conjecture that $S\subseteq T$ A hint of a proof may be to much to ask for, but are there computational results that support or contradicts the conjecture? S comes from gaps of consecutive palindromes in the $100,000,000$ first palindromes >0.

论坛问题帖 | has_answers | Lehs | 2021-04-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019161 | MSE-4132665

Follow-up to question 3121498, asked in February 2019

Let $n = p^k m^2$ be an odd perfect number given in Eulerian form (i.e. p is the special/Euler prime satisfying $p \equiv k \equiv 1 \pmod 4$ and $\gcd(p,m)=1$). Denote the classical sum of divisors of the positive integer x by $\sigma(x)=\sigma_1(x)$. This question (and the answer contained therein) presented a proof of the following proposition, the first conditional on the hypothesis that $\sigma(m^2)/p^k$ is a square, and the second one was unconditional: If $n = p^k m^2$ is an odd perfect number with special prime p, then $m^2 - p^k$ is not a square. In both proofs, the equation $p^k + 1 = 2m$ was obtained, and was shown to lead to a contradiction. Here are my questions: (1) Does the contradiction imply that the inequation $p^k + 1 \neq 2m$ holds for all odd perfect numbers $n = p^k m^2$, in general? (2) If the answer to Question (1) is YES, which of the following inequalities hold in general for an odd perfect number? (a) $p^k + 1 < 2m$; or (b) $p^k + 1 > 2m$? MY ATTEMPT I figured this much from this closely related MathOverflow question, where we tried to disprove the Dris Conjecture (that $p^k < m$) and therefore the Descartes-Frenicle-Sorli Conjecture (that $k=1$) in one fell swoop. However, we were unsuccessful, as Pascal Ochem pointed out that our method required that $p^k \geq 2m - 1$.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2021-05-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019162 | MSE-4142531

Checking a conjecture on prime numbers

Thinking on Oppermann and Goldbach conjectures, it came to me the following conjecture that mixes some aspects of both of them: Let $n\notin\mathbb{P}$; then, there exist some prime numbers p and q such that $n^2-n<p<n^2<q<n^2+n$ such that $p+q=2n^2$ I have been checking first cases "by hand", and I would like to know if there exists an "easy" way to check if the conjecture is true or has some inmediate counterexample. I tried programming something in Excel but it was too messy. Thanks in advance!

论坛问题帖 | has_answers | Juan Moreno | 2021-05-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019163 | MSE-4154926

Subtraction puzzle conjecture

I recently learned about a puzzle called a "Diffy". You take four numbers positioned at the vertices of a square. On the midpoint of each side, you write the difference of the two numbers on the vertices of that side. Once you have done this for all four sides, you connect the four new numbers with a square and repeat. My conjecture is that it always comes to four zeroes, if we repeat to infinity. I have not yet been able to prove this, though I tried with variables w, x, y, z as the corners of the square. Unfortunately, I ended up with math mush. Can someone help me prove or disprove this?

论坛问题帖 | has_answers | Mathemagician314 | 2021-05-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019164 | MSE-4157034

Prove conjecture about circle pairing centers

I'm desperately trying to prove a statement which I need for some other results in my bachelor final project (I already spent several hours on this and it's driving me crazy). I would be very happy if you could help me, and if you can you will get credited in my bachelor final project report. Here are some definitions, theorems and terminology. Definition: a circle pairing f is a Möbius transformation which maps the exterior of a circle C_F to the interior of a circle C_f, and it maps the interior of C_F to the exterior of C_f (and from this follows that it maps C_F onto C_f). Terminology: for a circle pairing f as above, the circle C_f has center C_1, and the circle C_F has center C_2. Theorem: for any given triple of points there is a unique Möbius transformation which maps these three points to any other given triple of points. Theorem: Möbius transformations map circles and lines to circles or lines. The statement I want to prove: For every circle pairing f the circles C_F and C_f which are paired as above can be chosen such that f maps the point at infinity to C_1, and maps C_2 to the point at infinity. Assume that the point at infinity lies in the exterior of both circles which are paired. My question is if you can prove this, either by a totally different way (maybe this is trivial if you look at it from some different perspective or something), or finish my incomplete proof below. Of course you would also answer my question if you can find a counterexample but I doubt you will be able to. Here is what I have so far: (Incomplete) proof: Fix a circle pairing f such that the point at infinity is in neither circle which is paired. In other words, fix a Möbius transformation f such that there exist 2 circles J_1, J_2 such that the point at infinity is in the exterior of both, and such that the exterior of J_2 is mapped to the interior of J_1, and the interior of J_2 is mapped to the exterior of J_1. Say f has coefficients a,b,c,d. We have that f maps the point at infinity to a/c, as is the case for all Möbius transformations. Denote the inverse of f as F, which clearly has coefficients $d,-b,-c,a$. We have that F maps the point at infinity to $-d/c$. This implies that f maps $-d/c$ to the point at infinity. Consider the line through a/c and $-d/c$, and choose some complex number H which lies on this line, between

a/c and $-d/c$ (so formally such that the distance from H to both a/c and $-d/c$ is strictly smaller than the distance between a/c and $-d/c$). Given that we already know that $inf->a/c, -d/c->inf$ and $H=/=a/c,-d/c$, we have that H is mapped to some complex number J (Side note: by a theorem above we have that all information about the circle pairing has to be contained in the fact that inf->a/c, $-d/c$->inf, H->J, so H and J would somehow have to capture information about circles around a/c and $-d/c$). Define the circle C_F through H with center $-d/c$. We have that C_F is mapped to some circle C_f through J (by one of the theorems above it is mapped to a line or circle, but because $-d/c$ is not on the circle it can only be mapped to a circle, and because $H->J$ it must be a circle through J; FYI a line is a circle through the point at infinity). If we can show that this image circle always has a/c as center we are done as by some theorem in topology it follows that then f is a circle pairing where the circles which are paired are C_F and C_f. Possible leads which might help with finishing the proof (from huge leads to less promising leads): Maybe we can argue using the inverse map F, which is also a circle pairing, hence it should behave similarly to f. As f maps C_F to some circle C_f through J we have that F maps C_f to C_F. Furthermore we have because F is also a circle pairing, that we can consider the circle C_g through J (confusing notation maybe) with as center a/c and say that it is mapped to some circle through H. Now we have to show that $C_g=C_f$. Maybe this reasoning is even better as it could imply something for f directly, but I'm not sure. Something related to symmetry. If it somehow follows that by symmetry there must be some other point on C_F which is mapped to the reflection of J in the line through a/c and $-d/c$, and another point on C_F is mapped to some other point on this line, then maybe that could help. Something related to the side note in my incomplete proof above. Maybe there is some shortcut by using this in combination with some other theorem I don't know about. It should be possible to argue that a/c must lie inside C_f by the fact that we are considering a circle pairing. I.e. use the fact that we know that at least some circles are paired, and because the point at infinity is in neither is is mapped to the interior of the other circle. Something related to the distance between H and the other intersection point of the line through a/c and $-d/c$ which is on C_F. This might give a clue on the diameter of the image circle. The line through a/c, $-d/c$ and H is mapped to a line through J and the image of a/c. For the image of a/c we know that it is mapped to some number W which has smaller distance to the attracting fixed point of f. The only line which can be preserved is the line through a/c and $-d/c$ (as the point at infinity lies on every line, is mapped to a/c and $-d/c$ is mapped to the point at infinity). If the attracting fixed point lies on the line through a/c and $-d/c$ then it is mapped to the line through J and that fixed point. Thanks in advance if you are willing to help :)

论坛问题帖 | has_answers | Joris | 2021-05-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019165 | MSE-4190318

Difference between hypotenuse and larger leg in a Pythagorean triple

I've been number crunching irreducible Pythagorean triples and this pattern came up: the difference between the hypotenuse and the larger leg seems to always be n^2 or $2n^2$ for some integer n. Moreover, every integer of the form n^2 or $2n^2$ is the difference between hypotenuse and the larger leg for some irreducible Pythagorean triple. Is there a simple proof for that? There is a result listed on Wikipedia that looks kinda, sorta related: that the area of a Pythagorean triangle can not be the square or twice the square of a natural number. EDIT: Actually, the statements are correct only for odd n^2 (but also by any $2n^2$ as stated) as John Omielan demonstrated below.

论坛问题帖 | has_answers | JCCyC | 2021-07-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019166 | MSE-4190740

Is it true that $\frac{a_1^2}{a_2}+\frac{a_2^2}{a_3}+\cdots+\frac{a_n^2}{a_1}\geq\sqrt{n(a_1^2+a_2^2+\cdots+a_n^2)}$ for positive real a_i?

This problem is my attempt to generalize an easier problem: Let $a_1,a_2,\dots,a_n$ be n positive real numbers. Prove or disprove that: $$\frac{a_1^2}{a_2} + \frac{a_2^2}{a_3} + \dots + \frac{a_n^2}{a_1} \geq \sqrt{n \left(a_1^2 + a_2^2 +\cdots+a_n^2 \right)}.$$ I was able to prove that the inequality holds for $n \leq 4$ using AM-GM and Cauchy-Schwarz but was stuck with $n \geq 5$ after much effort. I don't know if this generalization is true and if it really is then it'll be great if someone can show me some hints to prove it. Edit: To show my effort, I have included my solution to the case $n=4$ (for $n \leq 3$, a similar method can be applied) For $n=4$, we need to prove that: $$ \frac{a_1^2}{a_2} + \frac{a_2^2}{a_3} + \frac{a_3^2}{a_4} + \frac{a_4^2}{a_1} \geq \sqrt{4 \left(a_1^2 +a_2^2 + a_3^2 + a_4^2 \right)} $$ By using the Cauchy-Schwarz inequality, we have: $$ \left(a_1^2a_2 + a_2^2a_3 + a_3^2a_4 + a_4^2a_1 \right) \left(\frac{a_1^2}{a_2} + \frac{a_2^2}{a_3} + \frac{a_3^2}{a_4} + \frac{a_4^2}{a_1} \right) \geq \left(a_1^2 + a_2^2 + a_3^2 + a_4^2 \right)^2 $$ We also have: $$ a_1^2a_2 + a_2^2a_3 + a_3^2a_4 + a_4^2a_1 \leq \sqrt{\left(a_1^2 + a_2^2 + a_3^2 + a_4^2 \right) \left(a_1^2a_2^2 + a_2^2a_3^2 + a_3^2a_4^2 + a_4^2a_1^2 \right) }$$ Note that: $$ a_1^2a_2^2 + a_2^2a_3^2 + a_3^2a_4^2 + a_4^2a_1^2 = \left(a_1^2 + a_3^2 \right) \left(a_2^2 + a_4^2 \right) \leq \frac{\left(a_1^2+a_2^2+a_3^2+a_4^2 \right) ^2}{4} $$ $$\Longrightarrow \sqrt{\left(a_1^2 + a_2^2 + a_3^2 + a_4^2 \right) \left(a_1^2a_2^2 + a_2^2a_3^2 + a_3^2a_4^2 + a_4^2a_1^2 \right) } \leq \sqrt{\frac{\left(a_1^2 + a_2^2 + a_3^2 + a_4^2 \right)^3}{4}}$$ Therefore: $$ \frac{a_1^2}{a_2} + \frac{a_2^2}{a_3} + \frac{a_3^2}{a_4} + \frac{a_4^2}{a_1} \geq \frac{\left(a_1^2 + a_2^2 + a_3^2 + a_4^2 \right)^2}{\sqrt{\dfrac{\left(a_1^2 + a_2^2 + a_3^2 + a_4^2 \right)^3}{4}}} = \sqrt{4 \left(a_1^2 +a_2^2 + a_3^2 + a_4^2 \right)} \text{ (QED)} $$

论坛问题帖 | has_answers | NVA | 2021-07-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019167 | MSE-4191843

Another 6 points circle associated with a triangle?

Consider a triangle ABC and a point D. For each of the triangles ABD, BCD, ACD construct the inscribed and circumscribed circles. Conjecture: For any triangle ABC there always exists a special location of D, so that six intersection points of these six circles are concyclic. Is this actually true or not? This construction gives us 12 intersection points , but we are only interested in the "orange" ones and not in the "green" ones. In other words, we only consider the first intersection points of the arches ADB,BDA,BDC,CDB,ADC,CDA with the inscribed circles. Geogebra dynamic sketch.

论坛问题帖 | has_answers | A Z | 2021-07-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019168 | MSE-4192815

Conjecture about the density of primes

Conjecture For any sufficiently large integer kn , the sequence representing the number of primes in each block obtained by splitting kn into k equal blocks, is a strictly decreasing sequence, i.e: $$\pi\left(n\right)>\pi\left(2n\right)-\pi\left(n\right)>\pi\left(3n\right)-\pi\left(2n\right)\ldots>\pi\left(kn\right)-\pi\left(\left(k-1\right)n\right)$$ My approach For any given k, we need to prove that for all sufficiently large n, the last block contains less primes than the block before:
$$\underbrace{\pi\left(\left(k-1\right)n\right)-\pi\left(\left(k-2\right)n\right)}_{\text{before last}}>\underbrace{\pi\left(kn\right)-\pi\left(\left(k-1\right)n\right)}_{\text{last}}$$ Let $f(k,n)$ denote the difference between the prime count of before last and last block of kn:
$$f\left(k,n\right)= \left(\pi\left(\left(k-1\right)n\right)-\pi\left(\left(k-2\right)n\right)\right)-\left(\pi\left(kn\right)-\pi\left(\left(k-1\right)n\right)\right)$$
$$=2\pi\left(\left(k-1\right)n\right)-\pi\left(\left(k-2\right)n\right)-\pi\left(kn\right)$$ Therefore, an equivalent formulation of the conjecture is that $f(k,n)>0$ for any integers k,n with sufficiently large n. Does anybody have ideas or references on how to prove this? $$$$ Computation of $f(k,n)$ I made some computation that strongly suggests the conjecture is true, at least for small k: Failing case for small n Let's take the case $k=2$. We see that the conjecture fails for $n=1,2,4$ & 10: I computed $f(k,n)$ for $2 ≤ k ≤ 30$ with $1 ≤ n ≤ 10^{8}$, in order to find what appears to be the last failing case of each k , after which $f(k,n) >0$ for all n:

论坛问题帖 | has_answers | François Huppé | 2021-07-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019169 | MSE-4193282

π found in Minesweeper - a new conjecture

Playing with Monte Carlo simulations over Minesweeper, I happened by chance to find π appearing in a pattern of the numbers relating to cells neighboring mines on $n\times n$ grids with n mines. My observations point to the following conjecture: We define S_n as the average sum of all numbers on a quadratic Minesweeper board with dimensions $n\times n$, and a number of mines equal to n. We define S_{n-1} as the average sum of all numbers on a quadratic Minesweeper board with dimensions $(n-1) \times (n-1)$, and a number of mines equal to $n-1$. Subtracting S_{n-1} from S_n yields approximately $7.85398 \approx 2.5\pi$. The full conjecture is given in a PDF here: Github link I, being a mere computer scientist, do not have enough insight into the topic, so I have discussed this with a professor of mathematics at my university. He thinks the edge effects are not that important, and suggests looking closer at the overlaps, further suggesting looking at the Minesweeper grid as a torus. What are your thoughts on this curious occurrence of π? Do you find any flaws in my conjecture?

论坛问题帖 | has_answers | Henrik Hillestad Løvold | 2021-07-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019170 | MSE-4197182

Does the following lower bound improve on $I(q^k) + I(n^2) > 3 - \frac{q-2}{q(q-1)}$, where $q^k n^2$ is an odd perfect number? - Part II

Preamble: This question is an offshoot of these earlier posts: (post1), (post2). Let $N = q^k n^2$ be an odd perfect number with special prime q satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. Define the abundancy index $$I(x)=\frac{\sigma(x)}{x}$$ where $\sigma(x)$ is the classical sum of divisors of x. Since q is prime, we have the bounds $$\frac{q+1}{q} \leq I(q^k) < \frac{q}{q-1},$$ which implies, since N is perfect, that $$\frac{2(q-1)}{q} < I(n^2) = \frac{2}{I(q^k)} \leq \frac{2q}{q+1}.$$ By considering the negative product $$\bigg(I(q^k) - \frac{2(q-1)}{q}\bigg)\bigg(I(n^2) - \frac{2(q-1)}{q}\bigg) < 0,$$ since we obviously have $$\frac{q}{q-1} < \frac{2(q-1)}{q},$$ then after some routine algebraic manipulations, we arrive at the lower bound $$I(q^k) + I(n^2) > 3 - \frac{q-2}{q(q-1)} = \frac{3q^2 - 4q + 2}{q(q - 1)}.$$ In (post2), I derived the following inequality: $$2qn^2 - q\sigma(n^2) < 2n^2 - 1.$$ Using my method, I then get the bounds $$I(q^k) < \dfrac{q}{q-1} < \dfrac{2(q-1)}{q} + \dfrac{1}{qn^2} < I(n^2)$$ which implies that the product $$(I(q^k) - x)(I(n^2) - x) < 0$$ is negative, where $$x = \dfrac{2(q-1)}{q} + \dfrac{1}{qn^2}.$$ Upon some algebra, I get $$I(q^k) + I(n^2) > \dfrac{3q^2 - 4q + 2}{q(q - 1)} - \dfrac{q}{(q - 1)(2qn^2 - 2n^2 + 1)} + \dfrac{1}{qn^2}.$$ But $$- \dfrac{q}{(q - 1)(2qn^2 - 2n^2 + 1)} + \dfrac{1}{qn^2} = \dfrac{qn^2 (q - 4) + q + 2n^2 - 1}{qn^2 (q - 1)(2n^2 (q - 1) + 1)} > 0,$$ since q is a prime satisfying $q \equiv 1 \pmod 4$ implies that $q \geq 5$. So it does appear that the inequality $$I(q^k) + I(n^2) > \dfrac{3q^2 - 4q + 2}{q(q - 1)} - \dfrac{q}{(q - 1)(2qn^2 - 2n^2 + 1)} + \dfrac{1}{qn^2}$$ is unconditionally true, which would mean that the new lower bound for $I(q^k) + I(n^2)$ improves on the old. Note that we were able to prove this analytically. And lastly: Based on this answer to a closely related question, since we appear to have obtained an improved lower bound for $I(q^k) + I(n^2)$, can we then say that there is indeed an integer a such that $k \leq a$?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2021-07-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019171 | MSE-4197351

Does $\sigma(n^2)/q \mid q^k n^2$ imply $\sigma(n^2)/q \mid n^2$, if $q^k n^2$ is an odd perfect number with special prime q?

Denote the classical sum of divisors of the positive integer x by $\sigma(x)=\sigma_1(x)$. Denote the abundancy index of x as $I(x)=\sigma(x)/x$. I discovered an interesting identity involving divisors of odd perfect numbers given in the Eulerian form $N = q^k n^2$ today (July 13, 2021). (Recall that an odd perfect number $N = q^k n^2$ is said to be given in Eulerian form if q is the special prime satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$.) The identity is: Proposition: If $N = q^k n^2$ is an odd perfect number with special prime q, then $$N\cdot\Bigg(I(n^2) - \frac{2(q - 1)}{q}\Bigg) = \frac{\sigma(n^2)}{q}.$$ Proof: Our starting point is the following blog post, where it is proved that $$\gcd(n^2, \sigma(n^2)) = 2(1 - q)n^2 + q\sigma(n^2).$$ However, note that we have $$\frac{\sigma(n^2)}{q^k} = \gcd(n^2, \sigma(n^2)).$$ These equations are equivalent to $$2(1 - q)n^2 + q\sigma(n^2) = \frac{\sigma(n^2)}{q^k}.$$ Factoring out qn^2 on the LHS, we obtain $$qn^2 \Bigg(I(n^2) - \frac{2(q - 1)}{q}\Bigg) = \frac{\sigma(n^2)}{q^k}.$$ Multiplying both sides of the last equation by q^{k-1}, we get $$N\cdot\Bigg(I(n^2) - \frac{2(q - 1)}{q}\Bigg) = \frac{\sigma(n^2)}{q}.$$ (Note that the RHS of the last equation is an odd integer.) This concludes our proof. QED. In particular, we have $$\frac{N}{\sigma(n^2)/q} = \frac{1}{\Bigg(I(n^2) - \frac{2(q - 1)}{q}\Bigg)}.$$ But we also know from the following MSE post that $$I(n^2) - \frac{2(q - 1)}{q} = \frac{2(q - 1)}{q\bigg(q^{k+1} - 1\bigg)}.$$ This means that we obtain $$\frac{1}{\Bigg(I(n^2) - \frac{2(q - 1)}{q}\Bigg)} = \frac{q\bigg(q^{k+1} - 1\bigg)}{2(q - 1)} = \frac{q\sigma(q^k)}{2}.$$ But $\sigma(q^k) \equiv k + 1 \pmod 4$, since $q \equiv 1 \pmod 4$, and since $k \equiv 1 \pmod 4$, then $$\sigma(q^k) \equiv k + 1 \equiv 2 \pmod 4.$$ This finding implies that $\sigma(n^2)/q$ divides $N = q^k n^2$. Here is my: QUESTION: Does $\sigma(n^2)/q \mid q^k n^2$ imply that $\sigma(n^2)/q \mid n^2$? If not, under what condition(s) does this implication hold? Note that a proof for $\sigma(n^2)/q \mid n^2$ would imply the Descartes-Frenicle-Sorli Conjecture that $k=1$.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2021-07-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019172 | MSE-4204865

Conjecture about the Carmichael function

Let $\ \lambda(n)\ $ denote the Carmichael-function and define $\ n(e)\ $ to be the number of solutions of $\ \lambda(m)=e\ $. For $\ e\ge 3\ $, we have $\ 4\mid n(e)\ $ I chose "$\ e\ $" because we can concentrate on then even numbers because clearly for odd $\ e\ge 3\ $, there is no solution of $\ \lambda(m)=e\ $, hence $\ n(e)=0\ $ in this case. For $\ e=2\ $, we have $\ 6\ $ solutions , the divisors of $\ 24\ $ except $\ 1\ $ and $\ 2\ $. But for $\ 4\le e\le 3\ 000\ $, we have $\ 4\mid n(e)\ $ as the conjecture states ($\ n(e)=0\ $ can occur as well). Can we prove this conjecture ? The possible prime factors $\ p\ $ of $\ m\ $ have the property $\ p-1\mid e\ $ and $\ p^k\mid m \implies p^{k-1}\cdot (p-1)\mid e\ $. This allows to find all the solutions. With this algorithm (there might be better ones , but I am not aware of anything better), I checked $\ n(e)\ $ upto $\ e=3\ 000\ $ for divisibility by $\ 4\ $. The conjecture would imply a solution to a problem similar to this open conjecture , namely with $\ \lambda(n)\ $ instead of $\ \varphi(n)\ $.

论坛问题帖 | has_answers | Peter | 2021-07-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019173 | MSE-4205210

Conjecture about divergent periodic summations of odd functions

$$f(x)=\lim_{m\to\infty}\frac{\sum_{n=-m}^m(x-n)^{1/3}}{\sum_{n=-m}^m(1-n)^{1/3}}\stackrel{?}{=}x$$ I had a 'proof' but I made the simple mistake of assuming 2 limits could be swapped, so I only really proved that $f(x)=x$ for $x\in[-1,0,1]$, which is pretty trivial. Here's the 'proof' that I had. This was a first draft so it may be confusing to read. The $(\text{stuff})^{1/3}$ in the numerator and denominator is really just $g(\text{stuff})$. Where g just needs to be any odd function and relatively smooth and have a few other properties that I don't remember.

论坛问题帖 | has_answers | William C. | 2021-07-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019174 | MSE-4211805

Why is $\ 100\ $ so special in this context?

Let $\ \lambda(k)\ $ be the Carmichael function and denote $\ N(n)\ $ to be the number of solutions of $\ \lambda(k)=n\ $ Can we prove that $\ N(n)=n\ $ only holds for $\ n=100\ $ (besides the uninteresting case $\ n=0\ $) ? Odd numbers can be ruled out since $\ N(1)=2\ $ and $\ N(u)=0\ $ for every odd $\ u\ge 3\ $. A number $\ n\ $ with $\ n\equiv 2\mod 4\ $ cannot be a solution because of $\ N(2)=6\ $ and for $\ n\ge 6\ $, we can easily show that $\ N(n)\ $ must be divisible by $\ 4\ $. Hence , we can concentrate on the multiples of $\ 4\ $. Maybe someone finds additional restrictions. Using recursive computation of $\ N(n)\ $, I verified the conjecture upto $\ n=10^7\ $. Does anyone have an idea why $\ 100\ $ is so special here ?

论坛问题帖 | no_answers_at_retrieval | Peter | 2021-07-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019175 | MSE-4212753

Is it true that $\frac{D(n^2)}{s(q^k)} > \frac{\sigma(q^k)}{2}$, if $q^k n^2$ is an odd perfect number given in Eulerian form?

Preamble: This post is an attempt to resolve the problem in this earlier question. Let $\sigma(x)$ denote the sum of divisors of the positive integer x. A number M is said to be perfect if $\sigma(M)=2M$. For example, 6 and 28 are perfect since $$\sigma(6) = 1 + 2 + 3 + 6 = 2\cdot{6}$$ and $$\sigma(28) = 1 + 2 + 4 + 7 + 14 + 28 = 2\cdot{28}.$$ It is currently unknown if there are infinitely many even perfect numbers. It is also an open problem whether any odd perfect numbers exist. It is widely believed that there are no odd perfect numbers. Euler proved that an odd perfect number N, if one exists, must necessarily have the so-called Eulerian form $$N = q^k n^2$$ where q is the special/Euler prime satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. UNDERLYING PROBLEM Here is the problem that I am considering: Can the odd perfect number N be of the form

$$\frac{q^k \sigma(q^k)}{2}\cdot{n}?$$ CONTEXT Slowak (1999) proved that the odd perfect number N must be of the form $$\frac{q^k \sigma(q^k)}{2}\cdot{d},$$ where $d > 1$. Dris (2017) showed further that d must have the form $$\frac{D(n^2)}{s(q^k)}=\gcd(n^2,\sigma(n^2))=\frac{\sigma(n^2)}{q^k}=\frac{n^2}{\sigma(q^k)/2},$$ where $D(x)=2x-\sigma(x)$ is the deficiency of x, and $s(x)=\sigma(x)-x$ is the aliquot sum of x. MY ATTEMPT Dris and San Diego (Theorem 5.6, page 22) proved that $$D(n^2) > \frac{q+1}{2}.$$ Since it is not true that $$\frac{D(n^2)}{s(q^k)} \leq D(n^2) \leq \frac{q+1}{2} \leq \frac{\sigma(q^k)}{2},$$ can we then say that $$\frac{D(n^2)}{s(q^k)} > \frac{\sigma(q^k)}{2}?$$ If so, this would imply that $$\frac{\sigma(n^2)}{q^k} = \frac{n^2}{\sigma(q^k)/2} > \frac{\sigma(q^k)}{2}$$ which would mean that $$n > \frac{\sigma(q^k)}{2}.$$ In particular, we have that $$\frac{2n}{\sigma(q^k)} > 1$$ which implies that $$\frac{\sigma(n^2)}{q^k} = \frac{2n^2}{\sigma(q^k)} > n > \frac{\sigma(q^k)}{2}.$$ This appears to prove that $\sigma(n^2)/q^k \neq n$. Here is my: QUESTION: Does this proof hold water? If not, how can it be mended so as to produce a valid proof?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2021-07-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019176 | MSE-4213383

Is this proof regarding the nonexistence of odd perfect numbers correct?

Preamble: This post is an offshoot of this earlier question, which was not so well-received in MathOverflow. Let $\sigma(x)=\sigma_1(x)$ denote the classical sum of divisors of the positive integer x. Denote the abundancy index of x by $I(x)=\sigma(x)/x$. If m is odd and $\sigma(m)=2m$, then m is called an odd perfect number. Euler proved that a hypothetical odd perfect number must necessarily have the form $$m = q^k n^2$$ where q is the special/Euler prime satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. As proved in MO, we have the following equation: $$I(n^2) - \frac{2(q - 1)}{q} = \frac{1}{q^{k+1}}\cdot{I(n^2)},$$ whereupon, starting from the lower bound $$I(n^2) > \frac{2(q - 1)}{q}$$ we get the recursive estimates $$I(n^2) > \frac{2(q - 1)}{q} + \frac{1}{q^{k+1}}\cdot\frac{2(q - 1)}{q}$$ $$I(n^2) > \frac{2(q - 1)}{q} + \frac{1}{q^{k+1}}\bigg(\frac{2(q - 1)}{q} + \frac{1}{q^{k+1}}\cdot\frac{2(q - 1)}{q}\bigg)$$ $$= \frac{2(q - 1)}{q}\cdot\Bigg(1 + \frac{1}{q^{k+1}} + \bigg(\frac{1}{q^{k+1}}\bigg)^2\Bigg)$$ $$I(n^2) > \frac{2(q - 1)}{q} + \frac{1}{q^{k+1}}\Bigg(\frac{2(q - 1)}{q}\cdot\Bigg(1 + \frac{1}{q^{k+1}} + \bigg(\frac{1}{q^{k+1}}\bigg)^2\Bigg)\Bigg)$$ $$= \frac{2(q - 1)}{q}\cdot\Bigg(1 + \frac{1}{q^{k+1}} + \bigg(\frac{1}{q^{k+1}}\bigg)^2 + \bigg(\frac{1}{q^{k+1}}\bigg)^3\Bigg)$$ $$\ldots$$ $$\ldots$$ $$\ldots$$ Repeating the process ad infinitum, we get: $$I(n^2) > \frac{2(q - 1)}{q}\cdot\Bigg(\sum_{i=0}^{\infty}{\bigg(\frac{1}{q^{k+1}}\bigg)^i}\Bigg).$$ But $$\sum_{i=0}^{\infty}{\bigg(\frac{1}{q^{k+1}}\bigg)^i}$$ is an infinite geometric series, with sum $$\frac{a_0}{1 - r}$$ where the first term $a_0 = 1$ and the common ratio $$r = \frac{1}{q^{k+1}}.$$ Hence, we obtain $$\sum_{i=0}^{\infty}{\bigg(\frac{1}{q^{k+1}}\bigg)^i} = \frac{1}{1 - \frac{1}{q^{k+1}}},$$ from which we finally get $$I(n^2) > \frac{2(q - 1)}{q}\cdot\frac{q^{k+1}}{q^{k+1} - 1}.$$ But we can simplify the RHS of the last inequality as follows: $$\frac{2(q - 1)}{q}\cdot\frac{q^{k+1}}{q^{k+1} - 1} = \frac{2q^k (q - 1)}{q^{k+1} - 1} = \frac{2q^k}{\sigma(q^k)} = \frac{2}{I(q^k)} = I(n^2).$$ We have therefore finally arrived at the contradiction $$I(n^2) > I(n^2).$$ We therefore conclude that there cannot be any odd perfect numbers. Here is my: QUESTION: Does this proof hold water? If it does not, where is the error in the argument, and can it be mended so as to produce a valid proof?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2021-07-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019177 | MSE-4213766

New method for constructing six-point circles?

I would like to present a circle associated with a triangle that can be constructed as follows: Let A' be the first intersection point of the bisector of angle A with the inscribed circle of the triangle ABC and define B' and C' cyclically. I is the incenter of the triangle ABC. Then circumcircles of the triangles A'B'I, B'C'I, A'C'I intersect the sides of the original triangle at six points, that are always concyclic: Geogebra dynamic sketch I wonder what triangle centres (if any) might be lying on this "red circle" and what is known about it? Point Z (the center of the "red circle" that is passing through six points) is not included into the Clark Kimberling's Encyclopedia... Mathworld.wolfram.com doesn't mention it as well. P.S. The same construction can be also applied to other remarkable circles: Geogebra dynamic sketch for the Taylor point/circle. Geogebra dynamic sketch for a Cevian circle of the Symmedian point. <slightly different construction principle was applied here, but overall it is the same idea. X(6) is not the center of the Symmedian circle> Geogebra dynamic sketch for the nine point circle So perhaps this principle holds true for all central circles and there might be some sort of a generalisation?

论坛问题帖 | has_answers | A Z | 2021-07-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019178 | MSE-4216498

Conjecture:$\int_{0}^{\infty} \frac{\sin^3(x)\sin(3\pi x)}{x^4} J_0(x)^2\text{d}x \overset{?}{=} \frac{\pi}{2}$

I found $$\int_{0}^{\infty} \frac{\sin^3(x)\sin(3\pi x)}{x^4} J_0(x)^2\text{d}x=1.570796... \overset{?}{=} \frac{\pi}{2}$$ Where $J_0(x)$ is the 0th Bessel functions of the first kind. I try to use the following identity, $$\int_{-\infty}^{\infty} J_0(x)e^{-i\omega x}\text{d}x =\frac{1}{\sqrt{1-\omega^2} }$$ And $$\int_{-\infty}^{\infty} f(x)g(x)e^{i\omega x}\text{d}x =\frac{1}{2\pi} \int_{-\infty}^{\infty} F(x)G(\omega-x)\text{d}x$$ Where $F(x)=\int_{-\infty}^{\infty} f(t)e^{-itx}\text{d}t, G(x)=\int_{-\infty}^{\infty} g(t)e^{-itx}\text{d}t$. But the integral $\int_{-\infty}^{\infty} J_0(x)^2e^{i\omega x}\text{d}x$ involves $K(k)$(Complete elliptic integral of the first kind).So it's not a good way to prove it. Now I don't have any idea to prove it. Are there any hints? Or what techniques should I use?

论坛问题帖 | has_answers | Sakup2485 | 2021-08-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019179 | MSE-4217718

Conjecture: there is only one configuration of d lines in n dimensional space where any pair have same angle and d is the max possible.

The sequence here: http://oeis.org/A002853 tells us the maximum number of lines that can be placed in d dimensional space in a way that any pair of lines has the same angle as any other pair and all of them pass through the origin. In general, there can be multiple configurations with n lines that satisfy this property. For example, when $d=3$ and we want to place 3 lines, we can have the coordinate system (x, y and z axes) or three lines at $3 \pi/2$ radians restricted to a plane. So, there are at least two such configurations. I conjecture that when we have the maximum possible number of lines, there will be only one configuration possible. Is it possible to prove this conjecture or refute with counter-example? Related: Number of configurations of lines in d dimensional space where any pair of them have the same angle. How to draw six lines in 3-d space such that angle between any pair of them is the same.

论坛问题帖 | no_answers_at_retrieval | Rohit Pandey | 2021-08-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019180 | MSE-4230387

What is known about this extension of the Erdös-Strauss-conjecture?

The Erdös-Strauss conjecture states that $$\frac{4}{n}=\frac{1}{a}+\frac{1}{b}+\frac{1}{c}$$ has a solution in positive integers for every integer $n>1$. What is known about this extension : The equation $$\frac{k}{n}=\frac{1}{a}+\frac{1}{b}+\frac{1}{c}$$ with $k=5,6$ or 7 and any integer $n\ge 3$ has a solution in positive integers. In the case $k=5$, I worked out that a counterexample is only possible, if n is of the form $20l+1$ and in this case, the conjecure holds upto $n\le 2\cdot 10^6$. I neither found counterexamples for $k=6$ and $k=7$. $k=8$ is no more possible , since $\frac{8}{11}$ has not a representation of the desired form. What is known about this extension ? Any ideas or references ?

论坛问题帖 | no_answers_at_retrieval | Peter | 2021-08-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019181 | MSE-4235579

Proving formula about the least number n such that $a|n$ and $b|(n+1)$, a,b coprime.

I have been looking for theorems that say something about relations of factors between consecutive numbers. With Bezout's identity it can be proved that if $\gcd(a,b)=1$ there is always a number n such that $a|n$ and $b|(n+1)$, which tells something about independency between factors of consecutive numbers. After a lot of work I found a simple formula that seems to determine the least non-negative number n as above, $n=b\cdot r^{-1}-1$, where r^{-1} is the number corresponding to the inverse of r in the multiplicative group of numbers coprime to a and r is the reminder when b is divided with a. Obviously $b|(n+1)$ but how to prove that $a|n$ and that there are no smaller number than n?

论坛问题帖 | has_answers | Lehs | 2021-08-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019182 | MSE-4236253

Prime Algorithim Conjecture

Let g be the greatest factor of a positive integer(a) where $g \neq a$ and l be the smallest factor such that $gl = a$. Let t_i be the ith term of the sequence where the starting term is $\frac{g_1!}{(a_1)^n}$ and n is the greatest positive integer such that $(a_i)^n|g_i!$. More specifically the sequence/recursive algorithim $t_i = \frac{g_{i}! }{(t_{i - 1})^n}$ For example, when a_1 is 10: $t_1 = \frac{5!}{10^n}$ Here $n = 1$, so $t_1 = \frac{5!}{10} \implies t_1 = 12$ Then $t_2 = \frac{6!}{12^n}$ In which we find $n = 2$, giving $t_2 = \frac{6!}{144} \implies t_2 = 5$ So here the sequence stops, as t_2 is prime and $g \neq a$. Its interesting to note that n is the number of pairs of the form $a = mn$ where m and n are factors of a. Now take a look at the number where a_1 is 14, $t_1=7!/141 \implies t_1 = 360 \implies t_2=180!/360^n$ You'll notice the explosive growth that it exhibits when put into the algorithm. This makes it seem skeptical this algorithm will always end at a prime number but consider the following: Lets say we have a number $t_{i -1}$ with over 10 factors after the fraction cancels out and each one has a very large prime factorization with many distinct primes in each. Now note that the prime factorization of each factor allows us to spread the primes across the other factors and combine them in many different ways. Lets say we had a function called $F(x)$ that determines the number of factors that can be possibly made from t_{i-1}. Then if we could prove that $F(x)$ at no point contains more than y of $g_i!$ factors($G(x)$) such that $t_{i} > t_{i - 1}$ than there will be numbers of a_{1} in which the algorithm will fail. But on the other hand if it does, and it can be proven that for some a_{i} for any a_1 $F(x) > \frac{G(x)}{y}$ (In this case $t_{i} < t_{i - 1}$ is true) than the algorithm can still end with a prime. Note that every consecutive term is co-prime and that we consider that a sequence terminates when t_i is prime. Given any value of g and l, where a_i is not a perfect square, this algorithm always terminates. Prove or disprove this.

论坛问题帖 | has_answers | SpoonedBread | 2021-08-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019183 | MSE-4241915

The 'inscribed-circumscribed' 6 point circle

Here is a construction I came upon recently: A'B'C' is the contact triangle. X(1) is the Incenter. A'',B'',C'' are the midpoints of the sides of the contact triangle. A''',B''',C''' are the midpoints of the segments AX(1), BX(1), CX(1). Then quadrilaterals A''A'''C''C''', B''B'''C''C''', A''A'''B''B''' are cyclic. Let Ob, Oa, Oc be the circumcenters of these 3 quadrilaterals. Finally by drawing circles with centers at Oa, Ob, Oc and passing through the incenter we get the 'inscribed-circumscribed' six point circle with its center at the midpoint of X(1) and X(3) - the point X(1385) in the ETC: There is an even easier way to get a smaller concentric six point circle: circumcircles of the quadrilaterals A''A'''C''C''', B''B'''C''C''', A''A'''B''B''' cut the sides of the triangle ABC at six concyclic points. The mere fact that the point X(1385) is already

described in the ETC supposedly makes the construction not particularly inspiring for a geometer. However as this 'X(1385)-circle' has never been previously mentioned (?) in the literature, there might still be a chance that it has some 'nice' properties of its own. Specifically I am interested whether any Kimberling center belongs to this 'inscribed-circumscribed' circle? Or, alternatively, does its central function correspond to any known center?

论坛问题帖 | has_answers | A Z | 2021-09-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019184 | MSE-4247808

Does Diophantine equation $1+n+n^2+\dots+n^k=2m^2$ have a solution for $n,k \geq 2$?

When studying properties of perfect numbers (specifically this post), I ran into the Diophantine equation $$ 1+n+n^2+\dots+n^k=2m^2, n\geq 2, k \geq 2. $$ Searching in range $n \leq 10^6$, $k \leq 10^2$ yields no solution. So I wonder if there are any solutions, and if not, is there some elementary reason for that? Or if it can be converted to some known open problem, that would do too... Some thoughts: Of course if we allow $n=1$ or $k=1$ we could get trivial solutions. We can also see by mod 2 that both n and k must be odd. Now we could try to solve some small cases such as $k=3$ or $k=5$... So set $k=3$ and try to solve $$1+n+n^2+n^3=2m^2.$$ Left side factors and hence we want to solve $(n+1)(n^2+1)=2m^2$. Now this imples $4 \mid 2m^2$ and so m is even. Also we can see that $4 \nmid n^2+1$ for any integer n (becase $n^2\equiv 0,1 \pmod 4 $). So all powers of 2 in $2m^2$ except one will divide $n+1$. So let $m=2^t r$ with $2 \nmid r$, then $2m^2=2^{2t+1}r^2$, $2^{2t} \mid n+1$, $2 \mid n^2+1$. So we can put $n=2^{2t}s-1$ with $2 \nmid s$, substitute it back, divide all powers of 2 and we have the equation $$ s(2^{4t-1}s^2-2^{2t}s+1)=r^2. $$ Now the two expressions in the product are coprime, so they both have to be square, and that is farthest I got so far. Also the expression $2^{4t-1}s^2-2^{2t}s+1$ being square has a solution $s=15,t=2$, and I am yet to find another (but of course s is not a square in this case so it does nothing for the original problem).

论坛问题帖 | has_answers | Sil | 2021-09-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019185 | MSE-4257045

$p = (u + v)^2 + uv$ for primes $p = 5k \pm 1$?

I came up with the following conjecture which I am unable to prove. For all primes p which can be represented as either $5k + 1$ or $5k - 1$ for some positive integer k, we can find positive integers u, v such that $p = (u + v)^2 + u \times v$ For example, $31 = 5 \times 6 + 1 = (2 + 3)^2 + 2 \times 3$ Using a computer, I have verified it for $p < 10000$. Is there a general proof for this?

论坛问题帖 | has_answers | MathsLion1234 | 2021-09-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019186 | MSE-4263964

On a possible proof for $p^k < m$, if $p^k m^2$ is an odd perfect number with special prime p

Let $\sigma(x)=\sigma_1(x)$ denote the classical sum of divisors of the positive integer x. If $\sigma(M)=2M$ and M is odd, then M is called an odd perfect number (hereinafter abbreviated as OPN). It is currently unknown if there are any OPNs. Euler proved that a hypothetical OPN, if one exists, must have the so-called Eulerian form $$M = p^k m^2$$ where p is the special/Euler prime satisfying $p \equiv k \equiv 1 \pmod 4$ and $\gcd(p,m)=1$. Dris conjectured in his M. Sc. thesis that $p^k < m$ always holds for an OPN M given in Eulerian form. Consider the following "proof argument" for $p^k < m$: Let $M = p^k m^2$ be an OPN given in Eulerian form. Suppose that we have $m^2 < 5p^k$. Then we have $$\Bigg(m^2 < 5p^k\Bigg) \iff \Bigg(\frac{m^2}{p^k} < 5 \leq p^k\Bigg) \implies \Bigg(m^2 < p^{2k}\Bigg) \iff \Bigg(m < p^k\Bigg).$$ However, in the paper titled Improving the Chen and Chen result for odd perfect numbers, Broughan et al. show that $\sigma(m^2)/p^k \geq 315$. This implies that $$\frac{p^k}{m^2} < \frac{2}{315},$$ which contradicts $$\frac{1}{5} < \frac{p^k}{m^2}.$$ Here is my question: Does this "proof" indeed show that $p^k < m$ holds? If it does not, can it be mended so as to produce a valid argument?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2021-09-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019187 | MSE-4271270

This is a conjecture, refined through previous responses, about sums of positive integers which define sums of the corresponding cubes.

Revising 4255316, the conjecture now is: If $$\sum_{i=1}^{i=n}{a_{i}}, i>2$$, divides $$3\sum{x_{i}x_{j}x_{k}}$$, where the product is summed over the$\binom{n}{3}$distinct triplets drawn from the n suffices, then $$\sum_{y=0}^{y=a}\sum_{i=1}^{i=n}{x_{i}+2yb}$$ divides $$\sum_{y=0}^{y=a}\sum_{i=1}^{i=n}{(x_{i}+2yb)^{3}}$$

论坛问题帖 | no_answers_at_retrieval | Paul Stephenson | 2021-10-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019188 | MSE-4284239

Maximum degree for this set of planar graphs where all neighbours of any point lie on a circle centered on that point

Consider an undirected simple planar graph, with these additional restrictions: The plane in which the graph is embedded has a Cartesian coordinate system, and there's an injective function mapping each node to its pair of coordinates. I.e., no two nodes lie on the same point of the plane. Each node maps to integer coordinates. I.e., the nodes lie on a grid. For any node, call it A, all neighbors of A lie on a circle centered on A. I.e., all neighbors of any node have the same distance to that node. I couldn't draw a graph like this with max degree greater than 4, so I'm wondering whether 4 is indeed the max degree for this set of graphs. If not 4, perhaps it's 5 or 6? Edit: now I mechanically found the maximal

degree of a graph from this set to be 16, so the conjecture is disproved. It would still be nice if someone gave an answer that explains this intuitively.

论坛问题帖 | has_answers | user2373145 | 2021-10-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019189 | MSE-4284406

Colinear and concurrency configuration in triangle involving arbitrary point

I found out this configuration while fooling around with Geogebra. I tried to solve it on my way back home from work but apparently it's not that easy. Is it a well-known theorem in elementary geometry? Let M is an arbitrary point inside the triangle ABC. AM, BM, CM intersects BC, CA, AB at D, E, F respectively. The line passing through A and parallel to BM intersects CM at H. Let K on AC such that $DK \parallel BM$. DH intersect AB at I. The parallel line to CM passed through A intersects BE at L. DL intersects AC at N, $O\in AB$ such that $DO \parallel CM$. Let P, Q be the intersections of HL and AB, AC respectively. EQ intersects PF at R; FN intersects IE at S. Let T be the midpoint of BC. Prove that: i- K, I, M; O, M, N are colinear. ii- AH, EF, KI; AL, EF, ON are concurrent. iii- A, R, S, T are colinear.

论坛问题帖 | no_answers_at_retrieval | Arsenaler | 2021-10-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019190 | MSE-4286427

Prove conjecture about classification of circle pairings

I need help proving or disproving a conjecture related to circle pairings, which I'm trying to prove for my bachelor final project. I first present some needed terminology and context. A circle pairing of circles C and D is a Möbius transformation which maps D to C, maps the exterior of D to the interior of C, and maps the interior of D to the exterior of C. The circles which are paired are not unique. I'm trying to classify circle pairings based on the absolute value of the trace of the corresponding matrices (the group of Möbius transformations is isomorphic to the group of 2x2 invertible matrices modulo plusminus). I have already found the following results: Theorem 1: the most general way any circle pairing that pairs $C(x_1, r_1)$ and $D(x_2, r_2)$ can be written as is $$g(z)=x_1+r_1\cfrac{\bar{v}(z-x_2)+r_2 \bar{u}}{u(z-x_2)+r_2 v},$$ where $u,v\in \mathbb{C}:|u|^2-|v|^2=1$, where x_1, x_2 are the centers and r_1, r_2 are the radii of C and D. Lemma: the most general way any circle pairing f that pairs circles $C(x_1, r_1)$ and $D(x_2, r_2)$ such that $f(x_2)=\infty$ and $f(\infty)=x_1$, can be written as is $$x_1 +\cfrac{(H-x_2)(J-x_1)}{z-x_2},$$ where $H\in D$ and $J\in C$ (so $|H-x_2|=r_2$ and $|J-x_1|=r_1$). Theorem 2: for every circle pairing g given as in Theorem 1, which maps $\tilde{x}_2\mapsto \infty$ and $\infty \mapsto \tilde{x}_1$ there exist circles $C(\tilde{x}_1, r_1/|u|)$ and $D(\tilde{x}_2,r_2/|u|)$ such that g pairs C and D. In other words, every circle pairing as in Theorem 1, can be written in the form of the lemma above. Proof: We denote $\tilde{x}_1=x_1+r_1 \bar{v}/u$, $\tilde{x}_2=x_2-r_2 v/u$, $\tilde{r}_2=r_2/|u|=|H-\tilde{x}_2|$ and $\tilde{r}_1=r_1/|u|=|J-\tilde{x}_1|$, where $H=x_2+\tilde{r}_2$ and $J=\tilde{x}_1+\tilde{r}_1$. We get $$\begin{equation*} \begin{split} \tilde{x}_1 +\cfrac{(H-\tilde{x}_2)(J-\tilde{x}_1)}{z-\tilde{x}_2} & = x_1+r_1 \frac{\bar{v}}{u}+\cfrac{r_1 r_2 / u^2 }{(z-x_2)+r_2 v/u} \\ & = x_1+r_1(\frac{\bar{v}}{u}\frac{1/u [(z-x_2)+r_2 v/u]}{1/u [(z-x_2)+r_2 v/u]}+\cfrac{r_2/u^2}{(z-x_2)+r_2 v/u}) \\ & = x_1+r_1\cfrac{\bar{v}(z-x_2)+\bar{v} r_2 v/u+r_2/u}{u(z-x_2)+r_2 v } \\ & = x_1+r_1 \cfrac{\bar{v}(z-x_2)+r_2(\bar{v}v+1)/u}{u(z-x_2)+r_2 v} \\ & = x_1+r_1\cfrac{\bar{v}(z-x_2)+r_2 \bar{u}}{u(z-x_2)+r_2 v}. \\ \end{split} \end{equation*}$$ The absolute value of the trace of a circle pairing f as in Theorem 1 is $|Tr f|=\cfrac{|\tilde{x}_2-\tilde{x}_1|}{\sqrt{r_1 r_2 /|u|^2}}$, where $f(\tilde{x}_2)=\infty$ and $f(\infty)=\tilde{x}_1$ (so $\tilde{x}_1=x_1+r_1 \bar{v}/u$ and $\tilde{x}_2=x_2-r_2 v/u$). I have shown that if $|Tr f|>2$, then the radii of the circles C and D as given the existence of by Theorem 2 can be chosen such that C and D are disjoint (in short, if the $R_1=r_1/|u|$ and $ R_2=r_2/|u|$ are the radii of the paired circles then it can be shown that sR_2, R_1/s, where $s>0$, are also paired by f). If $|Tr f|=2$, then these circles can be chosen to touch (i.e. intersect in precisely one point), but never be disjoint. If $|Tr f|<2$, then these circles can only be chosen to overlap. The reason this is true is that the value for s for which these circles are the 'most disjoint' is such that the new radii $\hat{R_1}$ and $\hat{R_2}$ are such that $\hat{R_1}=\hat{R_2}=\sqrt{R_1 R_2}$. Circles are disjoint if $|x_1-x_2|>\text{sum of possible radii} \geq 2\sqrt{R_1 R_2}$, from which these conditions can be derived. We call a circle pairing f disjoint if $|Tr f|>2$, touching if $|Tr f|=2$ and overlapping if $|Tr f|<2$. Conjecture: if a circle pairing g is touching, then there exist no disjoint circles C and D such that g pairs these. If a circle pairing g is overlapping, then there exist no disjoint and no touching circles C and D such that g pairs these. In other words, the freedom of how disjoint the circles as given by Theorem 2 can be chosen, is such that it is the 'most disjoint' case of all possible circles that can be chosen to be paired. Theoretically, for some circle pairing g as given by Theorem 1, the points which are mapped from and to ∞ have to just be somewhere in the interiors of $C(x_1,r_1)$ and $D(x_2,r_2)$ which are paired by g. If they are the centers x_1 and x_2, then the classification of the conjecture is trivially true, but it is not clear what happens if they are not. Some things which may help to prove the conjecture which I have already considered but have not succeeded with: Note that the product of the radii of the circles as given by Theorem 2, namely $r_1 r_2 /|u|^2$, is smaller than that of the original radii r_1 and r_2 as given by Theorem 1, as $|u|>1$. Note that each of the fixed point(s) $f_{1,2}=1/2[\tilde{x}_1+\tilde{x}_2\pm \sqrt{4 r_1 r_2/u^2+(\tilde{x}_1-\tilde{x}_2)^2}]$, where again $\tilde{x}_1=x_1+r_1 \bar{v}/u$ and $\tilde{x}_2=x_2-r_2 v/u$, must either be in the interior of PRECISELY one of the circles that are paired, or be the intersection point(s) of these circles (this follows from the definition of circle pairings). Maybe the equation of g as given by Theorem 1 can be rewritten such that it equals itself, but where the values of x_1, x_2, r_1, r_2, u, and v are different. This should theoretically be possible for every pair of circles which g can pair. Maybe this could give insight into the freedom of possible circles which can be chosen to be paired, and why hypothetically the case where the centers are the points that map from and to ∞, would be the 'most disjoint' case. Maybe $|Tr f|$ can be related directly to the 'disjointness' of circles which can be chosen to be paired by g, similarly to how I did it already for the case for which the centers are the points which are mapped from and to ∞. In other words, maybe the logic itself I used to come up with the classification for the simple case, can be generalized to the general case. Hopefully anyone can find a proof or a counterexample.

Thanks in advance!

论坛问题帖 | has_answers | Joris | 2021-10-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019191 | MSE-4290297

New trigamma identity for $\Psi_1(\frac3{20})+6\,\Psi_1(\frac15)+10\,\Psi_1(\frac25)-\Psi_1(\frac1{20})$

I play with Maple, and I find this relation for the trigamma function: $$\begin{align} \Psi_1\left({\frac{3}{20}}\right)+6\,\Psi_1\left(\frac15\right)+ 10\,\Psi_1\left(\frac25 \right)-\Psi_1\left(\frac1{20}\right) &=-96\,{G}-{\frac{24\, \pi^2\sqrt{5}}{5}}+16\,\pi^2\\[0.5em] &\quad-2\pi^2{\frac{15+\sqrt{5}}{\sqrt{10+2\sqrt{5}}}} \end{align}$$ where G is the Catalan's constant. But I don't know if this relation is well-known or not. Please suggest how to prove it. Thanks.

论坛问题帖 | has_answers | Dens | 2021-10-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019192 | MSE-4293214

Is 3 the only prime that is both a Mersenne prime and a Fermat prime?

A Mersenne prime is a prime of the form 2^n-1. A Fermat prime is a prime of the form 2^n+1. Despite the two being superficially very similar, it is conjectured that there are infinitely many Mersenne primes but only finitely many Fermat primes. Here is my inquiry: QUESTION: Is 3 the only prime that is both a Mersenne prime and a Fermat prime? MY ATTEMPT The prime 3 is of the form $$3 = 2^2 - 1 = 2^{2^0} + 1$$ which shows that it is both a Mersenne prime and a Fermat prime. However, I am having trouble proving that it is the only such prime. Edited in response to an earlier closure of this question due to lack of context: (February 08, 2022 - 12:34 AM) My main motivation for having asked this question was to check whether an odd perfect number could be divisible by some prime that is both a Mersenne prime and a Fermat prime. As the excellent answers so show, 3 is the only such prime. However, (and this is the main caveat why I did not include this context earlier) it is currently unknown whether 3, or whether any prime below 105 for that matter, does divide an odd perfect number. To quote (or paraphrase, as I am currently unsure of the exact wording of) another mathematician's remark over at MathOverflow: "Such a result would be a major breakthrough in our understanding of odd perfect numbers."

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2021-11-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019193 | MSE-4293888

A conjecture about non-nilpotent groups

A finite group, G, is nilpotent if its upper central series terminates (at $i \in \mathbb{N}$) with $Z^i(G)=G$, where $Z^i(G)$ is it's i-th center which can be described as $\{x \in G\mid \forall y \in G : [x,y]\in Z_{i-1}(G)\}$ where $[x,y]$ is the commutator of x and y in G (in other words, it is the set of elements that commute with one another up to an element of the $(i-1)$-th center). A finite group, G, is non-nilpotent if it is not nilpotent. In this case, we still have, by finiteness of G that the upper central series will terminate after finitely many steps, but now we have an $i \in \mathbb{N}$ at where $Z^i(G)=Z^{i+1}(G)$, and yet $Z^i(G)\neq G$, but rather a proper subgroup of G. In this case, since the upper centers are all characteristic and thus normal, we can consider the quotient group, $Q=G/Z^i(G)$. This group must be centerless, else we could take the pullback of $Z(Q)$ under the natural projection to obtain $Z^{i+1}(G)\supset Z^i(G)$ with equality not holding, contradicting our assumption that the upper central series stabilized at i. This gives a rather elegant characterization of non-nilpotent finite groups: A non-nilpotent finite group G can be viewed as an extension of a nilpotent normal subgroup, $U=Z^i(G)$ by a centerless quotient group, Q. In the most extreme case, the central series terminates immediately in the trivial group, and you obtain a centerless $G=Q$. This shows that the centerless groups are the primitive elements of the class of non-nilpotent finite groups with the rest being constructed from them by extensions of nilpotent groups by centerless groups. A (finite) group, G, is complete if it is centerless and isomorphic to its automorphism group $G\simeq\text{Aut}(G)$. Conjecture: Suppose we have a non-nilpotent finite group G, whose upper central series terminates at $i>0$, meaning that G is not centerless, and $Z^i(G)=Z^{i+1}(G)$. Let $U=Z^i(G)$, and suppose further that the centerless quotient $Q=G/U$ is complete, meaning $Q\simeq\text{Aut}(Q)$. Lastly, suppose that the upper central series for $\text{Aut}(G)$ terminates at some $i_1\geq i$, that is, the upper central series for $\text{Aut}(G)$ is at least as long as the upper central series for G. Note in particularly that $\text{Aut}(G)$ is non-nilpotent, and so has a centerless quotient Q_1. Then the claim is that the centerless quotients are isomorphic, $Q_1\simeq Q$. I came up with the conjecture based on computer searches in GAP looking at the automorphism series for various small finite groups as far as I could construct them in GAP (many eventually reach very large groups and run into issues with memory or the size of the computation). Looking at the upper central series for each term of the automorphism series and comparing how they changed as you step through the automorphism series lead me to propose that the above conjecture holds in general for finite groups. Naturally a counterexample will be enough to disprove the claim, but I didn't find any looking at over 50 different automorphism series, so it may not be easy to find if it exists. I'm not really sure how to approach trying to prove this either. I don't really know enough about the relationship between a group and its automorphism group to leverage the givens in this conjecture productively. Edit 1: This is a response to David A. Craven's answer: I took the liberty of constructing the proposed group in GAP for $n=7$. To do this, I needed to put all three of the G_i's into permutation groups so that the direct product operation in GAP would also return a permutation group, so that fast algorithms would be available for subsequent calculations. Specifically I used: G1:=SymmetricGroup(7); G2:=Group((1,2,3,4),(1,2)(3,4)); #This is D8 G3:=Group((1,2,3),(4,5,6),(7,8,9)); #This (Z_3)^3 #That G2 and G3 are as claimed can be checked by IdSmallGroup, is should return (8,3) for G2 and (27,5) for G3. #Now we can call DirectProduct and the result will be a permutation group G:=DirectProduct(G1,G2,G3); #Because this is a permutation group, computation of the upper central series is 'easy' for GAP, trying this without doing the above preparations lead to a computation that was taking minutes and still hadn't finished U:=UpperCentralSeriesOfGroup(G); From the above I obtained that G has upper central series of length 2. With $Z(G)\simeq (54,15)$, and $Z^2(G)\simeq (216,151)$ (Where (x,y) is the GAP library id for order x, id y.). GAP

StructureDescription for U[1] confirms that $Z^2(G)\simeq\mathbb{Z}_3^3\times D_8$. Next I constructed the quotient and confirmed it was the symmetric group (necessarily S_7 by an easy order argument). After a computation that took a few seconds I got the automorphism group, $A1$, but it was not in the form of a permutation group, so I used NiceMonomorphism and then NiceObject to obtain a permutation group isomorphic to the automorphism group. This allows for fast(er) algorithms to used for computing its upper central series, which I found to be of length 2 with $Z(A1)\simeq (8,5)$ and $Z^2(A1)\simeq (32,46)$ (again using GAP ids for the groups). The size of $A1$ is 905748480 which is twice that predicted in the answer (Which would be 452874240 if it had the structure given in the answer). I was able to get GAP to give me a structure description for $A1$, which was $\mathbb{Z}_2^2\times D_8\times S_7\times \text{PSL}(3,3)$. I suspect the comment that the proposed automorphism group is, in fact, a subgroup of index 2 in the full automorphism group is correct. The quotient $Q1$ is of order 28304640 (and so obviously not isomorphic to S_7), GAP gives a structure description of $S_7\times \text{PSL}(3,3)$ for this quotient. (So it is still a counter example for $i_1=i$ case).

论坛问题帖 | has_answers | Justin Benfield | 2021-11-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019194 | MSE-4294505

Irreducible polynomials in $\mathbb Z[X]$ with no primes in the image.

It seems intuitive that if an irreducible polynomial $q\in\mathbb Z[X]$ has no primes in it's image, i.e. $n\in\mathbb Z\implies q(n)\notin \mathbb P$, then there is a non unit integer $m\in\mathbb Z$ such that $m|q(n)$ for all $n\in \mathbb Z$. But how to prove it? Example: $q(x)=4x^5-7x^4-4x^3-4x^2+7x+8$ is irreducible and $2|q(n)$ all n. I didn't have a single idea how to manage this, but I had a feeling that a theorem or conjecture was appropriate, but I had no key word, and thats the reason why I tagged reference-request

论坛问题帖 | has_answers | Lehs | 2021-11-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019195 | MSE-4300742

Does $\gcd$ of values of n-th degree primitive integer polynomial always divide $n!$?

Let's consider the following conjecture: Let $f(x)=a_0+a_1x+a_2x^2+\dots +a_nx^n$ be a primitive non-zero polynomial with integer coefficients and let $g=\gcd(\{f(k), k \in \mathbb{Z}\})$, then g divides $n!$. Or equivalently just: Let $0\not\equiv f(x)\in \mathbb{Z}[x]$, then $\gcd(f(k)) \mid c(f)\cdot (\deg f)!$. Here $c(f)$ is the content of a polynomial. Can we prove/disprove this? Edit: Found literature reference Today I found that g is called the fixed divisor of a polynomial and with this I was able to trace the same claim to the article Uber ganzwertige ganze funktionen by George Pólya from 1915. Unfortunately the article is in german so I am not sure how the proof goes (various articles mention this claim and cite this one without providing the proof themselves). Some thoughts I found this based on observation for small degrees. For $n=2$ and $\gcd(a_0,a_1,a_2)=1$ we can prove $g \mid 2$ (see below), and similarly for $n=3$ we can show $g \mid 6=3!$. I also wrote a simple script in Maple to verify over small degrees and coefficients range and haven't found a counterexample yet, so I wonder if this can be proven to hold in general. It can be useful that $\gcd(\{f(k), k \in \mathbb{Z}\})=\gcd(f(0),f(1),\dots,f(n))$ (or any consecutive $n+1$ integers for that matter). Currently I am thinking that Lagrange interpolation could be perhaps useful in generic case, I worked out that if we know f on consecutive integers $i=0,1,\dots,n$, then the Lagrange interpolation implies $$ f(x)=\sum_{i=0}^{n} (-1)^{n-i}\binom{x}{i}\binom{x-i-1}{n-i}f(i). $$ So I guess if we assume $\gcd(f(i))=d$ where $d \not\mid n!$, we should be able to use the above to reach contradiction with $f(x)$ having integer and coprime coefficients, but I wasn't able to. Proof for $n=2$ For $f(x)=a_0+a_1x+a_2x^2$ and $\gcd(a_0,a_1,a_2)=1$, we want to show $\gcd(f(-1),f(0),f(1)) \mid 2$. Now $f(-1)=a_0-a_1+a_2$, $f(0)=a_0$, $f(1)=a_0+a_1+a_2$ and so \begin{align} \gcd(f(-1),f(0),f(1)) &=\gcd(a_0-a_1+a_2,a_0+a_1+a_2,a_0)\\ &=\gcd(-a_1+a_2,a_1+a_2,a_0)\\ &=\gcd(2a_2,a_1+a_2,a_0)\\ \end{align} Now $g=\gcd(2a_2,a_1+a_2,a_0) \mid 2a_2$, but a_2 is coprime to g (otherwise we had a prime $p\mid a_2, p\mid a_1+a_2, p \mid a_0$ and so $p \mid \gcd(a_2,a_1,a_0)=1$, impossible). Hence $\gcd(2a_2,a_1+a_2,a_0) \mid 2$.

论坛问题帖 | has_answers | Sil | 2021-11-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019196 | MSE-4301413

Primes of the form n^2+n+1

Let $f(n)=n^2+n+1$. While experimenting I found that Given $m,n\in\mathbb N, \: n>1$. If $f(2n)\in\mathbb P$ and $f((3m+1)n)\in\mathbb P$ then $3|n$. It has ben tested for $0\le m <1000$ and $1<n<1000$. I've tried to let $n=3k+i$ looking for polynomials of k for $i=1,2$ but it didn't work well. My hope is that someone can prove the conjecture.

论坛问题帖 | has_answers | Lehs | 2021-11-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019197 | MSE-4302235

Can two circles intersect each other at right angles, such that one circle passes through the center of the other circle?

More precisely, do there exist two intersecting circles such that the tangent lines at the intersection points make an angle of 90 degrees, and one of the circles passes through the center of the other? I think the answer is no, and that this can only be the case if the circle passing through the center of the other circle is actually a line. I'm not sure how to show this though. Can someone give a proof or a counterexample?

论坛问题帖 | has_answers | Joris | 2021-11-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019198 | MSE-4303264

Conjectured closed form for ${\it {Li_2}} \left(1-{\frac {\sqrt {2}}{2}}-i \left(1-{\frac {\sqrt { 2}}{2}} \right) \right)$

With Maple i find this closed form: ${\it {Li_2}} \left(1-{\frac {\sqrt {2}}{2}}-i \left(1-{\frac {\sqrt { 2}}{2}} \right) \right)$=$-{\frac {{\pi}^{2}}{64}}-{\frac { \left(\ln \left(1+\sqrt {2} \right) \right) ^{2}}{2}}-{\frac {{\it {Li_2}} \left(-1-\sqrt {2} \right) }{2}}+{\frac {i}{32}} \left(4\,\pi\,\ln \left(1+\sqrt {2} \right) -24\,G \right)$ where G is the Catalan's constant and Li_2 is the dilogarithm function or Spence's function. If you have got any ideas please for proving this identity.

论坛问题帖 | no_answers_at_retrieval | Dens | 2021-11-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019199 | MSE-4308296

Real number version of the complex conjugate theorem?

Recently, I have encountered a question where a polynomial and one of its irrational roots $a+b\sqrt{c}$ where $a,b,c\in\mathbb{Q}, \sqrt{c}\in\mathbb{R}$ were given, and I was asked to find all roots. I immediately assumed that there existed a root $a-b\sqrt{c}$, and this was a correct assumption. Looking back, however, I am not entirely sure if this is true. I know that the complex conjugate theorem states that for any polynomial $p(x)$ with real coefficients, if the polynomial has a complex root $a+bi$ then it must also have a root $a-bi$. My question is if there is a real number version of such a theorem, which might go along the lines of something like: For a polynomial $p(x)=a_1x^n+a_2x^{n-1}+a_3x^{n-2}+\dots+a_{n-1}x+a_n$ with real coefficients a_n, if it has a root $a+b\sqrt{c}$ then $a-b\sqrt{c}$ must also be a root of $p(x)$. Or maybe instead of real coefficients it could be integer coefficients?

论坛问题帖 | has_answers | Max0815 | 2021-11-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019200 | MSE-4310657

Generalizing the result $|a\times b|=|a||b||\sin \theta|$ to arbitrary dimensions.

The cross product can be generalized to arbitrary dimensions as done below or here. I'm trying to state and prove the general analogue (for arbitrary dimensions) of the equation $$|a\times b| = |a||b||\sin \theta |$$ where θ is the angle between a and b. Definition: let V be an inner product space of dimension n. The cross product of the vectors $v_1, \ldots ,v_{n-1}$ of V is defined as the unique vector v such that $$\det (u \ v_1 \ldots v_{n-1}) = u\cdot v$$ for every $u\in V$. It is denoted as $[v_1 \ldots v_{n-1}]$. Let $v_0, v_1, \ldots , v_{n-1}$ be vectors of an inner product space V (of dimension $n+1$). It is easy to show that the cross product $[v_0, v_1, \ldots ,v_{n-1}]$ is perpendicular to each v_i, and has a magnitude equal to the nth dimensional "area" of the parallelotope with edges $v_0, v_1, \ldots , v_{n-1}$. Let A be the $(n-1)$ dimensional area of the parallelotope with edges $v_1, \ldots , v_{n-1}$. That is, let $$A:=\sqrt{| \det G |}$$ where G is the Gram matrix of the vectors $v_1, \ldots, v_{n-1}$ (i.e. $G_{i,j}=v_i\cdot v_j$). Then the n-dimensional area of the parallelotope with edges $v_0, v_1, \ldots , v_{n-1}$ (namely $|[v_0, v_1, \ldots ,v_{n-1}]|$) should be $$A|v_0||\sin \theta |$$ where θ is the angle between v_0 and its projection v_0' onto the space $\langle v_1, \ldots , v_{n-1} \rangle$, namely $$v_0':=\sum_{i=1}^{n-1}\frac{v_0\cdot v_i}{|v_i|^2}v_i.$$ Is it true that $$|[v_0 \ v_1 \ldots v_{n-1}]| = A|v_0||\sin \theta| \ ?$$ If true, how could one prove it? If it isn't true, what am I getting wrong in my attempt to generalize the result stated in the introduction? My attempt at proving it: the expression $\sin \theta$ can be expanded by using the following definition of angle: $$\theta := \arccos \left(\frac{v_0\cdot v_0'}{|v_0||v_0'|}\right)$$ together with the fact that $\sin (\arccos (x))=\sqrt{1-x^2}$. Now A can be expanded in multiple ways (i.e. using Leibniz's definition of the determinant or the Laplace expansion). Finally, the LHS can be expanded by noting that $$[v_0 \ v_1 \ldots v_{n-1}] = \sum_{i=1}^{n+1}(-1)^{i+1}M_{1,i}e_i$$ where $\{ e_1, \ldots , e_{n+1}\}$ is a basis of V and $M_{1,i}$ is the determinant of the submatrix obtained by removing the ith row and the jth column of the matrix $$[1 \ v_0 \ v_1 \ldots v_{n-1}] .$$ I've been playing around with the resulting expression trying to force an equality, with no avail.

论坛问题帖 | has_answers | Sam | 2021-11-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019201 | MSE-4322129

All solutions to $1/a+1/b=1/c$?

Conjecture: Given integers $a>b>c>1$ such that $\gcd(a,b,c)=1$. Then all the positive integer solutions to $$\frac{1}{a}+\frac{1}{b}=\frac{1}{c}$$ is given by: c any number and for each divisor $d<\sqrt{c}\:$ of $\,c\,$ with $\gcd(d,c/d)=1$ $$ \left\{ \begin{array}{l} a_d=c+d^2 \\ b_d=\frac{c}{d}\cdot(\frac{c}{d}+d)=c\cdot(c+d^2)/d^2 \end{array} \right. $$ It is easy to show that a_d,b_d,c all are solutions, but how to prove they are the only solutions?

论坛问题帖 | has_answers | Lehs | 2021-12-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019202 | MSE-4327136

On the inequality $I(q^k)+I(n^2) \leq \frac{3q^{2k} + 2q^k + 1}{q^k (q^k + 1)}$ where $q^k n^2$ is an odd perfect number

(Note: This post is an offshoot of this earlier MSE question.) Let $N = q^k n^2$ be an odd perfect number with special prime q satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. Define the abundancy index $$I(x)=\frac{\sigma(x)}{x}$$ where $\sigma(x)$ is the classical sum of divisors of x. Since q is prime, we have the bounds $$1 < \frac{q+1}{q} \leq I(q^k) < \frac{q}{q-1} \leq \frac{5}{4},$$ which implies, since N is perfect, that $$\frac{8}{5} \leq \frac{2(q-1)}{q} < I(n^2) = \frac{2}{I(q^k)} \leq \frac{2q}{q+1} < 2.$$ By considering the nonnegative product $$\bigg(I(q^k) - \frac{q+1}{q}\bigg)\bigg(I(n^2) - \frac{q+1}{q}\bigg) \geq 0,$$ then after some routine algebraic manipulations, we arrive at the upper bound $$I(q^k) + I(n^2) \leq 3 - \Bigg(\frac{q-1}{q(q+1)}\Bigg) = \frac{3q^2 + 2q + 1}{q(q + 1)}.$$ Here, I shall derive the following slightly different bound: If $q^k n^2$ is an odd perfect number with special prime q, then $$I(q^k) + I(n^2) \leq \frac{3q^{2k} + 2q^k + 1}{q^k (q^k + 1)}.$$

PROOF: Suppose that $q^k n^2$ is an odd perfect number with special prime q. Then we have $$\sigma(q^k) \geq q^k + 1$$ so that $$\frac{1}{4} \geq \frac{1}{q-1} > I(q^k) - 1 \geq \frac{1}{q^k}$$ and $$1 > \frac{q-1}{q+1} \geq I(n^2) - 1 > \frac{q-2}{q} \geq \frac{3}{5}$$ so that we obtain $$\Bigg(\bigg(I(q^k) - 1\bigg) - \frac{1}{q^k}\Bigg)\Bigg(\bigg(I(n^2) - 1\bigg) - \frac{1}{q^k}\Bigg)$$ is nonnegative. Thus, we have $$\Bigg(\bigg(I(q^k) - 1\bigg) - \frac{1}{q^k}\Bigg)\Bigg(\bigg(I(n^2) - 1\bigg) - \frac{1}{q^k}\Bigg) \geq 0$$ which is equivalent to $$\bigg(I(q^k) - 1\bigg)\bigg(I(n^2) - 1\bigg) + \frac{1}{q^{2k}} \geq \frac{1}{q^k}\Bigg(I(q^k) + I(n^2) - 2\Bigg)$$ $$\Bigg(3 - \bigg(I(q^k) + I(n^2)\bigg)\Bigg) + \frac{1}{q^{2k}} \geq \frac{1}{q^k}\Bigg(I(q^k) + I(n^2) - 3\Bigg) + \frac{1}{q^k}$$ $$\Bigg(\frac{q^k + 1}{q^k}\Bigg)\Bigg(3 - \bigg(I(q^k) + I(n^2)\bigg)\Bigg) = \Bigg(1 + \frac{1}{q^k}\Bigg)\Bigg(3 - \bigg(I(q^k) + I(n^2)\bigg)\Bigg) \geq \frac{1}{q^k} - \frac{1}{q^{2k}} = \frac{q^k - 1}{q^{2k}}$$ $$3 - \bigg(I(q^k) + I(n^2)\bigg) \geq \frac{q^k - 1}{q^k (q^k + 1)}.$$ Hence, we finally get $$I(q^k) + I(n^2) \leq 3 - \Bigg(\frac{q^k - 1}{q^k (q^k + 1)}\Bigg) = \frac{3q^{2k} + 2q^k + 1}{q^k (q^k + 1)}.$$ QED. Notice the striking similarity between the upper bound $$I(q^k) + I(n^2) \leq 3 - \Bigg(\frac{q-1}{q(q+1)}\Bigg) = \frac{3q^2 + 2q + 1}{q(q + 1)},$$ and the upper bound $$I(q^k) + I(n^2) \leq 3 - \Bigg(\frac{q^k - 1}{q^k (q^k + 1)}\Bigg) = \frac{3q^{2k} + 2q^k + 1}{q^k (q^k + 1)}.$$ Here are my: QUESTIONS: (1) Does this argument imply that $k=1$? (i.e. in the sense that k is taken to be a placeholder for 1?) (2) Similar to the train of thought in this post, can you think of a way to show that $$I(q^k) + I(n^2) > 3 - \Bigg(\frac{q^k - 2}{q^k (q^k - 1)}\Bigg) = \frac{3q^{2k} - 4q^k + 2}{q^k (q^k - 1)}?$$ (3) If it is not possible to prove the inequality in (2), can you explain why?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2021-12-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019203 | MSE-4330941

Primes being a square sum of an odd number and a less number

For natural numbers a,b, let a be odd and let $f(a)$ be the number of $\,b<a\,$ such that a^2+b^2 is prime. In the diagram f is blue, the function $g(n)=\frac{n}{\ln n}$ is red and $h(n)=\frac{1}{\pi}\cdot\frac{n}{\ln n}$ is green. Since the blue curve is nearly limited by the red and the green curves it seems reasonable to make the conjecture: There are real numbers A and B such that $A\cdot\frac{n} {\ln n}<f(n)<B\cdot\frac{n}{\ln n}$, where n is odd. What would be the intuition behind this?

论坛问题帖 | no_answers_at_retrieval | Lehs | 2021-12-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019204 | MSE-4358856

If $\gcd(n^2, \sigma(n^2)) = q\sigma(n^2) - 2(q - 1) n^2$, does it follow that $q n^2$ is perfect?

Let q be an (odd) prime, and let $\gcd(q,n)=1$. Denote the classical sum of divisors of the positive integer x by $\sigma(x)=\sigma_1(x)$. A number N is said to be perfect if $\sigma(N)=2N$. Here is my question: If $\gcd(n^2, \sigma(n^2)) = q\sigma(n^2) - 2(q - 1) n^2$, does it follow that $q n^2$ is perfect? CONTEXT It is known that if $p^k m^2$ is perfect, where p is an odd prime satisfying $\gcd(p,m)=1$, then we obtain $$i(p):=\dfrac{\sigma(m^2)}{p^k}=\dfrac{2m^2}{\sigma(p^k)}=\dfrac{D(m^2)}{s(p^k)}, \tag{*}$$ where $D(x)=2x-\sigma(x)$ is the deficiency of x and $s(x)=\sigma(x)-x$ is the aliquot sum of x. (Note that $i(p)$ is odd.) We then obtain $$\sigma(m^2) = p^k {i(p)}$$ and $$m^2 = \dfrac{\sigma(p^k)}{2}\cdot{i(p)},$$ so that $$\gcd(m^2, \sigma(m^2)) = i(p)\cdot\gcd\Bigg(p^k, \dfrac{\sigma(p^k)}{2}\Bigg) = i(p).$$ Additionally, we can also rewrite Equation (*) as $$i(p) = \dfrac{\sigma(m^2)}{p^k} = \dfrac{D(m^2)}{s(p^k)} = \dfrac{(p-1)D(m^2)}{p^k - 1} = \dfrac{\sigma(m^2) - (p-1)D(m^2)}{p^k - (p^k - 1)}$$ $$= p\sigma(m^2) - 2(p-1)m^2.$$ Specializing to $k=1$, we obtain the implication $$qn^2 \text{ is perfect with } \gcd(q,n)=1 \implies \gcd(n^2, \sigma(n^2))=q\sigma(n^2) - 2(q-1)n^2.$$ SANITY CHECK Let qn^2 be an even perfect number. Then $q = 2^t - 1$ and $n^2 = 2^{t-1}$, for some prime number t. We compute $$\gcd(n^2, \sigma(n^2)) = \gcd(2^{t-1}, 2^t - 1) = 1$$ and $$q\sigma(n^2) - 2(q - 1)n^2 = (2^t - 1)(2^t - 1) - 2(2^t - 2){2^{t-1}} = 2^{2t} - 2^{t+1} + 1 - 2^{2t} + 2^{t+1} = 1,$$ whence we have equality between $\gcd(n^2, \sigma(n^2))$ and $q\sigma(n^2) - 2(q - 1)n^2$. (Note that these computations also "work" for the even perfect number 6, even though it is squarefree.) PROBLEM This takes care of one direction. Can you come up with a proof for the other direction? Alas, this is where I get stuck.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2022-01-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019205 | MSE-4377085

Is the following infinite product of fractions of linear factors equal to an exponential function or not?

Is the following infinite product: $$ \prod_{\substack{(a,b) \in \mathbb{Z}^2 \\ a > b}} \frac{x+b}{x+a} $$ defined? If so, does it simplify to an exponential function of x? The subscript is over all integer pairs a and b such that $a>b $ Motivation/suggestions: When plotting a graph of a finite truncation of the product : it resembles a negative exponential function. However such an approximation appears to not be accurate for small values of x (where it has vertical asymptotes). Can it be transformed in such a way that it becomes an exponential function, by modifying each term individually so that the whole function is finite and the asymptotes lie at infinite distance in the limit? Or can we show that the derivative of this infinite product with respect to x is a constant factor of itself or not, thus proving that it is an exponential? I believe it should be exponential because it was obtained by raising e to the power a modified version of an integral of a constant function.

论坛问题帖 | has_answers | AmoebaProteus | 2022-02-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019206 | MSE-4379015

How much Base Proof is required?

If I were looking to use the Theory of Strong Induction, to prove the statement that: The Set of All Even Numbers is equal to the Set of All Odd Numbers multiplied by 2^n, where $n \geq 1$ $$ \mathbb{E} = \mathbb{O} \cdot \{2^n \mid n \geq 1 \} $$ How much Base Proof would be required to show or infer that it is true? $2 = 1 \cdot 2^1$ $4 = 1 \cdot 2^2$ $6 =

3 \cdot 2^1$ $8 = 1 \cdot 2^4$ $10 = 5 \cdot 2^1 \dots$ I get that for Strong Induction you have to show all the base proofs up to a certain point. But in this case, what would that minimum requirement be? Is there a minimum amount required, or is this what makes a Conjecture? Because the base proof requires an infinite number of base proofs, to prove the statement and relies more on a formula for its proof over Strong Induction?

论坛问题帖 | has_answers | Houshou Rattengod | 2022-02-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019207 | MSE-4388714

What is the distribution of ax^2+bx+c as $\{a,b,c,x\}$ vary over $\mathbb Z$? (re: quadratic primes)

I realize this may be one of those questions where it's ill-posed and can't be answered without further constraints. But if not, what's the distribution like? In particular, is every integer equally often represented? On the one hand, that appears likely based on the observation that regardless of what effect $\{a,b,x\}$ might have, c can serve to exactly map one $\{a,b,x\}$ tuple to every integer. On the other hand, this feels suspiciously like one of those things that turns out to depend on how exactly you enumerate a,b,c,x as you walk towards infinity. I am also suspicious because of how I arrived at this question, via basic thoughts on the infinitude of quadratic primes. I reasoned as follows: Suppose there are only finitely many quadratic primes, and let $f(x)=ax^2+bx+c$ be some function that returns primes for finitely many values of x. Let m,n be the largest and smallest x such that $f(x)$ is prime, and let k be any integer outside of the interval $[n,m]$. Then $f'(x)=f(x+k)=ax^2+bx+c+ak^2+bk+2axk$ will be a quadratic function which is irreducible but always composite. Moreover, since there are infinitely many k values yielding never-prime quadratics compared to finitely many k that are sometimes prime, and the same would apply to any quadratic function used, it seems to follow that almost all quadratic functions must yield no primes. And that brings us full circle, since if ax^2+bx+c could be said to be "evenly distributed" in some meaningful way, and if my logic above holds, we'd have a contradiction proving infinitude of quadratic primes. Since we don't, quadratics must not be evenly distributed, and/or my logic above is flawed. If someone could point out where I've gone astray, I'll consider my question answered. Edit: Okay, let's do the naive thing and increase some $i\to\infty$ and consider a,b,c,x in range $[-i,i]$.

论坛问题帖 | has_answers | Trevor | 2022-02-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019208 | MSE-4388965

For any composition sequence s of maps $h(X)=X/2, \ f(X)=(3X + 1)/2$, there exists an integer X such that its Collatz sequence contains s

Let $h(X) = X/2$, and $f(X) = (3X + 1)/2$. Then clearly every iteration $g^i(X), X \in \Bbb{Z}$ the Collatz mapping $$g(X) = \begin{cases} X/2, \ X=0\pmod 2\\ \dfrac{3X + 1}{2}, \ X = 1\pmod 2 \end{cases}$$ can be modelled by compositions of f, h. For example, $g^7(3) = 1 = hhhff(3)$ where $g(X) = f(X)$ for odd numbers and $g(X) = h(X)$ for even numbers. Under iterations of g, each number $X \in \Bbb{Z}$ induces a unique composition of f and h functions. This should be clear because a number is either even or odd but not both, so at each stage in an iteration there is one and only one function (f or h) to apply. Conjecture. For any sequence $s \in \{f,h\}^*$ (Kleene closure under $\circ$) there exists an integer X such that $g^i(X) = s(X)$ for some $i \geq 0$. Can we prove this or is it just as difficult as Collatz? Clearly we have that it's true for the sublanguage $\{h\}^*$ since we can take $g^n(2^n) = h^n(X)$ for all $n \geq 0$. Clearly we have that it's true for the sublanguage $\{h\}^*$ since we can take $g^n(2^n) = h^n(X)$ for all $n \geq 0$. It's also true for $f\{h\}^*$ since $X = 2^n O$ works, where O is an odd number. And it's true for $\{h\}^* f$ since $3O + 1 = 2^ny$ has a solution if and only if $30 + 1 = 0 \pmod 2^n \iff O = -3^{-1} \pmod 2^n$. So in fact we've proven an infinitude of solutions namely: $-3^{-1} + 2^n\Bbb{Z}$ for any $h^n f \in \{h\}^* f$. Let these two proofs comprise a base case for induction on the number of f's appearing in s.

论坛问题帖 | has_answers | TotoposDiagramChaseApp | 2022-02-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019209 | MSE-440859

Why does one counterexample disprove a conjecture?

Can't a conjecture be correct about most solutions except maybe a family of solutions? For example, a few centuries ago it was widely believed that $2^{2^n}+1$ is a prime number for any n . For $n=0$ we get 3 , for $n=1$ we get 5 , for $n=2$ we get 17 , for $n=3$ we get 257 , but for $n=4$ it was too difficult to find if this was a prime, until Euler was able to find a factor of it. It seems like this conjecture stopped after that. But what if this conjecture isn't true only when n satisfies a certain equation, or when n is a power of 2 $\ge$ 4 , or something like that? Did anybody bother to check? I am not asking about this conjecture specifically, but as to why we consider one counterexample as proof that a conjecture is totally wrong. P.S. Andre Nicolas pointed out that Euler found a factor when $n=5$, not 4 .

论坛问题帖 | has_answers | Ovi | 2013-07-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019210 | MSE-4417988

What can we say about the colourings of some subset of vertices of a graph

I will clarify the question by defining some straightforward concepts. Let G = (V, E) be a simple graph with chromatic number k. We call a subset $V'\subset V$ fully chromatic if every proper k-colouring of G uses also k distinct colors on V'. We say such a fully chromatic set is minimal, if every subset of V' is not fully chromatic. Now I conjecture that for every G=(V, E) and every minimally chromatic set $V'\subset V$, there exists some graph $G^* = (V^*, E^*) $ that has the same proper k-colourings on V^* as all proper k-colourings of G restricted to V'. Furtermore, I conjecture that G has G^* as a minor

论坛问题帖 | no_answers_at_retrieval | AndroidBeginner | 2022-04-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019211 | MSE-4419646

Proof for strict inequality $\pi(ab) > \pi(a)\pi(b)$?

I asked about the very similar $\pi(ab)\geq \pi(a)\pi(b)$ a while ago, and this is indeed a proven result for $a,b\geq \sqrt{53}$. Empirically, the stricter $\pi(ab)>\pi(a)\pi(b)$ looks true so long as either a or b is ≥ 23. It also seems to pretty immediately fall out of the PNT, a la $$\pi(ab)\sim\frac{ab}{\log(ab)}>\frac{ab}{\log(a)\log(b)}\sim \pi(a)\pi(b)$$ given sufficiently large a,b. If someone can point me to a paper, give pointers/work out on how this proof could be done, or if applicable, explain why this is probably not provable, I'll consider this answered. If you can verify my proof below or give some solid feedback, that would work too. Okay, I have a first attempt. For $x\geq 17$, Wikipedia gives the bounds $$\frac x {\log x} < \pi(x) \tag{1}$$ $$\pi(x) < 1.25506 \frac x {\log x}. \tag{2}$$ From applying (2) twice, we have $$\pi(a)\pi(b) < 1.25506^2 \cdot \frac{ab}{\log(a)\log(b)}.\tag{3}$$ From (1), we get $$\frac{ab}{\log(ab)} < \pi(ab).\tag{4}$$ It only remains to connect them: $$1.25506^2 \cdot \frac{ab}{\log(a)\log(b)} < \frac{ab}{\log(ab)}.\tag{5}$$ From trial and error, letting $a,b \geq 24$ seems sufficient to satisfy (5). We put the pieces together and get the final inequality chain $$\pi(a)\pi(b) < 1.25506^2 \cdot \frac{ab}{\log(a)\log(b)} <\frac{ab}{\log(ab)} < \pi(ab). \tag{6}$$ This gives $\pi(a)\pi(b)<\pi(ab)$ for $a,b\geq 24$. Have I gone awry somewhere or does this work?

论坛问题帖 | has_answers | Trevor | 2022-04-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019212 | MSE-4420871

Asymmetric graphs where all adjacent vertices have different degree

I conjecture the following : If G is a graph so that every pair of adjacent vertices have different degree, then G is not asymmetric graph. I will remind that a graph is a asymmetric if there are no automorphisms, except for the trivial automorphisms. It is possible that this statement is incorrect. Can someone find a counterexample? And a similar conjecture: If G is a 2-connected chordal graph, then it contains a triangle where at least two vertices are of the same degree.

论坛问题帖 | has_answers | AndroidBeginner | 2022-04-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019213 | MSE-4429396

I conjecture that the circumference of a graph is larger than its average degree

Let $G = (V,E)$ be a simple loop-less graph. The circumference refers to the length of the largest cycle of G. The average degree is $avg(G) = |E|*2 / |V|$. I conjecture that if the avg(G) > 2, then $circum(G) > avg(G)$. I can prove that the circumference is larger than the minimum degree of G. This can be quite easily seen by looking at the maximum path of G. The neighbors of the endpoints must then all lie within the path. Connecting one endpoint with it's neighbor that lies the furthest on the path would then result in a cycle of at least the degree of the endpoint + 1. Any idea how to prove my stronger claim?, or a counter-example?

论坛问题帖 | has_answers | AndroidBeginner | 2022-04-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019214 | MSE-4430444

Is there anything known about the tree width of the direct sum of two graphs?

So, if you have two graphs (both with a designated edge), you could take the direct sum of those two graphs, where you glue those two graphs to each other along the designated edge. I think different literature use different definitions. Anyways, see the picture below as an example of gluing two cycles to each other of length 6 along some edge. I was wondering if there is anything known about what happens to the tree width of the new graph, given the tree widths of the original two graphs? Is there some literature about it. I thought perhaps that $tw(G \oplus H) = \max({tw(G), tw(H)})$, where $\oplus$ denotes the operation I described above, and tw being the tree width. I'm not even sure if this is true, and if it'd be true, where I could read more about this, or whether there are known proofs I could refer to. If not, maybe something is known about tree widths of graphs under different kind of operations. If something is unclear about my question, let me know. Thank you!

论坛问题帖 | no_answers_at_retrieval | AkatsukiMaliki | 2022-04-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019215 | MSE-4433059

I have a conjecture regarding vertex coloring

Let $G = (V, E)$ be a graph with chromatic number $n+1$, and let there be some vertex $v* \in V$, so that deleting this vertex results in a graph $G/{v*}$ that has chromatic number n. Now, assign to $G/{v*}$ any proper n-coloring, then color $v*$ in the $(n+1)-th$ color. Rainbow Conjecture: Choose at least 2 (and at most n) distinct colors from the first n colors, and order these in any way. I conjecture that there exists a cycle in G that contains $v*$, and so that the colors we come across when we traverse this cycle starting from $v*$, are precisely (one or multiple) repetitions of the chosen pattern. So cycles are for example colored like: $v*$ - blue - green - red - ($v*$) , or $v*$- blue- red - blue - red - ($v*$). And these cycles exist for all proper n-colorings of $G/{v*}$ and any combination and order of the n colors. The reason why I think this might be true is because if there would be an ordering that doesn't have a matching path that starts at $v*$, for example: if we can't go from $v*$ to a red vertex and then a green vertex, then all red neighbors of $v*$ don't have green neighbors, so we could change them all to green, but then we could change $v*$ to red, which would result in a n-coloring of G. But this doesn't really explain why the path must also come back to $v*$. My conjecture seems quite optimistic, however, I haven't been able to find a counter example. What do you guys think? Thanks for your time, and I'd love to see if someone can find a counterexample (or a proof even better). I guess counterexamples would have generally high girth or low clique.

论坛问题帖 | has_answers | AndroidBeginner | 2022-04-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019216 | MSE-4433274

Why does e/π (or $1/2$) arise in this limit?

(Related to this prior post.) Let $f(n)$ for $n\in\mathbb N$ be a function that increases the prime index of each prime factor of n (with multiplicity) by 1. e.g. $f(20)=f(2^2\cdot 5)=f(p_1^2\cdot p_3)=p_{2}^2\cdot p_{4}=3^2\cdot 7=63$. Consider $\frac{n}{f(n)}$ as $n\to\infty$. The mean of these terms has a clear limit: $$\lim_{n\to\infty} \left(\frac{1}{n}\sum_{i=1}^n \frac{i}{f(i)}\right)=\frac{1}{2}.$$ But more interesting is that if you let a be a count of how many terms through n have $\frac{i}{f(i)}>\frac{1}{2}$, and therefore $n-a$ be the count of terms with $\frac{i}{f(i)}<\frac{1}{2}$, then it appears that $$\lim_{n\to\infty} \frac{a}{n-a}=\frac{e}{\pi}.$$ This is a strictly empirical result, but it seems to hold within a margin of error of around $\frac{1}{\sqrt{a}}$. So far, I've checked this for $n\leq 10^7$, and while the results aren't strong enough to rule out coincidence yet, it seems too suggestive to ignore. Does anyone know why this might arise? And if it's correct, might it imply that $\frac{e}{\pi}$ is irrational? It seems to me that $\frac{a}{n-a}$ can't be rational in the limit since it will never cease to fluctuate in finer pseudorandom detail based on the distribution of the primes, but perhaps the limit negates that somehow. If anyone can explain why the limit $\frac{e}{\pi}$ would appear here (or can show that I'm mistaken), I'll consider this answered. Update The more I look, the less convinced I am that I had this right; I still believe there's a limit, but not e/π, as it seems to run about 0.002 higher pretty consistently. As for the $1/2$ limit, I'm quite sure that's solid. Empirical data leaves essentially no doubt, and it's very much in line with the behavior of f, which has properties like for any n, you'll get $f(k)> n$ for exactly half of the naturals $k\in [1,n]$. (See this post for more.) That said, I don't know how to prove the $1/2$ limit, so I'd accept that as an answer here as well if someone can demonstrate it. Basic notes on $\frac{i}{f(i)} > \frac{1}{2}$ properties Let $S_n = \{1 \leq s \leq n : f(s) > 2s\}$. (Note $f(s) > 2s$ is equivalent to $\frac{s}{f(s)} < \frac{1}{2}$, but cleaner looking.) Regardless of choice of n, $|S_n|$ will be slightly larger than $n/2$. Example $S_{128}=\{4, 6, 8, 9, 10, 12, 14, 15, 16, 18, 20, 21, 24, 27, 28, 30,\\ 32, 35, 36, 39, 40, 42, 44, 45, 48, 49, 50, 52, 54, 56, 57, 60, 63, 64, 66,\\ 68, 69, 70, 72, 75, 76, 78, 80, 81, 84, 88, 90, 91, 92, 96, 98, 99, 100,\\ 102, 104, 105, 108, 110, 112, 114, 116, 117, 120, 124, 125, 126, 128\}.$ $|S_{128}|=67$. This set can be substantially compressed by eliminating multiples, leaving $\{4, 6, 9, 10, 14, 15, 21, 35, 39, 49, 57, 69, 91, 125\}$. Any integer divisible by any of these values will also have the property $f(i) > 2i$. It is clear with a little inspection that there will be infinitely many of these values as n is made arbitrarily large. Various patterns abound, e.g. $5^2 \cdot p_i$ will be one of these minimal elements for all $i \geq 3$. The overall distribution appears complex, similar to the primes themselves in many ways. All integers in these S sets are necessarily composite, as having a prime would violate Bertrand's postulate. On the whole, elements in these sets are "especially" composite, with larger factor counts and smaller factors. Contrast this to the complementary set R_n comprising those $r\leq n$ such that $f(r) < 2r$: Example $R_{128}=\{1, 2, 3, 5, 7, 11, 13, 17, 19, 22, 23, 25, 26, 29, 31, 33,\\ 34, 37, 38, 41, 43, 46, 47, 51, 53, 55, 58, 59, 61, 62, 65, 67, 71, 73,\\ 74, 77, 79, 82, 83, 85, 86, 87, 89, 93, 94, 95, 97, 101, 103, 106,\\ 107, 109, 111, 113, 115, 118, 119, 121, 122, 123, 127\}.$ $|R_{128}|=61.$ R_n necessarily contains all the primes $\leq n$ as well as most $2p$, p^2, and other semiprime forms; in general, it is populated by high-prime-index low-factor-count integers, as expected from the complement to S_n.

论坛问题帖 | no_answers_at_retrieval | Trevor | 2022-04-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019217 | MSE-4445236

fundamental solution of Pell’s equation under special conditions

I make the following conjecture from an answer. conjecture Suppose N could be expressed as $$ N= p^2\pm q,\ \ \ (p,q \in \mathbb{N},\ 1 \lt q \lt p,\ q\mid 2p) $$ In this case, the fundamental solution of $x^2-Ny^2=1$ is obtained by the following expression. (double sign in same order) $$ x_0=\frac{2p^2}{q} \pm 1,\ \ y_0=\frac{2p}{q} $$ (x_0,y_0) is one of the solutions of $x^2-Ny^2=1$ because it can be calculated as follows. \begin{eqnarray*} x_0^2-Ny_0^2&=& \left(\frac{2p^2}{q} \pm 1\right)^2-(p^2\pm q)\cdot \left(\frac{2p}{q}\right)^2\\ &=& \frac{4p^4}{q^2} \pm \frac{4p^2}{q} +1 -\frac{4p^4}{q^2} \mp\frac{4p^2}{q}\\ &=& 1 \end{eqnarray*} But, I don't know how to prove that it is the fundamental solution. concrete example Let N be 75. The expression using the square number of the nearest neighbors is $$ 75 = 9^2-6 $$ $p=9,\ q=6$ satisfy $1 \lt q \lt p$ and $\ q\mid 2p$. Therefore, \begin{eqnarray*} x_0&=&\frac{2p^2}{q} - 1,\ \ \ y_0&=&\frac{2p}{q} \\ &=& \frac{2\cdot 9^2}{6} -1 &=& \frac{2\cdot 9}{6} \\ &=& 26 &=& 3 \end{eqnarray*} $(x_0, y_0) = (26, 3)$ is the fundamental solution of $x^2-75y^2=1$.

论坛问题帖 | no_answers_at_retrieval | isato | 2022-05-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019218 | MSE-4453332

Conjecture about areas of circular segment and polygon with equal perimeter sharing a side

I was playing around with shapes and have formed a conjecture. Length of the red circular arc $=$ total length of the n green line segments Conjecture: $$\sup{\left(\frac{\text{Area}_1}{\text{Area}_2}\right)}=\sqrt{1-\frac{1}{n^2}}$$ Is my conjecture true, and if so, is it a known thing? I have proven the case with $n=2$ algebraically, and (I think) I have verified the cases with $n=3$ and $n=4$, using desmos. The case with $n=2$: The ratio of the areas is maximized when the two green segments are the same length (this can be seen by considering the ellipse whose focal points are the ends of the black line, passing through the point where the two green segments meet). I let the central angle of the arc be 2θ, and got $$\frac{\text{Area}_1}{\text{Area}_2}=\frac{2(\sin{\theta})\sqrt{\theta^2-\sin^2{\theta}}}{2\theta-\sin{2\theta}}$$ Basic calculus shows that this function is decreasing in $0<\theta<\pi$, and the limit as $\theta\to0^+$ is indeed $\frac{\sqrt{3}}{2}$. Applying a similar method to larger n seems quite daunting. (My background: high school math teacher.)

论坛问题帖 | has_answers | Dan | 2022-05-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019219 | MSE-4456826

A function asymptotical equivalent with the prime counting function?

Let p_n be the n-th prime number and $Q_a(N)$ be the number of primes of the form p_n^2+a where $1\leq n\leq N$ and a is positive and even. For some a like $26,56$ it seems that no solutions to $p_n^2+a\in\mathbb P$ exist, for some a like $2,8$ there seems to be a unique solution and for some a like $4,12$ there seems to be an unlimited number of solutions. The diagrams below is of $Q_4(N)$ (blue) and the function $\frac{N}{\ln N}$ (red). My question, is it possible to prove that $Q_4(N)\sim\frac{N}{\ln N}$?

论坛问题帖 | has_answers | Lehs | 2022-05-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019220 | MSE-4461630

Conjecture that $\sum\limits_{n=1}^{\infty}\big(\frac 1{q^{p_n}-1}-\sum\limits_{k=1}^{n-1}\frac 1{q^{p_n p_{n-k}}-1}\big)=\frac 1{q(q-1)}$

I have had this conjecture for a while: Let $q\in\mathbb R\setminus [-1,1]$ and let p_n be the n^th prime. Then: $$\sum_{n=1}^{\infty}\bigg(\frac 1{q^{p_n}-1}-\sum_{k=1}^{n-1}\frac 1{q^{p_n p_{n-k}}-1}\bigg)=\frac 1{q(q-1)}$$ The sum looks like this when we expand it out: $$\frac 1{q^2-1}+\bigg(\frac 1{q^3-1}-\frac 1{q^{3\times 2}-1}\bigg)+\bigg(\frac 1{q^5-1}-\frac 1{q^{5\times 3}-1}-\frac 1{q^{5\times 2}-1}\bigg)+\cdots=\frac 1{q(q-1)}$$ I don't know how to show something like this. I can pull out a factor of $1/(q-1)$ but I don't think that will help. Is this even true? I just had it written down in my notebook. (Sorry for not having any good proof attempts of my own.)

论坛问题帖 | has_answers | Mr Pie | 2022-05-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019221 | MSE-4473633

Extending a simple 1-dimensional observation to 2-dimensions.

Consider the following (easy to prove) statemnt. Statement 1. There is an absolute constant c_1 such that whenever $n\geq 1$ and $x_1, \ldots, x_n$ are real numbers such that $|x_i| \leq 1$ for all i and $\sum_{i=1}^n x_i = 0$, one can find a permutation σ of $\{1, \ldots, n\}$ such that each of the numbers in the sequence $$ x_{\sigma(1)},\ x_{\sigma(1)} + x_{\sigma(2)},\ x_{\sigma(1)} + x_{\sigma(2)} + x_{\sigma(3)}, \ldots, x_{\sigma(1)} + \cdots + x_{\sigma(n)} $$ has magnitude at most c_1. In fact, one can easily show that $c_1 = 1$ works. Pick x_1 and set $\sigma(1) = 1$. If $x_1>0$, find a negative x_i and set $\sigma(2) = i$ and continue greedily. The following is the two dimensional version of the above statement, but it is quite unclear to me if it is true of not. Statement 2. There is an absolute constant c_2 such that whenever $n\geq 1$ and $x_1, \ldots, x_n$ and $y_1, \ldots, y_n$ are two sequences of real numbers such that $|x_i|, |y_i|\leq 1$ for all i, and $\sum_{i=1}^n x_i = \sum_{i=1}^n y_i = 0$, one can find a permutation σ of $\{1, \ldots, n\}$ such that for each $1\leq k\leq n$, the magnitude of both of $$ x_{\sigma(1)} + x_{\sigma(2)} + x_{\sigma(3)} + \cdots + x_{\sigma(k)}, \quad y_{\sigma(1)} + y_{\sigma(2)} + y_{\sigma(3)} + \cdots + y_{\sigma(k)} $$ is at most c_2.

论坛问题帖 | has_answers | caffeinemachine | 2022-06-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019222 | MSE-4488136

Strong induction on the maximum size of a set

I'm currently working on an advancement of the Union-closed sets conjcture, and I have a whole cascading series of implications that follow from one claim, and try as I may, I have not been able to prove this claim (I will provide intuition for why it could be true later on) Given a union closed family, F, on an m-universe, that is, F a collection of subsets of the set: $$ U:= \{1, 2,...,m\} $$ with the property that the union of any two sets in F is also in F Then define the k^{th} layer of of F as $L_k(F) = \{X \in F: |X| = k \}$ The claim that I am trying to prove is that if every element of U appears in exactly $$ 2^{m-1} - \sum_{i=1}^n \binom{m-1}{i-1}$$ Sets in F, for some $n<m$ then the maximum size of F is $$2^{m} - \sum_{i=1}^n \binom{m}{i}$$ Those quantities arise precisely from removing the first n layers of the powerset of U, $\mathcal{P}(U)$ and so what the claim is saying is that this is the least amount of sets you can hope to remove so that each element appears exactly that amount of times. For the powerset itself, i.e when $n=0$ this is indeed true, and every element appears 2^m times, and for $n=1$ it seems that not only is it true but this is in fact the only removal which results in every element appearing exactly $2^m -1$ times I tried attacking this with a strong induction, assuming that for every n up to k, this was the maximum size of F, and then showing that it must also be the maximum size for $n=k+1$ the intuition being that if you were to remove some sets from some of the "higher" layers of $\mathcal{P}(U)$, you would necessarily need to remove some sets from all the layers "below" it, to preserve the union-closed property, and indeed, if you remove only one set X_k from $L_k(\mathcal{P}(U))$ then you have to remove at least $k-1$ sets from $L_{k-1}(\mathcal{P}(U))$ to preserve the union-closed property, and the single removal of X_k can be substituted for an equivalent removal of some X_{k-1} from the layer below it, along with the previously removed $k-1 $ sets from this layer (equivalent in terms of the minimum decrease in appearance of all the elements in X_k) and so removing only from the layer $L_{k-1}(\mathcal{P}(U))$ is at least as effective as removing from both layers. This leads one to think that all the ingredients for an inductive step are present, but I haven't been able to complete it, since it gets considerably more complicated to "keep track of things" once you start removing more than simply one set from an otherwise full layer. Any help would be appreciated, if this were to be proben it would be a significant step forward for the conjecture (sorry,thought it too space consuming to include the description of the original conjecture too), accordingly, any complete answers will get at least an acknowledgment.

论坛问题帖 | no_answers_at_retrieval | Carlyle | 2022-07-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019223 | MSE-4490088

Three hyperbolas related to a triangle concur at two points

I found this interesting fact (which I believe is non-trivial): Let $\triangle ABC$ be a triangle. Let $\mathcal{H}_A$ be the hyperbola with foci at B and C, passing through A. Define $\mathcal{H}_B$ and $\mathcal{H}_C$ similarly. Then $\mathcal{H}_A,\mathcal{H}_B,\mathcal{H}_C$ meet at exactly two points unless $\triangle ABC$ is equilateral, in which they would meet at one point (which is the orthocenter=incenter=circumcenter in that case). I'm not very well-versed with conics, so I'm not sure how I would go about proving this. I tried using the basic definition of a hyperbola, and you get a system of three distance equations, which are quite tricky to solve. I then supposed that X is some point such that $|XA-XB|=|a-b|, |XB-XC|=|b-c|$ (the intersection of two of the hyperbolas), and then tossing this on the complex plane, setting X to be the origin. This didn't go so well. I don't believe any bashing approaches work, I think this problem requires some sort of projective transformation. Also, it's quite a simple conjecture, so it's likely well known, but I haven't been able to find it on the internet. All help is appreciated.

论坛问题帖 | has_answers | TheBestMagician | 2022-07-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019224 | MSE-4497061

Can every even number be written as $(p-q)/6$ where p,q are primes?

For n even, assume $\large n = \frac{q-p}{6}$ For $a,b \in\mathbb{N}$, primes p,q can be written as $q = 6a\pm1$ $p = 6b\pm1$ There are 4 possible cases Case 1 $n = \frac{(6a-1)-(6b-1)}{6}$, then $n=a-b$ Case 2 $n = \frac{(6a-1)-(6b+1)}{6}$, then $n = a-b-1/3$ Case 3 $n = \frac{(6a+1)-(6b-1)}{6}$, then $n = a-b+1/3$ Case 4 $n = \frac{(6a+1)-(6b+1)}{6}$, then $n=a-b$ For Cases 2 and 3, n cannot be even because of $\pm 1/3$. For Cases 1 and 4, both result in $n=a-b$. Not sure how to proceed from here?

论坛问题帖 | no_answers_at_retrieval | vengy | 2022-07-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019225 | MSE-4498168

On the eighth powers $A^8+B^8=C^8+D^8$. Are there non-trivial solutions to it?

It's conjectured that there are no non-trivial solutions to the Diophantine equation: $$ A^8+B^8=C^8+D^8 $$ I was trying to play around with it using some substitutions.. In particular I first write it as: $A^8-D^8=C^8-B^8$. Then I replace (A,D,C,B) with $(p+q,p-q,r+s,r-s)$ to obtein the following: $$ p q (p^2 + q^2) (p^4 + 6 p^2 q^2 + q^4) = r s (r^2 + s^2) (r^4 + 6 r^2 s^2 + s^4) $$ Then I replaced (p,q,r,s) with $(ax+by,ax-by,cx+dy,cx-dy)$ to obtein: $$ a^8 x^8 - b^8 y^8 = c^8 x^8 - d^8 y^8 $$ or: $$ \frac{x^8}{y^8} = \frac{b^8-d^8}{a^8-c^8} $$ I was trying to substitute some values but I only manage to get trivial solutions.. I would like to find some values such that $(A,B)\not=(C,D)$. Any kind of suggestions would be much appreciated. Thanks in advance for your help! Some references I have already looked at: https://arxiv.org/pdf/math/0505629v2.pdf, https://www.jstor.org/stable/pdf/2007700.pdf?refreqid=excelsior%3A139ca229e2c0550603c138b5ada591c3&ab_segments=&origin=&acceptTC=1, https://sites.google.com/site/tpiezas/013 . EDIT 1: My last substituion is useless. I have just written (A,D,C,B) as (ax,by,cx,dy) in the end.. And there are no non-trivial solution in this case.

论坛问题帖 | has_answers | MathRevenge | 2022-07-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019226 | MSE-4498966

For every odd prime number $p \gt 3$ there exists another prime number $q \lt p$ such that $p - q = 2^n$ for some $n \geq 1$. Can you prove it?

Conjecture. Let p be an odd prime number greater than 3. Then there exists another odd prime number $q \lt p$ such that $p - q = 2^n$ for some positive exponent n. Can we prove this or is it another one of those simple to state, but almost impossible to prove, statements about prime numbers?

论坛问题帖 | has_answers | TotoposDiagramChaseApp | 2022-07-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019227 | MSE-4501927

Find a Stirling-formula like for $(x!)!$

Hi working again on the gamma function I find that : let $0<x\leq 1$ and $1\leq k\leq \infty$ then define : $$f(x)=(x!)!,g(x)=\left(\left(x!\right)!\right)^{\frac{1}{e^{x^{k}}}}$$ Then a conjecture : It seems $\exists k\in(1,\infty)$ such that $g(x)$ admits an asymptote as $x\to \infty$. Then see here Trying to prove the Stirling approximation using concavity for $x\geq 1$ we can squeeze the function $f(x)$ and remains to determine some constant . If my conjecture is true how to achieve this ?

论坛问题帖 | has_answers | Barackouda | 2022-07-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019228 | MSE-4507749

Are there conjectured closed forms for factorial rational zeta series?

Context At the bottom of this page from Wolfram Mathworld on the Riemann zeta function, the following constants are defined in equations (133), (137), and (141), respectively: \begin{align*} C_{1} &:= \sum_{n=2}^{\infty} \frac{\zeta(n)}{n!} \\ & \approx 1.07818872957581848275826543676983238170721 9, \\\qquad C_{2} &:= \sum_{n=1}^{\infty} \frac{\zeta(2n)}{n!} \\ & \approx 2.40744655479032851470948665622302272558 2266, \text{ and} \\ C_{3} &:= \sum_{n=1}^{\infty} \frac{\zeta(2n)}{(2n)!} \\ &\approx 0.86900199196290899881105480556139568889 2494.\end{align*} It is also stated that "These sums have no known closed-form expression". However, it could be the case that there are conjectured closed forms for these constants that correspond to the actual values with a high degree of accuracy. Question: Are there any of such conjectured closed forms for

C_{1}, C_{2}, and C_{3} ?

论坛问题帖 | no_answers_at_retrieval | Max Lonysa Muller | 2022-08-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019229 | MSE-4508811

Proof-verification request: On the equation $\gcd(n^2,\sigma(n^2)) = D(n^2)/s(q^k)$

(Preamble: This question is an offshoot of this earlier post.) In what follows, denote the classical sum of divisors of the positive integer x by $\sigma(x)=\sigma_1(x)$, the deficiency of x by $D(x)=2x-\sigma(x)$, and the aliquot sum of x by $s(x)=\sigma(x)-x$. Let $N = q^k n^2$ be an odd perfect number with special prime q satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. Descartes, Frenicle, and subsequently Sorli conjectured that $k=1$ always holds. (In this proof-verification request, we shall attempt to prove this conjecture.) Using the fundamental equation $$\sigma(q^k)\sigma(n^2)=\sigma(q^k n^2)=\sigma(N)=2N=2q^k n^2$$ and using the fact that q^k and $\sigma(q^k)$ are relatively prime (that is, $\gcd(q^k,\sigma(q^k))=1$), then we get that $q^k \mid \sigma(n^2)$, whereupon we obtain that $$\frac{\sigma(n^2)}{q^k}=\frac{2n^2}{\sigma(q^k)}$$ is an odd integer. From the hyperlinked MSE question above, we have obtained the identities $$\gcd\bigl(n^2,\sigma(n^2)\bigr)=\frac{\sigma(n^2)}{q^k}=\frac{2n^2}{\sigma(q^k)}=\frac{D(n^2)}{s(q^k)}=\frac{2s(n^2)}{D(q^k)}.$$ Now, in a comment underneath MSE user Aravind's answer to the hyperlinked MSE question above, we derived that $$\gcd\bigl(n^2,\sigma(n^2)\bigr)=n^2 \bigl(q^k t - 2(q - 1)\bigr) + \sigma(n^2) \bigl(-(\sigma(q^k)/2)\cdot t + q \bigr), \tag{1}$$ where t is necessarily an integer to be determined by using the identity $$\gcd\bigl(n^2,\sigma(n^2)\bigr)=\frac{D(n^2)}{s(q^k)}. \tag{2}$$ Equating (1) and (2), we obtain $$\frac{D(n^2)}{s(q^k)}=n^2 \bigl(q^k t - 2(q - 1)\bigr) + \sigma(n^2) \bigl(-(\sigma(q^k)/2)\cdot t + q \bigr). \tag{3}$$ Multiplying both sides of (3) by $s(q^k)$, we obtain $$2n^2 - \sigma(n^2) = D(n^2) = n^2 \bigl(q^k s(q^k) t - 2(q - 1)s(q^k)\bigr) + \sigma(n^2) \bigl(- (\sigma(q^k)/2) s(q^k) \cdot t + q s(q^k) \bigr). \tag{4}$$ Equating coefficients for n^2 and $\sigma(n^2)$ in (4), we have that $$\begin{cases} 2 = q^k s(q^k) t - 2(q - 1)s(q^k) \\ -1 = - (\sigma(q^k)/2) s(q^k) \cdot t + q s(q^k) \end{cases} \tag{5} $$ The system in (5) is a set of two simultaneous "linear" equations in t, of the form $$\begin{cases} A t + B = 2 \\ C t + D = -1 \end{cases} \tag{6} $$ where $$A = q^k s(q^k), B = - 2(q - 1)s(q^k)$$ and $$C = - (\sigma(q^k)/2) s(q^k), D = q s(q^k).$$ We now solve the system (6) for t. We thereby get $$\begin{cases} 2D = ADt + BD \\ B = -BCt - BD \end{cases} \tag{7} $$ Adding the two equations in (7), we obtain $$2D + B = (AD - BC)t$$ $$t = \frac{2D + B}{AD - BC}.$$ But $$2D + B = 2qs(q^k) - 2(q - 1)s(q^k) = s(q^k)\bigl(2q - 2(q - 1)\bigr) = 2s(q^k) \tag{8}$$ and $$AD - BC = q^{k+1} \bigl(s(q^k)\bigr)^2 - (q - 1)\sigma(q^k) \bigl(s(q^k)\bigr)^2 = \bigl(s(q^k)\bigr)^2 \bigl(q^{k+1} - (q - 1)\sigma(q^k)\bigr) $$ $$= \bigl(s(q^k)\bigr)^2 \bigl(q^{k+1} - (q^{k+1} - 1)\bigr) = \bigl(s(q^k)\bigr)^2. \tag{9}$$ Consequently, from (8) and (9), the variable t (which is necessarily an integer) must have the value $$t = \frac{2D + B}{AD - BC} = \frac{2s(q^k)}{\bigl(s(q^k)\bigr)^2} = \frac{2}{s(q^k)}.$$ Since $s(q^k) \geq 1$ holds in general, then t is an integer implies that either $s(q^k) = 1$ or $s(q^k) = 2$. Want to show: $s(q^k) = 1$ Assume to the contrary that $s(q^k) = 2$. This is equivalent to $$q^k - 1 = 2(q - 1)$$ $$q^k - 2q = -1$$ $$2q - q^k = 1$$ $$q\bigl(2 - q^{k-1}\bigr) = 1$$ This implies that $q = 1$ AND $2 - q^{k - 1} = 1$, from which we obtain $1 = q^{k - 1}$ and $q = 1$. The first equation implies that $k = 1$, while the second condition contradicts q being the special prime satisfying $q \equiv 1 \pmod 4$ (so that $q \geq 5$). Hence, necessarily we have $s(q^k) = 1$, or $k = 1$. Here is my question: Is the method of equating coefficients for n^2 and $\sigma(n^2)$ in (5) a valid approach?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2022-08-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019230 | MSE-4512740

Show that: $e^{\pi}-\pi^{e} >\frac12$ without calculator

Show that: $$e^{\pi}-\pi^{e} >\frac12$$ I am not sure, if anything can be done using elementary methods. Because e and π are not algebraic numbers. Therefore, I find it impossible to prove this with elementary techniques. Maybe using the elementary formula $\left(1+\frac 1n\right)^n\rightarrow e$ for e would work. But there is no such formula for π. I guess there is no other way than using series, right?

论坛问题帖 | has_answers | User | 2022-08-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019231 | MSE-4520132

Express any positive integer as $\lfloor (\sqrt\ )^m (4(!^n))\rfloor$?

For some reason, I recently recalled a math puzzle column from many years ago about expressing integers in terms of some number of fours. The column noted that there was a conjecture to the effect that every positive integer can be expressed as $\left\lfloor\overbrace{\sqrt{\ \ldots \sqrt{4\underbrace{!\ \ldots\ !}_{n\text{ factorials}}}}}^{m\text{ square roots}}\right\rfloor\tag*{}$ for some non-negative (or was it positive?) integers m and n (possibly in more than one way). For instance, $5 = \left\lfloor\sqrt{\sqrt{\sqrt{\sqrt{\sqrt{4!!\,}}}}}\right\rfloor.\tag*{}$ Has any progress been made on this conjecture in the last 40 or so years? How would one even approach the general case? What about if 4 is replaced by some other integer greater than 2?

论坛问题帖 | has_answers | Ted Hopp | 2022-08-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019232 | MSE-4521411

An inequality holding for odd primes

Let $P_{\Sigma}(n)$ be the sum of all prime factors of n. Then there seems to be an inequality $$p\ge P_{\Sigma}(rad(p^2-1))$$ for all odd primes p. Tested for the first million primes. Is this a known fact?

论坛问题帖 | has_answers | Lehs | 2022-08-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019233 | MSE-4522813

A useful way of characterizing primes?

It's proved that if $P(n)$ is the sum of all different prime factors of n, then for all odd primes p there is an inequality $$p\ge P(p^2-1)$$ An inequality holding for odd primes Therefore there is an injection $j:\mathbb P \to \mathbb Q$, defined as $j(p)=\frac{p}{P(p^2-1)}$. Now to me the following conjecture seems very natural: Given $\varepsilon >0$ there are two primes p,q such that $|j(p)-j(q)|<\varepsilon$ The idea is to characterizing primes p by the factorizations of their neighbors $p-1,p+1$. A metric on the prime numbers? $d(p,q)=|j(p)-j(q)|$ seems to fulfill the triangle inequality. Tested for all combinations of the 100 first primes. I now realize that d must be a metric because j is an injection. It is possible to define convergence of sequences of primes.

论坛问题帖 | no_answers_at_retrieval | Lehs | 2022-09-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019234 | MSE-4524551

Every onto map f such that $f(xy)=f(x)+f(y)+f(x)f(y)$,$f : \Bbb{Z}\setminus 0 \to\Bbb{Z}/2$ is represented by some prime number p and $f=(p\mid-)$

Let $(n \mid x)$ mean 1 if $n \mid x$ and 0 otherwise. Let $f = (p\mid -)$ then $f: \Bbb{Z}\setminus 0 \to \Bbb{Z}/2$ is surjective and only satisfies $$f(xy) = f(x) + f(y) + f(x)f(y), \ \forall x,y \in \Bbb{Z}\setminus 0$$ because p is a prime number. I.e. if p were not prime then it would fail to hold that when $p\mid xy$ then $p \mid x$ or $p\mid y$ in general. The surjectivity part rules out $p = 0, 1$. I'm wondering if a converse is true, i.e. for any such surjective map f, does there exist a prime p such that $f = (p \mid -)$?

论坛问题帖 | has_answers | TotoposDiagramChaseApp | 2022-09-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019235 | MSE-4533307

If p can divide $a^n+b^n+c^n$, can p^k divide it as well?

Related to this Is there a method to decide whether a given function of the form $f(n)=a^n+b^n+c^n$ (a,b,c fixed positive integers , n running over the positive integers) satisfies the following property : If p is a prime number with $p\mid f(s)$ for some positive integer s and k is any positive integer , then there is a positive integer t with $p^k\mid f(t)$. In other words , if a prime can divide this expression, every power of this prime can it as well. I am particular interested in the case $(a,b,c)=(5,6,10)$, so $f(n)=5^n+6^n+10^n$ because in this case my prime search revealed a probable prime quite late (for $n=3168$) , but no other probable prime upto $n=60\ 000$. Furthermore I try to factor numbers of this form the last months (see the linked question) I tried to use that $p\mid f(n)$ is equivalent to $p\mid u^n+2^n+1$ where u is one of $\frac{p+6}{5},\frac{2p+6}{5},\frac{3p+6}{5},\frac{4p+6}{5}$ depending on which of those is an integer. But this led to nowhere. For this special case $(a,b,c)=(5,6,10)$ I found out the following : For $p\le 2\cdot 10^5$ and $k=2$ the conjecture is true. For $p\le 14\ 000$ and $3\le k\le 5$ the conjecture is true.

论坛问题帖 | has_answers | Peter | 2022-09-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019236 | MSE-4545592

Is this conjecture well-formed?

For each positive integer n, one of these is true, where $a,b,c,...etc.$ are natural numbers: $$ \begin{array}{l} 3^0\cdot{n}=2^a\\ 3^1\cdot{n}=2^a-(3^0\cdot2^b)\\ 3^2\cdot{n}=2^a-(3^0\cdot2^b)-(3^1\cdot2^c)\\ 3^3\cdot{n}=2^a-(3^0\cdot2^b)-(3^1\cdot2^c)-(3^2\cdot2^d)\\ ... etc. \end{array} $$ (The specific values of $2^a, 2^b, 2^c, ...etc.$ dependent on n, e.g. all powers of two satisfy $3^0\cdot{n}=2^a$ for some value of a) Or, more succinctly: $$ \forall{n}\in\mathbb{N}^+\exists{x}\in\mathbb{N}:n={2^{a_x}+\sum\limits_{y=0}^{x-1}-(3^y\cdot2^{a_y})\over3^x} $$

论坛问题帖 | no_answers_at_retrieval | Michael | 2022-10-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019237 | MSE-4552013

$\sum{x^{(i,n)}}+1$ is irreducible?

A simple calculation using Burnside lemma shows that the number of distinct "necklaces" with n balls of x colors is $$\frac{F_n(x)}{n} = \frac{\sum_{i=1}^n x^{(i,n)}}{n} = \frac{\sum_{d \mid n} x^d \varphi\left(\frac{n}{d}\right)}{n}.$$ It's natural to ask more about this polynomial F_n. A few attemptions shows that $F_n + 1$ might be irreducible over $\mathbb Q$: $$ F_2(x) + 1 = x^2 + x + 1,\\ F_3(x) + 1 = x^3 + 2x + 1,\\ F_4(x) + 1 = x^4 + x^2 + 2x + 1,\\ \cdots $$ In fact, a computer search up to F_{2000} shows that F_i is irreducible for all $i \le 2000$. QUESTION. Is F_n+1 irreducible over $\mathbb Q$ for all $n \ge 2$?

论坛问题帖 | has_answers | atzlt | 2022-10-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019238 | MSE-4563618

Prove or disprove: If $f(x)$ is continuous in $(0,1]$ and $f(x)\to\infty$ as $x\to 0^+$, then $\lim_{n\to\infty}\sum_{k=1}^n f(k/n)$ does not exist.

I'm trying to prove or disprove the following conjecture: If $f(x)$ is continuous in $(0,1]$ and $f(x)\to\infty$ as $x\to 0^+$ then $L=\lim\limits_{n\to\infty}\sum\limits_{k=1}^n f\left(\frac{k}{n}\right)$ does not exist. My attempt I tried proof by contradiction. Asssume L exists. $L=\lim\limits_{n\to\infty}n(\frac{1}{n})\sum\limits_{k=1}^n f\left(\frac{k}{n}\right)=\left(\lim\limits_{n\to\infty}n\right)\int_{0}^1 f(x)dx$ (EDIT: As mentioned by @FShrike in the comments, the previous step is not valid.) $\therefore \int_{0}^1 f(x)dx=0$ There are functions $f(x)$, continuous in $(0,1]$, such that $f(x)\to\infty$ as $x\to 0^+$ and $\int_{0}^1 f(x)dx=0$. For example, $f(x)=-\ln{x}-1$, in which case L does not exist, by Stirling's approximation. But I do not see why all such functions $f(x)$ would imply that L does not exist. Context: I am interested in geometrical infinite products (example1, example2). The conjecture in this question, via the substitution $f(x)=-\ln{g(x)}$, is equivalent to: If $g(x)$ is continuous in $(0,1]$ and $\lim\limits_{x\to 0^+}g(x)=0$ then

$\lim\limits_{n\to\infty}\prod\limits_{k=1}^ng\left(\frac{k}{n}\right)$ either equals 0 or does not exist, which stands in interesting contrast with the fact that infinite products of lengths or areas, that tend to 0, can equal a positive number. EDIT2: I'm not sure if this is helpful, but I have noticed that $L_2=\lim\limits_{n\to\infty}\sum\limits_{k=1}^n f\left(\frac{k-1/2}{n}\right)$ can exist. For example, $\lim\limits_{n\to\infty}\sum\limits_{k=1}^n \left(-\ln{\left(\frac{k-1/2}{n}\right)}-1\right)=-\frac{\ln{2}}{2}$. (Another question of mine yielded methods for dealing with the sum $\sum\limits_{k=1}^n \ln{(k-\frac12)}$.) I do not understand why replacing k with $k-\frac12$ seems to make the limit existable (if that's a word).

论坛问题帖 | has_answers | Dan | 2022-10-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019239 | MSE-4567232

Find conjugate function

$A\in R^{m\times n},\ rankA = m,\ f(y) -\text{ Convex function,} \ f:R^m\rightarrow R.$ Let $g(x) = f(Ax)$ Find $g^*(y)$. Because$x\rightarrow Ax$ - Affine transformation $\implies g(x)$ - is convex as well. Because $rank{A}=m \implies g:R^m \rightarrow R$. Because $g(x)$ is convex upwards $-g(x)$ is convex downwards and has sup (only one). Let's notice that $<x,y> - g(x)$ is also convex downwards. $$k(x,y)=<x,y>-g(x)$$ $$k_x'(x,y)=\bar{y}-\nabla g(x)=\bar{y}- A^T\nabla f(Ax)=0$$ $$\nabla f(Ax) = (A^T)^{-1}y$$ $$g^*(y) = \sup_{R^m}{k(x,y)}=<x,y>-...$$ How can I precede starting from this moment?

论坛问题帖 | has_answers | Johnaaa11 | 2022-11-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019240 | MSE-4567888

Calculation of generalized Artin's constants

Let $T(p)$ be the period of the decimal expansion of $1/p$, for prime p (e.g. $1/7=0.\overline{142857}\rightarrow T(7)=6$). It is known that $$T(p)=\frac{p-1}{t}$$ for some integer t. Then, Artin's conjecture states that the fraction of primes for which $t=1$ is positive and is given by Artin's constant: $$ A_1=\lim_{x\rightarrow\infty}\frac{\#\{p:\frac{p-1}{T(p)}=1\text{ } \ {\rm and } \ p\leq x\}}{\#\{p:p\leq x\}}=\prod_{p} \left(1-\frac{1}{p(p-1)}\right)=0.3739558136... $$ Artin's constant can be generalized by considering the fraction A_t of primes that satisfy some arbitrary t. My question is: Is there some known formula to calculate it for any t? By searching about the subject, I found some articles that seem to give expressions for that, but they are quite hard to read and understand. Ram Murty in this article and Wagstaff in this one seem to imply that $$A_t=\frac{A_1}{t^2}\prod_{p\mid t}\left(\frac{p^2-1}{p^2-p-1}\right)$$ But by considering the first 1,000,000 primes, it doesn't seem to give the correct result for some values of t, such as $t=40$. So, in addition to the first question, what are the limitations of the above formula?

论坛问题帖 | has_answers | ordptt | 2022-11-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019241 | MSE-4569603

How proof The shape of the two hearts in $\mathcal{D}^b(A_n)$ is the same if the graded undergrphas of the Ext quiver of heart of are equal

According to Yu Qiu 's Ext-quivers of hearts of A-type and the orientation of associahedron, We know hearts in $\mathcal{D}^b(A_n)$ one by one corresponds to the Ext quivers and are precisely the associated quivers of graded gentle trees with n vertices. And it's not hard to noticed if two hearts if the shapes are equal then the inverse proposition of the proposition in the title holds. Absolutely,the proposition and inverse proposition is trivia and I have proven the inverse proposition. But I can't find a simple proof to describe the title. Example in $\mathcal{D}^b(A_3)$,A_3 has straight orientation, $\mathcal{H}_1=<S_1,S_2,S_3>$,$\mathcal{Q}(\mathcal{H}_1)=S_1\xrightarrow{1}S_2\xrightarrow{1}S_3$ $\mathcal{H}_2=<S_1[-1],P_1,S_3[1]>$,$\mathcal{Q}(\mathcal{H}_1)=S_3[1]\xrightarrow{1}P_1\xrightarrow{1}S_1[-1]$ the under graph of ext quiver are $\bullet\xrightarrow{1}\bullet\xrightarrow{1}\bullet$, The two heart has the same shape (A positive triangle, an inverted triangle) Question: My main problem is that when n is large enough, Ext quiver itself will start to get complicated, and simply using Hom discussion is extremely tedious.Do we have a good idea or lemma to solve it Thanks for everyone! Remark: Let $\mathcal{H}$ be a finite heart in a derived category $\mathcal{D}^b(A_n)$. The Ext-quiver $\mathcal{Q}(\mathcal{H})$ is the (positively) graded quiver whose vertices are the simples of $\mathcal{H}$ and whose degree k arrows $S_i\rightarrow S_j$ correspond to a basis of $\mathrm{Hom}_\mathcal{D}(S_i,S_j[k])$.

论坛问题帖 | no_answers_at_retrieval | muyun | 2022-11-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019242 | MSE-4572630

Is this stronger conjecture about near-rep digit prime numbers true?

Inspired by this post, I established an even stronger conjecture: If we have the number $1\ldots 1$ with $k\ge 3$ digits 1 in decimal expansion, then we can replace a digit which is neither the first nor the last by one of $2,3,\ldots ,9$, to get a prime number. More mathematically formulated Let $k\ge 3$ be an integer. Then, there exists an integer j with $1\le j\le k-2$ and an integer m with $1\le m\le 8$, such that $$\frac{10^k-1}{9}+m\cdot 10^j$$ is prime (which is a near-rep digit prime number as desired in the question). This is a significant strengthening of what is asked in the question. Heuristic: The desired prime is roughly 10^{k-1}, a random number with approximately this magnitude is prime with probability about $\frac{1}{\ln(10)\cdot (k-1)}$ and since we have the choice of $8\cdot (k-2)$ numbers, for $k\ge 1000$, the expected number of primes is about 3.47. Computational result: The conjecture is true upto $k=1000$ (I proved the primality of the found examples until $k=500$, for the rest of the range the very realiable BPSW-test was used). Any idea, reference or extension of the search limit would be very welcome.

论坛问题帖 | no_answers_at_retrieval | Peter | 2022-11-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019243 | MSE-4574363

Any "reasonable" and "natural" false graph theory conjectures for which the Petersen graph is not a counterexample?

I heard that the Petersen graph is a good test of a graph theory conjecture. As in, if you want to know whether a graph theory conjecture is false, the Petersen graph is likely to be a counterexample. So, with that in mind, are there any "reasonable" and "natural" graph theory conjectures, which is false, but for which the Petersen graph is NOT a counterexample? I would like a list of several such conjectures. Note, I do not have a formal definition of "reasonable" or "natural", but I know it when I see it.

论坛问题帖 | no_answers_at_retrieval | user107952 | 2022-11-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019244 | MSE-4576393

If n is square-free and $n+1 \mid \sigma(n)$, is n a prime?

As the title says, is n being square-free together with $n+1 \mid \sigma(n)$ sufficient to show n is prime? It is well-known that if $\varphi(n)\mid n-1$ and $n+1 \mid \sigma(n)$, then n is a prime (see e.g. An integer is prime iff $\phi(n) \mid n-1$ and $n+1 \mid \sigma(n)$). Also it is a well-known open problem - Lehmer's totient problem - whether $\varphi(n)\mid n-1$ itself implies that n is a prime. Since $\varphi(n)\mid n-1$ implies that n is square-free, this makes me wondering what is known about the problem in between these two, i.e. n being square-free together with $n+1 \mid \sigma(n)$. I've checked by brute force that the conjecture holds up for n being a product of any of the first 31 primes. As for the proof, we can rule out product of two primes easily as $n=pq$ and $pq+1\mid \sigma(n)=(p+1)(q+1)=pq+p+q+1$ implies $pq+1\mid p+q$ and then we show $pq+1>p+q$ contradicts this result. But it does not seem to generalize to more primes that simply. It might be also that this problem is open, in that case I have yet to find a reference. Update: I've found a proof that only even n that satisfies the conditions is $n=2$, pushing the possible counterexample to odd numbers only. To see this let $n=2\prod_{i=1}^{k} p_i$ where $p_i=2^{e_i}d_i-1$ with $e_i \geq 1$ and d_i odd. Now we have two possibilities, either $e_i=1$, or $e_i \geq 2$. If $e_i=1$, then $d_i \geq 3$, and we have $$ 2^{e_i}d_i-1 \geq 2d_i-1\geq \sqrt[k]{\frac{3}{2}} d_i $$ where the last inequality follows from $(2-\sqrt[k]{\frac{3}{2}})d_i\geq (2-\sqrt[k]{\frac{3}{2}})3\geq 1$, which holds for $k \geq 1$. If on the other hand $e_i\geq 2$, then $$ 2^{e_i}d_i-1 \geq 4d_i-1\geq \sqrt[k]{\frac{3}{2}} d_i $$ this time because $(4-\sqrt[k]{\frac{3}{2}})d_i\geq (4-\sqrt[k]{\frac{3}{2}})\cdot 1\geq 1$, which again holds for $k \geq 1$. Either way we get $2^{e_i}d_i-1\geq \sqrt[k]{\frac{3}{2}} d_i$. But then $$ n+1=2\prod_{i=1}^{k}(2^{e_i}d_i-1)+1\geq 2 \prod_{i=1}^k \sqrt[k]{\frac{3}{2}} d_i +1=3\prod_{i=1}^{k}d_i+1 $$ and at the same time $$ n+1\mid\sigma(n)=(2+1)\prod_{i=1}^{k} 2^{e_i}d_i=3\cdot 2^{\sum e_i}\prod_{i=1}^k d_i. $$ Since $n+1$ is odd, we have $n+1 \mid 3\prod_{i=1}^k d_i$ and so $n+1 \leq 3\prod_{i=1}^k d_i$. But these results are contradictory as we have $$ 3\prod_{i=1}^{k}d_i+1\leq n+1 \leq 3\prod_{i=1}^k d_i. $$ Update 2: Furthermore any counterexample must have $n \equiv 3 \pmod 4$. This follows from C.S.'s argument in the linked answer: If $4\mid n-1$, then $2\mid n+1$ and $4\nmid n+1$. So $v_2(n+1)=1$. If n has k odd prime factors then $2^k \mid \sigma(n)$. But then $2^{k-1}\mid \frac{\sigma(n)}{n+1}$ and in turn $$ 2^{k-1}\leq \frac{\sigma(n)}{n+1}<\frac{\sigma(n)}{n}=\prod_{p\mid n} \frac{p+1}{p}\leq \prod (1+\frac{1}{3})=\Big(\frac{4}{3}\Big)^k $$ which does not hold for $k \geq 2$, so $k=1$ and n is a prime.

论坛问题帖 | has_answers | Sil | 2022-11-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019245 | MSE-4576520

Conjecture: Prove that, $\forall (m,n,k)\in\Bbb Z_{>0}^{3}$ there exists $x\in \Bbb Z_{>0}$, such that: $m^x \equiv x^n\pmod k$

Conjecture: Prove that, $\forall (m,n,k)\in\Bbb Z_{>0}^{3}$ there exists $x\in \Bbb Z_{>0}$, such that: $$m^x \equiv x^n\pmod k$$ The conjecture seems correct. Also this conjecture is possible generalization of the known conjecture. If $m=1$. Then the conjecture is trivial. We have $x^n-1\pmod n$. Then the smallest x equals to $x=n+1$. Because, $x-1\mid x^n-1^n$. The $n=1$ case has been proven before. I tried to solve the general case using the induction, but the induction didn't worked. But, I have many computational results. I chose big prime numbers to check the conjecture. Let $m=31$, $n=17, k=59$. I don't have a software. I've experimented with Wolfram as much as I can. Here is the my input: Table (31^x-x^17)/59 , x=1 to 100 I found that the smallest x equals to 99. These are the things I can do.

论坛问题帖 | has_answers | User | 2022-11-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019246 | MSE-4577030

Is 650 the only solution not fitting in the family?

Inspired by this question The linked question conjectures that $\frac{\sigma(n)}{n+1}$ (where $\sigma(n)$ denotes the divisor-sum function) is not an integer for any squarefree composite number. If we drop the condition that n is squarefree , there are still not many solutions. There is a (perhaps infinite) family of solutions , namely $$n=2^m\cdot (2^{m+1}-3)=2^{2m+1}-3\cdot 2^m$$ where $2^{m+1}-3$ is prime. In this case we have $$\sigma(n)=(2^{m+1}-1)(2^{m+1}-2)=2^{2m+2}-3\cdot 2^{m+1}+2=2(n+1)$$ The only solution I found so far that does not fit into this family is $$650=2\cdot 5^2\cdot 13$$. Current search limit is $n=10^9$ Is this the only such solution ? Since the linked question is still open , this won't be easy , but can we at least solve the case that n is not squarefree ? Has this perhaps already been verified ? If yes, a reference would be nice.

论坛问题帖 | no_answers_at_retrieval | Peter | 2022-11-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019247 | MSE-4577337

Filling space with polycube snakes

One of Martin Gardner's "Mathematical Games" columns (Scientific American June 1981, pp24-29; reprinted in The Last Recreations (1997), pp274-283, and The Colossal Book of Mathematics (2001) pp203-211) describes the following problem due to Scott Kim: First we must define a snake. It is a single connected chain of identical unit cubes joined at their faces in such a way that each cube (except for a cube at the end of a chain) is attached face to face to exactly two other cubes. The snake may twist in any possible direction, provided no internal cube abuts the face of any cube other than its two immediate neighbors. The snake may, however, twist so that any number of its cubes touch along edges or at corners. A polycube snake may [...] have just one end and be infinite in length, or it may be infinite and endless in both directions. We now ask a deceptively simple question. What is the smallest number of snakes needed to fill all space? We can put it another way: Imagine space to be completely packed with an infinite number of unit cubes. What is the smallest number of snakes into which it can be dissected by cutting along the planes that define the cubes? [...] Kim has found a way of twisting four infinitely long one-ended snakes into a structure of interlocked helical shapes that fill all space. The method is too complicated to explain in a limited space; you will have to take my word that it can be done. [...] Kim has conjectured that in a space of n dimensions the minimum number of snakes that completely fill it is $2(n-1)$, but the guess is still a shaky one. In the reprints (I can't find this ever printed in the Scientific American back issues, but maybe I missed it), Gardner adds: Dr. Koh Chor Jin, a physicist at the National University of Singapore, sent a clever proof that given a finite volume of space it is possible to cover it with two of Kim's cube-connected snakes. However, as Kim pointed out, Jin's construction does not approach all of space as a limit [...] So my question for StackExchange is: What was Scott Kim's (or, what is a) four-snake space-filling construction? And what was Koh Chor-Jin's (or, what is a) way to cover any finite volume with two snakes? This guy is also looking for information on the problem. Footnote: At first I wondered if $2(n-1)$ was a typo for 2^{n-1}, since obviously it takes one snake (not zero snakes) to fill 1-space. But it's consistently typeset as $2(n-1)$ in all three primary sources, so I think the "for $n\ge 2$" is implied.

论坛问题帖 | has_answers | Quuxplusone | 2022-11-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019248 | MSE-4590337

Regular polygon of radius 1 with diagonals: mysterious ring of radius $1/e$?

I was playing with a geogebra applet that shows regular n-gons of radius 1 with their diagonals. For example, here is the 12-gon with its diagonals: For any value of n, when I shrink the image, the image becomes darkened (due to the thickness of the lines), but sometimes there remains a faint single white ring, indicating a ring of exceptionally large cells. For example: $n=20$ $n=45$ $n=50$ Now here's the interesting thing: The radius of the white ring always seems to be approximately $1/e$. (I used Perfect Screen Ruler.) For some n-values, I cannot perceive a single white ring; I guess it still exists but is not perceivable due to limitations in pixelation and/or visual acuity. I can formalize my conjecture as follows: In a regular n-gon of radius 1 with it diagonals, if $d_n=$ distance between the centre and the centroid of one of the cells with the greatest area (excluding the centre cell when n is odd), then $$\lim\limits_{n\to\infty}d_n=\frac{1}{e}$$ Question: Is my conjecture true? (This question was inspired by another question: Distribution of areas in regular n-gon with diagonals, as $n\to\infty$.)

论坛问题帖 | has_answers | Dan | 2022-12-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019249 | MSE-4599735

Why is it not known that $\sum_{n>0}\frac{(-1)^nn}{p_n}$ converges?

No one knows if this sum $$\sum_{n>0}\frac{(-1)^nn}{p_n}$$converges (where p_n is the nth prime). But what makes proving this so hard? We know that $$\bigg|\sum_{n>0}\frac{(-1)^n}{p_n}\bigg|<\frac{1}{2}$$converges and is similar to the other sum. Remark: Apparently: $$\sum_{n>2}\frac{(-1)^n}{\pi(n)}$$converges, where $\pi(n)$ is the prime counting function.

论坛问题帖 | no_answers_at_retrieval | Kamal Saleh | 2022-12-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019250 | MSE-4606406

Conjecture about the representation of a constant $C=0.6516...$

It's a follow up of my previous question How to find the constant C such that $f(x)\geq Cx$: We start with : $$\left|\exp\left(1-\prod_{k=1}^{1000}\left(1-\frac{1}{2^{k}(k+1)}\right)\right)-\sqrt{2}-\frac{1}{2^{9}}-\frac{1}{2^{11}}-\frac{1}{2^{14}}-\frac{1}{2^{16}}+\frac{1}{2^{18}}-\frac{1}{2^{20}}-\frac{1}{2^{21}}-\frac{1}{2^{24}}+\frac{1}{2^{27}}+\frac{1}{2^{30}}-\frac{1}{2^{33}}+\frac{1}{2^{36}}+\frac{1}{2^{38}}+\frac{1}{2^{39}}-\frac{1}{2^{41}}\right|<4*10^{-13}$$ Conjecture If $a_n\in N_{\geq 9}$: $$\exp\left(1-\prod_{k=1}^{\infty}\left(1-\frac{1}{2^{k}(k+1)}\right)\right)=\sqrt{2}+\sum_{n=1}^{\infty}\pm\frac{1}{2^{a_n}}$$ I want to show that : $$a_{n+1}>a_n$$ $$10>a_{n+1}-a_n$$ The $\pm$ means it's either positive or negative . I have tried until $n=50$ without a counter-example . As in my answer using the exponent I don't find it in oeis . The a_n seems also steady enought to believe in this conjecture .I have tried also to represent the power of two with an integral without a significant result .For example we have : $$\int_{0}^{1}-\ln\left(x^{\frac{1}{2^{n+1}}}\left(1-x\right)^{\frac{1}{2^{n+1}}}\right)dx=1/2^n$$ We can also speak about the (ir)rationality of C. Question : How to (dis)prove it ? Thanks in advance .

论坛问题帖 | has_answers | Barackouda | 2022-12-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019251 | MSE-4608748

How can I determine the reason for the mismatch between the value of this infinite continued fraction and the expected result?

How can I determine the reason for the mismatch between the value of this infinite continued fraction and the expected result? $$6.9431070487= \lim\limits_{n\rightarrow\infty}7-\frac{7}{129- \dfrac{4096}{761- \dfrac{177147}{\ddots- \dfrac{n^{10}+6n^{9}}{n^{5}+ \left (n+ 1 \right)^{5}+ 6\left (n+ 1 \right)^{4}}}}}$$ This is a conjecture got from Ramanujan Machine and 6.9431070487 is also the expected result I am trying to compare to the value of the continued fraction. I hope to receive any relevant details or background information for help with.

论坛问题帖 | no_answers_at_retrieval | user1135362 | 2022-12-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019252 | MSE-4614502

Given the 2^n primorial divisors, $D(5\#) = \{ 1, 2, 3, 5, 6, 10, 15, 30\}$, why are there always exactly 2^n different divisor indicator vectors?

Define $D(m)$ to be the (ordered) set of divisors of $m \in \Bbb{N}$. Let $m = p_n\# = 1 \cdot 2 \cdot 3 \cdot 5 \cdots p_n$. Clearly, $|D(m)| = $ the number of divisors of $p_n\# = 2^n$. Now define a divisor indicator vector at x to be $\langle (d \mid x) : d \in D(m) \rangle$ where $(d \mid x) = \begin{cases} 1, \text{ if } d \mid x \\ 0, \text{ if } d \nmid x \end{cases}$; for any given $x \in \Bbb{Z}$. Conjecture 1. The number of possible divisor indicator vectors is precisely 2^n. For example: 1 2 3 5 6 10 15 30 1 0 0 0 0 0 0 0 1 1 0 0 0 0 0 0 1 0 1 0 0 0 0 0 1 0 0 1 0 0 0 0 1 1 1 0 1 0 0 0 1 1 0 1 0 1 0 0 1 0 1 1 0 0 1 0 1 1 1 1 1 1 1 1 I.e. there are 8 divisors so the number of divisor indicator vectors is also 8. Clearly it's since if $15 \mid x$ then both $3 \mid x$ and $5\mid x$ and vise-versa, since $\gcd(3,5) = 1$, but how would the argument go for the general case? My next question which I think is better is about this conjecture: Conjecture 2. The set of all possible divisor indicator vectors is linearly independent over any field. I say any field because every field by definition contains the values of the divisor indicator $(d \mid \cdot)$ or 0 and 1. So for each field F you'd define $(d\mid \cdot) : \Bbb{Z} \to \{0, 1\} \subset F$. Now this might have something to do with the first conjecture since our degrees of freedom are precisely determined by the n primes involved and taken together (just the prime components): 2 3 5 1 0 0 0 1 0 0 0 1 we have clear linear independence. What is the easiest & most elegant way you can think of to prove both of these conjectures?

论坛问题帖 | has_answers | TotoposDiagramChaseApp | 2023-01-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019253 | MSE-4624875

Graphs with all vertices having more neighbors than 2nd order neighbors

This is the conjecture that inspired my question: https://en.wikipedia.org/wiki/Second_neighborhood_problem For any vertex v, let $N(v)$ be the set of vertices adjacent to v and let $N^2(v)$ the set 2nd order neighbors, thus all vertices that have a distance of exactly 2 relative to v. The conjecture concerns directed graphs, but I was interested in undirected simple graphs with each vertex having strictly more neighbors than second order neighbors. A basic example would be a graph of 4 vertices, completely connected except for one edge. I believe such graphs are exactly the ones that have the following characteristic: for each vertex v, more than half of all possible 2-step walks starting from v have an endpoint in $N(v)$ So far I haven't encountered a counter example but neither can I prove it, it's somewhat intuitive but not completely. Would love to hear your thoughts.

论坛问题帖 | has_answers | AndroidBeginner | 2023-01-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019254 | MSE-4625084

Do all Turing-complete systems converge on a single universal function w.r.t. description redundancy?

Let L be any Turing-complete language. Let $d_L(n)$ be the number of distinct algorithms expressible in L using at most n bits. I'm not sure how to properly define "distinct algorithm", but broadly it's what you would expect: multiple halting programs returning the same value are considered equivalent to one another, as are cyclic programs stepping through the same sequence, as are runaway programs exhibiting identical growth. I assert that $$\frac{d_L(n)}{n} \to D(n)\qquad\text{as }n \to \infty,$$ where D is some unknown but very real function. I would imagine D is nearly or completely uncomputable, and perhaps monotonically increasing with n. Most importantly, I assert this holds regardless of choice of L, in contrast to Ω_L and related functions. The short version of my reasoning for this claim is that it is an immediate consequence of the no free lunch theorem, which states that over a uniform distribution of all possible binary strings, no compression algorithm will outperform any other. My understanding is that it's equivalent or similar to the principle that there is no such thing an ideal optimizing compiler (the "full-employment theorem"). In this case, since any candidate language L is aggregating its results over all possible binary strings up to length n, no one language should be able to come out ahead in terms of economy of representation. This should also be apparent by a simple counting argument. This means that, in general, any choice of L will encode the same total number of bits after running it over all $2^n -1$ possible descriptions. However, I think it may be the case that that only holds perfectly when the encoding is perfectly efficient, as for example with universal prefix-free languages. Even if I've gotten that wrong, thanks to Kolmogorov invariance there will be some constant c (given a finite set of languages) such that $$d_{L_1}(n) \leq d_{L_2}(n) \leq d_{L_1}(n+c)$$ for any choice of L_1,L_2. This provides a bound of $$2^{-c} \leq \frac{d_{L_2}(n)}{d_{L_1}(n)} \leq 2^c.$$ Since c should become negligible as n grows without bound, I would expect that to converge to 1 in the long run, which would imply that the number of distinct descriptions between languages do indeed all approach some single curve $D(n)$ in the limit. While I can't back this argument up formally yet, I see only one potential issue with it; D may be dependent on the set of languages being used. I'm not clear on whether that is likely to be true, but even if it is, so long as you don't let the set of languages grow without bound, the distinct-description proportions should still converge to a single curve. My question: Does this sound generally right, or am I off track? If right, I assume it's a known result and would appreciate a keyword to hunt for. If not, where did I go awry or what is unclear?

论坛问题帖 | no_answers_at_retrieval | Trevor | 2023-01-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019255 | MSE-4632595

Conjecture about difference of Fibonacci numbers and primes

I'm curious to see if this conjecture is true: if $ m > 4 $ is a positive integer not divisible by $ 2 $ or $ 3 $, it's possible to find a positive integer $ n $ such that the difference of the $ 2 $ Fibonacci numbers $ F_{m+n}-F_n $ is a prime number. Various example of $ [m, n] $, with the smallest n, are $[5,3],[7,3], [11,4], [13,5], [17,3],..., [619,1353]...$ I don't know if this could be of any help: I converted the primes I found that way in the Zeckendorf representation and they are an alternating sequence of $ 1 $ and $ 0 $ ending with two adjacent $ 0 $ and $ 1 $ (and eventually just various $ 0 $) Edit: I posted this problem on Mathoverflow too

论坛问题帖 | has_answers | user967210 | 2023-02-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019256 | MSE-4642118

Conjecture. If $n \in \Bbb{N}\setminus 1$ is not a twin prime average, then $n^2 - 1$ is not square-free.

From the evidence: from sympy import * N = 1000 for n in range(2, N): if not isprime(n + 1) or not isprime(n-1): s = n**2 - 1 if rad(s) != s: print(f"True for {n} = n, n^2 - 1 = {s}") else: print(f"False for {n} = n, n^2 - 1 = {s}") Prints all true: ..... True for 963 = n, n^2 - 1 = 927368 True for 964 = n, n^2 - 1 = 929295 True for 965 = n, n^2 - 1 = 931224 True for 966 = n, n^2 - 1 = 933155 True for 967 = n, n^2 - 1 = 935088 True for 968 = n, n^2 - 1 = 937023 True for 969 = n, n^2 - 1 = 938960 True for 970 = n, n^2 - 1 = 940899 True for 971 = n, n^2 - 1 = 942840 True for 972 = n, n^2 - 1 = 944783 True for 973 = n, n^2 - 1 = 946728 True for 974 = n, n^2 - 1 = 948675 True for 975 = n, n^2 - 1 = 950624 True for 976 = n, n^2 - 1 = 952575 True for 977 = n, n^2 - 1 = 954528 True for 978 = n, n^2 - 1 = 956483 True for 979 = n, n^2 - 1 = 958440 True for 980 = n, n^2 - 1 = 960399 True for 981 = n, n^2 - 1 = 962360 True for 982 = n, n^2 - 1 = 964323 True for 983 = n, n^2 - 1 = 966288 True for 984 = n, n^2 - 1 = 968255 True for 985 = n, n^2 - 1 = 970224 True for 986 = n, n^2 - 1 = 972195 True for 987 = n, n^2 - 1 = 974168 True for 988 = n, n^2 - 1 = 976143 True for 989 = n, n^2 - 1 = 978120 True for 990 = n, n^2 - 1 = 980099 True for 991 = n, n^2 - 1 = 982080 True for 992 = n, n^2 - 1 = 984063 True for 993 = n, n^2 - 1 = 986048 True for 994 = n, n^2 - 1 = 988035 True for 995 = n, n^2 - 1 = 990024 True for 996 = n, n^2 - 1 = 992015 True for 997 = n, n^2 - 1 = 994008 True for 998 = n, n^2 - 1 = 996003 True for 999 = n, n^2 - 1 = 998000 It seems that every non-twin prime average $n\gt 1$ is such that $n^2 - 1$ (the product of its would-be primes) is not square-free. How can we prove this by elementary means? from sympy import * N = 100000 k = 2 for n in range(3, N): if not isprime(n + k) or not isprime(n - k): s = n**2 - k**2 if rad(s) != s: print(f"True for {n} = n, n^2 - k^2 = {s}") else: print(f"False for {n} = n, n^2 - k^2 = {s}") assert(0) According to the last code output, I conclude with this conjecture: Generalized Conjecture. For all fixed $k \geq 1$, for sufficiently large $n \in \Bbb{N},$ namely about $n - k \gt 0$, we have that $n \pm k \in \Bbb{P}$ or in other words n is a $2k$-separated prime average if and only if $n^2 - k^2$ is a square-free integer! How can we prove it or has this already been proved?

论坛问题帖 | has_answers | TotoposDiagramChaseApp | 2023-02-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019257 | MSE-4644698

Every twin prime average $x \gt 6$ is the sum of two twin prime averages (Code checked up to $x \leq 1,000,000$).

If p,q are a pair of twin primes, then $x = \dfrac{p+ q}{2} = q-1 = p+1$ is their twin prime average. Conjecture. Every twin prime average $x \gt 6$ is the sum of two smaller twin prime averages, $x = x' + x''$. This doesn't require infinitude. So wondering if we can prove it. 12 = 6 + 6 18 = 6 + 12 30 = 12 + 18 42 = 12 + 30 60 = 18 + 42 72 = 12 + 60 102 = 30 + 72 108 = 6 + 102 138 = 30 + 108 150 = 12 + 138 180 = 30 + 150 192 = 12 + 180 198 = 6 + 192 228 = 30 + 198 240 = 12 + 228 270 = 30 + 240 282 = 12 + 270 312 = 30 + 282 348 = 108 + 240 420 = 72 + 348 432 = 12 + 420 462 = 30 + 432 522 = 60 + 462 570 = 108 + 462 600 = 30 + 570 618 = 18 + 600 642 = 42 + 600 660 = 18 + 642 810 = 150 + 660 822 = 12 + 810 828 = 6 + 822 858 = 30 + 828 882 = 60 + 822 1020 = 138 + 882 1032 = 12 + 1020 1050 = 18 + 1032 1062 = 12 + 1050 1092 = 30 + 1062 1152 = 60 + 1092 1230 = 138 + 1092 1278 = 228 + 1050 1290 = 12 + 1278 1302 = 12 + 1290 1320 = 18 + 1302 1428 = 108 + 1320 1452 = 150 + 1302 1482 = 30 + 1452 1488 = 6 + 1482 1608 = 180 + 1428 1620 = 12 + 1608 1668 = 60 + 1608 1698 = 30 + 1668 1722 = 102 + 1620 1788 = 180 + 1608 1872 = 150 + 1722 1878 = 6 + 1872 1932 = 60 + 1872 1950 = 18 + 1932 1998 = 570 + 1428 2028 = 30 + 1998 2082 = 150 + 1932 2088 = 6 + 2082 2112 = 30 + 2082 2130 = 18 + 2112 2142 = 12 + 2130 2238 = 108 + 2130 2268 = 30 + 2238 2310 = 42 + 2268 2340 = 30 + 2310 2382 = 42 + 2340 2550 = 240 + 2310 2592 = 42 + 2550 2658 = 108 + 2550 2688 = 30 + 2658 2712 = 570 + 2142 2730 = 18 + 2712 2790 = 60 + 2730 2802 = 12 + 2790 2970 = 180 + 2790 3000 = 30 + 2970 3120 = 150 + 2970 3168 = 198 + 2970 3252 = 282 + 2970 3258 = 6 + 3252 3300 = 42 + 3258 3330 = 30 + 3300 3360 = 30 + 3330 3372 = 12 + 3360 3390 = 18 + 3372 3462 = 72 + 3390 3468 = 6 + 3462 3528 = 60 + 3468 3540 = 12 + 3528 3558 = 18 + 3540 3582 = 42 + 3540 3672 = 282 + 3390 3768 = 228 + 3540 3822 = 150 + 3672 3852 = 30 + 3822 3918 = 150 + 3768 3930 = 12 + 3918 4002 = 72 + 3930 4020 = 18 + 4002 4050 = 30 + 4020 4092 = 42 + 4050 4128 = 108 + 4020 4158 = 30 + 4128 4218 = 60 + 4158 4230 = 12 + 4218 4242 = 12 + 4230 4260 = 18 + 4242 4272 = 12 + 4260 4338 = 108 + 4230 4422 = 150 + 4272 4482 = 60 + 4422 4518 = 180 + 4338 4548 = 30 + 4518 4638 = 420 + 4218 4650 = 12 + 4638 4722 = 72 + 4650 4788 = 138 + 4650 4800 = 12 + 4788 4932 = 282 + 4650 4968 = 180 + 4788 5010 = 42 + 4968 5022 = 12 + 5010 5100 = 312 + 4788 5232 = 432 + 4800 5280 = 180 + 5100 5418 = 138 + 5280 5442 = 420 + 5022 5478 = 60 + 5418 5502 = 60 + 5442 5520 = 18 + 5502 5640 = 138 + 5502 5652 = 12 + 5640 5658 = 6 + 5652 5742 = 102 + 5640 5850 = 108 + 5742 5868 = 18 + 5850 5880 = 12 + 5868 6090 = 240 + 5850 6132 = 42 + 6090 6198 = 108 + 6090 6270 = 72 + 6198 6300 = 30 + 6270 6360 = 60 + 6300 6450 = 150 + 6300 6552 = 102 + 6450 6570 = 18 + 6552 6660 = 108 + 6552 6690 = 30 + 6660 6702 = 12 + 6690 6762 = 60 + 6702 6780 = 18 + 6762 6792 = 12 + 6780 6828 = 138 + 6690 6870 = 42 + 6828 6948 = 858 + 6090 6960 = 12 + 6948 7128 = 180 + 6948 7212 = 420 + 6792 7308 = 180 + 7128 7332 = 462 + 6870 7350 = 18 + 7332 7458 = 108 + 7350 7488 = 30 + 7458 7548 = 60 + 7488 7560 = 12 + 7548 7590 = 30 + 7560 7758 = 198 + 7560 7878 = 420 + 7458 7950 = 72 + 7878 8010 = 60 + 7950 8088 = 138 + 7950 8220 = 270 + 7950 8232 = 12 + 8220 8292 = 60 + 8232 8388 = 828 + 7560 8430 = 42 + 8388 8538 = 108 + 8430

8598 = 60 + 8538 8628 = 30 + 8598 8820 = 192 + 8628 8838 = 18 + 8820 8862 = 42 + 8820 8970 = 108 + 8862 9000 = 30 + 8970 9012 = 12 + 9000 9042 = 30 + 9012 9240 = 198 + 9042 9282 = 42 + 9240 9342 = 60 + 9282 9420 = 138 + 9282 9432 = 12 + 9420 9438 = 6 + 9432 9462 = 30 + 9432 9630 = 192 + 9438 9678 = 240 + 9438 9720 = 42 + 9678 9768 = 138 + 9630 9858 = 138 + 9720 9930 = 72 + 9858 How I would use this data is this. Don't try to prove this, but we can safely assume that given a complete list of twin primes $\leq x_n$ each in the list can be assumed to be the sum of two prior. I would, given such a list $X = x_0 \lt x_1 \lt \dots \lt x_n$, assume that for each $k = 1..n$ that the sublist $x_0 \lt \dots \lt x_k$ also has this summatory property. Perhaps from there we can form a contradiction if for each $x, y \in X$, we have that: $$ (x + y)^2 - 1 = 0 \pmod q, \\ \text{ for some prime } q \leq \sqrt{x + y + 1} $$ in a Sieve-of-Eratosthenese way. We must also then note that if $y = \max X$, then $\sqrt{2y + 1} \leq y - 2$ for sufficiently large y, namely $y \geq 6$. That means that the primes involved in showing that for at least one prime q we have $(x+y)^2 - 1= 0 \pmod q$ can be taken to be the same list Q by which each $x \in X$ is such that $x \pm 1 \neq 0 \pmod q$ for some $q \in Q$, i.e. namely all $q \leq y - 2$.

论坛问题帖 | no_answers_at_retrieval | TotoposDiagramChaseApp | 2023-02-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019258 | MSE-4645153

A question about 3-smooth differences between primes

There are an infinite number of primes p such that there are no prime $q<p$ such that $p-q$ is 2-smooth (a power of two) and the smallest such numbers are $p=2,127,149,...$. https://oeis.org/A065381. The same idea for 3-smooth differences yields the question if there are prime numbers $p>2$ such that it doesn't exists a prime $q<p$ such that $p-q$ is 3-smooth? Due to my tests such a prime is greater than 10^{10} if it exists. I do think there is such a failing prime but that my computation skills and computer systems are too limited to find out. Question: is there a failing prime or can it be proved that 3-smooth differences always exists? Comment shows there should be a lot of failing primes, but which is the smallest greater than 2? If $F=\{k\in\mathbb N_{>1} |2^{k-1}\equiv 1\pmod k\}$ one can ask for which $n\in F$ it doesn't exist a $m\in F,\; m<n$ such that $n-m$ is 3-smooth. For all $n\in \{t\in F | 10^{10}\leq t < 2\cdot10^{10}\}$ there exist one such number 19115108601, which is not a prime. A big gap from p downwards to previous prime should decrease the probabillity for 3-smooth gaps. I tested all so far known records of gaps Prime gaps Wikipedia without find an exception. So I add the conjecture tag. Below a diagram of the number of 3-smooth differences as a function of the prime p. The red curve is $\ln p$

论坛问题帖 | no_answers_at_retrieval | Lehs | 2023-02-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019259 | MSE-4649148

Are there an infinity of consecutive primes that have no common digit?

Examples: 2 & 3, 3 & 5, 5 & 7, 59 & 61, 99999989 & 100000007. I was inspired by the fact that, in English, all consecutive numbers share a common letter, such as seven & eight that share 'e'. See Alex Bellos's Math Puzzle section https://www.theguardian.com/science/2023/feb/20/did-you-solve-it-thats-mathematics. Instead I consider consecutive primes and their base representation. Except for 2 & 3, 3 & 5 and 5 & 7 all such base-10 consecutive primes are of the form: $a \times 10^n - b$ and $a \times 10^n + c$ with $1 \le a\le 8$. It is easy to show that a cannot be 9. I found 701 cases below 10^{500}. A large example is $1 \times 10^{101}-203\ \ $ & $\ \ 1 \times 10^{101}+3$ and a larger example (with probable primes) is $8 \times 10^{5002}-6243\ \ $ & $\ \ 8 \times 10^{5002}+14481$. I am pessimistic that the conjecture can be easily proved. Landau's conjecture that there are infinitely many primes p of the form $p=n^2+1$ remains open. For other bases: in base 2 there are no such consecutive primes. In base 3 it is easy to show that there are none, except 2 & 3. Base 4 is the first interesting base. All cases must be of the form $4^n - b$ and $4^n + c$ (except for 2 & 3). I found only 5 cases with n = 1, 4, 28, 83 and 1816. EDIT: Replaced "disjoint" with the more standard "have no common digits". See the result for squares in oeis.org/A156981

论坛问题帖 | has_answers | Lewis Baxter | 2023-03-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019260 | MSE-4652573

A graph where for each pair of vertices, there exist a vértice adjacent to both and a vértice adjacent to neither

Can we construct a simple $G=(V, E) $ so that for any pair $u, v\in V $, there is a vértix x adjacent to both, as well as a vertix y that is not adjacent to either u, v. So, $ux, vx \in E$ and $uy, vy\notin E $. I believe this to be impossible, and therefore conjecture that if every pair of people have a mutual friend, then one pair doesn't have a mutual stranger. Can someone find a counterexample?

论坛问题帖 | has_answers | AndroidBeginner | 2023-03-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019261 | MSE-4653200

Every bipartite graph is an induced subgraph of a hypercube graph?

UPDATE: THIS IS FALSE. I came up with this question some day and have been working on it for some time: Every bipartite graph G is an induced subgraph of a hypercube graph Q_n. Here the hypercube graph is a graph defined as follows: The hypercube graph $Q_n(V_n,E_n)$ with $V_n=\{\text{all subsets of }\{1, 2, ... , n\}\}$, $E_n=\{\{u,v\}:|u\triangle v|=1\}$, where $A\triangle B=(A\setminus B)\cup(B\setminus A)$. With some small examples I believe this is correct. Here's what I've trying to do: by block decomposition we reduce to the biconnected scenario. Every biconnected graph can be constructed as follows: starting from a cycle, and adding paths to the graph. If we can present a way to add the paths in such a way that the add path itself is longer than the original distance of the two ends (in the graph before adding this path), we can always label a subset of $\mathbb{Z}_+$ on every vertices added such that two vertices are incident if and

only if the subset labelled on them has the difference of exactly one element (i.e. $|A\triangle B|=1$), because every cycle is even. (This should be true, I think) In this way we can embed this graph to some Q_n. Now it remains to show the "if" is true. I tried to add the shortest path every time but this doesn't seem to work.

论坛问题帖 | has_answers | atzlt | 2023-03-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019262 | MSE-4655700

Certain primes p which seems to be $\equiv 1\pmod{10}$

$A=\{5,61,181,1741,36721,60901,135721,431521,531481,552301,685621,2834581,3567121,3674761,3696481,4503001,6121501,6811741,9456901,11002741,11524801,12495001,15629641,16068781,18611101,20218441,20231161,25625641,25999261,28042561,33874681,35540881,38377561,42698041\}$ is a set of the smallest primes p such that $\frac{p^2+1}{2}$ is prime and such that there is a prime q with $p=\frac{q^2+1}{2}$. It seems that for all $p\in A$ except $p=5$, $p\equiv 1\pmod{10}$. Can this be proved?

论坛问题帖 | has_answers | Lehs | 2023-03-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019263 | MSE-4657141

Conjectures with the strongest numerical evidence later proven to be false

For a number of conjectures in number theory, the truth of some statement has been checked up to extremely large values. E.g. There are no odd perfect numbers below 10^{1500} Are there any examples where a conjecture eas believed to be true, and also had huge empirical evidence that it should be true, but was later proven to be false?

论坛问题帖 | no_answers_at_retrieval | Davis Yoshida | 2023-03-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019264 | MSE-4657217

Strengthening of FLT

FLT is that there are no positive nontrivial integer solutions to the Diophantine equation $a^n + b^n = c^n$ for $n>2$. What about the conjecture that there are no nontrivial solutions to the DE $a^n + b^n = c^n + d^n$ for $n>4$; is there an explicit name for it? I know that it’s a case of Lander, Parkin, Selfridge, but does this particular case have its own name? It’s so natural that it must have its own name, one would figure.

论坛问题帖 | no_answers_at_retrieval | Benjamin L. Warren | 2023-03-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019265 | MSE-4659494

Friendship theorem for bipartite graphs

The friendship theorem states that if every pair of people has exactly one common friend, then one person is friends with everybody. Which is unfortunately not easy to actually proof. I came up with the following variation on this theorem regarding bipartite graphs: if every pair of females has exactly one mutual male friend, and every pair of males has exactly one mutual female friend (the bipartite friendship condition), but the amount of females and males are not equal, then one male is friends with every girl and one girl is friends with every boy. If the amount of females and males are equal and more than 3, then there is another possible configuration: there is a popular male and popular female, who are friends with everyone from the other gender , except with each other. Everyone else has exactly 2 friends of opposite gender. Can we prove that these are the only bipartite graphs satisfying the bipartite friendship condition?

论坛问题帖 | has_answers | AndroidBeginner | 2023-03-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019266 | MSE-4659512

Formula for the ratio of Numbers not divisible by the first n primes

I was playing around with prime numbers, and specifically how many numbers we can exclude from an interval from being primes. It is easy to see, that after 2, the ratio of numbers is $1-\frac{1}{2}$, as we know every second number to be divisible by 2 and, as such be impossible to be a prime. After we include 3, the ratio would be $1-\frac{1}{2}-\frac{1}{3}$, except that every number divisible by 6 would be excluded once by the $\frac{1}{2}$ and a second time by the $\frac{1}{3}$. Therefore in the actual ratio we have to add every sixth number again. So the actual ratio is $1-\frac{1}{2}-\frac{1}{3}+\frac{1}{6}$. And naturally I wondered how this sequence continues once we include 5. However, the number of terms grows rapidly. This is because 1: We have to add back every tenth and fiveteenth number, again, as we would otherwise subtract them twice by the $\frac{1}{2}$ and $\frac{1}{5}$ or the $\frac{1}{3}$ and $\frac{1}{5}$ respectively. 2: We have to subtract every 30eth Number, as we subtract them thrice from the $2,3,5$ and add them back thrice from the $6,10,15$, so as to not ignore these, we subtract them. This is the sequence I have arrived at, although I could be missing something $1-\frac{1}{2}-\frac{1}{3}+\frac{1}{6}-\frac{1}{5}+\frac{1}{10}+\frac{1}{15}-\frac{1}{30}$ Conjectured sequence If we look at the number of terms, the sequence $1,2,4,8$ appears, furthermore, if we look at the latter half of our sequence for the first 3 prime numbers $(-\frac{1}{5}+\frac{1}{10}+\frac{1}{15}-\frac{1}{30})$, it is just the first half $(1-\frac{1}{2}-\frac{1}{3}+\frac{1}{6})$ multiplied by $-\frac{1}{5}$. So here is my conjecture stated a bit more mathematically: Let p_n be the n-th prime number, $r(n)$ the ratio of natural numbers divisible by the first n prime numbers to the natural numbers, then \begin{multline} r(0)=1,\\ r(n+1)=r(n)-\frac{1}{p_{n+1}}r(n)\\ =r(n)\cdot\left(1-\frac{1}{p_{n+1}}\right)\\ =r(0)\cdot\left(1-\frac{1}{p_1}\right)\left(1-\frac{1}{p_2}\right)...\left(1-\frac{1}{p_{n+1}}\right)\\ =\prod_{i=1}^{n+1}\left(1-\frac{1}{p_i}\right) \end{multline} My question is: is this true? It seems true but again, as the number of terms grow so rapidly I could have missed

something. I am also very certain, that I was not the first to have this idea, but I failed to find any theorem like this on google.

论坛问题帖 | has_answers | Brummi | 2023-03-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019267 | MSE-4660062

Conjecture about the minimum of the Gamma function

Problem/Conjecture: Let the function : $$f(x)=\frac{((x+x_{\min})!-(x_{\min})!)^{\frac{1}{x}}}{x^{\frac{1}{x^2}}}$$ Where $x_\min$ denotes the minimum abscissa of the Gamma function near by 0.4616 Then do we have : $$f'(1-x_{\min})<^?0$$ An computation show it could be true. And what is : $$\lim_{x\to \infty}(f(x)-f(x-1))=?$$ Is it true? If yes is it trivial or well-know ?

论坛问题帖 | has_answers | Barackouda | 2023-03-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019268 | MSE-4662757

Conjecture: Shallow diagonals in Pascal's triangle form polynomials whose roots are all real, distinct, and in $(-2,2)$.

Here are the first few "shallow diagonals" in Pascal's triangle. We can use these shallow diagonals to make polynomials. Note that the even degree polynomials have an extra $\pm x$ at the end. x x^2-x x^3-x x^4-2x^2+x x^5-3x^3+x $x^6-4x^4+3x^2-x$ $x^7-5x^5+6x^3-x$ $x^8-6x^6+10x^4-4x^2+x$ $\dots$ Here is an explicit description (you can skip this part if you already see the pattern): In the nth row of Pascal's triangle (first row is $n=1$), let the leftmost number be the leftmost number of a shallow diagonal. If n is odd, say $2k-1$, call the numbers in this shallow diagonal, from left to right, $a_1, a_3, \dots, a_{2k-1}$, and let $P_{2k-1}(x)=\sum\limits_{i=1}^{k} (-1)^{i-1}a_{2i-1}x^{2(k-i)+1}$. If n is even, say $2k$, call the numbers in this shallow diagonal, from left to right, $a_2, a_4, \dots, a_{2k}$, and let $P_{2k}(x)=\sum\limits_{i=1}^{k} (-1)^{i-1}a_{2i}x^{2(k+1-i)}+(-1)^{k}x$. Conjecture: For each of these polynomials, all of the roots are real, distinct, and in $(-2,2)$. Is my conjecture true? I've been picking random shallow diagonals, and so far I have not found a counter-example. Here are more shallow diagonals, and more even numbered shallow diagonals. Context: I formulated this conjecture when trying to answer another question of mine: For given n, how to find integers $a_0, ..., a_n$ so that $\sum_{k=0}^n a_k x^k$ has n distinct real roots and $\sum_{k=0}^n |a_k|$ is minimized?

论坛问题帖 | has_answers | Dan | 2023-03-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019269 | MSE-4669807

Possible connection between binary numbers and π

Here is the Desmos if you want to follow along: https://www.desmos.com/calculator/b4vtzruupm In messing around with binary numbers, I created a function $f(x)$ in Desmos that generated a list of digits of the binary representation of the input x in reverse. For example, $f(10)$ would output $[0,1,0,1]$. I then created a function $g(z)$ that created a list of points $(z,\text{indexes of all 1s in f(z)}))$. For example, $g(10)$ would output $(10,2),(10,4)$. $(10,1)$ and $(10,3)$ were removed because the first and third numbers in $f(10)$ were 0. Then, I created a list h made up of lists of $g(x)$ where $x\in\mathbb{N}$. After testing a few lists of natural numbers for x. I created a python program that let me put a little more than a thousand values of x in $g(x)$ without having to manually type them in. I then did a regression $h.y=\log_n(h.x)$ where $h.x$ denotes the list of x-values in h and $h.y$ denotes the list of y-values in h. After doing thorough testing with lists such as $[1,...,1000]$ and $[1000,...,1600]$, it seemed like as if I were to use the set of all natural numbers, the value of n from the regression would be equal to π. For example, the range $[600,...,1500]$ gave me $n=3.1562889032$. I used many other ranges too. Conjecture: Pick a positive integer N, and construct a set S of tuples consisting of all pairs $(\log m,i)$ for which $1\le m\le N$ and m has a one in the 2^{i-1}'s place when written in binary. Find the line through the origin of best fit for the points S. Then the slope, as $N\to\infty$, tends to $\log\pi$. It would be much appreciated if anyone could help prove, disprove, and/or explain this. I would also really appreciate if someone helped me edit my post to make it clearer and easier to understand as I struggled with trying to explain what I was doing at points and think it is a little confusing. Thanks in advance.

论坛问题帖 | has_answers | Dylan Levine | 2023-03-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019270 | MSE-4672840

Equivalent definition of convergence of a sequence in $\mathbb{R}$

While reading over the definition of a Cauchy sequence in $\mathbb{R}$, I conjectured that the following condition $\forall \epsilon>0, \exists N\in\mathbb{N}: \forall n>N, |a_n-a_N|<\epsilon$ is necessary and sufficient for a sequence in $\mathbb{R}$ to converge. Is this true? I haven't been able to find any counterexamples or proofs of this conjecture on this site. If this is indeed true, I'm guessing that a proof would involve subsequences and/or the Cauchy sequence condition.

论坛问题帖 | has_answers | FabrizzioMuzz | 2023-04-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019271 | MSE-4678913

Differences between primes which are powers of two

Not for every prime p there is smaller prime q such that $p-q=2^n$ for some natural number n and $n=127$ is an example. But what about the dual question: For each prime number $p>2$ there is a natural number n such that $p+2^n$ is a prime number. What is known about this?

论坛问题帖 | has_answers | Lehs | 2023-04-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019272 | MSE-4690948

Does an odd perfect number have a divisor (other than 1) which must necessarily be almost perfect?

Let $\sigma(x)=\sigma_1(x)$ be the classical sum of divisors of the positive integer x. Denote the aliquot sum of x by $s(x)=\sigma(x)-x$ and the deficiency of x by $d(x)=2x-\sigma(x)$. Finally, denote the abundancy index of x by $I(x)=\sigma(x)/x$. A number P is said to be perfect if $\sigma(P)=2P$. On the other hand, if a number A satisfies $\sigma(A)=2A-1$, then A is said to be almost perfect. (The only known examples for A are the powers of two, including 1.) Lastly, a number D is said to be deficient-perfect if the deficiency $d(D)$ of D divides D. Even Perfect Numbers The three smallest examples for P are $$6=2\cdot{3}={2^1}\cdot\left(2^2 - 1\right)$$ $$28=4\cdot{7}={2^2}\cdot\left(2^3 - 1\right)$$ $$496={16}\cdot{31}={2^4}\cdot\left(2^5 - 1\right).$$ The Euclid-Euler Theorem states that an even number E is perfect if and only if E has the form $$E={2^{p-1}}\cdot\left(2^p - 1\right) = \frac{M_p (M_p + 1)}{2},$$ where $M_p = 2^p - 1$ is a Mersenne prime. (Note that if M_p is prime, then p is also prime. The converse does not necessarily hold.) Only 51 Mersenne primes are known, corresponding to an equal number of (even) perfect numbers. (See this GIMPS page.) It is currently unknown whether there are finitely many such numbers E. It has been conjectured that there are infinitely many even perfect numbers. Note that we have the equations $$1=\gcd\left(M_p,\frac{M_p + 1}{2}\right)=\sigma\left(\frac{M_p + 1}{2}\right)/M_p=\frac{M_p + 1}{\sigma(M_p)}=\frac{d\left(\frac{M_p + 1}{2}\right)}{s(M_p)}=\frac{2s\left(\frac{M_p + 1}{2}\right)}{d(M_p)}$$ so that $$d\left(\frac{M_p + 1}{2}\right)d(M_p)=2s\left(\frac{M_p + 1}{2}\right)s(M_p).$$ (Note that, trivially, we have $(M_p + 1)/2 < M_p$.) Odd Perfect Numbers Much less is known about odd perfect numbers O. We do not know an actual value for even just a single divisor of O, apart from the trivial factor 1. This is because we do not have an example for O, unlike the case for E. (In a letter to Mersenne dated November 15, 1638, Descartes showed that $$\mathscr{D} = {{3}^2}\cdot{{7}^2}\cdot{{11}^2}\cdot{{13}^2}\cdot{22021} = 198585576189$$ would be an odd perfect number if 22021 were prime.) Nonetheless, Euler did prove that O must necessarily have the so-called Eulerian form $$O = q^k n^2$$ where q is the special prime satisfying $q \equiv k \equiv 1 \pmod 4$ and $\gcd(q,n)=1$. Note that we have the equations $$1 < \gcd(n^2,\sigma(n^2))=\frac{\sigma(n^2)}{q^k}=\frac{2n^2}{\sigma(q^k)}=\frac{d(n^2)}{s(q^k)}=\frac{2s(n^2)}{d(q^k)} \tag{*}$$ so that $$d(q^k)d(n^2)=2s(q^k)s(n^2).$$ (It is known that $q^k < n^2$.) Odd Perfect Numbers Must Have a Deficient-Perfect Divisor Remark 1. We may define a deficient-perfect number D to be a positive integer having a proper divisor e satisfying $$\sigma(D) = 2D - e.$$ The divisor e is said to be the deficient divisor of D. Remark 2. It is also easy to show that the deficient divisor e of a deficient-perfect number D is the greatest common divisor of D and $\sigma(D)$. That is, we have $$e=\gcd(D,\sigma(D))$$ if $σ(D) = 2D - e$ and $e \mid D$, $e \neq D$. Holdener and Rachfal (2019) proved the following assertion: THEOREM 1. Every odd perfect number has a deficient-perfect divisor. In particular, if $O = q^k n^2$ is an odd perfect number given in Eulerian form, then the divisor $D_O = q^j n^2$ is deficient-perfect, with deficient divisor $$e_O=\frac{2q^j n^2}{q^{j+1} + 1},$$ where $j=(k-1)/2$. In particular, by Remark 1 and Remark 2, we obtain $$\gcd\left(D_O,\sigma(D_O)\right)=e_O=2D_O-\sigma(D_O)=d(D_O).$$ Following in the footsteps of this short ResearchGate article, titled "Gcd in Odd perfect number" by Devansh Singh (B.Tech, M.E., I.E.T. Lucknow), we proceed as follows: From the Eulerian form $O=q^k n^2$, we get $$O={q^{(k+1)/2}}\cdot\left(q^{(k-1)/2} n^2\right)={q^{(k+1)/2}}\cdot{D_O}.$$ Since $\gcd\left(q^{(k+1)/2},q^{(k-1)/2} n^2\right)=1$ if and only if $k=1$, then we are unable to tie up, in full generality, the value of $$\sigma(O/q^k)/q^{j+1}=\sigma(n^2)/q^{(k+1)/2}$$ with that of $$\gcd(D_O,\sigma(D_O))=\gcd\left(q^{(k-1)/2} n^2,\sigma(q^{(k-1)/2} n^2)\right),$$ by using Singh's approach. This is getting too long already, so let me ask my question (as is) in the title: QUESTION. Does an odd perfect number have a divisor (other than 1) which must necessarily be almost perfect? MY ATTEMPT Note from the definitions that, if a number is almost perfect, then it is automatically deficient-perfect. Suppose to the contrary that the deficiency $d(D_O)$ of the deficient-perfect divisor $D_O=q^j n^2$ of an odd perfect number $O=q^k n^2$ satisfies $d(D_O)=1$ (where $j=(k-1)/2$). (Notice that D_O is a square.) In other words, assume that D_O is almost perfect. Then we have, by a criterion in this paper, the following inequality $$\frac{2D_O}{D_O + 1}< I(D_O) < \frac{2D_O + 1}{D_O + 1}$$ since $D_O > 1$ and D_O is almost perfect. Using the fact that the abundancy index function I is multiplicative, we obtain $$I(D_O)=I(q^j n^2)=I(q^j)I(n^2).$$ If $k=1$, then using the equations and inequality in $(*)$, we obtain $$1<\gcd(n^2,\sigma(n^2))=d(n^2),$$ which contradicts our earlier assumption that $d(D_O)=1$. Thus, it suffices to consider $k \neq 1$. Since $k \equiv 1 \pmod 4$ holds, then $k \geq 5$. It follows that $j=(k-1)/2 \geq 2$. In particular, we have $$I\left((qn)^2\right) = I(q^2)I(n^2) \leq I(D_O) < \frac{2q^j n^2 + 1}{q^j n^2 + 1} \tag{1}$$ and $$\frac{2q^j n^2}{q^j n^2 + 1} < I(D_O) = I(q^j)I(n^2) = \left(\frac{q^{j+1} - 1}{q^j (q - 1)}\right)\cdot{I(n^2)} < \left(\frac{q}{q - 1}\right)\cdot\left(\frac{2q}{q + 1}\right) = \frac{2q^2}{q^2 - 1}. \tag{2}$$ Inequality (2) results to $$0 < 2\left(q^j n^2 + q^2\right)$$ which is trivial. On the other hand, we derive the following estimate from Inequality (1): $$I(q^2)I(n^2)=\left(\frac{\sigma(q^2)}{q^2}\right)\cdot{I(n^2)}=\left(\frac{q^2 + q + 1}{q^2}\right)\cdot{I(n^2)}$$ $$>\left(\frac{q^2 + q + 1}{q^2}\right)\cdot\left(\frac{2(q-1)}{q} + \frac{1}{qn^2}\right).$$ We now test whether the inequality $$\left(\frac{q^2 + q + 1}{q^2}\right)\cdot\left(\frac{2(q-1)}{q} + \frac{1}{qn^2}\right) < \frac{2q^j n^2 + 1}{q^j n^2 + 1} \tag{3}$$ indeed holds. Using some help from WolframAlpha, Inequality (3) holds when $q < n$. To conclude, $D_O = q^j n^2$ might be an almost perfect divisor of the odd perfect number $O=q^k n^2$, where $j=(k-1)/2$, as we were not able to derive a contradiction from assuming that D_O is indeed almost perfect, and then using our criterion for almost perfect numbers. I will stop here for the time being, as I have got some more work to do.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2023-05-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019273 | MSE-4690996

The Connell Sequence and the Number of Ones in Redheffer Matrices

The divisor matrix $D=(d_{r,s})_{r,s\in\mathbf{N}}$ is defined by $d_{r,s}=1$ if r divides s and 0 otherwise. Raymond Redheffer considered a finite truncation of the divisor matrix. For each natural number n he considered the matrix, denoted R_{n}, obtained from the upper left $n\times n$ submatrix of D by setting each entry in the first column equal to 1. I say R_{n} is even if its number of nonzero entries is even, otherwise I say it is odd. Consider the following table: \begin{array}{ l | l | l } n & \text{No. of nonzero entries of R_{n}} & \text{parity of R_{n}} \\ \hline 1 & 1 & \text{odd} \\ 2 & 4 & \text{even} \\ 3 & 7 & \text{odd} \\ 4 & 11 & \text{odd} \\ 5 & 14 & \text{even} \\ 6 & 19 & \text{odd} \\ 7 & 22 & \text{even} \\ 8 & 27 & \text{odd} \\ 9 & 31 & \text{odd} \\ 10 & 36 & \text{even} \\ 11 & 39 & \text{odd} \\ 12 & 46 & \text{even} \\ 13 & 49 & \text{odd} \\ 14 & 54 & \text{even} \\ 15 & 59 & \text{odd} \\ \end{array} The values n for which the matrices R_{n}, are even in parity are the numbers $2,5,7,10,12,14,\ldots$ Similarly the values n for which the matrices R_{n} are odd in parity are the numbers $1,3,4,6,8,9,11,13,15,\ldots$ Writing in 1959 to the journal American Mathematical Monthly Ian Connell challenged readers to find a closed form to an unusual sequence. The sequence, which now bears his name, is made as follows: begin by writing the first odd number, 1, then write the next two even numbers after 1: $2, 4$ and then write the next three odd numbers after 4: $5, 7, 9$ and then write the next four even numbers after 9: $10, 12, 14, 16$, and so on. The first few terms are $1, 2, 4, 5, 7, 9, 10, 12, 14, 16, 17,\ldots$ A solution to Connell's challenge appeared in the same journal in 1960, where it was shown the n-th term of the Connell sequence, a_{n}, has the closed form $2n - \lfloor1/2(1 + \sqrt{8n - 7})\rfloor$. The sequences is found in the Online Encyclopedia of Integer Sequences (OEIS) by looking up the entry A001614. A modification to the Connell Sequence can be made by replacing the first term of the Connell sequence with the even number 0 and then proceed by writing down runs of n consecutive numbers of a given parity. In particular begin by writing the first even number 0, then write the next two odd numbers after 0: $1, 3$ then write the next three even numbers after 3: $4, 6 ,8$ and write the next four odd numbers after 8: $9,11,13, 15$, and so on. The first few terms are $0,1,3,4,6,8,9,11,13,15,\ldots$ The modified Connell sequence is found in the OEIS by looking up the entry A133280. There one finds a closed formula for the n-th term, a_{n}, namely $1+2n-\lfloor 1/2+\sqrt{2n + 2}\rfloor$. I say a number is a Connell number if it belongs to the Connell sequence. I say a number is a modified Connell number if it belongs to the modified Connell sequence. The Connell sequence and their modifications span the natural numbers and share the perfect squares between them. It is straightforward to show that the average order of the Connell numbers and their modification is $2,$ that is $\lim_{n\to\infty}a_{n}/n=2.$ Here is the Question: Are the following items true? R_{n} is even if and only if n is a Connell number that is not a perfect square. R_{n} is odd if and only if n is a modified Connell number. For example $n=19$ is a Connell number that is not a perfect square, and so we should expect R_{19} to have an even number on nonzero entries. Indeed, the number of nonzero entries in R_{19} is 78. Similarly, if $n=36$, a modified Connell number then we should expect R_{36} to have an odd number of nonzero entries. Sure enough, R_{36} has 175 nonzero entries.

论坛问题帖 | has_answers | Anthony | 2023-05-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019274 | MSE-4696409

Conjecture about expected distance between two points in a regular polygon

Point P is a uniformly random point on the perimeter of a regular polygon. Point C is a fixed point somewhere on the circle inscribed in the polygon. Is the following conjecture true: The expected distance between C and P is maximized when C is where the circle touches the polygon. Numerical investigation shows that the conjecture is true for a square. Suppose the square has vertices $(\pm1,\pm1)$. C has coordinates $(\cos\theta, \sin\theta)$. $E(PC)=\frac18\left(\int_{-1}^1 \sqrt{(\cos\theta-x)^2+(\sin\theta+1)^2}dx+\int_{-1}^1 \sqrt{(\cos\theta-1)^2+(\sin\theta-y)^2}dy+\int_{-1}^1 \sqrt{(\cos\theta-x)^2+(\sin\theta-1)^2}dx+\int_{-1}^1 \sqrt{(\cos\theta+1)^2+(\sin\theta-y)^2}dy\right)$ Here is the graph of $E(PC)$ against θ for $0\le \theta \le \pi/2$. The graph shows that $E(PC)$ is maximized when $\theta=0$, i.e. when C is where the circle touches the square. Is my conjecture true for all regular polygons?

论坛问题帖 | has_answers | Dan | 2023-05-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019275 | MSE-4698455

Reference Request - Online copy of R. Graham's "On Finite Sums of Rational Numbers"

Does anyone know where I could find a copy of Ron Graham's PhD dissertation, titled "On Finite Sums of Rational Numbers"? I found this paper, with a similar title, but I could not find Professor Graham's dissertation. I need to double-check some detail(s) in Professor Graham's dissertation which may not be in the published version. This is in consideration of my (current) approach to odd perfect numbers via Egyptian fractions.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2023-05-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019276 | MSE-4698554

Who discovered the largest known 3-perfect number in 1643?

Multiperfect numbers probably need no introduction. (These numbers are defined in Wikipedia and MathWorld.) I need the answer to the following question as additional context for a research article that I am currently writing: Who discovered (in the year 1643) the largest known 3-perfect number to date? Here are the six known 3-perfect numbers, with the last being discovered more than 360 years ago, in 1643: \begin{align*} 120 &= 2^3 \times 3 \times 5, \\ 672 &= 2^5 \times 3 \times 7, \\ 523776 &= 2^9 \times 3 \times 11 \times 31, \\ 459818240 &= 2^8 \times 5 \times 7 \times 19 \times 37 \times 73, \\ 1476304896 &= 2^{13} \times 3 \times 11 \times 43 \times 127, \\ 51001180160 &= 2^{14} \times 5 \times 7 \times 19 \times 31 \times 151. \\ \end{align*} Note that N is an odd perfect number if and only if $2N$ is a (necessarily) even 3-perfect number. (My profuse thanks to Professor V. M. Misyakov for pointing out my mistake in ResearchGate.)

This link asserts that it was René Descartes who discovered the following multiperfect numbers: \begin{array}{|c|c|c|c|} \hline \text{index} &\text{smallest} &\text{name} & \text{ found by } \\ \hline 4 &30240 &4-\text{perfect} &\text{Descartes (1638)} \\ \hline 5 & 14182439040 & 5-\text{perfect} &\text{Descartes (1638)} \\ \hline \end{array} Perhaps \begin{align*} 51001180160 &= 2^{14} \times 5 \times 7 \times 19 \times 31 \times 151 \end{align*} was also discovered by Descartes in 1643, judging from the timing. Did he?

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2023-05-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019277 | MSE-4699163

When is $\frac{5^n - 1}{4}$ squarefree?

While researching the topic of odd perfect numbers, I came across the following related subproblem: PROBLEM: Determine congruence conditions (on $n > 1$) such that $$\frac{5^n - 1}{4}$$ is squarefree. MY ATTEMPT Set $$m_n := \frac{5^n - 1}{4}.$$ I noticed that $m = m_1 = 1$ is squarefree for $n = 1$. Next, I considered the prime factorizations of m_n for the first fifteen values (i.e., $2 \leq n \leq 16$): $$\begin{array}{|c|c|c|} \hline \text{Value of } n &\text{Value of } m_n & \text{Prime Factorization of } m_n \\ \hline 2 & 6 & 2 \times 3 \\ \hline 3 & 31 & 31 \\ \hline 4 & 156 & {2}^2 \times 3 \times 13 \\ \hline 5 & 781 & 11 \times 71 \\ \hline 6 & 3906 & 2 \times {3}^2 \times 7 \times 31 \\ \hline 7 & 19531 & 19531 \\ \hline 8 & 97656 & {2^3} \times 3 \times 13 \times 313 \\ \hline 9 & 488281 & 19 \times 31 \times 829 \\ \hline 10 & 2441406 & 2 \times 3 \times 11 \times 71 \times 521 \\ \hline 11 & 12207031 & 12207031 \\ \hline 12 & 61035156 & 2^2 \times 3^2 \times 7 \times 13 \times 31 \times 601 \\ \hline 13 & 305175781 & 305175781 \\ \hline 14 & 1525878906 & 2 \times 3 \times 29 \times 449 \times 19531 \\ \hline 15 & 7629394531 & 11 \times 31 \times 71 \times 181 \times 1741 \\ \hline 16 & 38146972656 & 2^4 \times 3 \times 13 \times 17 \times 313 \times 11489 \\ \hline \end{array}$$ From this initial data sample, I predict the truth of the following conjectures: CONJECTURE 1: If $n \equiv 0 \pmod 6$, then m_n is not squarefree. CONJECTURE 2: If $n \equiv 0 \pmod 5$, then m_n is squarefree. CONJECTURE 3: If $n \equiv 0 \pmod 3$ and n is odd, then m_n is squarefree. OBSERVATION - If m_n is a prime number, then n must also prime. I skimmed through OEIS sequence A005117 and did not find any references to these conjectures. ATTEMPTING TO RESOLVE CONJECTURE 1 I searched for counterexamples to Conjecture 1 using Pari-GP in Sage Cell Server, I did not get any output in the range $2 \leq n \leq 100$: This gives further computational evidence for Conjecture 1. RESOLVING CONJECTURE 2 I searched for counterexamples to Conjecture 2 using Pari-GP in Sage Cell Server, I got the following output in the range $2 \leq n \leq 100$: 20[2, 2; 3, 1; 11, 1; 13, 1; 41, 1; 71, 1; 521, 1; 9161, 1] 30[2, 1; 3, 2; 7, 1; 11, 1; 31, 1; 61, 1; 71, 1; 181, 1; 521, 1; 1741, 1; 7621, 1] 40[2, 3; 3, 1; 11, 1; 13, 1; 41, 1; 71, 1; 241, 1; 313, 1; 521, 1; 9161, 1; 632133361, 1] 55[11, 2; 71, 1; 103511, 1; 511831, 1; 12207031, 1; 65628751, 1; 190295821, 1] 60[2, 2; 3, 2; 7, 1; 11, 1; 13, 1; 31, 1; 41, 1; 61, 1; 71, 1; 181, 1; 521, 1; 601, 1; 1741, 1; 2281, 1; 7621, 1; 9161, 1; 69566521, 1] 80[2, 4; 3, 1; 11, 1; 13, 1; 17, 1; 41, 1; 71, 1; 241, 1; 313, 1; 521, 1; 9161, 1; 11489, 1; 25601, 1; 632133361, 1; 9094568478143344O1, 1] 90[2, 1; 3, 3; 7, 1; 11, 1; 19, 1; 31, 1; 61, 1; 71, 1; 181, 1; 521, 1; 829, 1; 1171, 1; 1741, 1; 5167, 1; 7621, 1; 169831, 1; 297315901, 1; 60081451169922001, 1] 100[2, 2; 3, 1; 11, 1; 13, 1; 41, 1; 71, 1; 101, 1; 251, 1; 401, 1; 521, 1; 1901, 1; 9161, 1; 239201, 1; 9384251, 1; 424256201, 1; 50150933101, 1; 89620825374601, 1] This means that Conjecture 2 is false. RESOLVING CONJECTURE 3 I searched for counterexamples to Conjecture 3 using Pari-GP in Sage Cell Server, I got the following output in the range $2 \leq n \leq 100$: 93[31, 2; 1861, 1; 148429, 1; 878851, 1; 625552508473588471, 1; 1729748124632393100247504109291, 1] This means that Conjecture 3 is false. RESOLVING AN OBSERVATION I searched for counterexamples to Conjecture 4 using Pari-GP in Sage Cell Server, I did not get any output in the range $2 \leq n \leq 888$. I did search for the primes n satisfying Conjecture 4, in the range $2 \leq n \leq 888$, here is the output: 3Mat([31, 1]) 7Mat([19531, 1]) 11Mat([12207031, 1]) 13Mat([305175781, 1]) 47Mat([17763568394002504646778106689453l, 1]) 127Mat([14693679385278593849609206715278070972733319459651094018859396328480215743184089660644531, 1]) 149Mat([35032461608120426773093239582247903282006548546912894293926707097244777067146515037165954709053039550781, 1]) 181Mat([81566305849981556583878676365706844446264553225862081846982955693371540557468577840286201585673353520178352482616901397705078l, 1]) 619Mat([11491393399729176129058599637568039590983414433049669131423473249238986267377190768602760303952289003868743764537870943345358977246251614482432860160834726163018294937275891867524154115188155135131886536077893354390566834817748709990440493780386105420979057745074407793143691403688802000506454789703520782599348592727930257167323682704508334081537013209268294956642082599042229570588087445297438560221081615964067168538020324707031, 1]) This means that the numbers $$\frac{5^3 - 1}{4}, \frac{5^7 - 1}{4}, \frac{5^{11} - 1}{4}, \frac{5^{13} - 1}{4},$$ $$\frac{5^{47} - 1}{4}, \frac{5^{127} - 1}{4}, \frac{5^{149} - 1}{4}, \frac{5^{181} - 1}{4}, \frac{5^{619} - 1}{4}$$ are all prime. We end this section with the following prediction: CONJECTURE 4: The divisibility condition $n \mid m_n$ holds when either (a) $5 \nmid n$; or (b) n is an odd prime. Alas, this where I get stuck, as I do not currently know how to prove Conjectures 1 and 4.

论坛问题帖 | has_answers | Jose Arnaldo Bebita Dris | 2023-05-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019278 | MSE-4703716

Conjecture improvement of Bunyakovsky integral inequality $\int_{0}^{I}g(t)dt\leq J$

It's a follow-up of my previous question answered by user Apass.Jack nicely. See this post for motivation and background. Conjecture : Let f be a positive increasing continuous function on $x\in[0,1]$ and g be an integrable decreasing positive continuous function on $x\in(0,\infty)$. Let $$J=\left(\int_{0}^{1}\left(f\left(x\right)\right)^{2}dx\int_{0}^{1}\left(g\left(x\right)\right)^{2}dx-\left(\int_{0}^{1}f\left(x\right)g\left(x\right)dx\right)^{2}\right)\max_{x\in\left[0,1\right]}\left(g\left(x\right)\right)$$ and $$I=\left(\int_{0}^{1}\left(f\left(x\right)\right)^{2}dx\int_{0}^{1}\left(g\left(x\right)\right)^{2}dx-\left(\int_{0}^{1}f\left(x\right)g\left(x\right)dx\right)^{2}\right).$$ Then $$\left(\int_{0}^{I}g\left(t\right)dt\right)\leq J.$$ Now the usual counterexample doesn't works @Apass.jack. Here is one example : Choosing $g(x)=\arccos(x),f(x)=\arcsin(x)$ we have : $$2\frac{\left(1/4(2(2+((1+\pi)(4-(\pi-3)\pi^{2}))^{\frac{1}{2}})+\pi((\pi-3)\pi\arccos(1/4(\pi-3)\pi^{2})-((1+\pi)(4-(\pi-3)\pi^{2}))^{\frac{1}{2}}))\right)}{\pi}< \left(\frac{1}{4}\left(\pi^{2}-8\right)\left(\pi-2\right)-\left(2-\frac{\pi}{2}\right)^{2}\right)$$ $$I=\int_{0}^{1}e^{-x^{2}}dx\int_{0}^{1}\left(1+\frac

{1}{2}e^{-x}\right)dx-\left(\int_{0}^{1}\sqrt{e^{-x^{2}}\left(1+\frac{1}{2}e^{-x}\right)}dx\right)^{2},$$ $$\left|\int_{0}^{I}e^{-t^{2}}dt-I\right|<6\cdot10^{-7}$$ As I have no idea for a proof, here is my question. How to (dis)prove my conjecture?

论坛问题帖 | has_answers | Barackouda | 2023-05-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019279 | MSE-4703905

Conjecture: Every shear transformation of the plane is congruent to a dilation

I conjecture that every shear transformation of the plane is geometrically congruent to an orthogonal dilation of the plane. That is, in Euclidean geometry of the plane, if I shear figure F, I can produce a geometrically congruent figure F' via orthogonal dilation. Is my conjecture correct? If so, what is the orthogonal dilation D congruent to $$\begin{bmatrix}1 & \tan \theta \\ 0 & 1\end{bmatrix}.$$ Since shears have determinant 1, I believe it must be of the form $$\begin{bmatrix}\alpha & 0 \\ 0 & \frac 1 \alpha \end{bmatrix},$$ but can't figure out α as a function of θ (or of λ in the general case above). Likewise, consider - what dilation, along any axes, can produce it? I haven't been able to sketch one. The support for this conjecture is via linear algebra; below are the definitions and support: Definitions Shear transformation means $S: \mathbb R^2 \to \mathbb R^2, S = \begin{bmatrix}1 & \lambda \\ 0 & 1\end{bmatrix}$ or $S = \begin{bmatrix}1 & 0 \\ \lambda & 1\end{bmatrix}$. Congruent is defined: a set of points F is congruent to F' if there is an isometry (preserving distance, angles, and colinearity) that sends $F \to F'$. Note that congruent does not imply matrix similarity: it a geometric property related by isometry. Orthogonal dilation means an arbitrary set of axes is chosen, each orthogonal to the rest, and the plane is scaled by a (perhaps different) constant along each. This is precisely the matrices of the form PDP^{-1} where P is orthogonal and D is diagonal. Conjecture I conjecture that for any shear S there exists an orthogonal dilation D such that for all figures F, $S(F) \cong D(F)$. Support Every real 2x2 matrix can be decomposed into a polar decomposition $A = PU$ where U is orthogonal diagonizable and P is an orthogonal matrix (and therefore an isometry). This implies the conjecture and also something related: Every dilation along non-orthogonal axes is congruent to a dilation along congruent axes. Background: While trying to determine Geometrically, what is unique about the mapping of an orthogonally diagonizable matrix (versus non-orthogonal)? , I came to the conclusion that there is no difference: every diagonalizable transform is geometrically congruent to an orthogonally diagonalizable geometric transform! Further background to this conjecture is at Is every symmetric matrix in $\mathbb R^n$ a (non-uniform) stretch? .

论坛问题帖 | has_answers | SRobertJames | 2023-05-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019280 | MSE-4712273

Conjecture: It is not true that 2 eventually always divides $f(x) = \sum_{d \mid p_{\sqrt{x+1}}\#} (d \mid x^2 - 1)$.

Lemma. Let $(d \mid \cdot) : \Bbb{Z} \to \{0,1\}$ be whether (1) or not (0) $\ d$ divides the input. There exists no $N \in \Bbb{N}$ such that $\forall x \geq N$ we have $$f(x) = \sum_{d \mid p_{\pi(\sqrt{x+1})}\#} (d \mid x^2 - 1) = 0 \pmod 2$$ This is a strange conjecture I've come up with that relates to the twin prime problem. Attempt. Suppose the opposite, then we may augment the summation bound of $\pi(\sqrt{x + 1})$ all the way to $\pi(x-2)$, based upon a sieve-of-Eratosthenes argument. Either way, you'll get $f(x) = 0$. There are a growing number of primes that occur between $\lfloor\sqrt{x + 1}\rfloor + 1 \dots x - 2$. If $g(x) =f(x)$ written with the next prime $q = p_{\pi(\sqrt{x + 1}) + 1}$, we have: $$ 0 = f(x) = g(x) = (1 + (q \mid x^2-1))f(x) = f(x) + (q\mid x^2 - 1) f(x) \\ \ = 0 + \sum_{d \mid p_n\#}(qd\mid x^2 - 1), \text{ where } n \equiv \pi(\sqrt{x + 1}). $$ But we can turn this summation back from whence it came into a product: $$ 0 = g(x) = \prod_{2 \leq p \leq p_n} (1 + (qp \mid x^2 - 1)) \pmod 2 $$ This implies that there exists $p \leq p_n$ such that $(qd \mid x^2-1)$ where $q = p_{\pi(p) + 1}$ is the next prime. Now repeat the procedure, to get: $$ h(x) = (1 + (r|x^2 -1))g(x) = 0 = g(x) + (r\mid x^2 - 1) g(x) \\ \ = \prod_{2 \leq p \leq p_n} (1 + (rqp\mid x^2 - 1)) \pmod 2 $$ which means that there exists $p \leq p_n$ such that $rqp = p_n$ where q, r respectively are the next two primes. Now do this repetitively until you use up all the primes in range $\lfloor \sqrt{x + 1} \rfloor + 1 \dots x -2$. All together, this means: There exists $N \in \Bbb{N}$ such that for all $x \geq N$ we have that: $$ p\cdot \prod_{i= \pi(\sqrt{x + 1}) + 1}^{\pi(x - 2)} p_{i} $$ divides $x^2 - 1$ for some prime $p \leq \sqrt{x + 1}$. Or in terms of primorial, we have that: $$ \exists N \in \Bbb{N}:\forall x \geq N, \exists p \leq \sqrt{x + 1}: $$ $$ p\frac{(x-2)\#}{\sqrt{x + 1}\#} \mid (x^2 -1) $$ To someone experienced with numbers, this seems like an absurdity, but I'm not quite sure how to prove it. The primorial notation is just $(x-2)\#:= p_1 p_2 \cdots p_{\pi(x-2)}$ for example. Proving (the opposite of) this requires estimating $\pi(x-2) - \pi(\sqrt{x + 1})$ perhaps. Or can we argue by divisibility that given $x = N$; $x^2 -1$, $(x + 1)^2 -1$ and $(x + 2)^2 - 1$ and so on can't all share the above property. Any ideas?

论坛问题帖 | has_answers | TotoposDiagramChaseApp | 2023-06-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019281 | MSE-4724177

Is some twin prime average the sum of two twin prime averages, two ways?

Accoring to this question and a linked duplicate, it's been verified empirically up to some number that all twin prime averages greater than six, are the sum of two smaller twin prime averages. I was curious whether or not these formed a tree (with twin prime pairs at the vertices and the sum relation the edges). In order for these to form a tree, for every twin prime average, we would only be able to find one pair of smaller twin prime averages which summed to it. Can you find a counterexample, namely any twin prime average which is a sum of two smaller twin prime averages, more than one way?

论坛问题帖 | has_answers | Robert Frost | 2023-06-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019282 | MSE-4727654

Help me (dis)prove a conjecture about square numbers

Let $S=k*m+p$ be any number where $k,m\in\mathbb{N}$ and p is a prime number. My claim is the following: There exists some nondecreasing function $f:\mathbb{N}\mapsto\mathbb{N}$ with $f(m)\to\infty$ and $\frac{f(m)}{m}\to 0$ for $m\to\infty$, such that for any $m\in\mathbb{N}$ the number of primes that satisfy $p\leq f(m)$ and for at least one value of $k\in\mathbb{N}$, S is a square number that is coprime to m is less than some constant $\Phi\in\mathbb{N}$. I posted this about an hour ago but it was written poorly with many mistakes and I hope that this is a much better way of presentation. My guess is that Φ could be huge, if the claim holds true. Although I think that there is probably a counter example that I am not seeing. If there is not, then I would suspect $f(m)=log(m)$ defined for $m\geq T$ for some $T\in\mathbb{N}$ to be a candidate for this kind of slowly growing function.

论坛问题帖 | no_answers_at_retrieval | user1114342 | 2023-06-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019283 | MSE-4728147

$f(x) + f(x/2) + f(x/3) + ... = x$ and conjecture A : $\pi(x/a) > \frac{\pi(x)}{a}$

Consider $f(x)$: $f(x) + f(x/2) + f(x/3) + f(x/4) + ... = x$ and $\lim_{n \to \infty} \frac{f(n)}{\pi(n)} = 1$? and the edit that contains conjecture A : $$\pi(x/a) > \frac{\pi(x)}{a}$$ Is there a name for conjecture A ? Is there a proof or good argument ? Any references or so ?

论坛问题帖 | has_answers | mick | 2023-06-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019284 | MSE-473187

Approximation to $\pi(x)$ conjecture.

A friend conjectured that $\left[\prod_{k=1}^{a_j <\sqrt{x}} \left(1-\frac{1}{a_k}\right)\right] x$ is usually closer to $\pi(x)$ than $\operatorname{Li}(x)$ is for some (fixed) sequence of integers a_k. Could this be true ?

论坛问题帖 | no_answers_at_retrieval | mick | 2013-08-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019285 | MSE-4732398

Triple integral of the great-circle distance function

Numerical integration suggests that $$\mathcal U=\int_0^\pi\int_0^\pi\int_0^\pi\arccos\left(\cos x\cdot\cos y+\sin x\cdot\sin y\cdot\cos z\right) dx dy dz\stackrel{\small\color{gray}?}{=}\frac{\pi^4}2\tag1$$ (note that the function being integrated represents the great-circle distance$^{[1]}$\!^{[2]}$\!^{[3]}$ in spherical coordinates). How can we prove it? Is it possible to find a closed form for this one? $$\mathcal W=\int_0^{\tfrac\pi2}\!\int_0^\pi\int_0^\pi\arccos\left(\cos x\cdot\cos y+\sin x\cdot\sin y\cdot\cos z\right) dx dy dz\tag2$$

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2023-07-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019286 | MSE-4733782

Can we prove that repdigit-numbers cannot be Carmichael-numbers?

A rep-digit number is of the form $$b\cdot \frac{10^a-1}{9}$$ with an integer b with $1\le b\le 9$ and a positive integer a. In the decimal expansion , digit b occurs a times and no other digit. Conjecture : A rep-digit number cannot be a Carmichael-number I checked upto $a=5\ 000$ ($5\ 000$ digits) without finding a Carmichael-number of this form (in fact , not even a Poulet-number). Can we prove that there actually is none ? We can rule out $b=9$ and b even since a Carmichael number must be odd and squarefree and $b=9$ implies divisor 9.

论坛问题帖 | has_answers | Peter | 2023-07-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019287 | MSE-4734330

What is the radius of the circle $|z - z_0| = \rho |z - z_1|$? A conjecture based on street fighting mathematics.

What is the radius of the circle $|z - z_0| = \rho |z - z_1|$? My conjecture is $$\text{radius} = |z_1 - z_0| \cdot \frac {\rho ^2}{(\rho -1)^2(\rho + 1)^2}$$ which I request help verifying, proving, or correcting. Note: Formulas exist; I'm asking for help with this approach, as described below. Let $d = |z_1 - z_0|$. Using the approach taught in Street Fighting Mathematics, I deduced the following: The radius is obviously a function of d and ρ. It must be undefined when $\rho = 1$. This also implies that ρ is unitless. It must be linear in d, since the radius is in units of linear distance, and d is as well (whereas ρ is unit less). It must be symmetric when switching z_0, z_1, implying $f(\rho) = f(\frac 1 \rho)$. It must approach 0 when $\rho \to 0$ or $\rho \to \infty$. It must approach ∞ when ρ approaches 1 (illustrated by Circle of Appolonius diagrams). Asking What function $f(x) = f(1/x)$ has limits $\lim_{x \to 0} f(x) = 0, \lim_{x \to \infty} f(x) = 0$, and $\lim_{x \to 1} f(x) = \infty$? , I got some candidate functions, took the simplest, realized I needed to square ρ to makes things work (which is intuitive given that distance requires taking a square root, and so e.g. the standard equation of circles is in r^2), and got this candidate: $$\text{radius} = \frac d {\rho^2 + \frac 1 {\rho^2} - 2}\\ = d \cdot \frac {\rho ^2}{(\rho -1)^2(\rho + 1)^2}.$$ Can you verify or correct this? Is there a way I could develop this approach into a proof?

论坛问题帖 | has_answers | SRobertJames | 2023-07-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019288 | MSE-4736928

Conjecture: $\,\lim\limits_{n\to\infty}\int_0^1 (1+|\sin{nx}|)^{-2}\mathrm dx=\frac{4}{3\pi}$

I was playing with integrals, and came up with $$L=\lim_{n\to\infty}\int_0^1 \frac{1}{(1+|\sin (nx)|)^2}dx.$$ Conjecture: $L=\dfrac{4}{3\pi}$ Is my conjecture true? Remarks on numerical investigation: Desmos and Wolfram don't do a good job with numerical investigation of this limit, but we can consider the series $f(n)=\dfrac{1}{n}\sum\limits_{k=1}^n \dfrac{1}{(1+|\sin{k}|)^2}$. $f(10^3)\approx0.999568\left(\frac{4}{3\pi}\right)$ $f(10^6)\approx0.999999635\left(\frac{4}{3\pi}\right)$ $f(10^9)\approx0.999999999807\left(\frac{4}{3\pi}\right)$ This suggests that $\lim\limits_{n\to\infty}f(n)=\frac{4}{3\pi}$. Using Riemann sums, we have $\lim\limits_{n\to\infty}f(n)=L$. My attempt: I tried to use $\,\sin nx = \dfrac{1}{2i}(e^{nxi}-e^{-nxi})\,$ in $\;\displaystyle\int_0^1 \dfrac{1}{(1+|\sin (nx)|)^2}\,\mathrm dx\;,\;\;$ to no avail. I also tried to use complex numbers in the series $f(n)$, as in answers to a question about $\sum_{n=1}^{\infty} \frac{\cos (n)}{n}$, to no avail.

论坛问题帖 | has_answers | Dan | 2023-07-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019289 | MSE-4737962

Conjecture about the mean value of an almost periodic function (the "Mountains of Guilin")

Consider the function $f_{p_j,n}(x)=|\sin (p_1x)+\sin (p_2x)+\dots+\sin (p_nx)|$ where p_j is any sequence such that no two p are rational multiples of each other where the p_j are linearly independent over $\mathbb{Q}$ (so the function is almost periodic). (After reading @mollyerin's answer, I changed the condition on p_j.) For example, here is the graph of $f_{2^{1/j},\color{red}{5}}(x)$: And here is the graph of $f_{2^{1/j},\color{red}{10}}(x)$: I call this kind of function $f_{p_j,n}(x)$, the "Mountains of Guilin". Now consider the mean value of $f_{p_j,n}(x)$. I have evidence that suggests that: For any given value of n, the mean value of $f_{p_j,n}(x)$ is independent of the choice of sequence p_j (as long as the p_j are linearly independent over $\mathbb{Q}$). The mean value of $f_{p_j,\color{red}{2}}(x)$ is $8/\pi^2$. I have not been able to find a general expression for the mean value of $f_{p_j,n}(x)$ in terms of n, but I have the following conjecture: $$\color{red}{\lim_{n\to\infty}\frac{\text{mean value of $f_{p_j,n}(x)$}}{\sqrt n}=\frac{1}{\sqrt \pi}}$$ Is my conjecture true? Numerical evidence for my conjecture: Using desmos, I approximated the mean value of $f_{2^{1/j},n}(x)$ as $M_n=\frac{1}{10^7}\sum\limits_{k=1}^{10^7}|\sin (2^1k)+\sin (2^{1/2}k)+\dots+\sin (2^{1/n}k)|$ Letting $L(n)=\dfrac{M_n}{\sqrt n}$, we have: $L(1)\approx 1.128379\left(\frac{1}{\sqrt \pi}\right)$ $L(2)\approx 1.015898\left(\frac{1}{\sqrt \pi}\right)$ $L(10)\approx 1.006373\left(\frac{1}{\sqrt \pi}\right)$ $L(100)\approx 1.000334\left(\frac{1}{\sqrt \pi}\right)$ This suggests that my conjecture may be true, but I don't know how to prove it. Visual representation of the mean value: If my conjecture is true, then there is a nice way to visually represent the mean value of the function. Draw a circle of area n with its centre on the x-axis. A horizontal line through the top of the circle represents, approximately, the average value of the function.

论坛问题帖 | has_answers | Dan | 2023-07-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019290 | MSE-4738210

Can you beat every Tax Collector game?

There's a one-player math game called Tax Collector, and recently I was intrigued by it. Let me explain the game, and then I'll ask my question. To start, choose a number to be your ceiling - the game will use all numbers from 1 to x, where x is your ceiling. Then, claim a number to add to your score. The Tax Collector then claims all the factors of that number, and adds them to their score. Claimed numbers cannot be taken again. Note that you cannot take a number with no unclaimed factors. This means that you can only take one prime number at most (with a factor of 1), and all the others are guaranteed to go to the Tax Collector. The game ends when you cannot claim any numbers, and all the unclaimed numbers go to the Tax Collector. Whoever has a greater total wins. I started wondering if this game could ever be impossible to win. Instantly, I realized it could be - if your ceiling is 1, then you can't take anything and the Tax Collector wins, 1 - 0. However, because you don't have a chance to make a move, I considered the game with a ceiling of 1 to not count as a real game. So the question moved to 2 and upwards. I was stopped once again by the game with a ceiling of 3, as it also proved impossible to win. Your two options were to pick 2 or 3 - picking 3 was the better choice. Then, you get 3 points, the Tax Collector gets 1 point, and the Tax Collector gets two as well, since it has not factors left. Final score: 3 - 3. However, after this, I tried greater numbers, and the games were always beatable, usually by a rather large margin. So here's my question: After the games with ceilings 1 and 3, are there any ceilings that leave the game impossible to beat, no matter how well you play? I would assume no, since the prime numbers get scarcer as the numbers get larger, and primes are the Tax Collector's main source of income. However, I don't know how to prove that all games to infinity are beatable - and maybe there's an unbeatable one out there! I would really appreciate help with this question. Maybe there's something obvious I'm not seeing, but I feel like this is more complicated than the game is itself.

论坛问题帖 | no_answers_at_retrieval | Mathemagician314 | 2023-07-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019291 | MSE-4738407

Conjecture about determinant of matrix with column j with all numbers not divisible by $j+1$

Consider an $n \times n$ matrix A_n with elements $a_{i,j}$ such that $a_{1,j},a_{2,j},a_{3,j},\ldots$ is the sequence of numbers not divisible by $j+1$ in increasing order starting from 1 (e.g. $1,2,4,5,7,8,10,\ldots$ for $j = 2$). For fun, I have computed the determinant for $n \le 8$ and then conjectured that: $$\lvert A_n \rvert = (-1)^{n+1}$$ Is the conjecture true? How would you prove it?

论坛问题帖 | has_answers | Fabius Wiesner | 2023-07-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019292 | MSE-4738421

Large gaps between small primes

Does there exist a positive integer $n>5$ such that the sum of the two largest primes less than n equals n? If yes, lovely! If not, what is the largest prime gap possible between the two largest primes less than n? The first question could also be rewritten as $2*p - n = d$ where p is the largest prime less than n and d is the difference between p and the second largest prime less than n.

论坛问题帖 | no_answers_at_retrieval | hefe | 2023-07-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019293 | MSE-4739571

Numbers with a unique partition as a sum of two squares

The well known Ramanujan tau function $\tau(n)$ is defined as the nth Fourier coefficient of the modular discriminant $\displaystyle \Delta(q)=q\prod_{m=1}^\infty (1-q^m)^{24} = \sum_{n=1}^\infty \tau(n)\,q^n\tag{1a}$ In the spirit of my previous posts(here (1) and here (2) for example) concerning the parity of certain coefficients of modular forms, I was naturally led to investigate this subsequent conjecture Let us introduce the following convolution identity, see here (3) $\displaystyle\Big(\sum_{n=1}^{\infty} \tau(n) q^n\Big)^2+\sum_{n=1}^{\infty} \tau^2(n) q^{2n}=2\sum_{n=0}^{\infty} \sum_{k=0}^{n} \tau(k+1) \Big(\tau(2n-k+1) q^{2n+2}+\tau(2n-k+2) q^{2n+3}\Big) \tag{1b}$ Think of it as a generalisation of the cauchy product and power sum symmetric polynomials. In generalc consider a sequence of complex terms $a_0, a_1, a_2,...$, then the ff convolution identity holds $\displaystyle\Big(\sum_{n=1}^{\infty} a_{n}\Big)^2+\sum_{n=1}^{\infty} a^2_{n}=2\sum_{n=0}^{\infty} c_{n}$ where $\displaystyle c_{n}=\sum_{k=0}^{n} a_{k+1} \Big(a_{2n-k+1}+a_{2n-k+2}\Big)$ given of course that the sums $\displaystyle\Big(\sum_{n=1}^{\infty} a_{n}\Big)^2$ and $\displaystyle\sum_{n=1}^{\infty} a^2_{n}$ converge. A little bit of experimentation leads one to the ff conjecture The nth coefficient in the partial sum of the series $\displaystyle \sum_{n=0}^{\infty} \sum_{k=0}^{n} \tau(k+1)\Big(\tau(2n-k+1) q^{2n+2}+\tau(2n-k+2) q^{2n+3}\Big)\tag{1c}$ is odd if and only if the power of its last term is $\stackrel{\mathrm{def}}{=}2z$ where $z\equiv 1\pmod{4}$ with a unique partition as a sum of two squares x^2+y^2, a sequence of integers $\color{orange}{1}, \color{orange}{5}, \color{orange}{9}, \color{orange}{13}, \color{orange}{17}, \color{orange}{29}, \color{orange}{37},....$ which can readily be found in this oeis entry $\displaystyle \sum_{n=0}^{\infty} \sum_{k=0}^{n} \tau(k+1)\Big(\tau(2n-k+1) q^{2n+2}+\tau(2n-k+2) q^{2n+3}\Big)=q^{2×\color{orange}{1}}-24q^3+828q^4-7520 q^5+103662q^6-492912q^7+3512352q^8-8147520q^9+25870905q^{2×\color{orange}{5}}+19335800q^{11}-193971120q^{12}+6341175146q^{13}-465787812q^{14}-2153273040q^{15}+12714981120q^{16}-11761115712q^{17}-6692143617q^{2×\color{orange}{9}}+68609297160q^{19}-....$ Question: How do we prove it? Modified conjecture Following Somos' proof, the original conjecture may be modified as follows The nth coefficient of the partial series of $(1c)$ is odd if and only if $r(n)$ the number of integer solutions $n=x^2+y^2$ where $0< x \le y$ is odd since $\displaystyle \sum_{n=0}^{\infty} \sum_{k=0}^{n} \tau(k+1)\Big(\tau(2n-k+1) q^{2n+2}+\tau(2n-k+2) q^{2n+3}\Big) \equiv \sum_{n=1}^\infty r(n)q^n \pmod 2$ By using the RHS of our convolution identity (1b) we can determine every nth coefficient of the LHS with ease. For example let us determine the counterexample(spotted by Somos) to the unmodified stated conjecture $\displaystyle \sum_{k=0}^{424} \tau(k+1)\tau(849-k)q^{850}=-905126639660151664169216877 75q^{850}$

论坛问题帖 | has_answers | Nicco | 2023-07-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019294 | MSE-4743372

Prime numbers in rearrangements of Smarandache numbers

Take the first n integers, and concatenate them to form the n_{th} Smarandache number. Then, I have the following Conjecture For $n>2$, the digits of each Smarandache number can be rearranged in blocks of digits, such that each block forms some prime number. For instance, consider $n=1234$; it can be rearranged as 2341, with blocks 23 and 41, which are both prime numbers. The blocks could also be 2, 3 and 41. It can be expected an increasing number of ways to form blocks as n grows. I have written a Python program to check the validity of the conjecture, but it seems to be too slow from $n=15$ on (I share it below the questions). Questions Has been this already studied somewhere? Can you share some reference? Is the conjecture plausible for big values of n? Can you share a faster way / program to check the validity of the conjecture? Or a possible strategy to prove it. As a side note, and incidentally, the Python program finds mainly blocks that involve "quasi-Smarandache" numbers, where a quasi-Smarandache number can be defined as some integer that looks similar to some Smarandache number but changing the order of some digit, or substracting some. (F.i., 1234657, 12345678109,...). It surprises me that they seem to be more abundant than one could "a priori" expect. Thanks in advance! EDIT As stated at the comments, a second interesting conjecture can be posed: Conjecture 2 For $n>2$, the digits of each Smarandache number can be rearranged in blocks of at least $\lfloor\log(n)\rfloor$ digits, such that each block forms some prime number. I have written another Python program to check the validity of the conjecture 2, but it seems to be too slow from $n=15$ on (I share it below the questions too). Any insight on Conjecture 2,or an improvement on the program developed to check it for higher values of n, would also be welcomed. Python code used for checking the Conjecture Created on Thu Jul 27 14:55:31 2023 @author: juanmoreno import itertools def is_prime(num): if num < 2: return False for i in range(2, int(num ** 0.5) + 1): if num % i == 0: return False return True def check_permutation(perm): blocks = [] current_block = [] for num in perm: current_block.append(num) number = int("".join(str(digit) for digit in current_block)) if is_prime(number): blocks.append(number) current_block = [] return len(current_block) == 0, blocks def permute_first_n_integers(n): for perm in itertools.permutations(range(1, n+1)): is_valid, blocks = check_permutation(perm) if is_valid: return blocks return None n = int(input("Enter the value of n: ")) for i in range(1,n+1): result = permute_first_n_integers(i) if result: print("Example of blocks of prime numbers formed with the digits of integers less than or equal to ",i,":", result) else: print(f"No permutation found for n = ",i) Python code used for checking the Conjecture 2 Created on Wed Aug 2 16:12:58 2023 @author: juanmoreno """

import itertools import math def is_prime(num): if num < 2: return False for i in range(2, int(math.sqrt(num)) + 1): if num % i == 0: return False return True def check_permutation_blocks(perm_digits, log_n): num_blocks = len(perm_digits) // log_n for num_combinations in range(num_blocks, 0, -1): for combination in itertools.combinations(range(log_n * num_blocks), num_combinations): blocks = [perm_digits[i:j] for i, j in zip((0,) + combination, combination + (None,))] if all(len(block) >= log_n and is_prime(int(''.join(block))) for block in blocks): return blocks return None def find_prime_blocks_permutation(n): log_n = int(math.log(n)) permutations = itertools.permutations(str(i) for i in range(1, n+1)) for perm in permutations: perm_digits = ''.join(perm) blocks = check_permutation_blocks(perm_digits, log_n) if blocks: print("Valid permutation:", ''.join(perm)) print("Blocks forming prime numbers:", blocks) return print("No permutation found with blocks forming prime numbers.") if name == "main": n = int(input("Enter the value of n: ")) find_prime_blocks_permutation(n)

论坛问题帖 | no_answers_at_retrieval | Juan Moreno | 2023-07-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019295 | MSE-4745174

Can all primes greater than 5 be decomposed into the difference of a primorial and a prime?

Conjecture: For all primes $p_n≥7$, there is at least one solution to the equation $$ p_n = p_k\# - p_m$$ where $p_k\#$ is any primorial, and p_m is any prime number. Has this been explored before? Can it been proven or disproven? I tried it numerically up to $ p_n =2322869$ Interestingly, the additive version of this, $$ p_n = p_k\# + p_m$$ seems to have solutions for most primes but not all; the lowest of which lacks a solution is $p_n=149$.

论坛问题帖 | has_answers | Daniel Gibson | 2023-07-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019296 | MSE-4745280

If 3 numbers of the form $\frac{1}{q^2+1}$ have a mean of $\frac{1}{2}$, will one of them always be $\frac{1}{2}$?

I recently came onto a set of numbers with an interesting pattern, but I can't prove it always holds true. I'd appreciate any help in figuring it out. Let the set S $$S = \left \{ \frac{1}{q^2+1} , q \in \mathbb{Q^+} \right \}$$ And three elements of that set $a,b,c \in S$ My conjecture is that if their average is equal to $\frac{1}{2}$, then at least one of them must be equal to $\frac{1}{2}$. $$\frac{a+b+c}{3} = \frac{1}{2} \implies (a =\frac{1}{2}) \lor (b=\frac{1}{2}) \lor (c=\frac{1}{2})$$ Do you have any idea on how to prove, or disprove that claim? Edit to provide examples: You can easily generate sets of numbers that verify the conjectured pattern. Let $f(q) = \frac{1}{q^2+1}$ for rational q. All triplets of the form $(f(1), f(q), f(\frac{1}{q}))$ have an average of $\frac{1}{2}$.

论坛问题帖 | has_answers | Pyrofoux | 2023-07-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019297 | MSE-4746555

Computing the number of monomials of a certain polynomial

I have computed the number of monomials of the polynomial: $$\prod_{0 \le k_1 \lt \ldots \lt k_{n-t} \le n-1} \left(\sum_{j=1}^{n-1} x^{k_j} \right)$$ with $t = 1$ for $n=2,3,4,5,6$ and found it equal to $1,5,11,19,29$ respectively. Each factor is a combination of $n-t$ monomials chosen among $1,x,x^2,\ldots,x^{n-1}$. The product is over all possible combinations. For example, with $n=3$ we have: $$P(x) = (1+x)(1+x^2)(x+x^2) = x^5 + 2 x^4 + 2 x^3 + 2 x^2 + x$$ and therefore there are 5 monomials. Looking at OEIS, one could for example conjecture that the number of monomials is: $$n-2+(n-1)^2 = n^2-n-1$$ Is the conjecture true? In general, is it possible to find a formula as a function of n for any given t such that $n \gt t \ge 1$?

论坛问题帖 | has_answers | Fabius Wiesner | 2023-08-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019298 | MSE-4753618

Does "all derivatives vanish" $\iff$ "the rate of change is smaller than any function in P"?

Let $f: \mathbb R \to \mathbb R$ be C^∞. Suppose f is not constant on any interval containing x_0. Are the following two statements equivalent: (i) All derivatives of f at x_0 vanish. (ii) The rate of change of f at x_0 is smaller than any polynomial function. More formally, if $g(h) := f(x_0 + h) - f(x)$, for any polynomial function $P(h)$, for any interval I containing x_0, there exists $x_1 \in I$ such that for all $h < |x_1 - x_0|$, $0 < |g(h)| < |P(h)|$. Background: I am exploring flat functions and conjecture that the "reason" their derivatives vanish despite the function being non-constant is that their rate of change is too small to be captured by any polynomial. I am drawing on asymptotic concepts such as O, o, etc. See also: Is there an intuitive way of visualizing how a function that is flat at some given point can be something other than a constant function?

论坛问题帖 | has_answers | SRobertJames | 2023-08-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019299 | MSE-4757741

How can you prove or disprove that half of all natural numbers can be expressed as the sum of three squares minus the product of their roots?

I do not have much formal mathematical education beyond high school, but I do like watching math YouTube videos. I recently read a comment on a video asking if anyone could prove that there are no positive integer solutions to the equation a2 + b2 + c2 - abc = 1. It took me a while, but I found a proof, and I then used a similar method to prove a more general claim. For any given positive integer n, there exists a set of three positive integers a, b, and c satisfying the equation a2 + b2 + c2 - abc = n if and only if there exists such a set where a2 + b2 ≤ n. For example, if there are no positive integer solutions to the equation a2 + b2 + c2 - abc = 6 where a2 + b2 ≤ 6, then there are no positive integer solutions to that equation at all. My proof is too long to paste into this post, so here is a link to a PDF of it: htt ps://www.dropbox.com/scl/fi/9wsm5b4zsz9xpc0r7p0rk/naturalnumberproblem.pdf?rlkey=0g

wqktvfu9215fskihlt2fvu3&dl=0 Since there are only finitely many pairs of positive integers who squares sum to a value less than or equal to a given number, you only need to check finitely many cases to determine whether there is a solution for a given n. For example, for n = 6, you only need to check {a = 1, b = 1} and {a = 1, b = 2}. Solving for c shows there are no positive integer solutions. After discovering this, I wrote a program to that checked whether there were solutions for each n from 1 to 100. I was hoping to find a pattern in the numbers that have or don't have solutions, but I didn't find much. However, I did notice that about half of the numbers had solutions. That intrigued me, so I increased the range of my search to see if this percentage held. Here are the results: From n = 100 to 999, the percentage of numbers with solutions stays between 49.3% and 52.5%. From n = 100 to 9999, the percentage of numbers with solutions stays between 49.8% and 50.4%. From n = 10000 to 99999, the percentage of numbers with solution stays between 50.1% and 50.7% My conjecture is that this percentage stays around 50% and that the deviations get smaller and smaller over time. I assume there is some way to state this more formally using limits. Does anyone know if what I found has already been discovered and if my conjecture has already proven or disproven? What methods would you even use to do this? As I said, I don't have much formal mathematical education. The math used in the proof linked above is fairly simple. Nothing more 'advanced' than modular arithmetic and proofs by contradiction. My intuition is that proving my conjecture would rely on math I've never even heard of, but maybe there is some (relatively) simple proof out there. I can prove that ~41.6% of all numbers will not have solutions. (I am speaking loosely. I know this needs to be stated more formally with limits.) This is because there are no solutions to the equation when n is congruent to 3 mod 4, 3 mod 9, or 6 mod 9. This was verified by a computer trying every possible solution working from mod 2 to mod 800. Let f(n) count the numbers less than or equal to n that are congruent to 3 mod 4, 3 mod 6, or 3 mod 9. There are no numbers that are congruent to both 3 mod 9 and 6 mod 9. The only numbers that are congruent to both 3 mod 4 and 3 mod 9 are those that are congruent to 3 mod 36, and the only one congruent to both 3 mod 4 and 6 mod 9 are those that congruent to 15 mod 36. Therefore, the limit of f(n)/n as n approaches infinity is equal to 1/4 + 1/9 + 1/9 - 1/36 - 1/36 = 5/12 = ~41.6% One more thing I know (proof in that PDF) is that for all n except n = 2, there are either no positive integer solutions to the equation a2 + b2 + c2 - abc = n or there are infinitely many such solutions. For n = 2, there is only one solution {a = b = c = 1}. I doubt that is relevant to the conjecture, but it does not hurt to mention it. One last thing: I'm tagging this as a number theory question and as a question about Diophantine equations. Let me know if that is incorrect. EDIT #1: I want to add that I looked up the first dozen n with solutions in the OEIS and didn't find anything. Here's a list of all n with solutions up to 100 in case it helps: 2, 4, 5, 8, 10, 13, 14, 17, 18, 20, 22, 25, 26, 28, 29, 32, 34, 37, 38, 40, 41, 44, 45, 49, 50, 52, 53, 54, 58, 61, 62, 64, 65, 68, 70, 72, 73, 74, 76, 77, 80, 82, 85, 88, 89, 90, 92, 94, 97, 98, 100 EDIT #2: My program just finished checking up to n = 1,000,000. From n = 100,000 to 1,000,000 the percentage of numbers with solutions stays between 50.7% and 51.6%. It's getting further from 50%, so I'm starting to doubt my conjecture is true. If anyone wants to run the program themselves, here's a link to txt version of it: https://www.dropbox.com/scl/fi/kb2p277sa6d29ieeedigk/mathprogram.txt?rlkey=sr5su9zny13y75qg0b5k1v0g4&dl=0 It's written in Python. I don't have any formal programming education either. Just started teaching myself Python a few weeks ago, so I apologize if it's confusing or inefficient code. Also, the program will not make much sense unless you've read the section in the linked proof called "A Note on Searching For Satisfactory Triplets." EDIT 3: There was a typo in my post. I said there are no solutions when N is 3 mod 6. I meant to write 6 mod 9. Sorry! EDIT 4: There were also a couple typos in my PDF where I used < when I meant to write ≤, but I have corrected them. (Unfortunately, this somehow created some new typos. It must be an issue with the conversion from .doc to .pdf but I don't think it effects the clarity much.) EDIT 5: Some errors in the proof were fixed, and a new link has been created. This one should be viewable by anyone. EDIT 6: Refined the approximation of how many numbers (less than N for sufficiently large N) don't have solutions from 40% to 41.6%.

论坛问题帖 | has_answers | Justin Teague | 2023-08-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019300 | MSE-4758544

Conjecture about non-squarefree numbers.

Conjecture : Let k be a non-negative integer. Then , there is a positive integer j , such that 10^j+k is NOT squarefree. The conjecture is true upto $k=10^6$. We can always find a positive integer $j\le 421$ as desired. If we demand that 10^j+k is not cubefree, then still up to $k=17493$ L, there is a $j\le 6376$ as desired. I am however currently mainly interested in the non-squarefree case. Another approach (which I have not yet tried) is that we determine the k for which $4\mid 10^j+k$, $9\mid 10^j+k$, $25\mid 10^j+k$ is possible and hope that we can "cover" all positive integers. If someone solves this problem for which we need discrete logarithms, an answer would be appreciated and I would gladly upvote and accept it. The non-cubefree case would be a nice bonus.

论坛问题帖 | has_answers | Peter | 2023-08-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019301 | MSE-4759973

Computing the least common multiple of the numbers 1 through n for incremental n

I notice a pattern when computing the least common multiple for the numbers 1 through n as we incrementally increase n. The pattern is that the list of factors does not increase unless the incremental next value we include is a prime number to a positive integer power. Also, if it does increase, it only ever increases by 1. Consider the following python code. Notice that when we include 13, the factor list begins to include 13 because 13 is a prime raised to the first power. Notice that when we include 16, the factor list begins to include an additional 2 because 16 is a prime raised to the fourth power. >>> factors(lcm(*range(1,13))) [2, 2, 2, 3, 3, 5, 7, 11] >>> factors(lcm(*range(1,14))) [2, 2, 2, 3, 3, 5, 7, 11, 13] >>> factors(lcm(*range(1,15))) [2, 2, 2, 3, 3, 5, 7, 11, 13] >>> factors(lcm(*range(1,16))) [2, 2, 2, 3, 3, 5, 7, 11, 13] >>> factors(lcm(*range(1,17))) [2, 2, 2, 2, 3, 3, 5, 7, 11, 13] >>> factors(lcm(*range(1,18))) [2, 2, 2, 2, 3, 3, 5, 7, 11, 13, 17] >>> factors(lcm(*range(1,19))) [2, 2, 2, 2, 3, 3, 5, 7, 11, 13, 17] >>> factors(lcm(*range(1,20))) [2, 2, 2, 2, 3, 3, 5, 7, 11, 13, 17, 19] Does the conjecture have a name, and is it true? Stated in another way, we can always construct n+1 by rummaging around among the factors of the numbers 1 through n unless n+1 is a prime power. Further, if it is a prime power, the prime factorization of n+1 will always have just one additional factor than our previous budget had (e.g. 2^4 vs 2^3). (for what it's worth, I was motivated to think about this because of a recent post from

Fermat's Library: https://www.linkedin.com/posts/fermatslibrary_heres-a-surprising-appearance-of-e-the-activity-7101876060991950848-D7T-)

论坛问题帖 | no_answers_at_retrieval | efthimio | 2023-08-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019302 | MSE-4761048

Possible connection between prime numbers in binary and π

I just realized I asked a question with a very similar title a while back. This is not a duplicate. It is another conjecture though. Conjecture: $\lim\limits_{x\to\infty}\mu \{ f(2), f(3), f(5), ... , f(x) \}=\frac{\pi}{2}-1$ where $x\in\mathbb{P}$, $f(x)$ equals the amount of 1s divided by the total number of digits in the binary representation of x. μ is the arithmetic mean and $\mathbb{P}$ represents the set of all primes. Example of $f(x)$ because of my poor explanation (I tried my best): $f(5)\rightarrow 101\rightarrow \frac{\text{number of 1s (or the digit sum)}}{\text{total number of digits}}\rightarrow\frac{2}{3}$ I have no idea why this conjecture would be true but it appears that it may converge, at least for the first 2000 primes. In the picture, the green line is $y=\frac{\pi}{2}-1$ and the black points are $\mu \{ f(2), f(3), f(5), ... , f(x) \}$. My question is whether or not my conjecture is true? Am I missing something obvious here? Also, I find it very interesting that it looks similar to a quasiperiodic function.

论坛问题帖 | has_answers | Dylan Levine | 2023-08-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019303 | MSE-4764583

Deterministically finding g such that $f(g(x)) \ge x$?

It's just some random thoughts, but I find it confusing. Here's the classical "prove" that every natural number can be described within eleven words: Suppose not, then there's a smallest counterexample n. Then n can be described as "the smallest natural number that cannot be described within eleven words", which is itself in eleven words, contradiction. Of course, the proof above is false. But when I tried to do the same thing mathematically, something confusing happened. Suppose f is an arbitrary injective function $f: \mathbb{N} \to \mathbb{N}$. Let h be a bijection from "all finite logical formula" to $\mathbb{N}$. Let φ_n denote the formula with $f(h(\varphi_n))=n$. For some $M \in \mathbb{N}$ let $p \in \mathbb{N}$ such that $$\varphi_p(x) = \neg\exists m\in\mathbb{N}(m<M \wedge \varphi_m(x) \wedge \neg\exists y\in\mathbb{N}(\varphi_m(y))) \wedge\\ \forall y\in\mathbb{N}(y<x \to \exists m\in\mathbb{N}(m<M \wedge \varphi_m(y) \wedge \neg\exists z\in\mathbb{N}(\varphi_m(z))))$$ (x is the smallest natural number such that there is no formula φ_m with $m<M$ such that φ_m "accurately describes" x, i.e. x is the only natural number that satisfies φ_m.) Let us first assume that $p < M$. If there is an $y \in \mathbb{N}$ that $\neg\exists m\in\mathbb{N}(m<M \wedge \varphi_m(y))$, let y be the smallest such number. Then $\varphi_p(y)$, but $p < M$, contradiction! Therefore every natural number can be accurately described with some formula φ_m with $m<M$, but the number of such formula is finite, contradiction! So $p \ge M$. What does this mean? We have found, constructively, a function $h^{-1} \circ p: \mathbb{N} \to \mathbb{N}$ injective, such that $f(h(h^{-1}(p(M)))) \ge M$ which is $f(p(M)) \ge M$, for every M. This is very counter-intuitive, since such a p is usually found inductively. Is there something wrong in the above construction?

论坛问题帖 | no_answers_at_retrieval | atzlt | 2023-09-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019304 | MSE-4765979

Conjecture: If matrix M has entries (left to right, then top to botom) $\sin 1,\sin 2,\sin 3,\dots,\sin (n^2)$, where $n\ge 3$, then $\det M = 0$.

According to my calculator, $\det\begin{bmatrix}\sin 1 & \sin 2 & \sin 3 \\ \sin 4 & \sin 5 & \sin 6 \\ \sin 7 & \sin 8 & \sin 9\end{bmatrix}=0$ $\det\begin{bmatrix}\sin 1 & \sin 2 & \sin 3 & \sin 4 \\ \sin 5 & \sin 6 & \sin 7 & \sin 8 \\ \sin 9 & \sin 10 & \sin 11 & \sin 12 \\ \sin 13 & \sin 14 & \sin 15 & \sin 16 \end{bmatrix}=0$ $\det\begin{bmatrix}\sin 1 & \sin 2 & \sin 3 & \sin 4 & \sin 5 \\ \sin 6 & \sin 7 & \sin 8 & \sin 9 & \sin 10 \\ \sin 11 & \sin 12 & \sin 13 & \sin 14 & \sin 15 \\ \sin 16 & \sin 17 & \sin 18 & \sin 19 & \sin 20 \\ \sin 21 & \sin 22 & \sin 23 & \sin 24 & \sin 25\end{bmatrix}=0$ I conjecture that, for $n\ge 3$, $\det \begin{bmatrix} \sin 1 & \sin 2 & \sin 3 & \dots & \sin n \\ \sin (n+1) & \sin (n+2) & \sin (n+3) & \dots & \sin (2n) \\ \sin (2n+1) & \sin (2n+2) & \sin (2n+3) & \dots & \sin(3n) \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \sin ((n-1)n+1) & \sin ((n-1)n+2) & \sin ((n-1)n+3) & \dots & \sin (n^2) \end{bmatrix}=0$ Is my conjecture true? I have only been able to prove the case with $n=3$. $\sin 5 + \sin 7 + [\sin 1 + \sin (-1)] + [\sin 3 + \sin (-3)]$ $=\sin 5 + \sin 7 + [\sin 1 + \sin (-1)] + [\sin 3 + \sin (-3)]$ Rearrange each side: $[\sin (-3) + \sin 5] + [\sin 1 + \sin 3] + [\sin (-1) + \sin 7]$ $=[\sin (-1) + \sin 3] + [\sin (-3) + \sin 7] + [\sin 1 + \sin 5]$ Use the product-to-sum formulas: $(\sin 1)(\cos 4) + (\sin 2)(\cos 1) + (\sin 3)(\cos 4)$ $=(\sin 1)(\cos 2) + (\sin 2)(\cos 5) + (\sin 3)(\cos 2)$ Subtract $(\sin 1)(\cos 14)+(\sin 2)(\cos 13)+(\sin 3)(\cos 12)$ from both sides: $(\sin 1)(\cos 4 - \cos 14) + (\sin 2)(\cos 1 - \cos 13) + (\sin 3)(\cos 4 - \cos 12)$ $=(\sin 1)(\cos 2 - \cos 14) + (\sin 2)(\cos 5 - \cos 13) + (\sin 3)(\cos 2 - \cos 12)$ Use the product-to-sum formulas again: $(\sin 1)(\sin 5)(\sin 9)+(\sin 2)(\sin 6)(\sin 7)+(\sin 3)(\sin 4)(\sin 8)$ $=(\sin 1)(\sin 6)(\sin 8)+(\sin 2)(\sin 4)(\sin 9)+(\sin 3)(\sin 5)(\sin 7)$ which is equivalent to $\det\begin{bmatrix}\sin 1 & \sin 2 & \sin 3 \\ \sin 4 & \sin 5 & \sin 6 \\ \sin 7 & \sin 8 & \sin 9\end{bmatrix}=0$

论坛问题帖 | has_answers | Dan | 2023-09-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019305 | MSE-4774206

A conjecture for the golden ration via continued fraction $\phi=\frac{1+\sqrt{5}}{2}$

Playing with my own question Got a factored version of the Taylor's series? I found that : Define : $$f(x)=\frac{1}{\sqrt{x-2/x}}$$ Then it seems we have : $$\phi=f(f(\cdots f(\phi)\cdots))$$ Numerical checking is agree so some question : edit : I'm really interested by an generalization the next polynomial are : $$x^4-2x^3+2x,x^5-3x^3+2x,...$$ How to (dis)prove it ? How fast is the convergence if true ? Is there a generalization if true ?

论坛问题帖 | has_answers | Barackouda | 2023-09-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019306 | MSE-4781942

Conjecture: If $a_1\in\mathbb{Z}$ and $a_n=\sec (a_{n-1})$ then the proportion of positive terms approaches $\frac{1}{\sqrt{2\pi}}$ as $n\to\infty$.

I made up the following family of sequences: $a_1\in\mathbb{Z}$ and $a_n=\sec (a_{n-1})$ for $n>1$. I conjecture that, for any $a_1\in\mathbb{Z}$, the proportion of positive terms approaches $\frac{1}{\sqrt {2\pi}}$ as $n\to\infty$. Is my conjecture true? Evidence for my conjecture For $a_1=1$, here is the graph of a_n against n for $1\le n \le 25$. Using Excel, it seems that, the proportion of positive terms approaches a limit as $n\to\infty$, and the limit is independent of the value of a_1. I found that, for $a_1=1, 2, 3, \dots, 25$, among the first 10000 terms, the average proportion of positive terms is 0.398940, which is very close to $\frac{1}{\sqrt {2\pi}}\approx 0.398942$. Another look at the sequence Here are the graphs of $y=\sec x$ and $y=x$. We start at $(1,\sec 1)$, then go horizontally to the line $y=x$, then go vertically to the curve $y=\sec x$, then go horizontally to the line $y=x$, etc. Among the points that we meet on the curve $y=\sec x$, about 0.39894 of them are above the x-axis.

论坛问题帖 | no_answers_at_retrieval | Dan | 2023-10-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019307 | MSE-4787606

How do I find the focal distance?

I’ll first put the question as it is on the assignment and then ask my question. The following describes certain characteristics of a parabolic radio telescope that an astronomical observatory would like to build. It is presented in two positions: pointing upward and to the horizon. In this first position, the focus reaches a height of 18.4 m and the edge of the reflector is situated 14.5 m from the floor. In the second position, the bottom of the reflector is 2 m from the floor and the top is at 18 m. You should know that the efficiency of a radio telescope depends on the diameter of its reflector. The greater this diameter, the more the radio telescope is able to capture waves coming from remote regions of space. However, the capture quality also depends on the focal distance, which is the distance between the focus and the vertex. This distance must neither be too great nor too small relative to the diameter. In fact, what is important is the ratio of $\frac{focaldistance}{diameter}$, which ideally should be close to 0.5. Then they go on asking, a) Determine the value of this ratio for the radio telescope presented. b) To increase the value of the ratio without reducing the diameter of the reflector, what changes should be made to the shape of the parabola? State a conjecture justifying your point of view, using an example or reasoning. So my question is: How do I find the focal distance of the mentioned parabola? I tried to use the formulas in the book but I was not able to find it. I only found the diameter (18 m - 2 m = 16 m). I have a feeling that the answer is right in front of me but I don’t see it.

论坛问题帖 | has_answers | Shakir | 2023-10-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019308 | MSE-4790960

Conjecture for Integrals of the Form $\int_{0}^{1}\frac{\ln^n\left(1-x^{2}\right)}{1+x}dx$

Recently I got interested in Logarithmic Integrals from this $\int_{0}^{1}\frac{\ln^4\left(1-x^{2}\right)}{1+x}dx$ and sought out to find Higher Power variations. Let, $$I_n=\int_{0}^{1}\frac{\ln^n\left(1-x^{2}\right)}{1+x}dx$$ Consider the following expression : $$E=\zeta(\alpha_1)\zeta(\alpha_2)...\zeta(\alpha_n)\ln^\alpha2$$ whose weight is $\alpha_1+\alpha_2+...+\alpha_n+\alpha.$ $$\boxed{\text{Conjecture: All I_n can be written as a Linear Combination involving Expression E of weight $n+1$.}}$$ Here are the first few examples : $$I_1=-\frac{1}{2}\zeta\left(2\right)+\ln^{2}2$$ $$I_2=2\zeta\left(3\right)-2\zeta\left(2\right)\ln 2+\frac{4}{3}\ln^32$$ $$I_3=-\frac{27}{4}\zeta\left(4\right)+12\zeta\left(3\right)\ln2-6\zeta\left(2\right)\ln^22+2\ln^42$$ $$I_4=72ζ\left(5\right)-24ζ\left(3\right)ζ\left(2\right)-54ζ\left(4\right)\ln2+48ζ\left(3\right)\ln^{2}2-16ζ\left(2\right)\ln^{3}2+\frac{16}{5}\ln^{5}2$$ $$I_5=-\frac{1185}{4}ζ\left(6\right)+120ζ^{2}\left(3\right)+720ζ\left(5\right)\ln2-240ζ\left(3\right)ζ\left(2\right)\ln2-270ζ\left(4\right)\ln^{2}2+160ζ\left(3\right)\ln^{3}2-40ζ\left(2\right)\ln^{4}2+\frac{16}{3}\ln^{6}2$$ $$I_6=6480ζ\left(7\right)-2160ζ\left(5\right)ζ\left(2\right)-1620ζ\left(4\right)ζ\left(3\right)-3555ζ\left(6\right)\ln2+1440ζ^{2}\left(3\right)\ln2+4320ζ\left(5\right)\ln^{2}2-1440ζ\left(3\right)ζ\left(2\right)\ln^{2}2-1080ζ\left(4\right)\ln^{3}2+480ζ\left(3\right)\ln^{4}2-96ζ\left(2\right)\ln^{5}2+\frac{64}{7}\ln^{7}2$$ Could the Conjecture be Proved or Disproved. I had made a Similar Conjecture for another Logarithmic Integral of Similar Form but it stopped working from Weight 8.

论坛问题帖 | has_answers | Miracle Invoker | 2023-10-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019309 | MSE-4791750

Are there any general results that relate the amount of solutions to a diophantine equations to the average growth rate of such solutions?

Background Recently, I asked a question about the average growth rate of the solutions to the following sequence of Egyptian fraction equation: $$ \sum_{k=1}^{n} \frac{1}{x_{k}} = q. \tag{1} $$ Here, we have $x_i \in \mathbb{N}_{>0} $ for all $ 1 \leq i \leq n $, and $ \ 0 <x_{1}<x_{2}<\cdots<x_{n} $. So far, the question hasn't received any answers. What is known, however, are bounds for the amount of solutions to (1). If we define $$\mathsf{E}_{n}(q)=\left\{ \{x_1 , x_2 , \dots , x_n \} \in \mathbb{N}^n_{>1}: \sum_{k=1}^{n} \frac{1}{x_{k}} = q, \ 0 <x_{1}<x_{2}<\cdots<x_{n} \ \right\}, $$ then according to p.8 of this article we have $$ e^{c \frac{n^{3}}{\log(n)}} < | \mathsf{E}_{n}(1) | < c_{0}^{(1+\epsilon)2^{n-1} }, \tag{2} $$ where $c_{0} \approx 1.264$. Question Let's leave aside the specific case of Egyptian fractions (as described above) for a moment, and consider a more general case. Let $a(n)$ be a set consisting of sequences of solutions to a sequence of diophantine equations $d(n)$, with $|a(n)| = o(f(n))$ and $|a(n)| = O (g(n))$. Can anything be said, in this general case, about the growth rate of the average size of the solutions to $d(n)$, and how this growth rate relates to $f(\cdot)$ and $g(\cdot)$? Are there any results on this question, or any conjectures?

论坛问题帖 | no_answers_at_retrieval | Max Lonysa Muller | 2023-10-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019310 | MSE-4800031

Who can prove or disprove this conjecture?

Let m be a positive integer such that $p:=8m^2+1$ is prime. Conjecture : We always have $$2^{2m^2}\equiv 1\pmod p$$ I could only establish $$2^{4m^2}\equiv 1\pmod p$$ following from Euler's criterion , but I could not rule out $2^{2m^2}\equiv -1\pmod p$ Motivation : A proof would show that $$(2m^2+1)(8m^2+1)$$ is a Poulet-number whenever both factors are prime. Verification : The conjecture is true upto $m=10^{10}$.

论坛问题帖 | has_answers | Peter | 2023-11-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019311 | MSE-4800888

Ec primes multiple of 67 and $1\pmod{67}$

Ec primes are primes of the form: $(2^n-1)\cdot 10^d+2^{n-1}-1=ec(n)$, where d Is the Number of decimal digits of 2^{n-1}. I conjecture that there are infinitely many primes of this form with n either $\equiv 0\pmod {67}$ or $\equiv 1\pmod {67}$ Verification: for $n=67, 6231,51456, 79798, 285019$ $ec(n)$ is probable prime. Is that another coincidence or law of small numbers? The multiples of 67, that Is 6231 and 51456 are also multiple of 3. And 79797, 285018 are divisible by 3 too. Is there a particular reason? $79797=201\cdot 397$ Is a palindromic Number Just a curio.

论坛问题帖 | no_answers_at_retrieval | user1225681 | 2023-11-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019312 | MSE-4801272

Does it make sense to divide mathematical theorems into those susceptible to experimental conjecture and those that are not?

I read the following in a good book: ”let's start from zero", the authors are Vinicio Villani and Maurizio Berni (pisan mathematicians) but I don't know if the book is also marketed outside Italy. I don't think I'm exaggerating when I say that the shocking discovery of the existence of incommensurable segments gave a decisive impulse to the development of all mathematics, understood as a theoretical, hypothetical-deductive discipline. In fact, the validity of other theorems of classical geometry (starting from those of Thales and Pythagoras) can be empirically verified on a drawing through experimental measurements, albeit approximate. Instead, the question of the commensurability or incommensurability of two segments eluded us then, as it eludes us today and will elude us forever to every attempt at experimental verification, given the inevitable imprecision of every physical measurement" When I first read it, in my mind I enthusiastically endorsed this statement. But then I thought about it and I had some doubts. The statement seemed to imply that there are two classes of theorems, the one whose truthfulness can be suggested by so-called experimental and numerical arguments (even if obviously without a demonstration we are always at zero), and those which are totally precluded by a such approach that can suggest conjectures. But does this make sense? Really "the theorem the root of 2 cannot be expressed as a ratio between integers" is in this sense qualitatively different from the Pythagorean theorem? It is true that I can set about constructing squares on right-angled triangles and make measurements that lead me to conjecture the validity of the theorem, but I can also patiently set about searching for two integers such that the square of their ratio is two, and after long labors conjecture that such a pair of integers does not exist. All this if I didn't have an easy proof available but it's not the point here now. In short, the question is: does the division of mathematical theorems into two classes, those that can be conjectured experimentally and those that cannot, make sense?

论坛问题帖 | has_answers | Fausto Vezzaro | 2023-11-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019313 | MSE-4804630

A curious coincidence: $\prod\limits_{k=1}^\infty\left(1+\int_{k}^{k+1}\left(\frac{\sin (\pi x)}{x}\right)^2\mathrm dx\right)\overset{?}{=}\pi/2$

It is known that $\int_0^\infty \frac{\sin x}{x} \mathrm dx=\dfrac{\pi}{2}$ (proof) and $\int_0^\infty\left(\frac{\sin x}{x}\right)^2\mathrm dx=\dfrac{\pi}{2}$ (proof). Numerical investigation suggests that we also have: $$\prod\limits_{k=1}^\infty\left(1+\int_{k}^{k+1}\left(\frac{\sin (\pi x)}{x}\right)^2\mathrm dx\right)=\dfrac{\pi}{2}$$ But I do not know how to prove this. $\int\left(\frac{\sin (\pi x)}{x}\right)^2\mathrm dx$ cannot be expressed in terms of elementary functions. Is this infinite product really equal to $\dfrac{\pi}{2}$?

论坛问题帖 | has_answers | Dan | 2023-11-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019314 | MSE-480904

Weaker Version of "Goldbach's Other Conjecture"

Taken from problem 46 on Project Euler: It was proposed by Christian Goldbach that every odd composite number can be written as the sum of a prime and twice a square. $9 = 7 + 2 \times 1^2$ $15 = 7 + 2 \times 2^2$ $21 = 3 + 2 \times 3^2$ ... It turns out that the conjecture was false. Now, suppose the phrase "twice a square" was replaced with "two squares." In other words, this revision of the conjecture would state that all odd composites C can be written in the following form: $C=p+a^2+b^2$ Where p is a prime and a and b can take on any non-negative integer values. Making the substitution $a=b$ reduces this problem to the classical conjecture. Now, is this weaker conjecture true? My Attempt at the Problem: Let $d=a^2+b^2$. A well-known theorem states that the sum of two squares can be rewritten as $2^{e_0} \left({p_1}^{e_1}...{p_n}^{e_n} \right) \left({q_1}^{2 o_1}...{q_n}^{2 o_n} \right)$, where p_i is a prime of the form $4k+1$ and q_i is a prime of the form $4k+3$. We can therefore write the conjecture as follows: $C=p+d$ $C=p+ 2^{e_0} \left({p_1}^{e_1}...{p_n}^{e_n} \right) \left({q_1}^{2 o_1}...{q_n}^{2 o_n} \right)$ $C=p+ 2^{e_0}g$ Where g is any integer such that $g=1 \left(\text{mod } 4 \right)$. Letting $p=2$ and $e_0=0$ shows us that the conjecture is true when C is of the form $4k+3$. From here I am not sure where to go. Any clues?

论坛问题帖 | has_answers | Ryan | 2013-08-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019315 | MSE-4810043

Conjecture: Given any $2n+1$ points, we can always draw n non-intersecting circles whose diameter endpoints are $2n$ of those points.

Is the following conjecture true or false: Given any $2n+1$ coplanar points, we can always draw n non-intersecting circles coplanar with the points, whose diameter endpoints are $2n$ of those points. We consider tangent circles to be non-intersecting. This is a generalization of my earlier question about the case $n=2$. That question is currently unanswered. I wonder if it is easier to prove the general case. Example: Another example: I cannot find a counter-example, nor can I prove the conjecture. I made a generator of seven (pseudo)random points. (If, instead, we were given $2n$ points, the conjecture would not be true: for example, if the $2n$ points were the vertices of a regular $(2n-1)$-gon plus the centre, then we could not draw n non-intersecting circles.)

论坛问题帖 | has_answers | Dan | 2023-11-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019316 | MSE-4810220

Poulet numbers of the form $3p$, where p is a palindrome

A Poulet-number is a composite number N satisfying $$2^{N-1}\equiv 1\mod N$$ A palindrome is a positive integer with a digit string in base 10 which remains the same if it is written down backwards. If $3p$ is a Poulet-number and p a palindrome , it seems that p must begin (and therefore end) with digit 7. The examples upto $p=10^{18}$ are : 23001 7667 [11, 1; 17, 1; 41, 1] 212421 70807 [11, 1; 41, 1; 157, 1] 23054601 7684867 [17, 1; 251, 1; 1801, 1] 235201461 78400487 [11, 1; 41, 1; 131, 1; 1327, 1] Why is p always of the form $7...7$? Why is p divisible by 41 , if it is divisible by 11 ? Any counter-example?

论坛问题帖 | no_answers_at_retrieval | user1250659 | 2023-11-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019317 | MSE-4810737

"Twin binomial coefficient conjecture": There are no binomial coefficients that differ by 2, excluding $\binom{n}{1}$ and $\binom{n}{n-1}$.

Consider the binomial coefficients, excluding those of the form $\binom{n}{1}$ and $\binom{n}{n-1}$. There are some that differ by 1, for example $\binom{7}{2}-\binom{6}{3}=21-20=1$. There are some that differ by 3, for example $\binom{31}{2}-\binom{11}{5}=465-462=3$. There are some that differ by 4, for example $\binom{5}{2}-\binom{4}{2}=10-6=4$. There are some that differ by 5, for example $\binom{6}{2}-\binom{5}{2}=15-10=5$. There are some that differ by 6, for example $\binom{9}{3}-\binom{13}{2}=84-78=6$. Etc. But I cannot find any that differ by 2. I checked the first 120 rows of Pascal's triangle. So, is the following conjecture true or false: There are no binomial coefficients that differ by 2, excluding those of the form $\binom{n}{1}$ and $\binom{n}{n-1}$. Call it the "Twin binomial coefficient conjecture", if you will. Context: I was playing with Pascal's triangle, trying to approximate the median of the numbers in the first n rows, and I noticed a lack of differences of 2.

论坛问题帖 | has_answers | Dan | 2023-11-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019318 | MSE-4812171

$\int_0^{\pi/2}\int_0^{\pi/2}\frac{(\tan\alpha)(\tan\beta)}{\tan\alpha+\tan\beta} d\alpha d\beta=(0.9999999913...)(\pi/2)$? Seriously?

In the diagram, α and β are independent uniformly random real numbers in $\left(0,\frac{\pi}{2}\right)$. What is $\mathbb{E}(h)$? Superimposing a cartesian coordinate system, the equations of the lines are $y=(\tan\alpha)x$ and $y=(-\tan\beta)(x-1)$, so the y-coordinate of their intersection is $\frac{(\tan\alpha)(\tan\beta)}{\tan\alpha+\tan\beta}$. So we have $$\mathbb{E}(h)=\frac{4}{\pi^2}\int_0^{\pi/2}\int_0^{\pi/2}\frac{(\tan\alpha)(\tan\beta)}{\tan\alpha+\tan\beta} d\alpha d\beta$$ Desmos says $I=\int_0^{\pi/2}\int_0^{\pi/2}\frac{(\tan\alpha)(\tan\beta)}{\tan\alpha+\tan\beta} d\alpha d\beta$ is $(0.9999999913...)(\frac{\pi}{2})$. Is that a computer error, and the result is exactly $\frac{\pi}{2}$? I don't know how to evaluate the integral. Wolfram evaluates the inside integral but doesn't evaluate both integrals. If Desmos is not very reliable, then maybe my earlier weird conjecture is actually true. (This question was inspired by a question about random points in a square.) Edit: In the comments, @G.Gare notes that Mathematica says the integral I is exactly $\pi/2$. Can we prove it? Maybe an intuitive geometrical argument? Edit2: I seek to generalize this result here.

论坛问题帖 | has_answers | Dan | 2023-11-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019319 | MSE-4812776

Conjecture: If x_k are random in $(0,\pi/2)$ then expectation of $\frac{\prod_{k=1}^n\tan x_k}{\sum_{k=1}^n\tan x_k}$ is $(\pi/2)^{2n-6}$ for $n>2$.

Let $E(n)=\text{expectation of }\dfrac{\prod_{k=1}^n\tan x_k}{\sum_{k=1}^n\tan x_k}$ where x_k are independent uniformly random real numbers in $\left(0,\frac{\pi}{2}\right)$. Is the following conjecture true: $E(n)=\left(\frac{\pi}{2}\right)^{2n-6}$ for $n>2$. Context Earlier I found that $E(2)=\left(\frac{2}{\pi}\right)^2\int_0^{\pi/2}\int_0^{\pi/2}\frac{(\tan x_1)(\tan x_2)}{\tan x_1+\tan x_2}dx_1dx_2=\frac{2}{\pi}$. Naturally, I wondered if $E(n)$ has closed forms for other n values. Basis of my conjecture $E(3)=\left(\frac{2}{\pi}\right)^3\int_0^{\pi/2}\int_0^{\pi/2}\int_0^{\pi/2}\frac{(\tan x_1)(\tan x_2)(\tan x_3)}{\tan x_1+\tan x_2+\tan x_3}dx_1dx_2dx_3=0.9999996\dots$, according to Desmos. I guess it's actually 1. For $n>3$, Desmos and Wolfram Cloud don't work, so I used Excel with about a million trials. I got: $E(4)\approx2.480$, whereas $\left(\frac{\pi}{2}\right)^{2(4)-6}\approx2.467$ $E(5)\approx6.047$, whereas $\left(\frac{\pi}{2}\right)^{2(5)-6}\approx6.088$ $E(6)\approx16.794$, whereas

$\left(\frac{\pi}{2}\right)^{2(6)-6}\approx15.022$ $E(7)\approx37.495$, whereas $\left(\frac{\pi}{2}\right)^{2(7)-6}\approx37.065$ As n increases, the values of $\dfrac{\prod_{k=1}^n\tan x_k}{\sum_{k=1}^n\tan x_k}$ fluctuate more and more, so this numerical approach becomes less reliable. If my conjecture is true, I can't imagine what a proof would look like.

论坛问题帖 | has_answers | Dan | 2023-11-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019320 | MSE-4814920

What is the change in angle of of a symmetric matrix when summed over a plane?

A symmetric matrix has orthogonal eigenvectors with real eigenvalues, and hence can be thought of as scaling along a particular orthogonal set of axes. Of course, not all vectors are (usually) eigenvectors, and therefore many individual vectors will have their direction changed under a symmetric matrix. It seems to me, however, that these direction changes will somehow "cancel out" in the sense that there is no net rotation by a symmetric matrix. I don't know how to state this more precisely, other than to observe that $A_{ij} - A_{ji}$ is a measure of the rotation in the i,j plane (see here). That is, if A is a symmetric matrix, for any plane P, the net direction change of Av over all vectors $v \in P$ is zero. Again, I'm not sure how state this more precisely, but perhaps it could be formulated as an integral. Is my intuition correct? And how can it be stated more precisely?

论坛问题帖 | no_answers_at_retrieval | SRobertJames | 2023-11-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019321 | MSE-4815410

Conjecture: In Pascal's triangle with n rows, the proportion of numbers less than the centre number approaches $e^{-1/\pi}$ as $n\to\infty$.

Consider Pascal's triangle with 30 rows (the top 1 is the 0th row). The centre number is the number in the middle of row $30\times \frac23=20$, which is $\binom{20}{10}=184756$. The proportion of the numbers in the triangle that are less than this centre number is $\frac{364}{496}\approx0.7339$. In the following graph, the horizontal axis is n, the number of rows in the triangle (n is a multiple of 3). The vertical axis is P, the proportion of numbers that are less than the centre number. It looks like P is approaching some limit. The rightmost data point is $n=138$ and $P=\frac{7078}{9730}\approx0.72744$. I put 0.72744 into Wolfram, and it suggested $e^{-1/\pi}\approx 0.72738$. Is the following conjecture true or false: Conjecture: In Pascal's triangle with n rows, where n is a multiple of 3, the proportion of numbers less than the centre number approaches $e^{-1/\pi}$ as $n\to\infty$. I would not be too surprised if this is in fact the limiting proportion, since e is related to Pascal's triangle and so is π. My attempt: The numbers in the top $\frac23$ of the triangle are all less than the centre number. So we just need to determine the proportion of numbers in the bottom $\frac13$ that are less than the centre number. I considered the row that is m rows below the centre number: I tried to work out the proportion of numbers in this row that are less than the centre number, to obtain an asympototic boundary curve, but I didn't succeed. Context: I was trying to approximate the median of the numbers in the first n rows of Pascal's triangle, and in my investigation I stumbled upon this possible result.

论坛问题帖 | has_answers | Dan | 2023-11-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019322 | MSE-4815847

Can this be done? Split Pascal's triangle (without the 1s) with a straight line into two regions of equal sums.

Consider Pascal's triangle with n rows, without the 1s, with each number corresponding to a vertex on a pyramid of equilateral triangles, as shown below with example $n=5$. Does there exist an n such that Pascal's triangle with n rows, without the 1s, can be divided by a straight line (that does not pass through any vertex) into two regions of equal sums? For example, with $n=5$, the red line below fails to divide the triangle into two regions of equal sums: the upper-left region has a sum of 58, and the bottom-right region has a sum of 56. I have tried to do this with different size triangles, without success. It seems that it cannot be done, but I don't know how to prove it. Remarks We exclude the 1s because the triangle with the 1s has a total sum of 2^n-1, an odd number, making an equal split obviously impossible. The sums of the all the terms in the first n rows (not just one row) are given by A145654 (ignoring the initial 0): $2,8,22,52,114,240,494\dots$ I considered the partial sums of a row of Pascal's triangle, but such expressions are not simple. Taking @Jean Marie's suggestion, I've added a left-justified version of the triangle, which may be easier to use. EDIT: Posted on MO

论坛问题帖 | no_answers_at_retrieval | Dan | 2023-11-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019323 | MSE-4816337

When does the magnitude of the gradient equal the surface area of the $dxdy$ patch?

Given a surface S in $\mathbb R^3$, what is the relationship between the gradient (when S is defined as a level curve of function $F: \mathbb R^3 \to \mathbb R$) and surface area? I noticed such a relationship when examining the surface area of the saddle surface $z = x^2 - y^2$. The surface area has integrand $\sqrt{1 + 4x^2 + 4y^2}$. And, when defined as a level curve $F(x,y,z) = x^2 - y^2 - z$, we get $\|\nabla F\| = \sqrt{1 + 4x^2 + 4y^2}$. What is the relationship between surface area and the gradient? Does it hold for $\mathbb R^n$ or only $\mathbb R^3$? My (incomplete) attempt is below. Partial Solution If S is a surface defined as the level curve of $F(x,y,z)$, and $\frac {\partial F}{\partial z} = \pm 1$, then the magnitude of ∇F equals the ratio of surface area of a portion of S to its projection on the xy plane, and $$\text{Surface Area } = \iint \|\nabla F\| dx dy.$$ Given a surface S, point $v \in S$, and plane P, define surface amplification of S over P as $$\lim_{\delta \to 0} \frac{\text{surface area of }R}{\text{area of }R \text{ projected onto } P}$$ where R is a region of S including point v with area δ. (Formalizing this limit is hard.) If S is parameterized by $(x, y, Z(x,y))$, then surface amplification over the xy plane equals $$\sqrt{1 + \frac{\partial Z}{\partial x}^2 + \frac{\partial Z}{\partial y}^2}$$ as can be easily seen from the standard surface area integral. If S is defined as the level surface of $F(x,y,z)$, and $\frac {\partial F}{\partial z} = \pm 1$, then surface amplification over the xy plane equals $\| \nabla F \|$, which follows from the definition of

gradient. (How do I show this clearly?) I conjecture this applies for any $\mathbb R^n$. This post may be related. Is this correct? How do we complete the proof? What is the geometric intuition behind this?

论坛问题帖 | has_answers | SRobertJames | 2023-11-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019324 | MSE-4817807

An infinite linear system of equations with an uncountable number A of equations

I will start with an example to make things clear and avoid confusion : Take all $x>0$ and $$\exp(x) = \sum_{-1<n} a_n x^n$$ Now finding a_n can be described as an infinite linear system of equations : $$a_0 + a_1 1 + a_2 1 + ... = e$$ $$a_0 + a_1 2 + a_2 4 + ... = e^2$$ $$...$$ and for every $x>0$ $$a_0 + a_1 x + a_2 x^2 + ... = e^x$$ So we have a countable number of variables and an uncountable number of equations. Notice there is only one unique solution set a_n here. This leads to : Conjecture : An infinite linear system of equations with an uncountable number A of equations, but a countable number B of variables has at most countable ordinal C * Linear independant * solutions, where $C < B$ and Linear independant means not a COUNTABLE LINEAR combination of others. Notice that (in contrast) a countable number of (linear) equations with a countable number of variables CAN have an uncountable number of solutions, who are linear independant. For instance the (formal) taylor series solution to $f(x) = f(\exp(x))$ has an uncountable number of solutions and it is equivalent to a system of linear equations with a countable number of variables (the taylor coefficients). Explicitly: $$t_i = \sum_n \frac{t_n n^i}{n!}$$ And examples without calculus, dynamics or functional equations exist as well.

论坛问题帖 | no_answers_at_retrieval | mick | 2023-11-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019325 | MSE-4818640

Take a random walk on Pascal's triangle, without revisits: Does the final number have infinite expectation?

Let's take a random walk on Pascal's triangle, starting at the top. Each number is in a regular hexagon. At each step, we can move to any adjacent hexagon with equal probability, but we cannot revisit a hexagon. The walk ends when we cannot move. Here is an example in which the final number is 15. Does the final number have infinite expectation? (Paradoxically, a random variable whose possible values are all finite, can have infinite expectation.) My thoughts I suspect the answer is yes, because as we move down Pascal's triangle, the numbers quickly become very large. But I don't know how this could be proved. After some step, if we find ourselves in "open territory" (no previously visited hexagons nearby), then there is a sequence of five moves that would end the walk. This is illustrated by this image provided by @Stef in the comments: (The first red line segment after the blue line segment could have been in any other direction, so we don't count that one.) The probability of such a sequence would be $\left(\frac{1}{5}\right)\left(\frac{1}{5}\right)\left(\frac{1}{5}\right)\left(\frac{1}{5}\right)\left(\frac{1}{4}\right)=\frac{1}{2500}$. Using the expectation of a geometric distribution, the expected number of steps in a walk would be roughly 2500. I believe, but am not sure, that the expected row number at the end would be on the order of $\sqrt{2500}=50$. (This rough analysis ignores several factors, for example the effect of the boundaries of the triangle.) But sometimes our walk would go much deeper than the expected depth. The numbers in the triangle increase very quickly. Generally speaking, if the numbers increase faster than the probabilities of reaching them decrease, then the expectation of the final number could be infinite. Context This is one of several questions about Pascal's triangle that I've asked recently, for example: Does the interior of Pascal's triangle contain three consecutive integers? I tried to kill the central binomial coefficients, but they came back Conjecture: In Pascal's triangle with n rows, the proportion of numbers less than the centre number approaches $e^{-1/\pi}$ as $n\to\infty$.. Sometimes I am more interested in the methods used to answer the questions, than the answers themselves (but I am still curious what the answer is!). Many of my questions initially seemed almost unapproachable to me, but then turned out to have a rational explanation.

论坛问题帖 | has_answers | Dan | 2023-12-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019326 | MSE-4819179

Conjecture: $\prod\limits_{k=0}^{n}\binom{2n}{k}$ is divisible by $\prod\limits_{k=0}^n\binom{2k}{k}$ only if $n=1,2,5$.

The diagram shows Pascal's triangle down to row 10. I noticed that the product of the blue numbers is divisible by the product of the orange numbers. That is (including the bottom centre number 252 in both products), $$\frac{\prod\limits_{k=0}^{5}\binom{10}{k}}{\prod\limits_{k=0}^5\binom{2k}{k}}=\frac{2857680000}{4233600}=675\in\mathbb{Z}.$$ Now let $$P=\frac{\prod\limits_{k=0}^{n}\binom{2n}{k}}{\prod\limits_{k=0}^n\binom{2k}{k}}$$ According to Wolfram, for $n=1$ to $n=50$, P is an integer only when $n=1,2,5$. Is the following conjecture true or false: P is an integer only when $n=1,2,5$. P can be rewritten as $$P=\prod\limits_{k=0}^n\frac{(2n)!k!}{(2k)!(2n-k)!}$$ but that doesn't seem to make the question easier to answer. I did a search on the OEIS using "$1,2,5$ Pascal divisible", and found no sequence with $1,2,5$ and the next number being greater than 50. Context: I have been trying to unravel some of the mysteries of Pascal's triangle recently, for example with questions about random walks, three consecutive numbers and killing.

论坛问题帖 | has_answers | Dan | 2023-12-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019327 | MSE-4821709

Conjecture: If $\sum_{i=1}^n \frac{1}{x_i}=1$ then $x_i | x_j$

Let $x_1,x_2,x_3,\cdots , x_n\in\mathbb{N}$ prove that if $\sum_{i=1}^n\frac{1}{x_i}=1$, then there exist x_j,x_i such that $x_j | x_i$. I read that I should avoid the no clue questions, but this is a conjecture I came up with and need to know if it's true to continue my work on some project. All I could figure out is that it's true for $n=2$ - and then $x_1=x_2=\frac{1}{2}$ Edit The user @deif proved the conjecture if there exist atleast one x_i which is prime. So to prove the conjecture entirely(if it's true) we just need to prove it with the assumption that $x_1,x_2,\cdots ,x_n$ are not primes!

论坛问题帖 | has_answers | Moaoly | 2023-12-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019328 | MSE-4822524

Does this recursive journey through Pascal's triangle always reach 1?

Let $p(k)$ be the k^th number in Pascal's triangle, numbered from left to right in each row, going down the rows. So, for example, $p(1)$ to $p(10)$ are $\binom{0}{0},\space \binom{1}{0},\space \binom{1}{1},\space \binom{2}{0},\space \binom{2}{1},\space \binom{2}{2},\space \binom{3}{0},\space \binom{3}{1},\space \binom{3}{2},\space \binom{3}{3}$, respectively. Now define sequence a_n as: $$a_1\text{ is a positive integer}$$ $$a_{n+1}=p(a_n)$$ Is the following conjecture true or false: Conjecture: $\lim\limits_{n\to\infty}a_n=1$ for all $a_1\in\mathbb{Z^+}$. For example, with $a_1=150$ we have: $a_2=p(150)=560$ $a_3=p(560)=32$ $a_4=p(32)=35$ $a_5=p(35)=7$ $a_6=p(7)=1$ $a_n=1$ for $n\ge 6$ With $a_1=100$ we have: $a_2=p(100)=1287$ $a_3=p(1287)=37353738800$ $a_4=p(37353738800)=\space ?$ If an a_n value is very large, then a_{n+1} is likely to be much larger than a_n. However, even if the terms become very large, it is possible that eventually one of them will be small enough so that the sequence will become an endless stream of 1s. I don't know how the sequence will play out. Here and here is the sequence $p(k)$, except their index is shifted down 1 from mine. So for example, my $p(5)$ is numbered as the 4th term is the linked sequence. Context: I have been playing with Pascal's triangle, investigating some of its mysterious, geometrical and humourous properties.

论坛问题帖 | has_answers | Dan | 2023-12-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019329 | MSE-4823017

A generalization of the Stolz-Cesàro: For $k\ge 1$, $\lim_{n\to\infty }\frac{a_{n+k}-a_n}{b_{n+k}-b_n}=l$ implies $\lim_{n\to\infty}\frac{a_n}{b_n}=l$

Stolz-Cesàro theorem case $\frac{*}{\infty}$:- If $b_n $ is a monotone increasing sequence and $\lim \limits_{n \to \infty} b_n = \infty $, and if $\lim \limits_{n \to \infty} \frac{a_{n+1}-a_n}{b_{n+1}- b_n}= l \in \overline{\mathbb{R}} $, then $\lim \limits_{n \to \infty} \frac{a_n }{b_n}=l$. Stolz-Cesàro theorem case $\frac{0}{0}$:- If $b_n $ is a monotone decreasing sequence and $\lim \limits_{n \to \infty} b_n = \lim \limits_{n \to \infty} a_n = 0 $, and if $\lim \limits_{n \to \infty} \frac{a_{n+1}-a_n}{b_{n+1}- b_n}= l \in \overline{\mathbb{R}} $, then $\lim \limits_{n \to \infty} \frac{a_n }{b_n}=l$. I was solving this limit :If $a_n \to a \in \mathbb{R}$ find $$\lim\limits_{n \to \infty } \frac{\sum\limits_{k=1}^n (-1)^{n-k} 2^k a^k}{2^n}$$ The numerator is a bit annoying to apply Stolz-Cesàro theorem to because of that $(-1)^{n-k}$. The numerator is $\sum\limits_{k=1}^n (-1)^{n-k} 2^k a^k$ when n is odd and $-\sum\limits_{k=1}^n (-1)^{n-k} 2^k a^k$ when n is even. It would have been much easier if $\lim\limits_{n \to \infty } \frac{a_{n +2}- a_n}{b_{2+n}- b_n}= l$ was true, and out of curiosity, I tried to use this statement here: $$\lim \limits_{n \to \infty} \frac{2^{n+2}a_{n+2}-2^{n+1}a_{n+1}}{2^{n+2}-2^{n}}=\lim \limits_{n \to \infty} \frac{4a_{n+2}-2a_{n+1}}{3} =\frac{2a}{3}$$ Which, in fact, is the correct limit! This got me wondering if there is a generalisation of the Stolz-Cesàro Theorem. Generalized Stolz-Cesàro theorem case $\frac{*}{\infty}$:- If $b_n $ is a monotone increasing sequence and $\lim \limits_{n \to \infty} b_n = \infty $, and $\exists k \in \mathbb{N}$ st $\lim \limits_{n \to \infty} \frac{a_{n+k}-a_n}{b_{n+k}- b_n}= l \in \overline{\mathbb{R}} $, then $\lim \limits_{n \to \infty} \frac{a_n }{b_n}=l$. Generalized Stolz-Cesàro theorem case $\frac{0}{0}$:- If $b_n $ is a monotone decreasing sequence and $\lim \limits_{n \to \infty} b_n = \lim \limits_{n \to \infty} a_n = 0 $, and if $\exists k \in \mathbb{N}$ st $\lim \limits_{n \to \infty} \frac{a_{n+k}-a_n}{b_{n+k}- b_n}= l \in \overline{\mathbb{R}} $, then $\lim \limits_{n \to \infty} \frac{a_n }{b_n}=l$. Example: $\lim_limits_{n \to \infty}\frac{(-1)^n}{n } =0$ but the ordinal Stolz-Cesàro Limit don't exist , if the generalisation was true then $\lim \limits_{n \to \infty} \frac{a_{n+2} -a_{n}}{b_{n+2}-b_{n}}= 0$ (I know this is a trivial example) If it is not true, then what are the sufficient conditions for a_n,b_n so that this generalisation of the Stolz-Cesàro theorem becomes true?

论坛问题帖 | has_answers | pie | 2023-12-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019330 | MSE-4828000

Conjecture: $\sum\limits_{k=1}^nk^m=S_3(n)\times\frac{P_{m-3}(n)}{N_m}$ for odd $m>1 \ ;\ =S_2(n)\times\frac{P_{m-2}'(n)}{N_m}$ for even m.

When I was in high school, I was fascinated by $\displaystyle\sum\limits_{k=1}^n k= \frac{n(n+1)}{2}$ so I tried to find the general solution for $\displaystyle\sum\limits_{k=1}^n k^m$ s.t $m \in \mathbb{N}$. I used a similar approach to my answer here My approach was turning n^m to $n(n-1)(n-2)\dots (n-m+1)+ f(n)$ and to use the fact that $$\displaystyle\dbinom{n}{k}+\dbinom{n}{k+1}= \dbinom{n+1}{k+1}$$ $f(n) $ will be combination of sums $\displaystyle\sum_{k=1}^n k^i$ st $i<m$ which I evaluated before I was able to to find the sum up to $m=6$. Here, I tried to search for a pattern to find the general solution of $\sum\limits_{k=1}^n k^m $ s.t $m \in \mathbb{N}$ which I failed to do, but I noticed this pattern: $\\[3pt]$ $$S_1(n):={\sum\limits_{k=1}^n k= \frac{n(n+1)}{2}}$$ $$S_2(n):={\sum\limits_{k=1}^n k^2= \color{blue}{\frac{n(2n+1)(n+1)}{6}}}$$ $$S_3(n):={\sum\limits_{k=1}^n k^3= \color{red}{\frac{\color{red}{n^2(n+1)^2}}{\color{red}{4}}}}$$ $$S_4(n):={\sum\limits_{k=1}^n k^4=\color{blue}{\frac{n(2n+1)(n+1)}{6}} \times \frac{3n^2+3n-1}{ 5}}$$ $$S_5(n):={\sum\limits_{k=1}^n k^5= \color{red}{\frac{\color{red}{n^2(n+1)^2}}{\color{red}{4}}} \times\frac{2n^2+2n-1}{ 3}}$$ $$S_6(n):={\sum\limits_{k=1}^n k^6= \color{blue}{\frac{n(2n+1)(n+1)}{6}} \times \frac{3n^4+6n^3-3n+1}{ 7 }}$$ $$S_7(n):={\sum\limits_{k=1}^n k^7= \color{red}{\frac{\color{red}{n^2(n+1)^2}}{\color{red}{4}} }\times\frac{3n^4+6n^3-n^2-4n+2}{ 6}}$$ $$S_8(n):={\sum\limits_{k=1}^n k^8= \color{blue}{\frac{n(2n+1)(n+1)}{6}} \times \frac{5n^6+15n^5+5n^4-15n^3-n^2+9n+3}{ 15}}$$ $$S_9(n):={\sum\limits_{k=1}^n k^9= \color{red}{\frac{\color{red}{n^2(n+1)^2}}{\color{red}{4}}} \times\frac{(n^2+n-1)(2n^4 +4n^3-n^3-3n^2+3)}{ 5}}$$ $$S_{10}(n):={\sum\limits_{k=1}^n k^{10}= \color{blue}{\frac{n(2n+1)(n+1)}{6}} \times \frac{ 3 n^8+ 12 n^7+ 8 n^6 - 18 n^5- 10 n^4+ 24 n^3 + 2 n^2 - 15 n +5}{ 11}}$$ $$S_{11}(n):={\sum\limits_{k=1}^n k^{11}=

\color{red}{\frac{\color{red}{n^2(n+1)^2}}{\color{red}{4}}} \times\frac{2n^8 +8n^7+4n^6-16n^5-5n^4+26n^3-3n^2-20n+10}{ 6}}$$ $\\[6pt]$ I noticed that: For odd $m>1$, $\displaystyle\sum\limits_{k=1}^n k^m= \color{red}{\frac{{n^2(n+1)^2}}{{4}}} \times \frac{P_{m-3}(n)}{N_m}$ s.t $P_{m-3}(n)$ is an $m-3$ polynomial with integer coefficients $ \{a_{m-3},\dots a_1,a_0 \}$ such that $\gcd \{ a_{m-3},\dots a_1,a_0 \}=1$, $N_m\in \mathbb {N}$. For even m, $\displaystyle\sum\limits_{k=1}^n k^m= \color{blue}{\frac{n(2n+1)(n+1)}{6}}\times \frac{P_{m-2}'(n)}{N_m}$ s.t $P_{m-2}'(n)$ is an $m-2$ polynomial with integer coefficients $\{ a_{m-2},\dots a_1,a_0 \}$ such that $\gcd \{ a_{m-2},\dots a_1,a_0 \}=1$, $N_m\in \mathbb {N}$. When I was in high school I couldn't prove this pattern, and this question reminded me of this observation that I had totally forgotten about. Now, after two years from my first attempt, I tried to prove this pattern again, but I couldn't. EDIT This question has been partly answered in this MathOverflow answer (The answer shows that $\displaystyle\sum\limits_{k=1}^n k^m$ is divisible by ${n^2(n+1)^2}$ for odd $m>1$ and $\displaystyle\sum\limits_{k=1}^n k^m$ is divisible by ${n(2n+1)(n+1)}$ for even m) the only missing part is to show that the denominator is a multiple of 4 if $m\in 2\mathbb{N}+1$, and the denominator is a multiple of 6 if $m \in 2\mathbb{N}$. Although this question has been almost solved before, I still would like to see if there is any other proofs or methods. Update: I asked this question on MO here

论坛问题帖 | has_answers | pie | 2023-12-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019331 | MSE-4830769

Does $\int_{1}^{2}\ln\left(x+\ln\left(x^2+\ln\left(x^3+\ln\left(\dots\right)\right)\right)\right)dx=1$?

I was messing around with the infinitely nested logarithm $f(x)=\ln\left(x+\ln\left(x^2+\ln\left(x^3+\ln\left(\dots\right)\right)\right)\right)$ on Desmos when I decided to take the integral from $x = 1$ to $x = 2$. Here's what I got where n represents the number of times the logarithm is nested. For example, for $n=1$, it is $f_1(x)=\ln\left(x\right)$ and for $n=2$, it is $f_2(x)=\ln\left(x+\ln\left(x^2\right)\right)$. $$\begin{array}{c|c} \text{n} & \int_{1}^{2}{f_n(x)}dx \\ \hline 1 & 0.38629436112 \\ \hline 2 & 0.769701324476 \\ \hline 3 & 0.923253197047 \\ \hline 4 & 0.968765188969 \\ \hline 5 & 0.982950647903 \\ \hline 10 & 0.992710069039 \\ \end{array}$$ $\therefore\text{I conjecture that it converges to 1}$ Can someone help me analytically prove or refute my conjecture? I lack the skills to do it. P.S. I could not figure out a closed form for the infinitely nested logarithm but I assume finding one would make this a lot easier to prove.

论坛问题帖 | has_answers | Dylan Levine | 2023-12-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019332 | MSE-4833672

Conjecture about boundary commuting with union

Given a Topological space (X,τ) and two disjoint open sets A,B, the following holds: $$\partial(A\cup B) = \partial A\cup \partial B$$ I am interested in a modified version of this. Construction: Let $\mathcal{R}$ be the collection of rays (one dimensional subspaces) of $\mathbb{R}^2$. We call a subset of $\mathbb{R}^2$ an open stick, if it is bounded, connected, contains $(0,0)$, is contained in some $r\in \mathcal{R}$, and is open in the subspace Topology of this r. Here is your typical stick (Desmos doesn't do open dots at the ends for some reason): Now suppose we have some stick family (s_r) indexed by $\mathcal{R}$. Also, for each stick s_r, denote by $\partial's_r$ its boundary in the subspace r. Question: When is it possible to conclude $$ \partial \bigcup_{r\in \mathcal{R}} s_r = \bigcup_{r\in \mathcal{R}}\partial' s_r \hspace{ 0.5cm } ?$$ I conjecture that it is precisely when $ \bigcup_{r\in \mathcal{R}}\partial' s_r \simeq S^1$, the unit circle! The intuition for this is rather simple, for this identity to hold, we need each stick to contribute only its endpoint to the boundary of the union, and to prevent a stick from contributing a point other than its endpoint, we need to ensure that there is no discontinuity in the length of the sticks, at that particular stick. But I am looking for help to turn this into a rigorous argument.

论坛问题帖 | no_answers_at_retrieval | Carlyle | 2023-12-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019333 | MSE-4835643

Conjecture: $\binom{n}{k } \mod m =0$ for all $k=1,2,3,\dots,n-1$ only when $m $ is a prime number and n is a power of m

While playing with Pascal's triangle, I observed that $\binom{4}{k } \mod 2 =0$ for $k=1,2,3$,and $\binom{8}{k } \mod 2 =0$ for $k=1,2,3,4,5,6,7$ This made me curious about the values of $n>1$ and $m>1$ that satisfy $\binom{n}{k } \mod m =0$ for all $k=1,2,3,\dots,n-1$ I asked my friend to write a $C++$ code that could test this condition for n up to 10000 and m up to 31 and I found out that $\binom{n}{k } \mod m =0$ for all $k=1,2,3,\dots,n-1$ only when $m $ is a prime number and n is a power of m. #include <bits/stdc++.h> using namespace std;` typedef long long ll; #define rep(i , st , ed) for(int i = st; i < ed; i++) #define f first #define s second const int N = 10000; const int mod = 31; ll nCr[N][N] , frq[N]; void gen(){ nCr[0][0] = 1; for (int i = 0; i < N; ++i) { nCr[i][0] = nCr[i][i] = 1; for (int j = 1; j < i; ++j) { nCr[i][j] = (nCr[i-1][j] + nCr[i-1][j-1]) % mod; if(nCr[i][j] == 0) frq[i]++; } } } ll get(int n , int r){ return (n >= r) ? nCr[n][r] : 0; } int main(){ ios::sync_with_stdio(0); cin.tie(NULL); cout.tie(0); gen(); for(int i = 1; i < N; ++i) if(frq[i] == i - 1){ cout << i << '\n'; } } After many attempts to prove or disprove this conjecture, I couldn’t find a solution. This statement is the same as $$\gcd\left(\binom{n}{1 }, \binom{n}{2 },\dots , \binom{n}{n-1 } \right)= \begin{cases} p, & \text{if n is a power of a prime p} \\[2ex] 1, & \text{else. } \end{cases}$$

论坛问题帖 | has_answers | pie | 2023-12-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019334 | MSE-4840933

Conjecture: If A,B,C are random points on a sphere, then $E\left(\frac{\text{Area}_{\triangle ABC}}{\text{Area}_{\bigcirc ABC}}\right)=\frac14$.

On (not in) a sphere, choose three independent uniformly random points A,B,C. Is the following conjecture true: The expectation of the ratio of the area of (planar) $\triangle ABC$ to the area of the (planar) disk with A,B,C on its perimeter, is $1/4$. My conjecture is based on experimentation: using sphere point picking, I ran 1.5×10^7 trials and the average ratio was about 0.2500075. Since $1/4$ is a simple ratio, I am wondering if there is an intuitive explanation (here is a recent example), but any proof is welcome. Remarks: Assume the sphere is a unit sphere. The expected area of the disk is $\frac{4\pi}{5}$ (proof). The expected area of the triangle seems to be $\frac{\pi}{5}$ based on experimentation, but I don't have a proof (Wolfram has a page called "Sphere triangle picking" but it's empty). Curiously, the expectation of the ratio seems to equal the ratio of the expectations. The probability that the ratio is greater than $1/4$ is about 0.534. The standard deviation of the ratio is about 0.10667 (could it be $8/75$?). The probability density function of the ratio looks roughly like this: The expectation of the reciprocal ratio (i.e. the ratio of the area of the disk to the area of the triangle) seems to be about 6.6, but this ratio flucuates a lot and I'm not sure its expectation even exists. My attempt: I tried to adopt the techniques in the proof that the area of the disk is $\frac{4\pi}{5}$ (with a unit sphere), but I don't know how to express the area of the triangle in a usable way. I tried to find a more intuitive explanation, possibly using symmetry, but failed. (In the $2D$ case, the expected ratio of areas is $\frac{3}{2\pi^2}\approx 0.15198$ (proof), which is not an intuition-friendly number, so it seems that an intuitive explanation would apply to the $3D$ case but not the $2D$ case.) Context: This question was inspired by a question about a random triangle inscribed in a circle.

论坛问题帖 | has_answers | Dan | 2024-01-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019335 | MSE-4841690

Values of $\Phi_n(-1)$

Let $$\Phi_n(x) = \prod_{0<k\leq n, \gcd(k,n)=1}(x-e^{\frac{2\pi i k}{n}})$$ be the n-th cyclotomic polynomial. By observation it seems that cyclotomic polynomials when evaluated at $x=-1$ give the following $$\Phi_n(-1) = \begin{cases} 2 \quad \text{if} \quad n = 2^k, k>1\in \mathbb{N}\\ p \quad \text{if} \quad n = 2p\\ 1 \quad \text{otherwise} \end{cases}$$ where p denotes a prime (> 2). The first few cases are $$\Phi_1(-1) = -2$$ $$\Phi_2(-1) = 0$$ $$\Phi_3(-1) = 1$$ $$\Phi_4(-1) = 2$$ $$\Phi_5(-1) = 1$$ $$\Phi_6(-1) = 3$$ $$\Phi_7(-1) = 1$$ $$\Phi_8(-1) = 2$$ however we will only consider $n>2$ for the conjecture. This I thought was quite weird at first as it seems like a very simple pattern for something as complicated as the coefficients of cyclotomic polynomials. I managed to prove the first two cases, and will share the short proofs below, but am unsure of how to tackle the last case. I don't have any way to check for high values of n other than calculation, and so I apologise in advance if there is a counterexample to this conjecture that can easily be checked with a computer. Below are the proofs for $n= 2^k$ and $n=2p$, $n = 2^k$ By Möbius inversion we get $$\Phi_n(x) = \prod_{d\mid n}(x^d-1)^{\mu\left(\frac{n}{d}\right)}$$ Since $\mu(2^n) = 0$ for $n>1$ with $\mu(2)=-1$, therefore $$\Phi_{2^n}(x) = (x^{2^n}-1)(x^{2^{n-1}}-1)^{-1} = x^{2^{n-1}}+1$$ $$\Phi_{2^n}(-1) = 2$$ $n = 2p$ We can use the following identity $$\Phi_{2p}(x) = \sum_{k=0}^{p-1}(-1)^kx^k$$ $$ = \sum_{k=0}^{p-1}1 = p$$ But for the 'otherwise' case, I don't know how to approach it. It seems to general to approach in any specific way using identities or by looking at the values of $\mu(d)$ as that would require more information about the divisors of n. Any help or a proof/counterexample is appreciated!

论坛问题帖 | has_answers | Mako | 2024-01-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019336 | MSE-4846986

Moments of Alternate Elliptic Integrals

Let K_s denote Ramanujan's Alternate Elliptic Integrals as follows: $$K_s:=K_s(k)=\frac{\cos(\pi s)}{2}\int_0^1\frac{t^{s-1/2}}{(1-t)^{1/2+s}(1-k^2t)^{1/2-s}}dt$$ Where in Hypergeometric Form it is: $$K_s=\frac{\pi}{2}{}_2F_1\left[\frac{1}{2}+s,\frac{1}{2}-s,1,k^2\right]$$ The following signatures are important in this theory: $$s=\left\{0,\frac{1}{3},\frac{1}{4},\frac{1}{6}\right\}$$ I have found a General Theorem to calculate the following Integral: $$\int_0^1kK_uK_vdk$$ As examples I will present the corollaries that follow from the Theorem while considering the combination of 4 signatures. This gives us the following 10 Integrals: Corollaries: $$\int_0^1kK^2dk=\frac{7\zeta(3)}{4}$$ $$\int_0^1kK^2_{1/6}dk=\frac{9\sqrt{3}}{8}\operatorname{Cl_2}(\pi/3)$$ $$\int_0^1kK^2_{1/3}dk=\frac{15\sqrt{3}}{16}\operatorname{Cl_2}(\pi/3)$$ $$\int_0^1kK^2_{1/4}dk=2\beta(2)$$ $$\int_0^1kKK_{1/6}dk=\frac{27\sqrt{3}}{4}\ln 3-9\sqrt{3}\ln 2$$ $$\int_0^1kKK_{1/3}dk=\frac{27}{16}\ln 3$$ $$\int_0^1kKK_{1/4}dk=2\sqrt{2}\ln 2$$ $$\int_0^1kK_{1/3}K_{1/6}dk=\frac{3\sqrt{3}}{2}\ln 2$$ $$\int_0^1kK_{1/4}K_{1/6}dk=\frac{27\sqrt{6}}{5}\ln 2-\frac{27\sqrt{6}}{10}\ln3$$ $$\int_0^1kK_{1/4}K_{1/3}dk=\frac{27\sqrt{2}}{14}\ln 3-\frac{9\sqrt{2}}{7}\ln 2$$ The very first Integral is known in this list as K_0 is just the Complete Elliptic Integral of the First Kind K. I have not seen the General Theorem and Other Integrals as well anywhere in Literature. So I was wondering whether these results are new or already known? This is only a Small List, one may produce as many Integrals as they want from the General Theorem. Like for example after Differentiating: $$\int_0^1kK\left[\frac{d}{dv}K_v\right]_{v=1/4}dk=2\sqrt{2}\left(8\beta(2)-\pi \ln2-8\ln 2\right)$$ This opens up new possibilities too. The following is for example already known: $$\int_0^1k^{2n+1}K^2dk=a_n\zeta(3)+b_n$$ where $a_n,b_n\in\mathbb{Q}$. This is found using "FindIntegerNullVector": Conjecture: $$\int_0^1k^{2n+1}K^2_{1/6}dk=a_n\sqrt{3}\operatorname{Cl_2}(\pi/3)+b_n$$ $$\int_0^1k^{2n+1}K^2_{1/3}dk=a_n\sqrt{3}\operatorname{Cl_2}(\pi/3)+b_n$$ $$\int_0^1k^{2n+1}K^2_{1/4}dk=a_n\beta(2)+b_n$$

$$\int_0^1k^{2n+1}KK_{1/3}dk=a_n\ln 3+b_n$$ where $a_n,b_n\in\mathbb{Q}$. The rest of the cases doesn't seem to continue the pattern. I wonder if there is any greater meaning behind this: $$\left[\begin{array}{c} \text{Integral}&\text{Basis} \\\\ \displaystyle\int_0^1k^{2n+1}K_0^2dk&\zeta(3)\\\\ \displaystyle\int_0^1k^{2n+1}K_{1/6}^2dk&\sqrt{3}\operatorname{Cl_2}(\pi/3)\\\\ \displaystyle\int_0^1k^{2n+1}K_{1/3}^2dk&\sqrt{3}\operatorname{Cl_2}(\pi/3)\\\\ \displaystyle\int_0^1k^{2n+1}K_{1/4}^2dk&\beta(2)\\\\ \displaystyle\int_0^1k^{2n+1}K_0K_{1/3}dk&\ln3\\\\ \end{array} \right]$$

论坛问题帖 | has_answers | Miracle Invoker | 2024-01-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019337 | MSE-4847015

Conjecture about average cluster size in a prime-like sequence

Consider the sequence $u_n=n^a+p_n$, where a is a real constant and p_n is the nth prime. Write down the terms in u_n increasing from left to right. From each term, draw a line segment connecting it to its numerically closer neighbor, choosing the lower in case of a tie, thus forming clusters. Here is an example with $a=1$. $\color{red}{3-5-8-11} \qquad \color{blue}{16-19} \qquad \color{brown}{24-27-32} \qquad \color{green}{39-42}\qquad \color{orange}{49-54-57-62-69} \qquad \color{purple}{76-79} \qquad \cdots$ The cluster sizes are $4,2,3,2,5,3,\dots$. Is the following conjecture true: Conjecture: The average cluster size is 3 if $a\le 2$, and infinity if $a>2$. My conjecture is based on numerical investigation with Excel, using the first 10^6 primes. My conjecture can be expressed more precisely as follows. For a given value of a, let $f(a,c)$ be the arithmetic mean of the sizes of the first c clusters. My conjecture is that, as $c\to\infty$, $f(a,c)$ approaches 3 if $a\le 2$, and $f(a,c)$ approaches infinity if $a>2$. Connection with Poisson distribution In a $1D$ Poisson process, the probability that a point is its nearest neighbor's nearest neighbor, is $p=2/3$, as shown here, so the average cluster size is $2/p=3$, as shown here. Coming back to this question, I guess that u_n resembles a $1D$ Poisson process in some critical way if $a\le 2$. This question was inspired by the question linked to above.

论坛问题帖 | no_answers_at_retrieval | Dan | 2024-01-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019338 | MSE-4852480

Three random points on $x^2+y^2=1$ are the vertices of a triangle. Is the probability that $(0,0)$ is inside the triangle's incircle exactly 0.13?

Three uniformly random points on the circle $x^2+y^2=1$ are the vertices of a triangle. What is the probability that $(0,0)$ is inside the triangle's incircle? (This a variation of the question "What is the probability that $(0,0)$ is inside the triangle?".) A simulation with 10^8 trials gives $P\approx 0.129963$. Is the probability exactly 0.13? If so, this would be the weirdest geometrical probability I have ever seen. My attempt: Let $d=$ distance between the circles' centres, and let $r=$ radius of the triangle's incircle. We are looking for $P(d<r)$. Euler's triangle formula tells us that $d=\sqrt{1-2r}$. So we are looking for $P(\sqrt{1-2r}<r)=P(r>\sqrt2-1)$. I do not know how to calculate this probability. I know that $r=\sqrt{\frac{(s-a)(s-b)(s-c)}{s}}$ where a,b,c are the side lengths of the triangle and $s=\frac{a+b+c}{2}$.

论坛问题帖 | has_answers | Dan | 2024-01-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019339 | MSE-4854445

Conjecture: Expected total area of a certain set of random triangles in a unit disk is $1/\pi$.

Choose $3n$ independent uniformly random points in a disk with perimeter $x^2+y^2=1$. Label the points $P_1,P_2,P_3,\dots,P_{3n}$ in order of increasing x-coordinates. Form triangles $\triangle P_1P_2P_3,\space \triangle P_4P_5P_6,\space \dots,\space \triangle P_{3n-2}P_{3n-1}P_{3n}$. Here is an example with $3n=21$. What does the total area of the triangles approach as $n\to\infty$? I did a simulation of 100 trials with $3n=99999$, and the average total area was 0.318343178. This number multiplied by π is approximately 1.00010, so I conjecture that the answer is $1/\pi$. My attempt I considered a simpler related question: What is the expected area of a triangle with vertices $(1,a_1),(2,a_2),(3,a_3)$ where the a values are independent uniformly random real numbers in $[0,1]$? The probability that a_2 is between a_1 and a_3 is $1/3$. We have $\mathbb{E}(\max (a_1,a_2,a_3)-\min (a_1,a_2,a_3))=\frac12$ (explanation). The left and right sub-triangles share a vertical side, which in this case has expected length $\frac18$. So the expected area of the triangle is $2\times\frac12\times\frac18\times 1=\frac18$. The probability that a_2 is not between a_1 and a_3 is $2/3$. Again we have $\mathbb{E}(\max (a_1,a_2,a_3)-\min (a_1,a_2,a_3))=\frac12$. The left and right sub-triangles share a vertical side, which in this case has expected length $\frac38$. So the expected area of the triangle is $2\times\frac12\times\frac38\times 1=\frac38$. So the overall expected area of the triangle is $\frac13\times\frac18+\frac23\times\frac38=\frac{7}{24}$. This value is supported by a simulation. In the original question, the triangles are not limited to a rectangular box of height 1, but rather are limited to a unit disk, which has an average height of $2\int_0^1\sqrt{1-x^2}\mathrm dx=\frac{\pi}{2}$. The rectangle and the disk have the same width, but in the disk there are horizontal gaps between neighboring triangles, so we multiply by a factor of $\frac23$. So the total area of the triangles in the disk should approach approximately $\frac{7}{24}\times\frac{\pi}{2}\times\frac23=\frac{7\pi}{72}\approx 0.30543$. This is slightly less than $\frac{1}{\pi}\approx 0.31831$. The difference may be due to the fact that the x-coordinates of the points in the disk do not have equal spacing. EDIT: I made a mistake in my simulation. Using the shoelace theorem, the area of the triangle with vertices $(x_1,y_1),(x_2,y_2),(x_3,y_3)$ is $\frac12|x_1y_2-x_2y_1+x_2y_3-x_3y_2-x_1y_3+x_3y_1|$, but I was accidentally using $\frac12|x_1y_2-x_2y_1+x_2y_3-x_3y_2-x_1y_3+x_\color{red}{1}y_1|$. After correcting, my simulations suggest the answer is approximately 0.320, which agrees with @Rei Henigman's answer. So I no longer think my conjecture is true, but I would still like to know if the limiting area of the triangles has a closed form. I also did a (correct) simulation using a square lamina instead of a disk, and it seems that the proportion covered by the triangles is about 0.102, which seems to match the proportion in the disk. Context This question was inspired by the question "A disc contains n random points. Each point is connected to its

nearest neighbor. What does the average cluster size approach as $n\to\infty$?".

论坛问题帖 | has_answers | Dan | 2024-01-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019340 | MSE-4856487

New conjecture? $(\varphi(n))! = -1 \pmod n \iff n$ is prime (nearly the same as Wilson's)

$$ (n-1)! = -1 \pmod n \text{ iff } n \text{ is prime}, \text{ is Wilson's theorem,} $$ But coincidentally for now the expression passed to factorial is $n - 1$ which is (iff n is prime) equal to $\varphi(n)$. So, I was thinking, don't we have this related conjecture: $$ \varphi(n)! = -1 \pmod n \text{ iff } n \text{ is prime} $$ Because the following example verification code seems to indicate as such: from sympy import * N = 10000 for n in range(2, N): r = factorial(totient(n)) % n if isprime(n): assert (r + 1) % n == 0 else: assert (r + 1) % n != 0 print(n, r) Which prints: 2 1 3 2 4 2 5 4 6 2 7 6 8 0 9 0 10 4 11 10 12 0 13 12 14 6 15 0 16 0 17 16 18 0 19 18 20 0 21 0 22 10 23 22 24 0 25 0 26 12 27 0 28 0 29 28 30 0 31 30 32 0 33 0 34 16 35 0 36 0 37 36 38 18 39 0 40 0 41 40 42 0 43 42 44 0 45 0 46 22 47 46 48 0 49 0 50 0 51 0 52 0 53 52 54 0 55 0 56 0 57 0 58 28 59 58 60 0 61 60 62 30 63 0 64 0 65 0 66 0 67 66 68 0 69 0 70 0 71 70 72 0 73 72 74 36 75 0 76 0 77 0 78 0 79 78 80 0 81 0 82 40 83 82 84 0 85 0 86 42 87 0 88 0 89 88 90 0 91 0 92 0 93 0 94 46 95 0 96 0 97 96 98 0 99 0 100 0 101 100 102 0 103 102 104 0 105 0 106 52 107 106 108 0 109 108 110 0 111 0 112 0 113 112 114 0 115 0 116 0 117 0 118 58 119 0 120 0 121 0 122 60 123 0 124 0 125 0 126 0 127 126 128 0 129 0 130 0 131 130 132 0 133 0 134 66 135 0 136 0 137 136 138 0 139 138 140 0 141 0 142 70 143 0 144 0 145 0 146 72 147 0 148 0 149 148 150 0 151 150 152 0 153 0 154 0 155 0 156 0 157 156 158 78 159 0 160 0 161 0 162 0 163 162 164 0 165 0 166 82 167 166 168 0 169 0 170 0 171 0 172 0 173 172 174 0 175 0 176 0 177 0 178 88 179 178 180 0 181 180 182 0 183 0 184 0 185 0 186 0 187 0 188 0 189 0 190 0 191 190 192 0 193 192 194 96 195 0 196 0 197 196 198 0 199 198 200 0 201 0 202 100 203 0 204 0 205 0 206 102 207 0 208 0 209 0 210 0 211 210 212 0 213 0 214 106 215 0 216 0 217 0 218 108 219 0 220 0 221 0 222 0 223 222 224 0 225 0 226 112 227 226 228 0 229 228 230 0 231 0 232 0 233 232 234 0 235 0 236 0 237 0 238 0 239 238 240 0 241 240 242 0 243 0 244 0 245 0 246 0 247 0 248 0 249 0 250 0 251 250 252 0 253 0 254 126 255 0 256 0 257 256 258 0 259 0 260 0 261 0 262 130 263 262 264 0 265 0 266 0 267 0 268 0 269 268 270 0 271 270 272 0 273 0 274 136 275 0 276 0 277 276 278 138 279 0 280 0 281 280 282 0 283 282 284 0 285 0 286 0 287 0 288 0 289 0 290 0 291 0 292 0 293 292 294 0 295 0 296 0 297 0 298 148 299 0 300 0 301 0 302 150 303 0 304 0 305 0 306 0 307 306 308 0 309 0 310 0 311 310 312 0 313 312 314 156 315 0 316 0 317 316 318 0 319 0 320 0 321 0 322 0 323 0 324 0 325 0 326 162 327 0 328 0 329 0 330 0 331 330 332 0 333 0 334 166 335 0 336 0 337 336 338 0 339 0 340 0 341 0 342 0 343 0 344 0 345 0 346 172 347 346 348 0 349 348 350 0 351 0 352 0 353 352 354 0 355 0 356 0 357 0 358 178 359 358 360 0 361 0 362 180 363 0 364 0 365 0 366 0 367 366 368 0 369 0 370 0 371 0 372 0 373 372 374 0 375 0 376 0 377 0 378 0 379 378 380 0 381 0 382 190 383 382 384 0 385 0 386 192 387 0 388 0 389 388 390 0 391 0 392 0 393 0 394 196 395 0 396 0 397 396 398 198 399 0 400 0 401 400 402 0 403 0 404 0 405 0 406 0 407 0 408 0 409 408 410 0 411 0 412 0 413 0 414 0 415 0 416 0 417 0 418 0 419 418 420 0 421 420 422 210 423 0 424 0 425 0 426 0 427 0 428 0 429 0 430 0 431 430 432 0 433 432 434 0 435 0 436 0 437 0 438 0 439 438 440 0 441 0 442 0 443 442 444 0 445 0 446 222 447 0 448 0 449 448 450 0 451 0 452 0 453 0 454 226 455 0 456 0 457 456 458 228 459 0 460 0 461 460 462 0 463 462 464 0 465 0 466 232 467 466 468 0 469 0 470 0 471 0 472 0 473 0 474 0 475 0 476 0 477 0 478 238 479 478 480 0 481 0 482 240 483 0 484 0 485 0 486 0 487 486 488 0 489 0 490 0 491 490 492 0 493 0 494 0 495 0 496 0 497 0 498 0 499 498 500 0 501 0 502 250 503 502 504 0 505 0 506 0 507 0 508 0 509 508 510 0 511 0 512 0 513 0 514 256 515 0 516 0 517 0 518 0 519 0 520 0 521 520 522 0 523 522 524 0 525 0 526 262 527 0 528 0 529 0 530 0 531 0 532 0 533 0 534 0 535 0 536 0 537 0 538 268 539 0 540 0 541 540 542 270 543 0 544 0 545 0 546 0 547 546 548 0 549 0 550 0 551 0 552 0 553 0 554 276 555 0 556 0 557 556 558 0 559 0 560 0 561 0 562 280 563 562 564 0 565 0 566 282 567 0 568 0 569 568 570 0 571 570 572 0 573 0 574 0 575 0 576 0 577 576 578 0 579 0 580 0 581 0 582 0 583 0 584 0 585 0 586 292 587 586 588 0 589 0 590 0 591 0 592 0 593 592 594 0 595 0 596 0 597 0 598 0 599 598 600 0 601 600 602 0 603 0 604 0 605 0 606 0 607 606 608 0 609 0 610 0 611 0 612 0 613 612 614 306 615 0 616 0 617 616 618 0 619 618 620 0 621 0 622 310 623 0 624 0 625 0 626 312 627 0 628 0 629 0 630 0 631 630 632 0 633 0 634 316 635 0 636 0 637 0 638 0 639 0 640 0 641 640 642 0 643 642 644 0 645 0 646 0 647 646 648 0 649 0 650 0 651 0 652 0 653 652 654 0 655 0 656 0 657 0 658 0 659 658 660 0 661 660 662 330 663 0 664 0 665 0 666 0 667 0 668 0 669 0 670 0 671 0 672 0 673 672 674 336 675 0 676 0 677 676 678 0 679 0 680 0 681 0 682 0 683 682 684 0 685 0 686 0 687 0 688 0 689 0 690 0 691 690 692 0 693 0 694 346 695 0 696 0 697 0 698 348 699 0 700 0 701 700 702 0 703 0 704 0 705 0 706 352 707 0 708 0 709 708 710 0 711 0 712 0 713 0 714 0 715 0 716 0 717 0 718 358 719 718 720 0 721 0 722 0 723 0 724 0 725 0 726 0 727 726 728 0 729 0 730 0 731 0 732 0 733 732 734 366 735 0 736 0 737 0 738 0 739 738 740 0 741 0 742 0 743 742 744 0 745 0 746 372 747 0 748 0 749 0 750 0 751 750 752 0 753 0 754 0 755 0 756 0 757 756 758 378 759 0 760 0 761 760 762 0 763 0 764 0 765 0 766 382 767 0 768 0 769 768 770 0 771 0 772 0 773 772 774 0 775 0 776 0 777 0 778 388 779 0 780 0 781 0 782 0 783 0 784 0 785 0 786 0 787 786 788 0 789 0 790 0 791 0 792 0 793 0 794 396 795 0 796 0 797 796 798 0 799 0 800 0 801 0 802 400 803 0 804 0 805 0 806 0 807 0 808 0 809 808 810 0 811 810 812 0 813 0 814 0 815 0 816 0 817 0 818 408 819 0 820 0 821 820 822 0 823 822 824 0 825 0 826 0 827 826 828 0 829 828 830 0 831 0 832 0 833 0 834 0 835 0 836 0 837 0 838 418 839 838 840 0 841 0 842 420 843 0 844 0 845 0 846 0 847 0 848 0 849 0 850 0 851 0 852 0 853 852 854 0 855 0 856 0 857 856 858 0 859 858 860 0 861 0 862 430 863 862 864 0 865 0 866 432 867 0 868 0 869 0 870 0 871 0 872 0 873 0 874 0 875 0 876 0 877 876 878 438 879 0 880 0 881 880 882 0 883 882 884 0 885 0 886 442 887 886 888 0 889 0 890 0 891 0 892 0 893 0 894 0 895 0 896 0 897 0 898 448 899 0 900 0 901 0 902 0 903 0 904 0 905 0 906 0 907 906 908 0 909 0 910 0 911 910 912 0 913 0 914 456 915 0 916 0 917 0 918 0 919 918 920 0 921 0 922 460 923 0 924 0 925 0 926 462 927 0 928 0 929 928 930 0 931 0 932 0 933 0 934 466 935 0 936 0 937 936 938 0 939 0 940 0 941 940 942 0 943 0 944 0 945 0 946 0 947 946 948 0 949 0 950 0 951 0 952 0 953 952 954 0 955 0 956 0 957 0 958 478 959 0 960 0 961 0 962 0 963 0 964 0 965 0 966 0 967 966 968 0 969 0 970 0 971 970 972 0 973 0 974 486 975 0 976 0 977 976 978 0 979 0 980 0 981 0 982 490 983 982 984 0 985 0 986 0 987 0 988 0 989 0 990 0 991 990 992 0 993 0 994 0 995 0 996 0 997 996 998 498 999 0 1000 0 1001 0 1002 0 1003 0 1004 0 1005 0 1006 502 1007 0 1008 0 1009 1008 1010 0 1011 0 1012 0 1013 1012 1014 0 1015 0 1016 0 1017 0 1018 508 1019 1018 1020 0 1021 1020 1022 0 1023 0 1024 0 1025 0 1026 0 1027 0 1028 0 1029 0 1030 0 1031 1030 1032 0 1033 1032 1034 0 1035 0 1036 0 1037 0 1038 0 1039 1038 1040 0 1041 0 1042 520 1043 0 1044 0 1045 0 1046 522 1047 0 1048 0 1049 1048 1050 0 1051 1050 1052 0 1053 0 1054 0 1055

0 1056 0 1057 0 1058 0 1059 0 1060 0 1061 1060 1062 0 1063 1062 1064 0 1065 0 1066 0 1067 0 1068 0 1069 1068 1070 0 1071 0 1072 0 1073 0 1074 0 1075 0 1076 0 1077 0 1078 0 1079 0 1080 0 1081 0 1082 540 1083 0 1084 0 1085 0 1086 0 1087 1086 1088 0 1089 0 1090 0 1091 1090 1092 0 1093 1092 1094 546 1095 0 1096 0 1097 1096 1098 0 1099 0 1100 0 1101 0 1102 0 1103 1102 1104 0 1105 0 1106 0 1107 0 1108 0 1109 1108 1110 0 1111 0 1112 0 1113 0 1114 556 1115 0 1116 0 1117 1116 1118 0 1119 0 1120 0 1121 0 1122 0 1123 1122 1124 0 1125 0 1126 562 1127 0 1128 0 1129 1128 1130 0 1131 0 1132 0 1133 0 1134 0 1135 0 1136 0 1137 0 1138 568 1139 0 1140 0 1141 0 1142 570 1143 0 1144 0 1145 0 1146 0 1147 0 1148 0 1149 0 1150 0 1151 1150 1152 0 1153 1152 1154 576 1155 0 1156 0 1157 0 1158 0 1159 0 1160 0 1161 0 1162 0 1163 1162 1164 0 1165 0 1166 0 1167 0 1168 0 1169 0 1170 0 1171 1170 1172 0 1173 0 1174 586 1175 0 1176 0 1177 0 1178 0 1179 0 1180 0 1181 1180 1182 0 1183 0 1184 0 1185 0 1186 592 1187 1186 1188 0 1189 0 1190 0 1191 0 1192 0 1193 1192 1194 0 1195 0 1196 0 1197 0 1198 598 1199 0 1200 0 1201 1200 1202 600 1203 0 1204 0 1205 0 1206 0 1207 0 1208 0 1209 0 1210 0 1211 0 1212 0 1213 1212 1214 606 1215 0 1216 0 1217 1216 1218 0 1219 0 1220 0 1221 0 1222 0 1223 1222 In other words, it does not halt at an assertion checkpoint! And the conjecture appears true. Questions. How can we prove this? Do you already have a proof in mind? Is this a new conjecture?

论坛问题帖 | has_answers | TotoposDiagramChaseApp | 2024-02-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019341 | MSE-4857166

Conjecture: For any prime n, at least one of $2n^2-1$, $2n^2+1$, or $\sqrt{2n^2-1}$ is prime.

In my spare time, I came up with this conjecture: For any prime n, at least one of $2n^2-1$, $2n^2+1$, or $\sqrt{2n^2-1}$ is prime. Examples: \begin{alignat}{2} 2 &\to\quad 2\cdot 2^2-1 &&\quad=7 \\ 3 &\to\quad 2\cdot 3^2-1 &&\quad=17 \\ 5 &\to\quad \sqrt{2\cdot 5^2-1} &&\quad=7 \\ 7 &\to\quad 2\cdot 7^2-1 &&\quad=97 \\ 11 &\to\quad 2\cdot 11^2-1 &&\quad=241 \\ 13 &\to\quad 2\cdot 13^2-1 &&\quad=337 \\ 17 &\to\quad 2\cdot 17^2-1 &&\quad=577 \end{alignat} Can somebody prove this?

论坛问题帖 | has_answers | user1286521 | 2024-02-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019342 | MSE-4858546

True or false: In a 2D Poisson process, for every point P, there exists a convex 1000-gon with Poisson points as vertices, that contains only P.

I made a Desmos graph that generates 30 uniformly random black points in a disk, with the centre of the disk in red. I asked myself, "Can I always draw a convex quadrilateral with four of the random points as vertices, such that its interior contains the centre of the disk but does not contain any of the random points?" The answer seems to be yes, based on experimentation. Then I asked myself the same question, except with "quadrilateral" replaced by pentagon, hexagon, or heptagon, and the answer still seems to be always yes. So my question is: True or false: In a 2D Poisson process, for every point P, there exists a convex 1000-gon with Poisson points as vertices, whose interior contains P and none of the other random points. I suspect the answer is yes. By making the polygon very long and narrow, maybe we can always find enough points to serve as vertices. But I'm not sure if this is true. Context: Recently I have been investigating inherent properties of the $2D$ Poisson process, for example here, here and here.

论坛问题帖 | has_answers | Dan | 2024-02-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019343 | MSE-4860350

A conjecture on representing $\sum\limits_{k=0} ^m (-1)^ka^{m-k}b^k$ as sum of powers of $(a+b)$.

UPATE: I asked this question on MO here. I was solving problem 1.2.52 in "An introduction to the theory of numbers by by Ivan Niven, Herbert S. Zuckerman, Hugh L. Montgomery" Show that if $\gcd(a, b)$ = 1 and p is an odd prime, then $\displaystyle \gcd\left(a+b, \frac{a^p+b^p}{a+b}\right)=1$ or p. So I tried to represent $\displaystyle \sum_{k=0} ^m (-1)^ka^{m-k}b^k$ as sum of powers of $a+b$ For any even number m that is one less than a prime number . $$\sum_{k=0} ^2 (-1)^k a^{2-k}b^k = (a+b)^2 -3ab$$ $$\sum_{k=0} ^4 (-1)^k a^{4-k}b^k = (a+b)^4 -5(ab)(a+b)^2 +5(ab)^2$$ $$\sum_{k=0} ^6 (-1)^k a^{6-k}b^k = (a+b)^6 -7(ab)(a+b)^4 +7 \cdot 2(ab)^2(a+b)^2 -7(ab)^3$$ I noticed that $\displaystyle \sum_{k=0} ^m (-1)^ka^{m-k}b^k=(a+b)^m +(m+1)\left(\sum_{k=1}^{\frac m 2} g_m(k)(-(ab))^k (a+b)^{m-2k}\right)$ where $g_m(k): \mathbb{N} \to \mathbb{Z}$ (I was not able to find a way to determine of $g_m(k)$) I couldn't prove this pattern or find $g_m(k)$ or prove that $\displaystyle \sum_{k=0} ^2 (-1)^k a^{2-k}b^k $ can be represented as sum of powers of $a+b$ For any even number m that is one less than a prime number. So my question is: How to prove that for any $m=p-1$ such that p is prime number $\displaystyle \sum_{k=0} ^m (-1)^ka^{m-k}b^k=(a+b)^m +(m+1)\left(\sum_{k=1}^{\frac m 2} g_m(k)(-(ab))^k (a+b)^{m-2k}\right)$ and determine $g_m(k)$. I don't think it is possible represent $\displaystyle \sum_{k=0} ^m (-1)^ka^{m-k}b^k$ in the form $\displaystyle(a+b)^m +(m+1)\left(\sum_{k=1}^{\frac m 2} g_m(k)(-(ab))^k (a+b)^{m-2k}\right)$. I don't think this will work for any even number because I noticed that in any even rows m that is above a prime row p i.e ($p=m+1$) in this image There is a very interesting property $\dbinom{m}{k}+(-1)^{k+1} |p $ for example $253+1|11 , \ \ 15-1 | 7$ and so on but for any other even row this doesn't hold.

论坛问题帖 | no_answers_at_retrieval | pie | 2024-02-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019344 | MSE-4860726

On the fractional parts of the roots of the Alternating Harmonic Numbers

We define $$\bar{H}_x=\ln2+\cos(\pi x)\left(\psi(x)-\psi\left(\frac x2\right)-\frac1x-\ln2\right)$$As the xth Alternating Harmonic Number (test out a few values to see why). Let x_n be the nth largest zero of the Alternating Harmonic Numbers, show that $\lim_{n\rightarrow\infty}\{x_n\}=\eta$ converges (where $\{x\}$ denotes the fractional part of x) or equivalently

$\lim_{n\rightarrow\infty}(x_n-x_{n+1})=1$. This definition of the alternating harmonic number originates from my terribly formatted paper where I generalize sigma notation for non-integer upper and lower bounds. Graphically speaking, the conjecture (I called it so in my paper because I couldn't solve it) seems to be true and $\eta\approx-0.431$. Some interesting approximations are $\frac{7\pi}{51}$ and this seemingly random integral $$\int_0^1\frac{\bar{H}_x\bar{H}_{-x}\sin(\pi x)}{\pi x}dx\approx0.43141591$$Not only does it hold for three digits, but there is a decimal representation of pi mixed in, up to 5 accurate decimal digits. Anyway, using the reflection formula derived for $\bar{H}_x$: $$\bar{H}_x-\bar{H}_{2-x}=\pi\cot(\pi x)-\left(\frac1{1-x}+\frac1x-\frac1{2-x}\right)\cos(\pi x)$$ we get that the conjecture is equivalent to finding the limit of the fractional part of the roots of $$\bar{H}_x-\bar{H}_{2-x}+\ln2=\ln2+\pi\cot(\pi x)-\left(\frac1{1-x}+\frac1x-\frac1{2-x}\right)\cos(\pi x)$$I got stuck here.

论坛问题帖 | has_answers | Kamal Saleh | 2024-02-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019345 | MSE-4862230

(dis)proof of conjecture on square unit fractions

Consider a finite set S of positive integers, and define $q(S) = \sum_{s \in S}{1/s^2}$. Letting $\rho = \pi^2/6$, we have $q(S)$ in the ranges $[0, \rho - 1), [1, \rho)$. I conjecture that for every rational r within those ranges, $\exists S: q(S) = r$. Can anyone prove or disprove this? Note that for plain unit fractions $u(S) = \sum_{s \in S}{1/s}$ it is straightforward to show by a greedy algorithm that $\exists S: u(S) = r \forall r \in \mathbb{Q}^+$; conversely for prime unit fractions $p(S) = \sum_{s \in S}{1/p_s}$ almost all rationals are unreachable, since for each S, $p(S)$ has a unique (squarefree) denominator. Given that the primes are denser than the squares, it would thus be a somewhat interesting result if the conjecture were to be proven. Computationally I have established existence of S for all rationals with denominators up to 50 within the range; see results on github. Update: added "finite" to clarify first sentence; corrected $\mathbb{R}$ to $\mathbb{Q}$.

论坛问题帖 | has_answers | Hugo van der Sanden | 2024-02-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019346 | MSE-4874949

For any square-free $n \geq 1$ and $a \in \Bbb{Z}/n$ including $\gcd(a,n) \neq 1$, then in the list $a, a^2, a^3, \dots$ either a or a^2 repeats?

For example, modulo 30 we have that $\gcd(5, 30) = 5$, but $5, 5^2, 5^3 = 5, 5^2, \dots$ goes the list, so both repeat. We know it's true when $\gcd(a,n) = 1$ because a cyclic group is formed in $\Bbb{Z}/n^{\times}$, s it's sufficient to prove the case when $\gcd(a, n) = d \neq 1$. Another example: $$ n = 15:\\ 3, 3^2, 3^3=-3,-3^2,(-3)^2=3^2, \dots \\ 6,6,6, \dots \\ 9,9^2=6,9, \dots\\ 5,10,5,10, \dots\\ $$ Is this hopelessly difficult to prove?

论坛问题帖 | has_answers | TotoposDiagramChaseApp | 2024-03-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019347 | MSE-4877902

Are there infinitely many primes that are a highly composite number ± 1?

I've looked at some highly composite numbers and realized that a lot of them are almost primes, i.e. differ only by 1 to the next closest prime. Here's a short list (I made) of highly composite numbers and whether they are almost prime: $$ 1 \text{, true $(+1)$} \\ 2 \text{, true $(+1)$} \\ 4 \text{, true (± 1)} \\ 6 \text{, true (± 1)} \\ 12 \text{, true (± 1)} \\ 24 \text{, true (-1)} \\ 36 \text{, true $(+1)$} \\ 48 \text{, true (-1)} \\ 60 \text{, true (± 1)} \\ 120 \text{, false} \\ 180 \text{, true (± 1)} \\ 240 \text{, true (± 1)} \\ \dots $$ I've thought about it, and it's actually not so surprising that many highly composite numbers (which I've checked) are almost prime. Since $n \pm 1$ won't share any prime divisors with n, it's likely for $n \pm 1$ to be prime if n has a high number of divisors. But are there infinitely many highly composite numbers that are ± 1 a prime? I don't know how I would prove something like this-I'm not even sure if this conjecture really holds. Also, if there are infinitely many such numbers, do they get rarer? And at what rate?

论坛问题帖 | no_answers_at_retrieval | SchellerSchatten | 2024-03-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019348 | MSE-4879782

Conjecture about Proving Primality of Fermat numbers by Elliptic Curves technic

In 2008 and 2009, Denomme-Savin and Tsumura provided 2 papers providing a Primality Test for Fermat numbers based on Elliptic Curves technic: $$ \text{Let } DST(x)= \frac{\displaystyle x^4+2x^2+1}{\displaystyle 4(x^3-x)} \ \text{ and } \ F_n = 2^{2^n}+1 $$ $$ \text{Let } x_1=F_1=\pm 5, \ x_{j+1} = DST(x_j)$$ $$ \text{If } x_{2^{n-1}} \equiv \mp 1 \pmod{F_n} \ , \ \text{then } F_n \text{ is prime.}$$ Now, can you prove the following conjecture (which makes use of a BigTree of the DiGraph): $$\text{Let } x_1 = \pm 4 \ , \ x_{j+1} = DST(x_j) $$ $$\text{If } x_{2^{n-1}} \equiv 0 \pmod{F_n} \ , \ \text{then } F_n \text{ is prime}.$$ And, then, can you prove the following conjecture (which makes use of a Cycle of the DiGraph): $$\text{Let } x_1 = \pm 1/3 \ , \ x_{j+1} = DST(x_j) $$ $$\text{If } x_{2^{n}-1} \equiv x_1 \pmod{F_n} \ , \ \text{then } F_n \text{ is prime}.$$ R. Denomme and G. Savin, “Elliptic curve primality tests for fermat and related primes,” Journal of Number Theory, vol. 128, no. 1, pp. 2398-2412, 2008. Y. Tsumura, “Elliptic curve primality tests for fermat and related primes” 2009.

论坛问题帖 | no_answers_at_retrieval | Tony Reix | 2024-03-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019349 | MSE-4881474

An extension of Brahmagupta's theorem.

As we know, there are conjectures that are easy to formulate but difficult to prove, and there are conjectures that are easy to prove but difficult to conceive. This conjecture is simple to conceive and probably has a simple proof, however I could not find any reference to it on the Internet so I quote it here: In the image below, the cyclic crossed quadrilateral A, B, C, D is orthodiagonal, in the sense that the lines passing through the vertices A, C and B, D intersect each other perpendicularly at a point M outside the circle. Prove that the perpendicular to a

side from the point M always bisects the opposite side (the intersecting sides AD and BC are considered opposite each other). Brahmagupta's theorem: https://en.m.wikipedia.org/wiki/Brahmagupta_theorem In Geogebra, the transition from Brahmagupta's theorem to the above conjecture is done naturally by moving the yellow control point in and out of the circle, respectively: https://www.geogebra.org/m/wthe5qt2

论坛问题帖 | has_answers | George Plousos | 2024-03-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019350 | MSE-4887134

Is $N=3\cdot 2^k+1$ prime if and only if $2^{N-1}\equiv 1 \pmod N$?

Is the following statement true? Let $k\geq 1$ be an integer and $N=3\cdot 2^k+1$. Then N is a prime prime if and only if $2^{N-1}\equiv 1 \pmod {N}$ One implication is simply a Fermat's Little Theorem, so the difficult part is proving or disproving that $2^{N-1}\equiv 1 \pmod {N}$ implies N is a prime. Or in other words showing whether there is a k such that N is a Fermat pseudoprime to base 2 (Poulet Number). I have checked the claim holds for $k\leq 15000$. As for the proof, we know complete factorization of $N-1$ so I was thinking of Lucas primality test, for which it would be sufficient to show that if $2^{N-1}\equiv 1 \pmod {N}$, then $2^{\frac{N-1}{3}}\not\equiv 1 \pmod {N}$ and $2^{\frac{N-1}{2}}\not\equiv 1 \pmod {N}$. Then I thought about Poclikngton primality test, but it requires a prime $p\mid N-1$ with $p>\sqrt{N}-1$ which we don't have (as $p=2,3$). As noted by Peter in comments, the last requirement can be removed in which case it's sufficient to prove $\gcd(2^{\frac{N-1}{3}}-1,N)=\gcd(2^{\frac{N-1}{2}}-1,N)=1$, however that seems stronger than what suffices for Lucas' test. I don't expect this result to be anything new or profound, I just have not found/proven it yet... Then again, I am not way too familiar with primality tests. As from where this comes from, I just did a computer search to look for similar criterion as Lucas-Lehmer test but for numbers of form $3\cdot 2^k+1$ and it found a match which when solving the recurrence to a closed form gave the statement above.

论坛问题帖 | no_answers_at_retrieval | Sil | 2024-03-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019351 | MSE-4890478

If $\lim\limits_{x\to\infty}\frac{f(x+1)-f(x)}{g(x+1)-g(x)}=l$ what are sufficient conditions to make $\lim\limits_{x\to\infty}\frac{f(x)}{g(x)}=l$?

In this question it is proved in the answers that If $f :[0 , \infty) \to \mathbb{R}$ and $f $ is bounded on every (a,b) such that $a<b <\infty$, prove that $\lim\limits_{x \to \infty }(f(x+1) - f(x))=l$ implies $\lim\limits_{x \to \infty }\frac{f(x)}{x}=l$. This condition looks suspiciously similar to Stolz-Cesaro theorem see this for a proof of using Stolz-Cesaro theorem . My question is : If If $f,g :[0 , \infty) \to \mathbb{R}$ $\lim\limits_{x\to\infty}\frac{f(x+1)-f(x)}{g(x+1)-g(x)}=l$ what are sufficient conditions for f,g to make $\lim\limits_{x\to\infty}\frac{f(x)}{g(x)}=l$? Of course when f and $g $ are both differentiable functions on $\mathbb{R}$ and if $\lim\limits_{x \to \infty } g(x)= \infty$ this condition sufficient to make $\lim\limits_{x\to\infty}\frac{f(x)}{g(x)}=l$, but this hypothesis is too strong is there a weaker hypothesis. and I want to generalise this result. I conjectured this more general hypothesis : If f is bounded on every (a,b) such that $a<b <\infty$, g is a continuous increasing function on $\mathbb{R}$ then $\lim\limits_{x\to\infty}\frac{f(x)}{g(x)}=l$. I couldn't prove this result but I think it is true. My question is this conjecture true ? If it was true is there a weaker hypothesis ? If it wasn't true, Is there a weaker hypothesis than f and $g $ are both differentiable functions on $\mathbb{R}$ and]$\lim\limits_{x \to \infty } g(x)= \infty$?

论坛问题帖 | no_answers_at_retrieval | pie | 2024-03-31

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019352 | MSE-4891208

Closed form for this generalisation of the gamma function. $f(x+1)=f(x)g(x+1) $

Just for curiosity I want to generalise the Pi function i.e $f(x+1) = f(x)g(x+1)$ for some differentiable function, I know this function probably has no closed form for general functions g as I couldn't find any but maybe it exist and more experienced user can find it. For a differentiable function g on $\mathbb{R}^+$ and $\exists M \in \mathbb{R}$ st $\forall x>M$ $g(x)\ge0$ with $g(1)\ne 0$ define $f(x+1) = f(x)g(x+1)$ To make it clear that f depends on x and g I will write it as $\Pi(x, g(t))$, $\Pi(x, t)$ is just the ordinary pi function. . I found analogous result to the Gauss product by a non rigorous approach. $$\Pi(x, g(t)) = \lim_{n \to \infty } g(n)^x \prod\limits_{1\le k\le n} \frac{g(k)}{g(k+x)} $$ With this It is easy to see that $\Pi(x, t^a) =\Pi(x, t)^a $, $\Pi(x, e^t)=1 $. As I said before, probably there is not a closed form for a general function g put if a closed form could be found when g is any polynomial satisfying the conditions of g this could be extended to power series which cover a wide range of functions.

论坛问题帖 | has_answers | pie | 2024-04-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019353 | MSE-4891945

How do you find constants in an k-Tuple conjectures?

By introducing modular objects associated to the sequel of rings $$(Z/2,Z/6,Z/30,Z/210,Z/2310,..., Z/p_n\#Z)$$ a sequence of coefficients is updated$$(2;\color{green}{\frac83}; 3.2;...$$ (see my answer to my proper question here) I've been running computers and I've observed that There are twice more primes p such that $p+6$ is also prime (sexys) than twins; $\color{green}{\frac83}$ more primes p such that $p+30$ is also prime than twins; 3.2 more p such that $p+210$ is also prime than twins; ... Until now, I had always had the greatest difficulty in deciphering how to find these constants in Wolphram Alpha's article about k-Tuples conjectures. For example, for $p+$30, where I'm looking for $\color{green}{\frac83}$, I understand that I am interested in what the article notes $$\boxed{C(15)=2\prod_{q\text{ prime}}^{}\frac{q(q-2)}{(q-1)^2}\prod_{q|15}^{}\frac{q-1}{q-2}}$$ In fact, the initial purpose of my post was even to ask for help to do so but I think I'm starting to understand since $$\prod_{q|15}^{}\frac{q-1}{q-2}=\frac{3-1}{3-2}\frac{5-1}{5-2}=2\times\frac{4}{3}=\color{green}{\frac{8}{3}}$$ 1.- I still have to think about it a little more, maybe get your help but it will do it. 2.- But, suddenly, a more interesting question arises for me: is this approach in modular arithmetic to highlight the Hardy and Littlewood coefficients, but

especially to give them a non-statistical modular meaning (see here) , known? Definitions, notations and examples: $p_1=2, p_2=3, p_3=5, p_4=7, p_5=11, ... , $ are the primes; $\forall n\in \mathbb N^*, p_n\#:=p_1\times p_2\times ....\times p_n$; examples : $p_1\#=2, p_2\#=p_1\times p_2=2\times 3=6, p_3\#=2.3.5=30, ...$

论坛问题帖 | no_answers_at_retrieval | Stéphane Jaouen | 2024-04-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019354 | MSE-4893052

Conjecture: Two different random triangles (both based on random points on a circle) have the same distribution of side length ratios.

On a circle, choose three uniformly random points A,B,C. Triangle T_1 has vertices A,B,C. The side lengths of T_1 are, in random order, a,b,c. Triangle T_2 is formed by drawing tangents to the circle at A,B,C. (T_2 may or may not lie inside the circle.) The side lengths of T_2 are, in random order, d,e,f. Is the following conjecture true: The distribution of ratios of a,b,c is the same as the distribution of ratios of d,e,f. So for example, if my conjecture is true, then we would have the following probability equations: $P(a^2+b^2<c^2)=P(d^2+e^2<f^2)$ $P(a^3+b^3<c^3)=P(d^3+e^3<f^3)$ $P(ab<c^2)=P(de<f^2)$ $P\left(\frac1a+\frac1b<\frac1c\right)=P\left(\frac1d+\frac1e<\frac1f\right)$ (My conjecture, if true, would not imply that $P(ab<c)=P(de<f)$, because these inequalities are not dimensionally homogeneous.) Basis for my conjecture I found that $P(ab<c^2)=\frac35$. Then I wondered what $P(de<f^2)$ equals, and a simulation suggests that it also equals $\frac35$. I found that $P\left(\frac1a+\frac1b<\frac1c\right)=\frac15$. Then I wondered what $P\left(\frac1d+\frac1e<\frac1f\right)$ equals, and a simulation suggests that it also equals $\frac15$. I tried some other probabilities, and simulations suggest that each probability has the same value for the two triangles. So I suspect that the two triangles have the same distribution of ratios of side lengths. But I don't know why.

论坛问题帖 | has_answers | Dan | 2024-04-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019355 | MSE-4895587

Consecutive numbers

Are there ever more consecutive composite numbers than there are primes up to that point? I imagine not, because the primes are the ones which cancel out multiples, so will inevitably have to fill in the gaps between multiples of them. Then again, the number of combinations possible grows faster than an exponential. It may be a matter of calculating how many products are within certain bounds. Additionally, I have that there is a known set of composite numbers, an infinite union of sets described by an arithmetic sequence. $\{9+6x\}\cup\{15+10x\}\cup\{21+14x\}\cup\{33+22x\}\cup\{39+26x\}\cup...$ which when removed from the odd numbers yields the prime numbers. Conversely, if we show that there is a point that many of these terms are lining up with common difference 2 we have shown some progress on the result.

论坛问题帖 | no_answers_at_retrieval | Talon Eaglefeathers | 2024-04-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019356 | MSE-4896790

What is the $\inf$ and $\sup$ of the area of quadrilateral given its sides length?

Now asked on MO here. Given the length of the sides of a quadrilateral a,b,c,d the area of the quadrilateral is less than or equal to $\frac{(a+b+c+d)^2}{16}$ i.e it is an upper bound of the area of any quadrilateral with side length a,b,c,d. A question that came to my mind is if $\frac{(a+b+c+d)^2}{16}$ is an upper bound Then the least upper bound exits so what is the infimum and supremum of the area of quadrilateral (0 is an oblivious lower bound so the infimum exists too)? I made a silly mistake thinking that $\frac{(a+b+c+d)^2}{16}$ is the supremum but it turns out I was wrong. If $a=b, \ c=d$ "a kite" or if $a=c, \ d=b$ "a parallelogram" then it is easy to prove the infimum is 0 and the supremum is just a^2 when it is a rhombus. The infimum depends on a,b,c,d in this examples the side lengths are $19,13,7,4$ and the minimum area is around 9.5 so it is not 0. Conjecture: If $a<b<c<d$ and the area of the quadrilateral is S then $\inf S $ is the area of the triangle $a,b,d-c$ which is $$\sqrt{\frac{a+b+d-c}{2}\cdot\frac{a-b+d-c}{2}\cdot\frac{-a+b+d-c}{2}\cdot\frac{a+b-d+c}{2}}$$ This also works when $a=b=c=d, \ \ \ a=b, \ c=d \ \ \ ,a=c, \ d=b $. I couldn't rigorously proof that claim but I am pretty sure it is correct. In the previous example when side lengths are $19,13,7,4$ the naximum area is around 73.5 but this time I have no Idea what kind of shape is that. How to determine the infimum and the supremum of the area given a,b,c,d.

论坛问题帖 | has_answers | pie | 2024-04-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019357 | MSE-4898900

The vertices of a hexagon are random points on a unit circle; a,b,c are the lengths of three random sides. Conjecture: $P(ab<c)=\frac35$.

The vertices of a hexagon are uniformly random points on a unit circle; a,b,c are the lengths of three distinct random sides. A simulation with 10^7 such random hexagons yielded a proportion of 0.60008 of them satisfying $ab<c$. Is the following conjecture true: $P(ab<c)=\dfrac35$. Probabilities involving the side lengths of random polygons inscribed in a circle are often simple rational numbers; for example here, here and here. Those examples involve triangles, which are determined by only two parameters (two random central angles, with the third angle determined by the others), so it was easy to set up an integral. But this question involves a hexagon, which involves five parameters (five random angles, with the sixth angle determined by the others), and I do not know how to set up an integral. Sometimes these kinds of questions can be answered without integration, using pure geometry, for example here. But I have been unable to find a purely geometric proof for this question. $P(ab<c)$ seems to be a simple rational number only for triangles and hexagons, among n-gons for $3\le n\le 12$, based on simulations. Distribution of a Here I derive $f(x)$, the density function of a, which is the same as the density function of b and c. However, I do not know if this is helpful, because a,b,c are not independent. Suppose the first chosen random point on the circle is O, and a is the length of the side immediately anticlockwise from O. $a>x$ if all the other chosen points are on the major

arc subtended by the red line segment, and the other chosen points are not all on the minor arc subtended by the green line segment. $\theta=\arcsin \frac{x}{2}$ $\begin{align} P(a<x)&=1-P(a>x)\\ &=1-\left(\left(1-\frac{1}{\pi}\arcsin\frac{x}{2}\right)^5-\left(\frac{1}{\pi}\arcsin\frac{x}{2}\right)^5\right) \end{align}$ $\begin{align} f(x)&=\frac{d}{dx}P(a<x)\\ &=\frac{5}{\pi\sqrt{4-x^2}}\left(\left(1-\frac{1}{\pi}\arcsin\frac{x}{2}\right)^4+\left(\frac{1}{\pi}\arcsin\frac{x}{2}\right)^4\right)\\ \end{align}$ I checked that $\int_0^2f(x)\mathrm dx=1$.

论坛问题帖 | has_answers | Dan | 2024-04-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019358 | MSE-4900158

The vertices of a pentagram are five random points on a circle. Conjecture: The probability that the pentagram contains the circle's centre is $3/8$.

The vertices of a pentagram are five uniformly random points on a circle. Is the following conjecture true: The probability that the pentagram contains the circle's centre is $\frac38$. (The pentagram is said to contain the circle's centre if the central pentagon, or any of the five triangles adjecent to the central pentagon, contains the circle's centre.) A simulation with 10^7 such random pentagrams yielded a proportion of $0.3750079\approx1.00002\times\frac38$ containing the circle's centre. Thus, my conjecture. How to set up a simulation Let the circle be $x^2+y^2=1$ with centre $O\space(0,0)$. Assume the first random point is $(1,0)$ and let the other four random points be, going anticlockwise from $(1,0)$: $(\cos\theta_1,\sin\theta_1)$, $(\cos\theta_2,\sin\theta_2)$, $(\cos\theta_3,\sin\theta_3)$, $(\cos\theta_4,\sin\theta_4)$. Let A,B,C,D,E be the five regions in the circle and outside the pentagram, going anticlockwise starting with the region between $(1,0)$ and $(\cos\theta_1,\sin\theta_1)$. O lies in A if and only if $\theta_2>\pi,$ and $\theta_4-\theta_1<\pi$. O lies in B if and only if $\theta_2>\pi,$ and $\theta_3-\theta_1>\pi$. O lies in C if and only if $\theta_3-\theta_1>\pi$, and $\theta_4-\theta_2>\pi$. O lies in D if and only if $\theta_3<\pi,$ and $\theta_4-\theta_2>\pi$. O lies in E if and only if $\theta_3<\pi,$ and $\theta_4-\theta_1<\pi$. The pentagram contains O just if O lies in none of A,B,C,D,E. If my conjecture is true, then, given the simplicity of the probability, there might be a proof based on some kind of symmetry.

论坛问题帖 | has_answers | Dan | 2024-04-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019359 | MSE-4900413

What is $\sup\{a\}$ such that for continuous non differentiable function f $\lim\limits_{h\to0}\frac{f(x+h)-f(x)}{h^a}$ exist?

Now asked on MO here The definition of the derivative of a function f at a point x is : $\lim\limits_{h \to 0 }\frac{f(x+h) -f(x)}{h}$ but what if we change that h to be any continuous function $g(h)$ that goes to 0 as h goes to 0? Lets focus our attention at powers of h so $g(h)=h^a$ for some a. If f is a differentiable function at x any function other than h the absolute value of the derivative will go to either 0 or ∞. I will refer to $\lim\limits_{h \to 0 }\frac{f(x+h) -f(x)}{g(h)}$ as $a-$derivative. A question that came to my mind Given any continuous function nowhere differentiable on $\mathbb{R}$ is there a g such that the absolute value of the $a-$derivative" exist almost everywhere and how to determine such functions ? The case where $g(x)$ is big enough to make this $a-$derivative 0 is annoying because one can say a very big number like $a=1/TREE(3) $ and say, for example, at this number the $a-$derivative exist and it is 0 for the Weierstrass function, so let me clear that : Given any continuous function nowhere differentiable what is the $\sup g(x)$ such that $\lim\limits_{h \to 0 }\frac{f(x+h) -f(x)}{g(h)}= 0$ for all x where the $\sup $ is taken of all real a does the absolute value of the $a-$derivative exist at that value ? Lets take L. van der Waerden's function for example Define $g(x)= |x|$ for $|x|\in [-1,1]$, $g(x+2)=g(x)$ $$f(x)= \sum_{n \ge 1} \frac{3^n g\left(4^n x\right) }{4^n}$$ Is an example of continuous function nowhere differentiable, There is a proof that the absolute value secant line's slope goes to infinity as it approaches the point so this made me wonder will the absolute value of the $a-$derivative exist if we decrease the power of h?

论坛问题帖 | no_answers_at_retrieval | pie | 2024-04-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019360 | MSE-4902760

A probability involving side lengths of a random triangle on a disk: Is it really $\frac37$?

Choose three uniformly random points on a disk, and let them be the vertices of a triangle. Call the side lengths, in random order, a,b,c. What is $P(a^2<bc)$? A simulation with 10^7 such random triangles yielded a proportion of $0.4285833\approx1.00003\times\frac{3}{7}$ satisfying $a^2<bc$, leading me to believe that the probability is $\frac37$. I find this alleged probability interesting because it seems that the number 7 rarely appears in the answers to natural geometry or probability questions. Context Recently I have learned that some probabilities related to circles, of the form $P(x^2<yz)$, have simple rational values (for example, a probability about a random triangle inscribed in a circle, and a probability about a random pentagram inscribed in a circle). So I wondered if there is a probability like this related to a disk. I may have found one. My attempt Let the boundary of the disk be $x^2+y^2=1$. Using disk point picking, let the three points be $C\left(\sqrt{r_1}\cos\theta_1,\sqrt{r_1}\sin\theta_1\right)$ $B\left(\sqrt{r_2}\cos\theta_2,\sqrt{r_2}\sin\theta_2\right)$ $A\left(\sqrt{r_3},0\right)$ where each r is a uniformly random real number from 0 to 1, and each θ is a uniformly random real number from 0 to 2π. Let $a=BC$, $b=AC$, $c=AB$. This gives: $$P(a^2<bc)=P\left(r_1+r_2-2\sqrt{r_1r_2}\cos(\theta_1-\theta_2)<\sqrt{\left(r_1-2\sqrt{r_1r_3}\cos\theta_1+r_3\right)\left(r_2-2\sqrt{r_2r_3}\cos\theta_2+r_3\right)}\right)$$ I do not know how to set up an integral, nor any other way to calculate the probability. Apparently, after the dust settles, we should be left with $\frac37$.

论坛问题帖 | has_answers | Dan | 2024-04-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019361 | MSE-4903514

For any positive integer $n\geq5$, there is at least one integer $m>n$ such that $ \pi\left(nm\right)=n+m $

Conjecture: For any positive integer $n\geq5$, there is at least one integer $m>n$ such that $ \pi\left(nm\right)=n+m $, where $\pi\left(x\right)$ is the prime counting function. Examples: $$ \pi(5\times9)=5+9=14 $$ $$ \pi(6\times7)=6+7=13 $$ $$ \pi(7\times368)=7+368=375 $$ $$ \pi(8\times998)=8+998=1006 $$ $$ \pi(9\times2551)=9+2551=2560 $$ I'm looking for either a proof of the conjecture, or at least an heuristic argument for big enough n. Computations: The nbr column denotes the number of solutions, while the min and max columns represent the first and last solutions. n nbr min max 1 0 2 0 3 0 4 0 5 22 9 41 6 9 7 148 7 14 368 409 8 1 998 998 9 12 2551 2597 10 6 6331 6424 11 7 15866 15889 12 25 39860 39970 13 20 100135 100192 14 9 251616 251637 15 19 636787 637063 16 2 1617099 1617110 17 13 4124188 4124272 18 9 10553076 10553278 19 13 27066256 27066824 20 28 69709243 69709557 21 17 179992154 179992524 22 50 465769460 465769748 23 13 1208198239 1208198402 24 43 3140421185 3140421406 25 20 8179001679 8179001739 26 8 21338684878 21338685037 27 68 55762147707 55762148901 28 1 145935688930 145935688930 29 95 382465571795 382465572544 Graph: Here is a graph of $\pi(nm)-(n+m)$ for $n=12$: And here is the same graph, but zoomed on the interval where it crosses the 0 axis: In this example, $n=12$, the line touches 25 times the 0 axis, and we can expect that as n gets bigger, the interval in which the line is close to 0 will also become larger.

论坛问题帖 | has_answers | François Huppé | 2024-04-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019362 | MSE-4906573

A continuous nowhere differentiable function with $\alpha-$derivative exists at $\sup\{a\}$.

In This question I asked: Define $g(x)= |x|$ for $|x|\in [-1,1]$, $g(x+2)=g(x)$ $$f(x)= \sum_{n \ge 1} \frac{3^n g\left(4^n x\right) }{4^n}$$ what is the $\sup \{\alpha\}$ such that $$ \lim\limits_{h \to 0 }\frac{f(x+h) -f(x)}{h^\alpha}= 0 $$ for all x? Does the absolute value of the $\alpha-$derivative exist at that value of α ? I expected that the absolute value of the $\alpha -$derivative exist at the $\sup \{\alpha\}$ but it turns out I was wrong as this answers shows (the supremum is $\frac{\ln\left(\frac 4 3\right)}{\ln 4}$ btw) . My question is: Is there a continuous nowhere differentiable function with the $\alpha-$derivative exist at that value of $\sup \{\alpha\}$ where is the sup is taken on all reals such that $ \lim\limits_{h \to 0 }\frac{f(x+h) -f(x)}{h^\alpha}= 0 $ for all x? If it is impossible to have such function how to prove that?

论坛问题帖 | has_answers | pie | 2024-04-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019363 | MSE-4915763

Conjectured connection between e and π in a semidisk.

A semidisk with diameter $\dfrac{e}{\pi}n$ is divided into n regions of equal area by line segments from a diameter endpoint. Here is an example with $n=6$. Consider the n arcs between neighboring line segment endpoints. Let $P(n)=\text{product of arc lengths}$. Is the following conjecture true: Conjecture: $\lim\limits_{n\to\infty}P(n)=\dfrac{\pi}{2}$ In other words, if the average arc length is $\color{red}{\dfrac{e}{2}}$ then the product of arc lengths approaches $\color{red}{\dfrac{\pi}{2}}$. Evidence for my conjecture I got the following approximations. $P(1)\approx 1.359$ $P(2)\approx 1.438$ $P(3)\approx 1.469$ $P(6)\approx 1.507$ $P(12)\approx 1.528$ $P(24)\approx 1.543$ $P(48)\approx 1.552$ $P(96)\approx 1.558$ It seems that if the average arc length is less than $\dfrac{e}{2}$ then the product approaches 0, and if the average arc length is greater than $\dfrac{e}{2}$ then the product approaches ∞. What makes this difficult What makes my conjecture difficult for me to prove or disprove, is that I cannot find exact expressions for the arc lengths. For example, in the example shown above with $n=6$, the length of the longest arc is $\dfrac{3ex}{\pi}$ where $x-\sin x=\frac{\pi}{6}$. I am aware of Kepler's equation, but that doesn't seem to help. Related questions This question is essentially the converse of my question, "Product of areas in a disk". I asked about the product of another kind of arc length related to a circle, in my question, "Another interesting property of $y=2^{n-1}\prod_{k=0}^n \left(x-\cos{\frac{k\pi}{n}}\right)$: product of arc lengths converges, but to what?". I recently asked a related question, "Product of lengths in a disk of area π/e".

论坛问题帖 | has_answers | Dan | 2024-05-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019364 | MSE-4922512

Convergence of sequences formed by orthocenters, incenters, and centroids in repeated triangle constructions

Now asked on MO here. Given a scalene triangle $A_1B_1C_1$, construct a triangle $A_{n+1}B_{n+1}C_{n+1}$ from the triangle $A_nB_nC_n$ where A_{n+1} is the orthocenter of $A_nB_nC_n$, B_{n+1} is the incenter of $A_nB_nC_n$ and C_{n+1} is the centroid of $A_nB_nC_n$, My question is If this process is repeated indefinitely would the sequences $\{A_n\}$, $\{B_n\}$, $\{C_n\} $ converge ? There are only four possible scenarios: The points will converge to a point. The points will eventually stuck on a loop. The points will completely diverge. The points will eventually make an isosceles triangle (or equilateral triangle) which will end the sequence. All of these scenarios would depend on the initial triangle. So given $\angle A_1$, $\angle B_1$, $\angle C_1, $ how to determine which of the four scenario would happen to the sequences $\{A_n\}$, $\{B_n\}$, $\{C_n\} $? I tried to draw the first few triangles and see if the point will converge or not. Graphing the first 25 triangles suggests that the sequence converge. Here is the graphing the first 30 triangle, If anyone needs them. If this is true that the three sequences will converge to a point this point will be unique to the triangle, I wonder what extra properties this point have. It seems that the limit point (assuming it exists) in not in the Encyclopedia of Triangle Centers, the distance between $A_{25} $ and $\overline {B_1C_1}$ (in the triangle 6-9-13) is

0.5360763749467, the distance between $B_{25} $ and $\overline {B_1C_1}$ is 0.5360798890437 and the distance between $C_{25} $ and $\overline {B_1C_1}$ is 0.5360789119326 so the distance between limit point and $\overline {B_1C_1}$ is around 0.53607

论坛问题帖 | no_answers_at_retrieval | pie | 2024-05-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019365 | MSE-4923238

Graph isomorphism checking/detection for directed acyclic graphs

The graph isomorphism problem is hard for an arbitrary graph, and certain classes of graphs have been proven to be "GI-complete", which as I understand means they can be reformulated in terms of a polynomial number of dual problems which themselves take polynomial time to solve. From wikipedia: In computational complexity theory, a polynomial-time reduction is a method for solving one problem using another. One shows that if a hypothetical subroutine solving the second problem exists, then the first problem can be solved by transforming or reducing it to inputs for the second problem and calling the subroutine one or more times. If both the time required to transform the first problem to the second, and the number of times the subroutine is called is polynomial, then the first problem is polynomial-time reducible to the second. I am wondering about direct acyclic graphs (a.k.a. feed-forward neural networks) specifically not because I have a research interest in them but because I happen to be confronted with them while studying the NEAT algorithm (video: video), which is a genetic/evolutionary algorithm that searches the space of different neural network architectures to find the "fittest" one for a particular task. The NEAT algorithm owes its success to the efficient way in which this exploration takes place: by uniquely labelling each edge (essentially keeping a bijective map in storage mapping a number, the "innovation number", to each possible edge) the algorithm ensures we don't create the "same" graph a second time. There is also this other thing called speciation, which is a measure of how different two graphs are based on the number of extra edges one graph has compared to another, the number of edges that appear in exactly one and not two of the graphs, and the average edge weight difference taken over all shared edges in the two graphs. But it seems that the NEAT algorithm does not address the possibility of graphs being isomorphic. Speciation looks at the "genomes" of the two graphs. BTW computer scientists use biology vocabulary, but: gene: edge gene: edge encoding (of the form [innovation number, (v1, v2), enabled_bool, edge_weight]) node gene: similar but node info genome: set of genes. Encodes a network. Any network produced by neat should have a unique genome phenotype: the actual graph (or network) produced by the genome Just wanted to get that out of the way. So, genomes are the thing that's used to manipulate candidate networks, and the measure of how different two graphs are is a function of their genomes. But my concern is that you could have two graphs that look very similar but with completely different genomes, or two genomes that are close with respect to the speciation formula, but whose phenotypes that are completely different. The type II error here would be to have very similar graphs (topologically speaking) but with a large speciation difference measure, leading to wasted compute resources. This may seem trivial, and I will look like an idiot if someone posts a small 3 node counter example in a comment, but if one wants to check that two DAGs (directed acyclic graphs) are not isomorphic isn't it enough to do the following: (Let $d^{(v)}_{in}$ and $d^{(v)}_{out}$ be the in-degree and out-degree of a given node v.) For each graph compute the set $\{(d^{(v)}_{in},d^{(v)}_{out})\ \ |\ \ v\in V\}$. If the two sets are the same then the graphs are isomorphic, and if they are not the same, the graphs are not isomorphic. I tried some examples by hand and it really seems like this degree-pairs set criterion is plausible.

论坛问题帖 | has_answers | John Cataldo | 2024-05-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019366 | MSE-4923419

Is there a function ϕ such that $\int_0^t\int_0^t \phi(u,v) dudv=\int_0^t\psi(s)ds$?

Let t be a positive real number and ψ a positive function. I am looking for a function ϕ such that: $$\int_0^t \int_0^t \phi(u,v) du\,dv=\int_0^t\psi(s)ds$$ Does such a function exist or not?

论坛问题帖 | has_answers | user1326164 | 2024-05-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019367 | MSE-4923435

Application of Bertrand's postulate

We can use Bertrand's Conjecture (that for any integer $n \not= 0$, there exists at least one prime number p with $n < p \leq 2n$) to demonstrate the following proposition: $\forall n \in \mathbb{N}\setminus\left\{0\right\}$, the rational number such that: $$ H_{2n} = \sum_{k = 1}^{2n}\frac{1}{k}\quad \mbox{is}\ \underline{never}\ \mbox{an integer} $$ What is the demonstration ?.

论坛问题帖 | has_answers | Alexis J | 2024-05-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019368 | MSE-4923678

is there a function ϕ such that $\int_R \int_R f(t)g(s)\phi(t,s)dsdt=\int_Rf(t)g(t)\rho(t)dt $?

Let ρ be a positive function, is there exist a function ϕ such that for any f,g: $$\int_R \int_R f(t)g(s)\phi(t,s)dsdt=\int_Rf(t)g(t)\rho(t)dt $$ of course both integrals are supposed finite here.

论坛问题帖 | no_answers_at_retrieval | user1326164 | 2024-05-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019369 | MSE-4924016

is $\int_0^t \int_0^s \delta(u - v) \, du \, dv = \min(t, s) $?

Is the following integral equation true? $ \int_0^t \int_0^s \delta(u - v) \, du \, dv = \min(t, s) $ where δ is the Dirac delta function.

论坛问题帖 | no_answers_at_retrieval | user1326164 | 2024-05-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019370 | MSE-4927378

Weaker than $X^2 + 1$ conjecture: $(a + bX)^2 + 1$ is a prime number at least once for any $\gcd(a,b) = 1$.

Let $\gcd(a,b) = 1$ where $a, b\in \Bbb{Z}$ and consider $f(X) = X^2 + 1$ and the forward image of the coset $a + b\Bbb{Z}$ under f. There is always at least one prime number in $f(a + b\Bbb{Z})$. Can we prove this using elementary methods?

论坛问题帖 | has_answers | TotoposDiagramChaseApp | 2024-06-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019371 | MSE-4929839

These constrained alternating series always satisfy an inequality.

Let $x_0 \in \Bbb{N}$ and suppose that $x_1 \lt \frac{x_0}{2}$, while $0 = x_n \leq \dots \leq x_3 \leq x_2\leq x_1$. Then is it possible that: $$ \sum_{i = 0}^n (-1)^i x_i \gt 1 $$ no matter what choices are made for $x_1, \dots x_{n-1}$ in $[0, \dots, \frac{x_0}{2}) \cap \Bbb{N}$? Note that keeping the variables $x_i \in \Bbb{N}$ instead of $\Bbb{R}$ has the advantage of having only a finite number of cases to check for each $x_0 \in \Bbb{N}$. For example: $$ x_0= 4, x_1 = 1, x_2=1, x_3=1, x_4 = 0 \\ 4 - 1 + 1 -1 + 0 = 3 \gt 1 \\ 4 - 1 + 1 - 0 + 0 = 4 \gt 1 \\ 4 - 1 + 0 - 0 + 0 = 3 \gt 1 \\ 4 - 0 + 0 - 0+0 = 4 \gt 1 \\ $$ And the reason we could start at 1 on the next term is because 1 is the largest integer $\lt \frac{4}{2} = 2$. Those are all the cases to check for $x_0 = 4$. [U+1F913] I'm wondering if in general this conjecture is true.

论坛问题帖 | has_answers | TotoposDiagramChaseApp | 2024-06-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019372 | MSE-4930407

Constructing a Curve from a Differentiable Function's Normal Lines at a Fixed Distance.

I've been experimenting with a graphing calculator and stumbled upon an interesting geometric construction involving functions and their normal lines. Here's the problem I'm trying to solve: Let's consider a differentiable function $ f$ defined on a closed interval. For a given positive real number $ r > 0 $, we perform the following construction for every point $ (x, f(x)) $ on the graph of $ f $: List item Draw the normal line to the graph of $ f $ at the point $ (x, f(x)) $. From the points on this normal line, select the one that is a distance $ r $ away from $ (x, f(x)) $ and has a $ y $-coordinate greater than $ f(x) $. If there are two such points with equal $ y $-coordinates, choose the one with the smaller $ x $-coordinate. The result of this process is a new curve, which I'll denote as $ F $. This curve is not necessarily a function. For instance, consider the function $ f(x) = (x-1)(x+2)(2x-1) $ with $ r = 0.1 $. The resulting curve from the above mapping is shown below: Additionally, here's an animation showing the function $ f(x) = (x-1)(x+2)(2x-1) $ with $ r $ varying between $ 0 $ and $ 2 $, and the corresponding curve $ F $: My question is: How can we mathematically determine the curve $ F $ given the function $ f $ and the distance $ r $? This problem appears to be quite challenging, especially for ugly functions like the antiderivative of the Weierstrass function. Any insights or methods to approach this would be greatly appreciated.

论坛问题帖 | no_answers_at_retrieval | pie | 2024-06-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019373 | MSE-4932367

Implications of having access to the Busy Beaver oracle

Apologies if I'm asking a naïve question as I've only recently learned about the concept. What would be the practical implications (if any) of having access to a magical black box providing the values of $\Sigma(n)$? Of course, this would readily imply access to the "answers" to long-standing open problems in math, but without proofs those probably wouldn't be particularly influential. So could we make any use at all of such an oracle? A follow-up question: Is there any relation between the values of $\Sigma(n)$ for different n which could allow one to check whether the oracle is providing self-consistent outputs?

论坛问题帖 | has_answers | mavzolej | 2024-06-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019374 | MSE-4932752

Finding measurable subsets of any given value?

I'm not sure if the following is true, but I would hope it is with the regularity properties of μ. Let X be a locally compact Hausdorff space with μ a nonzero Radon measure on X. Then given any $\alpha \in (0,\mu(X))$, there exists measurable $E \subset X$ with $\mu(E)=\alpha$. My idea was to define something like $\Lambda = \{A \subset X \: | \: \text{A measurable and $\mu(A) \leq \alpha$}\}$ and try to use sequences in Λ or even Zorn's lemma to get such a thing (Zorn's lemma seems harder to use as measurability only plays nice with countable unions). However I'm kind of stumped on how to actually go about showing this. I know that for $X = \mathbb{R}$ and $\mu=m$, you can do this trivially.

论坛问题帖 | has_answers | Isochron | 2024-06-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019375 | MSE-4933747

A conjecture about the sum of Faulhaber polynomials mod 2.

Context: In this question I asked about "Finding a polynomial function for alternating m numbers of odds and even numbers in sequences" I noticed that all Faulhaber polynomials are solutions for f_2 where $f_m(x)$ is apolynomial where $f_m(x)$ is odd for the first m natural numbers and $f_m(x)$ is even for the second m natural numbers and so on that made me do some experiments with $\sum\limits_{1\le k\le n} P_k(x)$. Let $P_n(x)$ be the $n-th$ Faulhaber polynomial let $Q_n(x)= \sum\limits_{1\le k\le n} P_k(x)$ let $r(n)\in \{0,1,2,3,\dots\}$ be the biggest number such that $\frac{Q_n(x)}{2^{r(n)}}$ is always integer when x is a natural number Now define $S_n(x)= \frac{Q_n(x)}{2^{r(n)}}$. I noticed that $S_n(x)$ when $x\in \mathbb{N}$ always result on a block of E even numbers followed by a block of O odd numbers. for example $S_1(x)=\frac{x(x+1)}{2}$ when $x \in \mathbb{N}$ the sequence would be the following $1,3,6,10,15,21,28,36$ so $E=O=2$ (i.e 2 evens followed by 2 odds and the cycle continue) The question is how to determine E,O for any S_n? Here is a table containing the first twelve of S_n (This is not a matrix): $\begin{bmatrix} & \text{Even(E)} &\text{Odd (O)} \\S_{1} & 2& 2 \\ S_{2}& 3 &1 \\ S_{3}& 2&2 \\ S_{4}& 4&4 \\ S_{5}& 2&2 \\ S_{6}& 3&1 \\ S_{7}& 2&2 \\ S_{8}& 4&4 \\ S_{9}& 2&2 \\ S_{10}& 3&1 \\ S_{11}& 2&2 \\ S_{12}& 4&4 \end{bmatrix}$ It seems that S_n will have a cycle $(2,2), \ (4,4), \ (2,2), \ (3,1)$ But I couldn't prove or disprove this, where (E,O) determine how many numbers even numbers and odd numbers appear in each cycle respectfully. It is easy to prove that for odd n, $(E,O)=(2,2)$ and $r(n)=0$. Also $r_{2n}=1$ for all even numbers that I tried, for some reason r_{2n} can't be greater than 1. Here is the first twelve Faulhaber polynomials: $$P_{1}(x)= \frac{x(x+1)}{2}$$ $$P_{2}(x)= \frac{x(2x+1)(x+1)}{6}$$ $$P_{3}(x)= \frac{x^2(x+1)^2}{4}$$ $$P_{4}(x)=\frac{x(2x+1)(x+1)}{6} \times \frac{3x^2+3x-1}{ 5}$$ $$P_{5}(x)= \frac{x^2(x+1)^2}{4} \times\frac{2x^2+2x-1}{ 3}$$ $$P_{6}(x)= \frac{x(2x+1)(x+1)}{6} \times \frac{3x^4+6x^3-3x+1}{ 7 }$$ $$P_{7}(x)= \frac{x^2(x+1)^2}{4 }\times\frac{3x^4+6x^3-x^2-4x+2}{ 6}$$ $$P_{8}(x)= \frac{x(2x+1)(x+1)}{6} \times \frac{5x^6+15x^5+5x^4-15x^3-x^2+9x-3}{ 15}$$ $$P_{9}(x)= \frac{x^2(x+1)^2}{4} \times\frac{(x^2+x-1)(2x^4 +4x^3-x^2-3x+3)}{ 5}$$ $$P_{10}(x)= \frac{x(2x+1)(x+1)}{6} \times \frac{ 3 x^8+ 12 x^7+ 8 x^6 - 18 x^5- 10 x^4+ 24 x^3 + 2 x^2 - 15 x +5}{ 11}$$ $$P_{11}(x)= \frac{x^2(x+1)^2}{4} \times\frac{2x^8 +8x^7+4x^6-16x^5-5x^4+26x^3-3x^2-20x+10}{ 6}$$ $$P_{12}(x)=\frac{x^{13}}{13}+\frac{x^{12}}{12}+x^{11}-\frac{11}{6}x^9+\frac{22x^7}{7}-\frac{33x^5}{10}+\frac{5x^3}{3}-\frac{961x}{2730}$$

论坛问题帖 | has_answers | pie | 2024-06-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019376 | MSE-4937560

Does the Sequence formed by Intersecting Angle Bisector in a Pentagon converges?

Now asked on MO here. Given a non-regular pentagon $A_1B_1C_1D_1E_1$ with no two adjacent angle having a sum of 360 degrees, from the pentagon $A_nB_nC_nD_nE_n$ construct the pentagon $A_{n+1}B_{n+1}C_{n+1}D_{n+1}E_{n+1}$ as follows: A_{n+1} is the intersection between the angle bisector of $\angle C_n $ and $\angle D_n$. B_{n+1} is the intersection between the angle bisectors of $\angle D_n$ and $\angle E_n$. C_{n+1} is the intersection between the angle bisectors of $\angle E_n$ and $\angle A_n$. D_{n+1} is the intersection between the angle bisectors of $\angle A_n$ and $\angle B_n$. E_{n+1} is the intersection between the angle bisectors of $\angle B_n$ and $\angle C_n$. (the two opposite angles) The reason why I chose this construction is that the point A_{n+1} is the only point that doesn't depend on A_n. I am allowing self-intersecting polygons in this constructions. My question is if this process is repeated indefinitely would the sequences ${A_n}$, ${B_n}$, ${C_n}$, $D_n $ and E_n converge ? There are only four possible scenarios: The points will converge. The points will eventually trapped on a loop. The points will diverge completely. Two or more points coincide, or two adjacent angle have a sum of 360 degrees ending the sequence. I tried to draw the first few pentagons to see if the point will converge or not. Here is the first 50 pentagons in a zoom-in animation: I conjecture that the sequences converge to a single point for all non-regular pentagons. If convergence occurs, how can we determine the limit point based on the initial pentagon? If the limit point exist what interesting properties does it have for the pentagon? Here is a Geogebra file that have the first 50 pentagons. This is a python code than can calculate A_n, B_n , C_n ,D_n ,E_n import math print("Number of pentagons") n=int(input()) m=max(n,50) m= min(m, 200) from decimal import Decimal import decimal decimal.getcontext().prec = m Decimal(10)**(-m) # enter your coordinates here A=[Decimal(0),Decimal(0)] B=[Decimal(0),Decimal(0)] C=[Decimal(0),Decimal(0)] D=[Decimal(0),Decimal(0)] E=[Decimal(0),Decimal(0)] print("Enter the coordinates of A_1") A[0]=Decimal(float (input())) A[1]=Decimal(float (input())) print("Enter the coordinates of B_1") B[0]=Decimal(float (input())) B[1]=Decimal(float (input())) print("Enter the coordinates of C_1") C[0]=Decimal(float (input())) C[1]=Decimal(float (input())) print("Enter the coordinates of D_1") D[0]=Decimal(float (input())) D[1]=Decimal(float (input())) print("Enter the coordinates of E_1") E[0]=Decimal(float (input())) E[1]=Decimal(float (input())) X=[Decimal(0),Decimal(0)] Y=[Decimal(0),Decimal(0)] Z=[Decimal(0),Decimal(0)] W=[Decimal(0),Decimal(0)] T=[Decimal(0),Decimal(0)] #defining some useful functions---------- def v(a,b): result= [b[0]-a[0], b[1]-a[1]] return result def crs(a, b): result = [-Decimal(0.5)*((Decimal(a[0]*b[1])-Decimal(a[1]*b[0]))*1)] return result def dis(a,b): result =[Decimal(math.sqrt((a[0]-b[0])**2+(a[1]-b[1])**2))] return result for i in range (2,n+1): AB= dis(A,B)[0] BC= dis(B,C)[0] CD= dis(C,D)[0] DE= dis(D,E)[0] EA= dis(E,A)[0] #Here to calculate A_n---------- x= CD*crs(v(D,C),v(D,E))[0] y= -CD*crs(v(E,D),v(E,B))[0]+ 2*DE*crs(v(C,B),v(C,D))[0] z= -CD*crs(v(C,B),v(C,E))[0]+ 2*BC*crs(v(D,C),v(D,E))[0] t= CD*crs(v(C,B),v(C,D))[0] X[0]=(x *B[0]+y*C[0]+z*D[0]+t*E[0])/(x+y+z+t) X[1]=(x *B[1]+y*C[1]+z*D[1]+t*E[1])/(x+y+z+t) #Here to calculate B_n---------- x= DE*crs(v(E,D),v(E,A))[0] y= -DE*crs(v(A,E),v(A,C))[0]+ 2*EA*crs(v(D,C),v(D,E))[0] z= -DE*crs(v(D,C),v(D,A))[0]+ 2*CD*crs(v(E,D),v(E,A))[0] t= DE*crs(v(D,C),v(D,E))[0] Y[0]=(x *C[0]+y*D[0]+z*E[0]+t*A[0])/(x+y+z+t) Y[1]=(x *C[1]+y*D[1]+z*E[1]+t*A[1])/(x+y+z+t) #Here to calculate C_n---------- x= EA*crs(v(A,E),v(A,B))[0] y= -EA*crs(v(B,A),v(B,D))[0]+ 2*AB*crs(v(E,D),v(E,A))[0] z= -EA*crs(v(E,D),v(E,B))[0]+ 2*DE*crs(v(A,E),v(A,B))[0] t= EA*crs(v(E,D),v(E,A))[0] Z[0]=(x *D[0]+y*E[0]+z*A[0]+t*B[0])/(x+y+z+t) Z[1]=(x *D[1]+y*E[1]+z*A[1]+t*B[1])/(x+y+z+t) #Here to calculate D_n---------- x= AB*crs(v(B,A),v(B,C))[0] y= -AB*crs(v(C,B),v(C,E))[0]+ 2*BC*crs(v(A,E),v(A,B))[0] z= -AB*crs(v(A,E),v(A,C))[0]+

2*EA*crs(v(B,A),v(B,C))[0] t= AB*crs(v(A,E),v(A,B))[0] W[0]=(x *E[0]+y*A[0]+z*B[0]+t*C[0])/(x+y+z+t) W[1]=(x *E[1]+y*A[1]+z*B[1]+t*C[1])/(x+y+z+t) #Here to calculate E_n---------- x= BC*crs(v(C,B),v(C,D))[0] y= -BC*crs(v(D,C),v(D,A))[0]+ 2*CD*crs(v(B,A),v(B,C))[0] z= -BC*crs(v(B,A),v(B,D))[0]+ 2*AB*crs(v(C,B),v(C,D))[0] t= BC*crs(v(B,A),v(B,C))[0] T[0]=(x *A[0]+y*B[0]+z*C[0]+t*D[0])/(x+y+z+t) T[1]=(x *A[1]+y*B[1]+z*C[1]+t*D[1])/(x+y+z+t) A=X B=Y C=Z D=W E=T print(f"A_{i}= {A}") print(f"B_{i}= {B}") print(f"C_{i}= {C}") print(f"D_{i}= {D}") print(f"E_{i}= {E}") print("\n") Another thing that supports the convergence of the series that no matter how many times I used that random function in python with the sequence the result always converge (suggesting it always converge except some zero measure set)

论坛问题帖 | no_answers_at_retrieval | pie | 2024-06-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019377 | MSE-4939773

Conjectures about the greatest common divisor of a vertical column of the pascal triangle.

I was playing around with pascal triangle I noticed an interesting property concerning the greatest common divisor gcd of binomial coefficients along a vertical line. Specifically, the line containing 2 is the only one where the $\gcd$ of all coefficients is greater than 1 (namely, 2). This made me wonder: What $n \in \mathbb{N}$ such that $\gcd _{k\ge1 }\left\{\binom{n+2k-2}{k}\right\}>1$ I tried every number between $2 $ and 10 by hand which all failed that made me conjecture that $n=2$ is the only number which has $\gcd _{k\ge1 }\left\{\binom{n+2k-2}{k}\right\}>1$ I wrote a simple python code to test that 2 is the only number that has this property for n in range(2,20_001): a=n for k in range (2,10_001): math.gcd(a, math.comb(n+2*k-2,k)) if math.gcd(a, math.comb(n+2*k-2,k))==1: break a=math.gcd(a, math.comb(n+2*k-2,k)) if k==10_000: print(n) print(a) print("\n") between 2 and $20,000$ the only number which has this property is 2. Another interesting thing about this sequence for $n\ne 2$ $\gcd _{1 \le k\le m}\left\{\binom{n+2k-2}{k}\right\}>1$ fail at a power of 2, in other words the for $ n \ne 2$ $\gcd _{1 \le k\le m}\left\{\binom{n+2k-2}{k}\right\}>1$ will first be broken when m is a power of 2. Here is the first 100 n and the first m that $\gcd _{1 \le k\le m}\left\{\binom{n+2k-2}{k}\right\}=1$ in the form "(n,m)" (3,2), (4,2), (5,2), (6,4), (7,2), (8,2), (9,2), (10,8), (11,2), (12,2), (13,2), (14,4), (15,2), (16,2), (17,2), (18,16), (19,2), (20,2), (21,2), (22,4), (23,2), (24,2), (25,2), (26,8), (27,2), (28,2), (29,2), (30,4), (31,2), (32,2), (33,2), (34,32), (35,2), (36,2), (37,2), (38,4), (39,2), (40,2), (41,2), (42,8), (43,2), (44,2), (45,2), (46,4), (47,2), (48,2), (49,2), (50,16), (51,2), (52,2), (53,2), (54,4), (55,2), (56,2), (57,2), (58,8), (59,2), (60,2), (61,2), (62,4), (63,2), (64,2), (65,2), (66,64), (67,2), (68,2), (69,2), (70,4), (71,2), (72,2), (73,2), (74,8), (75,2), (76,2), (77,2), (78,4), (79,2), (80,2), (81,2), (82,16), (83,2), (84,2), (85,2), (86,4), (87,2), (88,2), (89,2), (90,8), (91,2), (92,2), (93,2), (94,4), (95,2), (96,2), (97,2), (98,32), (99,2), (100,2), This sequence is also interesting because of its pattern i.e Three 2s followed by a power of 2 greater than 2. My questions are: How to prove that $n=2$ is the only $n \in \mathbb{N}$ such that $\gcd _{k\ge1 }\left\{\binom{n+2k-2}{k}\right\}>1$ For $n \ne 2$ how to prove that the first $m \in \mathbb{N}$ such that $\gcd _{1 \le k\le m}\left\{\binom{n+2k-2}{k}\right\}=1$ is always a power of 2 What is the pattern for such m? With the help of @ronno observation I was able to write a code that checked for all $n \le 10^6$ and none of them had a gcd >1 except 2

论坛问题帖 | has_answers | pie | 2024-06-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019378 | MSE-4942219

Lower bound of n th Taxi-cab number $N = a^3 + b^3 = x^3 + y^3$

Let N,a,b,x,y be distinct positive integers such that $$N = a^3 + b^3 = x^3 + y^3$$ Also known as Taxicab numbers or Taxi-cab numbers. see also : https://oeis.org/A001235 let $t(n)$ be the n th Taxi-cab number. So the first one is $t(1) = 1729$ because $1729 = 12^3 + 1^3 = 10^3 + 9^3$. Let $\pi(n)$ be the counting function of the primes up to n. And $\zeta(s)$ is the regular Riemann zeta function. Then it appears that for all positive n $$ \lfloor(\frac{\pi(n)}{\zeta(3)})^{T}\rfloor< t(n)$$ with $$T = 1+\dfrac{2+\dfrac{3+\dfrac{4+\cdots}{5+\cdots}}{4+\dfrac{5+\cdots}{6+\cdots}}} {3+\dfrac{4+\dfrac{5+\cdots}{6+\cdots}}{5+\dfrac{6+\cdots}{7+\cdots}}} $$ (The pattern is $+1$ when going up and $+2$ when going down). Or even the slightly stronger version : $$ \lfloor(\frac{\pi(n)}{\zeta(3)})^{\sqrt 3}\rfloor< t(n)$$ Notice $T = 1.728..$ (not 1.729 lol) and $\sqrt 3 = 1.732..$ (No closed form is know for T as far as I know. I incorrectly assumed it was $\sqrt 3$ here : Simplifying a complicated continued fraction expression.) Let $w(n)$ be the counting function of the Taxi-cab numbers. So $(w(t(n)) = t(w(n)) = n$. Then by looking at families of parametric solutions to $$N = a^3 + b^3 = x^3 + y^3$$ it seems manageable to prove that for large n we have $$w(n) < \sqrt n + \sqrt{\sqrt n} + \sqrt{\sqrt{\sqrt n}} $$ but this estimates $t(n)$ around n^2 what appears too high for large n. I tried some statistical ideas and sieve ideas but I did not get beyond this high estimate $t(n)$ around n^2. I tried some analytic number theory with a generalized theta function $f(s)^k = (\sum_{n>0} s^{n^3})^k$ but also without succes. How to prove one of these conjectures or get closer ? Or any good arguments pro or contra ? edit : Maybe this helps : The complete solution in positive integers to, $$x_1^3+x_2^3 = x_3^3+x_4^3$$ was given by Choudhry's On Equal Sums of Cubes (1998). For positive integers a,b,c, $$\begin{aligned} d\,x_1 &= (a^2 + a b + b^2)^2 + (2a + b)c^3\\ d\,x_2 &= (-a^3 + b^3 + c^3)c\\ d\,x_3 &= (a^2 + a b + b^2)^2 - (a - b)c^3\\ d\,x_4 &= (a^3 + (a + b)^3 + c^3)c\end{aligned}$$ where, $$(a^3-b^3)^{1/3}<\,c\,<\frac{(a^3-b^3)^{2/3}}{a-b}$$ and $d=1$, or chosen such that $\text{GCD}(a,b,c)=1$. P.S. For Choudhry's complete solution in positive integers to $x_1^3+x_2^3+ x_3^3=x_4^3$, see this post. edit 2 A simple heuristic to get a power estimate is this : $$a^3 + b^3$$ has the density $\sqrt n^3 = n^{\frac{3}{2}}$. so $$N = a^3 + b^3 = x^3 + y^3$$ has the density $ n^{(\frac{3}{2})^2} = n^{\frac{9}{4}}$. or $n^{2.25}$ But that is much larger than $n^{\sqrt 3}$ and it also much larger than what the data suggests under some logical assumptions and extrapolation. Say we estimate $t(n)$ as $C n^a $ or more precisely $1729 * n^a$. Then we have $t(30000) = 520890296211 = 1729 * 30000^{1.89..}$ I ofcourse assumed the constant $C = 1729$ is not estimated to high. see https://oeis.org/A001235/b001235.txt If you estimate the constant C much smaller such as 43 we get $$ 43 * 30000^{2.25} <t(30000) < 44 * 30000^{2.25}$$ So the final words have not been said. But from the data and arguments it seems the exponent

is going down for large n. We can also say $t(30000)$ is close to $9165 * 30000^{\sqrt 3}$ where 9165 is going down as n grows.

论坛问题帖 | no_answers_at_retrieval | mick | 2024-07-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019379 | MSE-4957665

Conjecture: Definite existence of a prime number in the range of ${[|3^{p_i}-k\times{p_i};3^{p_i}+k\times{p_i}|]}$

My old post was closed while I had clearly tried to explain at my best. So I'll retry without posting outer links and reexplaining at most details what this conjecture is about, which was a research that I did 7 years ago when I was a college student for fun. But since I wasn't in math field I couldn't finish the work so I hope if someone is interested, who's got interest in prime numbers, to take it over if he find this explanation below convincing. Note: this post What is the appropriate way to post conjectures on Math.SE clearly says that I can post conjecture here if I understand, as long as I provide enough details to make it interesting. Origin and problematic The spark of idea came from Bertrand's postulate, which are these 3 formulas: ${\forall n\in \mathbb {N};n > 3;\exists p\in \mathbb {P} :\qquad n<p<2n-2.}$ ${\forall n\in \mathbb {N};n > 1;\exists p\in \mathbb {P} :\qquad n<p<2n.}$ ${for~~{n}\geqslant 1:\qquad p_{n+1}<2p_n.}$ What I noticed, back at that time and if I wasn't wrong since I wasn't that versed in maths. Is that this theorem was the most precise theorem for ensuring that there exist primes in a certain range. I take n = 200, I'm sure I'll find primes between 200 and 400 I take n = ${2^{10}}$, I'm sure I'll find primes between ${2^{10}}$ and ${2\times2^{10}}$ Now the problem is when the scale become higher, which means the digits are growing, 100 digits, 10 to power of a huge n digits, etc. I can take a number ${a}$ which has like 100 digits, and according to the theorem, I'm sure to find a prime between ${a}$ and ${2a}$. But I have no idea where that next prime is, it could be the next 2 numbers after ${a}$, it could be the next 10k number, it could be after 1 million number(well I doubt), etc ... Because the search range is so big. We can sumarize this into two issues: the maximum range search is too big there is no minimum range search Note : While writing this, just found out that the theorem got some precision improvements, which gives a better search range but still it's considered a bit big. Example for using ${\displaystyle x<p\leq \left(1+{\frac {1}{5\,000\ln ^{2}{x}}}\right)x}$. I can input a number ${468,991,632,168,991,632}$ which has 18 digits, and the other side will give me approximately ${468,991,688,823,352,400}$ which has 18 digits. The search range here is ${56,654,360,768}$ numbers. Too much for introducing the problematic, let me share with you some few examples of what I did research: Observations Back at the time I wanted to narrow my research only on ${prime^{prime}}$ to find out of there are any special relationships, I ended up only testing values of ${3^{prime}}$ because it took a huge time. (now creating the table and copying values from wikiversity to here is such a pain) prime number ${p_i}$ ${3^{p_i}}$ distance from next prime next prime distance from previous prime previous prime 2 9 2 11 2 7 3 27 2 29 4 23 5 243 8 251 2 241 7 2187 16 2203 8 2179 11 117 147 16 117 163 14 117 133 13 1 594 323 8 1 594 331 22 1 594 301 17 129 140 163 34 129 140 197 4 129 140 159 19 1 162 261 467 56 1 162 261 523 14 1 162 261 453 23 ..... 178 827 32 ...178 859 20 .178 807 29 ..... 364 883 30 ...365 013 14 ...364 869 31 ..... 283 947 16 ...283 963 4 ...283 943 37 ..... 997 363 50 ...997 413 2 ...997 361 41 ..... 786 403 70 ...786 473 2 ...786 401 43 ..... 077 627 52 ...077 679 74 .077 553 47 ..... 287 787 52 ...287 839 46 ..287 741 53 ..... 796 723 26 ...796 749 4 ...796 719 59 ..... 811 067 64 ...811 131 38 ...811 029 61 ..... 299 603 34 ...299 637 74 ...299 529 67 ..... 410 587 230 ...410 817 298 ...410 289 71 ..... 257 547 20 ...257 567 20 ...257 527 Note: I couldn't put all what I tested in wikiversity, it was a true pain to already calculate and compare at that time so all the other tests I've done were with pen and paper and online tools to calculate. I have tested all powers from ${3^{2}}$ till ${3^{257}}$. The last one has like between 120 and 128 digits. Even the last one in this table above has 34 digits During all these tests, I have concluded these observations: I could definitely, from ${3^{2}}$ till ${3^{257}}$, find a prime number in a range of ${[|3^{p}-3{p}~~;~~3^{p}+3{p}|]}$. Except for ${3^{67}}$ which was ${[|3^{p}-4{p}~~;~~3^{p}+4{p}|]}$ so that means, for a huge number like ${3^{257}}$ which has 123 digits, I can find at least one prime in a range of ${[|3^{257}-3\times{257}~~;~~3^{257}+3\times{257}|]}$, which is a search range of 1542 numbers, and that's for a very huge number Hypotheses Now I would have been happier if ${3^{67}}$ didn't interfere that badly so that the multiplier could be stuck at 3, sadly. So I can put 2 hypotheses: The first hypothese : The multiplier, at it's minimum range, can be considered 3. If multiple occurences after ${3^{257}}$ denies that possibility. That means either we increment the multiplier value (named ${k}$ by one everytime, like going from ${[|3^{p}-3{p}~~;~~3^{p}+3{p}|]}$ to ${[|3^{p}-4{p}~~;~~3^{p}+4{p}|]}$ then ${[|3^{p}-5{p}~~;~~3^{p}+5{p}|]}$. Or that there could be a condition for the k to be incremented to a certain number The second hypothese : I can maximise, definitely, until proven wrong, the value of ${k}$ to be the given prime number. Which means that the maximum range would be ${[|3^{p}-p\times{p}~~;~~3^{p}+p\times{p}|]}$ => ${[|3^{p}-{p^2}~~;~~3^{p}+{p^2}|]}$. Taking the ${3^{257}}$ and supposing that I didn't find the minimum. I can assume that max range would be ${[|3^{257}-{257^2}~~;~~3^{257}+{257^2}|]}$. ${257^2}$ = 66 049 so that means the search range would be 132 098 which is so incredible as a search range for a 123 digits number In a nutshell: Using the min value : ${\displaystyle \forall p_i\in \mathbb {P} ;\exists {p}\in \mathbb {P} ;\exists {k}\in \mathbb {N^*}; (2 < k < p_i):~~ p\in [|3^{p_i}-k{p_i}~~;~~3^{p_i}+k{p_i}|]}$ Using the max value : ${\displaystyle \forall p_i\in \mathbb {P} ;\exists {p}\in \mathbb {P} :~~ p\in [|3^{p_i}-{p_i}^2~~;~~3^{p_i}+{p_i}^2|]}$ Like I've said, I was able to test only the powers of 3. So I wonder if maybe other primes to primes powers could have possibly, at least that max search range, based on the given prime. So finally, why do I think that this research may be valuable: Having a good search range and existence of a minimum prime number, based on primes numbers. especially for huge numbers Possibility of application of these idea to other primes to the power of primes. Unlocking another prime to prime relationship Minimising the search range for prime numbers that are huge You who are far more proficients in Math than I, and me who forgot a lot of advanced maths because I'm in another career. I really think this conjecture has a potential and would like to know if you think that this can be ever needed in math or no. Thanks for reading, if you have any questions or remarks, don't hesitate. Although like I've said I've forgotten most of the advanced stuffs

论坛问题帖 | has_answers | akutagawa akutagawa | 2024-08-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019380 | MSE-4959652

Do there exist integers that can be written as the sum of two powers of the same prime in two ways?

Prof. Peter Cameron has a list of conjectures on his website, one of which is the following (unresolved) one: There are only six positive integers greater than 2 which can be written as the sum of two powers of the same prime in more than two ways I fail to see how an integer can be written as the sum of two powers of the same prime in even two ways:- Suppose it is true for some integer k. Then, there exists a prime p and positive integers m, n, g, h, such that $m \neq n$ and: $k = p^m + p^{m + g} = p^n + p^{n + h} \Rightarrow p^{m - n} = \dfrac{1 + p^h}{1 + p^g}$, but this is not possible as p does not divide $1 + p^h$.

论坛问题帖 | no_answers_at_retrieval | drumjo | 2024-08-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019381 | MSE-4965811

Is it possible to prove that iterating this hailstone-resembling algorithm will always end in 1?

I've been messing with a rule I came up with recently, that resembles the Collatz function in formulation but shares very few practical similarities. Using p_n to represent the n^{th} prime number and c_n to represent the n^{th} composite number, the formulation is as such: \begin{cases} \text{if } n \text{ is prime:} & p_n \to p_{n+1} - p_n \\ \text{if } n \text{ is composite:} & c_n \to c_{n-1} + c_n \\ \end{cases} In words, this means, "If the number is prime, take the difference between the number and the next prime number, and if the number is composite, take the sum of the number and the previous composite number." The sequence ends when it gets to 1. (Because 4 is the first composite number, we treat c_0 as 1, meaning that 4 goes to 5.) Let's use 6 as an example: because 6 is composite, it goes to $6 + 4 = 10$; because 10 is composite, it goes to $10 + 9 = 19$; because 19 is prime, it goes to $23 - 19 =4$; and continuing this process, the sequence unfolds as: $$6 \to 10 \to 19 \to 4 \to 5 \to 2 \to 1.$$ These sequences have a tendency to act very erratically. For example, the sequence beginning with 32 demonstrates many features of this process: $${\small32 \to 62 \to 122 \to 243 \to 485 \to 969 \to 1937 \to 3873 \to 7745 \to 15489 \to 30977 \to 6 \to 10 \to 19 \to 4 \to 5 \to 2 \to 1.}$$ We can notice several things; first, when we reach an odd number, all the numbers thereafter will be odd until we reach a prime number (since the nearest composite number below an odd number will always be even). In the above sequence, all the numbers from 243 to 15489 are composite, and only because 30977 is prime do we return to an even number. Also, we can see, as the numbers get higher, they grow faster, and the chance of them being prime is much rarer. In fact, when a sequence starts growing, it looks like it'll never stop, at least for a while; the sequence starting with 95 has 595 terms, the largest of whom is the 178-digit prime number $${\scriptsize14879396952621968769079831664541974952513501961928904289230033454548697062408957128966234687844381586574195912989130942655378120463894152791647576690929892981863063412465740021 77.}$$ The next prime number is 152 greater than that, so the next item in the sequence is 152, after which the sequence rapidly goes to 1 (through the primes 4817 and then 101). Here's a picture of the start of the sequence (easier to see if you open it in a new tab): It is easy to see that these sequences can rapidly skyrocket, at a rate just barely less than 2^n. The question is: Is there a sequence that never ends (that never goes to 1)? Is there any rigorous way to prove that either all sequences go to 1 or that one sequence will never reach 1? I conjecture that they all go to 1, but I can't imagine how to go about proving it -- any help would be much appreciated!

论坛问题帖 | has_answers | Mathemagician314 | 2024-09-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019382 | MSE-4976603

Is it actually really easy to state conjectures, which are open and on the first view really hard to prove?

The longer I study mathematics, the more I think that we actually don’t know that much. I mean of course we don’t know that much, because theoretically mathematics is an infinitely big field of research and we as a species supposedly will only live for a finite time. So it would be reasonable to say that we don‘t know anything in mathematics. But that’s a little bit too philosophical for my question. Now I always wondered how some mathematicians even came up with theorems that are really hard to prove. But now I think it is really easy to come up with a conjecture. The only difference would be that this conjecture probably won’t have many interesting consequences. So what do I mean by that? Let me state the following: Let p be a prime number. p is called the n-th counting prime, if the sequence $123...n$ is somewhere in it’s decimal representation. So 11 would be the first counting prime and 127 the second counting prime. My conjecture would now be: for any $n \in \mathbb{N}$ the $n-th$ counting prime number does exist. So this just came to my mind and I think it would be an open problem. If it is then I would like to call it the Matthias Conjecture. And in here you could replace prime numbers with powers of two, three, seven or the round off of the imaginary part of non-trivial solutions to $\zeta(s) = 0$, where ζ is the Riemann zeta function.... Or even the real part of these solutions :) I am a little bit frustrated with this thought, because for the most part I had the feeling that we can do so much with mathematics today, but if even I could state something which is unknown and hard to prove, I rather feel like we really don’t know anything... So what do you think? I don’t really know if this fits in with this site; if not I can delete it.

论坛问题帖 | has_answers | MathMatthias | 2024-09-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019383 | MSE-4977548

Conjecture: Every Non-Zero Integer as the Difference of a Semiprime and a Prime

I would like to propose the following conjecture and seek insights or references related to it: Conjecture: Every non-zero integer n can be expressed as the difference of a semiprime q and a prime p, that is, $$n=q-p,$$ where: q is a semiprime (the product of two primes, not necessarily distinct), and p is a prime number. Numerical Verification: To gather empirical support for this conjecture, I conducted extensive numerical experiments within the range $-10,000,000≤n≤10,000,000$ (excluding $n=0$). In every case within this range, I successfully found pairs of p and q satisfying the equation $n=q-p$. No

counterexamples were observed. Example: For instance, when $n=15$: Choose p=7 (a prime), Then q=7+15=22, which is a semiprime (22=2×11). Thus, $15=22-7$. Questions: In seeking a proof or counterexample for this conjecture, what approaches or existing theories would be effective? Especially, I would appreciate advice from the perspectives of additive number theory or multiplicative number theory. Additional Information: Implementation Details: The numerical experiments were conducted using a Python program optimized with the Sieve of Eratosthenes for prime generation and parallel processing to handle large ranges efficiently. While I can provide the code upon request, the focus here is on the mathematical aspect of the conjecture. Exclusion of Zero: The case n=0 is excluded since it would require q=p, and while p is prime, q would need to be semiprime, which only occurs if p itself is a semiprime. Given that primes greater than 2 are not semiprimes, n=0 serves as an inherent exception to the conjecture. Graphical Representation: Upon observation, there is a tendency for n to stick to the upper range when it is a multiple of 6 (Especially multiples of 30), and to the lower range when n is a prime number. Motivation The motivation for this conjecture stems from the robustness of semiprimes used in RSA encryption. Specifically, semiprimes are known for their resistance to division by prime numbers, as they rarely divide evenly. This led me to consider a different approach: what if, instead of division, we use subtraction? Given that semiprimes do not divide cleanly, I hypothesized that their differences with primes might display some kind of pattern or periodicity, resulting in jumps to other numbers. From this idea, I further speculated that these "jumps" could cover all integers in a somewhat uniform manner. However, recognizing that semiprimes and primes belong to different classes, it became clear that expressing zero was not feasible. Thus, zero was excluded from the conjecture's conditions. If proven, this conjecture could offer new perspectives in number theory, with possible connections to cryptographic systems like RSA encryption. Additionally, this conjecture may inherently include basic memory management functionality. I appreciate any feedback, references, or discussions that can shed light on the validity and potential avenues for this conjecture. Thank you!

论坛问题帖 | no_answers_at_retrieval | Akira Sukigi | 2024-09-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019384 | MSE-4978314

Conjecture: Every Non-Negative Integer as the Difference of a Multiple of a Prime and a Prime

In a recent post, I proposed a conjecture suggesting that every non-zero integer can be expressed as the difference between a semiprime and a prime. Conjecture: Every Non-Zero Integer as the Difference of a Semiprime and a Prime To explore this idea further, I examined a related but smaller problem. This exploration led to the following new conjecture, now including zero and focusing on non-negative integers: (Note: The range of the conjecture was incorrect, so the title and content have been revised.) Conjecture: For any prime number p, every non-negative integer k (including zero) can be expressed in the form: $$ k = p \cdot m - q $$ where m is a positive integer and q is a prime number (which may be the same as or different from p). Computational Evidence To investigate this conjecture, I conducted a large-scale computational experiment using GPU acceleration. The steps were as follows: 1.Prime Generation: Generated all primes up to a very large limit (p_{max}) and used this set for the multiples p. Created a prime bitmask for rapid prime checks up to $q_{\text{limit}}=101,010,000$. 2.Expressibility Check: For each integer k in the range $[0,100,000,000]$ (including zero), checked whether there exists a prime p (from 2 up to p_{max}) and an integer $m≥1$ such that $k=p·m-q$ for some prime q. Utilized GPU parallel processing to handle the extensive computations. Results: Prime Generation: === Result === All integers in the range [0, 100,000,000], including zero, can be expressed as p * m - q , where p is any prime between 2 and 383,393, m is a positive integer, and q is any prime. This computational evidence suggests that the conjecture holds for integers from 0 to $100,000,000$ and for any prime p from 2 to $383,393$. Although this does not constitute a formal proof, it provides a strong numerical basis for considering the conjecture's validity in general. Questions and Discussion Points Theoretical Implications: What mathematical properties of primes might underlie the ability to express every non-negative integer, including zero, in this form? Could this conjecture be connected to or derived from existing results in number theory, such as prime gaps or properties of prime distributions? Relationship to Semiprimes: While this conjecture does not directly involve semiprimes, understanding why any prime p allows for such an expression could provide insights into the original conjecture involving semiprimes. Formal Proof or Counterexamples: What approaches might one take to prove or disprove this conjecture for all primes p? Are there known counterexamples or classes of numbers that are not expressible in this form? Further Computational Work: Should further computational experiments extend beyond the current range or explore additional properties of p and q to gather more evidence? Conclusion This numerical investigation, including the ability to express zero and all positive integers in the specified form, hints at a potentially deep relationship between integers, prime multiples, and prime differences. While the computational evidence is strong, a formal proof or theoretical framework is necessary to fully understand this phenomenon. I am curious to hear thoughts from the community on possible approaches to explore or prove this conjecture.

论坛问题帖 | has_answers | Akira Sukigi | 2024-09-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019385 | MSE-4986942

Is there a counterexample to the No-Loop Conjecture when A is not an Artin algebra?

Let A be an Artin algebra, i.e., A is an algebra over a commutative Artin ring R that is a finitely generated R-module. The No-Loop Conjecture states that when A is an Artin algebra (or more generally an Artinian ring), if A has finite global dimension, then $$ \text{Ext}^1(S, S) = 0 $$ for all simple A-modules S. My question is: when A is not an Artin algebra, i.e., A is a general ring, is there a counterexample to the No-Loop Conjecture?

论坛问题帖 | no_answers_at_retrieval | Liang Chen | 2024-10-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019386 | MSE-4987006

Are cage graphs Hamiltonian?

An (r, g)-cage graph is an r-regular graph with the least amount of vertices having girth g. While looking up examples of cage graphs I couldn't help myself but noticing they all seem to be Hamiltonian, with the exception of the Petersen graph. The Wikipedia article on cage graphs list the following as notable examples: and in fact, all of them but the Petersen graph are Hamiltonian (you can clearly see this in they're particular articles as their main diagram have their vertices arranged in one big circle). The $(r, 3)$ and $(r, 4)$-cages are K_r and $K_{r,r}$, respectively, both of which are clearly Hamiltonian for $r\ge 3$ (notice cage graphs only exist for $r\ge3$). Also, cage graphs are not necessarily unique for a fixed pair (r, g) and there are three $(3, 10)$-cages, the Balaban 10-cage, the Harries graph and the Harries-Wong graph (as listed by Wikipedia), all of them Hamiltonian. I looked upon some more examples and decided that was enough evidence gathering, so now I wonder how should one proceed about proving this. The amount of vertices in an (r, g)-cage is much larger than r or g in general (this is a theorem) so Bondy-Chvátal Theorem does not apply. This question has been asked before here, but it was also asking about a second (and also very interesting) conjecture on cage graphs and hasn't got much attention. This might be an open problem as far as I know, but I haven't seen anyone proposing it beyond the previous post. Update: The paper The smallest cubic graphs of girth nine (access here) lists every single one of the 18 $(3, 9)$-cages and notices they are all Hamiltonian.

论坛问题帖 | no_answers_at_retrieval | Alma Arjuna | 2024-10-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019387 | MSE-4989540

Conjecture: the integer cosets of ideals are Noetherian

I have strong evidence that this fact will help prove continuity of $\pi_2(n)$ the twin prime average counter on the naturals. Continuity with respect to the Furstenberg topology that is. See my recent post history under continuity for more details. Let $U_1 \subset U_2 \subset \dots$. This sequence must eventually stabilize where $U_i=a_i + b_i\Bbb{Z}$ are integer ring cosets of ideals. It makes sense but how do we prove that b_{i+1} divides b_i? If $a +b\Bbb{Z} \subset c + d\Bbb{Z}$ then $b = \text{lcm}(b,d)$ by intersecting the two cosets. Ok then so d must divide $a-c$. Or $a= c \pmod d$. Since the ideals obviously stabilize, so too must the taken standard residue and we’re done.[U+2705] So my question now becomes are either finite unions of cosets or arbitrary unions there of also Noetherian? If arbitrary unions stabilize then I can work with open sets in Furstenberg topology. But if it’s only finite unions that become stable when in the inclusion chain [U+26D3][U+1F4A5] then I will have to do something with closed sets. So really hoping for Noetherian arbitrary unions of cosets. Clarification. Let τ be a topology that is generated by cosets found in the Furstenberg topology, which is $\sigma = \langle a + b\Bbb{Z}: a,b \in \Bbb{Z}, b \neq 0 \rangle$. In other words, τ forms a topology itself using some subset of the Furstenberg basis. Now let $U_i = \bigcup_{j = 1}^{n_i} (a_j^i + b_j^i\Bbb{Z}), \ i \geq 1$ be a sequence of finite unions of basic sets such that $U_{i} \subset U_{i+1}$ for all $i \geq 1$. Question. Does the sequence $U_1 \subset U_2 \subset U_3 \dots$ stabilize to U_n for some $n \geq 1$? I.e. for all $m \geq n$ we have that $U_m = U_n$? In other words, are finite unions of basic open sets in τ Noetherian? Attempt. I would guess the answer is a resounding "No!", because even though some cosets within the unions would stabilize, we could just keep adding in more random cosets so that the sequence just increases to some open set, not necessarily an entry in the chain. Luckily, though I thought I needed this at one point in my Topo-NT problem, I found out that approach had other unrelated errors. It is still kind of cool that coset chains are Noetherian.

论坛问题帖 | no_answers_at_retrieval | TotoposDiagramChaseApp | 2024-10-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019388 | MSE-4991666

Can a slice from the edge of a regular tetrahedron form a right triangle?

Let $PABC$ be a regular tetrahedron. Fix a point K on PB, and consider $\triangle AKC$. This triangle is isoceles (proven below). Can it be right? Proof that $\triangle AKC$ is isoceles: Observe that $\triangle ABK \cong \triangle CBK$ by SAS: since $PABC$ is regular, $AB \cong CB$, and $\angle ABK \cong \angle ABP \cong \angle CBP \cong \angle CPK$. Thus, $AK \cong CK$. If $K=B$ or $K=P$, this triangle is equilateral; otherwise, $\angle AKC > \angle KAC = \angle KCA$. For $|CK|$ is at its minimum when it is the altitude of $\triangle PCB$, which in an isoceles triangle happens when CK bisects $\angle PCB$, and a maximum when K is incident with P or B. When $|CK|$ is at a minimum, $\angle CKA$ is at a maximum, by Euclid I.21. My hunch is this happens when $\angle CKA$ is right, but I can't prove this. I also can't determine the angle from KC to the plane ABC.

论坛问题帖 | has_answers | SRobertJames | 2024-10-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019389 | MSE-4993224

New property of positive integers and prime numbers?

While playing with numbers, I found the following properties of positive integers and prime numbers. Properties Every positive integer n can be written in the following form \begin{align*} n = \frac{a^2 + b^2}{a + b} \end{align*} with at least 3 integer solutions. Mostly, the number of solutions is (along with probabilities of occurrence): $$\begin{matrix} 11 & 46\,\% \\ 3 & 39.4\,\% \\ 35 & 9.48\,\% \\ 19 & 3.2\,\% & \text{factorization contains square}\\ 59 & 1\,\% & \text{factorization contains 5^2}\\ 27 & 0.6\,\% & \text{factorization contains cube}\\ 107 & 0.2\,\% \\ 83 & 0.08\,\% \\ \end{matrix}$$ Moreover, every prime number p can be written in the following form \begin{align*} p = \frac{a^2 + b^2}{a + b} \end{align*} with either 3 or 11 integer solutions. The number of solutions is (along with probabilities of occurrence): $$\begin{matrix} 3 & 50.953682\,\% \\ 11 & 49.046322\,\% \\ \end{matrix}$$ Example For \begin{align*} \frac{a^2 + b^2}{a + b} = 127 \end{align*} there is exactly 3 integer solutions: $(a = 0, b = 127)$, $(a = 127, b = 0)$, and $(a = 127, b = 127)$. Question Can you confirm this result? Is this known? Any feedback is welcome.

论坛问题帖 | has_answers | DaBler | 2024-11-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019390 | MSE-4994966

Does $\prod_{n\ge1}\frac{\cos\frac{x}{n}}{x^{n^{2}}+1}=1-x$ for $-1<x\le1$?

I was just typing random things into Desmos and I came across this. WolframAlpha didn't give a closed form for the numerator or denominator and I don't recall coming across any formula like this before. Here is my attempt to solve it analytically: $$\prod_{n\ge1}\frac{\cos\frac{x}{n}}{x^{n^{2}}+1}=\exp\sum_{n\ge1}^{ }\ln\frac{\cos\frac{x}{n}}{x^{n^{2}}+1}$$ As you can see, I didn't really get anywhere. I also tried to expand different parts into their series expansions but it didn't seem to help. Is my conjecture that $\prod_{n\ge1}\frac{\cos\frac{x}{n}}{x^{n^{2}}+1}=1-x$ on $-1<x\le1$ true?

论坛问题帖 | has_answers | Dylan Levine | 2024-11-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019391 | MSE-5015220

Conjecture about cyclic hexagons and tangent circles

Is the following conjecture true: In every cyclic convex hexagon, there exists a point P such that for every pair of opposite sides s and t, the circle through P and the endpoints of s, is tangent to the circle through P and the endpoints of t. In the diagram, circles associated with opposite sides are the same color. For brevity, I shall call such circles "opposite circles". The conjecture seems to be true, based on experiments using Geometer's Sketchpad. The conjecture is not generally true if: the hexagaon is non-cyclic the hexagon is inscribed in a non-circular ellipse if we change "hexagon" to "$2n$-gon" with $n>3$ My attempt I considered: Lines through the centres of opposite circles Lines through P tangent to opposite circles Lines that are tangent to opposite circles like an ice cream cone with two scoops of ice cream Labelling the vertices anticlockwise $ABCDEF$, then drawing AE extended, and BD extended But I could not find anything useful. Any proof must use the fact that the polygon has six sides, and is cyclic; but how? Context Recently I discovered that, in any tangential quadrilateral $ABCD$ with incentre O, the circle through A,B,O is tangent to the circle through C,D,O. (This is easily proved by extending opposite sides to form a triangle, and using the incentre-excentre lemma.) I tried to generalize this fact. I didn't generalize it, but instead came up with the conjecture here.

论坛问题帖 | has_answers | Dan | 2024-12-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019392 | MSE-5018417

Lemniscate constant as infinite product

I am looking for a hint on how to prove the following conjecture: $$\dfrac{\varpi}{\sqrt{2}}=\displaystyle\prod_{n=1}^{\infty} \dfrac{2n}{\left(2n +(-1)^n \right)}$$ where ϖ denotes lemniscate constant. PARI/GP code.

论坛问题帖 | has_answers | Pedja | 2025-01-02

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019393 | MSE-5019675

Conjecture: For n circles in a triangle, the maximum number of tangent points is $3n$.

Is the following conjecture true: For n non-overlapping circles (or disks) in a triangle, the maximum number of tangent points is $3n$. The tangent points include points where two circles are tangent, and points where a circle is tangent to a side of the triangle. For example, here are 7 circles and 21 tangent points. My thoughts The conjecture is not true for other polygons: for example, a single circle inscribed in a square has four tangent points. Could this be related to the fact that a circle can be surrounded by at most six other circles of the same size? I am not very familiar with graph theory, but I suspect that may be used here. Context I was playing with circles in triangles, and I started to wonder about he maximum number of tangent points.

论坛问题帖 | has_answers | Dan | 2025-01-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019394 | MSE-5026301

New property of composite Mersenne numbers?

While I was playing around with Mersenne numbers, I came across an interesting thing: For p prime, $2^p - 1$ composite, $2 < b < 2^p - 1$, $\gcd(b,2^p - 1) = 1$, there exists k such $(b^k)^2 \equiv 1 \pmod{2^p - 1}$, where $b^k \not\equiv 1 \pmod{2^p - 1}$. Then the integer factorization of $b^k - 1$ contains a factor of $2^p - 1$, whereas the factorization of $b^k + 1$ contains a cofactor (i.e. another factor of $2^p - 1$). I will demonstrate this with an example: Start with $p = 11$, thus $2^{11} - 1$ is composite integer (Mersenne composite) with the factorization 23×89. Next, we choose arbitrary base b, here for example $b = 3$. By exhaustive search, we can find that $b^k = 3^{44}$, and $3^{44} \equiv 622 \pmod{2^{11} - 1}$. Note that $622^2 \equiv 1 \pmod{2^{11} - 1}$. Now we have factorization of 621 a 623 as follows. $621 = 3 \times 3 \times 3 \times \mathbf{23}$ $623 = 7 \times \mathbf{89}$ Therefore if we find $x = b^k$ such that $x^2 \equiv 1 \pmod{2^{p} - 1}$, where $x \not\equiv 1 \pmod{2^{p} - 1}$, we have got immediately the factorization of $2^{p} - 1$. Another example: Let $p = 23$, thus $2^{23} - 1$ is a composite Mersenne number with the factorization 47×178481. We choose $b = 3$. We find $x \equiv -3034178 \equiv 3^{89240} \pmod{2^{23} - 1}$ such that $x^2 \equiv 1 \pmod{2^{23} - 1}$. Now we factorize the previous and next number. $(2^{23} - 1) -3034178 -1 = 5354428 = 2 \times 2 \times 19 \times \mathbf{47} \times 1499$ $(2^{23} - 1) -3034178 +1 = 5354430 = 2 \times 3 \times 5 \times \mathbf{178481}$ And we have got the factorization of $2^{23} - 1$. My questions: Is this procedure known? Can you replicate the procedure and confirm its correctness? Is there a quick way to find non-trivial x such that $x^2 \equiv 1 \pmod{2^{p} - 1}$, where $2^{p} - 1$ is composite?

论坛问题帖 | has_answers | DaBler | 2025-01-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019395 | MSE-5032746

A closed form for $\int \frac{dx}{1+x^{4n}}$.

If $f(x)$ is polynomial then $\bar{f}(x)$ is obtained from f by changing the sign of coefficient x. I was trying to find a pattern in the anti derivative of $\frac{1}{1+x^m}$ and I noticed an interesting pattern when m is a multiple of 4 or equivalently $m=4n, \ n \in \mathbb{N}$. -- First Observation: For $n \in \mathbb{N}$ it seems that there always exist second degree polynomials $p_{1,n}, p_{2,n}, p_{3,n}, \dots, p_{n,n}$ $a_{k,n}$ the the coefficient of x in $p_{k,n}$ and first degree polynomials $q_{1,n}, q_{2,n}, q_{3,n}, \dots, q_{n,n}$ be with $b_{k,n}$ the the coefficient of x in $q_{k,n}$ then $$\int \frac{dx}{1+x^{4n}}= \sum_{k=1}^n \left(a_{k,n}\frac{\ln(p_{k,n}(x))-\ln(\bar{p}_{k,n}(x))}{8n}\right)+\sum_{k=1}^n \left(\frac{\arctan(q_{k,n}(x))-\arctan(\bar{q}_{k,n}(x))}{4b_{k,n}n}\right)$$ Second observation: It seems to be that $b_{n,k}=\frac{2}{a_{k,n}}$ I also believe that $q_{k,n}$ can be obtained from $p_{k,n}$ but I couldn't find a way to obtain $q_{n,k}$'s absolute term. Questions Is there a way to obtain $q_{n,k}$ from $p_{n,k}$? How to prove that for all $n \in \mathbb{N}$ there exist second degree polynomial $p_{1,n}, p_{2,n}, p_{3,n}, \dots, p_{n,n}$ $a_{k,n}$ the the coefficient of x in $p_{k,n}$ and there exist a first degree polynomial $q_{1,n}, q_{2,n}, q_{3,n}, \dots, q_{n,n}$ be with $b_{k,n}$ the the coefficient of x in $q_{k,n}$ then $$\int \frac{dx}{1+x^{4n}}= \sum_{k=1}^n \left(a_{k,n}\frac{\ln(p_{k,n}(x))-\ln(\bar{p}_{k,n}(x))}{8n}\right)+\sum_{k=1}^n \left(\frac{\arctan(q_{k,n}(x))-\arctan(\bar{q}_{k,n}(x))}{4b_{k,n}n}\right)$$ How to find $p_{n,k}$ and $q_{n,k}$?

论坛问题帖 | no_answers_at_retrieval | pie | 2025-02-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019396 | MSE-5033452

How many times a number n is repeated in pascal's triangle?

I was playing with Pascal's triangle and noticed an interesting pattern: most numbers (except 1) appear only once (e.g., 2, 3, 4) or twice (e.g., $6 = \binom{4}{2} = \binom{6}{1}$), when excluding duplicates within the same row. To investigate this further, I wrote a simple Python script to count how many times each number appears in Pascal's triangle: import math import matplotlib.pyplot as plt import numpy as np a = [] b = [] for n in range(2, 10000): count = 0 for k in range(n + 1): for j in range(math.floor(k / 2) + 1): if math.comb(k, j) > n: break if math.comb(k, j) == n: count += 1 b.append(count) a.append(n) plt.plot(np.array(a), np.array(b)) plt.show() The first number that appears three times is 210: $$ 210 = \binom{10}{4} = \binom{21}{2} = \binom{210}{1} $$ The first number to appear four times is (3003): $$ 3003 = \binom{14}{6} = \binom{15}{5} = \binom{78}{2} = \binom{3003}{1} $$ No number is repeated five times in (1,30000) I conjecture that for any $ N \in \mathbb{N} $, there exists a number $ n $ that appears exactly $ N $ times in Pascal’s triangle. However, I have been unable to prove or disprove this. Is this conjecture true? If false, what is the maximum number of times a number can appear in Pascal’s triangle? Additionally, Assuming this conjecture holds I am curious about the sequence $ a_n $, where $ a_n $ is the first number to appear $ n $ times in Pascal’s triangle. The known values seem to be: $$ 2, 6, 210, 3003, \dots $$ This sequence does not appear in the OEIS, and finding a closed formula seems difficult due to the chaotic nature of the sequence. Can we find an upper bound on $ a_n $? Empirically, $ a_n $ seems to grow rapidly. can we find a function f such that $a_n =O(f(n))$

论坛问题帖 | no_answers_at_retrieval | pie | 2025-02-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019397 | MSE-5039304

General prime gap counter function remains "invariant under scaling" of three of its parameters by square-free number coprime to two of its parameters

Definitions. Let all variables be integers. When we sum over the set of residues r such that $r^2 = k^2 \mod \text{lcm}(\ell,d)$, we're allowed to take any residue $r' \in r + \text{lcm}(\ell,d)\Bbb{Z}$ in place of r in the formula. This is true via construction (basic inequality solving / modular arithmetic), and the ceiling term takes care of anything that would otherwise be "too large" in magnitude. For square-free $l, n \geq 1, \ k \geq 0, x\in \Bbb{Z}$, we define the following counting formula. $$ H_n(k,\ell,x) =\sum_{d \mid \text{rad}(n)}\mu(d)\sum_{r^2 = k^2 \mod \text{lcm}(\ell,d)}\left(\left\lfloor\frac{x - r}{\text{lcm}(\ell,d)}\right\rfloor - \left\lceil\frac{-r}{\text{lcm}(\ell,d)} \right\rceil + 1\right) $$ It is known that for $x \geq 0$, this counts EXACTLY the number of $z \in [0, x]$ such that $z^2 \neq k^2 \mod \text{lcm}(\ell, p)$ for any p prime dividing n. And therefore $H_{1,1}(p_n\#, x)$ for $x \in [p_n +2, p_{n+1}^2-2]$ (sieve of Eratosthenes bounds) will count EXACTLY the number of $z \in [p_n +2, x]$ that are twin prime averages. The reason for "$p_n + 2$" there is that the lower averages got sieved out as we incremented the primorial. Conjecture. Fix $n\geq 1$. Let $l\geq 1$ with $\gcd(l,n) =1$, $k\geq 0$, and $x \in \Bbb{Z}$. For any prime q such that $q \nmid n\ell$ we have that, the $H_n(k,\ell,x)$ expression remains invariant under scaling of the parameters k, ℓ, x by q. In other words $$ H_n(qk,q\ell, qx) = H_n(k,\ell,x) $$. Attempt. I figured out a nicer, more expressive proof method. If we set $A(k,d) = \{r \in \Bbb{Z}/d : r^2 = k^2\}$, then we can easily prove that $A(qk, q\ell d) \simeq A(k, \ell d)$ as sets (of solutions). Using an unnamed property of modular arithmetic, that should be familiar to most modular arithmetic students. Namely, the "cancelling out a factor in the modulus": $$ r^2 = k^2 \mod \ell d \\ \iff \\ r^2-k^2 = 0 \mod \ell d\\ \iff \\ q^2(r^2 - k^2) = 0 \mod q\ell d \\ \iff \\ (qr)^2 = (qk)^2 \mod q\ell d $$ Then if $\text{res}(A(k, \ell d))\subset \Bbb{Z}$ is a choice of residues for the solutions $r \in A(k, \ell d)$, then $q\cdot \text{res}(A(k,\ell d)) = \text{res}(A(qk,q\ell d))$ is a possible choice of residues for $r \in A(qk, q\ell d)$, because $r^2 = k^2 \mod \ell d \iff (qr)^2 = (qk)^2 \mod q\ell d$. Thus, we may transform the expression for $H_n(qk, q\ell, qx)$ in one step. Edit. I was finally able to prove this on my own (for the most part). But there is still an unanswered question that I have about it. Proof of Conjecture. Since $\gcd(q, \ell d) = \gcd(\ell, d) = 1$ we may replace every lcm in the formula by multiplication. $$ H_n(qk,q\ell,qx) = \sum_{d \mid \text{rad}(n)}\mu(d)\sum_{r^2 = (qk)^2 \mod q\ell d}\left(\left\lfloor\frac{qx - r}{q\ell d} \right\rfloor - \left\lceil\frac{-r}{q\ell d} \right\rceil + 1 \right) \\ = \sum_{d \mid \text{rad}(n)}\mu(d)\sum_{r^2 = k^2 \mod \ell d}\left(\left\lfloor\frac{qx - qr}{\ell d} \right\rfloor - \left\lceil\frac{-qr}{\ell d} \right\rceil + 1 \right) \\ = H_n(k,\ell, x) $$ The summation rearrangement is via the above lemma. Question. Why must q be a prime

number in this proof strategy, and not just coprime (to ℓn)? I know from code-testing that any coprime q, i.e. any $q\in \Bbb{N}$ such that $\gcd(q, \ell n) = 1$ doesn't work. But when q is a prime, the test code runs to completion without an assert (counter-example). Conjecture Verification Code. from sympy import * def H(k, l, n, x): S = 0 for d in divisors(n): T = 0 for r in range(l*d): if (r ** 2 - k ** 2) % (l * d) == 0: T += floor((x-(r % (l*d)))/(l*d)) - ceiling((-r % (l*d))/(l*d)) + 1 S += totient(d) * T return S N = 8 M = 16 x = Symbol('x') for n in range(1, N): print(f"n={n}:") P = primorial(n) for k in range(0, M): for l in range(1, M): if gcd(l, P) == 1: for m in range(n+1, M): # Already coprime to p_n# q = prime(m) if gcd(q, l) == 1: h_k_l_x = H(k, l, P, x) h_qk_ql_qx = H(q*k, q*l, P, q*x) print(f" H({q}*{k},{q}*{l},p_{n}#,{q}*{x}) = {h_qk_ql_qx}") print(" =? ") print(f" H({k},{l},p_{n}#,{x}) = {h_k_l_x}") if h_k_l_x != h_qk_ql_qx: print("CONJECTURE FAILURE") assert 0 print('------------------------') Extended Conjecture. For any $k \geq 0$ and $\ell \geq 1$ such that $\gcd(\ell, p_n\#) = 1$, we have that, for $\gcd(q,\ell p_n\#) = 1$: $$ H_n(k,q\ell,x) \leq H_n(k,\ell, x) = H_n(qk,q\ell,qx) $$ Conjecture verification was for small cases using the SymPy code and a few extra lines. Note. I didn't want to make another question for this, so perhaps there is some proof mechanism by which this extended conjecture also goes through. I think by construction (examining the set-wise inclusion-exclusion formula) we have the inequality.

论坛问题帖 | no_answers_at_retrieval | TotoposDiagramChaseApp | 2025-02-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019398 | MSE-5042113

Does $(a-c)>(b-a)$ imply that $(s-a)^2 + c^2 < (s-b)^2 + a^2$ for integer valued $0 \leq a, b, c \leq s.$

In a paper on theoretical voting systems, I have come to the need to prove (or disprove) the following statement: If \begin{equation} (a-c)>(b-a), \end{equation} then \begin{equation} (s-a)^2 + c^2 < (s-b)^2 + a^2, \end{equation} for all non-negative integer values $0 \leq a, b, c \leq s.$ This holds for all of the combinations of $a, b, c,$ and s that I have tried. I have tried to do a proof by contradiction but am unable to derive a contradiction. Any help would be greatly appreciated!

论坛问题帖 | has_answers | fauxfaux | 2025-03-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019399 | MSE-5042750

Congruences between CM Modular forms of different weight for special primes

While playing around with modular forms with CM, I found something interesting: Set $f_{27.2.a.a}:=q\prod_{n=1}^\infty (1-q^{3n})^2(1-q^{9n})^2=\sum_{n=1}^\infty \alpha_n q^n$ to be a cusp form of $\Gamma_0(27)$, weight 2 with $\mathbb Q(\sqrt{-3})$ CM. Set $f_{9.4.a.a}:=q\prod_{n=1}^\infty (1-q^{3n})^8=\sum_{n=1}^\infty \beta_n q^n$ to be a cusp form of $\Gamma_0(9)$, weight 4 with $\mathbb Q(\sqrt{-3})$ CM. It is easy to see when $n\equiv 0,2\bmod 3$ we have $\alpha_n=\beta_n=0$, so we are only interested in the case $n\equiv 1\bmod 3$, especially when $n=p\equiv 1\bmod 3$ is a prime. I try to compare $\alpha_p,\beta_p\bmod p$ for $p\le 5000$. And I conjecture that $\alpha_p\equiv\beta_p\bmod p$ if and only if $(4p-1)/27$ is an odd perfect square. These p-s form the OEIS sequence A275878, namely standard Jacobi primes. Is there some number theory facts behind this?

论坛问题帖 | has_answers | cybcat | 2025-03-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019400 | MSE-5047374

Computing $\int_0^{\pi(1-1/N)}\mathrm{d}\phi\,\left(\frac{\sin^N(\phi+\pi/N) -\sin^N(\phi)}{\sin^N(\phi+\pi/N) +\sin^N(\phi)}\right)^2$

I have managed to come up with the conjecture $$I_2(N) = \int_0^{\pi(1-1/N)}\mathrm{d}\phi\,\left(\frac{\sin^N(\phi+\pi/N) -\sin^N(\phi)}{\sin^N(\phi+\pi/N) +\sin^N(\phi)}\right)^2 = \pi \left(\frac{N-1}{N}\right)\left(\frac{2N-1}{3N}\right).$$ This holds trivially for $N=2$. In the limit of $N\rightarrow\infty$, the conjecture also holds: by performing an expansion, keeping only the lowest order terms, and substituting $u=\cot\phi$, I can write $$\lim_{N\rightarrow\infty}I_2(N) = \int_{-\infty}^\infty \mathrm{d}u \, \frac{\tanh^2(u\pi/2)}{1+u^2},$$ which I can solve using contour integration. For $N=4$, I can also show consistency with the ansatz. Expanding $\sin(\phi+\pi/N)$ with the $\sin$ addition formula and making the substitution $u=1+\cot\phi$, I can write $$I_2(4) = \int_0^\infty \mathrm{d}u\,\frac{1}{u^2-2u+2}\left(\frac{4-u^4}{4+u^4}\right)^2 =\int_0^\infty \mathrm{d}u\, f(u).$$ Annoyingly, this is not the integral of even function. Contour integration is therefore not easy. However, I can split the integrand into odd and even components $f(u) = f_e(u) + f_o(u)$. For $N=4$, these components are fortunately possible to evaluate with contour integration and some minor manipulations. I have attempted something similar for general N, but have so far been unsuccessful, as $f_o(u)$ no longer takes such a convenient form. I am very confident that this conjecture is true for $N>1$. I have performed numerical integration up to $N=50$, and it is in perfect agreement with my conjecture even for non-integer values of N (although I am specifically interested in the solutions for positive integers $N>2$), as you can see in the figure. A general proof of this, however, has been elusive, and I would appreciate any suggestions. I have also tried differentiation of my expression with respect to N to see if I can show equality that way. Unfortunately, as far as I can tell, the integral only becomes more unpleasant. N." />

论坛问题帖 | has_answers | scqueen | 2025-03-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019401 | MSE-5047802

If matrix A has a left inverse C, is AC orthogonally diagonalizable with all eigenvalues 0 or 1?

Let A,C be matrices such that $CA = I$. Then AC is defined. I conjecture the following: AC is orthogonally diagonalizable. Its eigenvalues are either 1 or 0. It is Hermitian Informally, if A has a left inverse C, then switching their order can either preserve information as is (eigenvalue 1) or lose it (eigenvalue 0), but can't do anything else with it (transform it otherwise). Is the above correct? Is this part of a theory?

论坛问题帖 | has_answers | SRobertJames | 2025-03-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019402 | MSE-5049175

Conjecture about gcd of Faulhaber's Polynomials.

Now asked on MO here Let $f_n $ be the $ n $-th Faulhaber polynomial. In this question, I observed the following divisibility properties: If $ n $ is even, then $ f_n $ is divisible by $ f_2 $. If $ n > 1 $ is odd, then $ f_n $ is divisible by $ f_3 $. However, it seems that Faulhaber polynomials do not share any additional common factors (ignoring the denominator). More precisely, for $ n \neq m $ and $n,m>1$, we have $$ \gcd(f_n, f_m) = \begin{cases} f_3, & \text{if both } n \text{ and } m \text{ are odd,} \\ f_2, & \text{if both } n \text{ and } m \text{ are even,} \\ f_1, & \text{otherwise.} \end{cases} $$ Beyond these cases, no additional common factors seem to exist. Added: As @sil mentioned in the comments $\gcd(f_9,f_{10})= n (n+1) \left(n^2+n-1\right)$ However for all $n,m\le100$ that is the only counter example that I found. I used this ugly Mathematica code to generate the gcds For[m = 1, m < 101, m++, For[j = 1, j < m + 1, j++, If[m != j, Print[m, j, Factor[PolynomialGCD[Sum[k^m, {k, 1, n}], Sum[k^j, {k, 1, n}]]]] ,]]] Added: Using the code in this answer I have checked that for $n,m<1000$ there are no other counter examples. Here are the first 14 Faulhaber's Polynomials: $$f_1=\frac{n(n+1)}{2}$$ $$ f_{2}=\frac16 n (1+n) (1+2 n)$$ $$f_{3}=\frac14 n^2 (1+n)^2$$ $$f_{4}=\frac1{30} n (1+n) (1+2 n) (-1+3 n+3 n^2)$$ $$f_{5}=\frac1{12} n^2 (1+n)^2 (-1+2 n+2 n^2)$$ $$f_{6}=\frac1{42} n (1+n) (1+2 n) (1-3 n+6 n^3+3 n^4)$$ $$f_{7}=\frac1{24} n^2 (1+n)^2 (2-4 n-n^2+6 n^3+3 n^4)$$ $$f_{8}=\frac1{90} n (1+n) (1+2 n) (-3+9 n-n^2-15 n^3+5 n^4+15 n^5+5 n^6)$$ $$f_{9}=\frac1{20} n^2 (1+n)^2 (-1+n+n^2) (3-3 n-n^2+4 n^3+2 n^4)$$ $$f_{10}=\frac1{66} n (1+n) (1+2 n) (-1+n+n^2) (-5+10 n+3 n^2-11 n^3+2 n^4+9 n^5+3 n^6)$$ $$f_{11}=\frac1{24} n^2 (1+n)^2 (10-20 n-3 n^2+26 n^3-5 n^4-16 n^5+4 n^6+8 n^7+2 n^8)$$ $$f_{12}=\frac1{2730}n (1+n) (1+2 n) (-691+2073 n-287 n^2-3285 n^3+1420 n^4+2310 n^5-1190 n^6-1050 n^7+525 n^8+525 n^9+105 n^{10})$$ $$f_{13}=\frac1{420} n^2 (1+n)^2 (-691+1382 n+202 n^2-1786 n^3+367 n^4+1052 n^5-326 n^6-400 n^7+125 n^8+150 n^9+30 n^{10})$$ $$f_{14}=\frac1{90} n (1+n) (1+2 n) (105-315 n+44 n^2+498 n^3-217 n^4-345 n^5+182 n^6+144 n^7-81 n^8-45 n^9+24 n^{10}+18 n^{11}+3 n^{12})$$

论坛问题帖 | has_answers | pie | 2025-03-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019403 | MSE-5050467

How to find all minimal upper bounds under graph minor relation?

Let $H\leq G$ denote that H is a minor of G. Given two graphs H_{1} and H_{2}, is there an algorithm to find all minimal upper bounds G of H_{1},H_{2} ? That is, find all graphs G such that $H_{1},H_{2}\leq G$ and $H_{1},H_{2}\leq H\leq G$ implies $H\cong G$. By the graph minor theorem, the set of such G is finite up to isomorphism. I conjectured that $|V(G)|\leq |V(H_{1})|\times |V(H_{2})|$, but I failed to prove it. There is an example that the branch sets of G corresponding to H_{1} and H_{2} share at least two vertices, but I can't construct a counter example to the conjecture either.

论坛问题帖 | no_answers_at_retrieval | 林浩誼 | 2025-03-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019404 | MSE-5050738

Radium Rabbit Conjecture (Version 2.0)

Radium Rabbit Conjecture (Version 2.0) Why This Version 2.0? The previous version of this conjecture led to some misunderstandings, and the discussion was closed before reaching a conclusion. In this revised version, I have clarified my observations and the open question I seek to answer. I hope this will allow for a more precise discussion and lead to a deeper understanding of the underlying mathematical structure. Radium Rabbit Conjecture The conjecture posits that, for a triangle formed by distinct prime numbers, the square of its area will always have a fractional part of $3/16=0.1875$. Specifically, the square of the area follows the form: $$S^2 = A + \frac{3}{16}$$ where A is some integer. Key Observations Modular Arithmetic Connection It has been established that the square of the area follows a modulo 16 pattern related to $3/16$. However, what has been proven so far is only that the result is a multiple of 3, not necessarily fixed at $3/16$ itself. I want to understand why the fraction is always exactly $3/16$ and not, for example, $6/16$ or $9/16$. Distinct Prime Triangles and Their Properties Such triangles do not form right-angled triangles due to the nature of prime numbers. They are neither congruent nor similar to each other, further suggesting a unique underlying property. Equilateral Triangle and a Possible Clue The squared area of an equilateral triangle follows a specific modular arithmetic pattern where the fractional part is $3/16$. This hints at a deeper structure that might explain why distinct prime triangles also conform to the same $3/16$ fraction. Open Question While I have observed and verified the $3/16$ pattern, I have yet to prove why this must always be the case. The key question I am seeking to answer is: Why does the squared area of a triangle formed by distinct primes always take the form , rather than any other multiple of $3/16$? I appreciate any insights or discussions that could help in proving this conjecture. Final Note I truly enjoy discussing mathematics with all of you, and I appreciate the opportunity to refine this idea further. Thank you!

论坛问题帖 | has_answers | Radium Rabbit | 2025-03-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019405 | MSE-5052714

Has anyone constructed a graph that shows implications between conjectures?

Does such a graph exist or is it considered redundant/unhelpful/infeasible? Such a graph could help fill in corollary results from a proof or its contrapositive. Relatedly, I have found some data aggregation for theorems and directed graphs for mathematical concepts or fields. I think that a 'conjectree' would also be useful for discovering bottlenecks, or for the deliberate exploration of problems that have fewer known consequences (leaves?).

论坛问题帖 | has_answers | Lem n | 2025-04-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019406 | MSE-5052749

Conjecture: For every prime p, there exists at least one prime in the range $[p^3-p, p^3+p]$

Conjecture: For every prime p, there exists at least one prime in the range $[p^3-p, p^3+p]$. Verified for all primes up to 1000. Let me know if it holds for all. Example Let’s work through the conjecture for all prime numbers up to 17, calculating p^3, the interval $[p^3 - p, p^3 + p]$, and confirming whether at least one prime exists within it, inclusive. The primes up to 17 are: $2, 3, 5, 7, 11, 13, 17$. Here’s the step-by-step verification: $p = 2$: $2^3 = 8, 8 - 2 = 6, 8 + 2 = 10$. Interval: $[6, 10] = \{6, 7, 8, 9, 10\}$. Primes: 7 Prime exists: Yes. $p = 3$: $3^3 = 27, 27 - 3 = 24, 27 + 3 = 30$. Interval: $[24, 30] = \{24, 25, 26, 27, 28, 29, 30\}$. Primes: 29 Prime exists: Yes. $p = 5$: $5^3 = 125, 125 - 5 = 120, 125 + 5 = 130$. Interval: $[120, 130] = \{120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130\}$. Primes: 127 Prime exists: Yes. $p = 7$: $7^3 = 343, 343 - 7 = 336, 343 + 7 = 350$. Interval: $[336, 350] = \{336, 337, 338, 339, 340, 341, 342, 343, 344, 345, 346, 347, 348, 349, 350\}$. Primes: $337, 347$ Prime exists: Yes. $p = 11$: $11^3 = 1331, 1331 - 11 = 1320, 1331 + 11 = 1342$. Interval: $[1320, 1342] = \{1320, 1321, 1322, ..., 1341, 1342\}$ (23 numbers). Primes: $1321, 1327$ Prime exists: Yes. $p = 13$: $13^3 = 2197, 2197 - 13 = 2184, 2197 + 13 = 2210$. Interval: $[2184, 2210] = \{2184, 2185, ..., 2209, 2210\}$ (27 numbers). Primes: $2203, 2207$ Prime exists: Yes. $p = 17$: $17^3 = 4913, 4913 - 17 = 4896, 4913 + 17 = 4920$. Interval: $[4896, 4920] = \{4896, 4897, ..., 4919, 4920\}$ (25 numbers). Primes: $4903, 4919$ Prime exists: Yes.

论坛问题帖 | has_answers | Arun Vangapalli | 2025-04-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019407 | MSE-5052753

Primality and the GCD of Stirling number of the second kind.

Let $ \left\{ \begin{matrix} n \\ k \end{matrix} \right\} $ denote the Stirling number of the second kind. $$ \sum_{k=0}^n {n \brace k} x^{\underline{k}} = x^n $$ where $ x^{\underline{k}} $ is the falling factorial. Also, the Stirling number can be expressed as: $$ \left\{\begin{array}{l} n \\ k \end{array}\right\}=\frac{1}{k!} \sum_{j=0}^k(-1)^{k-j}\binom{k}{j} j^n=\sum_{j=0}^k \frac{(-1)^{k-j} j^n}{(k-j)!j!} $$ $$x=x^{\underline{1}}$$ $$x^2 = x^{\underline{1}}+x^{\underline{2}} $$ $$x^{\color{red}{3}} = x^{\underline{1}} + \color{red}{3}x^{\underline{2}} +x^{\underline{3}} $$ $$x^4=x^{\underline{1}} +7 x^{\underline{2}} + 6x^{\underline{3}} +x^{\underline{4}} $$ $$x^{\color{red}{5}}=x^{\underline{1}}+\color{red}{15}x^{\underline{2} } +\color{red}{25}x^{\underline{3}} +\color{red}{10}x^{\underline{4}} +x^{\underline{5}}$$ $$x^6=x^{\underline{1}}+31x^{\underline{2} }+90x^{\underline{3}} +65x^{\underline{4}}+15x^{\underline{5}}+ x^{\underline{6}}$$ $$x^{\color{red}{7}}=x^{\underline{1}}+\color{red}{63}x^{\underline{2} }+\color{red}{301}x^{\underline{3}} +\color{red}{350}x^{\underline{4}}+\color{red}{140}x^{\underline{5}}+ \color{red}{21}x^{\underline{6}}+x^{\underline{7}}$$ I came across an interesting conjecture: For $n>2$ $$ \gcd\left(\left\{ \begin{matrix} n \\ 2 \end{matrix} \right\},\left\{ \begin{matrix} n \\ 3 \end{matrix} \right\},\dots, \left\{ \begin{matrix} n \\ n-1 \end{matrix} \right\} \right) > 1 \quad \text{if and only if} \quad n \text{ is prime}. $$ Furthermore, if $ n $ is prime, then $$ \gcd\left(\left\{ \begin{matrix} n \\ 2 \end{matrix} \right\},\left\{ \begin{matrix} n \\ 3 \end{matrix} \right\},\dots, \left\{ \begin{matrix} n \\ n-1 \end{matrix} \right\} \right) = n. $$ I was unable to prove this result rigorously, but I wrote a Python script to check it computationally, and the conjecture holds up to $ n = 10,000 $. Here is my implementation: import math stop=1 def is_prime(n): if n <= 1: return False for i in range(2, int(math.sqrt(n)) + 1): if n % i == 0: return False return True a=[1] max=10_000 nmax=int(max*(max+1)/2) for i in range (2,max+1): for n in range (1,nmax+1): if n<math.floor((i*(i-1))/2)+1: continue elif n==math.floor((i*(i-1))/2)+1 : a.append(1) elif n==math.floor((i*(i+1))/2) : a.append(1) break else: m=n-(math.floor((i*(i-1))/2)) s=math.floor(((i-2)*(i-1))/2)+m-1 a.append(a[s-1] +(m)*a[s]) for i in range (3,max+1): if stop ==0: print("false") break s=a[math.floor((i*(i-1))/2)+1] for n in range (1,nmax+1): if n<=math.floor((i*(i-1))/2)+1: continue elif n>=math.floor((i*(i+1))/2) : if s>1 and s==i : if is_prime(i)==True: print(i) else: stop=0 break else: s=math.gcd(s,a[n-1])

论坛问题帖 | has_answers | pie | 2025-04-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019408 | MSE-5056284

A Generalization of Hilbert's Nullstellensatz?

This question comes from the (intuitive) observation, that the failure of Hilbert's Nullstellensatz comes from the failure of the base field k to "detect" all roots of a polynomial. In the case of an algebraically closed field, all roots of any polynomial can be successfully detected. Therefore it seems to be natural to ask the following question, a generalization of the Nullstellensatz on non-algebraically closed fields: Question: if k is any characteristic zero field, $S \subseteq k[x_1,\cdots,x_n]$ a set of polynomials that can be factored into linear factors in $k[x_1,...,x_n]$, does it hold that $I(Z(S))=\sqrt{\langle S \rangle}$? Notations: $Z(S)$ is the set of zero points; $I(X)$ is the set of polynomials that vanishes on X; $\langle S \rangle$ is the ideal generated by S; $\sqrt I$ is the (nilpotent) radical. Edit: it's actually not a generalization.

论坛问题帖 | has_answers | atzlt | 2025-04-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019409 | MSE-5061293

A conjecture about multiplying odd numbers by 3 in binary

Setup : Let $I_n=\underbrace{11\cdots1}_{n \text{ times}}$ and $Z_n=\underbrace{00\cdots0}_{n \text{ times}}$ Example: $x=11001=I_2 \space Z_2 \space I_1$ Note: We are dealing only with odd numbers ! Motivation behind the problem : To compute $3 \times x$, the classical way is: $2x+x$ But it requires addition and bit-shifting ! So, I asked myself : Can we find an algorithm that computes $3\times x$ with no adders, no carry logic ? Here's the algorithm I've made to answer this question ! My algorithm is purely a rearrangement of the input bits under fixed rules. My algorithm and the conjecture : First, let

$\color{blue}{\zeta(n)} = \begin{cases} 01Z_{n-2}10 \space \text{if } n\ge2 \\ \\ 100 \text{ if } n=1\end{cases}$ and $x=I_{n_k}Z_{m_k} \cdots I_{n_2} Z_{m_1}I_{n_1}$ such that $\forall i>0: n_i,m_i>0$ To compute $3x$, first prepare the table below : $\color{red}{10}\space\space\space I_{n_k-2} \space\space \color{blue}{\zeta(m_k)}$ $\space\space\space\space\space\space\space I_{n_{k-1}-2} \space\space \color{blue}{\zeta(m_{k-1})}$ $\space\space\space\space\space\space\space \cdots$ $\space\space\space\space\space\space\space I_{n_2-2} \space\space \color{blue}{\zeta(m_1)}$ $\space\space\space\space\space\space\space I_{n_1-2} \space\space \color{red}{01}$ To start the algorithm, we assemble the digits from $\color{red}{01}$ up to $\color{red}{10}$ The behaviour of I_{-1} : If I_{-1} was before 1, we delete it and that's it ! If I_{-1} was before 0, we delete it and apply the inv function The inv func works as follows: If I_{-1} was after $\color{red}{10}$, he converts it to $\color{red}{1}$ and that's it ! If I_{-1} was after $\color{blue}{100}$, he converts it to $\color{blue}{011}$ and that's it ! If I_{-1} was after $\color{blue}{01Z_{m_{i}-2}10}$, he converts it to $\color{blue}{01Z_{m_{i}-2}01}$ and that's it ! Example: Suppose $x=11001=I_2 \space Z_2 \space I_1$ First let's prepare our table ! $\color{red}{10}\space\space\space I_{2-2}=I_0 \space\space \color{blue}{\zeta(2)}=0110$ $\space\space\space\space\space\space\space I_{1-2}=I_{-1} \space\space \color{red}{01}$ Starting the algorithm from $\color{red}{01}$ up to $\color{red}{10}$ we get: $\color{red}{10}\space\space\space I_0 \space\space \color{blue}{\operatorname{inv}(}0110\color{blue})=0101$ $\space\space\space\space\space\space\space .. \space\space \color{red}{1}$ Assembling, we get $3x=\color{red}{10} \space 0101 \space \color{red}{1}$ $\Rightarrow 3x=1001011$ Another example: $x=10011101=I_1 \space Z_2 \space I_3 \space Z_1 \space I_1$ First let's prepare our table ! $\color{red}{10}\space\space\space I_{1-2}=I_{-1} \space\space \color{blue}{\zeta(2)}=0110$ $\space\space\space\space\space\space\space I_{3-2}=I_{1} \space\space \color{blue}{\zeta(1)}=100$ $\space\space\space\space\space\space\space I_{1-2}=I_{-1} \space\space \color{red}{01}$ Starting the algorithm from $\color{red}{01}$ up to $\color{red}{10}$ we get: $\color{red}{1}\space\space\space .. \space\space 110$ $\space\space\space\space\space\space\space I_{1} \space\space \color{blue}{\operatorname{inv}(}100\color{blue})=011$ $\space\space\space\space\space\space\space .. \space\space \color{red}{1}$ Assembling, we get $3x=\color{red}{1} \space 110 \space 1 \space 011 \space \color{red}{1}$ $\Rightarrow 3x=111010111$ Now the conjecture is : $\forall x \in 2\mathbb{N}+1,$ my algorithm will always produce $3x$? Thanks for help !

论坛问题帖 | has_answers | Lhachimi | 2025-05-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019410 | MSE-5062159

Conjecture about a chain of six circles whose tangent points lie on a circle

Six black circles (not necessarily the same size) form a closed chain, in which each circle is externally tangent to its two neighbors. The six tangent points all lie on a red circle. Three line segments connect the centres of opposite black circles. Based on experimentation, I conjecture that the three line segments must intersect at a single point. Is my conjecture true? I tried to use circle inversion, but that does not seem to make the problem any easier. Note that Brianchon's Theorem does not apply here, because the red circle is not necessarily normal to the black circles, so the red circle is not necessarily tangent to the sides of the hexagon with vertices at the black circles' centres. If there are eight black circles instead of six, the four line segments do not necessarily intersect at a single point. This question was inspired by this question about a chain of four circles inscribed in a circle. Edit A theorem similar to my conjecture is Dao's Theorem. But they are different: In my conjecture, each of the six circles must be tangent to its two neighbors, but this is not required in Dao's theorem. (At the end of page 40 in the link, it says "Let (O_{ijk}) meets (O_{jkh}) again at P_{jk}'". Later in "Problem 6", they make use of the points P_{jk}'.) In Dao's theorem, each of the six circles goes through P_{ij},P_{jk},P_{ik} for some consecutive i,j,k, but this is not required in my conjecture.

论坛问题帖 | has_answers | Dan | 2025-05-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019411 | MSE-5076199

What is the status of the Palis conjecture?

Jacob Palis (born in 1940) sadly passed away last month. I read the lead of his Wikipedia article, and it says: "He proposed the Palis conjecture, which influenced the development of the theory of dynamical systems, and also of its applications to other sciences. He was a world leader in chaos theory research. Palis was an influential figure in the development of mathematics in Brazil." (emphasis is mine) What is the Palis conjecture? In which paper was it proposed? Is it still open? What is the progress that have been made so far? Why is it important? I am only an undergraduate student of math, interested in dynamical systems theory, and curious about this. EDIT: Judging by other topics in MathOverFlow, it's not only one conjecture, but many conjectures ("the Palis conjectures" or "Palis' Program"). I am now very confused about this.

论坛问题帖 | no_answers_at_retrieval | Viktor K. | 2025-06-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019412 | MSE-5077906

$n \mid 5\times2^{v_2(3n+1)}+9$, has no solutions ? ($\forall n>1$ and $n\equiv 1\pmod 4$)

The Conjecture: Let $n\in \mathbb{N^*}$ then $n,v_2(3n+1)>1$$\Rightarrow n \not\mid 5\times2^{v_2(3n+1)}+9$ Notice that, $v_2(3n+1)>1 \Leftrightarrow 4 \mid 3n+1 \Leftrightarrow n\equiv 1\pmod 4$ My attempt : $v_2(3n+1)=2 \Leftrightarrow 3n +1 \equiv 4 \pmod 8 \Leftrightarrow n\equiv 1 \pmod 8$ Suppose $n\mid 5\times2^{v_2(3n+1)} +9=5\times2^2+9=29$ $\Rightarrow n = 29$, but $29\equiv 5 \pmod 8$ $v_2(3n+1)=3 \Leftrightarrow 3n +1 \equiv 8 \pmod {16} \Leftrightarrow n\equiv 13 \pmod {16}$ Suppose $n\mid 5\times2^{v_2(3n+1)} +9=5\times2^3+9=49$ $\Rightarrow n = \begin{cases} 7 ,

\text{but it's not } 13 \pmod {16} \\ \\ 49, \text{but } 49\equiv 1 \pmod{16} \end{cases}$ In general, I managed to find the following congruences : $v_2(3n+1)=p \Leftrightarrow n \equiv \begin{cases} \displaystyle \frac{2^p-1}{3} \pmod {2^{p+1}} \text{ , if } p \equiv 0 \pmod 2 \\ \\ 3+10 \times \displaystyle \frac{2^{p-1}-1}{3} \pmod {2^{p+1}} \text{ , if } p \equiv 1 \pmod 2 \end{cases}$ Now suppose $n < 2^{p+1}$, Case one: p even ! $\Rightarrow n = \displaystyle \frac{2^p-1}{3} \Rightarrow 2^p-1\mid 3(5\times2^p +9) = 3(5\times(2^p - 1) + 14) \Rightarrow 2^p-1\mid 21 \Rightarrow n \mid 7 \Rightarrow n=7$ but $2^p-1=21 \Rightarrow 2^p=22$, which is a contradiction ! Case two: p odd ! $\Rightarrow n = 3+10 \times \displaystyle \frac{2^{p-1}-1}{3} \Rightarrow 5\times 2^p -10 + 9 \mid 3(5\times2^p+9)$ $\Rightarrow 5\times2^p -1 \mid 3(5\times2^p-1 + 10) \Rightarrow 5\times2^p -1 \mid 3 $ but $5\times2^p -1=3 \Rightarrow 5\times2^p=4$, which is a contradiction ! Thus, one have to check my conjecture only $\forall n>2^{p+1}$. I've also checked the conjecture $\forall n<10^6$, by python : def val_2(n): return (n & -n).bit_length() - 1 for n in range(5,pow(10,6),4): if (5*pow(2,val_2(3*n+1))+9)%n == 0: print(f'Contradiction at n={n}') Thanks for help !

论坛问题帖 | has_answers | Lhachimi | 2025-06-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019413 | MSE-5077942

Conjecture : $\forall d>1, d \mid 3^{n+1}-2^n \Rightarrow \log_2(3d+1) \not\in \mathbb{Z}$?

The conjecture : Let $n \in \mathbb{N^*} \Rightarrow \forall k >1, \displaystyle \frac{4^k-1}{3} \not\mid 3^{n+1}-2^n $ Note that : the statement above is equivalent to the title by considering $d=\displaystyle\frac{4^k-1}{3}$ we are dealing with a d such that $\gcd(d,3)=1$ because $3\mid d \Rightarrow 3\mid 3^{n+1}-2^n$, an immediate contradiction ! $\Rightarrow v_3(4^k-1)=1$, so that $\gcd(\displaystyle\frac{4^k-1}{3},3)=1$ Also, by LTE : $v_3(4^k-1)=v_3(4-1)+v_3(k)=1+v_3(k)=1, \Rightarrow v_3(k)=0$ In other words, $\gcd(d,3)=1 \Leftrightarrow \gcd(k,3)=1$ My attempt : $\displaystyle\frac{4^k-1}{3} \mid 3^{n+1}-2^n \Rightarrow 2\mid n$ and $k\equiv 1 \pmod 3$ Suppose n is odd, we will use the Jacobi symbol : $\displaystyle \left(\frac{3^{n+1}}{d} \right)$ $=$ $\displaystyle \left(\frac{2^n}{d} \right)\Rightarrow \displaystyle \left(\frac{2}{d} \right) = +1$ On the other hand, we have this identity :$\displaystyle \left(\frac{2}{d} \right) = (-1)^{\displaystyle \frac{d^2-1}{8}} = (-1)^{\displaystyle \frac{d-1}{4}\times \displaystyle \frac{d+1}{2}} $ $d-1=\displaystyle \frac{4^k-1}{3}-1 = \displaystyle \frac{4^k-4}{3} \Rightarrow \displaystyle \frac{d-1}{4} = \frac{4^{k-1}-1}{3}$, an odd integer $d+1=\displaystyle \frac{4^k-1}{3}+1 =\displaystyle \frac{4^k-2}{3} \Rightarrow \displaystyle \frac{d+1}{2} = \frac{2^{2k-1}-1}{3}$, an odd integer $\Rightarrow \displaystyle \left(\frac{2}{d} \right) = -1 $, which is a contradiction ! $\Rightarrow n$ is even As n is even $\Rightarrow \displaystyle \left(\frac{3}{d} \right) = +1 \Rightarrow \left(\frac{d}{3} \right) = (-1)^{\displaystyle \frac{d-1}{2} \times \frac{3-1}{2}} = +1$ Notice that, $d=\displaystyle \sum_{i=0}^{k-1} 4^i \equiv \displaystyle \sum_{i=0}^{k-1} 1^i \pmod 3 $, since $4 \equiv 1 \pmod 3$ $\Rightarrow d \equiv k \pmod 3 \Rightarrow k \equiv 1 \pmod 3$, since $\displaystyle \left(\frac{d}{3} \right)=+1$ Thanks for help !

论坛问题帖 | has_answers | Lhachimi | 2025-06-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019414 | MSE-5078510

Conjecture : $\forall n>1, d\mid 3^n-2^n \Rightarrow v_2(d+1)<n$

The conjecture : $\forall n>1, d\mid 3^n-2^n \Rightarrow v_2(d+1)<n$ where $v_2(x)$ denotes the 2-adic valuation of x, i.e the highest power of 2 that divides x e.g : $v_2(12)=2$ since $12=2^2 \cdot 3$ Initially, let's check the case when $d=3^n-2^n$ $\Rightarrow v_2(d+1)=v_2(3^n+1-2^n)$ Notice that, $v_2(3^n+1)=\begin{cases} 1 \text{ , if } n \equiv 0 \pmod 2 \\ \\ 2 \text{ , if } n \equiv 1 \pmod 2 \end{cases}$ $\Rightarrow v_2(d+1) =\begin{cases} v_2(2^1\times(\displaystyle \frac{3^n+1}{2}-2^{n-1})) = 1 \text{ , if } n \equiv 0 \pmod 2 \\ \\ v_2(2^2\times(\displaystyle \frac{3^n+1}{2^2}-2^{n-2}))=2 \text{ , if } n \equiv 1 \pmod 2 \end{cases} $ You can take $n>2 \Rightarrow v_2(d+1) < n$, the case $n=2$ is correct since 2 is even and $2>1$ Also, the case $d=1$ is trivial since $v_2(d+1)=v_2(1+1)=1<2<n$ @Aphelli noticed that $d\mid 3^n-2^n \Leftrightarrow \displaystyle \left(\frac{3}{2}\right)^n \equiv 1 \pmod d $ So it's enough to prove the following : $\gcd(d,6)=1 \Rightarrow v_2(d+1) < \text{ord}_d \displaystyle \left(\frac{3}{2}\right)$ My attempt : Firstly, we clearly notice that $d\mid 3^n-2^n \Rightarrow d$ is odd $ \Rightarrow 2\mid d+1$ Let's assume by contradiction, that $v_2(d+1)\ge n \Rightarrow 2^n \mid d+1 \Rightarrow d=2^nk-1$ $\Rightarrow 2^nk-1 \mid3^n-2^n$, (for some integer $k \ge 1$) e.g : for $k=1 \Rightarrow 2^n -1 \mid 3^n-2^n=3^n-(2^n-1)-1$ $\Rightarrow 2^n-1 \mid 3^n-1 \Rightarrow n$ is odd , (since $2\mid n \Rightarrow 3\mid 3^n-1$) Let's use the Jacobi symbol : $\displaystyle \left(\frac{3}{2^n-1} \right)=1 \Rightarrow \displaystyle \left(\frac{2^n-1}{3} \right) = (-1)^{\displaystyle\frac{3-1}{2} \times \frac{2^n-1-1}{2}}=-1$ On the other hand, $2^n-1\equiv (-1)^n-1 \equiv -2 \equiv 1 \pmod 3$ $\Rightarrow \displaystyle \left(\frac{2^n-1}{3} \right) = \displaystyle \left(\frac{1}{3} \right) = +1$, which is a contradiction ! From A Conjecture about the upper bound of $v_2(3^n+d)$ (proved by Joris Nieuwveld) we conclude the following : Fix $k \in \mathbb{N^*} \implies 2^nk-1 \mid3^n-2^n$ has only finite solutions for $n>1$ $2^nk-1 \mid3^n-2^n \Rightarrow (2^nk-1)\lambda = 3^n-2^n$, (for some integer $\lambda \ge 1$) $\Rightarrow 2^n(\lambda k +1)=3^n+\lambda \Rightarrow v_2(3^n+\lambda) \ge n$ which won't last longer, since $\forall d \in \mathbb{N^*}, \exists n_0 \in \mathbb{N^*}$ such that : $\forall n>n_0 : v_2(3^n+d)<n$, where n is a positive integer. Also, if someone managed to optimize the constant C_n mentioned on : Analyzing Yu, K.: p-adic logarithmic forms and group varieties we may deduce some tighter bounds on n or λ, which would contradict the fact that $\lambda < 3^n - 2^n$ I've also checked the conjecture $\forall n<10^2$, by python : from sympy import divisors def val_2(n): return (n & -n).bit_length() - 1 for n in range(2,pow(10,2)): for d in divisors(pow(3,n)-pow(2,n)): if val_2(d+1) >= n : print(f'Contradiction at n={n} , d={d}') break Also, setting G_n to be the set of gaps $n-v_2(d+1)$, $\forall d\mid 3^n-2^n $ and $g(n)=\min(G_n)$ we remark that $g(n) > 0$ and $g(n)=\mathcal{O}(n)$ So, as n grows up $v_2(d+1)$ became much smaller than n which is assured by our conjecture ! from sympy import divisors import matplotlib.pyplot as plt def val_2(n): return (n & -n).bit_length() - 1 plt.grid(True) plt.xlabel("n") plt.ylabel("g(n)") for n in range(2,pow(10,2)): G=[] for d in divisors(pow(3,n)-pow(2,n)): G.append(n-val_2(d+1)) plt.scatter(n,min(G),c='black') plt.show() Thanks for help !

论坛问题帖 | has_answers | Lhachimi | 2025-06-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019415 | MSE-5079500

Conjecture : $\forall n>1 , \exists a,b \in \mathbb{N^2}$ such that : $\tau(3^n-2^n)=2^a \cdot3^b$

Motivation : I wanted to understand more the behaviour of the divisors of 3^n-2^n, to solve this conjecture : Conjecture : $\forall n>1, d\mid 3^n-2^n \Rightarrow v_2(d+1)<n$ So I tried to examine initially: $\tau(3^n-2^n)$ (the number of divisors d of 3^n-2^n) My attempt: I literally have no idea of how we can prove this result, but I discover it from my python research routine ! from sympy import divisor_count from sympy.ntheory import factorint for n in range(2,100): print(n,factorint(divisor_count(pow(3,n)-pow(2,n)))) Here's a chart of the factorization of $\tau(3^n-2^n)$, ($\forall n < 10^2 $)

$\color{blue}{2}[\color{green}{2}\color{red}:1]$
$\color{blue}{3}[\color{green}{2}\color{red}:1]$
$\color{blue}{4}[\color{green}{2}\color{red}:2]$
$\color{blue}{5}[\color{green}{2}\color{red}:1]$
$\color{blue}{6}[\color{green}{2}\color{red}:3]$
$\color{blue}{7}[\color{green}{2}\color{red}:2]$
$\color{blue}{8}[\color{green}{2}\color{red}:3]$
$\color{blue}{9}[\color{green}{2}\color{red}:2]$
$\color{blue}{10}[\color{green}{2}\color{red}:2,\color{green}{3}\color{red}:1]$
$\color{blue}{11}[\color{green}{2}\color{red}:1,\color{green}{3}\color{red}:1]$
$\color{blue}{12}[\color{green}{2}\color{red}:5]$
$\color{blue}{13}[\color{green}{2}\color{red}:2]$
$\color{blue}{14}[\color{green}{2}\color{red}:4]$
$\color{blue}{15}[\color{green}{2}\color{red}:3]$
$\color{blue}{16}[\color{green}{2}\color{red}:5]$
$\color{blue}{17}[\color{green}{2}\color{red}:1]$
$\color{blue}{18}[\color{green}{2}\color{red}:5]$
$\color{blue}{19}[\color{green}{2}\color{red}:2]$
$\color{blue}{20}[\color{green}{2}\color{red}:4,\color{green}{3}\color{red}:1]$
$\color{blue}{21}[\color{green}{2}\color{red}:5]$
$\color{blue}{22}[\color{green}{2}\color{red}:3,\color{green}{3}\color{red}:1]$
$\color{blue}{23}[\color{green}{2}\color{red}:2]$
$\color{blue}{24}[\color{green}{2}\color{red}:7]$
$\color{blue}{25}[\color{green}{2}\color{red}:3]$
$\color{blue}{26}[\color{green}{2}\color{red}:5]$
$\color{blue}{27}[\color{green}{2}\color{red}:3]$
$\color{blue}{28}[\color{green}{2}\color{red}:6]$
$\color{blue}{29}[\color{green}{2}\color{red}:1]$
$\color{blue}{30}[\color{green}{2}\color{red}:7,\color{green}{3}\color{red}:1]$
$\color{blue}{31}[\color{green}{2}\color{red}:1]$
$\color{blue}{32}[\color{green}{2}\color{red}:7]$
$\color{blue}{33}[\color{green}{2}\color{red}:4,\color{green}{3}\color{red}:1]$
$\color{blue}{34}[\color{green}{2}\color{red}:4]$
$\color{blue}{35}[\color{green}{2}\color{red}:5]$
$\color{blue}{36}[\color{green}{2}\color{red}:10]$
$\color{blue}{37}[\color{green}{2}\color{red}:2]$
$\color{blue}{38}[\color{green}{2}\color{red}:5]$
$\color{blue}{39}[\color{green}{2}\color{red}:4]$
$\color{blue}{40}[\color{green}{2}\color{red}:8,\color{green}{3}\color{red}:1]$
$\color{blue}{41}[\color{green}{2}\color{red}:4]$
$\color{blue}{42}[\color{green}{2}\color{red}:8,\color{green}{3}\color{red}:1]$
$\color{blue}{43}[\color{green}{2}\color{red}:3]$
$\color{blue}{44}[\color{green}{2}\color{red}:5,\color{green}{3}\color{red}:1]$
$\color{blue}{45}[\color{green}{2}\color{red}:5]$
$\color{blue}{46}[\color{green}{2}\color{red}:6]$
$\color{blue}{47}[\color{green}{2}\color{red}:3]$
$\color{blue}{48}[\color{green}{2}\color{red}:10]$
$\color{blue}{49}[\color{green}{2}\color{red}:4]$
$\color{blue}{50}[\color{green}{2}\color{red}:8]$
$\color{blue}{51}[\color{green}{2}\color{red}:5]$
$\color{blue}{52}[\color{green}{2}\color{red}:7,\color{green}{3}\color{red}:1]$
$\color{blue}{53}[\color{green}{2}\color{red}:1]$
$\color{blue}{54}[\color{green}{2}\color{red}:7]$
$\color{blue}{55}[\color{green}{2}\color{red}:4,\color{green}{3}\color{red}:1]$
$\color{blue}{56}[\color{green}{2}\color{red}:9]$
$\color{blue}{57}[\color{green}{2}\color{red}:4,\color{green}{3}\color{red}:1]$
$\color{blue}{58}[\color{green}{2}\color{red}:5]$
$\color{blue}{59}[\color{green}{2}\color{red}:1]$
$\color{blue}{60}[\color{green}{2}\color{red}:12,\color{green}{3}\color{red}:1]$
$\color{blue}{61}[\color{green}{2}\color{red}:3]$
$\color{blue}{62}[\color{green}{2}\color{red}:5]$
$\color{blue}{63}[\color{green}{2}\color{red}:9]$
$\color{blue}{64}[\color{green}{2}\color{red}:9]$
$\color{blue}{65}[\color{green}{2}\color{red}:6]$
$\color{blue}{66}[\color{green}{2}\color{red}:9,\color{green}{3}\color{red}:1]$
$\color{blue}{67}[\color{green}{2}\color{red}:3]$
$\color{blue}{68}[\color{green}{2}\color{red}:6]$
$\color{blue}{69}[\color{green}{2}\color{red}:8]$
$\color{blue}{70}[\color{green}{2}\color{red}:8,\color{green}{3}\color{red}:1]$
$\color{blue}{71}[\color{green}{2}\color{red}:2]$
$\color{blue}{72}[\color{green}{2}\color{red}:15]$
$\color{blue}{73}[\color{green}{2}\color{red}:3]$
$\color{blue}{74}[\color{green}{2}\color{red}:5]$
$\color{blue}{75}[\color{green}{2}\color{red}:8]$
$\color{blue}{76}[\color{green}{2}\color{red}:7]$
$\color{blue}{77}[\color{green}{2}\color{red}:6,\color{green}{3}\color{red}:1]$
$\color{blue}{78}[\color{green}{2}\color{red}:10]$
$\color{blue}{79}[\color{green}{2}\color{red}:4]$

$\color{blue}{80}[\color{green}{2}\color{red}:11,\color{green}{3}\color{red}:1]$
$\color{blue}{81}[\color{green}{2}\color{red}:6]$
$\color{blue}{82}[\color{green}{2}\color{red}:9]$
$\color{blue}{83}[\color{green}{2}\color{red}:3]$
$\color{blue}{84}[\color{green}{2}\color{red}:13,\color{green}{3}\color{red}:1]$
$\color{blue}{85}[\color{green}{2}\color{red}:4]$
$\color{blue}{86}[\color{green}{2}\color{red}:9]$
$\color{blue}{87}[\color{green}{2}\color{red}:5]$
$\color{blue}{88}[\color{green}{2}\color{red}:9,\color{green}{3}\color{red}:1]$
$\color{blue}{89}[\color{green}{2}\color{red}:2]$
$\color{blue}{90}[\color{green}{2}\color{red}:12,\color{green}{3}\color{red}:1]$
$\color{blue}{91}[\color{green}{2}\color{red}:8]$
$\color{blue}{92}[\color{green}{2}\color{red}:9]$
$\color{blue}{93}[\color{green}{2}\color{red}:4]$
$\color{blue}{94}[\color{green}{2}\color{red}:7]$
$\color{blue}{95}[\color{green}{2}\color{red}:6]$
$\color{blue}{96}[\color{green}{2}\color{red}:15]$
$\color{blue}{97}[\color{green}{2}\color{red}:2]$
$\color{blue}{98}[\color{green}{2}\color{red}:9]$
$\color{blue}{99}[\color{green}{2}\color{red}:7,\color{green}{3}\color{red}:1]$ Thanks for help !

论坛问题帖 | has_answers | Lhachimi | 2025-06-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019416 | MSE-5082437

Proving (the lower bound that) N exceeds product of primes under special congruence constraints

Let $S = \{p_1, p_2, \dots, p_k\}$ be a finite set of at least two distinct odd primes, i.e., $k \ge 2$. Consider the system: $ N \equiv p_i \pmod{p_i^2 (p_i - 1)} \quad \text{for all } p_i \in S, $ where N is square-free. Then how can we prove the following claim: $N > \prod_{i=1}^{k} p_i$ I observed this empirically, but I am unable to give a correct rigorous proof for it (tried 3-4 different methods, getting stuck every time). Can anybody please help me out?

论坛问题帖 | has_answers | Math_Maven | 2025-07-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019417 | MSE-5085637

Conjecture: In a "happy triangle", the expected product of side lengths is π.

Definition of happy triangle On a unit circle, choose three uniformly random independent points A,B,C. Inside the circle, choose a uniformly random independent point D. Triangle ABC is called "happy" if it contains D. Conjecture Is the following conjecture true: In a happy triangle, the expected product of the three side lengths is π. I simulated 2×10^7 happy triangles, and the average product was $3.14184188\approx1.00007933\pi$. My thoughts Letting $X=\text{product of the three side lengths}$, I tried (but failed) to find the probability distribution function $P(X<x)$ so that I could find the probability density function and thus $E(X)$. The expected product of the three side lengths is not the cube of the expected side length, because the three side lengths are not independent random variables. (A simulation suggests that the expected side length is approximately 1.509.) Other elegant properties? A happy triangle seems to have many elegant properties.

论坛问题帖 | has_answers | Dan | 2025-07-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019418 | MSE-5086607

Proving a number of base 2 non-recurring if converted to base 10 having the same decimal point position from the right of the number

I have a precision P_2 and P_{10} defined as the position of the decimal point from the right of the number. P_2 for base 2, P_{10} for base 10, for example: $$ \begin{align*} \\ P_2(110.101_2) &= 3_{10} \\ P_2(1.10101_2) &= 5_{10} \\ \\ P_{10}(110.101_2) &= P_{10}(6.625_{10}) &= 3_{10} \\ P_{10}(1.10101_2) &= P_{10}(1.65625_{10}) &= 5_{10} \end{align*} $$ I would like to prove that $P_2(x)=P_{10}(x)$ for any number of x that can be represented in base 2 without recurring number, e.g. $0.0011..._2$ At the moment , I cannot compute an exception to prove a contradiction, so I am left with this conjecture. I tried to breakdown input as an expanded base 10 equation (unsubscripted terms are in base 10): $$ \begin{align*} a_2 &= (... b2 \cdot 2^{2} + b1 \cdot 2^{1} + b0 \cdot 2^{0}) \times 2^{-p1} \\ a_{10} &= (... b2 \cdot 10^{2} + b1 \cdot 10^{1} + b0 \cdot 10^{0}) \times 10^{-p2} \end{align*} $$ Now, I am left with how to proof if $a_2=a_{10}$, then $p1 = p2$ is the same in both equation. I am not sure if I can work on the factorisation here like: $$ \begin{align*} 10^{-p2} &= (2 \times 5)^{-p2} \\ 10^{-p2} &= 2^{-p2} \times 5^{-p2} \\ \frac{10^{-p2}}{5^{-p2}} &= 2^{-p2} \end{align*} $$ Then, I could substitute 2^{-p2} in that a_2 equation, but I have to handle $\frac{1}{5^{-p2}}$ every other terms in the infinite sum. I have a feeling I am making things more complicated here. What approach can I use to proof this equation? Or is there any properties that I use as a reference that they are equal?

论坛问题帖 | has_answers | dee cue | 2025-07-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019419 | MSE-5093066

There exists no mult. subsemigroup $S \subset\Bbb{Z}$ of a ring such that $S + 1$ is never prime?

By prime number we mean $\Bbb{P} = \{ \pm 1, \pm q : q \text{ is prime in the usual sense}\}$. Conjecture. There exist no mulitiplicative subsemigroup S of $\Bbb{Z}$, such that $S + 1$ is never prime. Any such S which are a counter-example of this are called semigroup spaceships [U+1F6F8] because they perfectly dodge all (quite randomly placed) primes ("astroids"). So in essence, adding 1 to a subsemigroup is never enough to build a nice spaceship, a subsemigroup is not complex enough. Can you provide a counter-example to this claim? Clearly if there exists such an S it must be infinite, because it contains $x

\neq \pm 1$, and so $x^2,x^3, \dots$ as well. It sufficient to prove there is no cyclic subsemigroup $S = \langle x \rangle$ such that $(S + 1) \cap \Bbb{P} = \varnothing$. Or in other words, for all $x \in \Bbb{Z}$ $$ x^n + 1 \in \Bbb{P} \text{ i.o. over } n \in \Bbb{N} $$

论坛问题帖 | has_answers | TotoposDiagramChaseApp | 2025-08-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019420 | MSE-5094437

Can we prove this conjecture related to the trisection of the angle?

I don't have much reference on trisecting an angle , but while trying to design a problem on this topic , I stumbled upon the following simple situation : In triangle ABC, the line segments CI and CJ are the two trisectors of $\angle ACB$. The circumscribed circle of triangle CIJ intersects the line segment CA at C, D, and the line segment BC at C, E. So here is my conjecture that I observed, after many constructions, and I don't know if it's already known or even true: the straight lines AB and DE are PARALLEL. I attempted an analytical approach that turned out to be too complicated, so I continue to search a proof. Any ideas are welcome.

论坛问题帖 | has_answers | Jamil Sanjakdar | 2025-09-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019421 | MSE-5094809

Steiner-Lehmus-like Conjecture: If corresponding trisectors of two angles of a triangle are congruent, then the triangle is isosceles

I am trying to find a property of trisectors that recalls the famous Steiner-Lehmus theorem on bisectors. So, here is, after several observations, a new conjecture that I wish to demonstrate: Conjecture. In ABC, AI and BJ are the two "upper" trisectors of angles A and B respectively, we know that $AI=BJ$, as shown in the figure, then ABC is isosceles. Here is my incomplete attempt at the proof: According to the sine law, we can easily find: $$\sin(3a)\sin(2a+3b) = \sin(3b)\sin(2b+3a) \tag1$$ and after developing and first transposition, we can write $$\cos(a-3b) - \cos(3a-b) = \cos(5a+3b) - \cos(5b+3a) \tag2$$ so $$\sin(a+b)\sin(2a-2b) = \sin(a-b)\sin(4a+4b) \tag3$$ and $$\sin(a-b)\left[\cos(a-b) - 2\cos(a+b)\cos(2a+2b)\right] = 0 \tag4$$ Finally, or else $a=b$, then ABC isosceles, or else $$\cos(a-b) - 2\cos(a+b)\cos(2a+2b) = 0 \tag5$$ that we can write as $$\cos(a-b) - \cos(a+b) - \cos (3a+3b) = 0 \tag6$$ and ultimately $$2\sin(a)\sin(b) + \cos(C) = 0\text{ with }C=\pi-3a-3b \tag7$$ This last equation is impossible if angle C is acute. So the conjecture is true in any acute ABC, but then, what about an obtuse triangle? Any suggestions?

论坛问题帖 | has_answers | Jamil Sanjakdar | 2025-09-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019422 | MSE-5096720

What is the probability that a plane through three random points on the surface of the Earth intersects the Moon?

Looking up at the night sky, I wonder: What is the probability that a plane through three random points on the surface of the Earth intersects the Moon? (The points are independent and uniform.) Let: $d=$ distance between centre of the Earth and centre of the Moon $r=$ radius of the Moon (Figure not to scale.) Simulations suggest that the probability is simply r/d, for any values of r and d, assuming that the two spheres are disjoint. (It seems that the radius of the Earth does not matter.) If this is true, then plugging in $r=1737$ km and $d=384400$ km, the probability is approximately $\frac{1}{221}$. Using sphere point picking, each random point has two variables, so there are a total of six variables. It seems that such a simple result should be provable without integrating over six variables, but I cannot find a way. The $2D$ case is not elegant. Fun fact The probability is indeed r/d. If we replace "moon" with "sun", the probability is approximately $\frac{7\times 10^5}{1.496\times 10^8}\approx\frac{1}{214}$. So there is a greater chance that the plane will intersect the sun, than the moon.

论坛问题帖 | has_answers | Dan | 2025-09-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019423 | MSE-5097283

Are $1,2,5,10$ the only solutions of $3^{n+1} \pmod {2^n} > 2^n-1.5^n$?

Introduction : Put $r_n=2^n \{ 1.5^n \} $, i.e : the remainder of $3^n \pmod {2^n}$. If the following inequality holds: $\forall n>1 , r_n<2^n-1.5^n$ (an open problem). It has been proved that the function $g(k)$ occurring in Waring's problem, is given by the formula : $g(k)= 2^k +\left \lfloor{\displaystyle 1.5^k} \right \rfloor -2$ This was noted by J. A. Euler, the son of Leonhard Euler, in about 1772 My attempt : I wondered what if we considered the next power 3^{n+1} , will the inequality stay true ? However , I checked the conjecture with PARI/GP , $\forall n < 10^5$: { for(n=1, 10^5, if (lift(Mod(3, 2^n)^(n+1)) > 2^n - (3/2)^n, print(n))) } It took around 41.44 seconds to finish , the only solutions were $\color{red}1,\color{red}2,\color{red}5,\color{red}{10}$.

论坛问题帖 | no_answers_at_retrieval | Lhachimi | 2025-09-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019424 | MSE-5099042

Conjecture: The largest root of $\sum_{k=0}^n\cos\left(\frac{k\pi}{2}\right)\binom{n}{k}x^k$ is approximately $\frac{n}{\pi}$ or $\frac{2n}{\pi}$.

Is the following conjecture true: For large n, the largest root of $\displaystyle\sum\limits_{k=0}^n\cos\left(\frac{k\pi}{2}\right)\binom{n}{k}x^k$ is approximately $\dfrac{n}{\pi}$ for odd n, or $\dfrac{2n}{\pi}$ for even n. For example, for $n=100$ we have $1-\binom{100}{2}x^2+\binom{100}{4}x^4-\binom{100}{6}x^6+\dots-\binom{100}{98}x^{98}+x^{100}$. The largest root is $x\approx 63.6567$, whereas $\frac{2(100)}{\pi}\approx63.6620$. Their ratio is approximately

0.99992. I do not know why $\frac{2(100)}{\pi}$ is close to a root, nor why that root is the largest root. Context: This question was inspired by my question "True or false: If you square the coefficients in the expansion of $(x+1)^n$, the resulting polynomial has n distinct real roots."

论坛问题帖 | has_answers | Dan | 2025-09-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019425 | MSE-5100277

Why does $\sum_{p\leq n}\sigma_{0}\left(p-1\right)\approx2n$?

I noticed that: $$\sum_{p\leq n}\sigma_{0}\left(p-1\right)\approx2n$$ where σ_{0} is the number-of-divisor function, and the sum is over primes. Any idea if and why this could hold for larger n ? And don't have any clue where to start, since it was a random observation while tracking the number of operations for some unrelated algorythm... I ran a script and got the following results: n SUM 10 10 100 160 1000 1804 10000 19069 100000 191593 1000000 1926814 10000000 19324386 100000000 193577243 1000000000 1937069910

论坛问题帖 | has_answers | François Huppé | 2025-10-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019426 | MSE-5101240

Conjecture on upper bound for sum of $\sum_{a \in S} 1/a$ for S satisfy $\mathrm{lcm}(a,b) > n$ for every $a \neq b \in S$

Recently I found an interesting problem on number theory which stated For a set S of integers ranging from 1 to n such that for every $a\neq b,~a,b \in S$ we have that $\mathrm{lcm}(a,b) > n$, prove that $\sum_{a \in S} 1/a < 3/2$. The solution for this problem is quite simple. Denote S_a the set of multiples less than n of a; then for every $a\neq b, a,b \in S$ we must have $S_a \cap S_b = \emptyset$ and obtain the "trivial" bound of $n \ge \sum |S_a| \ge \sum (n/a - 1)$. However when I want to make the constant $3/2$ lower the same strategy no longer works. I believe that the best constant possible is something around $\ln 2$ as n approaches infinity and the set not containing 1. A few examples are: the set of the least power of p greater than $\sqrt n$ for every prime $p \le n,$ or the set of all numbers greater than $n/2$, both result in a sum of $\ln 2$ (as n approaches infinity, since for $n=5$ there is already a counter example of $S=\{2,3,5\}$). A very similar problem has been conjectured by Erdős before, namely problem of finding the best upper bound for $\sum_{a\in S} 1/a$, but for the set S satisfying a is not divisible by b for every $a,b \in S$. But I couldn't find a source of this problem online. If anyone could share on how to tackle this problem or just the name of it, I'd be very appreciative.

论坛问题帖 | no_answers_at_retrieval | jackdean | 2025-10-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019427 | MSE-5101812

Do the area, semiperimeter and longest side of a cyclic quadrilateral always satisfy the triangle inequality?

My experimental data supports the following conjecture. Can this be proved? Conjecture: Let a convex quadrilateral be inscribed in a circle of radius $R>0$. Let a denote the longest side, s the semiperimeter, and A the area of the quadrilateral. Then the numbers $a,s,\frac{A}{R}$ satisfy the triangle inequality. Update 17-Oct-2025: I think I have a proof, posting it as answer below. Generalization of the question for cyclic polygons.

论坛问题帖 | has_answers | Nilotpal Kanti Sinha | 2025-10-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019428 | MSE-5103692

A similar sequence to OEIS-A098021

Let $U_0=0$, we make U_n by the following rules : If $\color{magenta}n=U_k$ for some $k \in \mathbb{N}$, $$U_{n+1} = U_{\color{magenta}n} + \color{red}5$$ $$U_{n+2} = U_{n+1} + \color{red}5$$ and $\forall \alpha \in \mathbb{N^*}$ such that $n+2\alpha +1 \le U_{k+1}$ we have : $$ U_{n+2\alpha+1}= U_{n+2\alpha}+\color{green}7 $$ $$ U_{n+2\alpha+2}= U_{n+2\alpha+1}+\color{blue}5 $$ That is : we add $\color{red}5$ twice if $\color{magenta}n=U_k$, otherwise we keep adding $\color{green}7$ and $\color{blue}5$ successively Here's how the first 20 terms looks like : $$ U_{0}=0$$ $$ U_1 =U_{\color{magenta}0}+\color{red}5 =5$$ $$ U_2 = U_1+\color{red}5=10$$ $$ U_3 = U_2+\color{green}7=17$$ $$ U_4 = U_3+\color{blue}5=22$$ $$ U_{5}=U_4+\color{green}7=29$$ $$ U_{6}=U_{\color{magenta}5}+\color{red}5=34$$ $$ U_{7}=U_{6}+\color{red}5=39$$ $$ U_{8}=U_{7}+\color{green}7=46$$ $$ U_{9}=U_{8}+\color{blue}5=51$$ $$ U_{10}=U_{9}+\color{green}7=58$$ $$ U_{11}=U_{\color{magenta}{10}}+\color{red}5=63$$ $$ U_{12}=U_{11}+\color{red}5=68$$ $$ U_{13}=U_{12}+\color{green}7=75$$ $$ U_{14}=U_{13}+\color{blue}5=80$$ $$ U_{15}=U_{14}+\color{green}7=87$$ $$ U_{16}=U_{15}+\color{blue}5=92$$ $$ U_{17}=U_{16}+\color{green}7=99$$ $$ U_{18}=U_{\color{magenta}{17}}+\color{red}5=104$$ $$ U_{19}=U_{18}+\color{red}5=109$$ $$ U_{20}=U_{19}+\color{green}7=116$$ plugging these terms to OEIS , I found that they're similar to A098021 ! So the question is , can you prove that : $$\bbox[1px,border:1px solid #0a0]{\bbox[8px,border:1px solid #0a0]{U_n=3n+2 \left \lfloor{\displaystyle n\sqrt2} \right \rfloor}}$$

论坛问题帖 | has_answers | Lhachimi | 2025-10-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019429 | MSE-5104244

Does decomposition of PDFs guarantee independence of random variables?

Is this conjecture correct? If not, can it be modified to a correct one: Let X,Y be continuous RVs with joint PDF $f(x,y)$. Then X,Y are independent iff there exists functions g, h such that $$f(x,y) = g(x)h(y).$$ This seems obviously true if g and h are the PDFs of X,Y. I conjecture this to be true without making that qualification: any g,h that multiply to f are

sufficient. But I don't know how to prove that.

论坛问题帖 | has_answers | SRobertJames | 2025-10-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019430 | MSE-5106839

A conjecture about a difference expression of n positive numbers

The starting point for this question is a set of n positive ordered numbers: $$ x_{1} \, \ge \, x_{2} \, \ge \, \ldots \, \ge x_{n} \, > \, 0 \; .$$ From these numbers difference expressions can be formed: $$ D \, = \, \sum_{i=1}^{n}\, c_{i} x_{i} \; .$$ Each c_{i} coefficient obeys $c_{i} \in \{-1, 1\}$ and so for a fixed n, there are 2^{n} possible difference expressions. I define the class of guaranteed non-negative difference expressions as follows. Definition A difference expression D is said to be guaranteed non-negative (gnn) if its value is non-negative regardless of the values of the x_{i} variables. For example, it is fairly easy to see that the following difference expressions are gnn: A difference expression D with the first $\lceil n/2 \rceil$ coefficients c_{i} equal to 1 and the remaining c_{i} equal to -1 is gnn. Hence, when $m = \lceil n/2 \rceil$, the following is gnn: $$ D_{1} \, = \, \sum_{i=1}^{m}\, x_{i} - \sum_{i=m+1}^{n}\, x_{i} $$ A difference expression where the c_{i} coefficients alternate between 1 and -1, starting with $c_{1} = 1$, is gnn. Hence, $$ D_{2} \, = \, \sum_{i=1}^{n}\, (-1)^{i+1} \, x_{i}$$ is gnn. My question concerns the following conjecture about difference expression D_{2}. Conjecture There does not exist a gnn difference expression $\hat{D}$ such that $D_{2} > \hat{D}$. My problem is to prove or disprove this conjecture. I believe it is correct. I have a couple of remarks. Remark 1 I think I have a proof that the following are necessary and sufficient conditions for a difference expression D to be gnn: $$\sum_{i =1}^{j} \, c_{i} \, \ge 0 \; , \; j = 1, 2, \ldots n $$ Remark 2 My question is related to the 2-way number partitioning problem. See the following: Cerquides, J. and Meseguer, P. 'Speeding Up 2-way Number Partitioning', ECAI, 2012. doi:10.3233/978-1-61499-098-7-223

论坛问题帖 | has_answers | F Cameron | 2025-11-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019431 | MSE-5109453

$a_1,\dots,a_n$ periodic sequences summing to $p_1,\dots,p_n$ over each of their resp. periods, then their sums synch. to some value $\leq\sum_i p_i$.

Conjecture. Let $(a_i(j))_{j \geq 0}$ be sequences of natural numbers ≥ 1. For example $a_1 = \overline{2} = 2,2,2,2, \dots$, is the constant 2, but $a_3 = \overline{2,1,2}$ is not. Define $B = B(a_1, \dots, a_n)$ to be the first point (minimum B) at which the sequences synchronize in sum, ie. where $\sum_{x = 0}^{k_i} a_i(x) = B,$ for some $k_i \in \Bbb{N} \cup 0$, $\forall i = 1..n$. Then, it's true that $$B \leq 2(p_1 + \dots + p_n)$$ is an upper bound for B, where $p_i = \sum_{x = 0}^{T_i - 1} a_i(x)$ is the sum of the sequence a_i over a single period $T_i =$ the period of a_i. Question. How can we prove this via any means (they don't have to be elementary style methods)? Attempt. I thought we could do it by induction on the number of sequences involved. So by induction we have: $$ \forall J \subset \{1, \dots, n\} $$ of size $|J| = n-1$, we would have $\sum_{j \in J} a_j^{k_j} = B(a_j : j \in J) \leq \sum_{j \in J} 2p_j$, where $p_j = \sum_{x = 0}^{T_j} a_j(x)$ and $T_j =$ the period of a_j. So for example if $n = 3$ we have: $$ B(a_1, a_2) \leq 2(p_1 + p_2)\\ B(a_2, a_3) \leq 2(p_2 + p_3) \\ B(a_1, a_3) \leq 2(p_1 + p_3) \\ $$ But I'm not sure how to conclude that $B(a_1, a_2, a_3) \leq 2(p_1 + p_2 + p_3)$ as a result... So perhaps maybe induction is out of the question?

论坛问题帖 | has_answers | TotoposDiagramChaseApp | 2025-11-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019432 | MSE-5109464

Rational points and sections on a family of genus-2 hyperelliptic curves

I am interested in the proof or disproof of some conjectures about rational points and sections over $\mathbb{Q}$ for the following family of genus-2 hyperelliptic curves: $$ C_t: f(x,a)\, g(x,a)\, h(x,a)\ = y^2, $$ Where, $$f(x,a)=\left(x^2+2ax-\left(a^2+4\right)\right)$$ $$g(x,a)=\left(a^2x^2-2ax+\left(2-a^2\right)\right)$$ $$h(x,a)=\left(\left(2-a^2\right)x^2+4ax-8\right)$$ And the parameter is, $$a(t)=\left(\frac{t^2+2t-1}{t^2+1}\right)$$ $\text{where } x, y, t \in \mathbb{Q}$ The following rational sections are known to lie on C_t , $$ S1:\left(x,y\right)=\left(\frac{(t^2+2t+3)(3t^2-2t+1)} {2(t^2+1)(t^2+2t-1)} ,\pm \frac{(-1-2t+t^2)^7} {8(1+t^2)^5(-1+2t+t^2)^2} \right) $$ $$ S2:\left(x,y\right)=\left(\frac{t^2-2t+3}{2(t-1)} ,\pm \frac{(t^2-2t-1)^3(t^2+2t-1)^2} {8(t-1)^3(t^2+1)^2} \right) $$ $$ S3:\left(x,y\right)=\left(\frac{3t^2+2t+1}{2t(t+1)} ,\pm \frac{(-1-2t+t^2)^3(-1+2t+t^2)^2} {8t^3(1+t)^3(1+t^2)^2} \right) $$ A rational section means a point $(x(t),y(t)) \in C_t(\mathbb{Q}(t))$ Edit The original form of Conjecture 2 stated that there were no rational points on the nonsingular fibres other than those arising from the three known sections. Arlo provided a counterexample showing this to be false. The conjecture has therefore been revised to include the lifting condition, which is the condition relevant to the original Diophantine problem. The conjectures are: Apart from the points at infinity of the smooth projective model, there are no other nontrivial affine rational sections of C_t apart from $S1$,$S2$ and $S3$. For nonsingular fibers $(a \neq \pm 1)$, there are no rational points on $C_t(\mathbb{Q})$ satisfying the lifting condition, other than those arising from the three known sections. Origin of the question: This family of genus-2 curves emerges from the 3x3 magic squares of squares problem. Hence, these conjectures have some correspondence with the problem. Clarification about the quotient curve The curve C_t arises from a quotient transformation of the original Diophantine problem. Therefore, non-trivial rational points on C_t give candidate solutions, but they must also satisfy the lifting condition to correspond to solutions of the original problem. The quotient coordinate is related to the original coordinate by $$ x=\frac{1}{k}+2k. $$ Hence a rational quotient point lifts to a rational k only when $$ 2k^2-xk+1=0 $$ has a rational solution, equivalently when $$ x^2-8 $$ is a square in $\mathbb{Q}$. Therefore, the intended question is whether there exist any additional rational points on C_t satisfying this lifting condition, beyond those obtained from the known sections. Evidence from a Specialisation at $t=-7$ I examined the specialized fibre at $t=-7$, $a = a(-7)$. Using Magma, the Jacobian of the specialized curve satisfies the rank bound $$ \operatorname{rank}\!\left(\mathrm{Jac}(C_{-7})(\mathbb{Q})\right) \le 4. $$ A search found only: the unique point at infinity; the specializations of the three sections $(S1, S2, S3)$, each with both choices of sign for y. No further rational points were found. This

computation is not intended as a proof of the conjectures, but provides some computational evidence that the three known sections may account for all rational points. I would especially appreciate suggestions on applicable methods (for example, Chabauty-Coleman, Mordell-Weil sieves, descent, or covering collections), or references to related computations.

论坛问题帖 | has_answers | Anonymous-math-guest | 2025-11-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019433 | MSE-5111248

A “block” generalization of factorions

I’ve been exploring a generalization of factorions-that is, numbers equal to the sum of factorials of their digits-and wanted some feedback on the concept. Let $n \ge 1$ where n is an integer. A number N is an n-factorion if you can split N into consecutive groups of n digits each, take the factorial of these n-digit blocks and sum them, and they all sum to N. Keep in mind this does require N to have a number of digits divisible by n. By this logic, a typical factorion would be considered a 1-factorion. For example, consider 2,763 as a candidate 2-factorion. It would turn into $27! + 63!$ which is equal to approximately 1.98×10^{88} which is pretty obviously way off. As you can see, 2-factorions alone would be monstrously big, the smallest candidate possible being 3,628,800 (10!). Clearly the smallest one would have to be hundreds or thousands of digits. I personally conjecture that there is no upper bound to how large n can be. The only thing holding regular factorions back was the fact that 9! can only work for so many digits, however we can keep theoretically increasing up to 99!, 999!, etc. I’d love feedback on whether this generalization is interesting, or if there are known results related to this topic. Are there any theoretical approaches for estimating or bounding the size of n-factorials without brute-force computation?

论坛问题帖 | has_answers | Sumaesioso | 2025-11-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019434 | MSE-5111817

Does there exist a number such that the first 16 numbers of its Collatz sequence mod 4 contain four 1s, 2s, 3s, and 4s?

I was setting a sudoku with a very unique constraint and I came across this. I thought it wouldn't be that hard to find a number, but after writing a python program to find one, I am doubtful of its existence. Let $C(n)$ be a 16 element sequence representing the first 16 numbers of n's Collatz sequence mod 4. Let $\DeclareMathOperator{Count}{Count}\Count(n)=(\Count_0(n),\Count_1(n),\Count_2(n),\Count_3(n))$ where $\Count_k(n)$ is the number of times k appears in $C(n)$. Using my program, I found that for $1\le n<100000$, if $\Count_3(n)=4$ and $\Count_k(n)\ge3$ for $0\le k\le2$, then $\Count(n)=(3,3,6,4)$. Since there are 3903 values of n that fit these constraints in this range and they all produce the same $C(n)$, I am tempted to conjecture that these conditions on n force $\Count(n)=(3,3,6,4)$. I've tried to reverse engineer this using my basic knowledge of modular arithmetic to no avail. Can anyone help me out? Edit: Here is the program I have been using: def collatz16(n): seq = [0]*16 seq[0] = n%4 for i in range(1, 16): n = (n >> 1) if (n & 1) == 0 else 3*n + 1 seq[i] = n%4 return seq c = 0 for i in range(100000): seq = collatz16(i) if((seq.count(0) >= 3) and (seq.count(1) >= 3) and (seq.count(2) >= 3) and (seq.count(3) == 4)): c+=1 print(f"{i} {[seq.count(0), seq.count(1), seq.count(2), seq.count(3)]} {seq}") print(c)

论坛问题帖 | has_answers | Dylan Levine | 2025-12-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019435 | MSE-5111840

If H, K and L are subgroup such that H commutes with K and L then is $H\ast K\ast L$ a subgroup?

If H and K are subgroup of a group $(G,\ast,e)$ then I know that $H\ast K$ is a subgroup of when H is commutable with K: so I am searching a counterexample showing that if a subgroup X commutes with the subgroup X, Y and Z then $X\ast Y\ast Z$ is not necessarly a subgroup when only X commutes with X and Y but I was not able to find it (I just tried to work with S_3 but I failed) so that I thought to put here a specific question. Could someone help me, please?

论坛问题帖 | has_answers | Antonio Maria Di Mauro | 2025-12-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019436 | MSE-5112131

Is the area of a cyclic polygon of nine or more sides greater than the square of the geometric mean of the sides?

The side lengths of a convex polygon do not uniquely determine the shape of the polygon but if the polygon is cyclic then the shape is uniquely determined by the side lengths. Consider a cyclic n-gon. For $n=3,4$ Heron's formula and Brahmagupta's formula give the area but for $n \ge 5$ there is no such simple formula in terms of the sides. So instead, we can ask given the side lengths $a_1, a_2,\ldots, a_n$, $n \ge 5$ what are the bounds on the area A. Let $\displaystyle s = \frac{\Sigma a_i}{2}$ be the semi-perimeter. Then the isoperimetric inequality gives a trivial upper bound $\displaystyle A \le \frac{s^2}{\pi}$. The lower bound is much more complicated however experimental results shows the possibility of an interesting an simple lower bound for $n \ge 9$. Definition: Let $a = (a_1 a_2 \cdots a_n)^{1/n}$ be the geometric mean of the sides. I estimated the probability $P_n := P\left(A \ge a^2\right)$. My experimental data gave $$ P_3 = 0 \\ P_4 \approx 0.5556 \\ P_5 \approx 0.9961 \\ P_6 \approx 0.99995 \\ P_7 \approx 0.9999995 \\ P_8 \approx 0.99999999 $$ For $n = 9$, even after 5×10^9, I could not find a single polygon where $A < a^2$. Based on this data, I conjecture that Conjecture: The area of a cyclic polygon of nine or more sides is greater than the square of the geometric mean of the sides. Can this conjecture be proved or disproved?

论坛问题帖 | has_answers | Nilotpal Kanti Sinha | 2025-12-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019437 | MSE-5112300

Conjecture: a constant-free equation is solvable in a group G if and only if it is solvable in its generating set B for any $G=\langle B\rangle$

Conjecture. Let G be a group and B any set of generators for G. That is to say $G = (G, \cdot) = \langle B \rangle$. Then for any equation $E=F$ in G that is constant-free, we have that $E=F$ is solvable in general elements $x_i \in G \iff$ it is solvable purely in the generators $b_i \in B$. An equation $E=F$ that is constant-free in G means that we merely use the group law $\cdot$ and/or the inverse operation $(\cdot)^{-1}$, and variables to form the equation. And no constant symbols are allowed! And one cannot "get rid" of 1 by rewriting with $yy^{-1} (\equiv 1)$; that's not allowed. In other words both sides of the equation $E = F$ must be "reduced words." That takes care of the the previous counterexample by Malady of $E:x=1$ having only 1 as a solution. For example, if $x^2y^3z^{-4} = 1$ has a solution $(x,y,z) \in G^3$ then there exists $(a,b,c) \in B^3 : a^2b^3c^{-4} = 1$ and (vise versa obviously). Where juxtaposition means the group operation $\cdot:G\times G \to G$. Question. Can you prove this or come up with a counterexample? If you can't prove it; can you come up with a modified version that you can prove (such as restricting to abelian groups)? Or alternatively, if you do come up with a counter-example, can you modify the statement in such a way that we bypass the given counterexample?

论坛问题帖 | has_answers | TotoposDiagramChaseApp | 2025-12-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019438 | MSE-5112523

Given an n-gon, by how much must we increase the length of each of k chosen sides so that they form a k-gon?

The polygon inequality states that the sum of any $n-1$ sides of a n-gon greater than the n-th side. Let $n \ge 4$ and $3 \le k < n$. Let a n-gon have positive side lengths $ a_1 \le a_2 \le \dots \le a_n$. Select a subset $b_1, b_2, \ldots, b_k$ of the n lengths $a_i, i = 1,2,\ldots, n$. These k sides may or may not form a k-gon themselves depending on weather they satisfy the k-gon inequality or not. Now we increase the length of each of these k by a fixed quantity t. As t increases, the variance of the resulting k increased lengths $b_i + t$ decreases and for sufficiently large t all the increased length will be approximately equal and we can form near regular k-gon. Thus we can ask, what is the minimum t so that the lengths $b_i + t$ will always form a polygon for every subset of sides $b_1, b_2, \ldots, b_k$. Experimental observation: Let s be the semiperimeter of the n-gon. If the k largest sides of the n-gon are selected then $$ t_{\min} = \frac{s(n-k)}{(k-2)(n-1)} \tag 1 $$ If we select any k side of the n-gon then $$ t_{\min} = \frac{s}{k-2} \tag 2 $$ Can these be proved or disproved? Corollary: As an example, $n=4, k=3$ implies that if a,b,c are the three largest sides of a quadrilateral whose semiperimeter is s then the three numbers $$ a + \frac{s}{3}, \quad b + \frac{s}{3}, \quad c + \frac{s}{3} $$ always satisfy the triangle inequality even if a,b,c do not but with any quantity lesser than $\displaystyle \frac{s}{3}$ the triangle inequality may fail.

论坛问题帖 | has_answers | Nilotpal Kanti Sinha | 2025-12-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019439 | MSE-5113220

Does $\sum_{n=0}^{\infty}\frac{(4n)!^2}{2^{8n}(n+1)^2(2n)!^4}$ have the proposed closed form?

Context While working with the $_4F_3(a,b,c,d;e,f,g;x)$ I arrived to: $$S=\frac{64}{9}\left(\hspace{.1cm} _4F_3(-\frac{3}{4},-\frac{3}{4},-\frac{1}{4},-\frac{1}{4};-\frac{1}{2},-\frac{1}{2},1;1)-1\right)=\sum_{n=0}^{\infty}\frac{(4n)!^2}{2^{8n}(n+1)^2(2n)!^4}\tag{1}.$$ Wich seems to evaluate to: $$S=\frac{16\sqrt{2}\Gamma^2{(\frac{1}{4})}}{27\pi^{3/2}}+\frac{32\sqrt{2\pi}}{3\Gamma^2{(\frac{1}{4})}}+\frac{352}{27\pi}-\frac{64}{9}\tag{2}.$$ Attempt Since it can be deduced from the Binomial Theorem: $$\sum_{n=0}^{\infty}\frac{x^{n}(4n)!}{2^{4n}(2n)!^2}=\frac{\sqrt{1+\sqrt{1-x}}}{\sqrt{2}\sqrt{1-x}},\hspace{.5cm}|x|<1.\tag{3}$$ Then integrating both sides of (3): $$\sum_{n=0}^{\infty}\frac{x^{n+1}(4n)!}{2^{4n}(n+1)(2n)!^2}=\frac{8}{3}-\frac{2\sqrt{2}(\sqrt{1-x}+1)^{3/2}}{3},\hspace{.5cm}|x|<1.\tag{4}$$ In view of the form of (1) suggests apply Parseval's Theorem to (4) when $x=-ke^{ix}$. Then: $$I(k)=\sum_{n=0}^{\infty}\left(\frac{(-k)^{n+1}(4n)!}{2^{4n}(n+1)(2n)!^2}\right)^2=\frac{1}{2\pi}\int_{-\pi}^{\pi}|\frac{8}{3}-\frac{2\sqrt{2}(\sqrt{1+ke^{ix}}+1)^{3/2}}{3}|^{2}dx.$$ I have some trouble with the last integral. Question How can we prove the proposed closed form? Thanks in advance for your efforts.

论坛问题帖 | has_answers | User-Refolio | 2025-12-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019440 | MSE-5114802

Is there a simple proof of this metric relation in this Sangaku?

Here's a sangaku problem I've been imagining but I don't know if it's already known : (ABCD) is a parallelogram whose diagonals intersection at O . If : a , b , c and d are the radii of the circles inscribed in triangles (BCD) ; (OAD) ; (OAB) and (ABC) respectively , then I conjecture the metric relation : ad = 2bc . which I was able to verify using GeoGebra by repeating the attached configuration several times. Is there a simple proof based on the basic relationships in triangles and parallelograms ? I wonder , and I'm searching

论坛问题帖 | has_answers | Jamil Sanjakdar | 2025-12-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019441 | MSE-5115533

Conjecture: If a tangential quadrilateral's incenter is its lamina centroid, then it must be a kite.

For symmetry reasons it is clear that rhombi do have the "incenter = lamina centroid" property. It is also not difficult to find some special kites having this property. And here the open problem begins: I am rather sure that no other quadrilaterals can have this property. I could not find a proof (neither a geometric nor an analytic one) for the following conjecture: Conjecture: If for a tangential quadrilateral "incenter = lamina centroid" holds, it must be a kite. In other words: In the world of tangential quadrilaterals, only kites can have the property "incenter = lamina centroid". Note that rhombi are special kites. Any proof with any means welcome!

论坛问题帖 | has_answers | Hans Humenberger | 2025-12-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019442 | MSE-5119528

A conjecture relating consecutive prime gaps using bounded powers of two

Let p_n denote the n-th prime number. For $n \ge 3$, define the consecutive prime gaps: $$ d_n = p_n - p_{n-1} $$ $$ d_{n-1} = p_{n-1} - p_{n-2} $$ Define the finite set: $$ E_n = \{ -1, 2, 3, 5, 7, d_{n-1} \} $$ I propose the following conjecture. Conjecture For every $n \ge 3$, there exist: $\varepsilon \in \{-1, 0, 1\}$, an integer $k \in \{0, 1, 2, 3, 4, 5, 6\}$, and a subset $S \subseteq E_n$ (each element used at most once), such that the following equality holds: $$ p_n + \varepsilon = p_{n-1} + \sum_{x \in S} x + 2^k $$ Equivalently, in terms of prime gaps: $$ d_n + \varepsilon = \sum_{x \in S} x + 2^k $$ Motivation This conjecture suggests that the current prime gap d_n can always be approximated (up to ± 1) by a combination of: the previous prime gap d_{n-1}, a small fixed set of primes $\{2, 3, 5, 7\}$, and a bounded power of two. I have tested this conjecture on many small values of n and found no counterexample so far. However, I do not have a proof, nor do I know whether a counterexample exists for large n. Question Is this conjecture known to be false or incompatible with known results on prime gaps? Are there known results suggesting that such a bounded additive control of consecutive prime gaps cannot hold for all n?

论坛问题帖 | has_answers | Luka Duval-Arnould | 2026-01-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019443 | MSE-5120935

Behavior of polynomial roots when removing a single term.

Now posted on MO here Let $n \in \mathbb{N}$ and let $C > 0$ be a fixed constant. Consider the class of monic polynomials of degree n with complex coefficients bounded by C: $$\mathcal{P}_n(C) = \left\{ P(z) = z^n + \sum_{j=0}^{n-1} a_j z^j \;\Bigg|\; a_j \in \mathbb{C}, \, |a_j| \le C \right\}$$ For a given polynomial $P(z) = \sum\limits_{j=0}^n a_j z^j$ (where $a_n=1$), we define a term deletion process. Let $P_k(z)$ denote the polynomial obtained by removing the k-th term of $P(z)$: $$P_k(z) = P(z) - a_k z^k \quad \text{for } k \in \{0, 1, \dots, n\}$$ Let $Z(P)$ denote the multiset of roots of a polynomial P. We are comparing the roots of the original polynomial, $Z(P)$, to the aggregate multiset of roots of all modified polynomials: $$\mathcal{Z}_{\text{mod} } = \bigcup_{k=0}^n Z(P_k)$$ Let $N_{\text{total}} = |\mathcal{Z}_{\text{mod}}|$ be the total number of roots generated by this process (counting multiplicity). For any root $\zeta \in \mathcal{Z}_{\text{mod}}$, we define its distance to the original root set as: $$\text{dist}(\zeta, Z(P)) = \min_{w \in Z(P)} |\zeta - w|$$ Proposed Property For "almost all" roots generated by term deletion, this distance becomes negligible as the degree increases. Specifically, for any fixed $\varepsilon > 0$, and for any sequence of polynomials $P^{(n)} \in \mathcal{P}_n(C)$, the proportion of modified roots that are "far" from the original roots tends to zero. Formally: $$\lim_{n \to \infty} \frac{\left| \left\{ \zeta \in \mathcal{Z}_{\text{mod}} : \text{dist}(\zeta, Z(P^{(n)})) > \varepsilon \right\} \right|}{N_{\text{total}}} = 0$$ Numerical Evidence To test the conjecture, a Python script was used with the following parameters: Inputs: Degree n and distance threshold ε. Generate a monic polynomial $P(z) = z^n + \sum_{j=0}^{n-1} a_j z^j$ is generated where each coefficient a_j is a random complex number. Output Data: The Generated Polynomial P(z) The total count of modified roots N_{total}. The count of "far" roots where $\text{dist}(\zeta, Z(P)) > \varepsilon$. A visualization plot. import numpy as np import matplotlib.pyplot as plt def format_complex(c): """Formats a complex number using 'i' instead of 'j'.""" real_part = c.real imag_part = c.imag sign = "+" if imag_part >= 0 else "-" return f"({real_part:.3f} {sign} {abs(imag_part):.3f}i)" def generate_and_plot(): try: n_input = input("Enter the degree (n): ") x0_input = input("Enter the distance threshold (ε): ") n = int(n_input) x0 = float(x0_input) except ValueError: print("Invalid input. Please enter an integer for n and a number for x0.") return coeffs = (np.random.uniform(-100, 100, n) + 1j * np.random.uniform(-100, 100, n)) coeffs_full = np.append(coeffs, 1.0 + 0j) original_roots = np.roots(coeffs_full[::-1]) print("\nGenerated Polynomial P(z):") poly_str = f"z^{n}" for i in range(n-1, -1, -1): c = coeffs_full[i] poly_str += f" + {format_complex(c)}z^{i}" print(poly_str) all_mod_roots = [] far_roots = [] for k in range(n + 1): temp_coeffs = coeffs_full.copy() temp_coeffs[k] = 0.0 + 0.0j k_roots = np.roots(temp_coeffs[::-1]) for r in k_roots: all_mod_roots.append(r) if len(original_roots) > 0: distances = np.abs(original_roots - r) min_dist = np.min(distances) if min_dist > x0: far_roots.append(r) else: far_roots.append(r) total_mod_roots_count = len(all_mod_roots) far_roots_count = len(far_roots) print("\n--- Statistics ---") print(f"Total original roots (Degree n): {n}") print(f"Total modified roots generated (Z_mod): {total_mod_roots_count}") print(f"Count of 'Far' roots (dist > {x0}): {far_roots_count}") if total_mod_roots_count > 0: print(f"Proportion of Far roots: {far_roots_count/total_mod_roots_count:.4f}") else: print("Proportion of Far roots: Undefined (No modified roots generated)") plt.figure(figsize=(10, 8)) if total_mod_roots_count > 0: mod_real = [z.real for z in all_mod_roots] mod_imag = [z.imag for z in all_mod_roots] plt.scatter(mod_real, mod_imag, c='gray', s=10, alpha=0.5, label='Modified Roots (P_k)') if far_roots_count > 0: far_real = [z.real for z in far_roots] far_imag = [z.imag for z in far_roots] plt.scatter(far_real, far_imag, c='blue', marker='x', s=40, label=f'Far Roots (> {x0})') orig_real = [z.real for z in original_roots] orig_imag = [z.imag for z in original_roots] plt.scatter(orig_real, orig_imag, c='red', s=80, edgecolors='black', label='Original Roots (P)') plt.title(f"Roots of P(z) and Term-Deleted Variations (n={n})") plt.xlabel("Re(z)") plt.ylabel("Im(z)") plt.axhline(0, color='black', linewidth=0.5, linestyle='--') plt.axvline(0, color='black', linewidth=0.5, linestyle='--') plt.legend() plt.grid(True, alpha=0.3) plt.show() if __name__ == "__main__": generate_and_plot() Generated Polynomial P(z): z^120 + (14.549 + 4.789i)z^119 + (-64.872 - 30.454i)z^118 + (-42.756 - 89.970i)z^117 + (33.601 - 46.041i)z^116 + (16.086 - 85.738i)z^115 + (-85.173 - 37.539i)z^114 + (49.884 - 46.903i)z^113 + (-12.823 - 89.226i)z^112 + (-94.270 - 50.121i)z^111 + (-83.669 - 35.626i)z^110 + (-52.091 + 75.844i)z^109 + (-61.266 - 95.643i)z^108 + (6.552 - 28.493i)z^107 + (12.750 - 62.763i)z^106 + (-55.874 + 1.230i)z^105 + (30.139 + 93.911i)z^104 + (-41.187 - 3.923i)z^103 + (-47.934 - 45.805i)z^102 + (25.322 - 66.519i)z^101 + (-31.942 + 74.200i)z^100 + (0.928 + 2.317i)z^99 + (6.079 + 66.727i)z^98 + (-6.494 - 5.094i)z^97 + (-90.498 - 46.044i)z^96 + (91.925 - 94.499i)z^95 + (6.824 - 43.377i)z^94 + (-78.000 + 42.320i)z^93 + (36.828 + 42.174i)z^92 + (-70.149 + 37.558i)z^91 + (-26.179 - 69.479i)z^90 + (-63.005 - 99.806i)z^89 + (29.172 + 36.844i)z^88 + (-71.685 - 63.670i)z^87 + (-67.593 - 77.329i)z^86 + (99.584 - 2.729i)z^85 + (55.739 - 24.792i)z^84 + (-43.818 - 89.606i)z^83 + (-1.526 + 23.966i)z^82 + (57.539 - 94.350i)z^81 + (43.114 - 2.197i)z^80 + (-66.952 + 5.514i)z^79 + (79.540 + 21.620i)z^78

+ (-50.353 - 81.357i)z^77 + (39.743 + 87.752i)z^76 + (55.435 + 9.258i)z^75 + (-81.340 + 4.640i)z^74 + (32.413 - 67.446i)z^73 + (-82.944 - 97.603i)z^72 + (56.134 - 44.344i)z^71 + (63.889 - 60.293i)z^70 + (-29.403 - 86.565i)z^69 + (19.351 - 52.593i)z^68 + (-51.060 + 43.591i)z^67 + (61.210 + 60.006i)z^66 + (41.264 - 49.092i)z^65 + (37.653 + 40.686i)z^64 + (-11.274 + 96.399i)z^63 + (88.072 + 89.275i)z^62 + (-84.852 + 64.667i)z^61 + (-75.787 + 72.384i)z^60 + (71.273 + 86.335i)z^59 + (-1.395 + 30.072i)z^58 + (-7.063 + 67.584i)z^57 + (-71.953 + 93.879i)z^56 + (77.405 - 24.546i)z^55 + (85.440 - 59.955i)z^54 + (-65.946 + 89.210i)z^53 + (-17.366 - 74.551i)z^52 + (-93.553 + 91.870i)z^51 + (-80.256 - 42.458i)z^50 + (-96.493 + 2.800i)z^49 + (41.449 - 58.326i)z^48 + (59.051 + 30.935i)z^47 + (15.497 - 36.403i)z^46 + (-7.093 - 70.560i)z^45 + (74.641 - 38.073i)z^44 + (51.600 + 85.893i)z^43 + (63.342 + 61.385i)z^42 + (49.426 - 8.851i)z^41 + (59.812 + 4.237i)z^40 + (36.973 + 18.570i)z^39 + (-61.769 + 90.691i)z^38 + (56.533 - 26.561i)z^37 + (-4.772 - 18.168i)z^36 + (-22.358 - 49.874i)z^35 + (89.394 - 85.242i)z^34 + (-77.224 - 1.868i)z^33 + (61.477 + 52.935i)z^32 + (13.016 - 32.324i)z^31 + (-67.614 + 78.254i)z^30 + (40.877 - 30.979i)z^29 + (30.987 - 20.667i)z^28 + (59.536 + 52.168i)z^27 + (42.817 + 47.837i)z^26 + (24.440 - 95.907i)z^25 + (-12.230 + 13.114i)z^24 + (-75.086 - 93.222i)z^23 + (1.178 + 56.078i)z^22 + (-36.735 + 84.915i)z^21 + (98.530 + 58.289i)z^20 + (46.365 + 90.056i)z^19 + (-75.194 + 90.001i)z^18 + (-33.897 - 34.966i)z^17 + (13.663 + 43.893i)z^16 + (-31.383 + 66.538i)z^15 + (47.984 + 60.144i)z^14 + (46.884 - 38.114i)z^13 + (21.700 - 60.699i)z^12 + (-14.108 - 51.566i)z^11 + (51.826 + 66.525i)z^10 + (3.964 + 71.961i)z^9 + (29.569 - 17.384i)z^8 + (-62.828 - 40.355i)z^7 + (27.712 - 69.163i)z^6 + (35.096 + 15.159i)z^5 + (14.742 - 22.328i)z^4 + (-75.001 - 95.746i)z^3 + (82.303 + 62.504i)z^2 + (-82.512 - 77.028i)z^1 + (-5.985 + 72.889i)z^0 $$ \begin{array}{|l|l|} \hline \text{Parameter} & \text{Value} \\ \hline \text{Degree } (n) & 120 \\ \text{Total Modified Roots } (N_{\text{total}}) & 14,519 \\ \text{Distance Threshold } (\epsilon) & 0.01 \\ \text{Count of "Far" Roots} & 538 \\ \text{Proportion of Far Roots} & 0.0371 \\ \hline \end{array} $$

论坛问题帖 | no_answers_at_retrieval | pie | 2026-01-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019444 | MSE-5121437

Is every sufficiently large integer the sum of two semiprimes with distinct prime factors?

I have been thinking about this conjecture lately: Every sufficiently large integer n can be written in the form $$ n = p_1 \cdot p_2 + p_3 \cdot p_4 $$ where p_1, p_2, p_3, p_4 are distinct prime numbers. Obviously, we need to separate the problem into dealing with even and odd numbers. Dealing with the even numbers, Chen’s theorem proves that every sufficiently large even number can be written as the sum of a prime and a semiprime. It seems logical that sufficiently large even numbers can be written as the sum of two semiprimes, as semiprimes are more dense than primes, but is there a rigorous proof out there? In the case of the odd numbers, I am not even sure which theorem could be the key to deal with this problem. Obviously, one of the prime numbers must be 2, so the conjecture takes up this form: $n_{\text{odd}}=2\cdot p_2+p_3\cdot p_4$. I think this might be related to Goldbach’s theorem. Can anyone give me some explanation if this statement can be proven from already existing theorems? Also, if this conjecture is true, there should exist an upper bound for integers that cannot be written in this form, are there any results covering this topic? I would appreciate any help or guidance, thank you in advance for your time!

论坛问题帖 | has_answers | LuckyTiger12 | 2026-01-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019445 | MSE-5124784

Is every sufficiently large integer is the sum of some integer n and the number of divisors of n

Is it true that $$ \forall N > N_{0}, \exists n : N = n + \tau(n) $$ where $\tau(n)$ is the number of divisors of n (with n included). I have been trying to solve that for several days without any success. I tried to use the fact that $\tau(n)$ is odd only for $n = a^2$. Also I tried to show that there are always two different numbers m and n such that $$ m + \tau(m) = n + \tau(n) $$ then we would (probably) show that there are infinitely many gaps. But I failed to prove that. First numbers which cannot be written in a such way are: 1, 3, 6, 8, 11

论坛问题帖 | no_answers_at_retrieval | LRDPRDX | 2026-02-16

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019446 | MSE-5125102

Bertrand's Postulate bound of $p_{n+1} \lt 2p_n$ can be lowered even further, down to $p_{n+1} \lt p_n + 2n$ - known conjecture?

Verification Code. from sympy import * def actual_count(n, x): count = 0 for a in range(prime(n) + 1, x+1): if isprime(a): count += 1 return count N = 800 for n in range(1, N+1): for x in range(1, 2*prime(n+1) + 1): A = actual_count(n, x) #F = formula(n, x) #assert A == F if A > 0: assert x <= prime(n) + 2 * n print(f"{x} <= p_{n} + 2*{n} = {prime(n) + 2 * n}") break Question. Bertrand's classical postulate implies that $p_{n+1} \lt 2p_n$ for all $n \geq 1$. Was wondering if anyone has conjectured or even proven the even sharper bound of $p_{n + 1} \lt p_n + 2n$. 3 <= p_1 + 2*1 = 4 <= 4 = 2*p_1 5 <= p_2 + 2*2 = 7 <= 6 = 2*p_2 7 <= p_3 + 2*3 = 11 <= 10 = 2*p_3 11 <= p_4 + 2*4 = 15 <= 14 = 2*p_4 13 <= p_5 + 2*5 = 21 <= 22 = 2*p_5 17 <= p_6 + 2*6 = 25 <= 26 = 2*p_6 19 <= p_7 + 2*7 = 31 <= 34 = 2*p_7 23 <= p_8 + 2*8 = 35 <= 38 = 2*p_8 29 <= p_9 + 2*9 = 41 <= 46 = 2*p_9 31 <= p_10 + 2*10 = 49 <= 58 = 2*p_10 37 <= p_11 + 2*11 = 53 <= 62 = 2*p_11 41 <= p_12 + 2*12 = 61 <= 74 = 2*p_12 43 <= p_13 + 2*13 = 67 <= 82 = 2*p_13 47 <= p_14 + 2*14 = 71 <= 86 = 2*p_14 53 <= p_15 + 2*15 = 77 <= 94 = 2*p_15 59 <= p_16 + 2*16 = 85 <= 106 = 2*p_16 61 <= p_17 + 2*17 = 93 <= 118 = 2*p_17 67 <= p_18 + 2*18 = 97 <= 122 = 2*p_18 71 <= p_19 + 2*19 = 105 <= 134 = 2*p_19 I think I can prove it using a combination of Bertrand's postulate itself and rank arguments involving a $\Bbb{Z}$-linearly independent basis of functions, which happens to be of size $2n$ functions in the basis. But that's off-topic for this site.

论坛问题帖 | has_answers | TotoposDiagramChaseApp | 2026-02-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019447 | MSE-5128694

Monotonicity of a sequence involving alternating binomial sums and logarithms: $a_p = \frac{\pi^2}{6} + p w_p - p^2 c_p^2$

Let $p \ge 1$ be an integer. Consider the following alternating binomial sums involving logarithms: $$c_p := -\frac{1}{p} \sum_{k=1}^p (-1)^k \binom{p}{k} \ln k$$ $$w_p := -\frac{1}{p} \sum_{k=1}^p (-1)^k \binom{p}{k} (\ln k)^2$$ Using these, define the sequence $(a_p)_{p \ge 1}$ as: $$a_p := \frac{\pi^2}{6} + p w_p - p^2 c_p^2$$ Can we prove that a_p is decreasing in p? I've computationally verified this for $p<100$. I've also considered using $\binom{p+1}{k} = \binom{p}{k} + \binom{p}{k-1}$, but this makes the expressions significantly more difficult to handle. These can be in the form of the Nörlund-Rice integrals, however I don't see how to move on from there (I'm no expert in complex analysis). Please, can a wizard help me? Edit: 16/3/2026 (Thanks to Daniel Goc's suggestions) Using: $$\ln(x) = \int_0^\infty \left(\frac{1}{1+t} - \frac{1}{x+t} \right) dt$$ We can pull the discrete sums inside the integral. As discovered, this beautifully collapses the binomial expansions into a single double integral over the interval $(0, \infty)^2$: $$(p+1)(a_p - a_{p+1}) = \int_0^\infty \int_0^\infty I(s,t) \, ds \, dt$$ where for $x > 0$: $$w(x) := \frac{(p+1)!}{(x+1)(x+2)\dots(x+p+1)} \in (0,1)$$ $$I(s,t) = \frac{w(s)(1-w(t))}{t} + \frac{w(t)(1-w(s))}{s} + \frac{w(t)-w(s)}{t-s} - \frac{w(s)w(t)}{p+1}$$ Let $$g(x) := \frac{1}{w(x)} = \prod_{k=1}^{p+1} \left(1 + \frac{x}{k}\right)$$ This is a polynomial of degree $p+1$, which we can write as: $$g(x) = \sum_{n=0}^{p+1} c_n x^n$$ Note that $c_0 = 1$ and $c_1 = \sum_{k=1}^{p+1} \frac{1}{k} = H_{p+1}$ which is the $(p+1)$th harmonic number. With a bit of work, we obtain: $$I(s,t) = w(s)w(t) \left[H_p - \sum_{n=3}^{p+1} c_n \sum_{k=1}^{n-2} t^{n-1-k} s^k \right]$$ Define the k-th moment of the $w(x)$ as $M_k := \int_0^\infty x^k w(x) \, dx$, integrating our factorized $I(s,t)$ over $(0, \infty)^2$, the separability of the polynomial terms allows us to distribute the integrals: $$\begin{aligned} \int_0^\infty \int_0^\infty I(s,t) \, ds \, dt &= \int_0^\infty \int_0^\infty w(s)w(t) \left[H_p - \sum_{n=3}^{p+1} c_n \sum_{k=1}^{n-2} t^{n-1-k} s^k \right] ds \, dt \\ &= H_p \left(\int_0^\infty w(s) ds\right) \left(\int_0^\infty w(t) dt\right) - \sum_{n=3}^{p+1} c_n \sum_{k=1}^{n-2} \left(\int_0^\infty w(s) s^k ds\right) \left(\int_0^\infty w(t) t^{n-1-k} dt\right) \end{aligned}$$ This gracefully collapses the tough problem into: $$(p+1)(a_p - a_{p+1}) = H_p M_0^2 - \sum_{n=3}^{p+1} c_n \sum_{k=1}^{n-2} M_k M_{n-1-k}$$ Where our goal is to prove that: $$H_p M_0^2 > \sum_{n=3}^{p+1} c_n \sum_{k=1}^{n-2} M_k M_{n-1-k}$$ I think this is not the way.

论坛问题帖 | no_answers_at_retrieval | Tveltzel | 2026-03-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019448 | MSE-5128701

Triangle inequality for distances from an interior point to three points on the edge of a rectangle

Definition: A closed curve S is said to have a completely triangular region if it has at least one interior point P such that for all triplets of points A,B,C on the edges of S the distances PA,PB,PC satisfy the triangle inequality. Example: For circle with radius R, its completely triangular region is a concentric circle with radius $R/3$. See proof by @lulu in the comments of this post. Question: Does a rectangle have a completely triangular region? My experimental data suggests that as a rectangle becomes thinner, its completely triangular region become smaller and at a critical point it vanishes. More specifically, Conjecture: If the ratio of the height to the width of a rectangle is greater than $\sqrt{3}$ then it has no completely triangular region. Can this be proved or disproved?

论坛问题帖 | has_answers | Nilotpal Kanti Sinha | 2026-03-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019449 | MSE-5129286

Is Euler's line tangent to the circle passing through the orthocenter and the two isogonic centers of a triangle?

By experimentations using GeoGebra and involving the two isogonic centers of triangle ABC, I observed that the circle passing through these two points, as well as through the orthocenter H of ABC, is tangent to the Euler line, precisely at H. Is this a known property, if it turns out to be true? If not, would a synthetic or computational proof be possible? Thank you for your answers and feedback.

论坛问题帖 | has_answers | Jamil Sanjakdar | 2026-03-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019450 | MSE-5129856

How small is $\Pr(TU=TV)$ for a random Toeplitz matrix over $\mathbb{Z}$?

How small is $\Pr(TU=TV)$ for a random Toeplitz matrix over $\mathbb{Z}$? Let h be a large integer and let $m = \alpha h \log h$ for some constant $\alpha > 0$. Let T be a random $(0,1)$-Toeplitz matrix of size $h \times m$, obtained by choosing its $h+m-1$ diagonal entries independently and uniformly from $\{0,1\}$. Let $U, V \in \{0,1\}^m$ be independent uniform random vectors, and consider the collision probability $$ \Pr(TU = TV), $$ where the product is taken over the integers (i.e., in $\mathbb{Z}^h$). Equivalently, writing $X := U - V \in \{-1,0,1\}^m$, this is $$ \Pr(TX = 0). $$ Basic bound A simple argument shows that $$ \Pr(TU = TV) \le 2^{-h} + 2^{-m}. $$ Indeed, conditioning on $X \neq 0$, one can expose the Toeplitz diagonals sequentially and see that each of the h equations forces a fresh random bit, giving probability at most 2^{-h}. Question / conjecture I am wondering whether a much stronger bound holds on average: Are there constants $\alpha, \beta > 0$ such that for $m = \alpha h \log h$, $$ \Pr_{T,U,V}(TU = TV) \le 2^{-\beta h \log h} \, ? $$ Heuristically, for a typical X, each row $(TX)_i$ is a random signed subset sum of about $m/2$ terms, so $$ \Pr((TX)_i = 0) \approx O(m^{-1/2}), $$ and if the h constraints behaved independently one would expect $$ \Pr(TX = 0) \approx m^{-h/2} = 2^{-\Theta(h \log h)}. $$ Difficulties The main issue is that the rows of a Toeplitz matrix are highly dependent (they are shifts of each other), so standard iid Littlewood--Offord arguments do not apply directly. Moreover, the bound cannot hold uniformly over all $X \in \{-1,0,1\}^m$ (e.g., very sparse or highly structured X can give much larger probabilities), so one needs to exploit the distribution of $X = U - V$. Question Is such a bound known? Are there relevant results on anti-concentration for random Toeplitz (or convolution-type) operators that could be used here?

论坛问题帖 | no_answers_at_retrieval | Simd | 2026-03-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019451 | MSE-5130226

Validity of the conjecture $ \ln (1^{a_1} 2^{a_2} \dots n^{a_n}) = C \left (n \ln (n) - n \right) +O(\ln n)$?

Heuristic proof: Consider the integral $$ \int_{0}^{b} \ln(x) dx = b \ln b -b $$ We define $$C = \lim_{s \to 1} (\sum_{r=1}^\infty \frac{a_r}{r^s} \times \frac{1}{\zeta(s)}) $$ and the $\epsilon = b/n$ and the Riemann limit but take a weighted sum (with positve numbers a_r): $$ C \int_{0}^{b} \ln(x) dx = \lim_{n \to \infty} (a_1 \ln \epsilon + a_2 \ln (2 \epsilon) + \dots + a_n \ln (n \epsilon)) \epsilon $$ Rearraging: $$ C \int_{0}^{b} \ln(x) dx = \lim_{n \to \infty} ((\ln (1^{a_1} 2^{a_2} \dots n^{a_n}) + (\frac{\sum a_r}{n} n \ln \epsilon) \epsilon$$ Now using asymptotics and cancelling some factors and terms: $$ C \left (n \ln (n) - n \right) \sim \ln (1^{a_1} 2^{a_2} \dots n^{a_n}) $$ Conjectured Claim: For $$C = \lim_{s \to 1} (\sum_{r=1}^\infty \frac{a_r}{r^s} \times \frac{1}{\zeta(s)})\neq 0$$ the following holds: $$ \ln (1^{a_1} 2^{a_2} \dots n^{a_n}) = C \left (n \ln (n) - n \right) +O(\ln n)$$ Is this correct? Can it be proven if so?

论坛问题帖 | has_answers | More Anonymous | 2026-03-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019452 | MSE-5130424

An angle in a triangle satisfying $a^3 + b^3 = c^3$

$\triangle ABC$ is a triangle. Let $|BC|= a$; $|AC|= b$; $|AB|= c$. We know that if: $a^2 + b^2 = c^2$, then angle C is equal to 90°. So, I asked myself the following question: What happens when, instead, we have $$a^3 + b^3 = c^3?$$ I conducted many experiments with GeoGebra which allowed me to conjecture that, in this case, we necessarily have $$78° < C < 90°.$$ Is this true? If so, can we prove this inequality with, perhaps, greater precision?

论坛问题帖 | has_answers | Jamil Sanjakdar | 2026-03-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019453 | MSE-5133875

Signs of polynomials in e and π

I am trying to prove this formula for all (or at least all but finitely many) n: $$ \begin{align} \tag{1} &\operatorname{sign}\!\left[\sum_{j=0}^{n+1} (-1)^j \binom{n+1}{2j} (2\pi)^{2j} \right]\\ &= \operatorname{sign}\!\left[-(e-1)^{n+1} n! + \sum_{k=0}^{n} (e-1)^{\,n-k} \sum_{j=0}^{k} (-1)^j \binom{k}{j} (k+1-j)^n \right] \end{align} $$ The LHS is a polynomial in π, let's call it $P_n(\pi)$. The RHS is a polynomial in e, say $Q_n(e)$. The full derivation is here: https://projecteuclid.org/journals/involve-a-journal-of-mathematics/volume-7/issue-1/Comparing-a-series-to-an-integral/10.2140/involve.2014.7.57.full. The polynomial in π is essentially $\cos(m \cdot \arctan(2\pi))$ scaled by a positive constant, which can be derived using Chebyshev polynomials. The polynomial in e is a negative integer order polylogarithm which can be computed explicitly using Stirling numbers of the second kind, which relates to the inclusion-exclusion principle mentioned by @ThomasAndrews. The idea for this conjecture essentially came from comparing the gamma function $\int_0^\infty t^x e^{-t}dt$ with the related sum $\sum_{k=1}^\infty k^x e^{-k}$. The difference can be written as an infinite series using the Poisson summation formula. It equals $$ \begin{align} -2 \Gamma(x+1) \cdot \sum_{k=1}^{\infty} \frac{ \cos \left((x+1)\arctan(2\pi k)\right) }{ \left(1 + 4\pi^2 k^2\right)^{(x+1)/2} } \end{align} $$ The series converges quickly, so I inserted integer values for x, truncated the sum after the first term and conjectured that this should give the correct sign (and asymptotic behaviour) of the full sum. The polynomials both change signs infinitely often (proven in the paper) and the conjecture would follow from the number $\arctan(2\pi)/\pi$ having finite irrationality measure. In the case where the irrationality measure is infinite (i.e., it's a Liouville number), the polynomial in π could come very close to zero infinitely many times which would make deciding the sign very difficult. I couldn't prove anything about the irrationality measure, however. There's one more thing I looked at, a recursion for the polynomials Q_n, defined as above (where $x=e-1$). They satisfies the recursion \begin{align} Q_{n+1}(x) &= (n+1)(x+1)\,Q_n(x) \\ &\quad - (x^2 + x)\,Q_n'(x) \\ &\quad - (n+1)!\,x^{n+2}. \end{align} This is not a linear recursion because the degree $n+1$ appears explicitely, but it still looks "rigid", in the sense that the sign sequence might be Sturmian, i.e. governed by $\cos(n\cdot c \pi)$ for fixed x and some irrational c.

论坛问题帖 | no_answers_at_retrieval | Leon | 2026-04-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019454 | MSE-5135135

Conjecture : $\forall n>2, \max_{d \mid (2^n + 3)} v_2(d+9)<3\log(n)$

Edit : $L(b,a,c)=\min(\{x>0 , a^x \equiv b \pmod c \})$ (like discrete log) Let me clarify, why discrete log shows up in this conjecture Since $d \mid 2^n+3$ we'll have $n \equiv L(-3,2,d) \pmod {\text{ord}_d(2)}$ As $n>0$, we'll have $n \ge L(-3,2,d)$ where $d=2^v\lambda-9$ for some odd integer λ So maybe using some lower bound of $L(-3,2,d)$ which involves d and hence v we'll get something like $v=O(\log(n))$, Can someone guide me toward papers involving nice estimates of L ? Let n be a positive integer , we call $\psi(n) = \max \left\{ v_2(d+9) \text{ s.t. } d \mid 2^n + 3 \right\}$ for instance : $2^8+3=7\times 37$ then $\left\{ v_2(\color{blue}d+9) \text{ s.t. } d \mid 2^8 + 3 \right\}=\{ v_2(\color{blue}1+9) , v_2(\color{blue}7+9),v_2(\color{blue}{37}+9),v_2(\color{blue}{7\times 37}+9)\}$ $=\{ 1,4,1,2 \}$, so $\psi(8)=\max\{ 1,4,1,2 \}=4$ my conjecture is : $\forall n>2 , \psi(n)<3\log(n)$ Remark: Here $\log$ denotes the natural logarithm (base e) here's a plot of $\psi(n)$ and $3\log(n)$ for $n\le300$, here's a chart of integers scattered in $\color{red}{\text{red}}$ where ψ is so big than usual, $$\psi(13)=6 < 3\log(13) \approx 7.694$$ $$\psi(22)=8 < 3\log(22) \approx 9.273$$ $$\psi(33)=10 < 3\log(33) \approx 10.489$$ $$\psi(248)=12 < 3\log(248) \approx 16.540$$ $$\psi(269)=14 < 3\log(269) \approx 16.784$$ My attempt : I found $\displaystyle \forall n >8$ we'll have $\displaystyle \psi(n)<\frac{n}{2}$, First note that , $\psi(n)≥1$ since $v_2(1+9)=1$ if $2^n+3 \in \mathbb{P}$ then $\displaystyle \forall n \ge 3 ,\psi(n)=2<\frac{n}{2}$ for $n > 4$ otherwise , take $d \mid 2^n+3$ s.t. $d<2^n+3$ and let $d=2^v\lambda-9$ where $v=v_2(d+9) \ge 1$ and λ an odd integer ≥1 since $\gcd(2^n+3,6)=1$, we'll

have $5d \le2^n+3 \implies 2^v \le 0.2\cdot2^n+9.6$ so $v \le \log_{2}\left(0.2\cdot2^{n}+9.6\right)<n$ for $n \ge 4$ thus $\forall n ≥4 , \psi(n)<n$ now note that $2^v\lambda \equiv 9 \pmod d$ and $2^n \equiv -3 \pmod d$ $\Rightarrow 2^v\lambda \equiv -3\cdot2^n \pmod d \Rightarrow 3\cdot2^{n-v}\equiv -\lambda \pmod d$ (since d is odd and $v<n$) so $2^v\lambda-9 \mid 3\cdot 2^{n-v}+\lambda$, if $2^v\lambda-9=3\cdot 2^{n-v}+\lambda$, I'll show $\displaystyle v<\frac{n}{2}$ we have : $(2^v-1)\lambda = 3(2^{n-v}+3)$ so $\lambda \mid 2^{n-v}+3$ (since $\gcd(\lambda,3)=1$) so $v_2(\lambda+9)\le\psi(n-v)<n-v$ for $n-v \ge 4$ and $2^v\lambda=\lambda+9+3\cdot2^{n-v} \Rightarrow v=v_2(\lambda+9)$ and $2^v \mid 3 \cdot 2^{n-v}$ so $2v \le n$ $\displaystyle \Rightarrow \lambda = \frac{\lambda+9}{2^v}+3\cdot2^{n-2v} \Rightarrow n-2v>0$ since λ and $\displaystyle \frac{\lambda+9}{2^v}$ are odd hence $\displaystyle v<\frac{n}{2}$, now if $n-v\le3$ we have $\lambda \mid 2^{n-v}+3$ $\displaystyle \implies \frac{2^{n-v}+3}{\lambda} \in \{ 1,5,7,11 \}$ since $2^{n-v}+3$ is prime for $n-v \le 3$ and $\displaystyle 2^v=1+\frac{3(2^{n-v}+3)}{\lambda}\le1+3\times 11=34 \Rightarrow v \le 5$ so $n \le 3+v \le 8$, but we're taking $n>8$ otherwise , $2^v\lambda-9<3\cdot 2^{n-v}+\lambda$ we will have $\displaystyle2^v\lambda-9 \le \frac{3\cdot 2^{n-v}+\lambda}{5}$, since $\gcd(3\cdot 2^{n-v}+\lambda,6)=1$ this will lead to $5X^2-46X-3\cdot 2^n \le0$ where $X=2^v$ (we got rid from λ , by using $\lambda \ge 1$) $\displaystyle \Rightarrow X \le \frac{23+\sqrt{529+15\cdot2^n}}{5} \Rightarrow v \le \log_{2}\left(\frac{23+\sqrt{529+15\cdot2^{n}}}{5}\right)<\frac{n}{2}$, for $n \ge 10$ and $\psi(9)=4<4.5 , \psi(10)=3 <5$ so we can take $n>8$ Thus $\displaystyle \forall n >8 , \psi(n) < \frac{n}{2}$ some further questions : Can we further optimize our estimate to ψ ? If so how can we do it ? How far can we generalize this to estimate valuations of shifted divisors ?

论坛问题帖 | has_answers | Lhachimi | 2026-04-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019455 | MSE-5137196

Conjecture: The probability that a line through two random points in a prism with convex bases intersects neither base is 1/2.

A line passes through two uniformly random points inside a prism with convex bases. Here is an example with triangular bases. Is the following conjecture true: The probability that the line intersects neither base is $1/2$. I have proven that if the bases are circles, the probability is $1/2$. Simulations suggest that if the bases are triangles or rectangles, the probability is also $1/2$. I am hoping for an intuitive proof based on symmetry, but any proof will be appreciated. (If an intuitive explanation cannot be found, I will add this question to the list.)

论坛问题帖 | has_answers | Dan | 2026-05-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019456 | MSE-5137610

A conjecture concerning three-way number partitioning

To start with, apologies for the length of this posting. I have a conjecture that I am interested in proving or disproving. Before I get to the conjecture, I need a fair amount of background material. The starting point for the situation is a multiset of n non-negative ordered numbers: \begin{equation} x_{1} \, \ge \, x_{2} \, \ge \, \ldots \, \ge x_{n} \, > \, 0 \; . \tag{1} \end{equation} I assume $n \ge 4$. I want to partition these numbers into 3 subsets. I will represent this partition by an ordered triplet of the indices $\{1,2, \ldots n\}$: \begin{equation} T \, = \ (P(1), P(2), P(3)) \, . \tag{2} \end{equation} As this is a partition, it has the following two properties: \begin{equation} P(1) \, \cup \, P(2) \, \cup \, P(3) \, = \, \{1, 2, \ldots \, n \} \quad \text{and} \quad P(i) \cap P(j) \, = \, \emptyset, \, i \ne j \tag{3} \end{equation} To prevent repeated partitions, I require the following montonicity property: Monontonic partition A partition will be monotonic when the smallest element in $P(i)$ is always less than the smallest element in $P(j)$, when $i < j$. There are three pairs of subsets that I am interested in: \begin{equation} \Psi \, = \, \{ (1,2), (2,3), (1,3) \} \; . \tag{4} \end{equation} Given a subset $P \subset \{1, 2, \, \ldots \, n\}$, I will use the shorthand $S(P)$ to represent the following sum: \begin{equation} S(P) \, = \, \sum_{i \in P} \, x_{i} \; . \tag{5} \end{equation} I am interested in the following three differences: \begin{equation} D(i,j) \, = \, S(P(i)) - S(P(j))\, = \, \sum_{r\in P(i)} x_{r} \, - \, \sum_{r\in P(j)} x_{r}, \; (i,j) \in \Psi . \tag{6} \end{equation} The objective in considering partitions is to minimize the largest absolute value of the three $D(i,j)$ differences: \begin{equation} J(P(1), P(2), P(3)) \, = \, \max\{ |D(i,j)| \, : \, (i,j) \in \Psi \} \; . \tag{7} \end{equation} In general, without knowing the x_{i}, I cannot know beforehand whether a difference $D(i,j)$ will be positive or negative. However, there is a class of differences, which I call assurdely non-negative (ANN), which can never be negative. ANN difference A difference $D(i,j)$ is assuredly non-negative (ANN), when $D(i,j) \ge 0$, regardless of the x_{i} values. There are conditions that are reasonably easy to compute for testing whether a $D(i,j)$ is ANN or not. (See https://math.stackexchange.com/a/5106854.) Correspondingly, I define an ANN partition. ANN partition A partition $(P(1), P(2), P(3))$ will be called an ANN partition when at least one of the differences $(D(1,2), D(2,3), D(1,3))$ is ANN. Given two partitions $(\bar{P}(1), \bar{P}(2), \bar{P}(3))$ and $(\hat{P}(1), \hat{P}(2), \hat{P}(3))$, it is possible that one dominates the other. Domination Let $(\hat{P}(1), \hat{P}(2), \hat{P}(3))$ and $(\bar{P}(1), \bar{P}(2), \bar{P}(3))$ be two partitions for which it can be proved that $J(\hat{P}(1), \hat{P}(2), \hat{P}(3))$ $\le$ $J(\bar{P}(1), \bar{P}(2), \bar{P}(3))$, regardless of the x_{i} numbers. In such a case I say that partition $(\hat{P}(1), \hat{P}(2), \hat{P}(3))$ dominates partition $(\bar{P}(1), \bar{P}(2), \bar{P}(3))$. Consider two partitions, $\bar{T} = (\bar{P}(1), \bar{P}(2), \bar{P}(3))$ and $\hat{T} = (\hat{P}(1), \hat{P}(2), \hat{P}(3))$, whose associated differences are $\bar{D}(i,j)$ and $\hat{D}(i,j)$ respectively. Partition $\bar{T}$ will be dominated by partition $\hat{T}$, when for each $\hat{D}(i,j), (i,j) \in \Psi$, the following two conditions are met: domination condition when $\hat{D}(i, j) \ge 0$: There exists $(k,\ell) \in \Psi$ such that $\bar{D}(k,\ell) - \hat{D}(i, j)$ is ANN or there exists $(p,q) \in \Psi$ such that $-\hat{D}(i, j) - \bar{D}(p,q)$ is ANN. domination condition when $\hat{D}(i, j) < 0$: There exists $(r,s) \in \Psi$ such that $\bar{D}(r,s) + \hat{D}(i, j)$ is ANN or there exists $(t,u) \in \Psi$ such that $\hat{D}(i, j) - \bar{D}(t,u)$ is ANN. For any given n, the following is an ANN partition, which I will call the base ANN partition:

\begin{equation} P(1) = \{1\}, \quad P(2)= \{2\}, \quad P(3)) = \{3,4, \, \ldots \, n \} \, . \tag{8} \end{equation} For the base partition, the difference $D(1,2) = x_{1} - x_{2}$ is ANN. I believe that I have been able to prove the following: Proposition 1 Apart from the base ANN partition (8), all other ANN partitions are dominated. What I wish to prove/disprove is the following: Conjecture 1 For a partition $(P(1), P(2), P(3))$ to be dominated, at least one of the differences $D(i,j), (i,j) \in \Psi$, must be ANN. In other words, a necessary condition for a partition to be dominated is that it is an ANN partition. I have tested computationally the conjecture and have found it to be true for $5 \le n \le 12$. Example The following table contains three monotonic partitions for $n=6$ and their corresponding differences $D(i,j)$: \begin{equation*} \begin{array}{c|ccc} \text{item} & \text{partition } T_{1} & \text{partition } T_{2} & \text{partition } T_{3} \\ \hline P(1) & \{1,6 \} & \{ 1,6 \} & \{1 \} \\ P(2) & \{2, 3 \} & \{2, 5 \} & \{2, 5 \} \\ P(3) & \{4, 5 \} & \{3, 4 \} & \{3, 4, 6 \} \\ D(1,2) & x_{1} - x_{2} - x_{3} + x_{6} \quad \quad & x_{1} - x_{2} - x_{5} + x_{6} \quad \quad & x_{1} - x_{2} - x_{5} \\ D(2,3) & x_{2} + x_{3} - x_{4} - x_{5} \quad \quad & x_{2} - x_{3} - x_{4} + x_{5} \quad \quad & x_{2} - x_{3} - x_{4} + x_{5} - x_{6}\\ D(1,3) & x_{1} - x_{4} - x_{5} + x_{6} \quad \quad & x_{1} - x_{3} - x_{4} + x_{6} \quad \quad & x_{1} - x_{3} - x_{4} - x_{6} \\ D(i,j) \text{ that are ANN} & D(2,3) & \text{--} & \text{--} \end{array} \end{equation*} Of these three, only T_{1} is an ANN partition. Let $D_{k}(i,j)$ be a difference associated with partition T_{k}. The following table contains the inequalities showing that T_{2} dominates T_{1}: \begin{equation*} \begin{array}{c|c} \text{$D_{2}(i,j)$ that determines } J(T_{2}) & \text{ANN showing } J(T_{2}) \le J(T_{1}) \\ \hline J(T_{2}) = |D_{2}(1,2)|, D_{2}(1,2) \ge 0 & D_{1}(1,3) - D_{2}(1,2) \, = \, x_{2} - x_{4} \ge 0 \\[1ex] J(T_{2}) = |D_{2}(1,2)|, D_{2}(1,2) < 0 & D_{2}(1,2) - D_{1}(1,2) \, = \, x_{3} - x_{5} \ge 0\\[1ex] J(T_{2}) = |D_{2}(2,3)|, D_{2}(2,3) \ge 0 & D_{1}(2,3) - D_{2}(2,3) \, = \, 2x_{3} - 2x_{5} \ge 0\\[1ex] J(T_{2}) = |D_{2}(2,3)|, D_{2}(2,3) < 0 & D_{2}(2,3) + D_{1}(2,3) \, = \, 2x_{2} - 2x_{4} \ge 0 \\[1ex] J(T_{2}) = |D_{2}(1,3)|, D_{2}(1,3) \ge 0 & D_{1}(1,3) - D_{2}(1,3) \, = \, x_{3} - x_{5} \ge 0\\[1ex] J(T_{2}) = |D_{2}(1,3)|, D_{2}(1,3) < 0 & D_{2}(1,3) - D_{1}(1,2) \, = \, x_{2} - x_{4} \ge 0 \end{array} \end{equation*} To show that T_{2} does not dominate T_{3}, I can show that when $J(T_{2}) = |D_{2}(1,2)|$ and $D_{2}(1,2) \ge 0$, none of the six possible expressions given in the conditions for domination are ANN. \begin{equation*} \begin{array}{c|cc} D_{3}(i,j) & D_{3}(i,j) - D_{2}(1,2) & -D_{3}(i,j) - D_{2}(1,2) \\ \hline D_{3}(1,2) & - x_{6} & - 2x_{1} + 2x_{2} + 2x_{5} - x_{6} \\[1ex] D_{3}(2,3) & -x_{1} + 2x_{2} - x_{3} - x_{4} + 2x_{5} - 2x_{6} & -x_{1} + x_{3} + x_{4} \\[1ex] D_{3}(1,3) & x_{2} - x_{3} - x_{4} + x_{5} - 2x_{6} & -2x_{1} + x_{2} + x_{3} + x_{4} + x_{5} \end{array} \end{equation*} Using a similar argument one can show that T_{3} does not dominate T_{2}. For example, it is possible to show that when $J(T_{3}) = |D_{3}(1,2)|$ and $D_{2}(1,2) < 0$, none of the six possible expressions given in the conditions for domination are ANN. EDIT I supply here some progress that I have made. I will start here with one particular case that is indicative of my difficulties. If you are interested in how I arrived at this case, then read until the end. I start with partitions $\hat{T} = (\hat{P}(1), \hat{P}(2), \hat{P}(3))$ and $\bar{T} = (\bar{P}(1), \bar{P}(2), \bar{P}(3))$ and their associated differences $\hat{D}(i,j)$ and $\bar{D}(i,j)$. Given that both $\hat{T}$ and $\bar{T}$ are non-ANN partitions, I want to prove that it is not possible to satisfy the following six conditions: \begin{equation*} \begin{split} \gamma_{84} \, = \, \{ \bar{D}(1,2) \ge_{ANN} \hat{D}(1,2), \quad \bar{D}(1,2) \ge_{ANN} \hat{D}(2,3), \quad \bar{D}(1,2) \ge_{ANN} \hat{D}(1,3), \\ \hat{D}(1,2) \ge_{ANN} \bar{D}(2,3), \quad \hat{D}(2,3) \ge_{ANN} \bar{D}(2,3), \quad \hat{D}(1,3) \ge_{ANN} \bar{D}(2,3) \} \end{split} \end{equation*} (The notation $A \ge_{ANN} B$ means $A-B$ is ANN.) Now for the details. Suppose we are considering the conditions for which $\hat{T}$ dominates $\bar{T}$. Since we cannot assume that any difference is ANN, to fulfill the domination conditions given in the original posting, we must consider the following 6 cases: \begin{equation*} \begin{array}{ccc} \text{(a)} \, \hat{D}(1,2) \ge 0 \quad \quad & \text{(b)} \, \hat{D}(1,2) < 0 \quad \quad & \text{(c)} \, \hat{D}(2,3) \ge 0 \quad \quad \\ \text{(d)} \, \hat{D}(2,3) < 0 \quad \quad & \text{(e)} \, \hat{D}(1,3) \ge 0 \quad \quad & \text{(f)} \, \hat{D}(1,3) < 0 \quad \quad \end{array} \end{equation*} The six possible conditions that ensure that $\hat{D}(1,2)$ will dominate some difference from partition $\bar{T}$ in case (a) are the following: \begin{equation*} \begin{split} \{\bar{D}(1,2) \ge_{ANN} \hat{D}(1,2), \quad \bar{D}(2,3) \ge_{ANN}\hat{D}(1,2), \quad \bar{D}(1,3) \ge_{ANN} \hat{D}(1,2), \\ -\bar{D}(1,2) \ge_{ANN} \hat{D}(1,2), \quad -\bar{D}(2,3) \ge_{ANN}\hat{D}(1,2), \quad -\bar{D}(1,3) \ge_{ANN} \hat{D}(1,2) \} \end{split} \end{equation*} (The notation $A \ge_{ANN} B$ means $A-B$ is ANN.) Of these 6, the following can never be satisfied: \begin{equation*} \begin{split} \bar{D}(2,3) \ge_{ANN}\hat{D}(1,2), \quad -\bar{D}(1,2) \ge_{ANN} \hat{D}(1,2), \\ -\bar{D}(2,3) \ge_{ANN}\hat{D}(1,2), \quad -\bar{D}(1,3) \ge_{ANN} \hat{D}(1,2) \end{split} \end{equation*} Consider condition $\bar{D}(2,3) \ge_{ANN}\hat{D}(1,2)$. In the difference $\bar{D}(2,3) - \hat{D}(1,2)$, the first nonzero will be $-x_{1}$, owing to the monotonicity of the partitions. As such, $\bar{D}(2,3) -\hat{D}(1,2)$ can never be ANN. Similar arguments apply for the other impossible conditions. After removing the 4 conditions that cannot be satisfied, we are left with two valid conditions for case (a): \begin{equation*} \Phi(\text{a}) \, = \, \{\bar{D}(1,2) \ge_{ANN} \hat{D}(1,2), \quad \bar{D}(1,3) \ge_{ANN} \hat{D}(1,2) \} \end{equation*} The corresponding condition sets for cases (b), (c), (d), (e) and (f) are as follows: \begin{equation*} \Phi(\text{b}) \, = \, \{\bar{D}(1,2) \ge_{ANN} \hat{D}(2,3), \quad \bar{D}(2,3) \ge_{ANN} \hat{D}(2,3), \quad \bar{D}(1,3) \ge_{ANN} \hat{D}(2,3) \} \end{equation*} \begin{equation*} \Phi(\text{c}) \, = \, \{\bar{D}(1,2) \ge_{ANN} \hat{D}(1,3), \quad \bar{D}(1,3) \ge_{ANN} \hat{D}(1,3) \} \end{equation*} \begin{equation*} \begin{split} \Phi(\text{d}) \, = \, \{\bar{D}(1,2) \ge_{ANN} -\hat{D}(1,2), \quad \bar{D}(2,3) \ge_{ANN}-\hat{D}(1,2), \quad \bar{D}(1,3) \ge_{ANN} -\hat{D}(1,2), \\ \hat{D}(1,2) \ge_{ANN} \bar{D}(1,2), \quad \hat{D}(1,2) \ge_{ANN} \bar{D}(2,3), \quad \hat{D}(1,2) \ge_{ANN} \bar{D}(1,3) \} \end{split} \end{equation*} \begin{equation*} \begin{split} \Phi(\text{e}) \, = \, \{\bar{D}(1,2) \ge_{ANN} -\hat{D}(2,3), \quad \bar{D}(2,3) \ge_{ANN}-\hat{D}(2,3), \\ \bar{D}(1,3) \ge_{ANN} -\hat{D}(2,3), \quad \hat{D}(2,3) \ge_{ANN} \bar{D}(2,3) \; \} \end{split} \end{equation*} \begin{equation*} \begin{split} \Phi(\text{f}) \, = \, \{\bar{D}(1,2) \ge_{ANN} -\hat{D}(1,3), \quad \bar{D}(2,3) \ge_{ANN}-\hat{D}(1,3), \quad \bar{D}(1,3) \ge_{ANN} -\hat{D}(1,3), \\ \hat{D}(1,3) \ge_{ANN} \bar{D}(1,2), \quad \hat{D}(1,3) \ge_{ANN} \bar{D}(2,3), \quad \hat{D}(1,3) \ge_{ANN} \bar{D}(1,3) \} \end{split} \end{equation*} To obtain all possible combinations of conditions ensuring that $\hat{T}$ dominates $\bar{T}$, we simply form the Cartesian product of these condition sets: \begin{equation*} \Gamma \, = \, \Phi(\text{a}) \, \times \, \Phi(\text{b})\, \times \, \Phi(\text{c}) \, \times \, \Phi(\text{d}) \, \times \, \Phi(\text{e}) \, \times \, \Phi(\text{f}) \; . \end{equation*} In total Γ contains $2 \cdot 3 \cdot 2 \cdot 6 \cdot 4 \cdot 6 = 1728$ tuples. One such tuple is the

following: \begin{equation*} \begin{split} \gamma_{1} \, = \, \{ \bar{D}(1,2) \ge_{ANN} \hat{D}(1,2), \quad \bar{D}(1,2) \ge_{ANN} \hat{D}(2,3), \quad \bar{D}(1,2) \ge_{ANN} \hat{D}(1,3), \\ \bar{D}(1,2) \ge_{ANN} -\hat{D}(1,2), \quad \bar{D}(1,2) \ge_{ANN} -\hat{D}(2,3), \quad \bar{D}(1,2) \ge_{ANN} -\hat{D}(1,3) \} \end{split} \end{equation*} We can express the first and the fourth of these as two ANN expressions: \begin{equation*} \bar{D}(1,2) - \hat{D}(1,2) \; \text{is ANN and } \, \bar{D}(1,2) + \hat{D}(1,2) \; \text{is ANN} \; . \end{equation*} The sum of any two ANN expressions is also an ANN expression. Hence, from the sum of these two we can conclude that $2\bar{D}(1,2)$ (or simply $\bar{D}(1,2)$) is an ANN expression. However, this contradicts the assumption that all $\bar{D}(i,j)$ are non-ANN. Hence, the conditions in γ_{1} can never result in partition $\hat{T}$ dominating $\bar{T}$. The contradiction obtained for tuple γ_{1} is only one of many possible contradictions. I have written messy code to test all 1728 condition tuples for such types of contradictions. Of the 1728 tuples, 1717 yield contradictions that I can detect. However, I cannot produce contradictions for the remaining 11 condition tuples. One of these remaining 11 tuples is what I provided above: \begin{equation*} \begin{split} \gamma_{84} \, = \, \{ \bar{D}(1,2) \ge_{ANN} \hat{D}(1,2), \quad \bar{D}(1,2) \ge_{ANN} \hat{D}(2,3), \quad \bar{D}(1,2) \ge_{ANN} \hat{D}(1,3), \\ \hat{D}(1,2) \ge_{ANN} \bar{D}(2,3), \quad \hat{D}(2,3) \ge_{ANN} \bar{D}(2,3), \quad \hat{D}(1,3) \ge_{ANN} \bar{D}(2,3) \} \end{split} \end{equation*} Note that from the conditions in γ_{84}, it is straightforward to obtain $\bar{D}(1,2) \ge_{ANN} \bar{D}(2,3)$. Several non-ANN partitions exist satisfying this. Here is one example: \begin{equation*} \bar{T}_{84} \, = \, (\bar{P}(1), \bar{P}(2), \bar{P}(3)) \, = \, (\{1, 2 \}, \; \{3, 6, 9 \}, \; \{4, 5, 7, 8 \} \,) \end{equation*} However, I repeat that partition $\bar{T}_{84}$ does not dominate any other non-ANN partition.

论坛问题帖 | no_answers_at_retrieval | F Cameron | 2026-05-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019457 | MSE-5138461

Conjecture: Is $n = 9$ the only positive integer for which both $2^n + n^2$ and $2^n - n^2$ are prime?

Conjecture. The only positive integer n such that both $2^n + n^2$ and $2^n - n^2$ are prime is $n = 9$. For $n = 9$: $$2^9 + 9^2 = 512 + 81 = \mathbf{593} \quad \text{(prime)}$$ $$2^9 - 9^2 = 512 - 81 = \mathbf{431} \quad \text{(prime)}$$ Computational Evidence I have verified computationally that $n = 9$ is the only solution with $n \leq 2000$. The table below shows selected candidates $n \equiv 3 \pmod{6}$ (see Corollary below for why only these need be checked): $$\begin{array}{|c|c|c|c|} \hline n & 2^n + n^2 & 2^n - n^2 & \text{Both prime?} \\ \hline 3 & 17 & -1 & - \\ 9 & 593 & 431 & \textbf{Yes} \\ 15 & 32993 & 32543 & \text{No } (32543 = 7 \times 4649) \\ 21 & 2097593 & 2096711 & \text{No } (2096711 = 641 \times 3271) \\ \hline \end{array}$$ Modular Restrictions Claim 1. If n is even and $n \geq 2$, then $2^n + n^2$ is composite. Proof. For even $n = 2k$ with $k \geq 1$, we have $2^n = 4^k$ and $n^2 = 4k^2$, so: $$2^n + n^2 = 4\!\left(4^{k-1} + k^2\right)$$ which is divisible by 4 and greater than 4. $\blacksquare$ Claim 2. If n is odd and $3 \nmid n$, then $3 \mid 2^n + n^2$. Proof. For odd n, we have $2^n \equiv 2 \pmod{3}$. If $3 \nmid n$, then $n^2 \equiv 1 \pmod{3}$. Thus: $$2^n + n^2 \equiv 2 + 1 \equiv 0 \pmod{3}$$ The only case where $2^n + n^2 = 3$ would require $n = 1$, but $2^1 - 1^2 = 1$ is not prime. $\blacksquare$ Corollary. Any solution must satisfy $n \equiv 3 \pmod{6}$. This reduces the search space by a factor of 6. Heuristic Argument For candidates $n \equiv 3 \pmod{6}$, both $2^n + n^2$ and $2^n - n^2$ are approximately 2^n. By the Prime Number Theorem, a number near 2^n is prime with probability $\sim \dfrac{1}{n \ln 2}$. Assuming independence, the probability that both are simultaneously prime is roughly $\dfrac{1}{(n \ln 2)^2}$. Summing over all remaining candidates: $$\sum_{\substack{n \,\equiv\, 3 \pmod{6} \\ n \,>\, 9}} \frac{1}{(n \ln 2)^2} \approx 0.054$$ Since this series converges and the expected number of solutions beyond $n = 9$ is well below 0.06, heuristically we expect at most finitely many solutions - and quite possibly none. Questions Can we prove that $n = 9$ is the only solution? Is there a theoretical obstruction - such as a covering congruence system - that rules out all $n \equiv 3 \pmod{6}$ beyond $n = 9$? Is there a larger solution? The heuristic suggests any additional solution would be a rare exception. Does this problem appear in the literature? It seems natural but I have not found it in standard references on exponential Diophantine equations involving primes. Pointers to related work would be welcome.

论坛问题帖 | has_answers | Rayhan Ahmed | 2026-05-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019458 | MSE-5139403

Finding the solution for a problem on game between 2 players on colouring a Polygon from SMMC 2017 Session A

The Problem Let us take a regular n-sided polygon, $n \geq 3$, drawn on a piece of paper with all its vertices connected. Two people play a game in which they take turns to colour one of these line segments. Here we consider only the lines between any vertices of this n-sided polygon to be a line segment, i.e., the player can only colour a full diagonal or a full side and not just a segment formed due to the intersection of the diagonals in the polygon. The first player colours line segments blue, while the second player colours line segments red. A player cannot colour a line segment that has already been coloured. A player wins if they are the first to create a triangle in their own colour, whose three vertices are also vertices of the regular pentagon. The game is declared a draw if all ten line segments have been coloured without a player winning. Determine whether the first player, the second player, or neither player can force a win. Comment The original problem only talked about a pentagon, but an interesting variation was put up in the comments of the original's solution, and I found myself with 3 conjectures that are potentially the solution that I cannot think of an approach to prove or, find a way or an example to use to disprove them. Definition(not part of the problem) Before the conjecture, I want to put up 2 definitions that I can reuse in the conjecture: If a player, in his consecutive turns colours 2 line segments that do not share a vertex, that may or may not be intersecting, then let such a sequence of moves be called the 'Duo move'. If a player, in his consecutive turns colours 3 line segments that share 1 single vertex, then let such a sequence of moves be called the 'Trio move'. The Conjectures So here are the conjectures: For all $n \geq 5$, n-sided polygons, the first player can always force a win. Any player who can make the duo move or a trio move first can force a win. For all $n \leq 4$, n-sided polygons, neither player can force a win. My Reasoning for the Cojectures Algorithm for forced wins: For the Duo Move: When we make a duo move, we, as the first player, will have 2 common lines that connect the lines we have already covered as options

for our third turn. So even if the opponent colours one of our options, we will have the third line. And when we colour one of the uncoloured lines, we will have 2 triangles that can be completed with one more turn on the next move, thus forcing a win. However this is only in the case of $n \geq 5$. When we make a trio move, however, it is quite obvious with the following images, as it has been taken from the SMMC solutions page for the pentagon. As you can see, when a trio move happens, we will have 2 triangles forming for us on the next turn, even if one of the three possible triangles due to the 3 covered lines is covered by our opponent. More Comments This is a problem in the Question Paper of the Simon Marais Mathematics Competition 2017, and I came up with these conjectures while solving the problem, and I need help finding a proper proof, or maybe a proper disproof of them.

论坛问题帖 | no_answers_at_retrieval | Parthiv Pedapati | 2026-06-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019459 | MSE-5140387

$\displaystyle \sum_{k=1}^n\mu(k)x^k+1$ is irreducible over $\mathbb{Q}$ iff $M(n)\not=-1$?

Let $n>0$, $\displaystyle M(n)=\sum_{k=1}^n \mu(k)$ where $\mu(k)$ is the Möbius function I was looking for values of n where $P_n(x)=\displaystyle \sum_{k=1}^n\mu(k)x^k+1$ is reducible over $\mathbb{Q}$, using PARI-GP for(n=1,100,l=polisirreducible(sum(k=1,n,moebius(k)*x^k)+1);if(l==0,print(n))) I found $3, 4, 6, 10, 15, 16, 22, 26, 27, 28, 35, \cdots$ which matches exactly the solutions to the equation $M(n) = -1$ Clearly If $M(n)=-1$, $\displaystyle P_n(1)=\sum_{k=1}^n\mu(k)+1=-1+1=0$ so P_n is reducible over $\mathbb{Q}$ but can you show the other way ? In general, how to deal with such similar polynomials like $\displaystyle\sum_{k=0}^nf(k)x^k$ where $f(k)$ is an arithmetic function like $\mu(k),\varphi(k),\chi(k),\cdots$? Thanks for help !

论坛问题帖 | has_answers | Lhachimi | 2026-06-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019460 | MSE-5140954

Proving exact formula of a coefficient

Let $$ P_n(X)=\prod_{\substack{\mu_1,\dots,\mu_n\in\{\pm1\}\\ \mu_1\cdots \mu_n=1}} \left(X-\sum_{k=1}^n \mu_k e^{k\pi i/n} \right) $$ be a polynomial of degree $d=2^{n-1}$. Let c_n'' be the coefficient in degree $d-2n$ of P_n. Conjecture. For every $n\ge 4$, $$c_n'' = (-1)^{n+1} 2^{2n-3} \left((n-1)!\right)^2 - \frac{2^{3n-3}(2^{2n}-1)B_{2n}}{n}$$where B_{2n} is the Bernoulli number with $B_2=1/6$. This answer proved $\nu_2(c''_n)\ge 6$, leaving the proof of $c_n'' \neq 0$ open, but validating this conjectured formula would help to show that $c_n'' \neq 0$.

论坛问题帖 | has_answers | user582761 | 2026-06-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019461 | MSE-5141236

About proof by contradiction of Morley's theorem in Euclidean geometry

I try to find a proof using ad absurdum of Morley's Miracle but find any. I tried also the conjecture : Let ABC be a triangle with angles 3α, 3β, 3γ, and let D, E, F be the intersections of its adjacent trisectors. Let ℓ be any line in the plane. Denote by (D^*, E^*, F^*) the image of (D, E, F) under Steiner symmetrization of DEF with respect to ℓ. Then the lines AD^*, BD^*, CE^*, AE^*, BF^*, CF^* are trisectors of the angles of ABC if and only if ℓ is an axis of symmetry of DEF. But cannot give a response to it . Some remarks : The map $z \mapsto z^3$ on the unit circle, viewed as a three-sheeted covering, has as its ramification curve (in the real plane) a limaçon of Pascal. The Steiner inellipse is to $z \mapsto z^2$ what Morley's triangle is to $z \mapsto z^3$: in both cases, the limaçon of Pascal is the transition curve that encodes the covering. Currently there a path to the Last Fermat theorem. $x^{2/3}+y^{2/3}=1$ The conjecture uses implicitly Morley's theorem. When a line (d) passes through the orthocenter (H) of a triangle, the orthopole (C) of (d), reflected with respect to (d), belongs to the nine-point circle. In this case, the lines joining (H) to the reflections of the vertices with respect to (d) meet the corresponding sides at three points which lie on a line (l) passing through (K). The point (K) is the projection of the circumcenter onto (d). The perpendicular drawn from (H) to (d) meets the circumcircle at two points; one of these points is the focus of the inscribed parabola tangent to (d). If, through the other point, one draws a line parallel to (d), then the reciprocal transversal of the resulting line passes through the point (A). Last page. Questions : Is it possible to prove Morley's Miracle using ad absurdum ? Have you a proof of the conjecture without Morley's theorem ? Related question : What is the link between the Steiner inellipse of ABC and the Morley triangle of ABC ? Reference : R. GOORMAGHTIGH Sur l’orthopole et certains limaçons de Pascal associés au triangle Nouvelles annales de mathématiques 4e série, tome 18(1918), p. 242-249 http://www.numdam.org/item?id=NAM_1918_4_18__242_0 Bogosel, B. (2017). A Geometric Proof of the Siebeck-Marden Theorem. The American Mathematical Monthly, 124(5), 459-463. https://doi.org/10.4169/amer.math.monthly.124.5.459 https://worldbuilding.stackexchange.com/users/145770/barackouda

论坛问题帖 | no_answers_at_retrieval | Barackouda | 2026-06-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019462 | MSE-5142361

Do there exist finitely many primes p such that there exists $k \in [1,p]$ with $\mathrm{ord}_p(k) = q$ such that $k^{p-1} \equiv 1 \pmod{p^2}$?

This is extended computational evidence for the Conjecture from this question, focusing on the $n=1$ case. Conjecture A for $n=1$ states: for all sufficiently large primes p with $q \mid p-1$, $$v_p\!\left(\prod_{k=1}^{p} \Phi_q(k)\right) = q-1.$$ Reduction to $v_p(k^{p-1}-1)$. By the Proposition in the linked question, the product receives p-adic contribution only from terms where $\mathrm{ord}_p(k) = q$, and for such k, $v_p(\Phi_q(k)) = v_p(k^{p-1}-1)$. There are exactly $q-1$ residue classes of order q mod p, each appearing exactly once in $[1,p]$, so $$v_p\!\left(\prod_{k=1}^{p}\Phi_q(k)\right) = \sum_{\substack{k=1 \\ \mathrm{ord}_p(k)=q}}^{p} v_p(k^{p-1}-1).$$ Now each term

is at least 1 by Fermat, giving the lower bound at least $q-1$. The conjecture says that for all sufficiently large p, every term is exactly 1 - which is to say no $k \in [1,p]$ with $\mathrm{ord}_p(k) = q$ satisfies $k^{p-1} \equiv 1 \pmod{p^2}$. We therefore check the following equivalent conjecture: For a fixed prime q and sufficiently large primes p, there exists no $k \in [1,p]$ such that $\mathrm{ord}_p(k) = q$ and $k^{p-1} \equiv 1 \pmod{p^2}$. Computational results. We searched all primes $p \equiv 1 \pmod{q}$ up to 10^{12} (for $q=11$) and 10^{11} (for $q=3$): $q=3$: no violations found. $q=11$: two violations, $p = 400263469$ with witness $k = 160169173$, and $p = 9544876463$ with witness $k = 3519317465$. No violations for $p > 9544876463$ up to 10^{12}, consistent with these being the finitely many exceptions below the "sufficiently large" threshold. Heuristic. The probability that a given k of order q satisfies $k^{p-1}\equiv 1\pmod{p^2}$ is heuristically $1/p$. Summing over all $q-1$ such k and all primes $p\leq X$, $$\mathbb{E}[\#\text{ violations up to }X] \;\approx\; \ln\ln X.$$ At $X=10^{12}$: predicted ≈ 3.3, observed 2 for $q=11$. This is the direct analogue of classical Wieferich primes, where only $p=1093$ and $p=3511$ would be known below 6.7×10^{15}. Sharpness. Allowing $k\leq p^{3/2}$ instead of $k\leq p$, violations become dense: a search over $p\in[10^6,10^8]$ finds 1271 violations for $q=3$ and 1100 for $q=11$. The $O(p)$ threshold is sharp. Question. For $q=11$, would $p = 400263469$ and $p = 9544876463$ be the only primes for which some $k \in [1,p]$ with $\mathrm{ord}_p(k) = 11$ satisfies $k^{p-1} \equiv 1 \pmod{p^2}$? More generally, for any fixed prime q, is the set of such exceptional primes finite? Remark. Our conjecture is logically independent of the classical Wieferich problem. For a fixed base $a > 1$, it is an open question whether there are infinitely many primes p with $a^{p-1} \equiv 1 \pmod{p^2}$; the heuristic predicts infinitely many but only finitely many are known for any base. Our conjecture makes no claim about any fixed base. Instead, for each prime p, we ask about the $q-1$ values of $[1,p]$ of order q mod p - a set of bases that changes entirely with p. Whether infinitely or finitely many primes p are Wieferich for some fixed k has no bearing on whether, for large p, the order-q elements of $[1,p]$ happen to include a Wieferich witness. The quantifier structures are different: the Wieferich problem fixes k and varies p; our conjecture fixes p and asks about all $k \in [1,p]$ of a given order simultaneously. Remark (sharpness of the $[1,p]$ threshold). The bound $k \leq p$ in the Conjecture is not arbitrary. The following computational data for $q=11$ shows a sharp transition precisely at $k = p$: For $k \in [1, p]$: searching all $p \equiv 1 \pmod{11}$ up to 10^{12}, only two violations appear, both at $p < 10^{10}$. No violations for $p > 9544876463$. For $k \in [1, p^{11/10}]$: four violations appear in $p \in [10^8, 10^{10}]$, including new ones at $p = 1788375733$ and $p = 8960980403$ that do not violate the Conjecture since $k > p$. For $k \in [1, p^{3/2}]$: 1100 violations in $p \in [10^6, 10^8]$. So violations are essentially absent inside $[1,p]$ for large p, appear almost immediately once k is allowed to exceed p, and become dense at $k \leq p^{3/2}$. The $O(p)$ threshold captures something structurally real, not a coincidence of the search range.

论坛问题帖 | no_answers_at_retrieval | John C | 2026-07-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019463 | MSE-5143502

For which composite s does $p^k - s$ hit a prime for small prime p and integer $k\ge 1$?

Let $s \ge 4$ be a composite integer with $s \ne 0 \pmod3$. Computationally, for every such $s \le 2000$, I can find a prime p and integer $k \ge 1$ such that $p^k - s$ is prime (usually with $p \in \{2,3\}$ and small k). Heuristically this seems unsurprising: for fixed small p, the values $p^k - s$ for $k = 1,\dots,20$ give many candidates, each prime with probability $\sim\frac{1}{\log(p^k - s)}$ by PNT, so the expected number of hits is large. My question: Is this a known/named result, or a straightforward consequence of an existing theorem (e.g. related to Bateman-Horn-type heuristics for exponential sequences, or sieve methods)? I haven't found a reference for primality in sequences of the form $({p^k - s})_k$ for fixed p, only for polynomial families. Any pointers to relevant literature would be appreciated. What I've tried: exhaustive verification for $s \le 2000$, and a robustness check (no counterexamples) using digit sums of primes up to $1.5×10^{11}$, though this doesn't extend the range of s tested since digit sums grow only like $O(\log P)$.

论坛问题帖 | has_answers | Zeyad Muhammad | 2026-07-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019464 | MSE-5143935

Does every odd prime determine a prime in an interval of length $\sqrt{p-2}$?

Let $p \geq 3$ be an odd prime. I would like to know whether the following conjecture is true. Conjecture For every odd prime $p \geq 3$, there exist integers a and b such that: $a+b+3$ is prime; $4a+2b+3=p$ $\gcd(a,b,3)=1$ $b^2\leq 4a$ $a\geq 1$. Here, $\mathbb{P}$ denotes the set of prime numbers. For example: all $$(a,b,p) \in \{(4,-4,3),(1,-1,5),(2,-2,7),(2,0,11)\}$$ work. In each case, all five conditions are satisfied. Equivalent formulation as a prime in a short interval Set $$ q:=a+b+3. $$ Then q is prime. Solving the system $$ q=a+b+3 $$ and $$ p=4a+2b+3 $$ for a and b, we obtain $$ a=\frac{p-2q+3}{2} $$ and $$ b=\frac{4q-p-9}{2}. $$ Since p and q are odd primes, both a and b are integers. The inequality $$ b^2\leq 4a $$ is equivalent to $$ \left(\frac{4q-p-9}{2}\right)^2 \leq 4\left(\frac{p-2q+3}{2}\right). $$ Multiplying by 4 gives $$ \left(4q-p-9\right)^2 \leq 8\left(p-2q+3\right). $$ After simplifying, this becomes $$ \left(4q-p-7\right)^2 \leq 4\left(p-2\right). $$ Therefore, $$ -2\sqrt{p-2} \leq 4q-p-7 \leq 2\sqrt{p-2}. $$ Equivalently, $$ \frac{p+7-2\sqrt{p-2}}{4} \leq q \leq \frac{p+7+2\sqrt{p-2}}{4}. $$ Define $$ L(p):= \frac{p+7-2\sqrt{p-2}}{4} $$ and $$ U(p):= \frac{p+7+2\sqrt{p-2}}{4}. $$ The length of this interval is $$ U(p)-L(p)=\sqrt{p-2}, $$ and its midpoint is $$ \frac{p+7}{4}. $$ Thus, the conjecture is essentially equivalent to the following statement: For every odd prime $p \geq 3$, the interval $$ \left[\frac{p+7-2\sqrt{p-2}}{4}, \frac{p+7+2\sqrt{p-2}}{4} \right] $$ contains a prime. For sufficiently large p, the condition $a\geq 1$ is automatic. Indeed, $$ a=\frac{p-2q+3}{2}\geq 1 $$ is equivalent to $$ q\leq \frac{p+1}{2}. $$ For $p\geq 11$, one has $$ U(p)\leq \frac{p+1}{2}, $$ so every prime $q\in[L(p),U(p)]$ automatically gives $a\geq 1$. The cases $p=3,5,7$ can be checked directly. The condition $$ \gcd(a,b,3)=1 $$ is also automatic. If $3\mid a$ and $3\mid b$, then $$ p=4a+2b+3 $$ and $$ q=a+b+3 $$ would both be divisible by 3. Since both are prime, this would force $$ p=q=3. $$ But then

$$ a+b=0 $$ and $$ 4a+2b=0. $$ Subtracting twice the first equation from the second gives $$ 2a=0, $$ so $a=0$, contradicting $a\geq 1$. Heuristic The interval is centered near $p/4$ and has length approximately $\sqrt{p}$. The prime number theorem therefore suggests that the expected number of primes in the interval is approximately $$ \frac{\sqrt{p}}{\log p}, $$ which tends to infinity as $p\to\infty$. Cramér's conjecture would imply that every sufficiently large interval of this form contains a prime because $$ \left(\log x\right)^2=o\left(\sqrt{x}\right). $$ However, known unconditional bounds for primes in short intervals do not appear to guarantee a prime in every interval of length comparable to $\sqrt{x}$. Questions Is this conjecture already known? Can one prove unconditionally that the interval $$ [L(p),U(p)] $$ contains a prime for every odd prime p? Does the fact that p itself is prime make the problem easier than the general problem of finding primes in every interval of length comparable to $\sqrt{x}$? Does this conjecture follow from a standard conjecture weaker than Cramér's conjecture? Important: What it needs to be proven is that for all primes $p \geq 3$ the interval $[L(p), U(p)]$ contains at least one prime. However, I proved that there are infinitely many primes $p \geq 3$ such that the interval contains at least one prime in $[L(p), U(p)]$.

论坛问题帖 | no_answers_at_retrieval | Yoyos Tutoring | 2026-07-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019465 | MSE-5144599

Possible counterexamples to Kaplansky's zero-divisors conjecture

Kaplansky's conjecture about zero divisors is formulated as follows: If G is a torsion-free group and K is a field, then the group ring $K[G]$ has no zero divisors. Some time ago, I came across the following article: В. Г. Бардаков, М. С. Петухова, “О потенциальных контрпримерах к проблеме делителей нуля”, Вестн. НГУ. Сер. матем., мех., информ., 15:3 (2015), 30-50; J. Math. Sci., 221:6 (2017), 778-797. The article talks about groups that Rips built in the hope that among them there would be a group without torsion, which would serve as a counterexample to the zero-divisor conjecture. I'll describe the construction itself here. For more detailed information, see the article. The group $G = G(\sigma, \tau, \lambda, \mu)$ is generated by $4n$, $n \geq 2$, elements $$ a_1, a_2, \dots, a_n, b_1, b_2, \dots, b_n, c_1, c_2, \dots, c_n, d_1, d_2, \dots, d_n $$ and is defined by the relations $$ a_i c_j = b_{\sigma(i,j)} d_{\tau(i,j)}, $$ $$ a_i d_j = b_{\lambda(i,j)} c_{\mu(i,j)} $$ for all (i, j) and some functions $$ \sigma, \tau, \lambda, \mu : \{1, 2, \dots, n\}^2 \to \{1, 2, \dots, n\} $$ such that their pairs $$ (\sigma, \tau), (\lambda, \mu) : \{1, 2, \dots, n\}^4 \to \{1, 2, \dots, n\}^2 $$ define bijections of the set $\{1, 2, \dots, n\}^2$, i.e., for any pair of indices (i', j') there exists a unique pair of indices (i, j) such that $i' = \sigma(i, j)$, $j' = \tau(i, j)$, and similarly for the functions (λ, μ). Then the ring $K[G]$ has zero divisors: $$ (a_1 + a_2 + \dots + a_n + b_1 + b_2 + \dots + b_n)(c_1 + c_2 + \dots + c_n - d_1 - d_2 - \dots - d_n) = 0. $$ The authors look at this class in the simplest case, $n=2$. For any n, there are $(n^2!)^2$ of such groups. The authors showed that for $n=2$, any group of this class contains a 2-torsion. They also provided a set of groups such that any group $G(\sigma, \tau, \lambda, \mu)$ is isomorphic to some group from this set. Indeed, for $n=2$, it's very easy to check for torsion. Using Tietze transformations, we eliminate the generators $$ a_1 = b_{\sigma(1,1)} d_{\tau(1,1)} c_1^{-1}, \quad a_2 = b_{\sigma(2,1)} d_{\tau(2,1)} c_1^{-1} $$ Then the group $$G(\sigma, \tau, \lambda, \mu)$$ will be generated by 6 elements $$ b_1, b_2, c_1, c_2, d_1, d_2 $$ and defined by 6 relations $$ b_{\sigma(1,1)} d_{\tau(1,1)} c_1^{-1} c_2 = b_{\sigma(1,2)} d_{\tau(1,2)}, $$ $$ b_{\sigma(2,1)} d_{\tau(2,1)} c_1^{-1} c_2 = b_{\sigma(2,2)} d_{\tau(2,2)}, $$ $$ b_{\sigma(1,1)} d_{\tau(1,1)} c_1^{-1} d_1 = b_{\lambda(1,1)} c_{\mu(1,1)}, $$ $$ b_{\sigma(1,1)} d_{\tau(1,1)} c_1^{-1} d_2 = b_{\lambda(1,2)} c_{\mu(1,2)}, $$ $$ b_{\sigma(2,1)} d_{\tau(2,1)} c_1^{-1} d_1 = b_{\lambda(2,1)} c_{\mu(2,1)}, $$ $$ b_{\sigma(2,1)} d_{\tau(2,1)} c_1^{-1} d_2 = b_{\lambda(2,2)} c_{\mu(2,2)}. $$ Next, using the SageMath package, I pretty quickly checked that a group with such relations contains torsion using the $simplified()$ function (it simplifies the presentation). In exactly the same way, by excluding a_1, a_2, a_3 in the case $n=3$, I checked a large number of these groups for torsion, and they all contained 2-torsion, but there were also groups with 3-torsion. But for larger n, I could only check random combinations, and the $simplified()$ function was getting slower and slower. All such groups contain simple torsion of the form $(xy^{-1})^2$ or $(x^{-1}y)^2$. For example, for $n=2$ consider $$σ: [2, 1, 2, 1] \quad τ: [1, 1, 2, 2] \quad λ: [1, 2, 1, 2] \quad μ: [2, 2, 1, 1]$$ For a pair of (i, j) equal to $(1, 1)$, we choose $\sigma(1, 1) = 2$, $\tau(1,1) = 1$, $\lambda(1,1) = 1$, $\mu(1,1) = 2$. For example, for $(2,1)$ we get $\sigma(2,1) = 2$, $\tau(2,1)=2$, $\lambda(2,1)=1$, $\mu(2,1) = 1$ and so for all (i, j). Functions $(i, j) \mapsto (\sigma((i, j)), \tau((i, j)))$ and $(i, j) \mapsto (\lambda((i, j)), \mu((i, j))$ are biective. Probably, in this context, it would be wiser to represent σ, τ, λ, μ as matrices of size $n \times n$, but my code works this way. Next, we build a group G with 6 generating and relations with these indices. I get $G.simplified()$ = Finitely presented group < b_1, c_1, c_2, d_1 | $(c_2^{-1}d_1)^2$, $d_1c_2^{-1}c_1d_1^{-1}c_1c_2^{-1}(d_1c_1^{-1})^2$ >. The presentation is simplified so that it always has a with torsion element. I did very little testing for $n \ge 4$, only random combinations, but almost all of them contain 2-torsion. A year ago we were discussing these examples with a friend, and he doesn't believe at all that they could provide a counterexample. Because of this, I have 2 questions: Question 1: Do you believe that among these groups there could (or could not) be a group without torsion? I'm not asking for a strict justification, any intuitive arguments are welcome. I also want to quote a question from an article posed by Roman Mikhailov: Question 2: Could there be a group among these with non-obvious torsion? (Not with torsion of the type $(xy^{-1})^k$ or $(x^{-1}y)^k$ for $k \ge 2$.)

论坛问题帖 | has_answers | Maximka123 | 2026-07-22

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019466 | MSE-5147523

Do first semiprime $\gcd$ occurrences have twin-prime cofactors?

Conjecture Let $S_1<S_2<S_3<\cdots$ be the sequence of semiprimes. For every prime $p\ge 5$, let k be the smallest index such that $$ \gcd(S_k,\,S_{k+1})=p. $$ Then the two cofactors are twin primes: $$ \frac{S_k}{p},\quad \frac{S_{k+1}}{p}. $$ Searching up to $S_{k+1}\le10^{14}$ gives the following results: $$ \begin{array}{|c|r|r|r|r|} \hline \large p & \large S_k & \large S_{k+1} & \large \frac{S_k}{p} & \large \frac{S_{k+1}}{p}\\ \hline 5 & 3295 & 3305 & 659 & 661\\ 7 & 15659 & 15673 & 2237 & 2239\\ 11 & 99121 & 99143 & 9011 & 9013\\ 13 & 1036061 & 1036087 & 79697 & 79699\\ 17 & 33607963 & 33607997 & 1976939 & 1976941\\ 19 & 17654933 & 17654971 & 929207 & 929209\\ 23 & 80419201 &

80419247 & 3496487 & 3496489\\ 29 & 777474949 & 777475007 & 26809481 & 26809483\\ 31 & 1942680317 & 1942680379 & 62667107 & 62667109\\ 37 & 8246671703 & 8246671777 & 222883019 & 222883021\\ 41 & 10406805361 & 10406805443 & 253824521 & 253824523\\ 43 & 22126251011 & 22126251097 & 514563977 & 514563979\\ 47 & 66521681287 & 66521681381 & 1415354921 & 1415354923\\ 53 & 1123008419353 & 1123008419459 & 21188838101 & 21188838103\\ 59 & 1485678418201 & 1485678418319 & 25180990139 & 25180990141\\ 61 & 3724394496341 & 3724394496463 & 61055647481 & 61055647483\\ 67 & 12472468041743 & 12472468041877 & 186156239429 & 186156239431\\ 71 & 15781281240031 & 15781281240173 & 222271566761 & 222271566763\\ 73 & 62345036200451 & 62345036200597 & 854041591787 & 854041591789\\ 79 & 16288457597003 & 16288457597161 & 206183007557 & 206183007559\\ \hline \end{array} $$ Question Why should the first occurrence of a prime p as the $\gcd$ of two consecutive semiprimes always have twin-prime cofactors?

论坛问题帖 | has_answers | vengy | 2026-08-21

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019467 | MSE-5148175

Does every triangle satisfy $\frac{1}{R}\left(\frac{a}{A}+\frac{b}{B}+\frac{c}{C}\right)+\frac{A^2+B^2+C^2}{\pi^2} \ge 5$

Let a, b, c be the sides of a triangle and A,B,C be the corresponding opposite angles and R be its circumradius. Experimental data shows that π^2 is the largest i.e. optimal constant for which the inequality $$ \frac{1}{R}\left(\frac{a}{A} + \frac{b}{B} + \frac{c}{C}\right) + \frac{A^2 + B^2 + C^2}{\pi^2} \ge 5. $$ for all triangles. Can this be proved? Update: We have a slightly stronger result. Let $4 <x \le 5$ then $$ \frac{1}{R}\left(\frac{a}{A} + \frac{b}{B} + \frac{c}{C}\right) + \frac{x-4}{\pi^2}\left(A^2 + B^2 + C^2\right) \ge x. $$ Related question. Update: Unfortunately, the question was closed. Reason: " ...why the question is relevant to you and our community. Some forms of context include: background and motivation, relevant definitions, source, possible strategies, your current progress, why the question is interesting or important, etc." So my honest answers are as follows: Motivation: The only motivation for me the beauty of math; I find an inequality not because I expect it to be useful or because it came out of a deeper theory etc. It may not be interesting/beautiful for some, but interest is relative and to me this is. Many of us do math just for the beauty of it, it does not have to need more motivation than this. Why is it relevant for the community: It is not a homework question; it is also not a deep research level MathOverflow type of question. It is a perfect example of a new inequality coming out of experimentation, so where else would it belong? Progress: I have discovered a plausible inequality which did not exist few hours back and I have done an extensive experimental verification. If you are of the type who only consider progress as only rigorous progress towards a proof, then I can't help it; to me experimental conjectures is also progress.

论坛问题帖 | has_answers | Nilotpal Kanti Sinha | 2026-08-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019468 | MSE-5148506

How large can the area be of a triangle whose side lengths are the angles of another triangle?

Let a,b,c be the side lengths of an acute triangle, with opposite angles A,B,C, and let K_a denote its area. Since the triangle is acute, the angles A,B,C themselves satisfy the triangle inequalities, and hence may be regarded as the side lengths of a second triangle. Let K_A denote the area of this angled triangle. This suggests the natural question: $$ \boxed{\text{How large can }K_A\text{ be relative to }K_a\text{?}} $$ More specifically, is there a sharp inequality relating the normalized areas of the original triangle and its angle triangle? Numerical evidence suggests that $$ \frac{K_A}{K_a} \le \frac{\pi^2}{(a+b+c)^2}. $$ The constant π^2 is best possible, and equality holds precisely for equilateral triangles. Furthermore, if the original triangle is inscribed in a unit circle (equivalently, its circumradius is $R=1$), then the stronger universal bound $$ \frac{K_A}{K_a}<\frac{\pi}{8} $$ holds. The constant $\pi/8$ and π^2 are best possible. Can this be proved?

论坛问题帖 | has_answers | Nilotpal Kanti Sinha | 2026-09-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019469 | MSE-5150395

Does the interval $(n^2, 4n^2)$ always contain a Mersenne prime exponent?

I have been analyzing the exponents p of known Mersenne primes $(2^p - 1)$ and noticed a pattern. I am conjecturing the following: For every $n \ge 1$, there exists a Mersenne prime exponent p such that: $n^2 < p < 4n^2$ This holds true for all 52 currently known Mersenne primes. The tightest squeeze occurs between $p = 127$ and $p = 521$: For $n = 11$, the interval is $(121, 484)$, catching $p = 127$. For $n = 12$, the interval is $(144, 576)$, catching $p = 521$. My Question: A formal proof is impossible right now since we don't even know whether or not there are infinitely many Mersenne primes. But from a probabilistic standpoint, does the Lenstra-Pomerance-Wagstaff heuristic suggest this pattern should hold globally, or is a massive gap statistically guaranteed to ruin it for a very large n?

论坛问题帖 | no_answers_at_retrieval | Mildred Kemunto | 2026-09-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019470 | MSE-5151707

Bijection related to formula for number of k length compositions of positive integer n and subsets of a set

From the book Enumerative Combinatorics 2nd ed. by R.P.Stanley, top of page 25 How does he come up with that $\binom{n-1}{k-1}$ answer and what does it have to do with bijection and maybe a precise answer as to what does bijection and it's inverse really mean ? I understand for example $(1+x)^n=\sum_{k=0}^n\binom {n}{k}x^k$ and for example that the number of solutions in k non negative integers to $n_1+n_2+...+n_k=n$ is $\binom{n+k-1}{k-1}$ so the number of solutions in strictly positive integers to $n_1+n_2+...+n_k=n+k$ is the same $\binom{n+k-1}{k-1}$ so then I could just put the

number of solutions to $n_1+n_2+...+n_k=n$ would just be to replace the n+k-1 in the first arg. of $\binom{n+k-1}{k-1}$ by just $n-1$ and would get his answer of $\binom{n-1}{k-1}$ but something tells me that is not a valid maneuver so not a valid way of proving it but just a coincidence ? And besides even if it is valid have not done it by method of bijection related to number of subsets of a set which is what he proclaims to be the way of envisioning such a conclusion since he states on the prior page 24 "There is a close connection between subsets of a set and compositions of a nonnegative integer." So does anyone have a clear way of describing all this and related subset of a set bijection ?

论坛问题帖 | has_answers | user158293 | 2026-10-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019471 | MSE-5152158

State of Baranovsky conjecture

Vladimir Baranovsky in his paper "The Variety of Pairs of Commuting Nilpotent Matrices is Irreducible", published in Transformation Groups, Vol 6, No.1, pages 3-8, 2001, posts a conjecture that all irreducible components of $\mathcal{N}_2(\mathfrak{g})$ (variety of nilpotent pairs of elements of complex semisimple Lie algebra $\mathfrak{g}$) have maximal dimension $\dim(\mathfrak{g})$, see end of Section 2 of the paper. He proves it for $\mathfrak{sl}_n$, i.e. for simple Lie algebras of type A. Now I wonder - Baranovsky paper is 25 years old and in meantime commuting nilpotent varieties got serious manpower. So, what is the state of his conjecture now?

论坛问题帖 | has_answers | Dibidus | 2026-10-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019472 | MSE-516263

A conjectural closed form for $\sum\limits_{n=0}^\infty\frac{n!\,(2n)!}{(3n+2)!}$

Let $$S=\sum\limits_{n=0}^\infty\frac{n!\,(2n)!}{(3n+2)!},\tag1$$ its numeric value is approximately $S \approx 0.517977853388534047...$${}^{[more\ digits]}$ S can be represented in terms of the generalized hypergeometric function: $$S={}_3F_2\left(\frac12,1,1;\ \frac43,\frac53;\ \frac4{27}\right)\cdot\frac12.\tag2$$ Let σ be the closed-form expression constructed from integers and elementary functions as follows: $$\sigma=3\,\alpha\,\ln(2\,\alpha+1)-\sqrt{\beta\,}\arccos\gamma,\tag3$$ where $$\alpha=\frac{\sqrt[3]{3\,}}{6\,\sqrt[3]{2\,}}\left(\sqrt[3]{9-\sqrt{69}}+\sqrt[3]{9+\sqrt{69}}\right),\tag4$$ $$\beta=\frac1{4\,\sqrt[3]{2\,}}\left(\sqrt[3]{25+3\,\sqrt{69}}+\sqrt[3]{25-3\,\sqrt{69}}\right)-\frac12,\tag5$$ $$\gamma=\frac1{6\,\sqrt[3]{2\,}}\left(\sqrt[3]{57\,\sqrt{69}-459}-\sqrt[3]{57\,\sqrt{69}+459}\right)+\frac12\tag6$$ are the unique real roots of the following cubic equations: $$8\,\alpha^3-2\,\alpha-1=0,\tag7$$ $$64\,\beta^3+96\,\beta^2+36\,\beta-23=0,\tag8$$ $$8\,\gamma^3-12\,\gamma^2+16\,\gamma+11=0.\tag9$$ Equivalently, $$\sigma = \frac{3\,p}{2}\,\ln\big(p+1\big)-\frac{1}{2}\sqrt{\frac{3-p}{p}}\arccos\Big(\frac{p-6}{6p+2}\Big)\tag{10}$$ where p is the plastic constant or the real root of $$p^3-p-1=0\tag{11}$$ It can be numerically checked that the following inequality holds: $$\Big|S-\sigma\Big|<10^{-10^5},\tag{12}$$ I conjecture that the actual difference is the exact zero, and thus S has an elementary closed form: $$\sum\limits_{n=0}^\infty\frac{n!\,(2n)!}{(3n+2)!}\stackrel?=3\,\alpha\,\ln(2\,\alpha+1)-\sqrt{\beta\,}\arccos\gamma,\tag{13}$$ I am asking for you help in proving this conjecture.

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2013-10-06

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019473 | MSE-520801

A conjecture $\int_{-\infty}^\infty\frac{\arctan e^x}{\cosh x}\cdot\frac{\tanh\frac{x}2}{x}dx\stackrel?=\frac\pi2\ln2$

I need to find a closed form for this integral: $$\mathcal{I}=\int_{-\infty}^\infty\frac{\arctan e^x}{\cosh x}\cdot\frac{\tanh\frac{x}2}{x}dx.$$ A numerical integration results in an approximation $\mathcal{I}\approx1.0887930451518...$, and WolframAlpha suggests a possible closed form for this number: $$\mathcal{I}\stackrel?=\frac\pi2\ln2$$ Is it the correct value of this integral? If so, how to prove it?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2013-10-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019474 | MSE-521685

Integral $\int_0^\infty\frac{dx}{\frac{x^4-1}{x\cos(\pi\ln x)+1}+2x^2+2}$

I need your help with this integral: $$\int_0^\infty\frac{dx}{\frac{x^4-1}{x\cos(\pi\ln x)+1}+2\,x^2+2}.$$ I wasn't able to evaluate it in a closed form, although an approximate numerical evaluation suggested its value could be $\frac{\pi}{4}$.

论坛问题帖 | has_answers | Marty Colos | 2013-10-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019475 | MSE-521993

Integral $\int_1^\infty\operatorname{arccot}\left(1+\frac{2\pi}{\operatorname{arcoth}x-\operatorname{arccsc}x}\right)\frac{\mathrm dx}{\sqrt{x^2-1}}$

Consider the following integral: $$\mathcal{I}=\int_1^\infty\operatorname{arccot}\left(1+\frac{2\,\pi}{\operatorname{arcoth}x\,-\,\operatorname{arccsc}x}\right)\frac{\mathrm dx}{\sqrt{x^2-1}}\,,$$ where arccsc is the inverse cosecant, arccot is the inverse cotangent and $\operatorname{arcoth}x$ is the inverse hyperbolic cotangent. Approximate numerical integration suggests a possible closed form: $$\mathcal{I}\stackrel?=\frac{\pi\,\ln\pi}4-\frac{3\,\pi\,\ln2}8.$$ I was not able to rigorously establish the equality, but the value is correct up to at least 900 decimal digits. Is it the correct exact value of the integral $\,\mathcal{I}$?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2013-10-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019476 | MSE-522764

Simplify $\frac{_3F_2\left(\frac{1}{2},\frac{3}{4},\frac{5}{4};1,\frac{3}{2};\frac{3}{4}\right)}{\Pi\left(\frac{1}{4}\big|\frac{1}{\sqrt{3}}\right)}$

Is it possible to simplify the ratio $$\mathcal{E}=\frac{_3F_2\left(\frac{1}{2},\frac{3}{4},\frac{5}{4};\ 1,\frac{3}{2};\ \frac{3}{4}\right)}{\Pi\left(\frac{1}{4}\Big|\frac{1}{\sqrt{3}}\right)},$$ where $\Pi(n|k)$ is the complete elliptic integral of the third kind? Its numeric value is approximately $\mathcal{E}\approx0.73510519389572273268...$, that looks like $\frac{4}{\pi\sqrt{3}}$.

论坛问题帖 | has_answers | Laila Podlesny | 2013-10-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019477 | MSE-522998

Conjecture $\int_0^1\frac{\ln\left(\ln^2x+\arccos^2x\right)}{\sqrt{1-x^2}}dx\stackrel?=\pi\,\ln\ln2$

$$\int_0^1\frac{\ln\left(\ln^2x+\arccos^2x\right)}{\sqrt{1-x^2}}dx\stackrel?=\pi\,\ln\ln2$$ Is it possible to prove this?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2013-10-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019478 | MSE-526990

What are the recent advancements in mathematics that an undergraduate can understand?

I am an undergraduate student of mathematics. I am interested to know the conjectures that are proved in the last, say 5 or 10 years. Or any other development. Preferably in pure mathematics. One of the motives to ask this question is to get a feel for the cutting edge research areas. (If you feel this is too broad, please suggest me in comments to improve this question.) EDIT: I am interested to understand (not just know the statement of) the theorems. In your answers, you could just state the theorems, with references where I can research. If proofs are also provided, that would of course be acknowledged with deeper gratitude.

论坛问题帖 | no_answers_at_retrieval | math | 2013-10-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019479 | MSE-530058

Totient summatory function

Let $\Phi(n) = \sum_{k=1}^n \phi(k)$ be the totient summatory function. Here is an interesting conjecture I've made: The ratio $\Phi(n^2)/\Phi(n)$ is an integer only for $n=1,2,3,5$ and 6. I made a program that shows that these are only solution for about $n<3000$. Is there some way to prove or disaprove it? Eventually, what about the general case: what conditions should satisfy numbers n and m so that ratio of their totient summatory functions is integer? Is there some paper that deals with it?

论坛问题帖 | has_answers | user101521 | 2013-10-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019480 | MSE-532089

What's wrong with my conjecture?

I was doing math homework, and I formulated the following conjecture from one of the questions: If $f(x)$, $g(x)$ and $h(x)$ are continuous functions and the equations $f(x) = h(x)$ and $g(x) = h(x)$ both have only one root, then the equation $f(x) = g(x)$ has only one root. So can anyone find a counterexample? I can't think of one, but maybe that's because I haven't read Counterexamples in Analysis yet.

论坛问题帖 | has_answers | 11Kilobytes | 2013-10-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019481 | MSE-532260

Prove $_4F_3(1/8,3/8,5/8,7/8;1/4,1/2,3/4;1/2)=\frac{\sqrt{2-\sqrt2+\sqrt{2-\sqrt2}}+\sqrt{2+\sqrt2+\sqrt{2+\sqrt2}}}{2\,\sqrt2}$

How do I prove $_4F_3\left(\frac18,\frac38,\frac58,\frac78;\ \frac14,\frac12,\frac34;\ \frac12\right)\stackrel?=\frac{\sqrt{2-\sqrt2+\sqrt{2-\sqrt2}}+\sqrt{2+\sqrt2+\sqrt{2+\sqrt2}}}{2\,\sqrt2}$$ I encountered this conjecture when working with Fourier transforms of Kelvin functions.

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2013-10-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019482 | MSE-533166

My conjecture on almost integers.

Here when I was studying almost integers , I made the following conjecture - Let x be a natural number then For sufficiently large n (Natural number) Let $$\Omega=(\sqrt x+\lfloor \sqrt x \rfloor)^n$$ then Ω is an almost integer . The value of n depends upon the difference between the number x and its nearest perfect square which is smaller than it.Can anyone prove this conjecture. Moreover, I can provide examples like $(\sqrt 5+2)^{25}=4721424167835364.0000000000000002$ $(\sqrt 27+5)^{15}=1338273218579200.000000000024486$

论坛问题帖 | has_answers | Shivam Patel | 2013-10-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019483 | MSE-538564

Conjecture $\int_0^1\frac{dx}{\sqrt[3]x\,\sqrt[6]{1-x}\,\sqrt{1-x\left(\sqrt{6}\sqrt{12+7\sqrt3}-3\sqrt3-6\right)^2}}=\frac\pi9(3+\sqrt2\sqrt[4]{27})$

Let $$\alpha=\sqrt{6}\ \sqrt{12+7\,\sqrt3}-3\,\sqrt3-6\,=\,\big(2+\sqrt{3}\big) \big(\sqrt{2}\ \sqrt[4]{27}-3\big)\,=\,\frac{3\sqrt{3}}{3+\sqrt2\ \sqrt[4]{27}}.\tag1$$ Note that α is the unique positive root of the polynomial equation $$\alpha^4+24\,\alpha^3+18\,\alpha^2-27=0.\tag2$$ Now consider the following integral: $$I=\int_0^1\frac{dx}{\sqrt[3]x\ \sqrt[6]{1-x}\ \sqrt{1-x\,\alpha^2}}.\tag3$$ I have a conjectured elementary value for it $$I\stackrel?=\frac\pi9\Big(3+\sqrt2\ \sqrt[4]{27}\Big)=\color{blue}{\frac{\pi}{\sqrt{3}\,\alpha}}.\tag4$$ Actually, the integral

I can be evaluated in an exact form using Mathematica or manually, using formula DLMF 15.6.1: $$I=\frac{4\,\sqrt\pi}{\sqrt3}\cdot\frac{\Gamma\left(\frac56\right)}{\Gamma\left(\frac13\right)}\cdot{_2F_1}\left(\frac12,\frac23;\ \frac32;\ \alpha^2\right),\tag5$$ but I could not find a way to simplify this result to (4). So, my conjecture can be restated in a different form: $${_2F_1}\left(\frac12,\frac23;\ \frac32;\ \alpha^2\right)\stackrel?=\frac{\sqrt2+\sqrt[4]3}{4\,\sqrt[4]{27}}\cdot\frac{\Gamma\left(\frac13\right)}{\Gamma\left(\frac56\right)}\cdot\sqrt\pi\tag6$$ or $$\sum_{n=0}^\infty\frac{\Gamma\left(n+\frac23\right)}{(2\,n+1)\,\Gamma(n+1)}\alpha^{2\,n}\stackrel?=\frac{3+\sqrt2\,\sqrt[4]{27}}{18}\cdot\frac{\pi^{3/2}}{\Gamma\left(\frac56\right)}.\tag7$$ The conjecture can also be given in terms of the incomplete beta function: $$B\left(\alpha^2;\ \frac12,\frac13\right)\stackrel?=\frac{\sqrt\pi}2\cdot\frac{\Gamma\left(\frac13\right)}{\Gamma\left(\frac56\right)}.\tag8$$ Question: Is the conjecture indeed true? Note: It holds numerically up to at least 10^4 decimal digits. Conjecture 2 Let $$ {\small \begin{multline} \beta=\frac{21}4+\frac{9\,\sqrt5}4-\frac{15}8\sqrt{750-330\,\sqrt5}-\frac{33}8\sqrt{150-66\,\sqrt5} \\ + \frac12\sqrt{3\left(165+75\,\sqrt5-46\,\sqrt{750-330\,\sqrt5}-103\,\sqrt{150-66\,\sqrt5}\right)}. \end{multline}}\tag9$$ Added later: We can simplify it to $$ \small\beta=\frac 3 4 \left(7+3 \sqrt 5-\sqrt[4] 5 \sqrt{66+30 \sqrt 5}\right)+\frac 1 2 \sqrt{495+225 \sqrt 5-3 \sqrt{6 \big(8545+3821 \sqrt 5\big)}}.\tag{$9'$} $$ I conjecture that: $$\int_0^1\frac{dx}{\sqrt[3]x\ \sqrt[6]{1-x}\ \sqrt{1-x\,\beta^2}}\stackrel?=\color{blue}{\frac{2\,\pi}{5\,\sqrt3\,\beta}}.\tag{10}$$ I can imagine that if these conjectures are true, then there are generalizations for some other algebraic numbers.

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2013-10-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019484 | MSE-541135

Conjecture $\int_0^1\frac{\mathrm dx}{\sqrt{1-x}\ \sqrt[4]x\ \sqrt[4]{2-x\,\sqrt3}}\stackrel?=\frac{2\,\sqrt2}{3\,\sqrt[8]3}\pi$

$$\int_0^1\frac{\mathrm dx}{\sqrt{1-x}\ \sqrt[4]x\ \sqrt[4]{2-x\,\sqrt3}}\stackrel?=\frac{2\,\sqrt2}{3\,\sqrt[8]3}\pi\tag1$$ The equality numerically holds up to at least 10^4 decimal digits. Can we prove that the equality is exact? An equivalent form of this conjecture is $$I\left(\frac{\sqrt3}2;\ \frac14,\frac14\right)\stackrel?=\frac23,\tag2$$ where $I\left(z;\ a,b\right)$ is the regularized beta function. Even simpler case: $$\int_0^1\frac{\mathrm dx}{\sqrt{1-x}\ \sqrt[6]{9-x}\ \sqrt[3]x}\stackrel?=\frac\pi{\sqrt3},\tag3$$ which is equivalent to $$I\left(\frac19;\ \frac16,\frac13\right)\stackrel?=\frac12.\tag4$$ A related question.

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2013-10-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019485 | MSE-541581

Existence of a prime between ap and $(a+1)p$ - generalization of Bertrand's postulate

Conjecture: There exists at least one prime number p_{m} : $ap_{n} < p_{m} < (a+1)p_{n}$, $\forall$ $a \in \mathbb{N}$ and $\forall$ p_{n} $\in \mathbb{P} $ if $(a+1)p_{n} < p_{n+1}^2$. Is there a name for this conjecture, and has it been proven or disproven?

论坛问题帖 | has_answers | Brad Graham | 2013-10-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019486 | MSE-541727

Prime gaps with respect to the squared primes

Conjecture If we have two consecutive prime numbers p_{a} and p_{a+1}, and two other consecutive primes p_n and p_{n+1}, so that $p_{a} < p_{a+1} < p^2_{n+1}$, then $p_{a+1} - p_{a} < 2p_{n} $. Are there any known counter examples and are there any known similar conjectures?

论坛问题帖 | has_answers | Brad Graham | 2013-10-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019487 | MSE-542220

Does there exist a prime number within the interval?

Conjecture $\forall p_{n}\in \mathbb{P} : n\geq3, \: \exists p_{m}\in \mathbb{P} : 3p_{n} - 4 \geq p_{m} > \sqrt{2(p^2_{n+1} - 1)} $ How would you go about proving/disproving this?

论坛问题帖 | has_answers | Brad Graham | 2013-10-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019488 | MSE-543442

Minimum amount of primes between squared primes

Conjecture: $\forall $ p_{n}, $p_{n+1} \in \mathbb{P}$, $\:$ $p^2_{n+1} = p^2_{n} +\omega_{n} p_{n} + \phi_{n} : \phi_{n} , \omega_{n} \in \mathbb{N} $ and $ \phi_{n} < p_{n}$, $\:$ $\Rightarrow$ $\:$$\exists$ atleast $ \omega_{n}$ prime numbers in the interval (p^2_{n},p^2_{n+1}) How would you prove this?

论坛问题帖 | no_answers_at_retrieval | Brad Graham | 2013-10-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019489 | MSE-544807

Conjecture: $\int_0^1\frac{3x^3-2x}{(1+x)\sqrt{1-x}}K\big(\!\frac{2x}{1+x}\!\big)\,dx\stackrel?=\frac\pi{5\sqrt2}$

$$\int_0^1\frac{3x^3-2x}{(1+x)\sqrt{1-x}}K\left(\frac{2x}{1+x}\right)\,dx\stackrel?=\frac\pi{5\sqrt2}$$ The integral above comes from the evaluation of the integral $A=\int_0^{\pi/2}\frac{f(\theta)}\pi\,d\theta$, where $$f(\theta)=\int_0^\pi\frac{(3\sin^2\theta-2)\sin\theta\,d\phi}{\sqrt{2+2\sin\theta\cos\phi}}=\frac{\sqrt2(3\sin^2\theta-2)}{\csc\theta\sqrt{1+\sin\theta}}\int_0^{\pi/2}\left(1-\frac{2\sin\theta}{1+\sin\theta}\sin^2\gamma\right)^{-1/2}d\gamma,$$ where $\gamma=\phi/2$, and the right side integral is

$K\big(\!\frac{2\sin\theta}{1+\sin\theta}\!\big)$ by definition. After substitutions $x=\sin\theta$ and $y=\frac{2x}{1+x}$ we get $$\begin{align}A&=\frac{\sqrt2}\pi\int_0^1\frac{3x^3-2x}{(1+x)\sqrt{1-x}}K\left(\frac{2x}{1+x}\right)\,dx\\ &=\frac1\pi\int_0^1\frac{y^3+8y^2-8y}{(2-y)^3\sqrt{y^2-3y+2}}K(y)\,dy\stackrel ?=\frac15,\end{align}$$ where the integral has been numerically evaluated to suggest the analytic result on the right to several thousand digits. Is there any way to prove this equality, and are there any generalizations of this conjecture to include other parameters? P.S. Not that it's overly relevant to the question, but for the interested reader, the original integral comes from the following physics problem: [Morin Intro to Classical Mechanics, Ex. 10.12] The earth bulges slightly at the equator, due to the centrifugal force in the earth's rotating frame. The goal of this exercise is to find the shape of the earth, first incorrectly, and then correctly. (a) The common incorrect method is to assume that the gravitation force from the slightly nonspherical earth points toward the center, and to then calculate the equipotential surface (incorporating both the gravitational and centrifugal forces). Show that this method leads to a surface whose height (relative to a spherical earth of the same volume) is given by $h(\theta)=R\big(\!\frac{R\omega^2}{6g}\!\big)(3\sin^2\theta-2)$, where θ is the polar angle (the angle down from the north pole), and R is the radius of the earth. (b) The above method is incorrect, because the slight distortion of the earth causes the gravitational force to not point toward the center of the earth (except at the equator and the poles). This tilt in the force direction then changes the slope of the equipotential surface, and it turns out (although this is by no means obvious) that this effect is of the same order as the slope of the surface found in part (a). Your task: Assuming that the density of the earth is constant, and that the correct height takes the form of some constant factor f times the result found in part (a), show that $f=5/2$. Do this by demanding that the potential at a pole equals the potential at the equator.

论坛问题帖 | has_answers | Mario Carneiro | 2013-10-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019490 | MSE-551337

is this conjecture true or false?

I want to know if this conjecture istrue or false $$\Large e^{\frac{ \ln x}{x}} \notin \mathbb{Z} $$ for every $x \in \mathbb{R} \setminus \{1,-1,0\} $

论坛问题帖 | has_answers | user95440 | 2013-11-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019491 | MSE-552593

Lines formed from vertices of n-gons equate to triangular numbers.

Noticed something neat tonight! The number of unique lines you can form by connecting the vertices of an n-gon is equal to the (n-1)th triangular number. (e.g. in a square all 4 veritices make 4 lines (4 sides to a square), and then I can also connect any verticie to another verticie that is not directly to the left or right, assuming that it hasn't already been connected to that one earlier. So on a square the top right can connect to the bottom left (later on the bottom left can't connect to the top right because it already has), and the top left and connect to the bottom right. 6 unique lines (the 3rd triangular number is 6!) This works for any n-gon too!) I'm wondering if this property has already been known, I couldn't find anything about it online. Also, if so, does it have a name? I can't think of any real application for it other than, if this is held to be true than those who believe a digon is a circle/ellipsoid/ovaloid would be wrong (and it is currently widely held that a digon is a circle/ellipsoid/ovaloid), because then the 1st triangle-number would be "2", but if a digon was rather just a straight line with 2 end points, then the first triangular number would be "1" (which it is!) A digon would be two lines (NOT CURVES) that overlap and form a closed polygon with area 0 and perimeter 2*s.

论坛问题帖 | has_answers | Albert Renshaw | 2013-11-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019492 | MSE-566404

An integral $\int_0^\infty P_s(x-1)\,e^{-x}\,dx$ involving Legendre functions

Let $P_s(x)$ denote the Legendre functions of the 1^{st} kind, i.e. the Legendre polynomial generalized to an arbitrary (not necessarily integer) order s. It can be expressed using the hypergeometric function: $$P_s(x)={_2F_1}\left(-s,s+1;\ 1;\ \frac{1-x}2\right).\tag1$$ Let's consider the definite integral $$\mathcal{J}(s)=\int_0^\infty P_s(x-1)\,e^{-x}\,dx.\tag2$$ It evaluates to integer values when $s\in\mathbb{Z}^+$. Using a computer-assisted search for the general term formula, I discovered the following conjecture, that I haven't yet been able to prove: $$\mathcal{J}(s)\stackrel?=\left(K_{3/2}(1)\cdot I_{s+1/2}(-1)-I_{3/2}(-1)\cdot K_{s+1/2}(1)\right)\sqrt{-1},\tag3$$ where $I_\nu(z)$ and $K_\nu(z)$ are the modified Bessel functions of the 1^{st} and 2^{nd} kind. Unfortunately, it only seems to hold for $s\in\mathbb{Z}^+$, and does not generalize to non-integer values of s. Can we prove the conjecture (3)? Can we find a more general formula that holds not only for integer values of s? Can we find (or at least conjecture) a closed form for $\mathcal{J}(1/2)$?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2013-11-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019493 | MSE-568615

Conjectured closed form for $\int_0^1x^{2\,q-1}\,K(x)^2dx$ where $K(x)$ is the complete elliptic integral of the 1st kind

I am interested in a general closed-form formula for integrals of the following form: $$\mathcal{J}_q=\int_0^1x^{2\,q-1}\,K(x)^2dx,\tag0$$ where $K(x)$ is the complete elliptic integral of the 1st kind: $$K(x)={_2F_1}\left(\frac12,\frac12;\ 1;\ x^2\right)\frac\pi2\\=\int_0^{\pi/2}\frac{d\phi}{\sqrt{1-x^2\sin^2\phi}}=\int_0^1\frac{dz}{\sqrt{1-x^2z^2\vphantom{|^2}}\sqrt{1-z^2\vphantom{|^2}}}.\tag1$$ I could not find a suitable integral in DLMF or Gradshteyn-Ryzhik or Prudnikov tables of integrals. Mathematica was not able to evaluate it either, even when the parameter q was fixed to an integer (Note: In Mathematica you would write EllipticK[x^2] for $K(x)$ because of a different convention for the modulus parameter). But, numerical integration and lookups in WolframAlpha and ISC+ suggest that there are closed forms for (at least some) postitive integer values of the parameter q. For example, $$\mathcal{J}_1\stackrel?=\frac{7\,\zeta(3)}4,\ \mathcal{J}_2\

stackrel?=\frac{7\,\zeta(3)}8+\frac14,\\\mathcal{J}_3\stackrel?=\frac{77\,\zeta(3)}{128}+\frac{17}{64},\
\mathcal{J}_4\stackrel?=\frac{119\,\zeta(3)}{256}+\frac{881}{3456},\tag2$$ where $\zeta(3)$ is the Apéry constant. It looks that the general formula for postitive integer values of q is $$\mathcal{J}_q\stackrel?=a_q\,\zeta(3)+b_q,\tag3$$ where $a_q,\,b_q$ are some rational coefficients. I tried to find general formulae for these coefficients, and actually discovered plausible candidates for both. Conjecturally, a_q can be expressed in terms of a generalized hypergeometric function: $$a_q\stackrel?=\frac{7\,\pi}4\cdot\frac{{_4F_3}\left(\begin{array}{c}\frac12,\,\frac12,\,1-q,\,1-q\\1,\,\frac32-q,\,\frac32-q\end{array}\middle|\ 1\right)}{\Gamma\left(\frac32-q\right)^2\ \Gamma(q)^2}.\tag4$$ Somewhat suprisingly, this scarish expression seems to evaluate only to rationals for all $q\in\mathbb{Z}^+$. For b_q I found only a conjectural recurrence relation: $$b_1\stackrel?=0,\ b_2\stackrel?=\frac14,\ b_q\stackrel?=\frac{2\,(2\,q-3)\,\left(2\,q^2-6\,q+5\right)\cdot b_{q-1}-4\,(q-2)^3\cdot b_{q-2}+1}{4\,(q-1)^3},\tag5$$ but I could not find a general term formula for it. Can we prove that the closed forms shown in (2) are correct? Can we prove that the formula (3) holds for all $q\in\mathbb{Z}^+$ with some rational coefficients $a_q,\,b_q$? Are my conjectured formulae for these coefficients $(4),\,(5)$ correct? Can the formula (4) be expanded in terms of simpler functions? Is there a general term formula for (5)? Can we find (or at least conjecture) a general closed-form formula for $\mathcal{J}_q$ where q is not necessarily an integer?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2013-11-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019494 | MSE-58638

open conjectures in real analysis targeting real valued functions of a single real variable

I am hoping that this question (if in acceptable form) be community wiki. Are there any open conjectures in real analysis primarily targeting real valued functions of a single real variable ? (it may involve other concepts but primarily targeting these type of functions and no involvement of number theory). Its not that i am going to attack them right away (not due to lack of interest but time) but am just curious to know them. Also how difficult it is get a new result in this area ?

论坛问题帖 | has_answers | Rajesh D | 2011-08-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019495 | MSE-588680

Conjecture $_2F_1\left(\frac14,\frac34;\,\frac23;\,\frac13\right)=\frac1{\sqrt{\sqrt{\frac4{\sqrt{2-\sqrt[3]4}}+\sqrt[3]{4}+4}-\sqrt{2-\sqrt[3]4}-2}}$

Using a numerical search on my computer I discovered the following inequality: $$\left|\,{_2F_1}\left(\frac14,\frac34;\,\frac23;\,\frac13\right)-\rho\,\right|<10^{-20000},\tag1$$ where ρ is the positive root of the polynomial equation
$$12\,\rho^8-12\,\rho^4-8\,\rho^2-1=0,\tag2$$ that can be expressed in radicals: $$\rho=\frac1{\sqrt{\sqrt{\frac4{\sqrt{2-\sqrt[3]{4\vphantom{\large1}}}}}+\sqrt[3]{4}+4}-\sqrt{2-\sqrt[3]4}-2}}.\tag3$$ Based on this inequality I conjecture that the actual difference is the exact zero, i.e.

$$\color{#808080}{_2F_1\left(\frac14,\frac34;\,\frac23;\,\frac13\right)=\rho}.\tag4$$ I looked up in DLMF and MathWorld, but did not find a known special value with exactly these parameters. It also appears that CAS like Maple or Mathematica do not know this identity. Could you please suggest any ideas how to prove the conjecture (4)? Update: I can propose even more general conjecture: $$\color{#808080}{27\,(x-1)^2\cdot{_2F_1}\left(\tfrac14,\tfrac34;\tfrac23;x\right)^8+18\,(x-1)\cdot{_2F_1}\left(\tfrac14,\tfrac34;\tfrac23;x\right)^4-8\cdot{_2F_1}\left(\tfrac14,\tfrac34;\tfrac23;x\right)^2=1}$$

论坛问题帖 | has_answers | HW[U+FFA0] | 2013-12-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019496 | MSE-600024

making mathematical conjectures

If a non-mathematician wanted to conjecture something and had strong numerical evidence to support the conjecture, how would he/she go about doing so? Would the mathematical community (a) take it seriously? (b) even look at it?

论坛问题帖 | has_answers | martin | 2013-12-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019497 | MSE-601386

Why is $2^a > a^3$?

I found this rather interesting and maybe, a bit too obvious for some people property about 2 raised to some power. $2^a > a^3$, if $a=0,a=1 \text{ or } a\ge 10$.($a \in N$) I seem to get a bit of intuition as to why this happens(although the intuition is in "micro units of intuition"). Somebody please provide me some intuition regarding it the mathematical/logic way.

论坛问题帖 | has_answers | Shaurya Gupta | 2013-12-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019498 | MSE-608304

If AC is false , is this statement about the halting problem true?

Assume AC is false. (AC = axiom of choice) Let n,m be positive integers. Let $f: \Bbb N \rightarrow \Bbb N$ and $f(m)=m$. Let $g(n,m)=1$ if the iterations $f(n),f(f(n)),...$ converges to m. If $g(n,m)=1$, let $G(n,g)=x$ be the (min) amount of iterations needed to arrive at m. Statement : If one can prove that $g(n,m)=1$ for all n then one can prove an upper boundary $T(n)$ for $G(n,g)$. For example if one could prove Collatz : $g(n,1)=1$ is true for all n <=> Collatz is true. $\implies G(n,g) < T(n)$ and $T(n)$ equals (for example) $2^{2^{2^n}}$ This statement comes from someone I know, and I was not able to find a counterexample. Im not sure if this statement is a named conjecture or theorem yet. And I do not know if its true of course. I do know however that ZFC is required to prove Goodstein's theorem and there is no upper bound known for $G(n,g)$ here. So I think if AC is true then the statement is false. I considered this statement to be an axiom too , but that would mean it cannot be proved or disproved which I doubt. But who knows.

论坛问题帖 | has_answers | mick | 2013-12-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019499 | MSE-620626

Proof on a conjecture involving $d(N)$

Let $d(n)$ denote the number of divisors of $n $. Then let N be a number such that $d(N)$ divides N . Also let $I= \frac{N}{d(N)}$ which is defined as the "Index of Beauty of N ". Then prove that - For every number I there exists a number N such that I is the index of beauty of N.

论坛问题帖 | has_answers | Shivam Patel | 2013-12-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019500 | MSE-620629

Are there infinitely many primes p such that $\frac{(p-1)! +1}{p}$ is prime?

Here I have the following conjecture: Let $$S_1(n)= \frac{(n-1)! +1}{n}.$$ Then there exist infinite prime numbers p for which $S_1(p)$ is prime. And I don't know how to prove it. EDIT Let $C_1(n)=[S_1(n)]$ where $[n]$ denotes the integral part. Then I have another conjecture that $[C_1(1) ,C_1(2),C_1(3) ,C_1(4),C_1(5) ,C_1(6)...]$ contains infinite many primes . Are there any ways to prove it?

论坛问题帖 | has_answers | Shivam Patel | 2013-12-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019501 | MSE-633495

A conjecture on $\phi(n)$

Let $\phi(n)$ denote the Euler totient function of $n $. Then let N be a number such that $\phi(N)$ divides N . Also let $I_1= \frac{N}{\phi(N)}$ which is defined as the "Second order Index of Beauty of N ". Then prove that - For every number I_1 there exists a number N such that I_1 is the second order index of beauty of N.

论坛问题帖 | has_answers | Shivam Patel | 2014-01-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019502 | MSE-63862

A conjecture about the form of some prime numbers

Let k be an odd number of the form $k=2p+1$,where p denote any prime number, then it is true that for each number k at least one of $6k-1$, $6k+1$ gives a prime number. Can someone prove or disprove this statement?

论坛问题帖 | has_answers | Pedja | 2011-09-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019503 | MSE-645415

Conjecture involving semi-prime numbers of the form $2^{x}-1$

Let x be a positive integer such that $(2^{x}-1)=pq$, where p and q are prime numbers. I want to show that either $p^{2} \bmod x \equiv 1$ or $q^{2} \bmod x \equiv 1$ (or both of course). Is there a simple way of proving this conjecture?

论坛问题帖 | has_answers | Obinna Okechukwu | 2014-01-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019504 | MSE-648404

Conjecture linking multiplicative order of 2 and semi-primes

Suppose we have a semi-prime $N=pq$, where $p \ne q$, and $p>2$, $q>2$ Let k be the multiplicative order of 2 mod N, then either $p^{2} \bmod k \equiv 1$ or $q^{2} \bmod k \equiv 1$ Is there a simple proof (disproof) of this fact? There are counterexamples (see answer).An interesting question is - Are there additional conditions that must hold for the conjecture to be true? This is somewhat related to a question I asked previously, which was simply a special case of this conjecture when $N = 2^{m}-1$, where the multiplicative order in this case is m. See Conjecture involving semi-prime numbers of the form $2^{x}-1$

论坛问题帖 | has_answers | Obinna Okechukwu | 2014-01-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019505 | MSE-648686

Proof envolving center of a group and conjugacy classes

I need to prove that every group G of order 9 is abelian. So, this is what I've done so far: There are three options for $[G:Z(G)]$: $[G:Z(G)]=1$, in this case we are done, since $G=Z(G)$ and that means G is abelian. $[G:Z(G)]=3$, in this case, we have that $G/Z(G)$ is cyclic, and that implies that G is abelian (I proved it earlier). $[G:Z(G)]=9$, since this means that $Z(G)=\left\{e\right\}$, we have that: $|G|=|Z(G)|+\sum_{x\notin Z(G)}|[x]|$ (while $[x]$ means the conjugacy class represented by x) and then $\sum_{x\notin Z(G)}|[x]|=8$, but I couldn't find the contradiction here. Any help would be appreciated.

论坛问题帖 | has_answers | so.very.tired | 2014-01-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019506 | MSE-657032

Are $ut + 1$ and $ut + t + 1$ both prime for some t for any u?

Conjecture : For any natural number u, there is a natural number t such that $ut + 1$ and $ut + t + 1$ are both prime. So we get a solution of the equation $$au - b(u+1) = -1$$ with prime numbers a and b by setting $a := ut + t + 1 , b := ut + 1$: $$(ut+t+1)u - (ut+1)(u+1) = u^2t+ut+u-u^2t-ut-u-1 = -1$$ Motivation : If the conjecture is true, then for any natural number n, there is a pair $(a,a+1)$ of consecutive squarefree numbers with exactly n distinct prime factors. I checked the conjecture with PARI and for $u\le10^6$, it is true. The largest number t necessary to produce the prime pair is 3420 for the number

829123 upto $u = 10^6$

论坛问题帖 | has_answers | Peter | 2014-01-30

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019507 | MSE-680559

All about a failed conjecture.

Some months ago I made the following conjecture - Let $d(n)$ denote the number of divisors of $n $. Then let N be a number such that $d(N)$ divides N . Also let $I= \frac{N}{d(N)}$ which is defined as the "Index of Beauty of N ". Then, For every number I there exists a number N such that I is the index of beauty of N. This conjecture was proved false by Greg Martin here. He said that it can be showed by exaustive computation that the following I fail the conjecture under 1000 are $\{18, 27, 30, 45, 63, 64, 72, 99, 105, 112, 117, 144, 153, 160, 162, 165, 171, 195, 207, 225, 243, 252, 255, 261, 279, 285, 288, 294, 320, 333, 336, 345, 352, 360, 369, 387, 396, 405, 416, 423, 435, 441, 465, 468, 477, 490, 504, 531, 544, 549, 555, 567, 576, 603, 608, 612, 615, 616, 625, 639, 645, 657, 684, 705, 711, 726, 728, 735, 736, 747, 792, 795, 801, 810, 828, 840, 873, 880, 885, 891, 909, 915, 927, 928, 936, 952, 960, 963, 981, 992\}$ Now what I am interested is that sequence of I that fails. (i) Is this sequence infinite?How? (ii)Is there any approximation which can tell the number of such failed I less than a fixed x (iii)If the sequence is infinite then are there canonical forms in which all of the values are in our list.

论坛问题帖 | no_answers_at_retrieval | Shivam Patel | 2014-02-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019508 | MSE-683711

Symmetry of bicycle-lock numbers

Suppose you have a combination bicycle lock of this sort: with n dials and k numbers on each dial. Let $m(n,k)$ denote the minimum number of turns that always suffice to open the lock from any starting position, where a turn consists of rotating any number of adjacent rings by one place. For example $m(2,10)=6$ and $m(3,10)=10$. I have found an efficient algorithm to compute these numbers, which reveals a symmetry I can’t currently explain: $m(n, k+1) = m(k, n+1)$ This is such a striking symmetry that I guess it has a simple explanation. Can anyone find one? Here’s the table of values for small n and k, exhibiting the conjectured symmetry: n\k| 2 3 4 5 6 7 8 9 10 ---+---------- 1 | 1 1 2 2 3 3 4 4 5 2 | 1 2 2 3 4 4 5 6 6 3 | 2 2 4 4 6 6 8 8 10 4 | 2 3 4 6 6 8 9 10 12 5 | 3 4 6 6 9 9 12 12 15 6 | 3 4 6 8 9 12 12 15 16 7 | 4 5 8 9 12 12 16 16 20 8 | 4 6 8 10 12 15 16 20 20 9 | 5 6 10 12 15 16 20 20 25 10 | 5 7 10 12 15 18 20 24 25

论坛问题帖 | has_answers | Robin Houston | 2014-02-20

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019509 | MSE-687370

Mertens conjecture - bounds

The disproven Mertens Conjecture states that $|M(n)|\leq \sqrt{n}$ If it is bounded at all, would the bounds $|M(n)|\leq \sqrt{2n\log(\log (n))}$ not be more realistic, and still consistent with the counterexamples suggested by Odlyzko and te Riele, owing to the law of the iterated logarithm?

论坛问题帖 | has_answers | martin | 2014-02-23

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019510 | MSE-691969

Closed form for $\int_{-1}^1\frac{\ln\left(2+x\,\sqrt3\right)}{\sqrt{1-x^2}\,\left(2+x\,\sqrt3\right)^n}dx$

I'm trying to find a closed form for the following integral: $$\mathcal{J}(n)=\int_{-1}^1\frac{\ln\left(2+x\,\sqrt3\right)}{\sqrt{1-x^2}\,\left(2+x\,\sqrt3\right)^n}dx\tag1$$ I have conjectured values of $\mathcal{J}(n)$ (supported by numerical inegration) for some integer values of n: $$\begin{align}&\mathcal{J}(1)\stackrel?=-\pi\ln\left(\frac32\right),\\&\mathcal{J}(2)\stackrel?=-\pi\left(1+2\ln\left(\frac32\right)\right),\\&\mathcal{J}(3)\stackrel?=-\pi\left(\frac{15}4+\frac{11}2\ln\left(\frac32\right)\right),\\&\mathcal{J}(4)\stackrel?=-\pi\left(\frac{77}6+17\ln\left(\frac32\right)\right).\end{align}\tag2$$ These values suggest that a general form for $n\in\mathbb{N}$ is $$\mathcal{J}(n)\stackrel?=-\pi\left(a_n+b_n\ln\left(\frac32\right)\right),\tag3$$ where a_n, b_n are some rational coefficients. Moreover, I conjecture that $$b_n\stackrel?={}_2F_1\left(1-n,n;\,1;\,-\frac12\right).\tag4$$ Are my conjectures true? Can we find a formula or recurrence relation for a_n? Can we find a general formula for $\mathcal{J}(z)$ for non-integer values of z?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2014-02-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019511 | MSE-704818

Conjecture about integral $\int_0^1 K\left(\sqrt{\vphantom1x}\right)\,K\left(\sqrt{1-x}\right)\,x^ndx$

I'm interested in the following integral: $$\mathcal J(n)=\int_0^1 K\left(\sqrt{\vphantom1x}\right)\,K\left(\sqrt{1-x}\right)\,x^ndx,\tag1$$ where $K(z)$ is the complete elliptic integral of the 1st kind: $$K(z)={}_2F_1\left(\begin{array}c\tfrac12,\tfrac12\\1\end{array}\middle|\, z^2\right)\cdot\frac\pi2.\tag2$$ Using numerical integration and inverse symbolic calculation techniques, I found a conjectural formula for this integral, that only though seems to work for $n\in\mathbb N_0$: $$\mathcal J(n)\stackrel?=\frac{\pi^3}{8\cdot4^n}\left(\frac{(2n-1)!!}{n!}\right)^2{}_4F_3\left(\begin{array}c\tfrac12,\tfrac12,-n,-n\\1,\tfrac12-n,\tfrac12-n\end{array}\middle|\, 1\right).\tag3$$ This formula produces the following sequence of results for $n=0,1,2,...$ $$\frac{\pi^3}8,\frac{\pi^3}{16},\frac{11\,\pi^3}{256},\frac{17\,\pi^3}{512},\frac{1787\,\pi^3}{65536},\frac{3047\,\pi^3}{131072},\frac{42631\,\pi^3}{2097152},\frac{75937\,\

pi^3}{4194304},...\tag4$$ Can we prove the conjecture (3) is correct? Is it possible to find a generalization that works for non-integer values of n as well?

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2014-03-09

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019512 | MSE-706488

Conjecture: When does $n=ab$, with $a\leq b\leq 2a$?

I conjecture that if this occurs, a and b are unique. Obviously if n is an odd prime, this does not occur, and if $n=a^2$, it does. In any case, what is the set of numbers such that this sort of factorization exists? For example, $510=17\cdot30$, and $17\leq 30\leq 34$. An equivalent formulation is that the set $S_n=\{(x,\frac nx)\mid\sqrt{\frac n2}\leq x\leq\sqrt{2n}\}$ has at most one integral point (ie., $|S_n\cap \mathbb Z\times\mathbb{Z}|\in\{0,1\}~\forall n$).

论坛问题帖 | has_answers | math.nb | 2014-03-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019513 | MSE-715479

Conjectured closed form of $G^{2~2}_{3~3}\left(1\middle|\begin{array}c1,1;b+1\\b,b;0\end{array}\right)$

In my answer to this question, I come across the following case of the Meijer G-function: $$F(b)=G^{2~2}_{3~3}\left(1\middle|\begin{array}c1,1;b+1\\b,b;0\end{array}\right), b>0$$ and based on my experiments, $F(b)$ have the following closed form: $$F(b) \stackrel?=\frac{\Gamma(b)}{b}\left(-\gamma-\psi(b)+\frac{2^{1-b}}{b}{_2F_1}\left(\begin{array}c1,1\\b+1\end{array}\middle|-1\right)+b{_3F_2}\left(\begin{array}c1,1,b+1\\2,2\end{array}\middle|-1\right)\right)$$ Is there any chance of proving this? Edit: Using the definition of the Meijer G-Function, $F(b)$ have the integral representation: $$F(b)=\frac{1}{2\pi}\int^{+\infty}_{-\infty}\frac{\Gamma(\tfrac{b}{2}+ix)\Gamma(\tfrac{b}{2}-ix)}{\tfrac{b^2}{4}+x^2}dx.$$ Edit 2: I've found a further generalization: $$F(b,z)=G^{2~2}_{3~3}\left(z\middle|\begin{array}c1,1;b+1\\b,b;0\end{array}\right)=\frac{1}{2\pi}\int^{+\infty}_{-\infty}\frac{\Gamma(\tfrac{b}{2}+ix)\Gamma(\tfrac{b}{2}-ix)}{\tfrac{b^2}{4}+x^2}e^{(b/2+ix)\log z}dx\\ \stackrel?=\frac{\Gamma(b)z^b}{b}\left(-\log z-\gamma-\psi(b)+\frac{(z+1)^{1-b}}{b}{_2F_1}\left(\begin{array}c1,1\\b+1\end{array}\middle|-z\right)+bz~{_3F_2}\left(\begin{array}c1,1,b+1\\2,2\end{array}\middle|-z\right)\right), b\not\in\{0,-1,-2,\dots\},z\neq0,-1.$$ Edit 3: Further simplified the ${_2F_1}$ part.

论坛问题帖 | has_answers | Chen Wang | 2014-03-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019514 | MSE-727965

Prove that $\lim\limits_{x\to\infty} \frac{\Gamma(x+1,x(1+\epsilon))}{x\Gamma(x)}=0$.

I conjecture that for any $\epsilon>0$, we have $$ \lim_{x\to\infty} \frac{\Gamma(x+1,x(1+\epsilon))}{x\Gamma(x)}=0 $$ where $\Gamma(x,a) = \int_a^\infty t^{x-1}e^{-t} \mathrm{d}t$ denotes the incomplete Gamma function, and $\Gamma(x) = \Gamma(x,0)$ is the Gamma function. I'm also happy with the weaker conjecture $$ \lim_{x\to\infty} \frac{\int_{x(1+\epsilon)}^\infty t^{x-1}\log(1+t)e^{-t} \mathrm{d}t}{x\Gamma(x)}=0 $$ but I doubt that it is any easier. Any clues? NB: the stronger conjecture is obtained from the weaker conjecture by upper-bounding $\log(1+t)$ by t

论坛问题帖 | has_answers | jens | 2014-03-26

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019515 | MSE-730871

Srinivasa Ramanujan conjectures

I searched internet for the whole list of conjectures by Srinivasa Ramanujan , but its not fruitful . I came to know that recently a book of Ramanujan was out and contains many conjectures related to summations . Please link the reference of Srinivasa Ramanujan conjectures . Thanks in advance :)

论坛问题帖 | has_answers | hanugm | 2014-03-29

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019516 | MSE-745018

I have a conjecture on local max/min , can any of you propose a contradiction?

If f is a non-piecewise function defined continuous on an interval I, and within that interval I, there exists a value x, such that $f`(x)$ (derivative of f) does not exist , then at that value x, is a local $\max/\min$ value.

论坛问题帖 | has_answers | Greg Dillon | 2014-04-08

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019517 | MSE-750469

Measure Theory Conjecture

While I was doing some math here, I made this conjecture. Let $f_n:X\rightarrow \mathbb{R}$ be a sequence of measurable functions from the measure space $(X,\mathcal{A},\mu)$ to the measurable space $(\mathbb{R},\mathcal{B}(\mathbb{R}))$, and let $\mu:\mathcal{A}\rightarrow[0,\infty]$ be a positive measure. Consider that for all $\varepsilon > 0$, the sets $A_n = \{x\in X: \ |f_n(x)| > \varepsilon\}$ are such that $$\limsup_{n\to\infty}\mu(A_n) = 0.$$ Then its true that $f_n\to0$, μ-almost everywhere. The thing is, I think it is true but I don't know how to prove it. You can consider μ as a probability measure if it makes more easier to prove. Thank you.

论坛问题帖 | has_answers | Integral | 2014-04-12

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019518 | MSE-758876

On extracting primes from coprimes

Proof or disprove the following statement - There exists infinitely many a and b which are pair of co-prime integers , either $ab+1$ or $ab-1$ is prime. Motivation- Looking at some twin prime pairs and calculating a lot - I guessed something like above.

论坛问题帖 | has_answers | Shivam Patel | 2014-04-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019519 | MSE-767114

Random conjecture

For any numbers a and b so that $a>b$ and $\text{gcd}(a, b)=1$, there exists a $c\in\mathbb{N^+}$ so that $a+bc$ is prime. I've only just tested this out with a few numbers, but I'm curious as to what you guys have to say. $$100+21\cdot3 \\97+96\cdot1 \\97+50\cdot2 \\482914+15435\cdot5 \\500009+500000\cdot4$$

论坛问题帖 | has_answers | user137794 | 2014-04-24

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019520 | MSE-77780

Infinitely many primes of the $\sum_{i=0}^{a} n^{i}+m\cdot\sum_{i=0}^{b} n^{i}$ form?

How to show that there is infinitely many prime numbers of the form: $p=\sum_{i=0}^{a} n^{i}+m\cdot\sum_{i=0}^{b} n^{i}$ where: $m\in \mathbb{Z}^{*}$, $a,b,n\in \mathbb{N}$, $\gcd(a+1,b+1)=1$ For example: $\sum_{i=0}^{6} 10^{335\cdot i}+8648\cdot \sum_{i=0}^{4} 10^{335\cdot i}$ is prime number.

论坛问题帖 | has_answers | Pedja | 2011-11-01

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019521 | MSE-78502

If $\gcd(a,b)=1$, and a is even and b is odd then $\gcd(2^{a}+1,2^{b}+1)=1$?

How to prove that: $\gcd(a,b)=1 \Rightarrow \gcd(2^{a}+1,2^{b}+1)=1$,where a is even and b is odd natural number For example: $\gcd(2^8+1,2^{13}+1)=1 , \gcd(2^{64}+1,2^{73}+1)=1$ I know that Knuth showed that: $\gcd(2^{a}-1,2^{b}-1)=2^{\gcd(a,b)}-1$ so: $\gcd(a,b)=1\Rightarrow \gcd(2^{a}-1,2^{b}-1)=1$ but I don't see whether this fact is useful.

论坛问题帖 | has_answers | Pedja | 2011-11-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019522 | MSE-78531

If a is even and b is odd then $\gcd(2^{a}+1,2^{b}+1)=1$

How to prove that: If a is even and b is odd then $\gcd(2^{a}+1,2^{b}+1)=1$ This statement is generalization of the statement from my previous question. I have checked for many (a,b) pairs using Maple, and $\gcd(2^{a}+1,2^{b}+1)$ is always 1 For example : $\gcd(2^{66}+1,2^{33}+1)=1 , \gcd(2^{4422}+1,2^{3333}+1)=1$ and so on.

论坛问题帖 | has_answers | Pedja | 2011-11-03

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019523 | MSE-788687

Prove that there exists an m such that for any $n>m$ there exists at least one prime between c_n and n

Let c_n be the n-th composite. Then the problem is to prove that- $\pi(c_n)-\pi(n)>0$ $\forall n>m$ I have tried to progress in the problem using an elementary approach. So far I have been able to prove the following trivial results- For any $n>7$ the composite c_n satisfies the inequality $2n>c_n>n$. $\pi(c_n)=c_n-n-1$ $\pi(x)\geq \pi(p)$ $\iff$ $x \geq p$. Where p is a prime. Notice that when the R.H.S always implies L.H.S. To prove the converse we note that $\pi(x) > \pi(p)$ $\implies$ $x>p$. The problem occurs when $\pi(x) = \pi(p)$. In that case if $x<p$ then it would imply that $\pi(x) < \pi(p)$, contradicting our hypothesis. Hence proved. The $\pi(c_n)$-th prime is strictly less then c_n for any $n>7$. $c_n$$_-$$_\pi$$_($$_n$$_)$$_-$$_1$ $=$ $n-1$ $\iff$ n is a prime and $c_n$$_-$$_\pi$$_($$_n$$_)$$_-$$_1$ $=$ n $\iff$ n is a composite. Using 5 it can be proved that c_n $\geq$ $n+\pi(n)+1$. But what I want is strict inequality. I have proved that the function $\Psi(n)=c_n-n-\pi(n)-1$ is increasing between two consecutive primes. Notice that the statement of the problem is equivalent to the following statement- Prove that the $\pi(c_n)$-th prime is strictly greater then n for any $n > m$. Actually I have borrowed the problem from the AoPS site. For the actual post see http://www.artofproblemsolving.com/Forum/viewtopic.php?f=52&t=588724.

论坛问题帖 | has_answers | William Hilbert | 2014-05-10

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019524 | MSE-793492

Pi and the sum of reciprocals of primes?

So I know that $$\sum_{\underset{\Large p\; prime}{p=1}}^{\infty}\frac{1}{p}$$ blows up. But doing some fun on mathematica I found out that when the sum isn't infinite, it was so close to 3 and I conjectured that there exists an ϵ such that $$\sum_{\underset{\Large p\; prime}{p=1}}^{\epsilon}\frac{1}{p}=\pi$$ Is this conjecture true or false and more importantly why?

论坛问题帖 | has_answers | user149842 | 2014-05-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019525 | MSE-800048

Tangent at average of two roots of cubic with one real and two complex roots

I was able to easily prove that the tangent at the average of two roots of a real cubic polynomial passed through the third root of the function. But I have only done this for functions with three distinct real roots using $y = (x-a) (x-b) (x-c)$ as the function. I calculated the derivative of the function in terms of (x= midpoint of a and b) and then just subbed in ($x = c$) in the equation of the tangent that I worked out to prove this conjecture. BUT!! Now, I am trying to prove the same for cubics with one real and two complex roots. I am using $y = (x -d) (x-(a+bi) (x- (a-bi))$ where d is the one real root. So far, I have found the derivative of this to be \begin{align} dy/dx &= 2(x-d)(x-a) + (x-a-bi)(x-a+bi)\\ &= 2(x-d)(x-a) + x^2 + a^2 - bi^2\\ &= 2(x-a) (x-d) (x^2 + a^2 + b^2) \end{align} The average of the two roots I have used is $((a+bi/2) + (a - bi/2))/2 = a$ I have already proved it for this average of roots! But!! I also tried the other possible average of roots $((d + a +bi)/2)$ where the Point of contact I used is $((a+bi+d)/2), -((a+bi-d)/2)^3$ Why is that the equation of the tangent I worked out for this second possible pair of roots using ($m =ystep/xstep$) is not equalling zero when I sub in the third root into the equation of tangent I am getting?? Maybe I have calculated some things wrong? Can you please help me prove the conjecture? Plus, another mini question: Can cubic polynomials be non-real? because when we say all cubic polynomials must have at least one real root, it implies that the cubic is real. So can I say that a cubic with an imaginary coefficient is non-real? I won't try to prove this conjecture for non-real cubics though. Can you please help me?! I would really appreciate even a bit of help Sorry and Thanks!! =)

论坛问题帖 | has_answers | angel | 2014-05-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019526 | MSE-821682

Prove the inequality for composite numbers

Is it true that c_m+c_n $>$ c_{m+n} for all m, n $\in$ $\mathbb{N}$? Though the result seems true, I can't get a solution. Even the bounds on c_n obtained from Prime Number Theorem isn't helping me. Is there any way to prove it?

论坛问题帖 | no_answers_at_retrieval | William Hilbert | 2014-06-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019527 | MSE-824054

Equal perimeters of squares and right angled isosceles triangles

Consider a square ABCD having length l and breadth. Now start folding the sides AB and AC so that the figure becomes something like this $$$ All the vertical and horizontal folds/stairs are equal in length. The perimeter of the figure is equal to the the perimeter of the square. As we increase the number of divisions, the length of each fold/stair decreases. Let the number of stairs/folds be n. As $n\rightarrow\infty$ the figure becomes a right angled isosceles triangle BCD. The perimeter of triangle BCD should be equal to the perimeter of square ABCD since as we increase the number of folds/stairs the perimeter remains the same. Can anyone please correct me where I have gone wrong?

论坛问题帖 | has_answers | Akshit | 2014-06-07

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019528 | MSE-830699

Decidability of $P = NP$?

(Please, don't sign this as duplicate of this question, they are not.) Is it possible, that the well-known $P=NP$ conjecture is undecidable in ZFC? Is there any result about this topic?

论坛问题帖 | no_answers_at_retrieval | peterh | 2014-06-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019529 | MSE-830854

Non-repeating decimals in 1/n

Here's a conjecture I made: The number of non repeating decimal places in the base-ten representation of the fraction 1/n, where n is an integer, is equal to whichever is higher: the exponent of 2 in the prime factorization of n or the exponent of 5 in the prime factorization of n. Is this true? If so, how can I prove it?

论坛问题帖 | no_answers_at_retrieval | user156582 | 2014-06-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019530 | MSE-832770

How to prove this??

Does the following inequality hold? $p(n)\leq 2^n,$ where $p(n)$ is the nth prime. If this is true then it follows that: If $p(n)=p(m)^x+p(o)^y$, then $\max[x,y] \le n$.

论坛问题帖 | no_answers_at_retrieval | user96172 | 2014-06-13

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019531 | MSE-834001

Find all n so that c_n $>$ $\pi(n^2)$

Find all n $\in$ $\mathbb{N}$ so that p_{c_n} $>$ n^2 where p_n denotes the n-th prime and c_n, the n-th composite. I have tried doing the problem using The stronger version of Roger's Theorem, i.e., by the inequality, $$\ln n + \ln \ln n > \dfrac {p_n}{n} > \ln n + \ln \ln n - 1$$ but in vain. I think that the problem is true for all sufficiently large values of n. Even a proof of only this claim will also be appreciated.

论坛问题帖 | has_answers | William Hilbert | 2014-06-14

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019532 | MSE-836945

Low Level Books on Conjectures/Famous Problems

I am currently an undergraduate math/CS major with coursework done in Linear Algebra, Vector Calculus (that covered a significant amount of Real 1 material), Discrete Math, and about to take courses in Algebra and Real Analysis 1. I was wondering if there are any books about famous problems (Riemann, Goldbach, Collatz, etc) and progress thus far that are accessible to me. If possible, I would prefer them to have more emphasis on the math rather than the history. As an example, I enjoyed reading Fermat's Enigma by Simon Singh but was disappointed in how it glossed over most of the math. If there are any suggestions I would greatly appreciate it.

论坛问题帖 | has_answers | Christopher Liu | 2014-06-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019533 | MSE-838678

Intuitively, what separates Mersenne primes from Fermat primes?

A Mersenne prime is a prime of the form 2^n-1. A Fermat prime is a prime of the form 2^n+1. Despite the two being superficially very similar, it is conjectured that there are infinitely many Mersenne primes but only finitely many Fermat primes. Is there an intuition that can help me appreciate the nature of that seemingly paradoxical difference?

论坛问题帖 | has_answers | user139000 | 2014-06-18

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019534 | MSE-888402

conjecture about prime numbers and distance between them

is there a name for this conjecture? Conjecture: given p_n a prime number sequence where $p_1=2,p_2=3,\cdots$, then for all $n\in\mathbb{N}^*$ and $k\in\mathbb{N}$, holds that $\displaystyle p_{n+2+k}-p_{n+1}\le\sum_{i=0}^{k}p_{n+i}$

论坛问题帖 | no_answers_at_retrieval | cand | 2014-08-05

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019535 | MSE-8921

A conjecture of parallelogram inside convex and central symmetric curve

Assume Q is a convex central symmetric curve, whose area is $\displaystyle S$. The area of the maximum parallelogram inside Q is $\displaystyle S'$. How to prove the conjecture that $\displaystyle \frac{S'}{S} \ge \frac{2}{\pi}=0.6366\dots$? For example, If Q is an ellipse, $\displaystyle S'=2ab$, $\displaystyle S=\pi ab$. If Q is a regular hexagon, $\displaystyle \frac{S'}{S}= \frac{2}{3}$. It's trivial that $\displaystyle \frac{S'}{S} \ge \frac{1}{2}$, and I know how to prove $\displaystyle \frac{S'}{S} \ge \frac{4}{4+\pi}=0.56\dots$ From many reason, I believe this conjecture is true. Denote MAP="Maximum Area Parallelogram": For any Q and any direction θ, let $P(Q,\theta)$ be the area of MAP which have a corner in this direction. $S'=\max\{P(Q,\theta)\}$. In order to make $\frac{S'}{S}$ smallest, We need keep the largest one of $\{P(Q,\theta)\}$ small while S is a constant. Ellipse just keeps everyone in $\{P(Q,\theta)\}$ average. This is very special, I don't think there will be other curve having this property. On the other hand, distribute equally always lead to the min-max in our knowledge. About the $\frac{4}{4+\pi}$ lowerbound, the idea is as follows: First, use polar function $r(\theta)$ to describe the curve. The condition is that $r(a)r(b)\sin|a-b|\leq C$, and we want to bound is $S=\int_{\theta}{r(\theta)}^2$. Second, Without lose of generality, We assume $r(0)=r(90)=1,C=1$, and assume Q is in the boundary of $Z=\{(x,y)|-1\leq x,y\leq 1\}$. Third, let $a=r(\theta)$ and $b=r(\theta+90)$ and find a bound for (a^2+b^2) by Cauchy-Inequality. and it will give a bound for the area S.

论坛问题帖 | has_answers | galois | 2010-11-04

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 2.5

V3-019536 | MSE-897967

Trigamma identity $4\,\psi_1\!\left(\frac15\right)+\psi_1\!\left(\frac25\right)-\psi_1\!\left(\frac{10}\right)=\frac{4\pi^2}{\phi\,\sqrt5}.$

I heuristically discovered the following identity for the trigamma function, that I could not find in any tables or papers or infer from existing formulae (e.g. [1], [2], [3], [4], [5], [6]): $$4\,\psi_1\!\left(\frac15\right)+\psi_1\!\left(\frac25\right)-\psi_1\!\left(\frac1{10}\right)=\frac{4\pi^2}{\phi\,\sqrt5}.\tag1$$ It also seems to be unknown to Mathematica, but numerically checks with at least 20000 decimal digits. It might be provable through some application of reflection and multiplication theorems, but I couldn't do this. Please suggest how to prove it. Update: Another identity is $$3\,\psi_1\!\left(\frac1{12}\right)-30\,\psi_1\!\left(\frac13\right)=120\,G+\left(6\sqrt3-8\right)\pi^2.\tag2$$

论坛问题帖 | has_answers | Vladimir Reshetnikov | 2014-08-15

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019537 | MSE-90538

Interesting Property of Numbers in English

I was playing with the letters in numbers written in English and I found something quite funny. I found that if you count the number of letters in the number and write this as a number and then count the number of letters in this new number and keep repeating the process, you will arrive at the number 4. I've confirmed this (using a computer program) for all numbers up to 999999 and was wondering if there's a way to prove this or to find a counter example for which it does not hold. Just to give an example of the above statement, let's start with thirty seven (I chose this randomly) Thirty seven has 11 letters in it, Eleven has 6 letters in it, Six has three letters in it, Three has 5 letters in it, Five has 4 letters in it. It may look like I just picked this number, so let me show this for another random number, say 999. Nine hundred and ninety nine has 24 letters in it, Twenty four has 10 letters in it, Ten has 3 letters in it, Three has 5 letters in it, Five has 4 letters in it. What are your thoughts on how to prove this? (Just a note: I only confirmed this for numbers written in the standard British way of writing numbers - for example 101 is one hundred and one)

论坛问题帖 | has_answers | Obinna Okechukwu | 2011-12-11

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019538 | MSE-908651

A Conjecture on The Generalization of Quadratic Reciprocity Law

Is there any way to prove the following conjecture regarding the Generalization of Quadratic Reciprocity Law. The statement being, $$ \left(\dfrac{a_1}{a_2}\right)\left(\dfrac{a_2}{a_3}\right) \ldots\left(\dfrac{a_n}{a_1}\right)=(-1)^{\left(\frac{a_1-1}{2}\right) \left(\frac{a_2-1}{2}\right) \ldots \left(\frac{a_n-1}{2}\right)}$$ where the symbols are Jacobi Symbols and $\operatorname{gcd}(a_i,a_j)=1$ for $i\neq j$ and $a_i>1$ forall i satisfying $1 \leq i \leq n$. One of my friend told me that the argument for $n=3$ and when all a_i are distinct primes, is a modification of the Eisenstein's argument for proving the Quadratic Reciprocity Law itself. Instead of considering a planer rectangle as in Eisenstein's Proof of Quadratic Reciprocity Law if we consider a cuboid and count the number of points (x,y,z) (where x,y and z are all integers) that is within this cuboid and not on any outer surface, then the proof would be done. And when the result is proved for primes it can be proved in general also. However the problem is that in this scheme we can't visualize an n-dimensional rectangle and thus can't forward his argument to n dimensions, though we think that it can be generalized. Is there any way to help visualizing it or is there any way to prove the result in a simpler way?

论坛问题帖 | has_answers | user170039 | 2014-08-25

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019539 | MSE-935459

At most n functions satisfy $f(x)=\bigl(x-f(a_1)\bigr)\bigl(x-f(a_2)\bigr)\dots\bigl(x-f(a_n)\bigr)$?

Some background: I was trying to solve the functional equation $f\bigl(f(x)\bigr)=\sin x$. I realized that $f(\pi n)$ is a root of f for all integers n, because $f\bigl(f(\pi n)\bigr)=\sin(\pi n)=0$. Thus, we can write f as $f(x)=A\bigl(x-f(0)\bigr)\bigl(x-f(\pi)\bigr)\bigl(x-f(-\pi)\bigr)\dots$. The problem now is to find all functions f that solve this. I couldn't even begin on this, so I decided to try letting the constant A equal 1, and to try some simpler versions first. E.g. For $f(x)=x-f(0)$ we find the unique solution $f(x)=x$; for $f(x)=\bigl(x-f(0)\bigr)\Bigl(x-f\left(\sqrt{2}\right)\Bigr)$ we find two solutions, namely $f(x)\in \left\{x^2-1,x^2-2\left(\sqrt{2}-1\right)x\right\}$. I solved many more examples, and then conjectured that: Let $a_i\in \mathbb{C}$ for $i=1,2,\dots,n$. Prove that the functional equation $f(x)=\bigl(x-f(a_1)\bigr)\bigl(x-f(a_2)\bigr)\dots\bigl(x-f(a_n)\bigr)$ has exactly n distinct solutions. Finally, I realized that if this is proved, then it would "follow" (I don't know if this is rigorous but I hope someone knows how to make it rigorous) that our original functional equation $f(x)=\bigl(x-f(0)\bigr)\bigl(x-f(\pi)\bigr)\bigl(x-f(-\pi)\bigr)\dots$ (I put $A=1$) has infinitely many solutions. Thanks for reading.

论坛问题帖 | has_answers | Is Ne | 2014-09-17

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 4.0

V3-019540 | MSE-937964

Can decimals/fractions be odd or even?

At school I asked the question and I kept wondering "Can fractions or decimals be odd or even?"

论坛问题帖 | has_answers | Shaelynn McCabe | 2014-09-19

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA; version depends on revision date

V3-019541 | MSE-948616

A variation of Fermat's little theorem in the form $a^{n-d}\equiv a$ (mod p).

Fermat's little theorem states that for n prime, $$ a^n \equiv a \pmod{n}. $$ The values of n for which this holds are the primes and the Carmichael numbers. If we modify the congruence slightly, $$ a^{n - 1} \equiv a \pmod{n}, $$ the values of n for which it holds are listed in A216090: 1, 2, 6, 10, 14, 22, 26, 30, 34, 38, 46, ... More generally, we can examine the values of n where $$ a^{n - d} \equiv a \pmod{n} $$ for all a and a given d. $d = 0$ gives the primes/Carmichael numbers, $d = 1$ gives the sequence A216090, and $d = 2$ gives the sequence 1, 2, 3, 15, 21, 33, 39, 51, 57, 69, 87, ... For $d = 3$, the sequence grows much more rapidly: 1, 2, 4, 6, 154, 4774, 23254, 179014, 187054, 1066054, 1168654, ... But for $d = 4$, the growth slows down again: 1, 2, 3, 5, 65, 85, 145, 165, 185, 205, 265, ... The first few d values which give fast-growing sequences for n are 3, 7, 8, 11, 15, 17, 19, 23, 24, and 26. In general, the sequence seems to grow particularly quickly when $d + 1$ is not squarefree. I've proven this generalization of Korselt's criterion that characterises the possible n values for a given d: Theorem (Generalized Korselt's criterion). Let d be an arbitrary integer, and n a positive integer such that and $n \ge d + 1$. Then $a^{n - d} \equiv a \pmod{n}$ for all $a \in \mathbb{Z}$ if and only if either $n = 1$ or 2; $n = d + 1$; or n is squarefree, and for each of n's prime divisors p_i, $(p_i - 1) \vert (n - d - 1)$. Additionally, I've observed the following pattern that would explain why the squarefreeness of $d + 1$ matters: Conjecture. When $d + 1$ is squarefree, many prime multiples of $d + 1$ are valid values for n. But if $d + 1$ is not squarefree, no multiples of it (except possibly itself) are feasible values for n. In either case, valid n values which are not multiples of $d + 1$ are much less abundant. Any ideas how to tackle this conjecture, or at least make it more explicit?

论坛问题帖 | has_answers | Tavian Barnes | 2014-09-27

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

V3-019542 | MSE-995589

A conjecture concerning primes and algebra

A monoid morphism $\psi:\mathbb Z_+\!\!\rightarrow\mathbb Z_+$ is defined by an arbitrary function $f:\mathbb Z_+\!\!\rightarrow\mathbb Z_+$ and defines a group homomorphism $\varphi:\mathbb Q_+\!\!\rightarrow\mathbb Q_+$: $\psi(a\cdot b)=\psi(a)\cdot\psi(b)\Rightarrow\psi(\prod p_i^{n_i})=\prod\psi(p_i)^{n_i}$, so any function $i\mapsto \psi(p_i)$ defines ψ. Further, since $\varphi(\frac{1}{n})=\frac{1}{\varphi(n)}$ a function f uniquely defines the group

homomorphism φ. For fun I intend to study the group homomorphism generated by the identity function $i\mapsto\omega(p_i)=i$. Then $\omega:\mathbb Z_+\!\!\rightarrow\mathbb Z_+$ seems to be an algebraic analogue to the analytic approach with $\pi(N)$ since also ω is a left inverse to the prime number function p_n. In the diagram ω is the blue curve and π the red and one can see (especially if the picture is loaded and viewed in a larger scale) that $\omega(1)>\pi(1),\; \omega(35)>\pi(35)$ and $\omega(49)>\pi(49)$. Of course ω and π coincide for all primes, but also for other values as $91,\;95$ and 133. However, I have tested all values less than 2^{16} and $\omega(N)\leq\pi(N)$ for $N>49$ in this interval. I have no idea how to prove the conjecture (if possible) but I will try to construct a program that (in principle) can test any interval of integers. Edit: I realize now that it is virtually impossible to make such a program, but with help of arrays of primes my program have tested all values $< 1\,000\,000$ and the only numbers for which $\omega(N)\geq\pi(N)$ (italic numbers stands for '$>$') are: $\mathit{1},\,9,\,15,\,21,\,25,\,\mathit{35},\,39,\,\mathit{49},\,57,\,65,\,91,\,95,\,133$. Conjecture: For $N=\prod p_k^{n_k}>49$ it holds that $\prod k^{n_k}\leq\pi(N)$. Thoughtful: from an analytic point of view ω seems to be very irregular, but algebraically it is an almost canonical morphism.

论坛问题帖 | has_answers | Lehs | 2014-10-28

问题帖文字；图像和回答未重印；有回答不等于猜想已解决 | 原始来源 | CC BY-SA 3.0

C03 OpenConjecture / arXiv | 394 条

V3-019543 | OC-10

A lift of the colored Jones polynomial of a knot | tqft.knots.habiro.25.tex:728-745

定位：tqft.knots.habiro.25.tex:728-745。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Stavros Garoufalidis; Campbell Wheeler | 2026-03-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019544 | OC-1015

Lattice points arising from regularity and v-number of Graphs: Whisker and Cameron-Walker | Arxiv.tex:980-983

定位：Arxiv.tex:980-983。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Prativa Biswas; Mousumi Mandal; Kamalesh Saha | 2026-03-05

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019545 | OC-1016

Stability conditions on noncommutative crepant resolutions of 3-dimensional isolated singularities | modi_stab_arxiv.v1.tex:578-580

定位：modi_stab_arxiv.v1.tex:578-580。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Wahei Hara; Yuki Hirano | 2026-03-05

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019546 | OC-1057

Coarse separation and splittings in right-angled Artin groups | CoarseSepRAAGs.tex:238-240

定位：CoarseSepRAAGs.tex:238-240。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Bensaid, Oussama; Genevois, Anthony; Tessera, Romain | 2026-03-25

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019547 | OC-1077

Stein degree of proper morphisms | 2026-iccm-talk-birkar-2.tex:589-599

定位：2026-iccm-talk-birkar-2.tex:589-599。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Birkar, Caucher | 2026-03-25

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019548 | OC-1084

The Perimeter Winternitz Theorem in a Triangle | PerWintThmTriangle.tex:381-383

定位：PerWintThmTriangle.tex:381-383。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Berele, Allan; Catoiu, Stefan | 2026-03-24

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019549 | OC-1085

Absolute values and tensor powers of irreducible characters | char-ratio_4.tex:300-302

定位：char-ratio_4.tex:300-302。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Kushkuley, Alexander | 2026-03-24

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019550 | OC-110

Invertible Calabi-Yau Orbifolds over Finite Fields II | main__1_.tex:239-245

定位：main__1_.tex:239-245。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Marco Aldi; Andrija Peruničić | 2026-02-26

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019551 | OC-1114

Linear spaces in Hessian loci of cubic hypersurfaces | LinearHessian.tex:1250-1253

定位：LinearHessian.tex:1250-1253。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Bricalli, Davide | 2026-03-24

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019552 | OC-1145

A strengthened (∞, n)-categorical pasting theorem | main.tex:408-410

定位：main.tex:408-410。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chanavat, Clémence | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019553 | OC-1153

Towards sparse optimization over convex loops: Equivalence of Square Root Velocity distance and Wasserstein-Fisher-Rao | main.tex:1288-1292

定位：main.tex:1288-1292。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Cristinelli, Giacomo; Iglesias, José A. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019554 | OC-1157

Moment inequalities for Chow polynomials of matroids and bounds on Chern numbers | main.tex:903-910

定位：main.tex:903-910。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Cheng, Ronnie; Lin, Wangyang | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019555 | OC-1158

Moment inequalities for Chow polynomials of matroids and bounds on Chern numbers | main.tex:972-974

定位：main.tex:972-974。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Cheng, Ronnie; Lin, Wangyang | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019556 | OC-1159

Individual-based stochastic model with unbounded growth, birth and death rates: a tightness result | BRODU_IBM.tex:771-774

定位：BRODU_IBM.tex:771-774。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Brodu, Virgile | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_sa

V3-019557 | OC-1161

The non-peripheral curve graph and divergence in big mapping class groups | divergence.tex:540-543

定位：divergence.tex:540-543。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Bar-Natan, Assaf; Qing, Yulan; Rafi, Kasra | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019558 | OC-1173

Asymptotic Geometry of Four-Dimensional Steady Solitons | 4d_steady_arxiv.tex:164-167

定位：4d_steady_arxiv.tex:164-167。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Hebbar, Aprameya Girish; Sesum, Natasa | 2026-03-22

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019559 | OC-1176

Generalizing Saito's Criterion for Nonfree Arrangements | main.tex:652-658

定位：main.tex:652-658。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chu, Junyan | 2026-03-22

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019560 | OC-1177

Generalizing Saito's Criterion for Nonfree Arrangements | main.tex:661-666

定位：main.tex:661-666。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chu, Junyan | 2026-03-22

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019561 | OC-1178

Generalizing Saito's Criterion for Nonfree Arrangements | main.tex:671-677

定位：main.tex:671-677。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chu, Junyan | 2026-03-22

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019562 | OC-1179

Generalizing Saito's Criterion for Nonfree Arrangements | main.tex:680-682

定位：main.tex:680-682。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chu, Junyan | 2026-03-22

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019563 | OC-1180

The Erd\H{o}s-Ginzburg-Ziv theorem constant of finite groups | EGZTheoremConstant.tex:95-97

定位：EGZTheoremConstant.tex:95-97。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhao, Yang; Wang, Guoqing | 2026-03-22

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: missing

V3-019564 | OC-1191

The first eigenvalue of embedded minimal hypersurfaces in the unit sphere | main.tex:148-150

定位：main.tex:148-150。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhao, Yuhang | 2026-03-21

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019565 | OC-1192

The first eigenvalue of embedded minimal hypersurfaces in the unit sphere | main.tex:182-184

定位：main.tex:182-184。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhao, Yuhang | 2026-03-21

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019566 | OC-1207

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:3516-3537

定位：main_holography.tex:3516-3537。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019567 | OC-1208

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:3729-3779

定位：main_holography.tex:3729-3779。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019568 | OC-1209

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:3791-3799

定位：main_holography.tex:3791-3799。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019569 | OC-1210

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:3801-3809

定位：main_holography.tex:3801-3809。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019570 | OC-1211

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:4130-4140

定位：main_holography.tex:4130-4140。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019571 | OC-1212

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:4182-4189

定位：main_holography.tex:4182-4189。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019572 | OC-1213

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:4195-4216

定位：main_holography.tex:4195-4216。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019573 | OC-1214

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:4239-4252

定位：main_holography.tex:4239-4252。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019574 | OC-1215

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:4254-4259

定位：main_holography.tex:4254-4259。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019575 | OC-1216

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:4543-4557

定位：main_holography.tex:4543-4557。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019576 | OC-1217

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:4838-4982

定位：main_holography.tex:4838-4982。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019577 | OC-1218

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:4985-5040

定位：main_holography.tex:4985-5040。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019578 | OC-1219

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:5098-5117

定位：main_holography.tex:5098-5117。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019579 | OC-1220

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:5236-5239

定位：main_holography.tex:5236-5239。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019580 | OC-1221

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:5650-5686

定位：main_holography.tex:5650-5686。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019581 | OC-1222

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:5688-5711

定位：main_holography.tex:5688-5711。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019582 | OC-1223

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:5724-5747

定位：main_holography.tex:5724-5747。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019583 | OC-1224

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:6001-6007

定位：main_holography.tex:6001-6007。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019584 | OC-1225

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:6016-6036

定位：main_holography.tex:6016-6036。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019585 | OC-1226

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:6083-6089

定位：main_holography.tex:6083-6089。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019586 | OC-1227

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:6095-6108

定位：main_holography.tex:6095-6108。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019587 | OC-1228

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:6111-6136

定位：main_holography.tex:6111-6136。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019588 | OC-1229

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:6270-6340

定位：main_holography.tex:6270-6340。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019589 | OC-1230

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:6456-6465

定位：main_holography.tex:6456-6465。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019590 | OC-1231

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:6819-6868

定位：main_holography.tex:6819-6868。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019591 | OC-1232

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:6989-7043

定位：main_holography.tex:6989-7043。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019592 | OC-1233

CayleyPy-4: AI-Holography. Towards analogs of holographic string dualities for AI tasks | main_holography.tex:7152-7329

定位：main_holography.tex:7152-7329。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chervov, A.; Levkovich-Maslyuk, F.; Smolensky, A.; Khafizov, F.; Kiselev, I.; Melnikov, D.; Koltsov, I.; Kudashev, S.; Shiltsov, D.; Obozov, M.; Krymskii, S.; Kirova, V.; Konstantinova, E. V.; Soibelman, A.; Galkin, S.; Grunwald, L.; Kotov, A.; Alexandrov, A.; Lytkin, S.; Fedoriaka, D.; Chevychelov, A.; Kogan, Z.; Natyrova, A.; Cheldieva, L.; Nikitina, O.; Fironov, S.; Vakhrushev, A.; Lukyanenko, A.; Ilin, V.; Gorodkov, D.; Bogachev, N.; Gaiur, I.; Zaitsev, M.; Petrov, F.; Petrov, L.; Gaintseva, T.; Gavrilova, A.; Smirnov, M. N.; Kalinin, N.; Khan, A.; Jung, K.; Mousset, H.; Isambert, H.; Debeaupuis, O. | 2026-03-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019593 | OC-1249

Typical Weak Mixing and Exceptional Spectral Properties for Interval Translation Mappings | WeakMixing.tex:207-210

定位：WeakMixing.tex:207-210。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Artigiani, Mauro; Avila, Artur; Ferenczi, Sébastien; Hubert, Pascal; Skripchenko, Alexandra | 2026-03-19

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019594 | OC-126

A uniform version of a theorem by Lindström | lind.tex:230-232

定位：lind.tex:230-232。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Gábor Hegedüs | 2026-02-26

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019595 | OC-1265

Novel Self-similar Finite-time Blowups with Singular Profiles of the 1D Hou-Luo Model and the 2D Boussinesq Equations: A Numerical Investigation | Self-similar_Blowups_with_Singular_Profiles_of_Hou--Luo_Model_and_Boussinesq_equations.tex:723-725

定位：Self-similar_Blowups_with_Singular_Profiles_of_Hou--Luo_Model_and_Boussinesq_equations.tex:723-725。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Bojin Chen; De Huang; Xiangyuan Li | 2026-04-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019596 | OC-1271

Edge localization and Lifshitz tails for graphs with Ahlfors regular volume growth | arXiv.tex:492-494

定位：arXiv.tex:492-494。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Laura Shou; Wei Wang; Shiwen Zhang | 2026-04-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019597 | OC-1280

On the Construction of Recursively Differentiable Quasigroups and an Example of a Recursive $[4,2,3]_{26}$-Code | main_eng.tex:130-132

定位：main_eng.tex:130-132。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Petr Klimov | 2026-04-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019598 | OC-1283

Edge-Isoperimetric Inequalities in Chamber Graphs of Hyperplane Arrangements | isoperimetric2.tex:1212-1227

定位：isoperimetric2.tex:1212-1227。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Tilen Marc | 2026-04-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019599 | OC-1293

Existence of free boundaries for overdetermined value problems: Sharp conditions, regularity, and physical applications | AppFB2026.tex:1072-1074

定位：AppFB2026.tex:1072-1074。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Mohammed Barkatou; Samira Khatmi | 2026-04-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: missing

V3-019600 | OC-13

Duflo-Serganova functors and Brundan-Goodwin's parabolic inductions | main.tex:265-279

定位：main.tex:265-279。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Shunsuke Hirota | 2026-03-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019601 | OC-1302

Applications of renormalisation to orthonormal Strichartz estimates and the NLS system on the circle | Arxiv_1.tex:605-608

定位：Arxiv_1.tex:605-608。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Sonae Hadama; Andrew Rout | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019602 | OC-1310

The relativistic p-adic sunscreen conjecture | p-adic-sunscreen.tex:202-206

定位：p-adic-sunscreen.tex:202-206。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Sean Howe | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019603 | OC-1312

Various conjectural series identities | P91h.tex:223-233

定位：P91h.tex:223-233。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019604 | OC-1313

Various conjectural series identities | P91h.tex:241-256

定位：P91h.tex:241-256。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019605 | OC-1314

Various conjectural series identities | P91h.tex:260-280

定位：P91h.tex:260-280。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019606 | OC-1315

Various conjectural series identities | P91h.tex:284-304

定位：P91h.tex:284-304。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019607 | OC-1316

Various conjectural series identities | P91h.tex:310-321

定位：P91h.tex:310-321。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019608 | OC-1317

Various conjectural series identities | P91h.tex:323-334

定位：P91h.tex:323-334。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019609 | OC-1318

Various conjectural series identities | P91h.tex:336-346

定位：P91h.tex:336-346。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019610 | OC-1319

Various conjectural series identities | P91h.tex:349-359

定位：P91h.tex:349-359。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019611 | OC-1320

Various conjectural series identities | P91h.tex:361-373

定位：P91h.tex:361-373。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019612 | OC-1321

Various conjectural series identities | P91h.tex:375-388

定位：P91h.tex:375-388。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019613 | OC-1322

Various conjectural series identities | P91h.tex:390-400

定位：P91h.tex:390-400。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019614 | OC-1323

Various conjectural series identities | P91h.tex:402-406

定位：P91h.tex:402-406。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019615 | OC-1324

Various conjectural series identities | P91h.tex:418-426

定位：P91h.tex:418-426。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019616 | OC-1325

Various conjectural series identities | P91h.tex:431-439

定位：P91h.tex:431-439。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019617 | OC-1326

Various conjectural series identities | P91h.tex:444-452

定位：P91h.tex:444-452。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019618 | OC-1327

Various conjectural series identities | P91h.tex:457-467

定位：P91h.tex:457-467。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019619 | OC-1328

Various conjectural series identities | P91h.tex:472-477

定位：P91h.tex:472-477。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019620 | OC-1329

Various conjectural series identities | P91h.tex:482-488

定位：P91h.tex:482-488。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019621 | OC-1330

Various conjectural series identities | P91h.tex:493-499

定位：P91h.tex:493-499。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019622 | OC-1331

Various conjectural series identities | P91h.tex:504-508

定位：P91h.tex:504-508。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019623 | OC-1332

Various conjectural series identities | P91h.tex:513-517

定位：P91h.tex:513-517。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019624 | OC-1333

Various conjectural series identities | P91h.tex:523-527

定位：P91h.tex:523-527。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019625 | OC-1334

Various conjectural series identities | P91h.tex:535-541

定位：P91h.tex:535-541。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019626 | OC-1335

Various conjectural series identities | P91h.tex:560-567

定位：P91h.tex:560-567。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019627 | OC-1336

Various conjectural series identities | P91h.tex:572-587

定位：P91h.tex:572-587。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019628 | OC-1337

Various conjectural series identities | P91h.tex:593-600

定位：P91h.tex:593-600。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019629 | OC-1338

Various conjectural series identities | P91h.tex:606-614

定位：P91h.tex:606-614。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019630 | OC-1339

Various conjectural series identities | P91h.tex:623-636

定位：P91h.tex:623-636。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019631 | OC-1340

Various conjectural series identities | P91h.tex:644-658

定位：P91h.tex:644-658。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019632 | OC-1341

Various conjectural series identities | P91h.tex:663-670

定位：P91h.tex:663-670。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019633 | OC-1342

Various conjectural series identities | P91h.tex:677-686

定位：P91h.tex:677-686。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019634 | OC-1343

Various conjectural series identities | P91h.tex:694-708

定位：P91h.tex:694-708。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019635 | OC-1344

Various conjectural series identities | P91h.tex:714-721

定位：P91h.tex:714-721。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019636 | OC-1345

Various conjectural series identities | P91h.tex:728-740

定位：P91h.tex:728-740。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019637 | OC-1346

Various conjectural series identities | P91h.tex:749-763

定位：P91h.tex:749-763。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019638 | OC-1347

Various conjectural series identities | P91h.tex:771-789

定位：P91h.tex:771-789。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhi-Wei Sun | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019639 | OC-1352

Analytic rank-one elliptic curves over function fields and their rank over certain ring class fields | larsen_conjecture_function_field_arxiv.tex:158-160

定位：larsen_conjecture_function_field_arxiv.tex:158-160。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Seokhyun Choi; Bo-Hae Im; Beomho Kim | 2026-03-31

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019640 | OC-143

Quadratic Equations over Graph Products and the Exponent of Periodicity | paper.tex:879-881

定位：paper.tex:879-881。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Volker Diekert; Silas Natterer; Alexander Thumm | 2026-02-25

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019641 | OC-1435

Galois representation of the product of two Drinfeld modules of generic characteristic | newmain.tex:211-218

定位：newmain.tex:211-218。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Lian Duan; Jiangxue Fang | 2026-03-29

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019642 | OC-1501

Derived Weil Representation and Relative Langlands Duality | main.tex:217-223

定位：main.tex:217-223。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Haoshuo Fu | 2026-03-27

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019643 | OC-1502

Finiteness of Disjoint Covering Systems with Precisely One Repeated Modulus | main.tex:872-878

定位：main.tex:872-878。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Yu Hashimoto | 2026-03-27

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019644 | OC-151

Concentration for random Euclidean combinatorial optimization | Concentration_new.tex:117-124

定位：Concentration_new.tex:117-124。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Matteo D'Achille; Francesco Mattesini; Dario Trevisan | 2026-02-25

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019645 | OC-152

Scalable Kernel-Based Distances for Statistical Inference and Integration | main.tex:6088-6095

定位：main.tex:6088-6095。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Masha Naslidnyk | 2026-02-25

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_sa

V3-019646 | OC-1537

Newton polytopes of immanants of some combinatorial matrices | SNP-immanants.tex:114-116

定位：SNP-immanants.tex:114-116。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Candice X. T. Zhang | 2026-04-08

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019647 | OC-1538

Newton polytopes of immanants of some combinatorial matrices | SNP-immanants.tex:168-170

定位：SNP-immanants.tex:168-170。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Candice X. T. Zhang | 2026-04-08

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019648 | OC-1545

Greedy sparsifications of sums of positive semidefinite matrices | Greedy_Rudelson_standalone.tex:574-598

定位：Greedy_Rudelson_standalone.tex:574-598。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Grigory Ivanov | 2026-04-07

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019649 | OC-1582

Equitable coloring of large bipartite graphs | main.tex:35-37

定位：main.tex:35-37。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Amir Nikabadi | 2026-04-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019650 | OC-1589

Gromov-Hausdorff limits of the Chern-Ricci flow on smooth Hermitian minimal models of general type | main.tex:141-151

定位：main.tex:141-151。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Haoyuan Sun | 2026-04-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019651 | OC-1596

Non-existence probabilities and lower tails in the critical regime via Belief Propagation | TreeUniqueness.tex:416-423

定位：TreeUniqueness.tex:416-423。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Matthew Jenssen; Will Perkins; Aditya Potukuchi; Michael Simkin | 2026-04-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019652 | OC-1598

On the structure of approximate rings | completversion5.tex:2069-2071

定位：completversion5.tex:2069-2071。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Krzysztof Krupiński; Simon Machado | 2026-04-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019653 | OC-1629

Statistical Inference for Fractional Diffusions | main.tex:1455-1467

定位：main.tex:1455-1467。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Pablo Ramses Alonso-Martin; Horatio Boedihardjo; Anastasia Papavasiliou | 2026-04-04

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019654 | OC-1667

Infinitesimal sl(2)-symmetries on the equivariant skein lasagna module | main.tex:2903-2913

定位：main.tex:2903-2913。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | You Qi; Louis-Hadrien Robert; Joshua Sussan; Emmanuel Wagner; Paul Wedrich | 2026-04-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019655 | OC-1679

A Note on Generalized Erdős-Rogers Problems | main.tex:111-113

定位：main.tex:111-113。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Longma Du; Xinyu Hu; Ruilong Liu; Guanghui Wang | 2026-04-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019656 | OC-1680

A Note on Generalized Erdős-Rogers Problems | main.tex:144-147

定位：main.tex:144-147。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Longma Du; Xinyu Hu; Ruilong Liu; Guanghui Wang | 2026-04-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019657 | OC-1686

A rational model for the fiberwise THH transfer I: Sullivan algebras | 1main.tex:105-107

定位：1main.tex:105-107。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Florian Naef; Robin Stoll | 2026-04-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019658 | OC-1695

Counting Theorems for Algebraic Relations | main.tex:201-208

定位：main.tex:201-208。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Binyamini, Gal; Hirata-Kohno, Noriko; Kawashima, Makoto; Salant, Yuval | 2026-04-16

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019659 | OC-1696

Counting Theorems for Algebraic Relations | main.tex:241-250

定位：main.tex:241-250。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Binyamini, Gal; Hirata-Kohno, Noriko; Kawashima, Makoto; Salant, Yuval | 2026-04-16

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019660 | OC-170

Distributions of unramified extensions of global fields | ggrou.tex:271-275

定位：ggrou.tex:271-275。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Will Sawin; Melanie Matchett Wood | 2026-02-24

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019661 | OC-1718

Formal degree of principal series of quasi-split groups | main.tex:190-193

定位：main.tex:190-193。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Ricci, Giulio | 2026-04-16

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019662 | OC-1720

Locally Equienergetic Graphs | arXiv_Version.tex:483-485

定位：arXiv_Version.tex:483-485。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Dede, Cahit; Popat, Kalpesh M. | 2026-04-16

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019663 | OC-1809

Growth in noncommutative algebras and entropy in derived categories | pe4.tex:246-260

定位：pe4.tex:246-260。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Piontkovski, Dmitri | 2026-04-15

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_sa

V3-019664 | OC-1814

Fractional Strict Degeneracy of Graphs | Paper1.tex:229-231

定位：Paper1.tex:229-231。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Dominik, Daniel; Mudrock, Jeffrey A. | 2026-04-14

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019665 | OC-1830

An upper bound on the smallest singular value of dense random combinatorial matrices | Proj_2_Upper_Bound_LSV_Dense_revision.tex:474-482

定位：Proj_2_Upper_Bound_LSV_Dense_revision.tex:474-482。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Li, Dongbin; Litvak, Alexander E.; Yu, Tingzhou | 2026-04-14

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019666 | OC-1832

Semiorthogonal decompositions and components of derived categories of orthogonal Grassmannian fibrations | cliffordalgebrasv3.tex:1612-1614

定位：cliffordalgebrasv3.tex:1612-1614。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Shah, Saket | 2026-04-13

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019667 | OC-185

Monodromic singularities of Brunella-Miari vector fields with two edges in the Newton diagram | GarGinMan26-final-arXiv.tex:705-707

定位：GarGinMan26-final-arXiv.tex:705-707。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Isaac A. García; Jaume Giné; Víctor Mañosa | 2026-02-24

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019668 | OC-1850

Paths of length five with equal-degree endpoints | main.tex:73-76

定位：main.tex:73-76。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Liu, Zhen; Zeng, Qinghou | 2026-04-13

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019669 | OC-1874

On the Sum of Squares of Irreducible Brauer Character Degrees in Blocks | On_the_Sum_of_Squares_of_Irreducible_Brauer_Character_Degrees_in_Blocks.tex:225-229

定位：On_the_Sum_of_Squares_of_Irreducible_Brauer_Character_Degrees_in_Blocks.tex:225-229。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Li, Hanxiao; Zhang, Kun | 2026-04-13

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019670 | OC-1875

On the Sum of Squares of Irreducible Brauer Character Degrees in Blocks | On_the_Sum_of_Squares_of_Irreducible_Brauer_Character_Degrees_in_Blocks.tex:232-234

定位：On_the_Sum_of_Squares_of_Irreducible_Brauer_Character_Degrees_in_Blocks.tex:232-234。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Li, Hanxiao; Zhang, Kun | 2026-04-13

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019671 | OC-1896

Fundamental fields in the deformed W-algebras | walg_copie.tex:255-262

定位：walg_copie.tex:255-262。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Assakaf, Hicham | 2026-04-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019672 | OC-1897

Fundamental fields in the deformed W-algebras | walg_copie.tex:304-306

定位：walg_copie.tex:304-306。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Assakaf, Hicham | 2026-04-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019673 | OC-1920

Counting sunflowers with restricted matching number | Counting_sunflowers_with_restricted_matching_number.tex:97-99

定位：Counting_sunflowers_with_restricted_matching_number.tex:97-99。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Haixiang Zhang; Mengyu Cao; Mei Lu | 2026-04-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019674 | OC-1921

The Representation Type of the Descent Algebras | Apr26-v2.tex:759-769

定位：Apr26-v2.tex:759-769。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Karin Erdmann; Kay Jin Lim | 2026-04-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019675 | OC-1922

On the hamiltonicity problem of bicirculants: a reduction to cyclic Haar graphs | main_3rd_paper_20260423.tex:135-137

定位：main_3rd_paper_20260423.tex:135-137。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Simona Bonvicini; Tomaž Pisanski; Arjana Žitnik | 2026-04-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019676 | OC-194

A note on the affine plank conjecture | main.tex:111-118

定位：main.tex:111-118。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Egor Bakaev; Amir Yehudayoff | 2026-02-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019677 | OC-1940

An inductive approach to the Diaz-Park sharpness conjecture | inductive_approach_to_the_diaz_park_sharpness_conjecture.tex:328-334

定位：inductive_approach_to_the_diaz_park_sharpness_conjecture.tex:328-334。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Marco Praderio Bova | 2026-04-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019678 | OC-1943

Strong factorization theorem for smooth vectors of exponential solvable Lie group representations | Strong_factorization_theorem_expoential_solvable_Lie_group_representations22_4_26.tex:137-139

定位：Strong_factorization_theorem_expoential_solvable_Lie_group_representations22_4_26.tex:137-139。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Santiago Chaves; Andreas Debrouwere; Alberto Hernández Alvarado; Jasson Vindas; Rafael Zamora | 2026-04-22

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019679 | OC-1997

Analysis of persistence thresholds for a nonlocal PDE--ODE model of bacterial persister cells | main.tex:870-873

定位：main.tex:870-873。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Li, Chongming; Meadows, Tyler; Day, Troy | 2026-04-20

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019680 | OC-2006

An elementary proof of Sierksma's conjecture for seven points in the plane | paper-version.tex:74-76

定位：paper-version.tex:74-76。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Soberón, Pablo | 2026-04-20

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019681 | OC-2007

Families of symplectic Galois representations over small parabolic eigenvarieties for Siegel cuspforms of genus 2 | main.tex:3537-3539

定位：main.tex:3537-3539。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Manji, Muhammad; Thøgersen, Frederick E.; Wu, Ju-Feng | 2026-04-20

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019682 | OC-2050

Resolutions of spaces of crystalline representations and modularity | Master.tex:3252-3257

定位：Master.tex:3252-3257。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Bartlett, Robin; Hung, Bao V. Le; Levin, Brandon | 2026-04-19

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019683 | OC-2051

Resolutions of spaces of crystalline representations and modularity | Master.tex:3302-3307

定位：Master.tex:3302-3307。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Bartlett, Robin; Hung, Bao V. Le; Levin, Brandon | 2026-04-19

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019684 | OC-2075

Hypergraphs without Subgraphs of Given Connectivity | main_arXiv_version.tex:94-96

定位：main_arXiv_version.tex:94-96。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Ma, Jie; Xie, Shengjie; Zheng, Zhiheng | 2026-04-18

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019685 | OC-2076

Hypergraphs without Subgraphs of Given Connectivity | main_arXiv_version.tex:1662-1668

定位：main_arXiv_version.tex:1662-1668。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Ma, Jie; Xie, Shengjie; Zheng, Zhiheng | 2026-04-18

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019686 | OC-2111

Bergman--Einstein Rigidity for Hartogs Domains over Bounded Homogeneous Domains | EinsteinBergmannHartogs.tex:244-251

定位：EinsteinBergmannHartogs.tex:244-251。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Mossa, Roberto | 2026-04-17

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019687 | OC-2135

Sufficient conditions for spanning k-trees in tough graphs | spanning-k-tree-and-tough_4.30_.tex:97-99

定位：spanning-k-tree-and-tough_4.30_.tex:97-99。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Caili Jia; Yong Lu | 2026-04-30

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: missing

V3-019688 | OC-2176

BPS spectra of $\operatorname{tr}[\Psi^p]$ matrix models for odd p | paper_BPS.tex:813-821

定位：paper_BPS.tex:813-821。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Tierz, Miguel | 2026-04-29

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019689 | OC-2206

The coordinate ring of the universal centralizer via Demazure operators | main.tex:505-507

定位：main.tex:505-507。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Gannon, Tom; Ginzburg, Victor | 2026-04-28

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019690 | OC-2216

Geometry of Logarithmic Topological Recursion: Dilaton Equations, Free Energies and Variational Formulas | FinalVersionLogTRGeometry.tex:788-790

定位：FinalVersionLogTRGeometry.tex:788-790。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Hock, Alexander; Marchal, Olivier; Orantin, Nicolas | 2026-04-28

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_sa

V3-019691 | OC-2232

Size-4 Counterexamples to the Sidon-Extension Conjecture | paper_arxiv.tex:154-158

定位：paper_arxiv.tex:154-158。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Niu, Tong | 2026-04-28

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019692 | OC-2233

Size-4 Counterexamples to the Sidon-Extension Conjecture | paper_arxiv.tex:162-168

定位：paper_arxiv.tex:162-168。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Niu, Tong | 2026-04-28

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019693 | OC-2270

Algebraic expansivity on abelian groups | expgrouphom260427_-_arxiv.tex:717-719

定位：expgrouphom260427_-_arxiv.tex:717-719。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Achigar, Mauricio | 2026-04-27

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019694 | OC-2282

Dimension of the motivic Galois group of a 1-motive | DimGalMot.tex:288-304

定位：DimGalMot.tex:288-304。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Bertolin, Cristiana | 2026-04-27

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019695 | OC-2304

Towards a conjecture on degree conditions for Ramsey goodness of paths | 4.24.tex:108-116

定位：4.24.tex:108-116。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | You, Chunlin | 2026-04-25

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019696 | OC-2311

Statistical Analysis of Markovian Generative Modeling | main.tex:2376-2380

定位：main.tex:2376-2380。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Aamari, Eddie; Stéphanovitch, Arthur | 2026-04-24

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019697 | OC-2332

Twisted traces and quantization of moduli stacks of 3d $\mathcal{N}=4$ Chern-Simons-matter theories | 3dN4CS_v1.tex:185-191

定位：3dN4CS_v1.tex:185-191。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Santilli, Leonardo | 2026-04-22

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019698 | OC-2333

Twisted traces and quantization of moduli stacks of 3d $\mathcal{N}=4$ Chern-Simons-matter theories | 3dN4CS_v1.tex:972-988

定位：3dN4CS_v1.tex:972-988。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Santilli, Leonardo | 2026-04-22

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019699 | OC-2334

Twisted traces and quantization of moduli stacks of 3d $\mathcal{N}=4$ Chern-Simons-matter theories | 3dN4CS_v1.tex:1029-1035

定位：3dN4CS_v1.tex:1029-1035。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Santilli, Leonardo | 2026-04-22

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019700 | OC-2340

Accessory Parameter of Confluent Heun Equations, Voros Periods and classical irregular conformal blocks | AP-VP-CCB-v1.tex:622-625

定位：AP-VP-CCB-v1.tex:622-625。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Iwaki, Kohei; Nagoya, Hajime; Shukuta, Ayato | 2026-05-07

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019701 | OC-2358

Fourier coefficients of continuous functions with sparse spectrum | lacunary_KKdL_final.tex:810-813

定位：lacunary_KKdL_final.tex:810-813。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Kulikov, Aleksei; Saucedo, Miquel; Tikhonov, Sergey | 2026-05-07

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019702 | OC-2360

Spaces of matrices with few eigenvalues (II) | feweigenvaluesIIarxiv.tex:420-423

定位：feweigenvaluesIIarxiv.tex:420-423。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Pazzis, Clément de Seguins | 2026-05-07

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019703 | OC-2361

Spaces of matrices with few eigenvalues (II) | feweigenvaluesIIarxiv.tex:436-440

定位：feweigenvaluesIIarxiv.tex:436-440。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Pazzis, Clément de Seguins | 2026-05-07

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019704 | OC-2381

Energy-minimizing measures supported near fractal 1-sets | submission.tex:647-649

定位：submission.tex:647-649。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Bongers, Rosemarie | 2026-05-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_sa

V3-019705 | OC-2385

Exponential sums over singular binary quintics | BinQuintOnly__2_.tex:118-122

定位：BinQuintOnly__2_.tex:118-122。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Ishitsuka, Yasuhiro | 2026-05-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019706 | OC-2386

Neural Discovery of Strichartz Extremizers | main_Neurips_2026v9.tex:335-341

定位：main_Neurips_2026v9.tex:335-341。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Valenzuela, Nicolás; Freire, Ricardo; Muñoz, Claudio | 2026-05-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019707 | OC-2387

Neural Discovery of Strichartz Extremizers | main_Neurips_2026v9.tex:1053-1063

定位：main_Neurips_2026v9.tex:1053-1063。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Valenzuela, Nicolás; Freire, Ricardo; Muñoz, Claudio | 2026-05-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019708 | OC-2390

An Intersection-Weighted Erd\H{o}s-Ko-Rado Theorem | tompkinssubmit.tex:1006-1015

定位：tompkinssubmit.tex:1006-1015。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Tompkins, Casey | 2026-05-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019709 | OC-2408

The pebbling number of Fibonacci cubes | paper_arxiv.tex:99-101

定位：paper_arxiv.tex:99-101。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Niu, Tong | 2026-05-05

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019710 | OC-2437

A note on cycles in cyclically 4-edge-connected cubic planar graphs | Paper.tex:130-132

定位：Paper.tex:130-132。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Lo, On-Hei Solomon | 2026-05-05

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019711 | OC-2442

Higher categories of bordisms with geometric structures | hcbgs.tex:5268-5277

定位：hcbgs.tex:5268-5277。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Grady, Daniel; Pavlov, Dmitri | 2026-05-05

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019712 | OC-2443

Extended field theories are local and have classifying spaces | bord_desc.tex:2881-2888

定位：bord_desc.tex:2881-2888。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Grady, Daniel; Pavlov, Dmitri | 2020-11-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019713 | OC-2447

Global dimension of the category of rational incomplete Mackey functors for a finite abelian group G | Paper_-_HA.tex:1526-1528

定位：Paper_-_HA.tex:1526-1528。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Barnes, David; Bohmann, Anna Marie; Hill, Michael A.; Kędziorek, Magdalena | 2026-05-04

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019714 | OC-246

A function field analogue of Ligozat's theorem for Drinfeld modular units | main.tex:127-129

定位：main.tex:127-129。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Sheng-Yang Kevin Ho | 2026-02-20

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019715 | OC-2462

A characterization for positive semi-definite matrix products | PosSD_MatrixProd.tex:316-318

定位：PosSD_MatrixProd.tex:316-318。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Garbe, Frederik; Wei, Fan | 2026-05-04

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019716 | OC-2495

Extremal Problems for the Family of k-Strongly Connected Digraphs | Extremal_Problems_for_the_Family_of_k-Strongly_Connected_Digraphs.tex:376-378

定位：Extremal_Problems_for_the_Family_of_k-Strongly_Connected_Digraphs.tex:376-378。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Wang, Qinglin; Tian, Yingzhi | 2026-05-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019717 | OC-2500

Improved Rate-versus-Distance Upper Bounds for LDPC Codes | main.tex:728-736

定位：main.tex:728-736。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Shangguan, Chong; Yang, Yulin | 2026-05-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019718 | OC-2508

Linear-Readout Floors and Threshold Recovery in Computation in Superposition | arxiv_v11.tex:1964-1995

定位：arxiv_v11.tex:1964-1995。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Borobia, Hector; Seguí-Mas, Elies; Tormo-Carbó, Guillermina | 2026-05-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019719 | OC-2520

On the Global Curve Attractor for polynomial gluing | FGA-W.tex:102-106

定位：FGA-W.tex:102-106。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Wu, Panjing | 2026-05-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019720 | OC-2528

An exact small-n computation of the minimum 2-coloring discrepancy of $K_n^{(3)}$ | paper_arxiv.tex:192-199

定位：paper_arxiv.tex:192-199。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Niu, Tong | 2026-05-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019721 | OC-2529

On ${\pi}$-systems of symmetrizable Kac-Moody algebras | tex-me-2023.tex:1752-1754

定位：tex-me-2023.tex:1752-1754。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Raghavan, K. N.; Roy, Krishanu; Viswanath, S. | 2026-05-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: missing

V3-019722 | OC-2535

Characterisation of Stability for Interval Translation Maps | main.tex:217-221

定位：main.tex:217-221。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Drach, Kostiantyn; Staresinic, Leon; van Strien, Sebastian | 2026-04-30

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019723 | OC-2539

Transversality for Interval Translation Maps | main.tex:225-229

定位：main.tex:225-229。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Drach, Kostiantyn; Staresinic, Leon; van Strien, Sebastian | 2026-04-30

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019724 | OC-2542

The density conjecture for activated random walk | main.tex:389-393

定位：main.tex:389-393。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Hoffman, Christopher; Johnson, Tobias; Junge, Matthew | 2024-06-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019725 | OC-2546

Elliptic Generalization of Cherednik-Macdonald-Mehta identities | IntegralIdentities.tex:357-368

定位：IntegralIdentities.tex:357-368。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Shamil Shakirov | 2026-05-25

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019726 | OC-2578

A Borodin-Okounkov-Geronimo-Case identity for tilted Toeplitz minors | tilted_Toeplitz.tex:2118-2134

定位：tilted_Toeplitz.tex:2118-2134。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Leonid Petrov | 2026-05-24

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019727 | OC-2581

A Classification of Unmixed Binomial Edge Ideals of König Type | A_Classification_of_Unmixed_Binomial_Edge_Ideals_of_Koenig_Type.tex:484-486

定位：A_Classification_of_Unmixed_Binomial_Edge_Ideals_of_Koenig_Type.tex:484-486。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | David Williams | 2026-05-24

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019728 | OC-2605

On exotic rationally integrable planar dual billiards I. Complex geometry and type of dynamics | int-rat-dual-cons2.tex:158-161

定位：int-rat-dual-cons2.tex:158-161。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Alexey Glutsyuk | 2026-05-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019729 | OC-2606

Framing Triangulations for Arbitrary Integer Flow Polytopes | main.tex:1151-1153

定位：main.tex:1151-1153。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Jonah Berggren | 2026-05-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019730 | OC-2631

Path-Minimality of p-Energy for Connected Graphs | combined_v16.tex:5216-5221

定位：combined_v16.tex:5216-5221。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Yinchen Liu; Quanyu Tang | 2026-05-21

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019731 | OC-2652

On Isotopies and hyperbolicity of weaves | ihw.tex:790-792

定位：ihw.tex:790-792。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Ken'ichi Yoshida | 2026-05-21

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019732 | OC-2660

Indecomposable Quasiconformal Maps of Manifolds | IndecomposableQC.tex:218-220

定位：IndecomposableQC.tex:218-220。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Benjamin B. McMillan | 2026-05-21

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019733 | OC-2694

On degree bounds of k-uniform hypergraphs with bounded matching number | degree_matching_5.20.tex:113-115

定位：degree_matching_5.20.tex:113-115。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Haixiang Zhang; Mengyu Cao; Mei Lu | 2026-05-20

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019734 | OC-2756

On hamiltonian cycles of 1-tough $(P_{2} \cup kP_{1})$-free graphs | P2kP1_GC_.tex:125-127

定位：P2kP1_GC_.tex:125-127。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Masahiro Sanka | 2026-05-19

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019735 | OC-2764

Sharp analysis of sketched least squares and randomized low-rank approximation | report.tex:1524-1550

定位：report.tex:1524-1550。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Ethan N. Epperly; Robert J. Webber | 2026-05-18

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019736 | OC-277

Dynamics of Nesterov's Accelerated Gradient Descent in Quadratic Games | main.tex:532-535

定位：main.tex:532-535。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Jay Paek | 2026-02-19

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019737 | OC-2802

A Fourier perspective on the learning dynamics of neural networks: from sample complexities to mechanistic insights | Fourier_arXiv.tex:759-761

定位：Fourier_arXiv.tex:759-761。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Fabiola Ricci; Claudia Merger; Sebastian Goldt | 2026-05-16

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019738 | OC-2804

On t-edge-balanced graphs | t-edge-balanced.tex:366-368

定位：t-edge-balanced.tex:366-368。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Yeow Meng Chee | 2026-05-16

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019739 | OC-2831

Liouville theorem for immersed minimal surfaces in any codimension | Liouville2D57.tex:291-297

定位：Liouville2D57.tex:291-297。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Tobias Holck Colding; William P. Minicozzi | 2026-05-14

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019740 | OC-2864

Brieskorn spheres and rational homology ball symplectic fillings | Arxiv.tex:161-162

定位：Arxiv.tex:161-162。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Antonio Alfieri; Alberto Cavallo; Irena Matkovič | 2026-05-13

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019741 | OC-2870

On p-Lie algebras of finite Morley rank | p-algebre.tex:488-490

定位：p-algebre.tex:488-490。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Samuel Zamour | 2026-05-13

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_sa

V3-019742 | OC-2893

ℓ^{p} improving estimates for multilinear forms motivated by distance graphs | PalssonSmucker.tex:485-493

定位：PalssonSmucker.tex:485-493。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Eyvindur Palsson; Jennifer Smucker | 2026-05-12

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019743 | OC-2898

Intrinsic volumes of ℓ_p-balls and a continuum of Maxwell--Poincaré--Borel laws for their curvature measures | intrinsic_vol_l_p_balls.tex:948-958

定位：intrinsic_vol_l_p_balls.tex:948-958。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zakhar Kabluchko; Alexander Marynych; Joscha Prochno | 2026-05-12

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019744 | OC-2923

Dimension-free Gaussian tail estimates for linear functionals on convex bodies | main.tex:1352-1362

定位：main.tex:1352-1362。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Brayden Letwin; Dan Mikulincer | 2026-05-11

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019745 | OC-2930

Logarithmic scaling of selective sweep curves: from tents to houses | final_house_3.tex:590-593

定位：final_house_3.tex:590-593。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Florin Boenkost; Felix Hermann; András Tóbiás; Anton Wakolbinger | 2026-05-11

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019746 | OC-2942

Two-dimensional Virasoro algebras | 2dGRR.tex:47-60

定位：2dGRR.tex:47-60。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhengping Gui; Brian R. Williams | 2026-05-11

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019747 | OC-3008

Quadratic linear strands of prime ideals | quadrics.tex:373-374

定位：quadrics.tex:373-374。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Giulio Caviglia; Alessandro De Stefani | 2026-05-09

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019748 | OC-3020

Universal Structure of Graph Product Kernels | main.tex:1136-1137

定位：main.tex:1136-1137。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Ian J. Leary; Nansen Petrosyan | 2026-05-08

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019749 | OC-3098

Positive density for Sun's 2^k+m conjecture | main.tex:90-93

定位：main.tex:90-93。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Songlin Han; Jinbo Yu | 2026-05-15

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019750 | OC-323

On certain subspaces of 2-configuration spaces of graphs | paper1Final.tex:2730-2732

定位：paper1Final.tex:2730-2732。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Byung Hee An; Sangrok Oh | 2026-02-17

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019751 | OC-3384

Rigidity in the Positive Mass Theorem with C^0 Decay | main.tex:166-168

定位：main.tex:166-168。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Liam Mazurowski; Xuan Yao | 2026-05-28

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019752 | OC-34

Beyond singular value gaps in randomized subspace approximation | WangTownsendSIMAX.tex:1052-1058

定位：WangTownsendSIMAX.tex:1052-1058。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Christopher Wang; Alex Townsend | 2026-03-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019753 | OC-3414

Dual Chow polynomials of matroids and posets | main.tex:2354-2356

定位：main.tex:2354-2356。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Giovanni Caiolo; Luis Ferroni; Elena Hoster | 2026-05-27

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019754 | OC-3431

Homological Nielsen realization for the manifolds $\#_n \mathbb{CP}^2$ | final_draft.tex:879-881

定位：final_draft.tex:879-881。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Ethan Pesikoff | 2026-05-26

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019755 | OC-3435

Explicit class of finite-dimensional polynomial algebras with Wronskians over $\mathbb{R}^d$ as N-ary Lie brackets: beyond $\mathfrak{sl}(2)$ | FinDim220526.tex:2077-2079

定位：FinDim220526.tex:2077-2079。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Markuss G. Ķēniņš; Arthemy V. Kiselev | 2026-05-26

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019756 | OC-3448

Absolutely indecomposable quasi-parabolic G-bundles and the multiplicity of irreducible characters | Absolutely_indecomposable_parabolic_G_bundles_and_multiplicity_-_arXiv.tex:497-499

定位：Absolutely_indecomposable_parabolic_G_bundles_and_multiplicity_-_arXiv.tex:497-499。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | GyeongHyeon Nam | 2026-05-26

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019757 | OC-3471

Varieties admitting a holomorphic symplectic form: LLV algebras and derived equivalences | chapters/introduction.tex:22-25

定位：chapters/introduction.tex:22-25。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Dion Leijnse | 2026-05-26

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019758 | OC-3472

Varieties admitting a holomorphic symplectic form: LLV algebras and derived equivalences | chapters/introduction.tex:27-29

定位：chapters/introduction.tex:27-29。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Dion Leijnse | 2026-05-26

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019759 | OC-3473

Varieties admitting a holomorphic symplectic form: LLV algebras and derived equivalences | chapters/summary.tex:17-19

定位：chapters/summary.tex:17-19。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Dion Leijnse | 2026-05-26

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019760 | OC-3476

Minimal surfaces, Knots, and Neural Networks | paper_imported.tex:185-189

定位：paper_imported.tex:185-189。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Tancredi Schettini Gherardini; Marco Usula | 2026-05-25

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019761 | OC-3477

Minimal surfaces, Knots, and Neural Networks | paper_imported.tex:869-876

定位：paper_imported.tex:869-876。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Tancredi Schettini Gherardini; Marco Usula | 2026-05-25

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019762 | OC-3554

Bipartite cuts in Ramsey-Turán style | arxiv1.tex:457-465

定位：arxiv1.tex:457-465。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | József Balogh; Ce Chen; Bernard Lidický | 2026-06-18

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019763 | OC-3559

Spectral and size conditions for spanning k-trees in tough graphs | Spectral_and_size_conditions_for_spanning_k-trees_in_tough_graphs.tex:80-82

定位：Spectral_and_size_conditions_for_spanning_k-trees_in_tough_graphs.tex:80-82。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Siyuan Liang; Tao Tian | 2026-06-18

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019764 | OC-3602

Rigidity of infinite exchangeable sequences with Gaussian marginals | main.tex:134-136

定位：main.tex:134-136。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Maximillian Newman | 2026-06-17

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: missing

V3-019765 | OC-3603

Interval hypergraphic polytopes (or deformed associahedra), Tamari interval posets, and weeping willows | simpleIntervalHypergraphicPolytopes.tex:1315-1318

定位：simpleIntervalHypergraphicPolytopes.tex:1315-1318。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Jose Bastidas; Félix Gélinas; Vincent Pilaud; Germain Poullot; Andrew Sack; Eleni Tzanaki | 2026-06-16

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019766 | OC-3630

Invariant complex structures for affine automorphisms: a cocycle viewpoint | main.tex:573-576

定位：main.tex:573-576。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Jiesong Zhang | 2026-06-16

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019767 | OC-3631

Invariant complex structures for affine automorphisms: a cocycle viewpoint | main.tex:582-585

定位：main.tex:582-585。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Jiesong Zhang | 2026-06-16

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019768 | OC-3657

KKM theorems and discrete geometry beyond matroids | v3-paper.tex:1624-1626

定位：v3-paper.tex:1624-1626。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Pablo Soberón | 2026-06-15

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019769 | OC-3670

Bootstrap Percolation and Partial Difference Equations | bpde-paper.tex:401-401

定位：bpde-paper.tex:401-401。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Nikolai Beluhov | 2026-06-15

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019770 | OC-3677

Testing the max-flow min-cut property and the replication conjecture | main.tex:171-173

定位：main.tex:171-173。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Ahmad Abdi; Tamás Schwarcz | 2026-06-15

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019771 | OC-3678

Testing the max-flow min-cut property and the replication conjecture | main.tex:207-209

定位：main.tex:207-209。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Ahmad Abdi; Tamás Schwarcz | 2026-06-15

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019772 | OC-3679

Testing the max-flow min-cut property and the replication conjecture | main.tex:1283-1285

定位：main.tex:1283-1285。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Ahmad Abdi; Tamás Schwarcz | 2026-06-15

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019773 | OC-3680

Testing the max-flow min-cut property and the replication conjecture | main.tex:1319-1321

定位：main.tex:1319-1321。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Ahmad Abdi; Tamás Schwarcz | 2026-06-15

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019774 | OC-3681

Testing the max-flow min-cut property and the replication conjecture | main.tex:1327-1329

定位：main.tex:1327-1329。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Ahmad Abdi; Tamás Schwarcz | 2026-06-15

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019775 | OC-3788

Trivariate Hypergeometric Series Formulas for Pure Partition Functions of Multiple 3-SLE$_κ$ | SLEkapparho1rho2.tex:3116-3122

定位：SLEkapparho1rho2.tex:3116-3122。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Dapeng Zhan | 2026-06-12

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019776 | OC-38

Operator symmetric moduli and sharp triangle inequalities | Operator_symmetric_moduli_and_sharp_triangle_inequalities.tex:187-193

定位：Operator_symmetric_moduli_and_sharp_triangle_inequalities.tex:187-193。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Teng Zhang | 2026-03-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019777 | OC-380

On the Non-Orientable 3- and 4-Genera of a Knot: Connections and Comparisons | non-ori-3d-4d.tex:229-231

定位：non-ori-3d-4d.tex:229-231。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Julia Knihs; Jeanette Patel; Joshua M. Sabloff; Thea Rugg | 2026-02-13

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019778 | OC-383

The failure of square at all uncountable cardinals is weaker than a Woodin limit of Woodin cardinals | nairian_2_main.tex:696-703

定位：nairian_2_main.tex:696-703。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Douglas Blue; Paul Larson; Grigor Sargsyan | 2026-02-13

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019779 | OC-3831

Compatibility of Higher-Order Slow-Manifold Reduction and Continuum Limits in Adaptive Networks | main.tex:3284-3291

定位：main.tex:3284-3291。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Christian Kuehn; Fergal Murphy; Jan-Eric Sulzbach | 2026-06-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019780 | OC-3898

A correlated refinement of the double double ramification cycle | main.tex:736-742

定位：main.tex:736-742。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Thomas Blomme; Francesca Carocci; Ajith Urundolil Kumaran | 2026-06-09

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019781 | OC-3924

Dimension lower bounds in random geometry via Lipschitz functions | main.tex:411-414

定位：main.tex:411-414。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Manan Bhatia; Ewain Gwynne; Brin Harper | 2026-06-09

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019782 | OC-3949

Clock spacing for two-sided Jacobi matrices | main.tex:297-310

定位：main.tex:297-310。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Benjamin Eichinger; Milivoje Lukić; Giorgio Young | 2026-06-08

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019783 | OC-3987

A complete characterization of a family of permutation trinomials over $\mathbb F_{p^2}$ | 33-T.tex:1947-1955

定位：33-T.tex:1947-1955。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Marco Timpanella | 2026-06-07

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019784 | OC-3989

Large point-degrees in intersecting families of finite vector spaces | main.tex:241-251

定位：main.tex:241-251。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Jicheng Ma; Hao Li | 2026-06-07

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019785 | OC-4

Some properties of minimally nonperfectly divisible graphs | main.tex:113-115

定位：main.tex:113-115。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Qiming Hu; Baogang Xu; Miaoxia Zhuang | 2026-03-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: missing

V3-019786 | OC-4005

Digit Mixing under Polynomial Maps | bernoulli.tex:225-231

定位：bernoulli.tex:225-231。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chokri Manai | 2026-06-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019787 | OC-4012

$\textbf{k}$-neighborhood ideals of graphs | k-neighborhood_ideals_of_graphs.tex:398-400

定位：k-neighborhood_ideals_of_graphs.tex:398-400。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Somayeh Moradi; Leila Sharifan | 2026-06-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019788 | OC-4025

The singular cohomology ring of a uniform matroid: combinatorics and Lefschetz properties | preprint.tex:1709-1711

定位：preprint.tex:1709-1711。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Kyle Binder | 2026-06-05

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019789 | OC-4029

A study of maximal shifts in the minimal graded free resolution of the residue field | main.tex:1119-1129

定位：main.tex:1119-1129。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Dinesh Limbu; Liana M. Şega; Adela Vraciu | 2026-06-05

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019790 | OC-405

Khovanov concordance minima and the (4,5) torus knot | t45.tex:211-214

定位：t45.tex:211-214。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Andrew Lobb | 2026-02-13

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019791 | OC-406

Khovanov concordance minima and the (4,5) torus knot | t45.tex:217-220

定位：t45.tex:217-220。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Andrew Lobb | 2026-02-13

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019792 | OC-4062

Tighter Bounds on the Degree-Truncated Choice Number of Planar Graphs | Tight_bounds_on_degree-truncated_choice_number.tex:204-207

定位：Tight_bounds_on_degree-truncated_choice_number.tex:204-207。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Huijuan Xu; Huan Zhou; Jialu Zhu; Xuding Zhu | 2026-06-04

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019793 | OC-407

Khovanov concordance minima and the (4,5) torus knot | t45.tex:222-225

定位：t45.tex:222-225。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Andrew Lobb | 2026-02-13

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019794 | OC-4084

The leading constant in Malle's conjecture | Malle_Survey.tex:1732-1744

定位：Malle_Survey.tex:1732-1744。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Daniel Loughran; Tim Santens | 2026-06-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019795 | OC-4085

The leading constant in Malle's conjecture | Malle_Survey.tex:1887-1895

定位：Malle_Survey.tex:1887-1895。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Daniel Loughran; Tim Santens | 2026-06-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019796 | OC-4086

The leading constant in Malle's conjecture | Malle_Survey.tex:1943-1950

定位：Malle_Survey.tex:1943-1950。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Daniel Loughran; Tim Santens | 2026-06-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019797 | OC-4087

The leading constant in Malle's conjecture | Malle_Survey.tex:2048-2054

定位：Malle_Survey.tex:2048-2054。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Daniel Loughran; Tim Santens | 2026-06-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019798 | OC-4088

The leading constant in Malle's conjecture | Malle_Survey.tex:2067-2074

定位：Malle_Survey.tex:2067-2074。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Daniel Loughran; Tim Santens | 2026-06-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019799 | OC-4089

The leading constant in Malle's conjecture | Malle_Survey.tex:2105-2120

定位：Malle_Survey.tex:2105-2120。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Daniel Loughran; Tim Santens | 2026-06-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019800 | OC-4090

The leading constant in Malle's conjecture | Malle_Survey.tex:2203-2206

定位：Malle_Survey.tex:2203-2206。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Daniel Loughran; Tim Santens | 2026-06-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019801 | OC-4091

The leading constant in Malle's conjecture | Malle_Survey.tex:2252-2257

定位：Malle_Survey.tex:2252-2257。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Daniel Loughran; Tim Santens | 2026-06-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019802 | OC-4092

The leading constant in Malle's conjecture | Malle_Survey.tex:2419-2426

定位：Malle_Survey.tex:2419-2426。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Daniel Loughran; Tim Santens | 2026-06-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019803 | OC-4093

The leading constant in Malle's conjecture | Malle_Survey.tex:2434-2443

定位：Malle_Survey.tex:2434-2443。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Daniel Loughran; Tim Santens | 2026-06-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019804 | OC-4108

Third order Einstein deformations for Kaehler-Einstein metrics | 3E-def-v8.tex:434-449

定位：3E-def-v8.tex:434-449。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Paul-Andi Nagy | 2026-06-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019805 | OC-4119

Primitive Positive Constructions Among Finite Permutation Groups | main.tex:1727-1746

定位：main.tex:1727-1746。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Sebastian Meyer | 2026-06-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_sa

V3-019806 | OC-4120

Primitive Positive Constructions Among Finite Permutation Groups | main.tex:1755-1765

定位：main.tex:1755-1765。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Sebastian Meyer | 2026-06-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_sa

V3-019807 | OC-4135

Classical Symmetry TFTs for Continuous Symmetries via Higher Symplectic Geometry | main.tex:87-136

定位：main.tex:87-136。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Hao Xu | 2026-06-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019808 | OC-4144

Syzygies of Isotropic Kalman Varieties | 2025-gkp-isotropic-kalman.tex:502-517

定位：2025-gkp-isotropic-kalman.tex:502-517。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Suhas Vadan Gondi; Sarah Kumar; Abhik Pal | 2026-06-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_sa

V3-019809 | OC-4145

Syzygies of Isotropic Kalman Varieties | 2025-gkp-isotropic-kalman.tex:2195-2205

定位：2025-gkp-isotropic-kalman.tex:2195-2205。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Suhas Vadan Gondi; Sarah Kumar; Abhik Pal | 2026-06-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_sa

V3-019810 | OC-4146

Syzygies of Isotropic Kalman Varieties | 2025-gkp-isotropic-kalman.tex:2220-2236

定位：2025-gkp-isotropic-kalman.tex:2220-2236。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Suhas Vadan Gondi; Sarah Kumar; Abhik Pal | 2026-06-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_sa

V3-019811 | OC-4148

On maximal Dynkin friezes | Maximal_Dynkin_friezes_2026-05-30.tex:591-599

定位：Maximal_Dynkin_friezes_2026-05-30.tex:591-599。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Robin Zhang | 2026-06-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019812 | OC-4208

Extremal graphs with no subgraph admitting $k+1$ edge-disjoint spanning trees | manuscript.tex:501-503

定位：manuscript.tex:501-503。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Qinglin Wang; Yingzhi Tian | 2026-06-26

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019813 | OC-422

Eigenvalue Estimates of the Hodge Laplacian Under Lower Ricci Curvature Bound | v0.tex:98-100

定位：v0.tex:98-100。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Anusha Bhattacharya; Soma Maity; Aditya Tiwari | 2026-02-12

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: missing

V3-019814 | OC-4256

Structural Properties and Applications of the Augmented Sombor Index | ASO.tex:3635-3642

定位：ASO.tex:3635-3642。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chunlei Xu; Kinkar Chandra Das; Jayanta Bera | 2026-06-25

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019815 | OC-4268

Variation on a theme of Beauville-Voisin | variation.tex:169-172

定位：variation.tex:169-172。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Robert Laterveer | 2026-06-24

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019816 | OC-4304

Almost Symmetric Linear Arc Monadic Datalog and Transitive Tournaments | main.tex:2875-2877

定位：main.tex:2875-2877。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Sebastian Meyer; Florian Starke | 2026-06-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_sa

V3-019817 | OC-4324

Compact locally conformal Kähler manifolds with constant Chern holomorphic sectional curvature | LCKv2.tex:97-100

定位：LCKv2.tex:97-100。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Zhuzhu Huang; Xueyuan Wan | 2026-06-23

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019818 | OC-4344

The excluded minors for Z_{3}-gainable and regular biased graphs | arXiv-v1.tex:207-211

定位：arXiv-v1.tex:207-211。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Nick Brettell; Rutger Campbell; Daryl Funk; Dillon Mayhew | 2026-06-22

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019819 | OC-4345

The excluded minors for Z_{3}-gainable and regular biased graphs | arXiv-v1.tex:260-264

定位：arXiv-v1.tex:260-264。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Nick Brettell; Rutger Campbell; Daryl Funk; Dillon Mayhew | 2026-06-22

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019820 | OC-439

Random chain complexes of real vector spaces | main.tex:663-667

定位：main.tex:663-667。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Ayat Ababneh; Matthew Kahle | 2026-02-11

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019821 | OC-4396

Saturation numbers for joins of graphs and characterization of extremal graphs | joins_of_graphs1111_260623_latest_version.tex:567-569

定位：joins_of_graphs1111_260623_latest_version.tex:567-569。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Xinying Hua; Yuejian Peng | 2026-06-20

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019822 | OC-4397

Geometry of $\mathbf{F}_1$ and Cuntz-Krieger algebras | fun-nik.tex:864-867

定位：fun-nik.tex:864-867。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Igor V. Nikolaev | 2026-06-20

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019823 | OC-440

On the Calderón sum formula for wavelet systems | main.tex:165-171

定位：main.tex:165-171。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Ulrik Enstad; Jordy Timo van Velthoven | 2026-02-11

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019824 | OC-4411

Twofold universality of large-N melonic random tensors | uniwersalnosc_melonow.tex:1714-1733

定位：uniwersalnosc_melonow.tex:1714-1733。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Carlos I. Perez-Sanchez | 2026-07-09

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019825 | OC-4438

A reverse Riesz estimate combined with a spectral gap implies a Poincaré inequality | Poincare-v11-arxiv.tex:1923-1930

定位：Poincare-v11-arxiv.tex:1923-1930。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Cédric Arhancet | 2026-07-09

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019826 | OC-4439

A reverse Riesz estimate combined with a spectral gap implies a Poincaré inequality | Poincare-v11-arxiv.tex:2733-2738

定位：Poincare-v11-arxiv.tex:2733-2738。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Cédric Arhancet | 2026-07-09

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019827 | OC-4442

On the Etzion-Silberstein conjecture for block Ferrers diagrams | main.tex:677-680

定位：main.tex:677-680。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Marco Calderini; Marta Messia; Alessandro Neri | 2026-07-09

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019828 | OC-4479

Multiset Partition Dimension of Graphs | EJGTATemplate.tex:786-791

定位：EJGTATemplate.tex:786-791。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Azzah Albejania; Yuqing Lin; Joe Ryan; Kiki A. Sugeng | 2026-07-08

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019829 | OC-4489

Quantitative hyperbolicity for complex manifolds via numerical invariants | main.tex:2807-2809

定位：main.tex:2807-2809。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Tien-Cuong Dinh; Duc-Bao Nguyen; Duc-Viet Vu | 2026-07-08

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019830 | OC-4491

A sharp isoperimetric inequality and the top order Q-curvature | iso_ratio_about_Q7.tex:1091-1094

定位：iso_ratio_about_Q7.tex:1091-1094。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Mingxiang Li; Xingwang Xu | 2026-07-08

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019831 | OC-4511

Divisorial Persistence and Asymptotic Homology of Analytic Pairs | ArXiv_PDH.tex:5585-5615

定位：ArXiv_PDH.tex:5585-5615。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Nivaldo Grulha | 2026-07-07

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019832 | OC-4512

Divisorial Persistence and Asymptotic Homology of Analytic Pairs | ArXiv_PDH.tex:5665-5687

定位：ArXiv_PDH.tex:5665-5687。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Nivaldo Grulha | 2026-07-07

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019833 | OC-4518

Generalized spectral closedness of $\mathcal{F}$-free graph classes | manuscriptrevised.tex:687-689

定位：manuscriptrevised.tex:687-689。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Wei Wang; Quanyu Tang | 2026-07-07

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019834 | OC-4524

Purity of extremal rays of Betti cones | v1-purity-arXiv.tex:486-488

定位：v1-purity-arXiv.tex:486-488。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | H. Ananthnarayan; Omkar Javadekar; Rajiv Kumar | 2026-07-07

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019835 | OC-4525

Purity of extremal rays of Betti cones | v1-purity-arXiv.tex:933-935

定位：v1-purity-arXiv.tex:933-935。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | H. Ananthnarayan; Omkar Javadekar; Rajiv Kumar | 2026-07-07

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019836 | OC-4526

The Siegel-Weil formula in geometry and arithmetic | main_new_layout.tex:1484-1490

定位：main_new_layout.tex:1484-1490。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Jan Hendrik Bruinier; Riccardo Zuffetti | 2026-07-07

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019837 | OC-4527

The Siegel-Weil formula in geometry and arithmetic | main_new_layout.tex:1504-1514

定位：main_new_layout.tex:1504-1514。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Jan Hendrik Bruinier; Riccardo Zuffetti | 2026-07-07

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019838 | OC-4544

Polarised noncrossing partititions and the coherent self-dual $ω$-equivalence | main.tex:1122-1140

定位：main.tex:1122-1140。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Amar Hadzihasanovic | 2026-07-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019839 | OC-4560

Toric decomposition in algebraic groups | toricDecomposition.tex:1194-1196

定位：toricDecomposition.tex:1194-1196。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Dávid R. Szabó | 2026-07-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019840 | OC-4561

Toric decomposition in algebraic groups | toricDecomposition.tex:1214-1221

定位：toricDecomposition.tex:1214-1221。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Dávid R. Szabó | 2026-07-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019841 | OC-4562

Toric decomposition in algebraic groups | toricDecomposition.tex:1225-1229

定位：toricDecomposition.tex:1225-1229。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Dávid R. Szabó | 2026-07-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019842 | OC-4583

Derived operadic centers in algebraic geometry and deformation quantization | main.tex:2921-2926

定位：main.tex:2921-2926。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Sonja Farr | 2026-07-05

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019843 | OC-4599

The Rigidity Theorems for Self-Shrinkers in the Mean Curvature Flow | 708-7.tex:201-215

定位：708-7.tex:201-215。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Fagui Li; Yuhang Zhao | 2026-07-05

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019844 | OC-46

The Gromov-Hausdorff distance between l^p-products of metric spaces | 01_GH_Product_EN.tex:459-465

定位：01_GH_Product_EN.tex:459-465。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Emin Abdullaev | 2026-02-28

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019845 | OC-4600

The Rigidity Theorems for Self-Shrinkers in the Mean Curvature Flow | 708-7.tex:242-252

定位：708-7.tex:242-252。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Fagui Li; Yuhang Zhao | 2026-07-05

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019846 | OC-4601

The Rigidity Theorems for Self-Shrinkers in the Mean Curvature Flow | 708-7.tex:1091-1097

定位：708-7.tex:1091-1097。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Fagui Li; Yuhang Zhao | 2026-07-05

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019847 | OC-4609

Algebra of the Infrared with Curve-Valued Potential | main.tex:330-342

定位：main.tex:330-342。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Longfei Li | 2026-07-04

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019848 | OC-4610

Algebra of the Infrared with Curve-Valued Potential | main.tex:4419-4459

定位：main.tex:4419-4459。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Longfei Li | 2026-07-04

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019849 | OC-4614

Constructing the $\mathbf{2^{2^5} + 1 = F_5}$-gon: the first step, the 641-gon | Constructing_the_F_5-gon,_first_step.tex:157-160

定位：Constructing_the_F_5-gon,_first_step.tex:157-160。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Helmut Ruhland | 2026-07-04

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019850 | OC-4619

Hölder maps under Pfaffian constraints | heis18_arxiv.tex:541-544

定位：heis18_arxiv.tex:541-544。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Armin Schikorra | 2026-07-04

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019851 | OC-4629

On Determinacy for Cut and Choose Games of Uncountable Length | cutandchoose-arxiv.tex:906-906

定位：cutandchoose-arxiv.tex:906-906。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Maxwell Levine | 2026-07-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019852 | OC-4644

On Ramsey-type problems for paths and cycles with few colour changes | colour_change_arXiv.tex:156-158

定位：colour_change_arXiv.tex:156-158。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Peter Allen; Julia Böttcher; Dennis Clemens; Fabian Hamann; Jozef Skokan; Anusch Taraz | 2026-07-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019853 | OC-4645

On Ramsey-type problems for paths and cycles with few colour changes | colour_change_arXiv.tex:729-731

定位：colour_change_arXiv.tex:729-731。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Peter Allen; Julia Böttcher; Dennis Clemens; Fabian Hamann; Jozef Skokan; Anusch Taraz | 2026-07-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019854 | OC-4646

On Ramsey-type problems for paths and cycles with few colour changes | colour_change_arXiv.tex:733-735

定位：colour_change_arXiv.tex:733-735。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Peter Allen; Julia Böttcher; Dennis Clemens; Fabian Hamann; Jozef Skokan; Anusch Taraz | 2026-07-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_nd

V3-019855 | OC-467

First explicit reciprocity law for unitary Friedberg--Jacquet periods | main.tex:216-223

定位：main.tex:216-223。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Murilo Corato-Zanarella | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019856 | OC-4680

Polynomial mixing for polygonal side matchings | chord_swapping_on_surfaces.tex:478-480

定位：chord_swapping_on_surfaces.tex:478-480。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Renan Gross; Anđela Šarković | 2026-07-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019857 | OC-47

A family of Non-Weierstrass Semigroups | NonWeierstrassExamples.tex:486-489

定位：NonWeierstrassExamples.tex:486-489。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | David Eisenbud; Frank-Olaf Schreyer | 2026-02-28

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_sa

V3-019858 | OC-4708

Feigin-Semikhatov duality at the critical level | FEIGIN-SEMIKHATOV_DUALITY_AT_THE_CRITICAL_LEVEL.tex:2591-2597

定位：FEIGIN-SEMIKHATOV_DUALITY_AT_THE_CRITICAL_LEVEL.tex:2591-2597。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Thomas Creutzig; Xuanzhong Dai; Bailin Song | 2026-07-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019859 | OC-4726

An absolute bound for generalized Diophantine tuples over polynomial rings | dio.tex:197-207

定位：dio.tex:197-207。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Kin Ming Tsang; Chi Hoi Yip | 2026-07-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019860 | OC-473

Arazy-type decomposition theorem for bounded linear operators and commutators on the trace class | Commutator-on-C1-tex.tex:169-172

定位：Commutator-on-C1-tex.tex:169-172。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Jinghao Huang; Fedor Sukochev; Zhizheng Yu | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019861 | OC-474

The minimum spectral radius of tP_4-saturated graphs | main.tex:100-106

定位：main.tex:100-106。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Junxue Zhang; Liwen Zhang | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019862 | OC-4741

Actions of lattices in S-arithmetic groups on manifolds | main.tex:127-134

定位：main.tex:127-134。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Segev Gonen Cohen | 2026-07-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019863 | OC-4742

Actions of lattices in S-arithmetic groups on manifolds | main.tex:149-151

定位：main.tex:149-151。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Segev Gonen Cohen | 2026-07-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019864 | OC-4743

Actions of lattices in S-arithmetic groups on manifolds | main.tex:155-163

定位：main.tex:155-163。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Segev Gonen Cohen | 2026-07-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019865 | OC-4744

Actions of lattices in S-arithmetic groups on manifolds | main.tex:194-198

定位：main.tex:194-198。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Segev Gonen Cohen | 2026-07-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019866 | OC-4747

For edge-color-critical graphs, non-r-partite spectral extremal graphs are edge extremal | arXiv2ColorCritical.tex:57-62

定位：arXiv2ColorCritical.tex:57-62。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Suil O; Jiadong Wu | 2026-07-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019867 | OC-4754

On the Peres--Schlag orthogonal projection problem and Kakeya-type sets | On_the_Peres--Schlag_orthogonal_projection_problem__On_the_Peres--Schlag_orthogonal_projection_problem___and_Kakeya-type_sets.tex:432-437

定位：On_the_Peres--Schlag_orthogonal_projection_problem__On_the_Peres--Schlag_orthogonal_projection_problem___and_Kakeya-type_sets.tex:432-437。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Guo-Dong Hong; Chong-Wei Liang; Chun-Yen Shen | 2026-07-01

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: missing

V3-019868 | OC-4791

Tamagawa ratios and unbounded Selmer moments | main.tex:177-183

定位：main.tex:177-183。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Peter Koymans; Alexander Smith | 2026-06-30

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019869 | OC-4792

Tamagawa ratios and unbounded Selmer moments | main.tex:2025-2039

定位：main.tex:2025-2039。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Peter Koymans; Alexander Smith | 2026-06-30

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019870 | OC-4793

Tamagawa ratios and unbounded Selmer moments | main.tex:2740-2743

定位：main.tex:2740-2743。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Peter Koymans; Alexander Smith | 2026-06-30

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019871 | OC-480

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6573-6576

定位：QuadraticArXiv_1.tex:6573-6576。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019872 | OC-481

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6578-6581

定位：QuadraticArXiv_1.tex:6578-6581。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019873 | OC-483

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6588-6594

定位：QuadraticArXiv_1.tex:6588-6594。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019874 | OC-4831

L^p-form of the KNRS conjecture | main.tex:116-123

定位：main.tex:116-123。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Yuqi Zhao | 2026-06-29

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019875 | OC-4845

Finite-core Volterra reductions for a Weyl-positive Riemann phase kernel | rh_weyl_positive_draft.tex:697-704

定位：rh_weyl_positive_draft.tex:697-704。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Marvin B. Freedman | 2026-06-28

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019876 | OC-4846

Finite-core Volterra reductions for a Weyl-positive Riemann phase kernel | rh_weyl_positive_draft.tex:1115-1119

定位：rh_weyl_positive_draft.tex:1115-1119。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Marvin B. Freedman | 2026-06-28

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019877 | OC-4850

Handle decompositions and the 1-dimensional inputs skein lasagna module | main.tex:500-502

定位：main.tex:500-502。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Imogen Montague; Ian A. Sullivan | 2026-06-28

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019878 | OC-4857

Volume Law and Universality of Entanglement Entropy in Random Graph Fermi Systems | Manuscript_arxiv_2.tex:925-934

定位：Manuscript_arxiv_2.tex:925-934。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Saikat Sur | 2026-06-28

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019879 | OC-488

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6619-6622

定位：QuadraticArXiv_1.tex:6619-6622。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019880 | OC-489

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6626-6629

定位：QuadraticArXiv_1.tex:6626-6629。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019881 | OC-490

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6631-6634

定位：QuadraticArXiv_1.tex:6631-6634。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019882 | OC-491

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6636-6639

定位：QuadraticArXiv_1.tex:6636-6639。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019883 | OC-492

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6641-6644

定位：QuadraticArXiv_1.tex:6641-6644。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019884 | OC-493

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6649-6652

定位：QuadraticArXiv_1.tex:6649-6652。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019885 | OC-494

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6654-6657

定位：QuadraticArXiv_1.tex:6654-6657。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019886 | OC-495

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6660-6663

定位：QuadraticArXiv_1.tex:6660-6663。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019887 | OC-496

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6665-6668

定位：QuadraticArXiv_1.tex:6665-6668。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019888 | OC-497

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6672-6675

定位：QuadraticArXiv_1.tex:6672-6675。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019889 | OC-498

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6677-6680

定位：QuadraticArXiv_1.tex:6677-6680。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019890 | OC-499

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6682-6685

定位：QuadraticArXiv_1.tex:6682-6685。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019891 | OC-500

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6687-6690

定位：QuadraticArXiv_1.tex:6687-6690。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019892 | OC-501

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6697-6700

定位：QuadraticArXiv_1.tex:6697-6700。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019893 | OC-502

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6702-6705

定位：QuadraticArXiv_1.tex:6702-6705。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019894 | OC-504

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6713-6716

定位：QuadraticArXiv_1.tex:6713-6716。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019895 | OC-506

Quadratic irrational analogues of Ramanujan's series for $1/π$ | QuadraticArXiv_1.tex:6723-6726

定位：QuadraticArXiv_1.tex:6723-6726。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | John M. Campbell; Shaun Cooper; Dongxi Ye | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019896 | OC-508

On integrals of non-autonomous dynamical systems in finite characteristic | main.tex:1684-1694

定位：main.tex:1684-1694。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Nalini Joshi; Pieter Roffelsen | 2026-02-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019897 | OC-511

Generic flatness of the cohomology of thickenings | lc_free_2026_02_09.tex:219-239

定位：lc_free_2026_02_09.tex:219-239。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Yairon Cid-Ruiz; Anurag K. Singh | 2026-02-09

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019898 | OC-512

Cohomological rigidity of solvable Lie algebras of maximal ran | submission_version.tex:1159-1164

定位：submission_version.tex:1159-1164。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | B. A. Omirov; G. O. Solijanova; G. Kh. Urazmatov | 2026-02-09

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019899 | OC-546

Uniform spectral gaps for random hyperbolic surfaces with not many cusps | main.tex:236-241

定位：main.tex:236-241。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Yuxin He; Yunhui Wu; Yuhao Xue | 2026-02-09

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019900 | OC-552

Almost-primes in Sun's x^2+ny^2 conjecture | sunconj.tex:127-130

定位：sunconj.tex:127-130。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Songlin Han; Jinbo Yu | 2026-02-09

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019901 | OC-563

Minimal nonintegrable models with three-site interactions | minimal_nonintegrable_models_with_three_site_interactions.tex:263-268

定位：minimal_nonintegrable_models_with_three_site_interactions.tex:263-268。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Wen-Ming Fan; Kun Hao; Xiao-Hui Wang; Kun Zhang; Vladimir Korepin | 2026-02-08

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019902 | OC-567

Motivic invariants of moduli stacks of Higgs bundles and bundles with connections: results and speculations | MotivicClassesOfIrregularHiggs-6.tex:471-473

定位：MotivicClassesOfIrregularHiggs-6.tex:471-473。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Roman Fedorov; Alexander Soibelman; Yan Soibelman | 2026-02-07

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_sa

V3-019903 | OC-6

On the expected value of energy in groups | main.tex:1152-1157

定位：main.tex:1152-1157。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Marco Barbieri; Marusa Lekše; Andoni Zozaya | 2026-03-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019904 | OC-649

Strong local nondeterminism for stochastic time-fractional slow and fast diffusion equations | main.tex:873-878

定位：main.tex:873-878。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Le Chen; Cheuk Yin Lee; Panqiu Xia | 2026-02-05

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019905 | OC-7

On the expected value of energy in groups | main.tex:1536-1543

定位：main.tex:1536-1543。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Marco Barbieri; Marusa Lekše; Andoni Zozaya | 2026-03-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019906 | OC-70

Diamond Determinants and Somos Sequences | ddss-paper.tex:103-103

定位：ddss-paper.tex:103-103。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Nikolai Beluhov | 2026-02-27

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019907 | OC-71

Diamond Determinants and Somos Sequences | ddss-paper.tex:105-105

定位：ddss-paper.tex:105-105。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Nikolai Beluhov | 2026-02-27

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019908 | OC-72

Diamond Determinants and Somos Sequences | ddss-paper.tex:527-527

定位：ddss-paper.tex:527-527。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Nikolai Beluhov | 2026-02-27

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019909 | OC-729

Existence for Stable Rotating Star-Planet Systems | main.tex:1553-1555

定位：main.tex:1553-1555。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Hangsheng Chen | 2026-02-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019910 | OC-73

Diamond Determinants and Somos Sequences | ddss-paper.tex:531-531

定位：ddss-paper.tex:531-531。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Nikolai Beluhov | 2026-02-27

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019911 | OC-74

Diamond Determinants and Somos Sequences | ddss-paper.tex:539-539

定位：ddss-paper.tex:539-539。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Nikolai Beluhov | 2026-02-27

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019912 | OC-75

Diamond Determinants and Somos Sequences | ddss-paper.tex:541-541

定位：ddss-paper.tex:541-541。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Nikolai Beluhov | 2026-02-27

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019913 | OC-775

Unimodality of independence polynomials of two family of trees | main-15.tex:118-120

定位：main-15.tex:118-120。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Grace M. X. Li | 2026-03-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019914 | OC-787

Cohen-Macaulayness of squarefree powers of edge ideals of whisker graphs | 20.tex:298-308

定位：20.tex:298-308。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Rakesh Ghosh; S Selvaraja | 2026-03-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019915 | OC-788

Cohen-Macaulayness of squarefree powers of edge ideals of whisker graphs | 20.tex:1921-1933

定位：20.tex:1921-1933。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Rakesh Ghosh; S Selvaraja | 2026-03-03

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019916 | OC-8

Approximate message passing for block-structured ecological systems | main_final.tex:268-284

定位：main_final.tex:268-284。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Maxime Clenet; Mohammed-Younes Gueddari | 2026-03-02

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: cc_by_nc_sa

V3-019917 | OC-816

A step towards the Erdős-Rogers problem | er.tex:115-118

定位：er.tex:115-118。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Longma Du; Xinyu Hu; Ruilong Liu; Guanghui Wang | 2026-03-13

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019918 | OC-842

Construction of Local Arthur Packets for Metaplectic Groups and the Adams Conjecture | document.tex:364-366

定位：document.tex:364-366。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Jiahe Chen | 2026-03-12

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019919 | OC-846

Structural and Polynomial-Time Results on Core and Corona in Odd-Bicyclic Graphs | Bi-Cyclic.tex:1086-1090

定位：Bi-Cyclic.tex:1086-1090。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Kevin Pereyra | 2026-03-12

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019920 | OC-848

A characterization of graphs with $\a{\corona G}+\a{\core G}=2α(G)+1$ | core+corona=2a+1.tex:2409-2420

定位：core+corona=2a+1.tex:2409-2420。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Kevin Pereyra | 2026-03-12

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019921 | OC-849

Bohr sets in sumsets III: expanding difference sets and almost Bohr sets | Bohr_expanding.tex:3191-3193

定位：Bohr_expanding.tex:3191-3193。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Pierre-Yves Bienvenu; John T. Griesmer; Anh N. Le; Thái Hoàng Lê | 2026-03-11

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019922 | OC-860

Violating the All-or-Nothing Picture of Local Charges in Non-Hermitian Bosonic Chains | main.tex:1541-1543

定位：main.tex:1541-1543。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Mizuki Yamaguchi; Naoto Shiraishi | 2026-03-11

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019923 | OC-862

The Chow motive of LSV hyper-Kälher manifolds | L-S-V_H-K.tex:125-125

定位：L-S-V_H-K.tex:125-125。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Claudio Pedrini | 2026-03-11

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019924 | OC-899

On the Maximal Size of Irredundant Generating Sets in Lie Groups and Algebraic Groups | Redundancy_Rank.tex:224-227

定位：Redundancy_Rank.tex:224-227。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Tal Cohen; Itamar Vigdorovich | 2026-03-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019925 | OC-900

On the Maximal Size of Irredundant Generating Sets in Lie Groups and Algebraic Groups | Redundancy_Rank.tex:277-284

定位：Redundancy_Rank.tex:277-284。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Tal Cohen; Itamar Vigdorovich | 2026-03-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019926 | OC-909

Frobenius structure on rigid connections and arithmetic applications | theta-connection-1.4.tex:2455-2463

定位：theta-connection-1.4.tex:2455-2463。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Daxin Xu; Lingfei Yi | 2026-03-10

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019927 | OC-915

A Note on a Theorem of Apter | a_note_on_a_theorem_of_apter_v1.tex:248-250

定位：a_note_on_a_theorem_of_apter_v1.tex:248-250。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Rahman Mohammadpour; Otto Rajala; Sebastiano Thei | 2026-03-09

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019928 | OC-917

Hierarchical threshold structure in Max-Cut with geometric edge weights | LexiCut-arXiv.tex:698-700

定位：LexiCut-arXiv.tex:698-700。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Nevena Marić | 2026-03-09

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019929 | OC-924

Homogeneous ideals with minimal singularity thresholds | main.tex:1854-1856

定位：main.tex:1854-1856。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Benjamin Baily | 2026-03-09

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019930 | OC-925

Homogeneous ideals with minimal singularity thresholds | main.tex:1873-1875

定位：main.tex:1873-1875。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Benjamin Baily | 2026-03-09

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019931 | OC-937

The triplication method for constructing strong starters | triplication_gen_2026-arxiv.tex:2219-2224

定位：triplication_gen_2026-arxiv.tex:2219-2224。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Oleg Ogandzhanyants; Sergey Sadov; Margo Kondratieva | 2026-03-09

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019932 | OC-95

Cyclic sieving for a class of rectangular domino tableaux | csp-domino.tex:1144-1149

定位：csp-domino.tex:1144-1149。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Laura Colmenarejo; Bridget Eileen Tenner; Camryn E. Thompson | 2026-02-26

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019933 | OC-980

Rational Preperiodic Points of Quadratic Rational Maps over $\mathbb{Q}$ with Nonabelian Automorphism Groups | main.tex:134-136

定位：main.tex:134-136。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Hasan Bilgili; Mohammad Sadek | 2026-03-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: missing

V3-019934 | OC-985

Classical and irregular Hodge numbers | arxiv.tex:669-672

定位：arxiv.tex:669-672。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Yichen Qin; Dingxin Zhang | 2026-03-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019935 | OC-988

Hausdorff dimension of images and graphs of some random complex series | Random-series.tex:1010-1017

定位：Random-series.tex:1010-1017。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chun-Kit Lai; Ka-Sing Lau; Peng-Fei Zhang | 2026-03-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

V3-019936 | OC-989

Hausdorff dimension of images and graphs of some random complex series | Random-series.tex:1022-1030

定位：Random-series.tex:1022-1030。发布数据中的猜想候选；本卷仅提供论文与源码位置。

仅来源定位 | source_machine_label_real_open_conjecture | Chun-Kit Lai; Ka-Sing Lau; Peng-Fei Zhang | 2026-03-06

仅定位元数据，没有正文 | 原始来源 | metadata only; original license: arxiv_nonexclusive_distrib

C04 图论论文定位 | 372 条

V3-019937 | GAX-AX_3_coloring_diameter_2_quasipolynomial

Faster 3-colouring algorithm for graphs of diameter 3 | Problem 1

论文中定位：Problem 1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Carla Groenland; Hidde Koerts; Sophie Spirkl | 2026-01-19

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019938 | GAX-AX_4_colorability_fixed_surface_complexity

Three-coloring triangle-free graphs on surfaces IV. Bounding face sizes of 4-critical graphs | 4-colorability on Fixed Surfaces (Open Step in Thomassen's Program)

论文中定位：4-colorability on Fixed Surfaces (Open Step in Thomassen's Program)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdenek Dvorak; Daniel Kral; Robin Thomas | 2020-08-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019939 | GAX-AX_4_colorability_fixed_surface_polynomial_time

Three-coloring triangle-free graphs on surfaces I. Extending a coloring to a disk with one triangle | Open Problem: polynomial-time 4-colorability on fixed surfaces

论文中定位：Open Problem: polynomial-time 4-colorability on fixed surfaces；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdenek Dvorak; Dan Kral; Robin Thomas | 2016-03-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019940 | GAX-AX_5_coloring_p4_rp3_free_complexity

Complexity dichotomy for List-5-Coloring with a forbidden induced subgraph | Open Problem: 5-Coloring complexity for H-free graphs with a P4 component

论文中定位：Open Problem: 5-Coloring complexity for H-free graphs with a P4 component；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Sepehr Hajebi; Yanjia Li; Sophie Spirkl | 2022-08-30

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019941 | GAX-AX_active_clustering_clique_algorithm_optimal

Active clustering for labeling training data | Conjecture 5

论文中定位：Conjecture 5；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Quentin Lutz; Élie de Panafieu; Alex Scott; Maya Stein | 2021-10-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019942 | GAX-AX_additive_chi_approx_kk_minor_free

Additive non-approximability of chromatic number in proper minor-closed classes | Open question: better additive approximation for K_{4k_0+1}-minor-free graphs

论文中定位：Open question: better additive approximation for K_{4k_0+1}-minor-free graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Ken-ichi Kawarabayashi | 2017-07-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019943 | GAX-AX_additive_chi_approx_minor_closed_triangle_free

Additive non-approximability of chromatic number in proper minor-closed classes | Open question: additive approximation for triangle-free graphs

论文中定位：Open question: additive approximation for triangle-free graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Ken-ichi Kawarabayashi | 2017-07-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019944 | GAX-AX_annulus_digraph_winding_one_crossing_free

When all directed cycles have length three | Conjecture 1.2

论文中定位：Conjecture 1.2；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Paul Seymour | 2025-02-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019945 | GAX-AX_arithmetic_hyperbolic_surface_diameter_log_volume

Small diameters and generators for arithmetic lattices in $\mathrm{SL}_2(\mathbb{R})$ and certain Ramanujan graphs | Informal conjecture on the actual diameter

论文中定位：Informal conjecture on the actual diameter；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Raphael S. Steiner | 2022-07-26

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019946 | GAX-AX_arithmetic_k_cycle_removal_optimal_exponent

A polynomial bound for the arithmetic k-cycle removal lemma in vector spaces | Question 1.4

论文中定位：Question 1.4；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; László Miklós Lovász; Lisa Sauermann | 2018-09-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019947 | GAX-AX_avd_edge_coloring_probabilistic_barrier

Progress on the adjacent vertex distinguishing edge colouring conjecture | Informal Conjecture: probabilistic methods insufficient for the $\Delta+2$ bound

论文中定位：Informal Conjecture: probabilistic methods insufficient for the $\Delta+2$ bound；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Gwenaël Joret; William Lochet | 2020-07-22

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019948 | GAX-AX_avoidable_non_path_family_existence

Avoidable paths in graphs | Question 4.1

论文中定位：Question 4.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Oscar Defrain; Meike Hatzel; Jocelyn Thiebaut | 2019-08-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019949 | GAX-AX_avoidable_paths_two_disjoint_pk

Avoidable paths in graphs | Question 1.12

论文中定位：Question 1.12；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Oscar Defrain; Meike Hatzel; Jocelyn Thiebaut | 2019-08-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019950 | GAX-AX_behrend_sets_avoiding_mixed_sign_patterns

Triforce and Corners | Informal question on Behrend-style sets avoiding affine patterns with mixed-sign coefficients

论文中定位：Informal question on Behrend-style sets avoiding affine patterns with mixed-sign coefficients；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Ashwin Sah; Mehtaab Sawhney; David Stoner; Yufei Zhao | 2019-07-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019951 | GAX-AX_bicritical_snarks_almost_group_connected

Open problems of the 33rd Workshop on Cycles and Colourings | Problem 7.1

论文中定位：Problem 7.1；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | János Barát; Zdeněk Dvořák; Penny Haxell; František Kardoš; Borut Lužar; Alfréd Onderko; Jozef Rajník; Roman Soták; Nikolay Ulyanov | 2025-11-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019952 | GAX-AX_block_stable_random_graph_diameter_bound

Random graphs from a block-stable class | Informal Conjecture (diameter bound improvement)

论文中定位：Informal Conjecture (diameter bound improvement)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Colin McDiarmid; Alex Scott | 2016-05-16

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019953 | GAX-AX_bounded_clique_large_chi_consecutive_holes

Induced subgraphs of graphs with large chromatic number. X. Holes of specific residue | Conjecture 1.8

论文中定位：Conjecture 1.8；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour | 2018-12-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019954 | GAX-AX_bounded_clique_large_chi_consecutive_holes_extension

Induced subgraphs of graphs with large chromatic number. IV. Consecutive holes | Informal Conjecture (generalization of 1.3 to bounded clique number)

论文中定位：Informal Conjecture (generalization of 1.3 to bounded clique number)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour | 2018-02-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019955 | GAX-AX_bounded_degree_treewidth_induced_wall

On the tree-width of even-hole-free graphs | Conjecture 3

论文中定位：Conjecture 3；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Pierre Aboulker; Isolde Adler; Eun Jung Kim; Ni Luh Dewi Sintiari; Nicolas Trotignon | 2020-08-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019956 | GAX-AX_broom_nondominating_copy_no_scattering

Polynomial bounds for chromatic number VI. Adding a four-vertex path | Conjecture 2.3

论文中定位：Conjecture 2.3；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Alex Scott; Paul Seymour; Sophie Spirkl | 2023-03-22

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019957 | GAX-AX_c4_free_counting_lemma_characterization

Which graphs can be counted in C_4-free graphs? | Question 1.1

论文中定位：Question 1.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox; Benny Sudakov; Yufei Zhao | 2021-06-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019958 | GAX-AX_c4_free_counting_petersen_dodecahedron

Which graphs can be counted in C_4-free graphs? | Open Problem 2.17

论文中定位：Open Problem 2.17；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox; Benny Sudakov; Yufei Zhao | 2021-06-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019959 | GAX-AX_c5_free_triangle_removal_exponent

The regularity method for graphs with few 4-cycles | Optimality of the $o(n^{3/2})$ bound in Corollary 1.4

论文中定位：Optimality of the $o(n^{3/2})$ bound in Corollary 1.4；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox; Benny Sudakov; Yufei Zhao | 2021-09-25

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019960 | GAX-AX_c_closed_graphs_maximal_cliques_exponent

Finding Cliques in Social Networks: A New Distribution-Free Model | Open Problem (exact exponent for maximal cliques in c-closed graphs)

论文中定位：Open Problem (exact exponent for maximal cliques in c-closed graphs)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Tim Roughgarden; C. Seshadhri; Fan Wei; Nicole Wein | 2018-04-20

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019961 | GAX-AX_capture_time_higher_genus_bounds

Topological directions in Cops and Robbers | Capture time bounds for higher genus graphs

论文中定位：Capture time bounds for higher genus graphs；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Anthony Bonato; Bojan Mohar | 2018-04-22

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019962 | GAX-AX_capture_time_planar_linear_tightness

Topological directions in Cops and Robbers | Tightness of linear capture time bound for planar graphs

论文中定位：Tightness of linear capture time bound for planar graphs；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Anthony Bonato; Bojan Mohar | 2018-04-22

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019963 | GAX-AX_chi_bounded_not_polynomially_chi_bounded

Graphs of bounded cliquewidth are polynomially $χ$-bounded | Open problem: χ-bounded vs. polynomially χ-bounded

论文中定位：Open problem: χ-bounded vs. polynomially χ-bounded；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Michał Pilipczuk | 2020-07-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019964 | GAX-AX_chromatic_threshold_digraph_cycle_orientations

Orientations of cycles in digraphs of high chromatic number and high minimum out-degree | Question 1.2

论文中定位：Question 1.2；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Hidde Koerts; Benjamin Moore; Sophie Spirkl | 2025-03-25

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019965 | GAX-AX_chromatic_threshold_digraph_fixed_subdigraph

Orientations of cycles in digraphs of high chromatic number and high minimum out-degree | Question 4.1

论文中定位：Question 4.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Hidde Koerts; Benjamin Moore; Sophie Spirkl | 2025-03-25

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019966 | GAX-AX_chromatic_threshold_digraph_minimum_epsilon

Orientations of cycles in digraphs of high chromatic number and high minimum out-degree | Question 4.2

论文中定位：Question 4.2；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Hidde Koerts; Benjamin Moore; Sophie Spirkl | 2025-03-25

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019967 | GAX-AX_chromatic_threshold_odd_cycles_undirected

Orientations of cycles in digraphs of high chromatic number and high minimum out-degree | Question 1.1

论文中定位：Question 1.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Hidde Koerts; Benjamin Moore; Sophie Spirkl | 2025-03-25

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019968 | GAX-AX_circular_drawing_digraph_forbidden_characterization

When all directed cycles have the same weight | Open Question on Circular Drawing Characterisation

论文中定位：Open Question on Circular Drawing Characterisation；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Eli Berger; Daniel Carter; Paul Seymour | 2026-01-19

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019969 | GAX-AX_clean_classes_finite_family_characterization

Induced subgraphs and tree-decompositions VII. Basic obstructions in H-free graphs | Question: finite families yielding clean H-free classes

论文中定位：Question: finite families yielding clean H-free classes；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Tara Abrishami; Bogdan Alecu; Maria Chudnovsky; Sepehr Hajebi; Sophie Spirkl | 2023-11-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019970 | GAX-AX_clique_chromatic_dense_gnp_tight_constant

Clique coloring of dense random graphs | Informal Conjecture on Tight Constant for Clique Chromatic Number

论文中定位：Informal Conjecture on Tight Constant for Clique Chromatic Number；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Michael Krivelevich | 2017-11-05

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019971 | GAX-AX_clique_width_h1_h2_free_boundedness

Graph Isomorphism for (H_1,H_2)-free Graphs: An Almost Complete Dichotomy | Open Cases: Clique-Width Classification for (H_1,H_2)-Free Graphs

论文中定位：Open Cases: Clique-Width Classification for (H_1,H_2)-Free Graphs；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Nicolas Bousquet; Konrad K. Dabrowski; Matthew Johnson; Daniël Paulusma; Théo Pierron | 2019-09-03

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019972 | GAX-AX_cluster_vertex_deletion_2_approximation

Improved approximation algorithms for hitting 3-vertex paths | 2-Approximation Conjecture for Cluster-VD

论文中定位：2-Approximation Conjecture for Cluster-VD；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Samuel Fiorini; Gwenaël Joret; Oliver Schaudt | 2019-02-22

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019973 | GAX-AX_coarse_menger_bounded_genus

Asymptotic structure. V. The coarse Menger conjecture in bounded path-width | Informal conjecture: coarse Menger for bounded genus

论文中定位：Informal conjecture: coarse Menger for bounded genus；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Tung Nguyen; Alex Scott; Paul Seymour | 2025-09-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019974 | GAX-AX_coarse_menger_bounded_genus_surfaces

Asymptotic structure. VI. Distant paths across a disc | Conjecture 1.2 (Coarse Menger conjecture for surfaces)

论文中定位：Conjecture 1.2 (Coarse Menger conjecture for surfaces)；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Tung Nguyen; Alex Scott; Paul Seymour | 2025-09-08

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019975 | GAX-AX_coarse_menger_distance_2_paths

Asymptotic structure. IV. A counterexample to the weak coarse Menger conjecture | Open case: Coarse Menger Conjecture for c=2

论文中定位：Open case: Coarse Menger Conjecture for c=2；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Tung Nguyen; Alex Scott; Paul Seymour | 2025-08-20

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019976 | GAX-AX_coarse_treewidth_additive_quasi_isometry

Asymptotic structure. I. Coarse tree-width | Open question on Theorem 1.2 with $L=1$

论文中定位：Open question on Theorem 1.2 with $L=1$；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Tung Nguyen; Alex Scott; Paul Seymour | 2025-09-05

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019977 | GAX-AX_coherent_ideal_forest_filleting

Pure pairs. II. Excluding all subdivisions of a graph | Conjecture 2.4

论文中定位：Conjecture 2.4；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Alex Scott; Paul Seymour; Sophie Spirkl | 2020-06-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019978 | GAX-AX_complete_topological_graph_linear_disjoint_edges

A structure theorem for pseudo-segments and its applications | Optimal disjoint-edges bound in complete simple topological graphs

论文中定位：Optimal disjoint-edges bound in complete simple topological graphs；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Janos Pach; Andrew Suk | 2023-12-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019979 | GAX-AX_cone_crossing_number_2_page_redrawing

The crossing number of the cone of a graph | Problem 4

论文中定位：Problem 4；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Carlos A. Alfaro; Alan Arroyo; Marek Derunár; Bojan Mohar | 2016-08-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019980 | GAX-AX_cone_crossing_number_minimum_function

The crossing number of the cone of a graph | Problem 1

论文中定位：Problem 1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Carlos A. Alfaro; Alan Arroyo; Marek Derunár; Bojan Mohar | 2016-08-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019981 | GAX-AX_cone_crossing_number_simple_graphs_asymptotics

The crossing number of the cone of a graph | Conjecture on $f_s(k)$ asymptotics

论文中定位：Conjecture on $f_s(k)$ asymptotics；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Carlos A. Alfaro; Alan Arroyo; Marek Derunár; Bojan Mohar | 2016-08-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019982 | GAX-AX_connectivity_code_torus_product_cycles

Connectivity Graph-Codes | Open Problem: m(C_t × C_s) for all t, s

论文中定位：Open Problem: m(C_t × C_s) for all t, s；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon | 2023-09-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019983 | GAX-AX_constricting_hole_lengths_bounded_gaps

Induced subgraphs of graphs with large chromatic number. IV. Consecutive holes | Conjecture 1.5

论文中定位：Conjecture 1.5；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour | 2018-02-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019984 | GAX-AX_constricting_hole_lengths_density_zero

Induced subgraphs of graphs with large chromatic number. IV. Consecutive holes | Problem 1.6

论文中定位：Problem 1.6；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour | 2018-02-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019985 | GAX-AX_controlled_triangle_free_chromatic_4_hole

Induced subgraphs of graphs with large chromatic number. IV. Consecutive holes | Informal Question (on 2.1 for $\ell = 4$)

论文中定位：Informal Question (on 2.1 for $\ell = 4$)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour | 2018-02-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019986 | GAX-AX_convex_drawings_vs_pseudocircular_drawings

Topological Drawings meet Classical Theorems from Convex Geometry | Open problem: convex drawings vs. pseudocircular drawings

论文中定位：Open problem: convex drawings vs. pseudocircular drawings；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Helena Bergold; Stefan Felsner; Manfred Scheucher; Felix Schröder; Raphael Steiner | 2024-07-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019987 | GAX-AX_convex_unit_distance_realization_gk

Convexly independent subsets of Minkowski sums of convex polygons | Open Question on Convex Unit Distance Realization of Graphs G_k

论文中定位：Open Question on Convex Unit Distance Realization of Graphs G_k；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Mateusz Skomra; Stéphan Thomassé | 2021-06-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019988 | GAX-AX_cop_number_genus_sqrt_g_growth

Notes on Cops and Robber game on graphs | Conjecture 8

论文中定位：Conjecture 8；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Bojan Mohar | 2017-10-31

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019989 | GAX-AX_cop_number_outerplanar_classification

Topological directions in Cops and Robbers | Classification of outerplanar graphs by cop number

论文中定位：Classification of outerplanar graphs by cop number；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Anthony Bonato; Bojan Mohar | 2018-04-22

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019990 | GAX-AX_cop_number_planar_classification

Topological directions in Cops and Robbers | Classification of planar graphs by cop number

论文中定位：Classification of planar graphs by cop number；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Anthony Bonato; Bojan Mohar | 2018-04-22

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019991 | GAX-AX_cop_number_riemannian_surfaces_area_supremum

Notes on Cops and Robber game on graphs | Question (Riemannian surfaces): supremum of cop number for geometrically bounded surfaces

论文中定位：Question (Riemannian surfaces): supremum of cop number for geometrically bounded surfaces；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Bojan Mohar | 2017-10-31

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019992 | GAX-AX_cop_number_riemannian_surfaces_constant_curvature

Notes on Cops and Robber game on graphs | Question (Riemannian surfaces): constant curvature surfaces

论文中定位：Question (Riemannian surfaces): constant curvature surfaces；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Bojan Mohar | 2017-10-31

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019993 | GAX-AX_cop_number_riemannian_surfaces_cop_convergence

Notes on Cops and Robber game on graphs | Question (Riemannian surfaces): cop converging to robber

论文中定位：Question (Riemannian surfaces): cop converging to robber；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Bojan Mohar | 2017-10-31

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019994 | GAX-AX_cop_number_riemannian_surfaces_genus_bounded

Notes on Cops and Robber game on graphs | Question (Riemannian surfaces): cop number bounded by genus

论文中定位：Question (Riemannian surfaces): cop number bounded by genus；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Bojan Mohar | 2017-10-31

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019995 | GAX-AX_cop_number_riemannian_surfaces_worst_genus

Notes on Cops and Robber game on graphs | Question (Riemannian surfaces): worst surfaces of fixed genus

论文中定位：Question (Riemannian surfaces): worst surfaces of fixed genus；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Bojan Mohar | 2017-10-31

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019996 | GAX-AX_cop_number_sublinear_treedepth

Cops and robber in graphs with bounded vertex cover number | Problem 9

论文中定位：Problem 9；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Prosenjit Bose; Louis Esperet; Jędrzej Hodor; Gwenaël Joret; Piotr Micek; Clément Rambaud | 2026-02-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019997 | GAX-AX_corners_popular_difference_abelian_groups

Triforce and Corners | Informal conjecture on extending Mandache's Theorem 1.1 to all abelian groups

论文中定位：Informal conjecture on extending Mandache's Theorem 1.1 to all abelian groups；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Ashwin Sah; Mehtaab Sawhney; David Stoner; Yufei Zhao | 2019-07-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019998 | GAX-AX_correlation_polytope_extension_complexity_treewidth

Extension Complexity of the Correlation Polytope | Conjecture 2

论文中定位：Conjecture 2；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Pierre Aboulker; Samuel Fiorini; Tony Huynh; Marco Macchia; Johanna Seif | 2018-10-18

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-019999 | GAX-AX_corruption_detection_expansion_threshold

Distributed Corruption Detection in Networks | Open Problem: Expansion vs. Corruption Detection

论文中定位：Open Problem: Expansion vs. Corruption Detection；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Elchanan Mossel; Robin Pemantle | 2020-03-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020000 | GAX-AX_counting_perfect_matchings_alpha_2_hard

Exact Matching in Graphs of Bounded Independence Number | Conjecture 3

论文中定位：Conjecture 3；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Nicolas El Maalouly; Raphael Steiner | 2022-07-13

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020001 | GAX-AX_cover_free_families_optimal_size

Lower bounds for graph reconstruction with maximal independent set queries | Problem 5.1 (cover-free families)

论文中定位：Problem 5.1 (cover-free families)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Lukas Michel; Alex Scott | 2024-04-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020002 | GAX-AX_crossing_critical_max_degree_optimal_bounds

Bounded maximum degree conjecture holds precisely for c-crossing-critical graphs with $c \leq 12$ | Optimal degree bounds D_c for small c

论文中定位：Optimal degree bounds D_c for small c；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Drago Bokal; Zdeněk Dvořák; Petr Hliněný; Jesús Leaños; Bojan Mohar; Tilo Wiedera | 2021-05-05

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020003 | GAX-AX_cube_complex_boundary_distance_rigidity

Boundary rigidity of 3D CAT(0) cube complexes | Question 1

论文中定位：Question 1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | John Haslegrave; Alex Scott; Youri Tamitegama; Jane Tan | 2023-09-08

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020004 | GAX-AX_cubic_2_homogeneous_coloring_4_colors

Open problems of the 33rd Workshop on Cycles and Colourings | Problem 5.1

论文中定位：Problem 5.1；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | János Barát; Zdeněk Dvořák; Penny Haxell; František Kardoš; Borut Lužar; Alfréd Onderko; Jozef Rajník; Roman Soták; Nikolay Ulyanov | 2025-11-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020005 | GAX-AX_cubic_2_homogeneous_coloring_finite_exceptions

Open problems of the 33rd Workshop on Cycles and Colourings | Problem 5.2

论文中定位：Problem 5.2；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | János Barát; Zdeněk Dvořák; Penny Haxell; František Kardoš; Borut Lužar; Alfréd Onderko; Jozef Rajník; Roman Soták; Nikolay Ulyanov | 2025-11-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020006 | GAX-AX_cubic_many_3_edge_colorings_reflexive

Reflexive coloring complexes for 3-edge-colorings of cubic graphs | Informal Conjecture on reflexive graphs and coloring abundance

论文中定位：Informal Conjecture on reflexive graphs and coloring abundance；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Fiachra Knox; Bojan Mohar; Nathan Singer | 2020-04-14

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020007 | GAX-AX_cubic_truncation_strong_6_edge_coloring

Open problems of the 33rd Workshop on Cycles and Colourings | Problem 4.1

论文中定位：Problem 4.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | János Barát; Zdeněk Dvořák; Penny Haxell; František Kardoš; Borut Lužar; Alfréd Onderko; Jozef Rajník; Roman Soták; Nikolay Ulyanov | 2025-11-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020008 | GAX-AX_cycles_union_k4_copies_4_colorable

Open problems of the 33rd Workshop on Cycles and Colourings | Problem 3.1

论文中定位：Problem 3.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | János Barát; Zdeněk Dvořák; Penny Haxell; František Kardoš; Borut Lužar; Alfréd Onderko; Jozef Rajník; Roman Soták; Nikolay Ulyanov | 2025-11-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020009 | GAX-AX_cyclic_clique_minor_min_degree_bound

Lollipops, dense cycles and chords | Question 1.2

论文中定位：Question 1.2；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Beatriz Martins; Stéphan Thomassé; Nicolas Trotignon | 2025-10-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020010 | GAX-AX_degenerate_graphs_flexibility_d_plus_1_lists

List coloring with requests | Problem 1

论文中定位：Problem 1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Sergey Norin; Luke Postle | 2018-11-17

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020011 | GAX-AX_degenerate_graphs_polynomial_kempe_sequences

Kempe changes in degenerate graphs | Polynomial Kempe sequences for degenerate graphs

论文中定位：Polynomial Kempe sequences for degenerate graphs；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Vincent Delecroix; Clément Legrand-Duchesne | 2021-12-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020012 | GAX-AX_degenerate_kr_free_extremal_order_exponential

Coloring small locally sparse degenerate graphs and related problems | Problem 6.3

论文中定位：Problem 6.3；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Domagoj Bradač; Jacob Fox; Raphael Steiner; Benny Sudakov; Shengtong Zhang | 2026-01-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020013 | GAX-AX_degenerate_kr_free_fractional_chromatic_max

Coloring small locally sparse degenerate graphs and related problems | Problem 4.1

论文中定位：Problem 4.1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Domagoj Bradač; Jacob Fox; Raphael Steiner; Benny Sudakov; Shengtong Zhang | 2026-01-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020014 | GAX-AX_degenerate_kr_free_fractional_chromatic_sublinear

Coloring small locally sparse degenerate graphs and related problems | Problem 6.4

论文中定位：Problem 6.4；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Domagoj Bradač; Jacob Fox; Raphael Steiner; Benny Sudakov; Shengtong Zhang | 2026-01-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020015 | GAX-AX_degenerate_triangle_free_subexponential_order_coloring

Coloring small locally sparse degenerate graphs and related problems | Problem 6.1

论文中定位：Problem 6.1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Domagoj Bradač; Jacob Fox; Raphael Steiner; Benny Sudakov; Shengtong Zhang | 2026-01-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020016 | GAX-AX_degree_bounded_classes_c4_free_polynomial

Induced C_4-free subgraphs with large average degree | Polynomial bound in k for C4-free subgraphs

论文中定位：Polynomial bound in k for C4-free subgraphs；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Xiying Du; António Girão; Zach Hunter; Rose McCarty; Alex Scott | 2023-11-01

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020017 | GAX-AX_degree_bounded_classes_polynomial_bounding_function

Induced C_4-free subgraphs with large average degree | Polynomial degree-bounding function for hereditary degree-bounded classes

论文中定位：Polynomial degree-bounding function for hereditary degree-bounded classes；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Xiying Du; António Girão; Zach Hunter; Rose McCarty; Alex Scott | 2023-11-01

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020018 | GAX-AX_digraph_forward_connected_pairs_approximation_ratio

Temporalizing digraphs via linear-size balanced bi-trees | Problem 1

论文中定位：Problem 1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Stéphane Bessy; Stéphan Thomassé; Laurent Viennot | 2024-01-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020019 | GAX-AX_digraph_forward_connected_pairs_request_approximation

Temporalizing digraphs via linear-size balanced bi-trees | Problem 4

论文中定位：Problem 4；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Stéphane Bessy; Stéphan Thomassé; Laurent Viennot | 2024-01-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020020 | GAX-AX_digraph_left_maximal_dfs_tree_complexity

Temporalizing digraphs via linear-size balanced bi-trees | Problem 3

论文中定位：Problem 3；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Stéphane Bessy; Stéphan Thomassé; Laurent Viennot | 2024-01-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020021 | GAX-AX_digraph_strong_balanced_bitree_linear_constant

Temporalizing digraphs via linear-size balanced bi-trees | Problem 2

论文中定位：Problem 2；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Stéphane Bessy; Stéphan Thomassé; Laurent Viennot | 2024-01-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020022 | GAX-AX_digraph_strong_forward_cover_log_n

Temporalizing digraphs via linear-size balanced bi-trees | Conjecture 1

论文中定位：Conjecture 1；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Stéphane Bessy; Stéphan Thomassé; Laurent Viennot | 2024-01-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020023 | GAX-AX_disjoint_paths_stability_two_np_complete

Disjoint paths in unions of tournaments | Informal Conjecture (vertex-disjoint paths, stability number two)

论文中定位：Informal Conjecture (vertex-disjoint paths, stability number two)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Alex Scott; Paul Seymour | 2018-12-23

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020024 | GAX-AX_distributed_coloring_sparse_sublinear_rounds

Distributed coloring in sparse graphs with fewer colors | Open problem on sublinear round complexity

论文中定位：Open problem on sublinear round complexity；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Pierre Aboulker; Marthe Bonamy; Nicolas Bousquet; Louis Esperet | 2018-12-19

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020025 | GAX-AX_distributed_delta_list_coloring_randomized_rounds

Distributed coloring in sparse graphs with fewer colors | Question on randomized list-coloring round complexity

论文中定位：Question on randomized list-coloring round complexity；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Pierre Aboulker; Marthe Bonamy; Nicolas Bousquet; Louis Esperet | 2018-12-19

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020026 | GAX-AX_divisible_subdivision_clique_minor_f_growth

Divisible subdivisions | Open Problem (asymptotic behavior of f(H,q))

论文中定位：Open Problem (asymptotic behavior of f(H,q))；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Michael Krivelevich | 2021-06-29

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020027 | GAX-AX_divisible_subdivision_g_q_linear

Divisible subdivisions | Informal Conjecture (linearity of g(q))

论文中定位：Informal Conjecture (linearity of g(q))；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Michael Krivelevich | 2021-06-29

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020028 | GAX-AX_domination_2_independence_tied_classes

Characterization of sparse monotone graph classes with bounded domination-to-2-independence ratio | Problem 2

论文中定位：Problem 2；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Zdeněk Dvořák; Lukas Michel; David Mikšaník | 2026-01-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020029 | GAX-AX_domination_packing_ratio_bounded_2_independence

Characterization of sparse monotone graph classes with bounded domination-to-2-independence ratio | Problem 1

论文中定位：Problem 1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Zdeněk Dvořák; Lukas Michel; David Mikšaník | 2026-01-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020030 | GAX-AX_domination_packing_ratio_bounded_characterization

On graph classes with constant domination-packing ratio | Question 9.1

论文中定位：Question 9.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Mónika Csikós; Anna Gujgiczer; Yelena Yuditsky | 2025-03-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020031 | GAX-AX_domination_packing_ratio_bounded_given_class

On graph classes with constant domination-packing ratio | Question 1.1

论文中定位：Question 1.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Mónika Csikós; Anna Gujgiczer; Yelena Yuditsky | 2025-03-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020032 | GAX-AX_ell_holed_graphs_structure_small_ell

Graphs with all holes the same length | Open Problem: characterisation of ℓ-holed graphs for $\ell \in \{4, 5, 6\}$

论文中定位：Open Problem: characterisation of ℓ-holed graphs for $\ell \in \{4, 5, 6\}$；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Linda Cook; Jake Horsfield; Myriam Preissmann; Cléophée Robin; Paul Seymour; Ni Luh Dewi Sintiari; Nicolas Trotignon; Kristina Vušković | 2023-12-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020033 | GAX-AX_epsilon_t_nets_size_computation

The $ε$-t-Net Problem | The ε-t-Net Problem

论文中定位：The ε-t-Net Problem；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Bruno Jartoux; Chaya Keller; Shakhar Smorodinsky; Yelena Yuditsky | 2020-03-16

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020034 | GAX-AX_erdos_ginzburg_ziv_constant_fixed_k

Erdős-Ginzburg-Ziv constants by avoiding three-term arithmetic progressions | Open Question on s((Z/kZ)^n) for fixed k

论文中定位：Open Question on s((Z/kZ)^n) for fixed k；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Lisa Sauermann | 2018-04-17

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020035 | GAX-AX_erdos_posa_planar_minor_constant_polynomial

A tight Erdős-Pósa function for planar minors | Open Problem (constant c vs. |H|)

论文中定位：Open Problem (constant c vs. |H|)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Wouter Cames van Batenburg; Tony Huynh; Gwenaël Joret; Jean-Florent Raymond | 2019-10-23

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020036 | GAX-AX_erdos_szekeres_collinear_variant_ell_dependence

Big line or big convex polygon | Open Problem (Introduction)

论文中定位：Open Problem (Introduction)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox; Xiaoyu He; Dhruv Mubayi; Andrew Suk; Jacques Verstraete | 2024-05-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020037 | GAX-AX_euclidean_ramsey_diameter_free_bound

Lines in Euclidean Ramsey theory | Open problem: removing diameter dependence from Theorem 1.1

论文中定位：Open problem: removing diameter dependence from Theorem 1.1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox | 2018-03-20

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020038 | GAX-AX_euclidean_ramsey_l3_vs_lm

Lines in Euclidean Ramsey theory | Open problem: $E^n \not\to (\ell_3, \ell_m)$ for large m

论文中定位：Open problem: $E^n \not\to (\ell_3, \ell_m)$ for large m；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox | 2018-03-20

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020039 | GAX-AX_euclidean_ramsey_unit_distance_vs_copy

Lines in Euclidean Ramsey theory | Question 1.1

论文中定位：Question 1.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox | 2018-03-20

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020040 | GAX-AX_even_hole_free_bounded_degree_treewidth

On the tree-width of even-hole-free graphs | Conjecture 2

论文中定位：Conjecture 2；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Pierre Aboulker; Isolde Adler; Eun Jung Kim; Ni Luh Dewi Sintiari; Nicolas Trotignon | 2020-08-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020041 | GAX-AX_even_hole_free_chordal_modulator

Induced subgraphs and tree decompositions VIII. Excluding a forest in (theta, prism)-free graphs | Informal Conjecture (chordal modulator characterization for even-hole-free graphs)

论文中定位：Informal Conjecture (chordal modulator characterization for even-hole-free graphs)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Tara Abrishami; Bogdan Alecu; Maria Chudnovsky; Sepehr Hajebi; Sophie Spirkl | 2023-01-28

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020042 | GAX-AX_even_hole_free_coloring_polynomial

Tree Independence Number IV. Even-hole-free Graphs | Open Problem: Polynomial-time Coloring on even-hole-free graphs

论文中定位：Open Problem: Polynomial-time Coloring on even-hole-free graphs；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Peter Gartland; Sepehr Hajebi; Daniel Lokshtanov; Sophie Spirkl | 2024-07-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020043 | GAX-AX_even_hole_free_independent_set_complexity

Maximum independent sets in (pyramid, even hole)-free graphs | Open problem: MIS complexity in even-hole-free graphs

论文中定位：Open problem: MIS complexity in even-hole-free graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Stéphan Thomassé; Nicolas Trotignon; Kristina Vušković | 2019-12-24

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020044 | GAX-AX_even_hole_free_independent_set_polynomial

Tree Independence Number IV. Even-hole-free Graphs | Open Problem: Polynomial-time MWIS on even-hole-free graphs

论文中定位：Open Problem: Polynomial-time MWIS on even-hole-free graphs；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Peter Gartland; Sepehr Hajebi; Daniel Lokshtanov; Sophie Spirkl | 2024-07-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020045 | GAX-AX_even_hole_k4_diamond_free_treewidth

(Theta, triangle)-free and (even hole, K_4)-free graphs. Part 2 : bounds on treewidth | Open Question: (even hole, K4, diamond)-free treewidth/cliquewidth

论文中定位：Open Question: (even hole, K4, diamond)-free treewidth/cliquewidth；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marcin Pilipczuk; Ni Luh Dewi Sintiari; Stéphan Thomassé; Nicolas Trotignon | 2020-10-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020046 | GAX-AX_f3n_subspace_free_sets_exponential_base

Popular progression differences in vector spaces II | Informal Question (correct exponential constant in r(n,m))

论文中定位：Informal Question (correct exponential constant in r(n,m))；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Huy Tuan Pham | 2019-11-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020047 | GAX-AX_fat_minor_quasi_isometry_trees_planar

Asymptotic structure. III. Excluding a fat tree | Informal question: validity of Conjecture 1.2 for trees and planar graphs

论文中定位：Informal question: validity of Conjecture 1.2 for trees and planar graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Tung Nguyen; Alex Scott; Paul Seymour | 2025-09-13

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020048 | GAX-AX_flip_distance_complexity_binary_trees_rotation

Shortest paths on polymatroids and hypergraphic polytopes | Complexity of Rotation Distance Between Binary Search Trees

论文中定位：Complexity of Rotation Distance Between Binary Search Trees；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jean Cardinal; Raphael Steiner | 2023-11-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020049 | GAX-AX_flip_distance_complexity_rectangulations

Shortest paths on polymatroids and hypergraphic polytopes | Complexity of Flip Distance Between Rectangulations

论文中定位：Complexity of Flip Distance Between Rectangulations；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jean Cardinal; Raphael Steiner | 2023-11-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020050 | GAX-AX_fo_minimization_approximation_bounded_expansion_ptas

Approximation metatheorems for classes with bounded expansion | Problem 6

论文中定位：Problem 6；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák | 2021-10-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020051 | GAX-AX_forest_of_lanterns_pervasive

Induced subgraphs of graphs with large chromatic number. V. Chandeliers and strings | Conjecture: every forest of lanterns is pervasive

论文中定位：Conjecture: every forest of lanterns is pervasive；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Alex Scott; Paul Seymour | 2021-05-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020052 | GAX-AX_frozen_coloring_glauber_delta_o_n

Frozen $(Δ+1)$-colourings of bounded degree graphs | Informal: Δ = o(n) condition not necessary for Corollary 2

论文中定位：Informal: Δ = o(n) condition not necessary for Corollary 2；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Nicolas Bousquet; Guillem Perarnau | 2018-11-30

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020053 | GAX-AX_frozen_coloring_probability_exponent

Frozen $(Δ+1)$-colourings of bounded degree graphs | Informal: Exponent in Theorem 1 not optimal

论文中定位：Informal: Exponent in Theorem 1 not optimal；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Nicolas Bousquet; Guillem Perarnau | 2018-11-30

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020054 | GAX-AX_generalized_coloring_number_cc_characterization

On generalized choice and coloring numbers | Open Problem: Characterization of parameters satisfying (CC)

论文中定位：Open Problem: Characterization of parameters satisfying (CC)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Jakub Pekárek; Jean-Sébastien Sereni | 2019-02-26

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020055 | GAX-AX_gnm_subgraph_moderate_deviations_sparse

Moderate deviations of subgraph counts in the Erdős-Rényi random graphs $G(n,m)$ and $G(n,p)$ | Remark 1.2 (Open Problem on Sparse Densities)

论文中定位：Remark 1.2 (Open Problem on Sparse Densities)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Christina Goldschmidt; Simon Griffiths; Alex Scott | 2020-02-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020056 | GAX-AX_gnp_chromatic_concentration_lower_bound

On the concentration of the chromatic number of random graphs | Open problem on lower bound for concentration interval

论文中定位：Open problem on lower bound for concentration interval；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott | 2017-10-18

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020057 | GAX-AX_gnp_h_free_clique_count_threshold

Many cliques in H-free subgraphs of random graphs | Motivating Question (Introduction)

论文中定位：Motivating Question (Introduction)；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Alexandr Kostochka; Clara Shikhelman | 2017-11-19

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020058 | GAX-AX_graph_codes_cube_length_interval_vectors

Graph-codes | Open problem: binary vectors with cube-length forbidden symmetric difference

论文中定位：Open problem: binary vectors with cube-length forbidden symmetric difference；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon | 2023-02-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020059 | GAX-AX_graph_codes_even_edges_density_vanishes

Graph-codes | Question 1.1

论文中定位：Question 1.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon | 2023-02-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020060 | GAX-AX_graph_codes_k4_density_vanishes

Graph-codes | Informal question: is $d_{K_4}(n) = o(1)$?

论文中定位：Informal question: is $d_{K_4}(n) = o(1)$?；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon | 2023-02-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020061 | GAX-AX_graph_codes_linear_even_h_bounds

Graph-codes | Open problem: tighter bounds for linear graph-codes

论文中定位：Open problem: tighter bounds for linear graph-codes；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon | 2023-02-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020062 | GAX-AX_graph_codes_odd_intersection_edge_coloring

Graph-codes | Open problem: minimum edge-coloring with odd intersection property

论文中定位：Open problem: minimum edge-coloring with odd intersection property；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon | 2023-02-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020063 | GAX-AX_graph_isomorphism_fpt_clique_width

Graph Isomorphism for (H_1,H_2)-free Graphs: An Almost Complete Dichotomy | Open Question: FPT Algorithm for Graph Isomorphism Parameterized by Clique-Width

论文中定位：Open Question: FPT Algorithm for Graph Isomorphism Parameterized by Clique-Width；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Nicolas Bousquet; Konrad K. Dabrowski; Matthew Johnson; Daniël Paulusma; Théo Pierron | 2019-09-03

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020064 | GAX-AX_graph_isomorphism_h1_h2_free_dichotomy

Graph Isomorphism for (H_1,H_2)-free Graphs: An Almost Complete Dichotomy | Research Question (Introduction)

论文中定位：Research Question (Introduction)；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Nicolas Bousquet; Konrad K. Dabrowski; Matthew Johnson; Daniël Paulusma; Théo Pierron | 2019-09-03

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020065 | GAX-AX_graph_reconstruction_mis_queries_adaptive_deterministic

Lower bounds for graph reconstruction with maximal independent set queries | Problem 5.3 (adaptive vs non-adaptive gap)

论文中定位：Problem 5.3 (adaptive vs non-adaptive gap)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Lukas Michel; Alex Scott | 2024-04-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020066 | GAX-AX_graph_reconstruction_mis_queries_randomized_nonadaptive

Lower bounds for graph reconstruction with maximal independent set queries | Problem 5.2 (randomised non-adaptive optimality)

论文中定位：Problem 5.2 (randomised non-adaptive optimality)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Lukas Michel; Alex Scott | 2024-04-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020067 | GAX-AX_grid_ramsey_equals_hypergraph_clique_star

Hypergraph Ramsey numbers of cliques versus stars | Informal Conjecture: r(K^(3)_4, S^(3)_n) and gr(G_{2x2}, K_n) have the same order

论文中定位：Informal Conjecture: r(K^(3)_4, S^(3)_n) and gr(G_{2x2}, K_n) have the same order；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox; Xiaoyu He; Dhruv Mubayi; Andrew Suk; Jacques Verstraete | 2022-10-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020068 | GAX-AX_grid_ramsey_rectangle_vs_clique

Hypergraph Ramsey numbers of cliques versus stars | Grid Ramsey Problem (gr(G_{2x2}, K_n))

论文中定位：Grid Ramsey Problem (gr(G_{2x2}, K_n))；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox; Xiaoyu He; Dhruv Mubayi; Andrew Suk; Jacques Verstraete | 2022-10-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020069 | GAX-AX_grundy_coloring_biclique_free_fpt

Grundy Coloring & friends, Half-Graphs, Bicliques | FPT on $K_{t,t}$-free graphs for Grundy Coloring

论文中定位：FPT on $K_{t,t}$-free graphs for Grundy Coloring；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Pierre Aboulker; Édouard Bonnet; Eun Jung Kim; Florian Sikora | 2020-01-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020070 | GAX-AX_grundy_coloring_half_graph_free_fpt

Grundy Coloring & friends, Half-Graphs, Bicliques | FPT on $H_{t,t}$-free graphs for Grundy Coloring and b-Chromatic Core

论文中定位：FPT on $H_{t,t}$-free graphs for Grundy Coloring and b-Chromatic Core；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Pierre Aboulker; Édouard Bonnet; Eun Jung Kim; Florian Sikora | 2020-01-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020071 | GAX-AX_h_free_clique_maximizers_polynomial_error

H-free subgraphs of dense graphs maximizing the number of cliques and their blow-ups | Informal Conjecture (error term in Proposition 1.4)

论文中定位：Informal Conjecture (error term in Proposition 1.4)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Clara Shikhelman | 2017-06-18

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020072 | GAX-AX_h_free_coloring_dichotomy_all_k

List-k-Coloring H-free graphs for all $k>4$ | Full dichotomy for k-Coloring on H-free graphs ($k\geq 3$)

论文中定位：Full dichotomy for k-Coloring on H-free graphs ($k\geq 3$)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Sepehr Hajebi; Sophie Spirkl | 2023-11-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020073 | GAX-AX_hat_guessing_number_complete_bipartite

The hat guessing number of graphs | Open question: exact value of $HG(K_{n,n})$

论文中定位：Open question: exact value of $HG(K_{n,n})$；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Omri Ben-Eliezer; Chong Shangguan; Itzhak Tamo | 2020-01-15

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020074 | GAX-AX_hat_guessing_number_degree_degeneracy_bounds

The hat guessing number of graphs | Problem 1.4

论文中定位：Problem 1.4；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Omri Ben-Eliezer; Chong Shangguan; Itzhak Tamo | 2020-01-15

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020075 | GAX-AX_hat_guessing_universal_vertex_addition

On the Hat Guessing Number of Graphs | Problem on Hat Guessing Number After Adding a Universal Vertex

论文中定位：Problem on Hat Guessing Number After Adding a Universal Vertex；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Jeremy Chizewer | 2021-07-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020076 | GAX-AX_hereditary_tau_d_bounded_implies_polynomial_esperet

Polynomial bounds for chromatic number. V. Excluding a tree of radius two and a complete multipartite graph | Question: Polynomial bound analogue of Esperet's conjecture for τ_d

论文中定位：Question: Polynomial bound analogue of Esperet's conjecture for τ_d；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour | 2023-01-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020077 | GAX-AX_heroes_complete_multipartite_delta_1_2_2

Heroes in oriented complete multipartite graphs | Question 1.4

论文中定位：Question 1.4；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Pierre Aboulker; Guillaume Aubian; Pierre Charbit | 2023-12-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020078 | GAX-AX_heroes_delta_1_2_2_k1_p2_free

On heroes in digraphs with forbidden induced forests | Open problem: hero status of $\Delta(1,2,2)$

论文中定位：Open problem: hero status of $\Delta(1,2,2)$；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alvaro Carbonero; Hidde Koerts; Benjamin Moore; Sophie Spirkl | 2023-06-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020079 | GAX-AX_heroes_k1_join_delta_1_1_h

Heroes in oriented complete multipartite graphs | Question 5.4

论文中定位：Question 5.4；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Pierre Aboulker; Guillaume Aubian; Pierre Charbit | 2023-12-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020080 | GAX-AX_heroes_oriented_star_delta_1_m_m

On heroes in digraphs with forbidden induced forests | Open problem: hero status of $\Delta(1,m,m')$ for degree-4 oriented stars

论文中定位：Open problem: hero status of $\Delta(1,m,m')$ for degree-4 oriented stars；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alvaro Carbonero; Hidde Koerts; Benjamin Moore; Sophie Spirkl | 2023-06-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020081 | GAX-AX_heroic_sets_bounded_dichromatic_finite_families

On heroes in digraphs with forbidden induced forests | Question 1.1

论文中定位：Question 1.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alvaro Carbonero; Hidde Koerts; Benjamin Moore; Sophie Spirkl | 2023-06-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020082 | GAX-AX_holes_length_divisible_3_detection

Detecting a long even hole | Open Question: Detecting Holes of Length Divisible by Three

论文中定位：Open Question: Detecting Holes of Length Divisible by Three；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Linda Cook; Paul Seymour | 2020-09-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020083 | GAX-AX_hypercube_coloring_1_ball_reconstruction_threshold

Shotgun reconstruction in the hypercube | Gap problem for 1-ball reconstruction threshold

论文中定位：Gap problem for 1-ball reconstruction threshold；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Michał Przykucki; Alexander Roberts; Alex Scott | 2019-07-16

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020084 | GAX-AX_hypercube_statistic_lambda_d_1_limit

On hypercube statistics | Conjecture on $\lambda(d,1)$

论文中定位：Conjecture on $\lambda(d,1)$；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Maria Axenovich; John Goldwasser | 2024-10-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020085 | GAX-AX_hypercube_statistic_lambda_tight_bounds

On hypercube statistics | Open problem on tight bounds for $\lambda(d,s)$

论文中定位：Open problem on tight bounds for $\lambda(d,s)$；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Maria Axenovich; John Goldwasser | 2024-10-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020086 | GAX-AX_hypercube_subcube_partitions_dimension_2_asymptotics

Partitioning the hypercube into smaller hypercubes | Problem 1.9

论文中定位：Problem 1.9；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Jozsef Balogh; Vladimir N. Potapov | 2024-11-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020087 | GAX-AX_hypercube_subcube_partitions_dimension_2_bound

Partitioning the hypercube into smaller hypercubes | Informal conjecture on $f_2(d)$

论文中定位：Informal conjecture on $f_2(d)$；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Jozsef Balogh; Vladimir N. Potapov | 2024-11-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020088 | GAX-AX_hypercube_subcube_partitions_matchings_counting

Partitioning the hypercube into smaller hypercubes | Problem 1.1

论文中定位：Problem 1.1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Jozsef Balogh; Vladimir N. Potapov | 2024-11-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020089 | GAX-AX_hypercube_subcube_partitions_ratio_growth

Partitioning the hypercube into smaller hypercubes | Problem 1.2

论文中定位：Problem 1.2；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Jozsef Balogh; Vladimir N. Potapov | 2024-11-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020090 | GAX-AX_hypercube_subcube_partitions_ratio_subexponential

Partitioning the hypercube into smaller hypercubes | Problem 1.10

论文中定位：Problem 1.10；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Jozsef Balogh; Vladimir N. Potapov | 2024-11-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020091 | GAX-AX_hypergraph_ramsey_k4_minus_edge_star

Hypergraph Ramsey numbers of cliques versus stars | Informal Conjecture: r(K^(3)_{4-e}, S^(3)_n) = Theta(n^2 / log n)

论文中定位：Informal Conjecture: r(K^(3)_{4-e}, S^(3)_n) = Theta(n^2 / log n)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox; Xiaoyu He; Dhruv Mubayi; Andrew Suk; Jacques Verstraete | 2022-10-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020092 | GAX-AX_identifying_codes_hereditary_vc_dimension_dichotomy

Identifying codes in hereditary classes of graphs and VC-dimension | VC-Dimension Dichotomy for Identifying Codes (Approximation)

论文中定位：VC-Dimension Dichotomy for Identifying Codes (Approximation)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Nicolas Bousquet; Aurélie Lagoutte; Zhentao Li; Aline Parreau; Stéphan Thomassé | 2017-04-14

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020093 | GAX-AX_implicit_representation_speed_n23_labels

Implicit representation of sparse hereditary families | Question (Section 3, implicit representation for speed $2^{n^{1+\varepsilon}}$)

论文中定位：Question (Section 3, implicit representation for speed $2^{n^{1+\varepsilon}}$)；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon | 2022-01-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020094 | GAX-AX_implicit_representation_speed_sqrt_n_labels

Implicit representation of sparse hereditary families | Question (Section 3, implicit representation for speed $2^{O(n\log n)}$)

论文中定位：Question (Section 3, implicit representation for speed $2^{O(n\log n)}$)；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon | 2022-01-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020095 | GAX-AX_independent_fvs_linear_forest_free_complexity

Independent Feedback Vertex Set for P_5-free Graphs | Complexity of Independent Feedback Vertex Set for H-free graphs (linear forest cases)

论文中定位：Complexity of Independent Feedback Vertex Set for H-free graphs (linear forest cases)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Konrad K. Dabrowski; Carl Feghali; Matthew Johnson; Daniel Paulusma | 2017-07-28

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020096 | GAX-AX_independent_set_apx_hard_nonsublinear_separators

Thin graph classes and polynomial-time approximation schemes | Open Problem (Section 1.2): APX-hardness on classes without sublinear separators

论文中定位：Open Problem (Section 1.2): APX-hardness on classes without sublinear separators；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák | 2017-04-01

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020097 | GAX-AX_independent_set_fpt_blownup_p4_free

When Maximum Stable Set can be solved in FPT time | Conjecture 1

论文中定位：Conjecture 1；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Édouard Bonnet; Nicolas Bousquet; Stéphan Thomassé; Rémi Watrigant | 2019-09-18

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020098 | GAX-AX_independent_set_fpt_blownup_path_free

When Maximum Stable Set can be solved in FPT time | Conjecture 2

论文中定位：Conjecture 2；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Édouard Bonnet; Nicolas Bousquet; Stéphan Thomassé; Rémi Watrigant | 2019-09-18

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020099 | GAX-AX_independent_set_fpt_h_free_candidates

When Maximum Stable Set can be solved in FPT time | Informal belief on FPT candidates

论文中定位：Informal belief on FPT candidates；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Édouard Bonnet; Nicolas Bousquet; Stéphan Thomassé; Rémi Watrigant | 2019-09-18

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020100 | GAX-AX_independent_set_h_free_fpt_dichotomy

When Maximum Stable Set can be solved in FPT time | Parameterized MIS Dichotomy

论文中定位：Parameterized MIS Dichotomy；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Édouard Bonnet; Nicolas Bousquet; Stéphan Thomassé; Rémi Watrigant | 2019-09-18

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020101 | GAX-AX_independent_set_h_free_polynomial_dichotomy

When Maximum Stable Set can be solved in FPT time | Classical MIS Dichotomy Conjecture

论文中定位：Classical MIS Dichotomy Conjecture；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Édouard Bonnet; Nicolas Bousquet; Stéphan Thomassé; Rémi Watrigant | 2019-09-18

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020102 | GAX-AX_induced_arithmetic_removal_abelian_groups

Induced arithmetic removal: complexity 1 patterns over finite fields | Informal Conjecture (Induced Arithmetic Removal for Arbitrary Abelian Groups)

论文中定位：Informal Conjecture (Induced Arithmetic Removal for Arbitrary Abelian Groups)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Jonathan Tidor; Yufei Zhao | 2019-11-08

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020103 | GAX-AX_induced_arithmetic_removal_complexity_1_polynomial

Induced arithmetic removal: complexity 1 patterns over finite fields | Informal Conjecture (Polynomial Quantitative Bound for Complexity 1 Systems)

论文中定位：Informal Conjecture (Polynomial Quantitative Bound for Complexity 1 Systems)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Jonathan Tidor; Yufei Zhao | 2019-11-08

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020104 | GAX-AX_induced_arithmetic_removal_higher_complexity

Induced arithmetic removal: complexity 1 patterns over finite fields | Open Problem (Extending Induced Arithmetic Removal to Higher Complexity and General Groups)

论文中定位：Open Problem (Extending Induced Arithmetic Removal to Higher Complexity and General Groups)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Jonathan Tidor; Yufei Zhao | 2019-11-08

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020105 | GAX-AX_induced_detour_path_excess_3

Detecting a long odd hole | Open Problem (induced st-path of excess length three)

论文中定位：Open Problem (induced st-path of excess length three)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Alex Scott; Paul Seymour | 2020-09-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020106 | GAX-AX_induced_detour_path_fixed_k

Finding an induced path that is not a shortest path | Question 1.5

论文中定位：Question 1.5；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Eli Berger; Paul Seymour; Sophie Spirkl | 2020-05-26

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020107 | GAX-AX_induced_disjoint_paths_variants_complexity_gap

Induced Disjoint Paths Without an Induced Minor | Informal Question on Complexity Gap Between Induced Disjoint Paths Variants

论文中定位：Informal Question on Complexity Gap Between Induced Disjoint Paths Variants；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Pierre Aboulker; Édouard Bonnet; Timothé Picavet; Nicolas Trotignon | 2025-02-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020108 | GAX-AX_induced_even_cycle_saturation_edge_addition

Halfway to induced saturation for even cycles | Question 1.7

论文中定位：Question 1.7；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Xinyue Fan; Sahab Hajebi; Sepehr Hajebi; Sophie Spirkl | 2025-06-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020109 | GAX-AX_induced_even_cycle_saturation_polynomial_size

Halfway to induced saturation for even cycles | Question 1.8

论文中定位：Question 1.8；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Xinyue Fan; Sahab Hajebi; Sepehr Hajebi; Sophie Spirkl | 2025-06-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020110 | GAX-AX_induced_subdivision_containment_subcubic_planar_dichotomy

Induced Disjoint Paths Without an Induced Minor | Conjecture 1.5

论文中定位：Conjecture 1.5；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Pierre Aboulker; Édouard Bonnet; Timothé Picavet; Nicolas Trotignon | 2025-02-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020111 | GAX-AX_interactive_sum_choice_strict_non_complete

The Interactive Sum Choice Number of Graphs | Conjecture 1.1

论文中定位：Conjecture 1.1；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Kitty Meeks | 2021-01-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020112 | GAX-AX_intersection_graphs_asymptotic_dimension_tight

Asymptotic dimension of intersection graphs | Optimal bound question

论文中定位：Optimal bound question；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Sergey Norin | 2022-10-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020113 | GAX-AX_intersection_graphs_polynomial_weak_coloring_numbers

Weak Coloring Numbers of Intersection Graphs | Question (polynomial weak coloring numbers for intersection graph classes)

论文中定位：Question (polynomial weak coloring numbers for intersection graph classes)；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Jakub Pekárek; Torsten Ueckerdt; Yelena Yuditsky | 2021-04-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020114 | GAX-AX_k5_minor_bipartite_average_degree_5

Bipartite graphs with no K_6 minor | Informal Conjecture: average degree ≥ 5 and max degree ≤ 6 forces K_5 minor in bipartite graphs

论文中定位：Informal Conjecture: average degree ≥ 5 and max degree ≤ 6 forces K_5 minor in bipartite graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Alex Scott; Paul Seymour; Sophie Spirkl | 2022-04-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020115 | GAX-AX_k6_minor_bipartite_min_degree_5

Bipartite graphs with no K_6 minor | Open Question: minimum degree five in Theorem 1.6

论文中定位：Open Question: minimum degree five in Theorem 1.6；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Alex Scott; Paul Seymour; Sophie Spirkl | 2022-04-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020116 | GAX-AX_k_colorable_deletion_clique_free_sharp

Making an H-Free Graph k-Colorable | Conjecture on sharpness of Theorem 1.1

论文中定位：Conjecture on sharpness of Theorem 1.1；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Zoe Himwich; Nitya Mani | 2021-03-19

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020117 | GAX-AX_k_colorable_deletion_mnk_asymptotics

Making an H-Free Graph k-Colorable | Open problem on $m(n,k)$ for fixed $k \geq 2$

论文中定位：Open problem on $m(n,k)$ for fixed $k \geq 2$；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Zoe Himwich; Nitya Mani | 2021-03-19

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020118 | GAX-AX_k_colorable_deletion_neighborhood_edge_cover

Making an H-Free Graph k-Colorable | Question 2.1

论文中定位：Question 2.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Zoe Himwich; Nitya Mani | 2021-03-19

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020119 | GAX-AX_k_colorable_deletion_odd_wheel_sharp

Making an H-Free Graph k-Colorable | Conjecture on sharpness of Theorem 1.5

论文中定位：Conjecture on sharpness of Theorem 1.5；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Zoe Himwich; Nitya Mani | 2021-03-19

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020120 | GAX-AX_kk_free_bounded_treewidth_max_chromatic

Triangle-free graphs of tree-width t are ceil((t + 3)/2)-colorable | Problem 3

论文中定位：Problem 3；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Ken-ichi Kawarabayashi | 2017-06-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020121 | GAX-AX_kromatic_claw_free_grassmannian_subvariety

The Kromatic Symmetric Function: A K-theoretic Analogue of X_G | Grassmannian Subvariety Construction Problem

论文中定位：Grassmannian Subvariety Construction Problem；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Logan Crew; Oliver Pechenik; Sophie Spirkl | 2023-05-18

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020122 | GAX-AX_kt_immersion_free_clique_count

On the number of cliques in graphs with a forbidden subdivision or immersion | Conjecture on maximum cliques with no K_t-immersion

论文中定位：Conjecture on maximum cliques with no K_t-immersion；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Fan Wei | 2018-08-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020123 | GAX-AX_kt_minor_free_list_chromatic_2t

Improved lower bound for the list chromatic number of graphs with no K_t minor | Problem 1

论文中定位：Problem 1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Raphael Steiner | 2021-10-18

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020124 | GAX-AX_kt_minor_free_universal_graph_separators

Universality in minor-closed graph classes | Open Problem (Section 1): Balanced Separator for U_t

论文中定位：Open Problem (Section 1): Balanced Separator for U_t；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Tony Huynh; Bojan Mohar; Robert Šámal; Carsten Thomassen; David R. Wood | 2021-09-01

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020125 | GAX-AX_kt_subdivision_free_clique_count_exponent

On the number of cliques in graphs with a forbidden subdivision or immersion | Conjecture on optimal exponential constant for K_t-subdivision clique count

论文中定位：Conjecture on optimal exponential constant for K_t-subdivision clique count；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Fan Wei | 2018-08-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020126 | GAX-AX_large_chromatic_dense_minor_edge_density

Dense minors of graphs with independence number two | Informal conjecture on improving the $g(t)$ lower bound

论文中定位：Informal conjecture on improving the $g(t)$ lower bound；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Sergey Norin; Paul Seymour | 2022-06-01

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020127 | GAX-AX_large_chromatic_induced_path_uniquely_covered

Induced subgraphs of graphs with large chromatic number. IX. Rainbow paths | Question (uniquely-covered vertices in an induced path)

论文中定位：Question (uniquely-covered vertices in an induced path)；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour | 2017-07-03

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020128 | GAX-AX_large_chromatic_rainbow_hole_consecutive_vertices

Induced subgraphs of graphs with large chromatic number. IX. Rainbow paths | Question (rainbow hole with s consecutive rainbow vertices)

论文中定位：Question (rainbow hole with s consecutive rainbow vertices)；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour | 2017-07-03

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020129 | GAX-AX_large_treewidth_unavoidable_induced_subgraphs

Induced subgraphs and tree decompositions XVI. Complete bipartite induced minors | Question 1.1

论文中定位：Question 1.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Sepehr Hajebi; Sophie Spirkl | 2026-02-18

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020130 | GAX-AX_largest_clique_minor_opt_approximation

A Polynomial-Time Approximation Algorithm for Complete Interval Minors | Open Problem: polytime $f(\mathrm{OPT})$-approximation for largest complete minor

论文中定位：Open Problem: polytime $f(\mathrm{OPT})$-approximation for largest complete minor；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Romain Bourneuf; Julien Cocquet; Chaoliang Tang; Stéphan Thomassé | 2025-05-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020131 | GAX-AX_lattice_cube_list_colorability_threshold

Mixing properties of colorings of the $\mathbb{Z}^d$ lattice | Optimal n for L^d_n-list-colorability

论文中定位：Optimal n for L^d_n-list-colorability；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Raimundo Briceño; Nishant Chandgotia; Alexander Magazinov; Yinon Spinka | 2019-12-25

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020132 | GAX-AX_layered_treewidth_bounded_queue_number

Minor-closed graph classes with bounded layered pathwidth | Open problem on queue-number of bounded layered treewidth graphs

论文中定位：Open problem on queue-number of bounded layered treewidth graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Vida Dujmović; David Eppstein; Gwenaël Joret; Pat Morin; David R. Wood | 2020-06-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020133 | GAX-AX_linear_equation_free_sets_sqrt_n

The regularity method for graphs with few 4-cycles | Improvement of the bound in Theorem 1.12

论文中定位：Improvement of the bound in Theorem 1.12；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox; Benny Sudakov; Yufei Zhao | 2021-09-25

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020134 | GAX-AX_lipschitz_bijection_dictator_xor_inverse

Lipschitz bijections between boolean functions | Informal Question (Section 2)

论文中定位：Informal Question (Section 2)；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Tom Johnston; Alex Scott | 2021-12-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020135 | GAX-AX_list_chromatic_fraction_delta_small_clique

Bounding $χ$ by a fraction of $Δ$ for graphs without large cliques | Question 1.5

论文中定位：Question 1.5；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Tom Kelly; Peter Nelson; Luke Postle | 2018-03-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020136 | GAX-AX_list_chromatic_reed_bound

Bounding $χ$ by a fraction of $Δ$ for graphs without large cliques | List-chromatic analogue of Reed's Conjecture

论文中定位：List-chromatic analogue of Reed's Conjecture；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Tom Kelly; Peter Nelson; Luke Postle | 2018-03-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020137 | GAX-AX_list_coloring_rp3_free_np_hard

Complexity dichotomy for List-5-Coloring with a forbidden induced subgraph | Open Question: NP-hardness of List-k-Coloring on rP3-free graphs

论文中定位：Open Question: NP-hardness of List-k-Coloring on rP3-free graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Sepehr Hajebi; Yanjia Li; Sophie Spirkl | 2022-08-30

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020138 | GAX-AX_list_decodable_zero_rate_codes_asymptotics

List-decodable zero-rate codes | Informal Conjecture on exact asymptotics of maxcode_L for even L

论文中定位：Informal Conjecture on exact asymptotics of maxcode_L for even L；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Boris Bukh; Yury Polyanskiy | 2018-05-14

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020139 | GAX-AX_list_ramsey_chromatic_family_exact

Multicolor list Ramsey numbers grow exponentially | Conjecture on $R_\ell(\mathcal{H}_s, k)$

论文中定位：Conjecture on $R_\ell(\mathcal{H}_s, k)$；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Xiaoyu He; Sammy Luo; Max Wenqiang Xu | 2022-01-23

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020140 | GAX-AX_list_ramsey_exponential_constant_k3

Multicolor list Ramsey numbers grow exponentially | Determine the exponential constant in $R_\ell(H,k)$

论文中定位：Determine the exponential constant in $R_\ell(H,k)$；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Xiaoyu He; Sammy Luo; Max Wenqiang Xu | 2022-01-23

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020141 | GAX-AX_local_search_ptas_monotone_fo_optimization

Baker game and polynomial-time approximation schemes | Open Problem (Section 1.1)

论文中定位：Open Problem (Section 1.1)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák | 2019-01-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020142 | GAX-AX_lollipop_active_path_complete_graph_criterion

Lollipops, dense cycles and chords | Question 4.1

论文中定位：Question 4.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Beatriz Martins; Stéphan Thomassé; Nicolas Trotignon | 2025-10-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020143 | GAX-AX_lollipop_active_vertices_optimal_cycle

Lollipops, dense cycles and chords | Question 4.2

论文中定位：Question 4.2；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Beatriz Martins; Stéphan Thomassé; Nicolas Trotignon | 2025-10-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020144 | GAX-AX_long_odd_hole_detection_running_time

Detecting a long odd hole | Open Question (running time improvement for long hole detection)

论文中定位：Open Question (running time improvement for long hole detection)；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Alex Scott; Paul Seymour | 2020-09-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020145 | GAX-AX_long_odd_hole_heavy_path_lemma

Detecting a long odd hole | Informal Conjecture (heavy path extension)

论文中定位：Informal Conjecture (heavy path extension)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Alex Scott; Paul Seymour | 2020-09-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020146 | GAX-AX_majority_coloring_digraphs_1_over_k

Majority Colourings of Digraphs | Conjecture 9

论文中定位：Conjecture 9；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Stephan Kreutzer; Sang-il Oum; Paul Seymour; Dominic van der Zypen; David R. Wood | 2016-08-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020147 | GAX-AX_majority_coloring_digraphs_2_colorable_recognition

Majority Colourings of Digraphs | Open Problem 5

论文中定位：Open Problem 5；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Stephan Kreutzer; Sang-il Oum; Paul Seymour; Dominic van der Zypen; David R. Wood | 2016-08-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020148 | GAX-AX_majority_coloring_digraphs_3_colors

Majority Colourings of Digraphs | Conjecture 2

论文中定位：Conjecture 2；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Stephan Kreutzer; Sang-il Oum; Paul Seymour; Dominic van der Zypen; David R. Wood | 2016-08-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020149 | GAX-AX_majority_coloring_digraphs_beta_3_colors

Majority Colourings of Digraphs | Open Problem 1

论文中定位：Open Problem 1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Stephan Kreutzer; Sang-il Oum; Paul Seymour; Dominic van der Zypen; David R. Wood | 2016-08-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020150 | GAX-AX_majority_coloring_digraphs_choosability_constant

Majority Colourings of Digraphs | Open Problem 7

论文中定位：Open Problem 7；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Stephan Kreutzer; Sang-il Oum; Paul Seymour; Dominic van der Zypen; David R. Wood | 2016-08-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020151 | GAX-AX_majority_coloring_digraphs_fractional_weight

Majority Colourings of Digraphs | Open Problem 8

论文中定位：Open Problem 8；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Stephan Kreutzer; Sang-il Oum; Paul Seymour; Dominic van der Zypen; David R. Wood | 2016-08-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020152 | GAX-AX_majority_coloring_eulerian_digraphs_3_colors

Majority Colourings of Digraphs | Open Problem 3

论文中定位：Open Problem 3；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Stephan Kreutzer; Sang-il Oum; Paul Seymour; Dominic van der Zypen; David R. Wood | 2016-08-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020153 | GAX-AX_majority_coloring_tournaments_3_colors

Majority Colourings of Digraphs | Open Problem 2

论文中定位：Open Problem 2；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Stephan Kreutzer; Sang-il Oum; Paul Seymour; Dominic van der Zypen; David R. Wood | 2016-08-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020154 | GAX-AX_meyniel_cop_number_k_uniform_hypergraph

Catching a robber on a random k-uniform hypergraph | Conjecture 1.4

论文中定位：Conjecture 1.4；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Joshua Erde; Mihyun Kang; Florian Lehner; Bojan Mohar; Dominik Schmid | 2024-04-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020155 | GAX-AX_min_degree_3_cycle_linear_chords

Lollipops, dense cycles and chords | Question 4.3

论文中定位：Question 4.3；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Beatriz Martins; Stéphan Thomassé; Nicolas Trotignon | 2025-10-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020156 | GAX-AX_minor_free_choosability_trivial_bound_characterization

On the choosability of H-minor-free graphs | Problem 1

论文中定位：Problem 1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Olivier Fischer; Raphael Steiner | 2023-04-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020157 | GAX-AX_minor_free_p_centered_coloring_polynomial

The grid-minor theorem revisited | Question 3

论文中定位：Question 3；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Vida Dujmović; Robert Hickingbotham; Jędrzej Hodor; Gwenaël Joret; Hoang La; Piotr Micek; Pat Morin; Clément Rambaud; David R. Wood | 2023-07-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020158 | GAX-AX_minor_free_underlying_treewidth_polynomial_treedepth

The grid-minor theorem revisited | Question 1

论文中定位：Question 1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Vida Dujmović; Robert Hickingbotham; Jędrzej Hodor; Gwenaël Joret; Hoang La; Piotr Micek; Pat Morin; Clément Rambaud; David R. Wood | 2023-07-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020159 | GAX-AX_mis_complexity_even_hole_k4_free

(Theta, triangle)-free and (even hole, K_4)-free graphs. Part 2 : bounds on treewidth | Maximum Independent Set complexity for (even hole, K4)-free and (theta, triangle)-free graphs

论文中定位：Maximum Independent Set complexity for (even hole, K4)-free and (theta, triangle)-free graphs；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marcin Pilipczuk; Ni Luh Dewi Sintiari; Stéphan Thomassé; Nicolas Trotignon | 2020-10-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020160 | GAX-AX_mis_complexity_subdivided_claw_free_p7

(Theta, triangle)-free and (even hole, K_4)-free graphs. Part 2 : bounds on treewidth | Maximum Independent Set complexity for H-free graphs when H contains P7, S1,1,3, or S1,2,2

论文中定位：Maximum Independent Set complexity for H-free graphs when H contains P7, S1,1,3, or S1,2,2；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marcin Pilipczuk; Ni Luh Dewi Sintiari; Stéphan Thomassé; Nicolas Trotignon | 2020-10-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020161 | GAX-AX_mis_planar_induced_minor_free_quasipolynomial

Maximum Independent Set when excluding an induced minor: $K_1 + tK_2$ and $tC_3 \uplus C_4$ | Question 2

论文中定位：Question 2；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Édouard Bonnet; Julien Duron; Colin Geniet; Stéphan Thomassé; Alexandra Wesolek | 2025-12-31

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020162 | GAX-AX_multicolor_ramsey_3_graph_every_exponent

The growth rate of multicolor Ramsey numbers of 3-graphs | Problem 4.1

论文中定位：Problem 4.1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Domagoj Bradač; Jacob Fox; Benny Sudakov | 2024-04-29

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020163 | GAX-AX_multicolor_ramsey_hypergraph_tower_height

The growth rate of multicolor Ramsey numbers of 3-graphs | Problem 1.1

论文中定位：Problem 1.1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Domagoj Bradač; Jacob Fox; Benny Sudakov | 2024-04-29

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020164 | GAX-AX_multidimensional_ramsey_r2_superexponential

A multidimensional Ramsey Theorem | Problem 4.1

论文中定位：Problem 4.1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | António Girão; Gal Kronenberg; Alex Scott | 2024-12-31

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020165 | GAX-AX_multidimensional_ramsey_r2_vs_erdos_szekeres

A multidimensional Ramsey Theorem | Problem 4.2

论文中定位：Problem 4.2；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | António Girão; Gal Kronenberg; Alex Scott | 2024-12-31

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020166 | GAX-AX_multiplicative_group_matrix_rank_sharp_bounds

Rank of matrices with entries from a multiplicative group | Open problem: improved quantitative rank bounds

论文中定位：Open problem: improved quantitative rank bounds；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Jozsef Solymosi | 2021-09-01

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020167 | GAX-AX_multitasking_capacity_average_degree_log_n

A Graph-Theoretic Approach to Multitasking | Open Problem (multitasker at average degree $\Theta(\log n)$)

论文中定位：Open Problem (multitasker at average degree $\Theta(\log n)$)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Jonathan D. Cohen; Biswadip Dey; Tom Griffiths; Sebastian Musslick; Kayhan Ozcimder; Daniel Reichman; Igor Shinkar; Tal Wagner | 2017-06-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020168 | GAX-AX_multitasking_interference_increases_with_depth

A Graph-Theoretic Approach to Multitasking | Open Problem (interference worsens with depth)

论文中定位：Open Problem (interference worsens with depth)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Jonathan D. Cohen; Biswadip Dey; Tom Griffiths; Sebastian Musslick; Kayhan Ozcimder; Daniel Reichman; Igor Shinkar; Tal Wagner | 2017-06-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020169 | GAX-AX_mwis_long_hole_prism_free_fpt

On the Maximum Weight Independent Set Problem in graphs without induced cycles of length at least five | FPT Algorithm for MWIS in (Long-Hole, k-Prism)-Free Graphs

论文中定位：FPT Algorithm for MWIS in (Long-Hole, k-Prism)-Free Graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Marcin Pilipczuk; Michał Pilipczuk; Stéphan Thomassé | 2020-01-16

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020170 | GAX-AX_mwis_p7_free_complexity

Induced subgraphs of bounded treewidth and the container method | MWIS in P7-free graphs

论文中定位：MWIS in P7-free graphs；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Tara Abrishami; Maria Chudnovsky; Marcin Pilipczuk; Paweł Rzążewski; Paul Seymour | 2020-03-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020171 | GAX-AX_mwis_qptas_forest_subdivision_free

Quasi-polynomial time approximation schemes for the Maximum Weight Independent Set Problem in H-free graphs | Conjecture 1.3

论文中定位：Conjecture 1.3；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Marcin Pilipczuk; Michał Pilipczuk; Stéphan Thomassé | 2023-11-14

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020172 | GAX-AX_mycielski_graphs_independence_ratio

Revisiting a theorem by Folkman on graph colouring | Open problem on minimum independence ratio of Mycielski graphs

论文中定位：Open problem on minimum independence ratio of Mycielski graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Pierre Charbit; Oscar Defrain; Gwenaël Joret; Aurélie Lagoutte; Vincent Limouzy; Lucas Pastor; Jean-Sébastien Sereni | 2020-03-23

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020173 | GAX-AX_negative_p_energy_path_minimum

Vertex Partitioning and p-Energy of Graphs | Conjecture 2

论文中定位：Conjecture 2；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Saieed Akbari; Hitesh Kumar; Bojan Mohar; Shivaramakrishna Pragada | 2025-06-20

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020174 | GAX-AX_neighborhood_complexity_kt_minor_free_polynomial

Neighborhood complexity of planar graphs | Problem 42

论文中定位：Problem 42；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Gwenaël Joret; Clément Rambaud | 2023-12-19

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020175 | GAX-AX_neighborhood_complexity_planar_tight_bound

Neighborhood complexity of planar graphs | Problem 41

论文中定位：Problem 41；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Gwenaël Joret; Clément Rambaud | 2023-12-19

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020176 | GAX-AX_non_averaging_sets_n_quarter_tight

Homogeneous structures in subset sums and non-averaging sets | Informal Conjecture on h(n)

论文中定位：Informal Conjecture on h(n)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox; Huy Tuan Pham | 2023-11-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020177 | GAX-AX_normal_graphs_cover_sizes_max_order

Minimal normal graph covers | Problem 1

论文中定位：Problem 1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Gajser; Bojan Mohar | 2016-01-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020178 | GAX-AX_normal_graphs_random_gnp_whp

Minimal normal graph covers | Open Question (normality of random graphs)

论文中定位：Open Question (normality of random graphs)；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Gajser; Bojan Mohar | 2016-01-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020179 | GAX-AX_ordered_complete_graph_monochromatic_nonnested_matching

Open problems of the 33rd Workshop on Cycles and Colourings | Problem 1.1

论文中定位：Problem 1.1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | János Barát; Zdeněk Dvořák; Penny Haxell; František Kardoš; Borut Lužar; Alfréd Onderko; Jozef Rajník; Roman Soták; Nikolay Ulyanov | 2025-11-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020180 | GAX-AX_ordered_pure_pairs_n_polylog

Pure pairs. VII. Homogeneous submatrices in 0/1-matrices with a forbidden submatrix | Informal belief on Conjecture 1.7 and polylog bound

论文中定位：Informal belief on Conjecture 1.7 and polylog bound；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour; Sophie Spirkl | 2021-01-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020181 | GAX-AX_ordered_ramsey_matching_exponent_gap

Ordered Ramsey numbers | Open problem: true ordered Ramsey number of matchings

论文中定位：Open problem: true ordered Ramsey number of matchings；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox; Choongbum Lee; Benny Sudakov | 2016-04-25

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020182 | GAX-AX_ordered_ramsey_matching_vs_triangle_order

Ordered Ramsey numbers | Open problem: ordered Ramsey number of matchings vs triangles

论文中定位：Open problem: ordered Ramsey number of matchings vs triangles；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox; Choongbum Lee; Benny Sudakov | 2016-04-25

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020183 | GAX-AX_oriented_chi_bounded_forbidden_oriented_star

$χ$-bounded families of oriented graphs | Conjecture 4

论文中定位：Conjecture 4；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Pierre Aboulker; Jørgen Bang-Jensen; Nicolas Bousquet; Pierre Charbit; Frédéric Havet; Frédéric Maffray; Jose Zamora | 2016-05-24

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020184 | GAX-AX_oriented_chi_bounded_forbidden_p4_orientations

$χ$-bounded families of oriented graphs | Conjecture 5

论文中定位：Conjecture 5；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Pierre Aboulker; Jørgen Bang-Jensen; Nicolas Bousquet; Pierre Charbit; Frédéric Havet; Frédéric Maffray; Jose Zamora | 2016-05-24

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020185 | GAX-AX_oriented_tree_avoiding_2_coloring_complexity

Colorings of oriented planar graphs avoiding a monochromatic subgraph | Open Complexity Problem (2-F-PFC for oriented trees of maximum degree ≥ 3)

论文中定位：Open Complexity Problem (2-F-PFC for oriented trees of maximum degree ≥ 3)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Helena Bergold; Winfried Hochstättler; Raphael Steiner | 2022-06-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020186 | GAX-AX_p_entropic_matroids_prime_representable

Entropic matroids and their representation | Open problem: p-entropic matroids linear for prime p

论文中定位：Open problem: p-entropic matroids linear for prime p；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Emmanuel Abbe; Sophie Spirkl | 2019-09-26

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020187 | GAX-AX_padded_strings_unavoidable_substring_sets

Induced subgraphs and tree decompositions XIII. Basic obstructions in $\mathcal{H}$-free graphs for finite $\mathcal{H}$ | Question 8.3

论文中定位：Question 8.3；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Bogdan Alecu; Maria Chudnovsky; Sepehr Hajebi; Sophie Spirkl | 2024-11-26

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020188 | GAX-AX_partial_grundy_coloring_parameterized_complexity

Grundy Coloring & friends, Half-Graphs, Bicliques | Parameterized complexity of Partial Grundy Coloring on general graphs

论文中定位：Parameterized complexity of Partial Grundy Coloring on general graphs；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Pierre Aboulker; Édouard Bonnet; Eun Jung Kim; Florian Sikora | 2020-01-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020189 | GAX-AX_path_free_polynomial_chi_tau_d

Polynomial bounds for chromatic number. V. Excluding a tree of radius two and a complete multipartite graph | Question: Extending Theorem 1.6 to paths

论文中定位：Question: Extending Theorem 1.6 to paths；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour | 2023-01-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020190 | GAX-AX_perfect_graphs_combinatorial_independent_set_algorithm

The independent set problem is FPT for even-hole-free graphs | Open problem: Combinatorial algorithm for MIS on perfect graphs

论文中定位：Open problem: Combinatorial algorithm for MIS on perfect graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Edin Husic; Stephan Thomasse; Nicolas Trotignon | 2019-10-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020191 | GAX-AX_perfect_graphs_combinatorial_mis_algorithm

On the Maximum Weight Independent Set Problem in graphs without induced cycles of length at least five | Combinatorial Polynomial-Time Algorithm for MIS/MWIS in Perfect Graphs

论文中定位：Combinatorial Polynomial-Time Algorithm for MIS/MWIS in Perfect Graphs；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Marcin Pilipczuk; Michał Pilipczuk; Stéphan Thomassé | 2020-01-16

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020192 | GAX-AX_perfect_graphs_combinatorial_mwis_algorithm

Induced subgraphs of bounded treewidth and the container method | Combinatorial polynomial-time algorithm for MWIS in perfect graphs

论文中定位：Combinatorial polynomial-time algorithm for MWIS in perfect graphs；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Tara Abrishami; Maria Chudnovsky; Marcin Pilipczuk; Paweł Rzążewski; Paul Seymour | 2020-03-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020193 | GAX-AX_perfect_graphs_tight_skew_partitions_count

Colouring perfect graphs with bounded clique number | Open Question on Number of Tight Skew Partitions

论文中定位：Open Question on Number of Tight Skew Partitions；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Aurélie Lagoutte; Paul Seymour; Sophie Spirkl | 2017-07-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020194 | GAX-AX_perfect_matching_flip_switchable_recognition

The Perfect Matching Reconfiguration Problem | Open Question: Recognition of Switchable Graphs

论文中定位：Open Question: Recognition of Switchable Graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Nicolas Bousquet; Marc Heinrich; Takehiro Ito; Yusuke Kobayashi; Arnaud Mary; Moritz Mühlenthaler; Kunihiro Wasa | 2019-04-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020195 | GAX-AX_permutation_property_testing_query_bounds

Fast property testing and metrics for permutations | Open Problem (better query complexity bounds for property testing)

论文中定位：Open Problem (better query complexity bounds for property testing)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Fan Wei | 2018-04-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020196 | GAX-AX_planar_c4_c5_fractional_chromatic_11_3

Planar graphs without cycles of length 4 or 5 are (11:3)-colorable | Informal conjecture on optimality of 11/3

论文中定位：Informal conjecture on optimality of 11/3；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Xiaolan Hu | 2019-07-14

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020197 | GAX-AX_planar_c4_c7_free_3_choosable

Correspondence coloring and its application to list-coloring planar graphs without cycles of lengths 4 to 8 | Open Question on 3-Choosability for Forbidden Cycles 4-7 and 4-6

论文中定位：Open Question on 3-Choosability for Forbidden Cycles 4-7 and 4-6；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdenek Dvorak; Luke Postle | 2016-10-08

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020198 | GAX-AX_planar_c4_c8_free_correspondence_coloring

Correspondence coloring and its application to list-coloring planar graphs without cycles of lengths 4 to 8 | Informal Conjecture on Correspondence Chromatic Number of Planar Graphs Without Cycles 4-8

论文中定位：Informal Conjecture on Correspondence Chromatic Number of Planar Graphs Without Cycles 4-8；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdenek Dvorak; Luke Postle | 2016-10-08

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020199 | GAX-AX_planar_cover_graph_poset_dimension_height

Planar posets have dimension at most linear in their height | Open problem: dimension of posets with planar cover graphs vs. height

论文中定位：Open problem: dimension of posets with planar cover graphs vs. height；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Gwenaël Joret; Piotr Micek; Veit Wiechert | 2017-09-23

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020200 | GAX-AX_planar_expansions_edge_erdos_posa

Erdős-Pósa from ball packing | Open Problem (edge-EP property characterization for planar graph expansions)

论文中定位：Open Problem (edge-EP property characterization for planar graph expansions)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Wouter Cames van Batenburg; Gwenaël Joret; Arthur Ulmer | 2020-06-08

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020201 | GAX-AX_planar_independence_number_n_over_4

Triangle-free planar graphs with small independence number | Open Problem (independence number of planar graphs)

论文中定位：Open Problem (independence number of planar graphs)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Jordan Venters | 2017-02-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020202 | GAX-AX_planar_recoloring_linear_diameter_min_colors

An update on reconfiguring 10-colorings of planar graphs | Problem 3

论文中定位：Problem 3；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Carl Feghali | 2020-02-13

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020203 | GAX-AX_planar_weak_coloring_number_quadratic_log

Improved bounds for weak coloring numbers | Conjecture 4

论文中定位：Conjecture 4；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Gwenaël Joret; Piotr Micek | 2022-03-25

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020204 | GAX-AX_plane_graphs_weak_diameter_2_coloring

Weak diameter coloring of graphs on surfaces | Question 6

论文中定位：Question 6；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Sergey Norin | 2021-11-13

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020205 | GAX-AX_plane_triangle_free_distant_precoloring_extension

Three-coloring triangle-free graphs on surfaces V. Coloring planar graphs with distant anomalies | Conjecture 1.4

论文中定位：Conjecture 1.4；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdenek Dvorak; Daniel Kral; Robin Thomas | 2020-04-15

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020206 | GAX-AX_poly_chi_bounded_forests_known_complete

Polynomial bounds for chromatic number. II. Excluding a star-forest | Informal remark on completeness of known forests satisfying Esperet's conjecture

论文中定位：Informal remark on completeness of known forests satisfying Esperet's conjecture；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour; Sophie Spirkl | 2021-07-25

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020207 | GAX-AX_poly_chi_bounded_odd_multihole_free

Polynomial bounds for chromatic number VII. Disjoint holes | Open problem (poly-χ-bounded for k-multihole with only odd cycles)

论文中定位：Open problem (poly-χ-bounded for k-multihole with only odd cycles)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Alex Scott; Paul Seymour; Sophie Spirkl | 2022-02-18

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020208 | GAX-AX_poly_chi_bounded_self_isolating_all

Polynomial bounds for chromatic number VII. Disjoint holes | Question (are all graphs self-isolating?)

论文中定位：Question (are all graphs self-isolating?)；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Alex Scott; Paul Seymour; Sophie Spirkl | 2022-02-18

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020209 | GAX-AX_poly_chi_bounded_trees_disjoint_union

Polynomial bounds for chromatic number VII. Disjoint holes | Open problem (disjoint union of good trees)

论文中定位：Open problem (disjoint union of good trees)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Alex Scott; Paul Seymour; Sophie Spirkl | 2022-02-18

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020210 | GAX-AX_polynomial_expansion_balanced_separator_constant_m

On classes of graphs with strongly sublinear separators | Informal Conjecture (constant-size M)

论文中定位：Informal Conjecture (constant-size M)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák | 2018-02-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020211 | GAX-AX_polynomial_expansion_bounded_twin_width_speculation

Twin-width I: tractable FO model checking | Informal Conjecture (polynomial expansion and bounded twin-width)

论文中定位：Informal Conjecture (polynomial expansion and bounded twin-width)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Édouard Bonnet; Eun Jung Kim; Stéphan Thomassé; Rémi Watrigant | 2021-10-25

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020212 | GAX-AX_popular_progression_tower_height_small_primes

Popular progression differences in vector spaces II | Conjecture (Theorem 2 extends to primes p < 19)

论文中定位：Conjecture (Theorem 2 extends to primes p < 19)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Huy Tuan Pham | 2019-11-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020213 | GAX-AX_positive_square_energy_lower_bound_4n_5

A Linear Lower Bound for the Square Energy of Graphs | Informal Conjecture (improved linear bound)

论文中定位：Informal Conjecture (improved linear bound)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Saieed Akbari; Hitesh Kumar; Bojan Mohar; Shivaramakrishna Pragada | 2024-09-26

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020214 | GAX-AX_prime_cayley_graph_finite_ring_characterization

On prime Cayley graphs | Question 4.28

论文中定位：Question 4.28；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Michal Cizek; Logan Crew; Ján Mináč; Tung T. Nguyen; Sophie Spirkl; Nguyên Duy Tân | 2024-01-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020215 | GAX-AX_prime_cayley_graph_joined_union_decomposition

On prime Cayley graphs | Question 1.1

论文中定位：Question 1.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Michal Cizek; Logan Crew; Ján Mináč; Tung T. Nguyen; Sophie Spirkl; Nguyên Duy Tân | 2024-01-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020216 | GAX-AX_prime_cayley_graph_tensor_product_complete

On prime Cayley graphs | Question 4.31

论文中定位：Question 4.31；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Michal Cizek; Logan Crew; Ján Mináč; Tung T. Nguyen; Sophie Spirkl; Nguyên Duy Tân | 2024-01-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020217 | GAX-AX_prism_pyramid_theta_turtle_free_separators

Maximum independent sets in (pyramid, even hole)-free graphs | Conjecture 2.2

论文中定位：Conjecture 2.2；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Stéphan Thomassé; Nicolas Trotignon; Kristina Vušković | 2019-12-24

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020218 | GAX-AX_prism_pyramid_theta_wheel_free_separators

Maximum independent sets in (pyramid, even hole)-free graphs | Open problem: polynomial separator property for (prism, pyramid, theta, even wheel)-free graphs

论文中定位：Open problem: polynomial separator property for (prism, pyramid, theta, even wheel)-free graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Stéphan Thomassé; Nicolas Trotignon; Kristina Vušković | 2019-12-24

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020219 | GAX-AX_properly_colored_kn_longest_rainbow_cycle

Random subgraphs of properly edge-coloured complete graphs and long rainbow cycles | Open Problem: error term for longest rainbow cycle

论文中定位：Open Problem: error term for longest rainbow cycle；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Alexey Pokrovskiy; Benny Sudakov | 2016-08-25

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020220 | GAX-AX_pt_free_3_coloring_polynomial_all_t

List-three-coloring graphs with no induced P_6+rP_3 | Question 1.1

论文中定位：Question 1.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Shenwei Huang; Sophie Spirkl; Mingxian Zhong | 2018-07-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020221 | GAX-AX_pt_free_3_coloring_polynomial_large_t

H-colouring P_t-free graphs in subexponential time | Open Problem: Complexity of 3-colourability of P_t-free graphs

论文中定位：Open Problem: Complexity of 3-colourability of P_t-free graphs；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Carla Groenland; Karolina Okrasa; Pawel Rzążewski; Alex Scott; Paul Seymour; Sophie Spirkl | 2019-03-22

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020222 | GAX-AX_pt_free_3_coloring_polynomial_linear_forests

List-k-Coloring H-free graphs for all $k>4$ | Converse to Theorem 1.1 (3-Coloring dichotomy for H-free graphs)

论文中定位：Converse to Theorem 1.1 (3-Coloring dichotomy for H-free graphs)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Sepehr Hajebi; Sophie Spirkl | 2023-11-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020223 | GAX-AX_pt_free_independent_set_polynomial_t_7

H-colouring P_t-free graphs in subexponential time | Open Problem: Polynomial-time maximum independent set on P_t-free graphs

论文中定位：Open Problem: Polynomial-time maximum independent set on P_t-free graphs；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Carla Groenland; Karolina Okrasa; Pawel Rzążewski; Alex Scott; Paul Seymour; Sophie Spirkl | 2019-03-22

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020224 | GAX-AX_pure_pairs_blockade_polynomial_ordered_graphs

Pure pairs. IX. Transversal trees | Informal: which graphs H admit polynomial-size pure pairs

论文中定位：Informal: which graphs H admit polynomial-size pure pairs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour; Sophie Spirkl | 2024-02-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020225 | GAX-AX_pure_pairs_blockade_strong_transversal_characterization

Pure pairs. IX. Transversal trees | Informal: characterization of graphs with the strong transversal property

论文中定位：Informal: characterization of graphs with the strong transversal property；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour; Sophie Spirkl | 2024-02-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020226 | GAX-AX_pure_pairs_blockade_transversal_triangle_polynomial

Pure pairs. IX. Transversal trees | Question 1.3

论文中定位：Question 1.3；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour; Sophie Spirkl | 2024-02-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020227 | GAX-AX_quantum_ergodicity_small_spectral_window

High-girth near-Ramanujan graphs with localized eigenvectors | Open problem on quantum ergodicity over small spectral windows (Remark 1.2)

论文中定位：Open problem on quantum ergodicity over small spectral windows (Remark 1.2)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Shirshendu Ganguly; Nikhil Srivastava | 2019-08-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020228 | GAX-AX_quantum_sorting_poset_lower_bounds_equivalent

Information-theoretic lower bounds for quantum sorting | Conjecture on LB and QLB equivalence

论文中定位：Conjecture on LB and QLB equivalence；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jean Cardinal; Gwenaël Joret; Jérémie Roland | 2019-02-18

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020229 | GAX-AX_quartic_eulerian_digraphs_diplanar_obstructions

Embedding Quartic Eulerian Digraphs on the Plane | Problem 1.2

论文中定位：Problem 1.2；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Dan Archdeacon; Paul Bonnington; Bojan Mohar | 2017-06-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020230 | GAX-AX_quasi_isometry_multiplicative_to_additive_classes

Asymptotic structure. II. Path-width and additive quasi-isometry | Conjecture 1.2

论文中定位：Conjecture 1.2；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Tung Nguyen; Alex Scott; Paul Seymour | 2025-10-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020231 | GAX-AX_rainbow_stacking_odd_complete_proper_colorings

Rainbow Stackings of Random Edge-Colorings | Question 3.3

论文中定位：Question 3.3；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Colin Defant; Noah Kravitz | 2024-05-23

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020232 | GAX-AX_rainbow_stacking_random_colorings_sharp_threshold

Rainbow Stackings of Random Edge-Colorings | Problem 3.1

论文中定位：Problem 3.1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Colin Defant; Noah Kravitz | 2024-05-23

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020233 | GAX-AX_ramsey_sqrt_edges_bound_multicolor

Ramsey numbers of hypergraphs of a given size | Informal Problem: extending the $\sqrt{m}$ Ramsey bound to more than two colors for graphs

论文中定位：Informal Problem: extending the $\sqrt{m}$ Ramsey bound to more than two colors for graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Domagoj Bradač; Jacob Fox; Benny Sudakov | 2023-08-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020234 | GAX-AX_random_4_regular_graph_polynomial_coefficient

Open problems of the 33rd Workshop on Cycles and Colourings | Problem 2.1

论文中定位：Problem 2.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | János Barát; Zdeněk Dvořák; Penny Haxell; František Kardoš; Borut Lužar; Alfréd Onderko; Jozef Rajník; Roman Soták; Nikolay Ulyanov | 2025-11-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020235 | GAX-AX_random_graph_hereditary_extremal_bipartite_asymptotics

Largest subgraph from a hereditary property in a random graph | Open Problem (accurate estimate in the bipartite case)

论文中定位：Open Problem (accurate estimate in the bipartite case)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Michael Krivelevich; Wojciech Samotij | 2022-10-23

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020236 | GAX-AX_random_graph_hereditary_extremal_polynomial_saving

Largest subgraph from a hereditary property in a random graph | Informal Conjecture (polynomial saving in the bipartite case)

论文中定位：Informal Conjecture (polynomial saving in the bipartite case)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Michael Krivelevich; Wojciech Samotij | 2022-10-23

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020237 | GAX-AX_random_graph_hereditary_extremal_probability_range

Largest subgraph from a hereditary property in a random graph | Open Problem (characterizing edge probabilities for Theorem 1.1)

论文中定位：Open Problem (characterizing edge probabilities for Theorem 1.1)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Michael Krivelevich; Wojciech Samotij | 2022-10-23

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020238 | GAX-AX_random_irregular_subgraph_property_range

On random irregular subgraphs | Informal Conjecture on Property (*) for d = o(n/log n)

论文中定位：Informal Conjecture on Property (*) for d = o(n/log n)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Sammy Luo; Huy Tuan Pham | 2022-07-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020239 | GAX-AX_regular_4_critical_no_critical_edge

Critical edge sets in vertex-critical graphs | Problem 5.2

论文中定位：Problem 5.2；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Ema Skottova; Raphael Steiner | 2025-08-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020240 | GAX-AX_regular_independence_edge_deletion_cubic_infinite

A Counterexample to a Conjecture of Lovász | Question 4.1

论文中定位：Question 4.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alexander Clow; Penny Haxell; Bojan Mohar | 2025-06-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020241 | GAX-AX_restricted_difference_3_ap_free_density

A note on arithmetic progressions with restricted differences | Problem 8

论文中定位：Problem 8；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox; Huy Tuan Pham | 2026-05-13

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020242 | GAX-AX_restricted_difference_triangle_removal_lemma

A note on arithmetic progressions with restricted differences | Problem 9

论文中定位：Problem 9；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox; Huy Tuan Pham | 2026-05-13

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020243 | GAX-AX_rodl_h_free_restricted_union_cover

Strengthening Rodl's theorem | Union cover variant (open problem)

论文中定位：Union cover variant (open problem)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Alex Scott; Paul Seymour; Sophie Spirkl | 2022-08-03

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020244 | GAX-AX_runsort_permuton_horizontal_uniformity_direct

The runsort permuton | Direct Proof of Horizontal Uniformity

论文中定位：Direct Proof of Horizontal Uniformity；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Colin Defant; Noah Kravitz | 2021-06-28

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020245 | GAX-AX_ryser_near_extremal_hypergraphs_projective_planes

A note on intersecting hypergraphs with large cover number | Informal Problem (projective plane structure in near-extremal constructions)

论文中定位：Informal Problem (projective plane structure in near-extremal constructions)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Penny Haxell; Alex Scott | 2017-10-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020246 | GAX-AX_separable_index_parameterized_complexity

Factoring Pattern-Free Permutations into Separable ones | Question 1.3

论文中定位：Question 1.3；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Édouard Bonnet; Romain Bourneuf; Colin Geniet; Stéphan Thomassé | 2023-08-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020247 | GAX-AX_separable_index_pattern_avoiding_polynomial

Factoring Pattern-Free Permutations into Separable ones | Question 1.2

论文中定位：Question 1.2；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Édouard Bonnet; Romain Bourneuf; Colin Geniet; Stéphan Thomassé | 2023-08-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020248 | GAX-AX_separating_path_system_rainbow_cover_linear

Separating the edges of a graph by a linear number of paths | Problem 4

论文中定位：Problem 4；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Fábio Botler; François Dross; Tássio Naia; Jozef Skokan | 2023-10-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020249 | GAX-AX_separating_path_system_strong_2n

Separating the edges of a graph by a linear number of paths | Problem 3

论文中定位：Problem 3；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Fábio Botler; François Dross; Tássio Naia; Jozef Skokan | 2023-10-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020250 | GAX-AX_set_coloring_ramsey_chromatic_variant_equality

Set-coloring Ramsey numbers via codes | Informal Conjecture on R(n;r,s) vs R'(n;r,s) near Turán density

论文中定位：Informal Conjecture on R(n;r,s) vs R'(n;r,s) near Turán density；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox; Xiaoyu He; Dhruv Mubayi; Andrew Suk; Jacques Verstraete | 2022-06-22

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020251 | GAX-AX_set_coloring_ramsey_zero_rate_equality

Set-coloring Ramsey numbers and error-correcting codes near the zero-rate threshold | Informal Conjecture (Equality in Theorem 1.1 near the zero-rate threshold)

论文中定位：Informal Conjecture (Equality in Theorem 1.1 near the zero-rate threshold)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox; Huy Tuan Pham; Yufei Zhao | 2023-08-14

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020252 | GAX-AX_set_mapping_theorem_l_equals_1

Short proofs of some extremal results II | Open Problem: Reducing l to 1 in the Set Mapping Theorem

论文中定位：Open Problem: Reducing l to 1 in the Set Mapping Theorem；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | David Conlon; Jacob Fox; Benny Sudakov | 2016-02-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020253 | GAX-AX_shortest_even_hole_detection

Detecting a long even hole | Open Problem: Finding a Shortest Even Hole

论文中定位：Open Problem: Finding a Shortest Even Hole；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Linda Cook; Paul Seymour | 2020-09-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020254 | GAX-AX_shotgun_assembly_second_threshold_all_radii

Shotgun assembly of random graphs | Informal Conjecture (second phase transition for all radii)

论文中定位：Informal Conjecture (second phase transition for all radii)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Tom Johnston; Gal Kronenberg; Alexander Roberts; Alex Scott | 2025-06-23

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020255 | GAX-AX_shotgun_assembly_second_threshold_radius_2

Shotgun assembly of random graphs | Question

论文中定位：Question；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Tom Johnston; Gal Kronenberg; Alexander Roberts; Alex Scott | 2025-06-23

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020256 | GAX-AX_single_conflict_coloring_degenerate_log_n

Single-conflict colouring | Informal conjecture on single-conflict chromatic number of degenerate graphs

论文中定位：Informal conjecture on single-conflict chromatic number of degenerate graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Louis Esperet; Ross J. Kang; Kenta Ozeki | 2020-10-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020257 | GAX-AX_single_conflict_coloring_euler_genus_sqrt

Single-conflict colouring | Conjecture 3

论文中定位：Conjecture 3；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Louis Esperet; Ross J. Kang; Kenta Ozeki | 2020-10-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020258 | GAX-AX_spanning_trees_pivot_gray_code

A Gray code for arborescences of tournaments | Problem 1

论文中定位：Problem 1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Michael Hoffmann; Clément Legrand-Duchesne; Günter Rote | 2026-03-30

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020259 | GAX-AX_sparse_neighborhoods_chromatic_below_delta

Colouring Graphs with Sparse Neighbourhoods: Bounds and Applications | Question 1.4

论文中定位：Question 1.4；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Thomas Perrett; Luke Postle | 2018-10-15

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020260 | GAX-AX_sparse_neighborhoods_critical_bounded_clique

Colouring Graphs with Sparse Neighbourhoods: Bounds and Applications | Question 1.3

论文中定位：Question 1.3；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Thomas Perrett; Luke Postle | 2018-10-15

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020261 | GAX-AX_spectrally_symmetric_graph_edge_density

Spectrally symmetric orientations of graphs | Problem 1

论文中定位：Problem 1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Saieed Akbari; Jonathan Aloni; Maxwell Levit; Bojan Mohar; Steven Xia | 2025-12-08

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020262 | GAX-AX_split_digraph_strong_2_kernel_half

Distant digraph domination | Open question on strong 2-kernels in split digraphs

论文中定位：Open question on strong 2-kernels in split digraphs；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Tung Nguyen; Alex Scott; Paul Seymour | 2024-09-08

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020263 | GAX-AX_strong_coloring_number_bramble_polynomial_expansion

Shallow brambles | Question 2

论文中定位：Question 2；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Nicolas Bousquet; Wouter Cames van Batenburg; Louis Esperet; Gwenaël Joret; Piotr Micek | 2025-09-15

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020264 | GAX-AX_strong_coloring_numbers_tame_representation

Sublinear separators in intersection graphs of convex shapes | Question 6

论文中定位：Question 6；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdenek Dvorak; Rose McCarty; Sergey Norin | 2020-01-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020265 | GAX-AX_strong_coloring_numbers_tame_representation_negative

Sublinear separators in intersection graphs of convex shapes | Informal conjecture: negative answer to Question 6

论文中定位：Informal conjecture: negative answer to Question 6；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdenek Dvorak; Rose McCarty; Sergey Norin | 2020-01-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020266 | GAX-AX_structured_graph_codes_f2c_odd_n

Structured Codes of Graphs | Open case of Theorem 3 for odd n

论文中定位：Open case of Theorem 3 for odd n；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Anna Gujgiczer; János Körner; Aleksa Milojević; Gábor Simonyi | 2022-04-01

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020267 | GAX-AX_subdigraph_half_vertices_min_outdegree_gap

Subdigraphs of prescribed size and outdegree | Open problem: closing the gap for $d(s)$

论文中定位：Open problem: closing the gap for $d(s)$；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Raphael Steiner | 2022-10-23

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020268 | GAX-AX_subdivision_free_chi_bounded_multigraph_characterization

Induced subgraphs of graphs with large chromatic number. VI. Banana trees | Open Question: first pervasiveness characterisation

论文中定位：Open Question: first pervasiveness characterisation；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour | 2019-08-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020269 | GAX-AX_sublinear_separator_expansion_exponent_exact

A note on sublinear separators and expansion | Informal open question on the exact value of b_ε

论文中定位：Informal open question on the exact value of b_ε；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák | 2020-07-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020270 | GAX-AX_sublinear_separator_expansion_exponent_one_sided

A note on sublinear separators and expansion | Informal conjecture $b'_\varepsilon = b_\varepsilon$ for $0<\varepsilon<\tfrac{1}{2}$

论文中定位：Informal conjecture $b'_\varepsilon = b_\varepsilon$ for $0<\varepsilon<\tfrac{1}{2}$；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák | 2020-07-07

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020271 | GAX-AX_sublinear_separators_fractional_treewidth_fragile

Approximation metatheorems for classes with bounded expansion | Informal Conjecture (fractional treewidth-fragility of sublinear-separator classes)

论文中定位：Informal Conjecture (fractional treewidth-fragility of sublinear-separator classes)；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák | 2021-10-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020272 | GAX-AX_sublinear_separators_product_structure_tree_depth

Product structure of graph classes with strongly sublinear separators | Conjecture (product structure with bounded tree-depth and tight clique size)

论文中定位：Conjecture (product structure with bounded tree-depth and tight clique size)；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; David R. Wood | 2023-09-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020273 | GAX-AX_sublinear_separators_product_structure_treewidth

Product structure of graph classes with strongly sublinear separators | Open Problem 6

论文中定位：Open Problem 6；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; David R. Wood | 2023-09-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020274 | GAX-AX_sun_free_graphs_chi_bounded

Suns in triangle-free graphs of large chromatic number | Informal conjecture on sun-free χ-boundedness without net exclusion

论文中定位：Informal conjecture on sun-free χ-boundedness without net exclusion；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Sepehr Hajebi; Sophie Spirkl | 2026-03-08

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020275 | GAX-AX_surface_excluded_minors_hoppers_exist

Excluded minors for the Klein Bottle I. Low connectivity case | Informal Conjecture on Existence of Hoppers for Large Euler Genus

论文中定位：Informal Conjecture on Existence of Hoppers for Large Euler Genus；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Bojan Mohar; Petr Škoda | 2020-02-01

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020276 | GAX-AX_surface_triangle_free_3_coloring_linear

Three-coloring triangle-free graphs on surfaces VII. A linear-time algorithm | Open Problem (linear-time 3-coloring output)

论文中定位：Open Problem (linear-time 3-coloring output)；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdenek Dvorak; Daniel Kral; Robin Thomas | 2020-11-05

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020277 | GAX-AX_tea_sharing_optimal_sequence_exponential_length

Sharing tea on a graph | Question 6.1

论文中定位：Question 6.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | J. Pascal Gollin; Kevin Hendrey; Hao Huang; Tony Huynh; Bojan Mohar; Sang-il Oum; Ningyuan Yang; Wei-Hsuan Yu; Xuding Zhu | 2025-09-22

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020278 | GAX-AX_ternary_graphs_3_colorable

Proof of the Kalai-Meshulam conjecture | Open Problem: 3-colourability of ternary graphs

论文中定位：Open Problem: 3-colourability of ternary graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Alex Scott; Paul Seymour; Sophie Spirkl | 2018-09-28

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020279 | GAX-AX_theta_triangle_free_bounded_degree_treewidth

(Theta, triangle)-free and (even hole, K_4)-free graphs. Part 2 : bounds on treewidth | Open Question: (theta, triangle)-free graphs of bounded degree

论文中定位：Open Question: (theta, triangle)-free graphs of bounded degree；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marcin Pilipczuk; Ni Luh Dewi Sintiari; Stéphan Thomassé; Nicolas Trotignon | 2020-10-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020280 | GAX-AX_thin_overlays_sublinear_separators_unbounded_degree

Thin graph classes and polynomial-time approximation schemes | Informal Conjecture (Section 1.2): dropping bounded-degree assumption

论文中定位：Informal Conjecture (Section 1.2): dropping bounded-degree assumption；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák | 2017-04-01

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020281 | GAX-AX_token_sliding_chordal_clique_tree_degree

Token Sliding on Chordal Graphs | Question 1

论文中定位：Question 1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marthe Bonamy; Nicolas Bousquet | 2016-05-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020282 | GAX-AX_topological_minor_free_treewidth_product_structure

The grid-minor theorem revisited | Question 2

论文中定位：Question 2；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Vida Dujmović; Robert Hickingbotham; Jędrzej Hodor; Gweanël Joret; Hoang La; Piotr Micek; Pat Morin; Clément Rambaud; David R. Wood | 2023-07-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020283 | GAX-AX_tournament_color_avoiding_path_non_transitive

Color-avoiding directed paths in tournaments | Problem 5.1

论文中定位：Problem 5.1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Benny Sudakov; Yuval Wigderson | 2026-01-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020284 | GAX-AX_tournament_path_powers_exponent_constant

Powers of paths in tournaments | Open Problem: exact constant in the exponent

论文中定位：Open Problem: exact constant in the exponent；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Nemanja Draganić; François Dross; Jacob Fox; António Girão; Frédéric Havet; Dániel Korándi; William Lochet; David Munhá Correia; Alex Scott; Benny Sudakov | 2021-02-16

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020285 | GAX-AX_tournament_strong_eh_backedge_forest

Pure pairs. X. Tournaments and the strong Erdos-Hajnal property | Conjecture 1.5

论文中定位：Conjecture 1.5；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Alex Scott; Paul Seymour; Sophie Spirkl | 2023-08-08

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020286 | GAX-AX_tournament_strong_eh_three_backedges_d5

Pure pairs. X. Tournaments and the strong Erdos-Hajnal property | Informal question: can the D5-freeness hypothesis in Theorem 1.6 be dropped?

论文中定位：Informal question: can the D5-freeness hypothesis in Theorem 1.6 be dropped?；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Maria Chudnovsky; Alex Scott; Paul Seymour; Sophie Spirkl | 2023-08-08

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020287 | GAX-AX_tree_decomposition_edge_connectivity_leaf_count

Edge-partitioning 3-edge-connected graphs into paths | Conjecture 5

论文中定位：Conjecture 5；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Tereza Klimošová; Stéphan Thomassé | 2019-07-26

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020288 | GAX-AX_tree_pathwidth_path_partition_height_2k

Pathwidth and nonrepetitive list coloring | Optimality of factor 2 in Lemma 4

论文中定位：Optimality of factor 2 in Lemma 4；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Adam Gągol; Gwenaël Joret; Jakub Kozik; Piotr Micek | 2016-10-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020289 | GAX-AX_treedepth_bound_treewidth_binary_tree_path

Tight bound on treedepth in terms of pathwidth and longest path | Open Problem: sharpness of the treedepth bound in Theorem 1

论文中定位：Open Problem: sharpness of the treedepth bound in Theorem 1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Meike Hatzel; Gwenaël Joret; Piotr Micek; Marcin Pilipczuk; Torsten Ueckerdt; Bartosz Walczak | 2023-11-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020290 | GAX-AX_treewidth_fragile_weighted_vertex_cover_ptas

Approximation schemes for bounded distance problems on fractionally treewidth-fragile graphs | Open Problem: PTAS for weighted minimization problems in fractionally treewidth-fragile classes

论文中定位：Open Problem: PTAS for weighted minimization problems in fractionally treewidth-fragile classes；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Abhiruk Lahiri | 2021-05-04

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020291 | GAX-AX_triangle_free_planar_3_coloring_requests

Do triangle-free planar graphs have exponentially many 3-colorings? | Problem 1

论文中定位：Problem 1；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Jean-Sébastien Sereni | 2017-09-19

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020292 | GAX-AX_triangle_free_planar_fractional_chromatic_bound

Triangle-free planar graphs with the smallest independence number | Fractional Chromatic Number Bound for Planar Triangle-Free Graphs

论文中定位：Fractional Chromatic Number Bound for Planar Triangle-Free Graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Zdeněk Dvořák; Tomáš Masařík; Jan Musílek; Ondřej Pangrác | 2016-06-20

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020293 | GAX-AX_triangle_removal_vs_dfl_growth

Removal lemmas and approximate homomorphisms | Informal Conjecture on similar growth of $N_{DFL}(\varepsilon)$ and $1/\delta_{TRL}(\varepsilon)$

论文中定位：Informal Conjecture on similar growth of $N_{DFL}(\varepsilon)$ and $1/\delta_{TRL}(\varepsilon)$；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Jacob Fox; Yufei Zhao | 2022-01-13

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020294 | GAX-AX_triangle_s123_free_cliquewidth

(Theta, triangle)-free and (even hole, K_4)-free graphs. Part 2 : bounds on treewidth | Open Question: (triangle, S1,2,3)-free graphs cliquewidth

论文中定位：Open Question: (triangle, S1,2,3)-free graphs cliquewidth；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Marcin Pilipczuk; Ni Luh Dewi Sintiari; Stéphan Thomassé; Nicolas Trotignon | 2020-10-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020295 | GAX-AX_turan_h_free_bounded_matching_number

Turán graphs with bounded matching number | Extension to H-free graphs

论文中定位：Extension to H-free graphs；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Peter Frankl | 2022-10-26

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020296 | GAX-AX_twin_width_approximation_unordered_graphs

Twin-width VI: the lens of contraction sequences | Open Challenge: Efficient Approximation of Twin-Width for Unordered Graphs

论文中定位：Open Challenge: Efficient Approximation of Twin-Width for Unordered Graphs；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Édouard Bonnet; Eun Jung Kim; Amadeus Reinald; Stéphan Thomassé | 2022-05-31

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020297 | GAX-AX_twin_width_connected_domination_no_kernel

Twin-width and polynomial kernels | Open problem: no-kernel lower bound at twin-width 4 for Connected/Total k-Dominating Set

论文中定位：Open problem: no-kernel lower bound at twin-width 4 for Connected/Total k-Dominating Set；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Édouard Bonnet; Eun Jung Kim; Amadeus Reinald; Stéphan Thomassé; Rémi Watrigant | 2021-09-14

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020298 | GAX-AX_twin_width_matrices_infinite_fields

Twin-width V: linear minors, modular counting, and matrix multiplication | Open question on twin-width over infinite fields

论文中定位：Open question on twin-width over infinite fields；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Édouard Bonnet; Ugo Giocanti; Patrice Ossona de Mendez; Stéphan Thomassé | 2022-09-24

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020299 | GAX-AX_twin_width_mis_dominating_approximability

Twin-width III: Max Independent Set, Min Dominating Set, and Coloring | Informal conjecture on MIS approximability vs. Min Dominating Set

论文中定位：Informal conjecture on MIS approximability vs. Min Dominating Set；类型：Informal。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Édouard Bonnet; Colin Geniet; Eun Jung Kim; Stéphan Thomassé; Rémi Watrigant | 2021-02-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020300 | GAX-AX_vantage_protrusive_ordering_5_points

Ordering Candidates via Vantage Points | Five-point protrusive-but-not-witnessed construction

论文中定位：Five-point protrusive-but-not-witnessed construction；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Colin Defant; Noah Kravitz; Daniel G. Zhu | 2023-08-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020301 | GAX-AX_vantage_sign_patterns_r_dependence_gap

Ordering Candidates via Vantage Points | Gap in r-dependence of sign-pattern bound

论文中定位：Gap in r-dependence of sign-pattern bound；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Colin Defant; Noah Kravitz; Daniel G. Zhu | 2023-08-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020302 | GAX-AX_vantage_square_roots_sum_components_gap

Ordering Candidates via Vantage Points | Gap in connected-component count for sums of square roots

论文中定位：Gap in connected-component count for sums of square roots；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Noga Alon; Colin Defant; Noah Kravitz; Daniel G. Zhu | 2023-08-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020303 | GAX-AX_vertex_critical_edge_robust_fixed_k

Vertex-critical graphs far from edge-criticality | Open case of Erdős's problem for fixed small k

论文中定位：Open case of Erdős's problem for fixed small k；类型：Problem。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Anders Martinsson; Raphael Steiner | 2023-10-19

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020304 | GAX-AX_weightable_digraphs_np_characterization

When all directed cycles have length three | Informal Question (NP-characterization of weightable digraphs)

论文中定位：Informal Question (NP-characterization of weightable digraphs)；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Paul Seymour | 2025-02-09

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020305 | GAX-AX_widespread_multigraph_all

Induced subgraphs of graphs with large chromatic number. VI. Banana trees | Conjecture 1.8

论文中定位：Conjecture 1.8；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour | 2019-08-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020306 | GAX-AX_widespread_multigraph_chi_2_subdivision_free

Induced subgraphs of graphs with large chromatic number. VI. Banana trees | Conjecture 1.10

论文中定位：Conjecture 1.10；类型：Conjecture。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Alex Scott; Paul Seymour | 2019-08-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020307 | GAX-AX_zombie_number_leaf_attachment_invariance

A note on deterministic zombies | Question 4.1

论文中定位：Question 4.1；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Valentin Bartier; Laurine Bénéteau; Marthe Bonamy; Hoang La; Jonathan Narboni | 2021-06-03

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020308 | GAX-AX_zombie_number_subdivision_increase

A note on deterministic zombies | Question 4.2

论文中定位：Question 4.2；类型：Question。本卷仅收录原文入口，不以自动抽取标记认证开放状态。

仅来源定位 | extraction_role_states | Valentin Bartier; Laurine Bénéteau; Marthe Bonamy; Hoang La; Jonathan Narboni | 2021-06-03

摘录，定义可能依赖原文 | 原始来源 | metadata only

C05 Epoch / OEIS 历史定位｜42 条

V3-020309 | EPOCH-A338238_infinitely_often_primorial_conjecture

A338238 | Minimum number of rotations for a second maximum cyclic autocorrelation of the first n terms of the characteristic function of primes.

历史提出者：Andres Cicuttin；提出日期：2020-10-17；元数据声明名：A338238_infinitely_often_primorial_conjecture。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Andres Cicuttin | 2020-10-17

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020310 | EPOCH-A343812_conjecture

A343812 | a(n) = Sum_{i<=n} (A007504(n) mod prime(i)).

历史提出者：J. M. Bergot and Robert Israel；提出日期：2021-04-30；元数据声明名：A343812_conjecture。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | J. M. Bergot and Robert Israel | 2021-04-30

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020311 | EPOCH-A344989_conjecture_heuristic_zero

A344989 | Smallest number whose number of partitions into n distinct primes is n, or zero if there are no such partitions.

历史提出者：David A. Corneth；提出日期：2025-08-21；元数据声明名：A344989_conjecture_heuristic_zero。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | David A. Corneth | 2025-08-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020312 | EPOCH-A381358_limit_exists

A381358 | Row sums of irregular triangle A381587.

历史提出者：Paul D. Hanna；提出日期：2025-03-03；元数据声明名：A381358_limit_exists。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Paul D. Hanna | 2025-03-03

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020313 | EPOCH-a_never_zero

A359634 | a(0)=1 and thereafter a(n) is the length of the longest contiguous group of terms in the sequence thus far that add up to n; if no such group exists, set a(n)=0.

历史提出者：Neal Gersh Tolunsky；提出日期：2023-01-10；元数据声明名：a_never_zero。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Neal Gersh Tolunsky | 2023-01-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020314 | EPOCH-oeis_145062_conjecture_0

A145062 | Generalized Bessel numbers.

历史提出者：N. J. A. Sloane；提出日期：2016-01-28；元数据声明名：oeis_145062_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | N. J. A. Sloane | 2016-01-28

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020315 | EPOCH-oeis_193279_conjecture_0

A193279 | Number of distinct sums of distinct proper divisors of n.

历史提出者：Michael Engling；提出日期：2011-07-20；元数据声明名：oeis_193279_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Michael Engling | 2011-07-20

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020316 | EPOCH-oeis_197630_conjecture_0

A197630 | Lerch quotients of odd primes: ((Sum_{k=1..p-1} q_p(k)) - w_p)/p, where q_p(k) = (k^(p-1)-1)/p is a Fermat quotient, w_p = ((p-1)!+1)/p is a Wilson quotient, and p is the n-th prime, with n > 1.

历史提出者：Jonathan Sondow；提出日期：2011-10-18；元数据声明名：oeis_197630_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Jonathan Sondow | 2011-10-18

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020317 | EPOCH-oeis_208326_conjecture_0

A208326 | n + [nr/t] + [ns/t], where []=floor, r=5, s=(1+sqrt(5))/2, t=1/s.

历史提出者：Clark Kimberling；提出日期：2012-02-26；元数据声明名：oeis_208326_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Clark Kimberling | 2012-02-26

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020318 | EPOCH-oeis_231830_conjecture_0

A231830 | a(0) = 1; for n > 0, a(n) = 1 + 4*Product_{i=1..n-1} a(i)^2.

历史提出者：Max Alekseyev；提出日期：2023-04-21；元数据声明名：oeis_231830_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Max Alekseyev | 2023-04-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020319 | EPOCH-oeis_241898_conjecture_0

A241898 | a(n) is the largest integer such that n = a(n)^2 + ... is a decomposition of n into a sum of at most four nondecreasing squares.

历史提出者：Moshe Shmuel Newman；提出日期：2014-05-16；元数据声明名：oeis_241898_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Moshe Shmuel Newman | 2014-05-16

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020320 | EPOCH-oeis_265709_conjecture_0

A265709 | a(n) = numerator of Sum_{d|n} 1/sigma(d).

历史提出者：Jaroslav Krizek；提出日期：2015-12-24；元数据声明名：oeis_265709_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Jaroslav Krizek | 2015-12-24

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020321 | EPOCH-oeis_265710_conjecture_0

A265710 | a(n) = denominator of Sum_{d|n} 1/sigma(d).

历史提出者：Jaroslav Krizek；提出日期：2015-12-24；元数据声明名：oeis_265710_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Jaroslav Krizek | 2015-12-24

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020322 | EPOCH-oeis_275768_conjecture_0

A275768 | a(n) is the number of ways to express n = (prime(i) + prime(j))/2 when (prime(i) - prime(j))/2 also is prime.

历史提出者：Michael De Vlieger；提出日期：2017-04-30；元数据声明名：oeis_275768_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Michael De Vlieger | 2017-04-30

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020323 | EPOCH-oeis_297707_conjecture_0

A297707 | a(n) = Product_{k=1..n-1} n!k, where n!k is k-tuple factorial of n.

历史提出者：Lechoslaw Ratajczak；提出日期：2018-01-03；元数据声明名：oeis_297707_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Lechoslaw Ratajczak | 2018-01-03

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020324 | EPOCH-oeis_316774_conjecture_4

A316774 | a(n) = n for n < 2, a(n) = freq(a(n-1),n) + freq(a(n-2),n) for n >= 2, where freq(i,j) is the number of times i appears in [a(0),a(1),...,a(j-1)].

历史提出者：None；提出日期：None；元数据声明名：oeis_316774_conjecture_4。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | None | None

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020325 | EPOCH-oeis_320146_conjecture_0

A320146 | a(n) = 2*prime(n) modulo (prime(n-1) + prime(n+1)).

历史提出者：Andres Cicuttin；提出日期：2018-10-12；元数据声明名：oeis_320146_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Andres Cicuttin | 2018-10-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020326 | EPOCH-oeis_321475_conjecture_0

A321475 | Zeroless factorials (version 2): a(0) = 1, and for any n > 0, a(n) = noz(1 * noz(2 * ... * noz((n-1) * n))), where noz(n) = A004719(n) omits the zeros from n.

历史提出者：Rémy Sigrist；提出日期：2018-11-11；元数据声明名：oeis_321475_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Rémy Sigrist | 2018-11-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020327 | EPOCH-oeis_334916_conjecture_0

A334916 | a(n) is the smallest number > 1 whose base n digits yield the original number when added and multiplied left to right; or 0 if no such number exists.

历史提出者：Matej Veselovac；提出日期：2020-05-16；元数据声明名：oeis_334916_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Matej Veselovac | 2020-05-16

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020328 | EPOCH-oeis_338483_conjecture_0

A338483 | a(n) is the smallest number having n smaller numbers with the same number of divisors.

历史提出者：Ivan N.
Ianakiev；提出日期：2020-10-30；元数据声明名：oeis_338483_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Ivan N. Ianakiev | 2020-10-30

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020329 | EPOCH-oeis_340592_conjecture_0

A340592 | a(n) is the concatenation of the prime factors (with multiplicity) of n mod n.

历史提出者：J. M. Bergot and Robert
Israel；提出日期：2021-01-12；元数据声明名：oeis_340592_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | J. M. Bergot and Robert Israel | 2021-01-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020330 | EPOCH-oeis_340976_conjecture_4

A340976 | a(n) = Sum_{1 < k < n} sigma(n) mod k, where sigma = A000203.

历史提出者：M. F.
Hasler；提出日期：2021-02-01；元数据声明名：oeis_340976_conjecture_4。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | M. F. Hasler | 2021-02-01

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020331 | EPOCH-oeis_341996_conjecture_0

A341996 | a(n) = 1 if there is at least one such prime p that p^p divides the arithmetic derivative of n, A003415(n); a(0) = a(1) = 0 by convention.

历史提出者：Antti
Karttunen；提出日期：2024-01-11；元数据声明名：oeis_341996_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Antti Karttunen | 2024-01-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020332 | EPOCH-oeis_347475_conjecture_0

A347475 | Numbers k such that k and the k-th triangular number T(k) = k*(k+1)/2 have only odd digits.

历史提出者：M. F.
Hasler；提出日期：2021-11-23；元数据声明名：oeis_347475_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | M. F. Hasler | 2021-11-23

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020333 | EPOCH-oeis_351442_conjecture_0

A351442 | a(n) = A003958(sigma(n)), where A003958 is multiplicative with a(p^e) = (p-1)^e and sigma is the sum of divisors function.

历史提出者：Antti
Karttunen；提出日期：2022-02-12；元数据声明名：oeis_351442_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Antti Karttunen | 2022-02-12

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020334 | EPOCH-oeis_352965_conjecture_0

A352965 | A variant of Van Eck's sequence where we only consider prime numbers: for n >= 0, if a(n) = a(n-p) for some prime number p, take the least such p and set a(n+1) = p; otherwise a(n+1) = 0. Start with a(1) = 0.

历史提出者：Rémy
Sigrist；提出日期：2022-04-15；元数据声明名：oeis_352965_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Rémy Sigrist | 2022-04-15

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020335 | EPOCH-oeis_355228_conjecture_0

A355228 | a(n) is the smallest integer m such that there exist n of its distinct divisors (d_1, d_2, ..., d_n) with the property that m = d_1 + d_2 + ... + d_n = lcm(d_1, d_2, ..., d_n), or 0 if no such number m exists.

历史提出者：Bernard
Schott；提出日期：2022-06-25；元数据声明名：oeis_355228_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Bernard Schott | 2022-06-25

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020336 | EPOCH-oeis_364176_conjecture_0

A364176 | a(n) = (15*n)!*(5*n/2)!*(2*n)!/((15*n/2)!*(6*n)!*(5*n)!*n!).

历史提出者：Peter
Bala；提出日期：2023-07-14；元数据声明名：oeis_364176_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Peter Bala | 2023-07-14

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020337 | EPOCH-oeis_364178_conjecture_0

A364178 | a(n) = (10*n)!*(3*n)!*(n/2)!/((6*n)!*(5*n)!*(3*n/2)!*n!).

历史提出者：Peter Bala；提出日期：2023-07-14；元数据声明名：oeis_364178_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Peter Bala | 2023-07-14

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020338 | EPOCH-oeis_369462_conjecture_0

A369462 | Number of representations of 12n-1 as a sum (p*q + p*r + q*r) with three odd primes p <= q <= r.

历史提出者：Antti Karttunen；提出日期：2024-01-23；元数据声明名：oeis_369462_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Antti Karttunen | 2024-01-23

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020339 | EPOCH-oeis_376462_conjecture_0

A376462 | a(n) = Sum_{k = 0..n} binomial(n, k)^2*binomial(n+k, k)*A108625(n, n-k).

历史提出者：Peter Bala；提出日期：2024-09-24；元数据声明名：oeis_376462_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Peter Bala | 2024-09-24

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020340 | EPOCH-oeis_376930_conjecture_0

A376930 | a(0)=0, a(1)=1; for n>1, a(n) = a(n-1)+a(n-2), except where a(n-1) is a prime greater than 2, in which case a(n) = a(n-1)-a(n-2).

历史提出者：Stuart Coe；提出日期：2024-10-11；元数据声明名：oeis_376930_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Stuart Coe | 2024-10-11

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020341 | EPOCH-oeis_381159_conjecture_0

A381159 | Numbers whose prime divisors all end in the same digit.

历史提出者：A. Chironov；提出日期：2025-02-16；元数据声明名：oeis_381159_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | A. Chironov | 2025-02-16

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020342 | EPOCH-oeis_383327_conjecture_0

A383327 | a(n) is the number of occurrences of n in A049802.

历史提出者：Miles Englezou；提出日期：2025-05-10；元数据声明名：oeis_383327_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Miles Englezou | 2025-05-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020343 | EPOCH-oeis_385391_conjecture_0

A385391 | a(n) is the smallest integer k such that A384237(k) = n.

历史提出者：Michel Marcus and Juri-Stepan Gerasimov；提出日期：2025-06-27；元数据声明名：oeis_385391_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Michel Marcus and Juri-Stepan Gerasimov | 2025-06-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020344 | EPOCH-oeis_385958_conjecture_0

A385958 | a(n) is the largest prime p such that b(n) = b(n-1)*(p+1)/(p-1) is an integer (A385959), where b(0) = 1.

历史提出者：Thomas Ordowski；提出日期：2025-07-14；元数据声明名：oeis_385958_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Thomas Ordowski | 2025-07-14

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020345 | EPOCH-oeis_386660_conjecture_0

A386660 | a(n) = Sum_{k=1..n} binomial(n, k) (mod 2^k).

历史提出者：Paul D. Hanna；提出日期：2025-07-27；元数据声明名：oeis_386660_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Paul D. Hanna | 2025-07-27

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020346 | EPOCH-oeis_a122589_conjecture_0

A122589 | Expansion of 1/(1 - 11*x + 45*x^2 - 84*x^3 + 70*x^4 - 21*x^5 + x^6).

历史提出者：None；提出日期：None；元数据声明名：oeis_a122589_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | None | None

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020347 | EPOCH-oeis_a189409_conjectures

A189409 | a(n) = prime(n)#^2 + 1, where prime(n)# is the n-th primorial (A002110).

历史提出者：John M. Campbell；提出日期：2011-04-21；元数据声明名：oeis_a189409_conjectures。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | John M. Campbell | 2011-04-21

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020348 | EPOCH-oeis_a354747_conjecture_0

A354747 | Start with 2*n-1; repeatedly triple and add 2 until reaching a prime. a(n) = number of steps until reaching a prime > 2*n-1, or 0 if no prime is ever reached.

历史提出者：Felix Fröhlich；提出日期：2022-06-06；元数据声明名：oeis_a354747_conjecture_0。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Felix Fröhlich | 2022-06-06

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020349 | EPOCH-oeis_a358340_conjecture_k4

A358340 | a(n) is the smallest n-digit number whose fourth power is zeroless.

历史提出者：Mohammed Yaseen；提出日期：2022-11-10；元数据声明名：oeis_a358340_conjecture_k4。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Mohammed Yaseen | 2022-11-10

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020350 | EPOCH-oeis_a374265_conjecture_1_boundedness

A374265 | Minimized zeroless factorials.

历史提出者：Bryle Morga；提出日期：2024-07-02；元数据声明名：oeis_a374265_conjecture_1_boundedness。请核对 OEIS 当前内容与修订历史。

仅来源定位 | AI_summarized_historical_metadata | Bryle Morga | 2024-07-02

摘录，定义可能依赖原文 | 原始来源 | metadata only

C06 TOPP 来源定位 | 3 条

V3-020351 | CBC-topp-0033

Sum of Square Roots

来源定位：TOPP problem 33 (Problems/P.000033)。本卷保留题名与来源指针。

仅来源定位 | open | 原编者及问题提出者见来源 | 2025-06-19

仅题名与位置 | 原始来源 | metadata only

V3-020352 | CBC-topp-0045

Smallest Universal Set of Points for Planar Graphs

来源定位：TOPP problem 45 (Problems/P.000045)。本卷保留题名与来源指针。

仅来源定位 | open | 原编者及问题提出者见来源 | 2025-06-19

仅题名与位置 | 原始来源 | metadata only

V3-020353 | CBC-topp-0075

Edge-Coloring Geometric Graphs

来源定位：TOPP problem 75 (Problems/P.000075)。本卷保留题名与来源指针。

仅来源定位 | open | 原编者及问题提出者见来源 | 2025-06-19

仅题名与位置 | 原始来源 | metadata only

C07 Bondy-Murty 附录定位 | 7 条

V3-020354 | GBM-bm-022

Internally disjoint odd paths: is it in co-NP?

《Graph Theory》附录问题 22；法文版书页 628，PDF 页 644。

仅来源定位 | historical_mention_unreviewed | C. Thomassen | 1980

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020355 | GBM-bm-037

Constructive exponential lower bound for diagonal Ramsey numbers

《Graph Theory》附录问题 37；法文版书页 629，PDF 页 645。

仅来源定位 | historical_mention_unreviewed | P. Erdős | 1969

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020356 | GBM-bm-038

Limit of $r(k,k)^{1/k}$

《Graph Theory》附录问题 38；法文版书页 630，PDF 页 646。

仅来源定位 | historical_mention_unreviewed | P. Erdős | 1947

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020357 | GBM-bm-042

Hajós conjecture for $k = 5$ and $k = 6$

《Graph Theory》附录问题 42；法文版书页 630，PDF 页 646。

仅来源定位 | historical_mention_unreviewed | G. Hajós; P.A. Catlin | 1979

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020358 | GBM-bm-085

Cantoni's conjecture

《Graph Theory》附录问题 85；法文版书页 633，PDF 页 649。

仅来源定位 | historical_mention_unreviewed | R. Cantoni | 1950

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020359 | GBM-bm-090

Hypohamiltonian graphs of minimum degree at least 4

《Graph Theory》附录问题 90；法文版书页 634，PDF 页 650。

仅来源定位 | historical_mention_unreviewed | C. Thomassen | 1978

摘录，定义可能依赖原文 | 原始来源 | metadata only

V3-020360 | GBM-bm-091

Grötschel's conjecture on bipartite hypotraceable graphs

《Graph Theory》附录问题 91；法文版书页 634，PDF 页 650。

仅来源定位 | historical_mention_unreviewed | M. Grötschel | 1978

摘录，定义可能依赖原文 | 原始来源 | metadata only

D 参数型开放问题实例：9,342 行

本附录的实例来自 ConjectureBench 对六个原始问题族的整理。记录给出具体参数和观察日的开放范围；数值事实不等同于一项有独立名称的猜想。因此，这 9,342 行没有计入封面的 20,360 条主目录。

D01 二元常权码 | 250 行

A(n,d,w) 是长度 n、重量 w、两两 Hamming 距离至少 d 的二进制码的最大大小。best_known 为已知构造大小，upper_bound 为来源给出的上界。

原始维护者：Andries E. Brouwer

The page asks for corrections and updates at aeb@cwi.nl and names its own sources: cite A. E. Brouwer, "Bounds for binary constant weight codes", https://www.win.tue.nl/~aeb/codes/Andw.html, together with A. E. Brouwer, J. B. Shearer, N. J. A. Sloane & W. D. Smith, "A new table of constant weight codes", IEEE Trans. Inform. Theory 36 (1990) 1334-1380, and E. Agrell, A. Vardy & K. Zeger, "Upper bounds for constant-weight codes", IEEE Trans. Inform. Theory 46 (2000) 2373-2395, for the entries taken from them.

原始问题族入口

V3P-000001 | aebcw-a-16-6-7

Constant weight code A(16, 6, 7)

a code of size 109 is known, and no such code can have more than 122 words.

来源状态 open | 2026-07-23 | 来源

V3P-000002 | aebcw-a-16-6-8

Constant weight code A(16, 6, 8)

a code of size 120 is known, and no such code can have more than 138 words.

来源状态 open | 2026-07-23 | 来源

V3P-000003 | aebcw-a-17-6-6

Constant weight code A(17, 6, 6)

a code of size 113 is known, and no such code can have more than 124 words.

来源状态 open | 2026-07-23 | 来源

V3P-000004 | aebcw-a-17-6-7

Constant weight code A(17, 6, 7)

a code of size 166 is known, and no such code can have more than 206 words.

来源状态 open | 2026-07-23 | 来源

V3P-000005 | aebcw-a-17-6-8

Constant weight code A(17, 6, 8)

a code of size 184 is known, and no such code can have more than 259 words.

来源状态 open | 2026-07-23 | 来源

V3P-000006 | aebcw-a-18-6-5

Constant weight code A(18, 6, 5)

a code of size 69 is known, and no such code can have more than 72 words.

来源状态 open | 2026-07-23 | 来源

V3P-000007 | aebcw-a-18-6-6

Constant weight code A(18, 6, 6)

a code of size 133 is known, and no such code can have more than 186 words.

来源状态 open | 2026-07-23 | 来源

V3P-000008 | aebcw-a-18-6-7

Constant weight code A(18, 6, 7)

a code of size 243 is known, and no such code can have more than 312 words.

来源状态 open | 2026-07-23 | 来源

V3P-000009 | aebcw-a-18-6-8

Constant weight code A(18, 6, 8)

a code of size 260 is known, and no such code can have more than 427 words.

来源状态 open | 2026-07-23 | 来源

V3P-000010 | aebcw-a-18-6-9

Constant weight code A(18, 6, 9)

a code of size 320 is known, and no such code can have more than 424 words.

来源状态 open | 2026-07-23 | 来源

V3P-000011 | aebcw-a-18-8-8

Constant weight code A(18, 8, 8)

a code of size 46 is known, and no such code can have more than 49 words.

来源状态 open | 2026-07-23 | 来源

V3P-000012 | aebcw-a-18-8-9

Constant weight code A(18, 8, 9)

a code of size 48 is known, and no such code can have more than 58 words.

来源状态 open | 2026-07-23 | 来源

V3P-000013 | aebcw-a-19-6-5

Constant weight code A(19, 6, 5)

a code of size 76 is known, and no such code can have more than 83 words.

来源状态 open | 2026-07-23 | 来源

V3P-000014 | aebcw-a-19-6-6

Constant weight code A(19, 6, 6)

a code of size 172 is known, and no such code can have more than 228 words.

来源状态 open | 2026-07-23 | 来源

V3P-000015 | aebcw-a-19-6-7

Constant weight code A(19, 6, 7)

a code of size 338 is known, and no such code can have more than 463 words.

来源状态 open | 2026-07-23 | 来源

V3P-000016 | aebcw-a-19-6-8

Constant weight code A(19, 6, 8)

a code of size 408 is known, and no such code can have more than 693 words.

来源状态 open | 2026-07-23 | 来源

V3P-000017 | aebcw-a-19-6-9

Constant weight code A(19, 6, 9)

a code of size 504 is known, and no such code can have more than 789 words.

来源状态 open | 2026-07-23 | 来源

V3P-000018 | aebcw-a-19-8-9

Constant weight code A(19, 8, 9)

a code of size 88 is known, and no such code can have more than 103 words.

来源状态 open | 2026-07-23 | 来源

V3P-000019 | aebcw-a-20-6-10

Constant weight code A(20, 6, 10)

a code of size 944 is known, and no such code can have more than 1420 words.

来源状态 open | 2026-07-23 | 来源

V3P-000020 | aebcw-a-20-6-5

Constant weight code A(20, 6, 5)

a code of size 90 is known, and no such code can have more than 100 words.

来源状态 open | 2026-07-23 | 来源

V3P-000021 | aebcw-a-20-6-6

Constant weight code A(20, 6, 6)

a code of size 232 is known, and no such code can have more than 276 words.

来源状态 open | 2026-07-23 | 来源

V3P-000022 | aebcw-a-20-6-7

Constant weight code A(20, 6, 7)

a code of size 462 is known, and no such code can have more than 651 words.

来源状态 open | 2026-07-23 | 来源

V3P-000023 | aebcw-a-20-6-8

Constant weight code A(20, 6, 8)

a code of size 588 is known, and no such code can have more than 1084 words.

来源状态 open | 2026-07-23 | 来源

V3P-000024 | aebcw-a-20-6-9

Constant weight code A(20, 6, 9)

a code of size 832 is known, and no such code can have more than 1363 words.

来源状态 open | 2026-07-23 | 来源

V3P-000025 | aebcw-a-20-8-10

Constant weight code A(20, 8, 10)

a code of size 176 is known, and no such code can have more than 206 words.

来源状态 open | 2026-07-23 | 来源

V3P-000026 | aebcw-a-20-8-9

Constant weight code A(20, 8, 9)

a code of size 160 is known, and no such code can have more than 173 words.

来源状态 open | 2026-07-23 | 来源

V3P-000027 | aebcw-a-21-10-10

Constant weight code A(21, 10, 10)

a code of size 38 is known, and no such code can have more than 42 words.

来源状态 open | 2026-07-23 | 来源

V3P-000028 | aebcw-a-21-10-9

Constant weight code A(21, 10, 9)

a code of size 27 is known, and no such code can have more than 35 words.

来源状态 open | 2026-07-23 | 来源

V3P-000029 | aebcw-a-21-6-10

Constant weight code A(21, 6, 10)

a code of size 1474 is known, and no such code can have more than 2685 words.

来源状态 open | 2026-07-23 | 来源

V3P-000030 | aebcw-a-21-6-5

Constant weight code A(21, 6, 5)

a code of size 108 is known, and no such code can have more than 126 words.

来源状态 open | 2026-07-23 | 来源

V3P-000031 | aebcw-a-21-6-6

Constant weight code A(21, 6, 6)

a code of size 273 is known, and no such code can have more than 350 words.

来源状态 open | 2026-07-23 | 来源

V3P-000032 | aebcw-a-21-6-7

Constant weight code A(21, 6, 7)

a code of size 570 is known, and no such code can have more than 828 words.

来源状态 open | 2026-07-23 | 来源

V3P-000033 | aebcw-a-21-6-8

Constant weight code A(21, 6, 8)

a code of size 775 is known, and no such code can have more than 1665 words.

来源状态 open | 2026-07-23 | 来源

V3P-000034 | aebcw-a-21-6-9

Constant weight code A(21, 6, 9)

a code of size 1186 is known, and no such code can have more than 2328 words.

来源状态 open | 2026-07-23 | 来源

V3P-000035 | aebcw-a-21-8-10

Constant weight code A(21, 8, 10)

a code of size 336 is known, and no such code can have more than 363 words.

来源状态 open | 2026-07-23 | 来源

V3P-000036 | aebcw-a-21-8-9

Constant weight code A(21, 8, 9)

a code of size 280 is known, and no such code can have more than 294 words.

来源状态 open | 2026-07-23 | 来源

V3P-000037 | aebcw-a-22-10-10

Constant weight code A(22, 10, 10)

a code of size 46 is known, and no such code can have more than 71 words.

来源状态 open | 2026-07-23 | 来源

V3P-000038 | aebcw-a-22-10-11

Constant weight code A(22, 10, 11)

a code of size 46 is known, and no such code can have more than 79 words.

来源状态 open | 2026-07-23 | 来源

V3P-000039 | aebcw-a-22-10-8

Constant weight code A(22, 10, 8)

a code of size 25 is known, and no such code can have more than 33 words.

来源状态 open | 2026-07-23 | 来源

V3P-000040 | aebcw-a-22-10-9

Constant weight code A(22, 10, 9)

a code of size 35 is known, and no such code can have more than 51 words.

来源状态 open | 2026-07-23 | 来源

V3P-000041 | aebcw-a-22-6-10

Constant weight code A(22, 6, 10)

a code of size 2183 is known, and no such code can have more than 4415 words.

来源状态 open | 2026-07-23 | 来源

V3P-000042 | aebcw-a-22-6-11

Constant weight code A(22, 6, 11)

a code of size 2636 is known, and no such code can have more than 5033 words.

来源状态 open | 2026-07-23 | 来源

V3P-000043 | aebcw-a-22-6-5

Constant weight code A(22, 6, 5)

a code of size 132 is known, and no such code can have more than 136 words.

来源状态 open | 2026-07-23 | 来源

V3P-000044 | aebcw-a-22-6-6

Constant weight code A(22, 6, 6)

a code of size 343 is known, and no such code can have more than 462 words.

来源状态 open | 2026-07-23 | 来源

V3P-000045 | aebcw-a-22-6-7

Constant weight code A(22, 6, 7)

a code of size 759 is known, and no such code can have more than 1100 words.

来源状态 open | 2026-07-23 | 来源

V3P-000046 | aebcw-a-22-6-8

Constant weight code A(22, 6, 8)

a code of size 1144 is known, and no such code can have more than 2277 words.

来源状态 open | 2026-07-23 | 来源

V3P-000047 | aebcw-a-22-6-9

Constant weight code A(22, 6, 9)

a code of size 1818 is known, and no such code can have more than 3736 words.

来源状态 open | 2026-07-23 | 来源

V3P-000048 | aebcw-a-22-8-9

Constant weight code A(22, 8, 9)

a code of size 292 is known, and no such code can have more than 440 words.

来源状态 open | 2026-07-23 | 来源

V3P-000049 | aebcw-a-23-10-10

Constant weight code A(23, 10, 10)

a code of size 54 is known, and no such code can have more than 116 words.

来源状态 open | 2026-07-23 | 来源

V3P-000050 | aebcw-a-23-10-11

Constant weight code A(23, 10, 11)

a code of size 65 is known, and no such code can have more than 135 words.

来源状态 open | 2026-07-23 | 来源

V3P-000051 | aebcw-a-23-10-8

Constant weight code A(23, 10, 8)

a code of size 33 is known, and no such code can have more than 46 words.

来源状态 open | 2026-07-23 | 来源

V3P-000052 | aebcw-a-23-10-9

Constant weight code A(23, 10, 9)

a code of size 45 is known, and no such code can have more than 78 words.

来源状态 open | 2026-07-23 | 来源

V3P-000053 | aebcw-a-23-6-10

Constant weight code A(23, 6, 10)

a code of size 2990 is known, and no such code can have more than 7521 words.

来源状态 open | 2026-07-23 | 来源

V3P-000054 | aebcw-a-23-6-11

Constant weight code A(23, 6, 11)

a code of size 3585 is known, and no such code can have more than 7953 words.

来源状态 open | 2026-07-23 | 来源

V3P-000055 | aebcw-a-23-6-5

Constant weight code A(23, 6, 5)

a code of size 147 is known, and no such code can have more than 170 words.

来源状态 open | 2026-07-23 | 来源

V3P-000056 | aebcw-a-23-6-6

Constant weight code A(23, 6, 6)

a code of size 399 is known, and no such code can have more than 521 words.

来源状态 open | 2026-07-23 | 来源

V3P-000057 | aebcw-a-23-6-7

Constant weight code A(23, 6, 7)

a code of size 969 is known, and no such code can have more than 1518 words.

来源状态 open | 2026-07-23 | 来源

V3P-000058 | aebcw-a-23-6-8

Constant weight code A(23, 6, 8)

a code of size 1439 is known, and no such code can have more than 3162 words.

来源状态 open | 2026-07-23 | 来源

V3P-000059 | aebcw-a-23-6-9

Constant weight code A(23, 6, 9)

a code of size 2278 is known, and no such code can have more than 5819 words.

来源状态 open | 2026-07-23 | 来源

V3P-000060 | aebcw-a-23-8-10

Constant weight code A(23, 8, 10)

a code of size 648 is known, and no such code can have more than 1012 words.

来源状态 open | 2026-07-23 | 来源

V3P-000061 | aebcw-a-23-8-6

Constant weight code A(23, 8, 6)

a code of size 77 is known, and no such code can have more than 80 words.

来源状态 open | 2026-07-23 | 来源

V3P-000062 | aebcw-a-23-8-9

Constant weight code A(23, 8, 9)

a code of size 412 is known, and no such code can have more than 662 words.

来源状态 open | 2026-07-23 | 来源

V3P-000063 | aebcw-a-24-10-10

Constant weight code A(24, 10, 10)

a code of size 72 is known, and no such code can have more than 170 words.

来源状态 open | 2026-07-23 | 来源

V3P-000064 | aebcw-a-24-10-11

Constant weight code A(24, 10, 11)

a code of size 95 is known, and no such code can have more than 222 words.

来源状态 open | 2026-07-23 | 来源

V3P-000065 | aebcw-a-24-10-12

Constant weight code A(24, 10, 12)

a code of size 123 is known, and no such code can have more than 246 words.

来源状态 open | 2026-07-23 | 来源

V3P-000066 | aebcw-a-24-10-8

Constant weight code A(24, 10, 8)

a code of size 38 is known, and no such code can have more than 60 words.

来源状态 open | 2026-07-23 | 来源

V3P-000067 | aebcw-a-24-10-9

Constant weight code A(24, 10, 9)

a code of size 56 is known, and no such code can have more than 116 words.

来源状态 open | 2026-07-23 | 来源

V3P-000068 | aebcw-a-24-6-10

Constant weight code A(24, 6, 10)

a code of size 4224 is known, and no such code can have more than 12186 words.

来源状态 open | 2026-07-23 | 来源

V3P-000069 | aebcw-a-24-6-5

Constant weight code A(24, 6, 5)

a code of size 168 is known, and no such code can have more than 192 words.

来源状态 open | 2026-07-23 | 来源

V3P-000070 | aebcw-a-24-6-6

Constant weight code A(24, 6, 6)

a code of size 532 is known, and no such code can have more than 680 words.

来源状态 open | 2026-07-23 | 来源

V3P-000071 | aebcw-a-24-6-7

Constant weight code A(24, 6, 7)

a code of size 1368 is known, and no such code can have more than 1786 words.

来源状态 open | 2026-07-23 | 来源

V3P-000072 | aebcw-a-24-6-8

Constant weight code A(24, 6, 8)

a code of size 1882 is known, and no such code can have more than 4554 words.

来源状态 open | 2026-07-23 | 来源

V3P-000073 | aebcw-a-24-6-9

Constant weight code A(24, 6, 9)

a code of size 3080 is known, and no such code can have more than 8432 words.

来源状态 open | 2026-07-23 | 来源

V3P-000074 | aebcw-a-24-8-10

Constant weight code A(24, 8, 10)

a code of size 960 is known, and no such code can have more than 1551 words.

来源状态 open | 2026-07-23 | 来源

V3P-000075 | aebcw-a-24-8-11

Constant weight code A(24, 8, 11)

a code of size 1378 is known, and no such code can have more than 2142 words.

来源状态 open | 2026-07-23 | 来源

V3P-000076 | aebcw-a-24-8-6

Constant weight code A(24, 8, 6)

a code of size 78 is known, and no such code can have more than 92 words.

来源状态 open | 2026-07-23 | 来源

V3P-000077 | aebcw-a-24-8-7

Constant weight code A(24, 8, 7)

a code of size 253 is known, and no such code can have more than 274 words.

来源状态 open | 2026-07-23 | 来源

V3P-000078 | aebcw-a-24-8-9

Constant weight code A(24, 8, 9)

a code of size 656 is known, and no such code can have more than 968 words.

来源状态 open | 2026-07-23 | 来源

V3P-000079 | aebcw-a-25-10-10

Constant weight code A(25, 10, 10)

a code of size 100 is known, and no such code can have more than 262 words.

来源状态 open | 2026-07-23 | 来源

V3P-000080 | aebcw-a-25-10-11

Constant weight code A(25, 10, 11)

a code of size 125 is known, and no such code can have more than 379 words.

来源状态 open | 2026-07-23 | 来源

V3P-000081 | aebcw-a-25-10-12

Constant weight code A(25, 10, 12)

a code of size 137 is known, and no such code can have more than 433 words.

来源状态 open | 2026-07-23 | 来源

V3P-000082 | aebcw-a-25-10-7

Constant weight code A(25, 10, 7)

a code of size 28 is known, and no such code can have more than 32 words.

来源状态 open | 2026-07-23 | 来源

V3P-000083 | aebcw-a-25-10-8

Constant weight code A(25, 10, 8)

a code of size 51 is known, and no such code can have more than 75 words.

来源状态 open | 2026-07-23 | 来源

V3P-000084 | aebcw-a-25-10-9

Constant weight code A(25, 10, 9)

a code of size 72 is known, and no such code can have more than 157 words.

来源状态 open | 2026-07-23 | 来源

V3P-000085 | aebcw-a-25-12-10

Constant weight code A(25, 12, 10)

a code of size 28 is known, and no such code can have more than 36 words.

来源状态 open | 2026-07-23 | 来源

V3P-000086 | aebcw-a-25-12-11

Constant weight code A(25, 12, 11)

a code of size 36 is known, and no such code can have more than 42 words.

来源状态 open | 2026-07-23 | 来源

V3P-000087 | aebcw-a-25-6-6

Constant weight code A(25, 6, 6)

a code of size 700 is known, and no such code can have more than 800 words.

来源状态 open | 2026-07-23 | 来源

V3P-000088 | aebcw-a-25-6-7

Constant weight code A(25, 6, 7)

a code of size 1900 is known, and no such code can have more than 2428 words.

来源状态 open | 2026-07-23 | 来源

V3P-000089 | aebcw-a-25-6-8

Constant weight code A(25, 6, 8)

a code of size 2590 is known, and no such code can have more than 5581 words.

来源状态 open | 2026-07-23 | 来源

V3P-000090 | aebcw-a-25-6-9

Constant weight code A(25, 6, 9)

a code of size 4260 is known, and no such code can have more than 12620 words.

来源状态 open | 2026-07-23 | 来源

V3P-000091 | aebcw-a-25-8-10

Constant weight code A(25, 8, 10)

a code of size 1248 is known, and no such code can have more than 2333 words.

来源状态 open | 2026-07-23 | 来源

V3P-000092 | aebcw-a-25-8-11

Constant weight code A(25, 8, 11)

a code of size 1702 is known, and no such code can have more than 3422 words.

来源状态 open | 2026-07-23 | 来源

V3P-000093 | aebcw-a-25-8-12

Constant weight code A(25, 8, 12)

a code of size 2610 is known, and no such code can have more than 4087 words.

来源状态 open | 2026-07-23 | 来源

V3P-000094 | aebcw-a-25-8-7

Constant weight code A(25, 8, 7)

a code of size 255 is known, and no such code can have more than 328 words.

来源状态 open | 2026-07-23 | 来源

V3P-000095 | aebcw-a-25-8-8

Constant weight code A(25, 8, 8)

a code of size 759 is known, and no such code can have more than 850 words.

来源状态 open | 2026-07-23 | 来源

V3P-000096 | aebcw-a-25-8-9

Constant weight code A(25, 8, 9)

a code of size 837 is known, and no such code can have more than 1366 words.

来源状态 open | 2026-07-23 | 来源

V3P-000097 | aebcw-a-26-10-10

Constant weight code A(26, 10, 10)

a code of size 130 is known, and no such code can have more than 397 words.

来源状态 open | 2026-07-23 | 来源

V3P-000098 | aebcw-a-26-10-11

Constant weight code A(26, 10, 11)

a code of size 168 is known, and no such code can have more than 565 words.

来源状态 open | 2026-07-23 | 来源

V3P-000099 | aebcw-a-26-10-12

Constant weight code A(26, 10, 12)

a code of size 208 is known, and no such code can have more than 691 words.

来源状态 open | 2026-07-23 | 来源

V3P-000100 | aebcw-a-26-10-13

Constant weight code A(26, 10, 13)

a code of size 225 is known, and no such code can have more than 754 words.

来源状态 open | 2026-07-23 | 来源

V3P-000101 | aebcw-a-26-10-7

Constant weight code A(26, 10, 7)

a code of size 30 is known, and no such code can have more than 36 words.

来源状态 open | 2026-07-23 | 来源

V3P-000102 | aebcw-a-26-10-8

Constant weight code A(26, 10, 8)

a code of size 55 is known, and no such code can have more than 104 words.

来源状态 open | 2026-07-23 | 来源

V3P-000103 | aebcw-a-26-10-9

Constant weight code A(26, 10, 9)

a code of size 91 is known, and no such code can have more than 213 words.

来源状态 open | 2026-07-23 | 来源

V3P-000104 | aebcw-a-26-12-10

Constant weight code A(26, 12, 10)

a code of size 33 is known, and no such code can have more than 37 words.

来源状态 open | 2026-07-23 | 来源

V3P-000105 | aebcw-a-26-12-11

Constant weight code A(26, 12, 11)

a code of size 41 is known, and no such code can have more than 66 words.

来源状态 open | 2026-07-23 | 来源

V3P-000106 | aebcw-a-26-12-12

Constant weight code A(26, 12, 12)

a code of size 54 is known, and no such code can have more than 83 words.

来源状态 open | 2026-07-23 | 来源

V3P-000107 | aebcw-a-26-12-13

Constant weight code A(26, 12, 13)

a code of size 58 is known, and no such code can have more than 91 words.

来源状态 open | 2026-07-23 | 来源

V3P-000108 | aebcw-a-26-6-7

Constant weight code A(26, 6, 7)

a code of size 2600 is known, and no such code can have more than 2971 words.

来源状态 open | 2026-07-23 | 来源

V3P-000109 | aebcw-a-26-6-8

Constant weight code A(26, 6, 8)

a code of size 3532 is known, and no such code can have more than 7891 words.

来源状态 open | 2026-07-23 | 来源

V3P-000110 | aebcw-a-26-8-10

Constant weight code A(26, 8, 10)

a code of size 1519 is known, and no such code can have more than 3496 words.

来源状态 open | 2026-07-23 | 来源

V3P-000111 | aebcw-a-26-8-11

Constant weight code A(26, 8, 11)

a code of size 2037 is known, and no such code can have more than 5208 words.

来源状态 open | 2026-07-23 | 来源

V3P-000112 | aebcw-a-26-8-12

Constant weight code A(26, 8, 12)

a code of size 3070 is known, and no such code can have more than 6741 words.

来源状态 open | 2026-07-23 | 来源

V3P-000113 | aebcw-a-26-8-13

Constant weight code A(26, 8, 13)

a code of size 3588 is known, and no such code can have more than 7080 words.

来源状态 open | 2026-07-23 | 来源

V3P-000114 | aebcw-a-26-8-7

Constant weight code A(26, 8, 7)

a code of size 259 is known, and no such code can have more than 371 words.

来源状态 open | 2026-07-23 | 来源

V3P-000115 | aebcw-a-26-8-8

Constant weight code A(26, 8, 8)

a code of size 763 is known, and no such code can have more than 1066 words.

来源状态 open | 2026-07-23 | 来源

V3P-000116 | aebcw-a-26-8-9

Constant weight code A(26, 8, 9)

a code of size 887 is known, and no such code can have more than 1901 words.

来源状态 open | 2026-07-23 | 来源

V3P-000117 | aebcw-a-27-10-10

Constant weight code A(27, 10, 10)

a code of size 162 is known, and no such code can have more than 555 words.

来源状态 open | 2026-07-23 | 来源

V3P-000118 | aebcw-a-27-10-11

Constant weight code A(27, 10, 11)

a code of size 243 is known, and no such code can have more than 871 words.

来源状态 open | 2026-07-23 | 来源

V3P-000119 | aebcw-a-27-10-12

Constant weight code A(27, 10, 12)

a code of size 351 is known, and no such code can have more than 1190 words.

来源状态 open | 2026-07-23 | 来源

V3P-000120 | aebcw-a-27-10-13

Constant weight code A(27, 10, 13)

a code of size 420 is known, and no such code can have more than 1406 words.

来源状态 open | 2026-07-23 | 来源

V3P-000121 | aebcw-a-27-10-7

Constant weight code A(27, 10, 7)

a code of size 36 is known, and no such code can have more than 48 words.

来源状态 open | 2026-07-23 | 来源

V3P-000122 | aebcw-a-27-10-8

Constant weight code A(27, 10, 8)

a code of size 66 is known, and no such code can have more than 121 words.

来源状态 open | 2026-07-23 | 来源

V3P-000123 | aebcw-a-27-10-9

Constant weight code A(27, 10, 9)

a code of size 118 is known, and no such code can have more than 291 words.

来源状态 open | 2026-07-23 | 来源

V3P-000124 | aebcw-a-27-12-10

Constant weight code A(27, 12, 10)

a code of size 39 is known, and no such code can have more than 58 words.

来源状态 open | 2026-07-23 | 来源

V3P-000125 | aebcw-a-27-12-11

Constant weight code A(27, 12, 11)

a code of size 54 is known, and no such code can have more than 90 words.

来源状态 open | 2026-07-23 | 来源

V3P-000126 | aebcw-a-27-12-12

Constant weight code A(27, 12, 12)

a code of size 82 is known, and no such code can have more than 139 words.

来源状态 open | 2026-07-23 | 来源

V3P-000127 | aebcw-a-27-12-13

Constant weight code A(27, 12, 13)

a code of size 86 is known, and no such code can have more than 155 words.

来源状态 open | 2026-07-23 | 来源

V3P-000128 | aebcw-a-27-14-12

Constant weight code A(27, 14, 12)

a code of size 19 is known, and no such code can have more than 20 words.

来源状态 open | 2026-07-23 | 来源

V3P-000129 | aebcw-a-27-6-5

Constant weight code A(27, 6, 5)

a code of size 260 is known, and no such code can have more than 280 words.

来源状态 open | 2026-07-23 | 来源

V3P-000130 | aebcw-a-27-6-8

Constant weight code A(27, 6, 8)

a code of size 4786 is known, and no such code can have more than 10027 words.

来源状态 open | 2026-07-23 | 来源

V3P-000131 | aebcw-a-27-8-10

Constant weight code A(27, 8, 10)

a code of size 1721 is known, and no such code can have more than 4986 words.

来源状态 open | 2026-07-23 | 来源

V3P-000132 | aebcw-a-27-8-11

Constant weight code A(27, 8, 11)

a code of size 2404 is known, and no such code can have more than 7754 words.

来源状态 open | 2026-07-23 | 来源

V3P-000133 | aebcw-a-27-8-12

Constant weight code A(27, 8, 12)

a code of size 3335 is known, and no such code can have more than 10460 words.

来源状态 open | 2026-07-23 | 来源

V3P-000134 | aebcw-a-27-8-13

Constant weight code A(27, 8, 13)

a code of size 4094 is known, and no such code can have more than 11897 words.

来源状态 open | 2026-07-23 | 来源

V3P-000135 | aebcw-a-27-8-5

Constant weight code A(27, 8, 5)

a code of size 31 is known, and no such code can have more than 32 words.

来源状态 open | 2026-07-23 | 来源

V3P-000136 | aebcw-a-27-8-6

Constant weight code A(27, 8, 6)

a code of size 130 is known, and no such code can have more than 135 words.

来源状态 open | 2026-07-23 | 来源

V3P-000137 | aebcw-a-27-8-7

Constant weight code A(27, 8, 7)

a code of size 303 is known, and no such code can have more than 500 words.

来源状态 open | 2026-07-23 | 来源

V3P-000138 | aebcw-a-27-8-8

Constant weight code A(27, 8, 8)

a code of size 769 is known, and no such code can have more than 1252 words.

来源状态 open | 2026-07-23 | 来源

V3P-000139 | aebcw-a-27-8-9

Constant weight code A(27, 8, 9)

a code of size 1025 is known, and no such code can have more than 2616 words.

来源状态 open | 2026-07-23 | 来源

V3P-000140 | aebcw-a-28-10-10

Constant weight code A(28, 10, 10)

a code of size 219 is known, and no such code can have more than 785 words.

来源状态 open | 2026-07-23 | 来源

V3P-000141 | aebcw-a-28-10-11

Constant weight code A(28, 10, 11)

a code of size 308 is known, and no such code can have more than 1351 words.

来源状态 open | 2026-07-23 | 来源

V3P-000142 | aebcw-a-28-10-12

Constant weight code A(28, 10, 12)

a code of size 390 is known, and no such code can have more than 1977 words.

来源状态 open | 2026-07-23 | 来源

V3P-000143 | aebcw-a-28-10-13

Constant weight code A(28, 10, 13)

a code of size 756 is known, and no such code can have more than 2438 words.

来源状态 open | 2026-07-23 | 来源

V3P-000144 | aebcw-a-28-10-14

Constant weight code A(28, 10, 14)

a code of size 812 is known, and no such code can have more than 2628 words.

来源状态 open | 2026-07-23 | 来源

V3P-000145 | aebcw-a-28-10-7

Constant weight code A(28, 10, 7)

a code of size 38 is known, and no such code can have more than 56 words.

来源状态 open | 2026-07-23 | 来源

V3P-000146 | aebcw-a-28-10-8

Constant weight code A(28, 10, 8)

a code of size 81 is known, and no such code can have more than 168 words.

来源状态 open | 2026-07-23 | 来源

V3P-000147 | aebcw-a-28-10-9

Constant weight code A(28, 10, 9)

a code of size 140 is known, and no such code can have more than 376 words.

来源状态 open | 2026-07-23 | 来源

V3P-000148 | aebcw-a-28-12-10

Constant weight code A(28, 12, 10)

a code of size 56 is known, and no such code can have more than 87 words.

来源状态 open | 2026-07-23 | 来源

V3P-000149 | aebcw-a-28-12-11

Constant weight code A(28, 12, 11)

a code of size 65 is known, and no such code can have more than 147 words.

来源状态 open | 2026-07-23 | 来源

V3P-000150 | aebcw-a-28-12-12

Constant weight code A(28, 12, 12)

a code of size 119 is known, and no such code can have more than 198 words.

来源状态 open | 2026-07-23 | 来源

V3P-000151 | aebcw-a-28-12-13

Constant weight code A(28, 12, 13)

a code of size 112 is known, and no such code can have more than 244 words.

来源状态 open | 2026-07-23 | 来源

V3P-000152 | aebcw-a-28-12-14

Constant weight code A(28, 12, 14)

a code of size 172 is known, and no such code can have more than 264 words.

来源状态 open | 2026-07-23 | 来源

V3P-000153 | aebcw-a-28-12-9

Constant weight code A(28, 12, 9)

a code of size 39 is known, and no such code can have more than 45 words.

来源状态 open | 2026-07-23 | 来源

V3P-000154 | aebcw-a-28-6-5

Constant weight code A(28, 6, 5)

a code of size 280 is known, and no such code can have more than 302 words.

来源状态 open | 2026-07-23 | 来源

V3P-000155 | aebcw-a-28-6-6

Constant weight code A(28, 6, 6)

a code of size 1170 is known, and no such code can have more than 1306 words.

来源状态 open | 2026-07-23 | 来源

V3P-000156 | aebcw-a-28-8-10

Constant weight code A(28, 8, 10)

a code of size 2067 is known, and no such code can have more than 7016 words.

来源状态 open | 2026-07-23 | 来源

V3P-000157 | aebcw-a-28-8-11

Constant weight code A(28, 8, 11)

a code of size 3773 is known, and no such code can have more than 11896 words.

来源状态 open | 2026-07-23 | 来源

V3P-000158 | aebcw-a-28-8-12

Constant weight code A(28, 8, 12)

a code of size 4927 is known, and no such code can have more than 17008 words.

来源状态 open | 2026-07-23 | 来源

V3P-000159 | aebcw-a-28-8-6

Constant weight code A(28, 8, 6)

a code of size 131 is known, and no such code can have more than 149 words.

来源状态 open | 2026-07-23 | 来源

V3P-000160 | aebcw-a-28-8-7

Constant weight code A(28, 8, 7)

a code of size 318 is known, and no such code can have more than 540 words.

来源状态 open | 2026-07-23 | 来源

V3P-000161 | aebcw-a-28-8-8

Constant weight code A(28, 8, 8)

a code of size 1057 is known, and no such code can have more than 1750 words.

来源状态 open | 2026-07-23 | 来源

V3P-000162 | aebcw-a-28-8-9

Constant weight code A(28, 8, 9)

a code of size 1333 is known, and no such code can have more than 3855 words.

来源状态 open | 2026-07-23 | 来源

V3P-000163 | aebcw-a-29-10-10

Constant weight code A(29, 10, 10)

a code of size 266 is known, and no such code can have more than 1090 words.

来源状态 open | 2026-07-23 | 来源

V3P-000164 | aebcw-a-29-10-11

Constant weight code A(29, 10, 11)

a code of size 406 is known, and no such code can have more than 2036 words.

来源状态 open | 2026-07-23 | 来源

V3P-000165 | aebcw-a-29-10-12

Constant weight code A(29, 10, 12)

a code of size 543 is known, and no such code can have more than 3091 words.

来源状态 open | 2026-07-23 | 来源

V3P-000166 | aebcw-a-29-10-13

Constant weight code A(29, 10, 13)

a code of size 756 is known, and no such code can have more than 4282 words.

来源状态 open | 2026-07-23 | 来源

V3P-000167 | aebcw-a-29-10-14

Constant weight code A(29, 10, 14)

a code of size 1458 is known, and no such code can have more than 4927 words.

来源状态 open | 2026-07-23 | 来源

V3P-000168 | aebcw-a-29-10-7

Constant weight code A(29, 10, 7)

a code of size 42 is known, and no such code can have more than 66 words.

来源状态 open | 2026-07-23 | 来源

V3P-000169 | aebcw-a-29-10-8

Constant weight code A(29, 10, 8)

a code of size 91 is known, and no such code can have more than 203 words.

来源状态 open | 2026-07-23 | 来源

V3P-000170 | aebcw-a-29-10-9

Constant weight code A(29, 10, 9)

a code of size 168 is known, and no such code can have more than 523 words.

来源状态 open | 2026-07-23 | 来源

V3P-000171 | aebcw-a-29-12-10

Constant weight code A(29, 12, 10)

a code of size 66 is known, and no such code can have more than 126 words.

来源状态 open | 2026-07-23 | 来源

V3P-000172 | aebcw-a-29-12-11

Constant weight code A(29, 12, 11)

a code of size 87 is known, and no such code can have more than 197 words.

来源状态 open | 2026-07-23 | 来源

V3P-000173 | aebcw-a-29-12-12

Constant weight code A(29, 12, 12)

a code of size 119 is known, and no such code can have more than 298 words.

来源状态 open | 2026-07-23 | 来源

V3P-000174 | aebcw-a-29-12-13

Constant weight code A(29, 12, 13)

a code of size 146 is known, and no such code can have more than 441 words.

来源状态 open | 2026-07-23 | 来源

V3P-000175 | aebcw-a-29-12-14

Constant weight code A(29, 12, 14)

a code of size 173 is known, and no such code can have more than 492 words.

来源状态 open | 2026-07-23 | 来源

V3P-000176 | aebcw-a-29-12-8

Constant weight code A(29, 12, 8)

a code of size 22 is known, and no such code can have more than 26 words.

来源状态 open | 2026-07-23 | 来源

V3P-000177 | aebcw-a-29-12-9

Constant weight code A(29, 12, 9)

a code of size 42 is known, and no such code can have more than 59 words.

来源状态 open | 2026-07-23 | 来源

V3P-000178 | aebcw-a-29-14-11

Constant weight code A(29, 14, 11)

a code of size 28 is known, and no such code can have more than 29 words.

来源状态 open | 2026-07-23 | 来源

V3P-000179 | aebcw-a-29-14-12

Constant weight code A(29, 14, 12)

a code of size 29 is known, and no such code can have more than 47 words.

来源状态 open | 2026-07-23 | 来源

V3P-000180 | aebcw-a-29-14-13

Constant weight code A(29, 14, 13)

a code of size 35 is known, and no such code can have more than 50 words.

来源状态 open | 2026-07-23 | 来源

V3P-000181 | aebcw-a-29-6-5

Constant weight code A(29, 6, 5)

a code of size 315 is known, and no such code can have more than 364 words.

来源状态 open | 2026-07-23 | 来源

V3P-000182 | aebcw-a-29-6-6

Constant weight code A(29, 6, 6)

a code of size 1197 is known, and no such code can have more than 1459 words.

来源状态 open | 2026-07-23 | 来源

V3P-000183 | aebcw-a-29-6-7

Constant weight code A(29, 6, 7)

a code of size 4680 is known, and no such code can have more than 5410 words.

来源状态 open | 2026-07-23 | 来源

V3P-000184 | aebcw-a-30-10-10

Constant weight code A(30, 10, 10)

a code of size 322 is known, and no such code can have more than 1569 words.

来源状态 open | 2026-07-23 | 来源

V3P-000185 | aebcw-a-30-10-11

Constant weight code A(30, 10, 11)

a code of size 504 is known, and no such code can have more than 2924 words.

来源状态 open | 2026-07-23 | 来源

V3P-000186 | aebcw-a-30-10-12

Constant weight code A(30, 10, 12)

a code of size 768 is known, and no such code can have more than 5008 words.

来源状态 open | 2026-07-23 | 来源

V3P-000187 | aebcw-a-30-10-13

Constant weight code A(30, 10, 13)

a code of size 935 is known, and no such code can have more than 6724 words.

来源状态 open | 2026-07-23 | 来源

V3P-000188 | aebcw-a-30-10-14

Constant weight code A(30, 10, 14)

a code of size 1458 is known, and no such code can have more than 8146 words.

来源状态 open | 2026-07-23 | 来源

V3P-000189 | aebcw-a-30-10-15

Constant weight code A(30, 10, 15)

a code of size 1458 is known, and no such code can have more than 8948 words.

来源状态 open | 2026-07-23 | 来源

V3P-000190 | aebcw-a-30-10-7

Constant weight code A(30, 10, 7)

a code of size 48 is known, and no such code can have more than 85 words.

来源状态 open | 2026-07-23 | 来源

V3P-000191 | aebcw-a-30-10-8

Constant weight code A(30, 10, 8)

a code of size 108 is known, and no such code can have more than 247 words.

来源状态 open | 2026-07-23 | 来源

V3P-000192 | aebcw-a-30-10-9

Constant weight code A(30, 10, 9)

a code of size 203 is known, and no such code can have more than 657 words.

来源状态 open | 2026-07-23 | 来源

V3P-000193 | aebcw-a-30-12-10

Constant weight code A(30, 12, 10)

a code of size 98 is known, and no such code can have more than 159 words.

来源状态 open | 2026-07-23 | 来源

V3P-000194 | aebcw-a-30-12-11

Constant weight code A(30, 12, 11)

a code of size 120 is known, and no such code can have more than 268 words.

来源状态 open | 2026-07-23 | 来源

V3P-000195 | aebcw-a-30-12-12

Constant weight code A(30, 12, 12)

a code of size 190 is known, and no such code can have more than 492 words.

来源状态 open | 2026-07-23 | 来源

V3P-000196 | aebcw-a-30-12-13

Constant weight code A(30, 12, 13)

a code of size 236 is known, and no such code can have more than 642 words.

来源状态 open | 2026-07-23 | 来源

V3P-000197 | aebcw-a-30-12-14

Constant weight code A(30, 12, 14)

a code of size 289 is known, and no such code can have more than 801 words.

来源状态 open | 2026-07-23 | 来源

V3P-000198 | aebcw-a-30-12-15

Constant weight code A(30, 12, 15)

a code of size 302 is known, and no such code can have more than 894 words.

来源状态 open | 2026-07-23 | 来源

V3P-000199 | aebcw-a-30-12-9

Constant weight code A(30, 12, 9)

a code of size 44 is known, and no such code can have more than 74 words.

来源状态 open | 2026-07-23 | 来源

V3P-000200 | aebcw-a-30-14-12

Constant weight code A(30, 14, 12)

a code of size 36 is known, and no such code can have more than 62 words.

来源状态 open | 2026-07-23 | 来源

V3P-000201 | aebcw-a-30-14-13

Constant weight code A(30, 14, 13)

a code of size 45 is known, and no such code can have more than 78 words.

来源状态 open | 2026-07-23 | 来源

V3P-000202 | aebcw-a-30-14-14

Constant weight code A(30, 14, 14)

a code of size 58 is known, and no such code can have more than 95 words.

来源状态 open | 2026-07-23 | 来源

V3P-000203 | aebcw-a-30-14-15

Constant weight code A(30, 14, 15)

a code of size 62 is known, and no such code can have more than 104 words.

来源状态 open | 2026-07-23 | 来源

V3P-000204 | aebcw-a-31-10-10

Constant weight code A(31, 10, 10)

a code of size 465 is known, and no such code can have more than 2036 words.

来源状态 open | 2026-07-23 | 来源

V3P-000205 | aebcw-a-31-10-11

Constant weight code A(31, 10, 11)

a code of size 651 is known, and no such code can have more than 4141 words.

来源状态 open | 2026-07-23 | 来源

V3P-000206 | aebcw-a-31-10-12

Constant weight code A(31, 10, 12)

a code of size 930 is known, and no such code can have more than 7259 words.

来源状态 open | 2026-07-23 | 来源

V3P-000207 | aebcw-a-31-10-13

Constant weight code A(31, 10, 13)

a code of size 1395 is known, and no such code can have more than 10530 words.

来源状态 open | 2026-07-23 | 来源

V3P-000208 | aebcw-a-31-10-14

Constant weight code A(31, 10, 14)

a code of size 1538 is known, and no such code can have more than 13519 words.

来源状态 open | 2026-07-23 | 来源

V3P-000209 | aebcw-a-31-10-15

Constant weight code A(31, 10, 15)

a code of size 1922 is known, and no such code can have more than 15031 words.

来源状态 open | 2026-07-23 | 来源

V3P-000210 | aebcw-a-31-10-7

Constant weight code A(31, 10, 7)

a code of size 62 is known, and no such code can have more than 109 words.

来源状态 open | 2026-07-23 | 来源

V3P-000211 | aebcw-a-31-10-8

Constant weight code A(31, 10, 8)

a code of size 124 is known, and no such code can have more than 322 words.

来源状态 open | 2026-07-23 | 来源

V3P-000212 | aebcw-a-31-10-9

Constant weight code A(31, 10, 9)

a code of size 232 is known, and no such code can have more than 822 words.

来源状态 open | 2026-07-23 | 来源

V3P-000213 | aebcw-a-31-12-10

Constant weight code A(31, 12, 10)

a code of size 103 is known, and no such code can have more than 229 words.

来源状态 open | 2026-07-23 | 来源

V3P-000214 | aebcw-a-31-12-11

Constant weight code A(31, 12, 11)

a code of size 186 is known, and no such code can have more than 415 words.

来源状态 open | 2026-07-23 | 来源

V3P-000215 | aebcw-a-31-12-12

Constant weight code A(31, 12, 12)

a code of size 310 is known, and no such code can have more than 679 words.

来源状态 open | 2026-07-23 | 来源

V3P-000216 | aebcw-a-31-12-13

Constant weight code A(31, 12, 13)

a code of size 400 is known, and no such code can have more than 958 words.

来源状态 open | 2026-07-23 | 来源

V3P-000217 | aebcw-a-31-12-14

Constant weight code A(31, 12, 14)

a code of size 510 is known, and no such code can have more than 1238 words.

来源状态 open | 2026-07-23 | 来源

V3P-000218 | aebcw-a-31-12-15

Constant weight code A(31, 12, 15)

a code of size 572 is known, and no such code can have more than 1435 words.

来源状态 open | 2026-07-23 | 来源

V3P-000219 | aebcw-a-31-12-9

Constant weight code A(31, 12, 9)

a code of size 50 is known, and no such code can have more than 94 words.

来源状态 open | 2026-07-23 | 来源

V3P-000220 | aebcw-a-31-14-11

Constant weight code A(31, 14, 11)

a code of size 35 is known, and no such code can have more than 46 words.

来源状态 open | 2026-07-23 | 来源

V3P-000221 | aebcw-a-31-14-12

Constant weight code A(31, 14, 12)

a code of size 45 is known, and no such code can have more than 77 words.

来源状态 open | 2026-07-23 | 来源

V3P-000222 | aebcw-a-31-14-13

Constant weight code A(31, 14, 13)

a code of size 60 is known, and no such code can have more than 129 words.

来源状态 open | 2026-07-23 | 来源

V3P-000223 | aebcw-a-31-14-14

Constant weight code A(31, 14, 14)

a code of size 66 is known, and no such code can have more than 167 words.

来源状态 open | 2026-07-23 | 来源

V3P-000224 | aebcw-a-31-14-15

Constant weight code A(31, 14, 15)

a code of size 72 is known, and no such code can have more than 183 words.

来源状态 open | 2026-07-23 | 来源

V3P-000225 | aebcw-a-32-10-10

Constant weight code A(32, 10, 10)

a code of size 500 is known, and no such code can have more than 2630 words.

来源状态 open | 2026-07-23 | 来源

V3P-000226 | aebcw-a-32-10-11

Constant weight code A(32, 10, 11)

a code of size 992 is known, and no such code can have more than 5696 words.

来源状态 open | 2026-07-23 | 来源

V3P-000227 | aebcw-a-32-10-12

Constant weight code A(32, 10, 12)

a code of size 1395 is known, and no such code can have more than 10446 words.

来源状态 open | 2026-07-23 | 来源

V3P-000228 | aebcw-a-32-10-13

Constant weight code A(32, 10, 13)

a code of size 1984 is known, and no such code can have more than 16755 words.

来源状态 open | 2026-07-23 | 来源

V3P-000229 | aebcw-a-32-10-14

Constant weight code A(32, 10, 14)

a code of size 2325 is known, and no such code can have more than 22213 words.

来源状态 open | 2026-07-23 | 来源

V3P-000230 | aebcw-a-32-10-15

Constant weight code A(32, 10, 15)

a code of size 2635 is known, and no such code can have more than 26361 words.

来源状态 open | 2026-07-23 | 来源

V3P-000231 | aebcw-a-32-10-16

Constant weight code A(32, 10, 16)

a code of size 3038 is known, and no such code can have more than 27429 words.

来源状态 open | 2026-07-23 | 来源

V3P-000232 | aebcw-a-32-10-7

Constant weight code A(32, 10, 7)

a code of size 64 is known, and no such code can have more than 139 words.

来源状态 open | 2026-07-23 | 来源

V3P-000233 | aebcw-a-32-10-8

Constant weight code A(32, 10, 8)

a code of size 145 is known, and no such code can have more than 402 words.

来源状态 open | 2026-07-23 | 来源

V3P-000234 | aebcw-a-32-10-9

Constant weight code A(32, 10, 9)

a code of size 304 is known, and no such code can have more than 1143 words.

来源状态 open | 2026-07-23 | 来源

V3P-000235 | aebcw-a-32-12-10

Constant weight code A(32, 12, 10)

a code of size 122 is known, and no such code can have more than 300 words.

来源状态 open | 2026-07-23 | 来源

V3P-000236 | aebcw-a-32-12-11

Constant weight code A(32, 12, 11)

a code of size 186 is known, and no such code can have more than 573 words.

来源状态 open | 2026-07-23 | 来源

V3P-000237 | aebcw-a-32-12-12

Constant weight code A(32, 12, 12)

a code of size 496 is known, and no such code can have more than 952 words.

来源状态 open | 2026-07-23 | 来源

V3P-000238 | aebcw-a-32-12-13

Constant weight code A(32, 12, 13)

a code of size 434 is known, and no such code can have more than 1497 words.

来源状态 open | 2026-07-23 | 来源

V3P-000239 | aebcw-a-32-12-14

Constant weight code A(32, 12, 14)

a code of size 900 is known, and no such code can have more than 2140 words.

来源状态 open | 2026-07-23 | 来源

V3P-000240 | aebcw-a-32-12-15

Constant weight code A(32, 12, 15)

a code of size 629 is known, and no such code can have more than 2641 words.

来源状态 open | 2026-07-23 | 来源

V3P-000241 | aebcw-a-32-12-16

Constant weight code A(32, 12, 16)

a code of size 1144 is known, and no such code can have more than 2870 words.

来源状态 open | 2026-07-23 | 来源

V3P-000242 | aebcw-a-32-12-9

Constant weight code A(32, 12, 9)

a code of size 60 is known, and no such code can have more than 110 words.

来源状态 open | 2026-07-23 | 来源

V3P-000243 | aebcw-a-32-14-11

Constant weight code A(32, 14, 11)

a code of size 40 is known, and no such code can have more than 68 words.

来源状态 open | 2026-07-23 | 来源

V3P-000244 | aebcw-a-32-14-12

Constant weight code A(32, 14, 12)

a code of size 55 is known, and no such code can have more than 118 words.

来源状态 open | 2026-07-23 | 来源

V3P-000245 | aebcw-a-32-14-13

Constant weight code A(32, 14, 13)

a code of size 70 is known, and no such code can have more than 183 words.

来源状态 open | 2026-07-23 | 来源

V3P-000246 | aebcw-a-32-14-14

Constant weight code A(32, 14, 14)

a code of size 96 is known, and no such code can have more than 233 words.

来源状态 open | 2026-07-23 | 来源

V3P-000247 | aebcw-a-32-14-15

Constant weight code A(32, 14, 15)

a code of size 96 is known, and no such code can have more than 277 words.

来源状态 open | 2026-07-23 | 来源

V3P-000248 | aebcw-a-32-14-16

Constant weight code A(32, 14, 16)

a code of size 122 is known, and no such code can have more than 295 words.

来源状态 open | 2026-07-23 | 来源

V3P-000249 | aebcw-a-32-16-13

Constant weight code A(32, 16, 13)

a code of size 25 is known, and no such code can have more than 27 words.

来源状态 open | 2026-07-23 | 来源

V3P-000250 | aebcw-a-35-18-16

Constant weight code A(35, 18, 16)

a code of size 22 is known, and no such code can have more than 28 words.

来源状态 open | 2026-07-23 | 来源

D02 二元线性码 | 965 行

给定 q=2、码长 n、维数 k，求线性 [n,k,d] 码能够达到的最大最小距离 d。

原始维护者：Markus Grassl; Andries E. Brouwer

Cite M. Grassl, "Bounds on the minimum distance of linear codes and quantum codes", online at http://www.codetables.de, and credit A. E. Brouwer's tables for the entries derived from them.

原始问题族入口

V3P-000251 | codetables-gf2-100-11

Minimum distance of a binary [100, 11] linear code codes are known with d = 44, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-000252 | codetables-gf2-100-12

Minimum distance of a binary [100, 12] linear code codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-000253 | codetables-gf2-100-13

Minimum distance of a binary [100, 13] linear code codes are known with d = 41, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-000254 | codetables-gf2-100-14

Minimum distance of a binary [100, 14] linear code codes are known with d = 40, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-000255 | codetables-gf2-100-9

Minimum distance of a binary [100, 9] linear code codes are known with d = 46, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-000256 | codetables-gf2-101-11

Minimum distance of a binary [101, 11] linear code codes are known with d = 44, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-000257 | codetables-gf2-101-12

Minimum distance of a binary [101, 12] linear code codes are known with d = 43, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-000258 | codetables-gf2-101-13

Minimum distance of a binary [101, 13] linear code codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-000259 | codetables-gf2-101-14

Minimum distance of a binary [101, 14] linear code codes are known with d = 40, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-000260 | codetables-gf2-101-9

Minimum distance of a binary [101, 9] linear code codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-000261 | codetables-gf2-102-11

Minimum distance of a binary [102, 11] linear code codes are known with d = 45, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-000262 | codetables-gf2-102-12

Minimum distance of a binary [102, 12] linear code codes are known with d = 44, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-000263 | codetables-gf2-102-13

Minimum distance of a binary [102, 13] linear code codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-000264 | codetables-gf2-102-14

Minimum distance of a binary [102, 14] linear code codes are known with d = 41, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-000265 | codetables-gf2-103-12

Minimum distance of a binary [103, 12] linear code codes are known with d = 44, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-000266 | codetables-gf2-103-13

Minimum distance of a binary [103, 13] linear code codes are known with d = 43, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-000267 | codetables-gf2-103-14

Minimum distance of a binary [103, 14] linear code codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-000268 | codetables-gf2-104-11

Minimum distance of a binary [104, 11] linear code codes are known with d = 46, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-000269 | codetables-gf2-104-12

Minimum distance of a binary [104, 12] linear code codes are known with d = 44, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-000270 | codetables-gf2-104-13

Minimum distance of a binary [104, 13] linear code

codes are known with d = 44, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-000271 | codetables-gf2-104-14

Minimum distance of a binary [104, 14] linear code

codes are known with d = 42, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-000272 | codetables-gf2-105-11

Minimum distance of a binary [105, 11] linear code

codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-000273 | codetables-gf2-105-12

Minimum distance of a binary [105, 12] linear code

codes are known with d = 45, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-000274 | codetables-gf2-105-13

Minimum distance of a binary [105, 13] linear code

codes are known with d = 44, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-000275 | codetables-gf2-105-14

Minimum distance of a binary [105, 14] linear code

codes are known with d = 43, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-000276 | codetables-gf2-106-12

Minimum distance of a binary [106, 12] linear code

codes are known with d = 46, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-000277 | codetables-gf2-106-13

Minimum distance of a binary [106, 13] linear code

codes are known with d = 44, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-000278 | codetables-gf2-106-14

Minimum distance of a binary [106, 14] linear code

codes are known with d = 44, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-000279 | codetables-gf2-107-10

Minimum distance of a binary [107, 10] linear code

codes are known with d = 48, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-000280 | codetables-gf2-107-12

Minimum distance of a binary [107, 12] linear code

codes are known with d = 46, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-000281 | codetables-gf2-107-13

Minimum distance of a binary [107, 13] linear code

codes are known with d = 45, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-000282 | codetables-gf2-107-14

Minimum distance of a binary [107, 14] linear code

codes are known with d = 44, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-000283 | codetables-gf2-108-10

Minimum distance of a binary [108, 10] linear code

codes are known with d = 49, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-000284 | codetables-gf2-108-11

Minimum distance of a binary [108, 11] linear code

codes are known with d = 48, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-000285 | codetables-gf2-108-12

Minimum distance of a binary [108, 12] linear code

codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-000286 | codetables-gf2-108-13

Minimum distance of a binary [108, 13] linear code

codes are known with d = 46, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-000287 | codetables-gf2-108-14

Minimum distance of a binary [108, 14] linear code

codes are known with d = 44, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-000288 | codetables-gf2-109-11

Minimum distance of a binary [109, 11] linear code

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-000289 | codetables-gf2-109-13

Minimum distance of a binary [109, 13] linear code

codes are known with d = 46, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-000290 | codetables-gf2-109-14

Minimum distance of a binary [109, 14] linear code

codes are known with d = 45, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-000291 | codetables-gf2-110-10

Minimum distance of a binary [110, 10] linear code

codes are known with d = 50, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-000292 | codetables-gf2-110-11

Minimum distance of a binary [110, 11] linear code

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-000293 | codetables-gf2-110-12

Minimum distance of a binary [110, 12] linear code

codes are known with d = 48, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-000294 | codetables-gf2-110-13

Minimum distance of a binary [110, 13] linear code

codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-000295 | codetables-gf2-110-14

Minimum distance of a binary [110, 14] linear code

codes are known with d = 46, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-000296 | codetables-gf2-111-10

Minimum distance of a binary [111, 10] linear code

codes are known with d = 51, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-000297 | codetables-gf2-111-11

Minimum distance of a binary [111, 11] linear code

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-000298 | codetables-gf2-111-12

Minimum distance of a binary [111, 12] linear code

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-000299 | codetables-gf2-111-13

Minimum distance of a binary [111, 13] linear code

codes are known with d = 48, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-000300 | codetables-gf2-111-14

Minimum distance of a binary [111, 14] linear code

codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-000301 | codetables-gf2-112-11

Minimum distance of a binary [112, 11] linear code

codes are known with d = 49, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-000302 | codetables-gf2-112-12

Minimum distance of a binary [112, 12] linear code

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-000303 | codetables-gf2-112-13

Minimum distance of a binary [112, 13] linear code

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-000304 | codetables-gf2-112-14

Minimum distance of a binary [112, 14] linear code

codes are known with d = 48, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-000305 | codetables-gf2-113-11

Minimum distance of a binary [113, 11] linear code

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-000306 | codetables-gf2-113-12

Minimum distance of a binary [113, 12] linear code

codes are known with d = 48, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-000307 | codetables-gf2-113-13

Minimum distance of a binary [113, 13] linear code

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-000308 | codetables-gf2-113-14

Minimum distance of a binary [113, 14] linear code

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-000309 | codetables-gf2-113-9

Minimum distance of a binary [113, 9] linear code

codes are known with d = 52, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-000310 | codetables-gf2-114-10

Minimum distance of a binary [114, 10] linear code

codes are known with d = 52, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-000311 | codetables-gf2-114-11

Minimum distance of a binary [114, 11] linear code

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-000312 | codetables-gf2-114-12

Minimum distance of a binary [114, 12] linear code

codes are known with d = 48, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-000313 | codetables-gf2-114-13

Minimum distance of a binary [114, 13] linear code

codes are known with d = 48, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-000314 | codetables-gf2-114-14

Minimum distance of a binary [114, 14] linear code

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-000315 | codetables-gf2-114-9

Minimum distance of a binary [114, 9] linear code

codes are known with d = 53, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-000316 | codetables-gf2-115-10

Minimum distance of a binary [115, 10] linear code

codes are known with d = 53, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-000317 | codetables-gf2-115-11

Minimum distance of a binary [115, 11] linear code

codes are known with d = 51, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-000318 | codetables-gf2-115-12

Minimum distance of a binary [115, 12] linear code

codes are known with d = 49, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-000319 | codetables-gf2-115-13

Minimum distance of a binary [115, 13] linear code

codes are known with d = 48, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-000320 | codetables-gf2-115-14

Minimum distance of a binary [115, 14] linear code

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-000321 | codetables-gf2-116-11

Minimum distance of a binary [116, 11] linear code

codes are known with d = 52, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-000322 | codetables-gf2-116-12

Minimum distance of a binary [116, 12] linear code

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-000323 | codetables-gf2-116-13

Minimum distance of a binary [116, 13] linear code

codes are known with d = 48, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-000324 | codetables-gf2-116-14

Minimum distance of a binary [116, 14] linear code

codes are known with d = 48, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-000325 | codetables-gf2-116-9

Minimum distance of a binary [116, 9] linear code

codes are known with d = 54, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-000326 | codetables-gf2-117-10

Minimum distance of a binary [117, 10] linear code

codes are known with d = 54, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-000327 | codetables-gf2-117-11

Minimum distance of a binary [117, 11] linear code

codes are known with d = 52, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-000328 | codetables-gf2-117-12

Minimum distance of a binary [117, 12] linear code

codes are known with d = 50, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-000329 | codetables-gf2-117-13

Minimum distance of a binary [117, 13] linear code

codes are known with d = 49, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-000330 | codetables-gf2-117-14

Minimum distance of a binary [117, 14] linear code

codes are known with d = 48, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-000331 | codetables-gf2-117-9

Minimum distance of a binary [117, 9] linear code

codes are known with d = 55, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-000332 | codetables-gf2-118-10

Minimum distance of a binary [118, 10] linear code

codes are known with d = 55, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-000333 | codetables-gf2-118-11

Minimum distance of a binary [118, 11] linear code

codes are known with d = 53, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-000334 | codetables-gf2-118-12

Minimum distance of a binary [118, 12] linear code

codes are known with d = 51, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-000335 | codetables-gf2-118-13

Minimum distance of a binary [118, 13] linear code

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-000336 | codetables-gf2-118-14

Minimum distance of a binary [118, 14] linear code

codes are known with d = 49, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-000337 | codetables-gf2-119-11

Minimum distance of a binary [119, 11] linear code

codes are known with d = 54, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-000338 | codetables-gf2-119-12

Minimum distance of a binary [119, 12] linear code

codes are known with d = 52, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-000339 | codetables-gf2-119-13

Minimum distance of a binary [119, 13] linear code

codes are known with d = 50, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-000340 | codetables-gf2-119-14

Minimum distance of a binary [119, 14] linear code

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-000341 | codetables-gf2-120-11

Minimum distance of a binary [120, 11] linear code

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-000342 | codetables-gf2-120-12

Minimum distance of a binary [120, 12] linear code

codes are known with d = 52, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-000343 | codetables-gf2-120-13

Minimum distance of a binary [120, 13] linear code

codes are known with d = 51, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-000344 | codetables-gf2-120-14

Minimum distance of a binary [120, 14] linear code

codes are known with d = 50, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-000345 | codetables-gf2-121-11

Minimum distance of a binary [121, 11] linear code

codes are known with d = 55, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-000346 | codetables-gf2-121-12

Minimum distance of a binary [121, 12] linear code

codes are known with d = 53, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-000347 | codetables-gf2-121-13

Minimum distance of a binary [121, 13] linear code

codes are known with d = 52, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-000348 | codetables-gf2-121-14

Minimum distance of a binary [121, 14] linear code

codes are known with d = 51, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-000349 | codetables-gf2-121-9

Minimum distance of a binary [121, 9] linear code

codes are known with d = 56, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-000350 | codetables-gf2-122-10

Minimum distance of a binary [122, 10] linear code

codes are known with d = 56, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-000351 | codetables-gf2-122-12

Minimum distance of a binary [122, 12] linear code

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-000352 | codetables-gf2-122-13

Minimum distance of a binary [122, 13] linear code

codes are known with d = 52, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-000353 | codetables-gf2-122-14

Minimum distance of a binary [122, 14] linear code

codes are known with d = 52, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-000354 | codetables-gf2-122-9

Minimum distance of a binary [122, 9] linear code

codes are known with d = 57, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-000355 | codetables-gf2-123-10

Minimum distance of a binary [123, 10] linear code

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-000356 | codetables-gf2-123-12

Minimum distance of a binary [123, 12] linear code

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-000357 | codetables-gf2-123-13

Minimum distance of a binary [123, 13] linear code

codes are known with d = 53, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-000358 | codetables-gf2-123-14

Minimum distance of a binary [123, 14] linear code

codes are known with d = 52, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-000359 | codetables-gf2-124-10

Minimum distance of a binary [124, 10] linear code

codes are known with d = 57, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-000360 | codetables-gf2-124-11

Minimum distance of a binary [124, 11] linear code

codes are known with d = 56, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-000361 | codetables-gf2-124-12

Minimum distance of a binary [124, 12] linear code

codes are known with d = 55, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-000362 | codetables-gf2-124-13

Minimum distance of a binary [124, 13] linear code

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-000363 | codetables-gf2-124-14

Minimum distance of a binary [124, 14] linear code

codes are known with d = 53, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-000364 | codetables-gf2-124-9

Minimum distance of a binary [124, 9] linear code

codes are known with d = 58, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-000365 | codetables-gf2-125-11

Minimum distance of a binary [125, 11] linear code

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-000366 | codetables-gf2-125-13

Minimum distance of a binary [125, 13] linear code

codes are known with d = 55, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-000367 | codetables-gf2-125-14

Minimum distance of a binary [125, 14] linear code

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-000368 | codetables-gf2-125-9

Minimum distance of a binary [125, 9] linear code

codes are known with d = 59, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000369 | codetables-gf2-126-10

Minimum distance of a binary [126, 10] linear code

codes are known with d = 58, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-000370 | codetables-gf2-126-11

Minimum distance of a binary [126, 11] linear code

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-000371 | codetables-gf2-126-12

Minimum distance of a binary [126, 12] linear code

codes are known with d = 56, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-000372 | codetables-gf2-126-14

Minimum distance of a binary [126, 14] linear code

codes are known with d = 55, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-000373 | codetables-gf2-127-10

Minimum distance of a binary [127, 10] linear code

codes are known with d = 59, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000374 | codetables-gf2-127-11

Minimum distance of a binary [127, 11] linear code

codes are known with d = 56, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-000375 | codetables-gf2-127-12

Minimum distance of a binary [127, 12] linear code

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-000376 | codetables-gf2-127-13

Minimum distance of a binary [127, 13] linear code

codes are known with d = 56, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-000377 | codetables-gf2-128-11

Minimum distance of a binary [128, 11] linear code

codes are known with d = 57, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000378 | codetables-gf2-128-12

Minimum distance of a binary [128, 12] linear code

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-000379 | codetables-gf2-128-13

Minimum distance of a binary [128, 13] linear code

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-000380 | codetables-gf2-128-14

Minimum distance of a binary [128, 14] linear code

codes are known with d = 56, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-000381 | codetables-gf2-128-9

Minimum distance of a binary [128, 9] linear code

codes are known with d = 60, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-000382 | codetables-gf2-129-11

Minimum distance of a binary [129, 11] linear code

codes are known with d = 58, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000383 | codetables-gf2-129-12

Minimum distance of a binary [129, 12] linear code

codes are known with d = 56, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-000384 | codetables-gf2-129-13

Minimum distance of a binary [129, 13] linear code

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-000385 | codetables-gf2-129-14

Minimum distance of a binary [129, 14] linear code

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-000386 | codetables-gf2-129-9

Minimum distance of a binary [129, 9] linear code

codes are known with d = 61, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-000387 | codetables-gf2-130-10

Minimum distance of a binary [130, 10] linear code

codes are known with d = 60, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-000388 | codetables-gf2-130-11

Minimum distance of a binary [130, 11] linear code

codes are known with d = 58, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000389 | codetables-gf2-130-12

Minimum distance of a binary [130, 12] linear code

codes are known with d = 56, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000390 | codetables-gf2-130-13

Minimum distance of a binary [130, 13] linear code

codes are known with d = 56, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-000391 | codetables-gf2-130-14

Minimum distance of a binary [130, 14] linear code

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-000392 | codetables-gf2-131-10

Minimum distance of a binary [131, 10] linear code

codes are known with d = 60, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-000393 | codetables-gf2-131-11

Minimum distance of a binary [131, 11] linear code

codes are known with d = 59, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000394 | codetables-gf2-131-12

Minimum distance of a binary [131, 12] linear code

codes are known with d = 57, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000395 | codetables-gf2-131-13

Minimum distance of a binary [131, 13] linear code

codes are known with d = 56, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000396 | codetables-gf2-131-14

Minimum distance of a binary [131, 14] linear code

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-000397 | codetables-gf2-132-10

Minimum distance of a binary [132, 10] linear code

codes are known with d = 61, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-000398 | codetables-gf2-132-11

Minimum distance of a binary [132, 11] linear code

codes are known with d = 60, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-000399 | codetables-gf2-132-12

Minimum distance of a binary [132, 12] linear code

codes are known with d = 58, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000400 | codetables-gf2-132-13

Minimum distance of a binary [132, 13] linear code

codes are known with d = 56, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000401 | codetables-gf2-132-14

Minimum distance of a binary [132, 14] linear code

codes are known with d = 56, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-000402 | codetables-gf2-133-10

Minimum distance of a binary [133, 10] linear code

codes are known with d = 62, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-000403 | codetables-gf2-133-11

Minimum distance of a binary [133, 11] linear code

codes are known with d = 60, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-000404 | codetables-gf2-133-12

Minimum distance of a binary [133, 12] linear code

codes are known with d = 58, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000405 | codetables-gf2-133-13

Minimum distance of a binary [133, 13] linear code

codes are known with d = 57, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000406 | codetables-gf2-133-14

Minimum distance of a binary [133, 14] linear code

codes are known with d = 56, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000407 | codetables-gf2-134-10

Minimum distance of a binary [134, 10] linear code

codes are known with d = 62, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-000408 | codetables-gf2-134-11

Minimum distance of a binary [134, 11] linear code

codes are known with d = 61, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-000409 | codetables-gf2-134-12

Minimum distance of a binary [134, 12] linear code

codes are known with d = 59, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-000410 | codetables-gf2-134-13

Minimum distance of a binary [134, 13] linear code

codes are known with d = 58, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000411 | codetables-gf2-134-14

Minimum distance of a binary [134, 14] linear code

codes are known with d = 57, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000412 | codetables-gf2-135-10

Minimum distance of a binary [135, 10] linear code

codes are known with d = 63, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-000413 | codetables-gf2-135-11

Minimum distance of a binary [135, 11] linear code

codes are known with d = 62, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-000414 | codetables-gf2-135-12

Minimum distance of a binary [135, 12] linear code

codes are known with d = 60, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-000415 | codetables-gf2-135-13

Minimum distance of a binary [135, 13] linear code

codes are known with d = 58, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-000416 | codetables-gf2-135-14

Minimum distance of a binary [135, 14] linear code

codes are known with d = 58, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-000417 | codetables-gf2-136-11

Minimum distance of a binary [136, 11] linear code

codes are known with d = 62, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-000418 | codetables-gf2-136-12

Minimum distance of a binary [136, 12] linear code

codes are known with d = 60, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-000419 | codetables-gf2-136-13

Minimum distance of a binary [136, 13] linear code

codes are known with d = 58, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-000420 | codetables-gf2-136-14

Minimum distance of a binary [136, 14] linear code

codes are known with d = 58, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-000421 | codetables-gf2-137-11

Minimum distance of a binary [137, 11] linear code

codes are known with d = 63, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-000422 | codetables-gf2-137-12

Minimum distance of a binary [137, 12] linear code

codes are known with d = 61, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-000423 | codetables-gf2-137-13

Minimum distance of a binary [137, 13] linear code

codes are known with d = 59, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-000424 | codetables-gf2-137-14

Minimum distance of a binary [137, 14] linear code

codes are known with d = 58, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-000425 | codetables-gf2-138-12

Minimum distance of a binary [138, 12] linear code

codes are known with d = 62, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-000426 | codetables-gf2-138-13

Minimum distance of a binary [138, 13] linear code

codes are known with d = 60, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-000427 | codetables-gf2-138-14

Minimum distance of a binary [138, 14] linear code

codes are known with d = 59, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-000428 | codetables-gf2-138-9

Minimum distance of a binary [138, 9] linear code

codes are known with d = 64, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-000429 | codetables-gf2-139-10

Minimum distance of a binary [139, 10] linear code

codes are known with d = 64, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-000430 | codetables-gf2-139-12

Minimum distance of a binary [139, 12] linear code

codes are known with d = 63, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-000431 | codetables-gf2-139-13

Minimum distance of a binary [139, 13] linear code

codes are known with d = 61, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-000432 | codetables-gf2-139-14

Minimum distance of a binary [139, 14] linear code

codes are known with d = 60, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-000433 | codetables-gf2-139-9

Minimum distance of a binary [139, 9] linear code

codes are known with d = 65, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-000434 | codetables-gf2-140-10

Minimum distance of a binary [140, 10] linear code

codes are known with d = 64, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-000435 | codetables-gf2-140-11

Minimum distance of a binary [140, 11] linear code

codes are known with d = 64, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-000436 | codetables-gf2-140-13

Minimum distance of a binary [140, 13] linear code

codes are known with d = 62, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-000437 | codetables-gf2-140-14

Minimum distance of a binary [140, 14] linear code

codes are known with d = 60, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-000438 | codetables-gf2-141-10

Minimum distance of a binary [141, 10] linear code

codes are known with d = 64, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-000439 | codetables-gf2-141-11

Minimum distance of a binary [141, 11] linear code

codes are known with d = 64, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-000440 | codetables-gf2-141-13

Minimum distance of a binary [141, 13] linear code

codes are known with d = 62, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-000441 | codetables-gf2-141-14

Minimum distance of a binary [141, 14] linear code

codes are known with d = 60, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-000442 | codetables-gf2-141-9

Minimum distance of a binary [141, 9] linear code

codes are known with d = 66, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-000443 | codetables-gf2-142-10

Minimum distance of a binary [142, 10] linear code

codes are known with d = 65, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-000444 | codetables-gf2-142-11

Minimum distance of a binary [142, 11] linear code

codes are known with d = 64, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-000445 | codetables-gf2-142-12

Minimum distance of a binary [142, 12] linear code

codes are known with d = 64, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-000446 | codetables-gf2-142-13

Minimum distance of a binary [142, 13] linear code

codes are known with d = 63, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-000447 | codetables-gf2-142-14

Minimum distance of a binary [142, 14] linear code

codes are known with d = 61, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-000448 | codetables-gf2-142-9

Minimum distance of a binary [142, 9] linear code

codes are known with d = 67, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000449 | codetables-gf2-143-10

Minimum distance of a binary [143, 10] linear code

codes are known with d = 66, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000450 | codetables-gf2-143-11

Minimum distance of a binary [143, 11] linear code

codes are known with d = 64, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-000451 | codetables-gf2-143-12

Minimum distance of a binary [143, 12] linear code

codes are known with d = 64, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-000452 | codetables-gf2-143-13

Minimum distance of a binary [143, 13] linear code

codes are known with d = 64, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-000453 | codetables-gf2-143-14

Minimum distance of a binary [143, 14] linear code

codes are known with d = 62, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-000454 | codetables-gf2-144-10

Minimum distance of a binary [144, 10] linear code

codes are known with d = 66, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000455 | codetables-gf2-144-11

Minimum distance of a binary [144, 11] linear code

codes are known with d = 65, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000456 | codetables-gf2-144-12

Minimum distance of a binary [144, 12] linear code

codes are known with d = 64, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-000457 | codetables-gf2-144-13

Minimum distance of a binary [144, 13] linear code

codes are known with d = 64, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-000458 | codetables-gf2-144-14

Minimum distance of a binary [144, 14] linear code

codes are known with d = 62, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-000459 | codetables-gf2-145-10

Minimum distance of a binary [145, 10] linear code

codes are known with d = 67, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000460 | codetables-gf2-145-11

Minimum distance of a binary [145, 11] linear code

codes are known with d = 66, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000461 | codetables-gf2-145-12

Minimum distance of a binary [145, 12] linear code

codes are known with d = 64, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-000462 | codetables-gf2-145-13

Minimum distance of a binary [145, 13] linear code

codes are known with d = 64, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-000463 | codetables-gf2-145-14

Minimum distance of a binary [145, 14] linear code

codes are known with d = 63, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-000464 | codetables-gf2-145-9

Minimum distance of a binary [145, 9] linear code

codes are known with d = 68, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-000465 | codetables-gf2-146-10

Minimum distance of a binary [146, 10] linear code

codes are known with d = 68, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-000466 | codetables-gf2-146-11

Minimum distance of a binary [146, 11] linear code

codes are known with d = 66, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000467 | codetables-gf2-146-12

Minimum distance of a binary [146, 12] linear code

codes are known with d = 64, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000468 | codetables-gf2-146-13

Minimum distance of a binary [146, 13] linear code

codes are known with d = 64, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-000469 | codetables-gf2-146-14

Minimum distance of a binary [146, 14] linear code

codes are known with d = 64, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-000470 | codetables-gf2-146-9

Minimum distance of a binary [146, 9] linear code

codes are known with d = 69, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-000471 | codetables-gf2-147-10

Minimum distance of a binary [147, 10] linear code

codes are known with d = 68, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-000472 | codetables-gf2-147-11

Minimum distance of a binary [147, 11] linear code

codes are known with d = 67, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000473 | codetables-gf2-147-12

Minimum distance of a binary [147, 12] linear code

codes are known with d = 65, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000474 | codetables-gf2-147-13

Minimum distance of a binary [147, 13] linear code

codes are known with d = 64, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000475 | codetables-gf2-147-14

Minimum distance of a binary [147, 14] linear code

codes are known with d = 64, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-000476 | codetables-gf2-148-10

Minimum distance of a binary [148, 10] linear code

codes are known with d = 69, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-000477 | codetables-gf2-148-11

Minimum distance of a binary [148, 11] linear code

codes are known with d = 68, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-000478 | codetables-gf2-148-12

Minimum distance of a binary [148, 12] linear code

codes are known with d = 66, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000479 | codetables-gf2-148-13

Minimum distance of a binary [148, 13] linear code

codes are known with d = 64, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000480 | codetables-gf2-148-14

Minimum distance of a binary [148, 14] linear code

codes are known with d = 64, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-000481 | codetables-gf2-148-9

Minimum distance of a binary [148, 9] linear code

codes are known with d = 70, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-000482 | codetables-gf2-149-10

Minimum distance of a binary [149, 10] linear code

codes are known with d = 70, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-000483 | codetables-gf2-149-11

Minimum distance of a binary [149, 11] linear code

codes are known with d = 68, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-000484 | codetables-gf2-149-12

Minimum distance of a binary [149, 12] linear code

codes are known with d = 66, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000485 | codetables-gf2-149-13

Minimum distance of a binary [149, 13] linear code

codes are known with d = 64, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000486 | codetables-gf2-149-14

Minimum distance of a binary [149, 14] linear code

codes are known with d = 64, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000487 | codetables-gf2-149-9

Minimum distance of a binary [149, 9] linear code

codes are known with d = 71, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-000488 | codetables-gf2-150-10

Minimum distance of a binary [150, 10] linear code

codes are known with d = 70, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-000489 | codetables-gf2-150-11

Minimum distance of a binary [150, 11] linear code

codes are known with d = 68, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-000490 | codetables-gf2-150-12

Minimum distance of a binary [150, 12] linear code

codes are known with d = 67, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-000491 | codetables-gf2-150-13

Minimum distance of a binary [150, 13] linear code

codes are known with d = 65, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000492 | codetables-gf2-150-14

Minimum distance of a binary [150, 14] linear code

codes are known with d = 64, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000493 | codetables-gf2-151-10

Minimum distance of a binary [151, 10] linear code

codes are known with d = 71, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-000494 | codetables-gf2-151-11

Minimum distance of a binary [151, 11] linear code

codes are known with d = 69, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-000495 | codetables-gf2-151-12

Minimum distance of a binary [151, 12] linear code

codes are known with d = 68, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-000496 | codetables-gf2-151-13

Minimum distance of a binary [151, 13] linear code

codes are known with d = 66, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-000497 | codetables-gf2-151-14

Minimum distance of a binary [151, 14] linear code

codes are known with d = 64, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-000498 | codetables-gf2-152-11

Minimum distance of a binary [152, 11] linear code

codes are known with d = 70, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-000499 | codetables-gf2-152-12

Minimum distance of a binary [152, 12] linear code

codes are known with d = 68, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-000500 | codetables-gf2-152-13

Minimum distance of a binary [152, 13] linear code

codes are known with d = 66, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-000501 | codetables-gf2-152-14

Minimum distance of a binary [152, 14] linear code

codes are known with d = 64, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-000502 | codetables-gf2-153-11

Minimum distance of a binary [153, 11] linear code

codes are known with d = 70, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-000503 | codetables-gf2-153-12

Minimum distance of a binary [153, 12] linear code

codes are known with d = 68, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-000504 | codetables-gf2-153-13

Minimum distance of a binary [153, 13] linear code

codes are known with d = 67, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-000505 | codetables-gf2-153-14

Minimum distance of a binary [153, 14] linear code

codes are known with d = 65, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-000506 | codetables-gf2-153-9

Minimum distance of a binary [153, 9] linear code

codes are known with d = 72, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-000507 | codetables-gf2-154-10

Minimum distance of a binary [154, 10] linear code

codes are known with d = 72, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-000508 | codetables-gf2-154-11

Minimum distance of a binary [154, 11] linear code

codes are known with d = 71, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-000509 | codetables-gf2-154-12

Minimum distance of a binary [154, 12] linear code

codes are known with d = 69, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-000510 | codetables-gf2-154-13

Minimum distance of a binary [154, 13] linear code

codes are known with d = 68, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-000511 | codetables-gf2-154-14

Minimum distance of a binary [154, 14] linear code

codes are known with d = 66, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-000512 | codetables-gf2-154-9

Minimum distance of a binary [154, 9] linear code

codes are known with d = 73, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-000513 | codetables-gf2-155-10

Minimum distance of a binary [155, 10] linear code

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-000514 | codetables-gf2-155-12

Minimum distance of a binary [155, 12] linear code

codes are known with d = 70, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-000515 | codetables-gf2-155-13

Minimum distance of a binary [155, 13] linear code

codes are known with d = 68, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-000516 | codetables-gf2-155-14

Minimum distance of a binary [155, 14] linear code

codes are known with d = 66, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-000517 | codetables-gf2-156-10

Minimum distance of a binary [156, 10] linear code

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-000518 | codetables-gf2-156-11

Minimum distance of a binary [156, 11] linear code

codes are known with d = 72, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-000519 | codetables-gf2-156-12

Minimum distance of a binary [156, 12] linear code

codes are known with d = 70, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-000520 | codetables-gf2-156-13

Minimum distance of a binary [156, 13] linear code

codes are known with d = 69, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-000521 | codetables-gf2-156-14

Minimum distance of a binary [156, 14] linear code

codes are known with d = 66, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-000522 | codetables-gf2-156-9

Minimum distance of a binary [156, 9] linear code

codes are known with d = 74, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-000523 | codetables-gf2-157-10

Minimum distance of a binary [157, 10] linear code

codes are known with d = 73, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-000524 | codetables-gf2-157-11

Minimum distance of a binary [157, 11] linear code

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-000525 | codetables-gf2-157-12

Minimum distance of a binary [157, 12] linear code

codes are known with d = 71, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-000526 | codetables-gf2-157-13

Minimum distance of a binary [157, 13] linear code

codes are known with d = 70, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-000527 | codetables-gf2-157-14

Minimum distance of a binary [157, 14] linear code

codes are known with d = 67, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-000528 | codetables-gf2-157-9

Minimum distance of a binary [157, 9] linear code

codes are known with d = 74, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000529 | codetables-gf2-158-10

Minimum distance of a binary [158, 10] linear code

codes are known with d = 74, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000530 | codetables-gf2-158-11

Minimum distance of a binary [158, 11] linear code

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-000531 | codetables-gf2-158-12

Minimum distance of a binary [158, 12] linear code

codes are known with d = 72, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-000532 | codetables-gf2-158-13

Minimum distance of a binary [158, 13] linear code

codes are known with d = 70, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-000533 | codetables-gf2-158-14

Minimum distance of a binary [158, 14] linear code

codes are known with d = 68, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-000534 | codetables-gf2-158-9

Minimum distance of a binary [158, 9] linear code

codes are known with d = 75, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000535 | codetables-gf2-159-10

Minimum distance of a binary [159, 10] linear code

codes are known with d = 74, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000536 | codetables-gf2-159-11

Minimum distance of a binary [159, 11] linear code

codes are known with d = 72, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-000537 | codetables-gf2-159-12

Minimum distance of a binary [159, 12] linear code

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-000538 | codetables-gf2-159-13

Minimum distance of a binary [159, 13] linear code

codes are known with d = 71, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-000539 | codetables-gf2-159-14

Minimum distance of a binary [159, 14] linear code

codes are known with d = 68, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-000540 | codetables-gf2-160-10

Minimum distance of a binary [160, 10] linear code

codes are known with d = 75, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000541 | codetables-gf2-160-11

Minimum distance of a binary [160, 11] linear code

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000542 | codetables-gf2-160-12

Minimum distance of a binary [160, 12] linear code

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-000543 | codetables-gf2-160-13

Minimum distance of a binary [160, 13] linear code

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-000544 | codetables-gf2-160-14

Minimum distance of a binary [160, 14] linear code

codes are known with d = 68, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-000545 | codetables-gf2-160-9

Minimum distance of a binary [160, 9] linear code

codes are known with d = 76, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-000546 | codetables-gf2-161-10

Minimum distance of a binary [161, 10] linear code

codes are known with d = 76, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-000547 | codetables-gf2-161-11

Minimum distance of a binary [161, 11] linear code

codes are known with d = 73, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000548 | codetables-gf2-161-12

Minimum distance of a binary [161, 12] linear code

codes are known with d = 72, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-000549 | codetables-gf2-161-13

Minimum distance of a binary [161, 13] linear code

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-000550 | codetables-gf2-161-14

Minimum distance of a binary [161, 14] linear code

codes are known with d = 69, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-000551 | codetables-gf2-161-9

Minimum distance of a binary [161, 9] linear code

codes are known with d = 77, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-000552 | codetables-gf2-162-10

Minimum distance of a binary [162, 10] linear code

codes are known with d = 76, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-000553 | codetables-gf2-162-11

Minimum distance of a binary [162, 11] linear code

codes are known with d = 74, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000554 | codetables-gf2-162-12

Minimum distance of a binary [162, 12] linear code

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000555 | codetables-gf2-162-13

Minimum distance of a binary [162, 13] linear code

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-000556 | codetables-gf2-162-14

Minimum distance of a binary [162, 14] linear code

codes are known with d = 70, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-000557 | codetables-gf2-163-10

Minimum distance of a binary [163, 10] linear code

codes are known with d = 76, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-000558 | codetables-gf2-163-11

Minimum distance of a binary [163, 11] linear code

codes are known with d = 74, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000559 | codetables-gf2-163-12

Minimum distance of a binary [163, 12] linear code

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000560 | codetables-gf2-163-13

Minimum distance of a binary [163, 13] linear code

codes are known with d = 72, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-000561 | codetables-gf2-163-14

Minimum distance of a binary [163, 14] linear code

codes are known with d = 70, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-000562 | codetables-gf2-163-9

Minimum distance of a binary [163, 9] linear code

codes are known with d = 78, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-000563 | codetables-gf2-164-10

Minimum distance of a binary [164, 10] linear code

codes are known with d = 77, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-000564 | codetables-gf2-164-11

Minimum distance of a binary [164, 11] linear code

codes are known with d = 75, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-000565 | codetables-gf2-164-12

Minimum distance of a binary [164, 12] linear code

codes are known with d = 73, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000566 | codetables-gf2-164-13

Minimum distance of a binary [164, 13] linear code

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000567 | codetables-gf2-164-14

Minimum distance of a binary [164, 14] linear code

codes are known with d = 71, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-000568 | codetables-gf2-164-9

Minimum distance of a binary [164, 9] linear code

codes are known with d = 79, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000569 | codetables-gf2-165-10

Minimum distance of a binary [165, 10] linear code

codes are known with d = 78, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-000570 | codetables-gf2-165-11

Minimum distance of a binary [165, 11] linear code

codes are known with d = 76, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-000571 | codetables-gf2-165-12

Minimum distance of a binary [165, 12] linear code

codes are known with d = 74, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000572 | codetables-gf2-165-13

Minimum distance of a binary [165, 13] linear code

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000573 | codetables-gf2-165-14

Minimum distance of a binary [165, 14] linear code

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000574 | codetables-gf2-166-10

Minimum distance of a binary [166, 10] linear code

codes are known with d = 78, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000575 | codetables-gf2-166-11

Minimum distance of a binary [166, 11] linear code

codes are known with d = 76, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-000576 | codetables-gf2-166-12

Minimum distance of a binary [166, 12] linear code

codes are known with d = 74, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-000577 | codetables-gf2-166-13

Minimum distance of a binary [166, 13] linear code

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000578 | codetables-gf2-166-14

Minimum distance of a binary [166, 14] linear code

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000579 | codetables-gf2-167-10

Minimum distance of a binary [167, 10] linear code

codes are known with d = 79, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000580 | codetables-gf2-167-11

Minimum distance of a binary [167, 11] linear code

codes are known with d = 77, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-000581 | codetables-gf2-167-12

Minimum distance of a binary [167, 12] linear code

codes are known with d = 75, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-000582 | codetables-gf2-167-13

Minimum distance of a binary [167, 13] linear code

codes are known with d = 73, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-000583 | codetables-gf2-167-14

Minimum distance of a binary [167, 14] linear code

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000584 | codetables-gf2-168-11

Minimum distance of a binary [168, 11] linear code

codes are known with d = 78, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000585 | codetables-gf2-168-12

Minimum distance of a binary [168, 12] linear code

codes are known with d = 76, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-000586 | codetables-gf2-168-13

Minimum distance of a binary [168, 13] linear code

codes are known with d = 74, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-000587 | codetables-gf2-168-14

Minimum distance of a binary [168, 14] linear code

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-000588 | codetables-gf2-169-11

Minimum distance of a binary [169, 11] linear code

codes are known with d = 78, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000589 | codetables-gf2-169-12

Minimum distance of a binary [169, 12] linear code

codes are known with d = 76, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-000590 | codetables-gf2-169-13

Minimum distance of a binary [169, 13] linear code

codes are known with d = 74, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-000591 | codetables-gf2-169-14

Minimum distance of a binary [169, 14] linear code

codes are known with d = 72, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-000592 | codetables-gf2-169-9

Minimum distance of a binary [169, 9] linear code

codes are known with d = 80, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-000593 | codetables-gf2-170-10

Minimum distance of a binary [170, 10] linear code

codes are known with d = 80, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-000594 | codetables-gf2-170-11

Minimum distance of a binary [170, 11] linear code

codes are known with d = 79, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000595 | codetables-gf2-170-12

Minimum distance of a binary [170, 12] linear code

codes are known with d = 77, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000596 | codetables-gf2-170-13

Minimum distance of a binary [170, 13] linear code

codes are known with d = 75, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-000597 | codetables-gf2-170-14

Minimum distance of a binary [170, 14] linear code

codes are known with d = 73, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-000598 | codetables-gf2-170-9

Minimum distance of a binary [170, 9] linear code

codes are known with d = 80, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-000599 | codetables-gf2-171-10

Minimum distance of a binary [171, 10] linear code

codes are known with d = 80, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-000600 | codetables-gf2-171-12

Minimum distance of a binary [171, 12] linear code

codes are known with d = 78, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000601 | codetables-gf2-171-13

Minimum distance of a binary [171, 13] linear code

codes are known with d = 76, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-000602 | codetables-gf2-171-14

Minimum distance of a binary [171, 14] linear code

codes are known with d = 74, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-000603 | codetables-gf2-171-9

Minimum distance of a binary [171, 9] linear code

codes are known with d = 81, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-000604 | codetables-gf2-172-10

Minimum distance of a binary [172, 10] linear code

codes are known with d = 80, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-000605 | codetables-gf2-172-11

Minimum distance of a binary [172, 11] linear code

codes are known with d = 80, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-000606 | codetables-gf2-172-12

Minimum distance of a binary [172, 12] linear code

codes are known with d = 78, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000607 | codetables-gf2-172-13

Minimum distance of a binary [172, 13] linear code

codes are known with d = 76, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000608 | codetables-gf2-172-14

Minimum distance of a binary [172, 14] linear code

codes are known with d = 74, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-000609 | codetables-gf2-172-9

Minimum distance of a binary [172, 9] linear code

codes are known with d = 82, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-000610 | codetables-gf2-173-10

Minimum distance of a binary [173, 10] linear code

codes are known with d = 81, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-000611 | codetables-gf2-173-11

Minimum distance of a binary [173, 11] linear code

codes are known with d = 80, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-000612 | codetables-gf2-173-12

Minimum distance of a binary [173, 12] linear code

codes are known with d = 79, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000613 | codetables-gf2-173-13

Minimum distance of a binary [173, 13] linear code

codes are known with d = 77, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000614 | codetables-gf2-173-14

Minimum distance of a binary [173, 14] linear code

codes are known with d = 75, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000615 | codetables-gf2-173-9

Minimum distance of a binary [173, 9] linear code

codes are known with d = 82, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000616 | codetables-gf2-174-10

Minimum distance of a binary [174, 10] linear code

codes are known with d = 82, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000617 | codetables-gf2-174-11

Minimum distance of a binary [174, 11] linear code

codes are known with d = 80, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-000618 | codetables-gf2-174-12

Minimum distance of a binary [174, 12] linear code

codes are known with d = 80, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-000619 | codetables-gf2-174-13

Minimum distance of a binary [174, 13] linear code

codes are known with d = 78, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000620 | codetables-gf2-174-14

Minimum distance of a binary [174, 14] linear code

codes are known with d = 76, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000621 | codetables-gf2-174-9

Minimum distance of a binary [174, 9] linear code

codes are known with d = 83, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000622 | codetables-gf2-175-10

Minimum distance of a binary [175, 10] linear code

codes are known with d = 82, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000623 | codetables-gf2-175-11

Minimum distance of a binary [175, 11] linear code

codes are known with d = 80, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-000624 | codetables-gf2-175-12

Minimum distance of a binary [175, 12] linear code

codes are known with d = 80, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-000625 | codetables-gf2-175-13

Minimum distance of a binary [175, 13] linear code

codes are known with d = 79, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000626 | codetables-gf2-175-14

Minimum distance of a binary [175, 14] linear code

codes are known with d = 76, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000627 | codetables-gf2-176-10

Minimum distance of a binary [176, 10] linear code

codes are known with d = 83, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000628 | codetables-gf2-176-11

Minimum distance of a binary [176, 11] linear code

codes are known with d = 81, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000629 | codetables-gf2-176-12

Minimum distance of a binary [176, 12] linear code

codes are known with d = 80, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-000630 | codetables-gf2-176-13

Minimum distance of a binary [176, 13] linear code

codes are known with d = 80, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-000631 | codetables-gf2-176-14

Minimum distance of a binary [176, 14] linear code

codes are known with d = 77, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-000632 | codetables-gf2-176-9

Minimum distance of a binary [176, 9] linear code

codes are known with d = 84, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-000633 | codetables-gf2-177-11

Minimum distance of a binary [177, 11] linear code

codes are known with d = 82, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000634 | codetables-gf2-177-12

Minimum distance of a binary [177, 12] linear code

codes are known with d = 80, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-000635 | codetables-gf2-177-13

Minimum distance of a binary [177, 13] linear code

codes are known with d = 80, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-000636 | codetables-gf2-177-14

Minimum distance of a binary [177, 14] linear code

codes are known with d = 78, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-000637 | codetables-gf2-177-9

Minimum distance of a binary [177, 9] linear code

codes are known with d = 85, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-000638 | codetables-gf2-178-10

Minimum distance of a binary [178, 10] linear code

codes are known with d = 84, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-000639 | codetables-gf2-178-11

Minimum distance of a binary [178, 11] linear code

codes are known with d = 82, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000640 | codetables-gf2-178-12

Minimum distance of a binary [178, 12] linear code

codes are known with d = 80, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000641 | codetables-gf2-178-13

Minimum distance of a binary [178, 13] linear code

codes are known with d = 80, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-000642 | codetables-gf2-178-14

Minimum distance of a binary [178, 14] linear code

codes are known with d = 78, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-000643 | codetables-gf2-179-10

Minimum distance of a binary [179, 10] linear code

codes are known with d = 84, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-000644 | codetables-gf2-179-11

Minimum distance of a binary [179, 11] linear code

codes are known with d = 83, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-000645 | codetables-gf2-179-12

Minimum distance of a binary [179, 12] linear code

codes are known with d = 81, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000646 | codetables-gf2-179-13

Minimum distance of a binary [179, 13] linear code

codes are known with d = 80, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-000647 | codetables-gf2-179-14

Minimum distance of a binary [179, 14] linear code

codes are known with d = 79, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-000648 | codetables-gf2-179-9

Minimum distance of a binary [179, 9] linear code

codes are known with d = 86, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-000649 | codetables-gf2-180-10

Minimum distance of a binary [180, 10] linear code

codes are known with d = 85, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-000650 | codetables-gf2-180-11

Minimum distance of a binary [180, 11] linear code

codes are known with d = 84, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-000651 | codetables-gf2-180-12

Minimum distance of a binary [180, 12] linear code

codes are known with d = 82, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000652 | codetables-gf2-180-13

Minimum distance of a binary [180, 13] linear code

codes are known with d = 80, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000653 | codetables-gf2-180-14

Minimum distance of a binary [180, 14] linear code

codes are known with d = 80, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-000654 | codetables-gf2-180-9

Minimum distance of a binary [180, 9] linear code

codes are known with d = 87, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-000655 | codetables-gf2-181-10

Minimum distance of a binary [181, 10] linear code

codes are known with d = 86, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-000656 | codetables-gf2-181-11

Minimum distance of a binary [181, 11] linear code

codes are known with d = 84, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-000657 | codetables-gf2-181-12

Minimum distance of a binary [181, 12] linear code

codes are known with d = 82, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000658 | codetables-gf2-181-13

Minimum distance of a binary [181, 13] linear code

codes are known with d = 81, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000659 | codetables-gf2-181-14

Minimum distance of a binary [181, 14] linear code

codes are known with d = 80, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000660 | codetables-gf2-182-10

Minimum distance of a binary [182, 10] linear code

codes are known with d = 86, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-000661 | codetables-gf2-182-11

Minimum distance of a binary [182, 11] linear code

codes are known with d = 85, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-000662 | codetables-gf2-182-12

Minimum distance of a binary [182, 12] linear code

codes are known with d = 83, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-000663 | codetables-gf2-182-13

Minimum distance of a binary [182, 13] linear code

codes are known with d = 82, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000664 | codetables-gf2-182-14

Minimum distance of a binary [182, 14] linear code

codes are known with d = 81, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000665 | codetables-gf2-183-10

Minimum distance of a binary [183, 10] linear code

codes are known with d = 87, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-000666 | codetables-gf2-183-11

Minimum distance of a binary [183, 11] linear code

codes are known with d = 86, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-000667 | codetables-gf2-183-12

Minimum distance of a binary [183, 12] linear code

codes are known with d = 84, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-000668 | codetables-gf2-183-13

Minimum distance of a binary [183, 13] linear code

codes are known with d = 82, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000669 | codetables-gf2-183-14

Minimum distance of a binary [183, 14] linear code

codes are known with d = 82, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000670 | codetables-gf2-184-11

Minimum distance of a binary [184, 11] linear code

codes are known with d = 86, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-000671 | codetables-gf2-184-12

Minimum distance of a binary [184, 12] linear code

codes are known with d = 84, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-000672 | codetables-gf2-184-13

Minimum distance of a binary [184, 13] linear code

codes are known with d = 83, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-000673 | codetables-gf2-184-14

Minimum distance of a binary [184, 14] linear code

codes are known with d = 82, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-000674 | codetables-gf2-184-9

Minimum distance of a binary [184, 9] linear code

codes are known with d = 88, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-000675 | codetables-gf2-185-11

Minimum distance of a binary [185, 11] linear code

codes are known with d = 87, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-000676 | codetables-gf2-185-12

Minimum distance of a binary [185, 12] linear code

codes are known with d = 85, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-000677 | codetables-gf2-185-13

Minimum distance of a binary [185, 13] linear code

codes are known with d = 84, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-000678 | codetables-gf2-185-14

Minimum distance of a binary [185, 14] linear code

codes are known with d = 83, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-000679 | codetables-gf2-185-9

Minimum distance of a binary [185, 9] linear code

codes are known with d = 89, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-000680 | codetables-gf2-186-10

Minimum distance of a binary [186, 10] linear code

codes are known with d = 88, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-000681 | codetables-gf2-186-12

Minimum distance of a binary [186, 12] linear code

codes are known with d = 86, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-000682 | codetables-gf2-186-13

Minimum distance of a binary [186, 13] linear code

codes are known with d = 84, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-000683 | codetables-gf2-186-14

Minimum distance of a binary [186, 14] linear code

codes are known with d = 84, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-000684 | codetables-gf2-187-10

Minimum distance of a binary [187, 10] linear code

codes are known with d = 89, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-000685 | codetables-gf2-187-12

Minimum distance of a binary [187, 12] linear code

codes are known with d = 86, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-000686 | codetables-gf2-187-13

Minimum distance of a binary [187, 13] linear code

codes are known with d = 85, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-000687 | codetables-gf2-187-14

Minimum distance of a binary [187, 14] linear code

codes are known with d = 84, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-000688 | codetables-gf2-188-12

Minimum distance of a binary [188, 12] linear code

codes are known with d = 87, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-000689 | codetables-gf2-188-13

Minimum distance of a binary [188, 13] linear code

codes are known with d = 86, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-000690 | codetables-gf2-188-14

Minimum distance of a binary [188, 14] linear code

codes are known with d = 85, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-000691 | codetables-gf2-189-10

Minimum distance of a binary [189, 10] linear code

codes are known with d = 90, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-000692 | codetables-gf2-189-13

Minimum distance of a binary [189, 13] linear code

codes are known with d = 86, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-000693 | codetables-gf2-189-14

Minimum distance of a binary [189, 14] linear code

codes are known with d = 86, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-000694 | codetables-gf2-190-10

Minimum distance of a binary [190, 10] linear code

codes are known with d = 91, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-000695 | codetables-gf2-190-12

Minimum distance of a binary [190, 12] linear code

codes are known with d = 88, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-000696 | codetables-gf2-190-13

Minimum distance of a binary [190, 13] linear code

codes are known with d = 87, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-000697 | codetables-gf2-190-14

Minimum distance of a binary [190, 14] linear code

codes are known with d = 86, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-000698 | codetables-gf2-191-12

Minimum distance of a binary [191, 12] linear code

codes are known with d = 88, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-000699 | codetables-gf2-191-14

Minimum distance of a binary [191, 14] linear code

codes are known with d = 87, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-000700 | codetables-gf2-192-12

Minimum distance of a binary [192, 12] linear code

codes are known with d = 88, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-000701 | codetables-gf2-192-13

Minimum distance of a binary [192, 13] linear code

codes are known with d = 88, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-000702 | codetables-gf2-193-12

Minimum distance of a binary [193, 12] linear code

codes are known with d = 88, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-000703 | codetables-gf2-193-13

Minimum distance of a binary [193, 13] linear code

codes are known with d = 88, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-000704 | codetables-gf2-193-14

Minimum distance of a binary [193, 14] linear code

codes are known with d = 88, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-000705 | codetables-gf2-194-12

Minimum distance of a binary [194, 12] linear code

codes are known with d = 88, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-000706 | codetables-gf2-194-13

Minimum distance of a binary [194, 13] linear code

codes are known with d = 88, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-000707 | codetables-gf2-194-14

Minimum distance of a binary [194, 14] linear code

codes are known with d = 88, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-000708 | codetables-gf2-195-12

Minimum distance of a binary [195, 12] linear code

codes are known with d = 89, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-000709 | codetables-gf2-195-13

Minimum distance of a binary [195, 13] linear code

codes are known with d = 88, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-000710 | codetables-gf2-195-14

Minimum distance of a binary [195, 14] linear code

codes are known with d = 88, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-000711 | codetables-gf2-196-12

Minimum distance of a binary [196, 12] linear code

codes are known with d = 90, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-000712 | codetables-gf2-196-13

Minimum distance of a binary [196, 13] linear code

codes are known with d = 88, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-000713 | codetables-gf2-196-14

Minimum distance of a binary [196, 14] linear code

codes are known with d = 88, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-000714 | codetables-gf2-197-12

Minimum distance of a binary [197, 12] linear code

codes are known with d = 90, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-000715 | codetables-gf2-197-13

Minimum distance of a binary [197, 13] linear code

codes are known with d = 89, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-000716 | codetables-gf2-197-14

Minimum distance of a binary [197, 14] linear code

codes are known with d = 88, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-000717 | codetables-gf2-198-12

Minimum distance of a binary [198, 12] linear code

codes are known with d = 90, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-000718 | codetables-gf2-198-13

Minimum distance of a binary [198, 13] linear code

codes are known with d = 90, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-000719 | codetables-gf2-198-14

Minimum distance of a binary [198, 14] linear code

codes are known with d = 89, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-000720 | codetables-gf2-199-12

Minimum distance of a binary [199, 12] linear code

codes are known with d = 91, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-000721 | codetables-gf2-199-13

Minimum distance of a binary [199, 13] linear code

codes are known with d = 90, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-000722 | codetables-gf2-199-14

Minimum distance of a binary [199, 14] linear code

codes are known with d = 90, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-000723 | codetables-gf2-200-12

Minimum distance of a binary [200, 12] linear code

codes are known with d = 92, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-000724 | codetables-gf2-200-13

Minimum distance of a binary [200, 13] linear code

codes are known with d = 90, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-000725 | codetables-gf2-200-14

Minimum distance of a binary [200, 14] linear code

codes are known with d = 90, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-000726 | codetables-gf2-201-12

Minimum distance of a binary [201, 12] linear code

codes are known with d = 92, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-000727 | codetables-gf2-201-13

Minimum distance of a binary [201, 13] linear code

codes are known with d = 91, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-000728 | codetables-gf2-201-14

Minimum distance of a binary [201, 14] linear code

codes are known with d = 90, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-000729 | codetables-gf2-201-9

Minimum distance of a binary [201, 9] linear code

codes are known with d = 96, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-000730 | codetables-gf2-202-12

Minimum distance of a binary [202, 12] linear code

codes are known with d = 92, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-000731 | codetables-gf2-202-13

Minimum distance of a binary [202, 13] linear code

codes are known with d = 92, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-000732 | codetables-gf2-202-14

Minimum distance of a binary [202, 14] linear code

codes are known with d = 91, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-000733 | codetables-gf2-202-9

Minimum distance of a binary [202, 9] linear code

codes are known with d = 97, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-000734 | codetables-gf2-203-10

Minimum distance of a binary [203, 10] linear code

codes are known with d = 96, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-000735 | codetables-gf2-203-12

Minimum distance of a binary [203, 12] linear code

codes are known with d = 93, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-000736 | codetables-gf2-203-13

Minimum distance of a binary [203, 13] linear code

codes are known with d = 92, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-000737 | codetables-gf2-203-14

Minimum distance of a binary [203, 14] linear code

codes are known with d = 92, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-000738 | codetables-gf2-204-10

Minimum distance of a binary [204, 10] linear code

codes are known with d = 96, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-000739 | codetables-gf2-204-11

Minimum distance of a binary [204, 11] linear code

codes are known with d = 96, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-000740 | codetables-gf2-204-12

Minimum distance of a binary [204, 12] linear code

codes are known with d = 94, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-000741 | codetables-gf2-204-13

Minimum distance of a binary [204, 13] linear code

codes are known with d = 93, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-000742 | codetables-gf2-204-14

Minimum distance of a binary [204, 14] linear code

codes are known with d = 92, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-000743 | codetables-gf2-204-9

Minimum distance of a binary [204, 9] linear code

codes are known with d = 98, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-000744 | codetables-gf2-205-10

Minimum distance of a binary [205, 10] linear code

codes are known with d = 97, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-000745 | codetables-gf2-205-11

Minimum distance of a binary [205, 11] linear code

codes are known with d = 96, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-000746 | codetables-gf2-205-12

Minimum distance of a binary [205, 12] linear code

codes are known with d = 94, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-000747 | codetables-gf2-205-13

Minimum distance of a binary [205, 13] linear code

codes are known with d = 94, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-000748 | codetables-gf2-205-14

Minimum distance of a binary [205, 14] linear code

codes are known with d = 92, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-000749 | codetables-gf2-205-9

Minimum distance of a binary [205, 9] linear code

codes are known with d = 98, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000750 | codetables-gf2-206-10

Minimum distance of a binary [206, 10] linear code

codes are known with d = 98, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-000751 | codetables-gf2-206-11

Minimum distance of a binary [206, 11] linear code

codes are known with d = 97, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-000752 | codetables-gf2-206-12

Minimum distance of a binary [206, 12] linear code

codes are known with d = 95, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-000753 | codetables-gf2-206-13

Minimum distance of a binary [206, 13] linear code

codes are known with d = 94, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-000754 | codetables-gf2-206-14

Minimum distance of a binary [206, 14] linear code

codes are known with d = 93, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-000755 | codetables-gf2-206-9

Minimum distance of a binary [206, 9] linear code

codes are known with d = 99, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000756 | codetables-gf2-207-10

Minimum distance of a binary [207, 10] linear code

codes are known with d = 98, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000757 | codetables-gf2-207-11

Minimum distance of a binary [207, 11] linear code

codes are known with d = 98, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-000758 | codetables-gf2-207-12

Minimum distance of a binary [207, 12] linear code

codes are known with d = 96, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-000759 | codetables-gf2-207-13

Minimum distance of a binary [207, 13] linear code

codes are known with d = 95, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-000760 | codetables-gf2-207-14

Minimum distance of a binary [207, 14] linear code

codes are known with d = 94, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-000761 | codetables-gf2-208-10

Minimum distance of a binary [208, 10] linear code

codes are known with d = 99, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000762 | codetables-gf2-208-11

Minimum distance of a binary [208, 11] linear code

codes are known with d = 98, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000763 | codetables-gf2-208-12

Minimum distance of a binary [208, 12] linear code

codes are known with d = 96, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-000764 | codetables-gf2-208-13

Minimum distance of a binary [208, 13] linear code

codes are known with d = 96, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-000765 | codetables-gf2-208-14

Minimum distance of a binary [208, 14] linear code

codes are known with d = 94, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-000766 | codetables-gf2-208-9

Minimum distance of a binary [208, 9] linear code

codes are known with d = 100, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-000767 | codetables-gf2-209-11

Minimum distance of a binary [209, 11] linear code

codes are known with d = 99, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000768 | codetables-gf2-209-12

Minimum distance of a binary [209, 12] linear code

codes are known with d = 96, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000769 | codetables-gf2-209-13

Minimum distance of a binary [209, 13] linear code

codes are known with d = 96, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-000770 | codetables-gf2-209-14

Minimum distance of a binary [209, 14] linear code

codes are known with d = 95, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-000771 | codetables-gf2-209-9

Minimum distance of a binary [209, 9] linear code

codes are known with d = 100, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-000772 | codetables-gf2-210-10

Minimum distance of a binary [210, 10] linear code

codes are known with d = 100, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-000773 | codetables-gf2-210-12

Minimum distance of a binary [210, 12] linear code

codes are known with d = 96, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000774 | codetables-gf2-210-13

Minimum distance of a binary [210, 13] linear code

codes are known with d = 96, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-000775 | codetables-gf2-210-14

Minimum distance of a binary [210, 14] linear code

codes are known with d = 96, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-000776 | codetables-gf2-210-9

Minimum distance of a binary [210, 9] linear code

codes are known with d = 101, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-000777 | codetables-gf2-211-10

Minimum distance of a binary [211, 10] linear code

codes are known with d = 100, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-000778 | codetables-gf2-211-11

Minimum distance of a binary [211, 11] linear code

codes are known with d = 100, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-000779 | codetables-gf2-211-12

Minimum distance of a binary [211, 12] linear code

codes are known with d = 96, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000780 | codetables-gf2-211-13

Minimum distance of a binary [211, 13] linear code

codes are known with d = 96, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-000781 | codetables-gf2-211-14

Minimum distance of a binary [211, 14] linear code

codes are known with d = 96, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-000782 | codetables-gf2-211-9

Minimum distance of a binary [211, 9] linear code

codes are known with d = 102, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-000783 | codetables-gf2-212-10

Minimum distance of a binary [212, 10] linear code

codes are known with d = 101, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-000784 | codetables-gf2-212-11

Minimum distance of a binary [212, 11] linear code

codes are known with d = 100, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-000785 | codetables-gf2-212-12

Minimum distance of a binary [212, 12] linear code

codes are known with d = 97, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000786 | codetables-gf2-212-13

Minimum distance of a binary [212, 13] linear code

codes are known with d = 96, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000787 | codetables-gf2-212-14

Minimum distance of a binary [212, 14] linear code

codes are known with d = 96, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-000788 | codetables-gf2-212-9

Minimum distance of a binary [212, 9] linear code

codes are known with d = 102, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-000789 | codetables-gf2-213-10

Minimum distance of a binary [213, 10] linear code

codes are known with d = 102, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-000790 | codetables-gf2-213-11

Minimum distance of a binary [213, 11] linear code

codes are known with d = 101, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-000791 | codetables-gf2-213-12

Minimum distance of a binary [213, 12] linear code

codes are known with d = 98, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-000792 | codetables-gf2-213-13

Minimum distance of a binary [213, 13] linear code

codes are known with d = 96, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000793 | codetables-gf2-213-14

Minimum distance of a binary [213, 14] linear code

codes are known with d = 96, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000794 | codetables-gf2-213-9

Minimum distance of a binary [213, 9] linear code

codes are known with d = 103, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-000795 | codetables-gf2-214-10

Minimum distance of a binary [214, 10] linear code

codes are known with d = 102, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-000796 | codetables-gf2-214-11

Minimum distance of a binary [214, 11] linear code

codes are known with d = 102, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-000797 | codetables-gf2-214-12

Minimum distance of a binary [214, 12] linear code

codes are known with d = 98, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-000798 | codetables-gf2-214-13

Minimum distance of a binary [214, 13] linear code

codes are known with d = 97, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000799 | codetables-gf2-214-14

Minimum distance of a binary [214, 14] linear code

codes are known with d = 96, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000800 | codetables-gf2-215-10

Minimum distance of a binary [215, 10] linear code

codes are known with d = 103, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-000801 | codetables-gf2-215-11

Minimum distance of a binary [215, 11] linear code

codes are known with d = 102, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-000802 | codetables-gf2-215-12

Minimum distance of a binary [215, 12] linear code

codes are known with d = 99, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-000803 | codetables-gf2-215-13

Minimum distance of a binary [215, 13] linear code

codes are known with d = 98, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-000804 | codetables-gf2-215-14

Minimum distance of a binary [215, 14] linear code

codes are known with d = 96, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000805 | codetables-gf2-216-11

Minimum distance of a binary [216, 11] linear code

codes are known with d = 103, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-000806 | codetables-gf2-216-12

Minimum distance of a binary [216, 12] linear code

codes are known with d = 100, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-000807 | codetables-gf2-216-13

Minimum distance of a binary [216, 13] linear code

codes are known with d = 98, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-000808 | codetables-gf2-216-14

Minimum distance of a binary [216, 14] linear code

codes are known with d = 97, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-000809 | codetables-gf2-216-9

Minimum distance of a binary [216, 9] linear code

codes are known with d = 104, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-000810 | codetables-gf2-217-12

Minimum distance of a binary [217, 12] linear code

codes are known with d = 100, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-000811 | codetables-gf2-217-13

Minimum distance of a binary [217, 13] linear code

codes are known with d = 99, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-000812 | codetables-gf2-217-14

Minimum distance of a binary [217, 14] linear code

codes are known with d = 98, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-000813 | codetables-gf2-217-9

Minimum distance of a binary [217, 9] linear code

codes are known with d = 105, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-000814 | codetables-gf2-218-10

Minimum distance of a binary [218, 10] linear code

codes are known with d = 104, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-000815 | codetables-gf2-218-12

Minimum distance of a binary [218, 12] linear code

codes are known with d = 101, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-000816 | codetables-gf2-218-13

Minimum distance of a binary [218, 13] linear code

codes are known with d = 100, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-000817 | codetables-gf2-218-14

Minimum distance of a binary [218, 14] linear code

codes are known with d = 98, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-000818 | codetables-gf2-219-10

Minimum distance of a binary [219, 10] linear code

codes are known with d = 105, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-000819 | codetables-gf2-219-11

Minimum distance of a binary [219, 11] linear code

codes are known with d = 104, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-000820 | codetables-gf2-219-12

Minimum distance of a binary [219, 12] linear code

codes are known with d = 102, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-000821 | codetables-gf2-219-13

Minimum distance of a binary [219, 13] linear code

codes are known with d = 100, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-000822 | codetables-gf2-219-14

Minimum distance of a binary [219, 14] linear code

codes are known with d = 98, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-000823 | codetables-gf2-219-9

Minimum distance of a binary [219, 9] linear code

codes are known with d = 106, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-000824 | codetables-gf2-220-11

Minimum distance of a binary [220, 11] linear code

codes are known with d = 105, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-000825 | codetables-gf2-220-12

Minimum distance of a binary [220, 12] linear code

codes are known with d = 102, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-000826 | codetables-gf2-220-13

Minimum distance of a binary [220, 13] linear code

codes are known with d = 100, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-000827 | codetables-gf2-220-14

Minimum distance of a binary [220, 14] linear code

codes are known with d = 99, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-000828 | codetables-gf2-220-9

Minimum distance of a binary [220, 9] linear code

codes are known with d = 107, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-000829 | codetables-gf2-221-10

Minimum distance of a binary [221, 10] linear code

codes are known with d = 106, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-000830 | codetables-gf2-221-12

Minimum distance of a binary [221, 12] linear code

codes are known with d = 103, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-000831 | codetables-gf2-221-13

Minimum distance of a binary [221, 13] linear code

codes are known with d = 101, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-000832 | codetables-gf2-221-14

Minimum distance of a binary [221, 14] linear code

codes are known with d = 100, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-000833 | codetables-gf2-222-10

Minimum distance of a binary [222, 10] linear code

codes are known with d = 107, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-000834 | codetables-gf2-222-11

Minimum distance of a binary [222, 11] linear code

codes are known with d = 106, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-000835 | codetables-gf2-222-12

Minimum distance of a binary [222, 12] linear code

codes are known with d = 104, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-000836 | codetables-gf2-222-13

Minimum distance of a binary [222, 13] linear code

codes are known with d = 102, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-000837 | codetables-gf2-222-14

Minimum distance of a binary [222, 14] linear code

codes are known with d = 100, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-000838 | codetables-gf2-223-11

Minimum distance of a binary [223, 11] linear code

codes are known with d = 107, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-000839 | codetables-gf2-223-12

Minimum distance of a binary [223, 12] linear code

codes are known with d = 104, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-000840 | codetables-gf2-223-13

Minimum distance of a binary [223, 13] linear code

codes are known with d = 102, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-000841 | codetables-gf2-223-14

Minimum distance of a binary [223, 14] linear code

codes are known with d = 100, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-000842 | codetables-gf2-223-9

Minimum distance of a binary [223, 9] linear code

codes are known with d = 108, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-000843 | codetables-gf2-224-12

Minimum distance of a binary [224, 12] linear code

codes are known with d = 105, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-000844 | codetables-gf2-224-13

Minimum distance of a binary [224, 13] linear code

codes are known with d = 103, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-000845 | codetables-gf2-224-14

Minimum distance of a binary [224, 14] linear code

codes are known with d = 101, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-000846 | codetables-gf2-224-9

Minimum distance of a binary [224, 9] linear code

codes are known with d = 109, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-000847 | codetables-gf2-225-10

Minimum distance of a binary [225, 10] linear code

codes are known with d = 108, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-000848 | codetables-gf2-225-12

Minimum distance of a binary [225, 12] linear code

codes are known with d = 106, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-000849 | codetables-gf2-225-13

Minimum distance of a binary [225, 13] linear code

codes are known with d = 104, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-000850 | codetables-gf2-225-14

Minimum distance of a binary [225, 14] linear code

codes are known with d = 102, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-000851 | codetables-gf2-226-10

Minimum distance of a binary [226, 10] linear code

codes are known with d = 109, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-000852 | codetables-gf2-226-11

Minimum distance of a binary [226, 11] linear code

codes are known with d = 108, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-000853 | codetables-gf2-226-12

Minimum distance of a binary [226, 12] linear code

codes are known with d = 106, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-000854 | codetables-gf2-226-13

Minimum distance of a binary [226, 13] linear code

codes are known with d = 104, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-000855 | codetables-gf2-226-14

Minimum distance of a binary [226, 14] linear code

codes are known with d = 102, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-000856 | codetables-gf2-227-11

Minimum distance of a binary [227, 11] linear code

codes are known with d = 109, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-000857 | codetables-gf2-227-12

Minimum distance of a binary [227, 12] linear code

codes are known with d = 107, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-000858 | codetables-gf2-227-13

Minimum distance of a binary [227, 13] linear code

codes are known with d = 104, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-000859 | codetables-gf2-227-14

Minimum distance of a binary [227, 14] linear code

codes are known with d = 103, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-000860 | codetables-gf2-228-10

Minimum distance of a binary [228, 10] linear code

codes are known with d = 110, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-000861 | codetables-gf2-228-13

Minimum distance of a binary [228, 13] linear code

codes are known with d = 104, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-000862 | codetables-gf2-228-14

Minimum distance of a binary [228, 14] linear code

codes are known with d = 104, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-000863 | codetables-gf2-229-10

Minimum distance of a binary [229, 10] linear code

codes are known with d = 111, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-000864 | codetables-gf2-229-12

Minimum distance of a binary [229, 12] linear code

codes are known with d = 108, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-000865 | codetables-gf2-229-13

Minimum distance of a binary [229, 13] linear code

codes are known with d = 105, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-000866 | codetables-gf2-229-14

Minimum distance of a binary [229, 14] linear code

codes are known with d = 104, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-000867 | codetables-gf2-230-12

Minimum distance of a binary [230, 12] linear code

codes are known with d = 109, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-000868 | codetables-gf2-230-13

Minimum distance of a binary [230, 13] linear code

codes are known with d = 106, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-000869 | codetables-gf2-230-14

Minimum distance of a binary [230, 14] linear code

codes are known with d = 104, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-000870 | codetables-gf2-231-13

Minimum distance of a binary [231, 13] linear code

codes are known with d = 106, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-000871 | codetables-gf2-231-14

Minimum distance of a binary [231, 14] linear code

codes are known with d = 104, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-000872 | codetables-gf2-232-13

Minimum distance of a binary [232, 13] linear code

codes are known with d = 106, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-000873 | codetables-gf2-232-14

Minimum distance of a binary [232, 14] linear code

codes are known with d = 104, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-000874 | codetables-gf2-232-9

Minimum distance of a binary [232, 9] linear code

codes are known with d = 112, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-000875 | codetables-gf2-233-13

Minimum distance of a binary [233, 13] linear code

codes are known with d = 107, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-000876 | codetables-gf2-233-14

Minimum distance of a binary [233, 14] linear code

codes are known with d = 104, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-000877 | codetables-gf2-233-9

Minimum distance of a binary [233, 9] linear code

codes are known with d = 112, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-000878 | codetables-gf2-234-10

Minimum distance of a binary [234, 10] linear code

codes are known with d = 112, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-000879 | codetables-gf2-234-13

Minimum distance of a binary [234, 13] linear code

codes are known with d = 108, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-000880 | codetables-gf2-234-14

Minimum distance of a binary [234, 14] linear code

codes are known with d = 104, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-000881 | codetables-gf2-234-9

Minimum distance of a binary [234, 9] linear code

codes are known with d = 113, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-000882 | codetables-gf2-235-10

Minimum distance of a binary [235, 10] linear code

codes are known with d = 112, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-000883 | codetables-gf2-235-11

Minimum distance of a binary [235, 11] linear code

codes are known with d = 112, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-000884 | codetables-gf2-235-13

Minimum distance of a binary [235, 13] linear code

codes are known with d = 108, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-000885 | codetables-gf2-235-14

Minimum distance of a binary [235, 14] linear code

codes are known with d = 105, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-000886 | codetables-gf2-235-9

Minimum distance of a binary [235, 9] linear code

codes are known with d = 114, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-000887 | codetables-gf2-236-10

Minimum distance of a binary [236, 10] linear code

codes are known with d = 112, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-000888 | codetables-gf2-236-11

Minimum distance of a binary [236, 11] linear code

codes are known with d = 112, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-000889 | codetables-gf2-236-13

Minimum distance of a binary [236, 13] linear code

codes are known with d = 108, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-000890 | codetables-gf2-236-14

Minimum distance of a binary [236, 14] linear code

codes are known with d = 106, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-000891 | codetables-gf2-236-9

Minimum distance of a binary [236, 9] linear code

codes are known with d = 114, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000892 | codetables-gf2-237-10

Minimum distance of a binary [237, 10] linear code

codes are known with d = 113, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-000893 | codetables-gf2-237-11

Minimum distance of a binary [237, 11] linear code

codes are known with d = 112, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-000894 | codetables-gf2-237-12

Minimum distance of a binary [237, 12] linear code

codes are known with d = 112, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-000895 | codetables-gf2-237-13

Minimum distance of a binary [237, 13] linear code

codes are known with d = 109, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-000896 | codetables-gf2-237-14

Minimum distance of a binary [237, 14] linear code

codes are known with d = 106, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-000897 | codetables-gf2-237-9

Minimum distance of a binary [237, 9] linear code

codes are known with d = 115, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000898 | codetables-gf2-238-10

Minimum distance of a binary [238, 10] linear code

codes are known with d = 114, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000899 | codetables-gf2-238-11

Minimum distance of a binary [238, 11] linear code

codes are known with d = 112, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-000900 | codetables-gf2-238-12

Minimum distance of a binary [238, 12] linear code

codes are known with d = 112, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-000901 | codetables-gf2-238-13

Minimum distance of a binary [238, 13] linear code

codes are known with d = 110, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-000902 | codetables-gf2-238-14

Minimum distance of a binary [238, 14] linear code

codes are known with d = 106, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-000903 | codetables-gf2-239-10

Minimum distance of a binary [239, 10] linear code

codes are known with d = 114, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000904 | codetables-gf2-239-11

Minimum distance of a binary [239, 11] linear code

codes are known with d = 113, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000905 | codetables-gf2-239-12

Minimum distance of a binary [239, 12] linear code

codes are known with d = 112, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-000906 | codetables-gf2-239-13

Minimum distance of a binary [239, 13] linear code

codes are known with d = 111, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-000907 | codetables-gf2-239-14

Minimum distance of a binary [239, 14] linear code

codes are known with d = 107, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-000908 | codetables-gf2-240-10

Minimum distance of a binary [240, 10] linear code

codes are known with d = 114, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000909 | codetables-gf2-240-11

Minimum distance of a binary [240, 11] linear code

codes are known with d = 114, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000910 | codetables-gf2-240-12

Minimum distance of a binary [240, 12] linear code

codes are known with d = 112, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-000911 | codetables-gf2-240-13

Minimum distance of a binary [240, 13] linear code

codes are known with d = 112, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-000912 | codetables-gf2-240-14

Minimum distance of a binary [240, 14] linear code

codes are known with d = 108, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-000913 | codetables-gf2-240-9

Minimum distance of a binary [240, 9] linear code

codes are known with d = 116, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-000914 | codetables-gf2-241-10

Minimum distance of a binary [241, 10] linear code

codes are known with d = 115, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-000915 | codetables-gf2-241-11

Minimum distance of a binary [241, 11] linear code

codes are known with d = 114, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000916 | codetables-gf2-241-12

Minimum distance of a binary [241, 12] linear code

codes are known with d = 112, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000917 | codetables-gf2-241-13

Minimum distance of a binary [241, 13] linear code

codes are known with d = 112, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-000918 | codetables-gf2-241-14

Minimum distance of a binary [241, 14] linear code

codes are known with d = 108, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-000919 | codetables-gf2-241-9

Minimum distance of a binary [241, 9] linear code

codes are known with d = 117, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-000920 | codetables-gf2-242-10

Minimum distance of a binary [242, 10] linear code

codes are known with d = 116, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-000921 | codetables-gf2-242-11

Minimum distance of a binary [242, 11] linear code

codes are known with d = 114, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000922 | codetables-gf2-242-12

Minimum distance of a binary [242, 12] linear code

codes are known with d = 113, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000923 | codetables-gf2-242-13

Minimum distance of a binary [242, 13] linear code

codes are known with d = 112, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-000924 | codetables-gf2-242-14

Minimum distance of a binary [242, 14] linear code

codes are known with d = 109, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-000925 | codetables-gf2-243-10

Minimum distance of a binary [243, 10] linear code

codes are known with d = 116, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-000926 | codetables-gf2-243-11

Minimum distance of a binary [243, 11] linear code

codes are known with d = 115, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-000927 | codetables-gf2-243-12

Minimum distance of a binary [243, 12] linear code

codes are known with d = 114, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000928 | codetables-gf2-243-13

Minimum distance of a binary [243, 13] linear code

codes are known with d = 112, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000929 | codetables-gf2-243-14

Minimum distance of a binary [243, 14] linear code

codes are known with d = 110, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-000930 | codetables-gf2-243-9

Minimum distance of a binary [243, 9] linear code

codes are known with d = 118, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-000931 | codetables-gf2-244-10

Minimum distance of a binary [244, 10] linear code

codes are known with d = 116, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-000932 | codetables-gf2-244-11

Minimum distance of a binary [244, 11] linear code

codes are known with d = 116, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-000933 | codetables-gf2-244-12

Minimum distance of a binary [244, 12] linear code

codes are known with d = 114, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000934 | codetables-gf2-244-13

Minimum distance of a binary [244, 13] linear code

codes are known with d = 112, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000935 | codetables-gf2-244-14

Minimum distance of a binary [244, 14] linear code

codes are known with d = 110, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-000936 | codetables-gf2-244-9

Minimum distance of a binary [244, 9] linear code

codes are known with d = 119, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000937 | codetables-gf2-245-10

Minimum distance of a binary [245, 10] linear code

codes are known with d = 117, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000938 | codetables-gf2-245-11

Minimum distance of a binary [245, 11] linear code

codes are known with d = 116, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-000939 | codetables-gf2-245-12

Minimum distance of a binary [245, 12] linear code

codes are known with d = 115, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-000940 | codetables-gf2-245-13

Minimum distance of a binary [245, 13] linear code

codes are known with d = 113, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000941 | codetables-gf2-245-14

Minimum distance of a binary [245, 14] linear code

codes are known with d = 110, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-000942 | codetables-gf2-246-10

Minimum distance of a binary [246, 10] linear code

codes are known with d = 118, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000943 | codetables-gf2-246-11

Minimum distance of a binary [246, 11] linear code

codes are known with d = 116, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-000944 | codetables-gf2-246-12

Minimum distance of a binary [246, 12] linear code

codes are known with d = 116, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-000945 | codetables-gf2-246-13

Minimum distance of a binary [246, 13] linear code

codes are known with d = 114, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-000946 | codetables-gf2-246-14

Minimum distance of a binary [246, 14] linear code

codes are known with d = 111, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-000947 | codetables-gf2-247-10

Minimum distance of a binary [247, 10] linear code

codes are known with d = 118, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000948 | codetables-gf2-247-11

Minimum distance of a binary [247, 11] linear code

codes are known with d = 117, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000949 | codetables-gf2-247-12

Minimum distance of a binary [247, 12] linear code

codes are known with d = 116, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-000950 | codetables-gf2-247-13

Minimum distance of a binary [247, 13] linear code

codes are known with d = 114, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-000951 | codetables-gf2-247-14

Minimum distance of a binary [247, 14] linear code

codes are known with d = 112, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-000952 | codetables-gf2-247-9

Minimum distance of a binary [247, 9] linear code

codes are known with d = 120, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-000953 | codetables-gf2-248-10

Minimum distance of a binary [248, 10] linear code

codes are known with d = 119, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000954 | codetables-gf2-248-11

Minimum distance of a binary [248, 11] linear code

codes are known with d = 118, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000955 | codetables-gf2-248-12

Minimum distance of a binary [248, 12] linear code

codes are known with d = 117, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000956 | codetables-gf2-248-13

Minimum distance of a binary [248, 13] linear code

codes are known with d = 114, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-000957 | codetables-gf2-248-14

Minimum distance of a binary [248, 14] linear code

codes are known with d = 112, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-000958 | codetables-gf2-248-9

Minimum distance of a binary [248, 9] linear code

codes are known with d = 121, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-000959 | codetables-gf2-249-10

Minimum distance of a binary [249, 10] linear code

codes are known with d = 120, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-000960 | codetables-gf2-249-11

Minimum distance of a binary [249, 11] linear code

codes are known with d = 118, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000961 | codetables-gf2-249-12

Minimum distance of a binary [249, 12] linear code

codes are known with d = 118, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000962 | codetables-gf2-249-13

Minimum distance of a binary [249, 13] linear code

codes are known with d = 115, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-000963 | codetables-gf2-249-14

Minimum distance of a binary [249, 14] linear code

codes are known with d = 112, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-000964 | codetables-gf2-250-10

Minimum distance of a binary [250, 10] linear code

codes are known with d = 120, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-000965 | codetables-gf2-250-11

Minimum distance of a binary [250, 11] linear code

codes are known with d = 119, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000966 | codetables-gf2-250-12

Minimum distance of a binary [250, 12] linear code

codes are known with d = 118, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000967 | codetables-gf2-250-13

Minimum distance of a binary [250, 13] linear code

codes are known with d = 116, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000968 | codetables-gf2-250-14

Minimum distance of a binary [250, 14] linear code

codes are known with d = 112, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-000969 | codetables-gf2-251-10

Minimum distance of a binary [251, 10] linear code

codes are known with d = 120, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-000970 | codetables-gf2-251-11

Minimum distance of a binary [251, 11] linear code

codes are known with d = 120, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-000971 | codetables-gf2-251-12

Minimum distance of a binary [251, 12] linear code

codes are known with d = 119, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000972 | codetables-gf2-251-13

Minimum distance of a binary [251, 13] linear code

codes are known with d = 116, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000973 | codetables-gf2-251-14

Minimum distance of a binary [251, 14] linear code

codes are known with d = 112, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000974 | codetables-gf2-252-10

Minimum distance of a binary [252, 10] linear code

codes are known with d = 121, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-000975 | codetables-gf2-252-11

Minimum distance of a binary [252, 11] linear code

codes are known with d = 120, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-000976 | codetables-gf2-252-13

Minimum distance of a binary [252, 13] linear code

codes are known with d = 116, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000977 | codetables-gf2-252-14

Minimum distance of a binary [252, 14] linear code

codes are known with d = 113, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000978 | codetables-gf2-253-10

Minimum distance of a binary [253, 10] linear code

codes are known with d = 122, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-000979 | codetables-gf2-253-11

Minimum distance of a binary [253, 11] linear code

codes are known with d = 120, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-000980 | codetables-gf2-253-12

Minimum distance of a binary [253, 12] linear code

codes are known with d = 120, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-000981 | codetables-gf2-253-13

Minimum distance of a binary [253, 13] linear code

codes are known with d = 117, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000982 | codetables-gf2-253-14

Minimum distance of a binary [253, 14] linear code

codes are known with d = 114, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000983 | codetables-gf2-254-10

Minimum distance of a binary [254, 10] linear code

codes are known with d = 122, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-000984 | codetables-gf2-254-11

Minimum distance of a binary [254, 11] linear code

codes are known with d = 120, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-000985 | codetables-gf2-254-12

Minimum distance of a binary [254, 12] linear code

codes are known with d = 120, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-000986 | codetables-gf2-254-13

Minimum distance of a binary [254, 13] linear code

codes are known with d = 118, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000987 | codetables-gf2-254-14

Minimum distance of a binary [254, 14] linear code

codes are known with d = 114, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000988 | codetables-gf2-255-10

Minimum distance of a binary [255, 10] linear code

codes are known with d = 123, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-000989 | codetables-gf2-255-11

Minimum distance of a binary [255, 11] linear code

codes are known with d = 120, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-000990 | codetables-gf2-255-12

Minimum distance of a binary [255, 12] linear code

codes are known with d = 120, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-000991 | codetables-gf2-255-13

Minimum distance of a binary [255, 13] linear code

codes are known with d = 119, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-000992 | codetables-gf2-255-14

Minimum distance of a binary [255, 14] linear code

codes are known with d = 115, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-000993 | codetables-gf2-256-11

Minimum distance of a binary [256, 11] linear code

codes are known with d = 121, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-000994 | codetables-gf2-256-12

Minimum distance of a binary [256, 12] linear code

codes are known with d = 120, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-000995 | codetables-gf2-256-13

Minimum distance of a binary [256, 13] linear code

codes are known with d = 120, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-000996 | codetables-gf2-256-14

Minimum distance of a binary [256, 14] linear code

codes are known with d = 116, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-000997 | codetables-gf2-32-14

Minimum distance of a binary [32, 14] linear code

codes are known with d = 8, and no code can have d greater than 9.

来源状态 open | 2026-07-23 | 来源

V3P-000998 | codetables-gf2-33-14

Minimum distance of a binary [33, 14] linear code

codes are known with d = 9, and no code can have d greater than 10.

来源状态 open | 2026-07-23 | 来源

V3P-000999 | codetables-gf2-35-10

Minimum distance of a binary [35, 10] linear code

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-001000 | codetables-gf2-36-10

Minimum distance of a binary [36, 10] linear code

codes are known with d = 13, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-001001 | codetables-gf2-36-11

Minimum distance of a binary [36, 11] linear code

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-001002 | codetables-gf2-37-11

Minimum distance of a binary [37, 11] linear code

codes are known with d = 13, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-001003 | codetables-gf2-37-12

Minimum distance of a binary [37, 12] linear code

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-001004 | codetables-gf2-38-12

Minimum distance of a binary [38, 12] linear code

codes are known with d = 13, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-001005 | codetables-gf2-39-13

Minimum distance of a binary [39, 13] linear code

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-001006 | codetables-gf2-40-13

Minimum distance of a binary [40, 13] linear code

codes are known with d = 13, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-001007 | codetables-gf2-40-14

Minimum distance of a binary [40, 14] linear code

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-001008 | codetables-gf2-41-12

Minimum distance of a binary [41, 12] linear code

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-001009 | codetables-gf2-41-14

Minimum distance of a binary [41, 14] linear code

codes are known with d = 12, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-001010 | codetables-gf2-42-12

Minimum distance of a binary [42, 12] linear code

codes are known with d = 15, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-001011 | codetables-gf2-42-14

Minimum distance of a binary [42, 14] linear code

codes are known with d = 13, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-001012 | codetables-gf2-43-10

Minimum distance of a binary [43, 10] linear code

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-001013 | codetables-gf2-44-10

Minimum distance of a binary [44, 10] linear code

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-001014 | codetables-gf2-44-11

Minimum distance of a binary [44, 11] linear code

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-001015 | codetables-gf2-45-11

Minimum distance of a binary [45, 11] linear code

codes are known with d = 16, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-001016 | codetables-gf2-46-10

Minimum distance of a binary [46, 10] linear code

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-001017 | codetables-gf2-46-11

Minimum distance of a binary [46, 11] linear code

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-001018 | codetables-gf2-46-12

Minimum distance of a binary [46, 12] linear code

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-001019 | codetables-gf2-47-10

Minimum distance of a binary [47, 10] linear code

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-001020 | codetables-gf2-47-12

Minimum distance of a binary [47, 12] linear code

codes are known with d = 16, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-001021 | codetables-gf2-47-13

Minimum distance of a binary [47, 13] linear code

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-001022 | codetables-gf2-48-11

Minimum distance of a binary [48, 11] linear code

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-001023 | codetables-gf2-48-12

Minimum distance of a binary [48, 12] linear code

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-001024 | codetables-gf2-48-13

Minimum distance of a binary [48, 13] linear code

codes are known with d = 16, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-001025 | codetables-gf2-48-14

Minimum distance of a binary [48, 14] linear code

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-001026 | codetables-gf2-49-11

Minimum distance of a binary [49, 11] linear code

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-001027 | codetables-gf2-49-12

Minimum distance of a binary [49, 12] linear code

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-001028 | codetables-gf2-49-13

Minimum distance of a binary [49, 13] linear code

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-001029 | codetables-gf2-49-14

Minimum distance of a binary [49, 14] linear code

codes are known with d = 16, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-001030 | codetables-gf2-50-12

Minimum distance of a binary [50, 12] linear code

codes are known with d = 18, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-001031 | codetables-gf2-50-13

Minimum distance of a binary [50, 13] linear code

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-001032 | codetables-gf2-50-14

Minimum distance of a binary [50, 14] linear code

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-001033 | codetables-gf2-51-10

Minimum distance of a binary [51, 10] linear code

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-001034 | codetables-gf2-51-12

Minimum distance of a binary [51, 12] linear code

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-001035 | codetables-gf2-51-13

Minimum distance of a binary [51, 13] linear code

codes are known with d = 18, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-001036 | codetables-gf2-52-10

Minimum distance of a binary [52, 10] linear code

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-001037 | codetables-gf2-52-11

Minimum distance of a binary [52, 11] linear code

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-001038 | codetables-gf2-52-13

Minimum distance of a binary [52, 13] linear code

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-001039 | codetables-gf2-52-14

Minimum distance of a binary [52, 14] linear code

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-001040 | codetables-gf2-53-11

Minimum distance of a binary [53, 11] linear code

codes are known with d = 20, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-001041 | codetables-gf2-53-12

Minimum distance of a binary [53, 12] linear code

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-001042 | codetables-gf2-53-14

Minimum distance of a binary [53, 14] linear code

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-001043 | codetables-gf2-54-10

Minimum distance of a binary [54, 10] linear code

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-001044 | codetables-gf2-54-11

Minimum distance of a binary [54, 11] linear code

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-001045 | codetables-gf2-54-12

Minimum distance of a binary [54, 12] linear code

codes are known with d = 20, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-001046 | codetables-gf2-55-10

Minimum distance of a binary [55, 10] linear code

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-001047 | codetables-gf2-55-11

Minimum distance of a binary [55, 11] linear code

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-001048 | codetables-gf2-55-12

Minimum distance of a binary [55, 12] linear code

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-001049 | codetables-gf2-55-13

Minimum distance of a binary [55, 13] linear code

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-001050 | codetables-gf2-56-11

Minimum distance of a binary [56, 11] linear code

codes are known with d = 22, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-001051 | codetables-gf2-56-13

Minimum distance of a binary [56, 13] linear code

codes are known with d = 20, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-001052 | codetables-gf2-56-14

Minimum distance of a binary [56, 14] linear code

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-001053 | codetables-gf2-57-11

Minimum distance of a binary [57, 11] linear code

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-001054 | codetables-gf2-57-12

Minimum distance of a binary [57, 12] linear code

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-001055 | codetables-gf2-57-13

Minimum distance of a binary [57, 13] linear code

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-001056 | codetables-gf2-57-14

Minimum distance of a binary [57, 14] linear code

codes are known with d = 20, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-001057 | codetables-gf2-58-12

Minimum distance of a binary [58, 12] linear code

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-001058 | codetables-gf2-58-13

Minimum distance of a binary [58, 13] linear code

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-001059 | codetables-gf2-58-14

Minimum distance of a binary [58, 14] linear code

codes are known with d = 20, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-001060 | codetables-gf2-59-10

Minimum distance of a binary [59, 10] linear code

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-001061 | codetables-gf2-59-13

Minimum distance of a binary [59, 13] linear code

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-001062 | codetables-gf2-59-14

Minimum distance of a binary [59, 14] linear code

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-001063 | codetables-gf2-60-10

Minimum distance of a binary [60, 10] linear code

codes are known with d = 25, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-001064 | codetables-gf2-60-14

Minimum distance of a binary [60, 14] linear code

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-001065 | codetables-gf2-61-11

Minimum distance of a binary [61, 11] linear code

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-001066 | codetables-gf2-61-14

Minimum distance of a binary [61, 14] linear code

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-001067 | codetables-gf2-62-10

Minimum distance of a binary [62, 10] linear code

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-001068 | codetables-gf2-62-11

Minimum distance of a binary [62, 11] linear code

codes are known with d = 25, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-001069 | codetables-gf2-62-12

Minimum distance of a binary [62, 12] linear code

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-001070 | codetables-gf2-63-10

Minimum distance of a binary [63, 10] linear code

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-001071 | codetables-gf2-63-12

Minimum distance of a binary [63, 12] linear code

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-001072 | codetables-gf2-63-13

Minimum distance of a binary [63, 13] linear code

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-001073 | codetables-gf2-64-11

Minimum distance of a binary [64, 11] linear code

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-001074 | codetables-gf2-64-12

Minimum distance of a binary [64, 12] linear code

codes are known with d = 25, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-001075 | codetables-gf2-64-13

Minimum distance of a binary [64, 13] linear code

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-001076 | codetables-gf2-64-14

Minimum distance of a binary [64, 14] linear code

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-001077 | codetables-gf2-65-11

Minimum distance of a binary [65, 11] linear code

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-001078 | codetables-gf2-65-12

Minimum distance of a binary [65, 12] linear code

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-001079 | codetables-gf2-65-13

Minimum distance of a binary [65, 13] linear code

codes are known with d = 25, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-001080 | codetables-gf2-65-14

Minimum distance of a binary [65, 14] linear code

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-001081 | codetables-gf2-65-9

Minimum distance of a binary [65, 9] linear code

codes are known with d = 28, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-001082 | codetables-gf2-66-12

Minimum distance of a binary [66, 12] linear code

codes are known with d = 26, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-001083 | codetables-gf2-66-13

Minimum distance of a binary [66, 13] linear code

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-001084 | codetables-gf2-66-14

Minimum distance of a binary [66, 14] linear code

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-001085 | codetables-gf2-66-9

Minimum distance of a binary [66, 9] linear code

codes are known with d = 29, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-001086 | codetables-gf2-67-12

Minimum distance of a binary [67, 12] linear code

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-001087 | codetables-gf2-67-13

Minimum distance of a binary [67, 13] linear code

codes are known with d = 26, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-001088 | codetables-gf2-67-14

Minimum distance of a binary [67, 14] linear code

codes are known with d = 24, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-001089 | codetables-gf2-68-11

Minimum distance of a binary [68, 11] linear code

codes are known with d = 28, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-001090 | codetables-gf2-68-13

Minimum distance of a binary [68, 13] linear code

codes are known with d = 26, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-001091 | codetables-gf2-68-14

Minimum distance of a binary [68, 14] linear code

codes are known with d = 25, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-001092 | codetables-gf2-69-11

Minimum distance of a binary [69, 11] linear code

codes are known with d = 29, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-001093 | codetables-gf2-69-12

Minimum distance of a binary [69, 12] linear code

codes are known with d = 28, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-001094 | codetables-gf2-69-13

Minimum distance of a binary [69, 13] linear code

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-001095 | codetables-gf2-69-14

Minimum distance of a binary [69, 14] linear code

codes are known with d = 26, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-001096 | codetables-gf2-70-12

Minimum distance of a binary [70, 12] linear code

codes are known with d = 28, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-001097 | codetables-gf2-70-14

Minimum distance of a binary [70, 14] linear code

codes are known with d = 26, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-001098 | codetables-gf2-71-12

Minimum distance of a binary [71, 12] linear code

codes are known with d = 29, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-001099 | codetables-gf2-71-13

Minimum distance of a binary [71, 13] linear code

codes are known with d = 28, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-001100 | codetables-gf2-71-14

Minimum distance of a binary [71, 14] linear code

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-001101 | codetables-gf2-72-13

Minimum distance of a binary [72, 13] linear code

codes are known with d = 28, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-001102 | codetables-gf2-72-14

Minimum distance of a binary [72, 14] linear code

codes are known with d = 28, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-001103 | codetables-gf2-73-13

Minimum distance of a binary [73, 13] linear code

codes are known with d = 29, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-001104 | codetables-gf2-73-14

Minimum distance of a binary [73, 14] linear code

codes are known with d = 28, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-001105 | codetables-gf2-74-13

Minimum distance of a binary [74, 13] linear code

codes are known with d = 30, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-001106 | codetables-gf2-74-14

Minimum distance of a binary [74, 14] linear code

codes are known with d = 28, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-001107 | codetables-gf2-74-9

Minimum distance of a binary [74, 9] linear code

codes are known with d = 32, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-001108 | codetables-gf2-75-10

Minimum distance of a binary [75, 10] linear code

codes are known with d = 32, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-001109 | codetables-gf2-75-13

Minimum distance of a binary [75, 13] linear code

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-001110 | codetables-gf2-75-14

Minimum distance of a binary [75, 14] linear code

codes are known with d = 29, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-001111 | codetables-gf2-75-9

Minimum distance of a binary [75, 9] linear code

codes are known with d = 33, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-001112 | codetables-gf2-76-10

Minimum distance of a binary [76, 10] linear code

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-001113 | codetables-gf2-76-13

Minimum distance of a binary [76, 13] linear code

codes are known with d = 31, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-001114 | codetables-gf2-76-14

Minimum distance of a binary [76, 14] linear code

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-001115 | codetables-gf2-77-10

Minimum distance of a binary [77, 10] linear code

codes are known with d = 33, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-001116 | codetables-gf2-77-11

Minimum distance of a binary [77, 11] linear code

codes are known with d = 32, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-001117 | codetables-gf2-77-14

Minimum distance of a binary [77, 14] linear code

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-001118 | codetables-gf2-77-9

Minimum distance of a binary [77, 9] linear code

codes are known with d = 34, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-001119 | codetables-gf2-78-10

Minimum distance of a binary [78, 10] linear code

codes are known with d = 34, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-001120 | codetables-gf2-78-11

Minimum distance of a binary [78, 11] linear code

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-001121 | codetables-gf2-78-12

Minimum distance of a binary [78, 12] linear code

codes are known with d = 32, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-001122 | codetables-gf2-78-14

Minimum distance of a binary [78, 14] linear code

codes are known with d = 31, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-001123 | codetables-gf2-78-9

Minimum distance of a binary [78, 9] linear code

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-001124 | codetables-gf2-79-10

Minimum distance of a binary [79, 10] linear code

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-001125 | codetables-gf2-79-11

Minimum distance of a binary [79, 11] linear code

codes are known with d = 33, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-001126 | codetables-gf2-79-12

Minimum distance of a binary [79, 12] linear code

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-001127 | codetables-gf2-79-13

Minimum distance of a binary [79, 13] linear code

codes are known with d = 32, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-001128 | codetables-gf2-80-10

Minimum distance of a binary [80, 10] linear code

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-001129 | codetables-gf2-80-11

Minimum distance of a binary [80, 11] linear code

codes are known with d = 34, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-001130 | codetables-gf2-80-12

Minimum distance of a binary [80, 12] linear code

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-001131 | codetables-gf2-80-13

Minimum distance of a binary [80, 13] linear code

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-001132 | codetables-gf2-80-14

Minimum distance of a binary [80, 14] linear code

codes are known with d = 32, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-001133 | codetables-gf2-81-11

Minimum distance of a binary [81, 11] linear code

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-001134 | codetables-gf2-81-12

Minimum distance of a binary [81, 12] linear code

codes are known with d = 32, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-001135 | codetables-gf2-81-13

Minimum distance of a binary [81, 13] linear code

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-001136 | codetables-gf2-81-14

Minimum distance of a binary [81, 14] linear code

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-001137 | codetables-gf2-81-9

Minimum distance of a binary [81, 9] linear code

codes are known with d = 36, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-001138 | codetables-gf2-82-11

Minimum distance of a binary [82, 11] linear code

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-001139 | codetables-gf2-82-12

Minimum distance of a binary [82, 12] linear code

codes are known with d = 33, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-001140 | codetables-gf2-82-13

Minimum distance of a binary [82, 13] linear code

codes are known with d = 32, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-001141 | codetables-gf2-82-14

Minimum distance of a binary [82, 14] linear code

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-001142 | codetables-gf2-82-9

Minimum distance of a binary [82, 9] linear code

codes are known with d = 37, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-001143 | codetables-gf2-83-12

Minimum distance of a binary [83, 12] linear code

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-001144 | codetables-gf2-83-13

Minimum distance of a binary [83, 13] linear code

codes are known with d = 32, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-001145 | codetables-gf2-83-14

Minimum distance of a binary [83, 14] linear code

codes are known with d = 32, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-001146 | codetables-gf2-84-11

Minimum distance of a binary [84, 11] linear code

codes are known with d = 36, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-001147 | codetables-gf2-84-12

Minimum distance of a binary [84, 12] linear code

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-001148 | codetables-gf2-84-13

Minimum distance of a binary [84, 13] linear code

codes are known with d = 33, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-001149 | codetables-gf2-84-14

Minimum distance of a binary [84, 14] linear code

codes are known with d = 32, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-001150 | codetables-gf2-84-9

Minimum distance of a binary [84, 9] linear code

codes are known with d = 38, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-001151 | codetables-gf2-85-11

Minimum distance of a binary [85, 11] linear code

codes are known with d = 36, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-001152 | codetables-gf2-85-12

Minimum distance of a binary [85, 12] linear code

codes are known with d = 36, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-001153 | codetables-gf2-85-13

Minimum distance of a binary [85, 13] linear code

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-001154 | codetables-gf2-85-14

Minimum distance of a binary [85, 14] linear code

codes are known with d = 32, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-001155 | codetables-gf2-85-9

Minimum distance of a binary [85, 9] linear code

codes are known with d = 39, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-001156 | codetables-gf2-86-11

Minimum distance of a binary [86, 11] linear code

codes are known with d = 37, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-001157 | codetables-gf2-86-12

Minimum distance of a binary [86, 12] linear code

codes are known with d = 36, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-001158 | codetables-gf2-86-13

Minimum distance of a binary [86, 13] linear code

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-001159 | codetables-gf2-86-14

Minimum distance of a binary [86, 14] linear code

codes are known with d = 33, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-001160 | codetables-gf2-87-12

Minimum distance of a binary [87, 12] linear code

codes are known with d = 36, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-001161 | codetables-gf2-87-13

Minimum distance of a binary [87, 13] linear code

codes are known with d = 35, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-001162 | codetables-gf2-87-14

Minimum distance of a binary [87, 14] linear code

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-001163 | codetables-gf2-88-12

Minimum distance of a binary [88, 12] linear code

codes are known with d = 37, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-001164 | codetables-gf2-88-13

Minimum distance of a binary [88, 13] linear code

codes are known with d = 36, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-001165 | codetables-gf2-88-14

Minimum distance of a binary [88, 14] linear code

codes are known with d = 34, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-001166 | codetables-gf2-89-12

Minimum distance of a binary [89, 12] linear code

codes are known with d = 38, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-001167 | codetables-gf2-89-13

Minimum distance of a binary [89, 13] linear code

codes are known with d = 36, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-001168 | codetables-gf2-89-14

Minimum distance of a binary [89, 14] linear code

codes are known with d = 35, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-001169 | codetables-gf2-89-9

Minimum distance of a binary [89, 9] linear code

codes are known with d = 40, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-001170 | codetables-gf2-90-12

Minimum distance of a binary [90, 12] linear code

codes are known with d = 39, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-001171 | codetables-gf2-90-13

Minimum distance of a binary [90, 13] linear code

codes are known with d = 36, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-001172 | codetables-gf2-90-14

Minimum distance of a binary [90, 14] linear code

codes are known with d = 36, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-001173 | codetables-gf2-90-9

Minimum distance of a binary [90, 9] linear code

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-001174 | codetables-gf2-91-10

Minimum distance of a binary [91, 10] linear code

codes are known with d = 40, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-001175 | codetables-gf2-91-13

Minimum distance of a binary [91, 13] linear code

codes are known with d = 37, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-001176 | codetables-gf2-91-14

Minimum distance of a binary [91, 14] linear code

codes are known with d = 36, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-001177 | codetables-gf2-91-9

Minimum distance of a binary [91, 9] linear code

codes are known with d = 41, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-001178 | codetables-gf2-92-10

Minimum distance of a binary [92, 10] linear code

codes are known with d = 41, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-001179 | codetables-gf2-92-11

Minimum distance of a binary [92, 11] linear code

codes are known with d = 40, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-001180 | codetables-gf2-92-13

Minimum distance of a binary [92, 13] linear code

codes are known with d = 38, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-001181 | codetables-gf2-92-14

Minimum distance of a binary [92, 14] linear code

codes are known with d = 36, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-001182 | codetables-gf2-92-9

Minimum distance of a binary [92, 9] linear code

codes are known with d = 42, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-001183 | codetables-gf2-93-11

Minimum distance of a binary [93, 11] linear code

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-001184 | codetables-gf2-93-12

Minimum distance of a binary [93, 12] linear code

codes are known with d = 40, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-001185 | codetables-gf2-93-13

Minimum distance of a binary [93, 13] linear code

codes are known with d = 38, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-001186 | codetables-gf2-93-14

Minimum distance of a binary [93, 14] linear code

codes are known with d = 36, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-001187 | codetables-gf2-93-9

Minimum distance of a binary [93, 9] linear code

codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-001188 | codetables-gf2-94-10

Minimum distance of a binary [94, 10] linear code

codes are known with d = 42, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-001189 | codetables-gf2-94-11

Minimum distance of a binary [94, 11] linear code

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-001190 | codetables-gf2-94-12

Minimum distance of a binary [94, 12] linear code

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-001191 | codetables-gf2-94-13

Minimum distance of a binary [94, 13] linear code

codes are known with d = 39, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-001192 | codetables-gf2-94-14

Minimum distance of a binary [94, 14] linear code

codes are known with d = 37, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-001193 | codetables-gf2-94-9

Minimum distance of a binary [94, 9] linear code

codes are known with d = 43, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-001194 | codetables-gf2-95-10

Minimum distance of a binary [95, 10] linear code

codes are known with d = 43, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-001195 | codetables-gf2-95-11

Minimum distance of a binary [95, 11] linear code

codes are known with d = 41, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-001196 | codetables-gf2-95-12

Minimum distance of a binary [95, 12] linear code

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-001197 | codetables-gf2-95-13

Minimum distance of a binary [95, 13] linear code

codes are known with d = 40, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-001198 | codetables-gf2-95-14

Minimum distance of a binary [95, 14] linear code

codes are known with d = 38, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-001199 | codetables-gf2-96-11

Minimum distance of a binary [96, 11] linear code

codes are known with d = 42, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-001200 | codetables-gf2-96-12

Minimum distance of a binary [96, 12] linear code

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-001201 | codetables-gf2-96-13

Minimum distance of a binary [96, 13] linear code

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-001202 | codetables-gf2-96-14

Minimum distance of a binary [96, 14] linear code

codes are known with d = 38, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-001203 | codetables-gf2-97-11

Minimum distance of a binary [97, 11] linear code

codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-001204 | codetables-gf2-97-12

Minimum distance of a binary [97, 12] linear code

codes are known with d = 40, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-001205 | codetables-gf2-97-13

Minimum distance of a binary [97, 13] linear code

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-001206 | codetables-gf2-97-14

Minimum distance of a binary [97, 14] linear code

codes are known with d = 39, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-001207 | codetables-gf2-97-9

Minimum distance of a binary [97, 9] linear code

codes are known with d = 44, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-001208 | codetables-gf2-98-11

Minimum distance of a binary [98, 11] linear code

codes are known with d = 43, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-001209 | codetables-gf2-98-12

Minimum distance of a binary [98, 12] linear code

codes are known with d = 41, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-001210 | codetables-gf2-98-13

Minimum distance of a binary [98, 13] linear code

codes are known with d = 40, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-001211 | codetables-gf2-98-14

Minimum distance of a binary [98, 14] linear code

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-001212 | codetables-gf2-98-9

Minimum distance of a binary [98, 9] linear code

codes are known with d = 45, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-001213 | codetables-gf2-99-12

Minimum distance of a binary [99, 12] linear code

codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-001214 | codetables-gf2-99-13

Minimum distance of a binary [99, 13] linear code

codes are known with d = 40, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-001215 | codetables-gf2-99-14

Minimum distance of a binary [99, 14] linear code

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

D03 非二元线性码 | 3,535 行

给定有限域大小 q、码长 n、维数 k，求线性 [n,k,d] 码能够达到的最大最小距离 d。

原始维护者：Markus Grassl; Andries E. Brouwer; Rumen Daskalov

Cite M. Grassl, "Bounds on the minimum distance of linear codes and quantum codes", online at http://www.codetables.de, and credit A. E. Brouwer's tables for the entries derived from them.

原始问题族入口

V3P-001216 | codetables-gf3-100-10

Minimum distance of a [100, 10] linear code over GF(3)

codes are known with d = 57, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-001217 | codetables-gf3-100-11

Minimum distance of a [100, 11] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-001218 | codetables-gf3-100-12

Minimum distance of a [100, 12] linear code over GF(3)

codes are known with d = 53, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-001219 | codetables-gf3-100-6

Minimum distance of a [100, 6] linear code over GF(3)

codes are known with d = 63, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-001220 | codetables-gf3-100-8

Minimum distance of a [100, 8] linear code over GF(3)

codes are known with d = 60, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-001221 | codetables-gf3-100-9

Minimum distance of a [100, 9] linear code over GF(3)

codes are known with d = 58, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-001222 | codetables-gf3-101-10

Minimum distance of a [101, 10] linear code over GF(3)

codes are known with d = 57, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-001223 | codetables-gf3-101-11

Minimum distance of a [101, 11] linear code over GF(3)

codes are known with d = 55, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-001224 | codetables-gf3-101-12

Minimum distance of a [101, 12] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-001225 | codetables-gf3-101-6

Minimum distance of a [101, 6] linear code over GF(3)

codes are known with d = 64, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-001226 | codetables-gf3-101-8

Minimum distance of a [101, 8] linear code over GF(3)

codes are known with d = 60, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-001227 | codetables-gf3-101-9

Minimum distance of a [101, 9] linear code over GF(3)

codes are known with d = 59, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-001228 | codetables-gf3-102-10

Minimum distance of a [102, 10] linear code over GF(3)

codes are known with d = 57, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-001229 | codetables-gf3-102-11

Minimum distance of a [102, 11] linear code over GF(3)

codes are known with d = 56, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-001230 | codetables-gf3-102-12

Minimum distance of a [102, 12] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-001231 | codetables-gf3-102-6

Minimum distance of a [102, 6] linear code over GF(3)

codes are known with d = 65, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-001232 | codetables-gf3-102-8

Minimum distance of a [102, 8] linear code over GF(3)

codes are known with d = 61, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-001233 | codetables-gf3-102-9

Minimum distance of a [102, 9] linear code over GF(3)

codes are known with d = 60, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-001234 | codetables-gf3-103-10

Minimum distance of a [103, 10] linear code over GF(3)

codes are known with d = 58, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-001235 | codetables-gf3-103-11

Minimum distance of a [103, 11] linear code over GF(3)

codes are known with d = 57, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-001236 | codetables-gf3-103-12

Minimum distance of a [103, 12] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-001237 | codetables-gf3-103-8

Minimum distance of a [103, 8] linear code over GF(3)

codes are known with d = 62, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-001238 | codetables-gf3-103-9

Minimum distance of a [103, 9] linear code over GF(3)

codes are known with d = 60, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-001239 | codetables-gf3-104-10

Minimum distance of a [104, 10] linear code over GF(3)

codes are known with d = 59, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-001240 | codetables-gf3-104-11

Minimum distance of a [104, 11] linear code over GF(3)

codes are known with d = 57, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-001241 | codetables-gf3-104-12

Minimum distance of a [104, 12] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-001242 | codetables-gf3-104-6

Minimum distance of a [104, 6] linear code over GF(3)

codes are known with d = 66, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-001243 | codetables-gf3-104-8

Minimum distance of a [104, 8] linear code over GF(3)

codes are known with d = 63, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-001244 | codetables-gf3-104-9

Minimum distance of a [104, 9] linear code over GF(3)

codes are known with d = 60, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-001245 | codetables-gf3-105-10

Minimum distance of a [105, 10] linear code over GF(3)

codes are known with d = 60, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-001246 | codetables-gf3-105-11

Minimum distance of a [105, 11] linear code over GF(3)

codes are known with d = 57, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-001247 | codetables-gf3-105-12

Minimum distance of a [105, 12] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-001248 | codetables-gf3-105-8

Minimum distance of a [105, 8] linear code over GF(3)

codes are known with d = 63, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-001249 | codetables-gf3-105-9

Minimum distance of a [105, 9] linear code over GF(3)

codes are known with d = 61, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-001250 | codetables-gf3-106-10

Minimum distance of a [106, 10] linear code over GF(3)

codes are known with d = 60, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-001251 | codetables-gf3-106-11

Minimum distance of a [106, 11] linear code over GF(3)

codes are known with d = 58, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-001252 | codetables-gf3-106-12

Minimum distance of a [106, 12] linear code over GF(3)

codes are known with d = 55, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-001253 | codetables-gf3-106-8

Minimum distance of a [106, 8] linear code over GF(3)

codes are known with d = 63, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-001254 | codetables-gf3-106-9

Minimum distance of a [106, 9] linear code over GF(3)

codes are known with d = 62, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-001255 | codetables-gf3-107-10

Minimum distance of a [107, 10] linear code over GF(3)

codes are known with d = 60, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-001256 | codetables-gf3-107-11

Minimum distance of a [107, 11] linear code over GF(3)

codes are known with d = 59, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-001257 | codetables-gf3-107-12

Minimum distance of a [107, 12] linear code over GF(3)

codes are known with d = 56, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-001258 | codetables-gf3-107-8

Minimum distance of a [107, 8] linear code over GF(3)

codes are known with d = 64, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-001259 | codetables-gf3-107-9

Minimum distance of a [107, 9] linear code over GF(3)

codes are known with d = 63, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-001260 | codetables-gf3-108-10

Minimum distance of a [108, 10] linear code over GF(3)

codes are known with d = 61, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-001261 | codetables-gf3-108-11

Minimum distance of a [108, 11] linear code over GF(3)

codes are known with d = 60, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-001262 | codetables-gf3-108-12

Minimum distance of a [108, 12] linear code over GF(3)

codes are known with d = 57, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-001263 | codetables-gf3-108-8

Minimum distance of a [108, 8] linear code over GF(3)

codes are known with d = 65, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-001264 | codetables-gf3-108-9

Minimum distance of a [108, 9] linear code over GF(3)

codes are known with d = 63, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-001265 | codetables-gf3-109-10

Minimum distance of a [109, 10] linear code over GF(3)

codes are known with d = 62, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-001266 | codetables-gf3-109-11

Minimum distance of a [109, 11] linear code over GF(3)

codes are known with d = 60, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-001267 | codetables-gf3-109-12

Minimum distance of a [109, 12] linear code over GF(3)

codes are known with d = 58, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-001268 | codetables-gf3-109-8

Minimum distance of a [109, 8] linear code over GF(3)

codes are known with d = 66, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-001269 | codetables-gf3-109-9

Minimum distance of a [109, 9] linear code over GF(3)

codes are known with d = 63, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-001270 | codetables-gf3-110-10

Minimum distance of a [110, 10] linear code over GF(3)

codes are known with d = 63, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-001271 | codetables-gf3-110-11

Minimum distance of a [110, 11] linear code over GF(3)

codes are known with d = 60, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-001272 | codetables-gf3-110-12

Minimum distance of a [110, 12] linear code over GF(3)

codes are known with d = 59, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-001273 | codetables-gf3-110-8

Minimum distance of a [110, 8] linear code over GF(3)

codes are known with d = 66, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-001274 | codetables-gf3-110-9

Minimum distance of a [110, 9] linear code over GF(3)

codes are known with d = 64, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-001275 | codetables-gf3-111-10

Minimum distance of a [111, 10] linear code over GF(3)

codes are known with d = 63, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-001276 | codetables-gf3-111-11

Minimum distance of a [111, 11] linear code over GF(3)

codes are known with d = 61, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-001277 | codetables-gf3-111-12

Minimum distance of a [111, 12] linear code over GF(3)

codes are known with d = 60, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-001278 | codetables-gf3-111-8

Minimum distance of a [111, 8] linear code over GF(3)

codes are known with d = 66, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-001279 | codetables-gf3-111-9

Minimum distance of a [111, 9] linear code over GF(3)

codes are known with d = 65, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-001280 | codetables-gf3-112-10

Minimum distance of a [112, 10] linear code over GF(3)

codes are known with d = 63, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-001281 | codetables-gf3-112-11

Minimum distance of a [112, 11] linear code over GF(3)

codes are known with d = 62, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-001282 | codetables-gf3-112-12

Minimum distance of a [112, 12] linear code over GF(3)

codes are known with d = 60, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-001283 | codetables-gf3-112-8

Minimum distance of a [112, 8] linear code over GF(3)

codes are known with d = 67, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-001284 | codetables-gf3-112-9

Minimum distance of a [112, 9] linear code over GF(3)

codes are known with d = 66, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-001285 | codetables-gf3-113-10

Minimum distance of a [113, 10] linear code over GF(3)

codes are known with d = 64, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-001286 | codetables-gf3-113-11

Minimum distance of a [113, 11] linear code over GF(3)

codes are known with d = 63, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-001287 | codetables-gf3-113-12

Minimum distance of a [113, 12] linear code over GF(3)

codes are known with d = 61, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-001288 | codetables-gf3-113-8

Minimum distance of a [113, 8] linear code over GF(3)

codes are known with d = 68, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-001289 | codetables-gf3-113-9

Minimum distance of a [113, 9] linear code over GF(3)

codes are known with d = 66, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-001290 | codetables-gf3-114-10

Minimum distance of a [114, 10] linear code over GF(3)

codes are known with d = 65, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-001291 | codetables-gf3-114-11

Minimum distance of a [114, 11] linear code over GF(3)

codes are known with d = 63, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-001292 | codetables-gf3-114-12

Minimum distance of a [114, 12] linear code over GF(3)

codes are known with d = 62, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-001293 | codetables-gf3-114-8

Minimum distance of a [114, 8] linear code over GF(3)

codes are known with d = 69, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-001294 | codetables-gf3-114-9

Minimum distance of a [114, 9] linear code over GF(3)

codes are known with d = 66, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-001295 | codetables-gf3-115-10

Minimum distance of a [115, 10] linear code over GF(3)

codes are known with d = 66, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-001296 | codetables-gf3-115-11

Minimum distance of a [115, 11] linear code over GF(3)

codes are known with d = 63, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-001297 | codetables-gf3-115-12

Minimum distance of a [115, 12] linear code over GF(3)

codes are known with d = 63, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-001298 | codetables-gf3-115-7

Minimum distance of a [115, 7] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-001299 | codetables-gf3-115-8

Minimum distance of a [115, 8] linear code over GF(3)

codes are known with d = 69, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-001300 | codetables-gf3-115-9

Minimum distance of a [115, 9] linear code over GF(3)

codes are known with d = 67, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-001301 | codetables-gf3-116-10

Minimum distance of a [116, 10] linear code over GF(3)

codes are known with d = 67, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-001302 | codetables-gf3-116-11

Minimum distance of a [116, 11] linear code over GF(3)

codes are known with d = 64, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-001303 | codetables-gf3-116-12

Minimum distance of a [116, 12] linear code over GF(3)

codes are known with d = 63, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-001304 | codetables-gf3-116-7

Minimum distance of a [116, 7] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-001305 | codetables-gf3-116-8

Minimum distance of a [116, 8] linear code over GF(3)

codes are known with d = 70, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-001306 | codetables-gf3-116-9

Minimum distance of a [116, 9] linear code over GF(3)

codes are known with d = 68, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-001307 | codetables-gf3-117-10

Minimum distance of a [117, 10] linear code over GF(3)

codes are known with d = 68, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-001308 | codetables-gf3-117-11

Minimum distance of a [117, 11] linear code over GF(3)

codes are known with d = 65, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-001309 | codetables-gf3-117-12

Minimum distance of a [117, 12] linear code over GF(3)

codes are known with d = 63, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-001310 | codetables-gf3-117-7

Minimum distance of a [117, 7] linear code over GF(3)

codes are known with d = 73, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-001311 | codetables-gf3-117-8

Minimum distance of a [117, 8] linear code over GF(3)

codes are known with d = 71, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-001312 | codetables-gf3-117-9

Minimum distance of a [117, 9] linear code over GF(3)

codes are known with d = 69, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-001313 | codetables-gf3-118-10

Minimum distance of a [118, 10] linear code over GF(3)

codes are known with d = 69, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-001314 | codetables-gf3-118-11

Minimum distance of a [118, 11] linear code over GF(3)

codes are known with d = 66, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-001315 | codetables-gf3-118-12

Minimum distance of a [118, 12] linear code over GF(3)

codes are known with d = 64, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-001316 | codetables-gf3-118-7

Minimum distance of a [118, 7] linear code over GF(3)

codes are known with d = 74, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-001317 | codetables-gf3-118-8

Minimum distance of a [118, 8] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-001318 | codetables-gf3-118-9

Minimum distance of a [118, 9] linear code over GF(3)

codes are known with d = 70, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-001319 | codetables-gf3-119-10

Minimum distance of a [119, 10] linear code over GF(3)

codes are known with d = 70, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-001320 | codetables-gf3-119-11

Minimum distance of a [119, 11] linear code over GF(3)

codes are known with d = 66, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-001321 | codetables-gf3-119-12

Minimum distance of a [119, 12] linear code over GF(3)

codes are known with d = 65, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-001322 | codetables-gf3-119-7

Minimum distance of a [119, 7] linear code over GF(3)

codes are known with d = 75, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-001323 | codetables-gf3-119-8

Minimum distance of a [119, 8] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-001324 | codetables-gf3-119-9

Minimum distance of a [119, 9] linear code over GF(3)

codes are known with d = 71, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-001325 | codetables-gf3-120-10

Minimum distance of a [120, 10] linear code over GF(3)

codes are known with d = 71, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-001326 | codetables-gf3-120-11

Minimum distance of a [120, 11] linear code over GF(3)

codes are known with d = 67, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-001327 | codetables-gf3-120-12

Minimum distance of a [120, 12] linear code over GF(3)

codes are known with d = 66, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-001328 | codetables-gf3-120-7

Minimum distance of a [120, 7] linear code over GF(3)

codes are known with d = 75, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-001329 | codetables-gf3-120-8

Minimum distance of a [120, 8] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-001330 | codetables-gf3-120-9

Minimum distance of a [120, 9] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-001331 | codetables-gf3-121-10

Minimum distance of a [121, 10] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-001332 | codetables-gf3-121-11

Minimum distance of a [121, 11] linear code over GF(3)

codes are known with d = 68, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-001333 | codetables-gf3-121-12

Minimum distance of a [121, 12] linear code over GF(3)

codes are known with d = 66, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-001334 | codetables-gf3-121-7

Minimum distance of a [121, 7] linear code over GF(3)

codes are known with d = 76, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-001335 | codetables-gf3-121-8

Minimum distance of a [121, 8] linear code over GF(3)

codes are known with d = 73, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-001336 | codetables-gf3-121-9

Minimum distance of a [121, 9] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-001337 | codetables-gf3-122-10

Minimum distance of a [122, 10] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-001338 | codetables-gf3-122-11

Minimum distance of a [122, 11] linear code over GF(3)

codes are known with d = 69, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-001339 | codetables-gf3-122-12

Minimum distance of a [122, 12] linear code over GF(3)

codes are known with d = 66, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-001340 | codetables-gf3-122-7

Minimum distance of a [122, 7] linear code over GF(3)

codes are known with d = 77, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-001341 | codetables-gf3-122-8

Minimum distance of a [122, 8] linear code over GF(3)

codes are known with d = 74, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-001342 | codetables-gf3-122-9

Minimum distance of a [122, 9] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-001343 | codetables-gf3-123-10

Minimum distance of a [123, 10] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-001344 | codetables-gf3-123-11

Minimum distance of a [123, 11] linear code over GF(3)

codes are known with d = 69, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-001345 | codetables-gf3-123-12

Minimum distance of a [123, 12] linear code over GF(3)

codes are known with d = 67, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-001346 | codetables-gf3-123-7

Minimum distance of a [123, 7] linear code over GF(3)

codes are known with d = 78, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-001347 | codetables-gf3-123-8

Minimum distance of a [123, 8] linear code over GF(3)

codes are known with d = 75, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-001348 | codetables-gf3-123-9

Minimum distance of a [123, 9] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-001349 | codetables-gf3-124-10

Minimum distance of a [124, 10] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-001350 | codetables-gf3-124-11

Minimum distance of a [124, 11] linear code over GF(3)

codes are known with d = 70, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-001351 | codetables-gf3-124-12

Minimum distance of a [124, 12] linear code over GF(3)

codes are known with d = 68, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-001352 | codetables-gf3-124-7

Minimum distance of a [124, 7] linear code over GF(3)

codes are known with d = 78, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-001353 | codetables-gf3-124-8

Minimum distance of a [124, 8] linear code over GF(3)

codes are known with d = 75, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-001354 | codetables-gf3-124-9

Minimum distance of a [124, 9] linear code over GF(3)

codes are known with d = 73, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-001355 | codetables-gf3-125-10

Minimum distance of a [125, 10] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-001356 | codetables-gf3-125-11

Minimum distance of a [125, 11] linear code over GF(3)

codes are known with d = 71, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-001357 | codetables-gf3-125-12

Minimum distance of a [125, 12] linear code over GF(3)

codes are known with d = 69, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-001358 | codetables-gf3-125-7

Minimum distance of a [125, 7] linear code over GF(3)

codes are known with d = 79, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-001359 | codetables-gf3-125-8

Minimum distance of a [125, 8] linear code over GF(3)

codes are known with d = 75, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-001360 | codetables-gf3-125-9

Minimum distance of a [125, 9] linear code over GF(3)

codes are known with d = 74, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-001361 | codetables-gf3-126-10

Minimum distance of a [126, 10] linear code over GF(3)

codes are known with d = 73, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-001362 | codetables-gf3-126-11

Minimum distance of a [126, 11] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-001363 | codetables-gf3-126-12

Minimum distance of a [126, 12] linear code over GF(3)

codes are known with d = 69, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-001364 | codetables-gf3-126-7

Minimum distance of a [126, 7] linear code over GF(3)

codes are known with d = 80, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-001365 | codetables-gf3-126-8

Minimum distance of a [126, 8] linear code over GF(3)

codes are known with d = 76, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-001366 | codetables-gf3-126-9

Minimum distance of a [126, 9] linear code over GF(3)

codes are known with d = 75, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-001367 | codetables-gf3-127-10

Minimum distance of a [127, 10] linear code over GF(3)

codes are known with d = 74, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-001368 | codetables-gf3-127-11

Minimum distance of a [127, 11] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-001369 | codetables-gf3-127-12

Minimum distance of a [127, 12] linear code over GF(3)

codes are known with d = 69, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-001370 | codetables-gf3-127-6

Minimum distance of a [127, 6] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-001371 | codetables-gf3-127-8

Minimum distance of a [127, 8] linear code over GF(3)

codes are known with d = 77, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-001372 | codetables-gf3-127-9

Minimum distance of a [127, 9] linear code over GF(3)

codes are known with d = 75, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-001373 | codetables-gf3-128-10

Minimum distance of a [128, 10] linear code over GF(3)

codes are known with d = 74, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-001374 | codetables-gf3-128-11

Minimum distance of a [128, 11] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-001375 | codetables-gf3-128-12

Minimum distance of a [128, 12] linear code over GF(3)

codes are known with d = 69, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-001376 | codetables-gf3-128-6

Minimum distance of a [128, 6] linear code over GF(3)

codes are known with d = 82, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-001377 | codetables-gf3-128-7

Minimum distance of a [128, 7] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-001378 | codetables-gf3-128-8

Minimum distance of a [128, 8] linear code over GF(3)

codes are known with d = 78, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-001379 | codetables-gf3-128-9

Minimum distance of a [128, 9] linear code over GF(3)

codes are known with d = 76, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-001380 | codetables-gf3-129-10

Minimum distance of a [129, 10] linear code over GF(3)

codes are known with d = 75, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-001381 | codetables-gf3-129-11

Minimum distance of a [129, 11] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-001382 | codetables-gf3-129-12

Minimum distance of a [129, 12] linear code over GF(3)

codes are known with d = 69, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-001383 | codetables-gf3-129-6

Minimum distance of a [129, 6] linear code over GF(3)

codes are known with d = 83, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-001384 | codetables-gf3-129-7

Minimum distance of a [129, 7] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-001385 | codetables-gf3-129-8

Minimum distance of a [129, 8] linear code over GF(3)

codes are known with d = 78, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-001386 | codetables-gf3-129-9

Minimum distance of a [129, 9] linear code over GF(3)

codes are known with d = 77, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-001387 | codetables-gf3-130-10

Minimum distance of a [130, 10] linear code over GF(3)

codes are known with d = 76, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-001388 | codetables-gf3-130-11

Minimum distance of a [130, 11] linear code over GF(3)

codes are known with d = 73, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-001389 | codetables-gf3-130-12

Minimum distance of a [130, 12] linear code over GF(3)

codes are known with d = 70, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-001390 | codetables-gf3-130-7

Minimum distance of a [130, 7] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-001391 | codetables-gf3-130-8

Minimum distance of a [130, 8] linear code over GF(3)

codes are known with d = 79, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-001392 | codetables-gf3-130-9

Minimum distance of a [130, 9] linear code over GF(3)

codes are known with d = 78, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-001393 | codetables-gf3-131-10

Minimum distance of a [131, 10] linear code over GF(3)

codes are known with d = 77, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-001394 | codetables-gf3-131-11

Minimum distance of a [131, 11] linear code over GF(3)

codes are known with d = 74, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-001395 | codetables-gf3-131-12

Minimum distance of a [131, 12] linear code over GF(3)

codes are known with d = 71, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-001396 | codetables-gf3-131-6

Minimum distance of a [131, 6] linear code over GF(3)

codes are known with d = 84, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-001397 | codetables-gf3-131-7

Minimum distance of a [131, 7] linear code over GF(3)

codes are known with d = 82, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-001398 | codetables-gf3-131-8

Minimum distance of a [131, 8] linear code over GF(3)

codes are known with d = 80, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-001399 | codetables-gf3-131-9

Minimum distance of a [131, 9] linear code over GF(3)

codes are known with d = 78, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-001400 | codetables-gf3-132-10

Minimum distance of a [132, 10] linear code over GF(3)

codes are known with d = 78, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-001401 | codetables-gf3-132-11

Minimum distance of a [132, 11] linear code over GF(3)

codes are known with d = 75, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-001402 | codetables-gf3-132-12

Minimum distance of a [132, 12] linear code over GF(3)

codes are known with d = 72, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-001403 | codetables-gf3-132-6

Minimum distance of a [132, 6] linear code over GF(3)

codes are known with d = 85, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-001404 | codetables-gf3-132-7

Minimum distance of a [132, 7] linear code over GF(3)

codes are known with d = 83, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-001405 | codetables-gf3-132-8

Minimum distance of a [132, 8] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-001406 | codetables-gf3-132-9

Minimum distance of a [132, 9] linear code over GF(3)

codes are known with d = 79, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-001407 | codetables-gf3-133-10

Minimum distance of a [133, 10] linear code over GF(3)

codes are known with d = 78, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-001408 | codetables-gf3-133-11

Minimum distance of a [133, 11] linear code over GF(3)

codes are known with d = 75, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-001409 | codetables-gf3-133-12

Minimum distance of a [133, 12] linear code over GF(3)

codes are known with d = 73, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-001410 | codetables-gf3-133-6

Minimum distance of a [133, 6] linear code over GF(3)

codes are known with d = 86, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-001411 | codetables-gf3-133-7

Minimum distance of a [133, 7] linear code over GF(3)

codes are known with d = 84, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-001412 | codetables-gf3-133-8

Minimum distance of a [133, 8] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-001413 | codetables-gf3-133-9

Minimum distance of a [133, 9] linear code over GF(3)

codes are known with d = 80, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-001414 | codetables-gf3-134-10

Minimum distance of a [134, 10] linear code over GF(3)

codes are known with d = 78, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-001415 | codetables-gf3-134-11

Minimum distance of a [134, 11] linear code over GF(3)

codes are known with d = 75, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-001416 | codetables-gf3-134-12

Minimum distance of a [134, 12] linear code over GF(3)

codes are known with d = 74, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-001417 | codetables-gf3-134-7

Minimum distance of a [134, 7] linear code over GF(3)

codes are known with d = 84, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-001418 | codetables-gf3-134-8

Minimum distance of a [134, 8] linear code over GF(3)

codes are known with d = 82, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-001419 | codetables-gf3-134-9

Minimum distance of a [134, 9] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-001420 | codetables-gf3-135-10

Minimum distance of a [135, 10] linear code over GF(3)

codes are known with d = 79, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-001421 | codetables-gf3-135-11

Minimum distance of a [135, 11] linear code over GF(3)

codes are known with d = 76, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-001422 | codetables-gf3-135-12

Minimum distance of a [135, 12] linear code over GF(3)

codes are known with d = 75, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-001423 | codetables-gf3-135-6

Minimum distance of a [135, 6] linear code over GF(3)

codes are known with d = 87, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-001424 | codetables-gf3-135-7

Minimum distance of a [135, 7] linear code over GF(3)

codes are known with d = 85, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-001425 | codetables-gf3-135-8

Minimum distance of a [135, 8] linear code over GF(3)

codes are known with d = 83, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-001426 | codetables-gf3-135-9

Minimum distance of a [135, 9] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-001427 | codetables-gf3-136-10

Minimum distance of a [136, 10] linear code over GF(3)

codes are known with d = 80, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-001428 | codetables-gf3-136-11

Minimum distance of a [136, 11] linear code over GF(3)

codes are known with d = 77, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-001429 | codetables-gf3-136-12

Minimum distance of a [136, 12] linear code over GF(3)

codes are known with d = 75, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-001430 | codetables-gf3-136-7

Minimum distance of a [136, 7] linear code over GF(3)

codes are known with d = 86, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-001431 | codetables-gf3-136-8

Minimum distance of a [136, 8] linear code over GF(3)

codes are known with d = 84, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-001432 | codetables-gf3-136-9

Minimum distance of a [136, 9] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-001433 | codetables-gf3-137-10

Minimum distance of a [137, 10] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-001434 | codetables-gf3-137-11

Minimum distance of a [137, 11] linear code over GF(3)

codes are known with d = 78, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-001435 | codetables-gf3-137-12

Minimum distance of a [137, 12] linear code over GF(3)

codes are known with d = 76, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-001436 | codetables-gf3-137-7

Minimum distance of a [137, 7] linear code over GF(3)

codes are known with d = 87, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-001437 | codetables-gf3-137-8

Minimum distance of a [137, 8] linear code over GF(3)

codes are known with d = 84, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-001438 | codetables-gf3-137-9

Minimum distance of a [137, 9] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-001439 | codetables-gf3-138-10

Minimum distance of a [138, 10] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-001440 | codetables-gf3-138-11

Minimum distance of a [138, 11] linear code over GF(3)

codes are known with d = 78, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-001441 | codetables-gf3-138-12

Minimum distance of a [138, 12] linear code over GF(3)

codes are known with d = 77, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-001442 | codetables-gf3-138-7

Minimum distance of a [138, 7] linear code over GF(3)

codes are known with d = 88, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-001443 | codetables-gf3-138-8

Minimum distance of a [138, 8] linear code over GF(3)

codes are known with d = 85, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-001444 | codetables-gf3-138-9

Minimum distance of a [138, 9] linear code over GF(3)

codes are known with d = 82, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-001445 | codetables-gf3-139-10

Minimum distance of a [139, 10] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-001446 | codetables-gf3-139-11

Minimum distance of a [139, 11] linear code over GF(3)

codes are known with d = 78, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-001447 | codetables-gf3-139-12

Minimum distance of a [139, 12] linear code over GF(3)

codes are known with d = 78, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-001448 | codetables-gf3-139-7

Minimum distance of a [139, 7] linear code over GF(3)

codes are known with d = 89, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-001449 | codetables-gf3-139-8

Minimum distance of a [139, 8] linear code over GF(3)

codes are known with d = 86, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-001450 | codetables-gf3-139-9

Minimum distance of a [139, 9] linear code over GF(3)

codes are known with d = 83, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-001451 | codetables-gf3-140-10

Minimum distance of a [140, 10] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-001452 | codetables-gf3-140-11

Minimum distance of a [140, 11] linear code over GF(3)

codes are known with d = 79, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-001453 | codetables-gf3-140-12

Minimum distance of a [140, 12] linear code over GF(3)

codes are known with d = 78, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-001454 | codetables-gf3-140-6

Minimum distance of a [140, 6] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-001455 | codetables-gf3-140-8

Minimum distance of a [140, 8] linear code over GF(3)

codes are known with d = 87, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-001456 | codetables-gf3-140-9

Minimum distance of a [140, 9] linear code over GF(3)

codes are known with d = 84, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-001457 | codetables-gf3-141-10

Minimum distance of a [141, 10] linear code over GF(3)

codes are known with d = 82, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-001458 | codetables-gf3-141-11

Minimum distance of a [141, 11] linear code over GF(3)

codes are known with d = 80, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-001459 | codetables-gf3-141-12

Minimum distance of a [141, 12] linear code over GF(3)

codes are known with d = 78, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-001460 | codetables-gf3-141-6

Minimum distance of a [141, 6] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-001461 | codetables-gf3-141-8

Minimum distance of a [141, 8] linear code over GF(3)

codes are known with d = 87, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-001462 | codetables-gf3-141-9

Minimum distance of a [141, 9] linear code over GF(3)

codes are known with d = 84, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-001463 | codetables-gf3-142-10

Minimum distance of a [142, 10] linear code over GF(3)

codes are known with d = 83, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-001464 | codetables-gf3-142-11

Minimum distance of a [142, 11] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-001465 | codetables-gf3-142-12

Minimum distance of a [142, 12] linear code over GF(3)

codes are known with d = 78, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-001466 | codetables-gf3-142-6

Minimum distance of a [142, 6] linear code over GF(3)

codes are known with d = 91, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-001467 | codetables-gf3-142-7

Minimum distance of a [142, 7] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-001468 | codetables-gf3-142-8

Minimum distance of a [142, 8] linear code over GF(3)

codes are known with d = 88, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-001469 | codetables-gf3-142-9

Minimum distance of a [142, 9] linear code over GF(3)

codes are known with d = 84, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-001470 | codetables-gf3-143-10

Minimum distance of a [143, 10] linear code over GF(3)

codes are known with d = 84, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-001471 | codetables-gf3-143-11

Minimum distance of a [143, 11] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-001472 | codetables-gf3-143-12

Minimum distance of a [143, 12] linear code over GF(3)

codes are known with d = 79, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-001473 | codetables-gf3-143-6

Minimum distance of a [143, 6] linear code over GF(3)

codes are known with d = 92, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-001474 | codetables-gf3-143-7

Minimum distance of a [143, 7] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-001475 | codetables-gf3-143-8

Minimum distance of a [143, 8] linear code over GF(3)

codes are known with d = 89, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-001476 | codetables-gf3-143-9

Minimum distance of a [143, 9] linear code over GF(3)

codes are known with d = 85, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-001477 | codetables-gf3-144-10

Minimum distance of a [144, 10] linear code over GF(3)

codes are known with d = 84, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-001478 | codetables-gf3-144-11

Minimum distance of a [144, 11] linear code over GF(3)

codes are known with d = 82, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-001479 | codetables-gf3-144-12

Minimum distance of a [144, 12] linear code over GF(3)

codes are known with d = 80, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-001480 | codetables-gf3-144-7

Minimum distance of a [144, 7] linear code over GF(3)

codes are known with d = 91, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-001481 | codetables-gf3-144-8

Minimum distance of a [144, 8] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-001482 | codetables-gf3-144-9

Minimum distance of a [144, 9] linear code over GF(3)

codes are known with d = 86, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-001483 | codetables-gf3-145-10

Minimum distance of a [145, 10] linear code over GF(3)

codes are known with d = 84, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-001484 | codetables-gf3-145-11

Minimum distance of a [145, 11] linear code over GF(3)

codes are known with d = 83, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-001485 | codetables-gf3-145-12

Minimum distance of a [145, 12] linear code over GF(3)

codes are known with d = 81, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-001486 | codetables-gf3-145-6

Minimum distance of a [145, 6] linear code over GF(3)

codes are known with d = 93, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-001487 | codetables-gf3-145-7

Minimum distance of a [145, 7] linear code over GF(3)

codes are known with d = 92, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-001488 | codetables-gf3-145-8

Minimum distance of a [145, 8] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-001489 | codetables-gf3-145-9

Minimum distance of a [145, 9] linear code over GF(3)

codes are known with d = 87, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-001490 | codetables-gf3-146-10

Minimum distance of a [146, 10] linear code over GF(3)

codes are known with d = 85, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-001491 | codetables-gf3-146-11

Minimum distance of a [146, 11] linear code over GF(3)

codes are known with d = 84, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-001492 | codetables-gf3-146-12

Minimum distance of a [146, 12] linear code over GF(3)

codes are known with d = 82, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-001493 | codetables-gf3-146-6

Minimum distance of a [146, 6] linear code over GF(3)

codes are known with d = 94, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-001494 | codetables-gf3-146-8

Minimum distance of a [146, 8] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-001495 | codetables-gf3-146-9

Minimum distance of a [146, 9] linear code over GF(3)

codes are known with d = 87, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-001496 | codetables-gf3-147-10

Minimum distance of a [147, 10] linear code over GF(3)

codes are known with d = 86, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-001497 | codetables-gf3-147-11

Minimum distance of a [147, 11] linear code over GF(3)

codes are known with d = 84, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-001498 | codetables-gf3-147-12

Minimum distance of a [147, 12] linear code over GF(3)

codes are known with d = 82, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-001499 | codetables-gf3-147-6

Minimum distance of a [147, 6] linear code over GF(3)

codes are known with d = 95, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-001500 | codetables-gf3-147-7

Minimum distance of a [147, 7] linear code over GF(3)

codes are known with d = 93, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-001501 | codetables-gf3-147-8

Minimum distance of a [147, 8] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-001502 | codetables-gf3-147-9

Minimum distance of a [147, 9] linear code over GF(3)

codes are known with d = 88, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-001503 | codetables-gf3-148-10

Minimum distance of a [148, 10] linear code over GF(3)

codes are known with d = 87, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-001504 | codetables-gf3-148-11

Minimum distance of a [148, 11] linear code over GF(3)

codes are known with d = 84, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-001505 | codetables-gf3-148-12

Minimum distance of a [148, 12] linear code over GF(3)

codes are known with d = 82, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-001506 | codetables-gf3-148-7

Minimum distance of a [148, 7] linear code over GF(3)

codes are known with d = 94, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-001507 | codetables-gf3-148-8

Minimum distance of a [148, 8] linear code over GF(3)

codes are known with d = 91, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-001508 | codetables-gf3-148-9

Minimum distance of a [148, 9] linear code over GF(3)

codes are known with d = 89, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-001509 | codetables-gf3-149-10

Minimum distance of a [149, 10] linear code over GF(3)

codes are known with d = 87, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-001510 | codetables-gf3-149-11

Minimum distance of a [149, 11] linear code over GF(3)

codes are known with d = 84, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-001511 | codetables-gf3-149-12

Minimum distance of a [149, 12] linear code over GF(3)

codes are known with d = 82, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-001512 | codetables-gf3-149-6

Minimum distance of a [149, 6] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-001513 | codetables-gf3-149-7

Minimum distance of a [149, 7] linear code over GF(3)

codes are known with d = 95, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-001514 | codetables-gf3-149-8

Minimum distance of a [149, 8] linear code over GF(3)

codes are known with d = 92, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-001515 | codetables-gf3-149-9

Minimum distance of a [149, 9] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-001516 | codetables-gf3-150-10

Minimum distance of a [150, 10] linear code over GF(3)

codes are known with d = 87, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-001517 | codetables-gf3-150-11

Minimum distance of a [150, 11] linear code over GF(3)

codes are known with d = 85, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-001518 | codetables-gf3-150-12

Minimum distance of a [150, 12] linear code over GF(3)

codes are known with d = 83, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-001519 | codetables-gf3-150-6

Minimum distance of a [150, 6] linear code over GF(3)

codes are known with d = 97, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-001520 | codetables-gf3-150-8

Minimum distance of a [150, 8] linear code over GF(3)

codes are known with d = 93, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-001521 | codetables-gf3-150-9

Minimum distance of a [150, 9] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-001522 | codetables-gf3-151-10

Minimum distance of a [151, 10] linear code over GF(3)

codes are known with d = 88, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-001523 | codetables-gf3-151-11

Minimum distance of a [151, 11] linear code over GF(3)

codes are known with d = 86, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-001524 | codetables-gf3-151-12

Minimum distance of a [151, 12] linear code over GF(3)

codes are known with d = 83, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-001525 | codetables-gf3-151-6

Minimum distance of a [151, 6] linear code over GF(3)

codes are known with d = 98, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-001526 | codetables-gf3-151-7

Minimum distance of a [151, 7] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-001527 | codetables-gf3-151-8

Minimum distance of a [151, 8] linear code over GF(3)

codes are known with d = 93, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-001528 | codetables-gf3-151-9

Minimum distance of a [151, 9] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-001529 | codetables-gf3-152-10

Minimum distance of a [152, 10] linear code over GF(3)

codes are known with d = 89, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-001530 | codetables-gf3-152-11

Minimum distance of a [152, 11] linear code over GF(3)

codes are known with d = 87, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-001531 | codetables-gf3-152-12

Minimum distance of a [152, 12] linear code over GF(3)

codes are known with d = 84, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-001532 | codetables-gf3-152-7

Minimum distance of a [152, 7] linear code over GF(3)

codes are known with d = 97, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-001533 | codetables-gf3-152-8

Minimum distance of a [152, 8] linear code over GF(3)

codes are known with d = 94, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-001534 | codetables-gf3-152-9

Minimum distance of a [152, 9] linear code over GF(3)

codes are known with d = 91, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-001535 | codetables-gf3-153-10

Minimum distance of a [153, 10] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-001536 | codetables-gf3-153-11

Minimum distance of a [153, 11] linear code over GF(3)

codes are known with d = 87, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-001537 | codetables-gf3-153-12

Minimum distance of a [153, 12] linear code over GF(3)

codes are known with d = 85, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-001538 | codetables-gf3-153-6

Minimum distance of a [153, 6] linear code over GF(3)

codes are known with d = 99, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-001539 | codetables-gf3-153-7

Minimum distance of a [153, 7] linear code over GF(3)

codes are known with d = 98, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-001540 | codetables-gf3-153-8

Minimum distance of a [153, 8] linear code over GF(3)

codes are known with d = 95, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-001541 | codetables-gf3-153-9

Minimum distance of a [153, 9] linear code over GF(3)

codes are known with d = 92, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-001542 | codetables-gf3-154-10

Minimum distance of a [154, 10] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-001543 | codetables-gf3-154-11

Minimum distance of a [154, 11] linear code over GF(3)

codes are known with d = 88, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-001544 | codetables-gf3-154-12

Minimum distance of a [154, 12] linear code over GF(3)

codes are known with d = 86, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-001545 | codetables-gf3-154-8

Minimum distance of a [154, 8] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-001546 | codetables-gf3-154-9

Minimum distance of a [154, 9] linear code over GF(3)

codes are known with d = 93, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-001547 | codetables-gf3-155-10

Minimum distance of a [155, 10] linear code over GF(3)

codes are known with d = 91, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-001548 | codetables-gf3-155-11

Minimum distance of a [155, 11] linear code over GF(3)

codes are known with d = 89, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-001549 | codetables-gf3-155-12

Minimum distance of a [155, 12] linear code over GF(3)

codes are known with d = 87, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-001550 | codetables-gf3-155-7

Minimum distance of a [155, 7] linear code over GF(3)

codes are known with d = 99, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-001551 | codetables-gf3-155-8

Minimum distance of a [155, 8] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-001552 | codetables-gf3-155-9

Minimum distance of a [155, 9] linear code over GF(3)

codes are known with d = 93, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-001553 | codetables-gf3-156-10

Minimum distance of a [156, 10] linear code over GF(3)

codes are known with d = 92, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-001554 | codetables-gf3-156-11

Minimum distance of a [156, 11] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-001555 | codetables-gf3-156-12

Minimum distance of a [156, 12] linear code over GF(3)

codes are known with d = 87, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-001556 | codetables-gf3-156-7

Minimum distance of a [156, 7] linear code over GF(3)

codes are known with d = 99, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-001557 | codetables-gf3-156-8

Minimum distance of a [156, 8] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-001558 | codetables-gf3-156-9

Minimum distance of a [156, 9] linear code over GF(3)

codes are known with d = 94, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-001559 | codetables-gf3-157-10

Minimum distance of a [157, 10] linear code over GF(3)

codes are known with d = 93, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-001560 | codetables-gf3-157-11

Minimum distance of a [157, 11] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-001561 | codetables-gf3-157-12

Minimum distance of a [157, 12] linear code over GF(3)

codes are known with d = 88, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-001562 | codetables-gf3-157-7

Minimum distance of a [157, 7] linear code over GF(3)

codes are known with d = 99, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-001563 | codetables-gf3-157-8

Minimum distance of a [157, 8] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-001564 | codetables-gf3-157-9

Minimum distance of a [157, 9] linear code over GF(3)

codes are known with d = 95, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-001565 | codetables-gf3-158-10

Minimum distance of a [158, 10] linear code over GF(3)

codes are known with d = 94, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-001566 | codetables-gf3-158-11

Minimum distance of a [158, 11] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-001567 | codetables-gf3-158-12

Minimum distance of a [158, 12] linear code over GF(3)

codes are known with d = 89, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-001568 | codetables-gf3-158-7

Minimum distance of a [158, 7] linear code over GF(3)

codes are known with d = 100, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-001569 | codetables-gf3-158-8

Minimum distance of a [158, 8] linear code over GF(3)

codes are known with d = 97, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-001570 | codetables-gf3-158-9

Minimum distance of a [158, 9] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-001571 | codetables-gf3-159-10

Minimum distance of a [159, 10] linear code over GF(3)

codes are known with d = 95, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-001572 | codetables-gf3-159-11

Minimum distance of a [159, 11] linear code over GF(3)

codes are known with d = 91, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-001573 | codetables-gf3-159-12

Minimum distance of a [159, 12] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-001574 | codetables-gf3-159-7

Minimum distance of a [159, 7] linear code over GF(3)

codes are known with d = 101, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-001575 | codetables-gf3-159-8

Minimum distance of a [159, 8] linear code over GF(3)

codes are known with d = 98, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-001576 | codetables-gf3-159-9

Minimum distance of a [159, 9] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-001577 | codetables-gf3-160-10

Minimum distance of a [160, 10] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-001578 | codetables-gf3-160-11

Minimum distance of a [160, 11] linear code over GF(3)

codes are known with d = 92, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-001579 | codetables-gf3-160-12

Minimum distance of a [160, 12] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-001580 | codetables-gf3-160-7

Minimum distance of a [160, 7] linear code over GF(3)

codes are known with d = 102, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-001581 | codetables-gf3-160-8

Minimum distance of a [160, 8] linear code over GF(3)

codes are known with d = 99, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-001582 | codetables-gf3-160-9

Minimum distance of a [160, 9] linear code over GF(3)

codes are known with d = 97, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-001583 | codetables-gf3-161-10

Minimum distance of a [161, 10] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-001584 | codetables-gf3-161-11

Minimum distance of a [161, 11] linear code over GF(3)

codes are known with d = 93, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-001585 | codetables-gf3-161-12

Minimum distance of a [161, 12] linear code over GF(3)

codes are known with d = 90, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-001586 | codetables-gf3-161-7

Minimum distance of a [161, 7] linear code over GF(3)

codes are known with d = 102, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-001587 | codetables-gf3-161-8

Minimum distance of a [161, 8] linear code over GF(3)

codes are known with d = 99, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-001588 | codetables-gf3-161-9

Minimum distance of a [161, 9] linear code over GF(3)

codes are known with d = 98, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-001589 | codetables-gf3-162-10

Minimum distance of a [162, 10] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-001590 | codetables-gf3-162-11

Minimum distance of a [162, 11] linear code over GF(3)

codes are known with d = 93, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-001591 | codetables-gf3-162-12

Minimum distance of a [162, 12] linear code over GF(3)

codes are known with d = 91, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-001592 | codetables-gf3-162-7

Minimum distance of a [162, 7] linear code over GF(3)

codes are known with d = 103, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-001593 | codetables-gf3-162-8

Minimum distance of a [162, 8] linear code over GF(3)

codes are known with d = 100, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-001594 | codetables-gf3-162-9

Minimum distance of a [162, 9] linear code over GF(3)

codes are known with d = 99, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-001595 | codetables-gf3-163-10

Minimum distance of a [163, 10] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-001596 | codetables-gf3-163-11

Minimum distance of a [163, 11] linear code over GF(3)

codes are known with d = 93, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-001597 | codetables-gf3-163-12

Minimum distance of a [163, 12] linear code over GF(3)

codes are known with d = 92, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-001598 | codetables-gf3-163-7

Minimum distance of a [163, 7] linear code over GF(3)

codes are known with d = 104, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-001599 | codetables-gf3-163-8

Minimum distance of a [163, 8] linear code over GF(3)

codes are known with d = 101, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-001600 | codetables-gf3-163-9

Minimum distance of a [163, 9] linear code over GF(3)

codes are known with d = 99, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-001601 | codetables-gf3-164-10

Minimum distance of a [164, 10] linear code over GF(3)

codes are known with d = 97, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-001602 | codetables-gf3-164-11

Minimum distance of a [164, 11] linear code over GF(3)

codes are known with d = 94, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-001603 | codetables-gf3-164-12

Minimum distance of a [164, 12] linear code over GF(3)

codes are known with d = 93, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-001604 | codetables-gf3-164-7

Minimum distance of a [164, 7] linear code over GF(3)

codes are known with d = 105, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-001605 | codetables-gf3-164-8

Minimum distance of a [164, 8] linear code over GF(3)

codes are known with d = 102, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-001606 | codetables-gf3-164-9

Minimum distance of a [164, 9] linear code over GF(3)

codes are known with d = 99, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-001607 | codetables-gf3-165-10

Minimum distance of a [165, 10] linear code over GF(3)

codes are known with d = 98, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-001608 | codetables-gf3-165-11

Minimum distance of a [165, 11] linear code over GF(3)

codes are known with d = 95, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-001609 | codetables-gf3-165-12

Minimum distance of a [165, 12] linear code over GF(3)

codes are known with d = 93, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-001610 | codetables-gf3-165-7

Minimum distance of a [165, 7] linear code over GF(3)

codes are known with d = 105, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-001611 | codetables-gf3-165-8

Minimum distance of a [165, 8] linear code over GF(3)

codes are known with d = 102, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-001612 | codetables-gf3-165-9

Minimum distance of a [165, 9] linear code over GF(3)

codes are known with d = 99, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-001613 | codetables-gf3-166-10

Minimum distance of a [166, 10] linear code over GF(3)

codes are known with d = 98, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-001614 | codetables-gf3-166-11

Minimum distance of a [166, 11] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-001615 | codetables-gf3-166-12

Minimum distance of a [166, 12] linear code over GF(3)

codes are known with d = 94, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-001616 | codetables-gf3-166-7

Minimum distance of a [166, 7] linear code over GF(3)

codes are known with d = 106, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-001617 | codetables-gf3-166-8

Minimum distance of a [166, 8] linear code over GF(3)

codes are known with d = 103, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-001618 | codetables-gf3-166-9

Minimum distance of a [166, 9] linear code over GF(3)

codes are known with d = 100, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-001619 | codetables-gf3-167-10

Minimum distance of a [167, 10] linear code over GF(3)

codes are known with d = 99, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-001620 | codetables-gf3-167-11

Minimum distance of a [167, 11] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-001621 | codetables-gf3-167-12

Minimum distance of a [167, 12] linear code over GF(3)

codes are known with d = 95, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-001622 | codetables-gf3-167-7

Minimum distance of a [167, 7] linear code over GF(3)

codes are known with d = 107, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-001623 | codetables-gf3-167-8

Minimum distance of a [167, 8] linear code over GF(3)

codes are known with d = 104, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-001624 | codetables-gf3-167-9

Minimum distance of a [167, 9] linear code over GF(3)

codes are known with d = 101, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-001625 | codetables-gf3-168-10

Minimum distance of a [168, 10] linear code over GF(3)

codes are known with d = 100, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-001626 | codetables-gf3-168-11

Minimum distance of a [168, 11] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-001627 | codetables-gf3-168-12

Minimum distance of a [168, 12] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-001628 | codetables-gf3-168-8

Minimum distance of a [168, 8] linear code over GF(3)

codes are known with d = 105, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-001629 | codetables-gf3-168-9

Minimum distance of a [168, 9] linear code over GF(3)

codes are known with d = 102, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-001630 | codetables-gf3-169-10

Minimum distance of a [169, 10] linear code over GF(3)

codes are known with d = 101, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-001631 | codetables-gf3-169-11

Minimum distance of a [169, 11] linear code over GF(3)

codes are known with d = 97, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-001632 | codetables-gf3-169-12

Minimum distance of a [169, 12] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-001633 | codetables-gf3-169-7

Minimum distance of a [169, 7] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-001634 | codetables-gf3-169-8

Minimum distance of a [169, 8] linear code over GF(3)

codes are known with d = 105, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-001635 | codetables-gf3-169-9

Minimum distance of a [169, 9] linear code over GF(3)

codes are known with d = 102, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-001636 | codetables-gf3-170-10

Minimum distance of a [170, 10] linear code over GF(3)

codes are known with d = 102, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-001637 | codetables-gf3-170-11

Minimum distance of a [170, 11] linear code over GF(3)

codes are known with d = 98, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-001638 | codetables-gf3-170-12

Minimum distance of a [170, 12] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-001639 | codetables-gf3-170-7

Minimum distance of a [170, 7] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-001640 | codetables-gf3-170-8

Minimum distance of a [170, 8] linear code over GF(3)

codes are known with d = 106, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-001641 | codetables-gf3-170-9

Minimum distance of a [170, 9] linear code over GF(3)

codes are known with d = 103, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-001642 | codetables-gf3-171-10

Minimum distance of a [171, 10] linear code over GF(3)

codes are known with d = 102, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-001643 | codetables-gf3-171-11

Minimum distance of a [171, 11] linear code over GF(3)

codes are known with d = 99, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-001644 | codetables-gf3-171-12

Minimum distance of a [171, 12] linear code over GF(3)

codes are known with d = 96, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-001645 | codetables-gf3-171-7

Minimum distance of a [171, 7] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-001646 | codetables-gf3-171-8

Minimum distance of a [171, 8] linear code over GF(3)

codes are known with d = 107, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-001647 | codetables-gf3-171-9

Minimum distance of a [171, 9] linear code over GF(3)

codes are known with d = 104, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-001648 | codetables-gf3-172-10

Minimum distance of a [172, 10] linear code over GF(3)

codes are known with d = 102, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-001649 | codetables-gf3-172-11

Minimum distance of a [172, 11] linear code over GF(3)

codes are known with d = 99, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-001650 | codetables-gf3-172-12

Minimum distance of a [172, 12] linear code over GF(3)

codes are known with d = 97, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-001651 | codetables-gf3-172-7

Minimum distance of a [172, 7] linear code over GF(3)

codes are known with d = 109, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-001652 | codetables-gf3-172-8

Minimum distance of a [172, 8] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-001653 | codetables-gf3-172-9

Minimum distance of a [172, 9] linear code over GF(3)

codes are known with d = 105, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-001654 | codetables-gf3-173-10

Minimum distance of a [173, 10] linear code over GF(3)

codes are known with d = 103, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-001655 | codetables-gf3-173-11

Minimum distance of a [173, 11] linear code over GF(3)

codes are known with d = 100, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-001656 | codetables-gf3-173-12

Minimum distance of a [173, 12] linear code over GF(3)

codes are known with d = 98, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-001657 | codetables-gf3-173-7

Minimum distance of a [173, 7] linear code over GF(3)

codes are known with d = 110, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-001658 | codetables-gf3-173-8

Minimum distance of a [173, 8] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-001659 | codetables-gf3-173-9

Minimum distance of a [173, 9] linear code over GF(3)

codes are known with d = 105, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-001660 | codetables-gf3-174-10

Minimum distance of a [174, 10] linear code over GF(3)

codes are known with d = 104, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-001661 | codetables-gf3-174-11

Minimum distance of a [174, 11] linear code over GF(3)

codes are known with d = 101, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-001662 | codetables-gf3-174-12

Minimum distance of a [174, 12] linear code over GF(3)

codes are known with d = 98, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-001663 | codetables-gf3-174-7

Minimum distance of a [174, 7] linear code over GF(3)

codes are known with d = 111, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-001664 | codetables-gf3-174-8

Minimum distance of a [174, 8] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-001665 | codetables-gf3-174-9

Minimum distance of a [174, 9] linear code over GF(3)

codes are known with d = 106, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-001666 | codetables-gf3-175-10

Minimum distance of a [175, 10] linear code over GF(3)

codes are known with d = 104, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-001667 | codetables-gf3-175-11

Minimum distance of a [175, 11] linear code over GF(3)

codes are known with d = 102, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-001668 | codetables-gf3-175-12

Minimum distance of a [175, 12] linear code over GF(3)

codes are known with d = 99, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-001669 | codetables-gf3-175-7

Minimum distance of a [175, 7] linear code over GF(3)

codes are known with d = 111, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-001670 | codetables-gf3-175-8

Minimum distance of a [175, 8] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-001671 | codetables-gf3-175-9

Minimum distance of a [175, 9] linear code over GF(3)

codes are known with d = 107, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-001672 | codetables-gf3-176-10

Minimum distance of a [176, 10] linear code over GF(3)

codes are known with d = 105, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-001673 | codetables-gf3-176-11

Minimum distance of a [176, 11] linear code over GF(3)

codes are known with d = 102, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-001674 | codetables-gf3-176-12

Minimum distance of a [176, 12] linear code over GF(3)

codes are known with d = 100, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-001675 | codetables-gf3-176-7

Minimum distance of a [176, 7] linear code over GF(3)

codes are known with d = 112, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-001676 | codetables-gf3-176-8

Minimum distance of a [176, 8] linear code over GF(3)

codes are known with d = 109, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-001677 | codetables-gf3-176-9

Minimum distance of a [176, 9] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-001678 | codetables-gf3-177-10

Minimum distance of a [177, 10] linear code over GF(3)

codes are known with d = 105, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-001679 | codetables-gf3-177-11

Minimum distance of a [177, 11] linear code over GF(3)

codes are known with d = 102, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-001680 | codetables-gf3-177-12

Minimum distance of a [177, 12] linear code over GF(3)

codes are known with d = 101, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-001681 | codetables-gf3-177-7

Minimum distance of a [177, 7] linear code over GF(3)

codes are known with d = 113, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-001682 | codetables-gf3-177-8

Minimum distance of a [177, 8] linear code over GF(3)

codes are known with d = 110, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-001683 | codetables-gf3-177-9

Minimum distance of a [177, 9] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-001684 | codetables-gf3-178-10

Minimum distance of a [178, 10] linear code over GF(3)

codes are known with d = 106, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-001685 | codetables-gf3-178-11

Minimum distance of a [178, 11] linear code over GF(3)

codes are known with d = 103, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-001686 | codetables-gf3-178-12

Minimum distance of a [178, 12] linear code over GF(3)

codes are known with d = 102, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-001687 | codetables-gf3-178-7

Minimum distance of a [178, 7] linear code over GF(3)

codes are known with d = 114, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-001688 | codetables-gf3-178-8

Minimum distance of a [178, 8] linear code over GF(3)

codes are known with d = 111, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-001689 | codetables-gf3-178-9

Minimum distance of a [178, 9] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-001690 | codetables-gf3-179-10

Minimum distance of a [179, 10] linear code over GF(3)

codes are known with d = 107, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-001691 | codetables-gf3-179-11

Minimum distance of a [179, 11] linear code over GF(3)

codes are known with d = 104, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-001692 | codetables-gf3-179-12

Minimum distance of a [179, 12] linear code over GF(3)

codes are known with d = 102, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-001693 | codetables-gf3-179-7

Minimum distance of a [179, 7] linear code over GF(3)

codes are known with d = 114, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-001694 | codetables-gf3-179-8

Minimum distance of a [179, 8] linear code over GF(3)

codes are known with d = 111, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-001695 | codetables-gf3-179-9

Minimum distance of a [179, 9] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-001696 | codetables-gf3-180-10

Minimum distance of a [180, 10] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-001697 | codetables-gf3-180-11

Minimum distance of a [180, 11] linear code over GF(3)

codes are known with d = 105, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-001698 | codetables-gf3-180-12

Minimum distance of a [180, 12] linear code over GF(3)

codes are known with d = 103, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-001699 | codetables-gf3-180-7

Minimum distance of a [180, 7] linear code over GF(3)

codes are known with d = 115, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-001700 | codetables-gf3-180-8

Minimum distance of a [180, 8] linear code over GF(3)

codes are known with d = 112, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-001701 | codetables-gf3-180-9

Minimum distance of a [180, 9] linear code over GF(3)

codes are known with d = 109, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-001702 | codetables-gf3-181-10

Minimum distance of a [181, 10] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-001703 | codetables-gf3-181-11

Minimum distance of a [181, 11] linear code over GF(3)

codes are known with d = 105, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-001704 | codetables-gf3-181-12

Minimum distance of a [181, 12] linear code over GF(3)

codes are known with d = 104, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-001705 | codetables-gf3-181-7

Minimum distance of a [181, 7] linear code over GF(3)

codes are known with d = 116, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-001706 | codetables-gf3-181-8

Minimum distance of a [181, 8] linear code over GF(3)

codes are known with d = 113, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-001707 | codetables-gf3-181-9

Minimum distance of a [181, 9] linear code over GF(3)

codes are known with d = 110, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-001708 | codetables-gf3-182-10

Minimum distance of a [182, 10] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-001709 | codetables-gf3-182-11

Minimum distance of a [182, 11] linear code over GF(3)

codes are known with d = 106, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-001710 | codetables-gf3-182-12

Minimum distance of a [182, 12] linear code over GF(3)

codes are known with d = 105, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-001711 | codetables-gf3-182-7

Minimum distance of a [182, 7] linear code over GF(3)

codes are known with d = 117, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-001712 | codetables-gf3-182-8

Minimum distance of a [182, 8] linear code over GF(3)

codes are known with d = 114, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-001713 | codetables-gf3-182-9

Minimum distance of a [182, 9] linear code over GF(3)

codes are known with d = 111, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-001714 | codetables-gf3-183-10

Minimum distance of a [183, 10] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-001715 | codetables-gf3-183-11

Minimum distance of a [183, 11] linear code over GF(3)

codes are known with d = 107, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-001716 | codetables-gf3-183-12

Minimum distance of a [183, 12] linear code over GF(3)

codes are known with d = 105, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-001717 | codetables-gf3-183-7

Minimum distance of a [183, 7] linear code over GF(3)

codes are known with d = 117, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-001718 | codetables-gf3-183-8

Minimum distance of a [183, 8] linear code over GF(3)

codes are known with d = 114, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-001719 | codetables-gf3-183-9

Minimum distance of a [183, 9] linear code over GF(3)

codes are known with d = 111, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-001720 | codetables-gf3-184-10

Minimum distance of a [184, 10] linear code over GF(3)

codes are known with d = 109, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-001721 | codetables-gf3-184-11

Minimum distance of a [184, 11] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-001722 | codetables-gf3-184-12

Minimum distance of a [184, 12] linear code over GF(3)

codes are known with d = 106, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-001723 | codetables-gf3-184-7

Minimum distance of a [184, 7] linear code over GF(3)

codes are known with d = 117, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-001724 | codetables-gf3-184-8

Minimum distance of a [184, 8] linear code over GF(3)

codes are known with d = 114, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-001725 | codetables-gf3-184-9

Minimum distance of a [184, 9] linear code over GF(3)

codes are known with d = 112, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-001726 | codetables-gf3-185-10

Minimum distance of a [185, 10] linear code over GF(3)

codes are known with d = 110, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-001727 | codetables-gf3-185-11

Minimum distance of a [185, 11] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-001728 | codetables-gf3-185-12

Minimum distance of a [185, 12] linear code over GF(3)

codes are known with d = 106, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-001729 | codetables-gf3-185-7

Minimum distance of a [185, 7] linear code over GF(3)

codes are known with d = 118, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-001730 | codetables-gf3-185-8

Minimum distance of a [185, 8] linear code over GF(3)

codes are known with d = 115, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-001731 | codetables-gf3-185-9

Minimum distance of a [185, 9] linear code over GF(3)

codes are known with d = 113, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-001732 | codetables-gf3-186-10

Minimum distance of a [186, 10] linear code over GF(3)

codes are known with d = 111, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-001733 | codetables-gf3-186-11

Minimum distance of a [186, 11] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-001734 | codetables-gf3-186-12

Minimum distance of a [186, 12] linear code over GF(3)

codes are known with d = 106, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-001735 | codetables-gf3-186-7

Minimum distance of a [186, 7] linear code over GF(3)

codes are known with d = 119, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-001736 | codetables-gf3-186-8

Minimum distance of a [186, 8] linear code over GF(3)

codes are known with d = 116, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-001737 | codetables-gf3-186-9

Minimum distance of a [186, 9] linear code over GF(3)

codes are known with d = 114, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-001738 | codetables-gf3-187-10

Minimum distance of a [187, 10] linear code over GF(3)

codes are known with d = 111, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-001739 | codetables-gf3-187-11

Minimum distance of a [187, 11] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-001740 | codetables-gf3-187-12

Minimum distance of a [187, 12] linear code over GF(3)

codes are known with d = 107, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-001741 | codetables-gf3-187-7

Minimum distance of a [187, 7] linear code over GF(3)

codes are known with d = 120, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-001742 | codetables-gf3-187-8

Minimum distance of a [187, 8] linear code over GF(3)

codes are known with d = 117, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-001743 | codetables-gf3-187-9

Minimum distance of a [187, 9] linear code over GF(3)

codes are known with d = 114, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-001744 | codetables-gf3-188-10

Minimum distance of a [188, 10] linear code over GF(3)

codes are known with d = 112, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-001745 | codetables-gf3-188-11

Minimum distance of a [188, 11] linear code over GF(3)

codes are known with d = 109, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-001746 | codetables-gf3-188-12

Minimum distance of a [188, 12] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-001747 | codetables-gf3-188-7

Minimum distance of a [188, 7] linear code over GF(3)

codes are known with d = 121, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-001748 | codetables-gf3-188-8

Minimum distance of a [188, 8] linear code over GF(3)

codes are known with d = 117, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-001749 | codetables-gf3-188-9

Minimum distance of a [188, 9] linear code over GF(3)

codes are known with d = 114, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-001750 | codetables-gf3-189-10

Minimum distance of a [189, 10] linear code over GF(3)

codes are known with d = 113, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-001751 | codetables-gf3-189-11

Minimum distance of a [189, 11] linear code over GF(3)

codes are known with d = 110, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-001752 | codetables-gf3-189-12

Minimum distance of a [189, 12] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-001753 | codetables-gf3-189-7

Minimum distance of a [189, 7] linear code over GF(3)

codes are known with d = 121, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-001754 | codetables-gf3-189-8

Minimum distance of a [189, 8] linear code over GF(3)

codes are known with d = 117, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-001755 | codetables-gf3-189-9

Minimum distance of a [189, 9] linear code over GF(3)

codes are known with d = 115, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-001756 | codetables-gf3-190-10

Minimum distance of a [190, 10] linear code over GF(3)

codes are known with d = 114, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-001757 | codetables-gf3-190-11

Minimum distance of a [190, 11] linear code over GF(3)

codes are known with d = 111, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-001758 | codetables-gf3-190-12

Minimum distance of a [190, 12] linear code over GF(3)

codes are known with d = 108, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-001759 | codetables-gf3-190-7

Minimum distance of a [190, 7] linear code over GF(3)

codes are known with d = 122, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-001760 | codetables-gf3-190-8

Minimum distance of a [190, 8] linear code over GF(3)

codes are known with d = 118, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-001761 | codetables-gf3-190-9

Minimum distance of a [190, 9] linear code over GF(3)

codes are known with d = 116, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-001762 | codetables-gf3-191-10

Minimum distance of a [191, 10] linear code over GF(3)

codes are known with d = 114, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-001763 | codetables-gf3-191-11

Minimum distance of a [191, 11] linear code over GF(3)

codes are known with d = 111, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-001764 | codetables-gf3-191-12

Minimum distance of a [191, 12] linear code over GF(3)

codes are known with d = 109, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-001765 | codetables-gf3-191-7

Minimum distance of a [191, 7] linear code over GF(3)

codes are known with d = 123, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-001766 | codetables-gf3-191-8

Minimum distance of a [191, 8] linear code over GF(3)

codes are known with d = 119, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-001767 | codetables-gf3-191-9

Minimum distance of a [191, 9] linear code over GF(3)

codes are known with d = 117, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-001768 | codetables-gf3-192-10

Minimum distance of a [192, 10] linear code over GF(3)

codes are known with d = 114, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-001769 | codetables-gf3-192-11

Minimum distance of a [192, 11] linear code over GF(3)

codes are known with d = 112, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-001770 | codetables-gf3-192-12

Minimum distance of a [192, 12] linear code over GF(3)

codes are known with d = 110, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-001771 | codetables-gf3-192-7

Minimum distance of a [192, 7] linear code over GF(3)

codes are known with d = 124, and no code can have d greater than 125.

来源状态 open | 2026-07-23 | 来源

V3P-001772 | codetables-gf3-192-8

Minimum distance of a [192, 8] linear code over GF(3)

codes are known with d = 120, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-001773 | codetables-gf3-192-9

Minimum distance of a [192, 9] linear code over GF(3)

codes are known with d = 117, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-001774 | codetables-gf3-193-10

Minimum distance of a [193, 10] linear code over GF(3)

codes are known with d = 115, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-001775 | codetables-gf3-193-11

Minimum distance of a [193, 11] linear code over GF(3)

codes are known with d = 113, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-001776 | codetables-gf3-193-12

Minimum distance of a [193, 12] linear code over GF(3)

codes are known with d = 111, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-001777 | codetables-gf3-193-7

Minimum distance of a [193, 7] linear code over GF(3)

codes are known with d = 125, and no code can have d greater than 126.

来源状态 open | 2026-07-23 | 来源

V3P-001778 | codetables-gf3-193-8

Minimum distance of a [193, 8] linear code over GF(3)

codes are known with d = 120, and no code can have d greater than 125.

来源状态 open | 2026-07-23 | 来源

V3P-001779 | codetables-gf3-193-9

Minimum distance of a [193, 9] linear code over GF(3)

codes are known with d = 117, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-001780 | codetables-gf3-194-10

Minimum distance of a [194, 10] linear code over GF(3)

codes are known with d = 116, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-001781 | codetables-gf3-194-11

Minimum distance of a [194, 11] linear code over GF(3)

codes are known with d = 114, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-001782 | codetables-gf3-194-12

Minimum distance of a [194, 12] linear code over GF(3)

codes are known with d = 111, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-001783 | codetables-gf3-194-6

Minimum distance of a [194, 6] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 127.

来源状态 open | 2026-07-23 | 来源

V3P-001784 | codetables-gf3-194-8

Minimum distance of a [194, 8] linear code over GF(3)

codes are known with d = 121, and no code can have d greater than 125.

来源状态 open | 2026-07-23 | 来源

V3P-001785 | codetables-gf3-194-9

Minimum distance of a [194, 9] linear code over GF(3)

codes are known with d = 118, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-001786 | codetables-gf3-195-10

Minimum distance of a [195, 10] linear code over GF(3)

codes are known with d = 117, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-001787 | codetables-gf3-195-11

Minimum distance of a [195, 11] linear code over GF(3)

codes are known with d = 114, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-001788 | codetables-gf3-195-12

Minimum distance of a [195, 12] linear code over GF(3)

codes are known with d = 111, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-001789 | codetables-gf3-195-6

Minimum distance of a [195, 6] linear code over GF(3)

codes are known with d = 127, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-001790 | codetables-gf3-195-7

Minimum distance of a [195, 7] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 127.

来源状态 open | 2026-07-23 | 来源

V3P-001791 | codetables-gf3-195-8

Minimum distance of a [195, 8] linear code over GF(3)

codes are known with d = 122, and no code can have d greater than 126.

来源状态 open | 2026-07-23 | 来源

V3P-001792 | codetables-gf3-195-9

Minimum distance of a [195, 9] linear code over GF(3)

codes are known with d = 119, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-001793 | codetables-gf3-196-10

Minimum distance of a [196, 10] linear code over GF(3)

codes are known with d = 117, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-001794 | codetables-gf3-196-11

Minimum distance of a [196, 11] linear code over GF(3)

codes are known with d = 115, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-001795 | codetables-gf3-196-12

Minimum distance of a [196, 12] linear code over GF(3)

codes are known with d = 112, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-001796 | codetables-gf3-196-6

Minimum distance of a [196, 6] linear code over GF(3)

codes are known with d = 128, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-001797 | codetables-gf3-196-7

Minimum distance of a [196, 7] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-001798 | codetables-gf3-196-8

Minimum distance of a [196, 8] linear code over GF(3)

codes are known with d = 123, and no code can have d greater than 126.

来源状态 open | 2026-07-23 | 来源

V3P-001799 | codetables-gf3-196-9

Minimum distance of a [196, 9] linear code over GF(3)

codes are known with d = 120, and no code can have d greater than 125.

来源状态 open | 2026-07-23 | 来源

V3P-001800 | codetables-gf3-197-10

Minimum distance of a [197, 10] linear code over GF(3)

codes are known with d = 117, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-001801 | codetables-gf3-197-11

Minimum distance of a [197, 11] linear code over GF(3)

codes are known with d = 115, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-001802 | codetables-gf3-197-12

Minimum distance of a [197, 12] linear code over GF(3)

codes are known with d = 113, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-001803 | codetables-gf3-197-7

Minimum distance of a [197, 7] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-001804 | codetables-gf3-197-8

Minimum distance of a [197, 8] linear code over GF(3)

codes are known with d = 124, and no code can have d greater than 127.

来源状态 open | 2026-07-23 | 来源

V3P-001805 | codetables-gf3-197-9

Minimum distance of a [197, 9] linear code over GF(3)

codes are known with d = 120, and no code can have d greater than 125.

来源状态 open | 2026-07-23 | 来源

V3P-001806 | codetables-gf3-198-10

Minimum distance of a [198, 10] linear code over GF(3)

codes are known with d = 118, and no code can have d greater than 125.

来源状态 open | 2026-07-23 | 来源

V3P-001807 | codetables-gf3-198-11

Minimum distance of a [198, 11] linear code over GF(3)

codes are known with d = 116, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-001808 | codetables-gf3-198-12

Minimum distance of a [198, 12] linear code over GF(3)

codes are known with d = 114, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-001809 | codetables-gf3-198-6

Minimum distance of a [198, 6] linear code over GF(3)

codes are known with d = 129, and no code can have d greater than 130.

来源状态 open | 2026-07-23 | 来源

V3P-001810 | codetables-gf3-198-7

Minimum distance of a [198, 7] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-001811 | codetables-gf3-198-8

Minimum distance of a [198, 8] linear code over GF(3)

codes are known with d = 125, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-001812 | codetables-gf3-198-9

Minimum distance of a [198, 9] linear code over GF(3)

codes are known with d = 121, and no code can have d greater than 126.

来源状态 open | 2026-07-23 | 来源

V3P-001813 | codetables-gf3-199-10

Minimum distance of a [199, 10] linear code over GF(3)

codes are known with d = 119, and no code can have d greater than 126.

来源状态 open | 2026-07-23 | 来源

V3P-001814 | codetables-gf3-199-11

Minimum distance of a [199, 11] linear code over GF(3)

codes are known with d = 117, and no code can have d greater than 125.

来源状态 open | 2026-07-23 | 来源

V3P-001815 | codetables-gf3-199-12

Minimum distance of a [199, 12] linear code over GF(3)

codes are known with d = 114, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-001816 | codetables-gf3-199-6

Minimum distance of a [199, 6] linear code over GF(3)

codes are known with d = 130, and no code can have d greater than 131.

来源状态 open | 2026-07-23 | 来源

V3P-001817 | codetables-gf3-199-7

Minimum distance of a [199, 7] linear code over GF(3)

codes are known with d = 127, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-001818 | codetables-gf3-199-8

Minimum distance of a [199, 8] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-001819 | codetables-gf3-199-9

Minimum distance of a [199, 9] linear code over GF(3)

codes are known with d = 122, and no code can have d greater than 127.

来源状态 open | 2026-07-23 | 来源

V3P-001820 | codetables-gf3-200-10

Minimum distance of a [200, 10] linear code over GF(3)

codes are known with d = 120, and no code can have d greater than 126.

来源状态 open | 2026-07-23 | 来源

V3P-001821 | codetables-gf3-200-11

Minimum distance of a [200, 11] linear code over GF(3)

codes are known with d = 118, and no code can have d greater than 126.

来源状态 open | 2026-07-23 | 来源

V3P-001822 | codetables-gf3-200-12

Minimum distance of a [200, 12] linear code over GF(3)

codes are known with d = 115, and no code can have d greater than 125.

来源状态 open | 2026-07-23 | 来源

V3P-001823 | codetables-gf3-200-6

Minimum distance of a [200, 6] linear code over GF(3)

codes are known with d = 131, and no code can have d greater than 132.

来源状态 open | 2026-07-23 | 来源

V3P-001824 | codetables-gf3-200-7

Minimum distance of a [200, 7] linear code over GF(3)

codes are known with d = 128, and no code can have d greater than 130.

来源状态 open | 2026-07-23 | 来源

V3P-001825 | codetables-gf3-200-8

Minimum distance of a [200, 8] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-001826 | codetables-gf3-200-9

Minimum distance of a [200, 9] linear code over GF(3)

codes are known with d = 123, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-001827 | codetables-gf3-201-10

Minimum distance of a [201, 10] linear code over GF(3)

codes are known with d = 120, and no code can have d greater than 127.

来源状态 open | 2026-07-23 | 来源

V3P-001828 | codetables-gf3-201-11

Minimum distance of a [201, 11] linear code over GF(3)

codes are known with d = 118, and no code can have d greater than 126.

来源状态 open | 2026-07-23 | 来源

V3P-001829 | codetables-gf3-201-12

Minimum distance of a [201, 12] linear code over GF(3)

codes are known with d = 116, and no code can have d greater than 125.

来源状态 open | 2026-07-23 | 来源

V3P-001830 | codetables-gf3-201-7

Minimum distance of a [201, 7] linear code over GF(3)

codes are known with d = 129, and no code can have d greater than 131.

来源状态 open | 2026-07-23 | 来源

V3P-001831 | codetables-gf3-201-8

Minimum distance of a [201, 8] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 130.

来源状态 open | 2026-07-23 | 来源

V3P-001832 | codetables-gf3-201-9

Minimum distance of a [201, 9] linear code over GF(3)

codes are known with d = 123, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-001833 | codetables-gf3-202-10

Minimum distance of a [202, 10] linear code over GF(3)

codes are known with d = 121, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-001834 | codetables-gf3-202-11

Minimum distance of a [202, 11] linear code over GF(3)

codes are known with d = 119, and no code can have d greater than 127.

来源状态 open | 2026-07-23 | 来源

V3P-001835 | codetables-gf3-202-12

Minimum distance of a [202, 12] linear code over GF(3)

codes are known with d = 117, and no code can have d greater than 126.

来源状态 open | 2026-07-23 | 来源

V3P-001836 | codetables-gf3-202-6

Minimum distance of a [202, 6] linear code over GF(3)

codes are known with d = 132, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-001837 | codetables-gf3-202-7

Minimum distance of a [202, 7] linear code over GF(3)

codes are known with d = 129, and no code can have d greater than 132.

来源状态 open | 2026-07-23 | 来源

V3P-001838 | codetables-gf3-202-8

Minimum distance of a [202, 8] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 131.

来源状态 open | 2026-07-23 | 来源

V3P-001839 | codetables-gf3-202-9

Minimum distance of a [202, 9] linear code over GF(3)

codes are known with d = 123, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-001840 | codetables-gf3-203-10

Minimum distance of a [203, 10] linear code over GF(3)

codes are known with d = 122, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-001841 | codetables-gf3-203-11

Minimum distance of a [203, 11] linear code over GF(3)

codes are known with d = 120, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-001842 | codetables-gf3-203-12

Minimum distance of a [203, 12] linear code over GF(3)

codes are known with d = 117, and no code can have d greater than 127.

来源状态 open | 2026-07-23 | 来源

V3P-001843 | codetables-gf3-203-7

Minimum distance of a [203, 7] linear code over GF(3)

codes are known with d = 130, and no code can have d greater than 132.

来源状态 open | 2026-07-23 | 来源

V3P-001844 | codetables-gf3-203-8

Minimum distance of a [203, 8] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 131.

来源状态 open | 2026-07-23 | 来源

V3P-001845 | codetables-gf3-203-9

Minimum distance of a [203, 9] linear code over GF(3)

codes are known with d = 124, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-001846 | codetables-gf3-204-10

Minimum distance of a [204, 10] linear code over GF(3)

codes are known with d = 123, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-001847 | codetables-gf3-204-11

Minimum distance of a [204, 11] linear code over GF(3)

codes are known with d = 120, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-001848 | codetables-gf3-204-12

Minimum distance of a [204, 12] linear code over GF(3)

codes are known with d = 117, and no code can have d greater than 127.

来源状态 open | 2026-07-23 | 来源

V3P-001849 | codetables-gf3-204-7

Minimum distance of a [204, 7] linear code over GF(3)

codes are known with d = 131, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-001850 | codetables-gf3-204-8

Minimum distance of a [204, 8] linear code over GF(3)

codes are known with d = 127, and no code can have d greater than 132.

来源状态 open | 2026-07-23 | 来源

V3P-001851 | codetables-gf3-204-9

Minimum distance of a [204, 9] linear code over GF(3)

codes are known with d = 125, and no code can have d greater than 130.

来源状态 open | 2026-07-23 | 来源

V3P-001852 | codetables-gf3-205-10

Minimum distance of a [205, 10] linear code over GF(3)

codes are known with d = 123, and no code can have d greater than 130.

来源状态 open | 2026-07-23 | 来源

V3P-001853 | codetables-gf3-205-11

Minimum distance of a [205, 11] linear code over GF(3)

codes are known with d = 121, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-001854 | codetables-gf3-205-12

Minimum distance of a [205, 12] linear code over GF(3)

codes are known with d = 118, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-001855 | codetables-gf3-205-7

Minimum distance of a [205, 7] linear code over GF(3)

codes are known with d = 132, and no code can have d greater than 134.

来源状态 open | 2026-07-23 | 来源

V3P-001856 | codetables-gf3-205-8

Minimum distance of a [205, 8] linear code over GF(3)

codes are known with d = 128, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-001857 | codetables-gf3-205-9

Minimum distance of a [205, 9] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 131.

来源状态 open | 2026-07-23 | 来源

V3P-001858 | codetables-gf3-206-10

Minimum distance of a [206, 10] linear code over GF(3)

codes are known with d = 124, and no code can have d greater than 131.

来源状态 open | 2026-07-23 | 来源

V3P-001859 | codetables-gf3-206-11

Minimum distance of a [206, 11] linear code over GF(3)

codes are known with d = 122, and no code can have d greater than 130.

来源状态 open | 2026-07-23 | 来源

V3P-001860 | codetables-gf3-206-12

Minimum distance of a [206, 12] linear code over GF(3)

codes are known with d = 119, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-001861 | codetables-gf3-206-7

Minimum distance of a [206, 7] linear code over GF(3)

codes are known with d = 132, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-001862 | codetables-gf3-206-8

Minimum distance of a [206, 8] linear code over GF(3)

codes are known with d = 129, and no code can have d greater than 134.

来源状态 open | 2026-07-23 | 来源

V3P-001863 | codetables-gf3-206-9

Minimum distance of a [206, 9] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 131.

来源状态 open | 2026-07-23 | 来源

V3P-001864 | codetables-gf3-207-10

Minimum distance of a [207, 10] linear code over GF(3)

codes are known with d = 125, and no code can have d greater than 131.

来源状态 open | 2026-07-23 | 来源

V3P-001865 | codetables-gf3-207-11

Minimum distance of a [207, 11] linear code over GF(3)

codes are known with d = 123, and no code can have d greater than 131.

来源状态 open | 2026-07-23 | 来源

V3P-001866 | codetables-gf3-207-12

Minimum distance of a [207, 12] linear code over GF(3)

codes are known with d = 120, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-001867 | codetables-gf3-207-7

Minimum distance of a [207, 7] linear code over GF(3)

codes are known with d = 133, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-001868 | codetables-gf3-207-8

Minimum distance of a [207, 8] linear code over GF(3)

codes are known with d = 129, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-001869 | codetables-gf3-207-9

Minimum distance of a [207, 9] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 132.

来源状态 open | 2026-07-23 | 来源

V3P-001870 | codetables-gf3-208-10

Minimum distance of a [208, 10] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 132.

来源状态 open | 2026-07-23 | 来源

V3P-001871 | codetables-gf3-208-11

Minimum distance of a [208, 11] linear code over GF(3)

codes are known with d = 123, and no code can have d greater than 131.

来源状态 open | 2026-07-23 | 来源

V3P-001872 | codetables-gf3-208-12

Minimum distance of a [208, 12] linear code over GF(3)

codes are known with d = 120, and no code can have d greater than 130.

来源状态 open | 2026-07-23 | 来源

V3P-001873 | codetables-gf3-208-6

Minimum distance of a [208, 6] linear code over GF(3)

codes are known with d = 135, and no code can have d greater than 136.

来源状态 open | 2026-07-23 | 来源

V3P-001874 | codetables-gf3-208-7

Minimum distance of a [208, 7] linear code over GF(3)

codes are known with d = 134, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-001875 | codetables-gf3-208-8

Minimum distance of a [208, 8] linear code over GF(3)

codes are known with d = 130, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-001876 | codetables-gf3-208-9

Minimum distance of a [208, 9] linear code over GF(3)

codes are known with d = 127, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-001877 | codetables-gf3-209-10

Minimum distance of a [209, 10] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-001878 | codetables-gf3-209-11

Minimum distance of a [209, 11] linear code over GF(3)

codes are known with d = 124, and no code can have d greater than 132.

来源状态 open | 2026-07-23 | 来源

V3P-001879 | codetables-gf3-209-12

Minimum distance of a [209, 12] linear code over GF(3)

codes are known with d = 120, and no code can have d greater than 131.

来源状态 open | 2026-07-23 | 来源

V3P-001880 | codetables-gf3-209-6

Minimum distance of a [209, 6] linear code over GF(3)

codes are known with d = 136, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-001881 | codetables-gf3-209-7

Minimum distance of a [209, 7] linear code over GF(3)

codes are known with d = 135, and no code can have d greater than 136.

来源状态 open | 2026-07-23 | 来源

V3P-001882 | codetables-gf3-209-8

Minimum distance of a [209, 8] linear code over GF(3)

codes are known with d = 131, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-001883 | codetables-gf3-209-9

Minimum distance of a [209, 9] linear code over GF(3)

codes are known with d = 128, and no code can have d greater than 134.

来源状态 open | 2026-07-23 | 来源

V3P-001884 | codetables-gf3-210-10

Minimum distance of a [210, 10] linear code over GF(3)

codes are known with d = 127, and no code can have d greater than 134.

来源状态 open | 2026-07-23 | 来源

V3P-001885 | codetables-gf3-210-11

Minimum distance of a [210, 11] linear code over GF(3)

codes are known with d = 125, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-001886 | codetables-gf3-210-12

Minimum distance of a [210, 12] linear code over GF(3)

codes are known with d = 121, and no code can have d greater than 132.

来源状态 open | 2026-07-23 | 来源

V3P-001887 | codetables-gf3-210-6

Minimum distance of a [210, 6] linear code over GF(3)

codes are known with d = 137, and no code can have d greater than 138.

来源状态 open | 2026-07-23 | 来源

V3P-001888 | codetables-gf3-210-7

Minimum distance of a [210, 7] linear code over GF(3)

codes are known with d = 135, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-001889 | codetables-gf3-210-8

Minimum distance of a [210, 8] linear code over GF(3)

codes are known with d = 132, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-001890 | codetables-gf3-210-9

Minimum distance of a [210, 9] linear code over GF(3)

codes are known with d = 129, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-001891 | codetables-gf3-211-10

Minimum distance of a [211, 10] linear code over GF(3)

codes are known with d = 127, and no code can have d greater than 134.

来源状态 open | 2026-07-23 | 来源

V3P-001892 | codetables-gf3-211-11

Minimum distance of a [211, 11] linear code over GF(3)

codes are known with d = 125, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-001893 | codetables-gf3-211-12

Minimum distance of a [211, 12] linear code over GF(3)

codes are known with d = 122, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-001894 | codetables-gf3-211-7

Minimum distance of a [211, 7] linear code over GF(3)

codes are known with d = 135, and no code can have d greater than 138.

来源状态 open | 2026-07-23 | 来源

V3P-001895 | codetables-gf3-211-8

Minimum distance of a [211, 8] linear code over GF(3)

codes are known with d = 133, and no code can have d greater than 136.

来源状态 open | 2026-07-23 | 来源

V3P-001896 | codetables-gf3-211-9

Minimum distance of a [211, 9] linear code over GF(3)

codes are known with d = 129, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-001897 | codetables-gf3-212-10

Minimum distance of a [212, 10] linear code over GF(3)

codes are known with d = 128, and no code can have d greater than 134.

来源状态 open | 2026-07-23 | 来源

V3P-001898 | codetables-gf3-212-11

Minimum distance of a [212, 11] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-001899 | codetables-gf3-212-12

Minimum distance of a [212, 12] linear code over GF(3)

codes are known with d = 123, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-001900 | codetables-gf3-212-6

Minimum distance of a [212, 6] linear code over GF(3)

codes are known with d = 138, and no code can have d greater than 139.

来源状态 open | 2026-07-23 | 来源

V3P-001901 | codetables-gf3-212-7

Minimum distance of a [212, 7] linear code over GF(3)

codes are known with d = 136, and no code can have d greater than 138.

来源状态 open | 2026-07-23 | 来源

V3P-001902 | codetables-gf3-212-8

Minimum distance of a [212, 8] linear code over GF(3)

codes are known with d = 134, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-001903 | codetables-gf3-212-9

Minimum distance of a [212, 9] linear code over GF(3)

codes are known with d = 130, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-001904 | codetables-gf3-213-10

Minimum distance of a [213, 10] linear code over GF(3)

codes are known with d = 129, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-001905 | codetables-gf3-213-11

Minimum distance of a [213, 11] linear code over GF(3)

codes are known with d = 127, and no code can have d greater than 134.

来源状态 open | 2026-07-23 | 来源

V3P-001906 | codetables-gf3-213-12

Minimum distance of a [213, 12] linear code over GF(3)

codes are known with d = 123, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-001907 | codetables-gf3-213-6

Minimum distance of a [213, 6] linear code over GF(3)

codes are known with d = 139, and no code can have d greater than 140.

来源状态 open | 2026-07-23 | 来源

V3P-001908 | codetables-gf3-213-7

Minimum distance of a [213, 7] linear code over GF(3)

codes are known with d = 137, and no code can have d greater than 139.

来源状态 open | 2026-07-23 | 来源

V3P-001909 | codetables-gf3-213-8

Minimum distance of a [213, 8] linear code over GF(3)

codes are known with d = 135, and no code can have d greater than 138.

来源状态 open | 2026-07-23 | 来源

V3P-001910 | codetables-gf3-213-9

Minimum distance of a [213, 9] linear code over GF(3)

codes are known with d = 131, and no code can have d greater than 136.

来源状态 open | 2026-07-23 | 来源

V3P-001911 | codetables-gf3-214-10

Minimum distance of a [214, 10] linear code over GF(3)

codes are known with d = 129, and no code can have d greater than 136.

来源状态 open | 2026-07-23 | 来源

V3P-001912 | codetables-gf3-214-11

Minimum distance of a [214, 11] linear code over GF(3)

codes are known with d = 128, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-001913 | codetables-gf3-214-12

Minimum distance of a [214, 12] linear code over GF(3)

codes are known with d = 124, and no code can have d greater than 134.

来源状态 open | 2026-07-23 | 来源

V3P-001914 | codetables-gf3-214-6

Minimum distance of a [214, 6] linear code over GF(3)

codes are known with d = 140, and no code can have d greater than 141.

来源状态 open | 2026-07-23 | 来源

V3P-001915 | codetables-gf3-214-7

Minimum distance of a [214, 7] linear code over GF(3)

codes are known with d = 138, and no code can have d greater than 140.

来源状态 open | 2026-07-23 | 来源

V3P-001916 | codetables-gf3-214-8

Minimum distance of a [214, 8] linear code over GF(3)

codes are known with d = 135, and no code can have d greater than 138.

来源状态 open | 2026-07-23 | 来源

V3P-001917 | codetables-gf3-214-9

Minimum distance of a [214, 9] linear code over GF(3)

codes are known with d = 132, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-001918 | codetables-gf3-215-10

Minimum distance of a [215, 10] linear code over GF(3)

codes are known with d = 130, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-001919 | codetables-gf3-215-11

Minimum distance of a [215, 11] linear code over GF(3)

codes are known with d = 128, and no code can have d greater than 136.

来源状态 open | 2026-07-23 | 来源

V3P-001920 | codetables-gf3-215-12

Minimum distance of a [215, 12] linear code over GF(3)

codes are known with d = 125, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-001921 | codetables-gf3-215-7

Minimum distance of a [215, 7] linear code over GF(3)

codes are known with d = 138, and no code can have d greater than 141.

来源状态 open | 2026-07-23 | 来源

V3P-001922 | codetables-gf3-215-8

Minimum distance of a [215, 8] linear code over GF(3)

codes are known with d = 135, and no code can have d greater than 139.

来源状态 open | 2026-07-23 | 来源

V3P-001923 | codetables-gf3-215-9

Minimum distance of a [215, 9] linear code over GF(3)

codes are known with d = 132, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-001924 | codetables-gf3-216-10

Minimum distance of a [216, 10] linear code over GF(3)

codes are known with d = 131, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-001925 | codetables-gf3-216-11

Minimum distance of a [216, 11] linear code over GF(3)

codes are known with d = 129, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-001926 | codetables-gf3-216-12

Minimum distance of a [216, 12] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 136.

来源状态 open | 2026-07-23 | 来源

V3P-001927 | codetables-gf3-216-6

Minimum distance of a [216, 6] linear code over GF(3)

codes are known with d = 141, and no code can have d greater than 142.

来源状态 open | 2026-07-23 | 来源

V3P-001928 | codetables-gf3-216-7

Minimum distance of a [216, 7] linear code over GF(3)

codes are known with d = 139, and no code can have d greater than 141.

来源状态 open | 2026-07-23 | 来源

V3P-001929 | codetables-gf3-216-8

Minimum distance of a [216, 8] linear code over GF(3)

codes are known with d = 136, and no code can have d greater than 140.

来源状态 open | 2026-07-23 | 来源

V3P-001930 | codetables-gf3-216-9

Minimum distance of a [216, 9] linear code over GF(3)

codes are known with d = 132, and no code can have d greater than 138.

来源状态 open | 2026-07-23 | 来源

V3P-001931 | codetables-gf3-217-10

Minimum distance of a [217, 10] linear code over GF(3)

codes are known with d = 132, and no code can have d greater than 138.

来源状态 open | 2026-07-23 | 来源

V3P-001932 | codetables-gf3-217-11

Minimum distance of a [217, 11] linear code over GF(3)

codes are known with d = 130, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-001933 | codetables-gf3-217-12

Minimum distance of a [217, 12] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-001934 | codetables-gf3-217-6

Minimum distance of a [217, 6] linear code over GF(3)

codes are known with d = 142, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-001935 | codetables-gf3-217-7

Minimum distance of a [217, 7] linear code over GF(3)

codes are known with d = 140, and no code can have d greater than 142.

来源状态 open | 2026-07-23 | 来源

V3P-001936 | codetables-gf3-217-8

Minimum distance of a [217, 8] linear code over GF(3)

codes are known with d = 137, and no code can have d greater than 141.

来源状态 open | 2026-07-23 | 来源

V3P-001937 | codetables-gf3-217-9

Minimum distance of a [217, 9] linear code over GF(3)

codes are known with d = 133, and no code can have d greater than 139.

来源状态 open | 2026-07-23 | 来源

V3P-001938 | codetables-gf3-218-10

Minimum distance of a [218, 10] linear code over GF(3)

codes are known with d = 133, and no code can have d greater than 139.

来源状态 open | 2026-07-23 | 来源

V3P-001939 | codetables-gf3-218-11

Minimum distance of a [218, 11] linear code over GF(3)

codes are known with d = 131, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-001940 | codetables-gf3-218-12

Minimum distance of a [218, 12] linear code over GF(3)

codes are known with d = 126, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-001941 | codetables-gf3-218-6

Minimum distance of a [218, 6] linear code over GF(3)

codes are known with d = 143, and no code can have d greater than 144.

来源状态 open | 2026-07-23 | 来源

V3P-001942 | codetables-gf3-218-7

Minimum distance of a [218, 7] linear code over GF(3)

codes are known with d = 141, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-001943 | codetables-gf3-218-8

Minimum distance of a [218, 8] linear code over GF(3)

codes are known with d = 138, and no code can have d greater than 141.

来源状态 open | 2026-07-23 | 来源

V3P-001944 | codetables-gf3-218-9

Minimum distance of a [218, 9] linear code over GF(3)

codes are known with d = 134, and no code can have d greater than 140.

来源状态 open | 2026-07-23 | 来源

V3P-001945 | codetables-gf3-219-10

Minimum distance of a [219, 10] linear code over GF(3)

codes are known with d = 134, and no code can have d greater than 140.

来源状态 open | 2026-07-23 | 来源

V3P-001946 | codetables-gf3-219-11

Minimum distance of a [219, 11] linear code over GF(3)

codes are known with d = 131, and no code can have d greater than 138.

来源状态 open | 2026-07-23 | 来源

V3P-001947 | codetables-gf3-219-12

Minimum distance of a [219, 12] linear code over GF(3)

codes are known with d = 127, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-001948 | codetables-gf3-219-7

Minimum distance of a [219, 7] linear code over GF(3)

codes are known with d = 141, and no code can have d greater than 144.

来源状态 open | 2026-07-23 | 来源

V3P-001949 | codetables-gf3-219-8

Minimum distance of a [219, 8] linear code over GF(3)

codes are known with d = 138, and no code can have d greater than 142.

来源状态 open | 2026-07-23 | 来源

V3P-001950 | codetables-gf3-219-9

Minimum distance of a [219, 9] linear code over GF(3)

codes are known with d = 135, and no code can have d greater than 141.

来源状态 open | 2026-07-23 | 来源

V3P-001951 | codetables-gf3-22-8

Minimum distance of a [22, 8] linear code over GF(3)

codes are known with d = 9, and no code can have d greater than 10.

来源状态 open | 2026-07-23 | 来源

V3P-001952 | codetables-gf3-220-10

Minimum distance of a [220, 10] linear code over GF(3)

codes are known with d = 135, and no code can have d greater than 140.

来源状态 open | 2026-07-23 | 来源

V3P-001953 | codetables-gf3-220-11

Minimum distance of a [220, 11] linear code over GF(3)

codes are known with d = 132, and no code can have d greater than 139.

来源状态 open | 2026-07-23 | 来源

V3P-001954 | codetables-gf3-220-12

Minimum distance of a [220, 12] linear code over GF(3)

codes are known with d = 128, and no code can have d greater than 138.

来源状态 open | 2026-07-23 | 来源

V3P-001955 | codetables-gf3-220-7

Minimum distance of a [220, 7] linear code over GF(3)

codes are known with d = 142, and no code can have d greater than 144.

来源状态 open | 2026-07-23 | 来源

V3P-001956 | codetables-gf3-220-8

Minimum distance of a [220, 8] linear code over GF(3)

codes are known with d = 139, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-001957 | codetables-gf3-220-9

Minimum distance of a [220, 9] linear code over GF(3)

codes are known with d = 136, and no code can have d greater than 141.

来源状态 open | 2026-07-23 | 来源

V3P-001958 | codetables-gf3-221-10

Minimum distance of a [221, 10] linear code over GF(3)

codes are known with d = 135, and no code can have d greater than 140.

来源状态 open | 2026-07-23 | 来源

V3P-001959 | codetables-gf3-221-11

Minimum distance of a [221, 11] linear code over GF(3)

codes are known with d = 133, and no code can have d greater than 140.

来源状态 open | 2026-07-23 | 来源

V3P-001960 | codetables-gf3-221-12

Minimum distance of a [221, 12] linear code over GF(3)

codes are known with d = 129, and no code can have d greater than 139.

来源状态 open | 2026-07-23 | 来源

V3P-001961 | codetables-gf3-221-6

Minimum distance of a [221, 6] linear code over GF(3)

codes are known with d = 144, and no code can have d greater than 145.

来源状态 open | 2026-07-23 | 来源

V3P-001962 | codetables-gf3-221-7

Minimum distance of a [221, 7] linear code over GF(3)

codes are known with d = 143, and no code can have d greater than 144.

来源状态 open | 2026-07-23 | 来源

V3P-001963 | codetables-gf3-221-8

Minimum distance of a [221, 8] linear code over GF(3)

codes are known with d = 140, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-001964 | codetables-gf3-221-9

Minimum distance of a [221, 9] linear code over GF(3)

codes are known with d = 137, and no code can have d greater than 141.

来源状态 open | 2026-07-23 | 来源

V3P-001965 | codetables-gf3-222-10

Minimum distance of a [222, 10] linear code over GF(3)

codes are known with d = 135, and no code can have d greater than 141.

来源状态 open | 2026-07-23 | 来源

V3P-001966 | codetables-gf3-222-11

Minimum distance of a [222, 11] linear code over GF(3)

codes are known with d = 134, and no code can have d greater than 140.

来源状态 open | 2026-07-23 | 来源

V3P-001967 | codetables-gf3-222-12

Minimum distance of a [222, 12] linear code over GF(3)

codes are known with d = 129, and no code can have d greater than 140.

来源状态 open | 2026-07-23 | 来源

V3P-001968 | codetables-gf3-222-6

Minimum distance of a [222, 6] linear code over GF(3)

codes are known with d = 145, and no code can have d greater than 146.

来源状态 open | 2026-07-23 | 来源

V3P-001969 | codetables-gf3-222-7

Minimum distance of a [222, 7] linear code over GF(3)

codes are known with d = 144, and no code can have d greater than 145.

来源状态 open | 2026-07-23 | 来源

V3P-001970 | codetables-gf3-222-8

Minimum distance of a [222, 8] linear code over GF(3)

codes are known with d = 141, and no code can have d greater than 144.

来源状态 open | 2026-07-23 | 来源

V3P-001971 | codetables-gf3-222-9

Minimum distance of a [222, 9] linear code over GF(3)

codes are known with d = 138, and no code can have d greater than 142.

来源状态 open | 2026-07-23 | 来源

V3P-001972 | codetables-gf3-223-10

Minimum distance of a [223, 10] linear code over GF(3)

codes are known with d = 136, and no code can have d greater than 142.

来源状态 open | 2026-07-23 | 来源

V3P-001973 | codetables-gf3-223-11

Minimum distance of a [223, 11] linear code over GF(3)

codes are known with d = 134, and no code can have d greater than 141.

来源状态 open | 2026-07-23 | 来源

V3P-001974 | codetables-gf3-223-12

Minimum distance of a [223, 12] linear code over GF(3)

codes are known with d = 130, and no code can have d greater than 140.

来源状态 open | 2026-07-23 | 来源

V3P-001975 | codetables-gf3-223-6

Minimum distance of a [223, 6] linear code over GF(3)

codes are known with d = 146, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-001976 | codetables-gf3-223-7

Minimum distance of a [223, 7] linear code over GF(3)

codes are known with d = 144, and no code can have d greater than 146.

来源状态 open | 2026-07-23 | 来源

V3P-001977 | codetables-gf3-223-8

Minimum distance of a [223, 8] linear code over GF(3)

codes are known with d = 142, and no code can have d greater than 144.

来源状态 open | 2026-07-23 | 来源

V3P-001978 | codetables-gf3-223-9

Minimum distance of a [223, 9] linear code over GF(3)

codes are known with d = 139, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-001979 | codetables-gf3-224-10

Minimum distance of a [224, 10] linear code over GF(3)

codes are known with d = 137, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-001980 | codetables-gf3-224-11

Minimum distance of a [224, 11] linear code over GF(3)

codes are known with d = 135, and no code can have d greater than 142.

来源状态 open | 2026-07-23 | 来源

V3P-001981 | codetables-gf3-224-12

Minimum distance of a [224, 12] linear code over GF(3)

codes are known with d = 131, and no code can have d greater than 141.

来源状态 open | 2026-07-23 | 来源

V3P-001982 | codetables-gf3-224-7

Minimum distance of a [224, 7] linear code over GF(3)

codes are known with d = 144, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-001983 | codetables-gf3-224-8

Minimum distance of a [224, 8] linear code over GF(3)

codes are known with d = 143, and no code can have d greater than 145.

来源状态 open | 2026-07-23 | 来源

V3P-001984 | codetables-gf3-224-9

Minimum distance of a [224, 9] linear code over GF(3)

codes are known with d = 140, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-001985 | codetables-gf3-225-10

Minimum distance of a [225, 10] linear code over GF(3)

codes are known with d = 137, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-001986 | codetables-gf3-225-11

Minimum distance of a [225, 11] linear code over GF(3)

codes are known with d = 136, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-001987 | codetables-gf3-225-12

Minimum distance of a [225, 12] linear code over GF(3)

codes are known with d = 132, and no code can have d greater than 142.

来源状态 open | 2026-07-23 | 来源

V3P-001988 | codetables-gf3-225-6

Minimum distance of a [225, 6] linear code over GF(3)

codes are known with d = 147, and no code can have d greater than 148.

来源状态 open | 2026-07-23 | 来源

V3P-001989 | codetables-gf3-225-7

Minimum distance of a [225, 7] linear code over GF(3)

codes are known with d = 144, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-001990 | codetables-gf3-225-8

Minimum distance of a [225, 8] linear code over GF(3)

codes are known with d = 144, and no code can have d greater than 146.

来源状态 open | 2026-07-23 | 来源

V3P-001991 | codetables-gf3-225-9

Minimum distance of a [225, 9] linear code over GF(3)

codes are known with d = 141, and no code can have d greater than 144.

来源状态 open | 2026-07-23 | 来源

V3P-001992 | codetables-gf3-226-10

Minimum distance of a [226, 10] linear code over GF(3)

codes are known with d = 138, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-001993 | codetables-gf3-226-11

Minimum distance of a [226, 11] linear code over GF(3)

codes are known with d = 137, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-001994 | codetables-gf3-226-12

Minimum distance of a [226, 12] linear code over GF(3)

codes are known with d = 132, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-001995 | codetables-gf3-226-6

Minimum distance of a [226, 6] linear code over GF(3)

codes are known with d = 148, and no code can have d greater than 149.

来源状态 open | 2026-07-23 | 来源

V3P-001996 | codetables-gf3-226-7

Minimum distance of a [226, 7] linear code over GF(3)

codes are known with d = 145, and no code can have d greater than 148.

来源状态 open | 2026-07-23 | 来源

V3P-001997 | codetables-gf3-226-8

Minimum distance of a [226, 8] linear code over GF(3)

codes are known with d = 144, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-001998 | codetables-gf3-226-9

Minimum distance of a [226, 9] linear code over GF(3)

codes are known with d = 141, and no code can have d greater than 145.

来源状态 open | 2026-07-23 | 来源

V3P-001999 | codetables-gf3-227-10

Minimum distance of a [227, 10] linear code over GF(3)

codes are known with d = 139, and no code can have d greater than 144.

来源状态 open | 2026-07-23 | 来源

V3P-002000 | codetables-gf3-227-11

Minimum distance of a [227, 11] linear code over GF(3)

codes are known with d = 138, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-002001 | codetables-gf3-227-12

Minimum distance of a [227, 12] linear code over GF(3)

codes are known with d = 132, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-002002 | codetables-gf3-227-6

Minimum distance of a [227, 6] linear code over GF(3)

codes are known with d = 149, and no code can have d greater than 150.

来源状态 open | 2026-07-23 | 来源

V3P-002003 | codetables-gf3-227-7

Minimum distance of a [227, 7] linear code over GF(3)

codes are known with d = 146, and no code can have d greater than 149.

来源状态 open | 2026-07-23 | 来源

V3P-002004 | codetables-gf3-227-8

Minimum distance of a [227, 8] linear code over GF(3)

codes are known with d = 144, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-002005 | codetables-gf3-227-9

Minimum distance of a [227, 9] linear code over GF(3)

codes are known with d = 142, and no code can have d greater than 146.

来源状态 open | 2026-07-23 | 来源

V3P-002006 | codetables-gf3-228-10

Minimum distance of a [228, 10] linear code over GF(3)

codes are known with d = 140, and no code can have d greater than 145.

来源状态 open | 2026-07-23 | 来源

V3P-002007 | codetables-gf3-228-11

Minimum distance of a [228, 11] linear code over GF(3)

codes are known with d = 139, and no code can have d greater than 144.

来源状态 open | 2026-07-23 | 来源

V3P-002008 | codetables-gf3-228-12

Minimum distance of a [228, 12] linear code over GF(3)

codes are known with d = 133, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-002009 | codetables-gf3-228-7

Minimum distance of a [228, 7] linear code over GF(3)

codes are known with d = 147, and no code can have d greater than 150.

来源状态 open | 2026-07-23 | 来源

V3P-002010 | codetables-gf3-228-8

Minimum distance of a [228, 8] linear code over GF(3)

codes are known with d = 144, and no code can have d greater than 148.

来源状态 open | 2026-07-23 | 来源

V3P-002011 | codetables-gf3-228-9

Minimum distance of a [228, 9] linear code over GF(3)

codes are known with d = 143, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-002012 | codetables-gf3-229-10

Minimum distance of a [229, 10] linear code over GF(3)

codes are known with d = 141, and no code can have d greater than 145.

来源状态 open | 2026-07-23 | 来源

V3P-002013 | codetables-gf3-229-11

Minimum distance of a [229, 11] linear code over GF(3)

codes are known with d = 139, and no code can have d greater than 145.

来源状态 open | 2026-07-23 | 来源

V3P-002014 | codetables-gf3-229-12

Minimum distance of a [229, 12] linear code over GF(3)

codes are known with d = 134, and no code can have d greater than 144.

来源状态 open | 2026-07-23 | 来源

V3P-002015 | codetables-gf3-229-6

Minimum distance of a [229, 6] linear code over GF(3)

codes are known with d = 150, and no code can have d greater than 151.

来源状态 open | 2026-07-23 | 来源

V3P-002016 | codetables-gf3-229-7

Minimum distance of a [229, 7] linear code over GF(3)

codes are known with d = 147, and no code can have d greater than 150.

来源状态 open | 2026-07-23 | 来源

V3P-002017 | codetables-gf3-229-8

Minimum distance of a [229, 8] linear code over GF(3)

codes are known with d = 144, and no code can have d greater than 149.

来源状态 open | 2026-07-23 | 来源

V3P-002018 | codetables-gf3-229-9

Minimum distance of a [229, 9] linear code over GF(3)

codes are known with d = 144, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-002019 | codetables-gf3-23-8

Minimum distance of a [23, 8] linear code over GF(3)

codes are known with d = 10, and no code can have d greater than 11.

来源状态 open | 2026-07-23 | 来源

V3P-002020 | codetables-gf3-23-9

Minimum distance of a [23, 9] linear code over GF(3)

codes are known with d = 9, and no code can have d greater than 10.

来源状态 open | 2026-07-23 | 来源

V3P-002021 | codetables-gf3-230-10

Minimum distance of a [230, 10] linear code over GF(3)

codes are known with d = 141, and no code can have d greater than 146.

来源状态 open | 2026-07-23 | 来源

V3P-002022 | codetables-gf3-230-11

Minimum distance of a [230, 11] linear code over GF(3)

codes are known with d = 140, and no code can have d greater than 145.

来源状态 open | 2026-07-23 | 来源

V3P-002023 | codetables-gf3-230-12

Minimum distance of a [230, 12] linear code over GF(3)

codes are known with d = 134, and no code can have d greater than 145.

来源状态 open | 2026-07-23 | 来源

V3P-002024 | codetables-gf3-230-7

Minimum distance of a [230, 7] linear code over GF(3)

codes are known with d = 148, and no code can have d greater than 151.

来源状态 open | 2026-07-23 | 来源

V3P-002025 | codetables-gf3-230-8

Minimum distance of a [230, 8] linear code over GF(3)

codes are known with d = 145, and no code can have d greater than 150.

来源状态 open | 2026-07-23 | 来源

V3P-002026 | codetables-gf3-230-9

Minimum distance of a [230, 9] linear code over GF(3)

codes are known with d = 144, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-002027 | codetables-gf3-231-10

Minimum distance of a [231, 10] linear code over GF(3)

codes are known with d = 142, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-002028 | codetables-gf3-231-11

Minimum distance of a [231, 11] linear code over GF(3)

codes are known with d = 141, and no code can have d greater than 146.

来源状态 open | 2026-07-23 | 来源

V3P-002029 | codetables-gf3-231-12

Minimum distance of a [231, 12] linear code over GF(3)

codes are known with d = 135, and no code can have d greater than 145.

来源状态 open | 2026-07-23 | 来源

V3P-002030 | codetables-gf3-231-7

Minimum distance of a [231, 7] linear code over GF(3)

codes are known with d = 149, and no code can have d greater than 151.

来源状态 open | 2026-07-23 | 来源

V3P-002031 | codetables-gf3-231-8

Minimum distance of a [231, 8] linear code over GF(3)

codes are known with d = 146, and no code can have d greater than 150.

来源状态 open | 2026-07-23 | 来源

V3P-002032 | codetables-gf3-231-9

Minimum distance of a [231, 9] linear code over GF(3)

codes are known with d = 144, and no code can have d greater than 148.

来源状态 open | 2026-07-23 | 来源

V3P-002033 | codetables-gf3-232-10

Minimum distance of a [232, 10] linear code over GF(3)

codes are known with d = 143, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-002034 | codetables-gf3-232-11

Minimum distance of a [232, 11] linear code over GF(3)

codes are known with d = 142, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-002035 | codetables-gf3-232-12

Minimum distance of a [232, 12] linear code over GF(3)

codes are known with d = 136, and no code can have d greater than 146.

来源状态 open | 2026-07-23 | 来源

V3P-002036 | codetables-gf3-232-7

Minimum distance of a [232, 7] linear code over GF(3)

codes are known with d = 150, and no code can have d greater than 152.

来源状态 open | 2026-07-23 | 来源

V3P-002037 | codetables-gf3-232-8

Minimum distance of a [232, 8] linear code over GF(3)

codes are known with d = 147, and no code can have d greater than 151.

来源状态 open | 2026-07-23 | 来源

V3P-002038 | codetables-gf3-232-9

Minimum distance of a [232, 9] linear code over GF(3)

codes are known with d = 145, and no code can have d greater than 149.

来源状态 open | 2026-07-23 | 来源

V3P-002039 | codetables-gf3-233-10

Minimum distance of a [233, 10] linear code over GF(3)

codes are known with d = 144, and no code can have d greater than 148.

来源状态 open | 2026-07-23 | 来源

V3P-002040 | codetables-gf3-233-11

Minimum distance of a [233, 11] linear code over GF(3)

codes are known with d = 143, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-002041 | codetables-gf3-233-12

Minimum distance of a [233, 12] linear code over GF(3)

codes are known with d = 137, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-002042 | codetables-gf3-233-6

Minimum distance of a [233, 6] linear code over GF(3)

codes are known with d = 153, and no code can have d greater than 154.

来源状态 open | 2026-07-23 | 来源

V3P-002043 | codetables-gf3-233-7

Minimum distance of a [233, 7] linear code over GF(3)

codes are known with d = 150, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-002044 | codetables-gf3-233-8

Minimum distance of a [233, 8] linear code over GF(3)

codes are known with d = 147, and no code can have d greater than 152.

来源状态 open | 2026-07-23 | 来源

V3P-002045 | codetables-gf3-233-9

Minimum distance of a [233, 9] linear code over GF(3)

codes are known with d = 146, and no code can have d greater than 149.

来源状态 open | 2026-07-23 | 来源

V3P-002046 | codetables-gf3-234-10

Minimum distance of a [234, 10] linear code over GF(3)

codes are known with d = 145, and no code can have d greater than 149.

来源状态 open | 2026-07-23 | 来源

V3P-002047 | codetables-gf3-234-11

Minimum distance of a [234, 11] linear code over GF(3)

codes are known with d = 144, and no code can have d greater than 148.

来源状态 open | 2026-07-23 | 来源

V3P-002048 | codetables-gf3-234-12

Minimum distance of a [234, 12] linear code over GF(3)

codes are known with d = 138, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-002049 | codetables-gf3-234-6

Minimum distance of a [234, 6] linear code over GF(3)

codes are known with d = 154, and no code can have d greater than 155.

来源状态 open | 2026-07-23 | 来源

V3P-002050 | codetables-gf3-234-7

Minimum distance of a [234, 7] linear code over GF(3)

codes are known with d = 151, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-002051 | codetables-gf3-234-8

Minimum distance of a [234, 8] linear code over GF(3)

codes are known with d = 148, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-002052 | codetables-gf3-234-9

Minimum distance of a [234, 9] linear code over GF(3)

codes are known with d = 147, and no code can have d greater than 150.

来源状态 open | 2026-07-23 | 来源

V3P-002053 | codetables-gf3-235-10

Minimum distance of a [235, 10] linear code over GF(3)

codes are known with d = 146, and no code can have d greater than 149.

来源状态 open | 2026-07-23 | 来源

V3P-002054 | codetables-gf3-235-11

Minimum distance of a [235, 11] linear code over GF(3)

codes are known with d = 145, and no code can have d greater than 149.

来源状态 open | 2026-07-23 | 来源

V3P-002055 | codetables-gf3-235-12

Minimum distance of a [235, 12] linear code over GF(3)

codes are known with d = 138, and no code can have d greater than 148.

来源状态 open | 2026-07-23 | 来源

V3P-002056 | codetables-gf3-235-7

Minimum distance of a [235, 7] linear code over GF(3)

codes are known with d = 152, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-002057 | codetables-gf3-235-8

Minimum distance of a [235, 8] linear code over GF(3)

codes are known with d = 149, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-002058 | codetables-gf3-235-9

Minimum distance of a [235, 9] linear code over GF(3)

codes are known with d = 147, and no code can have d greater than 151.

来源状态 open | 2026-07-23 | 来源

V3P-002059 | codetables-gf3-236-10

Minimum distance of a [236, 10] linear code over GF(3)

codes are known with d = 147, and no code can have d greater than 150.

来源状态 open | 2026-07-23 | 来源

V3P-002060 | codetables-gf3-236-11

Minimum distance of a [236, 11] linear code over GF(3)

codes are known with d = 146, and no code can have d greater than 149.

来源状态 open | 2026-07-23 | 来源

V3P-002061 | codetables-gf3-236-12

Minimum distance of a [236, 12] linear code over GF(3)

codes are known with d = 138, and no code can have d greater than 149.

来源状态 open | 2026-07-23 | 来源

V3P-002062 | codetables-gf3-236-7

Minimum distance of a [236, 7] linear code over GF(3)

codes are known with d = 153, and no code can have d greater than 154.

来源状态 open | 2026-07-23 | 来源

V3P-002063 | codetables-gf3-236-8

Minimum distance of a [236, 8] linear code over GF(3)

codes are known with d = 150, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-002064 | codetables-gf3-236-9

Minimum distance of a [236, 9] linear code over GF(3)

codes are known with d = 148, and no code can have d greater than 152.

来源状态 open | 2026-07-23 | 来源

V3P-002065 | codetables-gf3-237-10

Minimum distance of a [237, 10] linear code over GF(3)

codes are known with d = 148, and no code can have d greater than 151.

来源状态 open | 2026-07-23 | 来源

V3P-002066 | codetables-gf3-237-11

Minimum distance of a [237, 11] linear code over GF(3)

codes are known with d = 147, and no code can have d greater than 150.

来源状态 open | 2026-07-23 | 来源

V3P-002067 | codetables-gf3-237-12

Minimum distance of a [237, 12] linear code over GF(3)

codes are known with d = 139, and no code can have d greater than 149.

来源状态 open | 2026-07-23 | 来源

V3P-002068 | codetables-gf3-237-7

Minimum distance of a [237, 7] linear code over GF(3)

codes are known with d = 153, and no code can have d greater than 155.

来源状态 open | 2026-07-23 | 来源

V3P-002069 | codetables-gf3-237-8

Minimum distance of a [237, 8] linear code over GF(3)

codes are known with d = 150, and no code can have d greater than 154.

来源状态 open | 2026-07-23 | 来源

V3P-002070 | codetables-gf3-237-9

Minimum distance of a [237, 9] linear code over GF(3)

codes are known with d = 149, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-002071 | codetables-gf3-238-10

Minimum distance of a [238, 10] linear code over GF(3)

codes are known with d = 149, and no code can have d greater than 151.

来源状态 open | 2026-07-23 | 来源

V3P-002072 | codetables-gf3-238-11

Minimum distance of a [238, 11] linear code over GF(3)

codes are known with d = 148, and no code can have d greater than 151.

来源状态 open | 2026-07-23 | 来源

V3P-002073 | codetables-gf3-238-12

Minimum distance of a [238, 12] linear code over GF(3)

codes are known with d = 140, and no code can have d greater than 150.

来源状态 open | 2026-07-23 | 来源

V3P-002074 | codetables-gf3-238-7

Minimum distance of a [238, 7] linear code over GF(3)

codes are known with d = 153, and no code can have d greater than 156.

来源状态 open | 2026-07-23 | 来源

V3P-002075 | codetables-gf3-238-8

Minimum distance of a [238, 8] linear code over GF(3)

codes are known with d = 151, and no code can have d greater than 155.

来源状态 open | 2026-07-23 | 来源

V3P-002076 | codetables-gf3-238-9

Minimum distance of a [238, 9] linear code over GF(3)

codes are known with d = 150, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-002077 | codetables-gf3-239-10

Minimum distance of a [239, 10] linear code over GF(3)

codes are known with d = 150, and no code can have d greater than 152.

来源状态 open | 2026-07-23 | 来源

V3P-002078 | codetables-gf3-239-11

Minimum distance of a [239, 11] linear code over GF(3)

codes are known with d = 149, and no code can have d greater than 151.

来源状态 open | 2026-07-23 | 来源

V3P-002079 | codetables-gf3-239-12

Minimum distance of a [239, 12] linear code over GF(3)

codes are known with d = 140, and no code can have d greater than 151.

来源状态 open | 2026-07-23 | 来源

V3P-002080 | codetables-gf3-239-7

Minimum distance of a [239, 7] linear code over GF(3)

codes are known with d = 154, and no code can have d greater than 156.

来源状态 open | 2026-07-23 | 来源

V3P-002081 | codetables-gf3-239-8

Minimum distance of a [239, 8] linear code over GF(3)

codes are known with d = 152, and no code can have d greater than 156.

来源状态 open | 2026-07-23 | 来源

V3P-002082 | codetables-gf3-239-9

Minimum distance of a [239, 9] linear code over GF(3)

codes are known with d = 151, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-002083 | codetables-gf3-24-10

Minimum distance of a [24, 10] linear code over GF(3)

codes are known with d = 9, and no code can have d greater than 10.

来源状态 open | 2026-07-23 | 来源

V3P-002084 | codetables-gf3-24-9

Minimum distance of a [24, 9] linear code over GF(3)

codes are known with d = 10, and no code can have d greater than 11.

来源状态 open | 2026-07-23 | 来源

V3P-002085 | codetables-gf3-240-10

Minimum distance of a [240, 10] linear code over GF(3)

codes are known with d = 151, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-002086 | codetables-gf3-240-11

Minimum distance of a [240, 11] linear code over GF(3)

codes are known with d = 150, and no code can have d greater than 152.

来源状态 open | 2026-07-23 | 来源

V3P-002087 | codetables-gf3-240-12

Minimum distance of a [240, 12] linear code over GF(3)

codes are known with d = 141, and no code can have d greater than 151.

来源状态 open | 2026-07-23 | 来源

V3P-002088 | codetables-gf3-240-7

Minimum distance of a [240, 7] linear code over GF(3)

codes are known with d = 154, and no code can have d greater than 157.

来源状态 open | 2026-07-23 | 来源

V3P-002089 | codetables-gf3-240-8

Minimum distance of a [240, 8] linear code over GF(3)

codes are known with d = 153, and no code can have d greater than 156.

来源状态 open | 2026-07-23 | 来源

V3P-002090 | codetables-gf3-240-9

Minimum distance of a [240, 9] linear code over GF(3)

codes are known with d = 152, and no code can have d greater than 154.

来源状态 open | 2026-07-23 | 来源

V3P-002091 | codetables-gf3-241-10

Minimum distance of a [241, 10] linear code over GF(3)

codes are known with d = 152, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-002092 | codetables-gf3-241-11

Minimum distance of a [241, 11] linear code over GF(3)

codes are known with d = 151, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-002093 | codetables-gf3-241-12

Minimum distance of a [241, 12] linear code over GF(3)

codes are known with d = 142, and no code can have d greater than 152.

来源状态 open | 2026-07-23 | 来源

V3P-002094 | codetables-gf3-241-7

Minimum distance of a [241, 7] linear code over GF(3)

codes are known with d = 155, and no code can have d greater than 158.

来源状态 open | 2026-07-23 | 来源

V3P-002095 | codetables-gf3-241-8

Minimum distance of a [241, 8] linear code over GF(3)

codes are known with d = 154, and no code can have d greater than 157.

来源状态 open | 2026-07-23 | 来源

V3P-002096 | codetables-gf3-241-9

Minimum distance of a [241, 9] linear code over GF(3)

codes are known with d = 153, and no code can have d greater than 155.

来源状态 open | 2026-07-23 | 来源

V3P-002097 | codetables-gf3-242-10

Minimum distance of a [242, 10] linear code over GF(3)

codes are known with d = 153, and no code can have d greater than 154.

来源状态 open | 2026-07-23 | 来源

V3P-002098 | codetables-gf3-242-11

Minimum distance of a [242, 11] linear code over GF(3)

codes are known with d = 152, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-002099 | codetables-gf3-242-12

Minimum distance of a [242, 12] linear code over GF(3)

codes are known with d = 143, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-002100 | codetables-gf3-242-7

Minimum distance of a [242, 7] linear code over GF(3)

codes are known with d = 156, and no code can have d greater than 159.

来源状态 open | 2026-07-23 | 来源

V3P-002101 | codetables-gf3-242-8

Minimum distance of a [242, 8] linear code over GF(3)

codes are known with d = 155, and no code can have d greater than 158.

来源状态 open | 2026-07-23 | 来源

V3P-002102 | codetables-gf3-242-9

Minimum distance of a [242, 9] linear code over GF(3)

codes are known with d = 153, and no code can have d greater than 156.

来源状态 open | 2026-07-23 | 来源

V3P-002103 | codetables-gf3-243-10

Minimum distance of a [243, 10] linear code over GF(3)

codes are known with d = 153, and no code can have d greater than 155.

来源状态 open | 2026-07-23 | 来源

V3P-002104 | codetables-gf3-243-11

Minimum distance of a [243, 11] linear code over GF(3)

codes are known with d = 153, and no code can have d greater than 154.

来源状态 open | 2026-07-23 | 来源

V3P-002105 | codetables-gf3-243-12

Minimum distance of a [243, 12] linear code over GF(3)

codes are known with d = 144, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-002106 | codetables-gf3-243-7

Minimum distance of a [243, 7] linear code over GF(3)

codes are known with d = 156, and no code can have d greater than 159.

来源状态 open | 2026-07-23 | 来源

V3P-002107 | codetables-gf3-243-8

Minimum distance of a [243, 8] linear code over GF(3)

codes are known with d = 156, and no code can have d greater than 158.

来源状态 open | 2026-07-23 | 来源

V3P-002108 | codetables-gf3-243-9

Minimum distance of a [243, 9] linear code over GF(3)

codes are known with d = 153, and no code can have d greater than 157.

来源状态 open | 2026-07-23 | 来源

V3P-002109 | codetables-gf3-25-10

Minimum distance of a [25, 10] linear code over GF(3)

codes are known with d = 10, and no code can have d greater than 11.

来源状态 open | 2026-07-23 | 来源

V3P-002110 | codetables-gf3-25-11

Minimum distance of a [25, 11] linear code over GF(3)

codes are known with d = 9, and no code can have d greater than 10.

来源状态 open | 2026-07-23 | 来源

V3P-002111 | codetables-gf3-26-10

Minimum distance of a [26, 10] linear code over GF(3)

codes are known with d = 10, and no code can have d greater than 11.

来源状态 open | 2026-07-23 | 来源

V3P-002112 | codetables-gf3-26-11

Minimum distance of a [26, 11] linear code over GF(3)

codes are known with d = 9, and no code can have d greater than 11.

来源状态 open | 2026-07-23 | 来源

V3P-002113 | codetables-gf3-26-12

Minimum distance of a [26, 12] linear code over GF(3)

codes are known with d = 9, and no code can have d greater than 10.

来源状态 open | 2026-07-23 | 来源

V3P-002114 | codetables-gf3-27-10

Minimum distance of a [27, 10] linear code over GF(3)

codes are known with d = 11, and no code can have d greater than 12.

来源状态 open | 2026-07-23 | 来源

V3P-002115 | codetables-gf3-27-11

Minimum distance of a [27, 11] linear code over GF(3)

codes are known with d = 10, and no code can have d greater than 11.

来源状态 open | 2026-07-23 | 来源

V3P-002116 | codetables-gf3-27-12

Minimum distance of a [27, 12] linear code over GF(3)

codes are known with d = 9, and no code can have d greater than 11.

来源状态 open | 2026-07-23 | 来源

V3P-002117 | codetables-gf3-28-11

Minimum distance of a [28, 11] linear code over GF(3)

codes are known with d = 11, and no code can have d greater than 12.

来源状态 open | 2026-07-23 | 来源

V3P-002118 | codetables-gf3-28-12

Minimum distance of a [28, 12] linear code over GF(3)

codes are known with d = 10, and no code can have d greater than 11.

来源状态 open | 2026-07-23 | 来源

V3P-002119 | codetables-gf3-28-9

Minimum distance of a [28, 9] linear code over GF(3)

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-002120 | codetables-gf3-29-10

Minimum distance of a [29, 10] linear code over GF(3)

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-002121 | codetables-gf3-29-12

Minimum distance of a [29, 12] linear code over GF(3)

codes are known with d = 11, and no code can have d greater than 12.

来源状态 open | 2026-07-23 | 来源

V3P-002122 | codetables-gf3-29-9

Minimum distance of a [29, 9] linear code over GF(3)

codes are known with d = 13, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-002123 | codetables-gf3-30-10

Minimum distance of a [30, 10] linear code over GF(3)

codes are known with d = 13, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-002124 | codetables-gf3-30-11

Minimum distance of a [30, 11] linear code over GF(3)

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-002125 | codetables-gf3-30-9

Minimum distance of a [30, 9] linear code over GF(3)

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-002126 | codetables-gf3-31-10

Minimum distance of a [31, 10] linear code over GF(3)

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-002127 | codetables-gf3-31-11

Minimum distance of a [31, 11] linear code over GF(3)

codes are known with d = 12, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-002128 | codetables-gf3-31-12

Minimum distance of a [31, 12] linear code over GF(3)

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-002129 | codetables-gf3-31-8

Minimum distance of a [31, 8] linear code over GF(3)

codes are known with d = 15, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-002130 | codetables-gf3-32-11

Minimum distance of a [32, 11] linear code over GF(3)

codes are known with d = 13, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-002131 | codetables-gf3-32-12

Minimum distance of a [32, 12] linear code over GF(3)

codes are known with d = 12, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-002132 | codetables-gf3-32-7

Minimum distance of a [32, 7] linear code over GF(3)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-002133 | codetables-gf3-32-8

Minimum distance of a [32, 8] linear code over GF(3)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-002134 | codetables-gf3-32-9

Minimum distance of a [32, 9] linear code over GF(3)

codes are known with d = 15, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-002135 | codetables-gf3-33-10

Minimum distance of a [33, 10] linear code over GF(3)

codes are known with d = 15, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-002136 | codetables-gf3-33-11

Minimum distance of a [33, 11] linear code over GF(3)

codes are known with d = 13, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-002137 | codetables-gf3-33-12

Minimum distance of a [33, 12] linear code over GF(3)

codes are known with d = 13, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-002138 | codetables-gf3-33-6

Minimum distance of a [33, 6] linear code over GF(3)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-002139 | codetables-gf3-33-8

Minimum distance of a [33, 8] linear code over GF(3)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-002140 | codetables-gf3-33-9

Minimum distance of a [33, 9] linear code over GF(3)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-002141 | codetables-gf3-34-10

Minimum distance of a [34, 10] linear code over GF(3)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-002142 | codetables-gf3-34-11

Minimum distance of a [34, 11] linear code over GF(3)

codes are known with d = 14, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-002143 | codetables-gf3-34-12

Minimum distance of a [34, 12] linear code over GF(3)

codes are known with d = 13, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-002144 | codetables-gf3-34-6

Minimum distance of a [34, 6] linear code over GF(3)

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-002145 | codetables-gf3-34-7

Minimum distance of a [34, 7] linear code over GF(3)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-002146 | codetables-gf3-34-9

Minimum distance of a [34, 9] linear code over GF(3)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-002147 | codetables-gf3-35-10

Minimum distance of a [35, 10] linear code over GF(3)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-002148 | codetables-gf3-35-11

Minimum distance of a [35, 11] linear code over GF(3)

codes are known with d = 15, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-002149 | codetables-gf3-35-12

Minimum distance of a [35, 12] linear code over GF(3)

codes are known with d = 14, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-002150 | codetables-gf3-35-7

Minimum distance of a [35, 7] linear code over GF(3)

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-002151 | codetables-gf3-36-11

Minimum distance of a [36, 11] linear code over GF(3)

codes are known with d = 16, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-002152 | codetables-gf3-36-12

Minimum distance of a [36, 12] linear code over GF(3)

codes are known with d = 15, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-002153 | codetables-gf3-36-8

Minimum distance of a [36, 8] linear code over GF(3)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-002154 | codetables-gf3-37-11

Minimum distance of a [37, 11] linear code over GF(3)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-002155 | codetables-gf3-37-12

Minimum distance of a [37, 12] linear code over GF(3)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-002156 | codetables-gf3-37-8

Minimum distance of a [37, 8] linear code over GF(3)

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-002157 | codetables-gf3-37-9

Minimum distance of a [37, 9] linear code over GF(3)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-002158 | codetables-gf3-38-10

Minimum distance of a [38, 10] linear code over GF(3)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-002159 | codetables-gf3-38-12

Minimum distance of a [38, 12] linear code over GF(3)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-002160 | codetables-gf3-38-8

Minimum distance of a [38, 8] linear code over GF(3)

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-002161 | codetables-gf3-38-9

Minimum distance of a [38, 9] linear code over GF(3)

codes are known with d = 18, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-002162 | codetables-gf3-39-10

Minimum distance of a [39, 10] linear code over GF(3)

codes are known with d = 18, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-002163 | codetables-gf3-39-11

Minimum distance of a [39, 11] linear code over GF(3)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-002164 | codetables-gf3-39-7

Minimum distance of a [39, 7] linear code over GF(3)

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-002165 | codetables-gf3-39-9

Minimum distance of a [39, 9] linear code over GF(3)

codes are known with d = 19, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-002166 | codetables-gf3-40-10

Minimum distance of a [40, 10] linear code over GF(3)

codes are known with d = 19, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-002167 | codetables-gf3-40-11

Minimum distance of a [40, 11] linear code over GF(3)

codes are known with d = 18, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-002168 | codetables-gf3-40-12

Minimum distance of a [40, 12] linear code over GF(3)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-002169 | codetables-gf3-40-7

Minimum distance of a [40, 7] linear code over GF(3)

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-002170 | codetables-gf3-40-8

Minimum distance of a [40, 8] linear code over GF(3)

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-002171 | codetables-gf3-40-9

Minimum distance of a [40, 9] linear code over GF(3)

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-002172 | codetables-gf3-41-10

Minimum distance of a [41, 10] linear code over GF(3)

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-002173 | codetables-gf3-41-11

Minimum distance of a [41, 11] linear code over GF(3)

codes are known with d = 18, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-002174 | codetables-gf3-41-12

Minimum distance of a [41, 12] linear code over GF(3)

codes are known with d = 18, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-002175 | codetables-gf3-41-7

Minimum distance of a [41, 7] linear code over GF(3)

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-002176 | codetables-gf3-41-8

Minimum distance of a [41, 8] linear code over GF(3)

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-002177 | codetables-gf3-41-9

Minimum distance of a [41, 9] linear code over GF(3)

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-002178 | codetables-gf3-42-10

Minimum distance of a [42, 10] linear code over GF(3)

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-002179 | codetables-gf3-42-11

Minimum distance of a [42, 11] linear code over GF(3)

codes are known with d = 19, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-002180 | codetables-gf3-42-12

Minimum distance of a [42, 12] linear code over GF(3)

codes are known with d = 18, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-002181 | codetables-gf3-42-8

Minimum distance of a [42, 8] linear code over GF(3)

codes are known with d = 22, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-002182 | codetables-gf3-42-9

Minimum distance of a [42, 9] linear code over GF(3)

codes are known with d = 21, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-002183 | codetables-gf3-43-10

Minimum distance of a [43, 10] linear code over GF(3)

codes are known with d = 21, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-002184 | codetables-gf3-43-11

Minimum distance of a [43, 11] linear code over GF(3)

codes are known with d = 20, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-002185 | codetables-gf3-43-12

Minimum distance of a [43, 12] linear code over GF(3)

codes are known with d = 18, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-002186 | codetables-gf3-43-7

Minimum distance of a [43, 7] linear code over GF(3)

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-002187 | codetables-gf3-43-8

Minimum distance of a [43, 8] linear code over GF(3)

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-002188 | codetables-gf3-43-9

Minimum distance of a [43, 9] linear code over GF(3)

codes are known with d = 22, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-002189 | codetables-gf3-44-10

Minimum distance of a [44, 10] linear code over GF(3)

codes are known with d = 21, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-002190 | codetables-gf3-44-11

Minimum distance of a [44, 11] linear code over GF(3)

codes are known with d = 21, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-002191 | codetables-gf3-44-12

Minimum distance of a [44, 12] linear code over GF(3)

codes are known with d = 19, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-002192 | codetables-gf3-44-7

Minimum distance of a [44, 7] linear code over GF(3)

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-002193 | codetables-gf3-44-8

Minimum distance of a [44, 8] linear code over GF(3)

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-002194 | codetables-gf3-44-9

Minimum distance of a [44, 9] linear code over GF(3)

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-002195 | codetables-gf3-45-10

Minimum distance of a [45, 10] linear code over GF(3)

codes are known with d = 22, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-002196 | codetables-gf3-45-11

Minimum distance of a [45, 11] linear code over GF(3)

codes are known with d = 21, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-002197 | codetables-gf3-45-12

Minimum distance of a [45, 12] linear code over GF(3)

codes are known with d = 20, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-002198 | codetables-gf3-45-7

Minimum distance of a [45, 7] linear code over GF(3)

codes are known with d = 25, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-002199 | codetables-gf3-45-8

Minimum distance of a [45, 8] linear code over GF(3)

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-002200 | codetables-gf3-45-9

Minimum distance of a [45, 9] linear code over GF(3)

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-002201 | codetables-gf3-46-10

Minimum distance of a [46, 10] linear code over GF(3)

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-002202 | codetables-gf3-46-11

Minimum distance of a [46, 11] linear code over GF(3)

codes are known with d = 21, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-002203 | codetables-gf3-46-12

Minimum distance of a [46, 12] linear code over GF(3)

codes are known with d = 21, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-002204 | codetables-gf3-46-6

Minimum distance of a [46, 6] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-002205 | codetables-gf3-46-7

Minimum distance of a [46, 7] linear code over GF(3)

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-002206 | codetables-gf3-46-8

Minimum distance of a [46, 8] linear code over GF(3)

codes are known with d = 25, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-002207 | codetables-gf3-46-9

Minimum distance of a [46, 9] linear code over GF(3)

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-002208 | codetables-gf3-47-10

Minimum distance of a [47, 10] linear code over GF(3)

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-002209 | codetables-gf3-47-11

Minimum distance of a [47, 11] linear code over GF(3)

codes are known with d = 22, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-002210 | codetables-gf3-47-12

Minimum distance of a [47, 12] linear code over GF(3)

codes are known with d = 21, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-002211 | codetables-gf3-47-8

Minimum distance of a [47, 8] linear code over GF(3)

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-002212 | codetables-gf3-47-9

Minimum distance of a [47, 9] linear code over GF(3)

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-002213 | codetables-gf3-48-10

Minimum distance of a [48, 10] linear code over GF(3)

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-002214 | codetables-gf3-48-11

Minimum distance of a [48, 11] linear code over GF(3)

codes are known with d = 23, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-002215 | codetables-gf3-48-12

Minimum distance of a [48, 12] linear code over GF(3)

codes are known with d = 21, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-002216 | codetables-gf3-48-7

Minimum distance of a [48, 7] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-002217 | codetables-gf3-48-9

Minimum distance of a [48, 9] linear code over GF(3)

codes are known with d = 25, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-002218 | codetables-gf3-49-10

Minimum distance of a [49, 10] linear code over GF(3)

codes are known with d = 24, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-002219 | codetables-gf3-49-11

Minimum distance of a [49, 11] linear code over GF(3)

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-002220 | codetables-gf3-49-12

Minimum distance of a [49, 12] linear code over GF(3)

codes are known with d = 22, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-002221 | codetables-gf3-49-7

Minimum distance of a [49, 7] linear code over GF(3)

codes are known with d = 28, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-002222 | codetables-gf3-49-8

Minimum distance of a [49, 8] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-002223 | codetables-gf3-49-9

Minimum distance of a [49, 9] linear code over GF(3)

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-002224 | codetables-gf3-50-10

Minimum distance of a [50, 10] linear code over GF(3)

codes are known with d = 25, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-002225 | codetables-gf3-50-11

Minimum distance of a [50, 11] linear code over GF(3)

codes are known with d = 24, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-002226 | codetables-gf3-50-12

Minimum distance of a [50, 12] linear code over GF(3)

codes are known with d = 23, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-002227 | codetables-gf3-50-7

Minimum distance of a [50, 7] linear code over GF(3)

codes are known with d = 29, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-002228 | codetables-gf3-50-8

Minimum distance of a [50, 8] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-002229 | codetables-gf3-50-9

Minimum distance of a [50, 9] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-002230 | codetables-gf3-51-10

Minimum distance of a [51, 10] linear code over GF(3)

codes are known with d = 26, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-002231 | codetables-gf3-51-11

Minimum distance of a [51, 11] linear code over GF(3)

codes are known with d = 24, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-002232 | codetables-gf3-51-12

Minimum distance of a [51, 12] linear code over GF(3)

codes are known with d = 24, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-002233 | codetables-gf3-51-8

Minimum distance of a [51, 8] linear code over GF(3)

codes are known with d = 28, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-002234 | codetables-gf3-51-9

Minimum distance of a [51, 9] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-002235 | codetables-gf3-52-10

Minimum distance of a [52, 10] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-002236 | codetables-gf3-52-11

Minimum distance of a [52, 11] linear code over GF(3)

codes are known with d = 25, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-002237 | codetables-gf3-52-12

Minimum distance of a [52, 12] linear code over GF(3)

codes are known with d = 24, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-002238 | codetables-gf3-52-7

Minimum distance of a [52, 7] linear code over GF(3)

codes are known with d = 30, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-002239 | codetables-gf3-52-8

Minimum distance of a [52, 8] linear code over GF(3)

codes are known with d = 29, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-002240 | codetables-gf3-52-9

Minimum distance of a [52, 9] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-002241 | codetables-gf3-53-10

Minimum distance of a [53, 10] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-002242 | codetables-gf3-53-11

Minimum distance of a [53, 11] linear code over GF(3)

codes are known with d = 26, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-002243 | codetables-gf3-53-12

Minimum distance of a [53, 12] linear code over GF(3)

codes are known with d = 24, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-002244 | codetables-gf3-53-7

Minimum distance of a [53, 7] linear code over GF(3)

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-002245 | codetables-gf3-53-8

Minimum distance of a [53, 8] linear code over GF(3)

codes are known with d = 30, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-002246 | codetables-gf3-53-9

Minimum distance of a [53, 9] linear code over GF(3)

codes are known with d = 28, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-002247 | codetables-gf3-54-10

Minimum distance of a [54, 10] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-002248 | codetables-gf3-54-11

Minimum distance of a [54, 11] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-002249 | codetables-gf3-54-12

Minimum distance of a [54, 12] linear code over GF(3)

codes are known with d = 25, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-002250 | codetables-gf3-54-7

Minimum distance of a [54, 7] linear code over GF(3)

codes are known with d = 31, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-002251 | codetables-gf3-54-8

Minimum distance of a [54, 8] linear code over GF(3)

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-002252 | codetables-gf3-54-9

Minimum distance of a [54, 9] linear code over GF(3)

codes are known with d = 29, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-002253 | codetables-gf3-55-10

Minimum distance of a [55, 10] linear code over GF(3)

codes are known with d = 28, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-002254 | codetables-gf3-55-11

Minimum distance of a [55, 11] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-002255 | codetables-gf3-55-12

Minimum distance of a [55, 12] linear code over GF(3)

codes are known with d = 26, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-002256 | codetables-gf3-55-7

Minimum distance of a [55, 7] linear code over GF(3)

codes are known with d = 32, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-002257 | codetables-gf3-55-8

Minimum distance of a [55, 8] linear code over GF(3)

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-002258 | codetables-gf3-55-9

Minimum distance of a [55, 9] linear code over GF(3)

codes are known with d = 30, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-002259 | codetables-gf3-56-10

Minimum distance of a [56, 10] linear code over GF(3)

codes are known with d = 29, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-002260 | codetables-gf3-56-11

Minimum distance of a [56, 11] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-002261 | codetables-gf3-56-12

Minimum distance of a [56, 12] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-002262 | codetables-gf3-56-7

Minimum distance of a [56, 7] linear code over GF(3)

codes are known with d = 33, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-002263 | codetables-gf3-56-8

Minimum distance of a [56, 8] linear code over GF(3)

codes are known with d = 31, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-002264 | codetables-gf3-56-9

Minimum distance of a [56, 9] linear code over GF(3)

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-002265 | codetables-gf3-57-10

Minimum distance of a [57, 10] linear code over GF(3)

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-002266 | codetables-gf3-57-11

Minimum distance of a [57, 11] linear code over GF(3)

codes are known with d = 28, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-002267 | codetables-gf3-57-12

Minimum distance of a [57, 12] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-002268 | codetables-gf3-57-7

Minimum distance of a [57, 7] linear code over GF(3)

codes are known with d = 33, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-002269 | codetables-gf3-57-8

Minimum distance of a [57, 8] linear code over GF(3)

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-002270 | codetables-gf3-57-9

Minimum distance of a [57, 9] linear code over GF(3)

codes are known with d = 30, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-002271 | codetables-gf3-58-10

Minimum distance of a [58, 10] linear code over GF(3)

codes are known with d = 30, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-002272 | codetables-gf3-58-11

Minimum distance of a [58, 11] linear code over GF(3)

codes are known with d = 29, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-002273 | codetables-gf3-58-12

Minimum distance of a [58, 12] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-002274 | codetables-gf3-58-7

Minimum distance of a [58, 7] linear code over GF(3)

codes are known with d = 34, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-002275 | codetables-gf3-58-8

Minimum distance of a [58, 8] linear code over GF(3)

codes are known with d = 33, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-002276 | codetables-gf3-58-9

Minimum distance of a [58, 9] linear code over GF(3)

codes are known with d = 31, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-002277 | codetables-gf3-59-10

Minimum distance of a [59, 10] linear code over GF(3)

codes are known with d = 30, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-002278 | codetables-gf3-59-11

Minimum distance of a [59, 11] linear code over GF(3)

codes are known with d = 30, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-002279 | codetables-gf3-59-12

Minimum distance of a [59, 12] linear code over GF(3)

codes are known with d = 27, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-002280 | codetables-gf3-59-7

Minimum distance of a [59, 7] linear code over GF(3)

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-002281 | codetables-gf3-59-8

Minimum distance of a [59, 8] linear code over GF(3)

codes are known with d = 33, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-002282 | codetables-gf3-59-9

Minimum distance of a [59, 9] linear code over GF(3)

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-002283 | codetables-gf3-60-10

Minimum distance of a [60, 10] linear code over GF(3)

codes are known with d = 31, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-002284 | codetables-gf3-60-11

Minimum distance of a [60, 11] linear code over GF(3)

codes are known with d = 30, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-002285 | codetables-gf3-60-12

Minimum distance of a [60, 12] linear code over GF(3)

codes are known with d = 28, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-002286 | codetables-gf3-60-6

Minimum distance of a [60, 6] linear code over GF(3)

codes are known with d = 36, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-002287 | codetables-gf3-60-8

Minimum distance of a [60, 8] linear code over GF(3)

codes are known with d = 33, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-002288 | codetables-gf3-60-9

Minimum distance of a [60, 9] linear code over GF(3)

codes are known with d = 33, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-002289 | codetables-gf3-61-10

Minimum distance of a [61, 10] linear code over GF(3)

codes are known with d = 32, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-002290 | codetables-gf3-61-11

Minimum distance of a [61, 11] linear code over GF(3)

codes are known with d = 31, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-002291 | codetables-gf3-61-12

Minimum distance of a [61, 12] linear code over GF(3)

codes are known with d = 29, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-002292 | codetables-gf3-61-6

Minimum distance of a [61, 6] linear code over GF(3)

codes are known with d = 37, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-002293 | codetables-gf3-61-7

Minimum distance of a [61, 7] linear code over GF(3)

codes are known with d = 36, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-002294 | codetables-gf3-61-8

Minimum distance of a [61, 8] linear code over GF(3)

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-002295 | codetables-gf3-61-9

Minimum distance of a [61, 9] linear code over GF(3)

codes are known with d = 33, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-002296 | codetables-gf3-62-10

Minimum distance of a [62, 10] linear code over GF(3)

codes are known with d = 33, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-002297 | codetables-gf3-62-11

Minimum distance of a [62, 11] linear code over GF(3)

codes are known with d = 31, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-002298 | codetables-gf3-62-12

Minimum distance of a [62, 12] linear code over GF(3)

codes are known with d = 30, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-002299 | codetables-gf3-62-6

Minimum distance of a [62, 6] linear code over GF(3)

codes are known with d = 38, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-002300 | codetables-gf3-62-7

Minimum distance of a [62, 7] linear code over GF(3)

codes are known with d = 36, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-002301 | codetables-gf3-62-8

Minimum distance of a [62, 8] linear code over GF(3)

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-002302 | codetables-gf3-62-9

Minimum distance of a [62, 9] linear code over GF(3)

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-002303 | codetables-gf3-63-10

Minimum distance of a [63, 10] linear code over GF(3)

codes are known with d = 33, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-002304 | codetables-gf3-63-11

Minimum distance of a [63, 11] linear code over GF(3)

codes are known with d = 32, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-002305 | codetables-gf3-63-12

Minimum distance of a [63, 12] linear code over GF(3)

codes are known with d = 30, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-002306 | codetables-gf3-63-7

Minimum distance of a [63, 7] linear code over GF(3)

codes are known with d = 37, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-002307 | codetables-gf3-63-8

Minimum distance of a [63, 8] linear code over GF(3)

codes are known with d = 36, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-002308 | codetables-gf3-63-9

Minimum distance of a [63, 9] linear code over GF(3)

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-002309 | codetables-gf3-64-10

Minimum distance of a [64, 10] linear code over GF(3)

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-002310 | codetables-gf3-64-11

Minimum distance of a [64, 11] linear code over GF(3)

codes are known with d = 33, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-002311 | codetables-gf3-64-12

Minimum distance of a [64, 12] linear code over GF(3)

codes are known with d = 31, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-002312 | codetables-gf3-64-6

Minimum distance of a [64, 6] linear code over GF(3)

codes are known with d = 39, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-002313 | codetables-gf3-64-7

Minimum distance of a [64, 7] linear code over GF(3)

codes are known with d = 38, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-002314 | codetables-gf3-64-8

Minimum distance of a [64, 8] linear code over GF(3)

codes are known with d = 37, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-002315 | codetables-gf3-64-9

Minimum distance of a [64, 9] linear code over GF(3)

codes are known with d = 36, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-002316 | codetables-gf3-65-10

Minimum distance of a [65, 10] linear code over GF(3)

codes are known with d = 35, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-002317 | codetables-gf3-65-11

Minimum distance of a [65, 11] linear code over GF(3)

codes are known with d = 33, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-002318 | codetables-gf3-65-12

Minimum distance of a [65, 12] linear code over GF(3)

codes are known with d = 32, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-002319 | codetables-gf3-65-6

Minimum distance of a [65, 6] linear code over GF(3)

codes are known with d = 40, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-002320 | codetables-gf3-65-7

Minimum distance of a [65, 7] linear code over GF(3)

codes are known with d = 39, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-002321 | codetables-gf3-65-8

Minimum distance of a [65, 8] linear code over GF(3)

codes are known with d = 37, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-002322 | codetables-gf3-65-9

Minimum distance of a [65, 9] linear code over GF(3)

codes are known with d = 36, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-002323 | codetables-gf3-66-10

Minimum distance of a [66, 10] linear code over GF(3)

codes are known with d = 36, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-002324 | codetables-gf3-66-11

Minimum distance of a [66, 11] linear code over GF(3)

codes are known with d = 33, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-002325 | codetables-gf3-66-12

Minimum distance of a [66, 12] linear code over GF(3)

codes are known with d = 33, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-002326 | codetables-gf3-66-6

Minimum distance of a [66, 6] linear code over GF(3)

codes are known with d = 41, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-002327 | codetables-gf3-66-7

Minimum distance of a [66, 7] linear code over GF(3)

codes are known with d = 39, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-002328 | codetables-gf3-66-8

Minimum distance of a [66, 8] linear code over GF(3)

codes are known with d = 38, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-002329 | codetables-gf3-66-9

Minimum distance of a [66, 9] linear code over GF(3)

codes are known with d = 36, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-002330 | codetables-gf3-67-10

Minimum distance of a [67, 10] linear code over GF(3)

codes are known with d = 36, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-002331 | codetables-gf3-67-11

Minimum distance of a [67, 11] linear code over GF(3)

codes are known with d = 34, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-002332 | codetables-gf3-67-12

Minimum distance of a [67, 12] linear code over GF(3)

codes are known with d = 33, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-002333 | codetables-gf3-67-7

Minimum distance of a [67, 7] linear code over GF(3)

codes are known with d = 39, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-002334 | codetables-gf3-67-8

Minimum distance of a [67, 8] linear code over GF(3)

codes are known with d = 39, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-002335 | codetables-gf3-67-9

Minimum distance of a [67, 9] linear code over GF(3)

codes are known with d = 37, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-002336 | codetables-gf3-68-10

Minimum distance of a [68, 10] linear code over GF(3)

codes are known with d = 36, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-002337 | codetables-gf3-68-11

Minimum distance of a [68, 11] linear code over GF(3)

codes are known with d = 35, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-002338 | codetables-gf3-68-12

Minimum distance of a [68, 12] linear code over GF(3)

codes are known with d = 33, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-002339 | codetables-gf3-68-6

Minimum distance of a [68, 6] linear code over GF(3)

codes are known with d = 42, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-002340 | codetables-gf3-68-7

Minimum distance of a [68, 7] linear code over GF(3)

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-002341 | codetables-gf3-68-8

Minimum distance of a [68, 8] linear code over GF(3)

codes are known with d = 39, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-002342 | codetables-gf3-68-9

Minimum distance of a [68, 9] linear code over GF(3)

codes are known with d = 38, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-002343 | codetables-gf3-69-10

Minimum distance of a [69, 10] linear code over GF(3)

codes are known with d = 36, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-002344 | codetables-gf3-69-11

Minimum distance of a [69, 11] linear code over GF(3)

codes are known with d = 36, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-002345 | codetables-gf3-69-12

Minimum distance of a [69, 12] linear code over GF(3)

codes are known with d = 34, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-002346 | codetables-gf3-69-6

Minimum distance of a [69, 6] linear code over GF(3)

codes are known with d = 42, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-002347 | codetables-gf3-69-7

Minimum distance of a [69, 7] linear code over GF(3)

codes are known with d = 41, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-002348 | codetables-gf3-69-8

Minimum distance of a [69, 8] linear code over GF(3)

codes are known with d = 39, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-002349 | codetables-gf3-69-9

Minimum distance of a [69, 9] linear code over GF(3)

codes are known with d = 39, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-002350 | codetables-gf3-70-10

Minimum distance of a [70, 10] linear code over GF(3)

codes are known with d = 37, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-002351 | codetables-gf3-70-11

Minimum distance of a [70, 11] linear code over GF(3)

codes are known with d = 36, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-002352 | codetables-gf3-70-12

Minimum distance of a [70, 12] linear code over GF(3)

codes are known with d = 35, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-002353 | codetables-gf3-70-6

Minimum distance of a [70, 6] linear code over GF(3)

codes are known with d = 43, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-002354 | codetables-gf3-70-7

Minimum distance of a [70, 7] linear code over GF(3)

codes are known with d = 42, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-002355 | codetables-gf3-70-8

Minimum distance of a [70, 8] linear code over GF(3)

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-002356 | codetables-gf3-70-9

Minimum distance of a [70, 9] linear code over GF(3)

codes are known with d = 39, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-002357 | codetables-gf3-71-10

Minimum distance of a [71, 10] linear code over GF(3)

codes are known with d = 38, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-002358 | codetables-gf3-71-11

Minimum distance of a [71, 11] linear code over GF(3)

codes are known with d = 36, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-002359 | codetables-gf3-71-12

Minimum distance of a [71, 12] linear code over GF(3)

codes are known with d = 36, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-002360 | codetables-gf3-71-6

Minimum distance of a [71, 6] linear code over GF(3)

codes are known with d = 44, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-002361 | codetables-gf3-71-7

Minimum distance of a [71, 7] linear code over GF(3)

codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-002362 | codetables-gf3-71-8

Minimum distance of a [71, 8] linear code over GF(3)

codes are known with d = 41, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-002363 | codetables-gf3-71-9

Minimum distance of a [71, 9] linear code over GF(3)

codes are known with d = 39, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-002364 | codetables-gf3-72-10

Minimum distance of a [72, 10] linear code over GF(3)

codes are known with d = 39, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-002365 | codetables-gf3-72-11

Minimum distance of a [72, 11] linear code over GF(3)

codes are known with d = 37, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-002366 | codetables-gf3-72-12

Minimum distance of a [72, 12] linear code over GF(3)

codes are known with d = 36, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-002367 | codetables-gf3-72-7

Minimum distance of a [72, 7] linear code over GF(3)

codes are known with d = 43, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-002368 | codetables-gf3-72-8

Minimum distance of a [72, 8] linear code over GF(3)

codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-002369 | codetables-gf3-72-9

Minimum distance of a [72, 9] linear code over GF(3)

codes are known with d = 40, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-002370 | codetables-gf3-73-10

Minimum distance of a [73, 10] linear code over GF(3)

codes are known with d = 39, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-002371 | codetables-gf3-73-11

Minimum distance of a [73, 11] linear code over GF(3)

codes are known with d = 38, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-002372 | codetables-gf3-73-12

Minimum distance of a [73, 12] linear code over GF(3)

codes are known with d = 37, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-002373 | codetables-gf3-73-6

Minimum distance of a [73, 6] linear code over GF(3)

codes are known with d = 45, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-002374 | codetables-gf3-73-7

Minimum distance of a [73, 7] linear code over GF(3)

codes are known with d = 44, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-002375 | codetables-gf3-73-8

Minimum distance of a [73, 8] linear code over GF(3)

codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-002376 | codetables-gf3-73-9

Minimum distance of a [73, 9] linear code over GF(3)

codes are known with d = 41, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-002377 | codetables-gf3-74-10

Minimum distance of a [74, 10] linear code over GF(3)

codes are known with d = 40, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-002378 | codetables-gf3-74-11

Minimum distance of a [74, 11] linear code over GF(3)

codes are known with d = 39, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-002379 | codetables-gf3-74-12

Minimum distance of a [74, 12] linear code over GF(3)

codes are known with d = 38, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-002380 | codetables-gf3-74-6

Minimum distance of a [74, 6] linear code over GF(3)

codes are known with d = 46, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-002381 | codetables-gf3-74-8

Minimum distance of a [74, 8] linear code over GF(3)

codes are known with d = 43, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-002382 | codetables-gf3-74-9

Minimum distance of a [74, 9] linear code over GF(3)

codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-002383 | codetables-gf3-75-10

Minimum distance of a [75, 10] linear code over GF(3)

codes are known with d = 41, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-002384 | codetables-gf3-75-11

Minimum distance of a [75, 11] linear code over GF(3)

codes are known with d = 40, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-002385 | codetables-gf3-75-12

Minimum distance of a [75, 12] linear code over GF(3)

codes are known with d = 39, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-002386 | codetables-gf3-75-6

Minimum distance of a [75, 6] linear code over GF(3)

codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-002387 | codetables-gf3-75-7

Minimum distance of a [75, 7] linear code over GF(3)

codes are known with d = 45, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-002388 | codetables-gf3-75-8

Minimum distance of a [75, 8] linear code over GF(3)

codes are known with d = 44, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-002389 | codetables-gf3-75-9

Minimum distance of a [75, 9] linear code over GF(3)

codes are known with d = 42, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-002390 | codetables-gf3-76-10

Minimum distance of a [76, 10] linear code over GF(3)

codes are known with d = 42, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-002391 | codetables-gf3-76-11

Minimum distance of a [76, 11] linear code over GF(3)

codes are known with d = 41, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-002392 | codetables-gf3-76-12

Minimum distance of a [76, 12] linear code over GF(3)

codes are known with d = 40, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-002393 | codetables-gf3-76-7

Minimum distance of a [76, 7] linear code over GF(3)

codes are known with d = 46, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-002394 | codetables-gf3-76-8

Minimum distance of a [76, 8] linear code over GF(3)

codes are known with d = 45, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-002395 | codetables-gf3-76-9

Minimum distance of a [76, 9] linear code over GF(3)

codes are known with d = 43, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-002396 | codetables-gf3-77-10

Minimum distance of a [77, 10] linear code over GF(3)

codes are known with d = 42, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-002397 | codetables-gf3-77-11

Minimum distance of a [77, 11] linear code over GF(3)

codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-002398 | codetables-gf3-77-12

Minimum distance of a [77, 12] linear code over GF(3)

codes are known with d = 41, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-002399 | codetables-gf3-77-6

Minimum distance of a [77, 6] linear code over GF(3)

codes are known with d = 48, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-002400 | codetables-gf3-77-7

Minimum distance of a [77, 7] linear code over GF(3)

codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-002401 | codetables-gf3-77-8

Minimum distance of a [77, 8] linear code over GF(3)

codes are known with d = 45, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-002402 | codetables-gf3-77-9

Minimum distance of a [77, 9] linear code over GF(3)

codes are known with d = 44, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-002403 | codetables-gf3-78-10

Minimum distance of a [78, 10] linear code over GF(3)

codes are known with d = 43, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-002404 | codetables-gf3-78-11

Minimum distance of a [78, 11] linear code over GF(3)

codes are known with d = 42, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-002405 | codetables-gf3-78-12

Minimum distance of a [78, 12] linear code over GF(3)

codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-002406 | codetables-gf3-78-6

Minimum distance of a [78, 6] linear code over GF(3)

codes are known with d = 49, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-002407 | codetables-gf3-78-8

Minimum distance of a [78, 8] linear code over GF(3)

codes are known with d = 46, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-002408 | codetables-gf3-78-9

Minimum distance of a [78, 9] linear code over GF(3)

codes are known with d = 45, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-002409 | codetables-gf3-79-10

Minimum distance of a [79, 10] linear code over GF(3)

codes are known with d = 44, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-002410 | codetables-gf3-79-11

Minimum distance of a [79, 11] linear code over GF(3)

codes are known with d = 43, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-002411 | codetables-gf3-79-12

Minimum distance of a [79, 12] linear code over GF(3)

codes are known with d = 42, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-002412 | codetables-gf3-79-6

Minimum distance of a [79, 6] linear code over GF(3)

codes are known with d = 50, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-002413 | codetables-gf3-79-8

Minimum distance of a [79, 8] linear code over GF(3)

codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-002414 | codetables-gf3-79-9

Minimum distance of a [79, 9] linear code over GF(3)

codes are known with d = 46, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-002415 | codetables-gf3-80-10

Minimum distance of a [80, 10] linear code over GF(3)

codes are known with d = 45, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-002416 | codetables-gf3-80-11

Minimum distance of a [80, 11] linear code over GF(3)

codes are known with d = 44, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-002417 | codetables-gf3-80-12

Minimum distance of a [80, 12] linear code over GF(3)

codes are known with d = 42, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-002418 | codetables-gf3-80-8

Minimum distance of a [80, 8] linear code over GF(3)

codes are known with d = 48, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-002419 | codetables-gf3-80-9

Minimum distance of a [80, 9] linear code over GF(3)

codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-002420 | codetables-gf3-81-10

Minimum distance of a [81, 10] linear code over GF(3)

codes are known with d = 45, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-002421 | codetables-gf3-81-11

Minimum distance of a [81, 11] linear code over GF(3)

codes are known with d = 45, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-002422 | codetables-gf3-81-12

Minimum distance of a [81, 12] linear code over GF(3)

codes are known with d = 42, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-002423 | codetables-gf3-81-8

Minimum distance of a [81, 8] linear code over GF(3)

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-002424 | codetables-gf3-81-9

Minimum distance of a [81, 9] linear code over GF(3)

codes are known with d = 48, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-002425 | codetables-gf3-82-10

Minimum distance of a [82, 10] linear code over GF(3)

codes are known with d = 46, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-002426 | codetables-gf3-82-11

Minimum distance of a [82, 11] linear code over GF(3)

codes are known with d = 45, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-002427 | codetables-gf3-82-12

Minimum distance of a [82, 12] linear code over GF(3)

codes are known with d = 43, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-002428 | codetables-gf3-82-8

Minimum distance of a [82, 8] linear code over GF(3)

codes are known with d = 49, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-002429 | codetables-gf3-82-9

Minimum distance of a [82, 9] linear code over GF(3)

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-002430 | codetables-gf3-83-10

Minimum distance of a [83, 10] linear code over GF(3)

codes are known with d = 47, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-002431 | codetables-gf3-83-11

Minimum distance of a [83, 11] linear code over GF(3)

codes are known with d = 46, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-002432 | codetables-gf3-83-12

Minimum distance of a [83, 12] linear code over GF(3)

codes are known with d = 44, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-002433 | codetables-gf3-83-8

Minimum distance of a [83, 8] linear code over GF(3)

codes are known with d = 50, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-002434 | codetables-gf3-83-9

Minimum distance of a [83, 9] linear code over GF(3)

codes are known with d = 49, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-002435 | codetables-gf3-84-10

Minimum distance of a [84, 10] linear code over GF(3)

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-002436 | codetables-gf3-84-11

Minimum distance of a [84, 11] linear code over GF(3)

codes are known with d = 47, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-002437 | codetables-gf3-84-12

Minimum distance of a [84, 12] linear code over GF(3)

codes are known with d = 45, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-002438 | codetables-gf3-84-8

Minimum distance of a [84, 8] linear code over GF(3)

codes are known with d = 51, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-002439 | codetables-gf3-84-9

Minimum distance of a [84, 9] linear code over GF(3)

codes are known with d = 50, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-002440 | codetables-gf3-85-10

Minimum distance of a [85, 10] linear code over GF(3)

codes are known with d = 48, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-002441 | codetables-gf3-85-11

Minimum distance of a [85, 11] linear code over GF(3)

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-002442 | codetables-gf3-85-12

Minimum distance of a [85, 12] linear code over GF(3)

codes are known with d = 45, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-002443 | codetables-gf3-85-8

Minimum distance of a [85, 8] linear code over GF(3)

codes are known with d = 51, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-002444 | codetables-gf3-85-9

Minimum distance of a [85, 9] linear code over GF(3)

codes are known with d = 51, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-002445 | codetables-gf3-86-10

Minimum distance of a [86, 10] linear code over GF(3)

codes are known with d = 49, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-002446 | codetables-gf3-86-11

Minimum distance of a [86, 11] linear code over GF(3)

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-002447 | codetables-gf3-86-12

Minimum distance of a [86, 12] linear code over GF(3)

codes are known with d = 45, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-002448 | codetables-gf3-86-8

Minimum distance of a [86, 8] linear code over GF(3)

codes are known with d = 52, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-002449 | codetables-gf3-86-9

Minimum distance of a [86, 9] linear code over GF(3)

codes are known with d = 51, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-002450 | codetables-gf3-87-10

Minimum distance of a [87, 10] linear code over GF(3)

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-002451 | codetables-gf3-87-11

Minimum distance of a [87, 11] linear code over GF(3)

codes are known with d = 49, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-002452 | codetables-gf3-87-12

Minimum distance of a [87, 12] linear code over GF(3)

codes are known with d = 46, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-002453 | codetables-gf3-87-6

Minimum distance of a [87, 6] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-002454 | codetables-gf3-87-8

Minimum distance of a [87, 8] linear code over GF(3)

codes are known with d = 53, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-002455 | codetables-gf3-87-9

Minimum distance of a [87, 9] linear code over GF(3)

codes are known with d = 52, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-002456 | codetables-gf3-88-10

Minimum distance of a [88, 10] linear code over GF(3)

codes are known with d = 51, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-002457 | codetables-gf3-88-11

Minimum distance of a [88, 11] linear code over GF(3)

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-002458 | codetables-gf3-88-12

Minimum distance of a [88, 12] linear code over GF(3)

codes are known with d = 47, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-002459 | codetables-gf3-88-6

Minimum distance of a [88, 6] linear code over GF(3)

codes are known with d = 55, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-002460 | codetables-gf3-88-7

Minimum distance of a [88, 7] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-002461 | codetables-gf3-88-9

Minimum distance of a [88, 9] linear code over GF(3)

codes are known with d = 53, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-002462 | codetables-gf3-89-10

Minimum distance of a [89, 10] linear code over GF(3)

codes are known with d = 51, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-002463 | codetables-gf3-89-11

Minimum distance of a [89, 11] linear code over GF(3)

codes are known with d = 51, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-002464 | codetables-gf3-89-12

Minimum distance of a [89, 12] linear code over GF(3)

codes are known with d = 47, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-002465 | codetables-gf3-89-6

Minimum distance of a [89, 6] linear code over GF(3)

codes are known with d = 56, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-002466 | codetables-gf3-89-7

Minimum distance of a [89, 7] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-002467 | codetables-gf3-89-8

Minimum distance of a [89, 8] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-002468 | codetables-gf3-90-10

Minimum distance of a [90, 10] linear code over GF(3)

codes are known with d = 51, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-002469 | codetables-gf3-90-11

Minimum distance of a [90, 11] linear code over GF(3)

codes are known with d = 51, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-002470 | codetables-gf3-90-12

Minimum distance of a [90, 12] linear code over GF(3)

codes are known with d = 48, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-002471 | codetables-gf3-90-7

Minimum distance of a [90, 7] linear code over GF(3)

codes are known with d = 55, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-002472 | codetables-gf3-90-8

Minimum distance of a [90, 8] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-002473 | codetables-gf3-91-10

Minimum distance of a [91, 10] linear code over GF(3)

codes are known with d = 52, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-002474 | codetables-gf3-91-11

Minimum distance of a [91, 11] linear code over GF(3)

codes are known with d = 51, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-002475 | codetables-gf3-91-12

Minimum distance of a [91, 12] linear code over GF(3)

codes are known with d = 48, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-002476 | codetables-gf3-91-6

Minimum distance of a [91, 6] linear code over GF(3)

codes are known with d = 57, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-002477 | codetables-gf3-91-7

Minimum distance of a [91, 7] linear code over GF(3)

codes are known with d = 56, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-002478 | codetables-gf3-91-8

Minimum distance of a [91, 8] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-002479 | codetables-gf3-91-9

Minimum distance of a [91, 9] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-002480 | codetables-gf3-92-10

Minimum distance of a [92, 10] linear code over GF(3)

codes are known with d = 53, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-002481 | codetables-gf3-92-11

Minimum distance of a [92, 11] linear code over GF(3)

codes are known with d = 52, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-002482 | codetables-gf3-92-12

Minimum distance of a [92, 12] linear code over GF(3)

codes are known with d = 48, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-002483 | codetables-gf3-92-6

Minimum distance of a [92, 6] linear code over GF(3)

codes are known with d = 58, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-002484 | codetables-gf3-92-7

Minimum distance of a [92, 7] linear code over GF(3)

codes are known with d = 57, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-002485 | codetables-gf3-92-8

Minimum distance of a [92, 8] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-002486 | codetables-gf3-92-9

Minimum distance of a [92, 9] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-002487 | codetables-gf3-93-10

Minimum distance of a [93, 10] linear code over GF(3)

codes are known with d = 53, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-002488 | codetables-gf3-93-11

Minimum distance of a [93, 11] linear code over GF(3)

codes are known with d = 53, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-002489 | codetables-gf3-93-12

Minimum distance of a [93, 12] linear code over GF(3)

codes are known with d = 49, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-002490 | codetables-gf3-93-6

Minimum distance of a [93, 6] linear code over GF(3)

codes are known with d = 59, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-002491 | codetables-gf3-93-7

Minimum distance of a [93, 7] linear code over GF(3)

codes are known with d = 57, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-002492 | codetables-gf3-93-8

Minimum distance of a [93, 8] linear code over GF(3)

codes are known with d = 55, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-002493 | codetables-gf3-93-9

Minimum distance of a [93, 9] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-002494 | codetables-gf3-94-10

Minimum distance of a [94, 10] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-002495 | codetables-gf3-94-11

Minimum distance of a [94, 11] linear code over GF(3)

codes are known with d = 53, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-002496 | codetables-gf3-94-12

Minimum distance of a [94, 12] linear code over GF(3)

codes are known with d = 50, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-002497 | codetables-gf3-94-7

Minimum distance of a [94, 7] linear code over GF(3)

codes are known with d = 58, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-002498 | codetables-gf3-94-8

Minimum distance of a [94, 8] linear code over GF(3)

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-002499 | codetables-gf3-94-9

Minimum distance of a [94, 9] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-002500 | codetables-gf3-95-10

Minimum distance of a [95, 10] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-002501 | codetables-gf3-95-11

Minimum distance of a [95, 11] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-002502 | codetables-gf3-95-12

Minimum distance of a [95, 12] linear code over GF(3)

codes are known with d = 51, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-002503 | codetables-gf3-95-6

Minimum distance of a [95, 6] linear code over GF(3)

codes are known with d = 60, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-002504 | codetables-gf3-95-7

Minimum distance of a [95, 7] linear code over GF(3)

codes are known with d = 59, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-002505 | codetables-gf3-95-8

Minimum distance of a [95, 8] linear code over GF(3)

codes are known with d = 57, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-002506 | codetables-gf3-95-9

Minimum distance of a [95, 9] linear code over GF(3)

codes are known with d = 55, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-002507 | codetables-gf3-96-10

Minimum distance of a [96, 10] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-002508 | codetables-gf3-96-11

Minimum distance of a [96, 11] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-002509 | codetables-gf3-96-12

Minimum distance of a [96, 12] linear code over GF(3)

codes are known with d = 51, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-002510 | codetables-gf3-96-8

Minimum distance of a [96, 8] linear code over GF(3)

codes are known with d = 57, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-002511 | codetables-gf3-96-9

Minimum distance of a [96, 9] linear code over GF(3)

codes are known with d = 56, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-002512 | codetables-gf3-97-10

Minimum distance of a [97, 10] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-002513 | codetables-gf3-97-11

Minimum distance of a [97, 11] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-002514 | codetables-gf3-97-12

Minimum distance of a [97, 12] linear code over GF(3)

codes are known with d = 51, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-002515 | codetables-gf3-97-7

Minimum distance of a [97, 7] linear code over GF(3)

codes are known with d = 60, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-002516 | codetables-gf3-97-8

Minimum distance of a [97, 8] linear code over GF(3)

codes are known with d = 57, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-002517 | codetables-gf3-97-9

Minimum distance of a [97, 9] linear code over GF(3)

codes are known with d = 56, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-002518 | codetables-gf3-98-10

Minimum distance of a [98, 10] linear code over GF(3)

codes are known with d = 55, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-002519 | codetables-gf3-98-11

Minimum distance of a [98, 11] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-002520 | codetables-gf3-98-12

Minimum distance of a [98, 12] linear code over GF(3)

codes are known with d = 51, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-002521 | codetables-gf3-98-7

Minimum distance of a [98, 7] linear code over GF(3)

codes are known with d = 61, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-002522 | codetables-gf3-98-8

Minimum distance of a [98, 8] linear code over GF(3)

codes are known with d = 58, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-002523 | codetables-gf3-98-9

Minimum distance of a [98, 9] linear code over GF(3)

codes are known with d = 57, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-002524 | codetables-gf3-99-10

Minimum distance of a [99, 10] linear code over GF(3)

codes are known with d = 56, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-002525 | codetables-gf3-99-11

Minimum distance of a [99, 11] linear code over GF(3)

codes are known with d = 54, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-002526 | codetables-gf3-99-12

Minimum distance of a [99, 12] linear code over GF(3)

codes are known with d = 52, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-002527 | codetables-gf3-99-7

Minimum distance of a [99, 7] linear code over GF(3)

codes are known with d = 62, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-002528 | codetables-gf3-99-8

Minimum distance of a [99, 8] linear code over GF(3)

codes are known with d = 59, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-002529 | codetables-gf3-99-9

Minimum distance of a [99, 9] linear code over GF(3)

codes are known with d = 57, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-002530 | codetables-gf4-100-5

Minimum distance of a [100, 5] linear code over GF(4)

codes are known with d = 72, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-002531 | codetables-gf4-100-6

Minimum distance of a [100, 6] linear code over GF(4)

codes are known with d = 70, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-002532 | codetables-gf4-100-7

Minimum distance of a [100, 7] linear code over GF(4)

codes are known with d = 68, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-002533 | codetables-gf4-100-8

Minimum distance of a [100, 8] linear code over GF(4)

codes are known with d = 66, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-002534 | codetables-gf4-100-9

Minimum distance of a [100, 9] linear code over GF(4)

codes are known with d = 64, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-002535 | codetables-gf4-101-5

Minimum distance of a [101, 5] linear code over GF(4)

codes are known with d = 73, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-002536 | codetables-gf4-101-6

Minimum distance of a [101, 6] linear code over GF(4)

codes are known with d = 71, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-002537 | codetables-gf4-101-7

Minimum distance of a [101, 7] linear code over GF(4)

codes are known with d = 68, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-002538 | codetables-gf4-101-8

Minimum distance of a [101, 8] linear code over GF(4)

codes are known with d = 67, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-002539 | codetables-gf4-101-9

Minimum distance of a [101, 9] linear code over GF(4)

codes are known with d = 64, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-002540 | codetables-gf4-102-6

Minimum distance of a [102, 6] linear code over GF(4)

codes are known with d = 72, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-002541 | codetables-gf4-102-7

Minimum distance of a [102, 7] linear code over GF(4)

codes are known with d = 69, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-002542 | codetables-gf4-102-8

Minimum distance of a [102, 8] linear code over GF(4)

codes are known with d = 68, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-002543 | codetables-gf4-102-9

Minimum distance of a [102, 9] linear code over GF(4)

codes are known with d = 65, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-002544 | codetables-gf4-103-5

Minimum distance of a [103, 5] linear code over GF(4)

codes are known with d = 74, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-002545 | codetables-gf4-103-6

Minimum distance of a [103, 6] linear code over GF(4)

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-002546 | codetables-gf4-103-7

Minimum distance of a [103, 7] linear code over GF(4)

codes are known with d = 70, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-002547 | codetables-gf4-103-8

Minimum distance of a [103, 8] linear code over GF(4)

codes are known with d = 68, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-002548 | codetables-gf4-103-9

Minimum distance of a [103, 9] linear code over GF(4)

codes are known with d = 66, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-002549 | codetables-gf4-104-5

Minimum distance of a [104, 5] linear code over GF(4)

codes are known with d = 75, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-002550 | codetables-gf4-104-6

Minimum distance of a [104, 6] linear code over GF(4)

codes are known with d = 73, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-002551 | codetables-gf4-104-7

Minimum distance of a [104, 7] linear code over GF(4)

codes are known with d = 71, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-002552 | codetables-gf4-104-8

Minimum distance of a [104, 8] linear code over GF(4)

codes are known with d = 68, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-002553 | codetables-gf4-104-9

Minimum distance of a [104, 9] linear code over GF(4)

codes are known with d = 67, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-002554 | codetables-gf4-105-6

Minimum distance of a [105, 6] linear code over GF(4)

codes are known with d = 74, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-002555 | codetables-gf4-105-7

Minimum distance of a [105, 7] linear code over GF(4)

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-002556 | codetables-gf4-105-8

Minimum distance of a [105, 8] linear code over GF(4)

codes are known with d = 69, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-002557 | codetables-gf4-105-9

Minimum distance of a [105, 9] linear code over GF(4)

codes are known with d = 68, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-002558 | codetables-gf4-106-6

Minimum distance of a [106, 6] linear code over GF(4)

codes are known with d = 75, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-002559 | codetables-gf4-106-7

Minimum distance of a [106, 7] linear code over GF(4)

codes are known with d = 72, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-002560 | codetables-gf4-106-8

Minimum distance of a [106, 8] linear code over GF(4)

codes are known with d = 70, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-002561 | codetables-gf4-106-9

Minimum distance of a [106, 9] linear code over GF(4)

codes are known with d = 68, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-002562 | codetables-gf4-107-6

Minimum distance of a [107, 6] linear code over GF(4)

codes are known with d = 76, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-002563 | codetables-gf4-107-7

Minimum distance of a [107, 7] linear code over GF(4)

codes are known with d = 73, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-002564 | codetables-gf4-107-8

Minimum distance of a [107, 8] linear code over GF(4)

codes are known with d = 71, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-002565 | codetables-gf4-107-9

Minimum distance of a [107, 9] linear code over GF(4)

codes are known with d = 68, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-002566 | codetables-gf4-108-6

Minimum distance of a [108, 6] linear code over GF(4)

codes are known with d = 76, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-002567 | codetables-gf4-108-7

Minimum distance of a [108, 7] linear code over GF(4)

codes are known with d = 74, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-002568 | codetables-gf4-108-8

Minimum distance of a [108, 8] linear code over GF(4)

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-002569 | codetables-gf4-108-9

Minimum distance of a [108, 9] linear code over GF(4)

codes are known with d = 69, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-002570 | codetables-gf4-109-6

Minimum distance of a [109, 6] linear code over GF(4)

codes are known with d = 77, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-002571 | codetables-gf4-109-7

Minimum distance of a [109, 7] linear code over GF(4)

codes are known with d = 75, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-002572 | codetables-gf4-109-8

Minimum distance of a [109, 8] linear code over GF(4)

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-002573 | codetables-gf4-109-9

Minimum distance of a [109, 9] linear code over GF(4)

codes are known with d = 70, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-002574 | codetables-gf4-110-6

Minimum distance of a [110, 6] linear code over GF(4)

codes are known with d = 78, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-002575 | codetables-gf4-110-7

Minimum distance of a [110, 7] linear code over GF(4)

codes are known with d = 76, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-002576 | codetables-gf4-110-8

Minimum distance of a [110, 8] linear code over GF(4)

codes are known with d = 73, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-002577 | codetables-gf4-110-9

Minimum distance of a [110, 9] linear code over GF(4)

codes are known with d = 71, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-002578 | codetables-gf4-111-5

Minimum distance of a [111, 5] linear code over GF(4)

codes are known with d = 80, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-002579 | codetables-gf4-111-6

Minimum distance of a [111, 6] linear code over GF(4)

codes are known with d = 79, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-002580 | codetables-gf4-111-7

Minimum distance of a [111, 7] linear code over GF(4)

codes are known with d = 76, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-002581 | codetables-gf4-111-8

Minimum distance of a [111, 8] linear code over GF(4)

codes are known with d = 74, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-002582 | codetables-gf4-111-9

Minimum distance of a [111, 9] linear code over GF(4)

codes are known with d = 72, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-002583 | codetables-gf4-112-5

Minimum distance of a [112, 5] linear code over GF(4)

codes are known with d = 81, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-002584 | codetables-gf4-112-7

Minimum distance of a [112, 7] linear code over GF(4)

codes are known with d = 76, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-002585 | codetables-gf4-112-8

Minimum distance of a [112, 8] linear code over GF(4)

codes are known with d = 75, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-002586 | codetables-gf4-112-9

Minimum distance of a [112, 9] linear code over GF(4)

codes are known with d = 72, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-002587 | codetables-gf4-113-5

Minimum distance of a [113, 5] linear code over GF(4)

codes are known with d = 82, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-002588 | codetables-gf4-113-6

Minimum distance of a [113, 6] linear code over GF(4)

codes are known with d = 80, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-002589 | codetables-gf4-113-7

Minimum distance of a [113, 7] linear code over GF(4)

codes are known with d = 77, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-002590 | codetables-gf4-113-8

Minimum distance of a [113, 8] linear code over GF(4)

codes are known with d = 76, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-002591 | codetables-gf4-113-9

Minimum distance of a [113, 9] linear code over GF(4)

codes are known with d = 73, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-002592 | codetables-gf4-114-5

Minimum distance of a [114, 5] linear code over GF(4)

codes are known with d = 83, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-002593 | codetables-gf4-114-6

Minimum distance of a [114, 6] linear code over GF(4)

codes are known with d = 81, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-002594 | codetables-gf4-114-7

Minimum distance of a [114, 7] linear code over GF(4)

codes are known with d = 78, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-002595 | codetables-gf4-114-8

Minimum distance of a [114, 8] linear code over GF(4)

codes are known with d = 76, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-002596 | codetables-gf4-114-9

Minimum distance of a [114, 9] linear code over GF(4)

codes are known with d = 74, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-002597 | codetables-gf4-115-6

Minimum distance of a [115, 6] linear code over GF(4)

codes are known with d = 82, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-002598 | codetables-gf4-115-7

Minimum distance of a [115, 7] linear code over GF(4)

codes are known with d = 79, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-002599 | codetables-gf4-115-8

Minimum distance of a [115, 8] linear code over GF(4)

codes are known with d = 76, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-002600 | codetables-gf4-115-9

Minimum distance of a [115, 9] linear code over GF(4)

codes are known with d = 75, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-002601 | codetables-gf4-116-6

Minimum distance of a [116, 6] linear code over GF(4)

codes are known with d = 83, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-002602 | codetables-gf4-116-7

Minimum distance of a [116, 7] linear code over GF(4)

codes are known with d = 80, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-002603 | codetables-gf4-116-8

Minimum distance of a [116, 8] linear code over GF(4)

codes are known with d = 77, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-002604 | codetables-gf4-116-9

Minimum distance of a [116, 9] linear code over GF(4)

codes are known with d = 76, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-002605 | codetables-gf4-117-7

Minimum distance of a [117, 7] linear code over GF(4)

codes are known with d = 80, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-002606 | codetables-gf4-117-8

Minimum distance of a [117, 8] linear code over GF(4)

codes are known with d = 78, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-002607 | codetables-gf4-117-9

Minimum distance of a [117, 9] linear code over GF(4)

codes are known with d = 77, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-002608 | codetables-gf4-118-6

Minimum distance of a [118, 6] linear code over GF(4)

codes are known with d = 84, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-002609 | codetables-gf4-118-7

Minimum distance of a [118, 7] linear code over GF(4)

codes are known with d = 81, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-002610 | codetables-gf4-118-8

Minimum distance of a [118, 8] linear code over GF(4)

codes are known with d = 79, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-002611 | codetables-gf4-118-9

Minimum distance of a [118, 9] linear code over GF(4)

codes are known with d = 78, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-002612 | codetables-gf4-119-6

Minimum distance of a [119, 6] linear code over GF(4)

codes are known with d = 85, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-002613 | codetables-gf4-119-7

Minimum distance of a [119, 7] linear code over GF(4)

codes are known with d = 82, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-002614 | codetables-gf4-119-8

Minimum distance of a [119, 8] linear code over GF(4)

codes are known with d = 80, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-002615 | codetables-gf4-119-9

Minimum distance of a [119, 9] linear code over GF(4)

codes are known with d = 79, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-002616 | codetables-gf4-120-6

Minimum distance of a [120, 6] linear code over GF(4)

codes are known with d = 86, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-002617 | codetables-gf4-120-7

Minimum distance of a [120, 7] linear code over GF(4)

codes are known with d = 83, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-002618 | codetables-gf4-120-8

Minimum distance of a [120, 8] linear code over GF(4)

codes are known with d = 80, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-002619 | codetables-gf4-120-9

Minimum distance of a [120, 9] linear code over GF(4)

codes are known with d = 80, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-002620 | codetables-gf4-121-6

Minimum distance of a [121, 6] linear code over GF(4)

codes are known with d = 87, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-002621 | codetables-gf4-121-7

Minimum distance of a [121, 7] linear code over GF(4)

codes are known with d = 84, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-002622 | codetables-gf4-121-8

Minimum distance of a [121, 8] linear code over GF(4)

codes are known with d = 81, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-002623 | codetables-gf4-121-9

Minimum distance of a [121, 9] linear code over GF(4)

codes are known with d = 80, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-002624 | codetables-gf4-122-5

Minimum distance of a [122, 5] linear code over GF(4)

codes are known with d = 88, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-002625 | codetables-gf4-122-7

Minimum distance of a [122, 7] linear code over GF(4)

codes are known with d = 84, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-002626 | codetables-gf4-122-8

Minimum distance of a [122, 8] linear code over GF(4)

codes are known with d = 82, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-002627 | codetables-gf4-122-9

Minimum distance of a [122, 9] linear code over GF(4)

codes are known with d = 80, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-002628 | codetables-gf4-123-5

Minimum distance of a [123, 5] linear code over GF(4)

codes are known with d = 89, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-002629 | codetables-gf4-123-6

Minimum distance of a [123, 6] linear code over GF(4)

codes are known with d = 88, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-002630 | codetables-gf4-123-7

Minimum distance of a [123, 7] linear code over GF(4)

codes are known with d = 85, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-002631 | codetables-gf4-123-8

Minimum distance of a [123, 8] linear code over GF(4)

codes are known with d = 83, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-002632 | codetables-gf4-123-9

Minimum distance of a [123, 9] linear code over GF(4)

codes are known with d = 81, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-002633 | codetables-gf4-124-5

Minimum distance of a [124, 5] linear code over GF(4)

codes are known with d = 90, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-002634 | codetables-gf4-124-6

Minimum distance of a [124, 6] linear code over GF(4)

codes are known with d = 88, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-002635 | codetables-gf4-124-7

Minimum distance of a [124, 7] linear code over GF(4)

codes are known with d = 86, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-002636 | codetables-gf4-124-8

Minimum distance of a [124, 8] linear code over GF(4)

codes are known with d = 84, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-002637 | codetables-gf4-124-9

Minimum distance of a [124, 9] linear code over GF(4)

codes are known with d = 82, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-002638 | codetables-gf4-125-5

Minimum distance of a [125, 5] linear code over GF(4)

codes are known with d = 91, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-002639 | codetables-gf4-125-6

Minimum distance of a [125, 6] linear code over GF(4)

codes are known with d = 89, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-002640 | codetables-gf4-125-7

Minimum distance of a [125, 7] linear code over GF(4)

codes are known with d = 87, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-002641 | codetables-gf4-125-8

Minimum distance of a [125, 8] linear code over GF(4)

codes are known with d = 85, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-002642 | codetables-gf4-125-9

Minimum distance of a [125, 9] linear code over GF(4)

codes are known with d = 83, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-002643 | codetables-gf4-126-6

Minimum distance of a [126, 6] linear code over GF(4)

codes are known with d = 90, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-002644 | codetables-gf4-126-7

Minimum distance of a [126, 7] linear code over GF(4)

codes are known with d = 88, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-002645 | codetables-gf4-126-8

Minimum distance of a [126, 8] linear code over GF(4)

codes are known with d = 86, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-002646 | codetables-gf4-126-9

Minimum distance of a [126, 9] linear code over GF(4)

codes are known with d = 84, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-002647 | codetables-gf4-127-5

Minimum distance of a [127, 5] linear code over GF(4)

codes are known with d = 92, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-002648 | codetables-gf4-127-6

Minimum distance of a [127, 6] linear code over GF(4)

codes are known with d = 91, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-002649 | codetables-gf4-127-7

Minimum distance of a [127, 7] linear code over GF(4)

codes are known with d = 88, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-002650 | codetables-gf4-127-8

Minimum distance of a [127, 8] linear code over GF(4)

codes are known with d = 87, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-002651 | codetables-gf4-127-9

Minimum distance of a [127, 9] linear code over GF(4)

codes are known with d = 84, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-002652 | codetables-gf4-128-5

Minimum distance of a [128, 5] linear code over GF(4)

codes are known with d = 93, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-002653 | codetables-gf4-128-6

Minimum distance of a [128, 6] linear code over GF(4)

codes are known with d = 92, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-002654 | codetables-gf4-128-7

Minimum distance of a [128, 7] linear code over GF(4)

codes are known with d = 88, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-002655 | codetables-gf4-128-8

Minimum distance of a [128, 8] linear code over GF(4)

codes are known with d = 88, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-002656 | codetables-gf4-128-9

Minimum distance of a [128, 9] linear code over GF(4)

codes are known with d = 85, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-002657 | codetables-gf4-129-5

Minimum distance of a [129, 5] linear code over GF(4)

codes are known with d = 94, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-002658 | codetables-gf4-129-6

Minimum distance of a [129, 6] linear code over GF(4)

codes are known with d = 93, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-002659 | codetables-gf4-129-7

Minimum distance of a [129, 7] linear code over GF(4)

codes are known with d = 89, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-002660 | codetables-gf4-129-8

Minimum distance of a [129, 8] linear code over GF(4)

codes are known with d = 88, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-002661 | codetables-gf4-129-9

Minimum distance of a [129, 9] linear code over GF(4)

codes are known with d = 86, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-002662 | codetables-gf4-130-5

Minimum distance of a [130, 5] linear code over GF(4)

codes are known with d = 95, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-002663 | codetables-gf4-130-6

Minimum distance of a [130, 6] linear code over GF(4)

codes are known with d = 94, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-002664 | codetables-gf4-130-7

Minimum distance of a [130, 7] linear code over GF(4)

codes are known with d = 90, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-002665 | codetables-gf4-130-8

Minimum distance of a [130, 8] linear code over GF(4)

codes are known with d = 88, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-002666 | codetables-gf4-130-9

Minimum distance of a [130, 9] linear code over GF(4)

codes are known with d = 86, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-002667 | codetables-gf4-131-6

Minimum distance of a [131, 6] linear code over GF(4)

codes are known with d = 95, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-002668 | codetables-gf4-131-7

Minimum distance of a [131, 7] linear code over GF(4)

codes are known with d = 91, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-002669 | codetables-gf4-131-8

Minimum distance of a [131, 8] linear code over GF(4)

codes are known with d = 89, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-002670 | codetables-gf4-131-9

Minimum distance of a [131, 9] linear code over GF(4)

codes are known with d = 87, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-002671 | codetables-gf4-132-7

Minimum distance of a [132, 7] linear code over GF(4)

codes are known with d = 92, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-002672 | codetables-gf4-132-8

Minimum distance of a [132, 8] linear code over GF(4)

codes are known with d = 90, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-002673 | codetables-gf4-132-9

Minimum distance of a [132, 9] linear code over GF(4)

codes are known with d = 88, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-002674 | codetables-gf4-133-5

Minimum distance of a [133, 5] linear code over GF(4)

codes are known with d = 96, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-002675 | codetables-gf4-133-7

Minimum distance of a [133, 7] linear code over GF(4)

codes are known with d = 92, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-002676 | codetables-gf4-133-8

Minimum distance of a [133, 8] linear code over GF(4)

codes are known with d = 91, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-002677 | codetables-gf4-133-9

Minimum distance of a [133, 9] linear code over GF(4)

codes are known with d = 88, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-002678 | codetables-gf4-134-5

Minimum distance of a [134, 5] linear code over GF(4)

codes are known with d = 97, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-002679 | codetables-gf4-134-6

Minimum distance of a [134, 6] linear code over GF(4)

codes are known with d = 96, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-002680 | codetables-gf4-134-7

Minimum distance of a [134, 7] linear code over GF(4)

codes are known with d = 93, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-002681 | codetables-gf4-134-8

Minimum distance of a [134, 8] linear code over GF(4)

codes are known with d = 92, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-002682 | codetables-gf4-134-9

Minimum distance of a [134, 9] linear code over GF(4)

codes are known with d = 88, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-002683 | codetables-gf4-135-5

Minimum distance of a [135, 5] linear code over GF(4)

codes are known with d = 98, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-002684 | codetables-gf4-135-6

Minimum distance of a [135, 6] linear code over GF(4)

codes are known with d = 96, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-002685 | codetables-gf4-135-7

Minimum distance of a [135, 7] linear code over GF(4)

codes are known with d = 94, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-002686 | codetables-gf4-135-8

Minimum distance of a [135, 8] linear code over GF(4)

codes are known with d = 92, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-002687 | codetables-gf4-135-9

Minimum distance of a [135, 9] linear code over GF(4)

codes are known with d = 88, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-002688 | codetables-gf4-136-5

Minimum distance of a [136, 5] linear code over GF(4)

codes are known with d = 99, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-002689 | codetables-gf4-136-6

Minimum distance of a [136, 6] linear code over GF(4)

codes are known with d = 97, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-002690 | codetables-gf4-136-7

Minimum distance of a [136, 7] linear code over GF(4)

codes are known with d = 94, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-002691 | codetables-gf4-136-8

Minimum distance of a [136, 8] linear code over GF(4)

codes are known with d = 93, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-002692 | codetables-gf4-136-9

Minimum distance of a [136, 9] linear code over GF(4)

codes are known with d = 89, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-002693 | codetables-gf4-137-6

Minimum distance of a [137, 6] linear code over GF(4)

codes are known with d = 98, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-002694 | codetables-gf4-137-7

Minimum distance of a [137, 7] linear code over GF(4)

codes are known with d = 95, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-002695 | codetables-gf4-137-8

Minimum distance of a [137, 8] linear code over GF(4)

codes are known with d = 94, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-002696 | codetables-gf4-137-9

Minimum distance of a [137, 9] linear code over GF(4)

codes are known with d = 90, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-002697 | codetables-gf4-138-5

Minimum distance of a [138, 5] linear code over GF(4)

codes are known with d = 100, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-002698 | codetables-gf4-138-6

Minimum distance of a [138, 6] linear code over GF(4)

codes are known with d = 99, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-002699 | codetables-gf4-138-7

Minimum distance of a [138, 7] linear code over GF(4)

codes are known with d = 96, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-002700 | codetables-gf4-138-8

Minimum distance of a [138, 8] linear code over GF(4)

codes are known with d = 94, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-002701 | codetables-gf4-138-9

Minimum distance of a [138, 9] linear code over GF(4)

codes are known with d = 91, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-002702 | codetables-gf4-139-5

Minimum distance of a [139, 5] linear code over GF(4)

codes are known with d = 101, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-002703 | codetables-gf4-139-6

Minimum distance of a [139, 6] linear code over GF(4)

codes are known with d = 100, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-002704 | codetables-gf4-139-7

Minimum distance of a [139, 7] linear code over GF(4)

codes are known with d = 96, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-002705 | codetables-gf4-139-8

Minimum distance of a [139, 8] linear code over GF(4)

codes are known with d = 95, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-002706 | codetables-gf4-139-9

Minimum distance of a [139, 9] linear code over GF(4)

codes are known with d = 92, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-002707 | codetables-gf4-140-5

Minimum distance of a [140, 5] linear code over GF(4)

codes are known with d = 102, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-002708 | codetables-gf4-140-6

Minimum distance of a [140, 6] linear code over GF(4)

codes are known with d = 100, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-002709 | codetables-gf4-140-7

Minimum distance of a [140, 7] linear code over GF(4)

codes are known with d = 97, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-002710 | codetables-gf4-140-8

Minimum distance of a [140, 8] linear code over GF(4)

codes are known with d = 96, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-002711 | codetables-gf4-140-9

Minimum distance of a [140, 9] linear code over GF(4)

codes are known with d = 92, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-002712 | codetables-gf4-141-5

Minimum distance of a [141, 5] linear code over GF(4)

codes are known with d = 103, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-002713 | codetables-gf4-141-6

Minimum distance of a [141, 6] linear code over GF(4)

codes are known with d = 101, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-002714 | codetables-gf4-141-7

Minimum distance of a [141, 7] linear code over GF(4)

codes are known with d = 98, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-002715 | codetables-gf4-141-8

Minimum distance of a [141, 8] linear code over GF(4)

codes are known with d = 96, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-002716 | codetables-gf4-141-9

Minimum distance of a [141, 9] linear code over GF(4)

codes are known with d = 93, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-002717 | codetables-gf4-142-6

Minimum distance of a [142, 6] linear code over GF(4)

codes are known with d = 102, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-002718 | codetables-gf4-142-7

Minimum distance of a [142, 7] linear code over GF(4)

codes are known with d = 99, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-002719 | codetables-gf4-142-8

Minimum distance of a [142, 8] linear code over GF(4)

codes are known with d = 97, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-002720 | codetables-gf4-142-9

Minimum distance of a [142, 9] linear code over GF(4)

codes are known with d = 94, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-002721 | codetables-gf4-143-5

Minimum distance of a [143, 5] linear code over GF(4)

codes are known with d = 104, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-002722 | codetables-gf4-143-6

Minimum distance of a [143, 6] linear code over GF(4)

codes are known with d = 103, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-002723 | codetables-gf4-143-7

Minimum distance of a [143, 7] linear code over GF(4)

codes are known with d = 100, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-002724 | codetables-gf4-143-8

Minimum distance of a [143, 8] linear code over GF(4)

codes are known with d = 98, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-002725 | codetables-gf4-143-9

Minimum distance of a [143, 9] linear code over GF(4)

codes are known with d = 95, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-002726 | codetables-gf4-144-5

Minimum distance of a [144, 5] linear code over GF(4)

codes are known with d = 105, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-002727 | codetables-gf4-144-6

Minimum distance of a [144, 6] linear code over GF(4)

codes are known with d = 104, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-002728 | codetables-gf4-144-7

Minimum distance of a [144, 7] linear code over GF(4)

codes are known with d = 100, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-002729 | codetables-gf4-144-8

Minimum distance of a [144, 8] linear code over GF(4)

codes are known with d = 99, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-002730 | codetables-gf4-144-9

Minimum distance of a [144, 9] linear code over GF(4)

codes are known with d = 96, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-002731 | codetables-gf4-145-5

Minimum distance of a [145, 5] linear code over GF(4)

codes are known with d = 106, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-002732 | codetables-gf4-145-6

Minimum distance of a [145, 6] linear code over GF(4)

codes are known with d = 104, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-002733 | codetables-gf4-145-7

Minimum distance of a [145, 7] linear code over GF(4)

codes are known with d = 100, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-002734 | codetables-gf4-145-8

Minimum distance of a [145, 8] linear code over GF(4)

codes are known with d = 100, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-002735 | codetables-gf4-145-9

Minimum distance of a [145, 9] linear code over GF(4)

codes are known with d = 96, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-002736 | codetables-gf4-146-5

Minimum distance of a [146, 5] linear code over GF(4)

codes are known with d = 107, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-002737 | codetables-gf4-146-6

Minimum distance of a [146, 6] linear code over GF(4)

codes are known with d = 105, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-002738 | codetables-gf4-146-7

Minimum distance of a [146, 7] linear code over GF(4)

codes are known with d = 101, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-002739 | codetables-gf4-146-8

Minimum distance of a [146, 8] linear code over GF(4)

codes are known with d = 100, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-002740 | codetables-gf4-146-9

Minimum distance of a [146, 9] linear code over GF(4)

codes are known with d = 96, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-002741 | codetables-gf4-147-6

Minimum distance of a [147, 6] linear code over GF(4)

codes are known with d = 106, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-002742 | codetables-gf4-147-7

Minimum distance of a [147, 7] linear code over GF(4)

codes are known with d = 102, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-002743 | codetables-gf4-147-8

Minimum distance of a [147, 8] linear code over GF(4)

codes are known with d = 100, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-002744 | codetables-gf4-147-9

Minimum distance of a [147, 9] linear code over GF(4)

codes are known with d = 97, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-002745 | codetables-gf4-148-5

Minimum distance of a [148, 5] linear code over GF(4)

codes are known with d = 108, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-002746 | codetables-gf4-148-6

Minimum distance of a [148, 6] linear code over GF(4)

codes are known with d = 106, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-002747 | codetables-gf4-148-7

Minimum distance of a [148, 7] linear code over GF(4)

codes are known with d = 103, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-002748 | codetables-gf4-148-8

Minimum distance of a [148, 8] linear code over GF(4)

codes are known with d = 100, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-002749 | codetables-gf4-148-9

Minimum distance of a [148, 9] linear code over GF(4)

codes are known with d = 98, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-002750 | codetables-gf4-149-5

Minimum distance of a [149, 5] linear code over GF(4)

codes are known with d = 109, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-002751 | codetables-gf4-149-6

Minimum distance of a [149, 6] linear code over GF(4)

codes are known with d = 107, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-002752 | codetables-gf4-149-7

Minimum distance of a [149, 7] linear code over GF(4)

codes are known with d = 104, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-002753 | codetables-gf4-149-8

Minimum distance of a [149, 8] linear code over GF(4)

codes are known with d = 101, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-002754 | codetables-gf4-149-9

Minimum distance of a [149, 9] linear code over GF(4)

codes are known with d = 99, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-002755 | codetables-gf4-150-5

Minimum distance of a [150, 5] linear code over GF(4)

codes are known with d = 110, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-002756 | codetables-gf4-150-6

Minimum distance of a [150, 6] linear code over GF(4)

codes are known with d = 108, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-002757 | codetables-gf4-150-7

Minimum distance of a [150, 7] linear code over GF(4)

codes are known with d = 104, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-002758 | codetables-gf4-150-8

Minimum distance of a [150, 8] linear code over GF(4)

codes are known with d = 102, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-002759 | codetables-gf4-150-9

Minimum distance of a [150, 9] linear code over GF(4)

codes are known with d = 100, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-002760 | codetables-gf4-151-5

Minimum distance of a [151, 5] linear code over GF(4)

codes are known with d = 111, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-002761 | codetables-gf4-151-6

Minimum distance of a [151, 6] linear code over GF(4)

codes are known with d = 108, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-002762 | codetables-gf4-151-7

Minimum distance of a [151, 7] linear code over GF(4)

codes are known with d = 105, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-002763 | codetables-gf4-151-8

Minimum distance of a [151, 8] linear code over GF(4)

codes are known with d = 103, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-002764 | codetables-gf4-151-9

Minimum distance of a [151, 9] linear code over GF(4)

codes are known with d = 100, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-002765 | codetables-gf4-152-6

Minimum distance of a [152, 6] linear code over GF(4)

codes are known with d = 109, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-002766 | codetables-gf4-152-7

Minimum distance of a [152, 7] linear code over GF(4)

codes are known with d = 106, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-002767 | codetables-gf4-152-8

Minimum distance of a [152, 8] linear code over GF(4)

codes are known with d = 104, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-002768 | codetables-gf4-152-9

Minimum distance of a [152, 9] linear code over GF(4)

codes are known with d = 101, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-002769 | codetables-gf4-153-6

Minimum distance of a [153, 6] linear code over GF(4)

codes are known with d = 110, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-002770 | codetables-gf4-153-7

Minimum distance of a [153, 7] linear code over GF(4)

codes are known with d = 107, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-002771 | codetables-gf4-153-8

Minimum distance of a [153, 8] linear code over GF(4)

codes are known with d = 104, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-002772 | codetables-gf4-153-9

Minimum distance of a [153, 9] linear code over GF(4)

codes are known with d = 101, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-002773 | codetables-gf4-154-6

Minimum distance of a [154, 6] linear code over GF(4)

codes are known with d = 111, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-002774 | codetables-gf4-154-7

Minimum distance of a [154, 7] linear code over GF(4)

codes are known with d = 108, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-002775 | codetables-gf4-154-8

Minimum distance of a [154, 8] linear code over GF(4)

codes are known with d = 105, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-002776 | codetables-gf4-154-9

Minimum distance of a [154, 9] linear code over GF(4)

codes are known with d = 102, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-002777 | codetables-gf4-155-6

Minimum distance of a [155, 6] linear code over GF(4)

codes are known with d = 112, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-002778 | codetables-gf4-155-7

Minimum distance of a [155, 7] linear code over GF(4)

codes are known with d = 108, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-002779 | codetables-gf4-155-8

Minimum distance of a [155, 8] linear code over GF(4)

codes are known with d = 106, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-002780 | codetables-gf4-155-9

Minimum distance of a [155, 9] linear code over GF(4)

codes are known with d = 103, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-002781 | codetables-gf4-156-5

Minimum distance of a [156, 5] linear code over GF(4)

codes are known with d = 114, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-002782 | codetables-gf4-156-6

Minimum distance of a [156, 6] linear code over GF(4)

codes are known with d = 112, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-002783 | codetables-gf4-156-7

Minimum distance of a [156, 7] linear code over GF(4)

codes are known with d = 109, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-002784 | codetables-gf4-156-8

Minimum distance of a [156, 8] linear code over GF(4)

codes are known with d = 107, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-002785 | codetables-gf4-156-9

Minimum distance of a [156, 9] linear code over GF(4)

codes are known with d = 104, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-002786 | codetables-gf4-157-5

Minimum distance of a [157, 5] linear code over GF(4)

codes are known with d = 115, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-002787 | codetables-gf4-157-6

Minimum distance of a [157, 6] linear code over GF(4)

codes are known with d = 113, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-002788 | codetables-gf4-157-7

Minimum distance of a [157, 7] linear code over GF(4)

codes are known with d = 110, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-002789 | codetables-gf4-157-8

Minimum distance of a [157, 8] linear code over GF(4)

codes are known with d = 108, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-002790 | codetables-gf4-157-9

Minimum distance of a [157, 9] linear code over GF(4)

codes are known with d = 105, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-002791 | codetables-gf4-158-6

Minimum distance of a [158, 6] linear code over GF(4)

codes are known with d = 114, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-002792 | codetables-gf4-158-7

Minimum distance of a [158, 7] linear code over GF(4)

codes are known with d = 111, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-002793 | codetables-gf4-158-8

Minimum distance of a [158, 8] linear code over GF(4)

codes are known with d = 108, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-002794 | codetables-gf4-158-9

Minimum distance of a [158, 9] linear code over GF(4)

codes are known with d = 106, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-002795 | codetables-gf4-159-6

Minimum distance of a [159, 6] linear code over GF(4)

codes are known with d = 115, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-002796 | codetables-gf4-159-7

Minimum distance of a [159, 7] linear code over GF(4)

codes are known with d = 112, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-002797 | codetables-gf4-159-8

Minimum distance of a [159, 8] linear code over GF(4)

codes are known with d = 108, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-002798 | codetables-gf4-159-9

Minimum distance of a [159, 9] linear code over GF(4)

codes are known with d = 107, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-002799 | codetables-gf4-160-6

Minimum distance of a [160, 6] linear code over GF(4)

codes are known with d = 116, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-002800 | codetables-gf4-160-7

Minimum distance of a [160, 7] linear code over GF(4)

codes are known with d = 112, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-002801 | codetables-gf4-160-8

Minimum distance of a [160, 8] linear code over GF(4)

codes are known with d = 109, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-002802 | codetables-gf4-160-9

Minimum distance of a [160, 9] linear code over GF(4)

codes are known with d = 108, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-002803 | codetables-gf4-161-6

Minimum distance of a [161, 6] linear code over GF(4)

codes are known with d = 116, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-002804 | codetables-gf4-161-7

Minimum distance of a [161, 7] linear code over GF(4)

codes are known with d = 112, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-002805 | codetables-gf4-161-8

Minimum distance of a [161, 8] linear code over GF(4)

codes are known with d = 110, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-002806 | codetables-gf4-161-9

Minimum distance of a [161, 9] linear code over GF(4)

codes are known with d = 109, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-002807 | codetables-gf4-162-6

Minimum distance of a [162, 6] linear code over GF(4)

codes are known with d = 117, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-002808 | codetables-gf4-162-7

Minimum distance of a [162, 7] linear code over GF(4)

codes are known with d = 113, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-002809 | codetables-gf4-162-8

Minimum distance of a [162, 8] linear code over GF(4)

codes are known with d = 111, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-002810 | codetables-gf4-162-9

Minimum distance of a [162, 9] linear code over GF(4)

codes are known with d = 110, and no code can have d greater than 115.

来源状态 open | 2026-07-23 | 来源

V3P-002811 | codetables-gf4-163-6

Minimum distance of a [163, 6] linear code over GF(4)

codes are known with d = 118, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-002812 | codetables-gf4-163-7

Minimum distance of a [163, 7] linear code over GF(4)

codes are known with d = 114, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-002813 | codetables-gf4-163-8

Minimum distance of a [163, 8] linear code over GF(4)

codes are known with d = 112, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-002814 | codetables-gf4-163-9

Minimum distance of a [163, 9] linear code over GF(4)

codes are known with d = 110, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-002815 | codetables-gf4-164-6

Minimum distance of a [164, 6] linear code over GF(4)

codes are known with d = 119, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-002816 | codetables-gf4-164-7

Minimum distance of a [164, 7] linear code over GF(4)

codes are known with d = 115, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-002817 | codetables-gf4-164-8

Minimum distance of a [164, 8] linear code over GF(4)

codes are known with d = 112, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-002818 | codetables-gf4-164-9

Minimum distance of a [164, 9] linear code over GF(4)

codes are known with d = 110, and no code can have d greater than 116.

来源状态 open | 2026-07-23 | 来源

V3P-002819 | codetables-gf4-165-6

Minimum distance of a [165, 6] linear code over GF(4)

codes are known with d = 120, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-002820 | codetables-gf4-165-7

Minimum distance of a [165, 7] linear code over GF(4)

codes are known with d = 116, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-002821 | codetables-gf4-165-8

Minimum distance of a [165, 8] linear code over GF(4)

codes are known with d = 112, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-002822 | codetables-gf4-165-9

Minimum distance of a [165, 9] linear code over GF(4)

codes are known with d = 111, and no code can have d greater than 117.

来源状态 open | 2026-07-23 | 来源

V3P-002823 | codetables-gf4-166-5

Minimum distance of a [166, 5] linear code over GF(4)

codes are known with d = 122, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-002824 | codetables-gf4-166-6

Minimum distance of a [166, 6] linear code over GF(4)

codes are known with d = 120, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-002825 | codetables-gf4-166-7

Minimum distance of a [166, 7] linear code over GF(4)

codes are known with d = 116, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-002826 | codetables-gf4-166-8

Minimum distance of a [166, 8] linear code over GF(4)

codes are known with d = 113, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-002827 | codetables-gf4-166-9

Minimum distance of a [166, 9] linear code over GF(4)

codes are known with d = 112, and no code can have d greater than 118.

来源状态 open | 2026-07-23 | 来源

V3P-002828 | codetables-gf4-167-5

Minimum distance of a [167, 5] linear code over GF(4)

codes are known with d = 123, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-002829 | codetables-gf4-167-6

Minimum distance of a [167, 6] linear code over GF(4)

codes are known with d = 120, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-002830 | codetables-gf4-167-7

Minimum distance of a [167, 7] linear code over GF(4)

codes are known with d = 117, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-002831 | codetables-gf4-167-8

Minimum distance of a [167, 8] linear code over GF(4)

codes are known with d = 114, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-002832 | codetables-gf4-167-9

Minimum distance of a [167, 9] linear code over GF(4)

codes are known with d = 113, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-002833 | codetables-gf4-168-6

Minimum distance of a [168, 6] linear code over GF(4)

codes are known with d = 121, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-002834 | codetables-gf4-168-7

Minimum distance of a [168, 7] linear code over GF(4)

codes are known with d = 118, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-002835 | codetables-gf4-168-8

Minimum distance of a [168, 8] linear code over GF(4)

codes are known with d = 115, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-002836 | codetables-gf4-168-9

Minimum distance of a [168, 9] linear code over GF(4)

codes are known with d = 114, and no code can have d greater than 119.

来源状态 open | 2026-07-23 | 来源

V3P-002837 | codetables-gf4-169-6

Minimum distance of a [169, 6] linear code over GF(4)

codes are known with d = 122, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-002838 | codetables-gf4-169-7

Minimum distance of a [169, 7] linear code over GF(4)

codes are known with d = 119, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-002839 | codetables-gf4-169-8

Minimum distance of a [169, 8] linear code over GF(4)

codes are known with d = 116, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-002840 | codetables-gf4-169-9

Minimum distance of a [169, 9] linear code over GF(4)

codes are known with d = 115, and no code can have d greater than 120.

来源状态 open | 2026-07-23 | 来源

V3P-002841 | codetables-gf4-170-6

Minimum distance of a [170, 6] linear code over GF(4)

codes are known with d = 122, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-002842 | codetables-gf4-170-7

Minimum distance of a [170, 7] linear code over GF(4)

codes are known with d = 120, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-002843 | codetables-gf4-170-8

Minimum distance of a [170, 8] linear code over GF(4)

codes are known with d = 117, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-002844 | codetables-gf4-170-9

Minimum distance of a [170, 9] linear code over GF(4)

codes are known with d = 116, and no code can have d greater than 121.

来源状态 open | 2026-07-23 | 来源

V3P-002845 | codetables-gf4-171-6

Minimum distance of a [171, 6] linear code over GF(4)

codes are known with d = 123, and no code can have d greater than 125.

来源状态 open | 2026-07-23 | 来源

V3P-002846 | codetables-gf4-171-7

Minimum distance of a [171, 7] linear code over GF(4)

codes are known with d = 120, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-002847 | codetables-gf4-171-8

Minimum distance of a [171, 8] linear code over GF(4)

codes are known with d = 118, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-002848 | codetables-gf4-171-9

Minimum distance of a [171, 9] linear code over GF(4)

codes are known with d = 117, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-002849 | codetables-gf4-172-6

Minimum distance of a [172, 6] linear code over GF(4)

codes are known with d = 124, and no code can have d greater than 126.

来源状态 open | 2026-07-23 | 来源

V3P-002850 | codetables-gf4-172-7

Minimum distance of a [172, 7] linear code over GF(4)

codes are known with d = 120, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-002851 | codetables-gf4-172-8

Minimum distance of a [172, 8] linear code over GF(4)

codes are known with d = 118, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-002852 | codetables-gf4-172-9

Minimum distance of a [172, 9] linear code over GF(4)

codes are known with d = 117, and no code can have d greater than 122.

来源状态 open | 2026-07-23 | 来源

V3P-002853 | codetables-gf4-173-6

Minimum distance of a [173, 6] linear code over GF(4)

codes are known with d = 124, and no code can have d greater than 127.

来源状态 open | 2026-07-23 | 来源

V3P-002854 | codetables-gf4-173-7

Minimum distance of a [173, 7] linear code over GF(4)

codes are known with d = 121, and no code can have d greater than 125.

来源状态 open | 2026-07-23 | 来源

V3P-002855 | codetables-gf4-173-8

Minimum distance of a [173, 8] linear code over GF(4)

codes are known with d = 119, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-002856 | codetables-gf4-173-9

Minimum distance of a [173, 9] linear code over GF(4)

codes are known with d = 117, and no code can have d greater than 123.

来源状态 open | 2026-07-23 | 来源

V3P-002857 | codetables-gf4-174-6

Minimum distance of a [174, 6] linear code over GF(4)

codes are known with d = 125, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-002858 | codetables-gf4-174-7

Minimum distance of a [174, 7] linear code over GF(4)

codes are known with d = 122, and no code can have d greater than 126.

来源状态 open | 2026-07-23 | 来源

V3P-002859 | codetables-gf4-174-8

Minimum distance of a [174, 8] linear code over GF(4)

codes are known with d = 120, and no code can have d greater than 125.

来源状态 open | 2026-07-23 | 来源

V3P-002860 | codetables-gf4-174-9

Minimum distance of a [174, 9] linear code over GF(4)

codes are known with d = 118, and no code can have d greater than 124.

来源状态 open | 2026-07-23 | 来源

V3P-002861 | codetables-gf4-175-5

Minimum distance of a [175, 5] linear code over GF(4)

codes are known with d = 128, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-002862 | codetables-gf4-175-6

Minimum distance of a [175, 6] linear code over GF(4)

codes are known with d = 126, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-002863 | codetables-gf4-175-7

Minimum distance of a [175, 7] linear code over GF(4)

codes are known with d = 123, and no code can have d greater than 127.

来源状态 open | 2026-07-23 | 来源

V3P-002864 | codetables-gf4-175-8

Minimum distance of a [175, 8] linear code over GF(4)

codes are known with d = 120, and no code can have d greater than 125.

来源状态 open | 2026-07-23 | 来源

V3P-002865 | codetables-gf4-175-9

Minimum distance of a [175, 9] linear code over GF(4)

codes are known with d = 118, and no code can have d greater than 125.

来源状态 open | 2026-07-23 | 来源

V3P-002866 | codetables-gf4-176-5

Minimum distance of a [176, 5] linear code over GF(4)

codes are known with d = 129, and no code can have d greater than 130.

来源状态 open | 2026-07-23 | 来源

V3P-002867 | codetables-gf4-176-6

Minimum distance of a [176, 6] linear code over GF(4)

codes are known with d = 127, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-002868 | codetables-gf4-176-7

Minimum distance of a [176, 7] linear code over GF(4)

codes are known with d = 124, and no code can have d greater than 127.

来源状态 open | 2026-07-23 | 来源

V3P-002869 | codetables-gf4-176-8

Minimum distance of a [176, 8] linear code over GF(4)

codes are known with d = 121, and no code can have d greater than 126.

来源状态 open | 2026-07-23 | 来源

V3P-002870 | codetables-gf4-176-9

Minimum distance of a [176, 9] linear code over GF(4)

codes are known with d = 119, and no code can have d greater than 125.

来源状态 open | 2026-07-23 | 来源

V3P-002871 | codetables-gf4-177-5

Minimum distance of a [177, 5] linear code over GF(4)

codes are known with d = 130, and no code can have d greater than 131.

来源状态 open | 2026-07-23 | 来源

V3P-002872 | codetables-gf4-177-6

Minimum distance of a [177, 6] linear code over GF(4)

codes are known with d = 128, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-002873 | codetables-gf4-177-7

Minimum distance of a [177, 7] linear code over GF(4)

codes are known with d = 124, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-002874 | codetables-gf4-177-8

Minimum distance of a [177, 8] linear code over GF(4)

codes are known with d = 122, and no code can have d greater than 127.

来源状态 open | 2026-07-23 | 来源

V3P-002875 | codetables-gf4-177-9

Minimum distance of a [177, 9] linear code over GF(4)

codes are known with d = 120, and no code can have d greater than 126.

来源状态 open | 2026-07-23 | 来源

V3P-002876 | codetables-gf4-178-5

Minimum distance of a [178, 5] linear code over GF(4)

codes are known with d = 131, and no code can have d greater than 132.

来源状态 open | 2026-07-23 | 来源

V3P-002877 | codetables-gf4-178-6

Minimum distance of a [178, 6] linear code over GF(4)

codes are known with d = 128, and no code can have d greater than 130.

来源状态 open | 2026-07-23 | 来源

V3P-002878 | codetables-gf4-178-7

Minimum distance of a [178, 7] linear code over GF(4)

codes are known with d = 125, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-002879 | codetables-gf4-178-8

Minimum distance of a [178, 8] linear code over GF(4)

codes are known with d = 122, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-002880 | codetables-gf4-178-9

Minimum distance of a [178, 9] linear code over GF(4)

codes are known with d = 120, and no code can have d greater than 127.

来源状态 open | 2026-07-23 | 来源

V3P-002881 | codetables-gf4-179-6

Minimum distance of a [179, 6] linear code over GF(4)

codes are known with d = 129, and no code can have d greater than 131.

来源状态 open | 2026-07-23 | 来源

V3P-002882 | codetables-gf4-179-7

Minimum distance of a [179, 7] linear code over GF(4)

codes are known with d = 126, and no code can have d greater than 130.

来源状态 open | 2026-07-23 | 来源

V3P-002883 | codetables-gf4-179-8

Minimum distance of a [179, 8] linear code over GF(4)

codes are known with d = 123, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-002884 | codetables-gf4-179-9

Minimum distance of a [179, 9] linear code over GF(4)

codes are known with d = 121, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-002885 | codetables-gf4-180-5

Minimum distance of a [180, 5] linear code over GF(4)

codes are known with d = 132, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-002886 | codetables-gf4-180-6

Minimum distance of a [180, 6] linear code over GF(4)

codes are known with d = 130, and no code can have d greater than 132.

来源状态 open | 2026-07-23 | 来源

V3P-002887 | codetables-gf4-180-7

Minimum distance of a [180, 7] linear code over GF(4)

codes are known with d = 127, and no code can have d greater than 130.

来源状态 open | 2026-07-23 | 来源

V3P-002888 | codetables-gf4-180-8

Minimum distance of a [180, 8] linear code over GF(4)

codes are known with d = 124, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-002889 | codetables-gf4-180-9

Minimum distance of a [180, 9] linear code over GF(4)

codes are known with d = 122, and no code can have d greater than 128.

来源状态 open | 2026-07-23 | 来源

V3P-002890 | codetables-gf4-181-5

Minimum distance of a [181, 5] linear code over GF(4)

codes are known with d = 133, and no code can have d greater than 134.

来源状态 open | 2026-07-23 | 来源

V3P-002891 | codetables-gf4-181-6

Minimum distance of a [181, 6] linear code over GF(4)

codes are known with d = 131, and no code can have d greater than 132.

来源状态 open | 2026-07-23 | 来源

V3P-002892 | codetables-gf4-181-7

Minimum distance of a [181, 7] linear code over GF(4)

codes are known with d = 128, and no code can have d greater than 131.

来源状态 open | 2026-07-23 | 来源

V3P-002893 | codetables-gf4-181-8

Minimum distance of a [181, 8] linear code over GF(4)

codes are known with d = 124, and no code can have d greater than 130.

来源状态 open | 2026-07-23 | 来源

V3P-002894 | codetables-gf4-181-9

Minimum distance of a [181, 9] linear code over GF(4)

codes are known with d = 122, and no code can have d greater than 129.

来源状态 open | 2026-07-23 | 来源

V3P-002895 | codetables-gf4-182-5

Minimum distance of a [182, 5] linear code over GF(4)

codes are known with d = 134, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-002896 | codetables-gf4-182-6

Minimum distance of a [182, 6] linear code over GF(4)

codes are known with d = 132, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-002897 | codetables-gf4-182-7

Minimum distance of a [182, 7] linear code over GF(4)

codes are known with d = 128, and no code can have d greater than 132.

来源状态 open | 2026-07-23 | 来源

V3P-002898 | codetables-gf4-182-8

Minimum distance of a [182, 8] linear code over GF(4)

codes are known with d = 125, and no code can have d greater than 130.

来源状态 open | 2026-07-23 | 来源

V3P-002899 | codetables-gf4-182-9

Minimum distance of a [182, 9] linear code over GF(4)

codes are known with d = 123, and no code can have d greater than 130.

来源状态 open | 2026-07-23 | 来源

V3P-002900 | codetables-gf4-183-5

Minimum distance of a [183, 5] linear code over GF(4)

codes are known with d = 135, and no code can have d greater than 136.

来源状态 open | 2026-07-23 | 来源

V3P-002901 | codetables-gf4-183-6

Minimum distance of a [183, 6] linear code over GF(4)

codes are known with d = 132, and no code can have d greater than 134.

来源状态 open | 2026-07-23 | 来源

V3P-002902 | codetables-gf4-183-7

Minimum distance of a [183, 7] linear code over GF(4)

codes are known with d = 129, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-002903 | codetables-gf4-183-8

Minimum distance of a [183, 8] linear code over GF(4)

codes are known with d = 126, and no code can have d greater than 131.

来源状态 open | 2026-07-23 | 来源

V3P-002904 | codetables-gf4-183-9

Minimum distance of a [183, 9] linear code over GF(4)

codes are known with d = 124, and no code can have d greater than 130.

来源状态 open | 2026-07-23 | 来源

V3P-002905 | codetables-gf4-184-6

Minimum distance of a [184, 6] linear code over GF(4)

codes are known with d = 133, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-002906 | codetables-gf4-184-7

Minimum distance of a [184, 7] linear code over GF(4)

codes are known with d = 130, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-002907 | codetables-gf4-184-8

Minimum distance of a [184, 8] linear code over GF(4)

codes are known with d = 127, and no code can have d greater than 132.

来源状态 open | 2026-07-23 | 来源

V3P-002908 | codetables-gf4-184-9

Minimum distance of a [184, 9] linear code over GF(4)

codes are known with d = 124, and no code can have d greater than 131.

来源状态 open | 2026-07-23 | 来源

V3P-002909 | codetables-gf4-185-5

Minimum distance of a [185, 5] linear code over GF(4)

codes are known with d = 136, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-002910 | codetables-gf4-185-6

Minimum distance of a [185, 6] linear code over GF(4)

codes are known with d = 134, and no code can have d greater than 136.

来源状态 open | 2026-07-23 | 来源

V3P-002911 | codetables-gf4-185-7

Minimum distance of a [185, 7] linear code over GF(4)

codes are known with d = 131, and no code can have d greater than 134.

来源状态 open | 2026-07-23 | 来源

V3P-002912 | codetables-gf4-185-8

Minimum distance of a [185, 8] linear code over GF(4)

codes are known with d = 128, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-002913 | codetables-gf4-185-9

Minimum distance of a [185, 9] linear code over GF(4)

codes are known with d = 125, and no code can have d greater than 132.

来源状态 open | 2026-07-23 | 来源

V3P-002914 | codetables-gf4-186-5

Minimum distance of a [186, 5] linear code over GF(4)

codes are known with d = 137, and no code can have d greater than 138.

来源状态 open | 2026-07-23 | 来源

V3P-002915 | codetables-gf4-186-6

Minimum distance of a [186, 6] linear code over GF(4)

codes are known with d = 135, and no code can have d greater than 136.

来源状态 open | 2026-07-23 | 来源

V3P-002916 | codetables-gf4-186-7

Minimum distance of a [186, 7] linear code over GF(4)

codes are known with d = 132, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-002917 | codetables-gf4-186-8

Minimum distance of a [186, 8] linear code over GF(4)

codes are known with d = 128, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-002918 | codetables-gf4-186-9

Minimum distance of a [186, 9] linear code over GF(4)

codes are known with d = 126, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-002919 | codetables-gf4-187-6

Minimum distance of a [187, 6] linear code over GF(4)

codes are known with d = 136, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-002920 | codetables-gf4-187-7

Minimum distance of a [187, 7] linear code over GF(4)

codes are known with d = 132, and no code can have d greater than 136.

来源状态 open | 2026-07-23 | 来源

V3P-002921 | codetables-gf4-187-8

Minimum distance of a [187, 8] linear code over GF(4)

codes are known with d = 129, and no code can have d greater than 134.

来源状态 open | 2026-07-23 | 来源

V3P-002922 | codetables-gf4-187-9

Minimum distance of a [187, 9] linear code over GF(4)

codes are known with d = 126, and no code can have d greater than 133.

来源状态 open | 2026-07-23 | 来源

V3P-002923 | codetables-gf4-188-6

Minimum distance of a [188, 6] linear code over GF(4)

codes are known with d = 136, and no code can have d greater than 138.

来源状态 open | 2026-07-23 | 来源

V3P-002924 | codetables-gf4-188-7

Minimum distance of a [188, 7] linear code over GF(4)

codes are known with d = 132, and no code can have d greater than 136.

来源状态 open | 2026-07-23 | 来源

V3P-002925 | codetables-gf4-188-8

Minimum distance of a [188, 8] linear code over GF(4)

codes are known with d = 130, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-002926 | codetables-gf4-188-9

Minimum distance of a [188, 9] linear code over GF(4)

codes are known with d = 127, and no code can have d greater than 134.

来源状态 open | 2026-07-23 | 来源

V3P-002927 | codetables-gf4-189-6

Minimum distance of a [189, 6] linear code over GF(4)

codes are known with d = 136, and no code can have d greater than 139.

来源状态 open | 2026-07-23 | 来源

V3P-002928 | codetables-gf4-189-7

Minimum distance of a [189, 7] linear code over GF(4)

codes are known with d = 133, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-002929 | codetables-gf4-189-8

Minimum distance of a [189, 8] linear code over GF(4)

codes are known with d = 131, and no code can have d greater than 136.

来源状态 open | 2026-07-23 | 来源

V3P-002930 | codetables-gf4-189-9

Minimum distance of a [189, 9] linear code over GF(4)

codes are known with d = 128, and no code can have d greater than 135.

来源状态 open | 2026-07-23 | 来源

V3P-002931 | codetables-gf4-19-8

Minimum distance of a [19, 8] linear code over GF(4)

codes are known with d = 8, and no code can have d greater than 9.

来源状态 open | 2026-07-23 | 来源

V3P-002932 | codetables-gf4-190-6

Minimum distance of a [190, 6] linear code over GF(4)

codes are known with d = 137, and no code can have d greater than 140.

来源状态 open | 2026-07-23 | 来源

V3P-002933 | codetables-gf4-190-7

Minimum distance of a [190, 7] linear code over GF(4)

codes are known with d = 134, and no code can have d greater than 138.

来源状态 open | 2026-07-23 | 来源

V3P-002934 | codetables-gf4-190-8

Minimum distance of a [190, 8] linear code over GF(4)

codes are known with d = 132, and no code can have d greater than 136.

来源状态 open | 2026-07-23 | 来源

V3P-002935 | codetables-gf4-190-9

Minimum distance of a [190, 9] linear code over GF(4)

codes are known with d = 128, and no code can have d greater than 136.

来源状态 open | 2026-07-23 | 来源

V3P-002936 | codetables-gf4-191-6

Minimum distance of a [191, 6] linear code over GF(4)

codes are known with d = 138, and no code can have d greater than 140.

来源状态 open | 2026-07-23 | 来源

V3P-002937 | codetables-gf4-191-7

Minimum distance of a [191, 7] linear code over GF(4)

codes are known with d = 135, and no code can have d greater than 139.

来源状态 open | 2026-07-23 | 来源

V3P-002938 | codetables-gf4-191-8

Minimum distance of a [191, 8] linear code over GF(4)

codes are known with d = 132, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-002939 | codetables-gf4-191-9

Minimum distance of a [191, 9] linear code over GF(4)

codes are known with d = 129, and no code can have d greater than 136.

来源状态 open | 2026-07-23 | 来源

V3P-002940 | codetables-gf4-192-6

Minimum distance of a [192, 6] linear code over GF(4)

codes are known with d = 139, and no code can have d greater than 141.

来源状态 open | 2026-07-23 | 来源

V3P-002941 | codetables-gf4-192-7

Minimum distance of a [192, 7] linear code over GF(4)

codes are known with d = 136, and no code can have d greater than 140.

来源状态 open | 2026-07-23 | 来源

V3P-002942 | codetables-gf4-192-8

Minimum distance of a [192, 8] linear code over GF(4)

codes are known with d = 132, and no code can have d greater than 138.

来源状态 open | 2026-07-23 | 来源

V3P-002943 | codetables-gf4-192-9

Minimum distance of a [192, 9] linear code over GF(4)

codes are known with d = 130, and no code can have d greater than 137.

来源状态 open | 2026-07-23 | 来源

V3P-002944 | codetables-gf4-193-6

Minimum distance of a [193, 6] linear code over GF(4)

codes are known with d = 140, and no code can have d greater than 142.

来源状态 open | 2026-07-23 | 来源

V3P-002945 | codetables-gf4-193-7

Minimum distance of a [193, 7] linear code over GF(4)

codes are known with d = 136, and no code can have d greater than 141.

来源状态 open | 2026-07-23 | 来源

V3P-002946 | codetables-gf4-193-8

Minimum distance of a [193, 8] linear code over GF(4)

codes are known with d = 133, and no code can have d greater than 139.

来源状态 open | 2026-07-23 | 来源

V3P-002947 | codetables-gf4-193-9

Minimum distance of a [193, 9] linear code over GF(4)

codes are known with d = 131, and no code can have d greater than 138.

来源状态 open | 2026-07-23 | 来源

V3P-002948 | codetables-gf4-194-6

Minimum distance of a [194, 6] linear code over GF(4)

codes are known with d = 140, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-002949 | codetables-gf4-194-7

Minimum distance of a [194, 7] linear code over GF(4)

codes are known with d = 137, and no code can have d greater than 142.

来源状态 open | 2026-07-23 | 来源

V3P-002950 | codetables-gf4-194-8

Minimum distance of a [194, 8] linear code over GF(4)

codes are known with d = 134, and no code can have d greater than 139.

来源状态 open | 2026-07-23 | 来源

V3P-002951 | codetables-gf4-194-9

Minimum distance of a [194, 9] linear code over GF(4)

codes are known with d = 132, and no code can have d greater than 139.

来源状态 open | 2026-07-23 | 来源

V3P-002952 | codetables-gf4-195-6

Minimum distance of a [195, 6] linear code over GF(4)

codes are known with d = 141, and no code can have d greater than 144.

来源状态 open | 2026-07-23 | 来源

V3P-002953 | codetables-gf4-195-7

Minimum distance of a [195, 7] linear code over GF(4)

codes are known with d = 138, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-002954 | codetables-gf4-195-8

Minimum distance of a [195, 8] linear code over GF(4)

codes are known with d = 135, and no code can have d greater than 140.

来源状态 open | 2026-07-23 | 来源

V3P-002955 | codetables-gf4-195-9

Minimum distance of a [195, 9] linear code over GF(4)

codes are known with d = 132, and no code can have d greater than 139.

来源状态 open | 2026-07-23 | 来源

V3P-002956 | codetables-gf4-196-6

Minimum distance of a [196, 6] linear code over GF(4)

codes are known with d = 142, and no code can have d greater than 144.

来源状态 open | 2026-07-23 | 来源

V3P-002957 | codetables-gf4-196-7

Minimum distance of a [196, 7] linear code over GF(4)

codes are known with d = 139, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-002958 | codetables-gf4-196-8

Minimum distance of a [196, 8] linear code over GF(4)

codes are known with d = 136, and no code can have d greater than 141.

来源状态 open | 2026-07-23 | 来源

V3P-002959 | codetables-gf4-196-9

Minimum distance of a [196, 9] linear code over GF(4)

codes are known with d = 133, and no code can have d greater than 140.

来源状态 open | 2026-07-23 | 来源

V3P-002960 | codetables-gf4-197-6

Minimum distance of a [197, 6] linear code over GF(4)

codes are known with d = 143, and no code can have d greater than 144.

来源状态 open | 2026-07-23 | 来源

V3P-002961 | codetables-gf4-197-7

Minimum distance of a [197, 7] linear code over GF(4)

codes are known with d = 140, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-002962 | codetables-gf4-197-8

Minimum distance of a [197, 8] linear code over GF(4)

codes are known with d = 136, and no code can have d greater than 142.

来源状态 open | 2026-07-23 | 来源

V3P-002963 | codetables-gf4-197-9

Minimum distance of a [197, 9] linear code over GF(4)

codes are known with d = 134, and no code can have d greater than 141.

来源状态 open | 2026-07-23 | 来源

V3P-002964 | codetables-gf4-198-6

Minimum distance of a [198, 6] linear code over GF(4)

codes are known with d = 144, and no code can have d greater than 145.

来源状态 open | 2026-07-23 | 来源

V3P-002965 | codetables-gf4-198-7

Minimum distance of a [198, 7] linear code over GF(4)

codes are known with d = 140, and no code can have d greater than 144.

来源状态 open | 2026-07-23 | 来源

V3P-002966 | codetables-gf4-198-8

Minimum distance of a [198, 8] linear code over GF(4)

codes are known with d = 137, and no code can have d greater than 142.

来源状态 open | 2026-07-23 | 来源

V3P-002967 | codetables-gf4-198-9

Minimum distance of a [198, 9] linear code over GF(4)

codes are known with d = 135, and no code can have d greater than 142.

来源状态 open | 2026-07-23 | 来源

V3P-002968 | codetables-gf4-199-6

Minimum distance of a [199, 6] linear code over GF(4)

codes are known with d = 144, and no code can have d greater than 146.

来源状态 open | 2026-07-23 | 来源

V3P-002969 | codetables-gf4-199-7

Minimum distance of a [199, 7] linear code over GF(4)

codes are known with d = 140, and no code can have d greater than 145.

来源状态 open | 2026-07-23 | 来源

V3P-002970 | codetables-gf4-199-8

Minimum distance of a [199, 8] linear code over GF(4)

codes are known with d = 138, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-002971 | codetables-gf4-199-9

Minimum distance of a [199, 9] linear code over GF(4)

codes are known with d = 136, and no code can have d greater than 142.

来源状态 open | 2026-07-23 | 来源

V3P-002972 | codetables-gf4-20-9

Minimum distance of a [20, 9] linear code over GF(4)

codes are known with d = 8, and no code can have d greater than 9.

来源状态 open | 2026-07-23 | 来源

V3P-002973 | codetables-gf4-200-6

Minimum distance of a [200, 6] linear code over GF(4)

codes are known with d = 145, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-002974 | codetables-gf4-200-7

Minimum distance of a [200, 7] linear code over GF(4)

codes are known with d = 141, and no code can have d greater than 146.

来源状态 open | 2026-07-23 | 来源

V3P-002975 | codetables-gf4-200-8

Minimum distance of a [200, 8] linear code over GF(4)

codes are known with d = 139, and no code can have d greater than 144.

来源状态 open | 2026-07-23 | 来源

V3P-002976 | codetables-gf4-200-9

Minimum distance of a [200, 9] linear code over GF(4)

codes are known with d = 136, and no code can have d greater than 143.

来源状态 open | 2026-07-23 | 来源

V3P-002977 | codetables-gf4-201-6

Minimum distance of a [201, 6] linear code over GF(4)

codes are known with d = 146, and no code can have d greater than 148.

来源状态 open | 2026-07-23 | 来源

V3P-002978 | codetables-gf4-201-7

Minimum distance of a [201, 7] linear code over GF(4)

codes are known with d = 142, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-002979 | codetables-gf4-201-8

Minimum distance of a [201, 8] linear code over GF(4)

codes are known with d = 140, and no code can have d greater than 145.

来源状态 open | 2026-07-23 | 来源

V3P-002980 | codetables-gf4-201-9

Minimum distance of a [201, 9] linear code over GF(4)

codes are known with d = 136, and no code can have d greater than 144.

来源状态 open | 2026-07-23 | 来源

V3P-002981 | codetables-gf4-202-6

Minimum distance of a [202, 6] linear code over GF(4)

codes are known with d = 147, and no code can have d greater than 148.

来源状态 open | 2026-07-23 | 来源

V3P-002982 | codetables-gf4-202-7

Minimum distance of a [202, 7] linear code over GF(4)

codes are known with d = 143, and no code can have d greater than 148.

来源状态 open | 2026-07-23 | 来源

V3P-002983 | codetables-gf4-202-8

Minimum distance of a [202, 8] linear code over GF(4)

codes are known with d = 140, and no code can have d greater than 145.

来源状态 open | 2026-07-23 | 来源

V3P-002984 | codetables-gf4-202-9

Minimum distance of a [202, 9] linear code over GF(4)

codes are known with d = 136, and no code can have d greater than 145.

来源状态 open | 2026-07-23 | 来源

V3P-002985 | codetables-gf4-203-6

Minimum distance of a [203, 6] linear code over GF(4)

codes are known with d = 148, and no code can have d greater than 149.

来源状态 open | 2026-07-23 | 来源

V3P-002986 | codetables-gf4-203-7

Minimum distance of a [203, 7] linear code over GF(4)

codes are known with d = 144, and no code can have d greater than 148.

来源状态 open | 2026-07-23 | 来源

V3P-002987 | codetables-gf4-203-8

Minimum distance of a [203, 8] linear code over GF(4)

codes are known with d = 140, and no code can have d greater than 146.

来源状态 open | 2026-07-23 | 来源

V3P-002988 | codetables-gf4-203-9

Minimum distance of a [203, 9] linear code over GF(4)

codes are known with d = 137, and no code can have d greater than 145.

来源状态 open | 2026-07-23 | 来源

V3P-002989 | codetables-gf4-204-6

Minimum distance of a [204, 6] linear code over GF(4)

codes are known with d = 148, and no code can have d greater than 150.

来源状态 open | 2026-07-23 | 来源

V3P-002990 | codetables-gf4-204-7

Minimum distance of a [204, 7] linear code over GF(4)

codes are known with d = 144, and no code can have d greater than 149.

来源状态 open | 2026-07-23 | 来源

V3P-002991 | codetables-gf4-204-8

Minimum distance of a [204, 8] linear code over GF(4)

codes are known with d = 141, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-002992 | codetables-gf4-204-9

Minimum distance of a [204, 9] linear code over GF(4)

codes are known with d = 138, and no code can have d greater than 146.

来源状态 open | 2026-07-23 | 来源

V3P-002993 | codetables-gf4-205-6

Minimum distance of a [205, 6] linear code over GF(4)

codes are known with d = 149, and no code can have d greater than 151.

来源状态 open | 2026-07-23 | 来源

V3P-002994 | codetables-gf4-205-7

Minimum distance of a [205, 7] linear code over GF(4)

codes are known with d = 145, and no code can have d greater than 150.

来源状态 open | 2026-07-23 | 来源

V3P-002995 | codetables-gf4-205-8

Minimum distance of a [205, 8] linear code over GF(4)

codes are known with d = 142, and no code can have d greater than 148.

来源状态 open | 2026-07-23 | 来源

V3P-002996 | codetables-gf4-205-9

Minimum distance of a [205, 9] linear code over GF(4)

codes are known with d = 139, and no code can have d greater than 147.

来源状态 open | 2026-07-23 | 来源

V3P-002997 | codetables-gf4-206-6

Minimum distance of a [206, 6] linear code over GF(4)

codes are known with d = 150, and no code can have d greater than 152.

来源状态 open | 2026-07-23 | 来源

V3P-002998 | codetables-gf4-206-7

Minimum distance of a [206, 7] linear code over GF(4)

codes are known with d = 146, and no code can have d greater than 151.

来源状态 open | 2026-07-23 | 来源

V3P-002999 | codetables-gf4-206-8

Minimum distance of a [206, 8] linear code over GF(4)

codes are known with d = 143, and no code can have d greater than 148.

来源状态 open | 2026-07-23 | 来源

V3P-003000 | codetables-gf4-206-9

Minimum distance of a [206, 9] linear code over GF(4)

codes are known with d = 140, and no code can have d greater than 148.

来源状态 open | 2026-07-23 | 来源

V3P-003001 | codetables-gf4-207-6

Minimum distance of a [207, 6] linear code over GF(4)

codes are known with d = 151, and no code can have d greater than 152.

来源状态 open | 2026-07-23 | 来源

V3P-003002 | codetables-gf4-207-7

Minimum distance of a [207, 7] linear code over GF(4)

codes are known with d = 147, and no code can have d greater than 152.

来源状态 open | 2026-07-23 | 来源

V3P-003003 | codetables-gf4-207-8

Minimum distance of a [207, 8] linear code over GF(4)

codes are known with d = 144, and no code can have d greater than 149.

来源状态 open | 2026-07-23 | 来源

V3P-003004 | codetables-gf4-207-9

Minimum distance of a [207, 9] linear code over GF(4)

codes are known with d = 141, and no code can have d greater than 148.

来源状态 open | 2026-07-23 | 来源

V3P-003005 | codetables-gf4-208-6

Minimum distance of a [208, 6] linear code over GF(4)

codes are known with d = 152, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-003006 | codetables-gf4-208-7

Minimum distance of a [208, 7] linear code over GF(4)

codes are known with d = 148, and no code can have d greater than 152.

来源状态 open | 2026-07-23 | 来源

V3P-003007 | codetables-gf4-208-8

Minimum distance of a [208, 8] linear code over GF(4)

codes are known with d = 144, and no code can have d greater than 150.

来源状态 open | 2026-07-23 | 来源

V3P-003008 | codetables-gf4-208-9

Minimum distance of a [208, 9] linear code over GF(4)

codes are known with d = 142, and no code can have d greater than 149.

来源状态 open | 2026-07-23 | 来源

V3P-003009 | codetables-gf4-209-6

Minimum distance of a [209, 6] linear code over GF(4)

codes are known with d = 152, and no code can have d greater than 154.

来源状态 open | 2026-07-23 | 来源

V3P-003010 | codetables-gf4-209-7

Minimum distance of a [209, 7] linear code over GF(4)

codes are known with d = 148, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-003011 | codetables-gf4-209-8

Minimum distance of a [209, 8] linear code over GF(4)

codes are known with d = 144, and no code can have d greater than 150.

来源状态 open | 2026-07-23 | 来源

V3P-003012 | codetables-gf4-209-9

Minimum distance of a [209, 9] linear code over GF(4)

codes are known with d = 143, and no code can have d greater than 150.

来源状态 open | 2026-07-23 | 来源

V3P-003013 | codetables-gf4-210-6

Minimum distance of a [210, 6] linear code over GF(4)

codes are known with d = 152, and no code can have d greater than 155.

来源状态 open | 2026-07-23 | 来源

V3P-003014 | codetables-gf4-210-7

Minimum distance of a [210, 7] linear code over GF(4)

codes are known with d = 148, and no code can have d greater than 154.

来源状态 open | 2026-07-23 | 来源

V3P-003015 | codetables-gf4-210-8

Minimum distance of a [210, 8] linear code over GF(4)

codes are known with d = 145, and no code can have d greater than 151.

来源状态 open | 2026-07-23 | 来源

V3P-003016 | codetables-gf4-210-9

Minimum distance of a [210, 9] linear code over GF(4)

codes are known with d = 144, and no code can have d greater than 150.

来源状态 open | 2026-07-23 | 来源

V3P-003017 | codetables-gf4-211-6

Minimum distance of a [211, 6] linear code over GF(4)

codes are known with d = 153, and no code can have d greater than 156.

来源状态 open | 2026-07-23 | 来源

V3P-003018 | codetables-gf4-211-7

Minimum distance of a [211, 7] linear code over GF(4)

codes are known with d = 149, and no code can have d greater than 155.

来源状态 open | 2026-07-23 | 来源

V3P-003019 | codetables-gf4-211-8

Minimum distance of a [211, 8] linear code over GF(4)

codes are known with d = 146, and no code can have d greater than 152.

来源状态 open | 2026-07-23 | 来源

V3P-003020 | codetables-gf4-211-9

Minimum distance of a [211, 9] linear code over GF(4)

codes are known with d = 144, and no code can have d greater than 151.

来源状态 open | 2026-07-23 | 来源

V3P-003021 | codetables-gf4-212-6

Minimum distance of a [212, 6] linear code over GF(4)

codes are known with d = 154, and no code can have d greater than 156.

来源状态 open | 2026-07-23 | 来源

V3P-003022 | codetables-gf4-212-7

Minimum distance of a [212, 7] linear code over GF(4)

codes are known with d = 150, and no code can have d greater than 156.

来源状态 open | 2026-07-23 | 来源

V3P-003023 | codetables-gf4-212-8

Minimum distance of a [212, 8] linear code over GF(4)

codes are known with d = 147, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-003024 | codetables-gf4-212-9

Minimum distance of a [212, 9] linear code over GF(4)

codes are known with d = 144, and no code can have d greater than 152.

来源状态 open | 2026-07-23 | 来源

V3P-003025 | codetables-gf4-213-6

Minimum distance of a [213, 6] linear code over GF(4)

codes are known with d = 155, and no code can have d greater than 157.

来源状态 open | 2026-07-23 | 来源

V3P-003026 | codetables-gf4-213-7

Minimum distance of a [213, 7] linear code over GF(4)

codes are known with d = 151, and no code can have d greater than 156.

来源状态 open | 2026-07-23 | 来源

V3P-003027 | codetables-gf4-213-8

Minimum distance of a [213, 8] linear code over GF(4)

codes are known with d = 148, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-003028 | codetables-gf4-213-9

Minimum distance of a [213, 9] linear code over GF(4)

codes are known with d = 145, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-003029 | codetables-gf4-214-6

Minimum distance of a [214, 6] linear code over GF(4)

codes are known with d = 156, and no code can have d greater than 158.

来源状态 open | 2026-07-23 | 来源

V3P-003030 | codetables-gf4-214-7

Minimum distance of a [214, 7] linear code over GF(4)

codes are known with d = 152, and no code can have d greater than 157.

来源状态 open | 2026-07-23 | 来源

V3P-003031 | codetables-gf4-214-8

Minimum distance of a [214, 8] linear code over GF(4)

codes are known with d = 148, and no code can have d greater than 154.

来源状态 open | 2026-07-23 | 来源

V3P-003032 | codetables-gf4-214-9

Minimum distance of a [214, 9] linear code over GF(4)

codes are known with d = 146, and no code can have d greater than 153.

来源状态 open | 2026-07-23 | 来源

V3P-003033 | codetables-gf4-215-6

Minimum distance of a [215, 6] linear code over GF(4)

codes are known with d = 156, and no code can have d greater than 159.

来源状态 open | 2026-07-23 | 来源

V3P-003034 | codetables-gf4-215-7

Minimum distance of a [215, 7] linear code over GF(4)

codes are known with d = 152, and no code can have d greater than 158.

来源状态 open | 2026-07-23 | 来源

V3P-003035 | codetables-gf4-215-8

Minimum distance of a [215, 8] linear code over GF(4)

codes are known with d = 149, and no code can have d greater than 155.

来源状态 open | 2026-07-23 | 来源

V3P-003036 | codetables-gf4-215-9

Minimum distance of a [215, 9] linear code over GF(4)

codes are known with d = 147, and no code can have d greater than 154.

来源状态 open | 2026-07-23 | 来源

V3P-003037 | codetables-gf4-216-6

Minimum distance of a [216, 6] linear code over GF(4)

codes are known with d = 157, and no code can have d greater than 160.

来源状态 open | 2026-07-23 | 来源

V3P-003038 | codetables-gf4-216-7

Minimum distance of a [216, 7] linear code over GF(4)

codes are known with d = 153, and no code can have d greater than 159.

来源状态 open | 2026-07-23 | 来源

V3P-003039 | codetables-gf4-216-8

Minimum distance of a [216, 8] linear code over GF(4)

codes are known with d = 150, and no code can have d greater than 156.

来源状态 open | 2026-07-23 | 来源

V3P-003040 | codetables-gf4-216-9

Minimum distance of a [216, 9] linear code over GF(4)

codes are known with d = 148, and no code can have d greater than 155.

来源状态 open | 2026-07-23 | 来源

V3P-003041 | codetables-gf4-217-6

Minimum distance of a [217, 6] linear code over GF(4)

codes are known with d = 158, and no code can have d greater than 160.

来源状态 open | 2026-07-23 | 来源

V3P-003042 | codetables-gf4-217-7

Minimum distance of a [217, 7] linear code over GF(4)

codes are known with d = 154, and no code can have d greater than 160.

来源状态 open | 2026-07-23 | 来源

V3P-003043 | codetables-gf4-217-8

Minimum distance of a [217, 8] linear code over GF(4)

codes are known with d = 151, and no code can have d greater than 156.

来源状态 open | 2026-07-23 | 来源

V3P-003044 | codetables-gf4-217-9

Minimum distance of a [217, 9] linear code over GF(4)

codes are known with d = 148, and no code can have d greater than 156.

来源状态 open | 2026-07-23 | 来源

V3P-003045 | codetables-gf4-218-6

Minimum distance of a [218, 6] linear code over GF(4)

codes are known with d = 159, and no code can have d greater than 160.

来源状态 open | 2026-07-23 | 来源

V3P-003046 | codetables-gf4-218-7

Minimum distance of a [218, 7] linear code over GF(4)

codes are known with d = 155, and no code can have d greater than 160.

来源状态 open | 2026-07-23 | 来源

V3P-003047 | codetables-gf4-218-8

Minimum distance of a [218, 8] linear code over GF(4)

codes are known with d = 152, and no code can have d greater than 157.

来源状态 open | 2026-07-23 | 来源

V3P-003048 | codetables-gf4-218-9

Minimum distance of a [218, 9] linear code over GF(4)

codes are known with d = 148, and no code can have d greater than 156.

来源状态 open | 2026-07-23 | 来源

V3P-003049 | codetables-gf4-219-6

Minimum distance of a [219, 6] linear code over GF(4)

codes are known with d = 160, and no code can have d greater than 161.

来源状态 open | 2026-07-23 | 来源

V3P-003050 | codetables-gf4-219-7

Minimum distance of a [219, 7] linear code over GF(4)

codes are known with d = 156, and no code can have d greater than 160.

来源状态 open | 2026-07-23 | 来源

V3P-003051 | codetables-gf4-219-8

Minimum distance of a [219, 8] linear code over GF(4)

codes are known with d = 152, and no code can have d greater than 158.

来源状态 open | 2026-07-23 | 来源

V3P-003052 | codetables-gf4-219-9

Minimum distance of a [219, 9] linear code over GF(4)

codes are known with d = 149, and no code can have d greater than 157.

来源状态 open | 2026-07-23 | 来源

V3P-003053 | codetables-gf4-22-6

Minimum distance of a [22, 6] linear code over GF(4)

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-003054 | codetables-gf4-220-6

Minimum distance of a [220, 6] linear code over GF(4)

codes are known with d = 160, and no code can have d greater than 162.

来源状态 open | 2026-07-23 | 来源

V3P-003055 | codetables-gf4-220-7

Minimum distance of a [220, 7] linear code over GF(4)

codes are known with d = 156, and no code can have d greater than 160.

来源状态 open | 2026-07-23 | 来源

V3P-003056 | codetables-gf4-220-8

Minimum distance of a [220, 8] linear code over GF(4)

codes are known with d = 153, and no code can have d greater than 159.

来源状态 open | 2026-07-23 | 来源

V3P-003057 | codetables-gf4-220-9

Minimum distance of a [220, 9] linear code over GF(4)

codes are known with d = 150, and no code can have d greater than 158.

来源状态 open | 2026-07-23 | 来源

V3P-003058 | codetables-gf4-221-6

Minimum distance of a [221, 6] linear code over GF(4)

codes are known with d = 160, and no code can have d greater than 163.

来源状态 open | 2026-07-23 | 来源

V3P-003059 | codetables-gf4-221-7

Minimum distance of a [221, 7] linear code over GF(4)

codes are known with d = 157, and no code can have d greater than 161.

来源状态 open | 2026-07-23 | 来源

V3P-003060 | codetables-gf4-221-8

Minimum distance of a [221, 8] linear code over GF(4)

codes are known with d = 154, and no code can have d greater than 159.

来源状态 open | 2026-07-23 | 来源

V3P-003061 | codetables-gf4-221-9

Minimum distance of a [221, 9] linear code over GF(4)

codes are known with d = 151, and no code can have d greater than 159.

来源状态 open | 2026-07-23 | 来源

V3P-003062 | codetables-gf4-222-6

Minimum distance of a [222, 6] linear code over GF(4)

codes are known with d = 161, and no code can have d greater than 164.

来源状态 open | 2026-07-23 | 来源

V3P-003063 | codetables-gf4-222-7

Minimum distance of a [222, 7] linear code over GF(4)

codes are known with d = 158, and no code can have d greater than 162.

来源状态 open | 2026-07-23 | 来源

V3P-003064 | codetables-gf4-222-8

Minimum distance of a [222, 8] linear code over GF(4)

codes are known with d = 154, and no code can have d greater than 160.

来源状态 open | 2026-07-23 | 来源

V3P-003065 | codetables-gf4-222-9

Minimum distance of a [222, 9] linear code over GF(4)

codes are known with d = 152, and no code can have d greater than 159.

来源状态 open | 2026-07-23 | 来源

V3P-003066 | codetables-gf4-223-6

Minimum distance of a [223, 6] linear code over GF(4)

codes are known with d = 162, and no code can have d greater than 164.

来源状态 open | 2026-07-23 | 来源

V3P-003067 | codetables-gf4-223-7

Minimum distance of a [223, 7] linear code over GF(4)

codes are known with d = 159, and no code can have d greater than 163.

来源状态 open | 2026-07-23 | 来源

V3P-003068 | codetables-gf4-223-8

Minimum distance of a [223, 8] linear code over GF(4)

codes are known with d = 155, and no code can have d greater than 161.

来源状态 open | 2026-07-23 | 来源

V3P-003069 | codetables-gf4-223-9

Minimum distance of a [223, 9] linear code over GF(4)

codes are known with d = 152, and no code can have d greater than 160.

来源状态 open | 2026-07-23 | 来源

V3P-003070 | codetables-gf4-224-6

Minimum distance of a [224, 6] linear code over GF(4)

codes are known with d = 163, and no code can have d greater than 165.

来源状态 open | 2026-07-23 | 来源

V3P-003071 | codetables-gf4-224-7

Minimum distance of a [224, 7] linear code over GF(4)

codes are known with d = 160, and no code can have d greater than 164.

来源状态 open | 2026-07-23 | 来源

V3P-003072 | codetables-gf4-224-8

Minimum distance of a [224, 8] linear code over GF(4)

codes are known with d = 156, and no code can have d greater than 162.

来源状态 open | 2026-07-23 | 来源

V3P-003073 | codetables-gf4-224-9

Minimum distance of a [224, 9] linear code over GF(4)

codes are known with d = 152, and no code can have d greater than 161.

来源状态 open | 2026-07-23 | 来源

V3P-003074 | codetables-gf4-225-6

Minimum distance of a [225, 6] linear code over GF(4)

codes are known with d = 164, and no code can have d greater than 166.

来源状态 open | 2026-07-23 | 来源

V3P-003075 | codetables-gf4-225-7

Minimum distance of a [225, 7] linear code over GF(4)

codes are known with d = 160, and no code can have d greater than 164.

来源状态 open | 2026-07-23 | 来源

V3P-003076 | codetables-gf4-225-8

Minimum distance of a [225, 8] linear code over GF(4)

codes are known with d = 156, and no code can have d greater than 162.

来源状态 open | 2026-07-23 | 来源

V3P-003077 | codetables-gf4-225-9

Minimum distance of a [225, 9] linear code over GF(4)

codes are known with d = 153, and no code can have d greater than 162.

来源状态 open | 2026-07-23 | 来源

V3P-003078 | codetables-gf4-226-6

Minimum distance of a [226, 6] linear code over GF(4)

codes are known with d = 164, and no code can have d greater than 167.

来源状态 open | 2026-07-23 | 来源

V3P-003079 | codetables-gf4-226-7

Minimum distance of a [226, 7] linear code over GF(4)

codes are known with d = 161, and no code can have d greater than 165.

来源状态 open | 2026-07-23 | 来源

V3P-003080 | codetables-gf4-226-8

Minimum distance of a [226, 8] linear code over GF(4)

codes are known with d = 157, and no code can have d greater than 163.

来源状态 open | 2026-07-23 | 来源

V3P-003081 | codetables-gf4-226-9

Minimum distance of a [226, 9] linear code over GF(4)

codes are known with d = 154, and no code can have d greater than 162.

来源状态 open | 2026-07-23 | 来源

V3P-003082 | codetables-gf4-227-6

Minimum distance of a [227, 6] linear code over GF(4)

codes are known with d = 165, and no code can have d greater than 168.

来源状态 open | 2026-07-23 | 来源

V3P-003083 | codetables-gf4-227-7

Minimum distance of a [227, 7] linear code over GF(4)

codes are known with d = 162, and no code can have d greater than 166.

来源状态 open | 2026-07-23 | 来源

V3P-003084 | codetables-gf4-227-8

Minimum distance of a [227, 8] linear code over GF(4)

codes are known with d = 158, and no code can have d greater than 164.

来源状态 open | 2026-07-23 | 来源

V3P-003085 | codetables-gf4-227-9

Minimum distance of a [227, 9] linear code over GF(4)

codes are known with d = 155, and no code can have d greater than 163.

来源状态 open | 2026-07-23 | 来源

V3P-003086 | codetables-gf4-228-6

Minimum distance of a [228, 6] linear code over GF(4)

codes are known with d = 166, and no code can have d greater than 168.

来源状态 open | 2026-07-23 | 来源

V3P-003087 | codetables-gf4-228-7

Minimum distance of a [228, 7] linear code over GF(4)

codes are known with d = 162, and no code can have d greater than 167.

来源状态 open | 2026-07-23 | 来源

V3P-003088 | codetables-gf4-228-8

Minimum distance of a [228, 8] linear code over GF(4)

codes are known with d = 159, and no code can have d greater than 165.

来源状态 open | 2026-07-23 | 来源

V3P-003089 | codetables-gf4-228-9

Minimum distance of a [228, 9] linear code over GF(4)

codes are known with d = 156, and no code can have d greater than 164.

来源状态 open | 2026-07-23 | 来源

V3P-003090 | codetables-gf4-229-6

Minimum distance of a [229, 6] linear code over GF(4)

codes are known with d = 167, and no code can have d greater than 169.

来源状态 open | 2026-07-23 | 来源

V3P-003091 | codetables-gf4-229-7

Minimum distance of a [229, 7] linear code over GF(4)

codes are known with d = 163, and no code can have d greater than 168.

来源状态 open | 2026-07-23 | 来源

V3P-003092 | codetables-gf4-229-8

Minimum distance of a [229, 8] linear code over GF(4)

codes are known with d = 160, and no code can have d greater than 165.

来源状态 open | 2026-07-23 | 来源

V3P-003093 | codetables-gf4-229-9

Minimum distance of a [229, 9] linear code over GF(4)

codes are known with d = 156, and no code can have d greater than 165.

来源状态 open | 2026-07-23 | 来源

V3P-003094 | codetables-gf4-23-7

Minimum distance of a [23, 7] linear code over GF(4)

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-003095 | codetables-gf4-230-6

Minimum distance of a [230, 6] linear code over GF(4)

codes are known with d = 168, and no code can have d greater than 170.

来源状态 open | 2026-07-23 | 来源

V3P-003096 | codetables-gf4-230-7

Minimum distance of a [230, 7] linear code over GF(4)

codes are known with d = 164, and no code can have d greater than 168.

来源状态 open | 2026-07-23 | 来源

V3P-003097 | codetables-gf4-230-8

Minimum distance of a [230, 8] linear code over GF(4)

codes are known with d = 160, and no code can have d greater than 166.

来源状态 open | 2026-07-23 | 来源

V3P-003098 | codetables-gf4-230-9

Minimum distance of a [230, 9] linear code over GF(4)

codes are known with d = 157, and no code can have d greater than 165.

来源状态 open | 2026-07-23 | 来源

V3P-003099 | codetables-gf4-231-6

Minimum distance of a [231, 6] linear code over GF(4)

codes are known with d = 168, and no code can have d greater than 171.

来源状态 open | 2026-07-23 | 来源

V3P-003100 | codetables-gf4-231-7

Minimum distance of a [231, 7] linear code over GF(4)

codes are known with d = 165, and no code can have d greater than 169.

来源状态 open | 2026-07-23 | 来源

V3P-003101 | codetables-gf4-231-8

Minimum distance of a [231, 8] linear code over GF(4)

codes are known with d = 161, and no code can have d greater than 167.

来源状态 open | 2026-07-23 | 来源

V3P-003102 | codetables-gf4-231-9

Minimum distance of a [231, 9] linear code over GF(4)

codes are known with d = 158, and no code can have d greater than 166.

来源状态 open | 2026-07-23 | 来源

V3P-003103 | codetables-gf4-232-6

Minimum distance of a [232, 6] linear code over GF(4)

codes are known with d = 169, and no code can have d greater than 172.

来源状态 open | 2026-07-23 | 来源

V3P-003104 | codetables-gf4-232-7

Minimum distance of a [232, 7] linear code over GF(4)

codes are known with d = 166, and no code can have d greater than 170.

来源状态 open | 2026-07-23 | 来源

V3P-003105 | codetables-gf4-232-8

Minimum distance of a [232, 8] linear code over GF(4)

codes are known with d = 162, and no code can have d greater than 168.

来源状态 open | 2026-07-23 | 来源

V3P-003106 | codetables-gf4-232-9

Minimum distance of a [232, 9] linear code over GF(4)

codes are known with d = 159, and no code can have d greater than 167.

来源状态 open | 2026-07-23 | 来源

V3P-003107 | codetables-gf4-233-6

Minimum distance of a [233, 6] linear code over GF(4)

codes are known with d = 170, and no code can have d greater than 172.

来源状态 open | 2026-07-23 | 来源

V3P-003108 | codetables-gf4-233-7

Minimum distance of a [233, 7] linear code over GF(4)

codes are known with d = 167, and no code can have d greater than 171.

来源状态 open | 2026-07-23 | 来源

V3P-003109 | codetables-gf4-233-8

Minimum distance of a [233, 8] linear code over GF(4)

codes are known with d = 163, and no code can have d greater than 168.

来源状态 open | 2026-07-23 | 来源

V3P-003110 | codetables-gf4-233-9

Minimum distance of a [233, 9] linear code over GF(4)

codes are known with d = 160, and no code can have d greater than 168.

来源状态 open | 2026-07-23 | 来源

V3P-003111 | codetables-gf4-234-6

Minimum distance of a [234, 6] linear code over GF(4)

codes are known with d = 171, and no code can have d greater than 173.

来源状态 open | 2026-07-23 | 来源

V3P-003112 | codetables-gf4-234-7

Minimum distance of a [234, 7] linear code over GF(4)

codes are known with d = 168, and no code can have d greater than 172.

来源状态 open | 2026-07-23 | 来源

V3P-003113 | codetables-gf4-234-8

Minimum distance of a [234, 8] linear code over GF(4)

codes are known with d = 164, and no code can have d greater than 169.

来源状态 open | 2026-07-23 | 来源

V3P-003114 | codetables-gf4-234-9

Minimum distance of a [234, 9] linear code over GF(4)

codes are known with d = 160, and no code can have d greater than 168.

来源状态 open | 2026-07-23 | 来源

V3P-003115 | codetables-gf4-235-6

Minimum distance of a [235, 6] linear code over GF(4)

codes are known with d = 172, and no code can have d greater than 174.

来源状态 open | 2026-07-23 | 来源

V3P-003116 | codetables-gf4-235-7

Minimum distance of a [235, 7] linear code over GF(4)

codes are known with d = 168, and no code can have d greater than 172.

来源状态 open | 2026-07-23 | 来源

V3P-003117 | codetables-gf4-235-8

Minimum distance of a [235, 8] linear code over GF(4)

codes are known with d = 164, and no code can have d greater than 170.

来源状态 open | 2026-07-23 | 来源

V3P-003118 | codetables-gf4-235-9

Minimum distance of a [235, 9] linear code over GF(4)

codes are known with d = 160, and no code can have d greater than 169.

来源状态 open | 2026-07-23 | 来源

V3P-003119 | codetables-gf4-236-6

Minimum distance of a [236, 6] linear code over GF(4)

codes are known with d = 172, and no code can have d greater than 175.

来源状态 open | 2026-07-23 | 来源

V3P-003120 | codetables-gf4-236-7

Minimum distance of a [236, 7] linear code over GF(4)

codes are known with d = 169, and no code can have d greater than 173.

来源状态 open | 2026-07-23 | 来源

V3P-003121 | codetables-gf4-236-8

Minimum distance of a [236, 8] linear code over GF(4)

codes are known with d = 164, and no code can have d greater than 171.

来源状态 open | 2026-07-23 | 来源

V3P-003122 | codetables-gf4-236-9

Minimum distance of a [236, 9] linear code over GF(4)

codes are known with d = 161, and no code can have d greater than 170.

来源状态 open | 2026-07-23 | 来源

V3P-003123 | codetables-gf4-237-6

Minimum distance of a [237, 6] linear code over GF(4)

codes are known with d = 173, and no code can have d greater than 176.

来源状态 open | 2026-07-23 | 来源

V3P-003124 | codetables-gf4-237-7

Minimum distance of a [237, 7] linear code over GF(4)

codes are known with d = 170, and no code can have d greater than 174.

来源状态 open | 2026-07-23 | 来源

V3P-003125 | codetables-gf4-237-8

Minimum distance of a [237, 8] linear code over GF(4)

codes are known with d = 165, and no code can have d greater than 171.

来源状态 open | 2026-07-23 | 来源

V3P-003126 | codetables-gf4-237-9

Minimum distance of a [237, 9] linear code over GF(4)

codes are known with d = 162, and no code can have d greater than 171.

来源状态 open | 2026-07-23 | 来源

V3P-003127 | codetables-gf4-238-5

Minimum distance of a [238, 5] linear code over GF(4)

codes are known with d = 176, and no code can have d greater than 177.

来源状态 open | 2026-07-23 | 来源

V3P-003128 | codetables-gf4-238-6

Minimum distance of a [238, 6] linear code over GF(4)

codes are known with d = 174, and no code can have d greater than 176.

来源状态 open | 2026-07-23 | 来源

V3P-003129 | codetables-gf4-238-7

Minimum distance of a [238, 7] linear code over GF(4)

codes are known with d = 171, and no code can have d greater than 175.

来源状态 open | 2026-07-23 | 来源

V3P-003130 | codetables-gf4-238-8

Minimum distance of a [238, 8] linear code over GF(4)

codes are known with d = 166, and no code can have d greater than 172.

来源状态 open | 2026-07-23 | 来源

V3P-003131 | codetables-gf4-238-9

Minimum distance of a [238, 9] linear code over GF(4)

codes are known with d = 163, and no code can have d greater than 171.

来源状态 open | 2026-07-23 | 来源

V3P-003132 | codetables-gf4-239-6

Minimum distance of a [239, 6] linear code over GF(4)

codes are known with d = 175, and no code can have d greater than 176.

来源状态 open | 2026-07-23 | 来源

V3P-003133 | codetables-gf4-239-7

Minimum distance of a [239, 7] linear code over GF(4)

codes are known with d = 172, and no code can have d greater than 176.

来源状态 open | 2026-07-23 | 来源

V3P-003134 | codetables-gf4-239-8

Minimum distance of a [239, 8] linear code over GF(4)

codes are known with d = 167, and no code can have d greater than 173.

来源状态 open | 2026-07-23 | 来源

V3P-003135 | codetables-gf4-239-9

Minimum distance of a [239, 9] linear code over GF(4)

codes are known with d = 164, and no code can have d greater than 172.

来源状态 open | 2026-07-23 | 来源

V3P-003136 | codetables-gf4-24-8

Minimum distance of a [24, 8] linear code over GF(4)

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-003137 | codetables-gf4-240-6

Minimum distance of a [240, 6] linear code over GF(4)

codes are known with d = 176, and no code can have d greater than 177.

来源状态 open | 2026-07-23 | 来源

V3P-003138 | codetables-gf4-240-7

Minimum distance of a [240, 7] linear code over GF(4)

codes are known with d = 173, and no code can have d greater than 176.

来源状态 open | 2026-07-23 | 来源

V3P-003139 | codetables-gf4-240-8

Minimum distance of a [240, 8] linear code over GF(4)

codes are known with d = 168, and no code can have d greater than 174.

来源状态 open | 2026-07-23 | 来源

V3P-003140 | codetables-gf4-240-9

Minimum distance of a [240, 9] linear code over GF(4)

codes are known with d = 164, and no code can have d greater than 173.

来源状态 open | 2026-07-23 | 来源

V3P-003141 | codetables-gf4-241-6

Minimum distance of a [241, 6] linear code over GF(4)

codes are known with d = 176, and no code can have d greater than 178.

来源状态 open | 2026-07-23 | 来源

V3P-003142 | codetables-gf4-241-7

Minimum distance of a [241, 7] linear code over GF(4)

codes are known with d = 174, and no code can have d greater than 176.

来源状态 open | 2026-07-23 | 来源

V3P-003143 | codetables-gf4-241-8

Minimum distance of a [241, 8] linear code over GF(4)

codes are known with d = 169, and no code can have d greater than 174.

来源状态 open | 2026-07-23 | 来源

V3P-003144 | codetables-gf4-241-9

Minimum distance of a [241, 9] linear code over GF(4)

codes are known with d = 165, and no code can have d greater than 174.

来源状态 open | 2026-07-23 | 来源

V3P-003145 | codetables-gf4-242-6

Minimum distance of a [242, 6] linear code over GF(4)

codes are known with d = 177, and no code can have d greater than 179.

来源状态 open | 2026-07-23 | 来源

V3P-003146 | codetables-gf4-242-7

Minimum distance of a [242, 7] linear code over GF(4)

codes are known with d = 175, and no code can have d greater than 177.

来源状态 open | 2026-07-23 | 来源

V3P-003147 | codetables-gf4-242-8

Minimum distance of a [242, 8] linear code over GF(4)

codes are known with d = 170, and no code can have d greater than 175.

来源状态 open | 2026-07-23 | 来源

V3P-003148 | codetables-gf4-242-9

Minimum distance of a [242, 9] linear code over GF(4)

codes are known with d = 166, and no code can have d greater than 174.

来源状态 open | 2026-07-23 | 来源

V3P-003149 | codetables-gf4-243-6

Minimum distance of a [243, 6] linear code over GF(4)

codes are known with d = 178, and no code can have d greater than 180.

来源状态 open | 2026-07-23 | 来源

V3P-003150 | codetables-gf4-243-7

Minimum distance of a [243, 7] linear code over GF(4)

codes are known with d = 175, and no code can have d greater than 178.

来源状态 open | 2026-07-23 | 来源

V3P-003151 | codetables-gf4-243-8

Minimum distance of a [243, 8] linear code over GF(4)

codes are known with d = 171, and no code can have d greater than 176.

来源状态 open | 2026-07-23 | 来源

V3P-003152 | codetables-gf4-243-9

Minimum distance of a [243, 9] linear code over GF(4)

codes are known with d = 167, and no code can have d greater than 175.

来源状态 open | 2026-07-23 | 来源

V3P-003153 | codetables-gf4-244-6

Minimum distance of a [244, 6] linear code over GF(4)

codes are known with d = 179, and no code can have d greater than 180.

来源状态 open | 2026-07-23 | 来源

V3P-003154 | codetables-gf4-244-7

Minimum distance of a [244, 7] linear code over GF(4)

codes are known with d = 176, and no code can have d greater than 179.

来源状态 open | 2026-07-23 | 来源

V3P-003155 | codetables-gf4-244-8

Minimum distance of a [244, 8] linear code over GF(4)

codes are known with d = 172, and no code can have d greater than 177.

来源状态 open | 2026-07-23 | 来源

V3P-003156 | codetables-gf4-244-9

Minimum distance of a [244, 9] linear code over GF(4)

codes are known with d = 168, and no code can have d greater than 176.

来源状态 open | 2026-07-23 | 来源

V3P-003157 | codetables-gf4-245-6

Minimum distance of a [245, 6] linear code over GF(4)

codes are known with d = 180, and no code can have d greater than 181.

来源状态 open | 2026-07-23 | 来源

V3P-003158 | codetables-gf4-245-7

Minimum distance of a [245, 7] linear code over GF(4)

codes are known with d = 177, and no code can have d greater than 180.

来源状态 open | 2026-07-23 | 来源

V3P-003159 | codetables-gf4-245-8

Minimum distance of a [245, 8] linear code over GF(4)

codes are known with d = 172, and no code can have d greater than 177.

来源状态 open | 2026-07-23 | 来源

V3P-003160 | codetables-gf4-245-9

Minimum distance of a [245, 9] linear code over GF(4)

codes are known with d = 168, and no code can have d greater than 176.

来源状态 open | 2026-07-23 | 来源

V3P-003161 | codetables-gf4-246-6

Minimum distance of a [246, 6] linear code over GF(4)

codes are known with d = 180, and no code can have d greater than 182.

来源状态 open | 2026-07-23 | 来源

V3P-003162 | codetables-gf4-246-7

Minimum distance of a [246, 7] linear code over GF(4)

codes are known with d = 178, and no code can have d greater than 180.

来源状态 open | 2026-07-23 | 来源

V3P-003163 | codetables-gf4-246-8

Minimum distance of a [246, 8] linear code over GF(4)

codes are known with d = 173, and no code can have d greater than 178.

来源状态 open | 2026-07-23 | 来源

V3P-003164 | codetables-gf4-246-9

Minimum distance of a [246, 9] linear code over GF(4)

codes are known with d = 169, and no code can have d greater than 177.

来源状态 open | 2026-07-23 | 来源

V3P-003165 | codetables-gf4-247-6

Minimum distance of a [247, 6] linear code over GF(4)

codes are known with d = 181, and no code can have d greater than 183.

来源状态 open | 2026-07-23 | 来源

V3P-003166 | codetables-gf4-247-7

Minimum distance of a [247, 7] linear code over GF(4)

codes are known with d = 179, and no code can have d greater than 181.

来源状态 open | 2026-07-23 | 来源

V3P-003167 | codetables-gf4-247-8

Minimum distance of a [247, 8] linear code over GF(4)

codes are known with d = 174, and no code can have d greater than 179.

来源状态 open | 2026-07-23 | 来源

V3P-003168 | codetables-gf4-247-9

Minimum distance of a [247, 9] linear code over GF(4)

codes are known with d = 170, and no code can have d greater than 178.

来源状态 open | 2026-07-23 | 来源

V3P-003169 | codetables-gf4-248-6

Minimum distance of a [248, 6] linear code over GF(4)

codes are known with d = 182, and no code can have d greater than 184.

来源状态 open | 2026-07-23 | 来源

V3P-003170 | codetables-gf4-248-7

Minimum distance of a [248, 7] linear code over GF(4)

codes are known with d = 180, and no code can have d greater than 182.

来源状态 open | 2026-07-23 | 来源

V3P-003171 | codetables-gf4-248-8

Minimum distance of a [248, 8] linear code over GF(4)

codes are known with d = 175, and no code can have d greater than 180.

来源状态 open | 2026-07-23 | 来源

V3P-003172 | codetables-gf4-248-9

Minimum distance of a [248, 9] linear code over GF(4)

codes are known with d = 171, and no code can have d greater than 179.

来源状态 open | 2026-07-23 | 来源

V3P-003173 | codetables-gf4-249-6

Minimum distance of a [249, 6] linear code over GF(4)

codes are known with d = 183, and no code can have d greater than 184.

来源状态 open | 2026-07-23 | 来源

V3P-003174 | codetables-gf4-249-7

Minimum distance of a [249, 7] linear code over GF(4)

codes are known with d = 181, and no code can have d greater than 183.

来源状态 open | 2026-07-23 | 来源

V3P-003175 | codetables-gf4-249-8

Minimum distance of a [249, 8] linear code over GF(4)

codes are known with d = 176, and no code can have d greater than 180.

来源状态 open | 2026-07-23 | 来源

V3P-003176 | codetables-gf4-249-9

Minimum distance of a [249, 9] linear code over GF(4)

codes are known with d = 172, and no code can have d greater than 180.

来源状态 open | 2026-07-23 | 来源

V3P-003177 | codetables-gf4-25-8

Minimum distance of a [25, 8] linear code over GF(4)

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-003178 | codetables-gf4-25-9

Minimum distance of a [25, 9] linear code over GF(4)

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-003179 | codetables-gf4-250-6

Minimum distance of a [250, 6] linear code over GF(4)

codes are known with d = 184, and no code can have d greater than 185.

来源状态 open | 2026-07-23 | 来源

V3P-003180 | codetables-gf4-250-7

Minimum distance of a [250, 7] linear code over GF(4)

codes are known with d = 182, and no code can have d greater than 184.

来源状态 open | 2026-07-23 | 来源

V3P-003181 | codetables-gf4-250-8

Minimum distance of a [250, 8] linear code over GF(4)

codes are known with d = 176, and no code can have d greater than 181.

来源状态 open | 2026-07-23 | 来源

V3P-003182 | codetables-gf4-250-9

Minimum distance of a [250, 9] linear code over GF(4)

codes are known with d = 172, and no code can have d greater than 180.

来源状态 open | 2026-07-23 | 来源

V3P-003183 | codetables-gf4-251-6

Minimum distance of a [251, 6] linear code over GF(4)

codes are known with d = 184, and no code can have d greater than 186.

来源状态 open | 2026-07-23 | 来源

V3P-003184 | codetables-gf4-251-7

Minimum distance of a [251, 7] linear code over GF(4)

codes are known with d = 183, and no code can have d greater than 184.

来源状态 open | 2026-07-23 | 来源

V3P-003185 | codetables-gf4-251-8

Minimum distance of a [251, 8] linear code over GF(4)

codes are known with d = 176, and no code can have d greater than 182.

来源状态 open | 2026-07-23 | 来源

V3P-003186 | codetables-gf4-251-9

Minimum distance of a [251, 9] linear code over GF(4)

codes are known with d = 173, and no code can have d greater than 181.

来源状态 open | 2026-07-23 | 来源

V3P-003187 | codetables-gf4-252-6

Minimum distance of a [252, 6] linear code over GF(4)

codes are known with d = 185, and no code can have d greater than 187.

来源状态 open | 2026-07-23 | 来源

V3P-003188 | codetables-gf4-252-7

Minimum distance of a [252, 7] linear code over GF(4)

codes are known with d = 184, and no code can have d greater than 185.

来源状态 open | 2026-07-23 | 来源

V3P-003189 | codetables-gf4-252-8

Minimum distance of a [252, 8] linear code over GF(4)

codes are known with d = 176, and no code can have d greater than 183.

来源状态 open | 2026-07-23 | 来源

V3P-003190 | codetables-gf4-252-9

Minimum distance of a [252, 9] linear code over GF(4)

codes are known with d = 174, and no code can have d greater than 182.

来源状态 open | 2026-07-23 | 来源

V3P-003191 | codetables-gf4-253-6

Minimum distance of a [253, 6] linear code over GF(4)

codes are known with d = 186, and no code can have d greater than 188.

来源状态 open | 2026-07-23 | 来源

V3P-003192 | codetables-gf4-253-7

Minimum distance of a [253, 7] linear code over GF(4)

codes are known with d = 185, and no code can have d greater than 186.

来源状态 open | 2026-07-23 | 来源

V3P-003193 | codetables-gf4-253-8

Minimum distance of a [253, 8] linear code over GF(4)

codes are known with d = 177, and no code can have d greater than 184.

来源状态 open | 2026-07-23 | 来源

V3P-003194 | codetables-gf4-253-9

Minimum distance of a [253, 9] linear code over GF(4)

codes are known with d = 175, and no code can have d greater than 183.

来源状态 open | 2026-07-23 | 来源

V3P-003195 | codetables-gf4-254-6

Minimum distance of a [254, 6] linear code over GF(4)

codes are known with d = 187, and no code can have d greater than 188.

来源状态 open | 2026-07-23 | 来源

V3P-003196 | codetables-gf4-254-7

Minimum distance of a [254, 7] linear code over GF(4)

codes are known with d = 186, and no code can have d greater than 187.

来源状态 open | 2026-07-23 | 来源

V3P-003197 | codetables-gf4-254-8

Minimum distance of a [254, 8] linear code over GF(4)

codes are known with d = 178, and no code can have d greater than 185.

来源状态 open | 2026-07-23 | 来源

V3P-003198 | codetables-gf4-254-9

Minimum distance of a [254, 9] linear code over GF(4)

codes are known with d = 176, and no code can have d greater than 184.

来源状态 open | 2026-07-23 | 来源

V3P-003199 | codetables-gf4-255-6

Minimum distance of a [255, 6] linear code over GF(4)

codes are known with d = 188, and no code can have d greater than 189.

来源状态 open | 2026-07-23 | 来源

V3P-003200 | codetables-gf4-255-7

Minimum distance of a [255, 7] linear code over GF(4)

codes are known with d = 187, and no code can have d greater than 188.

来源状态 open | 2026-07-23 | 来源

V3P-003201 | codetables-gf4-255-8

Minimum distance of a [255, 8] linear code over GF(4)

codes are known with d = 179, and no code can have d greater than 186.

来源状态 open | 2026-07-23 | 来源

V3P-003202 | codetables-gf4-255-9

Minimum distance of a [255, 9] linear code over GF(4)

codes are known with d = 176, and no code can have d greater than 185.

来源状态 open | 2026-07-23 | 来源

V3P-003203 | codetables-gf4-256-6

Minimum distance of a [256, 6] linear code over GF(4)

codes are known with d = 188, and no code can have d greater than 190.

来源状态 open | 2026-07-23 | 来源

V3P-003204 | codetables-gf4-256-8

Minimum distance of a [256, 8] linear code over GF(4)

codes are known with d = 180, and no code can have d greater than 187.

来源状态 open | 2026-07-23 | 来源

V3P-003205 | codetables-gf4-256-9

Minimum distance of a [256, 9] linear code over GF(4)

codes are known with d = 176, and no code can have d greater than 186.

来源状态 open | 2026-07-23 | 来源

V3P-003206 | codetables-gf4-26-7

Minimum distance of a [26, 7] linear code over GF(4)

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-003207 | codetables-gf4-26-8

Minimum distance of a [26, 8] linear code over GF(4)

codes are known with d = 13, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-003208 | codetables-gf4-26-9

Minimum distance of a [26, 9] linear code over GF(4)

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-003209 | codetables-gf4-27-7

Minimum distance of a [27, 7] linear code over GF(4)

codes are known with d = 15, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-003210 | codetables-gf4-27-8

Minimum distance of a [27, 8] linear code over GF(4)

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-003211 | codetables-gf4-27-9

Minimum distance of a [27, 9] linear code over GF(4)

codes are known with d = 13, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-003212 | codetables-gf4-28-8

Minimum distance of a [28, 8] linear code over GF(4)

codes are known with d = 15, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-003213 | codetables-gf4-28-9

Minimum distance of a [28, 9] linear code over GF(4)

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-003214 | codetables-gf4-29-7

Minimum distance of a [29, 7] linear code over GF(4)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-003215 | codetables-gf4-29-9

Minimum distance of a [29, 9] linear code over GF(4)

codes are known with d = 15, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-003216 | codetables-gf4-30-6

Minimum distance of a [30, 6] linear code over GF(4)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-003217 | codetables-gf4-30-7

Minimum distance of a [30, 7] linear code over GF(4)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-003218 | codetables-gf4-30-8

Minimum distance of a [30, 8] linear code over GF(4)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-003219 | codetables-gf4-31-6

Minimum distance of a [31, 6] linear code over GF(4)

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-003220 | codetables-gf4-31-7

Minimum distance of a [31, 7] linear code over GF(4)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-003221 | codetables-gf4-31-8

Minimum distance of a [31, 8] linear code over GF(4)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-003222 | codetables-gf4-31-9

Minimum distance of a [31, 9] linear code over GF(4)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-003223 | codetables-gf4-32-7

Minimum distance of a [32, 7] linear code over GF(4)

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-003224 | codetables-gf4-32-8

Minimum distance of a [32, 8] linear code over GF(4)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-003225 | codetables-gf4-32-9

Minimum distance of a [32, 9] linear code over GF(4)

codes are known with d = 16, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-003226 | codetables-gf4-33-6

Minimum distance of a [33, 6] linear code over GF(4)

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-003227 | codetables-gf4-33-8

Minimum distance of a [33, 8] linear code over GF(4)

codes are known with d = 18, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-003228 | codetables-gf4-33-9

Minimum distance of a [33, 9] linear code over GF(4)

codes are known with d = 17, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-003229 | codetables-gf4-34-6

Minimum distance of a [34, 6] linear code over GF(4)

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-003230 | codetables-gf4-34-7

Minimum distance of a [34, 7] linear code over GF(4)

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-003231 | codetables-gf4-34-8

Minimum distance of a [34, 8] linear code over GF(4)

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-003232 | codetables-gf4-34-9

Minimum distance of a [34, 9] linear code over GF(4)

codes are known with d = 18, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-003233 | codetables-gf4-35-6

Minimum distance of a [35, 6] linear code over GF(4)

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-003234 | codetables-gf4-35-7

Minimum distance of a [35, 7] linear code over GF(4)

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-003235 | codetables-gf4-35-8

Minimum distance of a [35, 8] linear code over GF(4)

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-003236 | codetables-gf4-35-9

Minimum distance of a [35, 9] linear code over GF(4)

codes are known with d = 18, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-003237 | codetables-gf4-36-6

Minimum distance of a [36, 6] linear code over GF(4)

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-003238 | codetables-gf4-36-7

Minimum distance of a [36, 7] linear code over GF(4)

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-003239 | codetables-gf4-36-8

Minimum distance of a [36, 8] linear code over GF(4)

codes are known with d = 20, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-003240 | codetables-gf4-36-9

Minimum distance of a [36, 9] linear code over GF(4)

codes are known with d = 19, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-003241 | codetables-gf4-37-7

Minimum distance of a [37, 7] linear code over GF(4)

codes are known with d = 22, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-003242 | codetables-gf4-37-8

Minimum distance of a [37, 8] linear code over GF(4)

codes are known with d = 21, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-003243 | codetables-gf4-37-9

Minimum distance of a [37, 9] linear code over GF(4)

codes are known with d = 20, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-003244 | codetables-gf4-38-6

Minimum distance of a [38, 6] linear code over GF(4)

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-003245 | codetables-gf4-38-7

Minimum distance of a [38, 7] linear code over GF(4)

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-003246 | codetables-gf4-38-8

Minimum distance of a [38, 8] linear code over GF(4)

codes are known with d = 21, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-003247 | codetables-gf4-38-9

Minimum distance of a [38, 9] linear code over GF(4)

codes are known with d = 20, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-003248 | codetables-gf4-39-6

Minimum distance of a [39, 6] linear code over GF(4)

codes are known with d = 25, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-003249 | codetables-gf4-39-7

Minimum distance of a [39, 7] linear code over GF(4)

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-003250 | codetables-gf4-39-8

Minimum distance of a [39, 8] linear code over GF(4)

codes are known with d = 22, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-003251 | codetables-gf4-39-9

Minimum distance of a [39, 9] linear code over GF(4)

codes are known with d = 21, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-003252 | codetables-gf4-40-6

Minimum distance of a [40, 6] linear code over GF(4)

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-003253 | codetables-gf4-40-7

Minimum distance of a [40, 7] linear code over GF(4)

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-003254 | codetables-gf4-40-8

Minimum distance of a [40, 8] linear code over GF(4)

codes are known with d = 23, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-003255 | codetables-gf4-40-9

Minimum distance of a [40, 9] linear code over GF(4)

codes are known with d = 22, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-003256 | codetables-gf4-41-7

Minimum distance of a [41, 7] linear code over GF(4)

codes are known with d = 25, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-003257 | codetables-gf4-41-8

Minimum distance of a [41, 8] linear code over GF(4)

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-003258 | codetables-gf4-41-9

Minimum distance of a [41, 9] linear code over GF(4)

codes are known with d = 22, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-003259 | codetables-gf4-42-7

Minimum distance of a [42, 7] linear code over GF(4)

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-003260 | codetables-gf4-42-8

Minimum distance of a [42, 8] linear code over GF(4)

codes are known with d = 25, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-003261 | codetables-gf4-42-9

Minimum distance of a [42, 9] linear code over GF(4)

codes are known with d = 23, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-003262 | codetables-gf4-43-6

Minimum distance of a [43, 6] linear code over GF(4)

codes are known with d = 28, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-003263 | codetables-gf4-43-7

Minimum distance of a [43, 7] linear code over GF(4)

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-003264 | codetables-gf4-43-8

Minimum distance of a [43, 8] linear code over GF(4)

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-003265 | codetables-gf4-43-9

Minimum distance of a [43, 9] linear code over GF(4)

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-003266 | codetables-gf4-44-5

Minimum distance of a [44, 5] linear code over GF(4)

codes are known with d = 30, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-003267 | codetables-gf4-44-6

Minimum distance of a [44, 6] linear code over GF(4)

codes are known with d = 29, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-003268 | codetables-gf4-44-7

Minimum distance of a [44, 7] linear code over GF(4)

codes are known with d = 27, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-003269 | codetables-gf4-44-8

Minimum distance of a [44, 8] linear code over GF(4)

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-003270 | codetables-gf4-44-9

Minimum distance of a [44, 9] linear code over GF(4)

codes are known with d = 24, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-003271 | codetables-gf4-45-7

Minimum distance of a [45, 7] linear code over GF(4)

codes are known with d = 28, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-003272 | codetables-gf4-45-8

Minimum distance of a [45, 8] linear code over GF(4)

codes are known with d = 27, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-003273 | codetables-gf4-45-9

Minimum distance of a [45, 9] linear code over GF(4)

codes are known with d = 25, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-003274 | codetables-gf4-46-6

Minimum distance of a [46, 6] linear code over GF(4)

codes are known with d = 30, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-003275 | codetables-gf4-46-7

Minimum distance of a [46, 7] linear code over GF(4)

codes are known with d = 29, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-003276 | codetables-gf4-46-8

Minimum distance of a [46, 8] linear code over GF(4)

codes are known with d = 27, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-003277 | codetables-gf4-46-9

Minimum distance of a [46, 9] linear code over GF(4)

codes are known with d = 26, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-003278 | codetables-gf4-47-6

Minimum distance of a [47, 6] linear code over GF(4)

codes are known with d = 31, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-003279 | codetables-gf4-47-7

Minimum distance of a [47, 7] linear code over GF(4)

codes are known with d = 30, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-003280 | codetables-gf4-47-8

Minimum distance of a [47, 8] linear code over GF(4)

codes are known with d = 28, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-003281 | codetables-gf4-47-9

Minimum distance of a [47, 9] linear code over GF(4)

codes are known with d = 27, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-003282 | codetables-gf4-48-7

Minimum distance of a [48, 7] linear code over GF(4)

codes are known with d = 31, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-003283 | codetables-gf4-48-8

Minimum distance of a [48, 8] linear code over GF(4)

codes are known with d = 29, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-003284 | codetables-gf4-48-9

Minimum distance of a [48, 9] linear code over GF(4)

codes are known with d = 28, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-003285 | codetables-gf4-49-6

Minimum distance of a [49, 6] linear code over GF(4)

codes are known with d = 32, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-003286 | codetables-gf4-49-8

Minimum distance of a [49, 8] linear code over GF(4)

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-003287 | codetables-gf4-49-9

Minimum distance of a [49, 9] linear code over GF(4)

codes are known with d = 29, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-003288 | codetables-gf4-50-6

Minimum distance of a [50, 6] linear code over GF(4)

codes are known with d = 33, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-003289 | codetables-gf4-50-7

Minimum distance of a [50, 7] linear code over GF(4)

codes are known with d = 32, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-003290 | codetables-gf4-50-8

Minimum distance of a [50, 8] linear code over GF(4)

codes are known with d = 31, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-003291 | codetables-gf4-50-9

Minimum distance of a [50, 9] linear code over GF(4)

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-003292 | codetables-gf4-51-6

Minimum distance of a [51, 6] linear code over GF(4)

codes are known with d = 34, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003293 | codetables-gf4-51-7

Minimum distance of a [51, 7] linear code over GF(4)

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-003294 | codetables-gf4-51-8

Minimum distance of a [51, 8] linear code over GF(4)

codes are known with d = 32, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-003295 | codetables-gf4-51-9

Minimum distance of a [51, 9] linear code over GF(4)

codes are known with d = 31, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-003296 | codetables-gf4-52-6

Minimum distance of a [52, 6] linear code over GF(4)

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-003297 | codetables-gf4-52-7

Minimum distance of a [52, 7] linear code over GF(4)

codes are known with d = 33, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003298 | codetables-gf4-52-8

Minimum distance of a [52, 8] linear code over GF(4)

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-003299 | codetables-gf4-52-9

Minimum distance of a [52, 9] linear code over GF(4)

codes are known with d = 32, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-003300 | codetables-gf4-53-7

Minimum distance of a [53, 7] linear code over GF(4)

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-003301 | codetables-gf4-53-8

Minimum distance of a [53, 8] linear code over GF(4)

codes are known with d = 32, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003302 | codetables-gf4-53-9

Minimum distance of a [53, 9] linear code over GF(4)

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-003303 | codetables-gf4-54-6

Minimum distance of a [54, 6] linear code over GF(4)

codes are known with d = 36, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-003304 | codetables-gf4-54-7

Minimum distance of a [54, 7] linear code over GF(4)

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-003305 | codetables-gf4-54-8

Minimum distance of a [54, 8] linear code over GF(4)

codes are known with d = 33, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003306 | codetables-gf4-54-9

Minimum distance of a [54, 9] linear code over GF(4)

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-003307 | codetables-gf4-55-6

Minimum distance of a [55, 6] linear code over GF(4)

codes are known with d = 36, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-003308 | codetables-gf4-55-7

Minimum distance of a [55, 7] linear code over GF(4)

codes are known with d = 36, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-003309 | codetables-gf4-55-8

Minimum distance of a [55, 8] linear code over GF(4)

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-003310 | codetables-gf4-55-9

Minimum distance of a [55, 9] linear code over GF(4)

codes are known with d = 32, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003311 | codetables-gf4-56-6

Minimum distance of a [56, 6] linear code over GF(4)

codes are known with d = 37, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-003312 | codetables-gf4-56-7

Minimum distance of a [56, 7] linear code over GF(4)

codes are known with d = 36, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-003313 | codetables-gf4-56-8

Minimum distance of a [56, 8] linear code over GF(4)

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-003314 | codetables-gf4-56-9

Minimum distance of a [56, 9] linear code over GF(4)

codes are known with d = 33, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-003315 | codetables-gf4-57-6

Minimum distance of a [57, 6] linear code over GF(4)

codes are known with d = 38, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-003316 | codetables-gf4-57-7

Minimum distance of a [57, 7] linear code over GF(4)

codes are known with d = 37, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-003317 | codetables-gf4-57-8

Minimum distance of a [57, 8] linear code over GF(4)

codes are known with d = 36, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-003318 | codetables-gf4-57-9

Minimum distance of a [57, 9] linear code over GF(4)

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-003319 | codetables-gf4-58-5

Minimum distance of a [58, 5] linear code over GF(4)

codes are known with d = 40, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-003320 | codetables-gf4-58-6

Minimum distance of a [58, 6] linear code over GF(4)

codes are known with d = 39, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-003321 | codetables-gf4-58-7

Minimum distance of a [58, 7] linear code over GF(4)

codes are known with d = 38, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-003322 | codetables-gf4-58-8

Minimum distance of a [58, 8] linear code over GF(4)

codes are known with d = 37, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-003323 | codetables-gf4-58-9

Minimum distance of a [58, 9] linear code over GF(4)

codes are known with d = 34, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-003324 | codetables-gf4-59-5

Minimum distance of a [59, 5] linear code over GF(4)

codes are known with d = 41, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-003325 | codetables-gf4-59-7

Minimum distance of a [59, 7] linear code over GF(4)

codes are known with d = 39, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-003326 | codetables-gf4-59-8

Minimum distance of a [59, 8] linear code over GF(4)

codes are known with d = 38, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-003327 | codetables-gf4-59-9

Minimum distance of a [59, 9] linear code over GF(4)

codes are known with d = 35, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-003328 | codetables-gf4-60-5

Minimum distance of a [60, 5] linear code over GF(4)

codes are known with d = 42, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-003329 | codetables-gf4-60-8

Minimum distance of a [60, 8] linear code over GF(4)

codes are known with d = 39, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-003330 | codetables-gf4-60-9

Minimum distance of a [60, 9] linear code over GF(4)

codes are known with d = 36, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-003331 | codetables-gf4-61-5

Minimum distance of a [61, 5] linear code over GF(4)

codes are known with d = 43, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-003332 | codetables-gf4-61-9

Minimum distance of a [61, 9] linear code over GF(4)

codes are known with d = 36, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-003333 | codetables-gf4-62-9

Minimum distance of a [62, 9] linear code over GF(4)

codes are known with d = 37, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-003334 | codetables-gf4-63-5

Minimum distance of a [63, 5] linear code over GF(4)

codes are known with d = 44, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-003335 | codetables-gf4-63-9

Minimum distance of a [63, 9] linear code over GF(4)

codes are known with d = 38, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-003336 | codetables-gf4-64-5

Minimum distance of a [64, 5] linear code over GF(4)

codes are known with d = 45, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-003337 | codetables-gf4-64-9

Minimum distance of a [64, 9] linear code over GF(4)

codes are known with d = 39, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-003338 | codetables-gf4-65-5

Minimum distance of a [65, 5] linear code over GF(4)

codes are known with d = 46, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-003339 | codetables-gf4-65-6

Minimum distance of a [65, 6] linear code over GF(4)

codes are known with d = 44, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-003340 | codetables-gf4-65-9

Minimum distance of a [65, 9] linear code over GF(4)

codes are known with d = 39, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-003341 | codetables-gf4-66-5

Minimum distance of a [66, 5] linear code over GF(4)

codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-003342 | codetables-gf4-66-6

Minimum distance of a [66, 6] linear code over GF(4)

codes are known with d = 45, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-003343 | codetables-gf4-66-7

Minimum distance of a [66, 7] linear code over GF(4)

codes are known with d = 44, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-003344 | codetables-gf4-66-9

Minimum distance of a [66, 9] linear code over GF(4)

codes are known with d = 40, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-003345 | codetables-gf4-67-6

Minimum distance of a [67, 6] linear code over GF(4)

codes are known with d = 46, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-003346 | codetables-gf4-67-7

Minimum distance of a [67, 7] linear code over GF(4)

codes are known with d = 45, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-003347 | codetables-gf4-67-8

Minimum distance of a [67, 8] linear code over GF(4)

codes are known with d = 44, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-003348 | codetables-gf4-67-9

Minimum distance of a [67, 9] linear code over GF(4)

codes are known with d = 41, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-003349 | codetables-gf4-68-6

Minimum distance of a [68, 6] linear code over GF(4)

codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-003350 | codetables-gf4-68-7

Minimum distance of a [68, 7] linear code over GF(4)

codes are known with d = 46, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-003351 | codetables-gf4-68-8

Minimum distance of a [68, 8] linear code over GF(4)

codes are known with d = 44, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-003352 | codetables-gf4-68-9

Minimum distance of a [68, 9] linear code over GF(4)

codes are known with d = 42, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-003353 | codetables-gf4-69-7

Minimum distance of a [69, 7] linear code over GF(4)

codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-003354 | codetables-gf4-69-8

Minimum distance of a [69, 8] linear code over GF(4)

codes are known with d = 45, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-003355 | codetables-gf4-69-9

Minimum distance of a [69, 9] linear code over GF(4)

codes are known with d = 43, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-003356 | codetables-gf4-70-6

Minimum distance of a [70, 6] linear code over GF(4)

codes are known with d = 48, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-003357 | codetables-gf4-70-8

Minimum distance of a [70, 8] linear code over GF(4)

codes are known with d = 46, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-003358 | codetables-gf4-70-9

Minimum distance of a [70, 9] linear code over GF(4)

codes are known with d = 44, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-003359 | codetables-gf4-71-6

Minimum distance of a [71, 6] linear code over GF(4)

codes are known with d = 49, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-003360 | codetables-gf4-71-7

Minimum distance of a [71, 7] linear code over GF(4)

codes are known with d = 48, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-003361 | codetables-gf4-71-8

Minimum distance of a [71, 8] linear code over GF(4)

codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-003362 | codetables-gf4-71-9

Minimum distance of a [71, 9] linear code over GF(4)

codes are known with d = 44, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-003363 | codetables-gf4-72-6

Minimum distance of a [72, 6] linear code over GF(4)

codes are known with d = 50, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-003364 | codetables-gf4-72-7

Minimum distance of a [72, 7] linear code over GF(4)

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-003365 | codetables-gf4-72-8

Minimum distance of a [72, 8] linear code over GF(4)

codes are known with d = 48, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-003366 | codetables-gf4-72-9

Minimum distance of a [72, 9] linear code over GF(4)

codes are known with d = 44, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-003367 | codetables-gf4-73-6

Minimum distance of a [73, 6] linear code over GF(4)

codes are known with d = 51, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-003368 | codetables-gf4-73-7

Minimum distance of a [73, 7] linear code over GF(4)

codes are known with d = 49, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-003369 | codetables-gf4-73-8

Minimum distance of a [73, 8] linear code over GF(4)

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-003370 | codetables-gf4-73-9

Minimum distance of a [73, 9] linear code over GF(4)

codes are known with d = 45, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-003371 | codetables-gf4-74-7

Minimum distance of a [74, 7] linear code over GF(4)

codes are known with d = 50, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-003372 | codetables-gf4-74-8

Minimum distance of a [74, 8] linear code over GF(4)

codes are known with d = 48, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-003373 | codetables-gf4-74-9

Minimum distance of a [74, 9] linear code over GF(4)

codes are known with d = 46, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-003374 | codetables-gf4-75-7

Minimum distance of a [75, 7] linear code over GF(4)

codes are known with d = 51, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-003375 | codetables-gf4-75-8

Minimum distance of a [75, 8] linear code over GF(4)

codes are known with d = 48, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-003376 | codetables-gf4-75-9

Minimum distance of a [75, 9] linear code over GF(4)

codes are known with d = 47, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-003377 | codetables-gf4-76-8

Minimum distance of a [76, 8] linear code over GF(4)

codes are known with d = 49, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-003378 | codetables-gf4-76-9

Minimum distance of a [76, 9] linear code over GF(4)

codes are known with d = 48, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-003379 | codetables-gf4-77-7

Minimum distance of a [77, 7] linear code over GF(4)

codes are known with d = 52, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-003380 | codetables-gf4-77-8

Minimum distance of a [77, 8] linear code over GF(4)

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-003381 | codetables-gf4-77-9

Minimum distance of a [77, 9] linear code over GF(4)

codes are known with d = 48, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-003382 | codetables-gf4-78-7

Minimum distance of a [78, 7] linear code over GF(4)

codes are known with d = 52, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-003383 | codetables-gf4-78-8

Minimum distance of a [78, 8] linear code over GF(4)

codes are known with d = 50, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-003384 | codetables-gf4-78-9

Minimum distance of a [78, 9] linear code over GF(4)

codes are known with d = 48, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-003385 | codetables-gf4-79-5

Minimum distance of a [79, 5] linear code over GF(4)

codes are known with d = 56, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-003386 | codetables-gf4-79-7

Minimum distance of a [79, 7] linear code over GF(4)

codes are known with d = 53, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-003387 | codetables-gf4-79-8

Minimum distance of a [79, 8] linear code over GF(4)

codes are known with d = 51, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-003388 | codetables-gf4-79-9

Minimum distance of a [79, 9] linear code over GF(4)

codes are known with d = 48, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-003389 | codetables-gf4-80-5

Minimum distance of a [80, 5] linear code over GF(4)

codes are known with d = 57, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-003390 | codetables-gf4-80-6

Minimum distance of a [80, 6] linear code over GF(4)

codes are known with d = 56, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-003391 | codetables-gf4-80-7

Minimum distance of a [80, 7] linear code over GF(4)

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-003392 | codetables-gf4-80-8

Minimum distance of a [80, 8] linear code over GF(4)

codes are known with d = 52, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-003393 | codetables-gf4-80-9

Minimum distance of a [80, 9] linear code over GF(4)

codes are known with d = 49, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-003394 | codetables-gf4-81-5

Minimum distance of a [81, 5] linear code over GF(4)

codes are known with d = 58, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-003395 | codetables-gf4-81-6

Minimum distance of a [81, 6] linear code over GF(4)

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-003396 | codetables-gf4-81-7

Minimum distance of a [81, 7] linear code over GF(4)

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-003397 | codetables-gf4-81-8

Minimum distance of a [81, 8] linear code over GF(4)

codes are known with d = 52, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-003398 | codetables-gf4-81-9

Minimum distance of a [81, 9] linear code over GF(4)

codes are known with d = 50, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-003399 | codetables-gf4-82-5

Minimum distance of a [82, 5] linear code over GF(4)

codes are known with d = 59, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-003400 | codetables-gf4-82-6

Minimum distance of a [82, 6] linear code over GF(4)

codes are known with d = 57, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-003401 | codetables-gf4-82-7

Minimum distance of a [82, 7] linear code over GF(4)

codes are known with d = 55, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-003402 | codetables-gf4-82-8

Minimum distance of a [82, 8] linear code over GF(4)

codes are known with d = 53, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-003403 | codetables-gf4-82-9

Minimum distance of a [82, 9] linear code over GF(4)

codes are known with d = 50, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-003404 | codetables-gf4-83-6

Minimum distance of a [83, 6] linear code over GF(4)

codes are known with d = 58, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-003405 | codetables-gf4-83-7

Minimum distance of a [83, 7] linear code over GF(4)

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-003406 | codetables-gf4-83-8

Minimum distance of a [83, 8] linear code over GF(4)

codes are known with d = 54, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-003407 | codetables-gf4-83-9

Minimum distance of a [83, 9] linear code over GF(4)

codes are known with d = 51, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-003408 | codetables-gf4-84-5

Minimum distance of a [84, 5] linear code over GF(4)

codes are known with d = 60, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-003409 | codetables-gf4-84-6

Minimum distance of a [84, 6] linear code over GF(4)

codes are known with d = 59, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-003410 | codetables-gf4-84-7

Minimum distance of a [84, 7] linear code over GF(4)

codes are known with d = 56, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-003411 | codetables-gf4-84-8

Minimum distance of a [84, 8] linear code over GF(4)

codes are known with d = 55, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-003412 | codetables-gf4-84-9

Minimum distance of a [84, 9] linear code over GF(4)

codes are known with d = 52, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-003413 | codetables-gf4-85-5

Minimum distance of a [85, 5] linear code over GF(4)

codes are known with d = 61, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-003414 | codetables-gf4-85-7

Minimum distance of a [85, 7] linear code over GF(4)

codes are known with d = 57, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-003415 | codetables-gf4-85-8

Minimum distance of a [85, 8] linear code over GF(4)

codes are known with d = 56, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-003416 | codetables-gf4-85-9

Minimum distance of a [85, 9] linear code over GF(4)

codes are known with d = 53, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-003417 | codetables-gf4-86-6

Minimum distance of a [86, 6] linear code over GF(4)

codes are known with d = 60, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-003418 | codetables-gf4-86-7

Minimum distance of a [86, 7] linear code over GF(4)

codes are known with d = 58, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-003419 | codetables-gf4-86-8

Minimum distance of a [86, 8] linear code over GF(4)

codes are known with d = 56, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-003420 | codetables-gf4-86-9

Minimum distance of a [86, 9] linear code over GF(4)

codes are known with d = 54, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-003421 | codetables-gf4-87-6

Minimum distance of a [87, 6] linear code over GF(4)

codes are known with d = 61, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-003422 | codetables-gf4-87-7

Minimum distance of a [87, 7] linear code over GF(4)

codes are known with d = 59, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-003423 | codetables-gf4-87-8

Minimum distance of a [87, 8] linear code over GF(4)

codes are known with d = 56, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-003424 | codetables-gf4-87-9

Minimum distance of a [87, 9] linear code over GF(4)

codes are known with d = 55, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-003425 | codetables-gf4-88-6

Minimum distance of a [88, 6] linear code over GF(4)

codes are known with d = 62, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-003426 | codetables-gf4-88-7

Minimum distance of a [88, 7] linear code over GF(4)

codes are known with d = 60, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-003427 | codetables-gf4-88-8

Minimum distance of a [88, 8] linear code over GF(4)

codes are known with d = 57, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-003428 | codetables-gf4-88-9

Minimum distance of a [88, 9] linear code over GF(4)

codes are known with d = 56, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-003429 | codetables-gf4-89-6

Minimum distance of a [89, 6] linear code over GF(4)

codes are known with d = 63, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-003430 | codetables-gf4-89-7

Minimum distance of a [89, 7] linear code over GF(4)

codes are known with d = 60, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-003431 | codetables-gf4-89-8

Minimum distance of a [89, 8] linear code over GF(4)

codes are known with d = 58, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-003432 | codetables-gf4-89-9

Minimum distance of a [89, 9] linear code over GF(4)

codes are known with d = 56, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-003433 | codetables-gf4-90-5

Minimum distance of a [90, 5] linear code over GF(4)

codes are known with d = 64, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-003434 | codetables-gf4-90-7

Minimum distance of a [90, 7] linear code over GF(4)

codes are known with d = 60, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-003435 | codetables-gf4-90-8

Minimum distance of a [90, 8] linear code over GF(4)

codes are known with d = 59, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-003436 | codetables-gf4-90-9

Minimum distance of a [90, 9] linear code over GF(4)

codes are known with d = 56, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-003437 | codetables-gf4-91-5

Minimum distance of a [91, 5] linear code over GF(4)

codes are known with d = 65, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-003438 | codetables-gf4-91-7

Minimum distance of a [91, 7] linear code over GF(4)

codes are known with d = 61, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-003439 | codetables-gf4-91-8

Minimum distance of a [91, 8] linear code over GF(4)

codes are known with d = 60, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-003440 | codetables-gf4-91-9

Minimum distance of a [91, 9] linear code over GF(4)

codes are known with d = 57, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-003441 | codetables-gf4-92-5

Minimum distance of a [92, 5] linear code over GF(4)

codes are known with d = 66, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-003442 | codetables-gf4-92-6

Minimum distance of a [92, 6] linear code over GF(4)

codes are known with d = 64, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-003443 | codetables-gf4-92-7

Minimum distance of a [92, 7] linear code over GF(4)

codes are known with d = 62, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-003444 | codetables-gf4-92-8

Minimum distance of a [92, 8] linear code over GF(4)

codes are known with d = 60, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-003445 | codetables-gf4-92-9

Minimum distance of a [92, 9] linear code over GF(4)

codes are known with d = 58, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-003446 | codetables-gf4-93-5

Minimum distance of a [93, 5] linear code over GF(4)

codes are known with d = 67, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-003447 | codetables-gf4-93-6

Minimum distance of a [93, 6] linear code over GF(4)

codes are known with d = 64, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-003448 | codetables-gf4-93-7

Minimum distance of a [93, 7] linear code over GF(4)

codes are known with d = 63, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-003449 | codetables-gf4-93-8

Minimum distance of a [93, 8] linear code over GF(4)

codes are known with d = 60, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-003450 | codetables-gf4-93-9

Minimum distance of a [93, 9] linear code over GF(4)

codes are known with d = 58, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-003451 | codetables-gf4-94-6

Minimum distance of a [94, 6] linear code over GF(4)

codes are known with d = 65, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-003452 | codetables-gf4-94-7

Minimum distance of a [94, 7] linear code over GF(4)

codes are known with d = 64, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-003453 | codetables-gf4-94-8

Minimum distance of a [94, 8] linear code over GF(4)

codes are known with d = 61, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-003454 | codetables-gf4-94-9

Minimum distance of a [94, 9] linear code over GF(4)

codes are known with d = 59, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-003455 | codetables-gf4-95-5

Minimum distance of a [95, 5] linear code over GF(4)

codes are known with d = 68, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-003456 | codetables-gf4-95-6

Minimum distance of a [95, 6] linear code over GF(4)

codes are known with d = 66, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-003457 | codetables-gf4-95-7

Minimum distance of a [95, 7] linear code over GF(4)

codes are known with d = 64, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-003458 | codetables-gf4-95-8

Minimum distance of a [95, 8] linear code over GF(4)

codes are known with d = 62, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-003459 | codetables-gf4-95-9

Minimum distance of a [95, 9] linear code over GF(4)

codes are known with d = 60, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-003460 | codetables-gf4-96-5

Minimum distance of a [96, 5] linear code over GF(4)

codes are known with d = 69, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-003461 | codetables-gf4-96-6

Minimum distance of a [96, 6] linear code over GF(4)

codes are known with d = 67, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-003462 | codetables-gf4-96-7

Minimum distance of a [96, 7] linear code over GF(4)

codes are known with d = 64, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-003463 | codetables-gf4-96-8

Minimum distance of a [96, 8] linear code over GF(4)

codes are known with d = 63, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-003464 | codetables-gf4-96-9

Minimum distance of a [96, 9] linear code over GF(4)

codes are known with d = 60, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-003465 | codetables-gf4-97-6

Minimum distance of a [97, 6] linear code over GF(4)

codes are known with d = 68, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-003466 | codetables-gf4-97-7

Minimum distance of a [97, 7] linear code over GF(4)

codes are known with d = 65, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-003467 | codetables-gf4-97-8

Minimum distance of a [97, 8] linear code over GF(4)

codes are known with d = 64, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-003468 | codetables-gf4-97-9

Minimum distance of a [97, 9] linear code over GF(4)

codes are known with d = 61, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-003469 | codetables-gf4-98-6

Minimum distance of a [98, 6] linear code over GF(4)

codes are known with d = 68, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-003470 | codetables-gf4-98-7

Minimum distance of a [98, 7] linear code over GF(4)

codes are known with d = 66, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-003471 | codetables-gf4-98-8

Minimum distance of a [98, 8] linear code over GF(4)

codes are known with d = 64, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-003472 | codetables-gf4-98-9

Minimum distance of a [98, 9] linear code over GF(4)

codes are known with d = 62, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-003473 | codetables-gf4-99-6

Minimum distance of a [99, 6] linear code over GF(4)

codes are known with d = 69, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-003474 | codetables-gf4-99-7

Minimum distance of a [99, 7] linear code over GF(4)

codes are known with d = 67, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-003475 | codetables-gf4-99-8

Minimum distance of a [99, 8] linear code over GF(4)

codes are known with d = 65, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-003476 | codetables-gf4-99-9

Minimum distance of a [99, 9] linear code over GF(4)

codes are known with d = 63, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-003477 | codetables-gf5-100-5

Minimum distance of a [100, 5] linear code over GF(5)

codes are known with d = 76, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-003478 | codetables-gf5-100-6

Minimum distance of a [100, 6] linear code over GF(5)

codes are known with d = 75, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-003479 | codetables-gf5-100-7

Minimum distance of a [100, 7] linear code over GF(5)

codes are known with d = 74, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-003480 | codetables-gf5-100-8

Minimum distance of a [100, 8] linear code over GF(5)

codes are known with d = 71, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-003481 | codetables-gf5-101-5

Minimum distance of a [101, 5] linear code over GF(5)

codes are known with d = 77, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-003482 | codetables-gf5-101-6

Minimum distance of a [101, 6] linear code over GF(5)

codes are known with d = 75, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-003483 | codetables-gf5-101-7

Minimum distance of a [101, 7] linear code over GF(5)

codes are known with d = 74, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-003484 | codetables-gf5-101-8

Minimum distance of a [101, 8] linear code over GF(5)

codes are known with d = 72, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-003485 | codetables-gf5-102-5

Minimum distance of a [102, 5] linear code over GF(5)

codes are known with d = 78, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-003486 | codetables-gf5-102-6

Minimum distance of a [102, 6] linear code over GF(5)

codes are known with d = 76, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-003487 | codetables-gf5-102-7

Minimum distance of a [102, 7] linear code over GF(5)

codes are known with d = 75, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-003488 | codetables-gf5-102-8

Minimum distance of a [102, 8] linear code over GF(5)

codes are known with d = 73, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-003489 | codetables-gf5-103-4

Minimum distance of a [103, 4] linear code over GF(5)

codes are known with d = 80, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-003490 | codetables-gf5-103-5

Minimum distance of a [103, 5] linear code over GF(5)

codes are known with d = 79, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-003491 | codetables-gf5-103-6

Minimum distance of a [103, 6] linear code over GF(5)

codes are known with d = 77, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-003492 | codetables-gf5-103-7

Minimum distance of a [103, 7] linear code over GF(5)

codes are known with d = 75, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-003493 | codetables-gf5-103-8

Minimum distance of a [103, 8] linear code over GF(5)

codes are known with d = 74, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-003494 | codetables-gf5-104-4

Minimum distance of a [104, 4] linear code over GF(5)

codes are known with d = 81, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-003495 | codetables-gf5-104-5

Minimum distance of a [104, 5] linear code over GF(5)

codes are known with d = 80, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-003496 | codetables-gf5-104-6

Minimum distance of a [104, 6] linear code over GF(5)

codes are known with d = 78, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-003497 | codetables-gf5-104-7

Minimum distance of a [104, 7] linear code over GF(5)

codes are known with d = 76, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-003498 | codetables-gf5-104-8

Minimum distance of a [104, 8] linear code over GF(5)

codes are known with d = 75, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-003499 | codetables-gf5-105-5

Minimum distance of a [105, 5] linear code over GF(5)

codes are known with d = 80, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-003500 | codetables-gf5-105-6

Minimum distance of a [105, 6] linear code over GF(5)

codes are known with d = 79, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-003501 | codetables-gf5-105-7

Minimum distance of a [105, 7] linear code over GF(5)

codes are known with d = 77, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-003502 | codetables-gf5-105-8

Minimum distance of a [105, 8] linear code over GF(5)

codes are known with d = 75, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-003503 | codetables-gf5-106-5

Minimum distance of a [106, 5] linear code over GF(5)

codes are known with d = 81, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-003504 | codetables-gf5-106-6

Minimum distance of a [106, 6] linear code over GF(5)

codes are known with d = 80, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-003505 | codetables-gf5-106-7

Minimum distance of a [106, 7] linear code over GF(5)

codes are known with d = 78, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-003506 | codetables-gf5-106-8

Minimum distance of a [106, 8] linear code over GF(5)

codes are known with d = 75, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-003507 | codetables-gf5-107-5

Minimum distance of a [107, 5] linear code over GF(5)

codes are known with d = 82, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-003508 | codetables-gf5-107-6

Minimum distance of a [107, 6] linear code over GF(5)

codes are known with d = 80, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-003509 | codetables-gf5-107-7

Minimum distance of a [107, 7] linear code over GF(5)

codes are known with d = 78, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-003510 | codetables-gf5-107-8

Minimum distance of a [107, 8] linear code over GF(5)

codes are known with d = 76, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-003511 | codetables-gf5-108-5

Minimum distance of a [108, 5] linear code over GF(5)

codes are known with d = 83, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-003512 | codetables-gf5-108-6

Minimum distance of a [108, 6] linear code over GF(5)

codes are known with d = 80, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-003513 | codetables-gf5-108-7

Minimum distance of a [108, 7] linear code over GF(5)

codes are known with d = 79, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-003514 | codetables-gf5-108-8

Minimum distance of a [108, 8] linear code over GF(5)

codes are known with d = 77, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-003515 | codetables-gf5-109-5

Minimum distance of a [109, 5] linear code over GF(5)

codes are known with d = 84, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-003516 | codetables-gf5-109-6

Minimum distance of a [109, 6] linear code over GF(5)

codes are known with d = 81, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-003517 | codetables-gf5-109-7

Minimum distance of a [109, 7] linear code over GF(5)

codes are known with d = 80, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-003518 | codetables-gf5-109-8

Minimum distance of a [109, 8] linear code over GF(5)

codes are known with d = 78, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-003519 | codetables-gf5-110-6

Minimum distance of a [110, 6] linear code over GF(5)

codes are known with d = 82, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-003520 | codetables-gf5-110-7

Minimum distance of a [110, 7] linear code over GF(5)

codes are known with d = 80, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-003521 | codetables-gf5-110-8

Minimum distance of a [110, 8] linear code over GF(5)

codes are known with d = 79, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-003522 | codetables-gf5-111-5

Minimum distance of a [111, 5] linear code over GF(5)

codes are known with d = 85, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-003523 | codetables-gf5-111-6

Minimum distance of a [111, 6] linear code over GF(5)

codes are known with d = 82, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-003524 | codetables-gf5-111-7

Minimum distance of a [111, 7] linear code over GF(5)

codes are known with d = 81, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-003525 | codetables-gf5-111-8

Minimum distance of a [111, 8] linear code over GF(5)

codes are known with d = 80, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-003526 | codetables-gf5-112-5

Minimum distance of a [112, 5] linear code over GF(5)

codes are known with d = 85, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-003527 | codetables-gf5-112-6

Minimum distance of a [112, 6] linear code over GF(5)

codes are known with d = 83, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-003528 | codetables-gf5-112-7

Minimum distance of a [112, 7] linear code over GF(5)

codes are known with d = 82, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-003529 | codetables-gf5-112-8

Minimum distance of a [112, 8] linear code over GF(5)

codes are known with d = 81, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-003530 | codetables-gf5-113-5

Minimum distance of a [113, 5] linear code over GF(5)

codes are known with d = 86, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-003531 | codetables-gf5-113-6

Minimum distance of a [113, 6] linear code over GF(5)

codes are known with d = 84, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-003532 | codetables-gf5-113-7

Minimum distance of a [113, 7] linear code over GF(5)

codes are known with d = 83, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-003533 | codetables-gf5-113-8

Minimum distance of a [113, 8] linear code over GF(5)

codes are known with d = 82, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-003534 | codetables-gf5-114-5

Minimum distance of a [114, 5] linear code over GF(5)

codes are known with d = 87, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-003535 | codetables-gf5-114-6

Minimum distance of a [114, 6] linear code over GF(5)

codes are known with d = 85, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-003536 | codetables-gf5-114-7

Minimum distance of a [114, 7] linear code over GF(5)

codes are known with d = 84, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-003537 | codetables-gf5-114-8

Minimum distance of a [114, 8] linear code over GF(5)

codes are known with d = 83, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-003538 | codetables-gf5-115-5

Minimum distance of a [115, 5] linear code over GF(5)

codes are known with d = 88, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-003539 | codetables-gf5-115-6

Minimum distance of a [115, 6] linear code over GF(5)

codes are known with d = 86, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-003540 | codetables-gf5-115-7

Minimum distance of a [115, 7] linear code over GF(5)

codes are known with d = 85, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-003541 | codetables-gf5-115-8

Minimum distance of a [115, 8] linear code over GF(5)

codes are known with d = 84, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-003542 | codetables-gf5-116-5

Minimum distance of a [116, 5] linear code over GF(5)

codes are known with d = 89, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-003543 | codetables-gf5-116-6

Minimum distance of a [116, 6] linear code over GF(5)

codes are known with d = 87, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-003544 | codetables-gf5-116-7

Minimum distance of a [116, 7] linear code over GF(5)

codes are known with d = 86, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-003545 | codetables-gf5-116-8

Minimum distance of a [116, 8] linear code over GF(5)

codes are known with d = 85, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-003546 | codetables-gf5-117-5

Minimum distance of a [117, 5] linear code over GF(5)

codes are known with d = 90, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-003547 | codetables-gf5-117-6

Minimum distance of a [117, 6] linear code over GF(5)

codes are known with d = 88, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-003548 | codetables-gf5-117-7

Minimum distance of a [117, 7] linear code over GF(5)

codes are known with d = 87, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-003549 | codetables-gf5-117-8

Minimum distance of a [117, 8] linear code over GF(5)

codes are known with d = 86, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-003550 | codetables-gf5-118-5

Minimum distance of a [118, 5] linear code over GF(5)

codes are known with d = 90, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-003551 | codetables-gf5-118-6

Minimum distance of a [118, 6] linear code over GF(5)

codes are known with d = 89, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-003552 | codetables-gf5-118-7

Minimum distance of a [118, 7] linear code over GF(5)

codes are known with d = 88, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-003553 | codetables-gf5-118-8

Minimum distance of a [118, 8] linear code over GF(5)

codes are known with d = 87, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-003554 | codetables-gf5-119-5

Minimum distance of a [119, 5] linear code over GF(5)

codes are known with d = 91, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-003555 | codetables-gf5-119-6

Minimum distance of a [119, 6] linear code over GF(5)

codes are known with d = 90, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-003556 | codetables-gf5-119-7

Minimum distance of a [119, 7] linear code over GF(5)

codes are known with d = 89, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-003557 | codetables-gf5-119-8

Minimum distance of a [119, 8] linear code over GF(5)

codes are known with d = 88, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-003558 | codetables-gf5-120-5

Minimum distance of a [120, 5] linear code over GF(5)

codes are known with d = 92, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-003559 | codetables-gf5-120-6

Minimum distance of a [120, 6] linear code over GF(5)

codes are known with d = 91, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-003560 | codetables-gf5-120-7

Minimum distance of a [120, 7] linear code over GF(5)

codes are known with d = 90, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-003561 | codetables-gf5-120-8

Minimum distance of a [120, 8] linear code over GF(5)

codes are known with d = 89, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-003562 | codetables-gf5-121-5

Minimum distance of a [121, 5] linear code over GF(5)

codes are known with d = 93, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-003563 | codetables-gf5-121-6

Minimum distance of a [121, 6] linear code over GF(5)

codes are known with d = 92, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-003564 | codetables-gf5-121-7

Minimum distance of a [121, 7] linear code over GF(5)

codes are known with d = 91, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-003565 | codetables-gf5-121-8

Minimum distance of a [121, 8] linear code over GF(5)

codes are known with d = 90, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-003566 | codetables-gf5-122-5

Minimum distance of a [122, 5] linear code over GF(5)

codes are known with d = 94, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-003567 | codetables-gf5-122-6

Minimum distance of a [122, 6] linear code over GF(5)

codes are known with d = 93, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-003568 | codetables-gf5-122-7

Minimum distance of a [122, 7] linear code over GF(5)

codes are known with d = 92, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-003569 | codetables-gf5-122-8

Minimum distance of a [122, 8] linear code over GF(5)

codes are known with d = 91, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-003570 | codetables-gf5-123-5

Minimum distance of a [123, 5] linear code over GF(5)

codes are known with d = 95, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-003571 | codetables-gf5-123-6

Minimum distance of a [123, 6] linear code over GF(5)

codes are known with d = 94, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-003572 | codetables-gf5-123-7

Minimum distance of a [123, 7] linear code over GF(5)

codes are known with d = 93, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-003573 | codetables-gf5-123-8

Minimum distance of a [123, 8] linear code over GF(5)

codes are known with d = 92, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-003574 | codetables-gf5-124-5

Minimum distance of a [124, 5] linear code over GF(5)

codes are known with d = 96, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-003575 | codetables-gf5-125-5

Minimum distance of a [125, 5] linear code over GF(5)

codes are known with d = 97, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-003576 | codetables-gf5-125-6

Minimum distance of a [125, 6] linear code over GF(5)

codes are known with d = 95, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-003577 | codetables-gf5-126-5

Minimum distance of a [126, 5] linear code over GF(5)

codes are known with d = 98, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-003578 | codetables-gf5-126-6

Minimum distance of a [126, 6] linear code over GF(5)

codes are known with d = 95, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-003579 | codetables-gf5-126-7

Minimum distance of a [126, 7] linear code over GF(5)

codes are known with d = 95, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-003580 | codetables-gf5-127-5

Minimum distance of a [127, 5] linear code over GF(5)

codes are known with d = 99, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-003581 | codetables-gf5-127-6

Minimum distance of a [127, 6] linear code over GF(5)

codes are known with d = 96, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-003582 | codetables-gf5-127-7

Minimum distance of a [127, 7] linear code over GF(5)

codes are known with d = 95, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-003583 | codetables-gf5-127-8

Minimum distance of a [127, 8] linear code over GF(5)

codes are known with d = 95, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-003584 | codetables-gf5-128-6

Minimum distance of a [128, 6] linear code over GF(5)

codes are known with d = 97, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-003585 | codetables-gf5-128-7

Minimum distance of a [128, 7] linear code over GF(5)

codes are known with d = 96, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-003586 | codetables-gf5-128-8

Minimum distance of a [128, 8] linear code over GF(5)

codes are known with d = 95, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-003587 | codetables-gf5-129-6

Minimum distance of a [129, 6] linear code over GF(5)

codes are known with d = 98, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-003588 | codetables-gf5-129-7

Minimum distance of a [129, 7] linear code over GF(5)

codes are known with d = 97, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-003589 | codetables-gf5-129-8

Minimum distance of a [129, 8] linear code over GF(5)

codes are known with d = 95, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-003590 | codetables-gf5-130-5

Minimum distance of a [130, 5] linear code over GF(5)

codes are known with d = 100, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-003591 | codetables-gf5-130-6

Minimum distance of a [130, 6] linear code over GF(5)

codes are known with d = 99, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-003592 | codetables-gf5-130-7

Minimum distance of a [130, 7] linear code over GF(5)

codes are known with d = 98, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-003593 | codetables-gf5-130-8

Minimum distance of a [130, 8] linear code over GF(5)

codes are known with d = 96, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-003594 | codetables-gf5-16-5

Minimum distance of a [16, 5] linear code over GF(5)

codes are known with d = 9, and no code can have d greater than 10.

来源状态 open | 2026-07-23 | 来源

V3P-003595 | codetables-gf5-16-6

Minimum distance of a [16, 6] linear code over GF(5)

codes are known with d = 8, and no code can have d greater than 9.

来源状态 open | 2026-07-23 | 来源

V3P-003596 | codetables-gf5-17-6

Minimum distance of a [17, 6] linear code over GF(5)

codes are known with d = 9, and no code can have d greater than 10.

来源状态 open | 2026-07-23 | 来源

V3P-003597 | codetables-gf5-17-7

Minimum distance of a [17, 7] linear code over GF(5)

codes are known with d = 8, and no code can have d greater than 9.

来源状态 open | 2026-07-23 | 来源

V3P-003598 | codetables-gf5-17-8

Minimum distance of a [17, 8] linear code over GF(5)

codes are known with d = 7, and no code can have d greater than 8.

来源状态 open | 2026-07-23 | 来源

V3P-003599 | codetables-gf5-18-7

Minimum distance of a [18, 7] linear code over GF(5)

codes are known with d = 9, and no code can have d greater than 10.

来源状态 open | 2026-07-23 | 来源

V3P-003600 | codetables-gf5-18-8

Minimum distance of a [18, 8] linear code over GF(5)

codes are known with d = 8, and no code can have d greater than 9.

来源状态 open | 2026-07-23 | 来源

V3P-003601 | codetables-gf5-19-8

Minimum distance of a [19, 8] linear code over GF(5)

codes are known with d = 9, and no code can have d greater than 10.

来源状态 open | 2026-07-23 | 来源

V3P-003602 | codetables-gf5-20-7

Minimum distance of a [20, 7] linear code over GF(5)

codes are known with d = 10, and no code can have d greater than 11.

来源状态 open | 2026-07-23 | 来源

V3P-003603 | codetables-gf5-20-8

Minimum distance of a [20, 8] linear code over GF(5)

codes are known with d = 9, and no code can have d greater than 10.

来源状态 open | 2026-07-23 | 来源

V3P-003604 | codetables-gf5-21-5

Minimum distance of a [21, 5] linear code over GF(5)

codes are known with d = 13, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-003605 | codetables-gf5-21-6

Minimum distance of a [21, 6] linear code over GF(5)

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-003606 | codetables-gf5-21-7

Minimum distance of a [21, 7] linear code over GF(5)

codes are known with d = 11, and no code can have d greater than 12.

来源状态 open | 2026-07-23 | 来源

V3P-003607 | codetables-gf5-21-8

Minimum distance of a [21, 8] linear code over GF(5)

codes are known with d = 10, and no code can have d greater than 11.

来源状态 open | 2026-07-23 | 来源

V3P-003608 | codetables-gf5-22-5

Minimum distance of a [22, 5] linear code over GF(5)

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-003609 | codetables-gf5-22-6

Minimum distance of a [22, 6] linear code over GF(5)

codes are known with d = 13, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-003610 | codetables-gf5-22-7

Minimum distance of a [22, 7] linear code over GF(5)

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-003611 | codetables-gf5-22-8

Minimum distance of a [22, 8] linear code over GF(5)

codes are known with d = 11, and no code can have d greater than 12.

来源状态 open | 2026-07-23 | 来源

V3P-003612 | codetables-gf5-23-6

Minimum distance of a [23, 6] linear code over GF(5)

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-003613 | codetables-gf5-23-7

Minimum distance of a [23, 7] linear code over GF(5)

codes are known with d = 13, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-003614 | codetables-gf5-23-8

Minimum distance of a [23, 8] linear code over GF(5)

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-003615 | codetables-gf5-24-7

Minimum distance of a [24, 7] linear code over GF(5)

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-003616 | codetables-gf5-24-8

Minimum distance of a [24, 8] linear code over GF(5)

codes are known with d = 13, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-003617 | codetables-gf5-25-8

Minimum distance of a [25, 8] linear code over GF(5)

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-003618 | codetables-gf5-26-5

Minimum distance of a [26, 5] linear code over GF(5)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-003619 | codetables-gf5-26-6

Minimum distance of a [26, 6] linear code over GF(5)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-003620 | codetables-gf5-26-8

Minimum distance of a [26, 8] linear code over GF(5)

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-003621 | codetables-gf5-27-5

Minimum distance of a [27, 5] linear code over GF(5)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-003622 | codetables-gf5-27-6

Minimum distance of a [27, 6] linear code over GF(5)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-003623 | codetables-gf5-27-7

Minimum distance of a [27, 7] linear code over GF(5)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-003624 | codetables-gf5-27-8

Minimum distance of a [27, 8] linear code over GF(5)

codes are known with d = 15, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-003625 | codetables-gf5-28-5

Minimum distance of a [28, 5] linear code over GF(5)

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-003626 | codetables-gf5-28-6

Minimum distance of a [28, 6] linear code over GF(5)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-003627 | codetables-gf5-28-7

Minimum distance of a [28, 7] linear code over GF(5)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-003628 | codetables-gf5-28-8

Minimum distance of a [28, 8] linear code over GF(5)

codes are known with d = 15, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-003629 | codetables-gf5-29-6

Minimum distance of a [29, 6] linear code over GF(5)

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-003630 | codetables-gf5-29-7

Minimum distance of a [29, 7] linear code over GF(5)

codes are known with d = 17, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-003631 | codetables-gf5-29-8

Minimum distance of a [29, 8] linear code over GF(5)

codes are known with d = 16, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-003632 | codetables-gf5-30-5

Minimum distance of a [30, 5] linear code over GF(5)

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-003633 | codetables-gf5-30-7

Minimum distance of a [30, 7] linear code over GF(5)

codes are known with d = 18, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-003634 | codetables-gf5-30-8

Minimum distance of a [30, 8] linear code over GF(5)

codes are known with d = 16, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-003635 | codetables-gf5-31-5

Minimum distance of a [31, 5] linear code over GF(5)

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-003636 | codetables-gf5-31-6

Minimum distance of a [31, 6] linear code over GF(5)

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-003637 | codetables-gf5-31-7

Minimum distance of a [31, 7] linear code over GF(5)

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-003638 | codetables-gf5-31-8

Minimum distance of a [31, 8] linear code over GF(5)

codes are known with d = 17, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-003639 | codetables-gf5-32-5

Minimum distance of a [32, 5] linear code over GF(5)

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-003640 | codetables-gf5-32-6

Minimum distance of a [32, 6] linear code over GF(5)

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-003641 | codetables-gf5-32-7

Minimum distance of a [32, 7] linear code over GF(5)

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-003642 | codetables-gf5-32-8

Minimum distance of a [32, 8] linear code over GF(5)

codes are known with d = 18, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-003643 | codetables-gf5-33-5

Minimum distance of a [33, 5] linear code over GF(5)

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-003644 | codetables-gf5-33-6

Minimum distance of a [33, 6] linear code over GF(5)

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-003645 | codetables-gf5-33-7

Minimum distance of a [33, 7] linear code over GF(5)

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-003646 | codetables-gf5-33-8

Minimum distance of a [33, 8] linear code over GF(5)

codes are known with d = 18, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-003647 | codetables-gf5-34-6

Minimum distance of a [34, 6] linear code over GF(5)

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-003648 | codetables-gf5-34-7

Minimum distance of a [34, 7] linear code over GF(5)

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-003649 | codetables-gf5-34-8

Minimum distance of a [34, 8] linear code over GF(5)

codes are known with d = 19, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-003650 | codetables-gf5-35-5

Minimum distance of a [35, 5] linear code over GF(5)

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-003651 | codetables-gf5-35-7

Minimum distance of a [35, 7] linear code over GF(5)

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-003652 | codetables-gf5-35-8

Minimum distance of a [35, 8] linear code over GF(5)

codes are known with d = 20, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-003653 | codetables-gf5-36-5

Minimum distance of a [36, 5] linear code over GF(5)

codes are known with d = 25, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-003654 | codetables-gf5-36-6

Minimum distance of a [36, 6] linear code over GF(5)

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-003655 | codetables-gf5-36-8

Minimum distance of a [36, 8] linear code over GF(5)

codes are known with d = 21, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-003656 | codetables-gf5-37-5

Minimum distance of a [37, 5] linear code over GF(5)

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-003657 | codetables-gf5-37-6

Minimum distance of a [37, 6] linear code over GF(5)

codes are known with d = 25, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-003658 | codetables-gf5-37-7

Minimum distance of a [37, 7] linear code over GF(5)

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-003659 | codetables-gf5-37-8

Minimum distance of a [37, 8] linear code over GF(5)

codes are known with d = 22, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-003660 | codetables-gf5-38-5

Minimum distance of a [38, 5] linear code over GF(5)

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-003661 | codetables-gf5-38-6

Minimum distance of a [38, 6] linear code over GF(5)

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-003662 | codetables-gf5-38-7

Minimum distance of a [38, 7] linear code over GF(5)

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-003663 | codetables-gf5-38-8

Minimum distance of a [38, 8] linear code over GF(5)

codes are known with d = 22, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-003664 | codetables-gf5-39-5

Minimum distance of a [39, 5] linear code over GF(5)

codes are known with d = 28, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-003665 | codetables-gf5-39-6

Minimum distance of a [39, 6] linear code over GF(5)

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-003666 | codetables-gf5-39-7

Minimum distance of a [39, 7] linear code over GF(5)

codes are known with d = 25, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-003667 | codetables-gf5-39-8

Minimum distance of a [39, 8] linear code over GF(5)

codes are known with d = 23, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-003668 | codetables-gf5-40-5

Minimum distance of a [40, 5] linear code over GF(5)

codes are known with d = 29, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-003669 | codetables-gf5-40-6

Minimum distance of a [40, 6] linear code over GF(5)

codes are known with d = 28, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-003670 | codetables-gf5-40-7

Minimum distance of a [40, 7] linear code over GF(5)

codes are known with d = 25, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-003671 | codetables-gf5-40-8

Minimum distance of a [40, 8] linear code over GF(5)

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-003672 | codetables-gf5-41-4

Minimum distance of a [41, 4] linear code over GF(5)

codes are known with d = 30, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-003673 | codetables-gf5-41-7

Minimum distance of a [41, 7] linear code over GF(5)

codes are known with d = 26, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-003674 | codetables-gf5-41-8

Minimum distance of a [41, 8] linear code over GF(5)

codes are known with d = 24, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-003675 | codetables-gf5-42-4

Minimum distance of a [42, 4] linear code over GF(5)

codes are known with d = 31, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-003676 | codetables-gf5-42-5

Minimum distance of a [42, 5] linear code over GF(5)

codes are known with d = 30, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-003677 | codetables-gf5-42-7

Minimum distance of a [42, 7] linear code over GF(5)

codes are known with d = 27, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-003678 | codetables-gf5-42-8

Minimum distance of a [42, 8] linear code over GF(5)

codes are known with d = 25, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-003679 | codetables-gf5-43-5

Minimum distance of a [43, 5] linear code over GF(5)

codes are known with d = 31, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-003680 | codetables-gf5-43-7

Minimum distance of a [43, 7] linear code over GF(5)

codes are known with d = 28, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-003681 | codetables-gf5-43-8

Minimum distance of a [43, 8] linear code over GF(5)

codes are known with d = 26, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-003682 | codetables-gf5-44-5

Minimum distance of a [44, 5] linear code over GF(5)

codes are known with d = 31, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-003683 | codetables-gf5-44-6

Minimum distance of a [44, 6] linear code over GF(5)

codes are known with d = 30, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-003684 | codetables-gf5-44-7

Minimum distance of a [44, 7] linear code over GF(5)

codes are known with d = 28, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-003685 | codetables-gf5-44-8

Minimum distance of a [44, 8] linear code over GF(5)

codes are known with d = 27, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-003686 | codetables-gf5-45-5

Minimum distance of a [45, 5] linear code over GF(5)

codes are known with d = 32, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-003687 | codetables-gf5-45-6

Minimum distance of a [45, 6] linear code over GF(5)

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-003688 | codetables-gf5-45-7

Minimum distance of a [45, 7] linear code over GF(5)

codes are known with d = 29, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-003689 | codetables-gf5-45-8

Minimum distance of a [45, 8] linear code over GF(5)

codes are known with d = 27, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-003690 | codetables-gf5-46-5

Minimum distance of a [46, 5] linear code over GF(5)

codes are known with d = 33, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-003691 | codetables-gf5-46-6

Minimum distance of a [46, 6] linear code over GF(5)

codes are known with d = 31, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-003692 | codetables-gf5-46-7

Minimum distance of a [46, 7] linear code over GF(5)

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-003693 | codetables-gf5-46-8

Minimum distance of a [46, 8] linear code over GF(5)

codes are known with d = 28, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-003694 | codetables-gf5-47-4

Minimum distance of a [47, 4] linear code over GF(5)

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-003695 | codetables-gf5-47-5

Minimum distance of a [47, 5] linear code over GF(5)

codes are known with d = 34, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003696 | codetables-gf5-47-6

Minimum distance of a [47, 6] linear code over GF(5)

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-003697 | codetables-gf5-47-7

Minimum distance of a [47, 7] linear code over GF(5)

codes are known with d = 30, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-003698 | codetables-gf5-47-8

Minimum distance of a [47, 8] linear code over GF(5)

codes are known with d = 29, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-003699 | codetables-gf5-48-4

Minimum distance of a [48, 4] linear code over GF(5)

codes are known with d = 36, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-003700 | codetables-gf5-48-5

Minimum distance of a [48, 5] linear code over GF(5)

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-003701 | codetables-gf5-48-6

Minimum distance of a [48, 6] linear code over GF(5)

codes are known with d = 33, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003702 | codetables-gf5-48-7

Minimum distance of a [48, 7] linear code over GF(5)

codes are known with d = 31, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-003703 | codetables-gf5-48-8

Minimum distance of a [48, 8] linear code over GF(5)

codes are known with d = 30, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-003704 | codetables-gf5-49-5

Minimum distance of a [49, 5] linear code over GF(5)

codes are known with d = 36, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-003705 | codetables-gf5-49-6

Minimum distance of a [49, 6] linear code over GF(5)

codes are known with d = 34, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003706 | codetables-gf5-49-7

Minimum distance of a [49, 7] linear code over GF(5)

codes are known with d = 32, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003707 | codetables-gf5-49-8

Minimum distance of a [49, 8] linear code over GF(5)

codes are known with d = 31, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-003708 | codetables-gf5-50-6

Minimum distance of a [50, 6] linear code over GF(5)

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-003709 | codetables-gf5-50-7

Minimum distance of a [50, 7] linear code over GF(5)

codes are known with d = 33, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003710 | codetables-gf5-50-8

Minimum distance of a [50, 8] linear code over GF(5)

codes are known with d = 31, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003711 | codetables-gf5-51-6

Minimum distance of a [51, 6] linear code over GF(5)

codes are known with d = 35, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-003712 | codetables-gf5-51-7

Minimum distance of a [51, 7] linear code over GF(5)

codes are known with d = 33, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-003713 | codetables-gf5-51-8

Minimum distance of a [51, 8] linear code over GF(5)

codes are known with d = 32, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003714 | codetables-gf5-52-6

Minimum distance of a [52, 6] linear code over GF(5)

codes are known with d = 36, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-003715 | codetables-gf5-52-7

Minimum distance of a [52, 7] linear code over GF(5)

codes are known with d = 34, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-003716 | codetables-gf5-52-8

Minimum distance of a [52, 8] linear code over GF(5)

codes are known with d = 33, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-003717 | codetables-gf5-53-6

Minimum distance of a [53, 6] linear code over GF(5)

codes are known with d = 37, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-003718 | codetables-gf5-53-7

Minimum distance of a [53, 7] linear code over GF(5)

codes are known with d = 35, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-003719 | codetables-gf5-53-8

Minimum distance of a [53, 8] linear code over GF(5)

codes are known with d = 33, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-003720 | codetables-gf5-54-6

Minimum distance of a [54, 6] linear code over GF(5)

codes are known with d = 38, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-003721 | codetables-gf5-54-7

Minimum distance of a [54, 7] linear code over GF(5)

codes are known with d = 36, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-003722 | codetables-gf5-54-8

Minimum distance of a [54, 8] linear code over GF(5)

codes are known with d = 34, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-003723 | codetables-gf5-55-5

Minimum distance of a [55, 5] linear code over GF(5)

codes are known with d = 40, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-003724 | codetables-gf5-55-6

Minimum distance of a [55, 6] linear code over GF(5)

codes are known with d = 39, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-003725 | codetables-gf5-55-7

Minimum distance of a [55, 7] linear code over GF(5)

codes are known with d = 36, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-003726 | codetables-gf5-55-8

Minimum distance of a [55, 8] linear code over GF(5)

codes are known with d = 35, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-003727 | codetables-gf5-56-5

Minimum distance of a [56, 5] linear code over GF(5)

codes are known with d = 41, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-003728 | codetables-gf5-56-6

Minimum distance of a [56, 6] linear code over GF(5)

codes are known with d = 40, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-003729 | codetables-gf5-56-7

Minimum distance of a [56, 7] linear code over GF(5)

codes are known with d = 37, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-003730 | codetables-gf5-56-8

Minimum distance of a [56, 8] linear code over GF(5)

codes are known with d = 36, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-003731 | codetables-gf5-57-5

Minimum distance of a [57, 5] linear code over GF(5)

codes are known with d = 42, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-003732 | codetables-gf5-57-6

Minimum distance of a [57, 6] linear code over GF(5)

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-003733 | codetables-gf5-57-7

Minimum distance of a [57, 7] linear code over GF(5)

codes are known with d = 38, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-003734 | codetables-gf5-57-8

Minimum distance of a [57, 8] linear code over GF(5)

codes are known with d = 36, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-003735 | codetables-gf5-58-5

Minimum distance of a [58, 5] linear code over GF(5)

codes are known with d = 43, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-003736 | codetables-gf5-58-6

Minimum distance of a [58, 6] linear code over GF(5)

codes are known with d = 41, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-003737 | codetables-gf5-58-7

Minimum distance of a [58, 7] linear code over GF(5)

codes are known with d = 39, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-003738 | codetables-gf5-58-8

Minimum distance of a [58, 8] linear code over GF(5)

codes are known with d = 37, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-003739 | codetables-gf5-59-5

Minimum distance of a [59, 5] linear code over GF(5)

codes are known with d = 44, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-003740 | codetables-gf5-59-6

Minimum distance of a [59, 6] linear code over GF(5)

codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-003741 | codetables-gf5-59-7

Minimum distance of a [59, 7] linear code over GF(5)

codes are known with d = 40, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-003742 | codetables-gf5-59-8

Minimum distance of a [59, 8] linear code over GF(5)

codes are known with d = 38, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-003743 | codetables-gf5-60-6

Minimum distance of a [60, 6] linear code over GF(5)

codes are known with d = 43, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-003744 | codetables-gf5-60-7

Minimum distance of a [60, 7] linear code over GF(5)

codes are known with d = 40, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-003745 | codetables-gf5-60-8

Minimum distance of a [60, 8] linear code over GF(5)

codes are known with d = 39, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-003746 | codetables-gf5-61-5

Minimum distance of a [61, 5] linear code over GF(5)

codes are known with d = 45, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-003747 | codetables-gf5-61-6

Minimum distance of a [61, 6] linear code over GF(5)

codes are known with d = 44, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-003748 | codetables-gf5-61-7

Minimum distance of a [61, 7] linear code over GF(5)

codes are known with d = 41, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-003749 | codetables-gf5-61-8

Minimum distance of a [61, 8] linear code over GF(5)

codes are known with d = 39, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-003750 | codetables-gf5-62-5

Minimum distance of a [62, 5] linear code over GF(5)

codes are known with d = 46, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-003751 | codetables-gf5-62-6

Minimum distance of a [62, 6] linear code over GF(5)

codes are known with d = 45, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-003752 | codetables-gf5-62-7

Minimum distance of a [62, 7] linear code over GF(5)

codes are known with d = 42, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-003753 | codetables-gf5-62-8

Minimum distance of a [62, 8] linear code over GF(5)

codes are known with d = 40, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-003754 | codetables-gf5-63-5

Minimum distance of a [63, 5] linear code over GF(5)

codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-003755 | codetables-gf5-63-6

Minimum distance of a [63, 6] linear code over GF(5)

codes are known with d = 45, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-003756 | codetables-gf5-63-7

Minimum distance of a [63, 7] linear code over GF(5)

codes are known with d = 43, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-003757 | codetables-gf5-63-8

Minimum distance of a [63, 8] linear code over GF(5)

codes are known with d = 41, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-003758 | codetables-gf5-64-5

Minimum distance of a [64, 5] linear code over GF(5)

codes are known with d = 48, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-003759 | codetables-gf5-64-6

Minimum distance of a [64, 6] linear code over GF(5)

codes are known with d = 45, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-003760 | codetables-gf5-64-7

Minimum distance of a [64, 7] linear code over GF(5)

codes are known with d = 44, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-003761 | codetables-gf5-64-8

Minimum distance of a [64, 8] linear code over GF(5)

codes are known with d = 42, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-003762 | codetables-gf5-65-5

Minimum distance of a [65, 5] linear code over GF(5)

codes are known with d = 49, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-003763 | codetables-gf5-65-6

Minimum distance of a [65, 6] linear code over GF(5)

codes are known with d = 46, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-003764 | codetables-gf5-65-7

Minimum distance of a [65, 7] linear code over GF(5)

codes are known with d = 45, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-003765 | codetables-gf5-65-8

Minimum distance of a [65, 8] linear code over GF(5)

codes are known with d = 43, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-003766 | codetables-gf5-66-6

Minimum distance of a [66, 6] linear code over GF(5)

codes are known with d = 47, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-003767 | codetables-gf5-66-7

Minimum distance of a [66, 7] linear code over GF(5)

codes are known with d = 45, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-003768 | codetables-gf5-66-8

Minimum distance of a [66, 8] linear code over GF(5)

codes are known with d = 44, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-003769 | codetables-gf5-67-5

Minimum distance of a [67, 5] linear code over GF(5)

codes are known with d = 50, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-003770 | codetables-gf5-67-6

Minimum distance of a [67, 6] linear code over GF(5)

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-003771 | codetables-gf5-67-7

Minimum distance of a [67, 7] linear code over GF(5)

codes are known with d = 46, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-003772 | codetables-gf5-67-8

Minimum distance of a [67, 8] linear code over GF(5)

codes are known with d = 45, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-003773 | codetables-gf5-68-5

Minimum distance of a [68, 5] linear code over GF(5)

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-003774 | codetables-gf5-68-6

Minimum distance of a [68, 6] linear code over GF(5)

codes are known with d = 49, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-003775 | codetables-gf5-68-7

Minimum distance of a [68, 7] linear code over GF(5)

codes are known with d = 47, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-003776 | codetables-gf5-68-8

Minimum distance of a [68, 8] linear code over GF(5)

codes are known with d = 46, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-003777 | codetables-gf5-69-5

Minimum distance of a [69, 5] linear code over GF(5)

codes are known with d = 51, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-003778 | codetables-gf5-69-6

Minimum distance of a [69, 6] linear code over GF(5)

codes are known with d = 49, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-003779 | codetables-gf5-69-7

Minimum distance of a [69, 7] linear code over GF(5)

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-003780 | codetables-gf5-69-8

Minimum distance of a [69, 8] linear code over GF(5)

codes are known with d = 47, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-003781 | codetables-gf5-70-5

Minimum distance of a [70, 5] linear code over GF(5)

codes are known with d = 52, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-003782 | codetables-gf5-70-6

Minimum distance of a [70, 6] linear code over GF(5)

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-003783 | codetables-gf5-70-7

Minimum distance of a [70, 7] linear code over GF(5)

codes are known with d = 49, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-003784 | codetables-gf5-70-8

Minimum distance of a [70, 8] linear code over GF(5)

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-003785 | codetables-gf5-71-5

Minimum distance of a [71, 5] linear code over GF(5)

codes are known with d = 53, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-003786 | codetables-gf5-71-6

Minimum distance of a [71, 6] linear code over GF(5)

codes are known with d = 51, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-003787 | codetables-gf5-71-7

Minimum distance of a [71, 7] linear code over GF(5)

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-003788 | codetables-gf5-71-8

Minimum distance of a [71, 8] linear code over GF(5)

codes are known with d = 49, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-003789 | codetables-gf5-72-5

Minimum distance of a [72, 5] linear code over GF(5)

codes are known with d = 54, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-003790 | codetables-gf5-72-6

Minimum distance of a [72, 6] linear code over GF(5)

codes are known with d = 51, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-003791 | codetables-gf5-72-7

Minimum distance of a [72, 7] linear code over GF(5)

codes are known with d = 50, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-003792 | codetables-gf5-72-8

Minimum distance of a [72, 8] linear code over GF(5)

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-003793 | codetables-gf5-73-6

Minimum distance of a [73, 6] linear code over GF(5)

codes are known with d = 52, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-003794 | codetables-gf5-73-7

Minimum distance of a [73, 7] linear code over GF(5)

codes are known with d = 50, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-003795 | codetables-gf5-73-8

Minimum distance of a [73, 8] linear code over GF(5)

codes are known with d = 50, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-003796 | codetables-gf5-74-6

Minimum distance of a [74, 6] linear code over GF(5)

codes are known with d = 53, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-003797 | codetables-gf5-74-7

Minimum distance of a [74, 7] linear code over GF(5)

codes are known with d = 51, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-003798 | codetables-gf5-74-8

Minimum distance of a [74, 8] linear code over GF(5)

codes are known with d = 50, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-003799 | codetables-gf5-75-6

Minimum distance of a [75, 6] linear code over GF(5)

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-003800 | codetables-gf5-75-7

Minimum distance of a [75, 7] linear code over GF(5)

codes are known with d = 52, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-003801 | codetables-gf5-75-8

Minimum distance of a [75, 8] linear code over GF(5)

codes are known with d = 50, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-003802 | codetables-gf5-76-6

Minimum distance of a [76, 6] linear code over GF(5)

codes are known with d = 55, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-003803 | codetables-gf5-76-7

Minimum distance of a [76, 7] linear code over GF(5)

codes are known with d = 52, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-003804 | codetables-gf5-76-8

Minimum distance of a [76, 8] linear code over GF(5)

codes are known with d = 51, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-003805 | codetables-gf5-77-6

Minimum distance of a [77, 6] linear code over GF(5)

codes are known with d = 55, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-003806 | codetables-gf5-77-7

Minimum distance of a [77, 7] linear code over GF(5)

codes are known with d = 53, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-003807 | codetables-gf5-77-8

Minimum distance of a [77, 8] linear code over GF(5)

codes are known with d = 51, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-003808 | codetables-gf5-78-6

Minimum distance of a [78, 6] linear code over GF(5)

codes are known with d = 56, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-003809 | codetables-gf5-78-7

Minimum distance of a [78, 7] linear code over GF(5)

codes are known with d = 54, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-003810 | codetables-gf5-78-8

Minimum distance of a [78, 8] linear code over GF(5)

codes are known with d = 52, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-003811 | codetables-gf5-79-6

Minimum distance of a [79, 6] linear code over GF(5)

codes are known with d = 57, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-003812 | codetables-gf5-79-7

Minimum distance of a [79, 7] linear code over GF(5)

codes are known with d = 55, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-003813 | codetables-gf5-79-8

Minimum distance of a [79, 8] linear code over GF(5)

codes are known with d = 53, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-003814 | codetables-gf5-80-5

Minimum distance of a [80, 5] linear code over GF(5)

codes are known with d = 60, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-003815 | codetables-gf5-80-6

Minimum distance of a [80, 6] linear code over GF(5)

codes are known with d = 58, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-003816 | codetables-gf5-80-7

Minimum distance of a [80, 7] linear code over GF(5)

codes are known with d = 56, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-003817 | codetables-gf5-80-8

Minimum distance of a [80, 8] linear code over GF(5)

codes are known with d = 54, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-003818 | codetables-gf5-81-5

Minimum distance of a [81, 5] linear code over GF(5)

codes are known with d = 61, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-003819 | codetables-gf5-81-6

Minimum distance of a [81, 6] linear code over GF(5)

codes are known with d = 59, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-003820 | codetables-gf5-81-7

Minimum distance of a [81, 7] linear code over GF(5)

codes are known with d = 57, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-003821 | codetables-gf5-81-8

Minimum distance of a [81, 8] linear code over GF(5)

codes are known with d = 54, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-003822 | codetables-gf5-82-5

Minimum distance of a [82, 5] linear code over GF(5)

codes are known with d = 62, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-003823 | codetables-gf5-82-6

Minimum distance of a [82, 6] linear code over GF(5)

codes are known with d = 60, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-003824 | codetables-gf5-82-7

Minimum distance of a [82, 7] linear code over GF(5)

codes are known with d = 57, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-003825 | codetables-gf5-82-8

Minimum distance of a [82, 8] linear code over GF(5)

codes are known with d = 55, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-003826 | codetables-gf5-83-5

Minimum distance of a [83, 5] linear code over GF(5)

codes are known with d = 63, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-003827 | codetables-gf5-83-6

Minimum distance of a [83, 6] linear code over GF(5)

codes are known with d = 60, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-003828 | codetables-gf5-83-7

Minimum distance of a [83, 7] linear code over GF(5)

codes are known with d = 58, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-003829 | codetables-gf5-83-8

Minimum distance of a [83, 8] linear code over GF(5)

codes are known with d = 56, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-003830 | codetables-gf5-84-5

Minimum distance of a [84, 5] linear code over GF(5)

codes are known with d = 64, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-003831 | codetables-gf5-84-6

Minimum distance of a [84, 6] linear code over GF(5)

codes are known with d = 61, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-003832 | codetables-gf5-84-7

Minimum distance of a [84, 7] linear code over GF(5)

codes are known with d = 59, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-003833 | codetables-gf5-84-8

Minimum distance of a [84, 8] linear code over GF(5)

codes are known with d = 57, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-003834 | codetables-gf5-85-6

Minimum distance of a [85, 6] linear code over GF(5)

codes are known with d = 62, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-003835 | codetables-gf5-85-7

Minimum distance of a [85, 7] linear code over GF(5)

codes are known with d = 60, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-003836 | codetables-gf5-85-8

Minimum distance of a [85, 8] linear code over GF(5)

codes are known with d = 58, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-003837 | codetables-gf5-86-5

Minimum distance of a [86, 5] linear code over GF(5)

codes are known with d = 65, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-003838 | codetables-gf5-86-6

Minimum distance of a [86, 6] linear code over GF(5)

codes are known with d = 63, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-003839 | codetables-gf5-86-7

Minimum distance of a [86, 7] linear code over GF(5)

codes are known with d = 61, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-003840 | codetables-gf5-86-8

Minimum distance of a [86, 8] linear code over GF(5)

codes are known with d = 59, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-003841 | codetables-gf5-87-5

Minimum distance of a [87, 5] linear code over GF(5)

codes are known with d = 66, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-003842 | codetables-gf5-87-6

Minimum distance of a [87, 6] linear code over GF(5)

codes are known with d = 64, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-003843 | codetables-gf5-87-7

Minimum distance of a [87, 7] linear code over GF(5)

codes are known with d = 61, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-003844 | codetables-gf5-87-8

Minimum distance of a [87, 8] linear code over GF(5)

codes are known with d = 59, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-003845 | codetables-gf5-88-5

Minimum distance of a [88, 5] linear code over GF(5)

codes are known with d = 67, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-003846 | codetables-gf5-88-6

Minimum distance of a [88, 6] linear code over GF(5)

codes are known with d = 65, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-003847 | codetables-gf5-88-7

Minimum distance of a [88, 7] linear code over GF(5)

codes are known with d = 62, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-003848 | codetables-gf5-88-8

Minimum distance of a [88, 8] linear code over GF(5)

codes are known with d = 60, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-003849 | codetables-gf5-89-5

Minimum distance of a [89, 5] linear code over GF(5)

codes are known with d = 68, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-003850 | codetables-gf5-89-6

Minimum distance of a [89, 6] linear code over GF(5)

codes are known with d = 65, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-003851 | codetables-gf5-89-7

Minimum distance of a [89, 7] linear code over GF(5)

codes are known with d = 63, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-003852 | codetables-gf5-89-8

Minimum distance of a [89, 8] linear code over GF(5)

codes are known with d = 61, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-003853 | codetables-gf5-90-5

Minimum distance of a [90, 5] linear code over GF(5)

codes are known with d = 69, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-003854 | codetables-gf5-90-6

Minimum distance of a [90, 6] linear code over GF(5)

codes are known with d = 66, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-003855 | codetables-gf5-90-7

Minimum distance of a [90, 7] linear code over GF(5)

codes are known with d = 64, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-003856 | codetables-gf5-90-8

Minimum distance of a [90, 8] linear code over GF(5)

codes are known with d = 62, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-003857 | codetables-gf5-91-6

Minimum distance of a [91, 6] linear code over GF(5)

codes are known with d = 67, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-003858 | codetables-gf5-91-7

Minimum distance of a [91, 7] linear code over GF(5)

codes are known with d = 65, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-003859 | codetables-gf5-91-8

Minimum distance of a [91, 8] linear code over GF(5)

codes are known with d = 62, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-003860 | codetables-gf5-92-5

Minimum distance of a [92, 5] linear code over GF(5)

codes are known with d = 70, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-003861 | codetables-gf5-92-6

Minimum distance of a [92, 6] linear code over GF(5)

codes are known with d = 68, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-003862 | codetables-gf5-92-7

Minimum distance of a [92, 7] linear code over GF(5)

codes are known with d = 66, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-003863 | codetables-gf5-92-8

Minimum distance of a [92, 8] linear code over GF(5)

codes are known with d = 63, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-003864 | codetables-gf5-93-5

Minimum distance of a [93, 5] linear code over GF(5)

codes are known with d = 70, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-003865 | codetables-gf5-93-6

Minimum distance of a [93, 6] linear code over GF(5)

codes are known with d = 69, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-003866 | codetables-gf5-93-7

Minimum distance of a [93, 7] linear code over GF(5)

codes are known with d = 67, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-003867 | codetables-gf5-93-8

Minimum distance of a [93, 8] linear code over GF(5)

codes are known with d = 64, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-003868 | codetables-gf5-94-5

Minimum distance of a [94, 5] linear code over GF(5)

codes are known with d = 71, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-003869 | codetables-gf5-94-6

Minimum distance of a [94, 6] linear code over GF(5)

codes are known with d = 70, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-003870 | codetables-gf5-94-7

Minimum distance of a [94, 7] linear code over GF(5)

codes are known with d = 68, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-003871 | codetables-gf5-94-8

Minimum distance of a [94, 8] linear code over GF(5)

codes are known with d = 65, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-003872 | codetables-gf5-95-5

Minimum distance of a [95, 5] linear code over GF(5)

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-003873 | codetables-gf5-95-6

Minimum distance of a [95, 6] linear code over GF(5)

codes are known with d = 70, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-003874 | codetables-gf5-95-7

Minimum distance of a [95, 7] linear code over GF(5)

codes are known with d = 69, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-003875 | codetables-gf5-95-8

Minimum distance of a [95, 8] linear code over GF(5)

codes are known with d = 66, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-003876 | codetables-gf5-96-5

Minimum distance of a [96, 5] linear code over GF(5)

codes are known with d = 73, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-003877 | codetables-gf5-96-6

Minimum distance of a [96, 6] linear code over GF(5)

codes are known with d = 71, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-003878 | codetables-gf5-96-7

Minimum distance of a [96, 7] linear code over GF(5)

codes are known with d = 70, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-003879 | codetables-gf5-96-8

Minimum distance of a [96, 8] linear code over GF(5)

codes are known with d = 67, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-003880 | codetables-gf5-97-5

Minimum distance of a [97, 5] linear code over GF(5)

codes are known with d = 74, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-003881 | codetables-gf5-97-6

Minimum distance of a [97, 6] linear code over GF(5)

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-003882 | codetables-gf5-97-7

Minimum distance of a [97, 7] linear code over GF(5)

codes are known with d = 71, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-003883 | codetables-gf5-97-8

Minimum distance of a [97, 8] linear code over GF(5)

codes are known with d = 68, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-003884 | codetables-gf5-98-5

Minimum distance of a [98, 5] linear code over GF(5)

codes are known with d = 75, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-003885 | codetables-gf5-98-6

Minimum distance of a [98, 6] linear code over GF(5)

codes are known with d = 73, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-003886 | codetables-gf5-98-7

Minimum distance of a [98, 7] linear code over GF(5)

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-003887 | codetables-gf5-98-8

Minimum distance of a [98, 8] linear code over GF(5)

codes are known with d = 69, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-003888 | codetables-gf5-99-5

Minimum distance of a [99, 5] linear code over GF(5)

codes are known with d = 75, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-003889 | codetables-gf5-99-6

Minimum distance of a [99, 6] linear code over GF(5)

codes are known with d = 74, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-003890 | codetables-gf5-99-7

Minimum distance of a [99, 7] linear code over GF(5)

codes are known with d = 73, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-003891 | codetables-gf5-99-8

Minimum distance of a [99, 8] linear code over GF(5)

codes are known with d = 70, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-003892 | codetables-gf7-100-5

Minimum distance of a [100, 5] linear code over GF(7)

codes are known with d = 81, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-003893 | codetables-gf7-100-6

Minimum distance of a [100, 6] linear code over GF(7)

codes are known with d = 78, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-003894 | codetables-gf7-100-7

Minimum distance of a [100, 7] linear code over GF(7)

codes are known with d = 75, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-003895 | codetables-gf7-18-7

Minimum distance of a [18, 7] linear code over GF(7)

codes are known with d = 9, and no code can have d greater than 10.

来源状态 open | 2026-07-23 | 来源

V3P-003896 | codetables-gf7-19-7

Minimum distance of a [19, 7] linear code over GF(7)

codes are known with d = 10, and no code can have d greater than 11.

来源状态 open | 2026-07-23 | 来源

V3P-003897 | codetables-gf7-20-6

Minimum distance of a [20, 6] linear code over GF(7)

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-003898 | codetables-gf7-20-7

Minimum distance of a [20, 7] linear code over GF(7)

codes are known with d = 11, and no code can have d greater than 12.

来源状态 open | 2026-07-23 | 来源

V3P-003899 | codetables-gf7-21-4

Minimum distance of a [21, 4] linear code over GF(7)

codes are known with d = 15, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-003900 | codetables-gf7-21-5

Minimum distance of a [21, 5] linear code over GF(7)

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-003901 | codetables-gf7-21-6

Minimum distance of a [21, 6] linear code over GF(7)

codes are known with d = 13, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-003902 | codetables-gf7-21-7

Minimum distance of a [21, 7] linear code over GF(7)

codes are known with d = 12, and no code can have d greater than 13.

来源状态 open | 2026-07-23 | 来源

V3P-003903 | codetables-gf7-22-4

Minimum distance of a [22, 4] linear code over GF(7)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-003904 | codetables-gf7-22-5

Minimum distance of a [22, 5] linear code over GF(7)

codes are known with d = 15, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-003905 | codetables-gf7-22-6

Minimum distance of a [22, 6] linear code over GF(7)

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-003906 | codetables-gf7-22-7

Minimum distance of a [22, 7] linear code over GF(7)

codes are known with d = 13, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-003907 | codetables-gf7-23-4

Minimum distance of a [23, 4] linear code over GF(7)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-003908 | codetables-gf7-23-5

Minimum distance of a [23, 5] linear code over GF(7)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-003909 | codetables-gf7-23-6

Minimum distance of a [23, 6] linear code over GF(7)

codes are known with d = 15, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-003910 | codetables-gf7-23-7

Minimum distance of a [23, 7] linear code over GF(7)

codes are known with d = 13, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-003911 | codetables-gf7-24-5

Minimum distance of a [24, 5] linear code over GF(7)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-003912 | codetables-gf7-24-6

Minimum distance of a [24, 6] linear code over GF(7)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-003913 | codetables-gf7-24-7

Minimum distance of a [24, 7] linear code over GF(7)

codes are known with d = 14, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-003914 | codetables-gf7-25-5

Minimum distance of a [25, 5] linear code over GF(7)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-003915 | codetables-gf7-25-6

Minimum distance of a [25, 6] linear code over GF(7)

codes are known with d = 16, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-003916 | codetables-gf7-25-7

Minimum distance of a [25, 7] linear code over GF(7)

codes are known with d = 15, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-003917 | codetables-gf7-26-5

Minimum distance of a [26, 5] linear code over GF(7)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-003918 | codetables-gf7-26-6

Minimum distance of a [26, 6] linear code over GF(7)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-003919 | codetables-gf7-26-7

Minimum distance of a [26, 7] linear code over GF(7)

codes are known with d = 16, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-003920 | codetables-gf7-27-5

Minimum distance of a [27, 5] linear code over GF(7)

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-003921 | codetables-gf7-27-6

Minimum distance of a [27, 6] linear code over GF(7)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-003922 | codetables-gf7-27-7

Minimum distance of a [27, 7] linear code over GF(7)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-003923 | codetables-gf7-28-4

Minimum distance of a [28, 4] linear code over GF(7)

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-003924 | codetables-gf7-28-5

Minimum distance of a [28, 5] linear code over GF(7)

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-003925 | codetables-gf7-28-6

Minimum distance of a [28, 6] linear code over GF(7)

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-003926 | codetables-gf7-28-7

Minimum distance of a [28, 7] linear code over GF(7)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-003927 | codetables-gf7-29-4

Minimum distance of a [29, 4] linear code over GF(7)

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-003928 | codetables-gf7-29-5

Minimum distance of a [29, 5] linear code over GF(7)

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-003929 | codetables-gf7-29-6

Minimum distance of a [29, 6] linear code over GF(7)

codes are known with d = 19, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-003930 | codetables-gf7-29-7

Minimum distance of a [29, 7] linear code over GF(7)

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-003931 | codetables-gf7-30-4

Minimum distance of a [30, 4] linear code over GF(7)

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-003932 | codetables-gf7-30-5

Minimum distance of a [30, 5] linear code over GF(7)

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-003933 | codetables-gf7-30-6

Minimum distance of a [30, 6] linear code over GF(7)

codes are known with d = 20, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-003934 | codetables-gf7-30-7

Minimum distance of a [30, 7] linear code over GF(7)

codes are known with d = 19, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-003935 | codetables-gf7-31-5

Minimum distance of a [31, 5] linear code over GF(7)

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-003936 | codetables-gf7-31-6

Minimum distance of a [31, 6] linear code over GF(7)

codes are known with d = 21, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-003937 | codetables-gf7-31-7

Minimum distance of a [31, 7] linear code over GF(7)

codes are known with d = 19, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-003938 | codetables-gf7-32-5

Minimum distance of a [32, 5] linear code over GF(7)

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-003939 | codetables-gf7-32-6

Minimum distance of a [32, 6] linear code over GF(7)

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-003940 | codetables-gf7-32-7

Minimum distance of a [32, 7] linear code over GF(7)

codes are known with d = 20, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-003941 | codetables-gf7-33-5

Minimum distance of a [33, 5] linear code over GF(7)

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-003942 | codetables-gf7-33-6

Minimum distance of a [33, 6] linear code over GF(7)

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-003943 | codetables-gf7-33-7

Minimum distance of a [33, 7] linear code over GF(7)

codes are known with d = 21, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-003944 | codetables-gf7-34-4

Minimum distance of a [34, 4] linear code over GF(7)

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-003945 | codetables-gf7-34-5

Minimum distance of a [34, 5] linear code over GF(7)

codes are known with d = 25, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-003946 | codetables-gf7-34-6

Minimum distance of a [34, 6] linear code over GF(7)

codes are known with d = 23, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-003947 | codetables-gf7-34-7

Minimum distance of a [34, 7] linear code over GF(7)

codes are known with d = 22, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-003948 | codetables-gf7-35-4

Minimum distance of a [35, 4] linear code over GF(7)

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-003949 | codetables-gf7-35-5

Minimum distance of a [35, 5] linear code over GF(7)

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-003950 | codetables-gf7-35-6

Minimum distance of a [35, 6] linear code over GF(7)

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-003951 | codetables-gf7-35-7

Minimum distance of a [35, 7] linear code over GF(7)

codes are known with d = 23, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-003952 | codetables-gf7-36-4

Minimum distance of a [36, 4] linear code over GF(7)

codes are known with d = 28, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-003953 | codetables-gf7-36-5

Minimum distance of a [36, 5] linear code over GF(7)

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-003954 | codetables-gf7-36-6

Minimum distance of a [36, 6] linear code over GF(7)

codes are known with d = 25, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-003955 | codetables-gf7-36-7

Minimum distance of a [36, 7] linear code over GF(7)

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-003956 | codetables-gf7-37-4

Minimum distance of a [37, 4] linear code over GF(7)

codes are known with d = 29, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-003957 | codetables-gf7-37-5

Minimum distance of a [37, 5] linear code over GF(7)

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-003958 | codetables-gf7-37-6

Minimum distance of a [37, 6] linear code over GF(7)

codes are known with d = 26, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-003959 | codetables-gf7-37-7

Minimum distance of a [37, 7] linear code over GF(7)

codes are known with d = 24, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-003960 | codetables-gf7-38-5

Minimum distance of a [38, 5] linear code over GF(7)

codes are known with d = 28, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-003961 | codetables-gf7-38-6

Minimum distance of a [38, 6] linear code over GF(7)

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-003962 | codetables-gf7-38-7

Minimum distance of a [38, 7] linear code over GF(7)

codes are known with d = 25, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-003963 | codetables-gf7-39-5

Minimum distance of a [39, 5] linear code over GF(7)

codes are known with d = 29, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-003964 | codetables-gf7-39-6

Minimum distance of a [39, 6] linear code over GF(7)

codes are known with d = 27, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-003965 | codetables-gf7-39-7

Minimum distance of a [39, 7] linear code over GF(7)

codes are known with d = 26, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-003966 | codetables-gf7-40-5

Minimum distance of a [40, 5] linear code over GF(7)

codes are known with d = 30, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-003967 | codetables-gf7-40-6

Minimum distance of a [40, 6] linear code over GF(7)

codes are known with d = 28, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-003968 | codetables-gf7-40-7

Minimum distance of a [40, 7] linear code over GF(7)

codes are known with d = 27, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-003969 | codetables-gf7-41-5

Minimum distance of a [41, 5] linear code over GF(7)

codes are known with d = 31, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-003970 | codetables-gf7-41-6

Minimum distance of a [41, 6] linear code over GF(7)

codes are known with d = 29, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-003971 | codetables-gf7-41-7

Minimum distance of a [41, 7] linear code over GF(7)

codes are known with d = 28, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-003972 | codetables-gf7-42-5

Minimum distance of a [42, 5] linear code over GF(7)

codes are known with d = 32, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-003973 | codetables-gf7-42-6

Minimum distance of a [42, 6] linear code over GF(7)

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-003974 | codetables-gf7-42-7

Minimum distance of a [42, 7] linear code over GF(7)

codes are known with d = 29, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-003975 | codetables-gf7-43-5

Minimum distance of a [43, 5] linear code over GF(7)

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-003976 | codetables-gf7-43-6

Minimum distance of a [43, 6] linear code over GF(7)

codes are known with d = 31, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-003977 | codetables-gf7-43-7

Minimum distance of a [43, 7] linear code over GF(7)

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-003978 | codetables-gf7-44-5

Minimum distance of a [44, 5] linear code over GF(7)

codes are known with d = 33, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003979 | codetables-gf7-44-6

Minimum distance of a [44, 6] linear code over GF(7)

codes are known with d = 31, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-003980 | codetables-gf7-44-7

Minimum distance of a [44, 7] linear code over GF(7)

codes are known with d = 30, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-003981 | codetables-gf7-45-5

Minimum distance of a [45, 5] linear code over GF(7)

codes are known with d = 34, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003982 | codetables-gf7-45-6

Minimum distance of a [45, 6] linear code over GF(7)

codes are known with d = 32, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003983 | codetables-gf7-45-7

Minimum distance of a [45, 7] linear code over GF(7)

codes are known with d = 31, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-003984 | codetables-gf7-46-5

Minimum distance of a [46, 5] linear code over GF(7)

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-003985 | codetables-gf7-46-6

Minimum distance of a [46, 6] linear code over GF(7)

codes are known with d = 33, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003986 | codetables-gf7-46-7

Minimum distance of a [46, 7] linear code over GF(7)

codes are known with d = 32, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003987 | codetables-gf7-47-5

Minimum distance of a [47, 5] linear code over GF(7)

codes are known with d = 36, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-003988 | codetables-gf7-47-6

Minimum distance of a [47, 6] linear code over GF(7)

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-003989 | codetables-gf7-47-7

Minimum distance of a [47, 7] linear code over GF(7)

codes are known with d = 33, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-003990 | codetables-gf7-48-5

Minimum distance of a [48, 5] linear code over GF(7)

codes are known with d = 37, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-003991 | codetables-gf7-48-6

Minimum distance of a [48, 6] linear code over GF(7)

codes are known with d = 35, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-003992 | codetables-gf7-48-7

Minimum distance of a [48, 7] linear code over GF(7)

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-003993 | codetables-gf7-49-5

Minimum distance of a [49, 5] linear code over GF(7)

codes are known with d = 37, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-003994 | codetables-gf7-49-6

Minimum distance of a [49, 6] linear code over GF(7)

codes are known with d = 36, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-003995 | codetables-gf7-49-7

Minimum distance of a [49, 7] linear code over GF(7)

codes are known with d = 34, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-003996 | codetables-gf7-50-5

Minimum distance of a [50, 5] linear code over GF(7)

codes are known with d = 38, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-003997 | codetables-gf7-50-6

Minimum distance of a [50, 6] linear code over GF(7)

codes are known with d = 37, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-003998 | codetables-gf7-50-7

Minimum distance of a [50, 7] linear code over GF(7)

codes are known with d = 35, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-003999 | codetables-gf7-51-5

Minimum distance of a [51, 5] linear code over GF(7)

codes are known with d = 39, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-004000 | codetables-gf7-51-6

Minimum distance of a [51, 6] linear code over GF(7)

codes are known with d = 37, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-004001 | codetables-gf7-51-7

Minimum distance of a [51, 7] linear code over GF(7)

codes are known with d = 35, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-004002 | codetables-gf7-52-5

Minimum distance of a [52, 5] linear code over GF(7)

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-004003 | codetables-gf7-52-6

Minimum distance of a [52, 6] linear code over GF(7)

codes are known with d = 38, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-004004 | codetables-gf7-52-7

Minimum distance of a [52, 7] linear code over GF(7)

codes are known with d = 36, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-004005 | codetables-gf7-53-5

Minimum distance of a [53, 5] linear code over GF(7)

codes are known with d = 41, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-004006 | codetables-gf7-53-6

Minimum distance of a [53, 6] linear code over GF(7)

codes are known with d = 39, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-004007 | codetables-gf7-53-7

Minimum distance of a [53, 7] linear code over GF(7)

codes are known with d = 37, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-004008 | codetables-gf7-54-5

Minimum distance of a [54, 5] linear code over GF(7)

codes are known with d = 42, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-004009 | codetables-gf7-54-6

Minimum distance of a [54, 6] linear code over GF(7)

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-004010 | codetables-gf7-54-7

Minimum distance of a [54, 7] linear code over GF(7)

codes are known with d = 38, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-004011 | codetables-gf7-55-5

Minimum distance of a [55, 5] linear code over GF(7)

codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-004012 | codetables-gf7-55-6

Minimum distance of a [55, 6] linear code over GF(7)

codes are known with d = 41, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-004013 | codetables-gf7-55-7

Minimum distance of a [55, 7] linear code over GF(7)

codes are known with d = 39, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-004014 | codetables-gf7-56-5

Minimum distance of a [56, 5] linear code over GF(7)

codes are known with d = 43, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-004015 | codetables-gf7-56-6

Minimum distance of a [56, 6] linear code over GF(7)

codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-004016 | codetables-gf7-56-7

Minimum distance of a [56, 7] linear code over GF(7)

codes are known with d = 40, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-004017 | codetables-gf7-57-5

Minimum distance of a [57, 5] linear code over GF(7)

codes are known with d = 44, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-004018 | codetables-gf7-57-6

Minimum distance of a [57, 6] linear code over GF(7)

codes are known with d = 42, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-004019 | codetables-gf7-57-7

Minimum distance of a [57, 7] linear code over GF(7)

codes are known with d = 41, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-004020 | codetables-gf7-58-5

Minimum distance of a [58, 5] linear code over GF(7)

codes are known with d = 45, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-004021 | codetables-gf7-58-6

Minimum distance of a [58, 6] linear code over GF(7)

codes are known with d = 43, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-004022 | codetables-gf7-58-7

Minimum distance of a [58, 7] linear code over GF(7)

codes are known with d = 41, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-004023 | codetables-gf7-59-4

Minimum distance of a [59, 4] linear code over GF(7)

codes are known with d = 48, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-004024 | codetables-gf7-59-5

Minimum distance of a [59, 5] linear code over GF(7)

codes are known with d = 46, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-004025 | codetables-gf7-59-6

Minimum distance of a [59, 6] linear code over GF(7)

codes are known with d = 44, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-004026 | codetables-gf7-59-7

Minimum distance of a [59, 7] linear code over GF(7)

codes are known with d = 42, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-004027 | codetables-gf7-60-5

Minimum distance of a [60, 5] linear code over GF(7)

codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-004028 | codetables-gf7-60-6

Minimum distance of a [60, 6] linear code over GF(7)

codes are known with d = 45, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-004029 | codetables-gf7-60-7

Minimum distance of a [60, 7] linear code over GF(7)

codes are known with d = 42, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-004030 | codetables-gf7-61-5

Minimum distance of a [61, 5] linear code over GF(7)

codes are known with d = 47, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-004031 | codetables-gf7-61-6

Minimum distance of a [61, 6] linear code over GF(7)

codes are known with d = 45, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-004032 | codetables-gf7-61-7

Minimum distance of a [61, 7] linear code over GF(7)

codes are known with d = 43, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-004033 | codetables-gf7-62-4

Minimum distance of a [62, 4] linear code over GF(7)

codes are known with d = 50, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-004034 | codetables-gf7-62-5

Minimum distance of a [62, 5] linear code over GF(7)

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-004035 | codetables-gf7-62-6

Minimum distance of a [62, 6] linear code over GF(7)

codes are known with d = 46, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-004036 | codetables-gf7-62-7

Minimum distance of a [62, 7] linear code over GF(7)

codes are known with d = 44, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-004037 | codetables-gf7-63-4

Minimum distance of a [63, 4] linear code over GF(7)

codes are known with d = 51, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-004038 | codetables-gf7-63-5

Minimum distance of a [63, 5] linear code over GF(7)

codes are known with d = 49, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-004039 | codetables-gf7-63-6

Minimum distance of a [63, 6] linear code over GF(7)

codes are known with d = 47, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-004040 | codetables-gf7-63-7

Minimum distance of a [63, 7] linear code over GF(7)

codes are known with d = 45, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-004041 | codetables-gf7-64-4

Minimum distance of a [64, 4] linear code over GF(7)

codes are known with d = 52, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-004042 | codetables-gf7-64-5

Minimum distance of a [64, 5] linear code over GF(7)

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-004043 | codetables-gf7-64-6

Minimum distance of a [64, 6] linear code over GF(7)

codes are known with d = 48, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-004044 | codetables-gf7-64-7

Minimum distance of a [64, 7] linear code over GF(7)

codes are known with d = 46, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-004045 | codetables-gf7-65-4

Minimum distance of a [65, 4] linear code over GF(7)

codes are known with d = 53, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-004046 | codetables-gf7-65-5

Minimum distance of a [65, 5] linear code over GF(7)

codes are known with d = 51, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-004047 | codetables-gf7-65-6

Minimum distance of a [65, 6] linear code over GF(7)

codes are known with d = 49, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-004048 | codetables-gf7-65-7

Minimum distance of a [65, 7] linear code over GF(7)

codes are known with d = 47, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-004049 | codetables-gf7-66-4

Minimum distance of a [66, 4] linear code over GF(7)

codes are known with d = 54, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-004050 | codetables-gf7-66-5

Minimum distance of a [66, 5] linear code over GF(7)

codes are known with d = 52, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-004051 | codetables-gf7-66-6

Minimum distance of a [66, 6] linear code over GF(7)

codes are known with d = 49, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-004052 | codetables-gf7-66-7

Minimum distance of a [66, 7] linear code over GF(7)

codes are known with d = 47, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-004053 | codetables-gf7-67-4

Minimum distance of a [67, 4] linear code over GF(7)

codes are known with d = 55, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-004054 | codetables-gf7-67-5

Minimum distance of a [67, 5] linear code over GF(7)

codes are known with d = 52, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-004055 | codetables-gf7-67-6

Minimum distance of a [67, 6] linear code over GF(7)

codes are known with d = 50, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-004056 | codetables-gf7-67-7

Minimum distance of a [67, 7] linear code over GF(7)

codes are known with d = 48, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-004057 | codetables-gf7-68-5

Minimum distance of a [68, 5] linear code over GF(7)

codes are known with d = 53, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-004058 | codetables-gf7-68-6

Minimum distance of a [68, 6] linear code over GF(7)

codes are known with d = 51, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-004059 | codetables-gf7-68-7

Minimum distance of a [68, 7] linear code over GF(7)

codes are known with d = 49, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-004060 | codetables-gf7-69-4

Minimum distance of a [69, 4] linear code over GF(7)

codes are known with d = 56, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-004061 | codetables-gf7-69-5

Minimum distance of a [69, 5] linear code over GF(7)

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-004062 | codetables-gf7-69-6

Minimum distance of a [69, 6] linear code over GF(7)

codes are known with d = 52, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-004063 | codetables-gf7-69-7

Minimum distance of a [69, 7] linear code over GF(7)

codes are known with d = 50, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-004064 | codetables-gf7-70-4

Minimum distance of a [70, 4] linear code over GF(7)

codes are known with d = 57, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-004065 | codetables-gf7-70-5

Minimum distance of a [70, 5] linear code over GF(7)

codes are known with d = 55, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-004066 | codetables-gf7-70-6

Minimum distance of a [70, 6] linear code over GF(7)

codes are known with d = 53, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-004067 | codetables-gf7-70-7

Minimum distance of a [70, 7] linear code over GF(7)

codes are known with d = 51, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-004068 | codetables-gf7-71-4

Minimum distance of a [71, 4] linear code over GF(7)

codes are known with d = 58, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-004069 | codetables-gf7-71-5

Minimum distance of a [71, 5] linear code over GF(7)

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-004070 | codetables-gf7-71-6

Minimum distance of a [71, 6] linear code over GF(7)

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-004071 | codetables-gf7-71-7

Minimum distance of a [71, 7] linear code over GF(7)

codes are known with d = 52, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-004072 | codetables-gf7-72-4

Minimum distance of a [72, 4] linear code over GF(7)

codes are known with d = 59, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-004073 | codetables-gf7-72-5

Minimum distance of a [72, 5] linear code over GF(7)

codes are known with d = 57, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-004074 | codetables-gf7-72-6

Minimum distance of a [72, 6] linear code over GF(7)

codes are known with d = 55, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-004075 | codetables-gf7-72-7

Minimum distance of a [72, 7] linear code over GF(7)

codes are known with d = 53, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-004076 | codetables-gf7-73-4

Minimum distance of a [73, 4] linear code over GF(7)

codes are known with d = 60, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-004077 | codetables-gf7-73-5

Minimum distance of a [73, 5] linear code over GF(7)

codes are known with d = 57, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-004078 | codetables-gf7-73-6

Minimum distance of a [73, 6] linear code over GF(7)

codes are known with d = 55, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-004079 | codetables-gf7-73-7

Minimum distance of a [73, 7] linear code over GF(7)

codes are known with d = 53, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-004080 | codetables-gf7-74-4

Minimum distance of a [74, 4] linear code over GF(7)

codes are known with d = 61, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-004081 | codetables-gf7-74-5

Minimum distance of a [74, 5] linear code over GF(7)

codes are known with d = 58, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-004082 | codetables-gf7-74-6

Minimum distance of a [74, 6] linear code over GF(7)

codes are known with d = 56, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-004083 | codetables-gf7-74-7

Minimum distance of a [74, 7] linear code over GF(7)

codes are known with d = 54, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-004084 | codetables-gf7-75-4

Minimum distance of a [75, 4] linear code over GF(7)

codes are known with d = 62, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-004085 | codetables-gf7-75-5

Minimum distance of a [75, 5] linear code over GF(7)

codes are known with d = 59, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-004086 | codetables-gf7-75-6

Minimum distance of a [75, 6] linear code over GF(7)

codes are known with d = 57, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-004087 | codetables-gf7-75-7

Minimum distance of a [75, 7] linear code over GF(7)

codes are known with d = 55, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-004088 | codetables-gf7-76-4

Minimum distance of a [76, 4] linear code over GF(7)

codes are known with d = 62, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-004089 | codetables-gf7-76-5

Minimum distance of a [76, 5] linear code over GF(7)

codes are known with d = 60, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-004090 | codetables-gf7-76-6

Minimum distance of a [76, 6] linear code over GF(7)

codes are known with d = 58, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-004091 | codetables-gf7-76-7

Minimum distance of a [76, 7] linear code over GF(7)

codes are known with d = 56, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-004092 | codetables-gf7-77-4

Minimum distance of a [77, 4] linear code over GF(7)

codes are known with d = 63, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-004093 | codetables-gf7-77-5

Minimum distance of a [77, 5] linear code over GF(7)

codes are known with d = 61, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-004094 | codetables-gf7-77-6

Minimum distance of a [77, 6] linear code over GF(7)

codes are known with d = 58, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-004095 | codetables-gf7-77-7

Minimum distance of a [77, 7] linear code over GF(7)

codes are known with d = 57, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-004096 | codetables-gf7-78-4

Minimum distance of a [78, 4] linear code over GF(7)

codes are known with d = 64, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-004097 | codetables-gf7-78-5

Minimum distance of a [78, 5] linear code over GF(7)

codes are known with d = 62, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-004098 | codetables-gf7-78-6

Minimum distance of a [78, 6] linear code over GF(7)

codes are known with d = 59, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-004099 | codetables-gf7-78-7

Minimum distance of a [78, 7] linear code over GF(7)

codes are known with d = 57, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-004100 | codetables-gf7-79-4

Minimum distance of a [79, 4] linear code over GF(7)

codes are known with d = 65, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-004101 | codetables-gf7-79-5

Minimum distance of a [79, 5] linear code over GF(7)

codes are known with d = 63, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-004102 | codetables-gf7-79-6

Minimum distance of a [79, 6] linear code over GF(7)

codes are known with d = 60, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-004103 | codetables-gf7-79-7

Minimum distance of a [79, 7] linear code over GF(7)

codes are known with d = 58, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-004104 | codetables-gf7-80-4

Minimum distance of a [80, 4] linear code over GF(7)

codes are known with d = 66, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-004105 | codetables-gf7-80-5

Minimum distance of a [80, 5] linear code over GF(7)

codes are known with d = 63, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-004106 | codetables-gf7-80-6

Minimum distance of a [80, 6] linear code over GF(7)

codes are known with d = 61, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-004107 | codetables-gf7-80-7

Minimum distance of a [80, 7] linear code over GF(7)

codes are known with d = 59, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-004108 | codetables-gf7-81-4

Minimum distance of a [81, 4] linear code over GF(7)

codes are known with d = 67, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-004109 | codetables-gf7-81-5

Minimum distance of a [81, 5] linear code over GF(7)

codes are known with d = 64, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-004110 | codetables-gf7-81-6

Minimum distance of a [81, 6] linear code over GF(7)

codes are known with d = 62, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-004111 | codetables-gf7-81-7

Minimum distance of a [81, 7] linear code over GF(7)

codes are known with d = 60, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-004112 | codetables-gf7-82-4

Minimum distance of a [82, 4] linear code over GF(7)

codes are known with d = 68, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-004113 | codetables-gf7-82-5

Minimum distance of a [82, 5] linear code over GF(7)

codes are known with d = 65, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-004114 | codetables-gf7-82-6

Minimum distance of a [82, 6] linear code over GF(7)

codes are known with d = 63, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-004115 | codetables-gf7-82-7

Minimum distance of a [82, 7] linear code over GF(7)

codes are known with d = 60, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-004116 | codetables-gf7-83-5

Minimum distance of a [83, 5] linear code over GF(7)

codes are known with d = 66, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-004117 | codetables-gf7-83-6

Minimum distance of a [83, 6] linear code over GF(7)

codes are known with d = 64, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-004118 | codetables-gf7-83-7

Minimum distance of a [83, 7] linear code over GF(7)

codes are known with d = 61, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-004119 | codetables-gf7-84-5

Minimum distance of a [84, 5] linear code over GF(7)

codes are known with d = 67, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-004120 | codetables-gf7-84-6

Minimum distance of a [84, 6] linear code over GF(7)

codes are known with d = 64, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-004121 | codetables-gf7-84-7

Minimum distance of a [84, 7] linear code over GF(7)

codes are known with d = 62, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-004122 | codetables-gf7-85-4

Minimum distance of a [85, 4] linear code over GF(7)

codes are known with d = 70, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-004123 | codetables-gf7-85-5

Minimum distance of a [85, 5] linear code over GF(7)

codes are known with d = 68, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-004124 | codetables-gf7-85-6

Minimum distance of a [85, 6] linear code over GF(7)

codes are known with d = 65, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-004125 | codetables-gf7-85-7

Minimum distance of a [85, 7] linear code over GF(7)

codes are known with d = 63, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-004126 | codetables-gf7-86-4

Minimum distance of a [86, 4] linear code over GF(7)

codes are known with d = 71, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-004127 | codetables-gf7-86-5

Minimum distance of a [86, 5] linear code over GF(7)

codes are known with d = 68, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-004128 | codetables-gf7-86-6

Minimum distance of a [86, 6] linear code over GF(7)

codes are known with d = 66, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-004129 | codetables-gf7-86-7

Minimum distance of a [86, 7] linear code over GF(7)

codes are known with d = 64, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-004130 | codetables-gf7-87-4

Minimum distance of a [87, 4] linear code over GF(7)

codes are known with d = 72, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-004131 | codetables-gf7-87-5

Minimum distance of a [87, 5] linear code over GF(7)

codes are known with d = 69, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-004132 | codetables-gf7-87-6

Minimum distance of a [87, 6] linear code over GF(7)

codes are known with d = 67, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-004133 | codetables-gf7-87-7

Minimum distance of a [87, 7] linear code over GF(7)

codes are known with d = 65, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-004134 | codetables-gf7-88-4

Minimum distance of a [88, 4] linear code over GF(7)

codes are known with d = 73, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-004135 | codetables-gf7-88-5

Minimum distance of a [88, 5] linear code over GF(7)

codes are known with d = 70, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-004136 | codetables-gf7-88-6

Minimum distance of a [88, 6] linear code over GF(7)

codes are known with d = 68, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-004137 | codetables-gf7-88-7

Minimum distance of a [88, 7] linear code over GF(7)

codes are known with d = 66, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-004138 | codetables-gf7-89-4

Minimum distance of a [89, 4] linear code over GF(7)

codes are known with d = 74, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-004139 | codetables-gf7-89-5

Minimum distance of a [89, 5] linear code over GF(7)

codes are known with d = 71, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-004140 | codetables-gf7-89-6

Minimum distance of a [89, 6] linear code over GF(7)

codes are known with d = 69, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-004141 | codetables-gf7-89-7

Minimum distance of a [89, 7] linear code over GF(7)

codes are known with d = 66, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-004142 | codetables-gf7-90-4

Minimum distance of a [90, 4] linear code over GF(7)

codes are known with d = 75, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-004143 | codetables-gf7-90-5

Minimum distance of a [90, 5] linear code over GF(7)

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-004144 | codetables-gf7-90-6

Minimum distance of a [90, 6] linear code over GF(7)

codes are known with d = 69, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-004145 | codetables-gf7-90-7

Minimum distance of a [90, 7] linear code over GF(7)

codes are known with d = 67, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-004146 | codetables-gf7-91-5

Minimum distance of a [91, 5] linear code over GF(7)

codes are known with d = 73, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-004147 | codetables-gf7-91-6

Minimum distance of a [91, 6] linear code over GF(7)

codes are known with d = 70, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-004148 | codetables-gf7-91-7

Minimum distance of a [91, 7] linear code over GF(7)

codes are known with d = 67, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-004149 | codetables-gf7-92-5

Minimum distance of a [92, 5] linear code over GF(7)

codes are known with d = 73, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-004150 | codetables-gf7-92-6

Minimum distance of a [92, 6] linear code over GF(7)

codes are known with d = 71, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-004151 | codetables-gf7-92-7

Minimum distance of a [92, 7] linear code over GF(7)

codes are known with d = 68, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-004152 | codetables-gf7-93-4

Minimum distance of a [93, 4] linear code over GF(7)

codes are known with d = 77, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-004153 | codetables-gf7-93-5

Minimum distance of a [93, 5] linear code over GF(7)

codes are known with d = 74, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-004154 | codetables-gf7-93-6

Minimum distance of a [93, 6] linear code over GF(7)

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-004155 | codetables-gf7-93-7

Minimum distance of a [93, 7] linear code over GF(7)

codes are known with d = 69, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-004156 | codetables-gf7-94-4

Minimum distance of a [94, 4] linear code over GF(7)

codes are known with d = 78, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-004157 | codetables-gf7-94-5

Minimum distance of a [94, 5] linear code over GF(7)

codes are known with d = 75, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-004158 | codetables-gf7-94-6

Minimum distance of a [94, 6] linear code over GF(7)

codes are known with d = 73, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-004159 | codetables-gf7-94-7

Minimum distance of a [94, 7] linear code over GF(7)

codes are known with d = 70, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-004160 | codetables-gf7-95-4

Minimum distance of a [95, 4] linear code over GF(7)

codes are known with d = 79, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-004161 | codetables-gf7-95-5

Minimum distance of a [95, 5] linear code over GF(7)

codes are known with d = 76, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-004162 | codetables-gf7-95-6

Minimum distance of a [95, 6] linear code over GF(7)

codes are known with d = 74, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-004163 | codetables-gf7-95-7

Minimum distance of a [95, 7] linear code over GF(7)

codes are known with d = 71, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-004164 | codetables-gf7-96-4

Minimum distance of a [96, 4] linear code over GF(7)

codes are known with d = 80, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-004165 | codetables-gf7-96-5

Minimum distance of a [96, 5] linear code over GF(7)

codes are known with d = 77, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-004166 | codetables-gf7-96-6

Minimum distance of a [96, 6] linear code over GF(7)

codes are known with d = 75, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-004167 | codetables-gf7-96-7

Minimum distance of a [96, 7] linear code over GF(7)

codes are known with d = 72, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-004168 | codetables-gf7-97-4

Minimum distance of a [97, 4] linear code over GF(7)

codes are known with d = 81, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-004169 | codetables-gf7-97-5

Minimum distance of a [97, 5] linear code over GF(7)

codes are known with d = 78, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-004170 | codetables-gf7-97-6

Minimum distance of a [97, 6] linear code over GF(7)

codes are known with d = 75, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-004171 | codetables-gf7-97-7

Minimum distance of a [97, 7] linear code over GF(7)

codes are known with d = 73, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-004172 | codetables-gf7-98-5

Minimum distance of a [98, 5] linear code over GF(7)

codes are known with d = 79, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-004173 | codetables-gf7-98-6

Minimum distance of a [98, 6] linear code over GF(7)

codes are known with d = 76, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-004174 | codetables-gf7-98-7

Minimum distance of a [98, 7] linear code over GF(7)

codes are known with d = 74, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-004175 | codetables-gf7-99-5

Minimum distance of a [99, 5] linear code over GF(7)

codes are known with d = 80, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-004176 | codetables-gf7-99-6

Minimum distance of a [99, 6] linear code over GF(7)

codes are known with d = 77, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-004177 | codetables-gf7-99-7

Minimum distance of a [99, 7] linear code over GF(7)

codes are known with d = 74, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-004178 | codetables-gf8-100-4

Minimum distance of a [100, 4] linear code over GF(8)

codes are known with d = 85, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-004179 | codetables-gf8-100-5

Minimum distance of a [100, 5] linear code over GF(8)

codes are known with d = 82, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-004180 | codetables-gf8-100-6

Minimum distance of a [100, 6] linear code over GF(8)

codes are known with d = 79, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-004181 | codetables-gf8-101-5

Minimum distance of a [101, 5] linear code over GF(8)

codes are known with d = 83, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-004182 | codetables-gf8-101-6

Minimum distance of a [101, 6] linear code over GF(8)

codes are known with d = 80, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-004183 | codetables-gf8-102-5

Minimum distance of a [102, 5] linear code over GF(8)

codes are known with d = 84, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-004184 | codetables-gf8-102-6

Minimum distance of a [102, 6] linear code over GF(8)

codes are known with d = 81, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-004185 | codetables-gf8-103-5

Minimum distance of a [103, 5] linear code over GF(8)

codes are known with d = 84, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-004186 | codetables-gf8-103-6

Minimum distance of a [103, 6] linear code over GF(8)

codes are known with d = 82, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-004187 | codetables-gf8-104-4

Minimum distance of a [104, 4] linear code over GF(8)

codes are known with d = 88, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-004188 | codetables-gf8-104-5

Minimum distance of a [104, 5] linear code over GF(8)

codes are known with d = 85, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-004189 | codetables-gf8-104-6

Minimum distance of a [104, 6] linear code over GF(8)

codes are known with d = 83, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-004190 | codetables-gf8-105-4

Minimum distance of a [105, 4] linear code over GF(8)

codes are known with d = 89, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-004191 | codetables-gf8-105-5

Minimum distance of a [105, 5] linear code over GF(8)

codes are known with d = 86, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-004192 | codetables-gf8-105-6

Minimum distance of a [105, 6] linear code over GF(8)

codes are known with d = 84, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-004193 | codetables-gf8-106-4

Minimum distance of a [106, 4] linear code over GF(8)

codes are known with d = 90, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-004194 | codetables-gf8-106-5

Minimum distance of a [106, 5] linear code over GF(8)

codes are known with d = 87, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-004195 | codetables-gf8-106-6

Minimum distance of a [106, 6] linear code over GF(8)

codes are known with d = 85, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-004196 | codetables-gf8-107-4

Minimum distance of a [107, 4] linear code over GF(8)

codes are known with d = 91, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-004197 | codetables-gf8-107-5

Minimum distance of a [107, 5] linear code over GF(8)

codes are known with d = 88, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-004198 | codetables-gf8-107-6

Minimum distance of a [107, 6] linear code over GF(8)

codes are known with d = 86, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-004199 | codetables-gf8-108-4

Minimum distance of a [108, 4] linear code over GF(8)

codes are known with d = 92, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-004200 | codetables-gf8-108-5

Minimum distance of a [108, 5] linear code over GF(8)

codes are known with d = 89, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-004201 | codetables-gf8-108-6

Minimum distance of a [108, 6] linear code over GF(8)

codes are known with d = 87, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-004202 | codetables-gf8-109-4

Minimum distance of a [109, 4] linear code over GF(8)

codes are known with d = 93, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-004203 | codetables-gf8-109-5

Minimum distance of a [109, 5] linear code over GF(8)

codes are known with d = 90, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-004204 | codetables-gf8-109-6

Minimum distance of a [109, 6] linear code over GF(8)

codes are known with d = 88, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-004205 | codetables-gf8-110-4

Minimum distance of a [110, 4] linear code over GF(8)

codes are known with d = 94, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-004206 | codetables-gf8-110-5

Minimum distance of a [110, 5] linear code over GF(8)

codes are known with d = 91, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-004207 | codetables-gf8-110-6

Minimum distance of a [110, 6] linear code over GF(8)

codes are known with d = 89, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-004208 | codetables-gf8-111-5

Minimum distance of a [111, 5] linear code over GF(8)

codes are known with d = 92, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-004209 | codetables-gf8-111-6

Minimum distance of a [111, 6] linear code over GF(8)

codes are known with d = 90, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-004210 | codetables-gf8-112-5

Minimum distance of a [112, 5] linear code over GF(8)

codes are known with d = 92, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-004211 | codetables-gf8-112-6

Minimum distance of a [112, 6] linear code over GF(8)

codes are known with d = 91, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-004212 | codetables-gf8-113-5

Minimum distance of a [113, 5] linear code over GF(8)

codes are known with d = 93, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-004213 | codetables-gf8-113-6

Minimum distance of a [113, 6] linear code over GF(8)

codes are known with d = 92, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-004214 | codetables-gf8-114-5

Minimum distance of a [114, 5] linear code over GF(8)

codes are known with d = 94, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-004215 | codetables-gf8-114-6

Minimum distance of a [114, 6] linear code over GF(8)

codes are known with d = 93, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-004216 | codetables-gf8-115-5

Minimum distance of a [115, 5] linear code over GF(8)

codes are known with d = 95, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-004217 | codetables-gf8-115-6

Minimum distance of a [115, 6] linear code over GF(8)

codes are known with d = 94, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-004218 | codetables-gf8-116-5

Minimum distance of a [116, 5] linear code over GF(8)

codes are known with d = 96, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-004219 | codetables-gf8-116-6

Minimum distance of a [116, 6] linear code over GF(8)

codes are known with d = 95, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-004220 | codetables-gf8-117-5

Minimum distance of a [117, 5] linear code over GF(8)

codes are known with d = 97, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-004221 | codetables-gf8-117-6

Minimum distance of a [117, 6] linear code over GF(8)

codes are known with d = 96, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-004222 | codetables-gf8-118-5

Minimum distance of a [118, 5] linear code over GF(8)

codes are known with d = 98, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-004223 | codetables-gf8-118-6

Minimum distance of a [118, 6] linear code over GF(8)

codes are known with d = 96, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-004224 | codetables-gf8-119-5

Minimum distance of a [119, 5] linear code over GF(8)

codes are known with d = 99, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-004225 | codetables-gf8-119-6

Minimum distance of a [119, 6] linear code over GF(8)

codes are known with d = 96, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-004226 | codetables-gf8-120-5

Minimum distance of a [120, 5] linear code over GF(8)

codes are known with d = 100, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-004227 | codetables-gf8-120-6

Minimum distance of a [120, 6] linear code over GF(8)

codes are known with d = 96, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-004228 | codetables-gf8-121-5

Minimum distance of a [121, 5] linear code over GF(8)

codes are known with d = 100, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-004229 | codetables-gf8-121-6

Minimum distance of a [121, 6] linear code over GF(8)

codes are known with d = 97, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-004230 | codetables-gf8-122-5

Minimum distance of a [122, 5] linear code over GF(8)

codes are known with d = 101, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-004231 | codetables-gf8-122-6

Minimum distance of a [122, 6] linear code over GF(8)

codes are known with d = 98, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-004232 | codetables-gf8-123-5

Minimum distance of a [123, 5] linear code over GF(8)

codes are known with d = 102, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-004233 | codetables-gf8-123-6

Minimum distance of a [123, 6] linear code over GF(8)

codes are known with d = 99, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-004234 | codetables-gf8-124-5

Minimum distance of a [124, 5] linear code over GF(8)

codes are known with d = 103, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-004235 | codetables-gf8-124-6

Minimum distance of a [124, 6] linear code over GF(8)

codes are known with d = 100, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-004236 | codetables-gf8-125-5

Minimum distance of a [125, 5] linear code over GF(8)

codes are known with d = 104, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-004237 | codetables-gf8-125-6

Minimum distance of a [125, 6] linear code over GF(8)

codes are known with d = 101, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-004238 | codetables-gf8-126-5

Minimum distance of a [126, 5] linear code over GF(8)

codes are known with d = 104, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-004239 | codetables-gf8-126-6

Minimum distance of a [126, 6] linear code over GF(8)

codes are known with d = 102, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-004240 | codetables-gf8-127-5

Minimum distance of a [127, 5] linear code over GF(8)

codes are known with d = 105, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-004241 | codetables-gf8-127-6

Minimum distance of a [127, 6] linear code over GF(8)

codes are known with d = 103, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-004242 | codetables-gf8-128-5

Minimum distance of a [128, 5] linear code over GF(8)

codes are known with d = 106, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-004243 | codetables-gf8-128-6

Minimum distance of a [128, 6] linear code over GF(8)

codes are known with d = 104, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-004244 | codetables-gf8-129-5

Minimum distance of a [129, 5] linear code over GF(8)

codes are known with d = 107, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-004245 | codetables-gf8-129-6

Minimum distance of a [129, 6] linear code over GF(8)

codes are known with d = 104, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-004246 | codetables-gf8-130-5

Minimum distance of a [130, 5] linear code over GF(8)

codes are known with d = 108, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-004247 | codetables-gf8-130-6

Minimum distance of a [130, 6] linear code over GF(8)

codes are known with d = 105, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-004248 | codetables-gf8-21-5

Minimum distance of a [21, 5] linear code over GF(8)

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-004249 | codetables-gf8-21-6

Minimum distance of a [21, 6] linear code over GF(8)

codes are known with d = 13, and no code can have d greater than 14.

来源状态 open | 2026-07-23 | 来源

V3P-004250 | codetables-gf8-22-5

Minimum distance of a [22, 5] linear code over GF(8)

codes are known with d = 15, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-004251 | codetables-gf8-22-6

Minimum distance of a [22, 6] linear code over GF(8)

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-004252 | codetables-gf8-23-5

Minimum distance of a [23, 5] linear code over GF(8)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-004253 | codetables-gf8-23-6

Minimum distance of a [23, 6] linear code over GF(8)

codes are known with d = 15, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-004254 | codetables-gf8-24-5

Minimum distance of a [24, 5] linear code over GF(8)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-004255 | codetables-gf8-24-6

Minimum distance of a [24, 6] linear code over GF(8)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-004256 | codetables-gf8-25-5

Minimum distance of a [25, 5] linear code over GF(8)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-004257 | codetables-gf8-25-6

Minimum distance of a [25, 6] linear code over GF(8)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-004258 | codetables-gf8-26-4

Minimum distance of a [26, 4] linear code over GF(8)

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-004259 | codetables-gf8-26-5

Minimum distance of a [26, 5] linear code over GF(8)

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-004260 | codetables-gf8-26-6

Minimum distance of a [26, 6] linear code over GF(8)

codes are known with d = 17, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-004261 | codetables-gf8-27-5

Minimum distance of a [27, 5] linear code over GF(8)

codes are known with d = 19, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-004262 | codetables-gf8-27-6

Minimum distance of a [27, 6] linear code over GF(8)

codes are known with d = 18, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-004263 | codetables-gf8-28-5

Minimum distance of a [28, 5] linear code over GF(8)

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-004264 | codetables-gf8-28-6

Minimum distance of a [28, 6] linear code over GF(8)

codes are known with d = 19, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-004265 | codetables-gf8-29-5

Minimum distance of a [29, 5] linear code over GF(8)

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-004266 | codetables-gf8-29-6

Minimum distance of a [29, 6] linear code over GF(8)

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-004267 | codetables-gf8-30-5

Minimum distance of a [30, 5] linear code over GF(8)

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-004268 | codetables-gf8-30-6

Minimum distance of a [30, 6] linear code over GF(8)

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-004269 | codetables-gf8-31-4

Minimum distance of a [31, 4] linear code over GF(8)

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-004270 | codetables-gf8-31-5

Minimum distance of a [31, 5] linear code over GF(8)

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-004271 | codetables-gf8-31-6

Minimum distance of a [31, 6] linear code over GF(8)

codes are known with d = 21, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-004272 | codetables-gf8-32-4

Minimum distance of a [32, 4] linear code over GF(8)

codes are known with d = 25, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-004273 | codetables-gf8-32-5

Minimum distance of a [32, 5] linear code over GF(8)

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-004274 | codetables-gf8-32-6

Minimum distance of a [32, 6] linear code over GF(8)

codes are known with d = 22, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-004275 | codetables-gf8-33-4

Minimum distance of a [33, 4] linear code over GF(8)

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-004276 | codetables-gf8-33-5

Minimum distance of a [33, 5] linear code over GF(8)

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-004277 | codetables-gf8-33-6

Minimum distance of a [33, 6] linear code over GF(8)

codes are known with d = 23, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-004278 | codetables-gf8-34-4

Minimum distance of a [34, 4] linear code over GF(8)

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-004279 | codetables-gf8-34-5

Minimum distance of a [34, 5] linear code over GF(8)

codes are known with d = 25, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-004280 | codetables-gf8-34-6

Minimum distance of a [34, 6] linear code over GF(8)

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-004281 | codetables-gf8-35-5

Minimum distance of a [35, 5] linear code over GF(8)

codes are known with d = 26, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-004282 | codetables-gf8-35-6

Minimum distance of a [35, 6] linear code over GF(8)

codes are known with d = 25, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-004283 | codetables-gf8-36-5

Minimum distance of a [36, 5] linear code over GF(8)

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-004284 | codetables-gf8-36-6

Minimum distance of a [36, 6] linear code over GF(8)

codes are known with d = 26, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-004285 | codetables-gf8-37-5

Minimum distance of a [37, 5] linear code over GF(8)

codes are known with d = 28, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-004286 | codetables-gf8-37-6

Minimum distance of a [37, 6] linear code over GF(8)

codes are known with d = 27, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-004287 | codetables-gf8-38-5

Minimum distance of a [38, 5] linear code over GF(8)

codes are known with d = 29, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-004288 | codetables-gf8-38-6

Minimum distance of a [38, 6] linear code over GF(8)

codes are known with d = 28, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-004289 | codetables-gf8-39-5

Minimum distance of a [39, 5] linear code over GF(8)

codes are known with d = 29, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-004290 | codetables-gf8-39-6

Minimum distance of a [39, 6] linear code over GF(8)

codes are known with d = 28, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-004291 | codetables-gf8-40-4

Minimum distance of a [40, 4] linear code over GF(8)

codes are known with d = 32, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-004292 | codetables-gf8-40-5

Minimum distance of a [40, 5] linear code over GF(8)

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-004293 | codetables-gf8-40-6

Minimum distance of a [40, 6] linear code over GF(8)

codes are known with d = 29, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-004294 | codetables-gf8-41-4

Minimum distance of a [41, 4] linear code over GF(8)

codes are known with d = 33, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-004295 | codetables-gf8-41-5

Minimum distance of a [41, 5] linear code over GF(8)

codes are known with d = 31, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-004296 | codetables-gf8-41-6

Minimum distance of a [41, 6] linear code over GF(8)

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-004297 | codetables-gf8-42-4

Minimum distance of a [42, 4] linear code over GF(8)

codes are known with d = 34, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-004298 | codetables-gf8-42-5

Minimum distance of a [42, 5] linear code over GF(8)

codes are known with d = 32, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-004299 | codetables-gf8-42-6

Minimum distance of a [42, 6] linear code over GF(8)

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-004300 | codetables-gf8-43-4

Minimum distance of a [43, 4] linear code over GF(8)

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-004301 | codetables-gf8-43-5

Minimum distance of a [43, 5] linear code over GF(8)

codes are known with d = 33, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-004302 | codetables-gf8-43-6

Minimum distance of a [43, 6] linear code over GF(8)

codes are known with d = 31, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-004303 | codetables-gf8-44-5

Minimum distance of a [44, 5] linear code over GF(8)

codes are known with d = 34, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-004304 | codetables-gf8-44-6

Minimum distance of a [44, 6] linear code over GF(8)

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-004305 | codetables-gf8-45-5

Minimum distance of a [45, 5] linear code over GF(8)

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-004306 | codetables-gf8-45-6

Minimum distance of a [45, 6] linear code over GF(8)

codes are known with d = 33, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-004307 | codetables-gf8-46-4

Minimum distance of a [46, 4] linear code over GF(8)

codes are known with d = 37, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-004308 | codetables-gf8-46-5

Minimum distance of a [46, 5] linear code over GF(8)

codes are known with d = 36, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-004309 | codetables-gf8-46-6

Minimum distance of a [46, 6] linear code over GF(8)

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-004310 | codetables-gf8-47-4

Minimum distance of a [47, 4] linear code over GF(8)

codes are known with d = 38, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-004311 | codetables-gf8-47-5

Minimum distance of a [47, 5] linear code over GF(8)

codes are known with d = 37, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-004312 | codetables-gf8-47-6

Minimum distance of a [47, 6] linear code over GF(8)

codes are known with d = 35, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-004313 | codetables-gf8-48-4

Minimum distance of a [48, 4] linear code over GF(8)

codes are known with d = 39, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-004314 | codetables-gf8-48-5

Minimum distance of a [48, 5] linear code over GF(8)

codes are known with d = 38, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-004315 | codetables-gf8-48-6

Minimum distance of a [48, 6] linear code over GF(8)

codes are known with d = 36, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-004316 | codetables-gf8-49-4

Minimum distance of a [49, 4] linear code over GF(8)

codes are known with d = 40, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-004317 | codetables-gf8-49-5

Minimum distance of a [49, 5] linear code over GF(8)

codes are known with d = 38, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-004318 | codetables-gf8-49-6

Minimum distance of a [49, 6] linear code over GF(8)

codes are known with d = 36, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-004319 | codetables-gf8-50-4

Minimum distance of a [50, 4] linear code over GF(8)

codes are known with d = 41, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-004320 | codetables-gf8-50-5

Minimum distance of a [50, 5] linear code over GF(8)

codes are known with d = 39, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-004321 | codetables-gf8-50-6

Minimum distance of a [50, 6] linear code over GF(8)

codes are known with d = 37, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-004322 | codetables-gf8-51-5

Minimum distance of a [51, 5] linear code over GF(8)

codes are known with d = 40, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-004323 | codetables-gf8-51-6

Minimum distance of a [51, 6] linear code over GF(8)

codes are known with d = 38, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-004324 | codetables-gf8-52-5

Minimum distance of a [52, 5] linear code over GF(8)

codes are known with d = 41, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-004325 | codetables-gf8-52-6

Minimum distance of a [52, 6] linear code over GF(8)

codes are known with d = 39, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-004326 | codetables-gf8-53-5

Minimum distance of a [53, 5] linear code over GF(8)

codes are known with d = 42, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-004327 | codetables-gf8-53-6

Minimum distance of a [53, 6] linear code over GF(8)

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-004328 | codetables-gf8-54-5

Minimum distance of a [54, 5] linear code over GF(8)

codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-004329 | codetables-gf8-54-6

Minimum distance of a [54, 6] linear code over GF(8)

codes are known with d = 41, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-004330 | codetables-gf8-55-5

Minimum distance of a [55, 5] linear code over GF(8)

codes are known with d = 43, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-004331 | codetables-gf8-55-6

Minimum distance of a [55, 6] linear code over GF(8)

codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-004332 | codetables-gf8-56-5

Minimum distance of a [56, 5] linear code over GF(8)

codes are known with d = 44, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-004333 | codetables-gf8-56-6

Minimum distance of a [56, 6] linear code over GF(8)

codes are known with d = 43, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-004334 | codetables-gf8-57-5

Minimum distance of a [57, 5] linear code over GF(8)

codes are known with d = 45, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-004335 | codetables-gf8-57-6

Minimum distance of a [57, 6] linear code over GF(8)

codes are known with d = 44, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-004336 | codetables-gf8-58-5

Minimum distance of a [58, 5] linear code over GF(8)

codes are known with d = 46, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-004337 | codetables-gf8-58-6

Minimum distance of a [58, 6] linear code over GF(8)

codes are known with d = 44, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-004338 | codetables-gf8-59-5

Minimum distance of a [59, 5] linear code over GF(8)

codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-004339 | codetables-gf8-59-6

Minimum distance of a [59, 6] linear code over GF(8)

codes are known with d = 45, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-004340 | codetables-gf8-60-5

Minimum distance of a [60, 5] linear code over GF(8)

codes are known with d = 48, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-004341 | codetables-gf8-60-6

Minimum distance of a [60, 6] linear code over GF(8)

codes are known with d = 46, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-004342 | codetables-gf8-61-5

Minimum distance of a [61, 5] linear code over GF(8)

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-004343 | codetables-gf8-61-6

Minimum distance of a [61, 6] linear code over GF(8)

codes are known with d = 47, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-004344 | codetables-gf8-62-5

Minimum distance of a [62, 5] linear code over GF(8)

codes are known with d = 49, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-004345 | codetables-gf8-62-6

Minimum distance of a [62, 6] linear code over GF(8)

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-004346 | codetables-gf8-63-5

Minimum distance of a [63, 5] linear code over GF(8)

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-004347 | codetables-gf8-63-6

Minimum distance of a [63, 6] linear code over GF(8)

codes are known with d = 49, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-004348 | codetables-gf8-64-5

Minimum distance of a [64, 5] linear code over GF(8)

codes are known with d = 51, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-004349 | codetables-gf8-64-6

Minimum distance of a [64, 6] linear code over GF(8)

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-004350 | codetables-gf8-65-5

Minimum distance of a [65, 5] linear code over GF(8)

codes are known with d = 52, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-004351 | codetables-gf8-65-6

Minimum distance of a [65, 6] linear code over GF(8)

codes are known with d = 51, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-004352 | codetables-gf8-66-5

Minimum distance of a [66, 5] linear code over GF(8)

codes are known with d = 53, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-004353 | codetables-gf8-66-6

Minimum distance of a [66, 6] linear code over GF(8)

codes are known with d = 52, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-004354 | codetables-gf8-67-5

Minimum distance of a [67, 5] linear code over GF(8)

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-004355 | codetables-gf8-67-6

Minimum distance of a [67, 6] linear code over GF(8)

codes are known with d = 52, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-004356 | codetables-gf8-68-5

Minimum distance of a [68, 5] linear code over GF(8)

codes are known with d = 55, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-004357 | codetables-gf8-68-6

Minimum distance of a [68, 6] linear code over GF(8)

codes are known with d = 53, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-004358 | codetables-gf8-69-5

Minimum distance of a [69, 5] linear code over GF(8)

codes are known with d = 56, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-004359 | codetables-gf8-69-6

Minimum distance of a [69, 6] linear code over GF(8)

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-004360 | codetables-gf8-70-5

Minimum distance of a [70, 5] linear code over GF(8)

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-004361 | codetables-gf8-70-6

Minimum distance of a [70, 6] linear code over GF(8)

codes are known with d = 55, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-004362 | codetables-gf8-71-5

Minimum distance of a [71, 5] linear code over GF(8)

codes are known with d = 57, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-004363 | codetables-gf8-71-6

Minimum distance of a [71, 6] linear code over GF(8)

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-004364 | codetables-gf8-72-5

Minimum distance of a [72, 5] linear code over GF(8)

codes are known with d = 58, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-004365 | codetables-gf8-72-6

Minimum distance of a [72, 6] linear code over GF(8)

codes are known with d = 57, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-004366 | codetables-gf8-73-5

Minimum distance of a [73, 5] linear code over GF(8)

codes are known with d = 59, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-004367 | codetables-gf8-73-6

Minimum distance of a [73, 6] linear code over GF(8)

codes are known with d = 58, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-004368 | codetables-gf8-74-5

Minimum distance of a [74, 5] linear code over GF(8)

codes are known with d = 60, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-004369 | codetables-gf8-74-6

Minimum distance of a [74, 6] linear code over GF(8)

codes are known with d = 58, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-004370 | codetables-gf8-75-5

Minimum distance of a [75, 5] linear code over GF(8)

codes are known with d = 60, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-004371 | codetables-gf8-75-6

Minimum distance of a [75, 6] linear code over GF(8)

codes are known with d = 58, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-004372 | codetables-gf8-76-5

Minimum distance of a [76, 5] linear code over GF(8)

codes are known with d = 61, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-004373 | codetables-gf8-76-6

Minimum distance of a [76, 6] linear code over GF(8)

codes are known with d = 59, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-004374 | codetables-gf8-77-5

Minimum distance of a [77, 5] linear code over GF(8)

codes are known with d = 62, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-004375 | codetables-gf8-77-6

Minimum distance of a [77, 6] linear code over GF(8)

codes are known with d = 60, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-004376 | codetables-gf8-78-5

Minimum distance of a [78, 5] linear code over GF(8)

codes are known with d = 63, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-004377 | codetables-gf8-78-6

Minimum distance of a [78, 6] linear code over GF(8)

codes are known with d = 61, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-004378 | codetables-gf8-79-5

Minimum distance of a [79, 5] linear code over GF(8)

codes are known with d = 64, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-004379 | codetables-gf8-79-6

Minimum distance of a [79, 6] linear code over GF(8)

codes are known with d = 62, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-004380 | codetables-gf8-80-5

Minimum distance of a [80, 5] linear code over GF(8)

codes are known with d = 65, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-004381 | codetables-gf8-80-6

Minimum distance of a [80, 6] linear code over GF(8)

codes are known with d = 63, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-004382 | codetables-gf8-81-4

Minimum distance of a [81, 4] linear code over GF(8)

codes are known with d = 68, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-004383 | codetables-gf8-81-5

Minimum distance of a [81, 5] linear code over GF(8)

codes are known with d = 66, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-004384 | codetables-gf8-81-6

Minimum distance of a [81, 6] linear code over GF(8)

codes are known with d = 64, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-004385 | codetables-gf8-82-4

Minimum distance of a [82, 4] linear code over GF(8)

codes are known with d = 69, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-004386 | codetables-gf8-82-5

Minimum distance of a [82, 5] linear code over GF(8)

codes are known with d = 66, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-004387 | codetables-gf8-82-6

Minimum distance of a [82, 6] linear code over GF(8)

codes are known with d = 64, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-004388 | codetables-gf8-83-4

Minimum distance of a [83, 4] linear code over GF(8)

codes are known with d = 70, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-004389 | codetables-gf8-83-5

Minimum distance of a [83, 5] linear code over GF(8)

codes are known with d = 67, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-004390 | codetables-gf8-83-6

Minimum distance of a [83, 6] linear code over GF(8)

codes are known with d = 65, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-004391 | codetables-gf8-84-4

Minimum distance of a [84, 4] linear code over GF(8)

codes are known with d = 71, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-004392 | codetables-gf8-84-5

Minimum distance of a [84, 5] linear code over GF(8)

codes are known with d = 68, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-004393 | codetables-gf8-84-6

Minimum distance of a [84, 6] linear code over GF(8)

codes are known with d = 66, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-004394 | codetables-gf8-85-5

Minimum distance of a [85, 5] linear code over GF(8)

codes are known with d = 69, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-004395 | codetables-gf8-85-6

Minimum distance of a [85, 6] linear code over GF(8)

codes are known with d = 66, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-004396 | codetables-gf8-86-4

Minimum distance of a [86, 4] linear code over GF(8)

codes are known with d = 72, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-004397 | codetables-gf8-86-5

Minimum distance of a [86, 5] linear code over GF(8)

codes are known with d = 70, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-004398 | codetables-gf8-86-6

Minimum distance of a [86, 6] linear code over GF(8)

codes are known with d = 67, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-004399 | codetables-gf8-87-4

Minimum distance of a [87, 4] linear code over GF(8)

codes are known with d = 73, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-004400 | codetables-gf8-87-5

Minimum distance of a [87, 5] linear code over GF(8)

codes are known with d = 71, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-004401 | codetables-gf8-87-6

Minimum distance of a [87, 6] linear code over GF(8)

codes are known with d = 68, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-004402 | codetables-gf8-88-4

Minimum distance of a [88, 4] linear code over GF(8)

codes are known with d = 74, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-004403 | codetables-gf8-88-5

Minimum distance of a [88, 5] linear code over GF(8)

codes are known with d = 72, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-004404 | codetables-gf8-88-6

Minimum distance of a [88, 6] linear code over GF(8)

codes are known with d = 69, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-004405 | codetables-gf8-89-4

Minimum distance of a [89, 4] linear code over GF(8)

codes are known with d = 75, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-004406 | codetables-gf8-89-5

Minimum distance of a [89, 5] linear code over GF(8)

codes are known with d = 73, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-004407 | codetables-gf8-89-6

Minimum distance of a [89, 6] linear code over GF(8)

codes are known with d = 70, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-004408 | codetables-gf8-90-4

Minimum distance of a [90, 4] linear code over GF(8)

codes are known with d = 76, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-004409 | codetables-gf8-90-5

Minimum distance of a [90, 5] linear code over GF(8)

codes are known with d = 74, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-004410 | codetables-gf8-90-6

Minimum distance of a [90, 6] linear code over GF(8)

codes are known with d = 71, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-004411 | codetables-gf8-91-4

Minimum distance of a [91, 4] linear code over GF(8)

codes are known with d = 77, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-004412 | codetables-gf8-91-5

Minimum distance of a [91, 5] linear code over GF(8)

codes are known with d = 74, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-004413 | codetables-gf8-91-6

Minimum distance of a [91, 6] linear code over GF(8)

codes are known with d = 72, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-004414 | codetables-gf8-92-4

Minimum distance of a [92, 4] linear code over GF(8)

codes are known with d = 78, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-004415 | codetables-gf8-92-5

Minimum distance of a [92, 5] linear code over GF(8)

codes are known with d = 75, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-004416 | codetables-gf8-92-6

Minimum distance of a [92, 6] linear code over GF(8)

codes are known with d = 72, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-004417 | codetables-gf8-93-5

Minimum distance of a [93, 5] linear code over GF(8)

codes are known with d = 76, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-004418 | codetables-gf8-93-6

Minimum distance of a [93, 6] linear code over GF(8)

codes are known with d = 73, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-004419 | codetables-gf8-94-5

Minimum distance of a [94, 5] linear code over GF(8)

codes are known with d = 77, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-004420 | codetables-gf8-94-6

Minimum distance of a [94, 6] linear code over GF(8)

codes are known with d = 74, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-004421 | codetables-gf8-95-4

Minimum distance of a [95, 4] linear code over GF(8)

codes are known with d = 80, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-004422 | codetables-gf8-95-5

Minimum distance of a [95, 5] linear code over GF(8)

codes are known with d = 78, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-004423 | codetables-gf8-95-6

Minimum distance of a [95, 6] linear code over GF(8)

codes are known with d = 75, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-004424 | codetables-gf8-96-4

Minimum distance of a [96, 4] linear code over GF(8)

codes are known with d = 81, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-004425 | codetables-gf8-96-5

Minimum distance of a [96, 5] linear code over GF(8)

codes are known with d = 78, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-004426 | codetables-gf8-96-6

Minimum distance of a [96, 6] linear code over GF(8)

codes are known with d = 76, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-004427 | codetables-gf8-97-4

Minimum distance of a [97, 4] linear code over GF(8)

codes are known with d = 82, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-004428 | codetables-gf8-97-5

Minimum distance of a [97, 5] linear code over GF(8)

codes are known with d = 79, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-004429 | codetables-gf8-97-6

Minimum distance of a [97, 6] linear code over GF(8)

codes are known with d = 77, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-004430 | codetables-gf8-98-4

Minimum distance of a [98, 4] linear code over GF(8)

codes are known with d = 83, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-004431 | codetables-gf8-98-5

Minimum distance of a [98, 5] linear code over GF(8)

codes are known with d = 80, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-004432 | codetables-gf8-98-6

Minimum distance of a [98, 6] linear code over GF(8)

codes are known with d = 77, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-004433 | codetables-gf8-99-4

Minimum distance of a [99, 4] linear code over GF(8)

codes are known with d = 84, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-004434 | codetables-gf8-99-5

Minimum distance of a [99, 5] linear code over GF(8)

codes are known with d = 81, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-004435 | codetables-gf8-99-6

Minimum distance of a [99, 6] linear code over GF(8)

codes are known with d = 78, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-004436 | codetables-gf9-100-4

Minimum distance of a [100, 4] linear code over GF(9)

codes are known with d = 86, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-004437 | codetables-gf9-100-5

Minimum distance of a [100, 5] linear code over GF(9)

codes are known with d = 83, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-004438 | codetables-gf9-100-6

Minimum distance of a [100, 6] linear code over GF(9)

codes are known with d = 80, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-004439 | codetables-gf9-101-4

Minimum distance of a [101, 4] linear code over GF(9)

codes are known with d = 87, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-004440 | codetables-gf9-101-5

Minimum distance of a [101, 5] linear code over GF(9)

codes are known with d = 84, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-004441 | codetables-gf9-101-6

Minimum distance of a [101, 6] linear code over GF(9)

codes are known with d = 81, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-004442 | codetables-gf9-102-4

Minimum distance of a [102, 4] linear code over GF(9)

codes are known with d = 88, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-004443 | codetables-gf9-102-5

Minimum distance of a [102, 5] linear code over GF(9)

codes are known with d = 84, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-004444 | codetables-gf9-102-6

Minimum distance of a [102, 6] linear code over GF(9)

codes are known with d = 82, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-004445 | codetables-gf9-103-4

Minimum distance of a [103, 4] linear code over GF(9)

codes are known with d = 88, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-004446 | codetables-gf9-103-5

Minimum distance of a [103, 5] linear code over GF(9)

codes are known with d = 85, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-004447 | codetables-gf9-103-6

Minimum distance of a [103, 6] linear code over GF(9)

codes are known with d = 83, and no code can have d greater than 87.

来源状态 open | 2026-07-23 | 来源

V3P-004448 | codetables-gf9-104-4

Minimum distance of a [104, 4] linear code over GF(9)

codes are known with d = 89, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-004449 | codetables-gf9-104-5

Minimum distance of a [104, 5] linear code over GF(9)

codes are known with d = 86, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-004450 | codetables-gf9-104-6

Minimum distance of a [104, 6] linear code over GF(9)

codes are known with d = 84, and no code can have d greater than 88.

来源状态 open | 2026-07-23 | 来源

V3P-004451 | codetables-gf9-105-4

Minimum distance of a [105, 4] linear code over GF(9)

codes are known with d = 90, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-004452 | codetables-gf9-105-5

Minimum distance of a [105, 5] linear code over GF(9)

codes are known with d = 87, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-004453 | codetables-gf9-105-6

Minimum distance of a [105, 6] linear code over GF(9)

codes are known with d = 84, and no code can have d greater than 89.

来源状态 open | 2026-07-23 | 来源

V3P-004454 | codetables-gf9-106-4

Minimum distance of a [106, 4] linear code over GF(9)

codes are known with d = 91, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-004455 | codetables-gf9-106-5

Minimum distance of a [106, 5] linear code over GF(9)

codes are known with d = 88, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-004456 | codetables-gf9-106-6

Minimum distance of a [106, 6] linear code over GF(9)

codes are known with d = 85, and no code can have d greater than 90.

来源状态 open | 2026-07-23 | 来源

V3P-004457 | codetables-gf9-107-4

Minimum distance of a [107, 4] linear code over GF(9)

codes are known with d = 92, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-004458 | codetables-gf9-107-5

Minimum distance of a [107, 5] linear code over GF(9)

codes are known with d = 89, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-004459 | codetables-gf9-107-6

Minimum distance of a [107, 6] linear code over GF(9)

codes are known with d = 86, and no code can have d greater than 91.

来源状态 open | 2026-07-23 | 来源

V3P-004460 | codetables-gf9-108-4

Minimum distance of a [108, 4] linear code over GF(9)

codes are known with d = 93, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-004461 | codetables-gf9-108-5

Minimum distance of a [108, 5] linear code over GF(9)

codes are known with d = 90, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-004462 | codetables-gf9-108-6

Minimum distance of a [108, 6] linear code over GF(9)

codes are known with d = 87, and no code can have d greater than 92.

来源状态 open | 2026-07-23 | 来源

V3P-004463 | codetables-gf9-109-4

Minimum distance of a [109, 4] linear code over GF(9)

codes are known with d = 94, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-004464 | codetables-gf9-109-5

Minimum distance of a [109, 5] linear code over GF(9)

codes are known with d = 91, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-004465 | codetables-gf9-109-6

Minimum distance of a [109, 6] linear code over GF(9)

codes are known with d = 88, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-004466 | codetables-gf9-110-4

Minimum distance of a [110, 4] linear code over GF(9)

codes are known with d = 94, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-004467 | codetables-gf9-110-5

Minimum distance of a [110, 5] linear code over GF(9)

codes are known with d = 92, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-004468 | codetables-gf9-110-6

Minimum distance of a [110, 6] linear code over GF(9)

codes are known with d = 89, and no code can have d greater than 93.

来源状态 open | 2026-07-23 | 来源

V3P-004469 | codetables-gf9-111-4

Minimum distance of a [111, 4] linear code over GF(9)

codes are known with d = 95, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-004470 | codetables-gf9-111-5

Minimum distance of a [111, 5] linear code over GF(9)

codes are known with d = 92, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-004471 | codetables-gf9-111-6

Minimum distance of a [111, 6] linear code over GF(9)

codes are known with d = 90, and no code can have d greater than 94.

来源状态 open | 2026-07-23 | 来源

V3P-004472 | codetables-gf9-112-4

Minimum distance of a [112, 4] linear code over GF(9)

codes are known with d = 96, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-004473 | codetables-gf9-112-5

Minimum distance of a [112, 5] linear code over GF(9)

codes are known with d = 93, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-004474 | codetables-gf9-112-6

Minimum distance of a [112, 6] linear code over GF(9)

codes are known with d = 90, and no code can have d greater than 95.

来源状态 open | 2026-07-23 | 来源

V3P-004475 | codetables-gf9-113-4

Minimum distance of a [113, 4] linear code over GF(9)

codes are known with d = 97, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-004476 | codetables-gf9-113-5

Minimum distance of a [113, 5] linear code over GF(9)

codes are known with d = 94, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-004477 | codetables-gf9-113-6

Minimum distance of a [113, 6] linear code over GF(9)

codes are known with d = 91, and no code can have d greater than 96.

来源状态 open | 2026-07-23 | 来源

V3P-004478 | codetables-gf9-114-4

Minimum distance of a [114, 4] linear code over GF(9)

codes are known with d = 98, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-004479 | codetables-gf9-114-5

Minimum distance of a [114, 5] linear code over GF(9)

codes are known with d = 95, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-004480 | codetables-gf9-114-6

Minimum distance of a [114, 6] linear code over GF(9)

codes are known with d = 92, and no code can have d greater than 97.

来源状态 open | 2026-07-23 | 来源

V3P-004481 | codetables-gf9-115-4

Minimum distance of a [115, 4] linear code over GF(9)

codes are known with d = 99, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-004482 | codetables-gf9-115-5

Minimum distance of a [115, 5] linear code over GF(9)

codes are known with d = 96, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-004483 | codetables-gf9-115-6

Minimum distance of a [115, 6] linear code over GF(9)

codes are known with d = 93, and no code can have d greater than 98.

来源状态 open | 2026-07-23 | 来源

V3P-004484 | codetables-gf9-116-4

Minimum distance of a [116, 4] linear code over GF(9)

codes are known with d = 100, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-004485 | codetables-gf9-116-5

Minimum distance of a [116, 5] linear code over GF(9)

codes are known with d = 97, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-004486 | codetables-gf9-116-6

Minimum distance of a [116, 6] linear code over GF(9)

codes are known with d = 94, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-004487 | codetables-gf9-117-4

Minimum distance of a [117, 4] linear code over GF(9)

codes are known with d = 101, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-004488 | codetables-gf9-117-5

Minimum distance of a [117, 5] linear code over GF(9)

codes are known with d = 98, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-004489 | codetables-gf9-117-6

Minimum distance of a [117, 6] linear code over GF(9)

codes are known with d = 95, and no code can have d greater than 99.

来源状态 open | 2026-07-23 | 来源

V3P-004490 | codetables-gf9-118-4

Minimum distance of a [118, 4] linear code over GF(9)

codes are known with d = 102, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-004491 | codetables-gf9-118-5

Minimum distance of a [118, 5] linear code over GF(9)

codes are known with d = 98, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-004492 | codetables-gf9-118-6

Minimum distance of a [118, 6] linear code over GF(9)

codes are known with d = 95, and no code can have d greater than 100.

来源状态 open | 2026-07-23 | 来源

V3P-004493 | codetables-gf9-119-4

Minimum distance of a [119, 4] linear code over GF(9)

codes are known with d = 102, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-004494 | codetables-gf9-119-5

Minimum distance of a [119, 5] linear code over GF(9)

codes are known with d = 99, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-004495 | codetables-gf9-119-6

Minimum distance of a [119, 6] linear code over GF(9)

codes are known with d = 96, and no code can have d greater than 101.

来源状态 open | 2026-07-23 | 来源

V3P-004496 | codetables-gf9-120-4

Minimum distance of a [120, 4] linear code over GF(9)

codes are known with d = 103, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-004497 | codetables-gf9-120-5

Minimum distance of a [120, 5] linear code over GF(9)

codes are known with d = 100, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-004498 | codetables-gf9-120-6

Minimum distance of a [120, 6] linear code over GF(9)

codes are known with d = 97, and no code can have d greater than 102.

来源状态 open | 2026-07-23 | 来源

V3P-004499 | codetables-gf9-121-4

Minimum distance of a [121, 4] linear code over GF(9)

codes are known with d = 104, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-004500 | codetables-gf9-121-5

Minimum distance of a [121, 5] linear code over GF(9)

codes are known with d = 101, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-004501 | codetables-gf9-121-6

Minimum distance of a [121, 6] linear code over GF(9)

codes are known with d = 98, and no code can have d greater than 103.

来源状态 open | 2026-07-23 | 来源

V3P-004502 | codetables-gf9-122-4

Minimum distance of a [122, 4] linear code over GF(9)

codes are known with d = 105, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-004503 | codetables-gf9-122-5

Minimum distance of a [122, 5] linear code over GF(9)

codes are known with d = 102, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-004504 | codetables-gf9-122-6

Minimum distance of a [122, 6] linear code over GF(9)

codes are known with d = 99, and no code can have d greater than 104.

来源状态 open | 2026-07-23 | 来源

V3P-004505 | codetables-gf9-123-4

Minimum distance of a [123, 4] linear code over GF(9)

codes are known with d = 106, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-004506 | codetables-gf9-123-5

Minimum distance of a [123, 5] linear code over GF(9)

codes are known with d = 103, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-004507 | codetables-gf9-123-6

Minimum distance of a [123, 6] linear code over GF(9)

codes are known with d = 100, and no code can have d greater than 105.

来源状态 open | 2026-07-23 | 来源

V3P-004508 | codetables-gf9-124-4

Minimum distance of a [124, 4] linear code over GF(9)

codes are known with d = 107, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-004509 | codetables-gf9-124-5

Minimum distance of a [124, 5] linear code over GF(9)

codes are known with d = 104, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-004510 | codetables-gf9-124-6

Minimum distance of a [124, 6] linear code over GF(9)

codes are known with d = 101, and no code can have d greater than 106.

来源状态 open | 2026-07-23 | 来源

V3P-004511 | codetables-gf9-125-4

Minimum distance of a [125, 4] linear code over GF(9)

codes are known with d = 108, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-004512 | codetables-gf9-125-5

Minimum distance of a [125, 5] linear code over GF(9)

codes are known with d = 105, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-004513 | codetables-gf9-125-6

Minimum distance of a [125, 6] linear code over GF(9)

codes are known with d = 102, and no code can have d greater than 107.

来源状态 open | 2026-07-23 | 来源

V3P-004514 | codetables-gf9-126-4

Minimum distance of a [126, 4] linear code over GF(9)

codes are known with d = 108, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-004515 | codetables-gf9-126-5

Minimum distance of a [126, 5] linear code over GF(9)

codes are known with d = 106, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-004516 | codetables-gf9-126-6

Minimum distance of a [126, 6] linear code over GF(9)

codes are known with d = 102, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-004517 | codetables-gf9-127-4

Minimum distance of a [127, 4] linear code over GF(9)

codes are known with d = 109, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-004518 | codetables-gf9-127-5

Minimum distance of a [127, 5] linear code over GF(9)

codes are known with d = 107, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-004519 | codetables-gf9-127-6

Minimum distance of a [127, 6] linear code over GF(9)

codes are known with d = 103, and no code can have d greater than 108.

来源状态 open | 2026-07-23 | 来源

V3P-004520 | codetables-gf9-128-4

Minimum distance of a [128, 4] linear code over GF(9)

codes are known with d = 110, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-004521 | codetables-gf9-128-5

Minimum distance of a [128, 5] linear code over GF(9)

codes are known with d = 107, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-004522 | codetables-gf9-128-6

Minimum distance of a [128, 6] linear code over GF(9)

codes are known with d = 104, and no code can have d greater than 109.

来源状态 open | 2026-07-23 | 来源

V3P-004523 | codetables-gf9-129-4

Minimum distance of a [129, 4] linear code over GF(9)

codes are known with d = 111, and no code can have d greater than 113.

来源状态 open | 2026-07-23 | 来源

V3P-004524 | codetables-gf9-129-5

Minimum distance of a [129, 5] linear code over GF(9)

codes are known with d = 108, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-004525 | codetables-gf9-129-6

Minimum distance of a [129, 6] linear code over GF(9)

codes are known with d = 105, and no code can have d greater than 110.

来源状态 open | 2026-07-23 | 来源

V3P-004526 | codetables-gf9-130-4

Minimum distance of a [130, 4] linear code over GF(9)

codes are known with d = 112, and no code can have d greater than 114.

来源状态 open | 2026-07-23 | 来源

V3P-004527 | codetables-gf9-130-5

Minimum distance of a [130, 5] linear code over GF(9)

codes are known with d = 109, and no code can have d greater than 112.

来源状态 open | 2026-07-23 | 来源

V3P-004528 | codetables-gf9-130-6

Minimum distance of a [130, 6] linear code over GF(9)

codes are known with d = 106, and no code can have d greater than 111.

来源状态 open | 2026-07-23 | 来源

V3P-004529 | codetables-gf9-18-3

Minimum distance of a [18, 3] linear code over GF(9)

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-004530 | codetables-gf9-19-3

Minimum distance of a [19, 3] linear code over GF(9)

codes are known with d = 15, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-004531 | codetables-gf9-20-3

Minimum distance of a [20, 3] linear code over GF(9)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-004532 | codetables-gf9-21-3

Minimum distance of a [21, 3] linear code over GF(9)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-004533 | codetables-gf9-22-5

Minimum distance of a [22, 5] linear code over GF(9)

codes are known with d = 15, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-004534 | codetables-gf9-22-6

Minimum distance of a [22, 6] linear code over GF(9)

codes are known with d = 14, and no code can have d greater than 15.

来源状态 open | 2026-07-23 | 来源

V3P-004535 | codetables-gf9-23-5

Minimum distance of a [23, 5] linear code over GF(9)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-004536 | codetables-gf9-23-6

Minimum distance of a [23, 6] linear code over GF(9)

codes are known with d = 15, and no code can have d greater than 16.

来源状态 open | 2026-07-23 | 来源

V3P-004537 | codetables-gf9-24-5

Minimum distance of a [24, 5] linear code over GF(9)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-004538 | codetables-gf9-24-6

Minimum distance of a [24, 6] linear code over GF(9)

codes are known with d = 16, and no code can have d greater than 17.

来源状态 open | 2026-07-23 | 来源

V3P-004539 | codetables-gf9-25-4

Minimum distance of a [25, 4] linear code over GF(9)

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-004540 | codetables-gf9-25-5

Minimum distance of a [25, 5] linear code over GF(9)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-004541 | codetables-gf9-25-6

Minimum distance of a [25, 6] linear code over GF(9)

codes are known with d = 17, and no code can have d greater than 18.

来源状态 open | 2026-07-23 | 来源

V3P-004542 | codetables-gf9-26-4

Minimum distance of a [26, 4] linear code over GF(9)

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-004543 | codetables-gf9-26-5

Minimum distance of a [26, 5] linear code over GF(9)

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-004544 | codetables-gf9-26-6

Minimum distance of a [26, 6] linear code over GF(9)

codes are known with d = 18, and no code can have d greater than 19.

来源状态 open | 2026-07-23 | 来源

V3P-004545 | codetables-gf9-27-4

Minimum distance of a [27, 4] linear code over GF(9)

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-004546 | codetables-gf9-27-5

Minimum distance of a [27, 5] linear code over GF(9)

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-004547 | codetables-gf9-27-6

Minimum distance of a [27, 6] linear code over GF(9)

codes are known with d = 19, and no code can have d greater than 20.

来源状态 open | 2026-07-23 | 来源

V3P-004548 | codetables-gf9-28-4

Minimum distance of a [28, 4] linear code over GF(9)

codes are known with d = 22, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-004549 | codetables-gf9-28-5

Minimum distance of a [28, 5] linear code over GF(9)

codes are known with d = 21, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-004550 | codetables-gf9-28-6

Minimum distance of a [28, 6] linear code over GF(9)

codes are known with d = 20, and no code can have d greater than 21.

来源状态 open | 2026-07-23 | 来源

V3P-004551 | codetables-gf9-29-3

Minimum distance of a [29, 3] linear code over GF(9)

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-004552 | codetables-gf9-29-4

Minimum distance of a [29, 4] linear code over GF(9)

codes are known with d = 23, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-004553 | codetables-gf9-29-5

Minimum distance of a [29, 5] linear code over GF(9)

codes are known with d = 21, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-004554 | codetables-gf9-29-6

Minimum distance of a [29, 6] linear code over GF(9)

codes are known with d = 20, and no code can have d greater than 22.

来源状态 open | 2026-07-23 | 来源

V3P-004555 | codetables-gf9-30-3

Minimum distance of a [30, 3] linear code over GF(9)

codes are known with d = 25, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-004556 | codetables-gf9-30-4

Minimum distance of a [30, 4] linear code over GF(9)

codes are known with d = 24, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-004557 | codetables-gf9-30-5

Minimum distance of a [30, 5] linear code over GF(9)

codes are known with d = 22, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-004558 | codetables-gf9-30-6

Minimum distance of a [30, 6] linear code over GF(9)

codes are known with d = 21, and no code can have d greater than 23.

来源状态 open | 2026-07-23 | 来源

V3P-004559 | codetables-gf9-31-3

Minimum distance of a [31, 3] linear code over GF(9)

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-004560 | codetables-gf9-31-4

Minimum distance of a [31, 4] linear code over GF(9)

codes are known with d = 25, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-004561 | codetables-gf9-31-5

Minimum distance of a [31, 5] linear code over GF(9)

codes are known with d = 23, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-004562 | codetables-gf9-31-6

Minimum distance of a [31, 6] linear code over GF(9)

codes are known with d = 21, and no code can have d greater than 24.

来源状态 open | 2026-07-23 | 来源

V3P-004563 | codetables-gf9-32-4

Minimum distance of a [32, 4] linear code over GF(9)

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-004564 | codetables-gf9-32-5

Minimum distance of a [32, 5] linear code over GF(9)

codes are known with d = 24, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-004565 | codetables-gf9-32-6

Minimum distance of a [32, 6] linear code over GF(9)

codes are known with d = 22, and no code can have d greater than 25.

来源状态 open | 2026-07-23 | 来源

V3P-004566 | codetables-gf9-33-5

Minimum distance of a [33, 5] linear code over GF(9)

codes are known with d = 25, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-004567 | codetables-gf9-33-6

Minimum distance of a [33, 6] linear code over GF(9)

codes are known with d = 23, and no code can have d greater than 26.

来源状态 open | 2026-07-23 | 来源

V3P-004568 | codetables-gf9-34-5

Minimum distance of a [34, 5] linear code over GF(9)

codes are known with d = 26, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-004569 | codetables-gf9-34-6

Minimum distance of a [34, 6] linear code over GF(9)

codes are known with d = 24, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-004570 | codetables-gf9-35-5

Minimum distance of a [35, 5] linear code over GF(9)

codes are known with d = 26, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-004571 | codetables-gf9-35-6

Minimum distance of a [35, 6] linear code over GF(9)

codes are known with d = 25, and no code can have d greater than 27.

来源状态 open | 2026-07-23 | 来源

V3P-004572 | codetables-gf9-36-5

Minimum distance of a [36, 5] linear code over GF(9)

codes are known with d = 27, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-004573 | codetables-gf9-36-6

Minimum distance of a [36, 6] linear code over GF(9)

codes are known with d = 26, and no code can have d greater than 28.

来源状态 open | 2026-07-23 | 来源

V3P-004574 | codetables-gf9-37-4

Minimum distance of a [37, 4] linear code over GF(9)

codes are known with d = 30, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-004575 | codetables-gf9-37-5

Minimum distance of a [37, 5] linear code over GF(9)

codes are known with d = 28, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-004576 | codetables-gf9-37-6

Minimum distance of a [37, 6] linear code over GF(9)

codes are known with d = 27, and no code can have d greater than 29.

来源状态 open | 2026-07-23 | 来源

V3P-004577 | codetables-gf9-38-4

Minimum distance of a [38, 4] linear code over GF(9)

codes are known with d = 31, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-004578 | codetables-gf9-38-5

Minimum distance of a [38, 5] linear code over GF(9)

codes are known with d = 29, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-004579 | codetables-gf9-38-6

Minimum distance of a [38, 6] linear code over GF(9)

codes are known with d = 27, and no code can have d greater than 30.

来源状态 open | 2026-07-23 | 来源

V3P-004580 | codetables-gf9-39-5

Minimum distance of a [39, 5] linear code over GF(9)

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-004581 | codetables-gf9-39-6

Minimum distance of a [39, 6] linear code over GF(9)

codes are known with d = 28, and no code can have d greater than 31.

来源状态 open | 2026-07-23 | 来源

V3P-004582 | codetables-gf9-40-5

Minimum distance of a [40, 5] linear code over GF(9)

codes are known with d = 31, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-004583 | codetables-gf9-40-6

Minimum distance of a [40, 6] linear code over GF(9)

codes are known with d = 29, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-004584 | codetables-gf9-41-5

Minimum distance of a [41, 5] linear code over GF(9)

codes are known with d = 32, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-004585 | codetables-gf9-41-6

Minimum distance of a [41, 6] linear code over GF(9)

codes are known with d = 30, and no code can have d greater than 32.

来源状态 open | 2026-07-23 | 来源

V3P-004586 | codetables-gf9-42-4

Minimum distance of a [42, 4] linear code over GF(9)

codes are known with d = 34, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-004587 | codetables-gf9-42-5

Minimum distance of a [42, 5] linear code over GF(9)

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-004588 | codetables-gf9-42-6

Minimum distance of a [42, 6] linear code over GF(9)

codes are known with d = 31, and no code can have d greater than 33.

来源状态 open | 2026-07-23 | 来源

V3P-004589 | codetables-gf9-43-4

Minimum distance of a [43, 4] linear code over GF(9)

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-004590 | codetables-gf9-43-5

Minimum distance of a [43, 5] linear code over GF(9)

codes are known with d = 33, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-004591 | codetables-gf9-43-6

Minimum distance of a [43, 6] linear code over GF(9)

codes are known with d = 32, and no code can have d greater than 34.

来源状态 open | 2026-07-23 | 来源

V3P-004592 | codetables-gf9-44-4

Minimum distance of a [44, 4] linear code over GF(9)

codes are known with d = 36, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-004593 | codetables-gf9-44-5

Minimum distance of a [44, 5] linear code over GF(9)

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-004594 | codetables-gf9-44-6

Minimum distance of a [44, 6] linear code over GF(9)

codes are known with d = 32, and no code can have d greater than 35.

来源状态 open | 2026-07-23 | 来源

V3P-004595 | codetables-gf9-45-4

Minimum distance of a [45, 4] linear code over GF(9)

codes are known with d = 37, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-004596 | codetables-gf9-45-5

Minimum distance of a [45, 5] linear code over GF(9)

codes are known with d = 35, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-004597 | codetables-gf9-45-6

Minimum distance of a [45, 6] linear code over GF(9)

codes are known with d = 33, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-004598 | codetables-gf9-46-4

Minimum distance of a [46, 4] linear code over GF(9)

codes are known with d = 38, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-004599 | codetables-gf9-46-5

Minimum distance of a [46, 5] linear code over GF(9)

codes are known with d = 36, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-004600 | codetables-gf9-46-6

Minimum distance of a [46, 6] linear code over GF(9)

codes are known with d = 34, and no code can have d greater than 36.

来源状态 open | 2026-07-23 | 来源

V3P-004601 | codetables-gf9-47-4

Minimum distance of a [47, 4] linear code over GF(9)

codes are known with d = 39, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-004602 | codetables-gf9-47-5

Minimum distance of a [47, 5] linear code over GF(9)

codes are known with d = 37, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-004603 | codetables-gf9-47-6

Minimum distance of a [47, 6] linear code over GF(9)

codes are known with d = 35, and no code can have d greater than 37.

来源状态 open | 2026-07-23 | 来源

V3P-004604 | codetables-gf9-48-4

Minimum distance of a [48, 4] linear code over GF(9)

codes are known with d = 40, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-004605 | codetables-gf9-48-5

Minimum distance of a [48, 5] linear code over GF(9)

codes are known with d = 38, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-004606 | codetables-gf9-48-6

Minimum distance of a [48, 6] linear code over GF(9)

codes are known with d = 36, and no code can have d greater than 38.

来源状态 open | 2026-07-23 | 来源

V3P-004607 | codetables-gf9-49-4

Minimum distance of a [49, 4] linear code over GF(9)

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-004608 | codetables-gf9-49-5

Minimum distance of a [49, 5] linear code over GF(9)

codes are known with d = 38, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-004609 | codetables-gf9-49-6

Minimum distance of a [49, 6] linear code over GF(9)

codes are known with d = 37, and no code can have d greater than 39.

来源状态 open | 2026-07-23 | 来源

V3P-004610 | codetables-gf9-50-4

Minimum distance of a [50, 4] linear code over GF(9)

codes are known with d = 41, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-004611 | codetables-gf9-50-5

Minimum distance of a [50, 5] linear code over GF(9)

codes are known with d = 39, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-004612 | codetables-gf9-50-6

Minimum distance of a [50, 6] linear code over GF(9)

codes are known with d = 37, and no code can have d greater than 40.

来源状态 open | 2026-07-23 | 来源

V3P-004613 | codetables-gf9-51-4

Minimum distance of a [51, 4] linear code over GF(9)

codes are known with d = 42, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-004614 | codetables-gf9-51-5

Minimum distance of a [51, 5] linear code over GF(9)

codes are known with d = 40, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-004615 | codetables-gf9-51-6

Minimum distance of a [51, 6] linear code over GF(9)

codes are known with d = 38, and no code can have d greater than 41.

来源状态 open | 2026-07-23 | 来源

V3P-004616 | codetables-gf9-52-4

Minimum distance of a [52, 4] linear code over GF(9)

codes are known with d = 43, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-004617 | codetables-gf9-52-5

Minimum distance of a [52, 5] linear code over GF(9)

codes are known with d = 41, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-004618 | codetables-gf9-52-6

Minimum distance of a [52, 6] linear code over GF(9)

codes are known with d = 39, and no code can have d greater than 42.

来源状态 open | 2026-07-23 | 来源

V3P-004619 | codetables-gf9-53-4

Minimum distance of a [53, 4] linear code over GF(9)

codes are known with d = 44, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-004620 | codetables-gf9-53-5

Minimum distance of a [53, 5] linear code over GF(9)

codes are known with d = 42, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-004621 | codetables-gf9-53-6

Minimum distance of a [53, 6] linear code over GF(9)

codes are known with d = 40, and no code can have d greater than 43.

来源状态 open | 2026-07-23 | 来源

V3P-004622 | codetables-gf9-54-4

Minimum distance of a [54, 4] linear code over GF(9)

codes are known with d = 45, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-004623 | codetables-gf9-54-5

Minimum distance of a [54, 5] linear code over GF(9)

codes are known with d = 43, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-004624 | codetables-gf9-54-6

Minimum distance of a [54, 6] linear code over GF(9)

codes are known with d = 41, and no code can have d greater than 44.

来源状态 open | 2026-07-23 | 来源

V3P-004625 | codetables-gf9-55-4

Minimum distance of a [55, 4] linear code over GF(9)

codes are known with d = 46, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-004626 | codetables-gf9-55-5

Minimum distance of a [55, 5] linear code over GF(9)

codes are known with d = 44, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-004627 | codetables-gf9-55-6

Minimum distance of a [55, 6] linear code over GF(9)

codes are known with d = 42, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-004628 | codetables-gf9-56-3

Minimum distance of a [56, 3] linear code over GF(9)

codes are known with d = 48, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-004629 | codetables-gf9-56-4

Minimum distance of a [56, 4] linear code over GF(9)

codes are known with d = 47, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-004630 | codetables-gf9-56-5

Minimum distance of a [56, 5] linear code over GF(9)

codes are known with d = 45, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-004631 | codetables-gf9-56-6

Minimum distance of a [56, 6] linear code over GF(9)

codes are known with d = 43, and no code can have d greater than 45.

来源状态 open | 2026-07-23 | 来源

V3P-004632 | codetables-gf9-57-3

Minimum distance of a [57, 3] linear code over GF(9)

codes are known with d = 49, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-004633 | codetables-gf9-57-4

Minimum distance of a [57, 4] linear code over GF(9)

codes are known with d = 48, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-004634 | codetables-gf9-57-5

Minimum distance of a [57, 5] linear code over GF(9)

codes are known with d = 45, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-004635 | codetables-gf9-57-6

Minimum distance of a [57, 6] linear code over GF(9)

codes are known with d = 43, and no code can have d greater than 46.

来源状态 open | 2026-07-23 | 来源

V3P-004636 | codetables-gf9-58-3

Minimum distance of a [58, 3] linear code over GF(9)

codes are known with d = 50, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-004637 | codetables-gf9-58-4

Minimum distance of a [58, 4] linear code over GF(9)

codes are known with d = 49, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-004638 | codetables-gf9-58-5

Minimum distance of a [58, 5] linear code over GF(9)

codes are known with d = 46, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-004639 | codetables-gf9-58-6

Minimum distance of a [58, 6] linear code over GF(9)

codes are known with d = 44, and no code can have d greater than 47.

来源状态 open | 2026-07-23 | 来源

V3P-004640 | codetables-gf9-59-3

Minimum distance of a [59, 3] linear code over GF(9)

codes are known with d = 51, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-004641 | codetables-gf9-59-4

Minimum distance of a [59, 4] linear code over GF(9)

codes are known with d = 49, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-004642 | codetables-gf9-59-5

Minimum distance of a [59, 5] linear code over GF(9)

codes are known with d = 47, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-004643 | codetables-gf9-59-6

Minimum distance of a [59, 6] linear code over GF(9)

codes are known with d = 45, and no code can have d greater than 48.

来源状态 open | 2026-07-23 | 来源

V3P-004644 | codetables-gf9-60-3

Minimum distance of a [60, 3] linear code over GF(9)

codes are known with d = 52, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-004645 | codetables-gf9-60-4

Minimum distance of a [60, 4] linear code over GF(9)

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-004646 | codetables-gf9-60-5

Minimum distance of a [60, 5] linear code over GF(9)

codes are known with d = 48, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-004647 | codetables-gf9-60-6

Minimum distance of a [60, 6] linear code over GF(9)

codes are known with d = 46, and no code can have d greater than 49.

来源状态 open | 2026-07-23 | 来源

V3P-004648 | codetables-gf9-61-3

Minimum distance of a [61, 3] linear code over GF(9)

codes are known with d = 53, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-004649 | codetables-gf9-61-4

Minimum distance of a [61, 4] linear code over GF(9)

codes are known with d = 51, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-004650 | codetables-gf9-61-5

Minimum distance of a [61, 5] linear code over GF(9)

codes are known with d = 49, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-004651 | codetables-gf9-61-6

Minimum distance of a [61, 6] linear code over GF(9)

codes are known with d = 47, and no code can have d greater than 50.

来源状态 open | 2026-07-23 | 来源

V3P-004652 | codetables-gf9-62-4

Minimum distance of a [62, 4] linear code over GF(9)

codes are known with d = 52, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-004653 | codetables-gf9-62-5

Minimum distance of a [62, 5] linear code over GF(9)

codes are known with d = 50, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-004654 | codetables-gf9-62-6

Minimum distance of a [62, 6] linear code over GF(9)

codes are known with d = 48, and no code can have d greater than 51.

来源状态 open | 2026-07-23 | 来源

V3P-004655 | codetables-gf9-63-4

Minimum distance of a [63, 4] linear code over GF(9)

codes are known with d = 53, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-004656 | codetables-gf9-63-5

Minimum distance of a [63, 5] linear code over GF(9)

codes are known with d = 51, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-004657 | codetables-gf9-63-6

Minimum distance of a [63, 6] linear code over GF(9)

codes are known with d = 49, and no code can have d greater than 52.

来源状态 open | 2026-07-23 | 来源

V3P-004658 | codetables-gf9-64-4

Minimum distance of a [64, 4] linear code over GF(9)

codes are known with d = 54, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-004659 | codetables-gf9-64-5

Minimum distance of a [64, 5] linear code over GF(9)

codes are known with d = 51, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-004660 | codetables-gf9-64-6

Minimum distance of a [64, 6] linear code over GF(9)

codes are known with d = 49, and no code can have d greater than 53.

来源状态 open | 2026-07-23 | 来源

V3P-004661 | codetables-gf9-65-4

Minimum distance of a [65, 4] linear code over GF(9)

codes are known with d = 55, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-004662 | codetables-gf9-65-5

Minimum distance of a [65, 5] linear code over GF(9)

codes are known with d = 52, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-004663 | codetables-gf9-65-6

Minimum distance of a [65, 6] linear code over GF(9)

codes are known with d = 50, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-004664 | codetables-gf9-66-3

Minimum distance of a [66, 3] linear code over GF(9)

codes are known with d = 57, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-004665 | codetables-gf9-66-4

Minimum distance of a [66, 4] linear code over GF(9)

codes are known with d = 56, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-004666 | codetables-gf9-66-5

Minimum distance of a [66, 5] linear code over GF(9)

codes are known with d = 53, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-004667 | codetables-gf9-66-6

Minimum distance of a [66, 6] linear code over GF(9)

codes are known with d = 51, and no code can have d greater than 54.

来源状态 open | 2026-07-23 | 来源

V3P-004668 | codetables-gf9-67-3

Minimum distance of a [67, 3] linear code over GF(9)

codes are known with d = 58, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-004669 | codetables-gf9-67-4

Minimum distance of a [67, 4] linear code over GF(9)

codes are known with d = 57, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-004670 | codetables-gf9-67-5

Minimum distance of a [67, 5] linear code over GF(9)

codes are known with d = 54, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-004671 | codetables-gf9-67-6

Minimum distance of a [67, 6] linear code over GF(9)

codes are known with d = 52, and no code can have d greater than 55.

来源状态 open | 2026-07-23 | 来源

V3P-004672 | codetables-gf9-68-3

Minimum distance of a [68, 3] linear code over GF(9)

codes are known with d = 59, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-004673 | codetables-gf9-68-4

Minimum distance of a [68, 4] linear code over GF(9)

codes are known with d = 58, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-004674 | codetables-gf9-68-5

Minimum distance of a [68, 5] linear code over GF(9)

codes are known with d = 55, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-004675 | codetables-gf9-68-6

Minimum distance of a [68, 6] linear code over GF(9)

codes are known with d = 53, and no code can have d greater than 56.

来源状态 open | 2026-07-23 | 来源

V3P-004676 | codetables-gf9-69-3

Minimum distance of a [69, 3] linear code over GF(9)

codes are known with d = 60, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-004677 | codetables-gf9-69-4

Minimum distance of a [69, 4] linear code over GF(9)

codes are known with d = 59, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-004678 | codetables-gf9-69-5

Minimum distance of a [69, 5] linear code over GF(9)

codes are known with d = 56, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-004679 | codetables-gf9-69-6

Minimum distance of a [69, 6] linear code over GF(9)

codes are known with d = 54, and no code can have d greater than 57.

来源状态 open | 2026-07-23 | 来源

V3P-004680 | codetables-gf9-70-3

Minimum distance of a [70, 3] linear code over GF(9)

codes are known with d = 61, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-004681 | codetables-gf9-70-4

Minimum distance of a [70, 4] linear code over GF(9)

codes are known with d = 60, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-004682 | codetables-gf9-70-5

Minimum distance of a [70, 5] linear code over GF(9)

codes are known with d = 57, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-004683 | codetables-gf9-70-6

Minimum distance of a [70, 6] linear code over GF(9)

codes are known with d = 55, and no code can have d greater than 58.

来源状态 open | 2026-07-23 | 来源

V3P-004684 | codetables-gf9-71-3

Minimum distance of a [71, 3] linear code over GF(9)

codes are known with d = 62, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-004685 | codetables-gf9-71-4

Minimum distance of a [71, 4] linear code over GF(9)

codes are known with d = 61, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-004686 | codetables-gf9-71-5

Minimum distance of a [71, 5] linear code over GF(9)

codes are known with d = 58, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-004687 | codetables-gf9-71-6

Minimum distance of a [71, 6] linear code over GF(9)

codes are known with d = 56, and no code can have d greater than 59.

来源状态 open | 2026-07-23 | 来源

V3P-004688 | codetables-gf9-72-4

Minimum distance of a [72, 4] linear code over GF(9)

codes are known with d = 62, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-004689 | codetables-gf9-72-5

Minimum distance of a [72, 5] linear code over GF(9)

codes are known with d = 58, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-004690 | codetables-gf9-72-6

Minimum distance of a [72, 6] linear code over GF(9)

codes are known with d = 57, and no code can have d greater than 60.

来源状态 open | 2026-07-23 | 来源

V3P-004691 | codetables-gf9-73-5

Minimum distance of a [73, 5] linear code over GF(9)

codes are known with d = 59, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-004692 | codetables-gf9-73-6

Minimum distance of a [73, 6] linear code over GF(9)

codes are known with d = 57, and no code can have d greater than 61.

来源状态 open | 2026-07-23 | 来源

V3P-004693 | codetables-gf9-74-5

Minimum distance of a [74, 5] linear code over GF(9)

codes are known with d = 60, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-004694 | codetables-gf9-74-6

Minimum distance of a [74, 6] linear code over GF(9)

codes are known with d = 58, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-004695 | codetables-gf9-75-5

Minimum distance of a [75, 5] linear code over GF(9)

codes are known with d = 61, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-004696 | codetables-gf9-75-6

Minimum distance of a [75, 6] linear code over GF(9)

codes are known with d = 59, and no code can have d greater than 62.

来源状态 open | 2026-07-23 | 来源

V3P-004697 | codetables-gf9-76-5

Minimum distance of a [76, 5] linear code over GF(9)

codes are known with d = 62, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-004698 | codetables-gf9-76-6

Minimum distance of a [76, 6] linear code over GF(9)

codes are known with d = 60, and no code can have d greater than 63.

来源状态 open | 2026-07-23 | 来源

V3P-004699 | codetables-gf9-77-5

Minimum distance of a [77, 5] linear code over GF(9)

codes are known with d = 63, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-004700 | codetables-gf9-77-6

Minimum distance of a [77, 6] linear code over GF(9)

codes are known with d = 61, and no code can have d greater than 64.

来源状态 open | 2026-07-23 | 来源

V3P-004701 | codetables-gf9-78-5

Minimum distance of a [78, 5] linear code over GF(9)

codes are known with d = 64, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-004702 | codetables-gf9-78-6

Minimum distance of a [78, 6] linear code over GF(9)

codes are known with d = 62, and no code can have d greater than 65.

来源状态 open | 2026-07-23 | 来源

V3P-004703 | codetables-gf9-79-5

Minimum distance of a [79, 5] linear code over GF(9)

codes are known with d = 64, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-004704 | codetables-gf9-79-6

Minimum distance of a [79, 6] linear code over GF(9)

codes are known with d = 63, and no code can have d greater than 66.

来源状态 open | 2026-07-23 | 来源

V3P-004705 | codetables-gf9-80-5

Minimum distance of a [80, 5] linear code over GF(9)

codes are known with d = 65, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-004706 | codetables-gf9-80-6

Minimum distance of a [80, 6] linear code over GF(9)

codes are known with d = 64, and no code can have d greater than 67.

来源状态 open | 2026-07-23 | 来源

V3P-004707 | codetables-gf9-81-5

Minimum distance of a [81, 5] linear code over GF(9)

codes are known with d = 66, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-004708 | codetables-gf9-81-6

Minimum distance of a [81, 6] linear code over GF(9)

codes are known with d = 65, and no code can have d greater than 68.

来源状态 open | 2026-07-23 | 来源

V3P-004709 | codetables-gf9-82-5

Minimum distance of a [82, 5] linear code over GF(9)

codes are known with d = 67, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-004710 | codetables-gf9-82-6

Minimum distance of a [82, 6] linear code over GF(9)

codes are known with d = 66, and no code can have d greater than 69.

来源状态 open | 2026-07-23 | 来源

V3P-004711 | codetables-gf9-83-5

Minimum distance of a [83, 5] linear code over GF(9)

codes are known with d = 68, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-004712 | codetables-gf9-83-6

Minimum distance of a [83, 6] linear code over GF(9)

codes are known with d = 66, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-004713 | codetables-gf9-84-5

Minimum distance of a [84, 5] linear code over GF(9)

codes are known with d = 69, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-004714 | codetables-gf9-84-6

Minimum distance of a [84, 6] linear code over GF(9)

codes are known with d = 67, and no code can have d greater than 70.

来源状态 open | 2026-07-23 | 来源

V3P-004715 | codetables-gf9-85-5

Minimum distance of a [85, 5] linear code over GF(9)

codes are known with d = 70, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-004716 | codetables-gf9-85-6

Minimum distance of a [85, 6] linear code over GF(9)

codes are known with d = 67, and no code can have d greater than 71.

来源状态 open | 2026-07-23 | 来源

V3P-004717 | codetables-gf9-86-5

Minimum distance of a [86, 5] linear code over GF(9)

codes are known with d = 71, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-004718 | codetables-gf9-86-6

Minimum distance of a [86, 6] linear code over GF(9)

codes are known with d = 68, and no code can have d greater than 72.

来源状态 open | 2026-07-23 | 来源

V3P-004719 | codetables-gf9-87-5

Minimum distance of a [87, 5] linear code over GF(9)

codes are known with d = 71, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-004720 | codetables-gf9-87-6

Minimum distance of a [87, 6] linear code over GF(9)

codes are known with d = 69, and no code can have d greater than 73.

来源状态 open | 2026-07-23 | 来源

V3P-004721 | codetables-gf9-88-5

Minimum distance of a [88, 5] linear code over GF(9)

codes are known with d = 72, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-004722 | codetables-gf9-88-6

Minimum distance of a [88, 6] linear code over GF(9)

codes are known with d = 70, and no code can have d greater than 74.

来源状态 open | 2026-07-23 | 来源

V3P-004723 | codetables-gf9-89-5

Minimum distance of a [89, 5] linear code over GF(9)

codes are known with d = 73, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-004724 | codetables-gf9-89-6

Minimum distance of a [89, 6] linear code over GF(9)

codes are known with d = 71, and no code can have d greater than 75.

来源状态 open | 2026-07-23 | 来源

V3P-004725 | codetables-gf9-90-5

Minimum distance of a [90, 5] linear code over GF(9)

codes are known with d = 74, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-004726 | codetables-gf9-90-6

Minimum distance of a [90, 6] linear code over GF(9)

codes are known with d = 71, and no code can have d greater than 76.

来源状态 open | 2026-07-23 | 来源

V3P-004727 | codetables-gf9-91-5

Minimum distance of a [91, 5] linear code over GF(9)

codes are known with d = 75, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-004728 | codetables-gf9-91-6

Minimum distance of a [91, 6] linear code over GF(9)

codes are known with d = 72, and no code can have d greater than 77.

来源状态 open | 2026-07-23 | 来源

V3P-004729 | codetables-gf9-92-5

Minimum distance of a [92, 5] linear code over GF(9)

codes are known with d = 76, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-004730 | codetables-gf9-92-6

Minimum distance of a [92, 6] linear code over GF(9)

codes are known with d = 73, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-004731 | codetables-gf9-93-4

Minimum distance of a [93, 4] linear code over GF(9)

codes are known with d = 80, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-004732 | codetables-gf9-93-5

Minimum distance of a [93, 5] linear code over GF(9)

codes are known with d = 77, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-004733 | codetables-gf9-93-6

Minimum distance of a [93, 6] linear code over GF(9)

codes are known with d = 74, and no code can have d greater than 78.

来源状态 open | 2026-07-23 | 来源

V3P-004734 | codetables-gf9-94-5

Minimum distance of a [94, 5] linear code over GF(9)

codes are known with d = 78, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-004735 | codetables-gf9-94-6

Minimum distance of a [94, 6] linear code over GF(9)

codes are known with d = 75, and no code can have d greater than 79.

来源状态 open | 2026-07-23 | 来源

V3P-004736 | codetables-gf9-95-4

Minimum distance of a [95, 4] linear code over GF(9)

codes are known with d = 81, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-004737 | codetables-gf9-95-5

Minimum distance of a [95, 5] linear code over GF(9)

codes are known with d = 78, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-004738 | codetables-gf9-95-6

Minimum distance of a [95, 6] linear code over GF(9)

codes are known with d = 76, and no code can have d greater than 80.

来源状态 open | 2026-07-23 | 来源

V3P-004739 | codetables-gf9-96-4

Minimum distance of a [96, 4] linear code over GF(9)

codes are known with d = 82, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-004740 | codetables-gf9-96-5

Minimum distance of a [96, 5] linear code over GF(9)

codes are known with d = 79, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-004741 | codetables-gf9-96-6

Minimum distance of a [96, 6] linear code over GF(9)

codes are known with d = 76, and no code can have d greater than 81.

来源状态 open | 2026-07-23 | 来源

V3P-004742 | codetables-gf9-97-4

Minimum distance of a [97, 4] linear code over GF(9)

codes are known with d = 83, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-004743 | codetables-gf9-97-5

Minimum distance of a [97, 5] linear code over GF(9)

codes are known with d = 80, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-004744 | codetables-gf9-97-6

Minimum distance of a [97, 6] linear code over GF(9)

codes are known with d = 77, and no code can have d greater than 82.

来源状态 open | 2026-07-23 | 来源

V3P-004745 | codetables-gf9-98-4

Minimum distance of a [98, 4] linear code over GF(9)

codes are known with d = 84, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-004746 | codetables-gf9-98-5

Minimum distance of a [98, 5] linear code over GF(9)

codes are known with d = 81, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

V3P-004747 | codetables-gf9-98-6

Minimum distance of a [98, 6] linear code over GF(9)

codes are known with d = 78, and no code can have d greater than 83.

来源状态 open | 2026-07-23 | 来源

V3P-004748 | codetables-gf9-99-4

Minimum distance of a [99, 4] linear code over GF(9)

codes are known with d = 85, and no code can have d greater than 86.

来源状态 open | 2026-07-23 | 来源

V3P-004749 | codetables-gf9-99-5

Minimum distance of a [99, 5] linear code over GF(9)

codes are known with d = 82, and no code can have d greater than 85.

来源状态 open | 2026-07-23 | 来源

V3P-004750 | codetables-gf9-99-6

Minimum distance of a [99, 6] linear code over GF(9)

codes are known with d = 79, and no code can have d greater than 84.

来源状态 open | 2026-07-23 | 来源

D04 覆盖设计 | 1,480 行

C(v,k,t) 是覆盖 v 元集全部 t 元子集所需 k 元块的最少数目。best_known 为已知覆盖所用块数，是上界；lower_bound 是来源宣称的下界。来源下界与另列的计算所得 Schönheim 界可能不同；条目据此将来源原称的 Schoenheim bound 转述为 source-reported lower bound，本卷没有重新证明较强下界。

原始维护者：Daniel M. Gordon

Cite the La Jolla Covering Repository, D. M. Gordon, https://ljcr.dmgordon.org/cover.html

原始问题族入口

V3P-004751 | ljcr-c-11-6-5

Covering design C(11, 6, 5)

the best known covering uses 100 blocks and the source-reported lower bound is 96.

来源状态 open | 2026-07-23 | 来源

V3P-004752 | ljcr-c-12-6-4

Covering design C(12, 6, 4)

the best known covering uses 41 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-004753 | ljcr-c-12-7-5

Covering design C(12, 7, 5)

the best known covering uses 59 blocks and the source-reported lower bound is 55.

来源状态 open | 2026-07-23 | 来源

V3P-004754 | ljcr-c-13-5-4

Covering design C(13, 5, 4)

the best known covering uses 157 blocks and the source-reported lower bound is 149.

来源状态 open | 2026-07-23 | 来源

V3P-004755 | ljcr-c-13-6-3

Covering design C(13, 6, 3)

the best known covering uses 21 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-004756 | ljcr-c-13-6-4

Covering design C(13, 6, 4)

the best known covering uses 66 blocks and the source-reported lower bound is 63.

来源状态 open | 2026-07-23 | 来源

V3P-004757 | ljcr-c-13-7-4

Covering design C(13, 7, 4)

the best known covering uses 30 blocks and the source-reported lower bound is 28.

来源状态 open | 2026-07-23 | 来源

V3P-004758 | ljcr-c-13-7-5

Covering design C(13, 7, 5)

the best known covering uses 78 blocks and the source-reported lower bound is 75.

来源状态 open | 2026-07-23 | 来源

V3P-004759 | ljcr-c-13-8-5

Covering design C(13, 8, 5)

the best known covering uses 42 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-004760 | ljcr-c-14-5-3

Covering design C(14, 5, 3)

the best known covering uses 43 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-004761 | ljcr-c-14-5-4

Covering design C(14, 5, 4)

the best known covering uses 229 blocks and the source-reported lower bound is 219.

来源状态 open | 2026-07-23 | 来源

V3P-004762 | ljcr-c-14-6-3

Covering design C(14, 6, 3)

the best known covering uses 25 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-004763 | ljcr-c-14-6-4

Covering design C(14, 6, 4)

the best known covering uses 80 blocks and the source-reported lower bound is 75.

来源状态 open | 2026-07-23 | 来源

V3P-004764 | ljcr-c-14-7-4

Covering design C(14, 7, 4)

the best known covering uses 44 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-004765 | ljcr-c-14-7-5

Covering design C(14, 7, 5)

the best known covering uses 138 blocks and the source-reported lower bound is 126.

来源状态 open | 2026-07-23 | 来源

V3P-004766 | ljcr-c-14-8-4

Covering design C(14, 8, 4)

the best known covering uses 24 blocks and the source-reported lower bound is 23.

来源状态 open | 2026-07-23 | 来源

V3P-004767 | ljcr-c-14-8-5

Covering design C(14, 8, 5)

the best known covering uses 55 blocks and the source-reported lower bound is 49.

来源状态 open | 2026-07-23 | 来源

V3P-004768 | ljcr-c-14-9-5

Covering design C(14, 9, 5)

the best known covering uses 32 blocks and the source-reported lower bound is 28.

来源状态 open | 2026-07-23 | 来源

V3P-004769 | ljcr-c-15-10-4

Covering design C(15, 10, 4)

the best known covering uses 14 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-004770 | ljcr-c-15-10-5

Covering design C(15, 10, 5)

the best known covering uses 27 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-004771 | ljcr-c-15-5-3

Covering design C(15, 5, 3)

the best known covering uses 55 blocks and the source-reported lower bound is 54.

来源状态 open | 2026-07-23 | 来源

V3P-004772 | ljcr-c-15-6-3

Covering design C(15, 6, 3)

the best known covering uses 31 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-004773 | ljcr-c-15-6-4

Covering design C(15, 6, 4)

the best known covering uses 117 blocks and the source-reported lower bound is 93.

来源状态 open | 2026-07-23 | 来源

V3P-004774 | ljcr-c-15-7-4

Covering design C(15, 7, 4)

the best known covering uses 57 blocks and the source-reported lower bound is 52.

来源状态 open | 2026-07-23 | 来源

V3P-004775 | ljcr-c-15-7-5

Covering design C(15, 7, 5)

the best known covering uses 189 blocks and the source-reported lower bound is 172.

来源状态 open | 2026-07-23 | 来源

V3P-004776 | ljcr-c-15-8-4

Covering design C(15, 8, 4)

the best known covering uses 30 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-004777 | ljcr-c-15-8-5

Covering design C(15, 8, 5)

the best known covering uses 89 blocks and the source-reported lower bound is 75.

来源状态 open | 2026-07-23 | 来源

V3P-004778 | ljcr-c-15-9-4

Covering design C(15, 9, 4)

the best known covering uses 20 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-004779 | ljcr-c-15-9-5

Covering design C(15, 9, 5)

the best known covering uses 42 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-004780 | ljcr-c-16-10-4

Covering design C(16, 10, 4)

the best known covering uses 18 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-004781 | ljcr-c-16-10-5

Covering design C(16, 10, 5)

the best known covering uses 34 blocks and the source-reported lower bound is 31.

来源状态 open | 2026-07-23 | 来源

V3P-004782 | ljcr-c-16-11-5

Covering design C(16, 11, 5)

the best known covering uses 22 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-004783 | ljcr-c-16-5-3

Covering design C(16, 5, 3)

the best known covering uses 65 blocks and the source-reported lower bound is 61.

来源状态 open | 2026-07-23 | 来源

V3P-004784 | ljcr-c-16-6-3

Covering design C(16, 6, 3)

the best known covering uses 38 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-004785 | ljcr-c-16-6-4

Covering design C(16, 6, 4)

the best known covering uses 152 blocks and the source-reported lower bound is 144.

来源状态 open | 2026-07-23 | 来源

V3P-004786 | ljcr-c-16-7-3

Covering design C(16, 7, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 23.

来源状态 open | 2026-07-23 | 来源

V3P-004787 | ljcr-c-16-7-4

Covering design C(16, 7, 4)

the best known covering uses 76 blocks and the source-reported lower bound is 69.

来源状态 open | 2026-07-23 | 来源

V3P-004788 | ljcr-c-16-8-5

Covering design C(16, 8, 5)

the best known covering uses 115 blocks and the source-reported lower bound is 104.

来源状态 open | 2026-07-23 | 来源

V3P-004789 | ljcr-c-16-9-4

Covering design C(16, 9, 4)

the best known covering uses 26 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-004790 | ljcr-c-16-9-5

Covering design C(16, 9, 5)

the best known covering uses 61 blocks and the source-reported lower bound is 52.

来源状态 open | 2026-07-23 | 来源

V3P-004791 | ljcr-c-17-10-4

Covering design C(17, 10, 4)

the best known covering uses 23 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-004792 | ljcr-c-17-10-5

Covering design C(17, 10, 5)

the best known covering uses 48 blocks and the source-reported lower bound is 41.

来源状态 open | 2026-07-23 | 来源

V3P-004793 | ljcr-c-17-11-4

Covering design C(17, 11, 4)

the best known covering uses 15 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-004794 | ljcr-c-17-11-5

Covering design C(17, 11, 5)

the best known covering uses 30 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-004795 | ljcr-c-17-6-3

Covering design C(17, 6, 3)

the best known covering uses 44 blocks and the source-reported lower bound is 43.

来源状态 open | 2026-07-23 | 来源

V3P-004796 | ljcr-c-17-6-4

Covering design C(17, 6, 4)

the best known covering uses 188 blocks and the source-reported lower bound is 173.

来源状态 open | 2026-07-23 | 来源

V3P-004797 | ljcr-c-17-7-3

Covering design C(17, 7, 3)

the best known covering uses 27 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-004798 | ljcr-c-17-7-4

Covering design C(17, 7, 4)

the best known covering uses 98 blocks and the source-reported lower bound is 85.

来源状态 open | 2026-07-23 | 来源

V3P-004799 | ljcr-c-17-8-3

Covering design C(17, 8, 3)

the best known covering uses 18 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-004800 | ljcr-c-17-8-4

Covering design C(17, 8, 4)

the best known covering uses 53 blocks and the source-reported lower bound is 49.

来源状态 open | 2026-07-23 | 来源

V3P-004801 | ljcr-c-17-8-5

Covering design C(17, 8, 5)

the best known covering uses 177 blocks and the source-reported lower bound is 147.

来源状态 open | 2026-07-23 | 来源

V3P-004802 | ljcr-c-17-9-4

Covering design C(17, 9, 4)

the best known covering uses 28 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-004803 | ljcr-c-17-9-5

Covering design C(17, 9, 5)

the best known covering uses 79 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-004804 | ljcr-c-18-10-3

Covering design C(18, 10, 3)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-004805 | ljcr-c-18-10-4

Covering design C(18, 10, 4)

the best known covering uses 26 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-004806 | ljcr-c-18-10-5

Covering design C(18, 10, 5)

the best known covering uses 54 blocks and the source-reported lower bound is 49.

来源状态 open | 2026-07-23 | 来源

V3P-004807 | ljcr-c-18-11-3

Covering design C(18, 11, 3)

the best known covering uses 10 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-004808 | ljcr-c-18-11-4

Covering design C(18, 11, 4)

the best known covering uses 19 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-004809 | ljcr-c-18-11-5

Covering design C(18, 11, 5)

the best known covering uses 42 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-004810 | ljcr-c-18-12-5

Covering design C(18, 12, 5)

the best known covering uses 24 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-004811 | ljcr-c-18-6-4

Covering design C(18, 6, 4)

the best known covering uses 236 blocks and the source-reported lower bound is 205.

来源状态 open | 2026-07-23 | 来源

V3P-004812 | ljcr-c-18-7-3

Covering design C(18, 7, 3)

the best known covering uses 32 blocks and the source-reported lower bound is 31.

来源状态 open | 2026-07-23 | 来源

V3P-004813 | ljcr-c-18-7-4

Covering design C(18, 7, 4)

the best known covering uses 126 blocks and the source-reported lower bound is 111.

来源状态 open | 2026-07-23 | 来源

V3P-004814 | ljcr-c-18-8-4

Covering design C(18, 8, 4)

the best known covering uses 66 blocks and the source-reported lower bound is 57.

来源状态 open | 2026-07-23 | 来源

V3P-004815 | ljcr-c-18-8-5

Covering design C(18, 8, 5)

the best known covering uses 233 blocks and the source-reported lower bound is 192.

来源状态 open | 2026-07-23 | 来源

V3P-004816 | ljcr-c-18-9-3

Covering design C(18, 9, 3)

the best known covering uses 16 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-004817 | ljcr-c-18-9-4

Covering design C(18, 9, 4)

the best known covering uses 38 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-004818 | ljcr-c-18-9-5

Covering design C(18, 9, 5)

the best known covering uses 113 blocks and the source-reported lower bound is 98.

来源状态 open | 2026-07-23 | 来源

V3P-004819 | ljcr-c-19-10-3

Covering design C(19, 10, 3)

the best known covering uses 14 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-004820 | ljcr-c-19-10-4

Covering design C(19, 10, 4)

the best known covering uses 32 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-004821 | ljcr-c-19-10-5

Covering design C(19, 10, 5)

the best known covering uses 83 blocks and the source-reported lower bound is 65.

来源状态 open | 2026-07-23 | 来源

V3P-004822 | ljcr-c-19-11-3

Covering design C(19, 11, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-004823 | ljcr-c-19-11-4

Covering design C(19, 11, 4)

the best known covering uses 23 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-004824 | ljcr-c-19-11-5

Covering design C(19, 11, 5)

the best known covering uses 49 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-004825 | ljcr-c-19-12-4

Covering design C(19, 12, 4)

the best known covering uses 17 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-004826 | ljcr-c-19-12-5

Covering design C(19, 12, 5)

the best known covering uses 36 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-004827 | ljcr-c-19-13-5

Covering design C(19, 13, 5)

the best known covering uses 21 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-004828 | ljcr-c-19-5-3

Covering design C(19, 5, 3)

the best known covering uses 108 blocks and the source-reported lower bound is 103.

来源状态 open | 2026-07-23 | 来源

V3P-004829 | ljcr-c-19-6-3

Covering design C(19, 6, 3)

the best known covering uses 60 blocks and the source-reported lower bound is 57.

来源状态 open | 2026-07-23 | 来源

V3P-004830 | ljcr-c-19-7-3

Covering design C(19, 7, 3)

the best known covering uses 35 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-004831 | ljcr-c-19-7-4

Covering design C(19, 7, 4)

the best known covering uses 151 blocks and the source-reported lower bound is 131.

来源状态 open | 2026-07-23 | 来源

V3P-004832 | ljcr-c-19-8-3

Covering design C(19, 8, 3)

the best known covering uses 27 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-004833 | ljcr-c-19-8-4

Covering design C(19, 8, 4)

the best known covering uses 84 blocks and the source-reported lower bound is 74.

来源状态 open | 2026-07-23 | 来源

V3P-004834 | ljcr-c-19-9-3

Covering design C(19, 9, 3)

the best known covering uses 17 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-004835 | ljcr-c-19-9-4

Covering design C(19, 9, 4)

the best known covering uses 48 blocks and the source-reported lower bound is 45.

来源状态 open | 2026-07-23 | 来源

V3P-004836 | ljcr-c-19-9-5

Covering design C(19, 9, 5)

the best known covering uses 149 blocks and the source-reported lower bound is 121.

来源状态 open | 2026-07-23 | 来源

V3P-004837 | ljcr-c-20-10-3

Covering design C(20, 10, 3)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-004838 | ljcr-c-20-10-4

Covering design C(20, 10, 4)

the best known covering uses 36 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-004839 | ljcr-c-20-10-5

Covering design C(20, 10, 5)

the best known covering uses 96 blocks and the source-reported lower bound is 90.

来源状态 open | 2026-07-23 | 来源

V3P-004840 | ljcr-c-20-11-3

Covering design C(20, 11, 3)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-004841 | ljcr-c-20-11-4

Covering design C(20, 11, 4)

the best known covering uses 28 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-004842 | ljcr-c-20-11-5

Covering design C(20, 11, 5)

the best known covering uses 64 blocks and the source-reported lower bound is 50.

来源状态 open | 2026-07-23 | 来源

V3P-004843 | ljcr-c-20-12-3

Covering design C(20, 12, 3)

the best known covering uses 10 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-004844 | ljcr-c-20-12-4

Covering design C(20, 12, 4)

the best known covering uses 20 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-004845 | ljcr-c-20-12-5

Covering design C(20, 12, 5)

the best known covering uses 42 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-004846 | ljcr-c-20-13-4

Covering design C(20, 13, 4)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-004847 | ljcr-c-20-13-5

Covering design C(20, 13, 5)

the best known covering uses 30 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-004848 | ljcr-c-20-14-5

Covering design C(20, 14, 5)

the best known covering uses 18 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-004849 | ljcr-c-20-5-3

Covering design C(20, 5, 3)

the best known covering uses 133 blocks and the source-reported lower bound is 124.

来源状态 open | 2026-07-23 | 来源

V3P-004850 | ljcr-c-20-6-3

Covering design C(20, 6, 3)

the best known covering uses 71 blocks and the source-reported lower bound is 64.

来源状态 open | 2026-07-23 | 来源

V3P-004851 | ljcr-c-20-7-3

Covering design C(20, 7, 3)

the best known covering uses 45 blocks and the source-reported lower bound is 43.

来源状态 open | 2026-07-23 | 来源

V3P-004852 | ljcr-c-20-7-4

Covering design C(20, 7, 4)

the best known covering uses 198 blocks and the source-reported lower bound is 164.

来源状态 open | 2026-07-23 | 来源

V3P-004853 | ljcr-c-20-8-4

Covering design C(20, 8, 4)

the best known covering uses 93 blocks and the source-reported lower bound is 83.

来源状态 open | 2026-07-23 | 来源

V3P-004854 | ljcr-c-20-9-3

Covering design C(20, 9, 3)

the best known covering uses 21 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-004855 | ljcr-c-20-9-4

Covering design C(20, 9, 4)

the best known covering uses 63 blocks and the source-reported lower bound is 54.

来源状态 open | 2026-07-23 | 来源

V3P-004856 | ljcr-c-20-9-5

Covering design C(20, 9, 5)

the best known covering uses 211 blocks and the source-reported lower bound is 165.

来源状态 open | 2026-07-23 | 来源

V3P-004857 | ljcr-c-21-10-3

Covering design C(21, 10, 3)

the best known covering uses 18 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-004858 | ljcr-c-21-10-4

Covering design C(21, 10, 4)

the best known covering uses 49 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-004859 | ljcr-c-21-10-5

Covering design C(21, 10, 5)

the best known covering uses 145 blocks and the source-reported lower bound is 114.

来源状态 open | 2026-07-23 | 来源

V3P-004860 | ljcr-c-21-11-3

Covering design C(21, 11, 3)

the best known covering uses 14 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-004861 | ljcr-c-21-11-4

Covering design C(21, 11, 4)

the best known covering uses 31 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-004862 | ljcr-c-21-11-5

Covering design C(21, 11, 5)

the best known covering uses 78 blocks and the source-reported lower bound is 62.

来源状态 open | 2026-07-23 | 来源

V3P-004863 | ljcr-c-21-12-4

Covering design C(21, 12, 4)

the best known covering uses 25 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-004864 | ljcr-c-21-12-5

Covering design C(21, 12, 5)

the best known covering uses 54 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-004865 | ljcr-c-21-13-4

Covering design C(21, 13, 4)

the best known covering uses 18 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-004866 | ljcr-c-21-13-5

Covering design C(21, 13, 5)

the best known covering uses 37 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-004867 | ljcr-c-21-14-4

Covering design C(21, 14, 4)

the best known covering uses 14 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-004868 | ljcr-c-21-14-5

Covering design C(21, 14, 5)

the best known covering uses 27 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-004869 | ljcr-c-21-15-5

Covering design C(21, 15, 5)

the best known covering uses 16 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-004870 | ljcr-c-21-5-3

Covering design C(21, 5, 3)

the best known covering uses 151 blocks and the source-reported lower bound is 147.

来源状态 open | 2026-07-23 | 来源

V3P-004871 | ljcr-c-21-6-3

Covering design C(21, 6, 3)

the best known covering uses 77 blocks and the source-reported lower bound is 74.

来源状态 open | 2026-07-23 | 来源

V3P-004872 | ljcr-c-21-7-3

Covering design C(21, 7, 3)

the best known covering uses 49 blocks and the source-reported lower bound is 48.

来源状态 open | 2026-07-23 | 来源

V3P-004873 | ljcr-c-21-7-4

Covering design C(21, 7, 4)

the best known covering uses 235 blocks and the source-reported lower bound is 192.

来源状态 open | 2026-07-23 | 来源

V3P-004874 | ljcr-c-21-8-3

Covering design C(21, 8, 3)

the best known covering uses 35 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-004875 | ljcr-c-21-8-4

Covering design C(21, 8, 4)

the best known covering uses 124 blocks and the source-reported lower bound is 113.

来源状态 open | 2026-07-23 | 来源

V3P-004876 | ljcr-c-21-9-3

Covering design C(21, 9, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-004877 | ljcr-c-21-9-4

Covering design C(21, 9, 4)

the best known covering uses 70 blocks and the source-reported lower bound is 66.

来源状态 open | 2026-07-23 | 来源

V3P-004878 | ljcr-c-22-10-3

Covering design C(22, 10, 3)

the best known covering uses 19 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-004879 | ljcr-c-22-10-4

Covering design C(22, 10, 4)

the best known covering uses 54 blocks and the source-reported lower bound is 47.

来源状态 open | 2026-07-23 | 来源

V3P-004880 | ljcr-c-22-10-5

Covering design C(22, 10, 5)

the best known covering uses 174 blocks and the source-reported lower bound is 146.

来源状态 open | 2026-07-23 | 来源

V3P-004881 | ljcr-c-22-11-3

Covering design C(22, 11, 3)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-004882 | ljcr-c-22-11-4

Covering design C(22, 11, 4)

the best known covering uses 38 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-004883 | ljcr-c-22-11-5

Covering design C(22, 11, 5)

the best known covering uses 108 blocks and the source-reported lower bound is 84.

来源状态 open | 2026-07-23 | 来源

V3P-004884 | ljcr-c-22-12-4

Covering design C(22, 12, 4)

the best known covering uses 28 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-004885 | ljcr-c-22-12-5

Covering design C(22, 12, 5)

the best known covering uses 63 blocks and the source-reported lower bound is 50.

来源状态 open | 2026-07-23 | 来源

V3P-004886 | ljcr-c-22-13-3

Covering design C(22, 13, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-004887 | ljcr-c-22-13-4

Covering design C(22, 13, 4)

the best known covering uses 22 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-004888 | ljcr-c-22-13-5

Covering design C(22, 13, 5)

the best known covering uses 47 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-004889 | ljcr-c-22-14-4

Covering design C(22, 14, 4)

the best known covering uses 17 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-004890 | ljcr-c-22-14-5

Covering design C(22, 14, 5)

the best known covering uses 33 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-004891 | ljcr-c-22-15-4

Covering design C(22, 15, 4)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-004892 | ljcr-c-22-15-5

Covering design C(22, 15, 5)

the best known covering uses 22 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-004893 | ljcr-c-22-16-5

Covering design C(22, 16, 5)

the best known covering uses 14 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-004894 | ljcr-c-22-17-5

Covering design C(22, 17, 5)

the best known covering uses 11 blocks and the source-reported lower bound is 10.

来源状态 open | 2026-07-23 | 来源

V3P-004895 | ljcr-c-22-5-3

Covering design C(22, 5, 3)

the best known covering uses 172 blocks and the source-reported lower bound is 163.

来源状态 open | 2026-07-23 | 来源

V3P-004896 | ljcr-c-22-7-3

Covering design C(22, 7, 3)

the best known covering uses 59 blocks and the source-reported lower bound is 54.

来源状态 open | 2026-07-23 | 来源

V3P-004897 | ljcr-c-22-8-3

Covering design C(22, 8, 3)

the best known covering uses 38 blocks and the source-reported lower bound is 36.

来源状态 open | 2026-07-23 | 来源

V3P-004898 | ljcr-c-22-8-4

Covering design C(22, 8, 4)

the best known covering uses 152 blocks and the source-reported lower bound is 132.

来源状态 open | 2026-07-23 | 来源

V3P-004899 | ljcr-c-22-9-3

Covering design C(22, 9, 3)

the best known covering uses 28 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-004900 | ljcr-c-22-9-4

Covering design C(22, 9, 4)

the best known covering uses 95 blocks and the source-reported lower bound is 79.

来源状态 open | 2026-07-23 | 来源

V3P-004901 | ljcr-c-23-10-3

Covering design C(23, 10, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-004902 | ljcr-c-23-10-4

Covering design C(23, 10, 4)

the best known covering uses 74 blocks and the source-reported lower bound is 63.

来源状态 open | 2026-07-23 | 来源

V3P-004903 | ljcr-c-23-10-5

Covering design C(23, 10, 5)

the best known covering uses 240 blocks and the source-reported lower bound is 182.

来源状态 open | 2026-07-23 | 来源

V3P-004904 | ljcr-c-23-11-4

Covering design C(23, 11, 4)

the best known covering uses 42 blocks and the source-reported lower bound is 36.

来源状态 open | 2026-07-23 | 来源

V3P-004905 | ljcr-c-23-11-5

Covering design C(23, 11, 5)

the best known covering uses 131 blocks and the source-reported lower bound is 99.

来源状态 open | 2026-07-23 | 来源

V3P-004906 | ljcr-c-23-12-3

Covering design C(23, 12, 3)

the best known covering uses 14 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-004907 | ljcr-c-23-12-4

Covering design C(23, 12, 4)

the best known covering uses 31 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-004908 | ljcr-c-23-12-5

Covering design C(23, 12, 5)

the best known covering uses 80 blocks and the source-reported lower bound is 62.

来源状态 open | 2026-07-23 | 来源

V3P-004909 | ljcr-c-23-13-4

Covering design C(23, 13, 4)

the best known covering uses 25 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-004910 | ljcr-c-23-13-5

Covering design C(23, 13, 5)

the best known covering uses 54 blocks and the source-reported lower bound is 43.

来源状态 open | 2026-07-23 | 来源

V3P-004911 | ljcr-c-23-14-3

Covering design C(23, 14, 3)

the best known covering uses 10 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-004912 | ljcr-c-23-14-4

Covering design C(23, 14, 4)

the best known covering uses 20 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-004913 | ljcr-c-23-14-5

Covering design C(23, 14, 5)

the best known covering uses 42 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-004914 | ljcr-c-23-15-4

Covering design C(23, 15, 4)

the best known covering uses 15 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-004915 | ljcr-c-23-15-5

Covering design C(23, 15, 5)

the best known covering uses 29 blocks and the source-reported lower bound is 23.

来源状态 open | 2026-07-23 | 来源

V3P-004916 | ljcr-c-23-16-4

Covering design C(23, 16, 4)

the best known covering uses 11 blocks and the source-reported lower bound is 10.

来源状态 open | 2026-07-23 | 来源

V3P-004917 | ljcr-c-23-16-5

Covering design C(23, 16, 5)

the best known covering uses 20 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-004918 | ljcr-c-23-5-3

Covering design C(23, 5, 3)

the best known covering uses 187 blocks and the source-reported lower bound is 180.

来源状态 open | 2026-07-23 | 来源

V3P-004919 | ljcr-c-23-6-2

Covering design C(23, 6, 2)

the best known covering uses 21 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-004920 | ljcr-c-23-7-2

Covering design C(23, 7, 2)

the best known covering uses 16 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-004921 | ljcr-c-23-7-3

Covering design C(23, 7, 3)

the best known covering uses 65 blocks and the source-reported lower bound is 63.

来源状态 open | 2026-07-23 | 来源

V3P-004922 | ljcr-c-23-8-3

Covering design C(23, 8, 3)

the best known covering uses 40 blocks and the source-reported lower bound is 38.

来源状态 open | 2026-07-23 | 来源

V3P-004923 | ljcr-c-23-8-4

Covering design C(23, 8, 4)

the best known covering uses 193 blocks and the source-reported lower bound is 156.

来源状态 open | 2026-07-23 | 来源

V3P-004924 | ljcr-c-23-9-3

Covering design C(23, 9, 3)

the best known covering uses 32 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-004925 | ljcr-c-23-9-4

Covering design C(23, 9, 4)

the best known covering uses 107 blocks and the source-reported lower bound is 92.

来源状态 open | 2026-07-23 | 来源

V3P-004926 | ljcr-c-24-10-4

Covering design C(24, 10, 4)

the best known covering uses 87 blocks and the source-reported lower bound is 70.

来源状态 open | 2026-07-23 | 来源

V3P-004927 | ljcr-c-24-11-3

Covering design C(24, 11, 3)

the best known covering uses 20 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-004928 | ljcr-c-24-11-4

Covering design C(24, 11, 4)

the best known covering uses 57 blocks and the source-reported lower bound is 46.

来源状态 open | 2026-07-23 | 来源

V3P-004929 | ljcr-c-24-11-5

Covering design C(24, 11, 5)

the best known covering uses 192 blocks and the source-reported lower bound is 138.

来源状态 open | 2026-07-23 | 来源

V3P-004930 | ljcr-c-24-12-4

Covering design C(24, 12, 4)

the best known covering uses 31 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-004931 | ljcr-c-24-12-5

Covering design C(24, 12, 5)

the best known covering uses 86 blocks and the source-reported lower bound is 72.

来源状态 open | 2026-07-23 | 来源

V3P-004932 | ljcr-c-24-13-3

Covering design C(24, 13, 3)

the best known covering uses 13 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-004933 | ljcr-c-24-13-4

Covering design C(24, 13, 4)

the best known covering uses 28 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-004934 | ljcr-c-24-13-5

Covering design C(24, 13, 5)

the best known covering uses 64 blocks and the source-reported lower bound is 50.

来源状态 open | 2026-07-23 | 来源

V3P-004935 | ljcr-c-24-14-3

Covering design C(24, 14, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-004936 | ljcr-c-24-14-4

Covering design C(24, 14, 4)

the best known covering uses 23 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-004937 | ljcr-c-24-14-5

Covering design C(24, 14, 5)

the best known covering uses 49 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-004938 | ljcr-c-24-15-4

Covering design C(24, 15, 4)

the best known covering uses 18 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-004939 | ljcr-c-24-15-5

Covering design C(24, 15, 5)

the best known covering uses 36 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-004940 | ljcr-c-24-16-4

Covering design C(24, 16, 4)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-004941 | ljcr-c-24-16-5

Covering design C(24, 16, 5)

the best known covering uses 24 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-004942 | ljcr-c-24-17-5

Covering design C(24, 17, 5)

the best known covering uses 18 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-004943 | ljcr-c-24-18-5

Covering design C(24, 18, 5)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-004944 | ljcr-c-24-5-3

Covering design C(24, 5, 3)

the best known covering uses 231 blocks and the source-reported lower bound is 222.

来源状态 open | 2026-07-23 | 来源

V3P-004945 | ljcr-c-24-6-2

Covering design C(24, 6, 2)

the best known covering uses 22 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-004946 | ljcr-c-24-6-3

Covering design C(24, 6, 3)

the best known covering uses 116 blocks and the source-reported lower bound is 112.

来源状态 open | 2026-07-23 | 来源

V3P-004947 | ljcr-c-24-7-3

Covering design C(24, 7, 3)

the best known covering uses 76 blocks and the source-reported lower bound is 69.

来源状态 open | 2026-07-23 | 来源

V3P-004948 | ljcr-c-24-8-3

Covering design C(24, 8, 3)

the best known covering uses 50 blocks and the source-reported lower bound is 45.

来源状态 open | 2026-07-23 | 来源

V3P-004949 | ljcr-c-24-8-4

Covering design C(24, 8, 4)

the best known covering uses 226 blocks and the source-reported lower bound is 189.

来源状态 open | 2026-07-23 | 来源

V3P-004950 | ljcr-c-24-9-3

Covering design C(24, 9, 3)

the best known covering uses 35 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-004951 | ljcr-c-24-9-4

Covering design C(24, 9, 4)

the best known covering uses 122 blocks and the source-reported lower bound is 102.

来源状态 open | 2026-07-23 | 来源

V3P-004952 | ljcr-c-25-10-3

Covering design C(25, 10, 3)

the best known covering uses 30 blocks and the source-reported lower bound is 28.

来源状态 open | 2026-07-23 | 来源

V3P-004953 | ljcr-c-25-10-4

Covering design C(25, 10, 4)

the best known covering uses 97 blocks and the source-reported lower bound is 80.

来源状态 open | 2026-07-23 | 来源

V3P-004954 | ljcr-c-25-11-3

Covering design C(25, 11, 3)

the best known covering uses 23 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-004955 | ljcr-c-25-11-4

Covering design C(25, 11, 4)

the best known covering uses 66 blocks and the source-reported lower bound is 55.

来源状态 open | 2026-07-23 | 来源

V3P-004956 | ljcr-c-25-11-5

Covering design C(25, 11, 5)

the best known covering uses 226 blocks and the source-reported lower bound is 160.

来源状态 open | 2026-07-23 | 来源

V3P-004957 | ljcr-c-25-12-3

Covering design C(25, 12, 3)

the best known covering uses 17 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-004958 | ljcr-c-25-12-4

Covering design C(25, 12, 4)

the best known covering uses 47 blocks and the source-reported lower bound is 38.

来源状态 open | 2026-07-23 | 来源

V3P-004959 | ljcr-c-25-12-5

Covering design C(25, 12, 5)

the best known covering uses 133 blocks and the source-reported lower bound is 96.

来源状态 open | 2026-07-23 | 来源

V3P-004960 | ljcr-c-25-13-3

Covering design C(25, 13, 3)

the best known covering uses 14 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-004961 | ljcr-c-25-13-4

Covering design C(25, 13, 4)

the best known covering uses 30 blocks and the source-reported lower bound is 28.

来源状态 open | 2026-07-23 | 来源

V3P-004962 | ljcr-c-25-13-5

Covering design C(25, 13, 5)

the best known covering uses 72 blocks and the source-reported lower bound is 58.

来源状态 open | 2026-07-23 | 来源

V3P-004963 | ljcr-c-25-14-3

Covering design C(25, 14, 3)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-004964 | ljcr-c-25-14-4

Covering design C(25, 14, 4)

the best known covering uses 26 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-004965 | ljcr-c-25-14-5

Covering design C(25, 14, 5)

the best known covering uses 56 blocks and the source-reported lower bound is 43.

来源状态 open | 2026-07-23 | 来源

V3P-004966 | ljcr-c-25-15-3

Covering design C(25, 15, 3)

the best known covering uses 10 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-004967 | ljcr-c-25-15-4

Covering design C(25, 15, 4)

the best known covering uses 21 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-004968 | ljcr-c-25-15-5

Covering design C(25, 15, 5)

the best known covering uses 44 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-004969 | ljcr-c-25-16-4

Covering design C(25, 16, 4)

the best known covering uses 17 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-004970 | ljcr-c-25-16-5

Covering design C(25, 16, 5)

the best known covering uses 32 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-004971 | ljcr-c-25-17-4

Covering design C(25, 17, 4)

the best known covering uses 11 blocks and the source-reported lower bound is 10.

来源状态 open | 2026-07-23 | 来源

V3P-004972 | ljcr-c-25-17-5

Covering design C(25, 17, 5)

the best known covering uses 20 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-004973 | ljcr-c-25-18-5

Covering design C(25, 18, 5)

the best known covering uses 16 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-004974 | ljcr-c-25-6-3

Covering design C(25, 6, 3)

the best known covering uses 130 blocks and the source-reported lower bound is 125.

来源状态 open | 2026-07-23 | 来源

V3P-004975 | ljcr-c-25-7-3

Covering design C(25, 7, 3)

the best known covering uses 83 blocks and the source-reported lower bound is 75.

来源状态 open | 2026-07-23 | 来源

V3P-004976 | ljcr-c-25-8-3

Covering design C(25, 8, 3)

the best known covering uses 57 blocks and the source-reported lower bound is 54.

来源状态 open | 2026-07-23 | 来源

V3P-004977 | ljcr-c-25-9-3

Covering design C(25, 9, 3)

the best known covering uses 38 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-004978 | ljcr-c-25-9-4

Covering design C(25, 9, 4)

the best known covering uses 162 blocks and the source-reported lower bound is 125.

来源状态 open | 2026-07-23 | 来源

V3P-004979 | ljcr-c-26-10-3

Covering design C(26, 10, 3)

the best known covering uses 33 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-004980 | ljcr-c-26-10-4

Covering design C(26, 10, 4)

the best known covering uses 117 blocks and the source-reported lower bound is 89.

来源状态 open | 2026-07-23 | 来源

V3P-004981 | ljcr-c-26-11-3

Covering design C(26, 11, 3)

the best known covering uses 26 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-004982 | ljcr-c-26-11-4

Covering design C(26, 11, 4)

the best known covering uses 80 blocks and the source-reported lower bound is 67.

来源状态 open | 2026-07-23 | 来源

V3P-004983 | ljcr-c-26-12-3

Covering design C(26, 12, 3)

the best known covering uses 18 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-004984 | ljcr-c-26-12-4

Covering design C(26, 12, 4)

the best known covering uses 51 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-004985 | ljcr-c-26-12-5

Covering design C(26, 12, 5)

the best known covering uses 173 blocks and the source-reported lower bound is 120.

来源状态 open | 2026-07-23 | 来源

V3P-004986 | ljcr-c-26-13-3

Covering design C(26, 13, 3)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-004987 | ljcr-c-26-13-4

Covering design C(26, 13, 4)

the best known covering uses 37 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-004988 | ljcr-c-26-13-5

Covering design C(26, 13, 5)

the best known covering uses 101 blocks and the source-reported lower bound is 76.

来源状态 open | 2026-07-23 | 来源

V3P-004989 | ljcr-c-26-14-3

Covering design C(26, 14, 3)

the best known covering uses 13 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-004990 | ljcr-c-26-14-4

Covering design C(26, 14, 4)

the best known covering uses 28 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-004991 | ljcr-c-26-14-5

Covering design C(26, 14, 5)

the best known covering uses 60 blocks and the source-reported lower bound is 52.

来源状态 open | 2026-07-23 | 来源

V3P-004992 | ljcr-c-26-15-3

Covering design C(26, 15, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-004993 | ljcr-c-26-15-4

Covering design C(26, 15, 4)

the best known covering uses 24 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-004994 | ljcr-c-26-15-5

Covering design C(26, 15, 5)

the best known covering uses 52 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-004995 | ljcr-c-26-16-4

Covering design C(26, 16, 4)

the best known covering uses 18 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-004996 | ljcr-c-26-16-5

Covering design C(26, 16, 5)

the best known covering uses 39 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-004997 | ljcr-c-26-17-4

Covering design C(26, 17, 4)

the best known covering uses 15 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-004998 | ljcr-c-26-17-5

Covering design C(26, 17, 5)

the best known covering uses 26 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-004999 | ljcr-c-26-18-5

Covering design C(26, 18, 5)

the best known covering uses 19 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005000 | ljcr-c-26-19-5

Covering design C(26, 19, 5)

the best known covering uses 14 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-005001 | ljcr-c-26-7-3

Covering design C(26, 7, 3)

the best known covering uses 94 blocks and the source-reported lower bound is 86.

来源状态 open | 2026-07-23 | 来源

V3P-005002 | ljcr-c-26-8-3

Covering design C(26, 8, 3)

the best known covering uses 64 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-005003 | ljcr-c-26-9-3

Covering design C(26, 9, 3)

the best known covering uses 39 blocks and the source-reported lower bound is 38.

来源状态 open | 2026-07-23 | 来源

V3P-005004 | ljcr-c-26-9-4

Covering design C(26, 9, 4)

the best known covering uses 188 blocks and the source-reported lower bound is 156.

来源状态 open | 2026-07-23 | 来源

V3P-005005 | ljcr-c-27-10-3

Covering design C(27, 10, 3)

the best known covering uses 35 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005006 | ljcr-c-27-10-4

Covering design C(27, 10, 4)

the best known covering uses 129 blocks and the source-reported lower bound is 103.

来源状态 open | 2026-07-23 | 来源

V3P-005007 | ljcr-c-27-11-3

Covering design C(27, 11, 3)

the best known covering uses 27 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005008 | ljcr-c-27-11-4

Covering design C(27, 11, 4)

the best known covering uses 94 blocks and the source-reported lower bound is 72.

来源状态 open | 2026-07-23 | 来源

V3P-005009 | ljcr-c-27-12-3

Covering design C(27, 12, 3)

the best known covering uses 22 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-005010 | ljcr-c-27-12-4

Covering design C(27, 12, 4)

the best known covering uses 63 blocks and the source-reported lower bound is 54.

来源状态 open | 2026-07-23 | 来源

V3P-005011 | ljcr-c-27-12-5

Covering design C(27, 12, 5)

the best known covering uses 208 blocks and the source-reported lower bound is 151.

来源状态 open | 2026-07-23 | 来源

V3P-005012 | ljcr-c-27-13-4

Covering design C(27, 13, 4)

the best known covering uses 42 blocks and the source-reported lower bound is 36.

来源状态 open | 2026-07-23 | 来源

V3P-005013 | ljcr-c-27-13-5

Covering design C(27, 13, 5)

the best known covering uses 125 blocks and the source-reported lower bound is 88.

来源状态 open | 2026-07-23 | 来源

V3P-005014 | ljcr-c-27-14-3

Covering design C(27, 14, 3)

the best known covering uses 14 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-005015 | ljcr-c-27-14-4

Covering design C(27, 14, 4)

the best known covering uses 31 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005016 | ljcr-c-27-14-5

Covering design C(27, 14, 5)

the best known covering uses 76 blocks and the source-reported lower bound is 58.

来源状态 open | 2026-07-23 | 来源

V3P-005017 | ljcr-c-27-15-4

Covering design C(27, 15, 4)

the best known covering uses 26 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005018 | ljcr-c-27-15-5

Covering design C(27, 15, 5)

the best known covering uses 56 blocks and the source-reported lower bound is 45.

来源状态 open | 2026-07-23 | 来源

V3P-005019 | ljcr-c-27-16-3

Covering design C(27, 16, 3)

the best known covering uses 10 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005020 | ljcr-c-27-16-4

Covering design C(27, 16, 4)

the best known covering uses 22 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005021 | ljcr-c-27-16-5

Covering design C(27, 16, 5)

the best known covering uses 46 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005022 | ljcr-c-27-17-4

Covering design C(27, 17, 4)

the best known covering uses 18 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005023 | ljcr-c-27-17-5

Covering design C(27, 17, 5)

the best known covering uses 35 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005024 | ljcr-c-27-18-4

Covering design C(27, 18, 4)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-005025 | ljcr-c-27-18-5

Covering design C(27, 18, 5)

the best known covering uses 25 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-005026 | ljcr-c-27-19-5

Covering design C(27, 19, 5)

the best known covering uses 17 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005027 | ljcr-c-27-7-3

Covering design C(27, 7, 3)

the best known covering uses 104 blocks and the source-reported lower bound is 93.

来源状态 open | 2026-07-23 | 来源

V3P-005028 | ljcr-c-27-8-2

Covering design C(27, 8, 2)

the best known covering uses 17 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005029 | ljcr-c-27-8-3

Covering design C(27, 8, 3)

the best known covering uses 69 blocks and the source-reported lower bound is 65.

来源状态 open | 2026-07-23 | 来源

V3P-005030 | ljcr-c-27-9-4

Covering design C(27, 9, 4)

the best known covering uses 215 blocks and the source-reported lower bound is 177.

来源状态 open | 2026-07-23 | 来源

V3P-005031 | ljcr-c-28-10-3

Covering design C(28, 10, 3)

the best known covering uses 36 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005032 | ljcr-c-28-10-4

Covering design C(28, 10, 4)

the best known covering uses 154 blocks and the source-reported lower bound is 110.

来源状态 open | 2026-07-23 | 来源

V3P-005033 | ljcr-c-28-11-3

Covering design C(28, 11, 3)

the best known covering uses 31 blocks and the source-reported lower bound is 28.

来源状态 open | 2026-07-23 | 来源

V3P-005034 | ljcr-c-28-11-4

Covering design C(28, 11, 4)

the best known covering uses 103 blocks and the source-reported lower bound is 84.

来源状态 open | 2026-07-23 | 来源

V3P-005035 | ljcr-c-28-12-3

Covering design C(28, 12, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005036 | ljcr-c-28-12-4

Covering design C(28, 12, 4)

the best known covering uses 74 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-005037 | ljcr-c-28-13-3

Covering design C(28, 13, 3)

the best known covering uses 19 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005038 | ljcr-c-28-13-4

Covering design C(28, 13, 4)

the best known covering uses 54 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-005039 | ljcr-c-28-13-5

Covering design C(28, 13, 5)

the best known covering uses 169 blocks and the source-reported lower bound is 117.

来源状态 open | 2026-07-23 | 来源

V3P-005040 | ljcr-c-28-14-3

Covering design C(28, 14, 3)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005041 | ljcr-c-28-14-4

Covering design C(28, 14, 4)

the best known covering uses 31 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-005042 | ljcr-c-28-14-5

Covering design C(28, 14, 5)

the best known covering uses 88 blocks and the source-reported lower bound is 72.

来源状态 open | 2026-07-23 | 来源

V3P-005043 | ljcr-c-28-15-3

Covering design C(28, 15, 3)

the best known covering uses 13 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005044 | ljcr-c-28-15-4

Covering design C(28, 15, 4)

the best known covering uses 29 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005045 | ljcr-c-28-15-5

Covering design C(28, 15, 5)

the best known covering uses 62 blocks and the source-reported lower bound is 51.

来源状态 open | 2026-07-23 | 来源

V3P-005046 | ljcr-c-28-16-3

Covering design C(28, 16, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005047 | ljcr-c-28-16-4

Covering design C(28, 16, 4)

the best known covering uses 24 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-005048 | ljcr-c-28-16-5

Covering design C(28, 16, 5)

the best known covering uses 52 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-005049 | ljcr-c-28-17-3

Covering design C(28, 17, 3)

the best known covering uses 10 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005050 | ljcr-c-28-17-4

Covering design C(28, 17, 4)

the best known covering uses 21 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005051 | ljcr-c-28-17-5

Covering design C(28, 17, 5)

the best known covering uses 42 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005052 | ljcr-c-28-18-4

Covering design C(28, 18, 4)

the best known covering uses 16 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005053 | ljcr-c-28-18-5

Covering design C(28, 18, 5)

the best known covering uses 32 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005054 | ljcr-c-28-19-4

Covering design C(28, 19, 4)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-005055 | ljcr-c-28-19-5

Covering design C(28, 19, 5)

the best known covering uses 23 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005056 | ljcr-c-28-21-5

Covering design C(28, 21, 5)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-005057 | ljcr-c-28-6-3

Covering design C(28, 6, 3)

the best known covering uses 185 blocks and the source-reported lower bound is 178.

来源状态 open | 2026-07-23 | 来源

V3P-005058 | ljcr-c-28-7-2

Covering design C(28, 7, 2)

the best known covering uses 22 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005059 | ljcr-c-28-7-3

Covering design C(28, 7, 3)

the best known covering uses 120 blocks and the source-reported lower bound is 108.

来源状态 open | 2026-07-23 | 来源

V3P-005060 | ljcr-c-28-8-3

Covering design C(28, 8, 3)

the best known covering uses 73 blocks and the source-reported lower bound is 70.

来源状态 open | 2026-07-23 | 来源

V3P-005061 | ljcr-c-28-9-3

Covering design C(28, 9, 3)

the best known covering uses 56 blocks and the source-reported lower bound is 50.

来源状态 open | 2026-07-23 | 来源

V3P-005062 | ljcr-c-29-10-3

Covering design C(29, 10, 3)

the best known covering uses 42 blocks and the source-reported lower bound is 41.

来源状态 open | 2026-07-23 | 来源

V3P-005063 | ljcr-c-29-10-4

Covering design C(29, 10, 4)

the best known covering uses 186 blocks and the source-reported lower bound is 145.

来源状态 open | 2026-07-23 | 来源

V3P-005064 | ljcr-c-29-11-3

Covering design C(29, 11, 3)

the best known covering uses 33 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005065 | ljcr-c-29-11-4

Covering design C(29, 11, 4)

the best known covering uses 115 blocks and the source-reported lower bound is 90.

来源状态 open | 2026-07-23 | 来源

V3P-005066 | ljcr-c-29-12-3

Covering design C(29, 12, 3)

the best known covering uses 27 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005067 | ljcr-c-29-12-4

Covering design C(29, 12, 4)

the best known covering uses 88 blocks and the source-reported lower bound is 68.

来源状态 open | 2026-07-23 | 来源

V3P-005068 | ljcr-c-29-13-3

Covering design C(29, 13, 3)

the best known covering uses 21 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005069 | ljcr-c-29-13-4

Covering design C(29, 13, 4)

the best known covering uses 59 blocks and the source-reported lower bound is 47.

来源状态 open | 2026-07-23 | 来源

V3P-005070 | ljcr-c-29-13-5

Covering design C(29, 13, 5)

the best known covering uses 215 blocks and the source-reported lower bound is 132.

来源状态 open | 2026-07-23 | 来源

V3P-005071 | ljcr-c-29-14-4

Covering design C(29, 14, 4)

the best known covering uses 43 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005072 | ljcr-c-29-14-5

Covering design C(29, 14, 5)

the best known covering uses 122 blocks and the source-reported lower bound is 81.

来源状态 open | 2026-07-23 | 来源

V3P-005073 | ljcr-c-29-15-4

Covering design C(29, 15, 4)

the best known covering uses 30 blocks and the source-reported lower bound is 28.

来源状态 open | 2026-07-23 | 来源

V3P-005074 | ljcr-c-29-15-5

Covering design C(29, 15, 5)

the best known covering uses 67 blocks and the source-reported lower bound is 58.

来源状态 open | 2026-07-23 | 来源

V3P-005075 | ljcr-c-29-16-3

Covering design C(29, 16, 3)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-005076 | ljcr-c-29-16-4

Covering design C(29, 16, 4)

the best known covering uses 27 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005077 | ljcr-c-29-16-5

Covering design C(29, 16, 5)

the best known covering uses 58 blocks and the source-reported lower bound is 46.

来源状态 open | 2026-07-23 | 来源

V3P-005078 | ljcr-c-29-17-3

Covering design C(29, 17, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005079 | ljcr-c-29-17-4

Covering design C(29, 17, 4)

the best known covering uses 23 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005080 | ljcr-c-29-17-5

Covering design C(29, 17, 5)

the best known covering uses 48 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-005081 | ljcr-c-29-18-4

Covering design C(29, 18, 4)

the best known covering uses 18 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005082 | ljcr-c-29-18-5

Covering design C(29, 18, 5)

the best known covering uses 39 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005083 | ljcr-c-29-19-4

Covering design C(29, 19, 4)

the best known covering uses 15 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-005084 | ljcr-c-29-19-5

Covering design C(29, 19, 5)

the best known covering uses 29 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005085 | ljcr-c-29-20-4

Covering design C(29, 20, 4)

the best known covering uses 11 blocks and the source-reported lower bound is 10.

来源状态 open | 2026-07-23 | 来源

V3P-005086 | ljcr-c-29-20-5

Covering design C(29, 20, 5)

the best known covering uses 21 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005087 | ljcr-c-29-21-5

Covering design C(29, 21, 5)

the best known covering uses 14 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-005088 | ljcr-c-29-22-5

Covering design C(29, 22, 5)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-005089 | ljcr-c-29-5-2

Covering design C(29, 5, 2)

the best known covering uses 43 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005090 | ljcr-c-29-6-2

Covering design C(29, 6, 2)

the best known covering uses 30 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005091 | ljcr-c-29-6-3

Covering design C(29, 6, 3)

the best known covering uses 217 blocks and the source-reported lower bound is 194.

来源状态 open | 2026-07-23 | 来源

V3P-005092 | ljcr-c-29-7-2

Covering design C(29, 7, 2)

the best known covering uses 23 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005093 | ljcr-c-29-7-3

Covering design C(29, 7, 3)

the best known covering uses 132 blocks and the source-reported lower bound is 116.

来源状态 open | 2026-07-23 | 来源

V3P-005094 | ljcr-c-29-8-3

Covering design C(29, 8, 3)

the best known covering uses 86 blocks and the source-reported lower bound is 77.

来源状态 open | 2026-07-23 | 来源

V3P-005095 | ljcr-c-29-9-3

Covering design C(29, 9, 3)

the best known covering uses 59 blocks and the source-reported lower bound is 55.

来源状态 open | 2026-07-23 | 来源

V3P-005096 | ljcr-c-30-10-3

Covering design C(30, 10, 3)

the best known covering uses 46 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005097 | ljcr-c-30-10-4

Covering design C(30, 10, 4)

the best known covering uses 209 blocks and the source-reported lower bound is 165.

来源状态 open | 2026-07-23 | 来源

V3P-005098 | ljcr-c-30-11-3

Covering design C(30, 11, 3)

the best known covering uses 37 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005099 | ljcr-c-30-11-4

Covering design C(30, 11, 4)

the best known covering uses 133 blocks and the source-reported lower bound is 112.

来源状态 open | 2026-07-23 | 来源

V3P-005100 | ljcr-c-30-12-3

Covering design C(30, 12, 3)

the best known covering uses 30 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005101 | ljcr-c-30-12-4

Covering design C(30, 12, 4)

the best known covering uses 96 blocks and the source-reported lower bound is 73.

来源状态 open | 2026-07-23 | 来源

V3P-005102 | ljcr-c-30-13-3

Covering design C(30, 13, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005103 | ljcr-c-30-13-4

Covering design C(30, 13, 4)

the best known covering uses 73 blocks and the source-reported lower bound is 58.

来源状态 open | 2026-07-23 | 来源

V3P-005104 | ljcr-c-30-13-5

Covering design C(30, 13, 5)

the best known covering uses 250 blocks and the source-reported lower bound is 157.

来源状态 open | 2026-07-23 | 来源

V3P-005105 | ljcr-c-30-14-4

Covering design C(30, 14, 4)

the best known covering uses 48 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-005106 | ljcr-c-30-14-5

Covering design C(30, 14, 5)

the best known covering uses 152 blocks and the source-reported lower bound is 101.

来源状态 open | 2026-07-23 | 来源

V3P-005107 | ljcr-c-30-15-3

Covering design C(30, 15, 3)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005108 | ljcr-c-30-15-4

Covering design C(30, 15, 4)

the best known covering uses 31 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-005109 | ljcr-c-30-15-5

Covering design C(30, 15, 5)

the best known covering uses 94 blocks and the source-reported lower bound is 68.

来源状态 open | 2026-07-23 | 来源

V3P-005110 | ljcr-c-30-16-3

Covering design C(30, 16, 3)

the best known covering uses 13 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005111 | ljcr-c-30-16-4

Covering design C(30, 16, 4)

the best known covering uses 29 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005112 | ljcr-c-30-16-5

Covering design C(30, 16, 5)

the best known covering uses 62 blocks and the source-reported lower bound is 53.

来源状态 open | 2026-07-23 | 来源

V3P-005113 | ljcr-c-30-17-4

Covering design C(30, 17, 4)

the best known covering uses 26 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005114 | ljcr-c-30-17-5

Covering design C(30, 17, 5)

the best known covering uses 55 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-005115 | ljcr-c-30-18-3

Covering design C(30, 18, 3)

the best known covering uses 10 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005116 | ljcr-c-30-18-4

Covering design C(30, 18, 4)

the best known covering uses 20 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005117 | ljcr-c-30-18-5

Covering design C(30, 18, 5)

the best known covering uses 42 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005118 | ljcr-c-30-19-4

Covering design C(30, 19, 4)

the best known covering uses 17 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005119 | ljcr-c-30-19-5

Covering design C(30, 19, 5)

the best known covering uses 35 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005120 | ljcr-c-30-20-4

Covering design C(30, 20, 4)

the best known covering uses 13 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-005121 | ljcr-c-30-20-5

Covering design C(30, 20, 5)

the best known covering uses 26 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-005122 | ljcr-c-30-21-5

Covering design C(30, 21, 5)

the best known covering uses 20 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005123 | ljcr-c-30-23-5

Covering design C(30, 23, 5)

the best known covering uses 11 blocks and the source-reported lower bound is 10.

来源状态 open | 2026-07-23 | 来源

V3P-005124 | ljcr-c-30-6-3

Covering design C(30, 6, 3)

the best known covering uses 225 blocks and the source-reported lower bound is 210.

来源状态 open | 2026-07-23 | 来源

V3P-005125 | ljcr-c-30-7-2

Covering design C(30, 7, 2)

the best known covering uses 25 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005126 | ljcr-c-30-7-3

Covering design C(30, 7, 3)

the best known covering uses 142 blocks and the source-reported lower bound is 125.

来源状态 open | 2026-07-23 | 来源

V3P-005127 | ljcr-c-30-8-3

Covering design C(30, 8, 3)

the best known covering uses 93 blocks and the source-reported lower bound is 83.

来源状态 open | 2026-07-23 | 来源

V3P-005128 | ljcr-c-30-9-3

Covering design C(30, 9, 3)

the best known covering uses 66 blocks and the source-reported lower bound is 60.

来源状态 open | 2026-07-23 | 来源

V3P-005129 | ljcr-c-31-10-3

Covering design C(31, 10, 3)

the best known covering uses 48 blocks and the source-reported lower bound is 47.

来源状态 open | 2026-07-23 | 来源

V3P-005130 | ljcr-c-31-11-3

Covering design C(31, 11, 3)

the best known covering uses 39 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-005131 | ljcr-c-31-11-4

Covering design C(31, 11, 4)

the best known covering uses 153 blocks and the source-reported lower bound is 119.

来源状态 open | 2026-07-23 | 来源

V3P-005132 | ljcr-c-31-12-3

Covering design C(31, 12, 3)

the best known covering uses 32 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005133 | ljcr-c-31-12-4

Covering design C(31, 12, 4)

the best known covering uses 110 blocks and the source-reported lower bound is 86.

来源状态 open | 2026-07-23 | 来源

V3P-005134 | ljcr-c-31-13-3

Covering design C(31, 13, 3)

the best known covering uses 26 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005135 | ljcr-c-31-13-4

Covering design C(31, 13, 4)

the best known covering uses 80 blocks and the source-reported lower bound is 60.

来源状态 open | 2026-07-23 | 来源

V3P-005136 | ljcr-c-31-14-3

Covering design C(31, 14, 3)

the best known covering uses 21 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-005137 | ljcr-c-31-14-4

Covering design C(31, 14, 4)

the best known covering uses 60 blocks and the source-reported lower bound is 47.

来源状态 open | 2026-07-23 | 来源

V3P-005138 | ljcr-c-31-14-5

Covering design C(31, 14, 5)

the best known covering uses 189 blocks and the source-reported lower bound is 129.

来源状态 open | 2026-07-23 | 来源

V3P-005139 | ljcr-c-31-15-5

Covering design C(31, 15, 5)

the best known covering uses 113 blocks and the source-reported lower bound is 77.

来源状态 open | 2026-07-23 | 来源

V3P-005140 | ljcr-c-31-16-4

Covering design C(31, 16, 4)

the best known covering uses 30 blocks and the source-reported lower bound is 28.

来源状态 open | 2026-07-23 | 来源

V3P-005141 | ljcr-c-31-16-5

Covering design C(31, 16, 5)

the best known covering uses 62 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-005142 | ljcr-c-31-17-3

Covering design C(31, 17, 3)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-005143 | ljcr-c-31-17-4

Covering design C(31, 17, 4)

the best known covering uses 28 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005144 | ljcr-c-31-17-5

Covering design C(31, 17, 5)

the best known covering uses 58 blocks and the source-reported lower bound is 50.

来源状态 open | 2026-07-23 | 来源

V3P-005145 | ljcr-c-31-18-3

Covering design C(31, 18, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005146 | ljcr-c-31-18-4

Covering design C(31, 18, 4)

the best known covering uses 23 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-005147 | ljcr-c-31-18-5

Covering design C(31, 18, 5)

the best known covering uses 52 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-005148 | ljcr-c-31-19-4

Covering design C(31, 19, 4)

the best known covering uses 19 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005149 | ljcr-c-31-19-5

Covering design C(31, 19, 5)

the best known covering uses 40 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005150 | ljcr-c-31-20-4

Covering design C(31, 20, 4)

the best known covering uses 16 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-005151 | ljcr-c-31-20-5

Covering design C(31, 20, 5)

the best known covering uses 31 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005152 | ljcr-c-31-21-4

Covering design C(31, 21, 4)

the best known covering uses 12 blocks and the source-reported lower bound is 10.

来源状态 open | 2026-07-23 | 来源

V3P-005153 | ljcr-c-31-21-5

Covering design C(31, 21, 5)

the best known covering uses 23 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005154 | ljcr-c-31-22-5

Covering design C(31, 22, 5)

the best known covering uses 18 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005155 | ljcr-c-31-23-5

Covering design C(31, 23, 5)

the best known covering uses 13 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-005156 | ljcr-c-31-24-5

Covering design C(31, 24, 5)

the best known covering uses 11 blocks and the source-reported lower bound is 10.

来源状态 open | 2026-07-23 | 来源

V3P-005157 | ljcr-c-31-7-3

Covering design C(31, 7, 3)

the best known covering uses 153 blocks and the source-reported lower bound is 138.

来源状态 open | 2026-07-23 | 来源

V3P-005158 | ljcr-c-31-8-3

Covering design C(31, 8, 3)

the best known covering uses 104 blocks and the source-reported lower bound is 93.

来源状态 open | 2026-07-23 | 来源

V3P-005159 | ljcr-c-31-9-2

Covering design C(31, 9, 2)

the best known covering uses 17 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005160 | ljcr-c-31-9-3

Covering design C(31, 9, 3)

the best known covering uses 73 blocks and the source-reported lower bound is 66.

来源状态 open | 2026-07-23 | 来源

V3P-005161 | ljcr-c-32-10-3

Covering design C(32, 10, 3)

the best known covering uses 60 blocks and the source-reported lower bound is 52.

来源状态 open | 2026-07-23 | 来源

V3P-005162 | ljcr-c-32-11-3

Covering design C(32, 11, 3)

the best known covering uses 40 blocks and the source-reported lower bound is 38.

来源状态 open | 2026-07-23 | 来源

V3P-005163 | ljcr-c-32-11-4

Covering design C(32, 11, 4)

the best known covering uses 176 blocks and the source-reported lower bound is 137.

来源状态 open | 2026-07-23 | 来源

V3P-005164 | ljcr-c-32-12-4

Covering design C(32, 12, 4)

the best known covering uses 121 blocks and the source-reported lower bound is 99.

来源状态 open | 2026-07-23 | 来源

V3P-005165 | ljcr-c-32-13-3

Covering design C(32, 13, 3)

the best known covering uses 29 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005166 | ljcr-c-32-13-4

Covering design C(32, 13, 4)

the best known covering uses 96 blocks and the source-reported lower bound is 72.

来源状态 open | 2026-07-23 | 来源

V3P-005167 | ljcr-c-32-14-3

Covering design C(32, 14, 3)

the best known covering uses 22 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-005168 | ljcr-c-32-14-4

Covering design C(32, 14, 4)

the best known covering uses 67 blocks and the source-reported lower bound is 51.

来源状态 open | 2026-07-23 | 来源

V3P-005169 | ljcr-c-32-14-5

Covering design C(32, 14, 5)

the best known covering uses 232 blocks and the source-reported lower bound is 138.

来源状态 open | 2026-07-23 | 来源

V3P-005170 | ljcr-c-32-15-3

Covering design C(32, 15, 3)

the best known covering uses 18 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005171 | ljcr-c-32-15-4

Covering design C(32, 15, 4)

the best known covering uses 49 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-005172 | ljcr-c-32-15-5

Covering design C(32, 15, 5)

the best known covering uses 153 blocks and the source-reported lower bound is 101.

来源状态 open | 2026-07-23 | 来源

V3P-005173 | ljcr-c-32-17-3

Covering design C(32, 17, 3)

the best known covering uses 14 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005174 | ljcr-c-32-17-4

Covering design C(32, 17, 4)

the best known covering uses 29 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005175 | ljcr-c-32-17-5

Covering design C(32, 17, 5)

the best known covering uses 62 blocks and the source-reported lower bound is 53.

来源状态 open | 2026-07-23 | 来源

V3P-005176 | ljcr-c-32-18-4

Covering design C(32, 18, 4)

the best known covering uses 26 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005177 | ljcr-c-32-18-5

Covering design C(32, 18, 5)

the best known covering uses 55 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-005178 | ljcr-c-32-19-3

Covering design C(32, 19, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005179 | ljcr-c-32-19-4

Covering design C(32, 19, 4)

the best known covering uses 22 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005180 | ljcr-c-32-19-5

Covering design C(32, 19, 5)

the best known covering uses 46 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-005181 | ljcr-c-32-20-4

Covering design C(32, 20, 4)

the best known covering uses 18 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005182 | ljcr-c-32-20-5

Covering design C(32, 20, 5)

the best known covering uses 34 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005183 | ljcr-c-32-21-4

Covering design C(32, 21, 4)

the best known covering uses 15 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-005184 | ljcr-c-32-21-5

Covering design C(32, 21, 5)

the best known covering uses 29 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-005185 | ljcr-c-32-22-4

Covering design C(32, 22, 4)

the best known covering uses 11 blocks and the source-reported lower bound is 10.

来源状态 open | 2026-07-23 | 来源

V3P-005186 | ljcr-c-32-22-5

Covering design C(32, 22, 5)

the best known covering uses 20 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005187 | ljcr-c-32-23-5

Covering design C(32, 23, 5)

the best known covering uses 16 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005188 | ljcr-c-32-24-5

Covering design C(32, 24, 5)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-005189 | ljcr-c-32-7-3

Covering design C(32, 7, 3)

the best known covering uses 165 blocks and the source-reported lower bound is 143.

来源状态 open | 2026-07-23 | 来源

V3P-005190 | ljcr-c-32-8-3

Covering design C(32, 8, 3)

the best known covering uses 106 blocks and the source-reported lower bound is 104.

来源状态 open | 2026-07-23 | 来源

V3P-005191 | ljcr-c-32-9-2

Covering design C(32, 9, 2)

the best known covering uses 18 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005192 | ljcr-c-32-9-3

Covering design C(32, 9, 3)

the best known covering uses 77 blocks and the source-reported lower bound is 72.

来源状态 open | 2026-07-23 | 来源

V3P-005193 | ljcr-c-33-10-2

Covering design C(33, 10, 2)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005194 | ljcr-c-33-10-3

Covering design C(33, 10, 3)

the best known covering uses 66 blocks and the source-reported lower bound is 57.

来源状态 open | 2026-07-23 | 来源

V3P-005195 | ljcr-c-33-11-3

Covering design C(33, 11, 3)

the best known covering uses 46 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005196 | ljcr-c-33-11-4

Covering design C(33, 11, 4)

the best known covering uses 216 blocks and the source-reported lower bound is 156.

来源状态 open | 2026-07-23 | 来源

V3P-005197 | ljcr-c-33-12-3

Covering design C(33, 12, 3)

the best known covering uses 36 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005198 | ljcr-c-33-12-4

Covering design C(33, 12, 4)

the best known covering uses 134 blocks and the source-reported lower bound is 105.

来源状态 open | 2026-07-23 | 来源

V3P-005199 | ljcr-c-33-13-3

Covering design C(33, 13, 3)

the best known covering uses 31 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-005200 | ljcr-c-33-13-4

Covering design C(33, 13, 4)

the best known covering uses 102 blocks and the source-reported lower bound is 82.

来源状态 open | 2026-07-23 | 来源

V3P-005201 | ljcr-c-33-14-3

Covering design C(33, 14, 3)

the best known covering uses 26 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005202 | ljcr-c-33-14-4

Covering design C(33, 14, 4)

the best known covering uses 78 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-005203 | ljcr-c-33-15-3

Covering design C(33, 15, 3)

the best known covering uses 20 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005204 | ljcr-c-33-15-4

Covering design C(33, 15, 4)

the best known covering uses 57 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005205 | ljcr-c-33-15-5

Covering design C(33, 15, 5)

the best known covering uses 180 blocks and the source-reported lower bound is 113.

来源状态 open | 2026-07-23 | 来源

V3P-005206 | ljcr-c-33-16-3

Covering design C(33, 16, 3)

the best known covering uses 16 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005207 | ljcr-c-33-16-4

Covering design C(33, 16, 4)

the best known covering uses 39 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-005208 | ljcr-c-33-16-5

Covering design C(33, 16, 5)

the best known covering uses 111 blocks and the source-reported lower bound is 81.

来源状态 open | 2026-07-23 | 来源

V3P-005209 | ljcr-c-33-17-4

Covering design C(33, 17, 4)

the best known covering uses 30 blocks and the source-reported lower bound is 28.

来源状态 open | 2026-07-23 | 来源

V3P-005210 | ljcr-c-33-17-5

Covering design C(33, 17, 5)

the best known covering uses 62 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-005211 | ljcr-c-33-18-4

Covering design C(33, 18, 4)

the best known covering uses 28 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005212 | ljcr-c-33-18-5

Covering design C(33, 18, 5)

the best known covering uses 59 blocks and the source-reported lower bound is 50.

来源状态 open | 2026-07-23 | 来源

V3P-005213 | ljcr-c-33-19-3

Covering design C(33, 19, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005214 | ljcr-c-33-19-4

Covering design C(33, 19, 4)

the best known covering uses 24 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-005215 | ljcr-c-33-19-5

Covering design C(33, 19, 5)

the best known covering uses 51 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-005216 | ljcr-c-33-20-3

Covering design C(33, 20, 3)

the best known covering uses 10 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005217 | ljcr-c-33-20-4

Covering design C(33, 20, 4)

the best known covering uses 20 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005218 | ljcr-c-33-20-5

Covering design C(33, 20, 5)

the best known covering uses 43 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005219 | ljcr-c-33-21-4

Covering design C(33, 21, 4)

the best known covering uses 17 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-005220 | ljcr-c-33-21-5

Covering design C(33, 21, 5)

the best known covering uses 33 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005221 | ljcr-c-33-22-4

Covering design C(33, 22, 4)

the best known covering uses 13 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-005222 | ljcr-c-33-22-5

Covering design C(33, 22, 5)

the best known covering uses 27 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-005223 | ljcr-c-33-23-5

Covering design C(33, 23, 5)

the best known covering uses 19 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005224 | ljcr-c-33-24-5

Covering design C(33, 24, 5)

the best known covering uses 15 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-005225 | ljcr-c-33-25-5

Covering design C(33, 25, 5)

the best known covering uses 12 blocks and the source-reported lower bound is 10.

来源状态 open | 2026-07-23 | 来源

V3P-005226 | ljcr-c-33-7-2

Covering design C(33, 7, 2)

the best known covering uses 30 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005227 | ljcr-c-33-7-3

Covering design C(33, 7, 3)

the best known covering uses 191 blocks and the source-reported lower bound is 180.

来源状态 open | 2026-07-23 | 来源

V3P-005228 | ljcr-c-33-8-2

Covering design C(33, 8, 2)

the best known covering uses 24 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005229 | ljcr-c-33-8-3

Covering design C(33, 8, 3)

the best known covering uses 127 blocks and the source-reported lower bound is 116.

来源状态 open | 2026-07-23 | 来源

V3P-005230 | ljcr-c-33-9-3

Covering design C(33, 9, 3)

the best known covering uses 80 blocks and the source-reported lower bound is 74.

来源状态 open | 2026-07-23 | 来源

V3P-005231 | ljcr-c-34-10-2

Covering design C(34, 10, 2)

the best known covering uses 16 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005232 | ljcr-c-34-10-3

Covering design C(34, 10, 3)

the best known covering uses 68 blocks and the source-reported lower bound is 62.

来源状态 open | 2026-07-23 | 来源

V3P-005233 | ljcr-c-34-11-3

Covering design C(34, 11, 3)

the best known covering uses 51 blocks and the source-reported lower bound is 44.

来源状态 open | 2026-07-23 | 来源

V3P-005234 | ljcr-c-34-11-4

Covering design C(34, 11, 4)

the best known covering uses 249 blocks and the source-reported lower bound is 177.

来源状态 open | 2026-07-23 | 来源

V3P-005235 | ljcr-c-34-12-3

Covering design C(34, 12, 3)

the best known covering uses 39 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-005236 | ljcr-c-34-12-4

Covering design C(34, 12, 4)

the best known covering uses 152 blocks and the source-reported lower bound is 119.

来源状态 open | 2026-07-23 | 来源

V3P-005237 | ljcr-c-34-13-3

Covering design C(34, 13, 3)

the best known covering uses 33 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005238 | ljcr-c-34-13-4

Covering design C(34, 13, 4)

the best known covering uses 109 blocks and the source-reported lower bound is 87.

来源状态 open | 2026-07-23 | 来源

V3P-005239 | ljcr-c-34-14-3

Covering design C(34, 14, 3)

the best known covering uses 27 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005240 | ljcr-c-34-14-4

Covering design C(34, 14, 4)

the best known covering uses 90 blocks and the source-reported lower bound is 64.

来源状态 open | 2026-07-23 | 来源

V3P-005241 | ljcr-c-34-15-3

Covering design C(34, 15, 3)

the best known covering uses 23 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-005242 | ljcr-c-34-15-4

Covering design C(34, 15, 4)

the best known covering uses 63 blocks and the source-reported lower bound is 50.

来源状态 open | 2026-07-23 | 来源

V3P-005243 | ljcr-c-34-15-5

Covering design C(34, 15, 5)

the best known covering uses 215 blocks and the source-reported lower bound is 134.

来源状态 open | 2026-07-23 | 来源

V3P-005244 | ljcr-c-34-16-3

Covering design C(34, 16, 3)

the best known covering uses 17 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005245 | ljcr-c-34-16-4

Covering design C(34, 16, 4)

the best known covering uses 45 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-005246 | ljcr-c-34-16-5

Covering design C(34, 16, 5)

the best known covering uses 129 blocks and the source-reported lower bound is 90.

来源状态 open | 2026-07-23 | 来源

V3P-005247 | ljcr-c-34-17-3

Covering design C(34, 17, 3)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005248 | ljcr-c-34-17-4

Covering design C(34, 17, 4)

the best known covering uses 34 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-005249 | ljcr-c-34-17-5

Covering design C(34, 17, 5)

the best known covering uses 80 blocks and the source-reported lower bound is 64.

来源状态 open | 2026-07-23 | 来源

V3P-005250 | ljcr-c-34-18-3

Covering design C(34, 18, 3)

the best known covering uses 13 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005251 | ljcr-c-34-18-4

Covering design C(34, 18, 4)

the best known covering uses 28 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005252 | ljcr-c-34-18-5

Covering design C(34, 18, 5)

the best known covering uses 60 blocks and the source-reported lower bound is 53.

来源状态 open | 2026-07-23 | 来源

V3P-005253 | ljcr-c-34-19-4

Covering design C(34, 19, 4)

the best known covering uses 27 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005254 | ljcr-c-34-19-5

Covering design C(34, 19, 5)

the best known covering uses 56 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-005255 | ljcr-c-34-20-3

Covering design C(34, 20, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005256 | ljcr-c-34-20-4

Covering design C(34, 20, 4)

the best known covering uses 23 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005257 | ljcr-c-34-20-5

Covering design C(34, 20, 5)

the best known covering uses 45 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005258 | ljcr-c-34-21-4

Covering design C(34, 21, 4)

the best known covering uses 19 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005259 | ljcr-c-34-21-5

Covering design C(34, 21, 5)

the best known covering uses 40 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005260 | ljcr-c-34-22-4

Covering design C(34, 22, 4)

the best known covering uses 15 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-005261 | ljcr-c-34-22-5

Covering design C(34, 22, 5)

the best known covering uses 30 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005262 | ljcr-c-34-23-4

Covering design C(34, 23, 4)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-005263 | ljcr-c-34-23-5

Covering design C(34, 23, 5)

the best known covering uses 24 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005264 | ljcr-c-34-24-5

Covering design C(34, 24, 5)

the best known covering uses 17 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005265 | ljcr-c-34-25-4

Covering design C(34, 25, 4)

the best known covering uses 9 blocks and the source-reported lower bound is 8.

来源状态 open | 2026-07-23 | 来源

V3P-005266 | ljcr-c-34-25-5

Covering design C(34, 25, 5)

the best known covering uses 14 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-005267 | ljcr-c-34-7-2

Covering design C(34, 7, 2)

the best known covering uses 31 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-005268 | ljcr-c-34-7-3

Covering design C(34, 7, 3)

the best known covering uses 209 blocks and the source-reported lower bound is 190.

来源状态 open | 2026-07-23 | 来源

V3P-005269 | ljcr-c-34-8-2

Covering design C(34, 8, 2)

the best known covering uses 25 blocks and the source-reported lower bound is 23.

来源状态 open | 2026-07-23 | 来源

V3P-005270 | ljcr-c-34-8-3

Covering design C(34, 8, 3)

the best known covering uses 139 blocks and the source-reported lower bound is 124.

来源状态 open | 2026-07-23 | 来源

V3P-005271 | ljcr-c-34-9-3

Covering design C(34, 9, 3)

the best known covering uses 95 blocks and the source-reported lower bound is 84.

来源状态 open | 2026-07-23 | 来源

V3P-005272 | ljcr-c-35-10-3

Covering design C(35, 10, 3)

the best known covering uses 77 blocks and the source-reported lower bound is 67.

来源状态 open | 2026-07-23 | 来源

V3P-005273 | ljcr-c-35-11-3

Covering design C(35, 11, 3)

the best known covering uses 56 blocks and the source-reported lower bound is 48.

来源状态 open | 2026-07-23 | 来源

V3P-005274 | ljcr-c-35-12-3

Covering design C(35, 12, 3)

the best known covering uses 40 blocks and the source-reported lower bound is 38.

来源状态 open | 2026-07-23 | 来源

V3P-005275 | ljcr-c-35-12-4

Covering design C(35, 12, 4)

the best known covering uses 169 blocks and the source-reported lower bound is 129.

来源状态 open | 2026-07-23 | 来源

V3P-005276 | ljcr-c-35-13-3

Covering design C(35, 13, 3)

the best known covering uses 36 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005277 | ljcr-c-35-13-4

Covering design C(35, 13, 4)

the best known covering uses 124 blocks and the source-reported lower bound is 100.

来源状态 open | 2026-07-23 | 来源

V3P-005278 | ljcr-c-35-14-3

Covering design C(35, 14, 3)

the best known covering uses 30 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005279 | ljcr-c-35-14-4

Covering design C(35, 14, 4)

the best known covering uses 98 blocks and the source-reported lower bound is 73.

来源状态 open | 2026-07-23 | 来源

V3P-005280 | ljcr-c-35-15-3

Covering design C(35, 15, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005281 | ljcr-c-35-15-4

Covering design C(35, 15, 4)

the best known covering uses 70 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-005282 | ljcr-c-35-15-5

Covering design C(35, 15, 5)

the best known covering uses 240 blocks and the source-reported lower bound is 150.

来源状态 open | 2026-07-23 | 来源

V3P-005283 | ljcr-c-35-16-3

Covering design C(35, 16, 3)

the best known covering uses 20 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005284 | ljcr-c-35-16-4

Covering design C(35, 16, 4)

the best known covering uses 57 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005285 | ljcr-c-35-16-5

Covering design C(35, 16, 5)

the best known covering uses 158 blocks and the source-reported lower bound is 110.

来源状态 open | 2026-07-23 | 来源

V3P-005286 | ljcr-c-35-17-4

Covering design C(35, 17, 4)

the best known covering uses 38 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-005287 | ljcr-c-35-17-5

Covering design C(35, 17, 5)

the best known covering uses 101 blocks and the source-reported lower bound is 77.

来源状态 open | 2026-07-23 | 来源

V3P-005288 | ljcr-c-35-18-4

Covering design C(35, 18, 4)

the best known covering uses 31 blocks and the source-reported lower bound is 28.

来源状态 open | 2026-07-23 | 来源

V3P-005289 | ljcr-c-35-18-5

Covering design C(35, 18, 5)

the best known covering uses 69 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-005290 | ljcr-c-35-19-3

Covering design C(35, 19, 3)

the best known covering uses 13 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005291 | ljcr-c-35-19-4

Covering design C(35, 19, 4)

the best known covering uses 28 blocks and the source-reported lower bound is 23.

来源状态 open | 2026-07-23 | 来源

V3P-005292 | ljcr-c-35-19-5

Covering design C(35, 19, 5)

the best known covering uses 58 blocks and the source-reported lower bound is 50.

来源状态 open | 2026-07-23 | 来源

V3P-005293 | ljcr-c-35-20-3

Covering design C(35, 20, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005294 | ljcr-c-35-20-4

Covering design C(35, 20, 4)

the best known covering uses 25 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-005295 | ljcr-c-35-20-5

Covering design C(35, 20, 5)

the best known covering uses 52 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-005296 | ljcr-c-35-21-3

Covering design C(35, 21, 3)

the best known covering uses 10 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005297 | ljcr-c-35-21-4

Covering design C(35, 21, 4)

the best known covering uses 21 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005298 | ljcr-c-35-21-5

Covering design C(35, 21, 5)

the best known covering uses 43 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005299 | ljcr-c-35-22-4

Covering design C(35, 22, 4)

the best known covering uses 18 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005300 | ljcr-c-35-22-5

Covering design C(35, 22, 5)

the best known covering uses 35 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005301 | ljcr-c-35-23-4

Covering design C(35, 23, 4)

the best known covering uses 14 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-005302 | ljcr-c-35-23-5

Covering design C(35, 23, 5)

the best known covering uses 28 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-005303 | ljcr-c-35-24-4

Covering design C(35, 24, 4)

the best known covering uses 11 blocks and the source-reported lower bound is 10.

来源状态 open | 2026-07-23 | 来源

V3P-005304 | ljcr-c-35-24-5

Covering design C(35, 24, 5)

the best known covering uses 22 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005305 | ljcr-c-35-25-5

Covering design C(35, 25, 5)

the best known covering uses 16 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005306 | ljcr-c-35-6-2

Covering design C(35, 6, 2)

the best known covering uses 42 blocks and the source-reported lower bound is 41.

来源状态 open | 2026-07-23 | 来源

V3P-005307 | ljcr-c-35-7-2

Covering design C(35, 7, 2)

the best known covering uses 32 blocks and the source-reported lower bound is 31.

来源状态 open | 2026-07-23 | 来源

V3P-005308 | ljcr-c-35-7-3

Covering design C(35, 7, 3)

the best known covering uses 233 blocks and the source-reported lower bound is 200.

来源状态 open | 2026-07-23 | 来源

V3P-005309 | ljcr-c-35-8-2

Covering design C(35, 8, 2)

the best known covering uses 26 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005310 | ljcr-c-35-8-3

Covering design C(35, 8, 3)

the best known covering uses 149 blocks and the source-reported lower bound is 132.

来源状态 open | 2026-07-23 | 来源

V3P-005311 | ljcr-c-35-9-3

Covering design C(35, 9, 3)

the best known covering uses 103 blocks and the source-reported lower bound is 90.

来源状态 open | 2026-07-23 | 来源

V3P-005312 | ljcr-c-36-10-2

Covering design C(36, 10, 2)

the best known covering uses 18 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005313 | ljcr-c-36-10-3

Covering design C(36, 10, 3)

the best known covering uses 81 blocks and the source-reported lower bound is 72.

来源状态 open | 2026-07-23 | 来源

V3P-005314 | ljcr-c-36-11-2

Covering design C(36, 11, 2)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005315 | ljcr-c-36-11-3

Covering design C(36, 11, 3)

the best known covering uses 64 blocks and the source-reported lower bound is 56.

来源状态 open | 2026-07-23 | 来源

V3P-005316 | ljcr-c-36-12-3

Covering design C(36, 12, 3)

the best known covering uses 40 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-005317 | ljcr-c-36-12-4

Covering design C(36, 12, 4)

the best known covering uses 180 blocks and the source-reported lower bound is 144.

来源状态 open | 2026-07-23 | 来源

V3P-005318 | ljcr-c-36-13-3

Covering design C(36, 13, 3)

the best known covering uses 37 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005319 | ljcr-c-36-13-4

Covering design C(36, 13, 4)

the best known covering uses 135 blocks and the source-reported lower bound is 106.

来源状态 open | 2026-07-23 | 来源

V3P-005320 | ljcr-c-36-14-3

Covering design C(36, 14, 3)

the best known covering uses 31 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005321 | ljcr-c-36-14-4

Covering design C(36, 14, 4)

the best known covering uses 104 blocks and the source-reported lower bound is 85.

来源状态 open | 2026-07-23 | 来源

V3P-005322 | ljcr-c-36-15-3

Covering design C(36, 15, 3)

the best known covering uses 26 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005323 | ljcr-c-36-15-4

Covering design C(36, 15, 4)

the best known covering uses 77 blocks and the source-reported lower bound is 60.

来源状态 open | 2026-07-23 | 来源

V3P-005324 | ljcr-c-36-16-3

Covering design C(36, 16, 3)

the best known covering uses 21 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-005325 | ljcr-c-36-16-4

Covering design C(36, 16, 4)

the best known covering uses 62 blocks and the source-reported lower bound is 48.

来源状态 open | 2026-07-23 | 来源

V3P-005326 | ljcr-c-36-16-5

Covering design C(36, 16, 5)

the best known covering uses 182 blocks and the source-reported lower bound is 133.

来源状态 open | 2026-07-23 | 来源

V3P-005327 | ljcr-c-36-17-3

Covering design C(36, 17, 3)

the best known covering uses 17 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005328 | ljcr-c-36-17-4

Covering design C(36, 17, 4)

the best known covering uses 47 blocks and the source-reported lower bound is 36.

来源状态 open | 2026-07-23 | 来源

V3P-005329 | ljcr-c-36-17-5

Covering design C(36, 17, 5)

the best known covering uses 138 blocks and the source-reported lower bound is 89.

来源状态 open | 2026-07-23 | 来源

V3P-005330 | ljcr-c-36-18-3

Covering design C(36, 18, 3)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005331 | ljcr-c-36-18-4

Covering design C(36, 18, 4)

the best known covering uses 33 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-005332 | ljcr-c-36-18-5

Covering design C(36, 18, 5)

the best known covering uses 80 blocks and the source-reported lower bound is 64.

来源状态 open | 2026-07-23 | 来源

V3P-005333 | ljcr-c-36-19-3

Covering design C(36, 19, 3)

the best known covering uses 14 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005334 | ljcr-c-36-19-4

Covering design C(36, 19, 4)

the best known covering uses 30 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005335 | ljcr-c-36-19-5

Covering design C(36, 19, 5)

the best known covering uses 64 blocks and the source-reported lower bound is 54.

来源状态 open | 2026-07-23 | 来源

V3P-005336 | ljcr-c-36-20-3

Covering design C(36, 20, 3)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-005337 | ljcr-c-36-20-4

Covering design C(36, 20, 4)

the best known covering uses 26 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005338 | ljcr-c-36-20-5

Covering design C(36, 20, 5)

the best known covering uses 54 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005339 | ljcr-c-36-21-3

Covering design C(36, 21, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005340 | ljcr-c-36-21-4

Covering design C(36, 21, 4)

the best known covering uses 23 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005341 | ljcr-c-36-21-5

Covering design C(36, 21, 5)

the best known covering uses 49 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-005342 | ljcr-c-36-22-3

Covering design C(36, 22, 3)

the best known covering uses 10 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005343 | ljcr-c-36-22-4

Covering design C(36, 22, 4)

the best known covering uses 19 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005344 | ljcr-c-36-22-5

Covering design C(36, 22, 5)

the best known covering uses 38 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-005345 | ljcr-c-36-23-4

Covering design C(36, 23, 4)

the best known covering uses 16 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005346 | ljcr-c-36-23-5

Covering design C(36, 23, 5)

the best known covering uses 33 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005347 | ljcr-c-36-24-4

Covering design C(36, 24, 4)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-005348 | ljcr-c-36-24-5

Covering design C(36, 24, 5)

the best known covering uses 24 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005349 | ljcr-c-36-25-4

Covering design C(36, 25, 4)

the best known covering uses 11 blocks and the source-reported lower bound is 10.

来源状态 open | 2026-07-23 | 来源

V3P-005350 | ljcr-c-36-25-5

Covering design C(36, 25, 5)

the best known covering uses 20 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005351 | ljcr-c-36-5-2

Covering design C(36, 5, 2)

the best known covering uses 66 blocks and the source-reported lower bound is 65.

来源状态 open | 2026-07-23 | 来源

V3P-005352 | ljcr-c-36-6-2

Covering design C(36, 6, 2)

the best known covering uses 47 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005353 | ljcr-c-36-7-2

Covering design C(36, 7, 2)

the best known covering uses 33 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-005354 | ljcr-c-36-8-2

Covering design C(36, 8, 2)

the best known covering uses 27 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005355 | ljcr-c-36-8-3

Covering design C(36, 8, 3)

the best known covering uses 162 blocks and the source-reported lower bound is 140.

来源状态 open | 2026-07-23 | 来源

V3P-005356 | ljcr-c-36-9-2

Covering design C(36, 9, 2)

the best known covering uses 21 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-005357 | ljcr-c-36-9-3

Covering design C(36, 9, 3)

the best known covering uses 112 blocks and the source-reported lower bound is 96.

来源状态 open | 2026-07-23 | 来源

V3P-005358 | ljcr-c-37-10-2

Covering design C(37, 10, 2)

the best known covering uses 19 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-005359 | ljcr-c-37-10-3

Covering design C(37, 10, 3)

the best known covering uses 84 blocks and the source-reported lower bound is 74.

来源状态 open | 2026-07-23 | 来源

V3P-005360 | ljcr-c-37-11-2

Covering design C(37, 11, 2)

the best known covering uses 16 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005361 | ljcr-c-37-11-3

Covering design C(37, 11, 3)

the best known covering uses 69 blocks and the source-reported lower bound is 58.

来源状态 open | 2026-07-23 | 来源

V3P-005362 | ljcr-c-37-12-3

Covering design C(37, 12, 3)

the best known covering uses 51 blocks and the source-reported lower bound is 44.

来源状态 open | 2026-07-23 | 来源

V3P-005363 | ljcr-c-37-12-4

Covering design C(37, 12, 4)

the best known covering uses 238 blocks and the source-reported lower bound is 173.

来源状态 open | 2026-07-23 | 来源

V3P-005364 | ljcr-c-37-13-3

Covering design C(37, 13, 3)

the best known covering uses 38 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-005365 | ljcr-c-37-13-4

Covering design C(37, 13, 4)

the best known covering uses 145 blocks and the source-reported lower bound is 111.

来源状态 open | 2026-07-23 | 来源

V3P-005366 | ljcr-c-37-14-3

Covering design C(37, 14, 3)

the best known covering uses 34 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-005367 | ljcr-c-37-14-4

Covering design C(37, 14, 4)

the best known covering uses 109 blocks and the source-reported lower bound is 90.

来源状态 open | 2026-07-23 | 来源

V3P-005368 | ljcr-c-37-15-3

Covering design C(37, 15, 3)

the best known covering uses 28 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005369 | ljcr-c-37-15-4

Covering design C(37, 15, 4)

the best known covering uses 93 blocks and the source-reported lower bound is 72.

来源状态 open | 2026-07-23 | 来源

V3P-005370 | ljcr-c-37-16-3

Covering design C(37, 16, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005371 | ljcr-c-37-16-4

Covering design C(37, 16, 4)

the best known covering uses 70 blocks and the source-reported lower bound is 56.

来源状态 open | 2026-07-23 | 来源

V3P-005372 | ljcr-c-37-16-5

Covering design C(37, 16, 5)

the best known covering uses 207 blocks and the source-reported lower bound is 139.

来源状态 open | 2026-07-23 | 来源

V3P-005373 | ljcr-c-37-17-3

Covering design C(37, 17, 3)

the best known covering uses 18 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005374 | ljcr-c-37-17-4

Covering design C(37, 17, 4)

the best known covering uses 51 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-005375 | ljcr-c-37-17-5

Covering design C(37, 17, 5)

the best known covering uses 164 blocks and the source-reported lower bound is 105.

来源状态 open | 2026-07-23 | 来源

V3P-005376 | ljcr-c-37-18-4

Covering design C(37, 18, 4)

the best known covering uses 39 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-005377 | ljcr-c-37-18-5

Covering design C(37, 18, 5)

the best known covering uses 105 blocks and the source-reported lower bound is 74.

来源状态 open | 2026-07-23 | 来源

V3P-005378 | ljcr-c-37-19-4

Covering design C(37, 19, 4)

the best known covering uses 31 blocks and the source-reported lower bound is 28.

来源状态 open | 2026-07-23 | 来源

V3P-005379 | ljcr-c-37-19-5

Covering design C(37, 19, 5)

the best known covering uses 68 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-005380 | ljcr-c-37-20-3

Covering design C(37, 20, 3)

the best known covering uses 13 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005381 | ljcr-c-37-20-4

Covering design C(37, 20, 4)

the best known covering uses 29 blocks and the source-reported lower bound is 23.

来源状态 open | 2026-07-23 | 来源

V3P-005382 | ljcr-c-37-20-5

Covering design C(37, 20, 5)

the best known covering uses 61 blocks and the source-reported lower bound is 50.

来源状态 open | 2026-07-23 | 来源

V3P-005383 | ljcr-c-37-21-3

Covering design C(37, 21, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005384 | ljcr-c-37-21-4

Covering design C(37, 21, 4)

the best known covering uses 25 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005385 | ljcr-c-37-21-5

Covering design C(37, 21, 5)

the best known covering uses 53 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-005386 | ljcr-c-37-22-3

Covering design C(37, 22, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005387 | ljcr-c-37-22-4

Covering design C(37, 22, 4)

the best known covering uses 22 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005388 | ljcr-c-37-22-5

Covering design C(37, 22, 5)

the best known covering uses 44 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005389 | ljcr-c-37-23-3

Covering design C(37, 23, 3)

the best known covering uses 9 blocks and the source-reported lower bound is 8.

来源状态 open | 2026-07-23 | 来源

V3P-005390 | ljcr-c-37-23-4

Covering design C(37, 23, 4)

the best known covering uses 18 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005391 | ljcr-c-37-23-5

Covering design C(37, 23, 5)

the best known covering uses 36 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005392 | ljcr-c-37-24-4

Covering design C(37, 24, 4)

the best known covering uses 16 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-005393 | ljcr-c-37-24-5

Covering design C(37, 24, 5)

the best known covering uses 31 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005394 | ljcr-c-37-25-5

Covering design C(37, 25, 5)

the best known covering uses 22 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005395 | ljcr-c-37-5-2

Covering design C(37, 5, 2)

the best known covering uses 68 blocks and the source-reported lower bound is 67.

来源状态 open | 2026-07-23 | 来源

V3P-005396 | ljcr-c-37-7-2

Covering design C(37, 7, 2)

the best known covering uses 35 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005397 | ljcr-c-37-8-3

Covering design C(37, 8, 3)

the best known covering uses 177 blocks and the source-reported lower bound is 148.

来源状态 open | 2026-07-23 | 来源

V3P-005398 | ljcr-c-37-9-3

Covering design C(37, 9, 3)

the best known covering uses 122 blocks and the source-reported lower bound is 103.

来源状态 open | 2026-07-23 | 来源

V3P-005399 | ljcr-c-38-10-3

Covering design C(38, 10, 3)

the best known covering uses 89 blocks and the source-reported lower bound is 80.

来源状态 open | 2026-07-23 | 来源

V3P-005400 | ljcr-c-38-11-2

Covering design C(38, 11, 2)

the best known covering uses 17 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005401 | ljcr-c-38-11-3

Covering design C(38, 11, 3)

the best known covering uses 74 blocks and the source-reported lower bound is 63.

来源状态 open | 2026-07-23 | 来源

V3P-005402 | ljcr-c-38-12-3

Covering design C(38, 12, 3)

the best known covering uses 54 blocks and the source-reported lower bound is 45.

来源状态 open | 2026-07-23 | 来源

V3P-005403 | ljcr-c-38-13-3

Covering design C(38, 13, 3)

the best known covering uses 40 blocks and the source-reported lower bound is 38.

来源状态 open | 2026-07-23 | 来源

V3P-005404 | ljcr-c-38-13-4

Covering design C(38, 13, 4)

the best known covering uses 176 blocks and the source-reported lower bound is 129.

来源状态 open | 2026-07-23 | 来源

V3P-005405 | ljcr-c-38-14-3

Covering design C(38, 14, 3)

the best known covering uses 35 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005406 | ljcr-c-38-14-4

Covering design C(38, 14, 4)

the best known covering uses 113 blocks and the source-reported lower bound is 95.

来源状态 open | 2026-07-23 | 来源

V3P-005407 | ljcr-c-38-15-3

Covering design C(38, 15, 3)

the best known covering uses 31 blocks and the source-reported lower bound is 28.

来源状态 open | 2026-07-23 | 来源

V3P-005408 | ljcr-c-38-15-4

Covering design C(38, 15, 4)

the best known covering uses 102 blocks and the source-reported lower bound is 76.

来源状态 open | 2026-07-23 | 来源

V3P-005409 | ljcr-c-38-16-3

Covering design C(38, 16, 3)

the best known covering uses 26 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005410 | ljcr-c-38-16-4

Covering design C(38, 16, 4)

the best known covering uses 81 blocks and the source-reported lower bound is 60.

来源状态 open | 2026-07-23 | 来源

V3P-005411 | ljcr-c-38-17-3

Covering design C(38, 17, 3)

the best known covering uses 22 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-005412 | ljcr-c-38-17-4

Covering design C(38, 17, 4)

the best known covering uses 63 blocks and the source-reported lower bound is 47.

来源状态 open | 2026-07-23 | 来源

V3P-005413 | ljcr-c-38-17-5

Covering design C(38, 17, 5)

the best known covering uses 183 blocks and the source-reported lower bound is 126.

来源状态 open | 2026-07-23 | 来源

V3P-005414 | ljcr-c-38-18-4

Covering design C(38, 18, 4)

the best known covering uses 41 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005415 | ljcr-c-38-18-5

Covering design C(38, 18, 5)

the best known covering uses 122 blocks and the source-reported lower bound is 85.

来源状态 open | 2026-07-23 | 来源

V3P-005416 | ljcr-c-38-19-3

Covering design C(38, 19, 3)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005417 | ljcr-c-38-19-4

Covering design C(38, 19, 4)

the best known covering uses 33 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-005418 | ljcr-c-38-19-5

Covering design C(38, 19, 5)

the best known covering uses 86 blocks and the source-reported lower bound is 64.

来源状态 open | 2026-07-23 | 来源

V3P-005419 | ljcr-c-38-20-3

Covering design C(38, 20, 3)

the best known covering uses 14 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005420 | ljcr-c-38-20-4

Covering design C(38, 20, 4)

the best known covering uses 29 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005421 | ljcr-c-38-20-5

Covering design C(38, 20, 5)

the best known covering uses 62 blocks and the source-reported lower bound is 54.

来源状态 open | 2026-07-23 | 来源

V3P-005422 | ljcr-c-38-21-4

Covering design C(38, 21, 4)

the best known covering uses 27 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005423 | ljcr-c-38-21-5

Covering design C(38, 21, 5)

the best known covering uses 59 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005424 | ljcr-c-38-22-3

Covering design C(38, 22, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005425 | ljcr-c-38-22-4

Covering design C(38, 22, 4)

the best known covering uses 23 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005426 | ljcr-c-38-22-5

Covering design C(38, 22, 5)

the best known covering uses 49 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-005427 | ljcr-c-38-23-3

Covering design C(38, 23, 3)

the best known covering uses 10 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005428 | ljcr-c-38-23-4

Covering design C(38, 23, 4)

the best known covering uses 20 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005429 | ljcr-c-38-23-5

Covering design C(38, 23, 5)

the best known covering uses 42 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005430 | ljcr-c-38-24-4

Covering design C(38, 24, 4)

the best known covering uses 17 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-005431 | ljcr-c-38-24-5

Covering design C(38, 24, 5)

the best known covering uses 34 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005432 | ljcr-c-38-25-4

Covering design C(38, 25, 4)

the best known covering uses 15 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-005433 | ljcr-c-38-25-5

Covering design C(38, 25, 5)

the best known covering uses 29 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-005434 | ljcr-c-38-7-2

Covering design C(38, 7, 2)

the best known covering uses 39 blocks and the source-reported lower bound is 38.

来源状态 open | 2026-07-23 | 来源

V3P-005435 | ljcr-c-38-8-3

Covering design C(38, 8, 3)

the best known covering uses 191 blocks and the source-reported lower bound is 157.

来源状态 open | 2026-07-23 | 来源

V3P-005436 | ljcr-c-38-9-2

Covering design C(38, 9, 2)

the best known covering uses 25 blocks and the source-reported lower bound is 23.

来源状态 open | 2026-07-23 | 来源

V3P-005437 | ljcr-c-38-9-3

Covering design C(38, 9, 3)

the best known covering uses 131 blocks and the source-reported lower bound is 119.

来源状态 open | 2026-07-23 | 来源

V3P-005438 | ljcr-c-39-10-3

Covering design C(39, 10, 3)

the best known covering uses 104 blocks and the source-reported lower bound is 90.

来源状态 open | 2026-07-23 | 来源

V3P-005439 | ljcr-c-39-11-2

Covering design C(39, 11, 2)

the best known covering uses 18 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005440 | ljcr-c-39-11-3

Covering design C(39, 11, 3)

the best known covering uses 77 blocks and the source-reported lower bound is 68.

来源状态 open | 2026-07-23 | 来源

V3P-005441 | ljcr-c-39-12-3

Covering design C(39, 12, 3)

the best known covering uses 57 blocks and the source-reported lower bound is 49.

来源状态 open | 2026-07-23 | 来源

V3P-005442 | ljcr-c-39-13-3

Covering design C(39, 13, 3)

the best known covering uses 40 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-005443 | ljcr-c-39-13-4

Covering design C(39, 13, 4)

the best known covering uses 195 blocks and the source-reported lower bound is 135.

来源状态 open | 2026-07-23 | 来源

V3P-005444 | ljcr-c-39-14-3

Covering design C(39, 14, 3)

the best known covering uses 38 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005445 | ljcr-c-39-14-4

Covering design C(39, 14, 4)

the best known covering uses 136 blocks and the source-reported lower bound is 106.

来源状态 open | 2026-07-23 | 来源

V3P-005446 | ljcr-c-39-15-3

Covering design C(39, 15, 3)

the best known covering uses 33 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005447 | ljcr-c-39-15-4

Covering design C(39, 15, 4)

the best known covering uses 105 blocks and the source-reported lower bound is 86.

来源状态 open | 2026-07-23 | 来源

V3P-005448 | ljcr-c-39-16-3

Covering design C(39, 16, 3)

the best known covering uses 28 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005449 | ljcr-c-39-16-4

Covering design C(39, 16, 4)

the best known covering uses 88 blocks and the source-reported lower bound is 69.

来源状态 open | 2026-07-23 | 来源

V3P-005450 | ljcr-c-39-17-3

Covering design C(39, 17, 3)

the best known covering uses 23 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-005451 | ljcr-c-39-17-4

Covering design C(39, 17, 4)

the best known covering uses 71 blocks and the source-reported lower bound is 51.

来源状态 open | 2026-07-23 | 来源

V3P-005452 | ljcr-c-39-17-5

Covering design C(39, 17, 5)

the best known covering uses 215 blocks and the source-reported lower bound is 138.

来源状态 open | 2026-07-23 | 来源

V3P-005453 | ljcr-c-39-18-3

Covering design C(39, 18, 3)

the best known covering uses 19 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005454 | ljcr-c-39-18-4

Covering design C(39, 18, 4)

the best known covering uses 54 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-005455 | ljcr-c-39-18-5

Covering design C(39, 18, 5)

the best known covering uses 147 blocks and the source-reported lower bound is 102.

来源状态 open | 2026-07-23 | 来源

V3P-005456 | ljcr-c-39-19-4

Covering design C(39, 19, 4)

the best known covering uses 35 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-005457 | ljcr-c-39-19-5

Covering design C(39, 19, 5)

the best known covering uses 98 blocks and the source-reported lower bound is 70.

来源状态 open | 2026-07-23 | 来源

V3P-005458 | ljcr-c-39-20-4

Covering design C(39, 20, 4)

the best known covering uses 31 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005459 | ljcr-c-39-20-5

Covering design C(39, 20, 5)

the best known covering uses 70 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-005460 | ljcr-c-39-21-3

Covering design C(39, 21, 3)

the best known covering uses 13 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005461 | ljcr-c-39-21-4

Covering design C(39, 21, 4)

the best known covering uses 28 blocks and the source-reported lower bound is 23.

来源状态 open | 2026-07-23 | 来源

V3P-005462 | ljcr-c-39-21-5

Covering design C(39, 21, 5)

the best known covering uses 60 blocks and the source-reported lower bound is 51.

来源状态 open | 2026-07-23 | 来源

V3P-005463 | ljcr-c-39-22-4

Covering design C(39, 22, 4)

the best known covering uses 26 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005464 | ljcr-c-39-22-5

Covering design C(39, 22, 5)

the best known covering uses 56 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-005465 | ljcr-c-39-23-3

Covering design C(39, 23, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005466 | ljcr-c-39-23-4

Covering design C(39, 23, 4)

the best known covering uses 22 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005467 | ljcr-c-39-23-5

Covering design C(39, 23, 5)

the best known covering uses 44 blocks and the source-reported lower bound is 28.

来源状态 open | 2026-07-23 | 来源

V3P-005468 | ljcr-c-39-24-3

Covering design C(39, 24, 3)

the best known covering uses 9 blocks and the source-reported lower bound is 8.

来源状态 open | 2026-07-23 | 来源

V3P-005469 | ljcr-c-39-24-4

Covering design C(39, 24, 4)

the best known covering uses 18 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005470 | ljcr-c-39-24-5

Covering design C(39, 24, 5)

the best known covering uses 39 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005471 | ljcr-c-39-25-4

Covering design C(39, 25, 4)

the best known covering uses 16 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-005472 | ljcr-c-39-25-5

Covering design C(39, 25, 5)

the best known covering uses 33 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005473 | ljcr-c-39-6-2

Covering design C(39, 6, 2)

the best known covering uses 54 blocks and the source-reported lower bound is 52.

来源状态 open | 2026-07-23 | 来源

V3P-005474 | ljcr-c-39-8-2

Covering design C(39, 8, 2)

the best known covering uses 32 blocks and the source-reported lower bound is 31.

来源状态 open | 2026-07-23 | 来源

V3P-005475 | ljcr-c-39-8-3

Covering design C(39, 8, 3)

the best known covering uses 218 blocks and the source-reported lower bound is 186.

来源状态 open | 2026-07-23 | 来源

V3P-005476 | ljcr-c-39-9-2

Covering design C(39, 9, 2)

the best known covering uses 26 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005477 | ljcr-c-39-9-3

Covering design C(39, 9, 3)

the best known covering uses 139 blocks and the source-reported lower bound is 126.

来源状态 open | 2026-07-23 | 来源

V3P-005478 | ljcr-c-40-10-2

Covering design C(40, 10, 2)

the best known covering uses 21 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-005479 | ljcr-c-40-10-3

Covering design C(40, 10, 3)

the best known covering uses 110 blocks and the source-reported lower bound is 96.

来源状态 open | 2026-07-23 | 来源

V3P-005480 | ljcr-c-40-11-3

Covering design C(40, 11, 3)

the best known covering uses 85 blocks and the source-reported lower bound is 73.

来源状态 open | 2026-07-23 | 来源

V3P-005481 | ljcr-c-40-12-2

Covering design C(40, 12, 2)

the best known covering uses 16 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005482 | ljcr-c-40-12-3

Covering design C(40, 12, 3)

the best known covering uses 66 blocks and the source-reported lower bound is 57.

来源状态 open | 2026-07-23 | 来源

V3P-005483 | ljcr-c-40-13-4

Covering design C(40, 13, 4)

the best known covering uses 219 blocks and the source-reported lower bound is 151.

来源状态 open | 2026-07-23 | 来源

V3P-005484 | ljcr-c-40-14-3

Covering design C(40, 14, 3)

the best known covering uses 39 blocks and the source-reported lower bound is 38.

来源状态 open | 2026-07-23 | 来源

V3P-005485 | ljcr-c-40-14-4

Covering design C(40, 14, 4)

the best known covering uses 150 blocks and the source-reported lower bound is 112.

来源状态 open | 2026-07-23 | 来源

V3P-005486 | ljcr-c-40-15-3

Covering design C(40, 15, 3)

the best known covering uses 35 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-005487 | ljcr-c-40-15-4

Covering design C(40, 15, 4)

the best known covering uses 114 blocks and the source-reported lower bound is 91.

来源状态 open | 2026-07-23 | 来源

V3P-005488 | ljcr-c-40-16-3

Covering design C(40, 16, 3)

the best known covering uses 28 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005489 | ljcr-c-40-16-4

Covering design C(40, 16, 4)

the best known covering uses 93 blocks and the source-reported lower bound is 73.

来源状态 open | 2026-07-23 | 来源

V3P-005490 | ljcr-c-40-17-3

Covering design C(40, 17, 3)

the best known covering uses 25 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005491 | ljcr-c-40-17-4

Covering design C(40, 17, 4)

the best known covering uses 76 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-005492 | ljcr-c-40-17-5

Covering design C(40, 17, 5)

the best known covering uses 243 blocks and the source-reported lower bound is 163.

来源状态 open | 2026-07-23 | 来源

V3P-005493 | ljcr-c-40-18-3

Covering design C(40, 18, 3)

the best known covering uses 20 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005494 | ljcr-c-40-18-4

Covering design C(40, 18, 4)

the best known covering uses 59 blocks and the source-reported lower bound is 43.

来源状态 open | 2026-07-23 | 来源

V3P-005495 | ljcr-c-40-18-5

Covering design C(40, 18, 5)

the best known covering uses 182 blocks and the source-reported lower bound is 114.

来源状态 open | 2026-07-23 | 来源

V3P-005496 | ljcr-c-40-19-3

Covering design C(40, 19, 3)

the best known covering uses 17 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005497 | ljcr-c-40-19-4

Covering design C(40, 19, 4)

the best known covering uses 46 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005498 | ljcr-c-40-19-5

Covering design C(40, 19, 5)

the best known covering uses 129 blocks and the source-reported lower bound is 83.

来源状态 open | 2026-07-23 | 来源

V3P-005499 | ljcr-c-40-20-4

Covering design C(40, 20, 4)

the best known covering uses 31 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-005500 | ljcr-c-40-20-5

Covering design C(40, 20, 5)

the best known covering uses 72 blocks and the source-reported lower bound is 64.

来源状态 open | 2026-07-23 | 来源

V3P-005501 | ljcr-c-40-21-3

Covering design C(40, 21, 3)

the best known covering uses 14 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005502 | ljcr-c-40-21-4

Covering design C(40, 21, 4)

the best known covering uses 30 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005503 | ljcr-c-40-21-5

Covering design C(40, 21, 5)

the best known covering uses 63 blocks and the source-reported lower bound is 56.

来源状态 open | 2026-07-23 | 来源

V3P-005504 | ljcr-c-40-22-3

Covering design C(40, 22, 3)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-005505 | ljcr-c-40-22-4

Covering design C(40, 22, 4)

the best known covering uses 27 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005506 | ljcr-c-40-22-5

Covering design C(40, 22, 5)

the best known covering uses 58 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005507 | ljcr-c-40-23-3

Covering design C(40, 23, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005508 | ljcr-c-40-23-4

Covering design C(40, 23, 4)

the best known covering uses 25 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-005509 | ljcr-c-40-23-5

Covering design C(40, 23, 5)

the best known covering uses 53 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-005510 | ljcr-c-40-24-3

Covering design C(40, 24, 3)

the best known covering uses 10 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005511 | ljcr-c-40-24-4

Covering design C(40, 24, 4)

the best known covering uses 20 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005512 | ljcr-c-40-24-5

Covering design C(40, 24, 5)

the best known covering uses 42 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005513 | ljcr-c-40-25-4

Covering design C(40, 25, 4)

the best known covering uses 18 blocks and the source-reported lower bound is 13.

来源状态 open | 2026-07-23 | 来源

V3P-005514 | ljcr-c-40-25-5

Covering design C(40, 25, 5)

the best known covering uses 36 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005515 | ljcr-c-40-5-2

Covering design C(40, 5, 2)

the best known covering uses 82 blocks and the source-reported lower bound is 80.

来源状态 open | 2026-07-23 | 来源

V3P-005516 | ljcr-c-40-6-2

Covering design C(40, 6, 2)

the best known covering uses 55 blocks and the source-reported lower bound is 54.

来源状态 open | 2026-07-23 | 来源

V3P-005517 | ljcr-c-40-8-2

Covering design C(40, 8, 2)

the best known covering uses 33 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-005518 | ljcr-c-40-8-3

Covering design C(40, 8, 3)

the best known covering uses 231 blocks and the source-reported lower bound is 195.

来源状态 open | 2026-07-23 | 来源

V3P-005519 | ljcr-c-40-9-2

Covering design C(40, 9, 2)

the best known covering uses 26 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005520 | ljcr-c-40-9-3

Covering design C(40, 9, 3)

the best known covering uses 159 blocks and the source-reported lower bound is 138.

来源状态 open | 2026-07-23 | 来源

V3P-005521 | ljcr-c-41-10-3

Covering design C(41, 10, 3)

the best known covering uses 121 blocks and the source-reported lower bound is 103.

来源状态 open | 2026-07-23 | 来源

V3P-005522 | ljcr-c-41-11-2

Covering design C(41, 11, 2)

the best known covering uses 19 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-005523 | ljcr-c-41-11-3

Covering design C(41, 11, 3)

the best known covering uses 87 blocks and the source-reported lower bound is 75.

来源状态 open | 2026-07-23 | 来源

V3P-005524 | ljcr-c-41-12-2

Covering design C(41, 12, 2)

the best known covering uses 17 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005525 | ljcr-c-41-12-3

Covering design C(41, 12, 3)

the best known covering uses 73 blocks and the source-reported lower bound is 62.

来源状态 open | 2026-07-23 | 来源

V3P-005526 | ljcr-c-41-13-3

Covering design C(41, 13, 3)

the best known covering uses 55 blocks and the source-reported lower bound is 45.

来源状态 open | 2026-07-23 | 来源

V3P-005527 | ljcr-c-41-14-3

Covering design C(41, 14, 3)

the best known covering uses 40 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-005528 | ljcr-c-41-14-4

Covering design C(41, 14, 4)

the best known covering uses 160 blocks and the source-reported lower bound is 118.

来源状态 open | 2026-07-23 | 来源

V3P-005529 | ljcr-c-41-15-3

Covering design C(41, 15, 3)

the best known covering uses 36 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005530 | ljcr-c-41-15-4

Covering design C(41, 15, 4)

the best known covering uses 122 blocks and the source-reported lower bound is 104.

来源状态 open | 2026-07-23 | 来源

V3P-005531 | ljcr-c-41-16-3

Covering design C(41, 16, 3)

the best known covering uses 31 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005532 | ljcr-c-41-16-4

Covering design C(41, 16, 4)

the best known covering uses 104 blocks and the source-reported lower bound is 82.

来源状态 open | 2026-07-23 | 来源

V3P-005533 | ljcr-c-41-17-3

Covering design C(41, 17, 3)

the best known covering uses 27 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005534 | ljcr-c-41-17-4

Covering design C(41, 17, 4)

the best known covering uses 83 blocks and the source-reported lower bound is 61.

来源状态 open | 2026-07-23 | 来源

V3P-005535 | ljcr-c-41-18-3

Covering design C(41, 18, 3)

the best known covering uses 23 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-005536 | ljcr-c-41-18-4

Covering design C(41, 18, 4)

the best known covering uses 69 blocks and the source-reported lower bound is 51.

来源状态 open | 2026-07-23 | 来源

V3P-005537 | ljcr-c-41-18-5

Covering design C(41, 18, 5)

the best known covering uses 209 blocks and the source-reported lower bound is 135.

来源状态 open | 2026-07-23 | 来源

V3P-005538 | ljcr-c-41-19-3

Covering design C(41, 19, 3)

the best known covering uses 18 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005539 | ljcr-c-41-19-4

Covering design C(41, 19, 4)

the best known covering uses 52 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-005540 | ljcr-c-41-19-5

Covering design C(41, 19, 5)

the best known covering uses 159 blocks and the source-reported lower bound is 93.

来源状态 open | 2026-07-23 | 来源

V3P-005541 | ljcr-c-41-20-4

Covering design C(41, 20, 4)

the best known covering uses 39 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-005542 | ljcr-c-41-20-5

Covering design C(41, 20, 5)

the best known covering uses 107 blocks and the source-reported lower bound is 70.

来源状态 open | 2026-07-23 | 来源

V3P-005543 | ljcr-c-41-21-4

Covering design C(41, 21, 4)

the best known covering uses 30 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005544 | ljcr-c-41-21-5

Covering design C(41, 21, 5)

the best known covering uses 67 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-005545 | ljcr-c-41-22-3

Covering design C(41, 22, 3)

the best known covering uses 13 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005546 | ljcr-c-41-22-4

Covering design C(41, 22, 4)

the best known covering uses 29 blocks and the source-reported lower bound is 23.

来源状态 open | 2026-07-23 | 来源

V3P-005547 | ljcr-c-41-22-5

Covering design C(41, 22, 5)

the best known covering uses 61 blocks and the source-reported lower bound is 51.

来源状态 open | 2026-07-23 | 来源

V3P-005548 | ljcr-c-41-23-4

Covering design C(41, 23, 4)

the best known covering uses 26 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005549 | ljcr-c-41-23-5

Covering design C(41, 23, 5)

the best known covering uses 57 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-005550 | ljcr-c-41-24-3

Covering design C(41, 24, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005551 | ljcr-c-41-24-4

Covering design C(41, 24, 4)

the best known covering uses 23 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005552 | ljcr-c-41-24-5

Covering design C(41, 24, 5)

the best known covering uses 47 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-005553 | ljcr-c-41-25-3

Covering design C(41, 25, 3)

the best known covering uses 9 blocks and the source-reported lower bound is 8.

来源状态 open | 2026-07-23 | 来源

V3P-005554 | ljcr-c-41-25-4

Covering design C(41, 25, 4)

the best known covering uses 20 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005555 | ljcr-c-41-25-5

Covering design C(41, 25, 5)

the best known covering uses 39 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-005556 | ljcr-c-41-6-2

Covering design C(41, 6, 2)

the best known covering uses 59 blocks and the source-reported lower bound is 56.

来源状态 open | 2026-07-23 | 来源

V3P-005557 | ljcr-c-41-7-2

Covering design C(41, 7, 2)

the best known covering uses 43 blocks and the source-reported lower bound is 41.

来源状态 open | 2026-07-23 | 来源

V3P-005558 | ljcr-c-41-8-2

Covering design C(41, 8, 2)

the best known covering uses 34 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005559 | ljcr-c-41-9-2

Covering design C(41, 9, 2)

the best known covering uses 28 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-005560 | ljcr-c-41-9-3

Covering design C(41, 9, 3)

the best known covering uses 168 blocks and the source-reported lower bound is 146.

来源状态 open | 2026-07-23 | 来源

V3P-005561 | ljcr-c-42-10-3

Covering design C(42, 10, 3)

the best known covering uses 130 blocks and the source-reported lower bound is 110.

来源状态 open | 2026-07-23 | 来源

V3P-005562 | ljcr-c-42-11-3

Covering design C(42, 11, 3)

the best known covering uses 91 blocks and the source-reported lower bound is 81.

来源状态 open | 2026-07-23 | 来源

V3P-005563 | ljcr-c-42-12-2

Covering design C(42, 12, 2)

the best known covering uses 17 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005564 | ljcr-c-42-12-3

Covering design C(42, 12, 3)

the best known covering uses 77 blocks and the source-reported lower bound is 63.

来源状态 open | 2026-07-23 | 来源

V3P-005565 | ljcr-c-42-13-3

Covering design C(42, 13, 3)

the best known covering uses 58 blocks and the source-reported lower bound is 49.

来源状态 open | 2026-07-23 | 来源

V3P-005566 | ljcr-c-42-14-3

Covering design C(42, 14, 3)

the best known covering uses 44 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005567 | ljcr-c-42-14-4

Covering design C(42, 14, 4)

the best known covering uses 193 blocks and the source-reported lower bound is 135.

来源状态 open | 2026-07-23 | 来源

V3P-005568 | ljcr-c-42-15-3

Covering design C(42, 15, 3)

the best known covering uses 38 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-005569 | ljcr-c-42-15-4

Covering design C(42, 15, 4)

the best known covering uses 131 blocks and the source-reported lower bound is 110.

来源状态 open | 2026-07-23 | 来源

V3P-005570 | ljcr-c-42-16-3

Covering design C(42, 16, 3)

the best known covering uses 34 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005571 | ljcr-c-42-16-4

Covering design C(42, 16, 4)

the best known covering uses 108 blocks and the source-reported lower bound is 87.

来源状态 open | 2026-07-23 | 来源

V3P-005572 | ljcr-c-42-17-3

Covering design C(42, 17, 3)

the best known covering uses 29 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005573 | ljcr-c-42-17-4

Covering design C(42, 17, 4)

the best known covering uses 95 blocks and the source-reported lower bound is 72.

来源状态 open | 2026-07-23 | 来源

V3P-005574 | ljcr-c-42-18-3

Covering design C(42, 18, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005575 | ljcr-c-42-18-4

Covering design C(42, 18, 4)

the best known covering uses 70 blocks and the source-reported lower bound is 52.

来源状态 open | 2026-07-23 | 来源

V3P-005576 | ljcr-c-42-18-5

Covering design C(42, 18, 5)

the best known covering uses 231 blocks and the source-reported lower bound is 143.

来源状态 open | 2026-07-23 | 来源

V3P-005577 | ljcr-c-42-19-3

Covering design C(42, 19, 3)

the best known covering uses 21 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005578 | ljcr-c-42-19-4

Covering design C(42, 19, 4)

the best known covering uses 60 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005579 | ljcr-c-42-19-5

Covering design C(42, 19, 5)

the best known covering uses 176 blocks and the source-reported lower bound is 113.

来源状态 open | 2026-07-23 | 来源

V3P-005580 | ljcr-c-42-20-4

Covering design C(42, 20, 4)

the best known covering uses 44 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005581 | ljcr-c-42-20-5

Covering design C(42, 20, 5)

the best known covering uses 129 blocks and the source-reported lower bound is 78.

来源状态 open | 2026-07-23 | 来源

V3P-005582 | ljcr-c-42-21-3

Covering design C(42, 21, 3)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005583 | ljcr-c-42-21-4

Covering design C(42, 21, 4)

the best known covering uses 34 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-005584 | ljcr-c-42-21-5

Covering design C(42, 21, 5)

the best known covering uses 87 blocks and the source-reported lower bound is 64.

来源状态 open | 2026-07-23 | 来源

V3P-005585 | ljcr-c-42-22-3

Covering design C(42, 22, 3)

the best known covering uses 14 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005586 | ljcr-c-42-22-4

Covering design C(42, 22, 4)

the best known covering uses 29 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005587 | ljcr-c-42-22-5

Covering design C(42, 22, 5)

the best known covering uses 63 blocks and the source-reported lower bound is 56.

来源状态 open | 2026-07-23 | 来源

V3P-005588 | ljcr-c-42-23-3

Covering design C(42, 23, 3)

the best known covering uses 12 blocks and the source-reported lower bound is 11.

来源状态 open | 2026-07-23 | 来源

V3P-005589 | ljcr-c-42-23-4

Covering design C(42, 23, 4)

the best known covering uses 28 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005590 | ljcr-c-42-23-5

Covering design C(42, 23, 5)

the best known covering uses 59 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005591 | ljcr-c-42-24-3

Covering design C(42, 24, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005592 | ljcr-c-42-24-4

Covering design C(42, 24, 4)

the best known covering uses 24 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-005593 | ljcr-c-42-24-5

Covering design C(42, 24, 5)

the best known covering uses 54 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-005594 | ljcr-c-42-25-3

Covering design C(42, 25, 3)

the best known covering uses 10 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005595 | ljcr-c-42-25-4

Covering design C(42, 25, 4)

the best known covering uses 21 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005596 | ljcr-c-42-25-5

Covering design C(42, 25, 5)

the best known covering uses 44 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005597 | ljcr-c-42-7-2

Covering design C(42, 7, 2)

the best known covering uses 46 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005598 | ljcr-c-42-8-2

Covering design C(42, 8, 2)

the best known covering uses 35 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005599 | ljcr-c-42-9-2

Covering design C(42, 9, 2)

the best known covering uses 29 blocks and the source-reported lower bound is 28.

来源状态 open | 2026-07-23 | 来源

V3P-005600 | ljcr-c-42-9-3

Covering design C(42, 9, 3)

the best known covering uses 176 blocks and the source-reported lower bound is 154.

来源状态 open | 2026-07-23 | 来源

V3P-005601 | ljcr-c-43-10-2

Covering design C(43, 10, 2)

the best known covering uses 26 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005602 | ljcr-c-43-10-3

Covering design C(43, 10, 3)

the best known covering uses 141 blocks and the source-reported lower bound is 121.

来源状态 open | 2026-07-23 | 来源

V3P-005603 | ljcr-c-43-11-3

Covering design C(43, 11, 3)

the best known covering uses 93 blocks and the source-reported lower bound is 83.

来源状态 open | 2026-07-23 | 来源

V3P-005604 | ljcr-c-43-12-2

Covering design C(43, 12, 2)

the best known covering uses 18 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005605 | ljcr-c-43-12-3

Covering design C(43, 12, 3)

the best known covering uses 77 blocks and the source-reported lower bound is 72.

来源状态 open | 2026-07-23 | 来源

V3P-005606 | ljcr-c-43-13-2

Covering design C(43, 13, 2)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005607 | ljcr-c-43-13-3

Covering design C(43, 13, 3)

the best known covering uses 61 blocks and the source-reported lower bound is 53.

来源状态 open | 2026-07-23 | 来源

V3P-005608 | ljcr-c-43-14-3

Covering design C(43, 14, 3)

the best known covering uses 47 blocks and the source-reported lower bound is 43.

来源状态 open | 2026-07-23 | 来源

V3P-005609 | ljcr-c-43-14-4

Covering design C(43, 14, 4)

the best known covering uses 218 blocks and the source-reported lower bound is 151.

来源状态 open | 2026-07-23 | 来源

V3P-005610 | ljcr-c-43-15-3

Covering design C(43, 15, 3)

the best known covering uses 40 blocks and the source-reported lower bound is 38.

来源状态 open | 2026-07-23 | 来源

V3P-005611 | ljcr-c-43-15-4

Covering design C(43, 15, 4)

the best known covering uses 150 blocks and the source-reported lower bound is 121.

来源状态 open | 2026-07-23 | 来源

V3P-005612 | ljcr-c-43-16-3

Covering design C(43, 16, 3)

the best known covering uses 35 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005613 | ljcr-c-43-16-4

Covering design C(43, 16, 4)

the best known covering uses 114 blocks and the source-reported lower bound is 95.

来源状态 open | 2026-07-23 | 来源

V3P-005614 | ljcr-c-43-17-3

Covering design C(43, 17, 3)

the best known covering uses 30 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-005615 | ljcr-c-43-17-4

Covering design C(43, 17, 4)

the best known covering uses 102 blocks and the source-reported lower bound is 74.

来源状态 open | 2026-07-23 | 来源

V3P-005616 | ljcr-c-43-18-3

Covering design C(43, 18, 3)

the best known covering uses 27 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005617 | ljcr-c-43-18-4

Covering design C(43, 18, 4)

the best known covering uses 81 blocks and the source-reported lower bound is 60.

来源状态 open | 2026-07-23 | 来源

V3P-005618 | ljcr-c-43-19-3

Covering design C(43, 19, 3)

the best known covering uses 22 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-005619 | ljcr-c-43-19-4

Covering design C(43, 19, 4)

the best known covering uses 66 blocks and the source-reported lower bound is 48.

来源状态 open | 2026-07-23 | 来源

V3P-005620 | ljcr-c-43-19-5

Covering design C(43, 19, 5)

the best known covering uses 198 blocks and the source-reported lower bound is 118.

来源状态 open | 2026-07-23 | 来源

V3P-005621 | ljcr-c-43-20-3

Covering design C(43, 20, 3)

the best known covering uses 18 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005622 | ljcr-c-43-20-4

Covering design C(43, 20, 4)

the best known covering uses 53 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-005623 | ljcr-c-43-20-5

Covering design C(43, 20, 5)

the best known covering uses 147 blocks and the source-reported lower bound is 91.

来源状态 open | 2026-07-23 | 来源

V3P-005624 | ljcr-c-43-21-4

Covering design C(43, 21, 4)

the best known covering uses 36 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-005625 | ljcr-c-43-21-5

Covering design C(43, 21, 5)

the best known covering uses 106 blocks and the source-reported lower bound is 70.

来源状态 open | 2026-07-23 | 来源

V3P-005626 | ljcr-c-43-22-4

Covering design C(43, 22, 4)

the best known covering uses 31 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005627 | ljcr-c-43-22-5

Covering design C(43, 22, 5)

the best known covering uses 73 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-005628 | ljcr-c-43-23-3

Covering design C(43, 23, 3)

the best known covering uses 13 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005629 | ljcr-c-43-23-4

Covering design C(43, 23, 4)

the best known covering uses 29 blocks and the source-reported lower bound is 23.

来源状态 open | 2026-07-23 | 来源

V3P-005630 | ljcr-c-43-23-5

Covering design C(43, 23, 5)

the best known covering uses 61 blocks and the source-reported lower bound is 51.

来源状态 open | 2026-07-23 | 来源

V3P-005631 | ljcr-c-43-24-4

Covering design C(43, 24, 4)

the best known covering uses 27 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005632 | ljcr-c-43-24-5

Covering design C(43, 24, 5)

the best known covering uses 57 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-005633 | ljcr-c-43-25-3

Covering design C(43, 25, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005634 | ljcr-c-43-25-4

Covering design C(43, 25, 4)

the best known covering uses 23 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005635 | ljcr-c-43-25-5

Covering design C(43, 25, 5)

the best known covering uses 48 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-005636 | ljcr-c-43-7-2

Covering design C(43, 7, 2)

the best known covering uses 49 blocks and the source-reported lower bound is 45.

来源状态 open | 2026-07-23 | 来源

V3P-005637 | ljcr-c-43-8-2

Covering design C(43, 8, 2)

the best known covering uses 37 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-005638 | ljcr-c-43-9-2

Covering design C(43, 9, 2)

the best known covering uses 30 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005639 | ljcr-c-43-9-3

Covering design C(43, 9, 3)

the best known covering uses 193 blocks and the source-reported lower bound is 163.

来源状态 open | 2026-07-23 | 来源

V3P-005640 | ljcr-c-44-10-2

Covering design C(44, 10, 2)

the best known covering uses 26 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005641 | ljcr-c-44-10-3

Covering design C(44, 10, 3)

the best known covering uses 149 blocks and the source-reported lower bound is 128.

来源状态 open | 2026-07-23 | 来源

V3P-005642 | ljcr-c-44-11-2

Covering design C(44, 11, 2)

the best known covering uses 21 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-005643 | ljcr-c-44-11-3

Covering design C(44, 11, 3)

the best known covering uses 110 blocks and the source-reported lower bound is 96.

来源状态 open | 2026-07-23 | 来源

V3P-005644 | ljcr-c-44-12-2

Covering design C(44, 12, 2)

the best known covering uses 19 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-005645 | ljcr-c-44-12-3

Covering design C(44, 12, 3)

the best known covering uses 77 blocks and the source-reported lower bound is 74.

来源状态 open | 2026-07-23 | 来源

V3P-005646 | ljcr-c-44-13-2

Covering design C(44, 13, 2)

the best known covering uses 16 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005647 | ljcr-c-44-13-3

Covering design C(44, 13, 3)

the best known covering uses 66 blocks and the source-reported lower bound is 58.

来源状态 open | 2026-07-23 | 来源

V3P-005648 | ljcr-c-44-14-3

Covering design C(44, 14, 3)

the best known covering uses 49 blocks and the source-reported lower bound is 44.

来源状态 open | 2026-07-23 | 来源

V3P-005649 | ljcr-c-44-14-4

Covering design C(44, 14, 4)

the best known covering uses 246 blocks and the source-reported lower bound is 167.

来源状态 open | 2026-07-23 | 来源

V3P-005650 | ljcr-c-44-15-3

Covering design C(44, 15, 3)

the best known covering uses 41 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-005651 | ljcr-c-44-15-4

Covering design C(44, 15, 4)

the best known covering uses 159 blocks and the source-reported lower bound is 127.

来源状态 open | 2026-07-23 | 来源

V3P-005652 | ljcr-c-44-16-3

Covering design C(44, 16, 3)

the best known covering uses 37 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005653 | ljcr-c-44-16-4

Covering design C(44, 16, 4)

the best known covering uses 127 blocks and the source-reported lower bound is 105.

来源状态 open | 2026-07-23 | 来源

V3P-005654 | ljcr-c-44-17-3

Covering design C(44, 17, 3)

the best known covering uses 32 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005655 | ljcr-c-44-17-4

Covering design C(44, 17, 4)

the best known covering uses 106 blocks and the source-reported lower bound is 86.

来源状态 open | 2026-07-23 | 来源

V3P-005656 | ljcr-c-44-18-3

Covering design C(44, 18, 3)

the best known covering uses 27 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005657 | ljcr-c-44-18-4

Covering design C(44, 18, 4)

the best known covering uses 92 blocks and the source-reported lower bound is 64.

来源状态 open | 2026-07-23 | 来源

V3P-005658 | ljcr-c-44-19-3

Covering design C(44, 19, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005659 | ljcr-c-44-19-4

Covering design C(44, 19, 4)

the best known covering uses 72 blocks and the source-reported lower bound is 51.

来源状态 open | 2026-07-23 | 来源

V3P-005660 | ljcr-c-44-19-5

Covering design C(44, 19, 5)

the best known covering uses 233 blocks and the source-reported lower bound is 139.

来源状态 open | 2026-07-23 | 来源

V3P-005661 | ljcr-c-44-20-3

Covering design C(44, 20, 3)

the best known covering uses 19 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005662 | ljcr-c-44-20-4

Covering design C(44, 20, 4)

the best known covering uses 54 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-005663 | ljcr-c-44-20-5

Covering design C(44, 20, 5)

the best known covering uses 162 blocks and the source-reported lower bound is 106.

来源状态 open | 2026-07-23 | 来源

V3P-005664 | ljcr-c-44-21-4

Covering design C(44, 21, 4)

the best known covering uses 45 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005665 | ljcr-c-44-21-5

Covering design C(44, 21, 5)

the best known covering uses 130 blocks and the source-reported lower bound is 78.

来源状态 open | 2026-07-23 | 来源

V3P-005666 | ljcr-c-44-22-3

Covering design C(44, 22, 3)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005667 | ljcr-c-44-22-4

Covering design C(44, 22, 4)

the best known covering uses 31 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-005668 | ljcr-c-44-22-5

Covering design C(44, 22, 5)

the best known covering uses 78 blocks and the source-reported lower bound is 64.

来源状态 open | 2026-07-23 | 来源

V3P-005669 | ljcr-c-44-23-3

Covering design C(44, 23, 3)

the best known covering uses 14 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005670 | ljcr-c-44-23-4

Covering design C(44, 23, 4)

the best known covering uses 30 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005671 | ljcr-c-44-23-5

Covering design C(44, 23, 5)

the best known covering uses 63 blocks and the source-reported lower bound is 56.

来源状态 open | 2026-07-23 | 来源

V3P-005672 | ljcr-c-44-24-4

Covering design C(44, 24, 4)

the best known covering uses 28 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005673 | ljcr-c-44-24-5

Covering design C(44, 24, 5)

the best known covering uses 60 blocks and the source-reported lower bound is 43.

来源状态 open | 2026-07-23 | 来源

V3P-005674 | ljcr-c-44-25-3

Covering design C(44, 25, 3)

the best known covering uses 11 blocks and the source-reported lower bound is 9.

来源状态 open | 2026-07-23 | 来源

V3P-005675 | ljcr-c-44-25-4

Covering design C(44, 25, 4)

the best known covering uses 25 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-005676 | ljcr-c-44-25-5

Covering design C(44, 25, 5)

the best known covering uses 54 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-005677 | ljcr-c-44-5-2

Covering design C(44, 5, 2)

the best known covering uses 99 blocks and the source-reported lower bound is 97.

来源状态 open | 2026-07-23 | 来源

V3P-005678 | ljcr-c-44-6-2

Covering design C(44, 6, 2)

the best known covering uses 67 blocks and the source-reported lower bound is 66.

来源状态 open | 2026-07-23 | 来源

V3P-005679 | ljcr-c-44-9-3

Covering design C(44, 9, 3)

the best known covering uses 210 blocks and the source-reported lower bound is 172.

来源状态 open | 2026-07-23 | 来源

V3P-005680 | ljcr-c-45-10-2

Covering design C(45, 10, 2)

the best known covering uses 27 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-005681 | ljcr-c-45-10-3

Covering design C(45, 10, 3)

the best known covering uses 158 blocks and the source-reported lower bound is 135.

来源状态 open | 2026-07-23 | 来源

V3P-005682 | ljcr-c-45-11-2

Covering design C(45, 11, 2)

the best known covering uses 22 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005683 | ljcr-c-45-11-3

Covering design C(45, 11, 3)

the best known covering uses 117 blocks and the source-reported lower bound is 103.

来源状态 open | 2026-07-23 | 来源

V3P-005684 | ljcr-c-45-12-2

Covering design C(45, 12, 2)

the best known covering uses 19 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-005685 | ljcr-c-45-12-3

Covering design C(45, 12, 3)

the best known covering uses 88 blocks and the source-reported lower bound is 75.

来源状态 open | 2026-07-23 | 来源

V3P-005686 | ljcr-c-45-13-2

Covering design C(45, 13, 2)

the best known covering uses 17 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005687 | ljcr-c-45-13-3

Covering design C(45, 13, 3)

the best known covering uses 73 blocks and the source-reported lower bound is 63.

来源状态 open | 2026-07-23 | 来源

V3P-005688 | ljcr-c-45-14-3

Covering design C(45, 14, 3)

the best known covering uses 59 blocks and the source-reported lower bound is 45.

来源状态 open | 2026-07-23 | 来源

V3P-005689 | ljcr-c-45-15-3

Covering design C(45, 15, 3)

the best known covering uses 42 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-005690 | ljcr-c-45-15-4

Covering design C(45, 15, 4)

the best known covering uses 166 blocks and the source-reported lower bound is 132.

来源状态 open | 2026-07-23 | 来源

V3P-005691 | ljcr-c-45-16-3

Covering design C(45, 16, 3)

the best known covering uses 38 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-005692 | ljcr-c-45-16-4

Covering design C(45, 16, 4)

the best known covering uses 133 blocks and the source-reported lower bound is 110.

来源状态 open | 2026-07-23 | 来源

V3P-005693 | ljcr-c-45-17-3

Covering design C(45, 17, 3)

the best known covering uses 34 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-005694 | ljcr-c-45-17-4

Covering design C(45, 17, 4)

the best known covering uses 111 blocks and the source-reported lower bound is 88.

来源状态 open | 2026-07-23 | 来源

V3P-005695 | ljcr-c-45-18-3

Covering design C(45, 18, 3)

the best known covering uses 30 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005696 | ljcr-c-45-18-4

Covering design C(45, 18, 4)

the best known covering uses 98 blocks and the source-reported lower bound is 73.

来源状态 open | 2026-07-23 | 来源

V3P-005697 | ljcr-c-45-19-3

Covering design C(45, 19, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005698 | ljcr-c-45-19-4

Covering design C(45, 19, 4)

the best known covering uses 74 blocks and the source-reported lower bound is 60.

来源状态 open | 2026-07-23 | 来源

V3P-005699 | ljcr-c-45-20-3

Covering design C(45, 20, 3)

the best known covering uses 22 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-005700 | ljcr-c-45-20-4

Covering design C(45, 20, 4)

the best known covering uses 65 blocks and the source-reported lower bound is 48.

来源状态 open | 2026-07-23 | 来源

V3P-005701 | ljcr-c-45-20-5

Covering design C(45, 20, 5)

the best known covering uses 185 blocks and the source-reported lower bound is 115.

来源状态 open | 2026-07-23 | 来源

V3P-005702 | ljcr-c-45-21-4

Covering design C(45, 21, 4)

the best known covering uses 48 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-005703 | ljcr-c-45-21-5

Covering design C(45, 21, 5)

the best known covering uses 148 blocks and the source-reported lower bound is 86.

来源状态 open | 2026-07-23 | 来源

V3P-005704 | ljcr-c-45-22-4

Covering design C(45, 22, 4)

the best known covering uses 37 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-005705 | ljcr-c-45-22-5

Covering design C(45, 22, 5)

the best known covering uses 107 blocks and the source-reported lower bound is 70.

来源状态 open | 2026-07-23 | 来源

V3P-005706 | ljcr-c-45-23-4

Covering design C(45, 23, 4)

the best known covering uses 30 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005707 | ljcr-c-45-23-5

Covering design C(45, 23, 5)

the best known covering uses 65 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-005708 | ljcr-c-45-24-3

Covering design C(45, 24, 3)

the best known covering uses 13 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005709 | ljcr-c-45-24-4

Covering design C(45, 24, 4)

the best known covering uses 29 blocks and the source-reported lower bound is 23.

来源状态 open | 2026-07-23 | 来源

V3P-005710 | ljcr-c-45-24-5

Covering design C(45, 24, 5)

the best known covering uses 61 blocks and the source-reported lower bound is 51.

来源状态 open | 2026-07-23 | 来源

V3P-005711 | ljcr-c-45-25-4

Covering design C(45, 25, 4)

the best known covering uses 27 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005712 | ljcr-c-45-25-5

Covering design C(45, 25, 5)

the best known covering uses 57 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-005713 | ljcr-c-45-6-2

Covering design C(45, 6, 2)

the best known covering uses 70 blocks and the source-reported lower bound is 68.

来源状态 open | 2026-07-23 | 来源

V3P-005714 | ljcr-c-45-8-2

Covering design C(45, 8, 2)

the best known covering uses 42 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-005715 | ljcr-c-45-9-2

Covering design C(45, 9, 2)

the best known covering uses 33 blocks and the source-reported lower bound is 32.

来源状态 open | 2026-07-23 | 来源

V3P-005716 | ljcr-c-45-9-3

Covering design C(45, 9, 3)

the best known covering uses 227 blocks and the source-reported lower bound is 195.

来源状态 open | 2026-07-23 | 来源

V3P-005717 | ljcr-c-46-10-2

Covering design C(46, 10, 2)

the best known covering uses 28 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-005718 | ljcr-c-46-10-3

Covering design C(46, 10, 3)

the best known covering uses 179 blocks and the source-reported lower bound is 148.

来源状态 open | 2026-07-23 | 来源

V3P-005719 | ljcr-c-46-11-2

Covering design C(46, 11, 2)

the best known covering uses 23 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005720 | ljcr-c-46-11-3

Covering design C(46, 11, 3)

the best known covering uses 128 blocks and the source-reported lower bound is 109.

来源状态 open | 2026-07-23 | 来源

V3P-005721 | ljcr-c-46-12-3

Covering design C(46, 12, 3)

the best known covering uses 90 blocks and the source-reported lower bound is 81.

来源状态 open | 2026-07-23 | 来源

V3P-005722 | ljcr-c-46-13-2

Covering design C(46, 13, 2)

the best known covering uses 17 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005723 | ljcr-c-46-13-3

Covering design C(46, 13, 3)

the best known covering uses 77 blocks and the source-reported lower bound is 64.

来源状态 open | 2026-07-23 | 来源

V3P-005724 | ljcr-c-46-14-2

Covering design C(46, 14, 2)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005725 | ljcr-c-46-14-3

Covering design C(46, 14, 3)

the best known covering uses 62 blocks and the source-reported lower bound is 50.

来源状态 open | 2026-07-23 | 来源

V3P-005726 | ljcr-c-46-15-3

Covering design C(46, 15, 3)

the best known covering uses 48 blocks and the source-reported lower bound is 43.

来源状态 open | 2026-07-23 | 来源

V3P-005727 | ljcr-c-46-15-4

Covering design C(46, 15, 4)

the best known covering uses 211 blocks and the source-reported lower bound is 138.

来源状态 open | 2026-07-23 | 来源

V3P-005728 | ljcr-c-46-16-3

Covering design C(46, 16, 3)

the best known covering uses 38 blocks and the source-reported lower bound is 36.

来源状态 open | 2026-07-23 | 来源

V3P-005729 | ljcr-c-46-16-4

Covering design C(46, 16, 4)

the best known covering uses 147 blocks and the source-reported lower bound is 115.

来源状态 open | 2026-07-23 | 来源

V3P-005730 | ljcr-c-46-17-3

Covering design C(46, 17, 3)

the best known covering uses 36 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005731 | ljcr-c-46-17-4

Covering design C(46, 17, 4)

the best known covering uses 116 blocks and the source-reported lower bound is 95.

来源状态 open | 2026-07-23 | 来源

V3P-005732 | ljcr-c-46-18-3

Covering design C(46, 18, 3)

the best known covering uses 31 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-005733 | ljcr-c-46-18-4

Covering design C(46, 18, 4)

the best known covering uses 105 blocks and the source-reported lower bound is 77.

来源状态 open | 2026-07-23 | 来源

V3P-005734 | ljcr-c-46-19-3

Covering design C(46, 19, 3)

the best known covering uses 27 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005735 | ljcr-c-46-19-4

Covering design C(46, 19, 4)

the best known covering uses 86 blocks and the source-reported lower bound is 61.

来源状态 open | 2026-07-23 | 来源

V3P-005736 | ljcr-c-46-20-3

Covering design C(46, 20, 3)

the best known covering uses 23 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-005737 | ljcr-c-46-20-4

Covering design C(46, 20, 4)

the best known covering uses 69 blocks and the source-reported lower bound is 51.

来源状态 open | 2026-07-23 | 来源

V3P-005738 | ljcr-c-46-20-5

Covering design C(46, 20, 5)

the best known covering uses 221 blocks and the source-reported lower bound is 138.

来源状态 open | 2026-07-23 | 来源

V3P-005739 | ljcr-c-46-21-3

Covering design C(46, 21, 3)

the best known covering uses 20 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005740 | ljcr-c-46-21-4

Covering design C(46, 21, 4)

the best known covering uses 59 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-005741 | ljcr-c-46-21-5

Covering design C(46, 21, 5)

the best known covering uses 170 blocks and the source-reported lower bound is 106.

来源状态 open | 2026-07-23 | 来源

V3P-005742 | ljcr-c-46-22-4

Covering design C(46, 22, 4)

the best known covering uses 38 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005743 | ljcr-c-46-22-5

Covering design C(46, 22, 5)

the best known covering uses 109 blocks and the source-reported lower bound is 78.

来源状态 open | 2026-07-23 | 来源

V3P-005744 | ljcr-c-46-23-3

Covering design C(46, 23, 3)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005745 | ljcr-c-46-23-4

Covering design C(46, 23, 4)

the best known covering uses 31 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-005746 | ljcr-c-46-23-5

Covering design C(46, 23, 5)

the best known covering uses 82 blocks and the source-reported lower bound is 64.

来源状态 open | 2026-07-23 | 来源

V3P-005747 | ljcr-c-46-24-3

Covering design C(46, 24, 3)

the best known covering uses 14 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005748 | ljcr-c-46-24-4

Covering design C(46, 24, 4)

the best known covering uses 30 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005749 | ljcr-c-46-24-5

Covering design C(46, 24, 5)

the best known covering uses 62 blocks and the source-reported lower bound is 56.

来源状态 open | 2026-07-23 | 来源

V3P-005750 | ljcr-c-46-25-3

Covering design C(46, 25, 3)

the best known covering uses 13 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005751 | ljcr-c-46-25-4

Covering design C(46, 25, 4)

the best known covering uses 28 blocks and the source-reported lower bound is 23.

来源状态 open | 2026-07-23 | 来源

V3P-005752 | ljcr-c-46-25-5

Covering design C(46, 25, 5)

the best known covering uses 59 blocks and the source-reported lower bound is 43.

来源状态 open | 2026-07-23 | 来源

V3P-005753 | ljcr-c-46-6-2

Covering design C(46, 6, 2)

the best known covering uses 73 blocks and the source-reported lower bound is 69.

来源状态 open | 2026-07-23 | 来源

V3P-005754 | ljcr-c-46-8-2

Covering design C(46, 8, 2)

the best known covering uses 43 blocks and the source-reported lower bound is 41.

来源状态 open | 2026-07-23 | 来源

V3P-005755 | ljcr-c-46-9-2

Covering design C(46, 9, 2)

the best known covering uses 34 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005756 | ljcr-c-46-9-3

Covering design C(46, 9, 3)

the best known covering uses 246 blocks and the source-reported lower bound is 205.

来源状态 open | 2026-07-23 | 来源

V3P-005757 | ljcr-c-47-10-3

Covering design C(47, 10, 3)

the best known covering uses 185 blocks and the source-reported lower bound is 156.

来源状态 open | 2026-07-23 | 来源

V3P-005758 | ljcr-c-47-11-2

Covering design C(47, 11, 2)

the best known covering uses 23 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005759 | ljcr-c-47-11-3

Covering design C(47, 11, 3)

the best known covering uses 139 blocks and the source-reported lower bound is 112.

来源状态 open | 2026-07-23 | 来源

V3P-005760 | ljcr-c-47-12-3

Covering design C(47, 12, 3)

the best known covering uses 92 blocks and the source-reported lower bound is 83.

来源状态 open | 2026-07-23 | 来源

V3P-005761 | ljcr-c-47-13-3

Covering design C(47, 13, 3)

the best known covering uses 79 blocks and the source-reported lower bound is 73.

来源状态 open | 2026-07-23 | 来源

V3P-005762 | ljcr-c-47-14-2

Covering design C(47, 14, 2)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005763 | ljcr-c-47-14-3

Covering design C(47, 14, 3)

the best known covering uses 66 blocks and the source-reported lower bound is 54.

来源状态 open | 2026-07-23 | 来源

V3P-005764 | ljcr-c-47-15-3

Covering design C(47, 15, 3)

the best known covering uses 52 blocks and the source-reported lower bound is 44.

来源状态 open | 2026-07-23 | 来源

V3P-005765 | ljcr-c-47-15-4

Covering design C(47, 15, 4)

the best known covering uses 235 blocks and the source-reported lower bound is 157.

来源状态 open | 2026-07-23 | 来源

V3P-005766 | ljcr-c-47-16-3

Covering design C(47, 16, 3)

the best known covering uses 41 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-005767 | ljcr-c-47-16-4

Covering design C(47, 16, 4)

the best known covering uses 166 blocks and the source-reported lower bound is 127.

来源状态 open | 2026-07-23 | 来源

V3P-005768 | ljcr-c-47-17-3

Covering design C(47, 17, 3)

the best known covering uses 37 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005769 | ljcr-c-47-17-4

Covering design C(47, 17, 4)

the best known covering uses 126 blocks and the source-reported lower bound is 100.

来源状态 open | 2026-07-23 | 来源

V3P-005770 | ljcr-c-47-18-3

Covering design C(47, 18, 3)

the best known covering uses 34 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005771 | ljcr-c-47-18-4

Covering design C(47, 18, 4)

the best known covering uses 108 blocks and the source-reported lower bound is 87.

来源状态 open | 2026-07-23 | 来源

V3P-005772 | ljcr-c-47-19-3

Covering design C(47, 19, 3)

the best known covering uses 29 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005773 | ljcr-c-47-19-4

Covering design C(47, 19, 4)

the best known covering uses 98 blocks and the source-reported lower bound is 65.

来源状态 open | 2026-07-23 | 来源

V3P-005774 | ljcr-c-47-20-3

Covering design C(47, 20, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005775 | ljcr-c-47-20-4

Covering design C(47, 20, 4)

the best known covering uses 70 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-005776 | ljcr-c-47-21-3

Covering design C(47, 21, 3)

the best known covering uses 22 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-005777 | ljcr-c-47-21-4

Covering design C(47, 21, 4)

the best known covering uses 65 blocks and the source-reported lower bound is 43.

来源状态 open | 2026-07-23 | 来源

V3P-005778 | ljcr-c-47-21-5

Covering design C(47, 21, 5)

the best known covering uses 187 blocks and the source-reported lower bound is 115.

来源状态 open | 2026-07-23 | 来源

V3P-005779 | ljcr-c-47-22-3

Covering design C(47, 22, 3)

the best known covering uses 18 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005780 | ljcr-c-47-22-4

Covering design C(47, 22, 4)

the best known covering uses 51 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-005781 | ljcr-c-47-22-5

Covering design C(47, 22, 5)

the best known covering uses 148 blocks and the source-reported lower bound is 86.

来源状态 open | 2026-07-23 | 来源

V3P-005782 | ljcr-c-47-23-5

Covering design C(47, 23, 5)

the best known covering uses 87 blocks and the source-reported lower bound is 68.

来源状态 open | 2026-07-23 | 来源

V3P-005783 | ljcr-c-47-24-4

Covering design C(47, 24, 4)

the best known covering uses 30 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005784 | ljcr-c-47-24-5

Covering design C(47, 24, 5)

the best known covering uses 63 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-005785 | ljcr-c-47-25-3

Covering design C(47, 25, 3)

the best known covering uses 13 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005786 | ljcr-c-47-25-4

Covering design C(47, 25, 4)

the best known covering uses 29 blocks and the source-reported lower bound is 23.

来源状态 open | 2026-07-23 | 来源

V3P-005787 | ljcr-c-47-25-5

Covering design C(47, 25, 5)

the best known covering uses 60 blocks and the source-reported lower bound is 51.

来源状态 open | 2026-07-23 | 来源

V3P-005788 | ljcr-c-47-7-2

Covering design C(47, 7, 2)

the best known covering uses 56 blocks and the source-reported lower bound is 54.

来源状态 open | 2026-07-23 | 来源

V3P-005789 | ljcr-c-47-8-2

Covering design C(47, 8, 2)

the best known covering uses 44 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005790 | ljcr-c-47-9-2

Covering design C(47, 9, 2)

the best known covering uses 35 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005791 | ljcr-c-48-10-2

Covering design C(48, 10, 2)

the best known covering uses 30 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005792 | ljcr-c-48-10-3

Covering design C(48, 10, 3)

the best known covering uses 192 blocks and the source-reported lower bound is 164.

来源状态 open | 2026-07-23 | 来源

V3P-005793 | ljcr-c-48-11-2

Covering design C(48, 11, 2)

the best known covering uses 26 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-005794 | ljcr-c-48-11-3

Covering design C(48, 11, 3)

the best known covering uses 150 blocks and the source-reported lower bound is 127.

来源状态 open | 2026-07-23 | 来源

V3P-005795 | ljcr-c-48-12-3

Covering design C(48, 12, 3)

the best known covering uses 92 blocks and the source-reported lower bound is 88.

来源状态 open | 2026-07-23 | 来源

V3P-005796 | ljcr-c-48-13-2

Covering design C(48, 13, 2)

the best known covering uses 19 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-005797 | ljcr-c-48-13-3

Covering design C(48, 13, 3)

the best known covering uses 81 blocks and the source-reported lower bound is 74.

来源状态 open | 2026-07-23 | 来源

V3P-005798 | ljcr-c-48-14-2

Covering design C(48, 14, 2)

the best known covering uses 17 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005799 | ljcr-c-48-14-3

Covering design C(48, 14, 3)

the best known covering uses 72 blocks and the source-reported lower bound is 62.

来源状态 open | 2026-07-23 | 来源

V3P-005800 | ljcr-c-48-15-3

Covering design C(48, 15, 3)

the best known covering uses 54 blocks and the source-reported lower bound is 45.

来源状态 open | 2026-07-23 | 来源

V3P-005801 | ljcr-c-48-16-3

Covering design C(48, 16, 3)

the best known covering uses 43 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-005802 | ljcr-c-48-16-4

Covering design C(48, 16, 4)

the best known covering uses 183 blocks and the source-reported lower bound is 132.

来源状态 open | 2026-07-23 | 来源

V3P-005803 | ljcr-c-48-17-3

Covering design C(48, 17, 3)

the best known covering uses 38 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-005804 | ljcr-c-48-17-4

Covering design C(48, 17, 4)

the best known covering uses 132 blocks and the source-reported lower bound is 111.

来源状态 open | 2026-07-23 | 来源

V3P-005805 | ljcr-c-48-18-3

Covering design C(48, 18, 3)

the best known covering uses 35 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-005806 | ljcr-c-48-18-4

Covering design C(48, 18, 4)

the best known covering uses 112 blocks and the source-reported lower bound is 91.

来源状态 open | 2026-07-23 | 来源

V3P-005807 | ljcr-c-48-19-3

Covering design C(48, 19, 3)

the best known covering uses 30 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-005808 | ljcr-c-48-19-4

Covering design C(48, 19, 4)

the best known covering uses 103 blocks and the source-reported lower bound is 74.

来源状态 open | 2026-07-23 | 来源

V3P-005809 | ljcr-c-48-20-3

Covering design C(48, 20, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005810 | ljcr-c-48-20-4

Covering design C(48, 20, 4)

the best known covering uses 86 blocks and the source-reported lower bound is 60.

来源状态 open | 2026-07-23 | 来源

V3P-005811 | ljcr-c-48-21-3

Covering design C(48, 21, 3)

the best known covering uses 23 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-005812 | ljcr-c-48-21-4

Covering design C(48, 21, 4)

the best known covering uses 69 blocks and the source-reported lower bound is 51.

来源状态 open | 2026-07-23 | 来源

V3P-005813 | ljcr-c-48-21-5

Covering design C(48, 21, 5)

the best known covering uses 204 blocks and the source-reported lower bound is 135.

来源状态 open | 2026-07-23 | 来源

V3P-005814 | ljcr-c-48-22-3

Covering design C(48, 22, 3)

the best known covering uses 18 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005815 | ljcr-c-48-22-4

Covering design C(48, 22, 4)

the best known covering uses 56 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-005816 | ljcr-c-48-22-5

Covering design C(48, 22, 5)

the best known covering uses 169 blocks and the source-reported lower bound is 94.

来源状态 open | 2026-07-23 | 来源

V3P-005817 | ljcr-c-48-23-3

Covering design C(48, 23, 3)

the best known covering uses 16 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005818 | ljcr-c-48-23-4

Covering design C(48, 23, 4)

the best known covering uses 43 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005819 | ljcr-c-48-23-5

Covering design C(48, 23, 5)

the best known covering uses 120 blocks and the source-reported lower bound is 74.

来源状态 open | 2026-07-23 | 来源

V3P-005820 | ljcr-c-48-24-5

Covering design C(48, 24, 5)

the best known covering uses 63 blocks and the source-reported lower bound is 62.

来源状态 open | 2026-07-23 | 来源

V3P-005821 | ljcr-c-48-25-3

Covering design C(48, 25, 3)

the best known covering uses 14 blocks and the source-reported lower bound is 12.

来源状态 open | 2026-07-23 | 来源

V3P-005822 | ljcr-c-48-25-4

Covering design C(48, 25, 4)

the best known covering uses 30 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005823 | ljcr-c-48-25-5

Covering design C(48, 25, 5)

the best known covering uses 62 blocks and the source-reported lower bound is 56.

来源状态 open | 2026-07-23 | 来源

V3P-005824 | ljcr-c-48-5-2

Covering design C(48, 5, 2)

the best known covering uses 117 blocks and the source-reported lower bound is 116.

来源状态 open | 2026-07-23 | 来源

V3P-005825 | ljcr-c-48-7-2

Covering design C(48, 7, 2)

the best known covering uses 56 blocks and the source-reported lower bound is 55.

来源状态 open | 2026-07-23 | 来源

V3P-005826 | ljcr-c-48-8-2

Covering design C(48, 8, 2)

the best known covering uses 46 blocks and the source-reported lower bound is 44.

来源状态 open | 2026-07-23 | 来源

V3P-005827 | ljcr-c-48-9-2

Covering design C(48, 9, 2)

the best known covering uses 37 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-005828 | ljcr-c-49-10-3

Covering design C(49, 10, 3)

the best known covering uses 198 blocks and the source-reported lower bound is 172.

来源状态 open | 2026-07-23 | 来源

V3P-005829 | ljcr-c-49-11-2

Covering design C(49, 11, 2)

the best known covering uses 27 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005830 | ljcr-c-49-11-3

Covering design C(49, 11, 3)

the best known covering uses 156 blocks and the source-reported lower bound is 130.

来源状态 open | 2026-07-23 | 来源

V3P-005831 | ljcr-c-49-12-2

Covering design C(49, 12, 2)

the best known covering uses 22 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005832 | ljcr-c-49-12-3

Covering design C(49, 12, 3)

the best known covering uses 116 blocks and the source-reported lower bound is 98.

来源状态 open | 2026-07-23 | 来源

V3P-005833 | ljcr-c-49-13-3

Covering design C(49, 13, 3)

the best known covering uses 82 blocks and the source-reported lower bound is 76.

来源状态 open | 2026-07-23 | 来源

V3P-005834 | ljcr-c-49-14-2

Covering design C(49, 14, 2)

the best known covering uses 17 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005835 | ljcr-c-49-14-3

Covering design C(49, 14, 3)

the best known covering uses 76 blocks and the source-reported lower bound is 63.

来源状态 open | 2026-07-23 | 来源

V3P-005836 | ljcr-c-49-15-2

Covering design C(49, 15, 2)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005837 | ljcr-c-49-15-3

Covering design C(49, 15, 3)

the best known covering uses 62 blocks and the source-reported lower bound is 49.

来源状态 open | 2026-07-23 | 来源

V3P-005838 | ljcr-c-49-16-3

Covering design C(49, 16, 3)

the best known covering uses 44 blocks and the source-reported lower bound is 41.

来源状态 open | 2026-07-23 | 来源

V3P-005839 | ljcr-c-49-16-4

Covering design C(49, 16, 4)

the best known covering uses 200 blocks and the source-reported lower bound is 138.

来源状态 open | 2026-07-23 | 来源

V3P-005840 | ljcr-c-49-17-3

Covering design C(49, 17, 3)

the best known covering uses 39 blocks and the source-reported lower bound is 36.

来源状态 open | 2026-07-23 | 来源

V3P-005841 | ljcr-c-49-17-4

Covering design C(49, 17, 4)

the best known covering uses 139 blocks and the source-reported lower bound is 116.

来源状态 open | 2026-07-23 | 来源

V3P-005842 | ljcr-c-49-18-3

Covering design C(49, 18, 3)

the best known covering uses 36 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005843 | ljcr-c-49-18-4

Covering design C(49, 18, 4)

the best known covering uses 123 blocks and the source-reported lower bound is 96.

来源状态 open | 2026-07-23 | 来源

V3P-005844 | ljcr-c-49-19-3

Covering design C(49, 19, 3)

the best known covering uses 32 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-005845 | ljcr-c-49-19-4

Covering design C(49, 19, 4)

the best known covering uses 106 blocks and the source-reported lower bound is 78.

来源状态 open | 2026-07-23 | 来源

V3P-005846 | ljcr-c-49-20-3

Covering design C(49, 20, 3)

the best known covering uses 28 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005847 | ljcr-c-49-20-4

Covering design C(49, 20, 4)

the best known covering uses 94 blocks and the source-reported lower bound is 64.

来源状态 open | 2026-07-23 | 来源

V3P-005848 | ljcr-c-49-21-3

Covering design C(49, 21, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-005849 | ljcr-c-49-21-4

Covering design C(49, 21, 4)

the best known covering uses 75 blocks and the source-reported lower bound is 52.

来源状态 open | 2026-07-23 | 来源

V3P-005850 | ljcr-c-49-22-3

Covering design C(49, 22, 3)

the best known covering uses 21 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-005851 | ljcr-c-49-22-4

Covering design C(49, 22, 4)

the best known covering uses 62 blocks and the source-reported lower bound is 43.

来源状态 open | 2026-07-23 | 来源

V3P-005852 | ljcr-c-49-22-5

Covering design C(49, 22, 5)

the best known covering uses 183 blocks and the source-reported lower bound is 114.

来源状态 open | 2026-07-23 | 来源

V3P-005853 | ljcr-c-49-23-3

Covering design C(49, 23, 3)

the best known covering uses 16 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005854 | ljcr-c-49-23-4

Covering design C(49, 23, 4)

the best known covering uses 49 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-005855 | ljcr-c-49-23-5

Covering design C(49, 23, 5)

the best known covering uses 134 blocks and the source-reported lower bound is 86.

来源状态 open | 2026-07-23 | 来源

V3P-005856 | ljcr-c-49-24-4

Covering design C(49, 24, 4)

the best known covering uses 37 blocks and the source-reported lower bound is 31.

来源状态 open | 2026-07-23 | 来源

V3P-005857 | ljcr-c-49-24-5

Covering design C(49, 24, 5)

the best known covering uses 102 blocks and the source-reported lower bound is 68.

来源状态 open | 2026-07-23 | 来源

V3P-005858 | ljcr-c-49-25-4

Covering design C(49, 25, 4)

the best known covering uses 30 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005859 | ljcr-c-49-25-5

Covering design C(49, 25, 5)

the best known covering uses 63 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-005860 | ljcr-c-49-5-2

Covering design C(49, 5, 2)

the best known covering uses 122 blocks and the source-reported lower bound is 118.

来源状态 open | 2026-07-23 | 来源

V3P-005861 | ljcr-c-49-8-2

Covering design C(49, 8, 2)

the best known covering uses 49 blocks and the source-reported lower bound is 44.

来源状态 open | 2026-07-23 | 来源

V3P-005862 | ljcr-c-49-9-2

Covering design C(49, 9, 2)

the best known covering uses 38 blocks and the source-reported lower bound is 36.

来源状态 open | 2026-07-23 | 来源

V3P-005863 | ljcr-c-50-10-3

Covering design C(50, 10, 3)

the best known covering uses 204 blocks and the source-reported lower bound is 180.

来源状态 open | 2026-07-23 | 来源

V3P-005864 | ljcr-c-50-11-2

Covering design C(50, 11, 2)

the best known covering uses 28 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-005865 | ljcr-c-50-11-3

Covering design C(50, 11, 3)

the best known covering uses 163 blocks and the source-reported lower bound is 137.

来源状态 open | 2026-07-23 | 来源

V3P-005866 | ljcr-c-50-12-2

Covering design C(50, 12, 2)

the best known covering uses 23 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-005867 | ljcr-c-50-12-3

Covering design C(50, 12, 3)

the best known covering uses 122 blocks and the source-reported lower bound is 105.

来源状态 open | 2026-07-23 | 来源

V3P-005868 | ljcr-c-50-13-3

Covering design C(50, 13, 3)

the best known covering uses 92 blocks and the source-reported lower bound is 81.

来源状态 open | 2026-07-23 | 来源

V3P-005869 | ljcr-c-50-14-2

Covering design C(50, 14, 2)

the best known covering uses 18 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005870 | ljcr-c-50-14-3

Covering design C(50, 14, 3)

the best known covering uses 77 blocks and the source-reported lower bound is 68.

来源状态 open | 2026-07-23 | 来源

V3P-005871 | ljcr-c-50-15-2

Covering design C(50, 15, 2)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005872 | ljcr-c-50-15-3

Covering design C(50, 15, 3)

the best known covering uses 66 blocks and the source-reported lower bound is 54.

来源状态 open | 2026-07-23 | 来源

V3P-005873 | ljcr-c-50-16-3

Covering design C(50, 16, 3)

the best known covering uses 53 blocks and the source-reported lower bound is 44.

来源状态 open | 2026-07-23 | 来源

V3P-005874 | ljcr-c-50-16-4

Covering design C(50, 16, 4)

the best known covering uses 236 blocks and the source-reported lower bound is 154.

来源状态 open | 2026-07-23 | 来源

V3P-005875 | ljcr-c-50-17-4

Covering design C(50, 17, 4)

the best known covering uses 150 blocks and the source-reported lower bound is 121.

来源状态 open | 2026-07-23 | 来源

V3P-005876 | ljcr-c-50-18-3

Covering design C(50, 18, 3)

the best known covering uses 37 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005877 | ljcr-c-50-18-4

Covering design C(50, 18, 4)

the best known covering uses 127 blocks and the source-reported lower bound is 100.

来源状态 open | 2026-07-23 | 来源

V3P-005878 | ljcr-c-50-19-3

Covering design C(50, 19, 3)

the best known covering uses 33 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005879 | ljcr-c-50-19-4

Covering design C(50, 19, 4)

the best known covering uses 109 blocks and the source-reported lower bound is 87.

来源状态 open | 2026-07-23 | 来源

V3P-005880 | ljcr-c-50-20-3

Covering design C(50, 20, 3)

the best known covering uses 29 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005881 | ljcr-c-50-20-4

Covering design C(50, 20, 4)

the best known covering uses 97 blocks and the source-reported lower bound is 65.

来源状态 open | 2026-07-23 | 来源

V3P-005882 | ljcr-c-50-21-3

Covering design C(50, 21, 3)

the best known covering uses 26 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005883 | ljcr-c-50-21-4

Covering design C(50, 21, 4)

the best known covering uses 79 blocks and the source-reported lower bound is 60.

来源状态 open | 2026-07-23 | 来源

V3P-005884 | ljcr-c-50-22-3

Covering design C(50, 22, 3)

the best known covering uses 22 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-005885 | ljcr-c-50-22-4

Covering design C(50, 22, 4)

the best known covering uses 66 blocks and the source-reported lower bound is 44.

来源状态 open | 2026-07-23 | 来源

V3P-005886 | ljcr-c-50-22-5

Covering design C(50, 22, 5)

the best known covering uses 204 blocks and the source-reported lower bound is 119.

来源状态 open | 2026-07-23 | 来源

V3P-005887 | ljcr-c-50-23-3

Covering design C(50, 23, 3)

the best known covering uses 19 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005888 | ljcr-c-50-23-4

Covering design C(50, 23, 4)

the best known covering uses 54 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-005889 | ljcr-c-50-23-5

Covering design C(50, 23, 5)

the best known covering uses 156 blocks and the source-reported lower bound is 94.

来源状态 open | 2026-07-23 | 来源

V3P-005890 | ljcr-c-50-24-4

Covering design C(50, 24, 4)

the best known covering uses 41 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005891 | ljcr-c-50-24-5

Covering design C(50, 24, 5)

the best known covering uses 117 blocks and the source-reported lower bound is 73.

来源状态 open | 2026-07-23 | 来源

V3P-005892 | ljcr-c-50-25-3

Covering design C(50, 25, 3)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005893 | ljcr-c-50-25-4

Covering design C(50, 25, 4)

the best known covering uses 33 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-005894 | ljcr-c-50-25-5

Covering design C(50, 25, 5)

the best known covering uses 79 blocks and the source-reported lower bound is 62.

来源状态 open | 2026-07-23 | 来源

V3P-005895 | ljcr-c-50-6-2

Covering design C(50, 6, 2)

the best known covering uses 87 blocks and the source-reported lower bound is 84.

来源状态 open | 2026-07-23 | 来源

V3P-005896 | ljcr-c-50-8-2

Covering design C(50, 8, 2)

the best known covering uses 49 blocks and the source-reported lower bound is 45.

来源状态 open | 2026-07-23 | 来源

V3P-005897 | ljcr-c-50-9-2

Covering design C(50, 9, 2)

the best known covering uses 40 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-005898 | ljcr-c-51-10-2

Covering design C(51, 10, 2)

the best known covering uses 35 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005899 | ljcr-c-51-10-3

Covering design C(51, 10, 3)

the best known covering uses 233 blocks and the source-reported lower bound is 199.

来源状态 open | 2026-07-23 | 来源

V3P-005900 | ljcr-c-51-11-2

Covering design C(51, 11, 2)

the best known covering uses 28 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-005901 | ljcr-c-51-11-3

Covering design C(51, 11, 3)

the best known covering uses 175 blocks and the source-reported lower bound is 140.

来源状态 open | 2026-07-23 | 来源

V3P-005902 | ljcr-c-51-12-2

Covering design C(51, 12, 2)

the best known covering uses 24 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005903 | ljcr-c-51-12-3

Covering design C(51, 12, 3)

the best known covering uses 129 blocks and the source-reported lower bound is 111.

来源状态 open | 2026-07-23 | 来源

V3P-005904 | ljcr-c-51-13-3

Covering design C(51, 13, 3)

the best known covering uses 95 blocks and the source-reported lower bound is 83.

来源状态 open | 2026-07-23 | 来源

V3P-005905 | ljcr-c-51-14-2

Covering design C(51, 14, 2)

the best known covering uses 19 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-005906 | ljcr-c-51-14-3

Covering design C(51, 14, 3)

the best known covering uses 81 blocks and the source-reported lower bound is 73.

来源状态 open | 2026-07-23 | 来源

V3P-005907 | ljcr-c-51-15-2

Covering design C(51, 15, 2)

the best known covering uses 16 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005908 | ljcr-c-51-15-3

Covering design C(51, 15, 3)

the best known covering uses 68 blocks and the source-reported lower bound is 55.

来源状态 open | 2026-07-23 | 来源

V3P-005909 | ljcr-c-51-16-3

Covering design C(51, 16, 3)

the best known covering uses 55 blocks and the source-reported lower bound is 45.

来源状态 open | 2026-07-23 | 来源

V3P-005910 | ljcr-c-51-17-3

Covering design C(51, 17, 3)

the best known covering uses 43 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-005911 | ljcr-c-51-17-4

Covering design C(51, 17, 4)

the best known covering uses 180 blocks and the source-reported lower bound is 132.

来源状态 open | 2026-07-23 | 来源

V3P-005912 | ljcr-c-51-18-3

Covering design C(51, 18, 3)

the best known covering uses 38 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-005913 | ljcr-c-51-18-4

Covering design C(51, 18, 4)

the best known covering uses 131 blocks and the source-reported lower bound is 111.

来源状态 open | 2026-07-23 | 来源

V3P-005914 | ljcr-c-51-19-3

Covering design C(51, 19, 3)

the best known covering uses 35 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-005915 | ljcr-c-51-19-4

Covering design C(51, 19, 4)

the best known covering uses 113 blocks and the source-reported lower bound is 92.

来源状态 open | 2026-07-23 | 来源

V3P-005916 | ljcr-c-51-20-3

Covering design C(51, 20, 3)

the best known covering uses 31 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-005917 | ljcr-c-51-20-4

Covering design C(51, 20, 4)

the best known covering uses 105 blocks and the source-reported lower bound is 74.

来源状态 open | 2026-07-23 | 来源

V3P-005918 | ljcr-c-51-21-3

Covering design C(51, 21, 3)

the best known covering uses 27 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005919 | ljcr-c-51-21-4

Covering design C(51, 21, 4)

the best known covering uses 89 blocks and the source-reported lower bound is 61.

来源状态 open | 2026-07-23 | 来源

V3P-005920 | ljcr-c-51-22-3

Covering design C(51, 22, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-005921 | ljcr-c-51-22-4

Covering design C(51, 22, 4)

the best known covering uses 71 blocks and the source-reported lower bound is 51.

来源状态 open | 2026-07-23 | 来源

V3P-005922 | ljcr-c-51-22-5

Covering design C(51, 22, 5)

the best known covering uses 225 blocks and the source-reported lower bound is 140.

来源状态 open | 2026-07-23 | 来源

V3P-005923 | ljcr-c-51-23-3

Covering design C(51, 23, 3)

the best known covering uses 20 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005924 | ljcr-c-51-23-4

Covering design C(51, 23, 4)

the best known covering uses 59 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-005925 | ljcr-c-51-23-5

Covering design C(51, 23, 5)

the best known covering uses 173 blocks and the source-reported lower bound is 98.

来源状态 open | 2026-07-23 | 来源

V3P-005926 | ljcr-c-51-24-3

Covering design C(51, 24, 3)

the best known covering uses 16 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005927 | ljcr-c-51-24-4

Covering design C(51, 24, 4)

the best known covering uses 50 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005928 | ljcr-c-51-24-5

Covering design C(51, 24, 5)

the best known covering uses 130 blocks and the source-reported lower bound is 85.

来源状态 open | 2026-07-23 | 来源

V3P-005929 | ljcr-c-51-25-4

Covering design C(51, 25, 4)

the best known covering uses 35 blocks and the source-reported lower bound is 31.

来源状态 open | 2026-07-23 | 来源

V3P-005930 | ljcr-c-51-25-5

Covering design C(51, 25, 5)

the best known covering uses 96 blocks and the source-reported lower bound is 68.

来源状态 open | 2026-07-23 | 来源

V3P-005931 | ljcr-c-51-6-2

Covering design C(51, 6, 2)

the best known covering uses 90 blocks and the source-reported lower bound is 85.

来源状态 open | 2026-07-23 | 来源

V3P-005932 | ljcr-c-51-9-2

Covering design C(51, 9, 2)

the best known covering uses 41 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-005933 | ljcr-c-52-10-2

Covering design C(52, 10, 2)

the best known covering uses 36 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005934 | ljcr-c-52-11-2

Covering design C(52, 11, 2)

the best known covering uses 30 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005935 | ljcr-c-52-11-3

Covering design C(52, 11, 3)

the best known covering uses 187 blocks and the source-reported lower bound is 156.

来源状态 open | 2026-07-23 | 来源

V3P-005936 | ljcr-c-52-12-2

Covering design C(52, 12, 2)

the best known covering uses 24 blocks and the source-reported lower bound is 23.

来源状态 open | 2026-07-23 | 来源

V3P-005937 | ljcr-c-52-12-3

Covering design C(52, 12, 3)

the best known covering uses 130 blocks and the source-reported lower bound is 117.

来源状态 open | 2026-07-23 | 来源

V3P-005938 | ljcr-c-52-13-2

Covering design C(52, 13, 2)

the best known covering uses 21 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-005939 | ljcr-c-52-13-3

Covering design C(52, 13, 3)

the best known covering uses 101 blocks and the source-reported lower bound is 88.

来源状态 open | 2026-07-23 | 来源

V3P-005940 | ljcr-c-52-14-2

Covering design C(52, 14, 2)

the best known covering uses 19 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-005941 | ljcr-c-52-14-3

Covering design C(52, 14, 3)

the best known covering uses 82 blocks and the source-reported lower bound is 75.

来源状态 open | 2026-07-23 | 来源

V3P-005942 | ljcr-c-52-15-2

Covering design C(52, 15, 2)

the best known covering uses 17 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-005943 | ljcr-c-52-15-3

Covering design C(52, 15, 3)

the best known covering uses 76 blocks and the source-reported lower bound is 63.

来源状态 open | 2026-07-23 | 来源

V3P-005944 | ljcr-c-52-16-3

Covering design C(52, 16, 3)

the best known covering uses 59 blocks and the source-reported lower bound is 46.

来源状态 open | 2026-07-23 | 来源

V3P-005945 | ljcr-c-52-17-3

Covering design C(52, 17, 3)

the best known covering uses 45 blocks and the source-reported lower bound is 41.

来源状态 open | 2026-07-23 | 来源

V3P-005946 | ljcr-c-52-17-4

Covering design C(52, 17, 4)

the best known covering uses 205 blocks and the source-reported lower bound is 138.

来源状态 open | 2026-07-23 | 来源

V3P-005947 | ljcr-c-52-18-3

Covering design C(52, 18, 3)

the best known covering uses 39 blocks and the source-reported lower bound is 38.

来源状态 open | 2026-07-23 | 来源

V3P-005948 | ljcr-c-52-18-4

Covering design C(52, 18, 4)

the best known covering uses 135 blocks and the source-reported lower bound is 116.

来源状态 open | 2026-07-23 | 来源

V3P-005949 | ljcr-c-52-19-3

Covering design C(52, 19, 3)

the best known covering uses 36 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005950 | ljcr-c-52-19-4

Covering design C(52, 19, 4)

the best known covering uses 119 blocks and the source-reported lower bound is 96.

来源状态 open | 2026-07-23 | 来源

V3P-005951 | ljcr-c-52-20-3

Covering design C(52, 20, 3)

the best known covering uses 31 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005952 | ljcr-c-52-20-4

Covering design C(52, 20, 4)

the best known covering uses 108 blocks and the source-reported lower bound is 78.

来源状态 open | 2026-07-23 | 来源

V3P-005953 | ljcr-c-52-21-3

Covering design C(52, 21, 3)

the best known covering uses 29 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005954 | ljcr-c-52-21-4

Covering design C(52, 21, 4)

the best known covering uses 99 blocks and the source-reported lower bound is 65.

来源状态 open | 2026-07-23 | 来源

V3P-005955 | ljcr-c-52-22-3

Covering design C(52, 22, 3)

the best known covering uses 25 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005956 | ljcr-c-52-22-4

Covering design C(52, 22, 4)

the best known covering uses 76 blocks and the source-reported lower bound is 52.

来源状态 open | 2026-07-23 | 来源

V3P-005957 | ljcr-c-52-23-3

Covering design C(52, 23, 3)

the best known covering uses 23 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-005958 | ljcr-c-52-23-4

Covering design C(52, 23, 4)

the best known covering uses 64 blocks and the source-reported lower bound is 44.

来源状态 open | 2026-07-23 | 来源

V3P-005959 | ljcr-c-52-23-5

Covering design C(52, 23, 5)

the best known covering uses 193 blocks and the source-reported lower bound is 116.

来源状态 open | 2026-07-23 | 来源

V3P-005960 | ljcr-c-52-24-3

Covering design C(52, 24, 3)

the best known covering uses 17 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005961 | ljcr-c-52-24-4

Covering design C(52, 24, 4)

the best known covering uses 51 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-005962 | ljcr-c-52-24-5

Covering design C(52, 24, 5)

the best known covering uses 147 blocks and the source-reported lower bound is 87.

来源状态 open | 2026-07-23 | 来源

V3P-005963 | ljcr-c-52-25-4

Covering design C(52, 25, 4)

the best known covering uses 42 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-005964 | ljcr-c-52-25-5

Covering design C(52, 25, 5)

the best known covering uses 120 blocks and the source-reported lower bound is 71.

来源状态 open | 2026-07-23 | 来源

V3P-005965 | ljcr-c-52-7-2

Covering design C(52, 7, 2)

the best known covering uses 68 blocks and the source-reported lower bound is 67.

来源状态 open | 2026-07-23 | 来源

V3P-005966 | ljcr-c-52-8-2

Covering design C(52, 8, 2)

the best known covering uses 54 blocks and the source-reported lower bound is 52.

来源状态 open | 2026-07-23 | 来源

V3P-005967 | ljcr-c-52-9-2

Covering design C(52, 9, 2)

the best known covering uses 44 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005968 | ljcr-c-53-10-2

Covering design C(53, 10, 2)

the best known covering uses 37 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-005969 | ljcr-c-53-11-2

Covering design C(53, 11, 2)

the best known covering uses 30 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-005970 | ljcr-c-53-11-3

Covering design C(53, 11, 3)

the best known covering uses 193 blocks and the source-reported lower bound is 164.

来源状态 open | 2026-07-23 | 来源

V3P-005971 | ljcr-c-53-12-2

Covering design C(53, 12, 2)

the best known covering uses 27 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-005972 | ljcr-c-53-12-3

Covering design C(53, 12, 3)

the best known covering uses 148 blocks and the source-reported lower bound is 129.

来源状态 open | 2026-07-23 | 来源

V3P-005973 | ljcr-c-53-13-3

Covering design C(53, 13, 3)

the best known covering uses 104 blocks and the source-reported lower bound is 94.

来源状态 open | 2026-07-23 | 来源

V3P-005974 | ljcr-c-53-14-3

Covering design C(53, 14, 3)

the best known covering uses 84 blocks and the source-reported lower bound is 76.

来源状态 open | 2026-07-23 | 来源

V3P-005975 | ljcr-c-53-15-2

Covering design C(53, 15, 2)

the best known covering uses 18 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-005976 | ljcr-c-53-15-3

Covering design C(53, 15, 3)

the best known covering uses 77 blocks and the source-reported lower bound is 64.

来源状态 open | 2026-07-23 | 来源

V3P-005977 | ljcr-c-53-16-2

Covering design C(53, 16, 2)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-005978 | ljcr-c-53-16-3

Covering design C(53, 16, 3)

the best known covering uses 64 blocks and the source-reported lower bound is 50.

来源状态 open | 2026-07-23 | 来源

V3P-005979 | ljcr-c-53-17-3

Covering design C(53, 17, 3)

the best known covering uses 46 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005980 | ljcr-c-53-17-4

Covering design C(53, 17, 4)

the best known covering uses 220 blocks and the source-reported lower bound is 144.

来源状态 open | 2026-07-23 | 来源

V3P-005981 | ljcr-c-53-18-4

Covering design C(53, 18, 4)

the best known covering uses 137 blocks and the source-reported lower bound is 121.

来源状态 open | 2026-07-23 | 来源

V3P-005982 | ljcr-c-53-19-3

Covering design C(53, 19, 3)

the best known covering uses 37 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005983 | ljcr-c-53-19-4

Covering design C(53, 19, 4)

the best known covering uses 127 blocks and the source-reported lower bound is 106.

来源状态 open | 2026-07-23 | 来源

V3P-005984 | ljcr-c-53-20-3

Covering design C(53, 20, 3)

the best known covering uses 35 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-005985 | ljcr-c-53-20-4

Covering design C(53, 20, 4)

the best known covering uses 111 blocks and the source-reported lower bound is 88.

来源状态 open | 2026-07-23 | 来源

V3P-005986 | ljcr-c-53-21-3

Covering design C(53, 21, 3)

the best known covering uses 30 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-005987 | ljcr-c-53-21-4

Covering design C(53, 21, 4)

the best known covering uses 104 blocks and the source-reported lower bound is 74.

来源状态 open | 2026-07-23 | 来源

V3P-005988 | ljcr-c-53-22-3

Covering design C(53, 22, 3)

the best known covering uses 27 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-005989 | ljcr-c-53-22-4

Covering design C(53, 22, 4)

the best known covering uses 85 blocks and the source-reported lower bound is 61.

来源状态 open | 2026-07-23 | 来源

V3P-005990 | ljcr-c-53-23-3

Covering design C(53, 23, 3)

the best known covering uses 23 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-005991 | ljcr-c-53-23-4

Covering design C(53, 23, 4)

the best known covering uses 67 blocks and the source-reported lower bound is 51.

来源状态 open | 2026-07-23 | 来源

V3P-005992 | ljcr-c-53-23-5

Covering design C(53, 23, 5)

the best known covering uses 213 blocks and the source-reported lower bound is 120.

来源状态 open | 2026-07-23 | 来源

V3P-005993 | ljcr-c-53-24-3

Covering design C(53, 24, 3)

the best known covering uses 21 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-005994 | ljcr-c-53-24-4

Covering design C(53, 24, 4)

the best known covering uses 60 blocks and the source-reported lower bound is 42.

来源状态 open | 2026-07-23 | 来源

V3P-005995 | ljcr-c-53-24-5

Covering design C(53, 24, 5)

the best known covering uses 166 blocks and the source-reported lower bound is 98.

来源状态 open | 2026-07-23 | 来源

V3P-005996 | ljcr-c-53-25-4

Covering design C(53, 25, 4)

the best known covering uses 45 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-005997 | ljcr-c-53-25-5

Covering design C(53, 25, 5)

the best known covering uses 126 blocks and the source-reported lower bound is 79.

来源状态 open | 2026-07-23 | 来源

V3P-005998 | ljcr-c-53-5-2

Covering design C(53, 5, 2)

the best known covering uses 140 blocks and the source-reported lower bound is 139.

来源状态 open | 2026-07-23 | 来源

V3P-005999 | ljcr-c-53-7-2

Covering design C(53, 7, 2)

the best known covering uses 70 blocks and the source-reported lower bound is 69.

来源状态 open | 2026-07-23 | 来源

V3P-006000 | ljcr-c-53-8-2

Covering design C(53, 8, 2)

the best known covering uses 55 blocks and the source-reported lower bound is 53.

来源状态 open | 2026-07-23 | 来源

V3P-006001 | ljcr-c-53-9-2

Covering design C(53, 9, 2)

the best known covering uses 45 blocks and the source-reported lower bound is 43.

来源状态 open | 2026-07-23 | 来源

V3P-006002 | ljcr-c-54-10-2

Covering design C(54, 10, 2)

the best known covering uses 38 blocks and the source-reported lower bound is 36.

来源状态 open | 2026-07-23 | 来源

V3P-006003 | ljcr-c-54-11-3

Covering design C(54, 11, 3)

the best known covering uses 198 blocks and the source-reported lower bound is 172.

来源状态 open | 2026-07-23 | 来源

V3P-006004 | ljcr-c-54-12-2

Covering design C(54, 12, 2)

the best known covering uses 27 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-006005 | ljcr-c-54-12-3

Covering design C(54, 12, 3)

the best known covering uses 159 blocks and the source-reported lower bound is 131.

来源状态 open | 2026-07-23 | 来源

V3P-006006 | ljcr-c-54-13-2

Covering design C(54, 13, 2)

the best known covering uses 23 blocks and the source-reported lower bound is 21.

来源状态 open | 2026-07-23 | 来源

V3P-006007 | ljcr-c-54-13-3

Covering design C(54, 13, 3)

the best known covering uses 122 blocks and the source-reported lower bound is 104.

来源状态 open | 2026-07-23 | 来源

V3P-006008 | ljcr-c-54-14-3

Covering design C(54, 14, 3)

the best known covering uses 85 blocks and the source-reported lower bound is 81.

来源状态 open | 2026-07-23 | 来源

V3P-006009 | ljcr-c-54-15-2

Covering design C(54, 15, 2)

the best known covering uses 18 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-006010 | ljcr-c-54-15-3

Covering design C(54, 15, 3)

the best known covering uses 79 blocks and the source-reported lower bound is 69.

来源状态 open | 2026-07-23 | 来源

V3P-006011 | ljcr-c-54-16-2

Covering design C(54, 16, 2)

the best known covering uses 16 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-006012 | ljcr-c-54-16-3

Covering design C(54, 16, 3)

the best known covering uses 67 blocks and the source-reported lower bound is 54.

来源状态 open | 2026-07-23 | 来源

V3P-006013 | ljcr-c-54-17-3

Covering design C(54, 17, 3)

the best known covering uses 57 blocks and the source-reported lower bound is 45.

来源状态 open | 2026-07-23 | 来源

V3P-006014 | ljcr-c-54-17-4

Covering design C(54, 17, 4)

the best known covering uses 241 blocks and the source-reported lower bound is 159.

来源状态 open | 2026-07-23 | 来源

V3P-006015 | ljcr-c-54-18-4

Covering design C(54, 18, 4)

the best known covering uses 141 blocks and the source-reported lower bound is 126.

来源状态 open | 2026-07-23 | 来源

V3P-006016 | ljcr-c-54-19-3

Covering design C(54, 19, 3)

the best known covering uses 38 blocks and the source-reported lower bound is 36.

来源状态 open | 2026-07-23 | 来源

V3P-006017 | ljcr-c-54-19-4

Covering design C(54, 19, 4)

the best known covering uses 128 blocks and the source-reported lower bound is 111.

来源状态 open | 2026-07-23 | 来源

V3P-006018 | ljcr-c-54-20-3

Covering design C(54, 20, 3)

the best known covering uses 35 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-006019 | ljcr-c-54-20-4

Covering design C(54, 20, 4)

the best known covering uses 114 blocks and the source-reported lower bound is 92.

来源状态 open | 2026-07-23 | 来源

V3P-006020 | ljcr-c-54-21-3

Covering design C(54, 21, 3)

the best known covering uses 31 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-006021 | ljcr-c-54-21-4

Covering design C(54, 21, 4)

the best known covering uses 107 blocks and the source-reported lower bound is 78.

来源状态 open | 2026-07-23 | 来源

V3P-006022 | ljcr-c-54-22-3

Covering design C(54, 22, 3)

the best known covering uses 27 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-006023 | ljcr-c-54-22-4

Covering design C(54, 22, 4)

the best known covering uses 94 blocks and the source-reported lower bound is 64.

来源状态 open | 2026-07-23 | 来源

V3P-006024 | ljcr-c-54-23-3

Covering design C(54, 23, 3)

the best known covering uses 25 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-006025 | ljcr-c-54-23-4

Covering design C(54, 23, 4)

the best known covering uses 72 blocks and the source-reported lower bound is 52.

来源状态 open | 2026-07-23 | 来源

V3P-006026 | ljcr-c-54-24-3

Covering design C(54, 24, 3)

the best known covering uses 21 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-006027 | ljcr-c-54-24-4

Covering design C(54, 24, 4)

the best known covering uses 63 blocks and the source-reported lower bound is 43.

来源状态 open | 2026-07-23 | 来源

V3P-006028 | ljcr-c-54-24-5

Covering design C(54, 24, 5)

the best known covering uses 174 blocks and the source-reported lower bound is 115.

来源状态 open | 2026-07-23 | 来源

V3P-006029 | ljcr-c-54-25-3

Covering design C(54, 25, 3)

the best known covering uses 19 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-006030 | ljcr-c-54-25-4

Covering design C(54, 25, 4)

the best known covering uses 54 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-006031 | ljcr-c-54-25-5

Covering design C(54, 25, 5)

the best known covering uses 156 blocks and the source-reported lower bound is 91.

来源状态 open | 2026-07-23 | 来源

V3P-006032 | ljcr-c-54-7-2

Covering design C(54, 7, 2)

the best known covering uses 73 blocks and the source-reported lower bound is 70.

来源状态 open | 2026-07-23 | 来源

V3P-006033 | ljcr-c-54-8-2

Covering design C(54, 8, 2)

the best known covering uses 57 blocks and the source-reported lower bound is 54.

来源状态 open | 2026-07-23 | 来源

V3P-006034 | ljcr-c-54-9-2

Covering design C(54, 9, 2)

the best known covering uses 46 blocks and the source-reported lower bound is 44.

来源状态 open | 2026-07-23 | 来源

V3P-006035 | ljcr-c-55-10-2

Covering design C(55, 10, 2)

the best known covering uses 39 blocks and the source-reported lower bound is 36.

来源状态 open | 2026-07-23 | 来源

V3P-006036 | ljcr-c-55-11-2

Covering design C(55, 11, 2)

the best known covering uses 31 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-006037 | ljcr-c-55-11-3

Covering design C(55, 11, 3)

the best known covering uses 206 blocks and the source-reported lower bound is 180.

来源状态 open | 2026-07-23 | 来源

V3P-006038 | ljcr-c-55-12-2

Covering design C(55, 12, 2)

the best known covering uses 28 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-006039 | ljcr-c-55-12-3

Covering design C(55, 12, 3)

the best known covering uses 166 blocks and the source-reported lower bound is 138.

来源状态 open | 2026-07-23 | 来源

V3P-006040 | ljcr-c-55-13-2

Covering design C(55, 13, 2)

the best known covering uses 24 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-006041 | ljcr-c-55-13-3

Covering design C(55, 13, 3)

the best known covering uses 130 blocks and the source-reported lower bound is 110.

来源状态 open | 2026-07-23 | 来源

V3P-006042 | ljcr-c-55-14-2

Covering design C(55, 14, 2)

the best known covering uses 21 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-006043 | ljcr-c-55-14-3

Covering design C(55, 14, 3)

the best known covering uses 99 blocks and the source-reported lower bound is 83.

来源状态 open | 2026-07-23 | 来源

V3P-006044 | ljcr-c-55-15-2

Covering design C(55, 15, 2)

the best known covering uses 19 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-006045 | ljcr-c-55-15-3

Covering design C(55, 15, 3)

the best known covering uses 81 blocks and the source-reported lower bound is 74.

来源状态 open | 2026-07-23 | 来源

V3P-006046 | ljcr-c-55-16-2

Covering design C(55, 16, 2)

the best known covering uses 16 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-006047 | ljcr-c-55-16-3

Covering design C(55, 16, 3)

the best known covering uses 72 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-006048 | ljcr-c-55-17-3

Covering design C(55, 17, 3)

the best known covering uses 59 blocks and the source-reported lower bound is 46.

来源状态 open | 2026-07-23 | 来源

V3P-006049 | ljcr-c-55-18-3

Covering design C(55, 18, 3)

the best known covering uses 48 blocks and the source-reported lower bound is 43.

来源状态 open | 2026-07-23 | 来源

V3P-006050 | ljcr-c-55-18-4

Covering design C(55, 18, 4)

the best known covering uses 196 blocks and the source-reported lower bound is 138.

来源状态 open | 2026-07-23 | 来源

V3P-006051 | ljcr-c-55-19-3

Covering design C(55, 19, 3)

the best known covering uses 38 blocks and the source-reported lower bound is 37.

来源状态 open | 2026-07-23 | 来源

V3P-006052 | ljcr-c-55-19-4

Covering design C(55, 19, 4)

the best known covering uses 131 blocks and the source-reported lower bound is 113.

来源状态 open | 2026-07-23 | 来源

V3P-006053 | ljcr-c-55-20-3

Covering design C(55, 20, 3)

the best known covering uses 36 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-006054 | ljcr-c-55-20-4

Covering design C(55, 20, 4)

the best known covering uses 118 blocks and the source-reported lower bound is 99.

来源状态 open | 2026-07-23 | 来源

V3P-006055 | ljcr-c-55-21-3

Covering design C(55, 21, 3)

the best known covering uses 33 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-006056 | ljcr-c-55-21-4

Covering design C(55, 21, 4)

the best known covering uses 110 blocks and the source-reported lower bound is 79.

来源状态 open | 2026-07-23 | 来源

V3P-006057 | ljcr-c-55-22-3

Covering design C(55, 22, 3)

the best known covering uses 30 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-006058 | ljcr-c-55-22-4

Covering design C(55, 22, 4)

the best known covering uses 99 blocks and the source-reported lower bound is 73.

来源状态 open | 2026-07-23 | 来源

V3P-006059 | ljcr-c-55-23-3

Covering design C(55, 23, 3)

the best known covering uses 25 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-006060 | ljcr-c-55-23-4

Covering design C(55, 23, 4)

the best known covering uses 78 blocks and the source-reported lower bound is 60.

来源状态 open | 2026-07-23 | 来源

V3P-006061 | ljcr-c-55-24-3

Covering design C(55, 24, 3)

the best known covering uses 23 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-006062 | ljcr-c-55-24-4

Covering design C(55, 24, 4)

the best known covering uses 68 blocks and the source-reported lower bound is 51.

来源状态 open | 2026-07-23 | 来源

V3P-006063 | ljcr-c-55-24-5

Covering design C(55, 24, 5)

the best known covering uses 195 blocks and the source-reported lower bound is 120.

来源状态 open | 2026-07-23 | 来源

V3P-006064 | ljcr-c-55-25-3

Covering design C(55, 25, 3)

the best known covering uses 20 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-006065 | ljcr-c-55-25-4

Covering design C(55, 25, 4)

the best known covering uses 58 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-006066 | ljcr-c-55-25-5

Covering design C(55, 25, 5)

the best known covering uses 169 blocks and the source-reported lower bound is 95.

来源状态 open | 2026-07-23 | 来源

V3P-006067 | ljcr-c-55-6-2

Covering design C(55, 6, 2)

the best known covering uses 105 blocks and the source-reported lower bound is 101.

来源状态 open | 2026-07-23 | 来源

V3P-006068 | ljcr-c-55-7-2

Covering design C(55, 7, 2)

the best known covering uses 76 blocks and the source-reported lower bound is 72.

来源状态 open | 2026-07-23 | 来源

V3P-006069 | ljcr-c-55-8-2

Covering design C(55, 8, 2)

the best known covering uses 57 blocks and the source-reported lower bound is 55.

来源状态 open | 2026-07-23 | 来源

V3P-006070 | ljcr-c-55-9-2

Covering design C(55, 9, 2)

the best known covering uses 49 blocks and the source-reported lower bound is 45.

来源状态 open | 2026-07-23 | 来源

V3P-006071 | ljcr-c-56-11-3

Covering design C(56, 11, 3)

the best known covering uses 221 blocks and the source-reported lower bound is 184.

来源状态 open | 2026-07-23 | 来源

V3P-006072 | ljcr-c-56-12-2

Covering design C(56, 12, 2)

the best known covering uses 28 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-006073 | ljcr-c-56-12-3

Covering design C(56, 12, 3)

the best known covering uses 175 blocks and the source-reported lower bound is 140.

来源状态 open | 2026-07-23 | 来源

V3P-006074 | ljcr-c-56-13-2

Covering design C(56, 13, 2)

the best known covering uses 25 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-006075 | ljcr-c-56-13-3

Covering design C(56, 13, 3)

the best known covering uses 136 blocks and the source-reported lower bound is 117.

来源状态 open | 2026-07-23 | 来源

V3P-006076 | ljcr-c-56-14-2

Covering design C(56, 14, 2)

the best known covering uses 21 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-006077 | ljcr-c-56-14-3

Covering design C(56, 14, 3)

the best known covering uses 103 blocks and the source-reported lower bound is 88.

来源状态 open | 2026-07-23 | 来源

V3P-006078 | ljcr-c-56-15-2

Covering design C(56, 15, 2)

the best known covering uses 19 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-006079 | ljcr-c-56-15-3

Covering design C(56, 15, 3)

the best known covering uses 83 blocks and the source-reported lower bound is 75.

来源状态 open | 2026-07-23 | 来源

V3P-006080 | ljcr-c-56-16-2

Covering design C(56, 16, 2)

the best known covering uses 17 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-006081 | ljcr-c-56-16-3

Covering design C(56, 16, 3)

the best known covering uses 73 blocks and the source-reported lower bound is 63.

来源状态 open | 2026-07-23 | 来源

V3P-006082 | ljcr-c-56-17-2

Covering design C(56, 17, 2)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-006083 | ljcr-c-56-17-3

Covering design C(56, 17, 3)

the best known covering uses 64 blocks and the source-reported lower bound is 50.

来源状态 open | 2026-07-23 | 来源

V3P-006084 | ljcr-c-56-18-3

Covering design C(56, 18, 3)

the best known covering uses 49 blocks and the source-reported lower bound is 44.

来源状态 open | 2026-07-23 | 来源

V3P-006085 | ljcr-c-56-18-4

Covering design C(56, 18, 4)

the best known covering uses 208 blocks and the source-reported lower bound is 144.

来源状态 open | 2026-07-23 | 来源

V3P-006086 | ljcr-c-56-19-3

Covering design C(56, 19, 3)

the best known covering uses 42 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-006087 | ljcr-c-56-19-4

Covering design C(56, 19, 4)

the best known covering uses 149 blocks and the source-reported lower bound is 127.

来源状态 open | 2026-07-23 | 来源

V3P-006088 | ljcr-c-56-20-3

Covering design C(56, 20, 3)

the best known covering uses 36 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-006089 | ljcr-c-56-20-4

Covering design C(56, 20, 4)

the best known covering uses 120 blocks and the source-reported lower bound is 104.

来源状态 open | 2026-07-23 | 来源

V3P-006090 | ljcr-c-56-21-3

Covering design C(56, 21, 3)

the best known covering uses 35 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-006091 | ljcr-c-56-21-4

Covering design C(56, 21, 4)

the best known covering uses 113 blocks and the source-reported lower bound is 88.

来源状态 open | 2026-07-23 | 来源

V3P-006092 | ljcr-c-56-22-3

Covering design C(56, 22, 3)

the best known covering uses 31 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-006093 | ljcr-c-56-22-4

Covering design C(56, 22, 4)

the best known covering uses 103 blocks and the source-reported lower bound is 74.

来源状态 open | 2026-07-23 | 来源

V3P-006094 | ljcr-c-56-23-3

Covering design C(56, 23, 3)

the best known covering uses 28 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-006095 | ljcr-c-56-23-4

Covering design C(56, 23, 4)

the best known covering uses 93 blocks and the source-reported lower bound is 61.

来源状态 open | 2026-07-23 | 来源

V3P-006096 | ljcr-c-56-24-3

Covering design C(56, 24, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-006097 | ljcr-c-56-24-4

Covering design C(56, 24, 4)

the best known covering uses 72 blocks and the source-reported lower bound is 52.

来源状态 open | 2026-07-23 | 来源

V3P-006098 | ljcr-c-56-24-5

Covering design C(56, 24, 5)

the best known covering uses 240 blocks and the source-reported lower bound is 140.

来源状态 open | 2026-07-23 | 来源

V3P-006099 | ljcr-c-56-25-3

Covering design C(56, 25, 3)

the best known covering uses 21 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-006100 | ljcr-c-56-25-4

Covering design C(56, 25, 4)

the best known covering uses 64 blocks and the source-reported lower bound is 43.

来源状态 open | 2026-07-23 | 来源

V3P-006101 | ljcr-c-56-25-5

Covering design C(56, 25, 5)

the best known covering uses 184 blocks and the source-reported lower bound is 115.

来源状态 open | 2026-07-23 | 来源

V3P-006102 | ljcr-c-56-6-2

Covering design C(56, 6, 2)

the best known covering uses 110 blocks and the source-reported lower bound is 103.

来源状态 open | 2026-07-23 | 来源

V3P-006103 | ljcr-c-56-8-2

Covering design C(56, 8, 2)

the best known covering uses 57 blocks and the source-reported lower bound is 56.

来源状态 open | 2026-07-23 | 来源

V3P-006104 | ljcr-c-56-9-2

Covering design C(56, 9, 2)

the best known covering uses 49 blocks and the source-reported lower bound is 46.

来源状态 open | 2026-07-23 | 来源

V3P-006105 | ljcr-c-57-10-2

Covering design C(57, 10, 2)

the best known covering uses 42 blocks and the source-reported lower bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-006106 | ljcr-c-57-11-2

Covering design C(57, 11, 2)

the best known covering uses 36 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-006107 | ljcr-c-57-11-3

Covering design C(57, 11, 3)

the best known covering uses 245 blocks and the source-reported lower bound is 208.

来源状态 open | 2026-07-23 | 来源

V3P-006108 | ljcr-c-57-12-3

Covering design C(57, 12, 3)

the best known covering uses 185 blocks and the source-reported lower bound is 148.

来源状态 open | 2026-07-23 | 来源

V3P-006109 | ljcr-c-57-13-2

Covering design C(57, 13, 2)

the best known covering uses 25 blocks and the source-reported lower bound is 24.

来源状态 open | 2026-07-23 | 来源

V3P-006110 | ljcr-c-57-13-3

Covering design C(57, 13, 3)

the best known covering uses 142 blocks and the source-reported lower bound is 119.

来源状态 open | 2026-07-23 | 来源

V3P-006111 | ljcr-c-57-14-3

Covering design C(57, 14, 3)

the best known covering uses 109 blocks and the source-reported lower bound is 90.

来源状态 open | 2026-07-23 | 来源

V3P-006112 | ljcr-c-57-15-3

Covering design C(57, 15, 3)

the best known covering uses 84 blocks and the source-reported lower bound is 76.

来源状态 open | 2026-07-23 | 来源

V3P-006113 | ljcr-c-57-16-2

Covering design C(57, 16, 2)

the best known covering uses 18 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-006114 | ljcr-c-57-16-3

Covering design C(57, 16, 3)

the best known covering uses 79 blocks and the source-reported lower bound is 65.

来源状态 open | 2026-07-23 | 来源

V3P-006115 | ljcr-c-57-17-2

Covering design C(57, 17, 2)

the best known covering uses 16 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-006116 | ljcr-c-57-17-3

Covering design C(57, 17, 3)

the best known covering uses 66 blocks and the source-reported lower bound is 51.

来源状态 open | 2026-07-23 | 来源

V3P-006117 | ljcr-c-57-18-3

Covering design C(57, 18, 3)

the best known covering uses 53 blocks and the source-reported lower bound is 45.

来源状态 open | 2026-07-23 | 来源

V3P-006118 | ljcr-c-57-18-4

Covering design C(57, 18, 4)

the best known covering uses 220 blocks and the source-reported lower bound is 159.

来源状态 open | 2026-07-23 | 来源

V3P-006119 | ljcr-c-57-19-3

Covering design C(57, 19, 3)

the best known covering uses 43 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-006120 | ljcr-c-57-19-4

Covering design C(57, 19, 4)

the best known covering uses 166 blocks and the source-reported lower bound is 132.

来源状态 open | 2026-07-23 | 来源

V3P-006121 | ljcr-c-57-20-3

Covering design C(57, 20, 3)

the best known covering uses 39 blocks and the source-reported lower bound is 36.

来源状态 open | 2026-07-23 | 来源

V3P-006122 | ljcr-c-57-20-4

Covering design C(57, 20, 4)

the best known covering uses 131 blocks and the source-reported lower bound is 112.

来源状态 open | 2026-07-23 | 来源

V3P-006123 | ljcr-c-57-21-3

Covering design C(57, 21, 3)

the best known covering uses 35 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-006124 | ljcr-c-57-21-4

Covering design C(57, 21, 4)

the best known covering uses 115 blocks and the source-reported lower bound is 93.

来源状态 open | 2026-07-23 | 来源

V3P-006125 | ljcr-c-57-22-3

Covering design C(57, 22, 3)

the best known covering uses 32 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-006126 | ljcr-c-57-22-4

Covering design C(57, 22, 4)

the best known covering uses 109 blocks and the source-reported lower bound is 78.

来源状态 open | 2026-07-23 | 来源

V3P-006127 | ljcr-c-57-23-3

Covering design C(57, 23, 3)

the best known covering uses 28 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-006128 | ljcr-c-57-23-4

Covering design C(57, 23, 4)

the best known covering uses 98 blocks and the source-reported lower bound is 65.

来源状态 open | 2026-07-23 | 来源

V3P-006129 | ljcr-c-57-24-3

Covering design C(57, 24, 3)

the best known covering uses 25 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-006130 | ljcr-c-57-24-4

Covering design C(57, 24, 4)

the best known covering uses 79 blocks and the source-reported lower bound is 60.

来源状态 open | 2026-07-23 | 来源

V3P-006131 | ljcr-c-57-25-3

Covering design C(57, 25, 3)

the best known covering uses 22 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-006132 | ljcr-c-57-25-4

Covering design C(57, 25, 4)

the best known covering uses 69 blocks and the source-reported lower bound is 44.

来源状态 open | 2026-07-23 | 来源

V3P-006133 | ljcr-c-57-25-5

Covering design C(57, 25, 5)

the best known covering uses 201 blocks and the source-reported lower bound is 119.

来源状态 open | 2026-07-23 | 来源

V3P-006134 | ljcr-c-57-5-2

Covering design C(57, 5, 2)

the best known covering uses 161 blocks and the source-reported lower bound is 160.

来源状态 open | 2026-07-23 | 来源

V3P-006135 | ljcr-c-57-9-2

Covering design C(57, 9, 2)

the best known covering uses 51 blocks and the source-reported lower bound is 47.

来源状态 open | 2026-07-23 | 来源

V3P-006136 | ljcr-c-58-10-2

Covering design C(58, 10, 2)

the best known covering uses 43 blocks and the source-reported lower bound is 41.

来源状态 open | 2026-07-23 | 来源

V3P-006137 | ljcr-c-58-11-2

Covering design C(58, 11, 2)

the best known covering uses 37 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-006138 | ljcr-c-58-12-2

Covering design C(58, 12, 2)

the best known covering uses 30 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-006139 | ljcr-c-58-12-3

Covering design C(58, 12, 3)

the best known covering uses 193 blocks and the source-reported lower bound is 165.

来源状态 open | 2026-07-23 | 来源

V3P-006140 | ljcr-c-58-13-2

Covering design C(58, 13, 2)

the best known covering uses 27 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-006141 | ljcr-c-58-13-3

Covering design C(58, 13, 3)

the best known covering uses 146 blocks and the source-reported lower bound is 130.

来源状态 open | 2026-07-23 | 来源

V3P-006142 | ljcr-c-58-14-3

Covering design C(58, 14, 3)

the best known covering uses 115 blocks and the source-reported lower bound is 100.

来源状态 open | 2026-07-23 | 来源

V3P-006143 | ljcr-c-58-15-3

Covering design C(58, 15, 3)

the best known covering uses 86 blocks and the source-reported lower bound is 82.

来源状态 open | 2026-07-23 | 来源

V3P-006144 | ljcr-c-58-16-2

Covering design C(58, 16, 2)

the best known covering uses 18 blocks and the source-reported lower bound is 17.

来源状态 open | 2026-07-23 | 来源

V3P-006145 | ljcr-c-58-16-3

Covering design C(58, 16, 3)

the best known covering uses 80 blocks and the source-reported lower bound is 69.

来源状态 open | 2026-07-23 | 来源

V3P-006146 | ljcr-c-58-17-2

Covering design C(58, 17, 2)

the best known covering uses 16 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-006147 | ljcr-c-58-17-3

Covering design C(58, 17, 3)

the best known covering uses 71 blocks and the source-reported lower bound is 55.

来源状态 open | 2026-07-23 | 来源

V3P-006148 | ljcr-c-58-18-3

Covering design C(58, 18, 3)

the best known covering uses 58 blocks and the source-reported lower bound is 46.

来源状态 open | 2026-07-23 | 来源

V3P-006149 | ljcr-c-58-18-4

Covering design C(58, 18, 4)

the best known covering uses 226 blocks and the source-reported lower bound is 165.

来源状态 open | 2026-07-23 | 来源

V3P-006150 | ljcr-c-58-19-3

Covering design C(58, 19, 3)

the best known covering uses 44 blocks and the source-reported lower bound is 41.

来源状态 open | 2026-07-23 | 来源

V3P-006151 | ljcr-c-58-19-4

Covering design C(58, 19, 4)

the best known covering uses 178 blocks and the source-reported lower bound is 138.

来源状态 open | 2026-07-23 | 来源

V3P-006152 | ljcr-c-58-20-3

Covering design C(58, 20, 3)

the best known covering uses 39 blocks and the source-reported lower bound is 38.

来源状态 open | 2026-07-23 | 来源

V3P-006153 | ljcr-c-58-20-4

Covering design C(58, 20, 4)

the best known covering uses 137 blocks and the source-reported lower bound is 114.

来源状态 open | 2026-07-23 | 来源

V3P-006154 | ljcr-c-58-21-3

Covering design C(58, 21, 3)

the best known covering uses 37 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-006155 | ljcr-c-58-21-4

Covering design C(58, 21, 4)

the best known covering uses 121 blocks and the source-reported lower bound is 100.

来源状态 open | 2026-07-23 | 来源

V3P-006156 | ljcr-c-58-22-3

Covering design C(58, 22, 3)

the best known covering uses 33 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-006157 | ljcr-c-58-22-4

Covering design C(58, 22, 4)

the best known covering uses 112 blocks and the source-reported lower bound is 80.

来源状态 open | 2026-07-23 | 来源

V3P-006158 | ljcr-c-58-23-3

Covering design C(58, 23, 3)

the best known covering uses 31 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-006159 | ljcr-c-58-23-4

Covering design C(58, 23, 4)

the best known covering uses 102 blocks and the source-reported lower bound is 74.

来源状态 open | 2026-07-23 | 来源

V3P-006160 | ljcr-c-58-24-3

Covering design C(58, 24, 3)

the best known covering uses 27 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-006161 | ljcr-c-58-24-4

Covering design C(58, 24, 4)

the best known covering uses 88 blocks and the source-reported lower bound is 61.

来源状态 open | 2026-07-23 | 来源

V3P-006162 | ljcr-c-58-25-3

Covering design C(58, 25, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 19.

来源状态 open | 2026-07-23 | 来源

V3P-006163 | ljcr-c-58-25-4

Covering design C(58, 25, 4)

the best known covering uses 72 blocks and the source-reported lower bound is 52.

来源状态 open | 2026-07-23 | 来源

V3P-006164 | ljcr-c-58-25-5

Covering design C(58, 25, 5)

the best known covering uses 225 blocks and the source-reported lower bound is 140.

来源状态 open | 2026-07-23 | 来源

V3P-006165 | ljcr-c-58-6-2

Covering design C(58, 6, 2)

the best known covering uses 117 blocks and the source-reported lower bound is 116.

来源状态 open | 2026-07-23 | 来源

V3P-006166 | ljcr-c-58-7-2

Covering design C(58, 7, 2)

the best known covering uses 84 blocks and the source-reported lower bound is 83.

来源状态 open | 2026-07-23 | 来源

V3P-006167 | ljcr-c-59-10-2

Covering design C(59, 10, 2)

the best known covering uses 46 blocks and the source-reported lower bound is 43.

来源状态 open | 2026-07-23 | 来源

V3P-006168 | ljcr-c-59-11-2

Covering design C(59, 11, 2)

the best known covering uses 37 blocks and the source-reported lower bound is 35.

来源状态 open | 2026-07-23 | 来源

V3P-006169 | ljcr-c-59-12-2

Covering design C(59, 12, 2)

the best known covering uses 31 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-006170 | ljcr-c-59-12-3

Covering design C(59, 12, 3)

the best known covering uses 198 blocks and the source-reported lower bound is 168.

来源状态 open | 2026-07-23 | 来源

V3P-006171 | ljcr-c-59-13-2

Covering design C(59, 13, 2)

the best known covering uses 28 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-006172 | ljcr-c-59-13-3

Covering design C(59, 13, 3)

the best known covering uses 161 blocks and the source-reported lower bound is 132.

来源状态 open | 2026-07-23 | 来源

V3P-006173 | ljcr-c-59-14-2

Covering design C(59, 14, 2)

the best known covering uses 24 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-006174 | ljcr-c-59-14-3

Covering design C(59, 14, 3)

the best known covering uses 128 blocks and the source-reported lower bound is 106.

来源状态 open | 2026-07-23 | 来源

V3P-006175 | ljcr-c-59-15-3

Covering design C(59, 15, 3)

the best known covering uses 89 blocks and the source-reported lower bound is 83.

来源状态 open | 2026-07-23 | 来源

V3P-006176 | ljcr-c-59-16-2

Covering design C(59, 16, 2)

the best known covering uses 19 blocks and the source-reported lower bound is 18.

来源状态 open | 2026-07-23 | 来源

V3P-006177 | ljcr-c-59-16-3

Covering design C(59, 16, 3)

the best known covering uses 81 blocks and the source-reported lower bound is 74.

来源状态 open | 2026-07-23 | 来源

V3P-006178 | ljcr-c-59-17-2

Covering design C(59, 17, 2)

the best known covering uses 17 blocks and the source-reported lower bound is 15.

来源状态 open | 2026-07-23 | 来源

V3P-006179 | ljcr-c-59-17-3

Covering design C(59, 17, 3)

the best known covering uses 73 blocks and the source-reported lower bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-006180 | ljcr-c-59-18-2

Covering design C(59, 18, 2)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-006181 | ljcr-c-59-18-3

Covering design C(59, 18, 3)

the best known covering uses 61 blocks and the source-reported lower bound is 47.

来源状态 open | 2026-07-23 | 来源

V3P-006182 | ljcr-c-59-19-3

Covering design C(59, 19, 3)

the best known covering uses 51 blocks and the source-reported lower bound is 44.

来源状态 open | 2026-07-23 | 来源

V3P-006183 | ljcr-c-59-19-4

Covering design C(59, 19, 4)

the best known covering uses 212 blocks and the source-reported lower bound is 143.

来源状态 open | 2026-07-23 | 来源

V3P-006184 | ljcr-c-59-20-3

Covering design C(59, 20, 3)

the best known covering uses 40 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-006185 | ljcr-c-59-20-4

Covering design C(59, 20, 4)

the best known covering uses 144 blocks and the source-reported lower bound is 121.

来源状态 open | 2026-07-23 | 来源

V3P-006186 | ljcr-c-59-21-3

Covering design C(59, 21, 3)

the best known covering uses 38 blocks and the source-reported lower bound is 34.

来源状态 open | 2026-07-23 | 来源

V3P-006187 | ljcr-c-59-21-4

Covering design C(59, 21, 4)

the best known covering uses 125 blocks and the source-reported lower bound is 107.

来源状态 open | 2026-07-23 | 来源

V3P-006188 | ljcr-c-59-22-3

Covering design C(59, 22, 3)

the best known covering uses 35 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-006189 | ljcr-c-59-22-4

Covering design C(59, 22, 4)

the best known covering uses 114 blocks and the source-reported lower bound is 92.

来源状态 open | 2026-07-23 | 来源

V3P-006190 | ljcr-c-59-23-3

Covering design C(59, 23, 3)

the best known covering uses 31 blocks and the source-reported lower bound is 26.

来源状态 open | 2026-07-23 | 来源

V3P-006191 | ljcr-c-59-23-4

Covering design C(59, 23, 4)

the best known covering uses 108 blocks and the source-reported lower bound is 75.

来源状态 open | 2026-07-23 | 来源

V3P-006192 | ljcr-c-59-24-3

Covering design C(59, 24, 3)

the best known covering uses 28 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-006193 | ljcr-c-59-24-4

Covering design C(59, 24, 4)

the best known covering uses 92 blocks and the source-reported lower bound is 64.

来源状态 open | 2026-07-23 | 来源

V3P-006194 | ljcr-c-59-25-3

Covering design C(59, 25, 3)

the best known covering uses 24 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-006195 | ljcr-c-59-25-4

Covering design C(59, 25, 4)

the best known covering uses 78 blocks and the source-reported lower bound is 53.

来源状态 open | 2026-07-23 | 来源

V3P-006196 | ljcr-c-59-6-2

Covering design C(59, 6, 2)

the best known covering uses 119 blocks and the source-reported lower bound is 118.

来源状态 open | 2026-07-23 | 来源

V3P-006197 | ljcr-c-59-7-2

Covering design C(59, 7, 2)

the best known covering uses 86 blocks and the source-reported lower bound is 85.

来源状态 open | 2026-07-23 | 来源

V3P-006198 | ljcr-c-59-9-2

Covering design C(59, 9, 2)

the best known covering uses 55 blocks and the source-reported lower bound is 54.

来源状态 open | 2026-07-23 | 来源

V3P-006199 | ljcr-c-60-10-2

Covering design C(60, 10, 2)

the best known covering uses 47 blocks and the source-reported lower bound is 44.

来源状态 open | 2026-07-23 | 来源

V3P-006200 | ljcr-c-60-11-2

Covering design C(60, 11, 2)

the best known covering uses 38 blocks and the source-reported lower bound is 36.

来源状态 open | 2026-07-23 | 来源

V3P-006201 | ljcr-c-60-12-2

Covering design C(60, 12, 2)

the best known covering uses 31 blocks and the source-reported lower bound is 30.

来源状态 open | 2026-07-23 | 来源

V3P-006202 | ljcr-c-60-12-3

Covering design C(60, 12, 3)

the best known covering uses 205 blocks and the source-reported lower bound is 175.

来源状态 open | 2026-07-23 | 来源

V3P-006203 | ljcr-c-60-13-2

Covering design C(60, 13, 2)

the best known covering uses 28 blocks and the source-reported lower bound is 27.

来源状态 open | 2026-07-23 | 来源

V3P-006204 | ljcr-c-60-13-3

Covering design C(60, 13, 3)

the best known covering uses 165 blocks and the source-reported lower bound is 139.

来源状态 open | 2026-07-23 | 来源

V3P-006205 | ljcr-c-60-14-2

Covering design C(60, 14, 2)

the best known covering uses 25 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-006206 | ljcr-c-60-14-3

Covering design C(60, 14, 3)

the best known covering uses 133 blocks and the source-reported lower bound is 112.

来源状态 open | 2026-07-23 | 来源

V3P-006207 | ljcr-c-60-15-2

Covering design C(60, 15, 2)

the best known covering uses 21 blocks and the source-reported lower bound is 20.

来源状态 open | 2026-07-23 | 来源

V3P-006208 | ljcr-c-60-15-3

Covering design C(60, 15, 3)

the best known covering uses 104 blocks and the source-reported lower bound is 88.

来源状态 open | 2026-07-23 | 来源

V3P-006209 | ljcr-c-60-16-3

Covering design C(60, 16, 3)

the best known covering uses 83 blocks and the source-reported lower bound is 75.

来源状态 open | 2026-07-23 | 来源

V3P-006210 | ljcr-c-60-17-2

Covering design C(60, 17, 2)

the best known covering uses 18 blocks and the source-reported lower bound is 16.

来源状态 open | 2026-07-23 | 来源

V3P-006211 | ljcr-c-60-17-3

Covering design C(60, 17, 3)

the best known covering uses 76 blocks and the source-reported lower bound is 64.

来源状态 open | 2026-07-23 | 来源

V3P-006212 | ljcr-c-60-18-2

Covering design C(60, 18, 2)

the best known covering uses 15 blocks and the source-reported lower bound is 14.

来源状态 open | 2026-07-23 | 来源

V3P-006213 | ljcr-c-60-18-3

Covering design C(60, 18, 3)

the best known covering uses 65 blocks and the source-reported lower bound is 50.

来源状态 open | 2026-07-23 | 来源

V3P-006214 | ljcr-c-60-19-3

Covering design C(60, 19, 3)

the best known covering uses 53 blocks and the source-reported lower bound is 45.

来源状态 open | 2026-07-23 | 来源

V3P-006215 | ljcr-c-60-19-4

Covering design C(60, 19, 4)

the best known covering uses 228 blocks and the source-reported lower bound is 149.

来源状态 open | 2026-07-23 | 来源

V3P-006216 | ljcr-c-60-20-3

Covering design C(60, 20, 3)

the best known covering uses 44 blocks and the source-reported lower bound is 39.

来源状态 open | 2026-07-23 | 来源

V3P-006217 | ljcr-c-60-20-4

Covering design C(60, 20, 4)

the best known covering uses 165 blocks and the source-reported lower bound is 132.

来源状态 open | 2026-07-23 | 来源

V3P-006218 | ljcr-c-60-21-3

Covering design C(60, 21, 3)

the best known covering uses 38 blocks and the source-reported lower bound is 36.

来源状态 open | 2026-07-23 | 来源

V3P-006219 | ljcr-c-60-21-4

Covering design C(60, 21, 4)

the best known covering uses 132 blocks and the source-reported lower bound is 112.

来源状态 open | 2026-07-23 | 来源

V3P-006220 | ljcr-c-60-22-3

Covering design C(60, 22, 3)

the best known covering uses 36 blocks and the source-reported lower bound is 33.

来源状态 open | 2026-07-23 | 来源

V3P-006221 | ljcr-c-60-22-4

Covering design C(60, 22, 4)

the best known covering uses 116 blocks and the source-reported lower bound is 93.

来源状态 open | 2026-07-23 | 来源

V3P-006222 | ljcr-c-60-23-3

Covering design C(60, 23, 3)

the best known covering uses 32 blocks and the source-reported lower bound is 29.

来源状态 open | 2026-07-23 | 来源

V3P-006223 | ljcr-c-60-23-4

Covering design C(60, 23, 4)

the best known covering uses 111 blocks and the source-reported lower bound is 79.

来源状态 open | 2026-07-23 | 来源

V3P-006224 | ljcr-c-60-24-3

Covering design C(60, 24, 3)

the best known covering uses 28 blocks and the source-reported lower bound is 25.

来源状态 open | 2026-07-23 | 来源

V3P-006225 | ljcr-c-60-24-4

Covering design C(60, 24, 4)

the best known covering uses 93 blocks and the source-reported lower bound is 65.

来源状态 open | 2026-07-23 | 来源

V3P-006226 | ljcr-c-60-25-3

Covering design C(60, 25, 3)

the best known covering uses 26 blocks and the source-reported lower bound is 22.

来源状态 open | 2026-07-23 | 来源

V3P-006227 | ljcr-c-60-25-4

Covering design C(60, 25, 4)

the best known covering uses 84 blocks and the source-reported lower bound is 60.

来源状态 open | 2026-07-23 | 来源

V3P-006228 | ljcr-c-60-6-2

Covering design C(60, 6, 2)

the best known covering uses 128 blocks and the source-reported lower bound is 120.

来源状态 open | 2026-07-23 | 来源

V3P-006229 | ljcr-c-60-7-2

Covering design C(60, 7, 2)

the best known covering uses 91 blocks and the source-reported lower bound is 86.

来源状态 open | 2026-07-23 | 来源

V3P-006230 | ljcr-c-60-9-2

Covering design C(60, 9, 2)

the best known covering uses 57 blocks and the source-reported lower bound is 54.

来源状态 open | 2026-07-23 | 来源

D05 阿贝尔差集 | 3,064 行

给定阿贝尔群及 (v,k,lambda)，问是否存在大小 k 的子集，使每个非零群元素恰好由 lambda 个有序差表示。通过必要条件检验不代表一定存在。

原始维护者：Daniel M. Gordon

The dataset ships a CITATION.cff asking that it be cited using that file's metadata: Daniel M. Gordon, "La Jolla Difference Sets Repository", dataset, version 1.0, 2024, https://doi.org/10.5281/zenodo.10775931. The CC-BY-4.0 licence also requires stating what was changed: only open parameter sets are kept here, and no difference sets are copied.

原始问题族入口

V3P-006231 | ljcr-ds-10005-2624-688-10005

(10005, 2624, 688)-difference set in Z_10005

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006232 | ljcr-ds-10011-3081-948-10011

(10011, 3081, 948)-difference set in Z_10011

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006233 | ljcr-ds-10021-3841-1472-10021

(10021, 3841, 1472)-difference set in Z_10021

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006234 | ljcr-ds-10043-5021-2510-10043

(10043, 5021, 2510)-difference set in Z_10043

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006235 | ljcr-ds-10075-1242-153-10075

(10075, 1242, 153)-difference set in Z_10075

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006236 | ljcr-ds-1008-266-70-12-84

(1008, 266, 70)-difference set in Z_12 x Z_84

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006237 | ljcr-ds-1008-266-70-2-2-2-126

(1008, 266, 70)-difference set in Z_2 x Z_2 x Z_2 x Z_126

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006238 | ljcr-ds-1008-266-70-2-2-252

(1008, 266, 70)-difference set in Z_2 x Z_2 x Z_252

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006239 | ljcr-ds-1008-266-70-2-2-6-42

(1008, 266, 70)-difference set in Z_2 x Z_2 x Z_6 x Z_42

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006240 | ljcr-ds-1008-266-70-2-504

(1008, 266, 70)-difference set in Z_2 x Z_504

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006241 | ljcr-ds-1008-266-70-2-6-84

(1008, 266, 70)-difference set in Z_2 x Z_6 x Z_84

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006242 | ljcr-ds-1008-266-70-4-252

(1008, 266, 70)-difference set in Z_4 x Z_252

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006243 | ljcr-ds-1008-266-70-6-168

(1008, 266, 70)-difference set in Z_6 x Z_168

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006244 | ljcr-ds-10089-2328-537-10089

(10089, 2328, 537)-difference set in Z_10089

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006245 | ljcr-ds-10107-5053-2526-10107

(10107, 5053, 2526)-difference set in Z_10107

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006246 | ljcr-ds-1012-337-112-1012

(1012, 337, 112)-difference set in Z_1012

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006247 | ljcr-ds-10143-462-21-21-483

(10143, 462, 21)-difference set in Z_21 x Z_483

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006248 | ljcr-ds-10143-462-21-3-3381

(10143, 462, 21)-difference set in Z_3 x Z_3381

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006249 | ljcr-ds-10143-462-21-7-1449

(10143, 462, 21)-difference set in Z_7 x Z_1449

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006250 | ljcr-ds-10171-2035-407-10171

(10171, 2035, 407)-difference set in Z_10171

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006251 | ljcr-ds-10179-728-52-10179

(10179, 728, 52)-difference set in Z_10179

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006252 | ljcr-ds-10179-728-52-3-3-1131

(10179, 728, 52)-difference set in Z_3 x Z_3 x Z_1131

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006253 | ljcr-ds-10179-728-52-3-3393

(10179, 728, 52)-difference set in Z_3 x Z_3393

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006254 | ljcr-ds-1021-340-113-1021

(1021, 340, 113)-difference set in Z_1021

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006255 | ljcr-ds-10225-4545-2020-10225

(10225, 4545, 2020)-difference set in Z_10225

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006256 | ljcr-ds-10255-5127-2563-10255

(10255, 5127, 2563)-difference set in Z_10255

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006257 | ljcr-ds-10261-1216-144-10261

(10261, 1216, 144)-difference set in Z_10261

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006258 | ljcr-ds-10263-2199-471-10263

(10263, 2199, 471)-difference set in Z_10263

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006259 | ljcr-ds-1029-257-64-7-147

(1029, 257, 64)-difference set in Z_7 x Z_147

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006260 | ljcr-ds-1029-257-64-7-7-21

(1029, 257, 64)-difference set in Z_7 x Z_7 x Z_21

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006261 | ljcr-ds-10296-1421-196-10296

(10296, 1421, 196)-difference set in Z_10296

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006262 | ljcr-ds-10297-936-85-10297

(10297, 936, 85)-difference set in Z_10297

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006263 | ljcr-ds-10305-4417-1893-10305

(10305, 4417, 1893)-difference set in Z_10305

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006264 | ljcr-ds-10339-1723-287-10339

(10339, 1723, 287)-difference set in Z_10339

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006265 | ljcr-ds-10351-5175-2587-10351

(10351, 5175, 2587)-difference set in Z_10351

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006266 | ljcr-ds-10366-3456-1152-10366

(10366, 3456, 1152)-difference set in Z_10366

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006267 | ljcr-ds-10385-4544-1988-10385

(10385, 4544, 1988)-difference set in Z_10385

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006268 | ljcr-ds-10431-2086-417-10431

(10431, 2086, 417)-difference set in Z_10431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006269 | ljcr-ds-10439-4913-2312-10439

(10439, 4913, 2312)-difference set in Z_10439

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006270 | ljcr-ds-10447-5223-2611-10447

(10447, 5223, 2611)-difference set in Z_10447

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006271 | ljcr-ds-10465-2944-828-10465

(10465, 2944, 828)-difference set in Z_10465

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006272 | ljcr-ds-10497-1025-100-10497

(10497, 1025, 100)-difference set in Z_10497

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006273 | ljcr-ds-10499-4525-1950-10499

(10499, 4525, 1950)-difference set in Z_10499

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006274 | ljcr-ds-10507-3400-1100-10507

(10507, 3400, 1100)-difference set in Z_10507

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006275 | ljcr-ds-10533-2633-658-10533

(10533, 2633, 658)-difference set in Z_10533

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006276 | ljcr-ds-10539-958-87-10539

(10539, 958, 87)-difference set in Z_10539

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006277 | ljcr-ds-10539-958-87-3-3513

(10539, 958, 87)-difference set in Z_3 x Z_3513

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006278 | ljcr-ds-10543-5271-2635-10543

(10543, 5271, 2635)-difference set in Z_10543

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006279 | ljcr-ds-10550-1233-144-10550

(10550, 1233, 144)-difference set in Z_10550

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006280 | ljcr-ds-10579-5289-2644-10579

(10579, 5289, 2644)-difference set in Z_10579

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006281 | ljcr-ds-10587-5293-2646-10587

(10587, 5293, 2646)-difference set in Z_10587

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006282 | ljcr-ds-10609-273-7-103-103

(10609, 273, 7)-difference set in Z_103 x Z_103

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006283 | ljcr-ds-10620-4551-1950-10620

(10620, 4551, 1950)-difference set in Z_10620

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006284 | ljcr-ds-10711-3060-874-10711

(10711, 3060, 874)-difference set in Z_10711

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006285 | ljcr-ds-10719-3496-1140-10719

(10719, 3496, 1140)-difference set in Z_10719

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006286 | ljcr-ds-10725-2681-670-10725

(10725, 2681, 670)-difference set in Z_10725

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006287 | ljcr-ds-10764-1833-312-10764

(10764, 1833, 312)-difference set in Z_10764

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006288 | ljcr-ds-10769-673-42-10769

(10769, 673, 42)-difference set in Z_10769

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006289 | ljcr-ds-10769-673-42-11-979

(10769, 673, 42)-difference set in Z_11 x Z_979

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006290 | ljcr-ds-10791-1080-108-10791

(10791, 1080, 108)-difference set in Z_10791

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006291 | ljcr-ds-10795-4369-1768-10795

(10795, 4369, 1768)-difference set in Z_10795

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006292 | ljcr-ds-10815-5407-2703-10815

(10815, 5407, 2703)-difference set in Z_10815

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006293 | ljcr-ds-10821-2165-433-10821

(10821, 2165, 433)-difference set in Z_10821

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006294 | ljcr-ds-1083-541-270-19-57

(1083, 541, 270)-difference set in Z_19 x Z_57

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006295 | ljcr-ds-10837-1548-221-10837

(10837, 1548, 221)-difference set in Z_10837

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006296 | ljcr-ds-10880-4301-1700-10880

(10880, 4301, 1700)-difference set in Z_10880

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006297 | ljcr-ds-10880-990-90-2-5440

(10880, 990, 90)-difference set in Z_2 x Z_5440

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006298 | ljcr-ds-10880-990-90-4-2720

(10880, 990, 90)-difference set in Z_4 x Z_2720

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006299 | ljcr-ds-10880-990-90-8-1360

(10880, 990, 90)-difference set in Z_8 x Z_1360

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006300 | ljcr-ds-10899-5449-2724-10899

(10899, 5449, 2724)-difference set in Z_10899

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006301 | ljcr-ds-10901-3925-1413-10901

(10901, 3925, 1413)-difference set in Z_10901

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006302 | ljcr-ds-10923-5461-2730-10923

(10923, 5461, 2730)-difference set in Z_10923

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006303 | ljcr-ds-1093-169-26-1093

(1093, 169, 26)-difference set in Z_1093

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006304 | ljcr-ds-10989-2748-687-10989

(10989, 2748, 687)-difference set in Z_10989

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006305 | ljcr-ds-1100-315-90-10-110

(1100, 315, 90)-difference set in Z_10 x Z_110

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006306 | ljcr-ds-1100-315-90-1100

(1100, 315, 90)-difference set in Z_1100

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006307 | ljcr-ds-1100-315-90-5-220

(1100, 315, 90)-difference set in Z_5 x Z_220

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006308 | ljcr-ds-11005-525-25-11005

(11005, 525, 25)-difference set in Z_11005

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006309 | ljcr-ds-11011-3670-1223-11011

(11011, 3670, 1223)-difference set in Z_11011

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006310 | ljcr-ds-11023-3675-1225-11023

(11023, 3675, 1225)-difference set in Z_11023

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006311 | ljcr-ds-11045-4016-1460-11045

(11045, 4016, 1460)-difference set in Z_11045

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006312 | ljcr-ds-11055-5527-2763-11055

(11055, 5527, 2763)-difference set in Z_11055

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006313 | ljcr-ds-11081-2216-443-11081

(11081, 2216, 443)-difference set in Z_11081

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006314 | ljcr-ds-1111-111-11-1111

(1111, 111, 11)-difference set in Z_1111

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006315 | ljcr-ds-1111-186-31-1111

(1111, 186, 31)-difference set in Z_1111

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006316 | ljcr-ds-11141-2785-696-11141

(11141, 2785, 696)-difference set in Z_11141

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006317 | ljcr-ds-11175-1813-294-11175

(11175, 1813, 294)-difference set in Z_11175

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006318 | ljcr-ds-11181-5161-2382-11181

(11181, 5161, 2382)-difference set in Z_11181

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006319 | ljcr-ds-11195-5597-2798-11195

(11195, 5597, 2798)-difference set in Z_11195

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006320 | ljcr-ds-11235-5617-2808-11235

(11235, 5617, 2808)-difference set in Z_11235

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006321 | ljcr-ds-1125-281-70-15-75

(1125, 281, 70)-difference set in Z_15 x Z_75

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006322 | ljcr-ds-1125-281-70-5-15-15

(1125, 281, 70)-difference set in Z_5 x Z_15 x Z_15

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006323 | ljcr-ds-1125-281-70-5-5-45

(1125, 281, 70)-difference set in Z_5 x Z_5 x Z_45

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006324 | ljcr-ds-112575-1463-19-112575

(112575, 1463, 19)-difference set in Z_112575

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006325 | ljcr-ds-11263-5631-2815-11263

(11263, 5631, 2815)-difference set in Z_11263

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006326 | ljcr-ds-11286-1221-132-11286

(11286, 1221, 132)-difference set in Z_11286

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006327 | ljcr-ds-11325-3725-1225-11325

(11325, 3725, 1225)-difference set in Z_11325

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006328 | ljcr-ds-11340-783-54-18-630

(11340, 783, 54)-difference set in Z_18 x Z_630

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006329 | ljcr-ds-11340-783-54-3-3-1260

(11340, 783, 54)-difference set in Z_3 x Z_3 x Z_1260

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006330 | ljcr-ds-11340-783-54-3-6-630

(11340, 783, 54)-difference set in Z_3 x Z_6 x Z_630

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006331 | ljcr-ds-11340-783-54-9-1260

(11340, 783, 54)-difference set in Z_9 x Z_1260

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006332 | ljcr-ds-11377-4977-2177-11377

(11377, 4977, 2177)-difference set in Z_11377

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006333 | ljcr-ds-11439-1806-285-11439

(11439, 1806, 285)-difference set in Z_11439

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006334 | ljcr-ds-11440-1395-170-11440

(11440, 1395, 170)-difference set in Z_11440

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006335 | ljcr-ds-11461-3820-1273-11461

(11461, 3820, 1273)-difference set in Z_11461

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006336 | ljcr-ds-11479-3826-1275-11479

(11479, 3826, 1275)-difference set in Z_11479

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006337 | ljcr-ds-11525-2881-720-11525

(11525, 2881, 720)-difference set in Z_11525

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006338 | ljcr-ds-11534-608-32-11534

(11534, 608, 32)-difference set in Z_11534

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006339 | ljcr-ds-11543-1393-168-11543

(11543, 1393, 168)-difference set in Z_11543

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006340 | ljcr-ds-1156-561-272-17-68

(1156, 561, 272)-difference set in Z_17 x Z_68

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006341 | ljcr-ds-1156-561-272-34-34

(1156, 561, 272)-difference set in Z_34 x Z_34

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006342 | ljcr-ds-11571-1691-247-11571

(11571, 1691, 247)-difference set in Z_11571

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006343 | ljcr-ds-1161-465-186-3-3-129

(1161, 465, 186)-difference set in Z_3 x Z_3 x Z_129

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006344 | ljcr-ds-1161-465-186-3-387

(1161, 465, 186)-difference set in Z_3 x Z_387

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006345 | ljcr-ds-11628-3927-1326-11628

(11628, 3927, 1326)-difference set in Z_11628

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006346 | ljcr-ds-11704-3901-1300-11704

(11704, 3901, 1300)-difference set in Z_11704

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006347 | ljcr-ds-11749-1068-97-11749

(11749, 1068, 97)-difference set in Z_11749

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006348 | ljcr-ds-11753-2600-575-11753

(11753, 2600, 575)-difference set in Z_11753

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006349 | ljcr-ds-11811-1181-118-11811

(11811, 1181, 118)-difference set in Z_11811

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006350 | ljcr-ds-11811-5905-2952-11811

(11811, 5905, 2952)-difference set in Z_11811

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006351 | ljcr-ds-1183-394-131-13-91

(1183, 394, 131)-difference set in Z_13 x Z_91

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006352 | ljcr-ds-11831-2366-473-11831

(11831, 2366, 473)-difference set in Z_11831

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006353 | ljcr-ds-1189-297-74-1189

(1189, 297, 74)-difference set in Z_1189

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006354 | ljcr-ds-1191-511-219-1191

(1191, 511, 219)-difference set in Z_1191

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006355 | ljcr-ds-11911-1191-119-11911

(11911, 1191, 119)-difference set in Z_11911

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006356 | ljcr-ds-11915-5957-2978-11915

(11915, 5957, 2978)-difference set in Z_11915

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006357 | ljcr-ds-11921-2385-477-11921

(11921, 2385, 477)-difference set in Z_11921

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006358 | ljcr-ds-11935-2431-495-11935

(11935, 2431, 495)-difference set in Z_11935

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006359 | ljcr-ds-11951-2151-387-11951

(11951, 2151, 387)-difference set in Z_11951

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006360 | ljcr-ds-11951-3825-1224-11951

(11951, 3825, 1224)-difference set in Z_11951

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006361 | ljcr-ds-1197-208-36-1197

(1197, 208, 36)-difference set in Z_1197

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006362 | ljcr-ds-1197-208-36-3-399

(1197, 208, 36)-difference set in Z_3 x Z_399

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006363 | ljcr-ds-1197-300-75-3-399

(1197, 300, 75)-difference set in Z_3 x Z_399

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006364 | ljcr-ds-1200-110-10-10-120

(1200, 110, 10)-difference set in Z_10 x Z_120

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006365 | ljcr-ds-1200-110-10-2-10-60

(1200, 110, 10)-difference set in Z_2 x Z_10 x Z_60

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006366 | ljcr-ds-1200-110-10-2-2-300

(1200, 110, 10)-difference set in Z_2 x Z_2 x Z_300

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006367 | ljcr-ds-1200-110-10-20-60

(1200, 110, 10)-difference set in Z_20 x Z_60

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006368 | ljcr-ds-1200-110-10-4-300

(1200, 110, 10)-difference set in Z_4 x Z_300

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006369 | ljcr-ds-12009-1976-325-12009

(12009, 1976, 325)-difference set in Z_12009

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006370 | ljcr-ds-12052-2575-550-12052

(12052, 2575, 550)-difference set in Z_12052

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006371 | ljcr-ds-12059-6029-3014-12059

(12059, 6029, 3014)-difference set in Z_12059

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006372 | ljcr-ds-12061-4356-1573-12061

(12061, 4356, 1573)-difference set in Z_12061

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006373 | ljcr-ds-12099-1841-280-12099

(12099, 1841, 280)-difference set in Z_12099

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006374 | ljcr-ds-12099-6049-3024-12099

(12099, 6049, 3024)-difference set in Z_12099

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006375 | ljcr-ds-12104-5929-2904-12104

(12104, 5929, 2904)-difference set in Z_12104

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006376 | ljcr-ds-12122-5797-2772-12122

(12122, 5797, 2772)-difference set in Z_12122

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006377 | ljcr-ds-12127-2538-531-12127

(12127, 2538, 531)-difference set in Z_12127

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006378 | ljcr-ds-12131-6065-3032-12131

(12131, 6065, 3032)-difference set in Z_12131

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006379 | ljcr-ds-12141-1821-273-12141

(12141, 1821, 273)-difference set in Z_12141

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006380 | ljcr-ds-12147-6073-3036-12147

(12147, 6073, 3036)-difference set in Z_12147

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006381 | ljcr-ds-12151-4375-1575-12151

(12151, 4375, 1575)-difference set in Z_12151

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006382 | ljcr-ds-12155-5665-2640-12155

(12155, 5665, 2640)-difference set in Z_12155

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006383 | ljcr-ds-12167-869-62-23-23-23

(12167, 869, 62)-difference set in Z_23 x Z_23 x Z_23

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006384 | ljcr-ds-12181-4060-1353-12181

(12181, 4060, 1353)-difference set in Z_12181

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006385 | ljcr-ds-12208-3757-1156-12208

(12208, 3757, 1156)-difference set in Z_12208

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006386 | ljcr-ds-1221-245-49-1221

(1221, 245, 49)-difference set in Z_1221

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006387 | ljcr-ds-1221-305-76-1221

(1221, 305, 76)-difference set in Z_1221

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006388 | ljcr-ds-1222-297-72-1222

(1222, 297, 72)-difference set in Z_1222

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006389 | ljcr-ds-12221-5265-2268-12221

(12221, 5265, 2268)-difference set in Z_12221

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006390 | ljcr-ds-1225-136-15-35-35

(1225, 136, 15)-difference set in Z_35 x Z_35

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006391 | ljcr-ds-1225-136-15-5-245

(1225, 136, 15)-difference set in Z_5 x Z_245

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006392 | ljcr-ds-1225-136-15-7-175

(1225, 136, 15)-difference set in Z_7 x Z_175

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006393 | ljcr-ds-12265-5257-2253-12265

(12265, 5257, 2253)-difference set in Z_12265

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006394 | ljcr-ds-12275-6137-3068-12275

(12275, 6137, 3068)-difference set in Z_12275

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006395 | ljcr-ds-12285-3072-768-12285

(12285, 3072, 768)-difference set in Z_12285

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006396 | ljcr-ds-12285-333-9-12285

(12285, 333, 9)-difference set in Z_12285

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006397 | ljcr-ds-12285-333-9-3-3-1365

(12285, 333, 9)-difference set in Z_3 x Z_3 x Z_1365

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006398 | ljcr-ds-12285-333-9-3-4095

(12285, 333, 9)-difference set in Z_3 x Z_4095

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006399 | ljcr-ds-12403-4824-1876-12403

(12403, 4824, 1876)-difference set in Z_12403

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006400 | ljcr-ds-12429-3108-777-12429

(12429, 3108, 777)-difference set in Z_12429

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006401 | ljcr-ds-12445-5185-2160-12445

(12445, 5185, 2160)-difference set in Z_12445

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006402 | ljcr-ds-12465-2337-438-12465

(12465, 2337, 438)-difference set in Z_12465

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006403 | ljcr-ds-12481-2496-499-12481

(12481, 2496, 499)-difference set in Z_12481

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006404 | ljcr-ds-12481-4096-1344-12481

(12481, 4096, 1344)-difference set in Z_12481

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006405 | ljcr-ds-12483-6241-3120-12483

(12483, 6241, 3120)-difference set in Z_12483

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006406 | ljcr-ds-1251-625-312-3-417

(1251, 625, 312)-difference set in Z_3 x Z_417

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006407 | ljcr-ds-12529-5104-2079-12529

(12529, 5104, 2079)-difference set in Z_12529

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006408 | ljcr-ds-12537-1568-196-12537

(12537, 1568, 196)-difference set in Z_12537

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006409 | ljcr-ds-12541-1881-282-12541

(12541, 1881, 282)-difference set in Z_12541

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006410 | ljcr-ds-12559-4186-1395-12559

(12559, 4186, 1395)-difference set in Z_12559

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006411 | ljcr-ds-12595-4198-1399-12595

(12595, 4198, 1399)-difference set in Z_12595

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006412 | ljcr-ds-12607-6303-3151-12607

(12607, 6303, 3151)-difference set in Z_12607

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006413 | ljcr-ds-12649-2449-474-12649

(12649, 2449, 474)-difference set in Z_12649

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006414 | ljcr-ds-12651-6325-3162-12651

(12651, 6325, 3162)-difference set in Z_12651

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006415 | ljcr-ds-12656-2531-506-12656

(12656, 2531, 506)-difference set in Z_12656

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006416 | ljcr-ds-12711-2666-559-12711

(12711, 2666, 559)-difference set in Z_12711

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006417 | ljcr-ds-12741-3185-796-12741

(12741, 3185, 796)-difference set in Z_12741

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006418 | ljcr-ds-12781-2556-511-12781

(12781, 2556, 511)-difference set in Z_12781

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006419 | ljcr-ds-12805-4753-1764-12805

(12805, 4753, 1764)-difference set in Z_12805

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006420 | ljcr-ds-12851-1800-252-12851

(12851, 1800, 252)-difference set in Z_12851

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006421 | ljcr-ds-1288-352-96-1288

(1288, 352, 96)-difference set in Z_1288

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006422 | ljcr-ds-1288-352-96-2-2-322

(1288, 352, 96)-difference set in Z_2 x Z_2 x Z_322

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006423 | ljcr-ds-1288-352-96-2-644

(1288, 352, 96)-difference set in Z_2 x Z_644

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006424 | ljcr-ds-12881-2576-515-12881

(12881, 2576, 515)-difference set in Z_12881

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006425 | ljcr-ds-12883-4408-1508-12883

(12883, 4408, 1508)-difference set in Z_12883

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006426 | ljcr-ds-1289-553-237-1289

(1289, 553, 237)-difference set in Z_1289

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006427 | ljcr-ds-12901-5376-2240-12901

(12901, 5376, 2240)-difference set in Z_12901

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006428 | ljcr-ds-12905-4840-1815-12905

(12905, 4840, 1815)-difference set in Z_12905

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006429 | ljcr-ds-12915-1761-240-12915

(12915, 1761, 240)-difference set in Z_12915

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006430 | ljcr-ds-1293-324-81-1293

(1293, 324, 81)-difference set in Z_1293

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006431 | ljcr-ds-12959-6138-2907-12959

(12959, 6138, 2907)-difference set in Z_12959

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006432 | ljcr-ds-1296-630-306-18-72

(1296, 630, 306)-difference set in Z_18 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006433 | ljcr-ds-1296-630-306-2-18-36

(1296, 630, 306)-difference set in Z_2 x Z_18 x Z_36

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006434 | ljcr-ds-1296-630-306-2-2-18-18

(1296, 630, 306)-difference set in Z_2 x Z_2 x Z_18 x Z_18

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006435 | ljcr-ds-1296-630-306-2-6-6-18

(1296, 630, 306)-difference set in Z_2 x Z_6 x Z_6 x Z_18

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006436 | ljcr-ds-1296-630-306-3-12-36

(1296, 630, 306)-difference set in Z_3 x Z_12 x Z_36

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006437 | ljcr-ds-1296-630-306-3-3-12-12

(1296, 630, 306)-difference set in Z_3 x Z_3 x Z_12 x Z_12

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006438 | ljcr-ds-1296-630-306-3-3-6-24

(1296, 630, 306)-difference set in Z_3 x Z_3 x Z_6 x Z_24

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006439 | ljcr-ds-1296-630-306-3-6-6-12

(1296, 630, 306)-difference set in Z_3 x Z_6 x Z_6 x Z_12

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006440 | ljcr-ds-1296-630-306-3-6-72

(1296, 630, 306)-difference set in Z_3 x Z_6 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006441 | ljcr-ds-1296-630-306-36-36

(1296, 630, 306)-difference set in Z_36 x Z_36

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006442 | ljcr-ds-1296-630-306-6-216

(1296, 630, 306)-difference set in Z_6 x Z_216

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006443 | ljcr-ds-1296-630-306-6-6-36

(1296, 630, 306)-difference set in Z_6 x Z_6 x Z_36

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006444 | ljcr-ds-1296-630-306-6-6-6-6

(1296, 630, 306)-difference set in Z_6 x Z_6 x Z_6 x Z_6

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006445 | ljcr-ds-12989-4012-1239-12989

(12989, 4012, 1239)-difference set in Z_12989

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006446 | ljcr-ds-12995-6497-3248-12995

(12995, 6497, 3248)-difference set in Z_12995

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006447 | ljcr-ds-12997-3249-812-12997

(12997, 3249, 812)-difference set in Z_12997

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006448 | ljcr-ds-13039-5248-2112-13039

(13039, 5248, 2112)-difference set in Z_13039

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006449 | ljcr-ds-13071-6535-3267-13071

(13071, 6535, 3267)-difference set in Z_13071

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006450 | ljcr-ds-13077-1401-150-13077

(13077, 1401, 150)-difference set in Z_13077

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006451 | ljcr-ds-13091-715-39-13091

(13091, 715, 39)-difference set in Z_13091

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006452 | ljcr-ds-1313-288-63-1313

(1313, 288, 63)-difference set in Z_1313

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006453 | ljcr-ds-13131-6565-3282-13131

(13131, 6565, 3282)-difference set in Z_13131

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006454 | ljcr-ds-13156-1755-234-13156

(13156, 1755, 234)-difference set in Z_13156

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006455 | ljcr-ds-13167-6583-3291-13167

(13167, 6583, 3291)-difference set in Z_13167

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006456 | ljcr-ds-13171-4390-1463-13171

(13171, 4390, 1463)-difference set in Z_13171

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006457 | ljcr-ds-13175-4705-1680-13175

(13175, 4705, 1680)-difference set in Z_13175

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006458 | ljcr-ds-13189-4396-1465-13189

(13189, 4396, 1465)-difference set in Z_13189

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006459 | ljcr-ds-13201-4225-1352-13201

(13201, 4225, 1352)-difference set in Z_13201

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006460 | ljcr-ds-13203-6601-3300-13203

(13203, 6601, 3300)-difference set in Z_13203

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006461 | ljcr-ds-13211-6605-3302-13211

(13211, 6605, 3302)-difference set in Z_13211

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006462 | ljcr-ds-13221-2645-529-13221

(13221, 2645, 529)-difference set in Z_13221

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006463 | ljcr-ds-13227-6613-3306-13227

(13227, 6613, 3306)-difference set in Z_13227

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006464 | ljcr-ds-1323-661-330-21-63

(1323, 661, 330)-difference set in Z_21 x Z_63

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006465 | ljcr-ds-1323-661-330-3-21-21

(1323, 661, 330)-difference set in Z_3 x Z_21 x Z_21

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006466 | ljcr-ds-1323-661-330-3-3-147

(1323, 661, 330)-difference set in Z_3 x Z_3 x Z_147

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006467 | ljcr-ds-1323-661-330-7-189

(1323, 661, 330)-difference set in Z_7 x Z_189

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006468 | ljcr-ds-13244-4675-1650-13244

(13244, 4675, 1650)-difference set in Z_13244

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006469 | ljcr-ds-13251-6625-3312-13251

(13251, 6625, 3312)-difference set in Z_13251

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006470 | ljcr-ds-13263-6631-3315-13263

(13263, 6631, 3315)-difference set in Z_13263

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006471 | ljcr-ds-13266-5685-2436-13266

(13266, 5685, 2436)-difference set in Z_13266

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006472 | ljcr-ds-13300-1365-140-13300

(13300, 1365, 140)-difference set in Z_13300

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006473 | ljcr-ds-1331-190-27-11-121

(1331, 190, 27)-difference set in Z_11 x Z_121

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006474 | ljcr-ds-1331-210-33-11-121

(1331, 210, 33)-difference set in Z_11 x Z_121

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006475 | ljcr-ds-1331-266-53-11-121

(1331, 266, 53)-difference set in Z_11 x Z_121

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006476 | ljcr-ds-1331-665-332-11-121

(1331, 665, 332)-difference set in Z_11 x Z_121

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006477 | ljcr-ds-13333-3333-833-13333

(13333, 3333, 833)-difference set in Z_13333

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006478 | ljcr-ds-13371-6685-3342-13371

(13371, 6685, 3342)-difference set in Z_13371

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006479 | ljcr-ds-13420-2485-460-13420

(13420, 2485, 460)-difference set in Z_13420

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006480 | ljcr-ds-13427-6713-3356-13427

(13427, 6713, 3356)-difference set in Z_13427

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006481 | ljcr-ds-13433-4600-1575-13433

(13433, 4600, 1575)-difference set in Z_13433

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006482 | ljcr-ds-1344-238-42-2-2-336

(1344, 238, 42)-difference set in Z_2 x Z_2 x Z_336

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006483 | ljcr-ds-1344-238-42-2-672

(1344, 238, 42)-difference set in Z_2 x Z_672

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006484 | ljcr-ds-1344-238-42-4-336

(1344, 238, 42)-difference set in Z_4 x Z_336

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006485 | ljcr-ds-13455-6727-3363-13455

(13455, 6727, 3363)-difference set in Z_13455

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006486 | ljcr-ds-13471-6286-2933-13471

(13471, 6286, 2933)-difference set in Z_13471

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006487 | ljcr-ds-13490-5453-2204-13490

(13490, 5453, 2204)-difference set in Z_13490

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006488 | ljcr-ds-1350-285-60-3-15-30

(1350, 285, 60)-difference set in Z_3 x Z_15 x Z_30

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006489 | ljcr-ds-13503-6751-3375-13503

(13503, 6751, 3375)-difference set in Z_13503

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006490 | ljcr-ds-1353-169-21-1353

(1353, 169, 21)-difference set in Z_1353

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006491 | ljcr-ds-13530-4565-1540-13530

(13530, 4565, 1540)-difference set in Z_13530

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006492 | ljcr-ds-135387-974-7-135387

(135387, 974, 7)-difference set in Z_135387

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006493 | ljcr-ds-135387-974-7-21-6447

(135387, 974, 7)-difference set in Z_21 x Z_6447

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006494 | ljcr-ds-135387-974-7-3-45129

(135387, 974, 7)-difference set in Z_3 x Z_45129

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006495 | ljcr-ds-135387-974-7-7-19341

(135387, 974, 7)-difference set in Z_7 x Z_19341

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006496 | ljcr-ds-13587-6793-3396-13587

(13587, 6793, 3396)-difference set in Z_13587

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006497 | ljcr-ds-13612-4537-1512-13612

(13612, 4537, 1512)-difference set in Z_13612

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006498 | ljcr-ds-13637-3409-852-13637

(13637, 3409, 852)-difference set in Z_13637

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006499 | ljcr-ds-13640-5337-2088-13640

(13640, 5337, 2088)-difference set in Z_13640

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006500 | ljcr-ds-13643-6821-3410-13643

(13643, 6821, 3410)-difference set in Z_13643

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006501 | ljcr-ds-13647-6823-3411-13647

(13647, 6823, 3411)-difference set in Z_13647

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006502 | ljcr-ds-13663-5589-2286-13663

(13663, 5589, 2286)-difference set in Z_13663

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006503 | ljcr-ds-13695-4510-1485-13695

(13695, 4510, 1485)-difference set in Z_13695

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006504 | ljcr-ds-13699-6849-3424-13699

(13699, 6849, 3424)-difference set in Z_13699

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006505 | ljcr-ds-13717-5589-2277-13717

(13717, 5589, 2277)-difference set in Z_13717

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006506 | ljcr-ds-13735-5887-2523-13735

(13735, 5887, 2523)-difference set in Z_13735

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006507 | ljcr-ds-1375-229-38-5-275

(1375, 229, 38)-difference set in Z_5 x Z_275

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006508 | ljcr-ds-1375-229-38-5-5-55

(1375, 229, 38)-difference set in Z_5 x Z_5 x Z_55

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006509 | ljcr-ds-13805-2465-440-13805

(13805, 2465, 440)-difference set in Z_13805

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006510 | ljcr-ds-13823-6911-3455-13823

(13823, 6911, 3455)-difference set in Z_13823

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006511 | ljcr-ds-13869-3468-867-13869

(13869, 3468, 867)-difference set in Z_13869

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006512 | ljcr-ds-1387-154-17-1387

(1387, 154, 17)-difference set in Z_1387

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006513 | ljcr-ds-13887-6943-3471-13887

(13887, 6943, 3471)-difference set in Z_13887

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006514 | ljcr-ds-13893-2416-420-13893

(13893, 2416, 420)-difference set in Z_13893

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006515 | ljcr-ds-13923-6961-3480-13923

(13923, 6961, 3480)-difference set in Z_13923

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006516 | ljcr-ds-13993-3817-1041-13993

(13993, 3817, 1041)-difference set in Z_13993

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006517 | ljcr-ds-14003-7001-3500-14003

(14003, 7001, 3500)-difference set in Z_14003

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006518 | ljcr-ds-14008-3451-850-14008

(14008, 3451, 850)-difference set in Z_14008

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006519 | ljcr-ds-14043-7021-3510-14043

(14043, 7021, 3510)-difference set in Z_14043

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006520 | ljcr-ds-1408-336-80-2-2-2-2-2-44

(1408, 336, 80)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_44

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006521 | ljcr-ds-1408-336-80-2-2-2-2-88

(1408, 336, 80)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_88

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006522 | ljcr-ds-1408-336-80-2-2-2-4-44

(1408, 336, 80)-difference set in Z_2 x Z_2 x Z_2 x Z_4 x Z_44

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006523 | ljcr-ds-1408-336-80-2-2-4-88

(1408, 336, 80)-difference set in Z_2 x Z_2 x Z_4 x Z_88

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006524 | ljcr-ds-1408-336-80-2-4-4-44

(1408, 336, 80)-difference set in Z_2 x Z_4 x Z_4 x Z_44

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006525 | ljcr-ds-1408-336-80-2-8-88

(1408, 336, 80)-difference set in Z_2 x Z_8 x Z_88

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006526 | ljcr-ds-1408-336-80-4-4-88

(1408, 336, 80)-difference set in Z_4 x Z_4 x Z_88

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006527 | ljcr-ds-14163-7081-3540-14163

(14163, 7081, 3540)-difference set in Z_14163

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006528 | ljcr-ds-14181-2836-567-14181

(14181, 2836, 567)-difference set in Z_14181

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006529 | ljcr-ds-14191-7095-3547-14191

(14191, 7095, 3547)-difference set in Z_14191

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006530 | ljcr-ds-14193-6209-2716-14193

(14193, 6209, 2716)-difference set in Z_14193

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006531 | ljcr-ds-14231-2846-569-14231

(14231, 2846, 569)-difference set in Z_14231

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006532 | ljcr-ds-14245-1188-99-14245

(14245, 1188, 99)-difference set in Z_14245

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006533 | ljcr-ds-14291-7145-3572-14291

(14291, 7145, 3572)-difference set in Z_14291

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006534 | ljcr-ds-14309-3577-894-14309

(14309, 3577, 894)-difference set in Z_14309

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006535 | ljcr-ds-1431-495-171-3-3-159

(1431, 495, 171)-difference set in Z_3 x Z_3 x Z_159

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006536 | ljcr-ds-14329-1792-224-14329

(14329, 1792, 224)-difference set in Z_14329

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006537 | ljcr-ds-14341-4780-1593-14341

(14341, 4780, 1593)-difference set in Z_14341

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006538 | ljcr-ds-14371-2395-399-14371

(14371, 2395, 399)-difference set in Z_14371

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006539 | ljcr-ds-14399-7199-3599-14399

(14399, 7199, 3599)-difference set in Z_14399

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006540 | ljcr-ds-14421-2576-460-14421

(14421, 2576, 460)-difference set in Z_14421

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006541 | ljcr-ds-14421-3605-901-14421

(14421, 3605, 901)-difference set in Z_14421

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006542 | ljcr-ds-14421-6181-2649-14421

(14421, 6181, 2649)-difference set in Z_14421

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006543 | ljcr-ds-1444-703-342-19-76

(1444, 703, 342)-difference set in Z_19 x Z_76

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006544 | ljcr-ds-1444-703-342-38-38

(1444, 703, 342)-difference set in Z_38 x Z_38

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006545 | ljcr-ds-14469-3617-904-14469

(14469, 3617, 904)-difference set in Z_14469

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006546 | ljcr-ds-14476-1351-126-14476

(14476, 1351, 126)-difference set in Z_14476

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006547 | ljcr-ds-14491-7245-3622-14491

(14491, 7245, 3622)-difference set in Z_14491

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006548 | ljcr-ds-14511-7255-3627-14511

(14511, 7255, 3627)-difference set in Z_14511

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006549 | ljcr-ds-14527-7263-3631-14527

(14527, 7263, 3631)-difference set in Z_14527

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006550 | ljcr-ds-14535-4902-1653-14535

(14535, 4902, 1653)-difference set in Z_14535

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006551 | ljcr-ds-14555-2299-363-14555

(14555, 2299, 363)-difference set in Z_14555

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006552 | ljcr-ds-14559-6526-2925-14559

(14559, 6526, 2925)-difference set in Z_14559

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006553 | ljcr-ds-1456-486-162-2-2-2-182

(1456, 486, 162)-difference set in Z_2 x Z_2 x Z_2 x Z_182

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006554 | ljcr-ds-1456-486-162-2-2-364

(1456, 486, 162)-difference set in Z_2 x Z_2 x Z_364

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006555 | ljcr-ds-1456-486-162-2-728

(1456, 486, 162)-difference set in Z_2 x Z_728

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006556 | ljcr-ds-1456-486-162-4-364

(1456, 486, 162)-difference set in Z_4 x Z_364

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006557 | ljcr-ds-14565-3641-910-14565

(14565, 3641, 910)-difference set in Z_14565

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006558 | ljcr-ds-14575-7287-3643-14575

(14575, 7287, 3643)-difference set in Z_14575

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006559 | ljcr-ds-14589-6252-2679-14589

(14589, 6252, 2679)-difference set in Z_14589

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006560 | ljcr-ds-14607-6432-2832-14607

(14607, 6432, 2832)-difference set in Z_14607

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006561 | ljcr-ds-14623-7311-3655-14623

(14623, 7311, 3655)-difference set in Z_14623

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006562 | ljcr-ds-14641-976-65-121-121

(14641, 976, 65)-difference set in Z_121 x Z_121

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006563 | ljcr-ds-14643-7321-3660-14643

(14643, 7321, 3660)-difference set in Z_14643

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006564 | ljcr-ds-1467-733-366-3-489

(1467, 733, 366)-difference set in Z_3 x Z_489

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006565 | ljcr-ds-14691-1470-147-14691

(14691, 1470, 147)-difference set in Z_14691

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006566 | ljcr-ds-14725-6544-2908-14725

(14725, 6544, 2908)-difference set in Z_14725

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006567 | ljcr-ds-1473-576-225-1473

(1473, 576, 225)-difference set in Z_1473

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006568 | ljcr-ds-1475-737-368-5-295

(1475, 737, 368)-difference set in Z_5 x Z_295

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006569 | ljcr-ds-14807-3366-765-14807

(14807, 3366, 765)-difference set in Z_14807

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006570 | ljcr-ds-14820-4235-1210-14820

(14820, 4235, 1210)-difference set in Z_14820

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006571 | ljcr-ds-1485-372-93-3-3-165

(1485, 372, 93)-difference set in Z_3 x Z_3 x Z_165

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006572 | ljcr-ds-1485-372-93-3-495

(1485, 372, 93)-difference set in Z_3 x Z_495

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006573 | ljcr-ds-1485-477-153-3-3-165

(1485, 477, 153)-difference set in Z_3 x Z_3 x Z_165

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006574 | ljcr-ds-1485-477-153-3-495

(1485, 477, 153)-difference set in Z_3 x Z_495

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006575 | ljcr-ds-14885-3721-930-14885

(14885, 3721, 930)-difference set in Z_14885

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006576 | ljcr-ds-14912-962-62-2-7456

(14912, 962, 62)-difference set in Z_2 x Z_7456

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006577 | ljcr-ds-14912-962-62-4-3728

(14912, 962, 62)-difference set in Z_4 x Z_3728

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006578 | ljcr-ds-14912-962-62-8-1864

(14912, 962, 62)-difference set in Z_8 x Z_1864

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006579 | ljcr-ds-14917-4521-1370-14917

(14917, 4521, 1370)-difference set in Z_14917

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006580 | ljcr-ds-14925-820-45-14925

(14925, 820, 45)-difference set in Z_14925

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006581 | ljcr-ds-14925-820-45-5-2985

(14925, 820, 45)-difference set in Z_5 x Z_2985

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006582 | ljcr-ds-14941-7056-3332-14941

(14941, 7056, 3332)-difference set in Z_14941

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006583 | ljcr-ds-14949-3737-934-14949

(14949, 3737, 934)-difference set in Z_14949

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006584 | ljcr-ds-14976-600-24-12-1248

(14976, 600, 24)-difference set in Z_12 x Z_1248

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006585 | ljcr-ds-14976-600-24-2-12-624

(14976, 600, 24)-difference set in Z_2 x Z_12 x Z_624

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006586 | ljcr-ds-14976-600-24-2-2-12-312

(14976, 600, 24)-difference set in Z_2 x Z_2 x Z_12 x Z_312

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006587 | ljcr-ds-14976-600-24-2-2-2-12-156

(14976, 600, 24)-difference set in Z_2 x Z_2 x Z_2 x Z_12 x Z_156

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006588 | ljcr-ds-14976-600-24-2-2-2-1872

(14976, 600, 24)-difference set in Z_2 x Z_2 x Z_2 x Z_1872

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006589 | ljcr-ds-14976-600-24-2-2-2-2-6-156

(14976, 600, 24)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_6 x Z_156

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006590 | ljcr-ds-14976-600-24-2-2-2-2-936

(14976, 600, 24)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_936

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006591 | ljcr-ds-14976-600-24-2-2-2-6-312

(14976, 600, 24)-difference set in Z_2 x Z_2 x Z_2 x Z_6 x Z_312

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006592 | ljcr-ds-14976-600-24-2-2-3744

(14976, 600, 24)-difference set in Z_2 x Z_2 x Z_3744

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006593 | ljcr-ds-14976-600-24-2-2-4-936

(14976, 600, 24)-difference set in Z_2 x Z_2 x Z_4 x Z_936

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006594 | ljcr-ds-14976-600-24-2-2-6-624

(14976, 600, 24)-difference set in Z_2 x Z_2 x Z_6 x Z_624

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006595 | ljcr-ds-14976-600-24-2-24-312

(14976, 600, 24)-difference set in Z_2 x Z_24 x Z_312

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006596 | ljcr-ds-14976-600-24-2-4-12-156

(14976, 600, 24)-difference set in Z_2 x Z_4 x Z_12 x Z_156

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006597 | ljcr-ds-14976-600-24-2-4-1872

(14976, 600, 24)-difference set in Z_2 x Z_4 x Z_1872

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006598 | ljcr-ds-14976-600-24-2-6-1248

(14976, 600, 24)-difference set in Z_2 x Z_6 x Z_1248

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006599 | ljcr-ds-14976-600-24-2-8-936

(14976, 600, 24)-difference set in Z_2 x Z_8 x Z_936

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006600 | ljcr-ds-14976-600-24-24-624

(14976, 600, 24)-difference set in Z_24 x Z_624

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006601 | ljcr-ds-14976-600-24-4-12-312

(14976, 600, 24)-difference set in Z_4 x Z_12 x Z_312

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006602 | ljcr-ds-14976-600-24-4-3744

(14976, 600, 24)-difference set in Z_4 x Z_3744

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006603 | ljcr-ds-14976-600-24-4-4-936

(14976, 600, 24)-difference set in Z_4 x Z_4 x Z_936

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006604 | ljcr-ds-14976-600-24-6-2496

(14976, 600, 24)-difference set in Z_6 x Z_2496

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006605 | ljcr-ds-14976-600-24-8-1872

(14976, 600, 24)-difference set in Z_8 x Z_1872

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006606 | ljcr-ds-14981-3745-936-14981

(14981, 3745, 936)-difference set in Z_14981

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006607 | ljcr-ds-14997-3749-937-14997

(14997, 3749, 937)-difference set in Z_14997

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006608 | ljcr-ds-15001-5001-1667-15001

(15001, 5001, 1667)-difference set in Z_15001

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006609 | ljcr-ds-15007-7381-3630-15007

(15007, 7381, 3630)-difference set in Z_15007

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006610 | ljcr-ds-15021-3756-939-15021

(15021, 3756, 939)-difference set in Z_15021

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006611 | ljcr-ds-15023-7511-3755-15023

(15023, 7511, 3755)-difference set in Z_15023

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006612 | ljcr-ds-15043-5451-1975-15043

(15043, 5451, 1975)-difference set in Z_15043

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006613 | ljcr-ds-15051-5075-1711-15051

(15051, 5075, 1711)-difference set in Z_15051

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006614 | ljcr-ds-15065-1345-120-15065

(15065, 1345, 120)-difference set in Z_15065

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006615 | ljcr-ds-15093-1716-195-15093

(15093, 1716, 195)-difference set in Z_15093

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006616 | ljcr-ds-15109-3777-944-15109

(15109, 3777, 944)-difference set in Z_15109

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006617 | ljcr-ds-15159-5512-2004-15159

(15159, 5512, 2004)-difference set in Z_15159

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006618 | ljcr-ds-15169-6400-2700-15169

(15169, 6400, 2700)-difference set in Z_15169

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006619 | ljcr-ds-1519-231-35-7-217

(1519, 231, 35)-difference set in Z_7 x Z_217

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006620 | ljcr-ds-1519-253-42-7-217

(1519, 253, 42)-difference set in Z_7 x Z_217

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006621 | ljcr-ds-1521-305-61-13-117

(1521, 305, 61)-difference set in Z_13 x Z_117

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006622 | ljcr-ds-1521-305-61-3-507

(1521, 305, 61)-difference set in Z_3 x Z_507

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006623 | ljcr-ds-1521-305-61-39-39

(1521, 305, 61)-difference set in Z_39 x Z_39

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006624 | ljcr-ds-15237-3809-952-15237

(15237, 3809, 952)-difference set in Z_15237

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006625 | ljcr-ds-15241-1905-238-15241

(15241, 1905, 238)-difference set in Z_15241

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006626 | ljcr-ds-15253-3937-1016-15253

(15253, 3937, 1016)-difference set in Z_15253

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006627 | ljcr-ds-15337-568-21-15337

(15337, 568, 21)-difference set in Z_15337

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006628 | ljcr-ds-15337-568-21-7-2191

(15337, 568, 21)-difference set in Z_7 x Z_2191

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006629 | ljcr-ds-15345-960-60-15345

(15345, 960, 60)-difference set in Z_15345

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006630 | ljcr-ds-15345-960-60-3-5115

(15345, 960, 60)-difference set in Z_3 x Z_5115

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006631 | ljcr-ds-15397-3849-962-15397

(15397, 3849, 962)-difference set in Z_15397

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006632 | ljcr-ds-15407-7703-3851-15407

(15407, 7703, 3851)-difference set in Z_15407

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006633 | ljcr-ds-15409-4816-1505-15409

(15409, 4816, 1505)-difference set in Z_15409

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006634 | ljcr-ds-15409-6849-3044-15409

(15409, 6849, 3044)-difference set in Z_15409

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006635 | ljcr-ds-15466-4125-1100-15466

(15466, 4125, 1100)-difference set in Z_15466

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006636 | ljcr-ds-15487-2581-430-15487

(15487, 2581, 430)-difference set in Z_15487

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006637 | ljcr-ds-15515-7757-3878-15515

(15515, 7757, 3878)-difference set in Z_15515

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006638 | ljcr-ds-15539-4114-1089-15539

(15539, 4114, 1089)-difference set in Z_15539

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006639 | ljcr-ds-15573-4352-1216-15573

(15573, 4352, 1216)-difference set in Z_15573

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006640 | ljcr-ds-15600-4105-1080-15600

(15600, 4105, 1080)-difference set in Z_15600

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006641 | ljcr-ds-1561-105-7-1561

(1561, 105, 7)-difference set in Z_1561

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006642 | ljcr-ds-15625-280-5-125-125

(15625, 280, 5)-difference set in Z_125 x Z_125

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006643 | ljcr-ds-15625-280-5-25-25-25

(15625, 280, 5)-difference set in Z_25 x Z_25 x Z_25

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006644 | ljcr-ds-15625-280-5-25-625

(15625, 280, 5)-difference set in Z_25 x Z_625

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006645 | ljcr-ds-15625-280-5-5-25-125

(15625, 280, 5)-difference set in Z_5 x Z_25 x Z_125

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006646 | ljcr-ds-15625-280-5-5-5-25-25

(15625, 280, 5)-difference set in Z_5 x Z_5 x Z_25 x Z_25

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006647 | ljcr-ds-15625-280-5-5-5-5-125

(15625, 280, 5)-difference set in Z_5 x Z_5 x Z_5 x Z_125

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006648 | ljcr-ds-15625-280-5-5-5-5-5-25

(15625, 280, 5)-difference set in Z_5 x Z_5 x Z_5 x Z_5 x Z_25

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006649 | ljcr-ds-15625-280-5-5-5-5-5-5-5

(15625, 280, 5)-difference set in Z_5 x Z_5 x Z_5 x Z_5 x Z_5 x Z_5

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006650 | ljcr-ds-15625-280-5-5-5-625

(15625, 280, 5)-difference set in Z_5 x Z_5 x Z_625

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006651 | ljcr-ds-15627-7813-3906-15627

(15627, 7813, 3906)-difference set in Z_15627

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006652 | ljcr-ds-15631-7815-3907-15631

(15631, 7815, 3907)-difference set in Z_15631

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006653 | ljcr-ds-15655-5218-1739-15655

(15655, 5218, 1739)-difference set in Z_15655

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006654 | ljcr-ds-15664-5221-1740-15664

(15664, 5221, 1740)-difference set in Z_15664

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006655 | ljcr-ds-15693-3924-981-15693

(15693, 3924, 981)-difference set in Z_15693

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006656 | ljcr-ds-15709-3213-657-15709

(15709, 3213, 657)-difference set in Z_15709

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006657 | ljcr-ds-15771-3155-631-15771

(15771, 3155, 631)-difference set in Z_15771

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006658 | ljcr-ds-15799-5266-1755-15799

(15799, 5266, 1755)-difference set in Z_15799

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006659 | ljcr-ds-15808-958-58-2-7904

(15808, 958, 58)-difference set in Z_2 x Z_7904

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006660 | ljcr-ds-15808-958-58-4-3952

(15808, 958, 58)-difference set in Z_4 x Z_3952

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006661 | ljcr-ds-15808-958-58-8-1976

(15808, 958, 58)-difference set in Z_8 x Z_1976

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006662 | ljcr-ds-15811-2635-439-15811

(15811, 2635, 439)-difference set in Z_15811

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006663 | ljcr-ds-15851-7925-3962-15851

(15851, 7925, 3962)-difference set in Z_15851

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006664 | ljcr-ds-15855-7927-3963-15855

(15855, 7927, 3963)-difference set in Z_15855

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006665 | ljcr-ds-15885-1805-205-15885

(15885, 1805, 205)-difference set in Z_15885

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006666 | ljcr-ds-15939-4291-1155-15939

(15939, 4291, 1155)-difference set in Z_15939

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006667 | ljcr-ds-15951-7975-3987-15951

(15951, 7975, 3987)-difference set in Z_15951

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006668 | ljcr-ds-15981-3196-639-15981

(15981, 3196, 639)-difference set in Z_15981

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006669 | ljcr-ds-1600-780-380-2-10-80

(1600, 780, 380)-difference set in Z_2 x Z_10 x Z_80

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006670 | ljcr-ds-1600-780-380-2-20-40

(1600, 780, 380)-difference set in Z_2 x Z_20 x Z_40

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006671 | ljcr-ds-1600-780-380-2-4-200

(1600, 780, 380)-difference set in Z_2 x Z_4 x Z_200

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006672 | ljcr-ds-1600-780-380-20-80

(1600, 780, 380)-difference set in Z_20 x Z_80

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006673 | ljcr-ds-1600-780-380-4-20-20

(1600, 780, 380)-difference set in Z_4 x Z_20 x Z_20

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006674 | ljcr-ds-1600-780-380-4-4-100

(1600, 780, 380)-difference set in Z_4 x Z_4 x Z_100

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006675 | ljcr-ds-1600-780-380-40-40

(1600, 780, 380)-difference set in Z_40 x Z_40

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006676 | ljcr-ds-1600-780-380-8-200

(1600, 780, 380)-difference set in Z_8 x Z_200

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006677 | ljcr-ds-16009-2553-407-16009

(16009, 2553, 407)-difference set in Z_16009

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006678 | ljcr-ds-1611-805-402-3-537

(1611, 805, 402)-difference set in Z_3 x Z_537

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006679 | ljcr-ds-16133-4033-1008-16133

(16133, 4033, 1008)-difference set in Z_16133

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006680 | ljcr-ds-16143-8071-4035-16143

(16143, 8071, 4035)-difference set in Z_16143

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006681 | ljcr-ds-1615-270-45-1615

(1615, 270, 45)-difference set in Z_1615

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006682 | ljcr-ds-16163-8081-4040-16163

(16163, 8081, 4040)-difference set in Z_16163

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006683 | ljcr-ds-16201-2025-253-16201

(16201, 2025, 253)-difference set in Z_16201

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006684 | ljcr-ds-16207-8103-4051-16207

(16207, 8103, 4051)-difference set in Z_16207

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006685 | ljcr-ds-16231-4870-1461-16231

(16231, 4870, 1461)-difference set in Z_16231

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006686 | ljcr-ds-16259-4434-1209-16259

(16259, 4434, 1209)-difference set in Z_16259

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006687 | ljcr-ds-1629-297-54-3-543

(1629, 297, 54)-difference set in Z_3 x Z_543

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006688 | ljcr-ds-1629-704-304-1629

(1629, 704, 304)-difference set in Z_1629

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006689 | ljcr-ds-1629-704-304-3-543

(1629, 704, 304)-difference set in Z_3 x Z_543

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006690 | ljcr-ds-16291-8145-4072-16291

(16291, 8145, 4072)-difference set in Z_16291

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006691 | ljcr-ds-16297-6888-2911-16297

(16297, 6888, 2911)-difference set in Z_16297

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006692 | ljcr-ds-16299-8149-4074-16299

(16299, 8149, 4074)-difference set in Z_16299

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006693 | ljcr-ds-1632-700-300-2-2-2-204

(1632, 700, 300)-difference set in Z_2 x Z_2 x Z_2 x Z_204

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006694 | ljcr-ds-1632-700-300-2-2-408

(1632, 700, 300)-difference set in Z_2 x Z_2 x Z_408

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006695 | ljcr-ds-1632-700-300-2-4-204

(1632, 700, 300)-difference set in Z_2 x Z_4 x Z_204

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006696 | ljcr-ds-1632-700-300-2-816

(1632, 700, 300)-difference set in Z_2 x Z_816

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006697 | ljcr-ds-1632-700-300-4-408

(1632, 700, 300)-difference set in Z_4 x Z_408

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006698 | ljcr-ds-16331-2485-378-16331

(16331, 2485, 378)-difference set in Z_16331

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006699 | ljcr-ds-163603-1342-11-163603

(163603, 1342, 11)-difference set in Z_163603

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006700 | ljcr-ds-16380-4467-1218-16380

(16380, 4467, 1218)-difference set in Z_16380

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006701 | ljcr-ds-16381-7281-3236-16381

(16381, 7281, 3236)-difference set in Z_16381

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006702 | ljcr-ds-16387-2731-455-16387

(16387, 2731, 455)-difference set in Z_16387

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006703 | ljcr-ds-16457-2057-257-16457

(16457, 2057, 257)-difference set in Z_16457

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006704 | ljcr-ds-16461-4116-1029-16461

(16461, 4116, 1029)-difference set in Z_16461

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006705 | ljcr-ds-16461-6585-2634-16461

(16461, 6585, 2634)-difference set in Z_16461

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006706 | ljcr-ds-1647-823-411-3-3-183

(1647, 823, 411)-difference set in Z_3 x Z_3 x Z_183

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006707 | ljcr-ds-1647-823-411-3-549

(1647, 823, 411)-difference set in Z_3 x Z_549

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006708 | ljcr-ds-16479-7063-3027-16479

(16479, 7063, 3027)-difference set in Z_16479

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006709 | ljcr-ds-16511-4446-1197-16511

(16511, 4446, 1197)-difference set in Z_16511

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006710 | ljcr-ds-16531-7221-3154-16531

(16531, 7221, 3154)-difference set in Z_16531

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006711 | ljcr-ds-16592-706-30-2-2-2-2074

(16592, 706, 30)-difference set in Z_2 x Z_2 x Z_2 x Z_2074

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006712 | ljcr-ds-16592-706-30-2-2-4148

(16592, 706, 30)-difference set in Z_2 x Z_2 x Z_4148

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006713 | ljcr-ds-16592-706-30-2-8296

(16592, 706, 30)-difference set in Z_2 x Z_8296

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006714 | ljcr-ds-16592-706-30-4-4148

(16592, 706, 30)-difference set in Z_4 x Z_4148

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006715 | ljcr-ds-16639-5782-2009-16639

(16639, 5782, 2009)-difference set in Z_16639

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006716 | ljcr-ds-16663-5554-1851-16663

(16663, 5554, 1851)-difference set in Z_16663

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006717 | ljcr-ds-16667-8333-4166-16667

(16667, 8333, 4166)-difference set in Z_16667

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006718 | ljcr-ds-16681-5560-1853-16681

(16681, 5560, 1853)-difference set in Z_16681

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006719 | ljcr-ds-16721-2641-417-16721

(16721, 2641, 417)-difference set in Z_16721

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006720 | ljcr-ds-16721-3136-588-16721

(16721, 3136, 588)-difference set in Z_16721

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006721 | ljcr-ds-16725-4181-1045-16725

(16725, 4181, 1045)-difference set in Z_16725

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006722 | ljcr-ds-16771-3225-620-16771

(16771, 3225, 620)-difference set in Z_16771

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006723 | ljcr-ds-16777-1864-207-16777

(16777, 1864, 207)-difference set in Z_16777

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006724 | ljcr-ds-16874-3949-924-16874

(16874, 3949, 924)-difference set in Z_16874

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006725 | ljcr-ds-16875-650-25-15-15-75

(16875, 650, 25)-difference set in Z_15 x Z_15 x Z_75

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006726 | ljcr-ds-16875-650-25-25-675

(16875, 650, 25)-difference set in Z_25 x Z_675

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006727 | ljcr-ds-16875-650-25-3-75-75

(16875, 650, 25)-difference set in Z_3 x Z_75 x Z_75

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006728 | ljcr-ds-16875-650-25-5-15-225

(16875, 650, 25)-difference set in Z_5 x Z_15 x Z_225

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006729 | ljcr-ds-16875-650-25-5-5-675

(16875, 650, 25)-difference set in Z_5 x Z_5 x Z_675

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006730 | ljcr-ds-16875-650-25-75-225

(16875, 650, 25)-difference set in Z_75 x Z_225

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006731 | ljcr-ds-16899-6461-2470-16899

(16899, 6461, 2470)-difference set in Z_16899

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006732 | ljcr-ds-16911-8455-4227-16911

(16911, 8455, 4227)-difference set in Z_16911

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006733 | ljcr-ds-16965-4241-1060-16965

(16965, 4241, 1060)-difference set in Z_16965

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006734 | ljcr-ds-17007-8503-4251-17007

(17007, 8503, 4251)-difference set in Z_17007

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006735 | ljcr-ds-17017-7800-3575-17017

(17017, 7800, 3575)-difference set in Z_17017

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006736 | ljcr-ds-17040-7745-3520-17040

(17040, 7745, 3520)-difference set in Z_17040

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006737 | ljcr-ds-1705-640-240-1705

(1705, 640, 240)-difference set in Z_1705

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006738 | ljcr-ds-17071-3415-683-17071

(17071, 3415, 683)-difference set in Z_17071

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006739 | ljcr-ds-17071-8535-4267-17071

(17071, 8535, 4267)-difference set in Z_17071

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006740 | ljcr-ds-17083-8541-4270-17083

(17083, 8541, 4270)-difference set in Z_17083

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006741 | ljcr-ds-17095-3367-663-17095

(17095, 3367, 663)-difference set in Z_17095

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006742 | ljcr-ds-17143-8571-4285-17143

(17143, 8571, 4285)-difference set in Z_17143

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006743 | ljcr-ds-17157-4289-1072-17157

(17157, 4289, 1072)-difference set in Z_17157

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006744 | ljcr-ds-17161-936-51-131-131

(17161, 936, 51)-difference set in Z_131 x Z_131

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006745 | ljcr-ds-17163-8581-4290-17163

(17163, 8581, 4290)-difference set in Z_17163

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006746 | ljcr-ds-17173-2916-495-17173

(17173, 2916, 495)-difference set in Z_17173

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006747 | ljcr-ds-17211-8605-4302-17211

(17211, 8605, 4302)-difference set in Z_17211

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006748 | ljcr-ds-1728-628-228-12-144

(1728, 628, 228)-difference set in Z_12 x Z_144

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006749 | ljcr-ds-1728-628-228-2-12-72

(1728, 628, 228)-difference set in Z_2 x Z_12 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006750 | ljcr-ds-1728-628-228-2-2-2-216

(1728, 628, 228)-difference set in Z_2 x Z_2 x Z_2 x Z_216

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006751 | ljcr-ds-1728-628-228-2-2-6-72

(1728, 628, 228)-difference set in Z_2 x Z_2 x Z_6 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006752 | ljcr-ds-1728-628-228-2-4-216

(1728, 628, 228)-difference set in Z_2 x Z_4 x Z_216

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006753 | ljcr-ds-1728-628-228-2-6-144

(1728, 628, 228)-difference set in Z_2 x Z_6 x Z_144

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006754 | ljcr-ds-1728-628-228-2-6-6-24

(1728, 628, 228)-difference set in Z_2 x Z_6 x Z_6 x Z_24

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006755 | ljcr-ds-1728-628-228-24-72

(1728, 628, 228)-difference set in Z_24 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006756 | ljcr-ds-1728-628-228-3-12-48

(1728, 628, 228)-difference set in Z_3 x Z_12 x Z_48

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006757 | ljcr-ds-1728-628-228-3-24-24

(1728, 628, 228)-difference set in Z_3 x Z_24 x Z_24

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006758 | ljcr-ds-1728-628-228-6-12-24

(1728, 628, 228)-difference set in Z_6 x Z_12 x Z_24

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006759 | ljcr-ds-1728-628-228-6-6-48

(1728, 628, 228)-difference set in Z_6 x Z_6 x Z_48

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006760 | ljcr-ds-1728-628-228-8-216

(1728, 628, 228)-difference set in Z_8 x Z_216

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006761 | ljcr-ds-17301-4325-1081-17301

(17301, 4325, 1081)-difference set in Z_17301

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006762 | ljcr-ds-17319-3711-795-17319

(17319, 3711, 795)-difference set in Z_17319

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006763 | ljcr-ds-17329-5776-1925-17329

(17329, 5776, 1925)-difference set in Z_17329

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006764 | ljcr-ds-17347-3186-585-17347

(17347, 3186, 585)-difference set in Z_17347

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006765 | ljcr-ds-17391-5735-1891-17391

(17391, 5735, 1891)-difference set in Z_17391

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006766 | ljcr-ds-17395-2899-483-17395

(17395, 2899, 483)-difference set in Z_17395

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006767 | ljcr-ds-17423-8711-4355-17423

(17423, 8711, 4355)-difference set in Z_17423

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006768 | ljcr-ds-17425-1089-68-17425

(17425, 1089, 68)-difference set in Z_17425

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006769 | ljcr-ds-17445-4361-1090-17445

(17445, 4361, 1090)-difference set in Z_17445

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006770 | ljcr-ds-17451-8376-4020-17451

(17451, 8376, 4020)-difference set in Z_17451

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006771 | ljcr-ds-17459-5075-1475-17459

(17459, 5075, 1475)-difference set in Z_17459

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006772 | ljcr-ds-17461-2620-393-17461

(17461, 2620, 393)-difference set in Z_17461

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006773 | ljcr-ds-17461-8245-3893-17461

(17461, 8245, 3893)-difference set in Z_17461

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006774 | ljcr-ds-17480-3179-578-17480

(17480, 3179, 578)-difference set in Z_17480

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006775 | ljcr-ds-17487-8743-4371-17487

(17487, 8743, 4371)-difference set in Z_17487

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006776 | ljcr-ds-17497-6561-2460-17497

(17497, 6561, 2460)-difference set in Z_17497

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006777 | ljcr-ds-1750-265-40-5-5-70

(1750, 265, 40)-difference set in Z_5 x Z_5 x Z_70

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006778 | ljcr-ds-17601-3201-582-17601

(17601, 3201, 582)-difference set in Z_17601

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006779 | ljcr-ds-17613-1037-61-17613

(17613, 1037, 61)-difference set in Z_17613

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006780 | ljcr-ds-17613-1037-61-3-5871

(17613, 1037, 61)-difference set in Z_3 x Z_5871

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006781 | ljcr-ds-1764-861-420-21-84

(1764, 861, 420)-difference set in Z_21 x Z_84

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006782 | ljcr-ds-1764-861-420-3-588

(1764, 861, 420)-difference set in Z_3 x Z_588

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006783 | ljcr-ds-1764-861-420-42-42

(1764, 861, 420)-difference set in Z_42 x Z_42

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006784 | ljcr-ds-1764-861-420-6-294

(1764, 861, 420)-difference set in Z_6 x Z_294

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006785 | ljcr-ds-17647-8823-4411-17647

(17647, 8823, 4411)-difference set in Z_17647

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006786 | ljcr-ds-17671-4960-1392-17671

(17671, 4960, 1392)-difference set in Z_17671

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006787 | ljcr-ds-17675-8837-4418-17675

(17675, 8837, 4418)-difference set in Z_17675

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006788 | ljcr-ds-1768-837-396-1768

(1768, 837, 396)-difference set in Z_1768

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006789 | ljcr-ds-1768-837-396-2-884

(1768, 837, 396)-difference set in Z_2 x Z_884

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006790 | ljcr-ds-1773-444-111-3-591

(1773, 444, 111)-difference set in Z_3 x Z_591

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006791 | ljcr-ds-17743-8871-4435-17743

(17743, 8871, 4435)-difference set in Z_17743

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006792 | ljcr-ds-17761-5920-1973-17761

(17761, 5920, 1973)-difference set in Z_17761

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006793 | ljcr-ds-17763-4067-931-17763

(17763, 4067, 931)-difference set in Z_17763

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006794 | ljcr-ds-17803-8901-4450-17803

(17803, 8901, 4450)-difference set in Z_17803

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006795 | ljcr-ds-17811-8905-4452-17811

(17811, 8905, 4452)-difference set in Z_17811

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006796 | ljcr-ds-17829-4457-1114-17829

(17829, 4457, 1114)-difference set in Z_17829

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006797 | ljcr-ds-17831-3566-713-17831

(17831, 3566, 713)-difference set in Z_17831

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006798 | ljcr-ds-1785-224-28-1785

(1785, 224, 28)-difference set in Z_1785

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006799 | ljcr-ds-17860-5953-1984-17860

(17860, 5953, 1984)-difference set in Z_17860

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006800 | ljcr-ds-17871-8935-4467-17871

(17871, 8935, 4467)-difference set in Z_17871

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006801 | ljcr-ds-1791-895-447-3-597

(1791, 895, 447)-difference set in Z_3 x Z_597

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006802 | ljcr-ds-17955-8977-4488-17955

(17955, 8977, 4488)-difference set in Z_17955

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006803 | ljcr-ds-17967-8983-4491-17967

(17967, 8983, 4491)-difference set in Z_17967

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006804 | ljcr-ds-1800-771-330-15-120

(1800, 771, 330)-difference set in Z_15 x Z_120

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006805 | ljcr-ds-1800-771-330-1800

(1800, 771, 330)-difference set in Z_1800

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006806 | ljcr-ds-1800-771-330-2-30-30

(1800, 771, 330)-difference set in Z_2 x Z_30 x Z_30

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006807 | ljcr-ds-1800-771-330-2-6-150

(1800, 771, 330)-difference set in Z_2 x Z_6 x Z_150

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006808 | ljcr-ds-1800-771-330-3-600

(1800, 771, 330)-difference set in Z_3 x Z_600

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006809 | ljcr-ds-1800-771-330-30-60

(1800, 771, 330)-difference set in Z_30 x Z_60

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006810 | ljcr-ds-1800-771-330-5-360

(1800, 771, 330)-difference set in Z_5 x Z_360

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006811 | ljcr-ds-1800-771-330-6-300

(1800, 771, 330)-difference set in Z_6 x Z_300

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006812 | ljcr-ds-1801-225-28-1801

(1801, 225, 28)-difference set in Z_1801

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006813 | ljcr-ds-18063-2464-336-18063

(18063, 2464, 336)-difference set in Z_18063

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006814 | ljcr-ds-18095-7304-2948-18095

(18095, 7304, 2948)-difference set in Z_18095

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006815 | ljcr-ds-18096-6721-2496-18096

(18096, 6721, 2496)-difference set in Z_18096

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006816 | ljcr-ds-18099-9049-4524-18099

(18099, 9049, 4524)-difference set in Z_18099

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006817 | ljcr-ds-18121-2265-283-18121

(18121, 2265, 283)-difference set in Z_18121

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006818 | ljcr-ds-18151-9075-4537-18151

(18151, 9075, 4537)-difference set in Z_18151

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006819 | ljcr-ds-18163-1009-56-18163

(18163, 1009, 56)-difference set in Z_18163

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006820 | ljcr-ds-18187-6496-2320-18187

(18187, 6496, 2320)-difference set in Z_18187

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006821 | ljcr-ds-18189-4548-1137-18189

(18189, 4548, 1137)-difference set in Z_18189

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006822 | ljcr-ds-18237-3008-496-18237

(18237, 3008, 496)-difference set in Z_18237

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006823 | ljcr-ds-18261-7305-2922-18261

(18261, 7305, 2922)-difference set in Z_18261

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006824 | ljcr-ds-18267-9133-4566-18267

(18267, 9133, 4566)-difference set in Z_18267

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006825 | ljcr-ds-18321-3665-733-18321

(18321, 3665, 733)-difference set in Z_18321

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006826 | ljcr-ds-18331-6345-2196-18331

(18331, 6345, 2196)-difference set in Z_18331

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006827 | ljcr-ds-18351-9175-4587-18351

(18351, 9175, 4587)-difference set in Z_18351

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006828 | ljcr-ds-1836-735-294-3-3-204

(1836, 735, 294)-difference set in Z_3 x Z_3 x Z_204

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006829 | ljcr-ds-1836-735-294-3-6-102

(1836, 735, 294)-difference set in Z_3 x Z_6 x Z_102

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006830 | ljcr-ds-1836-735-294-3-612

(1836, 735, 294)-difference set in Z_3 x Z_612

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006831 | ljcr-ds-1836-735-294-6-306

(1836, 735, 294)-difference set in Z_6 x Z_306

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006832 | ljcr-ds-18363-9181-4590-18363

(18363, 9181, 4590)-difference set in Z_18363

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006833 | ljcr-ds-18403-3067-511-18403

(18403, 3067, 511)-difference set in Z_18403

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006834 | ljcr-ds-18415-9207-4603-18415

(18415, 9207, 4603)-difference set in Z_18415

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006835 | ljcr-ds-18441-2305-288-18441

(18441, 2305, 288)-difference set in Z_18441

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006836 | ljcr-ds-18447-9223-4611-18447

(18447, 9223, 4611)-difference set in Z_18447

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006837 | ljcr-ds-1849-561-170-43-43

(1849, 561, 170)-difference set in Z_43 x Z_43

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006838 | ljcr-ds-18501-4625-1156-18501

(18501, 4625, 1156)-difference set in Z_18501

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006839 | ljcr-ds-18507-3896-820-18507

(18507, 3896, 820)-difference set in Z_18507

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006840 | ljcr-ds-18507-9253-4626-18507

(18507, 9253, 4626)-difference set in Z_18507

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006841 | ljcr-ds-18538-6513-2288-18538

(18538, 6513, 2288)-difference set in Z_18538

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006842 | ljcr-ds-18571-9285-4642-18571

(18571, 9285, 4642)-difference set in Z_18571

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006843 | ljcr-ds-18585-6969-2613-18585

(18585, 6969, 2613)-difference set in Z_18585

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006844 | ljcr-ds-18681-1401-105-18681

(18681, 1401, 105)-difference set in Z_18681

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006845 | ljcr-ds-18723-9361-4680-18723

(18723, 9361, 4680)-difference set in Z_18723

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006846 | ljcr-ds-18733-1561-130-18733

(18733, 1561, 130)-difference set in Z_18733

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006847 | ljcr-ds-18735-9367-4683-18735

(18735, 9367, 4683)-difference set in Z_18735

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006848 | ljcr-ds-18747-9373-4686-18747

(18747, 9373, 4686)-difference set in Z_18747

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006849 | ljcr-ds-18755-9377-4688-18755

(18755, 9377, 4688)-difference set in Z_18755

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006850 | ljcr-ds-18762-6425-2200-18762

(18762, 6425, 2200)-difference set in Z_18762

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006851 | ljcr-ds-1881-705-264-1881

(1881, 705, 264)-difference set in Z_1881

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006852 | ljcr-ds-1881-705-264-3-627

(1881, 705, 264)-difference set in Z_3 x Z_627

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006853 | ljcr-ds-1886-261-36-1886

(1886, 261, 36)-difference set in Z_1886

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006854 | ljcr-ds-1887-575-175-1887

(1887, 575, 175)-difference set in Z_1887

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006855 | ljcr-ds-18877-5929-1862-18877

(18877, 5929, 1862)-difference set in Z_18877

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006856 | ljcr-ds-18915-6370-2145-18915

(18915, 6370, 2145)-difference set in Z_18915

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006857 | ljcr-ds-18928-702-26-2-2-4732

(18928, 702, 26)-difference set in Z_2 x Z_2 x Z_4732

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006858 | ljcr-ds-18928-702-26-2-26-364

(18928, 702, 26)-difference set in Z_2 x Z_26 x Z_364

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006859 | ljcr-ds-18928-702-26-26-728

(18928, 702, 26)-difference set in Z_26 x Z_728

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006860 | ljcr-ds-18928-702-26-4-4732

(18928, 702, 26)-difference set in Z_4 x Z_4732

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006861 | ljcr-ds-18928-702-26-52-364

(18928, 702, 26)-difference set in Z_52 x Z_364

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006862 | ljcr-ds-18931-9465-4732-18931

(18931, 9465, 4732)-difference set in Z_18931

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006863 | ljcr-ds-1897-633-211-1897

(1897, 633, 211)-difference set in Z_1897

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006864 | ljcr-ds-18981-4745-1186-18981

(18981, 4745, 1186)-difference set in Z_18981

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006865 | ljcr-ds-1899-949-474-3-633

(1899, 949, 474)-difference set in Z_3 x Z_633

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006866 | ljcr-ds-18991-9495-4747-18991

(18991, 9495, 4747)-difference set in Z_18991

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006867 | ljcr-ds-19003-6334-2111-19003

(19003, 6334, 2111)-difference set in Z_19003

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006868 | ljcr-ds-19033-4576-1100-19033

(19033, 4576, 1100)-difference set in Z_19033

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006869 | ljcr-ds-1904-693-252-1904

(1904, 693, 252)-difference set in Z_1904

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006870 | ljcr-ds-1904-693-252-2-952

(1904, 693, 252)-difference set in Z_2 x Z_952

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006871 | ljcr-ds-19045-4761-1190-19045

(19045, 4761, 1190)-difference set in Z_19045

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006872 | ljcr-ds-19093-6364-2121-19093

(19093, 6364, 2121)-difference set in Z_19093

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006873 | ljcr-ds-19117-5428-1541-19117

(19117, 5428, 1541)-difference set in Z_19117

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006874 | ljcr-ds-19131-9565-4782-19131

(19131, 9565, 4782)-difference set in Z_19131

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006875 | ljcr-ds-19177-6393-2131-19177

(19177, 6393, 2131)-difference set in Z_19177

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006876 | ljcr-ds-19177-8977-4202-19177

(19177, 8977, 4202)-difference set in Z_19177

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006877 | ljcr-ds-1920-304-48-2-2-2-2-120

(1920, 304, 48)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_120

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006878 | ljcr-ds-1920-304-48-2-2-2-2-2-2-30

(1920, 304, 48)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_30

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006879 | ljcr-ds-1920-304-48-2-2-2-2-2-60

(1920, 304, 48)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_60

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006880 | ljcr-ds-1920-304-48-2-2-2-240

(1920, 304, 48)-difference set in Z_2 x Z_2 x Z_2 x Z_240

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006881 | ljcr-ds-1920-304-48-2-2-2-4-60

(1920, 304, 48)-difference set in Z_2 x Z_2 x Z_2 x Z_4 x Z_60

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006882 | ljcr-ds-1920-304-48-2-2-4-120

(1920, 304, 48)-difference set in Z_2 x Z_2 x Z_4 x Z_120

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006883 | ljcr-ds-1920-304-48-2-4-240

(1920, 304, 48)-difference set in Z_2 x Z_4 x Z_240

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006884 | ljcr-ds-1920-304-48-2-4-4-60

(1920, 304, 48)-difference set in Z_2 x Z_4 x Z_4 x Z_60

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006885 | ljcr-ds-1920-304-48-2-8-120

(1920, 304, 48)-difference set in Z_2 x Z_8 x Z_120

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006886 | ljcr-ds-1920-304-48-4-4-120

(1920, 304, 48)-difference set in Z_4 x Z_4 x Z_120

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006887 | ljcr-ds-1920-304-48-8-240

(1920, 304, 48)-difference set in Z_8 x Z_240

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006888 | ljcr-ds-19200-1315-90-19200

(19200, 1315, 90)-difference set in Z_19200

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006889 | ljcr-ds-19201-5376-1505-19201

(19201, 5376, 1505)-difference set in Z_19201

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006890 | ljcr-ds-19215-9607-4803-19215

(19215, 9607, 4803)-difference set in Z_19215

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006891 | ljcr-ds-19228-8671-3910-19228

(19228, 8671, 3910)-difference set in Z_19228

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006892 | ljcr-ds-19241-7696-3078-19241

(19241, 7696, 3078)-difference set in Z_19241

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006893 | ljcr-ds-1925-260-35-1925

(1925, 260, 35)-difference set in Z_1925

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006894 | ljcr-ds-1925-260-35-5-385

(1925, 260, 35)-difference set in Z_5 x Z_385

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006895 | ljcr-ds-19273-6424-2141-19273

(19273, 6424, 2141)-difference set in Z_19273

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006896 | ljcr-ds-19293-8268-3543-19293

(19293, 8268, 3543)-difference set in Z_19293

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006897 | ljcr-ds-1935-967-483-3-645

(1935, 967, 483)-difference set in Z_3 x Z_645

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006898 | ljcr-ds-1936-946-462-22-88

(1936, 946, 462)-difference set in Z_22 x Z_88

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006899 | ljcr-ds-1936-946-462-44-44

(1936, 946, 462)-difference set in Z_44 x Z_44

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006900 | ljcr-ds-19399-4453-1022-19399

(19399, 4453, 1022)-difference set in Z_19399

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006901 | ljcr-ds-19407-9703-4851-19407

(19407, 9703, 4851)-difference set in Z_19407

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006902 | ljcr-ds-19429-4857-1214-19429

(19429, 4857, 1214)-difference set in Z_19429

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006903 | ljcr-ds-19443-9721-4860-19443

(19443, 9721, 4860)-difference set in Z_19443

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006904 | ljcr-ds-19449-1496-115-19449

(19449, 1496, 115)-difference set in Z_19449

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006905 | ljcr-ds-19475-3745-720-19475

(19475, 3745, 720)-difference set in Z_19475

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006906 | ljcr-ds-19475-5993-1844-19475

(19475, 5993, 1844)-difference set in Z_19475

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006907 | ljcr-ds-19481-8280-3519-19481

(19481, 8280, 3519)-difference set in Z_19481

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006908 | ljcr-ds-1953-672-231-3-651

(1953, 672, 231)-difference set in Z_3 x Z_651

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006909 | ljcr-ds-19531-7750-3075-19531

(19531, 7750, 3075)-difference set in Z_19531

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006910 | ljcr-ds-19550-5537-1568-19550

(19550, 5537, 1568)-difference set in Z_19550

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006911 | ljcr-ds-19552-6175-1950-19552

(19552, 6175, 1950)-difference set in Z_19552

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006912 | ljcr-ds-1957-816-340-1957

(1957, 816, 340)-difference set in Z_1957

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006913 | ljcr-ds-19599-9799-4899-19599

(19599, 9799, 4899)-difference set in Z_19599

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006914 | ljcr-ds-19611-9805-4902-19611

(19611, 9805, 4902)-difference set in Z_19611

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006915 | ljcr-ds-19647-5358-1461-19647

(19647, 5358, 1461)-difference set in Z_19647

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006916 | ljcr-ds-19647-7239-2667-19647

(19647, 7239, 2667)-difference set in Z_19647

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006917 | ljcr-ds-19665-6145-1920-19665

(19665, 6145, 1920)-difference set in Z_19665

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006918 | ljcr-ds-19685-4921-1230-19685

(19685, 4921, 1230)-difference set in Z_19685

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006919 | ljcr-ds-19695-9847-4923-19695

(19695, 9847, 4923)-difference set in Z_19695

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006920 | ljcr-ds-19717-4929-1232-19717

(19717, 4929, 1232)-difference set in Z_19717

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006921 | ljcr-ds-1975-330-55-5-395

(1975, 330, 55)-difference set in Z_5 x Z_395

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006922 | ljcr-ds-19780-7981-3220-19780

(19780, 7981, 3220)-difference set in Z_19780

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006923 | ljcr-ds-19781-4945-1236-19781

(19781, 4945, 1236)-difference set in Z_19781

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006924 | ljcr-ds-19783-9891-4945-19783

(19783, 9891, 4945)-difference set in Z_19783

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006925 | ljcr-ds-19811-8491-3639-19811

(19811, 8491, 3639)-difference set in Z_19811

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006926 | ljcr-ds-19833-3552-636-19833

(19833, 3552, 636)-difference set in Z_19833

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006927 | ljcr-ds-19836-7935-3174-19836

(19836, 7935, 3174)-difference set in Z_19836

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006928 | ljcr-ds-1984-661-220-2-2-496

(1984, 661, 220)-difference set in Z_2 x Z_2 x Z_496

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006929 | ljcr-ds-1984-661-220-2-992

(1984, 661, 220)-difference set in Z_2 x Z_992

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006930 | ljcr-ds-1984-661-220-4-496

(1984, 661, 220)-difference set in Z_4 x Z_496

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006931 | ljcr-ds-19881-1065-57-141-141

(19881, 1065, 57)-difference set in Z_141 x Z_141

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006932 | ljcr-ds-19881-1065-57-47-423

(19881, 1065, 57)-difference set in Z_47 x Z_423

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006933 | ljcr-ds-19895-7889-3128-19895

(19895, 7889, 3128)-difference set in Z_19895

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006934 | ljcr-ds-19899-9949-4974-19899

(19899, 9949, 4974)-difference set in Z_19899

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006935 | ljcr-ds-19903-6634-2211-19903

(19903, 6634, 2211)-difference set in Z_19903

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006936 | ljcr-ds-1991-200-20-1991

(1991, 200, 20)-difference set in Z_1991

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006937 | ljcr-ds-19921-7056-2499-19921

(19921, 7056, 2499)-difference set in Z_19921

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006938 | ljcr-ds-19933-7249-2636-19933

(19933, 7249, 2636)-difference set in Z_19933

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006939 | ljcr-ds-19983-9991-4995-19983

(19983, 9991, 4995)-difference set in Z_19983

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006940 | ljcr-ds-19985-1249-78-19985

(19985, 1249, 78)-difference set in Z_19985

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006941 | ljcr-ds-19988-7821-3060-19988

(19988, 7821, 3060)-difference set in Z_19988

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006942 | ljcr-ds-20007-1429-102-20007

(20007, 1429, 102)-difference set in Z_20007

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006943 | ljcr-ds-20031-10015-5007-20031

(20031, 10015, 5007)-difference set in Z_20031

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006944 | ljcr-ds-20033-2817-396-20033

(20033, 2817, 396)-difference set in Z_20033

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006945 | ljcr-ds-20037-5009-1252-20037

(20037, 5009, 1252)-difference set in Z_20037

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006946 | ljcr-ds-20083-10041-5020-20083

(20083, 10041, 5020)-difference set in Z_20083

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006947 | ljcr-ds-20083-6694-2231-20083

(20083, 6694, 2231)-difference set in Z_20083

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006948 | ljcr-ds-20101-5025-1256-20101

(20101, 5025, 1256)-difference set in Z_20101

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006949 | ljcr-ds-2016-156-12-12-168

(2016, 156, 12)-difference set in Z_12 x Z_168

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006950 | ljcr-ds-2016-156-12-2-6-168

(2016, 156, 12)-difference set in Z_2 x Z_6 x Z_168

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006951 | ljcr-ds-2016-156-12-6-336

(2016, 156, 12)-difference set in Z_6 x Z_336

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006952 | ljcr-ds-2016-651-210-12-168

(2016, 651, 210)-difference set in Z_12 x Z_168

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006953 | ljcr-ds-2016-651-210-2-1008

(2016, 651, 210)-difference set in Z_2 x Z_1008

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006954 | ljcr-ds-2016-651-210-2-12-84

(2016, 651, 210)-difference set in Z_2 x Z_12 x Z_84

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006955 | ljcr-ds-2016-651-210-2-2-2-6-42

(2016, 651, 210)-difference set in Z_2 x Z_2 x Z_2 x Z_6 x Z_42

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006956 | ljcr-ds-2016-651-210-2-2-504

(2016, 651, 210)-difference set in Z_2 x Z_2 x Z_504

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006957 | ljcr-ds-2016-651-210-2-2-6-84

(2016, 651, 210)-difference set in Z_2 x Z_2 x Z_6 x Z_84

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006958 | ljcr-ds-2016-651-210-2-6-168

(2016, 651, 210)-difference set in Z_2 x Z_6 x Z_168

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006959 | ljcr-ds-2016-651-210-3-672

(2016, 651, 210)-difference set in Z_3 x Z_672

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006960 | ljcr-ds-2016-651-210-4-504

(2016, 651, 210)-difference set in Z_4 x Z_504

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006961 | ljcr-ds-2016-651-210-6-336

(2016, 651, 210)-difference set in Z_6 x Z_336

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006962 | ljcr-ds-2016-806-322-12-168

(2016, 806, 322)-difference set in Z_12 x Z_168

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006963 | ljcr-ds-2016-806-322-2-1008

(2016, 806, 322)-difference set in Z_2 x Z_1008

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006964 | ljcr-ds-2016-806-322-2-2-504

(2016, 806, 322)-difference set in Z_2 x Z_2 x Z_504

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006965 | ljcr-ds-2016-806-322-2-6-168

(2016, 806, 322)-difference set in Z_2 x Z_6 x Z_168

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006966 | ljcr-ds-2016-806-322-4-504

(2016, 806, 322)-difference set in Z_4 x Z_504

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006967 | ljcr-ds-2016-806-322-6-336

(2016, 806, 322)-difference set in Z_6 x Z_336

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006968 | ljcr-ds-20163-10081-5040-20163

(20163, 10081, 5040)-difference set in Z_20163

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006969 | ljcr-ds-20167-3361-560-20167

(20167, 3361, 560)-difference set in Z_20167

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006970 | ljcr-ds-20173-6724-2241-20173

(20173, 6724, 2241)-difference set in Z_20173

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006971 | ljcr-ds-20196-1155-66-20196

(20196, 1155, 66)-difference set in Z_20196

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006972 | ljcr-ds-20197-5049-1262-20197

(20197, 5049, 1262)-difference set in Z_20197

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006973 | ljcr-ds-20197-5644-1577-20197

(20197, 5644, 1577)-difference set in Z_20197

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006974 | ljcr-ds-20207-10103-5051-20207

(20207, 10103, 5051)-difference set in Z_20207

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006975 | ljcr-ds-20229-5057-1264-20229

(20229, 5057, 1264)-difference set in Z_20229

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006976 | ljcr-ds-20240-7659-2898-20240

(20240, 7659, 2898)-difference set in Z_20240

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006977 | ljcr-ds-20241-7360-2676-20241

(20241, 7360, 2676)-difference set in Z_20241

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006978 | ljcr-ds-20257-6753-2251-20257

(20257, 6753, 2251)-difference set in Z_20257

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006979 | ljcr-ds-20273-6336-1980-20273

(20273, 6336, 1980)-difference set in Z_20273

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006980 | ljcr-ds-2029-508-127-2029

(2029, 508, 127)-difference set in Z_2029

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006981 | ljcr-ds-20306-4061-812-20306

(20306, 4061, 812)-difference set in Z_20306

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006982 | ljcr-ds-20383-3871-735-20383

(20383, 3871, 735)-difference set in Z_20383

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006983 | ljcr-ds-20384-1309-84-20384

(20384, 1309, 84)-difference set in Z_20384

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006984 | ljcr-ds-20389-5097-1274-20389

(20389, 5097, 1274)-difference set in Z_20389

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006985 | ljcr-ds-20419-3403-567-20419

(20419, 3403, 567)-difference set in Z_20419

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006986 | ljcr-ds-20421-5105-1276-20421

(20421, 5105, 1276)-difference set in Z_20421

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006987 | ljcr-ds-2043-1021-510-3-681

(2043, 1021, 510)-difference set in Z_3 x Z_681

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006988 | ljcr-ds-20437-3537-612-20437

(20437, 3537, 612)-difference set in Z_20437

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006989 | ljcr-ds-20447-10223-5111-20447

(20447, 10223, 5111)-difference set in Z_20447

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006990 | ljcr-ds-20461-6820-2273-20461

(20461, 6820, 2273)-difference set in Z_20461

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006991 | ljcr-ds-20475-2118-219-20475

(20475, 2118, 219)-difference set in Z_20475

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006992 | ljcr-ds-20520-4047-798-20520

(20520, 4047, 798)-difference set in Z_20520

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006993 | ljcr-ds-20527-10263-5131-20527

(20527, 10263, 5131)-difference set in Z_20527

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006994 | ljcr-ds-20549-5137-1284-20549

(20549, 5137, 1284)-difference set in Z_20549

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006995 | ljcr-ds-20555-10277-5138-20555

(20555, 10277, 5138)-difference set in Z_20555

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006996 | ljcr-ds-20581-4116-823-20581

(20581, 4116, 823)-difference set in Z_20581

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006997 | ljcr-ds-20599-3433-572-20599

(20599, 3433, 572)-difference set in Z_20599

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006998 | ljcr-ds-2061-516-129-3-687

(2061, 516, 129)-difference set in Z_3 x Z_687

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-006999 | ljcr-ds-20671-8905-3836-20671

(20671, 8905, 3836)-difference set in Z_20671

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007000 | ljcr-ds-2068-637-196-2068

(2068, 637, 196)-difference set in Z_2068

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007001 | ljcr-ds-20680-3051-450-20680

(20680, 3051, 450)-difference set in Z_20680

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007002 | ljcr-ds-20735-7406-2645-20735

(20735, 7406, 2645)-difference set in Z_20735

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007003 | ljcr-ds-20749-5188-1297-20749

(20749, 5188, 1297)-difference set in Z_20749

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007004 | ljcr-ds-20751-10375-5187-20751

(20751, 10375, 5187)-difference set in Z_20751

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007005 | ljcr-ds-20791-9856-4672-20791

(20791, 9856, 4672)-difference set in Z_20791

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007006 | ljcr-ds-20799-10399-5199-20799

(20799, 10399, 5199)-difference set in Z_20799

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007007 | ljcr-ds-2080-540-140-2-2-2-260

(2080, 540, 140)-difference set in Z_2 x Z_2 x Z_2 x Z_260

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007008 | ljcr-ds-2080-540-140-2-2-520

(2080, 540, 140)-difference set in Z_2 x Z_2 x Z_520

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007009 | ljcr-ds-2080-540-140-2-4-260

(2080, 540, 140)-difference set in Z_2 x Z_4 x Z_260

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007010 | ljcr-ds-2080-540-140-4-520

(2080, 540, 140)-difference set in Z_4 x Z_520

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007011 | ljcr-ds-20809-2313-257-20809

(20809, 2313, 257)-difference set in Z_20809

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007012 | ljcr-ds-20809-2601-325-20809

(20809, 2601, 325)-difference set in Z_20809

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007013 | ljcr-ds-20815-10407-5203-20815

(20815, 10407, 5203)-difference set in Z_20815

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007014 | ljcr-ds-20819-7435-2655-20819

(20819, 7435, 2655)-difference set in Z_20819

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007015 | ljcr-ds-20823-2872-396-20823

(20823, 2872, 396)-difference set in Z_20823

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007016 | ljcr-ds-20847-10423-5211-20847

(20847, 10423, 5211)-difference set in Z_20847

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007017 | ljcr-ds-2091-760-276-2091

(2091, 760, 276)-difference set in Z_2091

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007018 | ljcr-ds-20911-10455-5227-20911

(20911, 10455, 5227)-difference set in Z_20911

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007019 | ljcr-ds-20945-6545-2045-20945

(20945, 6545, 2045)-difference set in Z_20945

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007020 | ljcr-ds-20945-8449-3408-20945

(20945, 8449, 3408)-difference set in Z_20945

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007021 | ljcr-ds-20955-10477-5238-20955

(20955, 10477, 5238)-difference set in Z_20955

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007022 | ljcr-ds-20995-3499-583-20995

(20995, 3499, 583)-difference set in Z_20995

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007023 | ljcr-ds-21029-3004-429-21029

(21029, 3004, 429)-difference set in Z_21029

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007024 | ljcr-ds-21043-10521-5260-21043

(21043, 10521, 5260)-difference set in Z_21043

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007025 | ljcr-ds-21051-10525-5262-21051

(21051, 10525, 5262)-difference set in Z_21051

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007026 | ljcr-ds-21055-9802-4563-21055

(21055, 9802, 4563)-difference set in Z_21055

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007027 | ljcr-ds-21065-2633-329-21065

(21065, 2633, 329)-difference set in Z_21065

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007028 | ljcr-ds-21069-5268-1317-21069

(21069, 5268, 1317)-difference set in Z_21069

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007029 | ljcr-ds-21091-2110-211-21091

(21091, 2110, 211)-difference set in Z_21091

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007030 | ljcr-ds-21113-4264-861-21113

(21113, 4264, 861)-difference set in Z_21113

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007031 | ljcr-ds-21113-5656-1515-21113

(21113, 5656, 1515)-difference set in Z_21113

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007032 | ljcr-ds-21137-9248-4046-21137

(21137, 9248, 4046)-difference set in Z_21137

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007033 | ljcr-ds-21141-756-27-27-783

(21141, 756, 27)-difference set in Z_27 x Z_783

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007034 | ljcr-ds-21141-756-27-3-3-3-783

(21141, 756, 27)-difference set in Z_3 x Z_3 x Z_3 x Z_783

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007035 | ljcr-ds-21141-756-27-3-3-9-261

(21141, 756, 27)-difference set in Z_3 x Z_3 x Z_9 x Z_261

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007036 | ljcr-ds-21141-756-27-3-9-783

(21141, 756, 27)-difference set in Z_3 x Z_9 x Z_783

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007037 | ljcr-ds-21141-756-27-9-9-261

(21141, 756, 27)-difference set in Z_9 x Z_9 x Z_261

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007038 | ljcr-ds-21151-7426-2607-21151

(21151, 7426, 2607)-difference set in Z_21151

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007039 | ljcr-ds-21157-5289-1322-21157

(21157, 5289, 1322)-difference set in Z_21157

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007040 | ljcr-ds-2116-1035-506-23-92

(2116, 1035, 506)-difference set in Z_23 x Z_92

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007041 | ljcr-ds-2116-1035-506-46-46

(2116, 1035, 506)-difference set in Z_46 x Z_46

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007042 | ljcr-ds-2117-529-132-2117

(2117, 529, 132)-difference set in Z_2117

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007043 | ljcr-ds-21183-10591-5295-21183

(21183, 10591, 5295)-difference set in Z_21183

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007044 | ljcr-ds-21231-10615-5307-21231

(21231, 10615, 5307)-difference set in Z_21231

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007045 | ljcr-ds-21291-10645-5322-21291

(21291, 10645, 5322)-difference set in Z_21291

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007046 | ljcr-ds-21315-10657-5328-21315

(21315, 10657, 5328)-difference set in Z_21315

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007047 | ljcr-ds-21321-4265-853-21321

(21321, 4265, 853)-difference set in Z_21321

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007048 | ljcr-ds-21327-10663-5331-21327

(21327, 10663, 5331)-difference set in Z_21327

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007049 | ljcr-ds-2133-533-133-3-3-237

(2133, 533, 133)-difference set in Z_3 x Z_3 x Z_237

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007050 | ljcr-ds-2133-533-133-3-711

(2133, 533, 133)-difference set in Z_3 x Z_711

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007051 | ljcr-ds-21373-3289-506-21373

(21373, 3289, 506)-difference set in Z_21373

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007052 | ljcr-ds-21387-10693-5346-21387

(21387, 10693, 5346)-difference set in Z_21387

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007053 | ljcr-ds-21403-1189-66-21403

(21403, 1189, 66)-difference set in Z_21403

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007054 | ljcr-ds-21421-4285-857-21421

(21421, 4285, 857)-difference set in Z_21421

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007055 | ljcr-ds-21421-6580-2021-21421

(21421, 6580, 2021)-difference set in Z_21421

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007056 | ljcr-ds-21421-7021-2301-21421

(21421, 7021, 2301)-difference set in Z_21421

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007057 | ljcr-ds-21424-7141-2380-21424

(21424, 7141, 2380)-difference set in Z_21424

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007058 | ljcr-ds-2145-737-253-2145

(2145, 737, 253)-difference set in Z_2145

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007059 | ljcr-ds-21450-5785-1560-21450

(21450, 5785, 1560)-difference set in Z_21450

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007060 | ljcr-ds-21523-7174-2391-21523

(21523, 7174, 2391)-difference set in Z_21523

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007061 | ljcr-ds-21541-7180-2393-21541

(21541, 7180, 2393)-difference set in Z_21541

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007062 | ljcr-ds-21553-6736-2105-21553

(21553, 6736, 2105)-difference set in Z_21553

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007063 | ljcr-ds-21571-4315-863-21571

(21571, 4315, 863)-difference set in Z_21571

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007064 | ljcr-ds-2160-255-30-12-180

(2160, 255, 30)-difference set in Z_12 x Z_180

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007065 | ljcr-ds-2160-255-30-2-1080

(2160, 255, 30)-difference set in Z_2 x Z_1080

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007066 | ljcr-ds-2160-255-30-2-6-180

(2160, 255, 30)-difference set in Z_2 x Z_6 x Z_180

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007067 | ljcr-ds-2160-255-30-2160

(2160, 255, 30)-difference set in Z_2160

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007068 | ljcr-ds-2160-255-30-3-12-60

(2160, 255, 30)-difference set in Z_3 x Z_12 x Z_60

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007069 | ljcr-ds-2160-255-30-3-3-240

(2160, 255, 30)-difference set in Z_3 x Z_3 x Z_240

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007070 | ljcr-ds-2160-255-30-3-6-120

(2160, 255, 30)-difference set in Z_3 x Z_6 x Z_120

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007071 | ljcr-ds-2160-255-30-3-720

(2160, 255, 30)-difference set in Z_3 x Z_720

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007072 | ljcr-ds-2160-255-30-6-360

(2160, 255, 30)-difference set in Z_6 x Z_360

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007073 | ljcr-ds-2160-255-30-6-6-60

(2160, 255, 30)-difference set in Z_6 x Z_6 x Z_60

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007074 | ljcr-ds-21605-5401-1350-21605

(21605, 5401, 1350)-difference set in Z_21605

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007075 | ljcr-ds-21637-5409-1352-21637

(21637, 5409, 1352)-difference set in Z_21637

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007076 | ljcr-ds-21659-10829-5414-21659

(21659, 10829, 5414)-difference set in Z_21659

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007077 | ljcr-ds-21661-8665-3466-21661

(21661, 8665, 3466)-difference set in Z_21661

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007078 | ljcr-ds-21673-4816-1070-21673

(21673, 4816, 1070)-difference set in Z_21673

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007079 | ljcr-ds-21679-3613-602-21679

(21679, 3613, 602)-difference set in Z_21679

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007080 | ljcr-ds-21681-4065-762-21681

(21681, 4065, 762)-difference set in Z_21681

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007081 | ljcr-ds-21701-5425-1356-21701

(21701, 5425, 1356)-difference set in Z_21701

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007082 | ljcr-ds-21707-10853-5426-21707

(21707, 10853, 5426)-difference set in Z_21707

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007083 | ljcr-ds-21721-3801-665-21721

(21721, 3801, 665)-difference set in Z_21721

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007084 | ljcr-ds-21731-6520-1956-21731

(21731, 6520, 1956)-difference set in Z_21731

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007085 | ljcr-ds-2176-726-242-2-1088

(2176, 726, 242)-difference set in Z_2 x Z_1088

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007086 | ljcr-ds-2176-726-242-2-2-2-272

(2176, 726, 242)-difference set in Z_2 x Z_2 x Z_2 x Z_272

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007087 | ljcr-ds-2176-726-242-2-2-4-136

(2176, 726, 242)-difference set in Z_2 x Z_2 x Z_4 x Z_136

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007088 | ljcr-ds-2176-726-242-2-2-544

(2176, 726, 242)-difference set in Z_2 x Z_2 x Z_544

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007089 | ljcr-ds-2176-726-242-2-4-272

(2176, 726, 242)-difference set in Z_2 x Z_4 x Z_272

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007090 | ljcr-ds-2176-726-242-2-8-136

(2176, 726, 242)-difference set in Z_2 x Z_8 x Z_136

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007091 | ljcr-ds-2176-726-242-4-4-136

(2176, 726, 242)-difference set in Z_4 x Z_4 x Z_136

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007092 | ljcr-ds-2176-726-242-4-544

(2176, 726, 242)-difference set in Z_4 x Z_544

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007093 | ljcr-ds-2176-726-242-8-272

(2176, 726, 242)-difference set in Z_8 x Z_272

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007094 | ljcr-ds-2177-256-30-2177

(2177, 256, 30)-difference set in Z_2177

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007095 | ljcr-ds-21793-4768-1043-21793

(21793, 4768, 1043)-difference set in Z_21793

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007096 | ljcr-ds-21825-1024-48-15-1455

(21825, 1024, 48)-difference set in Z_15 x Z_1455

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007097 | ljcr-ds-21825-1024-48-21825

(21825, 1024, 48)-difference set in Z_21825

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007098 | ljcr-ds-21825-1024-48-3-7275

(21825, 1024, 48)-difference set in Z_3 x Z_7275

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007099 | ljcr-ds-21825-1024-48-5-4365

(21825, 1024, 48)-difference set in Z_5 x Z_4365

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007100 | ljcr-ds-2185-105-5-2185

(2185, 105, 5)-difference set in Z_2185

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007101 | ljcr-ds-21868-7881-2840-21868

(21868, 7881, 2840)-difference set in Z_21868

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007102 | ljcr-ds-21896-3625-600-21896

(21896, 3625, 600)-difference set in Z_21896

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007103 | ljcr-ds-21897-7889-2842-21897

(21897, 7889, 2842)-difference set in Z_21897

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007104 | ljcr-ds-21901-876-35-11-1991

(21901, 876, 35)-difference set in Z_11 x Z_1991

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007105 | ljcr-ds-21945-7385-2485-21945

(21945, 7385, 2485)-difference set in Z_21945

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007106 | ljcr-ds-21951-10975-5487-21951

(21951, 10975, 5487)-difference set in Z_21951

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007107 | ljcr-ds-21967-10983-5491-21967

(21967, 10983, 5491)-difference set in Z_21967

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007108 | ljcr-ds-2197-793-286-13-169

(2197, 793, 286)-difference set in Z_13 x Z_169

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007109 | ljcr-ds-21979-5698-1477-21979

(21979, 5698, 1477)-difference set in Z_21979

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007110 | ljcr-ds-22041-4960-1116-22041

(22041, 4960, 1116)-difference set in Z_22041

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007111 | ljcr-ds-2205-609-168-21-105

(2205, 609, 168)-difference set in Z_21 x Z_105

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007112 | ljcr-ds-2205-609-168-3-735

(2205, 609, 168)-difference set in Z_3 x Z_735

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007113 | ljcr-ds-2205-609-168-7-315

(2205, 609, 168)-difference set in Z_7 x Z_315

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007114 | ljcr-ds-22063-7354-2451-22063

(22063, 7354, 2451)-difference set in Z_22063

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007115 | ljcr-ds-2209-576-150-2209

(2209, 576, 150)-difference set in Z_2209

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007116 | ljcr-ds-22093-5524-1381-22093

(22093, 5524, 1381)-difference set in Z_22093

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007117 | ljcr-ds-22100-2255-230-22100

(22100, 2255, 230)-difference set in Z_22100

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007118 | ljcr-ds-22141-10045-4557-22141

(22141, 10045, 4557)-difference set in Z_22141

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007119 | ljcr-ds-22155-7315-2415-22155

(22155, 7315, 2415)-difference set in Z_22155

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007120 | ljcr-ds-22181-5545-1386-22181

(22181, 5545, 1386)-difference set in Z_22181

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007121 | ljcr-ds-22189-5548-1387-22189

(22189, 5548, 1387)-difference set in Z_22189

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007122 | ljcr-ds-22203-11101-5550-22203

(22203, 11101, 5550)-difference set in Z_22203

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007123 | ljcr-ds-22223-11111-5555-22223

(22223, 11111, 5555)-difference set in Z_22223

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007124 | ljcr-ds-2223-606-165-2223

(2223, 606, 165)-difference set in Z_2223

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007125 | ljcr-ds-2223-606-165-3-741

(2223, 606, 165)-difference set in Z_3 x Z_741

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007126 | ljcr-ds-22231-495-11-22231

(22231, 495, 11)-difference set in Z_22231

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007127 | ljcr-ds-22244-6903-2142-22244

(22244, 6903, 2142)-difference set in Z_22244

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007128 | ljcr-ds-22277-5569-1392-22277

(22277, 5569, 1392)-difference set in Z_22277

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007129 | ljcr-ds-2233-217-21-2233

(2233, 217, 21)-difference set in Z_2233

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007130 | ljcr-ds-22351-3576-572-22351

(22351, 3576, 572)-difference set in Z_22351

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007131 | ljcr-ds-22359-9583-4107-22359

(22359, 9583, 4107)-difference set in Z_22359

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007132 | ljcr-ds-22360-5655-1430-22360

(22360, 5655, 1430)-difference set in Z_22360

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007133 | ljcr-ds-22363-11181-5590-22363

(22363, 11181, 5590)-difference set in Z_22363

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007134 | ljcr-ds-22381-5596-1399-22381

(22381, 5596, 1399)-difference set in Z_22381

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007135 | ljcr-ds-22383-11191-5595-22383

(22383, 11191, 5595)-difference set in Z_22383

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007136 | ljcr-ds-22387-6643-1971-22387

(22387, 6643, 1971)-difference set in Z_22387

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007137 | ljcr-ds-2241-960-411-3-747

(2241, 960, 411)-difference set in Z_3 x Z_747

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007138 | ljcr-ds-22431-6730-2019-22431

(22431, 6730, 2019)-difference set in Z_22431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007139 | ljcr-ds-22441-7480-2493-22441

(22441, 7480, 2493)-difference set in Z_22441

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007140 | ljcr-ds-22461-5616-1404-22461

(22461, 5616, 1404)-difference set in Z_22461

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007141 | ljcr-ds-22485-6644-1963-22485

(22485, 6644, 1963)-difference set in Z_22485

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007142 | ljcr-ds-22495-3750-625-22495

(22495, 3750, 625)-difference set in Z_22495

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007143 | ljcr-ds-22537-7200-2300-22537

(22537, 7200, 2300)-difference set in Z_22537

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007144 | ljcr-ds-22565-5641-1410-22565

(22565, 5641, 1410)-difference set in Z_22565

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007145 | ljcr-ds-22579-3763-627-22579

(22579, 3763, 627)-difference set in Z_22579

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007146 | ljcr-ds-22587-11293-5646-22587

(22587, 11293, 5646)-difference set in Z_22587

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007147 | ljcr-ds-22603-7534-2511-22603

(22603, 7534, 2511)-difference set in Z_22603

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007148 | ljcr-ds-22607-11303-5651-22607

(22607, 11303, 5651)-difference set in Z_22607

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007149 | ljcr-ds-22629-5657-1414-22629

(22629, 5657, 1414)-difference set in Z_22629

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007150 | ljcr-ds-22633-8856-3465-22633

(22633, 8856, 3465)-difference set in Z_22633

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007151 | ljcr-ds-22671-6801-2040-22671

(22671, 6801, 2040)-difference set in Z_22671

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007152 | ljcr-ds-22687-11343-5671-22687

(22687, 11343, 5671)-difference set in Z_22687

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007153 | ljcr-ds-22715-11357-5678-22715

(22715, 11357, 5678)-difference set in Z_22715

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007154 | ljcr-ds-22731-11365-5682-22731

(22731, 11365, 5682)-difference set in Z_22731

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007155 | ljcr-ds-22767-11383-5691-22767

(22767, 11383, 5691)-difference set in Z_22767

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007156 | ljcr-ds-22771-4555-911-22771

(22771, 4555, 911)-difference set in Z_22771

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007157 | ljcr-ds-22777-7593-2531-22777

(22777, 7593, 2531)-difference set in Z_22777

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007158 | ljcr-ds-22781-4556-911-22781

(22781, 4556, 911)-difference set in Z_22781

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007159 | ljcr-ds-22789-5697-1424-22789

(22789, 5697, 1424)-difference set in Z_22789

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007160 | ljcr-ds-22791-2280-228-22791

(22791, 2280, 228)-difference set in Z_22791

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007161 | ljcr-ds-22793-2849-356-22793

(22793, 2849, 356)-difference set in Z_22793

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007162 | ljcr-ds-22793-9472-3936-22793

(22793, 9472, 3936)-difference set in Z_22793

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007163 | ljcr-ds-22803-11401-5700-22803

(22803, 11401, 5700)-difference set in Z_22803

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007164 | ljcr-ds-22857-2857-357-22857

(22857, 2857, 357)-difference set in Z_22857

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007165 | ljcr-ds-22863-11431-5715-22863

(22863, 11431, 5715)-difference set in Z_22863

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007166 | ljcr-ds-22911-3161-436-22911

(22911, 3161, 436)-difference set in Z_22911

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007167 | ljcr-ds-22919-1638-117-22919

(22919, 1638, 117)-difference set in Z_22919

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007168 | ljcr-ds-22925-9900-4275-22925

(22925, 9900, 4275)-difference set in Z_22925

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007169 | ljcr-ds-22940-3277-468-22940

(22940, 3277, 468)-difference set in Z_22940

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007170 | ljcr-ds-22945-3585-560-22945

(22945, 3585, 560)-difference set in Z_22945

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007171 | ljcr-ds-22951-6426-1799-22951

(22951, 6426, 1799)-difference set in Z_22951

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007172 | ljcr-ds-22960-5103-1134-22960

(22960, 5103, 1134)-difference set in Z_22960

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007173 | ljcr-ds-22971-4595-919-22971

(22971, 4595, 919)-difference set in Z_22971

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007174 | ljcr-ds-22991-11495-5747-22991

(22991, 11495, 5747)-difference set in Z_22991

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007175 | ljcr-ds-22991-7866-2691-22991

(22991, 7866, 2691)-difference set in Z_22991

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007176 | ljcr-ds-23002-5577-1352-23002

(23002, 5577, 1352)-difference set in Z_23002

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007177 | ljcr-ds-2301-276-33-2301

(2301, 276, 33)-difference set in Z_2301

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007178 | ljcr-ds-2301-576-144-2301

(2301, 576, 144)-difference set in Z_2301

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007179 | ljcr-ds-23023-11511-5755-23023

(23023, 11511, 5755)-difference set in Z_23023

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007180 | ljcr-ds-23071-3846-641-23071

(23071, 3846, 641)-difference set in Z_23071

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007181 | ljcr-ds-23103-11551-5775-23103

(23103, 11551, 5775)-difference set in Z_23103

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007182 | ljcr-ds-23115-7112-2188-23115

(23115, 7112, 2188)-difference set in Z_23115

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007183 | ljcr-ds-23121-8960-3472-23121

(23121, 8960, 3472)-difference set in Z_23121

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007184 | ljcr-ds-23129-8968-3477-23129

(23129, 8968, 3477)-difference set in Z_23129

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007185 | ljcr-ds-23141-5785-1446-23141

(23141, 5785, 1446)-difference set in Z_23141

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007186 | ljcr-ds-23151-11575-5787-23151

(23151, 11575, 5787)-difference set in Z_23151

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007187 | ljcr-ds-23155-10897-5128-23155

(23155, 10897, 5128)-difference set in Z_23155

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007188 | ljcr-ds-23161-11001-5225-23161

(23161, 11001, 5225)-difference set in Z_23161

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007189 | ljcr-ds-23163-11581-5790-23163

(23163, 11581, 5790)-difference set in Z_23163

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007190 | ljcr-ds-23211-11605-5802-23211

(23211, 11605, 5802)-difference set in Z_23211

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007191 | ljcr-ds-23215-11607-5803-23215

(23215, 11607, 5803)-difference set in Z_23215

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007192 | ljcr-ds-23221-6021-1561-23221

(23221, 6021, 1561)-difference set in Z_23221

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007193 | ljcr-ds-23241-9961-4269-23241

(23241, 9961, 4269)-difference set in Z_23241

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007194 | ljcr-ds-23257-2584-287-23257

(23257, 2584, 287)-difference set in Z_23257

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007195 | ljcr-ds-23269-5817-1454-23269

(23269, 5817, 1454)-difference set in Z_23269

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007196 | ljcr-ds-23281-4560-893-23281

(23281, 4560, 893)-difference set in Z_23281

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007197 | ljcr-ds-23290-5545-1320-23290

(23290, 5545, 1320)-difference set in Z_23290

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007198 | ljcr-ds-23317-5829-1457-23317

(23317, 5829, 1457)-difference set in Z_23317

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007199 | ljcr-ds-23323-3381-490-23323

(23323, 3381, 490)-difference set in Z_23323

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007200 | ljcr-ds-23323-8281-2940-23323

(23323, 8281, 2940)-difference set in Z_23323

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007201 | ljcr-ds-23331-4666-933-23331

(23331, 4666, 933)-difference set in Z_23331

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007202 | ljcr-ds-23331-7000-2100-23331

(23331, 7000, 2100)-difference set in Z_23331

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007203 | ljcr-ds-23343-11671-5835-23343

(23343, 11671, 5835)-difference set in Z_23343

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007204 | ljcr-ds-2337-512-112-2337

(2337, 512, 112)-difference set in Z_2337

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007205 | ljcr-ds-23401-6201-1643-23401

(23401, 6201, 1643)-difference set in Z_23401

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007206 | ljcr-ds-23401-8425-3033-23401

(23401, 8425, 3033)-difference set in Z_23401

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007207 | ljcr-ds-23421-4685-937-23421

(23421, 4685, 937)-difference set in Z_23421

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007208 | ljcr-ds-23453-7216-2220-23453

(23453, 7216, 2220)-difference set in Z_23453

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007209 | ljcr-ds-23485-1236-65-23485

(23485, 1236, 65)-difference set in Z_23485

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007210 | ljcr-ds-2349-588-147-3-3-261

(2349, 588, 147)-difference set in Z_3 x Z_3 x Z_261

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007211 | ljcr-ds-2349-588-147-3-3-3-87

(2349, 588, 147)-difference set in Z_3 x Z_3 x Z_3 x Z_87

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007212 | ljcr-ds-2349-588-147-9-261

(2349, 588, 147)-difference set in Z_9 x Z_261

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007213 | ljcr-ds-2351-376-60-2351

(2351, 376, 60)-difference set in Z_2351

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007214 | ljcr-ds-23511-2351-235-23511

(23511, 2351, 235)-difference set in Z_23511

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007215 | ljcr-ds-23520-812-28-14-1680

(23520, 812, 28)-difference set in Z_14 x Z_1680

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007216 | ljcr-ds-23520-812-28-2-11760

(23520, 812, 28)-difference set in Z_2 x Z_11760

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007217 | ljcr-ds-23520-812-28-2-14-840

(23520, 812, 28)-difference set in Z_2 x Z_14 x Z_840

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007218 | ljcr-ds-23520-812-28-2-28-420

(23520, 812, 28)-difference set in Z_2 x Z_28 x Z_420

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007219 | ljcr-ds-23520-812-28-28-840

(23520, 812, 28)-difference set in Z_28 x Z_840

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007220 | ljcr-ds-23521-7840-2613-23521

(23521, 7840, 2613)-difference set in Z_23521

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007221 | ljcr-ds-23533-10176-4400-23533

(23533, 10176, 4400)-difference set in Z_23533

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007222 | ljcr-ds-23541-10165-4389-23541

(23541, 10165, 4389)-difference set in Z_23541

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007223 | ljcr-ds-23551-3925-654-23551

(23551, 3925, 654)-difference set in Z_23551

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007224 | ljcr-ds-23557-5889-1472-23557

(23557, 5889, 1472)-difference set in Z_23557

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007225 | ljcr-ds-23607-9658-3951-23607

(23607, 9658, 3951)-difference set in Z_23607

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007226 | ljcr-ds-23685-5921-1480-23685

(23685, 5921, 1480)-difference set in Z_23685

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007227 | ljcr-ds-23695-11847-5923-23695

(23695, 11847, 5923)-difference set in Z_23695

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007228 | ljcr-ds-23715-11857-5928-23715

(23715, 11857, 5928)-difference set in Z_23715

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007229 | ljcr-ds-23723-11861-5930-23723

(23723, 11861, 5930)-difference set in Z_23723

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007230 | ljcr-ds-23737-2968-371-23737

(23737, 2968, 371)-difference set in Z_23737

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007231 | ljcr-ds-23737-7912-2637-23737

(23737, 7912, 2637)-difference set in Z_23737

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007232 | ljcr-ds-23767-3961-660-23767

(23767, 3961, 660)-difference set in Z_23767

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007233 | ljcr-ds-23791-10920-5012-23791

(23791, 10920, 5012)-difference set in Z_23791

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007234 | ljcr-ds-23791-11895-5947-23791

(23791, 11895, 5947)-difference set in Z_23791

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007235 | ljcr-ds-23791-7320-2252-23791

(23791, 7320, 2252)-difference set in Z_23791

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007236 | ljcr-ds-2380-793-264-2-1190

(2380, 793, 264)-difference set in Z_2 x Z_1190

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007237 | ljcr-ds-2380-793-264-2380

(2380, 793, 264)-difference set in Z_2380

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007238 | ljcr-ds-23821-11116-5187-23821

(23821, 11116, 5187)-difference set in Z_23821

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007239 | ljcr-ds-23823-11911-5955-23823

(23823, 11911, 5955)-difference set in Z_23823

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007240 | ljcr-ds-23851-11025-5096-23851

(23851, 11025, 5096)-difference set in Z_23851

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007241 | ljcr-ds-23871-8030-2701-23871

(23871, 8030, 2701)-difference set in Z_23871

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007242 | ljcr-ds-23877-4700-925-23877

(23877, 4700, 925)-difference set in Z_23877

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007243 | ljcr-ds-23881-10945-5016-23881

(23881, 10945, 5016)-difference set in Z_23881

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007244 | ljcr-ds-23883-11941-5970-23883

(23883, 11941, 5970)-difference set in Z_23883

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007245 | ljcr-ds-23895-2757-318-23895

(23895, 2757, 318)-difference set in Z_23895

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007246 | ljcr-ds-23899-7966-2655-23899

(23899, 7966, 2655)-difference set in Z_23899

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007247 | ljcr-ds-23909-5977-1494-23909

(23909, 5977, 1494)-difference set in Z_23909

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007248 | ljcr-ds-23931-11965-5982-23931

(23931, 11965, 5982)-difference set in Z_23931

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007249 | ljcr-ds-23940-1295-70-23940

(23940, 1295, 70)-difference set in Z_23940

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007250 | ljcr-ds-23941-5985-1496-23941

(23941, 5985, 1496)-difference set in Z_23941

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007251 | ljcr-ds-23971-7990-2663-23971

(23971, 7990, 2663)-difference set in Z_23971

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007252 | ljcr-ds-23980-7993-2664-23980

(23980, 7993, 2664)-difference set in Z_23980

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007253 | ljcr-ds-24101-6025-1506-24101

(24101, 6025, 1506)-difference set in Z_24101

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007254 | ljcr-ds-24111-9645-3858-24111

(24111, 9645, 3858)-difference set in Z_24111

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007255 | ljcr-ds-2413-1072-476-2413

(2413, 1072, 476)-difference set in Z_2413

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007256 | ljcr-ds-24147-12073-6036-24147

(24147, 12073, 6036)-difference set in Z_24147

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007257 | ljcr-ds-24155-12077-6038-24155

(24155, 12077, 6038)-difference set in Z_24155

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007258 | ljcr-ds-24171-4835-967-24171

(24171, 4835, 967)-difference set in Z_24171

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007259 | ljcr-ds-24197-6049-1512-24197

(24197, 6049, 1512)-difference set in Z_24197

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007260 | ljcr-ds-24227-12113-6056-24227

(24227, 12113, 6056)-difference set in Z_24227

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007261 | ljcr-ds-24241-9696-3878-24241

(24241, 9696, 3878)-difference set in Z_24241

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007262 | ljcr-ds-2425-505-105-5-485

(2425, 505, 105)-difference set in Z_5 x Z_485

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007263 | ljcr-ds-24273-9472-3696-24273

(24273, 9472, 3696)-difference set in Z_24273

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007264 | ljcr-ds-24289-11616-5555-24289

(24289, 11616, 5555)-difference set in Z_24289

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007265 | ljcr-ds-243-121-60-3-9-9

(243, 121, 60)-difference set in Z_3 x Z_9 x Z_9

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007266 | ljcr-ds-24303-12151-6075-24303

(24303, 12151, 6075)-difference set in Z_24303

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007267 | ljcr-ds-24311-9945-4068-24311

(24311, 9945, 4068)-difference set in Z_24311

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007268 | ljcr-ds-2432-936-360-8-304

(2432, 936, 360)-difference set in Z_8 x Z_304

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007269 | ljcr-ds-24363-12181-6090-24363

(24363, 12181, 6090)-difference set in Z_24363

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007270 | ljcr-ds-24397-8988-3311-24397

(24397, 8988, 3311)-difference set in Z_24397

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007271 | ljcr-ds-24421-8140-2713-24421

(24421, 8140, 2713)-difference set in Z_24421

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007272 | ljcr-ds-24431-12215-6107-24431

(24431, 12215, 6107)-difference set in Z_24431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007273 | ljcr-ds-24431-4886-977-24431

(24431, 4886, 977)-difference set in Z_24431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007274 | ljcr-ds-24435-3858-609-24435

(24435, 3858, 609)-difference set in Z_24435

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007275 | ljcr-ds-24439-8146-2715-24439

(24439, 8146, 2715)-difference set in Z_24439

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007276 | ljcr-ds-24467-7528-2316-24467

(24467, 7528, 2316)-difference set in Z_24467

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007277 | ljcr-ds-24493-2041-170-24493

(24493, 2041, 170)-difference set in Z_24493

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007278 | ljcr-ds-24493-8164-2721-24493

(24493, 8164, 2721)-difference set in Z_24493

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007279 | ljcr-ds-24517-6129-1532-24517

(24517, 6129, 1532)-difference set in Z_24517

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007280 | ljcr-ds-24521-4905-981-24521

(24521, 4905, 981)-difference set in Z_24521

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007281 | ljcr-ds-24559-4093-682-24559

(24559, 4093, 682)-difference set in Z_24559

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007282 | ljcr-ds-24568-1293-68-24568

(24568, 1293, 68)-difference set in Z_24568

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007283 | ljcr-ds-2457-921-345-3-819

(2457, 921, 345)-difference set in Z_3 x Z_819

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007284 | ljcr-ds-24570-4977-1008-24570

(24570, 4977, 1008)-difference set in Z_24570

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007285 | ljcr-ds-24577-8193-2731-24577

(24577, 8193, 2731)-difference set in Z_24577

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007286 | ljcr-ds-24583-8194-2731-24583

(24583, 8194, 2731)-difference set in Z_24583

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007287 | ljcr-ds-24585-10976-4900-24585

(24585, 10976, 4900)-difference set in Z_24585

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007288 | ljcr-ds-24611-2461-246-24611

(24611, 2461, 246)-difference set in Z_24611

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007289 | ljcr-ds-24619-8206-2735-24619

(24619, 8206, 2735)-difference set in Z_24619

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007290 | ljcr-ds-24637-8212-2737-24637

(24637, 8212, 2737)-difference set in Z_24637

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007291 | ljcr-ds-24640-9933-4004-24640

(24640, 9933, 4004)-difference set in Z_24640

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007292 | ljcr-ds-24641-385-6-24641

(24641, 385, 6)-difference set in Z_24641

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007293 | ljcr-ds-24645-6161-1540-24645

(24645, 6161, 1540)-difference set in Z_24645

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007294 | ljcr-ds-24661-6165-1541-24661

(24661, 6165, 1541)-difference set in Z_24661

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007295 | ljcr-ds-24704-3529-504-24704

(24704, 3529, 504)-difference set in Z_24704

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007296 | ljcr-ds-24727-2536-260-24727

(24727, 2536, 260)-difference set in Z_24727

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007297 | ljcr-ds-24753-8177-2701-24753

(24753, 8177, 2701)-difference set in Z_24753

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007298 | ljcr-ds-24781-6196-1549-24781

(24781, 6196, 1549)-difference set in Z_24781

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007299 | ljcr-ds-24831-1911-147-24831

(24831, 1911, 147)-difference set in Z_24831

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007300 | ljcr-ds-2484-573-132-2-1242

(2484, 573, 132)-difference set in Z_2 x Z_1242

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007301 | ljcr-ds-2484-573-132-3-3-276

(2484, 573, 132)-difference set in Z_3 x Z_3 x Z_276

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007302 | ljcr-ds-2484-573-132-3-6-138

(2484, 573, 132)-difference set in Z_3 x Z_6 x Z_138

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007303 | ljcr-ds-2484-573-132-3-828

(2484, 573, 132)-difference set in Z_3 x Z_828

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007304 | ljcr-ds-2484-573-132-6-414

(2484, 573, 132)-difference set in Z_6 x Z_414

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007305 | ljcr-ds-24869-6217-1554-24869

(24869, 6217, 1554)-difference set in Z_24869

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007306 | ljcr-ds-24919-4153-692-24919

(24919, 4153, 692)-difference set in Z_24919

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007307 | ljcr-ds-24931-11080-4924-24931

(24931, 11080, 4924)-difference set in Z_24931

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007308 | ljcr-ds-24957-1836-135-24957

(24957, 1836, 135)-difference set in Z_24957

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007309 | ljcr-ds-2496-500-100-2-1248

(2496, 500, 100)-difference set in Z_2 x Z_1248

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007310 | ljcr-ds-2496-500-100-2-2-2-2-156

(2496, 500, 100)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_156

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007311 | ljcr-ds-2496-500-100-2-2-2-312

(2496, 500, 100)-difference set in Z_2 x Z_2 x Z_2 x Z_312

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007312 | ljcr-ds-2496-500-100-2-2-4-156

(2496, 500, 100)-difference set in Z_2 x Z_2 x Z_4 x Z_156

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007313 | ljcr-ds-2496-500-100-2-2-624

(2496, 500, 100)-difference set in Z_2 x Z_2 x Z_624

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007314 | ljcr-ds-2496-500-100-2-4-312

(2496, 500, 100)-difference set in Z_2 x Z_4 x Z_312

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007315 | ljcr-ds-2496-500-100-4-4-156

(2496, 500, 100)-difference set in Z_4 x Z_4 x Z_156

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007316 | ljcr-ds-2496-500-100-4-624

(2496, 500, 100)-difference set in Z_4 x Z_624

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007317 | ljcr-ds-2496-500-100-8-312

(2496, 500, 100)-difference set in Z_8 x Z_312

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007318 | ljcr-ds-24961-9985-3994-24961

(24961, 9985, 3994)-difference set in Z_24961

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007319 | ljcr-ds-24997-6249-1562-24997

(24997, 6249, 1562)-difference set in Z_24997

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007320 | ljcr-ds-25011-12505-6252-25011

(25011, 12505, 6252)-difference set in Z_25011

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007321 | ljcr-ds-25011-2501-250-25011

(25011, 2501, 250)-difference set in Z_25011

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007322 | ljcr-ds-25020-9653-3724-25020

(25020, 9653, 3724)-difference set in Z_25020

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007323 | ljcr-ds-2507-896-320-2507

(2507, 896, 320)-difference set in Z_2507

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007324 | ljcr-ds-25071-12535-6267-25071

(25071, 12535, 6267)-difference set in Z_25071

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007325 | ljcr-ds-25081-11001-4825-25081

(25081, 11001, 4825)-difference set in Z_25081

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007326 | ljcr-ds-25081-3136-392-25081

(25081, 3136, 392)-difference set in Z_25081

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007327 | ljcr-ds-25093-7668-2343-25093

(25093, 7668, 2343)-difference set in Z_25093

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007328 | ljcr-ds-25099-12549-6274-25099

(25099, 12549, 6274)-difference set in Z_25099

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007329 | ljcr-ds-2511-1005-402-3-837

(2511, 1005, 402)-difference set in Z_3 x Z_837

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007330 | ljcr-ds-2511-1005-402-9-279

(2511, 1005, 402)-difference set in Z_9 x Z_279

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007331 | ljcr-ds-25135-12567-6283-25135

(25135, 12567, 6283)-difference set in Z_25135

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007332 | ljcr-ds-25135-4189-698-25135

(25135, 4189, 698)-difference set in Z_25135

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007333 | ljcr-ds-25141-420-7-25141

(25141, 420, 7)-difference set in Z_25141

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007334 | ljcr-ds-25159-8386-2795-25159

(25159, 8386, 2795)-difference set in Z_25159

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007335 | ljcr-ds-25189-6297-1574-25189

(25189, 6297, 1574)-difference set in Z_25189

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007336 | ljcr-ds-25194-5369-1144-25194

(25194, 5369, 1144)-difference set in Z_25194

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007337 | ljcr-ds-25221-6305-1576-25221

(25221, 6305, 1576)-difference set in Z_25221

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007338 | ljcr-ds-25311-10125-4050-25311

(25311, 10125, 4050)-difference set in Z_25311

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007339 | ljcr-ds-25312-8437-2812-25312

(25312, 8437, 2812)-difference set in Z_25312

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007340 | ljcr-ds-25323-12661-6330-25323

(25323, 12661, 6330)-difference set in Z_25323

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007341 | ljcr-ds-25327-12663-6331-25327

(25327, 12663, 6331)-difference set in Z_25327

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007342 | ljcr-ds-25351-12000-5680-25351

(25351, 12000, 5680)-difference set in Z_25351

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007343 | ljcr-ds-25376-1625-104-25376

(25376, 1625, 104)-difference set in Z_25376

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007344 | ljcr-ds-25415-10088-4004-25415

(25415, 10088, 4004)-difference set in Z_25415

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007345 | ljcr-ds-25441-6625-1725-25441

(25441, 6625, 1725)-difference set in Z_25441

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007346 | ljcr-ds-25453-6364-1591-25453

(25453, 6364, 1591)-difference set in Z_25453

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007347 | ljcr-ds-2548-567-126-14-182

(2548, 567, 126)-difference set in Z_14 x Z_182

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007348 | ljcr-ds-2548-567-126-7-364

(2548, 567, 126)-difference set in Z_7 x Z_364

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007349 | ljcr-ds-25509-6377-1594-25509

(25509, 6377, 1594)-difference set in Z_25509

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007350 | ljcr-ds-2551-1225-588-2551

(2551, 1225, 588)-difference set in Z_2551

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007351 | ljcr-ds-25521-4785-897-25521

(25521, 4785, 897)-difference set in Z_25521

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007352 | ljcr-ds-25541-6385-1596-25541

(25541, 6385, 1596)-difference set in Z_25541

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007353 | ljcr-ds-25555-8518-2839-25555

(25555, 8518, 2839)-difference set in Z_25555

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007354 | ljcr-ds-25575-4712-868-25575

(25575, 4712, 868)-difference set in Z_25575

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007355 | ljcr-ds-25591-8530-2843-25591

(25591, 8530, 2843)-difference set in Z_25591

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007356 | ljcr-ds-25599-12799-6399-25599

(25599, 12799, 6399)-difference set in Z_25599

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007357 | ljcr-ds-25609-3201-400-25609

(25609, 3201, 400)-difference set in Z_25609

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007358 | ljcr-ds-25615-12807-6403-25615

(25615, 12807, 6403)-difference set in Z_25615

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007359 | ljcr-ds-25671-5135-1027-25671

(25671, 5135, 1027)-difference set in Z_25671

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007360 | ljcr-ds-25675-12837-6418-25675

(25675, 12837, 6418)-difference set in Z_25675

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007361 | ljcr-ds-25711-12855-6427-25711

(25711, 12855, 6427)-difference set in Z_25711

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007362 | ljcr-ds-25729-1408-77-25729

(25729, 1408, 77)-difference set in Z_25729

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007363 | ljcr-ds-25741-6436-1609-25741

(25741, 6436, 1609)-difference set in Z_25741

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007364 | ljcr-ds-25743-12871-6435-25743

(25743, 12871, 6435)-difference set in Z_25743

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007365 | ljcr-ds-2575-495-95-5-515

(2575, 495, 95)-difference set in Z_5 x Z_515

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007366 | ljcr-ds-2575-793-244-5-515

(2575, 793, 244)-difference set in Z_5 x Z_515

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007367 | ljcr-ds-25807-12903-6451-25807

(25807, 12903, 6451)-difference set in Z_25807

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007368 | ljcr-ds-25807-6325-1550-25807

(25807, 6325, 1550)-difference set in Z_25807

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007369 | ljcr-ds-25807-9384-3412-25807

(25807, 9384, 3412)-difference set in Z_25807

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007370 | ljcr-ds-25821-5165-1033-25821

(25821, 5165, 1033)-difference set in Z_25821

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007371 | ljcr-ds-25843-8614-2871-25843

(25843, 8614, 2871)-difference set in Z_25843

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007372 | ljcr-ds-25879-8626-2875-25879

(25879, 8626, 2875)-difference set in Z_25879

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007373 | ljcr-ds-2591-666-171-2591

(2591, 666, 171)-difference set in Z_2591

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007374 | ljcr-ds-25915-4320-720-25915

(25915, 4320, 720)-difference set in Z_25915

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007375 | ljcr-ds-25935-12967-6483-25935

(25935, 12967, 6483)-difference set in Z_25935

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007376 | ljcr-ds-2597-649-162-7-371

(2597, 649, 162)-difference set in Z_7 x Z_371

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007377 | ljcr-ds-25981-5196-1039-25981

(25981, 5196, 1039)-difference set in Z_25981

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007378 | ljcr-ds-25991-10396-4158-25991

(25991, 10396, 4158)-difference set in Z_25991

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007379 | ljcr-ds-25991-12995-6497-25991

(25991, 12995, 6497)-difference set in Z_25991

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007380 | ljcr-ds-26031-13015-6507-26031

(26031, 13015, 6507)-difference set in Z_26031

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007381 | ljcr-ds-26059-2323-207-26059

(26059, 2323, 207)-difference set in Z_26059

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007382 | ljcr-ds-26061-9121-3192-26061

(26061, 9121, 3192)-difference set in Z_26061

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007383 | ljcr-ds-26101-4176-668-26101

(26101, 4176, 668)-difference set in Z_26101

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007384 | ljcr-ds-26131-8710-2903-26131

(26131, 8710, 2903)-difference set in Z_26131

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007385 | ljcr-ds-26133-6533-1633-26133

(26133, 6533, 1633)-difference set in Z_26133

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007386 | ljcr-ds-26145-9633-3549-26145

(26145, 9633, 3549)-difference set in Z_26145

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007387 | ljcr-ds-26191-13095-6547-26191

(26191, 13095, 6547)-difference set in Z_26191

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007388 | ljcr-ds-26197-6549-1637-26197

(26197, 6549, 1637)-difference set in Z_26197

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007389 | ljcr-ds-26239-8746-2915-26239

(26239, 8746, 2915)-difference set in Z_26239

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007390 | ljcr-ds-26265-8576-2800-26265

(26265, 8576, 2800)-difference set in Z_26265

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007391 | ljcr-ds-26283-13141-6570-26283

(26283, 13141, 6570)-difference set in Z_26283

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007392 | ljcr-ds-26287-8425-2700-26287

(26287, 8425, 2700)-difference set in Z_26287

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007393 | ljcr-ds-26291-9560-3476-26291

(26291, 9560, 3476)-difference set in Z_26291

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007394 | ljcr-ds-26309-6577-1644-26309

(26309, 6577, 1644)-difference set in Z_26309

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007395 | ljcr-ds-26312-3487-462-26312

(26312, 3487, 462)-difference set in Z_26312

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007396 | ljcr-ds-26323-13161-6580-26323

(26323, 13161, 6580)-difference set in Z_26323

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007397 | ljcr-ds-26335-2926-325-26335

(26335, 2926, 325)-difference set in Z_26335

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007398 | ljcr-ds-26367-13183-6591-26367

(26367, 13183, 6591)-difference set in Z_26367

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007399 | ljcr-ds-2640-638-154-2-1320

(2640, 638, 154)-difference set in Z_2 x Z_1320

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007400 | ljcr-ds-26404-8125-2500-26404

(26404, 8125, 2500)-difference set in Z_26404

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007401 | ljcr-ds-2641-1200-545-2641

(2641, 1200, 545)-difference set in Z_2641

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007402 | ljcr-ds-26431-5286-1057-26431

(26431, 5286, 1057)-difference set in Z_26431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007403 | ljcr-ds-26437-6609-1652-26437

(26437, 6609, 1652)-difference set in Z_26437

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007404 | ljcr-ds-26445-1804-123-26445

(26445, 1804, 123)-difference set in Z_26445

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007405 | ljcr-ds-2645-661-165-23-115

(2645, 661, 165)-difference set in Z_23 x Z_115

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007406 | ljcr-ds-26467-4411-735-26467

(26467, 4411, 735)-difference set in Z_26467

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007407 | ljcr-ds-26501-6625-1656-26501

(26501, 6625, 1656)-difference set in Z_26501

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007408 | ljcr-ds-26511-13255-6627-26511

(26511, 13255, 6627)-difference set in Z_26511

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007409 | ljcr-ds-26565-8932-3003-26565

(26565, 8932, 3003)-difference set in Z_26565

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007410 | ljcr-ds-26575-13287-6643-26575

(26575, 13287, 6643)-difference set in Z_26575

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007411 | ljcr-ds-26597-6649-1662-26597

(26597, 6649, 1662)-difference set in Z_26597

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007412 | ljcr-ds-26623-13311-6655-26623

(26623, 13311, 6655)-difference set in Z_26623

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007413 | ljcr-ds-26629-7609-2174-26629

(26629, 7609, 2174)-difference set in Z_26629

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007414 | ljcr-ds-26676-4851-882-26676

(26676, 4851, 882)-difference set in Z_26676

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007415 | ljcr-ds-26689-8896-2965-26689

(26689, 8896, 2965)-difference set in Z_26689

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007416 | ljcr-ds-26701-6676-1669-26701

(26701, 6676, 1669)-difference set in Z_26701

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007417 | ljcr-ds-26709-6677-1669-26709

(26709, 6677, 1669)-difference set in Z_26709

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007418 | ljcr-ds-26741-9360-3276-26741

(26741, 9360, 3276)-difference set in Z_26741

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007419 | ljcr-ds-26747-13373-6686-26747

(26747, 13373, 6686)-difference set in Z_26747

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007420 | ljcr-ds-26761-12265-5621-26761

(26761, 12265, 5621)-difference set in Z_26761

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007421 | ljcr-ds-26761-7360-2024-26761

(26761, 7360, 2024)-difference set in Z_26761

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007422 | ljcr-ds-26763-13381-6690-26763

(26763, 13381, 6690)-difference set in Z_26763

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007423 | ljcr-ds-26789-6697-1674-26789

(26789, 6697, 1674)-difference set in Z_26789

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007424 | ljcr-ds-26791-4465-744-26791

(26791, 4465, 744)-difference set in Z_26791

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007425 | ljcr-ds-26796-8855-2926-26796

(26796, 8855, 2926)-difference set in Z_26796

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007426 | ljcr-ds-26797-1596-95-26797

(26797, 1596, 95)-difference set in Z_26797

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007427 | ljcr-ds-26797-8701-2825-26797

(26797, 8701, 2825)-difference set in Z_26797

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007428 | ljcr-ds-26803-13401-6700-26803

(26803, 13401, 6700)-difference set in Z_26803

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007429 | ljcr-ds-26811-2681-268-26811

(26811, 2681, 268)-difference set in Z_26811

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007430 | ljcr-ds-26866-3781-532-26866

(26866, 3781, 532)-difference set in Z_26866

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007431 | ljcr-ds-26885-6721-1680-26885

(26885, 6721, 1680)-difference set in Z_26885

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007432 | ljcr-ds-26895-13447-6723-26895

(26895, 13447, 6723)-difference set in Z_26895

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007433 | ljcr-ds-26909-2884-309-26909

(26909, 2884, 309)-difference set in Z_26909

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007434 | ljcr-ds-26909-3844-549-26909

(26909, 3844, 549)-difference set in Z_26909

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007435 | ljcr-ds-2691-270-27-3-897

(2691, 270, 27)-difference set in Z_3 x Z_897

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007436 | ljcr-ds-26929-8416-2630-26929

(26929, 8416, 2630)-difference set in Z_26929

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007437 | ljcr-ds-26981-6745-1686-26981

(26981, 6745, 1686)-difference set in Z_26981

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007438 | ljcr-ds-2699-1065-420-2699

(2699, 1065, 420)-difference set in Z_2699

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007439 | ljcr-ds-27003-13501-6750-27003

(27003, 13501, 6750)-difference set in Z_27003

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007440 | ljcr-ds-27007-3858-551-27007

(27007, 3858, 551)-difference set in Z_27007

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007441 | ljcr-ds-27031-4506-751-27031

(27031, 4506, 751)-difference set in Z_27031

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007442 | ljcr-ds-27031-9010-3003-27031

(27031, 9010, 3003)-difference set in Z_27031

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007443 | ljcr-ds-27037-12016-5340-27037

(27037, 12016, 5340)-difference set in Z_27037

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007444 | ljcr-ds-27064-5239-1014-27064

(27064, 5239, 1014)-difference set in Z_27064

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007445 | ljcr-ds-27071-13535-6767-27071

(27071, 13535, 6767)-difference set in Z_27071

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007446 | ljcr-ds-27077-6769-1692-27077

(27077, 6769, 1692)-difference set in Z_27077

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007447 | ljcr-ds-27087-13543-6771-27087

(27087, 13543, 6771)-difference set in Z_27087

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007448 | ljcr-ds-2709-677-169-2709

(2709, 677, 169)-difference set in Z_2709

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007449 | ljcr-ds-2709-677-169-3-903

(2709, 677, 169)-difference set in Z_3 x Z_903

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007450 | ljcr-ds-27099-13549-6774-27099

(27099, 13549, 6774)-difference set in Z_27099

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007451 | ljcr-ds-27133-2737-276-27133

(27133, 2737, 276)-difference set in Z_27133

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007452 | ljcr-ds-271467-1379-7-271467

(271467, 1379, 7)-difference set in Z_271467

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007453 | ljcr-ds-27151-13575-6787-27151

(27151, 13575, 6787)-difference set in Z_27151

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007454 | ljcr-ds-27171-13585-6792-27171

(27171, 13585, 6792)-difference set in Z_27171

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007455 | ljcr-ds-27211-9070-3023-27211

(27211, 9070, 3023)-difference set in Z_27211

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007456 | ljcr-ds-27235-8722-2793-27235

(27235, 8722, 2793)-difference set in Z_27235

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007457 | ljcr-ds-27253-6057-1346-27253

(27253, 6057, 1346)-difference set in Z_27253

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007458 | ljcr-ds-27269-6817-1704-27269

(27269, 6817, 1704)-difference set in Z_27269

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007459 | ljcr-ds-27280-3465-440-27280

(27280, 3465, 440)-difference set in Z_27280

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007460 | ljcr-ds-2733-684-171-2733

(2733, 684, 171)-difference set in Z_2733

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007461 | ljcr-ds-27343-13671-6835-27343

(27343, 13671, 6835)-difference set in Z_27343

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007462 | ljcr-ds-27357-11725-5025-27357

(27357, 11725, 5025)-difference set in Z_27357

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007463 | ljcr-ds-2737-1216-540-2737

(2737, 1216, 540)-difference set in Z_2737

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007464 | ljcr-ds-27391-13695-6847-27391

(27391, 13695, 6847)-difference set in Z_27391

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007465 | ljcr-ds-27391-2905-308-27391

(27391, 2905, 308)-difference set in Z_27391

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007466 | ljcr-ds-27395-13697-6848-27395

(27395, 13697, 6848)-difference set in Z_27395

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007467 | ljcr-ds-27405-6852-1713-27405

(27405, 6852, 1713)-difference set in Z_27405

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007468 | ljcr-ds-27495-9087-3003-27495

(27495, 9087, 3003)-difference set in Z_27495

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007469 | ljcr-ds-27511-8646-2717-27511

(27511, 8646, 2717)-difference set in Z_27511

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007470 | ljcr-ds-2759-1183-507-2759

(2759, 1183, 507)-difference set in Z_2759

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007471 | ljcr-ds-27601-11776-5024-27601

(27601, 11776, 5024)-difference set in Z_27601

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007472 | ljcr-ds-27601-2001-145-27601

(27601, 2001, 145)-difference set in Z_27601

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007473 | ljcr-ds-27601-9201-3067-27601

(27601, 9201, 3067)-difference set in Z_27601

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007474 | ljcr-ds-27603-13801-6900-27603

(27603, 13801, 6900)-difference set in Z_27603

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007475 | ljcr-ds-27608-2907-306-27608

(27608, 2907, 306)-difference set in Z_27608

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007476 | ljcr-ds-27627-13813-6906-27627

(27627, 13813, 6906)-difference set in Z_27627

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007477 | ljcr-ds-27657-3457-432-27657

(27657, 3457, 432)-difference set in Z_27657

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007478 | ljcr-ds-27711-13855-6927-27711

(27711, 13855, 6927)-difference set in Z_27711

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007479 | ljcr-ds-27721-12321-5476-27721

(27721, 12321, 5476)-difference set in Z_27721

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007480 | ljcr-ds-27721-3025-330-27721

(27721, 3025, 330)-difference set in Z_27721

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007481 | ljcr-ds-27747-13873-6936-27747

(27747, 13873, 6936)-difference set in Z_27747

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007482 | ljcr-ds-27779-646-15-27779

(27779, 646, 15)-difference set in Z_27779

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007483 | ljcr-ds-27781-6945-1736-27781

(27781, 6945, 1736)-difference set in Z_27781

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007484 | ljcr-ds-27799-9267-3089-27799

(27799, 9267, 3089)-difference set in Z_27799

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007485 | ljcr-ds-27867-13933-6966-27867

(27867, 13933, 6966)-difference set in Z_27867

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007486 | ljcr-ds-27877-6969-1742-27877

(27877, 6969, 1742)-difference set in Z_27877

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007487 | ljcr-ds-27891-13945-6972-27891

(27891, 13945, 6972)-difference set in Z_27891

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007488 | ljcr-ds-27899-13949-6974-27899

(27899, 13949, 6974)-difference set in Z_27899

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007489 | ljcr-ds-27913-9304-3101-27913

(27913, 9304, 3101)-difference set in Z_27913

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007490 | ljcr-ds-27931-931-31-27931

(27931, 931, 31)-difference set in Z_27931

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007491 | ljcr-ds-27949-9316-3105-27949

(27949, 9316, 3105)-difference set in Z_27949

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007492 | ljcr-ds-27949-9453-3197-27949

(27949, 9453, 3197)-difference set in Z_27949

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007493 | ljcr-ds-2795-1144-468-2795

(2795, 1144, 468)-difference set in Z_2795

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007494 | ljcr-ds-27966-9401-3160-27966

(27966, 9401, 3160)-difference set in Z_27966

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007495 | ljcr-ds-2800-622-138-10-280

(2800, 622, 138)-difference set in Z_10 x Z_280

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007496 | ljcr-ds-2800-622-138-2-10-140

(2800, 622, 138)-difference set in Z_2 x Z_10 x Z_140

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007497 | ljcr-ds-2800-622-138-2-1400

(2800, 622, 138)-difference set in Z_2 x Z_1400

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007498 | ljcr-ds-2800-622-138-2-2-10-70

(2800, 622, 138)-difference set in Z_2 x Z_2 x Z_10 x Z_70

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007499 | ljcr-ds-2800-622-138-2-2-2-350

(2800, 622, 138)-difference set in Z_2 x Z_2 x Z_2 x Z_350

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007500 | ljcr-ds-2800-622-138-2-2-700

(2800, 622, 138)-difference set in Z_2 x Z_2 x Z_700

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007501 | ljcr-ds-2800-622-138-20-140

(2800, 622, 138)-difference set in Z_20 x Z_140

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007502 | ljcr-ds-2800-622-138-4-700

(2800, 622, 138)-difference set in Z_4 x Z_700

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007503 | ljcr-ds-28047-14023-7011-28047

(28047, 14023, 7011)-difference set in Z_28047

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007504 | ljcr-ds-28069-7017-1754-28069

(28069, 7017, 1754)-difference set in Z_28069

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007505 | ljcr-ds-28079-12928-5952-28079

(28079, 12928, 5952)-difference set in Z_28079

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007506 | ljcr-ds-28115-14057-7028-28115

(28115, 14057, 7028)-difference set in Z_28115

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007507 | ljcr-ds-28127-3773-506-28127

(28127, 3773, 506)-difference set in Z_28127

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007508 | ljcr-ds-2821-565-113-2821

(2821, 565, 113)-difference set in Z_2821

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007509 | ljcr-ds-28211-2821-282-28211

(28211, 2821, 282)-difference set in Z_28211

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007510 | ljcr-ds-28217-3528-441-28217

(28217, 3528, 441)-difference set in Z_28217

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007511 | ljcr-ds-28231-8470-2541-28231

(28231, 8470, 2541)-difference set in Z_28231

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007512 | ljcr-ds-28241-11296-4518-28241

(28241, 11296, 4518)-difference set in Z_28241

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007513 | ljcr-ds-28245-7061-1765-28245

(28245, 7061, 1765)-difference set in Z_28245

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007514 | ljcr-ds-28263-8697-2676-28263

(28263, 8697, 2676)-difference set in Z_28263

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007515 | ljcr-ds-28321-10561-3938-28321

(28321, 10561, 3938)-difference set in Z_28321

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007516 | ljcr-ds-28337-3312-387-28337

(28337, 3312, 387)-difference set in Z_28337

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007517 | ljcr-ds-2835-546-105-3-3-3-105

(2835, 546, 105)-difference set in Z_3 x Z_3 x Z_3 x Z_105

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007518 | ljcr-ds-2835-546-105-3-3-315

(2835, 546, 105)-difference set in Z_3 x Z_3 x Z_315

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007519 | ljcr-ds-2835-546-105-3-945

(2835, 546, 105)-difference set in Z_3 x Z_945

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007520 | ljcr-ds-2835-546-105-9-315

(2835, 546, 105)-difference set in Z_9 x Z_315

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007521 | ljcr-ds-2835-872-268-3-3-315

(2835, 872, 268)-difference set in Z_3 x Z_3 x Z_315

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007522 | ljcr-ds-2835-872-268-3-945

(2835, 872, 268)-difference set in Z_3 x Z_945

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007523 | ljcr-ds-2835-872-268-9-315

(2835, 872, 268)-difference set in Z_9 x Z_315

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007524 | ljcr-ds-28399-9466-3155-28399

(28399, 9466, 3155)-difference set in Z_28399

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007525 | ljcr-ds-28405-6049-1288-28405

(28405, 6049, 1288)-difference set in Z_28405

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007526 | ljcr-ds-28413-7104-1776-28413

(28413, 7104, 1776)-difference set in Z_28413

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007527 | ljcr-ds-28421-7105-1776-28421

(28421, 7105, 1776)-difference set in Z_28421

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007528 | ljcr-ds-28427-11651-4775-28427

(28427, 11651, 4775)-difference set in Z_28427

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007529 | ljcr-ds-2849-801-225-2849

(2849, 801, 225)-difference set in Z_2849

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007530 | ljcr-ds-2850-925-300-5-570

(2850, 925, 300)-difference set in Z_5 x Z_570

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007531 | ljcr-ds-28517-7129-1782-28517

(28517, 7129, 1782)-difference set in Z_28517

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007532 | ljcr-ds-28533-12229-5241-28533

(28533, 12229, 5241)-difference set in Z_28533

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007533 | ljcr-ds-28539-1502-79-28539

(28539, 1502, 79)-difference set in Z_28539

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007534 | ljcr-ds-28543-1207-51-28543

(28543, 1207, 51)-difference set in Z_28543

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007535 | ljcr-ds-28651-9550-3183-28651

(28651, 9550, 3183)-difference set in Z_28651

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007536 | ljcr-ds-28653-7164-1791-28653

(28653, 7164, 1791)-difference set in Z_28653

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007537 | ljcr-ds-28673-4096-585-28673

(28673, 4096, 585)-difference set in Z_28673

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007538 | ljcr-ds-2869-240-20-2869

(2869, 240, 20)-difference set in Z_2869

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007539 | ljcr-ds-2871-616-132-3-957

(2871, 616, 132)-difference set in Z_3 x Z_957

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007540 | ljcr-ds-2875-480-80-5-5-115

(2875, 480, 80)-difference set in Z_5 x Z_5 x Z_115

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007541 | ljcr-ds-2875-480-80-5-575

(2875, 480, 80)-difference set in Z_5 x Z_575

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007542 | ljcr-ds-28768-6021-1260-28768

(28768, 6021, 1260)-difference set in Z_28768

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007543 | ljcr-ds-28777-11881-4905-28777

(28777, 11881, 4905)-difference set in Z_28777

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007544 | ljcr-ds-28800-930-30-10-2880

(28800, 930, 30)-difference set in Z_10 x Z_2880

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007545 | ljcr-ds-28800-930-30-12-2400

(28800, 930, 30)-difference set in Z_12 x Z_2400

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007546 | ljcr-ds-28800-930-30-120-240

(28800, 930, 30)-difference set in Z_120 x Z_240

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007547 | ljcr-ds-28800-930-30-2-14400

(28800, 930, 30)-difference set in Z_2 x Z_14400

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007548 | ljcr-ds-28800-930-30-20-1440

(28800, 930, 30)-difference set in Z_20 x Z_1440

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007549 | ljcr-ds-28800-930-30-24-1200

(28800, 930, 30)-difference set in Z_24 x Z_1200

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007550 | ljcr-ds-28800-930-30-30-960

(28800, 930, 30)-difference set in Z_30 x Z_960

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007551 | ljcr-ds-28800-930-30-4-7200

(28800, 930, 30)-difference set in Z_4 x Z_7200

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007552 | ljcr-ds-28800-930-30-40-720

(28800, 930, 30)-difference set in Z_40 x Z_720

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007553 | ljcr-ds-28800-930-30-6-4800

(28800, 930, 30)-difference set in Z_6 x Z_4800

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007554 | ljcr-ds-28800-930-30-60-480

(28800, 930, 30)-difference set in Z_60 x Z_480

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007555 | ljcr-ds-28800-930-30-8-3600

(28800, 930, 30)-difference set in Z_8 x Z_3600

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007556 | ljcr-ds-28807-4801-800-28807

(28807, 4801, 800)-difference set in Z_28807

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007557 | ljcr-ds-28813-7204-1801-28813

(28813, 7204, 1801)-difference set in Z_28813

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007558 | ljcr-ds-28821-5765-1153-28821

(28821, 5765, 1153)-difference set in Z_28821

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007559 | ljcr-ds-28837-7209-1802-28837

(28837, 7209, 1802)-difference set in Z_28837

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007560 | ljcr-ds-28841-10816-4056-28841

(28841, 10816, 4056)-difference set in Z_28841

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007561 | ljcr-ds-28911-14455-7227-28911

(28911, 14455, 7227)-difference set in Z_28911

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007562 | ljcr-ds-28923-14461-7230-28923

(28923, 14461, 7230)-difference set in Z_28923

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007563 | ljcr-ds-28947-14473-7236-28947

(28947, 14473, 7236)-difference set in Z_28947

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007564 | ljcr-ds-28971-8691-2607-28971

(28971, 8691, 2607)-difference set in Z_28971

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007565 | ljcr-ds-28975-14487-7243-28975

(28975, 14487, 7243)-difference set in Z_28975

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007566 | ljcr-ds-28992-13675-6450-28992

(28992, 13675, 6450)-difference set in Z_28992

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007567 | ljcr-ds-29023-4837-806-29023

(29023, 4837, 806)-difference set in Z_29023

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007568 | ljcr-ds-29029-12096-5040-29029

(29029, 12096, 5040)-difference set in Z_29029

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007569 | ljcr-ds-29029-7257-1814-29029

(29029, 7257, 1814)-difference set in Z_29029

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007570 | ljcr-ds-29043-1118-43-29043

(29043, 1118, 43)-difference set in Z_29043

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007571 | ljcr-ds-29047-14523-7261-29047

(29047, 14523, 7261)-difference set in Z_29047

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007572 | ljcr-ds-29069-12169-5094-29069

(29069, 12169, 5094)-difference set in Z_29069

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007573 | ljcr-ds-2912-1066-390-2-1456

(2912, 1066, 390)-difference set in Z_2 x Z_1456

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007574 | ljcr-ds-2912-1066-390-4-728

(2912, 1066, 390)-difference set in Z_4 x Z_728

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007575 | ljcr-ds-29140-13245-6020-29140

(29140, 13245, 6020)-difference set in Z_29140

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007576 | ljcr-ds-29179-1621-90-29179

(29179, 1621, 90)-difference set in Z_29179

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007577 | ljcr-ds-29205-7301-1825-29205

(29205, 7301, 1825)-difference set in Z_29205

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007578 | ljcr-ds-29221-7305-1826-29221

(29221, 7305, 1826)-difference set in Z_29221

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007579 | ljcr-ds-29223-13073-5848-29223

(29223, 13073, 5848)-difference set in Z_29223

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007580 | ljcr-ds-29227-14613-7306-29227

(29227, 14613, 7306)-difference set in Z_29227

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007581 | ljcr-ds-29231-8770-2631-29231

(29231, 8770, 2631)-difference set in Z_29231

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007582 | ljcr-ds-2925-732-183-15-195

(2925, 732, 183)-difference set in Z_15 x Z_195

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007583 | ljcr-ds-2925-732-183-3-975

(2925, 732, 183)-difference set in Z_3 x Z_975

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007584 | ljcr-ds-2925-732-183-5-585

(2925, 732, 183)-difference set in Z_5 x Z_585

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007585 | ljcr-ds-29260-13005-5780-29260

(29260, 13005, 5780)-difference set in Z_29260

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007586 | ljcr-ds-29275-2296-180-29275

(29275, 2296, 180)-difference set in Z_29275

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007587 | ljcr-ds-29293-7324-1831-29293

(29293, 7324, 1831)-difference set in Z_29293

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007588 | ljcr-ds-29311-7816-2084-29311

(29311, 7816, 2084)-difference set in Z_29311

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007589 | ljcr-ds-29319-3425-400-29319

(29319, 3425, 400)-difference set in Z_29319

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007590 | ljcr-ds-29331-14665-7332-29331

(29331, 14665, 7332)-difference set in Z_29331

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007591 | ljcr-ds-29331-5866-1173-29331

(29331, 5866, 1173)-difference set in Z_29331

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007592 | ljcr-ds-29355-7904-2128-29355

(29355, 7904, 2128)-difference set in Z_29355

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007593 | ljcr-ds-29381-8476-2445-29381

(29381, 8476, 2445)-difference set in Z_29381

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007594 | ljcr-ds-29389-2449-204-29389

(29389, 2449, 204)-difference set in Z_29389

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007595 | ljcr-ds-29407-14703-7351-29407

(29407, 14703, 7351)-difference set in Z_29407

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007596 | ljcr-ds-29411-2941-294-29411

(29411, 2941, 294)-difference set in Z_29411

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007597 | ljcr-ds-29431-5886-1177-29431

(29431, 5886, 1177)-difference set in Z_29431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007598 | ljcr-ds-29445-7361-1840-29445

(29445, 7361, 1840)-difference set in Z_29445

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007599 | ljcr-ds-29445-8228-2299-29445

(29445, 8228, 2299)-difference set in Z_29445

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007600 | ljcr-ds-29487-14743-7371-29487

(29487, 14743, 7371)-difference set in Z_29487

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007601 | ljcr-ds-29503-1639-91-29503

(29503, 1639, 91)-difference set in Z_29503

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007602 | ljcr-ds-29523-4582-711-29523

(29523, 4582, 711)-difference set in Z_29523

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007603 | ljcr-ds-29546-3421-396-29546

(29546, 3421, 396)-difference set in Z_29546

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007604 | ljcr-ds-29581-7396-1849-29581

(29581, 7396, 1849)-difference set in Z_29581

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007605 | ljcr-ds-29583-14791-7395-29583

(29583, 14791, 7395)-difference set in Z_29583

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007606 | ljcr-ds-29591-12375-5175-29591

(29591, 12375, 5175)-difference set in Z_29591

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007607 | ljcr-ds-29599-4933-822-29599

(29599, 4933, 822)-difference set in Z_29599

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007608 | ljcr-ds-2960-539-98-2-1480

(2960, 539, 98)-difference set in Z_2 x Z_1480

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007609 | ljcr-ds-2960-539-98-2-2-2-370

(2960, 539, 98)-difference set in Z_2 x Z_2 x Z_2 x Z_370

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007610 | ljcr-ds-2960-539-98-2-2-740

(2960, 539, 98)-difference set in Z_2 x Z_2 x Z_740

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007611 | ljcr-ds-2960-539-98-2960

(2960, 539, 98)-difference set in Z_2960

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007612 | ljcr-ds-2960-539-98-4-740

(2960, 539, 98)-difference set in Z_4 x Z_740

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007613 | ljcr-ds-29637-7409-1852-29637

(29637, 7409, 1852)-difference set in Z_29637

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007614 | ljcr-ds-29669-7417-1854-29669

(29669, 7417, 1854)-difference set in Z_29669

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007615 | ljcr-ds-29731-9910-3303-29731

(29731, 9910, 3303)-difference set in Z_29731

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007616 | ljcr-ds-29737-14337-6912-29737

(29737, 14337, 6912)-difference set in Z_29737

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007617 | ljcr-ds-29755-2755-255-29755

(29755, 2755, 255)-difference set in Z_29755

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007618 | ljcr-ds-29756-5951-1190-29756

(29756, 5951, 1190)-difference set in Z_29756

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007619 | ljcr-ds-2976-476-76-2-1488

(2976, 476, 76)-difference set in Z_2 x Z_1488

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007620 | ljcr-ds-2976-476-76-2-2-2-2-186

(2976, 476, 76)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_186

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007621 | ljcr-ds-2976-476-76-2-2-2-372

(2976, 476, 76)-difference set in Z_2 x Z_2 x Z_2 x Z_372

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007622 | ljcr-ds-2976-476-76-2-2-744

(2976, 476, 76)-difference set in Z_2 x Z_2 x Z_744

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007623 | ljcr-ds-2976-476-76-2-4-372

(2976, 476, 76)-difference set in Z_2 x Z_4 x Z_372

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007624 | ljcr-ds-2976-476-76-4-744

(2976, 476, 76)-difference set in Z_4 x Z_744

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007625 | ljcr-ds-29763-14881-7440-29763

(29763, 14881, 7440)-difference set in Z_29763

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007626 | ljcr-ds-29767-12342-5117-29767

(29767, 12342, 5117)-difference set in Z_29767

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007627 | ljcr-ds-29771-14885-7442-29771

(29771, 14885, 7442)-difference set in Z_29771

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007628 | ljcr-ds-29785-11680-4580-29785

(29785, 11680, 4580)-difference set in Z_29785

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007629 | ljcr-ds-29843-14921-7460-29843

(29843, 14921, 7460)-difference set in Z_29843

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007630 | ljcr-ds-29871-14935-7467-29871

(29871, 14935, 7467)-difference set in Z_29871

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007631 | ljcr-ds-2990-245-20-2990

(2990, 245, 20)-difference set in Z_2990

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007632 | ljcr-ds-29951-14975-7487-29951

(29951, 14975, 7487)-difference set in Z_29951

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007633 | ljcr-ds-2997-428-61-2997

(2997, 428, 61)-difference set in Z_2997

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007634 | ljcr-ds-2997-428-61-3-3-3-111

(2997, 428, 61)-difference set in Z_3 x Z_3 x Z_3 x Z_111

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007635 | ljcr-ds-2997-428-61-3-3-333

(2997, 428, 61)-difference set in Z_3 x Z_3 x Z_333

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007636 | ljcr-ds-2997-428-61-3-999

(2997, 428, 61)-difference set in Z_3 x Z_999

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007637 | ljcr-ds-2997-428-61-9-333

(2997, 428, 61)-difference set in Z_9 x Z_333

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007638 | ljcr-ds-29989-7497-1874-29989

(29989, 7497, 1874)-difference set in Z_29989

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007639 | ljcr-ds-30015-5934-1173-30015

(30015, 5934, 1173)-difference set in Z_30015

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007640 | ljcr-ds-3003-475-75-3003

(3003, 475, 75)-difference set in Z_3003

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007641 | ljcr-ds-30041-6760-1521-30041

(30041, 6760, 1521)-difference set in Z_30041

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007642 | ljcr-ds-30063-15031-7515-30063

(30063, 15031, 7515)-difference set in Z_30063

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007643 | ljcr-ds-3008-776-200-2-1504

(3008, 776, 200)-difference set in Z_2 x Z_1504

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007644 | ljcr-ds-3008-776-200-2-2-752

(3008, 776, 200)-difference set in Z_2 x Z_2 x Z_752

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007645 | ljcr-ds-3008-776-200-2-4-376

(3008, 776, 200)-difference set in Z_2 x Z_4 x Z_376

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007646 | ljcr-ds-3008-776-200-4-4-188

(3008, 776, 200)-difference set in Z_4 x Z_4 x Z_188

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007647 | ljcr-ds-3008-776-200-4-752

(3008, 776, 200)-difference set in Z_4 x Z_752

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007648 | ljcr-ds-3008-776-200-8-376

(3008, 776, 200)-difference set in Z_8 x Z_376

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007649 | ljcr-ds-30096-3705-456-30096

(30096, 3705, 456)-difference set in Z_30096

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007650 | ljcr-ds-30109-13897-6414-30109

(30109, 13897, 6414)-difference set in Z_30109

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007651 | ljcr-ds-30143-10766-3845-30143

(30143, 10766, 3845)-difference set in Z_30143

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007652 | ljcr-ds-30175-11985-4760-30175

(30175, 11985, 4760)-difference set in Z_30175

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007653 | ljcr-ds-30181-7545-1886-30181

(30181, 7545, 1886)-difference set in Z_30181

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007654 | ljcr-ds-30195-3410-385-30195

(30195, 3410, 385)-difference set in Z_30195

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007655 | ljcr-ds-30200-7475-1850-30200

(30200, 7475, 1850)-difference set in Z_30200

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007656 | ljcr-ds-30253-2521-210-30253

(30253, 2521, 210)-difference set in Z_30253

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007657 | ljcr-ds-30291-15145-7572-30291

(30291, 15145, 7572)-difference set in Z_30291

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007658 | ljcr-ds-30319-5053-842-30319

(30319, 5053, 842)-difference set in Z_30319

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007659 | ljcr-ds-30321-7201-1710-30321

(30321, 7201, 1710)-difference set in Z_30321

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007660 | ljcr-ds-30341-7585-1896-30341

(30341, 7585, 1896)-difference set in Z_30341

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007661 | ljcr-ds-30371-15185-7592-30371

(30371, 15185, 7592)-difference set in Z_30371

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007662 | ljcr-ds-30381-1520-76-30381

(30381, 1520, 76)-difference set in Z_30381

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007663 | ljcr-ds-30383-15191-7595-30383

(30383, 15191, 7595)-difference set in Z_30383

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007664 | ljcr-ds-30383-2762-251-30383

(30383, 2762, 251)-difference set in Z_30383

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007665 | ljcr-ds-3040-1014-338-2-1520

(3040, 1014, 338)-difference set in Z_2 x Z_1520

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007666 | ljcr-ds-3040-1014-338-4-760

(3040, 1014, 338)-difference set in Z_4 x Z_760

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007667 | ljcr-ds-30411-12165-4866-30411

(30411, 12165, 4866)-difference set in Z_30411

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007668 | ljcr-ds-30415-5070-845-30415

(30415, 5070, 845)-difference set in Z_30415

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007669 | ljcr-ds-30433-10144-3381-30433

(30433, 10144, 3381)-difference set in Z_30433

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007670 | ljcr-ds-3045-761-190-3045

(3045, 761, 190)-difference set in Z_3045

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007671 | ljcr-ds-30501-7625-1906-30501

(30501, 7625, 1906)-difference set in Z_30501

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007672 | ljcr-ds-30511-15255-7627-30511

(30511, 15255, 7627)-difference set in Z_30511

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007673 | ljcr-ds-30535-5089-848-30535

(30535, 5089, 848)-difference set in Z_30535

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007674 | ljcr-ds-30541-10180-3393-30541

(30541, 10180, 3393)-difference set in Z_30541

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007675 | ljcr-ds-30559-10186-3395-30559

(30559, 10186, 3395)-difference set in Z_30559

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007676 | ljcr-ds-30571-15285-7642-30571

(30571, 15285, 7642)-difference set in Z_30571

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007677 | ljcr-ds-30571-6115-1223-30571

(30571, 6115, 1223)-difference set in Z_30571

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007678 | ljcr-ds-30591-13111-5619-30591

(30591, 13111, 5619)-difference set in Z_30591

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007679 | ljcr-ds-30591-4256-592-30591

(30591, 4256, 592)-difference set in Z_30591

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007680 | ljcr-ds-30601-13600-6044-30601

(30601, 13600, 6044)-difference set in Z_30601

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007681 | ljcr-ds-30613-10204-3401-30613

(30613, 10204, 3401)-difference set in Z_30613

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007682 | ljcr-ds-30621-6125-1225-30621

(30621, 6125, 1225)-difference set in Z_30621

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007683 | ljcr-ds-30628-3403-378-30628

(30628, 3403, 378)-difference set in Z_30628

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007684 | ljcr-ds-30629-7657-1914-30629

(30629, 7657, 1914)-difference set in Z_30629

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007685 | ljcr-ds-30661-7665-1916-30661

(30661, 7665, 1916)-difference set in Z_30661

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007686 | ljcr-ds-30669-14433-6792-30669

(30669, 14433, 6792)-difference set in Z_30669

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007687 | ljcr-ds-30681-6136-1227-30681

(30681, 6136, 1227)-difference set in Z_30681

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007688 | ljcr-ds-30681-8320-2256-30681

(30681, 8320, 2256)-difference set in Z_30681

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007689 | ljcr-ds-3069-768-192-3-1023

(3069, 768, 192)-difference set in Z_3 x Z_1023

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007690 | ljcr-ds-30699-15349-7674-30699

(30699, 15349, 7674)-difference set in Z_30699

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007691 | ljcr-ds-30721-4096-546-30721

(30721, 4096, 546)-difference set in Z_30721

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007692 | ljcr-ds-30721-6145-1229-30721

(30721, 6145, 1229)-difference set in Z_30721

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007693 | ljcr-ds-30731-10976-3920-30731

(30731, 10976, 3920)-difference set in Z_30731

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007694 | ljcr-ds-30740-3127-318-30740

(30740, 3127, 318)-difference set in Z_30740

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007695 | ljcr-ds-30757-7689-1922-30757

(30757, 7689, 1922)-difference set in Z_30757

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007696 | ljcr-ds-30767-15383-7691-30767

(30767, 15383, 7691)-difference set in Z_30767

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007697 | ljcr-ds-30784-993-32-4-7696

(30784, 993, 32)-difference set in Z_4 x Z_7696

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007698 | ljcr-ds-30784-993-32-8-3848

(30784, 993, 32)-difference set in Z_8 x Z_3848

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007699 | ljcr-ds-30799-15399-7699-30799

(30799, 15399, 7699)-difference set in Z_30799

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007700 | ljcr-ds-3081-1001-325-3081

(3081, 1001, 325)-difference set in Z_3081

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007701 | ljcr-ds-30821-805-21-30821

(30821, 805, 21)-difference set in Z_30821

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007702 | ljcr-ds-30821-805-21-7-4403

(30821, 805, 21)-difference set in Z_7 x Z_4403

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007703 | ljcr-ds-30857-3249-342-30857

(30857, 3249, 342)-difference set in Z_30857

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007704 | ljcr-ds-30891-15445-7722-30891

(30891, 15445, 7722)-difference set in Z_30891

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007705 | ljcr-ds-30895-15447-7723-30895

(30895, 15447, 7723)-difference set in Z_30895

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007706 | ljcr-ds-30907-5050-825-30907

(30907, 5050, 825)-difference set in Z_30907

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007707 | ljcr-ds-30913-14721-7010-30913

(30913, 14721, 7010)-difference set in Z_30913

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007708 | ljcr-ds-30933-4884-771-30933

(30933, 4884, 771)-difference set in Z_30933

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007709 | ljcr-ds-30961-15265-7526-30961

(30961, 15265, 7526)-difference set in Z_30961

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007710 | ljcr-ds-30997-12096-4720-30997

(30997, 12096, 4720)-difference set in Z_30997

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007711 | ljcr-ds-30997-7749-1937-30997

(30997, 7749, 1937)-difference set in Z_30997

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007712 | ljcr-ds-31003-15501-7750-31003

(31003, 15501, 7750)-difference set in Z_31003

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007713 | ljcr-ds-31021-6205-1241-31021

(31021, 6205, 1241)-difference set in Z_31021

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007714 | ljcr-ds-31031-4495-651-31031

(31031, 4495, 651)-difference set in Z_31031

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007715 | ljcr-ds-31031-4815-747-31031

(31031, 4815, 747)-difference set in Z_31031

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007716 | ljcr-ds-31039-10347-3449-31039

(31039, 10347, 3449)-difference set in Z_31039

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007717 | ljcr-ds-31081-10360-3453-31081

(31081, 10360, 3453)-difference set in Z_31081

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007718 | ljcr-ds-31141-7785-1946-31141

(31141, 7785, 1946)-difference set in Z_31141

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007719 | ljcr-ds-31167-15583-7791-31167

(31167, 15583, 7791)-difference set in Z_31167

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007720 | ljcr-ds-31175-10791-3735-31175

(31175, 10791, 3735)-difference set in Z_31175

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007721 | ljcr-ds-31231-6940-1542-31231

(31231, 6940, 1542)-difference set in Z_31231

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007722 | ljcr-ds-31237-7809-1952-31237

(31237, 7809, 1952)-difference set in Z_31237

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007723 | ljcr-ds-31249-1953-122-31249

(31249, 1953, 122)-difference set in Z_31249

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007724 | ljcr-ds-31261-10420-3473-31261

(31261, 10420, 3473)-difference set in Z_31261

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007725 | ljcr-ds-31301-7825-1956-31301

(31301, 7825, 1956)-difference set in Z_31301

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007726 | ljcr-ds-31311-3535-399-31311

(31311, 3535, 399)-difference set in Z_31311

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007727 | ljcr-ds-3136-210-14-14-224

(3136, 210, 14)-difference set in Z_14 x Z_224

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007728 | ljcr-ds-3136-210-14-2-14-112

(3136, 210, 14)-difference set in Z_2 x Z_14 x Z_112

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007729 | ljcr-ds-3136-210-14-28-112

(3136, 210, 14)-difference set in Z_28 x Z_112

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007730 | ljcr-ds-3136-760-184-14-224

(3136, 760, 184)-difference set in Z_14 x Z_224

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007731 | ljcr-ds-3136-760-184-2-14-112

(3136, 760, 184)-difference set in Z_2 x Z_14 x Z_112

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007732 | ljcr-ds-3136-760-184-28-112

(3136, 760, 184)-difference set in Z_28 x Z_112

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007733 | ljcr-ds-31383-9231-2715-31383

(31383, 9231, 2715)-difference set in Z_31383

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007734 | ljcr-ds-3141-785-196-3-1047

(3141, 785, 196)-difference set in Z_3 x Z_1047

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007735 | ljcr-ds-3141-785-196-3141

(3141, 785, 196)-difference set in Z_3141

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007736 | ljcr-ds-31416-7931-2002-31416

(31416, 7931, 2002)-difference set in Z_31416

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007737 | ljcr-ds-31491-15745-7872-31491

(31491, 15745, 7872)-difference set in Z_31491

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007738 | ljcr-ds-31511-12605-5042-31511

(31511, 12605, 5042)-difference set in Z_31511

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007739 | ljcr-ds-31531-6306-1261-31531

(31531, 6306, 1261)-difference set in Z_31531

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007740 | ljcr-ds-31536-12615-5046-31536

(31536, 12615, 5046)-difference set in Z_31536

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007741 | ljcr-ds-31549-11473-4172-31549

(31549, 11473, 4172)-difference set in Z_31549

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007742 | ljcr-ds-3157-789-197-3157

(3157, 789, 197)-difference set in Z_3157

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007743 | ljcr-ds-31571-6930-1521-31571

(31571, 6930, 1521)-difference set in Z_31571

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007744 | ljcr-ds-31599-15799-7899-31599

(31599, 15799, 7899)-difference set in Z_31599

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007745 | ljcr-ds-31621-10540-3513-31621

(31621, 10540, 3513)-difference set in Z_31621

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007746 | ljcr-ds-31621-3876-475-31621

(31621, 3876, 475)-difference set in Z_31621

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007747 | ljcr-ds-31621-7905-1976-31621

(31621, 7905, 1976)-difference set in Z_31621

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007748 | ljcr-ds-31683-15841-7920-31683

(31683, 15841, 7920)-difference set in Z_31683

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007749 | ljcr-ds-3171-635-127-3171

(3171, 635, 127)-difference set in Z_3171

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007750 | ljcr-ds-31711-11326-4045-31711

(31711, 11326, 4045)-difference set in Z_31711

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007751 | ljcr-ds-31715-15857-7928-31715

(31715, 15857, 7928)-difference set in Z_31715

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007752 | ljcr-ds-31741-13225-5510-31741

(31741, 13225, 5510)-difference set in Z_31741

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007753 | ljcr-ds-31747-1222-47-31747

(31747, 1222, 47)-difference set in Z_31747

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007754 | ljcr-ds-31747-13728-5936-31747

(31747, 13728, 5936)-difference set in Z_31747

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007755 | ljcr-ds-31749-7937-1984-31749

(31749, 7937, 1984)-difference set in Z_31749

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007756 | ljcr-ds-31787-15893-7946-31787

(31787, 15893, 7946)-difference set in Z_31787

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007757 | ljcr-ds-31791-15895-7947-31791

(31791, 15895, 7947)-difference set in Z_31791

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007758 | ljcr-ds-31820-3385-360-31820

(31820, 3385, 360)-difference set in Z_31820

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007759 | ljcr-ds-31837-7581-1805-31837

(31837, 7581, 1805)-difference set in Z_31837

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007760 | ljcr-ds-31971-15985-7992-31971

(31971, 15985, 7992)-difference set in Z_31971

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007761 | ljcr-ds-31977-2856-255-31977

(31977, 2856, 255)-difference set in Z_31977

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007762 | ljcr-ds-31981-7996-1999-31981

(31981, 7996, 1999)-difference set in Z_31981

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007763 | ljcr-ds-32037-8009-2002-32037

(32037, 8009, 2002)-difference set in Z_32037

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007764 | ljcr-ds-32043-16021-8010-32043

(32043, 16021, 8010)-difference set in Z_32043

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007765 | ljcr-ds-32047-16023-8011-32047

(32047, 16023, 8011)-difference set in Z_32047

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007766 | ljcr-ds-32077-2916-265-32077

(32077, 2916, 265)-difference set in Z_32077

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007767 | ljcr-ds-32101-6100-1159-32101

(32101, 6100, 1159)-difference set in Z_32101

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007768 | ljcr-ds-32131-14365-6422-32131

(32131, 14365, 6422)-difference set in Z_32131

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007769 | ljcr-ds-32137-14833-6846-32137

(32137, 14833, 6846)-difference set in Z_32137

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007770 | ljcr-ds-32151-4501-630-32151

(32151, 4501, 630)-difference set in Z_32151

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007771 | ljcr-ds-32175-16087-8043-32175

(32175, 16087, 8043)-difference set in Z_32175

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007772 | ljcr-ds-32179-10726-3575-32179

(32179, 10726, 3575)-difference set in Z_32179

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007773 | ljcr-ds-32209-10065-3145-32209

(32209, 10065, 3145)-difference set in Z_32209

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007774 | ljcr-ds-32275-13366-5535-32275

(32275, 13366, 5535)-difference set in Z_32275

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007775 | ljcr-ds-32281-13720-5831-32281

(32281, 13720, 5831)-difference set in Z_32281

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007776 | ljcr-ds-32311-3231-323-32311

(32311, 3231, 323)-difference set in Z_32311

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007777 | ljcr-ds-32341-10780-3593-32341

(32341, 10780, 3593)-difference set in Z_32341

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007778 | ljcr-ds-32385-10880-3655-32385

(32385, 10880, 3655)-difference set in Z_32385

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007779 | ljcr-ds-32395-5400-900-32395

(32395, 5400, 900)-difference set in Z_32395

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007780 | ljcr-ds-32409-12153-4557-32409

(32409, 12153, 4557)-difference set in Z_32409

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007781 | ljcr-ds-3241-1296-518-3241

(3241, 1296, 518)-difference set in Z_3241

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007782 | ljcr-ds-32411-3241-324-32411

(32411, 3241, 324)-difference set in Z_32411

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007783 | ljcr-ds-32431-16215-8107-32431

(32431, 16215, 8107)-difference set in Z_32431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007784 | ljcr-ds-32445-8112-2028-32445

(32445, 8112, 2028)-difference set in Z_32445

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007785 | ljcr-ds-32449-10816-3605-32449

(32449, 10816, 3605)-difference set in Z_32449

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007786 | ljcr-ds-32453-5796-1035-32453

(32453, 5796, 1035)-difference set in Z_32453

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007787 | ljcr-ds-32473-5248-848-32473

(32473, 5248, 848)-difference set in Z_32473

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007788 | ljcr-ds-32485-8121-2030-32485

(32485, 8121, 2030)-difference set in Z_32485

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007789 | ljcr-ds-32488-1275-50-32488

(32488, 1275, 50)-difference set in Z_32488

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007790 | ljcr-ds-32509-15309-7209-32509

(32509, 15309, 7209)-difference set in Z_32509

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007791 | ljcr-ds-32551-10200-3196-32551

(32551, 10200, 3196)-difference set in Z_32551

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007792 | ljcr-ds-32551-2976-272-32551

(32551, 2976, 272)-difference set in Z_32551

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007793 | ljcr-ds-32571-6515-1303-32571

(32571, 6515, 1303)-difference set in Z_32571

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007794 | ljcr-ds-32640-10795-3570-32640

(32640, 10795, 3570)-difference set in Z_32640

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007795 | ljcr-ds-32641-13056-5222-32641

(32641, 13056, 5222)-difference set in Z_32641

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007796 | ljcr-ds-32641-1921-113-32641

(32641, 1921, 113)-difference set in Z_32641

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007797 | ljcr-ds-32645-8161-2040-32645

(32645, 8161, 2040)-difference set in Z_32645

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007798 | ljcr-ds-32667-16333-8166-32667

(32667, 16333, 8166)-difference set in Z_32667

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007799 | ljcr-ds-32671-6535-1307-32671

(32671, 6535, 1307)-difference set in Z_32671

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007800 | ljcr-ds-32713-4089-511-32713

(32713, 4089, 511)-difference set in Z_32713

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007801 | ljcr-ds-32723-16361-8180-32723

(32723, 16361, 8180)-difference set in Z_32723

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007802 | ljcr-ds-32725-7777-1848-32725

(32725, 7777, 1848)-difference set in Z_32725

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007803 | ljcr-ds-32731-16365-8182-32731

(32731, 16365, 8182)-difference set in Z_32731

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007804 | ljcr-ds-32731-6546-1309-32731

(32731, 6546, 1309)-difference set in Z_32731

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007805 | ljcr-ds-32733-4509-621-32733

(32733, 4509, 621)-difference set in Z_32733

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007806 | ljcr-ds-3276-525-84-2-1638

(3276, 525, 84)-difference set in Z_2 x Z_1638

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007807 | ljcr-ds-3276-525-84-3-1092

(3276, 525, 84)-difference set in Z_3 x Z_1092

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007808 | ljcr-ds-3276-525-84-3276

(3276, 525, 84)-difference set in Z_3276

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007809 | ljcr-ds-3276-525-84-6-546

(3276, 525, 84)-difference set in Z_6 x Z_546

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007810 | ljcr-ds-32760-1599-78-32760

(32760, 1599, 78)-difference set in Z_32760

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007811 | ljcr-ds-32763-16381-8190-32763

(32763, 16381, 8190)-difference set in Z_32763

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007812 | ljcr-ds-32767-2667-217-32767

(32767, 2667, 217)-difference set in Z_32767

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007813 | ljcr-ds-32767-8128-2016-32767

(32767, 8128, 2016)-difference set in Z_32767

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007814 | ljcr-ds-32767-8256-2080-32767

(32767, 8256, 2080)-difference set in Z_32767

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007815 | ljcr-ds-32773-10924-3641-32773

(32773, 10924, 3641)-difference set in Z_32773

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007816 | ljcr-ds-32787-16393-8196-32787

(32787, 16393, 8196)-difference set in Z_32787

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007817 | ljcr-ds-32795-16397-8198-32795

(32795, 16397, 8198)-difference set in Z_32795

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007818 | ljcr-ds-32847-16423-8211-32847

(32847, 16423, 8211)-difference set in Z_32847

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007819 | ljcr-ds-32869-8217-2054-32869

(32869, 8217, 2054)-difference set in Z_32869

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007820 | ljcr-ds-32871-14706-6579-32871

(32871, 14706, 6579)-difference set in Z_32871

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007821 | ljcr-ds-32887-4698-671-32887

(32887, 4698, 671)-difference set in Z_32887

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007822 | ljcr-ds-32890-4617-648-32890

(32890, 4617, 648)-difference set in Z_32890

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007823 | ljcr-ds-32891-12650-4865-32891

(32891, 12650, 4865)-difference set in Z_32891

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007824 | ljcr-ds-32891-14950-6795-32891

(32891, 14950, 6795)-difference set in Z_32891

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007825 | ljcr-ds-32891-3290-329-32891

(32891, 3290, 329)-difference set in Z_32891

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007826 | ljcr-ds-32891-7866-1881-32891

(32891, 7866, 1881)-difference set in Z_32891

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007827 | ljcr-ds-32896-3655-406-32896

(32896, 3655, 406)-difference set in Z_32896

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007828 | ljcr-ds-32923-4248-548-32923

(32923, 4248, 548)-difference set in Z_32923

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007829 | ljcr-ds-32969-2536-195-32969

(32969, 2536, 195)-difference set in Z_32969

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007830 | ljcr-ds-32971-10990-3663-32971

(32971, 10990, 3663)-difference set in Z_32971

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007831 | ljcr-ds-32981-7276-1605-32981

(32981, 7276, 1605)-difference set in Z_32981

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007832 | ljcr-ds-33031-11745-4176-33031

(33031, 11745, 4176)-difference set in Z_33031

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007833 | ljcr-ds-33041-14161-6069-33041

(33041, 14161, 6069)-difference set in Z_33041

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007834 | ljcr-ds-33055-5509-918-33055

(33055, 5509, 918)-difference set in Z_33055

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007835 | ljcr-ds-33079-11026-3675-33079

(33079, 11026, 3675)-difference set in Z_33079

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007836 | ljcr-ds-33089-6017-1094-33089

(33089, 6017, 1094)-difference set in Z_33089

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007837 | ljcr-ds-33099-1274-49-33099

(33099, 1274, 49)-difference set in Z_33099

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007838 | ljcr-ds-33103-10422-3281-33103

(33103, 10422, 3281)-difference set in Z_33103

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007839 | ljcr-ds-33103-16551-8275-33103

(33103, 16551, 8275)-difference set in Z_33103

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007840 | ljcr-ds-33111-2366-169-33111

(33111, 2366, 169)-difference set in Z_33111

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007841 | ljcr-ds-33111-5160-804-33111

(33111, 5160, 804)-difference set in Z_33111

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007842 | ljcr-ds-3312-946-270-12-276

(3312, 946, 270)-difference set in Z_12 x Z_276

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007843 | ljcr-ds-3312-946-270-2-1656

(3312, 946, 270)-difference set in Z_2 x Z_1656

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007844 | ljcr-ds-3312-946-270-4-828

(3312, 946, 270)-difference set in Z_4 x Z_828

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007845 | ljcr-ds-3312-946-270-6-552

(3312, 946, 270)-difference set in Z_6 x Z_552

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007846 | ljcr-ds-33139-14203-6087-33139

(33139, 14203, 6087)-difference set in Z_33139

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007847 | ljcr-ds-33156-16055-7774-33156

(33156, 16055, 7774)-difference set in Z_33156

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007848 | ljcr-ds-33165-8292-2073-33165

(33165, 8292, 2073)-difference set in Z_33165

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007849 | ljcr-ds-33181-6636-1327-33181

(33181, 6636, 1327)-difference set in Z_33181

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007850 | ljcr-ds-33187-16593-8296-33187

(33187, 16593, 8296)-difference set in Z_33187

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007851 | ljcr-ds-33229-4473-602-33229

(33229, 4473, 602)-difference set in Z_33229

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007852 | ljcr-ds-33285-8321-2080-33285

(33285, 8321, 2080)-difference set in Z_33285

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007853 | ljcr-ds-33297-2081-130-33297

(33297, 2081, 130)-difference set in Z_33297

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007854 | ljcr-ds-33299-16649-8324-33299

(33299, 16649, 8324)-difference set in Z_33299

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007855 | ljcr-ds-33306-6661-1332-33306

(33306, 6661, 1332)-difference set in Z_33306

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007856 | ljcr-ds-33313-11104-3701-33313

(33313, 11104, 3701)-difference set in Z_33313

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007857 | ljcr-ds-33317-8329-2082-33317

(33317, 8329, 2082)-difference set in Z_33317

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007858 | ljcr-ds-3333-1225-450-3333

(3333, 1225, 450)-difference set in Z_3333

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007859 | ljcr-ds-3333-833-208-3333

(3333, 833, 208)-difference set in Z_3333

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007860 | ljcr-ds-33331-6666-1333-33331

(33331, 6666, 1333)-difference set in Z_33331

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007861 | ljcr-ds-33333-1924-111-33333

(33333, 1924, 111)-difference set in Z_33333

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007862 | ljcr-ds-33345-3648-399-33345

(33345, 3648, 399)-difference set in Z_33345

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007863 | ljcr-ds-33349-8337-2084-33349

(33349, 8337, 2084)-difference set in Z_33349

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007864 | ljcr-ds-33401-12025-4329-33401

(33401, 12025, 4329)-difference set in Z_33401

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007865 | ljcr-ds-33405-5965-1065-33405

(33405, 5965, 1065)-difference set in Z_33405

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007866 | ljcr-ds-33423-16711-8355-33423

(33423, 16711, 8355)-difference set in Z_33423

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007867 | ljcr-ds-33443-16721-8360-33443

(33443, 16721, 8360)-difference set in Z_33443

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007868 | ljcr-ds-33445-8361-2090-33445

(33445, 8361, 2090)-difference set in Z_33445

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007869 | ljcr-ds-33464-15043-6762-33464

(33464, 15043, 6762)-difference set in Z_33464

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007870 | ljcr-ds-33465-15041-6760-33465

(33465, 15041, 6760)-difference set in Z_33465

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007871 | ljcr-ds-33471-16735-8367-33471

(33471, 16735, 8367)-difference set in Z_33471

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007872 | ljcr-ds-33481-10665-3397-33481

(33481, 10665, 3397)-difference set in Z_33481

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007873 | ljcr-ds-33507-13707-5607-33507

(33507, 13707, 5607)-difference set in Z_33507

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007874 | ljcr-ds-33511-16755-8377-33511

(33511, 16755, 8377)-difference set in Z_33511

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007875 | ljcr-ds-33535-16767-8383-33535

(33535, 16767, 8383)-difference set in Z_33535

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007876 | ljcr-ds-33537-5633-946-33537

(33537, 5633, 946)-difference set in Z_33537

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007877 | ljcr-ds-33579-14833-6552-33579

(33579, 14833, 6552)-difference set in Z_33579

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007878 | ljcr-ds-33583-16791-8395-33583

(33583, 16791, 8395)-difference set in Z_33583

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007879 | ljcr-ds-33619-11206-3735-33619

(33619, 11206, 3735)-difference set in Z_33619

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007880 | ljcr-ds-33643-15309-6966-33643

(33643, 15309, 6966)-difference set in Z_33643

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007881 | ljcr-ds-33649-14721-6440-33649

(33649, 14721, 6440)-difference set in Z_33649

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007882 | ljcr-ds-33691-11230-3743-33691

(33691, 11230, 3743)-difference set in Z_33691

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007883 | ljcr-ds-33699-1827-99-33699

(33699, 1827, 99)-difference set in Z_33699

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007884 | ljcr-ds-33771-16885-8442-33771

(33771, 16885, 8442)-difference set in Z_33771

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007885 | ljcr-ds-33797-8449-2112-33797

(33797, 8449, 2112)-difference set in Z_33797

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007886 | ljcr-ds-33825-1057-33-5-6765

(33825, 1057, 33)-difference set in Z_5 x Z_6765

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007887 | ljcr-ds-33829-8457-2114-33829

(33829, 8457, 2114)-difference set in Z_33829

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007888 | ljcr-ds-33903-16951-8475-33903

(33903, 16951, 8475)-difference set in Z_33903

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007889 | ljcr-ds-33915-7658-1729-33915

(33915, 7658, 1729)-difference set in Z_33915

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007890 | ljcr-ds-33916-14365-6084-33916

(33916, 14365, 6084)-difference set in Z_33916

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007891 | ljcr-ds-33971-6795-1359-33971

(33971, 6795, 1359)-difference set in Z_33971

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007892 | ljcr-ds-33989-8497-2124-33989

(33989, 8497, 2124)-difference set in Z_33989

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007893 | ljcr-ds-34021-8505-2126-34021

(34021, 8505, 2126)-difference set in Z_34021

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007894 | ljcr-ds-34045-14185-5910-34045

(34045, 14185, 5910)-difference set in Z_34045

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007895 | ljcr-ds-34056-7645-1716-34056

(34056, 7645, 1716)-difference set in Z_34056

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007896 | ljcr-ds-34063-17031-8515-34063

(34063, 17031, 8515)-difference set in Z_34063

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007897 | ljcr-ds-34069-11356-3785-34069

(34069, 11356, 3785)-difference set in Z_34069

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007898 | ljcr-ds-34100-3355-330-34100

(34100, 3355, 330)-difference set in Z_34100

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007899 | ljcr-ds-34105-7105-1480-34105

(34105, 7105, 1480)-difference set in Z_34105

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007900 | ljcr-ds-34117-8529-2132-34117

(34117, 8529, 2132)-difference set in Z_34117

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007901 | ljcr-ds-34131-10240-3072-34131

(34131, 10240, 3072)-difference set in Z_34131

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007902 | ljcr-ds-34141-11380-3793-34141

(34141, 11380, 3793)-difference set in Z_34141

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007903 | ljcr-ds-34188-14077-5796-34188

(34188, 14077, 5796)-difference set in Z_34188

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007904 | ljcr-ds-3421-1216-432-3421

(3421, 1216, 432)-difference set in Z_3421

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007905 | ljcr-ds-3421-1521-676-3421

(3421, 1521, 676)-difference set in Z_3421

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007906 | ljcr-ds-34221-15341-6877-34221

(34221, 15341, 6877)-difference set in Z_34221

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007907 | ljcr-ds-34221-6845-1369-34221

(34221, 6845, 1369)-difference set in Z_34221

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007908 | ljcr-ds-34227-17113-8556-34227

(34227, 17113, 8556)-difference set in Z_34227

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007909 | ljcr-ds-34255-14014-5733-34255

(34255, 14014, 5733)-difference set in Z_34255

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007910 | ljcr-ds-34271-13410-5247-34271

(34271, 13410, 5247)-difference set in Z_34271

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007911 | ljcr-ds-34279-5713-952-34279

(34279, 5713, 952)-difference set in Z_34279

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007912 | ljcr-ds-34287-13826-5575-34287

(34287, 13826, 5575)-difference set in Z_34287

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007913 | ljcr-ds-3429-857-214-3-1143

(3429, 857, 214)-difference set in Z_3 x Z_1143

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007914 | ljcr-ds-3429-857-214-3-3-381

(3429, 857, 214)-difference set in Z_3 x Z_3 x Z_381

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007915 | ljcr-ds-343-171-85-7-49

(343, 171, 85)-difference set in Z_7 x Z_49

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007916 | ljcr-ds-34365-13916-5635-34365

(34365, 13916, 5635)-difference set in Z_34365

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007917 | ljcr-ds-3439-1719-859-3439

(3439, 1719, 859)-difference set in Z_3439

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007918 | ljcr-ds-34393-11464-3821-34393

(34393, 11464, 3821)-difference set in Z_34393

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007919 | ljcr-ds-34413-14749-6321-34413

(34413, 14749, 6321)-difference set in Z_34413

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007920 | ljcr-ds-34417-3825-425-34417

(34417, 3825, 425)-difference set in Z_34417

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007921 | ljcr-ds-34441-2625-200-34441

(34441, 2625, 200)-difference set in Z_34441

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007922 | ljcr-ds-34501-8625-2156-34501

(34501, 8625, 2156)-difference set in Z_34501

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007923 | ljcr-ds-34511-11340-3726-34511

(34511, 11340, 3726)-difference set in Z_34511

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007924 | ljcr-ds-34531-12661-4642-34531

(34531, 12661, 4642)-difference set in Z_34531

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007925 | ljcr-ds-34531-5755-959-34531

(34531, 5755, 959)-difference set in Z_34531

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007926 | ljcr-ds-34565-8641-2160-34565

(34565, 8641, 2160)-difference set in Z_34565

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007927 | ljcr-ds-34575-17287-8643-34575

(34575, 17287, 8643)-difference set in Z_34575

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007928 | ljcr-ds-34591-5766-961-34591

(34591, 5766, 961)-difference set in Z_34591

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007929 | ljcr-ds-34595-17297-8648-34595

(34595, 17297, 8648)-difference set in Z_34595

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007930 | ljcr-ds-34615-14743-6279-34615

(34615, 14743, 6279)-difference set in Z_34615

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007931 | ljcr-ds-34623-17311-8655-34623

(34623, 17311, 8655)-difference set in Z_34623

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007932 | ljcr-ds-34681-16185-7553-34681

(34681, 16185, 7553)-difference set in Z_34681

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007933 | ljcr-ds-34681-7225-1505-34681

(34681, 7225, 1505)-difference set in Z_34681

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007934 | ljcr-ds-34684-13663-5382-34684

(34684, 13663, 5382)-difference set in Z_34684

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007935 | ljcr-ds-34711-13351-5135-34711

(34711, 13351, 5135)-difference set in Z_34711

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007936 | ljcr-ds-34711-17355-8677-34711

(34711, 17355, 8677)-difference set in Z_34711

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007937 | ljcr-ds-34715-16337-7688-34715

(34715, 16337, 7688)-difference set in Z_34715

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007938 | ljcr-ds-3473-1457-611-3473

(3473, 1457, 611)-difference set in Z_3473

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007939 | ljcr-ds-3479-518-77-7-497

(3479, 518, 77)-difference set in Z_7 x Z_497

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007940 | ljcr-ds-34811-17405-8702-34811

(34811, 17405, 8702)-difference set in Z_34811

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007941 | ljcr-ds-34813-15472-6876-34813

(34813, 15472, 6876)-difference set in Z_34813

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007942 | ljcr-ds-34813-6769-1316-34813

(34813, 6769, 1316)-difference set in Z_34813

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007943 | ljcr-ds-34816-1056-32-32-1088

(34816, 1056, 32)-difference set in Z_32 x Z_1088

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007944 | ljcr-ds-34821-8705-2176-34821

(34821, 8705, 2176)-difference set in Z_34821

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007945 | ljcr-ds-34861-13945-5578-34861

(34861, 13945, 5578)-difference set in Z_34861

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007946 | ljcr-ds-34863-17431-8715-34863

(34863, 17431, 8715)-difference set in Z_34863

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007947 | ljcr-ds-34901-8725-2181-34901

(34901, 8725, 2181)-difference set in Z_34901

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007948 | ljcr-ds-34949-8737-2184-34949

(34949, 8737, 2184)-difference set in Z_34949

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007949 | ljcr-ds-34987-11662-3887-34987

(34987, 11662, 3887)-difference set in Z_34987

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007950 | ljcr-ds-34989-8748-2187-34989

(34989, 8748, 2187)-difference set in Z_34989

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007951 | ljcr-ds-34991-17495-8747-34991

(34991, 17495, 8747)-difference set in Z_34991

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007952 | ljcr-ds-34999-5833-972-34999

(34999, 5833, 972)-difference set in Z_34999

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007953 | ljcr-ds-35001-15001-6429-35001

(35001, 15001, 6429)-difference set in Z_35001

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007954 | ljcr-ds-35003-12728-4628-35003

(35003, 12728, 4628)-difference set in Z_35003

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007955 | ljcr-ds-3501-876-219-3-1167

(3501, 876, 219)-difference set in Z_3 x Z_1167

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007956 | ljcr-ds-35041-4161-494-35041

(35041, 4161, 494)-difference set in Z_35041

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007957 | ljcr-ds-35101-8100-1869-35101

(35101, 8100, 1869)-difference set in Z_35101

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007958 | ljcr-ds-35113-13377-5096-35113

(35113, 13377, 5096)-difference set in Z_35113

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007959 | ljcr-ds-35149-8788-2197-35149

(35149, 8788, 2197)-difference set in Z_35149

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007960 | ljcr-ds-35151-17575-8787-35151

(35151, 17575, 8787)-difference set in Z_35151

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007961 | ljcr-ds-35156-13351-5070-35156

(35156, 13351, 5070)-difference set in Z_35156

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007962 | ljcr-ds-35156-7031-1406-35156

(35156, 7031, 1406)-difference set in Z_35156

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007963 | ljcr-ds-35161-11721-3907-35161

(35161, 11721, 3907)-difference set in Z_35161

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007964 | ljcr-ds-35181-7036-1407-35181

(35181, 7036, 1407)-difference set in Z_35181

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007965 | ljcr-ds-3520-460-60-2-1760

(3520, 460, 60)-difference set in Z_2 x Z_1760

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007966 | ljcr-ds-3520-460-60-2-2-2-2-2-110

(3520, 460, 60)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_110

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007967 | ljcr-ds-3520-460-60-2-2-2-2-220

(3520, 460, 60)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_220

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007968 | ljcr-ds-3520-460-60-2-2-2-440

(3520, 460, 60)-difference set in Z_2 x Z_2 x Z_2 x Z_440

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007969 | ljcr-ds-3520-460-60-2-2-4-220

(3520, 460, 60)-difference set in Z_2 x Z_2 x Z_4 x Z_220

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007970 | ljcr-ds-3520-460-60-2-2-880

(3520, 460, 60)-difference set in Z_2 x Z_2 x Z_880

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007971 | ljcr-ds-3520-460-60-2-4-440

(3520, 460, 60)-difference set in Z_2 x Z_4 x Z_440

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007972 | ljcr-ds-3520-460-60-4-4-220

(3520, 460, 60)-difference set in Z_4 x Z_4 x Z_220

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007973 | ljcr-ds-3520-460-60-4-880

(3520, 460, 60)-difference set in Z_4 x Z_880

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007974 | ljcr-ds-3520-460-60-8-440

(3520, 460, 60)-difference set in Z_8 x Z_440

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007975 | ljcr-ds-35224-10149-2924-35224

(35224, 10149, 2924)-difference set in Z_35224

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007976 | ljcr-ds-35267-15114-6477-35267

(35267, 15114, 6477)-difference set in Z_35267

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007977 | ljcr-ds-35275-5880-980-35275

(35275, 5880, 980)-difference set in Z_35275

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007978 | ljcr-ds-35295-15127-6483-35295

(35295, 15127, 6483)-difference set in Z_35295

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007979 | ljcr-ds-35299-15688-6972-35299

(35299, 15688, 6972)-difference set in Z_35299

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007980 | ljcr-ds-3535-589-98-3535

(3535, 589, 98)-difference set in Z_3535

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007981 | ljcr-ds-3536-910-234-4-884

(3536, 910, 234)-difference set in Z_4 x Z_884

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007982 | ljcr-ds-35431-7086-1417-35431

(35431, 7086, 1417)-difference set in Z_35431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007983 | ljcr-ds-35449-4432-554-35449

(35449, 4432, 554)-difference set in Z_35449

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007984 | ljcr-ds-35461-8865-2216-35461

(35461, 8865, 2216)-difference set in Z_35461

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007985 | ljcr-ds-35473-11824-3941-35473

(35473, 11824, 3941)-difference set in Z_35473

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007986 | ljcr-ds-35475-17737-8868-35475

(35475, 17737, 8868)-difference set in Z_35475

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007987 | ljcr-ds-35487-12904-4692-35487

(35487, 12904, 4692)-difference set in Z_35487

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007988 | ljcr-ds-35511-8375-1975-35511

(35511, 8375, 1975)-difference set in Z_35511

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007989 | ljcr-ds-35519-14750-6125-35519

(35519, 14750, 6125)-difference set in Z_35519

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007990 | ljcr-ds-35526-4901-676-35526

(35526, 4901, 676)-difference set in Z_35526

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007991 | ljcr-ds-35533-8884-2221-35533

(35533, 8884, 2221)-difference set in Z_35533

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007992 | ljcr-ds-35551-3951-439-35551

(35551, 3951, 439)-difference set in Z_35551

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007993 | ljcr-ds-35588-13111-4830-35588

(35588, 13111, 4830)-difference set in Z_35588

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007994 | ljcr-ds-35589-15253-6537-35589

(35589, 15253, 6537)-difference set in Z_35589

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007995 | ljcr-ds-35631-12216-4188-35631

(35631, 12216, 4188)-difference set in Z_35631

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007996 | ljcr-ds-35671-5946-991-35671

(35671, 5946, 991)-difference set in Z_35671

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007997 | ljcr-ds-35673-16121-7285-35673

(35673, 16121, 7285)-difference set in Z_35673

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007998 | ljcr-ds-35685-8921-2230-35685

(35685, 8921, 2230)-difference set in Z_35685

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-007999 | ljcr-ds-35733-8933-2233-35733

(35733, 8933, 2233)-difference set in Z_35733

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008000 | ljcr-ds-3577-448-56-7-511

(3577, 448, 56)-difference set in Z_7 x Z_511

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008001 | ljcr-ds-35779-11926-3975-35779

(35779, 11926, 3975)-difference set in Z_35779

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008002 | ljcr-ds-35791-17895-8947-35791

(35791, 17895, 8947)-difference set in Z_35791

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008003 | ljcr-ds-35823-17911-8955-35823

(35823, 17911, 8955)-difference set in Z_35823

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008004 | ljcr-ds-35861-8965-2241-35861

(35861, 8965, 2241)-difference set in Z_35861

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008005 | ljcr-ds-35893-12961-4680-35893

(35893, 12961, 4680)-difference set in Z_35893

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008006 | ljcr-ds-3597-900-225-3597

(3597, 900, 225)-difference set in Z_3597

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008007 | ljcr-ds-35971-13080-4756-35971

(35971, 13080, 4756)-difference set in Z_35971

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008008 | ljcr-ds-35981-15421-6609-35981

(35981, 15421, 6609)-difference set in Z_35981

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008009 | ljcr-ds-35987-12312-4212-35987

(35987, 12312, 4212)-difference set in Z_35987

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008010 | ljcr-ds-36001-8001-1778-36001

(36001, 8001, 1778)-difference set in Z_36001

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008011 | ljcr-ds-36019-6670-1235-36019

(36019, 6670, 1235)-difference set in Z_36019

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008012 | ljcr-ds-36023-12866-4595-36023

(36023, 12866, 4595)-difference set in Z_36023

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008013 | ljcr-ds-36027-18013-9006-36027

(36027, 18013, 9006)-difference set in Z_36027

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008014 | ljcr-ds-36036-14415-5766-36036

(36036, 14415, 5766)-difference set in Z_36036

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008015 | ljcr-ds-3605-901-225-3605

(3605, 901, 225)-difference set in Z_3605

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008016 | ljcr-ds-36069-9017-2254-36069

(36069, 9017, 2254)-difference set in Z_36069

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008017 | ljcr-ds-36111-18055-9027-36111

(36111, 18055, 9027)-difference set in Z_36111

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008018 | ljcr-ds-36135-7743-1659-36135

(36135, 7743, 1659)-difference set in Z_36135

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008019 | ljcr-ds-36151-6025-1004-36151

(36151, 6025, 1004)-difference set in Z_36151

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008020 | ljcr-ds-36165-9041-2260-36165

(36165, 9041, 2260)-difference set in Z_36165

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008021 | ljcr-ds-36169-12057-4019-36169

(36169, 12057, 4019)-difference set in Z_36169

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008022 | ljcr-ds-36169-4521-565-36169

(36169, 4521, 565)-difference set in Z_36169

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008023 | ljcr-ds-3619-1675-775-3619

(3619, 1675, 775)-difference set in Z_3619

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008024 | ljcr-ds-36211-8875-2175-36211

(36211, 8875, 2175)-difference set in Z_36211

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008025 | ljcr-ds-36229-9057-2264-36229

(36229, 9057, 2264)-difference set in Z_36229

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008026 | ljcr-ds-36271-13455-4991-36271

(36271, 13455, 4991)-difference set in Z_36271

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008027 | ljcr-ds-36279-1650-75-36279

(36279, 1650, 75)-difference set in Z_36279

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008028 | ljcr-ds-36281-4536-567-36281

(36281, 4536, 567)-difference set in Z_36281

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008029 | ljcr-ds-36297-11168-3436-36297

(36297, 11168, 3436)-difference set in Z_36297

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008030 | ljcr-ds-36301-3025-252-36301

(36301, 3025, 252)-difference set in Z_36301

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008031 | ljcr-ds-36303-18151-9075-36303

(36303, 18151, 9075)-difference set in Z_36303

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008032 | ljcr-ds-36319-12106-4035-36319

(36319, 12106, 4035)-difference set in Z_36319

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008033 | ljcr-ds-36323-18161-9080-36323

(36323, 18161, 9080)-difference set in Z_36323

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008034 | ljcr-ds-36366-16625-7600-36366

(36366, 16625, 7600)-difference set in Z_36366

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008035 | ljcr-ds-36381-7276-1455-36381

(36381, 7276, 1455)-difference set in Z_36381

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008036 | ljcr-ds-36385-7201-1425-36385

(36385, 7201, 1425)-difference set in Z_36385

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008037 | ljcr-ds-36389-9097-2274-36389

(36389, 9097, 2274)-difference set in Z_36389

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008038 | ljcr-ds-36437-9109-2277-36437

(36437, 9109, 2277)-difference set in Z_36437

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008039 | ljcr-ds-36493-9124-2281-36493

(36493, 9124, 2281)-difference set in Z_36493

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008040 | ljcr-ds-36495-18247-9123-36495

(36495, 18247, 9123)-difference set in Z_36495

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008041 | ljcr-ds-36507-18253-9126-36507

(36507, 18253, 9126)-difference set in Z_36507

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008042 | ljcr-ds-36519-6727-1239-36519

(36519, 6727, 1239)-difference set in Z_36519

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008043 | ljcr-ds-36581-9145-2286-36581

(36581, 9145, 2286)-difference set in Z_36581

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008044 | ljcr-ds-36583-5629-866-36583

(36583, 5629, 866)-difference set in Z_36583

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008045 | ljcr-ds-36583-6097-1016-36583

(36583, 6097, 1016)-difference set in Z_36583

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008046 | ljcr-ds-36585-10761-3165-36585

(36585, 10761, 3165)-difference set in Z_36585

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008047 | ljcr-ds-36591-18295-9147-36591

(36591, 18295, 9147)-difference set in Z_36591

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008048 | ljcr-ds-36594-16169-7144-36594

(36594, 16169, 7144)-difference set in Z_36594

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008049 | ljcr-ds-3661-976-260-3661

(3661, 976, 260)-difference set in Z_3661

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008050 | ljcr-ds-36645-9161-2290-36645

(36645, 9161, 2290)-difference set in Z_36645

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008051 | ljcr-ds-36661-12220-4073-36661

(36661, 12220, 4073)-difference set in Z_36661

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008052 | ljcr-ds-36673-192-1-13-2821

(36673, 192, 1)-difference set in Z_13 x Z_2821

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008053 | ljcr-ds-36685-13248-4784-36685

(36685, 13248, 4784)-difference set in Z_36685

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008054 | ljcr-ds-3669-393-42-3669

(3669, 393, 42)-difference set in Z_3669

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008055 | ljcr-ds-36709-9177-2294-36709

(36709, 9177, 2294)-difference set in Z_36709

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008056 | ljcr-ds-36720-4527-558-36720

(36720, 4527, 558)-difference set in Z_36720

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008057 | ljcr-ds-36723-18361-9180-36723

(36723, 18361, 9180)-difference set in Z_36723

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008058 | ljcr-ds-36741-9185-2296-36741

(36741, 9185, 2296)-difference set in Z_36741

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008059 | ljcr-ds-36765-15757-6753-36765

(36765, 15757, 6753)-difference set in Z_36765

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008060 | ljcr-ds-36775-1135-35-36775

(36775, 1135, 35)-difference set in Z_36775

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008061 | ljcr-ds-36801-576-9-3-12267

(36801, 576, 9)-difference set in Z_3 x Z_12267

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008062 | ljcr-ds-36801-576-9-3-3-4089

(36801, 576, 9)-difference set in Z_3 x Z_3 x Z_4089

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008063 | ljcr-ds-36837-9209-2302-36837

(36837, 9209, 2302)-difference set in Z_36837

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008064 | ljcr-ds-36871-12291-4097-36871

(36871, 12291, 4097)-difference set in Z_36871

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008065 | ljcr-ds-36871-7375-1475-36871

(36871, 7375, 1475)-difference set in Z_36871

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008066 | ljcr-ds-36891-15470-6487-36891

(36891, 15470, 6487)-difference set in Z_36891

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008067 | ljcr-ds-36895-12298-4099-36895

(36895, 12298, 4099)-difference set in Z_36895

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008068 | ljcr-ds-36895-14664-5828-36895

(36895, 14664, 5828)-difference set in Z_36895

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008069 | ljcr-ds-36895-9504-2448-36895

(36895, 9504, 2448)-difference set in Z_36895

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008070 | ljcr-ds-36897-1153-36-36897

(36897, 1153, 36)-difference set in Z_36897

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008071 | ljcr-ds-36925-8688-2044-36925

(36925, 8688, 2044)-difference set in Z_36925

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008072 | ljcr-ds-36939-13432-4884-36939

(36939, 13432, 4884)-difference set in Z_36939

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008073 | ljcr-ds-37051-5226-737-37051

(37051, 5226, 737)-difference set in Z_37051

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008074 | ljcr-ds-37056-7411-1482-37056

(37056, 7411, 1482)-difference set in Z_37056

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008075 | ljcr-ds-37061-9265-2316-37061

(37061, 9265, 2316)-difference set in Z_37061

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008076 | ljcr-ds-37081-13905-5214-37081

(37081, 13905, 5214)-difference set in Z_37081

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008077 | ljcr-ds-37081-6696-1209-37081

(37081, 6696, 1209)-difference set in Z_37081

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008078 | ljcr-ds-37087-17661-8410-37087

(37087, 17661, 8410)-difference set in Z_37087

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008079 | ljcr-ds-37103-18551-9275-37103

(37103, 18551, 9275)-difference set in Z_37103

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008080 | ljcr-ds-37107-18553-9276-37107

(37107, 18553, 9276)-difference set in Z_37107

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008081 | ljcr-ds-37115-5544-828-37115

(37115, 5544, 828)-difference set in Z_37115

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008082 | ljcr-ds-37128-12467-4186-37128

(37128, 12467, 4186)-difference set in Z_37128

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008083 | ljcr-ds-37131-18565-9282-37131

(37131, 18565, 9282)-difference set in Z_37131

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008084 | ljcr-ds-37147-18573-9286-37147

(37147, 18573, 9286)-difference set in Z_37147

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008085 | ljcr-ds-37157-9289-2322-37157

(37157, 9289, 2322)-difference set in Z_37157

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008086 | ljcr-ds-37219-12406-4135-37219

(37219, 12406, 4135)-difference set in Z_37219

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008087 | ljcr-ds-37221-9305-2326-37221

(37221, 9305, 2326)-difference set in Z_37221

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008088 | ljcr-ds-37231-6205-1034-37231

(37231, 6205, 1034)-difference set in Z_37231

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008089 | ljcr-ds-37259-12897-4464-37259

(37259, 12897, 4464)-difference set in Z_37259

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008090 | ljcr-ds-37261-11340-3451-37261

(37261, 11340, 3451)-difference set in Z_37261

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008091 | ljcr-ds-37261-12880-4452-37261

(37261, 12880, 4452)-difference set in Z_37261

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008092 | ljcr-ds-37264-12421-4140-37264

(37264, 12421, 4140)-difference set in Z_37264

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008093 | ljcr-ds-37309-12436-4145-37309

(37309, 12436, 4145)-difference set in Z_37309

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008094 | ljcr-ds-37315-3456-320-37315

(37315, 3456, 320)-difference set in Z_37315

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008095 | ljcr-ds-37317-9329-2332-37317

(37317, 9329, 2332)-difference set in Z_37317

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008096 | ljcr-ds-37327-18663-9331-37327

(37327, 18663, 9331)-difference set in Z_37327

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008097 | ljcr-ds-37345-7392-1463-37345

(37345, 7392, 1463)-difference set in Z_37345

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008098 | ljcr-ds-37371-18685-9342-37371

(37371, 18685, 9342)-difference set in Z_37371

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008099 | ljcr-ds-37431-12806-4381-37431

(37431, 12806, 4381)-difference set in Z_37431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008100 | ljcr-ds-37467-8646-1995-37467

(37467, 8646, 1995)-difference set in Z_37467

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008101 | ljcr-ds-37468-9775-2550-37468

(37468, 9775, 2550)-difference set in Z_37468

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008102 | ljcr-ds-37489-12496-4165-37489

(37489, 12496, 4165)-difference set in Z_37489

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008103 | ljcr-ds-3751-625-104-11-341

(3751, 625, 104)-difference set in Z_11 x Z_341

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008104 | ljcr-ds-37531-14040-5252-37531

(37531, 14040, 5252)-difference set in Z_37531

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008105 | ljcr-ds-37541-9385-2346-37541

(37541, 9385, 2346)-difference set in Z_37541

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008106 | ljcr-ds-37549-9388-2347-37549

(37549, 9388, 2347)-difference set in Z_37549

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008107 | ljcr-ds-37551-18775-9387-37551

(37551, 18775, 9387)-difference set in Z_37551

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008108 | ljcr-ds-37583-17158-7833-37583

(37583, 17158, 7833)-difference set in Z_37583

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008109 | ljcr-ds-37631-18815-9407-37631

(37631, 18815, 9407)-difference set in Z_37631

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008110 | ljcr-ds-37635-18817-9408-37635

(37635, 18817, 9408)-difference set in Z_37635

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008111 | ljcr-ds-37637-9409-2352-37637

(37637, 9409, 2352)-difference set in Z_37637

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008112 | ljcr-ds-37669-12556-4185-37669

(37669, 12556, 4185)-difference set in Z_37669

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008113 | ljcr-ds-37669-14965-5945-37669

(37669, 14965, 5945)-difference set in Z_37669

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008114 | ljcr-ds-37669-7009-1304-37669

(37669, 7009, 1304)-difference set in Z_37669

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008115 | ljcr-ds-37711-18855-9427-37711

(37711, 18855, 9427)-difference set in Z_37711

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008116 | ljcr-ds-3775-1480-580-3775

(3775, 1480, 580)-difference set in Z_3775

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008117 | ljcr-ds-37763-18881-9440-37763

(37763, 18881, 9440)-difference set in Z_37763

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008118 | ljcr-ds-37808-6875-1250-37808

(37808, 6875, 1250)-difference set in Z_37808

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008119 | ljcr-ds-37861-9465-2366-37861

(37861, 9465, 2366)-difference set in Z_37861

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008120 | ljcr-ds-37879-6313-1052-37879

(37879, 6313, 1052)-difference set in Z_37879

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008121 | ljcr-ds-37881-2841-213-37881

(37881, 2841, 213)-difference set in Z_37881

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008122 | ljcr-ds-3789-948-237-3-1263

(3789, 948, 237)-difference set in Z_3 x Z_1263

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008123 | ljcr-ds-3789-948-237-3789

(3789, 948, 237)-difference set in Z_3789

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008124 | ljcr-ds-37899-18949-9474-37899

(37899, 18949, 9474)-difference set in Z_37899

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008125 | ljcr-ds-37961-10440-2871-37961

(37961, 10440, 2871)-difference set in Z_37961

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008126 | ljcr-ds-37981-3376-300-37981

(37981, 3376, 300)-difference set in Z_37981

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008127 | ljcr-ds-37983-5426-775-37983

(37983, 5426, 775)-difference set in Z_37983

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008128 | ljcr-ds-38011-12670-4223-38011

(38011, 12670, 4223)-difference set in Z_38011

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008129 | ljcr-ds-38017-9856-2555-38017

(38017, 9856, 2555)-difference set in Z_38017

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008130 | ljcr-ds-38031-11410-3423-38031

(38031, 11410, 3423)-difference set in Z_38031

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008131 | ljcr-ds-38031-19015-9507-38031

(38031, 19015, 9507)-difference set in Z_38031

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008132 | ljcr-ds-3807-693-126-3-3-3-141

(3807, 693, 126)-difference set in Z_3 x Z_3 x Z_3 x Z_141

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008133 | ljcr-ds-3807-693-126-3-3-423

(3807, 693, 126)-difference set in Z_3 x Z_3 x Z_423

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008134 | ljcr-ds-3807-693-126-9-423

(3807, 693, 126)-difference set in Z_9 x Z_423

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008135 | ljcr-ds-38115-1122-33-38115

(38115, 1122, 33)-difference set in Z_38115

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008136 | ljcr-ds-38115-5016-660-38115

(38115, 5016, 660)-difference set in Z_38115

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008137 | ljcr-ds-38117-9529-2382-38117

(38117, 9529, 2382)-difference set in Z_38117

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008138 | ljcr-ds-38121-7625-1525-38121

(38121, 7625, 1525)-difference set in Z_38121

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008139 | ljcr-ds-38130-14665-5640-38130

(38130, 14665, 5640)-difference set in Z_38130

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008140 | ljcr-ds-38181-9545-2386-38181

(38181, 9545, 2386)-difference set in Z_38181

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008141 | ljcr-ds-38185-9288-2259-38185

(38185, 9288, 2259)-difference set in Z_38185

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008142 | ljcr-ds-38187-19093-9546-38187

(38187, 19093, 9546)-difference set in Z_38187

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008143 | ljcr-ds-38191-13870-5037-38191

(38191, 13870, 5037)-difference set in Z_38191

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008144 | ljcr-ds-38191-16416-7056-38191

(38191, 16416, 7056)-difference set in Z_38191

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008145 | ljcr-ds-38191-9045-2142-38191

(38191, 9045, 2142)-difference set in Z_38191

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008146 | ljcr-ds-38192-1267-42-38192

(38192, 1267, 42)-difference set in Z_38192

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008147 | ljcr-ds-38195-19097-9548-38195

(38195, 19097, 9548)-difference set in Z_38195

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008148 | ljcr-ds-38227-6648-1156-38227

(38227, 6648, 1156)-difference set in Z_38227

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008149 | ljcr-ds-3825-240-15-15-255

(3825, 240, 15)-difference set in Z_15 x Z_255

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008150 | ljcr-ds-3825-240-15-3-1275

(3825, 240, 15)-difference set in Z_3 x Z_1275

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008151 | ljcr-ds-3825-240-15-5-765

(3825, 240, 15)-difference set in Z_5 x Z_765

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008152 | ljcr-ds-38331-19165-9582-38331

(38331, 19165, 9582)-difference set in Z_38331

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008153 | ljcr-ds-38411-8350-1815-38411

(38411, 8350, 1815)-difference set in Z_38411

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008154 | ljcr-ds-38415-19207-9603-38415

(38415, 19207, 9603)-difference set in Z_38415

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008155 | ljcr-ds-38451-19225-9612-38451

(38451, 19225, 9612)-difference set in Z_38451

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008156 | ljcr-ds-38473-8016-1670-38473

(38473, 8016, 1670)-difference set in Z_38473

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008157 | ljcr-ds-38501-9625-2406-38501

(38501, 9625, 2406)-difference set in Z_38501

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008158 | ljcr-ds-38505-14440-5415-38505

(38505, 14440, 5415)-difference set in Z_38505

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008159 | ljcr-ds-3851-1001-260-3851

(3851, 1001, 260)-difference set in Z_3851

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008160 | ljcr-ds-38523-19261-9630-38523

(38523, 19261, 9630)-difference set in Z_38523

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008161 | ljcr-ds-38533-9633-2408-38533

(38533, 9633, 2408)-difference set in Z_38533

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008162 | ljcr-ds-38569-12856-4285-38569

(38569, 12856, 4285)-difference set in Z_38569

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008163 | ljcr-ds-38571-7715-1543-38571

(38571, 7715, 1543)-difference set in Z_38571

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008164 | ljcr-ds-38597-9649-2412-38597

(38597, 9649, 2412)-difference set in Z_38597

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008165 | ljcr-ds-38601-6176-988-38601

(38601, 6176, 988)-difference set in Z_38601

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008166 | ljcr-ds-38608-14383-5358-38608

(38608, 14383, 5358)-difference set in Z_38608

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008167 | ljcr-ds-38629-9657-2414-38629

(38629, 9657, 2414)-difference set in Z_38629

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008168 | ljcr-ds-3871-990-253-3871

(3871, 990, 253)-difference set in Z_3871

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008169 | ljcr-ds-3871-990-253-7-553

(3871, 990, 253)-difference set in Z_7 x Z_553

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008170 | ljcr-ds-38760-9605-2380-38760

(38760, 9605, 2380)-difference set in Z_38760

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008171 | ljcr-ds-38803-12934-4311-38803

(38803, 12934, 4311)-difference set in Z_38803

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008172 | ljcr-ds-38817-13344-4587-38817

(38817, 13344, 4587)-difference set in Z_38817

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008173 | ljcr-ds-3887-871-195-13-299

(3887, 871, 195)-difference set in Z_13 x Z_299

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008174 | ljcr-ds-3888-507-66-18-216

(3888, 507, 66)-difference set in Z_18 x Z_216

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008175 | ljcr-ds-3888-507-66-2-18-108

(3888, 507, 66)-difference set in Z_2 x Z_18 x Z_108

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008176 | ljcr-ds-3888-507-66-2-2-18-54

(3888, 507, 66)-difference set in Z_2 x Z_2 x Z_18 x Z_54

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008177 | ljcr-ds-3888-507-66-2-6-18-18

(3888, 507, 66)-difference set in Z_2 x Z_6 x Z_18 x Z_18

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008178 | ljcr-ds-3888-507-66-2-6-6-54

(3888, 507, 66)-difference set in Z_2 x Z_6 x Z_6 x Z_54

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008179 | ljcr-ds-3888-507-66-3-12-108

(3888, 507, 66)-difference set in Z_3 x Z_12 x Z_108

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008180 | ljcr-ds-3888-507-66-3-18-72

(3888, 507, 66)-difference set in Z_3 x Z_18 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008181 | ljcr-ds-3888-507-66-3-3-12-36

(3888, 507, 66)-difference set in Z_3 x Z_3 x Z_12 x Z_36

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008182 | ljcr-ds-3888-507-66-3-3-3-12-12

(3888, 507, 66)-difference set in Z_3 x Z_3 x Z_3 x Z_12 x Z_12

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008183 | ljcr-ds-3888-507-66-3-3-3-144

(3888, 507, 66)-difference set in Z_3 x Z_3 x Z_3 x Z_144

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008184 | ljcr-ds-3888-507-66-3-3-3-3-48

(3888, 507, 66)-difference set in Z_3 x Z_3 x Z_3 x Z_3 x Z_48

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008185 | ljcr-ds-3888-507-66-3-3-3-6-24

(3888, 507, 66)-difference set in Z_3 x Z_3 x Z_3 x Z_6 x Z_24

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008186 | ljcr-ds-3888-507-66-3-3-432

(3888, 507, 66)-difference set in Z_3 x Z_3 x Z_432

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008187 | ljcr-ds-3888-507-66-3-3-6-6-12

(3888, 507, 66)-difference set in Z_3 x Z_3 x Z_6 x Z_6 x Z_12

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008188 | ljcr-ds-3888-507-66-3-3-6-72

(3888, 507, 66)-difference set in Z_3 x Z_3 x Z_6 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008189 | ljcr-ds-3888-507-66-3-36-36

(3888, 507, 66)-difference set in Z_3 x Z_36 x Z_36

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008190 | ljcr-ds-3888-507-66-3-6-216

(3888, 507, 66)-difference set in Z_3 x Z_6 x Z_216

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008191 | ljcr-ds-3888-507-66-3-6-6-36

(3888, 507, 66)-difference set in Z_3 x Z_6 x Z_6 x Z_36

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008192 | ljcr-ds-3888-507-66-3-6-6-6-6

(3888, 507, 66)-difference set in Z_3 x Z_6 x Z_6 x Z_6 x Z_6

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008193 | ljcr-ds-3888-507-66-3-9-144

(3888, 507, 66)-difference set in Z_3 x Z_9 x Z_144

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008194 | ljcr-ds-3888-507-66-36-108

(3888, 507, 66)-difference set in Z_36 x Z_108

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008195 | ljcr-ds-3888-507-66-6-18-36

(3888, 507, 66)-difference set in Z_6 x Z_18 x Z_36

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008196 | ljcr-ds-3888-507-66-6-6-108

(3888, 507, 66)-difference set in Z_6 x Z_6 x Z_108

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008197 | ljcr-ds-3888-507-66-6-6-6-18

(3888, 507, 66)-difference set in Z_6 x Z_6 x Z_6 x Z_18

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008198 | ljcr-ds-3888-507-66-6-648

(3888, 507, 66)-difference set in Z_6 x Z_648

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008199 | ljcr-ds-3888-507-66-9-432

(3888, 507, 66)-difference set in Z_9 x Z_432

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008200 | ljcr-ds-38893-9261-2205-38893

(38893, 9261, 2205)-difference set in Z_38893

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008201 | ljcr-ds-38911-19455-9727-38911

(38911, 19455, 9727)-difference set in Z_38911

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008202 | ljcr-ds-38917-9729-2432-38917

(38917, 9729, 2432)-difference set in Z_38917

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008203 | ljcr-ds-38920-12973-4324-38920

(38920, 12973, 4324)-difference set in Z_38920

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008204 | ljcr-ds-38931-19465-9732-38931

(38931, 19465, 9732)-difference set in Z_38931

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008205 | ljcr-ds-38991-19495-9747-38991

(38991, 19495, 9747)-difference set in Z_38991

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008206 | ljcr-ds-39045-9761-2440-39045

(39045, 9761, 2440)-difference set in Z_39045

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008207 | ljcr-ds-39051-19525-9762-39051

(39051, 19525, 9762)-difference set in Z_39051

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008208 | ljcr-ds-39051-2201-124-39051

(39051, 2201, 124)-difference set in Z_39051

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008209 | ljcr-ds-39055-19527-9763-39055

(39055, 19527, 9763)-difference set in Z_39055

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008210 | ljcr-ds-3906-781-156-3-1302

(3906, 781, 156)-difference set in Z_3 x Z_1302

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008211 | ljcr-ds-39109-9777-2444-39109

(39109, 9777, 2444)-difference set in Z_39109

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008212 | ljcr-ds-3911-460-54-3911

(3911, 460, 54)-difference set in Z_3911

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008213 | ljcr-ds-3913-489-61-3913

(3913, 489, 61)-difference set in Z_3913

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008214 | ljcr-ds-39147-19573-9786-39147

(39147, 19573, 9786)-difference set in Z_39147

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008215 | ljcr-ds-39181-9796-2449-39181

(39181, 9796, 2449)-difference set in Z_39181

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008216 | ljcr-ds-39215-16807-7203-39215

(39215, 16807, 7203)-difference set in Z_39215

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008217 | ljcr-ds-39267-19633-9816-39267

(39267, 19633, 9816)-difference set in Z_39267

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008218 | ljcr-ds-39269-9817-2454-39269

(39269, 9817, 2454)-difference set in Z_39269

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008219 | ljcr-ds-39275-19637-9818-39275

(39275, 19637, 9818)-difference set in Z_39275

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008220 | ljcr-ds-39292-18711-8910-39292

(39292, 18711, 8910)-difference set in Z_39292

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008221 | ljcr-ds-39295-4366-485-39295

(39295, 4366, 485)-difference set in Z_39295

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008222 | ljcr-ds-39301-9825-2456-39301

(39301, 9825, 2456)-difference set in Z_39301

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008223 | ljcr-ds-39312-18621-8820-39312

(39312, 18621, 8820)-difference set in Z_39312

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008224 | ljcr-ds-39325-13108-4369-39325

(39325, 13108, 4369)-difference set in Z_39325

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008225 | ljcr-ds-39361-13120-4373-39361

(39361, 13120, 4373)-difference set in Z_39361

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008226 | ljcr-ds-3937-1600-650-3937

(3937, 1600, 650)-difference set in Z_3937

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008227 | ljcr-ds-39411-15765-6306-39411

(39411, 15765, 6306)-difference set in Z_39411

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008228 | ljcr-ds-39412-18271-8470-39412

(39412, 18271, 8470)-difference set in Z_39412

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008229 | ljcr-ds-39429-9857-2464-39429

(39429, 9857, 2464)-difference set in Z_39429

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008230 | ljcr-ds-39431-11830-3549-39431

(39431, 11830, 3549)-difference set in Z_39431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008231 | ljcr-ds-39471-19735-9867-39471

(39471, 19735, 9867)-difference set in Z_39471

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008232 | ljcr-ds-39501-11376-3276-39501

(39501, 11376, 3276)-difference set in Z_39501

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008233 | ljcr-ds-39501-9876-2469-39501

(39501, 9876, 2469)-difference set in Z_39501

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008234 | ljcr-ds-3951-1501-570-3951

(3951, 1501, 570)-difference set in Z_3951

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008235 | ljcr-ds-39525-11809-3528-39525

(39525, 11809, 3528)-difference set in Z_39525

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008236 | ljcr-ds-39560-17901-8100-39560

(39560, 17901, 8100)-difference set in Z_39560

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008237 | ljcr-ds-39571-7915-1583-39571

(39571, 7915, 1583)-difference set in Z_39571

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008238 | ljcr-ds-39577-12513-3956-39577

(39577, 12513, 3956)-difference set in Z_39577

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008239 | ljcr-ds-39583-19791-9895-39583

(39583, 19791, 9895)-difference set in Z_39583

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008240 | ljcr-ds-39585-3712-348-39585

(39585, 3712, 348)-difference set in Z_39585

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008241 | ljcr-ds-39589-3300-275-39589

(39589, 3300, 275)-difference set in Z_39589

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008242 | ljcr-ds-39603-19801-9900-39603

(39603, 19801, 9900)-difference set in Z_39603

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008243 | ljcr-ds-39613-13204-4401-39613

(39613, 13204, 4401)-difference set in Z_39613

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008244 | ljcr-ds-39649-5664-809-39649

(39649, 5664, 809)-difference set in Z_39649

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008245 | ljcr-ds-39691-19845-9922-39691

(39691, 19845, 9922)-difference set in Z_39691

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008246 | ljcr-ds-39711-1805-82-39711

(39711, 1805, 82)-difference set in Z_39711

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008247 | ljcr-ds-39753-4969-621-39753

(39753, 4969, 621)-difference set in Z_39753

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008248 | ljcr-ds-39771-19885-9942-39771

(39771, 19885, 9942)-difference set in Z_39771

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008249 | ljcr-ds-39776-17501-7700-39776

(39776, 17501, 7700)-difference set in Z_39776

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008250 | ljcr-ds-39805-2140-115-39805

(39805, 2140, 115)-difference set in Z_39805

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008251 | ljcr-ds-39877-9969-2492-39877

(39877, 9969, 2492)-difference set in Z_39877

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008252 | ljcr-ds-39901-13300-4433-39901

(39901, 13300, 4433)-difference set in Z_39901

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008253 | ljcr-ds-3991-1330-443-3991

(3991, 1330, 443)-difference set in Z_3991

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008254 | ljcr-ds-39919-19959-9979-39919

(39919, 19959, 9979)-difference set in Z_39919

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008255 | ljcr-ds-39919-6654-1109-39919

(39919, 6654, 1109)-difference set in Z_39919

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008256 | ljcr-ds-39941-9985-2496-39941

(39941, 9985, 2496)-difference set in Z_39941

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008257 | ljcr-ds-39951-19176-9204-39951

(39951, 19176, 9204)-difference set in Z_39951

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008258 | ljcr-ds-39953-2497-156-39953

(39953, 2497, 156)-difference set in Z_39953

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008259 | ljcr-ds-39977-16569-6867-39977

(39977, 16569, 6867)-difference set in Z_39977

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008260 | ljcr-ds-39999-19999-9999-39999

(39999, 19999, 9999)-difference set in Z_39999

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008261 | ljcr-ds-400-190-90-10-40

(400, 190, 90)-difference set in Z_10 x Z_40

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008262 | ljcr-ds-400-190-90-2-10-20

(400, 190, 90)-difference set in Z_2 x Z_10 x Z_20

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008263 | ljcr-ds-400-190-90-20-20

(400, 190, 90)-difference set in Z_20 x Z_20

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008264 | ljcr-ds-4000-775-150-10-400

(4000, 775, 150)-difference set in Z_10 x Z_400

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008265 | ljcr-ds-4000-775-150-2-10-200

(4000, 775, 150)-difference set in Z_2 x Z_10 x Z_200

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008266 | ljcr-ds-4000-775-150-2-20-100

(4000, 775, 150)-difference set in Z_2 x Z_20 x Z_100

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008267 | ljcr-ds-4000-775-150-20-200

(4000, 775, 150)-difference set in Z_20 x Z_200

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008268 | ljcr-ds-40007-7471-1395-40007

(40007, 7471, 1395)-difference set in Z_40007

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008269 | ljcr-ds-40009-5001-625-40009

(40009, 5001, 625)-difference set in Z_40009

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008270 | ljcr-ds-40021-10005-2501-40021

(40021, 10005, 2501)-difference set in Z_40021

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008271 | ljcr-ds-40021-18676-8715-40021

(40021, 18676, 8715)-difference set in Z_40021

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008272 | ljcr-ds-40021-5221-681-40021

(40021, 5221, 681)-difference set in Z_40021

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008273 | ljcr-ds-40027-7624-1452-40027

(40027, 7624, 1452)-difference set in Z_40027

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008274 | ljcr-ds-40033-13761-4730-40033

(40033, 13761, 4730)-difference set in Z_40033

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008275 | ljcr-ds-40037-10009-2502-40037

(40037, 10009, 2502)-difference set in Z_40037

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008276 | ljcr-ds-4005-1001-250-3-1335

(4005, 1001, 250)-difference set in Z_3 x Z_1335

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008277 | ljcr-ds-40051-11125-3090-40051

(40051, 11125, 3090)-difference set in Z_40051

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008278 | ljcr-ds-40051-11926-3551-40051

(40051, 11926, 3551)-difference set in Z_40051

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008279 | ljcr-ds-40051-801-16-11-3641

(40051, 801, 16)-difference set in Z_11 x Z_3641

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008280 | ljcr-ds-40051-801-16-40051

(40051, 801, 16)-difference set in Z_40051

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008281 | ljcr-ds-40111-13371-4457-40111

(40111, 13371, 4457)-difference set in Z_40111

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008282 | ljcr-ds-40135-13378-4459-40135

(40135, 13378, 4459)-difference set in Z_40135

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008283 | ljcr-ds-40136-10121-2552-40136

(40136, 10121, 2552)-difference set in Z_40136

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008284 | ljcr-ds-40139-14336-5120-40139

(40139, 14336, 5120)-difference set in Z_40139

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008285 | ljcr-ds-40153-15057-5646-40153

(40153, 15057, 5646)-difference set in Z_40153

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008286 | ljcr-ds-40205-16929-7128-40205

(40205, 16929, 7128)-difference set in Z_40205

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008287 | ljcr-ds-40223-11662-3381-40223

(40223, 11662, 3381)-difference set in Z_40223

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008288 | ljcr-ds-4025-504-63-4025

(4025, 504, 63)-difference set in Z_4025

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008289 | ljcr-ds-4025-504-63-5-805

(4025, 504, 63)-difference set in Z_5 x Z_805

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008290 | ljcr-ds-40261-10065-2516-40261

(40261, 10065, 2516)-difference set in Z_40261

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008291 | ljcr-ds-40267-20133-10066-40267

(40267, 20133, 10066)-difference set in Z_40267

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008292 | ljcr-ds-40271-8055-1611-40271

(40271, 8055, 1611)-difference set in Z_40271

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008293 | ljcr-ds-40291-6715-1119-40291

(40291, 6715, 1119)-difference set in Z_40291

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008294 | ljcr-ds-4032-696-120-12-336

(4032, 696, 120)-difference set in Z_12 x Z_336

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008295 | ljcr-ds-4032-696-120-2-12-168

(4032, 696, 120)-difference set in Z_2 x Z_12 x Z_168

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008296 | ljcr-ds-4032-696-120-2-2-1008

(4032, 696, 120)-difference set in Z_2 x Z_2 x Z_1008

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008297 | ljcr-ds-4032-696-120-2-2-2-504

(4032, 696, 120)-difference set in Z_2 x Z_2 x Z_2 x Z_504

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008298 | ljcr-ds-4032-696-120-2-2-6-168

(4032, 696, 120)-difference set in Z_2 x Z_2 x Z_6 x Z_168

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008299 | ljcr-ds-4032-696-120-2-2016

(4032, 696, 120)-difference set in Z_2 x Z_2016

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008300 | ljcr-ds-4032-696-120-2-4-504

(4032, 696, 120)-difference set in Z_2 x Z_4 x Z_504

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008301 | ljcr-ds-4032-696-120-2-6-336

(4032, 696, 120)-difference set in Z_2 x Z_6 x Z_336

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008302 | ljcr-ds-4032-696-120-24-168

(4032, 696, 120)-difference set in Z_24 x Z_168

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008303 | ljcr-ds-4032-696-120-4-1008

(4032, 696, 120)-difference set in Z_4 x Z_1008

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008304 | ljcr-ds-4032-696-120-6-672

(4032, 696, 120)-difference set in Z_6 x Z_672

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008305 | ljcr-ds-4032-696-120-8-504

(4032, 696, 120)-difference set in Z_8 x Z_504

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008306 | ljcr-ds-40381-10096-2524-40381

(40381, 10096, 2524)-difference set in Z_40381

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008307 | ljcr-ds-40399-20199-10099-40399

(40399, 20199, 10099)-difference set in Z_40399

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008308 | ljcr-ds-40411-4041-404-40411

(40411, 4041, 404)-difference set in Z_40411

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008309 | ljcr-ds-40443-9418-2193-40443

(40443, 9418, 2193)-difference set in Z_40443

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008310 | ljcr-ds-40470-13585-4560-40470

(40470, 13585, 4560)-difference set in Z_40470

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008311 | ljcr-ds-40489-11809-3444-40489

(40489, 11809, 3444)-difference set in Z_40489

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008312 | ljcr-ds-40489-7953-1562-40489

(40489, 7953, 1562)-difference set in Z_40489

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008313 | ljcr-ds-40495-20247-10123-40495

(40495, 20247, 10123)-difference set in Z_40495

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008314 | ljcr-ds-40531-8106-1621-40531

(40531, 8106, 1621)-difference set in Z_40531

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008315 | ljcr-ds-40533-10133-2533-40533

(40533, 10133, 2533)-difference set in Z_40533

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008316 | ljcr-ds-40563-20281-10140-40563

(40563, 20281, 10140)-difference set in Z_40563

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008317 | ljcr-ds-4061-1856-848-4061

(4061, 1856, 848)-difference set in Z_4061

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008318 | ljcr-ds-40612-13537-4512-40612

(40612, 13537, 4512)-difference set in Z_40612

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008319 | ljcr-ds-40645-10161-2540-40645

(40645, 10161, 2540)-difference set in Z_40645

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008320 | ljcr-ds-40671-15190-5673-40671

(40671, 15190, 5673)-difference set in Z_40671

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008321 | ljcr-ds-40671-8135-1627-40671

(40671, 8135, 1627)-difference set in Z_40671

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008322 | ljcr-ds-40687-20343-10171-40687

(40687, 20343, 10171)-difference set in Z_40687

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008323 | ljcr-ds-40693-13564-4521-40693

(40693, 13564, 4521)-difference set in Z_40693

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008324 | ljcr-ds-40755-13490-4465-40755

(40755, 13490, 4465)-difference set in Z_40755

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008325 | ljcr-ds-40767-2398-141-40767

(40767, 2398, 141)-difference set in Z_40767

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008326 | ljcr-ds-40831-12250-3675-40831

(40831, 12250, 3675)-difference set in Z_40831

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008327 | ljcr-ds-40847-9426-2175-40847

(40847, 9426, 2175)-difference set in Z_40847

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008328 | ljcr-ds-40849-17872-7819-40849

(40849, 17872, 7819)-difference set in Z_40849

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008329 | ljcr-ds-40911-20455-10227-40911

(40911, 20455, 10227)-difference set in Z_40911

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008330 | ljcr-ds-40921-13641-4547-40921

(40921, 13641, 4547)-difference set in Z_40921

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008331 | ljcr-ds-40921-16896-6976-40921

(40921, 16896, 6976)-difference set in Z_40921

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008332 | ljcr-ds-40921-8185-1637-40921

(40921, 8185, 1637)-difference set in Z_40921

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008333 | ljcr-ds-4095-1335-435-4095

(4095, 1335, 435)-difference set in Z_4095

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008334 | ljcr-ds-40964-3289-264-40964

(40964, 3289, 264)-difference set in Z_40964

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008335 | ljcr-ds-40965-10241-2560-40965

(40965, 10241, 2560)-difference set in Z_40965

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008336 | ljcr-ds-41005-18225-8100-41005

(41005, 18225, 8100)-difference set in Z_41005

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008337 | ljcr-ds-41031-11936-3472-41031

(41031, 11936, 3472)-difference set in Z_41031

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008338 | ljcr-ds-41047-13683-4561-41047

(41047, 13683, 4561)-difference set in Z_41047

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008339 | ljcr-ds-41047-6841-1140-41047

(41047, 6841, 1140)-difference set in Z_41047

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008340 | ljcr-ds-41053-3421-285-41053

(41053, 3421, 285)-difference set in Z_41053

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008341 | ljcr-ds-41071-13690-4563-41071

(41071, 13690, 4563)-difference set in Z_41071

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008342 | ljcr-ds-41071-20535-10267-41071

(41071, 20535, 10267)-difference set in Z_41071

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008343 | ljcr-ds-41073-19329-9096-41073

(41073, 19329, 9096)-difference set in Z_41073

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008344 | ljcr-ds-41103-20551-10275-41103

(41103, 20551, 10275)-difference set in Z_41103

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008345 | ljcr-ds-4113-257-16-3-1371

(4113, 257, 16)-difference set in Z_3 x Z_1371

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008346 | ljcr-ds-4113-257-16-4113

(4113, 257, 16)-difference set in Z_4113

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008347 | ljcr-ds-41161-5145-643-41161

(41161, 5145, 643)-difference set in Z_41161

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008348 | ljcr-ds-41182-13357-4332-41182

(41182, 13357, 4332)-difference set in Z_41182

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008349 | ljcr-ds-41189-10297-2574-41189

(41189, 10297, 2574)-difference set in Z_41189

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008350 | ljcr-ds-41211-4121-412-41211

(41211, 4121, 412)-difference set in Z_41211

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008351 | ljcr-ds-41263-8464-1736-41263

(41263, 8464, 1736)-difference set in Z_41263

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008352 | ljcr-ds-41275-11466-3185-41275

(41275, 11466, 3185)-difference set in Z_41275

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008353 | ljcr-ds-41301-10325-2581-41301

(41301, 10325, 2581)-difference set in Z_41301

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008354 | ljcr-ds-41375-7399-1323-41375

(41375, 7399, 1323)-difference set in Z_41375

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008355 | ljcr-ds-41381-10345-2586-41381

(41381, 10345, 2586)-difference set in Z_41381

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008356 | ljcr-ds-41407-20703-10351-41407

(41407, 20703, 10351)-difference set in Z_41407

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008357 | ljcr-ds-41419-16875-6875-41419

(41419, 16875, 6875)-difference set in Z_41419

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008358 | ljcr-ds-41420-2135-110-41420

(41420, 2135, 110)-difference set in Z_41420

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008359 | ljcr-ds-41421-8285-1657-41421

(41421, 8285, 1657)-difference set in Z_41421

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008360 | ljcr-ds-41427-20713-10356-41427

(41427, 20713, 10356)-difference set in Z_41427

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008361 | ljcr-ds-4145-1296-405-4145

(4145, 1296, 405)-difference set in Z_4145

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008362 | ljcr-ds-41464-4608-512-41464

(41464, 4608, 512)-difference set in Z_41464

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008363 | ljcr-ds-41481-13176-4185-41481

(41481, 13176, 4185)-difference set in Z_41481

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008364 | ljcr-ds-41487-20743-10371-41487

(41487, 20743, 10371)-difference set in Z_41487

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008365 | ljcr-ds-41528-15851-6050-41528

(41528, 15851, 6050)-difference set in Z_41528

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008366 | ljcr-ds-41551-20775-10387-41551

(41551, 20775, 10387)-difference set in Z_41551

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008367 | ljcr-ds-41557-10389-2597-41557

(41557, 10389, 2597)-difference set in Z_41557

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008368 | ljcr-ds-41581-1540-57-41581

(41581, 1540, 57)-difference set in Z_41581

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008369 | ljcr-ds-416-166-66-2-2-104

(416, 166, 66)-difference set in Z_2 x Z_2 x Z_104

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008370 | ljcr-ds-416-166-66-2-208

(416, 166, 66)-difference set in Z_2 x Z_208

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008371 | ljcr-ds-416-166-66-4-104

(416, 166, 66)-difference set in Z_4 x Z_104

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008372 | ljcr-ds-41705-12832-3948-41705

(41705, 12832, 3948)-difference set in Z_41705

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008373 | ljcr-ds-41751-20375-9943-41751

(41751, 20375, 9943)-difference set in Z_41751

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008374 | ljcr-ds-4176-501-60-12-348

(4176, 501, 60)-difference set in Z_12 x Z_348

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008375 | ljcr-ds-4176-501-60-2-2-1044

(4176, 501, 60)-difference set in Z_2 x Z_2 x Z_1044

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008376 | ljcr-ds-4176-501-60-2-2088

(4176, 501, 60)-difference set in Z_2 x Z_2088

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008377 | ljcr-ds-4176-501-60-2-6-348

(4176, 501, 60)-difference set in Z_2 x Z_6 x Z_348

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008378 | ljcr-ds-4176-501-60-3-1392

(4176, 501, 60)-difference set in Z_3 x Z_1392

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008379 | ljcr-ds-4176-501-60-4-1044

(4176, 501, 60)-difference set in Z_4 x Z_1044

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008380 | ljcr-ds-4176-501-60-4176

(4176, 501, 60)-difference set in Z_4176

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008381 | ljcr-ds-4176-501-60-6-696

(4176, 501, 60)-difference set in Z_6 x Z_696

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008382 | ljcr-ds-41831-15041-5408-41831

(41831, 15041, 5408)-difference set in Z_41831

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008383 | ljcr-ds-41867-20933-10466-41867

(41867, 20933, 10466)-difference set in Z_41867

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008384 | ljcr-ds-41899-20949-10474-41899

(41899, 20949, 10474)-difference set in Z_41899

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008385 | ljcr-ds-41957-10489-2622-41957

(41957, 10489, 2622)-difference set in Z_41957

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008386 | ljcr-ds-41967-20983-10491-41967

(41967, 20983, 10491)-difference set in Z_41967

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008387 | ljcr-ds-41976-5475-714-41976

(41976, 5475, 714)-difference set in Z_41976

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008388 | ljcr-ds-41987-20993-10496-41987

(41987, 20993, 10496)-difference set in Z_41987

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008389 | ljcr-ds-4201-1576-591-4201

(4201, 1576, 591)-difference set in Z_4201

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008390 | ljcr-ds-42021-8405-1681-42021

(42021, 8405, 1681)-difference set in Z_42021

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008391 | ljcr-ds-42031-21015-10507-42031

(42031, 21015, 10507)-difference set in Z_42031

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008392 | ljcr-ds-42031-8406-1681-42031

(42031, 8406, 1681)-difference set in Z_42031

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008393 | ljcr-ds-42043-14014-4671-42043

(42043, 14014, 4671)-difference set in Z_42043

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008394 | ljcr-ds-42075-9265-2040-42075

(42075, 9265, 2040)-difference set in Z_42075

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008395 | ljcr-ds-42123-21061-10530-42123

(42123, 21061, 10530)-difference set in Z_42123

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008396 | ljcr-ds-42127-21063-10531-42127

(42127, 21063, 10531)-difference set in Z_42127

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008397 | ljcr-ds-42144-11323-3042-42144

(42144, 11323, 3042)-difference set in Z_42144

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008398 | ljcr-ds-42151-15175-5463-42151

(42151, 15175, 5463)-difference set in Z_42151

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008399 | ljcr-ds-42155-21077-10538-42155

(42155, 21077, 10538)-difference set in Z_42155

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008400 | ljcr-ds-42161-8960-1904-42161

(42161, 8960, 1904)-difference set in Z_42161

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008401 | ljcr-ds-42175-10935-2835-42175

(42175, 10935, 2835)-difference set in Z_42175

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008402 | ljcr-ds-42211-7371-1287-42211

(42211, 7371, 1287)-difference set in Z_42211

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008403 | ljcr-ds-42259-14086-4695-42259

(42259, 14086, 4695)-difference set in Z_42259

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008404 | ljcr-ds-42275-920-20-5-8455

(42275, 920, 20)-difference set in Z_5 x Z_8455

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008405 | ljcr-ds-42277-10569-2642-42277

(42277, 10569, 2642)-difference set in Z_42277

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008406 | ljcr-ds-42333-8912-1876-42333

(42333, 8912, 1876)-difference set in Z_42333

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008407 | ljcr-ds-42351-18151-7779-42351

(42351, 18151, 7779)-difference set in Z_42351

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008408 | ljcr-ds-42361-16945-6778-42361

(42361, 16945, 6778)-difference set in Z_42361

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008409 | ljcr-ds-42383-21191-10595-42383

(42383, 21191, 10595)-difference set in Z_42383

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008410 | ljcr-ds-42447-21223-10611-42447

(42447, 21223, 10611)-difference set in Z_42447

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008411 | ljcr-ds-42449-4928-572-42449

(42449, 4928, 572)-difference set in Z_42449

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008412 | ljcr-ds-42471-12741-3822-42471

(42471, 12741, 3822)-difference set in Z_42471

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008413 | ljcr-ds-42484-11271-2990-42484

(42484, 11271, 2990)-difference set in Z_42484

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008414 | ljcr-ds-425-160-60-5-85

(425, 160, 60)-difference set in Z_5 x Z_85

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008415 | ljcr-ds-42507-21253-10626-42507

(42507, 21253, 10626)-difference set in Z_42507

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008416 | ljcr-ds-42511-21255-10627-42511

(42511, 21255, 10627)-difference set in Z_42511

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008417 | ljcr-ds-42515-21257-10628-42515

(42515, 21257, 10628)-difference set in Z_42515

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008418 | ljcr-ds-42541-17725-7385-42541

(42541, 17725, 7385)-difference set in Z_42541

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008419 | ljcr-ds-42543-21271-10635-42543

(42543, 21271, 10635)-difference set in Z_42543

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008420 | ljcr-ds-4256-1036-252-2-2128

(4256, 1036, 252)-difference set in Z_2 x Z_2128

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008421 | ljcr-ds-4256-1036-252-4-1064

(4256, 1036, 252)-difference set in Z_4 x Z_1064

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008422 | ljcr-ds-42597-10649-2662-42597

(42597, 10649, 2662)-difference set in Z_42597

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008423 | ljcr-ds-42602-3277-252-42602

(42602, 3277, 252)-difference set in Z_42602

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008424 | ljcr-ds-42613-9381-2065-42613

(42613, 9381, 2065)-difference set in Z_42613

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008425 | ljcr-ds-42615-1937-88-42615

(42615, 1937, 88)-difference set in Z_42615

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008426 | ljcr-ds-42625-1665-65-42625

(42625, 1665, 65)-difference set in Z_42625

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008427 | ljcr-ds-42637-14212-4737-42637

(42637, 14212, 4737)-difference set in Z_42637

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008428 | ljcr-ds-42673-16129-6096-42673

(42673, 16129, 6096)-difference set in Z_42673

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008429 | ljcr-ds-42735-21367-10683-42735

(42735, 21367, 10683)-difference set in Z_42735

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008430 | ljcr-ds-42745-8632-1743-42745

(42745, 8632, 1743)-difference set in Z_42745

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008431 | ljcr-ds-42747-21373-10686-42747

(42747, 21373, 10686)-difference set in Z_42747

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008432 | ljcr-ds-42805-9513-2114-42805

(42805, 9513, 2114)-difference set in Z_42805

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008433 | ljcr-ds-42813-10704-2676-42813

(42813, 10704, 2676)-difference set in Z_42813

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008434 | ljcr-ds-42829-10708-2677-42829

(42829, 10708, 2677)-difference set in Z_42829

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008435 | ljcr-ds-42883-7147-1191-42883

(42883, 7147, 1191)-difference set in Z_42883

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008436 | ljcr-ds-42927-21463-10731-42927

(42927, 21463, 10731)-difference set in Z_42927

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008437 | ljcr-ds-42949-10737-2684-42949

(42949, 10737, 2684)-difference set in Z_42949

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008438 | ljcr-ds-42951-12026-3367-42951

(42951, 12026, 3367)-difference set in Z_42951

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008439 | ljcr-ds-43023-21511-10755-43023

(43023, 21511, 10755)-difference set in Z_43023

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008440 | ljcr-ds-43025-2689-168-43025

(43025, 2689, 168)-difference set in Z_43025

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008441 | ljcr-ds-43029-10757-2689-43029

(43029, 10757, 2689)-difference set in Z_43029

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008442 | ljcr-ds-43033-4401-450-43033

(43033, 4401, 450)-difference set in Z_43033

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008443 | ljcr-ds-43036-3021-212-43036

(43036, 3021, 212)-difference set in Z_43036

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008444 | ljcr-ds-43057-13248-4076-43057

(43057, 13248, 4076)-difference set in Z_43057

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008445 | ljcr-ds-43087-21543-10771-43087

(43087, 21543, 10771)-difference set in Z_43087

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008446 | ljcr-ds-43111-19161-8516-43111

(43111, 19161, 8516)-difference set in Z_43111

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008447 | ljcr-ds-43111-6706-1043-43111

(43111, 6706, 1043)-difference set in Z_43111

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008448 | ljcr-ds-43183-21591-10795-43183

(43183, 21591, 10795)-difference set in Z_43183

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008449 | ljcr-ds-43213-3601-300-43213

(43213, 3601, 300)-difference set in Z_43213

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008450 | ljcr-ds-43263-2231-115-43263

(43263, 2231, 115)-difference set in Z_43263

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008451 | ljcr-ds-43267-21633-10816-43267

(43267, 21633, 10816)-difference set in Z_43267

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008452 | ljcr-ds-43282-12825-3800-43282

(43282, 12825, 3800)-difference set in Z_43282

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008453 | ljcr-ds-43291-9621-2138-43291

(43291, 9621, 2138)-difference set in Z_43291

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008454 | ljcr-ds-43307-18350-7775-43307

(43307, 18350, 7775)-difference set in Z_43307

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008455 | ljcr-ds-43311-21655-10827-43311

(43311, 21655, 10827)-difference set in Z_43311

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008456 | ljcr-ds-43321-14440-4813-43321

(43321, 14440, 4813)-difference set in Z_43321

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008457 | ljcr-ds-43369-12649-3689-43369

(43369, 12649, 3689)-difference set in Z_43369

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008458 | ljcr-ds-43401-15625-5625-43401

(43401, 15625, 5625)-difference set in Z_43401

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008459 | ljcr-ds-43407-21703-10851-43407

(43407, 21703, 10851)-difference set in Z_43407

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008460 | ljcr-ds-43435-17272-6868-43435

(43435, 17272, 6868)-difference set in Z_43435

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008461 | ljcr-ds-43435-6859-1083-43435

(43435, 6859, 1083)-difference set in Z_43435

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008462 | ljcr-ds-43439-12914-3839-43439

(43439, 12914, 3839)-difference set in Z_43439

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008463 | ljcr-ds-43439-21719-10859-43439

(43439, 21719, 10859)-difference set in Z_43439

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008464 | ljcr-ds-43471-15526-5545-43471

(43471, 15526, 5545)-difference set in Z_43471

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008465 | ljcr-ds-43471-6210-887-43471

(43471, 6210, 887)-difference set in Z_43471

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008466 | ljcr-ds-43485-20189-9373-43485

(43485, 20189, 9373)-difference set in Z_43485

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008467 | ljcr-ds-43493-8384-1616-43493

(43493, 8384, 1616)-difference set in Z_43493

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008468 | ljcr-ds-43511-17405-6962-43511

(43511, 17405, 6962)-difference set in Z_43511

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008469 | ljcr-ds-43520-6217-888-43520

(43520, 6217, 888)-difference set in Z_43520

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008470 | ljcr-ds-43525-10881-2720-43525

(43525, 10881, 2720)-difference set in Z_43525

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008471 | ljcr-ds-43540-4417-448-43540

(43540, 4417, 448)-difference set in Z_43540

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008472 | ljcr-ds-4355-2177-1088-4355

(4355, 2177, 1088)-difference set in Z_4355

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008473 | ljcr-ds-43563-5682-741-43563

(43563, 5682, 741)-difference set in Z_43563

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008474 | ljcr-ds-43611-21805-10902-43611

(43611, 21805, 10902)-difference set in Z_43611

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008475 | ljcr-ds-43617-10528-2541-43617

(43617, 10528, 2541)-difference set in Z_43617

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008476 | ljcr-ds-4367-1888-816-4367

(4367, 1888, 816)-difference set in Z_4367

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008477 | ljcr-ds-4370-2057-968-4370

(4370, 2057, 968)-difference set in Z_4370

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008478 | ljcr-ds-43711-13020-3878-43711

(43711, 13020, 3878)-difference set in Z_43711

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008479 | ljcr-ds-43717-10929-2732-43717

(43717, 10929, 2732)-difference set in Z_43717

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008480 | ljcr-ds-43729-14577-4859-43729

(43729, 14577, 4859)-difference set in Z_43729

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008481 | ljcr-ds-43749-10937-2734-43749

(43749, 10937, 2734)-difference set in Z_43749

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008482 | ljcr-ds-43773-10944-2736-43773

(43773, 10944, 2736)-difference set in Z_43773

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008483 | ljcr-ds-43781-10945-2736-43781

(43781, 10945, 2736)-difference set in Z_43781

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008484 | ljcr-ds-43785-8841-1785-43785

(43785, 8841, 1785)-difference set in Z_43785

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008485 | ljcr-ds-43804-19468-8652-43804

(43804, 19468, 8652)-difference set in Z_43804

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008486 | ljcr-ds-43841-17536-7014-43841

(43841, 17536, 7014)-difference set in Z_43841

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008487 | ljcr-ds-43883-21941-10970-43883

(43883, 21941, 10970)-difference set in Z_43883

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008488 | ljcr-ds-43897-5488-686-43897

(43897, 5488, 686)-difference set in Z_43897

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008489 | ljcr-ds-4393-793-143-4393

(4393, 793, 143)-difference set in Z_4393

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008490 | ljcr-ds-43951-14650-4883-43951

(43951, 14650, 4883)-difference set in Z_43951

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008491 | ljcr-ds-4401-176-7-3-1467

(4401, 176, 7)-difference set in Z_3 x Z_1467

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008492 | ljcr-ds-4401-176-7-3-3-489

(4401, 176, 7)-difference set in Z_3 x Z_3 x Z_489

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008493 | ljcr-ds-4401-176-7-4401

(4401, 176, 7)-difference set in Z_4401

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008494 | ljcr-ds-44023-12673-3648-44023

(44023, 12673, 3648)-difference set in Z_44023

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008495 | ljcr-ds-44092-3267-242-44092

(44092, 3267, 242)-difference set in Z_44092

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008496 | ljcr-ds-44111-17645-7058-44111

(44111, 17645, 7058)-difference set in Z_44111

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008497 | ljcr-ds-44116-21625-10600-44116

(44116, 21625, 10600)-difference set in Z_44116

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008498 | ljcr-ds-44145-21360-10335-44145

(44145, 21360, 10335)-difference set in Z_44145

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008499 | ljcr-ds-44157-4732-507-44157

(44157, 4732, 507)-difference set in Z_44157

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008500 | ljcr-ds-4417-1473-491-4417

(4417, 1473, 491)-difference set in Z_4417

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008501 | ljcr-ds-44177-19328-8456-44177

(44177, 19328, 8456)-difference set in Z_44177

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008502 | ljcr-ds-4421-1105-276-4421

(4421, 1105, 276)-difference set in Z_4421

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008503 | ljcr-ds-44220-12635-3610-44220

(44220, 12635, 3610)-difference set in Z_44220

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008504 | ljcr-ds-44243-22121-11060-44243

(44243, 22121, 11060)-difference set in Z_44243

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008505 | ljcr-ds-44252-20825-9800-44252

(44252, 20825, 9800)-difference set in Z_44252

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008506 | ljcr-ds-44287-22143-11071-44287

(44287, 22143, 11071)-difference set in Z_44287

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008507 | ljcr-ds-44289-6401-925-44289

(44289, 6401, 925)-difference set in Z_44289

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008508 | ljcr-ds-4431-886-177-4431

(4431, 886, 177)-difference set in Z_4431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008509 | ljcr-ds-44330-20565-9540-44330

(44330, 20565, 9540)-difference set in Z_44330

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008510 | ljcr-ds-44357-11089-2772-44357

(44357, 11089, 2772)-difference set in Z_44357

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008511 | ljcr-ds-44401-10176-2332-44401

(44401, 10176, 2332)-difference set in Z_44401

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008512 | ljcr-ds-44415-8799-1743-44415

(44415, 8799, 1743)-difference set in Z_44415

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008513 | ljcr-ds-44431-22215-11107-44431

(44431, 22215, 11107)-difference set in Z_44431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008514 | ljcr-ds-4446-1905-816-4446

(4446, 1905, 816)-difference set in Z_4446

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008515 | ljcr-ds-44463-22231-11115-44463

(44463, 22231, 11115)-difference set in Z_44463

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008516 | ljcr-ds-44491-14830-4943-44491

(44491, 14830, 4943)-difference set in Z_44491

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008517 | ljcr-ds-44499-9368-1972-44499

(44499, 9368, 1972)-difference set in Z_44499

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008518 | ljcr-ds-44517-6821-1045-44517

(44517, 6821, 1045)-difference set in Z_44517

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008519 | ljcr-ds-44527-22263-11131-44527

(44527, 22263, 11131)-difference set in Z_44527

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008520 | ljcr-ds-44541-3145-222-44541

(44541, 3145, 222)-difference set in Z_44541

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008521 | ljcr-ds-44549-11137-2784-44549

(44549, 11137, 2784)-difference set in Z_44549

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008522 | ljcr-ds-4455-918-189-3-1485

(4455, 918, 189)-difference set in Z_3 x Z_1485

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008523 | ljcr-ds-4455-918-189-9-495

(4455, 918, 189)-difference set in Z_9 x Z_495

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008524 | ljcr-ds-44556-7505-1264-44556

(44556, 7505, 1264)-difference set in Z_44556

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008525 | ljcr-ds-44575-7429-1238-44575

(44575, 7429, 1238)-difference set in Z_44575

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008526 | ljcr-ds-44604-19929-8904-44604

(44604, 19929, 8904)-difference set in Z_44604

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008527 | ljcr-ds-44605-19117-8193-44605

(44605, 19117, 8193)-difference set in Z_44605

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008528 | ljcr-ds-44611-7435-1239-44611

(44611, 7435, 1239)-difference set in Z_44611

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008529 | ljcr-ds-44643-18786-7905-44643

(44643, 18786, 7905)-difference set in Z_44643

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008530 | ljcr-ds-4465-496-55-4465

(4465, 496, 55)-difference set in Z_4465

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008531 | ljcr-ds-44650-19845-8820-44650

(44650, 19845, 8820)-difference set in Z_44650

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008532 | ljcr-ds-44655-22327-11163-44655

(44655, 22327, 11163)-difference set in Z_44655

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008533 | ljcr-ds-44660-4403-434-44660

(44660, 4403, 434)-difference set in Z_44660

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008534 | ljcr-ds-44671-8935-1787-44671

(44671, 8935, 1787)-difference set in Z_44671

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008535 | ljcr-ds-44691-22345-11172-44691

(44691, 22345, 11172)-difference set in Z_44691

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008536 | ljcr-ds-44721-16641-6192-44721

(44721, 16641, 6192)-difference set in Z_44721

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008537 | ljcr-ds-44721-8385-1572-44721

(44721, 8385, 1572)-difference set in Z_44721

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008538 | ljcr-ds-4473-1248-348-4473

(4473, 1248, 348)-difference set in Z_4473

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008539 | ljcr-ds-44733-10972-2691-44733

(44733, 10972, 2691)-difference set in Z_44733

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008540 | ljcr-ds-44741-11185-2796-44741

(44741, 11185, 2796)-difference set in Z_44741

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008541 | ljcr-ds-448-150-50-2-2-112

(448, 150, 50)-difference set in Z_2 x Z_2 x Z_112

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008542 | ljcr-ds-448-150-50-2-224

(448, 150, 50)-difference set in Z_2 x Z_224

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008543 | ljcr-ds-448-150-50-2-4-56

(448, 150, 50)-difference set in Z_2 x Z_4 x Z_56

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008544 | ljcr-ds-448-150-50-4-112

(448, 150, 50)-difference set in Z_4 x Z_112

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008545 | ljcr-ds-448-150-50-8-56

(448, 150, 50)-difference set in Z_8 x Z_56

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008546 | ljcr-ds-44881-12496-3479-44881

(44881, 12496, 3479)-difference set in Z_44881

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008547 | ljcr-ds-44883-22441-11220-44883

(44883, 22441, 11220)-difference set in Z_44883

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008548 | ljcr-ds-4489-561-70-67-67

(4489, 561, 70)-difference set in Z_67 x Z_67

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008549 | ljcr-ds-44919-19425-8400-44919

(44919, 19425, 8400)-difference set in Z_44919

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008550 | ljcr-ds-44943-22471-11235-44943

(44943, 22471, 11235)-difference set in Z_44943

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008551 | ljcr-ds-44959-7494-1249-44959

(44959, 7494, 1249)-difference set in Z_44959

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008552 | ljcr-ds-44973-11244-2811-44973

(44973, 11244, 2811)-difference set in Z_44973

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008553 | ljcr-ds-44991-22495-11247-44991

(44991, 22495, 11247)-difference set in Z_44991

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008554 | ljcr-ds-45003-22501-11250-45003

(45003, 22501, 11250)-difference set in Z_45003

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008555 | ljcr-ds-45017-10592-2492-45017

(45017, 10592, 2492)-difference set in Z_45017

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008556 | ljcr-ds-45018-19293-8268-45018

(45018, 19293, 8268)-difference set in Z_45018

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008557 | ljcr-ds-45051-22525-11262-45051

(45051, 22525, 11262)-difference set in Z_45051

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008558 | ljcr-ds-45069-11268-2817-45069

(45069, 11268, 2817)-difference set in Z_45069

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008559 | ljcr-ds-45103-22551-11275-45103

(45103, 22551, 11275)-difference set in Z_45103

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008560 | ljcr-ds-45136-7021-1092-45136

(45136, 7021, 1092)-difference set in Z_45136

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008561 | ljcr-ds-45141-11285-2821-45141

(45141, 11285, 2821)-difference set in Z_45141

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008562 | ljcr-ds-45152-2771-170-45152

(45152, 2771, 170)-difference set in Z_45152

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008563 | ljcr-ds-45193-5649-706-45193

(45193, 5649, 706)-difference set in Z_45193

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008564 | ljcr-ds-45257-5657-707-45257

(45257, 5657, 707)-difference set in Z_45257

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008565 | ljcr-ds-45267-1742-67-45267

(45267, 1742, 67)-difference set in Z_45267

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008566 | ljcr-ds-45277-9604-2037-45277

(45277, 9604, 2037)-difference set in Z_45277

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008567 | ljcr-ds-45288-18975-7950-45288

(45288, 18975, 7950)-difference set in Z_45288

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008568 | ljcr-ds-45319-16185-5780-45319

(45319, 16185, 5780)-difference set in Z_45319

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008569 | ljcr-ds-45323-20554-9321-45323

(45323, 20554, 9321)-difference set in Z_45323

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008570 | ljcr-ds-45325-1260-35-45325

(45325, 1260, 35)-difference set in Z_45325

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008571 | ljcr-ds-45327-22663-11331-45327

(45327, 22663, 11331)-difference set in Z_45327

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008572 | ljcr-ds-4533-825-150-4533

(4533, 825, 150)-difference set in Z_4533

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008573 | ljcr-ds-45356-5405-644-45356

(45356, 5405, 644)-difference set in Z_45356

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008574 | ljcr-ds-45391-22695-11347-45391

(45391, 22695, 11347)-difference set in Z_45391

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008575 | ljcr-ds-45401-19976-8789-45401

(45401, 19976, 8789)-difference set in Z_45401

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008576 | ljcr-ds-45409-15136-5045-45409

(45409, 15136, 5045)-difference set in Z_45409

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008577 | ljcr-ds-45431-15576-5340-45431

(45431, 15576, 5340)-difference set in Z_45431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008578 | ljcr-ds-45431-8261-1502-45431

(45431, 8261, 1502)-difference set in Z_45431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008579 | ljcr-ds-45431-9086-1817-45431

(45431, 9086, 1817)-difference set in Z_45431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008580 | ljcr-ds-45445-11361-2840-45445

(45445, 11361, 2840)-difference set in Z_45445

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008581 | ljcr-ds-45451-22725-11362-45451

(45451, 22725, 11362)-difference set in Z_45451

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008582 | ljcr-ds-45477-11369-2842-45477

(45477, 11369, 2842)-difference set in Z_45477

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008583 | ljcr-ds-45505-6400-900-45505

(45505, 6400, 900)-difference set in Z_45505

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008584 | ljcr-ds-45511-4551-455-45511

(45511, 4551, 455)-difference set in Z_45511

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008585 | ljcr-ds-45531-22765-11382-45531

(45531, 22765, 11382)-difference set in Z_45531

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008586 | ljcr-ds-45541-11385-2846-45541

(45541, 11385, 2846)-difference set in Z_45541

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008587 | ljcr-ds-4555-990-215-4555

(4555, 990, 215)-difference set in Z_4555

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008588 | ljcr-ds-45563-10792-2556-45563

(45563, 10792, 2556)-difference set in Z_45563

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008589 | ljcr-ds-4557-68-1-7-651

(4557, 68, 1)-difference set in Z_7 x Z_651

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008590 | ljcr-ds-45585-19537-8373-45585

(45585, 19537, 8373)-difference set in Z_45585

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008591 | ljcr-ds-45585-2849-178-45585

(45585, 2849, 178)-difference set in Z_45585

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008592 | ljcr-ds-45594-18669-7644-45594

(45594, 18669, 7644)-difference set in Z_45594

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008593 | ljcr-ds-45601-7201-1137-45601

(45601, 7201, 1137)-difference set in Z_45601

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008594 | ljcr-ds-45631-9126-1825-45631

(45631, 9126, 1825)-difference set in Z_45631

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008595 | ljcr-ds-4565-1793-704-4565

(4565, 1793, 704)-difference set in Z_4565

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008596 | ljcr-ds-45683-22841-11420-45683

(45683, 22841, 11420)-difference set in Z_45683

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008597 | ljcr-ds-45711-20896-9552-45711

(45711, 20896, 9552)-difference set in Z_45711

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008598 | ljcr-ds-45711-22855-11427-45711

(45711, 22855, 11427)-difference set in Z_45711

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008599 | ljcr-ds-45731-13720-4116-45731

(45731, 13720, 4116)-difference set in Z_45731

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008600 | ljcr-ds-45731-21251-9875-45731

(45731, 21251, 9875)-difference set in Z_45731

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008601 | ljcr-ds-45751-7626-1271-45751

(45751, 7626, 1271)-difference set in Z_45751

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008602 | ljcr-ds-4576-550-66-2-2-1144

(4576, 550, 66)-difference set in Z_2 x Z_2 x Z_1144

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008603 | ljcr-ds-4576-550-66-2-2-2-572

(4576, 550, 66)-difference set in Z_2 x Z_2 x Z_2 x Z_572

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008604 | ljcr-ds-4576-550-66-2-2288

(4576, 550, 66)-difference set in Z_2 x Z_2288

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008605 | ljcr-ds-4576-550-66-2-4-572

(4576, 550, 66)-difference set in Z_2 x Z_4 x Z_572

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008606 | ljcr-ds-4576-550-66-4-1144

(4576, 550, 66)-difference set in Z_4 x Z_1144

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008607 | ljcr-ds-45771-22885-11442-45771

(45771, 22885, 11442)-difference set in Z_45771

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008608 | ljcr-ds-45775-22887-11443-45775

(45775, 22887, 11443)-difference set in Z_45775

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008609 | ljcr-ds-45787-22893-11446-45787

(45787, 22893, 11446)-difference set in Z_45787

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008610 | ljcr-ds-45793-18657-7601-45793

(45793, 18657, 7601)-difference set in Z_45793

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008611 | ljcr-ds-45795-22897-11448-45795

(45795, 22897, 11448)-difference set in Z_45795

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008612 | ljcr-ds-45807-22903-11451-45807

(45807, 22903, 11451)-difference set in Z_45807

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008613 | ljcr-ds-45850-18445-7420-45850

(45850, 18445, 7420)-difference set in Z_45850

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008614 | ljcr-ds-45867-22933-11466-45867

(45867, 22933, 11466)-difference set in Z_45867

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008615 | ljcr-ds-45871-17931-7009-45871

(45871, 17931, 7009)-difference set in Z_45871

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008616 | ljcr-ds-45881-3256-231-45881

(45881, 3256, 231)-difference set in Z_45881

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008617 | ljcr-ds-45903-10503-2403-45903

(45903, 10503, 2403)-difference set in Z_45903

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008618 | ljcr-ds-45936-18375-7350-45936

(45936, 18375, 7350)-difference set in Z_45936

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008619 | ljcr-ds-45949-15316-5105-45949

(45949, 15316, 5105)-difference set in Z_45949

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008620 | ljcr-ds-45963-22981-11490-45963

(45963, 22981, 11490)-difference set in Z_45963

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008621 | ljcr-ds-45969-17408-6592-45969

(45969, 17408, 6592)-difference set in Z_45969

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008622 | ljcr-ds-45989-11497-2874-45989

(45989, 11497, 2874)-difference set in Z_45989

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008623 | ljcr-ds-46020-10829-2548-46020

(46020, 10829, 2548)-difference set in Z_46020

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008624 | ljcr-ds-46021-11505-2876-46021

(46021, 11505, 2876)-difference set in Z_46021

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008625 | ljcr-ds-46063-23031-11515-46063

(46063, 23031, 11515)-difference set in Z_46063

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008626 | ljcr-ds-46079-23039-11519-46079

(46079, 23039, 11519)-difference set in Z_46079

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008627 | ljcr-ds-4608-272-16-16-288

(4608, 272, 16)-difference set in Z_16 x Z_288

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008628 | ljcr-ds-4608-272-16-2-16-144

(4608, 272, 16)-difference set in Z_2 x Z_16 x Z_144

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008629 | ljcr-ds-4608-272-16-2-2-12-96

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_12 x Z_96

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008630 | ljcr-ds-4608-272-16-2-2-2-12-48

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_12 x Z_48

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008631 | ljcr-ds-4608-272-16-2-2-2-2-12-24

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_12 x Z_24

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008632 | ljcr-ds-4608-272-16-2-2-2-2-2-12-12

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_12 x Z_12

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008633 | ljcr-ds-4608-272-16-2-2-2-2-2-144

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_144

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008634 | ljcr-ds-4608-272-16-2-2-2-2-2-2-36

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_36

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008635 | ljcr-ds-4608-272-16-2-2-2-2-2-2-6-12

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_6 x Z_12

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008636 | ljcr-ds-4608-272-16-2-2-2-2-2-2-72

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008637 | ljcr-ds-4608-272-16-2-2-2-2-2-4-36

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_4 x Z_36

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008638 | ljcr-ds-4608-272-16-2-2-2-2-2-6-24

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_6 x Z_24

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008639 | ljcr-ds-4608-272-16-2-2-2-2-288

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_288

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008640 | ljcr-ds-4608-272-16-2-2-2-2-4-72

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_4 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008641 | ljcr-ds-4608-272-16-2-2-2-2-6-48

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_6 x Z_48

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008642 | ljcr-ds-4608-272-16-2-2-2-24-24

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_24 x Z_24

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008643 | ljcr-ds-4608-272-16-2-2-2-4-12-12

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_4 x Z_12 x Z_12

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008644 | ljcr-ds-4608-272-16-2-2-2-4-144

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_4 x Z_144

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008645 | ljcr-ds-4608-272-16-2-2-2-4-4-36

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_4 x Z_4 x Z_36

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008646 | ljcr-ds-4608-272-16-2-2-2-6-96

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_6 x Z_96

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008647 | ljcr-ds-4608-272-16-2-2-2-8-72

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_2 x Z_8 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008648 | ljcr-ds-4608-272-16-2-2-24-48

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_24 x Z_48

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008649 | ljcr-ds-4608-272-16-2-2-4-12-24

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_4 x Z_12 x Z_24

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008650 | ljcr-ds-4608-272-16-2-2-4-288

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_4 x Z_288

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008651 | ljcr-ds-4608-272-16-2-2-4-4-72

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_4 x Z_4 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008652 | ljcr-ds-4608-272-16-2-2-8-144

(4608, 272, 16)-difference set in Z_2 x Z_2 x Z_8 x Z_144

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008653 | ljcr-ds-4608-272-16-2-24-96

(4608, 272, 16)-difference set in Z_2 x Z_24 x Z_96

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008654 | ljcr-ds-4608-272-16-2-4-12-48

(4608, 272, 16)-difference set in Z_2 x Z_4 x Z_12 x Z_48

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008655 | ljcr-ds-4608-272-16-2-4-24-24

(4608, 272, 16)-difference set in Z_2 x Z_4 x Z_24 x Z_24

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008656 | ljcr-ds-4608-272-16-2-4-4-12-12

(4608, 272, 16)-difference set in Z_2 x Z_4 x Z_4 x Z_12 x Z_12

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008657 | ljcr-ds-4608-272-16-2-4-4-144

(4608, 272, 16)-difference set in Z_2 x Z_4 x Z_4 x Z_144

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008658 | ljcr-ds-4608-272-16-2-4-4-4-36

(4608, 272, 16)-difference set in Z_2 x Z_4 x Z_4 x Z_4 x Z_36

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008659 | ljcr-ds-4608-272-16-2-4-8-72

(4608, 272, 16)-difference set in Z_2 x Z_4 x Z_8 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008660 | ljcr-ds-4608-272-16-2-48-48

(4608, 272, 16)-difference set in Z_2 x Z_48 x Z_48

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008661 | ljcr-ds-4608-272-16-2-8-288

(4608, 272, 16)-difference set in Z_2 x Z_8 x Z_288

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008662 | ljcr-ds-4608-272-16-4-12-96

(4608, 272, 16)-difference set in Z_4 x Z_12 x Z_96

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008663 | ljcr-ds-4608-272-16-4-24-48

(4608, 272, 16)-difference set in Z_4 x Z_24 x Z_48

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008664 | ljcr-ds-4608-272-16-4-4-12-24

(4608, 272, 16)-difference set in Z_4 x Z_4 x Z_12 x Z_24

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008665 | ljcr-ds-4608-272-16-4-4-288

(4608, 272, 16)-difference set in Z_4 x Z_4 x Z_288

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008666 | ljcr-ds-4608-272-16-4-4-4-72

(4608, 272, 16)-difference set in Z_4 x Z_4 x Z_4 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008667 | ljcr-ds-4608-272-16-4-8-144

(4608, 272, 16)-difference set in Z_4 x Z_8 x Z_144

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008668 | ljcr-ds-4608-272-16-48-96

(4608, 272, 16)-difference set in Z_48 x Z_96

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008669 | ljcr-ds-4608-272-16-8-24-24

(4608, 272, 16)-difference set in Z_8 x Z_24 x Z_24

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008670 | ljcr-ds-4608-272-16-8-8-72

(4608, 272, 16)-difference set in Z_8 x Z_8 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008671 | ljcr-ds-46085-11521-2880-46085

(46085, 11521, 2880)-difference set in Z_46085

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008672 | ljcr-ds-46091-4610-461-46091

(46091, 4610, 461)-difference set in Z_46091

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008673 | ljcr-ds-46093-15364-5121-46093

(46093, 15364, 5121)-difference set in Z_46093

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008674 | ljcr-ds-46110-4705-480-46110

(46110, 4705, 480)-difference set in Z_46110

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008675 | ljcr-ds-46159-8478-1557-46159

(46159, 8478, 1557)-difference set in Z_46159

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008676 | ljcr-ds-46177-14209-4372-46177

(46177, 14209, 4372)-difference set in Z_46177

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008677 | ljcr-ds-46181-11545-2886-46181

(46181, 11545, 2886)-difference set in Z_46181

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008678 | ljcr-ds-46189-14113-4312-46189

(46189, 14113, 4312)-difference set in Z_46189

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008679 | ljcr-ds-46191-23095-11547-46191

(46191, 23095, 11547)-difference set in Z_46191

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008680 | ljcr-ds-46211-4621-462-46211

(46211, 4621, 462)-difference set in Z_46211

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008681 | ljcr-ds-46221-9245-1849-46221

(46221, 9245, 1849)-difference set in Z_46221

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008682 | ljcr-ds-46249-13161-3745-46249

(46249, 13161, 3745)-difference set in Z_46249

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008683 | ljcr-ds-4627-514-57-4627

(4627, 514, 57)-difference set in Z_4627

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008684 | ljcr-ds-46271-19831-8499-46271

(46271, 19831, 8499)-difference set in Z_46271

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008685 | ljcr-ds-46277-11569-2892-46277

(46277, 11569, 2892)-difference set in Z_46277

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008686 | ljcr-ds-46284-8959-1734-46284

(46284, 8959, 1734)-difference set in Z_46284

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008687 | ljcr-ds-46291-10801-2520-46291

(46291, 10801, 2520)-difference set in Z_46291

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008688 | ljcr-ds-46313-6616-945-46313

(46313, 6616, 945)-difference set in Z_46313

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008689 | ljcr-ds-46323-23161-11580-46323

(46323, 23161, 11580)-difference set in Z_46323

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008690 | ljcr-ds-46351-8550-1577-46351

(46351, 8550, 1577)-difference set in Z_46351

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008691 | ljcr-ds-46355-12299-3263-46355

(46355, 12299, 3263)-difference set in Z_46355

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008692 | ljcr-ds-46376-11501-2852-46376

(46376, 11501, 2852)-difference set in Z_46376

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008693 | ljcr-ds-46387-23193-11596-46387

(46387, 23193, 11596)-difference set in Z_46387

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008694 | ljcr-ds-46399-16170-5635-46399

(46399, 16170, 5635)-difference set in Z_46399

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008695 | ljcr-ds-46411-8841-1684-46411

(46411, 8841, 1684)-difference set in Z_46411

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008696 | ljcr-ds-46447-7741-1290-46447

(46447, 7741, 1290)-difference set in Z_46447

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008697 | ljcr-ds-465-145-45-465

(465, 145, 45)-difference set in Z_465

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008698 | ljcr-ds-46531-15840-5392-46531

(46531, 15840, 5392)-difference set in Z_46531

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008699 | ljcr-ds-46531-9306-1861-46531

(46531, 9306, 1861)-difference set in Z_46531

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008700 | ljcr-ds-46543-15514-5171-46543

(46543, 15514, 5171)-difference set in Z_46543

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008701 | ljcr-ds-46543-23271-11635-46543

(46543, 23271, 11635)-difference set in Z_46543

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008702 | ljcr-ds-46565-19957-8553-46565

(46565, 19957, 8553)-difference set in Z_46565

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008703 | ljcr-ds-46585-648-9-11-11-385

(46585, 648, 9)-difference set in Z_11 x Z_11 x Z_385

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008704 | ljcr-ds-46585-648-9-11-4235

(46585, 648, 9)-difference set in Z_11 x Z_4235

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008705 | ljcr-ds-46585-648-9-46585

(46585, 648, 9)-difference set in Z_46585

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008706 | ljcr-ds-46593-10752-2481-46593

(46593, 10752, 2481)-difference set in Z_46593

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008707 | ljcr-ds-46611-23305-11652-46611

(46611, 23305, 11652)-difference set in Z_46611

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008708 | ljcr-ds-46627-7771-1295-46627

(46627, 7771, 1295)-difference set in Z_46627

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008709 | ljcr-ds-46653-11664-2916-46653

(46653, 11664, 2916)-difference set in Z_46653

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008710 | ljcr-ds-46659-18778-7557-46659

(46659, 18778, 7557)-difference set in Z_46659

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008711 | ljcr-ds-46663-7777-1296-46663

(46663, 7777, 1296)-difference set in Z_46663

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008712 | ljcr-ds-46671-23335-11667-46671

(46671, 23335, 11667)-difference set in Z_46671

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008713 | ljcr-ds-46671-9335-1867-46671

(46671, 9335, 1867)-difference set in Z_46671

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008714 | ljcr-ds-46683-17850-6825-46683

(46683, 17850, 6825)-difference set in Z_46683

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008715 | ljcr-ds-46707-23353-11676-46707

(46707, 23353, 11676)-difference set in Z_46707

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008716 | ljcr-ds-46717-22900-11225-46717

(46717, 22900, 11225)-difference set in Z_46717

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008717 | ljcr-ds-46789-5013-537-46789

(46789, 5013, 537)-difference set in Z_46789

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008718 | ljcr-ds-4681-2080-924-4681

(4681, 2080, 924)-difference set in Z_4681

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008719 | ljcr-ds-46831-7806-1301-46831

(46831, 7806, 1301)-difference set in Z_46831

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008720 | ljcr-ds-46873-17361-6430-46873

(46873, 17361, 6430)-difference set in Z_46873

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008721 | ljcr-ds-469-208-92-469

(469, 208, 92)-difference set in Z_469

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008722 | ljcr-ds-46905-18656-7420-46905

(46905, 18656, 7420)-difference set in Z_46905

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008723 | ljcr-ds-46907-23453-11726-46907

(46907, 23453, 11726)-difference set in Z_46907

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008724 | ljcr-ds-46911-23455-11727-46911

(46911, 23455, 11727)-difference set in Z_46911

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008725 | ljcr-ds-46915-7819-1303-46915

(46915, 7819, 1303)-difference set in Z_46915

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008726 | ljcr-ds-46921-15640-5213-46921

(46921, 15640, 5213)-difference set in Z_46921

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008727 | ljcr-ds-46928-17703-6678-46928

(46928, 17703, 6678)-difference set in Z_46928

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008728 | ljcr-ds-46931-3250-225-46931

(46931, 3250, 225)-difference set in Z_46931

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008729 | ljcr-ds-46971-15555-5151-46971

(46971, 15555, 5151)-difference set in Z_46971

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008730 | ljcr-ds-46971-4026-345-46971

(46971, 4026, 345)-difference set in Z_46971

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008731 | ljcr-ds-46981-14500-4475-46981

(46981, 14500, 4475)-difference set in Z_46981

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008732 | ljcr-ds-46981-20736-9152-46981

(46981, 20736, 9152)-difference set in Z_46981

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008733 | ljcr-ds-46981-9396-1879-46981

(46981, 9396, 1879)-difference set in Z_46981

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008734 | ljcr-ds-46999-9325-1850-46999

(46999, 9325, 1850)-difference set in Z_46999

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008735 | ljcr-ds-47009-21697-10014-47009

(47009, 21697, 10014)-difference set in Z_47009

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008736 | ljcr-ds-47043-23521-11760-47043

(47043, 23521, 11760)-difference set in Z_47043

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008737 | ljcr-ds-47163-23581-11790-47163

(47163, 23581, 11790)-difference set in Z_47163

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008738 | ljcr-ds-4719-337-24-11-429

(4719, 337, 24)-difference set in Z_11 x Z_429

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008739 | ljcr-ds-47235-23617-11808-47235

(47235, 23617, 11808)-difference set in Z_47235

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008740 | ljcr-ds-47239-7873-1312-47239

(47239, 7873, 1312)-difference set in Z_47239

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008741 | ljcr-ds-47247-23623-11811-47247

(47247, 23623, 11811)-difference set in Z_47247

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008742 | ljcr-ds-4726-1701-612-4726

(4726, 1701, 612)-difference set in Z_4726

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008743 | ljcr-ds-47269-11817-2954-47269

(47269, 11817, 2954)-difference set in Z_47269

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008744 | ljcr-ds-47311-10375-2275-47311

(47311, 10375, 2275)-difference set in Z_47311

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008745 | ljcr-ds-47311-11040-2576-47311

(47311, 11040, 2576)-difference set in Z_47311

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008746 | ljcr-ds-47323-4302-391-47323

(47323, 4302, 391)-difference set in Z_47323

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008747 | ljcr-ds-47383-23691-11845-47383

(47383, 23691, 11845)-difference set in Z_47383

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008748 | ljcr-ds-47389-15796-5265-47389

(47389, 15796, 5265)-difference set in Z_47389

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008749 | ljcr-ds-47397-11849-2962-47397

(47397, 11849, 2962)-difference set in Z_47397

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008750 | ljcr-ds-4741-396-33-4741

(4741, 396, 33)-difference set in Z_4741

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008751 | ljcr-ds-47425-14080-4180-47425

(47425, 14080, 4180)-difference set in Z_47425

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008752 | ljcr-ds-47425-15808-5269-47425

(47425, 15808, 5269)-difference set in Z_47425

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008753 | ljcr-ds-47425-19969-8408-47425

(47425, 19969, 8408)-difference set in Z_47425

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008754 | ljcr-ds-47429-11857-2964-47429

(47429, 11857, 2964)-difference set in Z_47429

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008755 | ljcr-ds-47461-19776-8240-47461

(47461, 19776, 8240)-difference set in Z_47461

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008756 | ljcr-ds-47461-5425-620-47461

(47461, 5425, 620)-difference set in Z_47461

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008757 | ljcr-ds-47503-23751-11875-47503

(47503, 23751, 11875)-difference set in Z_47503

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008758 | ljcr-ds-47523-23761-11880-47523

(47523, 23761, 11880)-difference set in Z_47523

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008759 | ljcr-ds-47541-16640-5824-47541

(47541, 16640, 5824)-difference set in Z_47541

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008760 | ljcr-ds-47551-15850-5283-47551

(47551, 15850, 5283)-difference set in Z_47551

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008761 | ljcr-ds-47551-23775-11887-47551

(47551, 23775, 11887)-difference set in Z_47551

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008762 | ljcr-ds-47565-11892-2973-47565

(47565, 11892, 2973)-difference set in Z_47565

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008763 | ljcr-ds-4759-2197-1014-4759

(4759, 2197, 1014)-difference set in Z_4759

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008764 | ljcr-ds-4761-120-3-3-1587

(4761, 120, 3)-difference set in Z_3 x Z_1587

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008765 | ljcr-ds-4761-120-3-69-69

(4761, 120, 3)-difference set in Z_69 x Z_69

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008766 | ljcr-ds-47619-10673-2392-47619

(47619, 10673, 2392)-difference set in Z_47619

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008767 | ljcr-ds-47631-23815-11907-47631

(47631, 23815, 11907)-difference set in Z_47631

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008768 | ljcr-ds-47642-17325-6300-47642

(47642, 17325, 6300)-difference set in Z_47642

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008769 | ljcr-ds-47661-11916-2979-47661

(47661, 11916, 2979)-difference set in Z_47661

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008770 | ljcr-ds-477-204-87-3-159

(477, 204, 87)-difference set in Z_3 x Z_159

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008771 | ljcr-ds-47727-23863-11931-47727

(47727, 23863, 11931)-difference set in Z_47727

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008772 | ljcr-ds-47740-15913-5304-47740

(47740, 15913, 5304)-difference set in Z_47740

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008773 | ljcr-ds-47743-7957-1326-47743

(47743, 7957, 1326)-difference set in Z_47743

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008774 | ljcr-ds-47749-15916-5305-47749

(47749, 15916, 5305)-difference set in Z_47749

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008775 | ljcr-ds-47771-23885-11942-47771

(47771, 23885, 11942)-difference set in Z_47771

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008776 | ljcr-ds-47851-15225-4844-47851

(47851, 15225, 4844)-difference set in Z_47851

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008777 | ljcr-ds-47851-17226-6201-47851

(47851, 17226, 6201)-difference set in Z_47851

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008778 | ljcr-ds-47853-11964-2991-47853

(47853, 11964, 2991)-difference set in Z_47853

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008779 | ljcr-ds-47867-14729-4532-47867

(47867, 14729, 4532)-difference set in Z_47867

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008780 | ljcr-ds-47881-5985-748-47881

(47881, 5985, 748)-difference set in Z_47881

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008781 | ljcr-ds-47887-7981-1330-47887

(47887, 7981, 1330)-difference set in Z_47887

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008782 | ljcr-ds-47899-23949-11974-47899

(47899, 23949, 11974)-difference set in Z_47899

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008783 | ljcr-ds-47941-13396-3743-47941

(47941, 13396, 3743)-difference set in Z_47941

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008784 | ljcr-ds-47949-11988-2997-47949

(47949, 11988, 2997)-difference set in Z_47949

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008785 | ljcr-ds-47971-23985-11992-47971

(47971, 23985, 11992)-difference set in Z_47971

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008786 | ljcr-ds-47983-15994-5331-47983

(47983, 15994, 5331)-difference set in Z_47983

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008787 | ljcr-ds-48027-24013-12006-48027

(48027, 24013, 12006)-difference set in Z_48027

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008788 | ljcr-ds-48037-12009-3002-48037

(48037, 12009, 3002)-difference set in Z_48037

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008789 | ljcr-ds-48097-21376-9500-48097

(48097, 21376, 9500)-difference set in Z_48097

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008790 | ljcr-ds-48127-17277-6202-48127

(48127, 17277, 6202)-difference set in Z_48127

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008791 | ljcr-ds-48133-12033-3008-48133

(48133, 12033, 3008)-difference set in Z_48133

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008792 | ljcr-ds-48147-10318-2211-48147

(48147, 10318, 2211)-difference set in Z_48147

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008793 | ljcr-ds-48147-24073-12036-48147

(48147, 24073, 12036)-difference set in Z_48147

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008794 | ljcr-ds-48175-24087-12043-48175

(48175, 24087, 12043)-difference set in Z_48175

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008795 | ljcr-ds-48199-10527-2299-48199

(48199, 10527, 2299)-difference set in Z_48199

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008796 | ljcr-ds-48199-16066-5355-48199

(48199, 16066, 5355)-difference set in Z_48199

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008797 | ljcr-ds-48204-10621-2340-48204

(48204, 10621, 2340)-difference set in Z_48204

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008798 | ljcr-ds-48247-8041-1340-48247

(48247, 8041, 1340)-difference set in Z_48247

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008799 | ljcr-ds-48323-24161-12080-48323

(48323, 24161, 12080)-difference set in Z_48323

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008800 | ljcr-ds-48349-19908-8197-48349

(48349, 19908, 8197)-difference set in Z_48349

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008801 | ljcr-ds-48357-1100-25-48357

(48357, 1100, 25)-difference set in Z_48357

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008802 | ljcr-ds-48361-10881-2448-48361

(48361, 10881, 2448)-difference set in Z_48361

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008803 | ljcr-ds-48367-24183-12091-48367

(48367, 24183, 12091)-difference set in Z_48367

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008804 | ljcr-ds-4837-1612-537-4837

(4837, 1612, 537)-difference set in Z_4837

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008805 | ljcr-ds-48381-10620-2331-48381

(48381, 10620, 2331)-difference set in Z_48381

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008806 | ljcr-ds-48381-12096-3024-48381

(48381, 12096, 3024)-difference set in Z_48381

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008807 | ljcr-ds-48391-19356-7742-48391

(48391, 19356, 7742)-difference set in Z_48391

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008808 | ljcr-ds-48399-24199-12099-48399

(48399, 24199, 12099)-difference set in Z_48399

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008809 | ljcr-ds-48400-13651-3850-48400

(48400, 13651, 3850)-difference set in Z_48400

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008810 | ljcr-ds-48409-2017-84-48409

(48409, 2017, 84)-difference set in Z_48409

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008811 | ljcr-ds-48411-19365-7746-48411

(48411, 19365, 7746)-difference set in Z_48411

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008812 | ljcr-ds-48419-6193-792-48419

(48419, 6193, 792)-difference set in Z_48419

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008813 | ljcr-ds-48439-14743-4487-48439

(48439, 14743, 4487)-difference set in Z_48439

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008814 | ljcr-ds-4845-1904-748-4845

(4845, 1904, 748)-difference set in Z_4845

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008815 | ljcr-ds-48469-16156-5385-48469

(48469, 16156, 5385)-difference set in Z_48469

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008816 | ljcr-ds-48469-19041-7480-48469

(48469, 19041, 7480)-difference set in Z_48469

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008817 | ljcr-ds-48499-24249-12124-48499

(48499, 24249, 12124)-difference set in Z_48499

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008818 | ljcr-ds-48509-22445-10385-48509

(48509, 22445, 10385)-difference set in Z_48509

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008819 | ljcr-ds-48529-21232-9289-48529

(48529, 21232, 9289)-difference set in Z_48529

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008820 | ljcr-ds-48541-12945-3452-48541

(48541, 12945, 3452)-difference set in Z_48541

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008821 | ljcr-ds-48545-7585-1185-48545

(48545, 7585, 1185)-difference set in Z_48545

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008822 | ljcr-ds-48553-23409-11286-48553

(48553, 23409, 11286)-difference set in Z_48553

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008823 | ljcr-ds-48581-12145-3036-48581

(48581, 12145, 3036)-difference set in Z_48581

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008824 | ljcr-ds-48631-9726-1945-48631

(48631, 9726, 1945)-difference set in Z_48631

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008825 | ljcr-ds-48643-19966-8195-48643

(48643, 19966, 8195)-difference set in Z_48643

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008826 | ljcr-ds-48657-3041-190-48657

(48657, 3041, 190)-difference set in Z_48657

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008827 | ljcr-ds-48677-12169-3042-48677

(48677, 12169, 3042)-difference set in Z_48677

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008828 | ljcr-ds-48699-24349-12174-48699

(48699, 24349, 12174)-difference set in Z_48699

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008829 | ljcr-ds-48709-12177-3044-48709

(48709, 12177, 3044)-difference set in Z_48709

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008830 | ljcr-ds-48769-16257-5419-48769

(48769, 16257, 5419)-difference set in Z_48769

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008831 | ljcr-ds-48775-8130-1355-48775

(48775, 8130, 1355)-difference set in Z_48775

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008832 | ljcr-ds-48781-12196-3049-48781

(48781, 12196, 3049)-difference set in Z_48781

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008833 | ljcr-ds-48783-24391-12195-48783

(48783, 24391, 12195)-difference set in Z_48783

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008834 | ljcr-ds-48811-16270-5423-48811

(48811, 16270, 5423)-difference set in Z_48811

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008835 | ljcr-ds-48825-2873-169-48825

(48825, 2873, 169)-difference set in Z_48825

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008836 | ljcr-ds-48837-12209-3052-48837

(48837, 12209, 3052)-difference set in Z_48837

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008837 | ljcr-ds-4887-350-25-3-1629

(4887, 350, 25)-difference set in Z_3 x Z_1629

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008838 | ljcr-ds-4887-350-25-3-3-543

(4887, 350, 25)-difference set in Z_3 x Z_3 x Z_543

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008839 | ljcr-ds-4887-350-25-4887

(4887, 350, 25)-difference set in Z_4887

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008840 | ljcr-ds-48895-24447-12223-48895

(48895, 24447, 12223)-difference set in Z_48895

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008841 | ljcr-ds-4896-980-196-12-408

(4896, 980, 196)-difference set in Z_12 x Z_408

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008842 | ljcr-ds-4896-980-196-2-2448

(4896, 980, 196)-difference set in Z_2 x Z_2448

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008843 | ljcr-ds-4896-980-196-4-1224

(4896, 980, 196)-difference set in Z_4 x Z_1224

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008844 | ljcr-ds-4896-980-196-6-816

(4896, 980, 196)-difference set in Z_6 x Z_816

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008845 | ljcr-ds-48965-12241-3060-48965

(48965, 12241, 3060)-difference set in Z_48965

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008846 | ljcr-ds-48983-16758-5733-48983

(48983, 16758, 5733)-difference set in Z_48983

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008847 | ljcr-ds-48987-24493-12246-48987

(48987, 24493, 12246)-difference set in Z_48987

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008848 | ljcr-ds-48997-12249-3062-48997

(48997, 12249, 3062)-difference set in Z_48997

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008849 | ljcr-ds-49039-17832-6484-49039

(49039, 17832, 6484)-difference set in Z_49039

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008850 | ljcr-ds-49039-6688-912-49039

(49039, 6688, 912)-difference set in Z_49039

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008851 | ljcr-ds-49063-11544-2716-49063

(49063, 11544, 2716)-difference set in Z_49063

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008852 | ljcr-ds-49063-16354-5451-49063

(49063, 16354, 5451)-difference set in Z_49063

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008853 | ljcr-ds-49125-12281-3070-49125

(49125, 12281, 3070)-difference set in Z_49125

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008854 | ljcr-ds-49135-6897-968-49135

(49135, 6897, 968)-difference set in Z_49135

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008855 | ljcr-ds-49149-14521-4290-49149

(49149, 14521, 4290)-difference set in Z_49149

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008856 | ljcr-ds-49157-12289-3072-49157

(49157, 12289, 3072)-difference set in Z_49157

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008857 | ljcr-ds-4921-2296-1071-4921

(4921, 2296, 1071)-difference set in Z_4921

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008858 | ljcr-ds-4921-985-197-4921

(4921, 985, 197)-difference set in Z_4921

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008859 | ljcr-ds-49229-14292-4149-49229

(49229, 14292, 4149)-difference set in Z_49229

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008860 | ljcr-ds-49261-12316-3079-49261

(49261, 12316, 3079)-difference set in Z_49261

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008861 | ljcr-ds-49263-24631-12315-49263

(49263, 24631, 12315)-difference set in Z_49263

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008862 | ljcr-ds-49271-22361-10148-49271

(49271, 22361, 10148)-difference set in Z_49271

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008863 | ljcr-ds-49283-24641-12320-49283

(49283, 24641, 12320)-difference set in Z_49283

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008864 | ljcr-ds-49321-9865-1973-49321

(49321, 9865, 1973)-difference set in Z_49321

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008865 | ljcr-ds-49347-24673-12336-49347

(49347, 24673, 12336)-difference set in Z_49347

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008866 | ljcr-ds-49371-24685-12342-49371

(49371, 24685, 12342)-difference set in Z_49371

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008867 | ljcr-ds-49371-9875-1975-49371

(49371, 9875, 1975)-difference set in Z_49371

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008868 | ljcr-ds-4940-2245-1020-4940

(4940, 2245, 1020)-difference set in Z_4940

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008869 | ljcr-ds-4941-780-123-3-3-549

(4941, 780, 123)-difference set in Z_3 x Z_3 x Z_549

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008870 | ljcr-ds-4941-780-123-9-549

(4941, 780, 123)-difference set in Z_9 x Z_549

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008871 | ljcr-ds-49411-4941-494-49411

(49411, 4941, 494)-difference set in Z_49411

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008872 | ljcr-ds-49423-24711-12355-49423

(49423, 24711, 12355)-difference set in Z_49423

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008873 | ljcr-ds-49427-24713-12356-49427

(49427, 24713, 12356)-difference set in Z_49427

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008874 | ljcr-ds-49454-23273-10952-49454

(49454, 23273, 10952)-difference set in Z_49454

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008875 | ljcr-ds-49455-16590-5565-49455

(49455, 16590, 5565)-difference set in Z_49455

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008876 | ljcr-ds-49476-11875-2850-49476

(49476, 11875, 2850)-difference set in Z_49476

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008877 | ljcr-ds-49477-12369-3092-49477

(49477, 12369, 3092)-difference set in Z_49477

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008878 | ljcr-ds-495-247-123-3-165

(495, 247, 123)-difference set in Z_3 x Z_165

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008879 | ljcr-ds-4950-1617-528-4950

(4950, 1617, 528)-difference set in Z_4950

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008880 | ljcr-ds-49505-24480-12105-49505

(49505, 24480, 12105)-difference set in Z_49505

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008881 | ljcr-ds-49506-9901-1980-49506

(49506, 9901, 1980)-difference set in Z_49506

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008882 | ljcr-ds-49507-8251-1375-49507

(49507, 8251, 1375)-difference set in Z_49507

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008883 | ljcr-ds-49531-9906-1981-49531

(49531, 9906, 1981)-difference set in Z_49531

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008884 | ljcr-ds-49549-12388-3097-49549

(49549, 12388, 3097)-difference set in Z_49549

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008885 | ljcr-ds-49583-24791-12395-49583

(49583, 24791, 12395)-difference set in Z_49583

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008886 | ljcr-ds-4959-1407-399-4959

(4959, 1407, 399)-difference set in Z_4959

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008887 | ljcr-ds-49591-4960-496-49591

(49591, 4960, 496)-difference set in Z_49591

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008888 | ljcr-ds-4960-2205-980-4960

(4960, 2205, 980)-difference set in Z_4960

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008889 | ljcr-ds-49711-16570-5523-49711

(49711, 16570, 5523)-difference set in Z_49711

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008890 | ljcr-ds-49743-24871-12435-49743

(49743, 24871, 12435)-difference set in Z_49743

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008891 | ljcr-ds-49743-7106-1015-49743

(49743, 7106, 1015)-difference set in Z_49743

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008892 | ljcr-ds-4977-2177-952-4977

(4977, 2177, 952)-difference set in Z_4977

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008893 | ljcr-ds-49770-16485-5460-49770

(49770, 16485, 5460)-difference set in Z_49770

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008894 | ljcr-ds-49771-14616-4292-49771

(49771, 14616, 4292)-difference set in Z_49771

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008895 | ljcr-ds-49777-11713-2756-49777

(49777, 11713, 2756)-difference set in Z_49777

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008896 | ljcr-ds-49781-22401-10080-49781

(49781, 22401, 10080)-difference set in Z_49781

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008897 | ljcr-ds-49799-21343-9147-49799

(49799, 21343, 9147)-difference set in Z_49799

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008898 | ljcr-ds-49811-4981-498-49811

(49811, 4981, 498)-difference set in Z_49811

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008899 | ljcr-ds-49829-12457-3114-49829

(49829, 12457, 3114)-difference set in Z_49829

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008900 | ljcr-ds-49831-16611-5537-49831

(49831, 16611, 5537)-difference set in Z_49831

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008901 | ljcr-ds-49849-12865-3320-49849

(49849, 12865, 3320)-difference set in Z_49849

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008902 | ljcr-ds-49879-22168-9852-49879

(49879, 22168, 9852)-difference set in Z_49879

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008903 | ljcr-ds-49903-24951-12475-49903

(49903, 24951, 12475)-difference set in Z_49903

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008904 | ljcr-ds-4991-1996-798-4991

(4991, 1996, 798)-difference set in Z_4991

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008905 | ljcr-ds-4992-806-130-2-2-1248

(4992, 806, 130)-difference set in Z_2 x Z_2 x Z_1248

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008906 | ljcr-ds-4992-806-130-2-4-624

(4992, 806, 130)-difference set in Z_2 x Z_4 x Z_624

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008907 | ljcr-ds-4992-806-130-2-8-312

(4992, 806, 130)-difference set in Z_2 x Z_8 x Z_312

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008908 | ljcr-ds-4992-806-130-4-1248

(4992, 806, 130)-difference set in Z_4 x Z_1248

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008909 | ljcr-ds-4992-806-130-4-4-312

(4992, 806, 130)-difference set in Z_4 x Z_4 x Z_312

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008910 | ljcr-ds-4992-806-130-8-624

(4992, 806, 130)-difference set in Z_8 x Z_624

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008911 | ljcr-ds-49947-24973-12486-49947

(49947, 24973, 12486)-difference set in Z_49947

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008912 | ljcr-ds-49981-2500-125-49981

(49981, 2500, 125)-difference set in Z_49981

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008913 | ljcr-ds-49981-2941-173-49981

(49981, 2941, 173)-difference set in Z_49981

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008914 | ljcr-ds-49985-3905-305-49985

(49985, 3905, 305)-difference set in Z_49985

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008915 | ljcr-ds-49995-21427-9183-49995

(49995, 21427, 9183)-difference set in Z_49995

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008916 | ljcr-ds-5016-2125-900-5016

(5016, 2125, 900)-difference set in Z_5016

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008917 | ljcr-ds-5032-2107-882-5032

(5032, 2107, 882)-difference set in Z_5032

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008918 | ljcr-ds-5041-225-10-71-71

(5041, 225, 10)-difference set in Z_71 x Z_71

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008919 | ljcr-ds-5041-721-103-71-71

(5041, 721, 103)-difference set in Z_71 x Z_71

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008920 | ljcr-ds-5041-945-177-71-71

(5041, 945, 177)-difference set in Z_71 x Z_71

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008921 | ljcr-ds-5083-2058-833-5083

(5083, 2058, 833)-difference set in Z_5083

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008922 | ljcr-ds-5220-307-18-3-1740

(5220, 307, 18)-difference set in Z_3 x Z_1740

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008923 | ljcr-ds-5220-307-18-6-870

(5220, 307, 18)-difference set in Z_6 x Z_870

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008924 | ljcr-ds-5225-1960-735-5225

(5225, 1960, 735)-difference set in Z_5225

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008925 | ljcr-ds-5229-1308-327-5229

(5229, 1308, 327)-difference set in Z_5229

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008926 | ljcr-ds-5253-1717-561-5253

(5253, 1717, 561)-difference set in Z_5253

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008927 | ljcr-ds-5256-1051-210-3-1752

(5256, 1051, 210)-difference set in Z_3 x Z_1752

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008928 | ljcr-ds-5256-1051-210-5256

(5256, 1051, 210)-difference set in Z_5256

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008929 | ljcr-ds-5256-1051-210-6-876

(5256, 1051, 210)-difference set in Z_6 x Z_876

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008930 | ljcr-ds-5271-1055-211-5271

(5271, 1055, 211)-difference set in Z_5271

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008931 | ljcr-ds-5285-1321-330-5285

(5285, 1321, 330)-difference set in Z_5285

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008932 | ljcr-ds-5291-530-53-5291

(5291, 530, 53)-difference set in Z_5291

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008933 | ljcr-ds-5317-1329-332-5317

(5317, 1329, 332)-difference set in Z_5317

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008934 | ljcr-ds-5373-2449-1116-5373

(5373, 2449, 1116)-difference set in Z_5373

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008935 | ljcr-ds-5395-435-35-5395

(5395, 435, 35)-difference set in Z_5395

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008936 | ljcr-ds-54133-1041-20-54133

(54133, 1041, 20)-difference set in Z_54133

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008937 | ljcr-ds-5431-2535-1183-5431

(5431, 2535, 1183)-difference set in Z_5431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008938 | ljcr-ds-5440-148-4-2-2-2-2-2-170

(5440, 148, 4)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_170

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008939 | ljcr-ds-5440-148-4-2-2-2-2-340

(5440, 148, 4)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_340

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008940 | ljcr-ds-5440-148-4-2-2-2-680

(5440, 148, 4)-difference set in Z_2 x Z_2 x Z_2 x Z_680

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008941 | ljcr-ds-5440-148-4-2-2-4-340

(5440, 148, 4)-difference set in Z_2 x Z_2 x Z_4 x Z_340

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008942 | ljcr-ds-5440-148-4-2-4-680

(5440, 148, 4)-difference set in Z_2 x Z_4 x Z_680

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008943 | ljcr-ds-5440-148-4-4-4-340

(5440, 148, 4)-difference set in Z_4 x Z_4 x Z_340

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008944 | ljcr-ds-5440-148-4-8-680

(5440, 148, 4)-difference set in Z_8 x Z_680

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008945 | ljcr-ds-5447-1946-695-5447

(5447, 1946, 695)-difference set in Z_5447

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008946 | ljcr-ds-5461-1716-539-5461

(5461, 1716, 539)-difference set in Z_5461

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008947 | ljcr-ds-5461-2640-1276-5461

(5461, 2640, 1276)-difference set in Z_5461

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008948 | ljcr-ds-5490-1497-408-5490

(5490, 1497, 408)-difference set in Z_5490

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008949 | ljcr-ds-5565-1820-595-5565

(5565, 1820, 595)-difference set in Z_5565

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008950 | ljcr-ds-5593-1632-476-5593

(5593, 1632, 476)-difference set in Z_5593

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008951 | ljcr-ds-56056-1111-22-56056

(56056, 1111, 22)-difference set in Z_56056

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008952 | ljcr-ds-5613-1404-351-5613

(5613, 1404, 351)-difference set in Z_5613

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008953 | ljcr-ds-5621-1125-225-5621

(5621, 1125, 225)-difference set in Z_5621

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008954 | ljcr-ds-5629-1876-625-5629

(5629, 1876, 625)-difference set in Z_5629

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008955 | ljcr-ds-5669-1417-354-5669

(5669, 1417, 354)-difference set in Z_5669

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008956 | ljcr-ds-5677-1936-660-5677

(5677, 1936, 660)-difference set in Z_5677

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008957 | ljcr-ds-5720-1463-374-5720

(5720, 1463, 374)-difference set in Z_5720

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008958 | ljcr-ds-576-276-132-2-2-144

(576, 276, 132)-difference set in Z_2 x Z_2 x Z_144

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008959 | ljcr-ds-576-276-132-2-2-2-72

(576, 276, 132)-difference set in Z_2 x Z_2 x Z_2 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008960 | ljcr-ds-576-276-132-2-4-72

(576, 276, 132)-difference set in Z_2 x Z_4 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008961 | ljcr-ds-576-276-132-4-144

(576, 276, 132)-difference set in Z_4 x Z_144

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008962 | ljcr-ds-576-276-132-8-72

(576, 276, 132)-difference set in Z_8 x Z_72

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008963 | ljcr-ds-5767-961-160-5767

(5767, 961, 160)-difference set in Z_5767

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008964 | ljcr-ds-5823-1066-195-3-1941

(5823, 1066, 195)-difference set in Z_3 x Z_1941

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008965 | ljcr-ds-5824-648-72-2-2-1456

(5824, 648, 72)-difference set in Z_2 x Z_2 x Z_1456

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008966 | ljcr-ds-5824-648-72-2-2-2-2-364

(5824, 648, 72)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_364

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008967 | ljcr-ds-5824-648-72-2-2-2-728

(5824, 648, 72)-difference set in Z_2 x Z_2 x Z_2 x Z_728

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008968 | ljcr-ds-5824-648-72-2-2-4-364

(5824, 648, 72)-difference set in Z_2 x Z_2 x Z_4 x Z_364

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008969 | ljcr-ds-5824-648-72-2-2912

(5824, 648, 72)-difference set in Z_2 x Z_2912

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008970 | ljcr-ds-5824-648-72-2-4-728

(5824, 648, 72)-difference set in Z_2 x Z_4 x Z_728

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008971 | ljcr-ds-5824-648-72-4-1456

(5824, 648, 72)-difference set in Z_4 x Z_1456

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008972 | ljcr-ds-5824-648-72-4-4-364

(5824, 648, 72)-difference set in Z_4 x Z_4 x Z_364

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008973 | ljcr-ds-5824-648-72-8-728

(5824, 648, 72)-difference set in Z_8 x Z_728

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008974 | ljcr-ds-5859-203-7-3-1953

(5859, 203, 7)-difference set in Z_3 x Z_1953

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008975 | ljcr-ds-5859-203-7-3-3-651

(5859, 203, 7)-difference set in Z_3 x Z_3 x Z_651

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008976 | ljcr-ds-5859-203-7-5859

(5859, 203, 7)-difference set in Z_5859

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008977 | ljcr-ds-5900-1735-510-5900

(5900, 1735, 510)-difference set in Z_5900

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008978 | ljcr-ds-5929-456-35-11-539

(5929, 456, 35)-difference set in Z_11 x Z_539

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008979 | ljcr-ds-5929-456-35-7-847

(5929, 456, 35)-difference set in Z_7 x Z_847

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008980 | ljcr-ds-5929-456-35-77-77

(5929, 456, 35)-difference set in Z_77 x Z_77

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008981 | ljcr-ds-5931-1186-237-5931

(5931, 1186, 237)-difference set in Z_5931

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008982 | ljcr-ds-5951-1701-486-5951

(5951, 1701, 486)-difference set in Z_5951

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008983 | ljcr-ds-5951-1751-515-5951

(5951, 1751, 515)-difference set in Z_5951

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008984 | ljcr-ds-5971-1195-239-5971

(5971, 1195, 239)-difference set in Z_5971

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008985 | ljcr-ds-5989-1497-374-5989

(5989, 1497, 374)-difference set in Z_5989

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008986 | ljcr-ds-6021-645-69-3-3-669

(6021, 645, 69)-difference set in Z_3 x Z_3 x Z_669

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008987 | ljcr-ds-603-301-150-3-201

(603, 301, 150)-difference set in Z_3 x Z_201

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008988 | ljcr-ds-6100-2889-1368-6100

(6100, 2889, 1368)-difference set in Z_6100

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008989 | ljcr-ds-6104-2873-1352-6104

(6104, 2873, 1352)-difference set in Z_6104

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008990 | ljcr-ds-6115-2038-679-6115

(6115, 2038, 679)-difference set in Z_6115

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008991 | ljcr-ds-6119-1197-234-6119

(6119, 1197, 234)-difference set in Z_6119

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008992 | ljcr-ds-6149-1537-384-6149

(6149, 1537, 384)-difference set in Z_6149

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008993 | ljcr-ds-6149-1856-560-6149

(6149, 1856, 560)-difference set in Z_6149

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008994 | ljcr-ds-6149-2756-1235-6149

(6149, 2756, 1235)-difference set in Z_6149

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008995 | ljcr-ds-6160-2053-684-6160

(6160, 2053, 684)-difference set in Z_6160

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008996 | ljcr-ds-6188-2691-1170-6188

(6188, 2691, 1170)-difference set in Z_6188

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008997 | ljcr-ds-621-156-39-3-207

(621, 156, 39)-difference set in Z_3 x Z_207

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008998 | ljcr-ds-621-156-39-3-3-69

(621, 156, 39)-difference set in Z_3 x Z_3 x Z_69

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-008999 | ljcr-ds-6231-1246-249-6231

(6231, 1246, 249)-difference set in Z_6231

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009000 | ljcr-ds-6232-1675-450-6232

(6232, 1675, 450)-difference set in Z_6232

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009001 | ljcr-ds-6245-1561-390-6245

(6245, 1561, 390)-difference set in Z_6245

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009002 | ljcr-ds-6255-2597-1078-6255

(6255, 2597, 1078)-difference set in Z_6255

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009003 | ljcr-ds-6271-2926-1365-6271

(6271, 2926, 1365)-difference set in Z_6271

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009004 | ljcr-ds-6321-80-1-7-903

(6321, 80, 1)-difference set in Z_7 x Z_903

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009005 | ljcr-ds-6325-528-44-5-1265

(6325, 528, 44)-difference set in Z_5 x Z_1265

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009006 | ljcr-ds-6325-528-44-6325

(6325, 528, 44)-difference set in Z_6325

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009007 | ljcr-ds-6381-1596-399-6381

(6381, 1596, 399)-difference set in Z_6381

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009008 | ljcr-ds-639-232-84-3-213

(639, 232, 84)-difference set in Z_3 x Z_213

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009009 | ljcr-ds-639-232-84-639

(639, 232, 84)-difference set in Z_639

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009010 | ljcr-ds-6405-1601-400-6405

(6405, 1601, 400)-difference set in Z_6405

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009011 | ljcr-ds-6433-1408-308-6433

(6433, 1408, 308)-difference set in Z_6433

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009012 | ljcr-ds-6461-476-35-6461

(6461, 476, 35)-difference set in Z_6461

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009013 | ljcr-ds-6480-342-18-18-360

(6480, 342, 18)-difference set in Z_18 x Z_360

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009014 | ljcr-ds-6480-342-18-2-18-180

(6480, 342, 18)-difference set in Z_2 x Z_18 x Z_180

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009015 | ljcr-ds-6480-342-18-3-12-180

(6480, 342, 18)-difference set in Z_3 x Z_12 x Z_180

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009016 | ljcr-ds-6480-342-18-3-3-12-60

(6480, 342, 18)-difference set in Z_3 x Z_3 x Z_12 x Z_60

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009017 | ljcr-ds-6480-342-18-3-3-6-120

(6480, 342, 18)-difference set in Z_3 x Z_3 x Z_6 x Z_120

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009018 | ljcr-ds-6480-342-18-3-6-360

(6480, 342, 18)-difference set in Z_3 x Z_6 x Z_360

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009019 | ljcr-ds-6480-342-18-3-6-6-60

(6480, 342, 18)-difference set in Z_3 x Z_6 x Z_6 x Z_60

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009020 | ljcr-ds-6480-342-18-36-180

(6480, 342, 18)-difference set in Z_36 x Z_180

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009021 | ljcr-ds-6480-342-18-6-1080

(6480, 342, 18)-difference set in Z_6 x Z_1080

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009022 | ljcr-ds-6480-342-18-6-6-180

(6480, 342, 18)-difference set in Z_6 x Z_6 x Z_180

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009023 | ljcr-ds-6480-342-18-6-6-6-30

(6480, 342, 18)-difference set in Z_6 x Z_6 x Z_6 x Z_30

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009024 | ljcr-ds-6480-837-108-18-360

(6480, 837, 108)-difference set in Z_18 x Z_360

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009025 | ljcr-ds-6480-837-108-2-18-180

(6480, 837, 108)-difference set in Z_2 x Z_18 x Z_180

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009026 | ljcr-ds-6480-837-108-3-12-180

(6480, 837, 108)-difference set in Z_3 x Z_12 x Z_180

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009027 | ljcr-ds-6480-837-108-3-3-720

(6480, 837, 108)-difference set in Z_3 x Z_3 x Z_720

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009028 | ljcr-ds-6480-837-108-3-6-360

(6480, 837, 108)-difference set in Z_3 x Z_6 x Z_360

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009029 | ljcr-ds-6480-837-108-36-180

(6480, 837, 108)-difference set in Z_36 x Z_180

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009030 | ljcr-ds-6480-837-108-6-6-180

(6480, 837, 108)-difference set in Z_6 x Z_6 x Z_180

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009031 | ljcr-ds-6480-837-108-9-720

(6480, 837, 108)-difference set in Z_9 x Z_720

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009032 | ljcr-ds-6493-541-45-6493

(6493, 541, 45)-difference set in Z_6493

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009033 | ljcr-ds-65001-625-6-65001

(65001, 625, 6)-difference set in Z_65001

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009034 | ljcr-ds-6528-428-28-2-2-1632

(6528, 428, 28)-difference set in Z_2 x Z_2 x Z_1632

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009035 | ljcr-ds-6528-428-28-2-2-2-2-2-204

(6528, 428, 28)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_2 x Z_204

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009036 | ljcr-ds-6528-428-28-2-2-2-2-408

(6528, 428, 28)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_408

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009037 | ljcr-ds-6528-428-28-2-2-2-4-204

(6528, 428, 28)-difference set in Z_2 x Z_2 x Z_2 x Z_4 x Z_204

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009038 | ljcr-ds-6528-428-28-2-2-2-816

(6528, 428, 28)-difference set in Z_2 x Z_2 x Z_2 x Z_816

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009039 | ljcr-ds-6528-428-28-2-2-4-408

(6528, 428, 28)-difference set in Z_2 x Z_2 x Z_4 x Z_408

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009040 | ljcr-ds-6528-428-28-2-4-4-204

(6528, 428, 28)-difference set in Z_2 x Z_4 x Z_4 x Z_204

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009041 | ljcr-ds-6528-428-28-2-4-816

(6528, 428, 28)-difference set in Z_2 x Z_4 x Z_816

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009042 | ljcr-ds-6528-428-28-2-8-408

(6528, 428, 28)-difference set in Z_2 x Z_8 x Z_408

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009043 | ljcr-ds-6528-428-28-4-1632

(6528, 428, 28)-difference set in Z_4 x Z_1632

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009044 | ljcr-ds-6528-428-28-4-4-408

(6528, 428, 28)-difference set in Z_4 x Z_4 x Z_408

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009045 | ljcr-ds-6528-428-28-8-816

(6528, 428, 28)-difference set in Z_8 x Z_816

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009046 | ljcr-ds-6531-1960-588-6531

(6531, 1960, 588)-difference set in Z_6531

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009047 | ljcr-ds-6532-2800-1200-6532

(6532, 2800, 1200)-difference set in Z_6532

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009048 | ljcr-ds-6533-1633-408-6533

(6533, 1633, 408)-difference set in Z_6533

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009049 | ljcr-ds-6553-2457-921-6553

(6553, 2457, 921)-difference set in Z_6553

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009050 | ljcr-ds-6559-1093-182-6559

(6559, 1093, 182)-difference set in Z_6559

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009051 | ljcr-ds-6565-1641-410-6565

(6565, 1641, 410)-difference set in Z_6565

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009052 | ljcr-ds-6579-1496-340-6579

(6579, 1496, 340)-difference set in Z_6579

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009053 | ljcr-ds-6601-2376-855-6601

(6601, 2376, 855)-difference set in Z_6601

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009054 | ljcr-ds-6669-1668-417-6669

(6669, 1668, 417)-difference set in Z_6669

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009055 | ljcr-ds-6721-2625-1025-6721

(6721, 2625, 1025)-difference set in Z_6721

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009056 | ljcr-ds-675-337-168-15-45

(675, 337, 168)-difference set in Z_15 x Z_45

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009057 | ljcr-ds-675-337-168-3-3-75

(675, 337, 168)-difference set in Z_3 x Z_3 x Z_75

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009058 | ljcr-ds-675-337-168-5-135

(675, 337, 168)-difference set in Z_5 x Z_135

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009059 | ljcr-ds-67551-1351-27-67551

(67551, 1351, 27)-difference set in Z_67551

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009060 | ljcr-ds-676-325-156-13-52

(676, 325, 156)-difference set in Z_13 x Z_52

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009061 | ljcr-ds-676-325-156-26-26

(676, 325, 156)-difference set in Z_26 x Z_26

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009062 | ljcr-ds-6799-2266-755-6799

(6799, 2266, 755)-difference set in Z_6799

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009063 | ljcr-ds-6813-3276-1575-6813

(6813, 3276, 1575)-difference set in Z_6813

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009064 | ljcr-ds-6889-616-55-83-83

(6889, 616, 55)-difference set in Z_83 x Z_83

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009065 | ljcr-ds-6930-1353-264-6930

(6930, 1353, 264)-difference set in Z_6930

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009066 | ljcr-ds-6931-2520-916-6931

(6931, 2520, 916)-difference set in Z_6931

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009067 | ljcr-ds-6933-1733-433-6933

(6933, 1733, 433)-difference set in Z_6933

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009068 | ljcr-ds-6937-289-12-6937

(6937, 289, 12)-difference set in Z_6937

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009069 | ljcr-ds-69433-264-1-7-9919

(69433, 264, 1)-difference set in Z_7 x Z_9919

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009070 | ljcr-ds-6971-3281-1544-6971

(6971, 3281, 1544)-difference set in Z_6971

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009071 | ljcr-ds-6973-2989-1281-6973

(6973, 2989, 1281)-difference set in Z_6973

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009072 | ljcr-ds-6975-1585-360-6975

(6975, 1585, 360)-difference set in Z_6975

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009073 | ljcr-ds-6976-900-116-8-872

(6976, 900, 116)-difference set in Z_8 x Z_872

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009074 | ljcr-ds-6985-873-109-6985

(6985, 873, 109)-difference set in Z_6985

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009075 | ljcr-ds-703-325-150-703

(703, 325, 150)-difference set in Z_703

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009076 | ljcr-ds-7033-2344-781-7033

(7033, 2344, 781)-difference set in Z_7033

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009077 | ljcr-ds-7053-2624-976-7053

(7053, 2624, 976)-difference set in Z_7053

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009078 | ljcr-ds-7053-861-105-7053

(7053, 861, 105)-difference set in Z_7053

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009079 | ljcr-ds-7077-1769-442-7077

(7077, 1769, 442)-difference set in Z_7077

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009080 | ljcr-ds-7138-2197-676-7138

(7138, 2197, 676)-difference set in Z_7138

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009081 | ljcr-ds-7161-896-112-7161

(7161, 896, 112)-difference set in Z_7161

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009082 | ljcr-ds-7175-1055-155-5-1435

(7175, 1055, 155)-difference set in Z_5 x Z_1435

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009083 | ljcr-ds-7200-1565-340-7200

(7200, 1565, 340)-difference set in Z_7200

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009084 | ljcr-ds-7209-425-25-3-2403

(7209, 425, 25)-difference set in Z_3 x Z_2403

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009085 | ljcr-ds-7209-425-25-3-3-3-267

(7209, 425, 25)-difference set in Z_3 x Z_3 x Z_3 x Z_267

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009086 | ljcr-ds-7209-425-25-3-3-801

(7209, 425, 25)-difference set in Z_3 x Z_3 x Z_801

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009087 | ljcr-ds-7209-425-25-9-801

(7209, 425, 25)-difference set in Z_9 x Z_801

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009088 | ljcr-ds-7221-1445-289-7221

(7221, 1445, 289)-difference set in Z_7221

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009089 | ljcr-ds-7245-1812-453-7245

(7245, 1812, 453)-difference set in Z_7245

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009090 | ljcr-ds-7280-754-78-2-2-1820

(7280, 754, 78)-difference set in Z_2 x Z_2 x Z_1820

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009091 | ljcr-ds-7280-754-78-2-3640

(7280, 754, 78)-difference set in Z_2 x Z_3640

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009092 | ljcr-ds-7280-754-78-4-1820

(7280, 754, 78)-difference set in Z_4 x Z_1820

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009093 | ljcr-ds-729-273-102-27-27

(729, 273, 102)-difference set in Z_27 x Z_27

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009094 | ljcr-ds-729-273-102-3-3-3-27

(729, 273, 102)-difference set in Z_3 x Z_3 x Z_3 x Z_27

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009095 | ljcr-ds-729-273-102-3-3-3-3-3-3

(729, 273, 102)-difference set in Z_3 x Z_3 x Z_3 x Z_3 x Z_3 x Z_3

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009096 | ljcr-ds-729-273-102-3-3-3-3-9

(729, 273, 102)-difference set in Z_3 x Z_3 x Z_3 x Z_3 x Z_9

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009097 | ljcr-ds-729-273-102-3-3-9-9

(729, 273, 102)-difference set in Z_3 x Z_3 x Z_9 x Z_9

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009098 | ljcr-ds-729-273-102-3-9-27

(729, 273, 102)-difference set in Z_3 x Z_9 x Z_27

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009099 | ljcr-ds-729-273-102-9-9-9

(729, 273, 102)-difference set in Z_9 x Z_9 x Z_9

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009100 | ljcr-ds-7311-646-57-7311

(7311, 646, 57)-difference set in Z_7311

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009101 | ljcr-ds-7315-2968-1204-7315

(7315, 2968, 1204)-difference set in Z_7315

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009102 | ljcr-ds-7344-1050-150-12-612

(7344, 1050, 150)-difference set in Z_12 x Z_612

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009103 | ljcr-ds-7344-1050-150-2-3672

(7344, 1050, 150)-difference set in Z_2 x Z_3672

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009104 | ljcr-ds-7344-1050-150-4-1836

(7344, 1050, 150)-difference set in Z_4 x Z_1836

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009105 | ljcr-ds-7344-1050-150-6-1224

(7344, 1050, 150)-difference set in Z_6 x Z_1224

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009106 | ljcr-ds-736-196-52-2-2-184

(736, 196, 52)-difference set in Z_2 x Z_2 x Z_184

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009107 | ljcr-ds-736-196-52-2-2-2-2-46

(736, 196, 52)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_46

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009108 | ljcr-ds-736-196-52-2-2-2-92

(736, 196, 52)-difference set in Z_2 x Z_2 x Z_2 x Z_92

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009109 | ljcr-ds-736-196-52-2-368

(736, 196, 52)-difference set in Z_2 x Z_368

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009110 | ljcr-ds-736-196-52-2-4-92

(736, 196, 52)-difference set in Z_2 x Z_4 x Z_92

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009111 | ljcr-ds-736-196-52-4-184

(736, 196, 52)-difference set in Z_4 x Z_184

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009112 | ljcr-ds-7381-820-91-11-671

(7381, 820, 91)-difference set in Z_11 x Z_671

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009113 | ljcr-ds-741-185-46-741

(741, 185, 46)-difference set in Z_741

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009114 | ljcr-ds-7461-1865-466-7461

(7461, 1865, 466)-difference set in Z_7461

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009115 | ljcr-ds-7461-2985-1194-7461

(7461, 2985, 1194)-difference set in Z_7461

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009116 | ljcr-ds-7579-2106-585-7579

(7579, 2106, 585)-difference set in Z_7579

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009117 | ljcr-ds-7597-1900-475-7597

(7597, 1900, 475)-difference set in Z_7597

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009118 | ljcr-ds-7645-2353-724-7645

(7645, 2353, 724)-difference set in Z_7645

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009119 | ljcr-ds-765-192-48-3-255

(765, 192, 48)-difference set in Z_3 x Z_255

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009120 | ljcr-ds-7656-1531-306-7656

(7656, 1531, 306)-difference set in Z_7656

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009121 | ljcr-ds-7657-2784-1012-7657

(7657, 2784, 1012)-difference set in Z_7657

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009122 | ljcr-ds-768-118-18-16-48

(768, 118, 18)-difference set in Z_16 x Z_48

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009123 | ljcr-ds-768-118-18-4-192

(768, 118, 18)-difference set in Z_4 x Z_192

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009124 | ljcr-ds-768-118-18-8-96

(768, 118, 18)-difference set in Z_8 x Z_96

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009125 | ljcr-ds-7701-1925-481-7701

(7701, 1925, 481)-difference set in Z_7701

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009126 | ljcr-ds-7751-3751-1815-7751

(7751, 3751, 1815)-difference set in Z_7751

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009127 | ljcr-ds-7773-1073-148-7773

(7773, 1073, 148)-difference set in Z_7773

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009128 | ljcr-ds-7781-1945-486-7781

(7781, 1945, 486)-difference set in Z_7781

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009129 | ljcr-ds-781-300-115-781

(781, 300, 115)-difference set in Z_781

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009130 | ljcr-ds-7821-3520-1584-7821

(7821, 3520, 1584)-difference set in Z_7821

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009131 | ljcr-ds-783-391-195-3-261

(783, 391, 195)-difference set in Z_3 x Z_261

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009132 | ljcr-ds-784-378-182-14-56

(784, 378, 182)-difference set in Z_14 x Z_56

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009133 | ljcr-ds-784-378-182-2-14-28

(784, 378, 182)-difference set in Z_2 x Z_14 x Z_28

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009134 | ljcr-ds-784-378-182-2-2-14-14

(784, 378, 182)-difference set in Z_2 x Z_2 x Z_14 x Z_14

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009135 | ljcr-ds-784-378-182-28-28

(784, 378, 182)-difference set in Z_28 x Z_28

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009136 | ljcr-ds-7840-469-28-14-560

(7840, 469, 28)-difference set in Z_14 x Z_560

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009137 | ljcr-ds-7840-469-28-2-14-280

(7840, 469, 28)-difference set in Z_2 x Z_14 x Z_280

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009138 | ljcr-ds-7840-469-28-2-2-14-140

(7840, 469, 28)-difference set in Z_2 x Z_2 x Z_14 x Z_140

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009139 | ljcr-ds-7840-469-28-2-28-140

(7840, 469, 28)-difference set in Z_2 x Z_28 x Z_140

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009140 | ljcr-ds-7840-469-28-2-3920

(7840, 469, 28)-difference set in Z_2 x Z_3920

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009141 | ljcr-ds-7840-469-28-28-280

(7840, 469, 28)-difference set in Z_28 x Z_280

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009142 | ljcr-ds-7840-469-28-7-1120

(7840, 469, 28)-difference set in Z_7 x Z_1120

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009143 | ljcr-ds-7840-469-28-7840

(7840, 469, 28)-difference set in Z_7840

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009144 | ljcr-ds-79255-630-5-11-7205

(79255, 630, 5)-difference set in Z_11 x Z_7205

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009145 | ljcr-ds-79255-630-5-79255

(79255, 630, 5)-difference set in Z_79255

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009146 | ljcr-ds-7959-3634-1659-7959

(7959, 3634, 1659)-difference set in Z_7959

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009147 | ljcr-ds-8057-1008-126-8057

(8057, 1008, 126)-difference set in Z_8057

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009148 | ljcr-ds-8085-517-33-7-1155

(8085, 517, 33)-difference set in Z_7 x Z_1155

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009149 | ljcr-ds-8085-517-33-8085

(8085, 517, 33)-difference set in Z_8085

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009150 | ljcr-ds-80869-1173-17-80869

(80869, 1173, 17)-difference set in Z_80869

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009151 | ljcr-ds-8107-966-115-11-737

(8107, 966, 115)-difference set in Z_11 x Z_737

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009152 | ljcr-ds-8107-966-115-8107

(8107, 966, 115)-difference set in Z_8107

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009153 | ljcr-ds-8132-3807-1782-8132

(8132, 3807, 1782)-difference set in Z_8132

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009154 | ljcr-ds-816-326-130-2-408

(816, 326, 130)-difference set in Z_2 x Z_408

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009155 | ljcr-ds-8176-1200-176-8176

(8176, 1200, 176)-difference set in Z_8176

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009156 | ljcr-ds-8179-2350-675-8179

(8179, 2350, 675)-difference set in Z_8179

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009157 | ljcr-ds-8190-1293-204-8190

(8190, 1293, 204)-difference set in Z_8190

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009158 | ljcr-ds-8191-2016-496-8191

(8191, 2016, 496)-difference set in Z_8191

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009159 | ljcr-ds-8191-2080-528-8191

(8191, 2080, 528)-difference set in Z_8191

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009160 | ljcr-ds-8216-2015-494-8216

(8216, 2015, 494)-difference set in Z_8216

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009161 | ljcr-ds-8233-2401-700-8233

(8233, 2401, 700)-difference set in Z_8233

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009162 | ljcr-ds-8253-2064-516-8253

(8253, 2064, 516)-difference set in Z_8253

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009163 | ljcr-ds-8271-1655-331-8271

(8271, 1655, 331)-difference set in Z_8271

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009164 | ljcr-ds-82720-910-10-2-41360

(82720, 910, 10)-difference set in Z_2 x Z_41360

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009165 | ljcr-ds-82720-910-10-4-20680

(82720, 910, 10)-difference set in Z_4 x Z_20680

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009166 | ljcr-ds-8277-2069-517-8277

(8277, 2069, 517)-difference set in Z_8277

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009167 | ljcr-ds-8289-4033-1962-8289

(8289, 4033, 1962)-difference set in Z_8289

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009168 | ljcr-ds-8341-2641-836-8341

(8341, 2641, 836)-difference set in Z_8341

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009169 | ljcr-ds-8357-2089-522-8357

(8357, 2089, 522)-difference set in Z_8357

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009170 | ljcr-ds-8360-1287-198-8360

(8360, 1287, 198)-difference set in Z_8360

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009171 | ljcr-ds-8365-697-58-8365

(8365, 697, 58)-difference set in Z_8365

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009172 | ljcr-ds-837-209-52-3-279

(837, 209, 52)-difference set in Z_3 x Z_279

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009173 | ljcr-ds-837-209-52-3-3-93

(837, 209, 52)-difference set in Z_3 x Z_3 x Z_93

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009174 | ljcr-ds-8401-2401-686-8401

(8401, 2401, 686)-difference set in Z_8401

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009175 | ljcr-ds-8401-2976-1054-8401

(8401, 2976, 1054)-difference set in Z_8401

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009176 | ljcr-ds-8431-2811-937-8431

(8431, 2811, 937)-difference set in Z_8431

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009177 | ljcr-ds-8437-2812-937-8437

(8437, 2812, 937)-difference set in Z_8437

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009178 | ljcr-ds-847-423-211-11-77

(847, 423, 211)-difference set in Z_11 x Z_77

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009179 | ljcr-ds-8527-784-72-8527

(8527, 784, 72)-difference set in Z_8527

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009180 | ljcr-ds-8533-2133-533-8533

(8533, 2133, 533)-difference set in Z_8533

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009181 | ljcr-ds-855-183-39-3-285

(855, 183, 39)-difference set in Z_3 x Z_285

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009182 | ljcr-ds-8555-3290-1265-8555

(8555, 3290, 1265)-difference set in Z_8555

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009183 | ljcr-ds-8568-1977-456-8568

(8568, 1977, 456)-difference set in Z_8568

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009184 | ljcr-ds-8569-3808-1692-8569

(8569, 3808, 1692)-difference set in Z_8569

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009185 | ljcr-ds-8581-2145-536-8581

(8581, 2145, 536)-difference set in Z_8581

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009186 | ljcr-ds-8591-4295-2147-8591

(8591, 4295, 2147)-difference set in Z_8591

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009187 | ljcr-ds-8645-2161-540-8645

(8645, 2161, 540)-difference set in Z_8645

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009188 | ljcr-ds-8711-871-87-8711

(8711, 871, 87)-difference set in Z_8711

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009189 | ljcr-ds-8713-1089-136-8713

(8713, 1089, 136)-difference set in Z_8713

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009190 | ljcr-ds-8764-3175-1150-8764

(8764, 3175, 1150)-difference set in Z_8764

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009191 | ljcr-ds-8779-2926-975-8779

(8779, 2926, 975)-difference set in Z_8779

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009192 | ljcr-ds-8781-2196-549-8781

(8781, 2196, 549)-difference set in Z_8781

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009193 | ljcr-ds-8800-420-20-10-880

(8800, 420, 20)-difference set in Z_10 x Z_880

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009194 | ljcr-ds-8800-420-20-2-10-440

(8800, 420, 20)-difference set in Z_2 x Z_10 x Z_440

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009195 | ljcr-ds-8800-420-20-2-2-10-220

(8800, 420, 20)-difference set in Z_2 x Z_2 x Z_10 x Z_220

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009196 | ljcr-ds-8800-420-20-2-2-2-10-110

(8800, 420, 20)-difference set in Z_2 x Z_2 x Z_2 x Z_10 x Z_110

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009197 | ljcr-ds-8800-420-20-2-2-2-1100

(8800, 420, 20)-difference set in Z_2 x Z_2 x Z_2 x Z_1100

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009198 | ljcr-ds-8800-420-20-2-2-2200

(8800, 420, 20)-difference set in Z_2 x Z_2 x Z_2200

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009199 | ljcr-ds-8800-420-20-2-20-220

(8800, 420, 20)-difference set in Z_2 x Z_20 x Z_220

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009200 | ljcr-ds-8800-420-20-2-4-1100

(8800, 420, 20)-difference set in Z_2 x Z_4 x Z_1100

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009201 | ljcr-ds-8800-420-20-2-4400

(8800, 420, 20)-difference set in Z_2 x Z_4400

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009202 | ljcr-ds-8800-420-20-20-440

(8800, 420, 20)-difference set in Z_20 x Z_440

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009203 | ljcr-ds-8800-420-20-4-2200

(8800, 420, 20)-difference set in Z_4 x Z_2200

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009204 | ljcr-ds-8815-1470-245-8815

(8815, 1470, 245)-difference set in Z_8815

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009205 | ljcr-ds-8835-3787-1623-8835

(8835, 3787, 1623)-difference set in Z_8835

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009206 | ljcr-ds-8835-4417-2208-8835

(8835, 4417, 2208)-difference set in Z_8835

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009207 | ljcr-ds-8855-3496-1380-8855

(8855, 3496, 1380)-difference set in Z_8855

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009208 | ljcr-ds-891-445-222-3-3-3-33

(891, 445, 222)-difference set in Z_3 x Z_3 x Z_3 x Z_33

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009209 | ljcr-ds-891-445-222-3-3-99

(891, 445, 222)-difference set in Z_3 x Z_3 x Z_99

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009210 | ljcr-ds-891-445-222-9-99

(891, 445, 222)-difference set in Z_9 x Z_99

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009211 | ljcr-ds-8943-527-31-8943

(8943, 527, 31)-difference set in Z_8943

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009212 | ljcr-ds-896-180-36-2-2-2-112

(896, 180, 36)-difference set in Z_2 x Z_2 x Z_2 x Z_112

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009213 | ljcr-ds-896-180-36-2-2-224

(896, 180, 36)-difference set in Z_2 x Z_2 x Z_224

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009214 | ljcr-ds-896-180-36-2-4-112

(896, 180, 36)-difference set in Z_2 x Z_4 x Z_112

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009215 | ljcr-ds-896-180-36-4-224

(896, 180, 36)-difference set in Z_4 x Z_224

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009216 | ljcr-ds-896-180-36-8-112

(896, 180, 36)-difference set in Z_8 x Z_112

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009217 | ljcr-ds-8960-868-84-16-560

(8960, 868, 84)-difference set in Z_16 x Z_560

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009218 | ljcr-ds-8960-868-84-2-2-2240

(8960, 868, 84)-difference set in Z_2 x Z_2 x Z_2240

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009219 | ljcr-ds-8960-868-84-2-4-1120

(8960, 868, 84)-difference set in Z_2 x Z_4 x Z_1120

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009220 | ljcr-ds-8960-868-84-2-4480

(8960, 868, 84)-difference set in Z_2 x Z_4480

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009221 | ljcr-ds-8960-868-84-2-8-560

(8960, 868, 84)-difference set in Z_2 x Z_8 x Z_560

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009222 | ljcr-ds-8960-868-84-4-2240

(8960, 868, 84)-difference set in Z_4 x Z_2240

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009223 | ljcr-ds-8960-868-84-4-4-560

(8960, 868, 84)-difference set in Z_4 x Z_4 x Z_560

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009224 | ljcr-ds-8960-868-84-8-1120

(8960, 868, 84)-difference set in Z_8 x Z_1120

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009225 | ljcr-ds-8961-4096-1872-8961

(8961, 4096, 1872)-difference set in Z_8961

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009226 | ljcr-ds-8991-465-24-3-3-3-3-111

(8991, 465, 24)-difference set in Z_3 x Z_3 x Z_3 x Z_3 x Z_111

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009227 | ljcr-ds-8991-465-24-3-3-3-333

(8991, 465, 24)-difference set in Z_3 x Z_3 x Z_3 x Z_333

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009228 | ljcr-ds-8991-465-24-3-9-333

(8991, 465, 24)-difference set in Z_3 x Z_9 x Z_333

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009229 | ljcr-ds-900-435-210-15-60

(900, 435, 210)-difference set in Z_15 x Z_60

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009230 | ljcr-ds-900-435-210-30-30

(900, 435, 210)-difference set in Z_30 x Z_30

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009231 | ljcr-ds-9009-2816-880-9009

(9009, 2816, 880)-difference set in Z_9009

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009232 | ljcr-ds-9073-1296-185-9073

(9073, 1296, 185)-difference set in Z_9073

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009233 | ljcr-ds-909-228-57-3-303

(909, 228, 57)-difference set in Z_3 x Z_303

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009234 | ljcr-ds-910-405-180-910

(910, 405, 180)-difference set in Z_910

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009235 | ljcr-ds-9100-675-50-10-910

(9100, 675, 50)-difference set in Z_10 x Z_910

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009236 | ljcr-ds-9100-675-50-5-1820

(9100, 675, 50)-difference set in Z_5 x Z_1820

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009237 | ljcr-ds-9131-1661-302-9131

(9131, 1661, 302)-difference set in Z_9131

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009238 | ljcr-ds-9135-4567-2283-9135

(9135, 4567, 2283)-difference set in Z_9135

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009239 | ljcr-ds-921-161-28-921

(921, 161, 28)-difference set in Z_921

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009240 | ljcr-ds-9215-4607-2303-9215

(9215, 4607, 2303)-difference set in Z_9215

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009241 | ljcr-ds-925-385-160-5-185

(925, 385, 160)-difference set in Z_5 x Z_185

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009242 | ljcr-ds-931-186-37-7-133

(931, 186, 37)-difference set in Z_7 x Z_133

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009243 | ljcr-ds-931-280-84-7-133

(931, 280, 84)-difference set in Z_7 x Z_133

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009244 | ljcr-ds-931-465-232-7-133

(931, 465, 232)-difference set in Z_7 x Z_133

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009245 | ljcr-ds-9331-1866-373-9331

(9331, 1866, 373)-difference set in Z_9331

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009246 | ljcr-ds-936-375-150-2-468

(936, 375, 150)-difference set in Z_2 x Z_468

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009247 | ljcr-ds-936-375-150-2-6-78

(936, 375, 150)-difference set in Z_2 x Z_6 x Z_78

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009248 | ljcr-ds-936-375-150-3-312

(936, 375, 150)-difference set in Z_3 x Z_312

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009249 | ljcr-ds-936-375-150-6-156

(936, 375, 150)-difference set in Z_6 x Z_156

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009250 | ljcr-ds-936-375-150-936

(936, 375, 150)-difference set in Z_936

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009251 | ljcr-ds-9360-1911-390-9360

(9360, 1911, 390)-difference set in Z_9360

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009252 | ljcr-ds-9367-3346-1195-9367

(9367, 3346, 1195)-difference set in Z_9367

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009253 | ljcr-ds-9373-3124-1041-9373

(9373, 3124, 1041)-difference set in Z_9373

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009254 | ljcr-ds-9405-2352-588-9405

(9405, 2352, 588)-difference set in Z_9405

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009255 | ljcr-ds-9423-4711-2355-9423

(9423, 4711, 2355)-difference set in Z_9423

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009256 | ljcr-ds-9436-3996-1692-9436

(9436, 3996, 1692)-difference set in Z_9436

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009257 | ljcr-ds-944-369-144-2-472

(944, 369, 144)-difference set in Z_2 x Z_472

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009258 | ljcr-ds-944-369-144-944

(944, 369, 144)-difference set in Z_944

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009259 | ljcr-ds-945-177-33-3-3-105

(945, 177, 33)-difference set in Z_3 x Z_3 x Z_105

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009260 | ljcr-ds-945-177-33-3-315

(945, 177, 33)-difference set in Z_3 x Z_315

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009261 | ljcr-ds-9471-1895-379-9471

(9471, 1895, 379)-difference set in Z_9471

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009262 | ljcr-ds-9472-616-40-16-592

(9472, 616, 40)-difference set in Z_16 x Z_592

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009263 | ljcr-ds-9472-616-40-2-2-2-2-592

(9472, 616, 40)-difference set in Z_2 x Z_2 x Z_2 x Z_2 x Z_592

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009264 | ljcr-ds-9472-616-40-2-2-4-592

(9472, 616, 40)-difference set in Z_2 x Z_2 x Z_4 x Z_592

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009265 | ljcr-ds-9472-616-40-2-8-592

(9472, 616, 40)-difference set in Z_2 x Z_8 x Z_592

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009266 | ljcr-ds-9472-616-40-4-4-592

(9472, 616, 40)-difference set in Z_4 x Z_4 x Z_592

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009267 | ljcr-ds-950-365-140-5-190

(950, 365, 140)-difference set in Z_5 x Z_190

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009268 | ljcr-ds-9559-649-44-11-869

(9559, 649, 44)-difference set in Z_11 x Z_869

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009269 | ljcr-ds-960-274-78-2-2-2-120

(960, 274, 78)-difference set in Z_2 x Z_2 x Z_2 x Z_120

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009270 | ljcr-ds-960-274-78-2-2-240

(960, 274, 78)-difference set in Z_2 x Z_2 x Z_240

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009271 | ljcr-ds-960-274-78-2-4-120

(960, 274, 78)-difference set in Z_2 x Z_4 x Z_120

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009272 | ljcr-ds-960-274-78-2-480

(960, 274, 78)-difference set in Z_2 x Z_480

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009273 | ljcr-ds-960-274-78-4-240

(960, 274, 78)-difference set in Z_4 x Z_240

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009274 | ljcr-ds-960-274-78-8-120

(960, 274, 78)-difference set in Z_8 x Z_120

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009275 | ljcr-ds-961-256-68-31-31

(961, 256, 68)-difference set in Z_31 x Z_31

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009276 | ljcr-ds-963-222-51-3-321

(963, 222, 51)-difference set in Z_3 x Z_321

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009277 | ljcr-ds-963-222-51-963

(963, 222, 51)-difference set in Z_963

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009278 | ljcr-ds-963-481-240-3-321

(963, 481, 240)-difference set in Z_3 x Z_321

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009279 | ljcr-ds-9637-2409-602-9637

(9637, 2409, 602)-difference set in Z_9637

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009280 | ljcr-ds-9637-3796-1495-9637

(9637, 3796, 1495)-difference set in Z_9637

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009281 | ljcr-ds-9641-3616-1356-9641

(9641, 3616, 1356)-difference set in Z_9641

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009282 | ljcr-ds-975-487-243-5-195

(975, 487, 243)-difference set in Z_5 x Z_195

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009283 | ljcr-ds-9765-2441-610-9765

(9765, 2441, 610)-difference set in Z_9765

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009284 | ljcr-ds-981-196-39-3-327

(981, 196, 39)-difference set in Z_3 x Z_327

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009285 | ljcr-ds-981-196-39-981

(981, 196, 39)-difference set in Z_981

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009286 | ljcr-ds-981-441-198-3-327

(981, 441, 198)-difference set in Z_3 x Z_327

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009287 | ljcr-ds-9815-4207-1803-9815

(9815, 4207, 1803)-difference set in Z_9815

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009288 | ljcr-ds-9871-3291-1097-9871

(9871, 3291, 1097)-difference set in Z_9871

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009289 | ljcr-ds-98869-1177-14-98869

(98869, 1177, 14)-difference set in Z_98869

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009290 | ljcr-ds-990-345-120-3-330

(990, 345, 120)-difference set in Z_3 x Z_330

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009291 | ljcr-ds-990-345-120-990

(990, 345, 120)-difference set in Z_990

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009292 | ljcr-ds-9933-3056-940-9933

(9933, 3056, 940)-difference set in Z_9933

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009293 | ljcr-ds-9961-2656-708-9961

(9961, 2656, 708)-difference set in Z_9961

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

V3P-009294 | ljcr-ds-9969-3472-1209-9969

(9969, 3472, 1209)-difference set in Z_9969

the parameters pass the standard nonexistence tests, and no example is known.

来源状态 open | 2026-07-23 | 来源

D06 二色 Ramsey 数 | 48 行

R(s,t) 为使任意红蓝二染色必含红 K_s 或蓝 K_t 的最小完全图顶点数。记录为来源观察日的下界与上界之间的间隔。

原始维护者：Stanisław Radziszowski; Daniel Munro (Leaps in Bounds)

Cite S. Radziszowski, "Small Ramsey Numbers", Electronic Journal of Combinatorics, Dynamic Survey DS1, https://doi.org/10.37236/21, and credit Leaps in Bounds (D. Munro, https://leapsinbounds.org/) for the per-constant pages.

原始问题族入口

V3P-009295 | ramsey-r-10-10

Two-colour Ramsey number R(10, 10)

colourings of K_797 avoiding both are known, so R(10, 10) >= 798, and the survey's best upper bound is 16064.

来源状态 open | 2026-07-23 | 来源

V3P-009296 | ramsey-r-3-10

Two-colour Ramsey number R(3, 10)

colourings of K_39 avoiding both are known, so R(3, 10) >= 40, and the survey's best upper bound is 41.

来源状态 open | 2026-07-23 | 来源

V3P-009297 | ramsey-r-3-11

Two-colour Ramsey number R(3, 11)

colourings of K_46 avoiding both are known, so R(3, 11) >= 47, and the survey's best upper bound is 50.

来源状态 open | 2026-07-23 | 来源

V3P-009298 | ramsey-r-3-12

Two-colour Ramsey number R(3, 12)

colourings of K_52 avoiding both are known, so R(3, 12) >= 53, and the survey's best upper bound is 59.

来源状态 open | 2026-07-23 | 来源

V3P-009299 | ramsey-r-3-14

Two-colour Ramsey number R(3, 14)

colourings of K_66 avoiding both are known, so R(3, 14) >= 67, and the survey's best upper bound is 77.

来源状态 open | 2026-07-23 | 来源

V3P-009300 | ramsey-r-3-15

Two-colour Ramsey number R(3, 15)

colourings of K_73 avoiding both are known, so R(3, 15) >= 74, and the survey's best upper bound is 87.

来源状态 open | 2026-07-23 | 来源

V3P-009301 | ramsey-r-4-10

Two-colour Ramsey number R(4, 10)

colourings of K_91 avoiding both are known, so R(4, 10) >= 92, and the survey's best upper bound is 135.

来源状态 open | 2026-07-23 | 来源

V3P-009302 | ramsey-r-4-11

Two-colour Ramsey number R(4, 11)

colourings of K_101 avoiding both are known, so R(4, 11) >= 102, and the survey's best upper bound is 170.

来源状态 open | 2026-07-23 | 来源

V3P-009303 | ramsey-r-4-12

Two-colour Ramsey number R(4, 12)

colourings of K_127 avoiding both are known, so R(4, 12) >= 128, and the survey's best upper bound is 210.

来源状态 open | 2026-07-23 | 来源

V3P-009304 | ramsey-r-4-6

Two-colour Ramsey number R(4, 6)

colourings of K_35 avoiding both are known, so R(4, 6) >= 36, and the survey's best upper bound is 40.

来源状态 open | 2026-07-23 | 来源

V3P-009305 | ramsey-r-4-7

Two-colour Ramsey number R(4, 7)

colourings of K_48 avoiding both are known, so R(4, 7) >= 49, and the survey's best upper bound is 58.

来源状态 open | 2026-07-23 | 来源

V3P-009306 | ramsey-r-4-8

Two-colour Ramsey number R(4, 8)

colourings of K_58 avoiding both are known, so R(4, 8) >= 59, and the survey's best upper bound is 79.

来源状态 open | 2026-07-23 | 来源

V3P-009307 | ramsey-r-4-9

Two-colour Ramsey number R(4, 9)

colourings of K_72 avoiding both are known, so R(4, 9) >= 73, and the survey's best upper bound is 105.

来源状态 open | 2026-07-23 | 来源

V3P-009308 | ramsey-r-5-10

Two-colour Ramsey number R(5, 10)

colourings of K_148 avoiding both are known, so R(5, 10) >= 149, and the survey's best upper bound is 381.

来源状态 open | 2026-07-23 | 来源

V3P-009309 | ramsey-r-5-11

Two-colour Ramsey number R(5, 11)

colourings of K_182 avoiding both are known, so R(5, 11) >= 183, and the survey's best upper bound is 511.

来源状态 open | 2026-07-23 | 来源

V3P-009310 | ramsey-r-5-12

Two-colour Ramsey number R(5, 12)

colourings of K_202 avoiding both are known, so R(5, 12) >= 203, and the survey's best upper bound is 672.

来源状态 open | 2026-07-23 | 来源

V3P-009311 | ramsey-r-5-13

Two-colour Ramsey number R(5, 13)

colourings of K_232 avoiding both are known, so R(5, 13) >= 233, and the survey's best upper bound is 860.

来源状态 open | 2026-07-23 | 来源

V3P-009312 | ramsey-r-5-14

Two-colour Ramsey number R(5, 14)

colourings of K_266 avoiding both are known, so R(5, 14) >= 267, and the survey's best upper bound is 1081.

来源状态 open | 2026-07-23 | 来源

V3P-009313 | ramsey-r-5-15

Two-colour Ramsey number R(5, 15)

colourings of K_274 avoiding both are known, so R(5, 15) >= 275, and the survey's best upper bound is 1341.

来源状态 open | 2026-07-23 | 来源

V3P-009314 | ramsey-r-5-5

Two-colour Ramsey number R(5, 5)

colourings of K_42 avoiding both are known, so R(5, 5) >= 43, and the survey's best upper bound is 46.

来源状态 open | 2026-07-23 | 来源

V3P-009315 | ramsey-r-5-6

Two-colour Ramsey number R(5, 6)

colourings of K_58 avoiding both are known, so R(5, 6) >= 59, and the survey's best upper bound is 85.

来源状态 open | 2026-07-23 | 来源

V3P-009316 | ramsey-r-5-7

Two-colour Ramsey number R(5, 7)

colourings of K_79 avoiding both are known, so R(5, 7) >= 80, and the survey's best upper bound is 133.

来源状态 open | 2026-07-23 | 来源

V3P-009317 | ramsey-r-5-8

Two-colour Ramsey number R(5, 8)

colourings of K_100 avoiding both are known, so R(5, 8) >= 101, and the survey's best upper bound is 193.

来源状态 open | 2026-07-23 | 来源

V3P-009318 | ramsey-r-5-9

Two-colour Ramsey number R(5, 9)

colourings of K_132 avoiding both are known, so R(5, 9) >= 133, and the survey's best upper bound is 282.

来源状态 open | 2026-07-23 | 来源

V3P-009319 | ramsey-r-6-10

Two-colour Ramsey number R(6, 10)

colourings of K_203 avoiding both are known, so R(6, 10) >= 204, and the survey's best upper bound is 944.

来源状态 open | 2026-07-23 | 来源

V3P-009320 | ramsey-r-6-11

Two-colour Ramsey number R(6, 11)

colourings of K_261 avoiding both are known, so R(6, 11) >= 262, and the survey's best upper bound is 1346.

来源状态 open | 2026-07-23 | 来源

V3P-009321 | ramsey-r-6-12

Two-colour Ramsey number R(6, 12)

colourings of K_293 avoiding both are known, so R(6, 12) >= 294, and the survey's best upper bound is 1855.

来源状态 open | 2026-07-23 | 来源

V3P-009322 | ramsey-r-6-13

Two-colour Ramsey number R(6, 13)

colourings of K_346 avoiding both are known, so R(6, 13) >= 347, and the survey's best upper bound is 2499.

来源状态 open | 2026-07-23 | 来源

V3P-009323 | ramsey-r-6-15

Two-colour Ramsey number R(6, 15)

colourings of K_400 avoiding both are known, so R(6, 15) >= 401, and the survey's best upper bound is 4305.

来源状态 open | 2026-07-23 | 来源

V3P-009324 | ramsey-r-6-6

Two-colour Ramsey number R(6, 6)

colourings of K_101 avoiding both are known, so R(6, 6) >= 102, and the survey's best upper bound is 160.

来源状态 open | 2026-07-23 | 来源

V3P-009325 | ramsey-r-6-7

Two-colour Ramsey number R(6, 7)

colourings of K_114 avoiding both are known, so R(6, 7) >= 115, and the survey's best upper bound is 270.

来源状态 open | 2026-07-23 | 来源

V3P-009326 | ramsey-r-6-8

Two-colour Ramsey number R(6, 8)

colourings of K_133 avoiding both are known, so R(6, 8) >= 134, and the survey's best upper bound is 423.

来源状态 open | 2026-07-23 | 来源

V3P-009327 | ramsey-r-6-9

Two-colour Ramsey number R(6, 9)

colourings of K_182 avoiding both are known, so R(6, 9) >= 183, and the survey's best upper bound is 651.

来源状态 open | 2026-07-23 | 来源

V3P-009328 | ramsey-r-7-10

Two-colour Ramsey number R(7, 10)

colourings of K_291 avoiding both are known, so R(7, 10) >= 292, and the survey's best upper bound is 2119.

来源状态 open | 2026-07-23 | 来源

V3P-009329 | ramsey-r-7-11

Two-colour Ramsey number R(7, 11)

colourings of K_404 avoiding both are known, so R(7, 11) >= 405, and the survey's best upper bound is 3197.

来源状态 open | 2026-07-23 | 来源

V3P-009330 | ramsey-r-7-12

Two-colour Ramsey number R(7, 12)

colourings of K_416 avoiding both are known, so R(7, 12) >= 417, and the survey's best upper bound is 4665.

来源状态 open | 2026-07-23 | 来源

V3P-009331 | ramsey-r-7-13

Two-colour Ramsey number R(7, 13)

colourings of K_510 avoiding both are known, so R(7, 13) >= 511, and the survey's best upper bound is 6653.

来源状态 open | 2026-07-23 | 来源

V3P-009332 | ramsey-r-7-7

Two-colour Ramsey number R(7, 7)

colourings of K_204 avoiding both are known, so R(7, 7) >= 205, and the survey's best upper bound is 492.

来源状态 open | 2026-07-23 | 来源

V3P-009333 | ramsey-r-7-8

Two-colour Ramsey number R(7, 8)

colourings of K_218 avoiding both are known, so R(7, 8) >= 219, and the survey's best upper bound is 832.

来源状态 open | 2026-07-23 | 来源

V3P-009334 | ramsey-r-7-9

Two-colour Ramsey number R(7, 9)

colourings of K_251 avoiding both are known, so R(7, 9) >= 252, and the survey's best upper bound is 1368.

来源状态 open | 2026-07-23 | 来源

V3P-009335 | ramsey-r-8-10

Two-colour Ramsey number R(8, 10)

colourings of K_342 avoiding both are known, so R(8, 10) >= 343, and the survey's best upper bound is 4402.

来源状态 open | 2026-07-23 | 来源

V3P-009336 | ramsey-r-8-11

Two-colour Ramsey number R(8, 11)

colourings of K_456 avoiding both are known, so R(8, 11) >= 457, and the survey's best upper bound is 7040.

来源状态 open | 2026-07-23 | 来源

V3P-009337 | ramsey-r-8-13

Two-colour Ramsey number R(8, 13)

colourings of K_816 avoiding both are known, so R(8, 13) >= 817, and the survey's best upper bound is 27485.

来源状态 open | 2026-07-23 | 来源

V3P-009338 | ramsey-r-8-15

Two-colour Ramsey number R(8, 15)

colourings of K_872 avoiding both are known, so R(8, 15) >= 873, and the survey's best upper bound is 63609.

来源状态 open | 2026-07-23 | 来源

V3P-009339 | ramsey-r-8-8

Two-colour Ramsey number R(8, 8)

colourings of K_281 avoiding both are known, so R(8, 8) >= 282, and the survey's best upper bound is 1518.

来源状态 open | 2026-07-23 | 来源

V3P-009340 | ramsey-r-8-9

Two-colour Ramsey number R(8, 9)

colourings of K_328 avoiding both are known, so R(8, 9) >= 329, and the survey's best upper bound is 2662.

来源状态 open | 2026-07-23 | 来源

V3P-009341 | ramsey-r-9-10

Two-colour Ramsey number R(9, 10)

colourings of K_580 avoiding both are known, so R(9, 10) >= 581, and the survey's best upper bound is 8675.

来源状态 open | 2026-07-23 | 来源

V3P-009342 | ramsey-r-9-9

Two-colour Ramsey number R(9, 9)

colourings of K_564 avoiding both are known, so R(9, 9) >= 565, and the survey's best upper bound is 4956.

来源状态 open | 2026-07-23 | 来源

E 可追溯性、排除日志与后续边界

正文编号校验目标：20,360 个 V3 编号；附录编号校验目标：9,342 个 V3P 编号。每条记录保留原来源主键，并与 PDF 起始页对应。附件包括保留记录和排除清单，便于追查一次筛查为何纳入或暂缓某条材料。

各公开下载包 SHA-256

bespokelabsai/conjecture-bench
0351d52972268961c1146e05728f7664dbbe4fdd1253f7f314bf0aa495f03c0b

davisrbr/conjectures-arxiv
25bd62b9c6ee61f2ceded4d4e28531d2e57571da02b7e234fa366cc3ae278598

epoch-research/LeanOpenProblems-results
2aa6be3d67e11d5ce9270999f8acf3125e479bd36459ebb6baabaf7a8bf78c61

graph-theory-AI/graph-conjectures
5a8fb47b7de400a322414c54a33ec3d1d7d3922fea46b8f790cb0cd2fcf541a7

oeis/oeisdata
1da549afe7b6dd95eff5ba4d22dbd38fcf7733dda2e7fa4ac7722519a3c94f2d

本卷仍未覆盖的部分

未系统取得所有付费全文、所有个人网页、所有中文/俄文等语言材料、未数字化书籍、邮件与口头交流；没有遍历 OEIS 与论坛的全部修订历史，也没有枚举一项问题族的所有可能实例。对无法打开的网页，不根据标题臆造猜想正文。

后续扩大收录时，应继续以三卷机器可读记录为基准，并优先补全原文和进行数学语义归并。没有重复的 URL、ID 或文字，并不能证明没有重复的数学问题。